



(51) International Patent Classification:

A61K 31/5377 (2006.01) *A61P 17/00* (2006.01)
A61K 31/047 (2006.01) *A61P 17/06* (2006.01)
A61K 31/4436 (2006.01) *A61N 5/06* (2006.01)
A61K 9/00 (2006.01)

(21) International Application Number:

PCT/US2019/021037

(22) International Filing Date:

06 March 2019 (06.03.2019)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/639,742 07 March 2018 (07.03.2018) US

(71) Applicants: **ATHENEX HK INNOVATIVE LIMITED** [CN/CN]; Unit 608-613 IC Development Centre, No 6 Science Park West Ave, Hong Kong Science Park, Hong Kong (CN). **AVALON BIOMEDICAL (MAN-**

AGEMENT) LIMITED [CN/CN]; 10/F Fung House, 19-20 Connaught Road Central, Hong Kong (CN).

(72) Inventor; and

(71) Applicant: **LAU, Johnson Yiu-Nam** [US/US]; 6230 Cedar Creek Drive, Houston, Texas 77057 (US).

(72) Inventors: **VERONE-BOYLE, Alissa Rae**; 350 Parkwood Ave, Kenmore, New Jersey 14217 (US). **WONG, Chun-Ho**; Flat 31D, Block 6, Jubilee Garden, Lok King Street, Fo Tan, Shatin, Hong Kong (CN). **BU, Yahao**; 36 Troy View Lane, Williamsville, New York 14221 (US). **CUTLER, Murray John**; 8 Phyllis Street, Fort Erie, Ontario L2A 3Y2 (CA). **BELKO, Krista Elizabeth**; 10115 Keller Road, Clarence Center, New York 14032 (US). **KWAN, Min-Fun Rudolf**; 9 Friar Tuck Circle, Summit, New Jersey 07901 (US).

(74) Agent: **BRILLHART, Kurt L.** et al.; Fish IP Law, LLP, 2603 Main Street, Ste 1000, Irvine, California 92614 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM,

(54) Title: COMPOSITIONS AND METHODS FOR TREATING HYPERPROLIFERATIVE SKIN DISORDERS




FIG. 1

(57) Abstract: Compositions and methods are provided that are effective in treating skin hyperproliferative disorders. These include a topical pharmaceutical preparation that includes a compound that has two or more of a SRC-kinase inhibiting activity, a tubulin polymerization inhibiting activity, an activity that arrests the cell cycle at G2/M, and an activity that induces apoptosis. The compound KX01 is provided as an example of such a compound. The topical pharmaceutical preparation further includes a vitamin D derivative or a retinoid, where the combination provides a synergistic effect. Use of the topical pharmaceutical preparation can be combined with exposure to blue, UVA, or UVB light, which provides a synergistic effect.



AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

- (84) Designated States** (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

— *with international search report (Art. 21(3))*

COMPOSITIONS AND METHODS FOR TREATING HYPERPROLIFERATIVE SKIN DISORDERS

[0001] This application claims the benefit of United States Provisional Application No. 62/639,742 filed on March 7, 2018. These and all other referenced extrinsic materials are incorporated herein by reference in their entirety. Where a definition or use of a term in a reference that is incorporated by reference is inconsistent or contrary to the definition of that term provided herein, the definition of that term provided herein is deemed to be controlling.

Field of the Invention

[0002] The field of the invention is combination therapies for treating hyperproliferative skin disorders.

Background

[0003] The following description includes information that may be useful in understanding the present invention. It is not an admission that any of the information provided herein is prior art or relevant to the presently claimed invention, or that any publication specifically or implicitly referenced is prior art.

[0004] The background description includes information that may be useful in understanding the present invention. It is not an admission that any of the information provided herein is prior art or relevant to the presently claimed invention, or that any publication specifically or implicitly referenced is prior art.

[0005] Cell proliferation disorders of the skin such as psoriasis, actinic keratosis, and non-melanoma skin cancers (e.g., basal cell carcinoma and squamous cell carcinoma) are common, but treatments often lack the desired effectiveness, have inconvenient dosing regimens, and/or have severe side effects.

[0006] Current pharmaceutical approaches to treating skin proliferation disorders (such as psoriasis) include the use of corticosteroids, vitamin D analogs, and retinoids. In some instances phototherapy is used to treat skin proliferation disorders, and can be used in combination with pharmaceutical interventions. Phototherapy typically involves exposure to UVB and/or UVA

light, and can include treatment with photosensitizers (e.g. using psoralen in combination with exposure to UVA light).

[0007] More recently, biologics have been used to treat skin proliferative disorders. These are typically monoclonal antibodies or antibody fragments that target specific components of the immune system or cytokines involved in inflammation. Examples of such biologics include Humira™, Amevive™, ABT-874, Enbrel™, Remicade™, and Stelara™. Unfortunately such biologics are and expensive, and the use of such biologics is associated with increased incidence of cancer and certain types of infection.

[0008] Actinic keratoses are common pre-cancerous skin lesions that show excessive proliferation of skin cells, and are the third most common reason for consulting a dermatologist. These can be treated by excision, freezing, and/or the application of pharmaceutical agents to the lesion. Drugs used to treat these lesions include 5-fluorouracil (5-FU), diclofenac, ingenol, and imiquimod. Unfortunately, side effects (which include discomfort, redness, and blistering) and/or inconvenient dosing regimens limit patient compliance, and therefore, the efficacy of these approaches.

[0009] Calcipotriol is an FDA-approved topical medication for treatment of skin hyperproliferative disorders, and is hypothesized to act via induction of thymic stromal lymphopoietin ("TSLP") expression. The immune effects of TSLP expression have been shown to include antitumor immunity in barrier-defective skin (see Trevor J. Cunningham et al., Randomized trial of calcipotriol combined with 5-fluorouracil for skin cancer precursor immunotherapy, 127(1) J. Clinical Investigation 106–16 (2017)). All publications identified herein are incorporated by reference to the same extent as if each individual publication or patent application were specifically and individually indicated to be incorporated by reference. Where a definition or use of a term in an incorporated reference is inconsistent or contrary to the definition of that term provided herein, the definition of that term provided herein applies and the definition of that term in the reference does not apply.

[0010] Calcipotriol has been used in combination with other drugs. For example, treatment with calcipotriol ointment combined with 5-fluorouracil cream can decrease the number of actinic keratoses compared to treatment with 5-fluorouracil alone. The combined use of calcipotriol

ointment and 5-fluorouracil cream on face, scalp, or upper extremities has been found to induce TSLP, HLA class II, and natural killer cell group 2D ligand expression in keratinocytes of these lesions. Calcipotriol and 5-fluorouracil are thought to activate a CD4+ T cell-mediated immunity against actinic keratosis. However, this treatment regimen maintains the limitations of 5-fluorouracil, including side effects that limit patient compliance.


[0011] Thus, there is still a need for treatments for cell hyperproliferation disorders that are both effective and well tolerated.

Summary of The Invention

[0012] The inventive subject matter provides apparatus, systems and methods in which a ---

[0013] One embodiment of the inventive concept is a pharmaceutical composition that includes a first compound having an SRC-kinase inhibition activity, an inhibition of tubulin polymerization activity, a cell cycle arrest at G2/M activity, and/or a induction of apoptosis activity, and a second compound this is a vitamin D analog or a retinoid, where the first compound and the second compound are present in amounts sufficient to treat a hyperproliferative disorder in an affected area. Such composition can be formulated for topical application, for example as a foam, a cream, a paste, an ointment, a gel, a solution, a liquid suspension, a droplet suspension, an aerosol, and/or a powder. The vitamin D analog can also have a G0/G1 arrest activity, an anti-proliferation activity, an anti-inflammation activity, an induction of keratinocyte differentiation activity, an activity that decreases the number of pro-inflammatory T-cells, an apoptosis inducing activity.

[0014] In such a compositions the first compound can have a structure according to Formula I.



or a salt, solvate, hydrate, or prodrug thereof, wherein: T is a bond;

Xy is CZ, CY, N, or N—O;

Xz is CZ, CY, N, or N—O;

at least one of Xy and Xz is CZ;

Y is selected from hydrogen, hydroxyl, halogen, (C₁, C₂, C₃, C₄, C₅, or C₆) alkyl, C₁, C₂, C₃, C₄, C₅, or C₆ alkoxy, O- (C₁, C₂, C₃, C₄, C₅, or C₆) alkylaryl, and O-benzyl;

Xa is CRa or N, or N—O;


Xb is CRb, N, or N—O;

Xc is CRc or N, or N—O;

Xd is CRd or N, or N—O;

Xe is CRe, N, or N—O

Ra, Rb, Rc, Rd, Re, R₄, R₅, and R₆ are, independently, hydrogen, hydroxyl, halogen, C₁, C₂, C₃, C₄, C₅, or C₆ alkoxy, O-lower (C₁, C₂, C₃, C₄, C₅, or C₆) alkyl-aryl, O-benzyl, C₁, C₂, C₃, C₄, C₅, or C₆ alkyl-OH, C₁, C₂, C₃, C₄, C₅, or C₆alkyl-O- (C₁, C₂, C₃, C₄, C₅, or C₆) alkyl, COOH, COO- (C₁, C₂, C₃, C₄, C₅, or C₆) alkyl, SO₂H, SO₂ (C₁, C₂, C₃, C₄, C₅, or C₆) alkyl,



where W is H, or C₁, C₂, C₃, C₄, C₅, or C₆ alkyl, C₁, C₂, C₃, C₄, C₅, or C₆ alkyl-aryl;

V is a bond, —CH₂—, —CH₂CH₂—, —CH₂CH₂CH₂—, —O—CH₂—, —OCH₂CH₂— or —OCH₂CH₂CH₂—;

R₁₂, R₁₃, R₁₄, R₁₅, R₁₆, R₁₇, and R₁₈, are, independently, H or C₁, C₂, C₃, C₄, C₅, or C₆ alkyl;


Z is: (CHR₁)_n—C(O)—NR₂(CHR₃)_m—Ar, where Ar is a substituted or unsubstituted aryl or nitrogen-containing heteroaryl group, R₁, R₂, and R₃ are independently H or C₁, C₂, C₃, C₄, C₅, or C₆ alkyl;

n is 0, 1, or 2; and

m is 1 or 2,

wherein at least one of Xa, Xb, Xc, Xd and Xe is N.

[0015] In a preferred embodiment the first compound can be KX01, which has the structure shown in Formula 2.



Formula 2

[0016] In such a composition the second compound can be a vitamin D3 analog, such as calcipotriol (calipotriene). The composition can be formulated to provide a molar ratio of the first compound to the second compound of from 160:3 to 5:243 at the affected area. In some embodiments the composition this molar ratio is selected to provide an effect at the affected area that exceeds the additive effect of the first compound and the second compound when applied individually at corresponding concentrations. In other embodiments the second compound is a retinoid, such as an acetylenic retinoid or tazarotene. In some embodiments the second compound is a vitamin D analog, and the composition also includes a retinoid.

[0017] Another embodiment of the inventive concept is a method of treating a skin condition (such as actinic keratosis or psoriasis) by providing a topical preparation (e.g. a foam, a cream, a paste, an ointment, a gel, a solution, a liquid suspension, a droplet suspension, an aerosol, and/or a powder) that includes a first compound and a second compound, where the first compound has at least two of an SRC-kinase inhibiting activity, a tubulin polymerization inhibiting activity, a cell cycle arrest at G2/M activity, and/or an activity that induces apoptosis, and where the second compound includes a vitamin D analog (such as a vitamin D3 analog or calcipotriol) or a retinoid (such as an acetylenic retinoid or tazarotene), where the first compound and the second compound are present in amounts sufficient to provide a reduction or elimination of a hyperproliferative disorder in an affected area. The molar ratio molar ratio of the first compound to the second compound can range from 160:3 to 5:243 or from 1:1 to 1:2,500 at the affected area. In some embodiments the first compound can be KX01 (as described above). In some

embodiments the second compound is a vitamin D3 analog or calcipotriol (calcipotriene). In some embodiments the second compound is a vitamin D analog, and the topical composition also includes a retinoid. The method also includes a step of applying the topical preparation to an area affected by the skin condition on a schedule that is effective to treat the skin condition (e.g. from once a week to four times a day). Skin conditions that can be treated in this fashion include actinic keratosis and psoriasis. The vitamin D analog can have a G0/G1 arresting activity, an anti-proliferation activity, an anti-inflammatory activity, an activity that induces keratinocyte differentiation, an activity that decreases the number of pro-inflammatory T-cells, and/or an activity that induces apoptosis. The retinoid can have an activity that induces keratinocytes proliferation, modulates epidermal differentiation, stimulates extracellular matrix production, and/or alters sebocyte differentiation. In a preferred embodiment the first topical composition includes the first compound in a first amount and the second topical composition comprises the second compound in a second amount, where the first amount and the second amount are selected to provide a combined therapeutic effect. This combined therapeutic effect is greater than the sum of the therapeutic effects provided by application of the first compound at the first amount and by application of the second compound at the second amount.

[0018] Another embodiment of the inventive concept is a kit for treating a skin cell hyperproliferation disorder, having a first topical composition that includes a first compound (such as KX01, as shown above) with two or more of a SRC-kinase inhibiting activity, a tubulin polymerization inhibiting activity, an activity that arrests the cell cycle at G2/M, and an activity that induces apoptosis, along with a second topical composition that includes a second compound such as a vitamin D analog (such as a vitamin D3 analog and/or calcipotriol) or a retinoid (such as an acetylenic retinoid and/or tazarotene). In some embodiments the kit includes both a vitamin D analog and a retinoid. The first compound and the second compound are present in their respective topical compositions in amounts sufficient to provide a reduction or elimination of the skin cell hyperproliferation disorder when the first and second topical compositions are applied individually according to schedule. The molar ratio of the first compound to the second compound can range from 160:3 to 5:243 or from 1:1 to 1:2,500. Such topical compositions can be provided as a foam, a cream, a paste, an ointment, a gel, a solution, a liquid suspension, a droplet suspension, an aerosol, and/or a powder. In a preferred embodiment the first topical composition includes the first compound in a first amount and the second topical

composition comprises the second compound in a second amount, where the first amount and the second amount are selected to provide a combined therapeutic effect. This combined therapeutic effect is greater than the sum of the therapeutic effects provided by application of the first compound at the first amount and by application of the second compound at the second amount.

[0019] Another embodiment of the inventive concept is a method of increasing two or more of a SRC-kinase inhibiting activity, a tubulin polymerization inhibition activity, an activity that arrests the cell cycle at G2/M, and an activity that induces apoptosis produced by a first compound in a pharmaceutical composition. This is accomplished by including a second compound (such as a vitamin D analog or a retinoid) in the pharmaceutical composition, where the molar ratio of the first compound and the second compound is selected to enhance the activities in more than an additive manner. The molar ratio of the first compound to the second compound can range from 160:3 to 5:243 or from 1:1 to 1:2,500. The first compound can be KX01 (as shown above). The vitamin D analog can be a vitamin D3 analog and/or calcipotriol. The retinoid can be an acetylenic retinoid and/or tazarotene. In embodiments where the second compound is a vitamin D analog the composition can further include a retinoid.

[0020] Another embodiment of the inventive concept is the use of a first compound (such as Kx)1, as shown above) and a second compound in preparing a medicament for treating a skin condition (such as actinic keratosis and/or psoriasis), where the first compound has two or more of a SRC-kinase inhibiting activity, a tubulin polymerization activity, an activity that arrests the cell cycle at G2/M, and an activity that induces apoptosis, and where the second compound includes a vitamin D analog (e.g. a vitamin D3 analog and/or calcipotriol) or a retinoid (e.g. an acetylenic retinoid and/or tazarotene). The first compound and the second compound are present in the medicament in amounts sufficient to provide a reduction or elimination of a hyperproliferative disorder by applying the medicament to an area affected by the skin condition on a schedule effective to treat the skin condition (e.g. at a frequency of once a week to four times a day). The molar ratio of the first compound to the second compound can range from 160:3 to 5:243 or from 1:1 to 1:2,500. The vitamin D analog can be selected to have a G0/G1 arresting activity, an anti-proliferation activity, an anti-inflammation activity, an activity that induces keratinocyte differentiation, an activity that decreases the number of pro-inflammatory T-cells, and/or an activity that induces apoptosis. In embodiments where the

second compound is a vitamin D analog the medicament can further include a retinoid. Such a medicament can be formulated to be used for topical application, for example as a foam, a cream, a paste, an ointment, a gel, a solution, a liquid suspension, a droplet suspension, an aerosol, and/or a powder. In a preferred embodiment the medicament includes the first compound in a first amount and the second compound in a second amount, where the first amount and the second amount are selected to provide a combined therapeutic effect. This combined therapeutic effect is greater than the sum of the therapeutic effects provided by application of the first compound at the first amount and by application of the second compound at the second amount.

[0021] Another embodiment of the inventive concept is a method of treating a proliferative condition (e.g. actinic keratosis and/or psoriasis) by providing a topical preparation that includes a first compound, where the first compound (e.g. KX01, as shown above) has two or more of a SRC-kinase inhibiting activity, a tubulin polymerization inhibiting activity, an activity that arrests the cell cycle at G2/M, and an activity that induces apoptosis and applying the topical preparation to an area affected by the skin condition on a schedule effective to treat the skin condition, in combination with directing a blue light (e.g. having a wavelength from 280nm to 500 nm, which can be provided as either a narrow or broad band emission) onto the area affected by the skin condition at a specified light flux and for a specified exposure time. Suitable light sources include LED/LCD and fluorescent lights, and can emit visible blue light (e.g. 311 nm to 460 nm), UVA, and/or UVB). The topical preparation can be formulated as formulated as a foam, a cream, a paste, an ointment, a gel, a solution, a liquid suspension, a droplet suspension, an aerosol, and/or a powder. In some embodiments the topical preparation further includes a vitamin D analog (e.g. a vitamin D3 analog and/or calcipotriol) and/or a retinoid (e.g. an acetylenic retinoid and/or tazarotene). In such embodiments the retinoid can have an activity that induces keratinocytes to proliferate, an activity that modulates epidermal differentiation, an activity that stimulates extracellular matrix production, and/or an activity that alters of sebocyte differentiation. In some embodiments the topical preparation includes the first compound (and second compound, if present) in an amount(s) that provides a first therapeutic effect and the specified light flux and exposure time are provided that can produce a second therapeutic effect, however the method provides a third therapeutic effect that is greater than the sum of a first therapeutic effect and the second therapeutic effect.

[0022] Various objects, features, aspects and advantages of the inventive subject matter will become more apparent from the following detailed description of preferred embodiments, along with the accompanying drawing figures in which like numerals represent like components.

Brief Description of The Drawings

[0023] FIG. 1: Results of an experiment comparing cell growth in response to KX01, Calcipotriol, and KX01 + calcipotriol at a 1:5 ratio.

[0024] FIG. 2: Results of an experiment comparing cell growth in response to KX01, Calcipotriol, and KX01 + calcipotriol at a 1:2 ratio.

[0025] FIG. 3: Results from two experiments comparing cell growth in response to KX01, Calcipotriol, and KX01 + calcipotriol at a 2:1 ratio.

[0026] FIG. 4: Inhibition of human keratinocyte growth versus KX01 concentration (treatment duration: 72 hours; cell growth measured using the MTT assay). The IC_{50} of KX01 = 41.9 nM.

[0027] FIG.5: Inhibition of human keratinocyte growth versus calcipotriol concentration (treatment duration: 72 hours; cell growth measured using the MTT assay). The IC_{50} of calcipotriol = 95.5 nM.

[0028] FIG. 6: Dose response surface of cell viability versus calcipotriol concentration versus KX01 concentration. Synergistic combinations of calcipotriol and KX01 are below the surface of additivity (indicating enhanced inhibition of growth), and antagonistic combinations are above the surface of additivity (indicative of reduced inhibition of growth).

[0029] FIG. 7: Inhibition of CCD1106 KERTr keratinocyte growth by tazarotene, as determined by staining.

[0030] FIG. 8: Inhibition of A431 cell growth by tazarotene, as determined by staining.

[0031] FIG. 9: Results of exposure of keratinocytes to a light source providing primarily visible blue light (top left) or to a light source providing primarily UVB light (top right).

[0032] FIG. 10: Improved reduction in viability of keratinocytes at increased blue light intensity.

[0033] FIG. 11: Effects of blue light exposure in combination with either KX01 or calcipotriol for from 0 hours (no blue light exposure) to 6 hours on keratinocyte proliferation.

[0034] FIG. 12: PSI values determined for combined effects of treatment with blue light, KX01, and/or calcipotriol on skin cell proliferation.

Detailed Description

[0035] The inventive subject matter provides compositions and methods in which a vitamin D analog synergistically enhances the antiproliferative effects of an antiproliferative compound for the treatment of cell hyperproliferation disorders, particularly dermatological disorders characterized by increased proliferation of skin cells (e.g. psoriasis, actinic keratosis, etc.).

[0036] Within the context of this application the term I_{max} refers to the maximal inhibitory concentration of a compound or therapy, the term IC_{50} refers to a value at which 50% of the maximum inhibitory effect realized is achieved, and the term H is related to the steepness of the slope of a calculated curve or surface derived from a mathematical model.

[0037] Various objects, features, aspects and advantages of the inventive subject matter will become more apparent from the following detailed description of preferred embodiments, along with the accompanying drawing figures.

[0038] The inventive subject matter includes pharmaceutical compositions that include a first compound that exhibits at least two of: SRC-kinase inhibition, inhibition of tubulin polymerization, cell cycle arrest at G2/M, and induction of apoptosis. The pharmaceutical composition further includes a second compound that is a vitamin D analog and/or a retinoid (i.e. a retinoid or retinoid analog). The first and second compounds are present in the pharmaceutical composition in amounts sufficient to treat a cell hyperproliferation disorder, such actinic keratosis or psoriasis. In preferred embodiments the first and second compounds are present in amounts that provide a synergistic effect on inhibition of proliferation/growth of skin cells relative to their individual contributions. In some embodiments such drug combinations can be

used in combination with controlled exposure to light, for example blue light (e.g. 280nm to 500 nm light), UV/A light, and/or UV/B light.

[0039] The recitation of ranges of values herein is merely intended to serve as a shorthand method of referring individually to each separate value falling within the range. Unless otherwise indicated herein, each individual value is incorporated into the specification as if it were individually recited herein. All methods described herein can be performed in any suitable order unless otherwise indicated herein or otherwise clearly contradicted by context. The use of any and all examples, or exemplary language (e.g. "such as") provided with respect to certain embodiments herein is intended merely to better illuminate the invention and does not pose a limitation on the scope of the invention otherwise claimed. No language in the specification should be construed as indicating any non-claimed element essential to the practice of the invention.


[0040] In some embodiments, the numbers expressing quantities of ingredients, properties such as concentration, reaction conditions, and so forth, used to describe and claim certain embodiments of the invention are to be understood as being modified in some instances by the term "about." Accordingly, in some embodiments, the numerical parameters set forth in the written description and attached claims are approximations that can vary depending upon the desired properties sought to be obtained by a particular embodiment. In some embodiments, the numerical parameters should be construed in light of the number of reported significant digits and by applying ordinary rounding techniques. Notwithstanding that the numerical ranges and parameters setting forth the broad scope of some embodiments of the invention are approximations, the numerical values set forth in the specific examples are reported as precisely as practicable. The numerical values presented in some embodiments of the invention may contain certain errors necessarily resulting from the standard deviation found in their respective testing measurements.

[0041] Unless the context dictates the contrary, all ranges set forth herein should be interpreted as being inclusive of their endpoints and open-ended ranges should be interpreted to include only commercially practical values. Similarly, all lists of values should be considered as inclusive of intermediate values unless the context indicates the contrary.

[0042] As used in the description herein and throughout the claims that follow, the meaning of "a," "an," and "the" includes plural reference unless the context clearly dictates otherwise. Also, as used in the description herein, the meaning of "in" includes "in" and "on" unless the context clearly dictates otherwise.

[0043] Groupings of alternative elements or embodiments of the invention disclosed herein are not to be construed as limitations. Each group member can be referred to and claimed individually or in any combination with other members of the group or other elements found herein. One or more members of a group can be included in, or deleted from, a group for reasons of convenience and/or patentability. When any such inclusion or deletion occurs, the specification is herein deemed to contain the group as modified thus fulfilling the written description of all Markush groups used in the appended claims.

[0044] The first compound can be an antiproliferative compound as described in U.S. 8,003,641 to David G. Hangauer, Jr., filed Apr. 26, 2007, which is incorporated herein by reference. Exemplary compounds have the structure(s) shown below. Specifically, suitable compounds include those having Formula I:



or a salt, solvate, hydrate, or prodrug thereof, wherein: T is a bond;

Xy is CZ, CY, N, or N—O;

Xz is CZ, CY, N, or N—O;

at least one of Xy and Xz is CZ;

Y is selected from hydrogen, hydroxyl, halogen, (C₁, C₂, C₃, C₄, C₅, or C₆) alkyl, C₁, C₂, C₃, C₄, C₅, or C₆ alkoxy, O- (C₁, C₂, C₃, C₄, C₅, or C₆) alkylaryl, and O-benzyl;

Xa is CRa or N, or N—O;


Xb is CRb, N, or N—O;

Xc is CRc or N, or N—O;

Xd is CRd or N, or N—O;

Xe is CRe, N, or N—O

Ra, Rb, Rc, Rd, Re, R₄, R₅, and R₆ are, independently, hydrogen, hydroxyl, halogen, C₁, C₂, C₃, C₄, C₅, or C₆ alkoxy, O-lower (C₁, C₂, C₃, C₄, C₅, or C₆) alkyl-aryl, O-benzyl, C₁, C₂, C₃, C₄, C₅, or C₆ alkyl-OH, C₁, C₂, C₃, C₄, C₅, or C₆alkyl-O- (C₁, C₂, C₃, C₄, C₅, or C₆) alkyl, COOH, COO- (C₁, C₂, C₃, C₄, C₅, or C₆) alkyl, SO₂H, SO₂ (C₁, C₂, C₃, C₄, C₅, or C₆) alkyl,



where W is H, or C₁, C₂, C₃, C₄, C₅, or C₆ alkyl, C₁, C₂, C₃, C₄, C₅, or C₆ alkyl-aryl;

V is a bond, —CH₂—, —CH₂CH₂—, —CH₂CH₂CH₂—, —O—CH₂—, —OCH₂CH₂— or —OCH₂CH₂CH₂—;

R₁₂, R₁₃, R₁₄, R₁₅, R₁₆, R₁₇, and R₁₈, are, independently, H or C₁, C₂, C₃, C₄, C₅, or C₆ alkyl;


Z is: (CHR₁)_n—C(O)—NR₂(CHR₃)_m—Ar, where Ar is a substituted or unsubstituted aryl or nitrogen-containing heteroaryl group, R₁, R₂, and R₃ are independently H or C₁, C₂, C₃, C₄, C₅, or C₆ alkyl;

n is 0, 1, or 2; and

m is 1 or 2,

wherein at least one of Xa, Xb, Xc, Xd and Xe is N.

[0045] In a preferred embodiment of the inventive subject matter, the first compound is KX01 (N-benzyl-2-(5-{4-[2-(morpholine-4-yl)ethoxy]phenyl}pyridine-2-yl)acetamide. The chemical structure of KX01 is represented by Formula 2:



Formula 2

KX01 has been shown to inhibit SRC-kinase, inhibits tubulin polymerization, and induces apoptosis by G2/M arrest.

[0046] Suitable vitamin D analogs include calcipotriol in preferred embodiments of the inventive subject matter. Calcipotriol is a vitamin D3 analog that has been observed to induce apoptosis in a variety of cell types. Without wishing to be bound by a particular theory, the inventors hypothesize that the combination of an antiproliferative compound with a vitamin D analog is beneficial, because they have different mechanisms of action that target two different phases of the cell cycle.

[0047] Surprisingly, the inventors identified synergistic (e.g., more than additive) effects in regards to inhibition of skin cell proliferation/growth when such compounds are used together. Advantageously, combining KX01 and calcipotriol expands the clinical utility of KX01 ointment to the treatment of hyperproliferative skin disorders, while synergistic effects can provide for increased effectiveness and/or reduced dosages.

[0048] Typical data from studies of the effects of compound KX01 provided in combination with calcipotriol on growth of human keratinocytes in culture (cell line CCD-1106 KERTr) are shown in FIGs. 1 to 3 and Tables 1 to 6. FIG. 1 and Tables 1 and 2 show the degree of growth inhibition found in a typical study performed with KX01 and calcipotriol at a 1:5 ratio.

nM	Log	KX01	Calcipotriol	KX01+Calcipotriol (1:5)
0.333	-0.47756	100.45		96.21622
1	0	97.0297		93.33333
3	0.477121	90.90909	94.0367	92.61261
9	0.954243	94.68947	89.90826	88.01802
27	1.431364	67.14671	70.45872	70.27027
81	1.908485	1.350135	49.3578	27.74775
243	2.385606	2.070207	47.33945	-2.79279

729	2.862728	6.660666	46.97248	-4.32432
2187	3.339849	8.910891	44.77064	-1.71171
6561	3.81697		19.17431	
19683	4.294091		-9.3578	

Table 1

Table 1 shows the results of cell viability studies comparing cell growth in response to KX01, calcipotriol, and KX01 + calcipotriol at a 1:5 ratio, as graphically depicted in FIG. 1A. Table 2 shows typical GI₅₀ and combination index (CI) values for KX01 and calcipotriol (1:5) derived from such studies.

	GI₅₀	CI (combination index)
KX01	32 nM	0.37
Calcipotriol	265 nM	
KX01+Calcipotriol (1:5)	43 nM (7.2 nM+35.8 nM)	

Table 2

[0049] FIG. 2 shows typical results of similar studies in which KX01 and calcipotriol are used in a 1:2 ratio. Typical numerical data from such a study is shown in Table 3. Table 4 shows typical GI₅₀ and combination index (CI) values for KX01 and calcipotriol (1:2) derived from such studies.

nM	Log	KX01	Calcipotriol	KX01+Calcipotriol (1:2)
0.333	-0.47756	102.3		100.7
1	0	99.9		97.2
3	0.477121	100.0	95.3	95.7
9	0.954243	100.2	88.8	93.8
27	1.431364	79.8	72.5	83.6
81	1.908485	4.1	53.2	20.8
243	2.385606	1.2	47.7	-5.4
729	2.862728	9.9	47.1	-4.8
2187	3.339849	14.9	46.7	2.5
6561	3.81697		25.2	
19683	4.294091		-13.4	

Table 3

	GI ₅₀	CI (combination index)
KX01	38 nM	0.54
Calcipotriol	310 nM	
KX01+Calcipotriol (1:2)	49 nM (16.3 nM+32.7 nM)	

Table 4

[0050] FIG. 3 shows typical results of studies in which KX01 and calcipotriol are used in a 2:1 ratio. Typical numerical data from such a study is shown in Table 5. Table 6 shows typical GI₅₀ and combination index (CI) values for KX01 and calcipotriol (1:5) derived from such studies.

nM	Log	KX01	Calcipotriol	KX01+Calcipotriol (2:1)
0.333	-0.47756	98.0		98.8
1	0	101.3		97.8
3	0.477121	96.6	96.1	96.0
9	0.954243	98.2	89.7	95.3
27	1.431364	80.1	70.1	80.2
81	1.908485	1.6	52.8	-4.6
243	2.385606	-0.1	51.0	-5.9
729	2.862728	7.3	48.7	0.8
2187	3.339849	9.5	48.3	5.8
6561	3.81697		26.5	
19683	4.294091		-15.7	

Table 5

	GI ₅₀	CI (combination index)
KX01	36 nM	0.59
Calcipotriol	339 nM	
KX01+Calcipotriol (2:1)	30 nM (20nM+10nM)	

Table 6

[0051] In order to determine the range of concentrations over which compounds such as KX01 and calcipotriol can inhibit keratinocyte growth and subsequent combinatorial effects, individual dose response curves were determined using a 72 hour exposure to either KX01 or calcipotriol in order to determine individual IC₅₀ values. Results are shown in FIG. 4 (which shows typical

results for KX01) and FIG. 5 (which shows typical results for calcipotriol). A human keratinocyte cell line was used in these studies. The IC_{50} for KX01 was determined to be 41.9 nM; the IC_{50} for calcipotriol was determined to be 95.5 nM.

[0052] Individual dose response curves for KX01 and calcipotriol as determined above can be used to generate dose response surfaces representative of the additive effects of these compounds on keratinocyte growth. Such dose response surfaces can be utilized in detailed combinatorial studies in order to determine if the effects such compounds in combination is antagonistic, merely additive, or showed positive synergistic effects. Synergistic effects can be expressed as "PSI" or Ψ , which has been used as a measure of synergy (see Ariens EJ, Van Rossum JM and Simonis AM, Affinity, intrinsic activity and drug interactions, *Pharmacol Rev.* 1957 Jun. 9(2): 218-36; Chakraborty A and Jusko WJ, Pharmacodynamic interaction of recombinant human interleukin-10 and prednisolone using in vitro whole blood lymphocyte proliferation, *J. Pharm. Sci.* 2002, 91: 1334). A PSI value of 1 corresponds to an additive effect. A PSI value of less than 1 indicates a synergistic effect, and a PSI value of greater than 1 indicates an antagonistic effect.

[0053] Cell viability was measured in a number of experiments. In Experiment 1, the concentrations of calcipotriol were: 0, 3, 9, 27, 81, and 243 nM, and the concentrations of KX01 were: 0, 5, 10, 20, 40, 60, 80, and 160 nM. In Experiment 2, the concentrations of calcipotriol were: 0, 5, 10, 20, 40, 80, 160, 320, and 1280 nM, and the concentrations of KX01 were: 1, 3, 9, 27, and 81 nM. The results are shown in FIG. 6, which shows the calculated additive response surface derived from individual dose response curves for each compound and individual data points representing the results of individual combinations. Synergistic effects are indicated when the observed response lies below the additive response surface; antagonistic effects are indicated when the observed response lies above the additive response surface. It is notable that unexpected synergy is observed across a range of concentrations and combinations.

[0054] Tables 7, 8, and 9 show the results for determination of individual IC_{50} and maximum inhibitory concentration values for KX01 and calcipotriol, as well as the PSI value for the drug combination. Table 7 shows values derived from the study depicted in FIG. 6. Tables 8 and 9 show values derived from subsequent replicate studies.

Parameters	Estimate (CV%)
PSI	0.571 (3.72)
Max Inhibition KX01	0.856 (1.31)
IC₅₀ KX01	41.3 (3.62)
Max Inhibition Calcipotriol	0.474 (2.74)
IC₅₀ Calcipotriol	112 (7.01)

Table 7

Parameters	Estimate (CV%)
PSI	0.626 (3.26)
Max Inhibition KX01	0.77 (2.11)
IC₅₀ KX01	32.8 (6.29)
Max Inhibition Calcipotriol	0.36 (3.00)
IC₅₀ Calcipotriol	59.3 (10.4)

Table 8

Parameters	Estimate (CV%)
PSI	0.787 (4.99)
Max Inhibition KX01	0.741 (5.21)
IC₅₀ KX01	47.0 (14.8)
Max Inhibition Calcipotriol	0.439 (4.99)
IC₅₀ Calcipotriol	112 (5.72)

Table 9

[0055] As shown above, a significant synergistic effect in inhibition of keratinocyte growth is consistently found when KX01 and calcipotriol are used in combination. In some embodiments such synergy is observed when the concentration of KX01 is about 1 to about 30 nM and the concentration of calcipotriol is about 3 to about 300 nM, and is present when KX01 (or a related compound) and calcipotriol (or a similar vitamin D derivative) are provided in a suitable formulation in amounts sufficient to generate such concentrations in the tissue to be treated. Such formulation can provide KX01 (or a similar compound) and calcipotriol (or a similar vitamin D derivative) in amounts that provide a ratio of these active compounds that in turn provides a synergistic effect in regards to inhibition of keratinocyte growth.

[0056] Inventors have also found that treatment with KX01 can be combined with a retinoid and/or retinoid analog to provide a synergistic (i.e. greater than additive) effect in the inhibition of cell proliferation/growth that is characteristic of hyperproliferative disorders. Examples of suitable retinoids include retinol, retinal, retinoic acid isotretinoin, alitreninoin, etretinate, acitretin, adapalene, bexarotene, and/or tazarotene. FIGs. 7 and 8 show the effects of tazarotene on growth of cells in culture, including CCD-1106 KERTr keratinocytes (FIG. 7) and A431 cells (epidermoid cells obtained from a human squamous cell carcinoma, FIG. 8). Numerical data from these studies is provided in Table 10, which shows data derived from CCD-1106 KERTr keratinocytes and from A431 cells.

Parameter	Estimate (CV%) (Keratinocytes)	Estimate (CV%) (A431)
R0	0.455 (1.07)	1.03 (1.94)
Maximum inhibition	0.903 (1.68)	1.09 (12.3)
IC₅₀	12.4 (4.12)	36.3 (19.8)
H	2.11 (7.93)	1.66 (19.9)

Table 10

[0057] Studies on the effects of combined treatment with KX01 and tazarotene were performed, and showed a clear and unexpected synergistic effect. Typical results are shown in Table 11.

	KX01 (nM)	Tazarotene (nM)
IC₅₀	45	13.3 μ M
Imax	0.92	1.1
H	3.3	1.0
	PSI value	95% Confidence Interval
	0.76 (3.3%CV)	0.7–0.8

Table 11

[0058] In addition to growth inhibition, Inventors also investigated the effect of KX01 on apoptosis in keratinocytes. CCD-1106 KERTr cells (1×10^6) were seeded in 60 mm dishes and treated with test compounds at varying concentrations (e.g. KX01: 20 to 500nM; calcipotriol 50 to 1000nM) for 24 hours. Cell apoptosis was assessed by using a FITC annexin V apoptosis detection kit (BD Biosciences™) according to manufacturer's instruction. Briefly, cells were collected in 1x binding buffer and incubated with fluorescein isothiocyanate (FITC)-conjugated annexin V and propidium iodide (PI) for 15 minutes at room temperature in the dark, followed by flow cytometry analysis using a LSRFortessa™ flow cytometer (BD Biosciences™). Results are summarized in Table 12, which shows the percentage of late apoptic cells identified for various KX01 and calcipotriol combinations.

[Calcipotriol], nM					
1,000	4.19	3.15	4.61	8.74	4.64
500	4.13	2.71	4.88	7.93	5.89
200	2.06	2.55	4.97	7.94	6.13

50	2.41	2.12	4.50	5.70	5.36	
0	2.08	2.09	3.50	6.33	4.34	
	0	20	40	100	500	[KX01], nM

Table 12

[0059] Synergy between KX01 and calcipotriol in regards to increasing the number of keratinocytes undergoing apoptosis is evident. For example the effect of 200 nM calcipotriol is not distinguishable from that of control cells receiving no drug treatment, however the same concentration of calcipotriol enhances the effect of KX01 by approximately 40%.

[0060] The inventive subject matter also includes using blue light (e.g. 280 nm to 500 nm), UV/A light, and/or UV/B light to illuminate skin during, prior to, or following treatment with KX01 (either alone or in combination with other compounds), particularly where synergistic effects permit reduction in blue light exposure and/or drug dosage relative to either light exposure or chemotherapy alone. Any suitable source of blue light can be used. Suitable sources of blue light include incandescent bulbs, fluorescent lights, LED light sources, and laser light sources. Such a light source can be a broad spectrum light sources (i.e. providing a range of wavelengths exceeding 20 nm in width) or a narrow spectrum light source (i.e. providing a range of wavelengths spanning 20 nm or less). In some embodiments the light source can produce primarily (i.e. >90%) blue light. In other embodiments the light source can produce both blue, and non-blue light, with non-blue light selectively blocked by an optical filter. In a preferred embodiment the light source for visible blue light is a panel or fluorescent light emitting a narrow bandwidth light centered at 460nm. In another preferred embodiment the light sources for UVB light is a fluorescent and/or LED/LCD panel light emitting a narrow bandwidth centered at 311nm.

[0061] It should be appreciated that exposure to blue light alone can reduce cell viability, and that different sources of blue light can provide different effects. An example of this is provided in FIG. 9, which shows the results of exposure of CCD 1106 KERTr human keratinocytes cells to different blue light sources (i.e. blue visible light or UVB) for 72 hours. It is apparent that UVB light is significantly more effective in reducing keratinocyte viability relative to visible blue light, although both are effective. As shown in FIG. 10, the overall effectiveness (in terms of absolute reduction in cell viability) of light sources providing primarily visible blue light can be improved by increasing its intensity.

[0062] The effects of combined treatment with KX01, a vitamin D derivative, and/or blue/UV light exposure were characterized using a cell proliferation assay. Cell proliferation was assessed by using an MTT assay. Cells were seeded in 96-well plate (3,000-5,000 cells per well) and treated with test compounds at varying concentrations (e.g. KX01: 5 to 80 nM; calcipotriol 5 to 1,280 nM) for 72 hours, either in the dark or exposed to blue light irradiation (for 2, 4, or 6 hours). To each well, 10 μ l of 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) (RPI Corporation™) solution (5 mg/ml in PBS) was added and incubated at 37 °C for 3 hours. Next, 100 μ l of 20% SDS was added to each well. After overnight incubation at 37 °C, the absorbance in each well was measured at 570 nm in a microplate reader (Molecular Devices™). The cell growth curve and GI₅₀s were calculate using GraphPad Prism™ 5 software.

[0063] Tables 13 and 14 show the results of cell growth inhibition studies (expressed as a percentage relative to control cell growth) for treatment with either KX01 or calcipotriol in combination with exposure to blue light.

Blue light (hours)					
6	50	56	70	88	87
4	41	47	64	87	92

2	26	27	63	92	97	
0	0	4	40	90	96	
	0	10	20	40	80	[KX01], nM

Table 13

Table 13 shows the results of a cell proliferation assay performed using KX01 and blue light and the CCD 1106 KERTr human keratinocyte cell line. The percentage of growth inhibition relative to untreated control is shown.

Blue light (hours)							
6	53	56	59	71	72	66	
4	48	53	58	70	71	69	
2	25	32	39	55	66	62	
0	0	9	24	45	48	43	
	0	5	20	80	320	1,280	[calcipotriol], nM

Table 14

Table 14 shows the results of a cell proliferation assay performed using calcipotriol and blue light and the CCD 1106 KERTr human keratinocyte cell line. The percentage of growth inhibition relative to untreated control is shown.

[0064] Tables 15 to 18 show the effects of combined treatment with KX01, calcipotriol, and exposure to blue light at 0 hours, 2 hours, 4 hours, and 6 hours of light exposure (respectively). Values are expressed as the percentage of growth inhibition relative to cells grown under control conditions (0 hours blue light, [KX01] = 0, [calcipotriol] = 0). It should be appreciated that synergy (i.e. greater than additive) effects between blue light exposure and combined KX01 and calcipotriol treatment are observed, for example at 2 hours of blue light exposure and combined treatment with 10 nM KX01 and 20 nM calcipotriol.

[Calcipotriol], nM						
1,280	43	60	86	95	98	
320	48	69	90	100	100	
80	45	67	89	100	100	
20	24	48	85	100	100	
5	9	22	80	100	100	
0	0	9	24	45	48	
	0	10	20	40	80	[KX01], nM

Table 15

[Calcipotriol], nM	
-------------------------------	--

1,280	62	79	91	95	100	
320	66	86	96	100	100	
80	55	84	96	100	100	
20	39	76	97	100	100	
5	32	47	92	100	100	
0	25	32	74	98	48	
	0	10	20	40	80	[KX01], nM

Table 16

[Calcipotriol], nM						
1,280	69	81	91	96	97	
320	71	85	93	99	100	
80	70	83	90	100	100	
20	58	69	86	98	99	
5	53	62	78	94	94	
0	48	56	69	92	93	
	0	10	20	40	80	[KX01], nM

Table 17

[Calcipotriol], nM						
1,280	66	80	91	97	99	
320	72	82	91	97	98	
80	71	82	88	95	97	
20	59	72	82	93	93	
5	56	67	75	88	87	
0	53	62	67	84	83	
	0	10	20	40	80	[KX01], nM

Table 18

[0065] Similar studies were performed using a squamous cell carcinoma cell line (SCC A431). Tables 19 and 20 show the results of cell growth inhibition studies (expressed as a percentage relative to control cell growth) for treatment with either KX01 or calcipotriol in combination with exposure to blue light.

Blue light (hours)					
6	66	79	81	81	82
4	45	66	76	73	75

2	40	40	97	97	95	
0	0	0	99	99	99	
	0	10	20	40	80	[KX01], nM

Table 19

Table 19 shows the results of a cell proliferation assay performed using KX01 and blue light and a squamous cell human carcinoma cell line (SCC A431). The percentage of growth inhibition relative to untreated control is shown.

Blue light (hours)							
6	66	67	69	67	67	68	
4	45	49	44	47	49	43	
2	18	26	36	41	35	38	
0	0	19	30	40	41	36	
	0	5	20	80	320	1,280	[calcipotriol], nM

Table 20

[0066] Tables 21 to 24 show the effects of combined treatment with KX01, calcipotriol, and exposure to blue light at 0 hours, 2 hours, 4 hours, and 6 hours of light exposure (respectively) on human squamous cell carcinoma cells. Values are expressed as the percentage of growth inhibition relative to cells grown under control conditions (0 hours blue light, [KX01] = 0,

[calcipotriol] = 0). The percentage of growth inhibition relative to an untreated control is shown. It should be appreciated that synergy (i.e. greater than additive) effects between blue light exposure and combined KX01 and calcipotriol treatment are observed, for example at 2 to 6 hours of blue light exposure and combined treatment with 10 nM KX01 and 5 nM calcipotriol, and at 2 hours of exposure to blue light and combined treatment with 10 nM KX01 and from 5 to 1,280 nM calcipotriol.

[Calcipotriol], nM						
1,280	36	31	75	96	100	
320	41	26	74	98	99	
80	40	29	75	97	100	
20	30	26	76	96	100	
5	19	1	77	96	100	
0	0	0	80	99	99	
	0	10	20	40	80	[KX01], nM

Table 21

[Calcipotriol], nM					
1,280	38	55	78	95	98

320	35	56	79	96	98	
80	41	56	80	97	98	
20	39	52	80	98	98	
5	26	46	78	96	98	
0	18	40	79	97	95	
	0	10	20	40	80	[KX01], nM

Table 22

[Calcipotriol], nM						
1,280	43	69	75	77	82	
320	49	72	77	79	83	
80	47	72	78	80	83	
20	44	68	74	78	82	
5	49	68	74	76	80	
0	45	66	69	73	75	
	0	10	20	40	80	[KX01], nM

Table 23

[Calcipotriol], nM						
1,280	68	73	88	93	94	
320	67	79	88	91	96	
80	67	81	88	89	91	
20	69	81	86	88	88	
5	67	79	79	84	86	
0	66	79	81	81	82	
	0	10	20	40	80	[KX01], nM

Table 24

[0067] As shown, KX01 and calcipotriol in combination exhibit synergistic inhibition of cell growth in keratinocytes *in vitro*, for example within the range of 1 to 30 nM KX01 and 3 to 300 nM calcipotriol. Without wishing to be bound by theory, evidence suggests that at least part of this apparent growth inhibition is due to induction of apoptosis by KX01 and calcipotriol. In addition, blue light shows additive effect on cell growth inhibition in keratinocytes *in vitro* that are exposed to KX01, calcipotriol, and/or a combination of KX01 and calcipotriol, indicating that controlled exposure to blue light can serve as a valuable adjunct to chemotherapies.

[0068] Additional studies were performed to evaluate potential synergistic effects of cell growth inhibition of human keratinocytes on exposure to blue light and to KX01 or calcipotriol. Results are shown below in Table 25 and Table 26.

Blue light

(hours)						
6	47	57	73	88	88	
4	44	50	68	86	90	
2	29	34	68	91	96	
0	0	6	49	90	95	
	0	10	20	40	80	[KX01], nM

Table 25

Blue light (hours)							
6	47	51	58	67	72	67	
4	44	47	52	64	67	63	
2	29	33	39	53	60	58	
0	0	6	17	35	48	37	
	0	5	20	80	320	1,280	[calcipotriol], nM

Table 26

Synergistic effects between blue light exposure and KX01 or calcipotriol on cell growth inhibition in human keratinocytes were observed. For example, synergy is evident at 6 hours of blue light exposure in combination with 10 nm KX01.

[0069] Studies were also performed to evaluate potential synergistic effects of cell growth inhibition of human keratinocytes on exposure to blue light and to KX01 in combination with

calcipotriol. Results are shown below in Tables 27 to Table 30, which show results from 0, 2, 4, and 6 hours of blue light exposure (respectively). Values are expressed as the percentage of growth inhibition relative to cells grown under control conditions (0 hours blue light, [KX01] = 0, [calcipotriol] = 0). It should be appreciated that synergy (i.e. greater than additive) effects between blue light exposure and combined KX01 and calcipotriol treatment are observed, and that the synergistic effect observed at 2 hours of blue light exposure and combined treatment with 10 nM KX01 and 20 nM calcipotriol is confirmed.

[Calcipotriol], nM						
1,280	37	56	85	93	95	
320	48	65	91	98	98	
80	35	57	91	98	98	
20	17	30	86	97	97	
5	6	13	72	95	97	
0	0	6	49	90	95	
	0	10	20	40	80	[KX01], nM

Table 27

[Calcipotriol], nM					
1,280	58	76	91	96	99
320	60	83	97	100	100

80	53	79	97	100	100	
20	39	63	91	99	100	
5	33	42	83	97	99	
0	29	34	68	91	96	
	0	10	20	40	80	[KX01], nM

Table 28

[Calcipotriol], nM						
1,280	63	76	90	95	96	
320	67	80	94	98	98	
80	64	76	90	98	98	
20	52	61	82	93	97	
5	47	54	74	90	94	
0	44	50	68	86	90	
	0	10	20	40	80	[KX01], nM

Table 29

[Calcipotriol], nM

1,280	67	79	90	95	96	
320	72	81	93	96	97	
80	67	78	90	95	97	
20	58	68	84	93	94	
5	51	61	80	92	91	
0	47	57	73	88	88	
	0	10	20	40	80	[KX01], nM

Table 30

[0070] Similar studies were performed to determine the synergistic effects on cell growth inhibition of squamous cell carcinoma cells (SCC A431) on exposure to blue light and KX01 or calcipotriol. Results are shown below in Tables 31 and 32, which show results for KX01 and calcipotriol (respectively). Values are expressed as the percentage of growth inhibition relative to cells grown under control conditions (0 hours blue light, [KX01] = 0, [calcipotriol] = 0).

Blue light (hours)					
6	65	77	83	86	88
4	48	66	74	80	84
2	23	44	83	99	98
0	0	11	79	100	99

	0	10	20	40	80	[KX01], nM
--	----------	-----------	-----------	-----------	-----------	-------------------

Table 31

Blue light (hours)							
6	65	62	63	61	60	58	
4	48	50	48	50	51	48	
2	23	27	36	40	37	39	
0	0	14	19	27	31	28	
	0	5	20	80	320	1,280	[calcipotriol], nM

Table 32

Synergistic effects between blue light exposure and KX01 on cell growth inhibition in squamous cell carcinoma cells are evident, for example at 10 nM KX01.

[0071] Studies were also performed to evaluate potential synergistic effects of cell growth inhibition of human squamous cell carcinoma cells on exposure to blue light and to KX01 in combination with calcipotriol. Results are shown below in Tables 33 to Table 36, which show results from 0, 2, 4, and 6 hours of blue light exposure (respectively). Values are expressed as the percentage of growth inhibition relative to cells grown under control conditions (0 hours blue light, [KX01] = 0, [calcipotriol] = 0). It should be appreciated that synergy (i.e. greater than additive) effects between blue light exposure and combined KX01 and calcipotriol treatment are observed, for example at 2 to 6 hours of blue light exposure and combined treatment with 10 nM KX01 and 5 nM calcipotriol, and at 2 hours of exposure to blue light and combined treatment with 10 nM KX01 and from 5 to 1,280 nM calcipotriol are confirmed.

[Calcipotriol], nM						
1,280	28	29	70	97	100	
320	31	29	69	97	99	
80	26	28	70	97	100	
20	19	23	72	97	100	
5	14	10	74	97	100	
0	0	11	79	100	99	
	0	10	20	40	80	[KX01], nM

Table 33

[Calcipotriol], nM						
1,280	39	57	79	96	98	
320	37	56	78	97	98	
80	40	55	79	98	99	
20	36	51	81	99	99	
5	27	45	81	98	99	
0	23	44	83	99	98	
	0	10	20	40	80	[KX01],

						nM
--	--	--	--	--	--	-----------

Table 34

[Calcipotriol], nM						
1,280	48	66	74	81	88	
320	51	67	76	81	87	
80	50	67	76	81	88	
20	48	66	74	81	87	
5	49	66	76	81	87	
0	48	66	74	80	84	
	0	10	20	40	80	[KX01], nM

Table 35

[Calcipotriol], nM					
1,280	57	70	89	96	96
320	60	74	89	94	95
80	61	75	88	92	94
20	63	76	87	91	93
5	62	75	83	89	91

0	65	77	83	86	88	
	0	10	20	40	80	[KX01], nM

Table 36

[0072] Additional studies were performed to determine PSI values for the inhibition of skin cell proliferation by blue light exposure in combination with KX01 and/or calcipotriol exposure. FIG. 11 shows results of keratinocyte cell proliferation on exposure to either KX01 or calcipotriol at various concentrations and exposure to blue light for from 0 to 6 hours. FIG. 12 shows the results of calculations of PSI values for combined treatment with KX01, calcipotriol, and blue light exposure, where a value of less than 1 indicates a synergistic effect. Synergy between KX01 and calcipotriol is evident in the absence of blue light (0 hours), but additional synergy (as indicated by a further reduced psi value) is seen on exposure to blue light with this drug combination. This is particularly evident at 2 hours of blue light exposure. Table 37 provides a summary of the numeric data presented in FIG. 12.

Time (hours)	PSI (95% confidence interval)
0	0.762 (0.696-0.828)
2	0.591 (0.514-0.668)
4	0.746 (0.595-0.897)
6	0.786 (0.660-0.913)

Table 37

[0073] Inventors believe that additional synergistic effects between exposure to blue light and KX01 and/or calcipotriol can be realized through optimization of the wavelength of blue light utilized, spectral range of the blue light utilized, and/or light intensity.


[0074] The inventive compositions and methods can treat dermatological hyperproliferative conditions, including malignant or benign hyperproliferative disorders. Malignant hyperproliferative epidermal pathologies include: squamous-cell carcinoma (SCC), basal-cell carcinoma (BCC) and other non-melanoma skin cancers (NMSCs). Benign hyperproliferative epidermal pathologies include psoriasis, other disturbances of keratinization, common warts, keratoacanthoma, seborrhoeic keratosis, seborrhea and ichthyosis. Contemplated conditions that may be treated using compositions and method according to the inventive subject matter includes: solar keratosis, ichthyosis, Grover's disease, common warts, keratoacanthoma, seborrhoeic keratosis, scleroderma,, seborrhea, HIV-associated dermatoses, hyperproliferation associated with wound healing, chronic dermatoses of autoimmune type (e.g., scleroderma and pemphigus vulgaris), alopecia, skin atrophy, photoageing, keratodermos, acne vulgaris,acne rosacea, lichen planus, cutaneous lupus erythematosus, pre-malignant conditions (e.g., melanocytic naevus, myelodysplastic syndrome, among others), Darriers disease, palmoplantar keratodermas, pityriasis rubra pilaris, epidermal naevoid syndromes, erythrokeratoderma variabilis, epidermolytic hyperkeratoses, non-bullous ichthyosiform erythroderma, cutaneous lupus erythematosus, and lichen planus.

[0075] It should be apparent to those skilled in the art that many more modifications besides those already described are possible without departing from the inventive concepts herein. The inventive subject matter, therefore, is not to be restricted except in the spirit of the appended claims. Moreover, in interpreting both the specification and the claims, all terms should be interpreted in the broadest possible manner consistent with the context. In particular, the terms “comprises” and “comprising” should be interpreted as referring to elements, components, or steps in a non-exclusive manner, indicating that the referenced elements, components, or steps may be present, or utilized, or combined with other elements, components, or steps that are not expressly referenced. Where the specification claims refers to at least one of something selected from the group consisting of A, B, C ... and N, the text should be interpreted as requiring only one element from the group, not A plus N, or B plus N, etc.

CLAIMS

What is claimed is:

1. A pharmaceutical composition comprising:
 - a first compound having an activity comprising two or more of the group consisting of SRC-kinase inhibition, inhibition of tubulin polymerization, cell cycle arrest at G2/M, and induction of apoptosis; and
 - a second compound comprising a vitamin D analog or a retinoid,wherein the first compound and the second compound are present in amounts sufficient to treat a hyperproliferative disorder in an affected area.
2. The composition of claim 1, wherein the composition is formulated for topical application.
3. The composition of claim 1 or 2, wherein the composition is formulated as one of the group consisting of a foam, a cream, a paste, an ointment, a gel, a solution, a liquid suspension, a droplet suspension, an aerosol, and a powder.
4. The composition of one of claims 1 to 3, wherein the vitamin D analog has at least one activity selected from the group consisting of G0/G1 arrest, anti-proliferation, anti-inflammation, induction of keratinocyte differentiation, decreasing the number of pro-inflammatory T-cells, and induction of apoptosis.
5. The composition of one of claims 1 to 4, wherein the first compound comprises a compound according to formula I.
6. The composition of one of claims 1 to 5, wherein the first compound is KX01 and has the formula:



7. The composition of one of claims 1 to 6, wherein the second compound is a vitamin D3 analog.
8. The composition of one of claims 1 to 7, wherein the second compound is calcipotriol (calcipotriene).
9. The composition of one of claims 1 to 8, wherein the composition is formulated to provide a molar ratio of the first compound to the second compound of from 160:3 to 5:243 at the affected area.
10. The composition of claim 9, wherein the molar ratio is selected to provide an effect at the affected area that exceeds the additive effect of the first compound and the second compound when applied individually at corresponding concentrations.
11. The composition of one of claims 1 to 10, wherein the first compound is KX01 and the second compound is calcipotriol.
12. The composition of one of claims 1 to 10, wherein the first compound is KX01 and the second compound is the retinoid.
13. The composition of one of claims 1 to 10, wherein the second compound is the vitamin D analog, wherein the composition further comprises the retinoid.
14. The composition of claim 13, wherein the first compound is KX01, the second compound is calcipotriol (calcipotriene).
15. The composition of one of claims 1 to 13, wherein the retinoid is an acetylenic retinoid.
16. The composition of one of claims 1 to 14, wherein the retinoid is tazarotene.
17. The composition of one of claims 1 to 16, wherein the first compound is present in a first amount and the second compound is present in a second amount, wherein the first amount and the second amount are selected to provide a first therapeutic effect, wherein the first therapeutic effect is greater than the sum of a second effect provided by application of the first compound at

the first amount and a third effect provided by application of the second compound at the second amount.

18. A method of treating a skin condition, comprising:

providing a topical preparation comprising a first compound and a second compound, wherein the first compound has an activity comprising two or more of the group consisting of SRC-kinase inhibition, inhibition of tubulin polymerization, cell cycle arrest at G2/M, and induction of apoptosis, and the second compound comprises a vitamin D analog or a retinoid, wherein the first compound and the second compound are present in amounts sufficient to provide a reduction or elimination of a hyperproliferative disorder in an affected area; and applying the topical preparation to an area affected by the skin condition on a schedule effective to treat the skin condition.


19. The method of claim 18, wherein the skin condition is selected from the group consisting of actinic keratosis and psoriasis.

20. The method of claim 18 or 19, wherein the schedule comprises application of the topical preparation to the area affected by the skin condition at a frequency of from once a week to four times a day.

21. The method of one of claims 18 to 20, wherein the topical composition is formulated as one of the group consisting of a foam, a cream, a paste, an ointment, a gel, a solution, a liquid suspension, a droplet suspension, an aerosol, and a powder.

22. The method of one of claims 18 to 21, wherein the vitamin D analog has at least one activity selected from the group consisting of G0/G1 arrest, anti-proliferation, anti-inflammation, induction of keratinocyte differentiation, decreasing the number of pro-inflammatory T-cells, and induction of apoptosis.

23. The method of one of claims 18 to 22, wherein the first compound is KX01 having the following structure:



24. The method of one of claims 18 to 23, wherein the second compound is a vitamin D3 analog.

25. The method of one of claims 18 to 24, wherein the second compound is calcipotriol (calcipotriene).

26. The method of one of claims 18 to 25, wherein the topical composition is formulated to provide a molar ratio of the first compound to the second compound of from 160:3 to 5:243 at the affected area.

27. The method of claim 18, wherein the topical preparation comprises the first compound in a first amount and the second compound in a second amount, wherein the first amount and the second amount are selected to provide a first therapeutic effect, wherein the first therapeutic effect is greater than the sum of a second effect provided by application of the first compound at the first amount and a third effect provided by application of the second compound at the second amount.

28. The method of one of claims 18 to 27, wherein the second compound is a vitamin D analog, and wherein the topical composition further comprises the retinoid.

29. The composition of one of claims 18 to 28, wherein the retinoid is an acetylenic retinoid.

30. The composition of one of claims 18 to 14, wherein the retinoid is tazarotene.

31. A method of treating a skin condition, comprising:

providing a topical preparation comprising a first compound and a second compound, wherein the first compound has an activity comprising two or more of the group consisting of SRC-kinase inhibition, inhibition of tubulin polymerization, cell cycle arrest at G2/M, and induction of apoptosis, and the second compound

comprises a vitamin D analog or a retinoid, wherein the first compound and the second compound are present in amounts sufficient to provide a reduction or elimination of a hyperproliferative disorder in an affected area; and applying the topical preparation to an area affected by the skin condition on a schedule effective to treat the skin condition.


32. The method of claim 31, wherein the skin condition is selected from the group consisting of actinic keratosis and psoriasis.

33. The method of claim 31 or 32, wherein the schedule comprises application of the topical preparation to the area affected by the skin condition at a frequency of once a week to four times a day.

34. The method of one of claims 31 to 33, wherein the topical composition is formulated as one of the group consisting of a foam, a cream, a paste, an ointment, a gel, a solution, a liquid suspension, a droplet suspension, an aerosol, and a powder.

35. The method of one of claims 31 to 34, wherein the retinoid has at least one activity selected from the group consisting of induction of keratinocytes proliferation, modulation of epidermal differentiation, stimulation of extracellular matrix production, and alteration of sebocyte differentiation.

36. The method of one of claims 31 to 35, wherein the first compound is KX01 having the formula:



37. The method of one of claims 31 to 35, wherein the second compound is an acetylenic retinoid.

38. The method of one of claims 31 to 37, wherein the second compound is tazarotene.

39. The method of one of claims 31 to 36, wherein the second compound is calcipotriol (calcipotriene).

40. The method of one of claims 31 to 39, wherein the topical composition is formulated to provide a molar ratio of the first compound to the second compound of from 1:1 to 1:2,500 at the affected area.

41. The method of one of claims 31 to 40, wherein the topical preparation comprises the first compound in a first amount and the second compound in a second amount, wherein the first amount and the second amount are selected to provide a first therapeutic effect, wherein the first therapeutic effect is greater than the sum of a second effect provided by application of the first compound at the first amount and a third effect provided by application of the second compound at the second amount.

42. The method of one of claims 31 to 41, wherein the second compound is the retinoid and the topical preparation further comprises the vitamin D analog.

43. The method of claim 45, wherein the vitamin D analog is calcipotriol.

44. A kit for treating a skin cell hyperproliferation disorder comprising:


a first topical composition comprising a first compound, wherein the first compound has an activity comprising two or more of the group consisting of SRC-kinase inhibition, inhibition of tubulin polymerization, cell cycle arrest at G2/M, and induction of apoptosis; and

a second topical composition comprising a second compound, wherein the second compound comprises a vitamin D analog or a retinoid,

wherein the first compound is present in the first topical composition, and the second compound is present in the second topical composition in amounts sufficient to provide a reduction or elimination of the skin cell hyperproliferation disorder in response to applying the first and second topical compositions individually according to schedule.

45. The kit of claim 44, wherein the first topical composition is formulated as one of the group consisting of a foam, a cream, a paste, an ointment, a gel, a solution, a liquid suspension, a droplet suspension, an aerosol, and a powder.

46. The kit of claim 44 or 45, wherein the first compound is KX01 having the formula:



47. The kit of one of claims 44 to 46, wherein the vitamin D analog comprises calcipotriol.

48. The kit of one of claims 44 to 47, wherein the retinoid is an acetylenic retinoid.

49. The composition of one of claims 44 to 48, wherein the retinoid is tazarotene.

50. The kit of one of claims 44 to 49, wherein the first and second topical compositions are formulated to provide a molar ratio of the first compound to the second compound of from 160:3 to 5:243 at the affected area when the first and second topical compositions are applied to the affected area according to the schedule.


51. The kit of one of claims 44 to 50, wherein the first topical composition comprises the first compound in a first amount and the second topical composition comprises the second compound in a second amount, wherein the first amount and the second amount are selected to provide a first therapeutic effect, wherein the first therapeutic effect is greater than the sum of a second effect provided by application of the first compound at the first amount and a third effect provided by application of the second compound at the second amount.

52. The kit of one of claims 44 to 51, wherein the second composition comprises the vitamin D analog, and wherein the kit further comprises the retinoid.

53. A method of increasing an activity comprising two or more of the group consisting of SRC-kinase inhibition, inhibition of tubulin polymerization, cell cycle arrest at G2/M, and induction of apoptosis of a first compound in a pharmaceutical composition, the method comprising:

including in the pharmaceutical composition second compound comprising a vitamin D analog or a retinoid, wherein the molar ratio of the first compound and the second compound is selected to enhance the activity in more than an additive manner.

54. The method of claim 53, wherein the first compound comprises KX01 having the formula:



55. The method of claim 53 or 54, wherein the vitamin D analog comprises calcipotriol.

56. The method of one of claims 53 to 55, wherein the pharmaceutical composition is formulated to provide a molar ratio of the first compound to the second compound of from 160:3 to 5:243 at the affected area.

57. The method of one of claims 53 to 56 wherein the second compound comprises the vitamin D analog, and wherein pharmaceutical composition further comprises the retinoid.

58. The method of one of claims 53 to 56 wherein the vitamin D analog is calcipotriol (calcipotriene).

59. The method of one of claims 53 to 57, wherein the retinoid is an acetylenic retinoid.

60. The method of one of claims 53 to 57, wherein the retinoid is tazarotene.

61. Use of a first compound and a second compound in preparing an medicament for treating a skin condition, wherein the first compound has an activity comprising two or more of the group consisting of SRC-kinase inhibition, inhibition of tubulin polymerization, cell cycle arrest at G2/M, and induction of apoptosis, and the second compound comprises a vitamin D analog or a

retinoid, wherein the first compound and the second compound are present a topical preparation in amounts sufficient to provide a reduction or elimination of a hyperproliferative disorder in an affected area in response to applying the topical preparation to an area affected by the skin condition on a schedule effective to treat the skin condition.


62. The use of claim 61, wherein the skin condition is selected from the group consisting of actinic keratosis and psoriasis.

63. The use of claim 61 or 62, wherein the schedule comprises application of the topical preparation to the area affected by the skin condition at a frequency of once a week to four times a day.

64. The use one of claims 61 to 63, wherein the topical composition is formulated as one of the group consisting of a foam, a cream, a paste, an ointment, a gel, a solution, a liquid suspension, a droplet suspension, an aerosol, and a powder.

65. The use of one of claims 61 to 64, wherein the vitamin D analog has at least one activity selected from the group consisting of G0/G1 arrest, anti-proliferation, anti-inflammation, induction of keratinocyte differentiation, decreasing the number of pro-inflammatory T-cells, and induction of apoptosis.

66. The use of one of claims 61 to 65, wherein the first compound is KX01 having the formula:



67. The use one of claims 61 to 66, wherein the vitamin D analog is a vitamin D3 analog.

68. The use one claims 61 to 67, wherein the vitamin D3 analog is calcipotriol.

69. The use of one of claims 61 to 65, wherein the retinoid is an acetylenic retinoid.

70. The use of one of claims 61 to 65, wherein the retinoid is tazarotene.

71. The use of one of claims 61 to 70, wherein the topical composition is formulated to provide a molar ratio of the first compound to the second compound of from 160:3 to 5:243 at the affected area.

72. The use of one of claims 61 to 71, wherein the wherein the first compound is in a first amount and the second compound in a second amount, wherein the first amount and the second amount are selected to provide a first therapeutic effect, wherein the first therapeutic effect is greater than the sum of a second effect provided by application of the first compound at the first amount and a third effect provided by application of the second compound at the second amount.

73. The use of one of claims 60 to 69, wherein the second compound is the vitamin D analog, wherein the topical composition further comprises the retinoid.

74. A method of treating a proliferative condition, comprising:

- providing a topical preparation comprising a first compound, wherein the first compound has an activity comprising two or more of the group consisting of SRC-kinase inhibition, inhibition of tubulin polymerization, cell cycle arrest at G2/M, and induction of apoptosis;
- applying the topical preparation to an area affected by the skin condition on a schedule effective to treat the skin condition;
- and directing a blue light comprising a wavelength from 280nm to 500 nm onto the area affected by the skin condition at a specified light flux and for a specified exposure time.

75. The method of claim 74, wherein the blue light is provided by a light source comprising a narrow band emission.

76. The method of claim 74, wherein the blue light is provided by a light source comprising a broad band emission.

77. The method of claim 75 or 76, wherein the light source comprises an LED/LCD or fluorescent light.

78. The method of one of claims 74 to 77, wherein the blue light comprises a wavelength selected from the group consisting of visible blue light, UVA, and UVB.

79. The method of one of claims 74 to 78, wherein the blue light has a peak wavelength of 311nm or 460 nm.

80. The method of one of claims 74 to 79, wherein the skin condition is selected from the group consisting of actinic keratosis and psoriasis.

81. The method of one of claims 74 to 80, wherein the topical composition is formulated as one of the group consisting of a foam, a cream, a paste, an ointment, a gel, a solution, a liquid suspension, a droplet suspension, an aerosol, and a powder.

82. The method of one of claims 74 to 81, wherein the topical composition further comprises a second compound selected from the group consisting of a vitamin D analog or a retinoid.

83. The method of claim 82, wherein the retinoid has at least one activity selected from the group consisting of induction of keratinocytes proliferation, modulation of epidermal differentiation, stimulation of extracellular matrix production, and alteration of sebocyte differentiation.

84. The method of claim 82 or 83, wherein the retinoid is an acetylenic retinoid.


84. The method of one of claims 82 to 84, wherein the retinoid is tazarotene.

86. The method of one of claims 74 to 82, wherein the vitamin D analog is a vitamin D3 analog.

87. The method of claim 86, wherein the vitamin D3 analog is calcipotriol.

88. The method of one of claims 82 to 87, wherein the second compound is the retinoid and the topical composition further comprises the vitamin D analog.

89. The method of one of claims 74 to 88, wherein the first compound is KX01 having the formula:



90. The method of one of claims 74 to 89, wherein the topical preparation comprises the first compound in a first amount that provides a first therapeutic effect, wherein the specified light flux and exposure time provide a second therapeutic effect, wherein the method provides a third therapeutic effect that is greater than the sum of a first therapeutic effect and the second therapeutic effect.




FIG. 1




FIG. 2




FIG. 3




FIG. 4




FIG. 5




FIG. 6




FIG. 7




FIG. 8






FIG. 9



Light intensity: 4659 $\mu\text{W}/\text{cm}^2$




FIG. 10

REPLACEMENT SHEET 7/8




FIG. 11




FIG. 12

INTERNATIONAL SEARCH REPORTInternational application No.
PCT/US2019/021037**Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)**

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. Claims Nos.: 18-43, 74-90
because they relate to subject matter not required to be searched by this Authority, namely:
Claims 18-43 and 74-90 pertain to methods for treatment of the human body by therapy and thus relate to a subject matter which this International Searching Authority is not required to search under PCT Article 17(2)(a)(i) and Rule 39.1(iv).
2. Claims Nos.: 10, 14, 27, 43, 83, 87
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
Claims 10, 14, 27, 43, 83 and 87 refer to claims which are not drafted in accordance with the sentence of Rule 6.4(a).
3. Claims Nos.: 4-9, 11-13, 15-17, 21-26, 28-30, 34-42, 47-52, 56-60, 64-73, 78-82, 84-86, 88-90
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. As all searchable claims could be searched without effort justifying an additional fees, this Authority did not invite payment of any additional fees.
3. As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:

4. No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

- Remark on Protest**
- The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.
- The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.
- No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US2019/021037**A. CLASSIFICATION OF SUBJECT MATTER****A61K 31/5377(2006.01)i, A61K 31/047(2006.01)i, A61K 31/4436(2006.01)i, A61K 9/00(2006.01)i, A61P 17/00(2006.01)i, A61P 17/06(2006.01)i, A61N 5/06(2006.01)i**

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

A61K 31/5377; A61K 31/53; A61K 31/047; A61K 31/4436; A61K 9/00; A61P 17/00; A61P 17/06; A61N 5/06

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Korean utility models and applications for utility models
Japanese utility models and applications for utility models

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

eKOMPASS(KIPO internal) & Keywords:SCR-kinase, tubulin polymerization, G2M, apoptosis, viatamin D analog, retinoid, hyperproliferative disorder

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	NEWS PROVIDED BY Kinex Pharmaceuticals, "Kinex Pharmceuticals announces allowance of the Investigational New Drug (IND) application for KX2-391 ointment by the United States Food and Drug Administration (FDA)", 24 July 2014, pp. 1-3 <URL: https://www.prnewswire.com/news-releases/kinex-pharmaceuticals-announces-allowance-of-the-investigational-new-drug-ind-application-for-kx2-391-ointment-by-the-united-states-food-and-drug-administration-fda-268466522.html > See page 1.	1-3, 44-46, 53-55 , 61-63
Y	SCOTT, L. J. et al., "Calcipotriol ointment", American Journal of Clinical Dermatology, 2001, Vol. 2, No. 2, pp. 95-120 See pages 96, 110.	1-3, 44-46, 53-55 , 61-63
Y	US 7939529 B2 (HANGAUER, JR. D. G. et al.) 10 May 2011 See columns 15, 17.	1-3, 44-46, 53-55 , 61-63
Y	KRAGBALLE, K. "Treatment of psoriasis by the topical application of the novel cholecalciferol analogue calcipotriol (MC 903)", Archives of Dermatology. 1989, Vol. 125, No. 12, pp. 1647-1652 See abstract.	1-3, 44-46, 53-55 , 61-63

 Further documents are listed in the continuation of Box C. See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

19 June 2019 (19.06.2019)

Date of mailing of the international search report

19 June 2019 (19.06.2019)

Name and mailing address of the ISA/KR

International Application Division
Korean Intellectual Property Office
189 Cheongsa-ro, Seo-gu, Daejeon, 35208, Republic of Korea

Facsimile No. +82-42-481-8578

Authorized officer

LEE, Ki Cheul

Telephone No. +82-42-481-3353



INTERNATIONAL SEARCH REPORTInternational application No.
PCT/US2019/021037

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	DUBERTRET, L. et al., "Efficacy and safety of calcipotriol (MC 903) ointment in psoriasis vulgaris", Journal of the American Academy of Dermatology, 1992, Vol. 27, No. 6, pp. 983-988 See abstract.	1-3,44-46,53-55 ,61-63

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No.

PCT/US2019/021037

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 7939529 B2	10/05/2011	CN 101616915 A	30/12/2009
		CN 101616915 B	17/12/2014
		CN 101687798 A	31/03/2010
		CN 101687798 B	15/08/2012
		CN 102657656 A	12/09/2012
		CN 104230793 A	24/12/2014
		EP 2114934 A1	11/11/2009
		EP 2114934 B1	21/10/2015
		EP 2155681 A1	24/02/2010
		EP 2155681 B1	02/05/2018
		JP 2010-514771 A	06/05/2010
		JP 2010-527361 A	12/08/2010
		JP 2014-037439 A	27/02/2014
		JP 2015-028068 A	12/02/2015
		JP 2015-232050 A	24/12/2015
		JP 2016-222733 A	28/12/2016
		JP 2017-193587 A	26/10/2017
		JP 5679811 B2	04/03/2015
		JP 6022657 B2	09/11/2016
		JP 6329871 B2	23/05/2018
		US 2008-0221102 A1	11/09/2008
		US 2008-0287436 A1	20/11/2008
		US 2009-0318450 A1	24/12/2009
		US 2011-0201611 A1	18/08/2011
		US 2011-0201612 A1	18/08/2011
		US 2013-0210822 A1	15/08/2013
		US 2015-0315147 A1	05/11/2015
		US 2017-0101378 A1	13/04/2017
		US 7851470 B2	14/12/2010
		US 7935697 B2	03/05/2011
		US 8293739 B2	23/10/2012
		US 8309549 B2	13/11/2012
		US 8901297 B2	02/12/2014
		US 9556120 B2	31/01/2017
		WO 2008-082637 A1	10/07/2008
		WO 2008-144045 A1	27/11/2008
		WO 2010-135429 A2	25/11/2010
		WO 2010-135429 A3	31/03/2011