

US00PP30753P3

(12) **United States Plant Patent**
Sakurai

(10) **Patent No.:** **US PP30,753 P3**

(45) **Date of Patent:** **Jul. 30, 2019**

(54) *VACCINIUM CORYMBOSUM* L. PLANT
NAMED 'RYOKU NH-12'

(50) Latin Name: *Vaccinium corymbosum* L.
Varietal Denomination: **RYOKU NH-12**

(71) Applicant: **NIPPON RYOKUSAN CO., LTD.**,
Matsumoto-shi (JP)

(72) Inventor: **Shigetaka Sakurai**, Matsumoto (JP)

(73) Assignee: **NIPPON RYOKUSAN CO., LTD.**,
Matsumoto-shi (JP)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.

(21) Appl. No.: **15/732,725**

(22) Filed: **Dec. 21, 2017**

(65) **Prior Publication Data**
US 2018/0199482 P1 Jul. 12, 2018

(30) **Foreign Application Priority Data**
Jan. 4, 2017 (JP) 31724

(51) **Int. Cl.**
A01H 5/08 (2018.01)

(52) **U.S. Cl.**
USPC **Plt./157**
CPC *A01H 5/08* (2013.01)

(58) **Field of Classification Search**
USPC **Plt./157**
See application file for complete search history.

Primary Examiner — Annette H Para
(74) *Attorney, Agent, or Firm* — Oblon, McClelland,
Maier & Neustadt, L.L.P.

(57) **ABSTRACT**

A new and distinct variety of *Vaccinium corymbosum* L. plant named 'RYOKU NH-12', characterized by having more compact plant size, comparatively early fruit ripening time, comparatively large and uniform fruit size, firmer and lower dehiscent fruits, sweeter fruits, better taste balance of sweetness and acidity, and better taste of fruits, as compared to other *Vaccinium corymbosum* L. varieties.

8 Drawing Sheets

1

The Latin name of the genus and species of the novel variety disclosed herein is: *Vaccinium corymbosum* L.

The novel variety of the *Vaccinium corymbosum* L. disclosed herein has been given the variety denomination: 'RYOKU NH-12'.

CROSS-REFERENCE TO RELATED APPLICATIONS

This application claims priority to Japanese Plant Breeders' Rights Application No. 31724, filed Jan. 4, 2017, the contents of which are incorporated herein by reference in their entirety.

BACKGROUND OF THE INVENTION

The present invention relates to a new and distinct hybrid variety of northern highbush blueberry (*Vaccinium corymbosum* L.) named 'RYOKU NH-12'. This novel variety was found by open pollination of 'Spartan', 'Duke' and 'Denies Blue' in the tests conducted for the period from 2004 to 2008 in Matsumoto-City, Nagano-prefecture, Japan. As stated below, 'RYOKU NH-12' has apparently different characteristics from those of the 'Spartan' and 'Duke', both being widely planted and important varieties in the Chubu district of Japan.

SUMMARY OF THE INVENTION

Blueberry variety 'RYOKU NH-12' exhibits outstanding and distinguishing characteristics when grown under normal

2

horticultural conditions in the area from Nagano-prefecture to the north of the Kanto in Japan, including:

- (1) more compact plant size;
- (2) comparatively early fruit ripening time (50% fruit ripening around late June of each year in Matsumoto, Nagano, Japan);
- (3) comparatively large and uniform fruit size;
- (4) firmer and lower dehiscent fruits;
- (5) sweeter fruits with better taste balance of sweetness and acidity; and
- (6) better taste of fruits.

BRIEF DESCRIPTION OF THE DRAWINGS

The accompanying colored photographs (FIGS. 1 to 8) show typical bush, flower, fruit and leaf characteristics of the new *Vaccinium corymbosum* L. plant, 'RYOKU NH-12'. Colors shown are as accurate as can be reasonably reproduced by photographic means. In some cases, the color might differ slightly from the colors of 'RYOKU NH-12' recited in the description.

FIG. 1 shows a tree body of 'RYOKU NH-12' (photographed date: Jul. 9, 2014; photographed location: Matsumoto-City, Nagano-prefecture, Japan).

FIG. 2 shows a panoramic view of the cultivation area of 'RYOKU NH-12' (photographed date: Sep. 7, 2012; photographed location: same as FIG. 1).

FIG. 3 shows whole flowers of 'RYOKU NH-12' (photographed date: May 6, 2014; photographed location: same as FIG. 1).

FIG. 4 shows broken-down parts of a flower of 'RYOKU NH-12' (photographed date: May 6, 2014; photographed location: same as FIG. 1).

FIG. 5 shows fruits of 'RYOKU NH-12' (photographed date: Jul. 15, 2014; photographed location: same as FIG. 1).

FIG. 6 shows a cross-section of the fruits of 'RYOKU NH-12' (photographed date: Jul. 15, 2014; photographed location: same as FIG. 1).

FIG. 7 shows an upper side of the leaves (10 sheets) of 'RYOKU NH-12' (photographed date: Aug. 3, 2012; photographed location: same as FIG. 1).

FIG. 8 shows a lower side of the leaves (10 sheets) of 'RYOKU NH-12' (photographed date: Aug. 3, 2012; photographed location: same as FIG. 1).

DETAILED BOTANICAL DESCRIPTION

A. Distinctive Characteristics of 'RYOKU NY-12'

As described above, 'RYOKU NH-12' was obtained by the open pollination of 'Spartan', 'Duke' and 'Denies Blue'. On Apr. 15, 2004, about 1,000 seeds obtained from the mixed fruits of 'Spartan', 'Duke' and 'Denies Blue', all of which were harvested in a field in Matsumoto-City, Nagano-prefecture, Japan, seeded in plant seeding trays, and then transplanted to pots. The resulting seedlings (about 300) were planted in cultivation pots on May 1, 2005. Fructification of the planted seedlings was started from Jun. 20, 2008 (on Year 3), and about 20 plant individuals were selected based on the characteristics, including larger fruit size, better taste balance of sweetness and acidity, higher yield per plant, earlier ripening time, large and uniform fruits, etc. For the about 20 plant individuals selected, test plots (each including about 20 cuttings per plant individual) were formed, where these plants were asexually propagated by cutting means. During the period from Jul. 20, 2010 to Dec. 25, 2016 and for 3 generations, the plants were propagated and examined for their characteristics based on the growth, and yield and fruits quality in each test plot. For 10 test plots, the characteristics of the plants were observed for the period between the flowering time and the ripening time yearly for 4 years, and since neither variant nor off-type plant was observed for the period, the characterization of 'RYOKU NH-12' was finished on Dec. 25, 2016 and the breeding was completed.

'RYOKU NH-12' is a blueberry clone distinguishable from the important blueberry varieties 'Spartan' and 'Duke', both of which are widely planted in the Chubu district of Japan, due to its characteristics including more compact plant size, comparatively early fruit ripening time, comparatively large and uniform fruit size, firmer and lower dehiscent fruits, sweeter fruits with better taste balance of sweetness and acidity, and better taste fruits. 20 plants of 'RYOKU NH-12' had been propagated by cutting means in Matsumoto, Nagano, Japan, and all the resulting plants were phenotypically indistinguishable from the original plant variety 'RYOKU NH-12'. In addition, comparing to 'Spartan' and 'Duke', the claimed plant 'RYOKU NH-12' has more compact plant size, about 1 week earlier fruit ripening time (on average, around June 25 to July 1 of each year in Matsumoto, Nagano, Japan), larger and more uniform fruit size, and firmer and sweeter fruit when compared to its related variety 'Spartan', and has more compact plant size, lower dehiscent fruits, sweeter fruits, a better balance of sweetness and acidity, and better tasting of fruits when compared to the 'Duke' variety (see Table 1 below).

The following data defining characteristics of 'RYOKU NH-12' were collected from the asexual propagation carried out in Matsumoto, Nagano, Japan. The plant history was

taken on a plot of 10 four-year-old plants growing in Matsumoto, Nagano, Japan. 'RYOKU NH-12' has not been observed under all possible environmental conditions, and the measurements provided might therefore vary if grown in different environments. Where averages are given, the sample size was 10.

B. Phenotypic Description of *Vaccinium corymbosum* L. ('RYOKU NH-12')

Characteristics of 'RYOKU NH-12' are further specifically described as follows:

1. Plant:

Plant vigor.—Medium.

Growth habit.—Semi-upright.

Plant size.—Small.

Plant height.—1.3 m on average for 4-year old plant.

Plant spread.—0.8 m on average for 4-year old plant.

Color of bark of plant.—Dark Red, 187-B (The R.H.S. Colour Chart).

Cold hardiness.—Survived in winter frost (below -10° C.) with minimum damage.

Ease of propagation.—Propagated from each of the dormant wood cutting and the softwood stem cutting, where the rooting percentage was greater than 70% and comparable to the other varieties.

2. Trunk and branches:

Suckering tendency.—Medium suckering tendency.

Surface texture (of 6-month-old shoots).—Medium smoothness.

Surface texture (of 3-year-old and older wood).—Medium smoothness.

Color of new twigs observed in the field.—Orangish yellow green.

Internode length.—16.2 mm on average.

3. Leaves:

Length including petiole.—54.0 mm on average.

Width of leaf at widest point.—33.0 mm on average.

Shape.—Ovate.

Leaf margin.—Entire.

Color.—Upper surface of leaves: Dark Yellowish Green, 139-A. Lower surface of leaves: Strong Yellow Green, 143-A (The R.H.S. Colour Chart).

Pubesence.—Upper Surface of leaves: Absent. Lower Surface of leaves: Absent. Margin: Absent.

Timing of vegetative bud burst.—Medium.

4. Flowers:

Shape.—Campanulate.

Color of opened flower.—Pale Yellow Green, 157-C (The R.H.S. Colour Chart).

Flowering period.—Mean date of 50% opening of flowers in Matsumoto-City, Nagano-prefecture, Japan is May 1 (2 days earlier than 'Spartan' variety).

Corolla.—Diameter: 7.4 mm on average. Length (from pedicel attachment point to corolla tip excluding the pedicel): 10.8 mm on average. Color: light green white. Anthocyanin coloration in corolla tube — Absent or very weak.

5. Reproductive organs:

Pollen.—Color: Yellow.

6. Fruit:

Mean date of 50% harvest in Matsumoto-City, Nagano-prefecture.—July 1 on average.

Diameter of calyx aperture on mature berry.—5.9 mm on average.

Size and shape of calyx lobe on mature berry.—Medium in size, outcurving, and having deep calyx basin.

Detachment force for ripe berries (easy, medium, hard).—Easy.

Fruit cluster density (sparse, medium, dense).—Medium.

Fruiting type.—On one-year old shoots only.

7. Berry:

Cluster (tight, medium, loose).—Medium.

Weight (on well-pruned plants).—3.79 g on average.

Height.—14.9 mm on average.

Width.—20.4 mm on average.

Shape.—Oblate.

Skin of fruit, with bloom.—Light Purplish Blue, 98-D (The R.H.S. Colour Chart).

Intensity of fruit bloom.—Medium.

Skin of fruit, without bloom.—Greyish Purplish Blue, 103-A (The R.H.S. Colour Chart).

Immature berry color, with bloom.—Medium green.

Immature berry color, without bloom.—Yellow green.

Flesh color.—Light Yellow Green, 150-D (The R.H.S. Colour Chart).

Peel color.—Medium blue.

Color of seeds.—Moderate Orange, N167-C (The R.H.S. Colour Chart).

Pedical scar.—Medium. 2.92 mm on average.

Firmness.—Medium.

Intensity of fruit sweetness.—High, Bx 14.5.

Intensity of fruit acidity.—High, pH 2.59.

Texture.—Somewhat soft pulp, very juicy, medium seeds.

8. Use: ‘RYOKU NH-12’ produce northern highbush blueberries suitable for fruit-picking farms, fresh fruit markets and processed fruit market, etc.

9. Resistance to disease, insects, and mites: ‘RYOKU NH-12’ grew vigorously and showed excellent bush survival in the field. It appears to be tolerant to stem blight (*Botryosphaeria* spp.) and root rot (*Phytophthora cinnamoni*), with very few young plants dying soon after planting. The response of ‘RYOKU NH-12’ to the various fungal species that cause summer leaf spots is typical of other northern highbush varieties, and fungicide applications may be needed after harvest in order to reduce foliar diseases and to retain leaves until autumn and make maximum flower bud set. Similarly, susceptibility to typical blueberry insect and mite pathogens, such as spotted wing drosophila (*Drosophila suzukii*), blueberry gall midge (*Dasineura oxycoccana*) and blueberry bud mite (*Acalitus vaccini*), is similar to other northern highbush cultivars.

TABLE 1-continued

(Comparison of characteristics among varieties)					
	4	3	PQ	One-year-old shoot: color	Color of middle part of shoot extended before dormant period
	5		QN	One-year-old shoot: length	Length of middle part of shoot extended before dormant period
	10	6	4	QN	One-year-old shoot: length of internode (upper half)
	15	7	5	QN	Leaf: length
			(*)		Length of internode of shoot extended before dormant period (upper half)
		8	6	QN	Leaf: width
		9	7	QN	Leaf: ratio length/width
	20				Ratio of leaf length to maximum width (leaf length/leaf width)
	10	8	PQ	Leaf: shape	Shape of mature leaf
	11		QN	Leaf: shape of tip	Shape of tip of mature leaf
	25	12	9	QL	Leaf: color of upper side
		13	10	QN	Only varieties with green leaf color
			(*)		Intensity of green color on surface of mature leaf
	30				Leaf: intensity of green color on upper side
	14	11	QL	Leaf: margin	Type of margin of mature leaf
	15	12	QN	Flower bud: anthocyanin coloration	Intensity of anthocyanin coloration of flower bud occurring to one year old shoot
	16	13	QN	Inflorescence: length (excluding peduncle)	Length of inflorescence at flowering time (excluding peduncle)
	17	14	PQ	Flower: shape of corolla	Shape of corolla at full bloom
	18		PQ	Flower: color of corolla	Color of corolla at full bloom
	19	15	QN	Flower: size of corolla tube	Size of corolla tube at full bloom
	20	16	QN	Flower: anthocyanin coloration of corolla tube	Intensity of anthocyanin coloration on surface of corolla tube
	21	17	QL	Flower: ridges on corolla tube	Presence or absence of ridges on corolla tube
	22	18	QN	Fruit cluster: density	Density of fruit per fruit cluster
	23	19	QN	Unripe fruit: intensity of green color	Intensity of green color of fruit before ripening
	24	20	QN	Fruit: size	Size of fruit at ripening
	60	21	PQ	Fruit: shape in longitudinal section	Shape in longitudinal section of fruit at ripening
	25		(*)		Size of stem scar of mature fruit
	26		QN	Fruit: size of scar	
	65		(+)		

TABLE 1

(Comparison of characteristics among varieties)				
Charact. No.	UPOV No.	Code	Characteristics	Definition
1	1	QN	Plant: vigor	Strength of growth level of plant
	(*)	(+)		
2		QN	Plant: size	Size of plant crown
3	2	QN	Plant: growth habit	Whole shape of plant without pruning during dormant period
	(*)	G		

TABLE 1-continued

TABLE 1-continued

(Comparison of characteristics among varieties)					(Comparison of characteristics among varieties)						
Charact. No.	Method	Class	State	Standard Variety (Ex. Var.)							
27	PQ (+)		Fruit: shape of calyx cavity	Shape of calyx cavity of mature fruit	5	5	Measurement mm (a)	3	5	short	5
28	22	QN	Fruit: attitude of sepals	Attitude of sepals relative to mature fruit	6	6	Observation (a) VG	3	3	short	Avonblue, Weymouth
29	23	QN	Fruit: type of sepals	Direction of curving of sepals	10	7	Measurement mm (b)	5	7	medium long	Jersey
30	24	QN	Fruit: diameter of calyx basin	Diameter of calyx basin of mature fruit	7	7	Measurement mm (b)	3	5	short medium	
31	25	QN	Fruit: depth of calyx basin	Depth of calyx basin of mature fruit	8	8	MS/VG	3	7	long	
32	26 (*)	QN	Fruit: intensity of bloom	Intensity of bloom on surface of mature fruit	15	15	Measurement mm (b)	3	5	narrow medium	
33	27 (*)	PQ G	Fruit: color of skin (after removal of bloom)	Color of skin of mature fruit after removal of bloom	9	9	MS/VG	3	7	small medium large	
34	28	QN (*)	Fruit: firmness	Firmness of mature fruit	20	10	Observation (b) VG	1	2	lanceolate ovate	Northland Berkeley, Collins, Coville
35		PQ	Fruit: color of flesh	Color of flesh of mature fruit				3	3	elliptic	
36	29 (*)	QN (+)	Fruit: sweetness	Sweetness of mature fruit		11	Observation (b) VG	4	3	oblong acute	Weymouth, Woodard
37	30 (*)	QN (+)	Fruit: acidity	Acidity of mature fruit	25	25	VG	5	5	medium	Earliblue, Tifblue
38	31 (*)	QL G	Plant: fruiting type	Shoots to which fruit adnate				7	7	obtuse	Berkeley, Climax, Southland
39		QN	Fruit: tendency of cracking	Tendency of cracking during harvest season	12	30	Observation (b) VG	1	2	yellow green	Bluechip, Bluecrop, Blueray
40	32 (*)	QN (+)	Time of vegetative bud burst	Time of beginning to burst first vegetative bud of each individual plant	13	13	Observation (b) VG	3	5	light medium dark	
41	33 (*)	QN (+) G	Time of beginning of flowering on one-year-old shoot	Time of 10% flowering occurring to one year old shoot	14	35	Observation (b) VG	1	2	entire serrate	
					15	15	Observation (a) VG	3	5	weak medium strong	
					16	40	Measurement mm (c)	3	5	short medium long	
1	Observation (a) VG	3	weak	Bluetta, Meader	17	17	Observation (c) VG	1	2	urceolate	Bluecrop, Jersey
		5	medium	Collins, Weymouth	45	45		2	2	campanulate	Northblue, Northsky
		7	strong	Berkeley, Homebell, Woodard				3	3	cylindrical white	Aliceblue, Bluetta, Briteblue
2	Observation (a) VG	3	small	Avonblue, Bluetta, Flordablue	18	18	Observation (c) VG	1	2	creamy white	Avonblue, Berkeley, Bluecrop
		5	medium	Bluecrop, Earliblue	50	50		2	2		Blueray, Berkeley, Bluecrop
		7	large	Dixi, Homebell, Tifblue				3	3	greenish white	Collins, Coville
3	Observation (a) VG	1	upright	Becyblue, Bluechip, June, Spartan	55	19	Observation (c) VG	3	5	light pink	Bluebell, Delite, Dixi, Tifblue
		2	semi-upright	Bluecrop, Lateblue				7	7	small medium	
		3	spreading	Northland, Weymouth	20	20	Observation (c) VG	1	7	large absent or very weak	
4	Observation (a) VG	1	green		60	60		3	5	weak medium	
		2	greenish red	Briteblue, Homebell				7	7	strong	
		3	greyish red	Berkeley, Dixi	21	21	Observation (c) VG	1	9	absent present	Herbert Aliceblue Berkeley, Dixi
		4	reddish yellow	Blueray, Darrow, Weymouth							
		5	reddish brown								
		6	dark red		65	65					

TABLE 1-continued

(Comparison of characteristics among varieties)				
22	Observation (d) VG	3	sparse	Homebell, Jersey, Woodard Bluechip, Bluecrop, Bluetta Darrow, Herbert, Patriot
		5	medium	
		7	dense	
23	Observation VG	3	light	
		5	medium	
24	Observation (d) VG	3	small	Homebell, June, Northblue Collins, Earliblue Berkeley, Bluecrop, Spartan
		5	medium	
25	Observation (d) VG	1	elliptic	Berkeley, Jersey, Sharpblue Earliblue, Harrison, Woodard
		2	round	
		3	oblate	
26	Observation (d) VG	3	small	
		5	medium	
		7	large	
27	Observation (d) VG	1	star	
		2	circular	
28	Observation (d) VG	1	erect	
		2	erect to	
		3	semi-erect	
		4	level	
29	Observation (d) VG	1	incurving	
		2	straight	
30	Observation (d) VG	3	reflexed	Avonblue, Bluechip, Sharpblue Blueray, Woodard Coville, Darrow, Homebell Bluecrop, Rancocas, Tifblue Earliblue, Jersey Blueray, Collins, Dixi
		5	small	
		7	medium	
31	Observation (d) VG	3	shallow	
		5	medium	
		7	deep	
32	Observation (d) VG	1	very weak	Dixi, Herbert, Sharpblue Collins, Coville Avonblue, Bluecrop, Tifblue
		3	weak	
		5	medium	
33	Observation (d) VG	1	strong	Berkeley, Bluechip, Tifblue Blueray, Jersey, June Dixi, Homebell
		2	light blue	
		3	medium blue	
34	Observation (d) VG/VS	3	dark blue	Herbert, Homebell, Spartan Collins, Dixi Coville, Southland
		4	blue red	
		3	soft	
		5	medium	
		7	firm	
		9	very firm	

TABLE 1-continued

(Comparison of characteristics among varieties)					
35	Observation (d) VG	1	white	Berkeley, Bluecrop, Blueray Earliblue Bluechip, Lateblue, Sharpblue Aliceblue, Delite, Homebell Avonblue, Bluechip	
		2	cream		
		3	light green		
36	Observation (d) VG	4	light purple		
		3	low		
37	Observation (d) VG	5	medium	Berkeley, Bluetta, Spartan Aliceblue, Bluecrop, Blueray Earliblue, Homebell Blueray, Herbert Collins, Elliott, Lateblue	
		7	high		
		3	low		
38	Observation (c) VG	5	medium		
		7	high		
39	Observation (d) VG	1	on one-year-old shoots only	Earliblue, Herbert, Spartan Avonblue, Berkeley, Bluechip Briteblue, Climax, Darrow Avonblue, Beckyblue Sharpblue Darrow, Weymouth Elliot, Lateblue	
		2	on one-year-old and current season's shoots less		
		3	less		
40	Measurement MG	3	early		
		5	medium		
41	Measurement MG	7	late		
		1	very early		
		3	early		
45		5	medium	Bluecrop, Collins, Woodard Dixi, Herbert, Lateblue	
		7	late		
		9	very late		
40	Charact.	The present variety		Control Varieties	
		RYOKU			
45	No.	NH-12	Spartan	Duke	
		1	5	6	7
50		2	3	5	
		3	2	1	2
55		4	5	4	
		5	7	7	4
60		6	(164 mm)	(222 mm)	(245 mm)
		7	4	5	5
65		8	(16.2 mm)	(17.0 mm)	(18.0 mm)
		9	3	5	5
		10	(54.0 mm)	(76.0 mm)	(71.0 mm)
		11	5	5	5
		12	(33.0 mm)	(37.0 mm)	(36.0 mm)
		13	4	5	5
		14	(1.64)	(2.05)	(1.97)
		15	2	3	3
		16	7	5	5
		17	2	2	2
		18	5	7	7
		19	1	1	1
		20	5	5	5
		21	7	5	5
		22	(35.7 mm)	(25.2 mm)	(26.4 mm)
		23	2	1-2	2
		24	2	2-3	2-3
		25	2	2-3	2-3

TABLE 1-continued

(Comparison of characteristics among varieties)			
19	5	5	6
20	1	1	3
21	9	9	9
22	5	5	5
23	5	3	5
24	7	7	7
	(3.79 g)	(4.56 g)	(3.96 g)
25	3	3	3
26	5	5	5
	(2.92 mm)	(2.30 mm)	(1.90 mm)
27	2	2	2
28	2	1	1
29	3	1	3
30	5	5	5
	(5.86 mm)	(5.80 mm)	(5.90 mm)
31	7	7	5
	(2.72 mm)	(2.20 mm)	(1.50 mm)
32	5	5	5
33	2	2	2
34	5	3	5

TABLE 1-continued

(Comparison of characteristics among varieties)				
	35	1	2-3	1-2
5	36	7	5	5
		(Bx 14.5)	(Bx 11.9)	(Bx11.1)
	37	7	5	3
		(pH 2.59)	(pH 3.12)	(pH 3.87)
	38	1	1	1
	39	3	3	5
10	40	5	7	5
		Apr. 6,	Apr. 8, 2016	Apr. 5, 2016
		2016		
	41	3	3	3
		Apr. 25,	Apr. 24, 2016	Apr. 24, 2016
		2016		

15
20

What is claimed is:

1. A new and distinct variety of *Vaccinium corymbosum* L. plant named 'RYOKU NH-12', as described and illustrated herein.

* * * * *

Fig. 1



Fig. 2



Fig. 3




Fig. 4




Fig. 5




Fig. 6






Fig. 7



RYOKU NH-12

Fig. 8



RYOKU NH-12