

(19) **DANMARK**

(10) **DK/EP 3091073 T4**



Patent- og
Varemærkestyrelsen

(12) **Oversættelse af ændret
europæisk patentskrift**

-
- (51) Int.Cl.: **C 12 N 9/24 (2006.01)** **C 12 P 19/02 (2006.01)** **C 12 P 19/14 (2006.01)**
- (45) Oversættelsen bekendtgjort den: **2023-06-06**
- (80) Dato for Den Europæiske Patentmyndigheds bekendtgørelse om opretholdelse af patentet i ændret form: **2023-03-08**
- (86) Europæisk ansøgning nr.: **16170376.4**
- (86) Europæisk indleveringsdag: **2012-08-01**
- (87) Den europæiske ansøgnings publiceringsdag: **2016-11-09**
- (30) Prioritet: **2011-08-04 EP 11250700**
- (62) Stamansøgningsnr: **12748613.2**
- (84) Designerede stater: **AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO RS SE SI SK SM TR**
- (73) Patenthaver: **Novozymes Inc., 1445 Drew Avenue, Davis, CA 95616, USA**
- (72) Opfinder: **SPODSBERG, Nikolaj, Vandkarsevej 62, 2880 Bagsværd, Danmark**
- (74) Fuldmægtig i Danmark: **Novozymes A/S, Krogshøjvej 36, 2880 Bagsværd, Danmark**
- (54) Benævnelse: **POLYPEPTIDER MED XYLANASEAKTIVITET OG POLYNUKLEOTIDER, DER KODER FOR DEM**
- (56) Fremdragne publikationer:
WO-A1-02/24926
WO-A1-2011/057140
WO-A2-2010/059424

DESCRIPTION

Reference to a Sequence Listing

[0001] This application contains a Sequence Listing in computer readable form, which is incorporated herein by reference.

Background of the Invention

Field of the Invention

[0002] The present invention relates to polypeptides having xylanase activity and polynucleotides encoding the polypeptides. The invention also relates to nucleic acid constructs, vectors, and host cells comprising the polynucleotides as well as methods of producing and using the polypeptides.

Description of the Related Art

[0003] Cellulose is a polymer of glucose linked by beta-1,4-bonds. Many microorganisms produce enzymes that hydrolyze beta-linked glucans. These enzymes include endoglucanases, cellobiohydrolases, and beta-glucosidases. Endoglucanases digest the cellulose polymer at random locations, opening it to attack by cellobiohydrolases. Cellobiohydrolases sequentially release molecules of cellobiose from the ends of the cellulose polymer. Cellobiose is a watersoluble beta-1,4-linked dimer of glucose. Beta-glucosidases hydrolyze cellobiose to glucose.

[0004] The conversion of lignocellulosic feedstocks into ethanol has the advantages of the ready availability of large amounts of feedstock, the desirability of avoiding burning or land filling the materials, and the cleanliness of the ethanol fuel. Wood, agricultural residues, herbaceous crops, and municipal solid wastes have been considered as feedstocks for ethanol production. These materials primarily consist of cellulose, hemicellulose, and lignin. Once the cellulose is converted to glucose, the glucose is easily fermented by yeast into ethanol. Xylanases degrade beta-1,4-xylan into xylose, thus breaking down hemicellulose, one of the major components of plant cell walls.

[0005] There is a need in the art to improve cellulolytic and hemicellulolytic enzyme compositions through supplementation with additional enzymes to increase efficiency and to provide cost-effective enzyme solutions for degradation of lignocellulose. Xylanases are known

from the prior art. A xylanase from *Penicillium canescens* (Uniprot. C3VEV9) has 76.1% identity to the xylanase disclosed as SEQ ID NO: 2. Another xylanase from *Talaromyces stipitatus* (Uniprot. B8M9H8) has 76.5% identity to the xylanase disclosed as SEQ ID NO: 4. Another xylanase from *Penicillium sp* (Uniprot. AYB51189) disclosed in US2010124769-A1 has 84.0% identity to the xylanase disclosed as SEQ ID NO: 6.

[0006] The present invention provides polypeptides having xylanase activity and polynucleotides encoding the polypeptides.

Summary of the Invention

[0007] The present invention relates to isolated polypeptides having xylanase activity selected from the group consisting of:

1. (a) a polypeptide having at least 85% sequence identity to the mature polypeptide of SEQ ID NO: 6;
2. (b) a polypeptide encoded by a polynucleotide having at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or at least 100% sequence identity to the mature polypeptide coding sequence of SEQ ID NO: 5; or the cDNA sequence thereof;
3. (c) a fragment of the polypeptide of (a), or (b) that has xylanase activity. Also described are isolated polypeptides comprising a catalytic domain selected from the group consisting of:
 1. (a) a catalytic domain having at least 77% sequence identity to the catalytic domain of SEQ ID NO: 2 (for example, amino acids 18 to 364 of SEQ ID NO: 2), a catalytic domain having at least 77% sequence identity to the catalytic domain of SEQ ID NO: 4 (for example, amino acids 17 to 326 of SEQ ID NO: 4), or a catalytic domain having at least 85% sequence identity to the catalytic domain of SEQ ID NO: 6 (for example, amino acids 21 to 337 of SEQ ID NO: 6);
 2. (b) a catalytic domain encoded by a polynucleotide having at least 60% sequence identity to the catalytic domain coding sequence of SEQ ID NO: 1 (for example, nucleotides 52-240, and 314-1165 of SEQ ID NO: 1), a catalytic domain encoded by a polynucleotide having at least 60% sequence identity to the catalytic domain coding sequence of SEQ ID NO: 3 (for example, nucleotides 49-241, 302-342, 404-452, 518-639, 707-852, 912-1019, 1088-1205, 1282-1347, and 1430-1516 of SEQ ID NO: 3), or a catalytic domain encoded by a polynucleotide having at least 60% sequence identity to the catalytic domain coding sequence of SEQ ID NO: 5 (for example, nucleotides 124-270, 342-474, 567-680, and 757-1313 of SEQ ID NO: 5);
 3. (c) a variant of a catalytic domain comprising a substitution, deletion, and/or insertion of one or more (several) amino acids of the catalytic domain of SEQ ID NO: 2, SEQ ID NO: 4, or SEQ ID NO: 6; and
 4. (d) a fragment of a catalytic domain of (a), (b), or (c), which has xylanase activity.

[0008] The present invention also relates to isolated polynucleotides encoding the polypeptides of the present invention; nucleic acid constructs; recombinant expression vectors; recombinant host cells comprising the polynucleotides; and methods of producing the polypeptides.

[0009] The present invention also relates to processes for degrading a cellulosic material or xylan-containing material, comprising: treating the cellulosic material or xylan-containing material with an enzyme composition in the presence of a polypeptide having xylanase activity of the present invention. In one aspect, the processes further comprise recovering the degraded or converted cellulosic material or xylan-containing material.

[0010] The present invention also relates to processes of producing a fermentation product, comprising: (a) saccharifying a cellulosic material or xylan-containing material with an enzyme composition in the presence of a polypeptide having xylanase activity of the present invention; (b) fermenting the saccharified cellulosic material or xylan-containing material with one or more (e.g., several) fermenting microorganisms to produce the fermentation product; and (c) recovering the fermentation product from the fermentation.

[0011] The present invention also relates to processes of fermenting a cellulosic material or xylan-containing material, comprising: fermenting the cellulosic material or xylan-containing material with one or more (e.g., several) fermenting microorganisms, wherein the cellulosic material or xylan-containing material is saccharified with an enzyme composition in the presence of a polypeptide having xylanase activity of the present invention. In one aspect, the fermenting of the cellulosic material or xylan-containing material produces a fermentation product. In another aspect, the processes further comprise recovering the fermentation product from the fermentation. Also described is a polynucleotide encoding a signal peptide comprising or consisting of amino acids 1 to 17 of SEQ ID NO: 2, amino acids 1 to 16 of SEQ ID NO: 4, or amino acids 1 to 20 of SEQ ID NO: 6, which is operably linked to a gene encoding a protein; nucleic acid constructs, expression vectors, and recombinant host cells comprising the polynucleotides; and methods of producing a protein.

Definitions

[0012] Xylanase: The term "xylanase" means a 1,4-beta-D-xylan-xylohydrolase (E.C. 3.2.1.8) that catalyzes the endohydrolysis of 1,4-beta-D-xylosidic linkages in xylans. For purposes of the present invention, xylan degrading activity is determined by measuring the increase in hydrolysis of birchwood xylan (Sigma Chemical Co., Inc., St. Louis, MO, USA) by xylan-degrading enzyme(s) under the following typical conditions: 1 ml reactions, 5 mg/ml substrate (total solids), 5 mg of xylanolytic protein/g of substrate, 50 mM sodium acetate pH 5, 50°C, 24 hours, sugar analysis using *p*-hydroxybenzoic acid hydrazide (PHBAH) assay as described by

Lever, 1972, A new reaction for colorimetric determination of carbohydrates, Anal. Biochem 47: 273-279. In one aspect the polypeptide of the present invention have at least 20%, e.g., at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, or at least 100% of the xylanase activity of the mature polypeptide of SEQ ID NO: 6.

[0013] Acetylxylan esterase: The term "acetylxylan esterase" means a carboxylesterase (EC 3.1.1.72) that catalyzes the hydrolysis of acetyl groups from polymeric xylan, acetylated xylose, acetylated glucose, alpha-naphthyl acetate, and *p*-nitrophenyl acetate. For purposes of the present invention, acetylxylan esterase activity is determined using 0.5 mM *p*-nitrophenylacetate as substrate in 50 mM sodium acetate pH 5.0 containing 0.01% TWEEN™ 20 (polyoxyethylene sorbitan monolaurate). One unit of acetylxylan esterase is defined as the amount of enzyme capable of releasing 1 µmole of *p*-nitrophenolate anion per minute at pH 5, 25°C.

[0014] Allelic variant: The term "allelic variant" means any of two or more alternative forms of a gene occupying the same chromosomal locus. Allelic variation arises naturally through mutation, and may result in polymorphism within populations. Gene mutations can be silent (no change in the encoded polypeptide) or may encode polypeptides having altered amino acid sequences. An allelic variant of a polypeptide is a polypeptide encoded by an allelic variant of a gene.

[0015] Alpha-L-arabinofuranosidase: The term "alpha-L-arabinofuranosidase" means an alpha-L-arabinofuranoside arabinofuranohydrolase (EC 3.2.1.55) that catalyzes the hydrolysis of terminal non-reducing alpha-L-arabinofuranoside residues in alpha-L-arabinosides. The enzyme acts on alpha-L-arabinofuranosides, alpha-L-arabinans containing (1,3)- and/or (1,5)-linkages, arabinoxylans, and arabinogalactans. Alpha-L-arabinofuranosidase is also known as arabinosidase, alpha-arabinosidase, alpha-L-arabinosidase, alpha-arabinofuranosidase, polysaccharide alpha-L-arabinofuranosidase, alpha-L-arabinofuranoside hydrolase, L-arabinosidase, or alpha-L-arabinanase. For purposes of the present invention, alpha-L-arabinofuranosidase activity is determined using 5 mg of medium viscosity wheat arabinoxylan (Megazyme International Ireland, Ltd., Bray, Co. Wicklow, Ireland) per ml of 100 mM sodium acetate pH 5 in a total volume of 200 µl for 30 minutes at 40°C followed by arabinose analysis by AMINEX® HPX-87H column chromatography (Bio-Rad Laboratories, Inc., Hercules, CA, USA).

[0016] Alpha-glucuronidase: The term "alpha-glucuronidase" means an alpha-D-glucosiduronate glucuronohydrolase (EC 3.2.1.139) that catalyzes the hydrolysis of an alpha-D-glucuronoside to D-glucuronate and an alcohol. For purposes of the present invention, alpha-glucuronidase activity is determined according to de Vries, 1998, J. Bacteriol. 180: 243-249. One unit of alpha-glucuronidase equals the amount of enzyme capable of releasing 1 µmole of glucuronic or 4-O-methylglucuronic acid per minute at pH 5, 40°C.

[0017] Beta-glucosidase: The term "beta-glucosidase" means a beta-D-glucoside glucosidase (E.C. 3.2.1.21) that catalyzes the hydrolysis of terminal non-reducing beta-D-

glucose residues with the release of beta-D-glucose. For purposes of the present invention, beta-glucosidase activity is determined using *p*-nitrophenyl-beta-D-glucopyranoside as substrate according to the procedure of Venturi et al., 2002, Extracellular beta-D-glucosidase from *Chaetomium thermophilum* var. *coprophilum*: production, purification and some biochemical properties, J. Basic Microbiol. 42: 55-66. One unit of beta-glucosidase is defined as 1.0 μ mole of *p*-nitrophenolate anion produced per minute at 25°C, pH 4.8 from 1 mM *p*-nitrophenyl-beta-D-glucopyranoside as substrate in 50 mM sodium citrate containing 0.01% TWEEN® 20.

[0018] Beta-xylosidase: The term "beta-xylosidase" means a beta-D-xyloside xylohydrolase (E.C. 3.2.1.37) that catalyzes the exo-hydrolysis of short beta (1→4)-xylooligosaccharides to remove successive D-xylose residues from non-reducing termini. For purposes of the present invention, one unit of beta-xylosidase is defined as 1.0 μ mole of *p*-nitrophenolate anion produced per minute at 40°C, pH 5 from 1 mM *p*-nitrophenyl-beta-D-xyloside as substrate in 100 mM sodium citrate containing 0.01% TWEEN® 20.

[0019] cDNA: The term "cDNA" means a DNA molecule that can be prepared by reverse transcription from a mature, spliced, mRNA molecule obtained from a eukaryotic or prokaryotic cell. cDNA lacks intron sequences that may be present in the corresponding genomic DNA. The initial, primary RNA transcript is a precursor to mRNA that is processed through a series of steps, including splicing, before appearing as mature spliced mRNA.

[0020] Cellobiohydrolase: The term "cellobiohydrolase" means a 1,4-beta-D-glucan cellobiohydrolase (E.C. 3.2.1.91) that catalyzes the hydrolysis of 1,4-beta-D-glucosidic linkages in cellulose, celooligosaccharides, or any beta-1,4-linked glucose containing polymer, releasing cellobiose from the reducing or non-reducing ends of the chain (Teeri, 1997, Crystalline cellulose degradation: New insight into the function of cellobiohydrolases, Trends in Biotechnology 15: 160-167; Teeri et al., 1998, Trichoderma reesei cellobiohydrolases: why so efficient on crystalline cellulose?, Biochem. Soc. Trans. 26: 173-178). Cellobiohydrolase activity is determined according to the procedures described by Lever et al., 1972, Anal. Biochem. 47: 273-279; van Tilbeurgh et al., 1982, FEBS Letters, 149: 152-156; van Tilbeurgh and Claeyssens, 1985, FEBS Letters, 187: 283-288; and Tomme et al., 1988, Eur. J. Biochem. 170: 575-581. In the present invention, the Tomme *et al.* method can be used to determine cellobiohydrolase activity.

[0021] Cellulosic material: The term "cellulosic material" means any material containing cellulose. The predominant polysaccharide in the primary cell wall of biomass is cellulose, the second most abundant is hemicellulose, and the third is pectin. The secondary cell wall, produced after the cell has stopped growing, also contains polysaccharides and is strengthened by polymeric lignin covalently cross-linked to hemicellulose. Cellulose is a homopolymer of anhydrocellobiose and thus a linear beta-(1-4)-D-glucan, while hemicelluloses include a variety of compounds, such as xylans, xyloglucans, arabinoxylans, and mannans in complex branched structures with a spectrum of substituents. Although generally

polymorphous, cellulose is found in plant tissue primarily as an insoluble crystalline matrix of parallel glucan chains. Hemicelluloses usually hydrogen bond to cellulose, as well as to other hemicelluloses, which help stabilize the cell wall matrix.

[0022] Cellulose is generally found, for example, in the stems, leaves, hulls, husks, and cobs of plants or leaves, branches, and wood of trees. The cellulosic material can be, but is not limited to, agricultural residue, herbaceous material (including energy crops), municipal solid waste, pulp and paper mill residue, waste paper, and wood (including forestry residue) (see, for example, Wiselogle et al., 1995, in Handbook on Bioethanol (Charles E. Wyman, editor), pp.105-118, Taylor & Francis, Washington D.C.; Wyman, 1994, Bioresource Technology 50: 3-16; Lynd, 1990, Applied Biochemistry and Biotechnology 24/25: 695-719; Mosier et al., 1999, Recent Progress in Bioconversion of Lignocellulosics, in Advances in Biochemical Engineering/Biotechnology, T. Scheper, managing editor, Volume 65, pp.23-40, Springer-Verlag, New York). It is understood herein that the cellulose may be in the form of lignocellulose, a plant cell wall material containing lignin, cellulose, and hemicellulose in a mixed matrix. In a preferred aspect, the cellulosic material is any biomass material. In another preferred aspect, the cellulosic material is lignocellulose, which comprises cellulose, hemicelluloses, and lignin.

[0023] In one aspect, the cellulosic material is agricultural residue. In another aspect, the cellulosic material is herbaceous material (including energy crops). In another aspect, the cellulosic material is municipal solid waste. In another aspect, the cellulosic material is pulp and paper mill residue. In another aspect, the cellulosic material is waste paper. In another aspect, the cellulosic material is wood (including forestry residue).

[0024] In another aspect, the cellulosic material is arundo. In another aspect, the cellulosic material is bagasse. In another aspect, the cellulosic material is bamboo. In another aspect, the cellulosic material is corn cob. In another aspect, the cellulosic material is corn fiber. In another aspect, the cellulosic material is corn stover. In another aspect, the cellulosic material is miscanthus. In another aspect, the cellulosic material is orange peel. In another aspect, the cellulosic material is rice straw. In another aspect, the cellulosic material is switchgrass. In another aspect, the cellulosic material is wheat straw.

[0025] In another aspect, the cellulosic material is aspen. In another aspect, the cellulosic material is eucalyptus. In another aspect, the cellulosic material is fir. In another aspect, the cellulosic material is pine. In another aspect, the cellulosic material is poplar. In another aspect, the cellulosic material is spruce. In another aspect, the cellulosic material is willow.

[0026] In another aspect, the cellulosic material is algal cellulose. In another aspect, the cellulosic material is bacterial cellulose. In another aspect, the cellulosic material is cotton linter. In another aspect, the cellulosic material is filter paper. In another aspect, the cellulosic material is microcrystalline cellulose. In another aspect, the cellulosic material is phosphoric-acid treated cellulose.

[0027] In another aspect, the cellulosic material is an aquatic biomass. As used herein the term "aquatic biomass" means biomass produced in an aquatic environment by a photosynthesis process. The aquatic biomass can be algae, emergent plants, floating-leaf plants, or submerged plants.

[0028] The cellulosic material may be used as is or may be subjected to pretreatment, using conventional methods known in the art, as described herein. In a preferred aspect, the cellulosic material is pretreated.

[0029] Cellulolytic enzyme or cellulase: The term "cellulolytic enzyme" or "cellulase" means one or more (*e.g.*, several) enzymes that hydrolyze a cellulosic material. Such enzymes include endoglucanase(s), cellobiohydrolase(s), beta-glucosidase(s), or combinations thereof. The two basic approaches for measuring cellulolytic activity include: (1) measuring the total cellulolytic activity, and (2) measuring the individual cellulolytic activities (endoglucanases, cellobiohydrolases, and beta-glucosidases) as reviewed in Zhang et al., Outlook for cellulase improvement: Screening and selection strategies, 2006, Biotechnology Advances 24: 452-481. Total cellulolytic activity is usually measured using insoluble substrates, including Whatman N°1 filter paper, microcrystalline cellulose, bacterial cellulose, algal cellulose, cotton, pretreated lignocellulose, *etc.* The most common total cellulolytic activity assay is the filter paper assay using Whatman N°1 filter paper as the substrate. The assay was established by the International Union of Pure and Applied Chemistry (IUPAC) (Ghose, 1987, Measurement of cellulase activities, Pure Appl. Chem. 59: 257-68).

[0030] For purposes of the present invention, cellulolytic enzyme activity is determined by measuring the increase in hydrolysis of a cellulosic material by cellulolytic enzyme(s) under the following conditions: 1-50 mg of cellulolytic enzyme protein/g of cellulose in PCS (or other pretreated cellulosic material) for 3-7 days at a suitable temperature, *e.g.*, 50°C, 55°C, or 60°C, compared to a control hydrolysis without addition of cellulolytic enzyme protein. Typical conditions are 1 ml reactions, washed or unwashed PCS, 5% insoluble solids, 50 mM sodium acetate pH 5, 1 mM MnSO₄, 50°C, 55°C, or 60°C, 72 hours, sugar analysis by AMINEX® HPX-87H column (Bio-Rad Laboratories, Inc., Hercules, CA, USA).

[0031] Coding sequence: The term "coding sequence" means a polynucleotide, which directly specifies the amino acid sequence of a polypeptide. The boundaries of the coding sequence are generally determined by an open reading frame, which begins with a start codon such as ATG, GTG, or TTG and ends with a stop codon such as TAA, TAG, or TGA. The coding sequence may be a genomic DNA, cDNA, synthetic DNA, or a combination thereof. The coding sequence may be from positions 1-240 and 314-1168 of SEQ ID NO: 1 or from positions 1-241, 302-342, 404-452, 518-639, 707-852, 912-1019, 1088-1205, 1282-1347, and 1430-1708 of SEQ ID NO: 3. In an embodiment the coding sequence is positions 1-50, 114-270, 342-474, 567-680, 757-1520 of SEQ ID NO: 5.

[0032] Control sequences: The term "control sequences" means nucleic acid sequences necessary for expression of a polynucleotide encoding a mature polypeptide of the present

invention. Each control sequence may be native (*i.e.*, from the same gene) or foreign (*i.e.*, from a different gene) to the polynucleotide encoding the polypeptide or native or foreign to each other. Such control sequences include, but are not limited to, a leader, polyadenylation sequence, propeptide sequence, promoter, signal peptide sequence, and transcription terminator. At a minimum, the control sequences include a promoter, and transcriptional and translational stop signals. The control sequences may be provided with linkers for the purpose of introducing specific restriction sites facilitating ligation of the control sequences with the coding region of the polynucleotide encoding a polypeptide.

[0033] Endoglucanase: The term "endoglucanase" means an endo-1,4-(1,3;1,4)-beta-D-glucan 4-glucanohydrolase (E.C. 3.2.1.4) that catalyzes endohydrolysis of 1,4-beta-D-glycosidic linkages in cellulose, cellulose derivatives (such as carboxymethyl cellulose and hydroxyethyl cellulose), lichenin, beta-1,4 bonds in mixed beta-1,3 glucans such as cereal beta-D-glucans or xyloglucans, and other plant material containing cellulosic components. Endoglucanase activity can be determined by measuring reduction in substrate viscosity or increase in reducing ends determined by a reducing sugar assay (Zhang et al., 2006, *Biotechnology Advances* 24: 452-481). For purposes of the present invention, endoglucanase activity is determined using carboxymethyl cellulose (CMC) as substrate according to the procedure of Ghose, 1987, *Pure and Appl. Chem.* 59: 257-268, at pH 5, 40°C.

[0034] Expression: The term "expression" includes any step involved in the production of a polypeptide including, but not limited to, transcription, post-transcriptional modification, translation, post-translational modification, and secretion.

[0035] Expression vector: The term "expression vector" means a linear or circular DNA molecule that comprises a polynucleotide encoding a polypeptide and is operably linked to control sequences that provide for its expression.

[0036] Family 61 glycoside hydrolase: The term "Family 61 glycoside hydrolase" or "Family GH61" or "GH61" means a polypeptide falling into the glycoside hydrolase Family 61 according to Henrissat, 1991, *A classification of glycosyl hydrolases based on amino-acid sequence similarities*, *Biochem. J.* 280: 309-316, and Henrissat and Bairoch, 1996, *Updating the sequence-based classification of glycosyl hydrolases*, *Biochem. J.* 316: 695-696. The enzymes in this family were originally classified as a glycoside hydrolase family based on measurement of very weak endo-1,4-beta-D-glucanase activity in one family member. The structure and mode of action of these enzymes are non-canonical and they cannot be considered as bona fide glycosidases. However, they are kept in the CAZy classification on the basis of their capacity to enhance the breakdown of lignocellulose when used in conjunction with a cellulase or a mixture of cellulases.

[0037] Feruloyl esterase: The term "feruloyl esterase" means a 4-hydroxy-3-methoxycinnamoyl-sugar hydrolase (EC 3.1.1.73) that catalyzes the hydrolysis of 4-hydroxy-3-methoxycinnamoyl (feruloyl) groups from esterified sugar, which is usually arabinose in "natural" substrates, to produce ferulate (4-hydroxy-3-methoxycinnamate). Feruloyl esterase is

also known as ferulic acid esterase, hydroxycinnamoyl esterase, FAE-III, cinnamoyl ester hydrolase, FAEA, cinnAE, FAE-I, or FAE-II. For purposes of the present invention, feruloyl esterase activity is determined using 0.5 mM *p*-nitrophenylferulate as substrate in 50 mM sodium acetate pH 5.0. One unit of feruloyl esterase equals the amount of enzyme capable of releasing 1 μ mole of *p*-nitrophenolate anion per minute at pH 5, 25°C.

[0038] Fragment: The term "fragment" means a polypeptide having one or more (e.g., several) amino acids absent from the amino and/or carboxyl terminus of a mature polypeptide main; wherein the fragment has xylanase activity. A fragment may contain at least amino acid residues 18-364 of SEQ ID NO: 2 or at least amino acid residues 17-326 of SEQ ID NO: 4. In an aspect a fragment contains at least amino acid residues 21-337 of SEQ ID NO: 6.

[0039] Hemicellulolytic enzyme or hemicellulase: The term "hemicellulolytic enzyme" or "hemicellulase" means one or more (e.g., several) enzymes that hydrolyze a hemicellulosic material. See, for example, Shallom and Shoham, 2003, Microbial hemicellulases. Current Opinion In Microbiology 6(3): 219-228). Hemicellulases are key components in the degradation of plant biomass. Examples of hemicellulases include, but are not limited to, an acetylmannan esterase, an acetylxylan esterase, an arabinanase, an arabinofuranosidase, a coumaric acid esterase, a feruloyl esterase, a galactosidase, a glucuronidase, a glucuronoyl esterase, a mannanase, a mannosidase, a xylanase, and a xylosidase. The substrates of these enzymes, the hemicelluloses, are a heterogeneous group of branched and linear polysaccharides that are bound via hydrogen bonds to the cellulose microfibrils in the plant cell wall, crosslinking them into a robust network. Hemicelluloses are also covalently attached to lignin, forming together with cellulose a highly complex structure. The variable structure and organization of hemicelluloses require the concerted action of many enzymes for its complete degradation. The catalytic modules of hemicellulases are either glycoside hydrolases (GHs) that hydrolyze glycosidic bonds, or carbohydrate esterases (CEs), which hydrolyze ester linkages of acetate or ferulic acid side groups. These catalytic modules, based on homology of their primary sequence, can be assigned into GH and CE families. Some families, with an overall similar fold, can be further grouped into clans, marked alphabetically (e.g., GH-A). A most informative and updated classification of these and other carbohydrate active enzymes is available in the Carbohydrate-Active Enzymes (CAZy) database. Hemicellulolytic enzyme activities can be measured according to Ghose and Bisaria, 1987, Pure & Appl. Chem. 59: 1739-1752, at a suitable temperature, e.g., 50°C, 55°C, or 60°C, and pH, e.g., 5.0 or 5.5.

[0040] High stringency conditions: The term "high stringency conditions" means for probes of at least 100 nucleotides in length, prehybridization and hybridization at 42°C in 5X SSPE, 0.3% SDS, 200 micrograms/ml sheared and denatured salmon sperm DNA, and 50% formamide, following standard Southern blotting procedures for 12 to 24 hours. The carrier material is finally washed three times each for 15 minutes using 2X SSC, 0.2% SDS at 65°C.

[0041] Host cell: The term "host cell" means any cell type that is susceptible to transformation, transfection, transduction, or the like with a nucleic acid construct or expression vector comprising a polynucleotide of the present invention. The term "host cell" encompasses

any progeny of a parent cell that is not identical to the parent cell due to mutations that occur during replication.

[0042] Isolated: The term "isolated" means a substance in a form or environment that does not occur in nature. Non-limiting examples of isolated substances include (1) any non-naturally occurring substance, (2) any substance including, but not limited to, any enzyme, variant, nucleic acid, protein, peptide or cofactor, that is at least partially removed from one or more or all of the naturally occurring constituents with which it is associated in nature; (3) any substance modified by the hand of man relative to that substance found in nature; or (4) any substance modified by increasing the amount of the substance relative to other components with which it is naturally associated (e.g., multiple copies of a gene encoding the substance; use of a stronger promoter than the promoter naturally associated with the gene encoding the substance). An isolated substance may be present in a fermentation broth sample.

[0043] Low stringency conditions: The term "low stringency conditions" means for probes of at least 100 nucleotides in length, prehybridization and hybridization at 42°C in 5X SSPE, 0.3% SDS, 200 micrograms/ml sheared and denatured salmon sperm DNA, and 25% formamide, following standard Southern blotting procedures for 12 to 24 hours. The carrier material is finally washed three times each for 15 minutes using 2X SSC, 0.2% SDS at 50°C.

[0044] Mature polypeptide: The term "mature polypeptide" means a polypeptide in its final form following translation and any post-translational modifications, such as N-terminal processing, C-terminal truncation, glycosylation, phosphorylation, etc. A mature polypeptide may be amino acids 18 to 364 of SEQ ID NO: 2 based on the SignalP program (Nielsen et al., 1997, Protein Engineering 10: 1-6) that predicts amino acids 1 to 17 of SEQ ID NO: 2 are a signal peptide. A mature polypeptide may also be amino acids 17 to 389 of SEQ ID NO: 4 based on the SignalP program that predicts amino acids 1 to 16 of SEQ ID NO: 4 are a signal peptide. In an aspect, the mature polypeptide is amino acids 21 to 405 of SEQ ID NO: 6 based on the SignalP program that predicts amino acids 1 to 20 of SEQ ID NO: 6 are a signal peptide. It is known in the art that a host cell may produce a mixture of two or more different mature polypeptides (*i.e.*, with a different C-terminal and/or N-terminal amino acid) expressed by the same polynucleotide.

[0045] Mature polypeptide coding sequence: The term "mature polypeptide coding sequence" means a polynucleotide that encodes a mature polypeptide having xylanase activity. A mature polypeptide coding sequence may be nucleotides 52 to 1165 of SEQ ID NO: 1 or the cDNA sequence thereof based on the SignalP program (Nielsen *et al.*, 1997, *supra*) that predicts nucleotides 1 to 51 of SEQ ID NO: 1 encode a signal peptide. A mature polypeptide coding sequence may also be nucleotides 49 to 1705 of SEQ ID NO: 3 or the cDNA sequence thereof based on the SignalP program that predicts nucleotides 1 to 48 of SEQ ID NO: 3 encode a signal peptide. In an aspect, the mature polypeptide coding sequence is nucleotides 124 to 1517 of SEQ ID NO: 5 or the cDNA sequence thereof based on the SignalP program that predicts nucleotides 1 to 123 of SEQ ID NO: 5 encode a signal peptide.

[0046] Catalytic domain: The term "catalytic domain" means the portion of an enzyme containing the catalytic machinery of the enzyme.

[0047] Cellulose binding domain: The term "cellulose binding domain" means the portion of an enzyme that mediates binding of the enzyme to amorphous regions of a cellulose substrate. The cellulose binding domain (CBD) is found either at the N-terminal or at the C-terminal extremity of an enzyme. A CBD is also referred to as a cellulose binding module or CBM. The CBM may be amino acids 354 to 389 of SEQ ID NO: 4. In one embodiment the CBM is amino acids 370 to 405 of SEQ ID NO: 6. The CBM is separated from the catalytic domain by a linker sequence. The linker may be amino acids 327 to 353 of SEQ ID NO: 4. The linker is in one embodiment amino acids 338 to 369 of SEQ ID NO: 6.

[0048] Medium stringency conditions: The term "medium stringency conditions" means for probes of at least 100 nucleotides in length, prehybridization and hybridization at 42°C in 5X SSPE, 0.3% SDS, 200 micrograms/ml sheared and denatured salmon sperm DNA, and 35% formamide, following standard Southern blotting procedures for 12 to 24 hours. The carrier material is finally washed three times each for 15 minutes using 2X SSC, 0.2% SDS at 55°C.

[0049] Medium-high stringency conditions: The term "medium-high stringency conditions" means for probes of at least 100 nucleotides in length, prehybridization and hybridization at 42°C in 5X SSPE, 0.3% SDS, 200 micrograms/ml sheared and denatured salmon sperm DNA, and either 35% formamide, following standard Southern blotting procedures for 12 to 24 hours. The carrier material is finally washed three times each for 15 minutes using 2X SSC, 0.2% SDS at 60°C.

[0050] Nucleic acid construct: The term "nucleic acid construct" means a nucleic acid molecule, either single- or double-stranded, which is isolated from a naturally occurring gene or is modified to contain segments of nucleic acids in a manner that would not otherwise exist in nature or which is synthetic, which comprises one or more control sequences.

[0051] Operably linked: The term "operably linked" means a configuration in which a control sequence is placed at an appropriate position relative to the coding sequence of a polynucleotide such that the control sequence directs expression of the coding sequence.

[0052] Polypeptide having cellulolytic enhancing activity: The term "polypeptide having cellulolytic enhancing activity" means a GH61 polypeptide that catalyzes the enhancement of the hydrolysis of a cellulosic material by enzyme having cellulolytic activity. For purposes of the present invention, cellulolytic enhancing activity is determined by measuring the increase in reducing sugars or the increase of the total of cellobiose and glucose from the hydrolysis of a cellulosic material by cellulolytic enzyme under the following conditions: 1-50 mg of total protein/g of cellulose in PCS, wherein total protein is comprised of 50-99.5% w/w cellulolytic enzyme protein and 0.5-50% w/w protein of a GH61 polypeptide having cellulolytic enhancing activity for 1-7 days at a suitable temperature, e.g., 50°C, 55°C, or 60°C, and pH, e.g., 5.0 or 5.5, compared to a control hydrolysis with equal total protein loading without cellulolytic

enhancing activity (1-50 mg of cellulolytic protein/g of cellulose in PCS). In a preferred aspect, a mixture of CELLUCLAST® 1.5L (Novozymes A/S, Bagsværd, Denmark) in the presence of 2-3% of total protein weight *Aspergillus oryzae* beta-glucosidase (recombinantly produced in *Aspergillus oryzae* according to WO 02/095014) or 2-3% of total protein weight *Aspergillus fumigatus* beta-glucosidase (recombinantly produced in *Aspergillus oryzae* as described in WO 02/095014) of cellulase protein loading is used as the source of the cellulolytic activity.

[0053] The GH61 polypeptides having cellulolytic enhancing activity enhance the hydrolysis of a cellulosic material catalyzed by enzyme having cellulolytic activity by reducing the amount of cellulolytic enzyme required to reach the same degree of hydrolysis preferably at least 1.01-fold, e.g., at least 1.05-fold, at least 1.10-fold, at least 1.25-fold, at least 1.5-fold, at least 2-fold, at least 3-fold, at least 4-fold, at least 5-fold, at least 10-fold, or at least 20-fold.

[0054] Pretreated corn stover: The term "PCS" or "Pretreated Corn Stover" means a cellulosic material derived from corn stover by treatment with heat and dilute sulfuric acid, alkaline pretreatment, or neutral pretreatment.

[0055] Sequence identity: The relatedness between two amino acid sequences or between two nucleotide sequences is described by the parameter "sequence identity".

[0056] For purposes of the present invention, the sequence identity between two amino acid sequences is determined using the Needleman-Wunsch algorithm (Needleman and Wunsch, 1970, J. Mol. Biol. 48: 443-453) as implemented in the Needle program of the EMBOSS package (EMBOSS: The European Molecular Biology Open Software Suite, Rice et al., 2000, Trends Genet. 16: 276-277), preferably version 5.0.0 or later. The parameters used are gap open penalty of 10, gap extension penalty of 0.5, and the EBLOSUM62 (EMBOSS version of BLOSUM62) substitution matrix. The output of Needle labeled "longest identity" (obtained using the -nobrief option) is used as the percent identity and is calculated as follows:

$$(\text{Identical Residues} \times 100) / (\text{Length of Alignment} - \text{Total Number of Gaps in Alignment})$$

[0057] For purposes of the present invention, the sequence identity between two deoxyribonucleotide sequences is determined using the Needleman-Wunsch algorithm (Needleman and Wunsch, 1970, *supra*) as implemented in the Needle program of the EMBOSS package (EMBOSS: The European Molecular Biology Open Software Suite, Rice et al., 2000, *supra*), preferably version 5.0.0 or later. The parameters used are gap open penalty of 10, gap extension penalty of 0.5, and the EDNAFULL (EMBOSS version of NCBI NUC4.4) substitution matrix. The output of Needle labeled "longest identity" (obtained using the -nobrief option) is used as the percent identity and is calculated as follows:

$$(\text{Identical Deoxyribonucleotides} \times 100) / (\text{Length of Alignment} - \text{Total Number of Gaps in Alignment})$$

[0058] Subsequence: The term "subsequence" means a polynucleotide having one or more

(e.g., several) nucleotides absent from the 5' and/or 3' end of a mature polypeptide coding sequence; wherein the subsequence encodes a fragment having xylanase activity. In one aspect, a subsequence corresponds to the polynucleotide encoding the catalytic domain.

[0059] Variant: The term "variant" means a polypeptide having xylanase activity comprising an alteration, *i.e.*, a substitution, insertion, and/or deletion, at one or more (e.g., several) positions. A substitution means replacement of the amino acid occupying a position with a different amino acid; a deletion means removal of the amino acid occupying a position; and an insertion means adding an amino acid adjacent to and immediately following the amino acid occupying a position.

[0060] Very high stringency conditions: The term "very high stringency conditions" means for probes of at least 100 nucleotides in length, prehybridization and hybridization at 42°C in 5X SSPE, 0.3% SDS, 200 micrograms/ml sheared and denatured salmon sperm DNA, and 50% formamide, following standard Southern blotting procedures for 12 to 24 hours. The carrier material is finally washed three times each for 15 minutes using 2X SSC, 0.2% SDS at 70°C.

[0061] Very low stringency conditions: The term "very low stringency conditions" means for probes of at least 100 nucleotides in length, prehybridization and hybridization at 42°C in 5X SSPE, 0.3% SDS, 200 micrograms/ml sheared and denatured salmon sperm DNA, and 25% formamide, following standard Southern blotting procedures for 12 to 24 hours. The carrier material is finally washed three times each for 15 minutes using 2X SSC, 0.2% SDS at 45°C.

[0062] Xylan-containing material: The term "xylan-containing material" means any material comprising a plant cell wall polysaccharide containing a backbone of beta-(1-4)-linked xylose residues. Xylans of terrestrial plants are heteropolymers possessing a beta-(1-4)-D-xylopyranose backbone, which is branched by short carbohydrate chains. They comprise D-glucuronic acid or its 4-O-methyl ether, L-arabinose, and/or various oligosaccharides, composed of D-xylose, L-arabinose, D- or L-galactose, and D-glucose. Xylan-type polysaccharides can be divided into homoxylans and heteroxylans, which include glucuronoxylans, (arabino)glucuronoxylans, (glucurono)arabinoxylans, arabinoxylans, and complex heteroxylans. See, for example, Ebringerova et al., 2005, Adv. Polym. Sci. 186: 1-67.

[0063] In the processes of the present invention, any material containing xylan may be used. In a preferred aspect, the xylan-containing material is lignocellulose.

[0064] Xylan degrading activity or xylanolytic activity: The term "xylan degrading activity" or "xylanolytic activity" means a biological activity that hydrolyzes xylan-containing material. The two basic approaches for measuring xylanolytic activity include: (1) measuring the total xylanolytic activity, and (2) measuring the individual xylanolytic activities (e.g., endoxylanases, beta-xylosidases, arabinofuranosidases, alpha-glucuronidases, acetylxylan esterases, feruloyl esterases, and alpha-glucuronyl esterases). Recent progress in assays of xylanolytic enzymes was summarized in several publications including Biely and Puchard, 2006, Recent progress in

the assays of xylanolytic enzymes, Journal of the Science of Food and Agriculture 86(11): 1636-1647; Spanikova and Biely, 2006, Glucuronoyl esterase - Novel carbohydrate esterase produced by Schizophyllum commune, FEBS Letters 580(19): 4597-4601; Herrmann et al., 1997, The beta-D-xylosidase of Trichoderma reesei is a multifunctional beta-D-xylan xylohydrolase, Biochemical Journal 321: 375-381.

[0065] Total xylan degrading activity can be measured by determining the reducing sugars formed from various types of xylan, including, for example, oat spelt, beechwood, and larchwood xylans, or by photometric determination of dyed xylan fragments released from various covalently dyed xylans. The most common total xylanolytic activity assay is based on production of reducing sugars from polymeric 4-O-methyl glucuronoxylan as described in Bailey et al., 1992, Interlaboratory testing of methods for assay of xylanase activity, Journal of Biotechnology 23(3): 257-270. Xylanase activity can also be determined with 0.2% AZCL-arabinoxylan as substrate in 0.01% TRITON® X-100 (4-(1,1,3,3-tetramethylbutyl)phenylpolyethylene glycol) and 200 mM sodium phosphate buffer pH 6 at 37°C. One unit of xylanase activity is defined as 1.0 µmole of azurine produced per minute at 37°C, pH 6 from 0.2% AZCL-arabinoxylan as substrate in 200 mM sodium phosphate pH 6 buffer.

[0066] For purposes of the present invention, xylan degrading activity is determined by measuring the increase in hydrolysis of birchwood xylan (Sigma Chemical Co., Inc., St. Louis, MO, USA) by xylan-degrading enzyme(s) under the following typical conditions: 1 ml reactions, 5 mg/ml substrate (total solids), 5 mg of xylanolytic protein/g of substrate, 50 mM sodium acetate pH 5, 50°C, 24 hours, sugar analysis using *p*-hydroxybenzoic acid hydrazide (PHBAH) assay as described by Lever, 1972, A new reaction for colorimetric determination of carbohydrates, Anal. Biochem 47: 273-279.

Detailed Description of the Invention

Polypeptides Having Xylanase Activity

[0067] Described are polypeptides having a sequence identity to the mature polypeptide of SEQ ID NO: 2 of at least 77%, e.g., at least 78%, at least 79%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100%, which have xylanase activity. Also described are isolated polypeptides having a sequence identity to the mature polypeptide of SEQ ID NO: 4 of at least 77%, e.g., at least 78%, at least 79%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100%, which have xylanase activity. In an embodiment, the present invention relates to isolated polypeptides having a sequence identity to the mature polypeptide of SEQ ID NO: 6 of at least 85%, e.g., at least 86%, at least 87%, at least 88%, at least 89%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at

least 97%, at least 98%, at least 99%, or 100%, which have xylanase activity. In one aspect, the polypeptides differ by no more than 10 amino acids, e.g., 1, 2, 3, 4, 5, 6, 7, 8, or 9, from the mature polypeptide of SEQ ID NO: 6.

[0068] A polypeptide of the present invention preferably comprises or consists of the amino acid sequence of SEQ ID NO: 6 or an allelic variant thereof; or is a fragment thereof having xylanase activity. In another aspect, the polypeptide comprises or consists of the mature polypeptide of SEQ ID NO: 6. In another aspect, the polypeptide comprises or consists of amino acids 21 to 405 of SEQ ID NO: 6. Also described is an isolated polypeptide having xylanase activity encoded by a polynucleotide that hybridizes under very low stringency conditions, or low stringency conditions, or medium stringency conditions, or medium-high stringency conditions, or high stringency conditions, or very high stringency conditions with (i) the mature polypeptide coding sequence of SEQ ID NO: 1, SEQ ID NO: 3, or SEQ ID NO: 5, (ii) the cDNA sequence thereof, or (iii) the full-length complement of (i) or (ii) (Sambrook et al., 1989, Molecular Cloning, A Laboratory Manual, 2d edition, Cold Spring Harbor, New York).

[0069] The polynucleotide of SEQ ID NO: 1, SEQ ID NO: 3, or SEQ ID NO: 5, or a subsequence thereof, as well as the polypeptide of SEQ ID NO: 2, SEQ ID NO: 4, or SEQ ID NO: 6, or a fragment thereof, may be used to design nucleic acid probes to identify and clone DNA encoding polypeptides having xylanase activity from strains of different genera or species according to methods well known in the art. In particular, such probes can be used for hybridization with the genomic DNA or cDNA of a cell of interest, following standard Southern blotting procedures, in order to identify and isolate the corresponding gene therein. Such probes can be considerably shorter than the entire sequence, but should be at least 15, e.g., at least 25, at least 35, or at least 70 nucleotides in length. Preferably, the nucleic acid probe is at least 100 nucleotides in length, e.g., at least 200 nucleotides, at least 300 nucleotides, at least 400 nucleotides, at least 500 nucleotides, at least 600 nucleotides, at least 700 nucleotides, at least 800 nucleotides, or at least 900 nucleotides in length. Both DNA and RNA probes can be used. The probes are typically labeled for detecting the corresponding gene (for example, with ^{32}P , ^3H , ^{35}S , biotin, or avidin). Such probes are encompassed by the present invention.

[0070] A genomic DNA or cDNA library prepared from such other strains may be screened for DNA that hybridizes with the probes described above and encodes a polypeptide having xylanase activity. Genomic or other DNA from such other strains may be separated by agarose or polyacrylamide gel electrophoresis, or other separation techniques. DNA from the libraries or the separated DNA may be transferred to and immobilized on nitrocellulose or other suitable carrier material. In order to identify a clone or DNA that hybridizes with SEQ ID NO: 1, SEQ ID NO: 3, or SEQ ID NO: 5 or a subsequence thereof, the carrier material is used in a Southern blot. Hybridization indicates that the polynucleotide hybridizes to a labeled nucleic acid probe corresponding to (i) SEQ ID NO: 1, SEQ ID NO: 3, or SEQ ID NO: 5; (ii) the mature polypeptide coding sequence of SEQ ID NO: 1, SEQ ID NO: 3, or SEQ ID NO: 5; (iii) the cDNA sequence thereof; (iv) the full-length complement thereof; or (v) a subsequence thereof; under very low to very high stringency conditions. Molecules to which the nucleic acid probe hybridizes under

these conditions can be detected using, for example, X-ray film or any other detection means known in the art. The nucleic acid probe is a polynucleotide that encodes the polypeptide of SEQ ID NO: 2, SEQ ID NO: 4, or SEQ ID NO: 6; the mature polypeptide thereof; or a fragment thereof. In another aspect, the nucleic acid probe is SEQ ID NO: 1, SEQ ID NO: 3, or SEQ ID NO: 5; or the cDNA sequence thereof. The nucleic acid probe may also be the polynucleotide contained in *Talaromyces leycettanus* Strain CBS 398.68, wherein the polynucleotide encodes a polypeptide having xylanase activity. Also described is an isolated polypeptide having xylanase activity encoded by a polynucleotide having a sequence identity to the mature polypeptide coding sequence of SEQ ID NO: 1, or the cDNA sequence thereof, of at least 60%, e.g., at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100%. Also described is an isolated polypeptide having xylanase activity encoded by a polynucleotide having a sequence identity to the mature polypeptide coding sequence of SEQ ID NO: 3, or the cDNA sequence thereof, of at least 60%, e.g., at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100%. In another embodiment, the present invention relates to an isolated polypeptide having xylanase activity encoded by a polynucleotide having a sequence identity to the mature polypeptide coding sequence of SEQ ID NO: 5, or the cDNA sequence thereof, of at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100%. Also described are variants of the mature polypeptide of SEQ ID NO: 2, SEQ ID NO: 4, or SEQ ID NO: 6 comprising a substitution, deletion, and/or insertion at one or more (e.g., several) positions. The number of amino acid substitutions, deletions and/or insertions introduced into the mature polypeptide of SEQ ID NO: 2, SEQ ID NO: 4, or SEQ ID NO: 6 is not more than 10, e.g., 1, 2, 3, 4, 5, 6, 7, 8 or 9. The amino acid changes may be of a minor nature, that is conservative amino acid substitutions or insertions that do not significantly affect the folding and/or activity of the protein; small deletions, typically of 1-30 amino acids; small amino- or carboxyl-terminal extensions, such as an amino-terminal methionine residue; a small linker peptide of up to 20-25 residues; or a small extension that facilitates purification by changing net charge or another function, such as a poly-histidine tract, an antigenic epitope or a binding domain.

[0071] Examples of conservative substitutions are within the groups of basic amino acids (arginine, lysine and histidine), acidic amino acids (glutamic acid and aspartic acid), polar amino acids (glutamine and asparagine), hydrophobic amino acids (leucine, isoleucine and valine), aromatic amino acids (phenylalanine, tryptophan and tyrosine), and small amino acids (glycine, alanine, serine, threonine and methionine). Amino acid substitutions that do not generally alter specific activity are known in the art and are described, for example, by H. Neurath and R.L. Hill, 1979, In, *The Proteins*, Academic Press, New York. Common substitutions are Ala/Ser, Val/Ile, Asp/Glu, Thr/Ser, Ala/Gly, Ala/Thr, Ser/Asn, Ala/Val, Ser/Gly, Tyr/Phe, Ala/Pro, Lys/Arg, Asp/Asn, Leu/Ile, Leu/Val, Ala/Glu, and Asp/Gly.

[0072] Alternatively, the amino acid changes are of such a nature that the physico-chemical properties of the polypeptides are altered. For example, amino acid changes may improve the

thermal stability of the polypeptide, alter the substrate specificity, change the pH optimum, and the like.

[0073] Essential amino acids in a polypeptide can be identified according to procedures known in the art, such as site-directed mutagenesis or alanine-scanning mutagenesis (Cunningham and Wells, 1989, *Science* 244: 1081-1085). In the latter technique, single alanine mutations are introduced at every residue in the molecule, and the resultant mutant molecules are tested for xylanase activity to identify amino acid residues that are critical to the activity of the molecule. See also, Hilton et al., 1996, *J. Biol. Chem.* 271: 4699-4708. The active site of the enzyme or other biological interaction can also be determined by physical analysis of structure, as determined by such techniques as nuclear magnetic resonance, crystallography, electron diffraction, or photoaffinity labeling, in conjunction with mutation of putative contact site amino acids. See, for example, de Vos et al., 1992, *Science* 255: 306-312; Smith et al., 1992, *J. Mol. Biol.* 224: 899-904; Wlodaver et al., 1992, *FEBS Lett.* 309: 59-64. The identity of essential amino acids can also be inferred from an alignment with a related polypeptide.

[0074] Single or multiple amino acid substitutions, deletions, and/or insertions can be made and tested using known methods of mutagenesis, recombination, and/or shuffling, followed by a relevant screening procedure, such as those disclosed by Reidhaar-Olson and Sauer, 1988, *Science* 241: 53-57; Bowie and Sauer, 1989, *Proc. Natl. Acad. Sci. USA* 86: 2152-2156; WO 95/17413; or WO 95/22625. Other methods that can be used include error-prone PCR, phage display (e.g., Lowman et al., 1991, *Biochemistry* 30: 10832-10837; U.S. Patent No. 5,223,409; WO 92/06204), and region-directed mutagenesis (Derbyshire et al., 1986, *Gene* 46: 145; Ner et al., 1988, *DNA* 7: 127).

[0075] Mutagenesis/shuffling methods can be combined with high-throughput, automated screening methods to detect activity of cloned, mutagenized polypeptides expressed by host cells (Ness et al., 1999, *Nature Biotechnology* 17: 893-896). Mutagenized DNA molecules that encode active polypeptides can be recovered from the host cells and rapidly sequenced using standard methods in the art. These methods allow the rapid determination of the importance of individual amino acid residues in a polypeptide.

[0076] The polypeptide may be a hybrid polypeptide in which a region of one polypeptide is fused at the N-terminus or the C-terminus of a region of another polypeptide.

[0077] The polypeptide may be a fusion polypeptide or cleavable fusion polypeptide in which another polypeptide is fused at the N-terminus or the C-terminus of the polypeptide of the present invention. A fusion polypeptide is produced by fusing a polynucleotide encoding another polypeptide to a polynucleotide of the present invention. Techniques for producing fusion polypeptides are known in the art, and include ligating the coding sequences encoding the polypeptides so that they are in frame and that expression of the fusion polypeptide is under control of the same promoter(s) and terminator. Fusion polypeptides may also be constructed using intein technology in which fusion polypeptides are created post-translationally (Cooper et al., 1993, *EMB J.* 12: 2575-2583; Dawson et al., 1994, *Science* 266:

776-779).

[0078] A fusion polypeptide can further comprise a cleavage site between the two polypeptides. Upon secretion of the fusion protein, the site is cleaved releasing the two polypeptides. Examples of cleavage sites include, but are not limited to, the sites disclosed in Martin et al., 2003, J. Ind. Microbiol. Biotechnol. 3: 568-576; Svetina et al., 2000, J. Biotechnol. 76: 245-251; Rasmussen-Wilson et al., 1997, Appl. Environ. Microbiol. 63: 3488-3493; Ward et al., 1995, Biotechnology 13: 498-503; and Contreras et al., 1991, Biotechnology 9: 378-381; Eaton et al., 1986, Biochemistry 25: 505-512; Collins-Racie et al., 1995, Biotechnology 13: 982-987; Carter et al., 1989, Proteins: Structure, Function, and Genetics 6: 240-248; and Stevens, 2003, Drug Discovery World 4: 35-48.

Sources of Polypeptides Having Xylanase Activity

[0079] A polypeptide having xylanase activity of the present invention may be obtained from microorganisms of any genus. For purposes of the present invention, the term "obtained from" as used herein in connection with a given source shall mean that the polypeptide encoded by a polynucleotide is produced by the source or by a strain in which the polynucleotide from the source has been inserted. In one aspect, the polypeptide obtained from a given source is secreted extracellularly.

[0080] The polypeptide may be a *Talaromyces* polypeptide.

[0081] In another aspect, the polypeptide is a *Talaromyces leycettanus* polypeptide, e.g., a polypeptide obtained from *Talaromyces leycettanus* Strain CBS398.68.

[0082] It will be understood that for the aforementioned species, the invention encompasses both the perfect and imperfect states, and other taxonomic equivalents, e.g., anamorphs, regardless of the species name by which they are known. Those skilled in the art will readily recognize the identity of appropriate equivalents.

[0083] Strains of these species are readily accessible to the public in a number of culture collections, such as the American Type Culture Collection (ATCC), Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH (DSMZ), Centraalbureau Voor Schimmelcultures (CBS), and Agricultural Research Service Patent Culture Collection, Northern Regional Research Center (NRRL).

[0084] The polypeptide may be identified and obtained from other sources including microorganisms isolated from nature (e.g., soil, composts, water, etc.) or DNA samples obtained directly from natural materials (e.g., soil, composts, water, etc.) using the above-mentioned probes. Techniques for isolating microorganisms and DNA directly from natural habitats are well known in the art. A polynucleotide encoding the polypeptide may then be obtained by similarly screening a genomic DNA or cDNA library of another microorganism or

mixed DNA sample. Once a polynucleotide encoding a polypeptide has been detected with the probe(s), the polynucleotide can be isolated or cloned by utilizing techniques that are known to those of ordinary skill in the art (see, e.g., Sambrook *et al.*, 1989, *supra*).

Catalytic Domains

[0085] The present invention also relates to isolated polypeptides comprising a catalytic domain, as defined in the accompanying claims. A catalytic domain may have at least 77% sequence identity to the catalytic domain of SEQ ID NO: 2 (for example, amino acids 18 to 364 of SEQ ID NO: 2), or at least 77% sequence identity to the catalytic domain of SEQ ID NO: 4 (for example, amino acids 17 to 326 of SEQ ID NO: 4). In one aspect of the present invention, the catalytic domain has xylanase activity and at least 90% sequence identity to the catalytic domain of SEQ ID NO: 6 (for example, amino acids 21 to 337 of SEQ ID NO: 6). The catalytic domain may also be encoded by a polynucleotide having at least 60% sequence identity to the catalytic domain coding sequence of SEQ ID NO: 1 (for example, nucleotides 52-240, and 314-1165 of SEQ ID NO: 1), a catalytic domain encoded by a polynucleotide having at least 60% sequence identity to the catalytic domain coding sequence of SEQ ID NO: 3 (for example, nucleotides 49-241, 302-342, 404-452, 518-639, 707-852, 912-1019, 1088-1205, 1282-1347, and 1430-1516 of SEQ ID NO: 3), or a catalytic domain encoded by a polynucleotide having at least 60% sequence identity to the catalytic domain coding sequence of SEQ ID NO: 5 (for example, nucleotides 124-270, 342-474, 567-680, and 757-1313 of SEQ ID NO: 5). A variant of a catalytic domain may comprise a substitution, deletion, and/or insertion of one or more (several) amino acids of the catalytic domain of SEQ ID NO: 2, SEQ ID NO: 4, or SEQ ID NO: 6. Furthermore are described fragments of the above described catalytic domains which have xylanase activity.

[0086] The catalytic domain preferably has a degree of sequence identity to the catalytic domain of SEQ ID NO: 2, of at least 60%, e.g. at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100%. The catalytic domain preferably has a degree of sequence identity to the catalytic domain of SEQ ID NO: 4, of at least 60%, e.g. at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100%. The catalytic domain preferably has a degree of sequence identity to the catalytic domain of SEQ ID NO: 6, of at least 60%, e.g. at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100%. The catalytic domain may comprise an amino acid sequence that differs by ten amino acids, e.g., by five amino acids, by four amino acids, by three amino acids, by two amino acids, and by one amino acid from the catalytic domain of SEQ ID NO: 2, SEQ ID NO: 4, or SEQ ID NO: 6.

[0087] The catalytic domain preferably comprises or consists of the catalytic domain of SEQ ID NO: 2 or an allelic variant thereof; or is a fragment thereof having xylanase activity, or the catalytic domain comprises or consists of amino acids 18 to 364 of SEQ ID NO: 2.

[0088] The catalytic domain preferably comprises or consists of the catalytic domain of SEQ ID NO: 4 or an allelic variant thereof; or is a fragment thereof having xylanase activity, or the catalytic domain comprises or consists of amino acids 17 to 326 of SEQ ID NO: 4.

[0089] The catalytic domain preferably comprises or consists of the catalytic domain of SEQ ID NO: 6 or an allelic variant thereof; or is a fragment thereof having xylanase activity. In another preferred aspect of the present invention, the catalytic domain comprises or consists of amino acids 21 to 337 of SEQ ID NO: 6. The catalytic domain may be encoded by a polynucleotide that hybridizes under very low stringency conditions, or low stringency conditions, or medium stringency conditions, or medium-high stringency conditions, or high stringency conditions, or very high stringency conditions (as defined above) with (i) the catalytic domain coding sequence of SEQ ID NO: 1, SEQ ID NO: 3, or SEQ ID NO: 5, (ii) the cDNA sequence contained in the catalytic domain coding sequence of SEQ ID NO: 1, SEQ ID NO: 3, or SEQ ID NO: 5, or (iii) the full-length complementary strand of (i) or (ii) (J. Sambrook *et al.*, 1989, *supra*).

[0090] The catalytic domain may be encoded by a polynucleotide having a degree of sequence identity to the catalytic domain coding sequence of SEQ ID NO: 1 of at least 60%, e.g. at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100%, which encode a polypeptide having xylanase activity.

[0091] The catalytic domain may be encoded by a polynucleotide having a degree of sequence identity to the catalytic domain coding sequence of SEQ ID NO: 3 of at least 60%, e.g. at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100%, which encode a polypeptide having xylanase activity.

[0092] The catalytic domain may be encoded by a polynucleotide having a degree of sequence identity to the catalytic domain coding sequence of SEQ ID NO: 5 of at least 60%, e.g. at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100%, which encode a polypeptide having xylanase activity. The polynucleotide encoding the catalytic domain comprises or consists of nucleotides 52 to 1165 of SEQ ID NO: 1 or the cDNA sequence thereof. In particular the polynucleotide encoding the catalytic domain comprises or consists of nucleotides 52-240, and 314-1165 of SEQ ID NO: 1. The polynucleotide encoding the catalytic domain comprises or consists of nucleotides 49 to 1516 of SEQ ID NO: 3 or the cDNA sequence thereof. In particular the polynucleotide encoding the catalytic domain comprises or consists of nucleotides 49-241, 302-342, 404-452, 518-639, 707-852, 912-1019, 1088-1205, 1282-1347, and 1430-1516 of SEQ ID NO: 3. The polynucleotide encoding the catalytic domain comprises or consists of nucleotides 124 to 1313 of SEQ ID NO: 5 or the cDNA sequence thereof. In particular the polynucleotide encoding the catalytic domain comprises or consists of nucleotides 124-270, 342-474, 567-680, and 757-1313 of SEQ ID NO: 5.

Polynucleotides

[0093] The present invention also relates to isolated polynucleotides encoding a polypeptide of the present invention, as described herein.

[0094] The techniques used to isolate or clone a polynucleotide are known in the art and include isolation from genomic DNA or cDNA, or a combination thereof. The cloning of the polynucleotides from genomic DNA can be effected, *e.g.*, by using the well known polymerase chain reaction (PCR) or antibody screening of expression libraries to detect cloned DNA fragments with shared structural features. See, *e.g.*, Innis et al., 1990, PCR: A Guide to Methods and Application, Academic Press, New York. Other nucleic acid amplification procedures such as ligase chain reaction (LCR), ligation activated transcription (LAT) and polynucleotide-based amplification (NASBA) may be used. The polynucleotides may be cloned from a strain of *Talaromyces*, or a related organism and thus, for example, may be an allelic or species variant of the polypeptide encoding region of the polynucleotide.

[0095] Modification of a polynucleotide encoding a polypeptide of the present invention may be necessary for synthesizing polypeptides substantially similar to the polypeptide. The term "substantially similar" to the polypeptide refers to non-naturally occurring forms of the polypeptide. These polypeptides may differ in some engineered way from the polypeptide isolated from its native source, *e.g.*, variants that differ in specific activity, thermostability, pH optimum, or the like. The variants may be constructed on the basis of the polynucleotide presented as the mature polypeptide coding sequence of SEQ ID NO: 1, SEQ ID NO: 3, or SEQ ID NO: 5, or the cDNA sequence thereof, *e.g.*, a subsequence thereof, and/or by introduction of nucleotide substitutions that do not result in a change in the amino acid sequence of the polypeptide, but which correspond to the codon usage of the host organism intended for production of the enzyme, or by introduction of nucleotide substitutions that may give rise to a different amino acid sequence. For a general description of nucleotide substitutions, see, *e.g.*, Ford et al., 1991, Protein Expression and Purification 2: 95-107.

Nucleic Acid Constructs

[0096] The present invention also relates to nucleic acid constructs comprising a polynucleotide of the present invention operably linked to one or more control sequences that direct the expression of the coding sequence in a suitable host cell under conditions compatible with the control sequences.

[0097] A polynucleotide may be manipulated in a variety of ways to provide for expression of the polypeptide. Manipulation of the polynucleotide prior to its insertion into a vector may be desirable or necessary depending on the expression vector. The techniques for modifying polynucleotides utilizing recombinant DNA methods are well known in the art.

[0098] The control sequence may be a promoter, a polynucleotide that is recognized by a host cell for expression of a polynucleotide encoding a polypeptide of the present invention. The promoter contains transcriptional control sequences that mediate the expression of the polypeptide. The promoter may be any polynucleotide that shows transcriptional activity in the host cell including mutant, truncated, and hybrid promoters, and may be obtained from genes encoding extracellular or intracellular polypeptides either homologous or heterologous to the host cell.

[0099] Examples of suitable promoters for directing transcription of the nucleic acid constructs of the present invention in a bacterial host cell are the promoters obtained from the *Bacillus amyloliquefaciens* alpha-amylase gene (*amyQ*), *Bacillus licheniformis* alpha-amylase gene (*amyL*), *Bacillus licheniformis* penicillinase gene (*penP*), *Bacillus stearothermophilus* maltogenic amylase gene (*amyM*), *Bacillus subtilis* levansucrase gene (*sacB*), *Bacillus subtilis* *xylA* and *xylB* genes, *Bacillus thuringiensis* *cryIIIA* gene (Agaisse and Lereclus, 1994, Molecular Microbiology 13: 97-107), *E. coli* *lac* operon, *E. coli* *trc* promoter (Egon et al., 1988, Gene 69: 301-315), *Streptomyces coelicolor* agarase gene (*dagA*), and prokaryotic beta-lactamase gene (Villa-Kamaroff et al., 1978, Proc. Natl. Acad. Sci. USA 75: 3727-3731), as well as the *tac* promoter (DeBoer et al., 1983, Proc. Natl. Acad. Sci. USA 80: 21-25). Further promoters are described in "Useful proteins from recombinant bacteria" in Gilbert et al., 1980, Scientific American 242: 74-94; and in Sambrook et al., 1989, *supra*. Examples of tandem promoters are disclosed in WO 99/43835.

[0100] Examples of suitable promoters for directing transcription of the nucleic acid constructs of the present invention in a filamentous fungal host cell are promoters obtained from the genes for *Aspergillus nidulans* acetamidase, *Aspergillus niger* neutral alpha-amylase, *Aspergillus niger* acid stable alpha-amylase, *Aspergillus niger* or *Aspergillus awamori* glucoamylase (*glaA*), *Aspergillus oryzae* TAKA amylase, *Aspergillus oryzae* alkaline protease, *Aspergillus oryzae* triose phosphate isomerase, *Fusarium oxysporum* trypsin-like protease (WO 96/00787), *Fusarium venenatum* amyloglucosidase (WO 00/56900), *Fusarium venenatum* Daria (WO 00/56900), *Fusarium venenatum* Quinn (WO 00/56900), *Rhizomucor miehei* lipase, *Rhizomucor miehei* aspartic proteinase, *Trichoderma reesei* beta-glucosidase, *Trichoderma reesei* cellobiohydrolase I, *Trichoderma reesei* cellobiohydrolase II, *Trichoderma reesei* endoglucanase I, *Trichoderma reesei* endoglucanase II, *Trichoderma reesei* endoglucanase III, *Trichoderma reesei* endoglucanase IV, *Trichoderma reesei* endoglucanase V, *Trichoderma reesei* xylanase I, *Trichoderma reesei* xylanase II, *Trichoderma reesei* beta-xylosidase, as well as the NA2-tpi promoter (a modified promoter from an *Aspergillus* neutral alpha-amylase gene in which the untranslated leader has been replaced by an untranslated leader from an *Aspergillus* triose phosphate isomerase gene; non-limiting examples include modified promoters from an *Aspergillus niger* neutral alpha-amylase gene in which the untranslated leader has been replaced by an untranslated leader from an *Aspergillus nidulans* or *Aspergillus oryzae* triose phosphate isomerase gene); and mutant, truncated, and hybrid promoters thereof.

[0101] In a yeast host, useful promoters are obtained from the genes for *Saccharomyces cerevisiae* enolase (ENO-1), *Saccharomyces cerevisiae* galactokinase (GAL1), *Saccharomyces cerevisiae* alcohol dehydrogenase/glyceraldehyde-3-phosphate dehydrogenase (ADH1, ADH2/GAP), *Saccharomyces cerevisiae* triose phosphate isomerase (TPI), *Saccharomyces cerevisiae* metallothionein (CUP1), and *Saccharomyces cerevisiae* 3-phosphoglycerate kinase. Other useful promoters for yeast host cells are described by Romanos et al., 1992, *Yeast* 8: 423-488.

[0102] The control sequence may also be a transcription terminator, which is recognized by a host cell to terminate transcription. The terminator is operably linked to the 3'-terminus of the polynucleotide encoding the polypeptide. Any terminator that is functional in the host cell may be used in the present invention.

[0103] Preferred terminators for bacterial host cells are obtained from the genes for *Bacillus clausii* alkaline protease (*aprH*), *Bacillus licheniformis* alpha-amylase (*amyL*), and *Escherichia coli* ribosomal RNA (*rrnB*).

[0104] Preferred terminators for filamentous fungal host cells are obtained from the genes for *Aspergillus nidulans* anthranilate synthase, *Aspergillus niger* glucoamylase, *Aspergillus niger* alpha-glucosidase, *Aspergillus oryzae* TAKA amylase, and *Fusarium oxysporum* trypsin-like protease.

[0105] Preferred terminators for yeast host cells are obtained from the genes for *Saccharomyces cerevisiae* enolase, *Saccharomyces cerevisiae* cytochrome C (CYC1), and *Saccharomyces cerevisiae* glyceraldehyde-3-phosphate dehydrogenase. Other useful terminators for yeast host cells are described by Romanos *et al.*, 1992, *supra*.

[0106] The control sequence may also be an mRNA stabilizer region downstream of a promoter and upstream of the coding sequence of a gene which increases expression of the gene.

[0107] Examples of suitable mRNA stabilizer regions are obtained from a *Bacillus thuringiensis cryIIIA* gene (WO 94/25612) and a *Bacillus subtilis* SP82 gene (Hue et al., 1995, *Journal of Bacteriology* 177: 3465-3471).

[0108] The control sequence may also be a leader, a nontranslated region of an mRNA that is important for translation by the host cell. The leader is operably linked to the 5'-terminus of the polynucleotide encoding the polypeptide. Any leader that is functional in the host cell may be used.

[0109] Preferred leaders for filamentous fungal host cells are obtained from the genes for *Aspergillus oryzae* TAKA amylase and *Aspergillus nidulans* triose phosphate isomerase.

[0110] Suitable leaders for yeast host cells are obtained from the genes for *Saccharomyces*

cerevisiae enolase (ENO-1), *Saccharomyces cerevisiae* 3-phosphoglycerate kinase, *Saccharomyces cerevisiae* alpha-factor, and *Saccharomyces cerevisiae* alcohol dehydrogenase/glyceraldehyde-3-phosphate dehydrogenase (ADH2/GAP).

[0111] The control sequence may also be a polyadenylation sequence, a sequence operably linked to the 3'-terminus of the polynucleotide and, when transcribed, is recognized by the host cell as a signal to add polyadenosine residues to transcribed mRNA. Any polyadenylation sequence that is functional in the host cell may be used.

[0112] Preferred polyadenylation sequences for filamentous fungal host cells are obtained from the genes for *Aspergillus nidulans* anthranilate synthase, *Aspergillus niger* glucoamylase, *Aspergillus niger* alpha-glucosidase *Aspergillus oryzae* TAKA amylase, and *Fusarium oxysporum* trypsin-like protease.

[0113] Useful polyadenylation sequences for yeast host cells are described by Guo and Sherman, 1995, Mol. Cellular Biol. 15: 5983-5990.

[0114] The control sequence may also be a signal peptide coding region that encodes a signal peptide linked to the N-terminus of a polypeptide and directs the polypeptide into the cell's secretory pathway. The 5'-end of the coding sequence of the polynucleotide may inherently contain a signal peptide coding sequence naturally linked in translation reading frame with the segment of the coding sequence that encodes the polypeptide. Alternatively, the 5'-end of the coding sequence may contain a signal peptide coding sequence that is foreign to the coding sequence. A foreign signal peptide coding sequence may be required where the coding sequence does not naturally contain a signal peptide coding sequence. Alternatively, a foreign signal peptide coding sequence may simply replace the natural signal peptide coding sequence in order to enhance secretion of the polypeptide. However, any signal peptide coding sequence that directs the expressed polypeptide into the secretory pathway of a host cell may be used.

[0115] Effective signal peptide coding sequences for bacterial host cells are the signal peptide coding sequences obtained from the genes for *Bacillus* NCIB 11837 maltogenic amylase, *Bacillus licheniformis* subtilisin, *Bacillus licheniformis* beta-lactamase, *Bacillus stearothermophilus* alpha-amylase, *Bacillus stearothermophilus* neutral proteases (*nprT*, *nprS*, *nprM*), and *Bacillus subtilis* *prsA*. Further signal peptides are described by Simonen and Palva, 1993, Microbiological Reviews 57: 109-137.

[0116] Effective signal peptide coding sequences for filamentous fungal host cells are the signal peptide coding sequences obtained from the genes for *Aspergillus niger* neutral amylase, *Aspergillus niger* glucoamylase, *Aspergillus oryzae* TAKA amylase, *Humicola insolens* cellulase, *Humicola insolens* endoglucanase V, *Humicola lanuginosa* lipase, and *Rhizomucor miehei* aspartic proteinase.

[0117] Useful signal peptides for yeast host cells are obtained from the genes for

Saccharomyces cerevisiae alpha-factor and *Saccharomyces cerevisiae* invertase. Other useful signal peptide coding sequences are described by Romanos *et al.*, 1992, *supra*.

[0118] The control sequence may also be a propeptide coding sequence that encodes a propeptide positioned at the N-terminus of a polypeptide. The resultant polypeptide is known as a proenzyme or propolypeptide (or a zymogen in some cases). A propolypeptide is generally inactive and can be converted to an active polypeptide by catalytic or autocatalytic cleavage of the propeptide from the propolypeptide. The propeptide coding sequence may be obtained from the genes for *Bacillus subtilis* alkaline protease (*aprE*), *Bacillus subtilis* neutral protease (*nprT*), *Myceliophthora thermophila* laccase (WO 95/33836), *Rhizomucor miehei* aspartic proteinase, and *Saccharomyces cerevisiae* alpha-factor.

[0119] Where both signal peptide and propeptide sequences are present, the propeptide sequence is positioned next to the N-terminus of a polypeptide and the signal peptide sequence is positioned next to the N-terminus of the propeptide sequence.

[0120] It may also be desirable to add regulatory sequences that regulate expression of the polypeptide relative to the growth of the host cell. Examples of regulatory systems are those that cause expression of the gene to be turned on or off in response to a chemical or physical stimulus, including the presence of a regulatory compound. Regulatory systems in prokaryotic systems include the *lac*, *tac*, and *trp* operator systems. In yeast, the ADH2 system or GAL1 system may be used. In filamentous fungi, the *Aspergillus niger* glucoamylase promoter, *Aspergillus oryzae* TKA alpha-amylase promoter, and *Aspergillus oryzae* glucoamylase promoter may be used. Other examples of regulatory sequences are those that allow for gene amplification. In eukaryotic systems, these regulatory sequences include the dihydrofolate reductase gene that is amplified in the presence of methotrexate, and the metallothionein genes that are amplified with heavy metals. In these cases, the polynucleotide encoding the polypeptide would be operably linked with the regulatory sequence.

Expression Vectors

[0121] The present invention also relates to recombinant expression vectors comprising a polynucleotide of the present invention, a promoter, and transcriptional and translational stop signals. The various nucleotide and control sequences may be joined together to produce a recombinant expression vector that may include one or more convenient restriction sites to allow for insertion or substitution of the polynucleotide encoding the polypeptide at such sites. Alternatively, the polynucleotide may be expressed by inserting the polynucleotide or a nucleic acid construct comprising the polynucleotide into an appropriate vector for expression. In creating the expression vector, the coding sequence is located in the vector so that the coding sequence is operably linked with the appropriate control sequences for expression.

[0122] The recombinant expression vector may be any vector (e.g., a plasmid or virus) that can be conveniently subjected to recombinant DNA procedures and can bring about

expression of the polynucleotide. The choice of the vector will typically depend on the compatibility of the vector with the host cell into which the vector is to be introduced. The vector may be a linear or closed circular plasmid.

[0123] The vector may be an autonomously replicating vector, *i.e.*, a vector that exists as an extrachromosomal entity, the replication of which is independent of chromosomal replication, *e.g.*, a plasmid, an extrachromosomal element, a minichromosome, or an artificial chromosome. The vector may contain any means for assuring self-replication. Alternatively, the vector may be one that, when introduced into the host cell, is integrated into the genome and replicated together with the chromosome(s) into which it has been integrated. Furthermore, a single vector or plasmid or two or more vectors or plasmids that together contain the total DNA to be introduced into the genome of the host cell, or a transposon, may be used.

[0124] The vector preferably contains one or more selectable markers that permit easy selection of transformed, transfected, transduced, or the like cells. A selectable marker is a gene the product of which provides for biocide or viral resistance, resistance to heavy metals, prototrophy to auxotrophs, and the like.

[0125] Examples of bacterial selectable markers are *Bacillus licheniformis* or *Bacillus subtilis* *dal* genes, or markers that confer antibiotic resistance such as ampicillin, chloramphenicol, kanamycin, neomycin, spectinomycin, or tetracycline resistance. Suitable markers for yeast host cells include, but are not limited to, ADE2, HIS3, LEU2, LYS2, MET3, TRP1, and URA3. Selectable markers for use in a filamentous fungal host cell include, but are not limited to, *amdS* (acetamidase), *argB* (ornithine carbamoyltransferase), *bar* (phosphinothricin acetyltransferase), *hph* (hygromycin phosphotransferase), *niaD* (nitrate reductase), *pyrG* (orotidine-5'-phosphate decarboxylase), *sC* (sulfate adenylyltransferase), and *trpC* (anthranilate synthase), as well as equivalents thereof. Preferred for use in an *Aspergillus* cell are *Aspergillus nidulans* or *Aspergillus oryzae* *amdS* and *pyrG* genes and a *Streptomyces hygroscopicus* *bar* gene.

[0126] The vector preferably contains an element(s) that permits integration of the vector into the host cell's genome or autonomous replication of the vector in the cell independent of the genome.

[0127] For integration into the host cell genome, the vector may rely on the polynucleotide's sequence encoding the polypeptide or any other element of the vector for integration into the genome by homologous or non-homologous recombination. Alternatively, the vector may contain additional polynucleotides for directing integration by homologous recombination into the genome of the host cell at a precise location(s) in the chromosome(s). To increase the likelihood of integration at a precise location, the integrational elements should contain a sufficient number of nucleic acids, such as 100 to 10,000 base pairs, 400 to 10,000 base pairs, and 800 to 10,000 base pairs, which have a high degree of sequence identity to the corresponding target sequence to enhance the probability of homologous recombination. The integrational elements may be any sequence that is homologous with the target sequence in

the genome of the host cell. Furthermore, the integrational elements may be non-encoding or encoding polynucleotides. On the other hand, the vector may be integrated into the genome of the host cell by non-homologous recombination.

[0128] For autonomous replication, the vector may further comprise an origin of replication enabling the vector to replicate autonomously in the host cell in question. The origin of replication may be any plasmid replicator mediating autonomous replication that functions in a cell. The term "origin of replication" or "plasmid replicator" means a polynucleotide that enables a plasmid or vector to replicate *in vivo*.

[0129] Examples of bacterial origins of replication are the origins of replication of plasmids pBR322, pUC19, pACYC177, and pACYC184 permitting replication in *E. coli*, and pUB110, pE194, pTA1060, and pAM β 1 permitting replication in *Bacillus*.

[0130] Examples of origins of replication for use in a yeast host cell are the 2 micron origin of replication, ARS1, ARS4, the combination of ARS1 and CEN3, and the combination of ARS4 and CEN6.

[0131] Examples of origins of replication useful in a filamentous fungal cell are AMA1 and ANS1 (Gems et al., 1991, Gene 98: 61-67; Cullen et al., 1987, Nucleic Acids Res. 15: 9163-9175; WO 00/24883). Isolation of the AMA1 gene and construction of plasmids or vectors comprising the gene can be accomplished according to the methods disclosed in WO 00/24883.

[0132] More than one copy of a polynucleotide of the present invention may be inserted into a host cell to increase production of a polypeptide. An increase in the copy number of the polynucleotide can be obtained by integrating at least one additional copy of the sequence into the host cell genome or by including an amplifiable selectable marker gene with the polynucleotide where cells containing amplified copies of the selectable marker gene, and thereby additional copies of the polynucleotide, can be selected for by cultivating the cells in the presence of the appropriate selectable agent.

[0133] The procedures used to ligate the elements described above to construct the recombinant expression vectors of the present invention are well known to one skilled in the art (see, e.g., Sambrook et al., 1989, *supra*).

Host Cells

[0134] The present invention also relates to recombinant host cells, comprising a polynucleotide of the present invention operably linked to one or more control sequences that direct the production of a polypeptide of the present invention. A construct or vector comprising a polynucleotide is introduced into a host cell so that the construct or vector is maintained as a chromosomal integrant or as a self-replicating extra-chromosomal vector as described earlier.

The term "host cell" encompasses any progeny of a parent cell that is not identical to the parent cell due to mutations that occur during replication. The choice of a host cell will to a large extent depend upon the gene encoding the polypeptide and its source.

[0135] The host cell may be any cell useful in the recombinant production of a polypeptide of the present invention, e.g., a prokaryote or a eukaryote.

[0136] The prokaryotic host cell may be any Gram-positive or Gram-negative bacterium. Gram-positive bacteria include, but are not limited to, *Bacillus*, *Clostridium*, *Enterococcus*, *Geobacillus*, *Lactobacillus*, *Lactococcus*, *Oceanobacillus*, *Staphylococcus*, *Streptococcus*, and *Streptomyces*. Gram-negative bacteria include, but are not limited to, *Campylobacter*, *E. coli*, *Flavobacterium*, *Fusobacterium*, *Helicobacter*, *Ilyobacter*, *Neisseria*, *Pseudomonas*, *Salmonella*, and *Ureaplasma*.

[0137] The bacterial host cell may be any *Bacillus* cell including, but not limited to, *Bacillus alkalophilus*, *Bacillus amyloliquefaciens*, *Bacillus brevis*, *Bacillus circulans*, *Bacillus clausii*, *Bacillus coagulans*, *Bacillus firmus*, *Bacillus lautus*, *Bacillus lentus*, *Bacillus licheniformis*, *Bacillus megaterium*, *Bacillus pumilus*, *Bacillus stearothermophilus*, *Bacillus subtilis*, and *Bacillus thuringiensis* cells.

[0138] The bacterial host cell may also be any *Streptococcus* cell including, but not limited to, *Streptococcus equisimilis*, *Streptococcus pyogenes*, *Streptococcus uberis*, and *Streptococcus equi subsp. Zooepidemicus* cells.

[0139] The bacterial host cell may also be any *Streptomyces* cell including, but not limited to, *Streptomyces achromogenes*, *Streptomyces avermitilis*, *Streptomyces coelicolor*, *Streptomyces griseus*, and *Streptomyces lividans* cells.

[0140] The introduction of DNA into a *Bacillus* cell may be effected by protoplast transformation (see, e.g., Chang and Cohen, 1979, Mol. Gen. Genet. 168: 111-115), competent cell transformation (see, e.g., Young and Spizizen, 1961, J. Bacteriol. 81: 823-829, or Dubnau and Davidoff-Abelson, 1971, J. Mol. Biol. 56: 209-221), electroporation (see, e.g., Shigekawa and Dower, 1988, Biotechniques 6: 742-751), or conjugation (see, e.g., Koehler and Thorne, 1987, J. Bacteriol. 169: 5271-5278). The introduction of DNA into an *E. coli* cell may be effected by protoplast transformation (see, e.g., Hanahan, 1983, J. Mol. Biol. 166: 557-580) or electroporation (see, e.g., Dower et al., 1988, Nucleic Acids Res. 16: 6127-6145). The introduction of DNA into a *Streptomyces* cell may be effected by protoplast transformation, electroporation (see, e.g., Gong et al., 2004, Folia Microbiol. (Praha) 49: 399-405), conjugation (see, e.g., Mazodier et al., 1989, J. Bacteriol. 171: 3583-3585), or transduction (see, e.g., Burke et al., 2001, Proc. Natl. Acad. Sci. USA 98: 6289-6294). The introduction of DNA into a *Pseudomonas* cell may be effected by electroporation (see, e.g., Choi et al., 2006, J. Microbiol. Methods 64: 391-397) or conjugation (see, e.g., Pinedo and Smets, 2005, Appl. Environ. Microbiol. 71: 51-57). The introduction of DNA into a *Streptococcus* cell may be effected by natural competence (see, e.g., Perry and Kuramitsu, 1981, Infect. Immun. 32: 1295-1297),

protoplast transformation (see, e.g., Catt and Jollick, 1991, *Microbios* 68: 189-207), electroporation (see, e.g., Buckley et al., 1999, *Appl. Environ. Microbiol.* 65: 3800-3804), or conjugation (see, e.g., Clewell, 1981, *Microbiol. Rev.* 45: 409-436). However, any method known in the art for introducing DNA into a host cell can be used.

[0141] The host cell may also be a eukaryote, such as a mammalian, insect, plant, or fungal cell.

[0142] The host cell may be a fungal cell. "Fungi" as used herein includes the phyla Ascomycota, Basidiomycota, Chytridiomycota, and Zygomycota as well as the Oomycota and all mitosporic fungi (as defined by Hawksworth et al., In, Ainsworth and Bisby's Dictionary of The Fungi, 8th edition, 1995, CAB International, University Press, Cambridge, UK).

[0143] The fungal host cell may be a yeast cell. "Yeast" as used herein includes ascosporogenous yeast (Endomycetales), basidiosporogenous yeast, and yeast belonging to the Fungi Imperfecti (Blastomycetes). Since the classification of yeast may change in the future, for the purposes of this invention, yeast shall be defined as described in Biology and Activities of Yeast (Skinner, Passmore, and Davenport, editors, Soc. App. Bacteriol. Symposium Series No. 9, 1980).

[0144] The yeast host cell may be a *Candida*, *Hansenula*, *Kluyveromyces*, *Pichia*, *Saccharomyces*, *Schizosaccharomyces*, or *Yarrowia* cell, such as a *Kluyveromyces lactis*, *Saccharomyces carlsbergensis*, *Saccharomyces cerevisiae*, *Saccharomyces diastaticus*, *Saccharomyces douglasii*, *Saccharomyces kluyveri*, *Saccharomyces norbensis*, *Saccharomyces oviformis*, or *Yarrowia lipolytica* cell.

[0145] The fungal host cell may be a filamentous fungal cell. "Filamentous fungi" include all filamentous forms of the subdivision Eumycota and Oomycota (as defined by Hawksworth *et al.*, 1995, *supra*). The filamentous fungi are generally characterized by a mycelial wall composed of chitin, cellulose, glucan, chitosan, mannan, and other complex polysaccharides. Vegetative growth is by hyphal elongation and carbon catabolism is obligately aerobic. In contrast, vegetative growth by yeasts such as *Saccharomyces cerevisiae* is by budding of a unicellular thallus and carbon catabolism may be fermentative.

[0146] The filamentous fungal host cell may be an *Acremonium*, *Aspergillus*, *Aureobasidium*, *Bjerkandera*, *Ceriporiopsis*, *Chrysosporium*, *Coprinus*, *Coriolus*, *Cryptococcus*, *Filibasidium*, *Fusarium*, *Humicola*, *Magnaporthe*, *Mucor*, *Myceliophthora*, *Neocallimastix*, *Neurospora*, *Paecilomyces*, *Penicillium*, *Phanerochaete*, *Phlebia*, *Piromyces*, *Pleurotus*, *Schizophyllum*, *Talaromyces*, *Thermoascus*, *Thielavia*, *Tolypocladium*, *Trametes*, or *Trichoderma* cell.

[0147] For example, the filamentous fungal host cell may be an *Aspergillus awamori*, *Aspergillus foetidus*, *Aspergillus fumigatus*, *Aspergillus japonicus*, *Aspergillus nidulans*, *Aspergillus niger*, *Aspergillus oryzae*, *Bjerkandera adusta*, *Ceriporiopsis aneirina*, *Ceriporiopsis caregiea*, *Ceriporiopsis gilvescens*, *Ceriporiopsis pannocinta*, *Ceriporiopsis rivulosa*,

Ceriporiopsis subrufa, *Ceriporiopsis subvermispora*, *Chrysosporium inops*, *Chrysosporium keratinophilum*, *Chrysosporium lucknowense*, *Chrysosporium merdarium*, *Chrysosporium pannicola*, *Chrysosporium queenslandicum*, *Chrysosporium tropicum*, *Chrysosporium zonatum*, *Coprinus cinereus*, *Coriolus hirsutus*, *Fusarium bactridioides*, *Fusarium cerealis*, *Fusarium crookwellense*, *Fusarium culmorum*, *Fusarium graminearum*, *Fusarium graminum*, *Fusarium heterosporum*, *Fusarium negundi*, *Fusarium oxysporum*, *Fusarium reticulatum*, *Fusarium roseum*, *Fusarium sambucinum*, *Fusarium sarcochroum*, *Fusarium sporotrichioides*, *Fusarium sulphureum*, *Fusarium torulosum*, *Fusarium trichothecioides*, *Fusarium venenatum*, *Humicola insolens*, *Humicola lanuginosa*, *Mucor miehei*, *Myceliophthora thermophila*, *Neurospora crassa*, *Penicillium purpurogenum*, *Phanerochaete chrysosporium*, *Phlebia radiata*, *Pleurotus eryngii*, *Thielavia terrestris*, *Trametes villosa*, *Trametes versicolor*, *Trichoderma harzianum*, *Trichoderma koningii*, *Trichoderma longibrachiatum*, *Trichoderma reesei*, or *Trichoderma viride* cell.

[0148] Fungal cells may be transformed by a process involving protoplast formation, transformation of the protoplasts, and regeneration of the cell wall in a manner known *per se*. Suitable procedures for transformation of *Aspergillus* and *Trichoderma* host cells are described in EP 238023, Yelton et al., 1984, Proc. Natl. Acad. Sci. USA 81: 1470-1474, and Christensen et al., 1988, Bio/Technology 6: 1419-1422. Suitable methods for transforming *Fusarium* species are described by Malardier et al., 1989, Gene 78: 147-156, and WO 96/00787. Yeast may be transformed using the procedures described by Becker and Guarente, In Abelson, J.N. and Simon, M.I., editors, Guide to Yeast Genetics and Molecular Biology, Methods in Enzymology, Volume 194, pp 182-187, Academic Press, Inc., New York; Ito et al., 1983, J. Bacteriol. 153: 163; and Hinnen et al., 1978, Proc. Natl. Acad. Sci. USA 75: 1920.

Methods of Production

[0149] The present invention also relates to methods of producing a polypeptide of the present invention, comprising (a) cultivating a cell, which in its wild-type form produces the polypeptide, under conditions conducive for production of the polypeptide; and (b) recovering the polypeptide. In a preferred aspect, the cell is a *Talaromyces* cell. In a more preferred aspect, the cell is a *Talaromyces leycettanus* cell. In a most preferred aspect, the cell is *Talaromyces leycettanus* Strain CBS398.68.

[0150] The present invention also relates to methods of producing a polypeptide of the present invention, comprising (a) cultivating a recombinant host cell of the present invention under conditions conducive for production of the polypeptide; and (b) recovering the polypeptide.

[0151] The host cells are cultivated in a nutrient medium suitable for production of the polypeptide using methods known in the art. For example, the cell may be cultivated by shake flask cultivation, or small-scale or large-scale fermentation (including continuous, batch, fed-batch, or solid state fermentations) in laboratory or industrial fermentors in a suitable medium and under conditions allowing the polypeptide to be expressed and/or isolated. The cultivation

takes place in a suitable nutrient medium comprising carbon and nitrogen sources and inorganic salts, using procedures known in the art. Suitable media are available from commercial suppliers or may be prepared according to published compositions (e.g., in catalogues of the American Type Culture Collection). If the polypeptide is secreted into the nutrient medium, the polypeptide can be recovered directly from the medium. If the polypeptide is not secreted, it can be recovered from cell lysates.

[0152] The polypeptide may be detected using methods known in the art that are specific for the polypeptides. These detection methods include, but are not limited to, use of specific antibodies, formation of an enzyme product, or disappearance of an enzyme substrate. For example, an enzyme assay may be used to determine the activity of the polypeptide.

[0153] The polypeptide may be recovered using methods known in the art. For example, the polypeptide may be recovered from the nutrient medium by conventional procedures including, but not limited to, collection, centrifugation, filtration, extraction, spray-drying, evaporation, or precipitation.

[0154] The polypeptide may be purified by a variety of procedures known in the art including, but not limited to, chromatography (e.g., ion exchange, affinity, hydrophobic, chromatofocusing, and size exclusion), electrophoretic procedures (e.g., preparative isoelectric focusing), differential solubility (e.g., ammonium sulfate precipitation), SDS-PAGE, or extraction (see, e.g., Protein Purification, Janson and Ryden, editors, VCH Publishers, New York, 1989) to obtain substantially pure polypeptides.

[0155] In an alternative aspect, the polypeptide is not recovered, but rather a host cell of the present invention expressing the polypeptide is used as a source of the polypeptide.

Plants

[0156] The present invention also relates to isolated plants, e.g., a transgenic plant, plant part, or plant cell, comprising a polynucleotide of the present invention so as to express and produce a polypeptide or domain in recoverable quantities. The polypeptide or domain may be recovered from the plant or plant part. Alternatively, the plant or plant part containing the polypeptide or domain may be used as such for improving the quality of a food or feed, e.g., improving nutritional value, palatability, and rheological properties, or to destroy an antinutritive factor.

[0157] The transgenic plant can be dicotyledonous (a dicot) or monocotyledonous (a monocot). Examples of monocot plants are grasses, such as meadow grass (blue grass, *Poa*), forage grass such as *Festuca*, *Lolium*, temperate grass, such as *Agrostis*, and cereals, e.g., wheat, oats, rye, barley, rice, sorghum, and maize (corn).

[0158] Examples of dicot plants are tobacco, legumes, such as lupins, potato, sugar beet, pea,

bean and soybean, and cruciferous plants (family Brassicaceae), such as cauliflower, rape seed, and the closely related model organism *Arabidopsis thaliana*.

[0159] Examples of plant parts are stem, callus, leaves, root, fruits, seeds, and tubers as well as the individual tissues comprising these parts, e.g., epidermis, mesophyll, parenchyme, vascular tissues, meristems. Specific plant cell compartments, such as chloroplasts, apoplasts, mitochondria, vacuoles, peroxisomes and cytoplasm are also considered to be a plant part. Furthermore, any plant cell, whatever the tissue origin, is considered to be a plant part. Likewise, plant parts such as specific tissues and cells isolated to facilitate the utilization of the invention are also considered plant parts, e.g., embryos, endosperms, aleurone and seed coats.

[0160] Also included within the scope of the present invention are the progeny of such plants, plant parts, and plant cells.

[0161] The transgenic plant or plant cell expressing the polypeptide or domain may be constructed in accordance with methods known in the art. In short, the plant or plant cell is constructed by incorporating one or more expression constructs encoding the polypeptide or domain into the plant host genome or chloroplast genome and propagating the resulting modified plant or plant cell into a transgenic plant or plant cell.

[0162] The expression construct is conveniently a nucleic acid construct that comprises a polynucleotide encoding a polypeptide or domain operably linked with appropriate regulatory sequences required for expression of the polynucleotide in the plant or plant part of choice. Furthermore, the expression construct may comprise a selectable marker useful for identifying plant cells into which the expression construct has been integrated and DNA sequences necessary for introduction of the construct into the plant in question (the latter depends on the DNA introduction method to be used).

[0163] The choice of regulatory sequences, such as promoter and terminator sequences and optionally signal or transit sequences, is determined, for example, on the basis of when, where, and how the polypeptide or domain is desired to be expressed. For instance, the expression of the gene encoding a polypeptide or domain may be constitutive or inducible, or may be developmental, stage or tissue specific, and the gene product may be targeted to a specific tissue or plant part such as seeds or leaves. Regulatory sequences are, for example, described by Tague et al., 1988, Plant Physiology 86: 506.

[0164] For constitutive expression, the 35S-CaMV, the maize ubiquitin 1, or the rice actin 1 promoter may be used (Franck et al., 1980, Cell 21: 285-294; Christensen et al., 1992, Plant Mol. Biol. 18: 675-689; Zhang et al., 1991, Plant Cell 3: 1155-1165). Organ-specific promoters may be, for example, a promoter from storage sink tissues such as seeds, potato tubers, and fruits (Edwards and Coruzzi, 1990, Ann. Rev. Genet. 24: 275-303), or from metabolic sink tissues such as meristems (Ito et al., 1994, Plant Mol. Biol. 24: 863-878), a seed specific promoter such as the glutelin, prolamin, globulin, or albumin promoter from rice (Wu et al.,

1998, Plant Cell Physiol. 39: 885-889), a *Vicia faba* promoter from the legumin B4 and the unknown seed protein gene from *Vicia faba* (Conrad et al., 1998, J. Plant Physiol. 152: 708-711), a promoter from a seed oil body protein (Chen et al., 1998, Plant Cell Physiol. 39: 935-941), the storage protein *napA* promoter from *Brassica napus*, or any other seed specific promoter known in the art, e.g., as described in WO 91/14772. Furthermore, the promoter may be a leaf specific promoter such as the *rbcS* promoter from rice or tomato (Kyoizuka et al., 1993, Plant Physiol. 102: 991-1000), the chlorella virus adenine methyltransferase gene promoter (Mitra and Higgins, 1994, Plant Mol. Biol. 26: 85-93), the *aldP* gene promoter from rice (Kagaya et al., 1995, Mol. Gen. Genet. 248: 668-674), or a wound inducible promoter such as the potato *pin2* promoter (Xu et al., 1993, Plant Mol. Biol. 22: 573-588). Likewise, the promoter may be induced by abiotic treatments such as temperature, drought, or alterations in salinity or induced by exogenously applied substances that activate the promoter, e.g., ethanol, oestrogens, plant hormones such as ethylene, abscisic acid, and gibberellic acid, and heavy metals.

[0165] A promoter enhancer element may also be used to achieve higher expression of a polypeptide or domain in the plant. For instance, the promoter enhancer element may be an intron that is placed between the promoter and the polynucleotide encoding a polypeptide or domain. For instance, Xu *et al.*, 1993, *supra*, disclose the use of the first intron of the rice actin 1 gene to enhance expression.

[0166] The selectable marker gene and any other parts of the expression construct may be chosen from those available in the art.

[0167] The nucleic acid construct is incorporated into the plant genome according to conventional techniques known in the art, including *Agrobacterium*-mediated transformation, virus-mediated transformation, microinjection, particle bombardment, biolistic transformation, and electroporation (Gasser et al., 1990, Science 244: 1293; Potrykus, 1990, Bio/Technology 8: 535; Shimamoto et al., 1989, Nature 338: 274).

[0168] *Agrobacterium tumefaciens*-mediated gene transfer is a method for generating transgenic dicots (for a review, see Hooykas and Schilperoort, 1992, Plant Mol. Biol. 19: 15-38) and for transforming monocots, although other transformation methods may be used for these plants. A method for generating transgenic monocots is particle bombardment (microscopic gold or tungsten particles coated with the transforming DNA) of embryonic calli or developing embryos (Christou, 1992, Plant J. 2: 275-281; Shimamoto, 1994, Curr. Opin. Biotechnol. 5: 158-162; Vasil et al., 1992, Bio/Technology 10: 667-674). An alternative method for transformation of monocots is based on protoplast transformation as described by Omirulleh et al., 1993, Plant Mol. Biol. 21: 415-428. Additional transformation methods include those described in U.S. Patent Nos. 6,395,966 and 7,151,204. Following transformation, the transformants having incorporated the expression construct are selected and regenerated into whole plants according to methods well known in the art. Often the transformation procedure is designed for the selective elimination of selection genes either during regeneration or in the following generations by using, for example, co-transformation with two separate T-DNA

constructs or site specific excision of the selection gene by a specific recombinase.

[0169] In addition to direct transformation of a particular plant genotype with a construct of the present invention, transgenic plants may be made by crossing a plant having the construct to a second plant lacking the construct. For example, a construct encoding a polypeptide or domain can be introduced into a particular plant variety by crossing, without the need for ever directly transforming a plant of that given variety. The present invention encompasses a plant directly regenerated from cells which have been transformed in accordance with the present invention. Progeny of such plants are also described. As used herein, progeny may refer to the offspring of any generation of a parent plant prepared in accordance with the present invention. Such progeny may include a DNA construct prepared in accordance with the present invention. Crossing results in the introduction of a transgene into a plant line by cross pollinating a starting line with a donor plant line. Non-limiting examples of such steps are described in U.S. Patent No. 7,151,204.

[0170] Plants may be generated through a process of backcross conversion. For example, plants include plants referred to as a backcross converted genotype, line, inbred, or hybrid.

[0171] Genetic markers may be used to assist in the introgression of one or more transgenes of the invention from one genetic background into another. Marker assisted selection offers advantages relative to conventional breeding in that it can be used to avoid errors caused by phenotypic variations. Further, genetic markers may provide data regarding the relative degree of elite germplasm in the individual progeny of a particular cross. For example, when a plant with a desired trait which otherwise has a non-agronomically desirable genetic background is crossed to an elite parent, genetic markers may be used to select progeny which not only possess the trait of interest, but also have a relatively large proportion of the desired germplasm. In this way, the number of generations required to introgress one or more traits into a particular genetic background is minimized.

[0172] The present invention also relates to methods of producing a polypeptide or domain of the present invention comprising (a) cultivating a transgenic plant or a plant cell comprising a polynucleotide encoding the polypeptide or domain under conditions conducive for production of the polypeptide or domain; and (b) recovering the polypeptide or domain.

Compositions

[0173] The present invention also relates to compositions comprising a polypeptide of the present invention. Preferably, the compositions are enriched in such a polypeptide. The term "enriched" indicates that the xylanase activity of the composition has been increased, e.g., with an enrichment factor of at least 1.1.

[0174] The composition may comprise a polypeptide of the present invention as the major enzymatic component, e.g., a mono-component composition. Alternatively, the composition

may comprise multiple enzymatic activities, such as one or more (e.g., several) enzymes selected from the group consisting of a cellulase, a hemicellulase, GH61 polypeptide, an expansin, an esterase, a laccase, a ligninolytic enzyme, a pectinase, a peroxidase, a protease, and a swollenin.

[0175] The polypeptide compositions may be prepared in accordance with methods known in the art and may be in the form of a liquid or a dry composition. For instance, the polypeptide composition may be in the form of a granulate or a microgranulate. The polypeptide to be included in the composition may be stabilized in accordance with methods known in the art.

[0176] Examples are given below of preferred uses of the polypeptide compositions of the invention. The dosage of the polypeptide composition of the invention and other conditions under which the composition is used may be determined on the basis of methods known in the art.

Uses

[0177] The present invention is also directed to the following processes for using the polypeptides having xylanase activity, or compositions thereof.

[0178] The present invention also relates to processes for degrading a cellulosic material or xylan-containing material, comprising: treating the cellulosic material or xylan-containing material with an enzyme composition in the presence of a polypeptide having xylanase activity of the present invention. In one aspect, the processes further comprise recovering the degraded or converted cellulosic material or xylan-containing material. Soluble products of degradation or conversion of the cellulosic material or xylan-containing material can be separated from insoluble cellulosic material or xylan-containing material using a method known in the art such as, for example, centrifugation, filtration, or gravity settling.

[0179] The present invention also relates to processes of producing a fermentation product, comprising: (a) saccharifying a cellulosic material or xylan-containing material with an enzyme composition in the presence of a polypeptide having xylanase activity of the present invention; (b) fermenting the saccharified cellulosic material or xylan-containing material with one or more (e.g., several) fermenting microorganisms to produce the fermentation product; and (c) recovering the fermentation product from the fermentation.

[0180] The present invention also relates to processes of fermenting a cellulosic material or xylan-containing material, comprising: fermenting the cellulosic material or xylan-containing material with one or more (e.g., several) fermenting microorganisms, wherein the cellulosic material or xylan-containing material is saccharified with an enzyme composition in the presence of a polypeptide having xylanase activity of the present invention. In one aspect, the fermenting of the cellulosic material or xylan-containing material produces a fermentation product. In another aspect, the processes further comprise recovering the fermentation

product from the fermentation.

[0181] The processes of the present invention can be used to saccharify the cellulosic material or xylan-containing material to fermentable sugars and to convert the fermentable sugars to many useful fermentation products, e.g., fuel, potable ethanol, and/or platform chemicals (e.g., acids, alcohols, ketones, gases, and the like). The production of a desired fermentation product from the cellulosic material or xylan-containing material typically involves pretreatment, enzymatic hydrolysis (saccharification), and fermentation.

[0182] The processing of the cellulosic material or xylan-containing material according to the present invention can be accomplished using methods conventional in the art. Moreover, the processes of the present invention can be implemented using any conventional biomass processing apparatus configured to operate in accordance with the invention.

[0183] Hydrolysis (saccharification) and fermentation, separate or simultaneous, include, but are not limited to, separate hydrolysis and fermentation (SHF); simultaneous saccharification and fermentation (SSF); simultaneous saccharification and co-fermentation (SSCF); hybrid hydrolysis and fermentation (HHF); separate hydrolysis and co-fermentation (SHCF); hybrid hydrolysis and co-fermentation (HHCF); and direct microbial conversion (DMC), also sometimes called consolidated bioprocessing (CBP). SHF uses separate process steps to first enzymatically hydrolyze the cellulosic material or xylan-containing material to fermentable sugars, e.g., glucose, cellobiose, and pentose monomers, and then ferment the fermentable sugars to ethanol. In SSF, the enzymatic hydrolysis of the cellulosic material or xylan-containing material and the fermentation of sugars to ethanol are combined in one step (Philippidis, G. P., 1996, Cellulose bioconversion technology, in Handbook on Bioethanol: Production and Utilization, Wyman, C. E., ed., Taylor & Francis, Washington, DC, 179-212). SSCF involves the co-fermentation of multiple sugars (Sheehan, J., and Himmel, M., 1999, Enzymes, energy and the environment: A strategic perspective on the U.S. Department of Energy's research and development activities for bioethanol, Biotechnol. Prog. 15: 817-827). HHF involves a separate hydrolysis step, and in addition a simultaneous saccharification and hydrolysis step, which can be carried out in the same reactor. The steps in an HHF process can be carried out at different temperatures, *i.e.*, high temperature enzymatic saccharification followed by SSF at a lower temperature that the fermentation strain can tolerate. DMC combines all three processes (enzyme production, hydrolysis, and fermentation) in one or more (e.g., several) steps where the same organism is used to produce the enzymes for conversion of the cellulosic material or xylan-containing material to fermentable sugars and to convert the fermentable sugars into a final product (Lynd et al., 2002, Microbial cellulose utilization: Fundamentals and biotechnology, Microbiol. Mol. Biol. Reviews 66: 506-577). It is understood herein that any method known in the art comprising pretreatment, enzymatic hydrolysis (saccharification), fermentation, or a combination thereof, can be used in the practicing the processes of the present invention.

[0184] A conventional apparatus can include a fed-batch stirred reactor, a batch stirred reactor, a continuous flow stirred reactor with ultrafiltration, and/or a continuous plug-flow

column reactor (Fernanda de Castilhos Corazza, Flávio Faria de Moraes, Gisella Maria Zanin and Ivo Neitzel, 2003, Optimal control in fed-batch reactor for the cellobiose hydrolysis, *Acta Scientiarum. Technology* 25: 33-38; Gusakov and Sinitsyn, 1985, Kinetics of the enzymatic hydrolysis of cellulose: 1. A mathematical model for a batch reactor process, *Enz. Microb. Technol.* 7: 346-352), an attrition reactor (Ryu and Lee, 1983, Bioconversion of waste cellulose by using an attrition bioreactor, *Biotechnol. Bioeng.* 25: 53-65), or a reactor with intensive stirring induced by an electromagnetic field (Gusakov et al., 1996, Enhancement of enzymatic cellulose hydrolysis using a novel type of bioreactor with intensive stirring induced by electromagnetic field, *Appl. Biochem. Biotechnol.* 56: 141-153). Additional reactor types include fluidized bed, upflow blanket, immobilized, and extruder type reactors for hydrolysis and/or fermentation.

[0185] Pretreatment. In practicing the processes of the present invention, any pretreatment process known in the art can be used to disrupt plant cell wall components of the cellulosic material or xylan-containing material (Chandra et al., 2007, Substrate pretreatment: The key to effective enzymatic hydrolysis of lignocellulosics?, *Adv. Biochem. Engin./Biotechnol.* 108: 67-93; Galbe and Zacchi, 2007, Pretreatment of lignocellulosic materials for efficient bioethanol production, *Adv. Biochem. Engin./Biotechnol.* 108: 41-65; Hendriks and Zeeman, 2009, Pretreatments to enhance the digestibility of lignocellulosic biomass, *Bioresource Technol.* 100: 10-18; Mosier et al., 2005, Features of promising technologies for pretreatment of lignocellulosic biomass, *Bioresource Technol.* 96: 673-686; Taherzadeh and Karimi, 2008, Pretreatment of lignocellulosic wastes to improve ethanol and biogas production: A review, *Int. J. of Mol. Sci.* 9: 1621-1651; Yang and Wyman, 2008, Pretreatment: the key to unlocking low-cost cellulosic ethanol, *Biofuels Bioproducts and Biorefining-Biofpr.* 2: 26-40).

[0186] The cellulosic material or xylan-containing material can also be subjected to particle size reduction, sieving, pre-soaking, wetting, washing, and/or conditioning prior to pretreatment using methods known in the art.

[0187] Conventional pretreatments include, but are not limited to, steam pretreatment (with or without explosion), dilute acid pretreatment, hot water pretreatment, alkaline pretreatment, lime pretreatment, wet oxidation, wet explosion, ammonia fiber explosion, organosolv pretreatment, and biological pretreatment. Additional pretreatments include ammonia percolation, ultrasound, electroporation, microwave, supercritical CO₂, supercritical H₂O, ozone, ionic liquid, and gamma irradiation pretreatments.

[0188] The cellulosic material or xylan-containing material can be pretreated before hydrolysis and/or fermentation. Pretreatment is preferably performed prior to the hydrolysis. Alternatively, the pretreatment can be carried out simultaneously with enzyme hydrolysis to release fermentable sugars, such as glucose, xylose, and/or cellobiose. In most cases the pretreatment step itself results in some conversion of biomass to fermentable sugars (even in absence of enzymes).

[0189] Steam Pretreatment. In steam pretreatment, the cellulosic material or xylan-containing

material is heated to disrupt the plant cell wall components, including lignin, hemicellulose, and cellulose to make the cellulose and other fractions, e.g., hemicellulose, accessible to enzymes. The cellulosic material or xylan-containing material is passed to or through a reaction vessel where steam is injected to increase the temperature to the required temperature and pressure and is retained therein for the desired reaction time. Steam pretreatment is preferably performed at 140-250°C, e.g., 160-200°C or 170-190°C, where the optimal temperature range depends on addition of a chemical catalyst. Residence time for the steam pretreatment is preferably 1-60 minutes, e.g., 1-30 minutes, 1-20 minutes, 3-12 minutes, or 4-10 minutes, where the optimal residence time depends on temperature range and addition of a chemical catalyst. Steam pretreatment allows for relatively high solids loadings, so that the cellulosic material or xylan-containing material is generally only moist during the pretreatment. The steam pretreatment is often combined with an explosive discharge of the material after the pretreatment, which is known as steam explosion, that is, rapid flashing to atmospheric pressure and turbulent flow of the material to increase the accessible surface area by fragmentation (Duff and Murray, 1996, *Bioresource Technology* 855: 1-33; Galbe and Zacchi, 2002, *Appl. Microbiol. Biotechnol.* 59: 618-628; U.S. Patent Application No. 2002/0164730). During steam pretreatment, hemicellulose acetyl groups are cleaved and the resulting acid autocatalyzes partial hydrolysis of the hemicellulose to monosaccharides and oligosaccharides. Lignin is removed to only a limited extent.

[0190] Chemical Pretreatment: The term "chemical treatment" refers to any chemical pretreatment that promotes the separation and/or release of cellulose, hemicellulose, and/or lignin. Such a pretreatment can convert crystalline cellulose to amorphous cellulose. Examples of suitable chemical pretreatment processes include, for example, dilute acid pretreatment, lime pretreatment, wet oxidation, ammonia fiber/freeze explosion (AFEX), ammonia percolation (APR), ionic liquid, and organosolv pretreatments.

[0191] A catalyst such as H₂SO₄ or SO₂ (typically 0.3 to 5% w/w) is often added prior to steam pretreatment, which decreases the time and temperature, increases the recovery, and improves enzymatic hydrolysis (Ballesteros et al., 2006, *Appl. Biochem. Biotechnol.* 129-132: 496-508; Varga et al., 2004, *Appl. Biochem. Biotechnol.* 113-116: 509-523; Sassner et al., 2006, *Enzyme Microb. Technol.* 39: 756-762). In dilute acid pretreatment, the cellulosic material or xylan-containing material is mixed with dilute acid, typically H₂SO₄, and water to form a slurry, heated by steam to the desired temperature, and after a residence time flashed to atmospheric pressure. The dilute acid pretreatment can be performed with a number of reactor designs, e.g., plug-flow reactors, counter-current reactors, or continuous counter-current shrinking bed reactors (Duff and Murray, 1996, *supra*; Schell et al., 2004, *Bioresource Technol.* 91: 179-188; Lee et al., 1999, *Adv. Biochem. Eng. Biotechnol.* 65: 93-115).

[0192] Several methods of pretreatment under alkaline conditions can also be used. These alkaline pretreatments include, but are not limited to, sodium hydroxide, lime, wet oxidation, ammonia percolation (APR), and ammonia fiber/freeze explosion (AFEX).

[0193] Lime pretreatment is performed with calcium oxide or calcium hydroxide at

temperatures of 85-150°C and residence times from 1 hour to several days (Wyman et al., 2005, *Bioresource Technol.* 96: 1959-1966; Mosier et al., 2005, *Bioresource Technol.* 96:673-686). WO 2006/110891, WO 2006/110899, WO 2006/110900, and WO 2006/110901 disclose pretreatment methods using ammonia.

[0194] Wet oxidation is a thermal pretreatment performed typically at 180-200°C for 5-15 minutes with addition of an oxidative agent such as hydrogen peroxide or over-pressure of oxygen (Schmidt and Thomsen, 1998, *Bioresource Technol.* 64: 139-151; Palonen et al., 2004, *Appl. Biochem. Biotechnol.* 117: 1-17; Varga et al., 2004, *Biotechnol. Bioeng.* 88: 567-574; Martin et al., 2006, *J. Chem. Technol. Biotechnol.* 81: 1669-1677). The pretreatment is performed preferably at 1-40% dry matter, e.g., 2-30% dry matter or 5-20% dry matter, and often the initial pH is increased by the addition of alkali such as sodium carbonate.

[0195] A modification of the wet oxidation pretreatment method, known as wet explosion (combination of wet oxidation and steam explosion) can handle dry matter up to 30%. In wet explosion, the oxidizing agent is introduced during pretreatment after a certain residence time. The pretreatment is then ended by flashing to atmospheric pressure (WO 2006/032282).

[0196] Ammonia fiber explosion (AFEX) involves treating the cellulosic material or xylan-containing material with liquid or gaseous ammonia at moderate temperatures such as 90-150°C and high pressure such as 17-20 bar for 5-10 minutes, where the dry matter content can be as high as 60% (Gollapalli et al., 2002, *Appl. Biochem. Biotechnol.* 98: 23-35; Chundawat et al., 2007, *Biotechnol. Bioeng.* 96: 219-231; Alizadeh et al., 2005, *Appl. Biochem. Biotechnol.* 121: 1133-1141; Teymouri et al., 2005, *Bioresource Technol.* 96: 2014-2018). During AFEX pretreatment cellulose and hemicelluloses remain relatively intact. Lignin-carbohydrate complexes are cleaved.

[0197] Organosolv pretreatment delignifies the cellulosic material or xylan-containing material by extraction using aqueous ethanol (40-60% ethanol) at 160-200°C for 30-60 minutes (Pan et al., 2005, *Biotechnol. Bioeng.* 90: 473-481; Pan et al., 2006, *Biotechnol. Bioeng.* 94: 851-861; Kurabi et al., 2005, *Appl. Biochem. Biotechnol.* 121: 219-230). Sulphuric acid is usually added as a catalyst. In organosolv pretreatment, the majority of hemicellulose and lignin is removed.

[0198] Other examples of suitable pretreatment methods are described by Schell et al., 2003, *Appl. Biochem. and Biotechnol.* 105-108: 69-85, and Mosier et al., 2005, *Bioresource Technology* 96: 673-686, and U.S. Published Application 2002/0164730.

[0199] In one aspect, the chemical pretreatment is preferably carried out as a dilute acid treatment, and more preferably as a continuous dilute acid treatment. The acid is typically sulfuric acid, but other acids can also be used, such as acetic acid, citric acid, nitric acid, phosphoric acid, tartaric acid, succinic acid, hydrogen chloride, or mixtures thereof. Mild acid treatment is conducted in the pH range of preferably 1-5, e.g., 1-4 or 1-2.5. In one aspect, the acid concentration is in the range from preferably 0.01 to 10 wt % acid, e.g., 0.05 to 5 wt % acid or 0.1 to 2 wt % acid. The acid is contacted with the cellulosic material or xylan-containing

material and held at a temperature in the range of preferably 140-200°C, *e.g.*, 165-190°C, for periods ranging from 1 to 60 minutes.

[0200] In another aspect, pretreatment takes place in an aqueous slurry. In preferred aspects, the cellulosic material or xylan-containing material is present during pretreatment in amounts preferably between 10-80 wt %, *e.g.*, 20-70 wt % or 30-60 wt %, such as around 40 wt %. The pretreated cellulosic material or xylan-containing material can be unwashed or washed using any method known in the art, *e.g.*, washed with water.

[0201] Mechanical Pretreatment or Physical Pretreatment: The term "mechanical pretreatment" or "physical pretreatment" refers to any pretreatment that promotes size reduction of particles. For example, such pretreatment can involve various types of grinding or milling (*e.g.*, dry milling, wet milling, or vibratory ball milling).

[0202] The cellulosic material or xylan-containing material can be pretreated both physically (mechanically) and chemically. Mechanical or physical pretreatment can be coupled with steaming/steam explosion, hydrothermolysis, dilute or mild acid treatment, high temperature, high pressure treatment, irradiation (*e.g.*, microwave irradiation), or combinations thereof. In one aspect, high pressure means pressure in the range of preferably about 100 to about 400 psi, *e.g.*, about 150 to about 250 psi. In another aspect, high temperature means temperatures in the range of about 100 to about 300°C, *e.g.*, about 140 to about 200°C. In a preferred aspect, mechanical or physical pretreatment is performed in a batch-process using a steam gun hydrolyzer system that uses high pressure and high temperature as defined above, *e.g.*, a Sands Hydrolyzer available from Sands Defibrator AB, Sweden. The physical and chemical pretreatments can be carried out sequentially or simultaneously, as desired.

[0203] Accordingly, in a preferred aspect, the cellulosic material or xylan-containing material is subjected to physical (mechanical) or chemical pretreatment, or any combination thereof, to promote the separation and/or release of cellulose, hemicellulose, and/or lignin.

[0204] Biological Pretreatment: The term "biological pretreatment" refers to any biological pretreatment that promotes the separation and/or release of cellulose, hemicellulose, and/or lignin from the cellulosic material or xylan-containing material. Biological pretreatment techniques can involve applying lignin-solubilizing microorganisms and/or enzymes (see, for example, Hsu, T.-A., 1996, Pretreatment of biomass, in Handbook on Bioethanol: Production and Utilization, Wyman, C. E., ed., Taylor & Francis, Washington, DC, 179-212; Ghosh and Singh, 1993, Physicochemical and biological treatments for enzymatic/microbial conversion of cellulosic biomass, *Adv. Appl. Microbiol.* 39: 295-333; McMillan, J. D., 1994, Pretreating lignocellulosic biomass: a review, in *Enzymatic Conversion of Biomass for Fuels Production*, Himmel, M. E., Baker, J. O., and Overend, R. P., eds., ACS Symposium Series 566, American Chemical Society, Washington, DC, chapter 15; Gong, C. S., Cao, N. J., Du, J., and Tsao, G. T., 1999, Ethanol production from renewable resources, in *Advances in Biochemical Engineering/Biotechnology*, Scheper, T., ed., Springer-Verlag Berlin Heidelberg, Germany, 65: 207-241; Olsson and Hahn-Hagerdal, 1996, Fermentation of lignocellulosic hydrolysates for

ethanol production, *Enz. Microb. Tech.* 18: 312-331; and Vallander and Eriksson, 1990, *Production of ethanol from lignocellulosic materials: State of the art*, *Adv. Biochem. Eng./Biotechnol.* 42: 63-95).

[0205] Saccharification. In the hydrolysis step, also known as saccharification, the cellulosic material or xylan-containing material, e.g., pretreated, is hydrolyzed to break down cellulose and/or hemicellulose to fermentable sugars, such as glucose, cellobiose, xylose, xylulose, arabinose, mannose, galactose, and/or soluble oligosaccharides. The hydrolysis is performed enzymatically by an enzyme composition in the presence of a polypeptide having xylanase activity of the present invention. The enzymes of the compositions can be added simultaneously or sequentially.

[0206] Enzymatic hydrolysis is preferably carried out in a suitable aqueous environment under conditions that can be readily determined by one skilled in the art. In one aspect, hydrolysis is performed under conditions suitable for the activity of the enzyme(s), *i.e.*, optimal for the enzyme(s). The hydrolysis can be carried out as a fed batch or continuous process where the cellulosic material or xylan-containing material is fed gradually to, for example, an enzyme containing hydrolysis solution.

[0207] The saccharification is generally performed in stirred-tank reactors or fermentors under controlled pH, temperature, and mixing conditions. Suitable process time, temperature and pH conditions can readily be determined by one skilled in the art. For example, the saccharification can last up to 200 hours, but is typically performed for preferably about 12 to about 120 hours, e.g., about 16 to about 72 hours or about 24 to about 48 hours. The temperature is in the range of preferably about 25°C to about 70°C, e.g., about 30°C to about 65°C, about 40°C to about 60°C, or about 50°C to about 55°C. The pH is in the range of preferably about 3 to about 8, e.g., about 3.5 to about 7, about 4 to about 6, or about 5.0 to about 5.5. The dry solids content is in the range of preferably about 5 to about 50 wt %, e.g., about 10 to about 40 wt % or about 20 to about 30 wt %.

[0208] The enzyme compositions can comprise any protein useful in degrading the cellulosic material or xylan-containing material.

[0209] In one aspect, the enzyme composition comprises or further comprises one or more (e.g., several) proteins selected from the group consisting of a cellulase, a GH61 polypeptide having cellulolytic enhancing activity, a hemicellulase, an esterase, an expansin, a laccase, a ligninolytic enzyme, a pectinase, a peroxidase, a protease, and a swollenin. In another aspect, the hemicellulase is preferably one or more (e.g., several) enzymes selected from the group consisting of an acetylmannan esterase, an acetylxylan esterase, an arabinanase, an arabinofuranosidase, a coumaric acid esterase, a feruloyl esterase, a galactosidase, a glucuronidase, a glucuronoyl esterase, a mannanase, a mannosidase, a xylanase, and a xylosidase. In another aspect, the cellulase is preferably one or more (e.g., several) enzymes selected from the group consisting of an endoglucanase, a cellobiohydrolase, and a beta-glucosidase.

[0210] In another aspect, the enzyme composition comprises one or more (e.g., several) cellulolytic enzymes. In another aspect, the enzyme composition comprises or further comprises one or more (e.g., several) hemicellulolytic enzymes. In another aspect, the enzyme composition comprises one or more (e.g., several) cellulolytic enzymes and one or more (e.g., several) hemicellulolytic enzymes. In another aspect, the enzyme composition comprises one or more (e.g., several) enzymes selected from the group of cellulolytic enzymes and hemicellulolytic enzymes.

[0211] In another aspect, the enzyme composition comprises an acetylmannan esterase. In another aspect, the enzyme composition comprises an acetylxylan esterase. In another aspect, the enzyme composition comprises an arabinanase (e.g., alpha-L-arabinanase). In another aspect, the enzyme composition comprises an arabinofuranosidase (e.g., alpha-L-arabinofuranosidase). In another aspect, the enzyme composition comprises a coumaric acid esterase. In another aspect, the enzyme composition comprises a feruloyl esterase. In another aspect, the enzyme composition comprises a galactosidase (e.g., alpha-galactosidase and/or beta-galactosidase). In another aspect, the enzyme composition comprises a glucuronidase (e.g., alpha-D-glucuronidase). In another aspect, the enzyme composition comprises a glucuronoyl esterase. In another aspect, the enzyme composition comprises a mannanase. In another aspect, the enzyme composition comprises a mannosidase (e.g., beta-mannosidase). In another aspect, the enzyme composition comprises a xylanase. In a preferred aspect, the xylanase is a Family 10 xylanase. In another aspect, the enzyme composition comprises a xylosidase (e.g., beta-xylosidase).

[0212] In another aspect, the enzyme composition comprises an endoglucanase. In another aspect, the enzyme composition comprises a cellobiohydrolase. In another aspect, the enzyme composition comprises a beta-glucosidase. In another aspect, the enzyme composition comprises a polypeptide having cellulolytic enhancing activity. In another aspect, the enzyme composition comprises an endoglucanase and a polypeptide having cellulolytic enhancing activity. In another aspect, the enzyme composition comprises a cellobiohydrolase and a polypeptide having cellulolytic enhancing activity. In another aspect, the enzyme composition comprises a beta-glucosidase and a polypeptide having cellulolytic enhancing activity. In another aspect, the enzyme composition comprises an endoglucanase and a cellobiohydrolase. In another aspect, the enzyme composition comprises an endoglucanase and a beta-glucosidase. In another aspect, the enzyme composition comprises a cellobiohydrolase and a beta-glucosidase. In another aspect, the enzyme composition comprises an endoglucanase, a cellobiohydrolase, and a polypeptide having cellulolytic enhancing activity. In another aspect, the enzyme composition comprises an endoglucanase, a beta-glucosidase, and a polypeptide having cellulolytic enhancing activity. In another aspect, the enzyme composition comprises a cellobiohydrolase, a beta-glucosidase, and a polypeptide having cellulolytic enhancing activity. In another aspect, the enzyme composition comprises an endoglucanase, a cellobiohydrolase, and a beta-glucosidase. In another aspect, the enzyme composition comprises an endoglucanase, a cellobiohydrolase, a beta-glucosidase, and a polypeptide having cellulolytic enhancing activity.

[0213] In another aspect, the enzyme composition comprises an esterase. In another aspect, the enzyme composition comprises an expansin. In another aspect, the enzyme composition comprises a laccase. In another aspect, the enzyme composition comprises a ligninolytic enzyme. In a preferred aspect, the ligninolytic enzyme is a manganese peroxidase. In another preferred aspect, the ligninolytic enzyme is a lignin peroxidase. In another preferred aspect, the ligninolytic enzyme is a H₂O₂-producing enzyme. In another aspect, the enzyme composition comprises a pectinase. In another aspect, the enzyme composition comprises a peroxidase. In another aspect, the enzyme composition comprises a protease. In another aspect, the enzyme composition comprises a swollenin

[0214] In the processes of the present invention, the enzyme(s) can be added prior to or during fermentation, *e.g.*, during saccharification or during or after propagation of the fermenting microorganism(s).

[0215] One or more (*e.g.*, several) components of the enzyme composition may be wild-type proteins, recombinant proteins, or a combination of wild-type proteins and recombinant proteins. For example, one or more (*e.g.*, several) components may be native proteins of a cell, which is used as a host cell to express recombinantly one or more (*e.g.*, several) other components of the enzyme composition. One or more (*e.g.*, several) components of the enzyme composition may be produced as monocomponents, which are then combined to form the enzyme composition. The enzyme composition may be a combination of multicomponent and monocomponent protein preparations.

[0216] The enzymes used in the processes of the present invention may be in any form suitable for use, such as, for example, a crude fermentation broth with or without cells removed, a cell lysate with or without cellular debris, a semi-purified or purified enzyme preparation, or a host cell as a source of the enzymes. The enzyme composition may be a dry powder or granulate, a non-dusting granulate, a liquid, a stabilized liquid, or a stabilized protected enzyme. Liquid enzyme preparations may, for instance, be stabilized by adding stabilizers such as a sugar, a sugar alcohol or another polyol, and/or lactic acid or another organic acid according to established processes.

[0217] The optimum amounts of the enzymes and polypeptides having xylanase activity depend on several factors including, but not limited to, the mixture of component cellulolytic enzymes, the cellulosic material or xylan-containing material, the concentration of cellulosic material or xylan-containing material, the pretreatment(s) of the cellulosic material or xylan-containing material, temperature, time, pH, and inclusion of fermenting organism (*e.g.*, yeast for Simultaneous Saccharification and Fermentation).

[0218] In one aspect, an effective amount of cellulolytic or hemicellulolytic enzyme to the cellulosic material or xylan-containing material is about 0.5 to about 50 mg, *e.g.*, about 0.5 to about 40 mg, about 0.5 to about 25 mg, about 0.75 to about 20 mg, about 0.75 to about 15 mg, about 0.5 to about 10 mg, or about 2.5 to about 10 mg per g of the cellulosic material or

xylan-containing material.

[0219] In another aspect, an effective amount of a polypeptide having xylanase activity to the cellulosic material or xylan-containing material is about 0.01 to about 50.0 mg, *e.g.*, about 0.01 to about 40 mg, about 0.01 to about 30 mg, about 0.01 to about 20 mg, about 0.01 to about 10 mg, about 0.01 to about 5 mg, about 0.025 to about 1.5 mg, about 0.05 to about 1.25 mg, about 0.075 to about 1.25 mg, about 0.1 to about 1.25 mg, about 0.15 to about 1.25 mg, or about 0.25 to about 1.0 mg per g of the cellulosic material or xylan-containing material.

[0220] In another aspect, an effective amount of a polypeptide having xylanase activity to cellulolytic or hemicellulolytic enzyme is about 0.005 to about 1.0 g, *e.g.*, about 0.01 to about 1.0 g, about 0.15 to about 0.75 g, about 0.15 to about 0.5 g, about 0.1 to about 0.5 g, about 0.1 to about 0.25 g, or about 0.05 to about 0.2 g per g of cellulolytic or hemicellulolytic enzyme.

[0221] The polypeptides having cellulolytic enzyme activity or hemicellulolytic enzyme activity as well as other proteins/polypeptides useful in the degradation of the cellulosic material or xylan-containing material, *e.g.*, GH61 polypeptides having cellulolytic enhancing activity (collectively hereinafter "polypeptides having enzyme activity") can be derived or obtained from any suitable origin, including, bacterial, fungal, yeast, plant, or mammalian origin. The term "obtained" also means herein that the enzyme may have been produced recombinantly in a host organism employing methods described herein, wherein the recombinantly produced enzyme is either native or foreign to the host organism or has a modified amino acid sequence, *e.g.*, having one or more (*e.g.*, several) amino acids that are deleted, inserted and/or substituted, *i.e.*, a recombinantly produced enzyme that is a mutant and/or a fragment of a native amino acid sequence or an enzyme produced by nucleic acid shuffling processes known in the art. Encompassed within the meaning of a native enzyme are natural variants and within the meaning of a foreign enzyme are variants obtained recombinantly, such as by site-directed mutagenesis or shuffling.

[0222] One or more (*e.g.*, several) components of the enzyme composition may be a recombinant component, *i.e.*, produced by cloning of a DNA sequence encoding the single component and subsequent cell transformed with the DNA sequence and expressed in a host (see, for example, WO 91/17243 and WO 91/17244). The host is preferably a heterologous host (enzyme is foreign to host), but the host may under certain conditions also be a homologous host (enzyme is native to host). Monocomponent cellulolytic proteins may also be prepared by purifying such a protein from a fermentation broth.

[0223] In one aspect, the one or more (*e.g.*, several) cellulolytic enzymes comprise a commercial cellulolytic enzyme preparation. Examples of commercial cellulolytic enzyme preparations suitable for use in the present invention include, for example, CELLIC[™] CTec (Novozymes A/S), CELLIC[™] CTec2 (Novozymes A/S), CELLUCLAST[™] (Novozymes A/S), NOVOZYME[™] 188 (Novozymes A/S), CELLUZYME[™] (Novozymes A/S), CEREFLO[™] (Novozymes A/S), and ULTRAFLO[™] (Novozymes A/S), ACCELERASE[™] (Genencor Int.),

LAMINEX™ (Genencor Int.), SPEZYME™ CP (Genencor Int.), FILTRASE® NL (DSM); METHAPLUS® S/L 100 (DSM). ROHAMENT™ 7069 W (Röhm GmbH), FIBREZYME® LDI (Dyadic International, Inc.), FIBREZYME® LBR (Dyadic International, Inc.), or VISCOSTAR® 150L (Dyadic International, Inc.). The cellulase enzymes are added in amounts effective from about 0.001 to about 5.0 wt % of solids, e.g., about 0.025 to about 4.0 wt % of solids or about 0.005 to about 2.0 wt % of solids.

[0224] Examples of bacterial endoglucanases that can be used in the processes of the present invention, include, but are not limited to, an *Acidothermus cellulolyticus* endoglucanase (WO 91/05039; WO 93/15186; U.S. Patent No. 5,275,944; WO 96/02551; U.S. Patent No. 5,536,655, WO 00/70031, WO 05/093050); *Thermobifida fusca* endoglucanase III (WO 05/093050); and *Thermobifida fusca* endoglucanase V (WO 05/093050).

[0225] Examples of fungal endoglucanases that can be used in the present invention, include, but are not limited to, a *Trichoderma reesei* endoglucanase I (Penttila et al., 1986, Gene 45: 253-263, *Trichoderma reesei* Cel7B endoglucanase I (GENBANK™ accession no. M15665), *Trichoderma reesei* endoglucanase II (Saloheimo, et al., 1988, Gene 63:11-22), *Trichoderma reesei* Cel5A endoglucanase II (GENBANK™ accession no. M19373), *Trichoderma reesei* endoglucanase III (Okada et al., 1988, Appl. Environ. Microbiol. 64: 555-563, GENBANK™ accession no. AB003694), *Trichoderma reesei* endoglucanase V (Saloheimo et al., 1994, Molecular Microbiology 13: 219-228, GENBANK™ accession no. Z33381), *Aspergillus aculeatus* endoglucanase (Ooi et al., 1990, Nucleic Acids Research 18: 5884), *Aspergillus kawachii* endoglucanase (Sakamoto et al., 1995, Current Genetics 27: 435-439), *Erwinia carotovora* endoglucanase (Saarilahti et al., 1990, Gene 90: 9-14), *Fusarium oxysporum* endoglucanase (GENBANK™ accession no. L29381), *Humicola grisea* var. *thermoidea* endoglucanase (GENBANK™ accession no. AB003107), *Melanocarpus albomyces* endoglucanase (GENBANK™ accession no. MAL515703), *Neurospora crassa* endoglucanase (GENBANK™ accession no. XM_324477), *Humicola insolens* endoglucanase V, *Myceliophthora thermophila* CBS 117.65 endoglucanase, basidiomycete CBS 495.95 endoglucanase, basidiomycete CBS 494.95 endoglucanase, *Thielavia terrestris* NRRL 8126 CEL6B endoglucanase, *Thielavia terrestris* NRRL 8126 CEL6C endoglucanase, *Thielavia terrestris* NRRL 8126 CEL7C endoglucanase, *Thielavia terrestris* NRRL 8126 CEL7E endoglucanase, *Thielavia terrestris* NRRL 8126 CEL7F endoglucanase, *Cladorrhinum foecundissimum* ATCC 62373 CEL7A endoglucanase, and *Trichoderma reesei* strain No. VTT-D-80133 endoglucanase (GENBANK™ accession no. M15665).

[0226] Examples of cellobiohydrolases useful in the present invention include, but are not limited to, *Aspergillus aculeatus* cellobiohydrolase II (WO 2011/059740), *Chaetomium thermophilum* cellobiohydrolase I, *Chaetomium thermophilum* cellobiohydrolase II, *Humicola insolens* cellobiohydrolase I, *Myceliophthora thermophila* cellobiohydrolase II (WO 2009/042871), *Thielavia hyrcanie* cellobiohydrolase II (WO 2010/141325), *Thielavia terrestris*

cellobiohydrolase II (CEL6A, WO 2006/074435), *Trichoderma reesei* cellobiohydrolase I, *Trichoderma reesei* cellobiohydrolase II, and *Trichophaea saccata* cellobiohydrolase II (WO 2010/057086).

[0227] Examples of beta-glucosidases useful in the present invention include, but are not limited to, beta-glucosidases from *Aspergillus aculeatus* (Kawaguchi et al., 1996, Gene 173: 287-288), *Aspergillus fumigatus* (WO 2005/047499), *Aspergillus niger* (Dan et al., 2000, J. Biol. Chem. 275: 4973-4980), *Aspergillus oryzae* (WO 02/095014), *Penicillium brasilianum* IBT 20888 (WO 2007/019442 and WO 2010/088387), *Thielavia terrestris* (WO 2011/035029), and *Trichophaea saccata* (WO 2007/019442).

[0228] The beta-glucosidase may be a fusion protein. In one aspect, the beta-glucosidase is an *Aspergillus oryzae* beta-glucosidase variant BG fusion protein (WO 2008/057637) or an *Aspergillus oryzae* beta-glucosidase fusion protein (WO 2008/057637).

[0229] Other useful endoglucanases, cellobiohydrolases, and beta-glucosidases are disclosed in numerous Glycosyl Hydrolase families using the classification according to Henrissat, 1991, A classification of glycosyl hydrolases based on amino-acid sequence similarities, Biochem. J. 280: 309-316, and Henrissat and Bairoch, 1996, Updating the sequence-based classification of glycosyl hydrolases, Biochem. J. 316: 695-696.

[0230] Other cellulolytic enzymes that may be used in the present invention are described in WO 98/13465, WO 98/15619, WO 98/15633, WO 99/06574, WO 99/10481, WO 99/25847, WO 99/31255, WO 02/101078, WO 2003/027306, WO 2003/052054, WO 2003/052055, WO 2003/052056, WO 2003/052057, WO 2003/052118, WO 2004/016760, WO 2004/043980, WO 2004/048592, WO 2005/001065, WO 2005/028636, WO 2005/093050, WO 2005/093073, WO 2006/074005, WO 2006/117432, WO 2007/071818, WO 2007/071820, WO 2008/008070, WO 2008/008793, U.S. Patent No. 5,457,046, U.S. Patent No. 5,648,263, and U.S. Patent No. 5,686,593.

[0231] In the processes of the present invention, any GH61 polypeptide having cellulolytic enhancing activity can be used.

[0232] Examples of GH61 polypeptides having cellulolytic enhancing activity useful in the processes of the present invention include, but are not limited to, GH61 polypeptides from *Thielavia terrestris* (WO 2005/074647, WO 2008/148131, and WO 2011/035027), *Thermoascus aurantiacus* (WO 2005/074656 and WO 2010/065830), *Trichoderma reesei* (WO 2007/089290), *Myceliophthora thermophila* (WO 2009/085935, WO 2009/085859, WO 2009/085864, WO 2009/085868), *Aspergillus fumigatus* (WO 2010/138754), GH61 polypeptides from *Penicillium pinophilum* (WO 2011/005867), *Thermoascus* sp. (WO 2011/039319), *Penicillium* sp. (WO 2011/041397), and *Thermoascus crustaceus* (WO 2011/041504).

[0233] In one aspect, the GH61 polypeptide having cellulolytic enhancing activity is used in the

presence of a soluble activating divalent metal cation according to WO 2008/151043, e.g., manganese sulfate.

[0234] In one aspect, the GH61 polypeptide having cellulolytic enhancing activity is used in the presence of a dioxy compound, a bicyclic compound, a heterocyclic compound, a nitrogen-containing compound, a quinone compound, a sulfur-containing compound, or a liquor obtained from a pretreated cellulosic material or xylan-containing material such as pretreated corn stover (PCS).

[0235] The dioxy compound may include any suitable compound containing two or more oxygen atoms. In some aspects, the dioxy compounds contain a substituted aryl moiety as described herein. The dioxy compounds may comprise one or more (e.g., several) hydroxyl and/or hydroxyl derivatives, but also include substituted aryl moieties lacking hydroxyl and hydroxyl derivatives. Non-limiting examples of the dioxy compounds include pyrocatechol or catechol; caffeic acid; 3,4-dihydroxybenzoic acid; 4-tert-butyl-5-methoxy-1,2-benzenediol; pyrogallol; gallic acid; methyl-3,4,5-trihydroxybenzoate; 2,3,4-trihydroxybenzophenone; 2,6-dimethoxyphenol; sinapinic acid; 3,5-dihydroxybenzoic acid; 4-chloro-1,2-benzenediol; 4-nitro-1,2-benzenediol; tannic acid; ethyl gallate; methyl glycolate; dihydroxyfumaric acid; 2-butyne-1,4-diol; (croconic acid; 1,3-propanediol; tartaric acid; 2,4-pentanediol; 3-ethoxy-1,2-propanediol; 2,4,4'-trihydroxybenzophenone; cis-2-butene-1,4-diol; 3,4-dihydroxy-3-cyclobutene-1,2-dione; dihydroxyacetone; acrolein acetal; methyl-4-hydroxybenzoate; 4-hydroxybenzoic acid; and methyl-3,5-dimethoxy-4-hydroxybenzoate; or a salt or solvate thereof.

[0236] The bicyclic compound may include any suitable substituted fused ring system as described herein. The compounds may comprise one or more (e.g., several) additional rings, and are not limited to a specific number of rings unless otherwise stated. In one aspect, the bicyclic compound is a flavonoid. In another aspect, the bicyclic compound is an optionally substituted isoflavonoid. In another aspect, the bicyclic compound is an optionally substituted flavylum ion, such as an optionally substituted anthocyanidin or optionally substituted anthocyanin, or derivative thereof. Non-limiting examples of the bicyclic compounds include epicatechin; quercetin; myricetin; taxifolin; kaempferol; morin; acacetin; naringenin; isorhamnetin; apigenin; cyanidin; cyanin; kuromanin; keracyanin; or a salt or solvate thereof.

[0237] The heterocyclic compound may be any suitable compound, such as an optionally substituted aromatic or non-aromatic ring comprising a heteroatom, as described herein. In one aspect, the heterocyclic is a compound comprising an optionally substituted heterocycloalkyl moiety or an optionally substituted heteroaryl moiety. In another aspect, the optionally substituted heterocycloalkyl moiety or optionally substituted heteroaryl moiety is an optionally substituted 5-membered heterocycloalkyl or an optionally substituted 5-membered heteroaryl moiety. In another aspect, the optionally substituted heterocycloalkyl or optionally substituted heteroaryl moiety is an optionally substituted moiety selected from pyrazolyl, furanyl, imidazolyl, isoxazolyl, oxadiazolyl, oxazolyl, pyrrolyl, pyridyl, pyrimidyl, pyridazinyl, thiazolyl, triazolyl, thienyl, dihydrothieno-pyrazolyl, thianaphthenyl, carbazolyl, benzimidazolyl,

benzothienyl, benzofuranyl, indolyl, quinoliny, benzotriazolyl, benzothiazolyl, benzooxazolyl, benzimidazolyl, isoquinoliny, isoindolyl, acridiny, benzoisazolyl, dimethylhydantoin, pyraziny, tetrahydrofuranyl, pyrroliny, pyrrolidinyl, morpholiny, indolyl, diazepiny, azepiny, thiepinyl, piperidinyl, and oxepiny. In another aspect, the optionally substituted heterocycloalkyl moiety or optionally substituted heteroaryl moiety is an optionally substituted furanyl. Non-limiting examples of the heterocyclic compounds include (1,2-dihydroxyethyl)-3,4-dihydroxyfuran-2(5H)-one; 4-hydroxy-5-methyl-3-furanone; 5-hydroxy-2(5H)-furanone; [1,2-dihydroxyethyl]furan-2,3,4(5H)-trione; α -hydroxy- γ -butyrolactone; ribonic γ -lactone; aldohexuronic acid γ -lactone; gluconic acid δ -lactone; 4-hydroxycoumarin; dihydrobenzofuran; 5-(hydroxymethyl)furfural; furoin; 2(5H)-furanone; 5,6-dihydro-2H-pyran-2-one; and 5,6-dihydro-4-hydroxy-6-methyl-2H-pyran-2-one; or a salt or solvate thereof.

[0238] The nitrogen-containing compound may be any suitable compound with one or more nitrogen atoms. In one aspect, the nitrogen-containing compound comprises an amine, imine, hydroxylamine, or nitroxide moiety. Non-limiting examples of the nitrogen-containing compounds include acetone oxime; violuric acid; pyridine-2-aldoxime; 2-aminophenol; 1,2-benzenediamine; 2,2,6,6-tetramethyl-1-piperidinyloxy; 5,6,7,8-tetrahydrobiopterin; 6,7-dimethyl-5,6,7,8-tetrahydropterine; and maleamic acid; or a salt or solvate thereof.

[0239] The quinone compound may be any suitable compound comprising a quinone moiety as described herein. Non-limiting examples of the quinone compounds include 1,4-benzoquinone; 1,4-naphthoquinone; 2-hydroxy-1,4-naphthoquinone; 2,3-dimethoxy-5-methyl-1,4-benzoquinone or coenzyme Q₀; 2,3,5,6-tetramethyl-1,4-benzoquinone or duroquinone; 1,4-dihydroxyanthraquinone; 3-hydroxy-1-methyl-5,6-indolinedione or adrenochrome; 4-tert-butyl-5-methoxy-1,2-benzoquinone; pyrroloquinoline quinone; or a salt or solvate thereof.

[0240] The sulfur-containing compound may be any suitable compound comprising one or more sulfur atoms. In one aspect, the sulfur-containing compound comprises a moiety selected from thionyl, thioether, sulfinyl, sulfonyl, sulfamide, sulfonamide, sulfonic acid, and sulfonic ester. Non-limiting examples of the sulfur-containing compounds include ethanethiol; 2-propanethiol; 2-propene-1-thiol; 2-mercaptoethanesulfonic acid; benzenethiol; benzene-1,2-dithiol; cysteine; methionine; glutathione; cystine; or a salt or solvate thereof.

[0241] In one aspect, an effective amount of such a compound described above to cellulosic material or xylan-containing material as a molar ratio to glucosyl units of cellulose is about 10^{-6} to about 10, e.g., about 10^{-6} to about 7.5, about 10^{-6} to about 5, about 10^{-6} to about 2.5, about 10^{-6} to about 1, about 10^{-5} to about 1, about 10^{-5} to about 10^{-1} , about 10^{-4} to about 10^{-1} , about 10^{-3} to about 10^{-1} , or about 10^{-3} to about 10^{-2} . In another aspect, an effective amount of such a compound described above is about 0.1 μ M to about 1 M, e.g., about 0.5 μ M to about 0.75 M, about 0.75 μ M to about 0.5 M, about 1 μ M to about 0.25 M, about 1 μ M to about 0.1 M, about 5 μ M to about 50 mM, about 10 μ M to about 25 mM, about 50 μ M to about 25 mM, about 10 μ M to about 10 mM, about 5 μ M to about 5 mM, or about 0.1 mM to about 1 mM.

[0242] The term "liquor" means the solution phase, either aqueous, organic, or a combination thereof, arising from treatment of a lignocellulose and/or hemicellulose material in a slurry, or monosaccharides thereof, e.g., xylose, arabinose, mannose, etc., under conditions as described herein, and the soluble contents thereof. A liquor for cellulolytic enhancement of a GH61 polypeptide can be produced by treating a lignocellulose or hemicellulose material (or feedstock) by applying heat and/or pressure, optionally in the presence of a catalyst, e.g., acid, optionally in the presence of an organic solvent, and optionally in combination with physical disruption of the material, and then separating the solution from the residual solids. Such conditions determine the degree of cellulolytic enhancement obtainable through the combination of liquor and a GH61 polypeptide during hydrolysis of a cellulosic substrate by a cellulase preparation. The liquor can be separated from the treated material using a method standard in the art, such as filtration, sedimentation, or centrifugation.

[0243] In one aspect, an effective amount of the liquor to cellulose is about 10^{-6} to about 10 g per g of cellulose, e.g., about 10^{-6} to about 7.5 g, about 10^{-6} to about 5, about 10^{-6} to about 2.5 g, about 10^{-6} to about 1 g, about 10^{-5} to about 1 g, about 10^{-5} to about 10^{-1} g, about 10^{-4} to about 10^{-1} g, about 10^{-3} to about 10^{-1} g, or about 10^{-3} to about 10^{-2} g per g of cellulose.

[0244] In one aspect, the one or more (e.g., several) hemicellulolytic enzymes comprise a commercial hemicellulolytic enzyme preparation. Examples of commercial hemicellulolytic enzyme preparations suitable for use in the present invention include, for example, SHEARZYME™ (Novozymes A/S), CELLIC™ HTec (Novozymes A/S), CELLIC™ HTec2 (Novozymes A/S), VISCOZYME® (Novozymes A/S), ULTRAFLO® (Novozymes A/S), PULPZYME® HC (Novozymes A/S), MULTIFECT® Xylanase (Genencor), ACCELLERASE® XY (Genencor), ACCELLERASE® XC (Genencor), ECOP-ULP® TX-200A (AB Enzymes), HSP 6000 Xylanase (DSM), DEPOL™ 333P (Biocatalysts Limit, Wales, UK), DEPOL™ 740L (Biocatalysts Limit, Wales, UK), and DEPOL™ 762P (Biocatalysts Limit, Wales, UK).

[0245] Examples of additional xylanases useful in the processes of the present invention include, but are not limited to, xylanases from *Aspergillus aculeatus* (GeneSeqP:AAR63790; WO 94/21785), *Aspergillus fumigatus* (WO 2006/078256), *Penicillium pinophilum* (WO 2011/041405), *Penicillium* sp. (WO 2010/126772), *Thielavia terrestris* NRRL 8126 (WO 2009/079210), and *Trichophaea saccata* GH10 (WO 2011/057083).

[0246] Examples of beta-xylosidases useful in the processes of the present invention include, but are not limited to, beta-xylosidases from *Neurospora crassa* (SwissProt accession number Q7SOW4), *Trichoderma reesei* (UniProtKB/TrEMBL accession number Q92458), and *Talaromyces emersonii* (SwissProt accession number Q8X212).

[0247] Examples of acetylxylan esterases useful in the processes of the present invention include, but are not limited to, acetylxylan esterases from *Aspergillus aculeatus* (WO 2010/108918), *Chaetomium globosum* (Uniprot accession number Q2GWX4), *Chaetomium*

gracile (GeneSeqP accession number AAB82124), *Humicola insolens* DSM 1800 (WO 2009/073709), *Hypocrea jecorina* (WO 2005/001036), *Myceliophthora thermophila* (WO 2010/014880), *Neurospora crassa* (UniProt accession number q7s259), *Phaeosphaeria nodorum* (Uniprot accession number Q0UJH1), and *Thielavia terrestris* NRRL 8126 (WO 2009/042846).

[0248] Examples of feruloyl esterases (ferulic acid esterases) useful in the processes of the present invention include, but are not limited to, feruloyl esterases from *Humicola insolens* DSM 1800 (WO 2009/076122), *Neosartorya fischeri* (UniProt Accession number A1D9T4), *Neurospora crassa* (UniProt accession number Q9HGR3), *Penicillium aurantiogriseum* (WO 2009/127729), and *Thielavia terrestris* (WO 2010/053838 and WO 2010/065448).

[0249] Examples of arabinofuranosidases useful in the processes of the present invention include, but are not limited to, arabinofuranosidases from *Aspergillus niger* (GeneSeqP accession number AAR94170), *Humicola insolens* DSM 1800 (WO 2006/114094 and WO 2009/073383), and *M. giganteus* (WO 2006/114094).

[0250] Examples of alpha-glucuronidases useful in the processes of the present invention include, but are not limited to, alpha-glucuronidases from *Aspergillus clavatus* (UniProt accession number alcc12), *Aspergillus fumigatus* (SwissProt accession number Q4WW45), *Aspergillus niger* (Uniprot accession number Q96WX9), *Aspergillus terreus* (SwissProt accession number Q0CJP9), *Humicola insolens* (WO 2010/014706), *Penicillium aurantiogriseum* (WO 2009/068565), *Talaromyces emersonii* (UniProt accession number Q8X211), and *Trichoderma reesei* (Uniprot accession number Q99024).

[0251] The polypeptides having enzyme activity used in the processes of the present invention may be produced by fermentation of the above-noted microbial strains on a nutrient medium containing suitable carbon and nitrogen sources and inorganic salts, using procedures known in the art (see, e.g., Bennett, J.W. and LaSure, L. (eds.), *More Gene Manipulations in Fungi*, Academic Press, CA, 1991). Suitable media are available from commercial suppliers or may be prepared according to published compositions (e.g., in catalogues of the American Type Culture Collection). Temperature ranges and other conditions suitable for growth and enzyme production are known in the art (see, e.g., Bailey, J.E., and Ollis, D.F., *Biochemical Engineering Fundamentals*, McGraw-Hill Book Company, NY, 1986).

[0252] The fermentation can be any method of cultivation of a cell resulting in the expression or isolation of an enzyme or protein. Fermentation may, therefore, be understood as comprising shake flask cultivation, or small- or large-scale fermentation (including continuous, batch, fed-batch, or solid state fermentations) in laboratory or industrial fermentors performed in a suitable medium and under conditions allowing the enzyme to be expressed or isolated. The resulting enzymes produced by the methods described above may be recovered from the fermentation medium and purified by conventional procedures.

[0253] Fermentation. The fermentable sugars obtained from the hydrolyzed cellulosic material

or xylan-containing material can be fermented by one or more (e.g., several) fermenting microorganisms capable of fermenting the sugars directly or indirectly into a desired fermentation product. "Fermentation" or "fermentation process" refers to any fermentation process or any process comprising a fermentation step. Fermentation processes also include fermentation processes used in the consumable alcohol industry (e.g., beer and wine), dairy industry (e.g., fermented dairy products), leather industry, and tobacco industry. The fermentation conditions depend on the desired fermentation product and fermenting organism and can easily be determined by one skilled in the art.

[0254] In the fermentation step, sugars, released from the cellulosic material or xylan-containing material as a result of the pretreatment and enzymatic hydrolysis steps, are fermented to a product, e.g., ethanol, by a fermenting organism, such as yeast. Hydrolysis (saccharification) and fermentation can be separate or simultaneous, as described herein.

[0255] Any suitable hydrolyzed cellulosic material or xylan-containing material can be used in the fermentation step in practicing the present invention. The material is generally selected based on the desired fermentation product, *i.e.*, the substance to be obtained from the fermentation, and the process employed, as is well known in the art.

[0256] The term "fermentation medium" is understood herein to refer to a medium before the fermenting microorganism(s) is(are) added, such as, a medium resulting from a saccharification process, as well as a medium used in a simultaneous saccharification and fermentation process (SSF).

[0257] "Fermenting microorganism" refers to any microorganism, including bacterial and fungal organisms, suitable for use in a desired fermentation process to produce a fermentation product. The fermenting organism can be hexose and/or pentose fermenting organisms, or a combination thereof. Both hexose and pentose fermenting organisms are well known in the art. Suitable fermenting microorganisms are able to ferment, *i.e.*, convert, sugars, such as glucose, xylose, xylulose, arabinose, maltose, mannose, galactose, and/or oligosaccharides, directly or indirectly into the desired fermentation product. Examples of bacterial and fungal fermenting organisms producing ethanol are described by Lin et al., 2006, Appl. Microbiol. Biotechnol. 69: 627-642.

[0258] Examples of fermenting microorganisms that can ferment hexose sugars include bacterial and fungal organisms, such as yeast. Preferred yeast includes strains of *Candida*, *Kluyveromyces*, and *Saccharomyces*, e.g., *Candida sonorensis*, *Kluyveromyces marxianus*, and *Saccharomyces cerevisiae*.

[0259] Examples of fermenting organisms that can ferment pentose sugars in their native state include bacterial and fungal organisms, such as some yeast. Preferred xylose fermenting yeast include strains of *Candida*, preferably *C. sheatae* or *C. sonorensis*; and strains of *Pichia*, preferably *P. stipitis*, such as *P. stipitis* CBS 5773. Preferred pentose fermenting yeast include strains of *Pachysolen*, preferably *P. tannophilus*. Organisms not capable of fermenting pentose

sugars, such as xylose and arabinose, may be genetically modified to do so by methods known in the art.

[0260] Examples of bacteria that can efficiently ferment hexose and pentose to ethanol include, for example, *Bacillus coagulans*, *Clostridium acetobutylicum*, *Clostridium thermocellum*, *Clostridium phytofermentans*, *Geobacillus* sp., *Thermoanaerobacter saccharolyticum*, and *Zymomonas mobilis* (Philippidis, 1996, *supra*).

[0261] Other fermenting organisms include strains of *Bacillus*, such as *Bacillus coagulans*; *Candida*, such as *C. sonorensis*, *C. methanosorbosa*, *C. diddensiae*, *C. parapsilosis*, *C. naedodendra*, *C. blankii*, *C. entomophila*, *C. brassicae*, *C. pseudotropicalis*, *C. boidinii*, *C. utilis*, and *C. scephatae*; *Clostridium*, such as *C. acetobutylicum*, *C. thermocellum*, and *C. phytofermentans*; *E. coli*, especially *E. coli* strains that have been genetically modified to improve the yield of ethanol; *Geobacillus* sp.; *Hansenula*, such as *Hansenula anomala*; *Klebsiella*, such as *K. oxytoca*; *Kluyveromyces*, such as *K. marxianus*, *K. lactis*, *K. thermotolerans*, and *K. fragilis*; *Schizosaccharomyces*, such as *S. pombe*; *Thermoanaerobacter*, such as *Thermoanaerobacter saccharolyticum*; and *Zymomonas*, such as *Zymomonas mobilis*.

[0262] In a preferred aspect, the yeast is a *Bretannomyces*. In a more preferred aspect, the yeast is *Bretannomyces clausenii*. In another preferred aspect, the yeast is a *Candida*. In another more preferred aspect, the yeast is *Candida sonorensis*. In another more preferred aspect, the yeast is *Candida boidinii*. In another more preferred aspect, the yeast is *Candida blankii*. In another more preferred aspect, the yeast is *Candida brassicae*. In another more preferred aspect, the yeast is *Candida diddensii*. In another more preferred aspect, the yeast is *Candida entomophiliia*. In another more preferred aspect, the yeast is *Candida pseudotropicalis*. In another more preferred aspect, the yeast is *Candida scephatae*. In another more preferred aspect, the yeast is *Candida utilis*. In another preferred aspect, the yeast is a *Clavispora*. In another more preferred aspect, the yeast is *Clavispora lusitaniae*. In another more preferred aspect, the yeast is *Clavispora opuntiae*. In another preferred aspect, the yeast is a *Kluyveromyces*. In another more preferred aspect, the yeast is *Kluyveromyces fragilis*. In another more preferred aspect, the yeast is *Kluyveromyces marxianus*. In another more preferred aspect, the yeast is *Kluyveromyces thermotolerans*. In another preferred aspect, the yeast is a *Pachysolen*. In another more preferred aspect, the yeast is *Pachysolen tannophilus*. In another preferred aspect, the yeast is a *Pichia*. In another more preferred aspect, the yeast is a *Pichia stipitis*. In another preferred aspect, the yeast is a *Saccharomyces* spp. In a more preferred aspect, the yeast is *Saccharomyces cerevisiae*. In another more preferred aspect, the yeast is *Saccharomyces distaticus*. In another more preferred aspect, the yeast is *Saccharomyces uvarum*.

[0263] In a preferred aspect, the bacterium is a *Bacillus*. In a more preferred aspect, the bacterium is *Bacillus coagulans*. In another preferred aspect, the bacterium is a *Clostridium*. In another more preferred aspect, the bacterium is *Clostridium acetobutylicum*. In another more preferred aspect, the bacterium is *Clostridium phytofermentans*. In another more preferred

aspect, the bacterium is *Clostridium thermocellum*. In another more preferred aspect, the bacterium is *Geobacillus* sp. In another more preferred aspect, the bacterium is a *Thermoanaerobacter*. In another more preferred aspect, the bacterium is *Thermoanaerobacter saccharolyticum*. In another preferred aspect, the bacterium is a *Zymomonas*. In another more preferred aspect, the bacterium is *Zymomonas mobilis*.

[0264] Commercially available yeast suitable for ethanol production include, e.g., BIOFERM™ AFT and XR (NABC - North American Bioproducts Corporation, GA, USA), ETHANOL RED™ yeast (Fermentis/Lesaffre, USA), FALI™ (Fleischmann's Yeast, USA), FERMIOL™ (DSM Specialties), GERT STRAND™ (Gert Strand AB, Sweden), and SUPER-START™ and THERMOSACC™ fresh yeast (Ethanol Technology, WI, USA).

[0265] In a preferred aspect, the fermenting microorganism has been genetically modified to provide the ability to ferment pentose sugars, such as xylose utilizing, arabinose utilizing, and xylose and arabinose co-utilizing microorganisms.

[0266] The cloning of heterologous genes into various fermenting microorganisms has led to the construction of organisms capable of converting hexoses and pentoses to ethanol (co-fermentation) (Chen and Ho, 1993, Cloning and improving the expression of *Pichia stipitis* xylose reductase gene in *Saccharomyces cerevisiae*, Appl. Biochem. Biotechnol. 39-40: 135-147; Ho et al., 1998, Genetically engineered *Saccharomyces* yeast capable of effectively cofermenting glucose and xylose, Appl. Environ. Microbiol. 64: 1852-1859; Kotter and Ciriacy, 1993, Xylose fermentation by *Saccharomyces cerevisiae*, Appl. Microbiol. Biotechnol. 38: 776-783; Walfridsson et al., 1995, Xylose-metabolizing *Saccharomyces cerevisiae* strains overexpressing the TKL1 and TAL1 genes encoding the pentose phosphate pathway enzymes transketolase and transaldolase, Appl. Environ. Microbiol. 61:4184-4190; Kuyper et al., 2004, Minimal metabolic engineering of *Saccharomyces cerevisiae* for efficient anaerobic xylose fermentation: a proof of principle, FEMS Yeast Research 4: 655-664; Beall et al., 1991, Parametric studies of ethanol production from xylose and other sugars by recombinant *Escherichia coli*, Biotech. Bioeng. 38: 296-303; Ingram et al., 1998, Metabolic engineering of bacteria for ethanol production, Biotechnol. Bioeng. 58: 204-214; Zhang et al., 1995, Metabolic engineering of a pentose metabolism pathway in ethanologenic *Zymomonas mobilis*, Science 267: 240-243; Deanda et al., 1996, Development of an arabinose-fermenting *Zymomonas mobilis* strain by metabolic pathway engineering, Appl. Environ. Microbiol. 62: 4465-4470; WO 2003/062430, xylose isomerase).

[0267] In a preferred aspect, the genetically modified fermenting microorganism is *Candida sonorensis*. In another preferred aspect, the genetically modified fermenting microorganism is *Escherichia coli*. In another preferred aspect, the genetically modified fermenting microorganism is *Klebsiella oxytoca*. In another preferred aspect, the genetically modified fermenting microorganism is *Kluyveromyces marxianus*. In another preferred aspect, the genetically modified fermenting microorganism is *Saccharomyces cerevisiae*. In another preferred aspect, the genetically modified fermenting microorganism is *Zymomonas mobilis*.

[0268] It is well known in the art that the organisms described above can also be used to produce other substances, as described herein.

[0269] The fermenting microorganism is typically added to the degraded cellulosic material or xylan-containing material or hydrolysate and the fermentation is performed for about 8 to about 96 hours, *e.g.*, about 24 to about 60 hours. The temperature is typically between about 26°C to about 60°C, *e.g.*, about 32°C or 50°C, and about pH 3 to about pH 8, *e.g.*, pH 4-5, 6, or 7.

[0270] In one aspect, the yeast and/or another microorganism are applied to the degraded cellulosic material or xylan-containing material and the fermentation is performed for about 12 to about 96 hours, such as typically 24-60 hours. In another aspect, the temperature is preferably between about 20°C to about 60°C, *e.g.*, about 25°C to about 50°C, about 32°C to about 50°C, or about 32°C to about 50°C, and the pH is generally from about pH 3 to about pH 7, *e.g.*, about pH 4 to about pH 7. However, some fermenting organisms, *e.g.*, bacteria, have higher fermentation temperature optima. Yeast or another microorganism is preferably applied in amounts of approximately 10^5 to 10^{12} , preferably from approximately 10^7 to 10^{10} , especially approximately 2×10^8 viable cell count per ml of fermentation broth. Further guidance in respect of using yeast for fermentation can be found in, *e.g.*, "The Alcohol Textbook" (Editors K. Jacques, T.P. Lyons and D.R. Kelsall, Nottingham University Press, United Kingdom 1999), which is hereby incorporated by reference.

[0271] For ethanol production, following the fermentation the fermented slurry is distilled to extract the ethanol. The ethanol obtained according to the processes of the invention can be used as, *e.g.*, fuel ethanol, drinking ethanol, *i.e.*, potable neutral spirits, or industrial ethanol.

[0272] A fermentation stimulator can be used in combination with any of the processes described herein to further improve the fermentation process, and in particular, the performance of the fermenting microorganism, such as, rate enhancement and ethanol yield. A "fermentation stimulator" refers to stimulators for growth of the fermenting microorganisms, in particular, yeast. Preferred fermentation stimulators for growth include vitamins and minerals. Examples of vitamins include multivitamins, biotin, pantothenate, nicotinic acid, meso-inositol, thiamine, pyridoxine, para-aminobenzoic acid, folic acid, riboflavin, and Vitamins A, B, C, D, and E. See, for example, Alfenore et al., Improving ethanol production and viability of *Saccharomyces cerevisiae* by a vitamin feeding strategy during fed-batch process, Springer-Verlag (2002), which is hereby incorporated by reference. Examples of minerals include minerals and mineral salts that can supply nutrients comprising P, K, Mg, S, Ca, Fe, Zn, Mn, and Cu.

[0273] Fermentation products: A fermentation product can be any substance derived from the fermentation. The fermentation product can be, without limitation, an alcohol (*e.g.*, arabinitol, n-butanol, isobutanol, ethanol, glycerol, methanol, ethylene glycol, 1,3-propanediol [propylene glycol], butanediol, glycerin, sorbitol, and xylitol); an alkane (*e.g.*, pentane, hexane, heptane, octane, nonane, decane, undecane, and dodecane), a cycloalkane (*e.g.*, cyclopentane,

cyclohexane, cycloheptane, and cyclooctane), an alkene (e.g. pentene, hexene, heptene, and octene); an amino acid (e.g., aspartic acid, glutamic acid, glycine, lysine, serine, and threonine); a gas (e.g., methane, hydrogen (H₂), carbon dioxide (CO₂), and carbon monoxide (CO)); isoprene; a ketone (e.g., acetone); an organic acid (e.g., acetic acid, acetonc acid, adipic acid, ascorbic acid, citric acid, 2,5-diketo-D-gluconic acid, formic acid, fumaric acid, glucaric acid, gluconic acid, glucuronic acid, glutaric acid, 3-hydroxypropionic acid, itaconic acid, lactic acid, malic acid, malonic acid, oxalic acid, oxaloacetic acid, propionic acid, succinic acid, and xylonic acid); and polyketide. The fermentation product can also be protein as a high value product.

[0274] In a preferred aspect, the fermentation product is an alcohol. It will be understood that the term "alcohol" encompasses a substance that contains one or more hydroxyl moieties. In a more preferred aspect, the alcohol is n-butanol. In another more preferred aspect, the alcohol is isobutanol. In another more preferred aspect, the alcohol is ethanol. In another more preferred aspect, the alcohol is methanol. In another more preferred aspect, the alcohol is arabinitol. In another more preferred aspect, the alcohol is butanediol. In another more preferred aspect, the alcohol is ethylene glycol. In another more preferred aspect, the alcohol is glycerin. In another more preferred aspect, the alcohol is glycerol. In another more preferred aspect, the alcohol is 1,3-propanediol. In another more preferred aspect, the alcohol is sorbitol. In another more preferred aspect, the alcohol is xylitol. See, for example, Gong, C. S., Cao, N. J., Du, J., and Tsao, G. T., 1999, Ethanol production from renewable resources, in *Advances in Biochemical Engineering/Biotechnology*, Scheper, T., ed., Springer-Verlag Berlin Heidelberg, Germany, 65: 207-241; Silveira, M. M., and Jonas, R., 2002, The biotechnological production of sorbitol, *Appl. Microbiol. Biotechnol.* 59: 400-408; Nigam and Singh, 1995, Processes for fermentative production of xylitol - a sugar substitute, *Process Biochemistry* 30(2): 117-124; Ezeji et al., 2003, Production of acetone, butanol and ethanol by *Clostridium beijerinckii* BA101 and in situ recovery by gas stripping, *World Journal of Microbiology and Biotechnology* 19(6): 595-603.

[0275] In another preferred aspect, the fermentation product is an alkane. The alkane can be an unbranched or a branched alkane. In another more preferred aspect, the alkane is pentane. In another more preferred aspect, the alkane is hexane. In another more preferred aspect, the alkane is heptane. In another more preferred aspect, the alkane is octane. In another more preferred aspect, the alkane is nonane. In another more preferred aspect, the alkane is decane. In another more preferred aspect, the alkane is undecane. In another more preferred aspect, the alkane is dodecane.

[0276] In another preferred aspect, the fermentation product is a cycloalkane. In another more preferred aspect, the cycloalkane is cyclopentane. In another more preferred aspect, the cycloalkane is cyclohexane. In another more preferred aspect, the cycloalkane is cycloheptane. In another more preferred aspect, the cycloalkane is cyclooctane.

[0277] In another preferred aspect, the fermentation product is an alkene. The alkene can be an unbranched or a branched alkene. In another more preferred aspect, the alkene is

pentene. In another more preferred aspect, the alkene is hexene. In another more preferred aspect, the alkene is heptene. In another more preferred aspect, the alkene is octene.

[0278] In another preferred aspect, the fermentation product is an amino acid. In another more preferred aspect, the organic acid is aspartic acid. In another more preferred aspect, the amino acid is glutamic acid. In another more preferred aspect, the amino acid is glycine. In another more preferred aspect, the amino acid is lysine. In another more preferred aspect, the amino acid is serine. In another more preferred aspect, the amino acid is threonine. See, for example, Richard and Margaritis, 2004, Empirical modeling of batch fermentation kinetics for poly(glutamic acid) production and other microbial biopolymers, *Biotechnology and Bioengineering* 87(4): 501-515.

[0279] In another preferred aspect, the fermentation product is a gas. In another more preferred aspect, the gas is methane. In another more preferred aspect, the gas is H₂. In another more preferred aspect, the gas is CO₂. In another more preferred aspect, the gas is CO. See, for example, Kataoka et al., 1997, Studies on hydrogen production by continuous culture system of hydrogen-producing anaerobic bacteria, *Water Science and Technology* 36(6-7): 41-47; and Gunaseelan, 1997, Anaerobic digestion of biomass for methane production: A review, *Biomass and Bioenergy*. 13(1-2): 83-114.

[0280] In another preferred aspect, the fermentation product is isoprene.

[0281] In another preferred aspect, the fermentation product is a ketone. It will be understood that the term "ketone" encompasses a substance that contains one or more ketone moieties. In another more preferred aspect, the ketone is acetone. See, for example, Qureshi and Blaschek, 2003, *supra*.

[0282] In another preferred aspect, the fermentation product is an organic acid. In another more preferred aspect, the organic acid is acetic acid. In another more preferred aspect, the organic acid is acetonic acid. In another more preferred aspect, the organic acid is adipic acid. In another more preferred aspect, the organic acid is ascorbic acid. In another more preferred aspect, the organic acid is citric acid. In another more preferred aspect, the organic acid is 2,5-diketo-D-gluconic acid. In another more preferred aspect, the organic acid is formic acid. In another more preferred aspect, the organic acid is fumaric acid. In another more preferred aspect, the organic acid is glucaric acid. In another more preferred aspect, the organic acid is gluconic acid. In another more preferred aspect, the organic acid is glucuronic acid. In another more preferred aspect, the organic acid is glutaric acid. In another preferred aspect, the organic acid is 3-hydroxypropionic acid. In another more preferred aspect, the organic acid is itaconic acid. In another more preferred aspect, the organic acid is lactic acid. In another more preferred aspect, the organic acid is malic acid. In another more preferred aspect, the organic acid is malonic acid. In another more preferred aspect, the organic acid is oxalic acid. In another more preferred aspect, the organic acid is propionic acid. In another more preferred aspect, the organic acid is succinic acid. In another more preferred aspect, the organic acid is xylonic acid. See, for example, Chen and Lee, 1997, Membrane-mediated extractive

fermentation for lactic acid production from cellulosic biomass, Appl. Biochem. Biotechnol. 63-65: 435-448.

[0283] In another preferred aspect, the fermentation product is polyketide.

[0284] Recovery. The fermentation product(s) can be optionally recovered from the fermentation medium using any method known in the art including, but not limited to, chromatography, electrophoretic procedures, differential solubility, distillation, or extraction. For example, alcohol is separated from the fermented cellulosic material or xylan-containing material and purified by conventional methods of distillation. Ethanol with a purity of up to about 96 vol.% can be obtained, which can be used as, for example, fuel ethanol, drinking ethanol, *i.e.*, potable neutral spirits, or industrial ethanol.

Signal Peptide

[0285] Also described is an isolated polynucleotide encoding a signal peptide comprising or consisting of amino acids 1 to 17 of SEQ ID NO: 2, amino acids 1 to 16 of SEQ ID NO: 4, or amino acids 1 to 20 of SEQ ID NO: 6. The polynucleotide may further comprise a gene encoding a protein, which is operably linked to the signal peptide. The protein is preferably foreign to the signal peptide. In one aspect, the polynucleotide encoding the signal peptide is nucleotides 1 to 51 of SEQ ID NO: 1. Alternatively, the polynucleotide encoding the signal peptide is nucleotides 1 to 48 of SEQ ID NO: 3, or the polynucleotide encoding the signal peptide is nucleotides 1 to 50, and 114 to 123 of SEQ ID NO: 5. Also described are nucleic acid constructs, expression vectors and recombinant host cells comprising such polynucleotides. Also described are methods of producing a protein, comprising (a) cultivating a recombinant host cell comprising such polynucleotide; and (b) recovering the protein.

[0286] The protein may be native or heterologous to a host cell. The term "protein" is not meant herein to refer to a specific length of the encoded product and, therefore, encompasses peptides, oligopeptides, and polypeptides. The term "protein" also encompasses two or more polypeptides combined to form the encoded product. The proteins also include hybrid polypeptides and fused polypeptides.

[0287] Preferably, the protein is a hormone, enzyme, receptor or portion thereof, antibody or portion thereof, or reporter. For example, the protein may be a hydrolase, isomerase, ligase, lyase, oxidoreductase, or transferase, *e.g.*, an aminopeptidase, amylase, carbohydrase, carboxypeptidase, catalase, cellobiohydrolase, cellulase, chitinase, cutinase, cyclodextrin glycosyltransferase, deoxyribonuclease, endoglucanase, esterase, alpha-galactosidase, beta-galactosidase, glucoamylase, alpha-glucosidase, beta-glucosidase, invertase, laccase, lipase, mannosidase, mutanase, oxidase, pectinolytic enzyme, peroxidase, phytase, polyphenoloxidase, proteolytic enzyme, ribonuclease, transglutaminase, xylanase, or beta-xylosidase.

[0288] The gene may be obtained from any prokaryotic, eukaryotic, or other source.

[0289] The present invention is further described by the following examples that should not be construed as limiting the scope of the invention.

Examples

Strains

[0290] *Talaromyces leycettanus* Strain CBS398.68 was used as the source of a polypeptide having xylanase activity. *Aspergillus oryzae* MT3568 strain was used for expression of the *Talaromyces leycettanus* gene encoding the polypeptide having xylanase activity. *A. oryzae* MT3568 is an *amdS* (acetamidase) disrupted gene derivative of *Aspergillus oryzae* JaL355 (WO 02/40694) in which *pyrG* auxotrophy was restored by disrupting the *A. oryzae* acetamidase (*amdS*) gene.

Media and Solutions

[0291] YP+2% glucose medium was composed of 1% yeast extract, 2% peptone and 2% glucose.

[0292] PDA agar plates were composed of potato infusion (potato infusion was made by boiling 300 g of sliced (washed but unpeeled) potatoes in water for 30 minutes and then decanting or straining the broth through cheesecloth. Distilled water was then added until the total volume of the suspension was one liter, followed by 20 g of dextrose and 20 g of agar powder. The medium was sterilized by autoclaving at 15 psi for 15 minutes (Bacteriological Analytical Manual, 8th Edition, Revision A, 1998).

[0293] LB plates were composed of 10 g of Bacto-Tryptone, 5 g of yeast extract, 10 g of sodium chloride, 15 g of Bacto-agar, and deionized water to 1 liter. The medium was sterilized by autoclaving at 15 psi for 15 minutes (Bacteriological Analytical Manual, 8th Edition, Revision A, 1998).

[0294] COVE sucrose plates were composed of 342 g Sucrose (Sigma S-9378), 20 g Agar powder, 20 ml Cove salt solution (26 g $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 26 g KCL, 26 g KH_2PO_4 , 50 ml Cove trace metal solution) and deionized water to 1 liter), and deionized water to 1 liter). The medium was sterilized by autoclaving at 15 psi for 15 minutes (Bacteriological Analytical Manual, 8th Edition, Revision A, 1998). The medium was cooled to 60°C and added 10 mM acetamide, 15 mM CsCl, Triton X-100 (50 μl /500 ml)).

[0295] Cove trace metal solution was composed of 0.04 g $\text{Na}_2\text{B}_4\text{O}_7 \cdot 10\text{H}_2\text{O}$, 0.4 g $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$, 1.2 g $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$, 0.7 g $\text{MnSO}_4 \cdot \text{H}_2\text{O}$, 0.8 g $\text{Na}_2\text{MoO}_4 \cdot 2\text{H}_2\text{O}$, 10 g $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$, and deionized water to 1 liter.

[0296] Dap-4C medium was composed of 20 g Dextrose, 10 g Maltose, 11 g $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 1 g KH_2PO_4 , 2 g Citric Acid, 5.2 g $\text{K}_3\text{PO}_4 \cdot \text{H}_2\text{O}$, 0.5 g Yeast Extract (Difco), 1 ml Dowfax 63N10 (Dow Chemical Company), 0.5 ml KU6 trace metals solution, 2.5 g CaCO_3 , and deionized water to 1 liter. The medium was sterilized by autoclaving at 15 psi for 15 minutes (Bacteriological Analytical Manual, 8th Edition, Revision A, 1998). Before use, Dap-4C medium was added 3.5 ml sterile 50 % $(\text{NH}_4)_2\text{HPO}_4$ and 5 ml sterile 20 % Lactic Acid per 150 ml medium.

[0297] KU6 trace metals solution was composed of 0.13 g NiCl_2 , 2.5 g $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$, 13.9 g $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$, 8.45 g $\text{MnSO}_4 \cdot \text{H}_2\text{O}$, 6.8 g ZnCl_2 , 3g Citric Acid, and deionized water to 1 liter.

Example 1: Source of DNA sequence information for *Talaromyces leycettanus* Strain CBS398.68

[0298] Genomic sequence information was generated by Illumina DNA sequencing at the Beijing Genome Institute (BGI) in Beijing, China from genomic DNA isolated from *Talaromyces leycettanus* Strain CBS398.68. A preliminary assembly of the genome was analyzed using the Pedant-Pro™ Sequence Analysis Suite (Biomax Informatics AG, Martinsried, Germany). Gene models constructed by the software were used as a starting point for detecting GH10 homologues in the genome. More precise gene models were constructed manually using multiple known GH10 protein sequences as a guide.

Example 2: *Talaromyces leycettanus* Strain CBS398.68 genomic DNA extraction

[0299] To generate genomic DNA for PCR amplification, *Talaromyces leycettanus* Strain CBS398.68 was propagated on PDA agar plates by growing at 26°C for 7 days. Spores harvested from the PDA plates were used to inoculate 25 ml of YP+2% glucose medium in a baffled shake flask and incubated at 26°C for 72 hours with agitation at 85 rpm.

[0300] Genomic DNA was isolated according to a modified DNeasy Plant Maxi kit protocol (Qiagen Denmark, Copenhagen, Denmark). The fungal material from the above culture was harvested by centrifugation at 14,000 x g for 2 minutes. The supernatant was removed and the 0.5 g of the pellet was frozen in liquid nitrogen with quartz sand and grinded to a fine powder in a pre-chilled mortar. The powder was transferred to a 15 ml centrifuge tube and added 5 ml buffer AP1 (preheated to 65°C) and 10 µl RNase A stock solution (100 mg/ml) followed by vigorous vortexing. After incubation for 10 minutes at 65°C with regular inverting of the tube,

1.8 ml buffer AP2 was added to the lysate by gentle mixing followed by incubation on ice for 10 min. The lysate was then centrifuged at 3000 x g for 5 minutes at room temperature and the supernatant was decanted into a QIAshredder maxi spin column placed in a 50 ml collection tube. This was followed by centrifugation at 3000 x g for 5 minutes at room temperature. The flow-through was transferred into a new 50 ml tube and added 1.5 volumes of buffer AP3/E followed by vortexing. 15 ml of the sample was transferred into a DNeasy Maxi spin column placed in a 50 ml collection tube and centrifuged at 3000 x g for 5 minutes at room temperature. The flow-through was discarded and 12 ml buffer AW was added to the DNeasy Maxi spin column placed in a 50 ml collection tube and centrifuged at 3000 x g for 10 minutes at room temperature. After discarding the flow-through, centrifugation was repeated to dispose of the remaining alcohol. The DNeasy Maxi spin column was transferred to a new 50 ml tube and 0.5 ml buffer AE (preheated to 70°C) was added. After incubation for 5 minutes at room temperature, the sample was eluted by centrifugation at 3000 x g for 5 minutes at room temperature. Elution was repeated with an additional 0.5 ml buffer AE and the eluates were combined. The concentration of the harvested DNA was measured by a UV spectrophotometer at 260 nm.

Example 3: Construction of an *Aspergillus oryzae* expression vector containing *Talaromyces leycettanus* Strain CBS398.68 genomic sequence encoding a Family GH10 polypeptide having xylanase activity

[0301] Two synthetic oligonucleotide primers shown below were designed to PCR amplify the *Talaromyces leycettanus* Strain CBS398.68 P24F5Z gene (SEQ ID NO: 1) from the genomic DNA prepared in Example 2. An IN-FUSION™ Cloning Kit (BD Biosciences, Palo Alto, CA, USA) was used to clone the fragment directly into the expression vector pDau109 (WO 2005/042735).

F-P24F5Z 5'- ACACAACTGGGGATCCACCATGGTCCATCTTTCTTCCCTGGCC -3' (SEQ ID NO: 7)

R-P24F5Z 5'- CCCTCTAGATCTCGAGTTACAGGCACTGGTAGTAGTAGGGATTTC -3' (SEQ ID NO: 8)

Bold letters represent gene sequence. The underlined sequence is homologous to the insertion sites of pDau109.

[0302] An MJ Research PTC-200 DNA engine was used to perform the PCR reaction. A Phusion® High-Fidelity PCR Kit (Finnzymes Oy, Espoo, Finland) was used for the PCR amplification. The PCR reaction was composed of 5 µl of 5X HF buffer (Finnzymes Oy, Espoo, Finland), 0.5 µl of dNTPs (10 mM), 0.5 µl of Phusion® DNA polymerase (0.2 units/µl) (Finnzymes Oy, Espoo, Finland), 1 µl of primer F-P24F5Z (5 µM), 1 µl of primer R-P24F5Z (5 µM), 0.5 µl of *Talaromyces leycettanus* genomic DNA (100 ng/µl), and 16.5 µl of deionized water in a total volume of 25 µl. The PCR conditions were 1 cycle at 95°C for 2 minutes. 35

cycles each at 98°C for 10 seconds, 60°C for 30 seconds, and 72°C for 2.5 minutes; and 1 cycle at 72°C for 10 minutes. The sample was then held at 12°C until removed from the PCR machine.

[0303] The reaction products were isolated by 1.0% agarose gel electrophoresis using 40 mM Tris base, 20 mM sodium acetate, 1 mM disodium EDTA (TAE) buffer where a 1513 bp product band was excised from the gel and purified using an illustra GFX[®] PCR DNA and Gel Band Purification Kit (GE Healthcare Life Sciences, Brondby, Denmark) according to the manufacturer's instructions. The fragment was then cloned into *Bam* HI and *Xho* I digested pDau109 using an IN-FUSION[™] Cloning Kit resulting in plasmid pP24F5Z. Cloning of the P24F5Z gene into *Bam* HI-*Xho* I digested pDau109 resulted in the transcription of the *Talaromyces leycettanus* P24F5Z gene under the control of a NA2-tpi double promoter. NA2-tpi is a modified promoter from the gene encoding the *Aspergillus niger* neutral alpha-amylase in which the untranslated leader has been replaced by an untranslated leader from the gene encoding the *Aspergillus nidulans* triose phosphate isomerase.

[0304] The cloning protocol was performed according to the IN-FUSION[™] Cloning Kit instructions generating a P24F5Z GH10 construct. The treated plasmid and insert were transformed into One Shot[®] TOP10F' Chemically Competent *E. coli* cells (Invitrogen, Carlsbad, CA, USA) according to the manufacturer's protocol and plated onto LB plates supplemented with 0.1 mg of ampicillin per ml. After incubating at 37°C overnight, colonies were seen growing under selection on the LB ampicillin plates. Four colonies transformed with the P24F5Z GH10 construct were cultivated in LB medium supplemented with 0.1 mg of ampicillin per ml and plasmid was isolated with a QIAprep Spin Miniprep Kit (QIAGEN Inc., Valencia, CA, USA) according to the manufacturer's protocol.

[0305] Isolated plasmids were sequenced with vector primers and P24F5Z gene specific primers in order to determine a representative plasmid expression clone that was free of PCR errors.

Example 4: Characterization of the *Talaromyces leycettanus* CBS398.68 genomic sequence encoding a P24F5Z GH10 polypeptide having xylanase activity

[0306] DNA sequencing of the *Talaromyces leycettanus* CBS398.68 P24F5Z GH10 genomic clone was performed with an Applied Biosystems Model 3700 Automated DNA Sequencer using version 3.1 BIG-DYE[™] terminator chemistry (Applied Biosystems, Inc., Foster City, CA, USA) and primer walking strategy. Nucleotide sequence data were scrutinized for quality and all sequences were compared to each other with assistance of PHRED/PHRAP software (University of Washington, Seattle, WA, USA). The sequence obtained was identical to the sequence from the JGI.

[0307] The nucleotide sequence and deduced amino acid sequence of the *Talaromyces*

leycettanus P24F5Z gene is shown in SEQ ID NO: 1 and SEQ ID NO: 2, respectively. The coding sequence is 1268 bp including the stop codon and is interrupted by one intron. The encoded predicted protein is 364 amino acids. Using the SignalP program (Nielsen et al., 1997, Protein Engineering 10: 1-6), a signal peptide of 17 residues was predicted. The predicted mature protein contains 347 amino acids with a predicted molecular mass of 38.7 kDa and an isoelectric pH of 4.6.

[0308] A comparative pairwise global alignment of amino acid sequences was determined using the Needleman and Wunsch algorithm (Needleman and Wunsch, 1970, J. Mol. Biol. 48: 443-453) with gap open penalty of 10, gap extension penalty of 0.5, and the EBLOSUM62 matrix. The alignment showed that the deduced amino acid sequence of the *Talaromyces leycettanus* gene encoding the P24F5Z GH10 polypeptide having xylanase activity shares 75.6% identity (excluding gaps) to the deduced amino acid sequence of a predicted GH10 family protein from *Penicillium canescens* (accession number UNIPROT:C3VEV9) with xylanase activity.

Example 5: Expression of the *Talaromyces leycettanus* GH10 xylanase P24F5Z

[0309] The expression plasmid pP24F5Z was transformed into *Aspergillus oryzae* MT3568. *Aspergillus oryzae* MT3568 is an AMDS (acetamidase) disrupted derivative of JaL355 (WO 02/40694) in which *pyrG* auxotrophy was restored in the process of knocking out the *A. oryzae* acetamidase (AMDS) gene. MT3568 protoplasts are prepared according to the method of European Patent No. 0238023, pages 14-15, which are incorporated herein by reference.

[0310] Transformants were purified on COVE sucrose selection plates through single conidia prior to sporulating them on PDA plates. Production of the *Talaromyces leycettanus* GH10 polypeptide by the transformants was analyzed from culture supernatants of 1 ml 96 deep well stationary cultivations at 30°C in YP+2% glucose medium. Expression was verified on a E-Page 8% SDS-PAGE 48 well gel (Invitrogen, Carlsbad, CA, USA) by Coomassie staining. One transformant was selected for further work and designated *Aspergillus oryzae* 41.4.3.

[0311] For larger scale production, *Aspergillus oryzae* 41.4.3 spores were spread onto a PDA plate and incubated for five days at 37°C. The confluent spore plate was washed twice with 5 ml of 0.01% TWEEN® 20 to maximize the number of spores collected. The spore suspension was then used to inoculate twenty five 500 ml flasks containing 100 ml of Dap-4C medium. The culture was incubated at 30°C with constant shaking at 100 rpm. At day four post-inoculation, the culture broth was collected by filtration through a bottle top MF75 Supor MachV 0.2 µm PES filter (Thermo Fisher Scientific, Roskilde, Denmark). Fresh culture broth from this transformant produced a band of GH10 protein of approximately 50 kDa with smearing indicating possible glycosylation. The identity of the prominent band as the *Talaromyces leycettanus* GH10 polypeptide was verified by peptide sequencing.

Example 6: Alternative method for producing the *Talaromyces leycettanus* GH10

xylanase P24F5Z

[0312] Based on the nucleotide sequence identified as SEQ ID NO: 1, a synthetic gene can be obtained from a number of vendors such as Gene Art (GENEART AG BioPark, Josef-Engert-Str. 11, 93053, Regensburg, Germany) or DNA 2.0 (DNA2.0, 1430 O'Brien Drive, Suite E, Menlo Park, CA 94025, USA). The synthetic gene can be designed to incorporate additional DNA sequences such as restriction sites or homologous recombination regions to facilitate cloning into an expression vector.

[0313] Using the two synthetic oligonucleotide primers F-P24F5Z and R-P24F5Z described above, a simple PCR reaction can be used to amplify the full-length open reading frame from the synthetic gene of SEQ ID NO: 1. The gene can then be cloned into an expression vector for example as described above and expressed in a host cell, for example in *Aspergillus oryzae* as described above.

Example 7: Purification of the *Talaromyces leycettanus* GH10 xylanase P24F5Z

[0314] 1000 ml broth of the *Aspergillus oryzae* expression strain 41.4.3 was adjusted to pH 7.0 and filtrated on 0.22 µm PES filter (Thermo Fisher Scientific, Roskilde, Denmark). Following, the filtrate was added 1.8 M ammonium sulphate. The filtrate was loaded onto a Phenyl Sepharose™ 6 Fast Flow column (high sub) (GE Healthcare, Piscataway, NJ, USA) (with a column volume of 60 mL) equilibrated with 1.8 M ammonium sulphate pH 7.0, 25 mM HEPES pH7.0. After application the column was washed with 3 column volumes of equilibration buffer followed by 7 column volumes of 1.0 M ammonium sulphate (the protein kept binding to the column) and the protein eluted following with 5 column volumes of 25 mM HEPES pH 7.0 at a flow rate of 15 ml/min. Fractions of 10 mL were collected and analyzed by SDS-page. The fractions were pooled and applied to a Sephadex™ G-25 (medium) (GE Healthcare, Piscataway, NJ, USA) column equilibrated in 25 mM HEPES pH 7.0. The fractions were applied to a SOURCE™ 15Q (GE Healthcare, Piscataway, NJ, USA) column equilibrated in 25 mM HEPES pH 7.0 (column volumes 60 mL). After application the column was washed with 5 column volumes equilibration buffer and bound proteins were eluted with a linear gradient over 20 column volumes from 0-1000 mM sodium chloride. Fractions of 10 ml were collected and analyzed by SDS-page, and fractions with the protein were pooled. With two distinct peaks in the chromatogram two pools, A and B, were made. The protein concentration was determined by A280/A260 absorbance. The protein identity was verified by MS/MS on in-gel digested sample confirming the identity of both fractions.

Example 8: Construction of an *Aspergillus oryzae* expression vector containing *Talaromyces leycettanus* Strain CBS398.68 genomic sequence encoding a Family GH10 polypeptide having xylanase activity

[0315] Two synthetic oligonucleotide primers shown below were designed to PCR amplify the *Talaromyces leycettanus* Strain CBS398.68 P24F61 gene (SEQ ID NO: 3) from the genomic DNA prepared in Example 2. An IN-FUSION™ Cloning Kit (BD Biosciences, Palo Alto, CA, USA) was used to clone the fragment directly into the expression vector pDau109 (WO 2005/042735).

F-P24F61 5'- ACACAACTGGGGATCCACCATGGTCCGTCTTTCCGCTGGA -3' (SEQ ID NO: 9)

R-P24F61 5'- CCCTCTAGATCTCGAGTTACAAGCACTGGGAGTACCACTGG -3' (SEQ ID NO: 10)

Bold letters represent gene sequence. The underlined sequence is homologous to the insertion sites of pDau109.

[0316] An MJ Research PTC-200 DNA engine was used to perform the PCR reaction. A Phusion® High-Fidelity PCR Kit (Finnzymes Oy, Espoo, Finland) was used for the PCR amplification. The PCR reaction was composed of 5 µl of 5X HF buffer (Finnzymes Oy, Espoo, Finland), 0.5 µl of dNTPs (10 mM), 0.5 µl of Phusion® DNA polymerase (0.2 units/µl) (Finnzymes Oy, Espoo, Finland), 1 µl of primer F-P24F61 (5 µM), 1 µl of primer R-P24F61 (5 µM), 0.5 µl of *Talaromyces leycettanus* genomic DNA (100 ng/µl), and 16.5 µl of deionized water in a total volume of 25 µl. The PCR conditions were 1 cycle at 95°C for 2 minutes, 35 cycles each at 98°C for 10 seconds, 60°C for 30 seconds, and 72°C for 2.5 minutes; and 1 cycle at 72°C for 10 minutes. The sample was then held at 12°C until removed from the PCR machine.

[0317] The reaction products were isolated by 1.0% agarose gel electrophoresis using 40 mM Tris base, 20 mM sodium acetate, 1 mM disodium EDTA (TAE) buffer where a 1513 bp product band was excised from the gel and purified using an illustra GFX® PCR DNA and Gel Band Purification Kit (GE Healthcare Life Sciences, Brøndby, Denmark) according to the manufacturer's instructions. The fragment was then cloned into *Bam* HI and *Xho* I digested pDau109 using an IN-FUSION™ Cloning Kit resulting in plasmid pP24F61. Cloning of the P24F61 gene into *Bam* HI-*Xho* I digested pDau109 resulted in the transcription of the *Talaromyces leycettanus* P24F61 gene under the control of a NA2-tpi double promoter. NA2-tpi is a modified promoter from the gene encoding the *Aspergillus niger* neutral alpha-amylase in which the untranslated leader has been replaced by an untranslated leader from the gene encoding the *Aspergillus nidulans* triose phosphate isomerase.

[0318] The cloning protocol was performed according to the IN-FUSION™ Cloning Kit instructions generating a P24F61 GH10 construct. The treated plasmid and insert were transformed into One Shot® TOP10F' Chemically Competent *E. coli* cells (Invitrogen, Carlsbad,

CA, USA) according to the manufacturer's protocol and plated onto LB plates supplemented with 0.1 mg of ampicillin per ml. After incubating at 37°C overnight, colonies were seen growing under selection on the LB ampicillin plates. Four colonies transformed with the P24F61 GH10 construct were cultivated in LB medium supplemented with 0.1 mg of ampicillin per ml and plasmid was isolated with a QIAprep Spin Miniprep Kit (QIAGEN Inc., Valencia, CA, USA) according to the manufacturer's protocol.

[0319] Isolated plasmids were sequenced with vector primers and P24F61 gene specific primers in order to determine a representative plasmid expression clone that was free of PCR errors.

Example 9: Characterization of the *Talaromyces leycettanus* CBS398.68 genomic sequence encoding a P24F61 GH10 polypeptide having xylanase activity

[0320] DNA sequencing of the *Talaromyces leycettanus* CBS398.68 P24F61 GH10 genomic clone was performed with an Applied Biosystems Model 3700 Automated DNA Sequencer using version 3.1 BIG-DYE™ terminator chemistry (Applied Biosystems, Inc., Foster City, CA, USA) and primer walking strategy. Nucleotide sequence data were scrutinized for quality and all sequences were compared to each other with assistance of PHRED/PHRAP software (University of Washington, Seattle, WA, USA). The sequence obtained was identical to the sequence from the JGI.

[0321] The nucleotide sequence and deduced amino acid sequence of the *Talaromyces leycettanus* P24F61 gene is shown in SEQ ID NO: 3 and SEQ ID NO: 4, respectively. The coding sequence is 1708 bp including the stop codon and is interrupted by eight introns. The encoded predicted protein is 389 amino acids. Using the SignalP program (Nielsen et al., 1997, Protein Engineering 10: 1-6), a signal peptide of 16 residues was predicted. The predicted mature protein contains 373 amino acids with a predicted molecular mass of 39.6 kDa and an isoelectric pH of 5.2.

[0322] A comparative pairwise global alignment of amino acid sequences was determined using the Needleman and Wunsch algorithm (Needleman and Wunsch, 1970, J. Mol. Biol. 48: 443-453) with gap open penalty of 10, gap extension penalty of 0.5, and the EBLOSUM62 matrix. The alignment showed that the deduced amino acid sequence of the *Talaromyces leycettanus* gene encoding the P24F61 GH10 polypeptide having xylanase activity shares 76.0% identity (excluding gaps) to the deduced amino acid sequence of a predicted GH10 family protein from *Talaromyces stipitatus* (accession number SWISSPROT:B8M9H8) with putative xylanase activity.

Example 10: Expression of the *Talaromyces leycettanus* GH10 xylanase P24F61

[0323] The expression plasmid pP24F61 was transformed into *Aspergillus oryzae* MT3568. *Aspergillus oryzae* MT3568 is an AMDS (acetamidase) disrupted derivative of JaL355 (WO 02/40694) in which pyrG auxotrophy was restored in the process of knocking out the *A. oryzae* acetamidase (AMDS) gene. MT3568 protoplasts are prepared according to the method of European Patent No. 0238023, pages 14-15, which are incorporated herein by reference.

[0324] Transformants were purified on COVE sucrose selection plates through single conidia prior to sporulating them on PDA plates. Production of the *Talaromyces leycettanus* GH10 polypeptide by the transformants was analyzed from culture supernatants of 1 ml 96 deep well stationary cultivations at 30°C in YP+2% glucose medium. Expression was verified on a E-Page 8% SDS-PAGE 48 well gel (Invitrogen, Carlsbad, CA, USA) by Coomassie staining. One transformant was selected for further work and designated *Aspergillus oryzae* 40.2.3.

[0325] For larger scale production, *Aspergillus oryzae* 40.2.3 spores were spread onto a PDA plate and incubated for five days at 37°C. The confluent spore plate was washed twice with 5 ml of 0.01% TWEEN® 20 to maximize the number of spores collected. The spore suspension was then used to inoculate twenty five 500 ml flasks containing 100 ml of Dap-4C medium. The culture was incubated at 30°C with constant shaking at 100 rpm. At day four post-inoculation, the culture broth was collected by filtration through a bottle top MF75 Supor MachV 0.2 µm PES filter (Thermo Fisher Scientific, Roskilde, Denmark). Fresh culture broth from this transformant produced a band of GH10 protein of approximately 40 kDa. The identity of this band as the *Talaromyces leycettanus* GH10 polypeptide was verified by peptide sequencing.

Example 11: Alternative method for producing the *Talaromyces leycettanus* GH10 xylanase P24F61

[0326] Based on the nucleotide sequence identified as SEQ ID NO: 3, a synthetic gene can be obtained from a number of vendors such as Gene Art (GENEART AG BioPark, Josef-Engert-Str. 11, 93053, Regensburg, Germany) or DNA 2.0 (DNA2.0, 1430 O'Brien Drive, Suite E, Menlo Park, CA 94025, USA). The synthetic gene can be designed to incorporate additional DNA sequences such as restriction sites or homologous recombination regions to facilitate cloning into an expression vector.

[0327] Using the two synthetic oligonucleotide primers F-P24F61 and R-P24F61 described above, a simple PCR reaction can be used to amplify the full-length open reading frame from the synthetic gene of SEQ ID NO: 3. The gene can then be cloned into an expression vector for example as described above and expressed in a host cell, for example in *Aspergillus oryzae* as described above.

Example 12: Purification of the *Talaromyces leycettanus* GH10 xylanase P24F61

[0328] 1000 ml broth of the *Aspergillus oryzae* expression strain 40.2.3 was adjusted to pH 7.0 and filtrated on 0.22 µm PES filter (Thermo Fisher Scientific, Roskilde, Denmark). Following, the filtrate was added 1.4 M ammonium sulphate. The filtrate was loaded onto a Phenyl Sepharose™ 6 Fast Flow column (high sub) (GE Healthcare, Piscataway, NJ, USA) (with a column volume of 60 mL) equilibrated with 1.4 M ammonium sulphate pH 7.0, 25 mM HEPES pH7.0. After application the column was washed with 3 column volumes of equilibration buffer followed by 7 column volumes of 0.8 M ammonium sulphate (the protein kept binding to the column) and the protein eluted following with 5 column volumes of 25 mM HEPES pH 7.0 at a flow rate of 15 ml/min. Fractions of 10 mL were collected and analyzed by SDS-page. The fractions were pooled and applied to a Sephadex™ G-25 (medium) (GE Healthcare, Piscataway, NJ, USA) column equilibrated in 25 mM HEPES pH 7.0. The fractions were applied to a SOURCE™ 15Q (GE Healthcare, Piscataway, NJ, USA) column equilibrated in 25 mM HEPES pH 7.0 (column volumes 60 mL). After application the column was washed with 5 column volumes equilibration buffer and bound proteins were eluted with a linear gradient over 20 column volumes from 0-1000 mM sodium chloride. Fractions of 10 ml were collected and analyzed by SDS-page. The enzyme was in the run through and in the first fractions and pooled accordingly. With two distinct peaks in the chromatogram two pools were made. The enzyme was concentrated using a centrifugal concentrator Vivaspın®20 MWCO 10,000 polyethersulfone membrane (Sartorius Stedim Biotech GmbH, 37070 Goettingen, Germany). The protein concentration was determined by A280/A260 absorbance. The protein identity was verified by MS/MS on in-gel digested sample.

Example 13: Construction of an *Aspergillus oryzae* expression vector containing *Talaromyces leycettanus* Strain CBS398.68 genomic sequence encoding a Family GH10 polypeptide having xylanase activity

[0329] Two synthetic oligonucleotide primers shown below were designed to PCR amplify the *Talaromyces leycettanus* Strain CBS398.68 P24F62 gene (SEQ ID NO: 5) from the genomic DNA prepared in Example 2. An IN-FUSION™ Cloning Kit (BD Biosciences, Palo Alto, CA, USA) was used to clone the fragment directly into the expression vector pDau109 (WO 2005/042735).

F-P24F62 5'- ACACAACTGGGGATCCACCATGCGTTTCTCCTTGGCCACTG-3' (SEQ ID NO: 11)

R-P24F62 5'- CCCTCTAGATCTCGAGCTAGCAGACGCTGCAGGCCT -3' (SEQ ID NO: 12)

Bold letters represent gene sequence. The underlined sequence is homologous to the insertion sites of pDau109.

[0330] An MJ Research PTC-200 DNA engine was used to perform the PCR reaction. A Phusion® High-Fidelity PCR Kit (Finnzymes Oy, Espoo, Finland) was used for the PCR

amplification. The PCR reaction was composed of 5 µl of 5X HF buffer (Finnzymes Oy, Espoo, Finland), 0.5 µl of dNTPs (10 mM), 0.5 µl of Phusion® DNA polymerase (0.2 units/µl) (Finnzymes Oy, Espoo, Finland), 1 µl of primer F-P24F62 (5 µM), 1 µl of primer R-P24F62 (5 µM), 0.5 µl of *Talaromyces leycettanus* genomic DNA (100 ng/µl), and 16.5 µl of deionized water in a total volume of 25 µl. The PCR conditions were 1 cycle at 95°C for 2 minutes. 35 cycles each at 98°C for 10 seconds, 60°C for 30 seconds, and 72°C for 2.5 minutes; and 1 cycle at 72°C for 10 minutes. The sample was then held at 12°C until removed from the PCR machine.

[0331] The reaction products were isolated by 1.0% agarose gel electrophoresis using 40 mM Tris base, 20 mM sodium acetate, 1 mM disodium EDTA (TAE) buffer where a 1513 bp product band was excised from the gel and purified using an illustra GFX® PCR DNA and Gel Band Purification Kit (GE Healthcare Life Sciences, Brøndby, Denmark) according to the manufacturer's instructions. The fragment was then cloned into *Bam* HI and *Xho* I digested pDau109 using an IN-FUSION™ Cloning Kit resulting in plasmid pP24F62. Cloning of the P24F62 gene into *Bam* HI-*Xho* I digested pDau109 resulted in the transcription of the *Talaromyces leycettanus* P24F62 gene under the control of a NA2-tpi double promoter. NA2-tpi is a modified promoter from the gene encoding the *Aspergillus niger* neutral alpha-amylase in which the untranslated leader has been replaced by an untranslated leader from the gene encoding the *Aspergillus nidulans* triose phosphate isomerase.

[0332] The cloning protocol was performed according to the IN-FUSION™ Cloning Kit instructions generating a P24F62 GH10 construct. The treated plasmid and insert were transformed into One Shot® TOP10F' Chemically Competent *E. coli* cells (Invitrogen, Carlsbad, CA, USA) according to the manufacturer's protocol and plated onto LB plates supplemented with 0.1 mg of ampicillin per ml. After incubating at 37°C overnight, colonies were seen growing under selection on the LB ampicillin plates. Four colonies transformed with the P24F62 GH10 construct were cultivated in LB medium supplemented with 0.1 mg of ampicillin per ml and plasmid was isolated with a QIAprep Spin Miniprep Kit (QIAGEN Inc., Valencia, CA, USA) according to the manufacturer's protocol.

[0333] Isolated plasmids were sequenced with vector primers and P24F62 gene specific primers in order to determine a representative plasmid expression clone that was free of PCR errors.

Example 14: Characterization of the *Talaromyces leycettanus* CBS398.68 genomic sequence encoding a P24F62 GH10 polypeptide having xylanase activity

[0334] DNA sequencing of the *Talaromyces leycettanus* CBS398.68 P24F62 GH10 genomic clone was performed with an Applied Biosystems Model 3700 Automated DNA Sequencer using version 3.1 BIG-DYE™ terminator chemistry (Applied Biosystems, Inc., Foster City, CA,

USA) and primer walking strategy. Nucleotide sequence data were scrutinized for quality and all sequences were compared to each other with assistance of PHRED/PHRAP software (University of Washington, Seattle, WA, USA). The sequence obtained was identical to the sequence from the JGI.

[0335] The nucleotide sequence and deduced amino acid sequence of the *Talaromyces leycettanus* P24F62 gene is shown in SEQ ID NO: 5 and SEQ ID NO: 6, respectively. The coding sequence is 1520 bp including the stop codon and is interrupted by four introns. The encoded predicted protein is 405 amino acids. Using the SignalP program (Nielsen et al., 1997, Protein Engineering 10: 1-6), a signal peptide of 20 residues was predicted. The predicted mature protein contains 385 amino acids with a predicted molecular mass of 41.6 kDa and an isoelectric pH of 4.7.

[0336] A comparative pairwise global alignment of amino acid sequences was determined using the Needleman and Wunsch algorithm (Needleman and Wunsch, 1970, J. Mol. Biol. 48: 443-453) with gap open penalty of 10, gap extension penalty of 0.5, and the EBLOSUM62 matrix. The alignment showed that the deduced amino acid sequence of the *Talaromyces leycettanus* gene encoding the P24F62 GH10 polypeptide having xylanase activity shares 83.0% identity (excluding gaps) to the deduced amino acid sequence of a predicted GH10 family protein from *Penicillium* sp. (accession number GENESEQP:AYL61291) with xylanase activity.

Example 15: Expression of the *Talaromyces leycettanus* GH10 xylanase P24F62

[0337] The expression plasmid pP24F62 was transformed into *Aspergillus oryzae* MT3568. *Aspergillus oryzae* MT3568 is an AMDS (acetamidase) disrupted derivative of JaL355 (WO 02/40694) in which pyrG auxotrophy was restored in the process of knocking out the A, *oryzae* acetamidase (AMDS) gene. MT3568 protoplasts are prepared according to the method of European Patent No. 0238023, pages 14-15, which are incorporated herein by reference.

[0338] Transformants were purified on COVE sucrose selection plates through single conidia prior to sporulating them on PDA plates. Production of the *Talaromyces leycettanus* GH10 polypeptide by the transformants was analyzed from culture supernatants of 1 ml 96 deep well stationary cultivations at 30°C in YP+2% glucose medium. Expression was verified on a E-Page 8% SDS-PAGE 48 well gel (Invitrogen, Carlsbad, CA, USA) by Coomassie staining. One transformant was selected for further work and designated *Aspergillus oryzae* 39.3.1.

[0339] For larger scale production, *Aspergillus oryzae* 39.3.1 spores were spread onto a PDA plate and incubated for five days at 37°C. The confluent spore plate was washed twice with 5 ml of 0.01% TWEEN® 20 to maximize the number of spores collected. The spore suspension was then used to inoculate twenty five 500 ml flasks containing 100 ml of Dap-4C medium. The culture was incubated at 30°C with constant shaking at 100 rpm. At day four post-inoculation,

the culture broth was collected by filtration through a bottle top MF75 Supor MachV 0.2 µm PES filter (Thermo Fisher Scientific, Roskilde, Denmark). Fresh culture broth from this transformant produced a band of GH10 protein of approximately 50 kDa. The identity of this band as the *Talaromyces leycettanus* GH10 polypeptide was verified by peptide sequencing.

Example 16: Alternative method for producing the *Talaromyces leycettanus* GH10 xylanase P24F62

[0340] Based on the nucleotide sequence identified as SEQ ID NO: 5, a synthetic gene can be obtained from a number of vendors such as Gene Art (GENEART AG BioPark, Josef-Engert-Str. 11, 93053, Regensburg, Germany) or DNA 2.0 (DNA2.0, 1430 O'Brien Drive, Suite E, Menlo Park, CA 94025, USA). The synthetic gene can be designed to incorporate additional DNA sequences such as restriction sites or homologous recombination regions to facilitate cloning into an expression vector.

[0341] Using the two synthetic oligonucleotide primers F-P24F62 and R-P24F62 described above, a simple PCR reaction can be used to amplify the full-length open reading frame from the synthetic gene of SEQ ID NO: 5. The gene can then be cloned into an expression vector for example as described above and expressed in a host cell, for example in *Aspergillus oryzae* as described above.

Example 17: Purification of the *Talaromyces leycettanus* GH10 xylanase P24F62

[0342] 1000 ml broth of the *Aspergillus oryzae* expression strain 39.3.1 was adjusted to pH 7.0 and filtrated on 0.22 µm PES filter (Thermo Fisher Scientific, Roskilde, Denmark). Following, the filtrate was added 1.4 M ammonium sulphate. The filtrate was loaded onto a Phenyl Sepharose™ 6 Fast Flow column (high sub) (GE Healthcare, Piscataway, NJ, USA) (with a column volume of 60 mL) equilibrated with 1.4 M ammonium sulphate pH 7.0, 25 mM HEPES pH7.0. After application the column was washed with 3 column volumes of equilibration buffer followed by 7 column volumes of 0.8 M ammonium sulphate (the protein kept binding to the column) and the protein eluted following with 5 column volumes of 25 mM HEPES pH 7.0 at a flow rate of 15 ml/min. Fractions of 10 mL were collected and analyzed by SDS-page. The fractions were pooled and applied to a Sephadex™ G-25 (medium) (GE Healthcare, Piscataway, NJ, USA) column equilibrated in 25 mM HEPES pH 7.0. The fractions were applied to a SOURCE™ 15Q (GE Healthcare, Piscataway, NJ, USA) column equilibrated in 25 mM HEPES pH 7.0 (column volumes 60 mL). After application the column was washed with 5 column volumes equilibration buffer and bound proteins were eluted with a linear gradient over 20 column volumes from 0-1000 mM sodium chloride. Fractions of 10 ml were collected and analyzed by SDS-page. The enzyme was in the run through and in the first fractions and pooled accordingly. With two distinct peaks in the chromatogram two pools were made. The

enzyme was concentrated using a centrifugal concentrator Vivaspin®20 MWCO 10,000 polyethersulfone membrane (Sartorius Stedim Biotech GmbH, 37070 Goettingen, Germany). The protein concentration was determined by A280/A260 absorbance. The protein identity was verified by MS/MS on in-gel digested sample.

Example 18: Determination of xylanase activity of the xylanases according to the invention

Enzyme activity of the *Talaromyces leycettanus* GH10 xylanase P24F5Z

[0343] Pool A and pool B of the purified xylanase were diluted in distilled water with 0.01% Triton X-100 (100 ppm) based on a dose-response curve to be in the linear range. The substrate was AZCL-arabinoxylan (Megazyme Wicklow, Ireland) in 0.2 % (w/v) in pH 6.0 in (50 mM phosphoric acid, 50 mM acetic acid, 50 mM boric acid), 50 mM KCl, 1 mM CaCl₂, 0.01% Triton X-100; pH adjusted with NaOH. The substrate was equilibrated to 37°C. 1000 µl 0.2 % (w/v) AZCL-arabinoxylan was mixed with 20 µl diluted enzyme. The tube was incubated at 37°C for 15 minutes on an Eppendorf Comfort thermomixer (Eppendorf AG, Hamburg, Germany) at 1.400 rpm. The reaction was stopped by placing the tube on ice for 5 minutes. The eppendorf tube was centrifuged 5 minutes with 10,000 rpm at 4°C. 200 microliter supernatant is transferred to a flat bottom MicroWell plate (NUNC, Roskilde, Denmark) and the absorbance was read at 595nm in a spectrophotometer. Relative to Shearzyme (Novozymes, Bagsvaerd, Denmark), the specific activity on AZCL-arabinoxylan was found to be 2 % for pool A and 4 % for pool B.

Enzyme activity of the *Talaromyces leycettanus* GH10 xylanase P24F61

[0344] The purified xylanase was diluted in distilled water with 0.01% Triton X-100 (100 ppm) based on a dose-response curve to be in the linear range. The substrate was AZCL-arabinoxylan (Megazyme Wicklow, Ireland) in 0.2 % (w/v) in pH 6.0 in (50 mM phosphoric acid, 50 mM acetic acid, 50 mM boric acid), 50 mM KCl, 1 mM CaCl₂, 0.01% Triton X-100; pH adjusted with NaOH. The substrate was equilibrated to 37°C. 1000 µl 0.2 % (w/v) AZCL-arabinoxylan was mixed with 20 µl diluted enzyme. The tube was incubated at 37°C for 15 minutes on an Eppendorf Comfort thermomixer (Eppendorf AG, Hamburg, Germany) at 1.400 rpm. The reaction was stopped by placing the tube on ice for 5 minutes. The eppendorf tube was centrifuged 5 minutes with 10,000 rpm at 4°C. 200 microliter supernatant is transferred to a flat bottom MicroWell plate (NUNC, Roskilde, Denmark) and the absorbance was read at 595nm in a spectrophotometer. Relative to Shearzyme (Novozymes, Bagsvaerd, Denmark), the specific activity on AZCL-arabinoxylan was found to be 830 % for the purified xylanase.

Enzyme activity of the *Talaromyces leycettanus* GH10 xylanase P24F62

[0345] The purified xylanase was diluted in distilled water with 0.01% Triton X-100 (100 ppm) based on a dose-response curve to be in the linear range. The substrate was AZCL-arabinoxylan (Megazyme Wicklow, Ireland) in 0.2 % (w/v) in pH 6.0 in (50 mM phosphoric acid, 50 mM acetic acid, 50 mM boric acid), 50 mM KCl, 1 mM CaCl₂, 0.01% Triton X-100; pH adjusted with NaOH. The substrate was equilibrated to 37°C. 1000 µl 0.2 % (w/v) AZCL-arabinoxylan was mixed with 20 µl diluted enzyme. The tube was incubated at 37°C for 15 minutes on an Eppendorf Comfort thermomixer (Eppendorf AG, Hamburg, Germany) at 1.400 rpm. The reaction was stopped by placing the tube on ice for 5 minutes. The eppendorf tube was centrifuged 5 minutes with 10,000 rpm at 4°C. 200 microliter supernatant is transferred to a flat bottom MicroWell plate (NUNC, Roskilde, Denmark) and the absorbance was read at 595nm in a spectrophotometer. Relative to Shearzyme (Novozymes, Bagsvaerd, Denmark), the specific activity on AZCL-arabinoxylan was found to be 950 % for the purified xylanase. In the case of conflict, the present disclosure including definitions will control. Also described are:

1. [1] An isolated polypeptide having xylanase activity, selected from the group consisting of:
 1. (a) a polypeptide having at least 77%, *e.g.*, at least 78%, at least 79%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to the mature polypeptide of SEQ ID NO: 2, or a polypeptide having at least 77%, *e.g.*, at least 78%, at least 79%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to the mature polypeptide of SEQ ID NO: 4, or a polypeptide having at least 85%, *e.g.*, at least 86%, at least 87%, at least 88%, at least 89%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to the mature polypeptide of SEQ ID NO: 6;
 2. (b) a polypeptide encoded by a polynucleotide that hybridizes under low stringency conditions, or medium stringency conditions, or medium-high stringency conditions, or high stringency conditions, or very high stringency conditions with (i) the mature polypeptide coding sequence of SEQ ID NO: 1, SEQ ID NO: 3, or SEQ ID NO: 5, (ii) the cDNA sequence thereof, or (iii) the full-length complement of (i) or (ii);
 3. (c) a polypeptide encoded by a polynucleotide having at least 60%, *e.g.*, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to the mature polypeptide coding sequence of SEQ ID NO: 1, SEQ ID NO: 3, or SEQ ID NO: 5; or the cDNA sequence thereof;
 4. (d) a variant of the mature polypeptide of SEQ ID NO: 2, SEQ ID NO: 4, or SEQ ID NO: 6 comprising a substitution, deletion, and/or insertion at one or more

positions; and

5. (e) a fragment of the polypeptide of (a), (b), (c), or (d) that has xylanase activity.
2. [2] The polypeptide of paragraph 1, having at least 77%, at least 78%, at least 79%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to the mature polypeptide of SEQ ID NO: 2; having at least 77%, at least 78%, at least 79%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to the mature polypeptide of SEQ ID NO: 4; or having at least 85%, at least 86%, at least 87%, at least 88%, at least 89%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% sequence identity to the mature polypeptide of SEQ ID NO: 6.
3. [3] The polypeptide of paragraph 1 or 2, which is encoded by a polynucleotide that hybridizes under low stringency conditions, or low-medium stringency conditions, or medium stringency conditions, or medium-high stringency conditions, or high stringency conditions, or very high stringency conditions with (i) the mature polypeptide coding sequence of SEQ ID NO: 1, SEQ ID NO: 3, or SEQ ID NO: 5, (ii) the cDNA sequence thereof, or (iii) the full-length complement of (i) or (ii).
4. [4] The polypeptide of any of paragraphs 1-3, which is encoded by a polynucleotide having at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99% or 100% sequence identity to the mature polypeptide coding sequence of SEQ ID NO: 1, SEQ ID NO: 3, or SEQ ID NO: 5 or the cDNA sequence thereof.
5. [5] The polypeptide of any of paragraphs 1-4, comprising or consisting of SEQ ID NO: 2, SEQ ID NO: 4, or SEQ ID NO: 6 or the mature polypeptide of SEQ ID NO: 2, SEQ ID NO: 4, or SEQ ID NO: 6.
6. [6] The polypeptide of paragraph 5, wherein the mature polypeptide is amino acids 18 to 364 of SEQ ID NO: 2, amino acids 17 to 389 of SEQ ID NO: 4, or amino acids 21 to 405 of SEQ ID NO: 6.
7. [7] The polypeptide of any of paragraphs 1-4, which is a variant of the mature polypeptide of SEQ ID NO: 2, SEQ ID NO: 4, or SEQ ID NO: 6 comprising a substitution, deletion, and/or insertion at one or more positions.
8. [8] The polypeptide of paragraph 1, which is a fragment of SEQ ID NO: 2, SEQ ID NO: 4, or SEQ ID NO: 6, wherein the fragment has xylanase activity.
9. [9] An isolated polypeptide comprising a catalytic domain selected from the group consisting of:
 1. (a) a catalytic domain having at least 77% sequence identity to the catalytic domain of SEQ ID NO: 2 or SEQ ID NO: 4, or a catalytic domain having xylanase activity and at least 90% sequence identity to the catalytic domain of SEQ ID NO: 6;
 2. (b) a catalytic domain encoded by a polynucleotide having at least 60% sequence identity to the catalytic domain coding sequence of SEQ ID NO: 1, SEQ ID NO: 3,

- or SEQ ID NO: 5;
3. (c) a variant of a catalytic domain comprising a substitution, deletion, and/or insertion of one or more (several) amino acids of the catalytic domain of SEQ ID NO: 2, SEQ ID NO: 4, or SEQ ID NO: 6; and
 4. (d) a fragment of a catalytic domain of (a), (b), or (c), which has xylanase activity.
10. [10] The polypeptide of paragraph 9, comprising or consisting of the catalytic domain of SEQ ID NO: 2, SEQ ID NO: 4, or SEQ ID NO: 6.
 11. [11] The polypeptide of paragraph 10, wherein the catalytic domain is amino acids 18 to 364 of SEQ ID NO: 2, amino acids 17 to 326 of SEQ ID NO: 4, or amino acids 21 to 337 of SEQ ID NO: 6.
 12. [12] The polypeptide of any of paragraphs 9-11, further comprising a cellulose binding domain.
 13. [13] The polypeptide of any of paragraphs 1-12, which is encoded by the polynucleotide contained in *Talaromyces leycettanus* Strain CBS398.68.
 14. [14] A composition comprising the polypeptide of any of paragraphs 1-13.
 15. [15] An isolated polynucleotide encoding the polypeptide of any of paragraphs 1-13.
 16. [16] A nucleic acid construct or expression vector comprising the polynucleotide of paragraph 15 operably linked to one or more control sequences that direct the production of the polypeptide in an expression host.
 17. [17] A recombinant host cell comprising the polynucleotide of paragraph 15 operably linked to one or more control sequences that direct the production of the polypeptide.
 18. [18] A method of producing the polypeptide of any of paragraphs 1-13, comprising:
 1. (a) cultivating a cell, which in its wild-type form produces the polypeptide, under conditions conducive for production of the polypeptide; and
 2. (b) recovering the polypeptide.
 19. [19] A method of producing a polypeptide having xylanase activity, comprising:
 1. (a) cultivating the host cell of paragraph 17 under conditions conducive for production of the polypeptide; and
 2. (b) recovering the polypeptide.
 20. [20] A transgenic plant, plant part or plant cell comprising a polynucleotide encoding the polypeptide of any of paragraphs 1-13.
 21. [21] A method of producing a polypeptide having xylanase activity, comprising:
 1. (a) cultivating the transgenic plant or plant cell of paragraph 20 under conditions conducive for production of the polypeptide; and
 2. (b) recovering the polypeptide.
 22. [22] An isolated polynucleotide encoding a signal peptide comprising or consisting of amino acids 1 to 17 of SEQ ID NO: 2, amino acids 1 to 16 of SEQ ID NO: 4, or amino acids 1 to 20 of SEQ ID NO: 6.
 23. [23] A nucleic acid construct or expression vector comprising a gene encoding a protein operably linked to the polynucleotide of paragraph 22, wherein the gene is foreign to the polynucleotide encoding the signal peptide.
 24. [24] A recombinant host cell comprising a gene encoding a protein operably linked to the polynucleotide of paragraph 22, wherein the gene is foreign to the polynucleotide encoding the signal peptide.

25. [25] A method of producing a protein, comprising:
 1. (a) cultivating a recombinant host cell comprising a gene encoding a protein operably linked to the polynucleotide of paragraph 22, wherein the gene is foreign to the polynucleotide encoding the signal peptide, under conditions conducive for production of the protein; and
 2. (b) recovering the protein.
26. [26] A process for degrading a cellulosic material or xylan-containing material, comprising: treating the cellulosic material or xylan-containing material with an enzyme composition in the presence of the polypeptide having xylanase activity of any of paragraphs 1-13.
27. [27] The process of paragraph 26, wherein the cellulosic material or xylan-containing material is pretreated.
28. [28] The process of paragraph 26 or 27, wherein the enzyme composition comprises one or more enzymes selected from the group consisting of a cellulase, a GH61 polypeptide having cellulolytic enhancing activity, a hemicellulase, an esterase, an expansin, a laccase, a ligninolytic enzyme, a pectinase, a peroxidase, a protease, and a swollenin.
29. [29] The process of paragraph 28, wherein the hemicellulase is one or more enzymes selected from the group consisting of a xylanase, an acetylxylan esterase, a feruloyl esterase, an arabinofuranosidase, a xylosidase, and a glucuronidase.
30. [30] The process of paragraph 28, wherein the cellulase is one or more enzymes selected from the group consisting of an endoglucanase, a cellobiohydrolase, and a beta-glucosidase.
31. [31] The process of any of paragraphs 26-30, further comprising recovering the degraded cellulosic material or xylan-containing material.
32. [32] The process of paragraph 31, wherein the degraded cellulosic material or xylan-containing material is a sugar.
33. [33] The process of paragraph 32, wherein the sugar is selected from the group consisting of glucose, xylose, mannose, galactose, and arabinose.
34. [34] A process for producing a fermentation product, comprising:
 1. (a) saccharifying a cellulosic material or xylan-containing material with an enzyme composition in the presence of the polypeptide having xylanase activity of any of paragraphs 1-13;
 2. (b) fermenting the saccharified cellulosic material or xylan-containing material with one or more fermenting microorganisms to produce the fermentation product; and
 3. (c) recovering the fermentation product from the fermentation.
35. [35] The process of paragraph 34, wherein the cellulosic material or xylan-containing material is pretreated.
36. [36] The process of paragraph 34 or 35, wherein the enzyme composition comprises one or more enzymes selected from the group consisting of a cellulase, a GH61 polypeptide having cellulolytic enhancing activity, a hemicellulase, an esterase, an expansin, a laccase, a ligninolytic enzyme, a pectinase, a peroxidase, a protease, and a swollenin.
37. [37] The process of paragraph 36, wherein the hemicellulase is one or more enzymes

- selected from the group consisting of a xylanase, an acetylxylan esterase, a feruloyl esterase, an arabinofuranosidase, a xylosidase, and a glucuronidase.
38. [38] The process of paragraph 36, wherein the cellulase is one or more enzymes selected from the group consisting of an endoglucanase, a cellobiohydrolase, and a beta-glucosidase.
 39. [39] The process of any of paragraphs 34-38, wherein steps (a) and (b) are performed simultaneously in a simultaneous saccharification and fermentation.
 40. [40] The process of any of paragraphs 34-39, wherein the fermentation product is an alcohol, an alkane, a cycloalkane, an alkene, an amino acid, a gas, isoprene, a ketone, an organic acid, or polyketide.
 41. [41] A process of fermenting a cellulosic material or xylan-containing material, comprising: fermenting the cellulosic material or xylan-containing material with one or more fermenting microorganisms, wherein the cellulosic material or xylan-containing material is saccharified with an enzyme composition in the presence of the polypeptide having xylanase activity of any of paragraphs 1-13.
 42. [42] The process of paragraph 41, wherein the fermenting of the cellulosic material or xylan-containing material produces a fermentation product.
 43. [43] The process of paragraph 42, further comprising recovering the fermentation product from the fermentation.
 44. [44] The process of any of paragraphs 41-43, wherein the cellulosic material or xylan-containing material is pretreated before saccharification.
 45. [45] The process of any of paragraphs 41-44, wherein the enzyme composition comprises one or more enzymes selected from the group consisting of a cellulase, a GH61 polypeptide having cellulolytic enhancing activity, a hemicellulase, an esterase, an expansin, a laccase, a ligninolytic enzyme, a pectinase, a peroxidase, a protease, and a swollenin.
 46. [46] The process of paragraph 45, wherein the hemicellulase is one or more enzymes selected from the group consisting of a xylanase, an acetylxylan esterase, a feruloyl esterase, an arabinofuranosidase, a xylosidase, and a glucuronidase.
 47. [47] The process of paragraph 45, wherein the cellulase is one or more enzymes selected from the group consisting of an endoglucanase, a cellobiohydrolase, and a beta-glucosidase.
 48. [48] The process of any of paragraphs 41-47, wherein the fermentation product is an alcohol, an alkane, a cycloalkane, an alkene, an amino acid, a gas, isoprene, a ketone, an organic acid, or polyketide.

SEQUENCE MISTING

[0346]

<110> Novozymes A/S

Spodsberg, Nikolaj

<120> Polypeptides having xylanase activity and polynucleotides encoding same

<130> 12262-WO-PCT

<160> 12

<170> PatentIn version 3.5

<210> 1

<211> 1168

<212> DNA

<213> *Talaromyces leycettanus*

<400> 1

```

atgCGtttct ccttggccac tgcagctctt ctCGctggcc ctgccctggc agCGccacca      60
gctcctCGtc acgatgacaa ggatgttggg ctcaacgccc tggcccagag agcaggcaag      120
ctctggttcg gcactgctgc tgatatcccc ggcaccgacg agacgaccga tgctgcgtac      180
ctaaaaatct tggaaaatcc cgccaacttc ggcgagatca cccctgccaa cgccatgaag      240
gtacggagct gcttgacaag tcggaggatc atgcctacga gcgaacaaac ttccgatgct      300
gacgcattga cagttcatgt acaccgagcc agagcagaac gtgttcaact acaccggcgg      360
tgactacgtc ctgaacctcg ccgagcgcca cggccagcgt gtccgctgcc acaacctcgt      420
ctgggccagc cagctgtccg acttcgtcaa caacggcaac tggaccaagg agtccctcac      480
ggcCGtgatg cggaaccaca tcttcacgt cgtccagcac ttCGccggc gctgctactc      540
gtgggatgtc gtcaacgagg ccctcaacgg cgcggcacc ttctcctcca gcatctggta      600
cgacaccatc ggcgaggact acttctacct cgccttccag tacgcccagg aggccctcgc      660
ggagatccac gccaacgacg tcaagctcta ctacaacgac tacggcatcg agaaccCCg      720
caccaaggcc gatccCGtc acaacctcgt caaggagctg cgcaagcgcg acatccgcat      780
cgacggcatc ggtctcgagt cccacttcga ggtcggtttc accccctccc tacaggacca      840
gctcagcacc aagcagggct acatcgCGct cggctctgac gtcgccatca ccgagctgga      900
cgtgcgcttc acccaggccc ctactacga tgccCGggc gagaagcagc aggcccagga      960
ctactatacc agCGtttcta gctgcacga ggcCGgcccc aagtgcacg gtatcacCGt      1020
ctgggacttc gatgacaagt actCGtgggt tccttacact ttCGccggcc aggggtggtgc      1080
agatatctac aatgctacct tgcaggccaa gcctgcctac tatgccattg ccgatgctct      1140
tcagggcaag gcctgcagcg tctgctag                                     1168

```

<210> 2

<211> 364

<212> PRT

<213> *Talaromyces leycettanus*

<400> 2

Met Arg Phe Ser Leu Ala Thr Ala Ala Leu Leu Ala Gly Pro Ala Leu

1		5		10		15													
Ala	Ala	Pro	Pro	Ala	Pro	Arg	His	Asp	Asp	Lys	Asp	Val	Gly	Leu	Asn				
		20						25					30						
Ala	Leu	Ala	Gln	Arg	Ala	Gly	Lys	Leu	Trp	Phe	Gly	Thr	Ala	Ala	Asp				
		35					40					45							
Ile	Pro	Gly	Thr	Asp	Glu	Thr	Thr	Asp	Ala	Ala	Tyr	Leu	Lys	Ile	Leu				
	50					55					60								
Glu	Asn	Pro	Ala	Asn	Phe	Gly	Glu	Ile	Thr	Pro	Ala	Asn	Ala	Met	Lys				
65					70					75				80					
Phe	Met	Tyr	Thr	Glu	Pro	Glu	Gln	Asn	Val	Phe	Asn	Tyr	Thr	Gly	Gly				
				85					90					95					
Asp	Tyr	Val	Leu	Asn	Leu	Ala	Glu	Arg	His	Gly	Gln	Arg	Val	Arg	Cys				
			100					105						110					
His	Asn	Leu	Val	Trp	Ala	Ser	Gln	Leu	Ser	Asp	Phe	Val	Asn	Asn	Gly				
		115					120						125						
Asn	Trp	Thr	Lys	Glu	Ser	Leu	Thr	Ala	Val	Met	Arg	Asn	His	Ile	Phe				
	130					135					140								
His	Val	Val	Gln	His	Phe	Gly	Arg	Arg	Cys	Tyr	Ser	Trp	Asp	Val	Val				
145					150					155					160				
Asn	Glu	Ala	Leu	Asn	Gly	Asp	Gly	Thr	Phe	Ser	Ser	Ser	Ile	Trp	Tyr				
				165					170					175					
Asp	Thr	Ile	Gly	Glu	Asp	Tyr	Phe	Tyr	Leu	Ala	Phe	Gln	Tyr	Ala	Gln				
			180					185					190						
Glu	Ala	Leu	Ala	Glu	Ile	His	Ala	Asn	Asp	Val	Lys	Leu	Tyr	Tyr	Asn				
		195					200					205							
Asp	Tyr	Gly	Ile	Glu	Asn	Pro	Gly	Thr	Lys	Ala	Asp	Ala	Val	His	Asn				
	210					215					220								
Leu	Val	Lys	Glu	Leu	Arg	Lys	Arg	Asp	Ile	Arg	Ile	Asp	Gly	Ile	Gly				
225					230				235					240					
Leu	Glu	Ser	His	Phe	Glu	Val	Gly	Phe	Thr	Pro	Ser	Leu	Gln	Asp	Gln				
				245					250					255					
Leu	Ser	Thr	Lys	Gln	Gly	Tyr	Ile	Ala	Leu	Gly	Leu	Asp	Val	Ala	Ile				
			260					265					270						
Thr	Glu	Leu	Asp	Val	Arg	Phe	Thr	Gln	Ala	Pro	Tyr	Tyr	Asp	Ala	Ala				
			275				280						285						
Gly	Glu	Lys	Gln	Gln	Ala	Gln	Asp	Tyr	Tyr	Thr	Ser	Val	Ser	Ser	Cys				
	290					295					300								

Ile Glu Ala Gly Pro Lys Cys Ile Gly Ile Thr Val Trp Asp Phe Asp
 305 310 315 320

Asp Lys Tyr Ser Trp Val Pro Tyr Thr Phe Ala Gly Gln Gly Gly Ala
 325 330 335

Asp Ile Tyr Asn Ala Thr Leu Gln Ala Lys Pro Ala Tyr Tyr Ala Ile
 340 345 350

Ala Asp Ala Leu Gln Gly Lys Ala Cys Ser Val Cys
 355 360

<210> 3

<211> 1708

<212> DNA

<213> Talaromyces leycettanus

<400> 3

atggtccgtc tttccgctgg acttatcgtc ctccccctcg tgtccgccgc ggccgctgat	60
ctccagagcc gccaggcggc acagagcatc aacaccctca tccaggccaa gggcaagaag	120
tactggggca cttgcgccga tgagggccga ttgaccgaga actcgcaaaa ccgggccatc	180
gccaaggcgg actttggcca ggtgacgccg gagaacagca tgaagtggga tgctactgag	240
cgttagtcag gatgtccatg tgcaatatat agatggatgg ctaactgctg ttgatgtgta	300
gcaagccagg gccagttcaa ctttgctcag gctgattggg ttgtaagtga agctggcatg	360
ttgttcagat cctagactgt acggctggct gactgttctc aaggtaact gggcgagca	420
aatggcaag ctgatccgag gccacaacct gggtagtcc tgttcatcat cacaatcgtc	480
atagtagcta cgaacacgct aacttatgtg tcaacagtgt ggcaactcca gctcccatcc	540
tggtgtgctg gtatcaccca caagacggca ttgaccaatg ccatgaccaa ccacatcacc	600
accctggtga gccgctataa ggggaagatc tatgcctggg taagtgtttt tcttttctct	660
actgatatca gttccagaga aacggaggcc actgacagtt gtacaggacg tcgtcaacga	720
acccttcaac gaagatggaa gtcttcgtca gacctgcttc tacaacgtca tcggacctga	780
ctacatcaag attgccttcc aaacagctcg tgcggccgat ccgaacgcga agctctacgt	840
caatgattac aagtaagact gatcaagtcc cagccttagc tgcgttaatc cgctgacatt	900
cataacctca gccttgactc cgcttcctac gccaaagcca ccggcgtggc gaaccaggtc	960
aagcagtgga ttgcacaggg tgtcccgatt gacggtattg gttctgagtc tcaccttagg	1020
tatagctgac tcccatgtct caagtcaatg ctagttgtga acgtgtactt acgactttca	1080
aaaacagcgc cgggtgcagga gcgggtgtgc cagctgccct gcaagtgtc gccaatccg	1140
gagtctccga ggtcgcgac accgaactcg atatcgcca agcctcatcc actgactatg	1200
acaacgtaag aagaactctt ctgatatcct ctttaacca taaagtgtcg tgtgactaac	1260
actctttttt tttcaacata ggtcgcccaa gcctgcctga acgtcgcgaa gtgtgttggt	1320
atcacttcct ggggtatctc tgataaggtc cgtttcatgc ctgcatttcc cttccctgaa	1380
agacaagatg tgtctttcca ttgattgctg acgcctaaat caataacagg actcctggcg	1440

```

ctccagcgag aacctgatc tcttcgacag caactatcag cccaaggctg cctacaatgc      1500
tcttgtgacc ttgctcgggtg gaagctccgg ctccggctct ggctctggct ctggctctgg      1560
ctctgggtca ggctcaggct caggctcagg ctccggccag gctcaacact ggggtcagtg      1620
tgggtggcgaa ggctggaccg gaccaacgtc ctgtgtctct ccatacactt gccagtagca      1680
gaaccagtgg tactcccagt gcttgtaa                                          1708

```

<210> 4

<211> 389

<212> PRT

<213> Talaromyces leycettanus

<400> 4

```

Met Val Arg Leu Ser Ala Gly Leu Ile Val Leu Pro Leu Val Ser Ala
1              5              10              15

```

```

Ala Ala Val Asp Leu Gln Ser Arg Gln Ala Ala Gln Ser Ile Asn Thr
20              25              30

```

```

Leu Ile Gln Ala Lys Gly Lys Lys Tyr Trp Gly Thr Cys Ala Asp Glu
35              40              45

```

```

Gly Arg Leu Thr Glu Asn Ser Gln Asn Pro Ala Ile Ala Lys Ala Asp
50              55              60

```

```

Phe Gly Gln Val Thr Pro Glu Asn Ser Met Lys Trp Asp Ala Thr Glu
65              70              75              80

```

```

Pro Ser Gln Gly Gln Phe Asn Phe Ala Gln Ala Asp Trp Leu Val Asn
85              90              95

```

```

Trp Ala Gln Gln Asn Gly Lys Leu Ile Arg Gly His Asn Leu Val Trp
100             105             110

```

```

His Ser Gln Leu Pro Ser Trp Val Cys Gly Ile Thr Asp Lys Thr Ala
115             120             125

```

```

Leu Thr Asn Ala Met Thr Asn His Ile Thr Thr Leu Val Ser Arg Tyr
130             135             140

```

```

Lys Gly Lys Ile Tyr Ala Trp Asp Val Val Asn Glu Pro Phe Asn Glu
145             150             155             160

```

```

Asp Gly Ser Leu Arg Gln Thr Cys Phe Tyr Asn Val Ile Gly Pro Asp
165             170             175

```

```

Tyr Ile Lys Ile Ala Phe Gln Thr Ala Arg Ala Ala Asp Pro Asn Ala
180             185             190

```

```

Lys Leu Tyr Val Asn Asp Tyr Asn Leu Asp Ser Ala Ser Tyr Ala Lys
195             200             205

```

```

Thr Thr Gly Val Ala Asn Gln Val Lys Gln Trp Ile Ala Gln Gly Val
210             215             220

```

Pro Ile Asp Gly Ile Gly Ser Glu Ser His Leu Ser Ala Gly Ala Gly
225 230 235 240

Ala Gly Val Pro Ala Ala Leu Gln Val Leu Ala Asn Ser Gly Val Ser
245 250 255

Glu Val Ala Ile Thr Glu Leu Asp Ile Ala Gln Ala Ser Ser Thr Asp
260 265 270

Tyr Asp Asn Val Ala Gln Ala Cys Leu Asn Val Ala Lys Cys Val Gly
275 280 285

Ile Thr Ser Trp Gly Ile Ser Asp Lys Asp Ser Trp Arg Ser Ser Glu
290 295 300

Asn Pro Asp Leu Phe Asp Ser Asn Tyr Gln Pro Lys Ala Ala Tyr Asn
305 310 315 320

Ala Leu Val Thr Leu Leu Gly Gly Ser Ser Gly Ser Gly Ser Gly Ser
325 330 335

Gly Ser Gly Ser Gly Ser Gly Ser Gly Ser Gly Ser Gly Ser Gly Ser
340 345 350

Gly Gln Ala Gln His Trp Gly Gln Cys Gly Gly Glu Gly Trp Thr Gly
355 360 365

Pro Thr Ser Cys Val Ser Pro Tyr Thr Cys Gln Tyr Gln Asn Gln Trp
370 375 380

Tyr Ser Gln Cys Leu
385

<210> 5

<211> 1520

<212> DNA

<213> Talaromyces leycettanus

<400> 5

atggtccatc tttcttccct ggccctggct ttggcgcgcg gctcgcagct gtatgtgac	60
catgccatga ctcgagaagt gctcccaaaa ctgactccaa gtctcaatct tagtgcccaa	120
gctgcaggtc ttaacactgc tgccaaagcg attggaaagc tctatttcgg taccgcaacc	180
gacaacccgg agctgtccga cagcacatac atgcaggaga cggataacac cgatgatttc	240
ggccaactca cccagctaa ctccatgaag gttcgtctgac atcttagttc cccccccctt	300
ttgggaatct gcgcggagat atgctgagcc ttcaaaacta gtgggatgcc accgagccct	360
ctcagaacac cttcaccttc accaacggtg atcagatcgc aaaccttgct aagagcaacg	420
gtcagatgct gagatgccac aacctggtgt ggtacaacca gttgcccagc tggggaagc	480
aaccggttct gttaatatca tcagcgtgac cgcacgatc gtattgcgcg gagattggaa	540
agatttgcaa gctaattgtca ctacagtcac cagcggatct tggaccaatg ccacgcttct	600
tgcggccatg aagaaccaca tcaccaacgt tgtgacccac tacaagggac agtgctacgc	660

```

ttgggatgtt gtcaacgaag gtacgtttcg attcggcttc cctcggaccg tatctgcagg      720
caaaaagggtc aatcaattga caatcgtgat cccagctct caacgatgat ggcacctacc      780
gatccaatgt cttctatcag tacatcggcg aggcatacat tccattgcc tttgcgaccg      840
ctgccgccgc cgatccaaac gcgaagctct actacaacga ctacaacatt gagtaccccg      900
ggccaaggc caccgccgc cagaacatcg tcaagatggt caaggcttac ggcgcaaaa      960
tcgacggtgt cggctctgaa tctcacttca tcgttggcag caccctagc cagagctccc     1020
agcagagcaa catggctgct ttcaccgcgc tcggcgctga ggtcgccatc accgaactgg     1080
atatccgcat gacgttgctt tccaccagtg ctctcttggc ccagcaatcc accgattacc     1140
agagcactgt gtcggcttgc gtgaacactc cgaagtgcac tggatcacc ctctgggact     1200

ggaccgacaa gtactcctgg gttcccaaca ccttctcgg ccaagggtgac gcctgccct      1260
gggattctaa ctaccagaag aagcctgcct actacggtat cttgactgcg ctcgaggga      1320
gcgttccac ctccaccacc accactctgg tgacctccac caggacttcg actacgacca      1380
gcacttcggc cacctccacg tctactggcg ttgctcagca ctggggccag tgcggtgta      1440
tcggctggac agggccgact acctgcgcta gccctacac ctgccaggaa ctgaatccct     1500
actactacca gtgcctgtaa                                           1520

```

<210> 6

<211> 405

<212> PRT

<213> Talaromyces leycettanus

<400> 6

```

Met Val His Leu Ser Ser Leu Ala Leu Ala Ala Gly Ser Gln
1           5           10          15

Leu Ala Gln Ala Ala Gly Leu Asn Thr Ala Ala Lys Ala Ile Gly Lys
20           25           30

Leu Tyr Phe Gly Thr Ala Thr Asp Asn Pro Glu Leu Ser Asp Ser Thr
35           40           45

Tyr Met Gln Glu Thr Asp Asn Thr Asp Asp Phe Gly Gln Leu Thr Pro
50           55           60

Ala Asn Ser Met Lys Trp Asp Ala Thr Glu Pro Ser Gln Asn Thr Phe
65           70           75           80

Thr Phe Thr Asn Gly Asp Gln Ile Ala Asn Leu Ala Lys Ser Asn Gly
85           90           95

Gln Met Leu Arg Cys His Asn Leu Val Trp Tyr Asn Gln Leu Pro Ser
100          105          110

Trp Val Thr Ser Gly Ser Trp Thr Asn Ala Thr Leu Leu Ala Ala Met
115          120          125

```

Lys Asn His Ile Thr Asn Val Val Thr His Tyr Lys Gly Gln Cys Tyr
 130 135 140

Ala Trp Asp Val Val Asn Glu Ala Leu Asn Asp Asp Gly Thr Tyr Arg
 145 150 155 160

Ser Asn Val Phe Tyr Gln Tyr Ile Gly Glu Ala Tyr Ile Pro Ile Ala
 165 170 175

Phe Ala Thr Ala Ala Ala Ala Asp Pro Asn Ala Lys Leu Tyr Tyr Asn
 180 185 190

Asp Tyr Asn Ile Glu Tyr Pro Gly Ala Lys Ala Thr Ala Ala Gln Asn
 195 200 205

Ile Val Lys Met Val Lys Ala Tyr Gly Ala Lys Ile Asp Gly Val Gly
 210 215 220

Leu Gln Ser His Phe Ile Val Gly Ser Thr Pro Ser Gln Ser Ser Gln
 225 230 235 240

Gln Ser Asn Met Ala Ala Phe Thr Ala Leu Gly Val Glu Val Ala Ile
 245 250 255

Thr Glu Leu Asp Ile Arg Met Thr Leu Pro Ser Thr Ser Ala Leu Leu
 260 265 270

Ala Gln Gln Ser Thr Asp Tyr Gln Ser Thr Val Ser Ala Cys Val Asn
 275 280 285

Thr Pro Lys Cys Ile Gly Ile Thr Leu Trp Asp Trp Thr Asp Lys Tyr
 290 295 300

Ser Trp Val Pro Asn Thr Phe Ser Gly Gln Gly Asp Ala Cys Pro Trp
 305 310 315 320

Asp Ser Asn Tyr Gln Lys Lys Pro Ala Tyr Tyr Gly Ile Leu Thr Ala
 325 330 335

Leu Gly Gly Ser Ala Ser Thr Ser Thr Thr Thr Thr Leu Val Thr Ser
 340 345 350

Thr Arg Thr Ser Thr Thr Thr Ser Thr Ser Ala Thr Ser Thr Ser Thr
 355 360 365

Gly Val Ala Gln His Trp Gly Gln Cys Gly Gly Ile Gly Trp Thr Gly
 370 375 380

Pro Thr Thr Cys Ala Ser Pro Tyr Thr Cys Gln Glu Leu Asn Pro Tyr
 385 390 395 400

Tyr Tyr Gln Cys Leu
 405

<210> 7

<211> 41

<212> DNA

<213> Artificial

<220>

<223> PCR primer

<400> 7

acacaactgg ggatccacca tgcgtttctc ctggccact g 41

<210> 8

<211> 36

<212> DNA

<213> Artificial

<220>

<223> PCR primer

<400> 8

ccctctagat ctcgagctag cagacgctgc aggcct 36

<210> 9

<211> 40

<212> DNA

<213> Artificial

<220>

<223> PCR primer

<400> 9

acacaactgg ggatccacca tggtcctct tccgctgga 40

<210> 10

<211> 41

<212> DNA

<213> Artificial

<220>

<223> PCR primer

<400> 10

ccctctagat ctcgagttac aagcactggg agtaccactg g 41

<210> 11

<211> 43

<212> DNA

<213> Artificial

<220>

<223> PCR primer

<400> 11

acacaactgg ggatccacca tggatcatct ttctccctg gcc 43

<210> 12

<211> 44

<212> DNA

<213> Artificial

<220>

<223> PCR primer

<400> 12

ccctctagat ctcgagttac aggcaactgg agtagtaggg attc 44

REFERENCES CITED IN THE DESCRIPTION

Cited references

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- [US2010124769A1 \[0005\]](#)
- [WO02095014A \[0052\] \[0052\] \[0227\]](#)
- [WO9517413A \[0074\]](#)
- [WO9522625A \[0074\]](#)
- [US5223409A \[0074\]](#)
- [WO9206204A \[0074\]](#)
- [WO9943835A \[0099\]](#)
- [WO9600787A \[0100\] \[0148\]](#)
- [WO0056900A \[0100\] \[0100\] \[0100\]](#)
- [WO9425612A \[0107\]](#)
- [WO9533836A \[0118\]](#)

- [WO0024883A \[0131\] \[0131\]](#)
- [EP238023A \[0148\]](#)
- [WO9114772A \[0164\]](#)
- [US6395966B \[0168\]](#)
- [US7151204B \[0168\] \[0169\]](#)
- [US20020164730A \[0189\] \[0198\]](#)
- [WO2006110891A \[0193\]](#)
- [WO2006110899A \[0193\]](#)
- [WO2006110900A \[0193\]](#)
- [WO2006110901A \[0193\]](#)
- [WO2006032282A \[0195\]](#)
- [WO9117243A \[0222\]](#)
- [WO9117244A \[0222\]](#)
- [WO9105039A \[0224\]](#)
- [WO9315186A \[0224\]](#)
- [US5275944A \[0224\]](#)
- [WO9602551A \[0224\]](#)
- [US5536655A \[0224\]](#)
- [WO0070031A \[0224\]](#)
- [WO05093050A \[0224\] \[0224\] \[0224\]](#)
- [WO2011059740A \[0226\]](#)
- [WO2009042871A \[0226\]](#)
- [WO2010141325A \[0226\]](#)
- [WO2006074435A \[0226\]](#)
- [WO2010057086A \[0226\]](#)
- [WO2005047499A \[0227\]](#)
- [WO2007019442A \[0227\] \[0227\]](#)
- [WO2010088387A \[0227\]](#)
- [WO2011035029A \[0227\]](#)
- [WO2008057637A \[0228\] \[0228\]](#)
- [WO9813465A \[0230\]](#)
- [WO9815619A \[0230\]](#)
- [WO9815633A \[0230\]](#)
- [WO9906574A \[0230\]](#)
- [WO9910481A \[0230\]](#)
- [WO9925847A \[0230\]](#)
- [WO9931255A \[0230\]](#)
- [WO02101078A \[0230\]](#)
- [WO2003027306A \[0230\]](#)
- [WO2003052054A \[0230\]](#)
- [WO2003052055A \[0230\]](#)
- [WO2003052056A \[0230\]](#)
- [WO2003052057A \[0230\]](#)
- [WO2003052118A \[0230\]](#)

- [WO2004016760A \[0230\]](#)
- [WO2004043980A \[0230\]](#)
- [WO2004048592A \[0230\]](#)
- [WO2005001065A \[0230\]](#)
- [WO2005028636A \[0230\]](#)
- [WO2005093050A \[0230\]](#)
- [WO2005093073A \[0230\]](#)
- [WO2006074005A \[0230\]](#)
- [WO2006117432A \[0230\]](#)
- [WO2007071818A \[0230\]](#)
- [WO2007071820A \[0230\]](#)
- [WO2008008070A \[0230\]](#)
- [WO2008008793A \[0230\]](#)
- [US5457046A \[0230\]](#)
- [US5648263A \[0230\]](#)
- [US5686593A \[0230\]](#)
- [WO2005074647A \[0232\]](#)
- [WO2008148131A \[0232\]](#)
- [WO2011035027A \[0232\]](#)
- [WO2005074656A \[0232\]](#)
- [WO2010065830A \[0232\]](#)
- [WO2007089290A \[0232\]](#)
- [WO2009085935A \[0232\]](#)
- [WO2009085859A \[0232\]](#)
- [WO2009085864A \[0232\]](#)
- [WO2009085868A \[0232\]](#)
- [WO2010138754A \[0232\]](#)
- [WO2011005867A \[0232\]](#)
- [WO2011039319A \[0232\]](#)
- [WO2011041397A \[0232\]](#)
- [WO2011041504A \[0232\]](#)
- [WO2008151043A \[0233\]](#)
- [WO9421785A \[0245\]](#)
- [WO2006078256A \[0245\]](#)
- [WO2011041405A \[0245\]](#)
- [WO2010126772A \[0245\]](#)
- [WO2009079210A \[0245\]](#)
- [WO2011057083A \[0245\]](#)
- [WO2010108918A \[0247\]](#)
- [WO2009073709A \[0247\]](#)
- [WO2005001036A \[0247\]](#)
- [WO2010014880A \[0247\]](#)
- [WO2009042846A \[0247\]](#)
- [WO2009076122A \[0248\]](#)

- [WO2009127729A \[0248\]](#)
- [WO2010053838A \[0248\]](#)
- [WO2010065448A \[0248\]](#)
- [WO2006114094A \[0249\] \[0249\]](#)
- [WO2009073383A \[0249\]](#)
- [WO2010014706A \[0250\]](#)
- [WO2009068565A \[0250\]](#)
- [WO2003062430A \[0266\]](#)
- [WO0240694A \[0290\] \[0309\] \[0323\] \[0337\]](#)
- [WO2005042735A \[0301\] \[0315\] \[0329\]](#)
- [EP0238023A \[0309\] \[0323\] \[0337\]](#)

Non-patent literature cited in the description

- **LEVERA** new reaction for colorimetric determination of carbohydrates *Anal. Biochem.*, 1972, vol. 47, 273-279 [\[0012\] \[0066\]](#)
- **DE VRIESJ.** *Bacteriol.*, 1998, vol. 180, 243-249 [\[0016\]](#)
- **VENTURI et al.** Extracellular beta-D-glucosidase from *Chaetomium thermophilum* var. *coprophilum*: production, purification and some biochemical properties *J. Basic Microbiol.*, 2002, vol. 42, 55-66 [\[0017\]](#)
- **TEERI** Crystalline cellulose degradation: New insight into the function of cellobiohydrolases *Trends in Biotechnology*, 1997, vol. 15, 160-167 [\[0020\]](#)
- **TEERI et al.** *Trichoderma reesei* cellobiohydrolases: why so efficient on crystalline cellulose? *Biochem. Soc. Trans.*, 1998, vol. 26, 173-178 [\[0020\]](#)
- **LEVER et al.** *Anal. Biochem.*, 1972, vol. 47, 273-279 [\[0020\]](#)
- **VAN TILBEURGH et al.** *FEBS Letters*, 1982, vol. 149, 152-156 [\[0020\]](#)
- **VAN TILBEURGH CLAEYSSENS** *FEBS Letters*, 1985, vol. 187, 283-288 [\[0020\]](#)
- **TOMME et al.** *Eur. J. Biochem.*, 1988, vol. 170, 575-581 [\[0020\]](#)
- **WISELOGEL et al.** *Handbook on Bioethanol* Taylor & Francis 1995 0000105-118 [\[0022\]](#)
- **WYMAN** *Bioresource Technology*, 1994, vol. 50, 3-16 [\[0022\]](#)
- **LYND** *Applied Biochemistry and Biotechnology*, 1990, vol. 24, 25695-719 [\[0022\]](#)
- Recent Progress in Bioconversion of Lignocellulosics **MOSIER ET AL.** *Advances in Biochemical Engineering/Biotechnology* Springer-Verlag 1999 0000 vol. 65, 23-40 [\[0022\]](#)
- **ZHANG et al.** Outlook for cellulase improvement: Screening and selection strategies *Biotechnology Advances*, 2006, vol. 24, 452-481 [\[0029\]](#)
- **GHOSE** Measurement of cellulase activities *Pure Appl. Chem.*, 1987, vol. 59, 257-68 [\[0029\]](#)
- **ZHANG et al.** *Biotechnology Advances*, 2006, vol. 24, 452-481 [\[0033\]](#)
- **GHOSE** *Pure and Appl. Chem.*, 1987, vol. 59, 257-268 [\[0033\]](#)
- **HENRISSATA** classification of glycosyl hydrolases based on amino-acid sequence

- similaritiesBiochem. J., 1991, vol. 280, 309-316 [0036] [0229]
- **HENRISSATBAIROCH**Updating the sequence-based classification of glycosyl hydrolasesBiochem. J., 1996, vol. 316, 695-696 [0036] [0229]
 - **SHALLOMSHOHAM**Microbial hemicellulasesCurrent Opinion In Microbiology, 2003, vol. 6, 3219-228 [0039]
 - **GHOSEBISARIA**Pure & Appl. Chem., 1987, vol. 59, 1739-1752 [0039]
 - **NIELSEN et al.**Protein Engineering, 1997, vol. 10, 1-6 [0044] [0307] [0321] [0335]
 - **NEEDLEMANWUNSCHJ.** Mol. Biol., 1970, vol. 48, 443-453 [0056] [0308] [0322] [0336]
 - **RICE et al.**EMBOSS: The European Molecular Biology Open Software SuiteTrends Genet., 2000, vol. 16, 276-277 [0056]
 - **RICE et al.**EMBOSS: The European Molecular Biology Open Software Suite, 2000, [0057]
 - **EBRINGEROVA et al.**Adv. Polym. Sci., 2005, vol. 186, 1-67 [0062]
 - **BIELYPUCHARD**Recent progress in the assays of xylanolytic enzymesJournal of the Science of Food and Agriculture, 2006, vol. 86, 111636-1647 [0064]
 - **SPANIKOVABIELY**Glucuronoyl esterase - Novel carbohydrate esterase produced by Schizophyllum communeFEBS Letters, 2006, vol. 580, 194597-4601 [0064]
 - **HERRMANN et al.**The beta-D-xylosidase of Trichoderma reesei is a multifunctional beta-D-xylan xylohydrolaseBiochemical Journal, 1997, vol. 321, 375-381 [0064]
 - **BAILEY et al.**Interlaboratory testing of methods for assay of xylanase activityJournal of Biotechnology, 1992, vol. 23, 3257-270 [0065]
 - **SAMBROOK et al.**Molecular Cloning, A Laboratory ManualCold Spring Harbor19890000 [0068]
 - **H. NEURATHR.L. HILL**The ProteinsAcademic Press19790000 [0071]
 - **CUNNINGHAMWELL**Science, 1989, vol. 244, 1081-1085 [0073]
 - **HILTON et al.**J. Biol. Chem., 1996, vol. 271, 4699-4708 [0073]
 - **DE VOS et al.**Science, 1992, vol. 255, 306-312 [0073]
 - **SMITH et al.**J. Mol. Biol., 1992, vol. 224, 899-904 [0073]
 - **WLODAVER et al.**FEBS Lett., 1992, vol. 309, 59-64 [0073]
 - **REIDHAAR-OLSONSAUER**Science, 1988, vol. 241, 53-57 [0074]
 - **BOWIESAUER**Proc. Natl. Acad. Sci. USA, 1989, vol. 86, 2152-2156 [0074]
 - **LOWMAN et al.**Biochemistry, 1991, vol. 30, 10832-10837 [0074]
 - **DERBYSHIRE et al.**Gene, 1986, vol. 46, 145- [0074]
 - **NER et al.**DNA, 1988, vol. 7, 127- [0074]
 - **NESS et al.**Nature Biotechnology, 1999, vol. 17, 893-896 [0075]
 - **COOPER et al.**EMB J., 1993, vol. 12, 2575-2583 [0077]
 - **DAWSON et al.**Science, 1994, vol. 266, 776-779 [0077]
 - **MARTIN et al.**J. Ind. Microbiol. Biotechnol., 2003, vol. 3, 568-576 [0078]
 - **SVETINA et al.**J. Biotechnol., 2000, vol. 76, 245-251 [0078]
 - **RASMUSSEN-WILSON et al.**Appl. Environ. Microbiol., 1997, vol. 63, 3488-3493 [0078]
 - **WARD et al.**Biotechnology, 1995, vol. 13, 498-503 [0078]
 - **CONTRERAS et al.**Biotechnology, 1991, vol. 9, 378-381 [0078]
 - **EATON et al.**Biochemistry, 1986, vol. 25, 505-512 [0078]
 - **COLLINS-RACIE et al.**Biotechnology, 1995, vol. 13, 982-987 [0078]

- **CARTER et al.**Proteins: Structure, Function, and Genetics, 1989, vol. 6, 240-248 [0078]
- **STEVENS**Drug Discovery World, 2003, vol. 4, 35-48 [0078]
- **INNIS et al.**PCR: A Guide to Methods and ApplicationAcademic Press19900000 [0094]
- **FORD et al.**Protein Expression and Purification, 1991, vol. 2, 95-107 [0095]
- **AGAISSELERECLUS**Molecular Microbiology, 1994, vol. 13, 97-107 [0099]
- **EGON et al.**Gene, 1988, vol. 69, 301-315 [0099]
- **VILLA-KAMAROFF et al.**Proc. Natl. Acad. Sci. USA, 1978, vol. 75, 3727-3731 [0099]
- **DEBOER et al.**Proc. Natl. Acad. Sci. USA, 1983, vol. 80, 21-25 [0099]
- **GILBERT et al.**Scientific American, 1980, vol. 242, 74-94 [0099]
- **ROMANOS et al.**Yeast, 1992, vol. 8, 423-488 [0101]
- **HUE et al.**Journal of Bacteriology, 1995, vol. 177, 3465-3471 [0107]
- **GUOSHHERMAN**Mol. Cellular Biol., 1995, vol. 15, 5983-5990 [0113]
- **SIMONENPALVA**Microbiological Reviews, 1993, vol. 57, 109-137 [0115]
- **GEMS et al.**Gene, 1991, vol. 98, 61-67 [0131]
- **CULLEN et al.**Nucleic Acids Res., 1987, vol. 15, 9163-9175 [0131]
- **CHANGCOHEN**Mol. Gen. Genet., 1979, vol. 168, 111-115 [0140]
- **YOUNGSPIZIZEN**J. Bacteriol., 1961, vol. 81, 823-829 [0140]
- **DUBNAUDAVIDOFF-ABELSON**J. Mol. Biol., 1971, vol. 56, 209-221 [0140]
- **SHIGEKAWADOWER**Biotechniques, 1988, vol. 6, 742-751 [0140]
- **KOEHLERTHORNE**J. Bacteriol., 1987, vol. 169, 5271-5278 [0140]
- **HANAHA**NJ. Mol. Biol., 1983, vol. 166, 557-580 [0140]
- **DOWER et al.**Nucleic Acids Res., 1988, vol. 16, 6127-6145 [0140]
- **GONG et al.**Folia Microbiol. (Praha), 2004, vol. 49, 399-405 [0140]
- **MAZODIER et al.**J. Bacteriol., 1989, vol. 171, 3583-3585 [0140]
- **BURKE et al.**Proc. Natl. Acad. Sci. USA, 2001, vol. 98, 6289-6294 [0140]
- **CHOI et al.**J. Microbiol. Methods, 2006, vol. 64, 391-397 [0140]
- **PINEDOSMETS**Appl. Environ. Microbiol., 2005, vol. 71, 51-57 [0140]
- **PERRYKURAMITSU**Infect. Immun., 1981, vol. 32, 1295-1297 [0140]
- **CATTJOLICK**Microbios, 1991, vol. 68, 189-207 [0140]
- **BUCKLEY et al.**Appl. Environ. Microbiol., 1999, vol. 65, 3800-3804 [0140]
- **CLEWELL**Microbiol. Rev., 1981, vol. 45, 409-436 [0140]
- **HAWKSWORTH et al.**Ainsworth and Bisby's Dictionary of The FungiCAB International, University Press19950000 [0142]
- **Biology and Activities of Yeast**Soc. App. Bacteriol. Symposium Series19800000 [0143]
- **YELTON et al.**Proc. Natl. Acad. Sci. USA, 1984, vol. 81, 1470-1474 [0148]
- **CHRISTENSEN et al.**Bio/Technology, 1988, vol. 6, 1419-1422 [0148]
- **MALARDIER et al.**Gene, 1989, vol. 78, 147-156 [0148]
- **BECKERGUARENTE**Guide to Yeast Genetics and Molecular Biology, Methods in EnzymologyAcademic Press, Inc.vol. 194, 182-187 [0148]
- **ITO et al.**J. Bacteriol., 1983, vol. 153, 163- [0148]
- **HINNEN et al.**Proc. Natl. Acad. Sci. USA, 1978, vol. 75, 1920- [0148]
- **Protein Purification**VCH Publishers19890000 [0154]
- **TAGUE et al.**Plant Physiology, 1988, vol. 86, 506- [0163]
- **FRANCK et al.**Cell, 1980, vol. 21, 285-294 [0164]

- **CHRISTENSEN et al.** Plant Mol. Biol., 1992, vol. 18, 675-689 [0164]
- **ZHANG et al.** Plant Cell, 1991, vol. 3, 1155-1165 [0164]
- **EDWARDSCORUZZI** Ann. Rev. Genet., 1990, vol. 24, 275-303 [0164]
- **ITO et al.** Plant Mol. Biol., 1994, vol. 24, 863-878 [0164]
- **WU et al.** Plant Cell Physiol., 1998, vol. 39, 885-889 [0164]
- **CONRAD et al.** J. Plant Physiol., 1998, vol. 152, 708-711 [0164]
- **CHEN et al.** Plant Cell Physiol., 1998, vol. 39, 935-941 [0164]
- **KYOZUKA et al.** Plant Physiol., 1993, vol. 102, 991-1000 [0164]
- **MITRAHIGGINS** Plant Mol. Biol., 1994, vol. 26, 85-93 [0164]
- **KAGAYA et al.** Mol. Gen. Genet., 1995, vol. 248, 668-674 [0164]
- **XU et al.** Plant Mol. Biol., 1993, vol. 22, 573-588 [0164]
- **GASSER et al.** Science, 1990, vol. 244, 1293- [0167]
- **POTRYKUS** Bio/Technology, 1990, vol. 8, 535- [0167]
- **SHIMAMOTO et al.** Nature, 1989, vol. 338, 274- [0167]
- **HOOYKASSCHILPEROORT** Plant Mol. Biol., 1992, vol. 19, 15-38 [0168]
- **CHRISTOU** Plant J., 1992, vol. 2, 275-281 [0168]
- **SHIMAMOTO** Curr. Opin. Biotechnol., 1994, vol. 5, 158-162 [0168]
- **VASIL et al.** Bio/Technology, 1992, vol. 10, 667-674 [0168]
- **OMIRULLEH et al.** Plant Mol. Biol., 1993, vol. 21, 415-428 [0168]
- Cellulose bioconversion technology **PHILIPPIDIS, G. P.** Handbook on Bioethanol: Production and Utilization Taylor & Francis 1996 0000179-212 [0183]
- **SHEEHAN, J. HIMMEL, M.** Enzymes, energy and the environment: A strategic perspective on the U.S. Department of Energy's research and development activities for bioethanol Biotechnol. Prog., 1999, vol. 15, 817-827 [0183]
- **LYND et al.** Microbial cellulose utilization: Fundamentals and biotechnology Microbiol. Mol. Biol. Reviews, 2002, vol. 66, 506-577 [0183]
- **FERNANDA DE CASTILHOS CORAZZA FLÁVIO FARIA DE MORAES GISELLA MARIA ZANINIVO NEITZEL** Optimal control in fed-batch reactor for the cellobiose hydrolysis, Acta Scientiarum Technology, 2003, vol. 25, 33-38 [0184]
- **GUSAKOV SINITSYN** Kinetics of the enzymatic hydrolysis of cellulose: 1. A mathematical model for a batch reactor process Enz. Microb. Technol., 1985, vol. 7, 346-352 [0184]
- **RYULEE** Bioconversion of waste cellulose by using an attrition bioreactor Biotechnol. Bioeng., 1983, vol. 25, 53-65 [0184]
- **GUSAKOV et al.** Enhancement of enzymatic cellulose hydrolysis using a novel type of bioreactor with intensive stirring induced by electromagnetic field Appl. Biochem. Biotechnol., 1996, vol. 56, 141-153 [0184]
- **CHANDRA et al.** Substrate pretreatment: The key to effective enzymatic hydrolysis of lignocellulosics? Adv. Biochem. Engin./Biotechnol., 2007, vol. 108, 67-93 [0185]
- **GALBEZACCHI** Pretreatment of lignocellulosic materials for efficient bioethanol production Adv. Biochem. Engin./Biotechnol., 2007, vol. 108, 41-65 [0185]
- **HENDRIKS ZEEMAN** Pretreatments to enhance the digestibility of lignocellulosic biomass Bioresource Technol., 2009, vol. 100, 10-18 [0185]
- **MOSIER et al.** Features of promising technologies for pretreatment of lignocellulosic biomass Bioresource Technol., 2005, vol. 96, 673-686 [0185]

- **TAHERZADEHKARIMI** Pretreatment of lignocellulosic wastes to improve ethanol and biogas production: A review *Int. J. of Mol. Sci.*, 2008, vol. 9, 1621-1651 [0185]
- **YANGWYMAN** Pretreatment: the key to unlocking low-cost cellulosic ethanol *Biofuels Bioproducts and Biorefining-Biofpr.*, 2008, vol. 2, 26-40 [0185]
- **DUFFMURRAY** *Bioresource Technology*, 1996, vol. 855, 1-33 [0189]
- **GALBEZACCHI** *Appl. Microbiol. Biotechnol.*, 2002, vol. 59, 618-628 [0189]
- **BALLESTEROS et al.** *Appl. Biochem. Biotechnol.*, 2006, vol. 129-132, 496-508 [0191]
- **VARGA et al.** *Appl. Biochem. Biotechnol.*, 2004, vol. 113-116, 509-523 [0191]
- **SASSNER et al.** *Enzyme Microb. Technol.*, 2006, vol. 39, 756-762 [0191]
- **SHELL et al.** *Bioresource Technol.*, 2004, vol. 91, 179-188 [0191]
- **LEE et al.** *Adv. Biochem. Eng. Biotechnol.*, 1999, vol. 65, 93-115 [0191]
- **WYMAN et al.** *Bioresource Technol.*, 2005, vol. 96, 1959-1966 [0193]
- **MOSIER et al.** *Bioresource Technol.*, 2005, vol. 96, 673-686 [0193]
- **SCHMIDTTHOMSEN** *Bioresource Technol.*, 1998, vol. 64, 139-151 [0194]
- **PALONEN et al.** *Appl. Biochem. Biotechnol.*, 2004, vol. 117, 1-17 [0194]
- **VARGA et al.** *Biotechnol. Bioeng.*, 2004, vol. 88, 567-574 [0194]
- **MARTIN et al.** *J. Chem. Technol. Biotechnol.*, 2006, vol. 81, 1669-1677 [0194]
- **GOLLAPALLI et al.** *Appl. Biochem. Biotechnol.*, 2002, vol. 98, 23-35 [0196]
- **CHUNDAWAT et al.** *Biotechnol. Bioeng.*, 2007, vol. 96, 219-231 [0196]
- **ALIZADEH et al.** *Appl. Biochem. Biotechnol.*, 2005, vol. 121, 1133-1141 [0196]
- **TEYMOURI et al.** *Bioresource Technol.*, 2005, vol. 96, 2014-2018 [0196]
- **PAN et al.** *Biotechnol. Bioeng.*, 2005, vol. 90, 473-481 [0197]
- **PAN et al.** *Biotechnol. Bioeng.*, 2006, vol. 94, 851-861 [0197]
- **KURABI et al.** *Appl. Biochem. Biotechnol.*, 2005, vol. 121, 219-230 [0197]
- **SHELL et al.** *Appl. Biochem. and Biotechnol.*, 2003, vol. 105-108, 69-85 [0198]
- **MOSIER ET AL.** *Bioresource Technology*, 2005, vol. 96, 673-686 [0198]
- Pretreatment of biomass **HSU, T.-A.** *Handbook on Bioethanol: Production and Utilization* Taylor & Francis 1996 0000179-212 [0204]
- **GHOSH SINGH** Physicochemical and biological treatments for enzymatic/microbial conversion of cellulosic biomass *Adv. Appl. Microbiol.*, 1993, vol. 39, 295-333 [0204]
- Pretreating lignocellulosic biomass: a review **MCMILLAN, J. D.** *Enzymatic Conversion of Biomass for Fuels Production* American Chemical Society 1994 0000 [0204]
- Ethanol production from renewable resources **GONG, C. S. CAO, N. J. DU, J. TSAO, G. T.** *Advances in Biochemical Engineering/Biotechnology* Springer-Verlag 1999 0000 vol. 65, 207-241 [0204] [0274]
- **OLSSON HAHN-HAGERDAL** Fermentation of lignocellulosic hydrolysates for ethanol production *Enz. Microb. Tech.*, 1996, vol. 18, 312-331 [0204]
- **VALLANDER ERIKSSON** Production of ethanol from lignocellulosic materials: State of the art *Adv. Biochem. Eng./Biotechnol.*, 1990, vol. 42, 63-95 [0204]
- **PENTTILA et al.** *Gene*, 1986, vol. 45, 253-263 [0225]
- **SALOHEIMO et al.** *Gene*, 1988, vol. 63, 11-22 [0225]
- **OKADA et al.** *Appl. Environ. Microbiol.*, 1988, vol. 64, 555-563 [0225]
- **SALOHEIMO et al.** *Molecular Microbiology*, 1994, vol. 13, 219-228 [0225]
- **OOI et al.** *Nucleic Acids Research*, 1990, vol. 18, 5884- [0225]

- **SAKAMOTO et al.**Current Genetics, 1995, vol. 27, 435-439 [0225]
- **SAARILAHTI et al.**Gene, 1990, vol. 90, 9-14 [0225]
- **KAWAGUCHI et al.**Gene, 1996, vol. 173, 287-288 [0227]
- **DAN et al.**J. Biol. Chem., 2000, vol. 275, 4973-4980 [0227]
- **More Gene Manipulations in Fungi**Academic Press19910000 [0251]
- **BAILEY, J.E.OLLIS, D.F.**Biochemical Engineering FundamentalsMcGraw-Hill Book Company19860000 [0251]
- **LIN et al.**Appl. Microbiol. Biotechnol., 2006, vol. 69, 627-642 [0257]
- **CHENHO**Cloning and improving the expression of Pichia stipitis xylose reductase gene in Saccharomyces cerevisiaeAppl. Biochem. Biotechnol., 1993, vol. 39-40, 135-147 [0266]
- **HO et al.**Genetically engineered Saccharomyces yeast capable of effectively cofermenting glucose and xyloseAppl. Environ. Microbiol., 1998, vol. 64, 1852-1859 [0266]
- **KOTTERCIRIACY**Xylose fermentation by Saccharomyces cerevisiaeAppl. Microbiol. Biotechnol., 1993, vol. 38, 776-783 [0266]
- **WALFRIDSSON et al.**Xylose-metabolizing Saccharomyces cerevisiae strains overexpressing the TKL1 and TAL1 genes encoding the pentose phosphate pathway enzymes transketolase and transaldolaseAppl. Environ. Microbiol., 1995, vol. 61, 4184-4190 [0266]
- **KUYPER et al.**Minimal metabolic engineering of Saccharomyces cerevisiae for efficient anaerobic xylose fermentation: a proof of principleFEMS Yeast Research, 2004, vol. 4, 655-664 [0266]
- **BEALL et al.**Parametric studies of ethanol production from xylose and other sugars by recombinant Escherichia coliBiotech. Bioeng., 1991, vol. 38, 296-303 [0266]
- **INGRAM et al.**Metabolic engineering of bacteria for ethanol productionBiotechnol. Bioeng., 1998, vol. 58, 204-214 [0266]
- **ZHANG et al.**Metabolic engineering of a pentose metabolism pathway in ethanologenic Zymomonas mobilisScience, 1995, vol. 267, 240-243 [0266]
- **DEANDA et al.**Development of an arabinose-fermenting Zymomonas mobilis strain by metabolic pathway engineeringAppl. Environ. Microbiol., 1996, vol. 62, 4465-4470 [0266]
- **The Alcohol Textbook**Nottingham University Press19990000 [0270]
- **ALFENORE et al.**Improving ethanol production and viability of Saccharomyces cerevisiae by a vitamin feeding strategy during fed-batch processSpringer-Verlag20020000 [0272]
- **SILVEIRA, M. M.JONAS, R.**The biotechnological production of sorbitolAppl. Microbiol. Biotechnol., 2002, vol. 59, 400-408 [0274]
- **NIGAMSINGH**Processes for fermentative production of xylitol - a sugar substituteProcess Biochemistry, 1995, vol. 30, 2117-124 [0274]
- **EZEJI et al.**Production of acetone, butanol and ethanol by Clostridium beijerinckii BA101 and in situ recovery by gas strippingWorld Journal of Microbiology and Biotechnology, 2003, vol. 19, 6595-603 [0274]
- **RICHARDMARGARITIS**Empirical modeling of batch fermentation kinetics for

poly(glutamic acid) production and other microbial biopolymers *Biotechnology and Bioengineering*, 2004, vol. 87, 4501-515 [\[0278\]](#)

- **KATAOKA et al.** Studies on hydrogen production by continuous culture system of hydrogen-producing anaerobic bacteria *Water Science and Technology*, 1997, vol. 36, 6-741-47 [\[0279\]](#)
- **GUNASEELAN** Anaerobic digestion of biomass for methane production: A review *Biomass and Bioenergy*, 1997, vol. 13, 1-283-114 [\[0279\]](#)
- **CHENLEE** Membrane-mediated extractive fermentation for lactic acid production from cellulosic biomass *Appl. Biochem. Biotechnol.*, 1997, vol. 63-65, 435-448 [\[0282\]](#)
- *Bacteriological Analytical Manual* 19980000 [\[0292\]](#) [\[0293\]](#) [\[0294\]](#) [\[0296\]](#)

Patentkrav

1. Isoleret polypeptid med xylanaseaktivitet, som er valgt fra gruppen bestående af:
 - (a) et polypeptid, som udviser mindst 85%, mindst 90%, mindst 91%, mindst 92%, mindst 93%, mindst 94%, mindst 95%, mindst 96%, mindst 97%, mindst 98%, mindst 99% eller 100% sekvensidentitet med det modne polypeptid ifølge SEQ ID NO: 6;
 - (b) et polypeptid, der kodes for af et polynukleotid, som udviser mindst 90%, mindst 91%, mindst 92%, mindst 93%, mindst 94%, mindst 95%, mindst 96%, mindst 97%, mindst 98%, mindst 99% eller 100% sekvensidentitet med det modne polypeptids kodningssekvens ifølge SEQ ID NO: 5 eller cDNA-sekvensen deraf;
 - (c) et fragment af polypeptidet ifølge (a) eller (b), der har xylanaseaktivitet.
2. Polypeptid ifølge krav 1, som omfatter eller består af SEQ ID NO: 6 eller det modne polypeptid ifølge SEQ ID NO: 6.
3. Polypeptid ifølge krav 2, hvor det modne polypeptid er aminosyrerne 21 til 405 ifølge SEQ ID NO: 6.
4. Polypeptid ifølge krav 1, som er et fragment af SEQ ID NO: 6, hvor fragmentet har xylanaseaktivitet.
5. Isoleret polypeptid, som omfatter et katalytisk domæne valgt fra gruppen bestående af:
 - (a) et katalytisk domæne, som udviser mindst 90% sekvensidentitet med det katalytiske domæne ifølge SEQ ID NO: 6; og
 - (b) et fragment af et katalytisk domæne ifølge (a), der har xylanaseaktivitet.
6. Polypeptid ifølge krav 5, som omfatter eller består af det katalytiske domæne ifølge SEQ ID NO: 6.
7. Polypeptid ifølge krav 6, hvor det katalytiske domæne er 21 til 337 ifølge SEQ ID NO: 6.
8. Polypeptid ifølge et hvilket som helst af kravene 5-7, som endvidere omfatter et cellulosebindende domæne.
9. Polypeptid ifølge et hvilket som helst af kravene 1-8, der kodes for af polynukleotidet, som er indeholdt i *Talaromyces leycettanus*-stammen CBS398.68.
10. Sammensætning, som omfatter polypeptidet ifølge et hvilket som helst af kravene 1-9.

11. Isoleret polynukleotid, der koder for polypeptidet ifølge et hvilket som helst af kravene 1-9.

12. Nukleinsyrekonstruktion eller ekspressionsvektor, som omfatter polynukleotidet ifølge krav 11, hvilket polynukleotid er operativt forbundet til en eller flere kontrolsekvenser, der styrer fremstillingen af polypeptidet i en ekspressionsvært.

13. Rekombinant værtscelle, som omfatter polynukleotidet ifølge krav 11, hvilket polynukleotid er operativt forbundet til en eller flere kontrolsekvenser, der styrer fremstillingen af polypeptidet.

14. Fremgangsmåde til fremstilling af et polypeptid med xylanaseaktivitet, hvilken fremgangsmåde omfatter:

(a) dyrkning af værtscellen ifølge krav 13 under betingelser, som er befordrende for fremstilling af polypeptidet; og

(b) indvinding af polypeptidet.

15. Transgen plante, plantedel eller plantecelle, som omfatter et polynukleotid, hvilket polynukleotid koder for polypeptidet ifølge et hvilket som helst af kravene 1-9.

16. Fremgangsmåde til fremstilling af et polypeptid med xylanaseaktivitet, hvilken fremgangsmåde omfatter:

(a) dyrkning af den transgene plante eller plantecelle ifølge krav 15 under betingelser, som er befordrende for fremstilling af polypeptidet; og

(b) indvinding af polypeptidet.

17. Fremgangsmåde til nedbrydning af et celluloseholdigt materiale eller xylanholdigt materiale, hvilken fremgangsmåde omfatter: behandling af det celluloseholdige materiale eller xylanholdige materiale med en enzymsammensætning i nærværelse af polypeptidet med xylanaseaktivitet ifølge et hvilket som helst af kravene 1-9.

18. Fremgangsmåde til fremstilling af et fermenteringsprodukt, hvilken fremgangsmåde omfatter:

(a) forsukring af et celluloseholdigt materiale eller xylanholdigt materiale med en enzymsammensætning i nærværelse af polypeptidet med xylanaseaktivitet ifølge et hvilket som helst af kravene 1-9;

(b) fermentering af det forsukrede celluloseholdige materiale eller xylanholdige materiale med en eller flere fermenterende mikroorganismer for at fremstille fermenteringsproduktet; og

(c) indvinding af fermenteringsproduktet fra fermenteringen.

19. Fremgangsmåde til fermentering af et celluloseholdigt materiale eller xylanholdigt materiale, hvilken fremgangsmåde omfatter: fermentering af det celluloseholdige materiale eller xylanholdige materiale med en eller flere fermenterende mikroorganismer, hvor det celluloseholdige materiale eller xylanholdige materiale er forsuret med en
- 5 enzymsammensætning i nærværelse af polypeptidet med xylanaseaktivitet ifølge et hvilket som helst af kravene 1-9.