

(19)



(11)

EP 3 338 793 A1

(12)

EUROPEAN PATENT APPLICATION

(43) Date of publication:

27.06.2018 Bulletin 2018/26

(51) Int Cl.:

A61K 39/00 ^(2006.01) **A61K 39/395** ^(2006.01)
C12P 21/08 ^(2006.01) **C07K 16/00** ^(2006.01)
C07K 16/18 ^(2006.01) **C07K 16/28** ^(2006.01)
G01N 33/574 ^(2006.01)

(21) Application number: **17203670.9**

(22) Date of filing: **27.08.2014**

(84) Designated Contracting States:

**AL AT BE BG CH CY CZ DE DK EE ES FI FR GB
GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO
PL PT RO RS SE SI SK SM TR**

Designated Extension States:

BA ME

- **BHEDDAH, Sheila**
San Francisco, CA 94110 (US)
- **LOPEZ-MOLINA, Javier**
New York, NY 10038 (US)
- **SAUNDERS, Laura**
San Francisco, CA 94107 (US)

(30) Priority: **28.08.2013 US 201361871289 P**

(62) Document number(s) of the earlier application(s) in
accordance with Art. 76 EPC:

14839298.8 / 3 038 634

(71) Applicant: **AbbVie Stemcentrx LLC**
North Chicago, IL 60064-6400 (US)

(72) Inventors:

- **LIU, David**
South San Francisco, CA 94112 (US)
- **ROKKAM, Deepti**
Sunnyvale, CA 94087 (US)

(74) Representative: **Evenson, Jane Harriet et al**
Mathys & Squire LLP
The Shard
32 London Bridge Street
London SE1 9SG (GB)

Remarks:

- Claims filed after the date of receipt of the divisional application (Rule 68(4) EPC).
- This application was filed on 24-11-2017 as a divisional application to the application mentioned under INID code 62.

(54) **NOVEL SEZ6 MODULATORS AND METHODS OF USE**

(57) Novel modulators, including antibodies and derivatives thereof, and methods of using such modulators to treat proliferative disorders are provided.

EP 3 338 793 A1

Description**CROSS REFERENCED APPLICATIONS**

5 **[0001]** This application claims priority from U. S. Provisional Patent Application Ser. No. 61/871,289 filed on August 28, 2013, which is incorporated herein by reference in its entirety.

SEQUENCE LISTING

10 **[0002]** The instant application contains a sequence listing which has been submitted in ASCII format via EFS-Web and is hereby incorporated by reference in its entirety. Said ASCII copy, created on August 27, 2014, is named "S69697 1130WO SEQL 082714 for filing" and is 462,073 bytes in size.

FIELD OF THE INVENTION

15 **[0003]** This application generally relates to novel compounds, compositions and methods of their use in diagnosing, preventing, treating or ameliorating proliferative disorders and any expansion, recurrence, relapse or metastasis thereof. In a broad aspect, the present invention relates to the use of seizure related 6 homolog (SEZ6) modulators, including anti-SEZ6 antibodies and fusion constructs, for the treatment, diagnosis or prophylaxis of neoplastic disorders. Selected
20 embodiments of the present invention provide for the use of such SEZ6 modulators, including antibody drug conjugates, for the immunotherapeutic treatment of malignancies preferably comprising a reduction in tumor initiating cell frequency. Particularly preferred embodiments of the invention comprise modulators that associate with specific epitopes and use of the disclosed modulators to treat certain patient populations such as those suffering from medullary thyroid cancer, small cell lung cancer and patients that are resistant to standard of care platinum based agents.

BACKGROUND OF THE INVENTION

25 **[0004]** Stem and progenitor cell differentiation and cell proliferation are normal ongoing processes that act in concert to support tissue growth during organogenesis and cell replacement and repair of most tissues during the lifetime of all living organisms. In the normal course of events cellular differentiation and proliferation is controlled by numerous factors and signals that are generally balanced to maintain cell fate decisions and tissue architecture. Thus, to a large extent it is this controlled microenvironment that regulates cell division and tissue maturation where signals are properly generated based on the needs of the organism. In this regard cell proliferation and differentiation normally occur only as necessary for the replacement of damaged or dying cells or for growth. Unfortunately, disruption of cell proliferation and/or differ-
30 entiation can result from a myriad of factors including, for example, the under- or overabundance of various signaling chemicals, the presence of altered microenvironments, genetic mutations or some combination thereof. When normal cellular proliferation and/or differentiation is disturbed or somehow disrupted it can lead to various diseases or disorders including proliferative disorders such as cancer.

35 **[0005]** Conventional treatments for cancer include chemotherapy, radiotherapy, surgery, immunotherapy (e.g., biological response modifiers, vaccines or targeted therapeutics) or combinations thereof. Unfortunately, certain cancers are non-responsive or minimally responsive to such treatments. For example, in some patients tumors exhibit gene mutations that render them non-responsive despite the general effectiveness of selected therapies. Moreover, depending on the type of cancer and what form it takes some available treatments, such as surgery, may not be viable alternatives. Limitations inherent in current standard of care therapeutics are particularly evident when attempting to treat patients
40 who have undergone previous treatments and have subsequently relapsed. In such cases the failed therapeutic regimens and resulting patient deterioration may contribute to refractory tumors which often manifest themselves as a relatively aggressive disease that ultimately proves to be incurable. Although there have been great improvements in the diagnosis and treatment of cancer over the years, overall survival rates for many solid tumors have remained largely unchanged due to the failure of existing therapies to prevent relapse, tumor recurrence and metastases. Thus, it remains a challenge
45 to develop more targeted and potent therapies for proliferative disorders.

SUMMARY OF THE INVENTION

50 **[0006]** These and other objectives are provided for by the present invention which, in a broad sense, is directed to methods, compounds, compositions and articles of manufacture that may be used in the treatment of SEZ6 associated disorders (e.g., proliferative disorders or neoplastic disorders). To that end, the present invention provides novel seizure related 6 homolog (or SEZ6) modulators that effectively target tumor cells and/or cancer stem cells and may be used to treat patients suffering from a wide variety of malignancies. As will be discussed in more detail herein, there are at
55

least two naturally occurring SEZ6 isoforms or variants and the disclosed modulators may comprise or associate selectively with one isoform or the other or with both. Moreover, in certain embodiments the disclosed SEZ6 modulators may further react with one or more SEZ family members (e.g., SEZ6L or SEZ6L2) or, in other embodiments, may be generated and selected for so as to exclusively associate or react with SEZ6 isoform(s). In preferred embodiments the invention is more particularly directed to isolated SEZ6 modulators comprising antibodies (i.e., antibodies that immunopreferentially bind, react with or associate with at least one isoform of SEZ6) that, in particularly preferred embodiments, are associated or conjugated to one or more cytotoxic agents or therapeutic moieties, for example, auristatins, amanitins and pyrrolobenzodiazepines. Moreover, as discussed extensively below, such modulators may be used to provide pharmaceutical compositions useful for the prophylaxis, diagnosis or treatment of proliferative disorders.

[0007] In selected embodiments of the invention, SEZ6 modulators may comprise a SEZ6 polypeptide or fragments thereof, either in an isolated form or fused or associated with other moieties (e.g., Fc-SEZ6, PEG-SEZ6 or SEZ6 associated with a targeting moiety). In other selected embodiments SEZ6 modulators may comprise SEZ6 antagonists which, for the purposes of the instant application, shall be held to mean any construct or compound that recognizes, competes, interacts, binds or associates with SEZ6 and neutralizes, eliminates, reduces, sensitizes, reprograms, inhibits or controls the growth of neoplastic cells including tumor initiating cells. In preferred embodiments the SEZ6 modulators of the instant invention comprise anti-SEZ6 antibodies, or fragments or derivatives thereof, that have unexpectedly been found to silence, neutralize, reduce, decrease, deplete, moderate, diminish, reprogram, eliminate, or otherwise inhibit the ability of tumor initiating cells to propagate, maintain, expand, proliferate or otherwise facilitate the survival, recurrence, regeneration and/or metastasis of neoplastic cells. In particularly preferred embodiments the antibodies or immunoreactive fragments may be associated with or conjugated to one or more anti-cancer agents (e.g., a cytotoxic agent).

[0008] With regard to such modulators it will be appreciated that compatible antibodies may take on any one of a number of forms including, for example, polyclonal and monoclonal antibodies, chimeric, CDR grafted, humanized and human antibodies and immunoreactive fragments and/or variants of each of the foregoing. Preferred embodiments will comprise antibodies that are relatively non-immunogenic such as humanized or fully human constructs. Of course, in view of the instant disclosure those skilled in the art could readily identify one or more complementarity determining regions (CDRs) associated with heavy and light chain variable regions of SEZ6 antibody modulators and use those CDRs to engineer or fabricate chimeric, humanized or CDR grafted antibodies without undue experimentation. Accordingly, in certain preferred embodiments the SEZ6 modulator comprises an antibody that incorporates one or more CDRs derived from the light (FIG. 10A) or heavy (FIG. 10B) contiguous chain murine variable regions (SEQ ID NOS: 20-169) set forth therein. Such CDR grafted variable regions having a human framework and variants thereof are also shown in FIG. 10 comprising SEQ ID NOS: 170-199. In preferred embodiments such antibodies will comprise monoclonal antibodies and, in even more preferred embodiments, will comprise chimeric, CDR grafted or humanized antibodies.

[0009] Exemplary nucleic acid sequences encoding each of the amino acid sequences set forth in FIGS. 10A and 10B are set forth in the appended sequence listing and comprise SEQ ID NOS: 220 to 399. In this respect it will be appreciated that the invention further comprises nucleic acid molecules (and associated constructs, vectors and host cells) encoding disclosed antibody variable region amino acid sequences including those set forth in the sequence listing.

[0010] More particularly, in selected embodiments compatible SEZ6 modulators may comprise an antibody having a light chain variable region and a heavy chain variable region wherein said light chain variable region comprises an amino acid sequence having at least 60% identity to an amino acid sequence selected from the group consisting of amino acid sequences as set forth in SEQ ID NO: 20, SEQ ID NO: 22, SEQ ID NO: 24, SEQ ID NO: 26, SEQ ID NO: 28, SEQ ID NO: 30, SEQ ID NO: 32, SEQ ID NO: 34, SEQ ID NO: 36, SEQ ID NO: 38, SEQ ID NO: 40, SEQ ID NO: 42, SEQ ID NO: 44, SEQ ID NO: 46, SEQ ID NO: 48, SEQ ID NO: 50, SEQ ID NO: 52, SEQ ID NO: 54, SEQ ID NO: 56, SEQ ID NO: 58, SEQ ID NO: 60, SEQ ID NO: 62, SEQ ID NO: 64, SEQ ID NO: 66, SEQ ID NO: 68, SEQ ID NO: 70, SEQ ID NO: 72, SEQ ID NO: 74, SEQ ID NO: 76, SEQ ID NO: 78, SEQ ID NO: 80, SEQ ID NO: 82, SEQ ID NO: 84, SEQ ID NO: 86, SEQ ID NO: 88, SEQ ID NO: 90, SEQ ID NO: 92, SEQ ID NO: 94, SEQ ID NO: 96, SEQ ID NO: 98, SEQ ID NO: 100, SEQ ID NO: 102, SEQ ID NO: 104, SEQ ID NO: 106, SEQ ID NO: 108, SEQ ID NO: 110, SEQ ID NO: 112, SEQ ID NO: 114, SEQ ID NO: 116, SEQ ID NO: 118, SEQ ID NO: 120, SEQ ID NO: 122, SEQ ID NO: 124, SEQ ID NO: 126, SEQ ID NO: 128, SEQ ID NO: 130, SEQ ID NO: 132, SEQ ID NO: 134, SEQ ID NO: 136, SEQ ID NO: 138, SEQ ID NO: 140, SEQ ID NO: 142, SEQ ID NO: 144, SEQ ID NO: 146, SEQ ID NO: 148, SEQ ID NO: 150, SEQ ID NO: 152, SEQ ID NO: 154, SEQ ID NO: 156, SEQ ID NO: 158, SEQ ID NO: 160, SEQ ID NO: 162, SEQ ID NO: 164, SEQ ID NO: 166 and SEQ ID NO: 168 and wherein said heavy chain variable region comprises an amino acid sequence having at least 60% identity to an amino acid sequence selected from the group consisting of amino acid sequences as set forth in SEQ ID NO: 21, SEQ ID NO: 23, SEQ ID NO: 25, SEQ ID NO: 27, SEQ ID NO: 29, SEQ ID NO: 31, SEQ ID NO: 33, SEQ ID NO: 35, SEQ ID NO: 37, SEQ ID NO: 39, SEQ ID NO: 41, SEQ ID NO: 43, SEQ ID NO: 45, SEQ ID NO: 47, SEQ ID NO: 49, SEQ ID NO: 51, SEQ ID NO: 53, SEQ ID NO: 55, SEQ ID NO: 57, SEQ ID NO: 59, SEQ ID NO: 61, SEQ ID NO: 63, SEQ ID NO: 65, SEQ ID NO: 67, SEQ ID NO: 69, SEQ ID NO: 71, SEQ ID NO: 73, SEQ ID NO: 75, SEQ ID NO: 77, SEQ ID NO: 79, SEQ ID NO: 81, SEQ ID NO: 83, SEQ ID NO: 85, SEQ ID NO: 87, SEQ ID NO: 89, SEQ ID NO: 91, SEQ ID NO: 93, SEQ ID NO: 95, SEQ ID NO: 97, SEQ ID NO: 99, SEQ ID NO: 101, SEQ

ID NO: 103, SEQ ID NO: 105, SEQ ID NO: 107, SEQ ID NO: 109, SEQ ID NO: 111, SEQ ID NO: 113, SEQ ID NO: 115, SEQ ID NO: 117, SEQ ID NO: 119, SEQ ID NO: 121, SEQ ID NO: 123, SEQ ID NO: 125, SEQ ID NO: 127, SEQ ID NO: 129, SEQ ID NO: 131, SEQ ID NO: 133, SEQ ID NO: 135, SEQ ID NO: 137, SEQ ID NO: 139, SEQ ID NO: 141, SEQ ID NO: 143, SEQ ID NO: 145, SEQ ID NO: 147, SEQ ID NO: 149, SEQ ID NO: 151, SEQ ID NO: 153, SEQ ID NO: 155, SEQ ID NO: 157, SEQ ID NO: 159, SEQ ID NO: 161, SEQ ID NO: 163, SEQ ID NO: 165, SEQ ID NO: 167 and SEQ ID NO: 169. In other preferred embodiments the selected modulators will comprise heavy and light chain variable regions that comprise 65, 70, 75 or 80% identity to the aforementioned murine sequences. In still other embodiments the modulators will comprise heavy and light chain variable regions that comprise 85, 90 or even 95% identity to the disclosed murine sequences.

[0011] Of course, in view of the instant disclosure those skilled in the art could readily identify CDRs associated with each of the aforementioned heavy and light chain variable regions and use those CDRs to engineer or fabricate chimeric, humanized or CDR grafted antibodies without undue experimentation. In this regard several website databases are available that automatically designate CDRs and framework regions (as per any of the commonly used numbering systems) upon entry of the subject heavy or light chain variable region nucleic acid or amino acid sequence. As such, in selected embodiments the present invention is directed to anti-SEZ6 antibodies comprising one or more CDRs derived from a variable region sequence set forth in FIG. 10A or FIG. 10B. In preferred embodiments such antibodies will comprise monoclonal antibodies and, in even more preferred embodiments will comprise chimeric, CDR grafted or humanized antibodies. As discussed in more detail below still other embodiments will comprise such antibodies conjugated or associated with one or more cytotoxic agents.

[0012] Another aspect of the invention comprises modulators obtained or derived from SC17.1, SC17.2, SC17.3, SC17.4, SC17.8, SC17.9, SC17.10, SC17.11, SC17.14, SC17.15, SC17.16, SC17.17, SC17.18, SC17.19, SC17.22, SC17.24, SC17.27, SC17.28, SC17.29, SC17.30, SC17.32, SC17.34, SC17.35, SC17.36, SC17.38, SC17.39, SC17.40, SC17.41, SC17.42, SC17.45, SC17.46, SC17.47, SC17.49, SC17.50, SC17.53, SC17.54, SC17.56, SC17.57, SC17.59, SC17.61, SC17.63, SC17.71, SC17.72, SC17.74, SC17.76, SC17.77, SC17.79, SC17.81, SC17.82, SC17.84, SC17.85, SC17.87, SC17.89, SC17.90, SC17.91, SC17.93, SC17.95, SC17.97, SC17.99, SC17.102, SC17.114, SC17.115, SC17.120, SC17.121, SC17.122, SC17.140, SC17.151, SC17.156, SC17.161, SC17.166, SC17.187, SC17.191, SC17.193, SC17.199 and SC17.200.

[0013] In yet other compatible embodiments the instant invention will comprise the CDR grafted or humanized SEZ6 modulators hSC17.16, hSC17.17, hSC 17.24, hSC 17.28, SC17.34, hSC17.46, SC17.151, SC17.155, SC17.156, SC17.161 and SC17.200. Still other embodiments are directed to a SEZ6 modulator comprising a humanized antibody wherein said humanized antibody comprises a light chain variable region and a heavy chain variable region wherein said light chain variable region comprises an amino acid sequence having at least 60% identity to an amino acid sequence selected from the group consisting of amino acid sequences as set forth in SEQ ID NO: 170, SEQ ID NO: 172, SEQ ID NO: 174, SEQ ID NO: 176, SEQ ID NO: 178, SEQ ID NO: 180, SEQ ID NO: 182, SEQ ID NO: 184, SEQ ID NO: 186, SEQ ID NO: 188 and SEQ ID NO: 190 and wherein said heavy chain variable region comprises an amino acid sequence having at least 60% identity to an amino acid sequence selected from the group consisting of amino acid sequences as set forth in SEQ ID NO: 171, SEQ ID NO: 173, SEQ ID NO: 175, SEQ ID NO: 177, SEQ ID NO: 179 and SEQ ID NO: 181, SEQ ID NO: 183, SEQ ID NO: 185, SEQ ID NO: 187, SEQ ID NO: 189 and SEQ ID NO: 191. Additionally, certain humanized variants of light (SEQ ID NO: 192) and heavy (SEQ ID NO: 193, SEQ ID NO: 194, SEQ ID NO: 195, SEQ ID NO: 196, SEQ ID NO: 197, SEQ ID NO: 198 and SEQ ID NO: 199) chain variable regions are provided in accordance with the teachings herein. Moreover, as described immediately above nucleic acid sequences encoding the exemplified humanized heavy and light chain variable regions are set forth in the appended sequence listing as SEQ ID NOS: 370 - 399.

[0014] Besides the aforementioned aspects, other preferred embodiments of the instant invention will comprise SEZ6 modulators associated or conjugated to one or more drugs or therapeutic moieties (e.g. auristatins, amanitins and pyrrolbenzodiazepines) to provide modulator conjugates that may be particularly effective in treating proliferative disorders (alone or in combination with other pharmaceutically active agents). More generally, once the modulators of the invention have been fabricated and selected they may be linked with, fused to, conjugated to (e.g., covalently or non-covalently) or otherwise associated with pharmaceutically active or diagnostic moieties or biocompatible modifiers. As used herein the term "conjugate" or "modulator conjugate" or "antibody conjugate" will be used broadly and held to mean any biologically active or detectable molecule or drug associated with the disclosed modulators regardless of the method of association. In this respect it will be understood that such conjugates may, in addition to the disclosed modulators, comprise peptides, polypeptides, proteins, prodrugs which are metabolized to an active agent in vivo, polymers, nucleic acid molecules, small molecules, binding agents, mimetic agents, synthetic drugs, inorganic molecules, organic molecules and radioisotopes. Moreover, as indicated above the selected drug or therapeutic moiety may be covalently or non-covalently associated with, or linked to, the modulator and exhibit various stoichiometric molar ratios depending, at least in part, on the method used to effect the conjugation.

[0015] Particularly preferred aspects of the instant invention will comprise antibody modulator conjugates or antibody-drug conjugates that may be used for the diagnosis and/or treatment of proliferative disorders. Such conjugates may

be represented by the formula M-[L-D]_n where M stands for a disclosed modulator or target binding moiety, L is an optional linker or linker unit, D is a compatible drug or prodrug and n is an integer from about 1 to about 20. It will be appreciated that, unless otherwise dictated by context, the terms "antibody-drug conjugate" or "ADC" or the formula M-[L-D]_n shall be held to encompass conjugates comprising both therapeutic and diagnostic moieties. In such embodiments antibody-drug conjugate compounds will typically comprise anti-SEZ6 as the modulator unit (M), a therapeutic or diagnostic moiety (D), and optionally a linker (L) that joins the drug and the antigen binding agent. Modulators can be directly or indirectly conjugated to a therapeutic or diagnostic moiety. Modulators that are joined to the therapeutic or diagnostic moiety with a linker are referred to as being "indirectly conjugated" whereas antibodies that are joined to the therapeutic or diagnostic moiety in the absence of a linker are referred to as being "directly conjugated". In a preferred embodiment, the antibody is a SEZ6 mAb comprising at least one CDR from the heavy and light chain variable regions as described above.

[0016] As previously indicated one aspect of the invention may comprise the unexpected association of SEZ6 polypeptides with cancer stem cells. Thus, in certain other embodiments the invention will comprise a SEZ6 modulator that reduces the frequency of tumor initiating cells upon administration to a subject. Preferably the reduction in frequency will be determined using *in vitro* or *in vivo* limiting dilution analysis. In particularly preferred embodiments such analysis may be conducted using *in vivo* limiting dilution analysis comprising transplant of live human tumor cells into immunocompromised mice. Alternatively, the limiting dilution analysis may be conducted using *in vitro* limiting dilution analysis comprising limiting dilution deposition of live human tumor cells into *in vitro* colony supporting conditions. In either case, the analysis, calculation or quantification of the reduction in frequency will preferably comprise the use of Poisson distribution statistics to provide an accurate accounting. It will be appreciated that, while such quantification methods are preferred, other, less labor intensive methodology such as flow cytometry or immunohistochemistry may also be used to provide the desired values and, accordingly, are expressly contemplated as being within the scope of the instant invention. In such cases the reduction in frequency may be determined using flow cytometric analysis or immunohistochemical detection of tumor cell surface markers known to enrich for tumor initiating cells.

[0017] As such, another preferred embodiment of the instant invention comprises a method of treating a SEZ6 associated disorder comprising administering a therapeutically effective amount of a SEZ6 modulator to a subject in need thereof whereby the frequency of tumor initiating cells is reduced. Preferably the SEZ6 associated disorder comprises a neoplastic disorder. Again, the reduction in the tumor initiating cell frequency will preferably be determined using *in vitro* or *in vivo* limiting dilution analysis.

[0018] In this regard it will be appreciated that the present invention is based, at least in part, upon the discovery that SEZ6 immunogens are associated with tumor perpetuating cells (i.e., cancer stem cells) that are involved in the etiology of various neoplasia. More specifically, the instant application unexpectedly demonstrates that the administration of various exemplary SEZ6 modulators can mediate, reduce, deplete, inhibit or eliminate tumorigenic signaling by tumor initiating cells (i.e., reduce the frequency of tumor initiating cells). This reduced signaling, whether by depletion, neutralization, reduction, elimination, reprogramming or silencing of the tumor initiating cells or by modifying tumor cell morphology (e.g., induced differentiation, niche disruption), in turn allows for the more effective treatment of SEZ6 associated disorders by inhibiting tumorigenesis, tumor maintenance, expansion and/or metastasis and recurrence.

[0019] Besides the aforementioned association with cancer stem cells, there is evidence that SEZ6 isoforms may be implicated in the growth, recurrence or metastatic potential of tumors comprising neuroendocrine features. For the purposes of the instant invention such tumors will comprise neuroendocrine tumors (e.g. small cell lung cancer and medullary thyroid tumors) and pseudo neuroendocrine tumors. Intervention in the proliferation of such tumorigenic cells using the novel SEZ6 modulators described herein, may thereby ameliorate or treat a disorder by more than one mechanism (i.e., tumor initiating cell reduction and disruption of oncogenic pathway signaling) to provide additive or synergistic effects. Still other preferred embodiments may take advantage of the cellular internalization of cell surface SEZ6 to deliver a modulator mediated anti-cancer agent. In this regard it will be appreciated that the present invention is not limited by any particular mechanism of action but rather encompasses the broad use of the disclosed modulators to treat SEZ6 associated disorders (including various neoplasia).

[0020] Thus, in other embodiments the present invention will comprise the use of the disclosed modulators to treat tumors comprising neuroendocrine features, e.g. small cell lung cancer and medullary thyroid tumors in a subject in need thereof. Of course the same modulators may be used for the prophylaxis, prognosis, diagnosis, theragnosis, inhibition or maintenance therapy of these same tumors.

[0021] It has further been discovered that the disclosed modulators are effective in treating patients that are suffering from small cell lung cancer or medullary thyroid cancer. Moreover, as discussed in more detail below and set forth in the Examples the anti-SEZ6 antibodies and antibody drug conjugates of the instant invention are particularly effective in treating certain patient populations such as those suffering from forms of small cell lung cancer that are resistant to standard of care platinum based agents (e.g., carboplatin, cisplatin or oxaliplatin). Accordingly, in selected embodiments the present invention comprises a method of treating a patient suffering from platinum resistant small cell lung cancer comprising the step of administering a SEZ6 modulator.

[0022] Other facets of the instant invention exploit the ability of the disclosed modulators to potentially disrupt oncogenic pathways while simultaneously silencing tumor initiating cells. Such multi-active SEZ6 modulators (e.g., SEZ6 antagonists) may prove to be particularly effective when used in combination with standard of care anti-cancer agents or debulking agents. Accordingly preferred embodiments of the instant invention comprise using the disclosed modulators as anti-metastatic agents for maintenance therapy following initial treatments. In addition, two or more SEZ6 antagonists (e.g. antibodies that specifically bind to two discrete epitopes on SEZ6) may be used in combination in accordance with the present teachings. Moreover, as discussed in some detail below, the SEZ6 modulators of the present invention may be used in a conjugated or unconjugated state and, optionally, as a sensitizing agent in combination with a variety of chemical or biological anti-cancer agents.

[0023] Accordingly another preferred embodiment of the instant invention comprises a method of sensitizing a tumor in a subject for treatment with an anti-cancer agent comprising the step of administering a SEZ6 modulator to said subject. Other embodiments comprise a method of reducing metastasis or tumor recurrence following treatment comprising administering a SEZ6 modulator to a subject in need thereof. In a particularly preferred aspect of the invention the SEZ6 modulator will specifically result in a reduction of tumor initiating cell frequency as determined using *in vitro* or *in vivo* limiting dilution analysis.

[0024] More generally preferred embodiments of the invention comprise a method of treating a SEZ6 associated disorder in a subject in need thereof comprising the step of administering a SEZ6 modulator to the subject. In particularly preferred embodiments the SEZ6 modulator will be associated (e.g., conjugated) with an anti-cancer agent. In yet other embodiments the SEZ6 modulator will internalize following association or binding with SEZ6 on or near the surface of the cell. Moreover the beneficial aspects of the instant invention, including any disruption of signaling pathways and collateral benefits, may be achieved whether the subject tumor tissue exhibits elevated levels of SEZ6 or reduced or depressed levels of SEZ6 as compared with normal adjacent tissue. Particularly preferred embodiments will comprise the treatment of disorders exhibiting elevated levels of SEZ6 on tumorigenic cells as compared to normal tissue or non-tumorigenic cells.

[0025] In yet another aspect the present invention will comprise a method of treating a subject suffering from a neoplastic disorder comprising the step of administering a therapeutically effective amount of at least one internalizing SEZ6 modulator. Preferred embodiments will comprise the administration of internalizing antibody modulators wherein, in other selected embodiments, the internalizing antibody modulators are conjugated or associated with a cytotoxic agent.

[0026] Other embodiments are directed to a method of treating a subject suffering from a SEZ6 associated disorder comprising the step of administering a therapeutically effective amount of at least one depleting SEZ6 modulator.

[0027] In yet another embodiment the present invention provides methods of maintenance therapy wherein the disclosed effectors or modulators are administered over a period of time following an initial procedure (e.g., chemotherapy, radiation or surgery) designed to remove at least a portion of the tumor mass. Such therapeutic regimens may be administered over a period of weeks, a period of months or even a period of years wherein the SEZ6 modulators may act prophylactically to inhibit metastasis and/or tumor recurrence. In yet other embodiments the disclosed modulators may be administered in concert with known debulking regimens to prevent or retard metastasis, tumor maintenance or recurrence.

[0028] It will further be appreciated that the SEZ6 modulators of the instant invention may be generated and selected to react with known isoform(s) of SEZ6 or a single isoform of the protein or, conversely, may comprise a pan-SEZ6 modulator that reacts or associates with at least one additional SEZ6 family member (e.g., SEZ6L or SEZ6L2 and isoforms thereof) in addition to SEZ6. More specifically, as disclosed herein preferred modulators such as antibodies may be generated and selected so that they react with domains (or epitopes therein) that are exhibited by SEZ6 only or with domains that are at least somewhat conserved across two or more of the SEZ6 family members.

[0029] In yet other preferred embodiments the modulators will associate or bind to a specific epitope, portion, motif or domain of SEZ6. As will be discussed in some detail below both SEZ6 isoforms incorporate an identical extracellular region (see FIG. 1E) comprising at least an N-terminal domain, two alternating Sushi and CUB domains, and three additional tandem Sushi domain repeats. In addition the SEZ6 protein comprises a transmembrane domain and a cytoplasmic domain. Accordingly, in certain embodiments the modulators will bind or associate with the N-terminal domain of SEZ6 (i.e. amino acids 1-335 in the mature protein) or to an epitope therein. Other aspects of the instant invention comprise modulators that associate or bind to a specific epitope located in a particular Sushi domain of SEZ6. In this regard the particular modulator may associate or bind to an epitope located in Sushi Domain 1 (amino acids 336-395), Sushi Domain 2 (amino acids 511-572), Sushi Domain 3 (amino acids 690-748), Sushi Domain 4 (amino acids 750-813) or Sushi Domain 5 (amino acids 817-878). Other aspects of the instant invention comprise modulators that associate or bind to a specific epitope located in a particular CUB-like domain of SEZ6. In this regard the particular modulator may associate or bind to an epitope located in CUB Domain 1 (amino acids 397-508) or CUB Domain 2 (amino acids 574-685). In a further embodiment the antibodies of the invention may bind to certain epitopes on SEZ6.

[0030] In one embodiment, the invention provides for an isolated antibody that specifically binds to an epitope on a SEZ6 protein, wherein the epitope comprises amino acid residues selected from the group consisting of (i) residues

R762, L764, Q777, I779, D781 and Q782; (ii) residues R342 and K389 and (iii) residues T352, S353 and H375.

[0031] In another embodiment the invention provides for an antibody drug conjugate comprising an antibody conjugated directly or indirectly to a therapeutic moiety, wherein the antibody specifically binds to an epitope on a SEZ6 protein, wherein the epitope comprises amino acid residues selected from the group consisting of (i) residues R762, L764, Q777, I779, D781 and Q782; (ii) residues R342 and K389 and (iii) residues T352, S353 and H375.

[0032] Of course it will be appreciated that each of the aforementioned domains may comprise more than one epitope and may be associated with more than one bin. With regard to modulator or antibody "bins" it will be appreciated that the SEZ6 antigen may be analyzed or mapped through competitive antibody binding using art recognized techniques to define specific bins located along the protein. While discussed in more detail herein and shown in Examples 9 and 10 below, two antibodies (one of which may be termed a "reference antibody," "bin delineating antibody" or "delineating antibody") may be considered to be in the same bin if they substantially compete with each other for binding to the target antigen. In such cases the subject antibody epitopes may be identical, substantially identical or close enough (either in a linear sense where they are separated by a few amino acids or conformationally) so that both antibodies are sterically or electrostatically inhibited or precluded from binding to the antigen. Such defined bins may be generally associated with certain SEZ6 domains (e.g. the reference antibody will bind with an epitope contained in a specific domain) though the correlation is not always precise (e.g., there may be more than one bin in a domain or the bin may be defined conformationally and comprise more than one domain). It will be appreciated that those skilled in the art can readily determine the relationship between the SEZ6 domains and empirically determined bins.

[0033] With regard to the present invention competitive binding analysis using art-recognized techniques (e.g., ELISA, surface plasmon resonance or bio-layer interferometry) defined at least seven distinct bins, each of which was found to contain a number of antibody modulators. For the purposes of the instant disclosure the seven bins were termed bins A-F and bin U. Bins A-F are unique bins and the antibodies contained in each of these bins compete with each other for binding to the SEZ6 protein. Bin U contains antibodies that do not compete with antibodies in Bins A-F, but may compete for binding with each other. Thus, in selected embodiments the present invention will comprise a modulator residing in a bin selected from the group consisting of bin A, bin B, bin C, bin D, bin E, bin F, and bin U. In other embodiments the present invention comprises a modulator residing in a bin defined by a reference antibody selected from the group consisting of SC17.1, SC17.2, SC17.3, SC17.4, SC17.8, SC17.9, SC17.10, SC17.11, SC17.14, SC17.15, SC17.16, SC17.17, SC17.18, SC17.19, SC17.22, SC17.24, SC17.27, SC17.28, SC17.29, SC17.30, SC17.32, SC17.34, SC17.35, SC17.36, SC17.38, SC17.39, SC17.40, SC17.41, SC17.42, SC17.45, SC17.46, SC17.47, SC17.49, SC17.50, SC17.53, SC17.54, SC17.56, SC17.57, SC17.59, SC17.61, SC17.63, SC17.71, SC17.72, SC17.74, SC17.76, SC17.77, SC17.79, SC17.81, SC17.82, SC17.84, SC17.85, SC17.87, SC17.89, SC17.90, SC17.91, SC17.93, SC17.95, SC17.97, SC17.99, SC17.102, SC17.114, SC17.115, SC17.120, SC17.121, SC17.122, SC17.140, SC17.151, SC17.156, SC17.161, SC17.166, SC17.187, SC17.191, SC17.193, SC17.199 and SC17.200. In still other embodiments the invention will comprise modulators from bin A, modulators from bin B, modulators from bin C, modulators from bin D, modulators from bin E, modulators from bin F or modulators from bin U. Yet other preferred embodiments will comprise a reference antibody modulator and any antibody that competes with the reference antibody.

[0034] The term "compete" or "competing antibody" when used in the context of the disclosed modulators means binding competition between antibodies as determined by an assay in which a reference antibody or immunologically functional fragment substantially prevents or inhibits (e.g., greater than 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85% or 90%.) specific binding of a test antibody to a common antigen. Compatible methods for determining such competition comprise art known techniques such as, for example, bio-layer interferometry, surface plasmon resonance, flow cytometry, competitive ELISA, etc.

[0035] In a selected embodiment the invention comprises a pan-SEZ6 modulator that associates with SEZ6 and at least one other SEZ6 family member (e.g., SEZ6L or SEZ6L2). In other selected embodiments the invention comprises a SEZ6 modulator that immunospecifically associates with one or more isoform of SEZ6 but does not immunospecifically associate with any other SEZ6 family member. In yet other embodiments the present invention comprises a method of treating a subject in need thereof comprising administering a therapeutically effective amount of a pan-SEZ6 modulator. Still other embodiments comprise a method of treating a subject in need thereof comprising administering a therapeutically effective amount of a SEZ6 modulator that immunospecifically associates with one or more isoforms of SEZ6 but does not immunospecifically associate with any other SEZ6 family member.

[0036] In one embodiment the invention is directed to a method of treating a subject suffering from cancer comprising administering a therapeutically effective amount of an antibody drug conjugate comprising an antibody conjugated directly or indirectly to a therapeutic moiety, wherein the antibody specifically binds to an epitope on a SEZ6 protein, wherein the epitope comprises amino acid residues selected from the group consisting of (i) residues R762, L764, Q777, I779, D781 and Q782; (ii) residues R342 and K389 and (iii) residues T352, S353 and H375. In preferred embodiments the therapeutic moiety will comprise auristatins, amanitins and pyrrolobenzodiazepines. In some cases the subject suffering from cancer may have previously been treated with a platinum based agent.

[0037] In another embodiment the invention is directed to a method of treating a subject suffering from medullary

thyroid cancer comprising administering a therapeutically effective amount of an antibody drug conjugate comprising an antibody conjugated directly or indirectly to a therapeutic moiety. In one embodiment the antibody drug conjugate used to treat a subject suffering from medullary thyroid cancer may comprise an antibody that specifically binds to an epitope on a SEZ6 protein, wherein the epitope comprises amino acid residues selected from the group consisting of (i) residues R762, L764, Q777, I779, D781 and Q782; (ii) residues R342 and K389 and (iii) residues T352, S353 and H375. In a preferred embodiment the therapeutic moiety may comprise auristatins, amanitins and pyrrolobenzodiazepines.

[0038] In a further embodiment the invention is directed to a method of treating a subject suffering from platinum resistant small cell lung cancer comprising administering a therapeutically effective amount of an antibody drug conjugate comprising an antibody conjugated directly or indirectly to a therapeutic moiety, wherein the antibody specifically binds to an epitope on a SEZ6 protein, wherein the epitope comprises amino acid residues selected from the group consisting of (i) residues R762, L764, Q777, I779, D781 and Q782; (ii) residues R342 and K389 and (iii) residues T352, S353 and H375. In preferred embodiments the patient suffering from platinum resistant small cell lung cancer have previously been treated with a platinum based agent. In another preferred embodiments the therapeutic moiety will comprise auristatins, amanitins and pyrrolobenzodiazepines.

[0039] Beyond the therapeutic uses discussed above it will also be appreciated that the modulators of the instant invention may be used to detect, diagnose or classify SEZ6 related disorders and, in particular, proliferative disorders. In some embodiments the modulator may be administered to the subject and detected or monitored *in vivo*. Those of skill in the art will appreciate that such modulators may be labeled or associated with markers or reporters as disclosed below and detected using any one of a number of standard techniques (e.g., MRI, CAT scan PET scan, etc.).

[0040] Thus, in some embodiments the invention will comprise a method of diagnosing, detecting or monitoring a SEZ6 associated disorder *in vivo* in a subject in need thereof comprising the step of administering a SEZ6 modulator.

[0041] In other instances the modulators may be used in an *in vitro* diagnostic setting using art-recognized procedures. As such, a preferred embodiment comprises a method of diagnosing cancer (e.g. medullary thyroid cancer or platinum resistant small cell lung cancer) in a subject comprising the steps of: (a) providing a tumor sample from a subject; (b) exposing the tumor sample to an anti-SEZ6 antibody labeled with a reporter wherein said anti-SEZ6 antibody associates with the tumor sample (e.g. a medullary thyroid tumor sample, a small cell lung cancer tumor sample or a platinum resistant small cell lung cancer tumor sample); and (c) detecting the reporter associated with the tumor sample. In some embodiments, the step of providing the tumor sample may be performed separately from the step of exposing the tumor sample to an anti-SEZ6 antibody or the step of detecting the reporter associated with the tumor sample. In some cases the reporter associated with the tumor sample will be detected *in vitro*, for example, using immunohistochemistry. In some cases the tumor sample will be exposed to an anti-SEZ6 antibody that binds to an epitope on a SEZ6 protein, wherein the epitope comprises amino acid residues R762, L764, Q777, I779, D781 and Q782.

[0042] Such methods may be easily discerned in conjunction with the instant application and may be readily performed using generally available commercial technology such as automatic plate readers, dedicated reporter systems, etc. In selected embodiments the SEZ6 modulator will be associated with tumor perpetuating cells present in the sample. In other preferred embodiments the detecting or quantifying step will comprise a reduction of tumor initiating cell frequency and detection thereof. Moreover, limiting dilution analysis may be conducted as previously alluded to above and will preferably employ the use of Poisson distribution statistics to provide an accurate accounting as to the reduction of frequency.

[0043] In a similar vein the present invention also provides kits or devices and associated methods that are useful in the diagnosis and monitoring of SEZ6 associated disorders such as cancer. To this end the present invention preferably provides an article of manufacture useful for diagnosing or treating SEZ6 associated disorders comprising a receptacle comprising a SEZ6 modulator and instructional materials for using said SEZ6 modulator to treat or diagnose the SEZ6 associated disorder. In selected embodiments the devices and associated methods will comprise the step of contacting at least one circulating tumor cell.

[0044] Other preferred embodiments of the invention also exploit the properties of the disclosed modulators as an instrument useful for identifying, characterizing, isolating, sectioning or enriching populations or subpopulations of tumor initiating cells through methods such as flow cytometric analysis including fluorescence activated cell sorting (FACS) or laser mediated sectioning.

[0045] As such, another preferred embodiment of the instant invention is directed to a method of identifying, isolating, sectioning or enriching a population of tumor initiating cells comprising the step of contacting said tumor initiating cells with a SEZ6 modulator.

[0046] The foregoing is a summary and thus contains, by necessity, simplifications, generalizations, and omissions of detail; consequently, those skilled in the art will appreciate that the summary is illustrative only and is not intended to be in any way limiting. Other aspects, features, and advantages of the methods, compositions and/or devices and/or other subject matter described herein will become apparent in the teachings set forth herein. The summary is provided to introduce a selection of concepts in a simplified form that are further described below in the Detailed Description. This

summary is not intended to identify key features or essential features of the claimed subject matter, nor is it intended to be used as an aid in determining the scope of the claimed subject matter.

BRIEF DESCRIPTION OF THE FIGURES

[0047]

FIGS. 1A - 1E are various representations of SEZ6 including nucleic acid or amino acid sequences pertaining to the SEZ6 modulators described herein. FIGS. 1A and 1B (SEQ ID NOS: 1 and 2) depict the full length mRNA sequence containing the open reading frames (ORFs) (underlined) encoding the SEZ6 variants 1 and 2, respectively. FIGS. 1C and 1D (SEQ ID NOS: 3 and 4) provide the corresponding amino acid sequences of the ORFs denoted in FIGS. 1A and 1B, respectively, with the amino acid residues indicating the predicted transmembrane spanning domain for each protein isoform underlined and the amino acid residues (residues 1-19) indicating the signal peptide in bold and underlined; FIG. 1E depicts the alignment of the two protein isoforms (SEQ ID NOS: 3 and 4) to illustrate the sequence differences in the cytoplasmic termini of each isoform, with the underlined residues indicating the differences between the two sequences; and FIG. 1F provides a schematic representation of the extracellular region of the SEZ6 protein illustrating the positions of the various domains.

FIGS. 2A - 2C provide a tabular representation of the percent identity at the protein level between the closest human isoform of SEZ6 and rhesus, cynomolgus, mouse or rat SEZ6 proteins (FIG. 2A); a tabular listing of various cDNA or protein sequence accessions for each of the reported isoforms of the SEZ6 family of genes (FIG. 2B); and the percent identity at the protein level between the longest isoforms of human SEZ6, SEZ6L, and SEZ6L2 proteins (FIG. 2C).

FIGS. 3A - 3C provide various representations of nucleic acid or amino acid sequences related to the production of the immunogens or cell lines used to generate or characterize the SEZ6 modulators described herein. For human SEZ6 a specific cDNA clone (FIG. 3A; SEQ ID NO: 5) encoding the complete mature human SEZ6 protein (FIG. 3B; SEQ ID NO: 6) was constructed from a commercial cDNA clone (BC146292; SEQ ID NO: 7) with known differences (FIG. 3C) from a database reference sequence, NP_849191 (SEQ ID NO: 3), for the SEZ6 protein.

FIGS. 4A and 4B provide a cDNA (FIG. 4A; SEQ ID NO: 8) used to express an Fc-SEZ6 construct in CHO-S cells and yield a protein immunogen (FIG. 4B; SEQ ID NO: 9), comprising the ECD of human SEZ6 fused to a human IgG2 Fc domain, in which the underlined sequences correspond to the human IgG2 Fc domain, the double underlined sequences correspond to the IgK signal peptide, and the amino acids in bold font correspond to residues contributed by the restriction sites used to clone the hSEZ6 fragment.

FIGS. 5A - 5J provide various representations of nucleic acid or amino acid sequences related to the production of the immunogens or cell lines used to generate or characterize the SEZ6 modulators described herein, wherein the underlined sequences denote the ECD of protein for the specific SEZ6 or SEZ6 family member being illustrated, and the figures comprise the cDNA sequences for the constructs encoding mature murine SEZ6 (FIG. 5A, SEQ ID NO: 10), mature rat SEZ6 (FIG. 5C, SEQ ID NO: 12), mature cynomolgus SEZ6 (FIG. 5E, SEQ ID NO: 14), mature ECD of the human SEZ6L protein (FIG. 5G, SEQ ID NO: 16), or the mature ECD of the human SEZ6L2 protein (FIG. 5I, SEQ ID NO: 18), or the corresponding proteins encoded by these cDNA constructs, namely mature murine SEZ6 (FIG. 5B, SEQ ID NO: 11), mature rat SEZ6 (FIG. 5D, SEQ ID NO: 13), mature cynomolgus SEZ6 (FIG. 5F, SEQ ID NO: 15), the mature ECD of the human SEZ6L protein (FIG. 5H, SEQ ID NO: 17), or the mature ECD of the human SEZ6L2 protein (FIG. 5J, SEQ ID NO: 19).

FIGS. 6A and 6B are depictions of mRNA expression levels of various genes as measured using whole transcriptome (SOLiD) sequencing of mRNA derived from tumor cell subpopulations or normal tissues. FIG. 6A is a tabular representation of genes associated with tumors having neuroendocrine features; and FIG. 6B is a graphical representation of SEZ6 mRNA expression in normal tissues and several non-traditional xenograft (NTX) tumors derived from lung cancers.

FIG. 7A - 7F depict mRNA expression levels analyzed using microarray. FIG. 7A is a graphical representation of unsupervised clustering of microarray profiles for 46 tumor lines and two normal tissues; FIGS. 7B and 7C are tabular representations of normalized intensity values corresponding to relative expression levels of selected genes related to neuroendocrine phenotypes (FIG. 7B) or the Notch signaling pathway (FIG. 7C) wherein unshaded cells and relatively low numbers indicate little to no expression and darker cells and relatively higher numbers indicate higher expression levels; FIG. 7D is a graphical representation showing relative expression levels of HES6 mRNA in various tumors and control tissues as measured using qRT-PCR; FIG. 7E is a tabular representation of normalized intensity values corresponding to relative expression levels of selected genes indicative of neurogenesis, neural commitment, or differentiation towards neural fates, with un-shaded cells indicating little to no expression and darker cells indicating higher expression levels; and FIG. 7F is a graphical representation of normalized intensity values corresponding to relative expression of SEZ6 in various NTX tumor lines.

FIGS. 8A and 8B are graphical representations showing relative expression levels of SEZ6 mRNA transcripts as measured by RT-PCR in a variety of RNA samples isolated from normal tissues or bulk neuroendocrine NTX tumors (FIG. 8A) and a variety of other NTX tumors (FIG. 8B).

FIGS. 9A and 9B are graphical representations showing the absolute (FIG. 9A) or normalized (FIG. 9B) mRNA expression levels of human SEZ6 as measured by RT-PCR in whole tumor specimens (grey dot) or matched normal adjacent tissue (NAT; white dot) from patients with one of eighteen different solid tumor types.

FIGS. 10A-10B provide the continuous amino acid sequences of light (FIG. 10A) and heavy chain (FIG. 10B) variable regions of a number of murine and humanized exemplary SEZ6 modulators isolated, cloned and engineered as described in the Examples herein. The corresponding nucleic acid sequences are set forth in the appended sequence listing. FIG. 10C sets out the full length amino acid sequences of the light and heavy chains of the humanized antibodies SC17.200 and SC17.200vL1.

FIG. 11 sets forth various characteristics of exemplary modulators of the invention. FIG. 11A shows the biochemical and immunological properties of exemplary SEZ6 modulators as represented in a tabular format; and Figure 11B provides a correlation between the domain to which an antibody binds and the antibody's efficacy in an *in vitro* killing assay.

FIGS. 12A and 12B show detection of expression of SEZ6 protein measured using an electrochemiluminescent assay. FIG. 12A shows SEZ6 expression in HEK-293T cells engineered to over-express human SEZ6 protein (h293T-HuSEZ6) using the anti-SEZ6 antibody SC17.33; FIG. 12B shows the relative protein expression of human SEZ6 in various NTX tumor and normal tissue lysates.

FIGS. 13A and 13B show detection by flow cytometry of SEZ6 protein expression on NTX tumor cells using various anti-SEZ6 antibodies (FIG. 13A); whereas FIG. 13B shows enhanced expression of SEZ6 protein in CSCs compared to NTG subpopulations using various anti-SEZ6 antibodies (FIG. 13B).

FIGS. 14A and 14B show that CSCs expressing SEZ6 exhibit enhanced tumorigenicity compared to CSCs that do not express SEZ6. FIG. 14A is a contour plot showing cell sorting by FACS of the cells in a lung tumor (LU37) on the basis of expression of CD324 (a marker of CSCs) and SEZ6; FIG. 14B is a graphical representation of the growth of tumor cells that are either CD324⁺SEZ6⁺ (black circles) or CD324⁺SEZ6⁻ (white circles) after implantation into immunocompromised mice. Tumor cells expressing both CD324 and SEZ6 exhibit enhanced tumorigenicity.

FIGS. 15A and 15B provide, respectively, a tabular and graphical representation illustrating that the disclosed modulators may effectively be used as targeting moieties to direct cytotoxic payloads to cells engineered to express SEZ6 (FIG. 15A) and NTX lung tumors (LU80, LU37 and LU100) grown *in vitro* (FIG. 15B) where the decrease in normalized relative luminescence units (RLU) is indicative of cell killing through internalization of the saporin toxin. FIG. 16 shows the quantification of SEZ6 expression in various SCLC and medullary thyroid tumors determined using immunohistochemistry (IHC), where FIG. 16A shows SEZ6 expression in NTX SCLC tumors, FIG. 16B shows SEZ6 expression in SCLC tumor microarrays and FIG. 16C shows SEZ6 expression in primary medullary thyroid tumors.

FIGS. 17A - 17B depict the ability of conjugated anti-SEZ6 mouse antibodies to retard *in vitro* and *in vivo* growth of NTX tumor cells. FIG. 17A shows the results of an *in vitro* killing assay using anti-SEZ6 ADCs on SCLC (LU64) and OV (OV26) NTX tumor cell lines; whereas FIG. 17B shows the effect of anti-SEZ6 ADCs on growth of SCLC (LU86) and LCNEC (LU50) tumors *in vivo*.

FIGS. 18A - 18D depict the ability of conjugated humanized anti-SEZ6 antibodies to retard growth of a thyroid cell line *in vitro* (FIG. 18A) and *in vivo* (FIG. 18B); and to retard growth of four SCLC tumors *in vivo* (LU80, LU64, LU111 and LU117) and achieve durable remission in immunodeficient mice (FIGS. 18C and 18D).

FIG. 19 shows dose response curves of oxalaplatin in the parental small cell lung cancer (SCLC) patient-derived xenograft (PDX) line, LU124p2 compared to the oxalaplatin-resistant cell line LU124OXAHlp2 and LU124OXAHlp3.

FIG. 20 shows mRNA expression of SEZ6 in parental SCLC PDX lines LU124p1 and LU124p3, and in oxalaplatin-resistant cell line LU124OXAHlp3.

FIG. 21 shows protein expression of SEZ6 in in parental SCLC PDX line LU124p3 and in oxalaplatin-resistant cell line LU124OXAHlp3.

DETAILED DESCRIPTION OF THE INVENTION

I. Introduction

[0048] While the present invention may be embodied in many different forms, disclosed herein are specific illustrative embodiments thereof that exemplify the principles of the invention. It should be emphasized that the present invention is not limited to the specific embodiments illustrated. Moreover, any section headings used herein are for organizational purposes only and are not to be construed as limiting the subject matter described. Finally, for the purposes of the instant disclosure all identifying sequence Accession numbers may be found in the NCBI Reference Sequence (RefSeq) data-

base and/or the NCBI GenBank® archival sequence database unless otherwise noted.

[0049] As previously alluded to, it has surprisingly been found that the expression of SEZ6 is associated with neoplastic growth and proliferative disorders, particularly in the instance of tumors with neuroendocrine features, and that SEZ6 and variants or isoforms thereof provide useful tumor markers which may be exploited in the treatment of related diseases. Moreover, as shown in the instant application it has unexpectedly been found that SEZ6 markers or determinants such as cell surface SEZ6 protein are associated with cancer stem cells (also known as tumor perpetuating cells) and may be effectively exploited to eliminate or silence the same. The ability to selectively reduce or eliminate cancer stem cells (e.g., through the use of conjugated SEZ6 modulators) is particularly surprising in that such cells are known to generally be resistant to many conventional treatments. That is, the effectiveness of traditional, as well as more recent targeted treatment methods, is often limited by the existence and/or emergence of resistant cancer stem cells that are capable of perpetuating the cancer even in the face of these diverse treatment methods. Further, determinants associated with cancer stem cells often make poor therapeutic targets due to low or inconsistent expression, failure to remain associated with the tumorigenic cell or failure to present at the cell surface. In sharp contrast to the teachings of the prior art, the instantly disclosed compounds and methods effectively overcome this inherent resistance to specifically eliminate, deplete, silence or promote the differentiation of such cancer stem cells thereby negating their ability to sustain or re-induce the underlying tumor growth.

[0050] More specifically, it has been discovered that SEZ6 modulators such as those disclosed herein may advantageously be used in the prognosis, diagnosis, theragnosis, treatment or prevention of proliferative disorders (e.g. neoplastic disorders) in subjects in need thereof. Accordingly, while preferred embodiments of the invention will be discussed extensively below, particularly in terms of particular domains, regions or epitopes or in the context of cancer stem cells or tumors comprising neuroendocrine features and their interactions with the disclosed modulators, those skilled in the art will appreciate that the scope of the instant invention is not limited by such exemplary embodiments. Rather, the most expansive embodiments of the present invention and the appended claims are broadly and expressly directed to SEZ6 modulators (including conjugated modulators) and their use in the prognosis, diagnosis, theragnosis, treatment or prevention of a variety of SEZ6 associated or mediated disorders, including neoplastic or proliferative disorders, regardless of any particular mechanism of action or specifically targeted tumor, cellular or molecular component.

[0051] To that end, and as demonstrated in the instant application, it has unexpectedly been found that the disclosed SEZ6 modulators can effectively be used to target and eliminate or otherwise incapacitate proliferative or tumorigenic cells and treat SEZ6 associated disorders (e.g., neoplasia). As used herein a "SEZ6 associated disorder" shall be held to mean any disorder or disease (including proliferative disorders) that is marked, diagnosed, detected or identified by a phenotypic or genotypic aberration of SEZ6 genetic components or expression during the course or etiology of the disease or disorder. In this regard a SEZ6 phenotypic aberration or determinant may, for example, comprise elevated or depressed levels of SEZ6 protein expression, abnormal SEZ6 protein expression on certain definable cell populations or abnormal SEZ6 protein expression at an inappropriate phase or stage of a cell lifecycle. Of course, it will be appreciated that similar expression patterns of genotypic determinants (e.g., mRNA transcription levels) of SEZ6 may also be used to classify or detect SEZ6 associated disorders.

[0052] As used herein the term "determinant" or "SEZ6 determinant" shall mean any detectable trait, property, marker or factor that is identifiably associated with, or specifically found in or on a particular cell, cell population or tissue including those identified in or on a tissue, cell or cell population affected by a SEZ6 associated disease or disorder. In selected preferred embodiments the SEZ6 modulators may associate, bind or react directly with the SEZ6 determinant (e.g., cell surface SEZ6 protein or SEZ6 mRNA) and thereby ameliorate the disorder. More generally determinants may be morphological, functional or biochemical in nature and may be genotypic or phenotypic. In other preferred embodiments the determinant is a cell surface antigen or genetic component that is differentially or preferentially expressed (or is not) by specific cell types (e.g., cancer stem cells) or by cells under certain conditions (e.g., during specific points of the cell cycle or cells in a particular niche). In still other preferred embodiments the determinant may comprise a gene or genetic entity that is differently regulated (up or down) in a specific cell or discrete cell population, a gene that is differentially modified with regard to its physical structure and chemical composition or a protein or collection of proteins physically associated with a gene that show differential chemical modifications. Determinants contemplated herein are specifically held to be positive or negative and may denote a cell, cell subpopulation or tissue (e.g., tumors) by its presence (positive) or absence (negative).

[0053] In a similar vein "SEZ6 modulators" of the invention broadly comprise any compound that recognizes, reacts, competes, antagonizes, interacts, binds, agonizes, or associates with a SEZ6 variant or isoform (or specific domains, regions or epitopes thereof) or its genetic component. By these interactions, the SEZ6 modulators may advantageously eliminate, reduce or moderate the frequency, activity, recurrence, metastasis or mobility of tumorigenic cells (e.g., tumor perpetuating cells or cancer stem cells). Exemplary modulators disclosed herein comprise nucleotides, oligonucleotides, polynucleotides, peptides or polypeptides. In certain preferred embodiments the selected modulators will comprise antibodies to a SEZ6 protein isoform or immunoreactive fragments or derivatives thereof. Such antibodies may be antagonistic or agonistic in nature and may optionally be conjugated or associated with a therapeutic or diagnostic agent.

Moreover, such antibodies or antibody fragments may comprise depleting, neutralizing or internalizing antibodies. In other embodiments, modulators within the instant invention will constitute a SEZ6 construct comprising a SEZ6 isoform or a reactive fragment thereof. It will be appreciated that such constructs may comprise fusion proteins and can include reactive domains from other polypeptides such as immunoglobulins or biological response modifiers. In still other aspects, the SEZ6 modulator will comprise a nucleic acid moiety (e.g. miRNA, siRNA, shRNA, antisense constructs, etc.) that exerts the desired effects at a genomic level. Still other modulators compatible with the instant teachings will be discussed in detail below.

[0054] More generally SEZ6 modulators of the present invention broadly comprise any compound that recognizes, reacts, competes, antagonizes, interacts, binds, agonizes, or associates with a SEZ6 determinant (genotypic or phenotypic) including cell surface SEZ6 protein. Whichever form of modulator is ultimately selected it will preferably be in an isolated and purified state prior to introduction into a subject. In this regard the term "isolated SEZ6 modulator" or "isolated SEZ6 antibody" shall be construed in a broad sense and in accordance with standard pharmaceutical practice to mean any preparation or composition comprising the modulator in a state substantially free of unwanted contaminants (biological or otherwise). Moreover these preparations may be purified and formulated as desired using various art recognized techniques. Of course, it will be appreciated that such "isolated" preparations may be intentionally formulated or combined with inert or active ingredients as desired to improve the commercial, manufacturing or therapeutic aspects of the finished product and provide pharmaceutical compositions. In a broader sense the same general considerations may be applied to an "isolated" SEZ6 isoform or variant or an "isolated" nucleic acid encoding the same.

[0055] Further, it has surprisingly been found that modulators interacting, associating or binding to particular SEZ6 domains, motifs or epitopes are especially effective in eliminating tumorigenic cells and/or silencing or attenuating cancer stem cell effects on tumor growth or propagation. That is, while modulators that react or associate with domains that are proximal to the cell surface (e.g. one of the Sushi or CUB-like domains) are effective in depleting or neutralizing tumorigenic cells it has unexpectedly been discovered that modulators associating or binding to domains, motifs or regions that are relatively more distal to the cell surface are also effective in eliminating, neutralizing, depleting or silencing tumorigenic cells. This is especially true of conjugated modulators such as, for example, anti-SEZ6 antibody drug conjugates comprising a cytotoxic agent.

[0056] While the present invention expressly contemplates the use of any SEZ6 modulator in the treatment of any SEZ6 disorder, including any type of neoplasia, in particularly preferred embodiments the disclosed modulators may be used to prevent, treat or diagnose tumors comprising neuroendocrine features (genotypic or phenotypic) including neuroendocrine tumors. True or "canonical neuroendocrine tumors" (NETs) arise from the dispersed endocrine system and are typically highly aggressive. Neuroendocrine tumors occur in the kidney, genitourinary tract (bladder, prostate, ovary, cervix, and endometrium), gastrointestinal tract (stomach, colon), thyroid (medullary thyroid cancer), and lung (small cell lung carcinoma and large cell neuroendocrine carcinoma). Moreover, the disclosed modulators may advantageously be used to treat, prevent or diagnose pseudo neuroendocrine tumors (pNETs) that genotypically or phenotypically mimic, comprise, resemble or exhibit common traits with canonical neuroendocrine tumors. "Pseudo neuroendocrine tumors" are tumors that arise from cells of the diffuse neuroendocrine system or from cells in which a neuroendocrine differentiation cascade has been aberrantly reactivated during the oncogenic process. Such pNETs commonly share certain genotypic, phenotypic or biochemical characteristics with traditionally defined neuroendocrine tumors, including the ability to produce subsets of biologically active amines, neurotransmitters, and peptide hormones. Accordingly, for the purposes of the instant invention the phrases "tumors comprising neuroendocrine features" or "tumors exhibiting neuroendocrine features" shall be held to comprise both neuroendocrine tumors and pseudo neuroendocrine tumors unless otherwise dictated by context.

[0057] In preferred embodiments the disclosed modulators will be used to treat small cell lung cancer and, in particularly preferred embodiments, will be used to treat platinum resistant small cell lung cancer in a subject in need thereof. As used herein and known in the art, the term "platinum resistant small cell lung cancer" means the presence of tumors or tumorigenic small cell lung cancer cells in a subject that are resistant or refractory to treatment with standard of care platinum based agents such as carboplatin, cisplatin and/or oxaloplatin. Such conditions are readily diagnosed using standard clinical procedures and their recognition is well within the purview of a clinician of ordinary skill in the art.

[0058] Besides the association with tumors generally discussed above, there are also indications of phenotypic or genotypic association between selected tumor initiating cells (TIC) and SEZ6 determinants. In this regard selected TICs (e.g., cancer stem cells) may express elevated levels of SEZ6 proteins when compared to normal tissue and non-tumorigenic cells (NTG), which together typically comprise much of a solid tumor. Thus, SEZ6 determinants may comprise a tumor associated marker (or antigen or immunogen) and the disclosed modulators may provide effective agents for the detection and suppression of TIC and associated neoplasia due to altered levels of the proteins on cell surfaces or in the tumor microenvironment. Accordingly, SEZ6 modulators, including immunoreactive antagonists and antibodies that associate, bind or react with the proteins, may effectively reduce the frequency of tumor initiating cells and could be useful in eliminating, depleting, incapacitating, reducing, promoting the differentiation of, or otherwise precluding or limiting the ability of these tumor-initiating cells to lie dormant and/or continue to fuel tumor growth, metastasis or

recurrence in a patient. In this regard those skilled in the art will appreciate that the present invention further provides SEZ6 modulators and their use in reducing the frequency of tumor initiating cells.

II. SEZ6 Physiology

[0059] SEZ6 (also known as seizure related 6 homolog) is a type I transmembrane protein originally cloned from mouse cerebrum cortex-derived cells treated with the convulsant pentylentetrazole (Shimizu-Nishikawa, 1995; PMID: 7723619). Representative SEZ6 protein orthologs include, but are not limited to, human (NP_849191; NP_001092105), chimpanzee (XP_511368, NP_001139913), mouse (NP_067261), and rat (NP_001099224). In humans, the SEZ6 gene consists of 17 exons spanning 51.1 kbp located on chromosome 17q11.2. Alternate splice acceptor sites only 16 base pairs apart within the last exon gives rise to two processed transcripts, one of approximately 4210 bases (NM_178860; FIG. 1A) and one of approximately 4194 bases (NM_001098635, FIG. 1B). The former transcript encodes a 994 amino acid protein (NP_849191; FIG. 1C), whereas the latter encodes a 993 amino acid protein (NP_001092105; FIG. 1D). These two protein isoforms of SEZ6 share overall 100% identity across their extracellular domains and their transmembrane domains, differing only in the final ten amino acid residues (FIG. 1E). A third splice variant has been reported to generate a secreted form of SEZ6 (Shimizu-Nishikawa, 1995; PMID: 7723619), however it has not been included in the RefSeqs associated within the NCBI database Gene page entry. The modulators of the invention may bind to any of the splice variants.

[0060] The biological relevance of the isoforms is unclear, although one study has suggested opposing actions for the membrane versus soluble proteins when their expression is restored in neurons from murine SEZ6 knockout mice (Gunnarsen *et al.* 2007, PMID: 18031681). Cross species protein sequence identity for the SEZ6 proteins are listed in FIG. 2A. In the human genome, there are two closely related genes- seizure related 6 homolog-like (SEZ6L) and seizure related 6 homolog like-2 (SEZ6L2), each of which has multiple splice variants encoding numerous isoforms (FIG. 2B). Percent identities for the longest protein of each of the members of this family of SEZ6-like proteins in humans are shown in FIG. 2C. Taken together SEZ6, SEZ6L and SEZ6L2, including their various isoforms, will be termed the SEZ6 family for the purposes of the instant application. SEZ6 modulators of the invention comprise modulators that are specific for each of SEZ6, SEZ6L or SEZ6L2. Alternatively, the modulators of the invention may cross react with SEZ6 and one or both of SEZ6L and/or SEZ6L2.

[0061] The mature SEZ6 protein is composed of a series of structural domains: a cytoplasmic domain, a transmembrane domain and an extracellular domain comprising a unique N-terminal domain, followed by two alternating Sushi and CUB-like domains, and three additional tandem Sushi domain repeats. Two isoforms of the SEZ6 antigen exist, and differ only on the extreme carboxy terminal, cytoplasmic domain.

[0062] FIG. 1F provides a schematic diagram of the extracellular region of the SEZ6 protein, illustrating the general juxtaposition of the Sushi and CUB domains, and the N-terminal domain. Generally, the domains are recognized as occurring at about amino acid residues 336-395 (Sushi Domain 1), 397-508 (CUB Domain 1), 511-572 (Sushi Domain 2), 574-685 (CUB Domain 2), 690-748 (Sushi Domain 3), 750-813 (Sushi Domain 4), 817-878 (Sushi Domain 5), with the N terminal domain at about amino acid residues 1-335, and a compositional bias of proline-rich residues at about amino acid residues 71-169.

[0063] The Sushi repeats are similar to the short consensus repeats found in the other human complement regulatory proteins (i.e., complement C3b/C4b binding sites). The CUB-like domains are similar to CUB domains found in other mammalian complement binding proteins which are associated with a wide range of proteins that participate in numerous biological processes other than complement activation, including but not limited to patterning, axon guidance, inflammation, and tumor suppression (Bork and Beckman, 1993, PMID: 8510165). Both the Sushi and CUB domains imply a function for SEZ6 involving binding of other proteins extracellularly. Proteins containing CUB domains also have been linked to cell signaling pathways, and consistent with this function, the SEZ6 C-terminal cytoplasmic domains contain the Asn-Pro-Thr-Tyr motif (SEQ ID NO: 200), which is a potential target for phosphorylation by Src tyrosine kinase family members. If true, this would link SEZ6 to a cellular signal transduction pathway leading to the activation of Ras, suggesting that SEZ6 may be a neurotrophic receptor.

[0064] Note that, the terms "mature protein" or "mature polypeptide" as used herein refers to the form(s) of the SEZ6 protein produced without the signal peptide of 19 amino acids that may be cleaved prior to cell surface expression. Unless otherwise indicated SEZ6 amino acid numbering (for domains, regions, epitopes, etc.) will be in the context of a mature protein without the leader.

[0065] SEZ6 is detectable by RT-PCR at low levels in kidney, liver, heart, lung and thymus of rodents, although strong protein expression was seen only in brain, with a significant level expressed in testis (Herbst and Nicklin, 1997, PMID: 9073173). Using polyclonal sera to SEZ6, protein expression was detected in day 13 of developing mouse forebrain. Strong staining was detected in the post-mitotic, maturing neurons of the developing cortical plate and sub-plate. This staining is diminished in the adult brain where the SEZ6 expression can be detected in other brain regions associated with ongoing morphological plasticity, such as the hippocampus, cerebellum, and olfactory bulb and in neurons of the

retina and spinal cord (Gunnarsen *et al.*, 2007, PMID: 18031681). The densest signals are found in regions with greatest concentration of neuronal cell bodies. In spite of widespread retinal expression of SEZ6, retinal function in the absence of SEZ6 was not affected (Gunnarsen *et al.*, 2009, PMID: 19662096). The SEZ6 staining pattern is closely tied with the emergence of the neocortical layers and hippocampus, and implies a forebrain-specific role for this gene during development. In human and mice SEZ6 was found to be differentially expressed in highly specific regions of the neocortex (Gunnarsen *et al.*, 2007, *supra*).

[0066] Mutations in the human SEZ6 gene have been linked to febrile seizures (FS), a convulsion associated with a rise in body temperature and the most common type of seizure in childhood (Yu *et al.*, 2007, PMID:17086543). FS may be classified as simple or complex, depending upon duration, recurrence, and extent of the body affected by the seizure. In a Chinese cohort, no mutations in SEZ6 were found in 15 healthy controls, but mutations were found in 21 of 60 patients with FS, with the most common type of mutation being a heterozygous, cytosine insertion (frame shift mutation) at position 1435 of the cDNA. The mutation incidence was significantly higher in patients with complex FS and in patients with a positive family history. As there is an 80% chance that children with complex FS will have seizures later in life, the authors suggest that screening for mutations in SEZ6 may be valuable in predicting FS recurrence or the development of epilepsy (Yu *et al.*, 2007, *supra*). Later studies have questioned the incidence, relevance, and ability of this study to have adequate power to imply causality, but do support that SEZ6 may be one gene among many that may play a role in seizure disorders (Mulley *et al.*, 2011, PMID: 21785725).

[0067] The specific molecular functions of SEZ6 remain unclear. As discussed above, analysis of the structural modules of the protein identified by homology and sequence analysis suggest a possible role in signaling, cell-cell communication, and neural development. The neuronal dendritic branching and connectivity that form the signaling networks that constitute the brain's circuitry arise and are specified both by intrinsic molecular programs in the neural cell as well as extrinsic signals. The process of dendritic growth in pyramidal neurons, the principal neuron in the mammalian forebrain, yields neurons with distinctive morphologies - a pyramidal cell body, and two distinct, complex dendritic trees: one emerging from the apex and the other from the base of the cell body. Gunnarsen *et al.* (2007, *supra*) have shown that SEZ6 null mice exhibit an excess of short dendrites in the dendritic trees of these neurons, yet display no increase in the overall dendritic field, the range of neurons with which a given neuron connects. Restoring the expression of the membrane bound SEZ6 isoforms in the knockout neurons results in an anti-branching effect. In behavioral tests the SEZ6 null mice display specific exploratory, motor, and cognitive deficits. These data suggest that SEZ6 is important for the achievement of the necessary balance between dendrite elongation and branching during the elaboration of a complex dendritic arbor during development.

[0068] Together, the studies above strongly suggest that the SEZ6 protein is important in the context of neural development, and is likely to have some role in cell-cell communication and signaling. Inappropriate reactivation of developmental signaling pathways or dysregulation of normal signaling pathways are commonly observed in tumors (Harris *et al.*, 2012). One collection of tumors sharing features indicative of partial reactivation of developmental programs are tumors with neuroendocrine phenotypes (Yao 2008; PMID: 18565894), in which various hormone and endocrine markers are expressed and/or secreted, and various neural markers indicative of neurogenesis, neural commitment, or differentiation towards neural fates are expressed. Tumors with neuroendocrine features arise infrequently in a wide range of primary sites, and while their exhaustive classification remains problematic (Yao; PMID: 18565894; Klimstra 2010; PMID: 20664470; Klöppel, 2011; PMID: 22005112), they may be classified into four major types: low grade benign carcinoids, low-grade well-differentiated neuroendocrine tumors with malignant behavior, tumors with mixed neuroendocrine and epithelial features, and high-grade poorly differentiated neuroendocrine carcinomas. Of these classifications, the poorly differentiated neuroendocrine carcinomas, which include small cell lung cancer (SCLC) and subsets of non-small cell lung cancer (NSCLC), are cancer types with dismal prognoses. It has been postulated that SCLC is bronchogenic in origin, arising in part from pulmonary neuroendocrine cells (Galluzzo and Bocchetta, 2011; PMID: 21504320). Whatever the cellular source of origin for these tumors, it is clear that they show a poorly differentiated endocrine phenotype, often are highly proliferative and aggressive, and frequently over-express neural proteins. Similarly, medullary thyroid cancers (MTC), a special neuroendocrine tumor type that arises from the calcitonin-secreting parafollicular C cells of the thyroid, show neuroendocrine phenotypes consistent with both their mature endocrine function and their derivation from neural crest tissue (Cook *et al.*, 2010; PMID: 20182588). While representing only about 3-5% of thyroid cancers, MTC results in up to 14% of all thyroid cancer deaths, is not very responsive to standard chemotherapy or radiation treatments, and even with newer molecularly targeted tyrosine kinase inhibitors such as cabozantinib and vandetanib, responds poorly to monotherapies (Haddad, 2013; PMID: 24002516). Given these examples of tumors with dismal prognoses, the resultant elevation of neural expression markers in these tumors that otherwise may be primarily restricted to the nervous system or show limited expression during development, of which SEZ6 may be an exemplar, may therefore offer a unique therapeutic target for tumors with the neuroendocrine phenotype.

III. Cancer Stem Cells

[0069] As alluded to above it has surprisingly been discovered that aberrant SEZ6 expression (genotypic and/or phenotypic) is associated with various tumorigenic cell subpopulations. In this respect the present invention provides SEZ6 modulators that may be particularly useful for targeting such cells, and especially tumor perpetuating cells, thereby facilitating the treatment, management or prevention of neoplastic disorders. Thus, in preferred embodiments modulators of SEZ6 determinants (phenotypic or genotypic) may be advantageously be used to reduce tumor initiating cell frequency in accordance with the present teachings and thereby facilitate the treatment or management of proliferative disorders.

[0070] For the purposes of the instant application the term "tumor initiating cell" (TIC) encompasses both "tumor perpetuating cells" (TPC; i.e., cancer stem cells or CSC) and highly proliferative "tumor progenitor cells" (termed TProg), which together generally comprise a unique subpopulation (i.e. 0.1-40%) of a bulk tumor or mass. For the purposes of the instant disclosure the terms "tumor perpetuating cells" and "cancer stem cells" or "neoplastic stem cells" are equivalent and may be used interchangeably herein. TPC differ from TProg in that TPC can completely recapitulate the composition of tumor cells existing within a tumor and have unlimited self-renewal capacity as demonstrated by serial transplantation (two or more passages through mice) of low numbers of isolated cells, whereas TProg will not display unlimited self-renewal capacity.

[0071] Those skilled in the art will appreciate that fluorescence-activated cell sorting (FACS) using appropriate cell surface markers is a reliable method to isolate highly enriched cancer stem cell subpopulations (e.g., > 99.5% purity) due, at least in part, to its ability to discriminate between single cells and clumps of cells (i.e. doublets, etc.). Using such techniques it has been shown that when low cell numbers of highly purified TProg cells are transplanted into immunocompromised mice they can fuel tumor growth in a primary transplant. However, unlike purified TPC subpopulations the TProg generated tumors do not completely reflect the parental tumor in phenotypic cell heterogeneity and are demonstrably inefficient at reinitiating serial tumorigenesis in subsequent transplants. In contrast, TPC subpopulations completely reconstitute the cellular heterogeneity of parental tumors and can efficiently initiate tumors when serially isolated and transplanted. Thus, those skilled in the art will recognize that a definitive difference between TPC and TProg, though both may be tumor generating in primary transplants, is the unique ability of TPC to perpetually fuel heterogeneous tumor growth upon serial transplantation at low cell numbers. Other common approaches to characterize TPC involve morphology and examination of cell surface markers, transcriptional profile, and drug response although marker expression may change with culture conditions and with cell line passage *in vitro*.

[0072] Accordingly, for the purposes of the instant invention tumor perpetuating cells, like normal stem cells that support cellular hierarchies in normal tissue, are preferably defined by their ability to self-renew indefinitely while maintaining the capacity for multilineage differentiation. Tumor perpetuating cells are thus capable of generating both tumorigenic progeny (i.e., tumor initiating cells: TPC and TProg) and non-tumorigenic (NTG) progeny. As used herein a "non-tumorigenic cell" (NTG) refers to a tumor cell that arises from tumor initiating cells, but does not itself have the capacity to self-renew or generate the heterogeneous lineages of tumor cells that comprise a tumor. Experimentally, NTG cells are incapable of reproducibly forming tumors in mice, even when transplanted in excess cell numbers.

[0073] As indicated, TProg are also categorized as tumor initiating cells (or TIC) due to their limited ability to generate tumors in mice. TProg are progeny of TPC and are typically capable of a finite number of non-self-renewing cell divisions. Moreover, TProg cells may further be divided into early tumor progenitor cells (ETP) and late tumor progenitor cells (LTP), each of which may be distinguished by phenotype (e.g., cell surface markers) and different capacities to recapitulate tumor cell architecture. In spite of such technical differences, both ETP and LTP differ functionally from TPC in that they are generally less capable of serially reconstituting tumors when transplanted at low cell numbers and typically do not reflect the heterogeneity of the parental tumor. Notwithstanding the foregoing distinctions, it has also been shown that various TProg populations can, on rare occasion, gain self-renewal capabilities normally attributed to stem cells and themselves become TPC (or CSC). In any event both types of tumor-initiating cells are likely represented in the typical tumor mass of a single patient and are subject to treatment with the modulators as disclosed herein. That is, the disclosed compositions are generally effective in reducing the frequency or altering the chemosensitivity of such SEZ6 positive tumor initiating cells regardless of the particular embodiment or mix represented in a tumor.

[0074] In the context of the instant invention, TPC are more tumorigenic, relatively more quiescent and often more chemoresistant than the TProg (both ETP and LTP), NTG cells and the tumor-infiltrating non-TPC derived cells (e.g., fibroblasts/stroma, endothelial & hematopoietic cells) that comprise the bulk of a tumor. Given that conventional therapies and regimens have, in large part, been designed to both debulk tumors and attack rapidly proliferating cells, TPC are likely to be more resistant to conventional therapies and regimens than the faster proliferating TProg and other bulk tumor cell populations. Further, TPC often express other characteristics that make them relatively chemoresistant to conventional therapies, such as increased expression of multi-drug resistance transporters, enhanced DNA repair mechanisms and anti-apoptotic proteins. These properties, each of which contribute to drug tolerance by TPC, constitute a key reason for the failure of standard oncology treatment regimens to ensure long-term benefit for most patients with advanced stage neoplasia; i.e. the failure to adequately target and eradicate those cells that fuel continued tumor growth

and recurrence (i.e. TPC or CSC).

[0075] Unlike many prior art treatments, the novel compositions of the present invention preferably reduce the frequency of tumor initiating cells upon administration to a subject regardless of the form or specific target (e.g., genetic material, SEZ6antibody or ligand fusion construct) of the selected modulator. As noted above, the reduction in tumor initiating cell frequency may occur as a result of a) elimination, depletion, sensitization, silencing or inhibition of tumor initiating cells; b) controlling the growth, expansion or recurrence of tumor initiating cells; c) interrupting the initiation, propagation, maintenance, or proliferation of tumor initiating cells; or d) by otherwise hindering the survival, regeneration and/or metastasis of the tumorigenic cells. In some embodiments, the reduction in the frequency of tumor initiating cells occurs as a result of a change in one or more physiological pathways. The change in the pathway, whether by reduction or elimination of the tumor initiating cells or by modifying their potential (e.g., induced differentiation, niche disruption) or otherwise interfering with their ability to influence the tumor environment or other cells, in turn allows for the more effective treatment of SEZ6 associated disorders by inhibiting tumorigenesis, tumor maintenance and/or metastasis and recurrence.

[0076] Among art-recognized methods that can be used to assess such a reduction in the frequency of tumor initiating cells is limiting dilution analysis either *in vitro* or *in vivo*, preferably followed by enumeration using Poisson distribution statistics or assessing the frequency of predefined definitive events such as the ability to generate tumors *in vivo* or not. While such limiting dilution analysis comprise preferred methods of calculating reduction of tumor initiating cell frequency other, less demanding methods, may also be used to effectively determine the desired values, albeit slightly less accurately, and are entirely compatible with the teachings herein. Thus, as will be appreciated by those skilled in the art, it is also possible to determine reduction of frequency values through well-known flow cytometric or immunohistochemical means. As to all the aforementioned methods see, for example, Dylla et al. 2008, PMID: 18560594 & Hoey et al. 2009, PMID: 19664991; each of which is incorporated herein by reference in its entirety and, in particular, for the disclosed methods.

[0077] With respect to limiting dilution analysis, *in vitro* enumeration of tumor initiating cell frequency may be accomplished by depositing either fractionated or unfractionated human tumor cells (e.g. from treated and untreated tumors, respectively) into *in vitro* growth conditions that foster colony formation. In this manner, colony forming cells might be enumerated by simple counting and characterization of colonies, or by analysis consisting of, for example, the deposition of human tumor cells into plates in serial dilutions and scoring each well as either positive or negative for colony formation at least 10 days after plating. *In vivo* limiting dilution experiments or analyses, which are generally more accurate in their ability to determine tumor initiating cell frequency encompass the transplantation of human tumor cells, from either untreated control or treated populations, for example, into immunocompromised mice in serial dilutions and subsequently scoring each mouse as either positive or negative for tumor formation at least 60 days after transplant. The derivation of cell frequency values by limiting dilution analysis *in vitro* or *in vivo* is preferably done by applying Poisson distribution statistics to the known frequency of positive and negative events, thereby providing a frequency for events fulfilling the definition of a positive event; in this case, colony or tumor formation, respectively.

[0078] As to other methods compatible with the instant invention that may be used to calculate tumor initiating cell frequency, the most common comprise quantifiable flow cytometric techniques and immunohistochemical staining procedures. Though not as precise as the limiting dilution analysis techniques described immediately above, these procedures are much less labor intensive and provide reasonable values in a relatively short time frame. Thus, it will be appreciated that a skilled artisan may use flow cytometric cell surface marker profile determination employing one or more antibodies or reagents that bind art-recognized cell surface proteins known to enrich for tumor initiating cells (e.g., potentially compatible markers as are set forth in PCT application 2012/031280 which is incorporated herein in its entirety) and thereby measure TIC levels from various samples. In still another compatible method one skilled in the art might enumerate TIC frequency *in situ* (e.g., in a tissue section) by immunohistochemistry using one or more antibodies or reagents that are able to bind cell surface proteins thought to demarcate these cells.

[0079] Those skilled in the art will recognize that numerous markers (or their absence) have been associated with various populations of cancer stem cells and used to isolate or characterize tumor cell subpopulations. In this respect exemplary cancer stem cell markers comprise OCT4, Nanog, STAT3, EPCAM, CD24, CD34, NB84, TrkA, GD2, CD133, CD20, CD56, CD29, B7H3, CD46, transferrin receptor, JAM3, carboxypeptidase M, ADAM9, oncostatin M, Lgr5, Lgr6, CD324, CD325, nestin, Sox1, Bmi-1, eed, easyh1, easyh2, mf2, yy1, smarcA3, smarcA5, smarcD3, smarcE1, mlt3, FZD1, FZD2, FZD3, FZD4, FZD6, FZD7, FZD8, FZD9, FZD10, WNT2, WNT2B, WNT3, WNT5A, WNT10B, WNT16, AXIN1, BCL9, MYC, (TCF4) SLC7A8, IL1RAP, TEM8, TMPRSS4, MUC16, GPRC5B, SLC6A14, SLC4A11, PPAP2C, CAV1, CAV2, PTPN3, EPHA1, EPHA2, SLC1A1, CX3CL1, ADORA2A, MPZL1, FLJ10052, C4.4A, EDG3, RARRES1, TMEPAI, PTS, CEACAM6, NID2, STEAP, ABCA3, CRIM1, IL1R1, OPN3, DAF, MUC1, MCP, CPD, NMA, ADAM9, GJA1, SLC19A2, ABCA1, PCDH7, ADCY9, SLC39A1, NPC1, ENPP1, N33, GPNMB, LY6E, CELSR1, LRP3, C20orf52, TMEPAI, FLVCR, PCDHA10, GPR54, TGFB3, SEMA4B, PCDHB2, ABCG2, CD166, AFP, BMP-4, β -catenin, CD2, CD3, CD9, CD14, CD31, CD38, CD44, CD45, CD74, CD90, CXCR4, decorin, EGFR, CD105, CD64, CD16, CD16a, CD16b, GLI1, GLI2, CD49b, and CD49f. See, for example, Schulenburg et al., 2010, PMID: 20185329, U.S.P.N.

7,632,678 and U.S.P.Ns. 2007/0292414, 2008/0175870, 2010/0275280, 2010/0162416 and 2011/0020221 each of which is incorporated herein by reference. It will further be appreciated that each of the aforementioned markers may also be used as a secondary target antigen in the context of the bispecific or multispecific antibodies of the instant invention.

[0080] Similarly, non-limiting examples of cell surface phenotypes associated with cancer stem cells of certain tumor types include CD44^{hi}CD24^{low}, ALDH⁺, CD133⁺, CD123⁺, CD34⁺CD38⁻, CD44⁺CD24⁻, CD46^{hi}CD324⁺CD66c⁻, CD133⁺CD34⁺CD10⁻CD19⁻, CD138⁻CD34⁺CD19⁺, CD133⁺RC2⁺, CD44⁺ $\alpha_2\beta_1$ ^{hi}CD133⁺, CD44⁺CD24⁺ESA⁺, CD271⁺, ABCB5⁺ as well as other cancer stem cell surface phenotypes that are known in the art. See, for example, Schulenburg et al., 2010, supra, Visvader et al., 2008, PMID: 18784658 and U.S.P.N. 2008/0138313, each of which is incorporated herein in its entirety by reference. Those skilled in the art will appreciate that marker phenotypes such as those exemplified immediately above may be used in conjunction with standard flow cytometric analysis and cell sorting techniques to characterize, isolate, purify or enrich TIC and/or TPC cells or cell populations for further analysis. Of interest with regard to the instant invention CD46, CD324 and, optionally, CD66c are either highly or heterogeneously expressed on the surface of many human colorectal ("CR"), breast ("BR"), non-small cell lung (NSCLC), small cell lung (SCLC), pancreatic ("PA"), melanoma ("Mel"), ovarian ("OV"), and head and neck cancer ("HN") tumor cells, regardless of whether the tumor specimens being analyzed were primary patient tumor specimens or patient-derived NTX tumors.

[0081] Using any of the above-referenced methods and selected markers as known in the art it is then possible to quantify the reduction in frequency of TIC (or the TPC therein) provided by the disclosed SEZ6 modulators (including those conjugated to cytotoxic agents) in accordance with the teachings herein. In some instances, the compounds of the instant invention may reduce the frequency of TIC or TPC (by a variety of mechanisms noted above, including elimination, induced differentiation, niche disruption, silencing, etc.) by 10%, 15%, 20%, 25%, 30% or even by 35%. In other embodiments, the reduction in frequency of TIC or TPC may be on the order of 40%, 45%, 50%, 55%, 60% or 65%. In certain embodiments, the disclosed compounds may reduce the frequency of TIC or TPC by 70%, 75%, 80%, 85%, 90% or even 95%. Of course it will be appreciated that any reduction of the frequency of the TIC or TPC likely results in a corresponding reduction in the tumorigenicity, persistence, recurrence and aggressiveness of the neoplasia.

IV. SEZ6 Modulators

[0082] In any event, the present invention is directed to the use of SEZ6 modulators, including SEZ6 antagonists, for the diagnosis, theragnosis, treatment and/or prophylaxis of various disorders including any one of a number of SEZ6 associated malignancies. The disclosed modulators may be used alone or in conjunction with a wide variety of anti-cancer compounds such as chemotherapeutic or immunotherapeutic agents (e.g., therapeutic antibodies) or biological response modifiers. In other selected embodiments, two or more discrete SEZ6 modulators may be used in combination to provide enhanced anti-neoplastic effects or may be used to fabricate multispecific constructs.

[0083] In certain embodiments, the SEZ6 modulators of the present invention will comprise nucleotides, oligonucleotides, polynucleotides, peptides or polypeptides. More particularly, exemplary modulators of the invention may comprise antibodies and antigen-binding fragments or derivatives thereof, proteins, peptides, glycoproteins, glycopeptides, glycolipids, polysaccharides, oligosaccharides, nucleic acids, antisense constructs, siRNA, miRNA, bioorganic molecules, peptidomimetics, pharmacological agents and their metabolites, transcriptional and translation control sequences, and the like. In certain embodiments the modulators will comprise soluble SEZ6 (sSEZ6) or a form, variant, derivative or fragment thereof including, for example, SEZ6 fusion constructs (e.g., SEZ6-Fc, SEZ6-targeting moiety, etc.) or SEZ6-conjugates (e.g., SEZ6-PEG, SEZ6-cytotoxic agent, SEZ6-brm, etc.). It will also be appreciated that, in other embodiments, the SEZ6 modulators comprise antibodies or immunoreactive fragments or derivatives thereof. In particularly preferred embodiments the modulators of the instant invention will comprise neutralizing antibodies or derivatives or fragments thereof. In other embodiments the SEZ6 modulators may comprise internalizing antibodies or fragments thereof. In still other embodiments the SEZ6 modulators may comprise depleting antibodies or fragments thereof. Moreover, as with the aforementioned fusion constructs, these antibody modulators may be conjugated, linked or otherwise associated with selected cytotoxic agents, polymers, biological response modifiers (BRMs) or the like to provide directed immunotherapies with various (and optionally multiple) mechanisms of action. As alluded to above such antibodies may be pan-SEZ6 antibodies and associate with two or more SEZ6 family members (e.g., SEZ6 and SEZ6L as shown in FIG. 11A) or immunospecific antibodies that selectively react with one or both isoforms of SEZ6. In yet other embodiments the modulators may operate on the genetic level and may comprise compounds as antisense constructs, siRNA, miRNA and the like that interact or associate with the genotypic component of a SEZ6 determinant.

[0084] It will further be appreciated that the disclosed SEZ6 modulators may deplete, silence, neutralize, eliminate or inhibit growth, propagation or survival of tumor cells, including TPC, and/or associated neoplasia through a variety of mechanisms, including agonizing or antagonizing selected pathways or eliminating specific cells depending, for example, on the form of SEZ6 modulator, any associated payload or dosing and method of delivery. Thus, while preferred embodiments disclosed herein are directed to the depletion, inhibition or silencing of specific tumor cell subpopulations such as tumor perpetuating cells or to modulators that interact with a specific epitope or domain, it must be emphasized

that such embodiments are merely illustrative and not limiting in any sense. Rather, as set forth in the appended claims, the present invention is broadly directed to SEZ6 modulators and their use in the treatment, management or prophylaxis of various SEZ6 associated disorders irrespective of any particular mechanism, binding region or target tumor cell population.

[0085] Regardless of the form of the modulator selected it will be appreciated that the chosen compound may be antagonistic in nature. As used herein an "antagonist" refers to a molecule capable of neutralizing, blocking, inhibiting, abrogating, reducing or interfering with the activities of a particular or specified target (e.g., SEZ6), including the binding of receptors to ligands or the interactions of enzymes with substrates. In this respect it will be appreciated that SEZ6 antagonists of the instant invention may comprise any ligand, polypeptide, peptide, fusion protein, antibody or immunologically active fragment or derivative thereof that recognizes, reacts, binds, combines, competes, associates or otherwise interacts with the SEZ6 protein or fragment thereof and eliminates, silences, reduces, inhibits, hinders, restrains or controls the growth of tumor initiating cells or other neoplastic cells including bulk tumor or NTG cell. Compatible antagonists may further include small molecule inhibitors, aptamers, antisense constructs, siRNA, miRNA and the like, receptor or ligand molecules and derivatives thereof which recognize or associate with a SEZ6 genotypic or phenotypic determinant thereby altering expression patterns or sequestering its binding or interaction with a substrate, receptor or ligand.

[0086] As used herein an antagonist refers to a molecule capable of neutralizing, blocking, inhibiting, abrogating, reducing or interfering with the activities of a particular or specified protein, including the binding of receptors to ligands or the interactions of enzymes with substrates. More generally antagonists of the invention may comprise antibodies and antigen-binding fragments or derivatives thereof, proteins, peptides, glycoproteins, glycopeptides, glycolipids, polysaccharides, oligosaccharides, nucleic acids, antisense constructs, siRNA, miRNA, bioorganic molecules, peptidomimetics, pharmacological agents and their metabolites, transcriptional and translation control sequences, and the like. Antagonists may also include small molecule inhibitors, fusion proteins, receptor molecules and derivatives which bind specifically to the protein thereby sequestering its binding to its substrate target, antagonist variants of the protein, antisense molecules directed to the protein, RNA aptamers, and ribozymes against the protein.

[0087] As used herein and applied to two or more molecules or compounds, the terms "recognizes" or "associates" shall be held to mean the reaction, binding, specific binding, combination, interaction, connection, linkage, uniting, coalescence, merger or joining, covalently or non-covalently, of the molecules whereby one molecule exerts an effect on the other molecule.

[0088] Moreover, as demonstrated in the examples herein (e.g., see FIG. 11), some modulators of human SEZ6 may, in certain cases, cross-react with SEZ6 from a species other than human (e.g., rat or cynomolgus monkey). In other cases exemplary modulators may be specific for one or more isoforms of human SEZ6 and will not exhibit cross-reactivity with SEZ6 orthologs from other species. Of course, in conjunction with the teachings herein such embodiments may comprise pan-SEZ6 antibodies that associate with two or more SEZ6 family members from a single species or antibodies that exclusively associate with SEZ6.

[0089] In any event, and as will be discussed in more detail below, those skilled in the art will appreciate that the disclosed modulators may be used in a conjugated or unconjugated form. That is, the modulator may be associated with or conjugated to (e.g. covalently or non-covalently) pharmaceutically active compounds, biological response modifiers, anti-cancer agents, cytotoxic or cytostatic agents, diagnostic moieties or biocompatible modifiers. In this respect it will be understood that such conjugates may comprise peptides, polypeptides, proteins, fusion proteins, nucleic acid molecules, small molecules, mimetic agents, synthetic drugs, inorganic molecules, organic molecules and radioisotopes. Moreover, as indicated herein the selected conjugate may be covalently or non-covalently linked to the SEZ6 modulator in various molar ratios depending, at least in part, on the method used to effect the conjugation.

V. Modulator Fabrication and Supply

A. Antibody Modulators

1. Overview

[0090] As previously alluded to particularly preferred embodiments of the instant invention comprise SEZ6 modulators in the form of antibodies that preferentially associate with one or more isoforms of SEZ6 (and, optionally, may cross-react with other SEZ6 family members). Those of ordinary skill in the art will appreciate the well developed knowledge base on antibodies such as set forth, for example, in Abbas et al., Cellular and Molecular Immunology, 6th ed., W.B. Saunders Company (2010) or Murphey et al., Janeway's Immunobiology, 8th ed., Garland Science (2011), each of which is incorporated herein by reference in its entirety.

[0091] The term "antibody" comprises polyclonal antibodies, multiclonal antibodies, monoclonal antibodies, chimeric antibodies, humanized and primatized antibodies, human antibodies, recombinantly produced antibodies, intrabodies, multispecific antibodies, bispecific antibodies, monovalent antibodies, multivalent antibodies, anti-idiotypic antibodies,

synthetic antibodies, including muteins and variants thereof; antibody fragments such as Fab fragments, F(ab') fragments, single-chain FvFcs, single-chain Fvs; and derivatives thereof including Fc fusions and other modifications, and any other immunologically active molecule so long as they exhibit the desired biological activity (i.e., antigen association or binding). Moreover, the term further comprises all classes of antibodies (i.e. IgA, IgD, IgE, IgG, and IgM) and all isotypes (i.e., IgG1, IgG2, IgG3, IgG4, IgA1, and IgA2), as well as variations thereof unless otherwise dictated by context. Heavy-chain constant domains that correspond to the different classes of antibodies are denoted by the corresponding lower case Greek letter α , δ , ϵ , γ , and μ , respectively. Light chains of the antibodies from any vertebrate species can be assigned to one of two clearly distinct types, called kappa (κ) and lambda (λ), based on the amino acid sequences of their constant domains.

[0092] While all such antibodies are within the scope of the present invention, preferred embodiments comprising the IgG class of immunoglobulin will be discussed in some detail herein solely for the purposes of illustration. It will be understood that such disclosure is, however, merely demonstrative of exemplary compositions and methods of practicing the present invention and not in any way limiting of the scope of the invention or the claims appended hereto.

[0093] As is well known, the variable domains of both the light (V_L) and heavy (V_H) chain portions determine antigen recognition and specificity and the constant domains of the light chain (C_L) and the heavy chain (C_H1 , C_H2 or C_H3) confer and regulate important biological properties such as secretion, transplacental mobility, circulation half-life, complement binding, and the like.

[0094] The "variable" region includes hypervariable sites that manifest themselves in three segments commonly termed complementarity determining regions (CDRs), in both the light-chain and the heavy-chain variable domains. The more highly conserved portions of variable domains flanking the CDRs are termed framework regions (FRs). For example, in naturally occurring monomeric immunoglobulin G (IgG) antibodies, the six CDRs present on each arm of the "Y" are short, non-contiguous sequences of amino acids that are specifically positioned to form the antigen binding site as the antibody assumes its three dimensional configuration in an aqueous environment. Thus, each naturally occurring IgG antibody comprises two identical binding sites proximal to the amino-terminus of each arm of the Y.

[0095] It will be appreciated that the position of CDRs can be readily identified by one of ordinary skill in the art using standard techniques. Also familiar to those in the art is the numbering system described in Kabat et al. (1991, NIH Publication 91-3242, National Technical Information Service, Springfield, Va.). In this regard Kabat et al. defined a numbering system for variable domain sequences that is applicable to any antibody. One of ordinary skill in the art can unambiguously assign this system of "Kabat numbering" to any variable domain sequence, without reliance on any experimental data beyond the sequence itself. Unless otherwise specified, references to the numbering of specific amino acid residue positions in an antibody are according to the Kabat numbering system.

[0096] Thus, according to Kabat, in the V_H , residues 31-35 comprise CDR1, residues 50-65 make up CDR2, and 95-102 comprise CDR3, while in the V_L , residues 24-34 are CDR1, 50-56 comprise CDR2, and 89-97 make up CDR3. For context, in a V_H , FR1 corresponds to the domain of the variable region encompassing amino acids 1-30; FR2 corresponds to the domain of the variable region encompassing amino acids 36-49; FR3 corresponds to the domain of the variable region encompassing amino acids 66-94, and FR4 corresponds to the domain of the variable region from amino acids 103 to the end of the variable region. The FRs for the light chain are similarly separated by each of the light chain variable region CDRs.

[0097] Note that CDRs vary considerably from antibody to antibody (and by definition will not exhibit homology with the Kabat consensus sequences). In addition, the identity of certain individual residues at any given Kabat site number may vary from antibody chain to antibody chain due to interspecies or allelic divergence. Alternative numbering is set forth in Chothia et al., J. Mol. Biol. 196:901-917 (1987) and MacCallum et al., J. Mol. Biol. 262:732-745 (1996), although as in Kabat, the FR boundaries are separated by the respective CDR termini as described above. See also Chothia et al., Nature 342, pp. 877-883 (1989) and S. Dubel, ed., Handbook of Therapeutic Antibodies, 3rd ed., WILEY-VCH Verlag GmbH and Co. (2007), where the definitions include overlapping or subsets of amino acid residues when compared against each other. Each of the aforementioned references is incorporated herein by reference in its entirety and the amino acid residues which comprise binding regions or CDRs as defined by each of the above cited references and are set forth for comparison in Table 1 below.

TABLE 1

CDR DEFINITIONS			
	Kabat ¹	Chothia ²	MacCallum ³
V_H CDR1	31-35	26-32	30-35
V_H CDR2	50-65	50-58	47-58
V_H CDR3	95-102	95-102	93-101

(continued)

CDR DEFINITIONS			
	Kabat ¹	Chothia ²	MacCallum ³
V _L CDR1	24-34	23-34	30-36
V _L CDR2	50-56	50-56	46-55
V _L CDR3	89-97	89-97	89-96
¹ Residue numbering follows the nomenclature of Kabat <i>et al.</i> , supra ² Residue numbering follows the nomenclature of Chothia <i>et al.</i> , supra ³ Residue numbering follows the nomenclature of MacCallum <i>et al.</i> , supra			

[0098] More practically variable regions and CDRs in an antibody sequence can be identified (i) according to general rules that have been developed in the art such as those discussed above or (ii) by aligning the sequences against a database of known variable regions. Methods for identifying these regions are described in Kontermann and Dubel, eds., *Antibody Engineering*, Springer, New York, NY, 2001, and Dinarello et al., *Current Protocols in Immunology*, John Wiley and Sons Inc., Hoboken, NJ, 2000. Exemplary databases of antibody sequences are described in, and can be accessed through, the "Abysis" website at www.bioinf.org.uk/abs (maintained by A.C. Martin in the Department of Biochemistry & Molecular Biology University College London, London, England) and VBASE2 website at www.vbase2.org, as described in Retter et al., *Nucl. Acids Res.*, 33 (Database issue): D671 - D674 (2005). In this regard the Abysis database website will automatically designate and annotate CDRs and framework regions (as per any of the commonly used numbering systems) upon entry of the subject heavy or light chain variable region nucleic acid or amino acid sequence. Moreover, the Abysis database website also includes general rules that have been developed for readily identifying CDRs which can be used in accordance with the teachings herein. In the context of the instant invention it will be appreciated that any of the disclosed light and heavy chain CDRs derived from the murine variable region amino acid sequences set forth in FIG. 10A or FIG. 10B may be combined or rearranged to provide optimized anti-SEZ6 (e.g. humanized, CDR grafted or chimeric anti-hSEZ6) antibodies in accordance with the instant teachings. That is, one or more of the CDRs derived from the light chain variable region amino acid sequences set forth in FIG. 10A (SEQ ID NOS: 20 - 168, even numbers) or the heavy chain variable region amino acid sequences set forth in FIG. 10B (SEQ ID NOS: 21 - 169, odd numbers) may be incorporated in a SEZ6 modulator and, in particularly preferred embodiments, in a CDR grafted or humanized antibody that immunospecifically associates with one or more SEZ6 isoforms. Examples of light (SEQ ID NOS: 170 - 192, even numbers) and heavy (SEQ ID NOS: 171 - 193, odd numbers and 194 - 199) chain variable region amino acid sequences of such humanized modulators are also set forth in FIGS. 10A and 10B.

[0099] Note that hSC17.200vL1 (SEQ ID NO: 192) is a variant of the humanized light chain construct hSC17.200 (SEQ ID NO: 190), hSC17.155vH1 - vH6 (SEQ ID NOS: 193-198) are variants of the heavy chain construct hSC.155 (SEQ ID NO: 184) which is derived from SC17.90 (SEQ ID NO: 127) and that hSC161vH1 (SEQ ID NO: 199) is a variant of the heavy chain construct hSC17.161 (SEQ ID NO: 189). As will be discussed in more detail below these variants were constructed and tested to optimize one or more biochemical properties of the parent antibody. The full length amino acid sequences of exemplary humanized antibodies, hSC17.200 and hSC17.200vL1 are set out in FIG. 10C as SEQ ID NOS: 400-402. The humanized antibody variant hSC17.200vL1 is derived from humanized antibody hSC17.200 and shares a common HC with the hSC17.200 antibody. Thus the full length LC and HC of hSC17.200 correspond to SEQ ID NOS: 400 and 401, respectively; and the full length LC and HC of hSC17.200vL1 correspond to SEQ ID NOS: 403 and 401, respectively.

[0100] Taken together these novel amino acid sequences depict seventy-five murine and eleven humanized exemplary modulators (along with reported variants) in accordance with the instant invention. Moreover, corresponding nucleic acid sequences of each of the seventy-five exemplary murine modulators and eleven humanized modulators and variants set forth in FIGS. 10A and 10B are included in the sequence listing of the instant application (SEQ ID NOS: 220 - 399).

[0101] In FIGS. 10A and 10B the annotated CDRs are defined using Kabat numbering. However, as discussed herein and demonstrated in Example 8 below, one skilled in the art could readily define, identify, derive and/or enumerate the CDRs as defined by Chothia et al., MacCallum et al. or one of the website databases such as Abysis or VBase2 for each respective heavy and light chain sequence set forth in FIG. 10A or FIG. 10B. Accordingly, each of the subject CDRs and antibodies comprising CDRs defined by all such nomenclature are expressly included within the scope of the instant invention. More broadly, the terms "variable region CDR amino acid residue" or more simply "CDR" includes amino acids in a CDR as identified using any sequence or structure based method as set forth above.

[0102] For any heavy chain constant region amino acid positions discussed in the present application, numbering is according to the Eu index first described in Edelman et al., 1969, *Proc. Natl. Acad. Sci. USA* 63(1): 78-85 describing the

amino acid sequence of myeloma protein Eu, which reportedly was the first human IgG1 sequenced. The Eu index of Edelman is also set forth in Kabat et al., 1991 (*supra.*). Thus, the terms "EU index as set forth in Kabat" or "EU index of Kabat" in the context of the heavy chain refers to the residue numbering system based on the human IgG1 Eu antibody of Edelman et al. as set forth in Kabat et al., 1991 (*supra.*). The numbering system used for the light chain constant region amino acid sequence is similarly set forth in Kabat 1991. Exemplary kappa C_L and IgG1 heavy chain constant region amino acid sequences compatible with the instant invention are set forth as SEQ ID NOS: 403 and 404 in the appended sequence listing. Those of skill in the art will appreciate that the disclosed constant region sequences may be joined with the disclosed heavy and light chain variable regions using standard molecular biology techniques to provide full-length antibodies that may be incorporated in the SEZ6 antibodies and ADCs of the instant invention.

[0103] As set forth below in the Examples, selected embodiments of the invention comprise murine antibodies that immunospecifically bind to SEZ6, which can be considered "source" antibodies or "reference" antibodies. In other embodiments, antibodies contemplated by the invention may be derived from such "source" or "reference" antibodies through optional modification of the constant region or the epitope-binding amino acid sequences of the source antibody. In one embodiment an antibody is "derived" from a source antibody if selected amino acids in the source antibody are altered through deletion, mutation, substitution, integration or combination. In another embodiment, a "derived" antibody is one in which fragments of the source antibody (e.g., one or more CDRs or the entire variable region) are combined with or incorporated into an acceptor antibody sequence to provide the derivative antibody (e.g. chimeric, CDR grafted or humanized antibodies). These "derived" (e.g. humanized or CDR-grafted) antibodies can be generated using standard molecular biology techniques for various reasons such as, for example, to improve affinity for the determinant; to improve production and yield in cell culture; to reduce immunogenicity *in vivo*; to reduce toxicity; to facilitate conjugation of an active moiety; or to create a multispecific antibody. Such antibodies may also be derived from source antibodies through modification of the mature molecule (e.g., glycosylation patterns or pegylation) by chemical means or post-translational modification. Examples of "source" murine antibodies of the invention are SC17.155, SC17.161 and SC17.200 and examples of antibodies that are derived from such source antibodies are hSC17.155, hSC17.155vH1-vH5 (derived from SC17.155); hSC17.161vL1 (derived from SC17.161) and hSC17.200vL1 (derived from hSC17.200).

2. Antibody Modulator Generation

a. Polyclonal antibodies

[0104] The production of polyclonal antibodies in various host animals, including rabbits, mice, rats, etc. is well known in the art. In some embodiments, polyclonal anti-SEZ6 antibody-containing serum is obtained by bleeding or sacrificing the animal. The serum may be used for research purposes in the form obtained from the animal or, in the alternative, the anti-SEZ6 antibodies may be partially or fully purified to provide immunoglobulin fractions or homogeneous antibody preparations.

[0105] Briefly the selected animal is immunized with a SEZ6 immunogen (e.g., soluble SEZ6 or sSEZ6) which may, for example, comprise selected isoforms, domains and/or peptides, or live cells or cell preparations expressing SEZ6 or immunoreactive fragments thereof. Art known adjuvants that may be used to increase the immunological response, depending on the inoculated species include, but are not limited to, Freund's (complete and incomplete), mineral gels such as aluminum hydroxide, surface active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, keyhole limpet hemocyanins, dinitrophenol, and potentially useful human adjuvants such as BCG (bacille Calmette-Guerin) and corynebacterium parvum. Such adjuvants may protect the antigen from rapid dispersal by sequestering it in a local deposit, or they may contain substances that stimulate the host to secrete factors that are chemotactic for macrophages and other components of the immune system. Preferably the immunization schedule will involve two or more administrations of the selected immunogen spread out over a predetermined period of time.

[0106] The amino acid sequence of a SEZ6 protein as shown in FIGS. 1C or 1D can be analyzed to select specific regions of the SEZ6 protein for generating antibodies. For example, hydrophobicity and hydrophilicity analyses of a SEZ6 amino acid sequence are used to identify hydrophilic regions in the SEZ6 structure. Regions of a SEZ6 protein that show immunogenic structure, as well as other regions and domains, can readily be identified using various other methods known in the art, such as Chou-Fasman, Garnier-Robson, Kyte-Doolittle, Eisenberg, Karplus-Schultz or Jameson-Wolf analysis. Average Flexibility profiles can be generated using the method of Bhaskaran R., Ponnuswamy P. K., 1988, Int. J. Pept. Protein Res. 32:242-255. Beta-turn profiles can be generated using the method of Deleage, G., Roux B., 1987, Protein Engineering 1:289-294. Thus, each SEZ6 region, domain or motif identified by any of these programs or methods is within the scope of the present invention and may be isolated or engineered to provide immunogens giving rise to modulators comprising desired properties. Preferred methods for the generation of SEZ6 antibodies are further illustrated by way of the Examples provided herein. Methods for preparing a protein or polypeptide for use as an immunogen are well known in the art. Also well known in the art are methods for preparing immunogenic conjugates of a protein with a carrier, such as BSA, KLH or other carrier protein. In some circumstances, direct conjugation using, for

example, carbodiimide reagents are used; in other instances linking reagents are effective. Administration of a SEZ6 immunogen is often conducted by injection over a suitable time period and with use of a suitable adjuvant, as is understood in the art. During the immunization schedule, titers of antibodies can be taken as described in the Examples below to determine adequacy of antibody formation.

b. Monoclonal antibodies

[0107] In addition, the invention contemplates use of monoclonal antibodies. As known in the art, the term "monoclonal antibody" (or mAb) refers to an antibody obtained from a population of substantially homogeneous antibodies, i.e., the individual antibodies comprising the population are identical except for possible mutations (e.g., naturally occurring mutations), that may be present in minor amounts. In certain embodiments, such a monoclonal antibody includes an antibody comprising a polypeptide sequence that binds or associates with an antigen wherein the antigen-binding polypeptide sequence was obtained by a process that includes the selection of a single target binding polypeptide sequence from a plurality of polypeptide sequences.

[0108] More generally, and as exemplified in Example 6 herein, monoclonal antibodies can be prepared using a wide variety of techniques known in the art including hybridoma, recombinant techniques, phage display technologies, transgenic animals (e.g., a XenoMouse®) or some combination thereof. For example, monoclonal antibodies can be produced using hybridoma and art-recognized biochemical and genetic engineering techniques such as described in more detail in An, Zhigiang (ed.) *Therapeutic Monoclonal Antibodies: From Bench to Clinic*, John Wiley and Sons, 1st ed. 2009; Shire et al. (eds.) *Current Trends in Monoclonal Antibody Development and Manufacturing*, Springer Science + Business Media LLC, 1st ed. 2010; Harlow et al., *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, 2nd ed. 1988; Hammerling, et al., in: *Monoclonal Antibodies and T-Cell Hybridomas* 563-681 (Elsevier, N.Y., 1981) each of which is incorporated herein in its entirety by reference. It should be understood that a selected binding sequence can be further altered, for example, to improve affinity for the target, to humanize the target binding sequence, to improve its production in cell culture, to reduce its immunogenicity in vivo, to create a multispecific antibody, etc., and that an antibody comprising the altered target binding sequence is also an antibody of this invention.

c. Chimeric antibodies

[0109] In another embodiment, the antibody of the invention may comprise chimeric antibodies derived from covalently joined protein segments from at least two different species or types of antibodies. As known in the art, the term "chimeric" antibodies is directed to constructs in which a portion of the heavy and/or light chain is identical with or homologous to corresponding sequences in antibodies derived from a particular species or belonging to a particular antibody class or subclass, while the remainder of the chain(s) is identical with or homologous to corresponding sequences in antibodies derived from another species or belonging to another antibody class or subclass, as well as fragments of such antibodies, so long as they exhibit the desired biological activity (U.S. P.N. 4,816,567; Morrison et al., *Proc. Natl. Acad. Sci. USA*, 81:6851-6855 (1984)).

[0110] In one embodiment, a chimeric antibody in accordance with the teachings herein may comprise murine V_H and V_L amino acid sequences and constant regions derived from human sources. In other compatible embodiments a chimeric antibody of the present invention may comprise a humanized antibody as described below. In another embodiment, the so-called "CDR-grafted" antibody, the antibody comprises one or more CDRs from a particular species or belonging to a particular antibody class or subclass, while the remainder of the antibody chain(s) is/are identical with or homologous to a corresponding sequence in antibodies derived from another species or belonging to another antibody class or subclass. For use in humans, selected rodent CDRs may be grafted into a human antibody, replacing one or more of the naturally occurring variable regions or CDRs of the human antibody. These constructs generally have the advantages of providing full strength modulator functions (e.g., CDC (complement dependent cytotoxicity), ADCC (antibody-dependent cell-mediated cytotoxicity), etc.) while reducing unwanted immune responses to the antibody by the subject.

d. Humanized antibodies

[0111] Similar to the CDR-grafted antibody is a "humanized" antibody. As used herein, "humanized" forms of non-human (e.g., murine) antibodies are chimeric antibodies that contain a minimal sequence derived from one or more non-human immunoglobulins. In one embodiment, a humanized antibody is a human immunoglobulin (recipient or acceptor antibody) in which residues from a CDR of the recipient are replaced by residues from a CDR of a non-human species (donor antibody) such as mouse, rat, rabbit, or nonhuman primate having the desired specificity, affinity, and/or capacity. In certain preferred embodiments, residues in one or more FRs in the variable domain of the human immunoglobulin are replaced by corresponding non-human residues from the donor antibody to help maintain the appropriate three-dimensional configuration of the grafted CDR(s) and thereby improve affinity. Furthermore, humanized antibodies may

comprise residues that are not found in the recipient antibody or in the donor antibody to, for example, further refine antibody performance.

[0112] CDR grafting and humanized antibodies are described, for example, in U.S.P.Ns. 6,180,370 and 5,693,762. The humanized antibody optionally may also comprise at least a portion of an immunoglobulin Fc, typically that of a human immunoglobulin. For further details, see, e.g., Jones et al., Nature 321:522-525 (1986); and U.S.P.Ns. 6,982,321 and 7,087,409. Still another method is termed "humaneering" which is described, for example, in U.S.P.N. 2005/0008625. Additionally, a non-human antibody may also be modified by specific deletion of human T-cell epitopes or "deimmunization" by the methods disclosed in WO 98/52976 and WO 00/34317. Each of the aforementioned references are incorporated herein in their entirety.

[0113] Humanized antibodies may also be bioengineered using common molecular biology techniques, such as isolating, manipulating, and expressing nucleic acid sequences that encode all or part of immunoglobulin variable regions from at least one of a heavy or light chain. In addition to the sources of such nucleic acid noted above, human germline sequences are available as disclosed, for example, in Tomlinson, I. A. et al. (1992) J. Mol. Biol. 227:776-798; Cook, G. P. et al. (1995) Immunol. Today 16: 237-242; Chothia, D. et al. (1992) J. Mol. Biol. 227:799-817; and Tomlinson et al. (1995) EMBO J 14:4628-4638. The V-BASE directory (VBASE2 - Retter et al., Nucleic Acid Res. 33: 671-674, 2005) provides a comprehensive directory of human immunoglobulin variable region sequences (compiled by Tomlinson, I. A. et al. MRC Centre for Protein Engineering, Cambridge, UK). Consensus human FRs can also be used, e.g., as described in U.S.P.N. 6,300,064.

[0114] In selected embodiments, and as detailed in Example 8 below, at least 60%, 65%, 70%, 75%, or 80% of the humanized antibody heavy or light chain variable region amino acid residues will correspond to those of the recipient FR and CDR sequences. In other embodiments at least 85% or 90% of the humanized antibody variable region residues will correspond to those of the recipient FR and CDR sequences. In a further preferred embodiment, greater than 95% of the humanized antibody variable region residues will correspond to those of the recipient FR and CDR sequences.

e. Human antibodies

[0115] In another embodiment, the antibodies may comprise fully human antibodies. The term "human antibody" refers to an antibody which possesses an amino acid sequence that corresponds to that of an antibody produced by a human and/or has been made using any of the techniques for making human antibodies.

[0116] Human antibodies can be produced using various techniques known in the art. One technique is phage display in which a library of (preferably human) antibodies is synthesized on phages, the library is screened with the antigen of interest or an antibody-binding portion thereof, and the phage that binds the antigen is isolated, from which one may obtain the immunoreactive fragments. Methods for preparing and screening such libraries are well known in the art and kits for generating phage display libraries are commercially available (e.g., the Pharmacia Recombinant Phage Antibody System, catalog no. 27-9400-01; and the Stratagene SurfZAP™ phage display kit, catalog no. 240612). There also are other methods and reagents that can be used in generating and screening antibody display libraries (see, e.g., U.S.P.N. 5,223,409; PCT Publication Nos. WO 92/18619, WO 91/17271, WO 92/20791, WO 92/15679, WO 93/01288, WO 92/01047, WO 92/09690; and Barbas et al., Proc. Natl. Acad. Sci. USA 88:7978-7982 (1991)).

[0117] In one embodiment, recombinant human antibodies may be isolated by screening a recombinant combinatorial antibody library prepared as above. In one embodiment, the library is a scFv phage display library, generated using human V_L and V_H cDNAs prepared from mRNA isolated from B-cells.

[0118] The antibodies produced by naive libraries (either natural or synthetic) can be of moderate affinity (K_a of about 10^6 to 10^7 m⁻¹), but affinity maturation can also be mimicked *in vitro* by constructing and reselecting from secondary libraries as described in the art. For example, mutation can be introduced at random *in vitro* by using error-prone polymerase (reported in Leung et al., Technique, 1: 11-15 (1989)). Additionally, affinity maturation can be performed by randomly mutating one or more CDRs, e.g. using PCR with primers carrying random sequence spanning the CDR of interest, in selected individual Fv clones and screening for higher-affinity clones. WO 9607754 described a method for inducing mutagenesis in a CDR of an immunoglobulin light chain to create a library of light chain genes. Another effective approach is to recombine the V_H or V_L domains selected by phage display with repertoires of naturally occurring V domain variants obtained from unimmunized donors and to screen for higher affinity in several rounds of chain reshuffling as described in Marks et al., Biotechnol., 10: 779-783 (1992). This technique allows the production of antibodies and antibody fragments with a dissociation constant K_D (k_{off}/k_{on}) of about 10^{-9} M or less.

[0119] In other embodiments, similar procedures may be employed using libraries comprising eukaryotic cells (e.g., yeast) that express binding pairs on their surface. See, for example, U.S.P.N. 7,700,302 and U.S.S.N. 12/404,059. In one embodiment, the human antibody is selected from a phage library, where that phage library expresses human antibodies (Vaughan et al. Nature Biotechnology 14:309-314 (1996); Sheets et al. Proc. Natl. Acad. Sci. USA 95:6157-6162 (1998). In other embodiments, human binding pairs may be isolated from combinatorial antibody libraries generated in eukaryotic cells such as yeast. See e.g., U.S.P.N. 7,700,302. Such techniques advantageously allow for

the screening of large numbers of candidate modulators and provide for relatively easy manipulation of candidate sequences (e.g., by affinity maturation or recombinant shuffling).

[0120] Human antibodies can also be made by introducing human immunoglobulin loci into transgenic animals, e.g., mice in which the endogenous immunoglobulin genes have been partially or completely inactivated and human immunoglobulin genes have been introduced. Upon challenge, human antibody production is observed, which closely resembles that seen in humans in all respects, including gene rearrangement, assembly, and antibody repertoire. This approach is described, for example, in U.S.P.Ns. 5,545,807; 5,545,806; 5,569,825; 5,625,126; 5,633,425; 5,661,016, and U.S.P.Ns. 6,075,181 and 6,150,584 regarding XenoMouse® technology; and Lonberg and Huszar, Intern. Rev. Immunol. 13:65-93 (1995). Alternatively, the human antibody may be prepared via immortalization of human B lymphocytes producing an antibody directed against a target antigen (such B lymphocytes may be recovered from an individual suffering from a neoplastic disorder or may have been immunized *in vitro*). See, e.g., Cole et al., Monoclonal Antibodies and Cancer Therapy, Alan R. Liss, p. 77 (1985); Boerner et al., J. Immunol, 147 (1):86-95 (1991); and U.S.P.N. 5,750,373.

3. Further Processing

[0121] No matter how obtained, modulator-producing cells (e.g., hybridomas, yeast colonies, etc.) may be selected, cloned and further screened for desirable characteristics including, for example, robust growth, high antibody production and, as discussed in more detail below, desirable antibody characteristics. Hybridomas can be expanded *in vivo* in syngeneic animals, in animals that lack an immune system, e.g., nude mice, or in cell culture *in vitro*. Methods of selecting, cloning and expanding hybridomas and/or colonies, each of which produces a discrete antibody species, are well known to those of ordinary skill in the art.

B. Recombinant Modulator Production

1. Overview

[0122] Once the source is perfected DNA encoding the desired SEZ6 modulators may be readily isolated and sequenced using conventional procedures (e.g., by using oligonucleotide probes that are capable of binding specifically to genes encoding antibody heavy and light chains). Isolated and subcloned hybridoma cells (or phage or yeast derived colonies) may serve as a preferred source of such DNA if the modulator is an antibody. If desired, the nucleic acid can further be manipulated as described herein to create agents including fusion proteins, or chimeric, humanized or fully human antibodies. More particularly, isolated DNA (which may be modified) can be used to clone constant and variable region sequences for the manufacture antibodies.

[0123] Accordingly, in exemplary embodiments antibodies may be produced recombinantly, using conventional procedures (such as those set forth in Al-Rubeai; An, and Shire et. al. all supra, and Sambrook J. & Russell D. Molecular Cloning: A Laboratory Manual, 3rd ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (2000); Ausubel et al., Short Protocols in Molecular Biology: A Compendium of Methods from Current Protocols in Molecular Biology, Wiley, John & Sons, Inc. (2002)) in which the isolated and subcloned hybridoma cells (or phage or yeast derived colonies) serve as a preferred source of nucleic acid molecules.

[0124] The term "nucleic acid molecule", as used herein, is intended to include DNA molecules and RNA molecules and artificial variants thereof (e.g., peptide nucleic acids), whether single-stranded or double-stranded. The nucleic acids may encode one or both chains of an antibody of the invention, or a fragment or derivative thereof. The nucleic acid molecules of the invention also include polynucleotides sufficient for use as hybridization probes, PCR primers or sequencing primers for identifying, analyzing, mutating or amplifying a polynucleotide encoding a polypeptide; anti-sense nucleic acids for inhibiting expression of a polynucleotide, and as well as complementary sequences. The nucleic acids can be any length. They can be, for example, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 75, 100, 125, 150, 175, 200, 250, 300, 350, 400, 450, 500, 750, 1,000, 1,500, 3,000, 5,000 or more nucleotides in length, and/or can comprise one or more additional sequences, for example, regulatory sequences, and/or be part of a larger nucleic acid, for example, a vector. It will be appreciated that such nucleic acid sequences can further be manipulated to create modulators including chimeric, humanized or fully human antibodies. More particularly, isolated nucleic acid molecules (which may be modified) can be used to clone constant and variable region sequences for the manufacture antibodies as described in U.S.P.N. 7,709,611.

[0125] The term "isolated nucleic acid" means a that the nucleic acid was (i) amplified *in vitro*, for example by polymerase chain reaction (PCR), (ii) recombinantly produced by cloning, (iii) purified, for example by cleavage and gel-electrophoretic fractionation, or (iv) synthesized, for example by chemical synthesis. An isolated nucleic acid is a nucleic acid that is available for manipulation by recombinant DNA techniques.

[0126] Whether the source of the nucleic acid encoding the desired immunoreactive portion of the antibody is obtained or derived from phage display technology, yeast libraries, hybridoma-based technology or synthetically, it is to be un-

derstood that the present invention encompasses the nucleic acid molecules and sequences encoding the antibodies or antigen-binding fragments or derivatives thereof. Further, the instant invention is directed to vectors and host cells comprising such nucleic acid molecules.

[0127] Once DNA fragments encoding V_H and V_L segments are obtained, these DNA fragments can be further manipulated by standard recombinant DNA techniques, for example to convert the variable region genes to full-length antibody chain genes, to Fab fragment genes or to a scFv gene. In these manipulations, a V_L - or V_H -encoding DNA fragment is operatively linked to another DNA fragment encoding another protein, such as an antibody constant region or a flexible linker. The term "operatively linked", as used in this context, means that the two DNA fragments are joined such that the amino acid sequences encoded by the two DNA fragments remain in-frame.

[0128] The isolated DNA encoding the V_H region can be converted to a full-length heavy chain gene by operatively linking the V_H -encoding DNA to another DNA molecule encoding heavy chain constant regions (C_H1 , C_H2 and C_H3). The sequences of human heavy chain constant region genes are known in the art (see e.g., Kabat, E. A., et al. (1991) Sequences of Proteins of Immunological Interest, Fifth Edition, U.S. Department of Health and Human Services, NIH Publication No. 91-3242) and DNA fragments encompassing these regions can be obtained by standard PCR amplification. The heavy chain constant region can be an IgG1, IgG2, IgG3, IgG4, IgA, IgE, IgM or IgD constant region, but most preferably is an IgG1 or IgG4 constant region. As discussed in more detail below an exemplary IgG1 constant region that is compatible with the teachings herein is set forth as SEQ ID NO: 403 in the appended sequence listing. For a Fab fragment heavy chain gene, the V_H -encoding DNA can be operatively linked to another DNA molecule encoding only the heavy chain $CH1$ constant region.

[0129] The isolated DNA encoding the V_L region can be converted to a full-length light chain gene (as well as a Fab light chain gene) by operatively linking the V_L -encoding DNA to another DNA molecule encoding the light chain constant region, C_L . The sequences of human light chain constant region genes are known in the art (see e.g., Kabat, E. A., et al. (1991) Sequences of Proteins of Immunological Interest, Fifth Edition, U.S. Department of Health and Human Services, NIH Publication No. 91-3242) and DNA fragments encompassing these regions can be obtained by standard PCR amplification. The light chain constant region can be a kappa or lambda constant region, but most preferably is a kappa constant region. In this respect an exemplary compatible kappa light chain constant region is set forth as SEQ ID NO: 404 in the appended sequence listing.

2. Hybridization and Sequence Identity

[0130] As indicated, the invention further provides nucleic acids that hybridize to other nucleic acids under particular hybridization conditions. More specifically the invention encompasses nucleic acids molecules that hybridize under moderate or high stringency hybridization conditions (e.g., as defined below), to the nucleic acid molecules of the invention. Methods for hybridizing nucleic acids are well-known in the art. As is well known, a moderately stringent hybridization conditions comprise a prewashing solution containing 5x sodium chloride/sodium citrate (SSC), 0.5% SDS, 1.0 mM EDTA (pH 8.0), hybridization buffer of about 50% formamide, 6xSSC, and a hybridization temperature of 55°C (or other similar hybridization solutions, such as one containing about 50% formamide, with a hybridization temperature of 42°C), and washing conditions of 60°C, in 0.5xSSC, 0.1% SDS. By way of comparison hybridization under highly stringent hybridization conditions comprise washing with 6xSSC at 45°C, followed by one or more washes in 0.1xSSC, 0.2% SDS at 68°C. Furthermore, one of skill in the art can manipulate the hybridization and/or washing conditions to increase or decrease the stringency of hybridization such that nucleic acids comprising nucleotide sequences that are at least 65%, 70%, 75%, 80%, 85%, 90%, 95%, 98% or 99% identical to each other typically remain hybridized to each other.

[0131] The invention also includes nucleic acid molecules that are "substantially identical" to the described nucleic acid molecules. In one embodiment, the term substantially identical with regard to a nucleic acid sequence means may be construed as a sequence of nucleic acid molecules exhibiting at least about 65%, 70%, 75%, 80%, 85%, or 90% sequence identity. In other embodiments, the nucleic acid molecules exhibit 95% or 98% sequence identity to the reference nucleic acid sequence.

[0132] The basic parameters affecting the choice of hybridization conditions and guidance for devising suitable conditions are set forth by, for example, Sambrook, Fritsch, and Maniatis (1989, Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., chapters 9 and 11; and Current Protocols in Molecular Biology, 1995, Ausubel et al., eds., John Wiley & Sons, Inc., sections 2.10 and 6.3-6.4), and can be readily determined by those having ordinary skill in the art based on, for example, the length and/or base composition of the nucleic acid.

[0133] Sequence similarity for polypeptides, which is also referred to as sequence identity, is typically measured using sequence analysis software. Protein analysis software matches similar sequences using measures of similarity assigned to various substitutions, deletions and other modifications, including conservative amino acid substitutions. For instance, the sequence analysis tool GCG (Accelrys Software Inc.) contains programs such as "GAP" and "BEST-FIT" which can be used with default parameters to determine sequence homology or sequence identity between closely related polypep-

tides, such as homologous polypeptides from different species of organisms or between a wild type protein and a mutein thereof. (See, e.g., GCG Version 6.1 or Durbin et. Al., *Biological Sequence Analysis: Probabilistic models of proteins and nucleic acids.*, Cambridge Press (1998)).

[0134] Polypeptide sequences can also be compared using FASTA using default or recommended parameters, a program in GCG Version 6.1. FASTA (e.g., FASTA2 and FASTA3) provides alignments and percent sequence identity of the regions of the best overlap between the query and search sequences (Pearson (2000) *supra*). Another preferred algorithm when comparing a sequence of the invention to a database containing a large number of sequences from different organisms is the computer program BLAST, especially blastp or tblastn, using default parameters. See, e.g., Altschul et al. (1990) *J. Mol. Biol.* 215: 403 410 and Altschul et al. (1997) *Nucleic Acids Res.* 25:3389 402, each of which is herein incorporated by reference.

[0135] In this regard the invention also includes nucleic acid molecules that encode polypeptides that are "substantially identical" with respect to an antibody variable region polypeptide sequence (e.g., either the donor light or heavy chain variable region, acceptor light or heavy chain variable region or resulting humanized construct). As applied to such polypeptides, the term "substantial identity" or "substantially identical" means that two peptide sequences, when optimally aligned, such as by the programs GAP or BEST-FIT using default gap weights, share at least 60% or 65% sequence identity, preferably at least 70%, 75%, 80%, 85%, or 90% sequence identity, even more preferably at least 93%, 95%, 98% or 99% sequence identity. Preferably, residue positions which are not identical differ by conservative amino acid substitutions. A "conservative amino acid substitution" is one in which an amino acid residue is substituted by another amino acid residue having a side chain (R group) with similar chemical properties (e.g., charge or hydrophobicity). In general, a conservative amino acid substitution will not substantially change the functional properties of a protein. In cases where two or more amino acid sequences differ from each other by conservative substitutions, the percent sequence identity or degree of similarity may be adjusted upwards to correct for the conservative nature of the substitution.

3. Expression

[0136] The varied processes of recombinant expression, i.e., the production of RNA or of RNA and protein/peptide, are well known as set forth, for example, in Berger and Kimmel, *Guide to Molecular Cloning Techniques, Methods in Enzymology* volume 152 Academic Press, Inc., San Diego, Calif.; Sambrook et al., *Molecular Cloning-A Laboratory Manual* (3rd Ed.), Vol. 1-3, Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y., (2000); and *Current Protocols in Molecular Biology*, F. M. Ausubel et al., eds., Current Protocols, a joint venture between Greene Publishing Associates, Inc. and John Wiley & Sons, Inc., (supplemented through 2006).

[0137] Certain terms of interest include "expression control sequence" which comprises promoters, ribosome binding sites, enhancers and other control elements which regulate transcription of a gene or translation of mRNA. As is well known, a "promoter" or "promoter region" relates to a nucleic acid sequence which generally is located upstream (5') to the nucleic acid sequence being expressed and controls expression of the sequence by providing a recognition and binding site for RNA-polymerase.

[0138] Exemplary promoters which are compatible according to the invention include promoters for SP6, T3 and T7 polymerase, human U6 RNA promoter, CMV promoter, and artificial hybrid promoters thereof (e.g. CMV) where a part or parts are fused to a part or parts of promoters of genes of other cellular proteins such as e.g. human GAPDH (glyceraldehyde-3-phosphate dehydrogenase), and including or not including (an) additional intron(s).

[0139] In certain embodiments, the nucleic acid molecule may be present in a vector, where appropriate with a promoter, which controls expression of the nucleic acid. The well known term "vector" comprises any intermediary vehicle for a nucleic acid which enables said nucleic acid, for example, to be introduced into prokaryotic and/or eukaryotic cells and, where appropriate, to be integrated into a genome. Methods of transforming mammalian cells are well known in the art. See, for example, U.S.P.Ns. 4,399,216, 4,912,040, 4,740,461, and 4,959,455. The vectors may include a nucleotide sequence encoding an antibody of the invention (e.g., a whole antibody, a heavy or light chain of an antibody, a V_H or V_L of an antibody, or a portion thereof, or a heavy- or light-chain CDR, a single chain Fv, or fragments or variants thereof), operably linked to a promoter (see, e.g., PCT Publication WO 86/05807; PCT Publication WO 89/01036; and U.S.P.N. 5,122,464).

[0140] A variety of host-expression vector systems are commercially available, and many are compatible with the teachings herein and may be used to express the modulators of the invention. Such systems include, but are not limited to, microorganisms such as bacteria (e.g., *E. coli*, *B. subtilis*, streptomycetes) transformed with recombinant bacteriophage DNA, plasmid DNA or cosmid DNA expression vectors containing modulator coding sequences; yeast (e.g., *Saccharomyces*, *Pichia*) transfected with recombinant yeast expression vectors containing modulator coding sequences; insect cell systems infected with recombinant virus expression vectors (e.g., baculovirus) containing modulator coding sequences; plant cell systems (e.g., *Nicotiana*, *Arabidopsis*, duckweed, corn, wheat, potato, etc.) infected with recombinant viral expression vectors (e.g., cauliflower mosaic virus; tobacco mosaic virus) or transfected with recombinant plasmid expression vectors (e.g., Ti plasmid) containing modulator coding sequences; or mammalian cell systems (e.g., COS,

CHO, BHK, 293, 3T3 cells, etc.) harboring recombinant expression constructs containing promoters derived from the genome of mammalian cells (e.g., metallothionein promoter) or from mammalian viruses (e.g., the adenovirus late promoter; the vaccinia virus 7.5K promoter).

[0141] As used herein, the term "host cell" covers any kind of cellular system which can be engineered to generate the polypeptides and antigen-binding molecules of the present invention. In one embodiment, the host cell is engineered to allow the production of an antigen binding molecule with modified glycoforms. In a preferred embodiment, the antigen binding molecule, or variant antigen binding molecule, is an antibody, antibody fragment, or fusion protein. In certain embodiments, the host cells have been further manipulated to express increased levels of one or more polypeptides having N-acetylglucosaminyltransferase III (GnT11) activity. Compatible host cells include cultured cells, e.g., mammalian cultured cells, such as CHO cells, BHK cells, NSO cells, SP2/0 cells, YO myeloma cells, P3X63 mouse myeloma cells, PER cells, PER.C6 cells or hybridoma cells, yeast cells, insect cells, and plant cells, to name only a few, but also cells comprised within a transgenic animal, transgenic plant or cultured plant or animal tissue.

[0142] For long-term, high-yield production of recombinant proteins stable expression is preferred. Accordingly, cell lines that stably express the selected modulator may be engineered using standard art-recognized techniques. Rather than using expression vectors that contain viral origins of replication, host cells can be transformed with DNA controlled by appropriate expression control elements (e.g., promoter, enhancer, sequences, transcription terminators, polyadenylation sites, etc.), and a selectable marker. Any of the selection systems well known in the art may be used, including the glutamine synthetase gene expression system (the GS system) which provides an efficient approach for enhancing expression under certain conditions. The GS system is discussed in whole or part in connection with EP patents 0 216 846, 0 256 055, 0 323 997 and 0 338 841 and U.S.P.N.s 5,591,639 and 5,879,936 each of which is incorporated herein by reference. Another preferred expression system, the Freedom™ CHO-S Kit is commercially provided by Life Technologies (Catalog Number A13696-01) also allows for the development of stable cell lines that may be used for modulator production.

[0143] Such host-expression systems represent vehicles by which the coding sequences of interest may be produced and subsequently purified, but also represent cells which may, when transformed or transfected with the appropriate nucleotide coding sequences, express a molecule of the invention in situ. The host cell may be co-transfected with two expression vectors of the invention, for example, the first vector encoding a heavy chain derived polypeptide and the second vector encoding a light chain derived polypeptide.

[0144] Thus, in certain embodiments, the present invention provides recombinant host cells allowing for the expression of antibodies or portions thereof. Antibodies produced by expression in such recombinant host cells are referred to herein as recombinant antibodies. The present invention also provides progeny cells of such host cells, and antibodies produced by the same.

C. Chemical Synthesis

[0145] In addition, the modulators may be chemically synthesized using techniques known in the art (e.g., see Creighton, 1983, *Proteins: Structures and Molecular Principles*, W.H. Freeman & Co., N.Y., and Hunkapiller, M., et al., 1984, *Nature* 310:105-111). Furthermore, if desired, nonclassical amino acids or chemical amino acid analogs (such as D-isomers of the common amino acids, 2,4-diaminobutyric acid, α-amino isobutyric acid, 4-aminobutyric acid, and the like) can be introduced as a substitution or addition into a polypeptide sequence.

D. Transgenic Systems

[0146] In other embodiments modulators may be produced transgenically through the generation of a mammal or plant that is transgenic for recombinant molecules such as the immunoglobulin heavy and light chain sequences and that produces the desired compounds in a recoverable form. This includes, for example, the production of protein modulators (e.g., antibodies) in, and recovery from, the milk of goats, cows, or other mammals. See, e.g., U.S.P.Ns. 5,827,690, 5,756,687, 5,750,172, and 5,741,957. In some embodiments, non-human transgenic animals that comprise human immunoglobulin loci are immunized to produce antibodies.

[0147] Other transgenic techniques are set forth in Hogan et al., *Manipulating the Mouse Embryo: A Laboratory Manual* 2nd ed., Cold Spring Harbor Press (1999); Jackson et al., *Mouse Genetics and Transgenics: A Practical Approach*, Oxford University Press (2000); and Pinkert, *Transgenic Animal Technology: A Laboratory Handbook*, Academic Press (1999) and U.S. P.N. 6,417,429. In some embodiments, the non-human animals are mice, rats, sheep, pigs, goats, cattle or horses, and the desired product is produced in blood, milk, urine, saliva, tears, mucus and other bodily fluids from which it is readily obtainable using art-recognized purification techniques.

[0148] Other compatible production systems include methods for making antibodies in plants such as described, for example, in U.S.P.Ns. 6,046,037 and 5,959,177 which are incorporated herein with respect to such techniques.

E. Isolation/Purification

[0149] Once a modulator of the invention has been produced by recombinant expression or any other of the disclosed techniques, it may be purified by any method known in the art for purification of immunoglobulins or proteins. In this respect the modulator may be "isolated" which means that it has been identified and separated and/or recovered from a component of its natural environment. Contaminant components of its natural environment are materials that would interfere with diagnostic or therapeutic uses for the polypeptide and may include enzymes, hormones, and other proteinaceous or nonproteinaceous solutes. Isolated modulators include a modulator *in situ* within recombinant cells because at least one component of the polypeptide's natural environment will not be present.

[0150] If the desired molecule is produced intracellularly, as a first step, the particulate debris, either host cells or lysed fragments, may be removed, for example, by centrifugation or ultrafiltration. Where the modulator is secreted into the medium, supernatants from such expression systems are generally first concentrated using a commercially available protein concentration filter, for example, an Amicon or Pellicon ultrafiltration unit (Millipore Corp.). Once the insoluble contaminants are removed the modulator preparation may be further purified using standard techniques such as, for example, hydroxylapatite chromatography, gel electrophoresis, dialysis, and affinity chromatography, with affinity chromatography of particular interest. In this regard protein A can be used to purify antibodies that are based on human IgG1, IgG2 or IgG4 heavy chains (Lindmark, et al., J Immunol Meth 62:1 (1983)) while protein G is recommended for all mouse isotypes and for human IgG3 (Guss, et al., EMBO J 5:1567 (1986)). Other techniques for protein purification such as fractionation on an ion-exchange column, ethanol precipitation, reverse phase HPLC, chromatography on silica, chromatography on heparin, sepharose chromatography on an anion or cation exchange resin (such as a polyaspartic acid column), chromatofocusing, SDS-PAGE and ammonium sulfate precipitation are also available depending on the antibody to be recovered. In particularly preferred embodiments the modulators of the instant invention will be purified, at least in part, using Protein A or Protein G affinity chromatography.

VI. SEZ6 Modulator Fragments and Derivatives

[0151] Whatever generation and production methodology is selected, modulators of the instant invention will react, bind, combine, complex, connect, attach, join, interact or otherwise associate with a target determinant (e.g., antigen) and thereby provide the desired results. Where the modulator comprises an antibody or fragment, construct or derivative thereof such associations may be through one or more "binding sites" or "binding components" expressed on the antibody, where a binding site comprises a region of a polypeptide that is responsible for selectively binding to a target molecule or antigen of interest. Binding domains comprise at least one binding site (e.g., an intact IgG antibody will have two binding domains and two binding sites). Exemplary binding domains include an antibody variable domain, a receptor-binding domain of a ligand, a ligand-binding domain of a receptor or an enzymatic domain.

A. Antibodies

[0152] As noted above, the term "antibody" is intended to cover, at least, polyclonal antibodies, monoclonal antibodies, chimeric antibodies, CDR grafted antibodies, humanized and primatized antibodies, human antibodies, recombinantly produced antibodies, intrabodies, multispecific antibodies, bispecific antibodies, monovalent antibodies, multivalent antibodies, anti-idiotypic antibodies, as well as synthetic antibodies.

B. Fragments

[0153] Regardless of which form of the modulator (e.g. chimeric, humanized, etc.) is selected to practice the invention it will be appreciated that immunoreactive fragments of the same may be used in accordance with the teachings herein. An "antibody fragment" comprises at least a portion of an intact antibody. As used herein, the term "fragment" of an antibody molecule includes antigen-binding fragments of antibodies, and the term "antigen-binding fragment" refers to a polypeptide fragment of an immunoglobulin or antibody that immunospecifically binds or reacts with a selected antigen or immunogenic determinant thereof or competes with the intact antibody from which the fragments were derived for specific antigen binding.

[0154] Exemplary fragments include: V_L , V_H , scFv, F(ab')₂ fragment, Fab fragment, Fd fragment, Fv fragment, single domain antibody fragments, diabodies, linear antibodies, single-chain antibody molecules and multispecific antibodies formed from antibody fragments. In addition, an active fragment comprises a portion of the antibody that retains its ability to interact with the antigen/substrates or receptors and modify them in a manner similar to that of an intact antibody (though maybe with somewhat less efficiency).

[0155] In other embodiments, an antibody fragment is one that comprises the Fc region and that retains at least one of the biological functions normally associated with the Fc region when present in an intact antibody, such as FcRn

binding, antibody half-life modulation, ADCC function and complement binding. In one embodiment, an antibody fragment is a monovalent antibody that has an *in vivo* half-life substantially similar to an intact antibody. For example, such an antibody fragment may comprise an antigen binding arm linked to an Fc sequence capable of conferring *in vivo* stability to the fragment.

[0156] As would be well recognized by those skilled in the art, fragments can be obtained via chemical or enzymatic treatment (such as papain or pepsin) of an intact or complete antibody or antibody chain or by recombinant means. See, e.g., *Fundamental Immunology*, W. E. Paul, ed., Raven Press, N.Y. (1999), for a more detailed description of antibody fragments.

C. Derivatives

[0157] The invention further includes immunoreactive modulator derivatives and antigen binding molecules comprising one or more modifications.

1. Multivalent Antibodies

[0158] In one embodiment, the modulators of the invention may be monovalent or multivalent (e.g., bivalent, trivalent, etc.). As used herein, the term "valency" refers to the number of potential target binding sites associated with an antibody. Each target binding site specifically binds one target molecule or specific position or locus on a target molecule. When an antibody is monovalent, each binding site of the molecule will specifically bind to a single antigen position or epitope. When an antibody comprises more than one target binding site (multivalent), each target binding site may specifically bind the same or different molecules (e.g., may bind to different ligands or different antigens, or different epitopes or positions on the same antigen). See, for example, U.S.P.N. 2009/0130105. In each case at least one of the binding sites will comprise an epitope, motif or domain associated with a SEZ6 isoform.

[0159] In one embodiment, the modulators are bispecific antibodies in which the two chains have different specificities, as described in Millstein et al., 1983, *Nature*, 305:537-539. Other embodiments include antibodies with additional specificities such as trispecific antibodies. Other more sophisticated compatible multispecific constructs and methods of their fabrication are set forth in U.S.P.N. 2009/0155255, as well as WO 94/04690; Suresh et al., 1986, *Methods in Enzymology*, 121:210; and WO96/27011.

[0160] As alluded to above, multivalent antibodies may immunospecifically bind to different epitopes of the desired target molecule or may immunospecifically bind to both the target molecule as well as a heterologous epitope, such as a heterologous polypeptide or solid support material. While preferred embodiments of the anti-SEZ6 antibodies only bind two antigens (i.e. bispecific antibodies), antibodies with additional specificities such as trispecific antibodies are also encompassed by the instant invention. Bispecific antibodies also include cross-linked or "heteroconjugate" antibodies. For example, one of the antibodies in the heteroconjugate can be coupled to avidin, the other to biotin. Such antibodies have, for example, been proposed to target immune system cells to unwanted cells (U.S.P.N. 4,676,980), and for treatment of HIV infection (WO 91/00360, WO 92/200373, and EP 03089). Heteroconjugate antibodies may be made using any convenient cross-linking methods. Suitable cross-linking agents are well known in the art, and are disclosed in U.S. P.N. 4,676,980, along with a number of cross-linking techniques.

[0161] In yet other embodiments, antibody variable domains with the desired binding specificities (antibody-antigen combining sites) are fused to immunoglobulin constant domain sequences, such as an immunoglobulin heavy chain constant domain comprising at least part of the hinge, C_H2, and/or C_H3 regions, using methods well known to those of ordinary skill in the art.

2. Fc Region Modifications

[0162] In addition to the various modifications, substitutions, additions or deletions to the variable or binding region of the disclosed modulators (e.g., Fc-SEZ6 or anti-SEZ6 antibodies) set forth above, those skilled in the art will appreciate that selected embodiments of the present invention may also comprise substitutions or modifications of the constant region (i.e. the Fc region). More particularly, it is contemplated that the SEZ6 modulators of the invention may contain *enter alia* one or more additional amino acid residue substitutions, mutations and/or modifications which result in a compound with preferred characteristics including, but not limited to: altered pharmacokinetics, increased serum half life, increase binding affinity, reduced immunogenicity, increased production, altered Fc ligand binding to an Fc receptor (FcR), enhanced or reduced "ADCC" (antibody-dependent cell mediated cytotoxicity) or "CDC" (complement-dependent cytotoxicity) activity, altered glycosylation and/or disulfide bonds and modified binding specificity. In this regard it will be appreciated that these Fc variants may advantageously be used to enhance the effective anti-neoplastic properties of the disclosed modulators.

[0163] To this end certain embodiments of the invention may comprise substitutions or modifications of the Fc region,

for example the addition of one or more amino acid residue, substitutions, mutations and/or modifications to produce a compound with enhanced or preferred Fc effector functions. For example, changes in amino acid residues involved in the interaction between the Fc domain and an Fc receptor (e.g., Fc γ RI, Fc γ RIIA and B, Fc γ RIII and FcRn) may lead to increased cytotoxicity and/or altered pharmacokinetics, such as increased serum half-life (see, for example, Ravetch and Kinet, *Annu. Rev. Immunol* 9:457-92 (1991); Capel et al., *Immunomethods* 4:25-34 (1994); and de Haas et al., *J. Lab. Clin. Med.* 126:330-41 (1995) each of which is incorporated herein by reference).

[0164] In selected embodiments, antibodies with increased *in vivo* half-lives can be generated by modifying (e.g., substituting, deleting or adding) amino acid residues identified as involved in the interaction between the Fc domain and the FcRn receptor (see, e.g., International Publication Nos. WO 97/34631; WO 04/029207; U.S.P.N. 6,737,056 and U.S.P.N. 2003/0190311. With regard to such embodiments, Fc variants may provide half-lives in a mammal, preferably a human, of greater than 5 days, greater than 10 days, greater than 15 days, preferably greater than 20 days, greater than 25 days, greater than 30 days, greater than 35 days, greater than 40 days, greater than 45 days, greater than 2 months, greater than 3 months, greater than 4 months, or greater than 5 months. The increased half-life results in a higher serum titer which thus reduces the frequency of the administration of the antibodies and/or reduces the concentration of the antibodies to be administered. Binding to human FcRn *in vivo* and serum half life of human FcRn high affinity binding polypeptides can be assayed, e.g., in transgenic mice or transfected human cell lines expressing human FcRn, or in primates to which the polypeptides with a variant Fc region are administered. WO 2000/42072 describes antibody variants with improved or diminished binding to FcRns. See also, e.g., Shields et al. *J. Biol. Chem.* 9(2):6591-6604 (2001).

[0165] In other embodiments, Fc alterations may lead to enhanced or reduced ADCC or CDC activity. As is known in the art, CDC refers to the lysing of a target cell in the presence of complement, and ADCC refers to a form of cytotoxicity in which secreted Ig bound onto FcRs present on certain cytotoxic cells (e.g., Natural Killer cells, neutrophils, and macrophages) enables these cytotoxic effector cells to bind specifically to an antigen-bearing target cell and subsequently kill the target cell with cytotoxins. In the context of the instant invention antibody variants are provided with "altered" FcR binding affinity, which is either enhanced or diminished binding as compared to a parent or unmodified antibody or to an antibody comprising a native sequence FcR. Such variants which display decreased binding may possess little or no appreciable binding, e.g., 0-20% binding to the FcR compared to a native sequence, e.g. as determined by techniques well known in the art. In other embodiments the variant will exhibit enhanced binding as compared to the native immunoglobulin Fc domain. It will be appreciated that these types of Fc variants may advantageously be used to enhance the effective anti-neoplastic properties of the disclosed antibodies. In yet other embodiments, such alterations lead to increased binding affinity, reduced immunogenicity, increased production, altered glycosylation and/or disulfide bonds (e.g., for conjugation sites), modified binding specificity, increased phagocytosis; and/or down regulation of cell surface receptors (e.g. B cell receptor; BCR), etc.

3. Altered Glycosylation

[0166] Still other embodiments comprise one or more engineered glycoforms, i.e., a SEZ6 modulator comprising an altered glycosylation pattern or altered carbohydrate composition that is covalently attached to the protein (e.g., in the Fc domain). See, for example, Shields, R. L. et al. (2002) *J. Biol. Chem.* 277:26733-26740. Engineered glycoforms may be useful for a variety of purposes, including but not limited to enhancing or reducing effector function, increasing the affinity of the modulator for a target or facilitating production of the modulator. In certain embodiments where reduced effector function is desired, the molecule may be engineered to express an aglycosylated form. Substitutions that may result in elimination of one or more variable region framework glycosylation sites to thereby eliminate glycosylation at that site are well known (see e.g. U.S. P.Ns. 5,714,350 and 6,350,861). Conversely, enhanced effector functions or improved binding may be imparted to the Fc containing molecule by engineering in one or more additional glycosylation sites.

[0167] Other embodiments include an Fc variant that has an altered glycosylation composition, such as a hypofucosylated antibody having reduced amounts of fucosyl residues or an antibody having increased bisecting GlcNAc structures. Such altered glycosylation patterns have been demonstrated to increase the ADCC ability of antibodies. Engineered glycoforms may be generated by any method known to one skilled in the art, for example by using engineered or variant expression strains, by co-expression with one or more enzymes (for example N-acetylglucosaminyltransferase III (GnT11)), by expressing a molecule comprising an Fc region in various organisms or cell lines from various organisms or by modifying carbohydrate(s) after the molecule comprising Fc region has been expressed (see, for example, WO 2012/117002).

4. Additional Processing

[0168] The modulators may be differentially modified during or after production, e.g., by glycosylation, acetylation,

phosphorylation, amidation, derivatization by known protecting/blocking groups, proteolytic cleavage, linkage to an antibody molecule or other cellular ligand, etc. Any of numerous chemical modifications may be carried out by known techniques, including but not limited, to specific chemical cleavage by cyanogen bromide, trypsin, chymotrypsin, papain, V8 protease, NaBH₄, acetylation, formylation, oxidation, reduction, metabolic synthesis in the presence of tunicamycin, etc.

[0169] Various post-translational modifications also encompassed by the invention include, for example, N-linked or O-linked carbohydrate chains, processing of N-terminal or C-terminal ends, attachment of chemical moieties to the amino acid backbone, chemical modifications of N-linked or O-linked carbohydrate chains, and addition or deletion of an N-terminal methionine residue as a result of prokaryotic host cell expression. Moreover, the modulators may also be modified with a detectable label, such as an enzymatic, fluorescent, radioisotopic or affinity label to allow for detection and isolation of the modulator.

VII. Modulator Characteristics

[0170] No matter how obtained or which of the aforementioned forms the modulator takes, various embodiments of the disclosed modulators may exhibit certain characteristics. In selected embodiments, antibody-producing cells (e.g., hybridomas or yeast colonies) may be selected, cloned and further screened for favorable properties including, for example, robust growth, high modulator production and, as discussed in more detail below, desirable modulator characteristics. In other cases characteristics of the modulator may be imparted or influenced by selecting a particular antigen (e.g., a specific SEZ6 isoform or fragment thereof) or immunoreactive fragment of the target antigen for inoculation of the animal. In still other embodiments the selected modulators may be engineered as described above to enhance or refine immunochemical characteristics such as affinity or pharmacokinetics.

A. Neutralizing Modulators

[0171] In certain embodiments, the modulators will comprise "neutralizing" antibodies or derivatives or fragments thereof. That is, the present invention may comprise antibody molecules that bind specific domains, motifs or epitopes and are capable of blocking, reducing or inhibiting the biological activity of SEZ6. More generally the term "neutralizing antibody" refers to an antibody that binds to or interacts with a target molecule or ligand and prevents binding or association of the target molecule to a binding partner such as a receptor or substrate, thereby interrupting a biological response that otherwise would result from the interaction of the molecules.

[0172] It will be appreciated that competitive binding assays known in the art may be used to assess the binding and specificity of an antibody or immunologically functional fragment or derivative thereof. With regard to the instant invention an antibody or fragment will be held to inhibit or reduce binding of SEZ6 to a binding partner or substrate (e.g., a neurotrophic ligand) when an excess of antibody reduces the quantity of binding partner bound to SEZ6 by at least about 20%, 30%, 40%, 50%, 60%, 70%, 80%, 85%, 90%, 95%, 97%, 99% or more as measured, for example, by impaired neurotrophic ligand activity or in an *in vitro* competitive binding assay. In the case of antibodies to SEZ6 for example, a neutralizing antibody or antagonist will preferably alter ligand activity by at least about 20%, 30%, 40%, 50%, 60%, 70%, 80%, 85%, 90%, 95%, 97%, 99% or more. It will be appreciated that this modified activity may be measured directly using art-recognized techniques or may be measured by the impact the altered activity has downstream (e.g., oncogenesis, cell survival or pathway activation).

B. Internalizing Modulators

[0173] While evidence indicates that SEZ6 or selected isoforms thereof may be present in a soluble form, at least some SEZ6 likely remains associated with the cell surface thereby allowing for internalization of the disclosed modulators. Accordingly, the anti-SEZ6 antibodies of the instant invention may be internalized, at least to some extent, by cells that express SEZ6. For example, an anti-SEZ6 antibody that binds to SEZ6 on the surface of a tumor-initiating cell may be internalized by the tumor-initiating cell. In particularly preferred embodiments such anti-SEZ6 antibodies may be associated with or conjugated to anti-cancer agents such as cytotoxic moieties that kill the cell upon internalization. In particularly preferred embodiments the modulator will comprise an internalizing antibody drug conjugate.

[0174] As used herein, a modulator that "internalizes" is one that is taken up (along with any payload) by the cell upon binding to an associated antigen or receptor. As will be appreciated, the internalizing modulator may, in preferred embodiments, comprise an antibody including antibody fragments and derivatives thereof, as well as antibody conjugates. Internalization may occur *in vitro* or *in vivo*. For therapeutic applications, internalization will preferably occur *in vivo* in a subject in need thereof. The number of antibody molecules internalized may be sufficient or adequate to kill an antigen-expressing cell, especially an antigen-expressing cancer stem cell. Depending on the potency of the antibody or antibody conjugate, in some instances, the uptake of a single antibody molecule into the cell is sufficient to kill the target cell to

which the antibody binds. For example, certain toxins are so highly potent that the internalization of a few molecules of the toxin conjugated to the antibody is sufficient to kill the tumor cell. Whether an antibody internalizes upon binding to a mammalian cell can be determined by various assays including those described in the Examples below (e.g., Example 15, 17 and 18). Methods of detecting whether an antibody internalizes into a cell are also described in U.S.P.N. 7,619,068 which is incorporated herein by reference in its entirety.

C. Depleting Modulators

[0175] In other embodiments the antibodies will comprise depleting antibodies or derivatives or fragments thereof. The term "depleting" antibody refers to an antibody that preferably binds to or associates with an antigen on or near the cell surface and induces, promotes or causes the death or elimination of the cell (e.g., by CDC, ADCC or introduction of a cytotoxic agent). In some embodiments, the selected depleting antibodies will be associated or conjugated to a cytotoxic agent.

[0176] Preferably a depleting antibody will be able to remove, incapacitate, eliminate or kill at least 20%, 30%, 40%, 50%, 60%, 70%, 80%, 85%, 90%, 95%, 97%, or 99% of SEZ6 tumorigenic cells in a defined cell population. In some embodiments the cell population may comprise enriched, sectioned, purified or isolated tumor perpetuating cells. In other embodiments the cell population may comprise whole tumor samples or heterogeneous tumor extracts that comprise tumor perpetuating cells. Those skilled in the art will appreciate that standard biochemical techniques as described in the Examples below (e.g., Examples 14 and 15) may be used to monitor and quantify the depletion of tumorigenic cells or tumor perpetuating cells in accordance with the teachings herein.

D. Binning and Epitope Binding

[0177] It will further be appreciated the disclosed anti-SEZ6 antibody modulators will associate with, or bind to, discrete epitopes or immunogenic determinants presented by the selected target or fragment thereof. In certain embodiments, epitope or immunogenic determinants include chemically active surface groupings of molecules such as amino acids, sugar side chains, phosphoryl groups, or sulfonyl groups, and, in certain embodiments, may have specific three-dimensional structural characteristics, and/or specific charge characteristics. Thus, as used herein the term "epitope" includes any protein determinant capable of specific binding to an immunoglobulin or T-cell receptor or otherwise interacting with a molecule. In certain embodiments, an antibody is said to specifically bind (or immunospecifically bind or react) an antigen when it preferentially recognizes its target antigen in a complex mixture of proteins and/or macromolecules. In preferred embodiments, an antibody is said to specifically bind an antigen when the equilibrium dissociation constant (K_D) is less than or equal to $10^{-6}M$ or less than or equal to $10^{-7}M$, more preferably when the equilibrium dissociation constant is less than or equal to $10^{-8}M$, and even more preferably when the dissociation constant is less than or equal to $10^{-9}M$.

[0178] More directly the term "epitope" is used in its common biochemical sense and refers to that portion of the target antigen capable of being recognized and specifically bound by a particular antibody modulator. When the antigen is a polypeptide such as SEZ6, epitopes may generally be formed from both contiguous amino acids and noncontiguous amino acids juxtaposed by tertiary folding of a protein ("conformational epitopes"). In such conformational epitopes the points of interaction occur across amino acid residues on the protein that are linearly separated from one another. Epitopes formed from contiguous amino acids (sometimes referred to as "linear" or "continuous" epitopes) are typically retained upon protein denaturing, whereas epitopes formed by tertiary folding are typically lost upon protein denaturing. In any event an antibody epitope typically includes at least 3, and more usually, at least 5 or 8-10 amino acids in a unique spatial conformation.

[0179] In this respect it will be appreciated that, in certain embodiments, an epitope may be associated with, or reside in, one or more regions, domains or motifs of the SEZ6 protein (e.g., amino acids 1-906 of mature isoform 1). As discussed in more detail herein the extracellular region of the SEZ6 protein comprises a series of generally recognized domains including five Sushi domains and two CUB domains along with an N-terminal domain. For the purposes of the instant disclosure the term "domain" will be used in accordance with its generally accepted meaning and will be held to refer to an identifiable or definable conserved structural entity within a protein that exhibits a distinctive secondary structure content. In many cases, homologous domains with common functions will usually show sequence similarities and be found in a number of disparate proteins (e.g., Sushi domains are reportedly found in a large number of different proteins). Similarly, the art-recognized term "motif" will be used in accordance with its common meaning and shall generally refer to a short, conserved region of a protein that is typically ten to twenty contiguous amino acid residues. As discussed throughout, selected embodiments comprise modulators that associate with or bind to an epitope within specific regions, domains or motifs of SEZ6.

[0180] In any event once a desired epitope on an antigen is determined, it is possible to generate antibodies to that epitope, e.g., by immunizing with a peptide comprising the epitope using techniques described in the present invention.

Alternatively, during the discovery process, the generation and characterization of antibodies may elucidate information about desirable epitopes located in specific domains or motifs. From this information, it is then possible to competitively screen antibodies for binding to the same epitope. An approach to achieve this is to conduct competition studies to find antibodies that competitively bind with one another, i.e. the antibodies compete for binding to the antigen. A high throughput process for binning antibodies based upon their cross-competition is described in WO 03/48731. Other methods of binning or domain level or epitope mapping comprising modulator competition or antigen fragment expression on yeast is set forth in Examples 9 and 10 below.

[0181] As used herein, the term "binning" refers to methods used to group or classify antibodies based on their antigen binding characteristics and competition. While the techniques are useful for defining and categorizing modulators of the instant invention, the bins do not always directly correlate with epitopes and such initial determinations of epitope binding may be further refined and confirmed by other art-recognized methodology as described herein. However, as discussed and shown in the Examples below, empirical assignment of antibody modulators to individual bins provides information that may be indicative of the therapeutic potential of the disclosed modulators.

[0182] More specifically, one can determine whether a selected reference antibody (or fragment thereof) binds to the same epitope or cross competes for binding with a second test antibody (i.e., is in the same bin) by using methods known in the art and set forth in the Examples herein. In one embodiment, a reference antibody modulator is associated with SEZ6 antigen under saturating conditions and then the ability of a secondary or test antibody modulator to bind to SEZ6 is determined using standard immunochemical techniques. If the test antibody is able to substantially bind to SEZ6 at the same time as the reference anti-SEZ6 antibody, then the secondary or test antibody binds to a different epitope than the primary or reference antibody. However, if the test antibody is not able to substantially bind to SEZ6 at the same time, then the test antibody binds to the same epitope, an overlapping epitope, or an epitope that is in close proximity (at least sterically) to the epitope bound by the primary antibody. That is, the test antibody competes for antigen binding and is in the same bin as the reference antibody.

[0183] The term "compete" or "competing antibody" when used in the context of the disclosed modulators means competition between antibodies as determined by an assay in which a test antibody or immunologically functional fragment under test prevents or inhibits specific binding of a reference antibody to a common antigen. Typically, such an assay involves the use of purified antigen (e.g., SEZ6 or a domain or fragment thereof) bound to a solid surface or cells bearing either of these, an unlabeled test immunoglobulin and a labeled reference immunoglobulin. Competitive inhibition is measured by determining the amount of label bound to the solid surface or cells in the presence of the test immunoglobulin. Usually the test immunoglobulin is present in excess and/or allowed to bind first. Antibodies identified by competition assay (competing antibodies) include antibodies binding to the same epitope as the reference antibody and antibodies binding to an adjacent epitope sufficiently proximal to the epitope bound by the reference antibody for steric hindrance to occur. Additional details regarding methods for determining competitive binding are provided in the Examples herein. Usually, when a competing antibody is present in excess, it will inhibit specific binding of a reference antibody to a common antigen by at least 30%, 40%, 45%, 50%, 55%, 60%, 65%, 70% or 75%. In some instance, binding is inhibited by at least 80%, 85%, 90%, 95%, or 97% or more.

[0184] Conversely, when the reference antibody is bound it will preferably inhibit binding of a subsequently added test antibody (i.e., a SEZ6 modulator) by at least 30%, 40%, 45%, 50%, 55%, 60%, 65%, 70% or 75%. In some instances binding of the test antibody is inhibited by at least 80%, 85%, 90%, 95%, or 97% or more.

[0185] With regard to the instant invention, and as set forth in the Examples 9 and 10 below, it has been determined (via surface plasmon resonance or bio-layer interferometry) that the extracellular domain of SEZ6 defines at least seven bins by competitive binding termed "bin A" to "bin F" and bin U herein.

[0186] In this respect, and as known in the art and detailed in the Examples below, the desired binning or competitive binding data can be obtained using solid phase direct or indirect radioimmunoassay (RIA), solid phase direct or indirect enzyme immunoassay (EIA or ELISA), sandwich competition assay, a Biacore™ 2000 system (i.e., surface plasmon resonance - GE Healthcare), a ForteBio® Analyzer (i.e., bio-layer interferometry - ForteBio, Inc.) or flow cytometric methodology. The term "surface plasmon resonance," as used herein, refers to an optical phenomenon that allows for the analysis of real-time specific interactions by detection of alterations in protein concentrations within a biosensor matrix. The term "bio-layer interferometry" refers to an optical analytical technique that analyzes the interference pattern of white light reflected from two surfaces: a layer of immobilized protein on a biosensor tip, and an internal reference layer. Any change in the number of molecules bound to the biosensor tip causes a shift in the interference pattern that can be measured in real-time. In particularly preferred embodiments the analysis (whether surface plasmon resonance, bio-layer interferometry or flow cytometry) is performed using a Biacore or ForteBio instrument or a flow cytometer (e.g., FACSARIA II) as demonstrated in the Examples below.

[0187] In order to further characterize the epitopes that the disclosed SEZ6 antibody modulators associate with or bind to, domain-level epitope mapping was performed using a modification of the protocol described by Cochran et al. (J Immunol Methods. 287 (1-2):147-158 (2004) which is incorporated herein by reference). Briefly, individual domains of SEZ6 comprising specific amino acid sequences were expressed on the surface of yeast and binding by each SEZ6

antibody was determined through flow cytometry. The results are discussed below in Example 10 and shown in FIGS. 14A and 14B.

[0188] Other compatible epitope mapping techniques include alanine scanning mutants, peptide blots (Reineke (2004) Methods Mol Biol 248:443-63) (herein specifically incorporated by reference in its entirety), or peptide cleavage analysis. In addition, methods such as epitope excision, epitope extraction and chemical modification of antigens can be employed (Tomer (2000) Protein Science 9: 487-496) (herein specifically incorporated by reference in its entirety). In other embodiments Modification-Assisted Profiling (MAP), also known as Antigen Structure-based Antibody Profiling (ASAP) provides a method that categorizes large numbers of monoclonal antibodies (mAbs) directed against the same antigen according to the similarities of the binding profile of each antibody to chemically or enzymatically modified antigen surfaces (U.S.P.N. 2004/0101920, herein specifically incorporated by reference in its entirety). Each category may reflect a unique epitope either distinctly different from or partially overlapping with epitope represented by another category. This technology allows rapid filtering of genetically identical antibodies, such that characterization can be focused on genetically distinct antibodies. It will be appreciated that MAP may be used to sort the hSEZ6 antibody modulators of the invention into groups of antibodies binding different epitopes

[0189] Agents useful for altering the structure of the immobilized antigen include enzymes such as proteolytic enzymes (e.g., trypsin, endoproteinase Glu-C, endoproteinase Asp-N, chymotrypsin, etc.). Agents useful for altering the structure of the immobilized antigen may also be chemical agents, such as, succinimidyl esters and their derivatives, primary amine-containing compounds, hydrazines and carbohydrazines, free amino acids, etc.

[0190] The antigen protein may be immobilized on either biosensor chip surfaces or polystyrene beads. The latter can be processed with, for example, an assay such as multiplex LUMINEX™ detection assay (Luminex Corp.). Because of the capacity of LUMINEX to handle multiplex analysis with up to 100 different types of beads, LUMINEX provides almost unlimited antigen surfaces with various modifications, resulting in improved resolution in antibody epitope profiling over a biosensor assay.

E. Modulator Binding Characteristics

[0191] Besides epitope specificity the disclosed antibodies may be characterized using physical characteristics such as, for example, binding affinities. In this regard the present invention further encompasses the use of antibodies that have a high binding affinity for one or more SEZ6 isoforms or, in the case of pan-antibodies, more than one member of the SEZ6 family.

[0192] The term " K_D ", as used herein, is intended to refer to the dissociation constant of a particular antibody-antigen interaction. An antibody of the invention is said to immunospecifically bind its target antigen when the dissociation constant K_D (k_{off}/k_{on}) is $\leq 10^{-7}M$. The antibody specifically binds antigen with high affinity when the K_D is $\leq 5 \times 10^{-9}M$, and with very high affinity when the K_D is $\leq 5 \times 10^{-10}M$. In one embodiment of the invention, the antibody has a K_D of $\leq 10^{-9}M$ and an off-rate of about $1 \times 10^{-4}/sec$. In one embodiment of the invention, the off-rate is $< 1 \times 10^{-5}/sec$. In other embodiments of the invention, the antibodies will bind to SEZ6 with a K_D of between about $10^{-7}M$ and $10^{-10}M$, and in yet another embodiment it will bind with a $K_D \leq 2 \times 10^{-10}M$. Still other selected embodiments of the present invention comprise antibodies that have a disassociation constant or K_D (k_{off}/k_{on}) of less than $10^{-2}M$, less than $5 \times 10^{-2}M$, less than $10^{-3}M$, less than $5 \times 10^{-3}M$, less than $10^{-4}M$, less than $5 \times 10^{-4}M$, less than $10^{-5}M$, less than $5 \times 10^{-5}M$, less than $10^{-6}M$, less than $5 \times 10^{-6}M$, less than $10^{-7}M$, less than $5 \times 10^{-7}M$, less than $10^{-8}M$, less than $5 \times 10^{-8}M$, less than $10^{-9}M$, less than $5 \times 10^{-9}M$, less than $10^{-10}M$, less than $5 \times 10^{-10}M$, less than $10^{-11}M$, less than $5 \times 10^{-11}M$, less than $10^{-12}M$, less than $5 \times 10^{-12}M$, less than $10^{-13}M$, less than $5 \times 10^{-13}M$, less than $10^{-14}M$, less than $5 \times 10^{-14}M$, less than $10^{-15}M$ or less than $5 \times 10^{-15}M$.

[0193] In specific embodiments, an antibody of the invention that immunospecifically binds to SEZ6 has an association rate constant or k_{on} (or k_a) rate (SEZ6 (Ab) + antigen (Ag) $\xrightarrow{k_{on}}$ Ab-Ag) of at least $10^5 M^{-1}s^{-1}$, at least $2 \times 10^5 M^{-1}s^{-1}$, at least $5 \times 10^5 M^{-1}s^{-1}$, at least $10^6 M^{-1}s^{-1}$, at least $5 \times 10^6 M^{-1}s^{-1}$, at least $10^7 M^{-1}s^{-1}$, at least $5 \times 10^7 M^{-1}s^{-1}$, or at least $10^8 M^{-1}s^{-1}$.

[0194] In another embodiment, an antibody of the invention that immunospecifically binds to SEZ6 has a disassociation rate constant or k_{off} (or k_d) rate (SEZ6 (Ab) + antigen (Ag) $\xrightarrow{k_{off}}$ Ab-Ag) of less than $10^{-1}s^{-1}$, less than $5 \times 10^{-1}s^{-1}$, less than $10^{-2}s^{-1}$, less than $5 \times 10^{-2}s^{-1}$, less than $10^{-3}s^{-1}$, less than $5 \times 10^{-3}s^{-1}$, less than $10^{-4}s^{-1}$, less than $5 \times 10^{-4}s^{-1}$, less than $10^{-5}s^{-1}$, less than $5 \times 10^{-5}s^{-1}$, less than $10^{-6}s^{-1}$, less than $5 \times 10^{-6}s^{-1}$, less than $10^{-7}s^{-1}$, less than $5 \times 10^{-7}s^{-1}$, less than $10^{-8}s^{-1}$, less than $5 \times 10^{-8}s^{-1}$, less than $10^{-9}s^{-1}$ less than $5 \times 10^{-9}s^{-1}$ or less than $10^{-10}s^{-1}$.

[0195] In other selected embodiments of the present invention anti-SEZ6 antibodies will have an affinity constant or K_a (k_{on}/k_{off}) of at least $10^2 M^{-1}$, at least $5 \times 10^2 M^{-1}$, at least $10^3 M^{-1}$, at least $5 \times 10^3 M^{-1}$, at least $10^4 M^{-1}$, at least $5 \times 10^4 M^{-1}$, at least $10^5 M^{-1}$, at least $5 \times 10^5 M^{-1}$, at least $10^6 M^{-1}$, at least $5 \times 10^6 M^{-1}$, at least $10^7 M^{-1}$, at least $5 \times 10^7 M^{-1}$, at least $10^8 M^{-1}$, at least $5 \times 10^8 M^{-1}$, at least $10^9 M^{-1}$, at least $5 \times 10^9 M^{-1}$, at least $10^{10} M^{-1}$, at least $5 \times 10^{10} M^{-1}$, at least $10^{11} M^{-1}$, at least $5 \times 10^{11} M^{-1}$, at least $10^{12} M^{-1}$, at least $5 \times 10^{12} M^{-1}$, at least $10^{13} M^{-1}$, at least $5 \times 10^{13} M^{-1}$, at least $10^{14} M^{-1}$, at least $5 \times 10^{14} M^{-1}$, at least $10^{15} M^{-1}$ or at least $5 \times 10^{15} M^{-1}$.

[0196] Besides the aforementioned modulator characteristics antibodies of the instant invention may further be characterized using additional physical characteristics including, for example, thermal stability (i.e., melting temperature; T_m),

and isoelectric points. (See, e.g., Bjellqvist et al., 1993, Electrophoresis 14:1023; Vermeer et al., 2000, Biophys. J. 78:394-404; Vermeer et al., 2000, Biophys. J. 79: 2150-2154 each of which is incorporated by reference).

VIII. Conjugated Modulators

A. Overview

[0197] Once the modulators of the invention have been generated and/or fabricated and selected according to the teachings herein they may be linked with, fused to, conjugated to (e.g., covalently or non-covalently) or otherwise associated with pharmaceutically active or diagnostic moieties, therapeutic moieties or biocompatible modifiers. The modulators (e.g. antibodies) of the invention may be conjugated directly or indirectly to therapeutic moieties. An antibody is "conjugated directly" to a therapeutic moiety or other moiety, for example, a reporter, when such antibody is associated, linked or fused with such moiety without using a linker (linkers are described in more detail below) to connect the antibody to the therapeutic moiety or other moiety. An antibody is "conjugated indirectly" to a therapeutic moiety or other moiety, for example, a reporter, when the antibody is associated, linked or fused to such moiety using a linker that connects the antibody to such moiety. As used herein the term "conjugate" or "modulator conjugate" or "antibody conjugate" will be used broadly and held to mean any biologically active or detectable molecule or drug associated with the disclosed modulators regardless of the method of association. In this respect it will be understood that such conjugates may, in addition to the disclosed modulators, comprise peptides, polypeptides, proteins, prodrugs which are metabolized to an active agent *in vivo*, polymers, nucleic acid molecules, small molecules, binding agents, mimetic agents, synthetic drugs, inorganic molecules, organic molecules and radioisotopes. Moreover, as indicated above the selected conjugate may be covalently or non-covalently associated with, or linked to, the modulator and exhibit various stoichiometric molar ratios depending, at least in part, on the method used to effect the conjugation.

[0198] Particularly preferred aspects of the instant invention will comprise antibody modulator conjugates or antibody-drug conjugates that may be used for the diagnosis and/or treatment of proliferative disorders. It will be appreciated that, unless otherwise dictated by context, the term "antibody-drug conjugate" or "ADC" or the formula $M-[L-D]_n$ shall be held to encompass conjugates comprising both therapeutic and diagnostic moieties. In such embodiments antibody-drug conjugate compounds will comprise a SEZ6 modulator (typically an anti-SEZ6 antibody) as the modulator or cellular binding unit (abbreviated as CBA, M, or Ab herein), a therapeutic (e.g., anti-cancer agent) or diagnostic moiety (D), and optionally a linker (L) that joins the drug and the antigen binding agent. For the purposes of the instant disclosure "n" shall be held to mean an integer from 1 to 20. In a preferred embodiment, the modulator is a SEZ6 mAb comprising at least one CDR from the heavy and light chain variable regions as described above.

[0199] Those skilled in the art will appreciate that a number of different reactions are available for the attachment or association of therapeutic or diagnostic moieties and/or linkers to binding agents. In selected embodiments this may be accomplished by reaction of the amino acid residues of the binding agent, e.g., antibody molecule, including the amine groups of lysine, the free carboxylic acid groups of glutamic and aspartic acid, the sulfhydryl groups of cysteine and the various moieties of the aromatic amino acids. One of the most commonly used non-specific methods of covalent attachment is the carbodiimide reaction to link a carboxy (or amino) group of a compound to amino (or carboxy) groups of the antibody. Additionally, bifunctional agents such as dialdehydes or imidoesters have been used to link the amino group of a compound to amino groups of an antibody molecule. Also available for attachment of drugs to binding agents is the Schiff base reaction. This method involves the periodate oxidation of a drug that contains glycol or hydroxy groups, thus forming an aldehyde which is then reacted with the binding agent. Attachment occurs via formation of a Schiff base with amino groups of the binding agent. Isothiocyanates and azlactones can also be used as coupling agents for covalently attaching drugs to binding agents.

[0200] In other embodiments the disclosed modulators of the invention may be conjugated or associated with proteins, polypeptides or peptides that impart selected characteristics (e.g., biotoxins, biomarkers, purification tags, etc.). In certain preferred embodiments the present invention encompasses the use of modulators or fragments thereof recombinantly fused or chemically conjugated (including both covalent and non-covalent conjugations) to a heterologous protein or peptide wherein the protein or peptide comprises at least 10, at least 20, at least 30, at least 40, at least 50, at least 60, at least 70, at least 80, at least 90 or at least 100 amino acids. The construct does not necessarily need to be directly linked, but may occur through amino acid linker sequences. For example, antibodies may be used to target heterologous polypeptides to particular cell types expressing SEZ6, either *in vitro* or *in vivo*, by fusing or conjugating the modulators of the present invention to antibodies specific for particular cell surface receptors to provide bispecific constructs. Moreover, modulators fused or conjugated to heterologous polypeptides may also be used in *in vitro* immunoassays and may be particularly compatible with purification methodology (e.g., his-tags) as is known in the art. See e.g., International publication No. WO 93/21232; European Patent No. EP 439,095; Naramura et al., 1994, Immunol. Lett. 39:91-99; U.S. Pat. No. 5,474,981; Gillies et al., 1992, PNAS 89:1428-1432; and Fell et al., 1991, J. Immunol. 146:2446-2452.

B. Linkers

[0201] Besides the aforementioned peptide linkers or spacers, it will be appreciated that several other varieties or types of linker may be used to associate the disclosed modulators with pharmaceutically active or diagnostic moieties or biocompatible modifiers. In some embodiments, the linker is cleavable under intracellular conditions, such that cleavage of the linker releases the drug unit from the antibody in the intracellular environment. In yet other embodiments, the linker unit is not cleavable and the drug is released, for example, by antibody degradation.


[0202] The linkers of the ADC are preferably stable extracellularly, prevent aggregation of ADC molecules and keep the ADC freely soluble in aqueous media and in a monomeric state. Before transport or delivery into a cell, the antibody-drug conjugate (ADC) is preferably stable and remains intact, i.e. the antibody remains linked to the drug moiety. The linkers are stable outside the target cell and may be cleaved at some efficacious rate inside the cell. An effective linker will: (i) maintain the specific binding properties of the antibody; (ii) allow intracellular delivery of the conjugate or drug moiety; (iii) remain stable and intact, i.e. not cleaved, until the conjugate has been delivered or transported to its targeted site; and (iv) maintain a cytotoxic, cell-killing effect or a cytostatic effect of the PBD drug moiety. Stability of the ADC may be measured by standard analytical techniques such as mass spectroscopy, HPLC, and the separation/analysis technique LC/MS. Covalent attachment of the antibody and the drug moiety requires the linker to have two reactive functional groups, i.e. bivalency in a reactive sense. Bivalent linker reagents which are useful to attach two or more functional or biologically active moieties, such as peptides, nucleic acids, drugs, toxins, antibodies, haptens, and reporter groups are known, and methods have been described their resulting conjugates (Hermanson, G.T. (1996) Bioconjugate Techniques; Academic Press: New York, p 234-242).

[0203] To this end certain embodiments of the invention comprise the use a linker that is cleavable by a cleaving agent that is present in the intracellular environment (e.g., within a lysosome or endosome or caveolae). The linker can be, for example, a peptidyl linker that is cleaved by an intracellular peptidase or protease enzyme, including, but not limited to, a lysosomal or endosomal protease. In some embodiments, the peptidyl linker is at least two amino acids long or at least three amino acids long. Cleaving agents can include cathepsins B and D and plasmin, each of which is known to hydrolyze dipeptide drug derivatives resulting in the release of active drug inside target cells. Exemplary peptidyl linkers that are cleavable by the thiol-dependent protease Cathepsin-B are peptides comprising Phe-Leu since Cathepsin-B has been found to be highly expressed in cancerous tissue. Other examples of such linkers are described, for example, in U.S.P.N. 6,214,345 and U.S.P.N. 2012/0078028 each of which incorporated herein by reference in its entirety. In a specific preferred embodiment, the peptidyl linker cleavable by an intracellular protease is a Val-Cit linker, an Ala-Val linker or a Phe-Lys linker such as is described in U.S.P.N. 6,214,345. One advantage of using intracellular proteolytic release of the therapeutic agent is that the agent is typically attenuated when conjugated and the serum stabilities of the conjugates are typically high.

[0204] In other embodiments, the cleavable linker is pH-sensitive, i.e., sensitive to hydrolysis at certain pH values. Typically, the pH-sensitive linker hydrolyzable under acidic conditions. For example, an acid-labile linker that is hydrolyzable in the lysosome (e.g., a hydrazone, oxime, semicarbazone, thiosemicarbazone, cis-aconitic amide, orthoester, acetal, ketal, or the like) can be used (See, e.g., U.S.P.N. 5,122,368; 5,824,805; 5,622,929). Such linkers are relatively stable under neutral pH conditions, such as those in the blood, but are unstable at below pH 5.5 or 5.0, the approximate pH of the lysosome.

[0205] In yet other embodiments, the linker is cleavable under reducing conditions (e.g., a disulfide linker). A variety of disulfide linkers are known in the art, including, for example, those that can be formed using SATA (N-succinimidyl-S-acetylthioacetate), SPDP (N-succinimidyl-3-(2-pyridyldithio)propionate), SPDB (N-succinimidyl-3-(2-pyridyldithio) butyrate) and SMPT (N-succinimidyl-oxycarbonyl-alpha-methyl-alpha-(2-pyridyl-dithio)toluene). In yet other specific embodiments, the linker is a malonate linker (Johnson et al., 1995, Anticancer Res. 15:1387-93), a maleimidobenzoyl linker (Lau et al., 1995, Bioorg-Med-Chem. 3(10):1299-1304), or a 3'-N-amide analog (Lau et al., 1995, Bioorg-Med-Chem. 3(10):1305-12). In yet other embodiments, the linker unit is not cleavable and the drug is released by antibody degradation. (See U.S. Publication No. 2005/0238649 incorporated by reference herein in its entirety and for all purposes).

[0206] More particularly, in preferred embodiments (set forth in U.S.P.N. 2011/0256157 which is incorporated herein by reference in its entirety) compatible linkers will comprise:



where the asterisk indicates the point of attachment to the cytotoxic agent, CBA is a cell binding agent/modulator, L^1 is a linker, A is a connecting group connecting L^1 to the cell binding agent, L^2 is a covalent bond or together with $-OC(=O)-$

forms a self-immolative linker, and L^1 or L^2 is a cleavable linker.

[0207] L^1 is preferably the cleavable linker, and may be referred to as a trigger for activation of the linker for cleavage.

[0208] The nature of L^1 and L^2 , where present, can vary widely. These groups are chosen on the basis of their cleavage characteristics, which may be dictated by the conditions at the site to which the conjugate is delivered. Those linkers that are cleaved by the action of enzymes are preferred, although linkers that are cleavable by changes in pH (e.g. acid or base labile), temperature or upon irradiation (e.g. photolabile) may also be used. Linkers that are cleavable under reducing or oxidising conditions may also find use in the present invention.

[0209] L^1 may comprise a contiguous sequence of amino acids. The amino acid sequence may be the target substrate for enzymatic cleavage, thereby allowing release of R^{10} from the N10 position.

[0210] In one embodiment, L^1 is cleavable by the action of an enzyme. In one embodiment, the enzyme is an esterase or a peptidase.

[0211] In one embodiment, L^2 is present and together with $-C(=O)O-$ forms a self-immolative linker. In one embodiment, L^2 is a substrate for enzymatic activity, thereby allowing release of R^{10} from the N10 position.

[0212] In one embodiment, where L^1 is cleavable by the action of an enzyme and L^2 is present, the enzyme cleaves the bond between L^1 and L^2 .

[0213] L^1 and L^2 , where present, may be connected by a bond selected from:

$-C(=O)NH-$, $-C(=O)O-$, $-NHC(=O)-$, $-OC(=O)-$, $-OC(=O)O-$, $-NHC(=O)O-$, $-OC(=O)NH-$, and $-NHC(=O)NH-$.


[0214] An amino group of L^1 that connects to L^2 may be the N-terminus of an amino acid or may be derived from an amino group of an amino acid side chain, for example a lysine amino acid side chain.

[0215] A carboxyl group of L^1 that connects to L^2 may be the C-terminus of an amino acid or may be derived from a carboxyl group of an amino acid side chain, for example a glutamic acid amino acid side chain.

[0216] A hydroxyl group of L^1 that connects to L^2 may be derived from a hydroxyl group of an amino acid side chain, for example a serine amino acid side chain.

[0217] The term "amino acid side chain" includes those groups found in: (i) naturally occurring amino acids such as alanine, arginine, asparagine, aspartic acid, cysteine, glutamine, glutamic acid, glycine, histidine, isoleucine, leucine, lysine, methionine, phenylalanine, proline, serine, threonine, tryptophan, tyrosine, and valine; (ii) minor amino acids such as ornithine and citrulline; (iii) unnatural amino acids, beta-amino acids, synthetic analogs and derivatives of naturally occurring amino acids; and (iv) all enantiomers, diastereomers, isomerically enriched, isotopically labelled (e.g. ^2H , ^3H , ^{14}C , ^{15}N), protected forms, and racemic mixtures thereof.

[0218] In one embodiment, $-C(=O)O-$ and L^2 together form the group:




where the asterisk indicates the point of attachment to the drug or cytotoxic agent position, the wavy line indicates the point of attachment to the linker L^1 , Y is $-N(H)-$, $-O-$, $-C(=O)N(H)-$ or $-C(=O)O-$, and n is 0 to 3. The phenylene ring is optionally substituted with one, two or three substituents as described herein. In one embodiment, the phenylene group is optionally substituted with halo, NO_2 , R or OR.

[0219] In one embodiment, Y is NH.

[0220] In one embodiment, n is 0 or 1. Preferably, n is 0.

[0221] Where Y is NH and n is 0, the self-immolative linker may be referred to as a p-aminobenzylcarbonyl linker (PABC).


[0222] The self-immolative linker will allow for release of the protected compound when a remote site is activated, proceeding along the lines shown below (for n=0):



where L^* is the activated form of the remaining portion of the linker. These groups have the advantage of separating the site of activation from the compound being protected. As described above, the phenylene group may be optionally substituted.


In one embodiment described herein, the group L^* is a linker L^1 as described herein, which may include a dipeptide group.

[0223] In another embodiment, $-C(=O)O-$ and L^2 together form a group selected from:



where the asterisk, the wavy line, Y, and n are as defined above. Each phenylene ring is optionally substituted with one, two or three substituents as described herein. In one embodiment, the phenylene ring having the Y substituent is optionally substituted and the phenylene ring not having the Y substituent is unsubstituted. In one embodiment, the phenylene ring having the Y substituent is unsubstituted and the phenylene ring not having the Y substituent is optionally substituted.

[0224] In another embodiment, $-C(=O)O-$ and L^2 together form a group selected from:



where the asterisk, the wavy line, Y, and n are as defined above, E is O, S or NR, D is N, CH, or CR, and F is N, CH, or CR.

[0225] In one embodiment, D is N.

[0226] In one embodiment, D is CH.

[0227] In one embodiment, E is O or S.

[0228] In one embodiment, F is CH.

[0229] In a preferred embodiment, the linker is a cathepsin labile linker.

[0230] In one embodiment, L^1 comprises a dipeptide. The dipeptide may be represented as $-NH-X_1-X_2-CO-$, where $-NH-$ and $-CO-$ represent the N- and C-terminals of the amino acid groups X_1 and X_2 respectively. The amino acids in the dipeptide may be any combination of natural amino acids. Where the linker is a cathepsin labile linker, the dipeptide may be the site of action for cathepsin-mediated cleavage.

[0231] Additionally, for those amino acids groups having carboxyl or amino side chain functionality, for example Glu and Lys respectively, CO and NH may represent that side chain functionality.

[0232] In one embodiment, the group $-X_1-X_2-$ in dipeptide, $-NH-X_1-X_2-CO-$, is selected from:

5 $-Phe-Lys-$, $-Val-Ala-$, $-Val-Lys-$, $-Ala-Lys-$, $-Val-Cit-$, $-Phe-Cit-$, $-Leu-Cit-$, $-Ile-Cit-$, $-Phe-Arg-$ and $-Trp-Cit-$ where Cit is citrulline.

[0233] Preferably, the group $-X_1-X_2-$ in dipeptide, $-NH-X_1-X_2-CO-$, is selected from:

10 $-Phe-Lys-$, $-Val-Ala-$, $-Val-Lys-$, $-Ala-Lys-$, and $-Val-Cit-$.

[0234] Most preferably, the group $-X_1-X_2-$ in dipeptide, $-NH-X_1-X_2-CO-$, is $-Phe-Lys-$ or $-Val-Ala-$.

[0235] Other dipeptide combinations may be used, including those described by Dubowchik et al., Bioconjugate Chemistry, 2002, 13,855-869, which is incorporated herein by reference.

15 **[0236]** In one embodiment, the amino acid side chain is derivatised, where appropriate. For example, an amino group or carboxy group of an amino acid side chain may be derivatised.

[0237] In one embodiment, an amino group NH_2 of a side chain amino acid, such as lysine, is a derivatised form selected from the group consisting of NHR and NRR' .

20 **[0238]** In one embodiment, a carboxy group $COOH$ of a side chain amino acid, such as aspartic acid, is a derivatised form selected from the group consisting of $COOR$, $CONH_2$, $CONHR$ and $CONRR'$.

[0239] In one embodiment, the amino acid side chain is chemically protected, where appropriate. The side chain protecting group may be a group as discussed below in relation to the group R^L . Protected amino acid sequences are cleavable by enzymes. For example, it has been established that a dipeptide sequence comprising a Boc side chain-protected Lys residue is cleavable by cathepsin.

25 **[0240]** Protecting groups for the side chains of amino acids are well known in the art and are described in the Novabiochem Catalog. Additional protecting group strategies are set out in Protective Groups in Organic Synthesis, Greene and Wuts.

[0241] Possible side chain protecting groups are shown below for those amino acids having reactive side chain functionality:

30 Arg: Z, Mtr, Tos;
 Asn: Trt, Xan;
 Asp: Bzl, t-Bu;
 Cys: Acn, Bzl, Bzl-OMe, Bzl-Me, Trt;
 Glu: Bzl, t-Bu;
 Gln: Trt, Xan;
 His: Boc, Dnp, Tos, Trt;
 Lys: Boc, Z-Cl, Fmoc, Z, Alloc;
 Ser: Bzl, TBDMS, TBDPS;
 Thr: Bz;
 Trp: Boc;
 Tyr: Bzl, Z, Z-Br.

45 **[0242]** In one embodiment, the side chain protection is selected to be orthogonal to a group provided as, or as part of, a capping group, where present. Thus, the removal of the side chain protecting group does not remove the capping group, or any protecting group functionality that is part of the capping group.


[0243] In other embodiments of the invention, the amino acids selected are those having no reactive side chain functionality. For example, the amino acids may be selected from: Ala, Gly, Ile, Leu, Met, Phe, Pro, and Val.

50 **[0244]** In one embodiment, the dipeptide is used in combination with a self-immolative linker. The self-immolative linker may be connected to $-X_2-$.

[0245] Where a self-immolative linker is present, $-X_2-$ is connected directly to the self-immolative linker. Preferably the group $-X_2-CO-$ is connected to Y, where Y is NH, thereby forming the group $-X_2-CO-NH-$.


55 $-NH-X_1-$ is connected directly to A. A may comprise the functionality $-CO-$ thereby to form an amide link with $-X_1-$.

[0246] In one embodiment, L^1 and L^2 together with $-OC(=O)-$ comprise the group $NH-X_1-X_2-CO-PABC-$. The PABC group is connected directly to the cytotoxic agent. Preferably, the self-immolative linker and the dipeptide together form the group $-NH-Phe-Lys-CO-NH-PABC-$, which is illustrated below:




where the asterisk indicates the point of attachment to the selected cytotoxic moiety, and the wavy line indicates the point of attachment to the remaining portion of the linker L¹ or the point of attachment to A. Preferably, the wavy line indicates the point of attachment to A. The side chain of the Lys amino acid may be protected, for example, with Boc, Fmoc, or Alloc, as described above.

[0247] Alternatively, the self-immolative linker and the dipeptide together form the group -NH-Val-Ala-CO-NH-PABC-, which is illustrated below:



where the asterisk and the wavy line are as defined above.

[0248] Alternatively, the self-immolative linker and the dipeptide together form the group -NH-Val-Cit-CO-NH-PABC-, which is illustrated below:



where the asterisk and the wavy line are as defined above.

[0249] In some embodiments of the present invention, it may be preferred that if the drug moiety contains an unprotected imine bond, e.g. if moiety B is present, then the linker does not contain a free amino (H₂N-) group. Thus if the linker has the structure -A-L¹-L²- then this would preferably not contain a free amino group. This preference is particularly relevant when the linker contains a dipeptide, for example as L¹; in this embodiment, it would be preferred that one of the two amino acids is not selected from lysine.

[0250] Without wishing to be bound by theory, the combination of an unprotected imine bond in the drug moiety and a free amino group in the linker can cause dimerisation of the drug-linker moiety which may interfere with the conjugation of such a drug-linker moiety to an antibody. The cross-reaction of these groups may be accelerated in the case the free amino group is present as an ammonium ion (H₃N⁺-), such as when a strong acid (e.g. TFA) has been used to deprotect the free amino group.

[0251] In one embodiment, A is a covalent bond. Thus, L¹ and the cell binding agent are directly connected. For example, where L¹ comprises a contiguous amino acid sequence, the N-terminus of the sequence may connect directly

to the cell binding agent.

[0252] Thus, where A is a covalent bond, the connection between the cell binding agent and L¹ may be selected from:

-C(=O)NH-, -C(=O)O-, -NHC(=O)-, -OC(=O)-, -OC(=O)O-, -NHC(=O)O-, -OC(=O)NH-, -NHC(=O)NH-,
-C(=O)NHC(=O)-, -S-, -S-S-, -CH₂C(=O)-, and =N-NH-.

[0253] An amino group of L¹ that connects to the SEZ6 modulator may be the N-terminus of an amino acid or may be derived from an amino group of an amino acid side chain, for example a lysine amino acid side chain.


[0254] A carboxyl group of L¹ that connects to the modulator may be the C-terminus of an amino acid or may be derived from a carboxyl group of an amino acid side chain, for example a glutamic acid amino acid side chain.

[0255] A hydroxyl group of L¹ that connects to the cell binding agent may be derived from a hydroxyl group of an amino acid side chain, for example a serine amino acid side chain.

[0256] A thiol group of L¹ that connects to a modulator agent may be derived from a thiol group of an amino acid side chain, for example a serine amino acid side chain.

[0257] The comments above in relation to the amino, carboxyl, hydroxyl and thiol groups of L¹ also apply to the cell binding agent.


[0258] In one embodiment, L² together with -OC(=O)- represents:



where the asterisk indicates the point of attachment to the N10 position, the wavy line indicates the point of attachment to L¹, n is 0 to 3, Y is a covalent bond or a functional group, and E is an activatable group, for example by enzymatic action or light, thereby to generate a self-immolative unit. The phenylene ring is optionally further substituted with one, two or three substituents as described herein. In one embodiment, the phenylene group is optionally further substituted with halo, NO₂, R or OR. Preferably n is 0 or 1, most preferably 0.

[0259] E is selected such that the group is susceptible to activation, e.g. by light or by the action of an enzyme. E may be -NO₂ or glucuronic acid. The former may be susceptible to the action of a nitroreductase, the latter to the action of a β -glucuronidase.

[0260] In this embodiment, the self-immolative linker will allow for release of the protected compound when E is activated, proceeding along the lines shown below (for n=0):



where the asterisk indicates the point of attachment to the N10 position, E* is the activated form of E, and Y is as described above. These groups have the advantage of separating the site of activation from the compound being protected. As described above, the phenylene group may be optionally further substituted.

[0261] The group Y may be a covalent bond to L¹.

[0262] The group Y may be a functional group selected from:

-C(=O)-, -NH-, -O-, -C(=O)NH-, -C(=O)O-, -NHC(=O)-, -OC(=O)-, -OC(=O)O-, -NHC(=O)O-, -OC(=O)NH-,
-NHC(=O)NH-, -NHC(=O)NH-, -C(=O)NHC(=O)-, and -S-.


[0263] Where L¹ is a dipeptide, it is preferred that Y is -NH- or -C(=O)-, thereby to form an amide bond between L¹ and Y. In this embodiment, the dipeptide sequence need not be a substrate for an enzymatic activity.

[0264] In another embodiment, A is a spacer group. Thus, L¹ and the cell binding agent are indirectly connected.

[0265] L¹ and A may be connected by a bond selected from:

-C(=O)NH-, -C(=O)O-, -NHC(=O)-, -OC(=O)-, -OC(=O)O-, -NHC(=O)O-, -OC(=O)NH-, and -NHC(=O)NH-.


[0266] Preferably, the linker contains an electrophilic functional group for reaction with a nucleophilic functional group on the modulator. Nucleophilic groups on antibodies include, but are not limited to: (i) N-terminal amine groups, (ii) side chain amine groups, e.g. lysine, (iii) side chain thiol groups, e.g. cysteine, and (iv) sugar hydroxyl or amino groups where the antibody is glycosylated. Amine, thiol, and hydroxyl groups are nucleophilic and capable of reacting to form covalent bonds with electrophilic groups on linker moieties and linker reagents including: (i) maleimide groups (ii) activated disulfides, (iii) active esters such as NHS (N-hydroxysuccinimide) esters, HOBt (N-hydroxybenzotriazole) esters, haloformates, and acid halides; (iv) alkyl and benzyl halides such as haloacetamides; and (v) aldehydes, ketones, carboxyl, and, some of which are exemplified as follows:



[0267] Certain antibodies have reducible interchain disulfides, i.e. cysteine bridges. Antibodies may be made reactive for conjugation with linker reagents by treatment with a reducing agent such as DTT (dithiothreitol). Each cysteine bridge will thus form, theoretically, two reactive thiol nucleophiles. Additional nucleophilic groups can be introduced into antibodies through the reaction of lysines with 2-iminothiolane (Traut's reagent) resulting in conversion of an amine into a thiol. Reactive thiol groups may be introduced into the antibody (or fragment thereof) by introducing one, two, three, four, or more cysteine residues (e.g., preparing mutant antibodies comprising one or more non-native cysteine amino acid residues). US 7521541 teaches engineering antibodies by introduction of reactive cysteine amino acids.


[0268] In some embodiments, a linker has a reactive nucleophilic group which is reactive with an electrophilic group present on an antibody. Useful electrophilic groups on an antibody include, but are not limited to, aldehyde and ketone carbonyl groups. The heteroatom of a nucleophilic group of a Linker can react with an electrophilic group on an antibody and form a covalent bond to an antibody unit. Useful nucleophilic groups on a linker include, but are not limited to, hydrazide, oxime, amino, hydroxyl, hydrazine, thiosemicarbazone, hydrazine carboxylate, and arylhydrazide. The electrophilic group on an antibody provides a convenient site for attachment to a Linker.

[0269] In one embodiment, the group A is:




where the asterisk indicates the point of attachment to L¹, the wavy line indicates the point of attachment to the cell binding agent, and n is 0 to 6. In one embodiment, n is 5.

[0270] In one embodiment, the group A is:




where the asterisk indicates the point of attachment to L^1 , the wavy line indicates the point of attachment to the cell binding agent, and n is 0 to 6. In one embodiment, n is 5.

[0271] In one embodiment, the group A is:



where the asterisk indicates the point of attachment to L^1 , the wavy line indicates the point of attachment to the cell binding agent, n is 0 or 1, and m is 0 to 30. In a preferred embodiment, n is 1 and m is 0 to 10, 1 to 8, preferably 4 to 8, and most preferably 4 or 8. In another embodiment, m is 10 to 30, and preferably 20 to 30. Alternatively, m is 0 to 50. In this embodiment, m is preferably 10-40 and n is 1.


[0272] In one embodiment, the group A is:



where the asterisk indicates the point of attachment to L^1 , the wavy line indicates the point of attachment to the cell binding agent, n is 0 or 1, and m is 0 to 30. In a preferred embodiment, n is 1 and m is 0 to 10, 1 to 8, preferably 4 to 8, and most preferably 4 or 8. In another embodiment, m is 10 to 30, and preferably 20 to 30. Alternatively, m is 0 to 50. In this embodiment, m is preferably 10-40 and n is 1.


[0273] In one embodiment, the connection between the cell binding agent and A is through a thiol residue of the cell binding agent and a maleimide group of A.

[0274] In one embodiment, the connection between the cell binding agent and A is:




where the asterisk indicates the point of attachment to the remaining portion of A and the wavy line indicates the point of attachment to the remaining portion of the cell binding agent. In this embodiment, the S atom is typically derived from the modulator.

[0275] In each of the embodiments above, an alternative functionality may be used in place of the maleimide-derived group shown below:



where the wavy line indicates the point of attachment to the cell binding agent as before, and the asterisk indicates the bond to the remaining portion of the A group.

[0276] In one embodiment, the maleimide-derived group is replaced with the group:




where the wavy line indicates point of attachment to the cell binding agent, and the asterisk indicates the bond to the remaining portion of the A group.

[0277] In one embodiment, the maleimide-derived group is replaced with a group, which optionally together with the cell binding agent, is selected from:

$-C(=O)NH-$, $-C(=O)O-$, $-NHC(=O)-$, $-OC(=O)-$, $-OC(=O)O-$, $-NHC(=O)O-$, $-OC(=O)NH-$, $-NHC(=O)NH-$, $-NHC(=O)NH-$, $-C(=O)NHC(=O)-$, $-S-$, $-S-S-$, $-CH_2C(=O)-$, $-C(=O)CH_2-$, $=N-NH-$ and $-NH-N=$.

[0278] In one embodiment, the maleimide-derived group is replaced with a group, which optionally together with the cell binding agent, is selected from:



where the wavy line indicates either the point of attachment to the cell binding agent or the bond to the remaining portion of the A group, and the asterisk indicates the other of the point of attachment to the cell binding agent or the bond to the remaining portion of the A group.

[0279] Other groups suitable for connecting L¹ to the selected modulator are described in WO 2005/082023.

[0280] In another preferred embodiment the modulators of the instant invention may be associated with biocompatible polymers comprising drug linker units. In this respect one such type of compatible polymer comprises Fleximer® polymers (Mersana Therapeutics). Such polymers are reportedly biodegradable, well tolerated and have been clinically validated. Moreover, such polymers are compatible with a number of customizable linker technologies and chemistries allowing for control of pharmacokinetics, localization of drug release and improved biodistribution.

[0281] The selected modulators can also be directly conjugated radioisotopes or may comprise macrocyclic chelators useful for conjugating radiometal ions (as described herein). In certain embodiments, the macrocyclic chelator is 1,4,7,10-tetraazacyclododecane-N,N',N'',N'''-tetraacetic acid (DOTA) which can be attached to the antibody via a linker molecule. Such linker molecules are commonly known in the art and described in Denardo et al., 1998, Clin Cancer Res. 4:2483; Peterson et al., 1999, Bioconjug. Chem. 10:553; and Zimmerman et al., 1999, Nucl. Med. Biol. 26:943.

[0282] More generally, techniques for conjugating therapeutic moieties or cytotoxic agents to modulators are well known. As discussed above moieties can be conjugated to modulators by any art-recognized method, including, but not limited to aldehyde/Schiff linkage, sulphhydryl linkage, acid-labile linkage, cis-aconityl linkage, hydrazone linkage, enzymatically degradable linkage (see generally Garnett, 2002, Adv Drug Deliv Rev 53:171). Also see, e.g., Amon et al., "Monoclonal Antibodies For Immunotargeting Of Drugs In Cancer Therapy", in Monoclonal Antibodies And Cancer Therapy, Reisfeld et al. (eds.), pp. 243-56 (Alan R. Liss, Inc. 1985); Hellstrom et al., "Antibodies For Drug Delivery", in Controlled Drug Delivery (2nd Ed.), Robinson et al. (eds.), pp. 623-53 (Marcel Dekker, Inc. 1987); Thorpe, "Antibody Carriers Of Cytotoxic Agents In Cancer Therapy: A Review", in Monoclonal Antibodies '84: Biological And Clinical Applications, Pinchera et al. (eds.), pp. 475-506 (1985); "Analysis, Results, And Future Prospective Of The Therapeutic Use Of Radiolabeled Antibody In Cancer Therapy", in Monoclonal Antibodies For Cancer Detection And Therapy, Baldwin et al. (eds.), pp. 303-16 (Academic Press 1985), and Thorpe et al., 1982, Immunol. Rev. 62:119. In preferred embodiments a SEZ6 modulator that is conjugated to a therapeutic moiety or cytotoxic agent may be internalized by a cell upon binding to a SEZ6 molecule associated with the cell surface thereby delivering the therapeutic payload.

C. Biocompatible Modifiers

[0283] In selected embodiments the modulators of the invention may be conjugated or otherwise associated with biocompatible modifiers that may be used to adjust, alter, improve or moderate modulator characteristics as desired. For example, antibodies or fusion constructs with increased *in vivo* half-lives can be generated by attaching relatively high molecular weight polymer molecules such as commercially available polyethylene glycol (PEG) or similar biocompatible polymers. Those skilled in the art will appreciate that PEG may be obtained in many different molecular weight and molecular configurations that can be selected to impart specific properties to the antibody (e.g. the half-life may be tailored). PEG can be attached to modulators or antibody fragments or derivatives with or without a multifunctional linker either through site-specific conjugation of the PEG to the N- or C-terminus of said antibodies or antibody fragments or via epsilon-amino groups present on lysine residues. Linear or branched polymer derivatization that results in minimal loss of biological activity may be used. The degree of conjugation can be closely monitored by SDS-PAGE and mass spectrometry to ensure optimal conjugation of PEG molecules to antibody molecules. Unreacted PEG can be separated from antibody-PEG conjugates by, e.g., size exclusion or ion-exchange chromatography. In a similar manner, the disclosed modulators can be conjugated to albumin in order to make the antibody or antibody fragment more stable *in vivo* or have a longer half life *in vivo*. The techniques are well known in the art, see e.g., International Publication Nos. WO 93/15199, WO 93/15200, and WO 01/77137; and European Patent No. 0 413, 622. Other biocompatible conjugates are evident to those of ordinary skill and may readily be identified in accordance with the teachings herein.

D. Diagnostic or Detection Agents

[0284] In other preferred embodiments, modulators of the present invention, or fragments or derivatives thereof, are conjugated to a diagnostic or detectable agent, marker or reporter which may be, for example, a biological molecule (e.g., a peptide or nucleotide), a small molecule, fluorophore, or radioisotope. Labeled modulators can be useful for monitoring the development or progression of a hyperproliferative disorder or as part of a clinical testing procedure to determine the efficacy of a particular therapy including the disclosed modulators (i.e. theragnostics) or to determine a future course of treatment. Such markers or reporters may also be useful in purifying the selected modulator, modulator analytics (e.g., epitope binding or antibody binning), separating or isolating TIC or in preclinical procedures or toxicology studies.

[0285] Such diagnosis analysis and/or detection can be accomplished by coupling the modulator to detectable substances including, but not limited to, various enzymes comprising for example horseradish peroxidase, alkaline phosphatase, beta-galactosidase, or acetylcholinesterase; prosthetic groups, such as but not limited to streptavidin/biotin and avidin/biotin; fluorescent materials, such as but not limited to, umbelliferone, fluorescein, fluorescein isothiocyanate, rhodamine, dichlorotriazinylamine fluorescein, dansyl chloride or phycoerythrin; luminescent materials, such as but not limited to, luminol; bioluminescent materials, such as but not limited to, luciferase, luciferin, and aequorin; radioactive materials, such as but not limited to iodine (¹³¹I, ¹²⁵I, ¹²³I, ¹²¹I), carbon (¹⁴C), sulfur (³⁵S), tritium (³H), indium (¹¹⁵In, ¹¹³In, ¹¹²In, ¹¹¹In), and technetium (⁹⁹Tc), thallium (²⁰¹Tl), gallium (⁶⁸Ga, ⁶⁷Ga), palladium (¹⁰³Pd), molybdenum (⁹⁹Mo), xenon (¹³³Xe), fluorine (¹⁸F), ¹⁵³Sm, ¹⁷⁷Lu, ¹⁵⁹Gd, ¹⁴⁹Pm, ¹⁴⁰La, ¹⁷⁵Yb, ¹⁶⁶Ho, ⁹⁰Y, ⁴⁷Sc, ¹⁸⁶Re, ¹⁸⁸Re, ¹⁴²Pr, ¹⁰⁵Rh, ⁹⁷Ru, ⁶⁸Ge, ⁵⁷Co, ⁶⁵Zn, ⁸⁵Sr, ¹⁴⁹Pm, ¹⁴⁰La, ¹⁷⁵Yb, ¹⁶⁶Ho, ⁹⁰Y, ⁴⁷Sc, ¹⁸⁶Re, ¹⁸⁸Re, ¹⁴²Pr, ¹⁰⁵Rh, ⁹⁷Ru, ⁶⁸Ge, ⁵⁷Co, ⁶⁵Zn, ⁸⁵Sr, ³²P, ¹⁵³Gd, ¹⁶⁹Yb, ⁵¹Cr, ⁵⁴Mn, ⁷⁵Se, ¹¹³Sn, and ¹¹⁷Tm; positron emitting metals using various positron emission tomographies, norradioactive paramagnetic metal ions, and molecules that are radiolabeled or conjugated to specific radioisotopes. In such embodiments appropriate detection methodology is well known in the art and readily available from numerous commercial sources.

[0286] As indicated above, in other embodiments the modulators or fragments thereof can be fused or conjugated to marker sequences or compounds, such as a peptide or fluorophore to facilitate purification or diagnostic or analytic procedures such as immunohistochemistry, bio-layer interferometry, surface plasmon resonance, flow cytometry, competitive ELISA, FACs, etc. In preferred embodiments, the marker comprises a his-tag such as that provided by the pQE vector (Qiagen), among others, many of which are commercially available. Other peptide tags useful for purification include, but are not limited to, the hemagglutinin "HA" tag, which corresponds to an epitope derived from the influenza hemagglutinin protein (Wilson et al., 1984, Cell 37:767) and the "flag" tag (U.S.P.N. 4,703,004).

E. Therapeutic Moieties

[0287] As previously alluded to the modulators or fragments or derivatives thereof may also be conjugated, linked or fused to or otherwise associated with a "therapeutic moiety" or "drug" such as an anti-proliferative or anti-cancer agent including, but not limited to, cytotoxic agents, cytostatic agents, anti-angiogenic agents, debulking agents, chemotherapeutic agents, radiotherapy and radiotherapeutic agents, targeted anti-cancer agents, BRMs, therapeutic antibodies, cancer vaccines, cytokines, hormone therapies, radiation therapy and anti-metastatic agents and immunotherapeutic


agents.

[0288] Preferred exemplary anti-cancer agents include cytochalasin B, gramicidin D, ethidium bromide, emetine, mitomycin, etoposide, teniposide, vincristine, vinblastine, colchicin, doxorubicin, daunorubicin, dihydroxy anthracin, maytansinoids such as DM-1 and DM-4 (Immunogen, Inc.), dione, mitoxantrone, mithramycin, actinomycin D, 1-dehydrotestosterone, glucocorticoids, procaine, tetracaine, lidocaine, propranolol, puromycin, epirubicin, and cyclophosphamide and analogs or homologs thereof. Additional compatible cytotoxins comprise dolastatins and auristatins, including monomethyl auristatin E (MMAE) and monomethyl auristatin F (MMAF) (Seattle Genetics, Inc.), amanitins such as alpha-amanitin, beta-amanitin, gamma-amanitin or epsilon-amanitin (Heidelberg Pharma AG), DNA minor groove binding agents such as duocarmycin derivatives (Syntarga, B.V.) and modified pyrrolobenzodiazepine dimers (Spirogen, Ltd.), splicing inhibitors such as meayamycin analogs or derivatives (e.g., FR901464 as set forth in U.S.P.N. 7,825,267), tubular binding agents such as epothilone analogs and paclitaxel and DNA damaging agents such as calicheamicins and esperamicins. Furthermore, in certain embodiments the SEZ6 modulators of the instant invention may be associated with anti-CD3 binding molecules to recruit cytotoxic T-cells and have them target the tumor initiating cells (BiTE technology; see e.g., Fuhrmann, S. et. al. Annual Meeting of AACR Abstract No. 5625 (2010) which is incorporated herein by reference).


[0289] Still additional compatible anti-cancer agents include, but are not limited to, antimetabolites (e.g., methotrexate, 6-mercaptopurine, 6-thioguanine, cytarabine, 5-fluorouracil decarbazine), alkylating agents (e.g., mechlorethamine, thioepa, chlorambucil, melphalan, carmustine (BCNU) and lomustine (CCNU), busulfan, dibromomannitol, streptozotocin, and cisdichlorodiamine platinum (II) (DDP) cisplatin), anthracyclines (e.g., daunorubicin (formerly daunomycin) and doxorubicin), antibiotics (e.g., dactinomycin (formerly actinomycin), bleomycin, and anthramycin (AMC)), and anti-mitotic agents (e.g., vincristine and vinblastine). A more extensive list of therapeutic moieties can be found in PCT publication WO 03/075957 and U.S.P.N. 2009/0155255 each of which is incorporated herein by reference.

[0290] As indicated above selected embodiments of the instant invention are directed to conjugated SEZ6 modulators such as anti-SEZ6 antibody drug conjugates that comprise pyrrolobenzodiazepine (PBD) as a cytotoxic agent. It will be appreciated that PBDs are alkylating agents that exert anti-tumor activity by covalently binding to DNA in the minor groove and inhibiting nucleic acid synthesis. In this respect PBDs have been shown to have potent antitumor properties while exhibiting minimal bone marrow depression. PBDs compatible with the present invention may be linked to the SEZ6 modulator using any one of several types of linker (e.g., a peptidyl linker comprising a maleimido moiety with a free sulfhydryl) and, in certain embodiments are dimeric in form (i.e., PBD dimers). Compatible PBDs (and optional linkers) that may be conjugated to the disclosed modulators are described, for example, in U.S.P.N.s 6,362,331, 7,049,311, 7,189,710, 7,429,658, 7,407,951, 7,741,319, 7,557,099, 8,034,808, 8,163,736 U.S.P.N. 2011/0256157 and PCT filings WO2011/130613, WO2011/128650 and WO2011/130616 each of which is incorporated herein by reference. Accordingly, in particularly preferred embodiments the modulator will comprise an anti SEZ6 antibody conjugated or associated with one or more PBD dimers (i.e., a SEZ6-PBD ADC).

[0291] In particularly preferred embodiments compatible PBDs that may be conjugated to the disclosed modulators are described in U.S.P.N. 2011/0256157. In this disclosure, PBD dimers, i.e. those comprising two PBD moieties may be preferred. Thus, preferred conjugates of the present invention are those having the formula (AB) or (AC):



AB



AC

wherein:

the dotted lines indicate the optional presence of a double bond between C1 and C2 or C2 and C3;

R² is independently selected from H, OH, =O, =CH₂, CN, R, OR, =CH-R^D, =C(R^D)₂, O-SO₂-R, CO₂R and COR, and optionally further selected from halo or dihalo;

where R^D is independently selected from R, CO₂R, COR, CHO, CO₂H, and halo;

R⁶ and R⁹ are independently selected from H, R, OH, OR, SH, SR, NH₂, NHR, NRR', NO₂, Me₃Sn and halo;

R⁷ is independently selected from H, R, OH, OR, SH, SR, NH₂, NHR, NRR', NO₂, Me₃Sn and halo;

R¹⁰ is a linker connected to a modulator or fragment or derivative thereof, as described above;

Q is independently selected from O, S and NH;

R¹¹ is either H, or R or, where Q is O, SO₃M, where M is a metal cation;

R and R' are each independently selected from optionally substituted C₁₋₁₂ alkyl, C₃₋₂₀ heterocyclyl and C₅₋₂₀ aryl groups, and optionally in relation to the group NRR', R and R' together with the nitrogen atom to which they are


attached form an optionally substituted 4-, 5-, 6- or 7-membered heterocyclic ring; and

wherein R², R⁶, R⁷, R⁹, X, Q and R¹¹ are as defined according to R², R⁶, R⁷, R⁹, X, Q and R¹¹ respectively, and R^C is a capping group.

Double Bond


[0292] In one embodiment, there is no double bond present between C 1 and C2, and C2 and C3.

[0293] In one embodiment, the dotted lines indicate the optional presence of a double bond between C2 and C3, as shown below:



[0294] In one embodiment, a double bond is present between C2 and C3 when R² is C₅₋₂₀ aryl or C₁₋₁₂ alkyl.

[0295] In one embodiment, the dotted lines indicate the optional presence of a double bond between C 1 and C2, as shown below:



[0296] In one embodiment, a double bond is present between C1 and C2 when R² is C₅₋₂₀ aryl or C₁₋₁₂ alkyl.

R²

[0297] In one embodiment, R² is independently selected from H, OH, =O, =CH₂, CN, R, OR, =CH-R^D, =C(R^D)₂, O-SO₂-R, CO₂R and COR, and optionally further selected from halo or dihalo.

[0298] In one embodiment, R² is independently selected from H, OH, =O, =CH₂, CN, R, OR, =CH-R^D, =C(R^D)₂, O-SO₂-R, CO₂R and COR.


[0299] In one embodiment, R² is independently selected from H, =O, =CH₂, R, =CH-R^D, and =C(R^D)₂.

[0300] In one embodiment, R² is independently H.

[0301] In one embodiment, R² is independently =O.

[0302] In one embodiment, R² is independently =CH₂.

[0303] In one embodiment, R² is independently =CH-R^D. Within the PBD compound, the group =CH-R^D may have either configuration shown below:



[0304] In one embodiment, the configuration is configuration (I).

[0305] In one embodiment, R² is independently =C(R^D)₂.

[0306] In one embodiment, R² is independently =CF₂.

[0307] In one embodiment, R² is independently R.

[0308] In one embodiment, R² is independently optionally substituted C₅₋₂₀ aryl.

[0309] In one embodiment, R² is independently optionally substituted C₁₋₁₂ alkyl.

[0310] In one embodiment, R² is independently optionally substituted C₅₋₂₀ aryl.

[0311] In one embodiment, R² is independently optionally substituted C₅₋₇ aryl.

[0312] In one embodiment, R² is independently optionally substituted C₈₋₁₀ aryl.

[0313] In one embodiment, R² is independently optionally substituted phenyl.

[0314] In one embodiment, R² is independently optionally substituted naphthyl.


[0315] In one embodiment, R² is independently optionally substituted pyridyl.

[0316] In one embodiment, R² is independently optionally substituted quinoliny or isoquinoliny.

[0317] In one embodiment, R² bears one to three substituent groups, with 1 and 2 being more preferred, and singly substituted groups being most preferred. The substituents may be any position.

[0318] Where R² is a C₅₋₇ aryl group, a single substituent is preferably on a ring atom that is not adjacent the bond to the remainder of the compound, i.e. it is preferably β or γ to the bond to the remainder of the compound. Therefore, where the C₅₋₇ aryl group is phenyl, the substituent is preferably in the meta- or para- positions, and more preferably is in the para-position.

[0319] In one embodiment, R² is selected from:



where the asterisk indicates the point of attachment.

[0320] Where R² is a C₈₋₁₀ aryl group, for example quinoliny or isoquinoliny, it may bear any number of substituents at any position of the quinoline or isoquinoline rings. In some embodiments, it bears one, two or three substituents, and these may be on either the proximal and distal rings or both (if more than one substituent).

[0321] In one embodiment, where R² is optionally substituted, the substituents are selected from those substituents given in the substituent section below.

[0322] Where R is optionally substituted, the substituents are preferably selected from:

Halo, Hydroxyl, Ether, Formyl, Acyl, Carboxy, Ester, Acyloxy, Amino, Amido, Acylamido, Aminocarbonyloxy, Ureido, Nitro, Cyano and Thioether.

[0323] In one embodiment, where R or R² is optionally substituted, the substituents are selected from the group consisting of R, OR, SR, NRR', NO₂, halo, CO₂R, COR, CONH₂, CONHR, and CONRR'.

[0324] Where R² is C₁₋₁₂ alkyl, the optional substituent may additionally include C₃₋₂₀ heterocyclyl and C₅₋₂₀ aryl groups.

[0325] Where R² is C₃₋₂₀ heterocyclyl, the optional substituent may additionally include C₁₋₁₂ alkyl and C₅₋₂₀ aryl groups.

[0326] Where R² is C₅₋₂₀ aryl groups, the optional substituent may additionally include C₃₋₂₀ heterocyclyl and C₁₋₁₂ alkyl groups.

[0327] It is understood that the term "alkyl" encompasses the sub-classes alkenyl and alkynyl as well as cycloalkyl. Thus, where R² is optionally substituted C₁₋₁₂ alkyl, it is understood that the alkyl group optionally contains one or more carbon-carbon double or triple bonds, which may form part of a conjugated system. In one embodiment, the optionally substituted C₁₋₁₂ alkyl group contains at least one carbon-carbon double or triple bond, and this bond is conjugated with a double bond present between C1 and C2, or C2 and C3. In one embodiment, the C₁₋₁₂ alkyl group is a group selected from saturated C₁₋₁₂ alkyl, C₂₋₁₂ alkenyl, C₂₋₁₂ alkynyl and C₃₋₁₂ cycloalkyl.

[0328] If a substituent on R² is halo, it is preferably F or Cl, more preferably Cl.

[0329] If a substituent on R² is ether, it may in some embodiments be an alkoxy group, for example, a C₁₋₇ alkoxy group (e.g. methoxy, ethoxy) or it may in some embodiments be a C₅₋₇ aryloxy group (e.g. phenoxy, pyridyloxy, furanyloxy).

[0330] If a substituent on R² is C₁₋₇ alkyl, it may preferably be a C₁₋₄ alkyl group (e.g. methyl, ethyl, propyl, butyl).


[0331] If a substituent on R² is C₃₋₇ heterocyclyl, it may in some embodiments be C₆ nitrogen containing heterocyclyl group, e.g. morpholino, thiomorpholino, piperidinyl, piperazinyl. These groups may be bound to the rest of the PBD moiety via the nitrogen atom. These groups may be further substituted, for example, by C₁₋₄ alkyl groups.

[0332] If a substituent on R² is bis-oxy-C₁₋₃ alkylene, this is preferably bis-oxy-methylene or bis-oxy-ethylene.


[0333] Particularly preferred substituents for R² include methoxy, ethoxy, fluoro, chloro, cyano, bis-oxy-methylene, methyl-piperazinyl, morpholino and methyl-thienyl.

[0334] Particularly preferred substituted R² groups include, but are not limited to, 4-methoxyphenyl, 3-methoxyphenyl, 4-ethoxy-phenyl, 3-ethoxy-phenyl, 4-fluoro-phenyl, 4-chlorophenyl, 3,4-bisoxymethylene-phenyl, 4-methylthienyl, 4-cyanophenyl, 4-phenoxyphenyl, quinolin-3-yl and quinolin-6-yl, isoquinolin-3-yl and isoquinolin-6-yl, 2-thienyl, 2-furanyl, methoxynaphthyl, and naphthyl.

[0335] In one embodiment, R² is halo or dihalo. In one embodiment, R² is -F or -F₂, which substituents are illustrated below as (III) and (IV) respectively:



(III)



(IV)

R^D

[0336] In one embodiment, R^D is independently selected from R, CO₂R, COR, CHO, CO₂H, and halo.

[0337] In one embodiment, R^D is independently R.

[0338] In one embodiment, R^D is independently halo.

R⁶

[0339] In one embodiment, R⁶ is independently selected from H, R, OH, OR, SH, SR, NH₂, NHR, NRR', NO₂, Me₃Sn- and Halo.

[0340] In one embodiment, R⁶ is independently selected from H, OH, OR, SH, NH₂, NO₂ and Halo.

[0341] In one embodiment, R⁶ is independently selected from H and Halo.

[0342] In one embodiment, R⁶ is independently H.

[0343] In one embodiment, R⁶ and R⁷ together form a group -O-(CH₂)_p-O-, where p is 1 or 2.

R⁷

[0344] R⁷ is independently selected from H, R, OH, OR, SH, SR, NH₂, NHR, NRR', NO₂, Me₃Sn and halo.

[0345] In one embodiment, R⁷ is independently OR.

[0346] In one embodiment, R⁷ is independently OR^{7A}, where R^{7A} is independently optionally substituted C₁₋₆ alkyl.

[0347] In one embodiment, R^{7A} is independently optionally substituted saturated C₁₋₆ alkyl.

[0348] In one embodiment, R^{7A} is independently optionally substituted C₂₋₄ alkenyl.

[0349] In one embodiment, R^{7A} is independently Me.

[0350] In one embodiment, R^{7A} is independently CH₂Ph.

[0351] In one embodiment, R^{7A} is independently allyl.

[0352] In one embodiment, the compound is a dimer where the R⁷ groups of each monomer form together a dimer bridge having the formula X-R''-X linking the monomers.

R⁸

[0353] In one embodiment, the compound is a dimer where the R⁸ groups of each monomer form together a dimer bridge having the formula X-R''-X linking the monomers.

[0354] In one embodiment, R⁸ is independently OR^{8A}, where R^{8A} is independently optionally substituted C₁₋₄ alkyl.

[0355] In one embodiment, R^{8A} is independently optionally substituted saturated C₁₋₆ alkyl or optionally substituted C₂₋₄ alkenyl.

[0356] In one embodiment, R^{8A} is independently Me.

[0357] In one embodiment, R^{8A} is independently CH₂Ph.

[0358] In one embodiment, R^{8A} is independently allyl.

[0359] In one embodiment, R⁸ and R⁷ together form a group -O-(CH₂)_p-O-, where p is 1 or 2.

[0360] In one embodiment, R⁸ and R⁹ together form a group -O-(CH₂)_p-O-, where p is 1 or 2.

R⁹

[0361] In one embodiment, R⁹ is independently selected from H, R, OH, OR, SH, SR, NH₂, NHR, NRR', NO₂, Me₃Sn and Halo.

[0362] In one embodiment, R⁹ is independently H.

[0363] In one embodiment, R⁹ is independently R or OR.

R and R'

[0364] In one embodiment, R is independently selected from optionally substituted C₁₋₁₂ alkyl, C₃₋₂₀ heterocyclyl and C₅₋₂₀ aryl groups. These groups are each defined in the substituents section below.

[0365] In one embodiment, R is independently optionally substituted C₁₋₁₂ alkyl.

[0366] In one embodiment, R is independently optionally substituted C₃₋₂₀ heterocyclyl.

[0367] In one embodiment, R is independently optionally substituted C₅₋₂₀ aryl.

[0368] In one embodiment, R is independently optionally substituted C₁₋₁₂ alkyl.

[0369] Described above in relation to R² are various embodiments relating to preferred alkyl and aryl groups and the identity and number of optional substituents. The preferences set out for R² as it applies to R are applicable, where appropriate, to all other groups R, for examples where R⁶, R⁷, R⁸ or R⁹ is R.

[0370] The preferences for R apply also to R'.

[0371] In some embodiments of the invention there is provided a compound having a substituent group -NRR'. In one embodiment, R and R' together with the nitrogen atom to which they are attached form an optionally substituted 4-, 5-, 6- or 7-membered heterocyclic ring. The ring may contain a further heteroatom, for example N, O or S.

[0372] In one embodiment, the heterocyclic ring is itself substituted with a group R. Where a further N heteroatom is present, the substituent may be on the N heteroatom.

R''

[0373] R'' is a C₃₋₁₂ alkylene group, which chain may be interrupted by one or more heteroatoms, e.g. O, S, N(H), NMe and/or aromatic rings, e.g. benzene or pyridine, which rings are optionally substituted.

[0374] In one embodiment, R'' is a C₃₋₁₂ alkylene group, which chain may be interrupted by one or more heteroatoms and/or aromatic rings, e.g. benzene or pyridine.

[0375] In one embodiment, the alkylene group is optionally interrupted by one or more heteroatoms selected from O,

S, and NMe and/or aromatic rings, which rings are optionally substituted.

[0376] In one embodiment, the aromatic ring is a C₅₋₂₀ arylene group, where arylene pertains to a divalent moiety obtained by removing two hydrogen atoms from two aromatic ring atoms of an aromatic compound, which moiety has from 5 to 20 ring atoms.

[0377] In one embodiment, R" is a C₃₋₁₂ alkylene group, which chain may be interrupted by one or more heteroatoms, e.g. O, S, N(H), NMe and/or aromatic rings, e.g. benzene or pyridine, which rings are optionally substituted by NH₂.

[0378] In one embodiment, R" is a C₃₋₁₂ alkylene group.

[0379] In one embodiment, R" is selected from a C₃, C₅, C₇, C₉ and a C₁₁ alkylene group.

[0380] In one embodiment, R" is selected from a C₃, C₅ and a C₇ alkylene group.

[0381] In one embodiment, R" is selected from a C₃ and a C₅ alkylene group.

[0382] In one embodiment, R" is a C₃ alkylene group.

[0383] In one embodiment, R" is a C₅ alkylene group.

[0384] The alkylene groups listed above may be optionally interrupted by one or more heteroatoms and/or aromatic rings, e.g. benzene or pyridine, which rings are optionally substituted.

[0385] The alkylene groups listed above may be optionally interrupted by one or more heteroatoms and/or aromatic rings, e.g. benzene or pyridine.

[0386] The alkylene groups listed above may be unsubstituted linear aliphatic alkylene groups.

X

[0387] In one embodiment, X is selected from O, S, or N(H).

[0388] Preferably, X is O.


R¹⁰

[0389] Preferably compatible linkers such as those described above attach a SEZ6 modulator (CBA/Ab/M), to a PBD drug moiety D through covalent bond(s) at the R¹⁰ position (i.e., N10). The linker is a bifunctional or multifunctional moiety which can be used to link one or more drug moiety (D) and a modulator (preferably an antibody) to form antibody-drug conjugates (ADC). The linker (L) may be stable outside a cell, i.e. extracellular, or it may be cleavable by enzymatic activity, hydrolysis, or other metabolic conditions. Antibody-drug conjugates (ADC) can be conveniently prepared using a linker having reactive functionality for binding to the drug moiety and to the antibody. A cysteine thiol, or an amine, e.g. N-terminus or amino acid side chain such as lysine, of the antibody (Ab) can form a bond with a functional group of a linker or spacer reagent, PBD drug moiety (D) or drug-linker reagent (D-L).

[0390] Many functional groups on the linker attached to the N10 position of the PBD moiety may be useful to react with the cell binding agent. For example, ester, thioester, amide, thioamide, carbamate, thiocarbamate, urea, thiourea, ether, thioether, or disulfide linkages may be formed from reaction of the linker-PBD drug intermediates and the cell binding agent.

[0391] In another embodiment, the linker may be substituted with groups that modulate aggregation, solubility or reactivity. For example, a sulfonate substituent may increase water solubility of the reagent and facilitate the coupling reaction of the linker reagent with the antibody or the drug moiety, or facilitate the coupling reaction of Ab-L with D, or D-L with Ab, depending on the synthetic route employed to prepare the ADC.

[0392] In one preferred embodiment, R¹⁰ is a group:



where the asterisk indicates the point of attachment to the N10 position, CBA is a cell binding agent/modulator, L¹ is a linker, A is a connecting group connecting L¹ to the cell binding agent, L² is a covalent bond or together with -OC(=O)- forms a self-immolative linker, and L¹ or L² is a cleavable linker.

[0393] L¹ is preferably the cleavable linker, and may be referred to as a trigger for activation of the linker for cleavage.

[0394] As discussed in the linker section above the nature of L¹ and L², where present, can vary widely. These groups are chosen on the basis of their cleavage characteristics, which may be dictated by the conditions at the site to which the conjugate is delivered. Those linkers that are cleaved by the action of enzymes are preferred, although linkers that are cleavable by changes in pH (e.g. acid or base labile), temperature or upon irradiation (e.g. photolabile) may also be used. Linkers that are cleavable under reducing or oxidizing conditions may also find use in the present invention.

[0395] L¹ may comprise a contiguous sequence of amino acids. The amino acid sequence may be the target substrate for enzymatic cleavage, thereby allowing release of R¹⁰ from the N10 position.

[0396] In one embodiment, L¹ is cleavable by the action of an enzyme. In one embodiment, the enzyme is an esterase or a peptidase.

[0397] In one embodiment, L² is present and together with -C(=O)O- forms a self-immolative linker. In one embodiment, L² is a substrate for enzymatic activity, thereby allowing release of R¹⁰ from the N10 position.

[0398] In one embodiment, where L¹ is cleavable by the action of an enzyme and L² is present, the enzyme cleaves the bond between L¹ and L².

[0399] With regard to attaching the chosen linker to a selected PBD the group R^C is removable from the N10 position of certain PBD moieties to leave an N10-C11 imine bond, a carbinolamine, a substituted carbinolamine, where QR¹¹ is OSO₃M, a bisulfite adduct, a thiocarbinolamine, a substituted thiocarbinolamine, or a substituted carbinolamine.

[0400] In one embodiment, R^C may be a protecting group that is removable to leave an N10-C11 imine bond, a carbinolamine, a substituted carbinolamine, or, where QR¹¹ is OSO₃M, a bisulfite adduct. In one embodiment, R^C is a protecting group that is removable to leave an N10-C11 imine bond.

[0401] The group R^C is intended to be removable under the same conditions as those required for the removal of the group R¹⁰, for example to yield an N10-C11 imine bond, a carbinolamine and so on. The capping group acts as a protecting group for the intended functionality at the N10 position. The capping group is intended not to be reactive towards a cell binding agent. For example, R^C is not the same as R^L.

[0402] Compounds having a capping group may be used as intermediates in the synthesis of dimers having an imine monomer. Alternatively, compounds having a capping group may be used as conjugates, where the capping group is removed at the target location to yield an imine, a carbinolamine, a substituted carbinolamine and so on. Thus, in this embodiment, the capping group may be referred to as a therapeutically removable nitrogen protecting group, as defined in WO 00/12507.

[0403] In one embodiment, the group R^C is removable under the conditions that cleave the linker R^L of the group R¹⁰. Thus, in one embodiment, the capping group is cleavable by the action of an enzyme.

[0404] In an alternative embodiment, the capping group is removable prior to the connection of the linker R^L to the modulator. In this embodiment, the capping group is removable under conditions that do not cleave the linker R^L.

[0405] Where a compound includes a functional group G¹ to form a connection to the cell binding agent, the capping group is removable prior to the addition or unmasking of G¹.

[0406] The capping group may be used as part of a protecting group strategy to ensure that only one of the monomer units in a dimer is connected to a cell binding agent.

[0407] The capping group may be used as a mask for a N10-C11 imine bond. The capping group may be removed at such time as the imine functionality is required in the compound. The capping group is also a mask for a carbinolamine, a substituted carbinolamine, and a bisulfite adduct, as described above.

[0408] In one embodiment, R^C is a carbamate protecting group.

[0409] In one embodiment, the carbamate protecting group is selected from:


Alloc, Fmoc, Boc, Troc, Teoc, Psec, Cbz and PNZ.

[0410] Optionally, the carbamate protecting group is further selected from Moc.

[0411] In one embodiment, R^C is a linker group R^L lacking the functional group for connection to the cell binding agent.

[0412] This application is particularly concerned with those R^C groups which are carbamates.

[0413] In one embodiment, R^C is a group:



where the asterisk indicates the point of attachment to the N10 position, G² is a terminating group, L³ is a covalent bond or a cleavable linker L¹, L² is a covalent bond or together with OC(=O) forms a self-immolative linker.

[0414] Where L³ and L² are both covalent bonds, G² and OC(=O) together form a carbamate protecting group as defined above.

[0415] L¹ is as defined above in relation to R¹⁰.

[0416] L² is as defined above in relation to R¹⁰.

[0417] Various terminating groups are described below, including those based on well known protecting groups.


[0418] In one embodiment L³ is a cleavable linker L¹, and L², together with OC(=O), forms a self-immolative linker. In this embodiment, G² is Ac (acetyl) or Moc, or a carbamate protecting group selected from: Alloc, Fmoc, Boc, Troc, Teoc,

Psec, Cbz and PNZ. Optionally, the carbamate protecting group is further selected from Moc.

[0419] In another embodiment, G^2 is an acyl group $-C(=O)G^3$, where G^3 is selected from alkyl (including cycloalkyl, alkenyl and alkynyl), heteroalkyl, heterocyclyl and aryl (including heteroaryl and carboaryl). These groups may be optionally substituted. The acyl group together with an amino group of L^3 or L^2 , where appropriate, may form an amide bond. The acyl group together with a hydroxy group of L^3 or L^2 , where appropriate, may form an ester bond.


[0420] In one embodiment, G^3 is heteroalkyl. The heteroalkyl group may comprise polyethylene glycol. The heteroalkyl group may have a heteroatom, such as O or N, adjacent to the acyl group, thereby forming a carbamate or carbonate group, where appropriate, with a heteroatom present in the group L^3 or L^2 , where appropriate.

[0421] In one embodiment, G^3 is selected from NH_2 , NHR and NRR' . Preferably, G^3 is NRR' . In one embodiment G^2 is the group:




where the asterisk indicates the point of attachment to L^3 , n is 0 to 6 and G^4 is selected from OH, OR, SH, SR, COOR, $CONH_2$, $CONHR$, $CONRR'$, NH_2 , NHR , NRR' , NO_2 , and halo. The groups OH, SH, NH_2 and NHR are protected. In one embodiment, n is 1 to 6, and preferably n is 5. In one embodiment, G^4 is OR, SR, COOR, $CONH_2$, $CONHR$, $CONRR'$, and NRR' . In one embodiment, G^4 is OR, SR, and NRR' . Preferably G^4 is selected from OR and NRR' , most preferably G^4 is OR. Most preferably G^4 is OMe.

[0422] In one embodiment, the group G^2 is:




where the asterisk indicates the point of attachment to L^3 , and n and G^4 are as defined above.

[0423] In one embodiment, the group G^2 is:




where the asterisk indicates the point of attachment to L^3 , n is 0 or 1, m is 0 to 50, and G^4 is selected from OH, OR, SH, SR, COOR, $CONH_2$, $CONHR$, $CONRR'$, NH_2 , NHR , NRR' , NO_2 , and halo. In a preferred embodiment, n is 1 and m is 0 to 10, 1 to 2, preferably 4 to 8, and most preferably 4 or 8. In another embodiment, n is 1 and m is 10 to 50, preferably 20 to 40. The groups OH, SH, NH_2 and NHR are protected. In one embodiment, G^4 is OR, SR, COOR, $CONH_2$, $CONHR$, $CONRR'$, and NRR' . In one embodiment, G^4 is OR, SR, and NRR' . Preferably G^4 is selected from OR and NRR' , most preferably G^4 is OR. Preferably G^4 is OMe.

[0424] In one embodiment, the group G^2 is:




where the asterisk indicates the point of attachment to L^3 and n , m and G^4 are as defined above.

[0425] In one embodiment, the group G^2 is:



where n is 1-20, m is 0-6, and G⁴ is selected from OH, OR, SH, SR, COOR, CONH₂, CONHR, CONRR', NH₂, NHR, NRR', NO₂, and halo. In one embodiment, n is 1-10. In another embodiment, n is 10 to 50, preferably 20 to 40. In one embodiment, n is 1. In one embodiment, m is 1. The groups OH, SH, NH₂ and NHR are protected. In one embodiment, G⁴ is OR, SR, COOR, CONH₂, CONHR, CONRR', and NRR'. In one embodiment, G⁴ is OR, SR, and NRR'. Preferably G⁴ is selected from OR and NRR', most preferably G⁴ is OR. Preferably G⁴ is OMe.

[0426] In one embodiment, the group G^2 is:



where the asterisk indicates the point of attachment to L^3 , and n , m and G^4 are as defined above.

[0427] In each of the embodiments above G⁴ may be OH, SH, NH₂, and NHR. These groups are preferably protected.

[0428] In one embodiment, OH is protected with Bzl, TBDMS, or TBDPS.


[0429] In one embodiment, SH is protected with Acn, Bzl, Bzl-OMe, Bzl-Me, or Trt.

[0430] In one embodiment, NH₂ or NHR are protected with Boc, Moc, Z-Cl, Fmoc, Z, or Alloc.

[0431] In one embodiment, the group G^2 is present in combination with a group L^3 , which group is a dipeptide.


[0432] The capping group is not intended for connection to the modulator. Thus, the other monomer present in the dimer serves as the point of connection to the modulator via a linker. Accordingly, it is preferred that the functionality present in the capping group is not available for reaction with a modulator. Thus, reactive functional groups such as OH, SH, NH₂, COOH are preferably avoided. However, such functionality may be present in the capping group if protected, as described above.

[0433] Thus, in accordance with the teachings herein one embodiment of the invention comprises a conjugate comprising a compound:



wherein CBA is a cell binding agent/modulator, and n is 0 or 1. L¹ is as previously defined, and R^E and R^{E'} are each independently selected from H or R^D.

[0434] In another embodiment, the conjugate comprises a compound:



wherein CBA is a cell binding agent/modulator, L^1 is as previously defined, Ar^1 and Ar^2 are each independently optionally substituted C_{5-20} aryl, and n is 0 or 1.

[0435] Those of skill in the art will appreciate that other symmetric and asymmetric PBD dimers and linkers are compatible with the instant invention and could be selected without undue experimentation based on the teachings herein and the prior art.

[0436] Another aspect of the invention includes ADCs comprising radioisotopes. Exemplary radioisotopes that may be compatible with such embodiments include, but are not limited to, iodine (^{131}I , ^{125}I , ^{123}I , ^{121}I), carbon (^{14}C), copper (^{62}Cu , ^{64}Cu , ^{67}Cu), sulfur (^{35}S), tritium (^3H), indium (^{115}In , ^{113}In , ^{112}In , ^{111}In), bismuth (^{212}Bi , ^{213}Bi), technetium (^{99}Tc), thallium (^{201}Tl), gallium (^{68}Ga , ^{67}Ga), palladium (^{103}Pd), molybdenum (^{99}Mo), xenon (^{133}Xe), fluorine (^{18}F), ^{153}Sm , ^{177}Lu , ^{159}Gd , ^{149}Pm , ^{140}La , ^{175}Yb , ^{166}Ho , ^{90}Y , ^{47}Sc , ^{186}Re , ^{188}Re , ^{142}Pr , ^{105}Rh , ^{97}Ru , ^{68}Ge , ^{57}Co , ^{65}Zn , ^{85}Sr , ^{32}P , ^{153}Gd , ^{169}Yb , ^{51}Cr , ^{54}Mn , ^{75}Se , ^{113}Sn , ^{117}Sn , ^{225}Ac , ^{76}Br , and ^{211}At . Other radionuclides are also available as diagnostic and therapeutic agents, especially those in the energy range of 60 to 4,000 keV. Depending on the condition to be treated and the desired therapeutic profile, those skilled in the art may readily select the appropriate radioisotope for use with the disclosed modulators.

[0437] SEZ6 modulators of the present invention may also be conjugated to a therapeutic moiety or drug that modifies a given biological response (e.g., biological response modifiers or BRMs). That is, therapeutic agents or moieties compatible with the instant invention are not to be construed as limited to classical chemical therapeutic agents. For example, in particularly preferred embodiments the drug moiety may be a protein or polypeptide or fragment thereof possessing a desired biological activity. Such proteins may include, for example, a toxin such as abrin, ricin A, Onconase (or another cytotoxic RNase), pseudomonas exotoxin, cholera toxin, or diphtheria toxin; a protein such as tumor necrosis factor, α -interferon, β -interferon, nerve growth factor, platelet derived growth factor, tissue plasminogen activator, an apoptotic agent, e.g., TNF- α , TNF- β , AIM I (see, International Publication No. WO 97/33899), AIM II (see, International Publication No. WO 97/34911), Fas Ligand (Takahashi et al., 1994, J. Immunol., 6:1567), and VEGF (see, International Publication No. WO 99/23105), a thrombotic agent or an anti-angiogenic agent, e.g., angiostatin or endostatin; or, a biological response modifier such as, for example, a lymphokine (e.g., interleukin-1 ("IL-1"), interleukin-2 ("IL-2"), interleukin-6 ("IL-6"), granulocyte macrophage colony stimulating factor ("GM-CSF"), and granulocyte colony stimulating factor ("G-CSF")), or a growth factor (e.g., growth hormone ("GH")). As set forth above, methods for fusing or conjugating modulators to polypeptide moieties are known in the art. In addition to the previously disclosed subject references see, e.g., U.S.P.Ns. 5,336,603; 5,622,929; 5,359,046; 5,349,053; 5,447,851, and 5,112,946; EP 307,434; EP 367,166; PCT Publications WO 96/04388 and WO 91/06570; Ashkenazi et al., 1991, PNAS USA 88:10535; Zheng et al., 1995, J Immunol 154:5590; and Vil et al., 1992, PNAS USA 89:11337 each of which is incorporated herein by reference. Moreover, as set forth above the association of a modulator with such moieties does not necessarily need to be direct, but may occur through linker sequences. As previously alluded to, such linker molecules are commonly known in the art and described in Denardo et al., 1998, Clin Cancer Res 4:2483; Peterson et al., 1999, Bioconjug Chem 10:553; Zimmerman et al., 1999, Nucl Med Biol 26:943; Garnett, 2002, Adv Drug Deliv Rev 53:171 each of which is incorporated herein.

IX. Diagnostics and Screening

A. Diagnostics

[0438] In yet other embodiments, the invention provides *in vitro* or *in vivo* methods for detecting, diagnosing or mon-

itoring proliferative disorders and methods of screening cells from a patient to identify tumorigenic cells including CSCs. Such methods include identifying an individual having cancer for treatment or monitoring progression of a cancer comprising contacting the patient or a sample obtained from a patient (i.e. either *in vivo* or *in vitro*) with a modulator as described herein and detecting presence or absence, or level of association, of the modulator to bound or free target molecules in the sample. In particularly preferred embodiments the modulator will comprise a detectable label or reporter molecule as described herein.

[0439] In some embodiments, the association of the modulator, such as an antibody, with particular cells in the sample likely denotes that the sample may contain CSCs, thereby indicating that the individual having cancer may be effectively treated with a modulator as described herein. The methods may further comprise a step of comparing the level of binding to a control. Conversely, when the modulator is a Fc-construct, the binding properties may be exploited and monitored (directly or indirectly, *in vivo* or *in vitro*) when in contact with the sample to provide the desired information.

[0440] Exemplary compatible assay methods include radioimmunoassays, enzyme immunoassays, competitive-binding assays, fluorescent immunoassay, immunoblot assays, Western Blot analysis, flow cytometry assays, and ELISA assays. Compatible *in vivo* theragnostics or diagnostics may comprise art-recognized imaging or monitoring techniques such as magnetic resonance imaging, computerized tomography (e.g. CAT scan), positron tomography (e.g., PET scan) radiography, ultrasound, etc., as would be known by those skilled in the art.

[0441] In another embodiment, the invention provides a method of analyzing cancer progression and/or pathogenesis *in vivo*. In another embodiment, analysis of cancer progression and/or pathogenesis *in vivo* comprises determining the extent of tumor progression. In another embodiment, analysis comprises the identification of the tumor. In another embodiment, analysis of tumor progression is performed on the primary tumor. In another embodiment, analysis is performed over time depending on the type of cancer as known to one skilled in the art. In another embodiment, further analysis of secondary tumors originating from metastasizing cells of the primary tumor is analyzed *in-vivo*. In another embodiment, the size and shape of secondary tumors are analyzed. In some embodiments, further *ex vivo* analysis is performed.

[0442] In another embodiment, the invention provides a method of analyzing cancer progression and/or pathogenesis *in vivo* including determining cell metastasis or detecting and quantifying the level of circulating tumor cells. In yet another embodiment, analysis of cell metastasis comprises determination of progressive growth of cells at a site that is discontinuous from the primary tumor. In another embodiment, the site of cell metastasis analysis comprises the route of neoplastic spread. In some embodiment, cells can disperse via blood vasculature, lymphatics, within body cavities or combinations thereof. In another embodiment, cell metastasis analysis is performed in view of cell migration, dissemination, extravasation, proliferation or combinations thereof.

[0443] Accordingly, in a particularly preferred embodiment the modulators of the instant invention may be used to detect and quantify SEZ6 levels in a patient sample (e.g., plasma or blood) which may, in turn, be used to detect, diagnose or monitor SEZ6 associated disorders including proliferative disorders. In related embodiments the modulators of the instant invention may be used to detect, monitor and/or quantify circulating tumor cells either *in vivo* or *in vitro* (see, for example, WO 2012/0128801 which is incorporated herein by reference). In still other preferred embodiments the circulating tumor cells may comprise cancer stem cells.

[0444] In certain examples, the tumorigenic cells in a subject or a sample from a subject may be assessed or characterized using the disclosed modulators prior to therapy or regimen to establish a baseline. In other examples the sample is derived from a subject that was treated. In some examples the sample is taken from the subject at least about 1, 2, 4, 6, 7, 8, 10, 12, 14, 15, 16, 18, 20, 30, 60, 90 days, 6 months, 9 months, 12 months, or >12 months after the subject begins or terminates treatment. In certain examples, the tumorigenic cells are assessed or characterized after a certain number of doses (e.g., after 2, 5, 10, 20, 30 or more doses of a therapy). In other examples, the tumorigenic cells are characterized or assessed after 1 week, 2 weeks, 1 month, 2 months, 1 year, 2 years, 3 years, 4 years or more after receiving one or more therapies.

[0445] In another aspect, and as discussed in more detail below, the present invention provides kits for detecting, monitoring or diagnosing a hyperproliferative disorder, identifying individual having such a disorder for possible treatment or monitoring progression (or regression) of the disorder in a patient, wherein the kit comprises a modulator as described herein, and reagents for detecting the impact of the modulator on a sample.

[0446] Yet another aspect of the instant invention comprises the use of labeled SEZ6 for immunohistochemistry (IHC). In this respect SEZ6 IHC may be used as a diagnostic tool to aid in the diagnosis of various proliferative disorders and to monitor the potential response to treatments including SEZ6 modulator therapy. Compatible diagnostic assays may be performed on tissues that have been chemically fixed (including but not limited to: formaldehyde, gluteraldehyde, osmium tetroxide, potassium dichromate, acetic acid, alcohols, zinc salts, mercuric chloride, chromium tetroxide and picric acid) and embedded (including but not limited to: glycol methacrylate, paraffin and resins) or preserved via freezing. As discussed in more detail below such assays could be used to guide treatment decisions and determine dosing regimens and timing.

[0447] In a preferred embodiment the invention is directed to a method of diagnosing platinum resistant small cell lung

cancer comprising the steps of: (a) providing a small cell lung cancer tumor sample from a subject; (b) exposing the tumor sample to an anti-SEZ6 antibody labeled with a reporter wherein said anti-SEZ6 antibody associates with the tumor sample; and (c) detecting the reporter associated with the tumor sample. In a preferred embodiment such reporter may be detected using an *in vitro* diagnostic method (e.g. IHC, *in situ* hybridization or flow cytometry). In some embodiments, the step of providing the tumor sample may be performed separately from the step of exposing the tumor sample to an anti-SEZ6 antibody or the step of detecting the reporter associated with the tumor sample.

[0448] In another embodiment the invention is directed to a method of diagnosing medullary thyroid cancer comprising the steps of: (a) providing a medullary thyroid tumor sample from a subject; (b) exposing the tumor sample to an anti-SEZ6 antibody labeled with a reporter wherein said anti-SEZ6 antibody associates with the tumor sample; and (c) detecting the reporter associated with the tumor sample. In a preferred embodiment such reporter may be detected using an *in vitro* diagnostic method (e.g. IHC, *in situ* hybridization or flow cytometry). In some embodiments, the step of providing the tumor sample may be performed separately from the step of exposing the tumor sample to an anti-SEZ6 antibody or the step of detecting the reporter associated with the tumor sample.

B. Screening

[0449] In certain embodiments, the modulators can also be used to screen for or identify compounds or agents (e.g., drugs) that alter a function or activity of tumorigenic cells or progeny thereof by interacting with an antigen (e.g., genotypic or phenotypic components thereof). Such compounds and agents can be drug candidates that are screened for the treatment of a proliferative disorder, for example. In one embodiment, a system or method includes tumorigenic cells comprising SEZ6 and a compound or agent (e.g., drug), wherein the cells and compound or agent are in contact with each other. In such embodiments the subject cells may have been identified, monitored and/or enriched using the disclosed modulators.

[0450] In yet another embodiment, a method includes contacting, directly or indirectly, tumorigenic cells or progeny thereof with a test agent or compound and determining if the test agent or compound modulates an activity or function of the antigen-associated tumorigenic cells. One example of a direct interaction is physical interaction, while an indirect interaction includes the action of a composition upon an intermediary molecule that, in turn, acts upon the referenced entity (e.g., cell or cell culture). Exemplary activities or functions that can be modulated include changes in cell morphology or viability, expression of a marker, differentiation or de-differentiation, cell respiration, mitochondrial activity, membrane integrity, maturation, proliferation, viability, apoptosis or cell death.

[0451] Methods of screening and identifying agents and compounds include those suitable for high throughput screening, which include arrays of cells (e.g., microarrays) positioned or placed, optionally at pre-determined locations or addresses. For example, cells can be positioned or placed (pre-seeded) on a culture dish, tube, flask, roller bottle or plate (e.g., a single multi-well plate or dish such as an 8, 16, 32, 64, 96, 384 and 1536 multi-well plate or dish). High-throughput robotic or manual handling methods can probe chemical interactions and determine levels of expression of many genes in a short period of time. Techniques have been developed that utilize molecular signals (e.g., via fluorophores) and automated analyses that process information at a very rapid rate (see, e.g., Pinhasov et al., *Comb. Chem. High Throughput Screen.* 7:133 (2004)). For example, microarray technology has been extensively used to probe the interactions of thousands of genes at once, while providing information for specific genes (see, e.g., Mocellin and Rossi, *Adv. Exp. Med. Biol.* 593:19 (2007)).

[0452] Libraries that can be screened include, for example, small molecule libraries, phage display libraries, fully human antibody yeast display libraries (Adimab, LLC), siRNA libraries, and adenoviral transfection vectors.

X. Pharmaceutical Preparations and Therapeutic Uses

A. Formulations and Routes of Administration

[0453] Depending on the form of the modulator along with any optional conjugate, the mode of intended delivery, the disease being treated or monitored and numerous other variables, compositions of the invention may be formulated as desired using art-recognized techniques. In some embodiments, the therapeutic compositions of the invention may be administered neat or with a minimum of additional components while others may optionally be formulated to contain suitable pharmaceutically acceptable carriers comprising excipients and auxiliaries that are well known in the art (see, e.g., Gennaro, *Remington: The Science and Practice of Pharmacy with Facts and Comparisons: Drugfacts Plus*, 20th ed. (2003); Ansel et al., *Pharmaceutical Dosage Forms and Drug Delivery Systems*, 7th ed., Lippencott Williams and Wilkins (2004); Kibbe et al., *Handbook of Pharmaceutical Excipients*, 3rd ed., Pharmaceutical Press (2000)). Various pharmaceutically acceptable carriers, which include vehicles, adjuvants, and diluents, are readily available from numerous commercial sources. Moreover, an assortment of pharmaceutically acceptable auxiliary substances, such as pH adjusting and buffering agents, tonicity adjusting agents, stabilizers, wetting agents and the like, are also available.

Certain non-limiting exemplary carriers include saline, buffered saline, dextrose, water, glycerol, ethanol, and combinations thereof.

[0454] More particularly it will be appreciated that, in some embodiments, the therapeutic compositions of the invention may be administered neat or with a minimum of additional components. Conversely the SEZ6 modulators of the present invention may optionally be formulated to contain suitable pharmaceutically acceptable carriers comprising excipients and auxiliaries that are well known in the art and are relatively inert substances that facilitate administration of the modulator or which aid processing of the active compounds into preparations that are pharmaceutically optimized for delivery to the site of action. For example, an excipient can give form or consistency or act as a diluent to improve the pharmacokinetics or stability of the modulator. Suitable excipients or additives include, but are not limited to, stabilizing agents, wetting and emulsifying agents, salts for varying osmolality, encapsulating agents, buffers, and skin penetration enhancers. In certain preferred embodiments the pharmaceutical compositions may be provided in a lyophilized form and reconstituted in, for example, buffered saline prior to administration.

[0455] Disclosed modulators for systemic administration may be formulated for enteral, parenteral or topical administration. Indeed, all three types of formulation may be used simultaneously to achieve systemic administration of the active ingredient. Excipients as well as formulations for parenteral and nonparenteral drug delivery are set forth in Remington, The Science and Practice of Pharmacy 20th Ed. Mack Publishing (2000). Suitable formulations for parenteral administration include aqueous solutions of the active compounds in water-soluble form, for example, water-soluble salts. In addition, suspensions of the active compounds as appropriate for oily injection suspensions may be administered. Suitable lipophilic solvents or vehicles include fatty oils, for example, hexylsubstituted poly(lactide), sesame oil, or synthetic fatty acid esters, for example, ethyl oleate or triglycerides. Aqueous injection suspensions may contain substances that increase the viscosity of the suspension and include, for example, sodium carboxymethyl cellulose, sorbitol, and/or dextran. Optionally, the suspension may also contain stabilizers. Liposomes can also be used to encapsulate the agent for delivery into the cell.

[0456] Suitable formulations for enteral administration include hard or soft gelatin capsules, pills, tablets, including coated tablets, elixirs, suspensions, syrups or inhalations and controlled release forms thereof.

[0457] In general the compounds and compositions of the invention, comprising SEZ6 modulators may be administered *in vivo*, to a subject in need thereof, by various routes, including, but not limited to, oral, intravenous, intra-arterial, subcutaneous, parenteral, intranasal, intramuscular, intracranial, intracardiac, intraventricular, intratracheal, buccal, rectal, intraperitoneal, intradermal, topical, transdermal, and intrathecal, or otherwise by implantation or inhalation. The subject compositions may be formulated into preparations in solid, semi-solid, liquid, or gaseous forms; including, but not limited to, tablets, capsules, powders, granules, ointments, solutions, suppositories, enemas, injections, inhalants, and aerosols. The appropriate formulation and route of administration may be selected according to the intended application and therapeutic regimen.

B. Dosages

[0458] Similarly, the particular dosage regimen, i.e., dose, timing and repetition, will depend on the particular individual and that individual's medical history, as well as empirical considerations such as pharmacokinetics (e.g., half-life, clearance rate, etc.). Frequency of administration may be determined and adjusted over the course of therapy, and is based on reducing the number of proliferative or tumorigenic cells, maintaining the reduction of such neoplastic cells, reducing the proliferation of neoplastic cells, or delaying the development of metastasis. In other embodiments the dosage administered may be adjusted or attenuated to manage potential side effects and/or toxicity. Alternatively, sustained continuous release formulations of a subject therapeutic composition may be appropriate.

[0459] In general, the modulators of the invention may be administered in various ranges. These include about 10 $\mu\text{g/kg}$ body weight to about 100 mg/kg body weight per dose; about 50 $\mu\text{g/kg}$ body weight to about 5 mg/kg body weight per dose; about 100 $\mu\text{g/kg}$ body weight to about 10 mg/kg body weight per dose. Other ranges include about 100 $\mu\text{g/kg}$ body weight to about 20 mg/kg body weight per dose and about 0.5 mg/kg body weight to about 20 mg/kg body weight per dose. In certain embodiments, the dosage is at least about 100 $\mu\text{g/kg}$ body weight, at least about 250 $\mu\text{g/kg}$ body weight, at least about 750 $\mu\text{g/kg}$ body weight, at least about 3 mg/kg body weight, at least about 5 mg/kg body weight, at least about 10 mg/kg body weight.

[0460] In selected embodiments the modulators will be administered at approximately 10, 20, 30, 40, 50, 60, 70, 80, 90 or 100 $\mu\text{g/kg}$ body weight per dose. Other embodiments will comprise the administration of modulators at 200, 300, 400, 500, 600, 700, 800 or 900 $\mu\text{g/kg}$ body weight per dose. In other preferred embodiments the disclosed modulators will be administered at 1, 2, 3, 4, 5, 6, 7, 8, 9 or 10 mg/kg . In still other embodiments the modulators may be administered at 12, 14, 16, 18 or 20 mg/kg body weight per dose. In yet other embodiments the modulators may be administered at 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 90 or 100 mg/kg body weight per dose. In accordance with the teachings herein it will be appreciated that the aforementioned dosages are applicable to both unconjugated modulators and modulators conjugated to a cytotoxic agent. One of skill in the art could readily determine appropriate dosages for various

conjugated and unconjugated modulators based on preclinical animal studies, clinical observations and standard medical and biochemical techniques and measurements.

[0461] With regard to conjugated modulators particularly preferred embodiments will comprise dosages of between about 50 $\mu\text{g/kg}$ to about 5 mg/kg body weight per dose. In this regard conjugated modulators may be administered at 50, 75 or 100 $\mu\text{g/kg}$ or at 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9 or 1 mg/kg body weight per dose. In other preferred embodiments the conjugated modulators of the instant invention may be administered at 1.25, 1.5, 1.75, 2, 2.25, 2.5, 2.75, 3, 3.25, 3.5, 3.75, 4, 4.25, 4.5, 4.75 or 5 mg/kg body weight per dose. In particularly preferred embodiments such conjugated modulator dosages will be administered intravenously over a period of time. Moreover, such dosages may be administered multiple times over a defined course of treatment.

[0462] Other dosing regimens may be predicated on Body Surface Area (BSA) calculations as disclosed in U.S.P.N. 7,744,877. As is well known, the BSA is calculated using the patient's height and weight and provides a measure of a subject's size as represented by the surface area of his or her body. In certain embodiments, the modulators may be administered in dosages from 10 mg/m^2 to 800 mg/m^2 , from 50 mg/m^2 to 500 mg/m^2 and at dosages of 100 mg/m^2 , 150 mg/m^2 , 200 mg/m^2 , 250 mg/m^2 , 300 mg/m^2 , 350 mg/m^2 , 400 mg/m^2 or 450 mg/m^2 .

[0463] It will also be appreciated that art recognized and empirical techniques may be used to determine appropriate dosage for conjugated modulators (i.e., ADCs).

[0464] In any event, SEZ6 modulators (both conjugated and unconjugated) are preferably administered as needed to subjects in need thereof. Determination of the frequency of administration may be made by persons skilled in the art, such as an attending physician based on considerations of the condition being treated, age of the subject being treated, severity of the condition being treated, general state of health of the subject being treated and the like. Generally, an effective dose of the SEZ6 modulator is administered to a subject one or more times. More particularly, an effective dose of the modulator is administered to the subject once a month, more than once a month, or less than once a month. In certain embodiments, the effective dose of the SEZ6 modulator may be administered multiple times, including for periods of at least a month, at least six months, at least a year, at least two years or a period of several years. In yet other embodiments, several days (2, 3, 4, 5, 6 or 7), several weeks (1, 2, 3, 4, 5, 6, 7 or 8) or several months (1, 2, 3, 4, 5, 6, 7 or 8) or even a year or several years may lapse between administration of the disclosed modulators.

[0465] In certain preferred embodiments the course of treatment involving conjugated modulators will comprise multiple doses of the selected drug product (i.e., an ADC) over a period of weeks or months. More specifically, conjugated modulators of the instant invention may administered once every day, every two days, every four days, every week, every ten days, every two weeks, every three weeks, every month, every six weeks, every two months, every ten weeks or every three months. In this regard it will be appreciated that the dosages may be altered or the interval may be adjusted based on patient response and clinical practices.

[0466] Dosages and regimens may also be determined empirically for the disclosed therapeutic compositions in individuals who have been given one or more administration(s). For example, individuals may be given incremental dosages of a therapeutic composition produced as described herein. In selected embodiments the dosage may be gradually increased or reduced or attenuated based respectively on empirically determined or observed side effects or toxicity. To assess efficacy of the selected composition, a marker of the specific disease, disorder or condition can be followed as described previously. In embodiments where the individual has cancer, these include direct measurements of tumor size via palpation or visual observation, indirect measurement of tumor size by x-ray or other imaging techniques; an improvement as assessed by direct tumor biopsy and microscopic examination of the tumor sample; the measurement of an indirect tumor marker (e.g., PSA for prostate cancer) or an antigen identified according to the methods described herein, a decrease in pain or paralysis; improved speech, vision, breathing or other disability associated with the tumor; increased appetite; or an increase in quality of life as measured by accepted tests or prolongation of survival. It will be apparent to one of skill in the art that the dosage will vary depending on the individual, the type of neoplastic condition, the stage of neoplastic condition, whether the neoplastic condition has begun to metastasize to other location in the individual, and the past and concurrent treatments being used.

C. Combination Therapies

[0467] Combination therapies may be particularly useful in decreasing or inhibiting unwanted neoplastic cell proliferation, decreasing the occurrence of cancer, decreasing or preventing the recurrence of cancer, or decreasing or preventing the spread or metastasis of cancer. In such cases the modulators of the instant invention may function as sensitizing or chemosensitizing agents by removing the CSCs that would otherwise prop up and perpetuate the tumor mass and thereby allow for more effective use of current standard of care debulking or anti-cancer agents. "Combination therapy", as used herein, means the administration of a combination comprising at least one SEZ6 modulator and at least one therapeutic moiety (e.g., anti-cancer agent) wherein the combination preferably has therapeutic synergy or improves the measurable therapeutic effects in the treatment of cancer (e.g. medullary thyroid cancer or SCLC) over (i) the SEZ6 modulator used alone, or (ii) the therapeutic moiety used alone, or (iii) the use of the therapeutic moiety in

combination with another therapeutic moiety without the addition of a SEZ6 modulator. The term "therapeutic synergy", as used herein, means the combination of a SEZ6 modulator and one or more therapeutic moiety(ies) having a therapeutic effect greater than the additive effect of the combination of the SEZ6 modulator and the one or more therapeutic moiety(ies).

[0468] In the context of "combination therapy" in the instant invention, the therapeutic moiety may include one or more anti-cancer agents that include, but are not limited to, cytotoxic agents, cytostatic agents, anti-angiogenic agents, debulking agents, chemotherapeutic agents, radiotherapy and radiotherapeutic agents, targeted anti-cancer agents (including both monoclonal antibodies and small molecule entities), BRMs, therapeutic antibodies, cancer vaccines, cytokines, hormone therapies, radiation therapy and anti-metastatic agents and immunotherapeutic agents, including both specific and non-specific approaches.

[0469] Desired outcomes of the disclosed combinations are quantified by comparison to a control or baseline measurement. As used herein, relative terms such as "improve," "increase," or "reduce" indicate values relative to a control, such as a measurement in the same individual prior to initiation of treatment described herein, or a measurement in a control individual (or multiple control individuals) in the absence of the SEZ6 modulator described herein but in the presence of other therapeutic moiety(ies) such as standard of care treatment. A representative control individual is an individual afflicted with the same form of cancer as the individual being treated, who is about the same age as the individual being treated (to ensure that the stages of the disease in the treated individual and the control individual are comparable.)

[0470] Changes or improvements in response to therapy are generally statistically significant. As used herein, the term "significance" or "significant" relates to a statistical analysis of the probability that there is a non-random association between two or more entities. To determine whether or not a relationship is "significant" or has "significance," a "p-value" can be calculated. P-values that fall below a user-defined cut-off point are regarded as significant. A p-value less than or equal to 0.1, less than 0.05, less than 0.01, less than 0.005, or less than 0.001 may be regarded as significant.

[0471] A synergistic therapeutic effect may be an effect of at least about two-fold greater than the therapeutic effect elicited by a single therapeutic moiety or SEZ6 modulator, or the sum of the therapeutic effects elicited by the SEZ6 modulator or the single therapeutic moiety(ies) of a given combination, or at least about five-fold greater, or at least about ten-fold greater, or at least about twenty-fold greater, or at least about fifty-fold greater, or at least about one hundred-fold greater. A synergistic therapeutic effect may also be observed as an increase in therapeutic effect of at least 10% compared to the therapeutic effect elicited by a single therapeutic moiety or SEZ6 modulator, or the sum of the therapeutic effects elicited by the SEZ6 modulator or the single therapeutic moiety(ies) of a given combination, or at least 20%, or at least 30%, or at least 40%, or at least 50%, or at least 60%, or at least 70%, or at least 80%, or at least 90%, or at least 100%, or more. A synergistic effect is also an effect that permits reduced dosing of therapeutic agents when they are used in combination.

[0472] In practicing combination therapy, the modulator and anti-cancer agent may be administered to the subject simultaneously, either in a single composition, or as two or more distinct compositions using the same or different administration routes. Alternatively, the modulator may precede, or follow, the anti-cancer agent treatment by, e.g., intervals ranging from minutes to weeks. The time period between each delivery is such that the anti-cancer agent and modulator are able to exert a combined effect on the tumor. In at least one embodiment, both the anti-cancer agent and the modulator are administered within about 5 minutes to about two weeks of each other. In yet other embodiments, several days (2, 3, 4, 5, 6 or 7), several weeks (1, 2, 3, 4, 5, 6, 7 or 8) or several months (1, 2, 3, 4, 5, 6, 7 or 8) may lapse between administration of the modulator and the anti-cancer agent.

[0473] The combination therapy may be administered once, twice or at least for a period of time until the condition is treated, palliated or cured. In some embodiments, the combination therapy is administered multiple times, for example, from three times daily to once every six months. The administering may be on a schedule such as three times daily, twice daily, once daily, once every two days, once every three days, once weekly, once every two weeks, once every month, once every two months, once every three months, once every six months or may be administered continuously via a minipump. The combination therapy may be administered via any route, as noted previously. The combination therapy may be administered at a site distant from the site of the tumor.

[0474] In one embodiment a modulator is administered in combination with one or more anti-cancer agents for a short treatment cycle to a subject in need thereof. The invention also contemplates discontinuous administration or daily doses divided into several partial administrations. The modulator and anti-cancer agent may be administered interchangeably, on alternate days or weeks; or a sequence of antibody treatments may be given, followed by one or more treatments of anti-cancer agent therapy. In any event, as will be understood by those of ordinary skill in the art, the appropriate doses of chemotherapeutic agents will be generally around those already employed in clinical therapies wherein the chemotherapeutics are administered alone or in combination with other chemotherapeutics.

[0475] In another preferred embodiment the SEZ6 modulators of the instant invention may be used in maintenance therapy to reduce or eliminate the chance of tumor recurrence following the initial presentation of the disease. Preferably the disorder will have been treated and the initial tumor mass eliminated, reduced or otherwise ameliorated so the patient

is asymptomatic or in remission. At such time the subject may be administered pharmaceutically effective amounts of the disclosed modulators one or more times even though there is little or no indication of disease using standard diagnostic procedures. In some embodiments, the modulators will be administered on a regular schedule over a period of time, such as weekly, every two weeks, monthly, every six weeks, every two months, every three months every six months or annually. Given the teachings herein, one skilled in the art could readily determine favorable dosages and dosing regimens to reduce the potential of disease recurrence. Moreover such treatments could be continued for a period of weeks, months, years or even indefinitely depending on the patient response and clinical and diagnostic parameters.

[0476] In yet another preferred embodiment the modulators of the present invention may be used to prophylactically or as an adjuvant therapy to prevent or reduce the possibility of tumor metastasis following a debulking procedure. As used in the instant disclosure a "debulking procedure" is defined broadly and shall mean any procedure, technique or method that eliminates, reduces, treats or ameliorates a tumor or tumor proliferation. Exemplary debulking procedures include, but are not limited to, surgery, radiation treatments (i.e., beam radiation), chemotherapy, immunotherapy or ablation. At appropriate times readily determined by one skilled in the art in view of the instant disclosure the disclosed modulators may be administered as suggested by clinical, diagnostic or theragnostic procedures to reduce tumor metastasis. The modulators may be administered one or more times at pharmaceutically effective dosages as determined using standard techniques. Preferably the dosing regimen will be accompanied by appropriate diagnostic or monitoring techniques that allow it to be modified.

[0477] Yet other embodiments of the invention comprise administering the disclosed modulators to subjects that are asymptomatic but at risk of developing a proliferative disorder. That is, the modulators of the instant invention may be used in a truly preventative sense and given to patients that have been examined or tested and have one or more noted risk factors (e.g., genomic indications, family history, *in vivo* or *in vitro* test results, etc.) but have not developed neoplasia. In such cases those skilled in the art would be able to determine an effective dosing regimen through empirical observation or through accepted clinical practices.

[0478] In some embodiments the SEZ6 modulator may be used in combination with various first line SCLC treatments such platinum based agents (e.g. carboplatin, cisplatin and/or oxaloplatin). In one embodiment the combination therapy comprises the use of a SEZ6 modulator and a platinum based agent (e.g. carboplatin, cisplatin and/or oxaloplatin) and optionally one or more other therapeutic moiety(ies). In another embodiment the SEZ6 modulator may be used in combination with cyclophosphamide and optionally doxorubicin and/or vincristine. In yet another embodiment the SEZ6 modulator may be used in combination with etoposide. In further embodiments the SEZ6 modulator may be used in combination with topotecan or paclitaxel.

[0479] In other embodiments the SEZ6 modulator may be used in combination with various first line medullary thyroid treatments such cabozantinib or vandetanib. In one embodiment the combination therapy comprises the use of a SEZ6 modulator and cabozantinib and optionally one or more other therapeutic moiety(ies). In another embodiment the combination therapy comprises the use of a SEZ6 modulator and vandetanib and optionally one or more other therapeutic moiety(ies).

[0480] The combination therapy may comprise a SEZ6 modulator in combination with another therapeutic moiety that is effective on a medullary thyroid tumor comprising a mutated or aberrantly expressed gene or protein (e.g. RET).

[0481] The invention also provides for the combination of SEZ6 modulator with radiotherapy. In other embodiments a SEZ6 modulator may be used in combination with one or more of the anti-cancer described below.

D. Anti-Cancer Agents

[0482] The term "anti-cancer agent" or "anti-proliferative agent" means any agent that can be used to treat a cell proliferative disorder such as cancer, and includes, but is not limited to, cytotoxic agents, cytostatic agents, anti-angiogenic agents, debulking agents, chemotherapeutic agents, radiotherapy and radiotherapeutic agents, targeted anti-cancer agents, BRMs, therapeutic antibodies, cancer vaccines, cytokines, hormone therapies, radiation therapy and anti-metastatic agents and immunotherapeutic agents. It will be appreciated that, in selected embodiments as discussed above, such anti-cancer agents may comprise conjugates and may be associated with modulators prior to administration. In certain embodiments the disclosed anti-cancer agent will be linked to a SEZ6 modulator to provide an ADC as set forth herein.

[0483] As used herein the term "cytotoxic agent" means a substance that is toxic to the cells and decreases or inhibits the function of cells and/or causes destruction of cells. Typically, the substance is a naturally occurring molecule derived from a living organism. Examples of cytotoxic agents include, but are not limited to, small molecule toxins or enzymatically active toxins of bacteria (e.g., Diphtheria toxin, Pseudomonas endotoxin and exotoxin, Staphylococcal enterotoxin A), fungal (e.g., α -sarcin, restrictocin), plants (e.g., abrin, ricin, modeccin, viscumin, pokeweed anti-viral protein, saporin, gelonin, momoridin, trichosanthin, barley toxin, Aleurites fordii proteins, dianthin proteins, Phytolacca mericana proteins (PAPI, PAPII, and PAP-S), Momordica charantia inhibitor, curcin, croton, saponaria officinalis inhibitor, gelonin, mitegellin, restrictocin, phenomycin, neomycin, and the tricothecenes) or animals, (e.g., cytotoxic RNases, such as extracellular

pancreatic RNases; DNase I, including fragments and/or variants thereof).

[0484] For the purposes of the instant invention a "chemotherapeutic agent" comprises a chemical compound that non-specifically decreases or inhibits the growth, proliferation, and/or survival of cancer cells (e.g., cytotoxic or cytostatic agents). Such chemical agents are often directed to intracellular processes necessary for cell growth or division, and are thus particularly effective against cancerous cells, which generally grow and divide rapidly. For example, vincristine depolymerizes microtubules, and thus inhibits cells from entering mitosis. In general, chemotherapeutic agents can include any chemical agent that inhibits, or is designed to inhibit, a cancerous cell or a cell likely to become cancerous or generate tumorigenic progeny (e.g., TIC). Such agents are often administered, and are often most effective, in combination, e.g., in regimens such as CHOP or FOLFIRI. Again, in selected embodiments such chemotherapeutic agents may be conjugated to the disclosed modulators.

[0485] Examples of anti-cancer agents that may be used in combination with (or conjugated to) the modulators of the present invention include, but are not limited to, alkylating agents, alkyl sulfonates, aziridines, ethylenimines and methylenelamines, acetogenins, a camptothecin, bryostatin, callistatin, CC-1065, cryptophycins, dolastatin, duocarmycin, eleutherobin, pancratistatin, a sarcodictyin, spongistatin, nitrogen mustards, antibiotics, enediyne antibiotics, dynemicin, bisphosphonates, esperamicin, chromoprotein enediyne antibiotic chromophores, aclacinomysins, actinomycin, a-thramycin, azaserine, bleomycins, cactinomycin, carabacin, carminomycin, carzinophilin, chromomycinis, dactinomycin, daunorubicin, detorubicin, 6-diazo-5-oxo-L-norleucine, ADRIAMYCIN® doxorubicin, epirubicin, esorubicin, idarubicin, marcellomycin, mitomycins, mycophenolic acid, nogalamycin, olivomycins, peplomycin, potfiromycin, puromycin, quelamycin, rodorubicin, streptonigrin, streptozocin, tubercidin, ubenimex, zinostatin, zorubicin; anti-metabolites, erlotinib, vemurafenib, crizotinib, sorafenib, ibrutinib, enzalutamide, folic acid analogues, purine analogs, androgens, anti-adrenals, folic acid replenisher such as frolinic acid, aceglatone, aldophosphamide glycoside, aminolevulinic acid, eniluracil, amsacrine, bestabucil, bisantrene, edatraxate, defofamine, demecolcine, diaziquone, elfornithine, elliptinium acetate, an epothilone, etoglucid, gallium nitrate, hydroxyurea, lentinan, lonidainine, maytansinoids, mitoguazone, mitoxantrone, mopidanmol, nitraerine, pentostatin, phenamet, pirarubicin, losoxantrone, podophyllinic acid, 2-ethylhydrazide, procarbazine, PSK® polysaccharide complex (JHS Natural Products, Eugene, OR), razoxane; rhizoxin; sizofiran; spirogermanium; tenuazonic acid; triaziquone; 2,2',2''-trichlorotriethylamine; trichothecenes (especially T-2 toxin, verrucurin A, roridin A and anguidine); urethan; vindesine; dacarbazine; mannomustine; mitobronitol; mitolactol; pipobroman; gacytosine; arabinoside ("Ara-C"); cyclophosphamide; thiotepe; taxoids, chloranbucil; GEMZAR® gemcitabine; 6-thioguanine; mercaptopurine; methotrexate; platinum analogs, vinblastine; platinum; etoposide (VP-16); ifosfamide; mitoxantrone; vincristine; NAVELBINE® vinorelbine; novantrone; teniposide; edatrexate; daunomycin; aminopterin; xeloda; ibandronate; irinotecan (Camptosar, CPT-11), topoisomerase inhibitor RFS 2000; difluoromethylornithine; retinoids; capecitabine; combretastatin; leucovorin; oxaliplatin; inhibitors of PKC- α , Raf, H-Ras, EGFR and VEGF-A that reduce cell proliferation and pharmaceutically acceptable salts, acids or derivatives of any of the above. Also included in this definition are anti-hormonal agents that act to regulate or inhibit hormone action on tumors such as anti-estrogens and selective estrogen receptor modulators, aromatase inhibitors that inhibit the enzyme aromatase, which regulates estrogen production in the adrenal glands, and anti-androgens; as well as troxacitabine (a 1,3- dioxolane nucleoside cytosine analog); antisense oligonucleotides, ribozymes such as a VEGF expression inhibitor and a HER2 expression inhibitor; vaccines, PROLEUKIN® rIL-2; LURTOTECAN® topoisomerase 1 inhibitor; ABARELIX® rmRH; Vinorelbine and Esperamicins and pharmaceutically acceptable salts, acids or derivatives of any of the above.

[0486] In other embodiments the modulators of the instant invention may be used in combination with any one of a number of antibodies (or immunotherapeutic agents) presently in clinical trials or commercially available. To this end the disclosed modulators may be used in combination with an antibody selected from the group consisting of abagovomab, adecatumumab, afutuzumab, alemtuzumab, altumomab, amatuximab, anatumomab, arcitumomab, bavituximab, bectumomab, bevacizumab, bivatuzumab, blinatumomab, brentuximab, cantuzumab, catumaxomab, cetuximab, citatuzumab, cixutumumab, clivatuzumab, conatumumab, daratumumab, drozitumab, duligotumab, dusigitumab, detumomab, dacetuzumab, dalotuzumab, ecomeximab, elotuzumab, ensituximab, ertumaxomab, etaracizumab, farletuzumab, ficlatuzumab, figitumumab, flanvotumab, futuximab, ganitumab, gemtuzumab, girentuximab, glembatumumab, ibritumomab, igovomab, imgatuzumab, indatuximab, inotuzumab, intetumumab, ipilimumab, iratumumab, labetuzumab, lexatumumab, lintuzumab, lorvotuzumab, lucatumumab, mapatumumab, matuzumab, milatuzumab, minretumomab, mitumomab, moxetumomab, narnatumab, naptumomab, necitumumab, nimotuzumab, nofetumomab, ocaratuzumab, ofatumumab, olaratumab, onartuzumab, oportuzumab, oregovomab, panitumumab, parsatuzumab, patritumab, pemtumomab, pertuzumab, pintumomab, pritumumab, racotumomab, radretumab, rilotumumab, rituximab, robatumumab, satumomab, sibrotuzumab, siltuximab, simtuzumab, solitomab, tacatuzumab, taplitumomab, tenatumomab, teprotumumab, tigatuzumab, tositumomab, trastuzumab, tucotuzumab, ublituximab, veltuzumab, vorsetuzumab, votumumab, zalutumumab, CC49, 3F8 and combinations thereof.

[0487] Still other particularly preferred embodiments will comprise the use of antibodies approved for cancer therapy including, but not limited to, rituximab, trastuzumab, gemtuzumab, ozogamcin, alemtuzumab, ibritumomab, tiuxetan, tositumomab, bevacizumab, cetuximab, panitumumab, ofatumumab, ipilimumab and brentuximab vedotin. Those skilled

in the art will be able to readily identify additional anti-cancer agents that are compatible with the teachings herein.

E. Radiotherapy

[0488] The present invention also provides for the combination of modulators with radiotherapy (i.e., any mechanism for inducing DNA damage locally within tumor cells such as gamma-irradiation, X-rays, UV-irradiation, microwaves, electronic emissions and the like). Combination therapy using the directed delivery of radioisotopes to tumor cells is also contemplated, and may be used in connection with a targeted anti-cancer agent or other targeting means. Typically, radiation therapy is administered in pulses over a period of time from about 1 to about 2 weeks. The radiation therapy may be administered to subjects having head and neck cancer for about 6 to 7 weeks. Optionally, the radiation therapy may be administered as a single dose or as multiple, sequential doses.

XI. Indications

[0489] It will be appreciated that the modulators of the instant invention may be used to diagnose, treat or inhibit the occurrence or recurrence of any SEZ6 associated disorder. Accordingly, whether administered alone or in combination with an anti-cancer agent or radiotherapy, the modulators of the invention are particularly useful for generally treating neoplastic conditions in patients or subjects which may include benign or malignant tumors (e.g., adrenal, liver, kidney, bladder, breast, gastric, ovarian, colorectal, prostate, pancreatic, lung, thyroid, hepatic, cervical, endometrial, esophageal and uterine carcinomas; sarcomas; glioblastomas; and various head and neck tumors); leukemias and lymphoid malignancies; other disorders such as neuronal, glial, astrocytic, hypothalamic and other glandular, macrophagal, epithelial, stromal and blastocoelic disorders; and inflammatory, angiogenic, immunologic disorders and disorders caused by pathogens. Particularly, key targets for treatment are neoplastic conditions comprising solid tumors, although hematologic malignancies are within the scope of the invention. Preferably the "subject" or "patient" to be treated will be human although, as used herein, the terms are expressly held to comprise any mammalian species.

[0490] More specifically, neoplastic conditions subject to treatment in accordance with the instant invention may be selected from the group including, but not limited to, adrenal gland tumors, AIDS-associated cancers, alveolar soft part sarcoma, astrocytic tumors, bladder cancer (squamous cell carcinoma and transitional cell carcinoma), bone cancer (adamantinoma, aneurismal bone cysts, osteochondroma, osteosarcoma), brain and spinal cord cancers, metastatic brain tumors, breast cancer, carotid body tumors, cervical cancer, chondrosarcoma, chordoma, chromophobe renal cell carcinoma, clear cell carcinoma, colon cancer, colorectal cancer, cutaneous benign fibrous histiocytomas, desmoplastic small round cell tumors, ependymomas, Ewing's tumors, extraskeletal myxoid chondrosarcoma, fibrogenesis imperfecta ossium, fibrous dysplasia of the bone, gallbladder and bile duct cancers, gestational trophoblastic disease, germ cell tumors, head and neck cancers, islet cell tumors, Kaposi's Sarcoma, kidney cancer (nephroblastoma, papillary renal cell carcinoma), leukemias, lipoma/benign lipomatous tumors, liposarcoma/malignant lipomatous tumors, liver cancer (hepatoblastoma, hepatocellular carcinoma), lymphomas, lung cancers (small cell carcinoma, adenocarcinoma, squamous cell carcinoma, large cell carcinoma etc.), medulloblastoma, melanoma, meningiomas, multiple endocrine neoplasia, multiple myeloma, myelodysplastic syndrome, neuroblastoma, neuroendocrine tumors, ovarian cancer, pancreatic cancers, papillary thyroid carcinomas, parathyroid tumors, pediatric cancers, peripheral nerve sheath tumors, pheochromocytoma, pituitary tumors, prostate cancer, posterior uveal melanoma, rare hematologic disorders, renal metastatic cancer, rhabdoid tumor, rhabdomyosarcoma, sarcomas, skin cancer, soft-tissue sarcomas, squamous cell cancer, stomach cancer, synovial sarcoma, testicular cancer, thymic carcinoma, thymoma, thyroid metastatic cancer, and uterine cancers (carcinoma of the cervix, endometrial carcinoma, and leiomyoma).

[0491] In certain preferred embodiments the proliferative disorder will comprise a solid tumor including, but not limited to, adrenal, liver, kidney, bladder, breast, gastric, ovarian, cervical, uterine, esophageal, colorectal, prostate, pancreatic, lung (both small cell and non-small cell), thyroid, carcinomas, sarcomas, glioblastomas and various head and neck tumors. In other preferred embodiments, and as shown in the Examples below, the disclosed modulators are especially effective at treating small cell lung cancer (SCLC) and non-small cell lung cancer (NSCLC) (e.g., squamous cell non-small cell lung cancer or squamous cell small cell lung cancer).

[0492] In one embodiment, the small cell or non-small cell lung cancer is refractory, relapsed or resistant to a taxane (e.g., docetaxel, paclitaxel, larotaxel or cabazitaxel) and/or a platinum based agent (e.g., carboplatin, cisplatin, oxaliplatin, topotecan, etc.). With regards to platinum based agents, the primary standard-of-care chemotherapy for SCLC is combined cisplatin and etoposide. Following a course of treatment, many patients become resistant to the platinum based agent (Stewart, 2004; PMID: 20047843). There are multiple mechanisms of resistance to platinum-based drugs including changes to the membrane composition, increased hypoxia, decreased expression of copper transporter genes, and increased expression of multidrug resistance genes, among other mechanisms. Given that most patients with SCLC will first be treated with the standard-of-care treatment, in some embodiments of the invention, the antibodies disclosed herein may be used to treat patients who have either failed to respond or relapsed. Initial platinum responsiveness is

high but the majority of cancer patients will eventually relapse with cisplatin-resistant disease. In particularly preferred embodiments the disclosed modulators may be used in a conjugated form to treat small cell lung cancer. In other preferred embodiments the disclosed modulators may be used in a conjugated form to treat platinum resistant small cell lung cancer in a subject in need thereof. With regard to small cell lung cancer particularly preferred embodiments will comprise the administration of conjugated modulators (ADCs). In selected embodiments the conjugated modulators will be administered to patients exhibiting limited stage disease. In other embodiments the disclosed modulators will be administered to patients exhibiting extensive stage disease. In other preferred embodiments the disclosed conjugated modulators will be administered to refractory patients (i.e., those who recur during or shortly after completing a course of initial therapy). Still other embodiments comprise the administration of the disclosed modulators to sensitive patients (i.e., those whose relapse is longer than 2-3 months after primary therapy. In each case it will be appreciated that compatible modulators may be in a conjugated or unconjugated state depending the selected dosing regimen and the clinical diagnosis.

[0493] As discussed above the disclosed modulators may further be used to prevent, treat or diagnose tumors with neuroendocrine features or phenotypes including neuroendocrine tumors. True or canonical neuroendocrine tumors (NETs) arising from the dispersed endocrine system are relatively rare, with an incidence of 2-5 per 100,000 people, but highly aggressive. Neuroendocrine tumors occur in the kidney, genitourinary tract (bladder, prostate, ovary, cervix, and endometrium), gastrointestinal tract (colon, stomach), thyroid (medullary thyroid cancer), and lung (small cell lung carcinoma and large cell neuroendocrine carcinoma). These tumors may secrete several hormones including serotonin and/or chromogranin A that can cause debilitating symptoms known as carcinoid syndrome. Such tumors can be denoted by positive immunohistochemical markers such as neuron-specific enolase (NSE, also known as gamma enolase, gene symbol = ENO2), CD56 (or NCAM1), chromogranin A (CHGA), and synaptophysin (SYP) or by genes known to exhibit elevated expression such as ASCL1. Unfortunately traditional chemotherapies have not been particularly effective in treating NETs and liver metastasis is a common outcome.

[0494] In some embodiments of the invention, antibody drug conjugates comprising an antibody conjugated directly or indirectly to a therapeutic moiety, wherein the antibody specifically binds to SEZ6, may be used to treat a subject suffering from cancer (e.g. medullary thyroid cancer). In preferred embodiments the antibody drug conjugate used to treat cancer (e.g. medullary thyroid cancer) will comprise an anti-SEZ6 antibody comprising a light chain variable region comprising three CDRs as set forth in FIG. 10A and a heavy chain variable region comprising three CDRs as set forth in FIG. 10B. In other embodiments the antibody drug conjugate used to treat a subject suffering from cancer (e.g. medullary thyroid cancer) may compete with the anti-SEZ6 antibodies comprising light and heavy chain variable regions set forth in FIGS. 10A and 10B. In yet another embodiment, the antibody drug conjugates used to treat a a subject suffering from cancer (e.g. medullary thyroid cancer) may comprise a humanized anti-SEZ6 antibody, for example, hSC17.16, hSC17.17, hSC17.24, hSC17.28, hSC17.34, hSC17.46, hSC17.151, hSC17.155, hSC17.156, hSC17.161 and hSC17.200 as set out in FIGS. 10A and 10B and any antibodies that compete with such humanized antibodies. In other embodiments of the invention, antibody drug conjugates comprising an antibody conjugated directly or indirectly to a therapeutic moiety, wherein the antibody specifically binds to SEZ6, may be used to treat a subject suffering from small cell lung cancer (e.g. platinum resistant small cell lung cancer). In preferred embodiments the antibody drug conjugate used to treat a subject suffering from small cell lung cancer (e.g. platinum resistant small cell lung cancer) may comprise a humanized anti-SEZ6 antibody, for example, hSC17.16, hSC17.17, hSC17.24, hSC17.28, hSC17.34, hSC17.46, hSC17.151, hSC17.155, hSC17.156, hSC17.161 and hSC17.200, as set out in FIGS. 10A and 10B and any antibodies that compete with such humanized antibodies. In preferred embodiments, the subject suffering from small cell lung cancer will have previously been treated with a platinum based agent.

[0495] While the disclosed modulators may be advantageously used to treat neuroendocrine tumors they may also be used to treat, prevent or diagnose pseudo neuroendocrine tumors (pNETs) that genotypically or phenotypically mimic, resemble or exhibit common traits with canonical neuroendocrine tumors. Pseudo neuroendocrine tumors or tumors with neuroendocrine features are tumors that arise from cells of the diffuse neuroendocrine system or from cells in which a neuroendocrine differentiation cascade has been aberrantly reactivated during the oncogenic process. Such pNETs commonly share certain phenotypic or biochemical characteristics with traditionally defined neuroendocrine tumors, including the ability to produce subsets of biologically active amines, neurotransmitters, and peptide hormones. Histologically, such tumors (NETs and pNETs) share a common appearance often showing densely connected small cells with minimal cytoplasm of bland cytopathology and round to oval stippled nuclei. For the purposes of the instant invention commonly expressed histological markers or genetic markers that may be used to define neuroendocrine and pseudo neuroendocrine tumors include, but are not limited to, chromogranin A, CD56, synaptophysin, PGP9.5, ASCL1 and neuron-specific enolase (NSE).

[0496] Accordingly the modulators of the instant invention may beneficially be used to treat both pseudo neuroendocrine tumors and canonical neuroendocrine tumors. In this regard the modulators may be used as described herein to treat neuroendocrine tumors (both NET and pNET) arising in the kidney, genitourinary tract (bladder, prostate, ovary, cervix, and endometrium), gastrointestinal tract (colon, stomach), thyroid (medullary thyroid cancer), and lung (small cell lung

carcinoma and large cell neuroendocrine carcinoma). Moreover, the modulators of the instant invention may be used to treat tumors expressing one or more markers selected from the group consisting of NSE, CD56, synaptophysin, chromogranin A, ASCL1 and PGP9.5 (UCHL1). That is, the present invention may be used to treat a subject suffering from a tumor that is NSE⁺ or CD56⁺ or PGP9.5⁺ or ASCL1⁺ or SYP⁺ or CHGA⁺ or some combination thereof.

[0497] With regard to hematologic malignancies it will be further appreciated that the compounds and methods of the present invention may be particularly effective in treating a variety of B-cell lymphomas, including low grade/NHL follicular cell lymphoma (FCC), mantle cell lymphoma (MCL), diffuse large cell lymphoma (DLCL), small lymphocytic (SL) NHL, intermediate grade/follicular NHL, intermediate grade diffuse NHL, high grade immunoblastic NHL, high grade lymphoblastic NHL, high grade small non-cleaved cell NHL, bulky disease NHL, Waldenstrom's Macroglobulinemia, lymphoplasmacytoid lymphoma (LPL), mantle cell lymphoma (MCL), follicular lymphoma (FL), diffuse large cell lymphoma (DLCL), Burkitt's lymphoma (BL), AIDS-related lymphomas, monocytic B cell lymphoma, angioimmunoblastic lymphadenopathy, small lymphocytic, follicular, diffuse large cell, diffuse small cleaved cell, large cell immunoblastic lymphoblastoma, small, non-cleaved, Burkitt's and non-Burkitt's, follicular, predominantly large cell; follicular, predominantly small cleaved cell; and follicular, mixed small cleaved and large cell lymphomas. See, Gaidono et al., "Lymphomas", IN CANCER: PRINCIPLES & PRACTICE OF ONCOLOGY, Vol. 2: 2131-2145 (DeVita et al., eds., 5^{sup}.th ed. 1997). It should be clear to those of skill in the art that these lymphomas will often have different names due to changing systems of classification, and that patients having lymphomas classified under different names may also benefit from the combined therapeutic regimens of the present invention.

[0498] The present invention also provides for a preventative or prophylactic treatment of subjects who present with benign or precancerous tumors. Beyond being a SEZ6 associated disorder it is not believed that any particular type of tumor or proliferative disorder should be excluded from treatment using the present invention. However, the type of tumor cells may be relevant to the use of the invention in combination with secondary therapeutic agents, particularly chemotherapeutic agents and targeted anti-cancer agents.

XII. Research Reagents

[0499] Other preferred embodiments of the invention also exploit the properties of the disclosed modulators as an instrument useful for identifying, monitoring, isolating, sectioning or enriching populations or subpopulations of tumor initiating cells through methods such as flow cytometry, fluorescent activated cell sorting (FACS), magnetic activated cell sorting (MACS) or laser mediated sectioning. Those skilled in the art will appreciate that the modulators may be used in several compatible techniques for the characterization and manipulation of TIC including cancer stem cells (e.g., see U.S.S.Ns. 12/686,359, 12/669,136 and 12/757,649 each of which is incorporated herein by reference in its entirety).

XIII. Articles of Manufacture

[0500] Pharmaceutical packs and kits comprising one or more containers, comprising one or more doses of a SEZ6 modulator are also provided. In certain embodiments, a unit dosage is provided wherein the unit dosage contains a predetermined amount of a composition comprising, for example, an anti-SEZ6 antibody, with or without one or more additional agents. For other embodiments, such a unit dosage is supplied in single-use prefilled syringe for injection. In still other embodiments, the composition contained in the unit dosage may comprise saline, sucrose, or the like; a buffer, such as phosphate, or the like; and/or be formulated within a stable and effective pH range. Alternatively, in certain embodiments, the composition may be provided as a lyophilized powder that may be reconstituted upon addition of an appropriate liquid, for example, sterile water. In certain preferred embodiments, the composition comprises one or more substances that inhibit protein aggregation, including, but not limited to, sucrose and arginine. Any label on, or associated with, the container(s) indicates that the enclosed composition is used for diagnosing or treating the disease condition of choice.

[0501] The present invention also provides kits for producing single-dose or multi-dose administration units of a SEZ6 modulator and, optionally, one or more anti-cancer agents. The kit comprises a container and a label or package insert on or associated with the container. Suitable containers include, for example, bottles, vials, syringes, etc. The containers may be formed from a variety of materials such as glass or plastic and contain a pharmaceutically effective amount of the disclosed modulators in a conjugated or unconjugated form. In other preferred embodiments the container(s) comprise a sterile access port (for example the container may be an intravenous solution bag or a vial having a stopper pierceable by a hypodermic injection needle). Such kits will generally contain in a suitable container a pharmaceutically acceptable formulation of the SEZ6 modulator and, optionally, one or more anti-cancer agents in the same or different containers. The kits may also contain other pharmaceutically acceptable formulations, either for diagnosis or combined therapy. For example, in addition to the SEZ6 modulator of the invention such kits may contain any one or more of a range of anti-cancer agents such as chemotherapeutic or radiotherapeutic drugs; anti-angiogenic agents; anti-metastatic agents; targeted anti-cancer agents; cytotoxic agents; and/or other anti-cancer agents. Such kits may also provide appropriate

reagents to conjugate the SEZ6 modulator with an anti-cancer agent or diagnostic agent (e.g., see U.S.P.N. 7,422,739 which is incorporated herein by reference in its entirety).

[0502] More specifically the kits may have a single container that contains the SEZ6 modulator, with or without additional components, or they may have distinct containers for each desired agent. Where combined therapeutics are provided for conjugation, a single solution may be pre-mixed, either in a molar equivalent combination, or with one component in excess of the other. Alternatively, the SEZ6 modulator and any optional anti-cancer agent of the kit may be maintained separately within distinct containers prior to administration to a patient. The kits may also comprise a second/third container means for containing a sterile, pharmaceutically acceptable buffer or other diluent such as bacteriostatic water for injection (BWFI), phosphate-buffered saline (PBS), Ringer's solution and dextrose solution.

[0503] When the components of the kit are provided in one or more liquid solutions, the liquid solution is preferably an aqueous solution, with a sterile aqueous solution being particularly preferred. However, the components of the kit may be provided as dried powder(s). When reagents or components are provided as a dry powder, the powder can be reconstituted by the addition of a suitable solvent. It is envisioned that the solvent may also be provided in another container.

[0504] As indicated briefly above the kits may also contain a means by which to administer the antibody and any optional components to an animal or patient, e.g., one or more needles or syringes, or even an eye dropper, pipette, or other such like apparatus, from which the formulation may be injected or introduced into the animal or applied to a diseased area of the body. The kits of the present invention will also typically include a means for containing the vials, or such like, and other component in close confinement for commercial sale, such as, e.g., injection or blow-molded plastic containers into which the desired vials and other apparatus are placed and retained. Any label or package insert indicates that the SEZ6 modulator composition is used for treating cancer, for example small cell lung cancer.

[0505] In other preferred embodiments the modulators of the instant invention may be used in conjunction with, or comprise, diagnostic or therapeutic devices useful in the diagnosis or treatment of proliferative disorders. For example, in on preferred embodiment the compounds and compositions of the instant invention may be combined with certain diagnostic devices or instruments that may be used to detect, monitor, quantify or profile cells or marker compounds involved in the etiology or manifestation of proliferative disorders. For selected embodiments the marker compounds may comprise NSE, CD56, synaptophysin, chromogranin A, and PGP9.5.

[0506] In particularly preferred embodiments the devices may be used to detect, monitor and/or quantify circulating tumor cells either *in vivo* or *in vitro* (see, for example, WO 2012/0128801 which is incorporated herein by reference). In still other preferred embodiments, and as discussed above, the circulating tumor cells may comprise cancer stem cells.

XIV. Miscellaneous

[0507] Unless otherwise defined herein, scientific and technical terms used in connection with the present invention shall have the meanings that are commonly understood by those of ordinary skill in the art. Further, unless otherwise required by context, singular terms shall include pluralities and plural terms shall include the singular. More specifically, as used in this specification and the appended claims, the singular forms "a," "an" and "the" include plural referents unless the context clearly dictates otherwise. Thus, for example, reference to "a protein" includes a plurality of proteins; reference to "a cell" includes mixtures of cells, and the like. In addition, ranges provided in the specification and appended claims include both end points and all points between the end points. Therefore, a range of 2.0 to 3.0 includes 2.0, 3.0, and all points between 2.0 and 3.0.

[0508] Generally, nomenclature used in connection with, and techniques of, cell and tissue culture, molecular biology, immunology, microbiology, genetics and protein and nucleic acid chemistry and hybridization described herein are those well known and commonly used in the art. The methods and techniques of the present invention are generally performed according to conventional methods well known in the art and as described in various general and more specific references that are cited and discussed throughout the present specification unless otherwise indicated. See, e.g., Abbas et al., Cellular and Molecular Immunology, 6th ed., W.B. Saunders Company (2010); Sambrook J. & Russell D. Molecular Cloning: A Laboratory Manual, 3rd ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (2000); Ausubel et al., Short Protocols in Molecular Biology: A Compendium of Methods from Current Protocols in Molecular Biology, Wiley, John & Sons, Inc. (2002); Harlow and Lane Using Antibodies: A Laboratory Manual, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1998); and Coligan et al., Short Protocols in Protein Science, Wiley, John & Sons, Inc. (2003). Enzymatic reactions and purification techniques are performed according to manufacturer's specifications, as commonly accomplished in the art or as described herein. The nomenclature used in connection with, and the laboratory procedures and techniques of, analytical chemistry, synthetic organic chemistry, and medicinal and pharmaceutical chemistry described herein are those well known and commonly used in the art. Moreover, any section headings used herein are for organizational purposes only and are not to be construed as limiting the subject matter described.

XV. SEZ6 References

[0509] All references or documents disclosed or cited within this specification are, without limitation, incorporated herein by reference in their entirety. Moreover, any section headings used herein are for organizational purposes only and are not to be construed as limiting the subject matter described.

1. Bork P, Beckmann G. (1993). The CUB domain. A widespread module in developmentally regulated proteins. *J Mol Biol.* 231:539-45. PMID: 8510165.
2. Cook M et al. (2010). Notch in the development of thyro C-cells and the treatment of medullary thyroid cancer. *Am J Transl Res.* 2:119-25. PMID:20182588
3. Galluzzo P, and Bocchetta M (2011). Notch signaling in lung cancer. *Expert Rev Anticancer Ther.* 11:533-40. PMID: 21504320.
4. Gunnarsen JM et al. (2007). Sez-6 proteins affect dendritic arborization patterns and excitability of cortical pyramidal neurons. *Neuron.* 56:621-39. PMID: 18031681.
5. Gunnarsen JM et al. (2009). Seizure-related gene 6 (Sez-6) in amacrine cells of the rodent retina and the consequence of gene deletion. *PLoS One.* 4:e6546. PMID:19662096.
6. Haddad RI (2013). How to incorporate new tyrosine kinase inhibitors in the treatment of patients with medullary thyroid cancer. *J Clin Oncolo.* 31:3618-20. PMID:24002516.
7. Herbst R, Nicklin MJ (1997). SEZ-6: promoter selectivity, genomic structure and localized expression in the brain. *Brain Res Mol Brain Res.* 44:309-22. PMID: 9073173.
8. Klimstra DS, et al. (2010). The pathologic classification of neuroendocrine tumors: a review of nomenclature, grading, and staging systems. *Pancreas.* 39:707-12. PMID: 20664470.
9. Klöppel G. (2011). Classification and pathology of gastroenteropancreatic neuroendocrine neoplasms. *Endocr Relat Cancer.* 18 Suppl 1:S1-16. PMID: 22005112.
10. Mulley JC et al. (2011). The Role of Seizure-Related SEZ6 as a Susceptibility Gene in Febrile Seizures. *Neurol Res Int.* 2011:917565. PMID: 21785725.
11. Shimizu-Nishikawa K et al., (1995). Cloning and expression of SEZ-6, a brain-specific and seizure-related cDNA. *Brain Res Mol Brain Res.* 28:201-10. PMID: 7723619.
12. Yao JC et al. (2008). One hundred years after "carcinoid": epidemiology of and prognostic factors for neuroendocrine tumors in 35,825 cases in the United States. *J Clin Oncol.* 26:3063-72. PMID: 18565894.
13. Yu ZL et al., (2007). Febrile seizures are associated with mutation of seizure-related (SEZ) 6, a brain-specific gene. *J Neurosci Res.* 85:166-72. PMID: 17086543.

XVI. Sequence Listing Summary

[0510] Appended to the instant application is a sequence listing comprising a number of nucleic acid and amino acid sequences. The following Table 2 provides a summary of the included sequences.

TABLE 2

SEQ ID NO.	Description
1	SEZ6 isoform 1 mRNA sequence
2	SEZ6 isoform 2 mRNA sequence
3	SEZ6 isoform 1 protein sequence
4	SEZ6 isoform 2 protein sequence
5	cDNA sequence of human SEZ6 ORF
6	Human SEZ6 protein
7	cDNA sequence of a commercial SEZ6 clone (BC146292)
8	Human SEZ6-Fc ORF
9	Human SEZ6-Fc protein
10	cDNA sequence of mouse SEZ6 ORF
11	Mouse SEZ6 protein
12	cDNA sequence of rat SEZ6 ORF

EP 3 338 793 A1

(continued)

SEQ ID NO.	Description
13	Rat SEZ6 protein
14	cDNA sequence of cynomolgus SEZ6 ORF
15	Cynomolgus SEZ6 protein
16	cDNA sequence of human SEZ6L ECD
17	Human SEZ6L ECD protein
18	cDNA sequence of human SEZ6L2 ECD
19	Human SEZ6L2 ECD protein
20	SC 17.1 VL protein
21	SC 17.1 VH protein
22-169	Additional murine VL and VH proteins as in SEQ ID NOs 20-21
170	hSC 17.16 VL protein
171	hSC 17.16 VH protein
172-199	Additional humanized VL and VH proteins as in SEQ ID NOs 170-171
200	Asn-Pro-Thr-Tyr (motif on the SEZ6 C-terminal cytoplasmic domain)
201	9-Histidine Tag
202-219	Reserve
220	SC 17.1 VL nucleic acid
221	SC 17.1 VH nucleic acid
222-369	Additional murine VL and VH nucleic acids as in SEQ ID NOs 220-221
370	hSC17.16 VL nucleic acid
371	hSC17.16 VH nucleic acid
372-399	Additional humanized VL and VH nucleic acids as in SEQ ID NOs 270-271
400	hSC 17.200 full length light chain amino acid sequence
401	hSC 17.200 full length heavy chain amino acid sequence
402	hSC17.200vL1 full length light chain amino acid sequence
403	Kappa constant region protein
404	IgG1 constant region protein
405, 406, 407	hSC17.16 CDRL1, CDRL2, CDRL3
408, 409, 410	hSC17.16 CDRH1, CDRH2, CDRH3
411, 412, 413	hSC17.17 CDRL1, CDRL2, CDRL3
414, 415, 416	hSC17.17 CDRH1, CDRH2, CDRH3
417, 418, 419	hSC17.24 CDRL1, CDRL2, CDRL3
420, 421, 422	hSC17.24 CDRH1, CDRH2, CDRH3
423, 424, 425	hSC17.28 CDRL1, CDRL2, CDRL3
426, 427, 428	hSC17.28 CDRH1, CDRH2, CDRH3
429, 430, 431	hSC17.34 CDRL1, CDRL2, CDRL3
432, 433, 434	hSC17.34 CDRH1, CDRH2, CDRH3
435, 436, 437	hSC17.46 CDRL1, CDRL2, CDRL3

(continued)

SEQ ID NO.	Description
438, 439, 440	hSC17.46 CDRH1, CDRH2, CDRH3
441, 442, 443	hSC17.151 CDRL1, CDRL2, CDRL3
444, 445, 446	hSC17.151 CDRH1, CDRH2, CDRH3
447, 448, 449	hSC17.155 CDRL1, CDRL2, CDRL3
450, 451, 452	hSC17.155 CDRH1, CDRH2, CDRH3
453, 454, 455	hSC17.156 CDRL1, CDRL2, CDRL3
456, 457, 458	hSC17.156 CDRH1, CDRH2, CDRH3
459, 460, 461	hSC17.161 CDRL1, CDRL2, CDRL3
462, 463, 464	hSC17.161 CDRH1, CDRH2, CDRH3
465, 466, 467	hSC17.200 CDRL1, CDRL2, CDRL3
468, 469, 470	hSC17.200 CDRH1, CDRH2, CDRH3
471	hSC17.155vH1FR1
472	hSC17.155vH2 FR1
473	hSC17.155vH3 CDRH1
474	hSC17.155vH4 CDRH2
475	hSC17.155vH5 CDRH2
476	hSC17.155vH6 CDRH2
477	hSC17.161vH1 FR1
478	hSC17.161vH1 FR2
479	hSC17.161vH1 FR3
480	hSC17.200vL1 CDRL1

XVII. Selected Embodiments of the Invention

[0511] In addition to the disclosure herein, the present invention is directed to selected embodiments specifically set forth immediately below.

EXAMPLES

[0512] The present invention, thus generally described above, will be understood more readily by reference to the following examples, which are provided by way of illustration and are not intended to be limiting of the instant invention. The examples are not intended to represent that the experiments below are all or the only experiments performed. Unless indicated otherwise, parts are parts by weight, molecular weight is weight average molecular weight, temperature is in degrees Centigrade, and pressure is at or near atmospheric.

EXAMPLE 1

IDENTIFICATION OF TUMORS HAVING NEUROENDOCRINE FEATURES AND ANALYSIS OF MARKER EXPRESSION USING WHOLE TRANSCRIPTOME SEQUENCING

[0513] Neuroendocrine tumors (NETs) arising from the dispersed endocrine system are rare, with an incidence of 2-5 per 100,000 people, but highly aggressive. Neuroendocrine tumors occur in the adrenal gland, kidney, genitourinary tract (bladder, prostate, ovary, cervix, and endometrium), pancreas, gastrointestinal tract (stomach and colon), thyroid (medullary thyroid cancer), and lung (small cell lung carcinoma, large cell neuroendocrine carcinoma, and carcinoid). These tumors may secrete several hormones including serotonin and/or chromogranin A that can cause debilitating

symptoms known as carcinoid syndrome. These tumors can be denoted by positive immunohistochemical markers such as neuron-specific enolase (NSE, also known as gamma enolase, gene symbol = ENO2), CD56/NCAM1, and synaptophysin. Traditional chemotherapies have not been successful in treating NETs, and mortality due to metastatic spread is a common outcome. Unfortunately, in most cases surgery is the only potential curative treatment, provided it takes place following early detection and prior to tumor metastasis. In this context work was undertaken to identify novel therapeutic targets associated with tumors comprising neuroendocrine features.

[0514] To identify and characterize such tumors as they exist in cancer patients a large non-traditional xenograft (NTX) tumor bank was developed and maintained using art-recognized techniques. The NTX tumor bank, comprising a substantial number of discrete tumor cell lines, was propagated in immunocompromised mice through multiple passages of heterogeneous tumor cells originally obtained from numerous cancer patients afflicted by a variety of solid tumor malignancies. (Note that in some of the Examples and FIGS. herein the passage number of the tested sample is indicated by p0-p# appended to the sample designation where p0 is indicative of an unpassaged sample obtained directly from a patient tumor and p# is indicative of the number of times the tumor has been passaged through a mouse prior to testing.) The continued availability of a large number of discrete early passage NTX tumor cell lines having well defined lineages greatly facilitate the identification and characterization of cells purified from the cell lines. In such work the use of minimally passaged NTX cell lines simplifies *in vivo* experimentation and provides readily verifiable results. Moreover, early passage NTX tumors respond to therapeutic agents such as irinotecan (i.e. Camptosar®) and Cisplatin/Etoposide regimens, which provides clinically relevant insights into underlying mechanisms driving tumor growth, resistance to current therapies and tumor recurrence.

[0515] As the NTX tumor cell lines were established, their phenotype was characterized in various ways to examine global gene expression. To identify which NTX lines in the bank might be NETs, gene expression profiles were generated by whole transcriptome sequencing and/or microarray analysis. Specifically, the data was examined to identify tumors expressing high levels of specific genes known to be elevated in NETs or used as histochemical markers of neuroendocrine differentiation (e.g., ASCL1, NCAM1, CHGA) as well as tumors with changes in NOTCH pathway genes indicative of suppression of NOTCH signaling (e.g., reduced levels of NOTCH receptors, and changes to ligands and effector molecules).

[0516] More particularly, upon establishing various NTX tumor cell lines as is commonly done for human tumors in severely immune compromised mice, the tumors were resected after reaching 800 - 2,000 mm³ and the cells were dissociated and dispersed into suspension using art-recognized enzymatic digestion techniques (see, for example, U.S.P.N. 2007/0292414 which is incorporated herein). The dissociated cell preparations from these NTX lines were then depleted of murine cells, and human tumor cell subpopulations were then further isolated by fluorescence activated cell sorting and lysed in RLTplus RNA lysis buffer (Qiagen). These lysates were then stored at -80°C until used. Upon thawing, total RNA was extracted using a RNeasy isolation kit (Qiagen) following the vendor's instructions and quantified on a Nanodrop spectrophotometer (Thermo Scientific) and a Bioanalyzer 2100 (Agilent Technologies) again using the manufacturer's protocols and recommended instrument settings. The resulting total RNA preparations were suitable for genetic sequencing and gene expression analysis.

[0517] Whole transcriptome sequencing using an Applied Biosystems (ABI) SOLiD (Sequencing by Oligo Ligation/Detection) 4.5 or SOLiD 5500xl next generation sequencing system (Life Technologies) was performed on RNA samples from NTX lines. cDNA was generated from total RNA samples using either a modified whole transcriptome (WT) protocol from ABI designed for low input total RNA or Ovation RNA-Seq System V2™ (NuGEN Technologies Inc.). The modified low input WT protocol uses 1.0 ng of total RNA to amplify mRNA at the 3' end which leads to a heavy 3' bias of mapped gene expression, while NuGen's system allows for a more consistent amplification throughout the transcript, and includes amplification of both mRNA and non-polyadenylated transcript cDNA using random hexamers. The cDNA library was fragmented, and barcodes adapters were added to allow pooling of fragment libraries from different samples.

[0518] ABI's SOLiD 4.5 and SOLiD 5500xl next generation sequencing platforms enables parallel sequencing of transcriptomes from multiple NTX lines and sorted populations. A cDNA library is constructed from each RNA sample, which is fragmented and barcoded. Barcodes on each fragment library allow multiple samples to be pooled at equal concentrations and run together while ensuring sample specificity. The samples are taken through emulsion PCR using ABI's SOLiD™ EZ Bead™ robotics system, which ensures sample consistency. Paired-end sequencing generates a 50 base read in the 5' to 3' direction and a 25 base read in the 3' to 5' direction for each clonally amplified fragment on a single bead that exists in the pool. In the case of the 5500xl platform, for every set of 8 samples pooled in the method mentioned above, beads are evenly deposited into 6 single channel lanes on a single chip. This will, on average, generate more than 50 million 50 base reads and 50 million 25 base reads for each of the 8 samples and generates a very accurate representation of mRNA transcript level in the tumor cells. Data generated by the SOLiD platform mapped to 34,609 genes as annotated by RefSeq version 47 using NCBI version hg19.2 of the published human genome and provided verifiable measurements of RNA levels in most samples.

[0519] The SOLiD platform is able to capture not only expression, but SNPs, known and unknown alternative splicing events, small non-coding RNAs, and potentially new exon discoveries based solely on read coverage (reads mapped

uniquely to previously un-annotated genomic locations). Thus, use of this next generation sequencing platform paired with proprietary data analysis and visualization software thus allowed for discovery of differential transcript expression as well as differences and/or preferences for specific splice variants of expressed mRNA transcripts. Sequencing data from the SOLiD platform is nominally represented as a transcript expression value using the metrics RPM (reads per million) and RPKM (read per kilobase per million), enabling basic differential expression analysis as is standard practice.

[0520] Whole transcriptome sequencing of four small cell lung cancer (SCLC) tumors (LU73, LU64, LU86 and LU95), one ovarian tumor (OV26) and a large cell neuroendocrine carcinoma (LCNEC; LU37) resulted in the determination of gene expression patterns commonly found in NETs (FIG. 6A). More specifically, these tumors had high expression of several NET markers (ASCL1, NCAM1, CHGA) as well as reduced levels of Notch receptors and effector molecules (e.g., HES1, HEY1) and elevated markers of Notch suppression (e.g., DLL3 and HES6). In contrast, 4 normal lung samples, 3 lung adenocarcinoma tumors (LU137, LU146 and LU153), and 3 squamous cell lung carcinomas (LU49, LU70 and LU76) all have expression of various Notch receptors and effector molecules, and do not show elevated expression of Notch suppressors such as HES6 and DLL3.

[0521] Moreover, as seen in FIG. 6B, an analysis of the whole transcriptome data comparing normal tissue samples to various lung NTX populations having neuroendocrine features, showed that SEZ6 was up-regulated at the mRNA transcript level in four lung cancer populations having neuroendocrine features (LU73, LU64, LU86 and LU95) compared to extremely low or no transcript expression in the normal tissues tested. These results suggest that SEZ6 may play an important role in the tumorigenesis and maintenance of particular cancers (including lung cancers with neuroendocrine features). On this basis, SEZ6 was selected for further analysis as a potential immunotherapeutic target

EXAMPLE 2

MICROARRAY AND RT-PCR ANALYSIS OF GENE EXPRESSION IN SELECTED NTX TUMORS WITH NEUROENDOCRINE FEATURES

[0522] In an effort to identify additional NETs in the aforementioned NTX bank beyond those for which SOLiD whole transcriptome data existed, a larger set of NTX lines was examined using microarray analysis. Specifically, 2-6 μ g of total RNA samples derived from whole tumors in 46 NTX lines or from 2 normal tissues were analyzed using a OneArray[®] microarray platform (Phalanx Biotech Group), which contains 29,187 probes designed against 19,380 genes in the human genome. More specifically, RNA samples were obtained (as described in Example 1) from forty-six patient derived whole NTX tumors comprising colorectal (CR), melanoma (SK), kidney (KD), lung (LU), ovarian (OV), endometrial (EM), breast (BR), liver (LIV), or pancreatic (PA) cancers. Normal colorectal (NormCR) and normal pancreas (NormPA) tissues were used as controls. Still more specifically, lung tumors were further subclassified as small cell lung cancers (SCLC), squamous cell cancers (SCC), or large cell neuroendocrine carcinoma (LCNEC). RNA samples were run in triplicate using the manufacturer's protocols and the resulting data was analyzed using standard industry practices for normalizing and transforming the measured intensity values obtained for the subject gene in each sample. An unbiased Pearson Spearman hierarchical clustering algorithm in the R/BioConductor suite of packages called hclust.2 was used to create a standard microarray dendrogram for these 48 samples. As known in the art R/BioConductor is an open-source, statistical programming language widely used in academia, finance and the pharmaceutical industry for data analysis. Generally the tumors were arranged and clustered based on gene expression patterns, expression intensity, etc.

[0523] As shown in FIG. 7A, the dendrogram derived from the 48 samples and across all 19380 genes, clustered NTX lines together based upon their tumor type or tissue of origin. Several tumors typically associated with neuroendocrine phenotypes clustered together on the branch denoted by (1); these included skin cancers, numerous lung cancers and other NETs. Interestingly, a sub-branch, denoted by (2), showed that two large cell lung cancers with neuroendocrine features (LU50.LCNEC and LU37.LCNEC) and a small cell lung cancer (LU102.SCLC) clustered with an ovarian (OV26) and a kidney (KD66) tumor (cluster C) suggesting these later tumors also possessed neuroendocrine phenotypes. Moreover, FIG. 7A shows cluster D which consists of 3 additional SCLC tumors, and to its right is a small cluster (cluster E) containing an additional SCLC tumor (LU100) and a neuroendocrine endometrial tumor (EM6). All of the tumors in clusters D and E are generally understood to possess some neuroendocrine features based on the academic literature and pathology experience in the clinic. The fact that cluster G, comprising SCC, can be found on a completely different branch of the dendrogram of FIG. 7A, indicates that the clustering is not driven exclusively by the organ of origin of the tumor.

[0524] Closer inspection of a collection of gene markers associated with NETs (FIG. 7B) shows that they are strongly expressed in tumors comprising clusters C and D, while they are minimally expressed in tumors in Cluster G (squamous cell carcinoma of the lung), suggesting clusters C and D represent NETs or tumors with a neuroendocrine phenotype. More specifically, cluster C NETs highly express ASCL1, CALCA, CHGA, SST and NKX2-1, while cluster D NETs highly express CHGA, ENO2, and NCAM1, and it is the expression of these neuroendocrine phenotype genes that is in part responsible for the clustering of these tumors. An interesting feature is the strong expression of KIT in cluster D, a gene

occasionally reported to be associated with neuroendocrine tumors, but clearly linked to oncogenesis in other contexts. This is in contrast to the SCC tumors in cluster G which lack strong expression any of these genes (FIG. 7B).

[0525] Tumors in cluster C show a phenotype consistent with a reduction in Notch signaling: a lack of expression of any Notch receptor, a relative lack of JAG1 and HES1 expression, and strong levels of ASCL1 expression (FIG. 7C). Interestingly, cluster D shows high expression of HES6, a transcription factor that can support ASCL1 activity by antagonizing HES1 activity through heterodimer formation.

[0526] In view of the aforementioned results, mRNA expression of HES6 was examined from various NTX lines and normal tissues using an Applied Biosystems 7900HT Machine (Life Technologies) to perform Taqman real-time quantitative RT-PCR (qRT-PCR) in accordance with the manufacturer's protocols. RNA was isolated as described above and checked to ensure quality was suitable for gene expression analysis. RNA from normal tissues was purchased (Agilent Technologies and Life Technologies). 200 ng of RNA was used for cDNA synthesis using the cDNA archive kit (Life Technologies). cDNA was used for qRT-PCR analysis on Taqman Low Density Arrays (TLDA; Life Technologies) which contained the HES6 Taqman assay to measure mRNA levels of HES6.

[0527] HES6 mRNA levels are shown for each NTX line or normal tissue sample (single dot on graph) after normalization to endogenous controls. Normalized values are plotted relative to the average expression in the normal tissues of toxicity concern (NormTox). This technique allowed for the rapid identification and characterization of a variety of tumors having neuroendocrine features from the NTX tumor bank through measurement of HES6 and other relevant markers. FIG. 7D illustrates general overexpression of HES6 in the sampled tumors with neuroendocrine features (e.g., LU-SCLC, LU-LCNEC) compared to normal tissues, breast, colon, liver and other selected tumors. Significantly these microarray and qPCR data show that at least some endometrial, kidney and ovarian tumors may exhibit neuroendocrine tumor features (FIGS. 7A and 7D).

[0528] The microarray data generated as described above not only showed that the tumors in clusters C, D and E exhibited various neuroendocrine markers, but also showed that the tumors in those clusters expressed markers indicative of neurogenesis, neural commitment, or differentiation towards neural fates (FIG. 7E). Of particular interest, the tumors in Cluster D frequently show a stronger and more consistent upregulation of many of these markers (e.g., BEX1 and BEX4, CD56, NRCAM, SEMA receptors, SOX and ZIC factors) and frequently reduced hormone upregulation versus other clusters, suggesting a more neural phenotype. Remarkably, the tumors in those same clusters also showed high levels of SEZ6 transcript, suggesting that SEZ6 is associated with tumors having neuroendocrine and neural features (FIG. 7F).

EXAMPLE 3

EXPRESSION OF SEZ6 MRNA IN TUMORS HAVING NEUROENDOCRINE AND NEURAL FEATURES

[0529] Various techniques were used to identify tumors exhibiting neuroendocrine features including whole transcriptome sequencing (Example 1) and microarray and qRT-PCR (Example 2). The data thus generated was further analyzed in order to find potential therapeutic targets that are highly expressed in neuroendocrine tumors when compared to non-neuroendocrine tumors and normal tissues. As discussed in Example 1 it was found that SEZ6, a single pass transmembrane protein that is mainly expressed in the normal brain, has high expression in many neuroendocrine tumors (FIG. 6B).

[0530] The microarray data generated in Example 2 showed that tumors located in clusters C, D and E expressed neuroendocrine markers (FIG. 7B) and neural markers (FIG. 7E). The tumors in those same clusters also showed high levels of SEZ6 transcript, suggesting that SEZ6 is associated with tumors having neuroendocrine and neural features (FIG. 7F). This is in line with the known role of SEZ6 in postnatal forebrain development and continued expression in the specific regions of the hippocampus in the adult. SEZ6 is thought to play important roles in cell-cell recognition and signaling. Often developmental pathways are inappropriately expressed in tumors.

[0531] In order to determine SEZ6 mRNA expression levels in various sample NTX tumor lines, qRT-PCR was performed using the SEZ6 Taqman assay essentially as described in Example 2 above. FIG. 8A shows SEZ6 expression relative to the average expression in normal tissues and normalized to expression of the endogenous control gene ALAS1. SEZ6 gene expression is elevated more than 10,000,000-fold in neuroendocrine NTX populations versus normal tissues. Five of the SCLC NTX lines shown in FIG. 8A are mRNA samples extracted directly from primary biopsies (p0). The expression of SEZ6 in these unpassaged tumors demonstrates that SEZ6 expression is not an artifact that results from growing human tumors in mice. Three subtypes of NSCLC are also represented in FIG. 8A: LU25 is spindle cell carcinoma, LU50 is a large cell neuroendocrine carcinoma (LCNEC), and LU85 is a squamous cell carcinoma (SCC). KDY66 and OV26, a kidney and ovarian tumor, respectively, clustered on the microarray with SCLC and LCNEC tumors (FIG. 7A), suggesting they also have neuroendocrine features.

[0532] To extend the analysis of SEZ6 expression to a wider array of tumor specimens, qRT-PCR was performed using the Fluidigm BioMark™ HD System. Briefly, 1 ng of RNA, prepared as described in Example 1, was converted to

cDNA using the cDNA archive kit (Life Technologies). The cDNA was pre-amplified using a SEZ6 specific Taqman assay and was then used to perform qRT-PCR. Expression in normal tissues (NormTox or Norm) was compared to expression in the following NTX lines, where the number in brackets indicates the number of unique NTX lines tested: BR (13), CR (24), KDY (10), OV (35), PA (21), lung adenocarcinoma (LU-Adeno) (23), LU-CAR (1), LU-LCC (8), SCC (12), SCLC (30) and medullary thyroid cancer (THY) (1) (FIG. 8B). SCLC, THY LU-CAR and LC-LCC NTX show the highest expression of SEZ6, although some expression of SEZ6 was also seen in OV, PA, BR, CR and LU-Adeno NTX lines compared to normal tissue samples.

[0533] "NormTox" represents the following samples of normal tissue: one adrenal gland, four colon, four kidney, four liver, three lung, three pancreas, three heart, three esophagus, one skeletal muscle, one skin, two dermal fibroblasts, two keratinocytes, three small intestine, one spleen, two stomach, and two trachea samples. Another set of normal tissues designated "Norm" represents the following samples of normal tissue: adipose, B cells, bladder, brain, breast, cervix, melanocytes, monocytes, NK cells, ovary, peripheral blood mononuclear cells, placenta, prostate, salivary gland, T cells, testes, thymus, and thyroid. Most normal tissues have no expression of SEZ6, while low expression is seen in pancreas, colon, liver and lung and high expression in brain. A different SEZ6 specific Taqman assay, using essentially the same method as above, was conducted on various NTX tumor lines. The number of tumor lines that were tested for each type of tumor is denoted as the denominator, whereas the number of tumors that expressed SEZ6 is denoted as the numerator: 2/6 CR, 2/4 GA, 1/1 GB (glioblastoma), 1/1 KDY, 2/6 SK, 2/5 LU-Adeno, 4/4 LCNEC, 2/7 LU-SCC, 9/9 SCLC and 1/2 OV (data not shown).

[0534] Taken together, these data suggest that SEZ6 is upregulated in tumors exhibiting neuroendocrine and neural features suggesting it may serve as a therapeutic target for treatment of these types of tumors.

EXAMPLE 4

EXPRESSION OF SEZ6 MRNA IN VARIOUS TUMOR AND NORMAL TISSUE SPECIMENS USING QRT-PCR

[0535] To extend the analysis of SEZ6 expression to a wider array of tumor specimens, Taqman qRT-PCR was performed substantially as described in the previous Examples on a TissueScan™ qPCR (Origene Technologies) 384-well array. This array enables comparison of gene expression across 18 different solid tumor types, with multiple patient derived samples for each tumor type and from normal adjacent tissue.

[0536] In this regard, FIGS. 9A and 9B show the absolute and relative gene expression levels, respectively, of SEZ6 in whole tumor specimens (grey dots) or normal adjacent tissue (NAT; white dots) from patients with one of eighteen different solid tumor types. More specifically, FIG. 9A shows the absolute mRNA expression level of SEZ6 in various whole tumor specimens or matched normal adjacent tissue. FIG. 9B shows the expression level of SEZ6 as normalized against β -action and plotted relative to expression in normal adjacent tissue for each tumor type analyzed. Specimens in which SEZ6 was not detected were assigned a Ct value of 50, which represents the last cycle of amplification in the experimental protocol. Each dot represents a single tissue specimen, with the geometric mean value represented as a black line.

[0537] Using this Origene Array, overexpression of SEZ6 was seen in a subset of adrenal, liver, lung, ovarian, and pancreatic cancer, many of which may represent neuroendocrine tumors or tumors with poorly differentiated neuroendocrine phenotypes.. This includes high expression in 1/1 medullary thyroid cancer and 1/3 papillary carcinoma of the thyroid follicular variant, 8/8 neuroendocrine pancreatic tumors, 4/4 pancreatic islet cell tumors, 1/2 large cell neuroendocrine lung carcinomas, and 3/3 lung carcinoid tumors. As shown by the absolute gene expression in FIG. 9A, normal testis and pancreas are the only normal tissues with high expression of SEZ6. This suggests that SEZ6 may play a role in tumorigenesis and/or tumor progression in a wide variety of tumors including but not limited to those with neuroendocrine and neural features.

EXAMPLE 5

CLONING AND EXPRESSION OF RECOMBINANT SEZ6 PROTEINS

Human SEZ6 (hSEZ6)

[0538] The amino acid sequences of the SEZ6 isoform 1 (SEQ ID NO: 3) and SEZ6 isoform 2 (SEQ ID NO: 4) proteins are set out in FIGS. 1C and 1D, respectively. The extracellular domain of each of the SEZ6 isoforms is identical. In FIGS. 1C and 1D the leader sequence, comprising nineteen amino acids, is in bold and underlined. The rest of the amino acid residues of each of SEQ ID NO: 3 and SEQ ID NO: 4 comprises the mature SEZ6 protein. In order to generate and develop all molecular and cellular materials required in the present invention pertaining to human SEZ6, cDNA (FIG. 3A; SEQ ID NO: 5) encoding the complete mature human SEZ6 protein (FIG. 3B, SEQ ID NO: 6) was created as follows.

A commercial human SEZ6 cDNA was purchased from Open Biosystems where this cDNA sequence corresponded to NCBI accession BC146292. Sequence alignments showed the protein encoded by BC146292 differed by several residues from that of RefSeq NP_849191 (see residues 414, 415 and 417, FIG. 3C), encoding the endogenous human SEZ6 protein. PCR was used to amplify two separate cDNA fragments from the BC 146292 clone, in which the primers used introduced the desired changes into the cDNA at residues 414-417 during the process of overlap PCR to create a cDNA encoding a mature SEZ6 protein with identical sequence to that encoded by NM_178860, the mRNA sequence encoding endogenous human SEZ6 protein. The repaired cDNA clone, termed hSEZ6 (FIG. 3A), was used for all subsequent engineering of constructs expressing the mature human SEZ6 protein or fragments thereof.

[0539] In order to generate immunoreactive or immunospecific modulators to the SEZ6 molecule, a chimeric fusion gene was generated in which the ECD portion of the human SEZ6 protein was fused to the human IgG2 Fc domain (FIG. 4A, SEQ ID NO: 8). This was done as follows: cDNA encoding the ECD of SEZ6 was PCR amplified from the hSEZ6 cDNA clone (FIG. 3A), and this PCR product was subsequently subcloned into a CMV driven expression vector in frame and downstream of an IgK signal peptide sequence and upstream of a human IgG2 Fc cDNA, using standard molecular techniques. The cDNA sequence encoding the hSEZ6-Fc fusion protein, termed hSEZ6-Fc ORF, is shown in FIG. 4A; the corresponding protein sequence encoded by hSEZ6-Fc ORF is shown in FIG. 4B (SEQ ID NO: 9). The underlined regions of the sequences correspond to the human IgG2 Fc. The bolded underlined regions correspond to the IgK signal peptide, and the sequences in bold font correspond to the portions of the fusion protein encoded by the cloning restriction sites flanking the SEZ6 ECD.

[0540] To generate recombinant hSEZ6 ECD protein a similar PCR-based strategy was used. The cDNA fragment encoding the ECD of SEZ6 was amplified from the hSEZ6 cDNA clone and subcloned into a CMV driven expression vector in frame and downstream of an IgK signal peptide sequence and in frame upstream of a sequence encoding a 9-Histidine epitope tag (SEQ ID NO: 201).

[0541] The CMV-driven expression vector permits high level transient expression in HEK-293T and/or CHO-S cells. Suspension or adherent cultures of HEK-293T cells, or suspension CHO-S cells were transfected with expression constructs encoding either the hSEZ6 ECD-Fc or hSEZ6-ECD-His proteins, using polyethylenimine polymer as the transfecting reagent. Three to five days after transfection, the hSEZ6 ECD-Fc or hSEZ6-ECD-His proteins were purified from clarified cell-supernatants using an AKTA explorer and either MabSelect SuRe™ Protein A (GE Healthcare Life Sciences) or Nickel-EDTA (Qiagen) columns, respectively.

[0542] A stable cell line overexpressing recombinant human SEZ6 was constructed using lentiviral vectors to transduce HEK-293T cells as follows: PCR amplification was performed using the human SEZ6 clone as a template in order to produce a cDNA fragment encoding the mature human SEZ6 protein. The fragment that was generated was subcloned in frame downstream of a sequence encoding an IgK signal peptide and DDK epitope tag previously engineered upstream of the multiple cloning site of pCDH-EF1-MCS-T2A-GFP (System Biosciences) using standard molecular cloning techniques. The resulting bicistronic lentiviral vector was used to engineer cell lines overexpressing a human SEZ6-T2A peptide-GFP polypeptide. The T2A sequence promotes ribosomal skipping of a peptide bond condensation, resulting in two independent proteins, in this case SEZ6 and GFP.

Mouse SEZ6 (mSEZ6)

[0543] A stable cell line overexpressing recombinant mouse SEZ6 was engineered essentially as described above for recombinant human SEZ6. HEK-293T cells were transduced with a lentiviral vector expressing murine SEZ6. The vector was engineered essentially as follows. A cDNA fragment (FIG. 5A; SEQ ID NO: 10) encoding the mature mouse SEZ6 protein listed as RefSeq NM_021286 in the NCBI database (FIG. 5B; SEQ ID NO: 11) was obtained by PCR amplification from a commercial mouse SEZ6 cDNA (Origene; #MC203634) and subcloned downstream of an IgK signal peptide sequence and DDK epitope tag sequence previously engineered upstream of the multiple cloning site of pCDH-EF1-MCS-IRES-RFP (System Biosciences) using standard molecular cloning techniques. This yielded a bicistronic lentiviral vector that was used to produce a HEK-293T cell line overexpressing mouse SEZ6 and RFP.

Rat SEZ6 (rSEZ6)

[0544] To generate and develop all molecular and cellular materials required in the present invention pertaining to rat SEZ6 proteins, cDNA (FIG. 5C, SEQ ID NO: 12) encoding the complete mature rat SEZ6 protein (FIG. 5D, SEQ ID NO: 13) was obtained as follows. A cDNA encoding the full length mature rat protein (i.e., the full length protein minus the wild-type signal peptide) was amplified from rat brain marathon-ready cDNA (Clontech #639412). Sequence alignments showed the ECD of the encoded protein to be homologous to the endogenous rat SEZ6 protein listed as RefSeq NP_001099224 in the NCBI database. This cDNA clone, termed rSEZ6 (FIG. 5D), was used for subsequent engineering of constructs expressing the rat SEZ6 protein fragments.

Cynomolgus SEZ6 (cSEZ6)

[0545] To generate and develop all molecular and cellular materials required in the present invention pertaining to cynomolgus SEZ6 proteins, cDNA (FIG. 5E, SEQ ID NO: 14) encoding the cynomolgus SEZ6 protein (FIG. 5F, SEQ ID NO: 15) was obtained as follows: A predicted cynomolgus SEZ6 ORF sequence was assembled by bioinformatics analysis in the following way: the ORF of the human SEZ6 gene was obtained from NCBI accession NM_178860 and compared, using the BLAST algorithm, to the whole genome shotgun sequencing contigs in the NCBI database. The BLAST results were then used to assemble a putative cynomolgus SEZ6 ORF. The sequence encoding the predicted wild-type signal peptide of cynomolgus SEZ6 was removed from this BLAST derived sequence, and replaced with a sequence encoding an IgK signal peptide sequence. After codon optimization for production in mammalian cells, this complete hybrid ORF sequence was ordered as a synthetic gene (GeneWiz). This optimized cDNA clone, termed cSEZ6 (FIG. 5E), was used for subsequent engineering of constructs expressing the cynomolgus SEZ6 protein fragments.

Human SEZ6L and SEZ6L2

[0546] In the human genome, there are two genes closely related to SEZ6- seizure related 6 homolog-like (SEZ6L) and seizure related 6 homolog like-2 (SEZ6L2). Although the overall percent identity is relatively low between the three proteins (~42%), there are smaller stretches of perfect identity between pairs or all three of the proteins. In order to investigate any possible cross reactivity of the anti-SEZ6 modulators with human SEZ6L and SEZ6L2 proteins, the open reading frames encoding the ECDs of human SEZ6L protein (NP_0066938) and human SEZ6L2 protein (NP_001230261) were codon optimized and synthesized (GeneWiz). These optimized cDNA sequences encoding the ECDs of human SEZ6L or SEZ6L2 proteins are shown in FIGS. 5G and 5I.

Material for cross-reactivity studies

[0547] Material was generated in order to study whether the SEZ6 modulators of the invention cross-reacted with rat and/or cynomolgus SEZ6 homologues, or with the closely related human SEZ6L and SEZ6L2 proteins. Chimeric fusion genes were designed in which the ECD portion of either the rat or the cynomolgus SEZ6 protein (underlined in FIGS. 5D and 5F, respectively) was fused to a 9-Histidine epitope tag (SEQ ID NO: 201). Using PCR, the cDNA fragment encoding the ECD of either rat or cynomolgus SEZ6 was amplified from either rSEZ6 or cSEZ6, respectively, and subcloned into a CMV driven expression vector in frame and downstream of an IgK signal peptide sequence and in frame and upstream of a sequence encoding a 9-Histidine epitope tag (SEQ ID NO: 201). Similarly, chimeric fusion genes were designed in which the open reading frame encoding the ECD portion of the human SEZ6L or SEZ6L2 proteins was subcloned into a CMV driven expression vector in frame and downstream of an IgK signal peptide sequence and in frame and upstream of a sequence encoding a 9-Histidine epitope tag (SEQ ID NO: 201). The resultant encoded protein sequences for these fusion proteins are shown in FIGS. 5H and 5J, respectively, with the underlined sequence representing the ECD of the particular protein under consideration.

[0548] The rat and cynomolgus SEZ6 ECD-His vectors generated above, were used to produce and purify recombinant rSEZ6-ECD-His protein and cSEZ6-ECD-His protein, respectively, as follows: using art-recognized techniques, suspension or adherent cultures of HEK-293T cells, or suspension CHO-S cells were transfected with the expression vectors encoding rSEZ6-ECD-His or cSEZ6-ECD-His protein. Polyethylenimine polymer was used as the transfecting reagent. Three to five days after transfection, the rSEZ6-ECD-His or cSEZ6-ECD-His protein was purified from clarified cell-supernatants using AKTA explorer and Nickel-EDTA (Qiagen) columns. Similarly, the human SEZ6L and SEZ6L2 ECD-His vectors were used to produce and purify recombinant human SEZ6L and human SEZ6L2 ECD-His proteins, as described for the rat and cynomolgus homologs.

EXAMPLE 6**GENERATION OF ANTI-SEZ6 MURINE MODULATORS**

[0549] SEZ6 modulators in the form of murine antibodies were produced in accordance with the teachings herein through inoculation with human SEZ6-Fc. In this regard three strains of mice were used to generate high affinity, murine, monoclonal antibody modulators that can be used to associate with and/or inhibit the action of SEZ6 for the prevention and/or treatment of various proliferative disorders. Specifically, Balb/c, CD-1 and FVB mouse strains were immunized with human recombinant SEZ6-Fc and used to produce Hybridomas.

[0550] The SEZ6-Fc antigen was purified from supernatant from CHO-S cells over expressing the construct SEZ6-Fc as set forth in Example 5 (FIGS. 4A and 4B). 10 µg of SEZ6-Fc immunogen was used for the first immunization, followed by 5 µg and 2.5 µg of SEZ6-Fc immunogen for the subsequent three immunizations and five immunizations, respectively.

All immunizations were performed with the immunogen emulsified with an equal volume of TITERMAX® Gold (CytRx Corporation) or alum adjuvant. Murine antibodies were generated by immunizing six female mice (two each of: Balb/c, CD-1, FVB) via footpad route for all injections.

[0551] Solid-phase ELISA assays were used to screen mouse sera for mouse IgG antibodies specific for human SEZ6. A positive signal above background was indicative of antibodies specific for SEZ6. Briefly, 96 well plates (VWR International, Cat. #610744) were coated with recombinant SEZ6-His at 0.5 µg/ml in ELISA coating buffer overnight. After washing with PBS containing 0.02% (v/v) Tween 20, the wells were blocked with 3% (w/v) BSA in PBS, 200 µL/well for 1 hour at room temperature (RT). Mouse serum was titrated (1:100, 1:200, 1:400, and 1:800) and added to the SEZ6 coated plates at 50 µL/well and incubated at RT for 1 hour. The plates are washed and then incubated with 50 µL/well HRP-labeled goat anti-mouse IgG diluted 1:10,000 in 3% BSA-PBS or 2% FCS in PBS for 1 hour at RT. Again the plates were washed and 40 µL/well of a TMB substrate solution (Thermo Scientific 34028) was added for 15 minutes at RT. After developing, an equal volume of 2N H₂SO₄ was added to stop substrate development and the plates were analyzed by spectrophotometer at OD 450.

[0552] Sera-positive immunized mice were sacrificed and draining lymph nodes (popliteal and inguinal, and medial iliac if enlarged) were dissected out and used as a source for antibody producing cells. A single cell suspension of B cells (228.9x10⁶ cells) was fused with non-secreting P3x63Ag8.653 myeloma cells (ATCC #CRL-1580) at a ratio of 1:1 by electrofusion. Electrofusion was performed using the BTX Hybrimmune™ System, (BTX Harvard Apparatus) as per the manufacturer's directions. After the fusion procedure the cells were resuspended in hybridoma selection medium supplemented with Azaserine (Sigma #A9666), high glucose DMEM medium with sodium pyruvate (Cellgro cat#15-017-CM) containing 15% Fetal Clone I serum (Hyclone), 10% BM Condimed (Roche Applied Sciences), 4 mM L-glutamine, 100 IU Penicillin-Streptomycin and 50 µM 2-mercaptoethanol and then plated in three T225 flasks in 90 mL selection medium per flask. The flasks were then placed in a humidified 37°C incubator containing 5% CO₂ and 95% air for 6-7 days.

[0553] After six to seven days of growth the library consisting of the cells grown in bulk in the T225s was plated at 1 cell per well in Falcon 96 well U-bottom plates using the Aria I cell sorter. The selected hybridomas were then grown in 200 µL of culture medium containing 15% Fetal Clone I serum (Hyclone), 10% BM-Condimed (Roche Applied Sciences), 1 mM sodium pyruvate, 4 mM L-glutamine, 100 IU Penicillin-Streptomycin, 50 µM 2-mercaptoethanol, and 100 µM hypoxanthine. Any remaining unused hybridoma library cells were frozen for future library testing. After ten to eleven days of growth supernatants from each well of the plated cells were assayed for antibodies reactive for SEZ6 by ELISA and FACS assays.

[0554] For screening by ELISA 96 well plates were coated with SEZ6-Fc at 0.3 µg/mL in PBS overnight at 4°C. The plates were washed and blocked with 3% BSA in PBS/Tween for one hour at 37°C and used immediately or kept at 4°C. Undiluted hybridoma supernatants were incubated on the plates for one hour at RT. The plates were washed and probed with HRP labeled goat anti-mouse IgG diluted 1:10,000 in 3% BSA-PBS for one hour at RT. The plates were then incubated with substrate solution as described above and read at OD 450. Wells containing immunoglobulin that preferentially bound human SEZ6, as determined by a signal above background, were transferred and expanded.

[0555] Selected growth positive hybridoma wells secreting murine immunoglobulin were also screened for human SEZ6 specificity and cynomolgus, rat and murine SEZ6 cross reactivity using a flow cytometry based assay with 293 cells engineered to over-express the selected antigen or constructs fabricated in the previous Example.

[0556] For the flow cytometry assays, 50x10⁴ h293 cells transduced respectively with human, cynomolgus, rat or murine SEZ6 were incubated for 30 minutes with 25-100 µL hybridoma supernatant. Cells were washed with PBS, 2% FCS, twice and then incubated with 50 µL of a goat-anti-mouse IgG Fc fragment specific secondary conjugated to DyLight 649 diluted 1:200 in PBS/2%FCS. After 15 minutes of incubation, cells were washed twice with PBS, 2%FCS, and re-suspended in the same buffer with DAPI and analyzed by flow cytometry using a FACSCanto II as per the manufacturer's instructions. Wells containing immunoglobulin that preferentially bound the SEZ6⁺ GFP⁺ cells were transferred and expanded. The resulting hSEZ6 specific clonal hybridomas were cryopreserved in CS-10 freezing medium (Biolife Solutions) and stored in liquid nitrogen. Antibodies that bound with human, cynomolgus, rat or murine SEZ6 cells were noted as cross-reactive (see FIG. 11A).

[0557] ELISA and flow cytometry analysis confirmed that purified antibody from most or all of these hybridomas bound SEZ6 in a concentration-dependent manner. Wells containing immunoglobulin that bound SEZ6 GFP cells were transferred and expanded. The resulting clonal hybridomas were cryopreserved in CS-10 freezing medium (Biolife Solutions) and stored in liquid nitrogen.

[0558] One fusion was performed and seeded in 48 plates (4608 wells at approximately 40% cloning efficiency). The initial screen yielded sixty-three murine antibodies that associated with human SEZ6. A second screen was subsequently performed and yielded 134 antibodies that associated with human SEZ6.

EXAMPLE 7

SEQUENCING OF SEZ6 MURINE MODULATORS

[0559] Based on the foregoing, a number of exemplary distinct monoclonal antibodies that bind immobilized human SEZ6 or h293-hSEZ6 cells with apparently high affinity were selected for sequencing and further analysis. As shown in a tabular fashion in FIGS. 10A and 10B, sequence analysis of the light chain variable regions (FIG. 10A) and heavy chain variable regions (FIG. 10B) from selected monoclonal antibodies generated in Example 6 confirmed that many had novel complementarity determining regions and often displayed novel VDJ arrangements. Note that the complementarity determining regions set forth in FIGS. 10A and 10B are defined as per Kabat et al., supra.

[0560] As a first step in sequencing exemplary modulators, the selected hybridoma cells were lysed in Trizol® reagent (Trizol Plus RNA Purification System, Life Technologies) to prepare the RNA. In this regard between 10^4 and 10^5 cells were resuspended in 1 mL Trizol and shaken vigorously after addition of 200 μ L of chloroform. Samples were then centrifuged at 4°C for 10 minutes and the aqueous phase was transferred to a fresh microfuge tube where an equal volume of isopropanol was added. The tubes were again shaken vigorously and allowed to incubate at RT for 10 minutes before being centrifuged at 4°C for 10 minutes. The resulting RNA pellets were washed once with 1 mL of 70% ethanol and dried briefly at RT before being resuspended in 40 μ L of DEPC-treated water. The quality of the RNA preparations was determined by fractionating 3 μ L in a 1% agarose gel before being stored at -80°C until used.

[0561] The variable region of the Ig heavy chain of each hybridoma was amplified using a 5' primer mix comprising thirty-two mouse specific leader sequence primers, designed to target the complete mouse V_H repertoire, in combination with a 3' mouse C_γ primer specific for all mouse Ig isotypes. A 400 bp PCR fragment of the V_H was sequenced from both ends using the same PCR primers. Similarly a mix of thirty-two 5' V_K leader sequence primers designed to amplify each of the V_K mouse families combined with a single reverse primer specific to the mouse kappa constant region were used to amplify and sequence the kappa light chain. The V_H and V_L transcripts were amplified from 100 ng total RNA using reverse transcriptase polymerase chain reaction (RT-PCR).

[0562] A total of eight RT-PCR reactions were run for each hybridoma: four for the V_K light chain and four for the V_γ gamma heavy chain ($\gamma 1$). The One Step RT-PCR kit was used for amplification (Qiagen). This kit provides a blend of Sensiscript and Omniscript Reverse Transcriptases, HotStarTaq DNA Polymerase, dNTP mix, buffer and Q-Solution, a novel additive that enables efficient amplification of "difficult" (e.g., GC-rich) templates. Reaction mixtures were prepared that included 3 μ L of RNA, 0.5 of 100 μ M of either heavy chain or kappa light chain primers (custom synthesized by IDT), 5 μ L of 5 $\times$ RT-PCR buffer, 1 μ L dNTPs, 1 μ L of enzyme mix containing reverse transcriptase and DNA polymerase, and 0.4 μ L of ribonuclease inhibitor RNasin (1 unit). The reaction mixture contains all of the reagents required for both reverse transcription and PCR. The thermal cycler program was set for an RT step 50°C for 30 minutes, 95°C for 15 minutes, followed by 30 cycles of PCR (95°C for 30 seconds, 48°C for 30 seconds, 72°C for one minute). There was then a final incubation at 72°C for 10 minutes.

[0563] To prepare the PCR products for direct DNA sequencing, they were purified using the QIAquick™ PCR Purification Kit (Qiagen) according to the manufacturer's protocol. The DNA was eluted from the spin column using 50 μ L of sterile water and then sequenced directly from both strands. The extracted PCR products were directly sequenced using specific V region primers. Nucleotide sequences were analyzed using IMGT to identify germline V, D and J gene members with the highest sequence homology. The derived sequences were further compared to known germline DNA sequences of the Ig V- and J-regions using V-BASE2 (Retter et al., supra) and by alignment of V_H and V_L genes to the mouse germline database to assist with fabrication of the humanized constructs as set forth below.

[0564] More specifically, FIG. 10A depicts the contiguous amino acid sequences of seventy-four unique novel murine light chain variable regions from anti-SEZ6 antibodies (SEQ ID NOS: 20 - 168, even numbers) and eleven humanized light chain variable regions (SEQ ID NOS: 170 - 192, even numbers) derived from representative murine light chains. Similarly, FIG. 10B depicts the contiguous amino acid sequences of seventy-four unique novel murine heavy chain variable regions (SEQ ID NOS: 21 - 169, odd numbers) from the same anti-SEZ6 antibodies and eleven humanized heavy chain variable regions ((SEQ ID NOS: 171 - 193, odd numbers) from the same murine antibodies providing the humanized light chains. Thus, taken together FIGS. 10A and 10B provide the annotated sequences of seventy-four unique operable murine anti-SEZ6 antibodies (termed SC17.1, SC17.2, SC17.3, SC17.4, SC17.8, SC17.9, SC17.10, SC17.11, SC17.14, SC17.15, SC17.16 (duplicate of SC17.6), SC17.17, SC17.18, SC17.19, SC17.22, SC17.24 (duplicate sequence of SC17.2), SC17.27, SC17.28, SC17.29, SC17.30, SC17.32, SC17.34, SC17.35, SC17.36, SC17.38, SC17.39, SC17.40, SC17.41, SC17.42, SC17.45, SC17.46, SC17.47, SC17.49, SC17.50, SC17.53, SC17.54, SC17.56, SC17.57, SC17.59, SC17.61, SC17.63, SC17.71, SC17.72, SC17.74, SC17.76, SC17.77, SC17.79, SC17.81, SC17.82, SC17.84, SC17.85, SC17.87, SC17.89, SC17.90, SC17.91, SC17.93, SC17.95, SC17.97, SC17.99, SC17.102, SC17.114, SC17.115, SC17.120, SC17.121, SC17.122, SC17.140, SC17.151, SC17.156, SC17.161, SC17.166, SC17.187, SC17.191, SC17.193, SC17.199 and SC17.200) and eleven humanized antibodies (termed hSC17.16, hSC17.17, hSC17.24, hSC17.28, hSC17.34, hSC17.46, hSC17.151, hSC17.155, hSC17.156, hSC17.161 and

hSC17.200). Note that these same designations may refer to the clone that produces the subject antibody and, as such, the use of any particular designation should be interpreted in the context of the surrounding disclosure.

[0565] Additionally, hSC17.200vL1 (SEQ ID NO: 192) is a variant of the humanized light chain construct hSC17.200 (SEQ ID NO: 190), hSC17.155vH1 - hSC17.155vH6 (SEQ ID NOS: 193-198) are variants of the heavy chain construct hSC.155 (SEQ ID NO: 184) which is derived from SC17.90 (SEQ ID NO: 127) and that hSC161vH1 (SEQ ID NO: 199) is a variant of the heavy chain construct hSC17.161 (SEQ ID NO: 189). As will be discussed in more detail below these variants were constructed and tested to optimize one or more biochemical properties of the parent antibody.

[0566] Further, corresponding nucleic acid sequences of each of the seventy-four exemplary murine modulators and eleven humanized modulators and variants set forth in FIGS. 10A and 10B are included in the sequence listing of the instant application (SEQ ID NOS: 220 - 399).

[0567] For the purposes of the instant application the SEQ ID NOS of each particular antibody are sequential. Thus mAb SC17.1 comprises SEQ ID NOS: 20 and 21 for the light and heavy chain variable regions respectively. In this regard SC17.2 comprises SEQ ID NOS: 22 and 23, SC17.9 comprises SEQ ID NOS: 24 and 25, and so on. Moreover, corresponding nucleic acid sequences for each antibody amino acid sequence in FIGS. 10A and 10B are set forth in the sequence listing. In the sequence listing the included nucleic acid sequences comprise SEQ ID NOS that are two hundred greater than the corresponding amino acid sequence (light or heavy chain). Thus, nucleic acid sequences encoding the light and heavy chain variable region amino acid sequences of mAb SC17.1 (i.e., SEQ ID NOS: 20 and 21) comprise SEQ ID NOS: 220 and 221. In this regard nucleic acid sequences encoding all of the disclosed light and heavy chain variable region amino acid sequences, including those encoding the humanized constructs and variants thereof, are numbered similarly and comprise SEQ ID NOS: 220 - 399.

EXAMPLE 8

GENERATION OF CHIMERIC AND HUMANIZED SEZ6 MODULATORS

[0568] As alluded to above, eleven of the murine antibodies from Example 7 were humanized using complementarity determining region (CDR) grafting. Human frameworks for heavy and light chains were selected based on sequence and structure similarity with respect to functional human germline genes. In this regard structural similarity was evaluated by comparing the mouse canonical CDR structure to human candidates with the same canonical structures as described in Chothia et al. (supra).

[0569] More particularly eleven murine antibodies SC17.16, SC17.17, SC17.24, SC17.28, SC17.34, SC17.46, SC17.151, SC17.155 (duplicate of SC17.90), SC17.156, SC17.161 and SC17.200 were humanized using a computer-aided CDR-grafting method (Abysis Database, UCL Business Plc.) and standard molecular engineering techniques to provide hSC17.16, hSC17.17, hSC17.24, hSC17.28, hSC17.34, hSC17.46, hSC17.151, hSC17.155, hSC17.156, hSC17.161 and hSC17.200 modulators. The human framework regions of the variable regions were selected based on their highest sequence homology to the subject mouse framework sequence and its canonical structure. For the purposes of the humanization analysis, the assignment of amino acids to each of the CDR domains is in accordance with Kabat et al. numbering (supra).

[0570] Molecular engineering procedures were conducted using art-recognized techniques. To that end total mRNA was extracted from the hybridomas and amplified as set forth in Example 7 immediately above.

[0571] From the nucleotide sequence information, data regarding V, D and J gene segments of the heavy and light chains of subject murine antibodies were obtained. Based on the sequence data new primer sets specific to the leader sequence of the Ig V_H and V_K light chain of the antibodies were designed for cloning of the recombinant monoclonal antibody. Subsequently the V-(D)-J sequences were aligned with mouse Ig germ line sequences. The resulting genetic arrangements for each of the eleven humanized constructs are shown in Table 3 immediately below.

TABLE 3

mAb	human VH	human DH	human JH	FW changes	human VK	human JK	FW changes
hSC17.16	IGHV1-2	IGHD3-16	JH5	none	IGKV-O2	JK1	none
hSC17.17	IGHV1-2	IGHD4-11	JH4	none	IGKV-L6	JK2	none
hSC 17.24	VH1-f	IGHD5-12	JH4	48I, 73K	VKB3	JK1	none
hSC17.28	IGHV1-2	IGHD3-16	JH4	none	IGKV-A10	JK4	none
hSC17.34	IGHV1-3	IGHD3-10	JH4	71V	IGKV-L1	JK1	71Y
hSC17.46	IGHV1-2	IGHD4-23	JH4	48I, 69L	IGKV-L11	JK1	87F

(continued)

mAb	human VH	human DH	human JH	FW changes	human VK	human JK	FW changes
hSC17.151	IGHV1-46	IGHD1-14	JH4	none	VKL6	JK2	none
hSC17.155	IGHV1-46	IGHD2-2	JH4	none	VKB3	JK1	none
hSC17.156	IGHV2-26	IGHD4-17	JH4	none	VKO1	JK4	none
hSC17.161	IGHV1-2	IGHD1-14	JH4	none	VKB3	JK2	none
hSC17.200	IGHV5-51	IGHD4-17	JH4	none	IGKV-L6	JK4	none

[0572] The humanized antibodies listed in Table 3 correspond to the annotated light and heavy chain sequences set forth in FIGS. 10A and 10B (SEQ ID NOS: 170-191). The corresponding nucleic acid sequences of the light and heavy chain variable regions are set forth in the appended sequence listing. Table 3 further demonstrates that very few framework changes were necessary to maintain the favorable properties of the binding modulators. In this respect framework changes or back mutations were only made in three of the heavy chain variable regions and only two framework modifications were undertaken in the light chain variable regions.

[0573] Note that, for some humanized light and heavy chain variable regions amino acid mutations were introduced in the FRs or CDRs to improve stability while maintaining antigen binding. In the case of SC17.155 six variants were produced, termed hSC17.155vH1-hSC17.155vH6. In each of these variants changes were made to the FR or CDRs of the heavy chain variable region of hSC17.155 while the light chain variable region was left unchanged. In the case of hSC17.155vH1 and hSC17.155vH2 a point mutation was introduced in FR1 (SEQ ID NOS: 471 and 472) of the heavy chain variable region of hSC17.155. In the case of hSC17.155vH3 a point mutation was introduced in CDRH1 (SEQ ID NO: 473) of the hSC17.155 heavy chain variable region. In the case of hSC17.155vH4 - hSC17.155vH6 point mutations were introduced in CDRH2 (SEQ ID NOS: 474, 475 and 476, respectively) of the hSC17.155 heavy chain variable region. In the case of hSC17.161 certain amino acids in FR1 (SEQ ID NO: 477); FR2 (SEQ ID NO: 478) and FR3 (SEQ ID NO: 479) of the heavy chain variable region were changed whereas the light chain variable region was not modified. Finally, in the case of hSC200vL1 a point mutation was introduced in CDRL1 (SEQ ID NO: 480) of the hSC17.200 light chain variable region and the heavy chain variable region was not modified. In each case, the binding affinity of the antibodies with modified CDRs or FRs was found to be equivalent to either the corresponding chimeric or murine antibody. The sequences of nine exemplary humanized variant chains (light and heavy,) are listed at the end of FIGS. 10A and 10B (SEQ ID NOS: 192 - 199) where they retain the designation of the humanized parent chain with notation to indicate they have been altered (e.g. hSC17.200vL1, hSC17.155vH1-6 and hSC17.161vH1). The full length amino acid sequences of exemplary humanized antibodies, hSC17.200 and hSC17.200vL1 are set out in FIG. 10C as SEQ ID NOS: 400-402. The humanized antibody variant hSC17.200vL1 is derived from humanized antibody hSC17.200 and shares a common HC with the hSC17.200 antibody. Thus the full length LC and HC of hSC17.200 correspond to SEQ ID NOS: 400 and 401, respectively; and the full length LC and HC of hSC17.200vL1 correspond to SEQ ID NOS: 403 and 401, respectively. Following humanization of all selected antibodies by CDR grafting, the resulting light and heavy chain variable region amino acid sequences were analyzed to determine their homology with regard to the murine donor and human acceptor light and heavy chain variable regions. The results, shown in Table 4 below, reveal that the humanized constructs consistently exhibited a higher homology with respect to the human acceptor sequences than with the murine donor sequences. More specifically, the humanized heavy and light chain variable regions generally show a higher percentage homology to a closest match of human germline genes (84%-95%) as compared to the homology of the humanized variable region sequences and the donor hybridoma protein sequences (74%-89%).

TABLE 4

mAb	Homology to Human (CDR acceptor)	Homology to Murine Parent (CDR donor)
hSC17.16 HC	91%	80%
hSC17.16 LC	86%	85%
hSC17.17 HC	93%	80%
hSC17.17 LC	87%	77%
hSC 17.24 HC	86%	79%
hSC 17.24 LC	93%	89%

(continued)

mAb	Homology to Human (CDR acceptor)	Homology to Murine Parent (CDR donor)
hSC17.28 HC	89%	77%
hSC 17.28 LC	92%	78%
hSC17.34 HC	85%	83%
hSC17.34 LC	84%	86%
hSC 17.46 HC	85%	83%
hSC 17.46 LC	84%	80%
hSC17.151 HC	90%	79%
hSC17.151 LC	87%	80%
hSC17.155 HC	90%	80%
hSC17.155 LC	95%	87%
hSC17.156 HC	89%	79%
hSC17.156 LC	86%	93%
hSC17.161 HC	89%	86%
hSC17.161 LC	93%	87%
hSC17.200 HC	90%	74%
hSC17.200 LC	88%	82%

[0574] Upon testing, and as will be discussed in Example 9, each of the humanized constructs exhibited favorable binding characteristics roughly comparable to those shown by the murine parent antibodies (Data not shown).

[0575] Whether humanized or murine, once the nucleic acid sequences of the variable regions are determined the antibodies of the instant invention may be expressed and isolated using art-recognized techniques. To that end synthetic DNA fragments of the chosen heavy chain (humanized or murine) variable region were cloned into a human IgG1 expression vector. Similarly the variable region light chain DNA fragment (again humanized or murine) was cloned into a human light chain expression vector. The selected antibody was then expressed by co-transfection of the derived heavy and the light chain nucleic acid constructs into CHO cells.

[0576] More particularly, one compatible method of antibody production comprised directional cloning of murine or humanized variable region genes (amplified using PCR) into selected human immunoglobulin expression vectors. All primers used in Ig gene-specific PCRs included restriction sites which allowed direct cloning into expression vectors containing human IgG1 heavy chain and light chain constant regions. In brief, PCR products were purified with Qiaquick PCR purification kit (Qiagen) followed by digestion with AgeI and XhoI (for the heavy chain) and XmaI and DraIII (for the light chain), respectively. Digested PCR products were purified prior to ligation into expression vectors. Ligation reactions were performed in a total volume of 10 μ L with 200U T4-DNA Ligase (New England Biolabs), 7.5 μ L of digested and purified gene-specific PCR product and 25ng linearized vector DNA. Competent E. coli DH10B bacteria (Life Technologies) were transformed via heat shock at 42°C with 3 μ L ligation product and plated onto ampicillin plates (100 μ g/mL). The AgeI-EcoRI fragment of the V_H region was then inserted into the same sites of pEE6.4HuIgG1 expression vector while the synthetic XmaI-DraIII VK insert was cloned into the XmaI-DraIII sites of the respective pEE12.4Hu-Kappa expression vector.

[0577] Cells producing the selected antibody were generated by transfection of HEK 293 cells with the appropriate plasmids using 293fectin. In this respect plasmid DNA was purified with QIAprep Spin columns (Qiagen). Human embryonic kidney (HEK) 293T (ATCC No CRL-11268) cells were cultured in 150mm plates (Falcon, Becton Dickinson) under standard conditions in Dulbecco's Modified Eagle's Medium (DMEM) supplemented with 10% heat inactivated FCS, 100 μ g/mL streptomycin, 100 U/mL penicillin G (all from Life Technologies).

[0578] For transient transfections cells were grown to 80% confluency. Equal amounts of IgH and corresponding IgL chain vector DNA (12.5 μ g of each) was added to 1.5 mL Opti-MEM mixed with 50 μ L HEK 293 transfection reagent in 1.5 mL opti-MEM. The mix was incubated for 30 min at room temperature and distributed evenly to the culture plate. Supernatants were harvested three days after transfection, replaced by 20 mL of fresh DMEM supplemented with 10% FBS and harvested again at day 6 after transfection. Culture supernatants were cleared of cell debris by centrifugation

at 800×g for 10 min and stored at 4°C. Recombinant chimeric and humanized antibodies were purified with Protein G beads (GE Healthcare) and stored under appropriate conditions.

EXAMPLE 9

CHARACTERISTICS OF SEZ6 MODULATORS

[0579] Various methods were used to analyze the binding and immunochemical characteristics of selected SEZ6 modulators generated as set forth above. Specifically, a number of the antibody modulators were characterized as to affinity, binning, and cross reactivity with regard to human, cynomolgus, rat and mouse SEZ6 antigen along with SEZ6L and SEZ6L2 proteins by art-recognized methods including flow cytometry. Affinities and kinetic constants k_{on} and k_{off} of the selected modulators were measured using bio-layer interferometry analysis on a ForteBio RED (ForteBio, Inc.) or surface plasmon resonance using a Biacore 2000 each according to the manufacturer's instructions.

[0580] The characterization results are set forth in tabular form in FIG. 11A where it may be seen that the selected modulators generally exhibited relatively high affinities in the nanomolar range and, in many cases, were cross-reactive with one or more SEZ6 orthologs. FIG. 11A further lists the empirically determined bin occupied by the subject modulator. Taken together, these data demonstrate the varied binding properties of the disclosed modulators as well as their potential suitability for pharmaceutical development based on their reactivity in animal models.

[0581] In this regard flow cytometry was performed using a FACSCanto II as per the manufacturer's instructions in order to confirm that selected SC17 antibody modulators can immunospecifically associate with human SEZ6 and to determine whether the same modulators cross-react with cynomolgus, rat and/or murine SEZ6 in addition to SEZ6L and SEZ6L2. More particularly modulators were tested for cross reactivity to murine SEZ6 and rat SEZ6 by flow cytometry against Neuro2a (ATCC Cat # CCL131), and RIN-m5F (ATCC cat # CRL-11605) cell lines which express mouse SEZ6 and rat SEZ6, respectively. For examining cross reactivity to cynomolgus SEZ6, yeast displaying the extracellular domain of cynomolgus SEZ6 (Boder *et al*, 1997) were used for flow cytometry analysis.

[0582] Briefly 1×10^5 cells per well of Neuro2a, RIN-5mF, or yeast displaying cynomolgus SEZ6 cells were incubated for 30 minutes with 50 μ L PBS (2%FCS) buffer with 5 μ g/mL antibody. Cells were washed twice with the same buffer and then incubated with 50 μ L per sample DyLight 649 labeled goat-anti-mouse IgG, Fc fragment specific secondary diluted 1:200 in PBS buffer. After incubating for 15 minutes cells were washed twice with the PBS buffer and re-suspended in the same with DAPI for flow cytometry analysis of Neuro2a and Rin-m5F or buffer without DAPI for flow cytometric analysis of yeast cells with cSEZ6. Antibodies that bound to the Neuro2a or RIN-m5F cell lines, or yeast displaying cynomolgus SEZ6 were considered to be cross reactive to murine SEZ6, rat SEZ6, or cynomolgus SEZ6, respectively. FIG. 11A shows the cross reactivity results. Six antibodies were cross reactive for human and mouse SEZ6 (SC17.6 (duplicate of SC17.16), SC17.7, SC17.19, SC17.24, SC17.26 and SC17.42); six for human and rat SEZ6 (SC17.6, SC17.17, SC17.19, SC17.26, SC17.28, SC17.34 and SC17.42); and six for human and cynomolgus SEZ6 (SC17.17, SC17.24, SC17.26, SC17.34, SC17.36 and SC17.45). Note that SC17.6 is duplicative of SC17.16 and exhibits the same binding characteristics.

[0583] To verify the cross reactivity data above for rat SEZ6 and to determine the affinity and kinetic constants k_{on} and k_{off} of the selected effectors, either bio-layer interferometry analysis on a ForteBio RED (ForteBio, Inc.) or surface plasmon resonance on a Biacore 2000 (GE Healthcare) were conducted. Affinities were determined to both human recombinant SEZ6-His and rat recombinant SEZ6-His generated in Example 5. As seen in FIG. 11A, a number of the antibodies tested, cross reacted with rat SEZ6. The selected modulators exhibited relatively high affinities for both rat and human SEZ6 in the nanomolar range.

[0584] To determine cross reactivity to family member proteins, SEZ6L and SEZ6L2, an ELISA-based assay was used. Plates were coated with SEZ6, SEZ6L, or SEZ6L2 proteins at 0.2 μ g/mL in PBS overnight. After washing with PBS containing 0.05% (v/v) Tween 20 (PBST), the wells were blocked with 2% (w/v) BSA in PBS (PBSA), 100 μ L/well for 1 hour at room temperature. Antibody was then added at 1 μ g/mL in 100 μ L PBSA for 1 hour at room temperature. After washing with PBST, 100 μ L/well HRP-labeled goat anti-mouse IgG diluted 1:2,000 in PBSA for 1 hour at room temperature. The plates were washed and 100 μ L/well of the TMB substrate solution (Thermo Scientific 34028) was added for 15 minutes at room temperature. After developing, an equal volume of 2M H_2SO_4 was added to stop substrate development and analyzed by spectrophotometer at OD 450. FIG. 11A shows that one antibody was cross reactive with SEZ6L (SC17.7) and five were cross-reactive with SEZ6L2 (SC17.6, SC17.7, SC17.19, SC17.26 and SC17.28). As discussed above, such pan-SEZ6 antibodies are compatible with the teachings herein and may be used in conjunction with the disclosed methods.

[0585] Binding characteristics of the following humanized constructs from Example 8, hSC17.16, hSC17.17, hSC17.24, hSC17.28, hSC17.34 and hSC17.46, were analyzed in order to determine whether the CDR grafting process had appreciably altered their binding characteristics. The humanized constructs (CDR grafted) were compared with "traditional" chimeric antibodies comprising the murine parent (or donor) heavy and light chain variable domains and a human

constant region substantially equivalent to that used in the humanized constructs. With these constructs surface plasmon resonance was conducted using a Biacore 2000 (GE Healthcare) to identify any subtle changes in rate constants brought about by the humanization process. In all cases, the humanized antibodies had binding affinity equivalent or better than the corresponding murine antibodies (Data not shown).

[0586] Antibody binning was determined for various SEZ6 modulators as shown in FIG. 11A. A ForteBio RED was used per manufacturer's instructions to identify competing antibodies that bound to the same or different bins. Briefly, a reference antibody (Ab1) was captured onto an anti-mouse capture chip, a high concentration of non-binding antibody was then used to block the chip and a baseline was collected. Monomeric, recombinant human SEZ6 (described in Example 5) was then captured by the specific antibody (Ab1) and the tip was dipped into a well with either the same antibody (Ab1) as a control or into a well with a different test antibody (Ab2). If additional binding was observed with a new antibody, then Ab1 and Ab2 were determined to be in a different bin. If no further binding occurred, as determined by comparing binding levels with the control Ab1, then Ab2 was determined to be in the same bin. As known in the art this process can be expanded to screen large libraries of unique antibodies using a full row of antibodies representing unique bins in a 96-well plate. In the instant case this binning process showed the screened antibodies bound to at least seven different bins on the SEZ6 protein. Bins A-F are unique bins and the antibodies contained in each of these bins compete with each other (but not with antibodies from other defined bins) for binding to the SEZ6 protein. Bin U contains antibodies that do not compete with antibodies in bins A-F, but may compete for binding with each other. FIG. 11B shows the correlation between the antibodies that bind a certain epitope and the ability of those groups of antibodies to kill HEK-293T cells overexpressing SEZ6 (described in more detail in Example 10 immediately below).

EXAMPLE 10

EPITOPE MAPPING OF SEZ6 MODULATORS

[0587] In order to characterize the epitopes that the disclosed SEZ6 antibody modulators associate with or bind to, domain-level epitope mapping was performed using a modification of the protocol described by Cochran et al. (J Immunol Methods. 287 (1-2):147-158 (2004)) which is incorporated herein by reference). Individual domains of SEZ6 were expressed on the surface of yeast and binding by each SEZ6 antibody was determined through flow cytometry.

[0588] Yeast display plasmid constructs were created for the expression of the following constructs: SEZ6 extracellular domain (amino acids 1-904); Sushi Domain 1 (amino acids 336-395), CUB Domain 1 (amino acids 297-508), Sushi Domain 2 (amino acids 511-572), CUB Domain 2 (amino acids 574-685), Sushi Domain 3 (amino acids 690-748), Sushi Domain 4 (amino acids 750-813), Sushi Domain 5 (amino acids 817-878), and Sushi Domain 5 + C-terminus (amino acids 817-904). Additionally, the N terminal domain (amino acids 1-335) was divided into 3 fragments termed N1 (amino acids 1-70), N2 (amino acids 71-169) and N3 (amino acids 169-335), each of which was cloned into the yeast display plasmid. Amino acid numbering does not include the 19 amino acid leader peptide. For domain information see generally UniProtKB/Swiss-Prot database entry Q53EL9. These plasmids were transformed into yeast, which were then grown and induced as described in Cochran *et al.* Note that all amino acid numbering is based on mature SEZ6 protein without the 19 amino acid leader sequence.

[0589] To test for binding to a particular construct, 200,000 induced yeast cells expressing the desired construct were washed twice in PBS + 1 mg/mL BSA (PBSA), and incubated in 50 μ L of PBSA with chicken anti c-myc (Life Technologies) at 0.1 μ g/mL and either 50 nM purified antibody or 1:2 dilution of unpurified supernatant from hybridomas cultured for 7 days. Cells were incubated for 90 minutes on ice and then washed twice in PBSA. Cells were then incubated in 50 μ L PBSA with the appropriate secondary antibodies: for murine antibodies, Alexa 488 conjugated anti-chicken, and Alexa 647 conjugated goat anti-mouse (both Life Technologies) were added at 1 μ g/mL each, and for humanized or chimeric antibodies, Alexa 647 conjugated anti-chicken (Life Technologies) and R-phycoerythrin conjugated goat anti-human (Jackson ImmunoResearch) were added at 1 μ g/mL each. After twenty minutes' incubation on ice, cells were washed twice with PBSA and analyzed on a FACS Canto II.

[0590] All modulators bound uniquely to a single domain expressed on yeast cells. In some cases, antibody clones bound specifically to yeast expressing Sushi Domain 5 + C-terminus but not to yeast expressing Sushi Domain 5. These antibody clones were concluded to bind to the C-terminal region only (amino acids 879-904).

[0591] Epitopes were classified either as conformational (i.e. discontinuous) or linear. Yeast displaying the SEZ6 ECD construct was heat treated for 30 minutes at 80°C in order to denature the antigen, washed twice in ice-cold PBSA and then subjected to the same staining protocol and flow cytometry analysis as described above. Antibodies that bound to both the denatured and native yeast were classified as binding to a linear epitope, whereas antibodies that bound native yeast but not denatured yeast were classified as conformationally specific.

[0592] A summary of the domain-level epitope mapping data of the antibodies tested is presented in TABLE 5 below. Antibodies that bind a linear epitope are underlined and antibodies that bind SEZ6 family members SEZ6L and SEZ6L2 are designated with an asterisk and/or a dagger, respectively.

TABLE 5

Domain	Antibody Clones
N1 (aa 1-70)	SC17.4, SC17.7 [†] *, SC17.9, SC17.56, SC17.81, SC17.101, SC17.114, SC17.120, SC17.134, SC17.151, SC17.162, SC17.177, SC17.182, SC17.185, SC17.196, SC17.197, SC17.199
N2 (aa 71-169)	SC17.24, SC17.49, SC17.104, SC17.144, SC17.149, SC17.168, SC17.176, SC17.198
N3 (aa 170-335)	SC17.26 [†] , SC17.42, SC17.83, SC17.85, SC17.88, SC17.91, SC17.92, SC17.99, SC17.125, SC17.128, SC17.130, SC17.137, SC17.145, SC17.161, SC17.192, SC17.195
Sushi Domain 1 (aa 336-395)	SC17.34, SC17.36, SC17.46, SC17.75, SC17.82, SC17.87, SC17.97, SC17.116, SC17.129, SC17.178, SC17.187, SC17.200
CUB Domain 1 (aa 397-508)	SC17.73, SC17.76, SC17.86, SC17.100, SC17.105, SC17.107, SC17.109, SC17.122, SC17.124, SC17.136, SC17.138, SC17.146, SC17.154, SC17.170, SC17.174, SC17.189, SC17.201, SC17.202
Sushi Domain 2 (aa 511-572)	SC17.90, SC17.108, SC17.112, SC17.135, SC17.167, SC17.173, SC17.179, SC17.184, SC17.203, SC17.204
CUB Domain 2 (aa 574-685)	SC17.6 [†] , SC17.28 [†] , SC17.103, SC17.109, SC17.119, SC17.181, SC17.186, SC17.194
Sushi Domain 3 (aa 690-748)	SC17.72, SC17.84, SC17.95, SC17.141, SC17.143, SC17.163
Sushi Domain 4 (aa 750-813)	SC17.17, SC17.19 [†] , SC17.93, SC17.102, SC17.121, SC17.140, SC17.156, SC17.159, SC17.166, SC17.175, SC17.180, SC17.191, SC17.193
Sushi Domain 5 (aa 817-878)	SC17.74, SC17.106, SC17.142, SC17.190
C terminus (aa 879-904)	SC17.96, SC17.132

[0593] An interesting and surprising trend was observed when an *in vitro* cell killing assay was performed using the domain-mapped SEZ6 antibody modulators described in this Example 10. The *in vitro* killing assay, performed essentially as described below in Example 14, determined the ability of a particular antibody to internalize and kill HEK-293 cells. FIG. 11B is a plot of efficacy of the tested antibodies versus the domains to which they bind. Antibodies that bind to certain domains including: N1, N3, Sushi Domain 1, and Sushi Domain 4, exhibited enhanced *in vitro* killing. The antibodies that associate with Sushi Domain 4, which are very effective at internalizing and killing cells, exhibit a strong correlation with the IGHV1-34 and IKV4-59 murine germline framework regions.

[0594] Fine epitope mapping was further performed on selected antibodies to determine the specific amino acids to which they bound. Antibodies that bound to a linear epitope were mapped using the Ph.D.-12 phage display peptide library kit (New England Biolabs E8110S). The antibody selected for epitope mapping was coated onto a Nunc MaxiSorp tube (Nunc) at 50 µg/mL in 3 mL 0.1 M sodium bicarbonate solution, pH 8 and incubated overnight. The tube was blocked with 3% BSA solution in bicarbonate solution. Then, 10¹¹ input phage in PBS + 0.1% Tween-20 was allowed to bind, followed by ten consecutive washes with 0.1 % Tween-20 to wash away non-binding phage. Remaining phage were eluted with 1 mL 0.2 M glycine for 10 minutes at room temperature with gentle agitation, followed by neutralization with 150 µL 1 M Tris-HCl pH 9. Eluted phage were amplified and panned again with 10¹¹ input phage, using 0.5% Tween-20 during the wash steps to increase selection stringency. DNA from 24 plaques of the eluted phage from the second round was isolated using the Qiaprep M13 Spin kit (Qiagen) and sequenced. Binding of clonal phage was confirmed using an ELISA assay, where the mapped antibody or a control antibody was coated onto an ELISA plate, blocked, and exposed to each clone phage. Phage binding was detected using horseradish peroxidase conjugated anti-M13 antibody (GE Healthcare), and the 1-Step Turbo TMB ELISA solution (Pierce). Phage peptide sequences from specifically binding phage were aligned using Vector NTI (Life Technologies) against the antigen ECD peptide sequence to determine the epitope of binding.

[0595] Selected antibodies that bound to a discontinuous epitope were mapped using the technique described by Chao *et al.* (2007). Libraries of SEZ6 ECD mutants were generated with error prone PCR using nucleotide analogues 8-oxo-2'-deoxyguanosine-5'-triphosphate and 2'-deoxy-p-nucleoside-5'-triphosphate (both from TriLink Bio) for a target

mutagenesis rate of one amino acid mutation per clone. These were transformed into a yeast display format. Using the technique described above for domain-level mapping, the library was stained for c-myc and antibody binding at 50 nM. Using a FACS Aria (BD), clones that exhibited a loss of binding compared to wild type SEZ6 ECD were sorted. These clones were re-grown, and subjected to another round of FACS sorting for loss of binding to the target antibody. Using the Zymoprep Yeast Plasmid Miniprep kit (Zymo Research), individual ECD clones were isolated and sequenced. Where necessary, mutations were reformatted as single-mutant ECD clones using the Quikchange site directed mutagenesis kit (Agilent).

[0596] Individual ECD clones were next screened to determine whether loss of binding was due to a mutation in the epitope, or a mutation that caused misfolding. Mutations that involved cysteine, proline, and stop codons were automatically discarded due to the high likelihood of a misfolding mutation. Remaining ECD clones were then screened for binding to a non-competing, conformationally specific antibody. ECD clones that lost binding to non-competing, conformationally specific antibodies were concluded to contain misfolding mutations, whereas ECD clones that retained equivalent binding as wild type SEZ6 ECD were concluded to be properly folded. Mutations in the ECD clones in the latter group were concluded to be in the epitope. Homology models of isolated domains were also constructed using MODELLER to confirm that residues identified to be in the epitope: 1) were localized in close proximity to each other in the folded homology model, and 2) had side chains that were solvent exposed, and not buried, since buried residues would have a higher chance of causing misfolding, and would unlikely be part of the epitope of binding. A summary of antibodies with their epitopes are listed in Table 6. The residues found to most significantly contribute to the structure of the epitope are underlined.

TABLE 6

Antibody Clone	Epitope	Discontinuous	SEQ ID NO:
SC17.4	Q12, P14, I16, E17, E18	No	401
SC17.17	<u>R762</u> , L764, Q777, I779, <u>D781</u> , <u>Q782</u>	Yes	NR
SC17.24	L73, P74, F75, Q76, P77, D78, P79	No	402
SC17.34	T352, S353, <u>H375</u>	Yes	NR
SC17.36	<u>T352</u> , S353, <u>H375</u> , <u>S359</u>	Yes	NR
SC17.46	<u>R342</u> , <u>K389</u>	Yes	NR
SC17.102	<u>R762</u> , L764, Q777, I779, <u>D781</u> , <u>Q782</u>	Yes	NR
SC17.121	<u>R762</u> , L764, Q777, I779, <u>D781</u> , <u>Q782</u>	Yes	NR
SC17.140	<u>R762</u> , L764, Q777, I779, D781, Q782	Yes	NR
SC17.156	R807, <u>K810</u>	Yes	NR
SC17.166	<u>R762</u> , L764, Q777, I779, <u>D781</u> , <u>Q782</u>	Yes	NR
SC17.191	<u>R762</u> , L764, Q777, I779, <u>D781</u> , <u>Q782</u>	Yes	NR
SC17.200	T352, S353, <u>H375</u>	Yes	NR

[0597] NR indicates that a SEQ ID NO was not assigned as the epitopes were discontinuous.

[0598] For the purpose of fine epitope mapping of SC17.34, SC17.36, SC17.200 and SC17.46, point mutations were constructed on the isolated domain, Sushi Domain 1, which was determined to be the domain of binding by domain-level epitope mapping. In some cases, candidate mutations for screening were determined independently of the yeast displayed library. For example, in the case of SC17.46, candidate mutations for screening were not identified in a library-based screen; rather they were identified on the basis of domain mapping, lack of cross reactivity to cynomolgus SEZ6 ECD and rat SEZ6 ECD, and sequence alignments of the different species to identify differences in the species' primary sequence. In the case of SC17.102, SC17.121, SC17.140, SC17.156, SC17.166, SC17.191, and SC17.200, candidate residues were identified using a combination of alanine scanning, and domain binding analysis. In all cases, candidate mutations were subjected to the same analysis as other antibodies whose candidate mutations were identified by a library based approach.

[0599] The present invention is directed to anti-SEZ6 antibodies and antibody drug conjugates that bind to an epitope on a SEZ6 protein (including, for example, a SEZ6 protein of SEQ ID NO: 3 or 4), wherein the epitope comprises amino acid residues selected from the group consisting of (i) residues Q12, P14, I16, E17, E18; (ii) residues R762, L764, Q777, I779, D781, Q782; (iii) residues L73, P74, F75, Q76, P77, D78, P79; (iv) residues T352, S353, H375; (v) residues T352,

S353, H375, S359; (vi) residues R342, K389; (vii) residues R762, L764, Q777, I779, D781, Q782; or (viii) residues R807, K810. In one embodiment, the invention is directed to anti-SEZ6 antibodies and antibody drug conjugates that bind to an epitope on a SEZ6 protein (including, for example, a SEZ6 protein of SEQ ID NO: 3 or 4), wherein the epitope comprises amino acid residues selected from the group consisting of (i) residues Q12, P14, I16, E17, E18; (ii) residues R762, L764, Q777, I779, D781, Q782; (iii) residues L73, P74, F75, Q76, P77, D78, P79; (iv) residues T352, S353, H375; (v) residues T352, S353, H375, S359; (vi) residues R342, K389; (vii) residues R762, L764, Q777, I779, D781, Q782; or (viii) residues R807, K810. In another embodiment the invention is directed to anti-SEZ6 antibodies or antibody drug conjugates that bind to the same epitope on a SEZ6 protein (including, for example, a SEZ6 protein of SEQ ID NO: 3 or 4) as any one of the following antibodies: SC17.4, SC17.17, SC17.24, SC17.34, SC17.36, SC17.46, SC17.102, SC17.121, SC17.140, SC17.156, SC17.166, SC17.191 and SC17.200.

EXAMPLE 11

DETECTION OF SEZ6 SURFACE EXPRESSION BY FLOW CYTOMETRY

[0600] Flow cytometry was used to assess the specificity of the anti-SEZ6 antibodies that were generated for detecting the presence of human SEZ6 protein on the surface of engineered HEK-293T cell lines, constructed as described in Example 5. Isotype-stained and fluorescence minus one (FMO) controls were employed to confirm staining specificity. Briefly, HEK-293T transduced with human SEZ6 and GFP (see Example 5) or harvested NTX tumor samples were dissociated and dispersed into suspension using art-recognized enzymatic digestion techniques (see, for example, U.S.P.N. 2007/0292414 which is incorporated herein), were incubated for 30 minutes with an anti-SEZ6 antibody. Cells were washed in PBS (2%FCS) twice and then incubated with 50 μ l per sample DyLight 649 labeled goat-anti-mouse IgG, Fc fragment specific secondary diluted 1:200 in PBS buffer. After a 15 minute incubation cells were washed twice with PBS and re-suspended in PBS with DAPI and analyzed by flow cytometry as previously discussed.

[0601] As demonstrated by the representative data shown in FIG.12A for SC17.33, the SEZ6 modulator strongly recognized HEK-293T-HuSEZ6 cells. These data demonstrate that modulators were produced that specifically recognized human SEZ6 expressed on the cell surface.

[0602] Human SEZ6 protein expression on the surface of selected NTX tumors was assessed by flow cytometry using several exemplary SC17 antibodies. The expression of SEZ6 in LU37, LU86 and KDY66 was tested using the SC17.6 (duplicate of SC17.16) antibody while expression of SEZ6 in LU50, LU100 and LU73 was tested using the SC17.10, SC17.42 and SC17.28 antibodies respectively. The results are set forth in FIG. 13A. NTX tumors were harvested, dissociated, and co-stained with commercially available anti-mouse CD45, anti-mouse H-2Kd, anti-human EpCAM and one the above-described mouse anti-human SEZ6 antibodies. Data shown in FIG. 13A was generated using cells that did not stain positively for the above mentioned anti-mouse antibodies but did stain positively for anti-human EpCAM. Similar to the HEK-293T-staining experiments described above, isotype-stained and fluorescence minus one (FMO) controls were employed to confirm staining specificity. As seen in FIG. 13A, anti-SEZ6 staining was higher than FMO in all of the human NTX tumor cells, as indicated by the fluorescent profile shift to the right, and by changes in the mean fluorescence intensity (MFI) values, for the lung NTX tumors LU37, LU50 and LU86 and kidney NTX tumor, KDY66. These data suggest that the SEZ6 protein is expressed on the surface of various NTX tumors and therefore amenable to modulation using an anti-SEZ6 antibody.

EXAMPLE 12

EXPRESSION OF SEZ6 PROTEIN IN VARIOUS TUMORS

[0603] Given the elevated SEZ6 mRNA transcript levels associated with various tumors, work was undertaken to demonstrate a corresponding increase in the expression of SEZ6 protein in NTX tumors. SEZ6 protein expression was detected with (i) an electrochemiluminescence SEZ6 sandwich ELISA assay using the MSD Discovery Platform (Meso Scale Discovery, LLC); and (ii) immunohistochemistry staining.

[0604] NTX tumors were excised from mice and flash frozen on dry ice/ethanol. Protein Extraction Buffer (Biochain Institute, Inc.) was added to the thawed tumor pieces and tumors were pulverized using a TissueLyser system (Qiagen). Lysates were cleared by centrifugation (20,000g, 20 minutes, 4°C) and the total protein concentration in each lysate was quantified using bicinchoninic acid. Protein lysates were stored at -80°C until assayed. Normal tissue lysates were purchased from a commercial source.

[0605] SEZ6 protein concentrations from the lysate samples were determined by interpolating the values from a standard protein concentration curve that was generated using purified recombinant SEZ6 protein (Example 5). The SEZ6 protein standard curve and protein quantification assay were conducted as follows:

MSD standard plates were coated overnight at 4°C with 30 µL of SC17.17 antibody at 2 µg/mL in PBS. Plates were washed in PBST and blocked in 150 µL MSD 3% Blocker A solution for one hour. Plates were again washed in PBST. 25 µL of 10x diluted lysate in MSD 1% Blocker A or serially diluted recombinant SEZ6 standard in MSD 1% Blocker A containing 10% Protein Extraction Buffer was also added to the wells and incubated for two hours. Plates were washed in PBST. The SC17.36 antibody was then conjugated to the MSD sulfo-tag and 25 µL of the tagged SC17.36 was added to the washed plates at 0.5 µg/mL in MSD 1% Blocker A. MSD Read Buffer T with surfactant was diluted to 1X in water and 150 µL was added to each well. Plates were read on a MSD Sector Imager 2400 using an integrated software analysis program to derive SEZ6 concentrations in NTX samples via interpolation from the standard curve. Values were then divided by total protein concentration to yield nanograms of SEZ6 per milligram of total lysate protein. The resulting concentrations are set forth in FIG. 12B wherein each spot represents SEZ6 protein concentrations derived from a single NTX tumor line. While each spot is derived from a single NTX line, in most cases multiple biological samples were tested from the same NTX line and values were averaged to provide the data point.

[0606] FIG. 12B shows that compared to normal tissue lysates, selected kidney, ovarian and LCNEC tumor samples exhibited moderate SEZ6 protein expression whereas the highest SEZ6 protein expression was seen in SCLC tumors. All normal tissue lysates were negative for SEZ6 protein expression with the exception of normal human brain and eye lysate.

[0607] Immunohistochemistry (IHC) was performed on PDX tumors to confirm that SEZ6 is expressed on the surface of certain PDX tumors; and in order to determine the location of the SEZ6 protein in the tumor architecture. IHC was performed on formalin fixed paraffin embedded (FFPE) tissue sections using an indirect detection method which included murine monoclonal primary antibody against SEZ6 (SC17.140), mouse specific biotin conjugated secondary antibodies, avidin/biotin complex coupled with horse radish peroxidase, and DAB detection (Nakene PK 1968; 16:557-60). SC17.140 was validated and confirmed to be appropriate for IHC by showing specific staining on FFPE sections of HEK-293T cell pellets overexpressing SEZ6 compared to naïve HEK-293T cell pellets, prepared as described in Example 5. Specificity was further confirmed by competing signal with a 5 molar excess of purified recombinant SEZ6 on HEK-293T cells overexpressing human SEZ6 and NTX tumors that were shown by IHC to express SEZ6 (data not shown).

[0608] SEZ6 expression was measured by IHC in SCLC NTX tumors, where 13 out of the 14 SCLC NTX tumors tested, expressed SEZ6 (FIG. 16A). Additional IHC analysis was performed on five tissue microarrays (TMA) generated using cored sections of 174 human tumors resected from SCLC patients (Vanderbilt University, Memorial Sloan Kettering Cancer Center and Biomax, Inc.). FIG. 16B shows SEZ6 expression measured by IHC in 174 primary patient SCLC tumors included in those five TMAs, wherein 122/174 of the SCLC samples, 2/3 SCLC/Adenocarcinoma samples and 1/1 SCLC/Spindle Cell samples were shown to express SEZ6. In contrast, the normal lung tissue samples that were tested did not express SEZ6 (FIG. 16B). The IHC results shown in FIGS. 16A and 16B were analyzed with an automated image analysis software package (Leica Biosystems) that quantifies the intensity of cell surface staining and provides a final "H-Score", which reflects the percentage of tumor cells stained at each intensity level (0 for no staining and 3 for intense staining). The H-Score is calculated as follows: (% at 0) * 0 + (% at 1+) * 1 + (% at 2+) * 2 + (% at 3+) * 3. Thus, the H-Score produces a continuous variable that ranges from 0 to 300.

[0609] IHC was also performed on FFPE sections from patients with medullary thyroid cancer. All of the patient samples that were tested were shown to express SEZ6 (FIG. 16C). The thyroid cancer samples were scored manually with a (-) for no expression, and a (+, ++ or +++) to denote increasing intensity of staining. The percentage of tumor cells in the FFPE section estimated to express SEZ6, is also shown in FIG. 16C.

[0610] RNA *in situ* hybridization (ISH) was used to verify the results of SEZ6 expression determined by IHC described immediately above. ISH was performed manually using an RNAscope® 2.0 Reagent Kit (Advanced Cell Diagnostics; Wang et al, 2012, PMID: 22166544). The RNAscope probe that was used was specific to the ECD of SEZ6. Each sample was quality controlled for RNA integrity with an RNAscope probe specific to Peptidylprolyl Isomerase B (PPIB), a cyclosporine-binding protein located within the endoplasmic reticulum of all cells. Background staining was determined using a probe specific to DiAminoPimelate (dapB) RNA. Briefly, 5 µm FFPE tissue sections from 3 of the SCLC tumor microarray that was used in FIG. 16B, were pretreated with heat and protease prior to hybridization with the SEZ6 RNA oligonucleotide probes. Preamplifier, amplifier and HRP labeled oligonucleotides were then hybridized sequentially, followed by chromogenic precipitate development with 3,3'-diaminobenzidine. Specific RNA staining signal was identified as brown, punctate dots. Samples were counterstained with Gill's Hematoxylin. FFPE sections were analyzed under a light microscope and staining was scored manually on a scale of 0 to 4, where 0 = no staining or less than 1 dot per 10 cells; 1 = 1-3 dots per cell; 2 = 4-10 dots per cell; 3 = more than 10 dots per cell and less than 10% dots found in clusters. In addition, a score of 0.5 was given if, for example, 5-30% of cells in a sample had a score of 1 and more than 70% of the cells had a score of 0. Of the tumor samples tested 5/32 (16%) did not express SEZ6; 17/32 (53%) had a score of 1; 8/32 (25%) had a score of 2 and 2/32 (6%) had a score of 3. Overall 84% of primary patient SCLC samples in the tissue microarrays expressed SEZ6 RNA at some level. This is largely in agreement with the results obtained from IHC.

[0611] The data, combined with the mRNA transcription data for SEZ6 expression set forth in Example 4, and cell surface protein expression of SEZ6 set forth in Example 11, strongly reinforces the proposition that SEZ6 determinants provide attractive targets for therapeutic intervention.

EXAMPLE 13

ENRICHMENT OF TUMOR INITIATING CELL POPULATIONS

[0612] Tumor cells can be divided broadly into two types of cell subpopulations: non-tumorigenic cells (NTG) and tumor initiating cells (TICs). TICs have the ability to form tumors when implanted into immunocompromised mice. Cancer stem cells (CSCs) are a subset of TICs and are able to self-replicate indefinitely while maintaining the capacity for multilineage differentiation. To determine whether SEZ6 expression could be correlated with enhanced tumorigenicity whole transcriptome sequencing, flow cytometry and a tumorigenicity assay were performed, all of which are described below.

[0613] Whole transcriptome analysis of SEZ6 expression in various tumor samples was performed as described in Example 1. CSCs were identified on the basis of expression of CD324 which has been shown to be a marker of stem cells in various tumors (see PCT application 2012/031280). The results in FIG. 6B show that SEZ6 mRNA expression was elevated in CSCs compared to NTG cells isolated from two SCLC NTX tumor lines (LU86 and LU95).

[0614] Flow cytometry was performed on cells from NTX lung tumors essentially as described in Example 11. LU86, LU117 and LU64 cells were co-stained with CD324, a marker of CSC populations (see PCT application 2012/031280), and the anti-SEZ6 antibody, SC17.10, SC17.28 or SC17.42, respectively to determine if SEZ6 is differentially expressed on these populations. As indicated in FIG. 13B, LU86, LU117 and LU64 cells staining positive for both CD324 and SEZ6 (solid black line) shift further to the right compared to cells staining positive for SEZ6 alone (dotted black line), indicating that SEZ6 is more highly expressed on CSCs compared to the NTG cell population. The bulk population isotype control is shown as a gray filled histogram (MOPC = IgG1).

[0615] To determine whether cell surface SEZ6 expression could be correlated with enhanced ability to generate tumors, a tumorigenicity study was conducted. NTX tumor samples were dissociated and dispersed into suspension using art-recognized enzymatic digestion techniques (see, for example, U.S.P.N. 2007/0292414 which is incorporated herein). The dissociated cell preparations from these NTX lines were stained with fluorescently conjugated antibodies specifically recognizing murine CD45, H2kD, human CD324, and human SEZ6, clone SC17.42. Two subsets of human cells, both identified based on the absence of staining with murine CD45 or H2kD (to deplete the cell preparations of murine cells) were isolated using a FACS Aria™ Flow Cytometer (BD Biosciences). One subset was isolated on the basis of a CD324 and SEZ6 co-expression, while the other subset was isolated on the basis of a CD324⁺SEZ6⁻ phenotype. The distinct marker-enriched subpopulations were subsequently transplanted into female NOD/SCID immunocompromised mice by subcutaneous injection into the mammary fat pad at a dose of approximately 50 cells per mouse.

[0616] FIGS. 14A and 14B illustrate the results of such experiments conducted using representative NTX cell lines derived from NSCLC tumors obtained from patients. FIG. 14A is a scatter plot (gated using CD324 and SEZ6) showing the distribution of mCD45⁻H2kD⁻ subset of the parent tumor and sorted putative tumorigenic cells. FIG. 14B graphically shows the measured tumor volume arising from the implantation of sorted cell subpopulations into immunocompromised mice. Values in parenthesis indicate the number of tumors generated per mice implanted.

[0617] Significantly, the data from FIG. 14 show that tumorigenicity was consistently associated with the subpopulation of cells expressing SEZ6 in combination with high levels of CD324. Conversely, these same data demonstrate that tumor cells expressing either no, or low levels of SEZ6 were much less tumorigenic than their high or positive counterparts. Based on the generated data it was surprisingly found that subpopulations of tumor cells expressing the CD324⁺SEZ6⁺ phenotype generally contain the vast majority of tumorigenic capability and suggest that SEZ6 may provide an effective therapeutic target for tumorigenic cell modulation.

EXAMPLE 14

SEZ6 MODULATORS FACILITATE DELIVERY OF CYTOTOXIC AGENTS TO SEZ6-EXPRESSING HEK-293T CELLS

[0618] To demonstrate that SEZ6 modulators of the instant invention are able to mediate the delivery of a cytotoxic agent to live cells, an *in vitro* cell killing assay was performed using selected SEZ6 antibody modulators bound to a saporin toxin. Saporin kills cells by deactivating ribosomes in the cytoplasm. Thus cell death using the following assay is an indication that the SEZ6 antibodies are able to internalize and deliver cytotoxic agents to the cytoplasm of a target cell.

[0619] An anti-Mouse IgG Fab fragment covalently linked to saporin ("Fab-Saporin") (Advanced Targeting Systems, #IT-48) was combined with unlabeled SEZ6 antibodies and incubated with HEK-293T cells expressing human SEZ6 (see Example 5). The ability of the resulting saporin complexes to internalize and kill cells was measured 72 hours later

by measuring cell viability.

[0620] Specifically, 500 cells per well in DMEM supplemented with 10% fetal bovine serum, were plated into 96 well tissue culture treated plates one day before the addition of antibodies and toxin. HEK-293T cells expressing human SEZ6 were treated with a control (IgG1, IgG2a or IgG2b) or purified murine SEZ6 modulators at a concentration of 100, 50 or 10 pM, together with 2 nM Fab-Saporin. The cells were cultured for three days, after which, viable cell numbers were enumerated using Cell Titer Glo® (Promega) as per manufacturer's instructions. Raw Luminescence Units (RLU) using cultures containing cells with the Saporin Fab fragment were set as 100% reference values and all other counts calculated accordingly (referred to as Normalized RLU or "% live cells"). FIG. 15A shows that many of the SEZ6 modulators tested mediated the killing of HEK-293T cells in a concentration dependent manner. Isotype controls (IgG2a, IgG2b, and IgG1) did not affect cell counts as shown by the results in the first three rows of FIG. 15A (ND = Not Determined).

[0621] This assay demonstrates that internalization may occur upon binding of the SEZ6-specific antibody to the cell surface, without the need for additional crosslinking or dimerization.

EXAMPLE 15

SEZ6 MODULATORS MEDIATE CYTOTOXICITY IN LUNG TUMOR CELLS *IN VITRO*

[0622] To corroborate the results of Example 14 and determine whether SEZ6 modulators can mediate toxin internalization and cell killing of human tumor cells (as opposed to engineered cells), mouse lineage-depleted NTX cells were plated and subsequently exposed to anti-SEZ6 antibodies and Fab-saporin.

[0623] NTX tumors were dissociated into a single cell suspension and plated on Primaria™ plates (BD Biosciences) in growth factor supplemented serum free media as is known in the art. After culturing the cells for one day at 37°C/5%CO₂/5%O₂, they were treated with a control (IgG1, IgG2a or IgG2b) or a murine SEZ6 modulator and Fab-saporin as described in Example 14. After seven days, the modulator-mediated saporin cytotoxicity was assessed by quantifying the remaining number of live cells using Cell Titer Glo.

[0624] As seen in FIG. 15B a reduction in the number of tumor cells was evident when LU37, a NSCLC tumor and LU80, a SCLC tumor were exposed to SC17.6 (duplicate sequence of SC17.16) and SC17.33 SEZ6 modulators. Similarly when LU100, a SCLC tumor was exposed to four SEZ6 modulators, SC17.6, SC17.19, SC17.33 and SC17.34, at 50 and 500 pM a reduction of tumor cells was effected. In contrast, isotype control antibodies did not impact the number of live cells after treatment.

[0625] Not only does this data demonstrate that exemplary antibodies described herein are able to bind SEZ6 antigen on the cell surface and facilitate the delivery of a cytotoxic payload resulting in cell death, but the above data also demonstrated that multiple anti-SEZ6 antibodies can mediate killing of various NTX tumor cells.

EXAMPLE 16

PREPARATION OF SEZ6 ANTIBODY-DRUG CONJUGATES

[0626] Based on the *in vitro* killing assays with saporin in Examples 14 and 15 and to further demonstrate the versatility of the instant invention, anti-SEZ6 antibody drug conjugates were prepared having the M-[L-D] structure as described above. That is, anti-SEZ6 antibody drug conjugates (SEZ6-ADCs) were prepared using covalently linked cytotoxic agents. More specifically, SEZ6-ADCs were prepared comprising a linker as described herein, or in the references immediately below, and selected pyrrolobenzodiazepine (PBD) dimers that were covalently attached to the disclosed modulators (see, e.g., U.S.P.Ns. 2011/0256157 and 2012/0078028 and U.S.P.N 6,214,345 each of which is incorporated herein by reference in its entirety).

[0627] PBD drug-linker combinations were synthesized and purified using art-recognized techniques in view of the cited references. While various PBD dimers and linkers were employed to fabricate the selected drug-linker combinations, each linker unit comprised a terminal maleimido moiety with a free sulfhydryl. Using these linkers, conjugations were prepared via partial reduction of the mAb with tris (2-carboxyethyl)-phosphine (TCEP) followed by reaction of reduced Cys residues with the maleimido-linker payload.

[0628] More particularly, the selected SEZ6 antibody modulator was reduced with 1.3 mol TCEP per mol mAb for 2 hr at 37°C in 25 mM Tris HCl pH 7.5 and 5 mM EDTA buffer. The reaction was allowed to cool to 15°C and the linker payload in DMSO was added at a ratio of 2.7 mol/mol mAb followed by an additional amount of DMSO to a final concentration of 6% (v/v). The reaction was allowed to proceed for 1 hour. The unreacted drug-linker was capped by addition of an excess of N-acetyl cysteine. The SEZ6-ADC (or SC17-ADC) was then purified by ion exchange column using an AKTA Explorer FPLC system (G.E. Healthcare) to remove aggregated high molecular weight antibody, co-solvent and small molecules. The eluted ADC was then buffer-exchanged by tangential flow filtration (TFF) into formulation buffer followed by concentration adjustment and addition of a detergent. The final ADC was analyzed for protein con-

centration (by measuring UV), aggregation (SEC), drug to antibody ratio (DAR) by reverse phase (RP) HPLC, presence of unconjugated antibody by hydrophobic interaction chromatography (HIC) HPLC, non-proteinaceous materials by RP HPLC and *in vitro* cytotoxicity using a SEZ6 expressing cell line.

[0629] Using the aforementioned procedure, or substantially similar methodology, a number of ADCs (i.e., M-[L-D]_n) comprising various SEZ6 modulators and PBD dimers were generated and tested in a variety of *in vivo* and *in vitro* models. For the purposes of these Examples and the instant disclosures, such ADCs may generally be termed SEZ6-ADCs or SC17-ADCs. Discrete ADCs will be named according to the antibody (e.g., SC17.17) and the specific linker-cytotoxic agent designation ADC1, ADC2, etc. Thus, exemplary modulators compatible with the instant invention may comprise SC17.17-ADC1 or SC17.24-ADC2 where ADC1 and ADC2 represent individual PBD dimer cytotoxic agents (and optionally a linker).

[0630] As an initial benchmark, the *in vitro* cytotoxicity of hSC17.17-ADC1 was measured at an IC₅₀ of 11nM when exposed to HEK293 cells overexpressing SEZ6 (data not shown).

EXAMPLE 17

CONJUGATED SEZ6 MODULATORS MEDIATE CYTOTOXICITY IN LUNG AND OVARIAN TUMOR CELLS *IN VITRO*

[0631] The ADCs generated in Example 16 above were tested to determine whether they were able to mediate toxin internalization and cell killing of primary human tumor cells *in vitro*.

[0632] Mouse lineage-depleted NTX tumor cells were exposed to anti-SEZ6 ADCs or a mouse isotype control (mIgG1) using the same method as described in Example 15, except that Fab-saporin was not added. When LU64, a SCLC tumor and OV26, a NET ovarian tumor, were treated with anti-SEZ6 ADCs (SC17.24-ADC2, SC17.28-ADC2 and SC17.34-ADC2), an increased reduction in percent viable cells was observed compared to the control mIgG1 (FIG 17A). While mIgG1 can be cytotoxic to cells at high concentrations, all three anti-SEZ6 ADCs tested were more potent, indicating an immunospecific response to SEZ6 rather than a general response to the PBD cytotoxin.

EXAMPLE 18

CONJUGATED SEZ6 MODULATORS SUPPRESS TUMOR GROWTH *IN VIVO*

[0633] The ADCs generated in Example 16 above were tested to demonstrate their ability to shrink and suppress human NTX tumor growth in immunodeficient mice.

[0634] Patient-derived NTX tumors were grown subcutaneously in the flanks of female NOD/SCID recipient mice using art-recognized techniques. Tumor volumes and mouse weights were monitored twice per week. When tumor volumes reached 150-250 mm³, mice were randomly assigned to treatment groups and injected intraperitoneally with SC17-ADC1 or a control MlgG1-ADC1. Mice were given three injections of 1 mg/kg (indicated by the vertical lines in FIG. 17B) over a period of seven days. Following treatment, tumor volumes and mouse weights were monitored until tumors exceeded 800 mm³ or mice became sick.

[0635] FIG. 17B shows that murine anti-SEZ6 ADCs are able to inhibit *in vivo* growth of a SCLC tumors (e.g. LU86) and a LCNEC tumors (e.g. LU50) in mice. In the case of LU86 the five ADCs tested (SC17.3-ADC1, SC17.24-ADC1, SC17.26-ADC1, SC17.28-ADC1 and SC17.34-ADC1) produced durable remissions lasting, in some cases, beyond 120 days posttreatment. In particular, SC17.34-ADC1 treatment inhibited tumor growth for the duration of the study at this dose, while SC17.24-ADC1 led to significant tumor growth inhibition with time to progression of greater than 50 days. Similarly, treatment of LU50 with five exemplary ADCs (SC17.3-ADC1, SC17.17-ADC1, SC17.24-ADC1, SC17.34-ADC1 and SC17.46-ADC1) resulted in tumor growth suppression lasting as long as 35 days with SC17.46. Moreover, mice treated with SC17-ADC1 did not exhibit adverse health effects beyond those typically seen in immunodeficient, tumor-bearing NOD/SCID mice. These results suggest that the disclosed ADCs may be used to effectively suppress tumor growth and that the particulars of SC17 modulator binding can have an impact on *in vivo* efficacy.

[0636] More directly the ability of a variety of conjugated modulators to dramatically retard or suppress tumor growth *in vivo* for extended periods further validates the use of SEZ6 as a therapeutic target for the treatment of proliferative disorders.

EXAMPLE 19

HUMANIZED CONJUGATED SEZ6 MODULATORS SUPPRESS TUMOR GROWTH

[0637] Given the impressive results obtained with murine anti-SEZ6 ADC modulators, additional experiments were performed to demonstrate the efficacy of exemplary humanized anti-SEZ6 ADC modulators in treating a thyroid cancer

cell line *in vitro* and *in vivo* and SCLC tumors *in vivo*.

[0638] A thyroid cancer cell line was purchased (ATCC, CRL-1803) and plated at 500 cells per well in a 96 well plate in F-12K Medium (ATCC; Catalog # 30-2004) supplemented with 10% fetal bovine serum (FBS) at 37 °C. 5 nM of either hSC17.200-ADC1, produced as set forth in Example 16, or control IgG1-ADC1 were added to the cells. After 8 days viable cell numbers were enumerated using Cell Titer Glo® (Promega) as per manufacturer's instructions. Efficacy was measured using Raw Luminescence Units (RLU), where the RLU of untreated cells were set as 100% reference values and all other RLU values were calculated relative to the reference values (referred to as Normalized RLU). The results in FIG. 18A show that the anti-SEZ6 antibody drug conjugate, hSC17.200-ADC1 suppressed the growth of thyroid cancer cells significantly more than the human IgG1-ADC1 control.

[0639] The ability of humanized anti-SEZ6 ADCs to reduce medullary thyroid tumor volumes *in vivo* was also tested. A thyroid cell line (ATCC, CRL-1803) was cultured in a 96 well plate at 500 cells per well in F-12K Medium supplemented with 10% FBS at 37 °C. Cells were harvested and implanted subcutaneously in five NOD/SCID mice. Once the tumors in the mice reached an average of 200 mm³ a single dose of 2 mg/kg hSC17.200-ADC1 was administered via intraperitoneal injection. Efficacy was monitored by weekly tumor volume measurements, with mean tumor volumes and standard error mean (SEM) plotted versus time (days). Following administration of hSC17.200-ADC1 a significant reduction in tumor volume was observed (FIG. 18B).

[0640] The ability of humanized anti-SEZ6 ADCs to reduce SCLC tumor volume was also tested. Anti-SEZ6 ADCs (hSC17.17, hSC17.24, hSC17.34 and hSC17.46), and the human IgG1 isotype control ADC (hulgG1) were administered to immunodeficient mice bearing various patient derived NTX tumors. Mice were given three injections of 1 mg/kg (indicated by the vertical lines in FIGS. 18C and 18D) over a period of seven days. Following treatment, tumor volumes and mouse weights were monitored until tumors exceeded 800 mm³ or mice became sick. The results of these experiments are presented in FIGS. 18C and 18D. Complete and durable elimination of tumor mass was achieved by the administration of humanized anti-SEZ6 ADCs in four SCLC tumors. FIG. 18C shows reduction of the LU80 tumor by hSC17.17-ADC1 and hSC17.46-ADC1; and elimination of the LU64 tumor by hSC17.17-ADC1, hSC17.34-ADC1 and hSC17.46-ADC1. FIG. 18D shows reduction of the LU117 tumor by hSC17.17-ADC1 and hSC17.46-ADC1; and reduction of the LU111 tumor by hSC17.34-ADC1 and hSC17.46-ADC1.

[0641] These results demonstrate the surprising applicability of a variety of humanized SEZ6 modulators to effectively retard the growth of different tumors.

EXAMPLE 20

GENERATION OF A CHEMORESISTANT PDX CELL LINE

[0642] A chemoresistant SCLC line was generated to test the expression of SEZ6 in tumors that had already undergone a first line of chemotherapeutic treatment. The chemoresistant cell line was developed from a SCLC patient-derived xenograft (PDX) tumor cell line obtained from a PDX tumor bank that was generated and maintained using art-recognized techniques. The PDX tumor bank comprises a substantial number of discrete tumor cell lines that were propagated in immunocompromised mice through multiple passages of heterogeneous tumor cells originally obtained from numerous cancer patients afflicted by a variety of solid tumor malignancies. The passage number of the tested sample is indicated by p0-p# where p0 is indicative of an unpassaged sample obtained directly from a patient tumor and p# is indicative of the number of times the tumor has been passaged through a mouse prior to testing. Early passage PDX tumors respond to therapeutic agents such as irinotecan (i.e. Camptosar®) and cisplatin/etoposide regimens, providing clinically relevant insights into underlying mechanisms driving tumor growth, resistance to current therapies and tumor recurrence.

[0643] Cisplatin and its derivatives, including oxaliplatin are the chemotherapeutic standard of care for SCLC. An oxaliplatin-resistant SCLC PDX line was developed due to concerns that cisplatin may not be stable in tissue culture medium (Schuldes et al., 1997, PMID: 9128988). The oxaliplatin-resistant SCLC line was generated by isolating human cells from LU124p2, a SCLC PDX line that had been passaged subcutaneously twice through NOD.SCID mice. LU124p2 tumors were resected from mice after reaching 800 - 2,000 mm³, dissociated into single cell suspensions using art-recognized enzymatic digestion techniques (see, for example, US2007/0292414), and depleted of murine cells. The cells were then washed and cultured in 5% oxygen and serum-free medium and immediately treated with 1 μM of oxaliplatin. After seven days of exposure to oxaliplatin, the cells were washed and allowed to recover for two weeks in standard serum-free media with fresh renewal of medium twice a week. After the recovery period, the cells were washed and re-plated into a new flask with 1 μM of oxaliplatin for seven more days, followed by a final wash and a two week recovery period. The resulting cell line was termed LU124OXAHlp2. Single cell suspensions were generated using Versene (Invitrogen). Some of the cells were injected into mice to propagate the LU124OXAHlp2 cell line, while other cells were cultured *in vitro* to test sensitivity to oxaliplatin.

[0644] Oxaliplatin sensitivity of LU124OXAHlp2 was tested as follows: LU124OXAHlp2 cells were plated onto 96-well plates and treated with a titration of oxaliplatin in serum-free medium as is standard in the art. After seven days in culture,

cells were harvested using Cell Titer Glo® (Promega) as per the manufacturer's instructions and dose response curves were obtained. LU124OXAHlp2 was more resistant to higher doses of oxaliplatin ($IC_{50} = 2\text{--}2.5\text{ }\mu\text{M}$) compared to the parental cell line (LU124p2) ($IC_{50} = 0.036\text{ }\mu\text{M}$) (FIG. 19). To determine whether the resistance was stable, LU124OXAHlp2 cells were passaged again through immunocompromised mice to yield the cell line called LU124OXAHlp3. Resistance was maintained in LU124OXAHlp3, which shows that even in the absence of selective pressure i.e. even when oxaliplatin is absent, resistance of the cell line was maintained.

EXAMPLE 21

SEZ6 EXPRESSION IN CHEMORESISTANT TUMOR CELLS USING MICROARRAY

[0645] Microarray expression profiling was used to identify potential cell surface targets that were upregulated on LU124OXAHlp3 compared to the chemo-sensitive parental line. In preparation for the assay, tumors from the LU124p3 and LU124OXAHlp3 PDX lines were resected, dissociated into single cell suspensions and depleted of murine cells as described in Example 1. The cells were lysed in RLTplus RNA lysis buffer per the manufacturer's instructions, stored at $-80\text{ }^{\circ}\text{C}$ and thawed for mRNA extraction. Upon thawing total mRNA was extracted using an RNeasy isolation kit (Qiagen) and quantified using a Nanodrop spectrophotometer (Thermo Scientific) and/or a Bioanalyzer 2100 (Agilent Technologies), using the manufacturer's protocols and recommended instrument settings. The resulting total mRNA preparations were assessed for suitability for genetic sequencing and gene expression analysis.

[0646] 1-2 μg of whole tumor total mRNA samples were analyzed using the Agilent SurePrint GE Human 8x60 v2 microarray platform which contains 50,599 biological probes designed against 27,958 genes and 7,419 lncRNAs in the human genome. Standard industry practices were used to normalize and transform the intensity values to quantify gene expression for each sample. SEZ6 mRNA (Agilent probe ID: A_23_P49849) was identified as being upregulated in LU124OXAHlp3 compared to the matched parental line, LU124p3, and also the parental line from an earlier murine passage, LU124p1 (FIG. 20).

EXAMPLE 22

SEZ6 PROTEIN EXPRESSION IN A CHEMORESISTANT TUMOR CELL LINE

[0647] Given the elevated SEZ6 mRNA transcript levels associated with the oxaliplatin-resistant cell line generated as described in Example 1, work was undertaken to test whether SEZ6 protein expression was also elevated in that cell line. To detect and quantify SEZ6 protein expression, an electrochemiluminescence SEZ6 sandwich ELISA assay was developed using the MSD Discovery Platform (Meso Scale Discovery).

[0648] LU124p3 cells and LU124OXAHlp3 cells were flash frozen on dry ice/ethanol. Protein Extraction Buffer (Biochain Institute) was added to the thawed cells, lysates were cleared by centrifugation (20,000g, 20 minutes, $4\text{ }^{\circ}\text{C}$) and the total protein concentration in each lysate was quantified using bicinchoninic acid. Protein lysates were stored at $-80\text{ }^{\circ}\text{C}$ until assayed. SEZ6 protein concentrations from the lysate samples were determined by interpolating the values from a standard protein concentration curve that was generated using purified recombinant SEZ6 protein. The SEZ6 protein standard curve and protein quantification assay were conducted as follows:

MSD standard plates were coated overnight at $4\text{ }^{\circ}\text{C}$ with 30 μL of the anti-SEZ6 antibody SC17.17 at 1 $\mu\text{g}/\text{mL}$ in phosphate buffered saline (PBS). Plates were washed in PBST and blocked in 150 μL MSD 3% Blocker A solution for one hour while shaking. Plates were again washed in PBST. 25 μL of 10x diluted lysate (or serially diluted recombinant SEZ6 standard) in MSD 1% Blocker A containing 10% Protein Extraction Buffer was also added to the wells and incubated for two hours while shaking. Plates were again washed in PBST. The anti-SEZ6 antibody SC17.36 antibody was sulfo-tagged and 25 μL of the tagged SC17.36 antibody was added to the washed plates at 0.5 $\mu\text{g}/\text{mL}$ in MSD 1% Blocker A for 1 hour at room temperature while shaking. Plates were washed in PBST. MSD Read Buffer T with surfactant was diluted to 1X in water and 150 μL was added to each well. Plates were read on an MSD Sector Imager 2400 using an integrated software analysis program to derive SEZ6 concentrations in PDX samples via interpolation from the standard curve. Values were then divided by total protein concentration to yield nanograms of SEZ6 per milligram of total lysate protein. The resulting concentrations are set forth in FIG. 21 wherein the bar graph represents SEZ6 protein concentrations derived from LU124p3 and the LU124OXAHlp3 cell line. Expression of SEZ6 in LU124 cell lysate was 5.8 ng per mg total protein while expression of SEZ6 in the LU124 oxaliplatin resistant line was 14.6 ng per mg total protein. FIG. 21 shows that the oxaliplatin-resistant LU124OXAHlp3 cell line exhibited high SEZ6 protein expression compared to parental LU124p3 cells. These data, combined with the mRNA transcription data for SEZ6 expression set forth in the examples above strongly reinforces the proposition that SEZ6 protein expression is upregulated in tumors exhibiting chemoresistance.

[0649] Those skilled in the art will further appreciate that the present invention may be embodied in other specific forms without departing from the spirit or central attributes thereof. In that the foregoing description of the present invention discloses only exemplary embodiments thereof, it is to be understood that other variations are contemplated as being within the scope of the present invention. Accordingly, the present invention is not limited to the particular embodiments that have been described in detail herein. Rather, reference should be made to the appended claims as indicative of the scope and content of the invention. Preferred embodiments of the present invention are as described below and referred to as embodiments E1 to E19.

E1. An isolated antibody that specifically binds to an epitope on a SEZ6 protein, wherein the epitope comprises amino acid residues selected from the group consisting of (i) residues R762, L764, Q777, I779, D781 and Q782; (ii) residues R342 and K389 and (iii) residues T352, S353 and H375.

E2. An antibody drug conjugate comprising an antibody conjugated directly or indirectly to a therapeutic moiety, wherein the antibody specifically binds to an epitope on a SEZ6 protein, wherein the epitope comprises amino acid residues selected from the group consisting of (i) residues R762, L764, Q777, I779, D781 and Q782; (ii) residues R342 and K389 and (iii) residues T352, S353 and H375.

E3. The antibody drug conjugate of E2, wherein the therapeutic moiety is selected from the group consisting of auristatins, amanitins and pyrrolobenzodiazepines.

E4. A method of treating a subject suffering from cancer comprising administering a therapeutically effective amount of an antibody drug conjugate comprising an antibody conjugated directly or indirectly to a therapeutic moiety, wherein the antibody specifically binds to an epitope on a SEZ6 protein, wherein the epitope comprises amino acid residues selected from the group consisting of (i) residues R762, L764, Q777, I779, D781 and Q782; (ii) residues R342 and K389 and (iii) residues T352, S353 and H375.

E5. The method of E4, wherein the therapeutic moiety is selected from the group consisting of auristatins, amanitins and pyrrolobenzodiazepines.

E6. The method of E4, wherein the subject has previously been treated with a platinum based agent.

E7. A method of treating a subject suffering from platinum resistant small cell lung cancer comprising administering a therapeutically effective amount of an antibody drug conjugate comprising an antibody conjugated directly or indirectly to a therapeutic moiety, wherein the antibody specifically binds to an epitope on a SEZ6 protein, wherein the epitope comprises amino acid residues selected from the group consisting of (i) residues R762, L764, Q777, I779, D781 and Q782; (ii) residues R342 and K389 and (iii) residues T352, S353 and H375.

E8. The method of E7, wherein the therapeutic moiety is selected from the group consisting of auristatins, amanitins and pyrrolobenzodiazepines.

E9. A method of treating a patient suffering from medullary thyroid cancer comprising administering a therapeutically effective amount of an antibody drug conjugate comprising an antibody conjugated directly or indirectly to a therapeutic moiety, wherein the antibody specifically binds to an epitope on a SEZ6 protein.

E10. A method of E9, wherein the epitope comprises amino acid residues selected from the group consisting of (i) residues R762, L764, Q777, I779, D781 and Q782; (ii) residues R342 and K389 and (iii) residues T352, S353 and H375.

E11. The method of E9, wherein the therapeutic moiety selected from the group consisting of auristatins, amanitins and pyrrolobenzodiazepines.

E12. A method of diagnosing platinum resistant small cell lung cancer comprising the steps of:

- a. providing a platinum resistant small cell lung cancer tumor sample from a subject;
- b. exposing the tumor sample to an anti-SEZ6 antibody labeled with a reporter wherein said anti-SEZ6 antibody associates with the tumor sample; and
- c. detecting the reporter associated with the tumor sample.

E13. The method of E12 wherein the reporter is detected *in vitro*.

E14. The method of E12 wherein the reporter is detected using immunohistochemistry.

E15. The method of any one of E12 to E14 wherein the anti-SEZ6 antibody binds to an epitope on a SEZ6 protein, wherein the epitope comprises amino acid residues R762, L764, Q777, I779, D781 and Q782.

E16. A method of diagnosing medullary thyroid cancer comprising the steps of:

- a. providing a medullary thyroid tumor sample from a subject;
- b. exposing the tumor sample to an anti-SEZ6 antibody labeled with a reporter wherein said anti-SEZ6 antibody associates with the tumor sample; and
- c. detecting the reporter associated with the tumor sample.

EP 3 338 793 A1

E17. The method of E16 wherein the reporter is detected *in vitro*.

E18. The method of E16 wherein the reporter is detected using immunohistochemistry.

E19. The method of any one of E16 to E18 wherein the anti-SEZ6 antibody binds to an epitope on a SEZ6 protein, wherein the epitope comprises amino acid residues R762, L764, Q777, I779, D781 and Q782.

5

10

15

20

25

30

35

40

45

50

55

SEQUENCE LISTING

<110> Stem CentRx, LLC

<120> NOVEL SEZ6 MODULATORS AND METHODS OF USE

<130> S69697 1130WO/sc1702.pct

<150> 61/871,289

<151> 2013-08-28

<160> 480

<170> PatentIn version 3.5

<210> 1

<211> 4249

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> SEZ6 isoform 1 mRNA

<400> 1

gatccccggc gccgtcgcca ggcgctggcc gtggtgctga ttctgtcagg cgctggcggc	60
ggcagcggcg gtgacggctg cggccccgct ccctctaccc ggccggaccc ggctctgccc	120
ccgcgcccac gccccaccaa gccccccgcc ctcccgcgcg ggtcccagcc cagggcgogg	180
ccgcaaccag caccatgcgc ccggtagccc tgctgctcct gccctcgctg ctggcgctcc	240
tggctcacgg actctcttta gaggcccaa ccgtggggaa aggacaagcc ccaggcatcg	300
aggagacaga tggcgagctg acagcagccc ccacacctga gcagccagaa cgaggcgtcc	360
actttgtcac aacagcccc accttgaagc tgctcaacca ccaccgctg cttgaggaat	420
tcctacaaga ggggctggaa aaggagatg aggagctgag gccagcactg cccttcacagc	480
ctgaccaccc tgcacccttc accccaagtc cccttccccg cctggccaac caggacagcc	540
gccctgtctt taccagcccc actccagcca tggctgcggt acccactcag cccagtcaca	600
aggagggacc ctggagtccg gagtcaagc ccctatgct tcgaatcaca gctcccctac	660
ctccagggcc cagcatggca gtgcccaccc taggcccagg ggagatagcc agcactacac	720
ccccagcag agcctggaca ccaaccacag agggctcctg agacatggga aggccgtggg	780
ttgcagaggt tgtgtcccag ggcgcaggga tcgggatcca ggggaccatc acctcctcca	840
cagcttcagg agatgatgag gagaccacca ctaccaccac catcatcacc accaccatca	900
ccacagtcca gacaccaggc ccttgtagct ggaatttctc aggccagag ggctctctgg	960
actcccctac agacctcagc tccccactg atgttggcct ggactgcttc ttctacatct	1020
ctgtctaccc tggctatggc gtggaaatca aggtccagaa tatcagcctc cgggaagggg	1080
agacagtgac tgtggaaggc ctgggggggc ctgaccact gccctggcc aaccagtctt	1140

EP 3 338 793 A1

	tcctgctgcg	gggccaagtc	atccgcagcc	ccaccaccca	agcggccctg	aggttccaga	1200
	gcctccccgc	accggctggc	cctggcacct	tccatttcca	ttaccaagcc	tatctcctga	1260
5	gctgccactt	tccccgtcgt	ccagcttatg	gagatgtgac	tgtcaccagc	ctccaccag	1320
	ggggtagtgc	ccgcttccat	tgtgccactg	gctaccagct	gaagggcgcc	aggcatctca	1380
	cctgtctcaa	tgccaccag	cccttctggg	attcaaagga	gcccgtctgc	atcgtgctt	1440
10	gcggcgaggt	gatccgcaat	gccaccaccg	gccgcacgt	ctctccaggc	ttcccgggca	1500
	actacagcaa	caacctcacc	tgtcactggc	tgtttgaggg	tcctgagggc	cagcggctac	1560
	acctgcactt	tgagaagggt	tccttggcag	aggatgatga	caggctcatc	attcgcaatg	1620
15	gggacaacgt	ggaggcccca	ccagtgtatg	attcctatga	ggtggaatac	ctgccattg	1680
	agggcctgct	cagctctggc	aaacacttct	ttgttgagct	cagtactgac	agcagcgggg	1740
20	cagctgcagg	catggccctg	cgctatgagg	ccttccagca	gggccattgc	tatgagccct	1800
	ttgtcaaata	cggtaacttc	agcagcagca	caccaccta	ccctgtgggt	accactgtgg	1860
	agttcagctg	cgaccctggc	tacaccctgg	agcagggctc	catcatcatc	gagtgtgttg	1920
25	acccccacga	ccccagtggt	aatgagacag	agccagcctg	ccgagccgtg	tgacaggggg	1980
	agatcacaga	ctcggtggc	gtggtactct	ctcccaactg	gccagagccc	tacggctcgtg	2040
	ggcaggattg	tatctgggggt	gtgcatgtgg	aagaggacaa	gcgcatcatg	ctggacatcc	2100
30	gagtgcctgc	cataggccct	ggtgatgtgc	ttaccttcta	tgatggggat	gacctgacgg	2160
	cccgggttct	gggccagtac	tcagggcccc	gtagccactt	caagctcttt	acctccatgg	2220
35	ctgatgtcac	cattcagttc	cagtcggacc	ccgggacctc	agtgcctggc	taccagcagg	2280
	gcttcgtcat	ccacttcttt	gaggtgcccc	gcaatgacac	atgtccggag	ctgcctgaga	2340
	tccccaatgg	ctggaagagc	ccatcgcagc	ctgagctagt	gcacggcacc	gtggtcactt	2400
40	accagtgcga	ccctggctac	caggtagtgg	gatccagtg	cctcatgtgc	cagtgggacc	2460
	taacttggag	tgaggacctg	ccctcatgcc	agaggggtgac	ttcctgccac	gatcctggag	2520
	atgtggagca	cagccgacgc	ctcatatcca	gccccaaagt	tcccgtgggg	gccaccgtgc	2580
45	aatatatctg	tgaccaggggt	tttgtgctga	tgggcagctc	catcctcacc	tgccatgatc	2640
	gccaggctgg	cagccccaaag	tgagtgacc	ggggccctaa	atgtctcctg	gaacagctca	2700
	agccatgccca	tgggtctcagt	gcccctgaga	atgggtgccc	aagtcctgag	aagcagctac	2760
50	accagcagg	ggccaccatc	cacttctcgt	gtgcccctgg	ctatgtgctg	aagggccagg	2820
	ccagcatcaa	gtgtgtgcct	gggcaccct	cgcattggag	tgacccccca	cccatctgta	2880
55	gggctgcctc	tctggatggg	ttctacaaca	gtcgcagcct	ggatgttgcc	aaggcacctg	2940
	ctgcctccag	cacctggat	gctgccaca	ttgcagctgc	catcttcttg	ccactgggtg	3000

EP 3 338 793 A1

	cgatggtggtt gttggttagga ggtgtatact tctacttctc caggctccag ggaaaaagct	3060
	ccctgcagct gccccgcccc cgcccccgcc cctacaaccg cattaccata gagtcagcgt	3120
5	ttgacaatcc aacttacgag actggatctc tttcctttgc aggagacgag agaatatgaa	3180
	gtctccatct aggtgggggc agtctagga agtcaactca gacttgcacc acagtccagc	3240
	agcaaggctc cttgcttcct gctgtccctc cacctcctgt atataccacc taggaggaga	3300
10	tgccaccaag ccctcaagaa gttgtgccct tccccgcctg cgatgccac catggcctat	3360
	tttcttggtg tcattgccca cttggggccc ttcattgggc ccatgtcagg gggcatctac	3420
	ctgtgggaag aacatagctg gagcacaagc atcaacagcc agcatcctga gcctcctcat	3480
15	gccctggacc agcctggaac acactagcag agcaggagta cctttctcca catgaccacc	3540
	atccccccct ggcattggcaa cctgcagcag gattaacttg accatggtgg gaactgcacc	3600
	agggtactcc tcacagcgca tcaccaatgg ccaaaaactcc tctcaacggt gacctctggg	3660
20	tagtcctggc atgccaacat cagcctcttg ggaggtctct agttctctaa agttctggac	3720
	agttctgcct cctgccctgt cccagtggag gcagtaattc taggagatcc taaggggttc	3780
25	aggggggacc tacccccacc tcagggttggg cttccctggg cactcatgct ccacacccaa	3840
	gcaggacacg ccattttcca ctgaccaccc tataccctga ggaaaggag actttcctcc	3900
	gatgtttatt tagctgttgc aaacatcttc accctaatag tccctcctcc aattccagcc	3960
30	acttgtcagg ctctcctctt gaccactgtg ttatgggata aggggagggg gtgggcatat	4020
	tctggagagg agcagaggtc caaggaccca ggaatttggc atggaacagg tggtaggaga	4080
	gccccaggga gacgccag agctggctga aagccacttt gtacatgtaa tgtattatat	4140
35	ggggtctggg ctccagccag agaacaatct tttatttctg ttgtttcctt attaaaatgg	4200
	tgttttttga aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa	4249
40	<210> 2 <211> 4234 <212> DNA <213> Homo sapiens	
45	<220> <221> misc_feature <223> SEZ6 isoform 2 mRNA	
50	<400> 2 gatccccggc gccgtcgcca ggcgctggcc gtggtgctga ttctgtcagg cgctggcggc	60
	ggcagcggcg gtgacggctg cggccccgct ccctctaccc ggccggaccc ggctctgccc	120
	ccgcgcccac gcccaccaa gcccccgcc ctcccgccgc ggtcccagcc cagggcgcg	180
55	ccgcaaccag caccatgcgc ccggtagccc tgctgctcct gccctcgtg ctggcgctcc	240
	tggctcacgg actctcttta gaggcccaa ccgtggggaa aggacaagcc ccaggcatcg	300

EP 3 338 793 A1

	aggagacaga	tggcgagctg	acagcagccc	ccacacctga	gcagccagaa	cgaggcgctcc	360
	actttgtcac	aacagccccc	accttgaagc	tgctcaacca	ccaccgcgtg	cttgaggaat	420
5	tcctacaaga	ggggctggaa	aaggagatg	aggagctgag	gccagcactg	cccttccagc	480
	ctgaccacc	tgcacccttc	acccaagtc	cccttccccg	cctggccaac	caggacagcc	540
	gccctgtctt	taccagcccc	actccagcca	tggctgcggt	accactcag	ccccagtcca	600
10	aggagggacc	ctggagtccg	gagtcagagt	cccctatgct	tcgaatcaca	gctcccctac	660
	ctccagggcc	cagcatggca	gtgcccaccc	taggcccagg	ggagatagcc	agcactacac	720
	ccccagcag	agcctggaca	ccaacccaag	agggctcctg	agacatggga	aggccgtggg	780
15	ttgcagaggt	tgtgtcccag	ggcgcaggga	tcgggatcca	ggggaccatc	acctcctcca	840
	cagcttcagg	agatgatgag	gagaccacca	ctaccaccac	catcatcacc	accaccatca	900
20	ccacagtcca	gacaccaggc	ccttgtagct	ggaatttctc	aggcccagag	ggctctctgg	960
	actcccctac	agacctcagc	tccccactg	atgttggcct	ggactgcttc	ttctacatct	1020
	ctgtctaccc	tggctatggc	gtggaaatca	aggtccagaa	tatcagcctc	cgggaagggg	1080
25	agacagtgac	tgtggaaggc	ctgggggggc	ctgaccact	gccctggcc	aaccagtctt	1140
	tcctgctgcg	gggccaagtc	atccgcagcc	ccaccacca	agcggccctg	aggttccaga	1200
	gcctcccgcc	accggctggc	cctggcacct	tccatttcca	ttaccaagcc	tatctcctga	1260
30	gctgccactt	tccccgtcgt	ccagcttatg	gagatgtgac	tgtcaccagc	ctccaccag	1320
	ggggtagtgc	ccgcttccat	tgtgccactg	gctaccagct	gaagggcgcc	aggcatctca	1380
35	cctgtctcaa	tgccaccag	cccttctggg	attcaaagga	gccgctctgc	atcgctgctt	1440
	gcgggcgagt	gatccgcaat	gccaccaccg	gccgcacgt	ctctccaggc	ttcccgggca	1500
	actacagcaa	caacctcacc	tgtcactggc	tgcttgaggc	tcctgagggc	cagcggctac	1560
40	acctgcactt	tgagaagggt	tccttggcag	aggatgatga	caggctcatc	attcgcaatg	1620
	gggacaacgt	ggaggcccca	ccagtgtatg	attcctatga	ggtggaatac	ctgccattg	1680
	agggcctgct	cagctctggc	aaacacttct	ttgttgagct	cagtactgac	agcagcgggg	1740
45	cagctgcagg	catggccctg	cgctatgagg	ccttcagca	gggccattgc	tatgagccct	1800
	ttgtcaaata	cggtaacttc	agcagcagca	caccaccta	ccctgtgggt	accactgtgg	1860
	agttcagctg	cgaccctggc	tacaccctgg	agcagggctc	catcatcatc	gagtgtgttg	1920
50	acccccacga	ccccagtg	aatgagacag	agccagcctg	ccgagccgtg	tgcagcgggg	1980
	agatcacaga	ctcggtggc	gtggtactct	ctcccaactg	gccagagccc	tacggctcgtg	2040
55	ggcaggattg	tatctggggg	gtgcatgtgg	aagaggacaa	gcgcatcatg	ctggacatcc	2100
	gagtgtgctg	cataggccct	ggtgatgtgc	ttaccttcta	tgatggggat	gacctgacgg	2160

EP 3 338 793 A1

	cccgggttct	gggccagtac	tcagggcccc	gtagccactt	caagctcttt	acctccatgg	2220
	ctgatgtcac	cattcagttc	cagtcggacc	ccgggacctc	agtgtctggg	taccagcagg	2280
5	gcttcgtcat	ccacttcttt	gaggtgcccc	gcaatgacac	atgtccggag	ctgcctgaga	2340
	tccccaatgg	ctggaagagc	ccatcgcagc	ctgagctagt	gcacggcacc	gtggtcactt	2400
	accagtgcta	ccctggctac	caggtagtgg	gatccagtgt	cctcatgtgc	cagtgggacc	2460
10	taacttggag	tgaggacctg	ccctcatgcc	agaggggtgac	ttcctgccac	gatcctggag	2520
	atgtggagca	cagccgacgc	ctcatatcca	gccccaaagt	tcccgtgggg	gccaccgtgc	2580
	aatatatctg	tgaccaggg	tttgtgtctga	tgggcagctc	catcctcacc	tgccatgac	2640
15	gccaggctgg	cagccccaag	tggagtgacc	gggcccctaa	atgtctcctg	gaacagctca	2700
	agccatgcca	tgggtctcagt	gcccctgaga	atggtgcccc	aagtcctgag	aagcagctac	2760
	accagcagg	ggccaccatc	cacttctcgt	gtgcccctgg	ctatgtgctg	aagggccagg	2820
20	ccagcatcaa	gtgtgtgcct	gggcacccct	cgcattggag	tgacccccca	cccatctgta	2880
	gggctgcctc	tctggatggg	ttctacaaca	gtcgcagcct	ggatgttgcc	aaggcacctg	2940
25	ctgcctccag	caccctggat	gtgcccaca	ttgcagctgc	catcttcttg	ccactggtgg	3000
	cgatggtgtt	gttggtagga	ggtgtatact	tctacttctc	caggctccag	ggaaaaagct	3060
	ccctgcagct	gccccgcccc	cgcgcccgcc	cctacaaccg	cattaccata	gagtcagcgt	3120
30	ttgacaatcc	aacttacgag	actggagaga	cagagagaata	tgaagtctcc	atctaggtgg	3180
	gggcagtcta	gggaagtcaa	ctcagacttg	caccacagtc	cagcagcaag	gtccttgct	3240
	tcctgctgtc	cctccacctc	ctgtatatac	cacctaggag	gagatgccac	caagccctca	3300
35	agaagtgtg	cccttccccg	cctgcgatgc	ccaccatggc	ctattttctt	ggtgtcattg	3360
	cccacttggg	gcccttcatt	gggcccattg	cagggggcat	ctacctgtgg	gaagaacata	3420
	gctggagcac	aagcatcaac	agccagcatc	ctgagcctcc	tcatgccctg	gaccagcctg	3480
40	gaacacacta	gcagagcagg	agtaccttct	tccacatgac	caccatcccc	ccctggcatg	3540
	gcaacctgca	gcaggattaa	cttgaccatg	gtgggaactg	caccagggta	ctcctcacag	3600
	cgccatcacc	aatggccaaa	actcctctca	acggtgacct	ctgggtagtc	ctggcatgcc	3660
45	aacatcagcc	tcttgggagg	tctctagtct	tctaaagttc	tggacagttc	tgctcctgc	3720
	cctgtcccag	tggaggcagt	aattctagga	gatcctaagg	ggttcagggg	gacctaccc	3780
	ccacctcagg	ttgggcttcc	ctgggcactc	atgctccaca	caaagcagg	acacgccatt	3840
50	ttccactgac	caccctatac	cctgaggaaa	gggagacttt	cctccgatgt	ttatttagct	3900
	gttgcaaaca	tcttcaccct	aatagtccct	cctccaattc	cagccacttg	tcaggctctc	3960
55	ctcttgacca	ctgtgttatg	ggataagggg	aggggggtgg	catattctgg	agaggagcag	4020
	aggtccaagg	accaggaat	ttggcatgga	acagggtgga	ggagagcccc	agggagacgc	4080

EP 3 338 793 A1

ccaggagctg gctgaaagcc actttgtaca tgtaatgtat tatatgggggt ctgggctcca 4140

gccagagaac aatcttttat ttctgttgtt tccttattaa aatggtgttt ttggaaaaaa 4200

5 aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaa 4234

<210> 3

<211> 994

<212> PRT

10 <213> Homo sapiens

<220>

<221> MISC_FEATURE

15 <223> SEZ6 isoform 1 precursor protein

<220>

<221> MISC_FEATURE

<222> (1) .. (19)

20 <223> Leader sequence

<220>

<221> MISC_FEATURE

<222> (20) .. (994)

25 <223> Mature protein

<400> 3

Met Arg Pro Val Ala Leu Leu Leu Leu Pro Ser Leu Leu Ala Leu Leu

1 5 10 15

30 Ala His Gly Leu Ser Leu Glu Ala Pro Thr Val Gly Lys Gly Gln Ala

20 25 30

35 Pro Gly Ile Glu Glu Thr Asp Gly Glu Leu Thr Ala Ala Pro Thr Pro

35 40 45

Glu Gln Pro Glu Arg Gly Val His Phe Val Thr Thr Ala Pro Thr Leu

50 55 60

40

Lys Leu Leu Asn His His Pro Leu Leu Glu Glu Phe Leu Gln Glu Gly

65 70 75 80

45 Leu Glu Lys Gly Asp Glu Glu Leu Arg Pro Ala Leu Pro Phe Gln Pro

85 90 95

Asp Pro Pro Ala Pro Phe Thr Pro Ser Pro Leu Pro Arg Leu Ala Asn

50 100 105 110

Gln Asp Ser Arg Pro Val Phe Thr Ser Pro Thr Pro Ala Met Ala Ala

115 120 125

55 Val Pro Thr Gln Pro Gln Ser Lys Glu Gly Pro Trp Ser Pro Glu Ser

EP 3 338 793 A1

	130		135		140
5	Glu Ser Pro Met Leu Arg Ile Thr Ala Pro Leu Pro Pro Gly Pro Ser	145	150	155	160
10	Met Ala Val Pro Thr Leu Gly Pro Gly Glu Ile Ala Ser Thr Thr Pro	165	170	175	
15	Pro Ser Arg Ala Trp Thr Pro Thr Gln Glu Gly Pro Gly Asp Met Gly	180	185	190	
20	Arg Pro Trp Val Ala Glu Val Val Ser Gln Gly Ala Gly Ile Gly Ile	195	200	205	
25	Gln Gly Thr Ile Thr Ser Ser Thr Ala Ser Gly Asp Asp Glu Glu Thr	210	215	220	
30	Thr Thr Thr Thr Thr Ile Ile Thr Thr Thr Ile Thr Thr Val Gln Thr	225	230	235	240
35	Pro Gly Pro Cys Ser Trp Asn Phe Ser Gly Pro Glu Gly Ser Leu Asp	245	250	255	
40	Ser Pro Thr Asp Leu Ser Ser Pro Thr Asp Val Gly Leu Asp Cys Phe	260	265	270	
45	Phe Tyr Ile Ser Val Tyr Pro Gly Tyr Gly Val Glu Ile Lys Val Gln	275	280	285	
50	Asn Ile Ser Leu Arg Glu Gly Glu Thr Val Thr Val Glu Gly Leu Gly	290	295	300	
55	Gly Pro Asp Pro Leu Pro Leu Ala Asn Gln Ser Phe Leu Leu Arg Gly	305	310	315	320
60	Gln Val Ile Arg Ser Pro Thr His Gln Ala Ala Leu Arg Phe Gln Ser	325	330	335	
65	Leu Pro Pro Pro Ala Gly Pro Gly Thr Phe His Phe His Tyr Gln Ala	340	345	350	
70	Tyr Leu Leu Ser Cys His Phe Pro Arg Arg Pro Ala Tyr Gly Asp Val	355	360	365	
75	Thr Val Thr Ser Leu His Pro Gly Gly Ser Ala Arg Phe His Cys Ala	370	375	380	

EP 3 338 793 A1

	Thr	Gly	Tyr	Gln	Leu	Lys	Gly	Ala	Arg	His	Leu	Thr	Cys	Leu	Asn	Ala	385	390	395	400
5	Thr	Gln	Pro	Phe	Trp	Asp	Ser	Lys	Glu	Pro	Val	Cys	Ile	Ala	Ala	Cys	405	410	415	
10	Gly	Gly	Val	Ile	Arg	Asn	Ala	Thr	Thr	Gly	Arg	Ile	Val	Ser	Pro	Gly	420	425	430	
	Phe	Pro	Gly	Asn	Tyr	Ser	Asn	Asn	Leu	Thr	Cys	His	Trp	Leu	Leu	Glu	435	440	445	
15	Ala	Pro	Glu	Gly	Gln	Arg	Leu	His	Leu	His	Phe	Glu	Lys	Val	Ser	Leu	450	455	460	
20	Ala	Glu	Asp	Asp	Asp	Arg	Leu	Ile	Ile	Arg	Asn	Gly	Asp	Asn	Val	Glu	465	470	475	480
	Ala	Pro	Pro	Val	Tyr	Asp	Ser	Tyr	Glu	Val	Glu	Tyr	Leu	Pro	Ile	Glu	485	490	495	
25	Gly	Leu	Leu	Ser	Ser	Gly	Lys	His	Phe	Phe	Val	Glu	Leu	Ser	Thr	Asp	500	505	510	
30	Ser	Ser	Gly	Ala	Ala	Ala	Gly	Met	Ala	Leu	Arg	Tyr	Glu	Ala	Phe	Gln	515	520	525	
35	Gln	Gly	His	Cys	Tyr	Glu	Pro	Phe	Val	Lys	Tyr	Gly	Asn	Phe	Ser	Ser	530	535	540	
	Ser	Thr	Pro	Thr	Tyr	Pro	Val	Gly	Thr	Thr	Val	Glu	Phe	Ser	Cys	Asp	545	550	555	560
40	Pro	Gly	Tyr	Thr	Leu	Glu	Gln	Gly	Ser	Ile	Ile	Ile	Glu	Cys	Val	Asp	565	570	575	
45	Pro	His	Asp	Pro	Gln	Trp	Asn	Glu	Thr	Glu	Pro	Ala	Cys	Arg	Ala	Val	580	585	590	
50	Cys	Ser	Gly	Glu	Ile	Thr	Asp	Ser	Ala	Gly	Val	Val	Leu	Ser	Pro	Asn	595	600	605	
	Trp	Pro	Glu	Pro	Tyr	Gly	Arg	Gly	Gln	Asp	Cys	Ile	Trp	Gly	Val	His	610	615	620	
55	Val	Glu	Glu	Asp	Lys	Arg	Ile	Met	Leu	Asp	Ile	Arg	Val	Leu	Arg	Ile	625	630	635	640

EP 3 338 793 A1

	Gly	Pro	Gly	Asp	Val	Leu	Thr	Phe	Tyr	Asp	Gly	Asp	Asp	Leu	Thr	Ala	
					645					650					655		
5	Arg	Val	Leu	Gly	Gln	Tyr	Ser	Gly	Pro	Arg	Ser	His	Phe	Lys	Leu	Phe	
				660					665					670			
	Thr	Ser	Met	Ala	Asp	Val	Thr	Ile	Gln	Phe	Gln	Ser	Asp	Pro	Gly	Thr	
10			675					680					685				
	Ser	Val	Leu	Gly	Tyr	Gln	Gln	Gly	Phe	Val	Ile	His	Phe	Phe	Glu	Val	
		690					695					700					
15	Pro	Arg	Asn	Asp	Thr	Cys	Pro	Glu	Leu	Pro	Glu	Ile	Pro	Asn	Gly	Trp	
	705					710					715					720	
	Lys	Ser	Pro	Ser	Gln	Pro	Glu	Leu	Val	His	Gly	Thr	Val	Val	Thr	Tyr	
20					725					730					735		
	Gln	Cys	Tyr	Pro	Gly	Tyr	Gln	Val	Val	Gly	Ser	Ser	Val	Leu	Met	Cys	
25				740					745					750			
	Gln	Trp	Asp	Leu	Thr	Trp	Ser	Glu	Asp	Leu	Pro	Ser	Cys	Gln	Arg	Val	
			755					760					765				
30	Thr	Ser	Cys	His	Asp	Pro	Gly	Asp	Val	Glu	His	Ser	Arg	Arg	Leu	Ile	
		770					775					780					
	Ser	Ser	Pro	Lys	Phe	Pro	Val	Gly	Ala	Thr	Val	Gln	Tyr	Ile	Cys	Asp	
35		785				790					795					800	
	Gln	Gly	Phe	Val	Leu	Met	Gly	Ser	Ser	Ile	Leu	Thr	Cys	His	Asp	Arg	
					805					810					815		
40	Gln	Ala	Gly	Ser	Pro	Lys	Trp	Ser	Asp	Arg	Ala	Pro	Lys	Cys	Leu	Leu	
				820					825					830			
	Glu	Gln	Leu	Lys	Pro	Cys	His	Gly	Leu	Ser	Ala	Pro	Glu	Asn	Gly	Ala	
45			835					840					845				
	Arg	Ser	Pro	Glu	Lys	Gln	Leu	His	Pro	Ala	Gly	Ala	Thr	Ile	His	Phe	
50		850					855					860					
	Ser	Cys	Ala	Pro	Gly	Tyr	Val	Leu	Lys	Gly	Gln	Ala	Ser	Ile	Lys	Cys	
	865					870					875					880	
55	Val	Pro	Gly	His	Pro	Ser	His	Trp	Ser	Asp	Pro	Pro	Pro	Ile	Cys	Arg	
				885						890					895		

EP 3 338 793 A1

	Ala	Ala	Ser	Leu	Asp	Gly	Phe	Tyr	Asn	Ser	Arg	Ser	Leu	Asp	Val	Ala	
				900					905					910			
5	Lys	Ala	Pro	Ala	Ala	Ser	Ser	Thr	Leu	Asp	Ala	Ala	His	Ile	Ala	Ala	
			915					920					925				
10	Ala	Ile	Phe	Leu	Pro	Leu	Val	Ala	Met	Val	Leu	Leu	Val	Gly	Gly	Val	
		930					935					940					
15	Tyr	Phe	Tyr	Phe	Ser	Arg	Leu	Gln	Gly	Lys	Ser	Ser	Leu	Gln	Leu	Pro	
	945					950					955					960	
20	Arg	Pro	Arg	Pro	Arg	Pro	Tyr	Asn	Arg	Ile	Thr	Ile	Glu	Ser	Ala	Phe	
					965					970					975		
25	Asp	Asn	Pro	Thr	Tyr	Glu	Thr	Gly	Ser	Leu	Ser	Phe	Ala	Gly	Asp	Glu	
				980					985					990			
30	Arg Ile																
35	<210> 4																
	<211> 993																
	<212> PRT																
	<213> Homo sapiens																
40	<220>																
	<221> MISC_FEATURE																
	<223> SEZ6 isoform 2 precursor protein																
45	<220>																
	<221> MISC_FEATURE																
	<222> (1) .. (19)																
	<223> Leader sequence																
50	<220>																
	<221> MISC_FEATURE																
	<222> (20) .. (993)																
	<223> Mature protein																
55	<400> 4																
	Met	Arg	Pro	Val	Ala	Leu	Leu	Leu	Leu	Pro	Ser	Leu	Leu	Ala	Leu	Leu	
	1				5					10				15			
60	Ala	His	Gly	Leu	Ser	Leu	Glu	Ala	Pro	Thr	Val	Gly	Lys	Gly	Gln	Ala	
				20					25					30			
65	Pro	Gly	Ile	Glu	Glu	Thr	Asp	Gly	Glu	Leu	Thr	Ala	Ala	Pro	Thr	Pro	
			35					40					45				

EP 3 338 793 A1

	Glu	Gln	Pro	Glu	Arg	Gly	Val	His	Phe	Val	Thr	Thr	Ala	Pro	Thr	Leu	
	50						55					60					
5	Lys	Leu	Leu	Asn	His	His	Pro	Leu	Leu	Glu	Glu	Phe	Leu	Gln	Glu	Gly	
	65					70					75					80	
	Leu	Glu	Lys	Gly	Asp	Glu	Glu	Leu	Arg	Pro	Ala	Leu	Pro	Phe	Gln	Pro	
10					85					90					95		
	Asp	Pro	Pro	Ala	Pro	Phe	Thr	Pro	Ser	Pro	Leu	Pro	Arg	Leu	Ala	Asn	
				100					105					110			
15	Gln	Asp	Ser	Arg	Pro	Val	Phe	Thr	Ser	Pro	Thr	Pro	Ala	Met	Ala	Ala	
			115					120					125				
	Val	Pro	Thr	Gln	Pro	Gln	Ser	Lys	Glu	Gly	Pro	Trp	Ser	Pro	Glu	Ser	
20							135					140					
	Glu	Ser	Pro	Met	Leu	Arg	Ile	Thr	Ala	Pro	Leu	Pro	Pro	Gly	Pro	Ser	
25	145					150					155					160	
	Met	Ala	Val	Pro	Thr	Leu	Gly	Pro	Gly	Glu	Ile	Ala	Ser	Thr	Thr	Pro	
					165					170					175		
30	Pro	Ser	Arg	Ala	Trp	Thr	Pro	Thr	Gln	Glu	Gly	Pro	Gly	Asp	Met	Gly	
				180					185						190		
	Arg	Pro	Trp	Val	Ala	Glu	Val	Val	Ser	Gln	Gly	Ala	Gly	Ile	Gly	Ile	
35				195				200					205				
	Gln	Gly	Thr	Ile	Thr	Ser	Ser	Thr	Ala	Ser	Gly	Asp	Asp	Glu	Glu	Thr	
	210						215					220					
40	Thr	Thr	Thr	Thr	Thr	Ile	Ile	Thr	Thr	Thr	Ile	Thr	Thr	Val	Gln	Thr	
	225					230					235					240	
	Pro	Gly	Pro	Cys	Ser	Trp	Asn	Phe	Ser	Gly	Pro	Glu	Gly	Ser	Leu	Asp	
45					245					250					255		
	Ser	Pro	Thr	Asp	Leu	Ser	Ser	Pro	Thr	Asp	Val	Gly	Leu	Asp	Cys	Phe	
50				260					265					270			
	Phe	Tyr	Ile	Ser	Val	Tyr	Pro	Gly	Tyr	Gly	Val	Glu	Ile	Lys	Val	Gln	
			275					280					285				
55	Asn	Ile	Ser	Leu	Arg	Glu	Gly	Glu	Thr	Val	Thr	Val	Glu	Gly	Leu	Gly	
	290						295					300					

EP 3 338 793 A1

	Gly	Pro	Asp	Pro	Leu	Pro	Leu	Ala	Asn	Gln	Ser	Phe	Leu	Leu	Arg	Gly	
	305					310					315					320	
5	Gln	Val	Ile	Arg	Ser	Pro	Thr	His	Gln	Ala	Ala	Leu	Arg	Phe	Gln	Ser	
					325					330					335		
10	Leu	Pro	Pro	Pro	Ala	Gly	Pro	Gly	Thr	Phe	His	Phe	His	Tyr	Gln	Ala	
				340					345					350			
15	Tyr	Leu	Leu	Ser	Cys	His	Phe	Pro	Arg	Arg	Pro	Ala	Tyr	Gly	Asp	Val	
		355						360					365				
20	Thr	Val	Thr	Ser	Leu	His	Pro	Gly	Gly	Ser	Ala	Arg	Phe	His	Cys	Ala	
		370					375					380					
25	Thr	Gly	Tyr	Gln	Leu	Lys	Gly	Ala	Arg	His	Leu	Thr	Cys	Leu	Asn	Ala	
	385					390					395					400	
30	Thr	Gln	Pro	Phe	Trp	Asp	Ser	Lys	Glu	Pro	Val	Cys	Ile	Ala	Ala	Cys	
				405					410						415		
35	Gly	Gly	Val	Ile	Arg	Asn	Ala	Thr	Thr	Gly	Arg	Ile	Val	Ser	Pro	Gly	
			420						425					430			
40	Phe	Pro	Gly	Asn	Tyr	Ser	Asn	Asn	Leu	Thr	Cys	His	Trp	Leu	Leu	Glu	
		435					440					445					
45	Ala	Pro	Glu	Gly	Gln	Arg	Leu	His	Leu	His	Phe	Glu	Lys	Val	Ser	Leu	
	450						455					460					
50	Ala	Glu	Asp	Asp	Asp	Arg	Leu	Ile	Ile	Arg	Asn	Gly	Asp	Asn	Val	Glu	
	465					470				475						480	
55	Ala	Pro	Pro	Val	Tyr	Asp	Ser	Tyr	Glu	Val	Glu	Tyr	Leu	Pro	Ile	Glu	
				485					490					495			
60	Gly	Leu	Leu	Ser	Ser	Gly	Lys	His	Phe	Phe	Val	Glu	Leu	Ser	Thr	Asp	
			500						505					510			
65	Ser	Ser	Gly	Ala	Ala	Ala	Gly	Met	Ala	Leu	Arg	Tyr	Glu	Ala	Phe	Gln	
		515					520					525					
70	Gln	Gly	His	Cys	Tyr	Glu	Pro	Phe	Val	Lys	Tyr	Gly	Asn	Phe	Ser	Ser	
	530						535					540					
75	Ser	Thr	Pro	Thr	Tyr	Pro	Val	Gly	Thr	Thr	Val	Glu	Phe	Ser	Cys	Asp	

EP 3 338 793 A1

	545		550		555		560
5	Pro Gly Tyr Thr	Leu Glu Gln Gly	Ser Ile Ile Ile	Glu Cys Val Asp			
		565		570		575	
10	Pro His Asp Pro	Gln Trp Asn Glu	Thr Glu Pro Ala	Cys Arg Ala Val			
		580		585		590	
15	Cys Ser Gly Glu	Ile Thr Asp Ser	Ala Gly Val Val	Leu Ser Pro Asn			
		595		600		605	
20	Trp Pro Glu Pro	Tyr Gly Arg Gly	Gln Asp Cys Ile	Trp Gly Val His			
		610		615		620	
25	Val Glu Glu Asp	Lys Arg Ile Met	Leu Asp Ile Arg	Val Leu Arg Ile			
		625		630		635	
30	Gly Pro Gly Asp	Val Leu Thr Phe	Tyr Asp Gly Asp	Asp Leu Thr Ala			
		645		650		655	
35	Arg Val Leu Gly	Gln Tyr Ser Gly	Pro Arg Ser His	Phe Lys Leu Phe			
		660		665		670	
40	Thr Ser Met Ala	Asp Val Thr Ile	Gln Phe Gln Ser	Asp Pro Gly Thr			
		675		680		685	
45	Ser Val Leu Gly	Tyr Gln Gln Gly	Phe Val Ile His	Phe Phe Glu Val			
		690		695		700	
50	Pro Arg Asn Asp	Thr Cys Pro Glu	Leu Pro Glu Ile	Pro Asn Gly Trp			
		705		710		715	
55	Lys Ser Pro Ser	Gln Pro Glu Leu	Val His Gly Thr	Val Val Thr Tyr			
		725		730		735	
60	Gln Cys Tyr Pro	Gly Tyr Gln Val	Val Gly Ser Ser	Val Leu Met Cys			
		740		745		750	
65	Gln Trp Asp Leu	Thr Trp Ser Glu	Asp Leu Pro Ser	Cys Gln Arg Val			
		755		760		765	
70	Thr Ser Cys His	Asp Pro Gly Asp	Val Glu His Ser	Arg Arg Leu Ile			
		770		775		780	
75	Ser Ser Pro Lys	Phe Pro Val Gly	Ala Thr Val Gln	Tyr Ile Cys Asp			
		785		790		795	
80							800

EP 3 338 793 A1

Gln Gly Phe Val Leu Met Gly Ser Ser Ile Leu Thr Cys His Asp Arg
 805 810 815
 5 Gln Ala Gly Ser Pro Lys Trp Ser Asp Arg Ala Pro Lys Cys Leu Leu
 820 825 830
 Glu Gln Leu Lys Pro Cys His Gly Leu Ser Ala Pro Glu Asn Gly Ala
 835 840 845
 10 Arg Ser Pro Glu Lys Gln Leu His Pro Ala Gly Ala Thr Ile His Phe
 850 855 860
 15 Ser Cys Ala Pro Gly Tyr Val Leu Lys Gly Gln Ala Ser Ile Lys Cys
 865 870 875 880
 Val Pro Gly His Pro Ser His Trp Ser Asp Pro Pro Pro Ile Cys Arg
 885 890 895
 Ala Ala Ser Leu Asp Gly Phe Tyr Asn Ser Arg Ser Leu Asp Val Ala
 900 905 910
 25 Lys Ala Pro Ala Ala Ser Ser Thr Leu Asp Ala Ala His Ile Ala Ala
 915 920 925
 Ala Ile Phe Leu Pro Leu Val Ala Met Val Leu Leu Val Gly Gly Val
 930 935 940
 Tyr Phe Tyr Phe Ser Arg Leu Gln Gly Lys Ser Ser Leu Gln Leu Pro
 945 950 955 960
 35 Arg Pro Arg Pro Arg Pro Tyr Asn Arg Ile Thr Ile Glu Ser Ala Phe
 965 970 975
 40 Asp Asn Pro Thr Tyr Glu Thr Gly Glu Thr Arg Glu Tyr Glu Val Ser
 980 985 990
 Ile
 45
 <210> 5
 <211> 2925
 <212> DNA
 50 <213> Homo sapiens
 <220>
 <221> misc_feature
 55 <223> cDNA sequence of hSCRx17 ORF
 <400> 5

EP 3 338 793 A1

	ctgagcctgg	aggccccaac	cgtggggaaa	ggacaagccc	caggcatcga	ggagacagat	60
	ggcgagctga	cagcagcccc	cacacctgag	cagccagaac	gaggcgtcca	ctttgtcaca	120
5	acagccccc	ccttgaagct	gctcaaccac	cacccgctgc	ttgaggaatt	cctacaagag	180
	gggctggaaa	agggagatga	ggagttgagg	ccagcactgc	ccttcagacc	tgacccacct	240
	gcacccttca	ccccaagtcc	ccttccccgc	ctggccaacc	aggacagccg	ccctgtcttt	300
10	accagcccca	ctccagccat	ggctgcggta	cccactcagc	cccagtccaa	ggagggaccc	360
	tggagtccgg	agtcagagtc	ccctatgctt	cgaatcacag	ctcccctacc	tccagggccc	420
	agcatggcag	tgccaccctt	aggcccaggg	gagatagcca	gcactacacc	ccccagcaga	480
15	gcctggacac	caaccacaaga	gggtcctgga	gacatgggaa	ggccgtgggt	tgacagaggtt	540
	gtgtcccagg	gcgcggggat	cgggatccag	gggaccatca	cctcctccac	agcttcagga	600
	gatgatgagg	agaccaccac	taccaccacc	atcatcacca	ccaccatcac	cacagtccag	660
20	acaccaggcc	cttgtagctg	gaattttctca	ggcccagagg	gctctctgga	ctcccctaca	720
	gacctcagct	ccccactga	tgttggcctg	gactgcttct	tctacatctc	tgtctacctt	780
25	ggctatggcg	tggaaatcaa	ggtccagaat	atcagcctcc	gggaagggga	gacagtgact	840
	gtggaaggcc	tggggggggc	cgaccactg	cccctggcca	accagtcttt	cctgctgcgg	900
	ggccaagtca	tccgcagccc	cacccaccaa	gcggccctga	ggttccagag	cctcccggca	960
30	ccggctggcc	ctggcacctt	ccatttccat	taccaagcct	atctcctgag	ctgccacttt	1020
	ccccgtcgtc	cagcttatgg	agatgtgact	gtcaccagcc	tccaccaggg	gggtagtgcc	1080
	cgcttccatt	gtgccactgg	ctaccagctg	aagggcgcca	ggcatctcac	ctgtctcaat	1140
35	gccaccacgc	ccttctggga	ttcaaaggag	cccgtctgca	tcgctgcttg	cggcggagtg	1200
	atccgcaatg	ccaccaccgg	cgcacatcgtc	tctccaggct	tcccggggaa	ctacagcaac	1260
	aacctcacct	gtcactggct	gcttgaggct	cctgagggcc	agcggctaca	cctgcacttt	1320
40	gagaaggttt	ccctggcaga	ggatgatgac	aggctcatca	ttcgcaatgg	ggacaacgtg	1380
	gaggccccc	cagtgtatga	ttcctatgag	gtggaatacc	tgcccattga	gggcctgctc	1440
	agctctggca	aacacttctt	tgttgagctc	agtactgaca	gcagcggggc	agctgcaggg	1500
45	atggccctgc	gctatgaggc	cttccagcag	ggccattgct	atgagccctt	tgtcaaatac	1560
	ggtaacttca	gcagcagcac	accacacct	cctgtgggta	ccactgtgga	gttcagctgc	1620
	gaccctggct	acaccctgga	gcagggctcc	atcatcatcg	agtgtgttga	ccccacgac	1680
50	ccccagtgga	atgagacaga	gccagcctgc	cgagccgtgt	gcagcgggga	gatcacagac	1740
	tcggctggcg	tggtactctc	tcccaactgg	ccagagccct	acggctcgtg	gcaggattgt	1800
55	atctgggggtg	tgcatgtgga	agaggacaag	cgcacatcgc	tggacatccg	agtgtgcgc	1860
	ataggccctg	gtgatgtgct	taccttctat	gatggggatg	acctgacggc	ccgggttctg	1920

EP 3 338 793 A1

ggccagtact cagggccccc tagccacttc aagctcttta cctccatggc tgatgtcacc 1980
 attcagttcc agtcggaccc cgggacctca gtgctgggct accagcaggg cttcgtcatc 2040
 5 cacttctttg aggtgccccg caatgacaca tgtccggagc tgcctgagat ccccaatggc 2100
 tggaagagcc catcgcagcc tgagctagtg cacggcaccg tggtcactta ccagtgttac 2160
 cctggctacc aggtagtggg atccagtgtc ctcatgtgcc agtgggacct aacttggagt 2220
 10 gaggacctgc cctcatgcca gagggtgact tcctgccacg atcctggaga tgtggagcac 2280
 agccgacgcc tcatatccag ccccaagttt cccgtggggg ccaccgtgca atatatctgt 2340
 gaccagggtt ttgtgtgat gggcagctcc atcctcacct gccatgatcg ccaggctggc 2400
 15 agccccaagt ggagtgaccg ggcccctaaa tgtctcctgg aacagctcaa gccatgccat 2460
 ggtctcagtg cccctgagaa tgggtcccga agtcctgaga agcagctaca cccagcaggg 2520
 gccaccatcc acttctctgtg tgcccctggc tatgtgtctga agggccaggc cagcatcaag 2580
 20 tgtgtgcctg ggcacccctc gcattggagt gacccccac ccatctgtag ggctgcctct 2640
 ctggatgggt tctacaacag tcgcagcctg gatgttgcca aggcacctgc tgctccagc 2700
 25 accctggatg ctgccacat tgcagctgcc atcttcttgc cactggtggc gatggtgttg 2760
 ttggtaggag gtgtatactt ctacttctcc aggtccagg gaaaaagctc cctgcagctg 2820
 ccccgcccc gccccgccc ctacaaccgc attaccatag agtcagcgtt tgacaatcca 2880
 30 acttacgaga ctggatctct ttcctttgca ggagacgaga gaata 2925

<210> 6
 <211> 975
 35 <212> PRT
 <213> Homo sapiens

<220>
 40 <221> MISC_FEATURE
 <223> hSCR α 17 protein

<400> 6
 45 Leu Ser Leu Glu Ala Pro Thr Val Gly Lys Gly Gln Ala Pro Gly Ile
 1 5 10 15
 Glu Glu Thr Asp Gly Glu Leu Thr Ala Ala Pro Thr Pro Glu Gln Pro
 20 25 30
 50 Glu Arg Gly Val His Phe Val Thr Thr Ala Pro Thr Leu Lys Leu Leu
 35 40 45
 55 Asn His His Pro Leu Leu Glu Glu Phe Leu Gln Glu Gly Leu Glu Lys
 50 55 60

EP 3 338 793 A1

	Gly	Asp	Glu	Glu	Leu	Arg	Pro	Ala	Leu	Pro	Phe	Gln	Pro	Asp	Pro	Pro	65	70	75	80
5	Ala	Pro	Phe	Thr	Pro	Ser	Pro	Leu	Pro	Arg	Leu	Ala	Asn	Gln	Asp	Ser	85	90	95	
10	Arg	Pro	Val	Phe	Thr	Ser	Pro	Thr	Pro	Ala	Met	Ala	Ala	Val	Pro	Thr	100	105	110	
15	Gln	Pro	Gln	Ser	Lys	Glu	Gly	Pro	Trp	Ser	Pro	Glu	Ser	Glu	Ser	Pro	115	120	125	
20	Met	Leu	Arg	Ile	Thr	Ala	Pro	Leu	Pro	Pro	Gly	Pro	Ser	Met	Ala	Val	130	135	140	
25	Pro	Thr	Leu	Gly	Pro	Gly	Glu	Ile	Ala	Ser	Thr	Thr	Pro	Pro	Ser	Arg	145	150	155	160
30	Ala	Trp	Thr	Pro	Thr	Gln	Glu	Gly	Pro	Gly	Asp	Met	Gly	Arg	Pro	Trp	165	170	175	
35	Val	Ala	Glu	Val	Val	Ser	Gln	Gly	Ala	Gly	Ile	Gly	Ile	Gln	Gly	Thr	180	185	190	
40	Ile	Thr	Ser	Ser	Thr	Ala	Ser	Gly	Asp	Asp	Glu	Glu	Thr	Thr	Thr	Thr	195	200	205	
45	Thr	Thr	Ile	Ile	Thr	Thr	Thr	Ile	Thr	Thr	Val	Gln	Thr	Pro	Gly	Pro	210	215	220	
50	Cys	Ser	Trp	Asn	Phe	Ser	Gly	Pro	Glu	Gly	Ser	Leu	Asp	Ser	Pro	Thr	225	230	235	240
55	Asp	Leu	Ser	Ser	Pro	Thr	Asp	Val	Gly	Leu	Asp	Cys	Phe	Phe	Tyr	Ile	245	250	255	
60	Ser	Val	Tyr	Pro	Gly	Tyr	Gly	Val	Glu	Ile	Lys	Val	Gln	Asn	Ile	Ser	260	265	270	
65	Leu	Arg	Glu	Gly	Glu	Thr	Val	Thr	Val	Glu	Gly	Leu	Gly	Gly	Pro	Asp	275	280	285	
70	Pro	Leu	Pro	Leu	Ala	Asn	Gln	Ser	Phe	Leu	Leu	Arg	Gly	Gln	Val	Ile	290	295	300	
75	Arg	Ser	Pro	Thr	His	Gln	Ala	Ala	Leu	Arg	Phe	Gln	Ser	Leu	Pro	Pro	305	310	315	320

EP 3 338 793 A1

	Pro	Ala	Gly	Pro	Gly	Thr	Phe	His	Phe	His	Tyr	Gln	Ala	Tyr	Leu	Leu	
					325					330					335		
5	Ser	Cys	His	Phe	Pro	Arg	Arg	Pro	Ala	Tyr	Gly	Asp	Val	Thr	Val	Thr	
				340					345					350			
10	Ser	Leu	His	Pro	Gly	Gly	Ser	Ala	Arg	Phe	His	Cys	Ala	Thr	Gly	Tyr	
			355					360					365				
15	Gln	Leu	Lys	Gly	Ala	Arg	His	Leu	Thr	Cys	Leu	Asn	Ala	Thr	Gln	Pro	
		370					375					380					
20	Phe	Trp	Asp	Ser	Lys	Glu	Pro	Val	Cys	Ile	Ala	Ala	Cys	Gly	Gly	Val	
	385					390					395					400	
25	Ile	Arg	Asn	Ala	Thr	Thr	Gly	Arg	Ile	Val	Ser	Pro	Gly	Phe	Pro	Gly	
					405					410					415		
30	Asn	Tyr	Ser	Asn	Asn	Leu	Thr	Cys	His	Trp	Leu	Leu	Glu	Ala	Pro	Glu	
				420					425					430			
35	Gly	Gln	Arg	Leu	His	Leu	His	Phe	Glu	Lys	Val	Ser	Leu	Ala	Glu	Asp	
			435					440					445				
40	Asp	Asp	Arg	Leu	Ile	Ile	Arg	Asn	Gly	Asp	Asn	Val	Glu	Ala	Pro	Pro	
		450					455					460					
45	Val	Tyr	Asp	Ser	Tyr	Glu	Val	Glu	Tyr	Leu	Pro	Ile	Glu	Gly	Leu	Leu	
	465					470					475					480	
50	Ser	Ser	Gly	Lys	His	Phe	Phe	Val	Glu	Leu	Ser	Thr	Asp	Ser	Ser	Gly	
					485					490					495		
55	Ala	Ala	Ala	Gly	Met	Ala	Leu	Arg	Tyr	Glu	Ala	Phe	Gln	Gln	Gly	His	
				500					505					510			
60	Cys	Tyr	Glu	Pro	Phe	Val	Lys	Tyr	Gly	Asn	Phe	Ser	Ser	Ser	Thr	Pro	
			515					520					525				
65	Thr	Tyr	Pro	Val	Gly	Thr	Thr	Val	Glu	Phe	Ser	Cys	Asp	Pro	Gly	Tyr	
		530					535					540					
70	Thr	Leu	Glu	Gln	Gly	Ser	Ile	Ile	Ile	Glu	Cys	Val	Asp	Pro	His	Asp	
	545					550					555					560	
75	Pro	Gln	Trp	Asn	Glu	Thr	Glu	Pro	Ala	Cys	Arg	Ala	Val	Cys	Ser	Gly	

EP 3 338 793 A1

[illegible]

EP 3 338 793 A1

Lys Pro Cys His Gly Leu Ser Ala Pro Glu Asn Gly Ala Arg Ser Pro
 820 825 830

5 Glu Lys Gln Leu His Pro Ala Gly Ala Thr Ile His Phe Ser Cys Ala
 835 840 845

10 Pro Gly Tyr Val Leu Lys Gly Gln Ala Ser Ile Lys Cys Val Pro Gly
 850 855 860

His Pro Ser His Trp Ser Asp Pro Pro Pro Ile Cys Arg Ala Ala Ser
 865 870 875 880

15 Leu Asp Gly Phe Tyr Asn Ser Arg Ser Leu Asp Val Ala Lys Ala Pro
 885 890 895

20 Ala Ala Ser Ser Thr Leu Asp Ala Ala His Ile Ala Ala Ile Phe
 900 905 910

25 Leu Pro Leu Val Ala Met Val Leu Leu Val Gly Gly Val Tyr Phe Tyr
 915 920 925

Phe Ser Arg Leu Gln Gly Lys Ser Ser Leu Gln Leu Pro Arg Pro Arg
 930 935 940

30 Pro Arg Pro Tyr Asn Arg Ile Thr Ile Glu Ser Ala Phe Asp Asn Pro
 945 950 955 960

35 Thr Tyr Glu Thr Gly Ser Leu Ser Phe Ala Gly Asp Glu Arg Ile
 965 970 975

<210> 7
 <211> 994
 <212> PRT
 40 <213> Homo sapiens

<220>
 <221> MISC_FEATURE
 45 <223> hSEZ6 BC146292 protein

<400> 7

50 Met Arg Pro Val Ala Leu Leu Leu Leu Pro Ser Leu Leu Ala Leu Leu
 1 5 10 15

Ala His Gly Leu Ser Leu Glu Ala Pro Thr Val Gly Lys Gly Gln Ala
 20 25 30

55 Pro Gly Ile Glu Glu Thr Asp Gly Glu Leu Thr Ala Ala Pro Thr Pro
 35 40 45

EP 3 338 793 A1

	Glu	Gln	Pro	Glu	Arg	Gly	Val	His	Phe	Val	Thr	Thr	Ala	Pro	Thr	Leu	
	50						55					60					
5	Lys	Leu	Leu	Asn	His	His	Pro	Leu	Leu	Glu	Glu	Phe	Leu	Gln	Glu	Gly	
	65					70					75					80	
	Leu	Glu	Lys	Gly	Asp	Glu	Glu	Leu	Arg	Pro	Ala	Leu	Pro	Phe	Gln	Pro	
10					85					90					95		
	Asp	Pro	Pro	Ala	Pro	Phe	Thr	Pro	Ser	Pro	Leu	Pro	Arg	Leu	Ala	Asn	
				100					105					110			
15	Gln	Asp	Ser	Arg	Pro	Val	Phe	Thr	Ser	Pro	Thr	Pro	Ala	Met	Ala	Ala	
			115					120					125				
	Val	Pro	Thr	Gln	Pro	Gln	Ser	Lys	Glu	Gly	Pro	Trp	Ser	Pro	Glu	Ser	
20							135					140					
	Glu	Ser	Pro	Met	Leu	Arg	Ile	Thr	Ala	Pro	Leu	Pro	Pro	Gly	Pro	Ser	
25	145					150					155					160	
	Met	Ala	Val	Pro	Thr	Leu	Gly	Pro	Gly	Glu	Ile	Ala	Ser	Thr	Thr	Pro	
					165					170					175		
30	Pro	Ser	Arg	Ala	Trp	Thr	Pro	Thr	Gln	Glu	Gly	Pro	Gly	Asp	Met	Gly	
				180					185					190			
	Arg	Pro	Trp	Val	Ala	Glu	Val	Val	Ser	Gln	Gly	Ala	Gly	Ile	Gly	Ile	
35				195				200					205				
	Gln	Gly	Thr	Ile	Thr	Ser	Ser	Thr	Ala	Ser	Gly	Asp	Asp	Glu	Glu	Thr	
	210					215						220					
40	Thr	Thr	Thr	Thr	Thr	Ile	Ile	Thr	Thr	Thr	Ile	Thr	Thr	Val	Gln	Thr	
	225					230					235					240	
	Pro	Gly	Pro	Cys	Ser	Trp	Asn	Phe	Ser	Gly	Pro	Glu	Gly	Ser	Leu	Asp	
45					245					250					255		
	Ser	Pro	Thr	Asp	Leu	Ser	Ser	Pro	Thr	Asp	Val	Gly	Leu	Asp	Cys	Phe	
50				260					265					270			
	Phe	Tyr	Ile	Ser	Val	Tyr	Pro	Gly	Tyr	Gly	Val	Glu	Ile	Lys	Val	Gln	
			275					280					285				
55	Asn	Ile	Ser	Leu	Arg	Glu	Gly	Glu	Thr	Val	Thr	Val	Glu	Gly	Leu	Gly	
	290						295					300					

EP 3 338 793 A1

	Gly	Pro	Asp	Pro	Leu	Pro	Leu	Ala	Asn	Gln	Ser	Phe	Leu	Leu	Arg	Gly	
	305					310					315					320	
5	Gln	Val	Ile	Arg	Ser	Pro	Thr	His	Gln	Ala	Ala	Leu	Arg	Phe	Gln	Ser	
					325					330					335		
10	Leu	Pro	Pro	Pro	Ala	Gly	Pro	Gly	Thr	Phe	His	Phe	His	Tyr	Gln	Ala	
				340					345					350			
15	Tyr	Leu	Leu	Ser	Cys	His	Phe	Pro	Arg	Arg	Pro	Ala	Tyr	Gly	Asp	Val	
		355						360					365				
20	Thr	Val	Thr	Ser	Leu	His	Pro	Gly	Gly	Ser	Ala	Arg	Phe	His	Cys	Ala	
		370					375					380					
25	Thr	Gln	Pro	Phe	Trp	Asp	Ser	Lys	Glu	Pro	Val	Cys	Ile	Gly	Glu	Cys	
				405					410						415		
30	Pro	Gly	Val	Ile	Arg	Asn	Ala	Thr	Thr	Gly	Arg	Ile	Val	Ser	Pro	Gly	
			420						425					430			
35	Phe	Pro	Gly	Asn	Tyr	Ser	Asn	Asn	Leu	Thr	Cys	His	Trp	Leu	Leu	Glu	
		435						440					445				
40	Ala	Pro	Glu	Gly	Gln	Arg	Leu	His	Leu	His	Phe	Glu	Lys	Val	Ser	Leu	
		450					455					460					
45	Ala	Glu	Asp	Asp	Asp	Arg	Leu	Ile	Ile	Arg	Asn	Gly	Asp	Asn	Val	Glu	
	465					470					475					480	
50	Ala	Pro	Pro	Val	Tyr	Asp	Ser	Tyr	Glu	Val	Glu	Tyr	Leu	Pro	Ile	Glu	
				485					490						495		
55	Gly	Leu	Leu	Ser	Ser	Gly	Lys	His	Phe	Phe	Val	Glu	Leu	Ser	Thr	Asp	
				500					505					510			
60	Ser	Ser	Gly	Ala	Ala	Ala	Gly	Met	Ala	Leu	Arg	Tyr	Glu	Ala	Phe	Gln	
			515					520					525				
65	Gln	Gly	His	Cys	Tyr	Glu	Pro	Phe	Val	Lys	Tyr	Gly	Asn	Phe	Ser	Ser	
		530					535					540					
70	Ser	Thr	Pro	Thr	Tyr	Pro	Val	Gly	Thr	Thr	Val	Glu	Phe	Ser	Cys	Asp	

EP 3 338 793 A1

	545		550		555		560
5	Pro Gly Tyr Thr	Leu Glu Gln Gly	Ser Ile Ile Ile	Glu Cys Val Asp			
		565		570		575	
10	Pro His Asp Pro	Gln Trp Asn Glu	Thr Glu Pro Ala	Cys Arg Ala Val			
		580		585		590	
15	Cys Ser Gly Glu	Ile Thr Asp Ser	Ala Gly Val Val	Leu Ser Pro Asn			
		595		600		605	
20	Trp Pro Glu Pro	Tyr Gly Arg Gly	Gln Asp Cys Ile	Trp Gly Val His			
		610		615		620	
25	Val Glu Glu Asp	Lys Arg Ile Met	Leu Asp Ile Arg	Val Leu Arg Ile			
		625		630		635	
30	Gly Pro Gly Asp	Val Leu Thr Phe	Tyr Asp Gly Asp	Asp Leu Thr Ala			
		645		650		655	
35	Arg Val Leu Gly	Gln Tyr Ser Gly	Pro Arg Ser His	Phe Lys Leu Phe			
		660		665		670	
40	Thr Ser Met Ala	Asp Val Thr Ile	Gln Phe Gln Ser	Asp Pro Gly Thr			
		675		680		685	
45	Ser Val Leu Gly	Tyr Gln Gln Gly	Phe Val Ile His	Phe Phe Glu Val			
		690		695		700	
50	Pro Arg Asn Asp	Thr Cys Pro Glu	Leu Pro Glu Ile	Pro Asn Gly Trp			
		705		710		715	
55	Lys Ser Pro Ser	Gln Pro Glu Leu	Val His Gly Thr	Val Val Thr Tyr			
		725		730		735	
60	Gln Cys Tyr Pro	Gly Tyr Gln Val	Val Gly Ser Ser	Val Leu Met Cys			
		740		745		750	
65	Gln Trp Asp Leu	Thr Trp Ser Glu	Asp Leu Pro Ser	Cys Gln Arg Val			
		755		760		765	
70	Thr Ser Cys His	Asp Pro Gly Asp	Val Glu His Ser	Arg Arg Leu Ile			
		770		775		780	
75	Ser Ser Pro Lys	Phe Pro Val Gly	Ala Thr Val Gln	Tyr Ile Cys Asp			
		785		790		795	
80							

EP 3 338 793 A1

Gln Gly Phe Val Leu Met Gly Ser Ser Ile Leu Thr Cys His Asp Arg
 805 810 815

5 Gln Ala Gly Ser Pro Lys Trp Ser Asp Arg Ala Pro Lys Cys Leu Leu
 820 825 830

10 Glu Gln Leu Lys Pro Cys His Gly Leu Ser Ala Pro Glu Asn Gly Ala
 835 840 845

Arg Ser Pro Glu Lys Gln Leu His Pro Ala Gly Ala Thr Ile His Phe
 850 855 860

15 Ser Cys Ala Pro Gly Tyr Val Leu Lys Gly Gln Ala Ser Ile Lys Cys
 865 870 875 880

20 Val Pro Gly His Pro Ser His Trp Ser Asp Pro Pro Pro Ile Cys Arg
 885 890 895

25 Ala Ala Ser Leu Asp Gly Phe Tyr Asn Ser Arg Ser Leu Asp Val Ala
 900 905 910

Lys Ala Pro Ala Ala Ser Ser Thr Leu Asp Ala Ala His Ile Ala Ala
 915 920 925

30 Ala Ile Phe Leu Pro Leu Val Ala Met Val Leu Leu Val Gly Gly Val
 930 935 940

35 Tyr Phe Tyr Phe Ser Arg Leu Gln Gly Lys Ser Ser Leu Gln Leu Pro
 945 950 955 960

Arg Pro Arg Pro Arg Pro Tyr Asn Arg Ile Thr Ile Glu Ser Ala Phe
 965 970 975

40 Asp Asn Pro Thr Tyr Glu Thr Gly Ser Leu Ser Phe Ala Gly Asp Glu
 980 985 990

45 Arg Ile

50 <210> 8
 <211> 3483
 <212> DNA
 <213> Homo sapiens

55 <220>
 <221> misc_feature
 <223> hSCRx17-Fc ORF

<400> 8

EP 3 338 793 A1

	atggagacag acacactcct gctatgggta ctgctgctct gggttcccgg gtccactggt	60
	gacggcgcg cctggatccct gagcctggag gcccacaccg tggggaaagg acaagcccca	120
5	ggcatcgagg agacagatgg cgagctgaca gcagcccccac cacctgagca gccagaacga	180
	ggcgtccact ttgtcacaac agccccccacc ttgaagctgc tcaaccacca cccgctgctt	240
	gaggaattcc tacaagaggg gctggaaaag ggagatgagg agttgaggcc agcactgccc	300
10	ttccagcctg acccacctgc acccttcacc ccaagtcccc ttccccgcct ggccaaccag	360
	gacagccgcc ctgtctttac cagccccact ccagccatgg ctgcggtacc cactcagccc	420
	cagtccaagg agggaccctg gagtccggag tcagagtccc ctatgcttcg aatcacagct	480
15	cccctacctc caggggccag catggcagtg cccaccctag gcccagggga gatagccagc	540
	actacacccc ccagcagagc ctggacacca acccaagagg gtcctggaga catgggaagg	600
	ccgtgggttg cagaggttgt gtcccagggc gcggggatcg ggatccaggg gaccatcacc	660
20	tcctccacag cttcaggaga tgatgaggag accaccacta ccaccaccat catcaccacc	720
	accatcacca cagtccagac accaggccct tgtagctgga atttctcagg cccagagggc	780
25	tctctggact cccctacaga cctcagctcc cccactgatg ttggcctgga ctgcttcttc	840
	tacatctctg tctaccctgg ctatggcgtg gaaatcaagg tccagaatat cagcctccgg	900
	gaaggggaga cagtgactgt ggaaggcctg ggggggcccg acccaactgcc cctggccaac	960
30	cagtctttcc tgctgcgggg ccaagtcac cgcagcccca cccaccaagc ggccctgagg	1020
	ttccagagcc tcccgccacc ggctggccct ggcaaccttc atttccatta ccaagcctat	1080
	ctcctgagct gccactttcc cgtcgtcca gcttatggag atgtgactgt caccagcctc	1140
35	caccaggggg gtagtgccc cttccattgt gccactggct accagctgaa gggcgccagg	1200
	catctcacct gtctcaatgc caccagccc ttctgggatt caaaggagcc cgtctgcac	1260
	gctgcttgcg gcggagtgat ccgcaatgcc accaccggcc gcatcgtctc tccaggcttc	1320
40	ccgggcaact acagcaacaa cctcacctgt cactggctgc ttgaggctcc tgagggccag	1380
	cggctacacc tgcactttga gaagggttcc ctggcagagg atgatgacag gctcatcatt	1440
	cgcaatgggg acaacgtgga ggccccacca gtgtatgatt cctatgaggt ggaataacctg	1500
45	cccattgagg gcctgctcag ctctggcaaa cacttctttg ttgagctcag tactgacagc	1560
	agcggggcag ctgcaggcat ggccctgcgc tatgaggcct tccagcaggg ccattgctat	1620
	gagccctttg tcaaatacgg taacttcagc agcagcacac ccacctaccc tgtgggtacc	1680
50	actgtggagt tcagctgcga ccctggctac accctggagc agggctccat catcatcgag	1740
	tgtgttgacc cccacgaccc ccagtggaat gagacagagc cagcctgccg agccgtgtgc	1800
55	agcggggaga tcacagactc ggctggcgtg gtactctctc ccaactggcc agagccctac	1860
	ggctcgtgggc aggattgtat ctgggggtgt catgtggaag aggacaagcg catcatgctg	1920

EP 3 338 793 A1

gacatccgag tgctgcgcat aggccctggt gatgtgctta ccttctatga tggggatgac 1980

ctgacggccc gggttctggg ccagtactca gggccccgta gccacttcaa gctctttacc 2040

5 tccatggctg atgtcaccat tcagttccag tcggaccccg ggacctcagt gctgggctac 2100

cagcagggct tcgtcatcca cttctttgag gtgccccgca atgacacatg tccggagctg 2160

cctgagatcc ccaatggctg gaagagccca tcgcagcctg agctagtga cggcaccgtg 2220

10 gtcacttacc agtgctaccc tggctaccag gtagtgggat ccagtgtcct catgtgccag 2280

tgggacctaa cttggagtga ggacctgccc tcatgccaga gggtgacttc ctgccacgat 2340

cctggagatg tggagcacag ccgacgcctc atatccagcc ccaagtttcc cgtggggggcc 2400

15 accgtgcaat atatctgtga ccagggtttt gtgctgatgg gcagctccat cctcacctgc 2460

catgatcgcc aggtcggcag cccaagtgg agtgaccggg cccctaaatg tctcctggaa 2520

20 cagctcaagc catgccatgg tctcagtgcc cctgagaatg gtgcccgaag tcctgagaag 2580

cagctacacc cagcaggggc caccatccac ttctcgtgtg cccctggcta tgtgctgaag 2640

ggccaggcca gcatcaagtg tgtgcctggg caccctcgc attggagtga cccccaccc 2700

25 atctgtaggg ctgcctctct ggatgggttc tacaacagtc gcagcctgga tgttgccaag 2760

gcacctgctg cctccagcac cctggatgct gccacctgg ccggccacag atctgtcgag 2820

tgcccaccgt gcccagcacc acctgtggca ggacctcag tcttcctctt cccccaaaa 2880

30 cccaaggaca ccctcatgat ctcccgacc cctgaggtca cgtgcgtggt ggtggacgtg 2940

agccacgaag accccgaggt ccagttcaac tggtagctgg acggcgtgga ggtgcataat 3000

gccaagacaa agccacggga ggagcagttc aacagcacgt tccgtgtggt cagcgtcctc 3060

35 accgttgtgc accaggactg gctgaacggc aaggagtaca agtgcaaggt ctccaacaaa 3120

ggcctcccag ccccatcga gaaaaccatc tccaaaacca aagggcagcc ccgagaacca 3180

40 caggtgtaca ccctgcccc atccaggag gagatgacca agaaccaggt cagcctgacc 3240

tgcttggtca aaggcttcta cccagcgac atcgccgtgg agtgggagag caatgggcag 3300

ccggagaaca actacaagac cagcctccc atgctggact ccgacggctc cttcttcctc 3360

45 tacagcaagc tcaccgtgga caagagcagg tggcagcagg ggaacgtctt ctcatgctcc 3420

gtgatgcatg aggtctctga caaccactac acgcagaaga gcctctccct gtctccgggt 3480

tga 3483

50

<210> 9

<211> 1160

<212> PRT

<213> Homo sapiens

55

<220>

EP 3 338 793 A1

<221> MISC_FEATURE
<223> hSCRx17-Fc protein

<400> 9

5

Met Glu Thr Asp Thr Leu Leu Leu Trp Val Leu Leu Leu Trp Val Pro
1 5 10 15

10

Gly Ser Thr Gly Asp Gly Ala Pro Gly Ser Leu Ser Leu Glu Ala Pro
20 25 30

15

Thr Val Gly Lys Gly Gln Ala Pro Gly Ile Glu Glu Thr Asp Gly Glu
35 40 45

Leu Thr Ala Ala Pro Thr Pro Glu Gln Pro Glu Arg Gly Val His Phe
50 55 60

20

Val Thr Thr Ala Pro Thr Leu Lys Leu Leu Asn His His Pro Leu Leu
65 70 75 80

25

Glu Glu Phe Leu Gln Glu Gly Leu Glu Lys Gly Asp Glu Glu Leu Arg
85 90 95

Pro Ala Leu Pro Phe Gln Pro Asp Pro Pro Ala Pro Phe Thr Pro Ser
100 105 110

30

Pro Leu Pro Arg Leu Ala Asn Gln Asp Ser Arg Pro Val Phe Thr Ser
115 120 125

35

Pro Thr Pro Ala Met Ala Ala Val Pro Thr Gln Pro Gln Ser Lys Glu
130 135 140

40

Gly Pro Trp Ser Pro Glu Ser Glu Ser Pro Met Leu Arg Ile Thr Ala
145 150 155 160

Pro Leu Pro Pro Gly Pro Ser Met Ala Val Pro Thr Leu Gly Pro Gly
165 170 175

45

Glu Ile Ala Ser Thr Thr Pro Pro Ser Arg Ala Trp Thr Pro Thr Gln
180 185 190

50

Glu Gly Pro Gly Asp Met Gly Arg Pro Trp Val Ala Glu Val Val Ser
195 200 205

Gln Gly Ala Gly Ile Gly Ile Gln Gly Thr Ile Thr Ser Ser Thr Ala
210 215 220

55

Ser Gly Asp Asp Glu Glu Thr Thr Thr Thr Thr Thr Thr Thr Thr Thr Thr
225 230 235 240

EP 3 338 793 A1

	Thr	Ile	Thr	Thr	Val	Gln	Thr	Pro	Gly	Pro	Cys	Ser	Trp	Asn	Phe	Ser	
					245					250					255		
5	Gly	Pro	Glu	Gly	Ser	Leu	Asp	Ser	Pro	Thr	Asp	Leu	Ser	Ser	Pro	Thr	
				260					265					270			
10	Asp	Val	Gly	Leu	Asp	Cys	Phe	Phe	Tyr	Ile	Ser	Val	Tyr	Pro	Gly	Tyr	
			275					280					285				
15	Gly	Val	Glu	Ile	Lys	Val	Gln	Asn	Ile	Ser	Leu	Arg	Glu	Gly	Glu	Thr	
		290					295					300					
20	Val	Thr	Val	Glu	Gly	Leu	Gly	Gly	Pro	Asp	Pro	Leu	Pro	Leu	Ala	Asn	
	305					310					315					320	
25	Gln	Ser	Phe	Leu	Leu	Arg	Gly	Gln	Val	Ile	Arg	Ser	Pro	Thr	His	Gln	
					325					330					335		
30	Ala	Ala	Leu	Arg	Phe	Gln	Ser	Leu	Pro	Pro	Pro	Ala	Gly	Pro	Gly	Thr	
				340					345					350			
35	Phe	His	Phe	His	Tyr	Gln	Ala	Tyr	Leu	Leu	Ser	Cys	His	Phe	Pro	Arg	
			355					360					365				
40	Arg	Pro	Ala	Tyr	Gly	Asp	Val	Thr	Val	Thr	Ser	Leu	His	Pro	Gly	Gly	
		370					375					380					
45	Ser	Ala	Arg	Phe	His	Cys	Ala	Thr	Gly	Tyr	Gln	Leu	Lys	Gly	Ala	Arg	
	385					390					395					400	
50	His	Leu	Thr	Cys	Leu	Asn	Ala	Thr	Gln	Pro	Phe	Trp	Asp	Ser	Lys	Glu	
					405					410					415		
55	Pro	Val	Cys	Ile	Ala	Ala	Cys	Gly	Gly	Val	Ile	Arg	Asn	Ala	Thr	Thr	
				420					425					430			
60	Gly	Arg	Ile	Val	Ser	Pro	Gly	Phe	Pro	Gly	Asn	Tyr	Ser	Asn	Asn	Leu	
			435					440					445				
65	Thr	Cys	His	Trp	Leu	Leu	Glu	Ala	Pro	Glu	Gly	Gln	Arg	Leu	His	Leu	
		450					455					460					
70	His	Phe	Glu	Lys	Val	Ser	Leu	Ala	Glu	Asp	Asp	Asp	Arg	Leu	Ile	Ile	
	465					470					475					480	
75	Arg	Asn	Gly	Asp	Asn	Val	Glu	Ala	Pro	Pro	Val	Tyr	Asp	Ser	Tyr	Glu	

EP 3 338 793 A1

	485								490				495			
5	Val	Glu	Tyr	Leu	Pro	Ile	Glu	Gly	Leu	Leu	Ser	Ser	Gly	Lys	His	Phe
				500					505					510		
	Phe	Val	Glu	Leu	Ser	Thr	Asp	Ser	Ser	Gly	Ala	Ala	Ala	Gly	Met	Ala
			515					520					525			
10	Leu	Arg	Tyr	Glu	Ala	Phe	Gln	Gln	Gly	His	Cys	Tyr	Glu	Pro	Phe	Val
		530					535					540				
15	Lys	Tyr	Gly	Asn	Phe	Ser	Ser	Ser	Thr	Pro	Thr	Tyr	Pro	Val	Gly	Thr
	545					550					555					560
	Thr	Val	Glu	Phe	Ser	Cys	Asp	Pro	Gly	Tyr	Thr	Leu	Glu	Gln	Gly	Ser
20					565					570					575	
	Ile	Ile	Ile	Glu	Cys	Val	Asp	Pro	His	Asp	Pro	Gln	Trp	Asn	Glu	Thr
				580					585					590		
25	Glu	Pro	Ala	Cys	Arg	Ala	Val	Cys	Ser	Gly	Glu	Ile	Thr	Asp	Ser	Ala
			595					600					605			
30	Gly	Val	Val	Leu	Ser	Pro	Asn	Trp	Pro	Glu	Pro	Tyr	Gly	Arg	Gly	Gln
		610					615					620				
	Asp	Cys	Ile	Trp	Gly	Val	His	Val	Glu	Glu	Asp	Lys	Arg	Ile	Met	Leu
35	625					630					635					640
	Asp	Ile	Arg	Val	Leu	Arg	Ile	Gly	Pro	Gly	Asp	Val	Leu	Thr	Phe	Tyr
				645						650					655	
40	Asp	Gly	Asp	Asp	Leu	Thr	Ala	Arg	Val	Leu	Gly	Gln	Tyr	Ser	Gly	Pro
				660					665					670		
45	Arg	Ser	His	Phe	Lys	Leu	Phe	Thr	Ser	Met	Ala	Asp	Val	Thr	Ile	Gln
			675					680					685			
	Phe	Gln	Ser	Asp	Pro	Gly	Thr	Ser	Val	Leu	Gly	Tyr	Gln	Gln	Gly	Phe
50		690					695					700				
	Val	Ile	His	Phe	Phe	Glu	Val	Pro	Arg	Asn	Asp	Thr	Cys	Pro	Glu	Leu
	705					710					715					720
55	Pro	Glu	Ile	Pro	Asn	Gly	Trp	Lys	Ser	Pro	Ser	Gln	Pro	Glu	Leu	Val
					725					730					735	

EP 3 338 793 A1

	His	Gly	Thr	Val	Val	Thr	Tyr	Gln	Cys	Tyr	Pro	Gly	Tyr	Gln	Val	Val	
				740					745					750			
5	Gly	Ser	Ser	Val	Leu	Met	Cys	Gln	Trp	Asp	Leu	Thr	Trp	Ser	Glu	Asp	
				755				760					765				
10	Leu	Pro	Ser	Cys	Gln	Arg	Val	Thr	Ser	Cys	His	Asp	Pro	Gly	Asp	Val	
		770					775					780					
15	Glu	His	Ser	Arg	Arg	Leu	Ile	Ser	Ser	Pro	Lys	Phe	Pro	Val	Gly	Ala	
	785					790					795					800	
20	Thr	Val	Gln	Tyr	Ile	Cys	Asp	Gln	Gly	Phe	Val	Leu	Met	Gly	Ser	Ser	
					805					810					815		
25	Ile	Leu	Thr	Cys	His	Asp	Arg	Gln	Ala	Gly	Ser	Pro	Lys	Trp	Ser	Asp	
				820					825					830			
30	Arg	Ala	Pro	Lys	Cys	Leu	Leu	Glu	Gln	Leu	Lys	Pro	Cys	His	Gly	Leu	
			835					840					845				
35	Ser	Ala	Pro	Glu	Asn	Gly	Ala	Arg	Ser	Pro	Glu	Lys	Gln	Leu	His	Pro	
		850					855					860					
40	Ala	Gly	Ala	Thr	Ile	His	Phe	Ser	Cys	Ala	Pro	Gly	Tyr	Val	Leu	Lys	
	865					870					875					880	
45	Gly	Gln	Ala	Ser	Ile	Lys	Cys	Val	Pro	Gly	His	Pro	Ser	His	Trp	Ser	
					885					890					895		
50	Asp	Pro	Pro	Pro	Ile	Cys	Arg	Ala	Ala	Ser	Leu	Asp	Gly	Phe	Tyr	Asn	
				900					905					910			
55	Ser	Arg	Ser	Leu	Asp	Val	Ala	Lys	Ala	Pro	Ala	Ala	Ser	Ser	Thr	Leu	
			915					920					925				
60	Asp	Ala	Ala	His	Leu	Ala	Gly	His	Arg	Ser	Val	Glu	Cys	Pro	Pro	Cys	
		930					935					940					
65	Pro	Ala	Pro	Pro	Val	Ala	Gly	Pro	Ser	Val	Phe	Leu	Phe	Pro	Pro	Lys	
	945					950					955					960	
70	Pro	Lys	Asp	Thr	Leu	Met	Ile	Ser	Arg	Thr	Pro	Glu	Val	Thr	Cys	Val	
					965					970					975		
75	Val	Val	Asp	Val	Ser	His	Glu	Asp	Pro	Glu	Val	Gln	Phe	Asn	Trp	Tyr	
				980					985					990			

EP 3 338 793 A1

Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu
995 1000 1005

5 Gln Phe Asn Ser Thr Phe Arg Val Val Ser Val Leu Thr Val Val
1010 1015 1020

10 His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser
1025 1030 1035

Asn Lys Gly Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Thr
1040 1045 1050

15 Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser
1055 1060 1065

20 Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val
1070 1075 1080

Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn
1085 1090 1095

25 Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Met Leu Asp
1100 1105 1110

30 Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys
1115 1120 1125

35 Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His
1130 1135 1140

Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser
1145 1150 1155

40 Pro Gly
1160

45 <210> 10
<211> 2919
<212> DNA
<213> Mus musculus

50 <220>
<221> misc_feature
<223> cDNA sequence encoding mature murine SEZ6

<400> 10
ctctcctcag aggtctccgat cacgggggaa ggatcatgcc cgggcatcag ggagacggat 60
ggggagctga ccgcagcccc tacacctgag cagtcagacc gaggcgtcca cttcgtcacc 120

EP 3 338 793 A1

	acagccccta ccctcaagct gctcaaccac caccacttc tggaagaatt tcttcaagag	180
	gggctagaaa gagaggaagc gccgcagcct gcaactgccct tccagccgga ctacactaca	240
5	cactttactc caagccccct ccccccgcctc accaaccagg acaaccgccc cgtctttacc	300
	agtccgactc cagccgtggc tgcagcacc cccagcccc actccaggga gaaaccttgg	360
	aacctagaat ccaaaccccc tgagctttct atcacatcgt cccttcctcc agggccgagt	420
10	atggcagtg cccactgct cccagaggac agaccagta ctacaccccc tagccaagca	480
	tggactccaa ctccaggagg tcctggagac atggacagac cttgggttcc agaggctcatg	540
	tctaagacca cagggttgg tgtcgaggga accattgcc cctccacagc ttcaggggat	600
15	gacgaagaga ccactaccac catcattacc actactgtca ccacagttca gccaccaggc	660
	ccctgtagct ggaatttctc aggccagag ggctctctgg attccccac ggccccagc	720
	tcaccctctg atgttggcct ggactgttcc tactatatct ctgtctaccc tggatatgga	780
20	gtagagatca aggtggagaa catcagcctt caggaagggg agaccatcac cgtggagggc	840
	ctggggggcc ccgatccact gcccttggct aaccagtcgt tcctgctgag gggccaggtc	900
25	atccgcagcc ccaccacca agcagccctg aggttccaga gcctcccgt acccgctggg	960
	cctggcactt tccatttccg ctaccaagcc tatctcctga gctgccactt tccccagct	1020
	ccagcgtatg gagatgtgac tgtcaccagt ctccaccag gaggcagcgc ccacttccat	1080
30	tgtgccactg gctaccagct caagggtgcc aggttcctca cctgtctcaa tgccaccag	1140
	cccttttggg attcccaaga gcctgtttgc attgctgctt gtggtggagt gattcggaat	1200
	gccaccactg gccgcattgt ctctcctggc ttcccgggga actacagcaa caacctcacc	1260
35	tgccactggt tgctagaggc tccagagagc cagcggctgc acctgcactt tgaaaaggtc	1320
	tccttggcag aagacgacga caggctcatc atccgcaatg gaaataacgt ggaggccccg	1380
40	ccggtgtacg actcctatga ggtggaatac ctgcccattg agggcctgct cagctctggc	1440
	agacacttct tcgtggagtt cagtactgac agcagtgggg cagctgcagg catggccctg	1500
	cgctatgagg ccttccagca aggacattgc tatgagccct ttgtcaaata cggcaacttc	1560
45	agcagcagt caccgtccta ccctgtgggt acaactgtgg agttcagctg tgacctggc	1620
	tacaccctgg agcagggctc catcatcatc gaatgcgtcg acctccacga ccccagtg	1680
	aatgagacag agccagcctg ccgagccgtg tgcagcgggg agatcacaga ctctgcaggc	1740
50	gtggtgctct ctccaaactg gccggagcct tatggccgag ggcaggactg catctggggt	1800
	gtgcatgtgg aggaggacaa gcgcatcatg ctggacatcc gagtgtgcg cataggctct	1860
55	ggggatgtac tgaccttcta cgatggggat gacctcacag cccgggtcct gggccaatac	1920
	tcaggggccc gtggccactt caagctcttt acctccatgg ccgatgtcac catccagttc	1980

EP 3 338 793 A1

cagtcagacc ctgggacctc ggcgctgggt taccagcaag gatttgtcat ccacttcttt 2040
 gaggttcccc gcaacgacac atgtccagag ctacccgaga tccccaacgg ctggaagaac 2100
 5 ccatcacagc ctgagctggt gcacggcacg gtggtcacct atcagtgcta ccctggttac 2160
 caggtggtgg gatccagtat tctcatgtgc cagtgggacc taagctggag tgaggacctg 2220
 ccttcatgcc agagagtgc atcttgccat gacccagggg atgtggagca cagccgacgc 2280
 10 ctcatatcca gcccgaagtt tcccgtagga gcaactgtgc aatatgtctg tgaccagggg 2340
 tttgtgctga cggggagtgc cattctcacc tgccatgatc ggcaagcagg cagtcccaag 2400
 tggagtgaca gggcccccac gtgtctcttg gaacaattca agccgtgcca tggcctcagc 2460
 15 gccccggaga atggtgcccg cagccctgag aagcggcttc acccagcagg ggccaccatc 2520
 cacttctcct gtgcccctgg ttatgtgctg aaggggcagg ccagcatcaa atgcgtgcct 2580
 ggacaccctt cgcattggag tgaccaccca cccatctgta gggctgcctc tctggatggg 2640
 20 ttctacaacg gccgtagcct ggatgttgcc aaggcacctg ccgcctccag tgccctggac 2700
 gctgctcacc tggctgctgc catcttccta ccattggtgg ccatggtggt gctggtggga 2760
 ggagtgtacc tctatctttc cagattccag gggaaaagtc ccctgcaact tccccgaact 2820
 25 catcctcgcc cctataaccg catcacggtg gagtcagcat ttgacaatcc aacttatgag 2880
 actggatctc tttcctttgc aggagacgag agaatatga 2919

30 <210> 11
 <211> 972
 <212> PRT
 <213> Mus musculus

35 <220>
 <221> MISC_FEATURE
 <223> mSCRx17 protein

40 <400> 11

Leu Ser Ser Glu Ala Pro Ile Thr Gly Glu Gly His Ala Thr Gly Ile
 1 5 10 15

45 Arg Glu Thr Asp Gly Glu Leu Thr Ala Ala Pro Thr Pro Glu Gln Ser
 20 25 30

Asp Arg Gly Val His Phe Val Thr Thr Ala Pro Thr Leu Lys Leu Leu
 35 40 45

50 Asn His His Pro Leu Leu Glu Glu Phe Leu Gln Glu Gly Leu Glu Arg
 50 55 60

55 Glu Glu Ala Pro Gln Pro Ala Leu Pro Phe Gln Pro Asp Ser Pro Thr
 65 70 75 80

EP 3 338 793 A1

	His	Phe	Thr	Pro	Ser	Pro	Leu	Pro	Arg	Leu	Thr	Asn	Gln	Asp	Asn	Arg	
					85					90					95		
5	Pro	Val	Phe	Thr	Ser	Pro	Thr	Pro	Ala	Val	Ala	Ala	Ala	Pro	Thr	Gln	
				100					105					110			
10	Pro	His	Ser	Arg	Glu	Lys	Pro	Trp	Asn	Leu	Glu	Ser	Lys	Pro	Pro	Glu	
			115					120					125				
15	Leu	Ser	Ile	Thr	Ser	Ser	Leu	Pro	Pro	Gly	Pro	Ser	Met	Ala	Val	Pro	
		130					135					140					
20	Thr	Leu	Leu	Pro	Glu	Asp	Arg	Pro	Ser	Thr	Thr	Pro	Pro	Ser	Gln	Ala	
	145					150					155					160	
25	Trp	Thr	Pro	Thr	Gln	Glu	Gly	Pro	Gly	Asp	Met	Asp	Arg	Pro	Trp	Val	
					165					170					175		
30	Pro	Glu	Val	Met	Ser	Lys	Thr	Thr	Gly	Leu	Gly	Val	Glu	Gly	Thr	Ile	
				180					185					190			
35	Ala	Thr	Ser	Thr	Ala	Ser	Gly	Asp	Asp	Glu	Glu	Thr	Thr	Thr	Thr	Ile	
			195					200					205				
40	Ile	Thr	Thr	Thr	Val	Thr	Thr	Val	Gln	Pro	Pro	Gly	Pro	Cys	Ser	Trp	
	210						215					220					
45	Asn	Phe	Ser	Gly	Pro	Glu	Gly	Ser	Leu	Asp	Ser	Pro	Thr	Ala	Pro	Ser	
	225					230					235					240	
50	Ser	Pro	Ser	Asp	Val	Gly	Leu	Asp	Cys	Phe	Tyr	Tyr	Ile	Ser	Val	Tyr	
				245					250						255		
55	Pro	Gly	Tyr	Gly	Val	Glu	Ile	Lys	Val	Glu	Asn	Ile	Ser	Leu	Gln	Glu	
				260					265					270			
60	Gly	Glu	Thr	Ile	Thr	Val	Glu	Gly	Leu	Gly	Gly	Pro	Asp	Pro	Leu	Pro	
			275					280					285				
65	Leu	Ala	Asn	Gln	Ser	Phe	Leu	Leu	Arg	Gly	Gln	Val	Ile	Arg	Ser	Pro	
		290					295					300					
70	Thr	His	Gln	Ala	Ala	Leu	Arg	Phe	Gln	Ser	Leu	Pro	Leu	Pro	Ala	Gly	
	305					310					315					320	
75	Pro	Gly	Thr	Phe	His	Phe	Arg	Tyr	Gln	Ala	Tyr	Leu	Leu	Ser	Cys	His	

EP 3 338 793 A1

	325	330	335
5	Phe Pro Arg Arg Pro Ala Tyr Gly Asp Val Thr Val Thr Ser Leu His 340 345 350		
10	Pro Gly Gly Ser Ala His Phe His Cys Ala Thr Gly Tyr Gln Leu Lys 355 360 365		
15	Gly Ala Arg Phe Leu Thr Cys Leu Asn Ala Thr Gln Pro Phe Trp Asp 370 375 380		
20	Ser Gln Glu Pro Val Cys Ile Ala Ala Cys Gly Gly Val Ile Arg Asn 385 390 395 400		
25	Ala Thr Thr Gly Arg Ile Val Ser Pro Gly Phe Pro Gly Asn Tyr Ser 405 410 415		
30	Asn Asn Leu Thr Cys His Trp Leu Leu Glu Ala Pro Glu Ser Gln Arg 420 425 430		
35	Leu His Leu His Phe Glu Lys Val Ser Leu Ala Glu Asp Asp Asp Arg 435 440 445		
40	Leu Ile Ile Arg Asn Gly Asn Asn Val Glu Ala Pro Pro Val Tyr Asp 450 455 460		
45	Ser Tyr Glu Val Glu Tyr Leu Pro Ile Glu Gly Leu Leu Ser Ser Gly 465 470 475 480		
50	Arg His Phe Phe Val Glu Phe Ser Thr Asp Ser Ser Gly Ala Ala Ala 485 490 495		
55	Gly Met Ala Leu Arg Tyr Glu Ala Phe Gln Gln Gly His Cys Tyr Glu 500 505 510		
60	Pro Phe Val Lys Tyr Gly Asn Phe Ser Ser Ser Ala Pro Ser Tyr Pro 515 520 525		
65	Val Gly Thr Thr Val Glu Phe Ser Cys Asp Pro Gly Tyr Thr Leu Glu 530 535 540		
70	Gln Gly Ser Ile Ile Ile Glu Cys Val Asp Leu His Asp Pro Gln Trp 545 550 555 560		
75	Asn Glu Thr Glu Pro Ala Cys Arg Ala Val Cys Ser Gly Glu Ile Thr 565 570 575		

EP 3 338 793 A1

	Asp	Ser	Ala	Gly	Val	Val	Leu	Ser	Pro	Asn	Trp	Pro	Glu	Pro	Tyr	Gly	
				580					585					590			
5	Arg	Gly	Gln	Asp	Cys	Ile	Trp	Gly	Val	His	Val	Glu	Glu	Asp	Lys	Arg	
			595					600					605				
	Ile	Met	Leu	Asp	Ile	Arg	Val	Leu	Arg	Ile	Gly	Ser	Gly	Asp	Val	Leu	
10		610					615					620					
	Thr	Phe	Tyr	Asp	Gly	Asp	Asp	Leu	Thr	Ala	Arg	Val	Leu	Gly	Gln	Tyr	
	625					630					635					640	
15	Ser	Gly	Pro	Arg	Gly	His	Phe	Lys	Leu	Phe	Thr	Ser	Met	Ala	Asp	Val	
					645					650					655		
	Thr	Ile	Gln	Phe	Gln	Ser	Asp	Pro	Gly	Thr	Ser	Ala	Leu	Gly	Tyr	Gln	
20				660					665					670			
	Gln	Gly	Phe	Val	Ile	His	Phe	Phe	Glu	Val	Pro	Arg	Asn	Asp	Thr	Cys	
25			675					680					685				
	Pro	Glu	Leu	Pro	Glu	Ile	Pro	Asn	Gly	Trp	Lys	Asn	Pro	Ser	Gln	Pro	
		690					695					700					
30	Glu	Leu	Val	His	Gly	Thr	Val	Val	Thr	Tyr	Gln	Cys	Tyr	Pro	Gly	Tyr	
	705					710					715					720	
	Gln	Val	Val	Gly	Ser	Ser	Ile	Leu	Met	Cys	Gln	Trp	Asp	Leu	Ser	Trp	
35					725					730					735		
	Ser	Glu	Asp	Leu	Pro	Ser	Cys	Gln	Arg	Val	Thr	Ser	Cys	His	Asp	Pro	
				740					745					750			
40	Gly	Asp	Val	Glu	His	Ser	Arg	Arg	Leu	Ile	Ser	Ser	Pro	Lys	Phe	Pro	
			755					760					765				
	Val	Gly	Ala	Thr	Val	Gln	Tyr	Val	Cys	Asp	Gln	Gly	Phe	Val	Leu	Thr	
45		770					775					780					
	Gly	Ser	Ala	Ile	Leu	Thr	Cys	His	Asp	Arg	Gln	Ala	Gly	Ser	Pro	Lys	
50		785				790					795					800	
	Trp	Ser	Asp	Arg	Ala	Pro	Lys	Cys	Leu	Leu	Glu	Gln	Phe	Lys	Pro	Cys	
					805					810					815		
55	His	Gly	Leu	Ser	Ala	Pro	Glu	Asn	Gly	Ala	Arg	Ser	Pro	Glu	Lys	Arg	
				820					825					830			

EP 3 338 793 A1

Leu His Pro Ala Gly Ala Thr Ile His Phe Ser Cys Ala Pro Gly Tyr
 835 840 845

5 Val Leu Lys Gly Gln Ala Ser Ile Lys Cys Val Pro Gly His Pro Ser
 850 855 860

10 His Trp Ser Asp Pro Pro Pro Ile Cys Arg Ala Ala Ser Leu Asp Gly
 865 870 875 880

Phe Tyr Asn Gly Arg Ser Leu Asp Val Ala Lys Ala Pro Ala Ala Ser
 885 890 895

15 Ser Ala Leu Asp Ala Ala His Leu Ala Ala Ala Ile Phe Leu Pro Leu
 900 905 910

20 Val Ala Met Val Leu Leu Val Gly Gly Val Tyr Leu Tyr Phe Ser Arg
 915 920 925

Phe Gln Gly Lys Ser Pro Leu Gln Leu Pro Arg Thr His Pro Arg Pro
 930 935 940

25 Tyr Asn Arg Ile Thr Val Glu Ser Ala Phe Asp Asn Pro Thr Tyr Glu
 945 950 955 960

30 Thr Gly Ser Leu Ser Phe Ala Gly Asp Glu Arg Ile
 965 970

35 <210> 12
 <211> 2919
 <212> DNA
 <213> Rattus rattus

40 <220>
 <221> misc_feature
 <223> cDNA sequence of rSCRx17 ORF

<400> 12
 ctctcctcag aggtccaat cactggggaa ggtcaagcca cgggcatcag ggagatggat 60

45 ggggagctga ccgcagcccc tacacctgag cagtcagacc gaggcgtcca cttcgtcacc 120

acagccccta ccctcaagct actcaaccac caccacttc tggaggaatt tcttcaagag 180

50 gggctagaag ggagagagga agctccgagg ccggcactgc cttccagcc agactcacct 240

acacccttta ctccaagccc ctttccccgc ctaccaacc aggacaaccg ccctgtcttt 300

accagtccga cgccagctgt agctgcggca cccacgcagc cccactccag aaagaaaccc 360

55 tggaaccag agtcagagcc cccggagctt tacatcacat ctcccctccc tccagggccg 420

agtatggcag tgcccacact gcaccagag gacagaccca gcactacacc cccagccaa 480

EP 3 338 793 A1

	gcatggactc	caaccagga	gggtcctgga	gacatgggca	gaccttgggt	tccagagatc	540
	atgtctaaga	ccacagggct	tggtatcgag	gggaccattg	ccacctccac	agcttcaggg	600
5	gatgacgaag	agaccaccac	caccaccatc	attaccaccg	tcaccacaat	tcagccacca	660
	ggcccctgta	gctggaatth	ctcaggcccc	gagggctctc	tggattcccc	tgcgggtcccc	720
	agcgtcccct	ctgatgttgg	cctggactgt	ctctactaca	tctctgtcta	ccctggatat	780
10	ggagtcgaga	tcaaggtgaa	gaacatcagc	cttcaggaag	gagagaccat	aaccgtggag	840
	ggcctggggg	ggcctgaccc	actgcccttg	gctaaccagt	ctttcctgct	gagggggccag	900
15	gtcatccgca	gccccaccca	ccaggcagcc	gtgaggttcc	aaagccttcc	acttcccgtc	960
	ggacctggta	ctttccatth	ccactaccaa	gcctatctcc	tgagctgcca	ctttcctcgg	1020
	cgtccagctt	atggagatgt	gactgtcacc	agcctccacc	caggaggcag	cgcgcgttc	1080
20	cactgtgcca	ctggctacca	gctaaaggt	gccaggttcc	tcacctgtct	caatgccacc	1140
	cagcccttht	gggattccca	agagcctgtc	tgcattgctg	cttgtggagg	agtgattcgg	1200
	aatgccacca	ctggccgcat	tgtctctcct	ggctttcccg	ggaactacag	caacaacctc	1260
25	acctgccact	ggctgctaga	agcccccgag	agccagcggc	tgacacctga	ctttgaaaag	1320
	gtctccctgg	cagaagatga	cgacaggctc	atcatccgta	acgggaataa	cgtggaggcc	1380
	ccgccagtgt	atgactccta	tgaggtggag	tacctgccca	ttgagggcct	gctcagttct	1440
30	ggcagacact	tcttcgtgga	gttcagtact	gacagcagcg	gggcagccgc	aggcatggca	1500
	ctgcgctatg	aggccttcca	gcaaggacat	tgctatgagc	cctttgtcaa	atacggtaac	1560
35	ttcagcagca	gcgcaccgtc	ctaccctgtg	ggtacgactg	tggagttcag	ctgtgaccct	1620
	ggctacaccc	tggagcaggg	ttccatcatc	atcgaatgcg	tcgacctccg	tgacccccag	1680
	tggaatgaga	cagaaccagc	ctgccgagcc	gtgtgcagcg	gggagatcac	agactctgca	1740
40	ggcgtggtgc	tctctccaaa	ctggccggag	ccttatggcc	gagggcagga	ctgcatctgg	1800
	ggtgtgcatg	tggaggagga	caagcgcata	atgctggaca	tccgagtgtc	gcgcataggc	1860
	tctggggatg	tactgacctt	ctacgatggg	gatgacctga	cagcccgggt	cctggggcaa	1920
45	tactcagggc	cccgtggcca	cttcaagctc	tttacctcca	tggctgatgt	caccattcag	1980
	ttccagtcag	accctgggac	gtcggcgctg	ggttaccagc	aaggatttgt	catccacttc	2040
	tttgaggtgc	cccgcaatga	cacatgtcca	gagcttcccg	agatcccca	cggctggaag	2100
50	aacccatcac	agcctgagct	ggtgcatggc	acggtggtca	cctatcagtg	ctaccccggt	2160
	taccaggtgg	tgggatccag	tattctcatg	tgccagtggg	acctgagctg	gagtgaggac	2220
55	ctgccctcat	gccagagagt	gacatcctgc	catgaccagc	gggatgtgga	gcacagccga	2280
	cgcctcatat	ccagcctcaa	gtttcctgtg	ggagcaactg	tgagtatata	ctgtgaccag	2340

EP 3 338 793 A1

ggtttttgtgc tcacgggtag cgccatcctt acttgccatg atcgtcaagc gggcagtccc 2400
 aagtggagtg acagggcccc caagtgtctc ttggaacagt tcaaaccatg tcatggcctc 2460
 5 agtggccctg agaatggtgc ccgcagccct gagaagaggc tccaccacgc aggggccacc 2520
 attcacttct cctgtgcccc tggttatgtg ctgaagggcc aggccagcat caaatgcgtg 2580
 cctggacacc cctcacattg gagtgatcct ccacccatct gtagggctgc ttctctggat 2640
 10 gggttctaca acggccgtag cctggatggt gccaaaggcac ctgccacctc cagtgccctg 2700
 gatgctgccc acatggcagc tgccatcttt ctaccattgg tggccatggt gttgctggtg 2760
 ggaggagtgt acctctatct ctccagactc cagggaaaaa gtcctctgca gcttcccga 2820
 15 actcatcctc gccctataa ccgtatcacg gtagagtcag catttgacaa tccaacttat 2880
 gagaccggat ctctttcctt tgcaggagac gagagaata 2919

 20 <210> 13
 <211> 973
 <212> PRT
 <213> Rattus rattus

 25 <220>
 <221> MISC_FEATURE
 <223> rSCRx17 protein

 <400> 13
 30 Leu Ser Ser Glu Ala Pro Ile Thr Gly Glu Gly Gln Ala Thr Gly Ile
 1 5 10 15

 Arg Glu Met Asp Gly Glu Leu Thr Ala Ala Pro Thr Pro Glu Gln Ser
 35 20 25 30

 Asp Arg Gly Val His Phe Val Thr Thr Ala Pro Thr Leu Lys Leu Leu
 40 35 40 45

 Asn His His Pro Leu Leu Glu Glu Phe Leu Gln Glu Gly Leu Glu Gly
 50 55 60

 Arg Glu Glu Ala Pro Arg Pro Ala Leu Pro Phe Gln Pro Asp Ser Pro
 45 65 70 75 80

 Thr Pro Phe Thr Pro Ser Pro Leu Pro Arg Leu Thr Asn Gln Asp Asn
 50 85 90 95

 Arg Pro Val Phe Thr Ser Pro Thr Pro Ala Val Ala Ala Ala Pro Thr
 100 105 110

 55 Gln Pro His Ser Arg Lys Lys Pro Trp Asn Pro Glu Ser Glu Pro Pro
 115 120 125

EP 3 338 793 A1

	Glu	Leu	Tyr	Ile	Thr	Ser	Pro	Leu	Pro	Pro	Gly	Pro	Ser	Met	Ala	Val	
	130						135					140					
5	Pro	Thr	Leu	His	Pro	Glu	Asp	Arg	Pro	Ser	Thr	Thr	Pro	Pro	Ser	Gln	
	145					150					155					160	
10	Ala	Trp	Thr	Pro	Thr	Gln	Glu	Gly	Pro	Gly	Asp	Met	Gly	Arg	Pro	Trp	
					165					170					175		
15	Val	Pro	Glu	Ile	Met	Ser	Lys	Thr	Thr	Gly	Leu	Gly	Ile	Glu	Gly	Thr	
				180					185					190			
20	Ile	Ala	Thr	Ser	Thr	Ala	Ser	Gly	Asp	Asp	Glu	Glu	Thr	Thr	Thr	Thr	
			195					200					205				
25	Thr	Ile	Ile	Thr	Thr	Val	Thr	Thr	Ile	Gln	Pro	Pro	Gly	Pro	Cys	Ser	
		210						215					220				
30	Trp	Asn	Phe	Ser	Gly	Pro	Glu	Gly	Ser	Leu	Asp	Ser	Pro	Ala	Val	Pro	
	225					230					235					240	
35	Ser	Val	Pro	Ser	Asp	Val	Gly	Leu	Asp	Cys	Leu	Tyr	Tyr	Ile	Ser	Val	
					245					250					255		
40	Tyr	Pro	Gly	Tyr	Gly	Val	Glu	Ile	Lys	Val	Lys	Asn	Ile	Ser	Leu	Gln	
				260					265					270			
45	Glu	Gly	Glu	Thr	Ile	Thr	Val	Glu	Gly	Leu	Gly	Gly	Pro	Asp	Pro	Leu	
			275					280					285				
50	Pro	Leu	Ala	Asn	Gln	Ser	Phe	Leu	Leu	Arg	Gly	Gln	Val	Ile	Arg	Ser	
		290					295					300					
55	Pro	Thr	His	Gln	Ala	Ala	Val	Arg	Phe	Gln	Ser	Leu	Pro	Leu	Pro	Ala	
	305					310					315					320	
60	Gly	Pro	Gly	Thr	Phe	His	Phe	His	Tyr	Gln	Ala	Tyr	Leu	Leu	Ser	Cys	
					325					330					335		
65	His	Phe	Pro	Arg	Arg	Pro	Ala	Tyr	Gly	Asp	Val	Thr	Val	Thr	Ser	Leu	
				340					345					350			
70	His	Pro	Gly	Gly	Ser	Ala	Arg	Phe	His	Cys	Ala	Thr	Gly	Tyr	Gln	Leu	
			355					360					365				
75	Lys	Gly	Ala	Arg	Phe	Leu	Thr	Cys	Leu	Asn	Ala	Thr	Gln	Pro	Phe	Trp	

EP 3 338 793 A1

	370		375		380												
5	Asp	Ser	Gln	Glu	Pro	Val	Cys	Ile	Ala	Ala	Cys	Gly	Gly	Val	Ile	Arg	
	385					390					395					400	
10	Asn	Ala	Thr	Thr	Gly	Arg	Ile	Val	Ser	Pro	Gly	Phe	Pro	Gly	Asn	Tyr	
					405					410					415		
15	Ser	Asn	Asn	Leu	Thr	Cys	His	Trp	Leu	Leu	Glu	Ala	Pro	Glu	Ser	Gln	
				420					425					430			
20	Arg	Leu	His	Leu	His	Phe	Glu	Lys	Val	Ser	Leu	Ala	Glu	Asp	Asp	Asp	
			435					440					445				
25	Arg	Leu	Ile	Ile	Arg	Asn	Gly	Asn	Asn	Val	Glu	Ala	Pro	Pro	Val	Tyr	
	450						455					460					
30	Asp	Ser	Tyr	Glu	Val	Glu	Tyr	Leu	Pro	Ile	Glu	Gly	Leu	Leu	Ser	Ser	
	465					470					475					480	
35	Gly	Arg	His	Phe	Phe	Val	Glu	Phe	Ser	Thr	Asp	Ser	Ser	Gly	Ala	Ala	
				485						490					495		
40	Ala	Gly	Met	Ala	Leu	Arg	Tyr	Glu	Ala	Phe	Gln	Gln	Gly	His	Cys	Tyr	
			500						505					510			
45	Glu	Pro	Phe	Val	Lys	Tyr	Gly	Asn	Phe	Ser	Ser	Ser	Ala	Pro	Ser	Tyr	
			515					520					525				
50	Pro	Val	Gly	Thr	Thr	Val	Glu	Phe	Ser	Cys	Asp	Pro	Gly	Tyr	Thr	Leu	
	530						535					540					
55	Glu	Gln	Gly	Ser	Ile	Ile	Ile	Glu	Cys	Val	Asp	Leu	Arg	Asp	Pro	Gln	
	545					550					555					560	
60	Trp	Asn	Glu	Thr	Glu	Pro	Ala	Cys	Arg	Ala	Val	Cys	Ser	Gly	Glu	Ile	
					565					570					575		
65	Thr	Asp	Ser	Ala	Gly	Val	Val	Leu	Ser	Pro	Asn	Trp	Pro	Glu	Pro	Tyr	
				580					585					590			
70	Gly	Arg	Gly	Gln	Asp	Cys	Ile	Trp	Gly	Val	His	Val	Glu	Glu	Asp	Lys	
			595					600					605				
75	Arg	Ile	Met	Leu	Asp	Ile	Arg	Val	Leu	Arg	Ile	Gly	Ser	Gly	Asp	Val	
	610						615					620					

EP 3 338 793 A1

	Leu	Thr	Phe	Tyr	Asp	Gly	Asp	Asp	Leu	Thr	Ala	Arg	Val	Leu	Gly	Gln	625	630	635	640
5	Tyr	Ser	Gly	Pro	Arg	Gly	His	Phe	Lys	Leu	Phe	Thr	Ser	Met	Ala	Asp	645	650	655	
10	Val	Thr	Ile	Gln	Phe	Gln	Ser	Asp	Pro	Gly	Thr	Ser	Ala	Leu	Gly	Tyr	660	665	670	
	Gln	Gln	Gly	Phe	Val	Ile	His	Phe	Phe	Glu	Val	Pro	Arg	Asn	Asp	Thr	675	680	685	
15	Cys	Pro	Glu	Leu	Pro	Glu	Ile	Pro	Asn	Gly	Trp	Lys	Asn	Pro	Ser	Gln	690	695	700	
20	Pro	Glu	Leu	Val	His	Gly	Thr	Val	Val	Thr	Tyr	Gln	Cys	Tyr	Pro	Gly	705	710	715	720
	Tyr	Gln	Val	Val	Gly	Ser	Ser	Ile	Leu	Met	Cys	Gln	Trp	Asp	Leu	Ser	725	730	735	
25	Trp	Ser	Glu	Asp	Leu	Pro	Ser	Cys	Gln	Arg	Val	Thr	Ser	Cys	His	Asp	740	745	750	
30	Pro	Gly	Asp	Val	Glu	His	Ser	Arg	Arg	Leu	Ile	Ser	Ser	Leu	Lys	Phe	755	760	765	
35	Pro	Val	Gly	Ala	Thr	Val	Gln	Tyr	Ile	Cys	Asp	Gln	Gly	Phe	Val	Leu	770	775	780	
	Thr	Gly	Ser	Ala	Ile	Leu	Thr	Cys	His	Asp	Arg	Gln	Ala	Gly	Ser	Pro	785	790	795	800
40	Lys	Trp	Ser	Asp	Arg	Ala	Pro	Lys	Cys	Leu	Leu	Glu	Gln	Phe	Lys	Pro	805	810	815	
45	Cys	His	Gly	Leu	Ser	Ala	Pro	Glu	Asn	Gly	Ala	Arg	Ser	Pro	Glu	Lys	820	825	830	
	Arg	Leu	His	Pro	Ala	Gly	Ala	Thr	Ile	His	Phe	Ser	Cys	Ala	Pro	Gly	835	840	845	
50	Tyr	Val	Leu	Lys	Gly	Gln	Ala	Ser	Ile	Lys	Cys	Val	Pro	Gly	His	Pro	850	855	860	
55	Ser	His	Trp	Ser	Asp	Pro	Pro	Pro	Ile	Cys	Arg	Ala	Ala	Ser	Leu	Asp	865	870	875	880

EP 3 338 793 A1

Gly Phe Tyr Asn Gly Arg Ser Leu Asp Val Ala Lys Ala Pro Ala Thr
885 890 895

5 Ser Ser Ala Leu Asp Ala Ala His Met Ala Ala Ala Ile Phe Leu Pro
900 905 910

10 Leu Val Ala Met Val Leu Leu Val Gly Gly Val Tyr Leu Tyr Phe Ser
915 920 925

Arg Leu Gln Gly Lys Ser Pro Leu Gln Leu Pro Gly Thr His Pro Arg
930 935 940

15 Pro Tyr Asn Arg Ile Thr Val Glu Ser Ala Phe Asp Asn Pro Thr Tyr
945 950 955 960

20 Glu Thr Gly Ser Leu Ser Phe Ala Gly Asp Glu Arg Ile
965 970

25 <210> 14
<211> 2997
<212> DNA
<213> *Macaca fascicularis*

30 <220>
<221> misc_feature
<223> cDNA sequence of cSCRx17 ORF

<400> 14
atggagacag acacactcct gctatgggta ctgctgctct ggggtccagg ttccactggt 60
35 gacggcgcg cactcagcag cgaagctccc acaatgggca agggacaggc ccccgaatt 120
gaagaaaccg atggcgaact caccgctgcc cctacccttg agcaaccgga aaggggagtg 180
cactttgtga ccaccgctcc caccctgaag ctgctcaatc accaccccct cctggaggag 240
40 tttctgcagg aaggcctgga aaaaggcgac gaggaactca gacctgccct gcccttccaa 300
cccgaccctc ctaccccctt tacacctagc cctctcccta gactggccaa ccaagactcc 360
agacctgtgt tcaccagccc tacacctgct acagctgccg tcctaccca acctcaatcc 420
45 aaggagggac cttggagcct cgagagcgag cctcccgtgc tgagaatcac agctcctctc 480
cctcctggcc cttccatggc tgtccccaca ctgggacctg gcgaaaggcc cagcacaaca 540
cccccctcca gagcctggac ccctacacaa gaaggccctg gcgacatggg aaggccttgg 600
50 gtccctgaag tcgtgagcca aggcgcccgc atcggaatcc agggaacat ccgcagctcc 660
acagccagcg gagacgatga ggaaacaacc accacaacca ccatcatcac caccacaatc 720
55 acaacagtcc agacccccgg cccttgacgc tggaattttt ccggccctga gggatccctg 780
gattccccca cagatctgtc ctcccctcct gacgtgggccc tcgactgttt cttctatatc 840

EP 3 338 793 A1

	tccgtgtatc	ctggctacgg	cgtcgaaatc	aaagtccaga	acatctccct	gagggagggc	900
	gaaacagtca	ccgtggaagg	actgggcgga	cccgtcctc	tgcctctcgc	caaccaatcc	960
5	ttcctcctca	ggggccaagt	gattagatcc	cccacacatc	aagctgctct	cagggttccaa	1020
	agcctccctc	cccccgctgg	acccggaacc	tttcaacttc	actaccaagc	ctatctcctc	1080
	agctgccatt	tccccacag	gcccgttat	ggagatgtca	cagtcacctc	cctgcatact	1140
10	ggcggctccg	ctagattcca	ctgcgctacc	ggataccaac	tcaagggcgc	caggcatctg	1200
	acatgtctca	atgctacca	gcccttctgg	gacagcaagg	agcccgctctg	cattgccgct	1260
	tgcggagggc	tcatcagaaa	tgccaccacc	ggcagaatcg	tgagccccgg	cttccctggc	1320
15	aactactcca	acaacctgac	atgccactgg	ctgctggaag	ctcctgaggg	ccagagactg	1380
	catctgcact	tcgagaaggt	cagcctggcc	gaagatgacg	acagactcat	catcaggaac	1440
20	ggcgacaacg	tggaggctcc	ccccgtctat	gattcctacg	aggtcgagta	cctccccatc	1500
	gagggactgc	tgtcctccgg	caagcatttt	ttcgtggagc	tgtccacaga	ttccagcgga	1560
	gctgccgccg	gaatggctct	caggtacgag	gctttccaac	agggccactg	ttacgagccc	1620
25	tttgtgaagt	acggcaactt	ctccagctcc	gctcctacct	accccgctcg	cacaaccgtc	1680
	gaatttagct	gcgacctgg	atacacactc	gagcaaggct	ccatcatcat	cgagtgtgtc	1740
	gacccccacg	acccccaatg	gaacgagaca	gagcccgct	gtagggccgt	gtgtagcgga	1800
30	gagattaccg	actccgccgg	agtgggtgtc	tcccctaatt	ggcctgaacc	ctacggcaga	1860
	ggacaagatt	gtatttgggg	cgtccatgtc	gaggaggaca	agaggattat	gctcgacgtg	1920
35	aggggtgctga	ggattggacc	tggcgacgtg	ctcacattct	atgacggcga	cgatctcacc	1980
	gccagagtcc	tgggacaata	ctccggccct	cacagccact	tcaagctgtt	caccagcatg	2040
	gctgacgtga	ccatccagtt	ccagtcgat	cctggaacat	ccgtgctggg	ataccagcag	2100
40	ggcttcgtca	tccacttctt	cgagggtccc	aggaacgaca	cctgccccga	actgcccag	2160
	attcccaacg	gctggaaatc	cccctcccaa	cctgatctcg	tgcacggcac	cgtcgtcacc	2220
	taccaatgct	accctggata	ccaagtcgtc	ggcagcagcg	tgctgatgtg	ccaatgggac	2280
45	ctcacctgga	gcgaggatct	gccctcctgc	cagagagtca	cctcctgcca	cgatcccggc	2340
	gatgtggaac	actccaggag	gctgattagc	tcccccaagt	tcctgtcgg	agccaccgtg	2400
	caatacatct	gcgaccaggg	ctttgtgctg	accggaacca	gcatcctcac	atgccacgac	2460
50	aggcaagctg	gatcccccaa	gtgggtccgat	agggccccca	aatgcctcct	ggaacagctg	2520
	aagccttgct	atggcctcag	cgctcctgaa	aacggcgcta	ggagccccga	aaagaggctc	2580
55	caccctgccg	gagccaccat	ccacttttcc	tgtgcccccg	gatactgtct	gaagggccag	2640
	gcctccatta	agtgcgtgcc	cggacatcct	tcccactgg	ccgaccccc	tcccatctgt	2700

EP 3 338 793 A1

aaagccgcct ccctggacgg attctataac agcagaagcc tggacgtcgc taaggcccct 2760
gctgcttcct ccaccctgga tgctgctcac atcgctgctg ccacctttct gcccctcgtc 2820
5 gccatggtgc tgctggtggg aggcgtctac ttctacttct ccaggctgca gggaaagagc 2880
tccctgcaac tgcctaggac aagaccagc ccctacaata ggatcacagt cgagagcgcc 2940
ttcgacaacc ccacatacga gacaggatcc ctgagctttg ccggagacga gagaatt 2997
10
<210> 15
<211> 999
<212> PRT
<213> Macaca fascicularis
15
<220>
<221> MISC_FEATURE
<223> cSCRx17 protein
20
<400> 15
Met Glu Thr Asp Thr Leu Leu Leu Trp Val Leu Leu Leu Trp Val Pro
1 5 10 15
25
Gly Ser Thr Gly Asp Gly Ala Pro Leu Ser Ser Glu Ala Pro Thr Met
20 25 30
30
Gly Lys Gly Gln Ala Pro Gly Ile Glu Glu Thr Asp Gly Glu Leu Thr
35 40 45
Ala Ala Pro Thr Pro Glu Gln Pro Glu Arg Gly Val His Phe Val Thr
50 55 60
35
Thr Ala Pro Thr Leu Lys Leu Leu Asn His His Pro Leu Leu Glu Glu
65 70 75 80
40
Phe Leu Gln Glu Gly Leu Glu Lys Gly Asp Glu Glu Leu Arg Pro Ala
85 90 95
45
Leu Pro Phe Gln Pro Asp Pro Pro Thr Pro Phe Thr Pro Ser Pro Leu
100 105 110
Pro Arg Leu Ala Asn Gln Asp Ser Arg Pro Val Phe Thr Ser Pro Thr
115 120 125
50
Pro Ala Thr Ala Ala Val Pro Thr Gln Pro Gln Ser Lys Glu Gly Pro
130 135 140
55
Trp Ser Leu Glu Ser Glu Pro Pro Val Leu Arg Ile Thr Ala Pro Leu
145 150 155 160

EP 3 338 793 A1

	Pro	Pro	Gly	Pro	Ser	Met	Ala	Val	Pro	Thr	Leu	Gly	Pro	Gly	Glu	Arg	
					165					170					175		
5	Pro	Ser	Thr	Thr	Pro	Pro	Ser	Arg	Ala	Trp	Thr	Pro	Thr	Gln	Glu	Gly	
				180					185					190			
	Pro	Gly	Asp	Met	Gly	Arg	Pro	Trp	Val	Pro	Glu	Val	Val	Ser	Gln	Gly	
10			195					200					205				
	Ala	Gly	Ile	Gly	Ile	Gln	Gly	Thr	Ile	Ala	Ser	Ser	Thr	Ala	Ser	Gly	
		210					215					220					
15	Asp	Asp	Glu	Glu	Thr	Thr	Thr	Thr	Thr	Thr	Ile	Ile	Thr	Thr	Thr	Ile	
	225					230					235					240	
	Thr	Thr	Val	Gln	Thr	Pro	Gly	Pro	Cys	Ser	Trp	Asn	Phe	Ser	Gly	Pro	
20					245					250					255		
	Glu	Gly	Ser	Leu	Asp	Ser	Pro	Thr	Asp	Leu	Ser	Ser	Pro	Pro	Asp	Val	
25				260					265					270			
	Gly	Leu	Asp	Cys	Phe	Phe	Tyr	Ile	Ser	Val	Tyr	Pro	Gly	Tyr	Gly	Val	
			275					280					285				
30	Glu	Ile	Lys	Val	Gln	Asn	Ile	Ser	Leu	Arg	Glu	Gly	Glu	Thr	Val	Thr	
		290					295					300					
	Val	Glu	Gly	Leu	Gly	Gly	Pro	Ala	Pro	Leu	Pro	Leu	Ala	Asn	Gln	Ser	
35	305					310					315					320	
	Phe	Leu	Leu	Arg	Gly	Gln	Val	Ile	Arg	Ser	Pro	Thr	His	Gln	Ala	Ala	
					325					330					335		
40	Leu	Arg	Phe	Gln	Ser	Leu	Pro	Pro	Pro	Ala	Gly	Pro	Gly	Thr	Phe	His	
				340					345					350			
	Phe	His	Tyr	Gln	Ala	Tyr	Leu	Leu	Ser	Cys	His	Phe	Pro	His	Arg	Pro	
45			355					360					365				
	Ala	Tyr	Gly	Asp	Val	Thr	Val	Thr	Ser	Leu	His	Pro	Gly	Gly	Ser	Ala	
50		370					375					380					
	Arg	Phe	His	Cys	Ala	Thr	Gly	Tyr	Gln	Leu	Lys	Gly	Ala	Arg	His	Leu	
	385					390					395					400	
55	Thr	Cys	Leu	Asn	Ala	Thr	Gln	Pro	Phe	Trp	Asp	Ser	Lys	Glu	Pro	Val	
				405						410					415		

EP 3 338 793 A1

	Cys	Ile	Ala	Ala	Cys	Gly	Gly	Val	Ile	Arg	Asn	Ala	Thr	Thr	Gly	Arg	
				420					425						430		
5	Ile	Val	Ser	Pro	Gly	Phe	Pro	Gly	Asn	Tyr	Ser	Asn	Asn	Leu	Thr	Cys	
			435					440					445				
	His	Trp	Leu	Leu	Glu	Ala	Pro	Glu	Gly	Gln	Arg	Leu	His	Leu	His	Phe	
10		450					455					460					
	Glu	Lys	Val	Ser	Leu	Ala	Glu	Asp	Asp	Asp	Arg	Leu	Ile	Ile	Arg	Asn	
	465					470					475					480	
15	Gly	Asp	Asn	Val	Glu	Ala	Pro	Pro	Val	Tyr	Asp	Ser	Tyr	Glu	Val	Glu	
					485					490					495		
	Tyr	Leu	Pro	Ile	Glu	Gly	Leu	Leu	Ser	Ser	Gly	Lys	His	Phe	Phe	Val	
20				500					505					510			
	Glu	Leu	Ser	Thr	Asp	Ser	Ser	Gly	Ala	Ala	Ala	Gly	Met	Ala	Leu	Arg	
25			515					520					525				
	Tyr	Glu	Ala	Phe	Gln	Gln	Gly	His	Cys	Tyr	Glu	Pro	Phe	Val	Lys	Tyr	
	530						535					540					
30	Gly	Asn	Phe	Ser	Ser	Ser	Ala	Pro	Thr	Tyr	Pro	Val	Gly	Thr	Thr	Val	
	545					550					555					560	
	Glu	Phe	Ser	Cys	Asp	Pro	Gly	Tyr	Thr	Leu	Glu	Gln	Gly	Ser	Ile	Ile	
35					565					570					575		
	Ile	Glu	Cys	Val	Asp	Pro	His	Asp	Pro	Gln	Trp	Asn	Glu	Thr	Glu	Pro	
				580					585					590			
40	Ala	Cys	Arg	Ala	Val	Cys	Ser	Gly	Glu	Ile	Thr	Asp	Ser	Ala	Gly	Val	
			595					600					605				
	Val	Leu	Ser	Pro	Asn	Trp	Pro	Glu	Pro	Tyr	Gly	Arg	Gly	Gln	Asp	Cys	
45		610					615					620					
	Ile	Trp	Gly	Val	His	Val	Glu	Glu	Asp	Lys	Arg	Ile	Met	Leu	Asp	Val	
50	625					630					635					640	
	Arg	Val	Leu	Arg	Ile	Gly	Pro	Gly	Asp	Val	Leu	Thr	Phe	Tyr	Asp	Gly	
					645					650					655		
55	Asp	Asp	Leu	Thr	Ala	Arg	Val	Leu	Gly	Gln	Tyr	Ser	Gly	Pro	His	Ser	
				660					665					670			

EP 3 338 793 A1

	His Phe Lys Leu Phe Thr Ser Met Ala Asp Val Thr Ile Gln Phe Gln	
	675 680 685	
5	Ser Asp Pro Gly Thr Ser Val Leu Gly Tyr Gln Gln Gly Phe Val Ile	
	690 695 700	
10	His Phe Phe Glu Val Pro Arg Asn Asp Thr Cys Pro Glu Leu Pro Glu	
	705 710 715 720	
15	Ile Pro Asn Gly Trp Lys Ser Pro Ser Gln Pro Asp Leu Val His Gly	
	725 730 735	
20	Thr Val Val Thr Tyr Gln Cys Tyr Pro Gly Tyr Gln Val Val Gly Ser	
	740 745 750	
25	Ser Val Leu Met Cys Gln Trp Asp Leu Thr Trp Ser Glu Asp Leu Pro	
	755 760 765	
30	Ser Cys Gln Arg Val Thr Ser Cys His Asp Pro Gly Asp Val Glu His	
	770 775 780	
35	Ser Arg Arg Leu Ile Ser Ser Pro Lys Phe Pro Val Gly Ala Thr Val	
	785 790 795 800	
40	Gln Tyr Ile Cys Asp Gln Gly Phe Val Leu Thr Gly Thr Ser Ile Leu	
	805 810 815	
45	Thr Cys His Asp Arg Gln Ala Gly Ser Pro Lys Trp Ser Asp Arg Ala	
	820 825 830	
50	Pro Lys Cys Leu Leu Glu Gln Leu Lys Pro Cys His Gly Leu Ser Ala	
	835 840 845	
55	Pro Glu Asn Gly Ala Arg Ser Pro Glu Lys Arg Leu His Pro Ala Gly	
	850 855 860	
60	Ala Thr Ile His Phe Ser Cys Ala Pro Gly Tyr Val Leu Lys Gly Gln	
	865 870 875 880	
65	Ala Ser Ile Lys Cys Val Pro Gly His Pro Ser His Trp Ser Asp Pro	
	885 890 895	
70	Pro Pro Ile Cys Lys Ala Ala Ser Leu Asp Gly Phe Tyr Asn Ser Arg	
	900 905 910	
75	Ser Leu Asp Val Ala Lys Ala Pro Ala Ala Ser Ser Thr Leu Asp Ala	

EP 3 338 793 A1

915

920

925

Ala His Ile Ala Ala Ala Ile Phe Leu Pro Leu Val Ala Met Val Leu
930 935 940

Leu Val Gly Gly Val Tyr Phe Tyr Phe Ser Arg Leu Gln Gly Lys Ser
945 950 955 960

Ser Leu Gln Leu Pro Arg Thr Arg Pro Arg Pro Tyr Asn Arg Ile Thr
965 970 975

Val Glu Ser Ala Phe Asp Asn Pro Thr Tyr Glu Thr Gly Ser Leu Ser
980 985 990

Phe Ala Gly Asp Glu Arg Ile
995

<210> 16
<211> 2781
<212> DNA
<213> Homo sapiens

<220>
<221> misc_feature
<223> cDNA sequence of human SEZ6L ECD

<400> 16
ctcgagaggg atgctctgcc tgagggagat gcttcccctc tcggacctta tctgctgccc 60
agcggagctc ctgagagggg atccccgga aaggagcatc ccgaagaaag agtggtcaca 120
gctcccccta gctccagcca gagcgctgag gtgctgggag aactggctct cgacggaaca 180
gccccttccg cccatcacga tattcctgcc ctcagccctc tcctccccga ggaagctagg 240
cctaaacacg ccctcccccc taaaaagaag ctgccttccc tcaagcaggt caattccgcc 300
aggaagcagc tcagacccaa ggccacctcc gctgctacag tccaaagagc tggatcccag 360
cctgccagcc agggactcga tctgctcagc agctccacag aaaaacctgg acctcctggc 420
gatcctgacc ctattgtggc cagcgaagaa gctagcgaag tccctctgtg gctggacagg 480
aaggagtccg ctgtccccac cacaccgct cctctccaga tcagcccctt cacctcccag 540
ccttatgtcg ctcatacct gcctcagagg cctgagcctg gcgaacctgg acctgacatg 600
gctcaggagg ctccctcagga ggacaccagc cctatggccc tgatggataa gggcgagaat 660
gaactgaccg gaagcgccag cgaggaaagc caggagacca ccaccagcac aatcatcacc 720
accaccgtca tcaccaccga acaggccccc gctctgtgtt ccgtgtcctt ttccaacccc 780
gagggctaca ttgacagcag cgattacccc ctgctccctc tcaacaactt cctcgagtgc 840
acctacaatg tgaccgtgta caccggctac ggagtcgaac tccaggtgaa gtccgtgaac 900

EP 3 338 793 A1

	ctctccgatg	gcgaactgct	ctccattagg	ggcgtcgatg	gccctacact	caccgtcctg	960
	gctaaccaaa	ccctgctcgt	cgaaggccag	gtgattaggt	ccccaccaa	caccatctcc	1020
5	gtctacttca	ggacctttca	agacgacgga	ctgggaacct	tccaactgca	ttaccaggcc	1080
	ttcatgctgt	cctgtaattt	cccaggaga	cccgactccg	gagacgtcac	cgtcatggat	1140
	ctgcactccg	gaggcgtggc	ccactttcat	tgtcacctcg	gctacgagct	ccagggcgcc	1200
10	aagatgctga	catgcatcaa	cgccagcaa	cctcactggg	ccagccagga	gcctatctgt	1260
	agcgtcctt	gcggcgaggc	cgtgcacaat	gctacaattg	gcagagtgt	cagcccttcc	1320
	taccctgaaa	acaccaacgg	ctcccagttc	tgcactctgga	caatcgaggc	ccccgaaggc	1380
15	caaaagctgc	acctgcactt	tgagaggctc	ctgctccacg	acaaagacag	gatgaccgtg	1440
	cactccggcc	agaccaataa	gtccgcccct	ctgtatgaca	gcctgcagac	agagtccgtc	1500
20	ccttttgaag	gcctgctgtc	cgagggcaat	accatcagga	ttgagttcac	atccgaccaa	1560
	gccagggctg	ctagcacctt	caacattagg	tttgaggctt	tcgaaaagg	acactgctac	1620
	gagccctata	ttcagaatgg	caatttcaca	acctccgacc	ccacctacaa	tatcggcaca	1680
25	attgtggagt	ttacctgcga	ccctggacac	agcctggagc	agggacctgc	catcatcgaa	1740
	tgcataacg	tcagggaccc	ctactggaac	gacacagaac	ctctgtgtag	ggctatgtgc	1800
	ggaggcgaac	tgagcgctgt	ggctggagtc	gtgctctccc	ctaactggcc	cgaaccctat	1860
30	gtggagggcg	aagattgcat	ctggaagatc	cacgtcggcg	aggaaaaaag	gatctttctg	1920
	gacatccagt	tcctgaatct	ctccaacagc	gacatcctga	ccatctacga	cggagatgag	1980
35	gtcatgcccc	acattctggg	ccagtatctc	ggaaactccg	gcccccaaaa	gctctactcc	2040
	tccacccccg	acctcacaat	ccaattccac	agcgatcctg	ctggcctcat	ctttggaaag	2100
	ggacaaggct	ttatcatgaa	ttacatcgag	gtcagcagaa	acgacagctg	ctccgacctg	2160
40	cctgagatcc	agaacggatg	gaagaccacc	tcccacaccg	agctcgtcag	gggagctagg	2220
	atcacatacc	agtgcgaccc	cggatacgac	atcgtcgggt	ccgataccct	gacatgccag	2280
	tgggatctga	gctggagctc	cgaccccccc	ttttgtgaga	agatcatgta	ctgcaccgac	2340
45	cccggcgaag	tcgatcatag	caccaggctc	atcagcgatc	ctgtgctgct	cgtcggcaca	2400
	accatccaat	acacctgtaa	ccccggattc	gtgctcgaag	gatcctccct	gctcacctgt	2460
	tacagcaggg	aaaccggcac	ccccatttgg	acatccaggc	tgcctcactg	cgtgtccgaa	2520
50	gagagcctgg	cttgcgataa	tcccggcctg	cctgagaacg	gataccagat	tctgtacaaa	2580
	aggctgtacc	tccccggcga	gtccctgacc	ttcatgtgct	acgaaggatt	cgagctcatg	2640
55	ggcgaagtca	ccatcagggtg	catcctcggc	cagccctccc	actggaacgg	acctctcccc	2700
	gtctgtaagg	tcaatcagga	ttccttcgag	cacgtctctg	aagtcgctga	ggctgccgcc	2760

2781

```

5      <210> 17
      <211> 964
      <212> PRT
      <213> Homo sapiens

10     <220>
      <221> MISC_FEATURE
      <223> Human SEZ6L ECD protein

      <400> 17

15     Met Glu Thr Asp Thr Leu Leu Leu Trp Val Leu Leu Leu Trp Val Pro
      1          5          10          15

20     Gly Ser Thr Gly Asp His Gly Ala Pro Leu Glu Arg Asp Ala Leu Pro
      20          25          30

25     Glu Gly Asp Ala Ser Pro Leu Gly Pro Tyr Leu Leu Pro Ser Gly Ala
      35          40          45

30     Pro Glu Arg Gly Ser Pro Gly Lys Glu His Pro Glu Glu Arg Val Val
      50          55          60

35     Thr Ala Pro Pro Ser Ser Ser Gln Ser Ala Glu Val Leu Gly Glu Leu
      65          70          75          80

40     Val Leu Asp Gly Thr Ala Pro Ser Ala His His Asp Ile Pro Ala Leu
      85          90          95

45     Ser Pro Leu Leu Pro Glu Glu Ala Arg Pro Lys His Ala Leu Pro Pro
      100         105         110

50     Lys Lys Lys Leu Pro Ser Leu Lys Gln Val Asn Ser Ala Arg Lys Gln
      115         120         125

55     Leu Arg Pro Lys Ala Thr Ser Ala Ala Thr Val Gln Arg Ala Gly Ser
      130         135         140

60     Gln Pro Ala Ser Gln Gly Leu Asp Leu Leu Ser Ser Ser Thr Glu Lys
      145         150         155         160

65     Pro Gly Pro Pro Gly Asp Pro Asp Pro Ile Val Ala Ser Glu Glu Ala
      165         170         175

70     Ser Glu Val Pro Leu Trp Leu Asp Arg Lys Glu Ser Ala Val Pro Thr
      180         185         190

```

EP 3 338 793 A1

	Thr	Pro	Ala	Pro	Leu	Gln	Ile	Ser	Pro	Phe	Thr	Ser	Gln	Pro	Tyr	Val	
			195					200					205				
5	Ala	His	Thr	Leu	Pro	Gln	Arg	Pro	Glu	Pro	Gly	Glu	Pro	Gly	Pro	Asp	
		210					215					220					
	Met	Ala	Gln	Glu	Ala	Pro	Gln	Glu	Asp	Thr	Ser	Pro	Met	Ala	Leu	Met	
10		225				230					235					240	
	Asp	Lys	Gly	Glu	Asn	Glu	Leu	Thr	Gly	Ser	Ala	Ser	Glu	Glu	Ser	Gln	
					245					250					255		
15	Glu	Thr	Thr	Thr	Ser	Thr	Ile	Ile	Thr	Thr	Thr	Val	Ile	Thr	Thr	Glu	
				260					265					270			
	Gln	Ala	Pro	Ala	Leu	Cys	Ser	Val	Ser	Phe	Ser	Asn	Pro	Glu	Gly	Tyr	
20			275					280					285				
	Ile	Asp	Ser	Ser	Asp	Tyr	Pro	Leu	Leu	Pro	Leu	Asn	Asn	Phe	Leu	Glu	
25		290					295					300					
	Cys	Thr	Tyr	Asn	Val	Thr	Val	Tyr	Thr	Gly	Tyr	Gly	Val	Glu	Leu	Gln	
		305				310					315					320	
30	Val	Lys	Ser	Val	Asn	Leu	Ser	Asp	Gly	Glu	Leu	Leu	Ser	Ile	Arg	Gly	
				325						330					335		
	Val	Asp	Gly	Pro	Thr	Leu	Thr	Val	Leu	Ala	Asn	Gln	Thr	Leu	Leu	Val	
35				340					345					350			
	Glu	Gly	Gln	Val	Ile	Arg	Ser	Pro	Thr	Asn	Thr	Ile	Ser	Val	Tyr	Phe	
			355					360					365				
40	Arg	Thr	Phe	Gln	Asp	Asp	Gly	Leu	Gly	Thr	Phe	Gln	Leu	His	Tyr	Gln	
		370					375					380					
	Ala	Phe	Met	Leu	Ser	Cys	Asn	Phe	Pro	Arg	Arg	Pro	Asp	Ser	Gly	Asp	
45		385				390					395					400	
	Val	Thr	Val	Met	Asp	Leu	His	Ser	Gly	Gly	Val	Ala	His	Phe	His	Cys	
50				405						410					415		
	His	Leu	Gly	Tyr	Glu	Leu	Gln	Gly	Ala	Lys	Met	Leu	Thr	Cys	Ile	Asn	
			420						425					430			
55	Ala	Ser	Lys	Pro	His	Trp	Ser	Ser	Gln	Glu	Pro	Ile	Cys	Ser	Ala	Pro	
			435					440					445				

EP 3 338 793 A1

	Cys	Gly	Gly	Ala	Val	His	Asn	Ala	Thr	Ile	Gly	Arg	Val	Leu	Ser	Pro	
	450						455					460					
5	Ser	Tyr	Pro	Glu	Asn	Thr	Asn	Gly	Ser	Gln	Phe	Cys	Ile	Trp	Thr	Ile	
	465					470					475					480	
	Glu	Ala	Pro	Glu	Gly	Gln	Lys	Leu	His	Leu	His	Phe	Glu	Arg	Leu	Leu	
10					485					490					495		
	Leu	His	Asp	Lys	Asp	Arg	Met	Thr	Val	His	Ser	Gly	Gln	Thr	Asn	Lys	
				500					505					510			
15	Ser	Ala	Leu	Leu	Tyr	Asp	Ser	Leu	Gln	Thr	Glu	Ser	Val	Pro	Phe	Glu	
			515					520					525				
	Gly	Leu	Leu	Ser	Glu	Gly	Asn	Thr	Ile	Arg	Ile	Glu	Phe	Thr	Ser	Asp	
20		530					535					540					
	Gln	Ala	Arg	Ala	Ala	Ser	Thr	Phe	Asn	Ile	Arg	Phe	Glu	Ala	Phe	Glu	
25		545				550					555					560	
	Lys	Gly	His	Cys	Tyr	Glu	Pro	Tyr	Ile	Gln	Asn	Gly	Asn	Phe	Thr	Thr	
				565						570					575		
30	Ser	Asp	Pro	Thr	Tyr	Asn	Ile	Gly	Thr	Ile	Val	Glu	Phe	Thr	Cys	Asp	
				580					585					590			
	Pro	Gly	His	Ser	Leu	Glu	Gln	Gly	Pro	Ala	Ile	Ile	Glu	Cys	Ile	Asn	
35			595					600					605				
	Val	Arg	Asp	Pro	Tyr	Trp	Asn	Asp	Thr	Glu	Pro	Leu	Cys	Arg	Ala	Met	
		610					615					620					
40	Cys	Gly	Gly	Glu	Leu	Ser	Ala	Val	Ala	Gly	Val	Val	Leu	Ser	Pro	Asn	
	625					630					635					640	
	Trp	Pro	Glu	Pro	Tyr	Val	Glu	Gly	Glu	Asp	Cys	Ile	Trp	Lys	Ile	His	
45					645					650					655		
	Val	Gly	Glu	Glu	Lys	Arg	Ile	Phe	Leu	Asp	Ile	Gln	Phe	Leu	Asn	Leu	
50				660					665					670			
	Ser	Asn	Ser	Asp	Ile	Leu	Thr	Ile	Tyr	Asp	Gly	Asp	Glu	Val	Met	Pro	
			675					680					685				
55	His	Ile	Leu	Gly	Gln	Tyr	Leu	Gly	Asn	Ser	Gly	Pro	Gln	Lys	Leu	Tyr	
	690						695					700					

EP 3 338 793 A1

	Ser	Ser	Thr	Pro	Asp	Leu	Thr	Ile	Gln	Phe	His	Ser	Asp	Pro	Ala	Gly	
	705					710					715					720	
5	Leu	Ile	Phe	Gly	Lys	Gly	Gln	Gly	Phe	Ile	Met	Asn	Tyr	Ile	Glu	Val	
					725					730					735		
10	Ser	Arg	Asn	Asp	Ser	Cys	Ser	Asp	Leu	Pro	Glu	Ile	Gln	Asn	Gly	Trp	
				740					745					750			
15	Lys	Thr	Thr	Ser	His	Thr	Glu	Leu	Val	Arg	Gly	Ala	Arg	Ile	Thr	Tyr	
			755					760					765				
20	Gln	Cys	Asp	Pro	Gly	Tyr	Asp	Ile	Val	Gly	Ser	Asp	Thr	Leu	Thr	Cys	
		770					775					780					
25	Gln	Trp	Asp	Leu	Ser	Trp	Ser	Ser	Asp	Pro	Pro	Phe	Cys	Glu	Lys	Ile	
	785					790					795					800	
30	Met	Tyr	Cys	Thr	Asp	Pro	Gly	Glu	Val	Asp	His	Ser	Thr	Arg	Leu	Ile	
					805					810					815		
35	Ser	Asp	Pro	Val	Leu	Leu	Val	Gly	Thr	Thr	Ile	Gln	Tyr	Thr	Cys	Asn	
				820					825					830			
40	Pro	Gly	Phe	Val	Leu	Glu	Gly	Ser	Ser	Leu	Leu	Thr	Cys	Tyr	Ser	Arg	
			835					840					845				
45	Glu	Thr	Gly	Thr	Pro	Ile	Trp	Thr	Ser	Arg	Leu	Pro	His	Cys	Val	Ser	
		850					855					860					
50	Glu	Glu	Ser	Leu	Ala	Cys	Asp	Asn	Pro	Gly	Leu	Pro	Glu	Asn	Gly	Tyr	
	865					870					875					880	
55	Gln	Ile	Leu	Tyr	Lys	Arg	Leu	Tyr	Leu	Pro	Gly	Glu	Ser	Leu	Thr	Phe	
					885					890					895		
60	Met	Cys	Tyr	Glu	Gly	Phe	Glu	Leu	Met	Gly	Glu	Val	Thr	Ile	Arg	Cys	
				900					905					910			
65	Ile	Leu	Gly	Gln	Pro	Ser	His	Trp	Asn	Gly	Pro	Leu	Pro	Val	Cys	Lys	
			915					920					925				
70	Val	Asn	Gln	Asp	Ser	Phe	Glu	His	Ala	Leu	Glu	Val	Ala	Glu	Ala	Ala	
		930					935					940					
75	Ala	Glu	Thr	Ser	Leu	Glu	Gly	Gly	Leu	Ala	Gly	His	His	His	His	His	

945

950

955

960

His His His His

5

<210> 18

<211> 2487

<212> DNA

10

<213> Homo sapiens

<220>

<221> misc_feature

15

<223> cDNA sequence of human SEZ6L2 ECD

<400> 18

ctgcctctca aagaggaaga gattctcccc gagcccgat ccgagacacc cacagtggct 60

tccgaagccc tcgctgaact gctgcacgga gccctcctga gaaggggacc tgaatgggc 120

20

tatctccctg gctccgacag agatcccaca ctgccacac ctctctgctgg acagaccctc 180

gctgtgcctt ccctgccag agccacagaa cccggaacag gccctctcac aacagctgtg 240

25

accctaacg gcgtcagagg agctggacct acagcccctg agctgctgac acctcctcct 300

ggcacaaccg ctctcctcc tccttcccct gctagccctg gacccctct cggacctgaa 360

ggaggcgagg aggagacaac caccaccatt attaccacca ccaccgtgac aaccacagtg 420

30

accagccctg tcctgtgcaa caacaacatc agcgaaggcg aaggctatgt ggaatcccct 480

gacctgggct cccctgtgtc cagaacactc ggccctcctgg attgcacata ctccattcac 540

gtgtaccccg gctacggaat cgagattcag gtgcagaccc tgaatctgtc ccaggaggag 600

35

gaactgctgg tgctggctgg cggaggaagc cctggcctcg ctcttagact cctcgctaac 660

tcctccatgc tcggcgaagg ccaggtcctc agatccccta ccaacaggct gtcctgcac 720

ttccagagcc ccagagtgcc tagaggaggc ggcttcagga ttactacca ggcctatctc 780

40

ctgagctgtg gattccctcc cagaccgct catggcgatg tctccgtcac cgacctccac 840

cccggaggaa cagccacctt ccactgtgat tccggatacc agctgcaagg cgaggagacc 900

ctgatttgcc tcaatggcac caggcccagc tggaacggag agacacctag ctgcatggct 960

45

agctgcggcg gaaccatcca taatgccacc ctccggcagga tcgtcagccc tgaacctggc 1020

ggagctgtgg gacctaacct cacatgcaga tgggtgatcg aagctgctga aggcaggaga 1080

ctccacctcc acttcgagag ggtgtccctg gacgaggaca acgacaggct catggtcaga 1140

50

agcggcggaa gccctctcag ccctgtgatt tacgacagcg acatggacga tgtgcctgag 1200

aggggcctca tctccgatgc ccaaagcctg tacgtggaac tcctctccga gaccccgct 1260

55

aacccctcc tcctgagcct cagattcgag gccttcgagg aggacagatg tttcgctcct 1320

tttctggccc atggcaacgt gaccacaacc gaccccgagt acagaccgg agctctggct 1380

EP 3 338 793 A1

accttcagct gtctgcctgg ctacgccctc gaacctcccg gacctcctaa tgccatcgaa 1440
 tgtgtggatc ccaccgaacc ccattggaac gacaccgagc ccgcttgtaa ggctatgtgc 1500
 5 ggcgggagaac tcagcgaacc tgccggagtg gtcctctccc ctgattggcc ccagagctat 1560
 tcccccgac aagactgtgt ctggggcgtg cacgtccagg aggaaaagag gatcctctc 1620
 caggtggaga ttctgaacgt cagagaggga gacatgctga ccctgttcga cggagacgga 1680
 10 ccttccgcca gagtcctcgc tcagctgaga ggccctcagc ccagaaggag actgctcagc 1740
 tccggccccg atctgacact ccagtttcag gccccccctg gccccccctaa tcctggcctg 1800
 ggacagggct tcgtgctcca cttcaaggag gtccccagga atgatacatg ccccgaaactg 1860
 15 cctcctcccg agtggggatg gaggacagct tcccatggcg acctgatcag gggaaccgtg 1920
 ctgacatatc agtgtgaacc cggctacgag ctgctgggaa gcgatatcct gacctgtcag 1980
 tgggatctct cctggagcgc tgctccccct gcctgtcaga aaatcatgac ctgcgctgac 2040
 20 cctggagaga tcgctaacgg ccacaggacc gcttccgacg ctggatttcc cgtgggctcc 2100
 cacgtgcaat acaggtgcct ccccgatac tccctcgaag gcgctgccat gctgacatgc 2160
 25 tacagcaggg acaccggcac acccaagtgg tccgacaggg tgcccaaagtg tgctctgaag 2220
 tacgagccct gtctcaatcc cggagtgccg gagaacggat accagaccct gtacaagcac 2280
 cactatcagg ccggcgcaatc cctgagattc ttctgctacg agggcttcga gctcatcggc 2340
 30 gaggtgacaa ttacctgtgt gcccggccat ccttcccagt ggaccagcca gccccctctc 2400
 tgtaaggtcg cctacgaaga gctgctcgac aataggaagc tggaggtcac ccagaccacc 2460
 35 gacccttcca gacaactgga aggcggc 2487

<210> 19
 <211> 865
 <212> PRT
 40 <213> Homo sapiens

<220>
 <221> MISC_FEATURE
 45 <223> Human SEZ6L2 ECD protein

<400> 19

Met Glu Thr Asp Thr Leu Leu Leu Trp Val Leu Leu Leu Trp Val Pro
 1 5 10 15

Gly Ser Thr Gly Asp Gly Ala Pro Leu Pro Leu Lys Glu Glu Glu Ile
 20 25 30

Leu Pro Glu Pro Gly Ser Glu Thr Pro Thr Val Ala Ser Glu Ala Leu
 35 40 45

EP 3 338 793 A1

	Ala	Glu	Leu	Leu	His	Gly	Ala	Leu	Leu	Arg	Arg	Gly	Pro	Glu	Met	Gly	
	50						55					60					
5	Tyr	Leu	Pro	Gly	Ser	Asp	Arg	Asp	Pro	Thr	Leu	Ala	Thr	Pro	Pro	Ala	
	65					70					75					80	
	Gly	Gln	Thr	Leu	Ala	Val	Pro	Ser	Leu	Pro	Arg	Ala	Thr	Glu	Pro	Gly	
10					85					90					95		
	Thr	Gly	Pro	Leu	Thr	Thr	Ala	Val	Thr	Pro	Asn	Gly	Val	Arg	Gly	Ala	
				100					105					110			
15	Gly	Pro	Thr	Ala	Pro	Glu	Leu	Leu	Thr	Pro	Pro	Pro	Gly	Thr	Thr	Ala	
				115				120					125				
	Pro	Pro	Pro	Pro	Ser	Pro	Ala	Ser	Pro	Gly	Pro	Pro	Leu	Gly	Pro	Glu	
20					130			135					140				
	Gly	Gly	Glu	Glu	Glu	Thr	Thr	Thr	Thr	Ile	Ile	Thr	Thr	Thr	Thr	Val	
25					145			150				155				160	
	Thr	Thr	Thr	Val	Thr	Ser	Pro	Val	Leu	Cys	Asn	Asn	Asn	Ile	Ser	Glu	
				165					170					175			
30	Gly	Glu	Gly	Tyr	Val	Glu	Ser	Pro	Asp	Leu	Gly	Ser	Pro	Val	Ser	Arg	
				180					185					190			
	Thr	Leu	Gly	Leu	Leu	Asp	Cys	Thr	Tyr	Ser	Ile	His	Val	Tyr	Pro	Gly	
35				195				200					205				
	Tyr	Gly	Ile	Glu	Ile	Gln	Val	Gln	Thr	Leu	Asn	Leu	Ser	Gln	Glu	Glu	
				210			215					220					
40	Glu	Leu	Leu	Val	Leu	Ala	Gly	Gly	Gly	Ser	Pro	Gly	Leu	Ala	Pro	Arg	
				225			230				235					240	
	Leu	Leu	Ala	Asn	Ser	Ser	Met	Leu	Gly	Glu	Gly	Gln	Val	Leu	Arg	Ser	
45				245					250					255			
	Pro	Thr	Asn	Arg	Leu	Leu	Leu	His	Phe	Gln	Ser	Pro	Arg	Val	Pro	Arg	
50				260				265					270				
	Gly	Gly	Gly	Phe	Arg	Ile	His	Tyr	Gln	Ala	Tyr	Leu	Leu	Ser	Cys	Gly	
				275			280					285					
55	Phe	Pro	Pro	Arg	Pro	Ala	His	Gly	Asp	Val	Ser	Val	Thr	Asp	Leu	His	
				290			295				300						

EP 3 338 793 A1

	Pro	Gly	Gly	Thr	Ala	Thr	Phe	His	Cys	Asp	Ser	Gly	Tyr	Gln	Leu	Gln	305	310	315	320
5	Gly	Glu	Glu	Thr	Leu	Ile	Cys	Leu	Asn	Gly	Thr	Arg	Pro	Ser	Trp	Asn	325	330	335	
10	Gly	Glu	Thr	Pro	Ser	Cys	Met	Ala	Ser	Cys	Gly	Gly	Thr	Ile	His	Asn	340	345	350	
15	Ala	Thr	Leu	Gly	Arg	Ile	Val	Ser	Pro	Glu	Pro	Gly	Gly	Ala	Val	Gly	355	360	365	
20	Pro	Asn	Leu	Thr	Cys	Arg	Trp	Val	Ile	Glu	Ala	Ala	Glu	Gly	Arg	Arg	370	375	380	
25	Leu	His	Leu	His	Phe	Glu	Arg	Val	Ser	Leu	Asp	Glu	Asp	Asn	Asp	Arg	385	390	395	400
	Leu	Met	Val	Arg	Ser	Gly	Gly	Ser	Pro	Leu	Ser	Pro	Val	Ile	Tyr	Asp	405	410	415	
30	Ser	Asp	Met	Asp	Asp	Val	Pro	Glu	Arg	Gly	Leu	Ile	Ser	Asp	Ala	Gln	420	425	430	
35	Ser	Leu	Tyr	Val	Glu	Leu	Leu	Ser	Glu	Thr	Pro	Ala	Asn	Pro	Leu	Leu	435	440	445	
40	Leu	Ser	Leu	Arg	Phe	Glu	Ala	Phe	Glu	Glu	Asp	Arg	Cys	Phe	Ala	Pro	450	455	460	
	Phe	Leu	Ala	His	Gly	Asn	Val	Thr	Thr	Thr	Asp	Pro	Glu	Tyr	Arg	Pro	465	470	475	480
45	Gly	Ala	Leu	Ala	Thr	Phe	Ser	Cys	Leu	Pro	Gly	Tyr	Ala	Leu	Glu	Pro	485	490	495	
50	Pro	Gly	Pro	Pro	Asn	Ala	Ile	Glu	Cys	Val	Asp	Pro	Thr	Glu	Pro	His	500	505	510	
	Trp	Asn	Asp	Thr	Glu	Pro	Ala	Cys	Lys	Ala	Met	Cys	Gly	Gly	Glu	Leu	515	520	525	
55	Ser	Glu	Pro	Ala	Gly	Val	Val	Leu	Ser	Pro	Asp	Trp	Pro	Gln	Ser	Tyr	530	535	540	
	Ser	Pro	Gly	Gln	Asp	Cys	Val	Trp	Gly	Val	His	Val	Gln	Glu	Glu	Lys				

EP 3 338 793 A1

	545		550		555		560									
5	Arg	Ile	Leu	Leu	Gln	Val	Glu	Ile	Leu	Asn	Val	Arg	Glu	Gly	Asp	Met
					565					570					575	
10	Leu	Thr	Leu	Phe	Asp	Gly	Asp	Gly	Pro	Ser	Ala	Arg	Val	Leu	Ala	Gln
				580					585					590		
15	Leu	Arg	Gly	Pro	Gln	Pro	Arg	Arg	Arg	Leu	Leu	Ser	Ser	Gly	Pro	Asp
			595					600					605			
20	Leu	Thr	Leu	Gln	Phe	Gln	Ala	Pro	Pro	Gly	Pro	Pro	Asn	Pro	Gly	Leu
		610					615					620				
25	Gly	Gln	Gly	Phe	Val	Leu	His	Phe	Lys	Glu	Val	Pro	Arg	Asn	Asp	Thr
	625					630					635					640
30	Cys	Pro	Glu	Leu	Pro	Pro	Pro	Glu	Trp	Gly	Trp	Arg	Thr	Ala	Ser	His
					645					650					655	
35	Gly	Asp	Leu	Ile	Arg	Gly	Thr	Val	Leu	Thr	Tyr	Gln	Cys	Glu	Pro	Gly
				660					665					670		
40	Tyr	Glu	Leu	Leu	Gly	Ser	Asp	Ile	Leu	Thr	Cys	Gln	Trp	Asp	Leu	Ser
			675					680					685			
45	Trp	Ser	Ala	Ala	Pro	Pro	Ala	Cys	Gln	Lys	Ile	Met	Thr	Cys	Ala	Asp
		690					695					700				
50	Pro	Gly	Glu	Ile	Ala	Asn	Gly	His	Arg	Thr	Ala	Ser	Asp	Ala	Gly	Phe
	705					710					715					720
55	Pro	Val	Gly	Ser	His	Val	Gln	Tyr	Arg	Cys	Leu	Pro	Gly	Tyr	Ser	Leu
					725					730					735	
60	Glu	Gly	Ala	Ala	Met	Leu	Thr	Cys	Tyr	Ser	Arg	Asp	Thr	Gly	Thr	Pro
				740					745					750		
65	Lys	Trp	Ser	Asp	Arg	Val	Pro	Lys	Cys	Ala	Leu	Lys	Tyr	Glu	Pro	Cys
			755					760					765			
70	Leu	Asn	Pro	Gly	Val	Pro	Glu	Asn	Gly	Tyr	Gln	Thr	Leu	Tyr	Lys	His
		770					775					780				
75	His	Tyr	Gln	Ala	Gly	Glu	Ser	Leu	Arg	Phe	Phe	Cys	Tyr	Glu	Gly	Phe
	785					790					795					800

EP 3 338 793 A1

	Glu	Leu	Ile	Gly	Glu	Val	Thr	Ile	Thr	Cys	Val	Pro	Gly	His	Pro	Ser	
					805					810					815		
5	Gln	Trp	Thr	Ser	Gln	Pro	Pro	Leu	Cys	Lys	Val	Ala	Tyr	Glu	Glu	Leu	
				820					825					830			
	Leu	Asp	Asn	Arg	Lys	Leu	Glu	Val	Thr	Gln	Thr	Thr	Asp	Pro	Ser	Arg	
10			835					840					845				
	Gln	Leu	Glu	Gly	Gly	Leu	Ala	Gly	His	His	His	His	His	His	His	His	
		850					855					860					
15	His																
	865																
	<210>	20															
20	<211>	107															
	<212>	PRT															
	<213>	Mus musculus															
	<220>																
25	<221>	MISC_FEATURE															
	<223>	Murine SC17.1 VL															
	<400>	20															
30	Gln	Ile	Val	Leu	Thr	Gln	Ser	Pro	Ala	Leu	Met	Ser	Ala	Ser	Pro	Gly	
	1				5					10					15		
	Glu	Lys	Val	Ser	Leu	Thr	Cys	Ser	Ala	Asn	Ser	Thr	Val	Ser	Phe	Met	
35			20						25					30			
	Tyr	Trp	Tyr	Gln	Gln	Lys	Pro	Arg	Ser	Ser	Pro	Thr	Pro	Trp	Ile	Tyr	
			35					40					45				
40	Leu	Thr	Ser	Asn	Leu	Ala	Ser	Gly	Val	Pro	Ala	Arg	Phe	Ser	Gly	Ser	
		50					55					60					
	Gly	Ser	Gly	Thr	Ser	Tyr	Ser	Leu	Thr	Ile	Ser	Ser	Met	Glu	Ala	Glu	
45		65				70					75					80	
	Asp	Ala	Ala	Thr	Tyr	Tyr	Cys	Gln	Gln	Trp	Ser	Ser	Asn	Ser	Pro	Ile	
				85						90					95		
50	Thr	Phe	Gly	Ala	Gly	Thr	Lys	Leu	Glu	Leu	Lys						
				100					105								
	<210>	21															
55	<211>	117															
	<212>	PRT															

EP 3 338 793 A1

<213> Mus musculus

<220>

<221> MISC_FEATURE

<223> Murine SC17.1 VH

<400> 21

Asp Val Gln Leu Gln Asp Ser Gly Pro Gly Leu Val Lys Pro Ser Gln
1 5 10 15

Ser Leu Ser Val Thr Cys Thr Val Thr Gly Tyr Ser Ile Thr Trp Gly
20 25 30

Tyr Tyr Trp Asn Trp Ile Arg Gln Phe Pro Gly Asn Lys Leu Glu Trp
35 40 45

Met Gly Asn Ile His Asn Ser Gly Gly Thr Asn Tyr Asn Pro Ser Leu
50 55 60

Lys Ser Arg Ile Ser Ile Thr Arg Asp Thr Ser Lys Asn Gln Phe Phe
65 70 75 80

Leu Gln Leu Asn Ser Val Thr Thr Glu Asp Thr Ala Thr Tyr Tyr Cys
85 90 95

Ala Thr Thr Asn Trp Asp Tyr Phe Asp Tyr Trp Gly Gln Gly Thr Thr
100 105 110

Leu Thr Val Ser Ser
115

<210> 22

<211> 112

<212> PRT

<213> Mus musculus

<220>

<221> MISC_FEATURE

<223> Murine SC17.2 VL

<400> 22

Asp Ile Val Met Ser Gln Ser Pro Ser Ser Leu Ala Val Ser Val Gly
1 5 10 15

Glu Lys Val Thr Met Ser Cys Lys Ser Ser Gln Ser Leu Leu Tyr Ser
20 25 30

Ser Asn Gln Lys Ser Tyr Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln
35 40 45

EP 3 338 793 A1

Ser Pro Lys Leu Leu Ile Tyr Trp Ala Ser Thr Arg Glu Ser Gly Val
50 55 60

5 Pro Asp Arg Phe Thr Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr
65 70 75 80

10 Ile Ser Ser Val Gln Ala Glu Asp Leu Ala Val Tyr Tyr Cys Lys Gln
85 90 95

Ser Tyr Asn Leu Arg Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys
100 105 110

15

<210> 23
<211> 117
<212> PRT
<213> Mus musculus

20

<220>
<221> MISC_FEATURE
<223> Murine SC17.2 VH

25

<400> 23

Gln Val Gln Leu Gln Gln Ser Asp Ala Glu Leu Val Lys Pro Gly Ala
1 5 10 15

30

Ser Val Lys Ile Ser Cys Lys Val Ser Gly Tyr Thr Phe Thr Asp His
20 25 30

35

Thr Ile His Trp Met Lys Gln Arg Pro Glu Gln Gly Leu Glu Trp Ile
35 40 45

Gly Tyr Ile Tyr Pro Arg Asp Gly Ser Thr Lys Tyr Asn Glu Glu Phe
50 55 60

40

Lys Gly Lys Ala Thr Leu Thr Ala Asp Lys Ser Ser Ser Thr Ala Tyr
65 70 75 80

45

Met Gln Leu Asn Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Phe Cys
85 90 95

50

Ala Arg Ser Tyr Ser Asn Tyr Phe Asp Tyr Trp Gly Gln Gly Thr Thr
100 105 110

Leu Thr Val Ser Ser
115

55

<210> 24

EP 3 338 793 A1

<211> 111
<212> PRT
<213> Mus musculus

<220>
<221> MISC_FEATURE
<223> Murine SC17.9 VL

<400> 24

Asp Ile Val Met Ser Gln Ser Pro Ser Ser Leu Ala Val Ser Val Gly
1 5 10 15

Glu Lys Val Thr Met Ser Cys Lys Ser Ser Gln Ser Leu Leu Tyr Ser
20 25 30

Ser Asn Gln Lys Asn Tyr Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln
35 40 45

Ser Pro Lys Leu Leu Ile Tyr Trp Ala Ser Thr Arg Glu Ser Gly Val
50 55 60

Pro Asp Arg Phe Thr Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr
65 70 75 80

Ile Ser Ser Val Lys Ala Glu Asp Leu Ala Val Tyr Phe Cys Gln Gln
85 90 95

Tyr Tyr Asn Tyr Pro Tyr Thr Phe Gly Gly Gly Thr Lys Leu Lys
100 105 110

<210> 25
<211> 119
<212> PRT
<213> Mus musculus

<220>
<221> MISC_FEATURE
<223> Murine SC17.9 VH

<400> 25

Gln Val Gln Leu Gln Gln Pro Gly Ala Glu Ile Val Arg Pro Gly Ala
1 5 10 15

Ser Val Lys Leu Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asp Tyr
20 25 30

Trp Met Asn Trp Val Lys Gln Arg Pro Gly Gln Gly Leu Glu Trp Ile
35 40 45

EP 3 338 793 A1

Gly Ala Ile Asp Pro Ser Asp Ser Tyr Thr Ser Tyr Asn Pro Lys Phe
50 55 60

5 Lys Gly Lys Ala Thr Leu Thr Val Asp Thr Ser Ser Ser Ser Ala Tyr
65 70 75 80

10 Met Gln Leu Ser Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Phe Cys
85 90 95

Ala Arg Arg Gly Thr Pro Gly Lys Pro Leu Val Tyr Trp Gly Gln Gly
100 105 110

15 Thr Leu Val Thr Val Ser Ala
115

20 <210> 26
<211> 107
<212> PRT
<213> Mus musculus

25 <220>
<221> MISC_FEATURE
<223> Murine SC17.16 VL

<400> 26

30 Asp Ile Gln Met Thr Gln Ser Pro Ala Ser Leu Ser Val Ser Val Gly
1 5 10 15

35 Glu Thr Val Thr Ile Thr Cys Arg Ala Ser Ala Asn Ile Asn Ser Asn
20 25 30

Leu Val Trp Tyr Gln Gln Lys Gln Gly Lys Ser Pro Gln Leu Leu Val
35 40 45

40 Tyr Ala Ala Thr Asn Leu Ala Asp Gly Val Pro Ser Arg Phe Ser Gly
50 55 60

45 Ser Gly Ser Gly Thr Gln Tyr Ser Leu Lys Ile Asn Ser Leu Gln Ser
65 70 75 80

Glu Asp Phe Gly Asn Tyr Tyr Cys Gln His Phe Trp Gly Thr Pro Arg
85 90 95

50 Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys
100 105

55 <210> 27
<211> 116
<212> PRT

EP 3 338 793 A1

<213> Mus musculus

<220>

<221> MISC_FEATURE

<223> Murine SC17.16 VH

<400> 27

Glu Val Gln Leu Gln Gln Ser Gly Pro Glu Leu Met Lys Pro Gly Ala
1 5 10 15

Ser Val Lys Met Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asp Tyr
20 25 30

Asn Met Tyr Trp Val Lys Gln Asn Gln Gly Lys Ser Leu Glu Trp Ile
35 40 45

Gly Glu Ile Asn Pro Asn Asn Gly Gly Thr Ala Tyr Asn Gln Lys Phe
50 55 60

Arg Gly Lys Ala Thr Leu Thr Val Asp Lys Ser Ser Ser Thr Ala Tyr
65 70 75 80

Met Glu Leu Arg Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys
85 90 95

Ala Arg Tyr Asp Lys Gly Phe Asp Tyr Trp Gly Gln Gly Thr Thr Leu
100 105 110

Thr Val Ser Ser
115

<210> 28

<211> 111

<212> PRT

<213> Mus musculus

<220>

<221> MISC_FEATURE

<223> Murine SC17.38 VL

<400> 28

Asp Ile Val Val Thr Gln Ser Pro Ala Ser Leu Ala Val Ser Leu Gly
1 5 10 15

Gln Arg Ala Thr Ile Ser Cys Arg Ala Ser Glu Ser Val Glu Tyr Tyr
20 25 30

Gly Thr Ser Leu Met Gln Trp Phe Gln Gln Lys Pro Gly Gln Pro Pro
35 40 45

EP 3 338 793 A1

Lys Leu Leu Ile Tyr Ala Ala Ser Asn Val Glu Ser Gly Val Pro Ala
50 55 60

5 Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Ser Leu Asn Ile His
65 70 75 80

10 Pro Val Glu Glu Asp Asp Ile Ala Met Tyr Phe Cys Gln Gln Asp Arg
85 90 95

Lys Val Pro Trp Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys
100 105 110

15

<210> 29
<211> 120
<212> PRT
<213> Mus musculus

20

<220>
<221> MISC_FEATURE
<223> Murine SC17.38 VH

25

<400> 29

Gln Val Thr Leu Lys Glu Ser Gly Pro Gly Ile Leu Gln Pro Ser Gln
1 5 10 15

30

Thr Leu Ser Leu Thr Cys Ser Phe Ser Gly Phe Ser Leu Asn Thr Ser
20 25 30

35

Gly Met Ser Val Gly Trp Val Arg Gln Pro Ser Gly Arg Gly Leu Glu
35 40 45

40

Trp Leu Ala Pro Ile Trp Trp Asn Gly Asp Lys Tyr Tyr Asn Pro Ala
50 55 60

Leu Lys Ser Arg Leu Thr Ile Ser Lys Asp Thr Ser Asn Asn Gln Val
65 70 75 80

45

Phe Leu Lys Ile Ala Ser Val Val Thr Ala Asp Thr Ala Thr Tyr Phe
85 90 95

50

Cys Ala Arg Ile Arg Gln Tyr Tyr Tyr Ala Met Asp Tyr Trp Gly Gln
100 105 110

Gly Thr Ser Val Thr Val Ser Ser
115 120

55

<210> 30

EP 3 338 793 A1

<211> 112
<212> PRT
<213> Mus musculus

<220>
<221> MISC_FEATURE
<223> Murine SC17.3 VL

<400> 30

Asp Ile Val Met Ser Gln Ser Pro Ser Ser Leu Ala Val Ser Val Gly
1 5 10 15

Glu Lys Val Thr Met Ser Cys Lys Ser Ser Gln Ser Leu Leu Tyr Ser
20 25 30

Ser Asn Gln Lys Asn Tyr Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln
35 40 45

Ser Pro Lys Leu Leu Ile Tyr Trp Ala Ser Thr Arg Glu Ser Gly Val
50 55 60

Pro Asp Arg Phe Thr Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr
65 70 75 80

Ile Ser Ser Val Lys Ala Glu Asp Leu Ala Val Tyr Tyr Cys Gln Gln
85 90 95

Tyr Tyr Ser Tyr Pro Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys
100 105 110

<210> 31
<211> 118
<212> PRT
<213> Mus musculus

<220>
<221> MISC_FEATURE
<223> Murine SC17.3 VH

<400> 31

Gln Val Gln Leu Gln Gln Pro Gly Ala Glu Leu Val Lys Pro Gly Ala
1 5 10 15

Ser Val Lys Leu Ser Cys Lys Ala Ser Gly Tyr Thr Phe Pro Ser Tyr
20 25 30

Trp Ile His Cys Val Lys Gln Arg Pro Gly Gln Gly Leu Glu Trp Ile
35 40 45

EP 3 338 793 A1

Gly Val Ile Asn Pro Ser Asn Gly Arg Thr Asn Tyr Asn Glu Lys Phe
50 55 60

5 Lys Asn Lys Ala Thr Leu Thr Val Asp Lys Ser Ser Ser Thr Ala Tyr
65 70 75 80

10 Met Gln Leu Ser Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys
85 90 95

Val Arg Gly Gly Thr Gly Tyr Thr Met Asp Tyr Trp Gly Gln Gly Thr
100 105 110

15 Ser Val Thr Val Ser Ser
115

20 <210> 32
<211> 107
<212> PRT
<213> Mus musculus

25 <220>
<221> MISC_FEATURE
<223> Murine SC17.4 VL

<400> 32

30 Asp Ile Lys Met Thr Gln Ser Pro Ser Ser Met Tyr Ala Ser Leu Gly
1 5 10 15

35 Glu Arg Val Thr Ile Thr Cys Lys Ala Ser Gln Asp Ile Asn Ser Tyr
20 25 30

Leu Thr Trp Phe Gln Gln Lys Pro Gly Lys Ser Pro Lys Thr Leu Ile
35 40 45

40 Tyr Arg Ala Asn Arg Leu Ile Asp Gly Val Pro Ser Arg Phe Ser Gly
50 55 60

45 Ser Gly Ser Gly Gln Asp Tyr Ser Leu Thr Ile Ser Ser Leu Asp Tyr
65 70 75 80

Glu Asp Met Gly Ile Tyr Tyr Cys Leu Gln Tyr Asp Asp Phe Pro Trp
85 90 95

50 Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys
100 105

55 <210> 33
<211> 116
<212> PRT

EP 3 338 793 A1

<213> Mus musculus

<220>

<221> MISC_FEATURE

<223> Murine SC17.4 VH

<400> 33

Gln Ile Gln Leu Val Gln Ser Gly Pro Glu Leu Lys Lys Pro Gly Glu
1 5 10 15

Thr Val Lys Ile Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asp Tyr
20 25 30

Ser Met His Trp Val Lys Gln Ala Pro Gly Lys Gly Leu Lys Trp Leu
35 40 45

Gly Trp Ile Asn Thr Glu Thr Gly Glu Pro Thr Tyr Ser Glu Asp Phe
50 55 60

Lys Gly Arg Phe Ala Phe Ser Leu Glu Thr Ser Ala Ser Thr Ala Tyr
65 70 75 80

Leu Gln Ile Asn Asn Leu Lys Asn Glu Asp Thr Ala Thr Tyr Phe Cys
85 90 95

Val Lys Asn Lys Gly Trp Phe Ala Tyr Trp Gly Gln Gly Thr Leu Val
100 105 110

Thr Val Ser Ala
115

<210> 34

<211> 112

<212> PRT

<213> Mus musculus

<220>

<221> MISC_FEATURE

<223> Murine SC17.8 VL

<400> 34

Asp Val Val Met Thr Gln Thr Pro Leu Ser Leu Pro Val Ser Leu Gly
1 5 10 15

Asp Gln Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Leu Val His Ser
20 25 30

Asn Gly Asp Thr Tyr Leu His Trp Tyr Leu Gln Lys Pro Gly Gln Ser
35 40 45

EP 3 338 793 A1

Pro Lys Leu Leu Ile Tyr Lys Val Ser Asn Arg Phe Ser Gly Val Pro
50 55 60

5 Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile
65 70 75 80

10 Ser Arg Val Glu Ala Glu Asp Leu Gly Leu Tyr Phe Cys Ser Gln Ser
85 90 95

Thr Leu Ile Pro Tyr Thr Phe Gly Gly Gly Thr Lys Leu Asp Ile Lys
100 105 110

15

<210> 35
<211> 114
<212> PRT
<213> Mus musculus

20

<220>
<221> MISC_FEATURE
<223> Murine SC17.8 VH

25

<400> 35

Gln Val His Leu Gln Gln Ser Gly Thr Glu Val Met Lys Pro Gly Ala
1 5 10 15

30

Ser Val Lys Ile Ser Cys Lys Ala Thr Gly Tyr Thr Phe Ser Ser Tyr
20 25 30

35

Trp Ile Glu Trp Ile Lys Gln Arg Pro Gly His Gly Leu Glu Trp Ile
35 40 45

Gly Glu Ile Leu Pro Gly Ser Gly Asn Thr Asn Asn Asn Glu Lys Phe
50 55 60

40

Lys Gly Lys Ala Thr Ile Thr Ala Asp Thr Ser Ser Asn Ile Ala Tyr
65 70 75 80

45

Ile Gln Leu Ser Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys
85 90 95

50

Ala Gly Gly Pro Ala Ala Tyr Trp Gly Gln Gly Thr Leu Val Thr Val
100 105 110

Ser Ala

55

<210> 36

EP 3 338 793 A1

<211> 113
<212> PRT
<213> Mus musculus

<220>
<221> MISC_FEATURE
<223> Murine SC17.10 VL

<400> 36

Asp Ile Val Met Ser Gln Ser Pro Ser Ser Leu Ala Val Ser Val Gly
1 5 10 15

Glu Lys Val Thr Met Ser Cys Lys Ser Ser Gln Ser Leu Leu Tyr Ser
20 25 30

Ser Asn Gln Lys Asn Tyr Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln
35 40 45

Ser Pro Lys Leu Leu Ile Tyr Trp Ala Ser Thr Arg Glu Ser Gly Val
50 55 60

Pro Asp Arg Phe Thr Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr
65 70 75 80

Ile Ser Ser Val Lys Thr Glu Asp Leu Ala Leu Tyr Tyr Cys Gln Gln
85 90 95

Tyr Tyr Trp Phe Pro Tyr Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile
100 105 110

Lys

<210> 37
<211> 113
<212> PRT
<213> Mus musculus

<220>
<221> MISC_FEATURE
<223> Murine SC17.10 VH

<400> 37

Glu Val Gln Leu Gln Gln Ser Gly Ala Glu Leu Val Lys Pro Gly Ala
1 5 10 15

Ser Val Lys Leu Ser Cys Thr Ala Ser Gly Phe Asn Ile Lys Asp Thr
20 25 30

EP 3 338 793 A1

	Tyr	Met	His	Trp	Val	Lys	Gln	Arg	Pro	Glu	Gln	Gly	Leu	Glu	Trp	Ile	
			35					40					45				
5	Gly	Arg	Ile	Asp	Pro	Ala	Asn	Val	Asn	Thr	Lys	Tyr	Asp	Pro	Lys	Phe	
		50					55					60					
	Gln	Gly	Lys	Ala	Thr	Ile	Thr	Ala	Asp	Thr	Ser	Ser	Asn	Thr	Ala	Tyr	
10	65					70					75					80	
	Leu	Gln	Leu	Ser	Ser	Leu	Thr	Ser	Glu	Asp	Thr	Ala	Val	Tyr	Tyr	Cys	
				85						90					95		
15	Val	Arg	Gly	Asn	Val	Tyr	Trp	Gly	Gln	Gly	Thr	Leu	Val	Thr	Val	Ser	
				100					105					110			
20	Ala																
	<210>	38															
	<211>	108															
25	<212>	PRT															
	<213>	Mus musculus															
	<220>																
	<221>	MISC_FEATURE															
30	<223>	Murine SC17.11 VL															
	<400>	38															
	Glu	Asn	Val	Leu	Thr	Gln	Ser	Pro	Ala	Ile	Met	Ser	Ala	Ser	Pro	Gly	
35	1				5					10					15		
	Glu	Lys	Val	Thr	Met	Thr	Cys	Arg	Ala	Ser	Ser	Ser	Val	Ser	Ser	Ser	
				20					25					30			
40	Tyr	Leu	His	Trp	Tyr	Gln	Gln	Lys	Ser	Gly	Ala	Ser	Pro	Lys	Leu	Trp	
			35					40					45				
	Ile	Tyr	Ser	Thr	Ser	Asn	Leu	Ala	Ser	Gly	Val	Pro	Ala	Arg	Phe	Ser	
45		50					55					60					
	Gly	Ser	Gly	Ser	Gly	Thr	Ser	Tyr	Ser	Leu	Thr	Ile	Ser	Ser	Val	Glu	
50	65					70					75					80	
	Ala	Glu	Asp	Ala	Ala	Thr	Tyr	Tyr	Cys	Gln	Gln	Tyr	Ser	Asp	Tyr	Pro	
				85						90					95		
55	Phe	Thr	Phe	Gly	Ser	Gly	Thr	Lys	Leu	Val	Ile	Lys					
				100					105								

EP 3 338 793 A1

<210> 39
 <211> 122
 <212> PRT
 <213> Mus musculus

5

<220>
 <221> MISC_FEATURE
 <223> Murine SC17.11 VH

10

<400> 39

Glu Val Gln Leu Gln Gln Ser Gly Pro Glu Leu Val Lys Pro Gly Ala
 1 5 10 15

15

Leu Val Met Met Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asp Tyr
 20 25 30

20

Tyr Met His Trp Val Lys Gln Ser His Gly Gln Ser Leu Glu Trp Ile
 35 40 45

25

Gly Glu Val Ile Pro Tyr Asn Asp Glu Thr Phe Tyr Asn Arg Lys Phe
 50 55 60

Lys Asp Lys Ala Thr Leu Thr Val Asp Lys Ser Ser Ser Thr Ala Tyr
 65 70 75 80

30

Met Glu Leu Arg Ser Leu Thr Ser Glu Asp Ser Ala Ile Tyr Tyr Cys
 85 90 95

35

Ala Arg Arg His Arg Tyr Asp Gly Phe Arg Tyr Ala Ile Asp Tyr Trp
 100 105 110

Gly Gln Gly Thr Ser Val Thr Val Ser Ser
 115 120

40

<210> 40
 <211> 112
 <212> PRT
 <213> Mus musculus

45

<220>
 <221> MISC_FEATURE
 <223> Murine SC17.14 VL

50

<400> 40

Asp Val Leu Met Thr Gln Thr Pro Leu Ser Leu Pro Val Ser Leu Gly
 1 5 10 15

55

Asp Gln Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Ile Val His Ser
 20 25 30

EP 3 338 793 A1

	Asn	Gly	Asn	Thr	Tyr	Leu	Glu	Trp	Phe	Leu	Gln	Lys	Pro	Gly	Gln	Ser	
			35					40					45				
5	Pro	Lys	Leu	Leu	Ile	Tyr	Lys	Val	Ser	Asn	Arg	Phe	Ser	Gly	Val	Pro	
		50					55					60					
10	Asp	Arg	Phe	Ser	Gly	Ser	Gly	Ser	Gly	Thr	Asp	Phe	Thr	Leu	Lys	Ile	
	65					70					75					80	
15	Ser	Arg	Val	Glu	Ala	Glu	Asp	Leu	Gly	Val	Tyr	Tyr	Cys	Phe	Gln	Gly	
				85						90					95		
20	Ser	His	Val	Pro	Tyr	Thr	Phe	Gly	Gly	Gly	Thr	Lys	Leu	Glu	Ile	Lys	
				100					105						110		
	<210>	41															
	<211>	118															
	<212>	PRT															
	<213>	Mus musculus															
25	<220>																
	<221>	MISC_FEATURE															
	<223>	Murine SC17.14 VH															
30	<400>	41															
	Glu	Val	Gln	Leu	Gln	Gln	Ser	Gly	Pro	Val	Leu	Val	Lys	Pro	Gly	Ala	
	1				5					10					15		
35	Ser	Val	Lys	Met	Ser	Cys	Lys	Ala	Ser	Gly	Tyr	Thr	Ile	Thr	Asp	Tyr	
				20					25						30		
40	Asn	Met	Asn	Trp	Val	Lys	Gln	Ser	His	Gly	Lys	Ser	Leu	Glu	Trp	Ile	
			35					40						45			
45	Gly	Val	Ile	Asn	Pro	Tyr	Asn	Gly	Asn	Thr	Arg	Tyr	Asn	Gln	Met	Phe	
		50					55					60					
50	Lys	Gly	Lys	Ala	Thr	Leu	Thr	Val	Asp	Lys	Ser	Ser	Ser	Thr	Ala	Tyr	
	65					70					75					80	
55	Met	Glu	Leu	Asn	Ser	Leu	Thr	Ser	Glu	Asp	Ser	Ala	Val	Tyr	Tyr	Cys	
				85						90					95		
	Thr	Arg	Trp	Gly	Thr	Thr	Val	Val	Gly	Ala	Asn	Trp	Gly	Gln	Gly	Thr	
				100					105					110			
	Thr	Leu	Thr	Val	Ser	Ser											

EP 3 338 793 A1

115

5 <210> 42
 <211> 106
 <212> PRT
 <213> Mus musculus

10 <220>
 <221> MISC_FEATURE
 <223> Murine SC17.15 VL

<400> 42

15 Gln Ile Val Leu Thr Gln Ser Pro Ala Leu Met Ser Ala Ser Pro Gly
 1 5 10 15

20 Glu Lys Val Thr Met Thr Cys Ser Ala Ser Ser Ser Val Asn Tyr Met
 20 25 30

Tyr Trp Tyr Gln Gln Lys Pro Arg Ser Ser Pro Lys Pro Trp Ile Tyr
 35 40 45

25 Leu Thr Ser Asn Leu Ala Ser Gly Val Pro Val Arg Phe Ser Gly Ser
 50 55 60

30 Gly Ser Gly Thr Ser Tyr Ser Leu Thr Ile Ser Ser Met Glu Ala Glu
 65 70 75 80

35 Asp Ala Ala Thr Tyr Tyr Cys Gln Gln Trp Ser Asn Asn Pro Pro Thr
 85 90 95

Phe Gly Ser Gly Thr Lys Leu Glu Leu Lys
 100 105

40 <210> 43
 <211> 122
 <212> PRT
 <213> Mus musculus

45 <220>
 <221> MISC_FEATURE
 <223> Murine SC17.15 VH

50 <400> 43

Asp Val Lys Leu Val Glu Ser Gly Gly Gly Leu Val Lys Leu Gly Gly
 1 5 10 15

55 Ser Leu Lys Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
 20 25 30

EP 3 338 793 A1

	Ala Met Ser Trp Val Arg Gln Thr Pro Glu Lys Arg Leu Glu Trp Val	
	35	40 45
5	Ala Thr Ile Thr Ser Gly Gly Gly Asn Thr Tyr Tyr Pro Asp Ser Val	
	50	55 60
10	Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Thr Leu Tyr	
	65	70 75 80
15	Leu Gln Met Ser Ser Leu Lys Ser Glu Asp Thr Ala Met Tyr Tyr Cys	
		85 90 95
20	Ala Arg Arg Asp Tyr Tyr Gly Ser Ser Tyr Val Met Phe Ala Tyr Trp	
		100 105 110
25	Gly Gln Gly Thr Leu Val Thr Val Ser Ala	
		115 120
30	<210> 44	
	<211> 106	
	<212> PRT	
	<213> Mus musculus	
35	<220>	
	<221> MISC_FEATURE	
	<223> Murine SC17.17 VL	
40	<400> 44	
45	Gln Ile Val Leu Thr Gln Ser Pro Ala Ile Met Ser Ala Ser Pro Gly	
	1	5 10 15
50	Glu Lys Val Thr Met Thr Cys Ser Ala Ser Ser Ser Val Ser Tyr Met	
		20 25 30
55	His Trp Tyr Gln Gln Lys Ser Gly Thr Ser Pro Lys Arg Trp Ile Tyr	
		35 40 45
60	Asp Thr Ser Lys Leu Pro Ser Gly Val Pro Ala Arg Phe Ser Gly Ser	
		50 55 60
65	Gly Ser Gly Thr Ser Tyr Ser Leu Thr Ile Ser Ser Met Glu Ala Glu	
	65	70 75 80
70	Asp Ala Ala Thr Tyr Tyr Cys Gln Gln Trp Ser Ser Thr Pro Pro Thr	
		85 90 95
75	Phe Gly Ala Gly Thr Lys Leu Glu Leu Lys	
		100 105

EP 3 338 793 A1

<210> 45
 <211> 120
 <212> PRT
 <213> Mus musculus

5

<220>
 <221> MISC_FEATURE
 <223> Murine SC17.17 VH

10

<400> 45

Glu Val Gln Leu Gln Gln Ser Gly Pro Glu Val Met Lys Pro Gly Ala
 1 5 10 15

15

Ser Val Lys Met Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asp Tyr
 20 25 30

20

Asn Met His Trp Val Lys Gln Asn Gln Gly Lys Ser Leu Glu Trp Ile
 35 40 45

25

Gly Glu Ile Asn Pro Asn Ile Gly Gly Thr Gly Tyr Asn Gln Lys Phe
 50 55 60

Lys Gly Lys Ala Thr Leu Thr Val His Lys Ser Ser Ser Thr Ala Tyr
 65 70 75 80

30

Met Glu Leu Arg Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys
 85 90 95

35

Ala Arg Thr Tyr Ser Tyr Tyr Ser Tyr Glu Phe Ala Tyr Trp Gly Gln
 100 105 110

Gly Thr Leu Val Thr Val Ser Ala
 115 120

40

<210> 46
 <211> 107
 <212> PRT
 <213> Mus musculus

45

<220>
 <221> MISC_FEATURE
 <223> Murine SC17.18 VL

50

<400> 46

Asp Ile Gln Met Thr Gln Ser Ser Ser Tyr Leu Ser Val Ser Leu Gly
 1 5 10 15

55

Gly Arg Val Thr Ile Thr Cys Lys Ala Ser Asp His Ile Asn Asn Trp
 20 25 30

EP 3 338 793 A1

Leu Ala Trp Tyr Gln Gln Lys Pro Gly Asn Ala Pro Arg Leu Leu Ile
35 40 45

5 Ser Gly Ala Thr Ser Leu Glu Thr Gly Val Pro Ser Arg Phe Ser Gly
50 55 60

10 Ser Gly Ser Gly Lys Asp Tyr Thr Leu Ser Ile Thr Ser Leu Gln Thr
65 70 75 80

Glu Asp Val Ala Thr Tyr Tyr Cys Gln Gln Tyr Trp Ser Ile Pro Leu
85 90 95

15 Thr Phe Gly Ala Gly Thr Lys Leu Glu Leu Lys
100 105

20 <210> 47
<211> 123
<212> PRT
<213> Mus musculus

25 <220>
<221> MISC_FEATURE
<223> Murine SC17.18 VH

<400> 47

30 Gln Val Thr Leu Lys Glu Ser Gly Pro Gly Ile Leu Gln Pro Ser Gln
1 5 10 15

35 Thr Leu Ser Leu Thr Cys Ser Phe Ser Gly Phe Ser Leu Ser Thr Ser
20 25 30

40 Thr Met Gly Val Gly Trp Ile Arg Gln Pro Ser Gly Lys Gly Leu Glu
35 40 45

Trp Leu Ala Asp Ile Trp Trp Asp Asp Ser Lys Tyr Tyr Asn Pro Ser
50 55 60

45 Leu Lys Ser Arg Leu Thr Ile Ser Lys Asp Thr Ser Ser Asn Gln Val
65 70 75 80

50 Phe Leu Lys Ile Thr Ser Val Asp Thr Ala Asp Thr Ala Thr Tyr Tyr
85 90 95

Cys Ala Arg Lys Gly Arg Thr Ala Arg Ala Thr Arg Gly Phe Ala Tyr
100 105 110

55 Trp Gly His Gly Thr Leu Val Thr Val Ser Ala

EP 3 338 793 A1

115

120

5 <210> 48
 <211> 109
 <212> PRT
 <213> Mus musculus

10 <220>
 <221> MISC_FEATURE
 <223> Murine SC17.19 VL

<400> 48

15 Asp Ile Val Leu Thr Gln Ser Pro Ala Ser Leu Ala Val Ser Leu Gly
 1 5 10 15

Gln Arg Ala Ala Ile Ser Cys Lys Pro Ser Gln Ser Val Asp Tyr Asp
 20 20 25 30

Gly Asp Ser Tyr Met Asn Trp Tyr Gln Gln Lys Pro Gly Gln Pro Pro
 35 40 45

25 Lys Leu Leu Ile Tyr Ala Ala Ser Asn Leu Glu Ser Gly Ile Pro Ala
 50 55 60

30 Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Asn Ile His
 65 70 75 80

Pro Val Glu Glu Glu Asp Ala Ala Thr Tyr Tyr Cys His Gln Ile Asn
 35 85 90 95

Asp Asp Pro Trp Thr Phe Gly Gly Gly Thr Lys Leu Lys
 100 105

40 <210> 49
 <211> 121
 <212> PRT
 <213> Mus musculus

45 <220>
 <221> MISC_FEATURE
 <223> Murine SC17.19 VH

50 <400> 49

Asp Val Gln Leu Gln Glu Ser Gly Pro Gly Leu Val Lys Pro Ser Gln
 1 5 10 15

55 Ser Leu Ser Val Thr Cys Thr Val Thr Gly Tyr Ser Ile Thr Ser Ser
 20 25 30

EP 3 338 793 A1

Tyr Thr Trp Asn Trp Ile Arg Gln Phe Pro Gly Asn Lys Leu Glu Trp
 35 40 45
 5 Met Gly Tyr Ile His Tyr Ser Gly Ser Thr Asn Tyr Asn Pro Ser Leu
 50 55 60
 10 Arg Ser Arg Ile Ser Ile Thr Arg Asp Thr Ser Lys Asn Gln Phe Phe
 65 70 75 80
 15 Leu Gln Leu Asn Ser Val Thr Thr Glu Asp Thr Ala Thr Tyr Tyr Cys
 85 90 95
 20 Ala Arg Ser Arg Tyr Tyr Tyr Asp Ala Tyr Gly Phe Ala Tyr Trp Gly
 100 105 110
 25 Gln Gly Thr Leu Val Thr Val Ser Ala
 115 120
 30 <210> 50
 <211> 112
 <212> PRT
 <213> Mus musculus
 35 Asp Val Val Leu Thr Gln Thr Pro Leu Ser Leu Pro Val Ser Leu Gly
 1 5 10 15
 40 Asp Gln Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Ile Val His Ile
 20 25 30
 45 Asn Arg His Thr Tyr Leu Gly Trp Tyr Leu Gln Lys Pro Gly Gln Ser
 35 40 45
 50 Leu Lys Leu Leu Ile Tyr Gly Val Ser Asn Arg Phe Ser Gly Val Pro
 50 55 60
 55 Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile
 65 70 75 80
 Ser Arg Val Glu Ala Glu Asp Met Gly Val Tyr Tyr Cys Phe Gln Gly
 85 90 95
 60 Thr His Val Pro Phe Thr Phe Gly Ser Gly Thr Lys Leu Glu Ile Lys
 100 105 110

EP 3 338 793 A1

<210> 51
 <211> 121
 <212> PRT
 <213> Mus musculus

5

<220>
 <221> MISC_FEATURE
 <223> Murine SC17.22 VH

10

<400> 51

Gln Ile Gln Met Met Gln Ser Gly Pro Glu Leu Lys Lys Pro Gly Glu
 1 5 10 15

15

Thr Val Lys Ile Ser Cys Lys Ala Ser Gly Tyr Ser Phe Thr Asn Tyr
 20 25 30

20

Gly Met Asn Trp Val Lys Gln Ala Pro Gly Lys Gly Leu Lys Trp Met
 35 40 45

25

Gly Trp Ile Asn Thr Tyr Thr Gly Glu Pro Thr Tyr Ala Asp Asp Phe
 50 55 60

Lys Gly Arg Phe Ala Phe Ser Leu Glu Thr Ser Ala Ser Thr Ala Tyr
 65 70 75 80

30

Leu Gln Ile Asn Asn Leu Lys Asn Glu Asp Met Ala Thr Tyr Phe Cys
 85 90 95

35

Thr Arg Gly Tyr Tyr Gly Ser Ser Tyr Asp Ala Leu Asp Tyr Trp Gly
 100 105 110

Gln Gly Thr Ser Val Thr Val Ser Ser
 115 120

40

<210> 52
 <211> 112
 <212> PRT
 <213> Mus musculus

45

<220>
 <221> MISC_FEATURE
 <223> Murine SC17.24 VL

50

<400> 52

Asp Ile Val Met Ser Gln Ser Pro Ser Ser Leu Ala Val Ser Val Gly
 1 5 10 15

55

Glu Lys Val Thr Met Ser Cys Lys Ser Ser Gln Ser Leu Leu Tyr Ser
 20 25 30

EP 3 338 793 A1

Ser Asn Gln Lys Ser Tyr Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln
 35 40 45
 5 Ser Pro Lys Leu Leu Ile Tyr Trp Ala Ser Thr Arg Glu Ser Gly Val
 50 55 60
 10 Pro Asp Arg Phe Thr Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr
 65 70 75 80
 15 Ile Ser Ser Val Gln Ala Glu Asp Leu Ala Val Tyr Tyr Cys Lys Gln
 85 90 95
 20 Ser Tyr Asn Leu Arg Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys
 100 105 110
 25 <210> 53
 <211> 117
 <212> PRT
 <213> Mus musculus
 30 <220>
 <221> MISC_FEATURE
 <223> Murine SC17.24 VH
 <400> 53
 35 Gln Val Gln Leu Gln Gln Ser Asp Ala Glu Leu Val Lys Pro Gly Ala
 1 5 10 15
 40 Ser Val Lys Ile Ser Cys Lys Val Ser Gly Tyr Thr Phe Thr Asp His
 20 25 30
 45 Thr Ile His Trp Met Lys Gln Arg Pro Glu Gln Gly Leu Glu Trp Ile
 35 40 45
 50 Gly Tyr Ile Tyr Pro Arg Asp Gly Ser Thr Lys Tyr Asn Glu Glu Phe
 50 55 60
 55 Lys Gly Lys Ala Thr Leu Thr Ala Asp Lys Ser Ser Ser Thr Ala Tyr
 65 70 75 80
 60 Met Gln Leu Asn Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Phe Cys
 85 90 95
 65 Ala Arg Ser Tyr Ser Asn Tyr Phe Asp Tyr Trp Gly Gln Gly Thr Thr
 100 105 110
 70 Leu Thr Val Ser Ser

EP 3 338 793 A1

115

5 <210> 54
 <211> 112
 <212> PRT
 <213> Mus musculus

10 <220>
 <221> MISC_FEATURE
 <223> Murine SC17.27 VL

<400> 54

15 Asp Val Val Met Thr Gln Thr Pro Leu Thr Leu Ser Val Thr Ile Gly
 1 5 10 15

Gln Pro Ala Ser Ile Ser Cys Lys Ser Ser Gln Ser Leu Leu Glu Ser
 20 20 25 30

Asp Gly Lys Thr Tyr Leu Asn Trp Leu Leu Gln Arg Pro Gly Gln Ser
 35 40 45

25 Pro Lys Arg Leu Ile Tyr Leu Val Ser Lys Leu Asp Ser Gly Val Pro
 50 55 60

30 Asp Arg Phe Thr Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile
 65 70 75 80

Ser Arg Val Glu Ala Glu Asp Leu Gly Val Tyr Tyr Cys Trp Gln Gly
 85 90 95

35 Ile Gln His Pro Arg Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys
 100 105 110

40 <210> 55
 <211> 113
 <212> PRT
 <213> Mus musculus

45 <220>
 <221> MISC_FEATURE
 <223> Murine SC17.27 VH

50 <400> 55

Gln Val Gln Leu Gln Gln Ser Gly Ala Glu Leu Val Arg Pro Gly Ala
 1 5 10 15

55 Ser Val Thr Leu Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asp Tyr
 20 25 30

EP 3 338 793 A1

Glu Met His Trp Val Lys Gln Thr Pro Val His Gly Leu Glu Trp Ile
35 40 45

5 Gly Gly Ile Asp Pro Glu Thr Gly Gly Thr Ala Tyr Asn Gln Lys Phe
50 55 60

10 Lys Gly Lys Ala Thr Leu Thr Ala Asp Lys Ser Ser Ser Thr Ala Tyr
65 70 75 80

Met Glu Leu Arg Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Phe Cys
85 90 95

15 Thr Arg Trp Phe Ser Tyr Trp Gly Pro Gly Thr Leu Val Thr Val Ser
100 105 110

20 Ala

25 <210> 56
<211> 107
<212> PRT
<213> Mus musculus

30 <220>
<221> MISC_FEATURE
<223> Murine SC17.28 VL

<400> 56

35 Asp Ile Leu Leu Thr Gln Ser Pro Ala Ile Leu Ser Val Ser Pro Gly
1 5 10 15

Glu Gly Val Ser Phe Ser Cys Arg Ala Ser Gln Ser Ile Gly Thr Ser
20 25 30

40 Ile His Trp Tyr Gln Gln Arg Thr Asn Gly Ser Pro Arg Leu Leu Ile
35 40 45

45 Lys Tyr Ala Ser Glu Ser Ile Ser Gly Ile Pro Ser Arg Phe Ser Gly
50 55 60

50 Ser Gly Ser Gly Thr Asp Phe Thr Leu Arg Ile Asn Ser Leu Glu Ser
65 70 75 80

Glu Asp Ile Ala Asp Tyr Tyr Cys Gln Gln Ser Asn Ser Trp Pro Leu
85 90 95

55 Thr Phe Gly Ala Gly Thr Lys Leu Glu Leu Lys
100 105

EP 3 338 793 A1

<210> 57
 <211> 117
 <212> PRT
 <213> Mus musculus

5

<220>
 <221> MISC_FEATURE
 <223> Murine SC17.28 VH

10

<400> 57

Gln Val His Leu Pro Gln Ser Arg Pro Glu Leu Val Lys Pro Gly Ala
 1 5 10 15

15

Ser Val Lys Ile Ser Cys Lys Ala Ser Gly Tyr Gly Phe Thr Arg Ser
 20 25 30

20

Tyr Ile His Trp Val Lys Gln Arg Pro Gly Gln Gly Leu Glu Trp Ile
 35 40 45

25

Gly Tyr Ile Ser Ser Gly Ser Gly Gly Thr Thr Tyr Asn Gln Lys Phe
 50 55 60

Lys Gly Lys Ala Ser Leu Thr Ala Asp Asn Pro Ser Ser Thr Ala Tyr
 65 70 75 80

30

Met His Leu Ser Ser Leu Thr Ser Glu Asp Ser Ala Ile Tyr Phe Cys
 85 90 95

35

Ala Arg Gly Gly Val Arg Tyr Phe Asp Val Trp Gly Ala Gly Thr Thr
 100 105 110

Val Thr Val Ser Ser
 115

40

<210> 58
 <211> 107
 <212> PRT
 <213> Mus musculus

45

<220>
 <221> MISC_FEATURE
 <223> Murine SC17.29 VL

50

<400> 58

Asp Ile Val Met Thr Gln Ser His Lys Phe Met Ser Thr Ser Val Gly
 1 5 10 15

55

Asp Arg Val Ser Ile Thr Cys Lys Ala Ser Gln Asp Val Gly Thr Asp
 20 25 30

EP 3 338 793 A1

	Val	Ala	Trp	Tyr	Gln	Gln	Lys	Pro	Gly	Gln	Ser	Pro	Lys	Leu	Leu	Ile
			35					40					45			
5	Tyr	Trp	Ala	Ser	Thr	Arg	His	Thr	Gly	Val	Pro	Asp	Arg	Phe	Thr	Gly
		50					55					60				
10	Ser	Gly	Ser	Gly	Thr	Asp	Phe	Thr	Leu	Thr	Ile	Ser	Asn	Val	Gln	Ser
	65					70					75				80	
15	Glu	Asp	Leu	Ala	Asp	Tyr	Phe	Cys	Gln	Gln	Tyr	Ser	Ser	Tyr	Pro	Tyr
					85					90					95	
20	Thr	Phe	Gly	Gly	Gly	Thr	Lys	Leu	Glu	Ile	Lys					
				100					105							
	<210>	59														
	<211>	116														
	<212>	PRT														
	<213>	Mus musculus														
25	<220>															
	<221>	MISC_FEATURE														
	<223>	Murine SC17.29 VH														
30	<400>	59														
	Glu	Val	Gln	Leu	Gln	Gln	Ser	Gly	Pro	Glu	Leu	Met	Lys	Pro	Gly	Ala
	1				5					10					15	
35	Ser	Val	Lys	Met	Ser	Cys	Lys	Ala	Ser	Gly	Tyr	Thr	Phe	Thr	Asp	Tyr
				20					25					30		
40	Asn	Met	His	Trp	Val	Lys	Gln	Asn	Gln	Gly	Lys	Ser	Leu	Glu	Trp	Ile
			35					40					45			
45	Gly	Glu	Ile	Asn	Pro	His	Asn	Gly	Gly	Thr	Gly	Tyr	Asn	Gln	Lys	Phe
		50					55					60				
50	Lys	Gly	Lys	Ala	Thr	Leu	Thr	Val	Asp	Lys	Ser	Ser	Ser	Thr	Ser	Tyr
	65					70					75				80	
55	Met	Glu	Leu	Arg	Ser	Leu	Thr	Ser	Glu	Asp	Ser	Ala	Val	Tyr	Tyr	Cys
					85					90					95	
	Ala	Gly	Gly	Tyr	Pro	Ala	Phe	Asp	Tyr	Trp	Gly	Gln	Gly	Thr	Thr	Leu
				100					105					110		
	Thr	Val	Ser	Ser												

EP 3 338 793 A1

115

5 <210> 60
 <211> 108
 <212> PRT
 <213> Mus musculus

10 <220>
 <221> MISC_FEATURE
 <223> Murine SC17.30 VL

<400> 60

15 Glu Asn Val Leu Thr Gln Ser Pro Ala Ile Val Ser Ala Ser Pro Gly
 1 5 10 15

20 Glu Lys Val Thr Met Thr Cys Arg Ala Ser Ser Ser Val Ile Ser Ser
 20 25 30

Tyr Leu His Trp Tyr Gln Gln Lys Ser Gly Ala Ser Pro Lys Leu Trp
 35 40 45

25 Ile Tyr Ser Thr Ser Asn Leu Ala Ser Gly Val Pro Ala Arg Phe Ser
 50 55 60

30 Gly Ser Ala Ser Gly Thr Ser Tyr Ser Leu Thr Ile Ser Ser Val Glu
 65 70 75 80

35 Ala Glu Asp Ala Ala Thr Tyr Tyr Cys Gln Gln Tyr Ser Gly Tyr Pro
 85 90 95

Leu Thr Phe Gly Ala Gly Thr Lys Leu Glu Leu Lys
 100 105

40 <210> 61
 <211> 124
 <212> PRT
 <213> Mus musculus

45 <220>
 <221> MISC_FEATURE
 <223> Murine SC17.30 VH

50 <400> 61

Glu Val Lys Leu Val Glu Ser Glu Gly Gly Leu Val Gln Pro Gly Ser
 1 5 10 15

55 Ser Met Lys Leu Ser Cys Thr Ala Ser Gly Phe Thr Phe Ser Asp Tyr
 20 25 30

EP 3 338 793 A1

Tyr Met Ala Trp Val Arg Gln Val Pro Glu Lys Gly Leu Glu Trp Val
 35 40 45
 5 Ala Asn Ile Asn Tyr Asp Gly Ser Ser Thr Tyr Tyr Leu Asp Ser Leu
 50 55 60
 10 Lys Ser Arg Phe Ile Ile Ser Arg Asp Asn Ala Lys Asn Ile Leu Tyr
 65 70 75 80
 Leu Gln Met Ser Ser Leu Lys Ser Glu Asp Thr Ala Thr Tyr Tyr Cys
 85 90 95
 15 Ala Arg Asp Asp Tyr Tyr Gly Ser Ser Pro Ser Tyr Trp Tyr Phe Asp
 100 105 110
 20 Val Trp Gly Ala Gly Thr Thr Val Thr Val Ser Ser
 115 120
 25 <210> 62
 <211> 107
 <212> PRT
 <213> Mus musculus
 30 <220>
 <221> MISC_FEATURE
 <223> Murine SC17.32 VL
 <400> 62
 35 Asp Ile Gln Met Thr Gln Ser Pro Ala Ser Leu Ser Ala Ser Val Gly
 1 5 10 15
 Glu Thr Val Thr Met Thr Cys Arg Ala Ser Gly Asn Ile His Asn Tyr
 20 25 30
 40 Leu Val Trp Tyr Gln Gln Lys Gln Gly Lys Ser Pro Gln Leu Leu Val
 35 40 45
 45 Tyr Asn Ala Lys Thr Leu Ala Asp Gly Val Pro Ser Arg Phe Ser Gly
 50 55 60
 50 Ser Gly Ser Gly Thr Gln Tyr Ser Leu Lys Ile Asn Ser Leu Gln Pro
 65 70 75 80
 Glu Asp Phe Gly Ser Tyr Tyr Cys Gln His Phe Trp Ser Thr Pro Pro
 85 90 95
 55 Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys
 100 105

EP 3 338 793 A1

<210> 63
 <211> 119
 <212> PRT
 <213> Mus musculus

<220>
 <221> MISC_FEATURE
 <223> Murine SC17.32 VH

<400> 63

Glu Val Lys Leu Glu Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
 1 5 10 15

Ser Met Lys Leu Ser Cys Val Ala Ser Gly Phe Thr Phe Ser Asn Tyr
 20 25 30

Trp Met Ser Trp Val Arg Gln Ser Pro Glu Lys Gly Leu Glu Trp Val
 35 40 45

Ala Glu Ile Arg Leu Lys Ser Asn Asn Tyr Ala Thr His Tyr Ala Glu
 50 55 60

Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asp Ser Lys Ser Ser
 65 70 75 80

Val Phe Leu Gln Met Asn Asn Leu Arg Thr Glu Asp Thr Gly Ile Tyr
 85 90 95

Tyr Cys Thr Arg His Tyr Tyr Tyr Ala Met Asp Tyr Trp Gly Gln Gly
 100 105 110

Thr Ser Val Thr Val Ser Ser
 115

<210> 64
 <211> 107
 <212> PRT
 <213> Mus musculus

<220>
 <221> MISC_FEATURE
 <223> Murine SC17.34 VL

<400> 64

Asp Ile Lys Met Thr Gln Ser Pro Ser Ser Met Tyr Ala Ser Leu Gly
 1 5 10 15

Glu Arg Val Thr Ile Thr Cys Lys Ala Ser Gln Asp Ile Asn Ser Tyr
 20 25 30

EP 3 338 793 A1

Leu Ser Trp Phe Gln Gln Lys Pro Gly Lys Ser Pro Lys Thr Leu Ile
 35 40 45
 5 Tyr Arg Ala Asn Arg Leu Val Asp Gly Val Pro Ser Arg Phe Ser Gly
 50 55 60
 10 Ser Gly Ser Gly Gln Asp Tyr Ser Leu Thr Ile Ser Ser Leu Glu Tyr
 65 70 75 80
 15 Glu Asp Met Gly Ile Tyr Tyr Cys Leu Gln Tyr Asp Glu Phe Pro Pro
 85 90 95
 20 Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys
 100 105
 <210> 65
 <211> 118
 <212> PRT
 <213> Mus musculus
 25 <220>
 <221> MISC_FEATURE
 <223> Murine SC17.34 VH
 <400> 65
 30 Glu Val Gln Leu Gln Gln Ser Gly Pro Glu Leu Val Lys Pro Gly Ser
 1 5 10 15
 35 Ser Val Lys Ile Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asp Tyr
 20 25 30
 40 Asn Met Asp Trp Val Lys Gln Ser His Gly Lys Arg Leu Glu Trp Ile
 35 40 45
 Gly Tyr Ile Tyr Pro Asp Asn Gly Gly Ala Gly Tyr Asn Gln Lys Phe
 50 55 60
 45 Lys Gly Lys Ala Thr Leu Thr Val Asp Lys Ser Ser Ser Thr Ala Tyr
 65 70 75 80
 50 Met Glu Leu Arg Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys
 85 90 95
 Ser Arg Ser Ile Thr Thr Ala Trp Phe Ala Tyr Trp Gly Gln Gly Thr
 100 105 110
 55 Leu Val Thr Val Ser Ala

EP 3 338 793 A1

115

5 <210> 66
 <211> 108
 <212> PRT
 <213> Mus musculus

10 <220>
 <221> MISC_FEATURE
 <223> Murine SC17.35 VL

<400> 66

15 Glu Asn Val Leu Thr Gln Ser Pro Ala Ile Met Ser Ala Ser Pro Gly
 1 5 10 15

20 Glu Lys Val Thr Leu Thr Cys Arg Ala Ser Ser Ser Met Ser Ser Ser
 20 25 30

Tyr Leu His Trp Tyr Gln Gln Lys Ser Gly Ala Ser Pro Lys Leu Trp
 35 40 45

25 Ile Tyr Ser Thr Ser Asn Leu Ala Ser Gly Val Pro Ala Arg Phe Ser
 50 55 60

30 Gly Ser Gly Ser Gly Thr Ser Tyr Ser Leu Thr Ile Ser Ser Val Glu
 65 70 75 80

35 Ala Glu Asp Ala Ala Thr Tyr Tyr Cys Gln Gln Tyr Ser Ala Tyr Pro
 85 90 95

Phe Thr Phe Gly Ser Gly Thr Lys Leu Glu Ile Lys
 100 105

40 <210> 67
 <211> 121
 <212> PRT
 <213> Mus musculus

45 <220>
 <221> MISC_FEATURE
 <223> Murine SC17.35 VH

50 <400> 67

Glu Val Gln Leu Gln Gln Ser Gly Pro Glu Leu Val Lys Pro Gly Ala
 1 5 10 15

55 Leu Val Lys Met Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asp Tyr
 20 25 30

EP 3 338 793 A1

Tyr Ile His Trp Val Lys Gln Ser His Gly Lys Ser Leu Glu Trp Ile
 35 40 45
 5 Gly Glu Ile Asn Pro Tyr Asn Gly Glu Thr Phe Tyr Asn Gln Lys Phe
 50 55 60
 10 Lys Gly Lys Ala Thr Leu Thr Val Asp Lys Ser Ser Ser Thr Ala Tyr
 65 70 75 80
 Met Glu Leu Arg Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys
 85 90 95
 15 Ala Arg Arg Gly Trp Tyr Leu Thr Gly Tyr Ala Met Asp Tyr Trp Gly
 100 105 110
 20 Gln Gly Thr Ser Val Thr Val Ser Ser
 115 120
 25 <210> 68
 <211> 106
 <212> PRT
 <213> Mus musculus
 30 <220>
 <221> MISC_FEATURE
 <223> Murine SC17.36 VL
 <400> 68
 35 Gln Ile Val Leu Thr Gln Ser Pro Ala Leu Met Ser Ala Ser Pro Gly
 1 5 10 15
 Glu Lys Val Thr Met Thr Cys Ser Ala Ser Ser Ser Val Ser Tyr Met
 20 25 30
 40 Tyr Trp Tyr Gln Gln Lys Pro Arg Ser Ser Pro Lys Pro Trp Ile Tyr
 35 40 45
 45 Leu Thr Ser Asn Leu Ala Ser Gly Val Pro Ala Arg Phe Ser Gly Ser
 50 55 60
 50 Gly Ser Gly Thr Ser Tyr Ser Leu Thr Ile Ser Ser Met Glu Ala Glu
 65 70 75 80
 Asp Ala Ala Thr Tyr Tyr Cys Gln Gln Trp Ser Ser Asn Pro Pro Thr
 85 90 95
 55 Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys
 100 105

EP 3 338 793 A1

<210> 69
 <211> 120
 <212> PRT
 <213> Mus musculus

<220>
 <221> MISC_FEATURE
 <223> Murine SC17.36 VH

<400> 69

Glu Val Gln Leu Gln Glu Ser Gly Pro Ser Leu Val Lys Pro Ser Gln
 1 5 10 15

Ser Gln Ser Leu Thr Cys Ser Val Thr Gly Asp Ser Ile Thr Ser Asp
 20 25 30

Tyr Trp Asn Trp Ile Arg Lys Phe Pro Gly Lys Lys Val Glu Tyr Met
 35 40 45

Gly Tyr Ile Asn Tyr Ser Gly Ser Thr Tyr Tyr Asn Pro Ser Leu Lys
 50 55 60

Ser Arg Ile Ser Ile Thr Arg Asp Thr Ser Lys Asn Gln Tyr Tyr Leu
 65 70 75 80

Gln Leu Asn Ser Val Thr Ser Glu Asp Thr Ala Thr Tyr Tyr Cys Ala
 85 90 95

Arg Thr Ser Tyr Tyr Asn Lys Phe Leu Pro Phe Ala Tyr Trp Gly Gln
 100 105 110

Gly Thr Leu Val Thr Val Ser Ala
 115 120

<210> 70
 <211> 112
 <212> PRT
 <213> Mus musculus

<220>
 <221> MISC_FEATURE
 <223> Murine SC17.39 VL

<400> 70

Asp Val Leu Met Thr Gln Thr Pro Leu Ser Leu Pro Val Ser Leu Gly
 1 5 10 15

Asp Gln Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Leu Val His Arg
 20 25 30

EP 3 338 793 A1

	Asn	Gly	Asn	Thr	Tyr	Phe	His	Trp	Tyr	Leu	Gln	Lys	Pro	Gly	Gln	Ser	
			35					40					45				
5	Pro	Lys	Leu	Leu	Ile	Tyr	Lys	Val	Ser	Asn	Arg	Phe	Ser	Gly	Val	Pro	
		50					55					60					
10	Asp	Arg	Phe	Ser	Gly	Ser	Gly	Ser	Gly	Thr	Asp	Phe	Thr	Leu	Lys	Ile	
	65					70					75					80	
15	Ser	Arg	Val	Glu	Ala	Glu	Asp	Leu	Gly	Val	Tyr	Phe	Cys	Ser	Gln	Ser	
				85						90					95		
20	Thr	Tyr	Val	Pro	Trp	Thr	Phe	Gly	Gly	Gly	Thr	Lys	Leu	Glu	Ile	Lys	
				100					105						110		
	<210>	71															
	<211>	117															
	<212>	PRT															
	<213>	Mus musculus															
25	<220>																
	<221>	MISC_FEATURE															
	<223>	Murine SC17.39 VH															
30	<400>	71															
	Glu	Val	Gln	Leu	Val	Glu	Ser	Gly	Gly	Gly	Leu	Val	Gln	Pro	Gly	Gly	
	1				5					10					15		
35	Ser	Arg	Lys	Leu	Ser	Cys	Ala	Ala	Ser	Gly	Phe	Thr	Phe	Ser	Ser	Tyr	
				20					25					30			
40	Gly	Met	His	Trp	Val	Arg	Gln	Ala	Pro	Glu	Lys	Gly	Leu	Glu	Trp	Val	
			35					40					45				
45	Ala	Tyr	Ile	Ser	Ser	Asn	Asp	Gly	Thr	Ile	Tyr	Tyr	Ala	Asp	Thr	Val	
		50					55					60					
50	Arg	Gly	Arg	Phe	Thr	Ile	Ser	Arg	Asp	Asn	Ala	Lys	Asn	Thr	Leu	Phe	
	65					70					75					80	
55	Leu	Gln	Met	Thr	Ser	Leu	Arg	Ser	Glu	Asp	Thr	Ala	Met	Tyr	Tyr	Cys	
					85					90					95		
	Ala	Arg	Pro	Ser	Asn	Trp	Val	Phe	Asp	Tyr	Trp	Gly	Gln	Gly	Thr	Thr	
					100				105					110			
	Leu	Thr	Val	Ser	Ser												

115

5 <210> 72
 <211> 112
 <212> PRT
 <213> Mus musculus

10 <220>
 <221> MISC_FEATURE
 <223> Murine SC17.40 VL
 <400> 72

15 Asp Val Val Met Thr Gln Thr Pro Leu Ser Arg Pro Val Thr Leu Gly
 1 5 10 15

20 Asp Gln Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Leu Val His Ser
 20 25 30

25 Asn Gly Asn Thr Tyr Leu His Trp Tyr Leu Gln Lys Pro Gly Gln Ser
 35 40 45

30 Pro Lys Leu Leu Ile Tyr Lys Val Ser Asn Arg Phe Ser Gly Val Pro
 50 55 60

35 Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile
 65 70 75 80

40 Ser Arg Val Glu Ala Glu Asp Leu Gly Val Tyr Phe Cys Ser Gln Asn
 85 90 95

45 Thr His Val Pro Phe Thr Phe Gly Ser Gly Thr Lys Leu Glu Ile Lys
 100 105 110

50 <210> 73
 <211> 116
 <212> PRT
 <213> Mus musculus

55 <220>
 <221> MISC_FEATURE
 <223> Murine SC17.40 VH
 <400> 73

60 Gln Val Gln Leu Gln Gln Pro Gly Ala Glu Ile Val Arg Pro Gly Ala
 1 5 10 15

65 Ser Val Lys Leu Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asp Tyr
 20 25 30

EP 3 338 793 A1

	Trp Met Asn Trp Val Lys Gln Arg Pro Gly Gln Gly Leu Glu Trp Ile	
	35 40 45	
5	Gly Thr Ile Asp Pro Ser Asp Ser Tyr Thr Arg Tyr Asn Gln Lys Phe	
	50 55 60	
10	Lys Gly Lys Ala Thr Leu Thr Val Asp Thr Ser Phe Ser Ser Ala Tyr	
	65 70 75 80	
15	Met Gln Leu Ser Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Phe Cys	
	85 90 95	
20	Ala Ser Gly Gly Arg Gly Phe Gly Tyr Trp Gly Gln Gly Thr Pro Val	
	100 105 110	
25	Thr Val Ser Val	
	115	
	<210> 74	
	<211> 106	
	<212> PRT	
	<213> Mus musculus	
30	<220>	
	<221> MISC_FEATURE	
	<223> Murine SC17.41 VL	
	<400> 74	
35	Gln Ile Val Leu Thr Gln Ser Pro Ala Leu Met Ser Ala Ser Pro Gly	
	1 5 10 15	
40	Glu Lys Val Thr Met Thr Cys Ser Ala Ser Ser Ser Val Ser Tyr Met	
	20 25 30	
45	Tyr Trp Tyr Gln Gln Lys Pro Arg Ser Ser Pro Lys Pro Trp Ile Tyr	
	35 40 45	
50	Leu Thr Ser Asn Leu Ala Ser Gly Val Pro Thr Arg Phe Ser Gly Ser	
	50 55 60	
	Gly Ser Gly Thr Ser Tyr Ser Leu Thr Ile Ser Ser Met Gly Ala Glu	
	65 70 75 80	
55	Asp Ala Ala Thr Tyr Tyr Cys Gln Gln Trp Asn Thr Asn Pro Pro Thr	
	85 90 95	
	Phe Gly Ala Gly Thr Lys Leu Glu Leu Lys	
	100 105	

EP 3 338 793 A1

<210> 75
 <211> 121
 <212> PRT
 <213> Mus musculus

<220>
 <221> MISC_FEATURE
 <223> Murine SC17.41 VH

<400> 75

Asp Val Lys Leu Val Glu Ser Gly Gly Gly Leu Val Lys Leu Gly Gly
 1 5 10 15

Ser Leu Lys Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
 20 25 30

Ala Met Ser Trp Val Arg Gln Thr Pro Glu Lys Arg Leu Glu Trp Val
 35 40 45

Ala Thr Ile Ser Ser Gly Gly Gly Asn Thr Tyr Tyr Pro Asp Ser Val
 50 55 60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Thr Leu Tyr
 65 70 75 80

Leu Gln Met Ser Ser Leu Lys Ser Glu Asp Thr Ala Met Tyr Tyr Cys
 85 90 95

Ala Arg Arg Asp Tyr Tyr Gly Thr Ser Tyr Val Met Phe Ala Tyr Trp
 100 105 110

Gly Gln Gly Thr Leu Val Thr Val Ser
 115 120

<210> 76
 <211> 106
 <212> PRT
 <213> Mus musculus

<220>
 <221> MISC_FEATURE
 <223> Murine SC17.42 VL

<400> 76

Glu Asn Val Leu Thr Gln Ser Pro Ala Ile Met Ser Ala Ser Pro Gly
 1 5 10 15

Glu Lys Val Thr Met Thr Cys Ser Ala Ser Ser Ser Val Asn Tyr Met
 20 25 30

EP 3 338 793 A1

Tyr Trp Tyr Gln Gln Lys Ser Ser Thr Ser Pro Lys Leu Trp Ile Tyr
 35 40 45
 5 Asp Thr Ser Lys Leu Thr Ser Gly Val Pro Gly Arg Phe Ser Gly Ser
 50 55 60
 10 Gly Ser Gly Asn Ser Tyr Ser Leu Thr Ile Ser Asn Met Glu Ala Glu
 65 70 75 80
 15 Asp Val Ala Thr Tyr Tyr Cys Phe Gln Gly Ser Gly Tyr Pro Leu Thr
 85 90 95
 20 Phe Gly Ser Gly Thr Lys Leu Glu Ile Lys
 100 105
 <210> 77
 <211> 112
 <212> PRT
 <213> Mus musculus
 25 <220>
 <221> MISC_FEATURE
 <223> Murine SC17.42 VH
 30 <400> 77
 Asp Val Lys Leu Val Glu Ser Gly Gly Gly Leu Val Arg Pro Gly Gly
 1 5 10 15
 35 Ser Leu Lys Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Arg Tyr
 20 25 30
 40 Thr Met Ser Trp Val Arg Gln Thr Pro Glu Lys Arg Leu Glu Trp Ala
 35 40 45
 45 Ala Thr Ile Asn Ser Gly Gly Ser Asn Thr Tyr Tyr Pro Asp Ser Val
 50 55 60
 50 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Thr Leu Phe
 65 70 75 80
 55 Leu Gln Met Ser Ser Leu Lys Ser Glu Asp Thr Ala Met Tyr Tyr Cys
 85 90 95
 Thr Asn Gly Asn His Trp Gly Gln Gly Thr Thr Leu Thr Val Ser Ser
 100 105 110
 <210> 78

EP 3 338 793 A1

<211> 106
<212> PRT
<213> Mus musculus

<220>
<221> MISC_FEATURE
<223> Murine SC17.45 VL

<400> 78

Glu Asn Val Leu Thr Gln Ser Pro Ala Ile Met Ser Ala Ser Pro Gly
1 5 10 15

Glu Lys Val Thr Met Thr Cys Ser Ala Ser Ser Ser Val Asn Tyr Met
20 25 30

Tyr Trp Tyr Gln Gln Lys Ser Ser Thr Ser Pro Lys Leu Trp Ile Tyr
35 40 45

Asp Thr Ser Lys Leu Thr Ser Gly Val Pro Gly Arg Phe Ser Gly Ser
50 55 60

Gly Ser Gly Asn Ser Tyr Ser Leu Thr Ile Ser Asn Met Glu Ala Glu
65 70 75 80

Asp Val Ala Thr Tyr Tyr Cys Phe Gln Gly Ser Gly Tyr Pro Leu Thr
85 90 95

Phe Gly Ser Gly Thr Lys Leu Glu Ile Lys
100 105

<210> 79
<211> 114
<212> PRT
<213> Mus musculus

<220>
<221> MISC_FEATURE
<223> Murine SC17.45 VH

<400> 79

Gln Val Gln Leu Gln Gln Pro Gly Ser Val Leu Val Arg Pro Gly Asp
1 5 10 15

Ser Val Lys Leu Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Ser Tyr
20 25 30

Trp Met His Trp Val Lys Gln Ser Pro Gly Gln Gly Leu Glu Trp Ile
35 40 45

EP 3 338 793 A1

Gly Glu Ile His Pro His Ser Gly Ser Thr Asn Tyr Asn Glu Lys Phe
50 55 60

5 Lys Gly Lys Ala Thr Leu Thr Val Asp Thr Ser Ser Ser Thr Ala Tyr
65 70 75 80

10 Val Asp Leu Ser Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys
85 90 95

Val Gly Gly His Tyr Asp Tyr Trp Gly Gln Gly Thr Thr Leu Thr Val
100 105 110

15 Ser Ser

<210> 80
<211> 107
<212> PRT
<213> Mus musculus

25 <220>
<221> MISC_FEATURE
<223> Murine SC17.46 VL

<400> 80

30 Ser Phe Val Met Thr Gln Thr Pro Lys Phe Leu Leu Val Ser Ala Gly
1 5 10 15

35 Asp Arg Val Thr Ile Thr Cys Lys Ala Ser Gln Ser Val Asn Asn Asp
20 25 30

Val Ala Trp Tyr Gln Gln Lys Pro Gly Gln Ser Pro Lys Leu Leu Ile
35 40 45

40 Tyr Tyr Ala Ser Asn Arg Tyr Thr Gly Val Pro Asp Arg Phe Thr Gly
50 55 60

45 Ser Gly Tyr Gly Thr Asp Phe Thr Phe Thr Ile Ser Thr Val Gln Ala
65 70 75 80

Glu Asp Leu Ala Val Tyr Phe Cys Gln Gln Asp Tyr Ser Ser Pro Arg
85 90 95

50 Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys
100 105

55 <210> 81
<211> 119
<212> PRT

EP 3 338 793 A1

<213> Mus musculus

<220>

<221> MISC_FEATURE

<223> Murine SC17.46 VH

<400> 81

Gln Val Gln Leu Gln Gln Pro Gly Ala Glu Leu Val Lys Pro Gly Ala
1 5 10 15

Ser Met Lys Leu Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Ser Tyr
20 25 30

Trp Ile Asn Trp Val Lys Gln Arg Pro Gly Gln Gly Leu Glu Trp Ile
35 40 45

Gly Asn Ile Phe Pro Asp Thr Thr Thr Thr Asn Tyr Asn Glu Lys Phe
50 55 60

Lys Ser Lys Ala Thr Leu Thr Val Asp Thr Ser Ser Ser Thr Ala Tyr
65 70 75 80

Met Gln Leu Ser Ser Leu Thr Ser Asp Asp Ser Ala Val Tyr Tyr Cys
85 90 95

Ala Arg Glu Tyr Tyr Asp Gly Thr Tyr Asp Ala Met Asp Tyr Trp Gly
100 105 110

Gln Gly Thr Ser Val Thr Val
115

<210> 82

<211> 106

<212> PRT

<213> Mus musculus

<220>

<221> MISC_FEATURE

<223> Murine SC17.47 VL

<400> 82

Gln Ile Val Leu Thr Gln Ser Pro Ala Ile Met Ser Ala Ser Pro Gly
1 5 10 15

Glu Lys Val Ser Met Thr Cys Ser Ala Ser Ser Ser Val Ser Tyr Met
20 25 30

His Trp Tyr Gln Gln Lys Ser Gly Thr Ser Pro Lys Arg Trp Ile Tyr
35 40 45

EP 3 338 793 A1

Asp Thr Ser Lys Leu Ala Ser Gly Val Pro Ala Arg Phe Ser Gly Ser
50 55 60

5 Gly Ser Gly Ser Ser Tyr Ser Leu Thr Ile Ser Ser Met Glu Ala Glu
65 70 75 80

10 Asp Ala Ala Thr Tyr Tyr Cys Gln Gln Trp Ser Ser Thr Pro Pro Thr
85 90 95

Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys
100 105

15 <210> 83
<211> 119
<212> PRT
<213> Mus musculus

20 <220>
<221> MISC_FEATURE
<223> Murine SC17.47 VH

25 <400> 83

Glu Val Gln Leu Gln Gln Ser Gly Pro Glu Leu Val Lys Pro Gly Ala
1 5 10 15

30 Ser Val Lys Ile Ser Cys Lys Ala Ser Gly Tyr Ser Phe Thr Asp Tyr
20 25 30

35 Tyr Met Arg Trp Val Lys Gln Ser Pro Glu Lys Ser Leu Glu Trp Ile
35 40 45

40 Gly Glu Ile Asn Pro Ser Thr Gly Gly Thr Thr Tyr Asn Gln Asn Phe
50 55 60

Lys Ala Lys Ala Thr Leu Thr Val Asp Lys Ser Ser Ser Thr Ala Tyr
65 70 75 80

45 Met Gln Leu Lys Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys
85 90 95

50 Ala Arg Gly Gly Tyr Phe Leu Tyr Tyr Phe Asp Tyr Trp Gly Gln Gly
100 105 110

Thr Thr Leu Thr Val Ser Ser
115

55 <210> 84

EP 3 338 793 A1

<211> 112
<212> PRT
<213> Mus musculus

<220>
<221> MISC_FEATURE
<223> Murine SC17.49 VL

<400> 84

Asp Val Val Met Thr Gln Thr Pro Leu Thr Leu Ser Val Thr Ile Gly
1 5 10 15

Gln Pro Ala Ser Ile Ser Cys Lys Ser Ser Gln Ser Leu Leu Glu Ser
20 25 30

Asp Gly Lys Thr Tyr Leu Asn Trp Leu Leu Gln Arg Pro Gly Gln Ser
35 40 45

Pro Lys Arg Leu Ile Tyr Leu Val Ser Lys Leu Asp Ser Gly Val Pro
50 55 60

Asp Arg Phe Thr Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile
65 70 75 80

Ser Arg Val Glu Ala Glu Asp Leu Gly Val Tyr Tyr Cys Trp Gln Gly
85 90 95

Ile Gln His Pro Arg Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys
100 105 110

<210> 85
<211> 113
<212> PRT
<213> Mus musculus

<220>
<221> MISC_FEATURE
<223> Murine SC17.49 VH

<400> 85

Gln Val Gln Leu Gln Gln Ser Gly Ala Glu Leu Val Arg Pro Gly Ala
1 5 10 15

Ser Val Thr Leu Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asp Tyr
20 25 30

Glu Met His Trp Val Lys Gln Thr Pro Val His Gly Leu Glu Trp Ile
35 40 45

EP 3 338 793 A1

Gly Gly Ile Asp Pro Glu Thr Gly Gly Thr Ala Tyr Asn Gln Lys Phe
50 55 60

5 Lys Gly Lys Ala Thr Leu Thr Ala Asp Lys Ser Ser Ser Thr Ala Tyr
65 70 75 80

10 Met Glu Leu Arg Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Phe Cys
85 90 95

Thr Arg Trp Phe Ser Tyr Trp Gly Pro Gly Thr Leu Val Thr Val Ser
100 105 110

15 Ala

<210> 86
<211> 111
<212> PRT
<213> Mus musculus

25 <220>
<221> MISC_FEATURE
<223> Murine SC17.50 VL

<400> 86

30 Asp Ile Val Leu Thr Gln Ser Pro Ala Ser Leu Ala Ala Ser Leu Gly
1 5 10 15

35 Gln Arg Ala Thr Ile Ser Cys Arg Ala Ser Gln Ser Val Ser Thr Ser
20 25 30

Ser Tyr Ser Tyr Met His Trp Tyr Gln Gln Lys Pro Gly Gln Pro Pro
35 40 45

40 Lys Leu Leu Ile Lys Tyr Ala Ser Asn Leu Glu Ser Gly Val Pro Ala
50 55 60

45 Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Asn Ile His
65 70 75 80

Pro Val Glu Glu Glu Asp Thr Ala Thr Tyr Tyr Cys Gln His Ser Trp
85 90 95

50 Glu Ile Pro Trp Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys
100 105 110

55 <210> 87
<211> 121
<212> PRT

EP 3 338 793 A1

<213> Mus musculus

<220>

<221> MISC_FEATURE

<223> Murine SC17.50 VH

<400> 87

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Lys Pro Gly Gly
1 5 10 15

Ser Leu Lys Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Asp Tyr
20 25 30

Gly Met His Trp Val Arg Gln Ala Pro Glu Lys Gly Leu Glu Trp Val
35 40 45

Ala Tyr Ile Ser Ser Gly Ser Arg Thr Ile Tyr Tyr Ala Asp Thr Val
50 55 60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Thr Leu Phe
65 70 75 80

Leu Gln Met Thr Ser Leu Arg Ser Glu Asp Thr Ala Met Tyr Tyr Cys
85 90 95

Ala Arg Val Tyr Tyr Gly Ser Thr Tyr Gly Tyr Phe Asp Val Trp Gly
100 105 110

Thr Gly Thr Thr Val Thr Val Ser Ser
115 120

<210> 88

<211> 111

<212> PRT

<213> Mus musculus

<220>

<221> MISC_FEATURE

<223> Murine SC17.53 VL

<400> 88

Asp Ile Val Leu Thr Gln Ser Pro Ala Ser Leu Ala Ala Ser Leu Gly
1 5 10 15

Gln Arg Ala Thr Ile Ser Cys Arg Ala Ser Gln Ser Val Ser Thr Ser
20 25 30

Ser Tyr Ser Tyr Met His Trp Tyr Gln Gln Lys Pro Gly His Pro Pro
35 40 45

EP 3 338 793 A1

Lys Leu Leu Ile Arg Tyr Ala Ser Asn Leu Glu Ser Gly Val Pro Ala
50 55 60

5 Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Asn Ile His
65 70 75 80

10 Pro Val Glu Glu Glu Asp Thr Ala Thr Tyr Tyr Cys Gln His Ser Trp
85 90 95

Glu Ile Pro Tyr Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys
100 105 110

15

<210> 89
<211> 119
<212> PRT
<213> Mus musculus

20

<220>
<221> MISC_FEATURE
<223> Murine SC17.53 VH

25

<400> 89

Glu Val Gln Leu Gln Gln Ser Gly Pro Glu Leu Val Lys Pro Gly Ala
1 5 10 15

30

Ser Val Lys Ile Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asp Tyr
20 25 30

35

Asn Met His Trp Val Lys Gln Ser His Gly Lys Arg Leu Glu Trp Ile
35 40 45

Gly Tyr Ile His Pro Tyr Asn Gly Gly Ser Gly Tyr Asn Gln Lys Phe
50 55 60

40

Lys Arg Lys Ala Thr Leu Thr Val Asp Asn Ser Ser Asn Thr Thr Tyr
65 70 75 80

45

Met Glu Leu Arg Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys
85 90 95

50

Ala Arg Ser Tyr Asp Tyr Asp Thr Trp Phe Gly Tyr Trp Gly Gln Gly
100 105 110

Thr Leu Val Thr Val Arg Ala
115

55

<210> 90

EP 3 338 793 A1

<211> 112
<212> PRT
<213> Mus musculus

<220>
<221> MISC_FEATURE
<223> Murine SC17.54 VL

<400> 90

Asp Val Val Leu Thr Gln Thr Pro Leu Thr Leu Ser Val Thr Ile Gly
1 5 10 15

Gln Pro Ala Ser Ile Ser Cys Lys Ser Ser Gln Ser Leu Leu Tyr Ser
20 25 30

Asp Gly Lys Thr Tyr Leu Asn Trp Leu Leu Gln Arg Pro Gly Gln Ser
35 40 45

Pro Lys Arg Leu Ile Tyr Leu Val Ser Lys Leu Asp Ser Gly Val Pro
50 55 60

Asp Arg Phe Thr Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile
65 70 75 80

Ser Arg Val Glu Ala Glu Asp Leu Gly Leu Tyr Tyr Cys Trp Gln Gly
85 90 95

Thr His Phe Pro Trp Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys
100 105 110

<210> 91
<211> 113
<212> PRT
<213> Mus musculus

<220>
<221> MISC_FEATURE
<223> Murine SC17.54 VH

<400> 91

Glu Val Lys Leu Glu Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1 5 10 15

Ser Met Lys Leu Ser Cys Val Ala Ser Gly Phe Thr Phe Ser Asn Tyr
20 25 30

Trp Ile Asn Trp Val Arg Gln Ser Pro Glu Lys Gly Leu Glu Trp Val
35 40 45

EP 3 338 793 A1

Ala Glu Ile Arg Met Lys Ser Asn Asn Tyr Ala Thr His Tyr Ala Glu
50 55 60

5 Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asp Ser Lys Ser Cys
65 70 75 80

10 Val Tyr Leu Gln Met Asn Asn Leu Arg Pro Glu Asp Thr Gly Ile Tyr
85 90 95

Tyr Cys Thr Arg Gly Gly Tyr Trp Gly Gln Gly Thr Thr Leu Thr Val
100 105 110

15 Ser

20 <210> 92
<211> 113
<212> PRT
<213> Mus musculus

25 <220>
<221> MISC FEATURE
<223> Murine SC17.56 VL

<400> 92

30 Asp Ile Val Met Ser Gln Ser Pro Ser Ser Leu Ala Val Ser Val Gly
1 5 10 15

35 Glu Lys Val Thr Met Ser Cys Lys Ser Ser Gln Ser Leu Leu Tyr Ser
20 25 30

Ser Asn Gln Lys Asn Tyr Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln
35 40 45

40 Ser Pro Lys Leu Leu Ile Tyr Trp Ala Ser Thr Arg Glu Ser Gly Val
50 55 60

45 Pro Asp Arg Phe Thr Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr
65 70 75 80

50 Ile Ser Ser Val Lys Ala Glu Asp Leu Ala Val Tyr Phe Cys Gln Gln
85 90 95

Tyr Tyr Asn Tyr Pro Tyr Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile
100 105 110

55 Lys

EP 3 338 793 A1

<210> 93
 <211> 116
 <212> PRT
 <213> Mus musculus

<220>
 <221> MISC_FEATURE
 <223> Murine SC17.56 VH

<400> 93

Gln Ile Gln Leu Val Gln Ser Gly Pro Glu Leu Lys Lys Pro Gly Glu
 1 5 10 15

Thr Val Lys Ile Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asn Tyr
 20 25 30

Gly Met Asn Trp Val Lys Gln Ala Pro Gly Lys Gly Leu Lys Trp Met
 35 40 45

Ala Trp Ile Asn Thr Tyr Thr Gly Glu Pro Thr Tyr Ala Asp Asp Phe
 50 55 60

Lys Gly Arg Phe Ala Phe Ser Leu Glu Thr Ser Ala Ser Thr Ala Ser
 65 70 75 80

Leu Gln Ile Ile Asn Leu Lys Asn Glu Asp Thr Ala Thr Tyr Phe Cys
 85 90 95

Ala Arg Ile Gly Asp Ser Ser Pro Ser Asp Tyr Trp Gly Gln Gly Thr
 100 105 110

Thr Leu Thr Val
 115

<210> 94
 <211> 108
 <212> PRT
 <213> Mus musculus

<220>
 <221> MISC_FEATURE
 <223> Murine SC17.57 VL

<400> 94

Gln Ile Val Leu Thr Gln Ser Pro Ala Ile Met Ser Ala Ser Leu Gly
 1 5 10 15

Glu Arg Val Thr Met Thr Cys Thr Ala Ser Ser Ser Val Ser Ser Ser
 20 25 30

EP 3 338 793 A1

Tyr Leu His Trp Tyr Gln Gln Lys Pro Gly Ser Ser Pro Lys Leu Trp
 35 40 45
 5 Ile Tyr Ser Thr Ser Asn Leu Ala Ser Gly Val Pro Pro Arg Phe Ser
 50 55 60
 10 Gly Ser Gly Ser Gly Thr Ser Tyr Ser Leu Thr Ile Ser Ser Met Glu
 65 70 75 80
 Ala Glu Asp Ala Ala Thr Tyr Tyr Cys His Gln Tyr His Arg Ser Pro
 85 90 95
 15 Pro Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys
 100 105
 20 <210> 95
 <211> 121
 <212> PRT
 <213> Mus musculus
 25 <220>
 <221> MISC FEATURE
 <223> Murine SC17.57 VH
 30 <400> 95
 Gln Ile Gln Leu Val Gln Ser Gly Pro Glu Leu Lys Lys Pro Gly Glu
 1 5 10 15
 35 Thr Val Lys Ile Ser Cys Lys Ala Ser Asp Tyr Thr Phe Thr Asp Phe
 20 25 30
 Ser Ile His Trp Val Arg Gln Ser Pro Gly Lys Gly Leu Lys Trp Met
 35 40 45
 40 Gly Trp Ile Asn Thr Glu Thr Gly Glu Pro Thr Val Ala Glu Asp Phe
 50 55 60
 45 Lys Gly Arg Phe Ala Phe Ser Leu Glu Thr Ser Ala Ser Thr Ala Phe
 65 70 75 80
 50 Leu Gln Ile Tyr Asn Leu Lys Asn Glu Asp Ser Ala Thr Tyr Phe Cys
 85 90 95
 Ala Arg Gly Arg Tyr Tyr Gly His Asp Tyr Ala Met Asp Tyr Trp Gly
 100 105 110
 55 Gln Gly Thr Ser Val Thr Val Ser Ser

EP 3 338 793 A1

115

120

5 <210> 96
 <211> 107
 <212> PRT
 <213> Mus musculus

10 <220>
 <221> MISC_FEATURE
 <223> Murine SC17.59 VL

<400> 96

15 Asp Ile Gln Met Thr Gln Ser Pro Ala Ser Leu Ser Ala Ser Val Gly
 1 5 10 15

20 Glu Thr Val Thr Ile Thr Cys Arg Ala Ser Gly Asn Leu His Asn Tyr
 20 25 30

25 Leu Ala Trp Tyr Gln Gln Lys Gln Gly Lys Ser Pro Gln Leu Leu Val
 35 40 45

30 Tyr Asn Ala Lys Thr Leu Ala Asp Gly Val Pro Ser Arg Phe Ser Gly
 50 55 60

35 Ser Gly Ser Gly Thr Gln Tyr Ser Leu Lys Ile Asn Ser Leu Gln Pro
 65 70 75 80

40 Glu Asp Phe Gly Thr Tyr Phe Cys Gln His Phe Trp Ser Ile Pro Pro
 85 90 95

45 Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys
 100 105

50 <210> 97
 <211> 119
 <212> PRT
 <213> Mus musculus

<220>
 <221> MISC_FEATURE
 <223> Murine SC17.59 VH

<400> 97

55 Glu Val Lys Leu Glu Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
 1 5 10 15

Ser Met Lys Leu Ser Cys Val Ala Ser Gly Phe Thr Phe Ser Asn Tyr
 20 25 30

EP 3 338 793 A1

Trp Met Asn Trp Val Arg Gln Ser Pro Glu Lys Gly Leu Glu Trp Val
35 40 45

5 Ala Glu Ile Arg Leu Lys Ser Asn Asn Tyr Ala Thr His Tyr Ala Glu
50 55 60

10 Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asp Ser Lys Ser Ser
65 70 75 80

15 Val Tyr Leu Gln Met Asn Asn Leu Arg Ala Glu Asp Thr Gly Ile Tyr
85 90 95

20 Tyr Cys Thr Arg Leu Trp Asp Phe Ala Met Asp Tyr Trp Gly Gln Gly
100 105 110

Thr Ser Val Thr Val Ser Ser
115

<210> 98
<211> 106
<212> PRT
<213> Mus musculus

<220>
<221> MISC_FEATURE
<223> Murine SC17.61 VL

<400> 98

35 Gln Ile Val Leu Thr Gln Ser Pro Ala Ile Met Ser Ala Ser Pro Gly
1 5 10 15

Glu Lys Val Thr Ile Ser Cys Ser Ala Ser Ser Ser Val Ser Tyr Ile
20 25 30

40 Tyr Trp Tyr Gln Gln Lys Pro Gly Ser Ser Pro Lys Pro Trp Ile Tyr
35 40 45

45 Arg Thr Ser Asn Leu Ala Ser Gly Val Pro Ala Arg Phe Ser Gly Ser
50 55 60

50 Gly Ser Gly Thr Ser Tyr Ser Leu Thr Ile Ser Ser Met Glu Ala Glu
65 70 75 80

Asp Ala Ala Thr Tyr Tyr Cys Gln Gln Tyr His Ser Tyr Pro Trp Thr
85 90 95

55 Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys
100 105

EP 3 338 793 A1

<210> 99
 <211> 125
 <212> PRT
 <213> Mus musculus

<220>
 <221> MISC_FEATURE
 <223> Murine SC17.61 VH

<400> 99

Gln Val Thr Leu Lys Glu Ser Gly Pro Gly Ile Leu Gln Pro Ser Gln
 1 5 10 15

Thr Leu Ser Leu Thr Cys Ser Phe Ser Gly Phe Ser Leu Ser Thr Phe
 20 25 30

Gly Met Gly Val Gly Trp Ile Arg Gln Pro Ser Gly Lys Gly Leu Glu
 35 40 45

Trp Leu Ala Gln Ile Trp Trp Asp Asp Tyr Lys Tyr Tyr Asn Pro Ala
 50 55 60

Leu Lys Ser Arg Leu Thr Ile Ser Lys Asp Thr Ser Lys Asn Gln Val
 65 70 75 80

Phe Leu Lys Ile Ala Asn Val Asp Thr Ala Asp Thr Ala Thr Tyr Tyr
 85 90 95

Cys Ala Arg Ile Gly Tyr Tyr Ser Gly Ser Ser Arg Cys Trp Tyr Phe
 100 105 110

Asp Val Trp Gly Thr Gly Ser Thr Val Thr Val Ser Ser
 115 120 125

<210> 100
 <211> 107
 <212> PRT
 <213> Mus musculus

<220>
 <221> MISC_FEATURE
 <223> Murine SC17.63 VL

<400> 100

Ser Ile Val Met Thr Gln Thr Pro Lys Phe Leu Leu Val Ser Ala Gly
 1 5 10 15

Asp Arg Val Ala Ile Thr Cys Lys Ala Ser Gln Ser Val Ser Asn Asp
 20 25 30

EP 3 338 793 A1

Val Ala Trp Tyr Gln Gln Lys Pro Gly Gln Ser Pro Thr Leu Leu Ile
35 40 45

5 Ser Tyr Ala Ser Asn Arg Tyr Thr Gly Val Pro Asp Arg Phe Thr Gly
50 55 60

10 Ser Gly Tyr Gly Thr Asp Phe Thr Phe Thr Ile Ser Thr Val Gln Ala
65 70 75 80

15 Glu Asp Leu Ala Val Tyr Phe Cys Gln Gln Gly Tyr Ser Ser Pro Phe
85 90 95

20 Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys
100 105

<210> 101
<211> 116
<212> PRT
<213> Mus musculus

25 <220>
<221> MISC_FEATURE
<223> Murine SC17.63 VH

30 <400> 101

Gln Val Gln Leu Gln Gln Ser Asp Ala Glu Leu Val Lys Pro Gly Ala
1 5 10 15

35 Ser Val Lys Ile Ser Cys Lys Ala Ala Gly Tyr Thr Phe Thr Asp Leu
20 25 30

40 Thr Ile His Trp Val Lys Gln Arg Pro Glu Gln Gly Leu Glu Trp Ile
35 40 45

Gly Tyr Ile Tyr Pro Gly Asp Ser Asn Thr Lys Tyr Asn Glu Lys Phe
50 55 60

45 Lys Gly Lys Ala Thr Leu Thr Ala Asp Lys Ser Ser Ser Thr Ala Tyr
65 70 75 80

50 Met Gln Leu Asn Ser Leu Thr Ser Glu Asp Ser Val Val Tyr Phe Cys
85 90 95

55 Ala Arg Met Ile Thr Pro Tyr Tyr Phe Asp Tyr Trp Gly Gln Gly Thr
100 105 110

Thr Leu Thr Val

EP 3 338 793 A1

115

5 <210> 102
 <211> 107
 <212> PRT
 <213> Mus musculus

10 <220>
 <221> MISC_FEATURE
 <223> Murine SC17.71 VL

<400> 102

15 Asp Ile Gln Met Thr Gln Ser Pro Ala Ser Leu Ser Ala Ser Val Gly
 1 5 10 15

Glu Thr Val Thr Ile Ala Cys Arg Ala Ser Gly Asn Ile His Asn Tyr
 20 20 25 30

Leu Thr Trp Tyr Gln Gln Arg Gln Gly Lys Ser Pro Gln Leu Leu Val
 35 40 45

25 Tyr Asn Ala Lys Thr Leu Ala Val Gly Val Pro Ser Arg Phe Ser Gly
 50 55 60

30 Ser Gly Ser Gly Thr Gln Tyr Ser Leu Lys Ile Asn Ser Leu Gln Pro
 65 70 75 80

Glu Asp Phe Gly Ser Tyr Tyr Cys Gln His Phe Trp Asn Thr Pro Pro
 35 85 90 95

Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys
 100 105

40 <210> 103
 <211> 119
 <212> PRT
 <213> Mus musculus

45 <220>
 <221> MISC_FEATURE
 <223> Murine SC17.71 VH

50 <400> 103

Glu Val Lys Leu Glu Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
 1 5 10 15

55 Ser Met Lys Leu Ser Cys Val Ala Ser Gly Ile Ile Phe Ser Asn Tyr
 20 25 30

EP 3 338 793 A1

Trp Met Asn Trp Val Arg Gln Ser Pro Glu Lys Gly Leu Glu Trp Val
35 40 45

5 Ala Glu Ile Arg Leu Lys Ser Asn Asn Tyr Ser Thr His Tyr Ala Glu
50 55 60

10 Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asp Ser Lys Ser Ser
65 70 75 80

15 Val Tyr Leu Gln Met Asn Asn Leu Arg Ala Glu Asp Thr Gly Ile Tyr
85 90 95

20 Tyr Cys Thr Arg His Tyr Tyr Tyr Ala Met Asp Tyr Trp Gly Gln Gly
100 105 110

Thr Ser Val Thr Val Ser Ser
115

<210> 104
<211> 107
<212> PRT
<213> Mus musculus

<220>
<221> MISC_FEATURE
<223> Murine SC17.72 VL

<400> 104

35 Asp Ile Gln Met Thr Gln Thr Thr Ser Ser Leu Ser Ala Ser Leu Gly
1 5 10 15

40 Asp Arg Val Thr Ile Ser Cys Ser Ala Ser Gln Gly Ile Ser Asn Tyr
20 25 30

45 Leu Asn Trp Tyr Gln Gln Lys Pro Asp Gly Thr Val Lys Leu Leu Ile
35 40 45

50 Tyr Tyr Thr Ser Ser Leu His Ser Gly Val Pro Ser Lys Phe Ser Gly
50 55 60

Ser Gly Ser Gly Thr Asp Tyr Ser Leu Thr Ile Ser Asn Leu Glu Pro
65 70 75 80

Glu Asp Ile Ala Thr Tyr Tyr Cys Gln Gln Tyr Ser Lys Leu Pro Tyr
85 90 95

55 Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys
100 105

EP 3 338 793 A1

<210> 105
<211> 119
<212> PRT
<213> Mus musculus

5

<220>
<221> MISC_FEATURE
<223> Murine SC17.72 VH

10

<400> 105

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Lys Pro Gly Gly
1 5 10 15

15

Ser Leu Lys Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
20 25 30

20

Gly Met Ser Trp Val Arg Gln Thr Pro Glu Lys Arg Leu Glu Trp Val
35 40 45

25

Ala Ala Ile Asn Ser Asn Gly Gly Ser Thr Tyr Tyr Pro Asp Thr Val
50 55 60

Lys Gly Arg Leu Thr Ile Ser Arg Asp Asn Gly Lys Asn Thr Leu Tyr
65 70 75 80

30

Leu Gln Met Ser Ser Leu Arg Ser Glu Asp Thr Ala Leu Tyr Tyr Cys
85 90 95

35

Val Arg Asp Asp Gly Tyr Tyr Val Phe Phe Ala Tyr Trp Gly Gln Gly
100 105 110

Thr Leu Val Thr Val Ser Ala
115

40

<210> 106
<211> 107
<212> PRT
<213> Mus musculus

45

<220>
<221> MISC_FEATURE
<223> Murine SC17.74 VL

50

<400> 106

Asp Ile Gln Met Thr Gln Ser Ser Ser Tyr Leu Ser Val Ser Leu Gly
1 5 10 15

55

Gly Arg Val Thr Ile Thr Cys Lys Ala Ser Asp His Ile Asn Asn Trp
20 25 30

EP 3 338 793 A1

Leu Ala Trp Tyr Gln Gln Lys Pro Gly Asn Ala Pro Arg Leu Leu Ile
 35 40 45
 5 Ser Gly Ala Thr Ser Leu Glu Thr Gly Val Pro Ser Arg Phe Ser Gly
 50 55 60
 10 Ser Gly Ser Gly Lys Asp Tyr Thr Leu Ser Ile Thr Ser Leu Gln Thr
 65 70 75 80
 15 Glu Asp Val Ala Thr Tyr Tyr Cys Gln Gln Tyr Trp Ser Thr Pro Pro
 85 90 95
 20 Thr Phe Gly Ala Gly Thr Lys Leu Glu Leu Lys
 100 105
 <210> 107
 <211> 119
 <212> PRT
 <213> Mus musculus
 25 <220>
 <221> MISC_FEATURE
 <223> Murine SC17.74 VH
 <400> 107
 30 Gln Val Gln Leu Lys Gln Ser Gly Pro Gly Leu Val Ala Pro Ser Gln
 1 5 10 15
 35 Ser Leu Ser Ile Thr Cys Thr Val Ser Gly Phe Ser Leu Thr Ser Tyr
 20 25 30
 40 Gly Val Asp Trp Val Arg Gln Ser Pro Gly Lys Gly Leu Glu Trp Leu
 35 40 45
 45 Gly Val Ile Trp Gly Gly Gly Ser Thr Asn Tyr Asn Ser Ala Leu Lys
 50 55 60
 50 Ser Arg Leu Ser Ile Thr Lys Asp Asn Ser Lys Ser Gln Val Phe Leu
 65 70 75 80
 55 Lys Met Asn Ser Leu Gln Thr Asp Asp Thr Ala Met Tyr Tyr Cys Ala
 85 90 95
 Ser Gly Asp Tyr Asp Gly Ser Leu Trp Phe Ala Tyr Trp Gly Gln Gly
 100 105 110
 Thr Leu Val Thr Val Ser Ala

EP 3 338 793 A1

115

5 <210> 108
 <211> 112
 <212> PRT
 <213> Mus musculus

10 <220>
 <221> MISC_FEATURE
 <223> Murine SC17.76 VL

<400> 108

15 Asp Ile Val Ile Thr Gln Asp Glu Leu Ser Asn Pro Val Thr Ser Gly
 1 5 10 15

Glu Ser Val Ser Ile Ser Cys Arg Ser Ser Lys Ser Leu Leu Tyr Lys
 20 20 25 30

Asp Gly Lys Thr Tyr Leu Asn Trp Phe Leu Gln Arg Pro Gly Gln Ser
 35 40 45

25 Pro Gln Leu Leu Ile Tyr Leu Met Ser Thr Arg Ala Ser Gly Val Ser
 50 55 60

30 Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Glu Ile
 65 70 75 80

Ser Arg Val Lys Ala Glu Asp Val Gly Val Tyr Tyr Cys Gln Gln Leu
 85 90 95

35 Val Glu Tyr Pro Arg Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys
 100 105 110

40 <210> 109
 <211> 113
 <212> PRT
 <213> Mus musculus

45 <220>
 <221> MISC_FEATURE
 <223> Murine SC17.76 VH

50 <400> 109

Glu Val Gln Leu Val Glu Ser Gly Gly Asp Leu Val Lys Pro Gly Gly
 1 5 10 15

55 Ser Leu Lys Leu Ser Cys Val Ala Ser Gly Phe Thr Phe Ser Ser Tyr
 20 25 30

EP 3 338 793 A1

Gly Met Ser Trp Val Arg Gln Thr Pro Asp Lys Arg Leu Glu Trp Val
 35 40 45
 5 Ala Thr Ile Ser Ser Gly Gly Thr Phe Thr Tyr Tyr Pro Asp Ser Val
 50 55 60
 10 Lys Gly Arg Phe Thr Val Ser Arg Asp Asn Ala Lys Asn Thr Leu Tyr
 65 70 75 80
 15 Leu Gln Met Ser Ser Leu Lys Ser Glu Asp Thr Ala Met Tyr Tyr Cys
 85 90 95
 20 Ser Arg His Gly Trp Gly Trp Gly Gln Gly Thr Leu Val Thr Val Ser
 100 105 110
 25 Ala
 30 <210> 110
 <211> 107
 <212> PRT
 <213> Mus musculus
 35 <220>
 <221> MISC_FEATURE
 <223> Murine SC17.77 VL
 40 <400> 110
 45 Asp Ile Gln Met Thr Gln Ser Pro Ala Ser Leu Ser Ala Ser Val Gly
 1 5 10 15
 50 Glu Thr Val Thr Ile Thr Cys Arg Ala Ser Gly Asn Ile His Asn Tyr
 20 25 30
 55 Leu Ala Trp Tyr Gln Gln Lys Gln Gly Lys Ser Pro Gln Leu Leu Val
 35 40 45
 60 Tyr Asn Ala Lys Ala Leu Ala Asp Gly Val Pro Ser Arg Phe Ser Gly
 50 55 60
 65 Ser Gly Ser Gly Thr Gln Tyr Ser Leu Lys Ile Asn Ser Leu Gln Pro
 65 70 75 80
 70 Glu Asp Phe Gly Ser Tyr Tyr Cys Gln His Phe Trp Ser Ile Pro Pro
 85 90 95
 75 Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys
 100 105

EP 3 338 793 A1

<210> 111
<211> 119
<212> PRT
<213> Mus musculus

<220>
<221> MISC_FEATURE
<223> Murine SC17.77 VH

<400> 111

Glu Val Lys Leu Glu Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1 5 10 15

Ser Met Lys Leu Ser Cys Val Ala Ser Gly Phe Thr Phe Ser Asn Tyr
20 25 30

Trp Met Asn Trp Val Arg Gln Ser Pro Glu Lys Gly Leu Glu Trp Val
35 40 45

Ala Glu Ile Arg Leu Lys Ser Asn Asn Tyr Ala Thr His Tyr Ala Glu
50 55 60

Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asp Ser Lys Ser Ser
65 70 75 80

Val Tyr Leu Gln Met Asn Asn Leu Arg Val Glu Asp Thr Ala Ile Tyr
85 90 95

Tyr Cys Thr Arg His Tyr Asp Tyr Ala Met Asp Tyr Trp Gly Gln Gly
100 105 110

Thr Ser Val Thr Val Ser Ser
115

<210> 112
<211> 107
<212> PRT
<213> Mus musculus

<220>
<221> MISC_FEATURE
<223> Murine SC17.79 VL

<400> 112

Asp Ile Gln Met Thr Gln Ser Pro Ala Ser Leu Ser Ala Ser Val Gly
1 5 10 15

Glu Thr Val Thr Ile Thr Cys Arg Ala Ser Gly Asn Ile His Asn Tyr
20 25 30

EP 3 338 793 A1

Leu Ala Trp Tyr Gln Gln Lys Gln Gly Lys Ser Pro Gln Leu Leu Val
35 40 45

5 Tyr Asn Ala Lys Thr Leu Ala Asp Gly Val Pro Ser Arg Phe Ser Gly
50 55 60

10 Ser Gly Ser Gly Thr Gln Tyr Ser Leu Arg Ile Asn Ser Leu Gln Pro
65 70 75 80

Glu Asp Phe Gly Ser Tyr Tyr Cys Gln His Phe Trp Ser Thr Pro Pro
85 90 95

15 Thr Phe Gly Gly Gly Thr Lys Val Glu Ile Lys
100 105

20 <210> 113
<211> 119
<212> PRT
<213> Mus musculus

25 <220>
<221> MISC_FEATURE
<223> Murine SC17.79 VH

<400> 113

30 Glu Val Lys Leu Glu Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1 5 10 15

35 Ser Met Lys Leu Ser Cys Val Ala Ser Gly Phe Thr Phe Ser Asp Tyr
20 25 30

Trp Met Asn Trp Val Arg Gln Ser Pro Glu Lys Gly Leu Glu Leu Val
35 40 45

40 Ala Glu Ile Arg Leu Ile Ser Asn Asn Tyr Ala Thr His Tyr Ala Glu
50 55 60

45 Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asp Ser Lys Ser Ser
65 70 75 80

50 Val Tyr Leu Gln Met Asn Asn Leu Arg Ala Glu Asp Thr Gly Ile Tyr
85 90 95

Tyr Cys Thr Arg His Tyr Tyr Tyr Ala Leu Asp Tyr Trp Gly Gln Gly
100 105 110

55 Thr Ser Val Thr Val Ser Ser

115

5 <210> 114
 <211> 113
 <212> PRT
 <213> Mus musculus

10 <220>
 <221> MISC_FEATURE
 <223> Murine SC17.81 VL
 <400> 114

15 Asp Ile Val Met Ser Gln Ser Pro Ser Ser Leu Thr Val Ser Val Gly
 1 5 10 15

20 Glu Lys Val Thr Leu Ser Cys Lys Ser Ser Gln Ser Leu Leu Tyr Ser
 20 25 30

25 Thr Asn Gln Lys Ile Tyr Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln
 35 40 45

30 Ser Pro Lys Leu Leu Ile Tyr Trp Ala Ser Thr Arg Glu Ser Gly Val
 50 55 60

35 Pro Asp Arg Phe Thr Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Ala
 65 70 75 80

40 Ile Ser Asn Val Lys Ala Glu Asp Leu Ala Val Tyr Tyr Cys Gln Gln
 85 90 95

45 Tyr Tyr Ser Tyr Pro Tyr Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile
 100 105 110

50 Lys

55 <210> 115
 <211> 113
 <212> PRT
 <213> Mus musculus

60 <220>
 <221> MISC_FEATURE
 <223> Murine SC17.81 VH
 <400> 115

65 Glu Val Gln Leu Gln Gln Ser Gly Ala Glu Leu Val Lys Pro Gly Ala
 1 5 10 15

EP 3 338 793 A1

Ser Val Lys Leu Ser Cys Thr Ala Ser Gly Phe Asn Ile Asn Asp Thr
20 25 30

5 Tyr Tyr His Trp Leu Lys Gln Arg Pro Glu Gln Gly Leu Glu Trp Ile
35 40 45

10 Gly Arg Ile Asp Pro Ala Asn Val Asn Thr Lys Tyr Asp Pro Lys Phe
50 55 60

Gln Gly Lys Ala Thr Leu Thr Ala Asp Thr Ser Ser Asn Thr Ala Tyr
65 70 75 80

15 Leu Gln Leu Ser Ser Leu Thr Ser Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95

20 Gly Arg Gly Asn Ala Tyr Trp Gly Gln Gly Thr Leu Val Thr Val Ser
100 105 110

Ala

25

<210> 116
<211> 104
<212> PRT
<213> Mus musculus

30

<220>
<221> MISC_FEATURE
<223> Murine SC17.82 VL

35

<400> 116

Gln Ile Val Leu Thr Gln Ser Pro Ala Ile Met Ser Ala Ser Leu Gly
1 5 10 15

40

Glu Glu Ile Thr Leu Thr Cys Ser Ala Ser Ser Ser Val Ser Tyr Met
20 25 30

45

His Trp Tyr Gln Gln Lys Ser Gly Thr Ser Pro Lys Leu Leu Ile Tyr
35 40 45

Ser Thr Ser Asn Leu Ala Ser Gly Val Pro Ser Arg Phe Ser Gly Ser
50 55 60

50

Gly Ser Gly Thr Phe Tyr Ser Leu Thr Ile Ser Ser Val Glu Ala Glu
65 70 75 80

55

Asp Ala Ala Asp Tyr Tyr Cys His Gln Trp Ser Ser Phe Thr Phe Gly
85 90 95

EP 3 338 793 A1

Ser Gly Thr Lys Leu Glu Ile Lys
100

5 <210> 117
<211> 120
<212> PRT
<213> Mus musculus

10 <220>
<221> MISC_FEATURE
<223> Murine SC17.82 VH

<400> 117
15 Glu Val Gln Leu Gln Gln Ser Gly Pro Glu Leu Val Lys Pro Gly Ala
1 5 10 15

20 Ser Val Lys Met Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asp Ser
20 25 30

25 Tyr Met Asn Trp Val Lys Gln Ser His Gly Lys Ser Leu Glu Trp Ile
35 40 45

Gly Arg Val Asn Pro Asn Asn Gly Gly Ala Ser Tyr Asn His Lys Phe
50 55 60

30 Lys Gly Lys Ala Thr Leu Thr Val Asp Lys Ser Leu Ser Thr Ala Tyr
65 70 75 80

35 Met Arg Leu Asn Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys
85 90 95

Ser Arg Ser Gly Asp Leu Tyr Tyr Tyr Ala Met Asp Tyr Trp Gly Gln
100 105 110

40 Gly Thr Ser Val Thr Val Ser Ser
115 120

45 <210> 118
<211> 106
<212> PRT
<213> Mus musculus

50 <220>
<221> MISC_FEATURE
<223> Murine SC17.84 VL

<400> 118
55 Gln Ile Val Leu Thr Gln Ser Pro Ala Ile Met Ser Ala Ser Pro Gly
1 5 10 15

EP 3 338 793 A1

Glu Lys Val Thr Met Thr Cys Ser Ala Ser Ser Ser Ile Ser Tyr Met
20 25 30

5 His Trp Tyr Gln Gln Lys Ser Gly Thr Ser Pro Lys Arg Trp Ile Tyr
35 40 45

10 Asp Thr Ser Lys Leu Ala Ser Gly Val Pro Ala Arg Phe Ser Gly Ser
50 55 60

Gly Ser Gly Thr Ser Tyr Ser Leu Thr Ile Ser Asn Met Glu Ala Glu
65 70 75 80

15 Asp Ala Ala Thr Tyr Tyr Cys Gln Gln Trp Ser Ser Thr Pro Pro Thr
85 90 95

20 Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys
100 105

<210> 119
<211> 117
25 <212> PRT
<213> Mus musculus

<220>
30 <221> MISC_FEATURE
<223> Murine SC17.84 VH

<400> 119

35 Glu Val Gln Leu Gln Gln Ser Gly Pro Glu Leu Met Lys Pro Gly Ala
1 5 10 15

Ser Val Lys Met Ser Cys Lys Ala Ser Gly Tyr Ile Phe Thr Asp Tyr
20 25 30

40 Asn Met His Trp Val Lys Gln Asn Gln Gly Lys Ser Leu Glu Trp Ile
35 40 45

45 Gly Glu Val Asn Pro Asn Thr Gly Gly Ile Gly Tyr Asn Gln Lys Phe
50 55 60

Lys Gly Lys Ala Thr Leu Thr Val Asp Lys Ser Ser Ser Thr Ala Tyr
50 65 70 75 80

Met Asp Leu Arg Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys
85 90 95

55 Ala Arg Asp Gly Asn Tyr Cys Phe Asp Tyr Trp Gly Gln Gly Thr Thr

EP 3 338 793 A1

	100	105	110
5	Leu Thr Val Ser Ser		
	115		
	<210> 120		
	<211> 111		
10	<212> PRT		
	<213> Mus musculus		
	<220>		
	<221> MISC_FEATURE		
15	<223> Murine SC17.85 VL		
	<400> 120		
20	Asp Ile Val Met Thr Gln Ala Ala Phe Ser Asn Pro Val Thr Leu Gly		
	1 5 10 15		
	Thr Ser Ala Ser Ile Ser Cys Arg Ser Ser Lys Ser Leu Leu His Ser		
	20 25 30		
25	Asn Gly Ile Thr Tyr Leu Tyr Trp Tyr Leu Gln Lys Pro Gly Gln Ser		
	35 40 45		
30	Pro Gln Leu Leu Ile Tyr Gln Met Ser Asn Leu Ala Ser Gly Val Pro		
	50 55 60		
35	Glu Arg Phe Ser Ser Ser Gly Ser Gly Ser Asp Phe Thr Leu Arg Ile		
	65 70 75 80		
	Ser Arg Val Glu Ala Glu Asp Val Gly Val Tyr Tyr Cys Ala Gln Asn		
	85 90 95		
40	Leu Glu His Pro Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys		
	100 105 110		
45	<210> 121		
	<211> 120		
	<212> PRT		
	<213> Mus musculus		
50	<220>		
	<221> MISC_FEATURE		
	<223> Murine SC17.85 VH		
	<400> 121		
55	Glu Val Gln Leu Val Glu Ser Gly Gly Asp Leu Val Lys Pro Gly Gly		
	1 5 10 15		

EP 3 338 793 A1

Ser Leu Lys Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Asn Tyr
20 25 30

5 Gly Met Ser Trp Val Arg Gln Thr Pro Asp Lys Arg Leu Glu Trp Val
35 40 45

10 Ala Thr Ile Ser Thr Gly Gly Thr Tyr Thr Tyr Tyr Pro Asp Ser Val
50 55 60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Thr Leu Tyr
65 70 75 80

15 Leu Gln Met Ser Ser Leu Lys Ser Glu Asp Thr Ala Met Tyr Tyr Cys
85 90 95

20 Val Gly Gln Ser Tyr Ser Asp Tyr Val Ser Phe Ala Tyr Trp Gly Gln
100 105 110

Gly Thr Gln Val Thr Val Ser Ala
115 120

25

<210> 122
<211> 112
<212> PRT
<213> Mus musculus

30

<220>
<221> MISC_FEATURE
<223> Murine SC17.87 VL

35

<400> 122

Asp Val Val Met Thr Gln Thr Pro Leu Ser Leu Pro Val Ser Leu Gly
1 5 10 15

40 Asp Gln Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Leu Val His Ser
20 25 30

45 Asn Gly Asn Thr Tyr Leu His Trp Tyr Leu Gln Lys Pro Gly Gln Ser
35 40 45

Pro Lys Leu Leu Ile Ser Lys Val Ser Asn Arg Phe Ser Gly Val Pro
50 55 60

Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile
65 70 75 80

55 Ser Arg Val Glu Ala Glu Asp Leu Gly Val Tyr Phe Cys Ser Gln Ser
85 90 95

EP 3 338 793 A1

Thr His Val Pro Pro Met Phe Gly Gly Gly Thr Arg Leu Glu Ile Lys
100 105 110

5
<210> 123
<211> 115
<212> PRT
<213> Mus musculus

10
<220>
<221> MISC_FEATURE
<223> Murine SC17.87 VH

<400> 123

15
Glu Val Gln Leu Gln Gln Ser Gly Ala Glu Leu Leu Lys Pro Gly Ala
1 5 10 15

20
Ser Val Lys Leu Ser Cys Thr Ala Ser Gly Leu Asn Ile Lys Asp Tyr
20 25 30

Tyr Ile His Trp Val Tyr Gln Arg Pro Glu Gln Gly Leu Glu Trp Ile
35 40 45

25
Gly Arg Ile Asp Pro Glu Ser Asp Asn Thr Leu Tyr Asp Pro Lys Phe
50 55 60

30
Gln Gly Lys Ala Ser Ile Thr Ala Asp Thr Ser Ser Asn Thr Ala Tyr
65 70 75 80

35
Leu Gln Leu Ser Ser Leu Thr Ser Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95

Thr Thr Asn Thr Pro Phe Ala Tyr Trp Gly Gln Gly Thr Leu Val Thr
100 105 110

40
Val Ser Thr
115

45
<210> 124
<211> 112
<212> PRT
<213> Mus musculus

50
<220>
<221> MISC_FEATURE
<223> Murine SC17.89 VL

<400> 124

55
Asp Val Leu Met Thr Gln Thr Pro Leu Ser Leu Pro Val Ser Leu Gly
1 5 10 15

EP 3 338 793 A1

Asp Gln Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Ile Val His Ser
 20 25 30

5 Asn Gly Asn Thr Tyr Leu Glu Trp Tyr Leu Gln Lys Pro Gly Gln Ser
 35 40 45

10 Pro Lys Leu Leu Ile Tyr Lys Val Ser Asn Arg Phe Ser Gly Val Pro
 50 55 60

15 Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile
 65 70 75 80

20 Ser Arg Val Glu Ala Glu Asp Leu Gly Val Tyr Tyr Cys Phe Gln Gly
 85 90 95

25 Ser His Val Pro Phe Thr Phe Gly Ser Gly Thr Lys Leu Glu Ile Lys
 100 105 110

30 <210> 125
 <211> 120
 <212> PRT
 <213> Mus musculus

35 <220>
 <221> MISC_FEATURE
 <223> Murine SC17.89 VH
 <400> 125

40 Gln Val Gln Leu Gln Gln Ser Gly Ala Glu Leu Val Arg Pro Gly Thr
 1 5 10 15

45 Ser Val Lys Val Ser Cys Lys Thr Ser Gly Tyr Ala Phe Thr Asn Tyr
 20 25 30

50 Leu Ile Glu Trp Val Lys Gln Arg Pro Gly Gln Gly Leu Glu Trp Ile
 35 40 45

55 Gly Val Ile Asn Pro Gly Ser Gly Gly Thr Asn Tyr Asn Glu Lys Phe
 50 55 60

60 Lys Val Lys Ala Thr Leu Thr Ala Asp Lys Ser Ser Ser Thr Ala Tyr
 65 70 75 80

65 Met Gln Leu Thr Ser Leu Thr Ser Asp Asp Ser Ala Val Tyr Phe Cys
 85 90 95

70 Thr Arg Arg Asp Gly Tyr Phe Phe Pro Trp Phe Ala Tyr Trp Gly Gln

EP 3 338 793 A1

	100	105	110
5	Gly Thr Leu Val Thr Val Ser Ala 115 120		
10	<210> 126 <211> 113 <212> PRT <213> Mus musculus		
15	<220> <221> MISC_FEATURE <223> Murine SC17.90 VL		
20	<400> 126 Asp Ile Val Met Ser Gln Ser Pro Ser Ser Leu Ala Val Ser Val Gly 1 5 10 15		
25	Glu Lys Val Thr Met Ser Cys Lys Ser Ser Gln Ser Leu Leu Tyr Ser 20 25 30		
30	Ser Asn Gln Lys Asn Tyr Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln 35 40 45		
35	Ser Pro Lys Leu Leu Ile Tyr Trp Ala Ser Thr Arg Lys Ser Gly Val 50 55 60		
40	Pro Asp Arg Phe Thr Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr 65 70 75 80		
45	Ile Ser Ser Val Lys Ala Glu Asp Leu Ala Val Tyr Tyr Cys His Gln 85 90 95		
50	Tyr Tyr Ser Tyr Pro Leu Thr Phe Ala Ala Gly Thr Lys Leu Glu Leu 100 105 110		
55	Lys <210> 127 <211> 115 <212> PRT <213> Mus musculus <220> <221> MISC_FEATURE <223> Murine SC17.90 VH <400> 127		

EP 3 338 793 A1

Gln Val Gln Leu Gln Gln Pro Gly Ser Val Leu Val Arg Pro Gly Ala
 1 5 10 15
 5 Ser Val Lys Leu Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Ser Tyr
 20 25 30
 10 Trp Met His Trp Val Lys Gln Arg Pro Gly Gln Gly Leu Glu Trp Ile
 35 40 45
 Gly Glu Ile His Pro Asn Asn Gly Ser Thr Asn Tyr Asn Glu Lys Phe
 50 55 60
 15 Lys Gly Lys Ala Thr Leu Thr Val Asp Thr Ser Ser Ser Thr Ala Tyr
 65 70 75 80
 20 Val Asp Leu Ser Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys
 85 90 95
 25 Ala Arg Trp Thr Leu Phe Thr Tyr Trp Gly Gln Gly Thr Leu Val Thr
 100 105 110
 Val Ser Ala
 115
 30 <210> 128
 <211> 112
 <212> PRT
 <213> Mus musculus
 35 <220>
 <221> MISC_FEATURE
 <223> Murine SC17.91 VL
 40 <400> 128
 Asp Val Val Met Thr Gln Thr Pro Leu Ser Leu Pro Val Ser Leu Gly
 1 5 10 15
 45 Asp Gln Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Leu Val His Ser
 20 25 30
 Asn Gly Asn Thr Tyr Leu Leu Trp Tyr Leu Gln Lys Pro Gly Gln Ser
 35 40 45
 50 Pro Lys Leu Leu Ile Tyr Lys Val Ser Asn Arg Phe Ser Gly Val Pro
 50 55 60
 55 Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Ala Leu Lys Ile
 65 70 75 80

EP 3 338 793 A1

Ser Arg Val Glu Ala Glu Asp Leu Gly Val Tyr Phe Cys Ser Gln Ser
85 90 95

5 Thr His Val Pro Trp Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys
100 105 110

10 <210> 129
<211> 117
<212> PRT
<213> Mus musculus

15 <220>
<221> MISC_FEATURE
<223> Murine SC17.91 VH

<400> 129

20 Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Lys Pro Gly Gly
1 5 10 15

25 Ser Arg Lys Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Asp Tyr
20 25 30

Gly Met His Trp Val Arg Gln Ala Pro Glu Lys Gly Leu Glu Trp Val
35 40 45

30 Ala Tyr Ile Ser Arg Gly Ser Ser Thr Ile His Tyr Ala Asp Thr Val
50 55 60

35 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Thr Leu Phe
65 70 75 80

Leu Gln Met Thr Ser Leu Arg Ser Glu Asp Thr Ala Met Tyr Tyr Cys
85 90 95

40 Ala Arg Pro Phe Asn Trp Tyr Phe Asp Val Trp Gly Ala Gly Thr Thr
100 105 110

45 Val Thr Val Ser Ser
115

50 <210> 130
<211> 113
<212> PRT
<213> Mus musculus

55 <220>
<221> MISC_FEATURE
<223> Murine SC17.93 VL

EP 3 338 793 A1

<400> 130

Asp Ile Val Met Ser Gln Ser Pro Ser Ser Leu Ala Val Ser Val Gly
1 5 10 15

Glu Lys Val Thr Met Thr Cys Lys Ser Ser Gln Ser Leu Leu Tyr Ser
20 25 30

Ser Asn Gln Lys Asn Tyr Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln
35 40 45

Ser Pro Lys Leu Leu Ile Tyr Trp Ala Ser Thr Arg Glu Ser Gly Val
50 55 60

Pro Asp Arg Phe Ile Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr
65 70 75 80

Ile Ser Ser Val Lys Ala Glu Asp Leu Ala Ile Tyr Tyr Cys Gln Gln
85 90 95

Tyr Tyr Arg Tyr Pro Leu Thr Phe Gly Ala Gly Thr Lys Leu Glu Leu
100 105 110

Lys

<210> 131

<211> 121

<212> PRT

<213> Mus musculus

<220>

<221> MISC_FEATURE

<223> Murine SC17.93 VH

<400> 131

Gln Val Gln Leu Gln Gln Pro Gly Ala Glu Leu Val Lys Pro Gly Ala
1 5 10 15

Ser Val Met Leu Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Ser Tyr
20 25 30

Trp Val His Trp Val Lys Gln Arg Pro Gly Gln Gly Leu Glu Trp Ile
35 40 45

Gly Val Ile Asn Pro Arg Asn Gly Arg Asn Asn Tyr Asn Glu Lys Phe
50 55 60

Lys Thr Lys Ala Thr Leu Thr Val Asp Lys Ser Ser Ser Thr Ala Tyr

EP 3 338 793 A1

	65		70		75		80									
5	Met	Gln	Leu	Ser	Ser	Pro	Thr	Ser	Glu	Asp	Ser	Ala	Val	Tyr	Tyr	Cys
				85					90					95		
10	Ala	Arg	Glu	Asp	Tyr	Asp	Gly	Gly	Asp	Tyr	Ala	Met	Asp	Tyr	Trp	Gly
			100						105					110		
15	Gln	Gly	Thr	Ser	Val	Thr	Val	Ser	Ser							
			115					120								
20	<210>	132														
	<211>	107														
	<212>	PRT														
	<213>	Mus musculus														
25	<220>															
	<221>	MISC_FEATURE														
	<223>	Murine SC17.95 VL														
	<400>	132														
30	Asp	Ile	Gln	Met	Thr	Gln	Thr	Thr	Ser	Ser	Leu	Ser	Ala	Ser	Leu	Gly
	1				5					10					15	
35	Asp	Arg	Val	Thr	Ile	Ser	Cys	Ser	Ala	Ser	Gln	Gly	Ile	Asn	Asn	Tyr
			20						25					30		
40	Leu	Asn	Trp	Tyr	Gln	Gln	Lys	Pro	Asp	Gly	Thr	Val	Thr	Leu	Leu	Ile
			35					40					45			
45	Tyr	Tyr	Thr	Ser	Ser	Leu	His	Ser	Gly	Val	Pro	Ser	Arg	Phe	Ser	Gly
		50					55					60				
50	Ser	Gly	Ser	Gly	Thr	Asp	Tyr	Ser	Leu	Thr	Ile	Ser	Asn	Leu	Glu	Pro
	65				70						75				80	
55	Glu	Asp	Ile	Ala	Thr	Tyr	Tyr	Cys	Gln	Gln	Tyr	Ser	Lys	Leu	Pro	Trp
				85					90					95		
60	Thr	Phe	Gly	Gly	Gly	Thr	Lys	Leu	Glu	Ile	Lys					
			100						105							
65	<210>	133														
	<211>	121														
	<212>	PRT														
	<213>	Mus musculus														
70	<220>															
	<221>	MISC_FEATURE														

EP 3 338 793 A1

<223> Murine SC17.95 VH

<400> 133

5 Glu Val Glu Leu Gln Gln Ser Gly Pro Glu Leu Val Lys Pro Gly Ala
1 5 10 15

10 Ser Val Lys Ile Ser Cys Lys Thr Ser Gly Asn Thr Tyr Thr Glu Tyr
20 25 30

Thr Met Gln Trp Val Lys Leu Ser His Gly Lys Ser Leu Glu Trp Ile
35 40 45

15 Gly Gly Ile Asn Pro Asn Asn Gly Ile Thr Ser Tyr Asn Gln Lys Phe
50 55 60

20 Lys Gly Lys Ala Thr Leu Thr Val Asp Lys Ser Ser Ser Thr Ala Tyr
65 70 75 80

25 Met Glu Leu Arg Ser Leu Lys Ser Glu Asp Ser Ala Val Tyr Tyr Cys
85 90 95

Ala Arg Ala Gly Leu Gly Asn Tyr Val Trp Ala Met Asp Tyr Trp Gly
100 105 110

30 Gln Gly Ala Ser Val Thr Val Ser Ser
115 120

<210> 134

35 <211> 112

<212> PRT

<213> Mus musculus

<220>

40 <221> MISC_FEATURE

<223> Murine SC17.97 VL

<400> 134

45 Asp Val Val Met Thr Gln Thr Pro Leu Ser Leu Pro Val Ser Leu Gly
1 5 10 15

50 Asp Gln Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Leu Val His Asn
20 25 30

Asn Gly Asn Thr Tyr Leu His Trp Tyr Leu Gln Lys Pro Gly Gln Ser
35 40 45

55 Pro Asn Leu Leu Ile Tyr Lys Val Ser Asn Arg Phe Ser Gly Val Pro
50 55 60

EP 3 338 793 A1

	Asp	Arg	Phe	Ser	Gly	Ser	Gly	Ser	Gly	Thr	Asp	Phe	Thr	Leu	Lys	Ile
	65					70					75				80	
5	Ser	Ile	Val	Glu	Ala	Glu	Asp	Leu	Gly	Leu	Tyr	Phe	Cys	Ser	Gln	Ser
				85						90					95	
10	Thr	His	Val	Pro	Arg	Thr	Phe	Gly	Gly	Gly	Thr	Lys	Leu	Glu	Ile	Lys
				100					105					110		
15	<210>	135														
	<211>	119														
	<212>	PRT														
	<213>	Mus musculus														
20	<220>															
	<221>	MISC_FEATURE														
	<223>	Murine SC17.97 VH														
	<400>	135														
25	Gln	Val	Gln	Leu	Pro	Gln	Ser	Gly	Ala	Glu	Leu	Ala	Lys	Pro	Gly	Ala
	1				5					10					15	
30	Ser	Val	Lys	Ile	Ser	Cys	Lys	Ala	Ser	Gly	Phe	Thr	Phe	Thr	Ser	Tyr
				20					25					30		
35	Trp	Met	His	Trp	Val	Lys	Gln	Arg	Pro	Gly	Gln	Gly	Leu	Glu	Trp	Ile
			35					40					45			
40	Gly	Tyr	Ile	Asn	Pro	Ser	Thr	Asp	Tyr	Thr	Glu	Tyr	Asn	Gln	Lys	Phe
	50						55					60				
45	Lys	Asp	Lys	Ala	Thr	Leu	Thr	Ala	Asp	Lys	Ser	Ser	Ser	Thr	Ala	Tyr
	65					70					75				80	
50	Met	Gln	Leu	Gly	Ser	Leu	Thr	Ser	Glu	Asp	Ser	Ala	Val	Tyr	Tyr	Cys
					85					90					95	
55	Ala	Arg	Ser	Ser	Tyr	Gly	Ser	Ser	Pro	Phe	Asp	Tyr	Trp	Gly	Gln	Gly
					100				105					110		
60	Ser	Thr	Leu	Thr	Val	Ser	Ser									
					115											
65	<210>	136														
	<211>	113														
	<212>	PRT														
	<213>	Mus musculus														

EP 3 338 793 A1

<220>
 <221> MISC_FEATURE
 <223> Murine SC17.99 VL

<400> 136

Asp Ile Val Met Ser Gln Ser Pro Ser Ser Leu Ala Val Ser Val Gly
 1 5 10 15

Glu Lys Val Thr Met Asn Cys Glu Ser Ser Gln Ser Leu Leu Tyr Ser
 20 25 30

Ser Asn Gln Lys Asn Tyr Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln
 35 40 45

Ser Pro Lys Leu Leu Ile Tyr Trp Ala Ser Thr Arg Asp Ser Gly Val
 50 55 60

Pro Asp Arg Phe Thr Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr
 65 70 75 80

Ile Ser Ser Val Arg Ala Glu Asp Pro Ala Val Tyr Tyr Cys Gln Gln
 85 90 95

Tyr Tyr Ser Tyr Pro Leu Thr Phe Gly Ala Gly Thr Lys Leu Glu Leu
 100 105 110

Arg

<210> 137
 <211> 116
 <212> PRT
 <213> Mus musculus

<220>
 <221> MISC_FEATURE
 <223> Murine SC17.99 VH

<400> 137

Glu Val Lys Leu Glu Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
 1 5 10 15

Ser Met Lys Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Asp Ala
 20 25 30

Trp Met Asp Trp Val Arg Gln Ser Pro Glu Lys Gly Leu Glu Trp Val
 35 40 45

Ala Glu Ile Arg Ser Lys Ala Asn Asn His Ala Thr Tyr Tyr Ala Glu

EP 3 338 793 A1

	50		55		60	
5	Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asp Ser Lys Ser Ser					
	65		70		75	80
	Ala Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Gly Ile Tyr					
		85		90		95
10	Tyr Cys Val Ser Thr Gly Thr Ser Tyr Trp Gly Gln Gly Thr Leu Val					
		100		105		110
15	Thr Val Ser Ala					
		115				
	<210>	138				
	<211>	106				
20	<212>	PRT				
	<213>	Mus musculus				
	<220>					
	<221>	MISC_FEATURE				
25	<223>	Murine SC17.102 VL				
	<400>	138				
30	Gln Ile Val Leu Thr Gln Ser Pro Ala Ile Met Ser Ala Ser Pro Gly					
	1	5		10		15
	Glu Lys Val Thr Met Thr Cys Ser Ala Ser Ser Ser Val Ser Tyr Met					
		20		25		30
35	His Trp Tyr Gln Gln Lys Ser Gly Thr Ser Pro Lys Arg Trp Ile Tyr					
		35		40		45
40	Asp Thr Ser Lys Leu Ala Ser Gly Val Pro Pro Arg Phe Ser Gly Arg					
		50		55		60
	Gly Ser Gly Thr Ser Tyr Ser Leu Thr Ile Ser Ser Met Glu Ala Glu					
	65	70		75		80
45	Asp Ala Ala Thr Tyr Tyr Cys Gln His Trp Ser Ser Asn Pro Pro Thr					
		85		90		95
50	Phe Gly Ala Gly Thr Lys Leu Glu Met Lys					
		100		105		
	<210>	139				
	<211>	116				
55	<212>	PRT				
	<213>	Mus musculus				

EP 3 338 793 A1

<220>
 <221> MISC_FEATURE
 <223> Murine SC17.102 VH

5

<400> 139

Glu Val Gln Leu Gln Gln Ser Gly Pro Glu Leu Met Lys Pro Gly Ala
 1 5 10 15

10

Ser Val Lys Met Ser Cys Lys Ala Ser Gly Asp Thr Phe Thr Asp Tyr
 20 25 30

15

Asn Ile His Trp Val Lys Gln Asn Gln Gly Lys Ser Leu Glu Trp Ile
 35 40 45

Gly Glu Val Asn Pro Asn Ile Gly Gly Ile Gly Tyr Asn Gln Lys Phe
 50 55 60

20

Lys Gly Lys Ala Thr Leu Thr Val Asp Lys Ser Ser Ser Thr Ala Tyr
 65 70 75 80

25

Met Glu Leu Arg Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys
 85 90 95

30

Ala Met Gly Arg Trp Tyr Phe Asp Val Trp Gly Ala Gly Thr Thr Val
 100 105 110

Thr Val Ser Ser
 115

35

<210> 140
 <211> 112
 <212> PRT
 <213> Mus musculus

40

<220>
 <221> MISC_FEATURE
 <223> Murine SC17.114 VL

45

<400> 140

Asp Val Val Met Thr Gln Ser Pro Leu Ser Leu Pro Val Ser Leu Gly
 1 5 10 15

50

Asp Gln Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Leu Val His Ser
 20 25 30

55

Asn Gly Asn Thr Tyr Leu His Trp Tyr Leu Gln Lys Pro Gly Gln Ser
 35 40 45

EP 3 338 793 A1

Pro Lys Leu Leu Ile Tyr Lys Val Ser Ser Arg Phe Ser Gly Val Pro
50 55 60

5 Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile
65 70 75 80

10 Ser Arg Val Glu Ala Glu Asp Leu Gly Val Tyr Phe Cys Ser Gln Ser
85 90 95

Thr His Val Pro Phe Thr Phe Gly Ser Gly Thr Lys Leu Glu Ile Lys
100 105 110

15 <210> 141
<211> 115
<212> PRT
<213> Mus musculus

20 <220>
<221> MISC_FEATURE
<223> Murine SC17.114 VH

25 <400> 141

Glu Val Gln Leu Gln Gln Ser Gly Pro Glu Met Val Lys Pro Gly Ala
1 5 10 15

30 Ser Val Lys Ile Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asp Tyr
20 25 30

35 Tyr Met His Trp Val Lys Gln Ser His Gly Lys Ser Leu Glu Trp Ile
35 40 45

Gly Arg Val Asn Thr Asn Asn Gly Gly Thr Ser Tyr Asp Gln Lys Phe
50 55 60

40 Glu Gly Lys Ala Thr Leu Thr Val Asp Lys Ser Ser Ser Thr Ala Tyr
65 70 75 80

45 Met Glu Leu Asn Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys
85 90 95

50 Val Ile Pro Ala Trp Phe Ala Tyr Trp Gly Gln Gly Thr Leu Val Thr
100 105 110

Val Ser Ala
115

55 <210> 142
<211> 112

EP 3 338 793 A1

<212> PRT
<213> Mus musculus

<220>
<221> MISC_FEATURE
<223> Murine SC17.115 VL

<400> 142

Asp Ile Val Met Thr Gln Ser Pro Leu Ser Leu Pro Val Ser Leu Gly
1 5 10 15

Asp Gln Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Leu Val His Ser
20 25 30

Asn Gly Asn Thr Tyr Leu His Trp Tyr Leu Gln Lys Pro Gly Gln Ser
35 40 45

Pro Lys Leu Leu Ile Tyr Arg Val Ser Asn Arg Phe Ser Gly Val Pro
50 55 60

Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile
65 70 75 80

Ser Arg Val Glu Ala Glu Asp Leu Gly Val Tyr Phe Cys Ser Gln Ser
85 90 95

Thr His Leu Pro Arg Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys
100 105 110

<210> 143
<211> 114
<212> PRT
<213> Mus musculus

<220>
<221> MISC_FEATURE
<223> Murine SC17.115 VH

<400> 143

Gln Val Gln Leu Gln Gln Ser Gly Ser Val Leu Val Arg Pro Gly Ala
1 5 10 15

Ser Val Lys Leu Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Ser Tyr
20 25 30

Trp Met His Trp Val Lys Gln Arg Pro Gly Gln Gly Leu Glu Trp Ile
35 40 45

Gly Glu Ile His Pro Asn Ser Gly Asn Thr Asn Tyr Asn Glu Lys Phe

EP 3 338 793 A1

	50		55		60	
5	Lys Gly Lys Ala Thr Leu Thr Val Asp Thr Ser Ser Ser Thr Ala Tyr	65	70	75	80	
	Val Asp Leu Ser Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys	85	90	95		
10	Ala Gly Gly Asn Tyr Asp Tyr Trp Gly Gln Gly Thr Thr Leu Thr Val	100	105	110		
15	Ser Ser					
	<210> 144					
	<211> 111					
20	<212> PRT					
	<213> Mus musculus					
	<220>					
	<221> MISC_FEATURE					
25	<223> Murine SC17.120 VL					
	<400> 144					
30	Asp Ile Val Leu Thr Gln Ser Pro Ala Ser Leu Ala Val Ser Leu Gly	1	5	10	15	
	Gln Arg Ala Thr Ile Ser Cys Arg Ala Ser Glu Ser Val Asp Ser Tyr	20	25	30		
35	Gly Asn Ser Phe Met His Trp Tyr Gln Gln Lys Pro Gly Gln Pro Pro	35	40	45		
40	Lys Val Leu Ile Tyr Arg Ala Ser Asn Leu Glu Ser Gly Ile Pro Ala	50	55	60		
	Arg Phe Ser Gly Ser Gly Ser Arg Thr Asp Phe Thr Leu Thr Ile Asn	65	70	75	80	
45	Pro Val Glu Asp Glu Asp Val Ala Thr Tyr Tyr Cys Gln Gln Ser Asn	85	90	95		
50	Glu Asp Pro Tyr Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys	100	105	110		
	<210> 145					
	<211> 117					
55	<212> PRT					
	<213> Mus musculus					

EP 3 338 793 A1

<220>
 <221> MISC_FEATURE
 <223> Murine SC17.120 VH

5

<400> 145

Glu Val Gln Leu Glu Gln Ser Gly Thr Val Leu Ala Arg Pro Gly Ala
 1 5 10 15

10

Ser Val Lys Met Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Ser Tyr
 20 25 30

15

Trp Met His Trp Val Lys Gln Arg Pro Gly Gln Gly Leu Glu Trp Ile
 35 40 45

Gly Ala Phe Tyr Pro Gly Asn Ser Gly Thr Tyr Tyr Asn Gln Lys Phe
 50 55 60

20

Lys Asp Lys Ala Lys Leu Thr Ala Val Thr Ser Ala Ser Thr Ala Tyr
 65 70 75 80

25

Met Glu Leu Ser Ser Leu Thr Asn Glu Asp Ser Ala Val Tyr Tyr Cys
 85 90 95

30

Ser Arg Ser Gly Ser Gly Arg Phe Ala Tyr Trp Gly Gln Gly Thr Leu
 100 105 110

Val Thr Val Ser Ala
 115

35

<210> 146
 <211> 106
 <212> PRT
 <213> Mus musculus

40

<220>
 <221> MISC_FEATURE
 <223> Murine SC17.121 VL

45

<400> 146

Gln Ile Val Leu Thr Gln Ser Pro Ala Ile Met Ser Ala Ser Pro Gly
 1 5 10 15

50

Glu Lys Val Thr Met Thr Cys Ser Ala Ser Ser Ser Val Ser Tyr Met
 20 25 30

55

His Trp Tyr Gln Gln Lys Ser Gly Thr Ser Pro Lys Arg Trp Ile Tyr
 35 40 45

EP 3 338 793 A1

Asp Thr Ser Lys Leu Ala Ser Gly Val Pro Ala Arg Phe Ser Gly Ser
50 55 60

5 Gly Ser Gly Thr Ser Tyr Ser Leu Thr Ile Ser Ser Met Glu Thr Glu
65 70 75 80

10 Asp Ala Ala Thr Tyr Tyr Cys Gln Gln Trp Ser Asn Thr Pro Pro Thr
85 90 95

Phe Gly Ser Val Thr Lys Leu Glu Ile Lys
100 105

15 <210> 147
<211> 116
<212> PRT
<213> Mus musculus

20 <220>
<221> MISC_FEATURE
<223> Murine SC17.121 VH

25 <400> 147

Glu Val Gln Leu Gln Gln Ser Gly Pro Glu Leu Met Lys Pro Gly Ala
1 5 10 15

30 Ser Val Lys Met Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asp His
20 25 30

35 Asn Ile His Trp Val Lys Gln His Gln Gly Lys Ser Leu Glu Trp Ile
35 40 45

Gly Glu Ile Asn Pro Asn Thr Gly Gly Thr Gly Tyr Asn Gln Lys Phe
50 55 60

40 Gln Gly Lys Ala Thr Met Thr Val Asp Lys Ser Ser Ser Thr Ala Tyr
65 70 75 80

45 Met Glu Leu Arg Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys
85 90 95

50 Val Arg Gly Leu Tyr Phe Phe Asp Tyr Trp Gly Gln Gly Thr Thr Leu
100 105 110

Thr Val Ser Ser
115

55 <210> 148
<211> 112

EP 3 338 793 A1

<212> PRT
<213> Mus musculus

<220>
<221> MISC_FEATURE
<223> Murine SC17.122 VL

<400> 148

Asp Ile Val Ile Thr Gln Asp Asp Leu Ser Asn Pro Val Thr Ser Gly
1 5 10 15

Glu Ser Val Ser Ile Ser Cys Arg Ser Ser Lys Ser Leu Leu Tyr Lys
20 25 30

Asp Gly Lys Thr Tyr Leu Asn Trp Phe Leu Gln Arg Pro Gly Gln Ser
35 40 45

Pro Gln Leu Leu Ile Tyr Leu Met Ser Thr Arg Ala Ser Gly Val Ser
50 55 60

Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Glu Ile
65 70 75 80

Ser Arg Val Lys Ala Glu Asp Val Gly Val Tyr Tyr Cys Gln Gln Leu
85 90 95

Val Glu Tyr Pro Arg Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys
100 105 110

<210> 149
<211> 113
<212> PRT
<213> Mus musculus

<220>
<221> MISC_FEATURE
<223> Murine SC17.122 VH

<400> 149

Glu Val His Leu Val Glu Ser Gly Gly Asp Leu Val Lys Pro Gly Gly
1 5 10 15

Ser Leu Lys Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
20 25 30

Gly Met Ser Trp Val Arg Gln Thr Pro Asp Lys Arg Leu Glu Trp Val
35 40 45

Ala Thr Ile Ser Ser Gly Gly Thr Tyr Thr Tyr Tyr Pro Asp Ser Val

EP 3 338 793 A1

50

55

60

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Thr Leu Tyr
65 70 75 80

Leu Gln Met Ser Ser Leu Lys Ser Glu Asp Thr Ala Met Tyr Tyr Cys
85 90 95

Ser Arg His Gly Trp Gly Trp Gly Gln Gly Thr Leu Val Thr Val Ser
100 105 110

Ala

<210> 150
<211> 106
<212> PRT
<213> Mus musculus

<220>
<221> MISC_FEATURE
<223> Murine SC17.140 VL

<400> 150

Gln Ile Val Leu Thr Gln Ser Pro Ala Ile Met Ser Ala Ser Pro Gly
1 5 10 15

Glu Lys Val Thr Met Thr Cys Ser Ala Ser Ser Ser Val Ser Tyr Met
20 25 30

His Trp Tyr Gln Gln Lys Ser Gly Thr Ser Pro Lys Arg Trp Ile Tyr
35 40 45

Asp Thr Ser Lys Leu Ala Ser Gly Val Pro Ala Arg Phe Ser Gly Ser
50 55 60

Gly Ser Gly Thr Ser Tyr Ser Leu Thr Ile Ser Ser Met Glu Ala Glu
65 70 75 80

Asp Ala Ala Thr Tyr Tyr Cys Gln Gln Trp Ser Ser Thr Pro Pro Thr
85 90 95

Phe Gly Ser Gly Thr Lys Leu Glu Ile Lys
100 105

<210> 151
<211> 120
<212> PRT
<213> Mus musculus

EP 3 338 793 A1

<220>
 <221> MISC_FEATURE
 <223> Murine SC17.140 VH

5

<400> 151

Glu Val Gln Leu Gln Gln Ser Gly Pro Glu Leu Met Lys Pro Gly Ala
 1 5 10 15

10

Ser Val Lys Met Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asp Tyr
 20 25 30

15

Asn Met His Trp Val Lys Gln Asn Gln Gly Lys Ser Leu Glu Trp Ile
 35 40 45

Gly Glu Ile Asn Pro Asn Thr Gly Gly Thr Gly Tyr Asn Gln Lys Phe
 50 55 60

20

Lys Gly Lys Ala Thr Leu Thr Val Asp Lys Phe Ser Ser Thr Ala Phe
 65 70 75 80

25

Ile Glu Leu Arg Ser Leu Thr Ser Glu Asp Ser Ala Ile Tyr Tyr Cys
 85 90 95

30

Thr Arg Gly Gly Tyr Asp His Tyr Trp Tyr Phe Asp Val Trp Gly Ala
 100 105 110

Gly Thr Thr Val Thr Val Ser Ser
 115 120

35

<210> 152
 <211> 111
 <212> PRT
 <213> Mus musculus

40

<220>
 <221> MISC_FEATURE
 <223> Murine SC17.151 VL

45

<400> 152

Asp Ile Val Leu Thr Gln Phe Pro Ala Ser Leu Ala Val Ser Leu Gly
 1 5 10 15

50

Gln Arg Ala Thr Ile Pro Cys Arg Ala Ser Glu Ser Val Asp Ser Tyr
 20 25 30

55

Gly Asn Ser Phe Met His Trp Phe Gln Gln Lys Pro Gly Gln Pro Pro
 35 40 45

EP 3 338 793 A1

Lys Leu Leu Ile Tyr Arg Ala Ser Asn Leu Glu Ser Glu Ile Pro Ala
50 55 60

5 Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Asn
65 70 75 80

10 Pro Val Glu Ala Asp Asp Val Ala Thr Tyr Tyr Cys Gln Gln Ser His
85 90 95

Glu Asp Pro Tyr Thr Phe Gly Gly Gly Thr Lys Met Glu Ile Lys
100 105 110

15
<210> 153
<211> 117
<212> PRT
<213> Mus musculus

20
<220>
<221> MISC_FEATURE
<223> Murine SC17.151 VH

25
<400> 153

Glu Val Gln Leu Gln Gln Ser Gly Thr Val Leu Ala Arg Pro Gly Ala
1 5 10 15

30 Ser Val Lys Met Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Ser Tyr
20 25 30

35 Trp Met His Trp Val Lys Gln Arg Pro Gly Gln Gly Leu Glu Trp Ile
35 40 45

Gly Ala Ile Tyr Pro Gly Lys Asn Asp Thr Thr Tyr Asn Gln Lys Phe
50 55 60

40 Lys Gly Lys Ala Lys Leu Thr Ala Val Thr Ser Ala Ser Thr Leu Tyr
65 70 75 80

45 Met Glu Leu Ser Ser Leu Thr Asn Glu Asp Ser Ala Val Tyr Tyr Cys
85 90 95

50 Thr Arg Ser Gly Lys Gly Tyr Phe Ala Tyr Trp Gly Gln Gly Thr Leu
100 105 110

Val Thr Val Ser Ala
115

55
<210> 154
<211> 112

EP 3 338 793 A1

<212> PRT
<213> Mus musculus

<220>
<221> MISC_FEATURE
<223> Murine SC17.156 VL

<400> 154

Asp Val Val Met Thr Gln Ser Pro Leu Ser Leu Pro Val Ser Leu Gly
1 5 10 15

Asp Gln Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Ile Val His Ser
20 25 30

Asn Gly Asn Thr Tyr Leu Glu Trp Tyr Leu Gln Lys Pro Gly Gln Ser
35 40 45

Pro Lys Leu Leu Ile Tyr Lys Val Ser Asn Arg Phe Ser Gly Val Pro
50 55 60

Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile
65 70 75 80

Ser Arg Val Glu Ala Glu Asp Leu Gly Val Tyr Tyr Cys Phe Gln Gly
85 90 95

Ser His Val Pro Pro Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys
100 105 110

<210> 155
<211> 118
<212> PRT
<213> Mus musculus

<220>
<221> MISC_FEATURE
<223> Murine SC17.156 VH

<400> 155

Gln Val Thr Leu Lys Glu Ser Gly Pro Gly Ile Leu Gln Pro Ser Gln
1 5 10 15

Thr Leu Ser Leu Thr Cys Ser Phe Ser Gly Phe Ser Leu Ser Thr Ser
20 25 30

Gly Met Gly Val Ser Trp Ile Arg Lys Thr Ser Gly Lys Gly Leu Glu
35 40 45

Trp Leu Ala His Ile Phe Trp Asp Asp Asp Lys Trp Tyr Asn Pro Ser

EP 3 338 793 A1

	50	55	60
5	Leu Lys Ser Arg Leu Thr Ile Ser Lys Ala Thr Ser Ser Asn Gln Val	65	70 75 80
10	Phe Leu Ile Leu Thr Ser Val Asp Thr Ala Asp Thr Ala Thr Tyr Tyr	85	90 95
15	Cys Ala Thr Phe Tyr Gly Leu Tyr Phe Ala Tyr Trp Gly Gln Gly Thr	100	105 110
20	Thr Leu Thr Val Ser Ser	115	
25	<210> 156		
	<211> 113		
	<212> PRT		
	<213> Mus musculus		
30	<220>		
	<221> MISC_FEATURE		
	<223> Murine SC17.161 VL		
	<400> 156		
35	Asp Ile Val Met Ser Gln Ser Pro Ser Ser Leu Ala Val Ser Val Gly	1	5 10 15
40	Glu Lys Val Thr Met Asn Cys Glu Ser Ser Gln Ser Leu Leu Tyr Asn	20	25 30
45	Ser Asn Gln Lys Asn Tyr Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln	35	40 45
50	Ser Pro Lys Leu Leu Ile Tyr Trp Ala Ser Thr Arg Asp Ser Gly Val	50	55 60
55	Pro Asp Arg Phe Thr Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr	65	70 75 80
	Ile Ser Ser Val Arg Ala Asp Asp Pro Ala Val Tyr Tyr Cys Gln Gln	85	90 95
	Tyr Phe Asn Tyr Pro Leu Thr Phe Gly Ala Gly Thr Lys Leu Glu Leu	100	105 110
	Lys		

EP 3 338 793 A1

<210> 157
 <211> 116
 <212> PRT
 <213> Mus musculus

<220>
 <221> MISC_FEATURE
 <223> Murine SC17.161 VH

<400> 157

Glu Val Lys Leu Glu Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
 1 5 10 15

Ser Met Lys Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Asp Ala
 20 25 30

Trp Met Asp Trp Val Arg Gln Ser Pro Glu Lys Gly Leu Glu Trp Val
 35 40 45

Ala Glu Ile Arg Ser Lys Pro Asn Asn His Ala Thr Tyr Tyr Ala Glu
 50 55 60

Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asp Ser Lys Ser Ser
 65 70 75 80

Ala Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Gly Ile Tyr
 85 90 95

Tyr Cys Val Ser Thr Gly Thr Ser Tyr Trp Gly Gln Gly Thr Leu Val
 100 105 110

Thr Val Ser Ala
 115

<210> 158
 <211> 106
 <212> PRT
 <213> Mus musculus

<220>
 <221> MISC_FEATURE
 <223> Murine SC17.166 VL

<400> 158

Gln Ile Val Leu Thr Gln Ser Pro Ala Ile Met Ser Ala Ser Pro Gly
 1 5 10 15

Glu Lys Val Thr Met Thr Cys Ser Ala Ser Ser Ser Ile Ser Tyr Met
 20 25 30

EP 3 338 793 A1

His Trp Tyr Gln Gln Lys Ser Gly Thr Ser Pro Lys Arg Trp Ile Tyr
35 40 45

5 Asp Thr Ser Lys Leu Ala Ser Gly Val Pro Ala Arg Phe Ser Gly Ser
50 55 60

10 Gly Ser Gly Thr Ser Tyr Ser Leu Thr Ile Ser Asn Met Glu Ala Glu
65 70 75 80

15 Asp Ala Ala Thr Tyr Tyr Cys Gln Gln Trp Ser Ser Thr Pro Pro Thr
85 90 95

20 Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys
100 105

<210> 159
<211> 117
<212> PRT
<213> Mus musculus

25 <220>
<221> MISC_FEATURE
<223> Murine SC17.166 VH

<400> 159

30 Glu Val Gln Leu Gln Gln Ser Gly Pro Glu Leu Met Lys Pro Gly Ala
1 5 10 15

35 Ser Val Lys Met Ser Cys Lys Ala Ser Gly Tyr Ile Phe Thr Asp Tyr
20 25 30

40 Asn Met His Trp Val Lys Gln Asn Gln Gly Lys Ser Leu Glu Trp Ile
35 40 45

Gly Glu Val Asn Pro Asn Thr Gly Gly Ile Gly Tyr Asn Gln Lys Phe
50 55 60

45 Lys Gly Lys Ala Thr Leu Thr Val Asp Lys Ser Ser Ser Thr Ala Tyr
65 70 75 80

50 Met Asp Leu Arg Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys
85 90 95

Ala Arg Asp Gly Asn Tyr Cys Phe Asp Tyr Trp Gly Gln Gly Thr Thr
100 105 110

55 Leu Thr Val Ser Ser
115

EP 3 338 793 A1

<210> 160
<211> 107
<212> PRT
<213> Mus musculus

<220>
<221> MISC FEATURE
<223> Murine SC17.187 VL

<400> 160

Asp Ile Lys Met Thr Gln Ser Pro Ser Ser Met Tyr Ala Ser Leu Gly
1 5 10 15

Glu Arg Val Thr Leu Thr Cys Lys Ala Ser Gln Asp Ile Asn Ser Tyr
20 25 30

Leu Ser Trp Phe Gln Gln Lys Pro Gly Lys Ser Pro Glu Thr Leu Ile
35 40 45

Tyr Arg Ala Asn Arg Leu Ile Asp Gly Val Pro Ser Arg Phe Ser Gly
50 55 60

Ser Gly Ser Gly Gln Asp Tyr Ser Leu Thr Ile Ser Ser Leu Glu Tyr
65 70 75 80

Glu Asp Met Gly Ile Tyr Tyr Cys Leu Gln Tyr Asp Glu Phe Pro Pro
85 90 95

Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys
100 105

<210> 161
<211> 118
<212> PRT
<213> Mus musculus

<220>
<221> MISC FEATURE
<223> Murine SC17.187 VH

<400> 161

Glu Val His Leu Gln Gln Ser Gly Pro Glu Leu Val Asn Pro Gly Ser
1 5 10 15

Ser Val Lys Ile Ser Cys Lys Ala Ala Gly Tyr Thr Phe Thr Asp Tyr
20 25 30

Asn Met Asp Trp Val Lys Gln Ser His Gly Lys Arg Leu Glu Trp Ile

EP 3 338 793 A1

	35	40	45
5	Gly Asn Ile Tyr Pro Asn Asn Gly Gly Ala Gly Tyr Asn Gln Asn Phe 50 55 60		
10	Lys Asp Lys Ala Thr Leu Thr Val Asp Lys Ser Ser Ser Thr Ala Tyr 65 70 75 80		
15	Met Glu Leu Arg Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys 85 90 95		
20	Ala Arg Ser Ile Thr Ala Ala Trp Phe Ala Tyr Trp Gly Gln Gly Thr 100 105 110		
25	Leu Val Thr Val Ser Ala 115		
30	<210> 162 <211> 106 <212> PRT <213> Mus musculus		
35	<220> <221> MISC_FEATURE <223> Murine SC17.191 VL <400> 162		
40	Gln Ile Val Leu Thr Gln Ser Pro Ala Ile Met Ser Ala Ser Pro Gly 1 5 10 15		
45	Glu Lys Val Thr Met Thr Cys Ser Ala Ser Ser Ser Val Ser Tyr Met 20 25 30		
50	His Trp Tyr Gln Gln Lys Ser Gly Thr Ser Pro Lys Arg Trp Ile Tyr 35 40 45		
55	Asp Thr Ser Lys Leu Ala Ser Gly Val Pro Ala Arg Phe Thr Gly Ser 50 55 60		
60	Gly Ser Gly Thr Ser Tyr Ser Leu Thr Ile Ser Ser Met Glu Ala Glu 65 70 75 80		
65	Asp Ala Ala Thr Tyr Tyr Cys Gln Gln Trp Ser Ser Ser Pro Pro Thr 85 90 95		
70	Phe Gly Ala Gly Thr Lys Leu Glu Leu Lys 100 105		

EP 3 338 793 A1

<210> 163
<211> 120
<212> PRT
<213> Mus musculus

<220>
<221> MISC_FEATURE
<223> Murine SC17.191 VH

<400> 163

Glu Val Gln Leu Gln Gln Ser Gly Pro Glu Leu Met Lys Pro Gly Ala
1 5 10 15

Ser Val Lys Met Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asp Tyr
20 25 30

Asn Met His Trp Val Lys Gln Asn Gln Gly Lys Ser Leu Glu Trp Ile
35 40 45

Gly Glu Ile Asn Pro Asn Thr Gly Gly Thr Gly Tyr Asn Gln Lys Phe
50 55 60

Lys Asp Lys Ala Thr Leu Thr Val Asp Lys Ser Ser Ser Thr Ala Tyr
65 70 75 80

Met Glu Leu Arg Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys
85 90 95

Ala Arg Ile Pro Ser Leu Arg Arg Tyr Tyr Phe Asp Tyr Trp Gly Gln
100 105 110

Gly Thr Thr Leu Thr Val Ser Ser
115 120

<210> 164
<211> 111
<212> PRT
<213> Mus musculus

<220>
<221> MISC_FEATURE
<223> Murine SC17.193 VL

<400> 164

Asp Leu Val Leu Thr Gln Ser Pro Ala Ser Leu Ala Val Ser Leu Gly
1 5 10 15

Gln Arg Ala Thr Ile Ser Cys Arg Ala Ser Glu Ser Val Ser Thr Ser
20 25 30

EP 3 338 793 A1

	Gly	Tyr	Ser	Tyr	Met	His	Trp	Tyr	Gln	Gln	Lys	Pro	Gly	Gln	Pro	Pro	
			35					40					45				
5	Lys	Leu	Leu	Ile	Tyr	Leu	Ala	Ser	Asn	Leu	Glu	Ser	Gly	Val	Pro	Ala	
		50					55					60					
	Arg	Phe	Ser	Gly	Ser	Gly	Ser	Gly	Thr	Asp	Phe	Thr	Leu	Asn	Ile	His	
10	65					70					75					80	
	Pro	Val	Glu	Glu	Glu	Asp	Ala	Thr	Thr	Tyr	Tyr	Cys	Gln	His	Ser	Arg	
					85					90					95		
15	Glu	Leu	Pro	Tyr	Thr	Phe	Gly	Gly	Gly	Thr	Lys	Leu	Glu	Ile	Lys		
				100					105					110			
20	<210>	165															
	<211>	122															
	<212>	PRT															
	<213>	Mus musculus															
25	<220>																
	<221>	MISC_FEATURE															
	<223>	Murine SC17.193 VH															
	<400>	165															
30	Gln	Val	Thr	Leu	Lys	Glu	Ser	Gly	Pro	Gly	Ile	Leu	Gln	Pro	Ser	Gln	
	1				5					10					15		
	Thr	Leu	Ser	Leu	Thr	Cys	Ser	Phe	Ser	Gly	Phe	Ser	Leu	Ile	Thr	Tyr	
35				20					25					30			
	Gly	Ile	Gly	Val	Gly	Trp	Ile	Arg	Gln	Pro	Ser	Gly	Lys	Gly	Leu	Glu	
			35					40					45				
40	Trp	Leu	Ala	His	Ile	Trp	Trp	Asn	Asp	Asn	Lys	Tyr	Tyr	Asn	Thr	Ala	
		50					55					60					
	Leu	Lys	Ser	Arg	Leu	Thr	Ile	Ser	Lys	Asp	Thr	Ser	Asn	Asn	Gln	Val	
45	65					70					75					80	
	Phe	Leu	Lys	Ile	Ala	Asn	Val	Asp	Thr	Ala	Asp	Thr	Ala	Thr	Tyr	Tyr	
50					85					90					95		
	Cys	Ala	Arg	Met	Val	Tyr	Tyr	Asp	Tyr	Asp	Gly	Gly	Phe	Ala	Tyr	Trp	
				100					105					110			
55	Gly	Gln	Gly	Thr	Leu	Val	Thr	Val	Ser	Ala							
			115					120									

EP 3 338 793 A1

<210> 166
 <211> 111
 <212> PRT
 <213> Mus musculus

5

<220>
 <221> MISC FEATURE
 <223> Murine SC17.199 VL

10

<400> 166

Asp Ile Val Leu Thr Gln Ser Pro Ala Ser Leu Ala Val Ser Leu Gly
 1 5 10 15

15

Gln Arg Ala Thr Ile Ser Cys Arg Ala Ser Glu Ser Val Asp Ser Tyr
 20 25 30

20

Gly Asn Ser Phe Met His Trp Tyr Gln Gln Lys Pro Gly Gln Pro Pro
 35 40 45

25

Lys Pro Leu Ile Tyr Arg Ala Ser Asn Leu Glu Ser Gly Ile Pro Ala
 50 55 60

Arg Phe Ser Gly Ser Gly Ser Arg Thr Asp Phe Thr Leu Thr Ile Asn
 65 70 75 80

30

Pro Val Glu Ala Asp Asp Val Ala Thr Tyr Tyr Cys Gln Gln Ser Asn
 85 90 95

35

Glu Asp Pro Tyr Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys
 100 105 110

<210> 167
 <211> 116
 <212> PRT
 <213> Mus musculus

40

<220>
 <221> MISC FEATURE
 <223> Murine SC17.199 VH

45

<400> 167

Glu Val Gln Leu Gln Gln Ser Gly Thr Val Leu Ala Arg Pro Gly Ala
 1 5 10 15

50

Ser Val Arg Met Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Ser Tyr
 20 25 30

55

Trp Met His Trp Val Lys Gln Arg Pro Gly Gln Gly Leu Glu Trp Ile

EP 3 338 793 A1

	35	40	45
5	Gly Ala Ile Tyr Pro Gly Asn Ser Asp Thr Ser Tyr Asn His Lys Phe 50 55 60		
10	Lys Gly Lys Ala Lys Leu Thr Ala Val Thr Ser Ala Ser Thr Ala Tyr 65 70 75 80		
15	Met Glu Leu Ser Ser Leu Thr Asn Glu Asp Ser Ala Val Tyr Tyr Cys 85 90 95		
20	Thr Arg Ser Gly Thr Gly Trp Phe Ala Tyr Trp Gly Gln Gly Thr Leu 100 105 110		
25	Val Thr Val Ser 115		
30	<210> 168 <211> 111 <212> PRT <213> Mus musculus		
35	<220> <221> MISC_FEATURE <223> Murine SC17.200 VL <400> 168		
40	Asp Ile Val Leu Thr Gln Ser Pro Ala Ser Leu Ala Val Ser Leu Gly 1 5 10 15		
45	Gln Arg Ala Thr Ile Phe Cys Arg Ala Ser Gln Ser Val Asp Tyr Asn 20 25 30		
50	Gly Ile Ser Tyr Met His Trp Phe Gln Gln Lys Pro Gly Gln Pro Pro 35 40 45		
55	Lys Leu Leu Ile Tyr Ala Ala Ser Asn Val Gln Ser Gly Ile Pro Ala 50 55 60		
60	Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Asn Ile His 65 70 75 80		
65	Pro Val Glu Glu Glu Asp Ala Ala Thr Phe Tyr Cys Gln Gln Ser Ile 85 90 95		
70	Glu Asp Pro Pro Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys 100 105 110		

EP 3 338 793 A1

<210> 169
<211> 115
<212> PRT
<213> Mus musculus

5

<220>
<221> MISC_FEATURE
<223> Murine SC17. VH

10

<220>
<221> MISC_FEATURE
<223> Murine SC17.200 VH

<400> 169

15

Gln Val Gln Leu Gln Gln Ser Gly Pro Glu Leu Val Lys Pro Gly Ala
1 5 10 15

20

Ser Val Lys Ile Ser Cys Lys Ala Ser Gly Tyr Ala Phe Ser Ser Ser
20 25 30

Trp Ile Asn Trp Val Lys Gln Arg Pro Gly Gln Gly Leu Glu Trp Ile
35 40 45

25

Gly Arg Ile Tyr Pro Gly Glu Gly Asp Thr Asn Tyr Ser Gly Asn Phe
50 55 60

30

Glu Gly Lys Ala Thr Leu Thr Ala Asp Lys Ser Ser Thr Thr Ala Tyr
65 70 75 80

Met Gln Leu Ser Ser Leu Thr Ser Val Asp Ser Ala Val Tyr Phe Cys
85 90 95

35

Thr Arg Gly Leu Val Met Asp Tyr Trp Gly Gln Gly Thr Ala Leu Thr
100 105 110

40

Val Ser Ser
115

45

<210> 170
<211> 107
<212> PRT
<213> Artificial Sequence

<220>
<223> Humanized hSC17.16 VL

50

<400> 170

Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
1 5 10 15

55

Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Ala Asn Ile Asn Ser Asn

EP 3 338 793 A1

		20		25		30	
5	Leu Val Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile	35	40	45			
10	Tyr Ala Ala Thr Asn Leu Ala Asp Gly Val Pro Ser Arg Phe Ser Gly	50	55	60			
15	Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro	65	70	75	80		
20	Glu Asp Phe Ala Thr Tyr Tyr Cys Gln His Phe Trp Gly Thr Pro Arg	85	90	95			
25	Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys	100	105				
30	<210> 171						
	<211> 116						
	<212> PRT						
	<213> Artificial Sequence						
35	<220>						
	<223> Humanized hSC17.16 VH						
40	<400> 171						
45	Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala	1	5	10	15		
50	Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asp Tyr	20	25	30			
55	Asn Met Tyr Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met	35	40	45			
60	Gly Glu Ile Asn Pro Asn Asn Gly Gly Thr Ala Tyr Asn Gln Lys Phe	50	55	60			
65	Arg Gly Lys Val Thr Met Thr Arg Asp Thr Ser Ile Ser Thr Ala Tyr	65	70	75	80		
70	Met Glu Leu Ser Arg Leu Arg Ser Asp Asp Thr Ala Val Tyr Tyr Cys	85	90	95			
75	Ala Arg Tyr Asp Lys Gly Phe Asp Tyr Trp Gly Gln Gly Thr Thr Val	100	105	110			
80	Thr Val Ser Ser	115					

EP 3 338 793 A1

<210> 172
<211> 106
<212> PRT
<213> Artificial Sequence

<220>
<223> Humanized hSC17.17 VL

<400> 172

Glu Ile Val Leu Thr Gln Ser Pro Ala Thr Leu Ser Leu Ser Pro Gly
1 5 10 15

Glu Arg Ala Thr Leu Ser Cys Ser Ala Ser Ser Ser Val Ser Tyr Met
20 25 30

His Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro Arg Leu Leu Ile Tyr
35 40 45

Asp Thr Ser Lys Leu Pro Ser Gly Ile Pro Ala Arg Phe Ser Gly Ser
50 55 60

Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Glu Pro Glu
65 70 75 80

Asp Phe Ala Val Tyr Tyr Cys Gln Gln Trp Ser Ser Thr Pro Pro Thr
85 90 95

Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys
100 105

<210> 173
<211> 120
<212> PRT
<213> Artificial Sequence

<220>
<223> Humanized hSC17.17 VH

<400> 173

Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala
1 5 10 15

Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asp Tyr
20 25 30

Asn Met His Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met
35 40 45

Gly Glu Ile Asn Pro Asn Ile Gly Gly Thr Gly Tyr Asn Gln Lys Phe

EP 3 338 793 A1

50

55

60

Lys Gly Arg Val Thr Met Thr Arg Asp Thr Ser Ile Ser Thr Ala Tyr
65 70 75 80

Met Glu Leu Ser Arg Leu Arg Ser Asp Asp Thr Ala Val Tyr Tyr Cys
85 90 95

Ala Arg Thr Tyr Ser Tyr Tyr Ser Tyr Glu Phe Ala Tyr Trp Gly Gln
100 105 110

Gly Thr Leu Val Thr Val Ser Ser
115 120

<210> 174
<211> 112
<212> PRT
<213> Artificial Sequence

<220>
<223> Humanized hSC17.24 VL

<400> 174

Asp Ile Val Met Thr Gln Ser Pro Asp Ser Leu Ala Val Ser Leu Gly
1 5 10 15

Glu Arg Ala Thr Ile Asn Cys Lys Ser Ser Gln Ser Leu Leu Tyr Ser
20 25 30

Ser Asn Gln Lys Ser Tyr Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln
35 40 45

Pro Pro Lys Leu Leu Ile Tyr Trp Ala Ser Thr Arg Glu Ser Gly Val
50 55 60

Pro Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr
65 70 75 80

Ile Ser Ser Leu Gln Ala Glu Asp Val Ala Val Tyr Tyr Cys Lys Gln
85 90 95

Ser Tyr Asn Leu Arg Thr Phe Gly Gly Gly Thr Lys Val Glu Ile Lys
100 105 110

<210> 175
<211> 117
<212> PRT
<213> Artificial Sequence

<220>

EP 3 338 793 A1

<223> Humanized hSC17.24 VH

<400> 175

5 Glu Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala
1 5 10 15

10 Thr Val Lys Ile Ser Cys Lys Val Ser Gly Tyr Thr Phe Thr Asp His
20 25 30

15 Thr Ile His Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Ile
35 40 45

Gly Tyr Ile Tyr Pro Arg Asp Gly Ser Thr Lys Tyr Asn Glu Glu Phe
50 55 60

20 Lys Gly Arg Val Thr Ile Thr Ala Asp Thr Ser Thr Asp Thr Ala Tyr
65 70 75 80

25 Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95

Ala Arg Ser Tyr Ser Asn Tyr Phe Asp Tyr Trp Gly Gln Gly Thr Thr
100 105 110

30 Val Thr Val Ser Ser
115

<210> 176

<211> 107

<212> PRT

<213> Artificial Sequence

<220>

<223> Humanized hSC17.28 VL

<400> 176

40 Glu Ile Val Leu Thr Gln Ser Pro Asp Phe Gln Ser Val Thr Pro Lys
1 5 10 15

45 Glu Lys Val Thr Ile Thr Cys Arg Ala Ser Gln Ser Ile Gly Thr Ser
20 25 30

50 Ile His Trp Tyr Gln Gln Lys Pro Asp Gln Ser Pro Lys Leu Leu Ile
35 40 45

Lys Tyr Ala Ser Glu Ser Ile Ser Gly Val Pro Ser Arg Phe Ser Gly
50 55 60

55 Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Asn Ser Leu Glu Ala

EP 3 338 793 A1

	65		70		75		80
5	Glu Asp Ala Ala Thr Tyr Tyr Cys Gln Gln Ser Asn Ser Trp Pro Leu	85		90		95	
10	Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys	100		105			
15	<210> 177						
	<211> 117						
	<212> PRT						
	<213> Artificial Sequence						
20	Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala	5		10		15	
25	Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Arg Ser	20		25		30	
30	Tyr Ile His Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met	35		40		45	
35	Gly Tyr Ile Ser Ser Gly Ser Gly Gly Thr Thr Tyr Asn Gln Lys Phe	50		55		60	
40	Lys Gly Arg Val Thr Ser Thr Arg Asp Thr Ser Ile Ser Thr Ala Tyr	65		70		75	80
45	Met Glu Leu Ser Arg Leu Arg Ser Asp Asp Thr Ala Val Tyr Tyr Cys	85		90		95	
50	Ala Arg Gly Gly Val Arg Tyr Phe Asp Val Trp Gly Gln Gly Thr Thr	100		105		110	
55	Val Thr Val Ser Ser	115					
	<210> 178						
	<211> 107						
	<212> PRT						
	<213> Artificial Sequence						
	<220>						
	<223> Humanized hSC17.34 VL						
	<400> 178						

EP 3 338 793 A1

Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
 1 5 10 15
 5 Asp Arg Val Thr Ile Thr Cys Lys Ala Ser Gln Asp Ile Asn Ser Tyr
 20 25 30
 10 Leu Ser Trp Phe Gln Gln Lys Pro Gly Lys Ala Pro Lys Ser Leu Ile
 35 40 45
 Tyr Arg Ala Asn Arg Leu Val Asp Gly Val Pro Ser Arg Phe Ser Gly
 50 55 60
 15 Ser Gly Ser Gly Thr Asp Tyr Thr Leu Thr Ile Ser Ser Leu Gln Pro
 65 70 75 80
 20 Glu Asp Phe Ala Thr Tyr Tyr Cys Leu Gln Tyr Asp Glu Phe Pro Pro
 85 90 95
 Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys
 100 105
 25
 <210> 179
 <211> 118
 <212> PRT
 <213> Artificial Sequence
 30
 <220>
 <223> Humanized hSC17.34 VH
 <400> 179
 35 Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala
 1 5 10 15
 Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asp Tyr
 20 25 30
 40 Asn Met Asp Trp Val Arg Gln Ala Pro Gly Gln Arg Leu Glu Trp Ile
 35 40 45
 45 Gly Tyr Ile Tyr Pro Asp Asn Gly Gly Ala Gly Tyr Asn Gln Lys Phe
 50 55 60
 50 Lys Gly Arg Val Thr Ile Thr Val Asp Thr Ser Ala Ser Thr Ala Tyr
 65 70 75 80
 Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95
 55 Ser Arg Ser Ile Thr Thr Ala Trp Phe Ala Tyr Trp Gly Gln Gly Thr

EP 3 338 793 A1

	100	105	110
5	Leu Val Thr Val Ser Ser		
	115		
	<210> 180		
	<211> 107		
10	<212> PRT		
	<213> Artificial Sequence		
	<220>		
	<223> Humanized hSC17.46 VL		
15	<400> 180		
	Ala Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly		
	1 5 10 15		
20	Asp Arg Val Thr Ile Thr Cys Lys Ala Ser Gln Ser Val Asn Asn Asp		
	20 25 30		
	Val Ala Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile		
25	35 40 45		
	Tyr Tyr Ala Ser Asn Arg Tyr Thr Gly Val Pro Ser Arg Phe Ser Gly		
	50 55 60		
30	Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro		
	65 70 75 80		
35	Glu Asp Phe Ala Thr Tyr Phe Cys Gln Gln Asp Tyr Ser Ser Pro Arg		
	85 90 95		
	Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys		
40	100 105		
	<210> 181		
	<211> 121		
45	<212> PRT		
	<213> Artificial Sequence		
	<220>		
	<223> Humanized hSC17.46 VH		
50	<400> 181		
	Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala		
	1 5 10 15		
55	Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Ser Tyr		
	20 25 30		

EP 3 338 793 A1

Trp Ile Asn Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Ile
35 40 45

5 Gly Asn Ile Phe Pro Asp Thr Thr Thr Thr Asn Tyr Asn Glu Lys Phe
50 55 60

10 Lys Gly Arg Val Thr Leu Thr Arg Asp Thr Ser Ile Ser Thr Ala Tyr
65 70 75 80

Met Glu Leu Ser Arg Leu Arg Ser Asp Asp Thr Ala Val Tyr Tyr Cys
85 90 95

15 Ala Arg Glu Tyr Tyr Asp Gly Thr Tyr Asp Ala Met Asp Tyr Trp Gly
100 105 110

20 Gln Gly Thr Leu Val Thr Val Ser Ser
115 120

<210> 182
<211> 111
<212> PRT
25 <213> Artificial Sequence

<220>
<223> Humanized hSC17.151 VL

30 <400> 182

Glu Ile Val Leu Thr Gln Ser Pro Ala Thr Leu Ser Leu Ser Pro Gly
1 5 10 15

35 Glu Arg Ala Thr Leu Ser Cys Arg Ala Ser Glu Ser Val Asp Ser Tyr
20 25 30

40 Gly Asn Ser Phe Met His Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro
35 40 45

Arg Leu Leu Ile Tyr Arg Ala Ser Asn Leu Glu Ser Gly Ile Pro Ala
50 55 60

45 Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser
65 70 75 80

50 Ser Leu Glu Pro Glu Asp Phe Ala Val Tyr Tyr Cys Gln Gln Ser His
85 90 95

Glu Asp Pro Tyr Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys
100 105 110

55 <210> 183

EP 3 338 793 A1

<211> 117
<212> PRT
<213> Artificial Sequence

<220>
<223> Humanized hSC17.151 VH

<400> 183

Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala
1 5 10 15

Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Ser Tyr
20 25 30

Trp Met His Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met
35 40 45

Gly Ala Ile Tyr Pro Gly Lys Ser Asp Thr Thr Tyr Asn Gln Lys Phe
50 55 60

Lys Gly Arg Val Thr Met Thr Arg Asp Thr Ser Thr Ser Thr Val Tyr
65 70 75 80

Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95

Ala Arg Ser Gly Lys Gly Tyr Phe Ala Tyr Trp Gly Gln Gly Thr Leu
100 105 110

Val Thr Val Ser Ser
115

<210> 184
<211> 113
<212> PRT
<213> Artificial Sequence

<220>
<223> Humanized hSC17.155 VL

<400> 184

Asp Ile Val Met Thr Gln Ser Pro Asp Ser Leu Ala Val Ser Leu Gly
1 5 10 15

Glu Arg Ala Thr Ile Asn Cys Lys Ser Ser Gln Ser Leu Leu Tyr Ser
20 25 30

Ser Asn Gln Lys Asn Tyr Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln
35 40 45

EP 3 338 793 A1

Pro Pro Lys Leu Leu Ile Tyr Trp Ala Ser Thr Arg Lys Ser Gly Val
50 55 60

5 Pro Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr
65 70 75 80

10 Ile Ser Ser Leu Gln Ala Glu Asp Val Ala Val Tyr Tyr Cys His Gln
85 90 95

Tyr Tyr Ser Tyr Pro Leu Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile
100 105 110

15 Lys

<210> 185
<211> 113
<212> PRT
<213> Artificial Sequence

<220>
20 <223> Humanized hSC17.155 VH
25 <400> 185

Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala
1 5 10 15

30 Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Asn Ser Tyr
20 25 30

35 Trp Met His Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met
35 40 45

40 Gly Glu Ile His Pro Asn Asn Gly Ser Thr Asn Tyr Asn Glu Lys Phe
50 55 60

Lys Gly Arg Val Thr Met Thr Arg Asp Thr Ser Thr Ser Thr Val Tyr
65 70 75 80

45 Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95

50 Ala Arg Trp Thr Leu Phe Thr Tyr Trp Gly Gln Gly Thr Leu Val Thr
100 105 110

Val

55 <210> 186

EP 3 338 793 A1

<211> 112
<212> PRT
<213> Artificial Sequence

<220>
<223> Humanized hSC17.156 VL

<400> 186

Asp Ile Val Met Thr Gln Thr Pro Leu Ser Leu Pro Val Thr Pro Gly
1 5 10 15

Glu Pro Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Ile Val His Ser
20 25 30

Asn Gly Asn Thr Tyr Leu Glu Trp Tyr Leu Gln Lys Pro Gly Gln Ser
35 40 45

Pro Gln Leu Leu Ile Tyr Lys Val Ser Asn Arg Phe Ser Gly Val Pro
50 55 60

Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile
65 70 75 80

Ser Arg Val Glu Ala Glu Asp Val Gly Val Tyr Tyr Cys Phe Gln Gly
85 90 95

Ser His Val Pro Pro Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys
100 105 110

<210> 187
<211> 118
<212> PRT
<213> Artificial Sequence

<220>
<223> Humanized hSC17.156 VH

<400> 187

Gln Val Thr Leu Lys Glu Ser Gly Pro Val Leu Val Lys Pro Thr Glu
1 5 10 15

Thr Leu Thr Leu Thr Cys Thr Val Ser Gly Phe Ser Leu Ser Thr Ser
20 25 30

Gly Met Gly Val Ser Trp Ile Arg Gln Pro Pro Gly Lys Ala Leu Glu
35 40 45

Trp Leu Ala His Ile Phe Trp Asp Asp Asp Lys Trp Tyr Asn Pro Ser
50 55 60

EP 3 338 793 A1

Leu Lys Ser Arg Leu Thr Ile Ser Lys Asp Thr Ser Lys Ser Gln Val
65 70 75 80

5 Val Leu Thr Met Thr Asn Met Asp Pro Val Asp Thr Ala Thr Tyr Tyr
85 90 95

10 Cys Ala Thr Phe Tyr Gly Leu Tyr Phe Ala Tyr Trp Gly Gln Gly Thr
100 105 110

Leu Val Thr Val Ser Ser
115

15 <210> 188
<211> 113
<212> PRT
<213> Artificial Sequence

20 <220>
<223> Humanized hSC17.161 VL

<400> 188

25 Asp Ile Val Met Thr Gln Ser Pro Asp Ser Leu Ala Val Ser Leu Gly
1 5 10 15

30 Glu Arg Ala Thr Ile Asn Cys Glu Ser Ser Gln Ser Leu Leu Tyr Asn
20 25 30

Ser Asn Gln Lys Asn Tyr Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln
35 40 45

35 Pro Pro Lys Leu Leu Ile Tyr Trp Ala Ser Thr Arg Glu Ser Gly Val
50 55 60

40 Pro Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr
65 70 75 80

Ile Ser Ser Leu Gln Ala Glu Asp Val Ala Val Tyr Tyr Cys Gln Gln
85 90 95

45 Tyr Phe Asn Tyr Pro Leu Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile
100 105 110

50 Lys

55 <210> 189
<211> 116
<212> PRT
<213> Artificial Sequence

EP 3 338 793 A1

<220>

<223> Humanized hSC17.161 VH

<400> 189

5

Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala
1 5 10 15

10

Ser Val Lys Val Ser Cys Lys Ala Ser Gly Phe Thr Phe Ser Asp Ala
20 25 30

15

Trp Met Asp Trp Val Arg Gln Ala Pro Gly Gln Arg Leu Glu Trp Met
35 40 45

Gly Glu Ile Arg Ser Lys Pro Asn Asn His Ala Thr Tyr Tyr Ala Glu
50 55 60

20

Ser Val Lys Gly Arg Val Thr Ile Thr Arg Asp Thr Ser Ala Ser Thr
65 70 75 80

25

Ala Tyr Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr
85 90 95

30

Tyr Cys Ala Arg Thr Gly Thr Ser Tyr Trp Gly Gln Gly Thr Leu Val
100 105 110

Thr Val Ser Ser
115

35

<210> 190

<211> 111

<212> PRT

<213> Artificial Sequence

40

<220>

<223> Humanized hSC17.200 VL

<400> 190

45

Glu Ile Val Leu Thr Gln Ser Pro Ala Thr Leu Ser Leu Ser Pro Gly
1 5 10 15

50

Glu Arg Ala Thr Leu Ser Cys Arg Ala Ser Gln Ser Val Asp Tyr Asn
20 25 30

Gly Ile Ser Tyr Met His Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro
35 40 45

55

Arg Leu Leu Ile Tyr Ala Ala Ser Asn Val Gln Ser Gly Ile Pro Ala
50 55 60

EP 3 338 793 A1

Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser
65 70 75 80

5 Ser Leu Glu Pro Glu Asp Phe Ala Val Tyr Tyr Cys Gln Gln Ser Ile
85 90 95

10 Glu Asp Pro Pro Thr Phe Gly Gly Gly Thr Lys Val Glu Ile Lys
100 105 110

15 <210> 191
<211> 115
<212> PRT
<213> Artificial Sequence

<220>
<223> Humanized hSC17.200 VH

20 <400> 191
Glu Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Glu
1 5 10 15

25 Ser Leu Lys Ile Ser Cys Lys Gly Ser Gly Tyr Ser Phe Thr Ser Ser
20 25 30

30 Trp Ile Asn Trp Val Arg Gln Met Pro Gly Lys Gly Leu Glu Trp Met
35 40 45

Gly Arg Ile Tyr Pro Gly Glu Gly Asp Thr Asn Tyr Ser Gly Asn Phe
50 55 60

35 Glu Gly Gln Val Thr Ile Ser Ala Asp Lys Ser Ile Ser Thr Ala Tyr
65 70 75 80

40 Leu Gln Trp Ser Ser Leu Lys Ala Ser Asp Thr Ala Met Tyr Tyr Cys
85 90 95

45 Thr Arg Gly Leu Val Met Asp Tyr Trp Gly Gln Gly Thr Leu Val Thr
100 105 110

Val Ser Ser
115

50 <210> 192
<211> 111
<212> PRT
<213> Artificial Sequence

<220>
<223> Humanized hSC17.200 VL

55 <400> 192

EP 3 338 793 A1

	Glu	Ile	Val	Leu	Thr	Gln	Ser	Pro	Ala	Thr	Leu	Ser	Leu	Ser	Pro	Gly	
	1				5					10					15		
5	Glu	Arg	Ala	Thr	Leu	Ser	Cys	Arg	Ala	Ser	Gln	Ser	Val	Asp	Tyr	Asp	
				20					25					30			
10	Gly	Ile	Ser	Tyr	Met	His	Trp	Tyr	Gln	Gln	Lys	Pro	Gly	Gln	Ala	Pro	
			35					40					45				
15	Arg	Leu	Leu	Ile	Tyr	Ala	Ala	Ser	Asn	Val	Gln	Ser	Gly	Ile	Pro	Ala	
		50					55					60					
20	Arg	Phe	Ser	Gly	Ser	Gly	Ser	Gly	Thr	Asp	Phe	Thr	Leu	Thr	Ile	Ser	
	65					70					75					80	
25	Ser	Leu	Glu	Pro	Glu	Asp	Phe	Ala	Val	Tyr	Tyr	Cys	Gln	Gln	Ser	Ile	
					85					90					95		
30	Glu	Asp	Pro	Pro	Thr	Phe	Gly	Gly	Gly	Thr	Lys	Val	Glu	Ile	Lys		
				100					105					110			
	<210>	193															
	<211>	113															
	<212>	PRT															
	<213>	Artificial Sequence															
	<220>																
	<223>	Humanized hSC17.155 VH1															
35	<400>	193															
40	Gln	Val	Gln	Leu	Val	Gln	Ser	Gly	Ala	Glu	Val	Lys	Lys	Pro	Gly	Ala	
	1				5					10					15		
45	Ser	Val	Lys	Val	Ser	Cys	Lys	Ala	Ser	Gly	Tyr	Thr	Phe	Asp	Ser	Tyr	
				20					25					30			
50	Trp	Met	His	Trp	Val	Arg	Gln	Ala	Pro	Gly	Gln	Gly	Leu	Glu	Trp	Met	
			35					40					45				
55	Gly	Glu	Ile	His	Pro	Asn	Asn	Gly	Ser	Thr	Asn	Tyr	Asn	Glu	Lys	Phe	
		50					55					60					
60	Lys	Gly	Arg	Val	Thr	Met	Thr	Arg	Asp	Thr	Ser	Thr	Ser	Thr	Val	Tyr	
	65					70					75					80	
65	Met	Glu	Leu	Ser	Ser	Leu	Arg	Ser	Glu	Asp	Thr	Ala	Val	Tyr	Tyr	Cys	
					85					90					95		

EP 3 338 793 A1

Ala Arg Trp Thr Leu Phe Thr Tyr Trp Gly Gln Gly Thr Leu Val Thr
100 105 110

5 Val

<210> 194
<211> 113
<212> PRT
<213> Artificial Sequence

10

<220>
<223> Humanized hSC17.155 VH2

15

<400> 194

Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala
1 5 10 15

20

Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Ser Tyr
20 25 30

25

Trp Met His Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met
35 40 45

30

Gly Glu Ile His Pro Asn Asn Gly Ser Thr Asn Tyr Asn Glu Lys Phe
50 55 60

Lys Gly Arg Val Thr Met Thr Arg Asp Thr Ser Thr Ser Thr Val Tyr
65 70 75 80

35

Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95

40

Ala Arg Trp Thr Leu Phe Thr Tyr Trp Gly Gln Gly Thr Leu Val Thr
100 105 110

Val

45

<210> 195
<211> 113
<212> PRT
<213> Artificial Sequence

50

<220>
<223> Humanized hSC17.155 VH3

<400> 195

55

Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala
1 5 10 15

EP 3 338 793 A1

Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Asn Tyr Tyr
20 25 30

5 Trp Met His Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met
35 40 45

10 Gly Glu Ile His Pro Asn Asn Gly Ser Thr Asn Tyr Asn Glu Lys Phe
50 55 60

Lys Gly Arg Val Thr Met Thr Arg Asp Thr Ser Thr Ser Thr Val Tyr
65 70 75 80

15 Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95

20 Ala Arg Trp Thr Leu Phe Thr Tyr Trp Gly Gln Gly Thr Leu Val Thr
100 105 110

25 Val

<210> 196
<211> 113
<212> PRT
30 <213> Artificial Sequence

<220>
<223> Humanized hSC17.155 VH4

35 <400> 196

Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala
1 5 10 15

40 Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Asn Ser Tyr
20 25 30

45 Trp Met His Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met
35 40 45

Gly Glu Ile His Pro Asn Asp Gly Ser Thr Asn Tyr Asn Glu Lys Phe
50 55 60

50 Lys Gly Arg Val Thr Met Thr Arg Asp Thr Ser Thr Ser Thr Val Tyr
65 70 75 80

55 Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95

EP 3 338 793 A1

Ala Arg Trp Thr Leu Phe Thr Tyr Trp Gly Gln Gly Thr Leu Val Thr
100 105 110

5 Val

<210> 197
<211> 113
<212> PRT
<213> Artificial Sequence

<220>
<223> Humanized hSC17.155 VH5

<400> 197

Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala
1 5 10 15

Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Asn Ser Tyr
20 25 30

Trp Met His Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met
35 40 45

Gly Glu Ile His Pro Asn Gly Gly Ser Thr Asn Tyr Asn Glu Lys Phe
50 55 60

Lys Gly Arg Val Thr Met Thr Arg Asp Thr Ser Thr Ser Thr Val Tyr
65 70 75 80

Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95

Ala Arg Trp Thr Leu Phe Thr Tyr Trp Gly Gln Gly Thr Leu Val Thr
100 105 110

Val

<210> 198
<211> 113
<212> PRT
<213> Artificial Sequence

<220>
<223> Humanized hSC17.155 VH6

<400> 198

Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala
1 5 10 15

EP 3 338 793 A1

Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Asn Ser Tyr
20 25 30

5 Trp Met His Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met
35 40 45

10 Gly Glu Ile His Pro Asn Ser Gly Ser Thr Asn Tyr Asn Glu Lys Phe
50 55 60

Lys Gly Arg Val Thr Met Thr Arg Asp Thr Ser Thr Ser Thr Val Tyr
65 70 75 80

15 Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95

20 Ala Arg Trp Thr Leu Phe Thr Tyr Trp Gly Gln Gly Thr Leu Val Thr
100 105 110

25 Val

<210> 199
<211> 116
<212> PRT
30 <213> Artificial Sequence

<220>
<223> Humanized hSC17.161 VH1

35 <400> 199

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1 5 10 15

40 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Asp Ala
20 25 30

45 Trp Met Asp Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
35 40 45

Gly Glu Ile Arg Ser Lys Pro Asn Asn His Ala Thr Tyr Tyr Ala Glu
50 55 60

50 Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asp Ser Lys Asn Ser
65 70 75 80

55 Leu Tyr Leu Gln Met Asn Ser Leu Lys Thr Glu Asp Thr Ala Val Tyr
85 90 95

EP 3 338 793 A1

Tyr Cys Ala Arg Thr Gly Thr Ser Tyr Trp Gly Gln Gly Thr Leu Val
100 105 110

5 Thr Val Ser Ser
115

10 <210> 200
<211> 4
<212> PRT
<213> Homo sapiens

15 <220>
<221> MISC_FEATURE
<223> C-terminal cytoplasmic domain motif
<400> 200

20 Asn Pro Thr Tyr
1

25 <210> 201
<211> 9
<212> PRT
<213> Artificial Sequence

<220>
<223> His tag

30 <400> 201

His His His His His His His His His
1 5

35 <210> 202
<400> 202
000

40 <210> 203
<400> 203
000

45 <210> 204
<400> 204
000

50 <210> 205
<400> 205
000

<210> 206
<400> 206
000

55 <210> 206

EP 3 338 793 A1

	<400> 206 000
5	<210> 207 <400> 207 000
10	<210> 208 <400> 208 000
15	<210> 209 <400> 209 000
20	<210> 210 <400> 210 000
25	<210> 211 <400> 211 000
30	<210> 212 <400> 212 000
35	<210> 213 <400> 213 000
40	<210> 214 <400> 214 000
45	<210> 215 <400> 215 000
50	<210> 216 <400> 216 000
55	<210> 217 <400> 217 000
	<210> 218 <400> 218 000

<210> 219
 <400> 219
 000

5

<210> 220
 <211> 321
 <212> DNA
 <213> Mus musculus

10

<220>
 <221> misc_feature
 <223> Murine SC17.1 VL

15

<400> 220
 caaattgttc tcacccagtc tccagcactc atgtctgcat ctccaggga aaaggtctcc 60
 ctgacctgca gtgccaactc aactgtaagt ttcatgtact ggtaccagca gaagccaaga 120
 tcctcccca caccctggat ttatctcaca tccaacctgg cttctggagt ccctgctcgc 180
 ttcaagtggca gtgggtctgg gacctcttac tctcttaca tcagcagcat ggaggctgaa 240
 gatgctgcca cttattactg ccagcagtgg agtagtaact caccatcac gttcgggtgct 300
 gggaccaagc tggagctgaa a 321

25

<210> 221
 <211> 351
 <212> DNA
 <213> Mus musculus

30

<220>
 <221> misc_feature
 <223> Murine SC17.1 VH

35

<400> 221
 gatgtgcagc ttcaggactc aggacctggc ctggtgaaac cttctcagtc tctgtccgtc 60
 acctgcactg tcaactggta ctccatcacc tggggttatt actggaactg gatccggcag 120
 tttccaggaa acaaactgga gtggatgggt aacatacaca acagtgggtg cactaactac 180
 aacctatctc tcaagagtgc aatctctatc actogagaca catccaagaa ccagttcttc 240
 ctgcagttga attctgtgac tactgaggac acagccacat attactgtgc aaccacaaac 300
 tgggactact ttgactactg gggccaaggc accactctca cagtctcctc a 351

45

<210> 222
 <211> 336
 <212> DNA
 <213> Mus musculus

50

<220>
 <221> misc_feature
 <223> Murine SC17.2 VL

55

EP 3 338 793 A1

<400> 222
gacattgtga tgtcacagtc tccatcctcc ctggctgtgt cagttggaga gaaggtcact 60
atgagctgca agtccagtca gaggctttta tatagtagca atcaaaagag ctacttggcc 120
5 tgggtaccagc agaaaccagg gcagtctcct aaactgttaa tctactgggc atccactagg 180
gaatctgggg tccctgaccg cttcacaggc agtggatcag ggacagattt cactctcacc 240
10 atcagcagtg tgcaggctga agacctggcc gtttattact gcaagcaatc ttataatctt 300
cggacgttcg gtggaggcac caagctggaa atcaaa 336

<210> 223
15 <211> 351
<212> DNA
<213> Mus musculus

<220>
20 <221> misc_feature
<223> Murine SC17.2 VH

<400> 223
25 caggttcagc tgcagcagtc tgacgctgag ttggtgaaac ctggagcttc agtgaagata 60
tcctgcaagg tttctggcta caccttcact gaccatacta ttcactggat gaagcagagg 120
cctgaacagg gcctggaatg gattggatat atttattccta gagatggtag tactaagtac 180
30 aatgaggagt tcaagggcaa ggccacattg actgcagaca aatcctccag cacagcctac 240
atgcagctca acagcctgac atctgaggac tctgcagtct atttctgtgc aagatcatat 300
agtaactact ttgactactg gggccaaggc accactctca cagtctcctc a 351
35

<210> 224
40 <211> 333
<212> DNA
<213> Mus musculus

<220>
45 <221> misc_feature
<223> Murine SC17.9 VL

<400> 224
gacattgtga tgtcacagtc tccatcctcc ctagctgtgt cagttggaga gaaggttact 60
atgagctgca agtccagtca gaggctttta tatagtagca atcaaaagaa ctacttggcc 120
50 tgggtaccagc agaaaccagg gcagtctcct aaactgctga tttactgggc atccactagg 180
gaatctgggg tccctgatcg cttcacaggc agtggatctg ggacagattt cactctcacc 240
atcagcagtg tgaaggctga agacctggca gtttatttct gtcagcaata ttataactat 300
55 ccgtacacgt tcggaggggg gaccaagctg aaa 333

EP 3 338 793 A1

<210> 225
<211> 357
<212> DNA
<213> Mus musculus

5

<220>
<221> misc_feature
<223> Murine SC17.9 VH

10

<400> 225
caggtccaac tgcagcaacc tggggctgaa attgtgaggc ctggggcttc agtgaagctg 60
tcctgcaagg cttctggcta cacctttacc gactattgga tgaactgggt aaaacagagg 120
cctggacaag gccttgagtg gatcgagca attgatacct ctgatagtta tactagctac 180
aatccaaaat tcaagggcaa ggccacattg actgtagaca cctcctccag ctgagcctac 240
atgcagctca gcagcctgac atctgaggac tctgcggtct atttctgtgc aagaagagga 300
acctctggta aacctcttgt ttactggggc caagggactc tggtcactgt ctctgca 357

15

20

<210> 226
<211> 321
<212> DNA
<213> Mus musculus

25

<220>
<221> misc_feature
<223> Murine SC17.16 VL

30

<400> 226
gacatccaga tgactcagtc tccagcctcc ctatctgtat ctgtgggaga aactgtcacc 60
atcacatgtc gagcaagtgc gaatattaac agtaatttag tatggtatca gcagaaacag 120
ggaaaatctc ctgagctcct ggtctatgct gcaacaaact tagcggatgg tgtgccatca 180
cggttcagtgc gcagtggatc aggacacag tattccctca agatcaacag cctgcagtct 240
gaagattttg ggaattacta ctgtcaacat ttttggggta ctcctcggac gttcgggtgga 300
ggcaccaagc tggaaatcaa a 321

35

40

<210> 227
<211> 348
<212> DNA
<213> Mus musculus

45

<220>
<221> misc_feature
<223> Murine SC17.16 VH

50

<400> 227
gaggtccagc tgcaacagtc tggacctgag ctaatgaagc ctggggcttc agtgaagatg 60
tcctgcaagg cttctggata cacattcact gactacaaca tgtactgggt gaagcagaac 120

55

EP 3 338 793 A1

caaggaaaga gcctagagtg gataggagaa attaataccta acaatgggtg tactgcctac 180

aaccagaagt tcagaggcaa ggccacgttg actgtagaca agtcctccag cacagcctac 240

5 atggagctcc gcagcctgac atctgaggac tctgcagtct attactgtgc aagatatgat 300

aaggggtttg actactgggg ccaaggcacc actctcacag tctcctca 348

10 <210> 228
<211> 333
<212> DNA
<213> Mus musculus

15 <220>
<221> misc_feature
<223> Murine SC17.38 VL

20 <400> 228
gacattgtgg tcacccaatc tccagcttct ttggctgtgt ctctggggca gagagccacc 60

atctcctgca gagccagtga aagtgttgaa tattatggca caagtttaat gcagtggttc 120

caacagaaac caggacagcc acccaaactc ctcatctatg ctgcatccaa cgtagaatct 180

25 ggggtccctg ccaggttttag tggcagtggg tctgggacag acttcagcct caacatccat 240

cctgtggagg aggatgatat tgcaatgtat ttctgtcagc aagataggaa ggttccttgg 300

acgttcggtg gaggcaccaa gctggaaatc aaa 333

30 <210> 229
<211> 360
<212> DNA
<213> Mus musculus

35 <220>
<221> misc_feature
<223> Murine SC17.38 VH

40 <400> 229
caggttactc tgaaagagtc tggccctggg atattgcagc cctcccagac cctcagtcta 60

acttgttctt tctctgggtt ttactgaac acatctggta tgagttagg ctgggttcgt 120

45 cagccttcag ggaggggtct ggaatggctg gccccattt ggtggaatgg tgataagtac 180

tataaccag ccctgaaaag ccggctcaca atctccaagg atacctccaa caaccaggtt 240

ttcctcaaga tcgccagtgt ggtcactgca gatactgcca catacttctg tgctcgaata 300

50 cggcaatatt actatgctat ggactactgg ggtcaaggaa cctcagtcac cgtctcctca 360

55 <210> 230
<211> 336
<212> DNA
<213> Mus musculus

<220>
 <221> misc_feature
 <223> Murine SC17.3 VL

5 <400> 230
 gacattgtga tgtcacagtc tccatcctcc ctagctgtgt cagttggaga gaaggttact 60
 atgagctgca agtccagtc gagcctttta tatagtagca atcaaaagaa ctacttggcc 120
 10 tgggtaccagc agaaaccagg gcagtctcct aaactgctga tttactgggc atccactagg 180
 gaatctgggg tccctgatcg cttcacaggc agtggatctg ggacagattt cactctcacc 240
 atcagcagtg tgaaggctga agacctggca gtttattact gtcagcaata ttatagctat 300
 15 ccgacgttcg gtggaggcac caagctggaa atcaaa 336

<210> 231
 <211> 354
 <212> DNA
 20 <213> Mus musculus

<220>
 <221> misc_feature
 25 <223> Murine SC17.3 VH
 <400> 231
 caggtccaac tgcagcagcc tggggctgag cttgtgaagc ctggggcttc agtgaagctg 60
 30 tcctgcaagg cttctggcta caccttcccc agctactgga tacactgtgt gaagcagagg 120
 cctggacaag gccttgagtg gattggagtg attaataccta gcaacggctg tactaactac 180
 aatgagaagt tcaagaacaa ggccacactg actgtagaca aatcctccag cacagcctac 240
 35 atgcaactca gcagcctgac atctgaggac tctgcggtct attactgtgt cagggggggg 300
 acgggctata ctatggacta ctggggctcaa ggaacctcag tcaccgtctc ctca 354

<210> 232
 40 <211> 321
 <212> DNA
 <213> Mus musculus

<220>
 <221> misc_feature
 45 <223> Murine SC17.4 VL
 <400> 232
 gacatcaaga tgaccagtc tccatcttcc atgtatgcct ctctaggaga gagagtcact 60
 50 atcacttgca aggcgagtc ggacattaat agctatttaa cctggttcca gcagaaacca 120
 gggaaatctc ctaagaccct gatctatcgt gcaaacagat tgatagatgg ggtcccatca 180
 55 aggttcagtg gcagtggatc tgggcaagat tattctctca ccatcagcag cctggattat 240
 gaagatatgg gaatttatta ttgtctacag tatgatgact ttccgtggac gttcgggtgga 300

ggcaccaagc tggaaatcaa a

321

5
 <210> 233
 <211> 348
 <212> DNA
 <213> Mus musculus

10
 <220>
 <221> misc_feature
 <223> Murine SC17.4 VH

<400> 233
 cagatccagt tgggtgcagtc tggacctgag ctgaagaagc ctggagagac agtcaagatc 60
 15
 tcctgcaagg cttctgggta taccttcaca gactattcaa tgcactgggt gaagcaggct 120
 ccaggaaagg gtttaaagtg gctgggctgg ataaacactg agactggcga gccaacatat 180
 tcagaagact tcaaggagc gtttgccctc tctttggaaa cctctgccag cactgcctat 240
 20
 ttgcagatca acaacctcaa aaatgaagac acggctactt atttctgtgt taaaaataag 300
 ggctgggttg cttattgggg ccaagggact ctggctactg tctctgca 348

25
 <210> 234
 <211> 336
 <212> DNA
 <213> Mus musculus

30
 <220>
 <221> misc_feature
 <223> Murine SC17.8 VL

<400> 234
 35
 gatgttgatga tgacccaaac tccactctcc ctgcctgtca gtcttggaga tcaagcctcc 60
 atctcttgca gatctagtca gagccttgta cacagtaatg gagacaccta ttacattgg 120
 tacctgcaga agccaggcca gtctccaaaa ctctgatct acaaagtttc caaccgattt 180
 40
 tctggggtcc cagacagggt cagtggcagt ggatcaggga cagatttcac actcaagatc 240
 agcagagtgg aggctgagga tctgggactt tatttctgct ctcaaagtac acttattccg 300
 tacacgttcg gaggggggac caagctggac ataaaa 336

45
 <210> 235
 <211> 342
 <212> DNA
 <213> Mus musculus

50
 <220>
 <221> misc_feature
 <223> Murine SC17.8 VH

55
 <400> 235
 caggttcacc tgcagcagtc tggaaactgaa gtgatgaagc ctggggcctc agtgaagata 60

EP 3 338 793 A1

	tctctgcaagg ctactggcta cacattcagt agctactgga tagagtggat aaagcagagg	120
	cctggacatg gccttgagtg gattggagag attttgcctg gaagtggtaa tactaacaac	180
5	aatgagaagt tcaagggcaa ggccacaatc actgcagata catcctccaa tatagcctac	240
	atacaattaa gcagcctgac atctgaggac tctgccgtct attactgtgc gggaggcccg	300
	gcggttact ggggccaagg gactctggtc actgtctctg ca	342
10	<210> 236 <211> 339 <212> DNA <213> Mus musculus	
15	<220> <221> misc_feature <223> Murine SC17.10 VL	
20	<400> 236	
	gacattgtga tgtcacagtc tccatcctcc ctagctgtgt cagttggaga gaaggttact	60
	atgagctgca agtccagtca gaggctttta tatagtagca atcaaaagaa ctacttggcc	120
25	tggtaccagc agaaaccagg gcagtctcct aaactgctga tttactgggc atccactagg	180
	gaatctgggg tccctgatcg cttcacaggc agtggatctg ggacagattt cactctcacc	240
	atcagcagtg tgaagactga agacctggca ctttattact gtcagcaata ttattggttt	300
30	ccgtacacgt tcggaggggg gaccaagctg gaaataaaa	339
35	<210> 237 <211> 339 <212> DNA <213> Mus musculus	
40	<220> <221> misc_feature <223> Murine SC17.10 VH	
	<400> 237	
	gaggttcagc tgcagcagtc tggggcagaa cttgtgaagc caggggcctc agtcaagttg	60
45	tcttgcacag cttctggctt caacattaaa gacacctata tgcactgggt gaagcagagg	120
	cctgaacagg gcctggagtg gattggaagg attgatcctg cgaatgttaa tactaaatat	180
	gaccogaagt tccagggcaa ggccactata acagcagaca catcctccaa cacagcctac	240
50	ctgcagctca gcagcctgac atctgaggac actgccgtct attactgtgt tagggggaat	300
	gtttactggg gccaaaggac tctggtcact gtctctgca	339
55	<210> 238 <211> 324 <212> DNA	

EP 3 338 793 A1

<213> Mus musculus

<220>

<221> misc_feature

<223> Murine SC17.11 VL

<400> 238

```

gaaaatgtgc tcacccagtc tccagcaatc atgtctgcat ctccagggga aaaggtcacc      60
atgacctgca gggccagctc aagtgttaagt tccagttact tgcactggta ccagcagaag      120
tcaggtgcct cccccaact ctggatttat agcacatcca acttggtctc tggagtccct      180
gctcgcttca gtggcagtg gtctgggacc tcttactctc tcacaatcag cagtgtggag      240
gctgaagatg ctgccactta ttactgccag cagtacagtg attacccatt cacgttcggc      300
tcggggacaa agtttgtaat aaaa                                           324

```

<210> 239

<211> 366

<212> DNA

<213> Mus musculus

<220>

<221> misc_feature

<223> Murine SC17.11 VH

<400> 239

```

gaggtccagc tgcagcagtc tggacctgag ctggtgaaac ctggggcttt agtgatgatg      60
tcctgcaagg cttctggata cacattcact gactactaca tgcactgggt gaagcagagc      120
catggacaga gccttgagtg gattggagag gttattcctt acaatgatga aactttctac      180
aaccggaagt tcaaggacaa ggccacattg actgtagaca aatcctctag tacagcctac      240
atggagctcc ggagcctgac atctgaggac tctgcaatct attattgtgc aagaagacat      300
aggtacgacg ggtttcgtta tgctatagac tactggggtc aaggaacctc agtcaccgtc      360
tcctca                                                                366

```

<210> 240

<211> 336

<212> DNA

<213> Mus musculus

<220>

<221> misc_feature

<223> Murine SC17.14 VL

<400> 240

```

gatgttttga tgacccaaac tccactctcc ctgcctgtca gtcttggaga tcaagcctcc      60
atctcttgca gatctagtca gagcattgtc catagtaatg gaaacaccta tttagagtgg      120
ttcctgcaga aaccaggcca gtctccaaag ctctctgatct acaaagtttc caaccgattt      180

```

EP 3 338 793 A1

tctgggggtcc cagacaggtt cagtggcagt ggatcagga cagatttcac actcaagatc 240
 agcagagtgg aggctgagga tctgggagtt tattactgct ttcaagggtc acatgttccg 300
 5 tacacgttcg gaggggggac caagctggaa ataaaa 336

<210> 241
 <211> 354
 <212> DNA
 10 <213> Mus musculus

<220>
 <221> misc_feature
 15 <223> Murine SC17.14 VH

<400> 241
 gaggtccagc tgcaacagtc tggacctgtg ctggtgaagc ctggggcttc agtgaagatg 60
 tcctgtaagg cttctggata cacaatcact gactacaata tgaactgggt gaagcagagc 120
 20 catggaaaga gccttgagtg gattggagtt attaatacctt acaacggtaa tactagatat 180
 aaccagatgt tcaagggcaa ggccacattg actgttgaca agtcctccag cacagcctac 240
 25 atggagctca acagcctgac atctgaggac tctgcagtct attactgtac aagatgggggt 300
 actacggtgg taggtgcgaa ctggggccaa ggcaccactc tcacagtctc ctca 354

<210> 242
 <211> 318
 <212> DNA
 30 <213> Mus musculus

<220>
 <221> misc_feature
 35 <223> Murine SC17.15 VL

<400> 242
 caaattgttc tcacccagtc tccagcactc atgtctgcat ctccagggga gaaggtcacc 60
 40 atgacctgca gtgccagctc aagtgtaaat tacatgtact ggtaccagca gaagccaaga 120
 tcctccccca aaccctggat ttatctcaca tccaacctgg cttctggagt ccctgttcgc 180
 45 ttcagtggca gtgggtctgg gacctcttac tctctcacia tcagcagcat ggaggctgaa 240
 gatgctgcca ctactactg ccagcagtgg agtaataacc caccacggtt cggttctggg 300
 accaagctgg agctgaaa 318

<210> 243
 <211> 366
 <212> DNA
 50 <213> Mus musculus

<220>

55

<221> misc_feature
<223> Murine SC17.15 VH

<400> 243
5 gacgtgaagc tcgtggagtc tgggggaggc ttagtgaagc ttggagggtc cctgaaactc 60
tcctgtgcag cctctggatt cactttcagt agctatgcc tgtcttgggt tcgccagact 120
ccggagaaga ggctggagtg ggtcgcaacc attactagt gtggtggtta cacctactat 180
10 ccagacagtg tgaagggtcg attcaccatc tccagagaca atgccaagaa caccctgtac 240
ctgcaaatga gcagtttgaa gtctgaggac acggccatgt attactgtgc aagaagggat 300
tactacggta gtagttacgt tatgtttgct tattggggcc aagggaactct ggtcactgtc 360
15 tctgca 366

<210> 244
<211> 318
20 <212> DNA
<213> Mus musculus

<220>
25 <221> misc_feature
<223> Murine SC17.17 VL

<400> 244
caaattgttc tcaccagtc tccagcaatc atgtctgcat ctccagggga gaaggtcacc 60
30 atgacctgca gtgccagctc aagtgttaagt tacatgcact ggtaccagca gaagtcaggc 120
acctccccca aaagatggat ttatgacaca tccaaactgc cttctggagt ccctgctcgc 180
ttcagtggca gtgggtctgg gacctcttac tctctcacia tcagcagcat ggaggctgaa 240
35 gatgctgcc cttattactg ccagcagtgg agtagtacct caccacggt cggtgctggg 300
accaagctgg agctgaaa 318

<210> 245
40 <211> 360
<212> DNA
<213> Mus musculus

<220>
45 <221> misc_feature
<223> Murine SC17.17 VH

<400> 245
50 gaggtccagc tgcaacagtc tggacctgag gtaatgaagc ctggggcttc agtgaagatg 60
tcctgcaagg cttctggata cacattcact gactacaaca tgcactgggt gaagcagaac 120
caaggaaaga gcctagagt gataggagaa attaataccta acattggtgg tactggctac 180
55 aaccagaagt tcaaaggcaa ggccacattg actgtacaca agtcctccag cacagcctac 240
atggagctcc gcagcctgac atctgaggac tctgcagtct attactgtgc aagaacctat 300

agttactata gttacgagtt tgcttactgg ggccaaggga ctctgggtcac tgtctctgca 360

5 <210> 246
 <211> 321
 <212> DNA
 <213> Mus musculus

10 <220>
 <221> misc_feature
 <223> Murine SC17.18 VL

<400> 246
 15 gacatccaga tgacacaatc ttcatacctac ttgtctgtat ctctaggagg cagagtcacc 60
 attacttgca aggcaagtga ccacattaat aattgggttag cctgggtatca gcagaaacca 120
 ggaaatgctc ctaggctctt aatatctggt gcaaccagtt tggaaactgg gggttccttca 180
 20 agattcagtg gcagtggatc tggaaaggat tacactctca gcattaccag tcttcagact 240
 gaagatgttg ctacttatta ctgtcaacag tattggagta ttccggtcac gttcgggtgcg 300
 gggaccaagc tggagctgaa a 321

25 <210> 247
 <211> 369
 <212> DNA
 <213> Mus musculus

30 <220>
 <221> misc_feature
 <223> Murine SC17.18 VH

<400> 247
 35 caggttactc tgaaagagtc tggccctggg atattgcagc cctcccagac cctcagtcctg 60
 acttggttctt tctctggggtt ttcactgagc acttctacta tgggtgtagg ctggattcgt 120
 cagccttcag gaaagggtct agagtggctg gcagacattt ggtgggatga cagtaagtac 180
 40 tataatccat ccctgaagag ccggctcaca atctccaagg atacctccag caaccaggta 240
 ttctctcaaga tcaccagtgt ggacactgca gatactgcc a ttactactg tgcgcgaaag 300
 45 ggaaggacag ctcgggctac gagagggttt gcttactggg gccacgggac tctgggtcact 360
 gtctctgca 369

50 <210> 248
 <211> 327
 <212> DNA
 <213> Mus musculus

55 <220>
 <221> misc_feature
 <223> Murine SC17.19 VL

EP 3 338 793 A1

<400> 248
 gacattgtgc tgaccaatc tccagcttct ttggctgtgt ctctagggca gagggccgcc 60
 atctcttgca agcccagcca aagtgttgat tatgatggtg atagttatat gaactggtac 120
 5 caacagaaac caggccagcc acccaaactc ctcatctatg ctgcatccaa tctagaatct 180
 gggatcccag ccagggttag tggcagtggg tctgggacag acttcaccct caacatccat 240
 10 cctgtggagg aggaggatgc tgcaacctat tactgtcacc aaattaatga cgatccgtgg 300
 acgttcggtg gaggcaccaa gctgaaa 327

<210> 249
 <211> 363
 <212> DNA
 <213> Mus musculus

<220>
 <221> misc_feature
 <223> Murine SC17.19 VH

<400> 249
 gatgtgcagc ttcaggagtc aggacctggc ctggtgaaac cttctcagtc tctgtctgtc 60
 25 acctgcactg tctactggcta ctccatcacc agtagttata cctggaactg gatccggcag 120
 tttccaggaa acaaactgga gtggatgggc tacatacatt acagtggtag cactaactac 180
 30 aacctatctc tcagaagtcg aatctctatt actcgagaca cgtccaagaa ccagttcttc 240
 ctgcagttga attctgtgac tactgaggac acagccacat attactgtgc aagatcccgt 300
 tattactacg atgcttacgg gtttgcttac tggggccaag ggactctggt cactgtctct 360
 35 gca 363

<210> 250
 <211> 336
 <212> DNA
 <213> Mus musculus

<220>
 <221> misc_feature
 <223> Murine SC17.22 VL

<400> 250
 gatgttgtgt tgacccaaac tccactctcc ctgcctgtca gtcttgagga tcaagcctcc 60
 atctcttgca gatctagtca gagcattgta cacattaata gacacaccta cttaggatgg 120
 50 tacctgcaga aaccaggcca gtcgctaaag ctccctgatat atgggggttc caaccgattt 180
 tctgggggtcc cagacagggt cagtggcagt ggatcaggga cagatttcac actcaagatc 240
 55 agcagagtgg aggtctagga tatgggagtt tattactgct ttcaaggtag acatgttcca 300
 ttcacgttcg gctcggggac aaagttggaa ataaaa 336

EP 3 338 793 A1

<210> 251
<211> 363
<212> DNA
<213> Mus musculus

5

<220>
<221> misc_feature
<223> Murine SC17.22 VH

10

<400> 251
cagatccaga tgatgcagtc tggacctgag ctgaagaagc ctggagagac agtcaagatc 60
tcctgcaagg cttctgggta ttccttcaca aactatggaa tgaactgggt gaagcaggct 120
ccaggaaagg gtttaaagtg gatgggctgg ataaacacct aactggaga gccaacatat 180
gctgatgact tcaagggacg gtttgccttc tctttggaaa cctctgccag cactgcctat 240
ttgcagatca acaacctcaa aaatgaggac atggctacat atttctgtac aagaggttac 300
tacggtagta gctacgatgc tttggactac tggggccaag gaacctcagt caccgtctcc 360
tca 363

15

25

<210> 252
<211> 336
<212> DNA
<213> Mus musculus

30

<220>
<221> misc_feature
<223> Murine SC17.24 VL

35

<400> 252
gacattgtga tgtcacagtc tccatcctcc ctggctgtgt cagttggaga gaaggctact 60
atgagctgca agtccagtca gagcctttta tatagtagca atcaaaagag ctacttggcc 120
tggtaccagc agaaaccagg gcagtctcct aaactgttaa tctactgggc atccactagg 180
gaatctgggg tccctgaccg cttcacaggc agtggatcag ggacagattt cactctcacc 240
atcagcagtg tgcaggctga agacctggcc gtttattact gcaagcaatc ttataatctt 300
cggacgttcg gtggaggcac caagctggaa atcaaa 336

45

<210> 253
<211> 351
<212> DNA
<213> Mus musculus

50

<220>
<221> misc_feature
<223> Murine SC17.24 VH

55

<400> 253

EP 3 338 793 A1

caggttcagc tgcagcagtc tgacgctgag ttggtgaaac ctggagcttc agtgaagata 60
tcttgcaagg tttctggcta caccttcact gaccatacta ttactggat gaagcagagg 120
5 cctgaacagg gcctggaatg gattggatat atttatccta gagatggtag tactaagtac 180
aatgaggagt tcaagggcaa ggccacattg actgcagaca aatcctccag cacagcctac 240
atgcagctca acagcctgac atctgaggac tctgcagtct atttctgtgc aagatcatat 300
10 agtaactact ttgactactg gggccaaggc accactctca cagtctcctc a 351

<210> 254
<211> 336
15 <212> DNA
<213> Mus musculus

<220>
<221> misc_feature
20 <223> Murine SC17.27 VL

<400> 254
gatgttgtga tgaccagac tccactcact ttgtcggta ccattggaca accagcctcc 60
atctcttgca agtcaagtca gagcctctta gaaagtgatg gaaagacata tttgaattgg 120
25 ttgttacaga ggccaggcca gtctccaaag cgcctaactct atctgggtgtc taaactggac 180
tctggagtcc ctgacaggtt caccgggcagt ggatcaggga cagatttcac actgaaaatc 240
agcagagtgg aggctgagga tttgggagtt tattattgct ggcaaggat acaacatcct 300
30 cggacgttcg gtggaggcac caagctggaa atcaaa 336

<210> 255
35 <211> 339
<212> DNA
<213> Mus musculus

<220>
40 <221> misc_feature
<223> Murine SC17.27 VH

<400> 255
caggttcaac tgcagcagtc tggggctgag ctggtgaggc ctggggcttc agtgacgctg 60
45 tcttgcaagg cttcgggcta cacatttact gactatgaaa tgactgggt gaagcagaca 120
cctgtgcatg gcctggaatg gattggaggt attgatcctg aaactgggtg tactgcctac 180
aatcagaagt tcaagggcaa ggccacactg actgcagaca aatcctccag cacagcctac 240
50 atggagctcc gcagcctgac atctgaggac tctgccgtct acttctgtac aagatggttt 300
tcttactggg gccagggac tctggtcact gtctctgca 339

55 <210> 256
<211> 321

EP 3 338 793 A1

<212> DNA
<213> Mus musculus

5 <220>
<221> misc_feature
<223> Murine SC17.28 VL

<400> 256
10 gacatcttgc tgactcagtc tccagccatc ctgtctgtga gtccaggaga aggagtcagt 60
ttctcctgca gggccagtca gagcattggc acaagcatac actggtatca gcaaagaaca 120
aatggttctc caagacttct cataaagtat gcttctgagt ctatctctgg gatcccttct 180
15 aggttttagtg gcagtgggtc agggacagat tttactcttc gcatcaacag tctggagtct 240
gaagatattg cagattatta ctgtcaacaa agtaatagct ggccactcac gttcggtgct 300
gggaccaagc tggagctgaa a 321

20 <210> 257
<211> 351
<212> DNA
<213> Mus musculus

25 <220>
<221> misc_feature
<223> Murine SC17.28 VH

30 <400> 257
caggctccacc tgccgcagtc tagacctgaa ctggtgaagc ctggagcttc agtgaagata 60
tcctgcaagg cttctggcta cggcttcaca cgcagctata tacactgggt gaagcagagg 120
35 cctggacagg gcctagagtg gattggatat atttcttctg gaagtgggtg tactacctac 180
aatcagaagt ttaaggggcaa ggcctcactg actgcagaca atccctccag cactgcctac 240
atgcatctca gtagcctgac atctgaggac tctgcgatct atttctgtgc aagagggggg 300
40 gtacgggtact tcgatgtctg gggcgcaggg accacgggtca ccgtctcctc a 351

45 <210> 258
<211> 321
<212> DNA
<213> Mus musculus

50 <220>
<221> misc_feature
<223> Murine SC17.29 VL

<400> 258
55 gacattgtga tgaccagtc tcacaaattc atgtccacat cagtaggaga cagggtcagc 60
atcacctgca aggccagtca ggatgtgggt actgatgtag cctggtatca acagaaacca 120
gggcaatctc ctaaactact gatttactgg gcatccaccc ggcaactgg agtccttgat 180

EP 3 338 793 A1

cgcttcacag gcagtggatc tgggacagat ttcactctca ccattagcaa tgtgcagtct 240

gaagacttgg cagattatct ctgtcagcaa tatagcagct atccgtacac gttcggaggg 300

5 gggacaaaagc tggaataaaa a 321

<210> 259
<211> 348
<212> DNA
10 <213> Mus musculus

<220>
<221> misc_feature
15 <223> Murine SC17.29 VH

<400> 259
gaggtccagc tgcaacagtc tggacctgag ctaatgaagc ctggggcttc agtgaagatg 60

tcctgcaagg cttctggata cacattcact gactacaaca tgcactgggt gaagcagaac 120

20 caaggaaaga gcctagagtg gattggagaa attaatactc acaatggtgg tactggctac 180

aaccagaagt tcaaaggcaa ggccacattg actgtagaca agtcctccag cacatcctac 240

25 atggagctcc gcagcctgac atctgaggac tctgcagtct attactgtgc aggcggttac 300

ccggcctttg actactgggg ccaaggcacc actctcacag tctcctca 348

<210> 260
30 <211> 324
<212> DNA
<213> Mus musculus

<220>
35 <221> misc_feature
<223> Murine SC17.30 VL

<400> 260
gaaaatgtgc tcacccagtc tccagcaatc gtgtctgcat ctccagggga aaaggtcacc 60

40 atgacctgca gggccagctc aagtgtatt tccagttact tgcactggta ccagcagaag 120

tcagggtgcct ccccaaaact ctggatttat agcacatcca acttggcttc tggagtccct 180

45 gctcgcttca gtggcagtc gtctgggacc tcttactctc tcacaatcag cagtgtggag 240

gctgaagatg ctgccactta ttactgccag cagtacagtg gttaccgct cacgttcggt 300

gctgggacca agctggagct gaaa 324

50 <210> 261
<211> 372
<212> DNA
<213> Mus musculus

55 <220>
<221> misc_feature

EP 3 338 793 A1

<223> Murine SC17.30 VH

<400> 261

gaagtgaagc tgggtggagtc tgagggaggc ttagtgcagc ctggaagttc catgaaactc 60
 5 tcctgcacag cctctggatt cactttcagt gactattaca tggcttgggt ccgccaggtt 120
 ccagaaaagg gtctagaatg ggttgcaaac attaattatg atggtagtag cacttactat 180
 10 ctggactcct tgaagagccg tttcatcatc tcgagagaca atgcaaagaa cattctatac 240
 ctgcaaatga gcagtctgaa gtctgaggac acagccacgt attactgtgc aagagatgat 300
 tattacggta gtagcccaag ctactggtac ttcgatgtct ggggcgcagg gaccacggtc 360
 15 accgtctcct ca 372

<210> 262

<211> 321

<212> DNA

<213> Mus musculus

<220>

<221> misc_feature

<223> Murine SC17.32 VL

<400> 262

gacatccaga tgactcagtc tccagcctcc ctatctgcat ctgtgggaga aactgtcacc 60
 30 atgacatgtc gagcaagtgg gaatattcac aattatttag tatggtatca gcagaaacag 120
 ggaaaatctc ctacgtcct ggtctataat gcaaaaacct tagcagatgg tgtgccatca 180
 aggttcagtg gcagtgatc aggaacacaa tattctctca agatcaacag cctgcagcct 240
 35 gaagattttg ggagttatta ctgtcaacat ttttgagta ctctccgac gttcgggtgga 300
 ggcaccaagc tggaatcaa a 321

<210> 263

<211> 357

<212> DNA

<213> Mus musculus

<220>

<221> misc_feature

<223> Murine SC17.32 VH

<400> 263

gaagtgaaac ttgaggagtc tggaggaggc ttggtgcaac ctggaggatc catgaaactc 60
 50 tcctgtgttg cctctggatt cactttcagt aactactgga tgagctgggt ccgccagtct 120
 ccagagaagg ggcttgagtg ggttgctgaa attagattga aatctaataa ttatgcaaca 180
 55 cattatgcgg agtctgtgaa agggagggtc accatctcaa gagacgattc caaaagtagt 240
 gtcttcctgc aaatgaacaa cttaagaact gaagacactg gcatttatta ctgtaccagg 300

cactattact atgctatgga ctactggggg caaggaacct cagtcaccgt ctctca 357

5 <210> 264
 <211> 321
 <212> DNA
 <213> Mus musculus

10 <220>
 <221> misc_feature
 <223> Murine SC17.34 VL

<400> 264
 gacatcaaga tgaccagtc tccatcttcc atgtatgcat ctctaggaga gagagtcact 60
 15 atcacctgca aggcgagtc ggacattaat agctatttaa gctgggtcca gcagaaacca 120
 gggaaatctc ctaagaccct gatctatcgt gcaaacagat tggtagatgg ggtcccatca 180
 aggttcagtg gcagtggatc tgggcaagat tattctctca ccatcagtag cctggagtat 240
 20 gaagatatgg gaatttatta ttgtctacag tatgatgagt ttcctccgac gttcgggtgga 300
 ggcaccaagc tggaatcaa a 321

25 <210> 265
 <211> 354
 <212> DNA
 <213> Mus musculus

30 <220>
 <221> misc_feature
 <223> Murine SC17.34 VH

<400> 265
 35 gaggtccagc tacaacagtc tggacctgag ctggtgaagc ctgggtcttc agtgaagata 60
 tcctgcaagg cttctggata cacattcact gactacaaca tggactgggt gaagcagagc 120
 catggaaaga gacttgagtg gattggatat atttattctg acaatggtgg tgctggctac 180
 40 aaccagaagt tcaagggcaa ggccacattg actgtagaca agtcctccag cacagcctac 240
 atggagctcc gcagcctgac atctgaggac tctgcagtct attactgttc aagatccatt 300
 actacggctt ggtttgctta ctggggccaa gggactctgg tcaactgtctc tgca 354

45 <210> 266
 <211> 324
 <212> DNA
 <213> Mus musculus

50 <220>
 <221> misc_feature
 <223> Murine SC17.35 VL

55 <400> 266
 gaaaatgtgc tcaccagtc tccagcaatc atgtctgcat ctccagggga aaaggtcacc 60

EP 3 338 793 A1

	ctgacctgca gggccagctc aagtatgagt tccagttact tgcactggta ccagcagaag	120
	tcaggtgcct cccccaaact ctggatttat agcacatcca acttggcttc tggagtcctt	180
5	gctcgcttca gtggcagtggt gtctgggacc tcttactctc tcacaatcag cagtgtggag	240
	gctgaagatg ctgccactta ttactgccag cagtacagtg cttacccatt cacgttcggc	300
	tcggggacaa agttggaaat aaaa	324
10	<210> 267 <211> 363 <212> DNA <213> Mus musculus	
15	<220> <221> misc_feature <223> Murine SC17.35 VH	
20	<400> 267	
	gaggtccagc tgcagcagtc tggacctgag ctagtgaaac ctggggcttt agtgaagatg	60
	tcctgcaagg cttctggata cacattcact gactactaca tacactgggt gaagcagagc	120
25	catggaaaga gccttgagtg gattggagaa attaatcctt acaatgggtga gactttctac	180
	aaccagaagt tcaagggcaa ggccacattg actgtagaca aatcctctag tacagcctac	240
	atggaactcc ggagcctgac atctgaggac tctgcagctc attattgtgc aagaagggga	300
30	tggatatctaa caggctatgc tatggactac tggggctcaag gaacctcagt caccgtctcc	360
	tca	363
35	<210> 268 <211> 318 <212> DNA <213> Mus musculus	
40	<220> <221> misc_feature <223> Murine SC17.36 VL	
45	<400> 268	
	caaattgttc tcaccagtc tccagcactc atgtctgcat ctccagggga gaaggtcacc	60
	atgacctgca gtgccagctc aagtgtgaagt tacatgtact ggtaccagca gaagccaaga	120
	tcctcccca aaccctggat ttatctcaca tccaacctgg cttctggagt ccctgctcgc	180
50	ttcagtgcca gtgggtctgg gacctcttac tctctcacia tcagcagcat ggaggctgaa	240
	gatgctgcc cttattactg ccagcagtggt agtagtaacc caccacggtt cggagggggg	300
	accaagctgg aaataaaa	318
55	<210> 269	

EP 3 338 793 A1

<211> 360
<212> DNA
<213> Mus musculus

5

<220>
<221> misc_feature
<223> Murine SC17.36 VH

10

<400> 269
gaggtgcagc ttcaggagtc aggacctagc ctctgtgaaac cttctcagtc tcagtccctc 60
acctgttctg tcaactggcg ctcctatcacc agtgattact ggaactggat ccggaattc 120
ccagggaaga aagttgagta catggggtac ataaactaca gtggtagcac ttactacaat 180
ccatctctca aaagtcgaat ctccatcact cgagacacat ccaagaacca gtactacctg 240
cagttgaact ctgtgacttc tgaggacaca gccacatatt actgtgcacg tacctcgtac 300
tataataagt ttctaccatt tgcttactgg ggccaagga ctctgggtcac tgtctctgca 360

20

<210> 270
<211> 336
<212> DNA
<213> Mus musculus

25

<220>
<221> misc_feature
<223> Murine SC17.39 VL

30

<400> 270
gatgttttaa tgacccaaac tccactctcc ctgcctgtca gtcttggaga tcaagcctcc 60
atctcttgca gatctagtca gagtcttgta cacagaaatg gaaacaccta ttttcattgg 120
tacctgcaga agccaggcca gtctccaaag ctctctgatct acaaagtttc caaccgattt 180
tctggggtcc cagacaggtt cagtggcagt ggatcaggga cagatttcac actcaagatc 240
agcagagtgg aggctgagga tctgggagtt tatttctgct ctcaaagtac atatgttccg 300
tggaacgttcg gtggaggcac caagctggaa atcaaa 336

40

<210> 271
<211> 351
<212> DNA
<213> Mus musculus

45

<220>
<221> misc_feature
<223> Murine SC17.39 VH

50

<400> 271
gaggtgcagc tgggtggagtc tgggggaggc ttagtgcagc ctggagggtc ccggaactc 60
tcctgtgcag cctctggatt cactttcagt agctatggaa tgcaactgggt ccgtcaggct 120
ccagagaagg ggctggagtg ggtcgcatat attagtagta acgatgggtac catctactat 180

55

EP 3 338 793 A1

gcagacacag tgaggggccg attcaccatc tccagagaca atgccaagaa caccctgttc 240
 ttgcaaataa ccagtctaag gtctgaggac acggccatgt attactgtgc aagaccttct 300
 5 aactgggtct ttgactactg gggccaaggc accactctca cagtctcctc a 351

<210> 272
 <211> 336
 <212> DNA
 10 <213> Mus musculus

<220>
 <221> misc_feature
 15 <223> Murine SC17.40 VL

<400> 272
 gatgttgtga tgacccaaac tccactctcc cggcctgtca ctcttggaga tcaagcctcc 60
 atctcttgca gatctagtca gaggcttgta cacagtaatg gaaacaccta ttacattgg 120
 20 tacctgcaga agccaggcca gtctccaaag ctctgatct acaaagtttc caaccgattt 180
 tctgggggtcc ctgacagggt cagtggcagt ggatcaggga cagatttcac actcaagatc 240
 25 agcagagtgg aggctgagga tctgggagtt tatttctgct ctcaaaatac acatgttcca 300
 ttcacgttcg gctcggggac aaagttggaa ataaaa 336

<210> 273
 <211> 348
 <212> DNA
 30 <213> Mus musculus

<220>
 <221> misc_feature
 35 <223> Murine SC17.40 VH

<400> 273
 cagggtccaac tgcagcagcc tggggctgaa attgtgaggc ctggggcttc agtgaagctg 60
 40 tcttgcaagg cttctggcta cacctttacc gactattgga tgaactgggt gaagcagagg 120
 cctggacaag gccttgagtg gatcggaaca attgatcctt ctgatagtta tactcgttac 180
 45 aatcaaaagt tcaagggcaa ggccacattg actgtagaca catccttcag ctgagcctac 240
 atgcagctca gcagcctgac atctgaggac tctgcggtct atttctgtgc aagtggggga 300
 cggggggttg gttactgggg ccaagggact ccgggtcactg tctctgta 348

<210> 274
 <211> 318
 <212> DNA
 50 <213> Mus musculus

<220>

EP 3 338 793 A1

<221> misc_feature
<223> Murine SC17.41 VL

<400> 274
 5 caaattgttc tcacccagtc tccagcactc atgtctgcat ctccagggga gaaggtcacc 60
 atgacctgca gtgccagctc aagtgttaagt tacatgtact ggtaccagca gaagccaaga 120
 tcctccccc aaccctggat ttatctcaca tccaacctgg cttctggagt ccctactcgc 180
 10 ttcagtggca gtgggtctgg gacctcttac tctctcaca tcagcagcat gggggctgaa 240
 gatgctgcc cttattactg ccagcagtgg aatactaacc caccacggt cggtgctggg 300
 accaagctgg agctgaaa 318

<210> 275
 <211> 363
 <212> DNA
 <213> Mus musculus

<220>
 <221> misc_feature
 <223> Murine SC17.41 VH

<400> 275
 25 gacgtgaagc tcgtggagtc tgggggaggc ttagtgaagc ttggagggtc cctgaaactc 60
 tcctgtgcag cctctggatt cactttcagt agctatgcca tgtcttgggt tcgccagact 120
 30 ccggagaaga ggctggagtg ggtcgcaacc attagtagtg gtggtggtaa cacctactat 180
 ccagacagtg tgaagggtcg attcaccatc tccagagaca atgccaagaa caccctgtac 240
 ctgcaaatga gcagtttgaa gtctgaggac acggccatgt attactgtgc aagaagggat 300
 35 tactacggta ctagctacgt tatgtttgct tactggggcc aagggactct ggtcactgtc 360
 tct 363

<210> 276
 <211> 318
 <212> DNA
 <213> Mus musculus

<220>
 <221> misc_feature
 <223> Murine SC17.42 VL

<400> 276
 50 gaaaatgttc tcacccagtc tccagcaatc atgtctgcat ctccagggga aaaggtcacc 60
 atgacctgta gtgccagctc aagtgtaaat tacatgtact ggtaccagca gaagtcaagc 120
 acctccccc aactctggat ttatgacaca tccaaactga cttctggagt ccaggtcgc 180
 55 ttcagtggca gtgggtctgg aaactcttac tctctcacga tcagcaacat ggaggctgaa 240
 gatgttgcca cttattactg ttttcagggg agtgggtacc cactcacgtt cggtcgggg 300

acaaaattgg aaataaaa

318

<210> 277
 <211> 336
 <212> DNA
 <213> Mus musculus

<220>
 <221> misc_feature
 <223> Murine SC17.42 VH

<400> 277
 gacgtgaagc tggaggagtc ggggggaggc ttagtgaggc ctggagggtc cctgaaactc 60
 tcctgtgcag cctctggatt cactttcagt agatatacca tgtcttgggt tcgccagaca 120
 ccggagaaga ggctggagtg ggccgcaacc attaatagtg gtggtagtaa cacctactat 180
 ccagacagtg tgaagggccg attcaccatc tccagagaca atgccaagaa caccctgttc 240
 ctgcaaataga gcagtctgaa gtctgaggac acagccatgt attactgtac aaatggtaac 300
 cactggggcc aaggcaccac tctcacagtc tcctca 336

<210> 278
 <211> 318
 <212> DNA
 <213> Mus musculus

<220>
 <221> misc_feature
 <223> Murine SC17.45 VL

<400> 278
 gaaaatgttc tcacccagtc tccagcaatc atgtctgcat ctccaggga aaaggtcacc 60
 atgacctgta gtgccagctc aagtgtaaat tacatgtact ggtaccagca gaagtcaagc 120
 acctcccca aactctggat ttatgacaca tccaaactga cttctggagt ccaggtcgc 180
 ttcagtggca gtgggtctgg aaactcttac tctctcacga tcagcaacat ggaggctgaa 240
 gatgttgcca cttattactg ttttcagggg agtgggtacc cactcacgtt cggctcgggg 300
 acaaaaattgg aaataaaa 318

<210> 279
 <211> 342
 <212> DNA
 <213> Mus musculus

<220>
 <221> misc_feature
 <223> Murine SC17.45 VH

<400> 279

EP 3 338 793 A1

	caggtgcaac tgcagcagcc tgggtctgtg ctggtgaggc ctggagattc agtgaagctg	60
	tcgtgcaagg cttctggcta cacattcacc agctactgga tgcactgggt gaagcagagc	120
5	cctggacaag gccttgagtg gattggagag attcatcctc atagtggtag tactaactac	180
	aatgagaagt tcaagggcaa ggccacactg actgtagaca catcctccag cacagcctac	240
	gtggatctca gcagcctgac atctgaggac tctgcggctc attactgtgt aggtggtcac	300
10	tacgactact ggggccaagg caccactctc acagtctcct ca	342
	<210> 280	
	<211> 321	
15	<212> DNA	
	<213> Mus musculus	
	<220>	
	<221> misc_feature	
20	<223> Murine SC17.46 VL	
	<400> 280	
	agttttgtga tgacccaaac tcccaaattc ctgcttgtat cagcaggaga cagggttacc	60
25	ataacctgca aggccagtca gagtgtgaat aatgatgtag cttggtacca acagaagcca	120
	gggcagtctc ctaaaactgct gatatactat gcatccaatc gctacactgg agtccctgat	180
	cgcttcaactg gcagtggata tgggacggat ttcaactttca ccatcagcac tgtgcaggct	240
30	gaagacctgg cagtttatatt ctgtcagcag gattatagct ctccctcggac gttcggtgga	300
	ggcaccaagc tggaaatcaa a	321
	<210> 281	
35	<211> 357	
	<212> DNA	
	<213> Mus musculus	
	<220>	
40	<221> misc_feature	
	<223> Murine SC17.46 VH	
	<400> 281	
45	caggtccaac tgcagcagcc tgggtgctgag cttgtgaagc ctggggcctc aatgaagctg	60
	tcctgcaagg cttctggcta cactttcacc agctactgga taaactgggt gaagcagagg	120
	cctggacaag gccttgagtg gattggaaat atttttcctg atactactac tactaactac	180
50	aatgagaagt tcaagagcaa ggccacactg actgtagaca catcctccag cacagcctat	240
	atgcagctca gcagcctgac atctgacgac tctgcggctc attattgtgc aaggaggtac	300
	tacgatggta cctacgatgc tatggattac tggggccaag gaacctcagt caccgtc	357
55	<210> 282	
	<211> 318	

EP 3 338 793 A1

<212> DNA
<213> Mus musculus

5 <220>
<221> misc_feature
<223> Murine SC17.47 VL

<400> 282
10 caaattgttc tcaccagtc tccagcaatc atgtctgcat ctccagggga gaaggtctcc 60
atgacctgca gtgccagctc aagtgttaagt tacatgcact ggtaccagca gaagtcaggc 120
acctcccca aaagatggat ttatgacaca tccaaactgg cttctggagt ccctgctcgc 180
15 ttcagtggca gtgggtctgg gtcctcttac tctctcacia tcagcagcat ggaggctgaa 240
gatgctgccca cttattactg ccagcagtggt agtagcacc ccccccacgtt cggagggggg 300
accaagctgg aaataaaa 318

20 <210> 283
<211> 357
<212> DNA
<213> Mus musculus

25 <220>
<221> misc_feature
<223> Murine SC17.47 VH

30 <400> 283
gagggtccagc tgcagcagtc tggacctgag ctggtgaagc ctggggcttc agtgaagata 60
tcctgcaagg cttctgggta ctattcact gactactaca tgcgctgggt gaagcaaagt 120
35 cctgaaaaga gccttgagtg gattggagag attaataccta gcactgggtg tactacctac 180
aaccagaact tcaaggccaa ggccacattg actgtagaca aatcctccag cacagcctac 240
atgcagctca agagcctgac atctgaggac tctgcagtct attactgtgc aagagggggg 300
40 tacttcttgt actactttga ctactggggc caaggcacca ctctcacagt ctctca 357

45 <210> 284
<211> 336
<212> DNA
<213> Mus musculus

50 <220>
<221> misc_feature
<223> Murine SC17.49 VL

<400> 284
gatgttgtga tgaccagac tccactcact ttgtcgggta ccattggaca accagcctcc 60
55 atctcttgca agtcaagtca gagcctctta gaaagtgatg gaaagacata tttgaattgg 120
ttgttacaga ggccaggcca gtctccaaag cgcctaactct atctgggtgc taaactggac 180

EP 3 338 793 A1

tctggagtcc ctgacaggtt cacgggcagt ggatcagga cagatttcac actgaaaatc 240
 agcagagtgg aggctgagga tttgggagtt tattattgct ggcaaggtat acaacatcct 300
 5 cggacgttcg gtggaggcac caagctggaa atcaaa 336

<210> 285
 <211> 339
 <212> DNA
 10 <213> Mus musculus

<220>
 <221> misc_feature
 15 <223> Murine SC17.49 VH

<400> 285
 caggttcaac tgcagcagtc tggggctgag ctggtgagga ctggggcttc agtgacgctg 60
 tcctgcaagg cttcgggcta cacatttact gactatgaaa tgcactgggt gaagcagaca 120
 20 cctgtgcatg gcttggaatg gattggaggt attgatcctg aaactgggtg tactgcctac 180
 aatcagaagt tcaagggcaa ggccacactg actgcagaca aatcctccag cacagcctac 240
 atggagctcc gcagcctgac atctgaggac tctgccgtct acttctgtac aagatggttt 300
 25 tcttactggg gccagggac tctggtcact gtctctgca 339

<210> 286
 <211> 333
 <212> DNA
 30 <213> Mus musculus

<220>
 <221> misc_feature
 35 <223> Murine SC17.50 VL

<400> 286
 gacattgtgc tgacacagtc tcctgcttcc ttagctgcat ctctggggca gagggccacc 60
 40 atctcatgca gggccagcca aagtgtcagt acatctagct atagttatat gcactggtac 120
 caacagaaac caggacagcc acccaaactc ctcatcaagt atgcatcaa cctagaatct 180
 ggggtccctg ccaggttcag tggcagtggt tctgggacag acttcaccct caacatocat 240
 45 cctgtggagg aggaggatac tgcaacatat tactgtcagc acagttggga gattccgtgg 300
 acgttcggtg gaggcaccaa gctggaaatc aaa 333

<210> 287
 <211> 363
 <212> DNA
 50 <213> Mus musculus

<220>
 <221> misc_feature
 55

EP 3 338 793 A1

<223> Murine SC17.50 VH

<400> 287

gaggtgcagc tgggtggagtc tgggggaggc ttagtgaagc ctggagggtc cctgaaactc 60

tcctgtgcag cctctggatt cactttcagt gactatggaa tgcactgggt tcgtcaggct 120

ccagagaagg ggctggagtg ggttgcatac attagtagtg gcagtagaac catctactat 180

gcagacacag tgaagggccg attcaccatc tccagagaca atgccaagaa caccctgttc 240

ctgcaaataa ccagtctgag gtctgaggac acggccatgt attactgtgc aagggtttac 300

tacggaagta cctacgggta tttcgatgtc tggggcacag ggaccacggt caccgtctcc 360

tca 363

<210> 288

<211> 333

<212> DNA

<213> Mus musculus

<220>

<221> misc_feature

<223> Murine SC17.53 VL

<400> 288

gacattgtgc tgacacagtc tcctgcttcc ttagctgcat ctctggggca gagggccacc 60

atctcatgca gggccagtca aagtgtcagt acatctagct atagttatat gcactggtac 120

caacagaagc caggacatcc acccaaactc ctcatcaggt atgcatccaa cctagagtct 180

ggggtccttg ccaggttcag tggcagtggt tctgggacag acttcaccct caacatccat 240

cctgtggagg aggaggatac tgcaacatat tactgtcagc acagttggga gattccgtac 300

acgttcggag gggggaccaa gctggaaata aaa 333

<210> 289

<211> 357

<212> DNA

<213> Mus musculus

<220>

<221> misc_feature

<223> Murine SC17.53 VH

<400> 289

gaggtccagc ttcagcagtc aggacctgag ctggtgaaac ctggggcctc agtgaagata 60

tcctgcaagg cttctggata cacattcact gactacaaca tgcaactgggt gaagcagagc 120

catggaaagc gccttgagtg gattggatat attcatcctt acaatggtgg tagtggctac 180

aaccagaagt tcaagaggaa ggccacattg actgtagaca attcctccaa cacaacctac 240

atggagctcc gcagcctgac atctgaggac tctgcagtct attactgtgc aagatcttat 300

EP 3 338 793 A1

gattacgaca cctggtttgg ttactggggc caagggactc tggtcactgt ccgtgca 357

<210> 290
 <211> 336
 <212> DNA
 <213> Mus musculus

<220>
 <221> misc_feature
 <223> Murine SC17.54 VL

<400> 290
 gatgttgtgc tgaccagac tccactcact ttgtcgggta ccattggaca accagcctcc 60
 atctcttgca agtcaagtca gagcctctta tatagtgatg gaaagacata tttgaattgg 120
 ttgttacaga ggccaggcca gtctccaaag cgcctaactct atctggtgtc taaactggac 180
 tctggagtcc ctgacagggt cactggcagt ggatcagga cagatttcac actgaaaatc 240
 agcagagtgg aggctgagga tttgggactt tattattgct ggcaaggtag acattttccg 300
 tggacgttcg gtggaggcac caagctggaa atcaaa 336

<210> 291
 <211> 339
 <212> DNA
 <213> Mus musculus

<220>
 <221> misc_feature
 <223> Murine SC17.54 VH

<400> 291
 gaagtgaac ttgaggagtc tggaggaggc ttggtgcaac ctggaggatc catgaaactc 60
 tcctgtgttg cctctggatt cactttcagt aactactgga taaactgggt ccgccagtct 120
 ccagagaagg ggcttgagtg ggttgctgaa atcagaatga aatctaataa ttatgcaaca 180
 cattatgcgg agtctgtgaa agggagggtc accatctcaa gagatgattc caaaagtgtg 240
 gtctacctgc aaatgaacaa cttaagacct gaagacactg gcatttatta ctgtaccagg 300
 gggggctact ggggccaagg caccactctc accgtctcc 339

<210> 292
 <211> 339
 <212> DNA
 <213> Mus musculus

<220>
 <221> misc_feature
 <223> Murine SC17.56 VL

<400> 292
 gacattgtga tgtcacagtc tccatcctcc ctagctgtgt cagttggaga gaaggttact 60

EP 3 338 793 A1

atgagctgca agtccagtca gaggcctttta tatagtagca atcaaaagaa ctacttggcc 120
 tgggtaccagc agaaaccagg gcagtctcct aaactgctga ttacttgggc atccactagg 180
 5 gaatctgggg tccctgatcg cttcacaggc agtggatctg ggacagattt cactctcacc 240
 atcagcagtg tgaaggctga agacctggca gtttatttct gtcagcaata ttataactat 300
 ccgtacacgt tcggaggggg gaccaagctg gaaataaaa 339
 10
 <210> 293
 <211> 348
 <212> DNA
 <213> Mus musculus
 15
 <220>
 <221> misc_feature
 <223> Murine SC17.56 VH
 20
 <400> 293
 cagatccagt tgggtgcagtc tggacctgaa ctgaagaagc ctggagagac agtcaagatc 60
 tcctgcaagg cttctgggta taccttcaca aactatggaa tgaactgggt gaagcaggct 120
 25 ccaggaaagg gtttaaagtg gatggcctgg ataaacacct aactggaga gccaacatat 180
 gctgatgact tcaagggacg gtttgccttc tctttggaaa cctctgccag cactgcctct 240
 ttgcagatca tcaacctcaa aaatgaggac acggctacat atttctgtgc aaggatcggc 300
 30 gatagtagtc cctctgacta ctggggccag ggcaccactc tcacagtc 348
 <210> 294
 <211> 324
 35 <212> DNA
 <213> Mus musculus
 <220>
 <221> misc_feature
 40 <223> Murine SC17.57 VL
 <400> 294
 caaattgttc tcacccagtc tccagcaatc atgtctgcat ctctagggga acgggtcacc 60
 45 atgacctgca ctgccagctc aagtgttaagt tccagttact tgcaactgta ccagcagaag 120
 ccaggatcct ccccaaaact ctggatttat agcacatcca acctggcttc tggagtcca 180
 cctcgcttca gtggcagtg gtctgggacc tcttactctc tcacaatcag cagcatggag 240
 50 gctgaagatg ctgccactta ttactgccac cagtatcatc gttccccacc gacgttcggt 300
 ggaggcacca agctggaaat caaa 324
 55 <210> 295
 <211> 363
 <212> DNA

<213> Mus musculus

<220>

<221> misc_feature

<223> Murine SC17.57 VH

<400> 295

cagatccagt tgggtgcagtc tggacctgaa ctgaagaagc ctggagagac agtcaagatc 60

tcctgcaagg cttctgatta taccttcaca gacttttcaa tacactgggt gaggcagtct 120

ccaggaaagg gtttaaagtg gatgggctgg ataaacactg agactgggtga gccaacagtt 180

gcagaagact tcaagggacg gtttgccttc tctttggaga cctctgccag cactgccttt 240

ttgcagatct acaacctcaa aaatgaggac tcggcaacat atttctgtgc tagggggcgt 300

tactacggcc atgactatgc tatggactac tggggtaag gaacctcagt caccgtctcc 360

tca 363

<210> 296

<211> 321

<212> DNA

<213> Mus musculus

<220>

<221> misc_feature

<223> Murine SC17.59 VL

<400> 296

gacatccaga tgactcagtc tccagcctcc ctatctgcat ctgtgggaga aactgtcacc 60

atcacatgtc gagcaagtgg gaatcttcac aattatttag catgggtatca gcagaaacag 120

ggaaaatctc ctcagctcct ggtctataat gcaaaaacct tagcagatgg tgtgccatca 180

aggttcagtg gcagtggatc aggaacacaa tattctctca agatcaacag cctgcagcct 240

gaagattttg ggacttattt ctgtcaacat ttttggagta ttcctccac gttcgggggg 300

gggaccaagc tggaataaaa a 321

<210> 297

<211> 357

<212> DNA

<213> Mus musculus

<220>

<221> misc_feature

<223> Murine SC17.59 VH

<400> 297

gaagtgaagc ttgaggagtc tggaggaggc ttggtgcaac ctgggggatc catgaaactc 60

tcctgtgttg cctctggatt cactttcagt aactattgga tgaactgggt ccgccagtct 120

ccagagaagg ggcttgagtg ggttgctgaa attagattga aatctaataa ttatgcaaca 180

EP 3 338 793 A1

cattatgcgg agtctgtgaa agggaggttc accatctcaa gagatgattc caaaagtagt 240
gtctacctgc aaatgaacaa cttaagagct gaagacactg gcatttatta ctgtaccaga 300
5 ctctgggact ttgctatgga ctactggggt caaggaacct cagtcaccgt ctccctca 357

<210> 298
<211> 318
10 <212> DNA
<213> Mus musculus

<220>
<221> misc_feature
15 <223> Murine SC17.61 VL

<400> 298
caaattgttc tcacccagtc tccagcaatc atgtctgcat ctccagggga gaaggtcacc 60
20 atatcctgca gtgccagctc aagtgttaagt tacatatact ggtaccagca gaagccagga 120
tcctccccc aacctggat ttatcgcaca tccaacctgg cttctggagt ccctgctcgc 180
ttcagtggca gtgggtctgg gacctcttac tctctcacia tcagcagcat ggaggctgaa 240
25 gatgctgcca cttattactg ccagcagtat catagttacc cgtggacgtt cgggtggaggc 300
accaagctgg aaatcaaa 318

<210> 299
<211> 375
30 <212> DNA
<213> Mus musculus

<220>
<221> misc_feature
35 <223> Murine SC17.61 VH

<400> 299
40 caggttactc tgaaagagtc tggccctggg atattgcagc cctcccagac cctcagtctg 60
acttgttctt tctctgggtt ttactgagc acttttggtg tgggtgtagg ctggattcgt 120
cagccttcag ggaagggctc ggagtggctg gcacagattt ggtgggatga ttataagtac 180
45 tataacccag ccctgaagag tcggctcaca atctccaagg atacctcaa aaaccaggta 240
ttcctcaaga tcgccaatgt ggacactgca gatactgcca catactactg tgctcgaatc 300
ggatattact ccggtagtag ccgttgctgg tacttcgatg tctggggcac agggagcacg 360
50 gtcaccgtct cctca 375

<210> 300
<211> 321
55 <212> DNA
<213> Mus musculus

<220>
 <221> misc_feature
 <223> Murine SC17.63 VL

5 <400> 300
 agtattgtga tgaccagac tcccaaattc ctgcttgtat cagcaggaga cagggttgcc 60
 ataacctgca aggccagtca gagtgtgagt aatgatgtag cttggtacca acagaagcca 120
 10 gggcagtctc ctacactgct gatatacctat gcatccaatc gctacactgg agtccctgat 180
 cgcttcactg gcagtggata tgggacggat ttcaactttca ccatcagcac tgtgcaggct 240
 gaagacctgg cagtttattt ctgtcagcag ggttatagct ctccgttcac gttcggaggg 300
 15 gggaccaagc tggaaataaa a 321

<210> 301
 <211> 348
 <212> DNA
 20 <213> Mus musculus

<220>
 <221> misc_feature
 25 <223> Murine SC17.63 VH
 <400> 301
 caggttcagc tgcaacagtc tgacgctgag ttggtgaaac ctggggcttc agtgaagata 60
 tcctgcaagg ctgctggcta caccttcact gaccttacta ttactgggt gaaacagagg 120
 30 cctgaacagg gcctggagtg gattggatat atttatcctg gagatagtaa tactaagtac 180
 aatgagaagt tcaagggcaa ggccacattg actgcagata aatcctccag cactgcctat 240
 35 atgcagctca acagcctgac atctgaggat tctgtagtgt atttctgtgc aagaatgatt 300
 actccttact actttgacta ctggggccaa ggcaccactc tcacagtc 348

<210> 302
 40 <211> 321
 <212> DNA
 <213> Mus musculus

<220>
 <221> misc_feature
 45 <223> Murine SC17.71 VL
 <400> 302
 gacatccaga tgactcagtc tccagcctcc ctatctgcct ctgtgggaga aactgtcacc 60
 50 atcgcatgtc gagcaagtgg gaatattcac aattatttaa catggtatca gcagagacag 120
 ggaaaatctc ctacagctcct ggtctataat gcaaaaacct tagcagttgg tgtgccatca 180
 55 aggttcagtg gcagtggctc aggaacacaa tattctctca agatcaacag cctgcagcct 240
 gaagattttg ggagttatta ctgtcaacat ttttgggaata ctccctccgac gttcgggtgga 300

ggcaccaagc tggaaatcaa a

321

<210> 303
 <211> 357
 <212> DNA
 <213> Mus musculus

<220>
 <221> misc_feature
 <223> Murine SC17.71 VH

<400> 303
 gaagtgaagc ttgaggagtc tggaggaggc ttggtgcaac ctggaggatc catgaaactc 60
 tcctgtgttg cctctggaat cattttcagt aactactgga tgaattgggt ccgccagtct 120
 ccagagaagg ggcttgagtg ggttgctgaa attagattga aatctaataa ttattcaaca 180
 cattatgcgg agtctgtgaa agggagggtc accatctcaa gagatgattc caaaagtagt 240
 gtctacctgc aaatgaacaa cttaagagct gaagacactg gcatttatta ctgtaccagg 300
 cactattact atgctatgga ctactggggt caaggaacct cagtcaccgt ctctca 357

<210> 304
 <211> 321
 <212> DNA
 <213> Mus musculus

<220>
 <221> misc_feature
 <223> Murine SC17.72 VL

<400> 304
 gatatccaga tgacacagac tacatcctcc ctgtctgcct ctctgggaga cagagtcacc 60
 atcagttgca gtgcaagtca gggcattagc aattatttaa actggtatca gcagaaacca 120
 gatggaactg ttaaactcct gatctattac acatcaagtt tacactcagg agtcccatca 180
 aagttcagtg gcagtgggtc tgggacagat tattctctca ccatcagcaa cctggaacct 240
 gaagatatcg ccacttacta ttgtcagcag tatagtaagc ttccgtacac gttcggaggg 300
 gggaccaagc tggaaataaa a 321

<210> 305
 <211> 357
 <212> DNA
 <213> Mus musculus

<220>
 <221> misc_feature
 <223> Murine SC17.72 VH

<400> 305

EP 3 338 793 A1

gaggtgcagc tgggtggagtc tgggggaggc ttagtgaagc ctggagggtc cctgaaactc 60
tctctgtcag cctctggatt cactttcagt agctatggca tgtcttgggt tcgccagact 120
5 ccggagaaga ggctggagtg ggtcgagcc attaatagta atggtggttag cacctactat 180
ccagacactg tgaagggccg actcaccatc tccagagaca atggcaagaa caccctgtac 240
ctgcaaatga gcagtctgag gtctgaggac acagccttgt attactgtgt aagggatgat 300
10 ggttactacg ttttctttgc ttactggggc caagggactc tggtcactgt ctctgca 357

<210> 306
<211> 321
15 <212> DNA
<213> Mus musculus

<220>
<221> misc_feature
20 <223> Murine SC17.74 VL

<400> 306
gacatccaga tgacacaatc ttcacacctac ttgtctgtat ctctaggagg cagagtcacc 60
attacttgca aggcaagtga ccacattaat aattgggttag cctgggtatca gcagaaacca 120
25 ggaaatgctc ctaggctctt aatatctggt gcaaccagtt tggaaactgg ggttccttca 180
agattcagtg gcagtggatc tggaaaggat tacactctca gcattaccag tcttcagact 240
gaagatgttg ctacttatta ctgtcaacag tattggagta ctccctccac gttcgggtgt 300
30 gggaccaagc tggagctgaa a 321

<210> 307
35 <211> 357
<212> DNA
<213> Mus musculus

<220>
40 <221> misc_feature
<223> Murine SC17.74 VH

<400> 307
caggtgcagc tgaagcagtc aggacctggc ctagtggcgc cctcacagag cctgtccatc 60
45 acatgcactg tctctgggtt ctcatcacc agctatggtg tagactgggt tcgccagtct 120
ccaggaaagg gtctggagtg gctgggagtg atatggggtg gtggaagcac aaattataat 180
tcagctctca aatccagact gagcatcacc aaggacaact ccaagagcca agttttctta 240
50 aaaatgaaca gtctgcaaac tgatgacaca gccatgtact actgtgccag tggagactac 300
gatggtagcc tctggtttgc ttactggggc caagggactc tggtcactgt ctctgca 357

55 <210> 308
<211> 336

EP 3 338 793 A1

<212> DNA
<213> Mus musculus

5 <220>
<221> misc_feature
<223> Murine SC17.76 VL

<400> 308
10 gatattgtga taaccagga tgaactctcc aatcctgtca cttctggaga atcagtttcc 60
atctcctgca ggtctagtaa gagtctccta tataaggatg ggaagacata cttgaattgg 120
tttctgcaga gaccaggaca atctcctcag ctctgatct atttgatgtc caccctgca 180
15 tcaggagtct cagaccggtt tagtggcagt gggtcaggaa cagatttcac cctggaaatc 240
agtagagtga aggtcgagga tgtgggtgtg tattactgtc aacaacttgt agagtatcct 300
cggacgttcg gtggaggcac caagctggaa atcaaa 336

20 <210> 309
<211> 339
<212> DNA
<213> Mus musculus

25 <220>
<221> misc_feature
<223> Murine SC17.76 VH

30 <400> 309
gagggtgcagc tgggtggagtc tgggggagac ttagtgaagc ctggagggtc cctgaaactc 60
tcctgtgtag cctctggatt cactttcagt agctatggca tgtcttgggt tcgccagact 120
35 ccagacaaga ggctggagtg ggtcgcaacc attagtagtg gtggtacttt cacctactat 180
ccagacagtg tgaaggggagc attcaccgtc tccagagaca atgccaagaa caccctgtac 240
ctgcaaataa gcagtctgaa gtctgaggac acagccatgt attactgttc aagacatggg 300
40 tggggctggg gccaaaggac tctggtcact gtctctgca 339

45 <210> 310
<211> 321
<212> DNA
<213> Mus musculus

50 <220>
<221> misc_feature
<223> Murine SC17.77 VL

<400> 310
55 gacatccaga tgactcagtc tccagcctcc ctatctgcat ctgtgggaga aactgtcacc 60
atcacatgtc gagcaagtgg gaatattcac aattatttag catggtatca gcagaaacag 120
ggaaaatctc ctacgtcct ggtctataat gcaaaagcct tagcagatgg tgtgccatca 180

EP 3 338 793 A1

aggttcagtgc gcagtggatc aggaacacaa tattctctca agatcaacag cctgcagcct 240

gaagattttg ggagttatta ctgtcaacat ttttggagta ttcctccgac gttcgggtgga 300

5 ggcaccaagc tggaaatcaa a 321

<210> 311
 <211> 357
 <212> DNA
 <213> Mus musculus

10

<220>
 <221> misc_feature
 <223> Murine SC17.77 VH

15

<400> 311
 gaagtgaagc ttgaggagtc tggaggaggc ttggtgcaac ctggaggatc catgaaactc 60
 tcctgtgttg cctctggatt cactttcagt aactactgga tgaactgggt ccgccagtct 120
 20 ccagagaagg ggcttgagtgc ggttgctgaa attagattga aatctaataa ttatgcaaca 180
 cattatgcgg agtctgtgaa agggagggtc accatctcaa gagatgattc caaaagtagt 240
 gtctacctgc aaatgaacaa cttaagagtt gaagacactg ccatttatta ctgtaccagg 300
 25 cactatgact atgctatgga ctactggggc caaggaacct cagtcaccgt ctccctca 357

<210> 312
 <211> 321
 <212> DNA
 <213> Mus musculus

30

<220>
 <221> misc_feature
 <223> Murine SC17.79 VL

35

<400> 312
 gacatccaga tgactcagtc tccagcctcc ctatctgcat ctgtgggaga aactgtcacc 60
 40 atcacatgtc gagcaagtgc gaatattcac aattatttag catggatatca gcagaaacag 120
 ggaaaatctc ctgagctcct ggtctataat gcaaaaacct tagcagatgg tgtgccatca 180
 aggttcagtgc gcagtggatc aggaacacaa tattctctca ggatcaacag cctgcagcct 240
 45 gaagattttg ggagttatta ctgtcaacat ttttggagta ttcctccgac gttcgggtgga 300
 ggcaccaagg tggaaatcaa a 321

50

<210> 313
 <211> 357
 <212> DNA
 <213> Mus musculus

55

<220>
 <221> misc_feature

EP 3 338 793 A1

<223> Murine SC17.79 VH

<400> 313

gaagtgaagc ttgaggagtc tggaggaggc ttggtacaac ctggaggatc catgaaactc 60
 5 tctctgtgttg cctctggatt cactttcagt gactactgga tgaactgggt ccgccagtct 120
 ccagagaagg ggcttgagtt ggttgctgaa attagattga tatctaataa ttatgcaaca 180
 10 cattatgcgg agtctgtgaa agggagggtc accatctcaa gagatgattc caaaagtagt 240
 gtctacctgc aaatgaacaa cttaagagct gaagacactg gcatttatta ctgtaccagg 300
 cactattact atgcttttga ctactggggc caaggaacct cagtcaccgt ctctca 357

<210> 314

<211> 339

<212> DNA

<213> Mus musculus

<220>

<221> misc_feature

<223> Murine SC17.81 VL

<400> 314

gacattgtga tgtcacagtc tccatcctcc ctaactgtgt cagttggaga gaaggttact 60
 ttgagctgca agtccagtc gagcctttta tatagtacca atcaaaagat ctacttggcc 120
 30 tgggtaccagc agaaaccagg gcagtctcct aaactgctga ttactgggc atccactagg 180
 gaatctgggg tccctgatcg cttcacaggc agtggatctg ggacagattt cactctcgcc 240
 atcagcaatg tgaaggctga agacctggca gtttattact gtcagcaata ttatagctat 300
 35 ccgtacacgt tcggaggggg gaccaagctg gaaataaaa 339

<210> 315

<211> 339

<212> DNA

<213> Mus musculus

<220>

<221> misc_feature

<223> Murine SC17.81 VH

<400> 315

gaggttcagc tgcagcagtc tggggcagag cttgtgaagc caggggcctc agtcaagttg 60
 50 tctgtcacag cttctggctt caacattaat gacacctatt accattgggt gaagcagagg 120
 cctgaacagg gcctggagtg gattggaagg attgatcctg cgaatgttaa tactaaatat 180
 gacccgaagt tccagggcaa ggccacttta acagcagaca catcctccaa cacagcctac 240
 55 ctgcagctca gcagcctgac atctgaggac actgccgtct attactgtgg tagggggaat 300
 gcttactggg gccaaaggac tctggtcact gtctctgca 339

<210> 316
 <211> 312
 <212> DNA
 <213> Mus musculus

5

<220>
 <221> misc_feature
 <223> Murine SC17.82 VL

10

<400> 316
 caaattgttc tcacccagtc tccagcaatc atgtctgcat ctctagggga ggagatcacc 60
 ctaacctgca gtgccagttc gagtgttaagt tacatgcact ggtaccagca gaagtcaggc 120
 acttctccca aactcttgat ttatagcaca tccaacctgg cttctggagt cccttctcgc 180
 ttcagtggca gtgggtctgg gaccttttat tctctcacia tcagcagtgt ggaggctgaa 240
 gatgctgccg attattactg ccatcagtgg agtagtttca cgttcggctc ggggacaaaag 300
 ttggaaataa aa 312

20

<210> 317
 <211> 360
 <212> DNA
 <213> Mus musculus

25

<220>
 <221> misc_feature
 <223> Murine SC17.82 VH

30

<400> 317
 gaggtccagc tgcaacagtc tggacctgag ctggtgaagc ctggggcttc agtgaagatg 60
 tcctgtaagg cttctggata cacattcact gactcctaca tgaactgggt gaagcagagt 120
 catggaaaga gccttgagtg gattggacgt gttaatccta acaatgggtg tgctagctac 180
 aaccacaagt tcaagggcaa ggccacattg acagtagaca aatccctcag cacagcctac 240
 atgcgcctca acagcctgac atctgaggac tctgcggctc attactgttc aagatctgga 300
 gacctttatt actatgctat ggactactgg ggtcaaggaa cctcagtcac cgtctcctca 360

40

<210> 318
 <211> 318
 <212> DNA
 <213> Mus musculus

45

<220>
 <221> misc_feature
 <223> Murine SC17.84 VL

50

<400> 318
 caaattgttc tcacccagtc tccagcaatc atgtctgcat ctccagggga gaaggtcacc 60
 atgacctgca gtgccagctc aagtataagt tacatgcact ggtaccagca gaagtcaggc 120

55

EP 3 338 793 A1

acctccccca aaagatggat ttatgacaca tccaaactgg cttctggagt ccctgctcgc 180

ttcagtggca gtgggtctgg gacctcttac tctctcacia tcagcaacat ggaggctgaa 240

5 gatgctgccca cttattactg ccagcagtgg agtagtacct caccacggt cggagggggg 300

accaagctgg aaataaaa 318

10 <210> 319
<211> 351
<212> DNA
<213> Mus musculus

15 <220>
<221> misc_feature
<223> Murine SC17.84 VH

20 <400> 319
gaggtccagt tgcaacagtc tggacctgag ctaatgaagc ctggggcttc agtgaagatg 60

tcctgcaagg cttctggata tatatttact gactacaaca tgcactgggt gaagcagaac 120

caaggaaaga gcctagagtg gataggagaa gttaatccta aactgggtgg tattggctac 180

25 aatcagaaat tcaaaggcaa ggccacattg actgtagaca agtcctccag cacagcctac 240

atggacctcc gcagcctgac atctgaggac tctgcagtct attactgtgc aagagatggc 300

aattattgct ttgactactg gggccaaggc accactctca cagtctcctc a 351

30 <210> 320
<211> 333
<212> DNA
<213> Mus musculus

35 <220>
<221> misc_feature
<223> Murine SC17.85 VL

40 <400> 320
gatattgtga tgacgcaggc tgcattctcc aatccagtca ctcttggaac atcagcttcc 60

atctcctgca ggtctagtaa gagtctccta catagtaatg gcatcactta tttgtattgg 120

45 tatctgcaga agccaggcca gtctcctcag ctcttgattt atcagatgtc caaccttgcc 180

tcaggagtcc cagagagggt cagtagcagt gggtcaggat ctgatttcac actgagaatc 240

agcagagtgg aggctgagga tgtgggtgtt tattactgtg ctcaaaatct agaacatccg 300

50 acgttcggtg gaggcaccaa gctggaaatc aaa 333

55 <210> 321
<211> 360
<212> DNA
<213> Mus musculus

<220>
 <221> misc_feature
 <223> Murine SC17.85 VH

5 <400> 321
 gaggtgcagc tgggtggagtc tgggggagac ttagtgaagc ctggaggggtc cctgaaactc 60
 tcctgtgcag cctctggatt cactttcagt aactatggca tgtcttgggt tcgccagact 120
 10 ccagacaaga ggctggagtg ggtcgcaacc attagtactg gtggtactta cacctactat 180
 ccagacagtg tgaaggggag attcaccatc tccagagaca atgccaagaa caccctgtac 240
 ctgcaaataa gcagtctgaa gtctgaggac acagccatgt attactgtgt aggacagtcc 300
 15 tatagtgact acgtctcggt tgcttattgg ggccaaggga ctcagggtcac tgtctctgca 360

<210> 322
 <211> 336
 <212> DNA
 20 <213> Mus musculus

<220>
 <221> misc_feature
 25 <223> Murine SC17.87 VL

<400> 322
 gatgtttgtga tgacccaaac tccactctcc ctgcctgtca gtcttggaga tcaagcctcc 60
 atctcttgca gatctagtca gagccttgta cacagtaatg gaaacaccta ttacattgg 120
 30 tacctgcaga agccaggcca gtctccaaag ctccctgatct ccaaagtttc caaccgattt 180
 tctgggggtcc cagacaggtt cagtggcagt ggatcaggga cagatttcac tctcaagatc 240
 35 agcagagtgg aggctgaaga tctgggagtt tatttctgct ctcaaagtac acatgttcct 300
 cccatgttcg gaggggggac caggctggaa ataaaa 336

<210> 323
 40 <211> 345
 <212> DNA
 <213> Mus musculus

<220>
 <221> misc_feature
 45 <223> Murine SC17.87 VH

<400> 323
 gaggttcagc tgcagcagtc tggggctgag cttctgaagc caggggcctc agtcaagttg 60
 50 tcctgcacag cttctggcct caacattaaa gactactata taaactgggt gtaccagagg 120
 cctgaacagg gcctggagtg gattggaagg attgatcctg agagtataa tactttatat 180
 55 gaccogaagt tccagggcaa ggccagtata acagcagaca catctccaa cacagcctac 240
 ctgcagctca gcagcctgac atctgaggac actgccgtct attactgtac tactaatacc 300

EP 3 338 793 A1

ccttttgctt actggggcca agggactctg gtcactgtct ctaca 345

5 <210> 324
<211> 336
<212> DNA
<213> Mus musculus

10 <220>
<221> misc_feature
<223> Murine SC17.89 VL

<400> 324
gatgttttga tgacccaaac tccactctcc ctgcctgtca gtcttggaga tcaagcctcc 60
15 atctcttgca gatctagtca gagcattgta catagtaatg gaaacaccta tttagaatgg 120
tacctgcaga aaccaggcca gtctccaaag ctctgatct acaaagtttc caaccgattt 180
20 tctgggggtcc cagacagggt cagtggcagt ggatcagga cagatttcac actcaagatc 240
agtagagtgg aggctgagga tctgggagtt tattattgct ttcaagggtc acatgttcca 300
ttcacgttcg gctcggggac aaagttggaa ataaaa 336

25 <210> 325
<211> 360
<212> DNA
<213> Mus musculus

30 <220>
<221> misc_feature
<223> Murine SC17.89 VH

<400> 325
35 cagggtccagt tgcaacagtc tggagctgaa ctggtaaggc ctgggacttc agtgaaggtg 60
tcctgcaaga cttctggata cgccttcact aattacttga tagagtgggt aaagcagagg 120
cctggacagg gccttgagtg gattgggggtg attaatacctg gaagtgggtg tactaactac 180
40 aatgagaagt tcaagggtcaa ggcaacactg actgcagaca aatcctccag cactgcctac 240
atgcagctca ccagcctgac atctgatgac tctgcggtct atttctgtac aagaagggat 300
45 ggttacttct ttccctggtt tgcttactgg ggccaagga ctctgggtcac tgtctctgca 360

50 <210> 326
<211> 339
<212> DNA
<213> Mus musculus

55 <220>
<221> misc_feature
<223> Murine SC17.90 VL

<400> 326

EP 3 338 793 A1

gacattgtga tgtcacagtc tccatcctcc ctagctgtgt cagttggaga gaaggttact 60
atgagctgca agtccagtca gagcctttta tatagtagca atcaaaagaa ctacttggcc 120
5 tggtagcagc agaaaccagg gcagtctcct aaactgctga tttactgggc atccactagg 180
aaatctgggg tccctgatcg cttcacaggc agtggatctg ggacagattt cactctcacc 240
atcagcagtg tgaaggctga agacctggca gtttattact gtcacataa ttatagctat 300
10 ccgctcacgt tcgctgctgg gaccaagctg gagctgaaa 339

<210> 327
<211> 345
15 <212> DNA
<213> Mus musculus

<220>
<221> misc_feature
20 <223> Murine SC17.90 VH

<400> 327
caggtgcaac tgcagcagcc tgggtctgtg ctggtgaggg ctggagcttc agtgaagctg 60
25 tcctgcaagg cttctggcta cacattcacc agctactgga tgcactgggt gaagcagagg 120
cctggacaag gccttgagtg gattggagag attcatccta ataatggtag tactaactac 180
aatgagaagt tcaagggcaa ggccacactg actgtagaca catcctccag cacagcctac 240
30 gtggatctca gcagcctgac atctgaggac tctgcggtct attactgtgc aagatggact 300
ttgtttactt actggggcca agggactctg gtcactgtct ctgca 345

<210> 328
35 <211> 336
<212> DNA
<213> Mus musculus

<220>
40 <221> misc_feature
<223> Murine SC17.91 VL

<400> 328
gatgttgtga tgacccaaac tccactctcc ctgcctgtca gtcttggaga tcaagcctcc 60
45 atctcttgca gatctagtca gagccttgta cacagtaatg gaaacaccta tttactttgg 120
tacctgcaga agccaggcca gtctccaaag ctctgatct acaaagtttc caaccgattt 180
50 tctggggtcc cagacaggtt cagtggcagt ggatcagggc cagatttcgc actcaagatc 240
agcagagtgg aggctgagga tctgggagtt tatttctgct ctcaaagtac acatgttcog 300
tggaagttcg gtggaggcac caagctggaa atcaaa 336

55 <210> 329
<211> 351

EP 3 338 793 A1

<212> DNA
<213> Mus musculus

<220>
<221> misc_feature
<223> Murine SC17.91 VH

<400> 329
gaggtgcagc tggatggagtc tgggggaggc ttagtgaagc ctggaggggc ccggaactc 60
tcctgtgcag cctctggatt cactttcagt gactatggaa tgcactgggt ccgtcaggct 120
ccagagaagg ggctggagtg ggttgcatat attagtcgtg gcagtagtac catccactat 180
gcagacacag tgaagggccg attcaccatc tccagagaca atgccaagaa caccctgttc 240
ctgcaaatga ccagtctaag gtctgaggac acagccatgt attactgtgc aaggcctttc 300
aactggtact tcgatgtctg gggcgaggg acaacgggtca ccgtctcttc a 351

<210> 330
<211> 339
<212> DNA
<213> Mus musculus

<220>
<221> misc_feature
<223> Murine SC17.93 VL

<400> 330
gacattgtga tgtcacagtc tccatcctcc ctagctgtgt cagttggaga gaaggttact 60
atgacctgca agtccagtca gagcctttta tatagtagca atcaaaagaa ctacttggtc 120
tggtaccagc agaaaccagg gcagtctcct aaactactaa ttactgggc atccactagg 180
gaatctgggg tccctgatcg cttcataggc agtggctctg ggacagattt cactctcacc 240
atcagcagtg tgaaggctga agacctggca atttattact gtcagcaata ttatcgctat 300
ccgctcacgt tcggtgctgg gaccaaactg gagctgaaa 339

<210> 331
<211> 363
<212> DNA
<213> Mus musculus

<220>
<221> misc_feature
<223> Murine SC17.93 VH

<400> 331
caggtccaac tgcagcagcc tggggctgag cttgtgaagc ctggggcttc agtgatgctg 60
tcctgcaagg cttctggcta caccttcacc agctactggg tacactgggt gaagcagagg 120
cctggacaag gccttgagtg gattggagtg attaataccta gaaacggctg taacaattac 180

EP 3 338 793 A1

aatgagaagt tcaagaccaa ggccacactg actgtagaca aatcatccag cacagcctac 240

atgcaactca gcagcccgac atctgaggac tctgcggtct attactgtgc acgagaggat 300

5 tacgacgggg gggactatgc tatggactac tggggtaag gaacctcagt caccgtctcc 360

tca 363

<210> 332

10 <211> 321

<212> DNA

<213> Mus musculus

<220>

15 <221> misc_feature

<223> Murine SC17.95 VL

<400> 332

20 gatatccaga tgacacagac tacatcctcc ctgtcggcct ctctgggaga cagggtcacc 60

atcagttgca gtgcaagtca gggcattaac aattatttaa actggtatca gcagaaacca 120

gatggaactg ttacactcct gatctattac acatcaagtt tacactcagg agtcccatcc 180

25 aggttcagtg gcagtgggtc tgggacagat tattctctca ccatcagcaa cctggaacct 240

gaagatattg ccacttacta ttgtcagcag tatagtaagc ttccgtggac gttcgggtgga 300

ggcaccaagc tggaaatcaa a 321

30 <210> 333

<211> 363

<212> DNA

<213> Mus musculus

35 <220>

<221> misc_feature

<223> Murine SC17.95 VH

<400> 333

40 gaggtcgagc tgcaacagtc tggacctgag ctggtgaagc cgggggcttc agtgaagata 60

tcctgcaaga cttccggaaa cacatacact gaatacacca tgcagtgggt gaagctgagc 120

catggaaaga gccttgagtg gattggaggt attaataccta acaatggtat tactagttac 180

45 aaccagaagt tcaagggcaa ggccacattg actgtagaca agtcctccag cacagcctac 240

atggagctcc gcagcctgaa atctgaggat tctgcagtct attactgtgc aagagcggga 300

50 cttggttaact acgtttgggc tatggactac tggggtaag gagcctcagt caccgtctcc 360

tca 363

<210> 334

55 <211> 336

<212> DNA

<213> Mus musculus

<220>
 <221> misc_feature
 <223> Murine SC17.97 VL

5 <400> 334
 gatgttgtga tgacccaaac tccactctcc ctgcctgtca gtcttgaga tcaagcctcc 60
 atctcttgca gatctagtca gagccttgta cacaataatg gaaacaccta tttacattgg 120
 10 tacctgcaga agccaggcca gtctccaaac ctctgatct acaaagtttc caaccgattt 180
 tctgggggtcc cagacaggtt cagtggcagt ggatcaggga cagatttcac actcaagatc 240
 agcatagtgg aggctgagga tctgggactt tatttctgct ctcaaagtac acatgttcct 300
 15 cggacgttcg gtggaggcac caagctggaa atcaaa 336

<210> 335
 <211> 357
 20 <212> DNA
 <213> Mus musculus

<220>
 <221> misc_feature
 25 <223> Murine SC17.97 VH
 <400> 335
 caggtccagc ttccgcagtc tggggctgaa ctggcaaaac ctggggcctc agtgaaaatc 60
 30 tcctgcaagg cttctggctt cacctttact tcctactgga tgactgggt aaaacagagg 120
 cctggacagg gtctggaatg gattggatac attaatacta gcactgatta tactgagtac 180
 aatcagaagt tcaaggacaa ggccacattg actgcagaca aatcctccag cacagcctac 240
 35 atgcaactgg gcagcctgac atctgaggac tctgcagtct attactgtgc aagatcttcc 300
 tacggtagta gcccctttga ttattggggc caaggctcca ctctcacagt ctctca 357

40 <210> 336
 <211> 339
 <212> DNA
 <213> Mus musculus

45 <220>
 <221> misc_feature
 <223> Murine SC17.99 VL
 50 <400> 336
 gacattgtga tgtcacagtc tccatcctcc ctagctgtgt ctggttgaga gaaggttact 60
 atgaactgcg agtccagtca gagcctttta tatagtagca atcaaaagaa ctacttggcc 120
 55 tggtagcagc agaaaccagg gcagtctcct aaactgctga tttactgggc atccactagg 180
 gattctgggg tccctgatcg cttcacaggc agtggatctg ggacagattt cactctcacc 240

EP 3 338 793 A1

atcagcagtg tgagggctga agacccggca gtttattact gtcagcaata ttatagctat 300
ccgctcacgt tcggtgctgg gaccaagctg gagctgaga 339

5

<210> 337
<211> 348
<212> DNA
<213> Mus musculus

10

<220>
<221> misc_feature
<223> Murine SC17.99 VH

15

<400> 337
gaagtgaagc ttgaggagtc tggaggaggc ttggtgcaac ctggaggatc catgaaactc 60
tcttgcgctg cctctggatt cacttttagt gacgcctgga tggactgggt ccgccagtct 120
ccagagaagg ggcttgagtg ggttgctgaa ataagaagca aagctaataa tcatgcaaca 180
tactatgctg agtctgtgaa agggagggtc accatctcaa gagatgattc caaaagtagt 240
gcctacctgc aaatgaacag ctttaagagct gaagacactg gcatttatta ttgtgtttca 300
acaggggactt cttactgggg ccaagggact ctggctactg tctctgca 348

25

<210> 338
<211> 318
<212> DNA
<213> Mus musculus

30

<220>
<221> misc_feature
<223> Murine SC17.102 VL

35

<400> 338
caaattgttc tcaccagtc tccagcaatc atgtctgcat ctccagggga gaaggtcacc 60
atgacctgca gtgccagtc aagtgttaagt tacatgcact ggtaccagca gaagtcaggc 120
acctcccca aaagatggat ttatgacaca tccaaactgg cttctggagt ccctcctcgc 180
ttcagtggcc gtgggtcttg gacctcttac tctctcacia tcagcagcat ggaggctgaa 240
gatgctgcc cttattactg ccagcattgg agtagtaacc caccacggt cggtgctggg 300
accaagctgg agatgaaa 318

45

<210> 339
<211> 348
<212> DNA
<213> Mus musculus

50

<220>
<221> misc_feature
<223> Murine SC17.102 VH

55

EP 3 338 793 A1

<400> 339
 gaggtccagc tgcaacagtc tggacctgag ctaatgaagc ctggggcttc agtgaagatg 60
 tcctgcaagg cttctggaga cacattcact gactacaaca tacactgggt gaagcagaac 120
 5 caaggaaaga gcctagagtg gataggagaa gttaatccta acattggtgg tattggctat 180
 aaccagaagt tcaaaggcaa ggccacattg actgtagaca agtcctccag cacagcctac 240
 10 atggagctcc gcagcctgac atctgaggac tctgcagtct attactgtgc aatggggagg 300
 tggctacttcg atgtctgggg cgcagggacc acggtcaccg tctcctca 348

 <210> 340
 <211> 336
 <212> DNA
 <213> Mus musculus

 <220>
 <221> misc_feature
 <223> Murine SC17.114 VL

 <400> 340
 gatgttgtga tgaccagtc tccactctcc ctgcctgtca gtcttggaga tcaagcctcc 60
 25 atctcttgca gatctagtca gagccttgta cacagtaatg gaaacaccta ttacattgg 120
 tacctgcaga agccaggcca gtctccaaag ctctgatct acaaagtttc cagccgattt 180
 tctggggtcc cagacaggtt cagtggcagt ggatcagga cagatttcac actcaagatc 240
 30 agcagagtgg aggctgagga tctgggagtt tatttctgct ctcaaagtac acatgttcca 300
 ttcacgttcg gctcggggac aaagttggaa ataaaa 336

 <210> 341
 <211> 345
 <212> DNA
 <213> Mus musculus

 <220>
 <221> misc_feature
 <223> Murine SC17.114 VH

 <400> 341
 45 gaggtccagc tgcagcagtc tggacctgag atggtgaagc ctggggcttc agtgaagata 60
 tcctgcaagg cttctggata cacattcact gactactaca tgactgggt gaaacagagc 120
 catggaaaga gccttgagtg gattggacgt gttaatacta acaatggtgg aactagctac 180
 50 gaccagaagt tcgagggcaa ggccacattg actgttgaca aatcttcag cacagcctac 240
 atggagctca acagcctgac atctgaggac tctgcggtct attactgtgt aatccctgcc 300
 tggtttgctt actggggcca agggactctg gtcactgtct ctgca 345
 55
 <210> 342

EP 3 338 793 A1

<211> 336
<212> DNA
<213> Mus musculus

5

<220>
<221> misc_feature
<223> Murine SC17.115 VL

10

<400> 342
gatattgtga tgacccagtc tccactctcc ctgcctgtca gtcttggaga tcaagcctcc 60
atctcttgca gatctagtca gagccttgta cacagtaatg gaaacaccta ttacattgg 120
tacctgcaga agccaggcca gtctccaaag ctccctgatct acagagtttc caaccgattt 180
tctgggggtcc cagacagggt cagtggcagt ggatcagga cagatttcac actcacgac 240
agcagagtgg aggctgagga tctgggagtt tatttctgct ctcaaagtac acatcttct 300
cggacgttcg gtggaggcac caagctggag atcaaa 336

20

<210> 343
<211> 342
<212> DNA
<213> Mus musculus

25

<220>
<221> misc_feature
<223> Murine SC17.115 VH

30

<400> 343
caggtgcaac tgcagcagtc tgggtctgtg ctggtgagga ctggagcttc agtgaagctg 60
tcctgcaagg cttctggcta cacattcacc agctactgga tgcactgggt gaagcagagg 120
cctggacaag gccttgagtg gattggagag attcatccta atagtgggaa tactaattac 180
aatgagaagt tcaagggcaa ggccacactg actgtagaca catcctccag cacagcctac 240
gtggatctca gcagcctgac atctgaggac tctgcggtct attattgtgc aggtggtaac 300
tacgactact ggggccaagg caccactctc acagtctcct ca 342

40

<210> 344
<211> 333
<212> DNA
<213> Mus musculus

45

<220>
<221> misc_feature
<223> Murine SC17.120 VL

50

<400> 344
gacattgtgc tgacccaatc tccagcttct ttggctgtat ctctagggca gagggccacc 60
atatcctgca gagccagtga aagtgttgat agttatggca atagttttat gactggtac 120
cagcagaaac caggacagcc acccaaagtc ctcatctatc gtgcatccaa cctagaatct 180

55

EP 3 338 793 A1

gggatccctg ccagggttcag tggcagtggtg tctaggacag acttcaccct caccattaat 240
 cctgtggagg atgaagatgt tgcaacctat tactgtcagc aaagtaatga ggatccgtac 300
 5 acgttcgggg gggggaccaa gctggaaata aaa 333

 <210> 345
 <211> 351
 <212> DNA
 10 <213> Mus musculus

 <220>
 <221> misc_feature
 15 <223> Murine SC17.120 VH

 <400> 345
 gaggttcagc tcgagcagtc tgggactgtg ctggcaaggc ctggggcttc agtgaagatg 60
 tcctgcaagg cttctggcta cacctttacc agctactgga tgcactgggt gaaacagagg 120
 20 cctggacagg gtctggaatg gattggcgtt ttttatcctg gaaacagtgg tacttattac 180
 aacaaaaaat tcaaggacaa ggccaaactg actgcagtca catctgccag cactgcctac 240
 25 atggagctca gcagcctgac aaatgaggac tctgcggtct attactgttc aagatcaggg 300
 tcaggaaggt ttgcttactg gggccaaggg actctgggtca ctgtctctgc a 351

 <210> 346
 <211> 318
 <212> DNA
 30 <213> Mus musculus

 <220>
 <221> misc_feature
 35 <223> Murine SC17.121 VL

 <400> 346
 caaattgttc tcacccagtc tccagcaatc atgtctgcat ctccagggga gaaggtcacc 60
 40 atgacctgca gtgccagctc aagtgtgagt tacatgcact ggtaccagca gaagtcaggc 120
 acctccccca aaagatggat ttatgacaca tccaaactgg cttctggagt ccctgctcgc 180
 45 ttcagtggca gtgggtcttg gacctcttac tctctcacia tcagcagcat ggagactgaa 240
 gatgctgcca cttattactg ccagcagtggt agtaataccc caccacggtt cggctcgggtg 300
 acaaagttgg aaataaaa 318

 50 <210> 347
 <211> 348
 <212> DNA
 <213> Mus musculus

 55 <220>

EP 3 338 793 A1

<221> misc_feature
<223> Murine SC17.121 VH

<400> 347
gaggtccagc tgcaacagtc tggacctgag ctaatgaagc ctggggcttc agtgaagatg 60
tcctgcaagg cttctggata cacattcact gaccacaaca tacactgggt gaaacagcac 120
caaggaaaga gcctagagtg gataggagaa attaataccta aactgggtgg tactggctac 180
aaccagaagt tccaaggcaa ggccacaatg actgtagaca agtcctccag cacagcctac 240
atggaactcc gcagcctgac atctgaggac tctgcagtct attactgtgt tagaggactg 300
tacttctttg actactgggg ccaaggcacc actctcacag tctcctca 348

<210> 348
<211> 336
<212> DNA
<213> Mus musculus

<220>
<221> misc_feature
<223> Murine SC17.122 VL

<400> 348
gatattgtga taaccagga tgatctctcc aatcctgtca cttctggaga atcagtttcc 60
atctcctgca ggtctagtaa gagtctccta tataaggatg ggaagacata cttgaattgg 120
tttctgcaga gaccaggaca atctcctcag ctctgatct atttgatgtc caccctgca 180
tcaggagtct cagaccggtt tagtggcagt gggtcaggaa cagatttcac cctggaaatc 240
agtagagtga aggctgagga tgtgggtgtg tattactgtc aacaacttgt agagtatcct 300
cggacgttcg gtggaggcac caagctggaa atcaaa 336

<210> 349
<211> 339
<212> DNA
<213> Mus musculus

<220>
<221> misc_feature
<223> Murine SC17.122 VH

<400> 349
gaggtgcacc tgggtggagtc tgggggagac ttagtgaagc ctggagggtc cctgaaactc 60
tcctgtgcag cctctggatt cactttcagt agctatggca tgtcttgggt tcgccagact 120
ccagacaaga ggctggagtg ggtcgcaacc attagtagtg gtggtactta cacctactat 180
ccagacagtg tgaaggggag attcaccatc tccagagaca atgccaagaa caccctgtat 240
ctgcaaatga gcagtctgaa gtctgaggac acagccatgt attactgttc aagacatggg 300
tggggctggg gccaaaggac tctggtcact gtctctgca 339

EP 3 338 793 A1

<210> 350
 <211> 318
 <212> DNA
 <213> Mus musculus

5

<220>
 <221> misc_feature
 <223> Murine SC17.140 VL

10

<400> 350
 caaattgttc tcacccagtc tccagcaatc atgtctgcat ctccagggga gaaggtcacc 60
 atgacctgca gtgccagctc aagtgttagt tacatgcact ggtaccagca gaagtcaggc 120
 acctcccca aaagatggat ttatgacaca tccaaactgg cttctggagt ccctgctcgc 180
 ttcagtggca gtgggtctgg gacctcttac tctctcacia tcagcagcat ggaggctgaa 240
 gatgctgcca cttattactg ccagcagtgg agtagtacct caccacggtt cggctcgggg 300
 acaaagttgg aaataaaa 318

20

<210> 351
 <211> 360
 <212> DNA
 <213> Mus musculus

25

<220>
 <221> misc_feature
 <223> Murine SC17.140 VH

30

<400> 351
 gaggtccagc tgcaacagtc tggacctgag ctaatgaagc ctggggcttc agtgaagatg 60
 tcttgcaagg cttctggata cacattcact gactacaaca tgcactgggt gaagcagaac 120
 caaggaaaga gcctagagtg gataggagaa attaatccca aactgggtgg tactggctac 180
 aaccagaagt tcaaaggcaa ggccacattg actgtagaca agttttccag cacagccttc 240
 attgagctcc gcagcctgac atctgaggac tctgcaatct attactgtac aagagggggg 300
 tacgaccact attggtactt cgatgtctgg ggcgcaggga ccacggtcac cgtctcctca 360

45

<210> 352
 <211> 333
 <212> DNA
 <213> Mus musculus

50

<220>
 <221> misc_feature
 <223> Murine SC17.151 VL

55

<400> 352
 gacattgtgc tgacccaatt tccagcttct ttggctgtgt ctctagggca gagggccacc 60

EP 3 338 793 A1

	ataccctgca gagccagtga aagtgttgat agttatggca atagttttat gcactgggtc	120
	cagcagaaac caggacagcc acccaaactc ctcatctatc gtgcatccaa cctagaatct	180
5	gagatccctg ccaggttcag tggcagtggt tctgggacag acttcaccct caccattaat	240
	cctgtggagg ctgatgatgt tgcaacctat tactgtcagc aaagtcatga ggatccgtac	300
	acgttcggag gggggaccaa gatggaaata aaa	333
10	<210> 353 <211> 351 <212> DNA <213> Mus musculus	
15	<220> <221> misc_feature <223> Murine SC17.151 VH	
20	<400> 353	
	gaggttcagc tgcagcagtc tgggactgtg ctggcaaggc ctgggggttc agtgaagatg	60
	tcctgcaagg cttctggcta tacctttacc agctactgga tgcactgggt aaaacagagg	120
25	cctggacagg gtctggaatg gattggcgct atttatacctg gaaagaatga tactacctac	180
	aaccagaagt tcaagggcaa ggccaaactg actgcagtca catctgccag cactttatac	240
	atggagctca gcagcctgac aaatgaggac tctgcggtct attactgtac aagatctgga	300
30	aagggttact ttgcttactg gggccaaggg actctggtca ctgtctctgc a	351
35	<210> 354 <211> 336 <212> DNA <213> Mus musculus	
40	<220> <221> misc_feature <223> Murine SC17.156 VL	
	<400> 354	
	gatgttgtga tgaccagtc tccactctcc ctgcctgtca gtcttgaga tcaagcctcc	60
45	atctcttgca gatctagtca gagcattgta catagtaatg gaaacaccta tttagaatgg	120
	tacctgcaga aaccaggcca gtctccaaag ctctgatct acaaagtttc caaccgattt	180
	tctggggtcc cagacagggt cagtggcagt ggatcaggga cagatttcac actcaagatc	240
50	agcagagtgg aggctgagga tctgggagtt tattactgct ttcaaggttc acatgttctt	300
	ccgacgttcg gtggaggcac caaactggaa atcaaa	336
55	<210> 355 <211> 354 <212> DNA <213> Mus musculus	

<220>
 <221> misc_feature
 <223> Murine SC17.156 VH

5 <400> 355
 caggttactc tgaaagagtc tggccctggg atattgcagc cctcccagac cctcagtctg 60
 acttgttctt tctctgggtt ttcactgagc acttctggta tgggtgtgag ctggattcgt 120
 10 aagacttcag gaaagggctc ggaatggctg gcacacattt tctgggatga tgacaagtgg 180
 tataatccat ccctgaagag ccggctcaca atctccaagg ctacctccag caaccaggta 240
 ttctcatac tcaccagtgt ggatactgcc gatactgcca catactactg tgctaccttc 300
 15 tatggtctct actttgccta ctggggccaa ggcaccactc tcacagtctc ctca 354

20 <210> 356
 <211> 339
 <212> DNA
 <213> Mus musculus

25 <220>
 <221> misc_feature
 <223> Murine SC17.161 VL
 <400> 356
 gacattgtga tgtcacagtc tccatcctcc ctagctgtgt ctggtggaga gaaggttact 60
 30 atgaactgcy agtccagtca gagcctttta tataatagca atcaaaagaa ctacttggcc 120
 tggtagcagc agaaaccagg gcagtctcct aaactgctga tttactgggc atccactagg 180
 gattctgggg tccctgatcg cttcacaggc agtggatctg ggacagattt cactctcacc 240
 35 atcagcagtg tgagggctga tgacccggca gtttattact gtcagcaata ttttaactat 300
 ccgctcacgt tcggtgctgg gaccaagctg gagctgaaa 339

40 <210> 357
 <211> 348
 <212> DNA
 <213> Mus musculus

45 <220>
 <221> misc_feature
 <223> Murine SC17.161 VH
 50 <400> 357
 gaagtgaagc ttgaggagtc tggaggaggc ttggtgcaac ctggaggatc catgaaactc 60
 tcttgcgctg cctctggatt cacttttagt gacgcctgga tggactgggt ccgccagtct 120
 ccagagaagg ggcttgagtg ggttgctgaa ataagaagca aacctaataa tcatgcaaca 180
 55 tactatgctg agtctgtgaa agggaggttc accatctcaa gagatgattc caaaagtagt 240

EP 3 338 793 A1

gcctacctgc aaatgaacag cttaagagct gaagacactg gcatttatta ctgtgtttca 300
acagggactt cttactgggg ccaagggact ctggtcactg tctctgca 348

5

<210> 358
<211> 318
<212> DNA
<213> Mus musculus

10

<220>
<221> misc_feature
<223> Murine SC17.166 VL

15

<400> 358
caaattgttc tcacccagtc tccagcaatc atgtctgcat ctccagggga gaaggtcacc 60
atgacctgca gtgccagctc aagtataagt tacatgcact ggtaccagca gaagtcaggc 120
acctccccc aaagatggat ttatgacaca tccaaactgg cttctggagt ccctgctcgc 180
ttcagtggca gtgggtcttg gacctcttac tctctcaca tcagcaacat ggaggctgaa 240
gatgctgcc cttattactg ccagcagtgg agtagtacc caccacggt cggagggggg 300
accaagctgg aaataaaa 318

25

<210> 359
<211> 351
<212> DNA
<213> Mus musculus

30

<220>
<221> misc_feature
<223> Murine SC17.166 VH

35

<400> 359
gaggtccagt tgcaacagtc tggacctgag ctaatgaagc ctggggcttc agtgaagatg 60
tcctgcaagg cttctggata tatatttact gactacaaca tgcaactggg gaagcagaac 120
caaggaaaga gcctagagtg gataggagaa gttaatccta aactggtgg tattggctac 180
aatcagaaat tcaaaggcaa ggccacattg actgtagaca agtcctccag cacagcctac 240
atggacctcc gcagcctgac atctgaggac tctgcagtct attactgtgc aagagatggc 300
aattattgct ttgactactg gggccaaggc accactctca cagtctctc a 351

45

<210> 360
<211> 321
<212> DNA
<213> Mus musculus

50

<220>
<221> misc_feature
<223> Murine SC17.187 VL

55

EP 3 338 793 A1

<400> 360
 gacatcaaga tgaccagtc tccatcttcc atgtatgcat ctctaggaga gagagtcact 60
 ctcaacttgca aggcgagtc gacattaat agctatttaa gctgggtcca gcagaaacca 120
 5 gggaaatctc ctgagaccct gatctatcgt gcaaacagat tgatagatgg ggtcccatca 180
 aggttcagtg gcagtggatc tgggcaagat tattctctca ccatcagcag cctggagtat 240
 10 gaagatatgg ggatttatta ttgtctacag tatgatgagt ttcctccgac gttcgggtgga 300
 ggcaccaagc tggaaatcaa a 321

 <210> 361
 <211> 354
 <212> DNA
 <213> Mus musculus

 <220>
 <221> misc_feature
 <223> Murine SC17.187 VH

 <400> 361
 gaggtccacc tacaacagtc tggacctgaa ctggtgaacc ctgggtcttc agtgaagata 60
 25 tcctgcaagg ctgctggata cacattcact gactacaaca tggactgggt gaagcagagc 120
 catggaaaga gacttgagtg gattggaaat atttatccta acaatggtgg tgctggatac 180
 aaccagaact tcaaggacaa ggccacattg actgtagaca agtcctccag cacagcctac 240
 30 atggagctcc gcagcctgac atctgaggac tctgcagtct attactgtgc aagatccatt 300
 actgcggctt ggtttgctta ctggggccaa gggactctgg tcaactgtctc tgca 354

 35 <210> 362
 <211> 318
 <212> DNA
 <213> Mus musculus

 40 <220>
 <221> misc_feature
 <223> Murine SC17.191 VL

 <400> 362
 45 caaattgttc tcaccagtc tccagcaatc atgtctgcat ctccagggga gaagtcacc 60
 atgacctgca gtgccagctc aagtgtgaat tacatgcact ggtaccagca gaagtcaggc 120
 acctccccc aaagatggat ttatgacaca tccaaactgg cttctggagt ccctgctcgc 180
 50 ttactggca gtgggtctgg gacctcttac tctctcacia tcagcagcat ggaggctgaa 240
 gatgctgcca cttattactg ccagcagtg agtagtagcc caccacggt cggtgctggg 300
 accaagctgg aactgaaa 318
 55
 <210> 363

EP 3 338 793 A1

<211> 360
<212> DNA
<213> Mus musculus

5

<220>
<221> misc_feature
<223> Murine SC17.191 VH

10

<400> 363
gaggtccagc tgcaacagtc tggacctgag ctaatgaagc ctggggcttc agtgaagatg 60
tcctgcaagg cttctggata cacattcact gactacaaca tgcactgggt gaagcagaac 120
caaggaaaga gcctagagtg gataggagaa attaataccta aactgggtgg tactggctac 180
aaccagaagt tcaaagacaa ggccacattg actgtagaca agtcctccag cacagcctac 240
atggagctcc gcagcctgac atctgaggac tctgcagtct attactgtgc aagaattccc 300
tccctgagac gatactactt tgactactgg ggccaaggca ccactctcac agtctcctca 360

20

<210> 364
<211> 333
<212> DNA
<213> Mus musculus

25

<220>
<221> misc_feature
<223> Murine SC17.193 VL

30

<400> 364
gacctgtgtc tgacacagtc tcctgcttcc ttagctgtat ctctggggca gagggccacc 60
atctcatgca gggccagcga aagtgtcagt acatctggct atagttatat gcactggtac 120
caacagaaac caggacagcc acccaaactc ctcatctatc ttgcatccaa cctcgaatct 180
gggggtccctg ccaggttcag tggcagtggt tctgggacag acttcaccct caacatccat 240
cctgtggagg aggaggatgc tacaacctat tactgtcagc acagtaggga gcttccgtac 300
acgttcggag gggggaccaa gctggaaata aaa 333

40

<210> 365
<211> 366
<212> DNA
<213> Mus musculus

45

<220>
<221> misc_feature
<223> Murine SC17.193 VH

50

<400> 365
caggttactc tgaaagagtc tggccctggg atattgcagc cctcccagac cctcagtctg 60
acttgttctt tctctgggtt ttactgatc acttatggta taggagtagg ctggattcgt 120
cagccttcag ggaagggtct ggagtggctg gcacacattt ggtggaatga taataagtac 180

55

EP 3 338 793 A1

tataacacag ccctgaagag cgggctcaca atctccaagg atacctccaa caaccaggta 240

ttcctcaaga togccaatgt ggacactgca gatactgccata catactactg tgctogaatg 300

5 gtctactatg attacgacgg ggggtttgct tactggggcc aagggactct ggtcactgtc 360

tctgca 366

10 <210> 366
<211> 333
<212> DNA
<213> Mus musculus

15 <220>
<221> misc_feature
<223> Murine SC17.199 VL

<400> 366

20 gacattgtgc tgacccaatc tccagcttct ttggctgtgt ctctagggca gagggccacc 60

atatacctgca gagccagtga aagtgttgat agttatggca atagttttat gcactggtac 120

cagcagaaac caggacagcc acccaaacc ctcatttatc gtgcatccaa cctagaatct 180

25 gggatccctg ccagattcag tggcagtggg tctaggacag acttcaccct caccattaat 240

cctgtggagg ctgatgatgt tgcaacctat tactgtcagc aaagtaatga ggatccgtac 300

acgttcggag gggggaccaa gctggaaata aaa 333

30 <210> 367
<211> 348
<212> DNA
<213> Mus musculus

35 <220>
<221> misc_feature
<223> Murine SC17.199 VH

40 <400> 367

gaggtgcagc tgcagcagtc tgggactgtg ctggcaaggc ctggggcttc agtaaggatg 60

tcctgcaagg cttctggcta cacctttacc agctactgga tgcactgggt aaaacaaagg 120

45 cctggacagg gtctggaatg gattggcgct atttatcctg gaaatagtga tactagctac 180

aaccataagt tcaagggcaa ggccaaactg actgcagtca catctgccag cactgcctac 240

atggagctca gcagcctgac aaatgaggac tctgcggctc attactgtac aagatctggg 300

50 acgggctggg ttgcttactg gggccaaggg actctggtca ctgtctct 348

<210> 368
<211> 333
<212> DNA
<213> Mus musculus

55

<220>
 <221> misc_feature
 <223> Murine SC17.200 VL

5 <400> 368
 gacattgtgc tgaccaatc tccagcttct ttggctgtgt ctctaggaca gagagccact 60
 atcttctgca gagccagcca gagtgtcgat tataatggaa ttagttatat gcactggttc 120
 10 caacaaaaac caggacagcc acccaactc ctcatctatg ctgcatccaa cgttcaatct 180
 gggatccctg ccaggttcag tggcagtggg tctgggacag acttcaccct caacatccat 240
 cctgtggagg aggaagatgc tgcaacctt tactgtcagc aaagtattga ggatcctccg 300
 15 acgttcggtg gaggcaccaa gctggaaatc aaa 333

<210> 369
 <211> 345
 <212> DNA
 20 <213> Mus musculus

<220>
 <221> misc_feature
 25 <223> Murine SC17.200 VH

<400> 369
 caggtccagc tgcagcagtc tggacctgag ctggtgaaac ctggggcctc agtgaagatt 60
 30 tcctgcaaag cttctggcta cgcatcagc agttcttggg ttaactgggt gaagcagagg 120
 cctggacagg gtcttgagtg gattggacgg atttatcctg gagaaggatga tactaactac 180
 agtgggaatt tcgagggcaa ggccacactg actgcagaca aatcctccac cacagcctac 240
 35 atgcagctca gcagtctgac ctctgtggac tctgcggtct atttctgtac aagaggacta 300
 gtcatggact actggggcca aggcaccgct ctcacagtct cctca 345

<210> 370
 40 <211> 321
 <212> DNA
 <213> Artificial Sequence

<220>
 45 <223> Humanized hSC17.16 VL

<400> 370
 gacatccaga tgaccagtc tccatcctcc ctgtctgcat ctgtaggaga cagagtcacc 60
 50 atcacttgcc gggcaagtgc gaacattaac agcaatttag tttggtatca gcagaaacca 120
 gggaaagccc ctaagctcct gatctatgct gcaaccaatt tggcagatgg ggtcccatca 180
 aggttcagtg gcagtggatc tgggacagat ttcactctca ccatcagcag tctgcaacct 240
 55 gaagattttg caacttacta ctgtcaacat ttttggggta ctccctcgac gttcgggtgga 300
 ggcaccaagc tggaatcaa a 321

<210> 371
 <211> 348
 <212> DNA
 <213> Artificial Sequence

5

<220>
 <223> Humanized hSC17.16 VH

10

<400> 371
 caggtgcagc tgggtgcagtc tggggctgag gtgaagaagc ctggggcctc agtgaaggtc 60
 tcctgcaagg cttctggata caccttcacc gactacaata tgtactgggt gcgacaggcc 120
 cctggacaag ggcttgagtg gatgggagag atcaacccta acaatgggtg cacagcctat 180
 aatcagaagt ttaggggcaa ggtcaccatg accagggaaca cgtccatcag cacagcctac 240
 atggagctga gcaggctgag atctgacgac acggccgtgt attactgtgc gagatatgat 300
 aaggggtttg actactgggg ccaaggcacc actgtcacag tctcctca 348

20

<210> 372
 <211> 318
 <212> DNA
 <213> Artificial Sequence

25

<220>
 <223> Humanized hSC17.17 VL

30

<400> 372
 gaaattgtgt tgacacagtc tccagccacc ctgtctttgt ctccagggga aagagccacc 60
 ctctcctgca gtgccagtag cagtgttagc tacatgcatt ggtaccaaca gaaacctggc 120
 caggctccca ggctcctcat ctatgatata tccaaattgc ccagtggcat cccagccagg 180
 ttcagtggca gtgggtctgg gacagacttc actctacca tcagcagcct agagcctgaa 240
 gatcttgcag tttattactg tcagcagtgg agtagtacc caccacgtt cggtcagggg 300
 accaagctgg agattaata 318

40

<210> 373
 <211> 360
 <212> DNA
 <213> Artificial Sequence

45

<220>
 <223> Humanized hSC17.17 VH

50

<400> 373
 caggtgcagc tgggtgcagtc tggggctgag gtgaagaagc ctggggcctc agtgaaggtc 60
 tcctgcaagg cttctggata caccttcacc gactacaata tgactgggt gcgacaggcc 120
 cctggacaag ggcttgagtg gatgggagag atcaacccta acattgggtg cacaggctat 180
 aaccagaagt ttaagggcag ggtcaccatg accagggaaca cgtccatcag cacagcctac 240

55

EP 3 338 793 A1

atggagctga gcaggctgag atctgacgac acggccgtgt attactgtgc gagaacctat 300

agttactata gttacgagtt tgcttactgg ggccaaggga ctctgggtcac tgtctcttca 360

5

<210> 374
<211> 336
<212> DNA
<213> Artificial Sequence

10

<220>
<223> Humanized hSC17.24 VL

<400> 374
gacatcgtga tgaccagtc tccagactcc ctggctgtgt ctctgggcga gagggccacc 60

15

atcaactgca agtccagcca gagtcttctc tacagctcca accagaagag ctacttagct 120

tggtaccagc agaaaccagg acagcctcct aagctgtctc ttactgggc atctaccgg 180

gaatccgggg tccctgaccg attcagtggc agcgggtctg ggacagattt cactctcacc 240

20

atcagcagcc tgcaggctga agatgtggca gtttattact gtaagcaatc ttataatctt 300

cggacgttcg gtggaggcac caaggtggaa atcaaa 336

25

<210> 375
<211> 351
<212> DNA
<213> Artificial Sequence

30

<220>
<223> Humanized hSC17.24 VH

<400> 375
gaagtgcagc tgggtgcagtc tggggctgag gtgaagaagc ctggggccac cgtgaagata 60

35

tcttgcaagg tgtctggata caccttcaca gaccacacta tacactgggt gcgacaggcc 120

cctggaaagg ggcttgagtg gattggatac atctaccctc gtgatggtag cacaaaatac 180

aacgaggagt tcaaaggcag agtcaccatc accgccgaca cgtccacgga cacagcctac 240

40

atggagctga gcagcctgag atctgaggac acggccgtgt attactgtgc gagatcatat 300

agtaactact ttgactactg gggccaaggc accactgtca cagtctcctc a 351

45

<210> 376
<211> 321
<212> DNA
<213> Artificial Sequence

50

<220>
<223> Humanized hSC17.28 VL

<400> 376
gaaattgtgc tgactcagtc tccagacttt cagtctgtga ctccaaagga gaaagtcacc 60

55

atcacctgcc gggccagtc gagcattggt actagcatac actggtacca gcagaaacca 120

gatcagtcctc caaagctcct catcaagtat gcttccgagt ccatctcagg ggtccctcgt 180

EP 3 338 793 A1

aggttcagtgc gcagtggatc tgggacagat ttcaccctca ccatcaatag cctggaagct 240
gaagatgctg caacgtatta ctgtcagcaa agtaatagct ggccactcac gttcgggtcaa 300
5 gggaccaagc tggagataaa a 321

<210> 377
<211> 351
10 <212> DNA
<213> Artificial Sequence

<220>
<223> Humanized hSC17.28 VH

15 <400> 377
caggtgcagc tgggtgcagtc tggggctgag gtgaagaagc ctggggcctc agtgaaggtc 60
tcctgcaagg cttctggata caccttcacc agaagctata tccactgggt gcgacaggcc 120
20 cctggacaag ggcttgagtgc gatgggatac atcagcagtg gcagtgggtg cacaacctat 180
aaccagaagt ttaagggcag ggtcaccagt accagggaca cgtccatcag cacagcctac 240
atggagctga gcaggctgag atctgacgac acggccgtgt attactgtgc gagagggggg 300
25 gtacggtact tcgatgtctg gggccaaggg accacggtca ccgtctcctc a 351

<210> 378
<211> 321
30 <212> DNA
<213> Artificial Sequence

<220>
<223> Humanized hSC17.34 VL

35 <400> 378
gacatccaga tgaccagtc tccatcctca ctgtctgcat ctgtaggaga cagagtcacc 60
atcacttgta aggcagtc gacattaat agttatttat cctggtttca gcagaaacca 120
40 gggaaagccc ctaagtcctt gatctataga gcaaacagat tggtagatgg ggtcccatca 180
aggttcagcg gcagtggatc tgggacagat tacactctca ccatcagcag cctgcagcct 240
gaagattttg caacttatta ctgcctacag tatgatgagt ttcctccgac gttcgggtcag 300
45 ggcaccaagc tggaaatcaa a 321

<210> 379
<211> 354
50 <212> DNA
<213> Artificial Sequence

<220>
<223> Humanized hSC17.34 VL

55 <400> 379
caggtccagc ttgtgcagtc tggggctgag gtgaagaagc ctggggcctc agtgaaggtt 60

EP 3 338 793 A1

tcctgcaagg cttctggata caccttcact gactataata tggattgggt gcgccaggcc 120
 cccggacaaa ggcttgagt gattggatac atctaccctg acaatgggtg cgcaggatat 180
 5 aatcagaagt tcaagggcag agtcaccatt accgtggaca catccgcgag cacagcctac 240
 atggagctga gcagcctgag atctgaagac acggctgtgt attactgttc aagatccatt 300
 actacggctt ggtttgctta ctggggccaa gggactctgg tcaactgtctc ttca 354
 10
 <210> 380
 <211> 321
 <212> DNA
 <213> Artificial Sequence
 15
 <220>
 <223> Humanized hSC17.46 VL
 <400> 380
 gccatccaga tgaccagtc tccatcctcc ctgtctgcat ctgtaggaga cagagtcacc 60
 20 atcacttgca aggcaagtca gagcgtaaat aatgatgtag cctgggtatca gcagaaacca 120
 gggaaagccc ctaagctcct gatctattat gcatccaatc gatatactgg ggtcccatca 180
 25 aggttcagcg gcagtggatc tggcacagat ttcactctca ccatcagcag cctgcagcct 240
 gaagattttg caactatatt ctgtcagcag gattatagct ctccctcggac gttcggtcag 300
 gggaccaagc tggaaataaa g 321
 30
 <210> 381
 <211> 363
 <212> DNA
 <213> Artificial Sequence
 35
 <220>
 <223> Humanized hSC17.46 VH
 <400> 381
 caggtgcagc tgggtgcagtc tggggctgag gtgaagaagc ctggggcctc agtgaaggtc 60
 40 tcctgcaagg cttctggata caccttcacc agctactgga tcaactgggt gcgacaggcc 120
 cctggacaag ggcttgagt gattggaaac atcttccctg acaactactac cacaaactat 180
 aacgagaagt ttaagggcag ggtcaccctg accagggaca cgtccatcag cacagcctac 240
 45 atggagctga gcaggctgag atctgacgac acggccgtgt attactgtgc gagagagtac 300
 tacgatggta cctacgatgc tatggattac tggggtaag gaaccctagt caccgtctcc 360
 tca 363
 50
 <210> 382
 <211> 333
 <212> DNA
 55 <213> Artificial Sequence
 <220>

EP 3 338 793 A1

<223> Humanized hSC17.151 VL

<400> 382

gagatcgtgc tgaccagag ccctgctaca ctgtccctgt cccctggaga gagggccaca 60
 ctctcctgca gggcttccga gtccgtggat tcctacggca actccttcat gactgggtac 120
 cagcagaaac ccggccaggc ccctaggctg ctgatctaca gggcctccaa cctggagtcc 180
 ggcatccctg ctaggttctc cggatccggc tccggcaccg actttaccct gaccatctcc 240
 tccctggagc ccgaggactt cggcgtgtac tactgccagc agtcccacga ggaccctac 300
 accttcggcc agggcaccaa gctggagatc aag 333

<210> 383

<211> 351

<212> DNA

<213> Artificial Sequence

<220>

<223> Humanized hSC17.151 VH

<400> 383

caggccagc tggcgcagag cggcgcctgag gtgaagaagc ctggcgccag cgtgaaggctg 60
 tcctgcaaag ccagcggcta caccttcacc tcctactgga tgcattgggt gaggcaggct 120
 cctggccaag gactggagtg gatgggcgcc atctaccccg gcaagtccga caccacctac 180
 aaccagaagt tcaagggcag ggtgacctg acacgggaca cctccacctc caccgtgtac 240
 atggagctgt cctccctgag gtccgaggac accgccgtgt actactgcgc caggccgggc 300
 aagggtatt tcgcctactg gggccagggc aactgggtga ccgtgtcctc c 351

<210> 384

<211> 339

<212> DNA

<213> Artificial Sequence

<220>

<223> Humanized hSC17.155 VL

<400> 384

gacatcgtga tgaccagtc tcagactcc ctggctgtgt ctctgggcga gagggccacc 60
 atcaactgca agtccagcca gagtttatta tacagctcca accaaaagaa ctacttagct 120
 tggtagcagc agaaaccagg acagcctcct aagctgtctc ttactgggc atctacccgg 180
 aaatccgggg tcctgaccg attcagtggc agcgggtctg ggacagattt cactctcacc 240
 atcagcagcc tgcaggctga agatgtggca gtttattact gtcacataa ttatagctat 300
 ccgtcacgt tcggtcaagg caccaagctg gaaatcaaa 339

<210> 385

<211> 339

<212> DNA

<213> Artificial Sequence

<220>

<223> Humanized hSC17.155 VH

5

<400> 385

caggtgcagc tgggtgcagtc tggggctgag gtgaagaagc ctggggcctc agtgaagggt 60

tcctgcaagg catctggata caccttcaac agctactgga tgcactgggt gcgacaggcc 120

10

cctggacaag ggcttgagtg gatgggagaa atccacccta ataatggtag cacaaactac 180

aacgagaagt tcaagggcag agtcaccatg accagggaca cgtccacgag cacagtctac 240

atggagctga gcagcctgag atctgaggac acggccgtgt attactgtgc gagatggact 300

15

ttgtttactt actggggcca agggactctg gtcactgtc 339

<210> 386

<211> 336

<212> DNA

<213> Artificial Sequence

<220>

<223> Humanized hSC17.156 VL

25

<400> 386

gacatcgtga tgaccagac ccctctgtcc ctgcctgtga cccctggaga acccgccagc 60

atctcctgca ggtcctccca gtccatcgtg cactccaacg gcaacaccta cctggagtgg 120

30

tacctgcaga agcccgagca gtccccccag ctgctgatct acaagggtgtc caataggttt 180

tccggagtgc ccgacagggt ctccggatcc ggatccggca ccgacttcac cctgaagatc 240

tccaggggtgg agggcgagga cgtgggagtg tactactgct tccagggcag ccacgtgcc 300

35

cctacattcg gaggcggcac caagctggag atcaag 336

<210> 387

<211> 354

<212> DNA

<213> Artificial Sequence

<220>

<223> Humanized hSC17.156 VH

45

<400> 387

caggtcacc cgaaggagtc cggccccgtg ctggtgaaac ccaccgagac cctcaccctg 60

acctgcaccg tctccggctt ctccctgtcc acctccggca tgggagtgtc ctggatcagg 120

50

cagccccctg gaaaggctct ggagtggctg gccacatct tctgggacga cgacaagtgg 180

tacaaccctt ccctgaagtc caggctgacc atctccaagg acacctcaa gtcccagggtg 240

gtgctgacca tgaccaacat ggacccccgtg gacaccgcca cctactactg cgctaccttc 300

55

tacggcctgt acttcgccta ctggggccag ggaaccctgg tgaccgtgtc ctcc 354

EP 3 338 793 A1

<210> 388
<211> 339
<212> DNA
<213> Artificial Sequence

<220>
<223> Humanized hSC17.161 VL

<400> 388
gacatcgtga tgaccagtc ccccgattcc ctggctgtga gcctgggaga gagggccacc 60
atcaactgcg agtcctccca gtccctgctg tacaactcca accagaagaa ctacctggcc 120
tggtaccagc agaagcccgg acagccccc aagctgctga tctactgggc ttccacaagg 180
gagtcgggag tgcccgatcg gttcagcggg tccggatccg gcaccgactt caccctcacc 240
atcagctccc tgcaagccga ggacgtggcc gtgtactact gccagcagta cttcaactac 300
cctctgacct tcggccaggg caccaagctg gagatcaag 339

<210> 389
<211> 348
<212> DNA
<213> Artificial Sequence

<220>
<223> Humanized hSC17.161 VH

<400> 389
caggtgcagc tgggtccagtc cggagctgag gtgaagaagc ccggcgccctc cgtgaagggtg 60
tcctgcaagg ccagcggctt caccttctcc gatgcctgga tggactgggt gaggcaggct 120
cctggccaaa ggctggagtg gatgggcgag atcaggtcca agcccaacaa ccacgccacc 180
tactacgccg agagcgtgaa gggcaggggtg accatcaciaa gggatacatc cgcctccacc 240
gcctacatgg agctgtcctc cctgaggtcc gaggacaccg ccgtgtacta ctgtgccagg 300
accggaacct cctactgggg ccagggcaca ctggtgaccg tgtcctcc 348

<210> 390
<211> 333
<212> DNA
<213> Artificial Sequence

<220>
<223> Humanized hSC17.200 VL

<400> 390
gaaattgtgt tgacacagtc tccagccacc ctgtctttgt ctccagggga aagagccacc 60
ctctcctgca gggccagtca gagtgttgac tataatggaa ttagctacat gcactggtac 120
caacagaaac ctggccaggc tcccaggctc ctcatctatg ctgcatccaa cgtgcagagt 180
ggcatcccag ccaggttcag tggcagtggt tctgggacag acttcactct caccatcagc 240
agcctagagc ctgaagattt tgcagtttat tactgtcagc agagtattga ggatcctccg 300

EP 3 338 793 A1

acgttcggtg gaggcaccaa ggtggaaatc aaa 333

5 <210> 391
<211> 345
<212> DNA
<213> Artificial Sequence

<220>
10 <223> Humanized hSC17.200 VH

<400> 391
gaggtgcagc tgggtgcagtc tggagcagag gtgaaaaagc ccggggagtc tctgaagatc 60
tcctgtaagg gttctggata cagctttacc agctcctgga tcaactgggt gcgccagatg 120
15 cccgggaaag gcctggagtg gatggggaga atctatcctg gtgaggggtga taccaactac 180
agcgggaact tcgaaggcca ggtcaccatc tcagccgaca agtccatcag caccgcctac 240
ctgcagtgga gcagcctgaa ggcctcggac accgccatgt attactgtac aagaggacta 300
20 gtcatggact actggggcca aggcaccctt gtcacagtct cgagc 345

<210> 392
25 <211> 333
<212> DNA
<213> Artificial Sequence

<220>
30 <223> Humanized hSC17.200 VL1

<400> 392
gaaattgtgt tgacacagtc tccagccacc ctgtctttgt ctccagggga aagagccacc 60
ctctcctgca gggccagtca gagtgttgac tatgatggaa ttagctacat gcactggtac 120
35 caacagaaac ctggccagtc tcccaggctc ctcatctatg ctgcatccaa cgtgcagagt 180
ggcatcccag ccaggttcag tggcagtggt tctgggacag acttcactct caccatcagc 240
agcctagagc ctgaagattt tgcagtttat tactgtcagc agagtattga ggatcctccg 300
40 acgttcggtg gaggcaccaa ggtggaaatc aaa 333

<210> 393
45 <211> 339
<212> DNA
<213> Artificial Sequence

<220>
50 <223> Humanized hSC17.155 VH1

<400> 393
caggtgcagc tgggtgcagtc tggggctgag gtgaagaagc ctggggcctc agtgaagggtt 60
tcctgcaagg catctggata caccttcgac agctactgga tgcactgggt gcgacaggcc 120
cctggacaag ggcttgagtg gatgggagaa atccacccta ataattgtag cacaaactac 180
55 aacgagaagt tcaagggcag agtcaccatg accagggaca cgtccacgag cacagtctac 240

EP 3 338 793 A1

atggagctga gcagcctgag atctgaggac acggccgtgt attactgtgc gagatggact 300
 ttgtttactt actggggcca agggactctg gtcactgtc 339

5
 <210> 394
 <211> 339
 <212> DNA
 <213> Artificial Sequence

10
 <220>
 <223> Humanized hSC17.155 VH2

<400> 394
 caggtgcagc tgggtgcagtc tggggctgag gtgaagaagc ctggggcctc agtgaagggtt 60
 15
 tcctgcaagg catctggata caccttcacc agctactgga tgcactgggt gcgacaggcc 120
 cctggacaag ggcttgagtg gatgggagaa atccacccta ataatggtag cacaaactac 180
 20
 aacgagaagt tcaagggcag agtcaccatg accagggaca cgtccacgag cacagtctac 240
 atggagctga gcagcctgag atctgaggac acggccgtgt attactgtgc gagatggact 300
 ttgtttactt actggggcca agggactctg gtcactgtc 339

25
 <210> 395
 <211> 339
 <212> DNA
 <213> Artificial Sequence

30
 <220>
 <223> Humanized hSC17.155 VH3

<400> 395
 caggtgcagc tgggtgcagtc tggggctgag gtgaagaagc ctggggcctc agtgaagggtt 60
 35
 tcctgcaagg catctggata caccttcaac tactactgga tgcactgggt gcgacaggcc 120
 cctggacaag ggcttgagtg gatgggagaa atccacccta ataatggtag cacaaactac 180
 40
 aacgagaagt tcaagggcag agtcaccatg accagggaca cgtccacgag cacagtctac 240
 atggagctga gcagcctgag atctgaggac acggccgtgt attactgtgc gagatggact 300
 ttgtttactt actggggcca agggactctg gtcactgtc 339

45
 <210> 396
 <211> 339
 <212> DNA
 <213> Artificial Sequence

50
 <220>
 <223> Humanized hSC17.155 VH4

<400> 396
 caggtgcagc tgggtgcagtc tggggctgag gtgaagaagc ctggggcctc agtgaagggtt 60
 55
 tcctgcaagg catctggata caccttcaac agctactgga tgcactgggt gcgacaggcc 120

EP 3 338 793 A1

cctggacaag ggcttgagtg gatgggagaa atccacccta atgatggtag cacaaactac 180

aacgagaagt tcaagggcag agtcaccatg accagggaca cgtccacgag cacagtctac 240

5 atggagctga gcagcctgag atctgaggac acggccgtgt attactgtgc gagatggact 300

ttgtttactt actggggcca agggactctg gtcactgtc 339

<210> 397

10 <211> 339

<212> DNA

<213> Artificial Sequence

<220>

15 <223> Humanized hSC17.155 VH5

<400> 397

caggtgcagc tgggtgcagtc tggggctgag gtgaagaagc ctggggcctc agtgaagggt 60

tcctgcaagg catctggata caccttcaac agctactgga tgcactgggt gcgacaggcc 120

20 cctggacaag ggcttgagtg gatgggagaa atccacccta atggtggtag cacaaactac 180

aacgagaagt tcaagggcag agtcaccatg accagggaca cgtccacgag cacagtctac 240

25 atggagctga gcagcctgag atctgaggac acggccgtgt attactgtgc gagatggact 300

ttgtttactt actggggcca agggactctg gtcactgtc 339

<210> 398

30 <211> 339

<212> DNA

<213> Artificial Sequence

<220>

35 <223> Humanized hSC17.155 VH6

<400> 398

caggtgcagc tgggtgcagtc tggggctgag gtgaagaagc ctggggcctc agtgaagggt 60

tcctgcaagg catctggata caccttcaac agctactgga tgcactgggt gcgacaggcc 120

40 cctggacaag ggcttgagtg gatgggagaa atccacccta atagtgttag cacaaactac 180

aacgagaagt tcaagggcag agtcaccatg accagggaca cgtccacgag cacagtctac 240

45 atggagctga gcagcctgag atctgaggac acggccgtgt attactgtgc gagatggact 300

ttgtttactt actggggcca agggactctg gtcactgtc 339

<210> 399

50 <211> 348

<212> DNA

<213> Artificial Sequence

<220>

55 <223> Humanized hSC17.161 VH1

<400> 399

gaggtgcagc tgggtggaatc cggaggcggc ctggtgcaac ctggaggatc cctcaggctg 60

EP 3 338 793 A1

tcctgtgccg cttccggatt caccttctcc gatgcctgga tggactgggt gaggcaggcc 120
 cctggcaaag gactggaatg ggtgggagag atcaggtcca aaccaacaa ccacgccacc 180
 5 tactacgccg agtccgtgaa gggcaggttc accatctcca gggacgactc caagaactcc 240
 ctgtacctgc agatgaactc cctgaagacc gaggaacacc ccgtgtacta ctgcgctagg 300
 accggcacct cctattgggg acagggcacc ctggtgaccg tgcctctcc 348
 10
 <210> 400
 <211> 218
 <212> PRT
 <213> Artificial Sequence
 15
 <220>
 <223> Humanized hSC17.200 light chain
 <400> 400
 20 Glu Ile Val Leu Thr Gln Ser Pro Ala Thr Leu Ser Leu Ser Pro Gly
 1 5 10 15
 Glu Arg Ala Thr Leu Ser Cys Arg Ala Ser Gln Ser Val Asp Tyr Asn
 25 20 25 30
 Gly Ile Ser Tyr Met His Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro
 35 35 40 45
 30 Arg Leu Leu Ile Tyr Ala Ala Ser Asn Val Gln Ser Gly Ile Pro Ala
 50 55 60
 Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser
 35 65 70 75 80
 Ser Leu Glu Pro Glu Asp Phe Ala Val Tyr Tyr Cys Gln Gln Ser Ile
 40 85 90 95
 Glu Asp Pro Pro Thr Phe Gly Gly Gly Thr Lys Val Glu Ile Lys Arg
 100 105 110
 45 Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln
 115 120 125
 Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr
 50 130 135 140
 Pro Arg Glu Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser
 145 150 155 160
 55 Gly Asn Ser Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr

EP 3 338 793 A1

	165	170	175
5	Tyr Ser Leu Ser Ser Thr Leu Thr	Leu Ser Lys Ala Asp	Tyr Glu Lys
	180	185	190
10	His Lys Val Tyr Ala Cys Glu Val Thr His Gln Gly	Leu Ser Ser Pro	
	195	200	205
15	Val Thr Lys Ser Phe Asn Arg Gly Glu Cys		
	210	215	
20	<210> 401		
	<211> 444		
	<212> PRT		
	<213> Artificial Sequence		
	<220>		
	<223> Humanized hSC17.200 heavy chain		
	<400> 401		
25	Glu Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Glu		
	1	5	10 15
30	Ser Leu Lys Ile Ser Cys Lys Gly Ser Gly Tyr Ser Phe Thr Ser Ser		
	20	25	30
35	Trp Ile Asn Trp Val Arg Gln Met Pro Gly Lys Gly Leu Glu Trp Met		
	35	40	45
40	Gly Arg Ile Tyr Pro Gly Glu Gly Asp Thr Asn Tyr Ser Gly Asn Phe		
	50	55	60
45	Glu Gly Gln Val Thr Ile Ser Ala Asp Lys Ser Ile Ser Thr Ala Tyr		
	65	70	75 80
50	Leu Gln Trp Ser Ser Leu Lys Ala Ser Asp Thr Ala Met Tyr Tyr Cys		
	85	90	95
55	Thr Arg Gly Leu Val Met Asp Tyr Trp Gly Gln Gly Thr Leu Val Thr		
	100	105	110
60	Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro		
	115	120	125
65	Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly Cys Leu Val		
	130	135	140
70	Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala		
	145	150	155 160

EP 3 338 793 A1

	Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly	165	170	175
5	Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly	180	185	190
10	Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys Pro Ser Asn Thr Lys	195	200	205
15	Val Asp Lys Lys Val Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys	210	215	220
20	Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser Val Phe Leu	225	230	235
25	Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu	245	250	255
30	Val Thr Cys Val Val Val Asp Val Ser His Glu Asp Pro Glu Val Lys	260	265	270
35	Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys	275	280	285
40	Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu	290	295	300
45	Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys	305	310	315
50	Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys	325	330	335
55	Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser	340	345	350
	Arg Asp Glu Leu Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys	355	360	365
	Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln	370	375	380
	Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly	385	390	395
	Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln			

EP 3 338 793 A1

```

5      Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn
      420                                425                                430

10     His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly
      435                                440

15     <210> 402
      <211> 218
      <212> PRT
      <213> Artificial Sequence

20     <220>
      <223> Humanized hSC17.200vL1 light chain

      <400> 402

25     Glu Ile Val Leu Thr Gln Ser Pro Ala Thr Leu Ser Leu Ser Pro Gly
      1           5           10           15

30     Glu Arg Ala Thr Leu Ser Cys Arg Ala Ser Gln Ser Val Asp Tyr Asp
      20           25           30

35     Gly Ile Ser Tyr Met His Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro
      35           40           45

40     Arg Leu Leu Ile Tyr Ala Ala Ser Asn Val Gln Ser Gly Ile Pro Ala
      50           55           60

45     Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser
      65           70           75           80

50     Ser Leu Glu Pro Glu Asp Phe Ala Val Tyr Tyr Cys Gln Gln Ser Ile
      85           90           95

55     Glu Asp Pro Pro Thr Phe Gly Gly Gly Thr Lys Val Glu Ile Lys Arg
      100          105          110

60     Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln
      115          120          125

65     Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr
      130          135          140

70     Pro Arg Glu Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser
      145          150          155          160

75     Gly Asn Ser Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr
      165          170          175

```

EP 3 338 793 A1

	Tyr	Ser	Leu	Ser	Ser	Thr	Leu	Thr	Leu	Ser	Lys	Ala	Asp	Tyr	Glu	Lys	
				180					185					190			
5	His	Lys	Val	Tyr	Ala	Cys	Glu	Val	Thr	His	Gln	Gly	Leu	Ser	Ser	Pro	
			195					200					205				
10	Val	Thr	Lys	Ser	Phe	Asn	Arg	Gly	Glu	Cys							
		210					215										
	<210>	403															
	<211>	107															
	<212>	PRT															
15	<213>	Homo sapiens															
	<220>																
	<221>	MISC_FEATURE															
20	<223>	Kappa light chain constant region															
	<400>	403															
25	Arg	Thr	Val	Ala	Ala	Pro	Ser	Val	Phe	Ile	Phe	Pro	Pro	Ser	Asp	Glu	
	1				5					10					15		
	Gln	Leu	Lys	Ser	Gly	Thr	Ala	Ser	Val	Val	Cys	Leu	Leu	Asn	Asn	Phe	
				20					25					30			
30	Tyr	Pro	Arg	Glu	Ala	Lys	Val	Gln	Trp	Lys	Val	Asp	Asn	Ala	Leu	Gln	
			35					40					45				
35	Ser	Gly	Asn	Ser	Gln	Glu	Ser	Val	Thr	Glu	Gln	Asp	Ser	Lys	Asp	Ser	
		50					55					60					
40	Thr	Tyr	Ser	Leu	Ser	Ser	Thr	Leu	Thr	Leu	Ser	Lys	Ala	Asp	Tyr	Glu	
	65					70					75					80	
	Lys	His	Lys	Val	Tyr	Ala	Cys	Glu	Val	Thr	His	Gln	Gly	Leu	Ser	Ser	
					85					90					95		
45	Pro	Val	Thr	Lys	Ser	Phe	Asn	Arg	Gly	Glu	Cys						
				100					105								
	<210>	404															
50	<211>	329															
	<212>	PRT															
	<213>	Homo sapiens															
	<220>																
	<221>	MISC_FEATURE															
55	<223>	IgG1 heavy chain constant region															

EP 3 338 793 A1

<400> 404

5	Ala	Ser	Thr	Lys	Gly	Pro	Ser	Val	Phe	Pro	Leu	Ala	Pro	Ser	Ser	Lys	1	5	10	15
10	Ser	Thr	Ser	Gly	Gly	Thr	Ala	Ala	Leu	Gly	Cys	Leu	Val	Lys	Asp	Tyr	20	25	30	
15	Phe	Pro	Glu	Pro	Val	Thr	Val	Ser	Trp	Asn	Ser	Gly	Ala	Leu	Thr	Ser	35	40	45	
20	Gly	Val	His	Thr	Phe	Pro	Ala	Val	Leu	Gln	Ser	Ser	Gly	Leu	Tyr	Ser	50	55	60	
25	Leu	Ser	Ser	Val	Val	Thr	Val	Pro	Ser	Ser	Ser	Leu	Gly	Thr	Gln	Thr	65	70	75	80
30	Tyr	Ile	Cys	Asn	Val	Asn	His	Lys	Pro	Ser	Asn	Thr	Lys	Val	Asp	Lys	85	90	95	
35	Lys	Val	Glu	Pro	Lys	Ser	Cys	Asp	Lys	Thr	His	Thr	Cys	Pro	Pro	Cys	100	105	110	
40	Pro	Ala	Pro	Glu	Leu	Leu	Gly	Gly	Pro	Ser	Val	Phe	Leu	Phe	Pro	Pro	115	120	125	
45	Lys	Pro	Lys	Asp	Thr	Leu	Met	Ile	Ser	Arg	Thr	Pro	Glu	Val	Thr	Cys	130	135	140	
50	Val	Val	Val	Asp	Val	Ser	His	Glu	Asp	Pro	Glu	Val	Lys	Phe	Asn	Trp	145	150	155	160
55	Tyr	Val	Asp	Gly	Val	Glu	Val	His	Asn	Ala	Lys	Thr	Lys	Pro	Arg	Glu	165	170	175	
	Glu	Gln	Tyr	Asn	Ser	Thr	Tyr	Arg	Val	Val	Ser	Val	Leu	Thr	Val	Leu	180	185	190	
	His	Gln	Asp	Trp	Leu	Asn	Gly	Lys	Glu	Tyr	Lys	Cys	Lys	Val	Ser	Asn	195	200	205	
	Lys	Ala	Leu	Pro	Ala	Pro	Ile	Glu	Lys	Thr	Ile	Ser	Lys	Ala	Lys	Gly	210	215	220	
	Gln	Pro	Arg	Glu	Pro	Gln	Val	Tyr	Thr	Leu	Pro	Pro	Ser	Arg	Asp	Glu	225	230	235	240

EP 3 338 793 A1

Leu Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr
245 250 255

5 Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn
260 265 270

10 Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe
275 280 285

Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn
290 295 300

15 Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr
305 310 315 320

20 Gln Lys Ser Leu Ser Leu Ser Pro Gly
325

<210> 405
<211> 11
<212> PRT
25 <213> Mus musculus

<220>
<221> MISC_FEATURE
30 <223> hSC17.16 CDRL1

<400> 405

Arg Ala Ser Ala Asn Ile Asn Ser Asn Leu Val
1 5 10

35

<210> 406
<211> 7
<212> PRT
40 <213> Mus musculus

<220>
<221> MISC_FEATURE
45 <223> hSC17.16 CDRL2

<400> 406

Ala Ala Thr Asn Leu Ala Asp
1 5

50

<210> 407
<211> 9
<212> PRT
55 <213> Mus musculus

<220>

EP 3 338 793 A1

<221> MISC_FEATURE
<223> hSC17.16 CDRL3

<400> 407

5

Gln His Phe Trp Gly Thr Pro Arg Thr
1 5

<210> 408
<211> 5
<212> PRT
<213> Mus musculus

10

<220>
<221> MISC_FEATURE
<223> hSC17.16 CDRH1

<400> 408

15

Asp Tyr Asn Met Tyr
1 5

20

<210> 409
<211> 17
<212> PRT
<213> Mus musculus

25

<220>
<221> MISC_FEATURE
<223> hSC17.16 CDRH2

<400> 409

30

Glu Ile Asn Pro Asn Asn Gly Gly Thr Ala Tyr Asn Gln Lys Phe Arg
1 5 10 15

35

Gly

40

<210> 410
<211> 7
<212> PRT
<213> Mus musculus

45

<220>
<221> MISC_FEATURE
<223> hSC17.16 CDRH3

<400> 410

50

Tyr Asp Lys Gly Phe Asp Tyr
1 5

55

<210> 411
<211> 10

EP 3 338 793 A1

```

<212>  PRT
<213>  Mus musculus

5      <220>
      <221>  MISC_FEATURE
      <223>  hSC17.17 CDRL1

      <400>  411

10     Ser Ala Ser Ser Ser Val Ser Tyr Met His
      1              5              10

      <210>  412
      <211>  7
      <212>  PRT
      <213>  Mus musculus

15     <220>
      <221>  MISC_FEATURE
      <223>  hSC17.17 CDRL2

      <400>  412

20     Asp Thr Ser Lys Leu Pro Ser
      1              5

      <210>  413
      <211>  9
      <212>  PRT
      <213>  Mus musculus

25     <220>
      <221>  MISC_FEATURE
      <223>  hSC17.17 CDRL3

      <400>  413

30     Gln Gln Trp Ser Ser Thr Pro Pro Thr
      1              5

      <210>  414
      <211>  5
      <212>  PRT
      <213>  Mus musculus

35     <220>
      <221>  MISC_FEATURE
      <223>  hSC17.17 CDRH1

      <400>  414

40     Asp Tyr Asn Met His
      1              5

45     <210>  415

```

EP 3 338 793 A1

<211> 17
<212> PRT
<213> Mus musculus

5

<220>
<221> MISC_FEATURE
<223> hSC17.17 CDRH2

10

<400> 415

Glu Ile Asn Pro Asn Ile Gly Gly Thr Gly Tyr Asn Gln Lys Phe Lys
1 5 10 15

15

Gly

<210> 416
<211> 11
<212> PRT
<213> Mus musculus

20

<220>
<221> MISC_FEATURE
<223> hSC17.17 CDRH3

25

<400> 416

Thr Tyr Ser Tyr Tyr Ser Tyr Glu Phe Ala Tyr
1 5 10

30

<210> 417
<211> 17
<212> PRT
<213> Mus musculus

35

<220>
<221> MISC_FEATURE
<223> hSC17.24 CDRL1

40

<400> 417

Lys Ser Ser Gln Ser Leu Leu Tyr Ser Ser Asn Gln Lys Ser Tyr Leu
1 5 10 15

45

Ala

<210> 418
<211> 7
<212> PRT
<213> Mus musculus

50

<220>
<221> MISC_FEATURE

55

EP 3 338 793 A1

<223> hSC17.24 CDRL2

<400> 418

5 Trp Ala Ser Thr Arg Glu Ser
1 5

<210> 419

<211> 8

10 <212> PRT

<213> Mus musculus

<220>

<221> MISC_FEATURE

15 <223> hSC17.24 CDRL3

<400> 419

20 Lys Gln Ser Tyr Asn Leu Arg Thr
1 5

<210> 420

<211> 5

25 <212> PRT

<213> Mus musculus

<220>

<221> MISC_FEATURE

30 <223> hSC17.24 CDRH1

<400> 420

35 Asp His Thr Ile His
1 5

<210> 421

<211> 17

40 <212> PRT

<213> Mus musculus

<220>

<221> MISC_FEATURE

45 <223> hSC17.24 CDRH2

<400> 421

50 Tyr Ile Tyr Pro Arg Asp Gly Ser Thr Lys Tyr Asn Glu Glu Phe Lys
1 5 10 15

Gly

<210> 422

<211> 8

55 <212> PRT

EP 3 338 793 A1

<213> Mus musculus

<220>

<221> MISC_FEATURE

<223> hSC17.24 CDRH3

<400> 422

Ser Tyr Ser Asn Tyr Phe Asp Tyr

1 5

<210> 423

<211> 11

<212> PRT

<213> Mus musculus

<220>

<221> MISC_FEATURE

<223> hSC17.28 CDRL1

<400> 423

Arg Ala Ser Gln Ser Ile Gly Thr Ser Ile His

1 5 10

<210> 424

<211> 7

<212> PRT

<213> Mus musculus

<220>

<221> MISC_FEATURE

<223> hSC17.28 CDRL2

<400> 424

Tyr Ala Ser Glu Ser Ile Ser

1 5

<210> 425

<211> 9

<212> PRT

<213> Mus musculus

<220>

<221> MISC_FEATURE

<223> hSC17.28 CDRL3

<400> 425

Gln Gln Ser Asn Ser Trp Pro Leu Thr

1 5

<210> 426

<211> 5

EP 3 338 793 A1

<212> PRT
 <213> Mus musculus

 5 <220>
 <221> MISC_FEATURE
 <223> hSC17.28 CDRH1

 <400> 426

 10 Arg Ser Tyr Ile His
 1 5

 <210> 427
 15 <211> 17
 <212> PRT
 <213> Mus musculus

 <220>
 20 <221> MISC_FEATURE
 <223> hSC17.28 CDRH2

 <400> 427

 25 Tyr Ile Ser Ser Gly Ser Gly Gly Thr Thr Tyr Asn Gln Lys Phe Lys
 1 5 10 15

 Gly

 30 <210> 428
 <211> 8
 <212> PRT
 <213> Mus musculus

 35 <220>
 <221> MISC_FEATURE
 <223> hSC17.28 CDRH3

 40 <400> 428

 Gly Gly Val Arg Tyr Phe Asp Val
 1 5

 45 <210> 429
 <211> 11
 <212> PRT
 <213> Mus musculus

 50 <220>
 <221> MISC_FEATURE
 <223> hSC17.34 CDRL1

 55 <400> 429

 Lys Ala Ser Gln Asp Ile Asn Ser Tyr Leu Ser

EP 3 338 793 A1

1 5 10

5
 <210> 430
 <211> 7
 <212> PRT
 <213> Mus musculus

10
 <220>
 <221> MISC_FEATURE
 <223> hSC17.34 CDRL2

<400> 430

15
 Arg Ala Asn Arg Leu Val Asp
 1 5

20
 <210> 431
 <211> 9
 <212> PRT
 <213> Mus musculus

25
 <220>
 <221> MISC_FEATURE
 <223> hSC17.34 CDRL1

30
 <220>
 <221> MISC_FEATURE
 <223> hSC17.34 CDRL3

<400> 431

35
 Leu Gln Tyr Asp Glu Phe Pro Pro Thr
 1 5

40
 <210> 432
 <211> 5
 <212> PRT
 <213> Mus musculus

45
 <220>
 <221> MISC_FEATURE
 <223> hSC17.34 CDRH1

<400> 432

50
 Asp Tyr Asn Met Asp
 1 5

55
 <210> 433
 <211> 17
 <212> PRT
 <213> Mus musculus

<220>
 <221> MISC_FEATURE

EP 3 338 793 A1

<223> hSC17.34 CDRH1

<400> 433

5 Tyr Ile Tyr Pro Asp Asn Gly Gly Ala Gly Tyr Asn Gln Lys Phe Lys
1 5 10 15

Gly

10

<210> 434

<211> 9

<212> PRT

15

<213> Mus musculus

<220>

<221> MISC_FEATURE

20

<223> hSC17.34 CDRH3

<400> 434

Ser Ile Thr Thr Ala Trp Phe Ala Tyr
1 5

25

<210> 435

<211> 11

<212> PRT

30

<213> Mus musculus

<220>

<221> MISC_FEATURE

35

<223> hSC17.46 CDRL1

<400> 435

Lys Ala Ser Gln Ser Val Asn Asn Asp Val Ala
1 5 10

40

<210> 436

<211> 7

<212> PRT

<213> Mus musculus

45

<220>

<221> MISC_FEATURE

<223> hSC17.46 CDRL2

<400> 436

50

Tyr Ala Ser Asn Arg Tyr Thr
1 5

55

<210> 437

<211> 9

<212> PRT

EP 3 338 793 A1

<213> Mus musculus

<220>
 5 <221> MISC_FEATURE
 <223> hSC17.46 CDRL3

<400> 437

10 Gln Gln Asp Tyr Ser Ser Pro Arg Thr
 1 5

<210> 438
 <211> 5
 15 <212> PRT
 <213> Mus musculus

<220>
 20 <221> MISC_FEATURE
 <223> hSC17.46 CDRH1

<400> 438

25 Ser Tyr Trp Ile Asn
 1 5

<210> 439
 <211> 17
 30 <212> PRT
 <213> Mus musculus

<220>
 35 <221> MISC_FEATURE
 <223> hSC17.46 CDRH2

<400> 439

40 Asn Ile Phe Pro Asp Thr Thr Thr Thr Asn Tyr Asn Glu Lys Phe Lys
 1 5 10 15

Gly

45 <210> 440
 <211> 12
 <212> PRT
 <213> Mus musculus

50 <220>
 <221> MISC_FEATURE
 <223> hSC17.46 CDRH1

<400> 440

55 Glu Tyr Tyr Asp Gly Thr Tyr Asp Ala Met Asp Tyr
 1 5 10

EP 3 338 793 A1

<210> 441
 <211> 15
 <212> PRT
 <213> Mus musculus

5

<220>
 <221> MISC_FEATURE
 <223> hSC17.151 CDRL1

10

<400> 441

Arg Ala Ser Glu Ser Val Asp Ser Tyr Gly Asn Ser Phe Met His
 1 5 10 15

15

<210> 442
 <211> 7
 <212> PRT
 <213> Mus musculus

20

<220>
 <221> MISC_FEATURE
 <223> hSC17.151 CDRL2

25

<400> 442

Arg Ala Ser Asn Leu Glu Ser
 1 5

30

<210> 443
 <211> 9
 <212> PRT
 <213> Mus musculus

35

<220>
 <221> MISC_FEATURE
 <223> hSC17.151 CDRL3

40

<400> 443

Gln Gln Ser His Glu Asp Pro Tyr Thr
 1 5

45

<210> 444
 <211> 5
 <212> PRT
 <213> Mus musculus

50

<220>
 <221> MISC_FEATURE
 <223> hSC17.151 CDRH1

55

<400> 444

Ser Tyr Trp Met His

EP 3 338 793 A1

1 5

<210> 445
<211> 17
<212> PRT
<213> Mus musculus

<220>
<221> MISC_FEATURE
<223> hSC17.151 CDRH2

<400> 445

Ala Ile Tyr Pro Gly Lys Ser Asp Thr Thr Tyr Asn Gln Lys Phe Lys
1 5 10 15

Gly

<210> 446
<211> 8
<212> PRT
<213> Mus musculus

<220>
<221> MISC_FEATURE
<223> hSC17.151 CDRH3

<400> 446

Ser Gly Lys Gly Tyr Phe Ala Tyr
1 5

<210> 447
<211> 17
<212> PRT
<213> Mus musculus

<220>
<221> MISC_FEATURE
<223> hSC17.155 and hSC17.155vH1-6 CDRL1

<400> 447

Lys Ser Ser Gln Ser Leu Leu Tyr Ser Ser Asn Gln Lys Asn Tyr Leu
1 5 10 15

Ala

<210> 448
<211> 7
<212> PRT
<213> Mus musculus

<220>
 <221> MISC_FEATURE
 <223> hSC17.155 and hSC17.155vH1-6 CDRL2

5

<400> 448

Trp Ala Ser Thr Arg Lys Ser
 1 5

10

<210> 449
 <211> 9
 <212> PRT
 <213> Mus musculus

15

<220>
 <221> MISC_FEATURE
 <223> hSC17.155 and hSC17.155vH1-6 CDRL3

20

<400> 449

His Gln Tyr Tyr Ser Tyr Pro Leu Thr
 1 5

25

<210> 450
 <211> 5
 <212> PRT
 <213> Mus musculus

30

<220>
 <221> MISC_FEATURE
 <223> hSC17.155 and hSC17.155vH1, vH2 and vH4-6 CDRH1

35

<400> 450

Ser Tyr Trp Met His
 1 5

40

<210> 451
 <211> 17
 <212> PRT
 <213> Mus musculus

45

<220>
 <221> MISC_FEATURE
 <223> hSC17.155 and hSC17.155vH1-3 CDRH2

50

<400> 451

Glu Ile His Pro Asn Asn Gly Ser Thr Asn Tyr Asn Glu Lys Phe Lys
 1 5 10 15

55

Gly

EP 3 338 793 A1

<210> 452
<211> 6
<212> PRT
<213> Mus musculus

5

<220>
<221> MISC_FEATURE
<223> hSC17.155 and hSC17.155vH1-6 CDRH3

10

<400> 452

Trp Thr Leu Phe Thr Tyr
1 5

15

<210> 453
<211> 16
<212> PRT
<213> Mus musculus

20

<220>
<221> MISC_FEATURE
<223> hSC17.156 CDRL1

25

<400> 453

Arg Ser Ser Gln Ser Ile Val His Ser Asn Gly Asn Thr Tyr Leu Glu
1 5 10 15

30

<210> 454
<211> 7
<212> PRT
<213> Mus musculus

35

<220>
<221> MISC_FEATURE
<223> hSC17.156 CDRL2

<400> 454

40

Lys Val Ser Asn Arg Phe Ser
1 5

45

<210> 455
<211> 9
<212> PRT
<213> Mus musculus

50

<220>
<221> MISC_FEATURE
<223> hSC17.156 CDRL3

<400> 455

55

Phe Gln Gly Ser His Val Pro Pro Thr
1 5

EP 3 338 793 A1

<210> 456
 <211> 7
 <212> PRT
 <213> Mus musculus

5

<220>
 <221> MISC_FEATURE
 <223> hSC17.156 CDRH1

10

<400> 456

Thr Ser Gly Met Gly Val Ser
 1 5

15

<210> 457
 <211> 16
 <212> PRT
 <213> Mus musculus

20

<220>
 <221> MISC_FEATURE
 <223> hSC17.156 CDRH2

25

<400> 457

His Ile Phe Trp Asp Asp Asp Lys Trp Tyr Asn Pro Ser Leu Lys Ser
 1 5 10 15

30

<210> 458
 <211> 8
 <212> PRT
 <213> Mus musculus

35

<220>
 <221> MISC_FEATURE
 <223> hSC17.156 CDRH3

40

<400> 458

Phe Tyr Gly Leu Tyr Phe Ala Tyr
 1 5

45

<210> 459
 <211> 17
 <212> PRT
 <213> Mus musculus

50

<220>
 <221> MISC_FEATURE
 <223> hSC17.161 and hSC17.161v1 CDRL1

55

<400> 459

Glu Ser Ser Gln Ser Leu Leu Tyr Asn Ser Asn Gln Lys Asn Tyr Leu

EP 3 338 793 A1

1 5 10 15

Ala

5

<210> 460
<211> 7
<212> PRT
<213> Mus musculus

10

<220>
<221> MISC_FEATURE
<223> hSC17.161 and hSC17.161vL1 CDRL2

15

<400> 460

Trp Ala Ser Thr Arg Glu Ser
1 5

20

<210> 461
<211> 9
<212> PRT
<213> Mus musculus

25

<220>
<221> MISC_FEATURE
<223> hSC17.161 and hSC17.161vL1 CDRL3

30

<400> 461

Gln Gln Tyr Phe Asn Tyr Pro Leu Thr
1 5

35

<210> 462
<211> 5
<212> PRT
<213> Mus musculus

40

<220>
<221> MISC_FEATURE
<223> hSC17.161 and hSC17.161vL1 CDRH1

45

<400> 462

Asp Ala Trp Met Asp
1 5

50

<210> 463
<211> 19
<212> PRT
<213> Mus musculus

55

<220>
<221> MISC_FEATURE

EP 3 338 793 A1

<223> hSC17.161 and hSC17.161vL1 CDRH2

<400> 463

5 Glu Ile Arg Ser Lys Pro Asn Asn His Ala Thr Tyr Tyr Ala Glu Ser
1 5 10 15

Val Lys Gly

10

<210> 464

<211> 5

<212> PRT

15

<213> Mus musculus

<220>

<221> MISC_FEATURE

20

<223> hSC17.161 and hSC17.161vL1 CDRH3

<400> 464

Thr Gly Thr Ser Tyr

1 5

25

<210> 465

<211> 15

<212> PRT

30

<213> Mus musculus

<220>

<221> MISC_FEATURE

<223> hSC17.200 CDRL1

35

<400> 465

Arg Ala Ser Gln Ser Val Asp Tyr Asn Gly Ile Ser Tyr Met His

1 5 10 15

40

<210> 466

<211> 7

<212> PRT

<213> Mus musculus

45

<220>

<221> MISC_FEATURE

<223> hSC17.200 and hSC17.200vL1 CDRL2

<400> 466

50

Ala Ala Ser Asn Val Gln Ser

1 5

55

<210> 467

<211> 9

<212> PRT

EP 3 338 793 A1

<213> Mus musculus

<220>

<221> MISC_FEATURE

<223> hSC17.200 and hSC17.200vL1 CDRL3

<400> 467

Gln Gln Ser Ile Glu Asp Pro Pro Thr

1 5

<210> 468

<211> 5

<212> PRT

<213> Mus musculus

<220>

<221> MISC_FEATURE

<223> hSC17.200 and hSC17.200vL1 CDRH1

<400> 468

Ser Ser Trp Ile Asn

1 5

<210> 469

<211> 17

<212> PRT

<213> Mus musculus

<220>

<221> MISC_FEATURE

<223> hSC17.200 and hSC17.200vL1 CDRH2

<400> 469

Arg Ile Tyr Pro Gly Glu Gly Asp Thr Asn Tyr Ser Gly Asn Phe Glu

1 5 10 15

Gly

<210> 470

<211> 6

<212> PRT

<213> Mus musculus

<220>

<221> MISC_FEATURE

<223> hSC17.200 and hSC17.200vL1 CDRH3

<400> 470

Gly Leu Val Met Asp Tyr

1 5

EP 3 338 793 A1

5
 <210> 471
 <211> 30
 <212> PRT
 <213> Homo sapiens

10
 <220>
 <221> MISC_FEATURE
 <223> hSC17.155vH1 FR1

15
 <400> 471
 Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala
 1 5 10 15

20
 Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Asp
 20 25 30

25
 <210> 472
 <211> 30
 <212> PRT
 <213> Homo sapiens

30
 <220>
 <221> MISC_FEATURE
 <223> hSC17.155vH2 FR1

35
 <400> 472
 Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala
 1 5 10 15

40
 Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr
 20 25 30

45
 <210> 473
 <211> 5
 <212> PRT
 <213> Mus musculus

50
 <220>
 <221> MISC_FEATURE
 <223> hSC17.155vH3 CDRH1

55
 <400> 473
 Tyr Tyr Trp Met His
 1 5

60
 <210> 474
 <211> 17
 <212> PRT
 <213> Mus musculus

EP 3 338 793 A1

<220>
 <221> MISC_FEATURE
 <223> hSC17.155vH4 CDRH2

5

<400> 474

Glu Ile His Pro Asn Asp Gly Ser Thr Asn Tyr Asn Glu Lys Phe Lys
 1 5 10 15

10

Gly

<210> 475
 <211> 17
 <212> PRT
 <213> Mus musculus

15

<220>
 <221> MISC_FEATURE
 <223> hSC17.155vH5 CDRH2

20

<400> 475

25

Glu Ile His Pro Asn Gly Gly Ser Thr Asn Tyr Asn Glu Lys Phe Lys
 1 5 10 15

Gly

30

<210> 476
 <211> 17
 <212> PRT
 <213> Mus musculus

35

<220>
 <221> MISC_FEATURE
 <223> hSC17.155vH6 CDRH2

40

<400> 476

Glu Ile His Pro Asn Ser Gly Ser Thr Asn Tyr Asn Glu Lys Phe Lys
 1 5 10 15

45

Gly

<210> 477
 <211> 30
 <212> PRT
 <213> Homo sapiens

50

<220>
 <221> MISC_FEATURE

55

EP 3 338 793 A1

<223> hSC17.161vH1 FR1

<400> 477

5 Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1 5 10 15

10 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser
20 25 30

<210> 478

<211> 14

<212> PRT

15 <213> Homo sapiens

<220>

<221> MISC_FEATURE

20 <223> hSC17.161vH1 FR2

<400> 478

25 Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val Gly
1 5 10

<210> 479

<211> 32

<212> PRT

30 <213> Homo sapiens

<220>

<221> MISC_FEATURE

35 <223> hSC17.161vH1 FR3

<400> 479

40 Arg Phe Thr Ile Ser Arg Asp Asp Ser Lys Asn Ser Leu Tyr Leu Gln
1 5 10 15

Met Asn Ser Leu Lys Thr Glu Asp Thr Ala Val Tyr Tyr Cys Ala Arg
20 25 30

<210> 480

<211> 15

<212> PRT

45 <213> Mus musculus

<220>

<221> MISC_FEATURE

50 <223> hSC17.200vL1 CDRL1

<400> 480

55 Arg Ala Ser Gln Ser Val Asp Tyr Asp Gly Ile Ser Tyr Met His
1 5 10 15

Claims

1. A pharmaceutical composition for use in treating (i) chemoresistant small cell lung cancer or (ii) medullary thyroid cancer, wherein the pharmaceutical composition comprises an anti-SEZ6 antibody conjugated, linked, or otherwise associated with a cytotoxic agent.
2. A kit comprising:
 - (a) one or more containers comprising an antibody drug conjugate, wherein the antibody drug conjugate comprises an anti-SEZ6 antibody conjugated, linked, or otherwise associated with a cytotoxic agent and
 - (b) a label or package insert on or associated with the one or more containers, wherein the label or package insert indicates that the antibody drug conjugate is used for treating (i) chemoresistant small cell lung cancer or (ii) medullary thyroid cancer.
3. A method of diagnosing (i) chemoresistant small cell lung cancer or (ii) medullary thyroid cancer, the method comprising:
 - (a) contacting a chemoresistant small cell lung cancer tumor sample or medullary thyroid tumor sample with an antibody that binds to human SEZ6; and
 - (b) detecting the antibody bound to the tumor sample.
4. The pharmaceutical composition for use according to claim 1 or the kit according to claim 2, wherein the antibody drug conjugate comprises the formula M-[L-D]_n, wherein
 - M comprises the anti-SEZ6 antibody;
 - L comprises an optional linker;
 - D comprises the cytotoxic agent; and
 - n is an integer from 1 to 20.
5. The pharmaceutical composition for use according to claim 1 or 4, the kit according to claim 2 or 4, or the method according to claim 3 or 4, wherein the antibody comprises or competes for binding to a human SEZ6 protein with an antibody comprising:
 - (a) a light chain variable region of SEQ ID NO: 190 or SEQ ID NO: 192, and a heavy chain variable region of SEQ ID NO: 191 or
 - (b) a light chain variable region of SEQ ID NO: 172 and a heavy chain variable region of SEQ ID NO: 173.
6. The pharmaceutical composition for use according to any one of claims 1, 4 or 5, the kit according to any one of claims 2, 4 or 5, or the method according to any one of claims 3-5, wherein the antibody comprises a monoclonal antibody, chimeric antibody, CDR-grafted antibody, humanized antibody, human antibody, multispecific antibody, bispecific antibody, monovalent antibody, multivalent antibody, Fab fragment, F(ab')₂ fragment, Fv fragment, or ScFv fragment; or an immunoreactive fragment thereof.
7. The pharmaceutical composition for use according to any one of claims 1 or 4-6, the kit according to any one of claims 2 or 4-6, or the method of any one of claims 3-6, wherein the antibody comprises a chimeric antibody, CDR-grafted antibody, or humanized antibody.
8. The pharmaceutical composition for use according to any one of claims 1 or 4-7, the kit according to any one of claims 2 or 4-7, or the method of any one of claims 3-7, wherein the antibody comprises:
 - (a) three complementarity determining regions of a light chain variable region of SEQ ID NO: 190 or SEQ ID NO: 192, and three complementarity determining regions of a heavy chain variable region of SEQ ID NO: 191; or
 - (b) three complementarity determining regions of a light chain variable region of SEQ ID NO: 172 and three complementarity determining regions of a heavy chain variable region of SEQ ID NO: 173.
9. The pharmaceutical composition for use according to any one of claims 1 or 4-8, the kit according to any one of claims 2 or 4-8, or the method of any one of claims 3-8, wherein the antibody comprises:
 - (a) residues 24-34 of SEQ ID NO: 190 for CDR-L1, residues 50-56 of SEQ ID NO: 190 for CDR-L2, residues

89-97 of SEQ ID NO: 190 for CDR-L3, residues 31-35 of SEQ ID NO: 191 for CDR-H1, residues 50-65 of SEQ ID NO: 191 for CDR-H2 and residues 95-102 of SEQ ID NO: 191 for CDR-H3, wherein the residues are numbered according to Kabat;

(b) residues 23-34 of SEQ ID NO: 190 for CDR-L1, residues 50-56 of SEQ ID NO: 190 for CDR-L2, residues 89-97 of SEQ ID NO: 190 for CDR-L3, residues 26-32 of SEQ ID NO: 191 for CDR-H1, residues 50-58 of SEQ ID NO: 191 for CDR-H2 and residues 95-102 of SEQ ID NO: 191 for CDR-H3, wherein the residues are numbered according to Chothia;

(c) residues 30-36 of SEQ ID NO: 190 for CDR-L1, residues 46-55 of SEQ ID NO: 190 for CDR-L2, residues 89-96 of SEQ ID NO: 190 for CDR-L3, residues 30-35 of SEQ ID NO: 191 for CDR-H1, residues 47-58 of SEQ ID NO: 191 for CDR-H2 and residues 93-101 of SEQ ID NO: 191 for CDR-H3, wherein the residues are numbered according to MacCallum;

(d) residues 24-34 of SEQ ID NO: 192 for CDR-L1, residues 50-56 of SEQ ID NO: 192 for CDR-L2, residues 89-97 of SEQ ID NO: 192 for CDR-L3, residues 31-35 of SEQ ID NO: 191 for CDR-H1, residues 50-65 of SEQ ID NO: 191 for CDR-H2 and residues 95-102 of SEQ ID NO: 191 for CDR-H3, wherein the residues are numbered according to Kabat;

(e) residues 23-34 of SEQ ID NO: 192 for CDR-L1, residues 50-56 of SEQ ID NO: 192 for CDR-L2, residues 89-97 of SEQ ID NO: 192 for CDR-L3, residues 26-32 of SEQ ID NO: 191 for CDR-H1, residues 50-58 of SEQ ID NO: 191 for CDR-H2 and residues 95-102 of SEQ ID NO: 191 for CDR-H3, wherein the residues are numbered according to Chothia;

(f) residues 30-36 of SEQ ID NO: 192 for CDR-L1, residues 46-55 of SEQ ID NO: 192 for CDR-L2, residues 89-96 of SEQ ID NO: 192 for CDR-L3, residues 30-35 of SEQ ID NO: 191 for CDR-H1, residues 47-58 of SEQ ID NO: 191 for CDR-H2 and residues 93-101 of SEQ ID NO: 191 for CDR-H3, wherein the residues are numbered according to MacCallum;

(g) residues 24-34 of SEQ ID NO: 172 for CDR-L1, residues 50-56 of SEQ ID NO: 172 for CDR-L2, residues 89-97 of SEQ ID NO: 172 for CDR-L3, residues 31-35 of SEQ ID NO: 173 for CDR-H1, residues 50-65 of SEQ ID NO: 173 for CDR-H2 and residues 95-102 of SEQ ID NO: 173 for CDR-H3, wherein the residues are numbered according to Kabat;

(h) residues 23-34 of SEQ ID NO: 172 for CDR-L1, residues 50-56 of SEQ ID NO: 172 for CDR-L2, residues 89-97 of SEQ ID NO: 172 for CDR-L3, residues 26-32 of SEQ ID NO: 173 for CDR-H1, residues 50-58 of SEQ ID NO: 173 for CDR-H2 and residues 95-102 of SEQ ID NO: 173 for CDR-H3, wherein the residues are numbered according to Chothia; or

(i) residues 30-36 of SEQ ID NO: 172 for CDR-L1, residues 46-55 of SEQ ID NO: 172 for CDR-L2, residues 89-96 of SEQ ID NO: 172 for CDR-L3, residues 30-35 of SEQ ID NO: 173 for CDR-H1, residues 47-58 of SEQ ID NO: 173 for CDR-H2 and residues 93-101 of SEQ ID NO: 173 for CDR-H3, wherein the residues are numbered according to MacCallum.

10. The pharmaceutical composition for use according to any one of claims 1 or 4-9, the kit according to any one of claims 2 or 4-9, or the method of any one of claims 3-9, wherein the antibody comprises:

(i) a light chain variable region set forth as SEQ ID NO: 190 or SEQ ID NO: 192, and a heavy chain variable region set forth as SEQ ID NO: 191 or

(ii) a light chain variable region set forth as SEQ ID NO: 172 and a heavy chain variable region set forth as SEQ ID NO: 173.

11. The pharmaceutical composition for use according to any one of claims 1 or 4-10 or the kit of any one of claims 2 or 4-10, wherein the antibody is conjugated to the cytotoxic agent via a linker.

12. The pharmaceutical composition for use according to claim 11 or the kit according to claim 11, wherein the linker comprises a non-cleavable linker.

13. The pharmaceutical composition for use according to any one of claims 1 or 4-12 or the kit according to any one of claims 2 or 4-12, wherein the cytotoxic agent comprises a duocarmycin, a calicheamicin, an auristatin, an amanitin, a pyrrolobenzodiazepine, or a radioisotope.

14. The pharmaceutical composition for use according to any one of claims 1 or 4-13 or the kit according to any one of claims 2 or 4-13, wherein the cytotoxic agent comprises a calicheamicin.

15. The method according to any one of claims 3 or 5-10, wherein the antibody is detected using immunohistochemistry.

16. The pharmaceutical composition for use according to any one of claims 1 or 4-14, wherein the cytotoxic agent comprises a calicheamicin and wherein the antibody comprises three complementarity determining regions of a light chain variable region of SEQ ID NO: 190 or SEQ ID NO: 192, and three complementarity determining regions of a heavy chain variable region of SEQ ID NO: 191.

5

17. The kit according to any one of claims 2 or 4-14, wherein the cytotoxic agent comprises a calicheamicin and wherein the antibody comprises three complementarity determining regions of a light chain variable region of SEQ ID NO: 190 or SEQ ID NO: 192, and three complementarity determining regions of a heavy chain variable region of SEQ ID NO: 191.

10

15

20

25

30

35

40

45

50

55

Homo sapiens seizure related 6 homolog (SEZ6), transcript variant 1, mRNA
 >gi|148839279|ref|NM_178860.4|

(SEQ ID NO: 1)

GATCCCCGGCGCCGTGCGCAGGCGCTGGCCGTGGTGTCTGATCTGTCTAGGCGCTGGCGGGCGCAGCGCGGTGACGGCTGCGG
 CCCCCTCCCTCTACCCGGCCGGACCCGGCTCTGCCCCGCGCCCAAGCCCCACCAAGCCCCCGCCCTCCCGCCGCGGTCCC
 AGCCAGGGCGCGCCGCAACCAGCACCATGCGCCCGGTAGCCCTGCTGCTCCTGCCCTCGCTGCTGGCGCTCCTGGCTCACG
 GACTCTCTTTAGAGGCCCCAACCCTGGGGAAAGGACAAGCCCCAGGCATCGAGGAGACAGATGGCGAGCTGACAGCAGCCCCC
 ACACCTGAGCAGCCAGAACGAGGCGTCCACTTTGTCAACAACAGCCCCACCTTGAAGCTGCTCAACCACCACCCGCTGCTTGA
 GGAATTCCTACAAGAGGGGCTGGAAAAGGGAGATGAGGAGCTGAGGCCAGCACTGCCCTTCCAGCCTGACCCACCTGCACCCT
 TCACCCCAAGTCCCCTTCCCCGCTGGCCAACCAGGACAGCCGCTGTCTTTACCAGCCCCACTCCAGCCATGGCTGCGGTA
 CCCACTCAGCCCCAGTCCAAGGAGGACCCTGGAGTCCGGAGTCAGAGTCCCTATGCTTCGAATCACAGCTCCCTACCTCC
 AGGGCCAGCATGGCAGTGCCACCCCTAGGCCAGGGGAGATAGCCAGCACTACACCCCCAGCAGAGCCTGGACACCAACCC
 AAGAGGGTCTTGAGACATGGGAAGGCCGTGGCTTCAGAGGTTGTGTCCAGGGCGCAGGGATCGGGATCCAGGGGACCATC
 ACCTCCTCCACAGCTTCAGGAGATGATGAGGAGACCACCCTACCACCACCATCATCACCACCACCATCACCACAGTCCAGAC
 ACCAGGCCCTTGTAGCTGGAATTTCTCAGGCCAGAGGGCTCTGTGACTCCCTACAGACCTCAGCTCCCCACTGATGTG
 GCCTGGACTGCTTCTTCTACATCTCTGTCTACCTGGCTATGGCGTGAAATCAAGGTCCAGAATATCAGCCTCCGGGAAGGG
 GAGACAGTGAATGTGAAGGCCCTGGGGGGGCTGACCCACTGCCCTGGCCAACCAGTCTTCTGCTGCGGGGCCAAGTCAT
 CCGCAGCCCCACCAAGCGCCCTGAGGTTCCAGAGCCTCCCGCCACGGCTGGCCCTGGCACCTTCCATTTCCATTACC
 AAGCCTATCTCCTGAGCTGCCACTTCCCCGCTCCAGCTTATGGAGATGTGACTGTACACAGCCTCCACCCAGGGGGTAGT
 GCCGCTTCCATTGTGCCACTGGCTACCAGCTGAAGGGCGCCAGGCATCTCACTGTCTCAATGCCACCCAGCCCTTCTGGGA
 TTCAAAGGAGCCCGTCTGCATCGCTGCTTGCAGCGAGTGATCCGCAATGCCACCACCGGCCGCATCGTCTCTCCAGGCTTCC
 CGGGCACTACAGCAACAACCTCACCTGTCACTGGCTGCTTGAAGCTCCTGAGGGCCAGCGGTACACCTGCACTTTGAGAAG
 GTTTCCTTGGCAGAGGATGATGACAGGCTCATCTTCGCAATGGGGACAACGTGGAGGCCCCACCAAGTGTATGATTCTATGA
 GGTGGAATACCTGCCCATTGAGGGCCTGCTCAGCTCTGGCAACACTTCTTTGTTGAGCTCAGTACTGACAGCAGCGGGGCGAG
 CTGCAGGCATGGCCCTGCGCTATGAGGCCTTCCAGCAGGGCCATGTCTATGAGCCCTTTGTCAAATACGGTAACTTCAGCAGC
 AGCACACCCACCTACCCTGTGGGTACCACTGTGGAGTTCAGCTGCCACCCCTGGCTACACCTGGAGCAGGGCTCCATCATCAT
 CGAGTGTGTTGACCCCCACGACCCCAAGTGGAAATGAGACAGAGCCAGCCTGCCGAGCCGTGTGCAGCGGGGAGATCACAGCT
 CGGCTGGCGTGGTACTCTCTCCCACTGGCCAGAGCCCTACGGTCTGGGCAGGATTGTATCTGGGGTGTGCATGTGGAAGAG
 GACAAGCGCATCATGTGGACATCCGAGTGTGCGCATAGGCCCTGGTGATGTGCTTACCTTCTATGATGGGGATGACCTGAC
 GGCCCGGGTCTTGGGCCAGTACTCAGGGCCCCGTAGCCACTTCAAGCTCTTTACCTCCATGGCTGATGTCAACATTAGTTC
 AGTCGAGCCCCGGGACCTCAGTGTGGGCTACCAAGCAGGGCTTCGTCTATCCACTTCTTTGAGGTGCCCGCAATGACACATGT
 CCGGAGCTGCCTGAGATCCCCAATGGCTGGAAGAGCCCATCGCAGCCTGAGCTAGTGCACGGCACCCGTGGTCACTTACCAGTG
 CTACCTTGGCTACCAAGTGTGGGATCCAGTGTCTCATGTGCCAGTGGGACCTAACTTGGAGTGAGGACCTGCCCTCATGCC
 AGAGGTGACTTCCGACGATCCTGGAGATGTGGAGACAGCCAGCCTCATATCCAGCCCCAGTTTCCCGTGGGGGCC
 ACCGTGCAATATATCTGTGACCAGGGTTTTGTGCTGATGGGCGAGCTCCATCTCACCTGCCATGATCGCCAGGCTGGCAGCCC
 CAAGTGGAGTGACCGGGCCCCCTAAATGTCTCTTGAACAGCTCAAGCCATGCCATGGTCTCAGTGGCCCTGAGAATGGTGGCC
 GAAGTCTTGAGAAGCAGCTACACCCAGCAGGGGCCACCATCCACTTCTCGTGTGCCCTGGCTATGTGCTGAAGGGCCAGGCC
 AGCATCAAGTGTGTGCTGGGCACCCCTCGCATTGGAGTGACCCCCACCCATCTGTAGGGCTGCCTCTCTGGATGGGTCTTA
 CAACAGTTCGAGCCTGGATGTTGCCAAGGCACCTGCTGCCCTCCAGCACCCCTGGATGCTGCCACATTGACAGCTGCCATCTTCT
 TGCCACTGGTGGCGATGGTGTGTTGGTAGGAGGTGATACCTTCTACTTCTCCAGGCTCCAGGGAAAAAGCTCCCTGCAGCTG
 CCCCCCCCCGCCCCGCCCTACAACCGCATTACCATAGAGTCAGCGTTTGACAATCCAATACGAGACTGGATCTCTTTC
 CTTTGCAGGAGACGAGAGAATATGAAGTCTCCATCTAGGTGGGGGCGAGCTAGGGAAGTCAACTCAGACTTGACACACAGTCC
 AGCAGCAAGGCTCCTTGTCTCCTGTGCTCCCTCCACCTCCTGTATATACCACCTAGGAGGAGATGCCACCAAGCCCTCAAGAA
 GTTGTGCCCTTCCCCGCTGCGATGCCACCATGGCCTATTTTCTTGGTGTCTATGCCCACCTGGGGCCCTTCATTGGGCCCCA
 TGTAGGGGGCATCTACCTGTGGGAAGAACATAGCTGGAGCACAAGCATCAACAGCCAGCATCCTGAGCCTCCTCATGCCCTG
 GACCAGCCTGGAACACACTAGCAGAGCAGGAGTACCTTTCTCCACATGACCACCATCCCGCCCTGGCATGGCAACCTGCAGCA
 GGATTAACCTTGACCATGGTGGGAAGTGCACCAGGCTACTCCTCACAGCGCATACCAATGGCCAAAACCTCCTCTCAACGGTGA
 CCTCTGGGTAGTCTTGGCATGCCAACATCAGCCTCTTGGGAGGTCTCTAGTTCTCTAAAGTTCTGGACAGTTCTGCCTCCTGC
 CCTGTCCAGTGGAGGCGATAATTTCTAGGAGATCCTAAGGGGTTTCAGGGGGACCTACCCACCTCAGGTTGGGCTTCCCTG
 GGCATCATGCTCCACACCAAGCAGGACACGCCATTTTCCACTGACACCCCTATACCTGAGGAAAGGGAGACTTTCTCTCCG
 ATGTTTATTTAGCTGTGCAACATCTTACCCTAAATAGTCCCTCCTCCAATTCAGCCACTTGTGAGGCTCTCCTCTTGACC
 ACTGTGTATGGGATAAGGGGAGGGGGTGGGCATATTCTGGAGAGGAGCAGAGGTCCAAGGACCCAGGAATTTGGCATGGAAC
 AGGTGGTAGGAGAGCCCCAGGGAGACGCCAGGAGCTGGCTGAAAGCCACTTTGTACATGTAATGTATTATATGGGGTCTGGG
 CTCCAGCCAGAGAACAATCTTTTATTTCTGTTGTTTCTTATTAATAATGGTGTTTTTTGAAAAA
 AAAAAAAAAAAAAAAAAA

FIG. 1A

Homo sapiens seizure related 6 homolog (SEZ6), transcript variant 2,
mRNA>gi|148839345|ref|NM_001098635.1|

(SEQ ID NO: 2)

GATCCCCGGCGCCGTCGCCAGGCGCTGGCCGTGGTGCTGATTCTGTTCAGGCGCTGGCGGCGGCAGCGCGGTGACGGCTGCGG
CCCCGCTCCCTCTACCCGGCCGACCCGGCTTGCCCCCGGCCCAAGCCCCACCAAGCCCCCGCCCTCCCGCCCGGGTCCC
AGCCAGGGCGCGCCGCAACCAGCACCATGCGCCCGGTAGCCCTGCTGCTCCTGCCCTCGCTGCTGGCGCTCCTGGCTCACG
GACTCTCTTTAGAGGCCCCAACCCTGGGGAAAGGACAAGCCCCAGGCATCGAGGAGACAGATGGCGAGCTGACAGCAGCCCCC
ACACCTGAGCAGCCAGAACGAGGCGTCCACTTTGTACAAACAGCCCCACCTTGAAGCTGCTCAACCACCACCCGCTGCTTGA
GGAATTCCTACAAGAGGGGCTGGAAAAGGGAGATGAGGAGCTGAGGCCAGCACTGCCCTTCCAGCCTGACCCACCTGCACCCCT
TCACCCCAAGTCCCTTCCCCGCTGGCCAACCAGGACAGCCGCCCTGTCTTTACCAGCCCCACTCCAGCCATGGCTGCGGTA
CCCACTCAGCCCCAGTCCAAGGAGGGACCTGGAGTCCGGAGTCAGAGTCCCCTATGCTTCGAATCACAGCTCCCTACCTCC
AGGGCCAGCATGGCAGTGCCCAACCTAGGCCCAGGGGAGATAGCCAGCACTACACCCCCAGCAGAGCCTGGACACCAACCC
AAGAGGGTCTGGAGACATGGGAAGGCCGTGGGTGTCAGAGGTTGTGTCCCAGGGCGCAGGGATCGGGATCCAGGGGACCATC
ACCTCCTCCACAGCTTCAGGAGATGATGAGGAGACCACCACTACCACCACCATCATCACCACCACCATCACCACAGTCCAGAC
ACCAGGCCCTTGTAGCTGGAATTTCTCAGGCCAGAGGGCTCTCTGGACTCCCCTACAGACCTCAGCTCCCCACTGATGTTG
GCCTGGACTGCTTCTCTACATCTCTGTCTACCCTGGCTATGGCGTGGAAATCAAGGTCCAGAATATCAGCCTCCGGGAAGGG
GAGACAGTGAAGGCTGGGGGGGCTGACCCACTGCCCCCTGGCCAACCAGTCTTTCCTGCTGCGGGGCCAAGTCAT
CCGCAGCCCCACCAAGCGGCCCTGAGGTTCCAGAGCCTCCGCCACCGGCTGGCCCTGGCACCTTCCATTTCCATTACC
AAGCTATCTCTGAGCTGCCACTTTCCCGTCGTCAGCTTATGGAGATGTGACTGTACAGGCTCCACCCAGGGGGTAGT
GCCCGCTTCCATTGCTGACTGCTACCAGCTGAAGGGCGCCAGGCATCTCACCTGTCTCAATGCCACCCAGCCCTTCTGGGA
TTCAAAGGAGCCGCTGTCATCGCTGCTTGGCGCGAGTGATCCGCAATGCCACCACCGGCCGATCGTCTCTCCAGGCTTCC
CGGGCAACTACAGCAACAACCTCACCTGTCACTGGCTGCTTGAGGCTCCTGAGGGCCAGCGGCTACACCTGCACCTTTGAGAAG
GTTTTCCCTGGCAGAGGATGATGACAGGCTCATATTGCAATGGGGACAACGTGGAGGCCCCACAGTGTATGATTCCTATGA
GGTGGAAATACCTGCCCATTGAGGGCCTGCTCAGCTCTGGCAAAACACTTCTTTGTTGAGCTCAGTACTGACAGCAGCGGGCAG
CTGCAGGCATGGCCCTGCGCTATGAGGCCTTCCAGCAGGGCCATTGCTATGAGCCCTTTGTCAAATACGGTAACCTCAGCAGC
AGCACACCCACCTACCCTGTGGGTACCCTGTGGAGTTCAGCTGCGACCTGGCTACACCTGGAGCAGGGCTCCATCATCAT
CGAGTGTGTTGACCCCCACGACCCCAAGTGGAAATGAGACAGAGCCAGCCGCTGCCGAGCCGTGTGACAGCGGGAGATCACAGACT
CGGCTGGCGTGGTACTCTCTCCCAACTGGCCAGAGCCCTACGGTCTGGGCGAGGATTGTATCTGGGGTGTGCATGTGGAAGAG
GACAAGCGCATCATGCTGGACATCCGAGTGTGCGCATAGGCCCTGGTGTGCTTACCTTCTATGATGGGGATGACCTGAC
GGCCCGGTTCTGGGCCAGTACTCAGGGCCCCGTAGCCACTCAAGCTCTTACCTCCATGGCTGATGTCACCATTCAGTTCC
AGTCCGACCCCGGACCTCAGTGTGGGCTACCAGCAGGGCTTCGTATCCACTTCTTTGAGGTGCCCCGCAATGACACATGT
CCGGAGCTGCCTGAGATCCCCAATGGCTGGAAGAGCCCATCGCAGCCTGAGCTAGTGACGGCACCCTGGTCACTTACCAGTG
CTACCTGGCTACCAGGTAGTGGGATCCAGTGTCTCATGTGCCAGTGGGACCTAAGTTGGAGTGAGGACCTGCCCTCATGCC
AGAAATATGAAGTCTCCATCTAGGTGGGGGCGAGTCTAGGGAAGTCAACTCAGACTTGCACCACAGTCCAGCAGCAAGGCTCCTT
GCTTCTGCTGTCCCTCCACCTCCTGTATATACCACCTAGGAGGAGATGCCACCAAGCCCTCAAGAAGTTGTGCCCTTCCCCG
CCTGCGATGCCACCATTGGCCTATTTTCTTGGTGTGCTTGGCCACTTGGGGCCCTTCATTGGGGCCATGTGAGGGGCATCTA
CCTGTGGGAAGAACATAGCTGGAGCACAAGCATCAACAGCCAGCATCCTGAGCCTCCTCATGCCCTGGACCAGCCTGGAACAC
ACTAGCAGAGCAGGAGTACCTTTCTCCACATGACCACCATCCCGCCCTGGCATGGCAACCTGCAGCAGGATTAACCTTGACCAT
GGTGGGAAGTGCACCAGGGTACTCCTCACAGCGCCATCACAATGGCCAAAACCTCCTCTCAACGGTGACCTCTGGGTAGTCCT
GGCATGCCAACATCAGCCTCTTGGGAGGTCTCTAGTTCCTAAAGTTCTGGACAGTTCTGCCTCCTGCCCTGTCCAGTGGAG
GCAGTAATTCTAGGAGATCCTAAGGGGTTTCAGGGGGACCTTACCCACCTCAGGTTGGGCTTCCCTGGGCACTCATGCTCCA
CACCAGCAGGACAGCCATTTTCCACTGACCACCTATACCCTGAGGAAAGGAGACTTTCTCCGATGTTTTATTATAGCTG
TTGCAAAACATCTTACCCTAATAGTCCCTCCTCAATTCAGGCCACTTGTGAGGCTCTCCTCTTGACCACTGTGTTATGGGAT
AAGGGGAGGGGTGGGCATATTCTGGAGAGGAGCAGAGGTCCAAGGACCCAGGAATTTGGCATGGAACAGGTGGTAGGAGAGC
CCCAGGGAGACGCCAGGAGCTGGCTGAAAGCCACTTTGTACATGTAATGTATTATATGGGGTCTGGGCTCCAGCCAGAGAAC
AATCTTTTATTTCTGTGTTTCTTATTAATAGGTGTTTTTGGAAAAA

FIG. 1B

Homo sapiens seizure protein 6 homolog isoform 1 precursor
>gi|148839280|ref|NP_849191.3|

(SEQ ID NO: 3)

MRPVALLLPSLLAHGHLSLEAPTVGKGQAPGIEETDGELTAAPTPEQPERGVHFVTTAPTTLKLLNHH
 PLLEEFLEQEGLEKDEELRPALFPQDPAPFTPSPLRLANQDSRPVFTSPTPAMAAVPTQPQSKEGP
 WSPSESPMLRITAPLPPGPSMAVPTLGPGEIASTTPPSRAWTPQTQEGPGDMGRPWWAEVVSQGAGI
 GIQGTITSTASGDDEETTTTITTTITTITVQTPGPCSWNFSGPEGSLDPTDLSSPTDVGLDCFFYISVYP
 GYGVEIKVQNISLREGETVTV EGLGGPDLPLANQSFLLRGQVIRSPTHQAALRFQSLPPPAGPGTFFHFH
 YQAYLLSCHFPRRPAYGDVTVTSLHPGGSARFHCATGYQLKGARHLTCLNATQPFWDSKEPVCIACGG
 VIRNATTGRIVSPGFPGNYSNNLTCHWLLLEAPEGQRLHLHFEKVSLEAEDDDRLIIRNGDNVEAPPVYDSY
 EVEYLP IEGLLSGKHFFVELSTDSSGAAAGMALRYEAFQQGHCHCEPFVKYGNFSSSTPTYPVGTTVEFS
 CDPGYTLEQGSIIIECVDPHPDQWNETEPACRAVCSGEITDSAGVVLSPNWPEPYGRGQDCIWGVHVE
 EDKRIMLDIRVLRIGPGDVLTFYDGGDDLTVLGGQYSGPRSHFKLFTSMADVTIQFQSDPGTSLVGYQQ
 GFVIHFFEVRNDTCPELPEIPNGWKSPSQPELVHGTVTYQCYPGYQVVGSSVLMCQWDLTWSEDL
 PSCQRVTSCHDPGDVEHSRRLISSPKFPVGATVQYICDQGGFVLMGSSILTCHDRQAGSPKWSDRAPKCL
 LEQLKPCHGLSAPENGARSPEKQLHPAGATIHFSAPGYVLKGQASIKCVPGHPSHWSDPPPICRAASL
 DGFYNSRSLDVAKAPAAASSTLDAAHIAAAIFLPLVAMVLLVGGVYFYFSRLQKGKSSLQLPRPRPRPNRITI
 ESAFDNPTYETGSLSFAGDERI

FIG. 1C

Homo sapiens seizure protein 6 homolog isoform 2 precursor

>gi|148839346|ref|NP_001092105.1|

(SEQ ID NO: 4)

MRPVALLLPSLALLAHGLSLEAPTVGKGQAPGIETDGELTAAPTEQPERGVHFVTTAPT
 LKLLNHHP¹LL²EEFLQEGLEK³GDEELR⁴PALFQ⁵PD⁶PPA⁷FT⁸PS⁹LP¹⁰RLANQ¹¹DS¹²RP¹³VFT¹⁴SPT¹⁵PAM¹⁶
 AAVPTQ¹⁷PQ¹⁸SK¹⁹EGP²⁰W²¹SE²²SP²³ML²⁴RI²⁵TAP²⁶LP²⁷PG²⁸PS²⁹MA³⁰VP³¹TL³²GP³³GE³⁴IA³⁵ST³⁶TP³⁷PS³⁸RA³⁹WT⁴⁰PT⁴¹Q⁴²EG⁴³
 PGDMGR⁴⁴PW⁴⁵VA⁴⁶EV⁴⁷VS⁴⁸QG⁴⁹AG⁵⁰IG⁵¹IG⁵²GT⁵³TS⁵⁴ST⁵⁵AS⁵⁶GD⁵⁷DE⁵⁸ET⁵⁹TTTT⁶⁰TTTT⁶¹TT⁶²TT⁶³TT⁶⁴TT⁶⁵TT⁶⁶TT⁶⁷TT⁶⁸TT⁶⁹TT⁷⁰TT⁷¹TT⁷²TT⁷³TT⁷⁴TT⁷⁵TT⁷⁶TT⁷⁷TT⁷⁸TT⁷⁹TT⁸⁰TT⁸¹TT⁸²TT⁸³TT⁸⁴TT⁸⁵TT⁸⁶TT⁸⁷TT⁸⁸TT⁸⁹TT⁹⁰TT⁹¹TT⁹²TT⁹³TT⁹⁴TT⁹⁵TT⁹⁶TT⁹⁷TT⁹⁸TT⁹⁹TT¹⁰⁰TT¹⁰¹TT¹⁰²TT¹⁰³TT¹⁰⁴TT¹⁰⁵TT¹⁰⁶TT¹⁰⁷TT¹⁰⁸TT¹⁰⁹TT¹¹⁰TT¹¹¹TT¹¹²TT¹¹³TT¹¹⁴TT¹¹⁵TT¹¹⁶TT¹¹⁷TT¹¹⁸TT¹¹⁹TT¹²⁰TT¹²¹TT¹²²TT¹²³TT¹²⁴TT¹²⁵TT¹²⁶TT¹²⁷TT¹²⁸TT¹²⁹TT¹³⁰TT¹³¹TT¹³²TT¹³³TT¹³⁴TT¹³⁵TT¹³⁶TT¹³⁷TT¹³⁸TT¹³⁹TT¹⁴⁰TT¹⁴¹TT¹⁴²TT¹⁴³TT¹⁴⁴TT¹⁴⁵TT¹⁴⁶TT¹⁴⁷TT¹⁴⁸TT¹⁴⁹TT¹⁵⁰TT¹⁵¹TT¹⁵²TT¹⁵³TT¹⁵⁴TT¹⁵⁵TT¹⁵⁶TT¹⁵⁷TT¹⁵⁸TT¹⁵⁹TT¹⁶⁰TT¹⁶¹TT¹⁶²TT¹⁶³TT¹⁶⁴TT¹⁶⁵TT¹⁶⁶TT¹⁶⁷TT¹⁶⁸TT¹⁶⁹TT¹⁷⁰TT¹⁷¹TT¹⁷²TT¹⁷³TT¹⁷⁴TT¹⁷⁵TT¹⁷⁶TT¹⁷⁷TT¹⁷⁸TT¹⁷⁹TT¹⁸⁰TT¹⁸¹TT¹⁸²TT¹⁸³TT¹⁸⁴TT¹⁸⁵TT¹⁸⁶TT¹⁸⁷TT¹⁸⁸TT¹⁸⁹TT¹⁹⁰TT¹⁹¹TT¹⁹²TT¹⁹³TT¹⁹⁴TT¹⁹⁵TT¹⁹⁶TT¹⁹⁷TT¹⁹⁸TT¹⁹⁹TT²⁰⁰TT²⁰¹TT²⁰²TT²⁰³TT²⁰⁴TT²⁰⁵TT²⁰⁶TT²⁰⁷TT²⁰⁸TT²⁰⁹TT²¹⁰TT²¹¹TT²¹²TT²¹³TT²¹⁴TT²¹⁵TT²¹⁶TT²¹⁷TT²¹⁸TT²¹⁹TT²²⁰TT²²¹TT²²²TT²²³TT²²⁴TT²²⁵TT²²⁶TT²²⁷TT²²⁸TT²²⁹TT²³⁰TT²³¹TT²³²TT²³³TT²³⁴TT²³⁵TT²³⁶TT²³⁷TT²³⁸TT²³⁹TT²⁴⁰TT²⁴¹TT²⁴²TT²⁴³TT²⁴⁴TT²⁴⁵TT²⁴⁶TT²⁴⁷TT²⁴⁸TT²⁴⁹TT²⁵⁰TT²⁵¹TT²⁵²TT²⁵³TT²⁵⁴TT²⁵⁵TT²⁵⁶TT²⁵⁷TT²⁵⁸TT²⁵⁹TT²⁶⁰TT²⁶¹TT²⁶²TT²⁶³TT²⁶⁴TT²⁶⁵TT²⁶⁶TT²⁶⁷TT²⁶⁸TT²⁶⁹TT²⁷⁰TT²⁷¹TT²⁷²TT²⁷³TT²⁷⁴TT²⁷⁵TT²⁷⁶TT²⁷⁷TT²⁷⁸TT²⁷⁹TT²⁸⁰TT²⁸¹TT²⁸²TT²⁸³TT²⁸⁴TT²⁸⁵TT²⁸⁶TT²⁸⁷TT²⁸⁸TT²⁸⁹TT²⁹⁰TT²⁹¹TT²⁹²TT²⁹³TT²⁹⁴TT²⁹⁵TT²⁹⁶TT²⁹⁷TT²⁹⁸TT²⁹⁹TT³⁰⁰TT³⁰¹TT³⁰²TT³⁰³TT³⁰⁴TT³⁰⁵TT³⁰⁶TT³⁰⁷TT³⁰⁸TT³⁰⁹TT³¹⁰TT³¹¹TT³¹²TT³¹³TT³¹⁴TT³¹⁵TT³¹⁶TT³¹⁷TT³¹⁸TT³¹⁹TT³²⁰TT³²¹TT³²²TT³²³TT³²⁴TT³²⁵TT³²⁶TT³²⁷TT³²⁸TT³²⁹TT³³⁰TT³³¹TT³³²TT³³³TT³³⁴TT³³⁵TT³³⁶TT³³⁷TT³³⁸TT³³⁹TT³⁴⁰TT³⁴¹TT³⁴²TT³⁴³TT³⁴⁴TT³⁴⁵TT³⁴⁶TT³⁴⁷TT³⁴⁸TT³⁴⁹TT³⁵⁰TT³⁵¹TT³⁵²TT³⁵³TT³⁵⁴TT³⁵⁵TT³⁵⁶TT³⁵⁷TT³⁵⁸TT³⁵⁹TT³⁶⁰TT³⁶¹TT³⁶²TT³⁶³TT³⁶⁴TT³⁶⁵TT³⁶⁶TT³⁶⁷TT³⁶⁸TT³⁶⁹TT³⁷⁰TT³⁷¹TT³⁷²TT³⁷³TT³⁷⁴TT³⁷⁵TT³⁷⁶TT³⁷⁷TT³⁷⁸TT³⁷⁹TT³⁸⁰TT³⁸¹TT³⁸²TT³⁸³TT³⁸⁴TT³⁸⁵TT³⁸⁶TT³⁸⁷TT³⁸⁸TT³⁸⁹TT³⁹⁰TT³⁹¹TT³⁹²TT³⁹³TT³⁹⁴TT³⁹⁵TT³⁹⁶TT³⁹⁷TT³⁹⁸TT³⁹⁹TT⁴⁰⁰TT⁴⁰¹TT⁴⁰²TT⁴⁰³TT⁴⁰⁴TT⁴⁰⁵TT⁴⁰⁶TT⁴⁰⁷TT⁴⁰⁸TT⁴⁰⁹TT⁴¹⁰TT⁴¹¹TT⁴¹²TT⁴¹³TT⁴¹⁴TT⁴¹⁵TT⁴¹⁶TT⁴¹⁷TT⁴¹⁸TT⁴¹⁹TT⁴²⁰TT⁴²¹TT⁴²²TT⁴²³TT⁴²⁴TT⁴²⁵TT⁴²⁶TT⁴²⁷TT⁴²⁸TT⁴²⁹TT⁴³⁰TT⁴³¹TT⁴³²TT⁴³³TT⁴³⁴TT⁴³⁵TT⁴³⁶TT⁴³⁷TT⁴³⁸TT⁴³⁹TT⁴⁴⁰TT⁴⁴¹TT⁴⁴²TT⁴⁴³TT⁴⁴⁴TT⁴⁴⁵TT⁴⁴⁶TT⁴⁴⁷TT⁴⁴⁸TT⁴⁴⁹TT⁴⁵⁰TT⁴⁵¹TT⁴⁵²TT⁴⁵³TT⁴⁵⁴TT⁴⁵⁵TT⁴⁵⁶TT⁴⁵⁷TT⁴⁵⁸TT⁴⁵⁹TT⁴⁶⁰TT⁴⁶¹TT⁴⁶²TT⁴⁶³TT⁴⁶⁴TT⁴⁶⁵TT⁴⁶⁶TT⁴⁶⁷TT⁴⁶⁸TT⁴⁶⁹TT⁴⁷⁰TT⁴⁷¹TT⁴⁷²TT⁴⁷³TT⁴⁷⁴TT⁴⁷⁵TT⁴⁷⁶TT⁴⁷⁷TT⁴⁷⁸TT⁴⁷⁹TT⁴⁸⁰TT⁴⁸¹TT⁴⁸²TT⁴⁸³TT⁴⁸⁴TT⁴⁸⁵TT⁴⁸⁶TT⁴⁸⁷TT⁴⁸⁸TT⁴⁸⁹TT⁴⁹⁰TT⁴⁹¹TT⁴⁹²TT⁴⁹³TT⁴⁹⁴TT⁴⁹⁵TT⁴⁹⁶TT⁴⁹⁷TT⁴⁹⁸TT⁴⁹⁹TT⁵⁰⁰TT⁵⁰¹TT⁵⁰²TT⁵⁰³TT⁵⁰⁴TT⁵⁰⁵TT⁵⁰⁶TT⁵⁰⁷TT⁵⁰⁸TT⁵⁰⁹TT⁵¹⁰TT⁵¹¹TT⁵¹²TT⁵¹³TT⁵¹⁴TT⁵¹⁵TT⁵¹⁶TT⁵¹⁷TT⁵¹⁸TT⁵¹⁹TT⁵²⁰TT⁵²¹TT⁵²²TT⁵²³TT⁵²⁴TT⁵²⁵TT⁵²⁶TT⁵²⁷TT⁵²⁸TT⁵²⁹TT⁵³⁰TT⁵³¹TT⁵³²TT⁵³³TT⁵³⁴TT⁵³⁵TT⁵³⁶TT⁵³⁷TT⁵³⁸TT⁵³⁹TT⁵⁴⁰TT⁵⁴¹TT⁵⁴²TT⁵⁴³TT⁵⁴⁴TT⁵⁴⁵TT⁵⁴⁶TT⁵⁴⁷TT⁵⁴⁸TT⁵⁴⁹TT⁵⁵⁰TT⁵⁵¹TT⁵⁵²TT⁵⁵³TT⁵⁵⁴TT⁵⁵⁵TT⁵⁵⁶TT⁵⁵⁷TT⁵⁵⁸TT⁵⁵⁹TT⁵⁶⁰TT⁵⁶¹TT⁵⁶²TT⁵⁶³TT⁵⁶⁴TT⁵⁶⁵TT⁵⁶⁶TT⁵⁶⁷TT⁵⁶⁸TT⁵⁶⁹TT⁵⁷⁰TT⁵⁷¹TT⁵⁷²TT⁵⁷³TT⁵⁷⁴TT⁵⁷⁵TT⁵⁷⁶TT⁵⁷⁷TT⁵⁷⁸TT⁵⁷⁹TT⁵⁸⁰TT⁵⁸¹TT⁵⁸²TT⁵⁸³TT⁵⁸⁴TT⁵⁸⁵TT⁵⁸⁶TT⁵⁸⁷TT⁵⁸⁸TT⁵⁸⁹TT⁵⁹⁰TT⁵⁹¹TT⁵⁹²TT⁵⁹³TT⁵⁹⁴TT⁵⁹⁵TT⁵⁹⁶TT⁵⁹⁷TT⁵⁹⁸TT⁵⁹⁹TT⁶⁰⁰TT⁶⁰¹TT⁶⁰²TT⁶⁰³TT⁶⁰⁴TT⁶⁰⁵TT⁶⁰⁶TT⁶⁰⁷TT⁶⁰⁸TT⁶⁰⁹TT⁶¹⁰TT⁶¹¹TT⁶¹²TT⁶¹³TT⁶¹⁴TT⁶¹⁵TT⁶¹⁶TT⁶¹⁷TT⁶¹⁸TT⁶¹⁹TT⁶²⁰TT⁶²¹TT⁶²²TT⁶²³TT⁶²⁴TT⁶²⁵TT⁶²⁶TT⁶²⁷TT⁶²⁸TT⁶²⁹TT⁶³⁰TT⁶³¹TT⁶³²TT⁶³³TT⁶³⁴TT⁶³⁵TT⁶³⁶TT⁶³⁷TT⁶³⁸TT⁶³⁹TT⁶⁴⁰TT⁶⁴¹TT⁶⁴²TT⁶⁴³TT⁶⁴⁴TT⁶⁴⁵TT⁶⁴⁶TT⁶⁴⁷TT⁶⁴⁸TT⁶⁴⁹TT⁶⁵⁰TT⁶⁵¹TT⁶⁵²TT⁶⁵³TT⁶⁵⁴TT⁶⁵⁵TT⁶⁵⁶TT⁶⁵⁷TT⁶⁵⁸TT⁶⁵⁹TT⁶⁶⁰TT⁶⁶¹TT⁶⁶²TT⁶⁶³TT⁶⁶⁴TT⁶⁶⁵TT⁶⁶⁶TT⁶⁶⁷TT⁶⁶⁸TT⁶⁶⁹TT⁶⁷⁰TT⁶⁷¹TT⁶⁷²TT⁶⁷³TT⁶⁷⁴TT⁶⁷⁵TT⁶⁷⁶TT⁶⁷⁷TT⁶⁷⁸TT⁶⁷⁹TT⁶⁸⁰TT⁶⁸¹TT⁶⁸²TT⁶⁸³TT⁶⁸⁴TT⁶⁸⁵TT⁶⁸⁶TT⁶⁸⁷TT⁶⁸⁸TT⁶⁸⁹TT⁶⁹⁰TT⁶⁹¹TT⁶⁹²TT⁶⁹³TT⁶⁹⁴TT⁶⁹⁵TT⁶⁹⁶TT⁶⁹⁷TT⁶⁹⁸TT⁶⁹⁹TT⁷⁰⁰TT⁷⁰¹TT⁷⁰²TT⁷⁰³TT⁷⁰⁴TT⁷⁰⁵TT⁷⁰⁶TT⁷⁰⁷TT⁷⁰⁸TT⁷⁰⁹TT⁷¹⁰TT⁷¹¹TT⁷¹²TT⁷¹³TT⁷¹⁴TT⁷¹⁵TT⁷¹⁶TT⁷¹⁷TT⁷¹⁸TT⁷¹⁹TT⁷²⁰TT⁷²¹TT⁷²²TT⁷²³TT⁷²⁴TT⁷²⁵TT⁷²⁶TT⁷²⁷TT⁷²⁸TT⁷²⁹TT⁷³⁰TT⁷³¹TT⁷³²TT⁷³³TT⁷³⁴TT⁷³⁵TT⁷³⁶TT⁷³⁷TT⁷³⁸TT⁷³⁹TT⁷⁴⁰TT⁷⁴¹TT⁷⁴²TT⁷⁴³TT⁷⁴⁴TT⁷⁴⁵TT⁷⁴⁶TT⁷⁴⁷TT⁷⁴⁸TT⁷⁴⁹TT⁷⁵⁰TT⁷⁵¹TT⁷⁵²TT⁷⁵³TT⁷⁵⁴TT⁷⁵⁵TT⁷⁵⁶TT⁷⁵⁷TT⁷⁵⁸TT⁷⁵⁹TT⁷⁶⁰TT⁷⁶¹TT⁷⁶²TT⁷⁶³TT⁷⁶⁴TT⁷⁶⁵TT⁷⁶⁶TT⁷⁶⁷TT⁷⁶⁸TT⁷⁶⁹TT⁷⁷⁰TT⁷⁷¹TT⁷⁷²TT⁷⁷³TT⁷⁷⁴TT⁷⁷⁵TT⁷⁷⁶TT⁷⁷⁷TT⁷⁷⁸TT⁷⁷⁹TT⁷⁸⁰TT⁷⁸¹TT⁷⁸²TT⁷⁸³TT⁷⁸⁴TT⁷⁸⁵TT⁷⁸⁶TT⁷⁸⁷TT⁷⁸⁸TT⁷⁸⁹TT⁷⁹⁰TT⁷⁹¹TT⁷⁹²TT⁷⁹³TT⁷⁹⁴TT⁷⁹⁵TT⁷⁹⁶TT⁷⁹⁷TT⁷⁹⁸TT⁷⁹⁹TT⁸⁰⁰TT⁸⁰¹TT⁸⁰²TT⁸⁰³TT⁸⁰⁴TT⁸⁰⁵TT⁸⁰⁶TT⁸⁰⁷TT⁸⁰⁸TT⁸⁰⁹TT⁸¹⁰TT⁸¹¹TT⁸¹²TT⁸¹³TT⁸¹⁴TT⁸¹⁵TT⁸¹⁶TT⁸¹⁷TT⁸¹⁸TT⁸¹⁹TT⁸²⁰TT⁸²¹TT⁸²²TT⁸²³TT⁸²⁴TT⁸²⁵TT⁸²⁶TT⁸²⁷TT⁸²⁸TT⁸²⁹TT⁸³⁰TT⁸³¹TT⁸³²TT⁸³³TT⁸³⁴TT⁸³⁵TT⁸³⁶TT⁸³⁷TT⁸³⁸TT⁸³⁹TT⁸⁴⁰TT⁸⁴¹TT⁸⁴²TT⁸⁴³TT⁸⁴⁴TT⁸⁴⁵TT⁸⁴⁶TT⁸⁴⁷TT⁸⁴⁸TT⁸⁴⁹TT⁸⁵⁰TT⁸⁵¹TT⁸⁵²TT⁸⁵³TT⁸⁵⁴TT⁸⁵⁵TT⁸⁵⁶TT⁸⁵⁷TT⁸⁵⁸TT⁸⁵⁹TT⁸⁶⁰TT⁸⁶¹TT⁸⁶²TT⁸⁶³TT⁸⁶⁴TT⁸⁶⁵TT⁸⁶⁶TT⁸⁶⁷TT⁸⁶⁸TT⁸⁶⁹TT⁸⁷⁰TT⁸⁷¹TT⁸⁷²TT⁸⁷³TT⁸⁷⁴TT⁸⁷⁵TT⁸⁷⁶TT⁸⁷⁷TT⁸⁷⁸TT⁸⁷⁹TT⁸⁸⁰TT⁸⁸¹TT⁸⁸²TT⁸⁸³TT⁸⁸⁴TT⁸⁸⁵TT⁸⁸⁶TT⁸⁸⁷TT⁸⁸⁸TT⁸⁸⁹TT⁸⁹⁰TT⁸⁹¹TT⁸⁹²TT⁸⁹³TT⁸⁹⁴TT⁸⁹⁵TT⁸⁹⁶TT⁸⁹⁷TT⁸⁹⁸TT⁸⁹⁹TT⁹⁰⁰TT⁹⁰¹TT⁹⁰²TT⁹⁰³TT⁹⁰⁴TT⁹⁰⁵TT⁹⁰⁶TT⁹⁰⁷TT⁹⁰⁸TT⁹⁰⁹TT⁹¹⁰TT⁹¹¹TT⁹¹²TT⁹¹³TT⁹¹⁴TT⁹¹⁵TT⁹¹⁶TT⁹¹⁷TT⁹¹⁸TT⁹¹⁹TT⁹²⁰TT⁹²¹TT⁹²²TT⁹²³TT⁹²⁴TT⁹²⁵TT⁹²⁶TT⁹²⁷TT⁹²⁸TT⁹²⁹TT⁹³⁰TT⁹³¹TT⁹³²TT⁹³³TT⁹³⁴TT⁹³⁵TT⁹³⁶TT⁹³⁷TT⁹³⁸TT⁹³⁹TT⁹⁴⁰TT⁹⁴¹TT⁹⁴²TT⁹⁴³TT⁹⁴⁴TT⁹⁴⁵TT⁹⁴⁶TT⁹⁴⁷TT⁹⁴⁸TT⁹⁴⁹TT⁹⁵⁰TT⁹⁵¹TT⁹⁵²TT⁹⁵³TT⁹⁵⁴TT⁹⁵⁵TT⁹⁵⁶TT⁹⁵⁷TT⁹⁵⁸TT⁹⁵⁹TT⁹⁶⁰TT⁹⁶¹TT⁹⁶²TT⁹⁶³TT⁹⁶⁴TT⁹⁶⁵TT⁹⁶⁶TT⁹⁶⁷TT⁹⁶⁸TT⁹⁶⁹TT⁹⁷⁰TT⁹⁷¹TT⁹⁷²TT⁹⁷³TT⁹⁷⁴TT⁹⁷⁵TT⁹⁷⁶TT⁹⁷⁷TT⁹⁷⁸TT⁹⁷⁹TT⁹⁸⁰TT⁹⁸¹TT⁹⁸²TT⁹⁸³TT⁹⁸⁴TT⁹⁸⁵TT⁹⁸⁶TT⁹⁸⁷TT⁹⁸⁸TT⁹⁸⁹TT⁹⁹⁰TT⁹⁹¹TT⁹⁹²TT⁹⁹³TT⁹⁹⁴TT⁹⁹⁵TT⁹⁹⁶TT⁹⁹⁷TT⁹⁹⁸TT⁹⁹⁹TT¹⁰⁰⁰TT¹⁰⁰¹TT¹⁰⁰²TT¹⁰⁰³TT¹⁰⁰⁴TT¹⁰⁰⁵TT¹⁰⁰⁶TT¹⁰⁰⁷TT¹⁰⁰⁸TT¹⁰⁰⁹TT¹⁰¹⁰TT¹⁰¹¹TT¹⁰¹²TT¹⁰¹³TT¹⁰¹⁴TT¹⁰¹⁵TT¹⁰¹⁶TT¹⁰¹⁷TT¹⁰¹⁸TT¹⁰¹⁹TT¹⁰²⁰TT¹⁰²¹TT¹⁰²²TT¹⁰²³TT¹⁰²⁴TT¹⁰²⁵TT¹⁰²⁶TT¹⁰²⁷TT¹⁰²⁸TT¹⁰²⁹TT¹⁰³⁰TT¹⁰³¹TT¹⁰³²TT¹⁰³³TT¹⁰³⁴TT¹⁰³⁵TT¹⁰³⁶TT¹⁰³⁷TT¹⁰³⁸TT¹⁰³⁹TT¹⁰⁴⁰TT¹⁰⁴¹TT¹⁰⁴²TT¹⁰⁴³TT¹⁰⁴⁴TT¹⁰⁴⁵TT¹⁰⁴⁶TT¹⁰⁴⁷TT¹⁰⁴⁸TT¹⁰⁴⁹TT¹⁰⁵⁰TT¹⁰⁵¹TT¹⁰⁵²TT¹⁰⁵³TT¹⁰⁵⁴TT¹⁰⁵⁵TT¹⁰⁵⁶TT¹⁰⁵⁷TT¹⁰⁵⁸TT¹⁰⁵⁹TT¹⁰⁶⁰TT¹⁰⁶¹TT¹⁰⁶²TT¹⁰⁶³TT¹⁰⁶⁴TT¹⁰⁶⁵TT¹⁰⁶⁶TT¹⁰⁶⁷TT¹⁰⁶⁸TT¹⁰⁶⁹TT¹⁰⁷⁰TT¹⁰⁷¹TT¹⁰⁷²TT¹⁰⁷³TT¹⁰⁷⁴TT¹⁰⁷⁵TT¹⁰⁷⁶TT¹⁰⁷⁷TT¹⁰⁷⁸TT¹⁰⁷⁹TT¹⁰⁸⁰TT¹⁰⁸¹TT¹⁰⁸²TT¹⁰⁸³TT¹⁰⁸⁴TT¹⁰⁸⁵TT¹⁰⁸⁶TT¹⁰⁸⁷TT¹⁰⁸⁸TT¹⁰⁸⁹TT¹⁰⁹⁰TT¹⁰⁹¹TT¹⁰⁹²TT¹⁰⁹³TT¹⁰⁹⁴TT¹⁰⁹⁵TT¹⁰⁹⁶TT¹⁰⁹⁷TT¹⁰⁹⁸TT¹⁰⁹⁹TT¹¹⁰⁰TT¹¹⁰¹TT¹¹⁰²TT¹¹⁰³TT¹¹⁰⁴TT¹¹⁰⁵TT¹¹⁰⁶TT¹¹⁰⁷TT¹¹⁰⁸TT¹¹⁰⁹TT¹¹¹⁰TT¹¹¹¹TT¹¹¹²TT¹¹¹³TT¹¹¹⁴TT¹¹¹⁵TT¹¹¹⁶TT¹¹¹⁷TT¹¹¹⁸TT¹¹¹⁹TT¹¹²⁰TT¹¹²¹TT¹¹²²TT¹¹²³TT¹¹²⁴TT¹¹²⁵TT¹¹²⁶TT¹¹²⁷TT¹¹²⁸TT¹¹²⁹TT¹¹³⁰TT¹¹³¹TT¹¹³²TT¹¹³³TT¹¹³⁴TT¹¹³⁵TT¹¹³⁶TT¹¹³⁷TT¹¹³⁸TT¹¹³⁹TT¹¹⁴⁰TT¹¹⁴¹TT¹¹⁴²TT¹¹⁴³TT¹¹⁴⁴TT¹¹⁴⁵TT¹¹⁴⁶TT¹¹⁴⁷TT¹¹⁴⁸TT¹¹⁴⁹TT¹¹⁵⁰TT¹¹⁵¹TT¹¹⁵²TT¹¹⁵³TT¹¹⁵⁴TT¹¹⁵⁵TT¹¹⁵⁶TT¹¹⁵⁷TT¹¹⁵⁸TT¹¹⁵⁹TT¹¹⁶⁰TT¹¹⁶¹TT¹¹⁶²TT¹¹⁶³TT¹¹⁶⁴TT¹¹⁶⁵TT¹¹⁶⁶TT¹¹⁶⁷TT¹¹⁶⁸TT¹¹⁶⁹TT¹¹⁷⁰TT¹¹⁷¹TT¹¹⁷²TT¹¹⁷³TT¹¹⁷⁴TT¹¹⁷⁵TT¹¹⁷⁶TT¹¹⁷⁷TT¹¹⁷⁸TT¹¹⁷⁹TT¹¹⁸⁰TT¹¹⁸¹TT¹¹⁸²TT¹¹⁸³TT¹¹⁸⁴TT¹¹⁸⁵TT¹¹⁸⁶TT¹¹⁸⁷TT¹¹⁸⁸TT¹¹⁸⁹TT¹¹⁹⁰TT¹¹⁹¹TT¹¹⁹²TT¹

Alignment of SEZ6 Isoforms

	1	80
hSEZ6v1 (NP_849191)	(1) MRPVALLLLPSLLALLAHLGLEAPTGVKQAPGIEETDGLTAAPTQPERGVHFTVTTAPTLKLNHHPLLEEFQEG	
hSEZ6v2 (NP_001092105)	(1) MRPVALLLLPSLLALLAHLGLEAPTGVKQAPGIEETDGLTAAPTQPERGVHFTVTTAPTLKLNHHPLLEEFQEG	
	81	160
hSEZ6v1 (NP_849191)	(81) LEKDEELRPALEFPQDPAPFTTSPPLRLANQDSRPVFTSPTPMAAAVPTQPSKEGWPSESESMRLITAPLPGPS	
hSEZ6v2 (NP_001092105)	(81) LEKDEELRPALEFPQDPAPFTTSPPLRLANQDSRPVFTSPTPMAAAVPTQPSKEGWPSESESMRLITAPLPGPS	
	161	240
hSEZ6v1 (NP_849191)	(161) MAVPTLGPGEIASTTPPSRAWTPTEGPGDMGRPWAEVVSQAGIGIQGTITSTASGDDEETITTTIIITTTITTVQT	
hSEZ6v2 (NP_001092105)	(161) MAVPTLGPGEIASTTPPSRAWTPTEGPGDMGRPWAEVVSQAGIGIQGTITSTASGDDEETITTTIIITTTITTVQT	
	241	320
hSEZ6v1 (NP_849191)	(241) PGPCSWNFSGPEGSLDPTDLSSPTDVGLDCFFYISVYPGYGVEIKVQNISLREGETVTVTEGLGGPDPPLPLANOSFLLRG	
hSEZ6v2 (NP_001092105)	(241) PGPCSWNFSGPEGSLDPTDLSSPTDVGLDCFFYISVYPGYGVEIKVQNISLREGETVTVTEGLGGPDPPLPLANOSFLLRG	
	321	400
hSEZ6v1 (NP_849191)	(321) QVIRSPTHQAALRFQSLPPPPAGPGTFFHFHYQAYLLSCHFPFRPPAYGDVTVTSLHPGGSARFHCATGYQLKGARHLTCINA	
hSEZ6v2 (NP_001092105)	(321) QVIRSPTHQAALRFQSLPPPPAGPGTFFHFHYQAYLLSCHFPFRPPAYGDVTVTSLHPGGSARFHCATGYQLKGARHLTCINA	
	401	480
hSEZ6v1 (NP_849191)	(401) TQPFWDSDKEPVCIAACGGVIRNATTGRIVSPGPPGNYSNNTLCHWLLLEAPEGQRLHLHFEKVSLEAEDDDRLLIIRNGDNVE	
hSEZ6v2 (NP_001092105)	(401) TQPFWDSDKEPVCIAACGGVIRNATTGRIVSPGPPGNYSNNTLCHWLLLEAPEGQRLHLHFEKVSLEAEDDDRLLIIRNGDNVE	
	481	560
hSEZ6v1 (NP_849191)	(481) APPVYDSYEVEYLPIEGLLSSGKHFFVELSTDSSGAAAGMALRYEAFQQGHCEYEPFVKYGNFSSSTPTYPVGTTFEFS	
hSEZ6v2 (NP_001092105)	(481) APPVYDSYEVEYLPIEGLLSSGKHFFVELSTDSSGAAAGMALRYEAFQQGHCEYEPFVKYGNFSSSTPTYPVGTTFEFS	
	561	640
hSEZ6v1 (NP_849191)	(561) PGYTLEQGSIIIECVDPHDHPQNNETEPACRAVCSGEITDSAGVLSPNWPEPYGRGQDCIHWGVHVEEDKRIMLDIRVLRI	
hSEZ6v2 (NP_001092105)	(561) PGYTLEQGSIIIECVDPHDHPQNNETEPACRAVCSGEITDSAGVLSPNWPEPYGRGQDCIHWGVHVEEDKRIMLDIRVLRI	
	641	720
hSEZ6v1 (NP_849191)	(641) GPGDVLTFYDGDLLTARVLGQYSGPRSHFKLFTSMADVTIQFOSDPGTSVLGYQQGFVIHFEFVPRNDTCPELPEIPNGW	
hSEZ6v2 (NP_001092105)	(641) GPGDVLTFYDGDLLTARVLGQYSGPRSHFKLFTSMADVTIQFOSDPGTSVLGYQQGFVIHFEFVPRNDTCPELPEIPNGW	
	721	800
hSEZ6v1 (NP_849191)	(721) KSPSQPELVHGTVVTYQCYPGYQVVGSSVLMCQWDLTWSEDLPCQRYTSCHPDGDVDEHSRRLISSPKFPVGATVQYICD	
hSEZ6v2 (NP_001092105)	(721) KSPSQPELVHGTVVTYQCYPGYQVVGSSVLMCQWDLTWSEDLPCQRYTSCHPDGDVDEHSRRLISSPKFPVGATVQYICD	
	801	880
hSEZ6v1 (NP_849191)	(801) QGFVLMGSSILTCHDRQAGSPKWSDRAPKCLLEQLKPCHLGSLAPENGARSPKQLHPAGATIHFSAPGYVLKGQASIKC	
hSEZ6v2 (NP_001092105)	(801) QGFVLMGSSILTCHDRQAGSPKWSDRAPKCLLEQLKPCHLGSLAPENGARSPKQLHPAGATIHFSAPGYVLKGQASIKC	
	881	960
hSEZ6v1 (NP_849191)	(881) VPGHPHSHWSDPPICRAASLDGFFNSRSLDVAKAPAAASSTLDAAHIAAAIFLPLVAMVLLVGGVYFYFSRLQKSSSLQLP	
hSEZ6v2 (NP_001092105)	(881) VPGHPHSHWSDPPICRAASLDGFFNSRSLDVAKAPAAASSTLDAAHIAAAIFLPLVAMVLLVGGVYFYFSRLQKSSSLQLP	
	961	994
hSEZ6v1 (NP_849191)	(961) RPRPRPNRITTESAFDNPTYETGSLSFAGDERI	
hSEZ6v2 (NP_001092105)	(961) RPRPRPNRITTESAFDNPTYETGSLSFAGDERI	

(SEQ ID NO: 3)
(SEQ ID NO: 4)

FIG. 1E

Schematic Representation of SEZ6 Protein Extracellular Domain




FIG. 1F

Percent Identity Between Full Length
Mature SEZ6 Proteins from Various Species

SEZ6	rhesus (XP_001110503)	Cynomolgus (herein)	mouse (NP_067261)	rat (NP_001099224)
Human 1 (NP_849191)		97.5%	90.5%	90.4%
Human 2 (NP_001092105)	91.7%			
rhesus (XP_001110503)		93.0%		
mouse (NP_067261)				96.7%

FIG. 2A

Homo Sapiens SEZ6, SEZ6L, and SEZ6L2
Sequence Accession Numbers - NCBI

	mRNA		protein
SEZ6	1	NM_178860	NP_849191
	2	NM_001098635	NP_001092105
SEZ6L	1	NM_021115	NP_066938
	2	NM_001184773	NP_001171702
	3	NM_001184774	NP_001171703
	4	NM_001184775	NP_001701704
	5	NM_001184776	NP_001171705
	6	NM_001184777	NP_001171706
SEZ6L2			
	1	NM_012410	NP_036542
	2	NM_201575	NP_963869
	3	NM_001114099	NP_001107571
	4	NM_001114100	NP_001107572
	5	NM_001243332	NP_001230261
	6	NM_001243333	NP_001230262

FIG. 2B

Percent Identity between Various
Human SEZ6, SEZ6L and SEZ6L2 Proteins

Complete protein	
Homo sapiens SEZ6v1 (NP_849191)	SEZ6Lv1 (NP_066938)
	SEZ6L2v5 (NP_001230261)
	42.5%
	42.9%
	41.1%

ECD	
Homo sapiens SEZ6v1 (NP_849191)	SEZ6Lv1 (NP_066938)
	SEZ6L2v5 (NP_001230261)
	42.4%
	43.2%
	40.5%

FIG. 2C

>cDNA Sequence of human SEZ6 ORF

(SEQ ID NO: 5)

CTGAGCTGGAGGCCCAACCGTGGGGAAGGACAAGCCCCAGGCATCGAGGAGACAGATGGCGAGCTGACAGCAGCCCCACACCTGAGCAGCCAGAAC
 GAGCGTCCACTTTGTACAACAGCCCCACCTTGAAGCTGCTCAACCAACACCCGCTGCTTGAAGAAATTCCTACAAGAGGGGCTGGAAGAGGAGATGAGG
 AGTTAGGCCAGCACTGCCCTTCAGCTGACCCACTGACCCCTTCAACCAAGTCCCTTCCCGCTGGCCAAACCAAGGACAGCGCCCTGTCTTTACCAGC
 CCACTCCAGCCATGGTGGGTACCCACTAGCCCCAGTCCAAGAGGAGACCTGGAGTCCGAGTCAGAGTCCCTATGCTTGAATCACAGTCCCCCTAC
 CTCAGGGCCAGCATGGAGTGGCACCCTAGGCCAGGGGAGATAGCCAGCACTACACCCCAAGAGCCTGGACACCAACCAAGAGGGTCTCTGGA
 GACATGGGAAGGCCGTGGTTGCAGAGTTGTGCCAGGGCGGGGATCGGGATCCAGGGACCATCACCTCTCCACAGCTTCAGGAGATGATGAGG
 AGACCACCACTACCACCATCATCACCAACCATCACACAGTCCAGACACCAAGGCCCTTGTAGCTGGAATTTCTCAGGCCCAAGAGGGCTCTCTGGACTCC
 CCTACAGACCTCAGCTCCCCCACTGATTTGGCCTGGACTGCTTCTTACATCTCTGTACCCCTGGCTATGGCTGGAATCAAGTCCAGAATATCAGCCT
 CCGGGAAGGGGAGACAGTGACTGTGAAGGCTGGGGGGCCGACCACTGCCCTGGCCAAACCAAGTCTTCTGCTGGGGGCCAAGTCATCCGCAGCC
 CCACCAACCAAGGGCCCTGAGGTTCCAGAGCCTCCCGCAACCGGCTGGCCCTGGCACCTTCCATTTCCATTACCAAGCCTATCTCCTGAGCTGCCACTTTCCCC
 GTCGTCAGCTTATGGAGATGTGACTGTACCAAGCTCCACCAAGGGGTAGTGGCCGCTTCCATTGTGCCACTGGCTACCACTGAAGGGCCGACGGCATCGT
 TCACCTGTCTCAATGCCACCCAGCCCTTCTGGATTCAAAGAGGCCGCTGCTGATCGCTGCTGGCGGAGTGATCCGCAATGCCACCAACCGCCGCGCATCGT
 CTCTCCAGGCTTCCCGGGCAACTACAGCAACAACCTCACCTGTCACTGGTCTTGAAGCTCTGAGGCCAGCGGCTACACCTGCACCTTTGAGAAGGTTTCC
 CTGGCAGAGGATGACAGGCTCATCTCGCAATGGGACAACGTGGAGGCCCAACAGTGATGATTCCTATGAGGTGGAATACCTGCCATTGAGGG
 CTTGCTCAGCTCTGGCAACACTTCTTTGTAGCTCAGTACTACAGCAGCGGGGAGCTGAGGCTGCGCTATGAGGCTTCCAGCAGGGGCCA
 TTGCTATGAGCCCTTTGTCAAATACGGTAACCTTCAGCAGCAGCACCACTACCTGTGGGTACCACTGTGGAGTTCAGCTGCGACCTGGCTACACCTG
 GAGCAGGGCTCCATCATCATCGAGTGTGTTGACCCCAAGACCCCAAGTGGAAATGAGACAGAGCCAGCTGCCGAGCCGTGTGCAGCGGGGAGATCACAGA
 CTCGGCTGGCGTGTACTCTCTCCCAACTGGCCAGAGCCCTACGGTCGTGGCAGGATTGTATCTGGGGTGTGCATGTGGAAGAGGACAAGCGCATCATGCT
 GGACATCCGAGTGTCTGCGCATAGGCCCTGGTGTATGCTTACCTTCTATGATGGGATGACCTGACGGCCCGGTTCTGGGCCAGTACTCAGGGCCCCCGTAG
 CCACCTCAAGCTCTTACCTCCATGGCTGATGTACCACTCAGTTCAGTCGGACCCCGGACCTCAGTGTGGGTACCAAGCAGGGCTTCGTCATCCACTTCT
 TTGAGGTGCCCCGCAATGACACATGTCCGGAGTGCCTGAGATCCCCAATGGCTGGAAGAGCCCATCGACCTGAGCTAGTGACGGCACCCGTGGTCACTT
 ACCAGTGTACCTGGTACCAGGTAGTGGATCCAGTGTCTCATGTGCCAGTGGGACCTAACTTGGAGTGAGGACCTGCCCTCATGCCAGGGTGACTT
 CCTGCCACGATCCTGGAGATGTGAGCACAGCCGACGCTCATATCCAGCCCCAAGTTCCTGGGGGCCACCGTCAATATATCTGTGACCAGGGTTTGT
 GCTGATGGGAGCTCCATCCTCACCTGCCATGATCGCCAGGCTGGCAGCCCCAAGTGGAGTGACCCGGCCCCTAATGTCTCTGGAACAGCTCAAGCCCATG
 CCATGGTCTCAGTGCCCTGAGAAATGTGCCGAAGTCTGAGAAGCAGCTACACCCAGGAGGCCACCTCACTTCTGTGTGCCCCCTGGCTATGTGCTG
 AAGGGCCAGGCCAGCATCAAGTGTGTGCTGGCAGCCCTCGCATGGAGTGACCCCACTCTGTAGGGTGCCTCTCTGGATGGGTTCTACAACAGT
 CGCAGCTGGATGTTGCCAAGGCACCTGCTGCCTCCAGCACCTGGATGCTGCCACATTGACGTGCCATCTTCTGCCACTGGTGGCGATGGTGTGTTGG
 TAGGAGGTGTATCTTACTTCTCAGGCTCCAGGGAAGAAAGCTCCCTGCAAGTGCCTCCCGCCCGCCCTACAACCGCATTACCATAGAGTCAGC
 GTTTGACAATCCAACCTTACGAGACTGGATCTCTTCTTTCAGGAGACGAGAGAATA

FIG. 3A

> Translation of human SEZ6 ORF

(SEQ ID NO: 6)

LSLEAPTVGKGQAPGIEETDGELTAAPTPEQPERGVHFVTTAPTLLNHHPLLEEFLEQEGLEKGD
ELRPALPFQDPDPAPFTPSPLRLANQDSRPVFTSPTPAMAAVPTQQSQKEGPWSPESPEMLRIT
APLPPGPSMAVPTLGPGEIASTTPSRAWPTQEGPGDMGRPWWAEVVSQAGAGIGIGTITSST
ASGDDEETTTTITTTITTTVQTPGPCSWNFSGPEGSLDSPTDLSSPTDVGLDCFFYISVYPGYGVE
IKVQNISLREGETVTVGLGGPDPLPLANQSFLLRGQVIRSPTHQAALRFQSLPPAGPGTFFHFY
QAYLLSCHFPRRPAYGDVTVTSLHPGGSARFHCATGYQLKGARHLTCLNATQPFWDSKEPVCIAA
CGGVIRNATTGRIVSPGFPNGYSNNLTCHWLLEAPEGQRLHLHFKEKVSLEDDDDRLIIRNGDNVE
APPVYDSYEVEYLPIEGLSSGKHFFVELSTDSSGAAAGMALRYEAFQQGHCHCYEPFVKYGNFSSST
PTYPVGTTVEFSCDPGYTLEQGSIIIECVDPHDPQWNETEPACRAVCSGEITDSAGVVLSPNWPEP
YGRGQDCIWGVHVEEDKRIMLDIRVLRIGPGDVLTFYDGGDLTARVLGQYSGPRSHFKLFTSMAD
VTIQFQSDPGTSVLGYQQGFVIHFFEVPRNDTCPELPEIPNGWKSPSQPELVHGTVTVTYQCYPGY
QVVGSSVLMCQWDLTWSEDLPSQCQRVTSCHPDGDVEHSRRLISSPKFPVGATVQYICDQGFVL
MGSSILTCHDRQAGSPKWSDRAPKCLLEQLKPKCHGLSAPENGARSPEKQLHPAGATIHFSCAPGY
VLKGQASIKCVPGHPSHWSDPPICRAASLDGFYNSRSLDVAKAPAAASSTLDAAHIAAIFLPLVA
MVLVGGVYFYFSRLQKGKSSLQLPRPRPNRITIESAFDNPTYETGSLSFAGDERI

FIG. 3B

Alignment of SEZ6 Variants

		1				80
hSEZ6 BC146292	(1)	MRPVALLPSLLALLAHGLSLEAPT	VGKQAPGIEETD	GELTAAPT	EQPERGVHFTTAPT	KLNNHPLLEEFLOEG
hSEZ6 NP_849191	(1)	MRPVALLPSLLALLAHGLSLEAPT	VGKQAPGIEETD	GELTAAPT	EQPERGVHFTTAPT	KLNNHPLLEEFLOEG
		81				160
hSEZ6 BC146292	(81)	LEKGEELRPALPFQDPDPAPFT	PSPLRLANQDSRPVFTS	PTPMAAAV	PTQPSKEGSP	SESESMRLRITAPLPGPS
hSEZ6 NP_849191	(81)	LEKGEELRPALPFQDPDPAPFT	PSPLRLANQDSRPVFTS	PTPMAAAV	PTQPSKEGSP	SESESMRLRITAPLPGPS
		161				240
hSEZ6 BC146292	(161)	MAVPTLGPGEIASTTPPSRAW	TTQEGPGDMGRPWAEVVSQAGIGI	QGTITSS	TASGDDEETTTT	ITTTITTVQT
hSEZ6 NP_849191	(161)	MAVPTLGPGEIASTTPPSRAW	TTQEGPGDMGRPWAEVVSQAGIGI	QGTITSS	TASGDDEETTTT	ITTTITTVQT
		241				320
hSEZ6 BC146292	(241)	PGPCSWNFSGPEGSLDSPTD	LSSTDVGLDCFFYISVYPGYVEIK	VQNISLREGET	VTVEGLGGDP	LPLANOSFLLRG
hSEZ6 NP_849191	(241)	PGPCSWNFSGPEGSLDSPTD	LSSTDVGLDCFFYISVYPGYVEIK	VQNISLREGET	VTVEGLGGDP	LPLANOSFLLRG
		321				400
hSEZ6 BC146292	(321)	QVIRSPTHQAALRFQSLPP	PAGPCTFHFYQAYLLSCHFP	RRPAYGDVTV	TSLHPGGSAR	FHCATGYQLKGARHLTCLNA
hSEZ6 NP_849191	(321)	QVIRSPTHQAALRFQSLPP	PAGPCTFHFYQAYLLSCHFP	RRPAYGDVTV	TSLHPGGSAR	FHCATGYQLKGARHLTCLNA
		401	** *			480
hSEZ6 BC146292	(401)	TQPFWDSEKPEVCIG	ECPGVIRNATTGRIVSP	FGPGNYNNLTCHW	LEAPEGQRLHLH	FEKVSLEAEDDDRLLIIRNGDNVE
hSEZ6 NP_849191	(401)	TQPFWDSEKPEVCIA	ACGGVIRNATTGRIVSP	FGPGNYNNLTCHW	LEAPEGQRLHLH	FEKVSLEAEDDDRLLIIRNGDNVE
		481				560
hSEZ6 BC146292	(481)	APPVYDSYEVEYLP	PIEGLSSGKHFFVEL	STDSSGAAAGMALRYEAF	QQGHCYEPFVKYGN	FSSTPTYPVGTTFEFS
hSEZ6 NP_849191	(481)	APPVYDSYEVEYLP	PIEGLSSGKHFFVEL	STDSSGAAAGMALRYEAF	QQGHCYEPFVKYGN	FSSTPTYPVGTTFEFS
		561				640
hSEZ6 BC146292	(561)	PGYTLEQGSIIIEC	VDPHDPQWNETEPACRAV	CSGEITDSAGV	VLSPNWPEPYGRG	QDCIIGVHVEEDKRIMLDIRVLR
hSEZ6 NP_849191	(561)	PGYTLEQGSIIIEC	VDPHDPQWNETEPACRAV	CSGEITDSAGV	VLSPNWPEPYGRG	QDCIIGVHVEEDKRIMLDIRVLR
		641				720
hSEZ6 BC146292	(641)	GGDVLTFYDGD	DLTARVLGQYSGPRSHFKL	FTSMADVT	IQFQSDPGTS	VLGYQQGFVIHFFVPRNDTCPELPEIPNGW
hSEZ6 NP_849191	(641)	GGDVLTFYDGD	DLTARVLGQYSGPRSHFKL	FTSMADVT	IQFQSDPGTS	VLGYQQGFVIHFFVPRNDTCPELPEIPNGW
		721				800
hSEZ6 BC146292	(721)	KSPSQPELVHGT	VTVTYQCYPGYQVVGSSV	LMCQWDLTWSEDL	PSQQRVT	SCHDPGDVEHSRRLISSPKFPVGATVQYICD
hSEZ6 NP_849191	(721)	KSPSQPELVHGT	VTVTYQCYPGYQVVGSSV	LMCQWDLTWSEDL	PSQQRVT	SCHDPGDVEHSRRLISSPKFPVGATVQYICD
		801				880
hSEZ6 BC146292	(801)	QGFVLMGSSIL	TCHDRQAGSPKWS	DRAPKCLLEQLK	PCHGLSAPENGARS	PEKQLHPAGATIHFSCAPGYVLKQASIKC
hSEZ6 NP_849191	(801)	QGFVLMGSSIL	TCHDRQAGSPKWS	DRAPKCLLEQLK	PCHGLSAPENGARS	PEKQLHPAGATIHFSCAPGYVLKQASIKC
		881				960
hSEZ6 BC146292	(881)	VPGHPSHWS	SDPPPICRAASLDGF	YNRSRLDVAKAPAA	SSTLDAAHIAAA	IIFLPLVANVLLVGGVYFYSRLQGKSSLQLP
hSEZ6 NP_849191	(881)	VPGHPSHWS	SDPPPICRAASLDGF	YNRSRLDVAKAPAA	SSTLDAAHIAAA	IIFLPLVANVLLVGGVYFYSRLQGKSSLQLP
		961				994
hSEZ6 BC146292	(961)	RRPRPRYNRIT	IESAFDNPTYETG	SLSFAGDERI		
hSEZ6 NP_849191	(961)	RRPRPRYNRIT	IESAFDNPTYETG	SLSFAGDERI		

(SEQ ID NO: 7)
(SEQ ID NO: 3)

FIG. 3C

>human SEZ6-Fc ORF

(SEQ ID NO: 8)

ATGGAGACAGACACTCCTGCTATGGGTACTGCTGCTCTGGGTTCCGGGTCACACTGGTGACGGGGCTGGATCCCTGAGCCTGGAGGGCCCCAACCGTGGGAAAGG
 ACAAGCCCAGGCATCGAGGAGACAGATGGGAGCTGACAGCAGCCCCCACACTGAGCAGCCAGAACGAGGGCTCCACTTTGTACAAACAGCCCCACCTTGAAGCTGCT
 CAACCAACCCGCTGCTTGAAGAAATCTACAAGAGGGGCTGAAAAGGGAGATGAGGAGTTGAGGCCAGACTGCCCTTCAGCCTGACCCACTGACCCCTTACCCCC
 AAGTCCCTTCCCGCTGCCAACACAGGACAGCGCCCTGCTTTACAGCCCCACTCCAGCCATGGCTGCGGTACCACTAGCCCCAGTCCAAGGAGGACCTGGAGT
 CCGGAGTCAGAGTCCCTATGCTTCGAATCACAGTCCCTACCTCCAGGGCCAGCATGCGAGTCCCCACCTAGCCCCAGGGGAGATAGCCAGCACTACACCCCCAGCA
 GAGCTGGACACCAACCAAGAGGTCTTGAGACATGGGAAGCCGTGGTTGCGAGAGTTGTGCCAGGGCGGGGATCGGGATCCAGGGGACCATCACTCTCTCC
 ACAGCTTCAGGAGATGATGAGGAGACCACTACCAACCATCATCAACACCATCATCAACACAGTCCAGACACCAAGGCCCTTGTAGCTGGAAATTTCTAGGCCCAGAGG
 GCTCTGGACTCCCTACAGACCTCAGCTCCCCACTGATGTGGCCTGGACTGCTTCTACATCTCTGTCTACCTGGCTATGGCGTGGAAATCAAGGTCCAGAAATATCA
 GCCTCCGGAAAGGGAGACAGTGAAGGCTGGGGGGCCCCGACCACTGCCCTGGCCAAACCACTTTCTCTGCTGGGGGCAAGTCATCCGCAAGCCCCACC
 CACCAAGGGCCCTGAGGTTCCAGAGCTCCCGCCACCGGCTGGCCCTGGCACCTTCCATTTCCATTACCAAGCTATCTCTGAGTGCCACTTTCCCGCTGCTCAATGCCACCCAGC
 GGAGATGTGACTGCACAGCTCCACCCAGGGGTAGTGCCCGTTCCATTGTGCCACTGGCTACAGCTGAAGGGGCCAGGCATCTCACCTGTCTCAATGCCACCCAGC
 CCTTCTGGGATTCAAAGGAGCCGCTGCTGCTGCTTGGCGGAGTGATCCGCAATGCCACCGGCCCATCTCTCTCCAGGCTTCCCGGGCAACTACAGCAACAA
 CCTCACTGTACTGGCTTGAGGCTCTGAGGGCCAGCGGCTACACTGCATTTTGAAGAGTTTCCCTGGCAGAGGATGATGACAGGCTCATTCGCAATGGGGAC
 AACGTGGAGGCCACAGTGATGATTCTATGAGTGGAAATACCTGCCATTGAGGGCTGCTCAGCTCTGGCAACACTTTTGTGAGCTCAGTACTGACAGCAGCG
 GGGCAGCTGACGGCTGCGCTATGAGGCCCTCCAGCAGGGCCATTGCTAGAGCCCTTTGTCAAATACGGTAACTTCAGCAGCAGCACACCCACTACCTCTGTGGG
 TACCACTGGAGTTCAGTGCAGCCCTGGCTACACCTGGAGCAGGGCTCCATCATCATGAGTGTGTGACCCACGACCCCACTGGAATGGAATGAGACAGAGCCGCTGC
 CGAGCCGTGTGACGCGGGAGATCACAGATCGGCTGGCGTGGTACTCTCTCCAACTGGCAGAGCCCTACGGTCTGTGGCAGGATTTGATCTGGGGTGTGCATGTGGA
 AGAGACAAGCGCATCATGCTGGACATCCGAGTCTGCGCATAGGCCCTGTGATGTCTTACTTCTATGATGGGATGACCTGACGGCCCGGTTCTGGGCCAGTACTC
 AGGGCCCCGTAGCCACTTCAAGCTTTACCTCATGGCTGATGCACCATTCAGTCCAGTCGACCCCGGACCTCAGTGTGGGTACCAAGCGGGCTTCGTCTATCCACT
 TCTTTGAGTGCCCCGCAATGACACATGTCCGAGCTGCCTGAGATCCCAATGGCTGGAAGACCCATCGCAGCTGAGCTAGTGACGCGCACCGTGGTCACTTACCAGT
 GCTACCTGGCTACCAGGTAGTGGATCCAGTGTCTCATGTGCCAGTGGGACCTAACTTGGAGTGGGACCTGCCCTCATGCCAGAGGGTGACTTCTGCCACGATCCTGG
 AGATGTGGAGCAGACCCGACGCTCATATCCAGCCCCAAGTTCCCGTGGGGGCCAACCGTGCAATATATCTGTGACCAAGGTTTGTGCTGATGGGCACTCCATCCTCACC
 TGCCATGATCGCCAGGCTGGACGCCCAAGTGGAGTACCGGGCCCCTAATGTCTCTGGAAACAGCTCAAGCCATGCCATGGTCTCAGTGCCCCGTGAGATGGTGGCCCCGA
 AGTCTGAGAAAGCAGCTACCCAGCAGGGGCCACCATCCACTTCTGCTGGCTGGCTATGTGCTGAAGGGCCAGGCCAGCATCAAGTGTGTGCTGGGCACTCCCTCG
 CATTTGGAGTGACCCCCACCATCTGTAGGGCTGCCTCTCTGGATGGTTCTACAACAGTGCAGCCTGGATGTGCCAAGGCACCTGTGCTCCAGCACCCCTGGATGTCTG
 CCCACTGGCCGGCCACAGATCTGTGAGTGGCCACCGTGGCCAGCACCACTGTGGCAGGACCGTCACTCTCTCTCCCGCAAAACCCCAAGGACACCCCTCATGATCCTC
 CGGACCCCTGAGGTCACGTGCGTGGTGGACGTGAGCCACGAAGACCCGAGGTCAGTTCAACTGGTAGCTGGACGGCGTGGAGGTGCAATAATGCCAAGACAAAGCC
 ACGGGAGGAGCAGTTCAACAGCAGCTTCCGTGTTGGTTCAGCGTCTCACCGTTGTGACCAAGGACTGGCTGAACGGCAAGGAGTACAAGTGAAGGTCTCCCAACAAAGGCT
 CCCAGCCCCATCGAGAAAACCATCTCCAAAACCAAAAGGGCAGCCCCGAGAACACAGGTGTACACCTCGCCCCATCCAGGGAGGAGATGACCAAGAACCCAGGTCAAGCT
 GACCTGCTGGTCAAAGGCTTCTACCCAGCGACATCGCGTGGAGTGGAGCAATGGGCGAGCGGAGAACTACAAGACACGCTCCCTCCATGTGCTGAGCTCCGACG
 GCTCCTCTTCTCTACAGCAAGCTCACCGTGGACAAGAGCAGGTGGCAGCAGGGGAACGCTTCTCTCATGTCTCCGTGATGATGAGGCTCTGCACAAACCACTACACGCAGAA
 GAGCCTCTCCCTGTCTCCGGGTTGA

FIG. 4A

(SEQ ID NO: 9)

METDTLLWVLLWVPGSTGD**GAPG**SLSEAPTVKGQAPGIEETDGELTAAPTEQPERGVHFVTTAPTTKLLN
HHPLLEEFLEQEGLEKGEELRPALPFQDPAPFTPSPLPRLANQDSRPVFTSPTPAMAAVPTQPSKEGPWSPES
ESPMRITAPLPPGSMNAVPTLGPEIASTTPPSRAWTPTEGPGDMGRPWWAEVVSQGAGIGIQGTITSSTASG
DDEETTTTTITTTITTVQTPGPCSWNFGPEGSLDSPTDLSSTDVGLDCFFYISVYPGYGVEIKVQNISLREGETV
TVEGLGGPDPLANQSFLLRGQVIRSPTHQAALRFQSLPPAGPGTFHFHYQAYLLSCHFFRRPAYGDVTVTSLH
PGGSARFHCATGYQLKGARHLTCLNATQPFWDSKEPVCIAACGGVIRNATTGRIVSPGFPNGYSNNLTCHWLL
PEGQRLHLHFEKVSLEAEDDDRLLIRNGDNVEAPPVYDSYEVEYPIEGLSSGKHFFVELSTDSSGAAAGMALRYEA
FQQGHCEYEPVKYGNFSSSTPYPVGTTVEFSCDPGYTLEQGSIIIECVDPHDPQWNETEPACRAVCSGEITDSAG
VVLSPNWPEPYGRGQDCIWGVHVEEDKRIMLDIRVLRIGPGDVLTIFYDGGDLTARVLGQYSGPRSHFKLFTSMA
DVTIQFQSDPGTSVLGYQQGVHFFFEVPRNDTCPELPEIPNGWKSPSQPELVHGTVVITYQCYPGYQVVGSSVLM
CQWDLTUSEDLPSCQRVTSCHDPGDVEHSRRLISSPKFPVGATVQYICDQGFVLMGSSILTCHDRQAGSPKWSD
RAPKCLLEQLKPCHGLSAPENGARSPEKQLHPAGATIHFSCAPGYVLKGQASIKCVPGHPSHWSDPPPICRAASLD
GFYNSRSLDVAKAPAAASSTLDAAH**L**AGHRSVECPPCAPPVAGPSVFLFPKPKDTLMISRTPEVTCVVVDVSHED
PEVQFNWVYVDGVEVHNAKTKPREEQFNSTFRVVSVLTVVHQDWLNGKEYCKVSNKGLPAPIEKTISKTKGQPR
EPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPMLDSGSGFFLYSKLTVDKSRWQQ
GNVFSCSVMHEALHNHYTQKSLSLSPG

FIG. 4B

>cDNA sequence encoding mature mouse SEZ6 protein

(SEQ ID NO: 10)

CTCTCTCAGAGGCTCCGATCACGGGGGAAGGTTCATGCCACGGGGCATCAGGGGAGCGGATGGGGAGCTGACCGCAGCCCTACACCTGAGCAGTCAGACC
 GAGCGTCCACTTCGTCACCACAGCCCTACCTCAAGCTCAAGCTGCTCAACCAACCACTCTGGAAGAAATTTCTTCAAGAGGGGCTAGAAAGAGAGGAAGCGC
 CGAGCTGCACTGCCCTTCCAGCGGACTCACCTACACATTTACTCCAAGCCCTTCCCCCGCTCACCAACCAAGGACAACCGCCCGTCTTACCAGTCCG
 ACTCAGCCGTGGCTGCAGACCCACCGCCACTCCAGGGAGAACTTGGAACTAGAACTCAAAACCCCTGAGCTTTCTATCATCTGTCCTTCTCTCC
 AGGGCCGAGTATGGCAGTCCACACTGCTCCAGAGGACAGACCCAGTACTACACCCCTAGCCAAGCATGGACTCCAACCTCAGGAGGGTCTCTGGAGACA
 TGGACAGACCTTGGTTCAGAGTTCATGTCTAAGACCACAGGGCTTGGTCTGAGGGAACTTGGCACTCCACAGCTTCAGGGATGACGAAGAGACC
 ACTACCACCATCATACCACTACTGTACCAAGTTCAAGCACAGGCCCTGTAGCTGGAATTTCTCAGGCCCAGAGGGCTCTCTGGATTCCCCACGGCCCC
 CAGCTACCCCTGTGCTGGCTGGACTGTTTCTACTATATCTGTCTACCCCTGGATATGGAGTAGAGATCAAGGTGGAGAACATCAGCCTTCAGGAAGG
 GAGACCATCACCGTGGAGGGCTGGGGGCCCGGATCCACTGCCCTTGGCTAACCAGTCTGTTCTGCTGAGGGGCCAGGTCTATCCGACGCCCAACCCACCA
 AGCAGCCCTGAGGTTCCAGAGCCCTCCGCTACCCGCTGGGCTGGCACTTCCATTTCCGCTACCAAGCCTATCTCTGAGCTGCCACTTTCCCGACGTCCAG
 CGTATGGAGATGTACTGTACCACTCCACCCAGGAGGCGGCCACTTCCATTGCGCTACCAAGCCTATCTCTGAGCTGCCACTTTCCCGACGTCCAGTGTCT
 CAATGCCACCCAGCCCTTTGGGATTTCCCAAGAGCCCTGTTGCACTGCTGTTGGTGGAGTATTCGGAATGCCACCACTGGCCGATGCTCTCTCTGGC
 TTCCCGGGAACACAGCAACAACCTCACCTGCCACTGGTTGCTAGAGGCTCCAGAGAGCCAGGGCTGCACCTTGCACTTTGAAAGGTCTCCCTGGCAGAA
 GACGACAGAGCTCATATCCGCAATGGAATAACGTGGAGGCCCGCGGTGTACGACTCTATGAGGTGGAATACCTGCCCTTCAGCAAGGACATTGCTATGA
 CTCTGGCAGACACTTCTCGTGGAGTTCAGTACTGACAGCAGTGGGGCAGCTGCAGGCATGGCCCTGCGCTATGAGGCCCTCCAGCAAGGACATTGCTATGA
 GCCCTTTGTCAAATACGGCAACTTCAGCAGCAGTGCACCGTCTACCCCTGTGGGTACAACTGTGGAGTTCAGCTGTGACCCCTGGCTACACCTGGAGCAGGG
 CTCCATCATATCGAATGCGTCGACCTCCACGACCCCAAGTGAATGAGACAGAGCCAGCTGCCGAGCCGTGTGCGAGCGGGGAGATCACAGACTCTGCAG
 GCGTGGTGTCTCTCCAACTGGCCGGAGCCCTATGGCCGAGGGCAGGACTGCATCTGGGGTGTGCTGTGGAGGAGGACAAGCGCATCATGCTGGACAT
 CCGAGTGTGCGCATAGGCTGTGGGATGTACTGACCTTCTACGATGGGATGACCTCACAGCCCGGTCTCTGGCCAATACTCAGGGCCCCGTGGCCACTT
 CAAGCTTTTACCTCCATGGCCGATGTCACCATCCAGTCCAGTCCAGACCTGGGACCTGGCGCTGGGTTACCAAGCAAGGATTGTATCCACTTCTTTGAG
 GTTCCCGCAACGACACATGTCCAGAGTACCGAGATCCCAACGGCTGGAAGAACCCATCACAGCTGAGCTGGTGCACGGCACGGTGGTCACCTATCAG
 TGCTACCTGTTACAGGTGGTGGATCCAGTATTTCTATGTCCAGTGGGACCTAAGCTGGAGTGGGACCTGCCTTCATGCCAGAGAGTGACATCTTGC
 CATGACCCAGGGGATGTGGAGCACAGCCGACGCTCATATCCAGCCCCAAGTTTCCCGTGGGAGCAACTGTGCAATATGTCTGTGACCAGGGTTTGTGCTG
 ACGGGGAGTGCCATTCTACCTGCCATGATCGGCAAGCAGGCGAGTCCCAAGTGGAGTGACAGGGCCCCCAAGTGTCTTGGAAACAATTCAAGCCGTGCCA
 TGGCTCAGCGCCCCGGAGAAATGGTCCCCGAGCCCTGAGAAGCGGCTTACCCAGCAGGGGCCACCATCCACTTCTCTGTGCCCTTGGTTATGTGCTGAA
 GGGCCAGGCCAGCATCAAAATGCGTGCCTGGACACCCCTCGCATTGGAGTGACCCACCACTCTGTAGGGCTGCCCTCTGGATGGGTCTACAACGGCCG
 TAGCTTGGATGTGCCAAGGACCTGCCGCTCAGTGCCTTGGACGCTGCTCACCTGGCTGCTGCCATCTTCTACCATTTGGTGGCCATGGTGTGCTGGTG
 GGAGGAGTGTAACCTCTATTTTCCAGATTCAGGGGAAAAGTCCCCTGCAACTTCCCCGAACTCATCTCGCCCCCTATAACCGCATCACGGTAGAGTCAGCAT
 TTGACAACTCAACTTATGAGACTGGATCTCTTTCCTTTGACGAGAGCAGAGAATATGA

FIG. 5A

>Translation of mouse SEZ6 ORF

(SEQ ID NO: 11)

LSSEAPITGEGHATGIRETDGELTAAPTPEQSDRGVHFVTTAPTLLKLNHHPLLEEFLEGLEREAPQPALPFQPDSPTHFT
PSPPLRLTNQDNRPVFTSPTPAVAAAPTQPHSREKPWNLESKPPELSITSSLPPGPSMAVPTLLPEDRPSTTPPSQAWTPT
QEGPGDMDRPWVPEVMSKTTGLGVEGTIATSTASGDDEETTTIITTTVTVQPPGPCSWNFSGPEGSLDSPTAPSSPSD
VGLDCFYYISVYPGYGVEIKVENISLQEGETITVEGLGGPDPLPLANQSFLLRGQVIRSPTHQAALRFQSLPLPAGPGTFHFR
YQAYLLSCHFRPRPAYGDVTVTSLHPGGSAHFHCATGYQLKGARFLTCLNATQPFWDSEQEPVCIAACGGVIRNATTGRIVS
PGFPGNYSNNLTCHWLLLEAPESQRLHLHFEKVSLEAEDDDRLLIIRNGNNEAPPVYDSYEVEYLPIEGLSSGRHFFVEFSTD
SSGAAAGMALRYEAFQQGHCHCEPFVKYGNFSSAPSYPVGTTVEFSCDPGYTLEQGSIIIECVDLHDPQWNETEPACRAV
CSGEITDSAGVVLSPNWPEPYGRGQDCIWGVHVEEDKRIMLDIRVLRIGSGDVLTFYDGGDLTARVLGQYSGPRGHFKLF
TSMADVTIQFQSDPGTSALGYQQGFVIHFFEVPRNDTCPELPEIPNGWKNPSQPELVHGTVTVTYQCYPGYQVVGSSILMC
QWDLWSSEDLPSCQQRVTSCHDPGDDVEHSRRLISSPKFPVGATVQYVCDQGFVLTGSALTCHDRQAGSPKWSDRAPKCLL
EQFKPCHGLSAPENGARSPEKRLHPAGATIHFSAPGYVLKGQASIKCVPGHPSHWSDPPIICRAASLDGFFYNGRSLDVAK
APAASSALDAAHLAAIFLPLVAMVLLVGGVYLYFSRFQGSPLQLPRTHPRPNRITVESAFDNPTYE TGSLSFAGDERI

FIG. 5B

>cDNA sequence of rat SEZ6 ORF

(SEQ ID NO: 12)

ctctctcagagagctccaatcacgggggaaagggtcaagccacgggcatcaggggagatggatggggagctgaaccgagccctacacactgagcagtcagacgagggcgtccactctgcacac
agccctaccctcaagctactcaaccaccacccactctcgagagaaatttctcaaggggggctagaaggagagagagaaagctccgagggccggcacctgcccctccaggccagactcacctacac
cctttactcaagcccccctccgcctcaccacaggacaaacggccctctttaccagtcgagccagctgtagctgcggcaccacgcaaggcccccactccagaaagaaacccctggaaacc
cagagtcagagggcccccggagctttacatcacatctccctccctccagggccggagtatgagcagtgcccacactgacccagagagacagaccacgactacacccccagccaagcatggact
ccaaccagggaggtcctggagagacatggggcagaccctgggtccagagatcatgtctaagaccacagggcttgggtatcggaggggaccattggccacctccacagcttcagggggatgacgaaga
gaccacaccacacatcattaccacogtcacacaattcagccacagggcccctgtagctggaaatttccagggcccggaggggctctctggattccctgggggtcccacggcgtccctctgatg
ttggcctggactgtctactacatctctgtctacocctggatatggagtcgagatcaagggtgaagaacatcagcccttcaggaagagagagaccataaacgctggagggcctggggggcctgacc
cactgcccctggctaacaggtcttctgctgagggggccaggctacccgagcccccacccacaggcagccggtagggttccaaaggccttccactcccgctgggacctgggtactttccatttccact
acaaaggcctatctcctgagctgccatttctcggcgtcagcttatggagatgtgtacacagcctccaccagggagcagcggccttccactgtgccactgggctaccagctaaagggggt
ggcagggttctcaacctgtctcaatgccaccagcccctttggggattcccaaggagcctgtctgtcattgtgtggagggatgtggaatggccaccactggggcgcattgtctctcctgggctttcc
cggggaactacagcaacaacctcacctgccactggctgtagaaagccccggagagccagcggctgcacctgcacttgaaagggtctccctggcagaagatgacgacagggtcatcatccgta
acggggaataacgtggagggcccccagctgtatgactctatgaggtggagtagacctgcccattgagggggcctgctcagttctggcagagacacttctctggagagttcagctactgacagcagcgggggc
agccgcaggcatggcactgctgtatgagggccttcacagcaaggacattgtctatgagccctttgtcaaatacggtaacttcagcagcagccgctcctacccctgtgggtacgactgtggaggttc
agctgtgaccctggctacaccctggagcaggggttccatcatcctgaatggctgcgacctcgtgacccccagtggaatgagacagaaaccagcctgcccggagccgggtgagcagcggggagatcac
agactcgcaggcgtgggtctctccaaactggccggagccttatggccggagggcaggactgcatctgggggtgtgcatgtggagggagggacaaaggcgcatactgctggagacatccgagtgctgcg
cataggctctggggatgtactgaccttacgatagggggagatgacctgagacagccgggtcttgggggccaatactacagggggcccggtggccacttcaaggctctttaacctccatgggtgatgtcacattc
agttccagtcagaccctgggagctcgggcgtgggttacagcaaggatttgtcatccacttcttggaggtgccccgcaatgacacatgtccagagcttccggagatcccccaacggctgggaagaa
cccatcacagcctgagctggcatggcaggggtggtcacctatcagtgctacccgggtaccaggtgggtggggtaccagttatctcatgtgcccagtggggacctggagctggagtgagggacctgscct
catggcagagagatgacatcctgaccatgacatgcaagccggagcacagccgcctcatatccagcctcaaatgttctcttggggagcaactgtgaggtatatctgtgacacaggggttttggctc
acgggtagcgccatccttacttgccatgatcgtcaagcggggcagtcaccaagtgagagtgacagggggcccccaaggtctcttggaaacagttcaaacatgtcatgggctcagtgccccctgagaatg
gtggccgcagcccctgagaagagggctccaccagcagggggccaccattcattctctgtgccccctgggtatgtgctggaaggccagggccagcagcatcaaatgctggctgggacaccccctcacattg
gagtgatcctccacccatctgtagggtgcttctctggatgggggtttacaaacggccgtagacctggatgttgcacagggacactgcccagatgctggcccatggcagctgcc
tcttttaccattgggcatgggtgtgctgggggagaggttaccttatcttcagactccaggggaaaaagttcctctgcagcttccgggaactcatctctcgccccctataacccgtatcacggt
agagtcagcatttgacaatcaacttatgagccggatctcttcttcttggaggagacgagagaata

FIG. 5C

>Translation of rat SEZ6 ORF**(SEQ ID NO: 13)**

LSSEAPITGEGQATGIREMDGELTAAPTPEQSDRGVHFVTTAPTLLKLNHHPLLEEFLEQEGLEGREEAPRPALPFQPDSPPTFTPS
 PLRLTNQDNRPVFTSPTPAVAAAPTQPHSRKKPWNPESEPPELYITSLPPGPSMAVPTLHPEDRPSSTTPPSQAWTPTQEGPG
DMGRPWWPEIMSKTTGLGIEGTIATSTASGDDEETTTTITTTTITQPPGPCSWNFSGPEGLSDSPAVPSVPSDVGLDCLYISV
YPGYGVEIKVNISLQEGETITVEGLGGPDPLPLANQSFLLRGQVIRSPTHQAAVRFQSLPLPAGPGTFHFHYQAYLLSCHFRRP
AYGDVTVTSLHPGGSARFHCATGYQLKGARFLTCLNATQFWDSEQEPVCIACGGVIRNATTGRIVSPGFPNGYNNLTCHWL
LEAPESQRLHLHFEKVSLAEDDDRLLIRNGNNVEAPPVYDSYEVEYLPIEGLSSGRHFFVEFSTDSSGAAAAGMALRYEAFQQGH
CYEPFVKYGNFSSAPSYPVGTTVEFSCDPGYTLEQGSIIIECVDLRDPQWNETEPACRAVCSGEITDSAGVVLSPNWPEPYGRG
QDCIWGVHVEEDKRIMLDIRVLRIGSGDVLTFYDGGDLTARVLGQYSGPRGHFKLFTSMADVTIQFQSDPGTSALGYQQGFVI
HFFEVPRNDTCPELPEIPNGWKNPSQPELVHGTVTYQCYPGYQVVGSSILMCQWDLWSWSEDLPSCQRVTSCHDPGDVEHS
RRLISSLKFPVGATVQYICDQGFVLTGSAITLCHDRQAGSPKWSDRAPKCLLEQFKPCHGLSAPENGARSPEKRLHPAGATIHFC
APGYVLKGQASIKCVPGHPSHWSDPPPICRAASLDGFYNGRSLDVAKAPATSSALDAAHMAAAIFLPLVAMVLLVGGVYLYFSR
 LQGKSPQLPLPGTHPRPNRITVESAFDNPTYETGSLSFAGDERI

FIG. 5D

(SEQ ID NO: 14)

[illegible]

>Translation of cynomolgus SEZ6 ORF

(SEQ ID NO: 15)

METDTLLLWVLLWVPGSTGDGAPLSEAPTMGKGQAPGIETDGELTAAPTEQPER
GVHFVTTAPTKLLNHHPLLEEFLQEGLEKGDEELPALPFQPDPPTPFTPSPLPRLANQ
DSRPVTSPTPATAAVPTQPQSKEGPWSLESEPPVLRITAPLPPGPSMAVPTLGPGERPS
TTPPSAWTPTQEGPDMGRPWVEVSQGAGIGIQGTIASSTASGDDEETTTTTTIITT
TITTVQTPGCSWNFSGPEGSLDSPTDLSSPPDVGLDCFFYISVYPGYGVEIKVQNISLR
EGETVTVEGLGPAPLANQSFLLRGQVIRSPTHQAALRFQSLPPPAGPGTFFHYQAY
LLSCHFPHRPAYGDVTVSLHPGSARFHCATGYQLKGARHLTCLNATQPFWDSKEPVCI
AACGGVIRNATTGRIVSPGFPGNYSNNLTCHWLLEAPEGQRLHLHFEKVSLAEDDDRLII
RNGDNEAPPVDSYEVELPIEGLLSSGKHFFELSTDSSGAAGMALRYEAFQQGHCY
EPFVKYGNFSSAPTYPVGTTVEFSCDPGYTLEQSIIECVDPHDPQWNETEPACRAVC
SGEITDSAGVVLSPNWPEPYGRGQDCIWGVHVEEDKRIMLDVRVLRIGPGDVLTFYDGD
LTARVLGQYSGPHSHFKLFTSMADVTIQFQSDPGTSVLGYQQGFVIHFFEVPRNDTCPEL
PEIPNGWKSPSQPDLVHGTVVTYQCPGYQVVGSSVLMCQWDLTWSEDLPSCQRVTSCHD
PGDVEHSRRLISSPKFPVGATVQICDQGFVLTGTSILTCHDRQAGSPKWSDRAPKCLLE
QLKPCHGLSAPENGARSPEKRLHPAGATIHFSCAPGVVLKGQASIKCVPGHPSHWSDPPP
ICKAASLDGFYNSRSLDVAKAPASSTLDAAHIAAIFLPVAMVLLVGGVYFSRLQG
KSSLQLPRTRPRPNITVESAFDNPTYETGSLSFAGDERI

FIG. 5F

>cDNA sequence of human SEZ6L ECD

(SEQ ID NO: 16)

ctcgaagagagatgctctgctgagggagagatgctccctctcggaaccttatctgctgcccagcggaagctcttgagagagggagatccccgggaaagggaagcatcccgaaagaaagtggtcaagct
ccccctagctccagccagagcgctgaggtgctgggagaaactggtcctgagcggaacagccccttcggcccatcaggtatctctgcccctcagccctctctcccccggaggaagctagggccttaaac
acgcccctccccctaaaaaagaagctgccttccctcaagcagaggtcaattccgcccaggaagcagctcagaccacaagggccacctccgctgctacagttccaaagaagctggatccccagcctgccagc
cagggagctcgtatgctcagcagctccacagaaaacactggagacctctggcgatctgacccctatttggtggccagcgaaagaaagctagcgaaagtccctctgtggtggtgacagggaaagaggtccgct
gtccccaccacacccgctctctccagatcagcccccttcacctccagcccttatgtgctcatcacactgctcagagggcctgagcctggcgaaacctggacctgacatgggctcagggagggctctctca
ggagagacacagcccctatggccctgatggataaaggggcggaatgaactgaccggcgaaggaagccagcggaagaaagccagagagaccaccacacagcacacaatcatcaccacccgctcatcaccac
cgaaacaggccccgctctgtgtccgtgtcttttccaaccccgagggggctactattgacagcagcgattacccctgctcctctcaaacacttctctgagtgacacctacaatgtgacctgtgtatcac
cggtacggaggtcgaaactccaggtgaagtccgtgaacctctcgatggcgaaactgctctccattaggggcggtcgatggccctacactcacctcctggctgtaaccaaaccttgcctgctgctggaaggc
caggtgattagggtcccccaacaacatctcgtctactttcaggacaccttcaagacgacgggactggggaaccttccaactgcatcaccagggccttcatgctgtctctgttaatttccccagaggagaccc
gactccggagacgtcacggtcatggatctgcactccggaggcggtggccactttcattgtcaactcggtacgagctcagggcgccaagatgctgacatgcatcaaacgcccagcaaacctcact
ggttccagccaggagacctatctgtaggctccttgcggcgaggcggtgcacaaatgctacaattggcagagtgctcagcccttctacccttgaaacacacacacgggctccccagttctgcatctgggac
aatcgagggcccccgaaaggccaaaaggctgcacctgcactttgagagagctctgctccacgacaaaagacaggaatgacccgtgcactccgggcccagaccaaataaagtccgctctctgtatgacagcc
tgacagacagagtcctcgctcttttgaaggccctgctgtccgagggcacaataccatcaggagttgagttcacatccgaccaaaggccagggtgctgtagcaccttcaacattaggttgagggtcttccgaaaaag
ggacacgtctacggagccctatatcagaatggcaatttcacaaacctccgacccccaccctacataatcggcacaaattgggaggttttaccctggacccttggaacacacagcctggagcagggacacctggcca
tcatcgaaatgcatcaacgtcaggggacccctacttggaacgacacagaaacctctgtgtaggggctatgtgcggagagcggaactgagcgctgtggtggtgagagctgcttccccctaaactgggccccggaac
cctatgtggaggcggaagattgcatctgggaagatccacgtcggcgagggaaaagaagagatctttctggacatccagttcctggaatctctccaaacagcgacatctctgacccatctacgacgggagatga
ggttcatgccccacattctggggccagttatctcggaaaactcggcccccaaaaggctctactctccaccccgacctcaaatccaatccacagcggtatcctgctgggacctcatcttgggaaaaggggac
aaggctttatcatgaattacatcgagggtcagcagaaaacgcagagctgctccgacctgctgagatccagaaacgggattggaaagaccacctcccaacacggagctcgtcaggggagagctagagatcacaa
taccagtgacccccggatacgcacatcgtcggtcccgatatccctggacatgccagtggggattctgagctggagctccgaccccccttttgtggagaaagatcatgtactgcaaccgaccccccgcggaag
tcgatcatagcaccagggctcatcagcgatctctgtgctgctgcgggcacaaaccatccaatacacctgtaacccccgggattcgtgctcgaaaggatctctccctgctcaactgtttacagcaggggaaaccc
ggcacccttctggacatccagggctgcctcactgcgtgtccgaagagagcctggcttgcgataatccccggcctgctcgagaaacgggataccagattctgtacaaaagggtgtaccttccccggcg
agttccctgaccttcatgtgtctacgaaggaattcggagctcatggcggaagatcaccatcagagtgcatctctcgccagccctcccactggaacgggacctctccccgtctgttaaggtcaatcagggaattcc
ttcgagcacgctctgggaagtcgtgagggtgcccgcggagagacaagcctgggaaggcgggc

FIG. 5G

>human SEZ6L ECD protein

(SEQ ID NO: 17)

METDTLLWVLLWVPGSTGDHGAPLERDALPEGDASPLGPYLLPSGAPERGSPGKEHPEERVVT
 APPSSQSAEVLGELVLDGTAPSAHHDIPALSPLLPEEARPKHALPPKKKLPSLKQVNSARKQLRPK
ATSAATVQRAGSQPASQGLDLLSSSTEKPGPPGDPDPIVASEEASEVPLWLDRKESAVPTTPAPLQI
SPFTSQPYVAHTLPQRPEPEGPDMAQEAPQEDTSPMALMDKGENELTGSASEEQETTTSTII
TTTVITTEQAPALCSVSFSNPEGYIDSSDYPLPLNNFLECTYNVTVTGYGVELQKSVNLSDGELL
SIRGVDGPTTLTVLANQITLLVEGQVIRSPTNTISVYFRTFQDDGLGTFQLHYQAFLSCNFRPRPDS
GDVTVMDLHSGGVAHFHCHLGYELQGAKMILTCINASKPHWSSQEPICSA PCGGAVHNATIGRV
LSPYPENTNGSQFCIWTIEAPEGQKLHLHFERLLHDKDRMTVHSGQTNKSALLYDSLQTESVPF
EGLLSEGNIRIEFTSDQARAASTFNIRFEAFEKGHCYEPYIQNGNFTTSDPTYNIGTIVEFTCDPGH
SLEQGPAIIECINVRDPYWNDEPLCRAMCGGELS AVAGVVLSPNWPEPYVEGEDCIWKIHVGEE
KRIFLDIQFLNLSNDILTIYDGDVMPHILGQYLGNSGPQKLYSSTPDLTIQFHS DPAGLIFGKGQG
FIMNYIEVSRNDSCSDLPEIQNGWKTTSHTELVRGARTYQCDPGYDIVGSDTLTCQWDLSSWSSD
PPFCEKIMYCTDPGEVDHSTRLISDPVLLVGTTIQYTCNPGFVLEGSSLLTCYSRETGTPIWTSRLPH
CVSEESLACDNPGLPENGYQILYKRLLYPGESLTFMCYEGFELMG EVTIRCILGQPSHWNGPLPVC
KVNQDSFEHALEVAEAAETSLEGGLAGHHHHHHHHHH

FIG. 5H

>cDNA sequence of human SEZ6L2 ECD

(SEQ ID NO: 18)

ctgcctctcaaaagagaagagatttctcccgagcccggtatccgagacacccacagtggttcggaagccctcgtgaactgctgcacggagccctctctgagaaaggggacctgaaatgggcta
tctccctggctccgacagagatccacactcgcacacactcgtggaagaccctcgtgtgcttccctgcccagagccacagaaacccgggaacagggccctctcaacagctgtgacccct
aacggcgctcagaggaagctggacacctacagccctgagctgctgacacctctctggcacaaacggctcctcctctcctcctgtagccctggacccctctcggacctggaaggagagcgagagg
agagagacaaccaccaccattataccaccaccaacggtagacaaccacagtgaccgacctgtctgtgcaacaacaacacatcagcgaaaggcggaagggtatgttggaatccccctgacctgggctcc
cctgtgtccagaacacactcggcctcctggattggcacatactcattcacgtgtaccccggtctacgggaatcgagattcagggtagacacctggaatctgtccagggagaggaaactgctggtgctgg
ctggcggaggaagacccctggcctcctgtagactcctgctaactcctcatgctcggcggaaggccagggctctcagatccccctaccacagggctgctcctgcacttccagagagccccagagagtggcc
tagagaggagcggttcaggattcaatacaggccctatctctgagctgtgggattcctccagaccggctcatgtagcgatgtctcgtcacccgacctcacccccgggaggaacagacccttcacat
gtgattccgggataaccagctgcaaggcggaaggagaccctgatttgccctaattggcacaggcccgagctggaaacgggagagacacctagctgcatggctagctggcggcggaacccatccataatggcc
accctcggcagggtatcgtcagccctgaacctggcgaggctgtgggaccttaacctcacatgcagatgggggtgctcgaaagctgctgaaaggcagaggagactccacctccacttcgagagagggtgtccctgg
gacgagaggacaacgacagagctcatggtcagaaagcggcggaagccctctcagccctgtgatttacgacagcgacatggaaacgcatgtgacctgagagaggggccctcatctccgatggccccaaagcctgtga
cgtggaaactcctctcggagaccccgctaaccctcctcctgagcctcagattcgaggcccttcgagaggagacagatgtttcgtcctcttctgtggcccatggccaacgtgacacacaacccgacccccg
agtacagagacccggagctctggctaccttcagctgtcctggctacgcccctggaacctccggagacctctaattgcccataatgtgtgggatccacccggaaccccatgggaacgacacacggagccc
cgcttgtaaggctatgtgcggcgaggagaactcagcgaacctcagcgaacctgtcctcctcctgattggccccacagagctattccccggacaagaaagactgtgtctggggcggtgcaactccagggagga
aaaggaggatcctcctccagggtggagattctgaaactcagagaggagagacatgctgacacctgttcgacggagagacggagaccttcggccagaggtcctcgtcagctgagagaggccctcagccccagaa
ggagagactgctcagctccggccccgatgacactccagtttcaggccccccctggcccccttaattccttggcctgggacagagggttctgctccacttcaaggagaggtccccagaggaatgatacatg
ccccgaaactgctcctccggagtggagatggagagacagcttcccatggcgacctgatacagggggaacccgtgctgacatatcagtgtagaacccggctacagagctgctgggaaagcggtatctcctgac
ctgtcagtgaggatctctcctggagcgtgctccctcctgctcctgtaagaaatcatgacctgctgacacctggagagagatcgctaaacggccccacagagaccgcttccgacgctgggatttccccgtgggct
cccacgtgcaatacagggctccctcccggtatctcctcgaaggcgtgcccattgacatgctacagcagggagacacccggcacaccccaagtggtccgacaggggtgccccaaatgtgtccttgaagt
acgagccctgtctcaatcccggaagtcccgagaaacgggataccagaccctgtacaagcaccactatcaggccggcggaatccctgagattcttctgtactacgaggggttcgaagctcctcggcgaggg
tgacaattacctgtgtgccccggccaaccttccagtggaaccagccagccccctctctgtaaagtgctgcctacgaaagagctgctcgacaatataggaaagctggagaggtcacccagaccaccgaccttc
cagacaactgggaaggcggc

FIG. 51

>human SEZ6L2 ECD protein

(SEQ ID NO: 19)

METDTLLWVLLWVPGSTGDGAPLPLKEEILPEPGSETPTVASEALAE¹LLHGALLRRGPEMGYL
PGSDRDPTLATPPAGQTLAVPSLPRATEPGTGPLTTAVTPNGVRGAGPTAPELLTPPGTTAPPPPS
PASGPPLGPEGGEEETTTIIITTTT²⁰VTSPVLCNNNISEGEGYVESPDLGSPVSR²¹TLGLLDCTY
SIHVYPGYGIEIQVQTLNLSQEEELLVLAGGGSPGLAPRLANSSMLGEGQVLRSP²²TNRLLLHFQSP
RVPRGGGFRIHYQAYLLSCGFPPRPAHGDVSVTDLHPGGTATFHCDSGYQLQGEETL²³CLNGTRPS
WNGETPSCMASCGGTIHNATLGRIVSPEPGGAVGPNLTCRWVIEAAEGRRLHLHFERVSLDEDN
DRLMVRSGSPLSPVIYDSDMDDVPERGLISDAQSLYVELLSETPANPLLLSLRFEAFEEDRCFAPF
LAHGNVTTTDP²⁴EYRPGALATFSCLPGYALEPPGPPNAIECVDPTEPHWNDTEPACKAMCGGELSE
PAGVVLSPDW²⁵PQSYSPGQDCVWGVHVQEEKRILLQVEILNVREGDM²⁶LTFLFDGDP²⁷SARVLAQL
RG²⁸QPRRRLSSGPDLT²⁹LQFQAPP³⁰GNPGLGQGQGVLFHFKEVPRNDTCPELPPPEWG³¹WRTASH
GDLIRGTVLTYQCEPGYELLGSDILTCQWDLSW³²SAAPPACQKIMTCADPGEIANGHRTASDAGFP
VGSHVQYRCLPGYSLEGAAMLTCYSRDTGTPKWSDRVPKCALKYEPCLNPGV³³VPENGYQTLYKHH
YQAGESLRF³⁴FCYEGFELIGEVTITCVPGHPSQWTSQPP³⁵LCKVAYEELL³⁶DNRKLEVTQT³⁷TDPSRQLEG
 GLAGHHHHHHHHH

FIG. 5J

Relative Expression Values for mRNA Transcripts Associated With Neuroendocrine Tumors in NTX Lines using Whole Transcriptome Sequencing

	DLL1	DLL3	DLL4	NOTCH1	NOTCH2	NOTCH3	NOTCH4	ASCL1	NCAM1	CHGA	HES1	HES6	HEY1
LU37p3 - LCNEC	86.2	93.7	4.5	0.1	14.4	7.8	0.7	2690.8	72.0	94.0	2.7	55.1	5.6
LU64p2 - SCLC	10.9	34.3	10.8	8.2	0.2	0.7	5.7	418.4	57.8	739.5	6.3	2605.8	2.0
LU73p1 - SCLC	176.1	227.6	16.6	4.2	0.0	33.0	79.7	3429.1	77.5	515.0	34.5	3270.6	33.9
LU86p3 - SCLC	4.7	11.9	12.2	18.7	176.8	14.5	0.5	0.4	394.9	17.7	13.9	285.2	9.9
LU95p2 - SCLC	2.4	16.0	1.6	2.1	0.4	8.5	12.2	273.2	171.5	18.2	2.8	72.6	9.3
LU137p0 - LU_Ad	1.8	0.0	4.3	10.6	247.7	27.5	0.4	0.0	0.0	0.0	54.6	1.5	1.7
LU146p0 - LU_Ad	0.0	0.0	0.3	5.6	56.8	37.9	3.9	0.0	0.8	0.0	42.8	0.8	0.3
LU153p0 - LU_Ad	0.8	0.0	5.7	8.1	195.1	8.0	6.5	0.2	4.4	0.0	4.8	0.2	1.4
LU49p4 - LU_SCC	2.6	0.7	0.0	7.6	104.2	0.0	1.3	0.0	0.0	0.0	62.1	4.3	0.0
LU70p4 - LU_SCC	4.7	0.0	0.8	12.6	123.2	1.8	0.1	0.0	0.0	0.3	42.3	2.1	0.0
LU76p5 - LU_SCC	0.8	0.0	4.8	20.0	32.5	0.1	0.0	3.4	0.3	0.3	23.0	0.8	0.0
OV26p3 - OV	34.2	65.4	15.7	0.0	101.0	17.4	0.6	2253.7	128.8	23.1	4.3	35.1	7.1
OV100p0 - OV	0.0	0.5	0.4	3.6	154.2	16.3	0.5	0.0	2.0	0.0	84.1	0.0	17.3
OV45p3 - OV	1.7	1.9	0.1	14.9	53.2	84.5	2.7	0.0	60.4	0.1	14.6	2.4	84.4
OV55p5 - OV	0.3	0.2	0.0	31.0	139.8	71.7	1.4	0.0	11.4	0.0	19.4	1.9	6.7
OV72METp0 - OV	0.0	0.1	0.2	1.6	303.1	46.8	0.2	0.3	34.5	0.1	17.1	1.9	2.0
OV91METp0 - OV	0.3	1.6	0.1	10.5	340.1	345.3	2.3	0.0	3.9	0.0	31.7	1.2	1.0
Normal Lung 1	1.7	0.0	5.7	8.2	85.9	33.1	11.4	0.4	3.4	0.0	13.8	0.1	11.4
Normal Lung 2	17.0	0.1	8.8	24.0	81.5	54.0	62.8	5.3	4.6	0.4	23.2	2.4	43.0
Normal Lung 3	26.9	0.2	41.6	146.0	25.6	239.3	91.7	0.8	1.8	1.3	11.9	8.1	40.2
Normal Lung 4	0.2	0.0	6.0	11.8	81.5	40.4	15.8	0.0	1.2	0.0	11.4	0.3	14.1
Normal Ovary	0.3	0.0	5.1	7.8	250.9	44.1	5.1	0.6	125.5	0.2	8.5	0.7	0.4

FIG. 6A

Relative Expression Values For SEZ6 Transcripts in Various Samples as Determined by Whole Transcriptome Sequencing




FIG. 6B

Microarray Data Showing Clusters
of Tumors with Neuroendocrine Features




FIG. 7A

Average Normalized Intensity Values
for Common Markers of Neuroendocrine Phenotype

	Gene Symbol	Median (48 samples)	Cluster C				Cluster D				Cluster E		Cluster G	
			KD66	OV26	LU102(SCLC)	LU37- qc(LCNEC)	LU86(SCLC)	LU64-qc(SCLC)	LU95(SCLC)	EM6	LU100(SCLC)	LU49(SCC)	LU85(SCC)	
Achaete-scute complex homolog 1	ASCL1	9	6589	8238	9382	12169	9664	11	3390	10366	14	261	8	5
	CALCA	73	15180	8352	10035	14633	13335	70	24	2477	53	45	52	39
	CALCB	143	2534	1147	1547	2584	2757	13106	433	1264	309	135	65	81
	KIT	343	5978	4907	3561	6416	6254	12766	11316	15189	712	948	34	301
CD117, Kit receptor	CHGA	53	6167	8902	3333	8206	7408	7950	10353	35913	9726	4378	1249	2364
	CHGB	22	1615	2152	1516	1456	1242	2365	1362	4151	4656	1833	32	24
Dopa decaroxylase	DDC	2441	31606	21515	25536	34434	25400	1498	4234	9297	85	334	134	286
Gamma (Neural) Enolase	ENO2	1910	2054	1881	1911	1573	1262	4043	4737	11133	6394	1824	2472	2241
GDNF family receptor alpha 1	GFRA1	9	263	29	37	146	133	90	4	6	77	47	9	4
CD56	NCAM1	82	551	875	801	999	727	2618	2519	3296	361	2899	425	106
	UCHL1	415	16115	13163	12633	22712	19977	13313	9749	25333	7789	4249	122	8251
Proopiomelanocortin	POMC	94	751	427	590	750	657	66	560	5204	68	84	213	160
Somatostatin	SST	67	27000	1316	10013	5869	12139	35	9	62	1992	3613	90	28
Somatostatin Receptor 5	SSTR5	613	733	906	875	636	633	907	344	622	916	662	918	401
Synaptophysin	SYP	19	15	15	9	17	15	27	53	91	49	19	11	19
TTF1	NKX2-1	18	3137	3508	2734	5180	3315	1258	3585	2229	27	311	85	8

FIG. 7B

Average Normalized Intensity Values for Selected Genes in the Notch Pathway and ASCL1

	Median (48 samples)	Cluster C				Cluster D				Cluster G			
		KD66	LU50(LCNEC)	OV26	LU102(SCLC)	LU37(LCNEC)	LU86(SCLC)	LU64(SCLC)	LU95(SCLC)	LU49(SCC)	LU85(SCC)		
ASCL1	9	6589	8238	9382	12169	9664	11	3390	10298	8	5		
DLL1	51	348	565	406	497	179	218	98	514	29	120		
DLL3	350	4584	3985	6232	5884	5233	1686	3137	5814	601	492		
DLL4	614	601	445	592	301	280	763	198	673	357	469		
HES1	670	128	129	160	92	82	551	137	335	2665	2024		
HES6	117	196	361	481	416	279	5456	2716	3535	28	33		
HEY1	89	86	101	116	103	77	1660	680	2502	3776	231		
HEYL	87	157	132	128	148	132	2645	102	267	333	80		
JAG1	630	159	114	110	95	111	743	521	311	9131	678		
JAG2	125	335	529	398	420	247	324	513	611	159	153		
NOTCH1	666	34	23	41	17	14	1039	381	202	4720	438		
NOTCH2	26	6	11	12	16	12	105	11	1	37	5		
NOTCH3	101	13	27	91	81	72	302	37	136	1474	322		
NOTCH4	14	6	7	13	9	5	14	15	69	14	7		
RBPJ	1289	1891	2255	1933	2717	2278	4502	2678	5167	1226	1029		

EP 3 338 793 A1

FIG. 7C

Expression of HES6 mRNA in
Neuroendocrine Tumors Derived by qRT-PCR




FIG. 7D

Levels of Selected Genes Linked To Neural Phenotypes

			Cluster C				Cluster D				Cluster E		Cluster G	
	Gene Symbol	Median (48 samples)	KD66	LU50 (LCNEC)	OV26	LU102 (SCLC)	LU37 (LCNEC)	LU86 (SCLC)	LU64 (SCLC)	LU95 (SCLC)	EM6	LU100 (SCLC)	LU49 (SCLC)	LU85 (SCLC)
Achaete-scute complex homolog 1	ASCL1	9	6589	8238	9382	9664	9664	11	3390	1079	14	261	8	5
	BEX1	62	2393	4403	5349	5387	5653	1574	1174	2139	1474	81	764	50
	NGFRAP1	8484	3367	4346	5975	5977	5222	2108	3743	6528	2113	197	511	406
	BEX4	36	8	8	36	14	6	2108	3743	6528	361	2899	425	106
CD56	NCAM1	82	551	875	801	999	727	2618	2519	3296	14	108	18	3
Claudin 11	CLDN11	7	670	739	2374	5820	3373	548	126	213	3736	1026	5283	2310
Cytokeratin 10	KRT10	2624	3096	2503	2855	5373	5057	2731	1642	4621	63	304	5283	5462
Cytokeratin 17	KRT17	5858	5436	7664	9874	1127	8933	2600	3701	5971	877	816	2060	2755
Cytokeratin 19	KRT19	19604	2081	2432	3214	2435	2532	445	317	7074	12	54	18	600
Cytokeratin 7	KRT7	123	249	348	396	515	312	23	9	333	146	162	29	120
Delta-like 1	DLL1	51	348	565	406	497	179	218	98	514	758	676	601	492
Delta-like 3	DLL3	350	4584	3985	6332	5864	5233	1686	3137	5814	118	528	2665	2024
Hairy and enhancer of split 1	HES1	670	128	129	160	92	82	551	137	335	1634	694	28	33
Hairy and enhancer of split 6	HES6	117	196	361	481	416	279	5485	2716	3535	121	672	34	26
Isl1	ISL1	48	18	23	22	33	16	1807	6671	1067	6243	1663	161	29
LIM homeobox 1	LHX2	21	9	12	29	8	1	6949	6671	1067	4917	1475	6	2
Neurogenic differentiation 1	NEUROD1	8	44	11	35	62	91	331	135	799	3019	450	61	65
Neurogenic differentiation 2	NEUROD2	59	49	40	52	32	25	808	231	416	1311	1619	5191	659
Neuronal cell adhesion molecule	NRCAM	66	721	797	870	1156	1409	3981	4389	4579	63	201	0	247
NK2 homeobox 2	NKX2-2	7	4	7	7	10	4	3725	206	1168	182	9992	72	271
Rhombotin-3	LMO3	145	6875	6571	7101	6912	6886	45	2252	2717	147	935	42	20
Secretogagin	SCGN	77	1950	2620	2468	1701	3052	1164	9386	2717	1002	3050	108	51
Secretogranin II	SCG2	23	445	299	423	1122	1265	37	693	4085	5935	3721	54	19
Secretogranin III	SCG3	26	1552	1961	1723	1870	1892	2331	3164	1011	2769	3426	5124	957
Semaphorin 4C	SEMA4C	1647	941	651	781	1123	742	9648	5482	6341	512	432	1106	107
Semaphorin 6C	SEMA6C	118	323	318	279	241	220	1313	1859	619	135	690	51	61
Seizure related 6 homolog	SEZ6	60	153	143	135	75	86	1087	435	2324	108	114	18	119
Seizure related 6 homolog like	SEZ6L	113	43	25	53	42	26	48	64	116	3793	4933	1491	3247
Seizure related 6 homolog like2	SEZ6L2	1464	1136	1667	1852	1336	1089	2317	3026	5631	10	1880	11	17
SRY (sex determining region Y): box 11	SOX11	13	426	360	361	597	768	3754	6124	981	0	2937	1376	348
SRY (sex determining region Y): box 2	SOX2	8	171	175	270	396	312	2816	74	3267	618	1612	6451	1367
SRY (sex determining region Y): box 4	SOX4	5404	5693	5641	4070	3672	5240	1112	921	565	2775	2488	37	27
SRY (sex determining region Y): box 4	SNAP25	10	1037	1040	1027	1108	1040	1150	1061	1112	27	311	11	10
Synaptosomal-associated protein, 25kDa	SYT4	26	1418	1237	2115	3958	3076	1258	3585	2229	697	279	85	8
Synaptotagmin IV	NKX2-1	18	3137	3508	2734	5180	3315	7883	5585	2940	93	227	93	227
TTF-1														
Zic family member 2	ZIC2	163	25	18	29	10	5							

FIG. 7E

Average Normalized Intensity
Values For SEZ6 Obtained Via Microarray Analysis




FIG. 7F

Expression of SEZ6 mRNA Transcripts in Selected Neuroendocrine NTX Tumor Lines Determined by qRT-PCR




FIG. 8A

Expression of SEZ6 mRNA Transcripts in
NTX Tumor Lines Determined by qRT-PCR




FIG. 8B

Absolute and Normalized SEZ6 mRNA Expression in Various Whole Tumor Samples Using qRT-PCR




FIG. 9A




FIG. 9B

Protein Sequences of Exemplary Mouse SEZ6 Modulator Light Chain Variable Regions

10A – LC AA SEQS (incl. Humanized)
 10B – HC AA SEQS (incl. Humanized)
 10C – Full length LC and HC AA SEQS of hSC17.200 and hSC17.200vL1
 Sequence listing – NA SEQS (including Humanized)

FIG. 10

Protein Sequences of Exemplary Murine SEZ6 Modulator Light Chain Variable Regions

Name	FR1	CDRL1	FR2	CDRL2	FR3	CDRL3	FR4	SEQ ID NO
SC17.1	QIVLTQSPALMSASPGEKVSLTIC	SAMSTVSEFMY	WYQQKPKRSSPTPWLY	LTSNLA	GVPARFSGSGSGTSYSLTISSWEAEDAATYYC	QQWSSNSPIT	FGAGTKLELK	20
SC17.2	DIWMSQSPSSLAVSVGEKVTMSC	KSSQSLLYSSNQKSYLA	WYQQKPGQSPKLLIY	WASTRES	GVPDRFTGSGSGTDFTLTISSVQAEDLAVYYC	KQSYNLRT	FGGGTKLEIK	22
SC17.9	DIWMSQSPSSLAVSVGEKVTMSC	KSSQSLLYSSNQKNYLA	WYQQKPGQSPKLLIY	WASTRES	GVPDRFTGSGSGTDFTLTISSVKAEDLAVYFC	QQYYNYPYT	FGGGTKLK	24
SC17.16	DIQMTQSPASLSVSVGETVTTIC	RASAINISNLV	WYQQKQKQSPQLLVY	AATNLAD	GVPDRFSGSGSGTQYSLKINSIQSEDFGNYYC	QHFNGTPT	FGGGTKLEIK	26
SC17.38	DIWVTQSPASLAVSLGQRATISC	RASESVEYVGTSLMQ	WFQQKPGQSPKLLIY	AASNVES	GVPARFSGSGSGTDFSLNIHPVEEDDIAMVFC	QQDRKVPMT	FGGGTKLEIK	28
SC17.3	DIWMSQSPSSLAVSVGEKVTMSC	KSSQSLLYSSNQKNYLA	WYQQKPGQSPKLLIY	WASTRES	GVPDRFTGSGSGTDFTLTISSVKAEDLAVYYC	QQYYSYPT	FGGGTKLEIK	30
SC17.4	DIKMTQSPSSMYASLGERVTTIC	KASQDINSYLT	WFQQKPGKSPKTLIY	RAIRLID	GVPDRFSGSGSGQDYSLTISSLDYEDMGYYC	LQYDDFPMT	FGGGTKLEIK	32
SC17.8	DVWMTQTPLSLPVSLGDAQASISC	RSSQSLVHSNGDTYLIH	WYLLQKPGQSPKLLIY	KVSNRFS	GVPDRFSGSGSGTDFTLKISRVEAEDLGLYFC	SQSTLIIPYT	FGGGTKLDIK	34
SC17.10	DIWMSQSPSSLAVSVGEKVTMSC	KSSQSLLYSSNQKNYLA	WYQQKPGQSPKLLIY	WASTRES	GVPDRFTGSGSGTDFTLTISSVKTEDLALYYC	QQYYWFPYT	FGGGTKLEIK	36
SC17.11	ENIVLTQSPALMSASPGEKVTMTIC	RASSSVSSSYLH	WYQQKSGASPKLWLY	STSNLAS	GVPARFSGSGSGTSYSLTISSVEAEDAATYYC	QQYSDYPPFT	FGSGTKLVIK	38
SC17.14	DVLMTQTPLSLPVSLGDAQASISC	RSSQSLVHSNGNTYLE	WFLLQKPGQSPKLLIY	KVSNRFS	GVPDRFSGSGSGTDFTLKISRVEAEDLGYYC	FQGSHPVYT	FGGGTKLEIK	40
SC17.15	QIVLTQSPALMSASPGEKVTMTIC	SASSSVNMY	WYQQKPKRSSPKPWLY	LTSNLA	GVPDRFSGSGSGTSYSLTISSWEAEDAATYYC	QQWSSNPPT	FGSGTKLELK	42
SC17.17	QIVLTQSPALMSASPGEKVTMTIC	SASSSVSYMH	WYQQKSGTSPKRWLY	DTSKLPS	GVPARFSGSGSGTSYSLTISSWEAEDAATYYC	QQWSSTPPT	FGAGTKLELK	44
SC17.18	DIQMTQSSSYLSVSLGGRVTITC	KASDHINWLA	WYQQKPGNAPRLLIS	GATSLET	GVPDRFSGSGSGKDYTLTSLTQTEDVATYYC	QQYWSIPLT	FGAGTKLELK	46
SC17.19	DIVLTQSPASLAVSLGQRAAISC	KPSQSDVDYDGDYMN	WYQQKPGQSPKLLIY	AASNLES	GVPARFSGSGSGTDFTLNIHPVEEEDAATYYC	HQINDDPMT	FGGGTKLK	48
SC17.22	DVWLTQTPLSLPVSLGDAQASISC	RSSQSLVHSNGRHTYLG	WYLLQKPGQSLKLLIY	GVSNRFS	GVPDRFSGSGSGTDFTLKISRVEAEDMGVYYC	FQGTHVPPT	FGSGTKLEIK	50

FIG. 10A

Protein Sequences of Exemplary Murine SEZ6 Modulator Light Chain Variable Regions

Name	FR1	CDRL1	FR2	CDRL2	FR3	CDRL3	FR4	SEQ ID NO
SC17.24	DI VNSQSPSSLAVSVGEKVTMSC	KSSQSLLYSSINQKSYLA	WYQQKPGQSPKLLIY	WASTRES	GVPDRFTGSGSGTDFTLTISSVQAEDLAVYVC	KQSYNLRT	FGGGTKLEIK	52
SC17.27	DV VMTQTPLTILSVTIGQPASISC	KSSQSLLIESDGKTYLN	WLLQRPGQSPKRLIY	LVSKLDS	GVPDRFTGSGSGTDFTLKISRVEAEDLGVYVC	WQGIQHPRT	FGGGTKLEIK	54
SC17.28	DILLTQSPAILLSVSPGEGVSFSC	RASQSIGTSTIH	WYQQRTINGSRLLIK	VASESIS	GIPSRFSGSGSGTDFTLRINSLSESDIADYVC	QQSNSWPLT	FGAGTKLELK	56
SC17.29	DIVMTQSHKFMSTSVGDRVSIIC	KASQDVGTDA	WYQQKPGQSPKLLIY	WASTRHT	GVPDRFTGSGSGTDFTLTISNVQSEDLDYFC	QQYSSYPYT	FGGGTKLEIK	58
SC17.30	ENVL TQSPALIVSASPGEKVTMTC	RASSSVTSSYLH	WYQQKSGASPKLWIY	STSNLAS	GVPARFSGSASGTSYSLTISSVEAEDAATYYC	QQYSGYPLT	FGAGTKLELK	60
SC17.32	DIQMTQSPASLSASVGETVTMTC	RASGNIIHIVLV	WYQQKQKQSPQLLVY	NAKTLAD	GVPDRFSGSGSGTQVSLKINSIQPEDFGSYVC	QHFMSTPPT	FGGGTKLEIK	62
SC17.34	DIKMTQSPSPMYASLGERTVITC	KASQDINSYLS	WFQQKPGKSPKTLIY	RAMRLVD	GVPDRFSGSGSGGQDYSLTISLSEVEDIMGYYC	LQYDEF PPT	FGGGTKLEIK	64
SC17.35	ENVL TQSPALIVSASPGEKVTLTC	RASSSMSSSYLH	WYQQKSGASPKLWIY	STSNLAS	GVPARFSGSGSGTSYSLTISSVEAEDAATYYC	QQYSAYPFT	FGSGTKLEIK	66
SC17.36	QIVLTQSPALMSASPGEKVTMTC	SASSSVSYMY	WYQQKPRSPKPIIY	LTSNLAS	GVPARFSGSGSGTSYSLTISSWEAEDAATYYC	QQWSSNPPT	FGGGTKLEIK	68
SC17.39	DVLMTQTPLSLPVSLGDAQASISC	RSSQSLVHRNGNTYFH	WYLLQKPGQSPKLLIY	KVSNRFS	GVPDRFSGSGSGTDFTLKISRVEAEDLGVYFC	SQSYTVPMIT	FGGGTKLEIK	70
SC17.40	DVVMTQTPLSRPVTLGDAQASISC	RSSQSLVHSNGNTYLV	WYLLQKPGQSPKLLIY	KVSNRFS	GVPDRFSGSGSGTDFTLKISRVEAEDLGVYFC	SQNTHVPFT	FGSGTKLEIK	72
SC17.41	QIVLTQSPALMSASPGEKVTMTC	SASSSVSYMY	WYQQKPRSPKPIIY	LTSNLAS	GVPTRFSGSGSGTSYSLTISSWGAEDAATYYC	QQWNTNPPT	FGAGTKLELK	74
SC17.42	ENVL TQSPALIVSASPGEKVTMTC	SASSSVNYMY	WYQQKSSTSPKLWIY	DTSKLTS	GVPGRFSGSGSGMSYSLTISSWEAEDVATYYC	FQSGGYPLT	FGSGTKLEIK	76
SC17.45	ENVL TQSPALIVSASPGEKVTMTC	SASSSVNYMY	WYQQKSSTSPKLWIY	DTSKLTS	GVPGRFSGSGSGMSYSLTISSWEAEDVATYYC	FQSGGYPLT	FGSGTKLEIK	78
SC17.46	STVMTQTPKFLLVSAQRVITTC	KASQSVNIDVA	WYQQKPGQSPKLLIY	YASNRYT	GVPDRITGSGVGTDTITTTISTVQAEDLAVYFC	QQDYSSPRT	FGGGTKLEIK	80
SC17.47	QIVLTQSPALIVSASPGEKVSMTC	SASSSVSYMYH	WYQQKSGTSPKRWIY	DTSKLAS	GVPARFSGSGSGSSYSLTISSWEAEDAATYYC	QQWSSTPPT	FGGGTKLEIK	82

FIG. 10A Cont.

Protein Sequences of Exemplary Murine SEZ6 Modulator Light Chain Variable Regions

Name	FR1	CDRL1	FR2	CDRL2	FR3	CDRL3	FR4	SEQ ID NO
SC17.49	DVWMTQTPLTILSVTIQGPASISC	KSSQSLLES DGKTYLN	WLLQRPQSPKRLIY	LVSKLDS	GVPDRFTGSGSGTDFTLKISRVEAEDLGVYVC	WQGIQHPRT	FGGGTKLEIK	84
SC17.50	DIVLTQSPASLAASLGQRATISC	RASQSVSTSSSYVMH	WYQQKPGQPPKLLIK	YASNLES	GVPARFSGSGSGTDFTLNHPVEEEDTATYYC	QHSMELPMT	FGGGTKLEIK	86
SC17.53	DIVLTQSPASLAASLGQRATISC	RASQSVSTSSSYVMH	WYQQKPGHPPKLLIR	YASNLES	GVPARFSGSGSGTDFTLNHPVEEEDTATYYC	QHSMELPVT	FGGGTKLEIK	88
SC17.54	DVVLTTQTPLTILSVTIQGPASISC	KSSQSLLYSDGKTYLN	WLLQRPQSPKRLIY	LVSKLDS	GVPDRFTGSGSGTDFTLKISRVEAEDLGLYYC	WQGTHFPMT	FGGGTKLEIK	90
SC17.56	DIVWSQSPSSLAVSVGEKVTMTC	KSSQSLLYSSINQKNYLA	WYQQKPGQSPKLLIY	WASTRES	GVPDRFTGSGSGTDFTLTISVKAEDLANVFC	QQYVNYPYT	FGGGTKLEIK	92
SC17.57	QIVLTQSPAIMASLGERVTMTC	TASSSVSSSYLH	WYQQKPGQSPKLLIY	STSNLAS	GVPDRFSGSGSGTSYSLTISSWEAEDAATYYC	HQYHRSPT	FGGGTKLEIK	94
SC17.59	DIQMTQSPASLSASVGETVTITC	RASGNLHNYLA	WYQQKQKSPQLLVY	NAKTLAD	GVPSRFSGSGSGTQYSLKINSLQPEDFGTYFC	QHFWSTPPT	FGGGTKLEIK	96
SC17.61	QIVLTQSPAIMASVGEKVTISC	SASSSVSYIY	WYQQKPGQSPKLLIY	RTSNLAS	GVPARFSGSGSGTSYSLTISSWEAEDAATYYC	QQYHSYPMT	FGGGTKLEIK	98
SC17.63	SIWMTQTPKFLVVSAGDRAITC	KASQSVSHDVA	WYQQKPGQSPPTLLIS	YASNRVT	GVPDRFTGSGYGTDFTFITSTVQAEDLANVFC	QQGYSSPPT	FGGGTKLEIK	100
SC17.71	DIQMTQSPASLSASVGETVTIAC	RASGNLHNYLT	WYQQKQKSPQLLVY	NAKTLAV	GVPSRFSGSGSGTQYSLKINSLQPEDFGSYVC	QHFWNTPPT	FGGGTKLEIK	102
SC17.72	DIQMTQTTSLSLASLGURVTISC	SASQGLSNYLN	WYQQKPDGTVKLLIY	VTSSLHS	GVPSKFSGSGSGTDYSLTISNLEPEDLATYYC	QQYSKLPYT	FGGGTKLEIK	104
SC17.74	DIQMTQSSSYLSVSLGGRVTITC	KASDHINWLA	WYQQKPGNAPRLLIS	GATSLET	GVPSRFSGSGSGKDYTLTSLTQTEDVATYYC	QQYWSPTPT	FGAGTKLELK	106
SC17.76	DIVITQDELSPVTSGESVSISC	RSSKSLLYKDGKTYLN	WFLQRPQSPQLLIY	LMSTRAS	GVSDRFSGSGSGTDFTLEISRKAEDGVNYVC	QQLVVEPRT	FGGGTKLEIK	108
SC17.77	DIQMTQSPASLSASVGETVTITC	RASGNLHNYLA	WYQQKQKSPQLLVY	NAKALAD	GVPSRFSGSGSGTQYSLKINSLQPEDFGSYVC	QHFWSTPPT	FGGGTKLEIK	110
SC17.79	DIQMTQSPASLSASVGETVTITC	RASGNLHNYLA	WYQQKQKSPQLLVY	NAKTLAD	GVPSRFSGSGSGTQYSLRINSLQPEDFGSYVC	QHFWSTPPT	FGGGTKLEIK	112
SC17.81	DIVWSQSPSSLTVSVGEKVTISC	KSSQSLLYSTMQKTYLA	WYQQKPGQSPKLLIY	WASTRES	GVPDRFTGSGSGTDFTLAISNVKAEDLANVVC	QQYYSYPYT	FGGGTKLEIK	114

FIG. 10A Cont.

Protein Sequences of Exemplary Murine SEZ6 Modulator Light Chain Variable Regions

Name	FR1	CDRL1	FR2	CDRL2	FR3	CDRL3	FR4	SEQ ID NO
SC17.82	QIVLTQSPAIMASLGEIITLTC	SASSSVSYMH	WYQKPGQSPKLLIY	STSNLAS	GVPSRFSGSGSGTFYSLTISSVEAEDAAYYC	HQMSFT	FGSGTKLEIK	116
SC17.84	QIVLTQSPAIMASPGKVTMTIC	SASSISYMH	WYQKSGTSPKRWIY	DTSKLAS	GVPARFSGSGSGTSYSLTISSNWEAEDAATYYC	QQMSSTPPT	FGGGTKLEIK	118
SC17.85	DIVMTQAAFSNPVTLTGTSASISC	RSSKSLIHSNGITYLY	WYLQPGQSPQLLIY	QMSNLAS	GVPERFSSGSGSDFTLTISRVEAEDVGVYVC	AQNLHPT	FGGGTKLEIK	120
SC17.87	DVVMQTQTPLSLPVSLGDAQASISC	RSSQSLVHSNGINTYLY	WYLQPGQSPKLLIS	KVSNRFS	GVPDFRSGSGSGTDFTLTISRVEAEDLGVYFC	SQSTHVPPT	FGGSTRLEIK	122
SC17.89	DVLMQTQTPLSLPVSLGDAQASISC	RSSQSLVHSNGINTYLE	WYLQPGQSPKLLIY	KVSNRFS	GVPDFRSGSGSGTDFTLTISRVEAEDLGVYVC	FQGSHPVPT	FGSGTKLEIK	124
SC17.90	DIVMSQSPSSLAVSVGEKVTMSC	KSSQSLLYSSINQKNYLA	WYQKPGQSPKLLIY	WASTRKS	GVPDFRTGSGSGTDFTLTISSVKAEDLAVYVC	HQYYSYPLT	FAAGTKLELK	126
SC17.91	DVVMQTQTPLSLPVSLGDAQASISC	RSSQSLVHSNGINTYLL	WYLQPGQSPKLLIY	KVSNRFS	GVPDFRSGSGSGTDFTLTISRVEAEDLGVYFC	SQSTHVPMT	FGGGTKLEIK	128
SC17.93	DIVMSQSPSSLAVSVGEKVTMTIC	KSSQSLLYSSINQKNYLA	WYQKPGQSPKLLIY	WASTRES	GVPDFRIGSGSGTDFTLTISSVKAEDLAIYYC	QQYVRYPLT	FGAGTKLELK	130
SC17.95	DIQMTQTTSLSLASLGDRVTISC	SASQGINNYLN	WYQKPDGTVTLIIY	YTSSLHS	GVPSRFSGSGSGTDYSLTISSNLEPEDIATYYC	QQYSKLPMT	FGGGTKLEIK	132
SC17.97	DVVMQTQTPLSLPVSLGDAQASISC	RSSQSLVHSNGINTYLY	WYLQPGQSPKLLIY	KVSNRFS	GVPDFRSGSGSGTDFTLTISRVEAEDLGLYFC	SQSTHVPRT	FGGGTKLEIK	134
SC17.99	DIVMSQSPSSLAVSVGEKVTMNC	ESSQSLLYSSINQKNYLA	WYQKPGQSPKLLIY	WASTRDS	GVPDFRTGSGSGTDFTLTISSVRAEDPAVYVC	QQYYSYPLT	FGAGTKLELR	136
SC17.102	QIVLTQSPAIMASPGKVTMTIC	SASSSVSYMH	WYQKSGTSPKRWIY	DTSKLAS	GVPPRFSGRSGSGTSYSLTISSWEAEDAATYYC	QHWSSNPPT	FGAGTKLEMK	138
SC17.114	DVVMQTQSPSLPVSLGDAQASISC	RSSQSLVHSNGINTYLY	WYLQPGQSPKLLIY	KVSSRFS	GVPDFRSGSGSGTDFTLTISRVEAEDLGVYFC	SQSTHVPPT	FGSGTKLEIK	140
SC17.115	DIVMTQSPSLSLPVSLGDAQASISC	RSSQSLVHSNGINTYLY	WYLQPGQSPKLLIY	KVSNRFS	GVPDFRSGSGSGTDFTLTISRVEAEDLGVYFC	SQSTHLPRT	FGGGTKLEIK	142
SC17.120	DIVLTQSPASLAVSLGQRATISC	RASEVDSYGNSTFMH	WYQKPGQPPKVLIIY	RASNLIS	GLPARFSGSGRTDFTLTINPVEDVDATYYC	QQSNEDPYT	FGGGTKLEIK	144
SC17.121	QIVLTQSPAIMASPGKVTMTIC	SASSSVSYMH	WYQKSGTSPKRWIY	DTSKLAS	GVPARFSGSGSGTSYSLTISSMETEDAATYYC	QQWSNPPT	FGSVTKLEIK	146

FIG. 10A Cont.

Protein Sequences of Exemplary Murine SEZ6 Modulator Light Chain Variable Regions

Name	FR1	CDRL1	FR2	CDRL2	FR3	CDRL3	FR4	SEQ ID NO
SC17.122	DIVITQDDLINPVTSGESVSISC	RSSKSLLYKDGKTYLN	WF LQRPQSQPLLII	LMSTRAS	GVSDFRFGSGSGTDFTLEISRKAEDVGYYC	QQLVEYPRT	FGGGTKLEIK	148
SC17.140	QIVLTQSPAIMVASPGKVTMTIC	SASSSVSYMH	WYQQKSGTSPKRWIY	DTSKLAS	GVPARFSGSGSGTSYSLTISSWEAEDAATYYC	QQWSSTPPT	FGSGTKLEIK	150
SC17.151	DIVLTQFPASLAVSLGQRATIPC	RASESVDSYGNSFMH	WFQQKPGQPKPLIIY	RASNLES	EIPARFSGSGSGTDFTLTINPVEADDDVATYYC	QQSHEDPYT	FGGGTKMEIK	152
SC17.156	DVVMTQSPSLPVSIGDQASISC	RSSQSIIVHSNGNTYLE	WYLQKPGQSPKLLIIY	KVSNRFS	GVPDRFSGSGSGTDFTLKISRVEAEDLGVYYC	FQGSHPVPT	FGGGTKLEIK	154
SC17.161	DIVMSQSPSSLAVSVGEEKVTMNC	ESSQSLLYNSINQKNYLA	WYQQKPGQSPKLLIIY	WASTRDS	GVPDRFTGSGSGTDFTLTISSVRADDPAVYYC	QQYFNYPLT	FGAGTKLELK	156
SC17.166	QIVLTQSPAIMVASPGKVTMTIC	SASSSVSYMH	WYQQKSGTSPKRWIY	DTSKLAS	GVPARFSGSGSGTSYSLTISSWEAEDAATYYC	QQWSSTPPT	FGGGTKLEIK	158
SC17.187	DIKMTQSPSPMYASLGERTVLTIC	KASQDINSYLS	WFQQKPGKSPETLLIY	RAMRLID	GVPSRFSGSGSGGQDYSLTISLSEYEDMGIIYYC	LQYDEFPT	FGGGTKLEIK	160
SC17.191	QIVLTQSPAIMVASPGKVTMTIC	SASSSVSYMH	WYQQKSGTSPKRWIY	DTSKLAS	GVPARFTGSGSGTSYSLTISSWEAEDAATYYC	QQWSSTPPT	FGAGTKLELK	162
SC17.193	DLVLTQSPASLAVSLGQRATISC	RASESVSTSGYSYMH	WYQQKPGQPKPLIIY	LASNLES	GVPARFSGSGSGTDFTLNHPVVEEEDATYYC	QHSRELPTY	FGGGTKLEIK	164
SC17.199	DIVLTQSPASLAVSLGQRATISC	RASESVDSYGNSFMH	WYQQKPGQPKPLIIY	RASNLES	GIIPARFSGSGSRDFTLTINPVEADDDVATYYC	QQSNEDPYT	FGGGTKLEIK	166
SC17.200	DIVLTQSPASLAVSLGQRATIFC	RASQSDYDNGISYMH	WFQQKPGQPKPLIIY	AASNVQS	GIIPARFSGSGSGTDFTLNHPVVEEEDAAATFYC	QQSLIEDPPT	FGGGTKLEIK	168

FIG. 10A Cont.

Protein Sequences of Exemplary Humanized SEZ6 Modulator Light Chain Variable Regions

Name	FR1	CDRL1	FR2	CDRL2	FR3	CDRL3	FR4	SEQ ID NO
hSC17.16	DIQMTQSPSSLSASVGDRTITC	RASANINLV	WYQQKPGKAPKLLIY	AATNLAD	GVPSRFSGSGSGTDFTLTISSLQPEDFATYYC	QHFMTGTPRT	FGGGTKLEIK	170
hSC17.17	EIVLTQSPATLSLSPGERATLSC	SASSSVSYMH	WYQQKPGQAPRLLIY	DTSKLPS	GIPARFSGSGSGTDFTLTISSLQPEDFATYYC	QQMSSTPPT	FQGTKLEIK	172
hSC17.24	DIVMTQSPDLSAVSLGERATINC	KSSQSLLYSSNQKSYLA	WYQQKPGQPPKLLIY	WASTRES	GVPSRFSGSGSGTDFTLTISSLQAEADVAVYYC	KQSYNLR	FGGGTKVEIK	174
hSC17.28	EIVLTQSPDFQSVTPKKEVTITC	RASQSIGTSH	WYQQKPDQSPKLLIK	YASESIS	GVPSRFSGSGSGTDFTLTINSLEAEADAATYYC	QQSNSWPLT	FQGTKLEIK	176
hSC17.34	DIQMTQSPSSLSASVGDRTITC	KASQDINSVLS	WFQQKPGKAPKSLIY	RAIRLVD	GVPSRFSGSGSGTDFTLTISSLQPEDFATYYC	LQYDEFPPT	FQGTKLEIK	178
hSC17.46	AIQMTQSPSSLSASVGDRTITC	KASQSVNDDVA	WYQQKPGKAPKLLIY	YASNRYT	GVPSRFSGSGSGTDFTLTISSLQPEDFATYFC	QQDYSSPRT	FQGTKLEIK	180
hSC17.151	EIVLTQSPATLSLSPGERATLSC	RASESVDSYGNFMH	WYQQKPGQAPRLLIY	RASNLES	GIPARFSGSGSGTDFTLTISSLQPEDFATYYC	QQSHEDPYT	FQGTKLEIK	182
hSC17.155	DIVMTQSPDLSAVSLGERATINC	KSSQSLLYSSNQKINVLA	WYQQKPGQPPKLLIY	WASTRKS	GVPSRFSGSGSGTDFTLTISSLQAEADVAVYYC	HQYYSYPLT	FQGTKLEIK	184
hSC17.156	DIVMTQTPLSLPVTPGEPASTSC	RSSQSLVHSNGNTYLE	WYLLQKPGQSPQLLIY	KVSNRFS	GVPSRFSGSGSGTDFTLTISRVEAEADVAVYYC	FQSHVPPT	FGGGTKLEIK	186
hSC17.161	DIVMTQSPDLSAVSLGERATINC	ESSQSLLYNSNQKINVLA	WYQQKPGQPPKLLIY	WASTRES	GVPSRFSGSGSGTDFTLTISSLQAEADVAVYYC	QQYFNYPPT	FQGTKLEIK	188
hSC17.200	EIVLTQSPATLSLSPGERATLSC	RASQSVYNGISYMH	WYQQKPGQAPRLLIY	AASNVOQS	GIPARFSGSGSGTDFTLTISSLQPEDFATYYC	QQSIEDPPT	FGGGTKVEIK	190
hSC17.200v1.1	EIVLTQSPATLSLSPGERATLSC	RASQSVYDYGISYMH	WYQQKPGQAPRLLIY	AASNVOQS	GIPARFSGSGSGTDFTLTISSLQPEDFATYYC	QQSIEDPPT	FGGGTKVEIK	192

FIG. 10A Cont.

Protein Sequences of Exemplary Murine SEZ6 Modulator Heavy Chain Variable Regions

Name	FR1	CDRH1	FR2	CDRH2	FR3	CDRH3	FR4	SEQ ID NO
SC17.1	DVQLQDSGPGLVKPSQSLSVTCTVTGYSIT	WGYYIN	WIRQFPGNKLEWMG	NIHNSGGTNYNPSLKS	RISITRDTSKNQVFLQLNSVTTEDTATYYCAT	TNIHDFDY	WGQGTTLTVSS	21
SC17.2	QVQLQDSDAELVKPGASVKISCKVSGYTF	DHTIH	WMKQPEQGLEWIG	YIYPRDSTKYNEEFKG	KATLTADKSSSTAYHQLNLSLTSEDSAVYFCAR	SYSNYFDY	WGQGTTLTVSS	23
SC17.9	QVQLQDQPGAEIVRPGASVKLSCKASGYTFT	DYWMN	WVKQRPQGQGLEWIG	AIDPSDSYTSYNPKFKG	KATLTVDTSSSSAYHQLSSLTSEDSAVYFCAR	RGTGKPLVY	WGQGTTLTVSA	25
SC17.16	EVQLQQSGPELWMKPGASVKWSCASGYTFT	DYNMY	WVKQNGKSLIEWIG	EINPNNGGTAYNQKFRG	KATLTVDKSSSTAYMELRSLTSEDSAVYYCAR	YDKGFDY	WGQGTTLTVSS	27
SC17.38	QVTLKESGPGILQPSQTLSLTCSFSGFSLN	TSGMSVG	WVRQPSGRGLEWLA	PIMMIGDKYYNPALKS	RLTISKDTSNNQVFLKIASVWTADTATYFCAR	IRQYYAMDY	WGQGTSLTVSS	29
SC17.3	QVQLQDQPGAEIVKPGASVKLSCKASGYTFP	SYWTH	CVKQRPQGQGLEWIG	VINPNSGRTNWNKFKNI	KATLTVDKSSSTAYHQLSSLTSEDSAVYYCVR	GGTGYTMDY	WGQGTSLTVSS	31
SC17.4	QIQLVQSGPELKKPGETVKISCKASGYTFT	DYSWH	WVKQAPGKGLKWLG	WINTETGEPTYSEDFKG	RFAFLETSASTAYLQINNLLKNETATYFCVK	NKGNFAY	WGQGTTLTVSA	33
SC17.8	QVHLQQSGTEVMKPGASVKISCKATGYTFS	SYWTE	WIKQRP-GHGLEWIG	EILPQSGNTNNEKFKG	KATITADTSSNIAYTQLSSLTSEDSAVYYCAG	GPAAY	WGQGTTLTVSA	35
SC17.10	EVQLQQSGAEIVKPGASVKLSCTASGFNIK	DTYWH	WVKQRP-EQGLEWIG	RIDPANVNTKYDPKFDG	KATITADTSSNTAYLQLSSLTSEDTAVYYCVR	GNVY	WGQGTTLTVSA	37
SC17.11	EVQLQDSGPELVKPGALVMWSCASGYTFT	DYYWH	WVKQSHGQSLEWIG	EVIPYNDETFFYNRKFKD	KATLTVDKSSSTAYMELRSLTSEDSAIYYCAR	RHRYDGFYRIDY	WGQGTSLTVSS	39
SC17.14	EVQLQDSGPGVLVVKPGASVKWSCASGYTIT	DYNWN	WVKQSHGKSLEWIG	VINPYNGNTRYNQWFKG	KATLTVDKSSSTAYMELRSLTSEDSAVYYCTR	WGTTVVGAN	WGQGTTLTVSS	41
SC17.15	DVKLVESGGGLVKLGSLKLSCAASGFTFS	SYAVS	WVRQTP-EKRLEWVA	TITSGGNGTYYPDSVKG	RFTISRDNAKNTLYLQWSSLKSEDTAMYYCAR	RDYYGSSVYMFAY	WGQGTTLTVSA	43
SC17.17	EVQLQDSGPEVMKPGASVKWSCASGYTFT	DYNWH	WVKQNGKSLEWIG	EINPNIGGTGYNQKFKG	KATLTVHKSSSTAYMELRSLTSEDSAVYYCAR	TYSYYSYEFAY	WGQGTTLTVSA	45
SC17.18	QVTLKESGPGILQPSQTLSLTCSFSGFSLS	TSTMVG	WIRQPSGKGLEWLA	DIWMDDSKYYNPALKS	RLTISKDTSNQVFLKITSVDTDATATYYCAR	KGRTARATRGFAV	WGHGTLTVSA	47
SC17.19	DVQLQESGPGLVKPSQSLSVTCTVTGYSIT	SSYTMN	WIRQFPGNKLEWMG	YIHYSGSTNYNPSLRS	RISITRDTSKNQVFLQLNSVTTEDTATYYCAR	SRYYYDAYGFAY	WGQGTTLTVSA	49
SC17.22	QIQWQSGPELKKPGETVKISCKASGYSFT	NYGWN	WVKQAPGKGLKWMG	WINTYTGEPYADDFKG	RFAFLETSASTAYLQINNLLKNEEDMATYFCTR	GYGSSYDAILY	WGQGTSLTVSS	51

FIG. 10B

Protein Sequences of Exemplary Murine SEZ6 Modulator Heavy Chain Variable Regions

Name	FR1	CDRH1	FR2	CDRH2	FR3	CDRH3	FR4	SEQ ID NO
SC17 . 24	QVQLQQSDAELVPGASVKISCKVSGYTFT	DHTIH	WVKQRPEQGLEWIG	YIYPRDGS TKYNEEF KG	KATLTADKSSSTAYMQLNSLTSEDSAVYFCAR	SYSNYFDY	WGQGTTLTVSS	53
SC17 . 27	QVQLQQSGAELVRPGASVTLSCKASGYTFT	DYEMH	WVKQTPVHGLEWIG	GIDPETGGTAYNQKF KG	KATLTADKSSSTAYMELRSLTSEDSAVYFCTR	WFSY	WGPGLVTVSA	55
SC17 . 28	QVHLPPQSRPELVKPGASVKISCKASGYGFT	RSYIH	WVKQRPQGQLEWIG	YISSGSGGTTTAYNQKF KG	KASLTADNPSTAYMHLSSLTSEDSAIYFCAR	GGVRYFDV	WGAGTTTVTVSS	57
SC17 . 29	EVQLQQSGPELMKPGASVKMSCKASGYTFT	DYNMH	WVKQMQCKSLEWIG	EINPHNGGTGYNQKF KG	KATLTVDKSSSTSYMELRSLTSEDSAVYVCAG	GYPAFDY	WGQGTTLTVSS	59
SC17 . 30	EVKLVESEGGGLVQPGSSMKLSCTASGFTFS	DYYMA	WVRQVPEKGLEWIA	NINYDGSSTYYLDLSKS	RFIISRDNAKNIIYLQWSSLSKSEDTATYYCAR	DDYVGSSPSYMYFDV	WGAGTTTVTVSS	61
SC17 . 32	EVKLEESGGGLVQPGGSMKLSCTASGFTFS	NYWMS	WVRQSPKQGLEWIA	EIRLKSNNYATHYAESVKG	RFTISRDDSKSSVFLQWNLRLTDEDTGIYCTR	HYYYAMDY	WGQGTSTVTVSS	63
SC17 . 34	EVQLQQSGPELVKPGSSVKISCKASGYTFT	DYNMD	WVKQSHGKRLWIG	YIYPDINGGAGYNQKF KG	KATLTVDKSSSTAYMELRSLTSEDSAVYCSR	SITTAWFAY	WGQGLVTVSA	65
SC17 . 35	EVQLQQSGPELVKPGALVKMSCKASGYTFT	DYYIH	WVKQSHGKSLEWIG	EINPYNGETFYNQKF KG	KATLTVDKSSSTAYMELRSLTSEDSAVYVCAR	RGMYLTGYAMDY	WGQGTSTVTVSS	67
SC17 . 36	EVQLQESGPSLVKPSQSQSLTCSVTGDSIT	SDYMN	WIRKFPQKKVEYWG	YINYSGYTYNPSLSKS	RISITRDTSKNQYYLQWMSVTSEDTATYYCAR	TSYNNKFLPFAY	WGQGLVTVSA	69
SC17 . 39	EVQLVESGGGLVQPGGSRKLSCAASGFTFS	SYGMH	WVRQAPKQGLEWIA	YISSNDGTIYVADTVRG	RFTISRDNAKNTLFLQWTSLSRSEDTANYVCAR	PSNMYFDY	WGQGTTLTVSS	71
SC17 . 40	QVQLQQPGAEIVRPGASVKLSCKASGYTFT	DYWMN	WVKQRPQGQLEWIG	TIDPSDSYTRYNQKF KG	KATLTVDTSTFSSAYMQLSSLTSEDSAVYFCAS	GGRGFGY	WGQGTPTVTVSV	73
SC17 . 41	DKKLVESEGGGLVKGGLKLSCAASGFTFS	SYAMS	WVRQTPKRLWIA	TISSGGNTYYPDSVKG	RFTISRDNAKNTLYLQWSSLSKSEDTANYVCAR	RDYNGTSVWMEAY	WGQGLVTVS	75
SC17 . 42	DKKLVESEGGGLVRPGGSLKLSCAASGFTFS	RVTMS	WVRQTPKRLWIAA	TINSGGSNTYYPDSVKG	RFTISRDNAKNTLFLQWSSLSKSEDTANYVCNTN	GNH	WGQGTTLTVSS	77
SC17 . 45	QVQLQQPGSVLVRPGDSVKLSCKASGYTFT	SYWMH	WVKQSPQGQLEWIG	EIHPSGTSNYNEKF KG	KATLTVDTSSSTAYVDLSSLTSEDSAVYVCVG	GHYDY	WGQGTTLTVSS	79
SC17 . 46	QVQLQQPGAELVKPGASMKLSCKASGYTFT	SYMIN	WVKQRPQGQLEWIG	NIFPDITTTNYNEKF KS	KATLTVDTSSSTAYMQLSSLTSDSDSAVYVCAR	EYYDGTVDANDY	WGQGTSTVTV	81
SC17 . 47	EVQLQQSGPELVKPGASVKISCKASGY5FT	DYYMR	WVKQSPKQLEWIG	EINPSTGGTTTAYNQMF KA	KATLTVDKSSSTAYMQLKSLTSEDSAVYVCAR	GGYFLYFDY	WGQGTTLTVSS	83

[Type text]

FIG. 10B Cont.

Protein Sequences of Exemplary Murine SEZ6 Modulator Heavy Chain Variable Regions

Name	FR1	CDRH1	FR2	CDRH2	FR3	CDRH3	FR4	SEQ ID NO
SC17.49	QVQLQQSGAELVRPGASVTLSCKASGYTFT	DYEMH	WVKQTPVHGLEWIG	GIDPETGGTAYNQKF KG	KATLTADKSSSTAYMELKSLTSEDSAVYFCTR	WFSY	WGPGTLVTVSA	85
SC17.50	EVQLVESGGGLVKPGGSLKLSCAASGFTFS	DYGMH	WVRQAPKGLEWVA	YISSGRTIYYADTVKG	RFTISRDNKNTLFLQMTSLRSEDTANYFCAR	VYGSYGYFDV	WGTGTTVTSS	87
SC17.53	EVQLQSGPELVKPGASVKISCKASGYTFT	DYNMH	WVKQSHGKRLIEWIG	YIHPYNGSGSYNQKF KR	KATLTVDNSSNTTYMELKSLTSEDSAVYFCAR	SYDYDTWFGY	WQGGTLVTVRA	89
SC17.54	EVKLEESGGGLVQPGGSMKLSVCASGFTFS	NYMIN	WVRQSPKGLEWVA	EIRKSNINYATHYAESVKG	RFTISRDDSKSCVYLQMNILRPEDTGYYCTR	GGY	WQGGTTLTVS	91
SC17.56	QIQLVQSGPELKKPGETVKISCKASGYTFT	NYGMN	WVKQAPGKGLKMWIA	WINTYTGEPTYADDF KG	RFASFLETSASTASLQIINILKNEDTATYFCAR	IGDSSPSDY	WQGGTTLTV	93
SC17.57	QIQLVQSGPELKKPGETVKISCKASDYTFT	DFSIH	WVRQSPKGLKMWG	WINTETGEPTVAEDF KG	RFASFLETSASTAF LQIYNILKNEDSATYFCAR	GRYYGHDYANDY	WQGGTSVTVSS	95
SC17.59	EVKLEESGGGLVQPGGSMKLSVCASGFTFS	NYWMN	WVRQSPKGLEWVA	EIRLKSNNYATHYAESVKG	RFTISRDDSKSSVYLQMNILRAEDTGYYCTR	LWDFAMDY	WQGGTSVTVSS	97
SC17.61	QVTLKESGPGILQP5QTL5LTC5F5GF5L5	TFGMGVG	WIRQPSGKGLEWLA	QIWMDDYKYNPFAL K5	RLTISKDTSKNQVFLKIANVDTADTATYFCAR	IGYYSG55RCWYFDV	WGTGTSVTVSS	99
SC17.63	QVQLQQSDAELVRPGASVKISCKAAGYTFT	DLTIH	WVKQRPEQGLEWIG	YIYPGDSNTKYNEKF KG	KATLTADKSSSTAYMQLMSLTSEDSVWYFCAR	MITPYVFDY	WQGGTTLTV	101
SC17.71	EVKLEESGGGLVQPGGSMKLSVCASGIIFS	NYWMN	WVRQSPKGLEWVA	EIRLKSNNYTHYAESVKG	RFTISRDDSKSSVYLQMNILRAEDTGYYCTR	HYYYAMDY	WQGGTSVTVSS	103
SC17.72	EVQLVESGGGLVKPGGSLKLSCAASGFTFS	SYGMS	WVRQTPEKRLWVA	AINSNGGSTYYPTDKG	RLTISRONGKNTLYLQWSSLRSEDTALYYCVR	DDGYYVFFAY	WQGGTLVTVSA	105
SC17.74	QVQLKQSGPGLVAPSQSLSITCTVSGFSLT	SYGVD	WVRQSPKGLEWLG	VTHGGGSTNYNSAL K5	RLSITKDNKSQVFLKWNLSLQTDDETANYFCAS	GDYDGSLLWFAY	WQGGTLVTVSA	107
SC17.76	EVQLVESGGDLVKPGGSLKLSVCASGFTFS	SYGMS	WVRQTPDKRLWVA	TISSGGGTYYPPDSVKG	RFTVSRDNKNTLYLQWSSLSKSEDTANYYCSR	HGWG	WQGGTLVTVSA	109
SC17.77	EVKLEESGGGLVQPGGSMKLSVCASGFTFS	NYWMN	WVRQSPKGLEWVA	EIRLKSNNYATHYAESVKG	RFTISRDDSKSSVYLQMNILRVEDTALYYCTR	HYDYAMDY	WQGGTSVTVSS	111
SC17.79	EVKLEESGGGLVQPGGSMKLSVCASGFTFS	DYWMN	WVRQSPKGLELVA	EIRLISNNYATHYAESVKG	RFTISRDDSKSSVYLQMNILRAEDTGYYCTR	HYYYALDY	WQGGTSVTVSS	113
SC17.81	EVQLQQSGAELVKPGASVKLSCTASGFNIN	DTYYH	WLKQRPEQGLEWIG	RIDPANWNTKYDPKFQG	KATLTADTSSNTAYLQL5SLTSEDTAVYCGR	GNAY	WQGGTLVTVSA	115

[Type text]

FIG. 10B Cont.

Protein Sequences of Exemplary Murine SEZ6 Modulator Heavy Chain Variable Regions

Name	FR1	CDRH1	FR2	CDRH2	FR3	CDRH3	FR4	SEQ ID NO
SC17.82	EVQLQQSGPELVKPGASVKMSCKASGYTFT	DSYMN	WVKQSHGKSLIEWIG	RVPNNGGASYNHKFKG	KATLTVDKSLSTAYMRLNSLTSEDSAVYYCSR	SGDLYYYAMDY	WGQGTSTVTSS	117
SC17.84	EVQLQQSGPELMKPGASVKMSCKASGYIFT	DYNMH	WVKQHQGKSLIEWIG	EVNPNTGGIGYNQKFKG	KATLTVDKSSSTAYMDRLSLTSEDSAVYYCAR	DGNVCFDY	WGQGTTLTVSS	119
SC17.85	EVQLVESGGDLVKPGGSLKLSCAASGFTFS	NYGMS	WVRQTPDKRLHWYA	TISTGGTYTYPPDSVKG	RFTISRDNAKNTLYLQWSSLKSEDTANYVCVG	QSYSDVVSFAY	WGQGTQVTSA	121
SC17.87	EVQLQQSGAELLKPGASVKLSCTASGLNIK	DYYIH	WVYQRPEQGLEWIG	RIDPESDNTLYDPKFKQG	KASITADTSSNTAYLQLSSLTSEDTAVYCTT	NTPFAY	WGQGTTLTVST	123
SC17.89	QVQLQQSGAELVRPGTTSVKVCKTSGYAFT	NVLIIE	WVKQRPQGQGLEWIG	VINPQSGGTNYNEKFKV	KATLTADKSSSTAYMQLTSLTSDSDSAVYFCTR	RDGYFFPWIFAY	WGQGTTLTVSA	125
SC17.90	QVQLQQPGSVLVRPGASVKLSCKASGYTFT	SYWMH	WVKQRPQGQGLEWIG	EIHPNNGSTNYNEKFKG	KATLTVDTSSSTAYVDLSLTSSEDSAVYYCAR	WTLFTY	WGQGTTLTVSA	127
SC17.91	EVQLVESGGGLVKPGGSRKLSCAASGFTFS	DYGMH	WVRQAPEKGLEWYA	YISRGSTTIHYADTVKG	RFTISRDNAKNTLFLQWTSLSRSEDTANYVCAR	PFNWFDV	WGAGTTVTSS	129
SC17.93	QVQLQQPGAELVKPGASVMSLCKASGYTFT	SYWMH	WVKQRPQGQGLEWIG	VINPRNGRNMYNEKFKT	KATLTVDKSSSTAYMQLSSPTSEDSAVYYCAR	EDYDGGDYANDY	WGQGTSTVTSS	131
SC17.95	EVELQQSGPELVKPGASVKLSCKTSGNTYT	EYTMQ	WVKLSHGKSLIEWIG	GINPNNGITSYNQKFKG	KATLTVDKSSSTAYMELKSLKSEDSAVYYCAR	AGLGNVWAMDY	WGQGASVTSS	133
SC17.97	QVQLPQSGAELAKPGASVKISCKASGFTFT	SYWMH	WVKQRPQGQGLEWIG	VINPSTDYTEYNQKFKD	KATLTADKSSSTAYMQLGSLTSEDSAVYYCAR	SSYGSSPFDY	WGQGSTLTVSS	135
SC17.99	EVKLEESGGGLVQPGGSMKLSCAASGFTFS	DAWMD	WVRSPEKGLEWYA	EIRSKANNHATYYAESVKG	RFTISRDDKSSAYLQWMSLRAEDTGIYYCVS	TGTSY	WGQGTTLTVSA	137
SC17.102	EVQLQQSGPELMKPGASVKMSCKASGDTFT	DYNIH	WVKQHQGKSLIEWIG	EVNPNIIGGIGYNQKFKG	KATLTVDKSSSTAYMELKSLTSEDSAVYYCAM	GRWYFDV	WGAGTTVTSS	139
SC17.114	EVQLQQSGPENVKPGASVKISCKASGYTFT	DYYMH	WVKQSHGKSLIEWIG	RVNTNNGGTSYDQKFEK	KATLTVDKSSSTAYMELNSLTSEDSAVYYCVI	PAMFAY	WGQGTTLTVSA	141
SC17.115	QVQLQQSGSVLVRPGASVKLSCKASGYTFT	SYWMH	WVKQRPQGQGLEWIG	EIHPNSGNTNYNEKFKG	KATLTVDTSSSTAYVDLSLTSSEDSAVYYCAG	QNVYD	WGQGTTLTVSS	143
SC17.120	EVQLQQSGTVLARPASVKMSCKASGYTFT	SYWMH	WVKQRPQGQGLEWIG	AFYPGNSGTYYNQKFKD	KAKLTAVTSASTAYMELSSLTNEDSAVYYCSR	SGSGRFAY	WGQGTTLTVSA	145
SC17.121	EVQLQQSGPELMKPGASVKMSCKASGYTFT	DHNIIH	WVKQHQGKSLIEWIG	EINPNTGGTGYNQKFKQ	KATMTVDKSSSTAYMELKSLTSEDSAVYYCVR	GLYFFDY	WGQGTTLTVSS	147

[Type text]

FIG. 10B Cont.

Protein Sequences of Exemplary Murine SEZ6 Modulator Heavy Chain Variable Regions

Name	FR1	CDRH1	FR2	CDRH2	FR3	CDRH3	FR4	SEQ ID NO
SC17.122	EVHIVESGDLVPGGSLKLSAASGFTFS	SYGMS	WVRQTPDKRLEWIA	TISSGGTYTYYPDSVKG	RFTISRDNAKNTLYLQMSLSKSEDTAMYYCSR	HGWG	WGQGTLLVTVSA	149
SC17.140	EVQLQQSGPELMPGASVKMSCKASGYTFT	DYNMH	WVKQNGKSLIEWIG	EINPTNGGTGYNQKF KG	KATLTVDKFSSTAFIELRSLTSEDSAIYYCTR	GGYDHYHYFDV	WGAGTTVTVSS	151
SC17.151	EVQLQQSGTVLARPASVKMSCKASGYTFT	SYMMH	WVKQRPQGGLIEWIG	AIYPGKNDDTTYNQKF KG	KAKLTAVTASSTLYMELSSLTNEEDSAVYYCTR	SGKGYFAY	WGQGTLLVTVSA	153
SC17.156	QVTLKESGPGILQPSQTLSTCSFSGFSL	TSGMGVS	WIRKTSKGLEWLA	HLFWDDDKWNPSLKS	RLTISKATSSNQVFLILTSDVDTADTATYYCAT	FYGLYFAY	WGQGTLLTVSS	155
SC17.161	EVKLEESGGGLVQPGGSMKLSAASGFTFS	DAWMD	WVRQSPKGLIEWIA	EIRSKPINHATYYAESVKG	RFTISRDDSKSSAYLQMNLSLRAEDTGIYYCVS	TGTSY	WGQGTLLVTVSA	157
SC17.166	EVQLQQSGPELMPGASVKMSCKASGYIFT	DYNMH	WVKQNGKSLIEWIG	EYNPTNGGIGYNQKF KG	KATLTVDKSSSTAYMDLRSLTSEDSAVYYCAR	DGNYCFDY	WGQGTLLTVSS	159
SC17.187	EVHLLQSGPELVNPGSSVKISCKAAGYTFT	DYNMD	WVKQSHGKRLEWIG	NIYPNNGGAGYNQNF KD	KATLTVDKSSSTAYMELRSLTSEDSAVYYCAR	SITAAWFAY	WGQGTLLVTVSA	161
SC17.191	EVQLQQSGPELMPGASVKMSCKASGYTFT	DYNMH	WVKQNGKSLIEWIG	EINPTNGGTGYNQKF KD	KATLTVDKSSSTAYMELRSLTSEDSAVYYCAR	IPSLRRYYFDY	WGQGTLLTVSS	163
SC17.193	QVTLKESGPGILQPSQTLSTCSFSGFSLI	TYGLGVG	WIRQPSKGLEWLA	HLWMMDKYYNTALKS	RLTISKOTSNNQVFLKIANVDTADTATYYCAR	MVYYDYDGGFAY	WGQGTLLVTVSA	165
SC17.199	EVQLQQSGTVLARPASVKMSCKASGYTFT	SYMMH	WVKQRPQGGLIEWIG	AIYPGNSDTSYNHKF KG	KAKLTAVTASSTAYMELSSLTNEEDSAVYYCTR	SGTGMFAY	WGQGTLLTVSS	167
SC17.200	QVQLQQSGPELVKPGASVKISCKASGYAFS	SSWIN	WVKQRPQGGLIEWIG	RIYPGEEDTNSGMFEG	KATLTADKSSTTAYMQLSSLTSDSAVYFCTR	GLVMDY	WGQGTALLTVSS	169

[Type text]

FIG. 10B Cont.

Protein Sequences of Exemplary Humanized SEZ6 Modulator Heavy Chain Variable Regions

Name	FR1	CDRH1	FR2	CDRH2	FR3	CDRH3	FR4	SEQ ID NO
hSC17 .16	QVQLVQ5GAEVKKPGASVKVSKKASGYTFT	DYNMY	WVRQAPQGQLEIMMG	EINPNNGGTAYNQKFRG	KVTMTSDTSISTAYMELSR LRSDDTAVYYCAR	YDKGFDY	WGQGTITVTVSS	171
hSC17 .17	QVQLVQ5GAEVKKPGASVKVSKKASGYTFT	DYNMH	WVRQAPQGQLEIMMG	EINPNJGGTGYNQKFKG	RVTMTSDTSISTAYMELSR LRSDDTAVYYCAR	TYSYYSYEFAY	WGQGTITVTVSS	173
hSC17 .24	EVQLVQ5GAEVKKPGATVKISCKVSGYTFT	DHTIH	WVRQAPGKGLEIMLG	YIYPRDGSITKYNEEFKG	RVTITADTSDTAYMELSS LRSDDTAVYYCAR	SYSINFDY	WGQGTITVTVSS	175
hSC17 .28	QVQLVQ5GAEVKKPGASVKVSKKASGYTFT	RSYIH	WVRQAPQGQLEIMMG	YISSGSGGTTYNQKFKG	RVTSTSDTSISTAYMELSR LRSDDTAVYYCAR	GGVRVFDV	WGQGTITVTVSS	177
hSC17 .34	QVQLVQ5GAEVKKPGASVKVSKKASGYTFT	DYNMD	WVRQAPGQRLIMJG	YIYPDNGGAGYNQKFKG	RVTITVSDTSASTAYMELSS LRSDDTAVYYCSR	SITTAMFAY	WGQGTITVTVSS	179
hSC17 .46	QVQLVQ5GAEVKKPGASVKVSKKASGYTFT	SYMIN	WVRQAPQGQLEIMJG	NIFPDTTTTNNEKFKG	RVTLTSDTSISTAYMELSR LRSDDTAVYYCAR	EYYDGTVDAMDY	WGQGTITVTVSS	181
hSC17 .151	QVQLVQ5GAEVKKPGASVKVSKKASGYTFT	SYMMH	WVRQAPQGQLEIMMG	ALYPGKSDTTYNQKFKG	RVTMTSDTSISTAYMELSS LRSDDTAVYYCAR	SGKGYFAY	WGQGTITVTVSS	183
hSC17 .155	QVQLVQ5GAEVKKPGASVKVSKKASGYTFN	SYMMH	WVRQAPQGQLEIMMG	EIHPNNGSTNNEKFKG	RVTMTSDTSISTAYMELSS LRSDDTAVYYCAR	WTLFTY	WGQGTITVTV	185
hSC17 .156	QVTLKE5GPVLVKPTETLTLTCTVSGFSL5	TSGNVS	WIRQPPGKALEMLA	HIFMDDDKWMPSLKS	RLTISKDSKSKQVVL TMTNMDPVDATYYCAT	FYGLVFAY	WGQGTITVTVSS	187
hSC17 .161	QVQLVQ5GAEVKKPGASVKVSKKASGFTFS	DAWMD	WVRQAPGQRLIMMG	EIRSKPNNHATYYAESVKG	RVTITSDTSASTAYMELSS LRSDDTAVYYCAR	TGTSY	WGQGTITVTVSS	189
hSC17 .200	EVQLVQ5GAEVKKPGESLKLISKSGSY5FT	SSMIN	WVRQMPGKGLEIMMG	RIYPGEGDTNYSGNFEG	QVTTISADKSI5TAYLQWSS LKASDTAMYYCTR	GLVMDY	WGQGTITVTVSS	191
hSC17 .155vH1	QVQLVQ5GAEVKKPGASVKVSKKASGYTFD	SYMMH	WVRQAPQGQLEIMMG	EIHPNNGSTNNEKFKG	RVTMTSDTSISTAYMELSS LRSDDTAVYYCAR	WTLFTY	WGQGTITVTV	193
hSC17 .155vH2	QVQLVQ5GAEVKKPGASVKVSKKASGYTFT	SYMMH	WVRQAPQGQLEIMMG	EIHPNNGSTNNEKFKG	RVTMTSDTSISTAYMELSS LRSDDTAVYYCAR	WTLFTY	WGQGTITVTV	194
hSC17 .155vH3	QVQLVQ5GAEVKKPGASVKVSKKASGYTFN	YYMMH	WVRQAPQGQLEIMMG	EIHPNNGSTNNEKFKG	RVTMTSDTSISTAYMELSS LRSDDTAVYYCAR	WTLFTY	WGQGTITVTV	195
hSC17 .155vH4	QVQLVQ5GAEVKKPGASVKVSKKASGYTFN	SYMMH	WVRQAPQGQLEIMMG	EIHPNNGSTNNEKFKG	RVTMTSDTSISTAYMELSS LRSDDTAVYYCAR	WTLFTY	WGQGTITVTV	196

FIG. 10B Cont.

Protein Sequences of Exemplary Humanized SEZ6 Modulator Heavy Chain Variable Regions

Name	FR1	CDRH1	FR2	CDRH2	FR3	CDRH3	FR4	SEQ ID NO
hSC17 .155vH5	QVQLVQSGAEVKKPGASVKVCKASGYTFN	SYWH	WVRQAPGQGLEWMG	EIHPNGGSTNYNKFKG	RVTMTTRDTSTSTVYMEISSLRSEDTAVYYCAR	WTLFTY	WGQGTLLTVV	197
hSC17 .155vH6	QVQLVQSGAEVKKPGASVKVCKASGYTFN	SYWH	WVRQAPGQGLEWMG	EIHPNSGSTNYNKFKG	RVTMTTRDTSTSTVYMEISSLRSEDTAVYYCAR	WTLFTY	WGQGTLLTVV	198
hSC17 .161vH1	EVQLVESGGGLVQPGGSLRLSCAASGFTFS	DAWMD	WVRQAPGKGLEWMG	EIRSKPNNHATYYAESVKG	RFTISRDDSKNSLYLQMMSLKTEDAVYYCAR	TGTSY	WGQGTLLTVSS	199

FIG. 10B Cont.

Amino Acid Sequences of Exemplary Full Length Humanized Anti-SEZ6 Antibody Light and Heavy Chains

Name	Chain	Full Sequence	SEQ ID NO
hSC17.200	Light	EIVLTQSPATLSLSPGERATLSCRASQSVVDYNGISYNHWYQQKPGQAPRLLIYAASNVQSGIPARFSGSGSGTDFTLTISLLEPEDFVYVYCCQSIEDPPTFGGGTKVEIKRTVAAPSVFIFPPSDEQLKSGTASVWCLLNNFYYPREAKVQWKVDNALQSGNSQESVTEQDSKDSYSTYSLSTLTLSKADYEKHKVYACEVTHQGLSPVTKSFNRGEC	400
hSC17.200	Heavy	EVQLVQSGAEVVKPGESLKISCCKGSGYSFTSSMIINWRQMPGKGLNMGRIYPGEGDNTVSGHFEQQVTSADKSIISTAYLQWESLKASDTAWMYCTRGLVMDYWGQGT LVTVSSASTKGPSVFFLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVIRQSGLYSLSVWTVPSSSLGTQTYICNWNHHPNNTKVDKQVEPKSCDKTHTCPPCPAPELIGGPVSFIFPPKPKDITLITSRTEPEVTCVWVDNSHEDPEVKFNMYVDGV EYHNAKTKPREEQYNSTYRVASVLTVLHQDLNGKEYCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEMESNGQPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNV FSCSVMEALHNHYTQKSLSLSPG	401
hSC17.200vL1	Light	EIVLTQSPATLSLSPGERATLSCRASQSVVDYNGISYNHWYQQKPGQAPRLLIYAASNVQSGIPARFSGSGSGTDFTLTISLLEPEDFVYVYCCQSIEDPPTFGGGTKVEIKRTVAAPSVFIFPPSDEQLKSGTASVWCLLNNFYYPREAKVQWKVDNALQSGNSQESVTEQDSKDSYSTYSLSTLTLSKADYEKHKVYACEVTHQGLSPVTKSFNRGEC	402

FIG. 10C

SEZ6 Modulator Characteristics

Clone	Bin	Hu Affinity (Kd, nM)	Mouse XR	Rat XR (Kd, nM)	Cyno XR	SEZ6L XR	SEZ6L2 XR
SC17.3	D	N.D.	No	No	N.D.	No	No
SC17.4	A	N.D.	No	No	N.D.	No	No
SC17.6	B	<1	Yes	Yes (<1)	N.D.	No	Yes
SC17.7	A	N.D.	Yes	No	No	Yes	Yes
SC17.17	U	1.8	No	Yes (5.0)	Yes	No	No
SC17.19	E	5.3	Yes	Yes	N.D.	No	Yes
SC17.24	C	5.1	Yes	No	Yes	No	No
SC17.26	F	62.0	Yes	Yes (17.4)	Yes	No	Yes
SC17.28	U	22.1	No	Yes (31)	N.D.	No	Yes
SC17.34	A	15.1	No	Yes (79.5)	Yes	No	No
SC17.36	A	6.3	No	No	Yes	No	No
SC17.42	A	2.7	Yes	Yes (3.0)	No	No	No
SC17.45	A	17.4	No	No	Yes	No	No
SC17.46	E	N.D.	No	No	No	No	No
SC17.49	U	23.1	No	No	N.D.	No	No

FIG. 11A

Correlation Between Domain-Level Mapping and *In Vitro* Efficacy




FIG. 11B

Detection of SEZ6 Surface Expression on Engineered HEK-293T Cells




FIG. 12A

SEZ6 Protein Expression is Upregulated in Certain NTX Tumors




FIG. 12B

Selected SEZ6 Modulators
Detect Surface Expression of SEZ6 in NTX Tumors




FIG. 13A

SEZ6 Protein is a Biomarker of CSCs as Determined by FACS




FIG. 13B

Tumor Cells Expressing SEZ6 Exhibit
Enhanced Tumorigenicity in a Lung Tumor (LU37)

EP 3 338 793 A1




FIG. 14A




FIG. 14B

SEZ6 Modulators Facilitate Delivery of Cytotoxic Agents to SEZ6-Expressing HEK-293T Cells

Clone	100pM	50pM	10pM
IgG1	103.1	ND	94.3
IgG2a	97.3	100.4	94.7
IgG2b	102.3	ND	96.9
SC17.1	26.3	ND	70.1
SC17.3	16.9	ND	38.5
SC17.4	18.7	ND	34.7
SC17.6	8.8	ND	14.3
SC17.7	23.9	ND	49.4
SC17.8	15.2	ND	22.8
SC17.9	13.8	ND	24.4
SC17.10	56.6	ND	92.2
SC17.11	23.8	ND	64.5
SC17.12	112.4	ND	111.7
SC17.13	107.0	ND	103.4
SC17.14	34.9	ND	71.8
SC17.15	95.1	ND	110.6
SC17.17	11.3	12.8	21.4
SC17.18	59.3	ND	89.6
SC17.19	9.2	ND	14.4
SC17.21	36.4	ND	62.7
SC17.22	46.9	ND	68.9
SC17.23	13.3	ND	38.0
SC17.24	49.5	ND	77.5
SC17.25	58.7	ND	88.8
SC17.26	13.4	ND	15.6
SC17.27	21.7	ND	33.6
SC17.28	12.8	ND	17.9

Clone	100pM	50pM	10pM
SC17.29	112.4	ND	103.3
SC17.30	22.6	ND	50.0
SC17.31	47.0	ND	87.4
SC17.32	104.7	ND	122.9
SC17.33	10.4	ND	17.6
SC17.34	8.8	ND	16.7
SC17.35	15.9	ND	41.8
SC17.36	9.8	ND	12.9
SC17.37	78.4	ND	92.2
SC17.38	104.9	ND	104.8
SC17.39	121.7	ND	121.8
SC17.40	9.6	ND	11.3
SC17.41	60.3	ND	98.1
SC17.42	58.5	ND	91.7
SC17.43	21.5	ND	36.8
SC17.44	47.3	ND	86.3
SC17.45	11.1	ND	12.6
SC17.46	12.7	16.3	23.6
SC17.47	15.0	ND	39.2
SC17.49	21.3	ND	39.3
SC17.50	102.7	ND	109.9
SC17.51	105.4	ND	102.1
SC17.52	8.2	ND	8.6
SC17.53	111.6	ND	99.9
SC17.54	77.0	ND	99.6
SC17.55	11.0	ND	28.9
SC17.56	12.4	ND	16.5

Clone	100pM	50pM	10pM
SC17.57	111.7	ND	103.0
SC17.58	102.3	ND	95.1
SC17.59	111.4	ND	114.5
SC17.60	110.8	ND	101.2
SC17.61	114.0	ND	113.7
SC17.72	ND	101.3	ND
SC17.73	ND	96.4	ND
SC17.74	ND	105.1	ND
SC17.75	ND	108.3	ND
SC17.76	ND	99.6	ND
SC17.81	ND	62.8	ND
SC17.82	ND	97.7	ND
SC17.83	ND	95.4	ND
SC17.84	ND	107.8	ND
SC17.86	ND	111.7	ND
SC17.87	ND	100.9	ND
SC17.88	ND	97.8	ND
SC17.89	ND	17.0	ND
SC17.90	ND	105.4	ND
SC17.91	ND	104.0	ND
SC17.92	ND	107.4	ND
SC17.93	ND	106.3	ND
SC17.95	ND	113.0	ND
SC17.96	ND	53.5	ND
SC17.97	ND	111.9	ND
SC17.99	ND	18.8	ND
SC17.100	ND	97.0	ND

FIG. 15A

SEZ6 Modulators Facilitate Delivery of
Cytotoxic Agents to SEZ6-Expressing HEK-293T Cells

Clone	100pM	50pM	10pM	Clone	100pM	50pM	10pM
SC17.101	ND	50.7	ND	SC17.136	ND	92.9	ND
SC17.102	ND	13.6	ND	SC17.137	ND	15.3	ND
SC17.103	ND	111.4	ND	SC17.138	ND	107.7	ND
SC17.104	ND	39.4	ND	SC17.140	ND	22.5	ND
SC17.105	ND	115.2	ND	SC17.142	ND	101.4	ND
SC17.106	ND	116.6	ND	SC17.143	ND	97.0	ND
SC17.107	ND	36.0	ND	SC17.145	ND	100.8	ND
SC17.108	ND	104.7	ND	SC17.146	ND	96.7	ND
SC17.109	ND	108.3	ND	SC17.147	ND	108.9	ND
SC17.112	ND	102.7	ND	SC17.149	ND	45.9	ND
SC17.114	ND	11.4	ND	SC17.151	ND	8.1	ND
SC17.115	ND	10.7	ND	SC17.154	ND	102.0	ND
SC17.116	ND	95.4	ND	SC17.155	ND	15.3	ND
SC17.117	ND	107.4	ND	SC17.156	ND	19.5	ND
SC17.119	ND	111.0	ND	SC17.161	ND	18.3	ND
SC17.120	ND	9.1	ND	SC17.162	ND	30.1	ND
SC17.121	ND	16.2	ND	SC17.163	ND	106.5	ND
SC17.122	ND	107.1	ND	SC17.166	ND	19.2	ND
SC17.124	ND	109.9	ND	SC17.167	ND	103.4	ND
SC17.125	ND	98.7	ND	SC17.168	ND	46.0	ND
SC17.129	ND	102.9	ND	SC17.169	ND	109.7	ND
SC17.130	ND	80.4	ND	SC17.170	ND	108.9	ND
SC17.133	ND	108.2	ND	SC17.173	ND	68.9	ND
SC17.134	ND	37.3	ND	SC17.175	ND	15.9	ND
SC17.135	ND	99.0	ND	SC17.176	ND	82.8	ND

Clone	100pM	50pM	10pM	Clone	100pM	50pM	10pM
SC17.177	ND	102.5	ND	SC17.184	ND	63.2	ND
SC17.178	ND	102.1	ND	SC17.186	ND	105.1	ND
SC17.179	ND	70.2	ND	SC17.187	ND	13.5	ND
SC17.180	ND	21.7	ND	SC17.188	ND	21.2	ND
SC17.181	ND	101.2	ND	SC17.189	ND	98.8	ND
SC17.182	ND	10.9	ND	SC17.190	ND	91.3	ND
SC17.184	ND	63.2	ND	SC17.191	ND	17.3	ND
SC17.186	ND	105.1	ND	SC17.192	ND	103.6	ND
SC17.187	ND	13.5	ND	SC17.193	ND	21.1	ND
SC17.188	ND	21.2	ND	SC17.195	ND	38.0	ND
SC17.189	ND	98.8	ND	SC17.196	ND	106.8	ND
SC17.190	ND	91.3	ND	SC17.197	ND	7.0	ND
SC17.191	ND	17.3	ND	SC17.198	ND	108.0	ND
SC17.192	ND	103.6	ND	SC17.199	ND	21.0	ND
SC17.193	ND	21.1	ND	SC17.200	ND	9.9	ND
SC17.195	ND	38.0	ND	SC17.201	ND	40.1	ND
SC17.196	ND	106.8	ND	SC17.203	ND	96.2	ND
SC17.197	ND	7.0	ND	SC17.204	ND	114.1	ND
SC17.198	ND	108.0	ND				
SC17.199	ND	21.0	ND				
SC17.200	ND	9.9	ND				
SC17.201	ND	40.1	ND				
SC17.203	ND	96.2	ND				
SC17.204	ND	114.1	ND				

FIG. 15A (Cont.)

Selected SEZ6 Modulators
Mediate Killing of NSCLC NTX Tumor Cells




FIG. 15B

IHC Measurements of SEZ6 Expression on SCLC NTX

Cellular H-Score For Membrane Staining




FIG. 16A

IHC Measurements of SEZ6 Expression in Lung Cancer Tissue Microarrays




FIG. 16B

IHC Measurement of SEZ6 Expression in Medullary Thyroid Tumors

	Mouse IgG2a	Mouse IgG1	SC17.140
Patient 1	-	-	+, 10%,
Patient 2	-	-	+/+++, 90%,
Patient 3	-	-	++, 40%,
Patient 4	-	-	+++, 80%,
Patient 5	-	-	++, 80%,
Patient 6	-	-	-/+, 10%,
Patient 8	-	-	++, 90%
Patient 9	-	-	++, 75%
Patient 10 (Thy1p0MET)	-	ND	+++, 95%,

FIG. 16C

Murine SEZ6 ADC Modulators Suppress
SCLC and OV NTX Tumor Growth *In Vitro*




FIG. 17A

Murine SEZ6 ADC Modulators Suppress SCLC and NSCLC NTX Tumor Growth *In Vivo*




FIG. 17B

Humanized SEZ6 ADC Modulators
Suppress Growth of Thyroid Cancer Cells *In Vitro*




FIG. 18A

Humanized SEZ6 ADC Modulators Suppress Growth of Thyroid Tumors *In Vivo*




FIG. 18B

Humanized SEZ6 ADC Modulators
Suppress SCLC NTX Tumor Growth *In Vivo*




FIG. 18C

Humanized SEZ6 ADC Modulators

Show Differential Efficacy on SCLC NTX Tumor Growth *In Vivo*




FIG. 18D

In vitro sensitivity of parental and oxaliplatin-resistant tumor
cell line




FIG. 19

SEZ6 mRNA expression determined by microarray




FIG. 20

SEZ6 protein expression in oxaliplatin-resistant tumor cell line




FIG. 21



EUROPEAN SEARCH REPORT

Application Number
EP 17 20 3670

5

10

15

20

25

30

35

40

45

50

55

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (IPC)
X,P	WO 2013/126810 A1 (STEM CENTRX INC [US]) 29 August 2013 (2013-08-29) * page 84, paragraph 5 - page 85, paragraph 1; examples 16-19 *	1-15,17	INV. A61K39/00 A61K39/395 C12P21/08 C07K16/00 C07K16/18 C07K16/28 G01N33/574
A	FRANÇOISE KRAEBER-BODÉRE ET AL: "Pretargeted radioimmunotherapy in rapidly progressing, metastatic, medullary thyroid cancer", CANCER., vol. 116, no. S4, 2 February 2010 (2010-02-02), pages 1118-1125, XP055469044, US ISSN: 0008-543X, DOI: 10.1002/cncr.24800 * the whole document *	16	
A	FROESCH B ET AL: "Combination of doxorubicin-immunoconjugates and molecular intervention in bcl-2 oncogene expression to overcome drug resistance in small cell lung cancer", PROCEEDINGS OF THE AMERICAN ASSOCIATION FOR CANCER RESEARCH ANNUAL MEETING, vol. 36, 1995, page 341, XP002780294, EIGHTY-SIXTH ANNUAL MEETING OF THE AMERICAN ASSOCIATION FOR CANCER RESEARCH; TORONTO, ONTARIO, CANADA; MARCH 18-22, 1995 ISSN: 0197-016X * abstract *	16	TECHNICAL FIELDS SEARCHED (IPC) A61K C07K
The present search report has been drawn up for all claims			
Place of search The Hague		Date of completion of the search 20 April 2018	Examiner Lonnoy, Olivier
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	

EPO FORM 1503 03.02 (P04C01)

**ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.**

EP 17 20 3670

5

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report.
The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

20-04-2018

10

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 2013126810 A1	29-08-2013	AU 2013203506 A1	12-09-2013
		AU 2016225828 A1	22-09-2016
		CA 2865415 A1	29-08-2013
		CL 2014002237 A1	15-05-2015
		CN 104334580 A	04-02-2015
		CO 7151485 A2	29-12-2014
		EP 2817339 A1	31-12-2014
		HK 1205517 A1	18-12-2015
		JP 2015509948 A	02-04-2015
		KR 20150002621 A	07-01-2015
		NZ 631197 A	26-05-2017
		PE 00912015 A1	16-02-2015
		PH 12014501914 A1	24-11-2014
		RU 2014138420 A	10-04-2016
		SG 11201405130U A	27-11-2014
		US 2015018531 A1	15-01-2015
		US 2017369571 A1	28-12-2017
		WO 2013126810 A1	29-08-2013

15

20

25

30

35

40

45

50

55

EPO FORM P0459

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- US 61871289 A [0001]
- US 7632678 B [0079]
- US 20070292414 A [0079] [0516] [0600] [0615] [0643]
- US 20080175870 A [0079]
- US 20100275280 A [0079]
- US 20100162416 A [0079]
- US 20110020221 A [0079]
- US 20080138313 A [0080]
- US 4816567 A [0109]
- US 6982321 B [0112]
- US 7087409 B [0112]
- US 20050008625 A [0112]
- WO 9852976 A [0112]
- WO 0034317 A [0112]
- US 6300064 B [0113]
- US 5223409 A [0116]
- WO 9218619 A [0116]
- WO 9117271 PCT [0116]
- WO 9220791 PCT [0116]
- WO 9215679 PCT [0116]
- WO 9301288 PCT [0116]
- WO 9201047 PCT [0116]
- WO 9209690 A [0116]
- WO 9607754 A [0118]
- US 7700302 B [0119]
- US 12404059 B [0119]
- US 5545807 A [0120]
- US 5545806 A [0120]
- US 5569825 A [0120]
- US 5625126 A [0120]
- US 5633425 A [0120]
- US 5661016 A [0120]
- US 6075181 A [0120]
- US 6150584 A [0120]
- US 5750373 A [0120]
- US 7709611 B [0124]
- US 4399216 A [0139]
- US 4912040 A [0139]
- US 4740461 A [0139]
- WO 8605807 A [0139]
- WO 8901036 A [0139]
- US 5122464 A [0139]
- EP 0216846 A [0142]
- EP 0256055 A [0142]
- EP 0323997 A [0142]
- EP 0338841 A [0142]
- US 5591639 A [0142]
- US 5879936 A [0142]
- US 5827690 A [0146]
- US 5756687 A [0146]
- US 5750172 A [0146]
- US 5741957 A [0146]
- US 6417429 B [0147]
- US 6046037 A [0148]
- US 5959177 A [0148]
- US 20090130105 A [0158]
- US 20090155255 A [0159] [0289]
- WO 9404690 A [0159]
- WO 9627011 A [0159]
- US 4676980 A [0160]
- WO 9100360 A [0160]
- WO 92200373 A [0160]
- EP 03089 A [0160]
- WO 9734631 A [0164]
- WO 04029207 A [0164]
- US 6737056 B [0164]
- US 20030190311 A [0164]
- WO 200042072 A [0164]
- WO 2012117002 A [0167]
- US 7619068 B [0174]
- WO 0348731 A [0180]
- US 20040101920 A [0188]
- WO 9321232 A [0200]
- EP 439095 A [0200]
- US 5474981 A [0200]
- US 6214345 B [0203] [0626]
- US 20120078028 A [0203] [0626]
- US 5122368 A [0204]
- US 5824805 A [0204]
- US 5622929 A [0204] [0437]
- US 20050238649 A [0205]
- US 20110256157 A [0206] [0290] [0291] [0626]
- US 7521541 B [0267]
- WO 2005082023 A [0279]
- WO 9315199 A [0283]
- WO 9315200 A [0283]
- WO 0177137 A [0283]
- EP 0413622 A [0283]
- US 4703004 A [0286]
- FR 901464 [0288]
- US 7825267 B [0288]
- WO 03075957 A [0289]
- US 6362331 B [0290]
- US 7049311 B [0290]
- US 7189710 B [0290]
- US 7429658 B [0290]
- US 7407951 B [0290]

- US 7741319 B [0290]
- US 7557099 B [0290]
- US 8034808 B [0290]
- US 8163736 B [0290]
- WO 2011130613 A [0290]
- WO 2011128650 A [0290]
- WO 2011130616 A [0290]
- WO 0012507 A [0402]
- WO 9733899 A [0437]
- WO 9734911 A [0437]
- WO 9923105 A [0437]
- US 5336603 A [0437]
- US 5359046 A [0437]
- US 5349053 A [0437]
- US 5447851 A [0437]
- US 5112946 A [0437]
- EP 307434 A [0437]
- EP 367166 A [0437]
- WO 9604388 A [0437]
- WO 9106570 A [0437]
- WO 20120128801 A [0443] [0506]
- US 7744877 B [0462]
- US 7422739 B [0501]
- WO 2012031280 A [0613]
- US 2012031280 W [0614]

Non-patent literature cited in the description

- **ABBAS et al.** Cellular and Molecular Immunology. W.B. Saunders Company, 2010 [0090] [0508]
- **MURPHEY et al.** Janeway's Immunobiology. Garland Science, 2011 [0090]
- **KABAT et al.** National Technical Information Service. NIH Publication, 1991, 91-3242 [0095]
- **CHOTHIA et al.** *J. Mol. Biol.*, 1987, vol. 196, 901-917 [0097]
- **MACCALLUM et al.** *J. Mol. Biol.*, 1996, vol. 262, 732-745 [0097]
- **CHOTHIA et al.** *Nature*, 1989, vol. 342, 877-883 [0097]
- Handbook of Therapeutic Antibodies. WILEY-VCH Verlag GmbH and Co, 2007 [0097]
- Antibody Engineering. Springer, 2001 [0098]
- **DINARELLO et al.** Current Protocols in Immunology. John Wiley and Sons Inc, 2000 [0098]
- **A.C. MARTIN.** Department of Biochemistry & Molecular Biology. University College London [0098]
- **RETTET et al.** *Nucl. Acids Res.*, 2005, vol. 33, D671-D674 [0098]
- **EDELMAN et al.** *Proc. Natl. Acad. Sci. USA*, 1969, vol. 63 (1), 78-85 [0102]
- **BHASKARAN R. ; PONNUSWAMY P. K.** *Int. J. Pept. Protein Res.*, 1988, vol. 32, 242-255 [0106]
- **DELEAGE, G. ; ROUX B.** *Protein Engineering*, 1987, vol. 1, 289-294 [0106]
- Therapeutic Monoclonal Antibodies: From Bench to Clinic. John Wiley and Sons, 2009 [0108]
- Current Trends in Monoclonal Antibody Development and Manufacturing. Springer Science + Business Media LLC, 2010 [0108]
- **HARLOW et al.** Antibodies: A Laboratory Manual. Cold Spring Harbor Laboratory Press, 1988 [0108]
- **HAMMERLING et al.** Monoclonal Antibodies and T-Cell Hybridomas. Elsevier, 1981, 563-681 [0108]
- **MORRISON et al.** *Proc. Natl. Acad. Sci. USA*, 1984, vol. 81, 6851-6855 [0109]
- **JONES et al.** *Nature*, 1986, vol. 321, 522-525 [0112]
- **TOMLINSON, I. A. et al.** *J. Mol. Biol.*, 1992, vol. 227, 776-798 [0113]
- **COOK, G. P. et al.** *Immunol. Today*, 1995, vol. 16, 237-242 [0113]
- **CHOTHIA, D. et al.** *J. Mol. Biol.*, 1992, vol. 227, 799-817 [0113]
- **TOMLINSON et al.** *EMBO J.*, 1995, vol. 14, 4628-4638 [0113]
- **RETTET et al.** *Nucleic Acid Res.*, 2005, vol. 33, 671-674 [0113]
- **TOMLINSON, I. A. et al.** *MRC Centre for Protein Engineering* [0113]
- **BARBAS et al.** *Proc. Natl. Acad. Sci. USA*, 1991, vol. 88, 7978-7982 [0116]
- **LEUNG et al.** *Technique*, 1989, vol. 1, 11-15 [0118]
- **MARKS et al.** *Biotechnol.*, 1992, vol. 10, 779-783 [0118]
- **VAUGHAN et al.** *Nature Biotechnology*, 1996, vol. 14, 309-314 [0119]
- **SHEETS et al.** *Proc. Natl. Acad. Sci. USA*, 1998, vol. 95, 6157-6162 [0119]
- **LONBERG ; HUSZAR.** *Intern. Rev. Immunol.*, 1995, vol. 13, 65-93 [0120]
- **COLE et al.** Monoclonal Antibodies and Cancer Therapy. Alan R. Liss, 1985, 77 [0120]
- **BOERNER et al.** *J. Immunol.*, 1991, vol. 147 (1), 86-95 [0120]
- **SAMBROOK J. ; RUSSELL D.** Molecular Cloning: A Laboratory Manual. Cold Spring Harbor Laboratory Press, 2000 [0123] [0508]
- **AUSUBEL et al.** Short Protocols in Molecular Biology: A Compendium of Methods from Current Protocols in Molecular Biology. Wiley, John & Sons, Inc, 2002 [0123] [0508]
- **KABAT, E. A. et al.** Sequences of Proteins of Immunological Interest. U.S. Department of Health and Human Services, 1991 [0128]
- **KABAT, E. A. et al.** Sequences of Proteins of Immunological Interest. Department of Health and Human Services, 1991 [0129]
- Molecular Cloning: A Laboratory Manual. Cold Spring Harbor Laboratory Press, 1989 [0132]

- Current Protocols in Molecular Biology. John Wiley & Sons, Inc, 1995 [0132]
- DURBIN. Biological Sequence Analysis: Probabilistic models of proteins and nucleic acids. Cambridge Press, 1998 [0133]
- ALTSCHUL et al. *J. Mol. Biol.*, 1990, vol. 215, 403-410 [0134]
- ALTSCHUL et al. *Nucleic Acids Res.*, 1997, vol. 25 (402), 3389 [0134]
- Guide to Molecular Cloning Techniques. BERGER ; KIMMEL. Methods in Enzymology. Academic Press, Inc, vol. 152 [0136]
- SAMBROOK et al. Molecular Cloning-A Laboratory Manual. Cold Spring Harbor Laboratory, 2000, vol. 1-3 [0136]
- Current Protocols in Molecular Biology. Greene Publishing Associates, Inc. and John Wiley & Sons, Inc, 2006 [0136]
- CREIGHTON. Proteins: Structures and Molecular Principles. W.H. Freeman & Co, 1983 [0145]
- HUNKAPILLER, M. et al. *Nature*, 1984, vol. 310, 105-111 [0145]
- HOGAN et al. Manipulating the Mouse Embryo: A Laboratory Manual. Cold Spring Harbor Press, 1999 [0147]
- JACKSON et al. Mouse Genetics and Transgenics: A Practical Approach. Oxford University Press, 2000 [0147]
- PINKERT. Transgenic Animal Technology: A Laboratory Handbook. Academic Press, 1999 [0147]
- LINDMARK et al. *J Immunol Meth*, 1983, vol. 62, 1 [0150]
- GUSS et al. *EMBO J*, 1986, vol. 5, 1567 [0150]
- Fundamental Immunology. Raven Press, 1999 [0156]
- MILLSTEIN et al. *Nature*, 1983, vol. 305, 537-539 [0159]
- SURESH et al. *Methods in Enzymology*, 1986, vol. 121, 210 [0159]
- RAVETCH ; KINET. *Annu. Rev. Immunol*, 1991, vol. 9, 457-92 [0163]
- CAPEL et al. *Immunomethods*, 1994, vol. 4, 25-34 [0163]
- HAAS et al. *J. Lab. Clin. Med.*, 1995, vol. 126, 330-41 [0163]
- SHIELDS et al. *J. Biol. Chem.*, 2001, vol. 9 (2), 6591-6604 [0164]
- SHIELDS, R. L. et al. *J. Biol. Chem.*, 2002, vol. 277, 26733-26740 [0166]
- COCHRAN et al. *J Immunol Methods.*, 2004, vol. 287 (1-2), 147-158 [0187] [0587]
- REINEKE. *Methods Mol Biol*, 2004, vol. 248, 443-63 [0188]
- TOMER. *Protein Science*, 2000, vol. 9, 487-496 [0188]
- BJELLQVIST et al. *Electrophoresis*, 1993, vol. 14, 1023 [0196]
- VERMEER et al. *Biophys. J.*, 2000, vol. 78, 394-404 [0196]
- VERMEER et al. *Biophys. J.*, 2000, vol. 79, 2150-2154 [0196]
- NARAMURA et al. *Immunol. Lett.*, 1994, vol. 39, 91-99 [0200]
- GILLIES et al. *PNAS*, 1992, vol. 89, 1428-1432 [0200]
- FELL et al. *J. Immunol.*, 1991, vol. 146, 2446-2452 [0200]
- HERMANSON, G.T. Bioconjugate Techniques. Academic Press, 1996, 234-242 [0202]
- JOHNSON et al. *Anticancer Res.*, 1995, vol. 15, 1387-93 [0205]
- LAU et al. *Bioorg-Med-Chem.*, 1995, vol. 3 (10), 1299-1304 [0205]
- LAU et al. *Bioorg-Med-Chem.*, 1995, vol. 3 (10), 1305-12 [0205]
- DUBOWCHIK et al. *Bioconjugate Chemistry*, 2002, vol. 13, 855-869 [0235]
- DENARDO et al. *Clin Cancer Res.*, 1998, vol. 4, 2483 [0281]
- PETERSON et al. *Bioconjug. Chem.*, 1999, vol. 10, 553 [0281]
- ZIMMERMAN et al. *Nucl. Med. Biol.*, 1999, vol. 26, 943 [0281]
- GARNETT. *Adv Drug Deliv Rev*, 2002, vol. 53, 171 [0282] [0437]
- Monoclonal Antibodies For Immunotargeting Of Drugs In Cancer Therapy. AMON et al. Monoclonal Antibodies And Cancer Therapy. Alan R. Liss, Inc, 1985, 243-56 [0282]
- Antibodies For Drug Delivery. HELLSTROM et al. Controlled Drug Delivery. Marcel Dekker, Inc, 1987, 623-53 [0282]
- Antibody Carriers Of Cytotoxic Agents In Cancer Therapy: A Review. THORPE et al. Monoclonal Antibodies '84: Biological And Clinical Applications. 1985, 475-506 [0282]
- Analysis, Results, And Future Prospective Of The Therapeutic Use Of Radiolabeled Antibody In Cancer Therapy. Monoclonal Antibodies For Cancer Detection And Therapy. Academic Press, 1985, 303-16 [0282]
- THORPE et al. *Immunol. Rev.*, 1982, vol. 62, 119 [0282]
- WILSON et al. *Cell*, 1984, vol. 37, 767 [0286]
- FUHRMANN, S. *Annual Meeting of AACR Abstract*, 2010 [0288]
- TAKAHASHI et al. *J. Immunol.*, 1994, vol. 6, 1567 [0437]
- ASHKENAZI et al. *PNAS USA*, 1991, vol. 88, 10535 [0437]
- ZHENG et al. *J Immunol*, 1995, vol. 154, 5590 [0437]
- VIL et al. *PNAS USA*, 1992, vol. 89, 11337 [0437]
- DENARDO et al. *Clin Cancer Res*, 1998, vol. 4, 2483 [0437]

- **PETERSON et al.** *Bioconjug Chem*, 1999, vol. 10, 553 [0437]
- **ZIMMERMAN et al.** *Nucl Med Biol*, 1999, vol. 26, 943 [0437]
- **PINHASOV et al.** *Comb. Chem. High Throughput Screen*, 2004, vol. 7, 133 [0451]
- **MOCELLIN ; ROSSI.** *Adv. Exp. Med. Biol.*, 2007, vol. 593, 19 [0451]
- **GENNARO.** *Remington: The Science and Practice of Pharmacy with Facts and Comparisons: Drugfacts Plus*, 2003 [0453]
- **ANSEL et al.** *Pharmaceutical Dosage Forms and Drug Delivery Systems*. Lippencott Williams and Wilkins, 2004 [0453]
- **KIBBE et al.** *Handbook of Pharmaceutical Excipients*. Pharmaceutical Press, 2000 [0453]
- **REMINGTON.** *The Science and Practice of Pharmacy*. Mack Publishing, 2000 [0455]
- **GAIDONO et al.** *Lymphomas. IN CANCER: PRINCIPLES & PRACTICE OF ONCOLOGY*, vol. 2, 2131-2145 [0497]
- **HARLOW ; LANE.** *Using Antibodies: A Laboratory Manual*. Cold Spring Harbor Laboratory Press, 1998 [0508]
- **COLIGAN et al.** *Short Protocols in Protein Science*. Wiley, John & Sons, Inc, 2003 [0508]
- **BORKP ; BECKMANN G.** *The CUB domain. A widespread module in developmentally regulated proteins. J Mol Biol.*, 1993, vol. 231, 539-45 [0509]
- **COOK M et al.** *Notch in the development of thryof C-cells and the treatment of medullary thyroid cancer. Am J Transl Res.*, 2010, vol. 2, 119-25 [0509]
- **GALLUZZO P ; BOCCHETTA M.** *Notch signaling in lung cancer. Expert Rev Anticancer Ther.*, 2011, vol. 11, 533-40 [0509]
- **GUNNERSEN JM et al.** *Sez-6 proteins affect dendritic arborization patterns and excitability of cortical pyramidal neurons. Neuron*, 2007, vol. 56, 621-39 [0509]
- **GUNNERSEN JM et al.** *Seizure-related gene 6 (Sez-6) in amacrine cells of the rodent retina and the consequence of gene deletion. PLoS One*, 2009, vol. 4, e6546 [0509]
- **HADDAD RI.** *How to incorporate new tyrosine kinase inhibitors in the treatment of patients with medullary thyroid cancer. J Clin Oncolo*, 2013, vol. 31, 3618-20 [0509]
- **HERBST R ; NICKLIN MJ.** *SEZ-6: promoter selectivity, genomic structure and localized expression in the brain. Brain Res Mol Brain Res.*, 1997, vol. 44, 309-22 [0509]
- **KLIMSTRA DS et al.** *The pathologic classification of neuroendocrine tumors: a review of nomenclature, grading, and staging systems. Pancreas*, 2010, vol. 39, 707-12 [0509]
- **KLÖPPEL G.** *Classification and pathology of gastroenteropancreatic neuroendocrine neoplasms. Endocr Relat Cancer*, 2011, vol. 18 (1), 1-16 [0509]
- **MULLEY JC et al.** *The Role of Seizure-Related SEZ6 as a Susceptibility Gene in Febrile Seizures. Neurol Res Int.*, 2011, vol. 2011, 917565 [0509]
- **SHIMIZU-NISHIKAWA K et al.** *Cloning and expression of SEZ-6, a brain-specific and seizure-related cDNA. Brain Res Mol Brain Res.*, 1995, vol. 28, 201-10 [0509]
- **YAO JC et al.** *One hundred years after "carcinoid": epidemiology of and prognostic factors for neuroendocrine tumors in 35,825 cases in the United States. J Clin Oncol.*, 2008, vol. 26, 3063-72 [0509]
- **YU ZL et al.** *Febrile seizures are associated with mutation of seizure-related (SEZ) 6, a brain-specific gene. J Neurosci Res.*, 2007, vol. 85, 166-72 [0509]
- **WANG et al.** *Advanced Cell Diagnostics*, 2012 [0610]

摘要

本发明提供了新型调节剂，包括抗体及其衍生物，以及使用这样的调节剂治疗增殖性病症的方法。