



Office de la Propriété
Intellectuelle
du Canada

Un organisme
d'Industrie Canada

Canadian
Intellectual Property
Office

An agency of
Industry Canada

CA 2573241 A1 2004/08/12

(21) **2 573 241**

(12) **DEMANDE DE BREVET CANADIEN
CANADIAN PATENT APPLICATION**

(13) **A1**

(22) Date de dépôt/Filing Date: 2004/01/30

(41) Mise à la disp. pub./Open to Public Insp.: 2004/08/12

(62) Demande originale/Original Application: 2 514 224

(30) Priorité/Priority: 2003/01/30 (US60/444,351)

(51) Cl.Int./Int.Cl. *A61K 31/473* (2006.01),
A61J 1/05 (2006.01), *A61K 47/26* (2006.01),
A61K 9/08 (2006.01), *A61P 1/08* (2006.01)

(71) Demandeur/Applicant:
HELSINN HEALTHCARE SA, CH

(72) Inventeurs/Inventors:
CALDERARI, GIORGIO, CH;
BONADEO, DANIELE, IT;
CANNELLA, ROBERTA, IT;
BRAGLIA, ENRICO, CH;
BRAGLIA, RICARDO, CH

(74) Agent: BULL, HOUSSEY & TUPPER LLP

(54) Titre : FORMULATIONS PHARMACEUTIQUES LIQUIDES DE PALONOSETRONE

(54) Title: LIQUID PHARMACEUTICAL FORMULATIONS OF PALONOSETRON

(57) **Abrégé/Abstract:**

The present invention relates to shelf-stable liquid formulations of palonosetron for reducing chemotherapy and radiotherapy induced emesis with palonosetron. The formulations are particularly useful in the preparation of intravenous and oral liquid medicaments.



ABSTRACT

The present invention relates to shelf-stable liquid formulations of palonosetron for reducing chemotherapy and radiotherapy induced emesis with palonosetron. The formulations are particularly useful in the preparation of intravenous and oral liquid medicaments.

5

LIQUID PHARMACEUTICAL FORMULATIONS OF PALONOSETRON

FIELD OF THE INVENTION

The present invention relates to shelf-life stable liquid formulations of
5 palonosetron that are especially useful in the preparation of injectable and oral
medicaments.

BACKGROUND OF THE INVENTION

Emesis is a devastating consequence of cytotoxic therapy, radiotherapy, and post-
10 operative environments that drastically affects the quality of life of people undergoing
such treatments. In recent years a class of drugs referred to as 5-HT₃ (5-
hydroxytryptamine) receptor antagonists has been developed that treat such emesis by
antagonizing cerebral functions associated with the 5-HT₃ receptor. *See Drugs Acting on*
5-Hydroxytryptamine Receptors: The Lancet Sep. 23, 1989 and references cited therein.
15 Drugs within this class include ondansetron, granisetron, alosetron, tropisetron, and
dolasetron. These 5-HT₃ antagonists are often administered intravenously shortly before
chemotherapy or radiotherapy is initiated, and can be administered more than once during
a cycle of chemotherapy or radiotherapy. In addition, they are often supplied as tablets or
oral elixirs to either supplement an intravenous administration, or to ease home usage of
20 the drug if the patient is self-administering the chemotherapeutic regimen.

Because some chemotherapeutic agents can induce emesis over extended periods
of several days even when they are administered only once, it would be desirable to
administer an emesis-inhibiting drug such as a 5-HT₃ antagonist every day until the risk
of emesis has substantially subsided. The present class of 5-HT₃ antagonists has not
25 proven especially helpful meeting this need, however, because the 5-HT₃ receptor
antagonists currently marketed have proven to be less effective in controlling delayed
nausea and vomiting than they are at controlling acute emesis. Sabra, K, *Choice of a*
5HT₃ Receptor Antagonist for the Hospital Formulary. EHP, Oct. 1996;2 (suppl 1):S19-
24.

Recently, clinical investigations have been made concerning palonosetron, a new 5-HT₃ receptor antagonist reported in U.S. Patent No. 5,202,333. These investigations have shown that the drug is an order of magnitude more potent than most existing 5-HT₃ receptor antagonists, has a surprising half-life of about 40 hours, and is effective to
 5 reduce delayed-onset nausea induced by chemotherapeutic agents. However, formulating palonosetron in liquid formulations has not proven an easy task, typically due to shelf-stability issues. U.S. Pat. No. 5,202,333 discloses an intravenous formulation of palonosetron in example 13 that contains the following ingredients:

Ingredient	Mg
Palonosetron HCl	10-100 mg.
Dextrose Monohydrate	q.s. to make Isotonic
Citric Acid Monohydrate	1.05 mg.
Sodium Hydroxide	0.18 mg.
WFJ	To 1.0 ml.

10

The formulation has a pH of 3.7 and a shelf stability of less than the 1-2 year time period required by health authorities in various countries.

Ondansetron, its uses, and medicaments made with ondansetron are disclosed in U.S. Patent Numbers 4,695,578, 4,753,789, 4,929,632, 5,240,954, 5,344,658, 5,578,628,
 15 5,578,632, 5,922,749, 5,622,720, 5,955,488, and 6,063,802. Commercially it is distributed by GlaxoSmithKline as Zofran® and is indicated for prevention of postoperative nausea and vomiting (PONV), cancer chemotherapy-induced nausea and vomiting (CINV), and radiotherapy-induced nausea and vomiting (RINV) and it is available as an injection, tablets and solution, and as Zofran ODT® (ondansetron) Orally
 20 Disintegrating Tablets.

Granisetron, its uses, and medicaments made with granisetron are disclosed in U.S. Patent Numbers 4,886,808, 4,937,247, 5,034,398 and 6,294,548. Commercially it is

distributed by Roche Laboratories Inc. as Kytril®, indicated for the prevention of nausea and vomiting associated with chemotherapy or radiation therapy, and is offered in tablet form, oral solution, and as an injection.

Alosetron, its uses, and medicaments made with alosetron are disclosed in U.S. Patent Numbers 5,360,800 and 6,284,770. Commercially it is distributed by GlaxoSmithKline as Lotronex®.

Tropisetron is commercially available as Navoban® (Novartis) CAS - 89565-68-4 (tropisetron); CAS - 105826-92-4 (tropisetron hydrochloride) and it is indicated for treatment of PONV and CINV.

Dolasetron, its uses, and medicaments made with ondansetron are disclosed in U.S. Patent Numbers 5,011,846, and 4,906,755. Commercially it is distributed by Aventis Pharmaceuticals Inc. as Anzemet®, indicated for prevention of both PONV and CINV, and it is offered in the form of a tablet or an intravenous solution.

Therefore, there exists a need for a palonosetron formulation with increased stability and thereby increased shelf life. There also exists a need for an appropriate range of concentrations for both the 5-HT₃ receptor antagonist and its pharmaceutically acceptable carriers that would facilitate making a formulation with this increased stability.

It is an object of the present invention to provide a formulation of Palonosetron hydrochloride with increased pharmaceutical stability for preventing and/or reducing emesis.

It is another object of the invention to provide an acceptable range of concentrations which will stabilize a formulation containing Palonosetron hydrochloride.

It is a further object of the invention to provide a formulation of Palonosetron which would allow for prolonged storage.

It is also an object of the invention to provide a formulation of Palonosetron which would allow terminal sterilization.

SUMMARY OF THE INVENTION

The inventors have made a series of discoveries that support a surprisingly effective and versatile formulation for the treatment and prevention of emesis using
5 palonosetron. These formulations are shelf stable for periods greater than 24 months at room temperature, and thus can be stored without refrigeration, and manufactured using non-aseptic, terminal sterilization processes.

In one aspect, the inventors have discovered that formulations which include the active ingredient palonosetron require in some instances only $1/10^{\text{th}}$ the amount of other
10 previously known compounds for treating emesis, which surprisingly allows the use of concentrations of palonosetron far below those that would ordinarily be expected. Thus, in one embodiment the invention provides a pharmaceutically stable solution for preventing or reducing emesis comprising a) from about 0.01 mg/mL to about 5 mg/mL palonosetron or a pharmaceutically acceptable salt thereof; and b) a pharmaceutically
15 acceptable carrier.

The inventors have further discovered that by adjusting the formulation's pH and/or excipient concentrations it is possible to increase the stability of palonosetron formulations. Therefore, in another embodiment, the invention provides a pharmaceutically stable solution for preventing or reducing emesis comprising a)
20 palonosetron or a pharmaceutically acceptable salt thereof; and b) a pharmaceutically acceptable carrier, at a pH from about 4.0 to about 6.0. In another embodiment the invention provides a pharmaceutically stable solution for preventing or reducing emesis comprising from about 0.01 to about 5.0 mg/ml palonosetron or a pharmaceutically acceptable salt thereof; from about 10 to about 100 millimoles citrate buffer; and from
25 about 0.005 to about 1.0 mg/ml EDTA.

The inventors have further discovered that the addition of mannitol and a chelating agent can increase the stability of palonosetron formulations. Therefore, in still another embodiment the invention provides a pharmaceutically stable solution for preventing or reducing emesis comprising a) palonosetron or a pharmaceutically

acceptable salt thereof and b) a pharmaceutically acceptable carrier, wherein the pharmaceutically acceptable carrier comprises a chelating agent and mannitol.

DETAILED DESCRIPTION OF THE INVENTION


5 DEFINITIONS

“Vial” means a small glass container sealed with the most suitable stopper and seal, other suitable primary containers may be used, for instance, but not limited to, pre-filled syringes. Vial also means a sealed container of medication that is used one time only, and includes breakable and non-breakable glass vials, breakable plastic vials, 10 miniature screw-top jars, and any other type of container of a size capable of holding only one unit dose of palonosetron (typically about 5 mls.).

Throughout this specification the word “comprise,” or variations such as “comprises” or “comprising,” will be understood to imply the inclusion of a stated element, integer or step, or group of elements, integers or steps, but not the exclusion of 15 any other element, integer or step, or group of elements, integers or steps

“Palonosetron” means (3aS)-2,3,3a,4,5,6-Hexahydro-2-[(S)-1-Azabicyclo[2.2.2]oct-3-yl]2,3,3a,4,5,6-hexahydro-1-oxo-1*H*benz[*de*]isoquinoline, and is preferably present as the monohydrochloride. Palonosetron monohydrochloride can be represented by the following chemical structure:

20



Concentrations -- When concentrations of palonosetron are given herein, the concentration is measured in terms of the weight of the free base. Concentrations of all other ingredients are given based on the weight of ingredient added to the solution.

5 "Pharmaceutically acceptable" means that which is useful in preparing a pharmaceutical composition that is generally safe, non-toxic and neither biologically nor otherwise undesirable and includes that which is acceptable for veterinary use as well as human pharmaceutical use.

"Pharmaceutically acceptable salts" means salts which are pharmaceutically acceptable, as defined above, and which possess the desired pharmacological activity.
10 Such salts include acid addition salts formed with inorganic acids such as hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid, and the like; or with organic acids such as acetic acid, propionic acid, hexanoic acid, heptanoic acid, cyclopentanepropionic acid, glycolic acid, pyruvic acid, lactic acid, malonic acid, succinic acid, malic acid, maleic acid, fumaric acid, tartaric acid, citric acid, benzoic acid,
15 o-(4-hydroxybenzoyl)benzoic acid, cinnamic acid, mandelic acid, methanesulfonic acid, ethanesulfonic acid, 1,2-ethanedisulfonic acid, 2-hydroxyethanesulfonic acid, benzenesulfonic acid p-chlorobenzenesulfonic acid, 2-naphthalenesulfonic acid, p-toluenesulfonic acid, camphorsulfonic acid, 4-methylbicyclo[2.2.2]oct-2-ene-1-carboxylic acid, glucoheptonic acid, 4,4'-methylenebis(3-hydroxy-2-ene-1-carboxylic
20 acid), 3-phenylpropionic acid, trimethylacetic acid, tertiary butylacetic acid, lauryl sulfuric acid, gluconic acid, glutamic acid, hydroxynaphthoic acid, salicylic acid, stearic acid, muconic acid, and the like.

In addition, pharmaceutically acceptable salts may be formed when an acidic proton present is capable of reacting with inorganic or organic bases. Acceptable
25 inorganic bases include sodium hydroxide, sodium carbonate, potassium hydroxide, aluminum hydroxide and calcium hydroxide. Acceptable organic bases include ethanolamine, diethanolamine, triethanolamine, tromethamine, N-methylglucamine and the like.

DISCUSSION

The fact that palonosetron can be formulated in some instances at concentrations of only about $1/10^{\text{th}}$ the amount of other previously known compounds for treating emesis, surprisingly allows the use of concentrations of palonosetron far below those that would ordinarily be expected. Thus, in one embodiment the invention provides a
5 pharmaceutically stable solution for preventing or reducing emesis comprising a) from about 0.01 mg/mL to about 5 mg/mL palonosetron or a pharmaceutically acceptable salt thereof; and b) a pharmaceutically acceptable carrier. Similarly, in another embodiment the invention provides a method of formulating a pharmaceutically stable solution of palonosetron comprising admixing from about 0.01 mg/mL to about 5 mg/mL
10 palonosetron or a pharmaceutically acceptable salt thereof; with a pharmaceutically acceptable carrier. In alternative embodiments, the formulation includes palonosetron or a pharmaceutically acceptable salt thereof in a concentration from about 0.02 mg/mL to about 1.0 mg/mL, from about 0.03 mg/mL to about 0.2 mg/mL, and most optimally about 0.05 mg/mL.

15 A particular advantage associated with the lower dosages of intravenous palonosetron is the ability to administer the drug in a single intravenous bolus over a short, discrete time period. This time period generally extends from about 10 to about 60 seconds, or about 10 to about 40 seconds, and most preferably is about 10 to 30 seconds. In one particular embodiment the palonosetron is supplied in vials that comprise 5 mL of
20 solution, which equates to about 0.25 mg of palonosetron at a concentration of about 0.05 mg/mL.

The inventors have further discovered that by adjusting the formulation's pH and/or excipient concentrations it is possible to increase the stability of palonosetron formulations. Therefore, in another embodiment, the invention provides a
25 pharmaceutically stable solution for preventing or reducing emesis comprising a) palonosetron or a pharmaceutically acceptable salt thereof; and b) a pharmaceutically acceptable carrier, at a pH from about 4.0 to about 6.0. Similarly, in another embodiment the invention provides a method of formulating a pharmaceutically stable solution of palonosetron comprising admixing a) palonosetron or a pharmaceutically acceptable salt
30 thereof; and b) a pharmaceutically acceptable carrier, at a pH from about 4.0 to about 6.0. In alternative embodiments, the pH is from about 4.5 to about 5.5, and most

optimally about 5.0. There are many examples to those of skill in the art of suitable solutions to adjust the pH of a formulation. Two exemplary solutions are sodium hydroxide and hydrochloric acid solution, either of which could be used to adjust the pH of the formulation.

5 In another embodiment the invention provides a pharmaceutically stable solution for preventing or reducing emesis comprising from about 0.01 to about 5.0 mg/ml palonosetron or a pharmaceutically acceptable salt thereof and (i) from about 10 to about 100 millimoles citrate buffer, and/or (ii) from about 0.005 to about 1.0 mg/ml EDTA. Similarly, in another embodiment the invention provides a method of formulating a
10 pharmaceutically stable solution of palonosetron comprising admixing from about 0.01 to about 5.0 mg/ml palonosetron or a pharmaceutically acceptable salt thereof and (i) from about 10 to about 100 millimoles citrate buffer, and/or (ii) from about 0.005 to about 1.0 mg/ml EDTA. The citrate buffer can be in the form of citric acid and/or a salt of citric acid such as trisodium citrate. In various embodiments, the ranges of one or more of the
15 foregoing ingredients can be modified as follows:

- The formulation may comprise palonosetron or a pharmaceutically acceptable salt thereof in a concentration from about 0.02 mg/mL to about 1.0 mg/mL, from about 0.03 mg/mL to about 0.2 mg/mL palonosetron hydrochloride, and most optimally about 0.05 mg/ml.
- 20 • The formulation may comprise citrate buffer in a concentration of from about 10 to about 40 millimoles, or 15-30 millimoles.
- The formulation may comprise EDTA in a concentration of from about 0.005 mg/ml to about 1.0 mg/ml, or about 0.3 to about 0.7 mg/ml, and most optimally about 0.5 mg/ml.

25 The inventors have further discovered that the addition of mannitol and a chelating agent can increase the stability of palonosetron formulations. Therefore, in still another embodiment the invention provides a pharmaceutically stable solution for preventing or reducing emesis comprising a) palonosetron or a pharmaceutically acceptable salt thereof and b) a pharmaceutically acceptable carrier, wherein the
30 pharmaceutically acceptable carrier comprises a chelating agent and mannitol. Similarly,

in another embodiment the invention provides a method of formulating a pharmaceutically stable solution of palonosetron comprising admixing a) palonosetron or a pharmaceutically acceptable salt thereof and b) a pharmaceutically acceptable carrier, wherein the pharmaceutically acceptable carrier comprises a chelating agent and mannitol. The chelating agent is preferably EDTA, and, in various embodiments the chelating agent is present in a concentration of from about 0.005 to about 1.0 mg/mL or from about 0.05 mg/mL to about 1.0 mg/mL or from about 0.3 to about 0.7 mg/ml, or most optimally about 0.5 mg/ml. In various embodiments the mannitol is present in a concentration of from about 10.0 mg/ml to about 80.0 mg/ml, from about 20.0 mg/mL to about 60.0 mg/ml, or from about 40.0 to about 45.0 mg/ml.

Injectable formulations are typically formulated as aqueous solutions in which water is the primary excipient. Oral formulations will differ from injectable formulations generally by the additional presence of flavoring agents, coloring agents, or viscosity agents. Natural or synthetic sweeteners include, among others, mannitol, sorbitol, saccharose, saccharine, aspartame, acelsulphame K, or cyclamate. These agents are generally present in concentrations in excess of 100 mg/ml or 250 mg/ml when used as sweetening agents, in contrast to the 41.5 mg/ml concentration of mannitol described in some of the embodiments of the invention, in which mannitol is acting simply as a tonicifying agent.

The formulations of the present invention are particularly suited for use in injectable and oral liquid formulations, but it will be understood that the solutions may have alternative uses. For example, they may be used as intermediates in the preparation of other pharmaceutical dosage forms. Similarly, they may have other routes of administration including intranasal or inhalation. Injectable formulations may take any route including intramuscular, intravenous or subcutaneous.

Still further embodiments relate to improvements in the ease with which the palonosetron formulation can be stored or manufactured. In particular, the inventors have discovered that the formulations of the present invention allow storage of the product for extended periods at room temperature. Thus, in yet another embodiment the invention provides a method of storing one or more containers in which are contained a solution of

palonosetron or a pharmaceutically acceptable salt thereof comprising: a) providing a room comprising said one or more containers; b) adjusting or maintaining the temperature of the room at greater than about ten, 15, or 20 degrees celcius; and c) storing said containers in said room for one month, 3 months, 6 months, one year, 18
5 months, 24 months or more (but preferably not exceeding 36 months), wherein (i) the palonosetron or pharmaceutical salt thereof is present in a concentration of from about 0.01 mg/mL to about 5.0 mg/mL, (ii) the pH of the solution is from about 4.0 to about 6.0, (iii) the solution comprises from about 0.01 to about 5.0 mg/ml palonosetron or a pharmaceutically acceptable salt thereof, from about 10 to about 100 millimoles citrate
10 buffer and from about 0.005 to about 1.0 mg/ml EDTA, (iv) the solution comprises a chelating agent, or (v) the solution comprises from about 10 to about 100 milliMoles of a citrate buffer.

The stability of the foregoing formulations also lends itself well to terminal sterilization processes in the manufacturing process. Therefore, in still another
15 embodiment the invention provides a method of filling a container in which is contained a solution of palonosetron or a pharmaceutically acceptable salt thereof comprising: a) providing one or more sterile open containers (preferably 5 ml. vials); b) filling said containers with a solution of palonosetron in a non-aseptic environment; c) sealing said filled containers; and d) sterilizing said sealed, filled containers, wherein (i) the
20 palonosetron or pharmaceutical salt thereof is present in a concentration of from about 0.01 mg/mL to about 5 mg/mL, (ii) the pH of the solution is from about 4.0 to about 6.0, (iii) the solution comprises from about 0.01 to about 5.0 mg/ml palonosetron or a pharmaceutically acceptable salt thereof, from about 10 to about 100 millimoles citrate buffer and from about 0.005 to about 1.0 mg/ml EDTA, (iv) the solution comprises a
25 chelating agent, or (v) the solution comprises from about 10 to about 100 milliMoles of a citrate buffer.

EXAMPLES**EXAMPLE 1: STABILIZING PH**

A study was conducted to determine the effect of pH on formulations containing palonosetron hydrochloride, measuring the stability at 80°C at pH 2.0, 5.0, 7.4, and 10.0.
5 The results indicated that palonosetron hydrochloride is most stable at pH 5.0.

EXAMPLE 2: STABILIZING CONCENTRATION RANGES

A formulation optimization study was performed using an experimental design software. Twenty-four lots of drug product were analyzed to investigate the appropriate concentration ranges for palonosetron hydrochloride (0.05 mg/mL to 5.0 mg/mL), citrate
10 buffer (0 to 80 mM) and EDTA (0 to 0.10%). The level of EDTA and citrate buffer were selected based on the optimal formulation, which was shown to be formulated with EDTA 0.05% and 20 mM citrate buffer at pH 5.0. The results of this study indicated that palonosetron concentration was also a critical factor in chemical stability, with greatest stability seen at the lowest palonosetron concentrations.

EXAMPLE 3: TONICIFYING AGENT

Formulations of palonosetron hydrochloride in citrate buffer were prepared including either a) sodium chloride or b) mannitol. The palonosetron hydrochloride formulation including mannitol showed superior stability. The optimum level of mannitol required for an isotonic solution was found to be 4.15%.

EXAMPLE 4: FORMULATION I

20 The following is a representative pharmaceutical formulation containing palonosetron that is useful for intravenous formulations, or other liquid formulations of the drug.

Ingredient	mg/mL
Palonosetron Hydrochloride	0.05*
Mannitol	41.5

EDTA	0.5
Trisodium citrate	3.7
Citric acid	1.56
WFJ	q.s. to 1 ml
Sodium hydroxide solution and/or hydrochloric acid solution	pH 5.0 $\pm$ 0.5

* calculated as a free base

EXAMPLE 5: FORMULATION II

The following is a representative pharmaceutical formulation containing palonosetron that is useful for oral formulations, or other liquid formulations of the drug.

5

Ingredient	mg/mL
Palonosetron Hydrochloride	0.05*
Mannitol	150
EDTA	0.5
Trisodium citrate	3.7
Citric acid	1.56
WFJ	q.s. to 1 ml
Sodium hydroxide solution and/or hydrochloric acid solution	pH 5.0 $\pm$ 0.5
Flavoring	q.s.

* calculated as a free base

EXAMPLE 6 — STABILITY OF PALONOSETRON WITHOUT DEXAMETHASONE

The physical and chemical stability of palonosetron HCl was studies in concentrations of 5 μ g/mL and 30 μ g/mL in 5% dextrose injection, 0.9% sodium chloride injection, 5% dextrose in 0.45% sodium chloride injection, and dextrose 5% in lactated Ringer's injection. The admixtures were evaluated over 14 days at 4 °C in the dark and for 48 hours at 23 °C under fluorescent light.

10

Test samples of palonosetron HCl were prepared in polyvinyl chloride (PVC) bags of the infusion solutions at concentrations of 5 and 30 µg/mL. Evaluations for physical and chemical stability were performed on samples taken initially and after 1, 3, 5, 7, and 14 days of storage at 4 °C and after 1, 4, 24, and 48 hours at 23 °C. Physical stability was assessed using visual observation in normal room light and using a high-intensity monodirectional light beam. In addition, turbidity and particle content were measured electronically. Chemical stability of the drug was evaluated by using a stability-indicating high performance liquid chromatographic (HPLC) analytical technique.

All samples were physically stable throughout the study. The solution remained clear, and little or no change in particulate burden and haze level were found. Additionally, little or no loss of palonosetron HCl occurred in any of the samples at either temperature throughout the entire study period.

EXAMPLE 7 -- STABILITY OF PALONOSETRON WITH DEXAMETHASONE

The physical and chemical stability of palonosetron HCl 0.25 mg admixed with dexamethasone (as sodium phosphate) 10 mg or 20 mg in 5% dextrose injection or 0.9% sodium chloride injection in polyvinyl chloride (PVC) minibags, and also admixed with dexamethasone (as sodium phosphate) 3.3 mg in 5% dextrose injection or 0.9% sodium chloride injection in polypropylene syringes at 4 °C in the dark for 14 days and at 23 °C exposed to normal laboratory fluorescent light over 48 hours, was studied.

Test samples of palonosetron HCl 5 µg/mL with dexamethasone (as sodium phosphate) 0.2 mg/mL and also 0.4 mg/mL were prepared in polyvinyl chloride (PVC) minibags of each infusion solution. Additionally, palonosetron HCl 25 µg/mL with dexamethasone (as sodium phosphate) 0.33 mg/mL in each infusion solution were prepared as 10 mL of test solution in 20-mL polypropylene syringes. Evaluations for

physical and chemical stability were performed on samples taken initially and after 1, 3, 7, and 14 days of storage at 4 °C and after 1, 4, 24, and 48 hours at 23 °C. Physical stability was assessed using visual observation in normal room light and using a high-intensity monodirectional light beam. In addition, turbidity and particle content were
5 measured electronically. Chemical stability of the drug was evaluated by using a stability-indicating high performance liquid chromatographic (HPLC) analytical technique.

All samples were physically compatible throughout the study. The solutions remained clear, and little or no change in particulate burden and haze level were found.
10 Additionally, little or no loss of palonosetron HCl and dexamethasone occurred in any of the samples at either temperature throughout the entire study period.

This invention has been described with reference to its preferred embodiments. Variations and modifications of the invention will be obvious to those skilled in the art
15 from the foregoing detailed description of the invention.

CLAIMS

What is claimed is:

- 1) A method of storing one or more containers in which are contained a solution of
 5 palonosetron or a pharmaceutically acceptable salt thereof comprising:
 - a) providing a room comprising said one or more containers;
 - b) adjusting or maintaining the temperature of the room at greater than about ten
 degrees celcius;
 - c) storing said containers in said room for one month or more,
 10 wherein (i) the palonosetron or pharmaceutical salt thereof is present in an intravenous
 formulation at a concentration of from about 0.01 mg/mL to about 5.0 mg/mL, (ii) the pH
 of the solution is from about 4.0 to about 6.0, (iii) the solution comprises from about 0.01
 to about 5.0 mg/ml palonosetron or a pharmaceutically acceptable salt thereof, from
 about 10 to about 100 millimoles citrate buffer and from about 0.005 to about 1.0 mg/ml
 15 EDTA, (iv) the solution comprises mannitol, or (v) the solution comprises from about 10
 to about 100 milliMoles of a citrate buffer.
- 2) The method of claim 1 comprising storing said containers in said room for three months
 or more.
- 3) The method of claim 1 wherein the palonosetron or pharmaceutically acceptable salt
 20 thereof is present in an intravenous formulation at a concentration of from about 0.01
 mg/mL to about 5.0 mg/mL.
- 4) The method of claim 1 wherein the pH of the solution is from about 4.0 to about 6.0.
- 5) The method of claim 1 wherein the solution comprises from about 0.01 to about 5.0
 mg/ml palonosetron or a pharmaceutically acceptable salt thereof, from about 10 to about
 25 100 millimoles citrate buffer and from about 0.005 to about 1.0 mg/ml EDTA.
- 6) The method of claim 1 wherein the solution comprises mannitol.
- 7) The method of claim 1 wherein the solution comprises from about 10 to about 100
 milliMoles of a citrate buffer.
- 8) The method of claim 1 wherein the solution comprises a chelating agent.

- 9) The method of claim 1 wherein the solution comprises palonosetron or pharmaceutically acceptable salt thereof at a concentration of from about 0.01 mg/mL to about 5.0 mg/mL at a pH of from 4 to 6.
- 10) A method of storing one or more containers in which are contained an isotonic sterile shelf stable intravenous solution of palonosetron or a pharmaceutically acceptable salt thereof comprising:
 - a) providing a room comprising said one or more containers;
 - b) adjusting or maintaining the temperature of the room at greater than about ten degrees celcius;
 - 10 c) storing said containers in said room for one month or more, wherein (i) the palonosetron or pharmaceutical salt thereof is present at a concentration of from about 0.01 mg/mL to about 5.0 mg/mL, (ii) the pH of the solution is from about 4.0 to about 6.0, (iii) the solution comprises from about 0.01 to about 5.0 mg/ml palonosetron or a pharmaceutically acceptable salt thereof, from about 10 to about 100 millimoles citrate buffer and from about 0.005 to about 1.0 mg/ml EDTA, (iv) the solution comprises mannitol, or (v) the solution comprises from about 10 to about 100 milliMoles of a citrate buffer.
- 11) The method of claim 10 comprising storing said containers in said room for three months or more.
- 20 12) The method of claim 10 wherein the palonosetron or pharmaceutically acceptable salt thereof is present in an intravenous formulation at a concentration of from about 0.01 mg/mL to about 5.0 mg/mL.
- 13) The method of claim 10 wherein the pH of the solution is from about 4.0 to about 6.0.
- 14) The method of claim 10 wherein the solution comprises from about 0.01 to about 5.0 mg/ml palonosetron or a pharmaceutically acceptable salt thereof, from about 10 to about 100 millimoles citrate buffer and from about 0.005 to about 1.0 mg/ml EDTA.
- 25 15) The method of claim 10 wherein the solution comprises mannitol.
- 16) The method of claim 10 wherein the solution comprises from about 10 to about 100 milliMoles of a citrate buffer.
- 30 17) The method of claim 10 wherein the solution comprises a chelating agent.

- 18) The method of claim 10 wherein the solution comprises palonosetron or pharmaceutically acceptable salt thereof at a concentration of from about 0.01 mg/mL to about 5.0 mg/mL at a pH of from 4 to 6.
- 19) A method of filling a container in which is contained a solution of palonosetron or a pharmaceutically acceptable salt thereof comprising:
- a) providing one or more sterile open containers;
 - b) filling said containers with a solution of palonosetron in a non-aseptic environment
 - c) sealing said filled containers; and
 - d) sterilizing said sealed, filled containers,
- wherein (i) the palonosetron or pharmaceutical salt thereof is present in an intravenous formulation at a concentration of from about 0.01 mg/mL to about 5.0 mg/mL, (ii) the pH of the solution is from about 4.0 to about 6.0, (iii) the solution comprises from about 0.01 to about 5.0 mg/ml palonosetron or a pharmaceutically acceptable salt thereof, from about 10 to about 100 millimoles citrate buffer and from about 0.005 to about 1.0 mg/ml EDTA, (iv) the solution comprises mannitol, or (v) the solution comprises from about 10 to about 100 milliMoles of a citrate buffer.
- 20) The method of claim 19 wherein the palonosetron or pharmaceutically acceptable salt thereof is present in an intravenous formulation at a concentration of from about 0.01 mg/mL to about 5.0 mg/mL.
- 21) The method of claim 19 wherein the pH of the solution is from about 4.0 to about 6.0.
- 22) The method of claim 19 wherein the solution comprises from about 0.01 to about 5.0 mg/ml palonosetron or a pharmaceutically acceptable salt thereof, from about 10 to about 100 millimoles citrate buffer and from about 0.005 to about 1.0 mg/ml EDTA.
- 23) The method of claim 19 wherein the solution comprises mannitol.
- 24) The method of claim 19 wherein the solution comprises from about 10 to about 100 milliMoles of a citrate buffer.
- 25) The method of claim 19 wherein the solution comprises a chelating agent.
- 26) The method of claim 19 wherein the solution comprises palonosetron or pharmaceutically acceptable salt thereof at a concentration of from about 0.01 mg/mL to about 5.0 mg/mL at a pH of from 4 to 6.

- 27) The method of claim 19 wherein said solution has a shelf stability when stored under ambient conditions of twelve months or more.
- 28) A method of filling a container in which is contained an isotonic sterile intravenous solution of palonosetron or a pharmaceutically acceptable salt thereof comprising:
- 5 a) providing one or more sterile open containers;
 - b) filling said containers with a solution of palonosetron in a non-aseptic environment
 - c) sealing said filled containers; and
 - d) sterilizing said sealed, filled containers,
- 10 wherein (i) the palonosetron or pharmaceutical salt thereof is present in an intravenous formulation at a concentration of from about 0.01 mg/mL to about 5.0 mg/mL, (ii) the pH of the solution is from about 4.0 to about 6.0, (iii) the solution comprises from about 0.01 to about 5.0 mg/ml palonosetron or a pharmaceutically acceptable salt thereof, from about 10 to about 100 millimoles citrate buffer and from about 0.005 to about 1.0 mg/ml
- 15 EDTA, (iv) the solution comprises mannitol, or (v) the solution comprises from about 10 to about 100 milliMoles of a citrate buffer.
- 29) The method of claim 28 wherein the palonosetron or pharmaceutically acceptable salt thereof is present in an intravenous formulation at a concentration of from about 0.01 mg/mL to about 5.0 mg/mL.
- 20 30) The method of claim 28 wherein the pH of the solution is from about 4.0 to about 6.0.
- 31) The method of claim 28 wherein the solution comprises from about 0.01 to about 5.0 mg/ml palonosetron or a pharmaceutically acceptable salt thereof, from about 10 to about 100 millimoles citrate buffer and from about 0.005 to about 1.0 mg/ml EDTA.
- 32) The method of claim 28 wherein the solution comprises mannitol.
- 25 33) The method of claim 28 wherein the solution comprises from about 10 to about 100 milliMoles of a citrate buffer.
- 34) The method of claim 28 wherein the solution comprises a chelating agent.
- 35) The method of claim 28 wherein the solution comprises palonosetron or pharmaceutically acceptable salt thereof at a concentration of from about 0.01 mg/mL to about 5.0 mg/mL
- 30 at a pH of from 4 to 6.

- 36) The method of claim 28 wherein said solution has a shelf stability when stored under ambient conditions of twelve months or more.