



Office de la Propriété

Intellectuelle
du Canada

Un organisme
d'Industrie Canada

Canadian
Intellectual Property
Office

An agency of
Industry Canada

CA 2633730 C 2014/09/09

(11)(21) **2 633 730**

(12) **BREVET CANADIEN**
CANADIAN PATENT

(13) **C**

(86) Date de dépôt PCT/PCT Filing Date: 2006/12/21
(87) Date publication PCT/PCT Publication Date: 2007/07/05
(45) Date de délivrance/Issue Date: 2014/09/09
(85) Entrée phase nationale/National Entry: 2008/06/18
(86) N° demande PCT/PCT Application No.: US 2006/049061
(87) N° publication PCT/PCT Publication No.: 2007/076027
(30) Priorité/Priority: 2005/12/22 (US60/752,979)

(51) Cl.Int./Int.Cl. *A01N 53/08* (2006.01),
A01N 25/30 (2006.01), *A01N 53/06* (2006.01),
A01P 7/04 (2006.01)

(72) Inventeurs/Inventors:
STAETZ, CHARLES A., US;
YANG, HUI S., US;
GARCIA, HYLSA, US

(73) Propriétaire/Owner:
FMC CORPORATION, US

(74) Agent: SMART & BIGGAR

(54) Titre : FORMULATIONS DE BIFENTHRINE ET DE CYPERMETHRINE ENRICHIE

(54) Title: FORMULATIONS OF BIFENTHRIN AND ENRICHED CYPERMETHRIN

(57) Abrégé/Abstract:

The present invention is directed to novel insecticidal compositions comprising bifenthrin and enriched cypermethrin, the compositions are physically stable when diluted with water.



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
5 July 2007 (05.07.2007)

PCT

(10) International Publication Number
WO 2007/076027 A2(51) International Patent Classification:
A01N 35/04 (2006.01)

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(21) International Application Number:
PCT/US2006/049061(22) International Filing Date:
21 December 2006 (21.12.2006)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/752,979 22 December 2005 (22.12.2005) US

(71) Applicant (for all designated States except US): FMC CORPORATION [US/US]; 1735 Market Street, Philadelphia, PA 19103 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): STAETZ, Charles, A. [US/US]; 85 Hickory Lane, Newtown, PA 18940 (US). YANG, Hui, S. [CN/US]; 6 Daisy Court, Plainsboro, NJ 08536 (US). GARCIA, Hylsa [US/US]; 918 McLain Street, Elizabeth, NJ 07202 (US).

(74) Agents: FAIR, Paul, A. et al.; FMC CORPORATION, 1735 Market Street, Philadelphia, PA 19103 (US).

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 2007/076027 A2

(54) Title: NOVEL FORMULATIONS OF BIFENTHRIN AND ENRICHED CYPERMETHRIN

(57) Abstract: The present invention is directed to novel insecticidal compositions comprising bifenthrin and enriched cypermethrin, the compositions are physically stable when diluted with water.

76785-46

FORMULATIONS OF BIFENTHRIN AND ENRICHED CYPERMETHRIN

5

FIELD OF THE INVENTION

10 The present invention relates to the field of insecticides and insecticidal compositions. In particular, the invention relates to novel insecticidal compositions comprising bifenthrin and enriched cypermethrin that are physically stable when diluted with water.

15 BACKGROUND OF THE INVENTION

To enable the efficient elimination or controlling of unwanted insects in agricultural and other applications, it is desirable to use effective chemical insecticides on these unwanted pests. Formulations containing multiple insecticides are desirable in order to broaden the spectrum of economically important insect, 20 mite, and other pest species killed or controlled and take advantage of the individual pesticidal characteristics of each of the active ingredients.

Compositions containing two or more insecticides have been practiced in the art, but problems with the physical stability of such mixtures in water have caused application and efficacy issues. When traditional insecticidal compositions are 25 combined, the combined components (surfactants, viscosity modifiers, wetting agents) of both may cause accelerated physical degradation (phase separation) of the mixture when diluted in low to moderately hard water. This physical degradation can occur in the mix tanks prior to application on plants or another locus where control is desired. Often this problem goes unnoticed and a uniform application of 30 the insecticide mixture is not achieved, yielding inadequate efficacy.

Typically, in commercial applications, the insecticidal formulation comprises less than one percent of the tank mixture to keep the cost of the formulation low, while still yielding spectrum and onset of visual symptomology benefits. The

76785-46

physical stability of the formulation when diluted with water is a key problem in the art.

SUMMARY OF THE INVENTION

In accordance with the present invention, novel liquid formulations comprising bifenthrin and enriched cypermethrin have been found which have significantly improved the 5 physical stability when diluted in water.

In one composition aspect, the invention relates to an insecticidal composition, comprising: (a) bifenthrin; (b) an enriched cypermethrin; (c) one or more aromatic or aliphatic solvents; (d) acetic acid; (e) a severely solvent refined light and heavy paraffinic petroleum oil; and (f) a surfactant blend, comprising: (i) an alkylbenzene sulfonate salt, (ii) an 10 ethoxylated castor oil, and (iii) a polyethylene glycol fatty acid ester or ethoxylated fatty acid, which composition is in the form of an emulsifiable concentrate.

In a further composition aspect, the invention relates to an insecticidal composition, comprising: (a) bifenthrin and *zeta*-cypermethrin present in a weight ratio of from 1/4 to 4/1 and in a total concentration of from 0.05% to 30%; (b) an aromatic solvent 15 present in a concentration of from 40% to 55%; (c) acetic acid present in a concentration of from 0.01% to 0.15%; (d) a severely solvent refined light and heavy paraffinic petroleum oil present in a concentration of from 20% to 30%; and (e) a surfactant blend, consisting of: (a) an alkylbenzene sulfonate salt present in a concentration of from 1.5% to 4.5%, (b) an ethoxylated castor oil in a concentration of from 1.5% to 4.5%, and (c) a polyethylene glycol 20 fatty acid ester present in a concentration of from 0.10% to 0.60%, wherein all % are % by weight based on the total weight of the composition, which composition is in the form of an emulsifiable concentrate.

76785-46

DETAILED DESCRIPTION OF THE INVENTION

The present invention is directed to novel insecticidal compositions comprising bifenthrin and enriched cypermethrin in an insecticidal formulation having significantly improved physical stability when the composition is diluted in water.

15 *zeta*-Cypermethrin is a potent and quick acting insecticide, which controls a broad spectrum of chewing, sucking and flying insects. In addition to controlling chewing, sucking and flying insects, the pyrethroid bifenthrin is also active against a number of key mite pests and exhibits a longer residual activity than *zeta*-cypermethrin.

The term "bifenthrin" means 2-methylbiphenyl-3-ylmethyl (Z)-(1RS)-*cis*-3-(2-chloro-3,3,3-trifluoroprop-1-enyl)-2,2-dimethyl cyclopropanecarboxylate.

20 The term "cypermethrin" means the synthetic pyrethroid (R,S)- α -cyano-3-phenoxybenzyl-(1RS)-*cis-trans*-3-(2,2-dichlorovinyl)-2,2-dimethylcyclopropanecarboxylate which consists of a mixture of eight isomers, each of which are present in approximately the same amounts. Since the discovery and commercialization of cypermethrin, products containing increased amounts of 25 certain isomers have been developed. As used herein, the term "enriched cypermethrin" refers to the following:

30 *alpha*-cypermethrin which is a racemate comprising (S)- α -cyano-3-phenoxybenzyl (1R)-*cis*-3-(2,2-dichlorovinyl)-2,2-dimethylcyclopropanecarboxylate and (R)- α -cyano-3-phenoxybenzyl (1S)-*cis*-3-(2,2-dichlorovinyl)-2,2-dimethylcyclopropanecarboxylate;

beta-cypermethrin which is a reaction mixture comprising two enantiomeric pairs in a ratio of about 2:3 (S)- α -cyano-3-phenoxybenzyl (1R)-*cis*-3-(2,2-

dichlorovinyl)-2,2-dimethylcyclopropanecarboxylate and (R)- α -cyano-3-phenoxylbenzyl (1S)-*cis*-3-(2,2-dichlorovinyl)-2,2-dimethylcyclopropanecarboxylate with (S)- α -cyano-3-phenoxylbenzyl (1R)-*trans*-3-(2,2-dichlorovinyl)-2,2-dimethylcyclopropanecarboxylate and (R)- α -cyano-3-phenoxylbenzyl (1S)-*trans*-3-(2,2-dichlorovinyl)-2,2-dimethylcyclopropanecarboxylate;

theta-cypermethrin which is a 1:1 mixture of the enantiomers (R)- α -cyano-3- phenoxybenzyl (1S)-*trans*-3-(2,2-dichlorovinyl)-2,2- dimethylcyclopropanecarboxylate and (S)- α -cyano-3-phenoxybenzyl (1R)-*trans*-3- (2,2-dichlorovinyl)-2,2-dimethylcyclopropanecarboxylate; and

A particular form of “*zeta*-cypermethrin” is (R,S)- α -cyano-3-phenoxybenzyl-(1RS)-*cis-trans*-3-(2,2-dichlorovinyl)-2,2-dimethylcyclopropanecarboxylate which has been enriched in the 1R-*cis*-S and 1R-*trans*-S isomers by the processes disclosed in patents U. S. 5,164,411, U. S. 5,028,731 and U. S. 4,997,970. A most preferred form of “*zeta*-cypermethrin” is the cypermethrin isomer mixture prepared by the process disclosed in U. S. 4,997,970 starting with a 55/45 *cis/trans* mixture of cypermethrin with a catalytic amount of tricaprylylammonium chloride (Aliquat[®] 336, Aldrich Chemical Co.) and sodium carbonate in n-heptane. This process and the subsequent isolation procedure produces *zeta*-cypermethrin containing a small amount, usually 0.6% to 1.3%, of the catalyst. The presence of this catalyst adds to the complexity of producing a formulation that is stable when diluted with water.

In a preferred embodiment of the present invention, the bifenthrin and enriched cypermethrin insecticides are present in a combined concentration of from 0.01% by weight to 40% by weight, more particularly, from 0.05% by weight to 30% by weight based upon the total weight of all components in the composition. The ratio of bifenthrin active ingredient (AI) to enriched cypermethrin AI may be from 1/99 to 99/1. Preferably the ratio of bifenthrin AI to enriched cypermethrin AI is from 1/4 to 4/1. More preferably the ratio is from 1/3 to 3/1.

In one aspect, the present invention is directed to an insecticidal composition comprising bifenthrin and enriched cypermethrin, an aromatic or aliphatic solvent,

76785-46

an acid, severely solvent refined light and heavy paraffinic petroleum oil and a surfactant blend comprising an alkylbenzene sulfonate salt, an ethoxylated castor oil, and an ethoxylated fatty acid or a polyethylene glycol fatty acid ester, wherein said bifenthrin and enriched cypermethrin insecticides are present, collectively, in an 5 insecticidally effective amount.

Preferably, the aromatic solvent is selected from alkylated naphthalene aromatic and alkylated naphthalene depleted aromatic fluids, for example Aromatic 200 ND Fluid and Aromatic 200 Fluid (both available from ExxonMobil Chemicals). The aliphatic solvent is selected from de-aromatized hydrocarbon 10 fluids, for example, Exxsol D series fluids (available from ExxonMobile Chemicals), isoparaffinic fluids, for example Isopar fluids (available from ExxonMobile Chemicals), and hydrocarbon fluids with very high normal paraffin content, for example, Norpar fluids (available from ExxonMobile Chemicals). The aromatic or aliphatic solvent is present in a concentration of from 10% to 90% by 15 weight and preferably, 40% to 55% by weight, based on the total weight of all components in the composition. The most preferred solvent is an alkylated naphthalene depleted aromatic fluid present in a concentration of from 44% to 47%.

Preferably, the alkylbenzene sulfonate salt is the amine or calcium salt of a branched or linear alkylbenzene sulfonate. More preferred, the alkylbenzene 20 dodecylbenzene sulfonate salt is a branched dodecylbenzene sulfonate aminoethylethanolamine or calcium salt, for example, Agnique ABS 70AE and 25 Agnique ABS 60 BC (available from Cognis Corporation) and Rhodacal 70 (available from Rhodia Corporation). The dodecylbenzene sulfonate, or salt thereof, may be present in a concentration of from 1.5% to 4.5% by weight, based on the total weight of all components in the composition.

The ethoxylated castor oil may be one or more ethoxylated castor oils selected from the group of ethoxylated castor oils having an EO number of 8 to 50. Preferably the ethoxylated castor oil has an EO number of 15 to 40 and most 30 preferred an EO number of 20 to 40, for example, Agnique CSO-25 and Agnique CSO-36 (available from Cognis Corporation). The ethoxylated castor oil may be present in a concentration of from 1.5% to 4.5% by weight, based on the total weight of all the components in the composition.

76785-46

The ethoxylated fatty acid is preferably a C₉ to C₂₀ ethoxylated fatty acid, for example, stearic acid monooleate, stearic acid dioleate, and stearic acid monosterate. The polyethylene glycol fatty acid ester is selected from polyethylene glycol monooleates having an average M_n of 100 to 800, preferably an average M_n of 300 to 500, and most preferred, an average M_n of 400, for example, Agnique PEG 400MO (available from Cognis Corporation). The ethoxylated fatty acid or the polyethylene glycol fatty acid ester may be present in a concentration of from 0.10% to 0.60% by weight, based on the total weight of all the components in the composition.

10 The severely solvent refined light and heavy paraffinic petroleum oil is a mixture of from 80% to 100% by volume light paraffinic petroleum oil and from 0% to 20% by volume of heavy paraffinic petroleum oil, for example, Sunpray 6N (available from Sunoco, Inc.), Orchex 796 (available from ExxonMobile USA) and BVA Spray 10 (available from BVA Oils). The severely solvent refined light and heavy paraffinic petroleum oil mixture may be present in a concentration of from 20% to 30% by weight of all the components in the total formulation, preferably in a concentration of from 24% to 26% by weight, based on the total weight of all the components in the composition.

15 An acid is used to buffer the formulation in order to stabilize the zeta-cypermethrin from epimerizing to less active isomers. The acid is preferably acetic acid or glacial acetic acid and is present in a concentration of from 0.01% to 0.15% by weight, based on the total weight of all the components in the composition.

20 The insecticidal formulation may further comprise additional components such as an anti-freeze agent, an anti-foam agent and/or a biocide.

25 A preferred embodiment of this aspect of the invention is an insecticidal emulsifiable concentrate formulation wherein said bifenthrin and enriched cypermethrin insecticides are present in a ratio of 1/4 to 4/1 and a concentration of from 0.05% to 30%, the aromatic solvent is present in a concentration of from 40% to 55%, the acetic acid is present in a concentration of from 0.01% to 0.15%, the

30 severely solvent refined light and heavy paraffinic petroleum oil is present in a concentration of from 20% to 30%, the branched dodecylbenzene sulfonate salt is present in a concentration of from 1.5% to 4.5%, the ethoxylated castor oil or mixture of castor oils is present in a concentration of from 1.% to 4.5% and the

76785-46

polyethylene glycol fatty acid ester is present in a concentration of from 0.10% to 0.60% wherein all %'s are % by weight based upon the total weight of the formulation.

Another aspect of the present invention is directed to an insecticidal

5 composition comprising an aqueous mixture of bifenthrin and enriched cypermethrin, an aromatic solvent, one or more nonionic polymeric surfactants, an antifoam agent, a preservative and glycerine, wherein the bifenthrin and enriched cypermethrin insecticides are present, at least collectively, in an insecticidally effective amount.

10 The aromatic solvent is selected from alkylated naphthalene aromatic and alkylated naphthalene depleted aromatic fluids, for example Aromatic 200 ND Fluid and Aromatic 200 Fluid (both available from ExxonMobil Chemicals). The aromatic solvent is present in a concentration of from 5% to 30% by weight, preferably 12% to 15% by weight, based on the total weight of all components in the

15 composition.

Preferably, the nonionic polymeric surfactant is one or more surfactants selected from the group consisting of an alkyd polyethylene glycol, for example, TM Atlox 4914 (available from Uniqema Corporation) and a polyalkylene glycol ether, TM for example, Atlas G-5000 (available from Uniqema Corporation). Preferably the 20 nonionic polymeric surfactant is present in a concentration of from 3% to 20% by weight preferably from 8% to 12% by weight, based on the total weight of all components in the composition.

The antifoam agent is preferably a silicone containing antifoam agent, for example, TM Rhodorsil emulsion 1824 antifoam (available from Rhodia Corporation) 25 and Dow Corning AF Emulsion (available from Dow Corning Corporation). Preferably the antifoam agent is present in a concentration of from 0.001% to 1.5% by weight, based on the total weight of all the components in the composition.

The biocide is an isothiazolone mixture, for example, TM Kathon CG/ICP preservative or Legend MK preservative (available from Rohm and Haas 30 Corporation). Preferably the biocide is present in a concentration of from 0.001% to 1.5% by weight, based on the total weight of all the components in the composition.

The glycerine is preferably refined glycerine, for example, TM Agnique GLY 99-U glycerine (available from Cognis Corporation). Glycerine is present in a

concentration of from 3% to 15% by weight, preferably from 5% to 10% by weight, based on the total weight of all the components in the composition.

Water is used as a diluent and preferably is purified water, for example, deionized or distilled water, and is present in an amount that would dilute the active ingredients to a desired concentration. Preferably water is present in a concentration of from 40% to 60% by weight, based on the total weight of all the components in the composition.

A preferred embodiment of this aspect of the invention is an insecticidal concentrated aqueous emulsion formulation wherein said bifenthrin and enriched cypermethrin insecticides are present in a ratio of from 1/4 to 4/1 and in a concentration of from 0.05% to 30%, the aromatic solvent is present in a concentration of from 12% to 15%, nonionic polymeric surfactant is present in a concentration of from 8% to 12%, the antifoam agent is present in an amount of from 0.001% to 1.5%, the preservative is present in a concentration of from 0.001% to 1.5%, glycerine is present in a concentration of from 5% to 10% and water is present in a concentration of from 40% to 60%, wherein all %'s are % by weight based upon the total weight of all the components in the formulation.

The term "ambient temperature" as utilized herein shall generally mean any suitable temperature found in a laboratory or other working quarter, and is generally not below about 15° C nor above about 30° C.

The formulations of the present invention are further illustrated by the examples below. The examples serve only to illustrate the invention and should not be interpreted as limiting since further modifications of the disclosed invention will be apparent to those skilled in the art. All such modifications are deemed to be within the scope of the invention as defined in the claims.

Example 1

This example illustrates one protocol for the preparation of a 2/1 ratio of bifenthrin to *zeta*-cypermethrin emulsifiable concentrate formulation of the present invention.

30

To 42.32 grams of Aromatic 200 ND was added 8.34 grams of melted bifenthrin (95.9% active ingredient) and 11.11 grams of *zeta*-cypermethrin (36% active ingredient, prepared by the process disclosed in U. S. Patent 4,997,970). The

mixture was stirred at ambient temperature with a mechanical stirrer until a homogenous solution formed, at which time 2.52 grams of a branched dodecylbenzene sulfonate salt (Agnique ABS 70AE), 0.28 gram of polyethylene glycol monooleate (Agnique PEG 400MO) 0.84 gram of ethoxylated castor oil

5 (Agnique CSO-36), 1.96 grams ethoxylated castor oil (Agnique CSO-25), 20.00 grams of severely solvent refined light and heavy paraffinic petroleum oil (Sunspray 6N available) and 0.08 gram of acetic acid were added. Upon completion of the addition, the agitation was continued for 10 minutes to obtain a yellowish homogeneous solution.

10 The following Table 1 sets forth additional formulation examples prepared in the manner of Example 1.

TABLE 1

Bifenthrin and *Zeta*-Cypermethrin Formulations Containing, By Weight Of All The Components In The Total Formulation: 52.9% Aromatic 200, 25.0% Sunspray 6N

15 and 0.09% Acetic Acid

Formulation Code	AI*	Agnique ABS70AE	Agnique PEG400MO	Agnique CSO-36	Agnique CSO-25
	(g)	(g)	(g)	(g)	(g)
F2	0.31/0.42	0.0945	0.0105	0.0315	0.0735
F3	0.31/0.42	0.021	0.042	0.084	0.063
F4	0.31/0.42	0.063	0.042	0.084	0.021
F5	0.31/0.42	0.021	0.0315	0.084	0.0735
F6	0.31/0.42	0.042	0.0105	0.084	0.0735
F7	0.31/0.42	0.1155	0.042	0.0315	0.021
F8	0.31/0.42	0.021	0.042	0.0735	0.0735
F9	0.31/0.42	0.063	0.042	0.0315	0.0735
F10	0.31/0.42	0.147	0.0105	0.0315	0.021
F11	0.31/0.42	0.0945	0.0105	0.084	0.021
F12	1.82/16.42	1.68	0.56	1.867	1.493
F13	16.42/1.82	3.92	0.112	0.871	0.697
F14	9.12/9.12	1.68	0.112	2.116	1.692
F15	9.12/9.12	1.68	0.56	1.867	1.493
F16	16.42/1.82	1.68	0.112	2.116	1.692
F17	1.82/16.42	1.68	0.112	2.116	1.692
F18	9.12/9.12	3.92	0.112	0.871	0.697
F19	1.82/16.42	3.92	0.112	0.871	0.697
F20	9.12/9.12	3.92	0.56	0.622	0.498
F21	16.42/1.82	1.68	0.56	1.867	1.493
F22	16.42/1.82	3.92	0.56	0.622	0.498
F23	1.82/16.42	3.92	0.56	0.622	0.498

Formulation Code	AI*	Agnique ABS70AE	Agnique PEG400MO	Agnique CSO-36	Agnique CSO-25
	(g)	(g)	(g)	(g)	(g)

* Bifenthrin grams (95.5% AI)/*zeta*-cypermethrin grams (36% AI, prepared by the process disclosed in U. S. Patent 4,997,970)

5

EXAMPLE 2

Dilution stability studies were conducted using 2.5 mL of the formulation of Example 1 added to 47.5 mL of water with 342 ppm hardness in a 50 mL Nessler tube. The Nessler tube was sealed with a rubber stopper and the contents mixed by inverting the tube ten times. The Nessler tube was placed in a tube rack at ambient 10 temperature and the percent separation was recorded at 2.0 hours and 4.0 hours. The percent separation was calculated by first measuring the height of the separation, if any, then the total height of the mixture. Dividing the separation height by the total height and multiplying by 100 provides the percent separation. Table 2 sets forth the dilution stability percent separation of the formulations described in Example 1 and 15 Table 1. The formulations are identified by the formulation code corresponding to those in Example 1 and in Table 1.

As a control experiment, 6.0 grams of a commercially available formulation of bifenthrin (Capture 2EC® available from FMC Corporation) and 1.2 grams of a commercially available formulation of *zeta*-cypermethrin (Mustang Max 0.8EC® 20 available from FMC Corporation) were blended together in 1.2 grams of Aromatic 200. The concentration of AI's was similar to the concentration of AI's in formulation code F2. A dilution stability test was performed using 2.5 mL of the control blend in 47.5 mL of water containing 342 ppm hardness as described above. The dilution stability percent separation results of the control experiment are 25 presented in Table 2.

TABLE 2

Percent Separation of Formulation in Water Containing 342 ppm Hardness

Formulation Code	% Separation	
	2.0 Hr	4.0 Hr
F1	0	0
F2	0	0
F3	3	5
F4	3	6
F5	3	6
F6	3	7
F7	3	6
F8	4	6
F9	2	4
F10	6	10
F11	0	0
F12	0.5	4
F13	6	12
F14	0	10
F15	0	6
F16	0	8
F17	0.5	4
F18	10	16
F19	10	14
F20	10	18
F21	0	4
F22	14	16
F23	10	16
Control	47	71

The novel formulations of the present invention are superior in maintaining the physical stability of a mixture of bifenthrin and *zeta*-cypermethrin in dilution stability tests when compared to the control dilution stability test.

Example 3

This example illustrates one protocol for the preparation of a 3/1 ratio of bifenthrin to *zeta*-cypermethrin emulsifiable concentrate formulation of the present invention

To 45.64 grams of Aromatic 200 was added 11.84 grams of melted bifenthrin (95.9% active ingredient) and 10.42 grams of *zeta*-cypermethrin (36% active ingredient, prepared by the process disclosed in U. S. Patent 4,997,970). The mixture was stirred at ambient temperature with a mechanical stirrer until a homogenous solution formed at which time 3.48 grams of a branched dodecylbenzene sulfonate salt (Agnique ABS 70AE), 0.35 gram of polyethylene glycol monooleate (Agnique PEG 400M) 1.05 grams of ethoxylated castor oil

(Agnique CSO-36), 2.12 grams ethoxylated castor oil (Agnique CSO-25), 25.00 grams of severely solvent refined light and heavy paraffinic petroleum oil (Sunspray 6N) and 0.1 gram of glacial acetic acid was added. Upon completion of the addition, the agitation was continued until a yellowish homogeneous solution was 5 obtained.

Example 4

A dilution stability study was conducted using 2.5 mL of the formulation prepared in Example 3 added to 47.5 mL of water with 342 ppm hardness in a 50 10 mL Nessler tube. The Nessler tube was sealed with a rubber stopper and the contents mixed by inverting the tube ten times. A total of three Nessler tubes were prepared in this manner. One Nessler tube was placed in a tube rack maintained at 0°C, one in a tube rack maintained at ambient temperature and one in a tube rack maintained at 30 °C, all were maintained for 24 hours. The percent separation after 15 24 hours was 0% for each temperature.

Example 5

This example illustrates one protocol for the preparation of a 3/1 ratio of bifenthrin 20 to *zeta*-cypermethrin concentrated aqueous emulsion formulation of the present invention

To a vessel equipped with a mechanical stirrer and a homogenizer was added 0.10 gram of a silicone containing antifoam agent (Rhodorsil emulsion 1824 antifoam), 6.0 grams of a polyalkylene glycol ether nonionic polymeric surfactant 25 (Atlas G5000) and 49.15 grams of deionized water. The mixture was stirred and heated at 50°C to 55°C. While maintaining this temperature, the homogenizer was turned on and a warm (50°C to 55°C) mixture of 13.63 grams of alkylated naphthalene depleted aromatic solvent (Aromatic 200 ND), 8.42 grams of *zeta*-cypermethrin (36% active ingredient, prepared by the process disclosed in U. S. 30 Patent 4,997,970), 10.10 grams of bifenthrin (95.9% active ingredient) and 4.00 grams of alkyd polyethylene glycol nonionic surfactant (Atlox 4914) was added slowly. The mixture was homogenized until a particle size of less than three microns at 90% tile was obtained. The homogenizer was turned off and the stirred

mixture was allowed to cool to less than 40°C at which time 8.5 grams of refined glycerine (Agnique GLY99-U) and 0.10 grams of preservative (Legend MK) were added. The resulting mixture was stirred until a homogenous solution was obtained while cooling to ambient temperature.

5

Example 6

A dilution stability study was conducted using 2.5 mL of the formulation prepared in Example 5 added to 47.5 mL of water with 342 ppm hardness in a 50 mL Nessler tube. The Nessler tube was sealed with a rubber stopper and the 10 contents mixed by inverting the tube ten times. A total of three Nessler tubes were prepared in this manner. One Nessler tube was placed in a tube rack maintained at 0°C, one in a tube rack maintained at ambient temperature and one in a tube rack maintained at 30 °C, all were maintained for 24 hours. The percent separation after 24 hours was 0% for each temperature.

15

While this invention has been described with an emphasis upon preferred embodiments, it will be understood by those of ordinary skill in the art that variations of the preferred embodiments may be used and that it is intended that the invention may be practiced otherwise than as specifically described herein. 20 Accordingly, this invention includes all modifications encompassed within the spirit and scope of the invention as defined by the following claims.

76785-46

CLAIMS:

1. An insecticidal composition, comprising:
 - (a) bifenthrin;
 - (b) an enriched cypermethrin;
 - 5 (c) one or more aromatic or aliphatic solvents;
 - (d) acetic acid;
 - (e) a severely solvent refined light and heavy paraffinic petroleum oil; and
 - (f) a surfactant blend, comprising:
 - (i) an alkylbenzene sulfonate salt,
 - 10 (ii) an ethoxylated castor oil, and
 - (iii) a polyethylene glycol fatty acid ester or ethoxylated fatty acid,
- which composition is in the form of an emulsifiable concentrate.
2. An insecticidal composition of claim 1, wherein the enriched cypermethrin is selected from the group consisting of *alpha*-cypermethrin, *beta*-cypermethrin, *theta*-cypermethrin and *zeta*-cypermethrin.
- 15 3. An insecticidal composition of claim 2, wherein the enriched cypermethrin is *zeta*-cypermethrin.
4. An insecticidal composition of claim 3, wherein bifenthrin and *zeta*-cypermethrin are present in a combined amount of from 0.05% to 30% by weight based on the
20 total weight of the composition.
5. An insecticidal composition of claim 4, wherein the weight ratio of bifenthrin to *zeta*-cypermethrin is from 1/4 to 4/1.

76785-46

6. An insecticidal composition of any one of claims 1 to 5, wherein the aromatic solvent is selected from the group consisting of an alkylated naphthalene aromatic solvent and an alkylated naphthalene depleted aromatic solvent.
7. An insecticidal composition of any one of claims 1 to 6, wherein the aromatic solvent is present in a concentration of from 40% to 55% by weight based on the total weight of the composition.
8. An insecticidal composition of any one of claims 1 to 7, wherein the alkyl benzene sulfonate salt is selected from the group consisting of a branched dodecylbenzene sulfonate, calcium salt and a branched dodecylbenzene sulfonate, aminoethylethanolamine salt.
9. An insecticidal composition of any one of claims 1 to 8, wherein the alkylbenzene sulfonate salt is present in a concentration of from 1.5% to 4.5% by weight based upon the total weight of the composition.
10. An insecticidal composition of any one of claims 1 to 9, wherein the ethoxylated castor oil is an ethoxylated castor oil having an EO number of from 20 to 40.
11. An insecticidal composition of any one of claims 1 to 10, wherein the ethoxylated castor oil is present in a concentration of from 1.5% to 4.5% by weight based on the total weight of the composition.
12. An insecticidal composition of any one of claims 1 to 11, wherein the polyethylene glycol fatty acid ester is a polyethylene glycol monooleate glycol having an average molecular weight of from 300 to 500.
13. An insecticidal composition of any one of claims 1 to 12, wherein the polyethylene glycol fatty acid ester is present in a concentration of from 0.10% to 0.60% by weight based on the total weight of the composition.

76785-46

14. An insecticidal composition of any one of claims 1 to 13, wherein the severely solvent refined light and heavy paraffinic petroleum oil is present in a concentration of from 20% to 30% by weight based on the total weight of the composition.

15. An insecticidal composition of any one of claims 1 to 14, wherein the acetic acid is present in a concentration of from 0.10% to 0.15% by weight based on the total weight of the composition.

16. An insecticidal composition, comprising:

(a) bifenthrin and *zeta*-cypermethrin present in a weight ratio of from 1/4 to 4/1 and in a total concentration of from 0.05% to 30%;

10 (b) an aromatic solvent present in a concentration of from 40% to 55%;

(c) acetic acid present in a concentration of from 0.01% to 0.15%;

(d) a severely solvent refined light and heavy paraffinic petroleum oil present in a concentration of from 20% to 30%; and

(e) a surfactant blend, consisting of:

15 (a) an alkylbenzene sulfonate salt present in a concentration of

from 1.5% to 4.5%,

(b) an ethoxylated castor oil in a concentration of from 1.5% to 4.5%, and

(c) a polyethylene glycol fatty acid ester present in a concentration of from 0.10% to 0.60%,

20 wherein all % are % by weight based on the total weight of the composition,

which composition is in the form of an emulsifiable concentrate.

76785-46

17. A composition of any one of claims 1 to 16, further comprising at least one additional component selected from the group consisting of an anti-freeze agent, an anti-foam agent and a biocide.