



US00PP19793P3

(12) **United States Plant Patent**  
**Pait et al.**

(10) **Patent No.:** **US PP19,793 P3**

(45) **Date of Patent:** **Mar. 3, 2009**

(54) **LOBLOLLY PINE TREE NAMED ‘CF Q7766’**

(58) **Field of Classification Search** ..... Plt./213  
See application file for complete search history.

(50) Latin Name: *Pinus taeda*  
Varietal Denomination: **CF Q7766**

(56) **References Cited**

(75) Inventors: **John Pait**, Atlanta, GA (US); **Plamen Denchev**, Victoria (CA); **Stephen Attree**, Victoria (CA); **Robert J. Weir**, Cary, SC (US); **Andy Benowicz**, Victoria (CA)

PUBLICATIONS

[http://web.archive.org/web/20051130221944.http://www.chesapeakebay.net/info/liblolly.cfm](http://web.archive.org/web/20051130221944/http://www.chesapeakebay.net/info/liblolly.cfm) (2 pages).\*  
<http://www.cnr.vt.edu/DENDRO/dendrology/syllabus/fact-sheet.cfm> (1 page).\*  
U.S. Appl. No. 11/635,703, Pait et al., filed Dec. 7, 2006.  
United States Patent Office Action for U.S. Appl. No. 11/635,703 dated Jan. 18, 2008.

(73) Assignee: **CellFor, Inc.**, Victoria, British Columbia (CA)

\* cited by examiner

(\* ) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

*Primary Examiner*—Annette H Para

*Assistant Examiner*—S. B. McCormick-Ewoldt

(21) Appl. No.: **11/635,199**

(74) *Attorney, Agent, or Firm*—Michael Best & Friedrich LLP

(22) Filed: **Dec. 7, 2006**

(57) **ABSTRACT**

(65) **Prior Publication Data**

US 2008/0141406 P1 Jun. 12, 2008

A new and distinctive variety of a loblolly pine tree which has been denominated varietally as ‘CF Q7766’ which is distinguished by great resistance to fusiform rust and pitch canker, exceptionally high growth rate, excellent stem straightness, distinctive very long internodes, narrow crown.

(51) **Int. Cl.**  
**A01H 7/00** (2006.01)

(52) **U.S. Cl.** ..... **Plt./213**

**2 Drawing Sheets**

**1**

**2**

Latin name: *Pinus taeda*.  
Variety denomination: ‘CF Q7766’.

Coastal Plains, and Mid-Continent regions of the United States.

**BACKGROUND**

A new variety of loblolly pine tree (*Pinus taeda*), has been discovered. This selection has been designated as ‘CF Q7766.’

5

This new variety is a progeny of a second generation selection pollinated by a first generation selection. Female parent is an open pollinated progeny of Georgetown County, S.C. first generation selection. Male parent is a first generation selection made in Barnwell County S.C.

10

Cross pollination occurred in early 1998 followed by induction and cryopreservation of embryogenic tissue in 1999. First somatic seedlings were produced in 2000 and planted in early 2001 in seven field experiments. A total of 47 ramets (independent members of a clone) were planted ranging from 4 to 8 ramets per field experiment. The field experiments are located in Mississippi, Florida, Georgia and South Carolina.

15

The *Pinus taeda* plants of this variety were asexually propagated using an advanced form of micropopagation called somatic embryogenesis carried out at CellFor’s production facility in Victoria, Canada. Somatic embryogenesis uses a complex process which relies on the splitting of one embryo into many identical embryos. Somatic embryos can then be grown into plants which are all identical genetically. The asexual propagation occurs at an earlier stage in the plant’s life cycle than most other micropopagated plants. The detailed methods for somatic embryogenesis used for asexually propagating conifers in general are described in U.S. Pat. No. 6,372,496 and for loblolly pine in particular in U.S. patent application 2004/0203150.

**BRIEF DESCRIPTION OF THE DRAWINGS**

The drawings are color photographs showing the new variety of loblolly pine including the long internode length.

20

FIG. 1 is a photograph showing ‘CF Q7766’ ramet # 7 planted in Holt, Ga. The picture was taken after five field growing seasons. The picture shows distinctive long stem sections of clearwood between the whorls (long internodes), small to medium branch size (relative to the stem size), flat to medium angle between the branches and the stem and great stem straightness.

**BRIEF SUMMARY**

A new and distinct cultivar of loblolly pine (*Pinus taeda*) is distinctly characterized by great resistance to fusiform rust and pitch canker, exceptionally high growth rate, excellent stem straightness, distinctive very long internodes, narrow crown, and which is mature for commercial harvesting sooner than conventionally grown trees under the ecological conditions prevailing in the Piedmont, Atlantic and Gulf

25

30

FIG. 2 is a photograph showing ‘CF Q7766’ ramet # 4 planted in Winokur, Ga. The picture was taken after five field growing seasons. The picture shows superiority of growth, narrow tree crown relative to the size of this tree and the size

of surrounding neighbors, and distinctive long internodes seen particularly well in the upper crown.

#### DETAILED BOTANICAL DESCRIPTION

The botanical details of this new and distinct variety of loblolly pine tree follow. All color descriptions are made in reference to The Royal Horticultural Society (R.H.S.) Colour Chart (2005).

Parentage: Female parent: (Unknown) open pollinated progeny of a first generation selection of Georgetown County, S.C. Male parent: (Unknown) first generation selection of Barnwell County, S.C.

Compared to unimproved loblolly pine trees, 'CF Q7766' is characterized by exceptionally high growth rate, great resistance to fusiform rust (caused by *Cronartium quercuum* (Berk.) Miyabe ex Shirai f. sp. *fusiforme* (Cumm.) Burds. et Snow), great resistance to pitch canker (caused by *Fusarium circinatum* Nirenburg et O'Donnell), distinctive very long internodes, excellent stem straightness, narrow crown, small to medium branch diameter, flat to medium branch angle and low incidence of forking.

#### Disease resistance:

*Percent stem fusiform rust infection at age 5.*—0.

*Percent branch fusiform rust infection at age 5.*—5.

*Percent branch and stem fusiform rust infection at age 5.*—0.

*Percent dead ramets due to fusiform rust infection at age 5.*—0.

*Percent stem fusiform rust infection in the USDA Resistance Screening Center (Asheville, N.C.) tests after artificial inoculation with rust spores.*—5% (compared to 78% infection in unimproved seedlings).

*Percent of seedlings with over 50% of tissue infected by pitch canker in the USDA Resistance Screening Center (Asheville, N.C.) tests after artificial inoculation with pitch canker spores.*—1% (compared to 99% infection in unimproved seedlings).

#### Botanical description:

*Shape.*—A medium to large tree may reach well over 100 feet tall, self-prunes well and develops a fairly straight trunk and an oval, somewhat open crown.

*Average height.*—28 ft after 5 field growing seasons.

*Maximum height.*—34 ft after 5 field growing seasons.

*Average trunk diameter at breast height (4.5 feet above the soil level).*—4.7 inches after 5 field growing seasons, 6.7 inches to 7.1 inches after seven field growing seasons.

*Maximum trunk diameter at breast height (4.5 feet above the soil level).*—6.4 inches after 5 field growing seasons.

*Percent stem forking at age 4.*—2.

*Percent stem straightness gain over unimproved trees.*—15.

*Bark.*—Initially red- to gray-brown and scaly; older trees are ridged and furrowed, with somewhat rounded scaly plates. The RHS color is 200C in shade and 187D on exposure to sun.

*Branch.*—Orange-brown in color, fine to moderately stout; buds are narrowly ovoid, light reddish brown.

*Percent ramets with ramicorn branches at age 4.*—2.

*Sampling of branch characteristics.*—In order to sample branches from a consistent position from one tree to the next the following methodology was uti-

lized. From a point nine feet from the base of the tree, the first complete whorl of limbs below was labeled "Whorl One" and the first complete whorl of limbs above labeled "Whorl Two". This sampling point was chosen because it is the midpoint of the basal sixteen foot log of each tree. A complete whorl was defined as one with at least three branches. An average whorl contained 2–7 branches. All measurements were taken commencing on the South side of the tree and progressing anticlockwise around the stem. When more than three branches were available for measurement on the whorl the largest three branches, by basal diameter, were used for sampling. The following branch characteristics were measured after seven field growing seasons.

*Branch diameter.*—Diameter of each measured branch was taken at its base. Using a caliper the diameter of the branch, over bark, was measured to the closest 100<sup>th</sup> of an inch. 'CF Q7766' has an average branch diameter of 1.00 inches at the base of the branch. Zygotic seedlings of the same genetic origin have an average branch diameter of 1.24 inches at the base of the branch.

*Branch Angle.*—Utilizing a large protractor, the angle of each branch was measured as its deviation from horizontal. Branch angles were recorded for the portion of the branch emerging from the stem of the tree with data rounded to the closest 10 degrees. 'CF Q7766' has an average branch angle of 35.0 degrees from horizontal. Zygotic seedlings of the same genetic origin have an average branch angle of 43.3 degrees from horizontal.

*Branch length.*—The length of each sampled branch was measured directly with a graduated measurement pole. Branch lengths were recorded to the closest 0.5 feet. 'CF Q7766' has an average branch length of 6.48 feet. Zygotic seedlings of the same genetic origin have an average branch length of 9.17 feet.

*Crown diameter.*—The width of the crown, at the point where branch measurements were taken, was directly measured with the use of a graduated measurement pole. A radial measurement was taken on the East and West side of each tree. Crown radius was measured to the closest 0.5 feet. Crown width data is presented as diameter of the crown. 'CF Q7766' has an average crown diameter of 7.8 feet. Zygotic seedlings of the same genetic origin have an average crown diameter of 11.5 feet.

*Internode length.*—In proximity to the area of the stem utilized for branch measurements the mean internode length was determined for each tree. Internode distances for the calculation of the mean were directly measured from the stem of the tree using a graduated measurement pole. 'CF Q7766' has an average internode length of 3.12 feet. Zygotic seedlings of the same genetic origin have an average internode length of 2.25 feet.

*Leaf.*—Evergreen needles, 6 to 9 inches long, with (usually) three yellow-green needles per fascicle. The color of the foliage was measured at age 1 year and age 7 years and was found not to vary significantly with age. The color of the foliage was RHS 137A (40%) and 137C (60%). Diameter of the fascicle was  $\frac{9}{100}$  of an inch and the average sheath length is 6.75 mm.

*Flower.*—Monoecious; males long cylindrical, red to yellow, in clusters at branch tips; females yellow to purple.

*Fruit*.—Ovoid to cylindrical, 3 to 6 inch red-brown cones; umbo is armed with a short spine, maturing in early fall. Cones are sporadic in 5–7 year old plants.

*Propagation*.—Propagated by somatic embryogenesis.

*Seeds*.—none produced yet at 5–7 years of age, plants are not yet mature. Expected seed production by 12–15 years.

*Use*.—High yield industrial plantations

Although the new variety of loblolly pine tree possesses the detailed characteristics noted above as a result of the growing conditions prevailing in the seven test locations, it is to be understood that the variations of the usual magnitude and characteristics incident to changes in growing conditions, irrigation, fertilization, pruning, pest control, climatic variations and the like are to be expected. An example of ‘CF Q7766’ resulting from asexual reproduction can be found at Plum Creek Oliver year 2001 line trial, Screven county, Ga.

COMPARISON WITH PARENTS BY  
MICROSATELLITE ANALYSIS

Microsatellite markers were used to generate a unique DNA fingerprint for the variety. Vegetative buds and/or foliar material from eight individuals each produced by controlled crossing among parents for DNA fingerprinting. The DNA extraction protocol of Doyle and Doyle (1987) was used after slight modifications. DNA fingerprinting of parents and their offspring was initially conducted using a set of nine microsatellite markers (Auckland et. al. 2002) and a final set of five primer pairs were selected for the two lines mentioned above (see Table 1, for sequences and conditions of SSR primers). Primer selection was based on their ability to produce unique alleles and the presence of a high level of polymorphism.

TABLE 1

| Sequences and conditions of SSR primers currently used in loblolly pine. |  |                                   |                        |           |           |
|--|--|-----------------------------------|------------------------|-----------|-----------|
| Primer ID  | SEQUENCE (5'→3')   | LABEL TAIL (F/R); E (end labeled) | MgCl <sub>2</sub> (mM) | Tm (° C.) | Size (bp) |
|  |  |                                   |                        |           |           |
| PtTX 3034  | F: CACGACGTTGTAAAAC GACTCAAAATGCAAAAG ACG; SEQ ID NO: 3<br>R: ATTAGGACTGGGGATG AT; SEQ ID NO: 4          | F                                 | 1.5                    | 55        | 225       |
| PtTX 3049  | F: GAAGTGATAATGGCAT AGCAAAAT; SEQ ID NO: 5<br>R: GCAGACCCGTGAAAGT AATAAACAT; SEQ ID NO: 6                | R                                 | 3                      | 55        | 330       |
| PtTX 3105  | F: TGTCGGTGGAGTTGGC AGTAGACT; SEQ ID NO: 7<br>R: GCCCAGCGTTTCCTG; SEQ ID NO: 8                           | E                                 | 2                      | 59        | 280       |
| PtTX 3116  | F: CACGACGTTGTAAAAC GACTCCCAAAGCCTAAA GAAT; SEQ ID NO: 9<br>R: CATAACAAGGCCTTATC TTACAGAA; SEQ ID NO: 10 | F                                 | 2.5                    | 59        | 165       |

Microsatellite products were detected by M13 tailed primer (Oetting et al., 1995) or infra-red dye (IRD)-labeled primer. The amplification products were electrophoresed on 5.5% Long Ranger polyacrylamide gels using a LiCor 4200 automated sequencer (LiCor Inc., Lincoln, Nebr.). For each family, the female and male parents, as well as eight offspring were genotyped.

The observed parental genotypes and their expected offspring's genotypes at five studied SSR loci of each family are presented in Table 2.

TABLE 2

| Primer    | Parental genotypes and their expected offspring's genotypes at five different SSR loci of each family. |         |                             |         |                 |
|-----------|--|---------|-----------------------------|---------|-----------------|
|           | Genotype   |         | Expected offspring genotype |         |                 |
|           | Female   | Male    |                             |         |                 |
| PtTX 2146 | 181/199  | 193/208 | 181/193                     | 181/208 | 193/199 199/208 |
| PtTX 3105 | 166/184  | 169/172 | 166/169                     | 166/172 | 169/184 172/184 |
| PtTX 3034 | 226/228  | 216/226 | 216/226                     | 226/226 | 226/228         |
| PtTX 3049 | 301/303  | 301/311 | 301/301                     | 301/303 | 301/311         |
| PtTX 3116 | 151/163  | 151/160 | 151/151                     | 151/160 | 160/163         |

In general, offspring genotypes segregated following expected simple Mendelian segregation (see Table 3, for offspring multi-locus genotypes).

TABLE 3

| Parents and offspring genotypes at 5 different SSR loci for two loblolly pine full-sib families. |                        |          |                        |          |           |          |
|--|------------------------|----------|------------------------|----------|-----------|----------|
| Sample ID  | PtTX 3034 <sup>a</sup> |          | PtTX 3049 <sup>a</sup> |          | PtTX 3116 |          |
|  | Allele 1               | Allele 2 | Allele 1               | Allele 2 | Allele 1  | Allele 2 |
| Offspring Q 7766   | 226                    | 226      | 301                    | 303      | 151       |          |
| ID   | PtTX 3116              |          | PtTX 2146 <sup>a</sup> |          | PtTX 3105 |          |
|  | Allele 2               | Allele 1 | Allele 2               | Allele 1 | Allele 1  | Allele 2 |
| Offspring  | 151                    | 181      | 193                    | 169      | 184       |          |

<sup>a</sup>Allelic sizes have LiCor primer tails.

References:

Auckland, L., T. Bui, Y. Zhou, M. Shepherd and C. Williams. 2002. Conifer Microsatellite Handbook Corporate Press, Raleigh, N.C., USA.

Doyle, J. J. and J. L. Doyle. 1987. A rapid DNA isolation procedure for small quantities of fresh tissue. *Phytochemical bulletin* 19:11–15.

Oetting, W. S., H. K. Lee, D. J. Flanders, G. L. Wiesner, T. A. Sellers and R. A. King. 1995. Linkage analysis with multiplexed short tandem repeat polymorphisms using infrared fluorescence and M13 tailed primers. *Genomics* 30:450–458.

## SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 10

<210> SEQ ID NO 1  
 <211> LENGTH: 43  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Oligonucleotide PtTX 2146 - F

<400> SEQUENCE: 1

cacgacgttg taaaacgacc tggggatttg gattgggtat ttg 43

<210> SEQ ID NO 2  
 <211> LENGTH: 25  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Oligonucleotide PtTX 2146 - R

<400> SEQUENCE: 2

atattttcct tgccccttec agaca 25

<210> SEQ ID NO 3  
 <211> LENGTH: 36  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Oligonucleotide PtTX 3034 - F

<400> SEQUENCE: 3

cacgacgttg taaaacgact caaaatgcaa aagacg 36

<210> SEQ ID NO 4  
 <211> LENGTH: 18  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Oligonucleotide PtTX 3034 - R

<400> SEQUENCE: 4

attaggactg gggatgat 18

<210> SEQ ID NO 5  
 <211> LENGTH: 24  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Oligonucleotide PtTX 3049 - F

<400> SEQUENCE: 5

gaagtgataa tggcatagca aaat 24

<210> SEQ ID NO 6  
 <211> LENGTH: 25  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Oligonucleotide PtTX 3049 - R

<400> SEQUENCE: 6

gcagaccctg gaaagtaata aacat 25

<210> SEQ ID NO 7

-continued

---

```

<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: Oligonucleotide PtTX 3105 - F

<400> SEQUENCE: 7

tgtcggtgga gttggcagta gact                24

<210> SEQ ID NO 8
<211> LENGTH: 15
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: Oligonucleotide PtTX 3105 - R

<400> SEQUENCE: 8

gcccagcgtt tctctg                15

<210> SEQ ID NO 9
<211> LENGTH: 38
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: Oligonucleotide PtTX 3116 - F

<400> SEQUENCE: 9

cacgacgttg taaaacgacc tcccaaagcc taaagaat    38

<210> SEQ ID NO 10
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: Oligonucleotide PtTX 3116 - R

<400> SEQUENCE: 10

catacaaggc cttatcttac agaa                24

```

---

We claim:

1. A new and distinct variety of loblolly pine tree named 'CF Q7766' substantially as described and illustrated.

\* \* \* \* \*



FIG. 1




FIG. 2