



US 20200071412A1

(19) **United States**(12) **Patent Application Publication**
Ravetch et al.(10) **Pub. No.: US 2020/0071412 A1**(43) **Pub. Date: Mar. 5, 2020**(54) **ANTIBODIES TO CD40 WITH ENHANCED
AGONIST ACTIVITY**(71) Applicants: **The Rockefeller University**, New York,
NY (US); **BRISTOL-MYERS
SQUIBB COMPANY**, Princeton, NJ
(US)(72) Inventors: **Jeffrey V. Ravetch**, New York, NY
(US); **Rony Dahan**, New York, NY
(US); **Bryan C. Barnhart**, San
Francisco, CA (US); **Brigitte Devaux**,
Palo Alto, CA (US); **Aaron P.
Yamniuk**, Lawrenceville, NJ (US);
Shannon L. Okada, Seattle, WA (US);
Brenda L. Stevens, Seattle, WA (US)(73) Assignees: **The Rockefeller University**, New York,
NY (US); **BRISTOL-MYERS
SQUIBB COMPANY**, Princeton, NJ
(US)(21) Appl. No.: **16/687,107**(22) Filed: **Nov. 18, 2019****Related U.S. Application Data**(62) Division of application No. 15/195,098, filed on Jun.
28, 2016, now Pat. No. 10,479,838.(60) Provisional application No. 62/303,838, filed on Mar.
4, 2016, provisional application No. 62/252,615, filed
on Nov. 9, 2015, provisional application No. 62/186,
076, filed on Jun. 29, 2015.**Publication Classification**(51) **Int. Cl.**
C07K 16/28 (2006.01)(52) **U.S. Cl.**
CPC **C07K 16/2878** (2013.01); **C07K 2317/92**
(2013.01); **C07K 2317/75** (2013.01); **A61K**
2039/505 (2013.01); **C07K 2317/34** (2013.01);
C07K 2317/24 (2013.01); **C07K 2317/21**
(2013.01); **C07K 2317/52** (2013.01)(57) **ABSTRACT**

Provided herein are agonistic antibodies, or antigen binding portions thereof, that bind to human CD40. Such antibodies optionally comprise Fc regions with enhanced specificity for FcγRIIb. The invention also provides methods of treatment of cancer or chronic infection by administering the antibodies of the invention to a subject in need thereof.

Specification includes a Sequence Listing.

Human IgG1f Constant Domain (SEQ ID NO: 65)

Renumbered 118 – 446

ASTKGPSVFP	LAPSSKSTSG	GTAALGCLVK	DYFPEPTVS	WNSGALTSGV	HTFPAVLQSS	177
GLYSLSSVVT	VPSSSLGTQT	YICNVNHKPS	NTKVDKRVET	KSCDKTHTCP	PCPAP ELLGG	237
				D	D	
PSVFLFPPKP	KDTLMISRT	P	ED	EG		297
STYRVVSVLT	VLHQDWLNGK	EYCKCKVSNKA	LP	API	ETIS KAKGQPREPQ	357
			F	R		
MTKNQVSLTC	LVKGFYPSDI	AVEWESNGQP	ENNYKTTPPV	LDSDGSFFLY	SKLTVDKSRW	417
QQGNVFSCSV	MHEALHNHYT	QKSLSLSPG				446

FIG. 1




FIG. 2




FIG. 3A




FIG. 3B




FIG. 4




FIG. 5




FIG. 6A




FIG. 6B




FIG. 6C




FIG. 6D

ANTIBODIES TO CD40 WITH ENHANCED AGONIST ACTIVITY

CROSS REFERENCE TO RELATED APPLICATIONS

[0001] This application is a Divisional of U.S. patent application Ser. No. 15/195,098, filed Jun. 28, 2016, which claims priority to U.S. Provisional Application Nos. 62/303,838, filed Mar. 4, 2016, 62/252,615, filed Nov. 9, 2015 and 62/186,076, filed Jun. 29, 2015, the disclosures of which are all incorporated herein by reference.

BACKGROUND

[0002] Recent research has revealed that human cancers and chronic infections may be treated with agents that modulate the patient's immune response to malignant or infected cells. See, e.g., Reck & Paz-Ares (2015) *Semin. Oncol.* 42:402. Agonistic anti-CD40 antibodies, such as CP-870893 and dacetuzumab (SGN-40) have been tried for treating cancer based on the belief that they may enhance such an immune response. See, e.g., Kirkwood et al. (2012) *CA Cancer J. Clin.* 62:309; Vanderheide & Glennie (2013) *Clin. Cancer Res.* 19:1035. Recent experiments in mice have revealed that anti-CD40 antibodies with enhanced specificity for the inhibitory Fc receptor FcγRIIb have increased anti-tumor efficacy. See, e.g., WO 2012/087928; Li & Ravetch (2011) *Science* 333:1030; Li & Ravetch (2012) *Proc. Nat'l Acad. Sci. (USA)* 109:10966; Wilson et al. (2011) *Cancer Cell* 19:101; White et al. (2011) *J. Immunol.* 187:1754.

[0003] The need exists for improved agonistic anti-human CD40 antibodies for treatment of cancer and chronic infections in human subjects. Such antibodies will preferably have enhanced specificity for the inhibitory Fc receptor FcγRIIb as compared to activating Fc receptors, and will exhibit enhanced anti-tumor and/or anti-infective activity.

SUMMARY OF THE INVENTION

[0004] Provided herein are isolated humanized murine monoclonal antibodies that specifically bind to human CD40 (the mature sequence of SEQ ID NO: 1), optionally having modified Fc regions that enhance specificity for binding to FcγRIIb receptor.

[0005] In certain embodiments, the invention relates to anti-huCD40 antibodies or antigen binding fragments thereof that compete for binding with, cross-block, or bind to the same epitope as, one or more of antibodies 12D6 (SEQ ID NOs: 3 and 4), 5F11 (SEQ ID NOs: 23 and 24), 8E8 (SEQ ID NOs: 40 and 41), 5G7 (SEQ ID NOs: 52 and 53), and 19G3 (SEQ ID NOs: 58 and 59), including human or humanized antibodies.

[0006] In certain embodiments, the anti-human CD40 antibodies of the present invention, or antigen binding fragments thereof, bind at an epitope comprising or consisting of one or more sequences selected from the group consisting of WGCLLTAVHPEPTACRE (residues 11-28 of SEQ ID NO: 1) (antibody 12D6), EPPTACREKQYLINS (residues 21-35 of SEQ ID NO: 1) (antibodies 12D6, 5G7 and 19G3), and ECLPCGESE (residues 58-66 of SEQ ID NO: 1) (antibody 5F11).

[0007] In some embodiments the antibody of the present invention comprises a heavy chain and a light chain, wherein the heavy chain comprises CDRH1, CDRH2 and CDRH3

sequences and the light chain comprises CDRL1, CDRL2 and CDRL3 sequences derived at least in part from the same mouse germline V region gene segments and J region gene segments as anti-huCD40 antibody 12D6, 5F11, 8E8, 5G7 or 19G3, as disclosed at Table 3. Specifically, the antibody may comprise CDR sequences derived from the same murine germlines as antibody 12D6 (heavy chain CDR sequences derived at least in part from murine V region germline VH1-39_01 and J region germline IGHJ4 and light chain CDR sequences derived at least in part from murine V region germline VK1-110_01 and J region germline IGKJ1), antibody 5F11 (heavy chain CDR sequences derived at least in part from murine V region germline VH1-4_02 and J region germline IGHJ3 and light chain CDR sequences derived at least in part from murine V region germline VK3-5_01 and J region germline IGKJ5), antibody 8E8 (heavy chain CDR sequences derived at least in part from murine V region germline VH1-80_01 and J region germline IGHJ2 and light chain CDR sequences derived at least in part from murine V region germline VK1-110_01 and J region germline IGKJ2), antibody 5G7 (heavy chain CDR sequences derived at least in part from murine V region germline VH1-18_01 and J region germline IGHJ4 and light chain CDR sequences derived at least in part from murine V region germline VK10-96_01 and J region germline IGKJ2), or antibody 19G3 (heavy chain CDR sequences derived at least in part from murine V region germline VH5-9-4_01 and J region germline IGHJ3 and light chain CDR sequences derived at least in part from murine V region germline VK1-117_01 and J region germline IGKJ2).

[0008] In various embodiments the antibody of the present invention comprises a heavy chain and a light chain, wherein the heavy chain comprises CDRH1, CDRH2 and CDRH3 sequences and the light chain comprises CDRL1, CDRL2 and CDRL3 sequences selected from the group consisting of: the CDRs of antibody 12D6-03 wherein CDRH1, CDRH2 and CDRH3 comprise residues 31-35, 50-66 and 99-108, respectively, of SEQ ID NO:5 and CDRL1, CDRL2 and CDRL3 comprise residues 24-39, 55-61 and 94-102, respectively, of SEQ ID NO:6; the CDRs of antibody 12D6-22 wherein CDRH1, CDRH2 and CDRH3 comprise residues 31-35, 50-66 and 99-108, respectively, of SEQ ID NO:7 and CDRL1, CDRL2 and CDRL3 comprise residues 24-39, 55-61 and 94-102, respectively, of SEQ ID NO:9; the CDRs of antibody 12D6-23 wherein CDRH1, CDRH2 and CDRH3 comprise residues 31-35, 50-66 and 99-108, respectively, of SEQ ID NO:10 and CDRL1, CDRL2 and CDRL3 comprise residues 24-39, 55-61 and 94-102, respectively, of SEQ ID NO:11; the CDRs of antibody 12D6-24 wherein CDRH1, CDRH2 and CDRH3 comprise residues 31-35, 50-66 and 99-108, respectively, of SEQ ID NO:12 and CDRL1, CDRL2 and CDRL3 comprise residues 24-39, 55-61 and 94-102, respectively, of SEQ ID NO:9; the CDRs of antibody 5F11-17 wherein CDRH1, CDRH2 and CDRH3 comprise residues 31-35, 50-66 and 99-106, respectively, of SEQ ID NO:25 and CDRL1, CDRL2 and CDRL3 comprise residues 24-38, 54-60 and 93-101, respectively, of SEQ ID NO:26; the CDRs of antibody 5F11-23 wherein CDRH1, CDRH2 and CDRH3 comprise residues 31-35, 50-66 and 99-106, respectively, of SEQ ID NO:27 and CDRL1, CDRL2 and CDRL3 comprise residues 24-38, 54-60 and 93-101, respectively, of SEQ ID NO:28; the CDRs of antibody 5F11-45 wherein CDRH1, CDRH2 and CDRH3 comprise residues 31-35, 50-66 and 99-106, respectively, of

SEQ ID NO:29 and CDRL1, CDRL2 and CDRL3 comprise residues 24-38, 54-60 and 93-101, respectively, of SEQ ID NO:30; the CDRs of antibody 8E8-56 wherein CDRH1, CDRH2 and CDRH3 comprise residues 31-35, 50-66 and 99-111, respectively, of SEQ ID NO:42 and CDRL1, CDRL2 and CDRL3 comprise residues 24-39, 55-61 and 94-102, respectively, of SEQ ID NO:43; the CDRs of antibody 8E8-62 wherein CDRH1, CDRH2 and CDRH3 comprise residues 31-35, 50-66 and 99-111, respectively, of SEQ ID NO:44 and CDRL1, CDRL2 and CDRL3 comprise residues 24-39, 55-61 and 94-102, respectively, of SEQ ID NO:45; the CDRs of antibody 8E8-67 wherein CDRH1, CDRH2 and CDRH3 comprise residues 31-35, 50-66 and 99-111, respectively, of SEQ ID NO:46 and CDRL1, CDRL2 and CDRL3 comprise residues 24-39, 55-61 and 94-102, respectively, of SEQ ID NO:47; the CDRs of antibody 8E8-70 wherein CDRH1, CDRH2 and CDRH3 comprise residues 31-35, 50-66 and 99-111, respectively, of SEQ ID NO:48 and CDRL1, CDRL2 and CDRL3 comprise residues 24-39, 55-61 and 94-102, respectively, of SEQ ID NO:49; the CDRs of antibody 8E8-71 wherein CDRH1, CDRH2 and CDRH3 comprise residues 31-35, 50-66 and 99-111, respectively, of SEQ ID NO:50 and CDRL1, CDRL2 and CDRL3 comprise residues 24-39, 55-61 and 94-102, respectively, of SEQ ID NO:51; the CDRs of antibody 5G7-22 wherein CDRH1, CDRH2 and CDRH3 comprise residues 31-35, 50-66 and 99-102, respectively, of SEQ ID NO:54 and CDRL1, CDRL2 and CDRL3 comprise residues 24-34, 50-56 and 89-97, respectively, of SEQ ID NO:55; the CDRs of antibody 5G7-25 wherein CDRH1, CDRH2 and CDRH3 comprise residues 31-35, 50-66 and 99-102, respectively, of SEQ ID NO:56 and CDRL1, CDRL2 and CDRL3 comprise residues 24-34, 50-56 and 89-97, respectively, of SEQ ID NO:57; the CDRs of antibody 19G3-11 wherein CDRH1, CDRH2 and CDRH3 comprise residues 31-35, 50-66 and 99-101, respectively, of SEQ ID NO:60 and CDRL1, CDRL2 and CDRL3 comprise residues 24-39, 55-61 and 94-102, respectively, of SEQ ID NO:62; and the CDRs of antibody 19G3-22 wherein CDRH1, CDRH2 and CDRH3 comprise residues 31-35, 50-66 and 99-101, respectively, of SEQ ID NO:63 and CDRL1, CDRL2 and CDRL3 comprise residues 24-39, 55-61 and 94-102, respectively, of SEQ ID NO:64.

[0009] In various embodiments the antibody of the present invention comprises a heavy chain comprising a variable domain selected from the group consisting of 12D6 (residues 1-119 of SEQ ID NO: 3), 5F11 (residues 1-117 of SEQ ID NO: 23), 8E8 (residues 1-122 of SEQ ID NO: 40), 5G7 (residues 1-113 of SEQ ID NO: 52) and 19G3 (residues 1-112 of SEQ ID NO: 58) with constant regions comprising FcγRIIb-specific Fc region selected from the group consisting of IgG1f (SEQ ID NO: 65), SE (SEQ ID NO: 66), SELF (SEQ ID NO: 67), P238D (SEQ ID NO: 68), V4 (SEQ ID NO: 69), V4 D270E (SEQ ID NO: 70), V7 (SEQ ID NO: 71), V8 (SEQ ID NO: 72), V9 (SEQ ID NO: 73), V9 D270E (SEQ ID NO: 74), V11 (SEQ ID NO: 75), and V12 (SEQ ID NO: 76).

[0010] In some embodiments the antibody comprises specific heavy chain variable domains and light chain variable domains selected from the group consisting of 12D6-03 (residues 1-119 and 1-112 of SEQ ID NO:5 and SEQ ID NO:6, respectively), 12D6-22 (residues 1-119 and 1-112 of SEQ ID NO:7 and SEQ ID NO:9, respectively), 12D6-23 (residues 1-119 and 1-112 of SEQ ID NO:10 and SEQ ID

NO:11, respectively), 12D6-24 (residues 1-119 and 1-112 of SEQ ID NO:12 and SEQ ID NO:9, respectively), 5F11-17 (residues 1-117 and 1-111 of SEQ ID NO:25 and SEQ ID NO:26, respectively), 5F11-23 (residues 1-117 and 1-111 of SEQ ID NO:27 and SEQ ID NO:28, respectively), 5F11-45 (residues 1-117 and 1-111 of SEQ ID NO:29 and SEQ ID NO:30), 8E8-56 (residues 1-122 and 1-112 of SEQ ID NO:42 and SEQ ID NO:43, respectively), 8E8-62 (residues 1-122 and 1-112 of SEQ ID NO:44 and SEQ ID NO:45, respectively), 8E8-67 (residues 1-122 and 1-112 of SEQ ID NO:46 and SEQ ID NO:47, respectively), 8E8-70 (residues 1-122 and 1-112 of SEQ ID NO:48 and SEQ ID NO:49), 8E8-71 (residues 1-122 and 1-112 of SEQ ID NO:50 and SEQ ID NO:51, respectively), 5G7-22 (residues 1-113 and 1-107 of SEQ ID NO:54 and SEQ ID NO:55, respectively), 5G7-25 (residues 1-113 and 1-107 of SEQ ID NO:56 and SEQ ID NO:57, respectively), 19G3-11 (residues 1-112 and 1-112 of SEQ ID NO:60 and SEQ ID NO:62, respectively), and 9G3-22 (residues 1-112 and 1-112 of SEQ ID NO:63 and SEQ ID NO:64, respectively). Any of these antibodies may further comprise a heavy chain constant region comprising an FcγRIIb-specific Fc region, said heavy chain constant region selected from the group consisting of IgG1f (SEQ ID NO: 65), SE (SEQ ID NO: 66), SELF (SEQ ID NO: 67), P238D (SEQ ID NO: 68), V4 (SEQ ID NO: 69), V4 D270E (SEQ ID NO: 70), V7 (SEQ ID NO: 71), V8 (SEQ ID NO: 72), V9 (SEQ ID NO: 73), V9 D270E (SEQ ID NO: 74), V11 (SEQ ID NO: 75), and V12 (SEQ ID NO: 76). Any of these antibodies may further comprise the light chain kappa constant region of SEQ ID NO: 77.

[0011] In specific embodiments, the antibody of the present invention comprises a humanized 12D6-24 antibody comprising a light chain of SEQ ID NO: 9 and a heavy chain selected from the group consisting of any of SEQ ID NOs: 13-22, or a humanized 5F11-45 antibody comprising a light chain of SEQ ID NO: 30 and a heavy chain selected from the group consisting of any of SEQ ID NOs: 31-39. Specific antibodies include 12D6-24 SE (SEQ ID NOs: 9 and 14), 12D6-24 SELF (SEQ ID NOs: 9 and 15), 12D6-24 P238D (SEQ ID NOs: 9 and 13), 12D6-24 V4 (SEQ ID NOs: 9 and 16), 12D6-24 V4 D270E (SEQ ID NOs: 9 and 17), 12D6-24 V8 (SEQ ID NOs: 9 and 18), 12D6-24 V9 (SEQ ID NOs: 9 and 19), 12D6-24 V9 D270E (SEQ ID NOs: 9 and 20), 12D6-24 V11 (SEQ ID NOs: 9 and 21), 12D6-24 V12 (SEQ ID NOs: 9 and 22), 5F11-45 SE (SEQ ID NOs: 30 and 31), 5F11-45 SELF (SEQ ID NOs: 30 and 32), 5F11-45 V4 (SEQ ID NOs: 30 and 33), 5F11-45 V4 D270E (SEQ ID NOs: 30 and 34), 5F11-45 V8 (SEQ ID NOs: 30 and 35), 5F11-45 V9 (SEQ ID NOs: 30 and 36), 5F11-45 V9 D270E (SEQ ID NOs: 30 and 37), 5F11-45 V11 (SEQ ID NOs: 30 and 38), and 5F11-45 V12 (SEQ ID NOs: 30 and 39), where sequences are provided for light and heavy chains, respectively.

[0012] In further embodiments the anti-huCD40 antibodies of the present comprise heavy and light chains sharing at least 80%, 85%, 90% and 95% sequence identity with the sequences of the heavy and light chains of 12D6-24 SE (SEQ ID NOs: 9 and 14), 12D6-24 SELF (SEQ ID NOs: 9 and 15), 12D6-24 P238D (SEQ ID NOs: 9 and 13), 12D6-24 V4 (SEQ ID NOs: 9 and 16), 12D6-24 V4 D270E (SEQ ID NOs: 9 and 17), 12D6-24 V8 (SEQ ID NOs: 9 and 18), 12D6-24 V9 (SEQ ID NOs: 9 and 19), 12D6-24 V9 D270E (SEQ ID NOs: 9 and 20), 12D6-24 V11 (SEQ ID NOs: 9 and 21), 12D6-24 V12 (SEQ ID NOs: 9 and 22), 5F11-45 SE

(SEQ ID NOs: 30 and 31), 5F11-45 SELF (SEQ ID NOs: 30 and 32), 5F11-45 V4 (SEQ ID NOs: 30 and 33), 5F11-45 V4 D270E (SEQ ID NOs: 30 and 34), 5F11-45 V8 (SEQ ID NOs: 30 and 35), 5F11-45 V9 (SEQ ID NOs: 30 and 36), 5F11-45 V9 D270E (SEQ ID NOs: 30 and 37), 5F11-45 V11 (SEQ ID NOs: 30 and 38), or 5F11-45 V12 (SEQ ID NOs: 30 and 39).

[0013] In yet further embodiments the anti-huCD40 antibodies of the present comprise heavy and light chains consisting essentially of the sequences of the heavy and light chains of 12D6-24 SE (SEQ ID NOs: 9 and 14), 12D6-24 SELF (SEQ ID NOs: 9 and 15), 12D6-24 P238D (SEQ ID NOs: 9 and 13), 12D6-24 V4 (SEQ ID NOs: 9 and 16), 12D6-24 V4 D270E (SEQ ID NOs: 9 and 17), 12D6-24 V8 (SEQ ID NOs: 9 and 18), 12D6-24 V9 (SEQ ID NOs: 9 and 19), 12D6-24 V9 D270E (SEQ ID NOs: 9 and 20), 12D6-24 V11 (SEQ ID NOs: 9 and 21), 12D6-24 V12 (SEQ ID NOs: 9 and 22), 5F11-45 SE (SEQ ID NOs: 30 and 31), 5F11-45 SELF (SEQ ID NOs: 30 and 32), 5F11-45 V4 (SEQ ID NOs: 30 and 33), 5F11-45 V4 D270E (SEQ ID NOs: 30 and 34), 5F11-45 V8 (SEQ ID NOs: 30 and 35), 5F11-45 V9 (SEQ ID NOs: 30 and 36), 5F11-45 V9 D270E (SEQ ID NOs: 30 and 37), 5F11-45 V11 (SEQ ID NOs: 30 and 38), or 5F11-45 V12 (SEQ ID NOs: 30 and 39).

[0014] In some embodiments, anti-huCD40 antibodies of the present invention that comprise V4 or V9 Fc sequence variants further comprise the D270E sequence variant. Such antibodies include a humanized 12D6-24 V4 D270E (SEQ ID NOs: 9 and 17), 12D6-24 V9 D270E (SEQ ID NOs: 9 and 20), 5F11-45 V4 D270E (SEQ ID NOs: 30 and 34), and 5F11-45 V9 D270E (SEQ ID NOs: 30 and 37), where sequences are provided for light and heavy chains, respectively. In alternative embodiments, anti-human CD40 antibodies of the present invention include antibodies comprising heavy and light chains consisting essentially of the sequences of these heavy and light chains, or comprise heavy and light chains sharing at least 80%, 85%, 90% and 95% sequence identity with these sequences. In some embodiments, the anti-huCD40 antibodies of the present invention comprise modified Fc regions with greater specificity for binding to FcγRIIb as opposed to binding to activating receptors than antibodies with naturally occurring Fc regions. In certain embodiments the A/I ratio for the anti-huCD40 antibody of the present invention is less than 5, and in preferred embodiments, less than 1.

[0015] In some embodiments the anti-huCD40 antibody of the present invention comprises one or more heavy chains and one or more light chains, such as two heavy chains and two light chains.

[0016] The present invention further provides nucleic acids encoding the heavy and/or light chain variable regions, of the anti-CD40 antibodies of the present invention, or antigen binding fragments thereof, expression vectors comprising the nucleic acid molecules, cells transformed with the expression vectors, and methods of producing the antibodies by expressing the antibodies from cells transformed with the expression vectors and recovering the antibody.

[0017] The present invention also provides pharmaceutical compositions comprising anti-huCD40 antibodies of the present invention, or antigen binding fragments thereof, and a carrier.

[0018] The present invention provides a method of enhancing an immune response in a subject comprising administering an effective amount of an anti-huCD40 anti-

body of the present invention, or antigen binding fragment thereof, to the subject such that an immune response in the subject is enhanced. In certain embodiments, the subject has a tumor and an immune response against the tumor is enhanced. In another embodiment, the subject has a viral infection, e.g. a chronic viral infection, and an anti-viral immune response is enhanced.

[0019] The present invention also provides a method of inhibiting the growth of tumors in a subject comprising administering to the subject an anti-huCD40 antibody of the present invention, or antigen binding fragment thereof, such that growth of the tumor is inhibited.

[0020] The present invention further provides a method of treating cancer, e.g., by immunotherapy, comprising administering to a subject in need thereof a therapeutically effective amount of an anti-huCD40 antibody of the present invention, or antigen binding fragment thereof, e.g. as a pharmaceutical composition, thereby treating the cancer. In certain embodiments, the cancer is bladder cancer, breast cancer, uterine/cervical cancer, ovarian cancer, prostate cancer, testicular cancer, esophageal cancer, gastrointestinal cancer, pancreatic cancer, colorectal cancer, colon cancer, kidney cancer, head and neck cancer, lung cancer, stomach cancer, germ cell cancer, bone cancer, liver cancer, thyroid cancer, skin cancer, neoplasm of the central nervous system, lymphoma, leukemia, myeloma, sarcoma, and virus-related cancer. In certain embodiments, the cancer is a metastatic cancer, refractory cancer, or recurrent cancer.

[0021] In certain embodiments, the methods of modulating immune function and methods of treatment described herein comprise administering an anti-huCD40 antibody of the present invention in combination with, or as a bispecific reagent with, one or more additional therapeutics, for example, an anti-PD1 antibody, an anti-PD-L1 antibody, an anti-LAG3 antibody, an anti-GITR antibody, an anti-OX40 antibody, an anti-CD73 antibody, an anti-TIGIT antibody, an anti-CD137 antibody, an anti-CD27 antibody, an anti-CSF-1R antibody, an anti-CTLA-4 antibody, a TLR agonist, or a small molecule antagonist of IDO or TGFβ. In specific embodiments, anti-huCD40 therapy is combined with anti-PD1 and/or anti-PD-L1 therapy, e.g. treatment with an antibody or antigen binding fragment thereof that binds to human PD1 or an antibody or antigen binding fragment thereof that binds to human PD-L1.

BRIEF DESCRIPTION OF THE DRAWINGS

[0022] FIG. 1 shows the sequence of human IgG1f constant domain (SEQ ID NO: 65) renumbered 118-446 to better illustrate the Fc sequence variants disclosed herein (Table 4). Residues subject to variation are in bold, and the altered amino acid is provided in bold below the residue. The D270E substitution is underlined. A C-terminal lysine (K) residue has been removed in FIG. 1 and SEQ ID NO: 65, as well as all other heavy chain and heavy chain constant domain sequences disclosed in the Sequence Listing. However, in other embodiments, especially nucleic acid constructs encoding the heavy chains and heavy chain constant domains of the anti-huCD40 antibodies of the present invention, these sequences include an additional lysine residue at the C-terminus of the protein or nucleotides encoding the extra lysine at the 3' end of the nucleic acid.

[0023] FIG. 2 is a Venn diagram illustrating the epitope groups ("bins") on human CD40 bound by the antibodies of the present invention, as well as blocking of CD40L binding.

Antibodies with overlapping ovals or circles compete for binding to human CD40, and antibodies falling within the rectangle block CD40L binding to human CD40.

[0024] FIGS. 3A and 3B show activation of dendritic cells, as measured by IL-6 secretion, by agonist anti-CD40 antibodies as a function of Fc sequence. See Example 7. A series of antibodies was constructed comprising an mAb 12D6-24 variable domain and various human IgG1f constant regions, including IgG1f, SE, SELF, P238D, V4, V8, V9 and V12 variants. FIG. 3A presents data obtained using cells from one donor, and FIG. 3B presents data obtained using cells from a different donor.

[0025] FIG. 4 shows activation of cells, as measured by cell surface CD54, by agonist anti-CD40 antibodies as a function of variable domain sequence. See Example 7. A series of antibodies was constructed comprising a human IgG1f-V12 constant region and variable domains from parental (murine) anti-CD40 mAbs 12D6, 5G7, 8E8, 19G3 and 5F11. Results are plotted as median fluorescence intensity (MFI) as a function of antibody concentration.

[0026] FIG. 5 shows percent FcγR binding by various antibodies of the present invention, including antibodies having D270 substitutions. See Example 8. Antibody names including “-sup” represent supernatants from antibody producing cells, whereas others are purified antibodies. Data are presented as percentages of a maximum receptor binding value for each combination of antibody and receptor, as measured in a FORTIBIO Octet system. See, e.g., Example 3. Each cluster of three bars represents, from left to right, binding to hCD32a/FcγRIIa-H131 (10 μM) (hatched bars), hCD32b/FcγRIIa-R131 (10 μM) (black bars), and hCD32b/FcγRIIb (1 μM) (white bars).

[0027] FIGS. 6A and 6B show the effects of selected anti-CD40 antibodies of the present invention on T cell activation and change in platelet count, respectively, in transgenic mice expressing human CD40 and human Fcγ receptors. See Example 9. FIG. 6A shows the percent of Tet-OVA reactive CD8⁺ T cells in animals treated with selected anti-CD40 antibodies of the present invention, as indicated. FIG. 6B shows the platelet count as a percentage of pre-treatment platelet count at 24 hours post-injection with antibody. Comparison of the figures shows that the level of activation correlates with reduction in platelet count, with antibody 12D6-V11 exhibiting the highest activation but also the greatest reduction in platelet count. See Example 9.

[0028] FIG. 6C shows the antitumor response of humanized CD40/FcγR mice that were inoculated with MC38 tumor cells and treated with Fc variants of anti-CD40 12D6-24 and 5F11-45 clones. Results presented as means±SEM. n=7 (12D6-24) or 6 (5F11-45). See Example 9.

[0029] FIG. 6D shows tumor free mice from the 12D6-24 group in the experiment described in C were re-challenged with MC38 cells subcutaneously and followed for tumor growth. Control group consists of naïve mice. Results presented as means±SEM. n=4. See Example 9.

DETAILED DESCRIPTION

[0030] The present invention provides isolated antibodies, particularly monoclonal antibodies, e.g., humanized or human monoclonal antibodies, that specifically bind to human CD40 (“huCD40”) and have agonist activity. Sequences are provided for various humanized murine anti-

huCD40 monoclonal antibodies. In certain embodiments, the antibodies described herein are derived from particular murine heavy and light chain germline sequences and/or comprise particular structural features such as CDR regions comprising particular amino acid sequences. In other embodiments antibodies compete for CD40 binding with, or bind to the same epitope as, the anti-CD40 antibodies for which sequences are provided herein. In some embodiments the sequence of the heavy chain Fc region is modified to specifically enhance binding to FcγRIIb.

[0031] Further provided herein are methods of making such antibodies, immunoconjugates and bispecific molecules comprising such antibodies or antigen-binding fragments thereof, and pharmaceutical compositions formulated to contain the antibodies or fragments. Also provided herein are methods of using the antibodies for immune response enhancement, alone or in combination with other immunostimulatory agents (e.g., antibodies) and/or cancer or anti-infective therapies. Accordingly, the anti-huCD40 antibodies described herein may be used in a treatment in a wide variety of therapeutic applications, including, for example, inhibiting tumor growth and treating chronic viral infections.

Definitions

[0032] In order that the present description may be more readily understood, certain terms are first defined. Additional definitions are set forth throughout the detailed description.

[0033] CD40 refers to “TNF receptor superfamily member 5” (TNFRSF5). Unless otherwise indicated, or clear from the context, references to CD40 herein refer to human CD40 (“huCD40”), and anti-CD40 antibodies refer to anti-human CD40 antibodies. Human CD40 is further described at GENE ID NO: 958 and MIM (Mendelian Inheritance in Man): 109535. The sequence of human CD40 (NP_001241.1), including 20 amino acid signal sequence, is provided at SEQ ID NO: 1.

[0034] CD40 interacts with CD40 ligand (CD40L), which is also referred to as TNFSF5, gp39 and CD154. Unless otherwise indicated, or clear from the context, references to CD40L herein refer to human CD40L (“huCD40L”). Human CD40L is further described at GENE ID NO: 959 and MIM: 300386. The sequence of human CD40L (NP_000065.1) is provided at SEQ ID NO: 2.

[0035] Unless otherwise indicated or clear from the context, the term “antibody” as used to herein may include whole antibodies and any antigen-binding fragments (i.e., “antigen-binding portions”) or single chains thereof. An “antibody” refers, in one embodiment, to a glycoprotein comprising at least two heavy (H) chains and two light (L) chains inter-connected by disulfide bonds, or an antigen binding fragment thereof. Each heavy chain is comprised of a heavy chain variable region (abbreviated herein as V_H) and a heavy chain constant region. In certain naturally occurring IgG, IgD and IgA antibodies, the heavy chain constant region is comprised of three domains, CH1, CH2 and CH3. In certain naturally occurring antibodies, each light chain is comprised of a light chain variable region (abbreviated herein as V_L) and a light chain constant region. The light chain constant region is comprised of one domain, CL. The V_H and V_L regions can be further subdivided into regions of hypervariability, termed complementarity determining regions (CDR), interspersed with regions that are more conserved, termed framework regions (FR). Each V_H and V_L

is composed of three CDRs and four framework regions (FRs), arranged from amino-terminus to carboxy-terminus in the following order: FR1, CDR1, FR2, CDR2, FR3, CDR3, FR4. The variable regions of the heavy and light chains contain a binding domain that interacts with an antigen. The constant regions of the antibodies may mediate the binding of the immunoglobulin to host tissues or factors, including various cells of the immune system (e.g., effector cells) and the first component (C1q) of the classical complement system.

[0036] Antibodies typically bind specifically to their cognate antigen with high affinity, reflected by a dissociation constant (K_D) of 10^{-7} to 10^{-11} M or less. Any K_D greater than about 10^{-6} M is generally considered to indicate non-specific binding. As used herein, an antibody that “binds specifically” to an antigen refers to an antibody that binds to the antigen and substantially identical antigens with high affinity, which means having a K_D of 10^{-7} M or less, preferably 10^{-8} M or less, even more preferably 5×10^{-9} M or less, and most preferably between 10^{-8} M and 10^{-10} M or less, but does not bind with high affinity to unrelated antigens. An antigen is “substantially identical” to a given antigen if it exhibits a high degree of sequence identity to the given antigen, for example, if it exhibits at least 80%, at least 90%, preferably at least 95%, more preferably at least 97%, or even more preferably at least 99% sequence identity to the sequence of the given antigen. By way of example, an antibody that binds specifically to human CD40 might also cross-react with CD40 from certain non-human primate species (e.g., cynomolgus monkey), but might not cross-react with CD40 from other species, or with an antigen other than CD40.

[0037] Unless otherwise indicated, an immunoglobulin may be from any of the commonly known isotypes, including but not limited to IgA, secretory IgA, IgG and IgM. The IgG isotype is divided in subclasses in certain species: IgG1, IgG2, IgG3 and IgG4 in humans, and IgG1, IgG2a, IgG2b and IgG3 in mice. Immunoglobulins, e.g., human IgG1, exist in several allotypes, which differ from each other in at most a few amino acids. Unless otherwise indicated, antibodies of the present invention comprise the IgG1f constant domain (SEQ ID NO: 65). Unless otherwise indicated, “antibody” may include, by way of example, monoclonal and polyclonal antibodies; chimeric and humanized antibodies; human and non-human antibodies; wholly synthetic antibodies; and single chain antibodies.

[0038] The term “antigen-binding portion” or “antigen binding fragment” of an antibody, as used herein, refers to one or more fragments of an antibody that retain the ability to specifically bind to an antigen (e.g., human CD40). Examples of binding fragments encompassed within the term “antigen-binding portion/fragment” of an antibody include (i) a Fab fragment—a monovalent fragment consisting of the VL, VH, CL and CH1 domains; (ii) a F(ab')₂ fragment—a bivalent fragment comprising two Fab fragments linked by a disulfide bridge at the hinge region; (iii) a Fd fragment consisting of the VH and CH1 domains; (iv) a Fv fragment consisting of the VL and VH domains of a single arm of an antibody, and (v) a dAb fragment (Ward et al., (1989) *Nature* 341:544-546) consisting of a VH domain. An isolated complementarity determining region (CDR), or a combination of two or more isolated CDRs joined by a synthetic linker, may comprise and antigen binding domain of an antibody if able to bind antigen.

[0039] Single chain antibody constructs are also included in the invention. Although the two domains of the Fv fragment, VL and VH, are coded for by separate genes, they can be joined, using recombinant methods, by a synthetic linker that enables them to be made as a single protein chain in which the VL and VH regions pair to form monovalent molecules known as single chain Fv (scFv); see e.g., Bird et al. (1988) *Science* 242:423-426; and Huston et al. (1988) *Proc. Natl. Acad. Sci. (USA)* 85:5879-5883). Such single chain antibodies are also intended to be encompassed within the term “antigen-binding portion/fragment” of an antibody. These and other potential constructs are described at Chan & Carter (2010) *Nat. Rev. Immunol.* 10:301. These antibody fragments are obtained using conventional techniques known to those with skill in the art, and the fragments are screened for utility in the same manner as are intact antibodies. Antigen-binding portions/fragments can be produced by recombinant DNA techniques, or by enzymatic or chemical cleavage of intact immunoglobulins.

[0040] Unless otherwise indicated, the word “fragment” when used with reference to an antibody, such as in a claim, refers to an antigen binding fragment of the antibody, such that “antibody or fragment” has the same meaning as “antibody or antigen binding fragment thereof.”

[0041] A “bispecific” or “bifunctional antibody” is an artificial hybrid antibody having two different heavy/light chain pairs, giving rise to two antigen binding sites with specificity for different antigens. Bispecific antibodies can be produced by a variety of methods including fusion of hybridomas or linking of Fab' fragments. See, e.g., Song-sivilai & Lachmann, (1990) *Clin. Exp. Immunol.* 79:315-321; Kostelny et al., (1992) *J. Immunol.* 148, 1547-1553.

[0042] The term “monoclonal antibody,” as used herein, refers to an antibody that displays a single binding specificity and affinity for a particular epitope or a composition of antibodies in which all antibodies display a single binding specificity and affinity for a particular epitope. Typically such monoclonal antibodies will be derived from a single cell or nucleic acid encoding the antibody, and will be propagated without intentionally introducing any sequence alterations. Accordingly, the term “human monoclonal antibody” refers to a monoclonal antibody that has variable and optional constant regions derived from human germline immunoglobulin sequences. In one embodiment, human monoclonal antibodies are produced by a hybridoma, for example, obtained by fusing a B cell obtained from a transgenic or transchromosomal non-human animal (e.g., a transgenic mouse having a genome comprising a human heavy chain transgene and a light chain transgene), to an immortalized cell.

[0043] The term “recombinant human antibody,” as used herein, includes all human antibodies that are prepared, expressed, created or isolated by recombinant means, such as (a) antibodies isolated from an animal (e.g., a mouse) that is transgenic or transchromosomal for human immunoglobulin genes or a hybridoma prepared therefrom, (b) antibodies isolated from a host cell transformed to express the antibody, e.g., from a transfectoma, (c) antibodies isolated from a recombinant, combinatorial human antibody library, and (d) antibodies prepared, expressed, created or isolated by any other means that involve splicing of human immunoglobulin gene sequences to other DNA sequences. Such recombinant human antibodies comprise variable and constant regions that utilize particular human germline

immunoglobulin sequences are encoded by the germline genes, but include subsequent rearrangements and mutations that occur, for example, during antibody maturation. As known in the art (see, e.g., Lonberg (2005) *Nature Biotech.* 23(9):1117-1125), the variable region contains the antigen binding domain, which is encoded by various genes that rearrange to form an antibody specific for a foreign antigen. In addition to rearrangement, the variable region can be further modified by multiple single amino acid changes (referred to as somatic mutation or hypermutation) to increase the affinity of the antibody to the foreign antigen. The constant region will change in further response to an antigen (i.e., isotype switch). Therefore, the rearranged and somatically mutated nucleic acid sequences that encode the light chain and heavy chain immunoglobulin polypeptides in response to an antigen may not be identical to the original germline sequences, but instead will be substantially identical or similar (i.e., have at least 80% identity).

[0044] A “human” antibody (HuMAb) refers to an antibody having variable regions in which both the framework and CDR regions are derived from human germline immunoglobulin sequences. Furthermore, if the antibody contains a constant region, the constant region also is derived from human germline immunoglobulin sequences. Human antibodies of the present invention may include amino acid residues not encoded by human germline immunoglobulin sequences (e.g., mutations introduced by random or site-specific mutagenesis in vitro or by somatic mutation in vivo). However, the term “human antibody,” as used herein, is not intended to include antibodies in which CDR sequences derived from the germline of another mammalian species, such as a mouse, have been grafted onto human framework sequences. The terms “human” antibodies and “fully human” antibodies are used synonymously.

[0045] A “humanized” antibody refers to an antibody in which some, most or all of the amino acids outside the CDR domains of a non-human antibody, e.g. a mouse antibody, are replaced with corresponding amino acids derived from human immunoglobulins. In one embodiment of a humanized form of an antibody, some, most or all of the amino acids outside the CDR domains have been replaced with amino acids from human immunoglobulins, whereas some, most or all amino acids within one or more CDR regions are unchanged. Small additions, deletions, insertions, substitutions or modifications of amino acids are permissible as long as they do not abrogate the ability of the antibody to bind to a particular antigen. A “humanized” antibody retains an antigenic specificity similar to that of the original antibody.

[0046] A “chimeric antibody” refers to an antibody in which the variable regions are derived from one species and the constant regions are derived from another species, such as an antibody in which the variable regions are derived from a mouse antibody and the constant regions are derived from a human antibody. A “hybrid” antibody refers to an

antibody having heavy and light chains of different types, such as a mouse (parental) heavy chain and a humanized light chain, or vice versa.

[0047] As used herein, “isotype” refers to the antibody class (e.g., IgG1, IgG2, IgG3, IgG4, IgM, IgA1, IgA2, IgD, and IgE antibody) that is encoded by the heavy chain constant region genes.

[0048] “Allotype” refers to naturally occurring variants within a specific isotype group, which variants differ in one or a few amino acids. See, e.g., Jefferis et al. (2009) *mAbs* 1:1.

[0049] The phrases “an antibody recognizing an antigen” and “an antibody specific for an antigen” are used interchangeably herein with the term “an antibody that binds specifically to an antigen.”

[0050] An “isolated antibody,” as used herein, refers to an antibody that is substantially free of other antibodies having different antigenic specificities (e.g., an isolated antibody that specifically binds to CD40 is substantially free of antibodies that specifically bind antigens other than CD40). An isolated antibody that specifically binds to an epitope of CD40 may, however, have cross-reactivity to other CD40 proteins from different species.

[0051] “Effector functions,” deriving from the interaction of an antibody Fc region with certain Fc receptors, include but are not necessarily limited to C1q binding, complement dependent cytotoxicity (CDC), Fc receptor binding, FcγR-mediated effector functions such as ADCC and antibody dependent cell-mediated phagocytosis (ADCP), and down regulation of a cell surface receptor (e.g., the B cell receptor; BCR). Such effector functions generally require the Fc region to be combined with an antigen binding domain (e.g., an antibody variable domain).

[0052] An “Fc receptor” or “FcR” is a receptor that binds to the Fc region of an immunoglobulin. FcRs that bind to an IgG antibody comprise receptors of the FcγR family, including allelic variants and alternatively spliced forms of these receptors. The FcγR family consists of three activating (FcγRI, FcγRIII, and FcγRIV in mice; FcγRIA, FcγRIIA, and FcγRIIA in humans) and one inhibitory (FcγRIIb, or equivalently FcγRIIB) receptor. Various properties of human FcγRs are summarized in Table 1. The majority of innate effector cell types co-express one or more activating FcγR and the inhibitory FcγRIIb, whereas natural killer (NK) cells selectively express one activating Fc receptor (FcγRIII in mice and FcγRIIA in humans) but not the inhibitory FcγRIIb in mice and humans. Human IgG1 binds to most human Fc receptors and is considered equivalent to murine IgG2a with respect to the types of activating Fc receptors that it binds to.

TABLE 1

Properties of Human FcγRs				
Fcγ	Allelic variants	Affinity for human IgG	Isotype preference	Cellular distribution
FcγRI	None described	High ($K_D \sim 10$ nM)	IgG1 = 3 > 4 >> 2	Monocytes, macrophages, activated neutrophils, dendritic cells?

TABLE 1-continued

Properties of Human FcγRs				
Fcγ	Allelic variants	Affinity for human IgG	Isotype preference	Cellular distribution
FcγRIIA	H131	Low to medium	IgG1 > 3 > 2 > 4	Neutrophils, monocytes, macrophages, eosinophils, dendritic cells, platelets
	R131	Low	IgG1 > 3 > 4 > 2	
FcγRIIA	V158	Medium	IgG1 = 3 >> 4 > 2	NK cells, monocytes, macrophages, mast cells, eosinophils, dendritic cells?
	F158	Low	IgG1 = 3 >> 4 > 2	
FcγRIIb	I232	Low	IgG1 = 3 = 4 > 2	B cells, monocytes, macrophages, dendritic cells, mast cells
	T232	Low	IgG1 = 3 = 4 > 2	

[0053] An “Fc region” (fragment crystallizable region) or “Fc domain” or “Fc” refers to the C-terminal region of the heavy chain of an antibody that mediates the binding of the immunoglobulin to host tissues or factors, including binding to Fc receptors located on various cells of the immune system (e.g., effector cells) or to the first component (C1q) of the classical complement system. Thus, an Fc region comprises the constant region of an antibody excluding the first constant region immunoglobulin domain (e.g., CH1 or CL). In IgG, IgA and IgD antibody isotypes, the Fc region comprises C_{H2} and C_{H3} constant domains in each of the antibody’s two heavy chains; IgM and IgE Fc regions comprise three heavy chain constant domains (C_H domains 2-4) in each polypeptide chain. For IgG, the Fc region comprises immunoglobulin domains Cγ2 and Cγ3 and the hinge between Cγ1 and Cγ2. Although the boundaries of the Fc region of an immunoglobulin heavy chain might vary, the human IgG heavy chain Fc region is usually defined to stretch from an amino acid residue at position C226 or P230 (or an amino acid between these two amino acids) to the carboxy-terminus of the heavy chain, wherein the numbering is according to the EU index as in Kabat. Kabat et al. (1991) *Sequences of Proteins of Immunological Interest*, National Institutes of Health, Bethesda, Md.; see also FIGS. 3c-3f of U.S. Pat. App. Pub. No. 2008/0248028. The C_{H2} domain of a human IgG Fc region extends from about amino acid 231 to about amino acid 340, whereas the C_{H3} domain is positioned on C-terminal side of a C_{H2} domain in an Fc region, i.e., it extends from about amino acid 341 to about amino acid 447 of an IgG (including a C-terminal lysine). As used herein, the Fc region may be a native sequence Fc, including any allotypic variant, or a variant Fc (e.g., a non-naturally occurring Fc). Fc may also refer to this region in isolation or in the context of an Fc-comprising protein polypeptide such as a “binding protein comprising an Fc region,” also referred to as an “Fc fusion protein” (e.g., an antibody or immunoadhesin).

[0054] A “native sequence Fc region” or “native sequence Fc” comprises an amino acid sequence that is identical to the amino acid sequence of an Fc region found in nature. Native sequence human Fc regions include a native sequence human IgG1 Fc region; native sequence human IgG2 Fc region; native sequence human IgG3 Fc region; and native sequence human IgG4 Fc region as well as naturally occurring variants thereof. Native sequence Fc include the various allotypes of Fcs. See, e.g., Jefferis et al. (2009) *mAbs* 1:1.

[0055] The term “epitope” or “antigenic determinant” refers to a site on an antigen (e.g., huCD40) to which an immunoglobulin or antibody specifically binds. Epitopes

within protein antigens can be formed both from contiguous amino acids (usually a linear epitope) or noncontiguous amino acids juxtaposed by tertiary folding of the protein (usually a conformational epitope). Epitopes formed from contiguous amino acids are typically, but not always, retained on exposure to denaturing solvents, whereas epitopes formed by tertiary folding are typically lost on treatment with denaturing solvents. An epitope typically includes at least 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14 or 15 amino acids in a unique spatial conformation.

[0056] The term “epitope mapping” refers to the process of identification of the molecular determinants on the antigen involved in antibody-antigen recognition. Methods for determining what epitopes are bound by a given antibody are well known in the art and include, for example, immunoblotting and immunoprecipitation assays, wherein overlapping or contiguous peptides from (e.g., from CD40) are tested for reactivity with a given antibody (e.g., anti-CD40 antibody); x-ray crystallography; 2-dimensional nuclear magnetic resonance; yeast display (see Example 6); and HDX-MS (see, e.g., Epitope Mapping Protocols in *Methods in Molecular Biology*, Vol. 66, G. E. Morris, Ed. (1996)) (see Example 5).

[0057] The term “binds to the same epitope” with reference to two or more antibodies means that the antibodies bind to the same segment of amino acid residues, as determined by a given method. Techniques for determining whether antibodies bind to the “same epitope on CD40” with the antibodies described herein include, for example, epitope mapping methods, such as, x-ray analyses of crystals of antigen:antibody complexes, which provides atomic resolution of the epitope, and hydrogen/deuterium exchange mass spectrometry (HDX-MS). Other methods monitor the binding of the antibody to antigen fragments (e.g. proteolytic fragments) or to mutated variations of the antigen where loss of binding due to a modification of an amino acid residue within the antigen sequence is often considered an indication of an epitope component, such as alanine scanning mutagenesis (Cunningham & Wells (1985) *Science* 244:1081) or yeast display of mutant target sequence variants (see Example 6). In addition, computational combinatorial methods for epitope mapping can also be used. These methods rely on the ability of the antibody of interest to affinity isolate specific short peptides from combinatorial phage display peptide libraries. Antibodies having the same or closely related VH and VL or the same CDR sequences are expected to bind to the same epitope.

[0058] Antibodies that “compete with another antibody for binding to a target” refer to antibodies that inhibit

(partially or completely) the binding of the other antibody to the target. Whether two antibodies compete with each other for binding to a target, i.e., whether and to what extent one antibody inhibits the binding of the other antibody to a target, may be determined using known competition experiments. In certain embodiments, an antibody competes with, and inhibits binding of another antibody to a target by at least 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90% or 100%. The level of inhibition or competition may be different depending on which antibody is the “blocking antibody” (i.e., the cold antibody that is incubated first with the target). Competition assays can be conducted as described, for example, in Ed Harlow and David Lane, Cold Spring Harb. Protoc.; 2006; doi:10.1101/pdb.prot4277 or in Chapter 11 of “Using Antibodies” by Ed Harlow and David Lane, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., (USA) 1999. Competing antibodies bind to the same epitope, an overlapping epitope or to adjacent epitopes (e.g., as evidenced by steric hindrance).

[0059] Other competitive binding assays include: solid phase direct or indirect radioimmunoassay (RIA), solid phase direct or indirect enzyme immunoassay (EIA), sandwich competition assay (see Stahl et al. (1983) *Methods in Enzymology* 9:242); solid phase direct biotin-avidin EIA (see Kirkland et al. (1986) *J. Immunol.* 137:3614); solid phase direct labeled assay, solid phase direct labeled sandwich assay (see Harlow and Lane (1988), *Antibodies: A Laboratory Manual*, Cold Spring Harbor Press); solid phase direct label RIA using I-125 label (see Morel et al. (1988) *Mol. Immunol.* 25(1):7); solid phase direct biotin-avidin EIA (Cheung et al. (1990) *Virology* 176:546); and direct labeled RIA. (Moldenhauer et al. (1990) *Scand. J. Immunol.* 32:77).

[0060] As used herein, the terms “specific binding,” “selective binding,” “selectively binds,” and “specifically binds,” refer to antibody binding to an epitope on a predetermined antigen but not to other antigens. Typically, the antibody (i) binds with an equilibrium dissociation constant (K_D) of approximately less than 10^{-7} M, such as approximately less than 10^{-8} M, 10^{-9} M or 10^{-10} M or even lower when determined by, e.g., surface plasmon resonance (SPR) technology in a BIACORE® 2000 surface plasmon resonance instrument using the predetermined antigen, e.g., recombinant human CD40, as the analyte and the antibody as the ligand, or Scatchard analysis of binding of the antibody to antigen positive cells, and (ii) binds to the predetermined antigen with an affinity that is at least two-fold greater than its affinity for binding to a non-specific antigen (e.g., BSA, casein) other than the predetermined antigen or a closely-related antigen. Accordingly, an antibody that “specifically binds to human CD40” refers to an antibody that binds to soluble or cell bound human CD40 with a K_D of 10^{-7} M or less, such as approximately less than 10^{-8} M, 10^{-9} M or 10^{-10} M or even lower. An antibody that “cross-reacts with cynomolgus CD40” refers to an antibody that binds to cynomolgus CD40 with a K_D of 10^{-7} M or less, such as approximately less than 10^{-8} M, 10^{-9} M or 10^{-10} M or even lower.

[0061] The term “kassoc” or “ K_A ”, as used herein, refers to the association rate constant of a particular antibody-antigen interaction, whereas the term “kdis” or “ K_D ”, as used herein, refers to the dissociation rate constant of a particular antibody-antigen interaction. The term “ K_D ”, as used herein, refers to the equilibrium dissociation constant, which is obtained from the ratio of K_D to K_A (i.e., K_D/K_A)

and is expressed as a molar concentration (M). K_D values for antibodies can be determined using methods well established in the art. A preferred method for determining the K_D of an antibody is biolayer interferometry (BLI) analysis, preferably using a FORTEBIO Octet RED device (see Example 3), surface plasmon resonance, preferably using a biosensor system such as a BIACORE® surface plasmon resonance system (see Example 4), or flow cytometry and Scatchard analysis.

[0062] The term “EC50” in the context of an in vitro or in vivo assay using an antibody or antigen binding fragment thereof, refers to the concentration of an antibody or an antigen-binding fragment thereof that induces a response that is 50% of the maximal response, i.e., halfway between the maximal response and the baseline.

[0063] The term “binds to immobilized CD40” refers to the ability of an antibody described herein to bind to CD40, for example, expressed on the surface of a cell or attached to a solid support.

[0064] The term “cross-reacts,” as used herein, refers to the ability of an antibody described herein to bind to CD40 from a different species. For example, an antibody described herein that binds human CD40 may also bind CD40 from another species (e.g., cynomolgus CD40). As used herein, cross-reactivity may be measured by detecting a specific reactivity with purified antigen in binding assays (e.g., SPR, ELISA) or binding to, or otherwise functionally interacting with, cells physiologically expressing CD40. Methods for determining cross-reactivity include standard binding assays as described herein, for example, by BIACORE® surface plasmon resonance (SPR) analysis using a BIACORE® 2000 SPR instrument (BIACORE AB, Uppsala, Sweden), or flow cytometric techniques.

[0065] The term “naturally-occurring” as used herein as applied to an object refers to the fact that an object can be found in nature. For example, a polypeptide or polynucleotide sequence that is present in an organism (including viruses) that can be isolated from a source in nature and which has not been intentionally modified by man in the laboratory is naturally-occurring.

[0066] A “polypeptide” refers to a chain comprising at least two consecutively linked amino acid residues, with no upper limit on the length of the chain. One or more amino acid residues in the protein may contain a modification such as, but not limited to, glycosylation, phosphorylation or a disulfide bond. A “protein” may comprise one or more polypeptides.

[0067] The term “nucleic acid molecule,” as used herein, is intended to include DNA molecules and RNA molecules. A nucleic acid molecule may be single-stranded or double-stranded, and may be cDNA.

[0068] Also provided are “conservative sequence modifications” to the antibody sequence provided herein, i.e. nucleotide and amino acid sequence modifications that do not abrogate the binding of the antibody encoded by the nucleotide sequence or containing the amino acid sequence, to the antigen. For example, modifications can be introduced by standard techniques known in the art, such as site-directed mutagenesis and PCR-mediated mutagenesis. Conservative sequence modifications include conservative amino acid substitutions, in which the amino acid residue is replaced with an amino acid residue having a similar side chain. Families of amino acid residues having similar side chains have been defined in the art. These families include

amino acids with basic side chains (e.g., lysine, arginine, histidine), acidic side chains (e.g., aspartic acid, glutamic acid), uncharged polar side chains (e.g., glycine, asparagine, glutamine, serine, threonine, tyrosine, cysteine, tryptophan), nonpolar side chains (e.g., alanine, valine, leucine, isoleucine, proline, phenylalanine, methionine), beta-branched side chains (e.g., threonine, valine, isoleucine) and aromatic side chains (e.g., tyrosine, phenylalanine, tryptophan, histidine). Thus, a predicted nonessential amino acid residue in an anti-CD40 antibody is preferably replaced with another amino acid residue from the same side chain family. Methods of identifying nucleotide and amino acid conservative substitutions that do not eliminate antigen binding are well-known in the art. See, e.g., Brummell et al., *Biochem.* 32:1180-1187 (1993); Kobayashi et al. *Protein Eng.* 12(10): 879-884 (1999); and Burks et al. *Proc. Natl. Acad. Sci. (USA)* 94:412-417 (1997).

[0069] Alternatively, in another embodiment, mutations can be introduced randomly along all or part of an anti-CD40 antibody coding sequence, such as by saturation mutagenesis, and the resulting modified anti-CD40 antibodies can be screened for improved binding activity.

[0070] For nucleic acids, the term “substantial homology” indicates that two nucleic acids, or designated sequences thereof, when optimally aligned and compared, are identical, with appropriate nucleotide insertions or deletions, in at least about 80% of the nucleotides, usually at least about 90% to 95%, and more preferably at least about 98% to 99.5% of the nucleotides. Alternatively, substantial homology exists when the segments will hybridize under selective hybridization conditions, to the complement of the strand.

[0071] For polypeptides, the term “substantial homology” indicates that two polypeptides, or designated sequences thereof, when optimally aligned and compared, are identical, with appropriate amino acid insertions or deletions, in at least about 80% of the amino acids, usually at least about 90% to 95%, and more preferably at least about 98% to 99.5% of the amino acids.

[0072] The percent identity between two sequences is a function of the number of identical positions shared by the sequences when the sequences are optimally aligned (i.e., % homology = # of identical positions / total # of positions × 100), with optimal alignment determined taking into account the number of gaps, and the length of each gap, which need to be introduced for optimal alignment of the two sequences. The comparison of sequences and determination of percent identity between two sequences can be accomplished using a mathematical algorithm, as described in the non-limiting examples below.

[0073] The percent identity between two nucleotide sequences can be determined using the GAP program in the GCG software package, using a NWSgapdna.CMP matrix and a gap weight of 40, 50, 60, 70, or 80 and a length weight of 1, 2, 3, 4, 5, or 6. The percent identity between two nucleotide or amino acid sequences can also be determined using the algorithm of E. Meyers and W. Miller (CABIOS, 4:11-17 (1989)) which has been incorporated into the ALIGN program (version 2.0), using a PAM120 weight residue table, a gap length penalty of 12 and a gap penalty of 4. In addition, the percent identity between two amino acid sequences can be determined using the Needleman and Wunsch (J. Mol. Biol. (48):444-453 (1970)) algorithm which has been incorporated into the GAP program in the GCG software package, using either a Blossum 62 matrix or

a PAM250 matrix, and a gap weight of 16, 14, 12, 10, 8, 6, or 4 and a length weight of 1, 2, 3, 4, 5, or 6.

[0074] The nucleic acid and protein sequences described herein can further be used as a “query sequence” to perform a search against public databases to, for example, identify related sequences. Such searches can be performed using the NBLAST and XBLAST programs (version 2.0) of Altschul, et al. (1990) J. Mol. Biol. 215:403-10. BLAST nucleotide searches can be performed with the NBLAST program, score=100, wordlength=12 to obtain nucleotide sequences homologous to the nucleic acid molecules described herein. BLAST protein searches can be performed with the XBLAST program, score=50, wordlength=3 to obtain amino acid sequences homologous to the protein molecules described herein. To obtain gapped alignments for comparison purposes, Gapped BLAST can be utilized as described in Altschul et al., (1997) *Nucleic Acids Res.* 25(17):3389-3402. When utilizing BLAST and Gapped BLAST programs, the default parameters of the respective programs (e.g., XBLAST and NBLAST) can be used.

[0075] The nucleic acids may be present in whole cells, in a cell lysate, or in a partially purified or substantially pure form. A nucleic acid is “isolated” or “rendered substantially pure” when purified away from other cellular components or other contaminants, e.g., other cellular nucleic acids (e.g., the other parts of the chromosome) or proteins, by standard techniques, including alkaline/SDS treatment, CsCl banding, column chromatography, agarose gel electrophoresis and others well known in the art. See, F. Ausubel, et al., ed. *Current Protocols in Molecular Biology*, Greene Publishing and Wiley Interscience, New York (1987).

[0076] The term “vector,” as used herein, is intended to refer to a nucleic acid molecule capable of transporting another nucleic acid to which it has been linked. One type of vector is a “plasmid,” which refers to a circular double stranded DNA loop into which additional DNA segments may be ligated. Another type of vector is a viral vector, wherein additional DNA segments may be ligated into the viral genome. Certain vectors are capable of autonomous replication in a host cell into which they are introduced (e.g., bacterial vectors having a bacterial origin of replication and episomal mammalian vectors). Other vectors (e.g., non-episomal mammalian vectors) can be integrated into the genome of a host cell upon introduction into the host cell, and thereby are replicated along with the host genome. Moreover, certain vectors are capable of directing the expression of genes to which they are operatively linked. Such vectors are referred to herein as “recombinant expression vectors” (or simply, “expression vectors”). In general, expression vectors of utility in recombinant DNA techniques are often in the form of plasmids. In the present specification, “plasmid” and “vector” may be used interchangeably as the plasmid is the most commonly used form of vector. However, also included are other forms of expression vectors, such as viral vectors (e.g., replication defective retroviruses, adenoviruses and adeno-associated viruses), which serve equivalent functions.

[0077] The term “recombinant host cell” (or simply “host cell”), as used herein, is intended to refer to a cell that comprises a nucleic acid that is not naturally present in the cell, and may be a cell into which a recombinant expression vector has been introduced. It should be understood that such terms are intended to refer not only to the particular subject cell but to the progeny of such a cell. Because certain

modifications may occur in succeeding generations due to either mutation or environmental influences, such progeny may not, in fact, be identical to the parent cell, but are still included within the scope of the term “host cell” as used herein.

[0078] An “immune response” refers to a biological response within a vertebrate against foreign agents, which response protects the organism against these agents and diseases caused by them. An immune response is mediated by the action of a cell of the immune system (for example, a T lymphocyte, B lymphocyte, natural killer (NK) cell, macrophage, eosinophil, mast cell, dendritic cell or neutrophil) and soluble macromolecules produced by any of these cells or the liver (including antibodies, cytokines, and complement) that results in selective targeting, binding to, damage to, destruction of, and/or elimination from the vertebrate’s body of invading pathogens, cells or tissues infected with pathogens, cancerous or other abnormal cells, or, in cases of autoimmunity or pathological inflammation, normal human cells or tissues. An immune reaction includes, e.g., activation or inhibition of a T cell, e.g., an effector T cell or a Th cell, such as a CD4+ or CD8+ T cell, or the inhibition or depletion of a Treg cell. “T effector” (“Teff”) cells refers to T cells (e.g., CD4+ and CD8+ T cells) with cytolytic activities as well as T helper (Th) cells, which secrete cytokines and activate and direct other immune cells, but does not include regulatory T cells (Treg cells).

[0079] As used herein, the term “T cell-mediated response” refers to a response mediated by T cells, including effector T cells (e.g., CD8+ cells) and helper T cells (e.g., CD4+ cells). T cell mediated responses include, for example, T cell cytotoxicity and proliferation.

[0080] As used herein, the term “cytotoxic T lymphocyte (CTL) response” refers to an immune response induced by cytotoxic T cells. CTL responses are mediated primarily by CD8+ T cells.

[0081] An “immunomodulator” or “immunoregulator” refers to an agent, e.g., a component of a signaling pathway that may be involved in modulating, regulating, or modifying an immune response. “Modulating,” “regulating,” or “modifying” an immune response refers to any alteration in a cell of the immune system or in the activity of such cell (e.g., an effector T cell). Such modulation includes stimulation or suppression of the immune system which may be manifested by an increase or decrease in the number of various cell types, an increase or decrease in the activity of these cells, or any other changes which can occur within the immune system. Both inhibitory and stimulatory immunomodulators have been identified, some of which may have enhanced function in a tumor microenvironment. In preferred embodiments, the immunomodulator is located on the surface of a T cell. An “immunomodulatory target” or “immunoregulatory target” is an immunomodulator that is targeted for binding by, and whose activity is altered by the binding of, a substance, agent, moiety, compound or molecule. Immunomodulatory targets include, for example, receptors on the surface of a cell (“immunomodulatory receptors”) and receptor ligands (“immunomodulatory ligands”).

[0082] “Immunotherapy” refers to the treatment of a subject afflicted with, or at risk of contracting or suffering a recurrence of, a disease by a method comprising inducing, enhancing, suppressing or otherwise modifying an immune response.

[0083] “Immunostimulating therapy” or “immunostimulatory therapy” refers to a therapy that results in increasing (inducing or enhancing) an immune response in a subject for, e.g., treating cancer.

[0084] “Potentiating an endogenous immune response” means increasing the effectiveness or potency of an existing immune response in a subject. This increase in effectiveness and potency may be achieved, for example, by overcoming mechanisms that suppress the endogenous host immune response or by stimulating mechanisms that enhance the endogenous host immune response.

[0085] As used herein, the term “linked” refers to the association of two or more molecules. The linkage can be covalent or non-covalent. The linkage also can be genetic (i.e., recombinantly fused). Such linkages can be achieved using a wide variety of art recognized techniques, such as chemical conjugation and recombinant protein production.

[0086] As used herein, “administering” refers to the physical introduction of a composition comprising a therapeutic agent to a subject, using any of the various methods and delivery systems known to those skilled in the art. Preferred routes of administration for antibodies described herein include intravenous, intraperitoneal, intramuscular, subcutaneous, spinal or other parenteral routes of administration, for example by injection or infusion. The phrase “parenteral administration” as used herein means modes of administration other than enteral and topical administration, usually by injection, and includes, without limitation, intravenous, intraperitoneal, intramuscular, intraarterial, intrathecal, intralymphatic, intralesional, intracapsular, intraorbital, intracardiac, intradermal, transtracheal, subcutaneous, subcuticular, intraarticular, subcapsular, subarachnoid, intraspinal, epidural and intrasternal injection and infusion, as well as in vivo electroporation. Alternatively, an antibody described herein can be administered via a non-parenteral route, such as a topical, epidermal or mucosal route of administration, for example, intranasally, orally, vaginally, rectally, sublingually or topically. Administering can also be performed, for example, once, a plurality of times, and/or over one or more extended periods.

[0087] As used herein, the terms “inhibits” or “blocks” are used interchangeably and encompass both partial and complete inhibition/blocking by at least about 50%, for example, at least about 60%, 70%, 80%, 90%, 95%, 99%, or 100%.

[0088] As used herein, “cancer” refers a broad group of diseases characterized by the uncontrolled growth of abnormal cells in the body. Unregulated cell division may result in the formation of malignant tumors or cells that invade neighboring tissues and may metastasize to distant parts of the body through the lymphatic system or bloodstream.

[0089] The terms “treat,” “treating,” and “treatment,” as used herein, refer to any type of intervention or process performed on, or administering an active agent to, the subject with the objective of reversing, alleviating, ameliorating, inhibiting, or slowing down or preventing the progression, development, severity or recurrence of a symptom, complication, condition or biochemical indicia associated with a disease. Prophylaxis refers to administration to a subject who does not have a disease, to prevent the disease from occurring or minimize its effects if it does.

[0090] The term “effective dose” or “effective dosage” is defined as an amount sufficient to achieve or at least partially achieve a desired effect. A “therapeutically effective amount” or “therapeutically effective dosage” of a drug or

therapeutic agent is any amount of the drug that, when used alone or in combination with another therapeutic agent, promotes disease regression evidenced by a decrease in severity of disease symptoms, an increase in frequency and duration of disease symptom-free periods, or a prevention of impairment or disability due to the disease affliction. A “prophylactically effective amount” or a “prophylactically effective dosage” of a drug is an amount of the drug that, when administered alone or in combination with another therapeutic agent to a subject at risk of developing a disease or of suffering a recurrence of disease, inhibits the development or recurrence of the disease. The ability of a therapeutic or prophylactic agent to promote disease regression or inhibit the development or recurrence of the disease can be evaluated using a variety of methods known to the skilled practitioner, such as in human subjects during clinical trials, in animal model systems predictive of efficacy in humans, or by assaying the activity of the agent in *in vitro* assays.

[0091] By way of example, an anti-cancer agent is a drug that slows cancer progression or promotes cancer regression in a subject. In preferred embodiments, a therapeutically effective amount of the drug promotes cancer regression to the point of eliminating the cancer. “Promoting cancer regression” means that administering an effective amount of the drug, alone or in combination with an anti-neoplastic agent, results in a reduction in tumor growth or size, necrosis of the tumor, a decrease in severity of at least one disease symptom, an increase in frequency and duration of disease symptom-free periods, a prevention of impairment or disability due to the disease affliction, or otherwise amelioration of disease symptoms in the patient. Pharmacological effectiveness refers to the ability of the drug to promote cancer regression in the patient. Physiological safety refers to an acceptably low level of toxicity, or other adverse physiological effects at the cellular, organ and/or organism level (adverse effects) resulting from administration of the drug.

[0092] By way of example for the treatment of tumors, a therapeutically effective amount or dosage of the drug preferably inhibits cell growth or tumor growth by at least about 20%, more preferably by at least about 40%, even more preferably by at least about 60%, and still more preferably by at least about 80% relative to untreated subjects. In the most preferred embodiments, a therapeutically effective amount or dosage of the drug completely inhibits cell growth or tumor growth, i.e., preferably inhibits cell growth or tumor growth by 100%. The ability of a compound to inhibit tumor growth can be evaluated using the assays described *infra*. Inhibition of tumor growth may not be immediate after treatment, and may only occur after a period of time or after repeated administration. Alternatively, this property of a composition can be evaluated by examining the ability of the compound to inhibit cell growth, such inhibition can be measured *in vitro* by assays known to the skilled practitioner. In other preferred embodiments described herein, tumor regression may be observed and may continue for a period of at least about 20 days, more preferably at least about 40 days, or even more preferably at least about 60 days.

[0093] “Combination” therapy, as used herein, unless otherwise clear from the context, is meant to encompass administration of two or more therapeutic agents in a coordinated fashion, and includes, but is not limited to, concurrent dosing. Specifically, combination therapy encompasses both

co-administration (e.g. administration of a co-formulation or simultaneous administration of separate therapeutic compositions) and serial or sequential administration, provided that administration of one therapeutic agent is conditioned in some way on administration of another therapeutic agent. For example, one therapeutic agent may be administered only after a different therapeutic agent has been administered and allowed to act for a prescribed period of time. See, e.g., Kohrt et al. (2011) *Blood* 117:2423.

[0094] The terms “patient” and “subject” refer to any human that receives either prophylactic or therapeutic treatment. For example, the methods and compositions described herein can be used to treat a subject having cancer.

[0095] Various aspects described herein are described in further detail in the following subsections.

I. Anti-CD40 Antibodies

[0096] The present application discloses agonistic anti-huCD40 antibodies having desirable properties for use as therapeutic agents in treating diseases such as cancers. These properties include one or more of the ability to bind to human CD40 with high affinity, acceptably low immunogenicity in human subjects, the ability to bind preferentially to FcγRIIb, and the absence of sequence liabilities that might reduce the chemical stability of the antibody.

[0097] The anti-CD40 antibodies disclosed herein by sequence bind to specific epitopes on human CD40, as may be determined as described in Examples 5 and 6. Other antibodies that bind to the same or closely related epitopes would likely share these desirable properties, and may be discovered doing competition experiments.

Anti-huCD40 Antibodies that Compete with Anti-huCD40 Antibodies Disclosed Herein

[0098] Anti-huCD40 antibodies that compete with the antibodies of the present invention for binding to huCD40 may be raised using immunization protocols similar to those described herein (Examples 1 and 2). Antibodies that compete for binding with the anti-huCD40 antibodies disclosed herein by sequence may also be generated by immunizing mice or other non-human animal with human CD40 or a construct comprising the extracellular domain thereof (residues 21-193 of SEQ ID NO: 1), or by immunizing with a fragment of human CD40 containing the epitope bound by the anti-huCD40 antibodies disclosed herein. The resulting antibodies can be screened for the ability to block binding of 12D6, 5F11, 8E8, 5G7 and/or 19G3 to human CD40 by methods well known in the art, for example blocking binding to fusion protein of the extracellular domain of CD40 and an immunoglobulin Fc domain in a ELISA, or blocking the ability to bind to cells expressing huCD40 on their surface, e.g. by FACS. In various embodiments, the test antibody is contacted with the CD40-Fc fusion protein (or to cells expressing huCD40 on their surface) prior to, at the same time as, or after the addition of 12D6, 5F11, 8E8, 5G7 or 19G3. For example, “binning” experiments may be performed (Example 4) to determine whether a test antibody falls into the same “bin” as an antibodies disclosed herein by sequence, with antibodies disclosed herein by sequence as the “reference” antibodies and the antibodies to be tested as the “test” antibodies. Antibodies that reduce binding of the antibodies disclosed herein by sequence to human CD40 (either as an Fc fusion or on a cell), particularly at roughly stoichiometric concentrations, are likely to bind at the same,

overlapping, or adjacent epitopes, and thus may share the desirable functional properties of 12D6, 5F11, 8E8, 5G7 or 19G3.

[0099] Accordingly, provided herein are anti-huCD40 antibodies that inhibit the binding of an anti-huCD40 antibodies described herein to huCD40 on cells by at least 10%, 20%, 30%, 40%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or by 100%, and/or whose binding to huCD40 on cells is inhibited by an anti-huCD40 antibodies described herein by at least 10%, 20%, 30%, 40%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or by 100%, e.g., as measured by ELISA or FACS, such as by using the assay described in the following paragraph.

[0100] An exemplary competition experiment to determine whether a test antibody blocks the binding of (i.e., “competes with”) a reference antibody, may be conducted as follows: cells expressing CD40 are seeded at 10^5 cells per sample well in a 96 well plate. The plate is set on ice followed by the addition of unconjugated test antibody at concentrations ranging from 0 to 50 $\mu\text{g/mL}$ (three-fold titration starting from a highest concentration of 50 $\mu\text{g/mL}$). An unrelated IgG may be used as an isotype control for the first antibody and added at the same concentrations (three-fold titration starting from a highest concentration of 50 $\mu\text{g/mL}$). A sample pre-incubated with 50 $\mu\text{g/mL}$ unlabeled reference antibody may be included as a positive control for complete blocking (100% inhibition) and a sample without antibody in the primary incubation may be used as a negative control (no competition; 0% inhibition). After 30 minutes of incubation, labeled, e.g., biotinylated, reference antibody is added at a concentration of 2 $\mu\text{g/mL}$ per well without washing. Samples are incubated for another 30 minutes on ice. Unbound antibodies are removed by washing the cells with FACS buffer. Cell-bound labeled reference antibody is detected with an agent that detects the label, e.g., PE conjugated streptavidin (INVITROGEN, catalog #521388) for detecting biotin. The samples are acquired on a FACS Calibur Flow Cytometer (BD, San Jose) and analyzed with FLOWJO software (TREE STAR, Inc, Ashland, Oreg.). The results may be represented as the % inhibition.

[0101] Typically, the same experiment is then conducted in the reverse, i.e., the test antibody is the reference antibody and the reference antibody is the test antibody. In certain embodiments, an antibody at least partially (e.g., at least 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, or 90%) or completely (100%) blocks the binding of the other antibody to the target, e.g. human CD40 or fragment thereof, and regardless of whether inhibition occurs when one or the other antibody is the reference antibody. A reference antibody and a test antibody “cross-block” binding of each other to the target when the antibodies compete with each other both ways, i.e., in competition experiments in which the reference antibody is added first and in competition experiments in which the test antibody is added first.

[0102] Anti-huCD40 antibodies are considered to compete with the anti-huCD40 antibodies disclosed herein if they inhibit binding of 12D6, 5F11, 8E8, 5G7 and/or 19G3 to human CD40 by at least 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, or by 100%, when present at roughly equal concentrations, for example in competition experiments like those described in Example 4. Unless otherwise indicated, an antibody will be considered to compete with an antibody

selected from the group consisting of the anti-CD40 antibodies of the present invention if it reduces binding of the selected antibody to human CD40 (SEQ ID NO: 1) by at least 20% when used at a roughly equal molar concentration with the selected antibody, as measured in competition ELISA experiments as outlined in the preceding two paragraphs.

Anti-huCD40 Antibodies that Bind to the Same Epitope

[0103] Anti-huCD40 antibodies that bind to the same or similar epitopes to the antibodies disclosed herein may be raised using immunization protocols similar to those described herein (Examples 1 and 2). The resulting antibodies can be screened for high affinity binding to human CD40 (Example 3). Selected antibodies can then be studied in yeast display assay in which sequence variants of huCD40 are presented on the surface of yeast cells (Example 6), or by hydrogen-deuterium exchange experiments (Example 5), to determine the precise epitope bound by the antibody.

[0104] Epitope determinations may be made by any method known in the art. In various embodiments, anti-huCD40 antibodies are considered to bind to the same epitope as an anti-huCD40 mAb disclosed herein if they make contact with one or more of the same residues within at least one region of huCD40; if they make contacts with a majority of the residues within at least one region of huCD40; if they make contacts with a majority of the residues within each region of huCD40; if they make contact with a majority of contacts along the entire length of huCD40; if they make contacts within all of the same distinct regions of human CD40; if they make contact with all of the residues at any one region on human CD40; or if they make contact with all of the same residues at all of the same regions. Epitope “regions” are clusters of residues along the primary sequence.

[0105] Techniques for determining antibodies that bind to the “same epitope on huCD40” with the antibodies described herein include x-ray analyses of crystals of antigen:antibody complexes, which provides atomic resolution of the epitope. Other methods monitor the binding of the antibody to antigen fragments or mutated variations of the antigen where loss of binding due to a modification of an amino acid residue within the antigen sequence is often considered an indication of an epitope component. Methods may also rely on the ability of an antibody of interest to affinity isolate specific short peptides (either in native three dimensional form or in denatured form) from combinatorial phage display peptide libraries or from a protease digest of the target protein. The peptides are then regarded as leads for the definition of the epitope corresponding to the antibody used to screen the peptide library. For epitope mapping, computational algorithms have also been developed that have been shown to map conformational discontinuous epitopes.

[0106] The epitope or region comprising the epitope can also be identified by screening for binding to a series of overlapping peptides spanning CD40. Alternatively, the method of Jespers et al. (1994) *Biotechnology* 12:899 may be used to guide the selection of antibodies having the same epitope and therefore similar properties to the an anti-CD40 antibodies described herein. Using phage display, first the heavy chain of the anti-CD40 antibody is paired with a repertoire of (preferably human) light chains to select a CD40-binding antibody, and then the new light chain is paired with a repertoire of (preferably human) heavy chains

to select a (preferably human) CD40-binding antibody having the same epitope or epitope region as an anti-huCD40 antibody described herein. Alternatively variants of an antibody described herein can be obtained by mutagenesis of cDNA encoding the heavy and light chains of the antibody.

[0107] Alanine scanning mutagenesis, as described by Cunningham & Wells (1989) *Science* 244: 1081, or some other form of point mutagenesis of amino acid residues in CD40 (such as the yeast display method provided at Example 6) may also be used to determine the functional epitope for an anti-CD40 antibody.

[0108] The epitope or epitope region (an "epitope region" is a region comprising the epitope or overlapping with the epitope) bound by a specific antibody may also be determined by assessing binding of the antibody to peptides comprising fragments of CD40. A series of overlapping peptides encompassing the sequence of CD40 (e.g., human CD40) may be synthesized and screened for binding, e.g. in a direct ELISA, a competitive ELISA (where the peptide is assessed for its ability to prevent binding of an antibody to CD40 bound to a well of a microtiter plate), or on a chip. Such peptide screening methods may not be capable of detecting some discontinuous functional epitopes, i.e. functional epitopes that involve amino acid residues that are not contiguous along the primary sequence of the CD40 polypeptide chain.

[0109] An epitope may also be identified by MS-based protein footprinting, such as hydrogen/deuterium exchange mass spectrometry (HDX-MS) and Fast Photochemical Oxidation of Proteins (FPOP). HDX-MS may be conducted, e.g., as further described at Wei et al. (2014) *Drug Discovery Today* 19:95, the methods of which are specifically incorporated by reference herein. See also Example 5. FPOP may be conducted as described, e.g., in Hambley & Gross (2005) *J. American Soc. Mass Spectrometry* 16:2057, the methods of which are specifically incorporated by reference herein.

[0110] The epitope bound by anti-CD40 antibodies may also be determined by structural methods, such as X-ray crystal structure determination (e.g., WO 2005/044853), molecular modeling and nuclear magnetic resonance (NMR) spectroscopy, including NMR determination of the H-D exchange rates of labile amide hydrogens in CD40 when free and when bound in a complex with an antibody of interest (Zinn-Justin et al. (1992) *Biochemistry* 31:11335; Zinn-Justin et al. (1993) *Biochemistry* 32:6884).

[0111] With regard to X-ray crystallography, crystallization may be accomplished using any of the known methods in the art (e.g. Giege et al. (1994) *Acta Crystallogr. D* 50:339; McPherson (1990) *Eur. J. Biochem.* 189:1), including microbatch (e.g. Chayen (1997) *Structure* 5:1269), hanging-drop vapor diffusion (e.g. McPherson (1976) *J. Biol. Chem.* 251:6300), seeding and dialysis. It is desirable to use a protein preparation having a concentration of at least about 1 mg/mL and preferably about 10 mg/mL to about 20 mg/mL. Crystallization may be best achieved in a precipitant solution containing polyethylene glycol 1000-20,000 (PEG; average molecular weight ranging from about 1000 to about 20,000 Da), preferably about 5000 to about 7000 Da, more preferably about 6000 Da, with concentrations ranging from about 10% to about 30% (w/v). It may also be desirable to include a protein stabilizing agent, e.g. glycerol at a concentration ranging from about 0.5% to about 20%. A suitable salt, such as sodium chloride, lithium chloride or sodium citrate may also be desirable in the precipitant

solution, preferably in a concentration ranging from about 1 mM to about 1000 mM. The precipitant is preferably buffered to a pH of from about 3.0 to about 5.0, preferably about 4.0. Specific buffers useful in the precipitant solution may vary and are well-known in the art (Scopes, Protein Purification: Principles and Practice, Third ed., (1994) Springer-Verlag, New York). Examples of useful buffers include, but are not limited to, HEPES, Tris, MES and acetate. Crystals may be grown at a wide range of temperatures, including 2° C., 4° C., 8° C. and 26° C.

[0112] Antibody:antigen crystals may be studied using well-known X-ray diffraction techniques and may be refined using computer software such as X-PLOR (Yale University, 1992, distributed by Molecular Simulations, Inc.; see e.g. Blundell & Johnson (1985) *Meth. Enzymol.* 114 & 115, H. W. Wyckoff et al., eds., Academic Press; U.S. Patent Application Publication No. 2004/0014194), and BUSTER (Bricogne (1993) *Acta Cryst. D* 49:37-60; Bricogne (1997) *Meth. Enzymol.* 276A:361-423, Carter & Sweet, eds.; Rovnersi et al. (2000) *Acta Cryst. D* 56:1313-1323), the disclosures of which are hereby incorporated by reference in their entireties.

[0113] Unless otherwise indicated, and with reference to the claims, the epitope bound by an antibody is the epitope as determined by HDX-MS methods, substantially as described in Example 5.

Anti-CD40 Antibodies that Bind with High Affinity

[0114] In some embodiments the anti-huCD40 antibodies of the present invention bind to huCD40 with high affinity, like the anti-huCD40 antibodies disclosed herein, increasing their likelihood of being effective therapeutic agents. In various embodiments anti-huCD40 antibodies of the present invention bind to huCD40 with a K_D of less than 10 nM, 5 nM, 2 nM, 1 nM, 300 pM or 100 pM. In other embodiments, the anti-huCD40 antibodies of the present invention bind to huCD40 with a K_D between 2 nM and 100 pM. Standard assays to evaluate the binding ability of the antibodies toward huCD40 include ELISAs, RIAs, Western blots, bio-layer interferometry (BLI) (see Example 3) and BIA-CORE® SPR analysis (see Example 4).

Anti-CD40 Antibody Sequence Variants

[0115] Some variability in the antibody sequences disclosed herein may be tolerated and still maintain the desirable properties of the antibody. The CDR regions are delineated using the Kabat system (Kabat, E. A., et al. (1991) *Sequences of Proteins of Immunological Interest*, Fifth Edition, U.S. Department of Health and Human Services, NIH Publication No. 91-3242). Accordingly, the present invention further provides anti-huCD40 antibodies comprising CDR sequences that are at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical to the CDR sequences of the antibodies disclosed herein (i.e. 12D6, 5F11, 8E8, 5G7 and 19G3 and humanized derivatives thereof). The present invention also provides anti-huCD40 antibodies comprising heavy and/or light chain variable domain sequences that are at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical to the heavy and/or light chain variable domain sequences of the antibodies disclosed herein (i.e. 12D6, 5F11, 8E8, 5G7 and 19G3 and humanized derivatives thereof).

Anti-CD40 Antibodies Sharing CDR Sequences or Derived from the Same Murine Germlines

[0116] Given that antigen-binding specificity is determined primarily by the CDRs, antibodies sharing CDRs sequences with antibodies disclosed herein (i.e. 12D6, 5F11, 8E8, 5G7 and 19G3) are likely to share their desirable properties. In some embodiments, anti-huCD40 antibodies of the present invention comprises heavy and light chain variable regions derived from the same murine V region and J region germline sequences as antibody 12D6, 5F11, 8E8, 5G7 or 19G3. Antibody 12D6 has a heavy chain derived from murine germlines VH1-39_01 and IGHJ4, and light chain germlines VK1-110_01 and IGKJ1. Antibody 5F11 has a heavy chain derived from murine germlines VH1-4_02 and IGHJ3, and light chain germlines VK3-5_01 and IGKJ5. Antibody 8E8 has a heavy chain derived from murine germlines VH1-80_01 and IGHJ2, and light chain germlines VK1-110_01 and IGKJ2. Antibody 5G7 has a heavy chain derived from murine germlines VH1-18_01 and IGHJ4, and light chain germlines VK10-96_01 and IGKJ2. Antibody 19G3 has a heavy chain derived from murine germlines VH5-9-4_01 and IGHJ3, and light chain germlines VK1-117_01 and IGKJ2. Heavy chain D region germline sequences (making up part of CDRH3) are not specified, as they are often difficult to assign given their high variability, and thus antibodies of the present invention may comprise heavy chains derived from the listed V and J region germlines and any D region germline. Other antibodies that bind to human CD40 and are derived from some or all of these germline sequences are likely to be closely related in sequence, particularly those derived from the same V-region genes, and thus would be expected to share the same desirable properties.

[0117] As used herein, a murine antibody comprises heavy or light chain variable regions that are “derived from” a particular germline sequence if the variable regions of the antibody are obtained from a system that uses murine germline immunoglobulin genes, and the antibody sequence is sufficiently related to the germline that it is more likely derived from the given germline than from any other. Such systems include immunizing a mouse with the antigen of interest. The murine germline immunoglobulin sequence(s) from which the sequence of an antibody is “derived” can be identified by comparing the amino acid sequence of the antibody to the amino acid sequences of murine germline immunoglobulins and selecting the germline immunoglobulin sequence that is closest in sequence (i.e., greatest % identity) to the sequence of the antibody. A murine antibody that is “derived from” a particular germline immunoglobulin sequence may contain amino acid differences as compared to the germline sequence due to, for example, naturally-occurring somatic mutations or intentional introduction of site-directed mutation. However, a selected murine antibody typically is at least 90% identical in amino acids sequence to an amino acid sequence encoded by a germline immunoglobulin gene (e.g. V regions). In certain cases, a murine antibody may be at least 95%, or even at least 96%, 97%, 98%, or 99% identical in amino acid sequence to the amino acid sequence encoded by the germline immunoglobulin gene (e.g. V regions). Typically, an antibody derived from a particular murine germline sequence will display no more than 10 amino acid differences from the amino acid sequence encoded by the germline immunoglobulin gene (e.g. V regions). In certain cases, the murine antibody may

comprise no more than 5, or even no more than 4, 3, 2, or 1 amino acid difference from the amino acid sequence encoded by the germline immunoglobulin gene (e.g. V regions).

II. Engineered and Modified Antibodies

VH and VL Regions

[0118] Also provided are engineered and modified antibodies that can be prepared using an antibody having one or more of the V_H and/or V_L sequences disclosed herein as starting material to engineer a modified antibody, which modified antibody may have altered properties from the starting antibody. An antibody can be engineered by modifying one or more residues within one or both variable regions (i.e., VH and/or VL), for example within one or more CDR regions and/or within one or more framework regions. Additionally or alternatively, an antibody can be engineered by modifying residues within the constant region (s), for example to alter the effector function(s) of the antibody.

[0119] One type of variable region engineering that can be performed is CDR grafting. Such grafting is of particular use in humanizing non-human anti-CD40 antibodies that compete for binding with the anti-huCD40 antibodies disclosed herein and/or bind to the same epitope as the anti-huCD40 antibodies disclosed herein. Antibodies interact with target antigens predominantly through amino acid residues that are located in the six heavy and light chain complementarity determining regions (CDRs). For this reason, the amino acid sequences within CDRs are more diverse between individual antibodies than sequences outside of CDRs. Because CDR sequences are responsible for most antibody-antigen interactions, it is possible to express recombinant antibodies that mimic the properties of specific reference antibodies by constructing expression vectors that include CDR sequences from the specific reference antibody grafted onto framework sequences from a different antibody with different properties (see, e.g., Riechmann, L. et al. (1998) *Nature* 332:323-327; Jones, P. et al. (1986) *Nature* 321:522-525; Queen, C. et al. (1989) *Proc. Natl. Acad. Sci. (USA)* 86:10029-10033; U.S. Pat. No. 5,225,539 to Winter, and U.S. Pat. Nos. 5,530,101; 5,585,089; 5,693,762 and 6,180,370 to Queen et al.)

[0120] Such framework sequences can be obtained from public DNA databases or published references that include germline antibody gene sequences. For example, germline DNA sequences for human heavy and light chain variable region genes can be found in the “VBase” human germline sequence database, as well as in Kabat, E. A., et al. (1991) *Sequences of Proteins of Immunological Interest*, Fifth Edition, U.S. Department of Health and Human Services, NIH Publication No. 91-3242; Tomlinson, I. M., et al. (1992) “The Repertoire of Human Germline VH Sequences Reveals about Fifty Groups of VH Segments with Different Hypervariable Loops” *J. Mol. Biol.* 227:776-798; and Cox, J. P. L. et al. (1994) “A Directory of Human Germ-line VH Segments Reveals a Strong Bias in their Usage” *Eur. J. Immunol.* 24:827-836; the contents of each of which are expressly incorporated herein by reference.

[0121] Preferred framework sequences for use in the antibodies described herein are those that are structurally similar to the framework sequences used by antibodies described herein. The VH CDR1, 2 and 3 sequences, and the VL CDR1, 2 and 3 sequences, can be grafted onto framework

regions that have the identical sequence as that found in the germline immunoglobulin gene from which the framework sequence derive, or the CDR sequences can be grafted onto framework regions that contain up to 20, preferably conservative, amino acid substitutions as compared to the germline sequences. For example, it has been found that in certain instances it is beneficial to mutate residues within the framework regions to maintain or enhance the antigen binding ability of the antibody (see e.g., U.S. Pat. Nos. 5,530,101; 5,585,089; 5,693,762 and 6,180,370 to Queen et al).

[0122] Engineered antibodies described herein include those in which modifications have been made to framework residues within V_H and/or V_L , e.g. to improve the properties of the antibody. Often such framework modifications are made to decrease the immunogenicity of the antibody. For example, one approach is to “backmutate” one or more framework residues to the corresponding germline sequence. More specifically, an antibody that has undergone somatic mutation may contain framework residues that differ from the germline sequence from which the antibody is derived. Such residues can be identified by comparing the antibody framework sequences to the germline sequences from which the antibody is derived. To return the framework region sequences to their germline configuration, the somatic mutations can be “backmutated” to the germline sequence by, for example, site-directed mutagenesis or PCR-mediated mutagenesis. Such “backmutated” antibodies are also intended to be encompassed.

[0123] Another type of framework modification involves mutating one or more residues within the framework region, or even within one or more CDR regions, to remove T cell epitopes to thereby reduce the potential immunogenicity of the antibody. This approach is also referred to as “deimmunization” and is described in further detail in U.S. Patent Publication No. 20030153043 by Carr et al.

[0124] Another type of variable region modification is to mutate amino acid residues within the CDR regions to improve one or more binding properties (e.g., affinity) of the antibody of interest. Site-directed mutagenesis or PCR-mediated mutagenesis can be performed to introduce the mutation(s) and the effect on antibody binding, or other functional property of interest. Preferably conservative modifications are introduced. The mutations may be amino acid additions, deletions, or preferably substitutions. Moreover, typically no more than one, two, three, four or five residues within a CDR region are altered.

[0125] Methionine residues in CDRs of antibodies can be oxidized, resulting in potential chemical degradation and consequent reduction in potency of the antibody. Accordingly, also provided are anti-CD40 antibodies that have one or more methionine residues in the heavy and/or light chain CDRs replaced with amino acid residues that do not undergo oxidative degradation. Similarly, deamidation sites may be removed from anti-CD40 antibodies, particularly in the CDRs. Potential glycosylation sites within the antigen binding domain are preferably eliminated to prevent glycosylation that may interfere with antigen binding. See, e.g., U.S. Pat. No. 5,714,350.

Targeted Antigen Binding

[0126] In various embodiments, the antibody of the present invention is modified to selectively block antigen binding in tissues and environments where antigen binding

would be detrimental, but allow antigen binding where it would be beneficial. In one embodiment, a blocking peptide “mask” is generated that specifically binds to the antigen binding surface of the antibody and interferes with antigen binding, which mask is linked to each of the binding arms of the antibody by a peptidase cleavable linker. See, e.g., U.S. Pat. No. 8,518,404 to CytomX. Such constructs are useful for treatment of cancers in which protease levels are greatly increased in the tumor microenvironment compared with non-tumor tissues. Selective cleavage of the cleavable linker in the tumor microenvironment allows disassociation of the masking/blocking peptide, enabling antigen binding selectively in the tumor, rather than in peripheral tissues in which antigen binding might cause unwanted side effects.

[0127] Alternatively, in a related embodiment, a bivalent binding compound (“masking ligand”) comprising two antigen binding domains is developed that binds to both antigen binding surfaces of the (bivalent) antibody and interfere with antigen binding, in which the two binding domains masks are linked to each other (but not the antibody) by a cleavable linker, for example cleavable by a peptidase. See, e.g., Int'l Pat. App. Pub. No. WO 2010/077643 to Tegopharm Corp. Masking ligands may comprise, or be derived from, the antigen to which the antibody is intended to bind, or may be independently generated. Such masking ligands are useful for treatment of cancers in which protease levels are greatly increased in the tumor microenvironment compared with non-tumor tissues. Selective cleavage of the cleavable linker in the tumor microenvironment allows disassociation of the two binding domains from each other, reducing the avidity for the antigen-binding surfaces of the antibody. The resulting dissociation of the masking ligand from the antibody enables antigen binding selectively in the tumor, rather than in peripheral tissues in which antigen binding might cause unwanted side effects.

Fcs and Modified Fcs

[0128] Antibodies of the present invention may comprise the variable domains of the invention combined with constant domains comprising different Fc regions, selected based on the biological activities (if any) of the antibody for the intended use. Salfeld (2007) *Nat. Biotechnol.* 25:1369. Human IgGs, for example, can be classified into four subclasses, IgG1, IgG2, IgG3, and IgG4, and each these of these comprises an Fc region having a unique profile for binding to one or more of Fcγ receptors (activating receptors FcγRI (CD64), FcγRIIA, FcγRIIC (CD32a,c); FcγRIIIA and FcγRIIIB (CD16a,b) and inhibiting receptor FcγRIIB (CD32b), and for the first component of complement (C1q). Human IgG1 and IgG3 bind to all Fcγ receptors; IgG2 binds to FcγRIIA_{H131}, and with lower affinity to FcγRIIA_{R131}; FcγRIIIA_{V158}; IgG4 binds to FcγRI, FcγRIIA, FcγRIIB, FcγRIIC, and FcγRIIIA_{V158}; and the inhibitory receptor FcγRIIB has a lower affinity for IgG1, IgG2 and IgG3 than all other Fcγ receptors. Bruhns et al. (2009) *Blood* 113:3716. Studies have shown that FcγRI does not bind to IgG2, and FcγRIIIB does not bind to IgG2 or IgG4. Id. In general, with regard to ADCC activity, human IgG1 □ IgG3 □ IgG4 □ IgG2. As a consequence, for example, an IgG1 constant domain, rather than an IgG2 or IgG4, might be chosen for use in a drug where ADCC is desired; IgG3 might be chosen if activation of FcγRIIIA-expressing NK cells, monocytes of macrophages; and IgG4 might be chosen if the antibody is

to be used to desensitize allergy patients. IgG4 may also be selected if it is desired that the antibody lack all effector function.

[0129] Anti-huCD40 variable regions described herein may be linked (e.g., covalently linked or fused) to an Fc, e.g., an IgG1, IgG2, IgG3 or IgG4 Fc, which may be of any allotype or isoallotype, e.g., for IgG1: G1m, G1m1(a), G1m2(x), G1m3(f), G1m17(z); for IgG2: G2m, G2m23(n); for IgG3: G3m, G3m21(g1), G3m28(g5), G3m11(b0), G3m5(b1), G3m13(b3), G3m14(b4), G3m10(b5), G3m15(s), G3m16(t), G3m6(c3), G3m24(c5), G3m26(u), G3m27(v). See, e.g., Jefferis et al. (2009) mAbs 1:1). Selection of allotype may be influenced by the potential immunogenicity concerns, e.g. to minimize the formation of anti-drug antibodies.

[0130] In preferred embodiments, anti-CD40 antibodies of the present invention have an Fc that binds to or has enhanced binding to FcγRIIb, which can provide enhanced agonism. See, e.g., WO 2012/087928; Li & Ravetch (2011) *Science* 333:1030; Wilson et al. (2011) *Cancer Cell* 19:101; White et al. (2011) *J. Immunol.* 187:1754. Variable regions described herein may be linked to Fc variants that enhance affinity for the inhibitory receptor FcγRIIb, e.g. to enhance apoptosis-inducing or adjuvant activity. Li & Ravetch (2012) *Proc. Nat'l Acad. Sci. (USA)* 109:10966; U.S. Pat. App. Pub. 2014/0010812. Such variants may provide an antibody with immunomodulatory activities related to FcγRIIb+ cells, including for example B cells and monocytes. In one embodiment, the Fc variants provide selectively enhanced affinity to FcγRIIb relative to one or more activating receptors. Such variants may also exhibit enhanced FcR-mediated cross-linking, resulting in enhanced therapeutic efficacy. Modifications for altering binding to FcγRIIb include one or more modifications at a position selected from the group consisting of 234, 235, 236, 237, 239, 266, 267, 268, 325, 326, 327, 328, and 332, according to the EU index. Exemplary substitutions for enhancing FcγRIIb affinity include but are not limited to 234D, 234E, 234F, 234W, 235D, 235F, 235R, 235Y, 236D, 236N, 237D, 237N, 239D, 239E, 266M, 267D, 267E, 268D, 268E, 327D, 327E, 328F, 328W, 328Y, and 332E. Exemplary substitutions include 235Y, 236D, 239D, 266M, 267E, 268D, 268E, 328F, 328W, and 328Y. Other Fc variants for enhancing binding to FcγRIIb include 235Y-267E, 236D-267E, 239D-268D, 239D-267E, 267E-268D, 267E-268E, and 267E-328F. Specifically, the S267E, G236D, S239D, L328F and I332E variants, including the S267E-L328F double variant, of human IgG1 are of particular value in specifically enhancing affinity for the inhibitory FcγRIIb receptor. Chu et al. (2008) *Mol. Immunol.* 45:3926; U.S. Pat. App. Pub. 2006/024298; WO 2012/087928. Enhanced specificity for FcγRIIb (as distinguished from FcγRIIIa_{R131}) may be obtained by adding the P238D substitution and other mutations (Mimoto et al. (2013) *Protein. Eng. Des. & Selection* 26:589; WO 2012/1152410), as well as V262E and V264E (Yu et al. (2013) *J. Am. Chem. Soc.* 135:9723, and WO 2014/184545. See Table 4 (supra).

Half-Life Extension

[0131] In certain embodiments, the antibody is modified to increase its biological half-life. Various approaches are possible. For example, this may be done by increasing the binding affinity of the Fc region for FcRn. In one embodiment, the antibody is altered within the CH1 or CL region to

contain a salvage receptor binding epitope taken from two loops of a CH2 domain of an Fc region of an IgG, as described in U.S. Pat. Nos. 5,869,046 and 6,121,022 by Presta et al. Other exemplary Fc variants that increase binding to FcRn and/or improve pharmacokinetic properties include substitutions at positions 259, 308, and 434, including for example 259I, 308F, 428L, 428M, 434S, 434H, 434F, 434Y, and 434M. Other variants that increase Fc binding to FcRn include: 250E, 250Q, 428L, 428F, 250Q/428L (Hinton et al., 2004, *J. Biol. Chem.* 279(8): 6213-6216, Hinton et al. 2006 *Journal of Immunology* 176:346-356), 256A, 272A, 305A, 307A, 311A, 312A, 378Q, 380A, 382A, 434A (Shields et al., *Journal of Biological Chemistry*, 2001, 276(9):6591-6604), 252F, 252Y, 252W, 254T, 256Q, 256E, 256D, 433R, 434F, 434Y, 252Y/254T/256E, 433K/434F/436H (Dall'Acqua et al., *Journal of Immunology*, 2002, 169:5171-5180, Dall'Acqua et al., 2006, *Journal of Biological Chemistry* 281:23514-23524). See U.S. Pat. No. 8,367,805.

[0132] Modification of certain conserved residues in IgG Fc (1253, H310, Q311, H433, N434), such as the N434A variant (Yeung et al. (2009) *J. Immunol.* 182:7663), have been proposed as a way to increase FcRn affinity, thus increasing the half-life of the antibody in circulation. WO 98/023289. The combination Fc variant comprising M428L and N434S has been shown to increase FcRn binding and increase serum half-life up to five-fold. Zalevsky et al. (2010) *Nat. Biotechnol.* 28:157. The combination Fc variant comprising T307A, E380A and N434A modifications also extends half-life of IgG1 antibodies. Petkova et al. (2006) *Int. Immunol.* 18:1759. In addition, combination Fc variants comprising M252Y-M428L, M428L-N434H, M428L-N434F, M428L-N434Y, M428L-N434A, M428L-N434M, and M428L-N434S variants have also been shown to extend half-life. WO 2009/086320.

[0133] Further, a combination Fc variant comprising M252Y, S254T and T256E, increases half-life-nearly 4-fold. Dall'Acqua et al. (2006) *J. Biol. Chem.* 281:23514. A related IgG1 modification providing increased FcRn affinity but reduced pH dependence (M252Y-S254T-T256E-H433K-N434F) has been used to create an IgG1 construct ("MST-HN Abdeg") for use as a competitor to prevent binding of other antibodies to FcRn, resulting in increased clearance of that other antibody, either endogenous IgG (e.g. in an autoimmune setting) or another exogenous (therapeutic) mAb. Vaccaro et al. (2005) *Nat. Biotechnol.* 23:1283; WO 2006/130834.

[0134] Other modifications for increasing FcRn binding are described in Yeung et al. (2010) *J. Immunol.* 182:7663-7671; 6,277,375; 6,821,505; WO 97/34631; WO 2002/060919.

[0135] In certain embodiments, hybrid IgG isotypes may be used to increase FcRn binding, and potentially increase half-life. For example, an IgG1/IgG3 hybrid variant may be constructed by substituting IgG1 positions in the CH2 and/or CH3 region with the amino acids from IgG3 at positions where the two isotypes differ. Thus a hybrid variant IgG antibody may be constructed that comprises one or more substitutions, e.g., 274Q, 276K, 300F, 339T, 356E, 358M, 384S, 392N, 397M, 422I, 435R, and 436F. In other embodiments described herein, an IgG1/IgG2 hybrid variant may be constructed by substituting IgG2 positions in the CH2 and/or CH3 region with amino acids from IgG1 at positions where the two isotypes differ. Thus a hybrid variant IgG antibody

may be constructed that comprises one or more substitutions, e.g., one or more of the following amino acid substitutions: 233E, 234L, 235L, -236G (referring to an insertion of a glycine at position 236), and 327A. See U.S. Pat. No. 8,629,113. A hybrid of IgG1/IgG2/IgG4 sequences has been generated that purportedly increases serum half-life and improves expression. U.S. Pat. No. 7,867,491 (sequence number 18 therein).

[0136] The serum half-life of the antibodies of the present invention can also be increased by pegylation. An antibody can be pegylated to, for example, increase the biological (e.g., serum) half-life of the antibody. To pegylate an antibody, the antibody, or fragment thereof, typically is reacted with a polyethylene glycol (PEG) reagent, such as a reactive ester or aldehyde derivative of PEG, under conditions in which one or more PEG groups become attached to the antibody or antibody fragment. Preferably, the pegylation is carried out via an acylation reaction or an alkylation reaction with a reactive PEG molecule (or an analogous reactive water-soluble polymer). As used herein, the term “polyethylene glycol” is intended to encompass any of the forms of PEG that have been used to derivatize other proteins, such as mono (C1-C10) alkoxy- or aryloxy-polyethylene glycol or polyethylene glycol-maleimide. In certain embodiments, the antibody to be pegylated is an aglycosylated antibody. Methods for pegylating proteins are known in the art and can be applied to the antibodies described herein. See for example, EP 0154316 by Nishimura et al. and EP 0401384 by Ishikawa et al.

[0137] Alternatively, under some circumstances it may be desirable to decrease the half-life of an antibody of the present invention, rather than increase it. Modifications such as I253A (Hornick et al. (2000) *J. Nucl. Med.* 41:355) and H435A/R, I253A or H310A (Kim et al. (2000) *Eur. J. Immunol.* 29:2819) in Fc of human IgG1 can decrease FcRn binding, thus decreasing half-life (increasing clearance) for use in situations where rapid clearance is preferred, such as medical imaging. See also Kenanova et al. (2005) *Cancer Res.* 65:622. Other means to enhance clearance include formatting the antigen binding domains of the present invention as antibody fragments lacking the ability to bind FcRn, such as Fab fragments. Such modification can reduce the circulating half-life of an antibody from a couple of weeks to a matter of hours. Selective PEGylation of antibody fragments can then be used to fine-tune (increase) the half-life of the antibody fragments if necessary. Chapman et al. (1999) *Nat. Biotechnol.* 17:780. Antibody fragments may also be fused to human serum albumin, e.g. in a fusion protein construct, to increase half-life. Yeh et al. (1992) *Proc. Nat'l Acad. Sci.* 89:1904. Alternatively, a bispecific antibody may be constructed with a first antigen binding domain of the present invention and a second antigen binding domain that binds to human serum albumin (HSA). See Int'l Pat. Appl. Pub. WO 2009/127691 and patent references cited therein. Alternatively, specialized polypeptide sequences can be added to antibody fragments to increase half-life, e.g. “XTEN” polypeptide sequences. Schellenberger et al. (2009) *Nat. Biotechnol.* 27:1186; Int'l Pat. Appl. Pub. WO 2010/091122.

Additional Fc Variants

[0138] When using an IgG4 constant domain, it is usually preferable to include the substitution S228P, which mimics the hinge sequence in IgG1 and thereby stabilizes IgG4

molecules, e.g. reducing Fab-arm exchange between the therapeutic antibody and endogenous IgG4 in the patient being treated. Labrijn et al. (2009) *Nat. Biotechnol.* 27:767; Reddy et al. (2000) *J. Immunol.* 164:1925.

[0139] A potential protease cleavage site in the hinge of IgG1 constructs can be eliminated by D221G and K222S modifications, increasing the stability of the antibody. WO 2014/043344.

[0140] The affinities and binding properties of an Fc variant for its ligands (Fc receptors) may be determined by a variety of in vitro assay methods (biochemical or immunological based assays) known in the art including but not limited to, equilibrium methods (e.g., enzyme-linked immunosorbent assay (ELISA), or radioimmunoassay (RIA)), or kinetics (e.g., BIACORE® SPR analysis), and other methods such as indirect binding assays, competitive inhibition assays, fluorescence resonance energy transfer (FRET), gel electrophoresis and chromatography (e.g., gel filtration). These and other methods may utilize a label on one or more of the components being examined and/or employ a variety of detection methods including but not limited to chromogenic, fluorescent, luminescent, or isotopic labels. A detailed description of binding affinities and kinetics can be found in Paul, W. E., ed., *Fundamental Immunology*, 4th Ed., Lippincott-Raven, Philadelphia (1999), which focuses on antibody-immunogen interactions.

[0141] In still other embodiments, the glycosylation of an antibody is modified to increase or decrease effector function. For example, an aglycosylated antibody can be made that lacks all effector function by mutating the conserved asparagine residue at position 297 (e.g. N297A), thus abolishing complement and FcγRI binding. Bolt et al. (1993) *Eur. J. Immunol.* 23:403. See also Tao & Morrison (1989) *J. Immunol.* 143:2595 (using N297Q in IgG1 to eliminate glycosylation at position 297).

[0142] Although aglycosylated antibodies generally lack effector function, mutations can be introduced to restore that function. Aglycosylated antibodies, e.g. those resulting from N297A/C/D or H mutations or produced in systems (e.g. *E. coli*) that do not glycosylate proteins, can be further mutated to restore FcγR binding, e.g. S298G and/or T299A/G or H (WO 2009/079242), or E382V and M428I (Jung et al. (2010) *Proc. Nat'l Acad. Sci. (USA)* 107:604).

[0143] Glycoengineering can also be used to modify the anti-inflammatory properties of an IgG construct by changing the α2,6 sialyl content of the carbohydrate chains attached at Asn297 of the Fc regions, wherein an increased proportion of α2,6 sialylated forms results in enhanced anti-inflammatory effects. See Nimmerjahn et al. (2008) *Ann. Rev. Immunol.* 26:513. Conversely, reduction in the proportion of antibodies having α2,6 sialylated carbohydrates may be useful in cases where anti-inflammatory properties are not wanted. Methods of modifying α2,6 sialylation content of antibodies, for example by selective purification of α2,6 sialylated forms or by enzymatic modification, are provided at U.S. Pat. Appl. Pub. No. 2008/0206246. In other embodiments, the amino acid sequence of the Fc region may be modified to mimic the effect of α2,6 sialylation, for example by inclusion of an F241A modification. WO 2013/095966.

III. Antibody Physical Properties

[0144] Antibodies described herein can contain one or more glycosylation sites in either the light or heavy chain

variable region. Such glycosylation sites may result in increased immunogenicity of the antibody or an alteration of the pK of the antibody due to altered antigen binding (Marshall et al. (1972) *Ann. Rev. Biochem.* 41:673-702; Gala and Morrison (2004) *J. Immunol.* 172:5489-94; Wallick et al. (1988) *J. Exp. Med.* 168:1099-109; Spiro (2002) *Glycobiology* 12:43R-56R; Parekh et al. (1985) *Nature* 316:452-7; Mimura et al. (2000) *Mol Immunol.* 37:697-706). Glycosylation has been known to occur at motifs containing an N—X-S/T sequence. In some instances, it is preferred to have an anti-huCD40 antibody that does not contain variable region glycosylation. This can be achieved either by selecting antibodies that do not contain the glycosylation motif in the variable region or by mutating residues within the glycosylation region.

[0145] In certain embodiments, the antibodies described herein do not contain asparagine isomerism sites. The deamidation of asparagine may occur on N-G or D-G sequences and result in the creation of an isoaspartic acid residue that introduces a kink into the polypeptide chain and decreases its stability (isoaspartic acid effect).

[0146] Each antibody will have a unique isoelectric point (pI), which generally falls in the pH range between 6 and 9.5. The pI for an IgG1 antibody typically falls within the pH range of 7-9.5 and the pI for an IgG4 antibody typically falls within the pH range of 6-8. There is speculation that antibodies with a pI outside the normal range may have some unfolding and instability under in vivo conditions. Thus, it is preferred to have an anti-CD40 antibody that contains a pI value that falls in the normal range. This can be achieved either by selecting antibodies with a pI in the normal range or by mutating charged surface residues.

[0147] Each antibody will have a characteristic melting temperature, with a higher melting temperature indicating greater overall stability in vivo (Krishnamurthy R and Manning M C (2002) *Curr. Pharm. Biotechnol.* 3:361-71). Generally, it is preferred that the TM1 (the temperature of initial unfolding) be greater than 60° C., preferably greater than 65° C., even more preferably greater than 70° C. The melting point of an antibody can be measured using differential scanning calorimetry (Chen et al. (2003) *Pharm Res* 20:1952-60; Ghirlando et al. (1999) *Immunol. Lett.* 68:47-52) or circular dichroism (Murray et al. (2002) *J. Chromatogr. Sci.* 40:343-9).

[0148] In a preferred embodiment, antibodies are selected that do not degrade rapidly. Degradation of an antibody can be measured using capillary electrophoresis (CE) and MALDI-MS (Alexander A J and Hughes D E (1995) *Anal Chem.* 67:3626-32).

[0149] In another preferred embodiment, antibodies are selected that have minimal aggregation effects, which can lead to the triggering of an unwanted immune response and/or altered or unfavorable pharmacokinetic properties. Generally, antibodies are acceptable with aggregation of 25% or less, preferably 20% or less, even more preferably 15% or less, even more preferably 10% or less and even more preferably 5% or less. Aggregation can be measured by several techniques, including size-exclusion column (SEC), high performance liquid chromatography (HPLC), and light scattering.

IV. Nucleic Acid Molecules

[0150] Another aspect described herein pertains to nucleic acid molecules that encode the antibodies described herein.

The nucleic acids may be present in whole cells, in a cell lysate, or in a partially purified or substantially pure form. A nucleic acid is “isolated” or “rendered substantially pure” when purified away from other cellular components or other contaminants, e.g., other cellular nucleic acids (e.g., other chromosomal DNA, e.g., the chromosomal DNA that is linked to the isolated DNA in nature) or proteins, by standard techniques, including alkaline/SDS treatment, CsCl banding, column chromatography, restriction enzymes, agarose gel electrophoresis and others well known in the art. See, F. Ausubel, et al., ed. (1987) *Current Protocols in Molecular Biology*, Greene Publishing and Wiley Interscience, New York. A nucleic acid described herein can be, for example, DNA or RNA and may or may not contain intronic sequences. In a certain embodiment, the nucleic acid is a cDNA molecule.

[0151] Nucleic acids described herein can be obtained using standard molecular biology techniques. For antibodies expressed by hybridomas (e.g., hybridomas prepared from transgenic mice carrying human immunoglobulin genes as described further below), cDNAs encoding the light and heavy chains of the antibody made by the hybridoma can be obtained by standard PCR amplification or cDNA cloning techniques. For antibodies obtained from an immunoglobulin gene library (e.g., using phage display techniques), nucleic acid encoding the antibody can be recovered from the library.

[0152] Once DNA fragments encoding VH and VL segments are obtained, these DNA fragments can be further manipulated by standard recombinant DNA techniques, for example to convert the variable region genes to full-length antibody chain genes, to Fab fragment genes or to a scFv gene. In these manipulations, a VL- or VH-encoding DNA fragment is operatively linked to another DNA fragment encoding another protein, such as an antibody constant region or a flexible linker. The term “operatively linked”, as used in this context, is intended to mean that the two DNA fragments are joined such that the amino acid sequences encoded by the two DNA fragments remain in-frame.

[0153] The isolated DNA encoding the VH region can be converted to a full-length heavy chain gene by operatively linking the VH-encoding DNA to another DNA molecule encoding heavy chain constant regions (hinge, CH1, CH2 and/or CH3). The sequences of human heavy chain constant region genes are known in the art (see e.g., Kabat, E. A., et al. (1991) *Sequences of Proteins of Immunological Interest*, Fifth Edition, U.S. Department of Health and Human Services, NIH Publication No. 91-3242) and DNA fragments encompassing these regions can be obtained by standard PCR amplification. The heavy chain constant region can be an IgG1, IgG2, IgG3, IgG4, IgA, IgE, IgM or IgD constant region, for example, an IgG1 region. For a Fab fragment heavy chain gene, the VH-encoding DNA can be operatively linked to another DNA molecule encoding only the heavy chain CH1 constant region.

[0154] The isolated DNA encoding the VL region can be converted to a full-length light chain gene (as well as a Fab light chain gene) by operatively linking the VL-encoding DNA to another DNA molecule encoding the light chain constant region, CL. The sequences of human light chain constant region genes are known in the art (see e.g., Kabat, E. A., et al. (1991) *Sequences of Proteins of Immunological Interest*, Fifth Edition, U.S. Department of Health and Human Services, NIH Publication No. 91-3242) and DNA

fragments encompassing these regions can be obtained by standard PCR amplification. The light chain constant region can be a kappa or lambda constant region.

[0155] To create a scFv gene, the VH- and VL-encoding DNA fragments are operatively linked to another fragment encoding a flexible linker, e.g., encoding the amino acid sequence (Gly4-Ser)₃, such that the VH and VL sequences can be expressed as a contiguous single-chain protein, with the VL and VH regions joined by the flexible linker (see e.g., Bird et al. (1988) *Science* 242:423-426; Huston et al. (1988) *Proc. Natl. Acad. Sci. (USA)* 85:5879-5883; McCafferty et al., (1990) *Nature* 348:552-554).

V. Antibody Generation

[0156] Various antibodies of the present invention, e.g., those that compete with or bind to the same epitope as the anti-human CD40 antibodies disclosed herein, can be produced using a variety of known techniques, such as the standard somatic cell hybridization technique described by Kohler and Milstein, *Nature* 256: 495 (1975). Although somatic cell hybridization procedures are preferred, in principle, other techniques for producing monoclonal antibodies also can be employed, e.g., viral or oncogenic transformation of B lymphocytes, phage display technique using libraries of human antibody genes.

[0157] The preferred animal system for preparing hybridomas is the murine system. Hybridoma production in the mouse is a very well-established procedure. Immunization protocols and techniques for isolation of immunized splenocytes for fusion are known in the art. Fusion partners (e.g., murine myeloma cells) and fusion procedures are also known.

[0158] Chimeric or humanized antibodies described herein can be prepared based on the sequence of a murine monoclonal antibody prepared as described above. DNA encoding the heavy and light chain immunoglobulins can be obtained from the murine hybridoma of interest and engineered to contain non-murine (e.g., human) immunoglobulin sequences using standard molecular biology techniques. For example, to create a chimeric antibody, the murine variable regions can be linked to human constant regions using methods known in the art (see e.g., U.S. Pat. No. 4,816,567 to Cabilly et al.). To create a humanized antibody, the murine CDR regions can be inserted into a human framework using methods known in the art (see e.g., U.S. Pat. No. 5,225,539 to Winter, and U.S. Pat. Nos. 5,530,101; 5,585,089; 5,693,762 and 6,180,370 to Queen et al.).

[0159] In one embodiment, the antibodies described herein are human monoclonal antibodies. Such human monoclonal antibodies directed against human CD40 can be generated using transgenic or transchromosomal mice carrying parts of the human immune system rather than the mouse system. These transgenic and transchromosomal mice include mice referred to herein as HuMAb mice and KM mice, respectively, and are collectively referred to herein as "human Ig mice."

[0160] The HuMAb Mouse® (MEDAREX, Inc.) contains human immunoglobulin gene miniloci that encode unrearranged human heavy (μ and γ) and κ light chain immunoglobulin sequences, together with targeted mutations that inactivate the endogenous μ and κ chain loci (see e.g., Lonberg, et al. (1994) *Nature* 368(6474): 856-859). Accordingly, the mice exhibit reduced expression of mouse IgM or κ , and in response to immunization, the introduced human

heavy and light chain transgenes undergo class switching and somatic mutation to generate high affinity human IgGk monoclonal (Lonberg, N. et al. (1994), supra; reviewed in Lonberg, N. (1994) *Handbook of Experimental Pharmacology* 113:49-101; Lonberg, N. and Huszar, D. (1995) *Intern. Rev. Immunol.* 13: 65-93, and Harding, F. and Lonberg, N. (1995) *Ann. N.Y. Acad. Sci.* 764:536-546). The preparation and use of HuMAb mice, and the genomic modifications carried by such mice, is further described in Taylor, L. et al. (1992) *Nucleic Acids Research* 20:6287-6295; Chen, J. et al. (1993) *International Immunology* 5: 647-656; Tuaillon et al. (1993) *Proc. Natl. Acad. Sci. (USA)* 90:3720-3724; Choi et al. (1993) *Nature Genetics* 4:117-123; Chen, J. et al. (1993) *EMBO J.* 12: 821-830; Tuaillon et al. (1994) *J. Immunol.* 152:2912-2920; Taylor, L. et al. (1994) *International Immunology* 6: 579-591; and Fishwild, D. et al. (1996) *Nature Biotechnology* 14: 845-851, the contents of all of which are hereby specifically incorporated by reference in their entirety. See further, U.S. Pat. Nos. 5,545,806; 5,569,825; 5,625,126; 5,633,425; 5,789,650; 5,877,397; 5,661,016; 5,814,318; 5,874,299; and 5,770,429; all to Lonberg and Kay; U.S. Pat. No. 5,545,807 to Surani et al.; PCT Publication Nos. WO 92/03918, WO 93/12227, WO 94/25585, WO 97/13852, WO 98/24884 and WO 99/45962, all to Lonberg and Kay; and PCT Publication No. WO 01/14424 to Korman et al.

[0161] In certain embodiments, antibodies described herein are raised using a mouse that carries human immunoglobulin sequences on transgenes and transchromosomes, such as a mouse that carries a human heavy chain transgene and a human light chain transchromosome. Such mice, referred to herein as "KM mice", are described in detail in PCT Publication WO 02/43478 to Ishida et al.

[0162] Still further, alternative transgenic animal systems expressing human immunoglobulin genes are available in the art and can be used to raise anti-huCD40 antibodies described herein. For example, an alternative transgenic system referred to as the Xenomouse (ABGENIX, Inc.) can be used; such mice are described in, for example, U.S. Pat. Nos. 5,939,598; 6,075,181; 6,114,598; 6,150,584 and 6,162,963 to Kucherlapati et al.

[0163] Moreover, alternative transchromosomal animal systems expressing human immunoglobulin genes are available in the art and can be used to raise anti-CD40 antibodies described herein. For example, mice carrying both a human heavy chain transchromosome and a human light chain transchromosome, referred to as "TC mice" can be used; such mice are described in Tomizuka et al. (2000) *Proc. Natl. Acad. Sci. (USA)* 97:722-727. Furthermore, cows carrying human heavy and light chain transchromosomes have been described in the art (Kuroiwa et al. (2002) *Nature Biotechnology* 20:889-894) and can be used to raise anti-huCD40 antibodies described herein.

[0164] Additional mouse systems described in the art for raising human antibodies, e.g., human anti-huCD40 antibodies, include (i) the VelocImmune® mouse (REGENERON Pharmaceuticals, Inc.), in which the endogenous mouse heavy and light chain variable regions have been replaced, via homologous recombination, with human heavy and light chain variable regions, operatively linked to the endogenous mouse constant regions, such that chimeric antibodies (human V/mouse C) are raised in the mice, and then subsequently converted to fully human antibodies using standard recombinant DNA techniques; and (ii) the MeMo®

mouse (Merus Biopharmaceuticals, Inc.), in which the mouse contains unrearranged human heavy chain variable regions but a single rearranged human common light chain variable region. Such mice, and use thereof to raise antibodies, are described in, for example, WO 2009/15777, US 2010/0069614, WO 2011/072204, WO 2011/097603, WO 2011/163311, WO 2011/163314, WO 2012/148873, US 2012/0070861 and US 2012/0073004.

[0165] Human monoclonal antibodies described herein can also be prepared using phage display methods for screening libraries of human immunoglobulin genes. Such phage display methods for isolating human antibodies are established in the art. See for example: U.S. Pat. Nos. 5,223,409; 5,403,484; and U.S. Pat. No. 5,571,698 to Ladner et al.; U.S. Pat. Nos. 5,427,908 and 5,580,717 to Dower et al.; U.S. Pat. Nos. 5,969,108 and 6,172,197 to McCafferty et al.; and U.S. Pat. Nos. 5,885,793; 6,521,404; 6,544,731; 6,555,313; 6,582,915 and 6,593,081 to Griffiths et al.

[0166] Human monoclonal antibodies described herein can also be prepared using SCID mice into which human immune cells have been reconstituted such that a human antibody response can be generated upon immunization. Such mice are described in, for example, U.S. Pat. Nos. 5,476,996 and 5,698,767 to Wilson et al.

Immunizations

[0167] To generate fully human antibodies to human CD40, mice or transgenic or transchromosomal mice containing human immunoglobulin genes (e.g., HCo12, HCo7 or KM mice) can be immunized with a purified or enriched preparation of the CD40 antigen and/or cells expressing CD40, as described for other antigens, for example, by Lonberg et al. (1994) *Nature* 368(6474): 856-859; Fishwild et al. (1996) *Nature Biotechnology* 14: 845-851 and WO 98/24884. Alternatively, mice can be immunized with DNA encoding human CD40. Preferably, the mice will be 6-16 weeks of age upon the first infusion. For example, a purified or enriched preparation (5-50 μ g) of the recombinant human CD40 antigen can be used to immunize the mice intraperitoneally. In the event that immunizations using a purified or enriched preparation of the CD40 antigen do not result in antibodies, mice can also be immunized with cells expressing CD40, e.g., a cell line, to promote immune responses.

[0168] Cumulative experience with various antigens has shown that the HuMAb transgenic mice respond best when initially immunized intraperitoneally (IP) or subcutaneously (SC) with antigen in Ribi's adjuvant, followed by every other week IP/SC immunizations (up to a total of 10) with antigen in Ribi's adjuvant. The immune response can be monitored over the course of the immunization protocol with plasma samples being obtained by retroorbital bleeds. The plasma can be screened by ELISA and FACS (as described below), and mice with sufficient titers of anti-CD40 human immunoglobulin can be used for fusions. Mice can be boosted intravenously with antigen 3 days before sacrifice and removal of the spleen and lymph nodes. It is expected that 2-3 fusions for each immunization may need to be performed. Between 6 and 24 mice are typically immunized for each antigen. Usually, HCo7, HCo12, and KM strains are used. In addition, both HCo7 and HCo12 transgene can be bred together into a single mouse having two different human heavy chain transgenes (HCo7/HCo12).

Generation of Hybridomas Producing Monoclonal Antibodies to CD40

[0169] To generate hybridomas producing monoclonal antibodies described herein, splenocytes and/or lymph node cells from immunized mice can be isolated and fused to an appropriate immortalized cell line, such as a mouse myeloma cell line. The resulting hybridomas can be screened for the production of antigen-specific antibodies. For example, single cell suspensions of splenic lymphocytes from immunized mice can be fused to Sp2/0 nonsecreting mouse myeloma cells (ATCC, CRL 1581) with 50% PEG. Cells are plated at approximately 2×10^5 in flat bottom microtiter plate, followed by a two week incubation in selective medium containing 10% fetal Clone Serum, 18% "653" conditioned media, 5% origen (IGEN), 4 mM L-glutamine, 1 mM sodium pyruvate, 5 mM HEPES, 0.055 mM 2-mercaptoethanol, 50 units/ml penicillin, 50 mg/ml streptomycin, 50 mg/ml gentamycin and 1 $\times$ HAT (SIGMA). After approximately two weeks, cells can be cultured in medium in which the HAT is replaced with HT. Individual wells can then be screened by ELISA for human monoclonal IgM and IgG antibodies. Once extensive hybridoma growth occurs, medium can be observed usually after 10-14 days. The antibody secreting hybridomas can be replated, screened again, and if still positive for human IgG, the monoclonal antibodies can be subcloned at least twice by limiting dilution. The stable subclones can then be cultured in vitro to generate small amounts of antibody in tissue culture medium for characterization.

[0170] To purify monoclonal antibodies, selected hybridomas can be grown in two-liter spinner-flasks for monoclonal antibody purification. Supernatants can be filtered and concentrated before affinity chromatography with protein A-SEPHAROSE (PHARMACIA, Piscataway, N.J.). Eluted IgG can be checked by gel electrophoresis and high performance liquid chromatography to ensure purity. The buffer solution can be exchanged into PBS, and the concentration can be determined by OD280 using 1.43 extinction coefficient. The monoclonal antibodies can be aliquoted and stored at -80° C.

VI. Antibody Manufacture

Generation of Transfectomas Producing Monoclonal Antibodies to CD40

[0171] Antibodies of the present invention, including both specific antibodies for which sequences are provided and other, related anti-CD40 antibodies, can be produced in a host cell transfectoma using, for example, a combination of recombinant DNA techniques and gene transfection methods as is well known in the art (Morrison, S. (1985) *Science* 229:1202).

[0172] For example, to express antibodies, or antibody fragments thereof, DNAs encoding partial or full-length light and heavy chains, can be obtained by standard molecular biology techniques (e.g., PCR amplification or cDNA cloning using a hybridoma that expresses the antibody of interest) and the DNAs can be inserted into expression vectors such that the genes are operatively linked to transcriptional and translational control sequences. In this context, the term "operatively linked" is intended to mean that an antibody gene is ligated into a vector such that transcriptional and translational control sequences within the vector

serve their intended function of regulating the transcription and translation of the antibody gene. The expression vector and expression control sequences are chosen to be compatible with the expression host cell used. The antibody light chain gene and the antibody heavy chain gene can be inserted into separate vector or both genes are inserted into the same expression vector. The antibody genes are inserted into the expression vector(s) by standard methods (e.g., ligation of complementary restriction sites on the antibody gene fragment and vector, or blunt end ligation if no restriction sites are present). The light and heavy chain variable regions of the antibodies described herein can be used to create full-length antibody genes of any antibody isotype by inserting them into expression vectors already encoding heavy chain constant and light chain constant regions of the desired isotype such that the V_H segment is operatively linked to the C_H segment(s) within the vector and the V_L segment is operatively linked to the C_L segment within the vector. Additionally or alternatively, the recombinant expression vector can encode a signal peptide that facilitates secretion of the antibody chain from a host cell. The antibody chain gene can be cloned into the vector such that the signal peptide is linked in-frame to the amino terminus of the antibody chain gene. The signal peptide can be an immunoglobulin signal peptide or a heterologous signal peptide (i.e., a signal peptide from a non-immunoglobulin protein).

[0173] In addition to the antibody chain genes, recombinant expression vectors may carry regulatory sequences that control the expression of the antibody chain genes in a host cell. The term “regulatory sequence” is intended to include promoters, enhancers and other expression control elements (e.g., polyadenylation signals) that control the transcription or translation of the antibody chain genes. Such regulatory sequences are described, for example, in Goeddel (Gene Expression Technology. Methods in Enzymology 185, Academic Press, San Diego, Calif. (1990)). It will be appreciated by those skilled in the art that the design of the expression vector, including the selection of regulatory sequences, may depend on such factors as the choice of the host cell to be transformed, the level of expression of protein desired, etc. Preferred regulatory sequences for mammalian host cell expression include viral elements that direct high levels of protein expression in mammalian cells, such as promoters and/or enhancers derived from cytomegalovirus (CMV), Simian Virus 40 (SV40), adenovirus, (e.g., the adenovirus major late promoter (AdMLP) and polyoma. Alternatively, nonviral regulatory sequences may be used, such as the ubiquitin promoter or β -globin promoter. Still further, regulatory elements composed of sequences from different sources, such as the SR α promoter system, which contains sequences from the SV40 early promoter and the long terminal repeat of human T cell leukemia virus type 1 (Takebe, Y. et al. (1988) *Mol. Cell. Biol.* 8:466-472).

[0174] In addition to the antibody chain genes and regulatory sequences, recombinant expression vectors may carry additional sequences, such as sequences that regulate replication of the vector in host cells (e.g., origins of replication) and selectable marker genes. The selectable marker gene facilitates selection of host cells into which the vector has been introduced (see, e.g., U.S. Pat. Nos. 4,399,216, 4,634,665 and 5,179,017, all by Axel et al.). For example, typically the selectable marker gene confers resistance to drugs, such as G418, hygromycin or methotrexate, on a host cell into

which the vector has been introduced. Preferred selectable marker genes include the dihydrofolate reductase (DHFR) gene (for use in dhfr-host cells with methotrexate selection/amplification) and the neo gene (for G418 selection).

[0175] For expression of the light and heavy chains, the expression vector(s) encoding the heavy and light chains is transfected into a host cell by standard techniques. The various forms of the term “transfection” are intended to encompass a wide variety of techniques commonly used for the introduction of exogenous DNA into a prokaryotic or eukaryotic host cell, e.g., electroporation, calcium-phosphate precipitation, DEAE-dextran transfection and the like. Although it is theoretically possible to express the antibodies described herein in either prokaryotic or eukaryotic host cells, expression of antibodies in eukaryotic cells, and most preferably mammalian host cells, is the most preferred because such eukaryotic cells, and in particular mammalian cells, are more likely than prokaryotic cells to assemble and secrete a properly folded and immunologically active antibody. Prokaryotic expression of antibody genes has been reported to be ineffective for production of high yields of active antibody (Boss, M. A. and Wood, C. R. (1985) *Immunology Today* 6:12-13). Antibodies of the present invention can also be produced in glycoengineered strains of the yeast *Pichia pastoris*. Li et al. (2006) *Nat. Biotechnol.* 24:210.

[0176] Preferred mammalian host cells for expressing the recombinant antibodies described herein include Chinese Hamster Ovary (CHO cells) (including dhfr-CHO cells, described in Urlaub and Chasin, (1980) *Proc. Natl. Acad. Sci. (USA)* 77:4216-4220, used with a DHFR selectable marker, e.g., as described in R. J. Kaufman and P. A. Sharp (1982) *Mol. Biol.* 159:601-621), NSO myeloma cells, COS cells and SP2 cells. In particular, for use with NSO myeloma cells, another preferred expression system is the GS gene expression system disclosed in WO 87/04462, WO 89/01036 and EP 338,841. When recombinant expression vectors encoding antibody genes are introduced into mammalian host cells, the antibodies are produced by culturing the host cells for a period of time sufficient to allow for expression of the antibody in the host cells or, more preferably, secretion of the antibody into the culture medium in which the host cells are grown. Antibodies can be recovered from the culture medium using standard protein purification methods.

[0177] The N- and C-termini of antibody polypeptide chains of the present invention may differ from the expected sequence due to commonly observed post-translational modifications. For example, C-terminal lysine residues are often missing from antibody heavy chains. Dick et al. (2008) *Biotechnol. Bioeng.* 100:1132. N-terminal glutamine residues, and to a lesser extent glutamate residues, are frequently converted to pyroglutamate residues on both light and heavy chains of therapeutic antibodies. Dick et al. (2007) *Biotechnol. Bioeng.* 97:544; Liu et al. (2011) *JBC* 286:11211; Liu et al. (2011) *J. Biol. Chem.* 286:11211.

[0178] Amino acid sequences for various agonist anti-huCD40 antibodies of the present invention are provided in the Sequence Listing, which is summarized at Table 8. For the reasons mentioned above, the C-terminal lysine is not included in any of sequences in the Sequence Listing for heavy chains or heavy chain constant domains. However, in an alternative embodiment, each heavy chain for the anti-huCD40 antibodies of the present invention, and/or genetic

construct encoding such antibodies or the heavy or light chains thereof, includes this additional lysine residue at the C-terminus of the heavy chain(s).

VII. Assays

[0179] Antibodies described herein can be tested for binding to CD40 by, for example, standard ELISA. Briefly, microtiter plates are coated with purified CD40 at 1-2 µg/ml in PBS, and then blocked with 5% bovine serum albumin in PBS. Dilutions of antibody (e.g., dilutions of plasma from CD40-immunized mice) are added to each well and incubated for 1-2 hours at 37° C. The plates are washed with PBS/Tween and then incubated with secondary reagent (e.g., for human antibodies, or antibodies otherwise having a human heavy chain constant region, a goat-anti-human IgG Fc-specific polyclonal reagent) conjugated to horseradish peroxidase (HRP) for 1 hour at 37° C. After washing, the plates are developed with ABTS substrate (MOSS Inc, product: ABTS-1000) and analyzed by a spectrophotometer at OD 415-495. Sera from immunized mice are then further screened by flow cytometry for binding to a cell line expressing human CD40, but not to a control cell line that does not express CD40. Briefly, the binding of anti-CD40 antibodies is assessed by incubating CD40 expressing CHO cells with the anti-CD40 antibody at 1:20 dilution. The cells are washed and binding is detected with a PE-labeled anti-human IgG Ab. Flow cytometric analyses are performed using a FACScan flow cytometry (BECTON DICKINSON, San Jose, Calif.). Preferably, mice that develop the highest titers will be used for fusions. Analogous experiments may be performed using anti-mouse detection antibodies if mouse anti-huCD40 antibodies are to be detected.

[0180] An ELISA as described above can be used to screen for antibodies and, thus, hybridomas that produce antibodies that show positive reactivity with the CD40 immunogen. Hybridomas that produce antibodies that bind, preferably with high affinity, to CD40 can then be subcloned and further characterized. One clone from each hybridoma, which retains the reactivity of the parent cells (by ELISA), can then be chosen for making a cell bank, and for antibody purification.

[0181] To purify anti-CD40 antibodies, selected hybridomas can be grown in two-liter spinner-flasks for monoclonal antibody purification. Supernatants can be filtered and concentrated before affinity chromatography with protein A-SEPHAROSE (PHARMACIA, Piscataway, N.J.). Eluted IgG can be checked by gel electrophoresis and high performance liquid chromatography to ensure purity. The buffer solution can be exchanged into PBS, and the concentration can be determined by OD280 using 1.43 extinction coefficient. The monoclonal antibodies can be aliquoted and stored at -80° C.

[0182] To determine if the selected anti-CD40 monoclonal antibodies bind to unique epitopes, each antibody can be biotinylated using commercially available reagents (PIERCE, Rockford, Ill.). Biotinylated MAb binding can be detected with a streptavidin labeled probe. Competition studies using unlabeled monoclonal antibodies and biotinylated monoclonal antibodies can be performed using CD40 coated-ELISA plates as described above.

[0183] To determine the isotype of purified antibodies, isotype ELISAs can be performed using reagents specific for antibodies of a particular isotype. For example, to determine the isotype of a human monoclonal antibody, wells of

microtiter plates can be coated with 1 µg/ml of anti-human immunoglobulin overnight at 4° C. After blocking with 1% BSA, the plates are reacted with 1 µg/ml or less of test monoclonal antibodies or purified isotype controls, at ambient temperature for one to two hours. The wells can then be reacted with either human IgG1 or human IgM-specific alkaline phosphatase-conjugated probes. Plates are developed and analyzed as described above.

[0184] To test the binding of monoclonal antibodies to live cells expressing CD40, flow cytometry can be used. Briefly, cell lines expressing membrane-bound CD40 (grown under standard growth conditions) are mixed with various concentrations of monoclonal antibodies in PBS containing 0.1% BSA at 4° C. for 1 hour. After washing, the cells are reacted with Phycoerythrin (PE)-labeled anti-IgG antibody under the same conditions as the primary antibody staining. The samples can be analyzed by FACScan instrument using light and side scatter properties to gate on single cells and binding of the labeled antibodies is determined. An alternative assay using fluorescence microscopy may be used (in addition to or instead of) the flow cytometry assay. Cells can be stained exactly as described above and examined by fluorescence microscopy. This method allows visualization of individual cells, but may have diminished sensitivity depending on the density of the antigen.

[0185] Anti-huCD40 antibodies can be further tested for reactivity with the CD40 antigen by Western blotting. Briefly, cell extracts from cells expressing CD40 can be prepared and subjected to sodium dodecyl sulfate polyacrylamide gel electrophoresis. After electrophoresis, the separated antigens will be transferred to nitrocellulose membranes, blocked with 20% mouse serum, and probed with the monoclonal antibodies to be tested. IgG binding can be detected using anti-IgG alkaline phosphatase and developed with BCIP/NBT substrate tablets (SIGMA Chem. Co., St. Louis, Mo.).

[0186] Methods for analyzing binding affinity, cross-reactivity, and binding kinetics of various anti-CD40 antibodies include standard assays known in the art, for example, Biolayer Interferometry (BLI) analysis, and BIACORE® surface plasmon resonance (SPR) analysis using a BIA-CORE® 2000 SPR instrument (BIACORE AB, Uppsala, Sweden).

[0187] In one embodiment, an antibody specifically binds to the extracellular region of human CD40. An antibody may specifically bind to a particular domain (e.g., a functional domain) within the extracellular domain of CD40. In certain embodiments, the antibody specifically binds to the extracellular region of human CD40 and the extracellular region of cynomolgus CD40. Preferably, an antibody binds to human CD40 with high affinity.

VIII. Bispecific Molecules

[0188] Antibodies described herein may be used for forming bispecific molecules. An anti-CD40 antibody, or antigen-binding fragments thereof, can be derivatized or linked to another functional molecule, e.g., another peptide or protein (e.g., another antibody or ligand for a receptor) to generate a bispecific molecule that binds to at least two different binding sites or target molecules. The antibody described herein may in fact be derivatized or linked to more than one other functional molecule to generate multispecific molecules that bind to more than two different binding sites and/or target molecules; such multispecific molecules are

also intended to be encompassed by the term “bispecific molecule” as used herein. To create a bispecific molecule described herein, an antibody described herein can be functionally linked (e.g., by chemical coupling, genetic fusion, noncovalent association or otherwise) to one or more other binding molecules, such as another antibody, antibody fragment, peptide or binding mimetic, such that a bispecific molecule results.

[0189] Accordingly, provided herein are bispecific molecules comprising at least one first binding specificity for CD40 and a second binding specificity for a second target epitope. In an embodiment described herein in which the bispecific molecule is multispecific, the molecule can further include a third binding specificity.

[0190] In one embodiment, the bispecific molecules described herein comprise a binding specificity at least one antibody, or an antibody fragment thereof, including, e.g., an Fab, Fab', F(ab')₂, Fv, or a single chain Fv. The antibody may also be a light chain or heavy chain dimer, or any minimal fragment thereof such as a Fv or a single chain construct as described in Ladner et al. U.S. Pat. No. 4,946,778, the contents of which is expressly incorporated by reference.

[0191] While human monoclonal antibodies are preferred, other antibodies that can be employed in the bispecific molecules described herein are murine, chimeric and humanized monoclonal antibodies.

[0192] The bispecific molecules described herein can be prepared by conjugating the constituent binding specificities using methods known in the art. For example, each binding specificity of the bispecific molecule can be generated separately and then conjugated to one another. When the binding specificities are proteins or peptides, a variety of coupling or cross-linking agents can be used for covalent conjugation. Examples of cross-linking agents include protein A, carbodiimide, N-succinimidyl-S-acetyl-thioacetate (SATA), 5,5'-dithiobis(2-nitrobenzoic acid) (DTNB), o-phenylenedimaleimide (oPDM), N-succinimidyl-3-(2-pyridyldithio)propionate (SPDP), and sulfo-succinimidyl 4-(N-maleimidomethyl) cyclohexane-1-carboxylate (sulfo-SMCC) (see e.g., Karpovsky et al. (1984) *J. Exp. Med.* 160:1686; Liu, M A et al. (1985) *Proc. Natl. Acad. Sci. (USA)* 82:8648). Other methods include those described in Paulus (1985) *Behring Ins. Mitt.* No. 78, 118-132; Brennan et al. (1985) *Science* 229:81-83, and Glennie et al. (1987) *J. Immunol.* 139: 2367-2375). Preferred conjugating agents are SATA and sulfo-SMCC, both available from PIERCE Chemical Co. (Rockford, Ill.).

[0193] When the binding specificities are antibodies, they can be conjugated via sulfhydryl bonding of the C-terminus hinge regions of the two heavy chains. In a particularly preferred embodiment, the hinge region is modified to contain an odd number of sulfhydryl residues, preferably one, prior to conjugation.

[0194] Alternatively, both binding specificities can be encoded in the same vector and expressed and assembled in the same host cell. This method is particularly useful where the bispecific molecule is a mAb×mAb, mAb×Fab, Fab×F(ab')₂ or ligand x Fab fusion protein. A bispecific molecule described herein can be a single chain molecule comprising one single chain antibody and a binding determinant, or a single chain bispecific molecule comprising two binding determinants. Bispecific molecules may comprise at least two single chain molecules. Methods for preparing bispe-

cific molecules are described for example in U.S. Pat. Nos. 5,260,203; 5,455,030; 4,881,175; 5,132,405; 5,091,513; 5,476,786; 5,013,653; 5,258,498; and 5,482,858.

[0195] Binding of the bispecific molecules to their specific targets can be confirmed using art-recognized methods, such as enzyme-linked immunosorbent assay (ELISA), radioimmunoassay (RIA), FACS analysis, bioassay (e.g., growth inhibition), or Western Blot assay. Each of these assays generally detects the presence of protein-antibody complexes of particular interest by employing a labeled reagent (e.g., an antibody) specific for the complex of interest.

IX. Compositions

[0196] Further provided are compositions, e.g., a pharmaceutical compositions, containing one or more anti-CD40 antibodies, or antigen-binding fragment(s) thereof, as described herein, formulated together with a pharmaceutically acceptable carrier. Such compositions may include one or a combination of (e.g., two or more different) antibodies, or immunoconjugates or bispecific molecules described herein. For example, a pharmaceutical composition described herein can comprise a combination of antibodies (or immunoconjugates or bispecifics) that bind to different epitopes on the target antigen or that have complementary activities.

[0197] In certain embodiments, a composition comprises an anti-CD40 antibody at a concentration of at least 1 mg/ml, 5 mg/ml, 10 mg/ml, 50 mg/ml, 100 mg/ml, 150 mg/ml, 200 mg/ml, or at 1-300 mg/ml or 100-300 mg/ml.

[0198] Pharmaceutical compositions described herein also can be administered in combination therapy, i.e., combined with other agents. For example, the combination therapy can include an anti-CD40 antibody described herein combined with at least one other anti-cancer and/or T-cell stimulating (e.g., activating) agent. Examples of therapeutic agents that can be used in combination therapy are described in greater detail below in the section on uses of the antibodies described herein.

[0199] In some embodiments, therapeutic compositions disclosed herein can include other compounds, drugs, and/or agents used for the treatment of cancer. Such compounds, drugs, and/or agents can include, for example, chemotherapy drugs, small molecule drugs or antibodies that stimulate the immune response to a given cancer. In some instances, therapeutic compositions can include, for example, one or more of an anti-CTLA-4 antibody, an anti-PD-1 antibody, an anti-PD-L1 antibody, an anti-TIGIT antibody, an anti-OX40 (also known as CD134, TNFRSF4, ACT35 and/or TXGP1L) antibody, an anti-LAG-3 antibody, an anti-CD73 antibody, an anti-CD137 antibody, an anti-CD27 antibody, an anti-CSF-1R antibody, a TLR agonist, or a small molecule antagonist of IDO or TGFβ.

[0200] As used herein, “pharmaceutically acceptable carrier” includes any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents, and the like that are physiologically compatible. Preferably, the carrier is suitable for intravenous, intramuscular, subcutaneous, parenteral, spinal or epidermal administration (e.g., by injection or infusion). Depending on the route of administration, the active compound, i.e., antibody, immunoconjugate, or bispecific molecule, may be coated in a material to protect the compound from the action of acids and other natural conditions that may inactivate the compound.

[0201] The pharmaceutical compounds described herein may include one or more pharmaceutically acceptable salts. A "pharmaceutically acceptable salt" refers to a salt that retains the desired biological activity of the parent compound and does not impart any undesired toxicological effects (see e.g., Berge, S. M., et al. (1977) *J. Pharm. Sci.* 66:1-19). Examples of such salts include acid addition salts and base addition salts. Acid addition salts include those derived from nontoxic inorganic acids, such as hydrochloric, nitric, phosphoric, sulfuric, hydrobromic, hydroiodic, phosphorous and the like, as well as from nontoxic organic acids such as aliphatic mono- and dicarboxylic acids, phenyl-substituted alkanolic acids, hydroxy alkanolic acids, aromatic acids, aliphatic and aromatic sulfonic acids and the like. Base addition salts include those derived from alkaline earth metals, such as sodium, potassium, magnesium, calcium and the like, as well as from nontoxic organic amines, such as N,N'-dibenzylethylenediamine, N-methylglucamine, chlorprocaine, choline, diethanolamine, ethylenediamine, procaine and the like.

[0202] A pharmaceutical composition described herein also may include a pharmaceutically acceptable anti-oxidant. Examples of pharmaceutically acceptable antioxidants include: (1) water soluble antioxidants, such as ascorbic acid, cysteine hydrochloride, sodium bisulfate, sodium metabisulfite, sodium sulfite and the like; (2) oil-soluble antioxidants, such as ascorbyl palmitate, butylated hydroxyanisole (BHA), butylated hydroxytoluene (BHT), lecithin, propyl gallate, alpha-tocopherol, and the like; and (3) metal chelating agents, such as citric acid, ethylenediamine tetraacetic acid (EDTA), sorbitol, tartaric acid, phosphoric acid, and the like.

[0203] Examples of suitable aqueous and nonaqueous carriers that may be employed in the pharmaceutical compositions described herein include water, ethanol, polyols (such as glycerol, propylene glycol, polyethylene glycol, and the like), and suitable mixtures thereof, vegetable oils, such as olive oil, and injectable organic esters, such as ethyl oleate. Proper fluidity can be maintained, for example, by the use of coating materials, such as lecithin, by the maintenance of the required particle size in the case of dispersions, and by the use of surfactants.

[0204] These compositions may also contain adjuvants such as preservatives, wetting agents, emulsifying agents and dispersing agents. Prevention of presence of microorganisms may be ensured both by sterilization procedures, supra, and by the inclusion of various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol sorbic acid, and the like. It may also be desirable to include isotonic agents, such as sugars, sodium chloride, and the like into the compositions. In addition, prolonged absorption of the injectable pharmaceutical form may be brought about by the inclusion of agents that delay absorption such as aluminum monostearate and gelatin.

[0205] Pharmaceutically acceptable carriers include sterile aqueous solutions or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersion. The use of such media and agents for pharmaceutically active substances is known in the art. Except insofar as any conventional media or agent is incompatible with the active compound, use thereof in the pharmaceutical compositions described herein is contemplated. Supplementary active compounds can also be incorporated into the compositions.

[0206] Therapeutic compositions typically must be sterile and stable under the conditions of manufacture and storage. The composition can be formulated as a solution, micro-emulsion, liposome, or other ordered structure suitable to high drug concentration. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (for example, glycerol, propylene glycol, and liquid polyethylene glycol, and the like), and suitable mixtures thereof. The proper fluidity can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of dispersion and by the use of surfactants. In many cases, it will be preferable to include isotonic agents, for example, sugars, polyalcohols such as mannitol, sorbitol, or sodium chloride in the composition. Prolonged absorption of the injectable compositions can be brought about by including in the composition an agent that delays absorption, for example, monostearate salts and gelatin.

[0207] Sterile injectable solutions can be prepared by incorporating the active compound in the required amount in an appropriate solvent with one or a combination of ingredients enumerated above, as required, followed by sterilization microfiltration. Generally, dispersions are prepared by incorporating the active compound into a sterile vehicle that contains a basic dispersion medium and the required other ingredients from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum drying and freeze-drying (lyophilization) that yield a powder of the active ingredient plus any additional desired ingredient from a previously sterile-filtered solution thereof.

[0208] The amount of active ingredient that can be combined with a carrier material to produce a single dosage form will vary depending upon the subject being treated, and the particular mode of administration. The amount of active ingredient that can be combined with a carrier material to produce a single dosage form will generally be that amount of the composition that produces a therapeutic effect. Generally, out of one hundred percent, this amount will range from about 0.01 percent to about ninety-nine percent of active ingredient, preferably from about 0.1 percent to about 70 percent, most preferably from about 1 percent to about 30 percent of active ingredient in combination with a pharmaceutically acceptable carrier.

[0209] Dosage regimens are adjusted to provide the optimum desired response (e.g., a therapeutic response). For example, a single bolus may be administered, several divided doses may be administered over time or the dose may be proportionally reduced or increased as indicated by the exigencies of the therapeutic situation. It is especially advantageous to formulate parenteral compositions in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form as used herein refers to physically discrete units suited as unitary dosages for the subjects to be treated; each unit contains a predetermined quantity of active compound calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier. The specification for the dosage unit forms described herein are dictated by and directly dependent on (a) the unique characteristics of the active compound and the particular therapeutic effect to be achieved, and (b) the limitations inherent in the art of compounding such an active compound for the treatment of sensitivity in individuals.

[0210] For administration of the antibody, the dosage ranges from about 0.0001 to 100 mg/kg, and more usually 0.01 to 5 mg/kg, of the host body weight. For example dosages can be 0.3 mg/kg body weight, 1 mg/kg body weight, 3 mg/kg body weight, 5 mg/kg body weight or 10 mg/kg body weight or within the range of 1-10 mg/kg. An exemplary treatment regime entails administration once per week, once every two weeks, once every three weeks, once every four weeks, once a month, once every 3 months or once every three to 6 months.

[0211] In some methods, two or more monoclonal antibodies with different binding specificities are administered simultaneously, in which case the dosage of each antibody administered falls within the ranges indicated. A therapeutic antibody is usually administered on multiple occasions. Intervals between single dosages can be, for example, weekly, monthly, every three months or yearly. Intervals can also be irregular as indicated by measuring blood levels of antibody to the target antigen in the patient. In some methods, dosage is adjusted to achieve a plasma antibody concentration of about 1-1000 $\mu\text{g/ml}$ and in some methods about 25-300 $\mu\text{g/ml}$.

[0212] An antibody can be administered as a sustained release formulation, in which case less frequent administration is required. Dosage and frequency vary depending on the half-life of the antibody in the patient. In general, human antibodies show the longest half-life, followed by humanized antibodies, chimeric antibodies, and nonhuman antibodies. The dosage and frequency of administration can vary depending on whether the treatment is prophylactic or therapeutic. In prophylactic applications, a relatively low dosage is administered at relatively infrequent intervals over a long period of time. Some patients continue to receive treatment for the rest of their lives. In therapeutic applications, a relatively high dosage at relatively short intervals is sometimes required until progression of the disease is reduced or terminated, and preferably until the patient shows partial or complete amelioration of symptoms of disease. Thereafter, the patient can optionally be administered a prophylactic regime, although in many immune-oncology indications continued treatment is not necessary.

[0213] Actual dosage levels of the active ingredients in the pharmaceutical compositions described herein may be varied so as to obtain an amount of the active ingredient that is effective to achieve the desired therapeutic response for a particular patient, composition, and mode of administration, without being toxic to the patient. The selected dosage level will depend upon a variety of pharmacokinetic factors including the activity of the particular compositions described herein employed, or the ester, salt or amide thereof, the route of administration, the time of administration, the rate of excretion of the particular compound being employed, the duration of the treatment, other drugs, compounds and/or materials used in combination with the particular compositions employed, the age, sex, weight, condition, general health and prior medical history of the patient being treated, and like factors well known in the medical arts.

[0214] A "therapeutically effective dosage" of an anti-CD40 antibody described herein preferably results in a decrease in severity of disease symptoms, an increase in frequency and duration of disease symptom-free periods, or a prevention of impairment or disability due to the disease affliction. In the context of cancer, a therapeutically effective

dose preferably prevents further deterioration of physical symptoms associated with cancer. Symptoms of cancer are well-known in the art and include, for example, unusual mole features, a change in the appearance of a mole, including asymmetry, border, color and/or diameter, a newly pigmented skin area, an abnormal mole, darkened area under nail, breast lumps, nipple changes, breast cysts, breast pain, death, weight loss, weakness, excessive fatigue, difficulty eating, loss of appetite, chronic cough, worsening breathlessness, coughing up blood, blood in the urine, blood in stool, nausea, vomiting, liver metastases, lung metastases, bone metastases, abdominal fullness, bloating, fluid in peritoneal cavity, vaginal bleeding, constipation, abdominal distension, perforation of colon, acute peritonitis (infection, fever, pain), pain, vomiting blood, heavy sweating, fever, high blood pressure, anemia, diarrhea, jaundice, dizziness, chills, muscle spasms, colon metastases, lung metastases, bladder metastases, liver metastases, bone metastases, kidney metastases, and pancreatic metastases, difficulty swallowing, and the like. Therapeutic efficacy may be observable immediately after the first administration of an agonistic anti-huCD40 mAb of the present invention, or it may only be observed after a period of time and/or a series of doses. Such delayed efficacy may only be observed after several months of treatment, up to 6, 9 or 12 months. It is critical not to decide prematurely that an agonistic anti-huCD40 mAb of the present invention lacks therapeutically efficacy in light of the delayed efficacy exhibited by some immune-oncology agents.

[0215] A therapeutically effective dose may prevent or delay onset of cancer, such as may be desired when early or preliminary signs of the disease are present. Laboratory tests utilized in the diagnosis of cancer include chemistries (including the measurement of soluble CD40 or CD40L levels) (Hock et al. (2006) *Cancer* 106:2148; Chung & Lim (2014) *J. Trans. Med.* 12:102), hematology, serology and radiology. Accordingly, any clinical or biochemical assay that monitors any of the foregoing may be used to determine whether a particular treatment is a therapeutically effective dose for treating cancer. One of ordinary skill in the art would be able to determine such amounts based on such factors as the subject's size, the severity of the subject's symptoms, and the particular composition or route of administration selected.

[0216] A composition described herein can be administered via one or more routes of administration using one or more of a variety of methods known in the art. As will be appreciated by the skilled artisan, the route and/or mode of administration will vary depending upon the desired results. Preferred routes of administration for antibodies described herein include intravenous, intramuscular, intradermal, intraperitoneal, subcutaneous, spinal or other parenteral routes of administration, for example by injection or infusion. The phrase "parenteral administration" as used herein means modes of administration other than enteral and topical administration, usually by injection, and includes, without limitation, intravenous, intramuscular, intraarterial, intrathecal, intracapsular, intraorbital, intracardiac, intradermal, intraperitoneal, transtracheal, subcutaneous, subcuticular, intraarticular, subcapsular, subarachnoid, intraspinal, epidural and intrasternal injection and infusion.

[0217] Alternatively, an antibody described herein can be administered via a non-parenteral route, such as a topical,

epidermal or mucosal route of administration, for example, intranasally, orally, vaginally, rectally, sublingually or topically.

[0218] The active compounds can be prepared with carriers that will protect the compound against rapid release, such as a controlled release formulation, including implants, transdermal patches, and microencapsulated delivery systems. Biodegradable, biocompatible polymers can be used, such as ethylene vinyl acetate, polyanhydrides, polyglycolic acid, collagen, polyorthoesters, and polylactic acid. Many methods for the preparation of such formulations are patented or generally known to those skilled in the art. See, e.g., *Sustained and Controlled Release Drug Delivery Systems*, J. R. Robinson, ed., Marcel Dekker, Inc., New York, 1978.

[0219] Therapeutic compositions can be administered with medical devices known in the art. For example, in a preferred embodiment, a therapeutic composition described herein can be administered with a needleless hypodermic injection device, such as the devices disclosed in U.S. Pat. Nos. 5,399,163; 5,383,851; 5,312,335; 5,064,413; 4,941,880; 4,790,824; or 4,596,556. Examples of well-known implants and modules for use with anti-huCD40 antibodies described herein include: U.S. Pat. No. 4,487,603, which discloses an implantable micro-infusion pump for dispensing medication at a controlled rate; U.S. Pat. No. 4,486,194, which discloses a therapeutic device for administering medicaments through the skin; U.S. Pat. No. 4,447,233, which discloses a medication infusion pump for delivering medication at a precise infusion rate; U.S. Pat. No. 4,447,224, which discloses a variable flow implantable infusion apparatus for continuous drug delivery; U.S. Pat. No. 4,439,196, which discloses an osmotic drug delivery system having multi-chamber compartments; and U.S. Pat. No. 4,475,196, which discloses an osmotic drug delivery system. These patents are incorporated herein by reference. Many other such implants, delivery systems, and modules are known to those skilled in the art.

[0220] In certain embodiments, the anti-huCD40 antibodies described herein can be formulated to ensure proper distribution in vivo. For example, the blood-brain barrier (BBB) excludes many highly hydrophilic compounds. To ensure that the therapeutic compounds described herein cross the BBB (if desired), they can be formulated, for example, in liposomes. For methods of manufacturing liposomes, see, e.g., U.S. Pat. Nos. 4,522,811; 5,374,548; and 5,399,331. The liposomes may comprise one or more moieties that are selectively transported into specific cells or organs, thus enhance targeted drug delivery (see, e.g., V. V. Ranade (1989) *J. Clin. Pharmacol.* 29:685). Exemplary targeting moieties include folate or biotin (see, e.g., U.S. Pat. No. 5,416,016 to Low et al.); mannositides (Umezawa et al., (1988) *Biochem. Biophys. Res. Commun.* 153:1038); antibodies (P. G. Bloeman et al. (1995) *FEBS Lett.* 357:140; M. Owais et al. (1995) *Antimicrob. Agents Chemother.* 39:180); surfactant protein A receptor (Briscoe et al. (1995) *Am. J. Physiol.* 1233:134); p 120 (Schreier et al. (1994) *J. Biol. Chem.* 269:9090); see also K. Keinänen; M. L. Laukkanen (1994) *FEBS Lett.* 346:123; J. J. Killian; I. J. Fidler (1994) *Immunomethods* 4:273.

X. Uses and Methods

[0221] The antibodies, antibody compositions and methods described herein have numerous in vitro and in vivo utilities involving, for example, enhancement of immune

response by agonizing CD40 signaling. In a preferred embodiment, the antibodies described herein are human or humanized antibodies. For example, anti-huCD40 antibodies described herein can be administered to cells in culture, in vitro or ex vivo, or to human subjects, e.g., in vivo, to enhance immunity in a variety of diseases. Accordingly, provided herein are methods of modifying an immune response in a subject comprising administering to the subject an antibody, or antigen-binding fragment thereof, described herein such that the immune response in the subject is enhanced, stimulated or up-regulated.

[0222] Preferred subjects include human patients in whom enhancement of an immune response would be desirable. The methods are particularly suitable for treating human patients having a disorder that can be treated by augmenting an immune response (e.g., the T-cell mediated immune response). In a particular embodiment, the methods are particularly suitable for treatment of cancer in vivo. To achieve antigen-specific enhancement of immunity, anti-huCD40 antibodies described herein can be administered together with an antigen of interest or the antigen may already be present in the subject to be treated (e.g., a tumor-bearing or virus-bearing subject). When antibodies to CD40 are administered together with another agent, the two can be administered separately or simultaneously.

[0223] Also encompassed are methods for detecting the presence of human CD40 antigen in a sample, or measuring the amount of human CD40 antigen, comprising contacting the sample, and a control sample, with a human monoclonal antibody, or an antigen binding fragment thereof, that specifically binds to human CD40, under conditions that allow for formation of a complex between the antibody or fragment thereof and human CD40. The formation of a complex is then detected, wherein a difference complex formation between the sample compared to the control sample is indicative the presence of human CD40 antigen in the sample. Moreover, the anti-CD40 antibodies described herein can be used to purify human CD40 via immunoaffinity purification.

[0224] Given the ability of anti-huCD40 antibodies described herein to enhance co-stimulation of T cell responses, e.g., antigen-specific T cell responses, provided herein are in vitro and in vivo methods of using the antibodies described herein to stimulate, enhance or upregulate antigen-specific T cell responses, e.g., anti-tumor T cell responses.

[0225] CD4+ and CD8+ T cell responses can be enhanced using anti-CD40 antibodies. The T cells can be Teff cells, e.g., CD4+ Teff cells, CD8+ Teff cells, T helper (Th) cells and T cytotoxic (Tc) cells.

[0226] Further encompassed are methods of enhancing an immune response (e.g., an antigen-specific T cell response) in a subject comprising administering an anti-huCD40 antibody described herein to the subject such that an immune response (e.g., an antigen-specific T cell response) in the subject is enhanced. In a preferred embodiment, the subject is a tumor-bearing subject and an immune response against the tumor is enhanced. A tumor may be a solid tumor or a liquid tumor, e.g., a hematological malignancy. In certain embodiments, a tumor is an immunogenic tumor. In certain embodiments, a tumor is non-immunogenic. In certain embodiments, a tumor is PD-L1 positive. In certain embodi-

ments a tumor is PD-L1 negative. A subject may also be a virus-bearing subject and an immune response against the virus is enhanced.

[0227] Further provided are methods for inhibiting growth of tumor cells in a subject comprising administering to the subject an anti-huCD40 antibody described herein such that growth of the tumor is inhibited in the subject. Also provided are methods of treating chronic viral infection in a subject comprising administering to the subject an anti-huCD40 antibody described herein such that the chronic viral infection is treated in the subject.

[0228] In certain embodiments, an anti-huCD40 antibody is given to a subject as an adjunctive therapy. Treatments of subjects having cancer with an anti-huCD40 antibody may lead to a long-term durable response relative to the current standard of care; long term survival of at least 1, 2, 3, 4, 5, 10 or more years, recurrence free survival of at least 1, 2, 3, 4, 5, or 10 or more years. In certain embodiments, treatment of a subject having cancer with an anti-huCD40 antibody prevents recurrence of cancer or delays recurrence of cancer by, e.g., 1, 2, 3, 4, 5, or 10 or more years. An anti-CD40 treatment can be used as a primary or secondary line of treatment.

[0229] These and other methods described herein are discussed in further detail below.

Cancer

[0230] Provided herein are methods for treating a subject having cancer, comprising administering to the subject an anti-huCD40 antibody described herein, such that the subject is treated, e.g., such that growth of cancerous tumors is inhibited or reduced and/or that the tumors regress. An anti-huCD40 antibody can be used alone to inhibit the growth of cancerous tumors. Alternatively, an anti-huCD40 antibody can be used in conjunction with another agent, e.g., other immunogenic agents, standard cancer treatments, or other antibodies, as described below. Combination with an inhibitor of PD-1, such as an anti-PD-1 or anti-PD-L1 antibody, is also provided. See, e.g., Ellmark et al. (2015) *Oncimmunology* 4:7 e1011484.

[0231] Accordingly, provided herein are methods of treating cancer, e.g., by inhibiting growth of tumor cells, in a subject, comprising administering to the subject a therapeutically effective amount of an anti-huCD40 antibody described herein, e.g., a humanized form of 12D6, 5F11, 8E8, 5G7 or 19G3, or antigen-binding fragment thereof. The antibody may be a humanized anti-huCD40 antibody (such as any of the humanized anti-huCD40 antibodies described herein), a human chimeric anti-huCD40 antibody, or a humanized non-human anti-huCD40 antibody, e.g., a human, chimeric or humanized anti-huCD40 antibody that competes for binding with, or binds to the same epitope as, at least one of the anti-huCD40 antibodies specifically described herein.

[0232] Cancers whose growth may be inhibited using the antibodies of the invention include cancers typically responsive to immunotherapy. Non-limiting examples of cancers for treatment include squamous cell carcinoma, small-cell lung cancer, non-small cell lung cancer, squamous non-small cell lung cancer (NSCLC), non NSCLC, glioma, gastrointestinal cancer, renal cancer (e.g. clear cell carcinoma), ovarian cancer, liver cancer, colorectal cancer, endometrial cancer, kidney cancer (e.g., renal cell carcinoma (RCC)), prostate cancer (e.g. hormone refractory prostate

adenocarcinoma), thyroid cancer, neuroblastoma, pancreatic cancer, glioblastoma (glioblastoma multiforme), cervical cancer, stomach cancer, bladder cancer, hepatoma, breast cancer, colon carcinoma, and head and neck cancer (or carcinoma), gastric cancer, germ cell tumor, pediatric sarcoma, sinonasal natural killer, melanoma (e.g., metastatic malignant melanoma, such as cutaneous or intraocular malignant melanoma), bone cancer, skin cancer, uterine cancer, cancer of the anal region, testicular cancer, carcinoma of the fallopian tubes, carcinoma of the endometrium, carcinoma of the cervix, carcinoma of the vagina, carcinoma of the vulva, cancer of the esophagus, cancer of the small intestine, cancer of the endocrine system, cancer of the parathyroid gland, cancer of the adrenal gland, sarcoma of soft tissue, cancer of the urethra, cancer of the penis, solid tumors of childhood, cancer of the ureter, carcinoma of the renal pelvis, neoplasm of the central nervous system (CNS), primary CNS lymphoma, tumor angiogenesis, spinal axis tumor, brain stem glioma, pituitary adenoma, Kaposi's sarcoma, epidermoid cancer, squamous cell cancer, T-cell lymphoma, environmentally-induced cancers including those induced by asbestos, virus-related cancers (e.g., human papilloma virus (HPV)-related tumor), and hematologic malignancies derived from either of the two major blood cell lineages, i.e., the myeloid cell line (which produces granulocytes, erythrocytes, thrombocytes, macrophages and mast cells) or lymphoid cell line (which produces B, T, NK and plasma cells), such as all types of leukemias, lymphomas, and myelomas, e.g., acute, chronic, lymphocytic and/or myelogenous leukemias, such as acute leukemia (ALL), acute myelogenous leukemia (AML), chronic lymphocytic leukemia (CLL), and chronic myelogenous leukemia (CML), undifferentiated AML (MO), myeloblastic leukemia (M1), myeloblastic leukemia (M2; with cell maturation), promyelocytic leukemia (M3 or M3 variant [M3V]), myelomonocytic leukemia (M4 or M4 variant with eosinophilia [M4E]), monocytic leukemia (M5), erythroleukemia (M6), megakaryoblastic leukemia (M7), isolated granulocytic sarcoma, and chloroma; lymphomas, such as Hodgkin's lymphoma (HL), non-Hodgkin's lymphoma (NHL), B-cell lymphomas, T-cell lymphomas, lymphoplasmacytoid lymphoma, monocytoid B-cell lymphoma, mucosa-associated lymphoid tissue (MALT) lymphoma, anaplastic (e.g., Ki 1+) large-cell lymphoma, adult T-cell lymphoma/leukemia, mantle cell lymphoma, angio immunoblastic T-cell lymphoma, angiocentric lymphoma, intestinal T-cell lymphoma, primary mediastinal B-cell lymphoma, precursor T-lymphoblastic lymphoma, T-lymphoblastic; and lymphoma/leukemia (T-Lbly/T-ALL), peripheral T-cell lymphoma, lymphoblastic lymphoma, post-transplantation lymphoproliferative disorder, true histiocytic lymphoma, primary central nervous system lymphoma, primary effusion lymphoma, lymphoblastic lymphoma (LBL), hematopoietic tumors of lymphoid lineage, acute lymphoblastic leukemia, diffuse large B-cell lymphoma, Burkitt's lymphoma, follicular lymphoma, diffuse histiocytic lymphoma (DHL), immunoblastic large cell lymphoma, precursor B-lymphoblastic lymphoma, cutaneous T-cell lymphoma (CTLC) (also called mycosis fungoides or Sezary syndrome), and lymphoplasmacytoid lymphoma (LPL) with Waldenstrom's macroglobulinemia; myelomas, such as IgG myeloma, light chain myeloma, nonsecretory myeloma, smoldering myeloma (also called indolent myeloma), solitary plasmacytoma, and multiple myelomas, chronic lymphocytic leu-

kemia (CLL), hairy cell lymphoma; hematopoietic tumors of myeloid lineage, tumors of mesenchymal origin, including fibrosarcoma and rhabdomyosarcoma; seminoma, teratocarcinoma, tumors of the central and peripheral nervous, including astrocytoma, schwannomas; tumors of mesenchymal origin, including fibrosarcoma, rhabdomyosarcoma, and osteosarcoma; and other tumors, including melanoma, xeroderma pigmentosum, keratoacanthoma, seminoma, thyroid follicular cancer and teratocarcinoma, hematopoietic tumors of lymphoid lineage, for example T-cell and B-cell tumors, including but not limited to T-cell disorders such as T-prolymphocytic leukemia (T-PLL), including of the small cell and cerebriform cell type; large granular lymphocyte leukemia (LGL) preferably of the T-cell type; a/d T-NHL hepatosplenic lymphoma; peripheral/post-thymic T cell lymphoma (pleomorphic and immunoblastic subtypes); angiocentric (nasal) T-cell lymphoma; cancer of the head or neck, renal cancer, rectal cancer, cancer of the thyroid gland; acute myeloid lymphoma, as well as any combinations of said cancers. The methods described herein may also be used for treatment of metastatic cancers, refractory cancers (e.g., cancers refractory to previous immunotherapy, e.g., with a blocking CTLA-4 or PD-1 antibody), and recurrent cancers.

[0233] Notwithstanding the above, the agonist anti-huCD40 antibodies of the present invention will not find use in treating hematologic cancers with CD40 expression, which might be exacerbated by treatment with a CD40 agonist. Certain cancers may be known to express CD40 and thus be subject to such exacerbation, and thus may be categorically excluded. In other embodiments specific tumor samples are tested for expression of CD40 and are excluded from therapy with the agonist anti-huCD40 antibodies of the present invention based on the test results.

[0234] An anti-huCD40 antibody can be administered as a monotherapy, or as the only immunostimulating therapy, or it can be combined with an immunogenic agent in a cancer vaccine strategy, such as cancerous cells, purified tumor antigens (including recombinant proteins, peptides, and carbohydrate molecules), cells, and cells transfected with genes encoding immune stimulating cytokines (He et al. (2004) *J. Immunol.* 173:4919-28). Non-limiting examples of tumor vaccines that can be used include peptides of melanoma antigens, such as peptides of gp100, MAGE antigens, Trp-2, MART1 and/or tyrosinase, or tumor cells transfected to express the cytokine GM-CSF. Many experimental strategies for vaccination against tumors have been devised (see Rosenberg, S., 2000, Development of Cancer Vaccines, ASCO Educational Book Spring: 60-62; Logothetis, C., 2000, ASCO Educational Book Spring: 300-302; Khayat, D. 2000, ASCO Educational Book Spring: 414-428; Foon, K. 2000, ASCO Educational Book Spring: 730-738; see also Restifo, N. and Sznol, M., Cancer Vaccines, Ch. 61, pp. 3023-3043 in DeVita et al. (eds.), 1997, Cancer: Principles and Practice of Oncology, Fifth Edition). In one of these strategies, a vaccine is prepared using autologous or allogeneic tumor cells. These cellular vaccines have been shown to be most effective when the tumor cells are transduced to express GM-CSF. GM-CSF has been shown to be a potent activator of antigen presentation for tumor vaccination. Dranoff et al. (1993) *Proc. Natl. Acad. Sci. (USA)* 90: 3539-43.

[0235] The study of gene expression and large scale gene expression patterns in various tumors has led to the defini-

tion of so called tumor specific antigens. Rosenberg, S A (1999) *Immunity* 10: 281-7. In many cases, these tumor specific antigens are differentiation antigens expressed in the tumors and in the cell from which the tumor arose, for example melanocyte antigens gp100, MAGE antigens, and Trp-2. More importantly, many of these antigens can be shown to be the targets of tumor specific T cells found in the host. CD40 agonists can be used in conjunction with a collection of recombinant proteins and/or peptides expressed in a tumor in order to generate an immune response to these proteins. These proteins are normally viewed by the immune system as self antigens and are therefore tolerant to them. The tumor antigen can include the protein telomerase, which is required for the synthesis of telomeres of chromosomes and which is expressed in more than 85% of human cancers and in only a limited number of somatic tissues (Kim et al. (1994) *Science* 266: 2011-2013). Tumor antigen can also be "neo-antigens" expressed in cancer cells because of somatic mutations that alter protein sequence or create fusion proteins between two unrelated sequences (i.e., bcr-abl in the Philadelphia chromosome), or idiotype from B cell tumors.

[0236] Other tumor vaccines can include the proteins from viruses implicated in human cancers such as Human Papilloma Viruses (HPV), Hepatitis Viruses (HBV and HCV) and Kaposi's Herpes Sarcoma Virus (KHSV). Another form of tumor specific antigen that can be used in conjunction with CD40 inhibition is purified heat shock proteins (HSP) isolated from the tumor tissue itself. These heat shock proteins contain fragments of proteins from the tumor cells and these HSPs are highly efficient at delivery to antigen presenting cells for eliciting tumor immunity (Suot & Srivastava (1995) *Science* 269:1585-1588; Tamura et al. (1997) *Science* 278: 117-120).

[0237] Dendritic cells (DC) are potent antigen presenting cells that can be used to prime antigen-specific responses. DC's can be produced ex vivo and loaded with various protein and peptide antigens as well as tumor cell extracts (Nestle et al. (1998) *Nature Medicine* 4: 328-332). DCs can also be transduced by genetic means to express these tumor antigens as well. DCs have also been fused directly to tumor cells for the purposes of immunization (Kugler et al. (2000) *Nature Medicine* 6:332-336). As a method of vaccination, DC immunization can be effectively combined with CD40 agonism to activate (unleash) more potent anti-tumor responses.

[0238] Agonism of CD40 can also be combined with standard cancer treatments (e.g., surgery, radiation, and chemotherapy). Agonism of CD40 can be effectively combined with chemotherapeutic regimes. In these instances, it may be possible to reduce the dose of chemotherapeutic reagent administered (Mokyr et al. (1998) *Cancer Research* 58: 5301-5304). An example of such a combination is an anti-huCD40 antibody in combination with decarbazine for the treatment of melanoma. Another example of such a combination is an anti-huCD40 antibody in combination with interleukin-2 (IL-2) for the treatment of melanoma. The scientific rationale behind the combined use of CD40 agonists and chemotherapy is that cell death, that is a consequence of the cytotoxic action of most chemotherapeutic compounds, should result in increased levels of tumor antigen in the antigen presentation pathway. Other combination therapies that may result in synergy with CD40 agonism through cell death are radiation, surgery, and hor-

mone deprivation. Each of these protocols creates a source of tumor antigen in the host. Angiogenesis inhibitors can also be combined with CD40 agonists. Inhibition of angiogenesis leads to tumor cell death which may feed tumor antigen into host antigen presentation pathways.

[0239] The anti-huCD40 antibodies described herein can also be used in combination with bispecific antibodies that target Fc α or Fc γ receptor-expressing effectors cells to tumor cells (see, e.g., U.S. Pat. Nos. 5,922,845 and 5,837,243). Bispecific antibodies can be used to target two separate antigens. For example anti-Fc receptor/anti tumor antigen (e.g., Her-2/neu) bispecific antibodies have been used to target macrophages to sites of tumor. This targeting may more effectively activate tumor specific responses. The T cell arm of these responses would be augmented by agonism of CD40. Alternatively, antigen may be delivered directly to DCs by the use of bispecific antibodies that bind to tumor antigen and a dendritic cell specific cell surface marker.

[0240] Tumors evade host immune surveillance by a large variety of mechanisms. Many of these mechanisms may be overcome by the inactivation of immunosuppressive proteins expressed by the tumors. These include among others TGF- β (Kehrl et al. (1986) *J. Exp. Med.* 163: 1037-1050), IL-10 (Howard & O'Garra (1992) *Immunology Today* 13: 198-200), and Fas ligand (Hahne et al. (1996) *Science* 274: 1363-1365). Antibodies to each of these entities can be used in combination with anti-huCD40 antibodies to counteract the effects of the immunosuppressive agent and favor tumor immune responses by the host.

[0241] Anti-CD40 antibodies are able to substitute effectively for T cell helper activity. Ridge et al. (1998) *Nature* 393: 474-478. Activating antibodies to T cell costimulatory molecules such as CTLA-4 (e.g., U.S. Pat. No. 5,811,097), OX-40 (Weinberg et al. (2000) *Immunol.* 164: 2160-2169), CD137/4-1BB (Melero et al. (1997) *Nature Medicine* 3: 682-685 (1997), and ICOS (Hutloff et al. (1999) *Nature* 397: 262-266) may also provide for increased levels of T cell activation. Inhibitors of PD1 or PD-L1 may also be used in conjunction with anti-huCD40 antibodies.

[0242] There are also several experimental treatment protocols that involve ex vivo activation and expansion of antigen specific T cells and adoptive transfer of these cells into recipients in order to stimulate antigen-specific T cells against tumor (Greenberg & Riddell (1999) *Science* 285: 546-51). These methods can also be used to activate T cell responses to infectious agents such as CMV. Ex vivo activation in the presence of anti-CD40 antibodies can increase the frequency and activity of the adoptively transferred T cells.

Chronic Viral Infections

[0243] In another aspect, the invention described herein provides a method of treating an infectious disease in a subject comprising administering to the subject an anti-huCD40 antibody, or antigen-binding fragment thereof, such that the subject is treated for the infectious disease.

[0244] Similar to its application to tumors as discussed above, antibody-mediated CD40 agonism can be used alone, or as an adjuvant, in combination with vaccines, to enhance the immune response to pathogens, toxins, and self-antigens. Examples of pathogens for which this therapeutic approach can be particularly useful, include pathogens for which there is currently no effective vaccine, or pathogens for which conventional vaccines are less than completely effective.

These include, but are not limited to HIV, Hepatitis (A, B, & C), Influenza, Herpes, Giardia, Malaria, *Leishmania*, *Staphylococcus aureus*, *Pseudomonas aeruginosa*. CD40 agonism is particularly useful against established infections by agents such as HIV that present altered antigens over the course of the infections. These novel epitopes are recognized as foreign at the time of anti-human CD40 antibody administration, thus provoking a strong T cell response.

[0245] Some examples of pathogenic viruses causing infections treatable by methods described herein include HIV, hepatitis (A, B, or C), herpes virus (e.g., VZV, HSV-1, HAV-6, HSV-II, and CMV, Epstein Barr virus), adenovirus, influenza virus, flaviviruses, echovirus, rhinovirus, coxsackie virus, coronavirus, respiratory syncytial virus, mumps virus, rotavirus, measles virus, rubella virus, parvovirus, vaccinia virus, HTLV virus, dengue virus, papillomavirus, molluscum virus, poliovirus, rabies virus, JC virus and arboviral encephalitis virus.

[0246] Some examples of pathogenic bacteria causing infections treatable by methods described herein include *chlamydia*, rickettsial bacteria, mycobacteria, staphylococci, streptococci, pneumonococci, meningococci and gonococci, *klebsiella*, *proteus*, *serratia*, *pseudomonas*, *legionella*, diphtheria, *salmonella*, bacilli, cholera, tetanus, botulism, anthrax, plague, leptospirosis, and Lyme disease bacteria.

[0247] Some examples of pathogenic fungi causing infections treatable by methods described herein include *Candida* (*albicans*, *krusei*, *glabrata*, *tropicalis*, etc.), *Cryptococcus neoformans*, *Aspergillus* (*fumigatus*, *niger*, etc.), Genus *Mucorales* (*mucor*, *absidia*, *rhizopus*), *Sporothrix schenckii*, *Blastomyces dermatitidis*, *Paracoccidioides brasiliensis*, *Coccidioides immitis* and *Histoplasma capsulatum*.

[0248] Some examples of pathogenic parasites causing infections treatable by methods described herein include *Entamoeba histolytica*, *Balantidium coli*, *Naegleria fowleri*, *Acanthamoeba* sp., *Giardia lamblia*, *Cryptosporidium* sp., *Pneumocystis carinii*, *Plasmodium vivax*, *Babesia microti*, *Trypanosoma brucei*, *Trypanosoma cruzi*, *Leishmania donovani*, *Toxoplasma gondii*, *Nippostrongylus brasiliensis*.

[0249] In all of the above methods, CD40 agonism can be combined with other forms of immunotherapy such as cytokine treatment (e.g., interferons, GM-CSF, G-CSF, IL-2), or bispecific antibody therapy, which provides for enhanced presentation of tumor antigens. See, e.g., Holliger (1993) *Proc. Natl. Acad. Sci. (USA)* 90:6444-6448; Poljak (1994) *Structure* 2:1121-1123.

Vaccine Adjuvants

[0250] Anti-huCD40 antibodies described herein can be used to enhance antigen-specific immune responses by co-administration of an anti-huCD40 antibody with an antigen of interest, e.g., a vaccine. Accordingly, provided herein are methods of enhancing an immune response to an antigen in a subject, comprising administering to the subject: (i) the antigen; and (ii) an anti-huCD40 antibody, or antigen-binding fragment thereof, such that an immune response to the antigen in the subject is enhanced. The antigen can be, for example, a tumor antigen, a viral antigen, a bacterial antigen or an antigen from a pathogen. Non-limiting examples of such antigens include those discussed in the sections above, such as the tumor antigens (or tumor vaccines) discussed above, or antigens from the viruses, bacteria or other pathogens described above.

[0251] Suitable routes of administering the antibody compositions (e.g., human monoclonal antibodies, multispecific and bispecific molecules and immunoconjugates) described herein in vivo and in vitro are well known in the art and can be selected by those of ordinary skill. For example, the antibody compositions can be administered by injection (e.g., intravenous or subcutaneous). Suitable dosages of the molecules used will depend on the age and weight of the subject and the concentration and/or formulation of the antibody composition.

[0252] As previously described, anti-huCD40 antibodies described herein can be co-administered with one or other more therapeutic agents, e.g., a cytotoxic agent, a radiotoxic agent or an immunosuppressive agent. The antibody can be linked to the agent (as an immuno-complex) or can be administered separate from the agent. In the latter case (separate administration), the antibody can be administered before, after or concurrently with the agent or can be co-administered with other known therapies, e.g., an anti-cancer therapy, e.g., radiation. Such therapeutic agents include, among others, anti-neoplastic agents such as doxorubicin (adriamycin), cisplatin bleomycin sulfate, carmustine, chlorambucil, dacarbazine and cyclophosphamide hydroxyurea which, by themselves, are only effective at levels which are toxic or subtoxic to a patient. Cisplatin is intravenously administered as a 100 mg/ml dose once every four weeks and adriamycin is intravenously administered as a 60-75 mg/ml dose once every 21 days. Co-administration of anti-CD40 antibodies, or antigen binding fragments thereof, described herein with chemotherapeutic agents provides two anti-cancer agents which operate via different mechanisms which yield a cytotoxic effect to human tumor cells. Such co-administration can solve problems due to development of resistance to drugs or a change in the antigenicity of the tumor cells that would render them unreactive with the antibody.

[0253] Also within the scope described herein are kits comprising the antibody compositions described herein (e.g., human antibodies, bispecific or multispecific molecules, or immunoconjugates) and instructions for use. The kit can further contain at least one additional reagent, or one or more additional human antibodies described herein (e.g., a human antibody having a complementary activity that binds to an epitope in CD40 antigen distinct from the first human antibody). Kits typically include a label indicating the intended use of the contents of the kit. The term label includes any writing, or recorded material supplied on or with the kit, or that otherwise accompanies the kit.

Combination Therapies

[0254] In addition to the combinations therapies provided above, anti-CD40 antibodies described herein can also be used in combination therapy, e.g., for treating cancer, as described below.

[0255] The present invention provides methods of combination therapy in which an anti-huCD40 antibody is co-administered with one or more additional agents, e.g., antibodies, that are effective in stimulating immune responses to thereby further enhance, stimulate or upregulate immune responses in a subject.

[0256] Generally, an anti-huCD40 antibody described herein can be combined with (i) an agonist of another co-stimulatory receptor and/or (ii) an antagonist of an inhibitory signal on T cells, either of which results in amplifying

antigen-specific T cell responses (immune checkpoint regulators). Most of the co-stimulatory and co-inhibitory molecules are members of the immunoglobulin super family (IgSF), and anti-CD40 antibodies described herein may be administered with an agent that targets a member of the IgSF family to increase an immune response. One important family of membrane-bound ligands that bind to co-stimulatory or co-inhibitory receptors is the B7 family, which includes B7-1, B7-2, B7-H1 (PD-L1), B7-DC (PD-L2), B7-H2 (ICOS-L), B7-H3, B7-H4, B7-H5 (VISTA), and B7-H6. Another family of membrane bound ligands that bind to co-stimulatory or co-inhibitory receptors is the TNF family of molecules that bind to cognate TNF receptor family members, which include CD40 and CD40L, OX-40, OX-40L, CD70, CD27L, CD30, CD30L, 4-1BBL, CD137/4-1BB, TRAIL/Apo2-L, TRAILR1/DR4, TRAILR2/DR5, TRAILR3, TRAILR4, OPG, RANK, RANKL, TWEAKR/Fn14, TWEAK, BAFFR, EDAR, XEDAR, TACI, APRIL, BCMA, LT β R, LIGHT, DcR3, HVEM, VEGI/TL1A, TRAMP/DR3, EDAR, EDA1, XEDAR, EDA2, TNFR1, Lymphotoxin α /TNF β , TNFR2, TNF α , LT β R, Lymphotoxin 1 (32, FAS, FASL, RELT, DR6, TROY, NGFR (see, e.g., Tansey (2009) *Drug Discovery Today* 00:1).

[0257] In another aspect, anti-huCD40 antibodies can be used in combination with antagonists of cytokines that inhibit T cell activation (e.g., IL-6, IL-10, TGF- β , VEGF; or other “immunosuppressive cytokines,” or cytokines that stimulate T cell activation, for stimulating an immune response, e.g., for treating proliferative diseases, such as cancer.

[0258] In one aspect, T cell responses can be stimulated by a combination of the anti-huCD40 *mAbs* of the present invention and one or more of (i) an antagonist of a protein that inhibits T cell activation (e.g., immune checkpoint inhibitors) such as CTLA-4, PD-1, PD-L1, PD-L2, LAG-3, TIM-3, Galectin 9, CEACAM-1, BTLA, CD69, Galectin-1, TIGIT, CD113, GPR56, VISTA, 2B4, CD48, GARP, PD1H, LAIR1, TIM-1, and TIM-4, and (ii) an agonist of a protein that stimulates T cell activation such as B7-1, B7-2, CD28, 4-1BB (CD137), 4-1BBL, ICOS, ICOS-L, OX40, OX40L, GITR, GITRL, CD70, CD27, CD40, DR3 and CD28H.

[0259] Exemplary agents that modulate one of the above proteins and may be combined with agonist anti-huCD40 antibodies, e.g., those described herein, for treating cancer, include: YERVOY®/ipilimumab or tremelimumab (to CTLA-4), galiximab (to B7.1), BMS-936558 (to PD-1), pidilizumab/CT-011 (to PD-1), KEYTRUDA®/pembrolizumab/MK-3475 (to PD-1), AMP224 (to B7-DC/PD-L2), BMS-936559 (to B7-H1), MPDL3280A (to B7-H1), MEDI-570 (to ICOS), AMG557 (to B7H2), MGA271 (to B7H3—WO 11/109400), IMP321 (to LAG-3), urelumab/BMS-663513 and PF-05082566 (to CD137/4-1BB), varilumab/CDX-1127 (to CD27), MEDI-6383 and MEDI-6469 (to OX40), RG-7888 (to OX40L—WO 06/029879), Atacicept (to TACI), muromonab-CD3 (to CD3), ipilimumab (to CTLA-4).

[0260] Other molecules that can be combined with agonist anti-huCD40 antibodies for the treatment of cancer include antagonists of inhibitory receptors on NK cells or agonists of activating receptors on NK cells. For example, agonist anti-huCD40 antibodies can be combined with antagonists of MR (e.g., lirilumab).

[0261] Yet other agents for combination therapies include agents that inhibit or deplete macrophages or monocytes,

including but not limited to CSF-1R antagonists such as CSF-1R antagonist antibodies including RG7155 (WO 11/70024, WO 11/107553, WO 11/131407, WO 13/87699, WO 13/119716, WO 13/132044) or FPA-008 (WO 11/140249; WO 13/169264; WO 14/036357).

[0262] Generally, agonist anti-huCD40 antibodies described herein can be used together with one or more of agonistic agents that ligate positive co-stimulatory receptors, blocking agents that attenuate signaling through inhibitory receptors, and one or more agents that increase systemically the frequency of anti-tumor T cells, agents that overcome distinct immune suppressive pathways within the tumor microenvironment (e.g., block inhibitory receptor engagement (e.g., PD-L1/PD-1 interactions), deplete or inhibit Tregs (e.g., using an anti-CD25 monoclonal antibody (e.g., daclizumab) or by ex vivo anti-CD25 bead depletion), inhibit metabolic enzymes such as IDO, or reverse/prevent T cell anergy or exhaustion) and agents that trigger innate immune activation and/or inflammation at tumor sites.

[0263] Provided herein are methods for stimulating an immune response in a subject comprising administering to the subject a CD40 agonist, e.g., an antibody, and one or more additional immunostimulatory antibodies, such as a PD-1 antagonist, e.g., antagonist antibody, a PD-L1 antagonist, e.g., antagonist antibody, a CTLA-4 antagonist, e.g., antagonist antibody and/or a LAG3 antagonist, e.g., an antagonist antibody, such that an immune response is stimulated in the subject, for example to inhibit tumor growth or to stimulate an anti-viral response. In one embodiment, the subject is administered an agonist anti-huCD40 antibody and an antagonist anti-PD-1 antibody. In one embodiment, the subject is administered an agonist anti-huCD40 antibody and an antagonist anti-PD-L1 antibody. In one embodiment, the subject is administered an agonist anti-huCD40 antibody and an antagonist anti-CTLA-4 antibody. In one embodiment, the at least one additional immunostimulatory antibody (e.g., an antagonist anti-PD-1, an antagonist anti-PD-L1, an antagonist anti-CTLA-4 and/or an antagonist anti-LAG3 antibody) is a human antibody. Alternatively, the at least one additional immunostimulatory antibody can be, for example, a chimeric or humanized antibody (e.g., prepared from a mouse anti-PD-1, anti-PD-L1, anti-CTLA-4 and/or anti-LAG3 antibody).

[0264] Provided herein are methods for treating a hyperproliferative disease (e.g., cancer), comprising administering an agonist anti-huCD40 antibody and an antagonist PD-1 antibody to a subject. In certain embodiments, the agonist anti-huCD40 antibody is administered at a subtherapeutic dose, the anti-PD-1 antibody is administered at a subtherapeutic dose, or both are administered at a subtherapeutic dose. Also provided herein are methods for altering an adverse event associated with treatment of a hyperproliferative disease with an immunostimulatory agent, comprising administering an agonist anti-huCD40 antibody and a subtherapeutic dose of anti-PD-1 antibody to a subject. In certain embodiments, the subject is human. In certain embodiments, the anti-PD-1 antibody is a human sequence monoclonal antibody and the agonist anti-huCD40 antibody is a humanized monoclonal antibody, such as an antibody comprising the CDRs or variable regions of the antibodies disclosed herein.

[0265] Suitable PD-1 antagonists for use in the methods described herein, include, without limitation, ligands, antibodies (e.g., monoclonal antibodies and bispecific antibodies),

and multivalent agents. In one embodiment, the PD-1 antagonist is a fusion protein, e.g., an Fc fusion protein, such as AMP-244. In one embodiment, the PD-1 antagonist is an anti-PD-1 or anti-PD-L1 antibody.

[0266] An exemplary anti-PD-1 antibody is OPDIVO®/nivolumab (BMS-936558) or an antibody that comprises the CDRs or variable regions of one of antibodies 17D8, 2D3, 4H1, 5C4, 7D3, 5F4 and 4A11 described in WO 2006/121168. In certain embodiments, an anti-PD-1 antibody is MK-3475 (KEYTRUDA®/pembrolizumab/formerly lam-brolizumab) described in WO 2012/145493; AMP-514/MEDI-0680 described in WO 2012/145493; and CT-011 (pidilizumab; previously CT-AcTibody or BAT; see, e.g., Rosenblatt et al. (2011) *J. Immunotherapy* 34:409). Further known PD-1 antibodies and other PD-1 inhibitors include those described in WO 2009/014708, WO 03/099196, WO 2009/114335, WO 2011/066389, WO 2011/161699, WO 2012/145493, U.S. Pat. Nos. 7,635,757 and 8,217,149, and U.S. Patent Publication No. 2009/0317368. Any of the anti-PD-1 antibodies disclosed in WO 2013/173223 may also be used. An anti-PD-1 antibody that competes for binding with, and/or binds to the same epitope on PD-1 as, as one of these antibodies may also be used in combination treatments.

[0267] In certain embodiments, the anti-PD-1 antibody binds to human PD-1 with a K_D of 5×10^{-8} M or less, binds to human PD-1 with a K_D of 1×10^{-8} M or less, binds to human PD-1 with a K_D of 5×10^{-9} M or less, or binds to human PD-1 with a K_D of between 1×10^{-8} M and 1×10^{-10} M or less.

[0268] Provided herein are methods for treating a hyperproliferative disease (e.g., cancer), comprising administering an agonist anti-huCD40 antibody and an antagonist PD-L1 antibody to a subject. In certain embodiments, the agonist anti-huCD40 antibody is administered at a subtherapeutic dose, the anti-PD-L1 antibody is administered at a subtherapeutic dose, or both are administered at a subtherapeutic dose. Provided herein are methods for altering an adverse event associated with treatment of a hyperproliferative disease with an immunostimulatory agent, comprising administering an agonist anti-huCD40 antibody and a subtherapeutic dose of anti-PD-L1 antibody to a subject. In certain embodiments, the subject is human. In certain embodiments, the anti-PD-L1 antibody is a human sequence monoclonal antibody and the agonist anti-huCD40 antibody is a humanized monoclonal antibody, such as an antibody comprising the CDRs or variable regions of the antibodies disclosed herein.

[0269] In one embodiment, the anti-PD-L1 antibody is BMS-936559 (referred to as 12A4 in WO 2007/005874 and U.S. Pat. No. 7,943,743), MSB0010718C (WO 2013/79174), or an antibody that comprises the CDRs or variable regions of 3G10, 12A4, 10A5, 5F8, 10H10, 1B12, 7H1, 11E6, 12B7 and 13G4, which are described in PCT Publication WO 07/005874 and U.S. Pat. No. 7,943,743. In certain embodiment an anti-PD-L1 antibody is MEDI4736 (also known as Anti-B7-H1) or MPDL3280A (also known as RG7446). Any of the anti-PD-L1 antibodies disclosed in WO 2013/173223, WO 2011/066389, WO 2012/145493, U.S. Pat. Nos. 7,635,757 and 8,217,149 and U.S. Publication No. 2009/145493 may also be used. Anti-PD-L1 antibodies that compete with and/or bind to the same epitope as that of any of these antibodies may also be used in combination treatments.

[0270] In yet further embodiment, the agonist anti-huCD40 antibody of the present invention is combined with an antagonist of PD-1/PD-L1 signaling, such as a PD-1 antagonist or a PD-L1 antagonist, in combination with a third immunotherapeutic agent. In one embodiment the third immunotherapeutic agent is a GITR antagonist or an OX-40 antagonist, such as the anti-GITR or anti-OX40 antibodies disclosed herein.

[0271] In another aspect, the immuno-oncology agent is a GITR agonist, such as an agonistic GITR antibody. Suitable GITR antibodies include, for example, BMS-986153, BMS-986156, TRX-518 (WO 06/105021, WO 09/009116) and MK-4166 (WO 11/028683).

[0272] In another aspect, the immuno-oncology agent is an IDO antagonist. Suitable IDO antagonists include, for example, INCB-024360 (WO 2006/122150, WO 07/75598, WO 08/36653, WO 08/36642), indoximod, or NLG-919 (WO 09/73620, WO 09/1156652, WO 11/56652, WO 12/142237).

[0273] Provided herein are methods for treating a hyperproliferative disease (e.g., cancer), comprising administering an agonist anti-huCD40 antibody described herein and a CTLA-4 antagonist antibody to a subject. In certain embodiments, the agonist anti-huCD40 antibody is administered at a subtherapeutic dose, the anti-CTLA-4 antibody is administered at a subtherapeutic dose, or both are administered at a subtherapeutic dose. Provided herein are methods for altering an adverse event associated with treatment of a hyperproliferative disease with an immunostimulatory agent, comprising administering an agonist anti-huCD40 antibody and a subtherapeutic dose of anti-CTLA-4 antibody to a subject. In certain embodiments, the subject is human. In certain embodiments, the anti-CTLA-4 antibody is an antibody selected from the group consisting of: YERVOY® (ipilimumab or antibody 10D1, described in PCT Publication WO 01/14424), tremelimumab (formerly ticilimumab, CP-675,206), and the anti-CTLA-4 antibodies described in the following publications: WO 98/42752; WO 00/37504; U.S. Pat. No. 6,207,156; Hurwitz et al. (1998) *Proc. Natl. Acad. Sci. (USA)* 95(17):10067-10071; Camacho et al. (2004) *J. Clin. Oncology* 22(145): Abstract No. 2505 (antibody CP-675206); and Mokyr et al. (1998) *Cancer Res.* 58:5301-5304. Any of the anti-CTLA-4 antibodies disclosed in WO 2013/173223 may also be used.

[0274] Provided herein are methods for treating a hyperproliferative disease (e.g., cancer), comprising administering an agonist anti-huCD40 antibody and an anti-LAG-3 antibody to a subject. In further embodiments, the agonist anti-huCD40 antibody is administered at a subtherapeutic dose, the anti-LAG-3 antibody is administered at a subtherapeutic dose, or both are administered at a subtherapeutic dose. Provided herein are methods for altering an adverse event associated with treatment of a hyperproliferative disease with an immunostimulatory agent, comprising administering an agonist anti-huCD40 antibody and a subtherapeutic dose of anti-LAG-3 antibody to a subject. In certain embodiments, the subject is human. In certain embodiments, the anti-LAG-3 antibody is a human sequence monoclonal antibody and the agonist anti-huCD40 antibody is a humanized monoclonal antibody, such as an antibody comprising the CDRs or variable regions of the antibodies disclosed herein. Examples of anti-LAG3 antibodies include antibodies comprising the CDRs or variable regions of antibodies 25F7, 26H10, 25E3, 8B7, 11F2 or 17E5, which are

described in U.S. Patent Publication No. US 2011/0150892 and WO 2014/008218. In one embodiment, an anti-LAG-3 antibody is BMS-986016. Other art recognized anti-LAG-3 antibodies that can be used include IMP731 described in US 2011/007023. IMP-321 may also be used. Anti-LAG-3 antibodies that compete with and/or bind to the same epitope as that of any of these antibodies may also be used in combination treatments.

[0275] In certain embodiments, the anti-LAG-3 antibody binds to human LAG-3 with a K_D of 5×10^{-8} M or less, binds to human LAG-3 with a K_D of 1×10^{-8} M or less, binds to human LAG-3 with a K_D of 5×10^{-9} M or less, or binds to human LAG-3 with a K_D of between 1×10^{-8} M and 1×10^{-10} M or less.

[0276] Administration of agonist anti-huCD40 antibodies described herein and antagonists, e.g., antagonist antibodies, to one or more second target antigens such as LAG-3 and/or CTLA-4 and/or PD-1 and/or PD-L1 can enhance the immune response to cancerous cells in the patient. Cancers whose growth may be inhibited using the antibodies of the instant disclosure include cancers typically responsive to immunotherapy. Representative examples of cancers for treatment with the combination therapy of the instant disclosure include those cancers specifically listed above in the discussion of monotherapy with agonist anti-huCD40 antibodies.

[0277] In certain embodiments, the combination of therapeutic antibodies discussed herein can be administered concurrently as a single composition in a pharmaceutically acceptable carrier, or concurrently as separate compositions with each antibody in a pharmaceutically acceptable carrier. In another embodiment, the combination of therapeutic antibodies can be administered sequentially. For example, an anti-CTLA-4 antibody and an agonist anti-huCD40 antibody can be administered sequentially, such as anti-CTLA-4 antibody being administered first and agonist anti-huCD40 antibody second, or agonist anti-huCD40 antibody being administered first and anti-CTLA-4 antibody second. Additionally or alternatively, an anti-PD-1 antibody and an agonist anti-huCD40 antibody can be administered sequentially, such as anti-PD-1 antibody being administered first and agonist anti-huCD40 antibody second, or agonist anti-huCD40 antibody being administered first and anti-PD-1 antibody second. Additionally or alternatively, an anti-PD-L1 antibody and an agonist anti-huCD40 antibody can be administered sequentially, such as anti-PD-L1 antibody being administered first and agonist anti-huCD40 antibody second, or agonist anti-huCD40 antibody being administered first and anti-PD-L1 antibody second. Additionally or alternatively, an anti-LAG-3 antibody and an agonist anti-huCD40 antibody can be administered sequentially, such as anti-LAG-3 antibody being administered first and agonist anti-huCD40 antibody second, or agonist anti-huCD40 antibody being administered first and anti-LAG-3 antibody second.

[0278] Furthermore, if more than one dose of the combination therapy is administered sequentially, the order of the sequential administration can be reversed or kept in the same order at each time point of administration, sequential administrations can be combined with concurrent administrations, or any combination thereof. For example, the first administration of a combination anti-CTLA-4 antibody and agonist anti-huCD40 antibody can be concurrent, the second administration can be sequential with anti-CTLA-4 antibody first

and agonist anti-huCD40 antibody second, and the third administration can be sequential with agonist anti-huCD40 antibody first and anti-CTLA-4 antibody second, etc. Additionally or alternatively, the first administration of a combination anti-PD-1 antibody and agonist anti-huCD40 antibody can be concurrent, the second administration can be sequential with anti-PD-1 antibody first and agonist anti-huCD40 antibody second, and the third administration can be sequential with agonist anti-huCD40 antibody first and anti-PD-1 antibody second, etc. Additionally or alternatively, the first administration of a combination anti-PD-L1 antibody and agonist anti-huCD40 antibody can be concurrent, the second administration can be sequential with anti-PD-L1 antibody first and agonist anti-huCD40 antibody second, and the third administration can be sequential with agonist anti-huCD40 antibody first and anti-PD-L1 antibody second, etc. Additionally or alternatively, the first administration of a combination anti-LAG-3 antibody and agonist anti-huCD40 antibody can be concurrent, the second administration can be sequential with anti-LAG-3 antibody first and agonist anti-huCD40 antibody second, and the third administration can be sequential with agonist anti-huCD40 antibody first and anti-LAG-3 antibody second, etc. Another representative dosing scheme can involve a first administration that is sequential with agonist anti-huCD40 first and anti-CTLA-4 antibody (and/or anti-PD-1 antibody and/or anti-PD-L1 antibody and/or anti-LAG-3 antibody) second, and subsequent administrations may be concurrent.

[0279] Optionally, an agonist anti-huCD40 as sole immunotherapeutic agent, or the combination of an agonist anti-huCD40 antibody and one or more additional immunotherapeutic antibodies (e.g., anti-CTLA-4 and/or anti-PD-1 and/or anti-PD-L1 and/or anti-LAG-3 blockade) can be further combined with an immunogenic agent, such as cancerous cells, purified tumor antigens (including recombinant proteins, peptides, and carbohydrate molecules), cells, and cells transfected with genes encoding immune stimulating cytokines (He et al. (2004) *J. Immunol.* 173:4919-28). Non-limiting examples of tumor vaccines that can be used include peptides of melanoma antigens, such as peptides of gp100, MAGE antigens, Trp-2, MART1 and/or tyrosinase, or tumor cells transfected to express the cytokine GM-CSF (discussed further below). A CD40 agonist and one or more additional antibodies (e.g., CTLA-4 and/or PD-1 and/or PD-L1 and/or LAG-3 blockade) can also be further combined with standard cancer treatments. For example, a CD40 agonist and one or more additional antibodies (e.g., CTLA-4 and/or PD-1 and/or PD-L1 and/or LAG-3 blockade) can be effectively combined with chemotherapeutic regimens. In these instances, it is possible to reduce the dose of other chemotherapeutic reagent administered with the combination of the instant disclosure (Mokyr et al. (1998) *Cancer Research* 58: 5301-5304). An example of such a combination is a combination of CD40 agonist antibody with or without an additional antibody, such as anti-CTLA-4 antibodies and/or anti-PD-1 antibodies and/or anti-PD-L1 antibodies and/or anti-LAG-3 antibodies) further in combination with decarbazine for the treatment of melanoma. Another example is a combination of agonist anti-huCD40 antibody with or without anti-CTLA-4 antibodies and/or anti-PD-1 antibodies and/or anti-PD-L1 antibodies and/or LAG-3 antibodies further in combination with interleukin-2 (IL-2) for the treatment of melanoma. The scientific rationale behind the combined use of CD40 agonism and

CTLA-4 and/or PD-1 and/or PD-L1 and/or LAG-3 blockade with chemotherapy is that cell death, which is a consequence of the cytotoxic action of most chemotherapeutic compounds, should result in increased levels of tumor antigen in the antigen presentation pathway. Other combination therapies that may result in synergy with a combined CD40 agonism with or without anti-CTLA-4 and/or PD-1 and/or PD-L1 and/or LAG-3 blockade through cell death include radiation, surgery, or hormone deprivation. Each of these protocols creates a source of tumor antigen in the host. Angiogenesis inhibitors can also be combined with a combined CD40 agonism and CTLA-4 and/or PD-1 and/or PD-L1 and/or LAG-3 blockade. Inhibition of angiogenesis leads to tumor cell death, which can be a source of tumor antigen fed into host antigen presentation pathways.

[0280] An agonist anti-huCD40 antibody as sole immunotherapeutic agent, or a combination of CD40 agonist and CTLA-4 and/or PD-1 and/or PD-L1 and/or LAG-3 blocking antibodies can also be used in combination with bispecific antibodies that target Fc α or Fc γ receptor-expressing effector cells to tumor cells. See, e.g., U.S. Pat. Nos. 5,922,845 and 5,837,243. Bispecific antibodies can be used to target two separate antigens. The T cell arm of these responses would be augmented by the use of a combined CD40 agonism and CTLA-4 and/or PD-1 and/or PD-L1 and/or LAG-3 blockade.

[0281] In another example, an agonistic anti-CD40 antibody as sole immunotherapeutic agent or a combination of an anti-CD40 antibody and additional immunostimulating agent, e.g., anti-CTLA-4 antibody and/or anti-PD-1 antibody and/or anti-PD-L1 antibody and/or LAG-3 agent, e.g., antibody, can be used in conjunction with an anti-neoplastic antibody, such as RITUXAN® (rituximab), HERCEPTIN® (trastuzumab), BEXXAR® (tositumomab), ZEVALIN® (ibritumomab), CAMPATH® (alemtuzumab), LYMPHOCIDE® (epratuzumab), AVASTIN® (bevacizumab), and TARCEVA® (erlotinib), and the like. By way of example and not wishing to be bound by theory, treatment with an anti-cancer antibody or an anti-cancer antibody conjugated to a toxin can lead to cancer cell death (e.g., tumor cells) which would potentiate an immune response mediated by the immunostimulating agent, e.g., CD40, TIGIT, CTLA-4, PD-1, PD-L1 or LAG-3 agent, e.g., antibody. In an exemplary embodiment, a treatment of a hyperproliferative disease (e.g., a cancer tumor) can include an anti-cancer agent, e.g., antibody, in combination with an agonist anti-huCD40 antibody and optionally an additional immunostimulating agent, e.g., anti-CTLA-4 and/or anti-PD-1 and/or anti-PD-L1 and/or anti-LAG-3 agent, e.g., antibody, concurrently or sequentially or any combination thereof, which can potentiate an anti-tumor immune responses by the host.

[0282] Provided herein are methods for altering an adverse event associated with treatment of a hyperproliferative disease (e.g., cancer) with an immunostimulatory agent, comprising administering an agonist anti-huCD40 antibody with or without a subtherapeutic dose of anti-CTLA-4 and/or anti-PD-1 and/or anti-PD-L1 and/or anti-LAG-3 agent, e.g., antibody, to a subject. For example, the methods described herein provide for a method of reducing the incidence of immunostimulatory therapeutic antibody-induced colitis or diarrhea by administering a non-absorbable steroid to the patient. As used herein, a "non-absorbable steroid" is a glucocorticoid that exhibits extensive first pass metabolism such that, following metabolism in the liver, the bioavail-

ability of the steroid is low, i.e., less than about 20%. In one embodiment described herein, the non-absorbable steroid is budesonide. Budesonide is a locally-acting glucocorticosteroid, which is extensively metabolized, primarily by the liver, following oral administration. ENTOCORT EC® (ASTRA-ZENECA) is a pH- and time-dependent oral formulation of budesonide developed to optimize drug delivery to the ileum and throughout the colon. ENTOCORT EC® is approved in the U.S. for the treatment of mild to moderate Crohn's disease involving the ileum and/or ascending colon. The usual oral dosage of ENTOCORT EC® for the treatment of Crohn's disease is 6 to 9 mg/day. ENTOCORT EC® is released in the intestines before being absorbed and retained in the gut mucosa. Once it passes through the gut mucosa target tissue, ENTOCORT EC® is extensively metabolized by the cytochrome P450 system in the liver to metabolites with negligible glucocorticoid activity. Therefore, the bioavailability is low (about 10%). The low bioavailability of budesonide results in an improved therapeutic ratio compared to other glucocorticoids with less extensive first-pass metabolism. Budesonide results in fewer adverse effects, including less hypothalamic-pituitary suppression, than systemically-acting corticosteroids. However, chronic administration of ENTOCORT EC® can result in systemic glucocorticoid effects such as hypercorticism and adrenal suppression. See PDR 58th ed. 2004; 608-610.

[0283] In still further embodiments, a CD40 agonist with or without CTLA-4 and/or PD-1 and/or PD-L1 and/or LAG-3 blockade (i.e., immunostimulatory therapeutic antibodies against CD40 and optionally anti-CTLA-4 and/or anti-PD-1 and/or anti-PD-L1 and/or anti-LAG-3 antibodies) in conjunction with a non-absorbable steroid can be further combined with a salicylate. Salicylates include 5-ASA agents such as, for example: sulfasalazine (AZULFIDINE®, PHARMACIA & UPJOHN); olsalazine (DIPENTUM®, PHARMACIA & UPJOHN); balsalazide (COLAZAL®, SALIX Pharmaceuticals, Inc.); and mesalamine (ASACOL®, PROCTER & GAMBLE Pharmaceuticals; PENTASA®, SHIRE US; CANASA®, AXCAN SCANDIPHARM, Inc.; ROWASA®, SOLVAY).

[0284] In accordance with the methods described herein, a salicylate administered in combination with agonist anti-huCD40 antibody with or without anti-CTLA-4 and/or anti-PD-1 and/or anti-PD-L1 and/or LAG-3 antibodies and a non-absorbable steroid can include any overlapping or sequential administration of the salicylate and the non-absorbable steroid for the purpose of decreasing the incidence of colitis induced by the immunostimulatory antibodies. Thus, for example, methods for reducing the incidence of colitis induced by the immunostimulatory antibodies described herein encompass administering a salicylate and a non-absorbable concurrently or sequentially (e.g., a salicylate is administered 6 hours after a non-absorbable steroid), or any combination thereof. Further, a salicylate and a non-absorbable steroid can be administered by the same route (e.g., both are administered orally) or by different routes (e.g., a salicylate is administered orally and a non-absorbable steroid is administered rectally), which may differ from the route(s) used to administer the anti-huCD40 and anti-CTLA-4 and/or anti-PD-1 and/or anti-PD-L1 and/or anti-LAG-3 antibodies.

[0285] The agonist anti-huCD40 antibodies and combination antibody therapies described herein may also be used in conjunction with other well known therapies that are

selected for their particular usefulness against the indication being treated (e.g., cancer). Combinations of the agonist anti-huCD40 antibodies described herein may be used sequentially with known pharmaceutically acceptable agent(s).

[0286] For example, the agonist anti-huCD40 antibodies and combination antibody therapies described herein can be used in combination (e.g., simultaneously or separately) with an additional treatment, such as irradiation, chemotherapy (e.g., using camptothecin (CPT-11), 5-fluorouracil (5-FU), cisplatin, doxorubicin, irinotecan, paclitaxel, gemcitabine, cisplatin, paclitaxel, carboplatin-paclitaxel (TAXOL), doxorubicin, 5-fu, or camptothecin+apo2l/TRAIL (a 6x combo)), one or more proteasome inhibitors (e.g., bortezomib or MG132), one or more Bcl-2 inhibitors (e.g., BH3I-2' (bcl-xl inhibitor), indoleamine dioxygenase-1 (IDO1) inhibitor (e.g., INCB24360), AT-101 (R-(-)-gossypol derivative), ABT-263 (small molecule), GX-15-070 (obatoclax), or MCL-1 (myeloid leukemia cell differentiation protein-1) antagonists), iAP (inhibitor of apoptosis protein) antagonists (e.g., smac7, smac4, small molecule smac mimetic, synthetic smac peptides (see Fulda et al., *Nat Med* 2002; 8:808-15), ISIS23722 (LY2181308), or AEG-35156 (GEM-640)), HDAC (histone deacetylase) inhibitors, anti-CD20 antibodies (e.g., rituximab), angiogenesis inhibitors (e.g., bevacizumab), anti-angiogenic agents targeting VEGF and VEGFR (e.g., AVASTIN®), synthetic triterpenoids (see Hyer et al., *Cancer Research* 2005; 65:4799-808), c-FLIP (cellular FLICE-inhibitory protein) modulators (e.g., natural and synthetic ligands of PPARγ (peroxisome proliferator-activated receptor γ), 5809354 or 5569100), kinase inhibitors (e.g., Sorafenib), trastuzumab, cetuximab, Temsirolimus, mTOR inhibitors such as rapamycin and temsirolimus, Bortezomib, JAK2 inhibitors, HSP90 inhibitors, PI3K-AKT inhibitors, Lenalidomide, GSK3β inhibitors, IAP inhibitors and/or genotoxic drugs.

[0287] The agonist anti-huCD40 antibodies and combination antibody therapies described herein can further be used in combination with one or more anti-proliferative cytotoxic agents. Classes of compounds that may be used as anti-proliferative cytotoxic agents include, but are not limited to, the following:

[0288] Alkylating agents (including, without limitation, nitrogen mustards, ethylenimine derivatives, alkyl sulfonates, nitrosoureas and triazines): Uracil mustard, Chloromethine, Cyclophosphamide (CYTOXAN™) fosfamide, Melphalan, Chlorambucil, Pipobroman, Triethylenemelamine, Triethylenethiophosphoramide, Busulfan, Carmustine, Lomustine, Streptozocin, Dacarbazine, and Temozolomide.

[0289] Antimetabolites (including, without limitation, folic acid antagonists, pyrimidine analogs, purine analogs and adenosine deaminase inhibitors): Methotrexate, 5-Fluorouracil, Floxuridine, Cytarabine, 6-Mercaptopurine, 6-Thioguanine, Fludarabine phosphate, Pentostatin, and Gemcitabine.

[0290] Suitable anti-proliferative agents for combining with agonist anti-huCD40 antibodies, without limitation, taxanes, paclitaxel (paclitaxel is commercially available as TAXOL™), docetaxel, discodermolide (DDM), dictyostatin (DCT), Peloruside A, epothilones, epothilone A, epothilone B, epothilone C, epothilone D, epothilone E, epothilone F, furanopothilone D, desoxyepothilone BI, [17]-dehydrodesoxyepothilone B, [18]dehydrodesoxyepothilones B, C12,13-

cyclopropyl-epothilone A, C6-C8 bridged epothilone A, trans-9,10-dehydroepothilone D, cis-9,10-dehydroepothilone D, 16-desmethylepothilone B, epothilone B10, discoderomolide, patupilone (EPO-906), KOS-862, KOS-1584, ZK-EPO, ABI-789, XAA296A (Discodermolide), TZT-1027 (soblidotin), ILX-651 (tasidotin hydrochloride), Halichondrin B, Eribulin mesylate (E-7389), Hemiasterlin (HTI-286), E-7974, Cyrtophycins, LY-355703, Maytansinoid immunoconjugates (DM-1), MKC-1, ABT-751, T1-38067, T-900607, SB-715992 (ispinesib), SB-743921, MK-0731, STA-5312, eleutherobin, 17 β -acetoxy-2-ethoxy-6-oxo-B-homo-estra-1,3,5(10)-trien-3-ol, cyclostreptin, isolaulimalide, laulimalide, 4-epi-7-dehydroxy-14,16-didemethyl-(+)-discodermolides, and cryptothilone 1, in addition to other microtubuline stabilizing agents known in the art.

[0291] In cases where it is desirable to render aberrantly proliferative cells quiescent in conjunction with or prior to treatment with agonist anti-huCD40 antibodies described herein, hormones and steroids (including synthetic analogs), such as 17 α -Ethinylestradiol, Diethylstilbestrol, Testosterone, Prednisone, Fluoxymesterone, Dromostanolone propionate, Testolactone, Megestrolacetate, Methylprednisolone, Methyl-testosterone, Prednisolone, Triamcinolone, Chlorotrianisene, Hydroxyprogesterone, Aminoglutethimide, Estramustine, Medroxyprogesteroneacetate, Leuprolide, Flutamide, Toremifene, ZOLADEX™, can also be administered to the patient. When employing the methods or compositions described herein, other agents used in the modulation of tumor growth or metastasis in a clinical setting, such as antimetotics, can also be administered as desired.

[0292] Methods for the safe and effective administration of chemotherapeutic agents are known to those skilled in the art. In addition, their administration is described in the standard literature. For example, the administration of many of the chemotherapeutic agents is described in the Physicians' Desk Reference (PDR), e.g., 1996 edition (Medical Economics Company, Montvale, N.J. 07645-1742, USA); the disclosure of which is incorporated herein by reference thereto.

[0293] The chemotherapeutic agent(s) and/or radiation therapy can be administered according to therapeutic protocols well known in the art. It will be apparent to those skilled in the art that the administration of the chemotherapeutic agent(s) and/or radiation therapy can be varied depending on the disease being treated and the known effects of the chemotherapeutic agent(s) and/or radiation therapy on that disease. Also, in accordance with the knowledge of the skilled clinician, the therapeutic protocols (e.g., dosage amounts and times of administration) can be varied in view of the observed effects of the administered therapeutic agents on the patient, and in view of the observed responses of the disease to the administered therapeutic agents.

XI. Characterization of Specific Agonist Anti-CD40 Antibodies of the Present Invention

[0294] Agonist anti-CD40 antibodies of the present invention were obtained as described in Example 1. Variable domains and CDR sequence regions of exemplary antibodies of the present invention are provided in the Sequence Listing, and are summarized at Table 2. Variable domain and CDR region numbering is the same for all antibodies derived

from the same original clone, i.e. the humanized variants provided herein do not include any insertions or deletions.

TABLE 2

Antibody Variable Domains and CDRs					
Clone	Chain	Variable Domain	CDR1	CDR2	CDR3
12D6	Heavy chain	1-119	31-35	50-66	99-108
12D6	Light chain	1-112	24-39	55-61	94-102
5F11	Heavy chain	1-117	31-35	50-66	99-106
5F11	Light chain	1-111	24-38	54-60	93-101
8E8	Heavy chain	1-122	31-35	50-66	99-111
8E8	Light chain	1-112	24-39	55-61	94-102
5G7	Heavy chain	1-113	31-35	50-66	99-102
5G7	Light chain	1-107	24-34	50-56	89-97
19G3	Heavy chain	1-112	31-35	50-66	99-101
19G3	Light chain	1-112	24-39	55-61	94-102

[0295] The invention also provides anti-huCD40 antibodies related to those disclosed herein by sequence by being derived from the same murine germline sequences, specifically the V and J region gene segments. The murine germline sequences for each of the antibodies disclosed herein are provided at Table 3.

TABLE 3

Mouse Germline Sequences for anti-huCD40 mAbs			
Clone	Chain	V region	J region
12D6	Heavy chain	VH1-39_01	IGHJ4
12D6	Light chain	VK1-110_01	IGKJ1
5F11	Heavy chain	VH1-4_02	IGHJ3
5F11	Light chain	VK3-5_01	IGKJ5
8E8	Heavy chain	VH1-80_01	IGHJ2
8E8	Light chain	VK1-110_01	IGKJ2
5G7	Heavy chain	VH1-18_01	IGHJ4
5G7	Light chain	VK10-96_01	IGKJ2
19G3	Heavy chain	VH5-9-4_01	IGHJ3
19G3	Light chain	VK1-117_01	IGKJ2

[0296] The invention further provides humanized anti-huCD40 antibodies derived from the murine parental antibodies disclosed herein by replacement of murine framework sequences with human framework sequences, with or without additional mutations ("back mutations") to restore antigen (human CD40) affinity that would otherwise be lost in the humanization or to remove sequence liabilities. See Example 3.

[0297] The invention also provides antibody constructs comprising the novel variable domain sequences disclosed herein and constant domains with modified Fc regions having enhanced affinity for Fc γ RIIb as compared with their affinity for other Fc receptors, i.e. activating receptors. Such agonistic anti-huCD40 antibodies with enhanced Fc γ RIIb-specificity are expected to exhibit superior efficacy in treatment of cancer and chronic infection. Li & Ravetch (2011) *Science* 333:1030; White et al. (2011) *J. Immunol.* 187:1754. Without intending to be limited by theory, such Fc γ RIIb-specific agonistic anti-CD40 mAbs may exhibit enhanced adjuvant effects by increasing the maturation of dendritic cell promoting expansion and activation of cytotoxic CD8⁺ T cells, leading to enhanced anti-tumor response. Id. Without intending to be limited by theory, FcR-mediated signal enhancement of agonist CD40 antibodies due to increased receptor clustering, or "cross-linking," of the present inven-

tion may be a major contributor to therapeutic efficacy. Cross-linking of CD40 agonist antibodies by FcR engagement by the Fc portion of the antibody may increase signal strength and thereby enhance activation of cells.

[0298] The relative binding affinity of antibodies for activating (A) versus inhibitory (I) Fc receptors can be expressed as the “A/I” ratio, and is typically a function of the structure of the Fc region of an IgG antibody. See WO 2012/087928. Antibodies having enhanced specificity for binding to inhibitory receptor FcγRIIb have lower A/I ratios. Preferred antibodies for the agonistic anti-huCD40 antibodies of the present invention have, for example, A/I ratios of less than 5, 4, 3, 2, 1, 0.5, 0.3, 0.1, 0.05, 0.03 or 0.01.

[0299] Human IgG1 constant domains comprising mutations to enhance FcγRIIb specificity are also provided in the Sequence Listing, and are summarized at Table 4 and illustrated at FIG. 1. Sequence variants are defined with reference to human IgG1f constant domain sequence provided at SEQ ID NO: 65. The nomenclature regarding positions (numbering) of mutations in the Fc region is according to the EU index as in Kabat et al. (1981) *Sequences of Proteins of Immunological Interest*, 5th Ed. Public Health Service, National Institutes of Health, Bethesda, Md.), which facilitates comparison of Fc sequences at equivalent positions in antibodies with differing variable domain lengths. See also Edelman et al. (1969) *Proc. Nat'l Acad. Sci. (USA)* 63:78; WO 2012/130831 (using the same numbering system). It does not match the sequence numbering in the Sequence Listing. FIG. 1 provides a graphical representation of the Fc sequence variants of Table 4, from which one of skill in the art could readily recognize the corresponding positions in the antibody sequences disclosed herein. SE and SELF variants are described at Chu et al. (2008) *Mol. Immunol.* 45:3926. P238D, V4, V7, V8, V9, V11 and V12 variants are described at Mimoto et al. (2013) *Protein Engineering Design & Selection* 26:589 (e.g. at table 1 therein).

TABLE 4

Fc Sequence Variants	
Designation	SEQ ID: Sequence Variants
IgG1f	65
SE	66 S267E
SELF	67 S267E L328F
P238D	68 P238D
V4	69 P238D P271G
V4-D270E	70 P238D P271G D270E
V7	71 E233D P238D P271G A330R
V8	72 G237D P238D H268D P271G A330R
V9	73 G237D P238D P271G A330R D270E
V9-D270E	74 G237D P238D P271G A330R
V11	75 G237D P238D H268D P271G A330R
V12	76 E233D G237D P238D H268D P271G A330R

[0300] Additional Fc sequence variants with enhanced affinity for FcγRIIb are disclosed at Yu et al. (2013) *J. Am. Chem. Soc.* 135:9723 (and WO 2014/184545), including V262E and V264E, e.g. for use in combination with S267E and L328F.

[0301] Additional variants with D270E mutations were produced to reduce the D270 isomerization that occurred in the V4 and V9 Fc sequence variants, which otherwise exhibited enhanced isomerization rates in accelerated degradation studies suggesting ~12-16% isomerization after

two years at 4° C. in PBS, compared with ~5-7% for other variants. See Table 5 (providing data from a single experiment but replicate experiments showed comparable values). Substitution of aspartic acid at position 270 with glutamic acid eliminated the possibility of such DG isomerization, resulting in antibodies that are more chemically stable and antibody preparations that are more homogenous. Other D270 substitutions of V9, such as D270A, D270Q, D270S and D270T, effectively eliminated binding to FcγRIIa and FcγRIIb receptors (K_D was greater than 5 μM). Although the D270E mutation reduced binding to FcγRIIb receptor by an order of magnitude, the V9-D270E variant maintained a favorable bias in affinity for FcγRIIb receptor compared with FcγRIIa, and acceptable absolute affinity for FcγRIIb. See Example 8.

TABLE 5

Fc Sequence Variant Affinity for Fcγ Receptors (K_D in nM)			
Designation	FcγRIIa-H131	FcγRIIa-R131	FcγRIIb
IgG1f	530	850	3900
SE	520	22	98
SELF	1100	2	11
P238D	>5000	>5000	950
V4	>5000	1900	150
V4-D270E	>5000	>5000	1800
V7	>5000	1600	84
V8	>5000	1400	93
V9	>5000	420	15
V9-D270A,Q,S,T	>5000	>5000	>5000
V9-D270E	>5000	>5000	150
V11	—	450	15
V12	>5000	490	21

[0302] Agonist activity of various anti-CD40 antibodies of the present invention was measured. See Example 7, and

FIGS. 3A, 3B and 4. Agonist activity was found to depend on both the variable domain sequences (mAb clone number), which determine antigen (human CD40) binding, and the sequence of the Fc region, which determines Fc receptor (FcγRIIb) binding.

[0303] The present disclosure is further illustrated by the following examples, which should not be construed as further limiting. The contents of all figures and all references, Genbank sequences, patents and published patent

applications cited throughout this application are expressly incorporated herein by reference.

EXAMPLES

Example 1

Generation of Mouse Monoclonal Antibodies Against Human CD40

[0304] Murine anti-human CD40 monoclonal antibodies were generated using wild type Balb/c mice (Charles Rivers Labs) that express mouse antibody genes, as follows.

[0305] Antigen

[0306] A huCD40-muFc soluble recombinant protein was used as the antigen for immunizations. The soluble fusion protein has a MW of 91.6 K_D and is composed of the extracellular portion of huCD40 linked to a mouse IgG2b Fc at its C-terminus. This fusion protein is referred to herein as “huCD40-muFc fusion protein”. The fusion protein was generated by standard recombinant DNA methods and expressed in transfected CHO cells, which secreted the soluble fusion protein into the culture supernatant. The CHO host cells used for transfection were obtained from INVITROGEN (Cat #11619-012). The secreted soluble fusion protein was purified for use as immunogen.

[0307] Immunization of Mice

[0308] To generate mouse monoclonal antibodies to human CD40, Balb/c mice were immunized with purified huCD40-muFc fusion protein. The mice were approximately 2-4 months of age upon the first infusion of antigen. Purified recombinant huCD40-muFc antigen preparation (10 µg purified from transfected mammalian cells expressing the fusion protein) was used to immunize the mice using two immunization protocols (A and B). Protocol A consisted of four weekly footpad (FP) immunizations and protocol B of seven weekly subcutaneous (SC)/intraperitoneal (IP)/Hock immunizations. In both cases, the immunogen was mixed 1:1 with RIBI adjuvant (SIGMA Cat # M6536).

[0309] All mice were bled one week after the fourth immunizations to assess antigen specific titers. Final bleeds were also obtained from each mouse at time of sacrifice. The immune response was monitored by retro orbital bleeds. The plasma was screened by ELISA analysis using the recombinant protein used for immunizations. Mice from protocol A received two final boosts intravenously (IV) and in the footpads (FP) with soluble antigen on days -2 and -3 before fusion. Mice from protocol B received two final boosts intravenously as well as IP and in the hock with the soluble antigen on days -2 and -3 before fusion.

[0310] All four mice from protocol A (IDs 291763, 291764, 291765, 291766) were sacrificed. Lymph node and spleen cells were extracted and mixed for fusion (fusions 3582 and 3583). Only two mice from protocol B (IDs 294286 and 294288) were sacrificed. Spleen and lymph nodes from each mouse were mixed and fused (fusions 3716 and 3717).

[0311] Generation of Hybridomas Producing Monoclonal Antibodies to Human CD40

[0312] Mouse splenocytes and lymph nodes isolated from high titer Balb/c mice were fused with a mouse myeloma fusion partner using the electric field based electrofusion Hybri-mune instrument and a large 0.9ml fusion chamber (BTX Harvard Apparatus, Inc., Holliston, Mass.). Single cell suspensions of lymphocytes from immunized mice were

fused to an equal number of P3X63 Ag8.6.53 (ATCC CRL 1580) non-secreting mouse myeloma cells. Resulting cells were plated at 2.0×10^4 cells/well in flat bottom microtiter plates in selective CLONACELL-HY Medium E (catalog #03805; STEMCELL Technologies Inc., Vancouver BC, Canada) with addition of Aminopterin to select for hybridomas. After about 7 days, the culture medium was replaced with Medium E (without aminopterin).

[0313] After 10 to 12 days, individual wells were screened for the presence of mouse IgG/mouse kappa light chain antibodies using a homogenous HTRF assay. In this assay, supernatants from 96 well fusion plates were mixed with custom labeled Terbium-Cryptate goat anti-mouse IgG (Fey specific) (CISBIO US Inc. Bedford, Mass.) and goat anti-mouse IgG Fab'2 labeled with AlexaFluor 647 (Jackson ImmunoResearch; Catalog #109-605-098). Incubation was for 1 hour. The plates were then read on a RUBYSTAR reader. Hybridoma cells from wells positive for mouse IgG/mouse kappa light chain antibodies were then screened either by FACS using CHO cells transfected with human CD40 and CHO untransfected cells as control (fusions 3582/3583) or by ELISA using recombinant protein followed by FACS on Daudi B cells and Jurkat T cells as negative control (fusions 3716/3717). FACS positive parental lines were transferred to 24-well plates. A few days later, cell supernatants from individual wells were rescreened by FACS to confirm IgG specificity to human CD40.

[0314] A panel of antigen specific hybridomas were cloned by serial dilution and rescreened by FACS using either CHO transfectants or Daudi cells. Nineteen antibodies from fusions 3582/3583 and twenty antibodies from fusions 3716/3717 were purified and tested for functional activity. From this panel of antibodies, five strong agonists were selected for second subcloning. These five antibodies, namely 1802.3582.19G3.F10.E1, 1802.3583.5G7.F12.G3, 1802.3583.8E8.C10.G2, 1802.3716.12D6.B1.E3, and 1802.3717.5F11.A11.E7 were subsequently submitted to sequencing and further analysis.

Example 2

Generation of Fully Human Anti-huCD40 Antibodies

[0315] Fully human anti-huCD40 monoclonal antibodies that bind to the same epitope and/or cross-block the binding of the humanized anti-CD40 antibodies disclosed herein may find use in methods of the present invention. Such antibodies may be generated using transgenic mice that express human antibody genes, as follows.

[0316] Antigen

[0317] A huCD40 soluble recombinant protein is used as the antigen for immunization. The soluble fusion protein is composed of the extracellular portion of huCD40 linked to a mouse IgG2a Fc at its C-terminus. This fusion protein is referred to herein as “huCD40-muFc fusion protein.” The fusion protein is generated by standard recombinant DNA methods and expressed in transfected CHO cells, which secrete the soluble fusion protein into the culture supernatant. The CHO host cells used for transfection are obtained from INVITROGEN (Cat #11619-012). The secreted soluble fusion protein is purified for use as immunogen. The sequence of full length human CD40 including signal sequence is provided at SEQ ID NO: 1.

[0318] Transgenic Mice

[0319] Fully human monoclonal antibodies to human CD40 are prepared using mice from the CMD++; JKD++; KCo5(9272)+⁺; SC20+ genotype (hereafter called KM® mice). Individual transgene designations are in parentheses, followed by line numbers for randomly integrated transgenes. The symbols ++ and + indicate homozygous or hemizygous; however, because the mice are routinely screened using a PCR-based assay that does not allow us to distinguish between heterozygosity and homozygosity for the randomly integrated human Ig transgenes, a + designation may be given to mice that are actually homozygous for these elements. In this strain, the endogenous mouse kappa light chain gene has been homozygously disrupted as described in Chen et al. (1993) *EMBO J.* 12:811-820 and the endogenous mouse heavy chain gene has been homozygously disrupted as described in example 1 of WO 2001/09187. Furthermore, this mouse strain carries a human kappa light chain transgene, KCo5, as described in Fishwild et al. (1996) *Nature Biotechnology* 14:845-851, a yeast artificial chromosome (YAC) carrying most of the human kappa light chain locus, as described in WO 2000/026373.

[0320] Immunization of Mice

[0321] To generate fully human monoclonal antibodies to human CD40, KM mice are immunized with purified huCD40-muFc fusion protein. General immunization schemes are described in Lonberg et al. (1994) *Nature* 368(6474): 856-859; Fishwild, D. et al. (1996) *Nature Biotechnology* 14: 845-851 and WO 98/24884. The mice are approximately 4 months of age upon the first infusion of antigen. Either purified recombinant huCD40-muFc antigen preparation (10 µg purified from transfected mammalian cells expressing the fusion protein) or 300-19 cells transfected with human CD40 are used to immunize the mice intraperitoneally and subcutaneously. The immunogens are mixed 1:1 with RIBI adjuvant (SIGMA Cat # M6536).

[0322] The mice are immunized five times at 5-7 day intervals. The first and second immunizations are performed with the recombinant protein. The third immunization is with the cells, the fourth immunization with the protein and the fifth immunization with the cells. Mice are bled one week after the last immunizations to assess antigen specific titers. The immune response is monitored by retro orbital bleeds. The plasma is screened by FACS analysis using the transfected 300-19 cells, and mice with highest titers for anti-human CD40 human IgG are used for fusions. Mice receive a final boost by intravenous (IV) and intraperitoneal (IP) injection of soluble antigen two days and transfected cells three days before sacrifice and removal of the spleen.

[0323] Generation of Hybridomas Producing Human Monoclonal Antibodies to Human CD40

[0324] Mouse splenocytes isolated from high titer KM mice and a mouse myeloma fusion partner are fused with an electric field based electrofusion using a Cyto Pulse large chamber cell fusion electroporator (Cyto Pulse Sciences, Inc., Glen Burnie, Md.). Single cell suspensions of splenic lymphocytes from immunized mice are fused to an equal number of P3X63 Ag8.6.53 (ATCC CRL 1580) non-secreting mouse myeloma cells (fusion number: 2541). Resulting cells are plated at 2.0×10^4 cells/well in flat bottom microtiter plates in selective DMEM medium containing high glucose (CELLGRO #10-013-CM) and 10% fetal calf serum (HYCLONE # SH30071.03), and supplemented with beta-mercaptoethanol (1000x, GIBCO #21985-023), 7 mM HEPES

(CELLGRO 25-060-C1), additional 2 mM L-glutamine (CELLGRO 25-005-C1), HAT (50x, SIGMA # H-0262), 5% Hybridoma Cloning Factor (BIOVERIS #210001), 10% P388DI (ATCC # CRL TIB-63) conditioned medium and Penicillin-Streptomycin (100x, CELLGRO #30-002-CI). After about 7 days, some of the medium containing HAT is replaced with medium containing HT (CELLGRO #25-047-CI).

[0325] After 10 to 12 days, individual wells are screened for the presence of human IgG/human kappa light chain antibodies using a homogenous HTRF assay. In this assay, supernatants from 96 well fusion plates are mixed with Europium-cryptate labeled goat anti-human IgG (Fc fragment specific), biotinylated goat anti-human kappa light chain (Bethyl # A80-115B), streptavidin-XLent and incubated for 1 hour. The plates are then read on a RUBYSTAR reader.

[0326] Hybridoma cells from wells positive for human IgG/human kappa light chain or human IgG/human lambda light chain antibodies are then screened by FACS using 300-19 cells transfected with human CD40 and 300-19 untransfected cells as control. FACS positive parental lines are transferred to 24-well plates. A few days later, cell supernatants from individual wells are rescreened by FACS to confirm IgG specificity to human CD40.

[0327] The hybridomas are cloned by serial dilution and re-screened by FACS.

Example 3**Humanization of Anti-huCD40 Antibodies**

[0328] Parental (murine) antibodies of the present invention were humanized for potential use as human therapeutics. The closest matching human germline region sequence was selected for each mouse variable domain, and these human framework regions were used to replace murine frameworks in the "CDR graft" versions of the antibodies. Additional amino acid substitutions, such as framework back-mutations, were then made as necessary to restore the binding affinity of the humanized antibody. Further amino acid substitutions were made to eliminate sequence liabilities. The sequence relationships between the various forms of the antibodies of the present invention are provided at Table 6. The binding of various agonistic anti-huCD40 antibodies to soluble human CD40 was determined by Biolayer Interferometry (BLI) analysis using a FORTEBIO Octet RED (Rapid system—Extended Detection) label-free interaction analysis instrument. All studies were performed in 10 mM Na_2PO_4 , 130 mM NaCl, 0.05% Surfactant P20 (pH 7.1) at 25° C. Briefly, anti-huCD40 antibody supernatants (diluted to 10 µg/ml or captured undiluted if supernatant concentration was less than 10 µg/ml) were captured on protein A coated biosensors (PALL FORTEBIO #18-5010) using a loading time of 90s and shake speed of 1000 rpm. Supernatants were first screened for binding to 1 µM recombinant human CD40 monomer (huCD40-monomer) using 180s association and dissociation times, at 1000 rpm, with two 15s conditioning steps using 10 mM glycine pH 1.5 in between binding cycles. All supernatants that demonstrated a binding signal in the 1 µM huCD40-monomer experiment were then tested for binding to seven different concentrations of huCD40-monomer in a three-fold dilution series, where the highest concentration was either 10 µM huCD40-monomer or 1 µM huCD40-monomer depending on the

strength of the binding signal in the 1 μ M screening experiment. Results for selected antibodies and sequence variants thereof are shown in Table 6.

TABLE 6

Binding Affinity of anti-CD40 Antibodies				
Antibody	Type	V _H	V _L	K _D (nM)
12D6	parental	mouse	mouse	3.6
12D6-03	humanized	VH1-18 CDR graft	VKII O11 CDR graft	4.8
12D6-22	humanized	VH1-18 CDR graft	VKII O11 <u>G29A</u>	3.6
12D6-23	humanized	VH1-18 <u>M100bL</u>	VKII O11 CDR graft	4.3
12D6-24	humanized	VH1-18 <u>M100bL</u>	VKII O11 <u>G29A</u>	5.6
5F11	parental	mouse	mouse	4.7
5F11-03	humanized	VH1-e CDR graft	VKIV B3 CDR graft	>5000
5F11-17	humanized	VH1-e G27Y S30T V37I R38K A40R M48I R66K V67T I69L	VKIV B3 CDR graft	8.4
5F11-23	humanized	VH1-e G27Y S30T V37I R38K A40R M48I R66K V67T I69L	VKIV B3 M4L N22S	4.9
5F11-45	humanized	VH1-e G27Y S30T R66K V67T I69L	VKIV B3 M4L	12.3*
19G3	parental	mouse	mouse	870
19G3-03	humanized	VH3-21 CDR graft	VKII O11 CDR graft	no binding
19G3-11	humanized	VH3-21 S94R	VKII O11 CDR graft	850
19G3-22	humanized	VH3-21 S94R	VKII O11 <u>N28Q</u>	99
19G3-23	humanized	VH3-21 S94R	VKII O11 <u>G29A</u>	1300
5G7	parental	mouse	mouse	89
5G7-03	humanized	VH1-18 CDR graft	VKI A20 CDR graft	340
5G7-22	humanized	VH1-18 <u>M69L</u> A93V	VKI A20 CDR graft	50
5G7-25	humanized	VH1-18 <u>M69L</u> T71V A93V <u>G55A</u>	VKI A20 CDR graft	69
5G7-28	humanized	VH1-18 <u>M69L</u> T71V A93V <u>G55A</u> <u>M96L</u>	VKI A20 CDR graft	230
8E8	parental	mouse	mouse	85
8E8-05	humanized	VH1-e CDR graft	VKII A19 CDR graft	no binding
8E8-56	humanized	VH1-e L4F G27Y S30T R66K V67A T68L I69L	VKII A19 CDR graft	82
8E8-62	humanized	VH1-e L4F G27Y S30T R66K V67A T68L I69L <u>D99E</u>	VKII A19 CDR graft	31
8E8-66	humanized	VH1-e L4F G27Y S30T R66K V67A T68L I69L	VKII A19 <u>G29A</u>	82
8E8-67	humanized	VH1-e L4F G27Y S30T R66K V67A T68L I69L <u>D99E</u>	VKII A19 <u>G29A</u>	28
8E8-70	humanized	VH1-e L4F G27Y S30T R66K V67A T68L I69L <u>D99E</u> <u>G55S</u>	VKII A19 CDR graft	low signal
8E8-71	humanized	VH1-e L4F G27Y S30T R66K V67A T68L I69L <u>D99E</u> <u>G55S</u>	VKII A19 <u>G29A</u>	low signal

*= for comparison purposes, the parental 5F11 exhibited a K_D of 8.2 nM (rather than 4.7 nM) when run in the same assay in parallel with the 5F11-45 antibody.

[0329] For humanized antibodies, V_H and V_L columns provide the human variable domain germlines on which framework regions are based. “CDR graft” refers to variable domains comprising unmodified parental (mouse) CDRs grafted directly onto the recited human framework sequences. Residue numbering for all sequence variants in

Table 6 is according to Kabat, and thus the positions of sequence changes in Table 6 does not match the residue numbering of the antibody sequences in the Sequence Listing. Underlining is used to indicate sequence modifications that are intended to correct potential sequence liabilities, i.e. residues that are subject to chemical modification and potential degradation, leading to product heterogeneity. Other sequence modifications are intended to restore affinity, typically a framework back mutation (reverting from the human germline framework sequence to the original murine framework residue). The M69L modification in the heavy chain of antibody 5G7 is both a framework back mutation and a liability correction.

Example 4

Anti-CD40 Antibody Epitope Binning Experiments

[0330] Epitope binning experiments may be conducted to determine which anti-human CD40 antibodies compete with which others for binding to huCD40, and thus bind to similar epitopes. Pairwise competition between anti-huCD40 antibodies is determined as follows, in which a reference antibody is bound to the surface of a sensor chip, a test antibody is pre-incubated with a huCD40 polypeptide construct in a mixture, and the pre-incubated mixture is flowed over the sensor chip to determine the degree to which the test antibody interferes with binding of the huCD40 polypeptide construct to the reference antibody on the chip surface. Such competition experiments may be performed using a BIA-CORE® SPR instrument. Briefly, a reference anti-huCD40 antibody is immobilized onto Sensor Chip CM5 chip (Series S, GE Healthcare CAT # BR-1005-30) surfaces, flowcell2, flowcell3 & flowcell4 (5000 RUs), and flowcell1 is used as a negative control. A test antibody is diluted to 120 μ g/mL (2 $\times$) at starting concentration. A series of dilutions of the test antibody is made by diluting 1:3 concentration of antibody with buffer for seven different concentrations and a control sample (with 0 μ g/mL) to obtain a titration curve. Each antibody concentration series is divided into two halves. In the first half of the concentration series, 40 nM (2 $\times$) human CD40 antigen (e.g. huCD40/Fc) is added to make the final concentration series (60 μ g/mL-0.0 μ g/mL) and 20 nM of final antigen concentration in each well. In the second half of the concentration series, in place of antigen, buffer is added to have the antibody diluted to the same concentration, and this half is treated as the blank. Complexes of the test anti-CD40 antibodies and huCD40/Fc are incubated for 2 hours. 40 μ L complexes are injected on the reference antibody-coated surface at a 30 μ L/min. A BIA-CORE® T200 surface plasmon resonance instrument is used and the running buffer is HBE-EP, GE Healthcare CAT # BR-1001-88, filtered, degassed, 0.01M HEPES, pH7.4, 0.15 NaCl, 3 mM EDTA, 0.005% Surfactant P20. The surface is regenerated with 25 mM NaOH (order code: BR-1003-58, GE Healthcare) at 100 μ L/min for 5 seconds. The data are analyzed using Microsoft Excel where the concentration of test antibodies is plotted against the corresponding response unit to obtain titration curves.

[0331] Results of such epitope binning experiments for the anti-CD40 antibodies of the present invention are provided at FIG. 2. Antibodies 5F11 and 8E8 block ligand (CD40L) binding. The five anti-CD40 antibodies of the present invention fall into three epitope groups—12D6/5G7/19G3, 5F11/5G7/19G3, and 8E8.

Example 5

Epitope Mapping by HDX

[0332] The epitopes for anti-huCD40 antibodies 12D6, 5G7, 19G3 and 5F11 of the present invention were determined by hydrogen/deuterium exchange mass spectrometry (HDX-MS). HDX-MS probes protein conformation and conformational dynamics in solution by monitoring the rate and extent of deuterium exchange of backbone amide hydrogen atoms. Huang & Chen (2014) *Anal. Bioanalytical Chem.* 406:6541; Wei et al. (2014) *Drug Disc. Today* 19:95. The level of HDX depends on the solvent accessibility of backbone amide hydrogen atoms and the protein hydrogen bonds. The mass increase of the protein upon HDX can be precisely measured by MS. When this technique is paired with enzymatic digestion, structure features at the peptide level can be resolved, enabling differentiation of surface exposed peptides from those folded inside, or from those sequestered at the interface of a protein-protein complex. Typically, the deuterium labeling and subsequent quenching experiments are performed, followed by enzymatic digestion, peptide separation, and MS analysis.

[0333] Prior to epitope mapping experiments, non-deuteriated experiments were carried out to generate a list of common peptides for recombinant human CD40 monomer (10 μM), and protein complexes of CD40 with mAbs 12D6, 5G7, 19G3, and 5F11 (1:1 molar ratio). In the HDX-MS experiment, 5 μL of each sample (CD40 or CD40 with mAbs 12D6, 5G7, 19G3, and 5F11 respectively) was diluted into 55 μL of D20 buffer (10 mM phosphate buffer, D20, pH7.0) to start the labeling reactions. The reactions were carried out for different periods of time: 20 sec, 1 min, 10 min and 240 min. By the end of each labeling reaction period, the reaction was quenched by adding quenching buffer (100 mM phosphate buffer with 4M GdnCl and 0.4M TCEP, pH 2.5, 1:1, v/v) and 50 μL of quenched sample was injected into Waters HDX-MS system for analysis. The deuterium uptake levels of common peptic peptides were monitored in the absence/presence of CD40 mAbs. The obtained sequence coverage was 82%.

[0334] The HDX epitopes for anti-CD40 mAbs 12D6, 5G7, 19G3, and 5F11 are provided at Table 7. Antibody 12D6 protected two peptides, one of which (residues 11-28) includes a portion of the signal sequence and is thus unlikely to be physiologically relevant per se, but indicates that 12D6 makes contacts in the region 21-28 that are not made by mAbs 5G7 and 19G3.

TABLE 7

HDX Epitopes		
Clone	CD40 residues (SEQ ID NO: 1)	Sequence
12D6	11-28	WGCLLTAVHPEPPTACRE
12D6	21-35	EPPTACREKQYLINS
5G7	21-35	EPPTACREKQYLINS
19G3	21-35	EPPTACREKQYLINS
5F11	58-66	ECLPCGESE

Example 6

Epitope Mapping by Yeast Display

[0335] The epitopes for selected chimeric or humanized anti-huCD40 antibodies of the present invention are determined by displaying randomly mutagenized huCD40 extracellular region variants on yeast, and sorting these yeast based on their failure to bind to particular antibodies. Selected yeast cells that fail to bind are amplified and subjected to additional rounds of selection based on their inability to bind to particular chimeric or humanized forms of the antibodies of the present invention. See, e.g., Chao et al. (2004) *J. Mol. Biol.* 342:539. Sequences for huCD40 variants are determined for the resulting yeast and analyzed for the effects of each residue on antibody binding. The binding epitope for the antibodies of the present invention is determined as the loci within the huCD40 sequence where single amino acid mutations disrupt binding to the anti-huCD40 antibodies of the present invention.

[0336] Briefly, error-prone PCR is used to clone human CD40-encoding DNA into constructs allowing expression of the huCD40 variants as the amino-terminal portions of fusion proteins further comprising a c-myc tag sequence and yeast cell wall protein Aga1p. Such constructs, when expressed in yeast (*Saccharomyces cerevisiae*), display the variant huCD40 polypeptides on the surface of yeast cells, anchored to the cell surface by the Aga1p polypeptide. The c-myc tag can optionally be used as a positive control for display of huCD40 fusion proteins on a given yeast cell. Yeast cells are sorted by FACS, and those that express as properly folded huCD40-fusion proteins (as determined by binding of a control mouse anti-huCD40 antibody detected by an allophycocyanin (APC)-labeled goat anti-mouse IgG secondary), but do not bind to the antibodies of the present invention (as determined by detection with a phycoerythrin (PE) labeled goat anti-human IgG as a secondary), are pooled, amplified, and used in subsequent rounds of selection. The huCD40 sequence is determined for constructs from yeast remaining after several rounds of selection. Control experiments without anti-huCD40 antibody selection are performed to confirm good mutant coverage at each position along the huCD40 sequence, and provide a baseline for normalizing the results obtained with the selected libraries.

Example 7

Agonist Activity of Anti-CD40 Antibodies

[0337] The effect of Fc sequence variation on the agonist activity of selected anti-CD40 antibodies of the present invention was assessed by measuring activation of immature dendritic cells (DC). Experiments were performed with anti-CD40 mAb 12D6-24 constructs having human IgG1f (control), SE, SELF, P238D, V4, V8 and V12 Fc sequences. Human Monocytes (CD14⁺) were isolated from healthy normal donors using plastic adherence or human CD14-micro beads (MILTENYI Biotec). Monocytes were cultured with 100 ng/mL GM-CSF (Miltenyi Biotec) and 100 ng/mL IL-4 (Miltenyi Biotec). Half of the medium was removed and replenished on day 2 and day 5. Immature dendritic cells were harvested at day 6-7. DC from two donors were incubated with the indicated concentration of antibodies

overnight at 37° C. Cell culture supernatants were collected and assayed for human-IL-6 production (CISBIO). See FIGS. 3A and 3B.

[0338] Control human IgG1f antibodies did not induce IL-6 secretion when used at up to 100 nM, and the P238D variant induced only weakly. In contrast, SE, SELF, V9 and V12 variants all dramatically enhanced IL-6 secretion, with V8 and V4 exhibiting intermediate effects. These results roughly correlate with binding affinity for FcγRIIb, and confirm that use of the proper Fc sequence variant can result in anti-CD40 antibodies with enhanced agonist activity.

[0339] In addition, the differences in agonist activity between antibodies having 8E8, 5G7, 12D6, 19G3 and 5F11 variable regions were evaluated in experiments using chimeric anti-CD40 mAb constructs having a common human IgG1f V12 constant domain. Activation was measured on immature dendritic cells in vitro (isolated as described in the preceding paragraph) by plating cells in a 96 well plate, adding antibodies as indicated, and incubating overnight at 37° C. Cells were then harvested and stained with a fluorescent anti-CD54 antibody, which was detected by fluorescence activated cell sorting (FACS). See FIG. 4.

[0340] Control human IgG1f antibody did not significantly cause CD54 upregulation, whereas 12D6 caused dramatic CD54 upregulation. Antibodies 19G3 and 8G8 were somewhat less effective than 12D6, and antibodies 5G7 and 5F11 were similar to each other and exhibited relatively low stimulation. The activation results do not necessarily correlate with binding affinity for CD40, since antibody 5F11 has nearly the highest binding affinity and yet is a weak agonist, whereas antibody 19G3 is the opposite. Regardless of the reason for the differences, these results confirm that use of the proper antigen binding domain sequence is important in obtaining anti-CD40 antibodies with enhanced agonist activity.

Example 8

Remediation of DG Isomerization

[0341] Antibodies of the present invention having V4 and V9 Fc variant sequences exhibit unacceptable levels of DG isomerization at position D270. Such isomerization is undesirable because it leads to product heterogeneity and potentially reduced efficacy. To reduce such isomerization, the aspartic acid residue at position 270 of V4 Fc was changed to glutamic acid (D270E), and position 270 of V9 Fc was changed to alanine, glutamic acid, glutamine, serine and threonine (D270A, D270E, D270Q, D270S, D270T). Supernatants were obtained from cells expressing these antibodies and assayed for binding to hCD32/FcγRII. Selected purified antibodies without mutations at position 270 were used as controls. These experiments were performed on antibodies with variable domains from mAbs 12D6-24 and 5F11-45. Results are provided at FIG. 5.

[0342] For both V4 and V9 Fc variants, the D270E substitution led to a modest decrease in receptor binding. Other V9 Fc variants tested (D270A, D270Q, D270S, D270T) led to a greater decrease in receptor binding. Results were independent of whether the antibody was 12D6 or 5F11. In all cases the relative (rank order) binding to the three receptors tested (FcγRIIa-H131, FcγRIIa-R131, and FcγRIIb) remained similar, with roughly equivalent binding to FcγRIIa-R131 and FcγRIIb, and significantly weaker binding to FcγRIIa-H131.

[0343] For both V4 and V9 variants, the D270E substitution retained acceptably high FcγR affinity, and maintained the specificity for FcγRIIb that is desired for enhancing the agonistic activity of the anti-CD40 antibodies of the present invention.

Example 9

Anti-CD40 Activity in Human CD40/Human FcγR Transgenic Mice

[0344] Selected agonist anti-CD40 antibodies of the present invention were administered to mice transgenic for human CD40 and human Fcγ receptors [CD40^{+/+}hCD40⁺Fcgra^{-/-}Fcgr1^{-/-}hFCGR1⁺hFCGRIIA⁺hFCGRIIB⁺hFCGRIIIA⁺hFCGRIIB⁺]. Mice transgenic for human Fcγ receptors are described at Smith et al. (2012) *Proc. Nat'l Acad. Sci. (USA)* 109(16):6181-6. Such transgenic human CD40/FcγR mice are well suited to study potential human therapeutic anti-CD40 antibodies, which necessarily bind to human CD40 and human FcγRs. Agonist activity was measured in the "DEC-OVA" model described in Li & Ravetch (2011) *Science* 333:1030, and platelet levels were also measured. Results are shown in FIGS. 6A and 6B. Antibodies showing the greatest agonist activity, such as 12D6-V11 and to a lesser degree 12D6-V4, also exhibited the greatest decrease in platelet count at 24 hours. The decrease in platelet count was a transient phenomenon, with levels returning to normal by 7 days post-injection (data not shown).

[0345] 5F11 antibody recognizes epitopes overlapping with the CD40L binding epitope, unlike 12D6 antibody that binds hCD40 distinct from its CD40L binding epitope, similar to CP-870,893. 12D6 antibody blocks CP-870,893 binding, but 5F11 antibody does not (data not shown). In vitro DC activation assays have demonstrated significant activity for the SE-Fc, and V11-Fc variants of 12D6 while 12D6 Fc variants exhibit enhanced activity compared to 5F11 variants. Although the in vitro activity of the 12D6-Fc variants is greater than the wild type IgG1, the in vitro studies do not distinguish between the activities of the different Fc variants. We therefore evaluated the activity of these antibodies and their Fc variants in the humanized CD40/human FcγR mice. As was observed for clone CP-870,893, the selective FcγRIIb-enhanced V11 Fc variant of these mAbs displayed superior activity in vivo compared to its FcγRIIa/IIb-enhanced SE variant and the wild type IgG1 subclass. Moreover, the magnitude of the in vivo activity of the newly developed 12D6-V11 is about 20-fold higher than the activity of CP-870,893 (IgG2).

[0346] The anti tumor activity of the CD40 agonistic Abs, 12D6-24 and 5F11-45 was evaluated. Wild type IgG1, SE and V11 mutants of each clone were tested for their activity in the MC38 tumor model. The FcγRIIb-selectively enhanced V11 variants of these antibodies have superior anti tumor activity compared to the other tested Fc variants, consistent with their in vivo T cell stimulation activity and with the anti tumor activity hierarchy observed for CP-870,893 derivatives. Results are shown in FIG. 6C. Tumor-free mice from the 12D6-24 treated group were re-challenged with MC38 tumor cells 30 days after their CD40 Ab treatment. Results are shown in FIG. 6D. All mice in this group rejected the tumors within two weeks of the re-challenge, demonstrating that long-term protection is mediated by single injection of 12D6-24.

[0347] These data indicate that the antitumor activity of agonistic human CD40 Abs can be enhanced by Fc engineering that provides selective enhancement of FcγRIIB-engagement and that the V11 Fc variant is particularly effective at enhancing agonism.

TABLE 8

Summary of Sequence Listing	
SEQ ID	Description
1	Human CD40 (NP_001241)
2	Human CD40L-gp39 (NP_000065.1)
3	12D6 Chimeric Heavy Chain
4	12D6 Chimeric Light Chain
5	12D6-03 Heavy Chain
6	12D6-03 Light Chain
7	12D6-22 Heavy Chain
8	12D6-22 V9 Heavy Chain
9	12D6-22/12D6-24 Light Chain
10	12D6-23 Heavy Chain
11	12D6-23 Light Chain
12	12D6-24 Heavy Chain
13	12D6-24 P238D Heavy Chain
14	12D6-24 SE Heavy Chain
15	12D6-24 SELF Heavy Chain
16	12D6-24 V4 Heavy Chain
17	12D6-24 V4 D270E Heavy Chain
18	12D6-24 V8 Heavy Chain
19	12D6-24 V9 Heavy Chain
20	12D6-24 V9 D270E Heavy Chain
21	12D6-24 V11 Heavy Chain
22	12D6-24 V12 Heavy Chain
23	5F11 Chimeric Heavy Chain
24	5F11 Chimeric Light Chain
25	5F11-17 Heavy Chain
26	5F11-17 Light Chain
27	5F11-23 Heavy Chain
28	5F11-23 Light Chain
29	5F11-45 Heavy Chain
30	5F11-45 Light Chain
31	5F11-45 SE Heavy Chain
32	5F11-45 SELF Heavy Chain
33	5F11-45 V4 Heavy Chain
34	5F11-45 D270E V4 Heavy Chain
35	5F11-45 V8 Heavy Chain
36	5F11-45 V9 Heavy Chain
37	5F11-45 V9 D270E Heavy Chain
38	5F11-45 V11 Heavy Chain
39	5F11-45 V12 Heavy Chain
40	8E8 Chimeric Heavy Chain
41	8E8 Chimeric Light Chain
42	8E8-56 Heavy Chain
43	8E8-56 Light Chain

TABLE 8-continued

Summary of Sequence Listing	
SEQ ID	Description
44	8E8-62 Heavy Chain
45	8E8-62 Light Chain
46	8E8-67 Heavy Chain
47	8E8-67 Light Chain
48	8E8-70 Heavy Chain
49	8E8-70 Light Chain
50	8E8-71 Heavy Chain
51	8E8-71 Light Chain
52	5G7 Chimeric Heavy Chain
53	5G7 Chimeric Light Chain
54	5G7-22 Heavy Chain
55	5G7-22 Light Chain
56	5G7-25 Heavy Chain
57	5G7-25 Light Chain
58	19G3 Chimeric Heavy Chain
59	19G3 Chimeric Light Chain
60	19G3-11 Heavy Chain
61	19G3-11 V9
62	19G3-11 Light Chain
63	19G3-22 Heavy Chain
64	19G3-22 Light Chain
65	Human Constant Domain IgG1f
66	SE Constant Domain
67	SELF Constant Domain
68	P238D Constant Domain
69	V4 Constant Domain
70	V4 D270E Constant Domain
71	V7 Constant Domain
72	V8 Constant Domain
73	V9 Constant Domain
74	V9 D270E Constant Domain
75	V11 Constant Domain
76	V12 Constant Domain
77	Light Chain Kappa Constant Domain
78	Signal Sequence

[0348] The Sequence Listing provides the sequences of the mature heavy and light chains (i.e., sequences do not include signal peptides). A signal sequence for production of the antibodies of the present invention, for example in human cells, is provided at SEQ ID NO: 78.

EQUIVALENTS

[0349] Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents of the specific embodiments disclosed herein. Such equivalents are intended to be encompassed by the following claims.

SEQUENCE LISTING

```

<160> NUMBER OF SEQ ID NOS: 78

<210> SEQ ID NO 1
<211> LENGTH: 277
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: SIGNAL
<222> LOCATION: (1) .. (20)
<220> FEATURE:
<221> NAME/KEY: PEPTIDE
<222> LOCATION: (21) .. (277)
<220> FEATURE:
<221> NAME/KEY: DOMAIN
<222> LOCATION: (21) .. (193)
<223> OTHER INFORMATION: extracellular domain

```

-continued

```

<220> FEATURE:
<221> NAME/KEY: DOMAIN
<222> LOCATION: (194)..(215)
<223> OTHER INFORMATION: transmembrane domain
<220> FEATURE:
<221> NAME/KEY: DOMAIN
<222> LOCATION: (216)..(277)
<223> OTHER INFORMATION: intracellular domain

<400> SEQUENCE: 1

Met Val Arg Leu Pro Leu Gln Cys Val Leu Trp Gly Cys Leu Leu Thr
1          5          10          15

Ala Val His Pro Glu Pro Pro Thr Ala Cys Arg Glu Lys Gln Tyr Leu
20        25        30

Ile Asn Ser Gln Cys Cys Ser Leu Cys Gln Pro Gly Gln Lys Leu Val
35        40        45

Ser Asp Cys Thr Glu Phe Thr Glu Thr Glu Cys Leu Pro Cys Gly Glu
50        55        60

Ser Glu Phe Leu Asp Thr Trp Asn Arg Glu Thr His Cys His Gln His
65        70        75        80

Lys Tyr Cys Asp Pro Asn Leu Gly Leu Arg Val Gln Gln Lys Gly Thr
85        90        95

Ser Glu Thr Asp Thr Ile Cys Thr Cys Glu Glu Gly Trp His Cys Thr
100       105       110

Ser Glu Ala Cys Glu Ser Cys Val Leu His Arg Ser Cys Ser Pro Gly
115       120       125

Phe Gly Val Lys Gln Ile Ala Thr Gly Val Ser Asp Thr Ile Cys Glu
130       135       140

Pro Cys Pro Val Gly Phe Phe Ser Asn Val Ser Ser Ala Phe Glu Lys
145       150       155       160

Cys His Pro Trp Thr Ser Cys Glu Thr Lys Asp Leu Val Val Gln Gln
165       170       175

Ala Gly Thr Asn Lys Thr Asp Val Val Cys Gly Pro Gln Asp Arg Leu
180       185       190

Arg Ala Leu Val Val Ile Pro Ile Ile Phe Gly Ile Leu Phe Ala Ile
195       200       205

Leu Leu Val Leu Val Phe Ile Lys Lys Val Ala Lys Lys Pro Thr Asn
210       215       220

Lys Ala Pro His Pro Lys Gln Glu Pro Gln Glu Ile Asn Phe Pro Asp
225       230       235       240

Asp Leu Pro Gly Ser Asn Thr Ala Ala Pro Val Gln Glu Thr Leu His
245       250       255

Gly Cys Gln Pro Val Thr Gln Glu Asp Gly Lys Glu Ser Arg Ile Ser
260       265       270

Val Gln Glu Arg Gln
275

<210> SEQ ID NO 2
<211> LENGTH: 261
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 2

Met Ile Glu Thr Tyr Asn Gln Thr Ser Pro Arg Ser Ala Ala Thr Gly
1          5          10          15

```

-continued

```

Leu Pro Ile Ser Met Lys Ile Phe Met Tyr Leu Leu Thr Val Phe Leu
      20                25                30

Ile Thr Gln Met Ile Gly Ser Ala Leu Phe Ala Val Tyr Leu His Arg
      35                40                45

Arg Leu Asp Lys Ile Glu Asp Glu Arg Asn Leu His Glu Asp Phe Val
      50                55                60

Phe Met Lys Thr Ile Gln Arg Cys Asn Thr Gly Glu Arg Ser Leu Ser
      65                70                75                80

Leu Leu Asn Cys Glu Glu Ile Lys Ser Gln Phe Glu Gly Phe Val Lys
      85                90                95

Asp Ile Met Leu Asn Lys Glu Glu Thr Lys Lys Glu Asn Ser Phe Glu
      100               105               110

Met Gln Lys Gly Asp Gln Asn Pro Gln Ile Ala Ala His Val Ile Ser
      115               120               125

Glu Ala Ser Ser Lys Thr Thr Ser Val Leu Gln Trp Ala Glu Lys Gly
      130               135               140

Tyr Tyr Thr Met Ser Asn Asn Leu Val Thr Leu Glu Asn Gly Lys Gln
      145               150               155               160

Leu Thr Val Lys Arg Gln Gly Leu Tyr Tyr Ile Tyr Ala Gln Val Thr
      165               170               175

Phe Cys Ser Asn Arg Glu Ala Ser Ser Gln Ala Pro Phe Ile Ala Ser
      180               185               190

Leu Cys Leu Lys Ser Pro Gly Arg Phe Glu Arg Ile Leu Leu Arg Ala
      195               200               205

Ala Asn Thr His Ser Ser Ala Lys Pro Cys Gly Gln Gln Ser Ile His
      210               215               220

Leu Gly Gly Val Phe Glu Leu Gln Pro Gly Ala Ser Val Phe Val Asn
      225               230               235               240

Val Thr Asp Pro Ser Gln Val Ser His Gly Thr Gly Phe Thr Ser Phe
      245               250               255

Gly Leu Leu Lys Leu
      260

```

```

<210> SEQ ID NO 3
<211> LENGTH: 448
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: chimeric antibody with mouse variable domain,
human constant domain
<220> FEATURE:
<221> NAME/KEY: DOMAIN
<222> LOCATION: (1)..(119)
<223> OTHER INFORMATION: variable domain
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (31)..(35)
<223> OTHER INFORMATION: CDRH1
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (50)..(66)
<223> OTHER INFORMATION: CDRH2
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (99)..(108)
<223> OTHER INFORMATION: CDRH3

<400> SEQUENCE: 3

```

```

Glu Val Gln Leu Gln Gln Ser Gly Pro Glu Leu Glu Lys Pro Gly Ala

```

-continued

1	5	10	15
Ser Val Lys Met Ser Cys Lys Ala Ser Gly Tyr Ser Phe Thr Gly Tyr	20	25	30
Asn Met Asn Trp Val Lys Gln Ser Asn Gly Lys Ser Leu Glu Trp Ile	35	40	45
Gly Asn Ile Asp Pro Tyr Tyr Gly Asn Thr Asn Tyr Asn Gln Lys Phe	50	55	60
Lys Gly Lys Ala Thr Leu Thr Val Asp Lys Ser Ser Ser Thr Ala Tyr	65	70	75
Leu Gln Leu Lys Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys	85	90	95
Ala Arg Leu Gly Leu Gln Leu Tyr Ala Met Asp Tyr Trp Gly Gln Gly	100	105	110
Thr Ser Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe	115	120	125
Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala Leu	130	135	140
Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp	145	150	155
Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu	165	170	175
Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser	180	185	190
Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys Pro	195	200	205
Ser Asn Thr Lys Val Asp Lys Arg Val Glu Pro Lys Ser Cys Asp Lys	210	215	220
Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly Pro	225	230	235
Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser	245	250	255
Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu Asp	260	265	270
Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn	275	280	285
Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val	290	295	300
Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu	305	310	315
Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu Lys	325	330	335
Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr	340	345	350
Leu Pro Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr	355	360	365
Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu	370	375	380
Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu	385	390	395
Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys	405	410	415

-continued

Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu
420 425 430

Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly
435 440 445

<210> SEQ ID NO 4
 <211> LENGTH: 219
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: chimeric antibody with mouse variable domain
 and human constant domain
 <220> FEATURE:
 <221> NAME/KEY: DOMAIN
 <222> LOCATION: (1)..(112)
 <223> OTHER INFORMATION: variable domain
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (24)..(39)
 <223> OTHER INFORMATION: CDRL1
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (55)..(61)
 <223> OTHER INFORMATION: CDRL2
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (94)..(102)
 <223> OTHER INFORMATION: CDRL3

<400> SEQUENCE: 4

Asp Val Val Met Thr Gln Thr Pro Leu Ser Leu Pro Val Ser Leu Gly
1 5 10 15

Asp Gln Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Leu Val His Ser
20 25 30

Asn Gly Asn Thr Tyr Leu His Trp Tyr Leu Gln Lys Pro Gly Gln Ser
35 40 45

Pro Asn Leu Leu Ile Tyr Lys Leu Thr Asn Arg Phe Phe Gly Val Pro
50 55 60

Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile
65 70 75 80

Ser Arg Val Glu Ala Glu Asp Leu Gly Val Tyr Phe Cys Ser Gln Ser
85 90 95

Ile His Val Pro Trp Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys
100 105 110

Arg Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu
115 120 125

Gln Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe
130 135 140

Tyr Pro Arg Glu Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln
145 150 155 160

Ser Gly Asn Ser Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser
165 170 175

Thr Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu
180 185 190

Lys His Lys Val Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser
195 200 205

Pro Val Thr Lys Ser Phe Asn Arg Gly Glu Cys
210 215

-continued

<210> SEQ ID NO 5
<211> LENGTH: 448
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: humanized antibody with mouse CDRs and human framework and constant regions
<220> FEATURE:
<221> NAME/KEY: DOMAIN
<222> LOCATION: (1)..(119)
<223> OTHER INFORMATION: variable domain
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (31)..(35)
<223> OTHER INFORMATION: CDRH1
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (50)..(66)
<223> OTHER INFORMATION: CDRH2
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (99)..(108)
<223> OTHER INFORMATION: CDRH3

<400> SEQUENCE: 5

Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala
1 5 10 15

Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Gly Tyr
20 25 30

Asn Met Asn Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met
35 40 45

Gly Asn Ile Asp Pro Tyr Tyr Gly Asn Thr Asn Tyr Asn Gln Lys Phe
50 55 60

Lys Gly Arg Val Thr Met Thr Thr Asp Thr Ser Thr Ser Thr Ala Tyr
65 70 75 80

Met Glu Leu Arg Ser Leu Arg Ser Asp Asp Thr Ala Val Tyr Tyr Cys
85 90 95

Ala Arg Leu Gly Leu Gln Leu Tyr Ala Met Asp Tyr Trp Gly Gln Gly
100 105 110

Thr Thr Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe
115 120 125

Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala Leu
130 135 140

Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp
145 150 155 160

Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu
165 170 175

Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser
180 185 190

Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys Pro
195 200 205

Ser Asn Thr Lys Val Asp Lys Arg Val Glu Pro Lys Ser Cys Asp Lys
210 215 220

Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly Pro
225 230 235 240

Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser
245 250 255

Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu Asp

-continued

260	265	270
Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn		
275	280	285
Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val		
290	295	300
Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu		
305	310	315
Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu Lys		
325	330	335
Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr		
340	345	350
Leu Pro Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr		
355	360	365
Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu		
370	375	380
Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu		
385	390	395
Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys		
405	410	415
Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu		
420	425	430
Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly		
435	440	445

<210> SEQ ID NO 6
 <211> LENGTH: 219
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: humanized antibody with mouse CDRs and human framework and constant regions
 <220> FEATURE:
 <221> NAME/KEY: DOMAIN
 <222> LOCATION: (1)..(112)
 <223> OTHER INFORMATION: variable domain
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (24)..(39)
 <223> OTHER INFORMATION: CDRL1
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (55)..(61)
 <223> OTHER INFORMATION: CDRL2
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (94)..(102)
 <223> OTHER INFORMATION: CDRL3

<400> SEQUENCE: 6

Asp Ile Val Met Thr Gln Thr Pro Leu Ser Leu Pro Val Thr Pro Gly
1 5 10 15
Glu Pro Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Leu Val His Ser
20 25 30
Asn Gly Asn Thr Tyr Leu His Trp Tyr Leu Gln Lys Pro Gly Gln Ser
35 40 45
Pro Gln Leu Leu Ile Tyr Lys Leu Thr Asn Arg Phe Phe Gly Val Pro
50 55 60
Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile
65 70 75 80

-continued

```

Ser Arg Val Glu Ala Glu Asp Val Gly Val Tyr Tyr Cys Ser Gln Ser
      85                      90                      95
Ile His Val Pro Trp Thr Phe Gly Gly Gly Thr Lys Val Glu Ile Lys
      100                      105                      110
Arg Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu
      115                      120                      125
Gln Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe
      130                      135                      140
Tyr Pro Arg Glu Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln
      145                      150                      155                      160
Ser Gly Asn Ser Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser
      165                      170                      175
Thr Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu
      180                      185                      190
Lys His Lys Val Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser
      195                      200                      205
Pro Val Thr Lys Ser Phe Asn Arg Gly Glu Cys
      210                      215

```

```

<210> SEQ ID NO 7
<211> LENGTH: 448
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: humanized antibody with mouse CDRs and human
      framework and constant regions
<220> FEATURE:
<221> NAME/KEY: DOMAIN
<222> LOCATION: (1)..(119)
<223> OTHER INFORMATION: variable domain
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (31)..(35)
<223> OTHER INFORMATION: CDRH1
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (50)..(66)
<223> OTHER INFORMATION: CDRH2
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (99)..(108)
<223> OTHER INFORMATION: CDRH3

```

```

<400> SEQUENCE: 7

```

```

Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala
1      5      10      15
Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Gly Tyr
20     25     30
Asn Met Asn Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met
35     40     45
Gly Asn Ile Asp Pro Tyr Tyr Gly Asn Thr Asn Tyr Asn Gln Lys Phe
50     55     60
Lys Gly Arg Val Thr Met Thr Thr Asp Thr Ser Thr Ser Thr Ala Tyr
65     70     75     80
Met Glu Leu Arg Ser Leu Arg Ser Asp Asp Thr Ala Val Tyr Tyr Cys
85     90     95
Ala Arg Leu Gly Leu Gln Leu Tyr Ala Met Asp Tyr Trp Gly Gln Gly
100    105    110

```

-continued

Thr	Thr	Val	Thr	Val	Ser	Ser	Ala	Ser	Thr	Lys	Gly	Pro	Ser	Val	Phe
		115					120					125			
Pro	Leu	Ala	Pro	Ser	Ser	Lys	Ser	Thr	Ser	Gly	Gly	Thr	Ala	Ala	Leu
	130					135					140				
Gly	Cys	Leu	Val	Lys	Asp	Tyr	Phe	Pro	Glu	Pro	Val	Thr	Val	Ser	Trp
145					150					155					160
Asn	Ser	Gly	Ala	Leu	Thr	Ser	Gly	Val	His	Thr	Phe	Pro	Ala	Val	Leu
				165					170					175	
Gln	Ser	Ser	Gly	Leu	Tyr	Ser	Leu	Ser	Ser	Val	Val	Thr	Val	Pro	Ser
			180					185					190		
Ser	Ser	Leu	Gly	Thr	Gln	Thr	Tyr	Ile	Cys	Asn	Val	Asn	His	Lys	Pro
		195					200					205			
Ser	Asn	Thr	Lys	Val	Asp	Lys	Arg	Val	Glu	Pro	Lys	Ser	Cys	Asp	Lys
	210					215					220				
Thr	His	Thr	Cys	Pro	Pro	Cys	Pro	Ala	Pro	Glu	Leu	Leu	Gly	Gly	Pro
225					230					235					240
Ser	Val	Phe	Leu	Phe	Pro	Pro	Lys	Pro	Lys	Asp	Thr	Leu	Met	Ile	Ser
				245					250					255	
Arg	Thr	Pro	Glu	Val	Thr	Cys	Val	Val	Val	Asp	Val	Ser	His	Glu	Asp
			260					265					270		
Pro	Glu	Val	Lys	Phe	Asn	Trp	Tyr	Val	Asp	Gly	Val	Glu	Val	His	Asn
		275					280					285			
Ala	Lys	Thr	Lys	Pro	Arg	Glu	Glu	Gln	Tyr	Asn	Ser	Thr	Tyr	Arg	Val
	290					295					300				
Val	Ser	Val	Leu	Thr	Val	Leu	His	Gln	Asp	Trp	Leu	Asn	Gly	Lys	Glu
305					310					315					320
Tyr	Lys	Cys	Lys	Val	Ser	Asn	Lys	Ala	Leu	Pro	Ala	Pro	Ile	Glu	Lys
				325					330					335	
Thr	Ile	Ser	Lys	Ala	Lys	Gly	Gln	Pro	Arg	Glu	Pro	Gln	Val	Tyr	Thr
			340					345					350		
Leu	Pro	Pro	Ser	Arg	Glu	Glu	Met	Thr	Lys	Asn	Gln	Val	Ser	Leu	Thr
		355					360					365			
Cys	Leu	Val	Lys	Gly	Phe	Tyr	Pro	Ser	Asp	Ile	Ala	Val	Glu	Trp	Glu
	370					375					380				
Ser	Asn	Gly	Gln	Pro	Glu	Asn	Asn	Tyr	Lys	Thr	Thr	Pro	Pro	Val	Leu
385					390					395					400
Asp	Ser	Asp	Gly	Ser	Phe	Phe	Leu	Tyr	Ser	Lys	Leu	Thr	Val	Asp	Lys
				405					410					415	
Ser	Arg	Trp	Gln	Gln	Gly	Asn	Val	Phe	Ser	Cys	Ser	Val	Met	His	Glu
			420					425					430		
Ala	Leu	His	Asn	His	Tyr	Thr	Gln	Lys	Ser	Leu	Ser	Leu	Ser	Pro	Gly
		435					440					445			

<210> SEQ ID NO 8

<211> LENGTH: 448

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: humanized antibody with mouse CDRs and human framework and constant regions

<220> FEATURE:

<221> NAME/KEY: DOMAIN

<222> LOCATION: (1)..(119)

<223> OTHER INFORMATION: variable domain

<220> FEATURE:

-continued

```

<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (31)..(35)
<223> OTHER INFORMATION: CDRH1
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (50)..(66)
<223> OTHER INFORMATION: CDRH2
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (99)..(108)
<223> OTHER INFORMATION: CDRH3

<400> SEQUENCE: 8

Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala
1          5          10          15

Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Gly Tyr
20          25          30

Asn Met Asn Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met
35          40          45

Gly Asn Ile Asp Pro Tyr Tyr Gly Asn Thr Asn Tyr Asn Gln Lys Phe
50          55          60

Lys Gly Arg Val Thr Met Thr Thr Asp Thr Ser Thr Ser Thr Ala Tyr
65          70          75          80

Met Glu Leu Arg Ser Leu Arg Ser Asp Asp Thr Ala Val Tyr Tyr Cys
85          90          95

Ala Arg Leu Gly Leu Gln Leu Tyr Ala Met Asp Tyr Trp Gly Gln Gly
100         105         110

Thr Thr Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe
115         120         125

Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala Leu
130         135         140

Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp
145         150         155         160

Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu
165         170         175

Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser
180         185         190

Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys Pro
195         200         205

Ser Asn Thr Lys Val Asp Lys Arg Val Glu Pro Lys Ser Cys Asp Lys
210         215         220

Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Asp Asp
225         230         235         240

Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser
245         250         255

Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu Asp
260         265         270

Gly Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn
275         280         285

Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val
290         295         300

Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu
305         310         315         320

Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Arg Pro Ile Glu Lys
325         330         335

```

-continued

Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr
 340 345 350
 Leu Pro Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr
 355 360 365
 Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu
 370 375 380
 Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu
 385 390 395 400
 Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys
 405 410 415
 Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu
 420 425 430
 Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly
 435 440 445

<210> SEQ ID NO 9
 <211> LENGTH: 219
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: humanized antibody with mouse CDRs and human
 framework and constant regions
 <220> FEATURE:
 <221> NAME/KEY: DOMAIN
 <222> LOCATION: (1)..(112)
 <223> OTHER INFORMATION: variable domain
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (24)..(39)
 <223> OTHER INFORMATION: CDRL1
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (55)..(61)
 <223> OTHER INFORMATION: CDRL2
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (94)..(102)
 <223> OTHER INFORMATION: CDRL3

<400> SEQUENCE: 9

Asp Ile Val Met Thr Gln Thr Pro Leu Ser Leu Pro Val Thr Pro Gly
 1 5 10 15
 Glu Pro Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Leu Val His Ser
 20 25 30
 Asn Ala Asn Thr Tyr Leu His Trp Tyr Leu Gln Lys Pro Gly Gln Ser
 35 40 45
 Pro Gln Leu Leu Ile Tyr Lys Leu Thr Asn Arg Phe Phe Gly Val Pro
 50 55 60
 Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile
 65 70 75 80
 Ser Arg Val Glu Ala Glu Asp Val Gly Val Tyr Tyr Cys Ser Gln Ser
 85 90 95
 Ile His Val Pro Trp Thr Phe Gly Gly Gly Thr Lys Val Glu Ile Lys
 100 105 110
 Arg Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu
 115 120 125
 Gln Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe
 130 135 140

-continued

```

Tyr Pro Arg Glu Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln
145                      150                      155                      160

Ser Gly Asn Ser Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser
                      165                      170                      175

Thr Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu
                      180                      185                      190

Lys His Lys Val Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser
                      195                      200                      205

Pro Val Thr Lys Ser Phe Asn Arg Gly Glu Cys
210                      215

```

```

<210> SEQ ID NO 10
<211> LENGTH: 448
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: humanized antibody with mouse CDRs and human
framework and constant regions
<220> FEATURE:
<221> NAME/KEY: DOMAIN
<222> LOCATION: (1)..(119)
<223> OTHER INFORMATION: variable domain
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (31)..(35)
<223> OTHER INFORMATION: CDRH1
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (50)..(66)
<223> OTHER INFORMATION: CDRH2
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (99)..(108)
<223> OTHER INFORMATION: CDRH3

```

```

<400> SEQUENCE: 10

```

```

Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala
1                      5                      10                      15

Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Gly Tyr
20                      25                      30

Asn Met Asn Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met
35                      40                      45

Gly Asn Ile Asp Pro Tyr Tyr Gly Asn Thr Asn Tyr Asn Gln Lys Phe
50                      55                      60

Lys Gly Arg Val Thr Met Thr Thr Asp Thr Ser Thr Ser Thr Ala Tyr
65                      70                      75                      80

Met Glu Leu Arg Ser Leu Arg Ser Asp Asp Thr Ala Val Tyr Tyr Cys
85                      90                      95

Ala Arg Leu Gly Leu Gln Leu Tyr Ala Leu Asp Tyr Trp Gly Gln Gly
100                      105                      110

Thr Thr Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe
115                      120                      125

Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala Leu
130                      135                      140

Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp
145                      150                      155                      160

Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu
165                      170                      175

Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser

```

-continued

180					185					190					
Ser	Ser	Leu	Gly	Thr	Gln	Thr	Tyr	Ile	Cys	Asn	Val	Asn	His	Lys	Pro
	195						200					205			
Ser	Asn	Thr	Lys	Val	Asp	Lys	Arg	Val	Glu	Pro	Lys	Ser	Cys	Asp	Lys
	210					215					220				
Thr	His	Thr	Cys	Pro	Pro	Cys	Pro	Ala	Pro	Glu	Leu	Leu	Gly	Gly	Pro
	225					230					235				240
Ser	Val	Phe	Leu	Phe	Pro	Pro	Lys	Pro	Lys	Asp	Thr	Leu	Met	Ile	Ser
			245						250					255	
Arg	Thr	Pro	Glu	Val	Thr	Cys	Val	Val	Val	Asp	Val	Ser	His	Glu	Asp
			260					265					270		
Pro	Glu	Val	Lys	Phe	Asn	Trp	Tyr	Val	Asp	Gly	Val	Glu	Val	His	Asn
	275						280					285			
Ala	Lys	Thr	Lys	Pro	Arg	Glu	Glu	Gln	Tyr	Asn	Ser	Thr	Tyr	Arg	Val
	290					295					300				
Val	Ser	Val	Leu	Thr	Val	Leu	His	Gln	Asp	Trp	Leu	Asn	Gly	Lys	Glu
	305					310					315				320
Tyr	Lys	Cys	Lys	Val	Ser	Asn	Lys	Ala	Leu	Pro	Ala	Pro	Ile	Glu	Lys
			325						330					335	
Thr	Ile	Ser	Lys	Ala	Lys	Gly	Gln	Pro	Arg	Glu	Pro	Gln	Val	Tyr	Thr
			340					345					350		
Leu	Pro	Pro	Ser	Arg	Glu	Glu	Met	Thr	Lys	Asn	Gln	Val	Ser	Leu	Thr
	355						360					365			
Cys	Leu	Val	Lys	Gly	Phe	Tyr	Pro	Ser	Asp	Ile	Ala	Val	Glu	Trp	Glu
	370					375					380				
Ser	Asn	Gly	Gln	Pro	Glu	Asn	Asn	Tyr	Lys	Thr	Thr	Pro	Pro	Val	Leu
	385					390					395				400
Asp	Ser	Asp	Gly	Ser	Phe	Phe	Leu	Tyr	Ser	Lys	Leu	Thr	Val	Asp	Lys
			405						410					415	
Ser	Arg	Trp	Gln	Gln	Gly	Asn	Val	Phe	Ser	Cys	Ser	Val	Met	His	Glu
			420					425					430		
Ala	Leu	His	Asn	His	Tyr	Thr	Gln	Lys	Ser	Leu	Ser	Leu	Ser	Pro	Gly
	435						440					445			

<210> SEQ ID NO 11

<211> LENGTH: 219

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: humanized antibody with mouse CDRs and human framework and constant regions

<220> FEATURE:

<221> NAME/KEY: DOMAIN

<222> LOCATION: (1)..(112)

<223> OTHER INFORMATION: variable domain

<220> FEATURE:

<221> NAME/KEY: MISC_FEATURE

<222> LOCATION: (24)..(39)

<223> OTHER INFORMATION: CDRL1

<220> FEATURE:

<221> NAME/KEY: MISC_FEATURE

<222> LOCATION: (55)..(61)

<223> OTHER INFORMATION: CDRL2

<220> FEATURE:

<221> NAME/KEY: MISC_FEATURE

<222> LOCATION: (94)..(102)

<223> OTHER INFORMATION: CDRL3

<400> SEQUENCE: 11

-continued

```

Asp Ile Val Met Thr Gln Thr Pro Leu Ser Leu Pro Val Thr Pro Gly
1           5           10           15
Glu Pro Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Leu Val His Ser
20           25           30
Asn Gly Asn Thr Tyr Leu His Trp Tyr Leu Gln Lys Pro Gly Gln Ser
35           40           45
Pro Gln Leu Leu Ile Tyr Lys Leu Thr Asn Arg Phe Phe Gly Val Pro
50           55           60
Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile
65           70           75           80
Ser Arg Val Glu Ala Glu Asp Val Gly Val Tyr Tyr Cys Ser Gln Ser
85           90           95
Ile His Val Pro Trp Thr Phe Gly Gly Gly Thr Lys Val Glu Ile Lys
100          105          110
Arg Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu
115          120          125
Gln Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe
130          135          140
Tyr Pro Arg Glu Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln
145          150          155          160
Ser Gly Asn Ser Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser
165          170          175
Thr Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu
180          185          190
Lys His Lys Val Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser
195          200          205
Pro Val Thr Lys Ser Phe Asn Arg Gly Glu Cys
210          215

```

```

<210> SEQ ID NO 12
<211> LENGTH: 448
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: humanized antibody with mouse CDRs and human
framework and constant regions
<220> FEATURE:
<221> NAME/KEY: DOMAIN
<222> LOCATION: (1)..(119)
<223> OTHER INFORMATION: variable domain
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (31)..(35)
<223> OTHER INFORMATION: CDRH1
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (50)..(66)
<223> OTHER INFORMATION: CDRH2
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (99)..(108)
<223> OTHER INFORMATION: CDRH3
<400> SEQUENCE: 12

```

```

Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala
1           5           10           15
Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Gly Tyr
20           25           30

```

-continued

Asn	Met	Asn	Trp	Val	Arg	Gln	Ala	Pro	Gly	Gln	Gly	Leu	Glu	Trp	Met
	35						40					45			
Gly	Asn	Ile	Asp	Pro	Tyr	Tyr	Gly	Asn	Thr	Asn	Tyr	Asn	Gln	Lys	Phe
50						55					60				
Lys	Gly	Arg	Val	Thr	Met	Thr	Thr	Asp	Thr	Ser	Thr	Ser	Thr	Ala	Tyr
65					70					75					80
Met	Glu	Leu	Arg	Ser	Leu	Arg	Ser	Asp	Asp	Thr	Ala	Val	Tyr	Tyr	Cys
				85					90					95	
Ala	Arg	Leu	Gly	Leu	Gln	Leu	Tyr	Ala	Leu	Asp	Tyr	Trp	Gly	Gln	Gly
			100					105					110		
Thr	Thr	Val	Thr	Val	Ser	Ser	Ala	Ser	Thr	Lys	Gly	Pro	Ser	Val	Phe
		115					120					125			
Pro	Leu	Ala	Pro	Ser	Ser	Lys	Ser	Thr	Ser	Gly	Gly	Thr	Ala	Ala	Leu
	130					135					140				
Gly	Cys	Leu	Val	Lys	Asp	Tyr	Phe	Pro	Glu	Pro	Val	Thr	Val	Ser	Trp
145					150					155					160
Asn	Ser	Gly	Ala	Leu	Thr	Ser	Gly	Val	His	Thr	Phe	Pro	Ala	Val	Leu
				165					170					175	
Gln	Ser	Ser	Gly	Leu	Tyr	Ser	Leu	Ser	Ser	Val	Val	Thr	Val	Pro	Ser
			180					185					190		
Ser	Ser	Leu	Gly	Thr	Gln	Thr	Tyr	Ile	Cys	Asn	Val	Asn	His	Lys	Pro
		195					200					205			
Ser	Asn	Thr	Lys	Val	Asp	Lys	Arg	Val	Glu	Pro	Lys	Ser	Cys	Asp	Lys
	210					215					220				
Thr	His	Thr	Cys	Pro	Pro	Cys	Pro	Ala	Pro	Glu	Leu	Leu	Gly	Gly	Pro
225					230					235					240
Ser	Val	Phe	Leu	Phe	Pro	Pro	Lys	Pro	Lys	Asp	Thr	Leu	Met	Ile	Ser
				245					250					255	
Arg	Thr	Pro	Glu	Val	Thr	Cys	Val	Val	Val	Asp	Val	Ser	His	Glu	Asp
			260					265					270		
Pro	Glu	Val	Lys	Phe	Asn	Trp	Tyr	Val	Asp	Gly	Val	Glu	Val	His	Asn
		275					280					285			
Ala	Lys	Thr	Lys	Pro	Arg	Glu	Glu	Gln	Tyr	Asn	Ser	Thr	Tyr	Arg	Val
	290					295					300				
Val	Ser	Val	Leu	Thr	Val	Leu	His	Gln	Asp	Trp	Leu	Asn	Gly	Lys	Glu
305					310					315					320
Tyr	Lys	Cys	Lys	Val	Ser	Asn	Lys	Ala	Leu	Pro	Ala	Pro	Ile	Glu	Lys
				325					330					335	
Thr	Ile	Ser	Lys	Ala	Lys	Gly	Gln	Pro	Arg	Glu	Pro	Gln	Val	Tyr	Thr
			340					345					350		
Leu	Pro	Pro	Ser	Arg	Glu	Glu	Met	Thr	Lys	Asn	Gln	Val	Ser	Leu	Thr
		355					360					365			
Cys	Leu	Val	Lys	Gly	Phe	Tyr	Pro	Ser	Asp	Ile	Ala	Val	Glu	Trp	Glu
	370					375					380				
Ser	Asn	Gly	Gln	Pro	Glu	Asn	Asn	Tyr	Lys	Thr	Thr	Pro	Pro	Val	Leu
				390						395					400
Asp	Ser	Asp	Gly	Ser	Phe	Phe	Leu	Tyr	Ser	Lys	Leu	Thr	Val	Asp	Lys
				405					410					415	
Ser	Arg	Trp	Gln	Gln	Gly	Asn	Val	Phe	Ser	Cys	Ser	Val	Met	His	Glu
			420					425					430		
Ala	Leu	His	Asn	His	Tyr	Thr	Gln	Lys	Ser	Leu	Ser	Leu	Ser	Pro	Gly

-continued

435	440	445
<210> SEQ ID NO 13		
<211> LENGTH: 448		
<212> TYPE: PRT		
<213> ORGANISM: Artificial Sequence		
<220> FEATURE:		
<223> OTHER INFORMATION: humanized antibody with mouse CDRs and human framework and constant regions		
<220> FEATURE:		
<221> NAME/KEY: DOMAIN		
<222> LOCATION: (1)..(119)		
<223> OTHER INFORMATION: variable domain		
<220> FEATURE:		
<221> NAME/KEY: MISC_FEATURE		
<222> LOCATION: (31)..(35)		
<223> OTHER INFORMATION: CDRH1		
<220> FEATURE:		
<221> NAME/KEY: MISC_FEATURE		
<222> LOCATION: (50)..(66)		
<223> OTHER INFORMATION: CDRH2		
<220> FEATURE:		
<221> NAME/KEY: MISC_FEATURE		
<222> LOCATION: (99)..(108)		
<223> OTHER INFORMATION: CDRH3		
<400> SEQUENCE: 13		
Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala		
1 5 10 15		
Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Gly Tyr		
20 25 30		
Asn Met Asn Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met		
35 40 45		
Gly Asn Ile Asp Pro Tyr Tyr Gly Asn Thr Asn Tyr Asn Gln Lys Phe		
50 55 60		
Lys Gly Arg Val Thr Met Thr Thr Asp Thr Ser Thr Ser Thr Ala Tyr		
65 70 75 80		
Met Glu Leu Arg Ser Leu Arg Ser Asp Asp Thr Ala Val Tyr Tyr Cys		
85 90 95		
Ala Arg Leu Gly Leu Gln Leu Tyr Ala Leu Asp Tyr Trp Gly Gln Gly		
100 105 110		
Thr Thr Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe		
115 120 125		
Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala Leu		
130 135 140		
Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp		
145 150 155 160		
Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu		
165 170 175		
Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser		
180 185 190		
Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys Pro		
195 200 205		
Ser Asn Thr Lys Val Asp Lys Arg Val Glu Pro Lys Ser Cys Asp Lys		
210 215 220		
Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly Asp		
225 230 235 240		
Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser		
245 250 255		

-continued

```

Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu Asp
      260              265              270

Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn
      275              280              285

Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val
      290              295              300

Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu
      305              310              315              320

Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu Lys
      325              330              335

Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr
      340              345              350

Leu Pro Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr
      355              360              365

Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu
      370              375              380

Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu
      385              390              395              400

Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys
      405              410              415

Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu
      420              425              430

Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly
      435              440              445

```

```

<210> SEQ ID NO 14
<211> LENGTH: 448
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: humanized antibody with mouse CDRs and human
      framework and constant regions
<220> FEATURE:
<221> NAME/KEY: DOMAIN
<222> LOCATION: (1)..(119)
<223> OTHER INFORMATION: variable domain
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (31)..(35)
<223> OTHER INFORMATION: CDRH1
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (50)..(66)
<223> OTHER INFORMATION: CDRH2
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (99)..(108)
<223> OTHER INFORMATION: CDRH3

<400> SEQUENCE: 14

```

```

Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala
1              5              10              15

Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Gly Tyr
      20              25              30

Asn Met Asn Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met
      35              40              45

Gly Asn Ile Asp Pro Tyr Tyr Gly Asn Thr Asn Tyr Asn Gln Lys Phe
50              55              60

```

-continued

Lys	Gly	Arg	Val	Thr	Met	Thr	Thr	Asp	Thr	Ser	Thr	Ser	Thr	Ala	Tyr	65	70	75	80
Met	Glu	Leu	Arg	Ser	Leu	Arg	Ser	Asp	Asp	Thr	Ala	Val	Tyr	Tyr	Cys	85	90	95	
Ala	Arg	Leu	Gly	Leu	Gln	Leu	Tyr	Ala	Leu	Asp	Tyr	Trp	Gly	Gln	Gly	100	105	110	
Thr	Thr	Val	Thr	Val	Ser	Ser	Ala	Ser	Thr	Lys	Gly	Pro	Ser	Val	Phe	115	120	125	
Pro	Leu	Ala	Pro	Ser	Ser	Lys	Ser	Thr	Ser	Gly	Gly	Thr	Ala	Ala	Leu	130	135	140	
Gly	Cys	Leu	Val	Lys	Asp	Tyr	Phe	Pro	Glu	Pro	Val	Thr	Val	Ser	Trp	145	150	155	160
Asn	Ser	Gly	Ala	Leu	Thr	Ser	Gly	Val	His	Thr	Phe	Pro	Ala	Val	Leu	165	170	175	
Gln	Ser	Ser	Gly	Leu	Tyr	Ser	Leu	Ser	Ser	Val	Val	Thr	Val	Pro	Ser	180	185	190	
Ser	Ser	Leu	Gly	Thr	Gln	Thr	Tyr	Ile	Cys	Asn	Val	Asn	His	Lys	Pro	195	200	205	
Ser	Asn	Thr	Lys	Val	Asp	Lys	Arg	Val	Glu	Pro	Lys	Ser	Cys	Asp	Lys	210	215	220	
Thr	His	Thr	Cys	Pro	Pro	Cys	Pro	Ala	Pro	Glu	Leu	Leu	Gly	Gly	Pro	225	230	235	240
Ser	Val	Phe	Leu	Phe	Pro	Pro	Lys	Pro	Lys	Asp	Thr	Leu	Met	Ile	Ser	245	250	255	
Arg	Thr	Pro	Glu	Val	Thr	Cys	Val	Val	Val	Asp	Val	Glu	His	Glu	Asp	260	265	270	
Pro	Glu	Val	Lys	Phe	Asn	Trp	Tyr	Val	Asp	Gly	Val	Glu	Val	His	Asn	275	280	285	
Ala	Lys	Thr	Lys	Pro	Arg	Glu	Glu	Gln	Tyr	Asn	Ser	Thr	Tyr	Arg	Val	290	295	300	
Val	Ser	Val	Leu	Thr	Val	Leu	His	Gln	Asp	Trp	Leu	Asn	Gly	Lys	Glu	305	310	315	320
Tyr	Lys	Cys	Lys	Val	Ser	Asn	Lys	Ala	Leu	Pro	Ala	Pro	Ile	Glu	Lys	325	330	335	
Thr	Ile	Ser	Lys	Ala	Lys	Gly	Gln	Pro	Arg	Glu	Pro	Gln	Val	Tyr	Thr	340	345	350	
Leu	Pro	Pro	Ser	Arg	Glu	Glu	Met	Thr	Lys	Asn	Gln	Val	Ser	Leu	Thr	355	360	365	
Cys	Leu	Val	Lys	Gly	Phe	Tyr	Pro	Ser	Asp	Ile	Ala	Val	Glu	Trp	Glu	370	375	380	
Ser	Asn	Gly	Gln	Pro	Glu	Asn	Asn	Tyr	Lys	Thr	Thr	Pro	Pro	Val	Leu	385	390	395	400
Asp	Ser	Asp	Gly	Ser	Phe	Phe	Leu	Tyr	Ser	Lys	Leu	Thr	Val	Asp	Lys	405	410	415	
Ser	Arg	Trp	Gln	Gln	Gly	Asn	Val	Phe	Ser	Cys	Ser	Val	Met	His	Glu	420	425	430	
Ala	Leu	His	Asn	His	Tyr	Thr	Gln	Lys	Ser	Leu	Ser	Leu	Ser	Pro	Gly	435	440	445	

<210> SEQ ID NO 15

<211> LENGTH: 448

<212> TYPE: PRT

-continued

<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: humanized antibody with mouse CDRs and human framework and constant regions
<220> FEATURE:
<221> NAME/KEY: DOMAIN
<222> LOCATION: (1)..(119)
<223> OTHER INFORMATION: variable domain
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (31)..(35)
<223> OTHER INFORMATION: CDRH1
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (50)..(66)
<223> OTHER INFORMATION: CDRH2
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (99)..(108)
<223> OTHER INFORMATION: CDRH3

<400> SEQUENCE: 15

Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala
1 5 10 15

Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Gly Tyr
20 25 30

Asn Met Asn Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met
35 40 45

Gly Asn Ile Asp Pro Tyr Tyr Gly Asn Thr Asn Tyr Asn Gln Lys Phe
50 55 60

Lys Gly Arg Val Thr Met Thr Thr Asp Thr Ser Thr Ser Thr Ala Tyr
65 70 75 80

Met Glu Leu Arg Ser Leu Arg Ser Asp Asp Thr Ala Val Tyr Tyr Cys
85 90 95

Ala Arg Leu Gly Leu Gln Leu Tyr Ala Leu Asp Tyr Trp Gly Gln Gly
100 105 110

Thr Thr Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe
115 120 125

Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala Leu
130 135 140

Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp
145 150 155 160

Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu
165 170 175

Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser
180 185 190

Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys Pro
195 200 205

Ser Asn Thr Lys Val Asp Lys Arg Val Glu Pro Lys Ser Cys Asp Lys
210 215 220

Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly Pro
225 230 235 240

Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser
245 250 255

Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Glu His Glu Asp
260 265 270

Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn
275 280 285

-continued

Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val
 290 295 300

Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu
 305 310 315 320

Tyr Lys Cys Lys Val Ser Asn Lys Ala Phe Pro Ala Pro Ile Glu Lys
 325 330 335

Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr
 340 345 350

Leu Pro Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr
 355 360 365

Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu
 370 375 380

Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu
 385 390 395 400

Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys
 405 410 415

Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu
 420 425 430

Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly
 435 440 445

<210> SEQ ID NO 16
 <211> LENGTH: 448
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: humanized antibody with mouse CDRs and human
 framework and constant regions
 <220> FEATURE:
 <221> NAME/KEY: DOMAIN
 <222> LOCATION: (1)..(119)
 <223> OTHER INFORMATION: variable domain
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (31)..(35)
 <223> OTHER INFORMATION: CDRH1
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (50)..(66)
 <223> OTHER INFORMATION: CDRH2
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (99)..(108)
 <223> OTHER INFORMATION: CDRH3

<400> SEQUENCE: 16

Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala
 1 5 10 15

Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Gly Tyr
 20 25 30

Asn Met Asn Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met
 35 40 45

Gly Asn Ile Asp Pro Tyr Tyr Gly Asn Thr Asn Tyr Asn Gln Lys Phe
 50 55 60

Lys Gly Arg Val Thr Met Thr Thr Asp Thr Ser Thr Ser Thr Ala Tyr
 65 70 75 80

Met Glu Leu Arg Ser Leu Arg Ser Asp Asp Thr Ala Val Tyr Tyr Cys
 85 90 95

-continued

Ala	Arg	Leu	Gly	Leu	Gln	Leu	Tyr	Ala	Leu	Asp	Tyr	Trp	Gly	Gln	Gly	100	105	110
Thr	Thr	Val	Thr	Val	Ser	Ser	Ala	Ser	Thr	Lys	Gly	Pro	Ser	Val	Phe	115	120	125
Pro	Leu	Ala	Pro	Ser	Ser	Lys	Ser	Thr	Ser	Gly	Gly	Thr	Ala	Ala	Leu	130	135	140
Gly	Cys	Leu	Val	Lys	Asp	Tyr	Phe	Pro	Glu	Pro	Val	Thr	Val	Ser	Trp	145	150	155
Asn	Ser	Gly	Ala	Leu	Thr	Ser	Gly	Val	His	Thr	Phe	Pro	Ala	Val	Leu	165	170	175
Gln	Ser	Ser	Gly	Leu	Tyr	Ser	Leu	Ser	Ser	Val	Val	Thr	Val	Pro	Ser	180	185	190
Ser	Ser	Leu	Gly	Thr	Gln	Thr	Tyr	Ile	Cys	Asn	Val	Asn	His	Lys	Pro	195	200	205
Ser	Asn	Thr	Lys	Val	Asp	Lys	Arg	Val	Glu	Pro	Lys	Ser	Cys	Asp	Lys	210	215	220
Thr	His	Thr	Cys	Pro	Pro	Cys	Pro	Ala	Pro	Glu	Leu	Leu	Gly	Gly	Asp	225	230	235
Ser	Val	Phe	Leu	Phe	Pro	Pro	Lys	Pro	Lys	Asp	Thr	Leu	Met	Ile	Ser	245	250	255
Arg	Thr	Pro	Glu	Val	Thr	Cys	Val	Val	Val	Asp	Val	Ser	His	Glu	Asp	260	265	270
Gly	Glu	Val	Lys	Phe	Asn	Trp	Tyr	Val	Asp	Gly	Val	Glu	Val	His	Asn	275	280	285
Ala	Lys	Thr	Lys	Pro	Arg	Glu	Glu	Gln	Tyr	Asn	Ser	Thr	Tyr	Arg	Val	290	295	300
Val	Ser	Val	Leu	Thr	Val	Leu	His	Gln	Asp	Trp	Leu	Asn	Gly	Lys	Glu	305	310	315
Tyr	Lys	Cys	Lys	Val	Ser	Asn	Lys	Ala	Leu	Pro	Ala	Pro	Ile	Glu	Lys	325	330	335
Thr	Ile	Ser	Lys	Ala	Lys	Gly	Gln	Pro	Arg	Glu	Pro	Gln	Val	Tyr	Thr	340	345	350
Leu	Pro	Pro	Ser	Arg	Glu	Glu	Met	Thr	Lys	Asn	Gln	Val	Ser	Leu	Thr	355	360	365
Cys	Leu	Val	Lys	Gly	Phe	Tyr	Pro	Ser	Asp	Ile	Ala	Val	Glu	Trp	Glu	370	375	380
Ser	Asn	Gly	Gln	Pro	Glu	Asn	Asn	Tyr	Lys	Thr	Thr	Pro	Pro	Val	Leu	385	390	395
Asp	Ser	Asp	Gly	Ser	Phe	Phe	Leu	Tyr	Ser	Lys	Leu	Thr	Val	Asp	Lys	405	410	415
Ser	Arg	Trp	Gln	Gln	Gly	Asn	Val	Phe	Ser	Cys	Ser	Val	Met	His	Glu	420	425	430
Ala	Leu	His	Asn	His	Tyr	Thr	Gln	Lys	Ser	Leu	Ser	Leu	Ser	Pro	Gly	435	440	445

<210> SEQ ID NO 17

<211> LENGTH: 448

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: humanized antibody with mouse CDRs and human framework and constant regions

<400> SEQUENCE: 17

-continued

Gln	Val	Gln	Leu	Val	Gln	Ser	Gly	Ala	Glu	Val	Lys	Lys	Pro	Gly	Ala	
1			5						10					15		
Ser	Val	Lys	Val	Ser	Cys	Lys	Ala	Ser	Gly	Tyr	Thr	Phe	Thr	Gly	Tyr	
		20						25					30			
Asn	Met	Asn	Trp	Val	Arg	Gln	Ala	Pro	Gly	Gln	Gly	Leu	Glu	Trp	Met	
		35					40					45				
Gly	Asn	Ile	Asp	Pro	Tyr	Tyr	Gly	Asn	Thr	Asn	Tyr	Asn	Gln	Lys	Phe	
	50					55					60					
Lys	Gly	Arg	Val	Thr	Met	Thr	Thr	Asp	Thr	Ser	Thr	Ser	Thr	Ala	Tyr	
65					70					75					80	
Met	Glu	Leu	Arg	Ser	Leu	Arg	Ser	Asp	Asp	Thr	Ala	Val	Tyr	Tyr	Cys	
			85					90						95		
Ala	Arg	Leu	Gly	Leu	Gln	Leu	Tyr	Ala	Leu	Asp	Tyr	Trp	Gly	Gln	Gly	
		100						105					110			
Thr	Thr	Val	Thr	Val	Ser	Ser	Ala	Ser	Thr	Lys	Gly	Pro	Ser	Val	Phe	
		115					120					125				
Pro	Leu	Ala	Pro	Ser	Ser	Lys	Ser	Thr	Ser	Gly	Gly	Thr	Ala	Ala	Leu	
	130					135					140					
Gly	Cys	Leu	Val	Lys	Asp	Tyr	Phe	Pro	Glu	Pro	Val	Thr	Val	Ser	Trp	
145					150					155					160	
Asn	Ser	Gly	Ala	Leu	Thr	Ser	Gly	Val	His	Thr	Phe	Pro	Ala	Val	Leu	
			165					170						175		
Gln	Ser	Ser	Gly	Leu	Tyr	Ser	Leu	Ser	Ser	Val	Val	Thr	Val	Pro	Ser	
			180					185					190			
Ser	Ser	Leu	Gly	Thr	Gln	Thr	Tyr	Ile	Cys	Asn	Val	Asn	His	Lys	Pro	
		195					200					205				
Ser	Asn	Thr	Lys	Val	Asp	Lys	Arg	Val	Glu	Pro	Lys	Ser	Cys	Asp	Lys	
	210					215					220					
Thr	His	Thr	Cys	Pro	Pro	Cys	Pro	Ala	Pro	Glu	Leu	Leu	Gly	Gly	Asp	
225					230					235					240	
Ser	Val	Phe	Leu	Phe	Pro	Pro	Lys	Pro	Lys	Asp	Thr	Leu	Met	Ile	Ser	
		245						250						255		
Arg	Thr	Pro	Glu	Val	Thr	Cys	Val	Val	Val	Asp	Val	Ser	His	Glu	Glu	
		260					265						270			
Gly	Glu	Val	Lys	Phe	Asn	Trp	Tyr	Val	Asp	Gly	Val	Glu	Val	His	Asn	
	275					280					285					
Ala	Lys	Thr	Lys	Pro	Arg	Glu	Glu	Gln	Tyr	Asn	Ser	Thr	Tyr	Arg	Val	
	290				295						300					
Val	Ser	Val	Leu	Thr	Val	Leu	His	Gln	Asp	Trp	Leu	Asn	Gly	Lys	Glu	
305					310					315					320	
Tyr	Lys	Cys	Lys	Val	Ser	Asn	Lys	Ala	Leu	Pro	Ala	Pro	Ile	Glu	Lys	
			325					330						335		
Thr	Ile	Ser	Lys	Ala	Lys	Gly	Gln	Pro	Arg	Glu	Pro	Gln	Val	Tyr	Thr	
		340					345						350			
Leu	Pro	Pro	Ser	Arg	Glu	Glu	Met	Thr	Lys	Asn	Gln	Val	Ser	Leu	Thr	
		355					360					365				
Cys	Leu	Val	Lys	Gly	Phe	Tyr	Pro	Ser	Asp	Ile	Ala	Val	Glu	Trp	Glu	
	370					375					380					
Ser	Asn	Gly	Gln	Pro	Glu	Asn	Asn	Tyr	Lys	Thr	Thr	Pro	Pro	Val	Leu	
385				390						395					400	

-continued

Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly
435 440 445

```
<210> SEQ ID NO 18
<211> LENGTH: 448
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: humanized antibody with mouse CDRs and human
framework and constant regions
<220> FEATURE:
<221> NAME/KEY: DOMAIN
<222> LOCATION: (1)..(119)
<223> OTHER INFORMATION: variable domain
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (31)..(35)
<223> OTHER INFORMATION: CDRH1
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (50)..(66)
<223> OTHER INFORMATION: CDRH2
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (99)..(108)
<223> OTHER INFORMATION: CDRH3
```

<400> SEQUENCE: 18

Ser Asn Thr Lys Val Asp Lys Arg Val Glu Pro Lys Ser Cys Asp Lys

-continued

210	215	220
Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Asp Asp		
225	230	235 240
Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser		
	245	250 255
Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser Asp Glu Asp		
	260	265 270
Gly Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn		
	275	280 285
Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val		
	290	295 300
Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu		
305	310	315 320
Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu Lys		
	325	330 335
Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr		
	340	345 350
Leu Pro Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr		
	355	360 365
Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu		
	370	375 380
Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu		
385	390	395 400
Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys		
	405	410 415
Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu		
	420	425 430
Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly		
	435	440 445

<210> SEQ ID NO 19

<211> LENGTH: 448

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: humanized antibody with mouse CDRs and human framework and constant regions

<220> FEATURE:

<221> NAME/KEY: DOMAIN

<222> LOCATION: (1)..(119)

<223> OTHER INFORMATION: variable domain

<220> FEATURE:

<221> NAME/KEY: MISC_FEATURE

<222> LOCATION: (31)..(35)

<223> OTHER INFORMATION: CDRH1

<220> FEATURE:

<221> NAME/KEY: MISC_FEATURE

<222> LOCATION: (50)..(66)

<223> OTHER INFORMATION: CDRH2

<220> FEATURE:

<221> NAME/KEY: MISC_FEATURE

<222> LOCATION: (99)..(108)

<223> OTHER INFORMATION: CDRH3

<400> SEQUENCE: 19

Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala
1 5 10 15

Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Gly Tyr
20 25 30

-continued

Asn	Met	Asn	Trp	Val	Arg	Gln	Ala	Pro	Gly	Gln	Gly	Leu	Glu	Trp	Met
	35						40					45			
Gly	Asn	Ile	Asp	Pro	Tyr	Tyr	Gly	Asn	Thr	Asn	Tyr	Asn	Gln	Lys	Phe
	50					55					60				
Lys	Gly	Arg	Val	Thr	Met	Thr	Thr	Asp	Thr	Ser	Thr	Ser	Thr	Ala	Tyr
	65				70					75					80
Met	Glu	Leu	Arg	Ser	Leu	Arg	Ser	Asp	Asp	Thr	Ala	Val	Tyr	Tyr	Cys
				85					90					95	
Ala	Arg	Leu	Gly	Leu	Gln	Leu	Tyr	Ala	Leu	Asp	Tyr	Trp	Gly	Gln	Gly
			100					105					110		
Thr	Thr	Val	Thr	Val	Ser	Ser	Ala	Ser	Thr	Lys	Gly	Pro	Ser	Val	Phe
		115					120					125			
Pro	Leu	Ala	Pro	Ser	Ser	Lys	Ser	Thr	Ser	Gly	Gly	Thr	Ala	Ala	Leu
	130					135					140				
Gly	Cys	Leu	Val	Lys	Asp	Tyr	Phe	Pro	Glu	Pro	Val	Thr	Val	Ser	Trp
	145				150					155					160
Asn	Ser	Gly	Ala	Leu	Thr	Ser	Gly	Val	His	Thr	Phe	Pro	Ala	Val	Leu
				165					170					175	
Gln	Ser	Ser	Gly	Leu	Tyr	Ser	Leu	Ser	Ser	Val	Val	Thr	Val	Pro	Ser
			180					185					190		
Ser	Ser	Leu	Gly	Thr	Gln	Thr	Tyr	Ile	Cys	Asn	Val	Asn	His	Lys	Pro
	195						200					205			
Ser	Asn	Thr	Lys	Val	Asp	Lys	Arg	Val	Glu	Pro	Lys	Ser	Cys	Asp	Lys
	210					215					220				
Thr	His	Thr	Cys	Pro	Pro	Cys	Pro	Ala	Pro	Glu	Leu	Leu	Gly	Asp	Asp
	225				230					235					240
Ser	Val	Phe	Leu	Phe	Pro	Pro	Lys	Pro	Lys	Asp	Thr	Leu	Met	Ile	Ser
			245						250					255	
Arg	Thr	Pro	Glu	Val	Thr	Cys	Val	Val	Val	Asp	Val	Ser	His	Glu	Asp
			260					265					270		
Gly	Glu	Val	Lys	Phe	Asn	Trp	Tyr	Val	Asp	Gly	Val	Glu	Val	His	Asn
	275					280						285			
Ala	Lys	Thr	Lys	Pro	Arg	Glu	Glu	Gln	Tyr	Asn	Ser	Thr	Tyr	Arg	Val
	290					295					300				
Val	Ser	Val	Leu	Thr	Val	Leu	His	Gln	Asp	Trp	Leu	Asn	Gly	Lys	Glu
	305				310					315					320
Tyr	Lys	Cys	Lys	Val	Ser	Asn	Lys	Ala	Leu	Pro	Arg	Pro	Ile	Glu	Lys
			325						330					335	
Thr	Ile	Ser	Lys	Ala	Lys	Gly	Gln	Pro	Arg	Glu	Pro	Gln	Val	Tyr	Thr
			340					345					350		
Leu	Pro	Pro	Ser	Arg	Glu	Glu	Met	Thr	Lys	Asn	Gln	Val	Ser	Leu	Thr
	355						360					365			
Cys	Leu	Val	Lys	Gly	Phe	Tyr	Pro	Ser	Asp	Ile	Ala	Val	Glu	Trp	Glu
	370					375					380				
Ser	Asn	Gly	Gln	Pro	Glu	Asn	Asn	Tyr	Lys	Thr	Thr	Pro	Pro	Val	Leu
	385				390					395					400
Asp	Ser	Asp	Gly	Ser	Phe	Phe	Leu	Tyr	Ser	Lys	Leu	Thr	Val	Asp	Lys
			405						410					415	
Ser	Arg	Trp	Gln	Gln	Gly	Asn	Val	Phe	Ser	Cys	Ser	Val	Met	His	Glu
			420					425					430		

-continued

Ala	Leu	His	Asn	His	Tyr	Thr	Gln	Lys	Ser	Leu	Ser	Leu	Ser	Pro	Gly
			435					440						445	

<210> SEQ ID NO 20

<211> LENGTH: 448

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: humanized antibody with mouse CDRs and human framework and constant regions

<400> SEQUENCE: 20

Gln	Val	Gln	Leu	Val	Gln	Ser	Gly	Ala	Glu	Val	Lys	Lys	Pro	Gly	Ala
1			5						10				15		
Ser	Val	Lys	Val	Ser	Cys	Lys	Ala	Ser	Gly	Tyr	Thr	Phe	Thr	Gly	Tyr
		20					25					30			
Asn	Met	Asn	Trp	Val	Arg	Gln	Ala	Pro	Gly	Gln	Gly	Leu	Glu	Trp	Met
		35				40						45			
Gly	Asn	Ile	Asp	Pro	Tyr	Tyr	Gly	Asn	Thr	Asn	Tyr	Asn	Gln	Lys	Phe
	50					55					60				
Lys	Gly	Arg	Val	Thr	Met	Thr	Thr	Asp	Thr	Ser	Thr	Ser	Thr	Ala	Tyr
65					70					75				80	
Met	Glu	Leu	Arg	Ser	Leu	Arg	Ser	Asp	Asp	Thr	Ala	Val	Tyr	Tyr	Cys
			85					90						95	
Ala	Arg	Leu	Gly	Leu	Gln	Leu	Tyr	Ala	Leu	Asp	Tyr	Trp	Gly	Gln	Gly
		100						105					110		
Thr	Thr	Val	Thr	Val	Ser	Ser	Ala	Ser	Thr	Lys	Gly	Pro	Ser	Val	Phe
		115					120					125			
Pro	Leu	Ala	Pro	Ser	Ser	Lys	Ser	Thr	Ser	Gly	Gly	Thr	Ala	Ala	Leu
	130					135					140				
Gly	Cys	Leu	Val	Lys	Asp	Tyr	Phe	Pro	Glu	Pro	Val	Thr	Val	Ser	Trp
145				150					155					160	
Asn	Ser	Gly	Ala	Leu	Thr	Ser	Gly	Val	His	Thr	Phe	Pro	Ala	Val	Leu
			165					170						175	
Gln	Ser	Ser	Gly	Leu	Tyr	Ser	Leu	Ser	Ser	Val	Val	Thr	Val	Pro	Ser
			180					185					190		
Ser	Ser	Leu	Gly	Thr	Gln	Thr	Tyr	Ile	Cys	Asn	Val	Asn	His	Lys	Pro
		195					200					205			
Ser	Asn	Thr	Lys	Val	Asp	Lys	Arg	Val	Glu	Pro	Lys	Ser	Cys	Asp	Lys
	210					215					220				
Thr	His	Thr	Cys	Pro	Pro	Cys	Pro	Ala	Pro	Glu	Leu	Leu	Gly	Asp	Asp
225				230						235				240	
Ser	Val	Phe	Leu	Phe	Pro	Pro	Lys	Pro	Lys	Asp	Thr	Leu	Met	Ile	Ser
			245					250						255	
Arg	Thr	Pro	Glu	Val	Thr	Cys	Val	Val	Val	Asp	Val	Ser	His	Glu	Glu
		260						265					270		
Gly	Glu	Val	Lys	Phe	Asn	Trp	Tyr	Val	Asp	Gly	Val	Glu	Val	His	Asn
		275					280					285			
Ala	Lys	Thr	Lys	Pro	Arg	Glu	Glu	Gln	Tyr	Asn	Ser	Thr	Tyr	Arg	Val
	290					295					300				
Val	Ser	Val	Leu	Thr	Val	Leu	His	Gln	Asp	Trp	Leu	Asn	Gly	Lys	Glu
305					310					315				320	
Tyr	Lys	Cys	Lys	Val	Ser	Asn	Lys	Ala	Leu	Pro	Arg	Pro	Ile	Glu	Lys
			325					330						335	

-continued

Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr
 340 345 350

Leu Pro Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr
 355 360 365

Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu
 370 375 380

Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu
 385 390 395 400

Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys
 405 410 415

Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu
 420 425 430

Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly
 435 440 445

<210> SEQ ID NO 21
 <211> LENGTH: 448
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: humanized antibody with mouse CDRs and human
 framework and constant regions
 <220> FEATURE:
 <221> NAME/KEY: DOMAIN
 <222> LOCATION: (1)..(119)
 <223> OTHER INFORMATION: variable domain
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (31)..(35)
 <223> OTHER INFORMATION: CDRH1
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (50)..(66)
 <223> OTHER INFORMATION: CDRH2
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (99)..(108)
 <223> OTHER INFORMATION: CDRH3

<400> SEQUENCE: 21

Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala
 1 5 10 15

Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Gly Tyr
 20 25 30

Asn Met Asn Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met
 35 40 45

Gly Asn Ile Asp Pro Tyr Tyr Gly Asn Thr Asn Tyr Asn Gln Lys Phe
 50 55 60

Lys Gly Arg Val Thr Met Thr Thr Asp Thr Ser Thr Ser Thr Ala Tyr
 65 70 75 80

Met Glu Leu Arg Ser Leu Arg Ser Asp Asp Thr Ala Val Tyr Tyr Cys
 85 90 95

Ala Arg Leu Gly Leu Gln Leu Tyr Ala Leu Asp Tyr Trp Gly Gln Gly
 100 105 110

Thr Thr Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe
 115 120 125

Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala Leu
 130 135 140

-continued

Gly	Cys	Leu	Val	Lys	Asp	Tyr	Phe	Pro	Glu	Pro	Val	Thr	Val	Ser	Trp
145					150					155					160
Asn	Ser	Gly	Ala	Leu	Thr	Ser	Gly	Val	His	Thr	Phe	Pro	Ala	Val	Leu
				165					170					175	
Gln	Ser	Ser	Gly	Leu	Tyr	Ser	Leu	Ser	Ser	Val	Val	Thr	Val	Pro	Ser
			180					185					190		
Ser	Ser	Leu	Gly	Thr	Gln	Thr	Tyr	Ile	Cys	Asn	Val	Asn	His	Lys	Pro
		195					200					205			
Ser	Asn	Thr	Lys	Val	Asp	Lys	Arg	Val	Glu	Pro	Lys	Ser	Cys	Asp	Lys
	210					215					220				
Thr	His	Thr	Cys	Pro	Pro	Cys	Pro	Ala	Pro	Glu	Leu	Leu	Gly	Asp	Asp
225					230					235					240
Ser	Val	Phe	Leu	Phe	Pro	Pro	Lys	Pro	Lys	Asp	Thr	Leu	Met	Ile	Ser
			245						250					255	
Arg	Thr	Pro	Glu	Val	Thr	Cys	Val	Val	Val	Asp	Val	Ser	Asp	Glu	Asp
			260					265					270		
Gly	Glu	Val	Lys	Phe	Asn	Trp	Tyr	Val	Asp	Gly	Val	Glu	Val	His	Asn
		275					280					285			
Ala	Lys	Thr	Lys	Pro	Arg	Glu	Glu	Gln	Tyr	Asn	Ser	Thr	Tyr	Arg	Val
	290					295					300				
Val	Ser	Val	Leu	Thr	Val	Leu	His	Gln	Asp	Trp	Leu	Asn	Gly	Lys	Glu
305					310					315					320
Tyr	Lys	Cys	Lys	Val	Ser	Asn	Lys	Ala	Leu	Pro	Arg	Pro	Ile	Glu	Lys
			325						330					335	
Thr	Ile	Ser	Lys	Ala	Lys	Gly	Gln	Pro	Arg	Glu	Pro	Gln	Val	Tyr	Thr
			340					345					350		
Leu	Pro	Pro	Ser	Arg	Glu	Glu	Met	Thr	Lys	Asn	Gln	Val	Ser	Leu	Thr
		355					360					365			
Cys	Leu	Val	Lys	Gly	Phe	Tyr	Pro	Ser	Asp	Ile	Ala	Val	Glu	Trp	Glu
	370					375					380				
Ser	Asn	Gly	Gln	Pro	Glu	Asn	Asn	Tyr	Lys	Thr	Thr	Pro	Pro	Val	Leu
385				390						395					400
Asp	Ser	Asp	Gly	Ser	Phe	Phe	Leu	Tyr	Ser	Lys	Leu	Thr	Val	Asp	Lys
			405						410					415	
Ser	Arg	Trp	Gln	Gln	Gly	Asn	Val	Phe	Ser	Cys	Ser	Val	Met	His	Glu
			420					425					430		
Ala	Leu	His	Asn	His	Tyr	Thr	Gln	Lys	Ser	Leu	Ser	Leu	Ser	Pro	Gly
		435					440					445			

<210> SEQ ID NO 22

<211> LENGTH: 448

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: humanized antibody with mouse CDRs and human framework and constant regions

<220> FEATURE:

<221> NAME/KEY: DOMAIN

<222> LOCATION: (1)..(119)

<223> OTHER INFORMATION: variable domain

<220> FEATURE:

<221> NAME/KEY: MISC_FEATURE

<222> LOCATION: (31)..(35)

<223> OTHER INFORMATION: CDRH1

<220> FEATURE:

<221> NAME/KEY: MISC_FEATURE

<222> LOCATION: (50)..(66)

-continued

<223> OTHER INFORMATION: CDRH2

<220> FEATURE:

<221> NAME/KEY: MISC_FEATURE

<222> LOCATION: (99)..(108)

<223> OTHER INFORMATION: CDRH3

<400> SEQUENCE: 22

Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala
 1 5 10 15
 Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Gly Tyr
 20 25 30
 Asn Met Asn Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met
 35 40 45
 Gly Asn Ile Asp Pro Tyr Tyr Gly Asn Thr Asn Tyr Asn Gln Lys Phe
 50 55 60
 Lys Gly Arg Val Thr Met Thr Thr Asp Thr Ser Thr Ser Thr Ala Tyr
 65 70 75 80
 Met Glu Leu Arg Ser Leu Arg Ser Asp Asp Thr Ala Val Tyr Tyr Cys
 85 90 95
 Ala Arg Leu Gly Leu Gln Leu Tyr Ala Leu Asp Tyr Trp Gly Gln Gly
 100 105 110
 Thr Thr Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe
 115 120 125
 Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala Leu
 130 135 140
 Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp
 145 150 155 160
 Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu
 165 170 175
 Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser
 180 185 190
 Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys Pro
 195 200 205
 Ser Asn Thr Lys Val Asp Lys Arg Val Glu Pro Lys Ser Cys Asp Lys
 210 215 220
 Thr His Thr Cys Pro Pro Cys Pro Ala Pro Asp Leu Leu Gly Asp Asp
 225 230 235 240
 Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser
 245 250 255
 Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser Asp Glu Asp
 260 265 270
 Gly Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn
 275 280 285
 Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val
 290 295 300
 Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu
 305 310 315 320
 Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Arg Pro Ile Glu Lys
 325 330 335
 Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr
 340 345 350
 Leu Pro Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr
 355 360 365

-continued

Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu
 370 375 380

Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu
 385 390 395 400

Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys
 405 410 415

Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu
 420 425 430

Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly
 435 440 445

<210> SEQ ID NO 23
 <211> LENGTH: 446
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: chimeric antibody with mouse variable domain
 and human constant domain
 <220> FEATURE:
 <221> NAME/KEY: DOMAIN
 <222> LOCATION: (1)..(117)
 <223> OTHER INFORMATION: variable domain
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (31)..(35)
 <223> OTHER INFORMATION: CDRH1
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (50)..(66)
 <223> OTHER INFORMATION: CDRH2
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (99)..(106)
 <223> OTHER INFORMATION: CDRH3

<400> SEQUENCE: 23

Gln Val Gln Leu Gln Gln Ser Ala Ala Glu Leu Ala Arg Pro Gly Ala
 1 5 10 15

Ser Val Lys Met Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asp Leu
 20 25 30

Ser Met His Trp Ile Lys Gln Arg Pro Gly Gln Gly Leu Glu Trp Ile
 35 40 45

Gly Tyr Ile Thr Pro Ser Ser Gly Tyr Thr Ala Tyr Asn Gln Lys Phe
 50 55 60

Lys Gly Lys Thr Thr Leu Thr Ala Asp Lys Ser Ser Ser Thr Ala Tyr
 65 70 75 80

Met Gln Leu Ser Ser Leu Thr Ser Glu Asp Ser Ala Val Tyr Tyr Cys
 85 90 95

Ala Arg Leu Ile Leu Gln Arg Gly Ala Tyr Trp Gly Gln Gly Thr Leu
 100 105 110

Val Thr Val Ser Ala Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu
 115 120 125

Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly Cys
 130 135 140

Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser
 145 150 155 160

Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu Gln Ser
 165 170 175

-continued

```

Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser
    180                      185                      190

Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys Pro Ser Asn
    195                      200                      205

Thr Lys Val Asp Lys Arg Val Glu Pro Lys Ser Cys Asp Lys Thr His
    210                      215                      220

Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser Val
    225                      230                      235                      240

Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr
    245                      250                      255

Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu Asp Pro Glu
    260                      265                      270

Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala Lys
    275                      280                      285

Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser
    290                      295                      300

Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys
    305                      310                      315                      320

Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile
    325                      330                      335

Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro
    340                      345                      350

Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr Cys Leu
    355                      360                      365

Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn
    370                      375                      380

Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser
    385                      390                      395                      400

Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg
    405                      410                      415

Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu
    420                      425                      430

His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly
    435                      440                      445

```

```

<210> SEQ ID NO 24
<211> LENGTH: 218
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: chimeric antibody with mouse variable domain
and human constant domain
<220> FEATURE:
<221> NAME/KEY: DOMAIN
<222> LOCATION: (1)..(111)
<223> OTHER INFORMATION: variable domain
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (24)..(38)
<223> OTHER INFORMATION: CDRL1
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (54)..(60)
<223> OTHER INFORMATION: CDRL2
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (93)..(101)
<223> OTHER INFORMATION: CDRL3

```

-continued

<400> SEQUENCE: 24

```

Asp Ile Val Leu Thr Gln Ser Pro Ala Ser Leu Ala Val Ser Leu Gly
1           5           10           15
Gln Arg Ala Thr Ile Ser Cys Arg Ala Ser Lys Asn Val Asp Ser Tyr
20          25          30
Gly Asn Ser Phe Met His Trp Tyr Gln Gln Lys Pro Gly Gln Pro Pro
35          40          45
Lys Leu Leu Ile Tyr Arg Ala Ser Asn Leu Glu Ser Gly Ile Pro Ala
50          55          60
Arg Phe Gly Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Asn
65          70          75          80
Pro Val Glu Ala Asp Asp Val Ala Thr Tyr Tyr Cys Gln Gln Ser Asn
85          90          95
Glu Asp Pro Leu Thr Phe Gly Ala Gly Thr Lys Leu Glu Leu Lys Arg
100         105         110
Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln
115         120         125
Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr
130         135         140
Pro Arg Glu Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser
145         150         155         160
Gly Asn Ser Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr
165         170         175
Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys
180         185         190
His Lys Val Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro
195         200         205
Val Thr Lys Ser Phe Asn Arg Gly Glu Cys
210         215

```

<210> SEQ ID NO 25

<211> LENGTH: 446

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: humanized antibody with mouse CDRs and human framework and constant regions

<220> FEATURE:

<221> NAME/KEY: DOMAIN

<222> LOCATION: (1)..(117)

<223> OTHER INFORMATION: variable domain

<220> FEATURE:

<221> NAME/KEY: MISC_FEATURE

<222> LOCATION: (31)..(35)

<223> OTHER INFORMATION: CDRH1

<220> FEATURE:

<221> NAME/KEY: MISC_FEATURE

<222> LOCATION: (50)..(66)

<223> OTHER INFORMATION: CDRH2

<220> FEATURE:

<221> NAME/KEY: MISC_FEATURE

<222> LOCATION: (99)..(106)

<223> OTHER INFORMATION: CDRH3

<400> SEQUENCE: 25

```

Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ser
1           5           10           15
Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asp Leu
20          25          30

```

-continued

Ser	Met	His	Trp	Ile	Lys	Gln	Arg	Pro	Gly	Gln	Gly	Leu	Glu	Trp	Ile	
	35						40					45				
Gly	Tyr	Ile	Thr	Pro	Ser	Ser	Gly	Tyr	Thr	Ala	Tyr	Asn	Gln	Lys	Phe	
	50					55					60					
Lys	Gly	Lys	Thr	Thr	Leu	Thr	Ala	Asp	Lys	Ser	Thr	Ser	Thr	Ala	Tyr	
	65				70				75						80	
Met	Glu	Leu	Ser	Ser	Leu	Arg	Ser	Glu	Asp	Thr	Ala	Val	Tyr	Tyr	Cys	
			85					90					95			
Ala	Arg	Leu	Ile	Leu	Gln	Arg	Gly	Ala	Tyr	Trp	Gly	Gln	Gly	Thr	Leu	
		100						105					110			
Val	Thr	Val	Ser	Ser	Ala	Ser	Thr	Lys	Gly	Pro	Ser	Val	Phe	Pro	Leu	
	115						120					125				
Ala	Pro	Ser	Ser	Lys	Ser	Thr	Ser	Gly	Gly	Thr	Ala	Ala	Leu	Gly	Cys	
	130					135					140					
Leu	Val	Lys	Asp	Tyr	Phe	Pro	Glu	Pro	Val	Thr	Val	Ser	Trp	Asn	Ser	
	145				150					155					160	
Gly	Ala	Leu	Thr	Ser	Gly	Val	His	Thr	Phe	Pro	Ala	Val	Leu	Gln	Ser	
			165					170						175		
Ser	Gly	Leu	Tyr	Ser	Leu	Ser	Ser	Val	Val	Thr	Val	Pro	Ser	Ser	Ser	
		180						185					190			
Leu	Gly	Thr	Gln	Thr	Tyr	Ile	Cys	Asn	Val	Asn	His	Lys	Pro	Ser	Asn	
	195					200					205					
Thr	Lys	Val	Asp	Lys	Arg	Val	Glu	Pro	Lys	Ser	Cys	Asp	Lys	Thr	His	
	210					215					220					
Thr	Cys	Pro	Pro	Cys	Pro	Ala	Pro	Glu	Leu	Leu	Gly	Gly	Pro	Ser	Val	
	225				230					235					240	
Phe	Leu	Phe	Pro	Pro	Lys	Pro	Lys	Asp	Thr	Leu	Met	Ile	Ser	Arg	Thr	
			245					250						255		
Pro	Glu	Val	Thr	Cys	Val	Val	Val	Asp	Val	Ser	His	Glu	Asp	Pro	Glu	
		260						265					270			
Val	Lys	Phe	Asn	Trp	Tyr	Val	Asp	Gly	Val	Glu	Val	His	Asn	Ala	Lys	
	275					280						285				
Thr	Lys	Pro	Arg	Glu	Glu	Gln	Tyr	Asn	Ser	Thr	Tyr	Arg	Val	Val	Ser	
	290					295					300					
Val	Leu	Thr	Val	Leu	His	Gln	Asp	Trp	Leu	Asn	Gly	Lys	Glu	Tyr	Lys	
	305				310					315					320	
Cys	Lys	Val	Ser	Asn	Lys	Ala	Leu	Pro	Ala	Pro	Ile	Glu	Lys	Thr	Ile	
			325					330						335		
Ser	Lys	Ala	Lys	Gly	Gln	Pro	Arg	Glu	Pro	Gln	Val	Tyr	Thr	Leu	Pro	
		340						345					350			
Pro	Ser	Arg	Glu	Glu	Met	Thr	Lys	Asn	Gln	Val	Ser	Leu	Thr	Cys	Leu	
		355					360					365				
Val	Lys	Gly	Phe	Tyr	Pro	Ser	Asp	Ile	Ala	Val	Glu	Trp	Glu	Ser	Asn	
	370					375					380					
Gly	Gln	Pro	Glu	Asn	Asn	Tyr	Lys	Thr	Thr	Pro	Pro	Val	Leu	Asp	Ser	
	385				390					395					400	
Asp	Gly	Ser	Phe	Phe	Leu	Tyr	Ser	Lys	Leu	Thr	Val	Asp	Lys	Ser	Arg	
			405					410						415		
Trp	Gln	Gln	Gly	Asn	Val	Phe	Ser	Cys	Ser	Val	Met	His	Glu	Ala	Leu	
			420					425					430			

-continued

His	Asn	His	Tyr	Thr	Gln	Lys	Ser	Leu	Ser	Leu	Ser	Pro	Gly
		435					440					445	

```

<210> SEQ ID NO 26
<211> LENGTH: 218
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: humanized antibody with mouse CDRs and human
        framework and constant regions
<220> FEATURE:
<221> NAME/KEY: DOMAIN
<222> LOCATION: (1)..(111)
<223> OTHER INFORMATION: variable domain
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (24)..(38)
<223> OTHER INFORMATION: CDRL1
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (54)..(60)
<223> OTHER INFORMATION: CDRL2
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (93)..(101)
<223> OTHER INFORMATION: CDRL3

```

```

<400> SEQUENCE: 26

```

```

Asp Ile Val Met Thr Gln Ser Pro Asp Ser Leu Ala Val Ser Leu Gly
1             5             10             15

Glu Arg Ala Thr Ile Asn Cys Arg Ala Ser Lys Asn Val Asp Ser Tyr
20             25             30

Gly Asn Ser Phe Met His Trp Tyr Gln Gln Lys Pro Gly Gln Pro Pro
35             40             45

Lys Leu Leu Ile Tyr Arg Ala Ser Asn Leu Glu Ser Gly Val Pro Asp
50             55             60

Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser
65             70             75             80

Ser Leu Gln Ala Glu Asp Val Ala Val Tyr Tyr Cys Gln Gln Ser Asn
85             90             95

Glu Asp Pro Leu Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys Arg
100            105            110

Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln
115            120            125

Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr
130            135            140

Pro Arg Glu Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser
145            150            155            160

Gly Asn Ser Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr
165            170            175

Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys
180            185            190

His Lys Val Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro
195            200            205

Val Thr Lys Ser Phe Asn Arg Gly Glu Cys
210            215

```

```

<210> SEQ ID NO 27
<211> LENGTH: 446
<212> TYPE: PRT

```

-continued

```

<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: humanized antibody with mouse CDRs and human
        framework and constant regions
<220> FEATURE:
<221> NAME/KEY: DOMAIN
<222> LOCATION: (1)..(117)
<223> OTHER INFORMATION: variable domain
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (31)..(35)
<223> OTHER INFORMATION: CDRH1
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (50)..(66)
<223> OTHER INFORMATION: CDRH2
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (99)..(106)
<223> OTHER INFORMATION: CDRH3

<400> SEQUENCE: 27

Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ser
1          5          10          15
Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asp Leu
20         25         30
Ser Met His Trp Ile Lys Gln Arg Pro Gly Gln Gly Leu Glu Trp Ile
35         40         45
Gly Tyr Ile Thr Pro Ser Ser Gly Tyr Thr Ala Tyr Asn Gln Lys Phe
50         55         60
Lys Gly Lys Thr Thr Leu Thr Ala Asp Lys Ser Thr Ser Thr Ala Tyr
65         70         75         80
Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Tyr Cys
85         90         95
Ala Arg Leu Ile Leu Gln Arg Gly Ala Tyr Trp Gly Gln Gly Thr Leu
100        105        110
Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu
115        120        125
Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly Cys
130        135        140
Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser
145        150        155        160
Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu Gln Ser
165        170        175
Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser
180        185        190
Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys Pro Ser Asn
195        200        205
Thr Lys Val Asp Lys Arg Val Glu Pro Lys Ser Cys Asp Lys Thr His
210        215        220
Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser Val
225        230        235        240
Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr
245        250        255
Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu Asp Pro Glu
260        265        270
Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala Lys
275        280        285

```

-continued

Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser
 290 295 300
 Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys
 305 310 315 320
 Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile
 325 330 335
 Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro
 340 345 350
 Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr Cys Leu
 355 360 365
 Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn
 370 375 380
 Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser
 385 390 395 400
 Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg
 405 410 415
 Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu
 420 425 430
 His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly
 435 440 445

<210> SEQ ID NO 28
 <211> LENGTH: 218
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: humanized antibody with mouse CDRs and human
 framework and constant regions
 <220> FEATURE:
 <221> NAME/KEY: DOMAIN
 <222> LOCATION: (1)..(111)
 <223> OTHER INFORMATION: variable domain
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (24)..(38)
 <223> OTHER INFORMATION: CDRL1
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (54)..(60)
 <223> OTHER INFORMATION: CDRL2
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (93)..(101)
 <223> OTHER INFORMATION: CDRL3

<400> SEQUENCE: 28

Asp Ile Val Leu Thr Gln Ser Pro Asp Ser Leu Ala Val Ser Leu Gly
 1 5 10 15
 Glu Arg Ala Thr Ile Ser Cys Arg Ala Ser Lys Asn Val Asp Ser Tyr
 20 25 30
 Gly Asn Ser Phe Met His Trp Tyr Gln Gln Lys Pro Gly Gln Pro Pro
 35 40 45
 Lys Leu Leu Ile Tyr Arg Ala Ser Asn Leu Glu Ser Gly Val Pro Asp
 50 55 60
 Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser
 65 70 75 80
 Ser Leu Gln Ala Glu Asp Val Ala Val Tyr Tyr Cys Gln Gln Ser Asn
 85 90 95

-continued

Glu	Asp	Pro	Leu	Thr	Phe	Gly	Gln	Gly	Thr	Lys	Leu	Glu	Ile	Lys	Arg
			100					105					110		
Thr	Val	Ala	Ala	Pro	Ser	Val	Phe	Ile	Phe	Pro	Pro	Ser	Asp	Glu	Gln
		115					120					125			
Leu	Lys	Ser	Gly	Thr	Ala	Ser	Val	Val	Cys	Leu	Leu	Asn	Asn	Phe	Tyr
	130					135					140				
Pro	Arg	Glu	Ala	Lys	Val	Gln	Trp	Lys	Val	Asp	Asn	Ala	Leu	Gln	Ser
145					150					155					160
Gly	Asn	Ser	Gln	Glu	Ser	Val	Thr	Glu	Gln	Asp	Ser	Lys	Asp	Ser	Thr
			165					170						175	
Tyr	Ser	Leu	Ser	Ser	Thr	Leu	Thr	Leu	Ser	Lys	Ala	Asp	Tyr	Glu	Lys
		180					185						190		
His	Lys	Val	Tyr	Ala	Cys	Glu	Val	Thr	His	Gln	Gly	Leu	Ser	Ser	Pro
	195						200					205			
Val	Thr	Lys	Ser	Phe	Asn	Arg	Gly	Glu	Cys						
	210					215									

<210> SEQ ID NO 29
 <211> LENGTH: 446
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: humanized antibody with mouse CDRs and human framework and constant regions
 <220> FEATURE:
 <221> NAME/KEY: DOMAIN
 <222> LOCATION: (1)..(117)
 <223> OTHER INFORMATION: variable domain
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (31)..(35)
 <223> OTHER INFORMATION: CDRH1
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (50)..(66)
 <223> OTHER INFORMATION: CDRH2
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (99)..(106)
 <223> OTHER INFORMATION: CDRH3
 <400> SEQUENCE: 29

Gln	Val	Gln	Leu	Val	Gln	Ser	Gly	Ala	Glu	Val	Lys	Lys	Pro	Gly	Ser
1			5						10					15	
Ser	Val	Lys	Val	Ser	Cys	Lys	Ala	Ser	Gly	Tyr	Thr	Phe	Thr	Asp	Leu
		20						25					30		
Ser	Met	His	Trp	Val	Arg	Gln	Ala	Pro	Gly	Gln	Gly	Leu	Glu	Trp	Met
	35					40						45			
Gly	Tyr	Ile	Thr	Pro	Ser	Ser	Gly	Tyr	Thr	Ala	Tyr	Asn	Gln	Lys	Phe
	50					55					60				
Lys	Gly	Lys	Thr	Thr	Leu	Thr	Ala	Asp	Lys	Ser	Thr	Ser	Thr	Ala	Tyr
65				70					75					80	
Met	Glu	Leu	Ser	Ser	Leu	Arg	Ser	Glu	Asp	Thr	Ala	Val	Tyr	Tyr	Cys
			85					90					95		
Ala	Arg	Leu	Ile	Leu	Gln	Arg	Gly	Ala	Tyr	Trp	Gly	Gln	Gly	Thr	Leu
		100					105					110			
Val	Thr	Val	Ser	Ser	Ala	Ser	Thr	Lys	Gly	Pro	Ser	Val	Phe	Pro	Leu
		115					120					125			
Ala	Pro	Ser	Ser	Lys	Ser	Thr	Ser	Gly	Gly	Thr	Ala	Ala	Leu	Gly	Cys

-continued

130	135	140
Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser		
145	150	155 160
Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu Gln Ser		
	165	170 175
Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser		
	180	185 190
Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys Pro Ser Asn		
	195	200 205
Thr Lys Val Asp Lys Arg Val Glu Pro Lys Ser Cys Asp Lys Thr His		
	210	215 220
Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser Val		
	225	230 235 240
Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr		
	245	250 255
Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu Asp Pro Glu		
	260	265 270
Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala Lys		
	275	280 285
Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser		
	290	295 300
Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys		
	305	310 315 320
Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile		
	325	330 335
Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro		
	340	345 350
Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr Cys Leu		
	355	360 365
Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn		
	370	375 380
Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser		
	385	390 395 400
Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg		
	405	410 415
Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu		
	420	425 430
His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly		
	435	440 445

<210> SEQ ID NO 30

<211> LENGTH: 218

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: humanized antibody with mouse CDRs and human framework and constant regions

<220> FEATURE:

<221> NAME/KEY: DOMAIN

<222> LOCATION: (1)..(111)

<223> OTHER INFORMATION: variable domain

<220> FEATURE:

<221> NAME/KEY: MISC_FEATURE

<222> LOCATION: (24)..(38)

<223> OTHER INFORMATION: CDRL1

<220> FEATURE:

-continued

<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (54)..(60)
<223> OTHER INFORMATION: CDRL2
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (93)..(101)
<223> OTHER INFORMATION: CDRL3

<400> SEQUENCE: 30

Asp Ile Val Leu Thr Gln Ser Pro Asp Ser Leu Ala Val Ser Leu Gly
1 5 10 15
Glu Arg Ala Thr Ile Asn Cys Arg Ala Ser Lys Asn Val Asp Ser Tyr
20 25 30
Gly Asn Ser Phe Met His Trp Tyr Gln Gln Lys Pro Gly Gln Pro Pro
35 40 45
Lys Leu Leu Ile Tyr Arg Ala Ser Asn Leu Glu Ser Gly Val Pro Asp
50 55 60
Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser
65 70 75 80
Ser Leu Gln Ala Glu Asp Val Ala Val Tyr Tyr Cys Gln Gln Ser Asn
85 90 95
Glu Asp Pro Leu Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys Arg
100 105 110
Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln
115 120 125
Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr
130 135 140
Pro Arg Glu Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser
145 150 155 160
Gly Asn Ser Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr
165 170 175
Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys
180 185 190
His Lys Val Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro
195 200 205
Val Thr Lys Ser Phe Asn Arg Gly Glu Cys
210 215

<210> SEQ ID NO 31
<211> LENGTH: 446
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: humanized antibody with mouse CDRs and human
framework and constant regions
<220> FEATURE:
<221> NAME/KEY: DOMAIN
<222> LOCATION: (1)..(117)
<223> OTHER INFORMATION: variable domain
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (31)..(35)
<223> OTHER INFORMATION: CDRH1
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (50)..(66)
<223> OTHER INFORMATION: CDRH2
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (99)..(106)
<223> OTHER INFORMATION: CDRH3

-continued

<400> SEQUENCE: 31

Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ser
 1 5 10 15
 Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asp Leu
 20 25 30
 Ser Met His Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met
 35 40 45
 Gly Tyr Ile Thr Pro Ser Ser Gly Tyr Thr Ala Tyr Asn Gln Lys Phe
 50 55 60
 Lys Gly Lys Thr Thr Leu Thr Ala Asp Lys Ser Thr Ser Thr Ala Tyr
 65 70 75 80
 Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95
 Ala Arg Leu Ile Leu Gln Arg Gly Ala Tyr Trp Gly Gln Gly Thr Leu
 100 105 110
 Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu
 115 120 125
 Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly Cys
 130 135 140
 Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser
 145 150 155 160
 Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu Gln Ser
 165 170 175
 Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser
 180 185 190
 Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys Pro Ser Asn
 195 200 205
 Thr Lys Val Asp Lys Arg Val Glu Pro Lys Ser Cys Asp Lys Thr His
 210 215 220
 Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser Val
 225 230 235 240
 Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr
 245 250 255
 Pro Glu Val Thr Cys Val Val Val Asp Val Glu His Glu Asp Pro Glu
 260 265 270
 Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala Lys
 275 280 285
 Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser
 290 295 300
 Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys
 305 310 315 320
 Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile
 325 330 335
 Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro
 340 345 350
 Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr Cys Leu
 355 360 365
 Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn
 370 375 380
 Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser

-continued

385	390	395	400
Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg	405	410	415
Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu	420	425	430
His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly	435	440	445

<210> SEQ ID NO 32
 <211> LENGTH: 446
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: humanized antibody with mouse CDRs and human framework and constant regions
 <220> FEATURE:
 <221> NAME/KEY: DOMAIN
 <222> LOCATION: (1)..(117)
 <223> OTHER INFORMATION: variable domain
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (31)..(35)
 <223> OTHER INFORMATION: CDRH1
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (50)..(66)
 <223> OTHER INFORMATION: CDRH2
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (99)..(106)
 <223> OTHER INFORMATION: CDRH3

<400> SEQUENCE: 32

Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ser	1	5	10	15
Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asp Leu	20	25	30	
Ser Met His Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met	35	40	45	
Gly Tyr Ile Thr Pro Ser Ser Gly Tyr Thr Ala Tyr Asn Gln Lys Phe	50	55	60	
Lys Gly Lys Thr Thr Leu Thr Ala Asp Lys Ser Thr Ser Thr Ala Tyr	65	70	75	80
Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Tyr Cys	85	90	95	
Ala Arg Leu Ile Leu Gln Arg Gly Ala Tyr Trp Gly Gln Gly Thr Leu	100	105	110	
Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu	115	120	125	
Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly Cys	130	135	140	
Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser	145	150	155	160
Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu Gln Ser	165	170	175	
Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser	180	185	190	
Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys Pro Ser Asn	195	200	205	

-continued

Thr Lys Val Asp Lys Arg Val Glu Pro Lys Ser Cys Asp Lys Thr His
 210 215 220
 Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser Val
 225 230 235 240
 Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr
 245 250 255
 Pro Glu Val Thr Cys Val Val Val Asp Val Glu His Glu Asp Pro Glu
 260 265 270
 Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala Lys
 275 280 285
 Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser
 290 295 300
 Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys
 305 310 315 320
 Cys Lys Val Ser Asn Lys Ala Phe Pro Ala Pro Ile Glu Lys Thr Ile
 325 330 335
 Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro
 340 345 350
 Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr Cys Leu
 355 360 365
 Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn
 370 375 380
 Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser
 385 390 395 400
 Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg
 405 410 415
 Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu
 420 425 430
 His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly
 435 440 445

<210> SEQ ID NO 33
 <211> LENGTH: 446
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: humanized antibody with mouse CDRs and human
 framework and constant regions
 <220> FEATURE:
 <221> NAME/KEY: DOMAIN
 <222> LOCATION: (1)..(117)
 <223> OTHER INFORMATION: variable domain
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (31)..(35)
 <223> OTHER INFORMATION: CDRH1
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (50)..(66)
 <223> OTHER INFORMATION: CDRH2
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (99)..(106)
 <223> OTHER INFORMATION: CDRH3

<400> SEQUENCE: 33

Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ser
 1 5 10 15

-continued

Ser	Val	Lys	Val	Ser	Cys	Lys	Ala	Ser	Gly	Tyr	Thr	Phe	Thr	Asp	Leu	20	25	30
Ser	Met	His	Trp	Val	Arg	Gln	Ala	Pro	Gly	Gln	Gly	Leu	Glu	Trp	Met	35	40	45
Gly	Tyr	Ile	Thr	Pro	Ser	Ser	Gly	Tyr	Thr	Ala	Tyr	Asn	Gln	Lys	Phe	50	55	60
Lys	Gly	Lys	Thr	Thr	Leu	Thr	Ala	Asp	Lys	Ser	Thr	Ser	Thr	Ala	Tyr	65	70	80
Met	Glu	Leu	Ser	Ser	Leu	Arg	Ser	Glu	Asp	Thr	Ala	Val	Tyr	Tyr	Cys	85	90	95
Ala	Arg	Leu	Ile	Leu	Gln	Arg	Gly	Ala	Tyr	Trp	Gly	Gln	Gly	Thr	Leu	100	105	110
Val	Thr	Val	Ser	Ser	Ala	Ser	Thr	Lys	Gly	Pro	Ser	Val	Phe	Pro	Leu	115	120	125
Ala	Pro	Ser	Ser	Lys	Ser	Thr	Ser	Gly	Gly	Thr	Ala	Ala	Leu	Gly	Cys	130	135	140
Leu	Val	Lys	Asp	Tyr	Phe	Pro	Glu	Pro	Val	Thr	Val	Ser	Trp	Asn	Ser	145	150	160
Gly	Ala	Leu	Thr	Ser	Gly	Val	His	Thr	Phe	Pro	Ala	Val	Leu	Gln	Ser	165	170	175
Ser	Gly	Leu	Tyr	Ser	Leu	Ser	Ser	Val	Val	Thr	Val	Pro	Ser	Ser	Ser	180	185	190
Leu	Gly	Thr	Gln	Thr	Tyr	Ile	Cys	Asn	Val	Asn	His	Lys	Pro	Ser	Asn	195	200	205
Thr	Lys	Val	Asp	Lys	Arg	Val	Glu	Pro	Lys	Ser	Cys	Asp	Lys	Thr	His	210	215	220
Thr	Cys	Pro	Pro	Cys	Pro	Ala	Pro	Glu	Leu	Leu	Gly	Gly	Asp	Ser	Val	225	230	235
Phe	Leu	Phe	Pro	Pro	Lys	Pro	Lys	Asp	Thr	Leu	Met	Ile	Ser	Arg	Thr	245	250	255
Pro	Glu	Val	Thr	Cys	Val	Val	Val	Asp	Val	Ser	His	Glu	Asp	Gly	Glu	260	265	270
Val	Lys	Phe	Asn	Trp	Tyr	Val	Asp	Gly	Val	Glu	Val	His	Asn	Ala	Lys	275	280	285
Thr	Lys	Pro	Arg	Glu	Glu	Gln	Tyr	Asn	Ser	Thr	Tyr	Arg	Val	Val	Ser	290	295	300
Val	Leu	Thr	Val	Leu	His	Gln	Asp	Trp	Leu	Asn	Gly	Lys	Glu	Tyr	Lys	305	310	315
Cys	Lys	Val	Ser	Asn	Lys	Ala	Leu	Pro	Ala	Pro	Ile	Glu	Lys	Thr	Ile	325	330	335
Ser	Lys	Ala	Lys	Gly	Gln	Pro	Arg	Glu	Pro	Gln	Val	Tyr	Thr	Leu	Pro	340	345	350
Pro	Ser	Arg	Glu	Glu	Met	Thr	Lys	Asn	Gln	Val	Ser	Leu	Thr	Cys	Leu	355	360	365
Val	Lys	Gly	Phe	Tyr	Pro	Ser	Asp	Ile	Ala	Val	Glu	Trp	Glu	Ser	Asn	370	375	380
Gly	Gln	Pro	Glu	Asn	Asn	Tyr	Lys	Thr	Thr	Pro	Val	Leu	Asp	Ser		385	390	395
Asp	Gly	Ser	Phe	Phe	Leu	Tyr	Ser	Lys	Leu	Thr	Val	Asp	Lys	Ser	Arg	405	410	415
Trp	Gln	Gln	Gly	Asn	Val	Phe	Ser	Cys	Ser	Val	Met	His	Glu	Ala	Leu			

-continued

420	425	430
His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly		
435	440	445
<210> SEQ ID NO 34		
<211> LENGTH: 446		
<212> TYPE: PRT		
<213> ORGANISM: Artificial Sequence		
<220> FEATURE:		
<223> OTHER INFORMATION: humanized antibody with mouse CDRs and human framework and constant regions		
<400> SEQUENCE: 34		
Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ser		
1	5	10
Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asp Leu		
20	25	30
Ser Met His Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met		
35	40	45
Gly Tyr Ile Thr Pro Ser Ser Gly Tyr Thr Ala Tyr Asn Gln Lys Phe		
50	55	60
Lys Gly Lys Thr Thr Leu Thr Ala Asp Lys Ser Thr Ser Thr Ala Tyr		
65	70	75
Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Tyr Cys		
85	90	95
Ala Arg Leu Ile Leu Gln Arg Gly Ala Tyr Trp Gly Gln Gly Thr Leu		
100	105	110
Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu		
115	120	125
Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly Cys		
130	135	140
Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser		
145	150	155
Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu Gln Ser		
165	170	175
Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser		
180	185	190
Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys Pro Ser Asn		
195	200	205
Thr Lys Val Asp Lys Arg Val Glu Pro Lys Ser Cys Asp Lys Thr His		
210	215	220
Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly Asp Ser Val		
225	230	235
Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr		
245	250	255
Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu Glu Gly Glu		
260	265	270
Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala Lys		
275	280	285
Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser		
290	295	300
Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys		
305	310	315
		320

-continued

Cys	Lys	Val	Ser	Asn	Lys	Ala	Leu	Pro	Ala	Pro	Ile	Glu	Lys	Thr	Ile
				325					330					335	
Ser	Lys	Ala	Lys	Gly	Gln	Pro	Arg	Glu	Pro	Gln	Val	Tyr	Thr	Leu	Pro
			340					345					350		
Pro	Ser	Arg	Glu	Glu	Met	Thr	Lys	Asn	Gln	Val	Ser	Leu	Thr	Cys	Leu
			355				360					365			
Val	Lys	Gly	Phe	Tyr	Pro	Ser	Asp	Ile	Ala	Val	Glu	Trp	Glu	Ser	Asn
	370					375					380				
Gly	Gln	Pro	Glu	Asn	Asn	Tyr	Lys	Thr	Thr	Pro	Pro	Val	Leu	Asp	Ser
385				390						395				400	
Asp	Gly	Ser	Phe	Phe	Leu	Tyr	Ser	Lys	Leu	Thr	Val	Asp	Lys	Ser	Arg
			405					410					415		
Trp	Gln	Gln	Gly	Asn	Val	Phe	Ser	Cys	Ser	Val	Met	His	Glu	Ala	Leu
			420					425				430			
His	Asn	His	Tyr	Thr	Gln	Lys	Ser	Leu	Ser	Leu	Ser	Pro	Gly		
		435					440					445			

<210> SEQ ID NO 35

<211> LENGTH: 446

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: humanized antibody with mouse CDRs and human framework and constant regions

<400> SEQUENCE: 35

Gln	Val	Gln	Leu	Val	Gln	Ser	Gly	Ala	Glu	Val	Lys	Lys	Pro	Gly	Ser
1			5					10					15		
Ser	Val	Lys	Val	Ser	Cys	Lys	Ala	Ser	Gly	Tyr	Thr	Phe	Thr	Asp	Leu
	20					25						30			
Ser	Met	His	Trp	Val	Arg	Gln	Ala	Pro	Gly	Gln	Gly	Leu	Glu	Trp	Met
	35				40						45				
Gly	Tyr	Ile	Thr	Pro	Ser	Ser	Gly	Tyr	Thr	Ala	Tyr	Asn	Gln	Lys	Phe
50					55					60					
Lys	Gly	Lys	Thr	Thr	Leu	Thr	Ala	Asp	Lys	Ser	Thr	Ser	Thr	Ala	Tyr
65				70					75					80	
Met	Glu	Leu	Ser	Ser	Leu	Arg	Ser	Glu	Asp	Thr	Ala	Val	Tyr	Tyr	Cys
		85						90					95		
Ala	Arg	Leu	Ile	Leu	Gln	Arg	Gly	Ala	Tyr	Trp	Gly	Gln	Gly	Thr	Leu
	100						105					110			
Val	Thr	Val	Ser	Ser	Ala	Ser	Thr	Lys	Gly	Pro	Ser	Val	Phe	Pro	Leu
	115					120						125			
Ala	Pro	Ser	Ser	Lys	Ser	Thr	Ser	Gly	Gly	Thr	Ala	Ala	Leu	Gly	Cys
	130				135					140					
Leu	Val	Lys	Asp	Tyr	Phe	Pro	Glu	Pro	Val	Thr	Val	Ser	Trp	Asn	Ser
145				150						155				160	
Gly	Ala	Leu	Thr	Ser	Gly	Val	His	Thr	Phe	Pro	Ala	Val	Leu	Gln	Ser
		165						170					175		
Ser	Gly	Leu	Tyr	Ser	Leu	Ser	Ser	Val	Val	Thr	Val	Pro	Ser	Ser	Ser
		180						185					190		
Leu	Gly	Thr	Gln	Thr	Tyr	Ile	Cys	Asn	Val	Asn	His	Lys	Pro	Ser	Asn
	195					200						205			
Thr	Lys	Val	Asp	Lys	Arg	Val	Glu	Pro	Lys	Ser	Cys	Asp	Lys	Thr	His
	210					215					220				

-continued

Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Asp Asp Ser Val
 225 230 235 240
 Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr
 245 250 255
 Pro Glu Val Thr Cys Val Val Val Asp Val Ser Asp Glu Asp Gly Glu
 260 265 270
 Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala Lys
 275 280 285
 Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser
 290 295 300
 Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys
 305 310 315 320
 Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile
 325 330 335
 Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro
 340 345 350
 Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr Cys Leu
 355 360 365
 Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn
 370 375 380
 Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser
 385 390 395 400
 Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg
 405 410 415
 Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu
 420 425 430
 His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly
 435 440 445

<210> SEQ ID NO 36
 <211> LENGTH: 446
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: humanized antibody with mouse CDRs and human
 framework and constant regions
 <220> FEATURE:
 <221> NAME/KEY: DOMAIN
 <222> LOCATION: (1)..(117)
 <223> OTHER INFORMATION: variable domain
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (31)..(35)
 <223> OTHER INFORMATION: CDRH1
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (50)..(66)
 <223> OTHER INFORMATION: CDRH2
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (99)..(106)
 <223> OTHER INFORMATION: CDRH3
 <400> SEQUENCE: 36

Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ser
 1 5 10 15
 Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asp Leu
 20 25 30

-continued

Ser	Met	His	Trp	Val	Arg	Gln	Ala	Pro	Gly	Gln	Gly	Leu	Glu	Trp	Met
		35					40					45			
Gly	Tyr	Ile	Thr	Pro	Ser	Ser	Gly	Tyr	Thr	Ala	Tyr	Asn	Gln	Lys	Phe
	50					55					60				
Lys	Gly	Lys	Thr	Thr	Leu	Thr	Ala	Asp	Lys	Ser	Thr	Ser	Thr	Ala	Tyr
65					70					75					80
Met	Glu	Leu	Ser	Ser	Leu	Arg	Ser	Glu	Asp	Thr	Ala	Val	Tyr	Tyr	Cys
			85						90					95	
Ala	Arg	Leu	Ile	Leu	Gln	Arg	Gly	Ala	Tyr	Trp	Gly	Gln	Gly	Thr	Leu
			100					105					110		
Val	Thr	Val	Ser	Ser	Ala	Ser	Thr	Lys	Gly	Pro	Ser	Val	Phe	Pro	Leu
			115					120					125		
Ala	Pro	Ser	Ser	Lys	Ser	Thr	Ser	Gly	Gly	Thr	Ala	Ala	Leu	Gly	Cys
	130					135					140				
Leu	Val	Lys	Asp	Tyr	Phe	Pro	Glu	Pro	Val	Thr	Val	Ser	Trp	Asn	Ser
145					150					155					160
Gly	Ala	Leu	Thr	Ser	Gly	Val	His	Thr	Phe	Pro	Ala	Val	Leu	Gln	Ser
				165					170					175	
Ser	Gly	Leu	Tyr	Ser	Leu	Ser	Ser	Val	Val	Thr	Val	Pro	Ser	Ser	Ser
		180						185					190		
Leu	Gly	Thr	Gln	Thr	Tyr	Ile	Cys	Asn	Val	Asn	His	Lys	Pro	Ser	Asn
		195					200					205			
Thr	Lys	Val	Asp	Lys	Arg	Val	Glu	Pro	Lys	Ser	Cys	Asp	Lys	Thr	His
	210					215					220				
Thr	Cys	Pro	Pro	Cys	Pro	Ala	Pro	Glu	Leu	Leu	Gly	Asp	Asp	Ser	Val
225					230						235				240
Phe	Leu	Phe	Pro	Pro	Lys	Pro	Lys	Asp	Thr	Leu	Met	Ile	Ser	Arg	Thr
					245				250					255	
Pro	Glu	Val	Thr	Cys	Val	Val	Val	Asp	Val	Ser	His	Glu	Asp	Gly	Glu
			260					265					270		
Val	Lys	Phe	Asn	Trp	Tyr	Val	Asp	Gly	Val	Glu	Val	His	Asn	Ala	Lys
		275					280					285			
Thr	Lys	Pro	Arg	Glu	Glu	Gln	Tyr	Asn	Ser	Thr	Tyr	Arg	Val	Val	Ser
	290					295					300				
Val	Leu	Thr	Val	Leu	His	Gln	Asp	Trp	Leu	Asn	Gly	Lys	Glu	Tyr	Lys
305					310						315				320
Cys	Lys	Val	Ser	Asn	Lys	Ala	Leu	Pro	Arg	Pro	Ile	Glu	Lys	Thr	Ile
				325					330					335	
Ser	Lys	Ala	Lys	Gly	Gln	Pro	Arg	Glu	Pro	Gln	Val	Tyr	Thr	Leu	Pro
			340					345					350		
Pro	Ser	Arg	Glu	Glu	Met	Thr	Lys	Asn	Gln	Val	Ser	Leu	Thr	Cys	Leu
		355					360					365			
Val	Lys	Gly	Phe	Tyr	Pro	Ser	Asp	Ile	Ala	Val	Glu	Trp	Glu	Ser	Asn
	370					375					380				
Gly	Gln	Pro	Glu	Asn	Asn	Tyr	Lys	Thr	Thr	Pro	Pro	Val	Leu	Asp	Ser
385					390					395					400
Asp	Gly	Ser	Phe	Phe	Leu	Tyr	Ser	Lys	Leu	Thr	Val	Asp	Lys	Ser	Arg
				405					410					415	
Trp	Gln	Gln	Gly	Asn	Val	Phe	Ser	Cys	Ser	Val	Met	His	Glu	Ala	Leu
			420					425					430		
His	Asn	His	Tyr	Thr	Gln	Lys	Ser	Leu	Ser	Leu	Ser	Pro	Gly		

-continued

435	440	445
<210> SEQ ID NO 37		
<211> LENGTH: 446		
<212> TYPE: PRT		
<213> ORGANISM: Artificial Sequence		
<220> FEATURE:		
<223> OTHER INFORMATION: humanized antibody with mouse CDRs and human framework and constant regions		
<400> SEQUENCE: 37		
Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ser		
1 5 10 15		
Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asp Leu		
20 25 30		
Ser Met His Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met		
35 40 45		
Gly Tyr Ile Thr Pro Ser Ser Gly Tyr Thr Ala Tyr Asn Gln Lys Phe		
50 55 60		
Lys Gly Lys Thr Thr Leu Thr Ala Asp Lys Ser Thr Ser Thr Ala Tyr		
65 70 75 80		
Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Tyr Cys		
85 90 95		
Ala Arg Leu Ile Leu Gln Arg Gly Ala Tyr Trp Gly Gln Gly Thr Leu		
100 105 110		
Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu		
115 120 125		
Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly Cys		
130 135 140		
Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser		
145 150 155 160		
Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu Gln Ser		
165 170 175		
Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser		
180 185 190		
Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys Pro Ser Asn		
195 200 205		
Thr Lys Val Asp Lys Arg Val Glu Pro Lys Ser Cys Asp Lys Thr His		
210 215 220		
Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Asp Asp Ser Val		
225 230 235 240		
Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr		
245 250 255		
Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu Glu Gly Glu		
260 265 270		
Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala Lys		
275 280 285		
Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser		
290 295 300		
Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys		
305 310 315 320		
Cys Lys Val Ser Asn Lys Ala Leu Pro Arg Pro Ile Glu Lys Thr Ile		
325 330 335		

-continued

Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro
 340 345 350
 Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr Cys Leu
 355 360 365
 Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn
 370 375 380
 Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser
 385 390 395 400
 Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg
 405 410 415
 Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu
 420 425 430
 His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly
 435 440 445

<210> SEQ ID NO 38
 <211> LENGTH: 446
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: humanized antibody with mouse CDRs and human
 framework and constant regions
 <220> FEATURE:
 <221> NAME/KEY: DOMAIN
 <222> LOCATION: (1)..(117)
 <223> OTHER INFORMATION: variable domain
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (31)..(35)
 <223> OTHER INFORMATION: CDRH1
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (50)..(66)
 <223> OTHER INFORMATION: CDRH2
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (99)..(106)
 <223> OTHER INFORMATION: CDRH3

<400> SEQUENCE: 38

Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ser
 1 5 10 15
 Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asp Leu
 20 25 30
 Ser Met His Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met
 35 40 45
 Gly Tyr Ile Thr Pro Ser Ser Gly Tyr Thr Ala Tyr Asn Gln Lys Phe
 50 55 60
 Lys Gly Lys Thr Thr Leu Thr Ala Asp Lys Ser Thr Ser Thr Ala Tyr
 65 70 75 80
 Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95
 Ala Arg Leu Ile Leu Gln Arg Gly Ala Tyr Trp Gly Gln Gly Thr Leu
 100 105 110
 Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu
 115 120 125
 Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly Cys
 130 135 140
 Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser

-continued

145	150	155	160
Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu Gln Ser	165	170	175
Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser	180	185	190
Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys Pro Ser Asn	195	200	205
Thr Lys Val Asp Lys Arg Val Glu Pro Lys Ser Cys Asp Lys Thr His	210	215	220
Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Asp Asp Ser Val	225	230	235
Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr	245	250	255
Pro Glu Val Thr Cys Val Val Val Asp Val Ser Asp Glu Asp Gly Glu	260	265	270
Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala Lys	275	280	285
Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser	290	295	300
Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys	305	310	315
Cys Lys Val Ser Asn Lys Ala Leu Pro Arg Pro Ile Glu Lys Thr Ile	325	330	335
Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro	340	345	350
Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr Cys Leu	355	360	365
Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn	370	375	380
Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser	385	390	395
Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg	405	410	415
Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu	420	425	430
His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly	435	440	445

<210> SEQ ID NO 39

<211> LENGTH: 446

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: humanized antibody with mouse CDRs and human framework and constant regions

<220> FEATURE:

<221> NAME/KEY: DOMAIN

<222> LOCATION: (1)..(117)

<223> OTHER INFORMATION: variable domain

<220> FEATURE:

<221> NAME/KEY: MISC_FEATURE

<222> LOCATION: (31)..(35)

<223> OTHER INFORMATION: CDRH1

<220> FEATURE:

<221> NAME/KEY: MISC_FEATURE

<222> LOCATION: (50)..(66)

<223> OTHER INFORMATION: CDRH2

-continued

<220> FEATURE:

<221> NAME/KEY: MISC_FEATURE

<222> LOCATION: (99)..(106)

<223> OTHER INFORMATION: CDRH3

<400> SEQUENCE: 39

Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ser
1 5 10 15
Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asp Leu
20 25 30
Ser Met His Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met
35 40 45
Gly Tyr Ile Thr Pro Ser Ser Gly Tyr Thr Ala Tyr Asn Gln Lys Phe
50 55 60
Lys Gly Lys Thr Thr Leu Thr Ala Asp Lys Ser Thr Ser Thr Ala Tyr
65 70 75 80
Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Tyr Cys
85 90 95
Ala Arg Leu Ile Leu Gln Arg Gly Ala Tyr Trp Gly Gln Gly Thr Leu
100 105 110
Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu
115 120 125
Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly Cys
130 135 140
Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser
145 150 155 160
Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu Gln Ser
165 170 175
Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser
180 185 190
Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys Pro Ser Asn
195 200 205
Thr Lys Val Asp Lys Arg Val Glu Pro Lys Ser Cys Asp Lys Thr His
210 215 220
Thr Cys Pro Pro Cys Pro Ala Pro Asp Leu Leu Gly Asp Asp Ser Val
225 230 235 240
Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr
245 250 255
Pro Glu Val Thr Cys Val Val Val Asp Val Ser Asp Glu Asp Gly Glu
260 265 270
Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala Lys
275 280 285
Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser
290 295 300
Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys
305 310 315 320
Cys Lys Val Ser Asn Lys Ala Leu Pro Arg Pro Ile Glu Lys Thr Ile
325 330 335
Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro
340 345 350
Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val Ser Leu Thr Cys Leu
355 360 365

-continued

Val	Lys	Gly	Phe	Tyr	Pro	Ser	Asp	Ile	Ala	Val	Glu	Trp	Glu	Ser	Asn
370						375					380				
Gly	Gln	Pro	Glu	Asn	Asn	Tyr	Lys	Thr	Thr	Pro	Pro	Val	Leu	Asp	Ser
385					390					395					400
Asp	Gly	Ser	Phe	Phe	Leu	Tyr	Ser	Lys	Leu	Thr	Val	Asp	Lys	Ser	Arg
			405						410					415	
Trp	Gln	Gln	Gly	Asn	Val	Phe	Ser	Cys	Ser	Val	Met	His	Glu	Ala	Leu
			420					425					430		
His	Asn	His	Tyr	Thr	Gln	Lys	Ser	Leu	Ser	Leu	Ser	Pro	Gly		
		435					440					445			

<210> SEQ ID NO 40
 <211> LENGTH: 451
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: chimeric antibody with mouse variable domain
 and human constant domain
 <220> FEATURE:
 <221> NAME/KEY: DOMAIN
 <222> LOCATION: (1)..(122)
 <223> OTHER INFORMATION: variable domain
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (31)..(35)
 <223> OTHER INFORMATION: CDRH1
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (50)..(66)
 <223> OTHER INFORMATION: CDRH2
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (99)..(111)
 <223> OTHER INFORMATION: CDRH3

<400> SEQUENCE: 40

Gln	Val	Gln	Phe	Gln	Gln	Ser	Gly	Ala	Glu	Leu	Ala	Arg	Pro	Gly	Ala
1			5					10						15	
Ser	Val	Lys	Leu	Ser	Cys	Lys	Ala	Ser	Gly	Tyr	Thr	Phe	Thr	Ser	Tyr
		20					25					30			
Trp	Met	Gln	Trp	Val	Lys	Gln	Arg	Pro	Gly	Gln	Gly	Leu	Glu	Trp	Ile
	35				40						45				
Gly	Thr	Ile	Tyr	Pro	Gly	Asp	Gly	Asp	Ser	Arg	Tyr	Asn	Gln	Lys	Phe
	50				55					60					
Lys	Gly	Lys	Ala	Leu	Leu	Thr	Ala	Asp	Lys	Ser	Ser	Ser	Ile	Ala	Tyr
65				70					75					80	
Met	Gln	Leu	Asn	Ser	Leu	Ala	Ser	Glu	Asp	Ser	Ala	Val	Tyr	Phe	Cys
		85						90					95		
Ala	Arg	Phe	Ser	Leu	Tyr	Asp	Gly	Tyr	Pro	Tyr	Tyr	Phe	Asp	Tyr	Trp
		100					105					110			
Gly	Gln	Gly	Thr	Thr	Leu	Thr	Val	Ser	Ser	Ala	Ser	Thr	Lys	Gly	Pro
	115					120						125			
Ser	Val	Phe	Pro	Leu	Ala	Pro	Ser	Ser	Lys	Ser	Thr	Ser	Gly	Gly	Thr
	130				135						140				
Ala	Ala	Leu	Gly	Cys	Leu	Val	Lys	Asp	Tyr	Phe	Pro	Glu	Pro	Val	Thr
145				150					155					160	
Val	Ser	Trp	Asn	Ser	Gly	Ala	Leu	Thr	Ser	Gly	Val	His	Thr	Phe	Pro
			165				170						175		
Ala	Val	Leu	Gln	Ser	Ser	Gly	Leu	Tyr	Ser	Leu	Ser	Ser	Val	Val	Thr

-continued

180					185					190					
Val	Pro	Ser	Ser	Ser	Leu	Gly	Thr	Gln	Thr	Tyr	Ile	Cys	Asn	Val	Asn
	195						200					205			
His	Lys	Pro	Ser	Asn	Thr	Lys	Val	Asp	Lys	Arg	Val	Glu	Pro	Lys	Ser
	210					215					220				
Cys	Asp	Lys	Thr	His	Thr	Cys	Pro	Pro	Cys	Pro	Ala	Pro	Glu	Leu	Leu
225					230					235				240	
Gly	Gly	Pro	Ser	Val	Phe	Leu	Phe	Pro	Pro	Lys	Pro	Lys	Asp	Thr	Leu
				245					250					255	
Met	Ile	Ser	Arg	Thr	Pro	Glu	Val	Thr	Cys	Val	Val	Val	Asp	Val	Ser
			260					265					270		
His	Glu	Asp	Pro	Glu	Val	Lys	Phe	Asn	Trp	Tyr	Val	Asp	Gly	Val	Glu
	275						280					285			
Val	His	Asn	Ala	Lys	Thr	Lys	Pro	Arg	Glu	Glu	Gln	Tyr	Asn	Ser	Thr
	290					295					300				
Tyr	Arg	Val	Val	Ser	Val	Leu	Thr	Val	Leu	His	Gln	Asp	Trp	Leu	Asn
305					310					315				320	
Gly	Lys	Glu	Tyr	Lys	Cys	Lys	Val	Ser	Asn	Lys	Ala	Leu	Pro	Ala	Pro
				325					330					335	
Ile	Glu	Lys	Thr	Ile	Ser	Lys	Ala	Lys	Gly	Gln	Pro	Arg	Glu	Pro	Gln
			340					345					350		
Val	Tyr	Thr	Leu	Pro	Pro	Ser	Arg	Glu	Glu	Met	Thr	Lys	Asn	Gln	Val
	355						360					365			
Ser	Leu	Thr	Cys	Leu	Val	Lys	Gly	Phe	Tyr	Pro	Ser	Asp	Ile	Ala	Val
	370					375					380				
Glu	Trp	Glu	Ser	Asn	Gly	Gln	Pro	Glu	Asn	Asn	Tyr	Lys	Thr	Thr	Pro
385					390					395				400	
Pro	Val	Leu	Asp	Ser	Asp	Gly	Ser	Phe	Phe	Leu	Tyr	Ser	Lys	Leu	Thr
				405					410					415	
Val	Asp	Lys	Ser	Arg	Trp	Gln	Gln	Gly	Asn	Val	Phe	Ser	Cys	Ser	Val
		420						425					430		
Met	His	Glu	Ala	Leu	His	Asn	His	Tyr	Thr	Gln	Lys	Ser	Leu	Ser	Leu
	435					440						445			
Ser	Pro	Gly													
	450														

<210> SEQ ID NO 41

<211> LENGTH: 219

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: chimeric antibody with mouse variable domain
and human constant domain

<220> FEATURE:

<221> NAME/KEY: DOMAIN

<222> LOCATION: (1)..(112)

<223> OTHER INFORMATION: variable domain

<220> FEATURE:

<221> NAME/KEY: MISC_FEATURE

<222> LOCATION: (24)..(39)

<223> OTHER INFORMATION: CDRL1

<220> FEATURE:

<221> NAME/KEY: MISC_FEATURE

<222> LOCATION: (55)..(61)

<223> OTHER INFORMATION: CDRL2

<220> FEATURE:

<221> NAME/KEY: MISC_FEATURE

<222> LOCATION: (94)..(102)

-continued

<223> OTHER INFORMATION: CDRL3

<400> SEQUENCE: 41

Asp Val Val Met Thr Gln Thr Pro Leu Ser Leu Pro Val Ser Leu Gly
1 5 10 15
Asp Gln Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Leu Val His Arg
20 25 30
Asn Gly Asn Thr Tyr Leu His Trp Tyr Leu Gln Lys Pro Gly Gln Ser
35 40 45
Pro Lys Leu Leu Ile Tyr Arg Val Ser Asn Arg Phe Ser Gly Val Pro
50 55 60
Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile
65 70 75 80
Ser Arg Val Glu Ala Glu Asp Leu Gly Ile Tyr Phe Cys Ser Gln Ser
85 90 95
Thr His Phe Pro Tyr Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys
100 105 110
Arg Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu
115 120 125
Gln Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe
130 135 140
Tyr Pro Arg Glu Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln
145 150 155 160
Ser Gly Asn Ser Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser
165 170 175
Thr Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu
180 185 190
Lys His Lys Val Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser
195 200 205
Pro Val Thr Lys Ser Phe Asn Arg Gly Glu Cys
210 215

<210> SEQ ID NO 42

<211> LENGTH: 451

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: humanized antibody with mouse CDRs and human framework and constant regions

<220> FEATURE:

<221> NAME/KEY: DOMAIN

<222> LOCATION: (1)..(122)

<223> OTHER INFORMATION: variable domain

<220> FEATURE:

<221> NAME/KEY: MISC_FEATURE

<222> LOCATION: (31)..(35)

<223> OTHER INFORMATION: CDRH1

<220> FEATURE:

<221> NAME/KEY: MISC_FEATURE

<222> LOCATION: (50)..(66)

<223> OTHER INFORMATION: CDRH2

<220> FEATURE:

<221> NAME/KEY: MISC_FEATURE

<222> LOCATION: (99)..(111)

<223> OTHER INFORMATION: CDRH3

<400> SEQUENCE: 42

Gln Val Gln Phe Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ser
1 5 10 15

-continued

Ser	Val	Lys	Val	Ser	Cys	Lys	Ala	Ser	Gly	Tyr	Thr	Phe	Thr	Ser	Tyr
		20						25					30		
Trp	Met	Gln	Trp	Val	Arg	Gln	Ala	Pro	Gly	Gln	Gly	Leu	Glu	Trp	Met
		35					40					45			
Gly	Thr	Ile	Tyr	Pro	Gly	Asp	Gly	Asp	Ser	Arg	Tyr	Asn	Gln	Lys	Phe
	50					55					60				
Lys	Gly	Lys	Ala	Leu	Leu	Thr	Ala	Asp	Lys	Ser	Thr	Ser	Thr	Ala	Tyr
	65				70					75					80
Met	Glu	Leu	Ser	Ser	Leu	Arg	Ser	Glu	Asp	Thr	Ala	Val	Tyr	Tyr	Cys
			85					90						95	
Ala	Arg	Phe	Ser	Leu	Tyr	Asp	Gly	Tyr	Pro	Tyr	Tyr	Phe	Asp	Tyr	Trp
			100					105					110		
Gly	Gln	Gly	Thr	Thr	Val	Thr	Val	Ser	Ser	Ala	Ser	Thr	Lys	Gly	Pro
		115					120					125			
Ser	Val	Phe	Pro	Leu	Ala	Pro	Ser	Ser	Lys	Ser	Thr	Ser	Gly	Gly	Thr
	130					135					140				
Ala	Ala	Leu	Gly	Cys	Leu	Val	Lys	Asp	Tyr	Phe	Pro	Glu	Pro	Val	Thr
	145				150					155					160
Val	Ser	Trp	Asn	Ser	Gly	Ala	Leu	Thr	Ser	Gly	Val	His	Thr	Phe	Pro
			165						170					175	
Ala	Val	Leu	Gln	Ser	Ser	Gly	Leu	Tyr	Ser	Leu	Ser	Ser	Val	Val	Thr
		180						185					190		
Val	Pro	Ser	Ser	Ser	Leu	Gly	Thr	Gln	Thr	Tyr	Ile	Cys	Asn	Val	Asn
		195					200					205			
His	Lys	Pro	Ser	Asn	Thr	Lys	Val	Asp	Lys	Arg	Val	Glu	Pro	Lys	Ser
	210					215					220				
Cys	Asp	Lys	Thr	His	Thr	Cys	Pro	Pro	Cys	Pro	Ala	Pro	Glu	Leu	Leu
	225				230					235					240
Gly	Gly	Pro	Ser	Val	Phe	Leu	Phe	Pro	Pro	Lys	Pro	Lys	Asp	Thr	Leu
			245					250						255	
Met	Ile	Ser	Arg	Thr	Pro	Glu	Val	Thr	Cys	Val	Val	Val	Asp	Val	Ser
			260					265					270		
His	Glu	Asp	Pro	Glu	Val	Lys	Phe	Asn	Trp	Tyr	Val	Asp	Gly	Val	Glu
		275					280					285			
Val	His	Asn	Ala	Lys	Thr	Lys	Pro	Arg	Glu	Glu	Gln	Tyr	Asn	Ser	Thr
	290					295					300				
Tyr	Arg	Val	Val	Ser	Val	Leu	Thr	Val	Leu	His	Gln	Asp	Trp	Leu	Asn
	305				310					315					320
Gly	Lys	Glu	Tyr	Lys	Cys	Lys	Val	Ser	Asn	Lys	Ala	Leu	Pro	Ala	Pro
			325						330					335	
Ile	Glu	Lys	Thr	Ile	Ser	Lys	Ala	Lys	Gly	Gln	Pro	Arg	Glu	Pro	Gln
		340						345					350		
Val	Tyr	Thr	Leu	Pro	Pro	Ser	Arg	Glu	Glu	Met	Thr	Lys	Asn	Gln	Val
		355					360					365			
Ser	Leu	Thr	Cys	Leu	Val	Lys	Gly	Phe	Tyr	Pro	Ser	Asp	Ile	Ala	Val
	370					375					380				
Glu	Trp	Glu	Ser	Asn	Gly	Gln	Pro	Glu	Asn	Asn	Tyr	Lys	Thr	Thr	Pro
	385				390					395					400
Pro	Val	Leu	Asp	Ser	Asp	Gly	Ser	Phe	Phe	Leu	Tyr	Ser	Lys	Leu	Thr
			405					410						415	
Val	Asp	Lys	Ser	Arg	Trp	Gln	Gln	Gly	Asn	Val	Phe	Ser	Cys	Ser	Val

-continued

420	425	430
-----	-----	-----

Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu
435 440 445

Ser Pro Gly
450

<210> SEQ ID NO 43
<211> LENGTH: 219
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: humanized antibody with mouse CDRs and human
framework and constant regions
<220> FEATURE:
<221> NAME/KEY: DOMAIN
<222> LOCATION: (1)..(112)
<223> OTHER INFORMATION: variable domain
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (24)..(39)
<223> OTHER INFORMATION: CDRL1
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (55)..(61)
<223> OTHER INFORMATION: CDRL2
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (94)..(102)
<223> OTHER INFORMATION: CDRL3

<400> SEQUENCE: 43

Asp Ile Val Met Thr Gln Ser Pro Leu Ser Leu Pro Val Thr Pro Gly
1 5 10 15

Glu Pro Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Leu Val His Arg
20 25 30

Asn Gly Asn Thr Tyr Leu His Trp Tyr Leu Gln Lys Pro Gly Gln Ser
35 40 45

Pro Gln Leu Leu Ile Tyr Arg Val Ser Asn Arg Phe Ser Gly Val Pro
50 55 60

Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile
65 70 75 80

Ser Arg Val Glu Ala Glu Asp Val Gly Val Tyr Tyr Cys Ser Gln Ser
85 90 95

Thr His Phe Pro Tyr Thr Phe Gly Gly Gly Thr Lys Val Glu Ile Lys
100 105 110

Arg Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu
115 120 125

Gln Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe
130 135 140

Tyr Pro Arg Glu Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln
145 150 155 160

Ser Gly Asn Ser Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser
165 170 175

Thr Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu
180 185 190

Lys His Lys Val Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser
195 200 205

Pro Val Thr Lys Ser Phe Asn Arg Gly Glu Cys
210 215

<400> SEQUENCE: 44

Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu

-continued

```

Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser
      260                      265                      270

His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu
      275                      280                      285

Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr
      290                      295                      300

Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn
      305                      310                      315                      320

Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro
      325                      330                      335

Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln
      340                      345                      350

Val Tyr Thr Leu Pro Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val
      355                      360                      365

Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val
      370                      375                      380

Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro
      385                      390                      395                      400

Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr
      405                      410                      415

Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val
      420                      425                      430

Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu
      435                      440                      445

Ser Pro Gly
      450

```

```

<210> SEQ ID NO 45
<211> LENGTH: 219
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: humanized antibody with mouse CDRs and human
      framework and constant regions
<220> FEATURE:
<221> NAME/KEY: DOMAIN
<222> LOCATION: (1)..(112)
<223> OTHER INFORMATION: variable domain
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (24)..(39)
<223> OTHER INFORMATION: CDRL1
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (55)..(61)
<223> OTHER INFORMATION: CDRL2
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (94)..(102)
<223> OTHER INFORMATION: CDRL3

<400> SEQUENCE: 45

```

```

Asp Ile Val Met Thr Gln Ser Pro Leu Ser Leu Pro Val Thr Pro Gly
1          5          10          15

Glu Pro Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Leu Val His Arg
      20          25          30

Asn Gly Asn Thr Tyr Leu His Trp Tyr Leu Gln Lys Pro Gly Gln Ser
      35          40          45

Pro Gln Leu Leu Ile Tyr Arg Val Ser Asn Arg Phe Ser Gly Val Pro

```

-continued

50	55	60
Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile		
65	70	75 80
Ser Arg Val Glu Ala Glu Asp Val Gly Val Tyr Tyr Cys Ser Gln Ser		
	85	90 95
Thr His Phe Pro Tyr Thr Phe Gly Gly Gly Thr Lys Val Glu Ile Lys		
	100	105 110
Arg Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu		
	115	120 125
Gln Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe		
	130	135 140
Tyr Pro Arg Glu Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln		
	145	150 155 160
Ser Gly Asn Ser Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser		
	165	170 175
Thr Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu		
	180	185 190
Lys His Lys Val Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser		
	195	200 205
Pro Val Thr Lys Ser Phe Asn Arg Gly Glu Cys		
	210	215

<210> SEQ ID NO 46
 <211> LENGTH: 451
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: humanized antibody with mouse CDRs and human framework and constant regions
 <220> FEATURE:
 <221> NAME/KEY: DOMAIN
 <222> LOCATION: (1)..(122)
 <223> OTHER INFORMATION: variable domain
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (31)..(35)
 <223> OTHER INFORMATION: CDRH1
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (50)..(66)
 <223> OTHER INFORMATION: CDRH2
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (99)..(111)
 <223> OTHER INFORMATION: CDRH3

<400> SEQUENCE: 46

Gln Val Gln Phe Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ser		
1	5	10 15
Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Ser Tyr		
	20	25 30
Trp Met Gln Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met		
	35	40 45
Gly Thr Ile Tyr Pro Gly Asp Gly Asp Ser Arg Tyr Asn Gln Lys Phe		
	50	55 60
Lys Gly Lys Ala Leu Leu Thr Ala Asp Lys Ser Thr Ser Thr Ala Tyr		
	65	70 75 80
Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Tyr Cys		
	85	90 95

-continued

Ala Arg Phe Ser Leu Tyr Glu Gly Tyr Pro Tyr Tyr Phe Asp Tyr Trp
 100 105 110
 Gly Gln Gly Thr Thr Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro
 115 120 125
 Ser Val Phe Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr
 130 135 140
 Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr
 145 150 155 160
 Val Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro
 165 170 175
 Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr
 180 185 190
 Val Pro Ser Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn
 195 200 205
 His Lys Pro Ser Asn Thr Lys Val Asp Lys Arg Val Glu Pro Lys Ser
 210 215 220
 Cys Asp Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu
 225 230 235 240
 Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu
 245 250 255
 Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser
 260 265 270
 His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu
 275 280 285
 Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr
 290 295 300
 Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn
 305 310 315 320
 Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro
 325 330 335
 Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln
 340 345 350
 Val Tyr Thr Leu Pro Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val
 355 360 365
 Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val
 370 375 380
 Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro
 385 390 395 400
 Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr
 405 410 415
 Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val
 420 425 430
 Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu
 435 440 445
 Ser Pro Gly
 450

<210> SEQ ID NO 47

<211> LENGTH: 219

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

-continued

<223> OTHER INFORMATION: humanized antibody with mouse CDRs and human framework and constant regions
<220> FEATURE:
<221> NAME/KEY: DOMAIN
<222> LOCATION: (1)..(112)
<223> OTHER INFORMATION: variable domain
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (24)..(39)
<223> OTHER INFORMATION: CDRL1
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (55)..(61)
<223> OTHER INFORMATION: CDRL2
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (94)..(102)
<223> OTHER INFORMATION: CDRL3

<400> SEQUENCE: 47

Asp Ile Val Met Thr Gln Ser Pro Leu Ser Leu Pro Val Thr Pro Gly
1 5 10 15

Glu Pro Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Leu Val His Arg
20 25 30

Asn Ala Asn Thr Tyr Leu His Trp Tyr Leu Gln Lys Pro Gly Gln Ser
35 40 45

Pro Gln Leu Leu Ile Tyr Arg Val Ser Asn Arg Phe Ser Gly Val Pro
50 55 60

Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile
65 70 75 80

Ser Arg Val Glu Ala Glu Asp Val Gly Val Tyr Tyr Cys Ser Gln Ser
85 90 95

Thr His Phe Pro Tyr Thr Phe Gly Gly Gly Thr Lys Val Glu Ile Lys
100 105 110

Arg Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu
115 120 125

Gln Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe
130 135 140

Tyr Pro Arg Glu Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln
145 150 155 160

Ser Gly Asn Ser Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser
165 170 175

Thr Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu
180 185 190

Lys His Lys Val Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser
195 200 205

Pro Val Thr Lys Ser Phe Asn Arg Gly Glu Cys
210 215

<210> SEQ ID NO 48
<211> LENGTH: 451
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: humanized antibody with mouse CDRs and human framework and constant regions
<220> FEATURE:
<221> NAME/KEY: DOMAIN
<222> LOCATION: (1)..(122)
<223> OTHER INFORMATION: variable domain
<220> FEATURE:

-continued

```

<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (31)..(35)
<223> OTHER INFORMATION: CDRH1
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (50)..(66)
<223> OTHER INFORMATION: CDRH2
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (99)..(111)
<223> OTHER INFORMATION: CDRH3

<400> SEQUENCE: 48

Gln Val Gln Phe Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ser
1          5          10          15

Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Ser Tyr
20          25          30

Trp Met Gln Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met
35          40          45

Gly Thr Ile Tyr Pro Gly Asp Ser Asp Ser Arg Tyr Asn Gln Lys Phe
50          55          60

Lys Gly Lys Ala Leu Leu Thr Ala Asp Lys Ser Thr Ser Thr Ala Tyr
65          70          75          80

Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Tyr Cys
85          90          95

Ala Arg Phe Ser Leu Tyr Glu Gly Tyr Pro Tyr Tyr Phe Asp Tyr Trp
100         105         110

Gly Gln Gly Thr Thr Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro
115         120         125

Ser Val Phe Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr
130         135         140

Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr
145         150         155         160

Val Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro
165         170         175

Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr
180         185         190

Val Pro Ser Ser Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn
195         200         205

His Lys Pro Ser Asn Thr Lys Val Asp Lys Arg Val Glu Pro Lys Ser
210         215         220

Cys Asp Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu
225         230         235         240

Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu
245         250         255

Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser
260         265         270

His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu
275         280         285

Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr
290         295         300

Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn
305         310         315         320

Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro
325         330         335

```

-continued

```

Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln
    340                      345                      350

Val Tyr Thr Leu Pro Pro Ser Arg Glu Glu Met Thr Lys Asn Gln Val
    355                      360                      365

Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val
    370                      375                      380

Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro
    385                      390                      395                      400

Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr
    405                      410                      415

Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val
    420                      425                      430

Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu
    435                      440                      445

Ser Pro Gly
    450

```

```

<210> SEQ ID NO 49
<211> LENGTH: 219
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: humanized antibody with mouse CDRs and human
    framework and constant regions
<220> FEATURE:
<221> NAME/KEY: DOMAIN
<222> LOCATION: (1)..(112)
<223> OTHER INFORMATION: variable domain
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (24)..(39)
<223> OTHER INFORMATION: CDRL1
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (55)..(61)
<223> OTHER INFORMATION: CDRL2
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (94)..(102)
<223> OTHER INFORMATION: CDRL3

<400> SEQUENCE: 49

```

```

Asp Ile Val Met Thr Gln Ser Pro Leu Ser Leu Pro Val Thr Pro Gly
1          5          10          15

Glu Pro Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Leu Val His Arg
20         25         30

Asn Gly Asn Thr Tyr Leu His Trp Tyr Leu Gln Lys Pro Gly Gln Ser
35         40         45

Pro Gln Leu Leu Ile Tyr Arg Val Ser Asn Arg Phe Ser Gly Val Pro
50         55         60

Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile
65         70         75         80

Ser Arg Val Glu Ala Glu Asp Val Gly Val Tyr Tyr Cys Ser Gln Ser
85         90         95

Thr His Phe Pro Tyr Thr Phe Gly Gly Gly Thr Lys Val Glu Ile Lys
100        105        110

Arg Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu
115        120        125

```

-continued

Gln Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe
 130 135 140

Tyr Pro Arg Glu Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln
 145 150 155 160

Ser Gly Asn Ser Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser
 165 170 175

Thr Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu
 180 185 190

Lys His Lys Val Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser
 195 200 205

Pro Val Thr Lys Ser Phe Asn Arg Gly Glu Cys
 210 215

<210> SEQ ID NO 50
 <211> LENGTH: 451
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: humanized antibody with mouse CDRs and human
 framework and constant regions
 <220> FEATURE:
 <221> NAME/KEY: DOMAIN
 <222> LOCATION: (1)..(122)
 <223> OTHER INFORMATION: variable domain
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (31)..(35)
 <223> OTHER INFORMATION: CDRH1
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (50)..(66)
 <223> OTHER INFORMATION: CDRH2
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (99)..(111)
 <223> OTHER INFORMATION: CDRH3

<400> SEQUENCE: 50

Gln Val Gln Phe Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ser
 1 5 10 15

Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Ser Tyr
 20 25 30

Trp Met Gln Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met
 35 40 45

Gly Thr Ile Tyr Pro Gly Asp Ser Asp Ser Arg Tyr Asn Gln Lys Phe
 50 55 60

Lys Gly Lys Ala Leu Leu Thr Ala Asp Lys Ser Thr Ser Thr Ala Tyr
 65 70 75 80

Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95

Ala Arg Phe Ser Leu Tyr Glu Gly Tyr Pro Tyr Tyr Phe Asp Tyr Trp
 100 105 110

Gly Gln Gly Thr Thr Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro
 115 120 125

Ser Val Phe Pro Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr
 130 135 140

Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr
 145 150 155 160

Val Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro

-continued

165					170					175					
Ala	Val	Leu	Gln	Ser	Ser	Gly	Leu	Tyr	Ser	Leu	Ser	Ser	Val	Val	Thr
		180						185					190		
Val	Pro	Ser	Ser	Ser	Leu	Gly	Thr	Gln	Thr	Tyr	Ile	Cys	Asn	Val	Asn
		195					200					205			
His	Lys	Pro	Ser	Asn	Thr	Lys	Val	Asp	Lys	Arg	Val	Glu	Pro	Lys	Ser
	210					215					220				
Cys	Asp	Lys	Thr	His	Thr	Cys	Pro	Pro	Cys	Pro	Ala	Pro	Glu	Leu	Leu
	225					230					235				240
Gly	Gly	Pro	Ser	Val	Phe	Leu	Phe	Pro	Pro	Lys	Pro	Lys	Asp	Thr	Leu
			245					250						255	
Met	Ile	Ser	Arg	Thr	Pro	Glu	Val	Thr	Cys	Val	Val	Val	Asp	Val	Ser
			260					265					270		
His	Glu	Asp	Pro	Glu	Val	Lys	Phe	Asn	Trp	Tyr	Val	Asp	Gly	Val	Glu
		275					280					285			
Val	His	Asn	Ala	Lys	Thr	Lys	Pro	Arg	Glu	Glu	Gln	Tyr	Asn	Ser	Thr
	290					295					300				
Tyr	Arg	Val	Val	Ser	Val	Leu	Thr	Val	Leu	His	Gln	Asp	Trp	Leu	Asn
	305					310					315				320
Gly	Lys	Glu	Tyr	Lys	Cys	Lys	Val	Ser	Asn	Lys	Ala	Leu	Pro	Ala	Pro
			325					330					335		
Ile	Glu	Lys	Thr	Ile	Ser	Lys	Ala	Lys	Gly	Gln	Pro	Arg	Glu	Pro	Gln
		340						345					350		
Val	Tyr	Thr	Leu	Pro	Pro	Ser	Arg	Glu	Glu	Met	Thr	Lys	Asn	Gln	Val
		355					360					365			
Ser	Leu	Thr	Cys	Leu	Val	Lys	Gly	Phe	Tyr	Pro	Ser	Asp	Ile	Ala	Val
	370					375					380				
Glu	Trp	Glu	Ser	Asn	Gly	Gln	Pro	Glu	Asn	Asn	Tyr	Lys	Thr	Thr	Pro
	385					390					395				400
Pro	Val	Leu	Asp	Ser	Asp	Gly	Ser	Phe	Phe	Leu	Tyr	Ser	Lys	Leu	Thr
			405					410						415	
Val	Asp	Lys	Ser	Arg	Trp	Gln	Gln	Gly	Asn	Val	Phe	Ser	Cys	Ser	Val
		420						425					430		
Met	His	Glu	Ala	Leu	His	Asn	His	Tyr	Thr	Gln	Lys	Ser	Leu	Ser	Leu
		435					440					445			
Ser	Pro	Gly													
		450													

<210> SEQ ID NO 51

<211> LENGTH: 219

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: humanized antibody with mouse CDRs and human framework and constant regions

<220> FEATURE:

<221> NAME/KEY: DOMAIN

<222> LOCATION: (1)..(112)

<223> OTHER INFORMATION: variable domain

<220> FEATURE:

<221> NAME/KEY: MISC_FEATURE

<222> LOCATION: (24)..(39)

<223> OTHER INFORMATION: CDRL1

<220> FEATURE:

<221> NAME/KEY: MISC_FEATURE

<222> LOCATION: (55)..(61)

<223> OTHER INFORMATION: CDRL2

-continued

<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (94)..(102)
<223> OTHER INFORMATION: CDRL3

<400> SEQUENCE: 51

Asp Ile Val Met Thr Gln Ser Pro Leu Ser Leu Pro Val Thr Pro Gly
1 5 10 15
Glu Pro Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Leu Val His Arg
20 25 30
Asn Ala Asn Thr Tyr Leu His Trp Tyr Leu Gln Lys Pro Gly Gln Ser
35 40 45
Pro Gln Leu Leu Ile Tyr Arg Val Ser Asn Arg Phe Ser Gly Val Pro
50 55 60
Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile
65 70 75 80
Ser Arg Val Glu Ala Glu Asp Val Gly Val Tyr Tyr Cys Ser Gln Ser
85 90 95
Thr His Phe Pro Tyr Thr Phe Gly Gly Gly Thr Lys Val Glu Ile Lys
100 105 110
Arg Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu
115 120 125
Gln Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe
130 135 140
Tyr Pro Arg Glu Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln
145 150 155 160
Ser Gly Asn Ser Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser
165 170 175
Thr Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu
180 185 190
Lys His Lys Val Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser
195 200 205
Pro Val Thr Lys Ser Phe Asn Arg Gly Glu Cys
210 215

<210> SEQ ID NO 52
<211> LENGTH: 442
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: chimeric antibody with mouse variable domain
and human constant domain
<220> FEATURE:
<221> NAME/KEY: DOMAIN
<222> LOCATION: (1)..(113)
<223> OTHER INFORMATION: variable domain
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (31)..(35)
<223> OTHER INFORMATION: CDRH1
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (50)..(66)
<223> OTHER INFORMATION: CDRH2
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (99)..(102)
<223> OTHER INFORMATION: CDRH3

<400> SEQUENCE: 52

-continued

Glu	Val	Leu	Leu	Gln	Gln	Ser	Gly	Pro	Glu	Leu	Val	Lys	Pro	Gly	Ala	1	5	10	15
Ser	Val	Lys	Ile	Pro	Cys	Lys	Ala	Ser	Gly	Tyr	Lys	Phe	Thr	Asp	Tyr	20	25	30	
Asn	Met	Asp	Trp	Val	Lys	Gln	Ser	His	Gly	Lys	Ser	Leu	Glu	Trp	Ile	35	40	45	
Gly	Asp	Ile	Asn	Pro	Lys	Asn	Gly	Gly	Thr	Ile	Tyr	Asn	Leu	Lys	Phe	50	55	60	
Lys	Gly	Lys	Ala	Thr	Leu	Thr	Val	Asp	Met	Ser	Ser	Ser	Thr	Ala	Tyr	65	70	75	80
Met	Glu	Leu	Arg	Ser	Leu	Thr	Ser	Glu	Asp	Thr	Ala	Val	Tyr	Tyr	Cys	85	90	95	
Val	Arg	Arg	Met	Asp	Tyr	Trp	Gly	Gln	Gly	Thr	Ser	Val	Thr	Val	Ser	100	105	110	
Ser	Ala	Ser	Thr	Lys	Gly	Pro	Ser	Val	Phe	Pro	Leu	Ala	Pro	Ser	Ser	115	120	125	
Lys	Ser	Thr	Ser	Gly	Gly	Thr	Ala	Ala	Leu	Gly	Cys	Leu	Val	Lys	Asp	130	135	140	
Tyr	Phe	Pro	Glu	Pro	Val	Thr	Val	Ser	Trp	Asn	Ser	Gly	Ala	Leu	Thr	145	150	155	160
Ser	Gly	Val	His	Thr	Phe	Pro	Ala	Val	Leu	Gln	Ser	Ser	Gly	Leu	Tyr	165	170	175	
Ser	Leu	Ser	Ser	Val	Val	Thr	Val	Pro	Ser	Ser	Ser	Leu	Gly	Thr	Gln	180	185	190	
Thr	Tyr	Ile	Cys	Asn	Val	Asn	His	Lys	Pro	Ser	Asn	Thr	Lys	Val	Asp	195	200	205	
Lys	Arg	Val	Glu	Pro	Lys	Ser	Cys	Asp	Lys	Thr	His	Thr	Cys	Pro	Pro	210	215	220	
Cys	Pro	Ala	Pro	Glu	Leu	Leu	Gly	Gly	Pro	Ser	Val	Phe	Leu	Phe	Pro	225	230	235	240
Pro	Lys	Pro	Lys	Asp	Thr	Leu	Met	Ile	Ser	Arg	Thr	Pro	Glu	Val	Thr	245	250	255	
Cys	Val	Val	Val	Asp	Val	Ser	His	Glu	Asp	Pro	Glu	Val	Lys	Phe	Asn	260	265	270	
Trp	Tyr	Val	Asp	Gly	Val	Glu	Val	His	Asn	Ala	Lys	Thr	Lys	Pro	Arg	275	280	285	
Glu	Glu	Gln	Tyr	Asn	Ser	Thr	Tyr	Arg	Val	Val	Ser	Val	Leu	Thr	Val	290	295	300	
Leu	His	Gln	Asp	Trp	Leu	Asn	Gly	Lys	Glu	Tyr	Lys	Cys	Lys	Val	Ser	305	310	315	320
Asn	Lys	Ala	Leu	Pro	Ala	Pro	Ile	Glu	Lys	Thr	Ile	Ser	Lys	Ala	Lys	325	330	335	
Gly	Gln	Pro	Arg	Glu	Pro	Gln	Val	Tyr	Thr	Leu	Pro	Pro	Ser	Arg	Glu	340	345	350	
Glu	Met	Thr	Lys	Asn	Gln	Val	Ser	Leu	Thr	Cys	Leu	Val	Lys	Gly	Phe	355	360	365	
Tyr	Pro	Ser	Asp	Ile	Ala	Val	Glu	Trp	Glu	Ser	Asn	Gly	Gln	Pro	Glu	370	375	380	
Asn	Asn	Tyr	Lys	Thr	Thr	Pro	Pro	Val	Leu	Asp	Ser	Asp	Gly	Ser	Phe	385	390	395	400
Phe	Leu	Tyr	Ser	Lys	Leu	Thr	Val	Asp	Lys	Ser	Arg	Trp	Gln	Gln	Gly				

-continued

405	410	415
Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr		
420	425	430
Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly		
435	440	

<210> SEQ ID NO 53
 <211> LENGTH: 214
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: chimeric antibody with mouse variable domain
 and human constant domain
 <220> FEATURE:
 <221> NAME/KEY: DOMAIN
 <222> LOCATION: (1)..(107)
 <223> OTHER INFORMATION: variable domain
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (24)..(34)
 <223> OTHER INFORMATION: CDRL1
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (50)..(56)
 <223> OTHER INFORMATION: CDRL2
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (89)..(97)
 <223> OTHER INFORMATION: CDRL3
 <400> SEQUENCE: 53

Asp Ile Gln Met Thr Gln Thr Thr Ser Ser Leu Ser Ala Ser Leu Gly		
1	5	10 15
Asp Arg Val Thr Ile Ser Cys Arg Ala Ser Gln Asp Ile Ser Asn Tyr		
20	25	30
Leu Asn Trp Tyr Gln Gln Lys Pro Asp Gly Thr Val Lys Leu Leu Ile		
35	40	45
Tyr Tyr Ile Ser Arg Leu His Ser Gly Val Pro Ser Arg Phe Arg Gly		
50	55	60
Ser Gly Ser Gly Thr Asp Phe Ser Leu Thr Ile Ser Asn Leu Glu Gln		
65	70	75 80
Glu Asp Ile Ala Thr Tyr Phe Cys Gln Gln Gly Asn Thr Leu Pro Tyr		
85	90	95
Thr Phe Gly Gly Gly Thr Asn Leu Glu Ile Lys Arg Thr Val Ala Ala		
100	105	110
Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly		
115	120	125
Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala		
130	135	140
Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln		
145	150	155 160
Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser		
165	170	175
Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val Tyr		
180	185	190
Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys Ser		
195	200	205
Phe Asn Arg Gly Glu Cys		
210		

-continued

<210> SEQ ID NO 54
<211> LENGTH: 442
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: humanized antibody with mouse CDRs and human framework and constant regions
<220> FEATURE:
<221> NAME/KEY: DOMAIN
<222> LOCATION: (1)..(113)
<223> OTHER INFORMATION: variable domain
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (31)..(35)
<223> OTHER INFORMATION: CDRH1
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (50)..(66)
<223> OTHER INFORMATION: CDRH2
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (99)..(102)
<223> OTHER INFORMATION: CDRH3

<400> SEQUENCE: 54

Gln	Val	Gln	Leu	Val	Gln	Ser	Gly	Ala	Glu	Val	Lys	Lys	Pro	Gly	Ala	
1			5						10					15		
Ser	Val	Lys	Val	Ser	Cys	Lys	Ala	Ser	Gly	Tyr	Thr	Phe	Thr	Asp	Tyr	
		20						25					30			
Asn	Met	Asp	Trp	Val	Arg	Gln	Ala	Pro	Gly	Gln	Gly	Leu	Glu	Trp	Met	
		35					40					45				
Gly	Asp	Ile	Asn	Pro	Lys	Asn	Gly	Gly	Thr	Ile	Tyr	Asn	Leu	Lys	Phe	
	50					55				60						
Lys	Gly	Arg	Val	Thr	Leu	Thr	Thr	Asp	Thr	Ser	Thr	Ser	Thr	Ala	Tyr	
	65				70				75					80		
Met	Glu	Leu	Arg	Ser	Leu	Arg	Ser	Asp	Asp	Thr	Ala	Val	Tyr	Tyr	Cys	
			85					90					95			
Val	Arg	Arg	Met	Asp	Tyr	Trp	Gly	Gln	Gly	Thr	Thr	Val	Thr	Val	Ser	
			100					105					110			
Ser	Ala	Ser	Thr	Lys	Gly	Pro	Ser	Val	Phe	Pro	Leu	Ala	Pro	Ser	Ser	
		115					120					125				
Lys	Ser	Thr	Ser	Gly	Gly	Thr	Ala	Ala	Leu	Gly	Cys	Leu	Val	Lys	Asp	
	130					135					140					
Tyr	Phe	Pro	Glu	Pro	Val	Thr	Val	Ser	Trp	Asn	Ser	Gly	Ala	Leu	Thr	
	145				150				155					160		
Ser	Gly	Val	His	Thr	Phe	Pro	Ala	Val	Leu	Gln	Ser	Ser	Gly	Leu	Tyr	
			165						170					175		
Ser	Leu	Ser	Ser	Val	Val	Thr	Val	Pro	Ser	Ser	Ser	Leu	Gly	Thr	Gln	
			180					185					190			
Thr	Tyr	Ile	Cys	Asn	Val	Asn	His	Lys	Pro	Ser	Asn	Thr	Lys	Val	Asp	
		195					200					205				
Lys	Arg	Val	Glu	Pro	Lys	Ser	Cys	Asp	Lys	Thr	His	Thr	Cys	Pro	Pro	
	210					215					220					
Cys	Pro	Ala	Pro	Glu	Leu	Leu	Gly	Gly	Pro	Ser	Val	Phe	Leu	Phe	Pro	
	225				230				235					240		
Pro	Lys	Pro	Lys	Asp	Thr	Leu	Met	Ile	Ser	Arg	Thr	Pro	Glu	Val	Thr	
			245					250						255		

```
<210> SEQ ID NO 55
<211> LENGTH: 214
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: humanized antibody with mouse CDRs and human
framework and constant regions
<220> FEATURE:
<221> NAME/KEY: DOMAIN
<222> LOCATION: (1)..(107)
<223> OTHER INFORMATION: variable domain
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (24)..(34)
<223> OTHER INFORMATION: CDRL1
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (50)..(56)
<223> OTHER INFORMATION: CDRL2
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (89)..(97)
<223> OTHER INFORMATION: CDRL3

<400> SEQUENCE: 55
```

Asp	Ile	Gln	Met	Thr	Gln	Ser	Pro	Ser	Ser	Leu	Ser	Ala	Ser	Val	Gly
1				5					10					15	
Asp	Arg	Val	Thr	Ile	Thr	Cys	Arg	Ala	Ser	Gln	Asp	Ile	Ser	Asn	Tyr
			20					25					30		
Leu	Asn	Trp	Tyr	Gln	Gln	Lys	Pro	Gly	Lys	Val	Pro	Lys	Leu	Leu	Ile
		35					40					45			
Tyr	Tyr	Ile	Ser	Arg	Leu	His	Ser	Gly	Val	Pro	Ser	Arg	Phe	Ser	Gly
	50					55					60				
Ser	Gly	Ser	Gly	Thr	Asp	Phe	Thr	Leu	Thr	Ile	Ser	Ser	Leu	Gln	Pro

-continued

65	70	75	80
Glu Asp Val Ala Thr Tyr Tyr Cys Gln Gln Gly Asn Thr Leu Pro Tyr	85	90	95
Thr Phe Gly Gly Gly Thr Lys Val Glu Ile Lys Arg Thr Val Ala Ala	100	105	110
Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly	115	120	125
Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala	130	135	140
Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln	145	150	155
Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser	165	170	175
Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val Tyr	180	185	190
Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys Ser	195	200	205
Phe Asn Arg Gly Glu Cys	210		

<210> SEQ ID NO 56
 <211> LENGTH: 442
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: humanized antibody with mouse CDRs and human framework and constant regions
 <220> FEATURE:
 <221> NAME/KEY: DOMAIN
 <222> LOCATION: (1)..(113)
 <223> OTHER INFORMATION: variable domain
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (31)..(35)
 <223> OTHER INFORMATION: CDRH1
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (50)..(66)
 <223> OTHER INFORMATION: CDRH2
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (99)..(102)
 <223> OTHER INFORMATION: CDRH3

<400> SEQUENCE: 56

Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala	1	5	10	15
Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe Thr Asp Tyr	20	25	30	
Asn Met Asp Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met	35	40	45	
Gly Asp Ile Asn Pro Lys Asn Ala Gly Thr Ile Tyr Asn Leu Lys Phe	50	55	60	
Lys Gly Arg Val Thr Leu Thr Val Asp Thr Ser Thr Ser Thr Ala Tyr	65	70	75	80
Met Glu Leu Arg Ser Leu Arg Ser Asp Asp Thr Ala Val Tyr Tyr Cys	85	90	95	
Val Arg Arg Met Asp Tyr Trp Gly Gln Gly Thr Thr Val Thr Val Ser	100	105	110	

-continued

```

Ser Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Ser Ser
   115                               120                               125

Lys Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly Cys Leu Val Lys Asp
   130                               135                               140

Tyr Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr
   145                               150                               155                               160

Ser Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr
                               165                               170                               175

Ser Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Gln
                               180                               185                               190

Thr Tyr Ile Cys Asn Val Asn His Lys Pro Ser Asn Thr Lys Val Asp
   195                               200                               205

Lys Arg Val Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro
   210                               215                               220

Cys Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro
   225                               230                               235                               240

Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr
                               245                               250                               255

Cys Val Val Val Asp Val Ser His Glu Asp Pro Glu Val Lys Phe Asn
   260                               265                               270

Trp Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg
   275                               280                               285

Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val
   290                               295                               300

Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser
   305                               310                               315                               320

Asn Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys
   325                               330                               335

Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Glu
   340                               345                               350

Glu Met Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe
   355                               360                               365

Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu
   370                               375                               380

Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe
   385                               390                               395                               400

Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly
   405                               410                               415

Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr
   420                               425                               430

Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly
   435                               440

```

<210> SEQ ID NO 57

<211> LENGTH: 214

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: humanized antibody with mouse CDRs and human framework and constant regions

<220> FEATURE:

<221> NAME/KEY: DOMAIN

<222> LOCATION: (1)..(107)

<223> OTHER INFORMATION: variable domain

-continued

```
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (24)..(34)
<223> OTHER INFORMATION: CDRL1
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (50)..(56)
<223> OTHER INFORMATION: CDRL2
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (89)..(97)
<223> OTHER INFORMATION: CDRL3

<400> SEQUENCE: 57

Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
1             5             10             15

Asp Arg Val Thr Ile Thr Cys Arg Ala Ser Gln Asp Ile Ser Asn Tyr
                20             25             30

Leu Asn Trp Tyr Gln Gln Lys Pro Gly Lys Val Pro Lys Leu Leu Ile
            35             40             45

Tyr Tyr Ile Ser Arg Leu His Ser Gly Val Pro Ser Arg Phe Ser Gly
50             55             60

Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro
65             70             75             80

Glu Asp Val Ala Thr Tyr Tyr Cys Gln Gln Gly Asn Thr Leu Pro Tyr
            85             90             95

Thr Phe Gly Gly Gly Thr Lys Val Glu Ile Lys Arg Thr Val Ala Ala
100            105            110

Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys Ser Gly
115            120            125

Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg Glu Ala
130            135            140

Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn Ser Gln
145            150            155            160

Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser Leu Ser
165            170            175

Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys Val Tyr
180            185            190

Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr Lys Ser
195            200            205

Phe Asn Arg Gly Glu Cys
210

<210> SEQ ID NO 58
<211> LENGTH: 441
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: chimeric antibody with mouse variable domain
and human constant domain
<220> FEATURE:
<221> NAME/KEY: DOMAIN
<222> LOCATION: (1)..(112)
<223> OTHER INFORMATION: variable domain
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (31)..(35)
<223> OTHER INFORMATION: CDRH1
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (50)..(66)
```

-continued

<223> OTHER INFORMATION: CDRH2

<220> FEATURE:

<221> NAME/KEY: MISC_FEATURE

<222> LOCATION: (99)..(101)

<223> OTHER INFORMATION: CDRH3

<400> SEQUENCE: 58

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Lys Pro Gly Gly
1 5 10 15
Ser Leu Lys Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
20 25 30
Ala Met Ser Trp Val Arg Gln Ser Pro Glu Lys Arg Leu Glu Trp Val
35 40 45
Ala Glu Ile Ser Gly Gly Gly Ser Tyr Thr Tyr Tyr Pro Asp Thr Val
50 55 60
Thr Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Thr Leu Tyr
65 70 75 80
Leu Glu Met Ser Ser Leu Arg Ser Glu Asp Thr Ala Met Tyr Tyr Cys
85 90 95
Ala Ser Arg Ala Tyr Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ala
100 105 110
Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Ser Ser Lys
115 120 125
Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr
130 135 140
Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser
145 150 155 160
Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser
165 170 175
Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Gln Thr
180 185 190
Tyr Ile Cys Asn Val Asn His Lys Pro Ser Asn Thr Lys Val Asp Lys
195 200 205
Arg Val Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys
210 215 220
Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro
225 230 235 240
Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys
245 250 255
Val Val Val Asp Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp
260 265 270
Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu
275 280 285
Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu
290 295 300
His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn
305 310 315 320
Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly
325 330 335
Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Glu Glu
340 345 350
Met Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr
355 360 365

-continued

```

Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn
 370                375                380

Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe
385                390                395                400

Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn
                405                410                415

Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr
                420                425                430

Gln Lys Ser Leu Ser Leu Ser Pro Gly
                435                440

```

```

<210> SEQ ID NO 59
<211> LENGTH: 219
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: chimeric antibody with mouse variable domain
and human constant domain
<220> FEATURE:
<221> NAME/KEY: DOMAIN
<222> LOCATION: (1)..(112)
<223> OTHER INFORMATION: variable domain
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (24)..(39)
<223> OTHER INFORMATION: CDRL1
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (55)..(61)
<223> OTHER INFORMATION: CDRL2
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (94)..(102)
<223> OTHER INFORMATION: CDRL3

```

```

<400> SEQUENCE: 59

```

```

Asp Val Leu Met Thr Gln Thr Pro Leu Ser Leu Pro Val Ser Leu Gly
 1                5                10                15

Asp Gln Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Ile Val His Ser
                20                25                30

Asn Gly Asn Thr Tyr Phe Glu Trp Tyr Leu Gln Lys Pro Gly Gln Ser
                35                40                45

Pro Lys Leu Leu Ile Tyr Lys Val Ser Asn Arg Phe Ser Gly Val Pro
                50                55                60

Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile
65                70                75                80

Ser Arg Val Glu Ala Glu Asp Leu Gly Val Tyr Tyr Cys Phe Gln Gly
                85                90                95

Ser His Val Pro Tyr Thr Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys
                100                105                110

Arg Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu
                115                120                125

Gln Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe
                130                135                140

Tyr Pro Arg Glu Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln
145                150                155                160

Ser Gly Asn Ser Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser
                165                170                175

```

-continued

Thr Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu
 180 185 190

Lys His Lys Val Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser
 195 200 205

Pro Val Thr Lys Ser Phe Asn Arg Gly Glu Cys
 210 215

<210> SEQ ID NO 60
 <211> LENGTH: 441
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: humanized antibody with mouse CDRs and human
 framework and constant regions
 <220> FEATURE:
 <221> NAME/KEY: DOMAIN
 <222> LOCATION: (1)..(112)
 <223> OTHER INFORMATION: variable domain
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (31)..(35)
 <223> OTHER INFORMATION: CDRH1
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (50)..(66)
 <223> OTHER INFORMATION: CDRH2
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (99)..(101)
 <223> OTHER INFORMATION: CDRH3

<400> SEQUENCE: 60

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Lys Pro Gly Gly
 1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
 20 25 30

Ala Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45

Ser Glu Ile Ser Gly Gly Gly Ser Tyr Thr Tyr Tyr Pro Asp Thr Val
 50 55 60

Thr Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Ser Leu Tyr
 65 70 75 80

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95

Ala Ser Arg Ala Tyr Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser
 100 105 110

Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Ser Ser Lys
 115 120 125

Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr
 130 135 140

Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser
 145 150 155 160

Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser
 165 170 175

Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Gln Thr
 180 185 190

Tyr Ile Cys Asn Val Asn His Lys Pro Ser Asn Thr Lys Val Asp Lys
 195 200 205

Arg Val Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys

-continued

210	215	220
Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro		
225	230	235 240
Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys		
	245	250 255
Val Val Val Asp Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp		
	260	265 270
Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu		
	275	280 285
Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu		
	290	295 300
His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn		
305	310	315 320
Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly		
	325	330 335
Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Glu Glu		
	340	345 350
Met Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr		
	355	360 365
Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn		
	370	375 380
Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe		
385	390	395 400
Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn		
	405	410 415
Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr		
	420	425 430
Gln Lys Ser Leu Ser Leu Ser Pro Gly		
	435	440

<210> SEQ ID NO 61

<211> LENGTH: 441

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: humanized antibody with mouse CDRs and human framework and constant regions

<220> FEATURE:

<221> NAME/KEY: DOMAIN

<222> LOCATION: (1)..(112)

<223> OTHER INFORMATION: variable domain

<220> FEATURE:

<221> NAME/KEY: MISC_FEATURE

<222> LOCATION: (31)..(35)

<223> OTHER INFORMATION: CDRH1

<220> FEATURE:

<221> NAME/KEY: MISC_FEATURE

<222> LOCATION: (50)..(66)

<223> OTHER INFORMATION: CDRH2

<220> FEATURE:

<221> NAME/KEY: MISC_FEATURE

<222> LOCATION: (99)..(101)

<223> OTHER INFORMATION: CDRH3

<400> SEQUENCE: 61

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Lys Pro Gly Gly
1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
20 25 30

Ala	Met	Ser	Trp	Val	Arg	Gln	Ala	Pro	Gly	Lys	Gly	Leu	Glu	Trp	Val
35						40						45			
Ser	Glu	Ile	Ser	Gly	Gly	Ser	Tyr	Thr	Tyr	Tyr	Pro	Asp	Thr	Val	
50				55						60					
Thr	Gly	Arg	Phe	Thr	Ile	Ser	Arg	Asp	Asn	Ala	Lys	Asn	Ser	Leu	Tyr
65				70						75				80	
Leu	Gln	Met	Asn	Ser	Leu	Arg	Ala	Glu	Asp	Thr	Ala	Val	Tyr	Tyr	Cys
				85				90						95	
Ala	Ser	Arg	Ala	Tyr	Trp	Gly	Gln	Gly	Thr	Leu	Val	Thr	Val	Ser	Ser
		100						105				110			
Ala	Ser	Thr	Lys	Gly	Pro	Ser	Val	Phe	Pro	Leu	Ala	Pro	Ser	Ser	Lys
		115				120						125			
Ser	Thr	Ser	Gly	Gly	Thr	Ala	Ala	Leu	Gly	Cys	Leu	Val	Lys	Asp	Tyr
130						135				140					
Phe	Pro	Glu	Pro	Val	Thr	Val	Ser	Trp	Asn	Ser	Gly	Ala	Leu	Thr	Ser
145				150						155				160	
Gly	Val	His	Thr	Phe	Pro	Ala	Val	Leu	Gln	Ser	Ser	Gly	Leu	Tyr	Ser
				165				170						175	
Leu	Ser	Ser	Val	Val	Thr	Val	Pro	Ser	Ser	Ser	Leu	Gly	Thr	Gln	Thr
		180						185				190			
Tyr	Ile	Cys	Asn	Val	Asn	His	Lys	Pro	Ser	Asn	Thr	Lys	Val	Asp	Lys
195						200						205			
Arg	Val	Glu	Pro	Lys	Ser	Cys	Asp	Lys	Thr	His	Thr	Cys	Pro	Pro	Cys
210						215				220					
Pro	Ala	Pro	Glu	Leu	Leu	Gly	Asp	Asp	Ser	Val	Phe	Leu	Phe	Pro	Pro
225				230						235				240	
Lys	Pro	Lys	Asp	Thr	Leu	Met	Ile	Ser	Arg	Thr	Pro	Glu	Val	Thr	Cys
				245				250						255	
Val	Val	Val	Asp	Val	Ser	His	Glu	Asp	Gly	Glu	Val	Lys	Phe	Asn	Trp
		260						265				270			
Tyr	Val	Asp	Gly	Val	Glu	Val	His	Asn	Ala	Lys	Thr	Lys	Pro	Arg	Glu
275						280						285			
Glu	Gln	Tyr	Asn	Ser	Thr	Tyr	Arg	Val	Val	Ser	Val	Leu	Thr	Val	Leu
290						295				300					
His	Gln	Asp	Trp	Leu	Asn	Gly	Lys	Glu	Tyr	Lys	Cys	Lys	Val	Ser	Asn
305				310						315				320	
Lys	Ala	Leu	Pro	Arg	Pro	Ile	Glu	Lys	Thr	Ile	Ser	Lys	Ala	Lys	Gly
				325				330						335	
Gln	Pro	Arg	Glu	Pro	Gln	Val	Tyr	Thr	Leu	Pro	Pro	Ser	Arg	Glu	Glu
		340						345				350			
Met	Thr	Lys	Asn	Gln	Val	Ser	Leu	Thr	Cys	Leu	Val	Lys	Gly	Phe	Tyr
		355				360						365			
Pro	Ser	Asp	Ile	Ala	Val	Glu	Trp	Glu	Ser	Asn	Gly	Gln	Pro	Glu	Asn
370						375				380					
Asn	Tyr	Lys	Thr	Thr	Pro	Pro	Val	Leu	Asp	Ser	Asp	Gly	Ser	Phe	Phe
385				390						395				400	
Leu	Tyr	Ser	Lys	Leu	Thr	Val	Asp</								

-continued

Gln Lys Ser Leu Ser Leu Ser Pro Gly
435 440

<210> SEQ ID NO 62
 <211> LENGTH: 219
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: humanized antibody with mouse CDRs and human
 framework and constant regions
 <220> FEATURE:
 <221> NAME/KEY: DOMAIN
 <222> LOCATION: (1)..(112)
 <223> OTHER INFORMATION: variable domain
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (24)..(39)
 <223> OTHER INFORMATION: CDRL1
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (55)..(61)
 <223> OTHER INFORMATION: CDRL2
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (94)..(102)
 <223> OTHER INFORMATION: CDRL3

<400> SEQUENCE: 62

Asp Ile Val Met Thr Gln Thr Pro Leu Ser Leu Pro Val Thr Pro Gly
1 5 10 15
 Glu Pro Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Ile Val His Ser
20 25 30
 Asn Gly Asn Thr Tyr Phe Glu Trp Tyr Leu Gln Lys Pro Gly Gln Ser
35 40 45
 Pro Gln Leu Leu Ile Tyr Lys Val Ser Asn Arg Phe Ser Gly Val Pro
50 55 60
 Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile
65 70 75 80
 Ser Arg Val Glu Ala Glu Asp Val Gly Val Tyr Tyr Cys Phe Gln Gly
85 90 95
 Ser His Val Pro Tyr Thr Phe Gly Gly Gly Thr Lys Val Glu Ile Lys
100 105 110
 Arg Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu
115 120 125
 Gln Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe
130 135 140
 Tyr Pro Arg Glu Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln
145 150 155 160
 Ser Gly Asn Ser Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser
165 170 175
 Thr Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu
180 185 190
 Lys His Lys Val Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser
195 200 205
 Pro Val Thr Lys Ser Phe Asn Arg Gly Glu Cys
210 215

<210> SEQ ID NO 63
 <211> LENGTH: 441
 <212> TYPE: PRT

-continued

```

<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: humanized antibody with mouse CDRs and human
        framework and constant regions
<220> FEATURE:
<221> NAME/KEY: DOMAIN
<222> LOCATION: (1)..(112)
<223> OTHER INFORMATION: variable domain
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (31)..(35)
<223> OTHER INFORMATION: CDRH1
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (50)..(66)
<223> OTHER INFORMATION: CDRH2
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (99)..(101)
<223> OTHER INFORMATION: CDRH3

<400> SEQUENCE: 63

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Lys Pro Gly Gly
1          5          10          15
Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
20         25         30
Ala Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
35         40         45
Ser Glu Ile Ser Gly Gly Gly Ser Tyr Thr Tyr Tyr Pro Asp Thr Val
50         55         60
Thr Gly Arg Phe Thr Ile Ser Arg Asp Asn Ala Lys Asn Ser Leu Tyr
65         70         75         80
Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
85         90         95
Ala Ser Arg Ala Tyr Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser
100        105        110
Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Ser Ser Lys
115        120        125
Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr
130        135        140
Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser
145        150        155        160
Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser
165        170        175
Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Gln Thr
180        185        190
Tyr Ile Cys Asn Val Asn His Lys Pro Ser Asn Thr Lys Val Asp Lys
195        200        205
Arg Val Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys
210        215        220
Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro
225        230        235        240
Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys
245        250        255
Val Val Val Asp Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp
260        265        270
Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu
275        280        285

```

-continued

Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu
 290 295 300
 His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn
 305 310 315 320
 Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly
 325 330 335
 Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Glu Glu
 340 345 350
 Met Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr
 355 360 365
 Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn
 370 375 380
 Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe
 385 390 395 400
 Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn
 405 410 415
 Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr
 420 425 430
 Gln Lys Ser Leu Ser Leu Ser Pro Gly
 435 440

<210> SEQ ID NO 64
 <211> LENGTH: 219
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: humanized antibody with mouse CDRs and human
 framework and constant regions
 <220> FEATURE:
 <221> NAME/KEY: DOMAIN
 <222> LOCATION: (1)..(112)
 <223> OTHER INFORMATION: variable domain
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (24)..(39)
 <223> OTHER INFORMATION: CDRL1
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (55)..(61)
 <223> OTHER INFORMATION: CDRL2
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (94)..(102)
 <223> OTHER INFORMATION: CDRL3

<400> SEQUENCE: 64

Asp Ile Val Met Thr Gln Thr Pro Leu Ser Leu Pro Val Thr Pro Gly
 1 5 10 15
 Glu Pro Ala Ser Ile Ser Cys Arg Ser Ser Gln Ser Ile Val His Ser
 20 25 30
 Gln Gly Asn Thr Tyr Phe Glu Trp Tyr Leu Gln Lys Pro Gly Gln Ser
 35 40 45
 Pro Gln Leu Leu Ile Tyr Lys Val Ser Asn Arg Phe Ser Gly Val Pro
 50 55 60
 Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Lys Ile
 65 70 75 80
 Ser Arg Val Glu Ala Glu Asp Val Gly Val Tyr Tyr Cys Phe Gln Gly
 85 90 95

-continued

Ser	His	Val	Pro	Tyr	Thr	Phe	Gly	Gly	Gly	Thr	Lys	Val	Glu	Ile	Lys
			100					105					110		
Arg	Thr	Val	Ala	Ala	Pro	Ser	Val	Phe	Ile	Phe	Pro	Pro	Ser	Asp	Glu
		115					120					125			
Gln	Leu	Lys	Ser	Gly	Thr	Ala	Ser	Val	Val	Cys	Leu	Leu	Asn	Asn	Phe
	130					135					140				
Tyr	Pro	Arg	Glu	Ala	Lys	Val	Gln	Trp	Lys	Val	Asp	Asn	Ala	Leu	Gln
145					150					155					160
Ser	Gly	Asn	Ser	Gln	Glu	Ser	Val	Thr	Glu	Gln	Asp	Ser	Lys	Asp	Ser
			165						170					175	
Thr	Tyr	Ser	Leu	Ser	Ser	Thr	Leu	Thr	Leu	Ser	Lys	Ala	Asp	Tyr	Glu
		180					185						190		
Lys	His	Lys	Val	Tyr	Ala	Cys	Glu	Val	Thr	His	Gln	Gly	Leu	Ser	Ser
		195					200					205			
Pro	Val	Thr	Lys	Ser	Phe	Asn	Arg	Gly	Glu	Cys					
	210					215									

<210> SEQ ID NO 65

<211> LENGTH: 329

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 65

Ala	Ser	Thr	Lys	Gly	Pro	Ser	Val	Phe	Pro	Leu	Ala	Pro	Ser	Ser	Lys
1			5						10				15		
Ser	Thr	Ser	Gly	Gly	Thr	Ala	Ala	Leu	Gly	Cys	Leu	Val	Lys	Asp	Tyr
		20					25					30			
Phe	Pro	Glu	Pro	Val	Thr	Val	Ser	Trp	Asn	Ser	Gly	Ala	Leu	Thr	Ser
	35					40					45				
Gly	Val	His	Thr	Phe	Pro	Ala	Val	Leu	Gln	Ser	Ser	Gly	Leu	Tyr	Ser
	50					55				60					
Leu	Ser	Ser	Val	Val	Thr	Val	Pro	Ser	Ser	Ser	Leu	Gly	Thr	Gln	Thr
65					70					75					80
Tyr	Ile	Cys	Asn	Val	Asn	His	Lys	Pro	Ser	Asn	Thr	Lys	Val	Asp	Lys
			85						90					95	
Arg	Val	Glu	Pro	Lys	Ser	Cys	Asp	Lys	Thr	His	Thr	Cys	Pro	Pro	Cys
		100						105					110		
Pro	Ala	Pro	Glu	Leu	Leu	Gly	Gly	Pro	Ser	Val	Phe	Leu	Phe	Pro	Pro
	115					120						125			
Lys	Pro	Lys	Asp	Thr	Leu	Met	Ile	Ser	Arg	Thr	Pro	Glu	Val	Thr	Cys
	130				135						140				
Val	Val	Val	Asp	Val	Ser	His	Glu	Asp	Pro	Glu	Val	Lys	Phe	Asn	Trp
145					150					155					160
Tyr	Val	Asp	Gly	Val	Glu	Val	His	Asn	Ala	Lys	Thr	Lys	Pro	Arg	Glu
			165						170					175	
Glu	Gln	Tyr	Asn	Ser	Thr	Tyr	Arg	Val	Val	Ser	Val	Leu	Thr	Val	Leu
			180					185					190		
His	Gln	Asp	Trp	Leu	Asn	Gly	Lys	Glu	Tyr	Lys	Cys	Lys	Val	Ser	Asn
		195					200					205			
Lys	Ala	Leu	Pro	Ala	Pro	Ile	Glu	Lys	Thr	Ile	Ser	Lys	Ala	Lys	Gly
	210					215					220				
Gln	Pro	Arg	Glu	Pro	Gln	Val	Tyr	Thr	Leu	Pro	Pro	Ser	Arg	Glu	Glu
225					230					235					240

-continued

```

Met Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr
      245                      250                      255

Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn
      260                      265                      270

Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe
      275                      280                      285

Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn
      290                      295                      300

Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr
      305                      310                      315                      320

Gln Lys Ser Leu Ser Leu Ser Pro Gly
      325

```

```

<210> SEQ ID NO 66
<211> LENGTH: 329
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: variant of human IgG1f constant domain

<400> SEQUENCE: 66

```

```

Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Ser Ser Lys
1      5      10      15

Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr
      20      25      30

Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser
      35      40      45

Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser
      50      55      60

Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Gln Thr
      65      70      75      80

Tyr Ile Cys Asn Val Asn His Lys Pro Ser Asn Thr Lys Val Asp Lys
      85      90      95

Arg Val Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys
      100     105     110

Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro
      115     120     125

Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys
      130     135     140

Val Val Val Asp Val Glu His Glu Asp Pro Glu Val Lys Phe Asn Trp
      145     150     155     160

Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu
      165     170     175

Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu
      180     185     190

His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn
      195     200     205

Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly
      210     215     220

Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Glu Glu
      225     230     235     240

Met Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr
      245                      250                      255

```

-continued

```

Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn
    260                      265                      270

Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe
    275                      280                      285

Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn
    290                      295                      300

Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr
    305                      310                      315                      320

Gln Lys Ser Leu Ser Leu Ser Pro Gly
    325

```

```

<210> SEQ ID NO 67
<211> LENGTH: 329
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: variant of human IgG1f constant domain

<400> SEQUENCE: 67

```

```

Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Ser Ser Lys
 1      5      10      15

Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr
 20     25     30

Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser
 35     40     45

Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser
 50     55     60

Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Gln Thr
 65     70     75     80

Tyr Ile Cys Asn Val Asn His Lys Pro Ser Asn Thr Lys Val Asp Lys
 85     90     95

Arg Val Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys
100    105    110

Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro
115    120    125

Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys
130    135    140

Val Val Val Asp Val Glu His Glu Asp Pro Glu Val Lys Phe Asn Trp
145    150    155    160

Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu
165    170    175

Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu
180    185    190

His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn
195    200    205

Lys Ala Phe Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly
210    215    220

Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Glu Glu
225    230    235    240

Met Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr
245    250    255

Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn
260                      265                      270

```

-continued

```

Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe
  275                               280               285

Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn
  290                               295               300

Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr
  305                               310               315               320

Gln Lys Ser Leu Ser Leu Ser Pro Gly
  325

```

```

<210> SEQ ID NO 68
<211> LENGTH: 329
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: variant of human IgG1f constant domain

```

```

<400> SEQUENCE: 68

```

```

Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Ser Ser Lys
  1      5      10      15

Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr
  20      25      30

Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser
  35      40      45

Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser
  50      55      60

Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Gln Thr
  65      70      75      80

Tyr Ile Cys Asn Val Asn His Lys Pro Ser Asn Thr Lys Val Asp Lys
  85      90      95

Arg Val Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys
  100     105     110

Pro Ala Pro Glu Leu Leu Gly Gly Asp Ser Val Phe Leu Phe Pro Pro
  115     120     125

Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys
  130     135     140

Val Val Val Asp Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp
  145     150     155     160

Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu
  165     170     175

Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu
  180     185     190

His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn
  195     200     205

Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly
  210     215     220

Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Glu Glu
  225     230     235     240

Met Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr
  245     250     255

Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn
  260     265     270

Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe
  275                               280               285

```

-continued

Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn
 290 295 300

Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr
 305 310 315 320

Gln Lys Ser Leu Ser Leu Ser Pro Gly
 325

<210> SEQ ID NO 69
 <211> LENGTH: 329
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: variant of human IgG1f constant domain
 <400> SEQUENCE: 69

Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Ser Ser Lys
 1 5 10 15

Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr
 20 25 30

Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser
 35 40 45

Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser
 50 55 60

Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Gln Thr
 65 70 75 80

Tyr Ile Cys Asn Val Asn His Lys Pro Ser Asn Thr Lys Val Asp Lys
 85 90 95

Arg Val Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys
 100 105 110

Pro Ala Pro Glu Leu Leu Gly Gly Asp Ser Val Phe Leu Phe Pro Pro
 115 120 125

Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys
 130 135 140

Val Val Val Asp Val Ser His Glu Asp Gly Glu Val Lys Phe Asn Trp
 145 150 155 160

Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu
 165 170 175

Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu
 180 185 190

His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn
 195 200 205

Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly
 210 215 220

Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Glu Glu
 225 230 235 240

Met Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr
 245 250 255

Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn
 260 265 270

Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe
 275 280 285

Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn
 290 295 300

-continued

Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr
305 310 315 320

Gln Lys Ser Leu Ser Leu Ser Pro Gly
325

<210> SEQ ID NO 70
<211> LENGTH: 329
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: variant of human IgG1f constant domain

<400> SEQUENCE: 70

Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Ser Ser Lys
1 5 10 15

Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr
20 25 30

Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser
35 40 45

Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser
50 55 60

Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Gln Thr
65 70 75 80

Tyr Ile Cys Asn Val Asn His Lys Pro Ser Asn Thr Lys Val Asp Lys
85 90 95

Arg Val Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys
100 105 110

Pro Ala Pro Glu Leu Leu Gly Gly Asp Ser Val Phe Leu Phe Pro Pro
115 120 125

Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys
130 135 140

Val Val Val Asp Val Ser His Glu Glu Gly Glu Val Lys Phe Asn Trp
145 150 155 160

Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu
165 170 175

Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu
180 185 190

His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn
195 200 205

Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly
210 215 220

Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Glu Glu
225 230 235 240

Met Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr
245 250 255

Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn
260 265 270

Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe
275 280 285

Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn
290 295 300

Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr
305 310 315 320

-continued

Gln Lys Ser Leu Ser Leu Ser Pro Gly
325

<210> SEQ ID NO 71
<211> LENGTH: 329
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: variant of human IgG1f constant domain

<400> SEQUENCE: 71

Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Ser Ser Lys
1 5 10 15
Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr
20 25 30
Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser
35 40 45
Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser
50 55 60
Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Gln Thr
65 70 75 80
Tyr Ile Cys Asn Val Asn His Lys Pro Ser Asn Thr Lys Val Asp Lys
85 90 95
Arg Val Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys
100 105 110
Pro Ala Pro Asp Leu Leu Gly Gly Asp Ser Val Phe Leu Phe Pro Pro
115 120 125
Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys
130 135 140
Val Val Val Asp Val Ser His Glu Asp Gly Glu Val Lys Phe Asn Trp
145 150 155 160
Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu
165 170 175
Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu
180 185 190
His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn
195 200 205
Lys Ala Leu Pro Arg Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly
210 215 220
Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Glu Glu
225 230 235 240
Met Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr
245 250 255
Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn
260 265 270
Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe
275 280 285
Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn
290 295 300
Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr
305 310 315 320
Gln Lys Ser Leu Ser Leu Ser Pro Gly
325

-continued

<210> SEQ ID NO 72
<211> LENGTH: 329
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: variant of human IgG1f constant domain

<400> SEQUENCE: 72

Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Ser Ser Lys
1 5 10 15
Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr
20 25 30
Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser
35 40 45
Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser
50 55 60
Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Gln Thr
65 70 75 80
Tyr Ile Cys Asn Val Asn His Lys Pro Ser Asn Thr Lys Val Asp Lys
85 90 95
Arg Val Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys
100 105 110
Pro Ala Pro Glu Leu Leu Gly Asp Asp Ser Val Phe Leu Phe Pro Pro
115 120 125
Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys
130 135 140
Val Val Val Asp Val Ser Asp Glu Asp Gly Glu Val Lys Phe Asn Trp
145 150 155 160
Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu
165 170 175
Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu
180 185 190
His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn
195 200 205
Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly
210 215 220
Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Glu Glu
225 230 235 240
Met Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr
245 250 255
Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn
260 265 270
Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe
275 280 285
Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn
290 295 300
Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr
305 310 315 320
Gln Lys Ser Leu Ser Leu Ser Pro Gly
325

<210> SEQ ID NO 73

-continued

<211> LENGTH: 329
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: variant of human IgG1f constant domain

<400> SEQUENCE: 73

```

Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Ser Ser Lys
1      5      10      15
Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr
      20      25      30
Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser
      35      40      45
Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser
      50      55      60
Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Gln Thr
      65      70      75      80
Tyr Ile Cys Asn Val Asn His Lys Pro Ser Asn Thr Lys Val Asp Lys
      85      90      95
Arg Val Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys
      100     105     110
Pro Ala Pro Glu Leu Leu Gly Asp Asp Ser Val Phe Leu Phe Pro Pro
      115     120     125
Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys
      130     135     140
Val Val Val Asp Val Ser His Glu Asp Gly Glu Val Lys Phe Asn Trp
      145     150     155     160
Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu
      165     170     175
Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu
      180     185     190
His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn
      195     200     205
Lys Ala Leu Pro Arg Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly
      210     215     220
Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Glu Glu
      225     230     235     240
Met Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr
      245     250     255
Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn
      260     265     270
Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe
      275     280     285
Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn
      290     295     300
Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr
      305     310     315     320
Gln Lys Ser Leu Ser Leu Ser Pro Gly
      325

```

<210> SEQ ID NO 74
 <211> LENGTH: 329
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence

-continued

```

<220> FEATURE:
<223> OTHER INFORMATION: variant of human IgG1f constant domain

<400> SEQUENCE: 74

Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Ser Ser Lys
1      5      10      15
Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr
20      25      30
Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser
35      40      45
Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser
50      55      60
Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Gln Thr
65      70      75      80
Tyr Ile Cys Asn Val Asn His Lys Pro Ser Asn Thr Lys Val Asp Lys
85      90      95
Arg Val Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys
100     105     110
Pro Ala Pro Glu Leu Leu Gly Asp Asp Ser Val Phe Leu Phe Pro Pro
115     120     125
Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys
130     135     140
Val Val Val Asp Val Ser His Glu Glu Gly Glu Val Lys Phe Asn Trp
145     150     155     160
Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu
165     170     175
Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu
180     185     190
His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn
195     200     205
Lys Ala Leu Pro Arg Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly
210     215     220
Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Glu Glu
225     230     235     240
Met Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr
245     250     255
Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn
260     265     270
Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe
275     280     285
Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn
290     295     300
Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr
305     310     315     320
Gln Lys Ser Leu Ser Leu Ser Pro Gly
325

<210> SEQ ID NO 75
<211> LENGTH: 329
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: variant of human IgG1f constant domain

```

-continued

<400> SEQUENCE: 75

```

Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Ser Ser Lys
1      5      10      15
Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr
20      25      30
Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser
35      40      45
Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser
50      55      60
Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Gln Thr
65      70      75      80
Tyr Ile Cys Asn Val Asn His Lys Pro Ser Asn Thr Lys Val Asp Lys
85      90      95
Arg Val Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys
100     105     110
Pro Ala Pro Glu Leu Leu Gly Asp Asp Ser Val Phe Leu Phe Pro Pro
115     120     125
Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys
130     135     140
Val Val Val Asp Val Ser Asp Glu Asp Gly Glu Val Lys Phe Asn Trp
145     150     155     160
Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu
165     170     175
Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu
180     185     190
His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn
195     200     205
Lys Ala Leu Pro Arg Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly
210     215     220
Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Glu Glu
225     230     235     240
Met Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr
245     250     255
Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn
260     265     270
Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe
275     280     285
Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn
290     295     300
Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr
305     310     315     320
Gln Lys Ser Leu Ser Leu Ser Pro Gly
325

```

<210> SEQ ID NO 76

<211> LENGTH: 329

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: variant of human IgG1f constant domain

<400> SEQUENCE: 76

```

Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Ser Ser Lys

```

-continued

1	5	10	15
Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr	20	25	30
Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser	35	40	45
Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser	50	55	60
Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Gln Thr	65	70	75
Tyr Ile Cys Asn Val Asn His Lys Pro Ser Asn Thr Lys Val Asp Lys	85	90	95
Arg Val Glu Pro Lys Ser Cys Asp Lys Thr His Thr Cys Pro Pro Cys	100	105	110
Pro Ala Pro Asp Leu Leu Gly Asp Asp Ser Val Phe Leu Phe Pro Pro	115	120	125
Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys	130	135	140
Val Val Val Asp Val Ser Asp Glu Asp Gly Glu Val Lys Phe Asn Trp	145	150	155
Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu	165	170	175
Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu	180	185	190
His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn	195	200	205
Lys Ala Leu Pro Arg Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly	210	215	220
Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Glu Glu	225	230	235
Met Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr	245	250	255
Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn	260	265	270
Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe	275	280	285
Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn	290	295	300
Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr	305	310	315
Gln Lys Ser Leu Ser Leu Ser Pro Gly	325		

<210> SEQ ID NO 77

<211> LENGTH: 107

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 77

Arg Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu	1	5	10	15
Gln Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe	20	25	30	

-continued

```

Tyr  Pro  Arg  Glu  Ala  Lys  Val  Gln  Trp  Lys  Val  Asp  Asn  Ala  Leu  Gln
      35              40              45

Ser  Gly  Asn  Ser  Gln  Glu  Ser  Val  Thr  Glu  Gln  Asp  Ser  Lys  Asp  Ser
      50              55              60

Thr  Tyr  Ser  Leu  Ser  Ser  Thr  Leu  Thr  Leu  Ser  Lys  Ala  Asp  Tyr  Glu
      65              70              75              80

Lys  His  Lys  Val  Tyr  Ala  Cys  Glu  Val  Thr  His  Gln  Gly  Leu  Ser  Ser
      85              90              95

Pro  Val  Thr  Lys  Ser  Phe  Asn  Arg  Gly  Glu  Cys
      100              105

```

```

<210> SEQ ID NO 78
<211> LENGTH: 17
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 78

```

```

Met  Arg  Ala  Trp  Ile  Phe  Phe  Leu  Leu  Cys  Leu  Ala  Gly  Arg  Ala  Leu
 1          5          10          15

```

Ala

We claim:

1. An isolated antibody, or antigen binding portion thereof, that specifically binds to human CD40 and competes for binding to human CD40 in a cross-blocking assay with one or more of antibodies selected from the group consisting of 12D6 (SEQ ID NOs: 3 and 4), 5F11 (SEQ ID NOs: 23 and 24), 8E8 (SEQ ID NOs: 40 and 41), 5G7 (SEQ ID NOs: 52 and 53), and 19G3 (SEQ ID NOs: 58 and 59).

2. The isolated antibody or antigen binding portion thereof of claim 1, wherein the competition in a cross-blocking assay comprises the ability to reduce binding of the selected antibody to human CD40 in a competition ELISA by at least 20% when used at an equal molar concentration with the selected antibody.

3. An isolated antibody, or antigen binding portion thereof, that specifically binds to human CD40 at:

a. an epitope comprising or consisting of the sequence EPPTACREKQYLINS (residues 21-35 of SEQ ID NO: 1) (antibodies 12D6, 5G7 and 19G3); or

b. an epitope comprising or consisting of the sequence ECLPCGESE (residues 58-66 of SEQ ID NO: 1) (antibody 5F11).

4. An isolated antibody, or antigen binding portion thereof, that specifically binds to human CD40 comprising:

a) heavy chain CDR sequences derived at least in part from murine V region germline VH1-39_01 and J region germline IGHJ4 and

light chain CDR sequences derived at least in part from murine V region germline VK1-110_01 and J region germline IGKJ1 (12D6);

b) heavy chain CDR sequences derived at least in part from murine V region germline VH1-4_02 and J region germline IGHJ3 and

light chain CDR sequences derived at least in part from murine V region germline VK3-5_01 and J region germline IGKJ5 (5F11);

c) heavy chain CDR sequences derived at least in part from murine V region germline VH1-80_01 and J region germline IGHJ2 and

light chain CDR sequences derived at least in part from murine V region germline VK1-110_01 and J region germline IGKJ2 (8E8);

d) heavy chain CDR sequences derived at least in part from murine V region germline VH1-18_01 and J region germline IGHJ4 and

light chain CDR sequences derived at least in part from murine V region germline VK10-96_01 and J region germline IGKJ2 (5G7); or

e) heavy chain CDR sequences derived at least in part from murine V region germline VH5-9-4_01 and J region germline IGHJ3 and

light chain CDR sequences derived at least in part from murine V region germline VK1-117_01 and J region germline IGKJ2 (19G3).

5. The isolated antibody or antigen binding portion thereof of claim 1 comprising a heavy chain and a light chain, wherein the heavy chain comprises CDRH1, CDRH2 and CDRH3 sequences and the light chain comprises CDRL1, CDRL2 and CDRL3 sequences selected from the group consisting of:

a) the CDRs of antibody 12D6-03 wherein CDRH1, CDRH2 and CDRH3 comprise residues 31-35, 50-66 and 99-108, respectively, of SEQ ID NO:5 and CDRL1, CDRL2 and CDRL3 comprise residues 24-39, 55-61 and 94-102, respectively, of SEQ ID NO:6;

b) the CDRs of antibody 12D6-22 wherein CDRH1, CDRH2 and CDRH3 comprise residues 31-35, 50-66 and 99-108, respectively, of SEQ ID NO:7 and CDRL1, CDRL2 and CDRL3 comprise residues 24-39, 55-61 and 94-102, respectively, of SEQ ID NO:9;

c) the CDRs of antibody 12D6-23 wherein CDRH1, CDRH2 and CDRH3 comprise residues 31-35, 50-66 and 99-108, respectively, of SEQ ID NO:10 and

- CDRL1, CDRL2 and CDRL3 comprise residues 24-39, 55-61 and 94-102, respectively, of SEQ ID NO:11;
- d) the CDRs of antibody 12D6-24 wherein CDRH1, CDRH2 and CDRH3 comprise residues 31-35, 50-66 and 99-108, respectively, of SEQ ID NO:12 and CDRL1, CDRL2 and CDRL3 comprise residues 24-39, 55-61 and 94-102, respectively, of SEQ ID NO:9;
- e) the CDRs of antibody 5F11-17 wherein CDRH1, CDRH2 and CDRH3 comprise residues 31-35, 50-66 and 99-106, respectively, of SEQ ID NO:25 and CDRL1, CDRL2 and CDRL3 comprise residues 24-38, 54-60 and 93-101, respectively, of SEQ ID NO:26;
- f) the CDRs of antibody 5F11-23 wherein CDRH1, CDRH2 and CDRH3 comprise residues 31-35, 50-66 and 99-106, respectively, of SEQ ID NO:27 and CDRL1, CDRL2 and CDRL3 comprise residues 24-38, 54-60 and 93-101, respectively, of SEQ ID NO:28;
- g) the CDRs of antibody 5F11-45 wherein CDRH1, CDRH2 and CDRH3 comprise residues 31-35, 50-66 and 99-106, respectively, of SEQ ID NO:29 and CDRL1, CDRL2 and CDRL3 comprise residues 24-38, 54-60 and 93-101, respectively, of SEQ ID NO:30;
- h) the CDRs of antibody 8E8-56 wherein CDRH1, CDRH2 and CDRH3 comprise residues 31-35, 50-66 and 99-111, respectively, of SEQ ID NO:42 and CDRL1, CDRL2 and CDRL3 comprise residues 24-39, 55-61 and 94-102, respectively, of SEQ ID NO:43;
- i) the CDRs of antibody 8E8-62 wherein CDRH1, CDRH2 and CDRH3 comprise residues 31-35, 50-66 and 99-111, respectively, of SEQ ID NO:44 and CDRL1, CDRL2 and CDRL3 comprise residues 24-39, 55-61 and 94-102, respectively, of SEQ ID NO:45;
- j) the CDRs of antibody 8E8-67 wherein CDRH1, CDRH2 and CDRH3 comprise residues 31-35, 50-66 and 99-111, respectively, of SEQ ID NO:46 and CDRL1, CDRL2 and CDRL3 comprise residues 24-39, 55-61 and 94-102, respectively, of SEQ ID NO:47;
- k) the CDRs of antibody 8E8-70 wherein CDRH1, CDRH2 and CDRH3 comprise residues 31-35, 50-66 and 99-111, respectively, of SEQ ID NO:48 and CDRL1, CDRL2 and CDRL3 comprise residues 24-39, 55-61 and 94-102, respectively, of SEQ ID NO:49;
- l) the CDRs of antibody 8E8-71 wherein CDRH1, CDRH2 and CDRH3 comprise residues 31-35, 50-66 and 99-111, respectively, of SEQ ID NO:50 and CDRL1, CDRL2 and CDRL3 comprise residues 24-39, 55-61 and 94-102, respectively, of SEQ ID NO:51;
- m) the CDRs of antibody 5G7-22 wherein CDRH1, CDRH2 and CDRH3 comprise residues 31-35, 50-66 and 99-102, respectively, of SEQ ID NO:54 and CDRL1, CDRL2 and CDRL3 comprise residues 24-34, 50-56 and 89-97, respectively, of SEQ ID NO:55;
- n) the CDRs of antibody 5G7-25 wherein CDRH1, CDRH2 and CDRH3 comprise residues 31-35, 50-66 and 99-102, respectively, of SEQ ID NO:56 and CDRL1, CDRL2 and CDRL3 comprise residues 24-34, 50-56 and 89-97, respectively, of SEQ ID NO:57;
- o) the CDRs of antibody 19G3-11 wherein CDRH1, CDRH2 and CDRH3 comprise residues 31-35, 50-66 and 99-101, respectively, of SEQ ID NO:60 and CDRL1, CDRL2 and CDRL3 comprise residues 24-39, 55-61 and 94-102, respectively, of SEQ ID NO:62; and
- p) the CDRs of antibody 19G3-22 wherein CDRH1, CDRH2 and CDRH3 comprise residues 31-35, 50-66 and 99-101, respectively, of SEQ ID NO:63 and CDRL1, CDRL2 and CDRL3 comprise residues 24-39, 55-61 and 94-102, respectively, of SEQ ID NO:64.
6. The antibody of claim 5 comprising heavy and light chain variable domain sequences selected from the groups consisting of:
- a) the heavy and light chain variable regions of antibody 12D6-03 comprising residues 1-119 and 1-112 of SEQ ID NO:5 and SEQ ID NO:6, respectively;
 - b) the heavy and light chain variable regions of antibody 12D6-22 comprising residues 1-119 and 1-112 of SEQ ID NO:7 and SEQ ID NO:9, respectively;
 - c) the heavy and light chain variable regions of antibody 12D6-23 comprising residues 1-119 and 1-112 of SEQ ID NO:10 and SEQ ID NO:11, respectively;
 - d) the heavy and light chain variable regions of antibody 12D6-24 comprising residues 1-119 and 1-112 of SEQ ID NO:12 and SEQ ID NO:9, respectively;
 - e) the heavy and light chain variable regions of antibody 5F11-17 comprising residues 1-117 and 1-111 of SEQ ID NO:25 and SEQ ID NO:26, respectively;
 - f) the heavy and light chain variable regions of antibody 5F11-23 comprising residues 1-117 and 1-111 of SEQ ID NO:27 and SEQ ID NO:28, respectively;
 - g) the heavy and light chain variable regions of antibody 5F11-45 comprising residues 1-117 and 1-111 of SEQ ID NO:29 and SEQ ID NO:30, respectively;
 - h) the heavy and light chain variable regions of antibody 8E8-56 comprising residues 1-122 and 1-112 of SEQ ID NO:42 and SEQ ID NO:43, respectively;
 - i) the heavy and light chain variable regions of antibody 8E8-62 comprising residues 1-122 and 1-112 of SEQ ID NO:44 and SEQ ID NO:45, respectively;
 - j) the heavy and light chain variable regions of antibody 8E8-67 comprising residues 1-122 and 1-112 of SEQ ID NO:46 and SEQ ID NO:47, respectively;
 - k) the heavy and light chain variable regions of antibody 8E8-70 comprising residues 1-122 and 1-112 of SEQ ID NO:48 and SEQ ID NO:49, respectively;
 - l) the heavy and light chain variable regions of antibody 8E8-71 comprising residues 1-122 and 1-112 of SEQ ID NO:50 and SEQ ID NO:51, respectively;
 - m) the heavy and light chain variable regions of antibody 5G7-22 comprising residues 1-113 and 1-107 of SEQ ID NO:54 and SEQ ID NO:55, respectively;
 - n) the heavy and light chain variable regions of antibody 5G7-25 comprising residues 1-113 and 1-107 of SEQ ID NO:56 and SEQ ID NO:57, respectively;
 - o) the heavy and light chain variable regions of antibody 19G3-11 comprising residues 1-112 and 1-112 of SEQ ID NO:60 and SEQ ID NO:62, respectively; and
 - p) the heavy and light chain variable regions of antibody 19G3-22 comprising residues 1-112 and 1-112 of SEQ ID NO:63 and SEQ ID NO:64, respectively.
7. The antibody of claim 1 wherein the antibody further comprises an Fc region modified to enhance specificity of binding to FcγRIIb.
8. The antibody of claim 7 exhibiting an A/I ratio of less than 5.
9. The antibody of claim 8 exhibiting an A/I ratio of less than 1.
10. The antibody of claim 7 wherein the modified Fc region is in a heavy chain constant region selected from the group consisting of SE (SEQ ID NO: 66), SELF (SEQ ID

NO: 67), P238D (SEQ ID NO: 68), V4 (SEQ ID NO: 69), V4 D270E (SEQ ID NO: 70), V7 (SEQ ID NO: 71), V8 (SEQ ID NO: 72), V9 (SEQ ID NO: 73), V9 D270E (SEQ ID NO: 74), V11 (SEQ ID NO: 75), and V12 (SEQ ID NO: 76).

11. The antibody of claim **5** comprising heavy and light chain sequences selected from the groups consisting of:

- a) the heavy and light chains of antibody 12D6-24-P238D comprising the sequences of SEQ ID NO:13 and SEQ ID NO:9, respectively;
- b) the heavy and light chains of antibody 12D6-24-SE comprising the sequences of SEQ ID NO:14 and SEQ ID NO:9, respectively;
- c) the heavy and light chains of antibody 12D6-24-SELF comprising the sequences of SEQ ID NO:15 and SEQ ID NO:9, respectively;
- d) the heavy and light chains of antibody 12D6-24-V4 comprising the sequences of SEQ ID NO:16 and SEQ ID NO:9, respectively;
- e) the heavy and light chains of antibody 12D6-24-V4 D270E comprising the sequences of SEQ ID NO:17 and SEQ ID NO:9, respectively;
- f) the heavy and light chains of antibody 12D6-24-V8 comprising the sequences of SEQ ID NO:18 and SEQ ID NO:9, respectively;
- g) the heavy and light chains of antibody 12D6-24-V9 comprising the sequences of SEQ ID NO:19 and SEQ ID NO:9, respectively;
- h) the heavy and light chains of antibody 12D6-24-V9 D270E comprising the sequences of SEQ ID NO:20 and SEQ ID NO:9, respectively;
- i) the heavy and light chains of antibody 12D6-24-V11 comprising the sequences of SEQ ID NO:21 and SEQ ID NO:9, respectively;
- j) the heavy and light chains of antibody 12D6-24-V12 comprising the sequences of SEQ ID NO:22 and SEQ ID NO:9, respectively;
- k) the heavy and light chains of antibody 5F11-45-SE comprising the sequences of SEQ ID NO:31 and SEQ ID NO:30, respectively;
- l) the heavy and light chains of antibody 5F11-45-SELF comprising the sequences of SEQ ID NO:32 and SEQ ID NO:30, respectively;
- m) the heavy and light chains of antibody 5F11-45-V4 comprising the sequences of SEQ ID NO:33 and SEQ ID NO:30, respectively;
- n) the heavy and light chains of antibody 5F11-45-V4 D270E comprising the sequences of SEQ ID NO:34 and SEQ ID NO:30, respectively;
- o) the heavy and light chains of antibody 5F11-45 V8 comprising the sequences of SEQ ID NO:35 and SEQ ID NO:30, respectively;
- p) the heavy and light chains of antibody 5F11-45-V9 comprising the sequences of SEQ ID NO:36 and SEQ ID NO:30, respectively;

q) the heavy and light chains of antibody 5F11-45-V9 D270E comprising the sequences of SEQ ID NO:37 and SEQ ID NO:30, respectively;

r) the heavy and light chains of antibody 5F11-45-V11 comprising the sequences of SEQ ID NO:38 and SEQ ID NO:30, respectively; and

s) the heavy and light chains of antibody 5F11-45-V12 comprising the sequences of SEQ ID NO:39 and SEQ ID NO:30, respectively.

12. A nucleic acid encoding the heavy and/or light chain variable region of the antibody, or antigen binding portion thereof, of claim **1**.

13. An expression vector comprising the nucleic acid molecule of claim **12**.

14. A cell transformed with an expression vector of claim **13**.

15. A method of preparing an anti-human CD40 antibody, or antigen binding portion thereof, comprising:

- a) expressing the antibody, or antigen binding portion thereof, in the cell of claim **14**; and
- b) isolating the antibody, or antigen binding portion thereof, from the cell.

16. A pharmaceutical composition comprising:

- a) the antibody, or antigen binding portion thereof, of claim **1**; and
- b) a carrier.

17. A method of stimulating an immune response in a subject comprising administering to the subject the pharmaceutical composition of claim **16**.

18. The method of claim **17**, wherein the subject has a tumor and an immune response against the tumor is stimulated.

19. The method of claim **17**, wherein the subject has a chronic viral infection and an immune response against the viral infection is stimulated.

20. A method of treating cancer comprising administering to a subject in need thereof a therapeutically effective amount of the pharmaceutical composition of claim **16**.

21. The method of claim **20**, wherein the cancer is selected from the group consisting of: bladder cancer, breast cancer, uterine/cervical cancer, ovarian cancer, prostate cancer, testicular cancer, esophageal cancer, gastrointestinal cancer, pancreatic cancer, colorectal cancer, colon cancer, kidney cancer, head and neck cancer, lung cancer, stomach cancer, germ cell cancer, bone cancer, liver cancer, thyroid cancer, skin cancer, neoplasm of the central nervous system, lymphoma, leukemia, myeloma, sarcoma, and virus-related cancer.

22. A method of treating a chronic viral infection comprising administering to a subject in need thereof a therapeutically effective amount of the pharmaceutical composition of claim **16**.

* * * * *