



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁶ : C12Q 1/68, C12N 9/18, C07K 14/33</p>	<p>A2</p>	<p>(11) International Publication Number: WO 95/17521</p> <p>(43) International Publication Date: 29 June 1995 (29.06.95)</p>
<p>(21) International Application Number: PCT/EP94/04292</p> <p>(22) International Filing Date: 22 December 1994 (22.12.94)</p> <p>(30) Priority Data: 08/172,026 22 December 1993 (22.12.93) US</p> <p>(71) Applicants (for all designated States except US): INSTITUT PASTEUR [FR/FR]; 28, rue du Docteur-Roux, F-75724 Paris Cédex 5 (FR). CENTRE NATIONAL D'ETUDES VETERINAIRES ET ALIMENTAIRES CNEVA [FR/FR]; 22, rue Pierre-Curie, F-94700 Maisons-Alfort (FR).</p> <p>(72) Inventors; and</p> <p>(75) Inventors/Applicants (for US only): FACH, Patrick [FR/FR]; 3, rue G.-Duhamel, F-94000 Créteil (FR). GUILLOU, Jean-Pierre [FR/FR]; 2, allée de la Mare-de-l'Orme, F-94430 Chennevières (FR). POPOFF, Michel [FR/FR]; 18 ter, rue B.-Galliera, F-92140 Clamart (FR).</p> <p>(74) Agent: PHELIP, Bruno; Cabinet Harle & Phelip, 21, rue de la Rochefoucauld, F-75009 Paris (FR).</p>		<p>(81) Designated States: CA, CN, JP, US, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).</p> <p>Published <i>Without international search report and to be republished upon receipt of that report.</i></p>
<p>(54) Title: PRIMERS FOR THE AMPLIFICATION OF GENES CODING FOR THE ENTEROTOXIN AND THE LECITHINASE OF CLOSTRIDIUM PERFRINGENS AND THEIR APPLICATION TO THE DETECTION AND NUMERATION OF THESE BACTERIAE</p>		
<p>(57) Abstract</p> <p>The presence of lecithinase or enterotoxin genes or the presence of Clostridium perfringens bacteriae in a sample is determined by polymerase chain reaction using specific primers. The bacteriae are detected in particular in food samples or fecal samples.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgystan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

Primers for the amplification of genes coding for the enterotoxin and the lecithinase of Clostridium perfringens and their application to the detection and numeration of these bacteriae

5 The present invention relates to primers for the amplification of genes coding for the enterotoxin and the lecithinase , also called phospholipase C, of Clostridium perfringens.

10 Another object of the invention is the application of these primers for the detection and the numeration of C. perfringens.

15 Clostridium perfringens type A is widely distributed (soil, sewage, intestinal tracts of humans and animals), and is a common causative agent of bacterial food poisoning outbreaks worldwide. The symptoms, predominantly diarrhea and abdominal pain, appear 6 to 24 hours after ingestion of contaminated food. Vomiting and fever are unusual. Death occurs occasionally among debilitated persons, particularly the elderly.

20 C. perfringens enterotoxin CPE which is produced during the sporulation phase has been shown to produce the symptoms associated with C. perfringens food poisoning. The illness is caused by the ingestion of food that contains larger number of vegetative enterotoxigenic C. perfringens (more than 10^5 organisms per g) . These bacteriae multiply and sporulate, releasing CPE into the intestine.

30 A C. perfringens count of more than 10^6 /g in fecal samples of ill persons is indicative of C. perfringens food poisoning . In addition, CPE detection directly in fecal samples is a valuable method confirming the diagnosis.

The epidemiological investigations involve *C. perfringens* numeration in suspected foods. The characterization of the enterotoxigenic *C. perfringens* strains is not used routinely, since *C. perfringens* sporulates poorly in usual culture media.

Recently, CPE and phospholipase C gene sequences have been determined (VAN DAMME et al. 1989, Ant. Van Leeuwen 56, 181-190; TSO J. and SIEBEL, 1989. Infect. Immun. 57: 468-476, TITBALL et al. 1989 Infect. Immun. 57:367-376). The phospholipase C gene is located on a variable region of the chromosomal DNA in all *C. perfringens* toxinotypes whereas the distribution of CPE gene is restricted. Only 6% of the *C. perfringens* isolates from various origins showed the presence of CPE gene by DNA-DNA hybridization. This ratio is higher (59%) among *C. perfringens* strains isolated from confirmed outbreaks of food poisoning. In the standard methods, the enterotoxin detection by biological or immunological tests requires previously the sporulation of *C. perfringens*. Several specific medium and protocols for *C. perfringens* sporulation have been described, but they are time consuming and many *C. perfringens* strains do not sporulate or very poorly (DUCAN and STRONG, 1968. Appl. Microbiol. 16: 82; PHILIPS, 1986, Lett. Appl. Microbiol 3: 77-79), which impairs the CPE detection.

A method for the detection of *C. perfringens* by polymerase chain reaction (PCR) has yet been disclosed during the third Congress of the French Society of Microbiology (April 21-24, 1992) through a poster of FACH et al.

The method was consisting in an amplification of the parts of the genes encoding the α -toxin, also called

lecithinase, and the enterotoxin of *C. perfringens* by using oligonucleotidic primers which were chosen in these genes. However, the sequence of the primers used for carrying out this method was not disclosed and the sensitivity mentioned by the authors was low, i.e. from 500 to 5000 bacteriae per gram of feces.

Moreover, the results were obtained on feces artificially contaminated and not on feces from contaminated patients or on contaminated food.

The inventors have thus sought to elaborate a sensitive and reliable method allowing the detection of low quantities of bacteriae in samples of different origins, such as in feces or food, in a raw form.

They have surprisingly shown that it is necessary to choose the primers in some well determined regions of the genes, as well as in the gene of the enterotoxin that in the one of the lecithinase.

Besides, they have carried out a process for the treatment of foods samples, allowing a specific, sensitive and reliable determination of the presence of the *C. perfringens* contained in these foods.

The present invention firstly relates to primers for the amplification of the gene encoding the lecithinase of *C. perfringens*, also called alphatoxin or phospholipase C, corresponding to a part of the sequence of the gene comprised between the nucleotides 1350 and 1850, and preferentially between nucleotides 1350 and 1470 or 1650 and 1850, and which can amplify at least a part of the said gene, by cooperating with other primers having similar features and having a reversed polarity.

The said primers comprise preferentially from 10 to 40 and more preferentially from 10 to 30 nucleotides and

their sequences can be one of the following ones :

SEQ ID1 (PL3) AAG TTA CCT TTG CTG CAT AAT CCC

SEQ ID2 (PL7) ATA GAT ACT CCA TAT CAT CCT GCT

SEQ ID3 (Plc) TCA AAA GAA TAT GCA AGA GGT

5 SEQ ID4 (PL1) TTCTAT CTT GGA GAGG CTATGCAC

SEQ ID5 (PL4) GCTACTAGTTCTTTTACATTCTTTCC.

The invention relates besides to primers for the amplification of the gene encoding enterotoxin of C. perfringens, corresponding to a part of the sequence of
10 the gene comprised between the nucleotides 450 and 950 and preferentially between the nucleotides 450 and 550, or the nucleotides 750 and 950 of said gene, and which can amplify at least a part of the said gene, by cooperating with other primers having similar features and having a
15 reversed polarity.

The said primers comprised preferentially from 10 to 40 and more preferentially from 10 to 30 nucleotides and their sequences can be one of the following ones:

SEQ ID6 (P145) GAA AGA TCT GTA TCT ACA ACT GCT GGT CC

20 SEQ ID7 (P146) GCT GGC TAA GAT TCT ATA TTT TTG TCC AGT

SEQ ID8(Ent A) GAA CGC CAA TCA TAT AAA TTT CCA GCT GGG

The present invention relates moreover to a process for the determination of the presence of the gene encoding the lecithinase, or of the gene encoding the enterotoxin,
25 in a sample, comprising the following steps:

- the DNA from the sample is isolated ,
- parts of the genes encoding the lecithinase, or the enterotoxin, are amplified by polymerization chain reaction (PCR) with the help of specific primers, such as
30 defined hereabove respectively for the lecithinase and for the enterotoxin,

- the amplification products are determined with

the help of known methods.

- The present invention is further directed to a new C. Perfringens beta toxin gene, called β_2 gene, as hereinbelow described.

5 Another object of the present invention is a process for the determination of the presence, and the numeration of C. perfringens in a sample, wherein:

- the DNA from the said sample is isolated ,
- parts of the genes encoding the lecithinase and
10 the enterotoxin are amplified by polymerase chain reaction with the help of specific primers such as defined hereabove for respectively the lecithinase and the enterotoxin ,

- the amplification products are determined with
15 the help of known methods.

The amplification products obtained by the process described hereabove can in particular be determined by electrophoresis on an agarose gel, followed by a transfert of the eluted DNA on "Nylon" membranes and by an
20 hybridation with a labelled probe specific of the amplified part of the gene.

Other methods known by the man skilled in the art can also be used , if such methods are sufficiently specific and sensitive.

25 Such methods are in particular described in "The molecular cloning : A Laboratory Manual; SANDERS et al., Cold Spring Harbor Editor", in which the man skilled in the art can refer for carrying out the processes according to the present invention.

30 The choice of the hereabove described primers did not obviously emerge from the genes sequences such as published. Indeed, the man skilled in the art well knows

that the choice of primers for the amplification of a given DNA part is difficult and that one can be faced with numerous difficulties.

5 For example, primers can lack of specificity or be weakly thermodynamicly stable.

The man skilled in the art can also be faced with other technical difficulties from different types. Thus, the choice that has to be made by the man skilled in the art in the sequence of the gene is difficult because of
10 the important number of thinkable combinations between the two primers corresponding respectively to sequences of the two strands, from the DNA molecule which has to be amplified.

The use of softwares help the man skilled in the art in his choice but does not constitute a method leading
15 automatically and obviously to primers permitting the sought amplification.

The primers such as defined hereabove are able to amplify the said genes if they cooperate with another
20 primer having similar features but of reversed polarity. Indeed, the other primer must be situated on the strand of reversed polarity, called non-sense strand, in such a way that the polymerization of the two strands, initiated from the two primers produce two types of single strand DNA
25 fragments which hybridate and form a double strand DNA molecule, called amplification product, which will be determined by known methods. The primer cooperating in the amplification of the fragment of the said gene have some similar features, i.e. that it is situated in the region
30 of the said gene but on the strand of reversed polarity.

Preferentially, the combinations of the primers can be the following : PL1 and PL4, PL3 and PL7, P145 and

P146, P145 and Ent A.

The processes hereabove described allow the detection of *C. perfringens* in a lot of samples, such as feces or food products, such as convenient foods, without necessitating a pretreatment of the sample.

In a general way, heating of the samples at temperature lysing the bacteriae and separating by centrifugation the DNA from bacterial fragments is sufficient.

In the case of food samples, in particular meats, it can be necessary to perform a pretreatment of the sample in order to eliminate substances which can interfere with the polymerase chain reaction.

Thus, the DNA can be isolated from a food sample by a process comprising the following steps:

- the sample is incubated in a medium allowing the growth of *C. perfringens*,

- the bacteriae are separated from the food particles by centrifugation,

- the bacteriae are put in contact with a resin which lyses them ,

- the DNA is separated from bacterial fragments.

The present invention also relates to a new *C. perfringens* beta toxine gene which has been identified, isolated and characterized.

Said gene, called β_2 gene, can be isolated from the *C. Perfringens* strains of C-type. Two corresponding plasmids were filed on December 12, 1994 at the CNCM Collection of Institut Pasteur and were registered respectively under the reference I-1499 for TGI/pMRP 109 and under the reference I-1500 for TGI/pMRP 126.

	1	AAA	GAA	ATG	GAT	GCG	TAT	AGG	AAA	GTG	ATG	GAG	AAT	TAT	CTT	AAT	GCT	40
	1	K	E	M	D	A	Y	R	K	V	M	E	N	Y	L	N	A	16
	49	TTA	AAA	AAC	TAC	GAT	ATT	AAT	ACA	GTT	GTA	AAC	ATT	TCA	GAA	GAT	GAA	96
	17	L	K	N	Y	D	I	N	T	V	V	N	I	S	E	D	E	32
5	97	AGA	GTA	AAT	AAT	GTT	GAA	CAG	TAT	AGA	GAA	ATG	TTA	GAA	GAT	TTT	AAA	144
	33	R	V	N	N	V	E	Q	Y	R	E	M	L	E	D	F	K	48
	145	TAT	GAT	CCT	AAC	CAA	CAA	CTG	AAA	TCT	TTT	GAA	ATA	CTT	AAT	TCA	CAA	192
	49	Y	D	F	N	Q	Q	L	K	S	F	E	I	L	N	S	Q	64
	193	AAG	AGC	GAT	AAT	AAA	GAA	ATA	TTT	AAT	GTA	AAA	ACT	GAA	TTT	TTA	AAT	240
	65	K	S	D	N	K	E	I	F	N	V	K	T	E	F	L	N	80
10	241	GGT	GCA	ATT	TAT	GAT	ATG	GAA	TTT	ACT	GTA	TCA	TCT	AAA	GAT	GGA	AAA	288
	81	G	A	I	Y	D	M	E	F	T	V	S	S	K	D	G	K	96
	289	TTA	ATA	GTA	TCT	GAT	ATG	GAA	AGA	ACA	AAA	GTT	GAG	AAT	GAA	GGA	AAA	336
	97	L	I	V	S	D	M	E	R	T	K	V	E	N	E	G	K	112
	337	TAT	ATT	TTA	ACA	CCA	TCA	TTT	AGA	ACT	CAA	GTT	TGT	ACA	TGG	GAT	GAT	384
	113	Y	I	L	T	P	S	F	R	T	Q	V	C	T	W	D	D	128
15	385	GAA	CTA	GCA	CAA	GCA	ATT	GGG	GGA	GTT	TAT	CCA	CAA	ACA	TAT	TCT	GAT	432
	129	E	L	A	Q	A	I	G	G	V	Y	P	Q	T	Y	S	D	144
	433	AGA	TTT	ACA	TAT	TAT	GCA	GAT	AAT	ATA	TTA	TTA	AAC	TTC	AGA	CAA	TAT	480
	145	R	F	T	Y	Y	A	D	N	I	L	L	N	F	R	Q	Y	160
	481	GCA	ACT	TCA	GGT	TCA	AGA	GAT	TTA	AAA	GTA	GAA	TAT	AGT	GTT	GTA	GAT	528
	161	A	T	S	G	S	R	D	L	K	V	E	Y	S	V	V	D	176
20	529	CAT	TGG	ATG	TGG	AAA	GAT	GAT	GTT	AAA	GCT	TCT	CAA	ATG	GTA	TAT	GGT	576
	177	H	W	M	W	K	D	D	V	K	A	S	Q	M	V	Y	G	192
	577	CAA	AAT	CCT	GAT	TCT	GCT	AGA	CAA	ATA	AGA	TTA	TAT	ATA	GAA	AAA	GGA	624
	193	Q	N	P	D	S	A	R	Q	I	R	L	Y	I	E	K	G	208
	625	CAA	TCT	TTC	TAT	AAA	TAT	AGA	ATA	AGA	ATT	AAA	AAC	TTT	ACA	CCT	GCA	672
	209	Q	S	F	Y	K	Y	R	I	R	I	K	N	F	T	F	A	224
25	673	TCA	AT															
	225	S																

pMRP109 is a recombinant plasmid containing the beginning of the beta toxin 2 gene.

pMRP109 was produced by insertion into the pUC19 vector restricted by SmaI of a DNA fragment amplified by PCR from the total DNA of strain CWC 245 of *C. perfringens* type C and primers deduced from the N-terminal protein

limited by the following examples in which figure 1 is an ethidium bromide stained agarose gel of amplification products obtained with enterotoxigenic *C. perfringens* type A strain 8-6 (1), type D strain IP76 (2), type E strain NCIB10748 (3), and non enterotoxigenic *C. perfringens* type A strain ATCC13124 (4), type B strain CN3922 (5) and type C strain CWC236 (6) (Panel A), and Southern hybridization with Plc (Panel B), and EntA (Panel C) probes.

MATERIALS AND METHODS

10 Bacterial strains

All bacterial strains used in this study are listed in table 1. The others *C. perfringens* strains were isolated from food poisoning outbreaks and feces of children suffering from diarrhoea (see table 1).

15 *Clostridium* were grown, under anaerobic conditions, in TYG medium: trypticase (30 g/l), yeast extract (20 g/l), glucose (5 g/l) and cysteine HCl (0.5 g/l), pH 7,2. Strains were confirmed as *C. perfringens* by lactose fermentation, nitrate reduction, gelatinase production and
20 absence of motility.

Standard bacteriological methods

For bacteriological food and feces analysis, 10g of the samplee were weighted aseptically into sterile stomacher bags and homogenized for 2 minutes with 90 ml of
25 peptoned water. One ml of decimal dilutions of the suspension was mixed with 9 ml SPS agar: Tryptone (15 g/l), yeast extract (10 g/l), feric citrate (0.5 g/l), sodium sulfite (0.5 g/l), sodium thioglycolate (0.1 g/l), tween 80 (0.005 g/l), polymyxine B sulfite (0.001 g/l),
30 disodic sulfadiazine (0.12 g/l) and agar (14 g/l pH 7). After a 18 h anaerobic incubation at 46°C the number of sulfito-reductor *Clostridium* was determined. Colonies

surrounded with the black characteristic precipitate were identified by biochemical test (lactose fermentation, nitrate reduction, gelatinase production and motility).

Anti C. perfringens enterotoxin immunoglobulins

5 C. perfringens enterotoxin was purified from C. perfringens strain 8-6 and rabbit anti C. perfringens enterotoxin antibodies were prepared as previously described (POPOFF, 1984, FEMS MICROBIOL Let.21: 1-5). Anti C. perfringens enterotoxin immunoglobulins were purified
10 by immunoaffinity. C. perfringens enterotoxin (3- 5 mg) was coupled to 1 g of cyanogen bromide - activated Sepharose 4B (Pharmacia, Paris, France) according to the instructions of the manufacturer. Five ml of rabbit anti C. perfringens enterotoxin serum were passed over the
15 immuno affinity column. The column was washed with phosphate-buffered saline (PBS) until no protein was detected, and eluted with 1 M acetic acid. Fractions of 200 μ l were collected into tubes containing 100 μ l TRIS 3 M pH 8. The fractions containing anti C. perfringens
20 enterotoxin immunoglobulins were dialyzed against PBS.

Preparation of latex beads

Coating of latex beads (Bacto Latex 0.81, Difco, USA) was performed as described (SHAHABADI et al. 1984, Clin. Microbiol. 20: 339-341). Latex suspension (2 - 5 ml)
25 was diluted in 15 ml of glycine buffer (0.1 M glycine, 0.15 M NaCl, pH 8.2) containing 200 μ g of specific anti C. perfringens enterotoxin immunoglobulins, and homogenized for 1 minute at room temperature. Then, 1 volume of PBS - 0.1 % bovine serum albumine (BSA) was added. The mixture
30 was shaken by vortex mixing, and stored at 4°C. The negative latex control was prepared as the same, but using non immune IgG (Sigma, Paris, France).

Slide latex agglutination test (SLAT) for C. perfringens enterotoxin detection.

5 SLAT was performed on a glass slide by mixing 25 μ l of coated latex and 25 μ l of serial ten fold dilutions of samples in PBS - 0,5 % BSA.

The mixture was gently rotated and the results were recorded after 3 minutes.

10 The enterotoxin detection in C. perfringens strains was carried out using SLAT and culture supernatant from cultures in the Duncan and Strong sporulation medium (DAMME-JONGSTEN et al. J. Clin. Microbiol. 28: 131-133).

15 Fecal specimens were diluted at 1 : 10 in PBS-0.5 % BSA, homogenized by vortexing, and centrifuged at 15 000 x g for 30 minutes. The clarified supernatant was tested by SLAT.

Primers used in duplex PCR

20 The oligonucleotides for PCR amplification and for hybridization used in this study, the primers PL3, PL7, PL1, PL4 and Plc were derived from the C. perfringens alpha-toxin gene (11), the primers P145 and P146 and the probe Ent A were from the enterotoxin gene (VAN DAMME et al. 1989, précédemment cités).

25 All the oligonucleotides were chemically synthesized by using a nucleic acid synthesizer (model 380 B; Applied Biosystems Inc. Foster City, Calif.).

Duplex PCR with broth culture

30 One ml of overnight bacterial cultures were transferred to a microcentrifuge tube and centrifuged at 12.000 x g for 3 minutes. The pellet was resuspended in 100 μ l water, heated at 100°C for 10 minutes, and centrifugated at 12000 x g for 10 minutes. Five μ l of the supernatant were submitted to PCR using a PTC-100

programmable Thermal Controller (M.J. Research Inc. Watertown, MA, USA). Each reaction tube contained 50 μ l of a solution of 10 mM Tris-HCl pH 8.3, 50 mM KCl, 1.5 mM MgCl₂, 0.1 mg/ml gelatine, 200 μ M of each of the deoxynucleotide (Boehringer Mannheim, Germany), 0.5 μ M of each primer and DNA sample. Evaporation within the tube was prevented by the addition of 100 μ l mineral oil (Sigma, St-Louis, MO, USA). The reaction mixtures were treated to 94°C for 5 minutes to denature the DNA and then 2.5 units of Taq polymerase (Boehringer) were added to each tube. All PCR reaction were performed using 45 cycles of the optimized thermal profile: 30 s denaturation at 94°C, 30 s primer-annealing at 55°C and 30 s primer-extension at 72°C. After the 45th cycle, the extension reaction was continued for another 10 minutes at 72°C, to ensure that the final extension steps was complete. Negative controls containing all reagents except template DNA were performed in each amplification set. To avoid contamination, sample preparation, PCR amplification and electrophoresis were performed in three different rooms.

Duplex PCR with stool samples

Stool samples (0,1 g) were weighed aseptically into sterile tubes and homogenized with 0,9 ml of water. Samples were centrifugated at 12.000 x g for 3 minutes. The pellet was resuspended in 100 μ l water, heated at 100°C for 10 minutes, centrifugated at 12000 x g for 10 minutes, and 5 μ l of the supernatant were PCR amplified as described above.

Pretreatment of food samples before the amplification reaction.

a) Incubation of the sample.

The detection of viable bacteriae is of great

importance in the food processes.

The discrimination between dead and alived bacteriae can be performed by cultivating the sample.

5 For the detection of *C. perfringens*, 25g of the sample are incubated in 225 ml of buffered peptonized water for 18 hours at 37°C in anaerobiosis and without shaking.

10 The spores of *C. perfringens* are selected with the help of a thermal shock at 80°C of the food culture before incubating.

b) Treatment of the sample for the PCR analysis.

After incubation of the food cultures, the food fragments are sedimentated and 1 ml of the enrichment medium is taken at the surface.

15 A low speed centrifugation (600 rpm) allows the elimination of the lowest food particles. The supernatant is recovered and centrifugated at a high speed (12 000 rpm) for 2 minutes . The supernatant is eliminated and the bacterial pellet is suspended in 2 ml of sterile water.
20 The tube is then centrifuged at 12 000 rpm and the supernatant is eliminated with the help of a Pasteur pipette .

25 200 µl of the "InstaGene" matrix (commercialized by Biorad) are added on the pellet and incubated at 56°C for 30 minutes. The tube is vortexed during 10 seconds and put in a water bath at 100°C during 8 minutes. The tube is then vortexed for 10 seconds and centrifuged at 12 000 rpm for 3 minutes. 3 µl of the supernatant are taken in order to perform the duplex PCR which is finally performed on a
30 volume of 50 µl.

Starting from 25 g of foods containing 10 or more than 10 *C. perfringens* allows the determination in the 15

μ l of the amplified solution of the amplification product.

Gel Electrophoresis

5 A 10 μ l aliquot of amplified DNA was examined after electrophoresis through a composite agarose gel consisting of 3% NuSieve agarose and 1 % SeaKem agarose (FMC BioProducts, Ropckland, (Me, USA) and containing 0.5 μ g/ml ethidium bromide, in 1 x Tris-borate-EDTA buffer (0.089 M Tris, 0.089 M boric acid, 0.002 M EDTA, pH 8). Electrophoresis was carried out for 1 hour at 80 V. The gel was examined visually with an ultra-violet transilluminator.

Hybridization procedure

DNAs electrophoresed on agarose gel were transferred to "Nylon" membranes (Boehringer, Paris, France) using 0.4 N NaOH and fixed by heating the membranes for 15 minutes at 120°C. Filters were pre-hybridized in 5 x Saline Sodium Citrate buffer (SSC), 0.1 % N-laurylsarcosine, 0.02 % Sodium Dodecyl Sulfate (SDS) in 1 % blocking reagent (Boehringer) at 65°C for 1 hour. Hybridization occurred with Ent A or Plc oligonucleotide 3'-endlabelled by digoxigenin 11-DUTP with terminal deoxynucleotidyl-transferase (Boehringer). The probe was added to 50 ng/ml in the pre-hybridization solution and filters were incubated at 40°C for 1 hour. After 2 x SSC, 0.1% SDS washing, membranes were washed to a final stringency of 50°C in 0.1 x SSC, 0.1 % SDS and detection was performed with alkaline phosphatase-labelled anti-digoxigenin antibody (Boehringer), nitroblue tetrayolium (NBT) and bromo-4-chloro-3-indolyl phosphate (BCIP) according to the instruction of the manufacturer (Boehringer).

EXAMPLE 1:

C. perfringens duplex PCR specificity and characterization of the enterotoxigenic strains.

5 The specificity of the duplex PCR with the two primers sets derived from the phospholipase C and CPE genes was assessed by using different Clostridium species and various bacterial species frequently encountered in foods (table 1). The 283 bp amplification product was observed in ethidium bromide stained agarose gel with all C. perfringens strains tested (table 1). This
10 amplification product had the expected size of the phospholipase C gene DNA fragment generated by PL3 and PL7 primers set. This result was confirmed by DNA-DNA hybridization with the internal primer Plc (figure 1).

15 The other bacterial strains tested showed no amplification products with the two primers sets PL3-PL7 and P145-P146 (table 1).

20 Seven C. perfringens strains out of 24 showed an additional 426 bp amplification product which corresponded to the expected size of the CPE DNA fragment delimited with P145 and P146. This 426 bp amplification product hybridized with the Ent A primer localized between P145 and P146 on the CPE DNA sequence and not with Plc primer derived from the phospholipase C gene (figure 1).

25 The C. perfringens strain 8.6 was found enterotoxinogenic by the duplex PCR amplification (table 1). This data is in agreement with the fact that this strain produced CPE using Vero cell cytotoxicity , mouse lethality and immunoprecipitation.

EXAMPLE 2:

30 Sensitivity of the duplex PCR using broth cultures.

The duplex PCR was performed directly with broth cultures of the enterotoxogenic C. perfringens strain 8.6,

in order to determine the sensitivity of the method. The number of bacteriae in culture samples was determined microscopically using a Petrov Chamber. The PCR amplification was carried out as described in material and methods. It was found that as few as 50 bacteria in the reaction volume yielded positive results after agarose gel electrophoresis and ethidium bromide staining. The sensitivity was 10 fold increased when the amplification products electrophoresed on agarose gel were transferred to nylon membrane and hybridized with digoxigenin labelled internal probes Ent A and Plc. In these condituions, the detection limit was approximately 5 bacteria in the reaction volume of the duplex PCR.

EXAMPLE 3

15 Application of the duplex PCR to stool samples.

Twenty three stool samples originated from a suspected *C. perfringens* food borne intoxication outbreak were investigated directly by the duplex PCR and standard microbiological methods for *C. perfringens* detection. Eighteen samples were found positive in the duplex PCR (table 2). All these samples yielded two amplifications products which has the expected size of the DNA fragments delimited by PL3-PL7 and P145 - P146, and hybridized with the internal primers Plc and Ent A respectively, signifying that they contained enterotoxigenic *C. perfringens*. The sulfito reductor bacteria counted by the standard bacteriological methods ranged from less than 10^4 to 10^8 bacteria per gram. Five feces samples were found negative in the duplex PCR and DNA-DNA hybridization test. The corresponding sulfito-reductor bacteria counting was less than 10^4 per gram for 3 samples, and 10^5 to $2 \cdot 10^5$ for the two other (table 2).

EXAMPLE 4C. perfringens enterotoxin SLAT in stool samples.

The CPE was detected in 18 stool samples out of 23 by SLAT as described in material and methods (table 2). Enterotoxigenic C. perfringens was identified by the duplex PCR in 17 of these 18 SLAT positive stool samples. Four samples were negative with both tests.

One stool sample was shown to contain enterotoxigenic C. perfringens by the duplex PCR and no CPE by SLAT. The low number of enterotoxigenic C. perfringens (less than 10^4 per gram) explained the absence of CPE in this example.

CPE was detected by SLAT in one stool sample (table 2) which was negative in the duplex PCR. The level of sulfite reductor bacteria in this sample was 10^5 per gram. However, twenty C. perfringens clones isolated on sheep blood agar, were identified as enterotoxigenic C. perfringens by the duplex PCR.

EXAMPLE 5:Application of the duplex PCR to naturally and artificially contaminated food samples.

The food sample which was responsible of the food intoxication outbreak was found to contain enterotoxigenic C. perfringens by the duplex PCR performed directly without enrichment culture. The counting of sulfite reductor bacteria was 10^5 per g, and no CPE was detected by SLAT.

In order to investigate the sensitivity of this method, the duplex PCR was assayed directly with artificially contaminated food samples. The detection limit was found to be 10^5 C. perfringens per g. The sensitivity was improved by using an overnight enrichment

culture as described in material and methods. Among 59 naturally contaminated food samples, 2 were found to contain $5 \cdot 10^5$ and 10^3 C. perfringens per g respectively by the classical method and were positive by the duplex PCR.

5 All the 57 food samples which did not contain sulfitoreducto bacteria and were artificially contaminated (10 C. perfringens per g) gave a positive result in duplex PCR after enrichment culture (table 3).

CONCLUSION

10 All the 24 C. perfringens strains tested yielded the 426 bp DNA fragment hybridizing with the probe Plc, which corresponds to the presence of the phospholipase C gene. Seven of these C. perfringens strains (2 type A reference strains, 3 type A strains isolated from food,
15 and 2 type D strains) showed an additional amplification product hybridizing with Ent A probe, which indicates the presence of the CPE gene.

The duplex PCR is specific of C. perfringens. No amplification product was observed with 27 Clostridium
20 strains different from C. perfringens and 20 other bacteria species frequently encountered in foods.

An important advantage of this method is to identify easily and quickly the enterotoxigenic C. perfringens strains.

25 The duplex PCR can be used with C. perfringens grown in regular culture medium for anaerobic bacteria without DNA purification. The sensitivity of the duplex PCR using pure culture is 50 bacteria by visualization of ethidium bromide stained amplification product in agarose
30 gel, and 5 bacteria in the reaction volume by DNA-DNA hybridization with the internal probes.

This duplex PCR DNA-DNA hybridization assay was

tested with biological and food samples. Enterotoxigenic *C. perfringens* were detected directly in 18 among 23 stools samples from patients suffering food borne intoxication. These results were in agreement with the
5 numeration of the sulfite reductor bacteria by the standard method and the detection of the CPE by SLAT, except for two stools samples. One stool sample containing 10^5 sulfite reductor bacteria per gram and CPE by SLAT was found negative by the duplex PCR. But, individual *C.*
10 *perfringens* clones isolated from this stool sample were identified as enterotoxigenic *C. perfringens* by the duplex PCR. The presence of PCR inhibitors in this sample accounts probably for this negative result. The other stool sample showed a low number of sulfite-reductor
15 bacteria and no detectable CPE, but was positive in the duplex PCR. A contamination of this sample could explain the positive result in duplex PCR.

The detection limit of enterotoxigenic *C. perfringens* in stools by the duplex PCR is 10^4 and 10^5 per
20 gram. This sensitivity is suitable for the diagnosis of *C. perfringens* food borne intoxication, since it has been reported that *C. perfringens* is recovered at high number (10^6 and more per g) in stools from patients suffering *C. perfringens* food borne intoxication.

25 Controls of food *C. perfringens* contamination constitute an important step in the prevention of food born intoxication by this microorganism. The standard method routinely used detects sulfite-reductor bacteria which encompass *C. perfringens* and also other
30 *Clostridium*. The duplex PCR is specific of *C. perfringens* and discriminates the enterotoxigenic strains which are the causative agents of *C. perfringens* food borne

intoxication. The sensitivity of this method (10 C. perfringens per g) performed with enrichment cultures of food samples is compatible with the prescribed detection levels for food controls. The limit of the duplex PCR directly with food samples without enrichment cultures is 10⁵ bacteria per g, the duplex PCR can be used directly when a C. perfringens food borne intoxication is suspected.

The duplex PCR is more rapid and simple than the standard procedure which needs additional bacteriological tests for the identification of the suspected clones. This method can be applied to food controls for the detection of C. perfringens and identification of the enterotoxigenic strains.

EXAMPLE 6:

Further experiments illustrating the application of the duplex PCR according to the invention to the detection of C. perfringens in foodstuffs, are presented hereinbelow.

1) ANIMAL FEED

Several samples of animal feed have been analyzed for the presence of C. perfringens. These feeds were made up of meat, fish or bone meals. The concentration of sulfite-reductor anaerobic organisms cultivated at 46°C have been determined in 44 naturally contaminated foods and more precisely the presence of C. perfringens spores has been ascertained by use of LS medium (BEERENS H, Romond Ch., Lepage C., Criquelion J. (1982). A liquid medium for the enumeration Clostridium perfringens in food and faeces. In "Isolation and identification methods for foods poisons organisms", pp. 137-149. Ed. Corry J.E.L., Roberts D. and Skinner F.A. Academic Press London) and by

the PCT duplex technique described previously. Spore selection was carried out by thermal shock treatment at 80°C for 10 minutes. The presence of inhibitors for the gene amplification reaction (PCR) led to use an extraction and purification procedure for the nucleic acids which involved the "Insta Gen" (Bio-Rad) matrix and which has been described previously.

On the 44 samples analyzed (table 4) 12 were positive according to the procedure using the LS medium and the PCR duplex test. 24 samples were negative with both methods. 36 samples out of the 44 analyzed gave the same results for the two methods, giving a correlation rate of over 80% between the bacteriological method and the genetic test. For 5 samples the PCR duplex method seemed to be more sensitive than the procedure using the LS medium. For these 5 samples it was possible to isolate, from the thioglycolate liquor incubated for 48 hours, at least one colony on TSC medium which had the biochemical characteristics of *C. perfringens*. The low concentration of *C. perfringens* (< 10/g) in the samples was only detected by the PCR technique, which was more sensitive than the procedure using the LS medium.

Three samples were found to be positive with the procedure using LS medium but stayed negative with the PCR duplex test. These samples contained respectively 7.2 and 1 ASR per gram. In order to ascertain whether the PCR duplex test had not been compromised by the presence of inhibitors remaining after the nucleic acid extraction phase, a few nanograms of *C. perfringens* DNA (2 ng) were added to the extracts obtained from the 3 samples and the gene amplification test was again carried out under the same conditions. The 3 tests then became positive showing

that no inhibitor effect had been observed initially. In addition, the 10 ASR colonies corresponding to these 3 samples were subjected to biochemical identification and a PCR duplex test. These 2 analyses still remained negative, showing evidence of the absence of *C. perfringens* in these samples. In fact, bearing in mind the difficulties in interpretation of the method described by BEERENS and al. (the presence of gas in 1/3 of the Durham belljar and the presence of a black precipitate at the bottom of the tube were sometimes evaluated differently according to the users) and the very doubtful results obtained for these 3 samples, it seems that these 3 samples correspond to false positives in the method described by BEERENS and al.

15 2) HUMAN FOODSTUFFS

Within the framework of systematic monitoring of food hygiene, the number of *C. perfringens* which can be identified is generally low ($< 10^2$ per gram). Various foods (cooked dishes, beef, cooked meats, fish, powdered milk) were analyzed using the PCR duplex technique and the results were compared with those obtained by the standard bacteriological method (standard method ISO 7937).

Only 3 samples were found to be positive according to the two methods, the other 27 containing no *C. perfringens*. The PCR test allowed the detection of *C. perfringens* in 24 hours, while the bacteriological method was much longer (48-72 h). The foods concerned were packs of a frozen ground beef and potato dish. They were contaminated by type A non-enterotoxinogenic *C. perfringens*.

30 EXAMPLE 7:

STUDY OF THE *C. PERFRINGENS* BETA TOXIN GENE

In order to apply a PCR type method to the identification of type C C. perfringens strains responsible for necrotic enteritis, it was necessary to determine the sequence of the beta toxin gene.

5 The applicant carried out the production and purification of the beta toxin from the CWC 245 strain of C. perfringens, then determined peptide sequences, produced oligonucleotide markers and finally carried out the cloning and sequencing of this gene.

10 (1) - Production and purification of the beta toxin.

Cultures were made on TGY (Trypticase 30 g/l, yeast extract 20 g/l, glucose 25 g/l, pH 7.5) in a 2 l fermenter with continuous nitrogen flow, moderate stirring and regulation of pH at 7.5.

15 After 5 hours at 37°C, cholesterol in alcoholic solution (20 mg/l) was added to the culture in order to precipitate the theta toxin. The culture was centrifuged at 10 000 g at 4°C for 20 minutes. The supernatant was precipitated by 60% saturated ammonium sulfate. After
20 centrifuging at 20 000 for 20 minutes, the residue was dissolved in PBS (pH 6.8) and reduced with 50 mM of dithiothreitol. The sample was added in 3 fractions to a biogel P4 (Bio-rad) column equilibrated in PBS. The eluted proteins were applied to a column of thiopropyl-Sysharon
25 6B (58g of resin expanded in PBS) and a flow of 5 ml/h. The column was washed with 100 ml of PBS. Under these conditions the beta and delta toxins were not retained on the column, while the theta and alpha toxins were retained. The effluent was concentrated by ultrafiltration
30 on a PM 10 (Millipore) membrane and submitted to preparative iso-electrofocalization (16 000 V for 16 hours) with a pH gradient of 3.5 to 8. The material

focalized between pH 5.3 and 5.5 was collected and concentrated by ultrafiltration.

The concentrated fraction was added to a column of Biogel P100 (Biorad) equilibrated in PBS and the different
5 fractions were tested by electrophoresis in polyacrylamide-SDS gel. In this way the beta toxin was purified to homogeneity (MW 28 000 and pH 5.4 - 5.5).

The purified beta toxin was lethal to mice (4 μ g, IV), dermonecrotic after injection into guinea-pig skin (2
10 μ g), and enteronecrotic after injection into rabbit intestinal loop (40 μ g). The enterotoxic effect was inhibited by incubation of the beta toxin with trypsin.

(2) Protein microsequencing of the beta toxin and oligonucleotides.

The purified beta toxin was electrophoresed on
15 polyacrylamide-SDS, then transferred onto Immobilon membrane and subjected to microsequencing of the N-terminal region.

The internal protein sequences were obtained by
20 tryptic digestion of the beta toxin, purification of the different peptides by HPLC chromatography and microsequencing of peptides 21, 23 and 13.

Oligonucleotides deduced from the protein sequences were synthesized taking into account of the normal codon
25 of Clostridium and with inosines (I) in the most degenerate positions (figure 2).

(3) Hybridization of the oligonucleotides.

The synthesized oligonucleotides were tested for hybridization with total DNA of CWC 245 digested by
30 different restriction enzymes. The hybridizations were carried out over 16 hours at 40°C with oligonucleotides labeled with ²²P by polynucleotide kinase and washings

were carried out in 6 X SSC at 40°C.

After autoradiography, these oligonucleotides did not clearly reveal a specific DNA fragment. However, clonings were carried out and the clones, identified with the help of the oligonucleotides, did not contain the DNA fragments sought for after verification by nucleotide sequencing.

(4) Amplification of a DNA fragment of the beta toxin gene by PCR.

10 Since the production of recombinant clones containing the beta toxin gene was not possible by conventional cloning methods, this gene was amplified by PCR. To do this, an oligonucleotide (P 279) deduced from the N-terminal protein sequence of the beta toxin and an
15 oligonucleotide (P 280) deduced from an internal protein sequence (peptide 13) but in the reverse direction were used.

A DNA fragment of about 600 bp was thus amplified specifically from the total DNA of CWC 245.

20 The 600 bp DNA fragment was cloned in the pUC19 vector (pMRP 109 recombinant plasmid) and then sequenced. It contained an open reading framework and the terminal portions corresponded to the protein sequences obtained by protein microsequencing (N-terminal and peptide 13),
25 showing that the cloned DNA fragment corresponded well to the gene sought for.

The 3' part of this gene was cloned in pUC 19 using the total DNA of CWC 245 digested by Sau 3A and identified with the P 311 marker whose sequence is localized in the
30 3' part of insert of pMRP 109. The pMRP 126 recombinant clone was sequenced and contained the stop codon of the beta toxin gene and a consensus transcription termination

sequence. The sequence of the beta toxin gene from strain CWC 245 and its translation into amino acids are given in figure 3.

5 (5) Comparison of the beta toxin gene from strain CWC 245 with that from strain NCTC 8533 and genetic analysis of the type C strains.

10 Titball et al (FEMS Microbiology Letters 97 (1992) 77-82) recently published the cloning and sequencing of the beta toxin gene from C. perfringens type B strain NCTC 8533.

15 Comparison of the protein sequence of the beta toxin from strain NCTC 8533 (provisionally referred to as beta 1) with that of the beta toxin gene from strain CWC 245 (provisionally referred to as beta 2) showed no significant homology (figure 4).

 The role of beta toxin 2 as a virulence factor has not yet been clearly established.

20 The presence of the beta 1 and beta 2 genes was searched for by PCR in a series of type B and C C. perfringens strains (Table 1).

 PCR conditions:

 - Buffer :

- 25 - TRIS 10 mM pH 8.2
 - KCl 50 mM
 - dNTP 100 mM
 - Bovine serum albumin 0.1 mg/ml
 - Beta-mercaptoethanol 10 mM
 - Primers 50 pmoles of each per reaction
 - TAQ polymerase (Amersham) 0.5 μ

30 - Primers:

- Primers for beta 2
 - P319 5'-GGAAAGTGATGGAGAATTATCTTAATGC-3'

- P320 5'-GCAGAATCAGGATTTTGACCATATAACC-3'
(reverse)

573 bp amplified fragment.

- Primers for beta 1

5 - BetatoxL 5'-AGGAGGTTTTTTTATTTTTTTTTTTGAAG-3'

- BetatoxR 5'-TCTAAATAGCTGTTACTTTGTG-3'

962 bp amplified fragment.

- Amplification cycles

- 10 - initial denaturation, 2 min at 94°C
- 30 cycles comprising -20 s at 94°C
- 20 s at 50°C
- 20 s at 72°C
- 5 minutes at 72°C

PREM III thermocycler apparatus (Flobio)

15 - Amplification products analysis by electrophoresis on 1 % agar gel containing ethidium bromide and detection by UV transillumination.

20 These strains were identified by the standard method on mice using anti *C. perfringens* neutralizing serum (Wellcome) according to the currently recognized nomenclature; all these strains produced the beta toxin.

25 Genetic analysis showed that the 3 B strains possessed the beta 1 gene and that the type C strains fell into 3 groups: those with the beta 1 gene, those with the beta 2 gene, and those with both (table 5).

The above experimental results show that the beta toxin gene from strain CMC245 differs from the beta toxin gene from strain NCTC 8533.

TABLE 1

Ability of duplex PCR to distinguish enterotoxigenic *C. perfringens* and *C. perfringens* between other *Clostridium* and other enterobacteria.

Bacteria	Strain	PCR Result (1)	Bacteria	Strain	PCR Result (1)
<i>Clostridium perfringens</i> type A	ATCC 13124	+	<i>Clostridium spiroformé</i>	247	-
<i>Clostridium perfringens</i> type B	CN 39.22	+	<i>Clostridium subterminale</i>	ATCC 25774	-
<i>Clostridium perfringens</i> type C	CWC 236	+	<i>Clostridium septicum</i>	ATCC 12464	-
<i>Clostridium perfringens</i> type D	2534	+	<i>Clostridium limosum</i>	384	-
<i>Clostridium perfringens</i> type D	250	+	<i>Clostridium beijerinckii</i>	VPI 5481	-
<i>Clostridium perfringens</i> type D	A0	+	<i>Clostridium mangenotii</i>	ATCC 25761	-
<i>Clostridium perfringens</i> type D	48	+	<i>Clostridium chauvei</i>	IP 91	-
<i>Clostridium perfringens</i> type D	76	+	<i>Bacillus thuringiensis</i>	14001	-
<i>Clostridium perfringens</i> type D	64/1	+	<i>Bacillus subtilis</i>	168	-
<i>Clostridium perfringens</i> type E	NCIB 10748	+	<i>Bacillus cereus</i>	LCHA 1203	-
<i>Clostridium perfringens</i>	8 - 6	+	<i>Listeria monocytogenes</i> 1/2 a	A3	-
<i>Clostridium perfringens</i>	1088.0	+	<i>Listeria monocytogenes</i> 1/2 b	B5	-
<i>Clostridium perfringens</i>	4012	+	<i>Listeria monocytogenes</i> 4 b	O3	-
<i>Clostridium perfringens</i>	4086	+	<i>Listeria innocua</i> 6 a	C3	-
<i>Clostridium perfringens</i>	4009	+	<i>Listeria welshimeri</i> 6 b	E3	-
<i>Clostridium perfringens</i>	4010	+	<i>Listeria seeligeri</i> 1/2 b	D3	-
<i>Clostridium perfringens</i>	4011	+			
<i>Clostridium perfringens</i>	1089.1	+			
<i>Clostridium perfringens</i>	1089.2	+			
<i>Clostridium perfringens</i>	1089.3	+			
<i>Clostridium perfringens</i>	1089.4	+			
<i>Clostridium perfringens</i>	1089.5	+			
<i>Clostridium perfringens</i>	1122	+			
<i>Clostridium perfringens</i>	1513	+			

TABLE 1 (continued)
 Ability of duplex PCR to distinguish enterotoxigenic *C. perfringens*
 and *C. perfringens* between other Clostridium and other enterobacteria

Bacteria	Strain	PCR Result (1)	Bacteria	Strain	PCR Result (1)
<i>Clostridium botulinum</i> type A	ATCC 25763	-	<i>Escherichia coli</i> O : 157 H7	LCHA P1	-
<i>Clostridium botulinum</i> type B	F 11	-	<i>Escherichia coli</i> O : 6 H16	LCHA 877	-
<i>Clostridium botulinum</i> type C	468	-	<i>Escherichia coli</i> O : 125 B15	LCHA 483	-
<i>Clostridium botulinum</i> type D	1875	-	<i>Salmonella typhimurium</i>	3612	-
<i>Clostridium botulinum</i> type E	9009	-	<i>Salmonella enteritidis</i>	5625	-
<i>Clostridium botulinum</i> type F	NCIB 10658	-	<i>Salmonella agona</i>	5628	-
<i>Clostridium botulinum</i> type G	NCIB 10714	-	<i>Proteus sp</i>	7208	-
<i>Clostridium baratii</i> (tox -)	ATCC 27639	-	<i>Kurthia sp</i>	C4	-
<i>Clostridium baratii</i> (tox F)	ATCC 43756	-	<i>Pseudomonas aeruginosa</i>	52	-
<i>Clostridium butyricum</i> (tox -)	CB 25-1	-	<i>Staphylococcus aureus</i>	C3-0	-
<i>Clostridium butyricum</i> (tox E)	ATCC 43181	-	<i>Yersinia enterocolitica</i>	S31	-
<i>Clostridium tetani</i>	E 19406	-			
<i>Clostridium sordellii</i>	IP 82	-			
<i>Clostridium bifermentans</i>	ATCC 638	-			
<i>Clostridium innocuum</i>	NCIB 10574	-			
<i>Clostridium sporogenes</i>	ATCC 3584	-			
<i>Clostridium paraputrificum</i>	ATCC 25780	-			
<i>Clostridium difficile</i>	ATCC 9639	-			
<i>Clostridium difficile</i>	203-1	-			
<i>Clostridium oedematiens</i>	ATCC 17861	-			

(1) Symbols used for PCR results :
 - : Negative reaction for alpha-toxin and enterotoxin genes.
 + : Positive reaction for alpha-toxin gene.
 + + : Positive reaction for alpha-toxin and enterotoxin genes.

Table 2: Stool samples investigation

Samples	Number of sulfite-reductor bacteria	Duplex PCR	SLAT
1	$< 10^4$	-	-
2	$< 10^4$	-	-
3	$< 10^4$	-	-
4	$< 10^4$	+	-
5	10^4	+	+
6	10^4	+	+
7	10^4	+	+
8	$2 \cdot 10^4$	+	+
9	$5 \cdot 10^4$	+	+
10	$5 \cdot 10^4$	+	+
11	10^5	-	+
12	10^5	+	+
13	10^5	+	+
14	$2 \cdot 10^5$	-	-
15	$2 \cdot 10^5$	+	+
16	$8 \cdot 10^5$	+	+
17	10^6	+	+
18	$2 \cdot 10^6$	+	+
19	$2 \cdot 10^6$	+	+
20	10^7	+	+
21	10^7	+	+
22	10^7	+	+
23	10^7	+	+

Table 3. *C. perfringens* investigation in food samples

Food sample	<i>C. perfringens</i> investigation in naturally contaminated food samples			<i>C. perfringens</i> investigation in artificially contaminated food samples	
	Total	Classical method	Duplex PCR	Total	Duplex PCR
Cooked food	31	1 (5×10^5)	1	30	30
Pork butchery	13	1 (10^3)	1	12	12
Raw meat	11	0	0	11	11
Milk product	2	0	0	2	2
Salad	2	0	0	2	2

TABLE 4:

Comparison between LS medium and the PCR
technique in the detection of C. perfringens

5	N = 44 animal foods (bone meal, meat or fish meal)			
			LS MEDIUM (Beerens)	
			+	-
10	PCR	+	12	5 *
		-	3**	24

* ASR concentration < 10/g

At least 1 colony identified as C. perfringens
after isolation on TSC medium.

15 ** No colony isolated on TSC medium was identified as C.
perfringens.

TABLE 5

5 PCR identification of the genes for beta toxin 1
 sequenced from strain NCTC 8533 and beta toxin 2
 sequenced from strain CWC 245 in different strains of type
 B and C *C. perfringens* identified by the standard method

	Beta 1	Beta 2
10 <i>C. perfringens</i> B		
NCTC8533	+	-
NCTC6121	+	-
IP278-220	+	-
CN3922	+	-
15 <i>C. perfringens</i> C		
ATCC3628	+	-
NCTC8081	+	-
NCTC3180	+	+
NCTC3182	+	+
20 NR866	+	+
NR316	+	+
CWC236	+	+
CWC239	+	+
CWC240	+	+
25 CWC241	+	+

TABLE 5 (continued)

5 PCR identification of the genes for beta toxin 1
sequenced from strain NCTC 8533 and beta toxin 2
sequenced from strain CWC 245 in different strains of type
B and C *C. perfringens* identified by the standard method

10

	Beta 1	Beta 2
CWC243	+	+
IP232-87	+	+
IP692-83	+	+
15 CWC235	-	+
CWC237	-	+
CWC245	-	+
CWC249	-	+
S608	-	+
20 IP93R	-	+

WHAT WE CLAIM IS:

1. A primer for the amplification of at least a part of a gene encoding the lecithinase from *C. perfringens*, characterized in that it corresponds to a part of the sequence comprised between nucleotides 1350 and 1850 and preferentially between nucleotides 1350 and 1470 or nucleotides 1650 and 1850 of the said gene, and in that it is able to amplify a part of the said gene, by cooperating with another primer having similar features and a reversed polarity.

2. A primer according to claim 1, characterized in that it comprises almost from 10 to 30 nucleotides.

3. A primer according to one of the claims 1 and 2, characterized in that it comprises one of the following sequences :

SEQ ID1 (PL3) AAG TTA CCT TTG CTG CAT AAT CCC
SEQ ID2 (PL7) ATA GAT ACT CCA TAT CAT CCT GCT
SEQ ID3 (Plc) TCA AAA GAA TAT GCA AGA GGT
SEQ ID4 (PL1) TTCTAT CTT GGA GAGG CTATGCAC
SEQ ID5 (PL4) GCTACTAGTTCTTTTACATTCTTTCC.

4. A primer for the amplification of at least one part of the gene encoding the enterotoxin of *C. perfringens*, characterized in that it corresponds to a part of the sequence comprised between the nucleotides 450 and 950 and preferentially between the nucleotides 450 and 550, or between the nucleotides 750 and 950 of the said gene, and in that it is able to amplify a part of the said gene, by cooperating with another primer having similar features and a reversed polarity.

5. A primer according to claim 4, characterized in that it comprises almost from 10 to 30 nucleotides.

6. A primer according to one of the claims 4 and 5

characterized in that it has one of the following sequences:

SEQ ID6 (P145) GAA AGA TCT GTA TCT ACA ACT GCT GGT CC

SEQ ID7 (P146) GCT GGC TAA GAT TCT ATA TTT TTG TCC AGT

5 SEQ ID8 (Ent A) GAA CGC CAA TCA TAT AAA TTT CCA GCT GGG

7. A process for the detection of the gene encoding the lecithinase in a sample, comprising the following steps:

- DNA from the sample is isolated ,
- 10 - the parts of the genes encoding the lecithinase are amplified by chain polymerisation with the help of specific primers, such as defined in one of the claims 1 to 3,
- the amplification products are determined with
- 15 the help of known methods,

8. A process for the detection of the gene encoding the enterotoxin in a sample, comprising the following steps:

- DNA is isolated from the sample,
- 20 - the parts of the genes encoding the lecithinase are amplified by chain polymerisation with the help of specific primers, such as defined in one of the claims 4 to 6,
- the amplification products are determined with
- 25 the help of known methods,

9. A process for the determination of the presence and the numeration of *C. perfringens* in a sample , comprising the following steps:

- DNA from the sample is isolated ,
- 30 - parts of the genes of *C. perfringens* encoding respectively the lecithinase and the enterotoxin are simultaneously amplified by chain polymerization with the

help of specific primers , such as defined in one of claims 1 to 6 ,

- the amplification products are determined with the help of known methods.

5 10. A process according to one the claims 6 to 9 wherein the DNA is isolated from a food sample according to the following steps:

- the sample is incubated in a medium allowing the growth of *C. perfringens*,

10 - the bacteriae are separated from the food particles par centrifugation,

- the bacteriae are put in contact with a resin lysing them, and

15 - the DNA is separated from the bacterial fragments.

11. The *C. perfringens* beta toxin gene, called β_2 gene, isolated from the *C. perfringens* strains of C type.

20 12. The plasmids corresponding to the gene of claim 11 and registered in the CNCM Collection under the respective references I-1499 and I-1500.




FIGURE 1

peptide 21 (N)-Y-D-I-N-T-V-N-I-S-E-D-E
 P221 5'-TAT GAT ATI AAT ACI GTI AAT ATI TCI GAA GAT GAA-3'

peptide 23 V-M-E-N-Y-L-N-A-L-K-Y-I-L-T-P-S-F
 GTI ATG GAA AAT TAT TTI AAT GCI TTI AAA
 P263 3'-CAI TAC CTT TTA ATA AAI TTA CGI AAI TP-5'

N-terminal K-E-I-D-A-Y-R-K-V-M-E-N-Y-L-N-A-L
 P268 5'-AAA GAA ATI GAT GCI TAT AGI AAA GTI ATG GAA AAT TAT TTI AAT GC-3'
 P279 5'-AAA GAA ATI GAT GCI TAT AGI AAA GTI AIG G-3'

peptide 13 N-F-T-P-A-S-I
 AAT TTI ACI CCI GCI TCI AT
 P280 3'-TTA AAA TGI CGI AGI TA-5'

FIGURE 2: Internal and N-terminal protein microsequencing of the purified beta toxin from strain CWC 245 and corresponding synthetic oligonucleotides.

FIGURE 3: Partial nucleotide sequence and translation into amino acids of beta 2 .

1	AAA	GAA	ATG	GAT	GCG	TAT	AGG	AAA	GTG	ATG	GAG	AAT	TAT	CTT	AAT	GCT	48
1	K	E	M	D	A	Y	R	K	V	M	E	N	Y	L	N	A	16
49	TTA	AAA	AAC	TAC	GAT	ATT	AAT	ACA	GTT	GTA	AAC	ATT	TCA	GAA	GAT	GAA	96
17	L	K	N	Y	D	I	N	T	V	V	N	I	S	E	D	E	32
97	AGA	GTA	AAT	AAT	GTT	GAA	CAG	TAT	AGA	GAA	ATG	TTA	GAA	GAT	TTT	AAA	144
33	R	V	N	N	V	E	Q	Y	R	E	M	L	E	D	F	K	48
145	TAT	GAT	CCT	AAC	CAA	CAA	CTG	AAA	TCT	TTT	GAA	ATA	CTT	AAT	TCA	CAA	192
49	Y	D	P	N	Q	Q	L	K	S	F	E	I	L	N	S	Q	64
193	AAG	AGC	GAT	AAT	AAA	GAA	ATA	TTT	AAT	GTA	AAA	ACT	GAA	TTT	TTA	AAT	240
65	K	S	D	N	K	E	I	F	N	V	K	T	E	F	L	N	80
241	GGT	GCA	ATT	TAT	GAT	ATG	GAA	TTT	ACT	GTA	TCA	TCT	AAA	GAT	GGA	AAA	288
81	G	A	I	Y	D	M	E	F	T	V	S	S	K	D	G	K	96
289	TTA	ATA	GTA	TCT	GAT	ATG	GAA	AGA	ACA	AAA	GTT	GAG	AAT	GAA	GGA	AAA	336
97	L	I	V	S	D	M	E	R	T	K	V	E	N	E	G	K	112
337	TAT	ATT	TTA	ACA	CCA	TCA	TTT	AGA	ACT	CAA	GTT	TGT	ACA	TGG	GAT	GAT	384
113	Y	I	L	T	P	S	F	R	T	Q	V	C	T	W	D	D	128
385	GAA	CTA	GCA	CAA	GCA	ATT	GGG	GGA	GTT	TAT	CCA	CAA	ACA	TAT	TCT	GAT	432
129	E	L	A	Q	A	I	G	G	V	Y	P	Q	T	Y	S	D	144
433	AGA	TTT	ACA	TAT	TAT	GCA	GAT	AAT	ATA	TTA	TTA	AAC	TTC	AGA	CAA	TAT	480
145	R	F	T	Y	Y	A	D	N	I	L	L	N	F	R	Q	Y	160
481	GCA	ACT	TCA	GGT	TCA	AGA	GAT	TTA	AAA	GTA	GAA	TAT	AGT	GTT	GTA	GAT	528
161	A	T	S	G	S	R	D	L	K	V	E	Y	S	V	V	D	176
529	CAT	TGG	ATG	TGG	AAA	GAT	GAT	GTT	AAA	GCT	TCT	CAA	ATG	GTA	TAT	GGT	576
177	H	W	M	W	K	D	D	V	K	A	S	Q	M	V	Y	G	192
577	CAA	AAT	CCT	GAT	TCT	GCT	AGA	CAA	ATA	AGA	TTA	TAT	ATA	GAA	AAA	GGA	624
193	Q	N	P	D	S	A	R	Q	I	R	L	Y	I	E	K	G	208
625	CAA	TCT	TTC	TAT	AAA	TAT	AGA	ATA	AGA	ATT	AAA	AAC	TTT	ACA	CCT	GCA	672
209	Q	S	F	Y	K	Y	R	I	R	I	K	N	F	T	P	A	224
673	TCA	ATT	AGA	GTA	TTT	GGT	GAA	GGG	TAT	TGT	GCA	TAG	AAA	AAA	ATA	TGA	720
225	S	I	R	V	F	G	E	G	Y	C	A	*					236
721	AGT	GAC	TTA	GTC	ACT	TCA	TAT	TTT	TTT	TAC	TAT	TAA	TTT	TAT	TAT	ATA	768
769	AAA	ACC	TAA	CAT	ACA	TGA	AAG	TAT	TCT	TAA	TAC	AGT	TAT	ATC	AAA	ATT	816
817	AAA	GTA	GGG	GAA	ATA	AAA	TAA	AAG	GCT	AAA	AAC	TAT	ATT	AAA	AAC	TAT	864
865	AAA	AAT	TAT	TAA	ATT	AGC											882

FIGURE 4: Comparison of amino acid sequences of the beta 2 (upper line) and beta 1 (lower line) toxins using the GCG Gap program .

Gap Weight: 3.000 Average Match: 0.540
Length Weight: 0.100 Average Mismatch: -0.396

Quality: 73.4 Length: 338
Ratio: 0.326 Gaps: 5
Percent Similarity: 38.117 Percent Identity: 15.695

```

1  .....KEMDAYRKVMENYLNALKN 19
1  MKKKFISLVIVSSLLNGCLLSPTLVYANDIGKTTTITRNKTS DGYTIITQ 50
20 YDINTVVNISEDERVNNVEQY.....REMLEDFKYDPNQQLKSFEIINS 63
51 NDKQIISYQSV DSSSKNEDGFTASIDARFIDDKYSSEMTTLINLTGFMSS 100
64 QKSDNKEIFNV.....KTEFLNGAIYD..... 85
101 KKEDVIKKYNLHDVTNSTAINFPVRY SISILNESINENVKIVDSIPKNTI 150
86 ....MEFTVSSKDGKLI VSDMERTKVENEGKYILTPSF..... 119
151 SQKTVSNTMGYKIGGSIEIEENKPKASIESEYAESSTIEYVQPDFSTIDT 200
120 ..RTQVCTWDEL AQAI GGVPQTYSDRFTYYADNILLNFRQYATSGSRD 167
201 DHSTSKASWDTKFETETTRGNYNLKSNNPV..YGNEMFMYGRYTNVPATEN 248
168 LKVEYSVVDHWMWKDDVKASQMVYQNPDSARQIRLYIEKGQSFYKYRIR 217
249 IIPDYQMSKLI TGG LNPNMSVVL TAPNGTEESIIKVKMERERN CYL NWN 298
218 IKNFTPAS..... 225
299 GANWVGQVYSRLAFDTPNVDSHIFTFKINWLTHKVTAI 336

```