



(51) International Patent Classification:

C07K 16/10 (2006.01) A61P 31/16 (2006.01)
A61K 39/42 (2006.01)

(21) International Application Number:

PCT/EP2012/063637

(22) International Filing Date:

12 July 2012 (12.07.2012)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

11173953.8 14 July 2011 (14.07.2011) EP
61/572,417 14 July 2011 (14.07.2011) US

(71) Applicant (for all designated States except US): **CRUCELL HOLLAND B.V.** [NL/NL]; Archimedesweg 4, NL-2333 CN Leiden (NL).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **KWAKS, Theodorus Hendrikus Jacobus** [NL/NL]; Archimedesweg 4-6, NL-2333 CN Leiden (NL). **ZUIJDEEST, David A.T.M.** [NL/NL]; Archimedesweg 4-6, NL-2333 CN Leiden (NL). **VOGELS, Ronald** [NL/NL]; Archimedesweg 4-6, NL-2333 CN Leiden (NL). **FRIESEN, Robert Heinz Edward** [NL/NL]; Archimedesweg 4-6, NL-2333 CN Leiden (NL).

(74) Agent: **MANTEN, Annemieke**; CRUCELL HOLLAND B.V., Archimedesweg 4-6, NL-2333 CN Leiden (NL).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN,

HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

- as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))
- as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii))
- of inventorship (Rule 4.17(iv))

Published:

- with international search report (Art. 21(3))
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))
- with (an) indication(s) in relation to deposited biological material furnished under Rule 13bis separately from the description (Rules 13bis.4(d)(i) and 48.2(a)(viii))

(54) Title: HUMAN BINDING MOLECULES CAPABLE OF NEUTRALIZING INFLUENZA A VIRUSES OF PHYLOGENETIC GROUP 1 AND PHYLOGENETIC GROUP 2 AND INFLUENZA B VIRUSES

(57) Abstract: The present invention relates to binding molecules, such as human monoclonal antibodies, that bind to an epitope in the stem region of hemagglutinin of influenza A viruses of phylogenetic group 1 and group 2, as well as influenza B viruses, and have a broad neutralizing activity against such influenza viruses. The disclosure provides nucleic acid molecules encoding the binding molecules, their sequences and compositions comprising the binding molecules. The binding molecules can be used in the diagnosis, prophylaxis and/or treatment of influenza A viruses of phylogenetic group 1 and 2, as well as influenza B viruses.



TITLE OF THE INVENTION

Human binding molecules capable of neutralizing influenza A viruses of phylogenetic group 1 and phylogenetic group 2 and influenza B viruses

FIELD OF THE INVENTION

5 The invention relates to medicine. The invention in particular relates to human binding molecules capable of neutralizing influenza A viruses of both phylogenetic group 1 and phylogenetic group 2. In particular, the invention relates to binding molecules capable of neutralizing influenza A viruses of both phylogenetic group 1 and
10 phylogenetic group 2, as well as influenza B viruses. The invention further relates to the diagnosis, prophylaxis and/or treatment of an infection caused by influenza A viruses of phylogenetic groups 1 and 2, and preferably also influenza B viruses.

BACKGROUND OF THE INVENTION

15 Influenza infection (also referred to as “influenza” or “the flu”) is one of the most common diseases known to man causing between three and five million cases of severe illness and between 250,000 and 500,000 deaths every year around the world. Influenza rapidly spreads in seasonal epidemics affecting 5-15% of the population and the burden on health care costs and lost productivity are extensive (World Healthcare Organization
20 (WHO)).

There are 3 types of influenza virus (types A, B and C) responsible for infectious pathologies in humans and animals. The type A and type B viruses are the agents responsible for the influenza seasonal epidemics and pandemics observed in humans.

Influenza A viruses can be classified into influenza virus subtypes based on
25 variations in antigenic regions of two genes that encode the surface glycoproteins hemagglutinin (HA) and neuraminidase (NA) which are required for viral attachment and cellular release. Currently, sixteen subtypes of HA (H1-H16) and nine NA (N1-N9) antigenic variants are known in influenza A virus. Influenza virus subtypes can further be classified by reference to their phylogenetic group. Phylogenetic analysis (Fouchier et al.,
30 2005) has demonstrated a subdivision of HAs comprising two main groups (Air, 1981): *inter alia* the H1, H2, H5 and H9 subtypes in phylogenetic group 1 (herein also referred

to as “group 1”) and *inter alia* the H3, H4 and H7 subtypes in phylogenetic group 2 (or “group 2”). Only some of the influenza A subtypes (i.e. H1N1, H1N2 and H3N2) circulate among people, but all combinations of the 16 HA and 9 NA subtypes have been identified in animals, in particular in avian species. Animals infected with influenza A
5 often act as a reservoir for the influenza viruses and certain subtypes have been shown to cross the species barrier to humans, such as the highly pathogenic influenza A strain H5N1.

The influenza type B virus strains are strictly human. The antigenic variations in HA within the influenza type B virus strains are weaker than those observed within the
10 type A strains. Two genetically and antigenically distinct lineages of influenza B virus are circulating in humans, as represented by the B/Yamagata/16/88 (also referred to as B/Yamagata) and B/Victoria/2/87 (B/Victoria) lineages (Ferguson et al., 2003). Although the spectrum of disease caused by influenza B viruses is generally milder than that caused by influenza A viruses, severe illness requiring hospitalization is still frequently observed
15 with influenza B infection.

Current approaches to dealing with annual influenza epidemics include annual vaccination, preferably generating heterotypic cross-protection. However, circulating influenza viruses in humans are subject to permanent antigenic changes which require annual adaptation of the influenza vaccine formulation to ensure the closest possible
20 match between the influenza vaccine strains and the circulating influenza strains. Although yearly vaccination with influenza vaccines is the best way to prevent influenza, antiviral drugs, such as oseltamivir (Tamiflu®) can be effective for prevention and treatment of influenza infection. The number of influenza virus strains showing resistance against antiviral drugs, such as oseltamivir is, however, increasing.

25 An alternative approach is the development of antibody-based prophylactic or therapeutic treatments to neutralize various seasonal and pandemic influenza viruses. The primary target of most neutralizing antibodies that protect against influenza virus infection is the globular head (HA1 part) of the viral HA protein which contains the receptor binding site, but which is subject to continuing genetic evolution with amino
30 acid substitutions in antibody-binding sites (antigenic drift).

Recently, broadly cross-neutralizing antibodies recognizing an epitope in the conserved stem region of hemagglutinin of influenza A viruses of phylogenetic group 1 (including e.g. the H1 and H5 influenza subtypes) have been identified (see e.g. WO2008/028946), as well as cross-neutralizing antibodies recognizing a highly
5 conserved epitope in the stem region of HA of influenza A viruses of phylogenetic group 2 (including e.g. H3 and H7 subtypes) (WO 2010/130636). The neutralizing activity of these antibodies is restricted to either group 1 or group 2 influenza viruses. In addition, these antibodies are not capable of binding to and neutralizing influenza B viruses.

Furthermore, WO 2010/010466 discloses a human antibody FI6 binding to
10 hemagglutinin and capable of binding to and neutralizing influenza A subtypes of group 1 (including H1 and H5 subtypes) and group 2 (including H3 and H7 subtypes). This antibody also does not bind HA from influenza B viruses.

In addition, US 2009/0092620 discloses a murine antibody recognizing an antigenic structure present in hemagglutinin of both the H1 and the H3 subtype and on
15 hemagglutinin of influenza B viruses belonging to the B/Victoria and B/Yamagata groups. The antibodies inhibit the hemagglutination activity of several H3N2 strains implicating that this antibody binds an epitope in the globular head of HA.

In view of the severity of the respiratory illness caused by influenza A and influenza B viruses, as well as the high economic impact of the seasonal epidemics, and
20 the continuing risk for pandemics, there is an ongoing need for effective means for the prevention and treatment of influenza A and B subtypes. There is thus a need for binding molecules, preferably broadly neutralizing human binding molecules, capable of cross-neutralizing influenza A viruses of both phylogenetic group 1 and phylogenetic group 2, and preferably also influenza B viruses.

25

SUMMARY OF THE INVENTION

The invention provides binding molecules capable of specifically binding to influenza A virus strains from both phylogenetic group 1 (including e.g. influenza viruses comprising HA of the H1 and H5 subtype) and influenza A virus strains from
30 phylogenetic group 2 (including e.g. influenza viruses comprising HA of the H3 and H7 subtype). In an embodiment, the binding molecules also have neutralizing activity against

influenza A virus strains from both phylogenetic group 1 and phylogenetic group 2. In an embodiment, the binding molecules are furthermore capable of specifically binding influenza B virus strains, including e.g. influenza B virus strains of the B/Yamagata and/or B/Victoria lineages. In an embodiment, the binding molecules are furthermore
5 capable of neutralizing influenza B virus strains, including e.g. influenza B virus strains of the B/Yamagata and/or B/Victoria lineages. In an embodiment, the binding molecules are capable of in vivo neutralizing influenza A and/or B virus strains. In an embodiment the binding molecules bind to a conserved epitope in the stem region of the HA protein of influenza A and B viruses. In an embodiment, the binding molecules have no
10 hemagglutination inhibiting (HI) activity.

The invention thus provides binding molecules that bind to an epitope in the stem region of the haemagglutinin protein that is shared between influenza A virus subtypes within the phylogenetic group 1 and influenza virus subtypes within phylogenetic group 2, as well as influenza B virus subtypes, and therefore relates to binding molecules that
15 cross-react between both group 1 and group 2 influenza A virus subtypes and influenza B viruses. The invention also pertains to nucleic acid molecules encoding at least the binding region of the human binding molecules.

The binding molecules and/or nucleic acid molecules of the invention are suitable for use as a universal prophylactic, diagnostic and/or treatment agent for influenza A
20 viruses and influenza B viruses, even irrespective of the causative influenza subtype.

It is surmised that the binding molecules according to the present invention bind to hitherto unknown and highly conserved epitopes that are not or much less prone to antigenic drift or shift. In particular, this epitope is shared between influenza viruses belonging to both phylogenetic group 1 and phylogenetic group 2, and influenza B
25 viruses. It is also encompassed to use the binding molecules of the invention to identify and/or characterize these epitopes.

The invention further provides the use of the human binding molecules and/or the nucleic acid molecules of the invention in the diagnosis, prophylaxis and/or treatment of a subject having, or at risk of developing, an influenza virus infection. Furthermore, the
30 invention pertains to the use of the human binding molecules and/or the nucleic acid molecules of the invention in the diagnosis/detection of such influenza infections.

DESCRIPTION OF THE FIGURES

- FIG. 1 shows the blocking of conformational change of H1, H5, H9, H3, and H7 HAs by CR9114. **(A)** FACS binding of CR9114 to various conformations – uncleaved precursor (HA0); neutral pH, cleaved (HA); fusion pH, cleaved (fusion pH) – of surface-expressed rHA of A/New Caledonia/20/1999 (H1) A/Viet Nam/1203/2004 (H5), A/Hong Kong/1073/1999 (H9), A/Wisconsin/67/2005 (H3), and A/Netherlands/219/2003 (H7). Binding is expressed as the percentage of binding to untreated rHA (HA0). **(B)** FACS binding of CR9114 to surface-expressed HA as above, except that mAb CR9114 was added before exposure of the cleaved HAs to a pH of 4.9.
- FIG. 2 shows that MAb CR9114 competes with CR6261 and CR8020 for binding to H1 and H3, respectively. Additional degree of binding of indicated mAbs to immobilized HA of A/New Caledonia/20/1999 (H1N1) saturated with 100 nM of CR6261 or CR9114 (panels **A** and **B**), or to immobilized HA of A/Wisconsin/67/2005 (H3N2) saturated with 100 nM of CR8020 or CR9114 (panels **C** and **D**), measured using biolayer interferometry.
- FIG. 3 demonstrates the prophylactic efficacy of CR9114 in the mouse lethal challenge model with influenza B (B/Florida/04/2006) virus. **A.** Kaplan-Meier survival curves of mice treated intravenously with either 15 mg/kg CR9114 or vehicle control on day -1 before challenge, followed by a challenge at day 0 of 25 LD B/Florida/04/2006. **B.** Mean bodyweight change (%) relative to day 0. Bars represent 95% CI of the mean. If a mouse died/was euthanized during the study, the last observed bodyweight was carried forward. **C.** Median Clinical scores. Bars represent interquartile ranges. Clinical score explanation: 0=no clinical signs; 1=rough coat; 2=rough coat, less reactive during handling; 3=rough coat, rolled up, laboured breathing, less reactive during handling; 4=rough coat, rolled up, laboured breathing, inactive response to manipulation/handlings.

DESCRIPTION OF THE INVENTION

Definitions of terms as used in the present invention are given below.

- The term “included” or “including” as used herein is deemed to be followed by the words “without limitation”.

As used herein the term “binding molecule” refers to an intact immunoglobulin including monoclonal antibodies, such as chimeric, humanized or human monoclonal antibodies, or to an antigen-binding and/or variable domain comprising fragment of an immunoglobulin that competes with the intact immunoglobulin for specific binding to the binding partner of the immunoglobulin, e.g. HA. Regardless of structure, the antigen-binding fragment binds with the same antigen that is recognized by the intact immunoglobulin. An antigen-binding fragment can comprise a peptide or polypeptide comprising an amino acid sequence of at least 2, 5, 10, 15, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 125, 150, 175, 200, or 250 contiguous amino acid residues of the amino acid sequence of the binding molecule.

The term “binding molecule”, as used herein includes all immunoglobulin classes and subclasses known in the art. Depending on the amino acid sequence of the constant domain of their heavy chains, binding molecules can be divided into the five major classes of intact antibodies: IgA, IgD, IgE, IgG, and IgM, and several of these may be further divided into subclasses (isotypes), e.g., IgA1, IgA2, IgG1, IgG2, IgG3 and IgG4.

Antigen-binding fragments include, inter alia, Fab, F(ab'), F(ab')₂, Fv, dAb, Fd, complementarity determining region (CDR) fragments, single-chain antibodies (scFv), bivalent single-chain antibodies, single-chain phage antibodies, diabodies, triabodies, tetrabodies, (poly)peptides that contain at least a fragment of an immunoglobulin that is sufficient to confer specific antigen binding to the (poly)peptide, etc. The above fragments may be produced synthetically or by enzymatic or chemical cleavage of intact immunoglobulins or they may be genetically engineered by recombinant DNA techniques. The methods of production are well known in the art and are described, for example, in *Antibodies: A Laboratory Manual*, Edited by: E. Harlow and D, Lane (1988), Cold Spring Harbor Laboratory, Cold Spring Harbor, New York, which is incorporated herein by reference. A binding molecule or antigen-binding fragment thereof may have one or more binding sites. If there is more than one binding site, the binding sites may be identical to one another or they may be different.

The binding molecule can be a naked or unconjugated binding molecule but can also be part of an immunoconjugate. A naked or unconjugated binding molecule is intended to refer to a binding molecule that is not conjugated, operatively linked or

otherwise physically or functionally associated with an effector moiety or tag, such as inter alia a toxic substance, a radioactive substance, a liposome, an enzyme. It will be understood that naked or unconjugated binding molecules do not exclude binding molecules that have been stabilized, multimerized, humanized or in any other way manipulated, other than by the attachment of an effector moiety or tag. Accordingly, all post-translationally modified naked and unconjugated binding molecules are included herewith, including where the modifications are made in the natural binding molecule-producing cell environment, by a recombinant binding molecule-producing cell, and are introduced by the hand of man after initial binding molecule preparation. Of course, the term naked or unconjugated binding molecule does not exclude the ability of the binding molecule to form functional associations with effector cells and/or molecules after administration to the body, as some of such interactions are necessary in order to exert a biological effect. The lack of associated effector group or tag is therefore applied in definition to the naked or unconjugated binding molecule in vitro, not in vivo.

As used herein, the term “biological sample” encompasses a variety of sample types, including blood and other liquid samples of biological origin, solid tissue samples such as a biopsy specimen or tissue cultures, or cells derived there from and the progeny thereof. The term also includes samples that have been manipulated in any way after their procurement, such as by treatment with reagents, solubilization, or enrichment for certain components, such as proteins or polynucleotides. The term encompasses various kinds of clinical samples obtained from any species, and also includes cells in culture, cell supernatants and cell lysates.

The term “complementarity determining regions” (CDR) as used herein means sequences within the variable regions of binding molecules, such as immunoglobulins, that usually contribute to a large extent to the antigen binding site which is complementary in shape and charge distribution to the epitope recognized on the antigen. The CDR regions can be specific for linear epitopes, discontinuous epitopes, or conformational epitopes of proteins or protein fragments, either as present on the protein in its native conformation or, in some cases, as present on the proteins as denatured, e.g., by solubilization in SDS. Epitopes may also consist of posttranslational modifications of proteins.

The term “deletion”, as used herein, denotes a change in either amino acid or nucleotide sequence in which one or more amino acid or nucleotide residues, respectively, are absent as compared to the reference, often the naturally occurring, molecule.

5 The term “expression-regulating nucleic acid sequence” as used herein refers to polynucleotide sequences necessary for and/or affecting the expression of an operably linked coding sequence in a particular host organism. The expression-regulating nucleic acid sequences, such as inter alia appropriate transcription initiation, termination, promoter, enhancer sequences; repressor or activator sequences; efficient RNA
10 processing signals such as splicing and polyadenylation signals; sequences that stabilize cytoplasmic mRNA; sequences that enhance translation efficiency (e.g., ribosome binding sites); sequences that enhance protein stability; and when desired, sequences that enhance protein secretion, can be any nucleic acid sequence showing activity in the host organism of choice and can be derived from genes encoding proteins, which are either
15 homologous or heterologous to the host organism. The identification and employment of expression-regulating sequences is routine to the person skilled in the art.

 The term “functional variant”, as used herein, refers to a binding molecule that comprises a nucleotide and/or amino acid sequence that is altered by one or more nucleotides and/or amino acids compared to the nucleotide and/or amino acid sequences
20 of the reference binding molecule and that is capable of competing for binding to the binding partner, i.e. the influenza virus, with the reference binding molecule. In other words, the modifications in the amino acid and/or nucleotide sequence of the reference binding molecule do not significantly affect or alter the binding characteristics of the binding molecule encoded by the nucleotide sequence or containing the amino acid
25 sequence, i.e. the binding molecule is still able to recognize and bind its target. The functional variant may have conservative sequence modifications including nucleotide and amino acid substitutions, additions and deletions. These modifications can be introduced by standard techniques known in the art, such as site-directed mutagenesis and random PCR-mediated mutagenesis, and may comprise natural as well as non-natural
30 nucleotides and amino acids.

Conservative amino acid substitutions include the ones in which the amino acid residue is replaced with an amino acid residue having similar structural or chemical properties. Families of amino acid residues having similar side chains have been defined in the art. These families include amino acids with basic side chains (e.g., lysine, arginine, histidine), acidic side chains (e.g., aspartic acid, glutamic acid), uncharged polar side chains (e.g., asparagine, glutamine, serine, threonine, tyrosine, cysteine, tryptophan), non-polar side chains (e.g., glycine, alanine, valine, leucine, isoleucine, proline, phenylalanine, methionine), beta-branched side chains (e.g., threonine, valine, isoleucine) and aromatic side chains (e.g., tyrosine, phenylalanine, tryptophan). It will be clear to the skilled artisan that also other classifications of amino acid residue families than the one used above can be employed. Furthermore, a variant may have non-conservative amino acid substitutions, e.g., replacement of an amino acid with an amino acid residue having different structural or chemical properties. Similar minor variations may also include amino acid deletions or insertions, or both. Guidance in determining which amino acid residues may be substituted, inserted, or deleted without abolishing immunological activity may be found using computer programs well known in the art.

A mutation in a nucleotide sequence can be a single alteration made at a locus (a point mutation), such as transition or transversion mutations, or alternatively, multiple nucleotides may be inserted, deleted or changed at a single locus. In addition, one or more alterations may be made at any number of loci within a nucleotide sequence. The mutations may be performed by any suitable method known in the art.

The term “influenza virus subtype” as used herein in relation to influenza A viruses refers to influenza A virus variants that are characterized by various combinations of the hemagglutinin (H) and neuramidase (N) viral surface proteins. According to the present invention influenza virus subtypes may be referred to by their H number, such as for example “influenza virus comprising HA of the H1 or H3 subtype”, or “H1 influenza virus” “H3 influenza virus”, or by a combination of a H number and an N number, such as for example “influenza virus subtype H3N2” or “H3N2”.

The term influenza virus “subtype” specifically includes all individual influenza virus “strains” within each subtype, which usually result from mutations and show different pathogenic profiles. Such strains may also be referred to as various “isolates” of

a viral subtype. Accordingly, as used herein, the terms “strains” and “isolates” may be used interchangeably. The current nomenclature for human influenza virus strains or isolates includes the geographical location of the first isolation, strain number and year of isolation, usually with the antigenic description of HA and NA given in brackets, e.g.

5 A/Moscow/10/00 (H3N2). Non-human strains also include the host of origin in the nomenclature.

The term “neutralizing” as used herein in relation to the binding molecules of the invention refers to binding molecules that inhibit an influenza virus from replicatively infecting a target cell, regardless of the mechanism by which neutralization is achieved.

10 Thus, neutralization can e.g. be achieved by inhibiting the attachment or adhesion of the virus to the cell surface, or by inhibition of the fusion of viral and cellular membranes following attachment of the virus to the target cell, and the like.

The term “cross-neutralizing” or “cross-neutralization” as used herein in relation to the binding molecules of the invention refers to the ability of the binding molecules of the invention to neutralize different subtypes of influenza A and/or B viruses.

The term “host”, as used herein, is intended to refer to an organism or a cell into which a vector such as a cloning vector or an expression vector has been introduced. The organism or cell can be prokaryotic or eukaryotic. Preferably, the hosts isolated host cells, e.g. host cells in culture. The term “host cells” merely signifies that the cells are modified for the (over)-expression of the binding molecules of the invention and include B-cells that originally express these binding molecules and which cells have been modified to over-express the binding molecule by immortalization, amplification, enhancement of expression etc. It should be understood that the term host is intended to refer not only to the particular subject organism or cell but to the progeny of such an organism or cell as well. Because certain modifications may occur in succeeding generations due to either mutation or environmental influences, such progeny may not, in fact, be identical to the parent organism or cell, but are still included within the scope of the term “host” as used herein.

The term “human”, when applied to binding molecules as defined herein, refers to molecules that are either directly derived from a human or based upon a human germ line sequence. When a binding molecule is derived from or based on a human sequence and

subsequently modified, it is still to be considered human as used throughout the specification. In other words, the term human, when applied to binding molecules is intended to include binding molecules having variable and constant regions derived from human germline immunoglobulin sequences or based on variable or constant regions occurring in a human or human lymphocyte and modified in some form. Thus, the human binding molecules may include amino acid residues not encoded by human germline immunoglobulin sequences, comprise substitutions and/or deletions (*e.g.*, mutations introduced by for instance random or site-specific mutagenesis *in vitro* or by somatic mutation *in vivo*). “Based on” as used herein refers to the situation that a nucleic acid sequence may be exactly copied from a template, or with minor mutations, such as by error-prone PCR methods, or synthetically made matching the template exactly or with minor modifications.

The term “insertion”, also known as the term “addition”, denotes a change in an amino acid or nucleotide sequence resulting in the addition of one or more amino acid or nucleotide residues, respectively, as compared to the parent sequence.

The term “isolated”, when applied to binding molecules as defined herein, refers to binding molecules that are substantially free of other proteins or polypeptides, particularly free of other binding molecules having different antigenic specificities, and are also substantially free of other cellular material and/or chemicals. For example, when the binding molecules are recombinantly produced, they are preferably substantially free of culture medium components, and when the binding molecules are produced by chemical synthesis, they are preferably substantially free of chemical precursors or other chemicals, *i.e.*, they are separated from chemical precursors or other chemicals which are involved in the synthesis of the protein. The term “isolated” when applied to nucleic acid molecules encoding binding molecules as defined herein, is intended to refer to nucleic acid molecules in which the nucleotide sequences encoding the binding molecules are free of other nucleotide sequences, particularly nucleotide sequences encoding binding molecules that bind other binding partners. Furthermore, the term “isolated” refers to nucleic acid molecules that are substantially separated from other cellular components that naturally accompany the native nucleic acid molecule in its natural host, *e.g.*, ribosomes, polymerases, or genomic sequences with which it is naturally associated.

Moreover, “isolated” nucleic acid molecules, such as cDNA molecules, can be substantially free of other cellular material, or culture medium when produced by recombinant techniques, or substantially free of chemical precursors or other chemicals when chemically synthesized.

5 The term “monoclonal antibody” as used herein refers to a preparation of antibody molecules of single specificity. A monoclonal antibody displays a single binding specificity and affinity for a particular epitope. Accordingly, the term “human monoclonal antibody” refers to an antibody displaying a single binding specificity which has variable and constant regions derived from or based on human germline
10 immunoglobulin sequences or derived from completely synthetic sequences. The method of preparing the monoclonal antibody is not relevant for the binding specificity.

 The term “naturally occurring” as used herein as applied to an object refers to the fact that an object or compound can be found in nature. For example, a polypeptide or polynucleotide sequence that is present in an organism that can be isolated from a source
15 in nature and which has not been intentionally modified by man in the laboratory is naturally occurring.

 The term “nucleic acid molecule” as used in the present invention refers to a polymeric form of nucleotides and includes both sense and anti-sense strands of RNA, cDNA, genomic DNA, and synthetic forms and mixed polymers of the above. A
20 nucleotide refers to a ribonucleotide, deoxynucleotide or a modified form of either type of nucleotide. The term also includes single- and double-stranded forms of DNA. In addition, a polynucleotide may include either or both naturally occurring and modified nucleotides linked together by naturally occurring and/or non-naturally occurring nucleotide linkages. The nucleic acid molecules may be modified chemically or
25 biochemically or may contain non-natural or derivatized nucleotide bases, as will be readily appreciated by those of skill in the art. Such modifications include, for example, labels, methylation, substitution of one or more of the naturally occurring nucleotides with an analogue, internucleotide modifications such as uncharged linkages (*e.g.*, methyl phosphonates, phosphotriesters, phosphoramidates, carbamates, etc.), charged linkages
30 (*e.g.*, phosphorothioates, phosphorodithioates, etc.), pendent moieties (*e.g.*, polypeptides), intercalators (*e.g.*, acridine, psoralen, etc.), chelators, alkylators, and modified linkages

(*e.g.*, alpha anomeric nucleic acids, etc.). The above term is also intended to include any topological conformation, including single-stranded, double-stranded, partially duplexed, triplex, hairpinned, circular and padlocked conformations. Also included are synthetic molecules that mimic polynucleotides in their ability to bind to a designated sequence via hydrogen bonding and other chemical interactions. Such molecules are known in the art and include, for example, those in which peptide linkages substitute for phosphate linkages in the backbone of the molecule. A reference to a nucleic acid sequence encompasses its complement unless otherwise specified. Thus, a reference to a nucleic acid molecule having a particular sequence should be understood to encompass its complementary strand, with its complementary sequence. The complementary strand is also useful, *e.g.*, for anti-sense therapy, hybridisation probes and PCR primers.

The term “operably linked” refers to two or more nucleic acid sequence elements that are usually physically linked and are in a functional relationship with each other. For instance, a promoter is operably linked to a coding sequence, if the promoter is able to initiate or regulate the transcription or expression of a coding sequence, in which case the coding sequence should be understood as being “under the control of” the promoter.

By “pharmaceutically acceptable excipient” is meant any inert substance that is combined with an active molecule such as a drug, agent, or binding molecule for preparing an agreeable or convenient dosage form. The “pharmaceutically acceptable excipient” is an excipient that is non-toxic to recipients at the used dosages and concentrations, and is compatible with other ingredients of the formulation comprising the drug, agent or binding molecule. Pharmaceutically acceptable excipients are widely applied and known in the art.

The term “specifically binding”, as used herein, in reference to the interaction of a binding molecule, *e.g.* an antibody, and its binding partner, *e.g.* an antigen, means that the interaction is dependent upon the presence of a particular structure, *e.g.* an antigenic determinant or epitope, on the binding partner. In other words, the antibody preferentially binds or recognizes the binding partner even when the binding partner is present in a mixture of other molecules or organisms. The binding may be mediated by covalent or non-covalent interactions or a combination of both. In yet other words, the term “specifically binding” means immunospecifically binding to an antigenic determinant or

epitope and not immunospecifically binding to other antigenic determinants or epitopes.

A binding molecule that immunospecifically binds to an antigen may bind to other peptides or polypeptides with lower affinity as determined by, *e.g.*, radioimmunoassays (RIA), enzyme-linked immunosorbent assays (ELISA), BIACORE, or other assays

5 known in the art. Binding molecules or fragments thereof that immunospecifically bind to an antigen may be cross-reactive with related antigens, carrying the same epitope.

Preferably, binding molecules or fragments thereof that immunospecifically bind to an antigen do not cross-react with other antigens.

A “substitution”, as used herein, denotes the replacement of one or more amino
10 acids or nucleotides by different amino acids or nucleotides, respectively.

The term “therapeutically effective amount” refers to an amount of the binding molecule as defined herein that is effective for preventing, ameliorating and/or treating a condition resulting from infection with an influenza B virus. Amelioration as used in
15 herein may refer to the reduction of visible or perceptible disease symptoms, viremia, or any other measurable manifestation of influenza infection.

The term “treatment” refers to therapeutic treatment as well as prophylactic or preventative measures to cure or halt or at least retard disease progress. Those in need of treatment include those already inflicted with a condition resulting from infection with influenza virus as well as those in which infection with influenza virus is to be prevented.
20 Subjects partially or totally recovered from infection with influenza virus might also be in need of treatment. Prevention encompasses inhibiting or reducing the spread of influenza virus or inhibiting or reducing the onset, development or progression of one or more of the symptoms associated with infection with influenza virus.

The term “vector” denotes a nucleic acid molecule into which a second nucleic
25 acid molecule can be inserted for introduction into a host where it will be replicated, and in some cases expressed. In other words, a vector is capable of transporting a nucleic acid molecule to which it has been linked. Cloning as well as expression vectors are contemplated by the term “vector”, as used herein. Vectors include, but are not limited to, plasmids, cosmids, bacterial artificial chromosomes (BAC) and yeast artificial
30 chromosomes (YAC) and vectors derived from bacteriophages or plant or animal (including human) viruses. Vectors comprise an origin of replication recognized by the

proposed host and in case of expression vectors, promoter and other regulatory regions recognized by the host. A vector containing a second nucleic acid molecule is introduced into a cell by transformation, transfection, or by making use of viral entry mechanisms. Certain vectors are capable of autonomous replication in a host into which they are introduced (e.g., vectors having a bacterial origin of replication can replicate in bacteria). Other vectors can be integrated into the genome of a host upon introduction into the host, and thereby are replicated along with the host genome.

DETAILED DESCRIPTION

10 In a first aspect the present invention encompasses binding molecules capable of specifically binding to hemagglutinin (HA) of influenza A virus subtypes of phylogenetic group 1 and influenza A virus subtypes of phylogenetic group 2. In an embodiment, the binding molecules are capable of neutralizing influenza A virus subtypes of both phylogenetic group 1 and phylogenetic group 2. The binding molecules of the invention thus are unique in that they are capable of cross-neutralizing group 1 influenza A virus strains and group 2 influenza A virus strains. In an embodiment, the binding molecules are capable of neutralizing at least one or more, preferably two or more, preferably three or more, preferably four or more, even more preferably five or more group 1 influenza A virus subtypes selected from the group consisting of the H1, H2, H5, H6, H8, H9 and H11 subtype, and at least one or more, preferably two or more, preferably three or more group 2 influenza A virus subtypes selected from the group consisting of the H3, H4, H7, and H10 subtype. In an embodiment, the binding molecules are capable of specifically binding to hemagglutinin (HA) of influenza B virus subtypes. In another embodiment, the binding molecules are capable of neutralizing influenza B viruses. In an embodiment, the binding molecules are capable of *in vivo* neutralizing influenza A and/or B viruses. The influenza A and B virus strains may be both human and non-human influenza virus strains (i.e. obtained from non-human animals, e.g. birds).

Preferably, the binding molecules are human binding molecules. In a preferred embodiment, the binding molecules are human antibodies, or antigen-binding fragments thereof.

In an embodiment, the binding molecules are derived from the VH1-69 germ line gene. Thus, the binding molecules all use the same VH1-69 germ line encoded framework.

In an embodiment, the binding interaction of the binding molecules, preferably the antibody, and HA is mediated exclusively by heavy chain variable sequences.

In an embodiment, the binding molecules comprise a heavy chain CDR1 comprising the amino acid sequence of SEQ ID NO: 133 or SEQ ID NO: 139, a heavy chain CDR2 comprising the amino acid sequence of SEQ ID NO: 134, SEQ ID NO: 140 or SEQ ID NO: 151, and a heavy chain CDR3 comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 135, SEQ ID NO: 141, SEQ ID NO: 145, SEQ ID NO: 152, SEQ ID NO: 161, and SEQ ID NO: 162. The CDR regions of binding molecules of the invention are shown in Table 7. CDR regions are according to Kabat *et al.* (1991) as described in Sequences of Proteins of Immunological Interest.

Influenza viruses infect cells by binding to sialic acid residues on the cell surface of target cells, and following transfer into endosomes, by fusing their membranes with the endosomal membranes and releasing the genome-transcriptase complex into the cell. Both receptor binding and membrane fusion process are mediated by the HA glycoprotein. The HA of influenza virus A comprises two structurally distinct regions, i.e. a globular head region, which contains a receptor binding site which is responsible for virus attachment to the target cell, and is involved in the haemagglutination activity of HA, and a stem region, containing a fusion peptide which is necessary for membrane fusion between the viral envelope and the endosomal membrane of the cell. The HA protein is a trimer in which each monomer consists of two disulphide - linked glycopolypeptides, HA1 and HA2, that are produced during infection by proteolytic cleavage of a precursor (HA0). Cleavage is necessary for virus infectivity since it is required to prime the HA for membrane fusion, to allow conformational change. Activation of the primed molecule occurs at low pH in endosomes, between pH5 and pH6, and requires extensive changes in HA structure. Each of the stages in the priming and activation of HA for its participation in the membrane fusion process, presents a different target for inhibition, e.g. by monoclonal antibodies. In an embodiment, the

binding molecules are capable of blocking the pH-induced conformational changes in HA associated with membrane fusion.

The binding molecules of the invention may be capable of specifically binding to the HA0, HA1 and/or HA2 subunit of the HA protein. They may be capable of specifically binding to linear or structural and/or conformational epitopes on the HA0, HA1 and/or HA2 subunit of the HA protein. The HA molecule may be purified from viruses or recombinantly produced and optionally isolated before use. Alternatively, HA may be expressed on the surface of cells. In an embodiment, the binding molecules of the invention are capable of specifically binding to an epitope in the stem region of HA. In an embodiment, the binding molecules bind to an epitope that is accessible in the pre-fusion conformation of HA.

The binding molecules of the invention may be capable of specifically binding to influenza viruses that are viable, living and/or infective or that are in inactivated/attenuated form. Methods for inactivating/attenuating virus, e.g. influenza viruses are well known in the art and include, but are not limited to, treatment with formalin, β -propiolactone (BPL), merthiolate, and/or ultraviolet light.

The binding molecules of the invention may also be capable of specifically binding to one or more fragments of the influenza viruses, such as *inter alia* a preparation of one or more proteins and/or (poly)peptides, derived from subtypes of influenza A and/or B viruses or one or more recombinantly produced proteins and/or polypeptides of influenza A and/or B viruses. The nucleotide and/or amino acid sequence of proteins of various influenza A and B strains can be found in the GenBank-database, NCBI Influenza Virus Sequence Database, Influenza Sequence Database (ISD), EMBL-database and/or other databases. It is well within the reach of the skilled person to find such sequences in the respective databases.

In another embodiment the binding molecules of the invention are capable of specifically binding to a fragment of the above-mentioned proteins and/or polypeptides, wherein the fragment at least comprises an epitope recognized by the binding molecules of the invention. An "epitope" as used herein is a moiety that is capable of binding to a binding molecule of the invention with sufficiently high affinity to form a detectable antigen-binding molecule complex.

The binding molecules of the invention may or may not be capable of specifically binding to the extracellular part of HA (also called herein soluble HA (sHA)).

The binding molecules of the invention can be intact immunoglobulin molecules such as polyclonal or monoclonal antibodies or the binding molecules can be antigen-binding fragments thereof, including, but not limited to, heavy and light chain variable regions, Fab, F(ab'), F(ab')₂, Fv, dAb, Fd, complementarity determining region (CDR) fragments, single-chain antibodies (scFv), bivalent single-chain antibodies, single-chain phage antibodies, diabodies, triabodies, tetrabodies, and (poly)peptides that contain at least a fragment of an immunoglobulin that is sufficient to confer specific antigen binding to influenza virus strains or a fragment thereof. In a preferred embodiment the binding molecules of the invention are human monoclonal antibodies, and/or antigen-binding fragments thereof. The binding molecules may also be Nanobodies, alphabodies, affibodies, FN3-domain scaffolds and other scaffolds based on domains in (human) repeat proteins like Adnectins, Anticalins, Darpins, etc, or other scaffolds comprising epitope binding sequences.

The binding molecules of the invention can be used in non-isolated or isolated form. Furthermore, the binding molecules of the invention can be used alone or in a mixture comprising at least one binding molecule (or variant or fragment thereof) of the invention, and/or with other binding molecules that bind to influenza and have influenza virus inhibiting effect. In other words, the binding molecules can be used in combination, *e.g.*, as a pharmaceutical composition comprising two or more binding molecules of the invention, variants or fragments thereof. For example, binding molecules having different, but complementary activities can be combined in a single therapy to achieve a desired prophylactic, therapeutic or diagnostic effect, but alternatively, binding molecules having identical activities can also be combined in a single therapy to achieve a desired prophylactic, therapeutic or diagnostic effect. Optionally, the mixture further comprises at least one other therapeutic agent. Preferably, the therapeutic agent such as, *e.g.*, M2 inhibitors (*e.g.*, amantidine, rimantadine) and/or neuraminidase inhibitors (*e.g.*, zanamivir, oseltamivir) is useful in the prophylaxis and/or treatment of an influenza virus infection

Typically, binding molecules according to the invention can bind to their binding partners, i.e. an influenza A virus of group 1 (such as H1N1) and an influenza A virus of group 2 (such as H3N2), and/or an influenza B virus, and/or fragments thereof, with an affinity constant (K_d -value) that is lower than 0.2×10^{-4} M, 1.0×10^{-5} M, 1.0×10^{-6} M,
5 1.0×10^{-7} M, preferably lower than 1.0×10^{-8} M, more preferably lower than 1.0×10^{-9} M, more preferably lower than 1.0×10^{-10} M, even more preferably lower than 1.0×10^{-11} M, and in particular lower than 1.0×10^{-12} M. The affinity constants can vary for antibody isotypes. For example, affinity binding for an IgM isotype refers to a binding affinity of at least about 1.0×10^{-7} M. Affinity constants can for instance be measured using surface
10 plasmon resonance, for example using the BIACORE system (Pharmacia Biosensor AB, Uppsala, Sweden).

The binding molecules of the invention exhibit neutralizing activity. Neutralizing activity can for instance be measured as described herein. Alternative assays measuring neutralizing activity are described in for instance WHO Manual on Animal Influenza
15 Diagnosis and Surveillance, Geneva: World Health Organisation, 2005, version 2002.5.

Typically, the binding molecules according to the invention have a neutralizing activity of $50 \mu\text{g/ml}$ or less, preferably $20 \mu\text{g/ml}$ or less, more preferably a neutralizing activity of $10 \mu\text{g/ml}$ or less, even more preferably $5 \mu\text{g/ml}$ or less, as determined in an *in vitro* virus neutralization assay (VNA) as described in Example 6. The binding molecules
20 according to the invention may bind to influenza virus or a fragment thereof in soluble form such as for instance in a sample or in suspension or may bind to influenza viruses or fragments thereof bound or attached to a carrier or substrate, e.g., microtiter plates, membranes and beads, etc. Carriers or substrates may be made of glass, plastic (e.g., polystyrene), polysaccharides, nylon, nitrocellulose, or Teflon, etc. The surface of such
25 supports may be solid or porous and of any convenient shape. Furthermore, the binding molecules may bind to influenza virus in purified/isolated or non-purified/non-isolated form.

As discussed above, the present invention relates to isolated human binding molecules that are able to recognize and bind to an epitope in the influenza
30 haemagglutinin protein (HA) wherein said binding molecules have neutralizing activity against influenza A viruses of phylogenetic group 1 and influenza A viruses of

phylogenetic group 2. According to the invention, it thus has been shown that the binding molecules of the present invention cross-neutralize influenza virus subtypes belonging to both phylogenetic groups. The skilled person, based on what has been disclosed herein, can determine whether an antibody indeed cross-reacts with HA proteins from different subtypes and can also determine whether they are able to neutralize influenza viruses of different subtypes *in vitro* and/or *in vivo*.

In an embodiment the binding molecule according to the present invention is selected from the group consisting of:

- a) a binding molecule comprising a heavy chain CDR1 region of SEQ ID NO:133, a heavy chain CDR2 region of SEQ ID NO:134, and a heavy chain CDR3 region of SEQ ID NO:135,
- b) a binding molecule comprising a heavy chain CDR1 region of SEQ ID NO:139, a heavy chain CDR2 region of SEQ ID NO:140, and a heavy chain CDR3 region of SEQ ID NO:141,
- c) a binding molecule comprising a heavy chain CDR1 region of SEQ ID NO:139, a heavy chain CDR2 region of SEQ ID NO:134, and a heavy chain CDR3 region of SEQ ID NO:145,
- d) a binding molecule comprising a heavy chain CDR1 region of SEQ ID NO:139, a heavy chain CDR2 region of SEQ ID NO:151, and a heavy chain CDR3 region of SEQ ID NO:152,
- e) a binding molecule comprising a heavy chain CDR1 region of SEQ ID NO:139, a heavy chain CDR2 region of SEQ ID NO:134, and a heavy chain CDR3 region of SEQ ID NO:152,
- f) a binding molecule comprising a heavy chain CDR1 region of SEQ ID NO:139, a heavy chain CDR2 region of SEQ ID NO:151, and a heavy chain CDR3 region of SEQ ID NO:161,
- g) a binding molecule comprising a heavy chain CDR1 region of SEQ ID NO:139, a heavy chain CDR2 region of SEQ ID NO:151, and a heavy chain CDR3 region of SEQ ID NO:162, and

h) a binding molecule comprising a heavy chain CDR1 region of SEQ ID NO:139, a heavy chain CDR2 region of SEQ ID NO:134, and a heavy chain CDR3 region of SEQ ID NO:141.

In a preferred embodiment, the binding molecule comprises a heavy chain CDR1 region comprising the amino acid sequence of SEQ ID NO:139, a heavy chain CDR2 region comprising an amino acid sequence of SEQ ID NO:134, and a heavy chain CDR3 region comprising the amino acid sequence of SEQ ID NO:145 or SEQ ID NO: 152.

In another embodiment, the human binding molecules according to the invention are selected from the group consisting of:

- 10 a) a binding molecule having a heavy chain CDR1 region of SEQ ID NO:133, a heavy chain CDR2 region of SEQ ID NO:134, and a heavy chain CDR3 region of SEQ ID NO:135, a light chain CDR1 region having the amino acid sequence of SEQ ID NO:136, a light chain CDR2 region having the amino acid sequence of SEQ ID NO:137, and a light chain CDR3 region having the amino acid sequence of SEQ ID NO:138,
- 15 b) a binding molecule comprising a heavy chain CDR1 region of SEQ ID NO:139, a heavy chain CDR2 region of SEQ ID NO:140, and a heavy chain CDR3 region of SEQ ID NO:141, a light chain CDR1 region having the amino acid sequence of SEQ ID NO:142, a light chain CDR2 region having the amino acid sequence of SEQ ID NO:143, and a light chain CDR3 region having the amino acid sequence of SEQ ID NO:144,
- 20 c) a binding molecule comprising a heavy chain CDR1 region of SEQ ID NO:139, a heavy chain CDR2 region of SEQ ID NO:134, and a heavy chain CDR3 region of SEQ ID NO:145, a light chain CDR1 region having the amino acid sequence of SEQ ID NO:146, a light chain CDR2 region having the amino acid sequence of SEQ ID NO:174, and a light chain CDR3 region having the amino acid sequence of SEQ ID NO:147,
- 25 d) a binding molecule comprising a heavy chain CDR1 region of SEQ ID NO:139, a heavy chain CDR2 region of SEQ ID NO:134, and a heavy chain CDR3 region of SEQ ID NO:145, a light chain CDR1 region having the amino acid sequence of SEQ ID NO:148, a light chain CDR2 region having the amino acid sequence of SEQ ID NO:149, and a light chain CDR3 region having the amino acid sequence of SEQ ID NO:150,
- 30 e) a binding molecule comprising a heavy chain CDR1 region of SEQ ID NO:139, a heavy chain CDR2 region of SEQ ID NO:151, and a heavy chain CDR3 region of SEQ

ID NO:152, a light chain CDR1 region having the amino acid sequence of SEQ ID NO:153, a light chain CDR2 region having the amino acid sequence of SEQ ID NO:154, and a light chain CDR3 region having the amino acid sequence of SEQ ID NO:155,

f) a binding molecule comprising a heavy chain CDR1 region of SEQ ID NO:139, a heavy chain CDR2 region of SEQ ID NO:134, and a heavy chain CDR3 region of SEQ ID NO:152, a light chain CDR1 region having the amino acid sequence of SEQ ID NO:148, a light chain CDR2 region having the amino acid sequence of SEQ ID NO:149, and a light chain CDR3 region having the amino acid sequence of SEQ ID NO:150,

g) a binding molecule comprising a heavy chain CDR1 region of SEQ ID NO:139, a heavy chain CDR2 region of SEQ ID NO:134, and a heavy chain CDR3 region of SEQ ID NO:152, a light chain CDR1 region having the amino acid sequence of SEQ ID NO:156, a light chain CDR2 region having the amino acid sequence of SEQ ID NO:157, and a light chain CDR3 region having the amino acid sequence of SEQ ID NO:158,

h) a binding molecule comprising a heavy chain CDR1 region of SEQ ID NO:139, a heavy chain CDR2 region of SEQ ID NO:134, and a heavy chain CDR3 region of SEQ ID NO:152, a light chain CDR1 region having the amino acid sequence of SEQ ID NO:148, a light chain CDR2 region having the amino acid sequence of SEQ ID NO:159, and a light chain CDR3 region having the amino acid sequence of SEQ ID NO:160,

i) a binding molecule comprising a heavy chain CDR1 region of SEQ ID NO:139, a heavy chain CDR2 region of SEQ ID NO:151, and a heavy chain CDR3 region of SEQ ID NO:161, a light chain CDR1 region having the amino acid sequence of SEQ ID NO:142, a light chain CDR2 region having the amino acid sequence of SEQ ID NO:143, and a light chain CDR3 region having the amino acid sequence of SEQ ID NO:144,

j) a binding molecule comprising a heavy chain CDR1 region of SEQ ID NO:139, a heavy chain CDR2 region of SEQ ID NO:151, and a heavy chain CDR3 region of SEQ ID NO:162, a light chain CDR1 region having the amino acid sequence of SEQ ID NO:163, a light chain CDR2 region having the amino acid sequence of SEQ ID NO:164, and a light chain CDR3 region having the amino acid sequence of SEQ ID NO:165,

k) a binding molecule comprising a heavy chain CDR1 region of SEQ ID NO:139, a heavy chain CDR2 region of SEQ ID NO:134, and a heavy chain CDR3 region of SEQ ID NO:152, a light chain CDR1 region having the amino acid sequence of SEQ ID

NO:166, a light chain CDR2 region having the amino acid sequence of SEQ ID NO:167, and a light chain CDR3 region having the amino acid sequence of SEQ ID NO:168,

l) a binding molecule comprising a heavy chain CDR1 region of SEQ ID NO:139, a heavy chain CDR2 region of SEQ ID NO:134, and a heavy chain CDR3 region of SEQ ID NO:152, a light chain CDR1 region having the amino acid sequence of SEQ ID NO:169, a light chain CDR2 region having the amino acid sequence of SEQ ID NO:149, and a light chain CDR3 region having the amino acid sequence of SEQ ID NO:150,

m) a binding molecule comprising a heavy chain CDR1 region of SEQ ID NO:139, a heavy chain CDR2 region of SEQ ID NO:134, and a heavy chain CDR3 region of SEQ ID NO:141, a light chain CDR1 region having the amino acid sequence of SEQ ID NO:163, a light chain CDR2 region having the amino acid sequence of SEQ ID NO:169, and a light chain CDR3 region having the amino acid sequence of SEQ ID NO:170,

n) a binding molecule comprising a heavy chain CDR1 region of SEQ ID NO:139, a heavy chain CDR2 region of SEQ ID NO:134, and a heavy chain CDR3 region of SEQ ID NO:152, a light chain CDR1 region having the amino acid sequence of SEQ ID NO:171, a light chain CDR2 region having the amino acid sequence of SEQ ID NO:164, and a light chain CDR3 region having the amino acid sequence of SEQ ID NO:172,

o) a binding molecule comprising a heavy chain CDR1 region of SEQ ID NO:139, a heavy chain CDR2 region of SEQ ID NO:134, and a heavy chain CDR3 region of SEQ ID NO:145, a light chain CDR1 region having the amino acid sequence of SEQ ID NO:142, a light chain CDR2 region having the amino acid sequence of SEQ ID NO:143, and a light chain CDR3 region having the amino acid sequence of SEQ ID NO:173, and

p) a binding molecule comprising a heavy chain CDR1 region of SEQ ID NO:139, a heavy chain CDR2 region of SEQ ID NO:134, and a heavy chain CDR3 region of SEQ ID NO:152, a light chain CDR1 region having the amino acid sequence of SEQ ID NO:142, a light chain CDR2 region having the amino acid sequence of SEQ ID NO:143, and a light chain CDR3 region having the amino acid sequence of SEQ ID NO:144.

In another embodiment, the human binding molecules according to the invention are selected from the group consisting of:

a) a binding molecule comprising a heavy chain CDR1 region of SEQ ID NO:139, a heavy chain CDR2 region of SEQ ID NO:134, and a heavy chain CDR3 region of SEQ

ID NO:145, a light chain CDR1 region having the amino acid sequence of SEQ ID NO:146, a light chain CDR2 region having the amino acid sequence of SEQ ID NO:174, and a light chain CDR3 region having the amino acid sequence of SEQ ID NO:147,
b) a binding molecule comprising a heavy chain CDR1 region of SEQ ID NO:139, a
5 heavy chain CDR2 region of SEQ ID NO:134, and a heavy chain CDR3 region of SEQ ID NO:152, a light chain CDR1 region having the amino acid sequence of SEQ ID NO:171, a light chain CDR2 region having the amino acid sequence of SEQ ID NO:164, and a light chain CDR3 region having the amino acid sequence of SEQ ID NO:172,
c) a binding molecule comprising a heavy chain CDR1 region of SEQ ID NO:139, a
10 heavy chain CDR2 region of SEQ ID NO:134, and a heavy chain CDR3 region of SEQ ID NO:145, a light chain CDR1 region having the amino acid sequence of SEQ ID NO:142, a light chain CDR2 region having the amino acid sequence of SEQ ID NO:143, and a light chain CDR3 region having the amino acid sequence of SEQ ID NO:173, and
d) a binding molecule comprising a heavy chain CDR1 region of SEQ ID NO:139, a
15 heavy chain CDR2 region of SEQ ID NO:134, and a heavy chain CDR3 region of SEQ ID NO:152, a light chain CDR1 region having the amino acid sequence of SEQ ID NO:142, a light chain CDR2 region having the amino acid sequence of SEQ ID NO:143, and a light chain CDR3 region having the amino acid sequence of SEQ ID NO:144.

In another embodiment, the binding molecule according to the invention is
20 selected from the group consisting of

- a) a binding molecule comprising a heavy chain variable region of SEQ ID NO: 2,
- b) a binding molecule comprising a heavy chain variable region of SEQ ID NO: 6,
- c) a binding molecule comprising a heavy chain variable region of SEQ ID NO: 10,
- d) a binding molecule comprising a heavy chain variable region of SEQ ID NO: 14,
- 25 e) a binding molecule comprising a heavy chain variable region of SEQ ID NO: 18,
- f) a binding molecule comprising a heavy chain variable region of SEQ ID NO: 22,
- g) a binding molecule comprising a heavy chain variable region of SEQ ID NO: 26,
- h) a binding molecule comprising a heavy chain variable region of SEQ ID NO: 30,
- i) a binding molecule comprising a heavy chain variable region of SEQ ID NO: 34,
- 30 j) a binding molecule comprising a heavy chain variable region of SEQ ID NO: 38,
- k) a binding molecule comprising a heavy chain variable region of SEQ ID NO: 42,

- l) a binding molecule comprising a heavy chain variable region of SEQ ID NO: 46,
- m) a binding molecule comprising a heavy chain variable region of SEQ ID NO: 50,
- n) a binding molecule comprising a heavy chain variable region of SEQ ID NO: 54,
- o) a binding molecule comprising a heavy chain variable region of SEQ ID NO: 58,
- 5 and
- p) a binding molecule comprising a heavy chain variable region of SEQ ID NO: 62.

In an embodiment, the binding molecule according to the invention is selected from the group consisting of a binding molecule comprising a heavy chain variable region of SEQ ID NO: 10 a binding molecule comprising a heavy chain variable region of SEQ ID NO: 54, a binding molecule comprising a heavy chain variable region of SEQ ID NO: 58, and a binding molecule comprising a heavy chain variable region of SEQ ID NO: 62.

In a further embodiment, the binding molecules according to the invention comprise a light chain variable region comprising an amino acid sequence selected from the group consisting of SEQ ID NO:4, SEQ ID NO:8, SEQ ID NO:12, SEQ ID NO:16, 15 SEQ ID NO:20, SEQ ID NO:24, SEQ ID NO:28, SEQ ID NO:32, SEQ ID NO:36, SEQ ID NO:40, SEQ ID NO:44, SEQ ID NO:48, SEQ ID NO:52, SEQ ID NO:56, SEQ ID NO:60, and SEQ ID NO:64.

In yet another embodiment, the binding molecule is selected from the group consisting of

- 20 a) a binding molecule comprising a heavy chain variable region of SEQ ID NO: 2 and a light chain variable region of SEQ ID NO: 4,
- b) a binding molecule comprising a heavy chain variable region of SEQ ID NO: 6 and a light chain variable region of SEQ ID NO: 8,
- c) a binding molecule comprising a heavy chain variable region of SEQ ID NO: 10 25 and a light chain variable region of SEQ ID NO: 12,
- d) a binding molecule comprising a heavy chain variable region of SEQ ID NO: 14 and a light chain variable region of SEQ ID NO: 16,
- e) a binding molecule comprising a heavy chain variable region of SEQ ID NO: 18 and a light chain variable region of SEQ ID NO: 20,
- 30 f) a binding molecule comprising a heavy chain variable region of SEQ ID NO: 22 and a light chain variable region of SEQ ID NO: 24,

- g) a binding molecule comprising a heavy chain variable region of SEQ ID NO: 26 and a light chain variable region of SEQ ID NO: 28,
- h) a binding molecule comprising a heavy chain variable region of SEQ ID NO: 30 and a light chain variable region of SEQ ID NO: 32,
- 5 i) a binding molecule comprising a heavy chain variable region of SEQ ID NO: 34 and a light chain variable region of SEQ ID NO: 36,
- j) a binding molecule comprising a heavy chain variable region of SEQ ID NO: 38 and a light chain variable region of SEQ ID NO: 40,
- k) a binding molecule comprising a heavy chain variable region of SEQ ID NO: 42 and a light chain variable region of SEQ ID NO: 44,
- 10 l) a binding molecule comprising a heavy chain variable region of SEQ ID NO: 46 and a light chain variable region of SEQ ID NO: 48,
- m) a binding molecule comprising a heavy chain variable region of SEQ ID NO: 50 and a light chain variable region of SEQ ID NO: 52,
- 15 n) a binding molecule comprising a heavy chain variable region of SEQ ID NO: 54 and a light chain variable region of SEQ ID NO: 56,
- o) a binding molecule comprising a heavy chain variable region of SEQ ID NO: 58 and a light chain variable region of SEQ ID NO: 60, and
- p) a binding molecule comprising a heavy chain variable region of SEQ ID NO: 62 and a light chain variable region of SEQ ID NO: 64.
- 20

In an embodiment, the human binding molecules according to the invention are selected from the group consisting of: a binding molecule comprising a heavy chain variable region of SEQ ID NO: 10 and a light chain variable region of SEQ ID NO: 12, a binding molecule comprising a heavy chain variable region of SEQ ID NO: 54 and a light chain variable region of SEQ ID NO: 56, a binding molecule comprising a heavy chain variable region of SEQ ID NO: 58 and a light chain variable region of SEQ ID NO: 60, and a binding molecule comprising a heavy chain variable region of SEQ ID NO: 62 and a light chain variable region of SEQ ID NO: 64.

- 25

In a preferred embodiment, the binding molecules according to the invention are for a use as a medicament, and preferably for use in the diagnostic, therapeutic and/or prophylactic treatment of influenza infection caused by influenza A and/or B viruses.

- 30

Preferably, the influenza virus that causes the influenza infection and that can be treated using the binding molecules of the present invention is an influenza A virus of phylogenetic group 1 and/or 2, and/or a influenza B virus. The present invention also relates to a pharmaceutical composition comprising at least one binding molecule according to the invention, and a pharmaceutically acceptable excipient.

In yet another embodiment the invention relates to the use of a binding molecule according to the invention in the preparation of a medicament for the diagnosis, prophylaxis, and/or treatment of an influenza virus infection. Such infections can occur in small populations, but can also spread around the world in seasonal epidemics or, worse, in global pandemics where millions of individuals are at risk. The invention provides binding molecules that can neutralize the infection of influenza strains that cause such seasonal epidemics, as well as potential pandemics. Importantly, protection and treatment can be envisioned now with the binding molecules of the present invention in relation to various influenza subtypes as it has been disclosed that the binding molecules of the present invention are capable of cross-neutralizing various influenza subtypes of both phylogenetic group 1, encompassing H1, H2, H5, H6, H8, H9 and H11 subtypes and phylogenetic group 2, encompassing subtypes H3, H4, H7 and H10 subtypes, as well as influenza B subtypes.

Another aspect of the invention includes functional variants of the binding molecules as defined herein. Molecules are considered to be functional variants of a binding molecule according to the invention, if the variants are capable of competing for specifically binding to an influenza virus or a fragment thereof with the “parental” or “reference” binding molecules. In other words, molecules are considered to be functional variants of a binding molecule according to the invention when the functional variants are still capable of binding to the same or overlapping epitope of the influenza virus or a fragment thereof. For the sake of this application “parental” and “reference” will be used as synonyms meaning that the information of the reference or parental molecule, or the physical molecule itself may form the basis for the variation. Functional variants include, but are not limited to, derivatives that are substantially similar in primary structural sequence, including those that have modifications in the Fc receptor or other regions involved with effector functions, and/or which contain *e.g. in vitro* or *in vivo*

modifications, chemical and/or biochemical, that are not found in the parental binding molecule. Such modifications include *inter alia* acetylation, acylation, covalent attachment of a nucleotide or nucleotide derivative, covalent attachment of a lipid or lipid derivative, cross-linking, disulfide bond formation, glycosylation, hydroxylation, methylation, oxidation, pegylation, proteolytic processing, phosphorylation, and the like. Alternatively, functional variants can be binding molecules as defined in the present invention comprising an amino acid sequence containing substitutions, insertions, deletions or combinations thereof of one or more amino acids compared to the amino acid sequences of the parental binding molecules. Furthermore, functional variants can comprise truncations of the amino acid sequence at either or both the amino or carboxyl termini. Functional variants according to the invention may have the same or different, either higher or lower, binding affinities compared to the parental binding molecule but are still capable of binding to the influenza virus or a fragment thereof. For instance, functional variants according to the invention may have increased or decreased binding affinities for an influenza virus or a fragment thereof compared to the parental binding molecules. Preferably, the amino acid sequences of the variable regions, including, but not limited to, framework regions, hypervariable regions, in particular the CDR3 regions, are modified. Generally, the light chain and the heavy chain variable regions comprise three hypervariable regions, comprising three CDRs, and more conserved regions, the so-called framework regions (FRs). The hypervariable regions comprise amino acid residues from CDRs and amino acid residues from hypervariable loops. Functional variants intended to fall within the scope of the present invention have at least about 50% to about 99%, preferably at least about 60% to about 99%, more preferably at least about 70% to about 99%, even more preferably at least about 80% to about 99%, most preferably at least about 90% to about 99%, in particular at least about 95% to about 99%, and in particular at least about 97% to about 99% amino acid sequence identity and/or homology with the parental binding molecules as defined herein. Computer algorithms such as *inter alia* Gap or Bestfit known to a person skilled in the art can be used to optimally align amino acid sequences to be compared and to define similar or identical amino acid residues. Functional variants can be obtained by altering the parental binding molecules or parts thereof by general molecular biology methods known in the art including, but not

limited to, error-prone PCR, oligonucleotide-directed mutagenesis, site-directed mutagenesis and heavy and/or light chain shuffling. In an embodiment the functional variants of the invention have neutralizing activity against influenza A viruses of group 1 and group 2, and/or influenza B viruses. The neutralizing activity may either be identical, or be higher or lower compared to the parental binding molecules. Henceforth, when the term (human) binding molecule is used, this also encompasses functional variants of the (human) binding molecule. Assays for verifying if a variant binding molecule has neutralizing activity are well known in the art (see WHO Manual on Animal Influenza Diagnosis and Surveillance, Geneva: World Health Organisation, 2005 version 2002.5).

10 In yet a further aspect, the invention includes immunoconjugates, *i.e.* molecules comprising at least one binding molecule as defined herein and further comprising at least one tag, such as *inter alia* a detectable moiety/agent. Also contemplated in the present invention are mixtures of immunoconjugates according to the invention or mixtures of at least one immunoconjugates according to the invention and another molecule, such as a therapeutic agent or another binding molecule or immunoconjugate. In a further embodiment, the immunoconjugates of the invention may comprise more than one tag. These tags can be the same or distinct from each other and can be joined/conjugated non-covalently to the binding molecules. The tag(s) can also be joined/conjugated directly to the human binding molecules through covalent bonding. Alternatively, the tag(s) can be joined/conjugated to the binding molecules by means of one or more linking compounds. Techniques for conjugating tags to binding molecules are well known to the skilled artisan.

The tags of the immunoconjugates of the present invention may be therapeutic agents, but they can also be detectable moieties/agents. Tags suitable in therapy and/or prevention may be toxins or functional parts thereof, antibiotics, enzymes, other binding molecules that enhance phagocytosis or immune stimulation. Immunoconjugates comprising a detectable agent can be used diagnostically to, for example, assess if a subject has been infected with an influenza virus or to monitor the development or progression of an influenza virus infection as part of a clinical testing procedure to, *e.g.*, determine the efficacy of a given treatment regimen. However, they may also be used for other detection and/or analytical and/or diagnostic purposes. Detectable moieties/agents

include, but are not limited to, enzymes, prosthetic groups, fluorescent materials, luminescent materials, bioluminescent materials, radioactive materials, positron emitting metals, and non-radioactive paramagnetic metal ions. The tags used to label the binding molecules for detection and/or analytical and/or diagnostic purposes depend on the specific detection/analysis/diagnosis techniques and/or methods used such as *inter alia* immunohistochemical staining of (tissue) samples, flow cytometric detection, scanning laser cytometric detection, fluorescent immunoassays, enzyme-linked immunosorbent assays (ELISAs), radioimmunoassays (RIAs), bioassays (*e.g.*, phagocytosis assays), Western blotting applications, etc. Suitable labels for the detection/analysis/diagnosis techniques and/or methods known in the art are well within the reach of the skilled artisan.

Furthermore, the human binding molecules or immunoconjugates of the invention can also be attached to solid supports, which are particularly useful for *in vitro* immunoassays or purification of influenza viruses or fragments thereof. Such solid supports might be porous or nonporous, planar or non-planar. The binding molecules of the present invention can be fused to marker sequences, such as a peptide to facilitate purification. Examples include, but are not limited to, the hexa-histidine tag, the hemagglutinin (HA) tag, the myc tag or the flag tag. Alternatively, an antibody can be conjugated to a second antibody to form an antibody heteroconjugate. In another aspect the binding molecules of the invention may be conjugated/attached to one or more antigens. Preferably, these antigens are antigens which are recognized by the immune system of a subject to which the binding molecule-antigen conjugate is administered. The antigens may be identical, but may also differ from each other. Conjugation methods for attaching the antigens and binding molecules are well known in the art and include, but are not limited to, the use of cross-linking agents. The binding molecules of the invention will bind to influenza virus HA and the antigens attached to the binding molecules will initiate a powerful T-cell attack on the conjugate, which will eventually lead to the destruction of the influenza virus.

Next to producing immunoconjugates chemically by conjugating, directly or indirectly, via for instance a linker, the immunoconjugates can be produced as fusion proteins comprising the binding molecules of the invention and a suitable tag. Fusion

proteins can be produced by methods known in the art such as, *e.g.*, recombinantly by constructing nucleic acid molecules comprising nucleotide sequences encoding the binding molecules in frame with nucleotide sequences encoding the suitable tag(s) and then expressing the nucleic acid molecules.

5 It is another aspect of the present invention to provide a nucleic acid molecule encoding at least a binding molecule, functional variant or immunoconjugate according to the invention. Such nucleic acid molecules can be used as intermediates for cloning purposes, *e.g.* in the process of affinity maturation as described above. In a preferred embodiment, the nucleic acid molecules are isolated or purified.

10 The skilled man will appreciate that functional variants of these nucleic acid molecules are also intended to be a part of the present invention. Functional variants are nucleic acid sequences that can be directly translated, using the standard genetic code, to provide an amino acid sequence identical to that translated from the parental nucleic acid molecules.

15 Preferably, the nucleic acid molecules encode binding molecules comprising the CDR regions as described above. In a further embodiment the nucleic acid molecules encode binding molecules comprising two, three, four, five or even all six CDR regions of the binding molecules of the invention.

 In another embodiment, the nucleic acid molecules encode binding molecules
20 comprising a heavy chain comprising the variable heavy chain sequences as described above. In another embodiment the nucleic acid molecules encode binding molecules comprising a light chain comprising the variable light chain sequences as described above. The nucleotide sequences and the amino acid sequences of the heavy and light chain variable regions of the binding molecules of the invention are given below.

25 It is another aspect of the invention to provide vectors, *i.e.* nucleic acid constructs, comprising one or more nucleic acid molecules according to the present invention. Vectors can be derived from plasmids such as *inter alia* F, R1, RP1, Col, pBR322, TOL, Ti, etc; cosmids; phages such as lambda, lambdoid, M13, Mu, P1, P22, Q β , T-even, T-odd, T2, T4, T7, etc; plant viruses. Vectors can be used for cloning and/or for expression
30 of the binding molecules of the invention and might even be used for gene therapy purposes. Vectors comprising one or more nucleic acid molecules according to the

invention operably linked to one or more expression-regulating nucleic acid molecules are also covered by the present invention. The choice of the vector is dependent on the recombinant procedures followed and the host used. Introduction of vectors in host cells can be effected by *inter alia* calcium phosphate transfection, virus infection, DEAE-dextran mediated transfection, lipofectamin transfection or electroporation. Vectors may be autonomously replicating or may replicate together with the chromosome into which they have been integrated. Preferably, the vectors contain one or more selection markers. The choice of the markers may depend on the host cells of choice, although this is not critical to the invention as is well known to persons skilled in the art. They include, but are not limited to, kanamycin, neomycin, puromycin, hygromycin, zeocin, thymidine kinase gene from Herpes simplex virus (HSV-TK), dihydrofolate reductase gene from mouse (dhfr). Vectors comprising one or more nucleic acid molecules encoding the human binding molecules as described above operably linked to one or more nucleic acid molecules encoding proteins or peptides that can be used to isolate the human binding molecules are also covered by the invention. These proteins or peptides include, but are not limited to, glutathione-S-transferase, maltose binding protein, metal-binding polyhistidine, green fluorescent protein, luciferase and beta-galactosidase.

Hosts containing one or more copies of the vectors mentioned above are an additional subject of the present invention. Preferably, the hosts are host cells. Host cells include, but are not limited to, cells of mammalian, plant, insect, fungal or bacterial origin. Bacterial cells include, but are not limited to, cells from Gram-positive bacteria or Gram-negative bacteria such as several species of the genera *Escherichia*, such as *E. coli*, and *Pseudomonas*. In the group of fungal cells preferably yeast cells are used. Expression in yeast can be achieved by using yeast strains such as *inter alia Pichia pastoris*, *Saccharomyces cerevisiae* and *Hansenula polymorpha*. Furthermore, insect cells such as cells from *Drosophila* and Sf9 can be used as host cells. Besides that, the host cells can be plant cells such as *inter alia* cells from crop plants such as forestry plants, or cells from plants providing food and raw materials such as cereal plants, or medicinal plants, or cells from ornamentals, or cells from flower bulb crops. Transformed (transgenic) plants or plant cells are produced by known methods, for example, *Agrobacterium*-mediated gene transfer, transformation of leaf discs, protoplast transformation by polyethylene glycol-

induced DNA transfer, electroporation, sonication, microinjection or bolistic gene transfer. Additionally, a suitable expression system can be a baculovirus system. Expression systems using mammalian cells, such as Chinese Hamster Ovary (CHO) cells, COS cells, BHK cells, NSO cells or Bowes melanoma cells are preferred in the present invention. Mammalian cells provide expressed proteins with posttranslational modifications that are most similar to natural molecules of mammalian origin. Since the present invention deals with molecules that may have to be administered to humans, a completely human expression system would be particularly preferred. Therefore, even more preferably, the host cells are human cells. Examples of human cells are *inter alia* HeLa, 911, AT1080, A549, 293 and HEK293T cells. In preferred embodiments, the human producer cells comprise at least a functional part of a nucleic acid sequence encoding an adenovirus E1 region in expressible format. In even more preferred embodiments, said host cells are derived from a human retina and immortalized with nucleic acids comprising adenoviral E1 sequences, such as 911 cells or the cell line deposited at the European Collection of Cell Cultures (ECACC), CAMR, Salisbury, Wiltshire SP4 OJG, Great Britain on 29 February 1996 under number 96022940 and marketed under the trademark PER.C6[®] (PER.C6 is a registered trademark of Crucell Holland B.V.). For the purposes of this application “PER.C6 cells” refers to cells deposited under number 96022940 or ancestors, passages up-stream or downstream as well as descendants from ancestors of deposited cells, as well as derivatives of any of the foregoing. Production of recombinant proteins in host cells can be performed according to methods well known in the art. The use of the cells marketed under the trademark PER.C6[®] as a production platform for proteins of interest has been described in WO 00/63403 the disclosure of which is incorporated herein by reference in its entirety.

In yet another embodiment, binding molecules of the present invention can also be produced in transgenic, non-human, mammals such as *inter alia* rabbits, goats or cows, and secreted into for instance the milk thereof.

In yet another alternative embodiment, binding molecules according to the present invention may be generated by transgenic non-human mammals, such as for instance transgenic mice or rabbits that express human immunoglobulin genes. Preferably, the transgenic non-human mammals have a genome comprising a human heavy chain

transgene and a human light chain transgene encoding all or a portion of the human binding molecules as described above. The transgenic non-human mammals can be immunized with a purified or enriched preparation of influenza virus or a fragment thereof. Protocols for immunizing non-human mammals are well established in the art.

5 See Using Antibodies: A Laboratory Manual, Edited by: E. Harlow, D. Lane (1998), Cold Spring Harbor Laboratory, Cold Spring Harbor, New York and Current Protocols in Immunology, Edited by: J.E. Coligan, A.M. Kruisbeek, D.H. Margulies, E.M. Shevach, W. Strober (2001), John Wiley & Sons Inc., New York, the disclosures of which are incorporated herein by reference. Immunization protocols often include multiple
10 immunizations, either with or without adjuvants such as Freund's complete adjuvant and Freund's incomplete adjuvant, but may also include naked DNA immunizations. In another embodiment, the human binding molecules are produced by B-cells, plasma and/or memory cells derived from the transgenic animals. In yet another embodiment, the human binding molecules are produced by hybridomas, which are prepared by fusion of
15 B-cells obtained from the above-described transgenic non-human mammals to immortalized cells. B-cells, plasma cells and hybridomas as obtainable from the above-described transgenic non-human mammals and human binding molecules as obtainable from the above-described transgenic non-human mammals, B-cells, plasma and/or memory cells and hybridomas are also a part of the present invention.

20 In yet a further aspect, the invention provides compositions comprising at least a binding molecule, preferably a human monoclonal antibody, according to the invention, at least a functional variant thereof, at least an immunoconjugate according to the invention and/or a combination thereof. In addition to that, the compositions may comprise *inter alia* stabilizing molecules, such as albumin or polyethylene glycol, or
25 salts. Preferably, the salts used are salts that retain the desired biological activity of the binding molecules and do not impart any undesired toxicological effects. If necessary, the human binding molecules of the invention may be coated in or on a material to protect them from the action of acids or other natural or non-natural conditions that may inactivate the binding molecules.

30 In yet a further aspect, the invention provides compositions comprising at least a nucleic acid molecule as defined in the present invention. The compositions may

comprise aqueous solutions such as aqueous solutions containing salts (*e.g.*, NaCl or salts as described above), detergents (*e.g.*, SDS) and/or other suitable components.

Furthermore, the present invention pertains to pharmaceutical compositions comprising at least a binding molecule such as a human monoclonal antibody of the invention (or functional fragment or variant thereof), at least an immunoconjugate
5 according to the invention, at least a composition according to the invention, or combinations thereof. The pharmaceutical composition of the invention further comprises at least one pharmaceutically acceptable excipient. Pharmaceutically acceptable excipients are well known to the skilled person. The pharmaceutical composition
10 according to the invention may further comprise at least one other therapeutic agent. Suitable agents are also well known to the skilled artisan.

In a preferred embodiment the pharmaceutical composition according to the invention comprises at least one additional binding molecule, *i.e.* the pharmaceutical composition can be a cocktail or mixture of binding molecules. The pharmaceutical
15 composition may comprise at least two binding molecules according to the invention, or at least one binding molecule according to the invention and at least one further influenza virus binding and/or neutralizing molecule, such as another antibody directed against the HA protein or against other antigenic structures present on influenza viruses, such as M2. In another embodiment the additional binding molecule may be formulated for
20 simultaneous separate or sequential administration.

In an embodiment the pharmaceutical compositions may comprise two or more binding molecules that have neutralizing activity against influenza A viruses and/or influenza B viruses. In an embodiment, the binding molecules exhibit synergistic neutralizing activity, when used in combination. As used herein, the term "synergistic"
25 means that the combined effect of the binding molecules when used in combination is greater than their additive effects when used individually. The synergistically acting binding molecules may bind to different structures on the same or distinct fragments of influenza virus. A way of calculating synergy is by means of the combination index. The concept of the combination index (CI) has been described by Chou and Talalay (1984).
30 The compositions may *e.g.* comprise one binding molecule having neutralizing activity

and one non-neutralizing binding molecule. The non-neutralizing and neutralizing binding molecules may also act synergistically in neutralizing influenza virus.

In an embodiment, the pharmaceutical composition may comprise at least one binding molecule according to the invention and at least one further influenza virus neutralizing binding molecule. The binding molecules in the pharmaceutical composition preferably are capable of reacting with influenza viruses of different subtypes. The binding molecules should be of high affinity and should have a broad specificity. Preferably, both binding molecules are cross-neutralizing molecules in that they each neutralize influenza viruses of different subtypes. In addition, preferably they neutralize as many strains of each of the different influenza virus subtypes as possible.

A pharmaceutical composition according to the invention can further comprise at least one other therapeutic, prophylactic and/or diagnostic agent. Preferably, the pharmaceutical composition comprises at least one other prophylactic and/or therapeutic agent. Preferably, said further therapeutic and/or prophylactic agents are agents capable of preventing and/or treating an influenza virus infection and/or a condition resulting from such an infection. Therapeutic and/or prophylactic agents include, but are not limited to, anti-viral agents. Such agents can be binding molecules, small molecules, organic or inorganic compounds, enzymes, polynucleotide sequences, anti-viral peptides, etc. Other agents that are currently used to treat patients infected with influenza viruses are M2 inhibitors (*e.g.*, amantadine, rimantadine) and/or neuraminidase inhibitors (*e.g.*, zanamivir, oseltamivir). These can be used in combination with the binding molecules of the invention. "In combination" herein means simultaneously, as separate formulations, or as one single combined formulation, or according to a sequential administration regimen as separate formulations, in any order. Agents capable of preventing and/or treating an infection with influenza virus and/or a condition resulting from such an infection that are in the experimental phase might also be used as other therapeutic and/or prophylactic agents useful in the present invention.

The binding molecules or pharmaceutical compositions of the invention can be tested in suitable animal model systems prior to use in humans. Such animal model systems include, but are not limited to, mouse, ferret and monkey.

Typically, pharmaceutical compositions must be sterile and stable under the conditions of manufacture and storage. The binding molecules, immunoconjugates, nucleic acid molecules or compositions of the present invention can be in powder form for reconstitution in the appropriate pharmaceutically acceptable excipient before or at the time of delivery. In the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum drying and freeze-drying (lyophilization) that yield a powder of the active ingredient plus any additional desired ingredient from a previously sterile-filtered solution thereof.

Alternatively, the binding molecules, immunoconjugates, nucleic acid molecules or compositions of the present invention can be in solution and the appropriate pharmaceutically acceptable excipient can be added and/or mixed before or at the time of delivery to provide a unit dosage injectable form. Preferably, the pharmaceutically acceptable excipient used in the present invention is suitable to high drug concentration, can maintain proper fluidity and, if necessary, can delay absorption.

The choice of the optimal route of administration of the pharmaceutical compositions will be influenced by several factors including the physicochemical properties of the active molecules within the compositions, the urgency of the clinical situation and the relationship of the plasma concentrations of the active molecules to the desired therapeutic effect. For instance, if necessary, the binding molecules of the invention can be prepared with carriers that will protect them against rapid release, such as a controlled release formulation, including implants, transdermal patches, and microencapsulated delivery systems. Biodegradable, biocompatible polymers can *inter alia* be used, such as ethylene vinyl acetate, polyanhydrides, polyglycolic acid, collagen, polyorthoesters, and polylactic acid. Furthermore, it may be necessary to coat the binding molecules with, or co-administer the binding molecules with, a material or compound that prevents the inactivation of the human binding molecules. For example, the binding molecules may be administered to a subject in an appropriate carrier, for example, liposomes or a diluent.

The routes of administration can be divided into two main categories, oral and parenteral administration. The preferred administration route is intravenous or by inhalation.

Oral dosage forms can be formulated *inter alia* as tablets, troches, lozenges, aqueous or oily suspensions, dispersible powders or granules, emulsions, hard capsules, soft gelatin capsules, syrups or elixirs, pills, dragees, liquids, gels, or slurries. These formulations can contain pharmaceutically excipients including, but not limited to, inert
5 diluents, granulating and disintegrating agents, binding agents, lubricating agents, preservatives, colouring, flavouring or sweetening agents, vegetable or mineral oils, wetting agents, and thickening agents.

The pharmaceutical compositions of the present invention can also be formulated for parenteral administration. Formulations for parenteral administration can be *inter alia*
10 in the form of aqueous or non-aqueous isotonic sterile non-toxic injection or infusion solutions or suspensions. The solutions or suspensions may comprise agents that are non-toxic to recipients at the dosages and concentrations employed such as 1,3-butanediol, Ringer's solution, Hank's solution, isotonic sodium chloride solution, oils, fatty acids, local anaesthetic agents, preservatives, buffers, viscosity or solubility increasing agents,
15 water-soluble antioxidants, oil-soluble antioxidants and metal chelating agents.

In a further aspect, the binding molecules such as human monoclonal antibodies (functional fragments and variants thereof), immunoconjugates, compositions, or pharmaceutical compositions of the invention can be used as a medicament. So, a method of diagnosis, treatment and/or prevention of an influenza virus infection using the binding
20 molecules, immunoconjugates, compositions, or pharmaceutical compositions of the invention is another part of the present invention. The above-mentioned molecules can *inter alia* be used in the diagnosis, prophylaxis, treatment, or combination thereof, of an influenza virus infection caused influenza viruses comprising HA of the H1, H2, H3, H4, H5, H6, H7, H8, H9, H10 and/or H11 subtype. In an embodiment, the above-mentioned
25 molecules can also be used in the diagnosis, prophylaxis, treatment or combination thereof of an influenza virus infection caused by an influenza B virus. They are suitable for treatment of yet untreated patients suffering from an influenza virus infection and patients who have been or are treated for an influenza virus infection.

The above-mentioned molecules or compositions may be employed in
30 conjunction with other molecules useful in diagnosis, prophylaxis and/or treatment. They can be used *in vitro*, *ex vivo* or *in vivo*. For instance, the binding molecules such as

human monoclonal antibodies (or functional variants thereof), immunoconjugates, compositions or pharmaceutical compositions of the invention can be co-administered with a vaccine against influenza virus (if available). Alternatively, the vaccine may also be administered before or after administration of the molecules of the invention. Instead
5 of a vaccine, anti-viral agents can also be employed in conjunction with the binding molecules of the present invention. Suitable anti-viral agents are mentioned above.

The molecules are typically formulated in the compositions and pharmaceutical compositions of the invention in a therapeutically or diagnostically effective amount. Alternatively, they may be formulated and administered separately. For instance the other
10 molecules such as the anti-viral agents may be applied systemically, while the binding molecules of the invention may be applied intravenously.

Treatment may be targeted at patient groups that are susceptible to influenza infection. Such patient groups include, but are not limited to e.g., the elderly (e.g. ≥ 50 years old, ≥ 60 years old, and preferably ≥ 65 years old), the young (e.g. ≤ 5 years old, \leq
15 1 year old), hospitalized patients and already infected patients who have been treated with an antiviral compound but have shown an inadequate antiviral response.

Dosage regimens can be adjusted to provide the optimum desired response (e.g., a therapeutic response). A suitable dosage range may for instance be 0.01-100 mg/kg body weight, preferably 0.1-50 mg/kg body weight, preferably 0.01-15 mg/kg body
20 weight. Furthermore, for example, a single bolus may be administered, several divided doses may be administered over time or the dose may be proportionally reduced or increased as indicated by the exigencies of the therapeutic situation. The molecules and compositions according to the present invention are preferably sterile. Methods to render these molecules and compositions sterile are well known in the art. The other molecules
25 useful in diagnosis, prophylaxis and/or treatment can be administered in a similar dosage regimen as proposed for the binding molecules of the invention. If the other molecules are administered separately, they may be administered to a patient prior to (e.g., 2 min, 5 min, 10 min, 15 min, 30 min, 45 min, 60 min, 2 hrs, 4 hrs, 6 hrs, 8 hrs, 10 hrs, 12 hrs, 14 hrs, 16 hrs, 18 hrs, 20 hrs, 22 hrs, 24 hrs, 2 days, 3 days, 4 days, 5 days, 7 days, 2 weeks,
30 4 weeks or 6 weeks before), concomitantly with, or subsequent to (e.g., 2 min, 5 min, 10 min, 15 min, 30 min, 45 min, 60 min, 2 hrs, 4 hrs, 6 hrs, 8 hrs, 10 hrs, 12 hrs, 14 hrs, 16

hrs, 18 hrs, 20 hrs, 22 hrs, 24 hrs, 2 days, 3 days, 4 days, 5 days, 7 days, 2 weeks, 4 weeks or 6 weeks after) the administration of one or more of the human binding molecules or pharmaceutical compositions of the invention. The exact dosing regimen is usually sorted out during clinical trials in human patients.

5 Human binding molecules and pharmaceutical compositions comprising the human binding molecules are particularly useful, and often preferred, when to be administered to human beings as *in vivo* therapeutic agents, since recipient immune response to the administered antibody will often be substantially less than that occasioned by administration of a monoclonal murine, chimeric or humanized binding molecule.

10 In another aspect, the invention concerns the use of the binding molecules such as neutralizing human monoclonal antibodies (functional fragments and variants thereof), immunoconjugates, nucleic acid molecules, compositions or pharmaceutical compositions according to the invention in the preparation of a medicament for the diagnosis, prophylaxis, treatment, or combination thereof, of an influenza virus infection, 15 in particular an influenza virus infection caused influenza viruses comprising HA of the H1, H2, H3, H4, H5, H6, H7, H8, H9, H10, and/or H11 subtype and/or influenza B viruses.

Next to that, kits comprising at least a binding molecule such as a neutralizing human monoclonal antibody (functional fragments and variants thereof), at least an 20 immunoconjugate, at least a nucleic acid molecule, at least a composition, at least a pharmaceutical composition, at least a vector, at least a host according to the invention or a combination thereof are also a part of the present invention. Optionally, the above-described components of the kits of the invention are packed in suitable containers and labelled for diagnosis, prophylaxis and/or treatment of the indicated conditions. The 25 above-mentioned components may be stored in unit or multi-dose containers as an aqueous, preferably sterile, solution or as a lyophilised, preferably sterile, formulation for reconstitution. The containers may be formed from a variety of materials such as glass or plastic and may have a sterile access port (for example, the container may be an intravenous solution bag or a vial having a stopper pierceable by a hypodermic injection 30 needle). The kit may further comprise more containers comprising a pharmaceutically acceptable buffer. It may further include other materials desirable from a commercial and

user standpoint, including other buffers, diluents, filters, needles, syringes, culture medium for one or more of the suitable hosts and, possibly, even at least one other therapeutic, prophylactic or diagnostic agent. Associated with the kits can be instructions customarily included in commercial packages of therapeutic, prophylactic or diagnostic products, that contain information about for example the indications, usage, dosage, manufacture, administration, contra-indications and/or warnings concerning the use of such therapeutic, prophylactic or diagnostic products.

The binding molecules according to the present invention can also be advantageously used as a diagnostic agent in an *in vitro* method for the detection of influenza virus. The invention thus further pertains to a method of detecting influenza virus phylogenetic group 1 or group 2, or influenza B subtype influenza virus in a sample, wherein the method comprises the steps of (a) contacting a sample with a diagnostically effective amount of a binding molecule (functional fragments and variants thereof) or an immunoconjugate according to the invention, and (b) determining whether the binding molecule or immunoconjugate specifically binds to a molecule of the sample. The sample may be a biological sample including, but not limited to blood, serum, stool, sputum, nasopharyngeal aspirates, bronchial lavages, urine, tissue or other biological material from (potentially) infected subjects, or a non-biological sample such as water, drink, etc. The (potentially) infected subjects may be human subjects, but also animals that are suspected as carriers of influenza virus might be tested for the presence of the virus using the human binding molecules or immunoconjugates of the invention. The sample may first be manipulated to make it more suitable for the method of detection. Manipulation means *inter alia* treating the sample suspected to contain and/or containing the virus in such a way that the virus will disintegrate into antigenic components such as proteins, (poly)peptides or other antigenic fragments. Preferably, the human binding molecules or immunoconjugates of the invention are contacted with the sample under conditions which allow the formation of an immunological complex between the human binding molecules and the virus or antigenic components thereof that may be present in the sample. The formation of an immunological complex, if any, indicating the presence of the virus in the sample, is then detected and measured by suitable means. Such methods include, *inter alia*, homogeneous and heterogeneous binding immunoassays, such as radio-

immunoassays (RIA), ELISA, immunofluorescence, immunohistochemistry, FACS, BIACORE and Western blot analyses.

Preferred assay techniques, especially for large-scale clinical screening of patient sera and blood and blood-derived products are ELISA and Western blot techniques.

5 ELISA tests are particularly preferred. For use as reagents in these assays, the binding molecules or immunoconjugates of the invention are conveniently bonded to the inside surface of microtiter wells. The binding molecules or immunoconjugates of the invention may be directly bonded to the microtiter well. However, maximum binding of the binding molecules or immunoconjugates of the invention to the wells might be accomplished by
10 pre-treating the wells with polylysine prior to the addition of the binding molecules or immunoconjugates of the invention. Furthermore, the binding molecules or immunoconjugates of the invention may be covalently attached by known means to the wells. Generally, the binding molecules or immunoconjugates are used in a concentration between 0.01 to 100 µg/ml for coating, although higher as well as lower amounts may
15 also be used. Samples are then added to the wells coated with the binding molecules or immunoconjugates of the invention.

Furthermore, binding molecules of the invention can be used to identify specific binding structures of influenza virus. The binding structures can be epitopes on proteins and/or polypeptides. They can be linear, but also structural and/or conformational. In one
20 embodiment, the binding structures can be analysed by means of PEPSCAN analysis (see *inter alia* WO 84/03564, WO 93/09872, Sloodstra *et al.*, 1996). Alternatively, a random peptide library comprising peptides from a protein of influenza virus can be screened for peptides capable of binding to the binding molecules of the invention.

The invention is further illustrated in the following examples and figures. The
25 examples are not intended to limit the scope of the invention in any way.

EXAMPLES

Example 1 *Construction of scFv phage display libraries using RNA extracted from*
30 *peripheral blood mononuclear cells*

Peripheral blood was collected from normal healthy donors by venapuncture in EDTA anti-coagulation sample tubes. scFv phage display libraries were obtained as described in WO 2008/028946, which is incorporated by reference herein. RNA was isolated from peripheral blood mononuclear cells and cDNA prepared. A two round PCR
5 amplification approach was applied using the primer sets shown in Tables 1 and 2 to isolate the immunoglobulin VH and VL regions from the respective donor repertoire.

First round amplification on the respective cDNA using the primer sets mentioned in Table 1 yielded 7, 6 and 9 products of about 650 base pairs for respectively VH, Vkappa and Vlamba regions. For IgM VH region amplification the OCM constant
10 primer was used in combination with OH1 to OH7. The thermal cycling program for first round amplifications was: 2 min 96°C (denaturation step), 30 cycles of 30 sec 96°C/ 30 sec 60°C/ 60 sec 72°C, 10 min 72°C final elongation and 6°C refrigeration. The products were loaded on and isolated from a 1% agarose gel using gel-extraction columns (Macherey Nagel) and eluted in 50 µl 5 mM Tris-HCl pH 8.0. Ten percent of first round
15 products (3 to 5 µl) was subjected to second round amplification using the primers mentioned in Table 2. These primers were extended with restriction sites enabling the directional cloning of the respective VL and VH regions into phage display vector PDV-C06. The PCR program for second round amplifications was as follows: 2 min 96°C (denaturation step), 30 cycles of 30 sec 96°C/ 30 sec 60°C/ 60 sec 72°C, 10 min 72°C
20 final elongation and 6°C refrigeration. The second round products (~350 base pairs) were first pooled according to natural occurrence of J segments found in immunoglobulin gene products, resulting in 7, 6 and 9 pools for respectively the VH, Vkappa and Vlamba variable regions (see Tables 3 and 4). To obtain a normalized distribution of immunoglobulin sequences in the immune library the 6 Vkappa and 9 Vlamba light
25 chain pools were mixed according to the percentages mentioned in Table 3. This single final VL pool (3 µg) was digested overnight with Sall and NotI restriction enzymes, loaded on and isolated from a 1% agarose gel (~350 base pairs) using Macherey Nagel gel-extraction columns and ligated in Sall-NotI cut PDV-C06 vector (~5000 base pairs) as follows: 10 µl PDV-C06 vector (50 ng/µl), 7 µl VL insert (10 ng/µl), 5 µl 10X ligation
30 buffer (NEB), 2.5 T4 DNA Ligase (400 U/µl) (NEB), 25.5 µl ultrapure water (vector to insert ratio was 1:2). Ligation was performed overnight in a water bath of 16°C. Next, the

volume was doubled with water, extracted with an equal volume of phenol-chloroform-isoamylalcohol (75:24:1) (Invitrogen) followed by chloroform (Merck) extraction and precipitated with 1 μ l Pellet Paint (Novogen), 10 μ l sodium acetate (3 M pH 5.0) and 100 μ l isopropanol for 2 hrs at -20°C. The obtained sample was subsequently centrifuged at 20.000xg for 30 min at 4°C. The obtained precipitate was washed with 70% ethanol and centrifuged for 10 min at 20.000xg at room temperature. Ethanol was removed by vacuum aspiration and the pellet was air dried for several min and then dissolved in 50 μ l buffer containing 10 mM Tris-HCl, pH 8.0. 2 μ l ligation mixture was used for the transformation of 40 μ l TG-1 electro-competent cells (Agilent) in a chilled 0.1 cm electroporation cuvette (Biorad) using a Genepulser II apparatus (Biorad) set at 1.7 kV, 200 Ohm, 25 μ F (time constant ~4,5 msec). Directly after pulse, the bacteria were flushed from the cuvette with 1000 μ l SOC medium (Invitrogen) containing 5% (w/v) glucose (Sigma) at 37°C and transferred to a 15 ml round bottom culture tube. Another 500 μ l SOC/glucose was used to flush residual bacteria from the cuvette and was added to the culture tube. Bacteria were recovered by culturing for exactly one hr at 37°C in a shaker incubator at 220 rpm. The transformed bacteria were plated over large 240 mm square petridishes (NUNC) containing 150 ml 2TY agar (16 g/l bacto-tryptone, 10 g/l bacto-yeast extract, 5 g/l NaCl, 15 g/l agar, pH 7.0) supplemented with 50 μ g/ml ampicillin and 5% (w/v) glucose (Sigma). A 1 to 1000 dilution was plated for counting purposes on 15 cm petridishes containing the same medium. This transformation procedure was repeated sequentially ten times and the complete each transformation was plated on a separate square petridish and grown overnight in a 37°C culture stove. Typically, around 1×10^7 cfu (1×10^6 per petridish) were obtained using the above protocol. The intermediate VL light chain library was harvested from the plates by mildly scraping the bacteria into 10 ml 2TY medium per plate. The cell mass was determined by OD600 measurement and two times 500 OD of bacteria was used for maxi plasmid DNA preparation using two P500 maxiprep columns (Macherey Nagel) according to manufacturer's instructions.

Analogous to the VL variable regions, the second round VH-JH products were first mixed together to obtain the normal J segment usage distribution (see Table 4), resulting in 7 VH subpools called PH1 to PH7. The pools were mixed to acquire a normalized sequence distribution using the percentages depicted in Table 4, obtaining

one VH fraction that was digested with SfiI and XhoI restriction enzymes and ligated in SfiI-XhoI cut PDV-VL intermediate library obtained as described above. The ligation set-up, purification method, subsequent transformation of TG1 and harvest of bacteria was essentially as described for the VL intermediate library (see above) with the exception that 20 transformations and 20 square petridishes were used. The final library (approximately 1×10^7 cfu) was checked for insert frequency with a colony PCR using a primer set flanking the inserted VH-VL regions. 90% of the colonies showed a correct length insert. The colony PCR products were used for subsequent DNA sequence analysis to check sequence variation and to assess the percentage of colonies showing a complete ORF. This was 76%. Finally, the library was rescued and amplified by using CT helper phages (see WO 02/103012) and was used for phage antibody selection by panning methods as described below.

Example 2

Selection of phages carrying single chain Fv fragments against Influenza A and Influenza B hemagglutinin

Antibody fragments were selected using antibody phage display libraries constructed essentially as described above and general phage display technology and MABSTRACT[®] technology essentially as described in US Patent Number 6,265,150 and in WO 98/15833 (both of which are incorporated by reference herein). Furthermore, the methods and helper phages as described in WO 02/103012 (which is incorporated by reference herein) were used in the present invention.

Selection was performed against recombinant hemagglutinin (HA) of influenza A subtype H1 (A/New Caledonia/20/99), H3 (A/Wisconsin/67/2005), H4 (A/Duck/Hong Kong/24/1976), H5 (A/Chicken/Vietnam/28/2003), H7 (A/Netherlands/219/2003) and H9 (A/HongKong/1073/99). HA antigens were diluted in PBS (5.0 $\mu\text{g/ml}$), added to MaxiSorp[™] Nunc-Immuno Tubes (Nunc) and incubated overnight at 4°C on a rotating wheel. The immunotubes were emptied and washed three times in block buffer (2% non-fat dry milk (ELK) in PBS). Subsequently, the immunotubes were filled completely with block buffer and incubated for 1-2 hrs at room temperature. Aliquots of phage display library (500-1000 μl , 0.5×10^{13} – 1×10^{13} cfu, amplified using CT helper phage (see WO

02/103012)) were blocked in blocking buffer supplemented with 10% non-heat inactivated fetal bovine serum and 2% mouse serum for 1-2 hrs at room temperature. The blocked phage library was added to the immunotubes, incubated for 2 hrs at room temperature, and washed with wash buffer (0.05% (v/v) Tween-20 in PBS) to remove unbound phages. Bound phages were eluted from the respective antigen by incubation with 1 ml of 100 mM triethylamine (TEA) for 10 min at room temperature. Subsequently, the eluted phages were mixed with 0.5 ml of 1 M Tris-HCl pH 7.5 to neutralize the pH. This mixture was used to infect 5 ml of an XL1-Blue *E.coli* culture that had been grown at 37°C to an OD 600 nm of approximately 0.3. The phages were allowed to infect the XL1-Blue bacteria for 30 min at 37°C. Then, the mixture was centrifuged for 10 min at 3000xg at room temperature and the bacterial pellet was resuspended in 0.5 ml 2-trypton yeast extract (2TY) medium. The obtained bacterial suspension was divided over two 2TY agar plates supplemented with tetracycline, ampicillin and glucose. After incubation overnight of the plates at 37°C, the colonies were scraped from the plates and used to prepare an enriched phage library, essentially as described by De Kruif *et al.* (1995) and WO 02/103012. Briefly, scraped bacteria were used to inoculate 2TY medium containing ampicillin, tetracycline and glucose and grown at a temperature of 37°C to an OD 600 nm of ~0.3. CT helper phages were added and allowed to infect the bacteria after which the medium was changed to 2TY containing ampicillin, tetracycline and kanamycin. Incubation was continued overnight at 30°C. The next day, the bacteria were removed from the 2TY medium by centrifugation after which the phages in the medium were precipitated using polyethylene glycol (PEG) 6000/NaCl. Finally, the phages were dissolved in 2 ml of PBS with 1% bovine serum albumin (BSA), filter-sterilized and used for the next round of selection. The second round of selection is performed either on the same HA subtype and/or on HA of a different subtype.

Two consecutive rounds of selections were performed before isolation of individual single-chain phage antibodies. After the second round of selection, individual *E.coli* colonies were used to prepare monoclonal phage antibodies. Essentially, individual colonies were grown to log-phase in 96 well plate format and infected with VCS-M13 helper phages after which phage antibody production was allowed to proceed overnight. Phagemids were sequence analysed and all unique phagemids were used for further

analysis. The supernatants containing phage antibodies were used directly in ELISA for binding to HA antigens. Alternatively, phage antibodies were PEG/NaCl-precipitated and filter-sterilized for both elisa and flow cytometry analysis.

Example 3

5 *Validation of the HA specific single-chain phage antibodies*

Selected supernatants containing single-chain phage antibodies that were obtained in the screenings described above were validated in ELISA for specificity, *i.e.* binding to different HA antigens. For this purpose, baculovirus expressed recombinant H1 (A/New Caledonia/20/99), H3 (A/Wisconsin/67/2005), H5 (A/Vietnam/1203/04) H7
10 (A/Netherlands/219/2003), and B (B/Ohio/01/2005) HAs (Protein Sciences, CT, USA) were coated to Maxisorp™ ELISA plates. After coating, the plates were washed three times with PBS containing 0.1% v/v Tween-20 and blocked in PBS containing 3% BSA or 2% ELK for 1 hr at room temperature. The selected single-chain phage antibodies were incubated for 1 hr in an equal volume of PBS containing 4% ELK to obtain blocked
15 phage antibodies. The plates were emptied, washed three times with PBS/0.1% Tween-20 and the blocked single-chain phage antibodies were added to the wells. Incubation was allowed to proceed for one hr, the plates were washed with PBS/0.1% Tween-20 and bound phage antibodies were detected (using OD 492nm measurement) using an anti-M13 antibody conjugated to peroxidase. As a control, the procedure was performed
20 simultaneously without single-chain phage antibody and with an unrelated negative control single-chain phage antibody. From the selections on the different HA antigens with the phage libraries, 13 unique single-chain phage antibodies specifically binding recombinant influenza A H1, H3, H5, H7 and influenza B HA were obtained (SC09-003, SC09-004, SC09-005, SC09-006, SC09-007, SC09-008, SC09-009, SC09-010, SC09-
25 011, SC09-030, SC09-112, SC09-113 and SC09-114). See Table 5.

Alternatively, PEG/NaCl-precipitated and filter-sterilized phage antibodies were used to validate binding and specificity by FACS analysis. For this purpose, full-length recombinant influenza A subtypes H1 (A/New Caledonia/20/1999), H3
(A/Wisconsin/67/2005) and H7 (A/Netherlands/219/2003) HAs were expressed on the
30 surface of PER.C6 cells. The cells were incubated with single-chain phage antibodies for

1 hr followed by three wash steps with PBS+0.1%BSA. Bound phages were detected using FITC conjugated M13-antibody. From the selections on the different HA antigens with the phage libraries, 14 single-chain phage antibodies specifically binding influenza A subtypes H1, H3 and H7 HA were found (SC09-003, SC09-004, SC09-005, SC09-006, SC09-007, SC09-008, SC09-009, SC09-010, SC09-011, SC09-012, SC09-030, SC09-112, SC09-113 and SC09-114). See table 6.

All 16 phage antibodies, SC09-003, SC09-004, SC09-005, SC09-006, SC09-007, SC09-008, SC09-009, SC09-010, SC09-011, SC09-012, SC09-029, SC09-030, SC09-031, SC09-112, SC09-113 and SC09-114, were used for construction of fully human immunoglobulins.

Example 4

Construction of fully human immunoglobulin molecules (human monoclonal antibodies) from the selected single chain Fvs

From the selected specific single-chain phage antibodies (scFv) clones plasmid DNA was obtained and nucleotide and amino acid sequences were determined according to standard techniques. Heavy and light chain variable regions of the scFvs were cloned directly by restriction digest for expression in the IgG expression vectors pIg-C911-HCgamma1 (see SEQ ID NO: 175), pIG-C909-Ckappa (see SEQ ID NO: 176), or pIg-C910-Clambda (see SEQ ID No: 177). The VH and VL gene identity (see Tomlinson IM et al. V-BASE Sequence Directory. Cambridge United Kingdom: MRC Centre for Protein Engineering (1997)) of the scFvs were determined (see Table 7).

Nucleotide sequences for all constructs were verified according to standard techniques known to the skilled artisan. The resulting expression constructs encoding the human IgG1 heavy and light chains were transiently expressed in combination in 293T cells and supernatants containing human IgG1 antibodies were obtained and produced using standard purification procedures.

The amino acid sequence of the CDRs of the heavy and light chains of the selected immunoglobulin molecules is given in Table 7.

The number of amino-acid differences and the % identity of all heavy and light chain variable domains is given in Table 8.

Example 5

Cross-binding reactivity of IgGs

5 A panel of five of the IgG antibodies described above, CR9005, CR9030, CR9112, CR9113 and CR9114, was validated in ELISA for binding specificity, *i.e.* binding to different HA antigens. For this purpose, baculovirus expressed recombinant H1 (A/New Caledonia/20/1999), H3 (A/Wisconsin/67/2005), H5 (A/Vietnam/1203/04, H7 (A/Netherlands/219/2003) and H9 (A/HongKong/1073/99) HA's (Protein Sciences, CT, USA) were coated to Maxisorp™ ELISA plates. After coating, the plates were washed three times with PBS containing 0.1% v/v Tween-20 and blocked in PBS containing 3% BSA or 2% ELK for 1 hr at room temperature. The plates were emptied, washed three times with PBS/0.1% Tween-20 and the IgG antibodies were added to the wells. Incubation was allowed to proceed for one hr, the plates were washed with
10 PBS/0.1% Tween-20 and bound antibodies were detected (using OD 492nm measurement) using an anti-human IgG antibody conjugated to peroxidase. As a control, an unrelated IgG CR4098 was used.

CR9005, CR9030, CR9112, CR9113 and CR9114 were shown to have heterosubtypic cross-binding activity to all the recombinant HAs tested. See table 9.

20 Additionally, the selected antibodies were used to test heterosubtypic binding by FACS analysis. For this purpose, full-length recombinant influenza A subtypes H1 (A/New Caledonia/20/1999), H3 (A/Wisconsin/67/2005) and H7 (A/Netherlands/219/2003) HAs were expressed on the surface of PER.C6 cells. The cells were incubated with IgG antibodies for 1 hr followed by three wash steps with
25 PBS+0.1%BSA. Bound antibodies were detected using PE conjugated anti-human antibody. As a control, untransfected PER.C6 cells were used. CR9005, CR9030, CR9112, CR9113 and CR9114 show cross-binding activity to influenza A subtypes H1, H3 and H7 HA but not wild-type PER.C6 cells. See table 9.

Example 6

Cross-neutralizing activity of IgGs

In order to determine whether the selected IgGs were capable of blocking multiple influenza A strains, additional *in vitro* virus neutralization assays (VNA) were performed. The VNA were performed on MDCK cells (ATCC CCL-34). MDCK cells were cultured
5 in MDCK cell culture medium (MEM medium supplemented with antibiotics, 20 mM HEPES and 0.15% (w/v) sodium bicarbonate (complete MEM medium), supplemented with 10% (v/v) fetal bovine serum). The H1 (A/WSN/33, A/New Caledonia/20/1999, A/Solomon Islands/IVR-145 (high-growth reassortant of A/Solomon Islands/3/2006), A/Brisbane/59/2007, A/NYMC/X-181 (high-growth reassortant of
10 A/California/07/2009), H2 (A/Env/MPU3156/05), H3 (A/Hong Kong/1/68, A/Johannesburg/33/94, A/Panama/2000/1999, A/Hiroshima/52/2005, A/Wisconsin/67/2005 and A/Brisbane/10/2007), H4 (A/WF/HK/MPA892/06), H5 (PR8-H5N1-HK97 (6:2 reassortant of A/Hong Kong/156/97 and A/PR/8/34) and A/Eurasian Wigeon/MPF461/07), H6 (A/Eurasian Wigeon/MPD411/07), H7 (NIBRG-60 (6:2
15 reassortant of A/Mallard/Netherlands/12/2000) and PR8-H7N7-NY (7:1 reassortant of A/New York/107/2003 (H7N7) and A/PR/8/34)), H8 (A/Eurasian Wigeon/MPH571/08) H9 (A/Hong Kong/1073/99 and A/Chick/HK/SSP176/09), H10 (A/Chick/Germany/N/49) and H14 (PR8-H14N5 (6:2 reassortant of A/mallard/Astrakhan/263/1982 (H14N5) and A/PR/8/34)) strains which were used in the assay were all diluted to a titer of $5,7 \times 10^3$
20 TCID₅₀/ml (50% tissue culture infective dose per ml), with the titer calculated according to the method of Spearman and Karber. The IgG preparations (200 µg/ml) were serially 2-fold diluted (1:2 - 1:512) in complete MEM medium in quadruplicate wells. 25 µl of the respective IgG dilution was mixed with 25 µl of virus suspension (100 TCID₅₀/25 µl) and incubated for one hr at 37°C. The suspension was then transferred in quadruplicate
25 onto 96-well plates containing confluent MDCK cultures in 50 µl complete MEM medium. Prior to use, MDCK cells were seeded at 3×10^4 cells per well in MDCK cell culture medium, grown until cells had reached confluence, washed with 300-350 µl PBS, pH 7.4 and finally 50 µl complete MEM medium was added to each well. The inoculated cells were cultured for 3-4 days at 37°C and observed daily for the development of
30 cytopathogenic effect (CPE). CPE was compared to the positive control.

CR9005, CR9112, CR9113 and CR9114 show heterosubtypic cross-neutralizing activity to representative strains of all tested influenza A subtypes H1, H2, H3, H4, H5, H6, H7, H8, H9 and H10 viruses. See table 10.

5 Example 7

Pan-influenza antibodies bind to the pre-fusion conformation of HA

In order to determine whether the selected IgGs were capable of binding the pre- or post-fusion conformation of the HA molecule, an in vitro pH-shift experiment was performed. For this purpose, full-length recombinant influenza A subtypes H1 (A/New
10 Caledonia/20/99), H3 (A/Wisconsin/67/2005), H5 (A/Vietnam/1203/04), H7 (A/Netherlands/219/03) and H9 (A/Hong Kong/1073/99) HA were expressed on the surface of PER.C6 cells. To measure mAb binding to different structural HA conformations, cells were detached from the plastic support using PBS-EDTA and subsequently treated with trypsin (TrypLE™ Select, Gibco) for 5 min at RT, washed (1%
15 BSA in PBS) and incubated for 15 min in citric acid–sodium phosphate buffer (pH 4.9). Cell samples were set aside after each processing step (untrypsinized/HA0; trypsinized/HA1-HA2; pH 4.9/fusion HA) and fractions of each treatment were incubated with mAb CR9114 for 1 hour. Cells were then incubated for 30 min with phycoerythrin-conjugated anti-human IgG (Southern Biotech) in 1% BSA. Stained cells were analysed
20 using a FACS Canto with FACS Diva software (Becton Dickinson). FACS binding of IgG1s to surface expressed HA was after sequential treatment with trypsin and pH 4.9 buffered medium and expressed as percentage binding to untreated HA (A). See Figure 1A.

Antibody CR9114 shows a marked decrease in binding after pH-shift indicating
25 specificity for an epitope present only before the low pH induced conformational change of the HA molecule.

Alternatively, to test whether the IgGs can block the low pH induced conformational change of HA, antibody CR9114 was added before the low pH step. Samples of consecutive treatments were split and stained with either phycoerythrin-
30 conjugated anti-human IgG (Southern Biotech). Stained cells were analysed using a FACS Canto with FACS Diva software (Becton Dickinson). See Fig 1B.

Antibody CR9114 shows a high level of residual binding to the various HAs after pH-shift indicating that when these antibodies are bound to the HA molecule, the low pH induced conformational change does not occur.

5 Example 8

Affinity measurements of Fabs on various influenza A and B HAs.

Recombinant soluble HA of A/New Caledonia/20/1999 (H1), A/Brisbane/59/2007 (H1), A/Wisconsin/67/2005 (H3), A/Brisbane/10/2007 (H3, B/Florida/4/2006 (B), B/Brisbane/60/2008 (B) and B/Malaysia/2506/2004 (B) produced using baculovirus
10 vectors in insect cells were purchased from Protein Sciences Corp (CT, USA) and biotinylated at room temperature (RT) for 40 min using EZ-link sulfo-NHS-LC-LC-biotin (Pierce). Buffer exchange step to PBS was performed using Amicon Ultra 0.5 ml Centrifugal Filters (Millipore). Biotinylated HA was bound to Streptavidin sensors at 37 °C for 1200 seconds. Association of Fab fragment of CR9005, CR9112, CR9113 and
15 CR9114 to HA was measured on Octet QK (ForteBio) for 700 seconds at 37 °C by exposing the sensors to 100 nM antibody in 1x kinetic buffer (ForteBio). Dissociation of the Fab fragments was assessed by exposing the sensors to 1x kinetic buffer for 9000 seconds at 37 °C. Fab fragments of CR9005, CR9112, CR9113 and CR9114 all bind with micro- to pico-molar affinities to H1, H3 and influenza B HA.

20

Example 9

Competition for binding with other stem binding antibodies

Recombinant soluble HA of A/New Caledonia/20/1999 (H1N1) and A/Wisconsin/67/2005 (H3N2) produced using baculovirus vectors in insect cells were
25 purchased from Protein Sciences Corp (CT, USA) and biotinylated at room temperature (RT) for 40 min using EZ-link sulfo-NHS-LC-LC-biotin (Pierce). Buffer exchange step to PBS was performed using Amicon Ultra 0.5 ml Centrifugal Filters (Millipore). Biotinylated HA was bound to Streptavidin sensors at 37 °C for 1200 seconds. Association of antibodies CR9114 and CR6261 to H1 HA was measured on Octet QK
30 (ForteBio) for 700 seconds at 37 °C by exposing the sensors to 100 nM antibody in 1x

kinetic buffer (ForteBio) after which the degree of additional binding was assessed by exposing the sensors to a second antibody (100 nM in 1x kinetic buffer) in the presence of the first antibody (100 nM) for 700 seconds at 37 °C. As a control, mAb CR9020, binding to the globular head of H1 was taken along. Association of antibodies CR9114 and CR8020 to H3 HA was measured on Octet QK (ForteBio) for 900 seconds at 37 °C by exposing the sensors to 100 nM antibody in 1x kinetic buffer (ForteBio) after which the degree of additional binding was assessed by exposing the sensors to a second antibody (100 nM in 1x kinetic buffer) in the presence of the first antibody (100 nM) for 900 seconds at 37 °C. As a control, mAb CR8057, binding to the globular head of H3 was taken along.

CR9114 competes for binding to H1 HA with CR6261 and to H3 HA with CR8020. CR9114 therefore likely binds an epitope overlapping with both the epitopes of CR6261 and CR8020 in the stem-region of HA. (See Fig. 2)

15 Example 10

Prophylactic activity of human IgG monoclonal antibody CR9114 against lethal influenza B challenge in vivo

A study was performed to test the prophylactic effect of the monoclonal antibody CR9114 against a lethal challenge with influenza B virus in vivo. MAb CR9114 was tested for prophylactic efficacy in a mouse lethal challenge model with mouse adapted influenza B/Florida/04/2006 virus (Central Veterinary Institute (CVI), Lelystad, The Netherlands). The B/Florida/04/2006 virus was adapted to mice after 5 lung-to-lung passages. The mouse adapted influenza B passage 5 virus was propagated in embryonated chicken eggs in CVI's laboratory. All mice (Balb/c, female, age 6-8 weeks, n=10 per group) were acclimatized and maintained for a period of at least 4 days prior to the start of the experiment. MAb CR9114 was dosed at 15 mg/kg intravenously in the tail vein (*vena coccygeus*) at day -1 before challenge, assuming an average weight of 18 g per mouse and a fixed dose volume of 0.2 mL. A control group was taken along dosed with vehicle control. The mice were then challenged at day 0 with 25 LD₅₀ B/Florida/04/2006 influenza B virus by intranasal inoculation. Clinical signs and body weights were determined daily from day-1 before challenge until day 8. Clinical signs were scored with

a scoring system (0=no clinical signs; 1=rough coat; 2=rough coat, less reactive during handling; 3=rough coat, rolled up, laboured breathing, less reactive during handling; 4=rough coat, rolled up, laboured breathing, inactive response to manipulation/handlings). At a score of 4 the animal was euthanized.

5 All mice were active and appeared healthy without showing signs of disease during the acclimatization period. Fig. 3A shows the survival rates of the mice, following mAb administration. Mice dosed with 15 mg/kg mAb CR9114 showed a survival rate of 100%, whereas in the control mAb group 50% survived.

10 In Figure 3B the mean body weight change of the mice during the 8 day study period following mAb administration is shown. In the mAb CR9114 group the mice did not loose weight over the 8 day study period, whereas in the vehicle control group weight loss was observed. Median clinical scores of the mice are depicted in Fig. 3C. Of the mice treated with 15 mg/kg mAb CR9114 at day -1 pre-challenge, all survived and none of the animals showed any clinical signs during the observation period (from day 0 to day 15 8 post infection). These results show that the human anti influenza antibody CR9114, identified and developed as disclosed herein, is able to provide protection against a lethal dose of influenza B virus *in vivo*. When administered one day prior to infection at a dose of 15 mg/kg or higher, mAb CR9114 was able to completely prevent clinical manifestation of influenza B infection in mice.

Table 1. First round Vkappa, Vlambda and VH amplifications

Primer name	Primer nucleotide sequence	SEQ ID NO:
OK1 (HuVK1B)	GAC ATC CAG WTG ACC CAG TCT CC	65
OK2 (HuVK2)	GAT GTT GTG ATG ACT CAG TCT CC	66
OK3 (HuVK2B2)	GAT ATT GTG ATG ACC CAG ACT CC	67
OK4 (HuVK3B)	GAA ATT GTG WTG ACR CAG TCT CC	68
OK5 (HuVK5)	GAA ACG ACA CTC ACG CAG TCT CC	69
OK6 (HuVK6)	GAA ATT GTG CTG ACT CAG TCT CC	70
OCK (HuCK)	ACA CTC TCC CCT GTT GAA GCT CTT	71
OL1 (HuVL1A) *	CAG TCT GTG CTG ACT CAG CCA CC	72
OL1 (HuVL1B) *	CAG TCT GTG YTG ACG CAG CCG CC	73
OL1 (HuVL1C) *	CAG TCT GTC GTG ACG CAG CCG CC	74
OL2 (HuVL2B)	CAG TCT GCC CTG ACT CAG CC	75
OL3 (HuVL3A)	TCC TAT GWG CTG ACT CAG CCA CC	76
OL4 (HuVL3B)	TCT TCT GAG CTG ACT CAG GAC CC	77
OL5 (HuVL4B)	CAG CYT GTG CTG ACT CAA TC	78
OL6 (HuVL5)	CAG GCT GTG CTG ACT CAG CCG TC	79
OL7 (HuVL6)	AAT TTT ATG CTG ACT CAG CCC CA	80
OL8 (HuVL7/8)	CAG RCT GTG GTG ACY CAG GAG CC	81
OL9 (HuVL9) #	CWG CCT GTG CTG ACT CAG CCM CC	82
OL9 (HuVL10) #	CAG GCA GGG CTG ACT CAG	83
OCL (HuCL2) X	TGA ACA TTC TGT AGG GGC CAC TG	84
OCL (HuCL7) X	AGA GCA TTC TGC AGG GGC CAC TG	85
OH1 (HuVH1B7A) +	CAG RTG CAG CTG GTG CAR TCT GG	86
OH1 (HuVH1C) +	SAG GTC CAG CTG GTR CAG TCT GG	87
OH2 (HuVH2B)	CAG RTC ACC TTG AAG GAG TCT GG	88
OH3 (HuVH3A)	GAG GTG CAG CTG GTG GAG	89
OH4 (HuVH3C)	GAG GTG CAG CTG GTG GAG WCY GG	90
OH5 (HuVH4B)	CAG GTG CAG CTA CAG CAG TGG GG	91
OH6 (HuVH4C)	CAG STG CAG CTG CAG GAG TCS GG	92
OH7 (HuVH6A)	CAG GTA CAG CTG CAG CAG TCA GG	93
OCM (HuCIgM)	TGG AAG AGG CAC GTT CTT TTC TTT	94

* Mix in 1:1:1 ratio

Mix in 1:1 ratio

X Mix in 1:1 ratio

+ Mix in 1:1 ratio

Table 2. Second round Vkappa, Vlambda and VH amplifications

Primer name	Primer nucleotide sequence	SEQ ID NO
OK1S (HuVK1B-SAL)	TGA GCA CAC AGG TCG ACG GAC ATC CAG WTG ACC CAG TCT CC	95
OK2S (HuVK2-SAL)	TGA GCA CAC AGG TCG ACG GAT GTT GTG ATG ACT CAG TCT CC	96
OK3S (HuVK2B2-SAL)	TGA GCA CAC AGG TCG ACG GAT ATT GTG ATG ACC CAG ACT CC	97
OK4S (HuVK3B-SAL)	TGA GCA CAC AGG TCG ACG GAA ATT GTG WTG ACR CAG TCT CC	98
OK5S (HuVK5-SAL)	TGA GCA CAC AGG TCG ACG GAA ACG ACA CTC ACG CAG TCT CC	99
OK6S (HuVK6-SAL)	TGA GCA CAC AGG TCG ACG GAA ATT GTG CTG ACT CAG TCT CC	100
OJK1 (HuJK1-NOT)	GAG TCA TTC TCG ACT TGC GGC CGC ACG TTT GAT TTC CAC CTT GGT CCC	101
OJK2 (HuJK2-NOT)	GAG TCA TTC TCG ACT TGC GGC CGC ACG TTT GAT CTC CAG CTT GGT CCC	102
OJK3 (HuJK3-NOT)	GAG TCA TTC TCG ACT TGC GGC CGC ACG TTT GAT ATC CAC TTT GGT CCC	103
OJK4 (HuJK4-NOT)	GAG TCA TTC TCG ACT TGC GGC CGC ACG TTT GAT CTC CAC CTT GGT CCC	104
OJK5 (HuJK5-NOT)	GAG TCA TTC TCG ACT TGC GGC CGC ACG TTT AAT CTC CAG TCG TGT CCC	105
OL1S (HuVL1A-SAL) *	TGA GCA CAC AGG TCG ACG CAG TCT GTG CTG ACT CAG CCA CC	106
OL1S (HuVL1B-SAL) *	TGA GCA CAC AGG TCG ACG CAG TCT GTG YTG ACG CAG CCG CC	107
OL1S (HuVL1C-SAL) *	TGA GCA CAC AGG TCG ACG CAG TCT GTC GTG ACG CAG CCG CC	108
OL2S (HuVL2B-SAL)	TGA GCA CAC AGG TCG ACG CAG TCT GCC CTG ACT CAG CC	109
OL3S (HuVL3A-SAL)	TGA GCA CAC AGG TCG ACG TCC TAT GWG CTG ACT CAG CCA CC	110
OL4S (HuVL3B-SAL)	TGA GCA CAC AGG TCG ACG TCT TCT GAG CTG ACT CAG GAC CC	111
OL5S (HuVL4B-SAL)	TGA GCA CAC AGG TCG ACG CAG CYT GTG CTG ACT CAA TC	112
OL6S (HuVL5-SAL)	TGA GCA CAC AGG TCG ACG CAG GCT GTG CTG ACT CAG CCG TC	113
OL7S (HuVL6-SAL)	TGA GCA CAC AGG TCG ACG AAT TTT ATG CTG ACT CAG CCC CA	114
OL8S (HuVL7/8-SAL)	TGA GCA CAC AGG TCG ACG CAG RCT GTG GTG ACY CAG GAG CC	115
OL9S (HuVL9-SAL) #	TGA GCA CAC AGG TCG ACG CWG CCT GTG CTG ACT CAG CCM CC	116
OL9S (HuVL10-SAL) #	TGA GCA CAC AGG TCG ACG CAG GCA GGG CTG ACT CAG	117
OJL1 (HuJL1-NOT)	GAG TCA TTC TCG ACT TGC GGC CGC ACC TAG GAC GGT GAC CTT GGT CCC	118
OJL2 (HuJL2/3-NOT)	GAG TCA TTC TCG ACT TGC GGC CGC ACC TAG GAC GGT CAG CTT GGT CCC	119
OJL3 (HuJL7-NOT)	GAG TCA TTC TCG ACT TGC GGC CGC ACC GAG GAC GGT CAG CTG GGT GCC	120

OH1S (HuVH1B-SFI)+	GTC CTC GCA ACT GCG GCC CAG CCG GCC ATG GCC CAG RTG CAG CTG GTG CAR TCT GG	121
OH1S (HuVH1C-SFI)+	GTC CTC GCA ACT GCG GCC CAG CCG GCC ATG GCC SAG GTC CAG CTG GTR CAG TCT GG	122
OH2S (HuVH2B-SFI)	GTC CTC GCA ACT GCG GCC CAG CCG GCC ATG GCC CAG RTC ACC TTG AAG GAG TCT GG	123
OH3S (HuVH3A-SFI)	GTC CTC GCA ACT GCG GCC CAG CCG GCC ATG GCC GAG GTG CAG CTG GTG GAG	124
OH4S (HuVH3C-SFI)	GTC CTC GCA ACT GCG GCC CAG CCG GCC ATG GCC GAG GTG CAG CTG GTG GAG WCY GG	125
OH5S (HuVH4B-SFI)	GTC CTC GCA ACT GCG GCC CAG CCG GCC ATG GCC CAG GTG CAG CTA CAG CAG TGG GG	126
OH6S (HuVH4C-SFI)	GTC CTC GCA ACT GCG GCC CAG CCG GCC ATG GCC CAG STG CAG CTG CAG GAG TCS GG	127
OH7S (HuVH6A-SFI)	GTC CTC GCA ACT GCG GCC CAG CCG GCC ATG GCC CAG GTA CAG CTG CAG CAG TCA GG	128
OJH1 (HuJH1/2-XHO)	GAG TCA TTC TCG ACT CGA GAC RGT GAC CAG GGT GCC	129
OJH2 (HuJH3-XHO)	GAG TCA TTC TCG ACT CGA GAC GGT GAC CAT TGT CCC	130
OJH3 (HuJH4/5-XHO)	GAG TCA TTC TCG ACT CGA GAC GGT GAC CAG GGT TCC	131
OJH4 (HuJH6-XHO)	GAG TCA TTC TCG ACT CGA GAC GGT GAC CGT GGT CCC	132

* Mix in 1:1:1 ratio

Mix in 1:1 ratio

+ Mix in 1:1 ratio

Table 3. Second round VL regions amplification overview

Template	5' primer	3' primer	Product	Share in PK/PL (%)	Pool	Share in VL (%)
K1	OK1S	OJK1	K1J1	25	PK1	30
	OK1S	OJK2	K1J2	25		
	OK1S	OJK3	K1J3	10		
	OK1S	OJK4	K1J4	25		
	OK1S	OJK5	K1J5	15		
K2	OK2S	OJK1	K2J1	25	PK2	4
	OK2S	OJK2	K2J2	25		
	OK2S	OJK3	K2J3	10		
	OK2S	OJK4	K2J4	25		
	OK2S	OJK5	K2J5	15		
K3	OK3S	OJK1	K3J1	25	PK3	1
	OK3S	OJK2	K3J2	25		
	OK3S	OJK3	K3J3	10		
	OK3S	OJK4	K3J4	25		
	OK3S	OJK5	K3J5	15		
K4	OK4S	OJK1	K4J1	25	PK4	19
	OK4S	OJK2	K4J2	25		
	OK4S	OJK3	K4J3	10		
	OK4S	OJK4	K4J4	25		
	OK4S	OJK5	K4J5	15		
	OK5S	OJK1	K5J1	25		

K5	OK5S	OJK2	K5J2	25	PK5	1
	OK5S	OJK3	K5J3	10		
	OK5S	OJK4	K5J4	25		
	OK5S	OJK5	K5J5	15		
K6	OK6S	OJK1	K6J1	25	PK6	5
	OK6S	OJK2	K6J2	25		
	OK6S	OJK3	K6J3	10		
	OK6S	OJK4	K6J4	25		
	OK6S	OJK5	K6J5	15		
L1	OL1S	OJL1	L1J1	30	PL1	14
	OL1S	OJL2	L1J2	60		
	OL1S	OJL3	L1J3	10		
L2	OL2S	OJL1	L2J1	30	PL2	10
	OL2S	OJL2	L2J2	60		
	OL2S	OJL3	L2J3	10		
L3	OL3S	OJL1	L3J1	30	PL3	10
	OL3S	OJL2	L3J2	60		
	OL3S	OJL3	L3J3	10		
L4	OL4S	OJL1	L4J1	30	PL4	1
	OL4S	OJL2	L4J2	60		
	OL4S	OJL3	L4J3	10		
L5	OL5S	OJL1	L5J1	30	PL5	1
	OL5S	OJL2	L5J2	60		
	OL5S	OJL3	L5J3	10		

L6	OL6S	OJL1	L6J1	30	PL6	1
	OL6S	OJL2	L6J2	60		
	OL6S	OJL3	L6J3	10		
L7	OL7S	OJL1	L7J1	30	PL7	1
	OL7S	OJL2	L7J2	60		
	OL7S	OJL3	L7J3	10		
L8	OL8S	OJL1	L8J1	30	PL8	1
	OL8S	OJL2	L8J2	60		
	OL8S	OJL3	L8J3	10		
L9	OL9S	OJL1	L9J1	30	PL9	1
	OL9S	OJL2	L9J2	60		
	OL9S	OJL3	L9J3	10		
					VL	100%

Table 4. Second round VH regions amplification overview

Template	5' primer	3' primer	Product	Share in PK/PL (%)	Pool	Share in VH (%)
H1	OH1S	OJH1	H1J1	10	PH1	25
	OH1S	OJH2	H1J2	10		
	OH1S	OJH3	H1J3	60		
	OH1S	OJH4	H1J4	20		
H2	OH2S	OJH1	H2J1	10	PH2	2
	OH2S	OJH2	H2J2	10		
	OH2S	OJH3	H2J3	60		
	OH2S	OJH4	H2J4	20		
H3	OH3S	OJH1	H3J1	10	PH3	25
	OH3S	OJH2	H3J2	10		
	OH3S	OJH3	H3J3	60		
	OH3S	OJH4	H3J4	20		
H4	OH4S	OJH1	H4J1	10	PH4	25
	OH4S	OJH2	H4J2	10		
	OH4S	OJH3	H4J3	60		
	OH4S	OJH4	H4J4	20		
H5	OH5S	OJH1	H5J1	10	PH5	2
	OH5S	OJH2	H5J2	10		
	OH5S	OJH3	H5J3	60		

	OH5S	OJH4	H5J4	20		
H6	OH6S	OJH1	H6J1	10	PH6	20
	OH6S	OJH2	H6J2	10		
	OH6S	OJH3	H6J3	60		
	OH6S	OJH4	H6J4	20		
H7	OH7S	OJH1	H7J1	10	PH7	1
	OH7S	OJH2	H7J2	10		
	OH7S	OJH3	H7J3	60		
	OH7S	OJH4	H7J4	20		
					VH	100%

- Table 5:** Cross-binding activity of PEG/NACl-precipitated and filter-sterilized single-chain phage antibodies to HA of different subtypes, as measured by ELISA. + = binding (>4x background); +/- = low binding (2-4x background) - = no detectable binding; H1= HA of influenza A H1 subtype; H3= HA of influenza A H3 subtype; H5=HA of influenza A H5 subtype; H7=HA of influenza A H7 subtype; B= HA of influenza virus B; Rabies=Glycoprotein of Rabies virus (negative control).

	Phage midi Elisa					
	H1	H3	H5	H7	B	Rabies
sc09-003	+	+	+	+	+	-
sc09-004	+	+	+	+	+	-
sc09-005	+	+	+	+	+	-
sc09-006	+	+	+	+	+	-
sc09-007	+	+/-	+	+	+/-	-
sc09-008	+	+/-	+	+	+/-	-
sc09-009	+	+/-	+	+	+/-	-
sc09-010	+	+	+	+	+/-	-
sc09-011	+	+	+	+	+	-
sc09-012	+	+	+	+	-	-
sc09-029	+	+/-	+	+	-	-
sc09-030	+	+	+	+	+	-
sc09-031	+	+/-	+	+	-	-
sc09-112	+	+	+	+	+	-
sc09-113	+	+	+	+	+	-
sc09-114	+	+	+	+	+	-

Table 6. FACS analysis of PEG/NaCl-precipitated and filter-sterilized phage antibodies. + = binding (>4x background); +/- = low binding (2-4x background) - = no detectable binding; PER.C6=untransfected PER.C6 cells (control); mH1, mH3, mH7= membrane bound HA of the subtypes H1, H3 and H7 subtypes, respectively.

5

	Phage midi Facs (% gated UL)			
	PerC6	mH1	mH3	mH7
sc09-003	-	+	+	+
sc09-004	-	+	+	+
sc09-005	-	+	+	+
sc09-006	-	+	+	+
sc09-007	-	+	+/-	+
sc09-008	-	+	+/-	+
sc09-009	-	+	+/-	+
sc09-010	-	+	+	+
sc09-011	-	+	+	+
sc09-012	-	+	+	+
sc09-029	-	+	-	+/-
sc09-030	-	+	+	+
sc09-031	-	+	-	+/-
sc09-112	-	+	+	+
sc09-113	-	+	+	+
sc09-114	-	+	+	+

Table 7. Data of the CDR regions of the HA specific immunoglobulins. The SEQ ID NO is given between brackets.

IgG#	VH	HC CDR1	HC CDR2	HC CDR3	VL	LC CDR1	LC CDR2	LC CDR3
CR9003	IGHV1-69*06	GGTSNNFG (133)	ISPIFGST (134)	ARHGNYFYSGMDL (135)	IGLV3-21*02	NVGSNS (136)	DDR (137)	QVWDSSSDHRV (138)
CR9004	IGHV1-69*06	GGTSNNYA (139)	VSPIFGST (140)	ARHGNYNYSMDV (141)	IGLV1-44*01	DSNIGRRS (142)	SND (143)	AAWDDSLKGAV (144)
CR9005	IGHV1-69*06	GGTSNNYA (139)	ISPIFGST (134)	ARHGNYYSMDL (145)	IGLV2-14*01	SSDVGGYNY (146)	DVS (174)	CSYAGSAKGV (147)
CR9006	IGHV1-69*06	GGTSNNYA (139)	ISPIFGST (134)	ARHGNYYSMDL (145)	IGLV3-21*02	NIGSKT (148)	GDS (149)	QVWDSSSDHPGAV (150)
CR9007	IGHV1-69*06	GGTSNNYA (139)	ISPIFGSA (151)	ARHGNYYSMDV (152)	IGLV1-44*01	SSNIGSNT (153)	GDD (154)	ATWDDSLNGHV (155)
CR9008	IGHV1-69*06	GGTSNNYA (139)	ISPIFGST (134)	ARHGNYYSMDV (152)	IGLV3-21*02	NIGSKT (148)	GDS (149)	QVWDSSSDHPGAV (150)
CR9009	IGHV1-69*06	GGTSNNYA (139)	ISPIFGST (134)	ARHGNYYSMDV (152)	IGKV1-12*01	QHSSW (156)	SAS (157)	QQANSFPLT (158)
CR9010	IGHV1-69*06	GGTSNNYA (139)	ISPIFGST (134)	ARHGNYYSMDV (152)	IGLV3-21*02	NIGSKT (148)	VDS (159)	QVWDSSSDHPGAV (160)
CR9011	IGHV1-69*06	GGTSNNYA (139)	ISPIFGSA (151)	ARHGNYYSMDV (161)	IGLV1-44*01	DSNIGRRS (142)	SND (143)	AAWDDSLKGAV (144)
CR9012	IGHV1-69*06	GGTSNNYA (139)	ISPIFGSA (151)	ARHGTYYSMDV (162)	IGLV1-40*02	SSNIGAGYD (163)	GNN (164)	QSYDQNLSEGV (165)
CR9029	IGHV1-69*06	GGTSNNYA (139)	ISPIFGST (134)	ARHGNYYSMDV (152)	IGKV3-20*01	QSVSSY (166)	GAS (167)	QQYGSSPFA (168)
CR9030	IGHV1-69*06	GGTSNNYA (139)	ISPIFGST (134)	ARHGNYYSMDV (152)	IGLV3-21*02	NIGSKS (169)	GDS (149)	QVWDSSSDHPGAV (150)
CR9031	IGHV1-69*06	GGTSNNYA (139)	ISPIFGST (134)	ARHGNYYSMDV (141)	IGLV1-40*01	SSNIGAGYD (163)	DNN (169)	QSYDSSLSPYV (170)
CR9112	IGHV1-69*06	GGTSNNYA (139)	ISPIFGST (134)	ARHGNYYSMDV (152)	IGLV1-40*01	SANIGAGYD (171)	GNN (164)	QSYDSSLGAL (172)
CR9113	IGHV1-69*06	GGTSNNYA (139)	ISPIFGST (134)	ARHGNYYSMDL (145)	IGLV1-44*01	DSNIGRRS (142)	SND (143)	AAWDASLSPV (173)
CR9114	IGHV1-69*06	GGTSNNYA (139)	ISPIFGST (134)	ARHGNYYSMDV (152)	IGLV1-44*01	DSNIGRRS (142)	SND (143)	AAWDDSLKGAV (144)

Table 8. Identity cross-tables of the amino acid sequences of the heavy and light chain variable domains.

A.

		Amino acid differences in Heavy Chain															
		SC09-007	SC09-011	SC09-112	SC09-010	SC09-029	SC09-008	SC09-030	SC09-114	SC09-009	SC09-004	SC09-031	SC09-005	SC09-006	SC09-012	SC09-113	SC09-003
Percentage identity	SC09-007		2	3	5	5	6	5	4	5	5	6	7	9	3	11	15
	SC09-011	98.4		5	5	5	6	7	6	7	7	8	9	9	3	13	15
	SC09-112	97.5	95.9		2	2	3	2	1	2	2	3	4	6	6	8	12
	SC09-010	95.9	95.9	98.4		0	3	4	3	4	4	5	6	4	6	10	10
	SC09-029	95.9	95.9	98.4	100.0		3	4	3	4	4	5	6	4	6	10	10
	SC09-008	95.0	95.0	97.5	97.5	97.5		3	2	3	5	6	5	5	7	9	11
	SC09-030	95.9	94.2	98.4	96.7	96.7	97.5		1	2	4	5	4	6	8	6	12
	SC09-114	96.7	95.0	99.2	97.5	97.5	98.4	99.2		1	3	4	3	5	7	7	11
	SC09-009	95.9	94.2	98.4	96.7	96.7	97.5	98.4	99.2		4	5	4	6	8	8	12
	SC09-004	95.9	94.2	98.4	96.7	96.7	95.9	96.7	97.5	96.7		3	6	8	8	10	14
	SC09-031	95.0	93.4	97.5	95.9	95.9	95.0	95.9	96.7	95.9	97.5		5	7	9	11	15
	SC09-005	94.2	92.6	96.7	95.0	95.0	95.9	96.7	97.5	96.7	95.0	95.9		2	8	6	10
	SC09-006	92.6	92.6	95.0	96.7	96.7	95.9	95.0	95.9	95.0	93.4	94.2	98.4		8	8	8
	SC09-012	97.5	97.5	95.0	95.0	95.0	94.2	93.4	94.2	93.4	93.4	92.6	93.4	93.4		12	14
	SC09-113	90.9	89.3	93.4	91.7	91.7	92.6	95.0	94.2	93.4	91.7	90.9	95.0	93.4	90.1		8
	SC09-003	87.6	87.6	90.1	91.7	91.7	90.9	90.1	90.9	90.1	88.4	87.6	91.7	93.4	88.4	93.4	

B.

		Amino acid differences in Light Chain															
		SC09-011	SC09-114	SC09-004	SC09-113	SC09-007	SC09-012	SC09-112	SC09-031	SC09-005	SC09-006	SC09-008	SC09-030	SC09-010	SC09-003	SC09-009	SC09-029
Percentage identity	SC09-011		0	2	7	14	29	26	34	44	47	47	45	52	47	62	64
	SC09-114	100.0		2	7	14	29	26	34	44	47	47	45	52	47	62	64
	SC09-004	98.2	98.2		5	16	27	24	32	42	49	49	47	54	49	62	64
	SC09-113	93.6	93.6	95.5		17	25	22	29	41	46	46	44	51	47	62	64
	SC09-007	87.3	87.3	85.5	84.6		26	25	32	42	41	41	41	47	43	61	61
	SC09-012	73.9	73.9	75.7	77.5	76.6		9	13	39	48	48	47	52	48	61	62
	SC09-112	76.6	76.6	78.4	80.2	77.5	91.9		13	37	45	45	44	51	45	60	60
	SC09-031	69.9	69.9	71.7	74.3	71.7	88.5	88.5		37	50	50	49	53	46	60	62
	SC09-005	60.4	60.4	62.2	63.1	62.2	64.9	66.7	67.3		55	55	54	56	46	64	63
	SC09-006	58.0	58.0	56.3	58.9	63.4	57.5	60.2	55.8	51.3		0	3	7	17	64	61
	SC09-008	58.0	58.0	56.3	58.9	63.4	57.5	60.2	55.8	51.3	100.0		3	7	17	64	61
	SC09-030	59.8	59.8	58.0	60.7	63.4	58.4	61.1	56.6	52.2	97.3	97.3		10	14	62	59
	SC09-010	53.6	53.6	51.8	54.5	58.0	54.0	54.9	53.1	50.4	93.6	93.6	90.9		22	67	67
	SC09-003	57.7	57.7	55.9	57.7	61.3	57.1	59.8	59.3	58.6	84.6	84.6	87.3	80.0		62	56
	SC09-009	45.1	45.1	45.1	45.1	46.0	46.5	47.4	47.4	43.4	42.9	42.9	44.6	40.2	44.1		34
	SC09-029	43.4	43.4	43.4	43.4	46.0	45.6	47.4	45.6	44.3	45.5	45.5	47.3	40.2	49.6	68.2	

Table 9. Cross-binding reactivity of IgGs, as measured by ELISA and FACS.

H1=soluble recombinant A/New Caledonia/20/1999 H1 HA; H3= soluble recombinant A/Wisconsin/67/2005 H3 HA; H5= soluble recombinant A/Vietnam/1203/04 H5 HA; H7= soluble recombinant A/Netherlands/219/2003 H7 HA; H9= soluble recombinant A/Hong Kong/1073/99 H9 HA; B= soluble recombinant B/Ohio/01/05 influenza B HA; Rabies= rabies glycoprotein; PER.C6=untransfected PER.C6 cells (control); mH1=PER.C6 expressed A/New Caledonia/20/1999 H1 HA; mH3= PER.C6 expressed A/Wisconsin/67/2005 H3 HA; mH7= PER.C6 expressed A/Netherlands/219/2003 H7 HA; ND=not done. + = binding (>10x background); +/- = low binding (2-10x background) - = no detectable binding.

	IgG Elisa							IgG Facs			
	H1	H3	H5	H7	H9	B	Rabies	PerC6	mH1	mH3	mH7
CR9005	+	+	+	+	+	+	-	-	+	+	+
CR9030	+	+	+	+	+	+/-	-	-	+	+	+
CR9112	+	+	+	+	+	+	-	-	+	+	+
CR9113	+	+	+	+	+	+	-	-	+	+	+
CR9114	+	+	+	+	+	+	-	-	+	+	+
CR4098	-	-	-	-	-	-	+	-	-	-	-

Table 10. Cross-neutralizing activity of IgGs; Titers (indicated in $\mu\text{g/ml}$) are geomean IC50 values as determined according to the Spearman-Kärber method of at least duplicate experiments; >100 = not neutralizing at highest tested concentration (100 $\mu\text{g/ml}$).

	Subtype	Strain	CR9005	CR9112	CR9113	CR9114
Group I	H1	A/WSN/33	1.1	0.9	1.1	1.1
		A/New Caledonia/20/99	2.6	1.9	4.4	3.7
		A/Solomon Islands/3/2006	1.4	1.3	2.2	1.8
		A/Brisbane/59/2007	3.4	2	3.1	2.6
		A/California/7/2009	0.7	0.5	0.3	0.3
	H2	A/Env/MPU3156/05	8.8	6.3	8.8	8.8
	H5	A/Hong Kong/156/97	0.8	0.7	0.9	0.4
		A/EW/MPF461/07	10.5	10.5	8.8	10.5
	H6	A/EW/MPD411/07	29.7	10.5	17.7	10.5
	H8	A/EW/MPH571/08	8.8	8.8	8.8	8.8
	H9	A/Hong Kong/1073/99	6.3	3.7	3.7	4.4
A/Ck/HK/SSP176/09		4.4	4.4	6.3	6.3	
Group II	H3	A/Hong Kong/1/68	42	27.6	22.3	19
		A/Johannesburg/33/94	17.7	13.8	32.4	21.9
		A/Panama/2007/1999	28.2	47.5	47.5	39.9
		A/Hiroshima/52/2005	22.9	10.5	13.6	12.5
		A/Wisconsin/67/2005	35.4	29.7	35.4	32.4
		A/Brisbane/10/2007	11.2	5.6	9.4	5.6
	H4	A/WF/MPA 892/06	1.2	0.8	1.3	0.8
	H7	A/Mallard/Netherlands/12/2000	9.6	6.3	6.3	4.8
		A/New York/107/2003	> 100	> 100	> 100	> 100
	H10	A/Chick/Germany/N/49	29.6	26.5	19.8	15.7
H14	A/Mallard/Astrakhan/263/1982	> 100	> 100	> 100	> 100	

SEQUENCES

>SC09-003 VH DNA (SEQ ID NO: 1)
GAGGTGCAGCTGGTGGAGTCTGGGGCTGAGGTCAAGAAGGCTGGGTCTCGGTGAAAGTCTCCTGCAAGTC
TTCTGGAGGCACCTCCAACAACCTTTGGTATCAGCTGGGTACGACAGGCCCCCTGGCCAAGGCCTTGAGTGGA
TGGGCGGGATCAGCCCAATCTTTGGTTCGACAGTCTACGCACAGAAATTTCCAGGGCAGAGTCACATATTTCC
GCGGACATATTTTTCACACACTGCCACATGGAGATGAACAGCCTGACATCTGAGGACACGGCCGTCTATTT
CTGTGCGAGGCACGGAAATTATTATTTCTACTCCGGTATGGACCTCTGGGGCCAAGGGACCACGGTCACC

>SC09-003 VH PROTEIN (SEQ ID NO: 2)
EVQLVESGAIEVKKAGSSVKVSKSSGGTSSNFGISWVRQAPQGLEWMGGISPIFGSTVYAQKFQGRVTIS
ADIFSHYAYMEMNSLTSEDYAVYFCARHGNYFYSGMDLWQGTTVT

>SC09-003 VL DNA (SEQ ID NO: 3)
TCCTATGTGCTGACTCAGCCACCTCGGTGTCAGTGGCCCCAGGACAGACGGCCACGATTTCTGTGGGGG
AGACAACGTTGGAAGTAACAGTGTGACTGTTACCAGCAGAAGCCAGGCCAGGCCCTGTGCTGGTCTGTCT
ATGATGATCGCGACCGACCTCAGGGATCCCTGAGCGATTCTCTGGCTCCAACCTCTGGGAACACGGCCACC
CTGACCATCAGCAGGGTCAAGCCGGGATGAGGCCGACTATTACTGTGAGGTGTGGGATAGTAGTAGTGA
TCATCGAGTCTTCGGAACGGGACCAAGTCCACCGTCCCTAG

>SC09-003 VL PROTEIN (SEQ ID NO: 4)
SYVLTQPPSVSVAPGQTATISCGGDNVGSNSVHWYQQKPGQAPVLLVYDDRDRPSGIPERFSGSNSGNTAT
LTISRVEAGDEADYICQVWDSSTDRHVFVGTGKVTVL

>SC09-004 VH DNA (SEQ ID NO: 5)
CAGGTCCAGCTGGTACAGTCTGGGGCTGAGGTGAAGAAGCCTGGGTCTCGGTGAAAGTCTCCTGCAAGTC
TTCTGGCGGCACCTCCAATAACTATGCCATCAGCTGGGTGCGACAGGCCCCCTGGACAAGGCCTTGACTGGA
TGGGCGGGGTGAGCCCTATCTTTGGTTCGACAGCCTACGCACAGAAAGTCCAGGGCAGAGTCACATATTTCC
GCGGACATATTTTTCGAACACAGCTACATGGAGCTGAACAGTCTGACATCTGAGGACACGGCCGTCTATTA
TTGTGCGAGACACGGGAATTATTATTACAACCTCCGGTATGGACGTCTGGGGCCAAGGGACCACGGTCACC

>SC09-004 VH PROTEIN (SEQ ID NO: 6)
QVQLVQSGAEVKKPKGSSVKVSKSSGGTSSNYAISWVRQAPQGLDWMGGVSPIFGSTAYAQKFQGRVTIS
ADIFSNYAYMELNSLTSEDYAVYYCARHGNYYSNGMDVWQGTTVT

>SC09-004 VL DNA (SEQ ID NO: 7)
CAGTCTGTGCTGACGCAGCCGCCCGCAGTGTCTGGGACCCCCGGGCAGAGGGTCAACATCTCGTGTTCCTGG
AAGTGATTCCAACATCGGGAGAAGAAGTGTAACTGGTACCAGCAGTCCCAGGAACGGCCCCCAAACCTCC
TCATCTATAGTAACGATCAGCGCCCTCAGTGGTCCCTGACCGATTCTCTGGCTCCAAGTCCGGCACCTCA
GCCTCCCTGGCCATCAGTGGGCTCCAGTCTGAAGATGAGGCCGAATATTACTGTGCAGCATGGGATGACAG
CCTGAAGGGGGCTGTGTTCGGAGGAGCACCCAGCTGACCGTCCCTCG

>SC09-004 VL PROTEIN (SEQ ID NO: 8)
QSVLTQPPAVSGTPGQRVTISCSGSDSNIGRRSVNHWYQQFPGTAPKLLIYSNDQRPSVVPDRFSGSKSGTS
ASLAISGLQSEDEAEYYCAAWDDSLKGAVFVGGTQLTVL

>SC09-005 VH DNA (SEQ ID NO: 9)
CAGGTGCAGCTGGTGAATCTGGGGCTGAGGTCAAGAGGCCTGGGTCTCGGTGAAAGTCTCCTGCAAGTC
TTCTGGAGGCACCTCCAATAACTATGCTATTAGTTGGGTGCGACAGGCCCCCTGGACAAGGCCTTGACTGGA
TGGGCGGGATCAGCCCTATCTTTGGTTCGACAGTCTACGCACAGAAATTTCCAGGGCAGAGTCACATATTTCC
GCGGACATATTTTTCGAACACAGCTACATGGAGCTGAACAGCCTGACATCTGAGGACACGGCCGTATATTT
CTGTGCGAGGCACGGAACTATTATTACTACTCCGGTATGGACCTCTGGGGCCAAGGGACCACGGTCACC

>SC09-005 VH PROTEIN (SEQ ID NO: 10)

QVQLVQSGAEVKRPGSSVKVSKSSGGTSNNYAISWVRQAPGQGLDWMGGISPIFGSTVYAQKFQGRVTIS
ADIFSN TAYMELNSLTSEDTAVYFCARHGNYYYYS GMDLWGQGT TTVT

>SC09-005 VL DNA (SEQ ID NO: 11)
CAGTCTGCCCTGACTCAGCCTGCCTCCGTGTCTGGGTCTCCTGGACAGTCGATCACCATCTCCTGCAC TGG
AACCAGCAGTGACGTCGGTGGTTATAACTATGTCTCCTGGTACCAACAACACCCAGGCAAAGCCCCCAAAC
TCCTGATTTTTGATGTCAGTGATCGGCCCTCAGGGGTTTCTGATCGCTTCTCTGGCTCCAAGTCTGCGGAC
ACGGCCTCCCTGACCATCTCTGGACTCCAGGCTCAGGACGAGGCTGATTATTACTGCTGCTCATATGCAGG
TAGTGCCAAGGGCGTCTTCGGAAC TGGGACCAAGGTCACCGTCTCTAG

>SC09-005 VL PROTEIN (SEQ ID NO: 12)
QSALTQPASVSGSPGQSITISCTGTSSDVGGYNYVSWYQQHPGKAPKLLIFDVSDRPSGVSDRFSGSKSAD
TASLTISGLQAQDEADYYCCSYAGSAKGVFVGTGTVTL

>SC09-006 VH DNA (SEQ ID NO: 13)
GAGGTGCAGCTGGTGGAGTCTGGGGCTGAGGTCAAGAGGCCTGGGTCTCGGTGAAAGTCTCCTGCAAGTC
TTCTGGAGGCACCTCCAATAACTATGCTATTAGTTGGGTGCGACAGGCCCTGGACAAGGCCTTGACTGGA
TGGGCGGGATCAGCCCTATCTTTGGTTCGACAGTCTACGCACAGAAATCCAGGGCAGAGTCAC TATTTCC
GCGGACATATTTTGAACACAGCTACATGGAGCTGAACAGCCTGACATCTGAGGACACGGCCGTATATTT
CTGTGCGAGGCACGGGAAC TATTATTACTACTCCGGTATGGACCTCTGGGGCCAAGGGACCACGGTCACC

>SC09-006 VH PROTEIN (SEQ ID NO: 14)
EVQLVESGAEVKRPGSSVKVSKSSGGTSNNYAISWVRQAPGQGLDWMGGISPIFGSTVYAQKFQGRVTIS
ADIFSN TAYMELNSLTSEDTAVYFCARHGNYYYYS GMDLWGQGT TTVT

>SC09-006 VL DNA (SEQ ID NO: 15)
TCCTATGTGCTGACTCAGCCACCTCGGTGTCTAGTGGCCCCAGGACAGACGGCCAGGATTACCTGTGGGGG
AAACAACATTGGAAGTAAACTGTGCATTTGGTACCAGCAGA AACTCAGGCCAGGCCCTGTGCTGGTCTGTCT
ATGGTGTATAGCGACCGGCCCTCAGGGATCCCTGAGCGATTCTCTGGCTCCA AACTCTGGGACCACGGCCACC
CTGACCATCAGCAGGCTCGAAGCCGGGGATGAGGCCGACTATTACTGTCTAGGTGTGGGATAGTAGTAGTGA
TCATCCCGGTGCTGTGTTCCGAGGAGGCACCCAGCTGACCGTCTCTCG

>SC09-006 VL PROTEIN (SEQ ID NO: 16)
SYVLTQPPSVSVAPGQTARITCGGNNIGSKTVHWYQQNSGQAPVLVYVYGDSDRPSGIPERFSGSNSGTTAT
LTISRVEAGDEADYYCQVWDSSSDHPGAVFVGGTQLTVL

>SC09-007 VH DNA (SEQ ID NO: 17)
CAGGTGCAGCTGGTGC AATCTGGAGCTGAGGTCAAGAAGCCTGGGTCTCGGTGAAAGTCTCCTGCAAGTC
TTCTGGAGGCACCTCCAATAACTATGCTATCAGCTGGGTGCGACAGGCCCTGGACAAGGCCTTGACTGGA
TGGGAGGGATCAGCCCTATCTTTGGTTCAGCAGCTACGCACAGAAAGTTCCAGGGCAGAGTCAC TATTTACC
GCGGACATATTTTGAACACAGTGTACATGGAGCTGAACAGCCTGACATCTGAGGACACGGCCGTGTATTA
CTGTGCGAGACACGGGAATTATTATTACTACTCCGGTATGGACGTCTGGGGCCAAGGGACCACGGTCACCC
TCTCGAGC

>SC09-007 VH PROTEIN (SEQ ID NO: 18)
QVQLVQSGAEVKKPGSSVKVSKSSGGTSNNYAISWVRQAPGQGLDWMGGISPIFGSAAYAQKFQGRVTIT
ADIFSN TVYMELNSLTSEDTAVYYCARHGNYYYYS GMDVWVGQGT TTVTVSS

>SC09-007 VL DNA (SEQ ID NO: 19)
TCCTATGTGCTGACTCAGCCACCTCAGCGTCTGGGACCCCCGGGCAGAGGGTCACCATCTCTTGTCTGG
AAGCAGTCCAACATCGGAAGTAACTGTAAACTGGTACCAGCAGGTCCCCGGAACGGCCCCCAAAC TCC
TCATCTATGGTGTATGATCAGCGGCCCTCAGGGGTCCCTGACCGATTCTCTGGCTCCAAGTCTGGCACCTCA
GCCTCCCTGGCCATCAGTGGGCTCCAGTCTGAGGATGAGGCTGATTATTACTGTGCAACATGGGATGACAG
CCTGAATGGTCATGTGTTCCGAGGAGGCACCCAGCTGACCGTCTCTCG

>SC09-007 VL PROTEIN (SEQ ID NO: 20)

SYVLTQPPSASGTPGQRVITISCSGSSSNIGSNTVNWYQQVPGTAPKLLIYGDDQRPSGVPDRFSGSKSGTS
ASLAISGLQSEDEADYYCATWDDSLNGHVFGGGTQLTVL

>SC09-008 VH DNA (SEQ ID NO: 21)

GAGGTCCAGCTGGTGCAGTCTGGGGCTGAGGTCAAGAAGCCTGGGTCTCGGTGAGAGTCTCCTGTAAGTC
TTCTGGAGGCACCTCCAATAACTATGCTATCAGCTGGGTGCGACAGGCCCTGGACAAGGCCTTGACTGGA
TGGGCGGGATCAGCCCTATCTTTGGTTCGACAGCCTACGCACAGAAGTTCCAGGGCAGAGTCACTATTTCC
GCGGACATATTTTGAACACAGCCTACATGGAGCTGAACAGCCTGACATCTGAGGACACGGCCGTATATTT
CTGTGCGAGGCACGGGAATTATTATTACTACTCCGGTATGGACGTCTGGGGCCAAGGGACCACGGTCACCG
TCTCGAGC

>SC09-008 VH PROTEIN (SEQ ID NO: 22)

EVQLVQSGAEVKKPGSSVRVSKSSGGTSNNYAIWVWRQAPGQGLDWMGGISPIFGSTAYAQKFQGRVTIS
ADIFSNAYMELNSLTSEDVAVYFCARHGNYYYSGMDVWVGQGTTVTVSS

>SC09-008 VL DNA (SEQ ID NO: 23)

TCCTATGTGCTGACTCAGCCACCTCGGTGTGAGTGGCCCCAGGACAGACGGCCAGGATTACCTGTGGGGG
AAACAACATTGGAAGTAAACTGTGCATTTGGTACCAGCAGAAGTCAAGGCCAGGCCCTGTGCTGGTCTCT
ATGGTGTATAGCGACCGGCCCTCAGGGATCCCTGAGCGATTCTCTGGCTCCAAGTCTGGGACCACGGCCACC
CTGACCATCAGCAGGGTCAAGCCGGGGATGAGGCCGACTATTACTGTGAGGTGTGGGATAGTAGTAGTGA
TCATCCCGGTGCTGTGTTCCGAGGAGGCACCCAGCTGACCGTCTCTCG

>SC09-008 VL PROTEIN (SEQ ID NO: 24)

SYVLTQPPSVSVAPGQTARITCGGNNIGSKTVHWYQQNSGQAPVLVVGSDRPSGIPERFSGSNSGTTAT
LTISRVEAGDEADYYCQVWDSSSDHPGAVFGGGTQLTVL

>SC09-009 VH DNA (SEQ ID NO: 25)

CAGGTGCAGCTGGTGCATCTGGGGCTGAGGTCAAGAAGCCTGGGTCTCGGTGAAAGTCTCCTGCAAGTC
TTCTGGAGGCACCTCCAATAACTATGCTATCAGCTGGGTGCGACAGGCCCTGGACAAGGCCTTGACTGGA
TGGGCGGGATCAGCCCTATCTTTGGTTCGACAGCCTACGCACAGAAATTCAGGGCAGAGTCACTATTTCC
GCGGACATATTTTGAACACAGCCTACATGGAGCTGAACAGCCTGGCATCTGAGGACACGGCCGTATATTT
CTGTGCGAGGCACGGGAATTATTATTACTACTCCGGTATGGACGTCTGGGGCCAAGGGACCACGGTCACCG
TCTCGAGC

>SC09-009 VH PROTEIN (SEQ ID NO: 26)

QVQLVQSGAEVKKPGSSVKVSKSSGGTSNNYAIWVWRQAPGQGLDWMGGISPIFGSTAYAQKFQGRVTIS
ADIFSNAYMELNSLASEDTAVYFCARHGNYYYSGMDVWVGQGTTVTVSS

>SC09-009 VL DNA (SEQ ID NO: 27)

GACATCCAGATGACCCAGTCTCCATCTTCCGTGCTGCATCTGTAGGAGACAGAGTCACCATCACTTGTGCG
GGCGAGTCAGCATATTAGCAGTTGGTTAGCTGGTATCAGCAGAAGCCAGGGAAAGGCCCTCAGCTCCTGA
TCTATTCTGCATCCCGTTTGCAAAGTGGGGTCCCATCAAGGTTTCAGCGGCAGTGGATCTGGGACAGATTTCC
ACTCTCACCATCAGCAGCCTGCAGCCTGAAGATTTTGCAACTTACTATTGTCAACAGGCTAACAGTTTCCC
CCTCACTTTCCGGCCCTGGGACCAAAGTGGATATCAAAC

>SC09-009 VL PROTEIN (SEQ ID NO: 28)

DIQMTQSPSSVSASVGRVTITCRASQHISSWLAWYQQKPGKGPQLLIYSASRLQSGVPSRFSGSGSGTDF
TLTISSLQPEDFATYYCQANSPFLTFGPGTKVDIK

>SC09-010 VH DNA (SEQ ID NO: 29)

GAGGTGCAGCTGGTGGAGTCCGGGGCTGAGGTCAAGAAGCCTGGGTCTCGGTGAAAGTCTCCTGCAAGTC
TTCTGGAGGCACCTCCAATAATTATGCTATCAGCTGGGTGCGACAGGCCCTGGACAAGGCCTTGACTGGA
TGGGCGGGATCAGCCCTATCTTTGGTTCGACAGCCTACGCACAGAAGTTCCAGGGCAGAGTCACTATTTCC
GCGGACATATTTTCCAACACAGCCTACATGGAGCTGAACAGCCTGACATCTGAGGACACGGCCGTATATTA
CTGTGCGAGGCACGGGAATTATTATTACTACTCCGGTATGGACGTCTGGGGCCAAGGGACCACGGTCACCG
TCTCGAGC

>SC09-010 VH PROTEIN (SEQ ID NO: 30)
 EVQLVESGAEVKKPGSSVKVSKSSGGTSNNYAIISWVRQAPGQGLDWMGGISPIFGSTAYAQKFQGRVTIS
 ADIFSN TAYMELNSLTSEDTAVYYCARHGNYYYYS GMDVWGQGT TTVTVSS

>SC09-010 VL DNA (SEQ ID NO: 31)
 TCCTATGTGCTGACTCAGCCACCC TCGGTGTCAGTGGCCCCAGGACAGACGGCCAGGATTACCTGTGGGGG
 AAACAACATYGAAGTAAACTGTGCATTGGTACCAGCAGAACTCAGGCCAGGCCCTGTGCTGGTCGTCT
 TTGTTGATAGCGACCGTCCCTCAGGGATCCATGAGCGATTCTGTGGCTCCA ACTCTGGGTCCACGGCCACC
 CTGACCATCAGCAGCGTCAAGCCGGGGATGAGGCCGACTATTACTGT CAGGTGTGGGATAGTAATAGCGA
 TCATCCCGGTGCTGTGTTCCGAGGAGGCACCCAGCTGACCGTCTCTCG

>SC09-010 VL PROTEIN (SEQ ID NO: 32)
 SYVLTQPPSVSVAPGQTARITCGGNNIGSKTVHWYQQNSGQAPVLVVFVDSDRPSGIHERFCGSNSGSTAT
 LTISSVEAGDEADYYCQVWDSNSDHPGAVFVGGGTQLTVL

>SC09-011 VH DNA (SEQ ID NO: 33)
 GAGGTCCAGCTGGTACAGTCTGGGGCTGAGGTCAAGAAGCCTGGGTCTCGGTGAAGGTCTCCTGCAAGTC
 TTCTGGAGGCACCTCCAATAACTATGCTATCAGCTGGGTGCGGCAGGCCCTTGACAAGGCCTTGACTGGA
 TGGGAGGGATCAGCCCTATCTTTGGTTCAGCAGCTACGCACAGAAGTTCAGGGCAGAGTCACTATTACC
 GCGGACATATTTTCGAACACAGTGTACATGGAGCTGAACAGCCTGACATCTGAGGACACGGCCGTGTATTA
 CTGTGCGAGACACGGGAATTATTATTACTACTCCGGTACGGACGTCTGGGGCCAAGGGACCACGGTCACCG
 TCTCGAGC

>SC09-011 VH PROTEIN (SEQ ID NO: 34)
 EVQLVQSGAEVKKPGSSVKVSKSSGGTSNNYAIISWVRQAPGQGLDWMGGISPIFGSAAYAQKFQGRVTIT
 ADIFSN TVYME LNSLTSEDTAVYYCARHGNYYYYS GTDVWGQGT TTVTVSS

>SC09-011 VL DNA (SEQ ID NO: 35)
 TCCTATGTGCTGACTCAGCCACCCG CAGTGTCTGGGACCCCCGGGCAGAGGGTCACCATCTCGTGTCTGG
 AAGTGATTCCAACATCGGGAGAAGAAGTGTA ACTGGTACCAGCAGTCCCAGGAACGGCCCCAACTCC
 TCATCTATAGTAACGATCAGCGGCCCTCAGTGGTCCCTGACCGATTCTCTGGCTCCAAGTCCGGCACCTCA
 GCCTCCCTGGCCATCAGTGGGCTCCAGTCTGAAGATGAGGCCGAATATTACTGTGCAGCATGGGATGACAG
 CCTGAAGGGGGCTGTGTTCCGAGGAGGCACCCAGCTGACCGTCTCTCG

>SC09-011 VL PROTEIN (SEQ ID NO: 36)
 SYVLTQPPAVSGTPGQRVTISCSGSDSNIGRRSVN WYQFPGTAPKLLIYSNDQRPSVVPDRFSGSKSGTS
 ASLAISGLQSEDEAEYYCAA WDDSLKGAVFVGGGTQLTVL

>SC09-012 VH DNA (SEQ ID NO: 37)
 GAGGTCCAGCTGGTACAGTCTGGGGCTGAGGTCAAGAAGCCTGGGTCTCGGTGAAGGTCTCCTGCAAGTC
 TTCTGGAGGCACCTCCAATAATTATGCTATCAGCTGGGTGCGACAGGCCCTTGACAAGGCCTTGACTGGA
 TGGGAGGGATCAGCCCTATTTTTGGTTCAGCAGTCTACGCACAGAAGTTCAGGGCAGAGTCACTATTACC
 GCGGACATATTTTCGAACACAGTGTACATGGAGCTGAACAGCCTGACATCTGAGGACACGGCCGTGTATTA
 CTGTGCGAGACACGGGACTTATTATTACTACTCCGGTATGGACGTCTGGGGCCAAGGGACCACGGTCACCG
 TCTCGAGC

>SC09-012 VH PROTEIN (SEQ ID NO: 38)
 EVQLVQSGAEVKKPGSSVKVSKSSGGTSNNYAIISWVRQAPGQGLDWMGGISPIFGSAVYAQKFQGRVTIT
 ADIFSN TVYME LNSLTSEDTAVYYCARHGTYYYYS GMDVWGQGT TTVTVSS

>SC09-012 VL DNA (SEQ ID NO: 39)
 CAGTCTGTCGTGACGCAGCCGCCCTCAGTGTCTGGGGCCCCAGGGCAGAGGGTCACCATCTCCTGC ACTGG
 GAGCAGCTCCAACATCGGGGCAGGTTATGATGTACTACTGGTACCAGCAGCTTCCAGGGACAGCCCCAAAC
 TCCTCATCTATGGTAACAACAATCGGCCCTCAGGGGTCCCTGACCGATTCTCTGGCTCCAAGTCTGGCACC
 TCAGCCTCCCTGGCCATCACTGGGCTCCAGGTTGAGGATGAGGCTGATTATTACTGCCAGTCTTATGACCA
 GAACCTGAGTGAGGGGGTCTTCGGCGGAGGGACCAAGCTGACCGTCTCTAG

>SC09-012 VL PROTEIN (SEQ ID NO: 40)
QSVVTQPPSVSGAPGQRVTISCTGSSSNIGAGYDVHWYQQLPGTAPKLLIYGNNRPSGVPDRFSGSKSGT
SASLAITGLQVEDEADYYCQSYDQNLSEGVFGGGTKLTVL

>SC09-029 VH DNA (SEQ ID NO: 41)
GAGGTGCAGCTGGTGGAGTCCGGGGCTGAGGTCAAGAAGCCTGGGTCCCTCGGTGAAAGTCTCCTGCAAGTC
TTCTGGAGGCACCTCCAATAACTATGCTATCAGCTGGGTGCGACAGGCCCTTGACAAGGCCTTGACTGGA
TGGGCGGGATCAGCCCTATCTTTGGTTCGACAGCCTACGCACAGAAGTCCAGGGCAGAGTCACTATTTCC
GCGGACATATTTTGAACACAGCCTACATGGAGCTGAACAGCCTGACATCTGAGGACACGGCCGTATATTA
CTGTGCGAGGCACGGGAATTATTATTACTACTCCGGTATGGACGTCTGGGGCCAAGGGACCACGGTCACCG
TCTCGAGC

>SC09-029 VH PROTEIN (SEQ ID NO: 42)
EVQLVESGAEVKPKGSSVKVSKSSGGTSNNYAIWVRQAPGQGLDWMGGISPIFGSTAYAQKFQGRVTIS
ADIFSNAYMELNSLTSEDVAVYYCARHGNYYYSGMDVWVGQGTTVTVSS

>SC09-029 VL DNA (SEQ ID NO: 43)
GAAATTGTGATGACGCAGTCTCCAGGCACCTGTCTTTGTCTCCTGGGGAAAGAGGCACCCCTCTCCTGCAG
GGCCAGTCAGAGTGTTAGCAGCTACTTAGCCTGGTACCAACAGAAACCTGGCCAGGCTCCCAGGCTCCTCA
TCTATGGTGCATCCACCAGGGCCACTGGCATCCAGACAGGTTCACTGGCAGTGGGTCTGGGACAGACTTC
ACTCTCACCATCAGCAGACTGGAGCCTGAAGATTTGCAAGTGTATTACTGTGTCAGCAGTATGGGAGCTCACC
ATTCGCTTTCGGCCCTGGGACCAAGGTGGAGATCAAA

>SC09-029 VL PROTEIN (SEQ ID NO: 44)
EIVMTQSPGTLVSLSPGERGTLSCRASQSVSSYLAWYQQKPGQAPRLLIYGASTRATGIPDRFTGSGSGTDF
TLTISRLEPEDFAVYYCQQYGGSSPFAFGPGTKVEIK

>SC09-030 VH DNA (SEQ ID NO: 45)
CAGATGCAGCTGGTGCAGTCTGGGGCTGAGGTCAAGAAGCCTGGGTCCCTCGGTGAAAGTCTCCTGCAAGTC
TTCTGGAGGCACCTCCAATAACTATGCTATCAGCTGGGTGCGACAGGCCCTTGACAAGGCCTTGACTGGA
TGGGCGGGATCAGCCCTATCTTTGGTTCGACAGCCTACGCACAGAAGTCCAGGGCAGAGTCACTATTTCC
GCGGACATATTTTGAACACAGCCTACATGGAGCTGAACAGCCTGACATCTGAGGACACGGCCGTATATTT
CTGTGCGAGGCACGGGAATTATTATTACTACTCCGGTATGGACGTCTGGGGCCAAGGGACCACGGTCACCG
TCTCGAGC

>SC09-030 VH PROTEIN (SEQ ID NO: 46)
QMQLVQSGAEVKKPKGSSVKVSKSSGGTSNNYAIWVRQAPGQGLDWMGGISPIFGSTAYAQKFQGRVTIS
ADIFSNAYMELNSLTSEDVAVYFCARHGNYYYSGMDVWVGQGTTVTVSS

>SC09-030 VL DNA (SEQ ID NO: 47)
TCCTATGTGCTGACTCAGCCACCTCGGTGTCAGTGGCCCCAGGACAGACGGCCAGGATTACCTGTGGGGG
AAACAACATTGGAAGTAAAAGTGTGCACTGGTACCAGCAGAAGCCAGGCCAGGCCCTGTGCTGGTCTGCT
ATGGTGTAGCGACCGGCCCTCAGGGATCCCTGAGCGATTCTCTGGCTCCAACCTCTGGGACCACGGCCACC
CTGACCATCAGCAGGTCGAAGCCGGGGATGAGGCCGACTATTACTGTGTCAGGTGTGGGATAGTAGTAGTGA
TCATCCCGGTGCTGTGTTCCGGAGGAGGCACCCAGCTGACCGTCTCTCG

>SC09-030 VL PROTEIN (SEQ ID NO: 48)
SYVLTQPPSVSVAPGQTARITCGNNGSKSVHWYQQKPGQAPVLLVYGDSDRPSGIPERFSGSNSGTTAT
LTISRVEAGDEADYYCQVWDSDDHPGAVFGGGTQLTVL

>SC09-031 VH DNA (SEQ ID NO: 49)
CAGGTCCAGCTGGTACAGTCTGGGGCTGAGGTGAGAGGCCTGGGTCCCTCGGTGAAAGTCTCCTGCAAGTC
TTCTGGCGGCACCTCCAATAACTATGCCATCAGCTGGGTGCGACAGGCCCTTGACAAGGCCTTGACTGGA
TGGGCGGGATCAGCCCTATCTTTGGTTCGACAGCCTACGCACAGAAGTCCAGGGCAGAGTCACTATTTCC
GCGGACATATTTTGAACACAGCCTACATGGAGCTGAACAGTCTGACATCTGAGGACACGGCCGTCTATTA

TTGTGCGAGACACGGGAATTATTATTACAACCTCCGGTATGGACGTCTGGGGCCAAGGGACCACGGTCACCG
TCTCGAGC

>SC09-031 VH PROTEIN (SEQ ID NO: 50)
QVQLVQSGAEVERPGSSVKVSKSSGGTSNNYAI SWVRQAPGQGLDWMGGISPIFGSTAYAQKFQGRVTIS
ADIFSN TAYMELNSLTSEDTAVYYCARHGNYYYNSGMDVWVGQGT TTVTVSS

>SC09-031 VL DNA (SEQ ID NO: 51)
CAGTCTGTGTTGACGCAGCCGCCCTCAGTGTCTGGGGCCCCAGGGCAGAGGGTCACCATCTCCTGCAC TGG
GAGCAGCTCCAACATCGGGGCAGGTTATGATGTACTGTTACCAGCAGCTTCCAGAAACAGCCCCCAAAC
TCCTCATTTATGATAACAACAATCGTCCCTCAGGGGTTTCTGACCGATTCTCTGGCTCCAAGTC TGGCACT
TCAGCCTCCCTGGCCATCAC TGGGCTCCAGGCTGAGGATGAGGCTGATTATTACTGCCAGTCC TATGACAG
CGGCCTGAGTGCTTCGCCTTATGTCTTCGGAGCTGGGACCAAGGTCACCGTCCTAG

>SC09-031 VL PROTEIN (SEQ ID NO: 52)
QSVLTQPPSVSGAPGQRVTISCTGSSNIGAGYDVHWYQQLPETAPKLLIYDNNRPSGVSDRFSGSKSGT
SASLAITGLQAEDEADYYCQSYDSGLSASPYVFGAGTKVTVL

>SC09-112 VH DNA (SEQ ID NO: 53)
CAGGTGCAGCTGGTGCAGTCTGGGGCTGAGGTCAAGAAGCCTGGGTCTCGGTGAAAGTCTCCTGCAAGTC
TTCTGGAGGCACCTCCAATAACTATGCTATCAGCTGGGTGCGACAGGCCCTTGACAAGGCCTTGACTGGA
TGGGCGGGATCAGCCCTATCTTTGGTTCGACAGCTACGCACAGAAGTCCAGGGCAGAGTCACTATTTCC
GCGGACATATTTTCAACACAGCCTACATGGAGCTGAACAGCCTGACATCTGAGGACACGGCCGTATATTA
CTGTGCGAGGCACGGGAATTATTATTACTACTCCGGTATGGACGTCTGGGGCCAAGGGACCACGGTCACCG
TCTCGAGC

>SC09-112 VH PROTEIN (SEQ ID NO: 54)
QVQLVQSGAEVKKPGSSVKVSKSSGGTSNNYAI SWVRQAPGQGLDWMGGISPIFGSTAYAQKFQGRVTIS
ADIFSN TAYMELNSLTSEDTAVYYCARHGNYYYNSGMDVWVGQGT TTVTVSS

>SC09-112 VL DNA (SEQ ID NO: 55)
CAGTCTGTGTTGACGCAGCCGCCCTCAGTGTCTGGGGCCCCAGGGCAGAGGGTCACCATCTCCTGCAC TGG
GAGCAGCGCCAACATCGGGGCAGGTTATGATGTCCACTGGTACCAGCAGTTTCCAGGAACAGCCCCCAAAC
TCCTCATCTATGGTAACAACAATCGGCCCTCAGGGGTCCTGACCGATTCTCTGGCTCCAAGTC TGGCACC
TCAGCCTCCCTGGCCATCAC TGGGCTCCAGGCTGAGGATGAGGCTGATTATTACTGCCAGTCC TATGACAG
CAGCCTGAGTGGTGCCTTATTCGGCGAGGGACCAAGCTGACCGTCCTAG

>SC09-112 VL PROTEIN (SEQ ID NO: 56)
QSVLTQPPSVSGAPGQRVTISCTGSSANIGAGYDVHWYQQFPGTAPKLLIYGNRPSGV PDRFSGSKSGT
SASLAITGLQAEDEADYYCQSYDSSLGALFGGGTKLTVL

>SC09-113 VH DNA (SEQ ID NO: 57)
CAGATGCAGCTGGTGCAGTCTGGGGCTGAGGTCAAGAAGGCTGGGTCTCGGTGAAAGTCTCCTGCAAGTC
TTCTGGAGGCACCTCCAATAACTATGCTATCAGCTGGGTGCGACAGGCCCTTGACAAGGCCTTGAGTGGA
TGGGCGGGATCAGTCCAATCTTTGGTTCGACAGTCTACGCACAGAAATTCAGGGCAGAGTCACTATTTCC
GCGGACATATTTTACACACTGCC TACATGGAGCTGAACAGCCTGACATCTGAGGACACGGCCGCATATTT
CTGTGCGAGGCACGGAACTATTATTACTACTCCGGTATGGACCTCTGGGGCCAAGGGACCACGGTCACCG
TCTCGAGC

>SC09-113 VH PROTEIN (SEQ ID NO: 58)
QMQLVQSGAEVKKAGSSVKVSKSSGGTSNNYAI SWVRQAPGQGLEWMGGISPIFGSTVYAQKFQGRVTIS
ADIFSH TAYMELNSLTSEDTAAYFCARHGNYYYNSGMDLWVGQGT TTVTVSS

>SC09-113 VL DNA (SEQ ID NO: 59)
CAGTCTGTGCTGACTCAGCCACCCGAGTGTCTGGGACCCCCGGGCAGAGGGTCACCATCTCGTGTCTGG
AAGTGATTCCAACATCGGGAGAAGAAGTG TAAACTGGTACCAGCAGTTCCCAGGAACGGCCCCCAAAC TCC
TCATCTATAGTAACGATCAGCGGCCCTCAGTGGTCCCTGACCGATTCTCTGGCTCCAAGTCCGGCACCTCA

GCCTCCCTGGCCATCAGTGGGCTCCAGGCTGAGGATGAGGCTGATTATTACTGTGCAGCATGGGATGCCAG
CCTGAGTGGTCTGTGTTCGGAGGAGGCACCCAGCTGACCGTCTCTCG

>SC09-113 VL PROTEIN (SEQ ID NO: 60)
QSVLTQPPAVSGTPGQRVTISCSGSDSNIGRRSVNWWYQQFPGTAPKLLIYSNDQRPSVVPDRFSGSKSGTS
ASLAISGLQAEDEADYYCAAWDASLSGPVFGGGTQLTVL

>SC09-114 VH DNA (SEQ ID NO: 61)
CAGGTGCAGCTGGTGCAATCTGGGGCTGAGGTCAAGAAGCCTGGGTCTCGGTGAAAGTCTCCTGCAAGTC
TTCTGGAGGCACCTCCAATAACTATGCTATCAGCTGGGTGCGACAGGCCCTTGACAAGGCCTTGACTGGA
TGGGCGGGATCAGCCCTATCTTTGGTTCGACAGCCTACGCACAGAAATCCAGGGCAGAGTCACTATTTCC
GCGGACATATTTTGAACACAGCCTACATGGAGCTGAACAGCCTGACATCTGAGGACACGGCCGTATATTT
CTGTGCGAGGCACGGGAATTATTATTACTACTCCGGTATGGACGTCTGGGGCCAAGGGACCACGGTACCCG
TCTCGAGC

>SC09-114 VH PROTEIN (SEQ ID NO: 62)
QVQLVQSGAEVKKPGSSVKVSKSSGGTSNNYAIISWVRQAPGQGLDWMGGISPIFGSTAYAQKFQGRVTIS
ADIFSN TAYMELNSLTSED TAVYFCARHGNYYYSGMDVWVGQGT TTVTVSS

>SC09-114 VL DNA (SEQ ID NO: 63)
TCCTATGTGCTGACTCAGCCACCCGAGTGTCTGGGACCCCCGGGCAGAGGGTCACCATCTCGTGTCTTGG
AAGTGATTCCAACATCGGGAGAAGAAGTGTAACCTGGTACCAGCAGTCCCAGGAACGGCCCCAAACTCC
TCATCTATAGTAACGATCAGCGCCCTCAGTGGTCCCTGACCGATTCTCTGGCTCCAAGTCCGGCACCTCA
GCCTCCCTGGCCATCAGTGGGCTCCAGTCTGAAGATGAGGCCGAATATTACTGTGCAGCATGGGATGACAG
CCTGAAGGGGGCTGTGTTCGGAGGAGGCACCCAGCTGACCGTCTCTCG

>SC09-114 VL PROTEIN (SEQ ID NO: 64)
SYVLTQPPAVSGTPGQRVTISCSGSDSNIGRRSVNWWYQQFPGTAPKLLIYSNDQRPSVVPDRFSGSKSGTS
ASLAISGLQSEDEAEYYCAAWDDSLKGAVFGGGTQLTVL

Vector pIg-C911-HCgamma1 (SEQ ID NO:175)

tgcagcgatc gggagatctc ccgatcccct atggtgcact ctcagtaca tctgctctga 60
tgccgcatag ttaagccagt atctgctccc tgcttgtgtg ttggagggtcg ctgagtagtg 120
cgcgagcaaa atttaagcta caacaaggca aggcttgacc gacaattgca tgaagaatct 180
gcttaggggtt aggcgttttg cgctgcttcg ctagggtggtc aatattggcc attagccata 240
ttattcattg gttatatagc ataaatcaat attggctatt ggccattgca tacgttgtat 300
ccatatcata atatgtacat ttatatattggc tcatgtccaa cattaccgcc atgttgacat 360
tgattattga ctagttatta atagtaatca attacggggt cattagttca tagcccatat 420
atggagttcc gcgttacata acttacggta aatggcccgc ctggctgacc gcccacacgac 480
ccccgccc at tgacgtcaat aatgacgtat gttccc atag taacgccaat agggactttc 540
cattgacgtc aatgggtgga gtatttacgg taaactgccc acttggcagt acatcaagtg 600
tatcatatgc caagtacgcc ccctattgac gtcaatgacg gtaa atggcc cgctggcat 660
tatgcccagt acatgacctt atgggacttt cctacttggc agtacatcta cgtattagtc 720
atcgctatta ccatggtgat gcggttttg cagtacatca atgggcgtgg atagcggttt 780
gactcacggg gatttccaag tctccacccc attgacgtca atgggagttt gttttggcac 840
caaaatcaac gggactttcc aaaatgtcgt aacaactccg cccattgac gcaa atgggc 900
ggtaggcgtg tacggtggga ggtctatata agcagagctc gtttagtgaa ccgtcagatc 960
gcctggagac gccatccacg ctgttttgac ctccatagaa gacaccggga ccgatccagc 1020
ctccgcggcc gggaacggtg cattggaagc tggcctggat atcctgactc tcttaggtag 1080
ccttgacagaa gttggtcgtg aggcactggg caggtaagta tcaaggttac aagacagggt 1140
taaggagatc aatagaaact gggcttgtcg agacagagaa gactccttgcg tttctgatag 1200
gcacctattg gtcttactga catccacttt gcctttctct ccacagggtg ccactcccag 1260
ttcaattaca gctcgcacc atgggatgga gctgtatcat cctcttcttg gtactgctgc 1320
tggcccagcc ggccagtac cttgaccggt gcaccacttt tgatgatggt caagctccta 1380
attacactca acatacttca tctatgaggg gggtttacta tctctgatgaa attttttagat 1440
cggacactct ttatttaact caggatttat ttcttccatt ttatttctaat gttacagggt 1500

ttcatactat	taatcatacg	tttggcaacc	ctgtcatacc	ttttaaggat	ggatatttatt	1560
ttgctgccac	agagaaatca	aatggtgtcc	gtgggtgggt	ttttggttct	accatgaaca	1620
acaagtcaca	gtcggtgatt	attattaaca	attctactaa	tgttgttata	cgagcatgta	1680
actttgaatt	gtgtgacaac	cctttccttg	ctgtttctaa	acccatgggt	acacagacac	1740
atactatgat	attcgataat	gcatttaatt	gcactttcga	gtacatatct	gatgcctttt	1800
cgcttgatgt	ttcagaaaag	tcaggtaatt	ttaaactctt	acgagagttt	gtgtttaaaa	1860
ataaagatgg	gtttctctat	gtttataagg	gctatcaacc	tatagatgta	gttcgtgatc	1920
taccttctgg	ttttaaacct	ttgaaaccta	tttttaagtt	gcctcttggg	attaacatta	1980
caaatttttag	agccattctt	acagcctttt	cacctgctca	agacatttgg	ggcacgtcag	2040
ctgcagccta	ttttgttggc	tatttaaagc	caactacatt	tatgctcaag	tatgatgaaa	2100
atggtacaat	cacagatgct	gttgattggt	ctcaaaatcc	acttgctgaa	ctcaaatgct	2160
ctgttaagag	ctttgagatt	gacaaaaggaa	tttaccagac	ctctaatttc	aggggtgttc	2220
cctcaggaga	tgttgtgaga	ttccctaata	ttacaaactt	gtgtcctttt	ggagaggttt	2280
ttaatgctac	taaattccct	tctgtctatg	catgggagag	aaaaaaaaatt	tctaattgtg	2340
ttgctgatta	ctctgtgctc	tacaactcaa	catttttttc	aacctttaag	tgctatggcg	2400
tttctgccac	taagttgaat	gatcctttgct	tctccaatgt	ctatgcagat	tctttttag	2460
tcaagggaga	tgatgtaaga	caaatagcgc	caggacaaac	tggtgttatt	gctgattata	2520
attataaatt	gccagatgat	ttcatgggtt	gtgtccttgc	ttggaatact	aggaacattg	2580
atgctacttc	aactggtaat	tataattata	aatataggta	tcttagacat	ggcaagctta	2640
ggccctttga	gagagacata	tctaattgtc	ctttctcccc	tgatggcaaa	ccttgcacc	2700
cacctgctct	taattgttat	tggccattaa	atgattatgg	tttttacacc	actactggca	2760
ttggctacca	acettacaga	gttgtagtac	tttcttttga	acttttaaat	gcaccggcca	2820
cggtttgtgg	acaaaaatta	tccactgacc	ttattaagaa	ccagtgtgtc	aatttttaatt	2880
ttaatggact	cactggtaact	ggtgtgttaa	ctccttcttc	aaagagattt	caaccatttc	2940
aacaatttgg	cggtgatggt	tctgatttca	ctgattccgt	tcgagatcct	aaaacatctg	3000
aaatattaga	catttcacct	tgctcttttg	ggggtgtaag	tgtaattaca	cctggaacaa	3060
atgcttcate	tgaagttgct	gttctatate	aagatgttaa	ctgcactgat	gtttctacag	3120
caattcatgc	agatcaactc	acaccagctt	ggcgcataata	ttctactgga	aacaatgtat	3180
tccagactca	ggcaggctgt	cttataggag	ctgagcatgt	cgacacttct	tatgagtgcg	3240
acattcctat	tgtagctggc	atttgtgcta	gttaccatac	agtttcttta	ttacgtagta	3300
ctagccaaaa	atctattgtg	gcttatacta	tgtcttttagg	tgctgatagt	tcaattgctt	3360
actctaataa	caccattgct	atacctaata	acttttcaat	tagcattact	acagaagtaa	3420
tgctgttttc	tatggctaaa	acctccgtag	attgtaatat	gtacatctgc	ggagattcta	3480
ctgaatgtgc	taatttgctt	ctccaatatg	gtagcttttg	cacacaacta	aatcgtgcac	3540
tctcaggat	tgctgctgaa	caggatcgca	acacacgtga	agtgttcgct	caagtcaaac	3600
aaatgtacaa	aacccaact	ttgaaatatt	ttggtggttt	taatttttca	caaataattac	3660
ctgaccctct	aaagccaact	aagaggtctt	ttattgagga	cttgcctctt	aataaggtga	3720
cactcgctga	tgctggcttc	atgaagcaat	atggcgaatg	cctaggtgat	attaatgcta	3780
gagatctcat	ttgtgcgcag	aagttcaatg	gacttacagt	gttgccacct	ctgctcactg	3840
atgatatgat	tgctgcctac	actgctgctc	tagttagtgg	tactgccact	gctggatgga	3900
catttggtgc	tggcgctgct	cttcaaatac	cttttgctat	gcaaattggca	tataggttca	3960
atggcattgg	agttacceaa	aatgttctct	atgagaacca	aaaacaaatc	gccaaccaat	4020
ttaacaaggc	gattagtcaa	attcaagaat	cacttacaac	aacatcaact	gcattgggca	4080
agctgcaaga	cgttgttaac	cagaatgctc	aagcattaaa	cacacttggt	aaacaactta	4140
gctctaattt	tggtgcaatt	tcaagtgtgc	taaatgatata	cctttcgcga	cttgataaag	4200
tcgagggcga	ggtacaaatt	gacaggttaa	ttacaggcag	acttcaaagc	cttcaaacct	4260
atgtaacaca	acaactaatc	agggtctgtg	aaatcagggc	ttctgctaat	cttgcctgcta	4320
ctaaaatgtc	tgagtgtggt	cttggacaat	caaaaagagt	tgacttttgt	ggaaagggct	4380
accaccttat	gtccttccca	caagcagccc	cgcatgggtg	tgtcttccca	catgtcacgt	4440
atgtgccatc	ccaggagagg	aacttcacca	cagcgcagc	aatttgcct	gaaggcaaac	4500
catacttccc	tcgtgaaggt	gtttttgtgt	ttaatggcac	ttcttggttt	attacacaga	4560
ggaacttctt	ttctccacaa	ataattacta	cagacaatac	atltgtctca	ggaaattgtg	4620
atgtcgttat	tggtcatcatt	aacaacacag	tttatgatcc	tctgcaacct	gagcttgact	4680
cattcaaaga	agagctggac	aagtacttca	aaaatcatac	atcaccagat	gttgattttg	4740
gcgacatctc	aggcattaac	gcttctgtcg	tcaacattca	aaaagaaatt	gaccgcctca	4800
atgaggtcgc	taaaaattta	aatgaatcac	tcattgacct	tcaagaactg	ggaaaatatg	4860
agcaatatat	taaatggcct	ctcgcagcaac	aaaaactcat	ctcagaagag	gatctgaatg	4920

ctgtgggcca	ggacacgcag	gaggtcatcg	tgggtgccaca	ctccttgccc	tttaaggtgg	4980
tgggtgatctc	agccatcctg	gccctggtgg	tgctcaccat	catctccctt	atcatcctca	5040
tcatgctttg	gcagaagaag	ccacgttagg	cggccgctcg	agtgctagca	ccaagggccc	5100
cagcgtgttc	cccctggccc	ccagcagcaa	gagcaccagc	ggcggcacag	ccgccctggg	5160
ctgcctggtg	aaggactact	tccccgagcc	cgtgaccgtg	agctggaaca	gcggcgcctt	5220
gaccagcggc	gtgcacacct	tccccgcctg	gctgcagagc	agcggcctgt	acagcctgag	5280
cagcgtggtg	accgtgcccc	gcagcagcct	gggcacccag	acctacatct	gcaacgtgaa	5340
ccacaagccc	agcaacacca	aggtggacaa	acgcgtggag	cccaagagct	gcgacaagac	5400
ccacacctgc	ccccctgccc	ctgccccga	gctgctgggc	ggaccctccg	tgttctctgtt	5460
cccccccaag	cccaaggaca	ccctcatgat	cagccggacc	cccgaggtga	cctgcgtggt	5520
ggtggacgtg	agccacgagg	accccagagt	gaagttcaac	tggtagctgg	acggcgtgga	5580
ggtgcacaac	gccaagacca	agccccggga	ggagcagtac	aacagcacct	accgggtggt	5640
gagcgtgctc	accgtgctgc	accaggactg	gctgaacggc	aaggagtaca	agtgcaaggt	5700
gagcaacaag	gcctgctctg	cccccatcga	gaagaccatc	agcaaggcca	agggccagcc	5760
ccgggagccc	caggtgtaca	ccctgcccc	cagccgggag	gagatgacca	agaaccaggt	5820
gtccctcacc	tgtctggtga	agggttcta	cccacgcgac	atcgccgtgg	agtgggagag	5880
caacggccag	cccgagaaca	actacaagac	cacccccct	gtgctggaca	gcgacggcag	5940
cttcttctctg	tacagcaagc	tcaccgtgga	caagagccgg	tggcagcagg	gcaacgtggt	6000
cagctgcagc	gtgatgcacg	aggccctgca	caaccactac	accagaaga	gcctgagcct	6060
gagccccggc	aagtgataat	ctagagggcc	cgtttaaacc	cgctgatcag	cctcgactgt	6120
gccttctagt	tgccagccat	ctggtgtttg	cccctcccc	gtgccttctt	tgaccctgga	6180
aggtgccaact	ccaactgtcc	tttctaata	aatgaggaa	attgcatcgc	attgtctgag	6240
taggtgtcat	tctattctgg	ggggtgggg	ggggcaggac	agcaaggggg	aggattggga	6300
agacaatagc	aggcatgctg	gggatgctgt	gggctctatg	gcttctgagg	cggaaagaac	6360
cagctggggc	tctaggggg	atccccacgc	gccctgtagc	ggcgcattaa	gcgcgccggg	6420
tgtggtggtt	acgcgcagcg	tgaccgctac	acttgccagc	gccctagcgc	ccgctccttt	6480
cgctttcttc	ccttccttc	tcgccacgtt	cgccgcttt	ccccgtcaag	ctctaaatcg	6540
ggggctccct	ttagggttcc	gatttagtgc	tttacggcac	ctcgacccca	aaaaacttga	6600
ttaggggtgat	ggttcacgta	gtgggccatc	gccctgatag	acggtttttc	gccctttgac	6660
gttggagtc	acgttcttta	atagtggact	cttgttccaa	actggaacaa	cactcaacc	6720
tatctcggtc	tattcttttg	atttataagg	gattttgcog	atctcggcct	attggttaaa	6780
aatgagctg	atttaacaaa	aatttaacgc	gaattaattc	tgtggaatgt	gtgtcagtta	6840
gggtgtggaa	agtccccagg	ctccccagca	ggcagaagta	tgcaaagcat	gcattctcaat	6900
tagtcagcaa	ccaggtgtgg	aaagtcccca	ggctccccag	caggcagaag	tatgcaaagc	6960
atgcatctca	attagtcagc	aaccatagtc	ccgcccctaa	ctccgcccct	cccgcacct	7020
actccgccc	gttccgccc	ttctccgccc	catggctgac	taatTTTTTT	tatttatgca	7080
gaggccgagg	cgccctctgc	ctctgagcta	ttccagaagt	agtgaggagg	ctTTTTTTgga	7140
ggcctaggct	tttgcaaaaa	gctcccggga	gcttgtatat	ccatTTTtcgg	atctgatcaa	7200
gagacaggat	gaggatcgtt	tcgcatgatt	gaacaagatg	gattgcacgc	aggttctccg	7260
gccgcttggg	tggagaggct	attcggctat	gactgggcac	aacagacaat	cggctgctct	7320
gatgccgccc	tgttccggct	gtcagcgcag	gggcgcccgg	ttctTTTTTgt	caagaccgac	7380
ctgtccgggtg	ccctgaatga	actgcaggac	gaggcagcgc	ggctatcgtg	gctggccacg	7440
acgggcgttc	cttgcgcagc	tgtgctcgac	gttgtcactg	aagcgggaag	ggactggctg	7500
ctattggggcg	aagtgccggg	gcaggatctc	ctgtcatctc	accttgctcc	tgccgagaaa	7560
gtatccatca	tggctgatgc	aatgcggcgg	ctgcatacgc	ttgatccggc	tacctgccc	7620
ttcgaccacc	aagcgaacaa	tcgcatcgag	cgagcacgta	ctcggatgga	agccggctct	7680
gtcgatcagg	atgatctgga	cgaagagcat	caggggctcg	cgccagccga	actgttccgc	7740
aggctcaagg	cgcgcatgcc	cgacggcgag	gatctcgtcg	tgacccatgg	cgatgacctgc	7800
ttgccgaata	tcatggtgga	aaatggccgc	ttttctggat	tcatcgactg	tggccggctg	7860
ggtgtggcgg	accgctatca	ggacatagcg	ttggctacct	gtgatattgc	tgaagagctt	7920
ggcggcgaat	gggctgaccg	cttctcgtg	ctttacggta	tcgccgctcc	cgattcgcag	7980
cgcctcgcct	tctatcgcct	tcttgacgag	ttcttctgag	cgggactctg	gggttcgaaa	8040
tgaccgacca	agcgacgccc	aacctgccat	cacgagattt	cgattccacc	gccgccttct	8100
atgaaaggtt	gggcttcgga	atcgTTTTcc	gggacgcccg	ctggatgate	ctccagcgcg	8160
gggatctcat	gctggagttc	ttcgcccacc	ccaacttgtt	tattgcagct	tataatggtt	8220
acaaataaag	caatagcatc	acaaatttca	caaataaagc	atTTTTTTca	ctgcattcta	8280
gttgtggttt	gtccaaactc	atcaatgtat	cttatcatgt	ctgtataaccg	tcgacctcta	8340

gctagagctt	ggcgtaatca	tggatcatagc	tgtttcctgt	gtgaaattgt	tatccgctca	8400
caattccaca	caacatacga	gccggaagca	taaagtgtaa	agcctggggg	gcctaattgag	8460
tgagctaact	caattaatt	gcgttgcgct	cactgcccgc	tttccagtcg	ggaaacctgt	8520
cgtgccagct	gcattaatga	atcggccaac	gcgcggggag	aggcggtttg	cgtattgggc	8580
gctcttccgc	ttctcgcctc	actgactcgc	tgcgctcggg	cgttcggctg	cggcgagcgg	8640
tatcagctca	ctcaaaggcg	gtaatacggg	tatccacaga	atcaggggat	aacgcaggaa	8700
agaacatgtg	agcaaaaggc	cagcaaaagg	ccaggaaccg	taaaaaggcc	gcggttgcgg	8760
cgTTTTTcca	taggctccgc	ccccctgacg	agcatcacia	aaatcgacgc	tcaagtcaga	8820
ggtggcgaaa	cccgcacagga	ctataaagat	accaggcggt	tccccctgga	agctccctcg	8880
tgcgctctcc	tgttccgacc	ctgcccgtta	ccggatacct	gtccgccttt	ctcccttcgg	8940
gaagcgtggc	gctttctcat	agctcacgct	gtaggtatct	cagttcgggtg	taggtcgttc	9000
gctccaagct	gggctgtgtg	cacgaacccc	ccgttcagcc	cgaccgctgc	gccttatccg	9060
gtaactatcg	tcttgagtcc	aacccggtaa	gacacgactt	atcgccactg	gcagcagcca	9120
ctggtaacag	gattagcaga	gcgaggtatg	taggcgggtg	tacagagttc	ttgaagtggg	9180
ggcctaacta	cggctacact	agaagaacag	tatttgggat	ctgcgctctg	ctgaagccag	9240
ttaccttcgg	aaaaagagtt	ggtagctctt	gatccggcaa	acaaaccacc	gctggtagcg	9300
gtTTTTTtgt	ttgcaagcag	cagattacgc	gcagaaaaaa	aggatctcaa	gaagatcctt	9360
tgatcttttc	tacggggctc	gacgctcagt	ggaacgaaaa	ctcacgtaa	gggattttgg	9420
tcatgagatt	atcaaaaagg	atcttcacct	agatcctttt	aaattaaaaa	tgaagtttta	9480
aatcaatcta	aagtataat	gagtaaactt	ggtctgacag	ttaccaatgc	ttaatcagtg	9540
aggcacctat	ctcagcgatc	tgtctatttc	gttcatccat	agttgcctga	ctccccgtcg	9600
tgtagataac	tacgatacgg	gagggcttac	catctggccc	cagtgtctga	atgataccgc	9660
gagacccacg	ctcaccggct	ccagatttat	cagcaataaa	ccagccagcc	ggaagggccg	9720
agcgcagaag	tggctctgca	actttatccg	cctccatcca	gtctattaat	tgttgccggg	9780
aagctagagt	aagtagttcg	ccagttaata	gtttgcgcaa	cgttgttgcc	attgctacag	9840
gcacgctggg	gtcacgctcg	tcgtttggtg	tggcttcatt	cagctccggg	tcccacgat	9900
caaggcgagt	tacatgatcc	cccattgtgt	gcaaaaaagc	ggttagctcc	ttcggctctc	9960
cgatcgttgt	cagaagtaag	ttggccgcag	tgttatcact	catggttatg	gcagcactgc	10020
ataattctct	tactgtcatg	ccatccgtaa	gatgcttttc	tgtgactggg	gagtactcaa	10080
ccaagtcatt	ctgagaatag	tgtatgcggc	gaccgagttg	ctcttgcccg	gcgtcaatac	10140
gggataatac	cgcgccacat	agcagaactt	taaaagtgtc	catcattgga	aaacgttctt	10200
cggggcgaaa	actctcaagg	atcttaccgc	tgttgagatc	cagttcagtg	taaccctctc	10260
gtgcacccaa	ctgatcttca	gcacttttta	ctttcaccag	cgtttctggg	tgagcaaaaa	10320
caggaaggca	aaatgccgca	aaaaagggaa	taagggcgac	acggaaatgt	tgaatactca	10380
tactcttctc	ttttcaatat	tattgaagca	tttatcaggg	ttattgtctc	atgagcggat	10440
acatatttga	atgtattttag	aaaaataaac	aaataggggt	tccgcgcaca	tttccccgaa	10500
aagtgccacc	tgacg					10515

Vector pIg-C909-Ckappa (SEQ ID NO:176)

tcgacggatc	gggagatctc	ccgatcccc	atggtgcact	ctcagtacaa	tctgctctga	60
tgccgcatag	ttaagccagt	atctgctccc	tgcttgtgtg	ttggagggtcg	ctgagtagtg	120
cgcgagcaaa	atttaagcta	caacaaggca	aggcttgacc	gacaattggt	aattaacatg	180
aagaatctgc	ttagggttag	gcgttttgcg	ctgcttcgct	agggtggtaaa	tattggccat	240
tagccatatt	attcattggg	tatatagcat	aaatcaatat	tggctatttg	ccattgcata	300
cgttgtatcc	atatcataat	atgtacattt	atattggctc	atgtccaaca	ttaccgccat	360
gttgacattg	attattgact	agttattaat	agtaatcaat	tacgggggtca	ttagttcata	420
gccccatata	ggagttccgc	gttacataac	ttacggtaaa	tggcccgcct	ggctgaccgc	480
ccaacgacct	ccgcccattg	acgtcaataa	tgacgtatgt	tcccatagta	acgccaatag	540
ggactttcca	ttgacgtcaa	tgggtggagt	atttacggta	aaactgccac	ttggcagtac	600
atcaagtgtg	tcatatgccg	agtacgccc	ctattgacgt	caatgacggg	aaatggcccg	660
cctggcatta	tgcccagtac	atgaccttat	gggactttcc	tacttggcag	tacatctacg	720
tattagtcac	cgctattacc	atgggtgatg	ggttttggca	gtacatcaat	gggctgggat	780
agcggtttga	ctcacgggga	tttccaagtc	tccaccccat	tgacgtcaat	gggagtttgt	840
tttggcacca	aaatcaacgg	gactttccaa	aatgtcgtaa	caactccgcc	ccattgacgc	900
aaatgggctg	taggctgtga	cggtgggagg	tctatataag	cagagctcgt	ttagtgaacc	960
gtcagatcgc	ctggagacgc	catccacgct	gttttgacct	ccatagaaga	caccgggacc	1020

gatccagcct	ccgcggccgg	gaacggtgca	ttggaatcga	tgactctctt	aggtagcctt	1080
gcagaagttg	gtcgtgaggc	actgggcagg	taagtatcaa	ggttacaaga	caggtttaag	1140
gagatcaata	gaaactgggc	ttgtcgagac	agagaagact	cttgcgtttc	tgataggcac	1200
ctattgggtct	tactgacatc	cactttgcct	ttctctccac	agggtgccac	tcccagttca	1260
attacagctc	gccaccatgc	ggctgcccgc	ccagctgctg	ggccttctca	tgctgtgggt	1320
gccccgctcg	agatctatcg	atgcatgcca	tggtaccaag	cttgccacca	tgagcagcag	1380
ctcttggctg	ctgctgagcc	tggtggccgt	gacagccgcc	cagagcacca	tcgaggagca	1440
ggccaagacc	ttcctggaca	agttcaacca	cgaggccgag	gacctgttct	accagagcag	1500
cctggccagc	tggaactaca	acaccaacat	caccgaggag	aacgtgcaga	acatgaacaa	1560
cgccggcgac	aagtggagcg	ccttcctgaa	ggagcagagc	acactggccc	agatgtacc	1620
cctgcaggag	atccagaacc	tgaccgtgaa	gctgcagctg	caggccctgc	agcagaacgg	1680
cagcagcgtg	ctgagcaggg	acaagagcaa	gcggtgaaac	accatcctga	acaccatgtc	1740
caccatctac	agcaccggca	aagtgtgcaa	ccccgacaac	ccccaggagt	gcctgctgct	1800
ggagcccggc	ctgaacgaga	tcatggccaa	cagcctggac	tacaacgagc	ggctgtgggc	1860
ctgggagagc	tggcggagcg	aagtgggcaa	gcagctgcgg	cccctgtacg	aggagtacgt	1920
ggtgctgaag	aacgagatgg	ccagggccaa	ccactacgag	gactacggcg	actactggag	1980
aggcgactac	gaagtgaacg	gcgtggacgg	ctacgactac	agcagaggcc	agctgatcga	2040
ggacgtggag	cacaccttcg	aggagatcaa	gcctctgtac	gagcacctgc	acgcctacgt	2100
gcgggccaag	ctgatgaacg	cctaccccag	ctacatcagc	cccatcggct	gcctgcccgc	2160
ccacctgctg	ggcgacatgt	ggggccgggt	ctggaccaac	ctgtacagcc	tgaccgtgcc	2220
cttcggccag	aagcccaaca	tcgacgtgac	cgacgccatg	gtggaccagg	cctgggacgc	2280
ccagcggatc	ttcaaggagg	ccgagaagtt	cttcgtgagc	gtgggcctgc	ccaacatgac	2340
ccagggcctt	tgggagaaca	gcatgctgac	cgaccccggc	aatgtgcaga	aggccgtgtg	2400
ccacccacc	gectgggacc	tgggcaaggg	cgacttccgg	atcctgatgt	gcaccaaagt	2460
gaccatggac	gacttctctga	ccgcccacca	cgagatgggc	cacatccagt	acgacatggc	2520
ctacgccgcc	cagcccttcc	tgtctcggaa	cggcgccaac	gagggctttc	acgaggccgt	2580
gggcgagatc	atgagcctga	gcgcccacc	ccccaaagc	ctgaagagca	tcggcctgct	2640
gagccccgac	ttccaggagg	acaacgagac	cgagatcaac	ttcctgctga	agcaggccct	2700
gaccatcgtg	ggcaccctgc	ccttcacct	catgctggag	aagtggcggg	ggatggtgtt	2760
taagggcgag	atcccccaag	accagtggat	gaagaagtgg	tgggagatga	agcgggagat	2820
cgtgggcgtg	gtggagcccc	tgccccacga	cgagacctac	tgcgaccccc	ccagcctggt	2880
ccacgtgagc	aacgactact	ccttcatccg	gtactacacc	cggaccctgt	accagttcca	2940
gttcaccagg	gcctgtgccc	aggccgcca	gcacgagggc	cccctgcaca	agtgcgacat	3000
cagcaacagc	accgaggccg	gacagaaact	gttcaacatg	ctgcggtctg	gcaagagcga	3060
gcccctggacc	ctggccctgg	agaatgtggt	gggcgccaag	aacatgaatg	tgcgccccct	3120
gctgaactac	ttcgagcccc	tgttcacctg	gctgaaggac	cagaacaaga	acagcttctg	3180
gggctgggagc	accgactgga	gcccctacgc	cgaccagagc	atcaaagtgc	ggatcagcct	3240
gaagagcgcc	ctgggcgaca	aggcctacga	gtggaacgac	aacgagatgt	acctgttccg	3300
gagcagcgtg	gcctatgcca	tgcggcagta	cttcctgaaa	gtgaagaacc	agatgatcct	3360
gttcggcgag	gaggacgtga	gagtggccaa	cctgaagccc	cggatcagct	tcaacttctt	3420
cgtgaccgcc	cccaagaacg	tgagcgacat	catccccggg	accgaagtgg	agaaggccat	3480
ccggatgagc	cggagccgga	tcaacgacgc	cttcgggctg	aacgacaact	ccctggagtt	3540
cctgggcate	cagcccaccc	tgggcccctc	caaccagccc	cccgtagca	tctggctgat	3600
cgtgtttggc	gtggtgatgg	gcgtgatcgt	ggtgggaatc	gtgatectga	tcttcaccgg	3660
catccgggac	cggagaaga	agaacaaggc	ccggagcggc	gagaaccctt	acgccagcat	3720
cgatatcagc	aagggcgaga	acaaccccgg	cttcagaac	accgacgacg	tgagaccag	3780
cttctgataa	tctagaacga	gctcgaattc	gaagcttctg	cagacgcgtc	gacgtcatat	3840
ggatccgata	tcgccgtggc	ggccgcaccc	agcgtgttca	tcttcccccc	ctccgacgag	3900
cagctgaaga	gcggcaccgc	cagcgtggtg	tgctgctgga	acaacttcta	cccccgggag	3960
gccaaggtgc	agtggaaggt	ggacaacgcc	ctgcagagcg	gcaacagcca	ggagagcgtg	4020
accgagcagg	acagcaagga	ctccacctac	agcctgagca	gcaccctcac	cctgagcaag	4080
gccgactacg	agaagcacia	ggtgtacgcc	tgcgagggtga	cccaccaggg	cctgagcagc	4140
cccgtgacca	agagcttcaa	ccggggcgag	tgtaaataga	cttaagttta	aaccgctgat	4200
cagcctcgac	tgtgccttct	agttgccagc	catctgttgt	ttgcccctcc	cccgtgcctt	4260
ccttgaccct	ggaaggtgcc	actcccactg	tcctttccta	ataaaatgag	gaaattgcat	4320
cgcattgtct	gagtaggtgt	cattctattc	tggggggtgg	ggtggggcag	gacagcaagg	4380
gggaggattg	ggaagacaat	agcaggcatg	ctggggatgc	ggtgggctct	atggcttctg	4440

aggcggaaag	aaccagctgg	ggctctaggg	ggtatcccca	cgcgcctgt	agcggcgc	4500
taagcgcggc	gggtgtgggt	gttacgcgca	gcgtgaccgc	tacacttgcc	agcgcctag	4560
cgcccgcctc	tttcgcttcc	ttcccttcc	ttctcgccac	gttcgcgggc	tttccccgtc	4620
aagctctaaa	tcgggggctc	cctttaggg	tccgatttag	tgctttacgg	cacctcgacc	4680
ccaaaaaact	tgattaggg	gatggttcac	gtagtggg	atcgccctga	tagacggttt	4740
ttcgcccttt	gacgttgag	tccacgttct	ttaatagtgg	actcttggtc	caaactggaa	4800
caacactcaa	ccctatctcg	gtctattctt	ttgatttata	agggattttg	gccatttcgg	4860
cctattgggt	aaaaaatgag	ctgatttaac	aaaaatttaa	cgcgaattaa	ttctgtggaa	4920
tgtgtgtcag	ttaggggtg	gaaagtcccc	aggctcccca	gcaggcagaa	gtatgcaaag	4980
catgcatctc	aattagtcag	caaccagggt	tggaaagtcc	ccaggctccc	cagcaggcag	5040
aagtatgcaa	agcatgcac	tcaattagtc	agcaaccata	gtcccgcgcc	taactccgcc	5100
catcccgc	ctaactccgc	ccagttccgc	ccattctcog	ccccatggct	gactaatttt	5160
ttttatttat	gcagaggccg	aggccgcctc	tgctctgag	ctattccaga	agtagtgagg	5220
aggctttttt	ggaggcctag	gcttttgcaa	aaagctccog	ggagcttgta	tatccatttt	5280
cggatctgat	cagcacgtga	tgaaaaagcc	tgaactcacc	gcgacgtctg	tcgagaagtt	5340
tctgatcgaa	aagttcgaca	gcgtctccga	cctgatgcag	ctctcggagg	gcgaagaatc	5400
tcgtgctttc	agcttcgatg	taggaggggc	tggatatgtc	ctgcccggta	atagctgcgc	5460
cgatggtttc	tacaaagatc	ggtatgttta	tcggcacttt	gcacggcccg	cgctcccgat	5520
tccggaagtg	cttgacattg	gggaattcag	cgagagcctg	acctattgca	tctcccgcgc	5580
tgcacaggg	gtcacgttgc	aagacctgcc	tgaaacccga	ctgcccgcctg	ttctgcagcc	5640
ggtcgcggag	gccatggatg	cgatcgctgc	ggccgatctt	agccagacga	gcgggttcgg	5700
cccattcggg	ccacaaggaa	tcgggtcaata	cactacatgg	cgtgatttca	tatgcgcgat	5760
tgctgatccc	catgtgtatc	actggcaaac	tgtgatggac	gacaccgtca	gtgctccgt	5820
cgcgcaggct	ctcgatgagc	tgatgctttg	ggccgaggac	tgccccgaag	tccggcacct	5880
cgtgcacgcg	gatttcggct	ccaacaatgt	cctgacggac	aatggccgca	taacagcgg	5940
cattgactgg	agcagggcga	tgttcgggga	ttccaatac	gaggtcgcca	acatcttctt	6000
ctggaggccg	tggttggctt	gtatggagca	gcagacgcgc	tacttcgagc	ggaggcatcc	6060
ggagcttgca	ggatcgccgc	ggctccgggc	gtatatgctc	cgcattggct	ttgaccaact	6120
ctatcagagc	ttggttgacg	gcaatttcga	tgatgcagct	tgggcccagg	gtcgatgcga	6180
cgcaatcgtc	cgatccggag	ccgggactgt	cgggcgtaca	caaatcgccc	gcagaagcgc	6240
ggccgtctgg	accgatggct	gtgtagaagt	actcgcgat	agtggaaacc	gacgcccag	6300
cactcgtccg	agggcaaagg	aatagcacgt	gctacgagat	ttcgattcca	ccgcgcctt	6360
ctatgaaagg	ttgggcttcg	gaatcgtttt	ccgggacgcc	ggctggatga	tcctccagcg	6420
cggggatctc	atgctggagt	tcttcgccc	ccccacttg	tttattgcag	cttataatgg	6480
ttacaaataa	agcaatagca	tcacaaat	cacaaataaa	gcattttttt	cactgcattc	6540
tagttgtgg	ttgtccaaac	tcatcaatgt	atcttatcat	gtctgtatac	cgctgcacct	6600
tagctagagc	ttggcgtaat	catggtcata	gctgtttcct	gtgtgaaatt	gttatccgct	6660
cacaattcca	cacaacatac	gagccggaag	cataaagtgt	aaagcctggg	gtgcctaattg	6720
agttagctaa	ctcacattaa	ttgcgttgcg	ctcactgcc	gctttccagt	cgggaaacct	6780
gtcgtgccag	ctgcattaat	gaatcggcc	acgcgcgggg	agaggcgggt	tgcgtattgg	6840
gcgctcttcc	gcttctctgc	tactgactc	gctgcgctcg	gtcgttcggc	tgcggcgagc	6900
ggtatcagct	cactcaaagg	cggtaatacg	gttatccaca	gaatcagggg	ataacgcagg	6960
aaagaacatg	tgagcaaaa	gccagcaaaa	ggccaggaac	cgtaaaaagg	ccgcgttgct	7020
ggcgtttttc	cataggetcc	gccccctga	cgagcatcac	aaaaatcgac	gctcaagtca	7080
gagggtggcg	aaccgacag	gactataaag	ataccaggcg	tttccccctg	gaagctccct	7140
cgtgcgctct	ctgttccga	ccctgccgct	taccggatac	ctgtccgctt	ttctcccttc	7200
gggaagcgtg	gcgctttctc	atagctcacg	ctgtaggtat	ctcagttcgg	tgtaggtcgt	7260
tcgctccaag	ctgggctgtg	tgcacgaacc	cccogttcag	cccgaccgct	gcgccttctc	7320
cggtaactat	cgtcttgagt	ccaacccggt	aagacacgac	ttatcgccac	tggcagcagc	7380
cactggtaac	aggattagca	gagcgaggta	tgtaggcgg	gctacagagt	tcttgaagtg	7440
gtggcctaac	tacggctaca	ctagaagaac	agtatttgg	atctgcgctc	tgctgaagcc	7500
agttaccttc	ggaaaaagag	ttggtagctc	ttgatccggc	aaacaaacca	ccgctggtag	7560
cggttttttt	gtttgcaagc	agcagattac	gcgcagaaaa	aaaggatctc	aagaagatcc	7620
tttgatcttt	tctacggggt	ctgacgctca	gtggaacgaa	aactcacggt	aagggatttt	7680
ggtcatgaga	ttatcaaaaa	ggatcttcac	ctagatcctt	ttaaatttaa	aatgaagttt	7740
taaatacaatc	taaagtatat	atgagtaaac	ttggtctgac	agttaccaat	gcttaatcag	7800
tgaggcacct	atctcagcga	tctgtctatt	tcgttcatcc	atagttgcct	gactccccgt	7860

cgtgtagata	actacgatac	gggagggcct	accatctggc	cccagtgctg	caatgatacc	7920
gcgagaccca	cgctcaccgg	ctccagattt	atcagcaata	aaccagccag	ccggaagggc	7980
cgagcgcaga	agtggctctg	caactttatc	cgctccatc	cagtctatta	attggtgccg	8040
ggaagctaga	gtaagtagtt	cgccagttaa	tagtttgccg	aacggtggtg	ccattgctac	8100
aggcatcgtg	gtgtcacgct	cgctcgtttg	tatggcttca	ttcagctccg	gttcccaacg	8160
atcaaggcga	gttacatgat	ccccatggt	gtgcaaaaaa	gcggttagct	ccttcgggtcc	8220
tccgatcgtt	gtcagaagta	agttggccgc	agtgttatca	ctcatgggta	tggcagcact	8280
gcataattct	cttactgtca	tgccatccgt	aagatgcttt	tctgtgactg	gtgagtactc	8340
aaccaagtca	ttctgagaat	agtgtatgcg	gcgaccgagt	tgctcttgcc	cggcgtcaat	8400
acgggataat	accgcgccac	atagcagaac	tttaaaagtg	ctcatcattg	gaaaacgttc	8460
ttcggggcga	aaactctcaa	ggatcttacc	gctgttgaga	tccagttcga	tgtaacccac	8520
tcgtgcaccc	aactgatctt	cagcatcttt	tactttcacc	agcgtttctg	ggtgagcaaa	8580
aacaggaagg	caaaatgccg	caaaaaagg	aataagggcg	acacggaaat	ggtgaatact	8640
catactcttc	ctttttcaat	attattgaag	catttatcag	ggttattgtc	tcatgagcgg	8700
atacatattt	gaatgtattt	agaaaaataa	acaaataggg	gttccgcgca	catttccccg	8760
aaaagtgccca	ctgacg					8777

Vector pIg-C910-Clambda (SEQ ID NO:177)

tcgacgggatc	gggagatctc	ccgatcccct	atgggtgcaact	ctcagtacaa	tctgctctga	60
tgccgcatag	ttaagccagt	atctgctccc	tgcttggtg	ttggaggctcg	ctgagtagtg	120
cgcgagcaaa	atttaagcta	caacaaggca	aggcttgacc	gacaattggt	aattaacatg	180
aagaatctgc	ttagggttag	gcgttttg	ctgcttcgct	agggtggtaaa	tattggccat	240
tagccatatt	attcattggt	tatatagcat	aatcaatat	tggctattgg	ccattgcata	300
cgttgatctc	atatcataat	atgtacattt	atattggctc	atgtccaaca	ttaccgccat	360
gttgacattg	attattgact	agttattaat	agtaataaat	tacgggggtca	ttagttcata	420
gccccatata	ggagttccgc	gttacataac	ttacggtaaa	tggcccgcct	ggctgaccgc	480
ccaacgaccc	cgccccattg	acgtcaataa	tgacgtatgt	tcccatagta	acgccaatag	540
ggactttcca	ttgacgtcaa	tgggtggagt	atttacggta	aactgcccac	ttggcagtac	600
atcaagtgtg	tcatatgcc	agtacgccc	ctattgacgt	caatgacggg	aatggccccg	660
cctggcatta	tgcccagtac	atgaccttat	gggactttcc	tacttggcag	tacatctacg	720
tattagtcac	cgctattacc	atgggtgatg	ggttttgcca	gtacatcaat	gggctggtgat	780
agcggtttga	ctcacgggga	tttccaagtc	tccaccccat	tgacgtcaat	gggagtttgt	840
tttggcacca	aatcaacgg	gactttccaa	aatgtcgtaa	caactccgcc	ccattgacgc	900
aatggggcgg	taggcgtgta	cggtgggagg	tctatataag	cagagctcgt	ttagtgaacc	960
gtcagatcgc	ctggagacgc	catccacgct	gttttgacct	ccatagaaga	caccgggacc	1020
gatccagcct	cgcgggccgg	gaacgggtgca	ttggaatcga	tgactctctt	aggtagcctt	1080
gcagaagttg	gtcgtgaggc	actgggcagg	taagtatcaa	ggttacaaga	caggtttaag	1140
gagatcaata	gaaactgggc	ttgtcgagac	agagaagact	cttgcgtttc	tgataggcac	1200
ctattggtct	tactgacatc	cactttgcct	ttctctccac	agggtgtccac	tcccagttca	1260
attacagctc	gccaccatgc	ggttctccgc	tcagctgctg	ggccttctgg	tgctgtggat	1320
tcccggcgtc	tcgagatcta	tcgatgcatg	ccatggtacc	aagcttgcca	ccatgagcag	1380
cagctcttgg	ctgctgctga	gcttgggtgg	cgtgacagcc	gcccagagca	ccatcgagga	1440
gcaggccaag	accttctctg	acaagttcaa	ccacgaggcc	gaggacctgt	tctaccagag	1500
cagcctggcc	agctggaact	acaacaccaa	catcaccgag	gagaacgtgc	agaacatgaa	1560
caacgcgggc	gacaagtgga	gcgcttccct	gaaggagcag	agcacactgg	cccagatgta	1620
ccccctgcag	gagatccaga	acctgaccgt	gaagctgcag	ctgcaggccc	tgcagcagaa	1680
cggcagcagc	gtgctgagcg	aggacaagag	caagcggctg	aacaccatcc	tgaacaccat	1740
gtccaccatc	tacagcaccg	gcaaagtgtg	caaccccagc	aacccccagg	agtgcctgct	1800
gctggagccc	ggcctgaacg	agatcatggc	caacagcctg	gactacaacg	agcggctgtg	1860
ggcctgggag	agctggcgga	gcgaagtggg	caagcagctg	cgccccctgt	acgaggagta	1920
cgtgggtgctg	agaacgaga	tggccagggc	caaccactac	gaggactacg	gcgactactg	1980
gagagggcag	tacgaagtga	acggcgtgga	cggctacgac	tacagcagag	gccagctgat	2040
cgaggacgtg	gagcacacct	tcgaggagat	caagcctctg	tacgagcacc	tgcacgccta	2100
cgtgcggggcc	aagctgatga	acgcctaccc	cagctacatc	agccccatcg	gctgcctgcc	2160
cgcccacctg	ctgggcgaca	tgtggggccg	gttctggacc	aacctgtaca	gacctgaccgt	2220
gcccttcggc	cagaagccca	acatcgacgt	gaccgacgcc	atgggtggacc	aggcctggga	2280
cgcccagcgg	atcttcaagg	aggccagaaa	gttcttctgtg	agcgtggggc	tgcccaacat	2340

gaccaggggc	ttttgggaga	acagcatgct	gaccgacccc	ggcaatgtgc	agaaggccgt	2400
gtgccacccc	accgcctggg	acctgggcaa	gggcgacttc	cggatcctga	tgtgcaccaa	2460
agtgaccatg	gacgacttcc	tgaccgccc	ccacgagatg	ggccacatcc	agtacgacat	2520
ggcctacgcc	gcccagccct	tctgtctgcg	gaacggcgcc	aacgagggct	ttcacgagge	2580
cgtgggcgag	atcatgagcc	tgagcgccgc	cacccccaa	cacctgaaga	gcatcggcct	2640
gctgagcccc	gacttccagg	aggacaacga	gaccgagatc	aacttctctg	tgaagcagge	2700
cctgaccatc	gtgggcaccc	tgcccttcac	ctacatgctg	gagaagtggc	ggtggatggt	2760
gtttaagggc	gagatcccca	aggaccagtg	gatgaagaag	tgggtgggaga	tgaagcggga	2820
gatcgtgggc	gtggtggagc	ccgtgcccc	cgacgagacc	tactgcgacc	ccgccagcct	2880
gttccacgtg	agcaacgact	actccttcat	ccggtactac	acccggaccc	tgtaccagtt	2940
ccagttccag	gaggccctgt	gccaggccgc	caagcacgag	ggccccctgc	acaagtgcga	3000
catcagcaac	agcaccgagg	ccggacagaa	actgttcaac	atgctgcggc	tgggcaagag	3060
cgagccctgg	accctggccc	tggagaatgt	ggtgggcgcc	aagaacatga	atgtgcgcc	3120
cctgctgaac	tacttcgagc	ccctgttcac	ctggctgaag	gaccagaaca	agaacagctt	3180
cgtgggctgg	agcaccgact	ggagccccta	cgccgaccag	agcatcaaag	tgcggatcag	3240
cctgaagagc	gcctggggcg	acaaggccta	cgagtggaac	gacaacgaga	tgtacctgtt	3300
ccggagcagc	gtggcctatg	ccatgcggca	gtacttctctg	aaagtgaaga	accagatgat	3360
cctgttcggc	gaggaggacg	tgagagtggc	caacctgaag	ccccggatca	gcttcaactt	3420
cttcgtgacc	gcccccaaga	acgtgagcga	catcatcccc	cggaccgaag	tggagaagge	3480
catccggatg	agccggagcc	ggatcaacga	cgcttccgg	ctgaacgaca	actccctgga	3540
gttctctggc	atccagccca	ccctgggccc	tcccaccag	ccccccgtga	gcatctggct	3600
gatcgtgttt	ggcgtggtga	tgggcgtgat	cgtggtggga	atcgtgatcc	tgatcttcac	3660
cggcatccgg	gaccggaaga	agaagaacaa	ggcccggagc	ggcgagaacc	cctacgccag	3720
catcgatatac	agcaagggcg	agaacaacc	cggcttccag	aacaccgacg	acgtgcagac	3780
cagcttctga	taatctagaa	cgagctcgaa	ttcgaagctt	ctgcagacgc	gtcagcgtca	3840
tatggatccg	atatcgccgt	ggcggccgca	ggccagccca	aggccgctcc	cagcgtgacc	3900
ctgttcccc	cctcctccga	ggagctgcag	gccaacaagg	ccaccctggt	gtgctcctc	3960
agcgacttct	accctggcgc	cgtgaccgtg	gcctggaagg	ccgacagcag	ccccgtgaag	4020
gccggcgtgg	agaccaccac	ccccagcaag	cagagcaaca	acaagtacgc	cgccagcagc	4080
tacctgagcc	tcacccccga	gcagtggaag	agccaccgga	gtacacagctg	ccaggtgacc	4140
cacgagggca	gcaccgtgga	gaagaccgtg	gccccaccg	agtgcagcta	atagacttaa	4200
gtttaaaccg	ctgatcagcc	tcgactgtgc	cttctagttg	ccagccatct	gttgtttgcc	4260
cctcccccg	gccttctctg	accctggaag	gtgccactcc	cactgtcctt	tcctaataaa	4320
atgaggaat	tgcctgcgat	tgtctgagta	ggtgtcattc	tattctgggg	ggtgggggtg	4380
ggcaggacag	caagggggag	gattgggaag	acaatagcag	gcatgctggg	gatgcgggtg	4440
gctctatggc	ttctgaggcg	gaaagaacca	gctggggctc	tagggggtat	ccccacgcgc	4500
cctgtagcgg	cgcattaagc	gcggcgggtg	tgggtggttac	gogcagcgtg	accgctacac	4560
ttgccagcgc	cctagcgcgc	gctcctttcg	ctttcttccc	ttcctttctc	gccacgttcg	4620
ccggctttcc	ccgtcaaget	ctaaatcggg	ggctcccctt	agggttccga	tttagtgctt	4680
tacggcacct	cgacccccaa	aaacttgatt	aggtgatggt	ttcacgtagt	gggccatcgc	4740
cctgatagac	ggtttttctg	cctttgacgt	tggagtccac	gttctttaat	agtggactct	4800
tgttccaaac	tggaacaaca	ctcaacccta	tctcggctca	ttcttttgat	ttataagggg	4860
ttttggccat	ttcggcctat	tggttaaaa	atgagctgat	ttaacaaaa	tttaacgcga	4920
attaattctg	tggaatgtgt	gtcagttagg	gtgtggaaag	tccccaggct	ccccagcagg	4980
cagaagtatg	caaagcatgc	atctcaatta	gtcagcaacc	aggtgtggaa	agtccccagg	5040
ctccccagca	ggcagaagta	tgcaaagcat	gcatctcaat	tagtcagcaa	ccatagtccc	5100
gcccctaact	cgccccatcc	cgcccctaac	tccgcccagt	tccgcccatt	ctccgcccc	5160
tggctgacta	atTTTTTTT	tttatgcaga	ggccgaggcc	gcctctgcct	ctgagctatt	5220
ccagaagtag	tgaggaggct	tttttgagg	cctaggcttt	tgcaaaaagc	tcccgggagc	5280
ttgtatatcc	atTTTtcggat	ctgatcagca	cgtgatgaaa	aagcctgaac	tcaccgcgac	5340
gtctgtcag	aagtttctga	tcgaaaagt	cgacagcgtc	tccgacctga	tgcagctctc	5400
ggagggcgaa	gaatctcgtg	ctttcagctt	cgatgtagga	gggctgggat	atgtcctgcg	5460
ggtaaatagc	tcgcgccgatg	gtttctacaa	agatcgttat	gtttatcggc	actttgcatc	5520
ggccgcgctc	ccgattccgg	aagtgtctga	cattggggaa	ttcagcgaga	gcctgacctc	5580
ttgcatctcc	cgccgtgcac	aggggtgtcac	gttgcaagac	ctgcctgaaa	ccgaactgcc	5640
cgtgttctcg	cagccggctc	cggaggccat	ggatgcgatc	gctgcccggc	atcttagcca	5700
gacgagcggg	ttcggcccat	tcggaccgca	aggaatcgg	caatacacta	catggcgtga	5760

tttcatatgc	gcgattgctg	atccccatgt	gtatcactgg	caaactgtga	tggacgacac	5820
cgtcagtgcg	tccgtcgcgc	aggetctcga	tgagctgatg	ctttgggccc	aggactgccc	5880
cgaagtccgg	cacctcgtgc	acgcggattt	cggctccaac	aatgtcctga	cggacaatgg	5940
ccgcataaca	gcggtcattg	actggagcga	ggcgatgttc	ggggattccc	aatacagagg	6000
cgccaacatc	tttctctgga	ggccgtggtt	ggcttgtatg	gagcagcaga	cgcgctactt	6060
cgagcggagg	catccggagc	ttgcaggatc	gccgcggctc	cgggcgtata	tgctccgcat	6120
tggctcttgac	caactctatc	agagcttggg	tgacggcaat	ttcgatgatg	cagcttgggc	6180
gcagggctga	tgcgacgcaa	tctgccgatc	cggagccggg	actgtcgggc	gtacacaaat	6240
cgcccgcaga	agcgcggccg	tctggaccga	tggctgtgta	gaagtactcg	ccgatagtgg	6300
aaaccgacgc	cccagcactc	gtccgagggc	aaaggaatag	cacgtgctac	gagatttcga	6360
ttccaccgcc	gccttctatg	aaaggttggg	cttcggaatc	gttttcgggg	acgccggctg	6420
gatgatcctc	cagcgcgggg	atctcatgct	ggagttcttc	gcccacccca	acttgtttat	6480
tgcagcttat	aatggttaca	aataaagcaa	tagcatcaca	aatttcacaa	ataaagcatt	6540
tttttactg	cattctagtt	gtggtttgct	caaactcact	aatgtatctt	atcatgtctg	6600
tataccgtcg	acctctagct	agagcttggc	gtaatcatgg	tcatagctgt	ttcctgtgtg	6660
aaattgttat	ccgctcacia	ttccacacaa	catacagacc	ggaagcataa	agtgtaaagc	6720
ctgggggtgcc	taatgagtga	gctaactcac	attaattgog	ttgcgctcac	tgcccgcctt	6780
ccagtcggga	aacctgtcgt	gccagctgca	ttaatgaatc	ggccaacgcg	cggggagagg	6840
cggtttgctg	attgggcgct	cttccgcctc	ctcgtcact	gactcgtctg	gctcggctct	6900
tccgctgcgg	cgagcgggat	cagctcactc	aaaggcggta	atacggttat	ccacagaatc	6960
aggggataac	gcaggaaaga	acatgtgagc	aaaaggccag	caaaaggcca	ggaaccgtaa	7020
aaaggccgcg	ttgctggcgt	ttttccatag	gctccgcccc	cctgacgagc	atcacaaaaa	7080
tgcagctca	agtcagaggt	ggcgaaacc	gacaggacta	taaagatacc	aggcgtttcc	7140
ccctggaagc	tcctcgtg	gctctcctgt	tccgaccctg	ccgcttaccg	gatacctgtc	7200
cgcctttctc	ccttcgggaa	gcgtggcgct	ttctcatagc	tcacgctgta	ggtatctcag	7260
ttcgggtgtag	gtcgttcgct	ccaagctggg	ctgtgtgcac	gaaccccccg	ttcagcccga	7320
ccgctgcgcc	ttatccggta	actatcgtct	tgagtccaac	ccggtaagac	acgacttatc	7380
gccactggca	gcagccactg	gtaacaggat	tagcagagcg	aggatgtag	gcgggtgctac	7440
agagttcttg	aagtgggtgg	ctaactacgg	ctacactaga	agaacagtat	ttggatctg	7500
cgtctgctg	aagccagtta	ccttcggaaa	aagagttggg	agctcttgat	ccggcaaa	7560
aaccaccgct	ggtagcggtt	tttttgtttg	caagcagcag	attacgcgca	gaaaaaaagg	7620
atctcaagaa	gaccttttga	tcttttctac	ggggtctgac	gctcagtgga	acgaaaactc	7680
acgtaaaggg	attttggtea	tgagattatc	aaaaaggatc	ttcacctaga	tccttttaaa	7740
ttaaaaatga	agttttaaat	caatctaaag	tatatatgag	taaacttggg	ctgacagtta	7800
ccaatgctta	atcagtgagg	cacctatctc	agcagatctgt	ctatctcgtt	catccatagt	7860
tgcttgactc	ccgctcgtgt	agataactac	gatacgggag	ggcttaccat	ctggccccag	7920
tgtgcaatg	ataccgcgag	acccacgctc	accggetcca	gatttatcag	caataaacca	7980
gccagccgga	agggccgagc	gcagaagtgg	tcctgcaact	ttatccgcct	ccatccagtc	8040
tattaattgt	tgccgggaag	ctagagtaa	tagttcgcca	gttaatagtt	tgcgcaacgt	8100
tgttgccatt	gctacaggca	tctgtgtgtc	acgctcgtcg	tttgggatgg	cttcattcag	8160
ctccggttcc	caacgatcaa	ggcgagttac	atgatcccc	atgttgtgca	aaaaagcggg	8220
tagctccttc	ggtcctccga	tctgtgtcag	aagtaagttg	gcccagctgt	tatcactcat	8280
ggttatggca	gcactgcata	attctcttac	tgtcatgcca	tcgtaagat	gcttttctgt	8340
gactggtgag	tactcaacca	agtcattctg	agaatagtgt	atgcccgcac	cgagttgctc	8400
ttgcccggcg	tcaatacggg	ataataccgc	gccacatagc	agaactttaa	aagtgtcat	8460
cattggaaaa	cgttcttcgg	ggcgaaaact	ctcaaggatc	ttaccgctgt	tgagatccag	8520
ttcgatgtaa	cccactcgtg	cacccaactg	atcttcagca	tcttttactt	tcaccagcgt	8580
ttctgggtga	gcaaaaacag	gaaggcaaaa	tgccgcaaaa	aagggaataa	gggagacacg	8640
gaaatgttga	atactcatac	tcttctttt	tcaatattat	tgaagcattt	atcagggtta	8700
ttgtctcatg	agcggataca	tatttgaatg	tatttagaaa	aataaaciaa	taggggttcc	8760
gcgcacattt	ccccgaaaag	tgccacctga	cg			8792

REFERENCES

Air MA (1981), Sequence relationships among the hemagglutinin genes of 12 subtypes of influenza A virus. Proc Natl Acad Sci USA 78(12):7639-7643.

De Kruif J et al. (1995), Rapid selection of cell subpopulation-specific human monoclonal antibodies from a synthetic phage antibody library. Proc Natl Acad Sci USA 92:3938.

Ferguson et al., (2003), Nature 422:428-443.

Fouchier AM et al. (2005), Characterization of a novel influenza A virus hemagglutinin subtype (H16) obtained from black-headed gulls. J Virol 79(5):2814-2822.

The World Health Organization Global Influenza Program Surveillance Network (2005), Evolution of H5N1 Avian Influenza Viruses in Asia. Emerg Infect Dis 11:1515-1521.

CLAIMS

1. An isolated binding molecule, capable of specifically binding to an epitope in the stem region of the hemagglutinin protein (HA) of influenza A virus subtypes of phylogenetic group 1 and influenza A virus subtypes of phylogenetic group 2 subtypes, and capable of neutralizing influenza A virus subtypes of phylogenetic group 1 and phylogenetic group 2, characterized in that the binding molecule is also capable of specifically binding to the hemagglutinin protein (HA) of influenza B virus subtypes.
2. A binding molecule according to claim 1, wherein the binding molecule is capable of neutralizing influenza B virus subtypes.
3. A binding molecule according to claim 1 or 2, wherein the binding molecule is capable of neutralizing at least one or more group 1 influenza A virus subtypes, selected from the group consisting of influenza A viruses comprising HA of the H1, H2, H5, H6, H8, H9 and H11 subtype, and at least one or more group 2 influenza A virus subtypes, selected from the group consisting of influenza A viruses comprising HA of the H3, H4, H7, and H10 subtype, and at least one or more influenza B virus subtypes.
4. A binding molecule according to any of the preceding claims, wherein the binding molecule has no hemagglutination inhibiting activity.
5. A binding molecule according to any of the preceding claims, wherein the binding molecules comprise a heavy chain CDR1 comprising the amino acid sequence of SEQ ID NO: 133 or SEQ ID NO: 139, a heavy chain CDR2 comprising the amino acid sequence of SEQ ID NO: 134, SEQ ID NO: 140 or SEQ ID NO: 151, and a heavy chain CDR3 comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 135, SEQ ID NO: 141, SEQ ID NO: 145, SEQ ID NO: 152, SEQ ID NO: 161, and SEQ ID NO: 162.

6. A binding molecule according to any of the preceding claims, wherein the binding molecule is selected from the group consisting of:

- a) a binding molecule comprising a heavy chain CDR1 region of SEQ ID NO:133, a heavy chain CDR2 region of SEQ ID NO:134, and a heavy chain CDR3 region of SEQ ID NO:135,
- b) a binding molecule comprising a heavy chain CDR1 region of SEQ ID NO:139, a heavy chain CDR2 region of SEQ ID NO:140, and a heavy chain CDR3 region of SEQ ID NO:141,
- c) a binding molecule comprising a heavy chain CDR1 region of SEQ ID NO:139, a heavy chain CDR2 region of SEQ ID NO:134, and a heavy chain CDR3 region of SEQ ID NO:145,
- d) a binding molecule comprising a heavy chain CDR1 region of SEQ ID NO:139, a heavy chain CDR2 region of SEQ ID NO:151, and a heavy chain CDR3 region of SEQ ID NO:152,
- e) a binding molecule comprising a heavy chain CDR1 region of SEQ ID NO:139, a heavy chain CDR2 region of SEQ ID NO:134, and a heavy chain CDR3 region of SEQ ID NO:152,
- f) a binding molecule comprising a heavy chain CDR1 region of SEQ ID NO:139, a heavy chain CDR2 region of SEQ ID NO:151, and a heavy chain CDR3 region of SEQ ID NO:161,
- g) a binding molecule comprising a heavy chain CDR1 region of SEQ ID NO:139, a heavy chain CDR2 region of SEQ ID NO:151, and a heavy chain CDR3 region of SEQ ID NO:162, and
- h) a binding molecule comprising a heavy chain CDR1 region of SEQ ID NO:139, a heavy chain CDR2 region of SEQ ID NO:134, and a heavy chain CDR3 region of SEQ ID NO:141.

7. A binding molecule according to any of the preceding claims, wherein the binding molecule comprises a heavy chain CDR1 region comprising the amino acid sequence of SEQ ID NO:139, a heavy chain CDR2 region comprising an amino acid sequence of

SEQ ID NO:134, and a heavy chain CDR3 region comprising the amino acid sequence of SEQ ID NO:145 or SEQ ID NO: 152.

8. A binding molecule according to any one of the preceding claims, wherein said binding molecule is a human monoclonal antibody, or an antigen-binding fragment thereof.

9. A nucleic acid molecule encoding a binding molecule according to any one of claims 1-8.

10. A binding molecule according to any one of claims 1-8, and/or a nucleic acid molecule according to claim 9, for use as a medicament and preferably for the diagnostic, therapeutic and/or prophylactic treatment of influenza infection.

11. A pharmaceutical composition comprising a binding molecule according to any one of claims 1-8, and/or a nucleic acid molecule according to claim 9, and a pharmaceutically acceptable excipient.

FIG. 1.




FIG. 2





FIG. 3



INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2012/063637

A. CLASSIFICATION OF SUBJECT MATTER INV. C07K16/10 A61K39/42 A61P31/16 ADD.		
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED		
Minimum documentation searched (classification system followed by classification symbols) C07K A61K		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) EPO-Internal, BIOSIS, EMBASE, WPI Data		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 2009/092620 A1 (MOSTE CATHERINE [FR] ET AL) 9 April 2009 (2009-04-09) cited in the application	1-11
Y	the whole document	1,4
Y	----- WO 2008/028946 A2 (CRUCCELL HOLLAND BV [NL]; VAN DEN BRINK EDWARD NORBERT [NL]; DE KRUIF C) 13 March 2008 (2008-03-13) cited in the application the whole document	1,4
	----- -/--	
<input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C. <input checked="" type="checkbox"/> See patent family annex.		
* Special categories of cited documents :		
"A" document defining the general state of the art which is not considered to be of particular relevance	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention	
"E" earlier application or patent but published on or after the international filing date	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone	
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art	
"O" document referring to an oral disclosure, use, exhibition or other means	"&" document member of the same patent family	
"P" document published prior to the international filing date but later than the priority date claimed		
Date of the actual completion of the international search 4 December 2012	Date of mailing of the international search report 18/12/2012	
Name and mailing address of the ISA/ European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Fax: (+31-70) 340-3016	Authorized officer Chapman, Rob	

INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2012/063637

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	<p>SUI JIANHUA ET AL: "Structural and functional bases for broad-spectrum neutralization of avian and human influenza A viruses", NATURE STRUCTURAL AND MOLECULAR BIOLOGY, NATURE PUBLISHING GROUP, US , vol. 16, no. 3 1 March 2009 (2009-03-01), pages 265-273, XP002538005, ISSN: 1545-9993, DOI: 10.1038/NSMB.1566 Retrieved from the Internet: URL:10.1038/NSMB.1566 [retrieved on 2009-02-22] the whole document</p>	1,4
Y	<p>LERNER RICHARD A: "Rare antibodies from combinatorial libraries suggests an SOS component of the human immunological repertoire", MOLECULAR BIOSYSTEMS, ROYAL SOCIETY OF CHEMISTRY, GB, vol. 7, no. 4, 1 January 2011 (2011-01-01), pages 1004-1012, XP009154610, ISSN: 1742-206X, DOI: 10.1039/COMB00310G [retrieved on 2011-02-04] the whole document</p>	1,4
Y	<p>EKIERT DAMIAN C ET AL: "Antibody Recognition of a Highly Conserved Influenza Virus Epitope", SCIENCE, AMERICAN ASSOCIATION FOR THE ADVANCEMENT OF SCIENCE, WASHINGTON, DC; US, vol. 324, no. 5924, 1 April 2009 (2009-04-01), pages 246-251, XP009144786, ISSN: 0036-8075 the whole document</p>	1,4
A	<p>SMIRNOV Y A ET AL: "An epitope shared by the hemagglutinins of H1, H2, H5, and H6 subtypes of influenza A virus", ACTA VIROLOGICA, ACADEMIA PRAGUE, PRAGUE, CS, vol. 43, no. 4, 1 August 1999 (1999-08-01), pages 237-244, XP009095964, ISSN: 0001-723X the whole document</p>	1-11
	----- -/--	

INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2012/063637

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>OKUNO Y ET AL: "A COMMON NEUTRALIZING EPI TOPE CONSERVED BETWEEN THE HEMAGGLUTININ SOF INFLUENZA A VIRUS H1 AND H2 STRAINS", JOURNAL OF VIROLOGY, THE AMERICAN SOCIETY FOR MICROBIOLOGY, US, vol. 67, no. 5, 1 May 1993 (1993-05-01), pages 2552-2558, XP001005475, ISSN: 0022-538X the whole document</p> <p style="text-align: center;">-----</p>	1-11
A	<p>RUDI KOFF S ET AL: "Single amino acid substitution altering antigen-binding specificity", PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF USA, NATIONAL ACADEMY OF SCIENCE, WASHINGTON, DC; US, vol. 79, 1 March 1982 (1982-03-01), pages 1979-1983, XP007901436, ISSN: 0027-8424, DOI: 10.1073/PNAS.79.6.1979 the whole document</p> <p style="text-align: center;">-----</p>	1-11
A	<p>CORTI DAVIDE ET AL: "Heterosubtypic neutralizing antibodies are produced by individuals immunized with a seasonal influenza vaccine", JOURNAL OF CLINICAL INVESTIGATION, AMERICAN SOCIETY FOR CLINICAL INVESTIGATION, US, vol. 120, no. 5, 3 May 2010 (2010-05-03), pages 1663-1673, XP009154686, ISSN: 0021-9738, DOI: 10.1172/JCI41902 [retrieved on 2010-04-12] the whole document</p> <p style="text-align: center;">-----</p>	1-11

INTERNATIONAL SEARCH REPORT

International application No.
PCT/EP2012/063637

Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

3. Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see additional sheet

1. As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.

2. As all searchable claims could be searched without effort justifying an additional fees, this Authority did not invite payment of additional fees.

3. As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:

1-11(partially)

4. No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.
- The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.
- No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. claims: 1-11(partially)

An isolated binding molecule, capable of specifically binding to an epitope in the stem region of the hemagglutinin protein (HA) of influenza A virus subtypes of phylogenetic group 1 and influenza A virus subtypes of phylogenetic group 2 subtypes, and capable of neutralizing influenza A virus subtypes of phylogenetic group 1 and phylogenetic group 2, characterized in that the binding molecule is also capable of specifically binding to the hemagglutinin protein (HA) of influenza B virus subtypes, and related subject-matter.

2-37. claims: 1-11(partially)

An isolated binding molecule, comprising a heavy chain CDR1 comprising the amino acid sequence of SEQ ID NO: 133 or SEQ ID NO: 139, a heavy chain CDR2 comprising the amino acid sequence of SEQ ID NO: 134, SEQ ID NO: 140 or SEQ ID NO: 151, and a heavy chain CDR3 comprising an amino acid sequence selected from the group consisting of SEQ ID NO: 135, SEQ ID NO: 141, SEQ ID NO: 145, SEQ ID NO: 152, SEQ ID NO: 161, and SEQ ID NO: 162, and related subject-matter.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/EP2012/063637

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 2009092620	A1	09-04-2009	CA 2700430 A1 30-04-2009
			CN 101821290 A 01-09-2010
			EP 2203477 A2 07-07-2010
			FR 2921928 A1 10-04-2009
			US 2009092620 A1 09-04-2009
			WO 2009053604 A2 30-04-2009

WO 2008028946	A2	13-03-2008	AU 2007293662 A1 13-03-2008
			CA 2663388 A1 13-03-2008
			EA 200970255 A1 30-10-2009
			EP 2059532 A2 20-05-2009
			EP 2450377 A1 09-05-2012
			JP 2010502207 A 28-01-2010
			KR 20090059121 A 10-06-2009
			US 2009311265 A1 17-12-2009
			US 2012093823 A1 19-04-2012
			US 2012276115 A1 01-11-2012
			WO 2008028946 A2 13-03-2008
