

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization

International Bureau



(10) International Publication Number
WO 2013/092616 A1

(43) International Publication Date
27 June 2013 (27.06.2013)

W I P O | P C T

(51) International Patent Classification:

C07D 401/04 (2006.01) *C07D 401/12* (2006.01)
C07D 403/04 (2006.01) *C07D 401/14* (2006.01)
C07D 405/04 (2006.01) *C07D 403/12* (2006.01)
C07D 407/04 (2006.01) *C07D 403/14* (2006.01)
C07D 413/04 (2006.01) *C07D 405/12* (2006.01)
A61K 31/397 (2006.01) *C07D 417/04* (2006.01)
A61K 31/4439 (2006.01) *C07D 417/12* (2006.01)
A61K 31/506 (2006.01)

(74) Agents: **SIMON, Elke** et al; Boehringer Ingelheim GmbH, Corporate Patents, Binger Strasse 173, 55216 Ingelheim am Rhein (DE).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(21) International Application Number:

PCT/EP2012/075996

(22) International Filing Date:

18 December 2012 (18.12.2012)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

11194277.7 19 December 2011 (19.12.2011) EP

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).


Declarations under Rule 4.17:

— as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(H))

Published:

— with international search report (Art. 21(3))

(54) Title: NEW AZIRIDINE COMPOUNDS, PHARMACEUTICAL COMPOSITIONS AND THEIR USE AS ACETYL-COA CARBOXYLASE INHIBITORS



(57) Abstract: The invention relates to new azetidine derivatives of the formula (I) wherein Ar¹, Ar², X, R, T and L are as defined in the description, to their use as medicaments, to methods for their therapeutic use and to pharmaceutical compositions containing them.

NEW AZIRIDINE COMPOUNDS, PHARMACEUTICAL COMPOSITIONS AND THEIR USE AS
ACETYL-COA CARBOXYLASE INHIBITORS

Field of the invention

This invention relates to new compounds, in particular azetidine derivatives, to
5 processes for preparing such compounds, to their use as inhibitors of acetyl-CoA
carboxylases, to methods for their therapeutic use, in particular in diseases and
conditions mediated by the inhibition of acetyl-CoA carboxylase(s), and to
pharmaceutical compositions comprising them.

10 **Background of the invention**

Obesity is major public health issues not only for the EU, USA, Japan but also for the
world in general. It is associated with a number of serious diseases including
diabetes, dyslipidemia, hypertension, cardiovascular and cerebrovascular diseases.
Although the underlying mechanisms are not yet fully understood, the impairment of
15 insulin action in target tissues by accumulation of excess lipids is generally regarded
as a key mechanism linking obesity to secondary pathologies (G. Wolf, Nutrition
Reviews Vol. 66(10):597-600; DB Savage, KF Petersen, GI Shulman, Physiol Rev.
2007;87:507-520). Therefore, understanding of cellular lipid metabolism in insulin
target tissues is crucial in order to elucidate the development of diseases associated
20 with obesity.

A central event in lipid metabolism is the generation of malonyl-CoA via carboxylation
of acetyl-CoA by the two mammalian ACC isoforms ACC1 (ACC-alpha, also termed
ACCA) and ACC2 (ACC-beta, also designated ACCB) (Saggesson D. Annu Rev Nutr.
2008;28:253-72). The malonyl-CoA generated is used for de novo fatty acid
25 synthesis and acts as inhibitor of CPT-1, thereby regulating mitochondrial fatty acid
oxidation. Furthermore, malonyl-CoA is also described to act centrally to control food
intake, and may play an important role in controlling insulin secretion from the
pancreas (GD Lopaschuk, JR Ussher, JS Jaswal. Pharmacol Rev. 2010;62(2):237-
64; D Saggesson Annu Rev Nutr. 2008;28:253-72), further coordinating the regulation
30 of intermediary metabolism.

Therefore ACC1 and ACC2 have been shown to be major regulators of fatty acid
metabolism and are presently considered as an attractive target to regulate the
human diseases of obesity, diabetes and cardiovascular complications (SJ Wakil and

LA Abu-Elheiga, J. Lipid Res. 2009. 50: S138-S143; L. Tong, HJ Harwood Jr. Journal of Cellular Biochemistry 99:1476-1488, 2006).

As a result of its unique position in intermediary metabolism, inhibition of ACC offers the ability to inhibit de novo fatty acid production in lipogenic tissues (liver and

5 adipose) while at the same time stimulating fatty acid oxidation in oxidative tissues (liver, heart, and skeletal muscle) and therefore offers an attractive modality for favorably affecting, in a concerted manner, a multitude of cardiovascular risk factors associated with obesity, diabetes, insulin resistance, and the metabolic syndrome (L. Tong, HJ Harwood Jr. Journal of Cellular Biochemistry 99:1476-1488, 2006; Corbett

10 JW, Harwood JH Jr., Recent Pat Cardiovasc Drug Discov. 2007 Nov;2(3):162-80).

In addition ACC inhibitors also have the potential to intervene in the progression of diseases that result from the rapid growth of malignant cells or invading organisms that are dependent on endogenous lipid synthesis to sustain their rapid proliferation.

The potential of ACC inhibitors as antifungal agents and as antibacterial agents is

15 well documented (L. Tong, HJ Harwood Jr. Journal of Cellular Biochemistry 99:1476-1488, 2006). Furthermore, de novo lipogenesis is known to be required for growth of many tumor cells and ACC up-regulation has been recognized in multiple human cancers, promoting lipogenesis to meet the need of cancer cells for rapid growth and proliferation (C Wang, S Rajput, K Watabe, DF Liao, D Cao Front Biosci 2010; 2:515-

20 26). This is further demonstrated in studies using ACC inhibitors which induced growth arrest and selective cytotoxicity in cancer cells and by RNA interference-mediated knock-down of ACC which inhibited growth and induced apoptosis in different types of cancer cells. Furthermore, ACC1 associates with and is regulated by the *breast cancer susceptibility gene 1* (BRCA1). Commonly occurring BRCA1 25 mutations lead to ACC1 activation and breast cancer susceptibility (C Wang, S Rajput, K Watabe, DF Liao, D Cao Front Biosci 2010; 2:515-26).

Furthermore in central nervous system disorders including but not limited to Alzheimer's disease, Parkinson disease and epilepsy, impairments in neuronal energy metabolism have been described (Ogawa M, Fukuyama H, Ouchi Y,

30 Yamauchi H, Kimura J, J Neurol Sci. 1996;139(1):78-82). Interventions targeting this metabolic defect may prove beneficial to the patients. One promising intervention is therefore to provide the glucose-compromised neuronscerebral brain neurons with ketone bodies as an alternative substrate (ST Henderson Neurotherapeutics, 2008, 5:470-480; LC Costantini, LJ Barr, JL Vogel, ST Henderson BMC Neurosci. 2008, 9

Suppl 2:S1 6; KW Baranano, AL Hartman. Curr Treat Options Neurol. 2008;10:410-9). ACC inhibition leading to increased fatty acid oxidation may thereby result in increases in the blood levels of ketone bodies thereby providing an alternative energy substrate for the brain.

5 Preclinical and clinical evidence indicates that ketone bodies can provide neuroprotective effects in models of Parkinson's disease, AD, hypoxia, ischemia, amyotrophic lateral sclerosis and glioma (LC Costantini, LJ Barr, JL Vogel, ST Henderson BMC Neurosci. 2008, 9 Suppl 2:S1 6) and improved cognitive scores in Alzheimers Diseases patients (MA Reger, ST Henderson, C Hale, B Cholerton, LD 10 Baker, GS Watson, K Hyde, D Chapman, S Craft Neurobiology of Aging 25 (2004) 311-314). The end result of increased ketone levels is an improvement in mitochondrial efficiency and reduction in the generation of reactive oxygen species (for reviews see LC Costantini, LJ Barr, JL Vogel, ST Henderson BMC Neurosci. 2008, 9 Suppl 2:S16; KW Baranano, AL Hartman. Curr Treat Options Neurol. 15 2008;10:410-9).

Aim of the present invention

The aim of the present invention is to provide new compounds, in particular new azetidine derivatives, which are active with regard to acetyl-CoA carboxylase(s).

20 Another aim of the present invention is to provide new compounds, in particular new azetidine derivatives, which are active with regard to ACC2

25 A further aim of the present invention is to provide new compounds, in particular new azetidine derivatives, which have an inhibitory effect on acetyl-CoA carboxylase(s) *in vitro* and/or *in vivo* and possess suitable pharmacological and pharmacokinetic properties to use them as medicaments.

30 A further aim of the present invention is to provide new compounds, in particular new azetidine derivatives, which have an inhibitory effect on ACC2 *in vitro* and/or *in vivo* and possess suitable pharmacological and pharmacokinetic properties to use them as medicaments.

A further aim of the present invention is to provide effective ACC inhibitors, in particular for the treatment of metabolic disorders, for example of obesity and/or diabetes.

5 A further aim of the present invention is to provide methods for treating a disease or condition mediated by the inhibition of acetyl-CoA carboxylase(s) in a patient.

A further aim of the present invention is to provide a pharmaceutical composition comprising at least one compound according to the invention.

10

A further aim of the present invention is to provide a combination of at least one compound according to the invention with one or more additional therapeutic agents.

15

A further aim of the present invention is to provide methods for the synthesis of the new compounds, in particular azetidine derivatives.

A further aim of the present invention is to provide starting and/or intermediate compounds suitable in methods for the synthesis of the new compounds.

20

Further aims of the present invention become apparent to the one skilled in the art by the description hereinbefore and in the following and by the examples.

Object of the Invention


25

Within the scope of the present invention it has now surprisingly been found that the new compounds of general formula (I) as described hereinafter exhibit an inhibiting activity with regard to acetyl-CoA carboxylase(s).

30

According to another aspect of the present invention it has been found that the new compounds of general formula (I) as described hereinafter exhibit an inhibiting activity with regard to ACC2.

Therefore, in a first aspect the present invention provides a compound of general formula (I)



wherein

Ar^1 is selected from the group $\text{Ar}^1\text{-G1}$ consisting of:

5 6- to 10-membered aryl and 5- to 10-membered heteroaryl, which may each
be substituted with one or more substituents R^1 ,

10 wherein two substituents R^1 linked to adjacent C-atoms of Ar^1 together
may form a C_{3-5} -alkylene bridge in which 1, 2 or 3 CH_2 -groups may be
replaced independently of each other by O, C(=O), S, S(=O), S(=O)₂, NH
or N(Ci-4-alkyl)-, wherein the alkylene bridge may optionally be substituted
by one or two F atoms or one or two Ci-3-alkyl groups; and

R^1 is selected from the group $\text{R}^1\text{-G1}$ consisting of:

15 H, F, Cl, Br, I, CN, OH, -NO₂, Ci-₆-alkyl, C_{2-6} -alkenyl, C_{2-6} -alkynyl, C_{3-10} -
carbocyclyl, C_{3-10} -carbocyclyl-Ci-3-alkyl, Ci-6-alkyl-O-, C_{3-6} -alkenyl-O-, C_{3-6} -
alkynyl-O-, C_{3-10} -carbocyclyl-O-, C_{3-10} -carbocyclyl-Ci-3-alkyl-O-, Ci-6-alkyl-S-,
Ci-6-alkyl-S(=O)-, Ci-6-alkyl-S(=O)₂-, C_{3-10} -carbocyclyl-S-, C_{3-10} -carbocyclyl-C-
 i-3 -alkyl-S-, Ci-₄-alkyl-C(=O)-, C_{3-10} -carbocyclyl-C(=O)-, $\text{R}^{\text{N1}}\text{R}^{\text{N2}}\text{N}$, $\text{R}^{\text{N1}}\text{R}^{\text{N2}}\text{N}$ -
20 C_{2-3} -alkyl-O-, $\text{R}^{\text{N1}}\text{R}^{\text{N2}}\text{N-C(=O)}$ -, $\text{R}^{\text{N1}}\text{R}^{\text{N2}}\text{N-S(=O)}$ ₂-, Ci-₆-alkyl-C(=O)-N R^{N1} , Ci-₆-
alkyl-S(=O)₂-N R^{N1} , Ci-₆-alkyl-C(=O)-N R^{N1} -Ci-₃-alkyl-, HO-C(=O)-, Ci-₆-
alkyl-O-C(=O)-, heterocyclyl, heterocyclyl-O-, heterocyclyl-Ci-3-alkyl,
heterocyclyl-Ci-3-alkyl-O-, heterocyclyl-C(=O)-, aryl, aryl-Ci-3-alkyl, aryl-O-,
aryl-Ci-3-alkyl-O-, heteroaryl, heteroaryl-Ci-3-alkyl, heteroaryl-O- and
25 heteroaryl-Ci-3-alkyl-O-,

wherein in each carbocyclyl and heterocyclyl a CH_2 -group may optionally be
replaced by -C(=O)- or -C(=CR^{N2})-, and

- 6 -

wherein each carbocydyl and heterocyclyl may be optionally substituted with one or more Ci₁₋₄-alkyl, which may be optionally substituted with one or more substituents R^c, and

5 wherein each alkyl, carbocydyl and heterocyclyl may be optionally substituted with one or more substituents R^c, and

wherein each heterocyclyl may be optionally substituted with aryl or heteroaryl, and

10 wherein each aryl and heteroaryl group may be optionally substituted with one or more substituents R²,

R^c is selected from the group R^c-G1 consisting of:

15 F, Cl, Br, CN, OH, Ci₁₋₄-alkyl-O-, C₃₋₇-cycloalkyl-O-, C₃₋₇-cycloalkyl-Ci₁₋₃-alkyl-O-, H₂N-, (Ci₁₋₄-alkyl)NH-, (Ci₁₋₄-alkyl)₂N-, Ci₁₋₄-alkyl-C(=O)-, Ci₁₋₄-alkyl-S(=O)₂-, HO-C(=O)- and Ci₁₋₄-alkyl-O-C(=O)-, wherein each alkyl or cycloalkyl may be optionally substituted with one or more substituents selected from F and OH, and

20 R^{N1} is selected from the group R^{N1}-G1 consisting of:
H, Ci-6-alkyl, C₃-io-carbocydyl, C₃-io-carbocydyl-Ci-3-alkyl, C₃-6-alkenyl, C₃-6-alkynyl, heterocyclyl, heterocyclyl-Ci-3-alkyl, aryl, aryl-Ci-3-alkyl, heteroaryl and heteroaryl-Ci-3-alkyl,

25 wherein each carbocydyl and heterocyclyl may be optionally substituted with one or more Ci₁₋₄-alkyl, and

30 wherein in each carbocydyl and heterocyclyl a CH₂-group may optionally be replaced by -C(=O)-, and

wherein each alkyl, carbocydyl and heterocyclyl may be optionally substituted with one or more substituents R^c, and

wherein each aryl and heteroaryl group may be optionally substituted with one or more substituents R²,

R^{N2} is selected from the group R^{N2}-G1 consisting of: H and Ci-6-alkyl, and

5

Ar² is selected from the group Ar²-G1 consisting of:

phenylene and a 5- or 6-membered monocyclic heteroaryl group containing 1, 2 or 3 heteroatoms selected from N, O, or S, wherein all of the before mentioned groups may be optionally substituted with one or more

10 substituents R², and

R² is selected from the group R²-G1 consisting of:

F, Cl, Br, CN, OH, Ci₄-alkyl-, Ci₄-alkyl-O-, Ci₄-alkyl-S-, Ci₄-alkyl-S(=O)₂-, H₂N-, (Ci₄-alkyl)NH-, (Ci₄-alkyl)₂N-, Ci₄-alkyl-C(=O)-NH- and heterocyclyl,

15

wherein each alkyl may be optionally substituted with one or more F-atoms and/ or with a substituent selected from OH, Ci-3-alkyl-O- and CN; and

20 wherein two substituents R² attached to an aryl or heteroaryl group may be linked to each other and form a C2-5-alkylene bridging group in which 1 or 2 - CH₂-groups may be replaced by a group independently of each other selected from O, S, NH and N(Ci₄-alkyl)-, wherein the C2-5-alkylene bridging group is optionally be substituted by 1 or 2 Ci-3-alkyl groups;

25 L is selected from the group L-G1 consisting of:

a Ci₄-alkylene group that is optionally substituted by one or two CH₃;

X is selected from the group X-G1 consisting of: O and S;

30 R is selected from the group R-G1 consisting of: H and Ci-3-alkyl; and

T is selected from the group T-G1 consisting of:

Ci-6-alkyl, C2-6-alkenyl, C2-6-alkynyl, C3-*io*-carbocyclyl, C3-*io*-carbocyclyl-Ci-3-alkyl, Ci-6-alkyl-O-, C3-*io*-carbocyclyl-O-, C3-*io*-carbocyclyl-Ci-3-alkyl-O-, Ci-6-

alkyl-S-, C₃-io-carbocyclyl-S-, C₃-io-carbocyclyl-Ci-3-alkyl-S-, Ci₄-alkyl-C(=O)-, Ci-4-alkyl-S(=O)₂-, R^{N1}R^{N2}-N-, R^{N1}R^{N2}-N-Ci₃-alkyl-, R^{N1}R^{N2}-N-CO-, Ci₄-alkyl-C(=O)-R^{N2}N-Ci-3-alkyl, heterocyclyl, aryl and heteroaryl,

5 wherein in each carbocyclyl and heterocyclyl a CH₂-group may optionally be replaced by -C(=O)-, and

10 wherein each carbocyclyl and heterocyclyl may be optionally substituted with one or more Ci₄-alkyl, which may be optionally substituted with one or more substituents R^c, and

wherein each alkyl, carbocyclyl and heterocyclyl may be optionally substituted with one or more substituents R^c, and

15 wherein each alkyl may optionally be substituted with a heteroaryl group; and

wherein each aryl and heteroaryl group may be optionally substituted with one or more substituents R²;

20 including any tautomers and stereoisomers thereof,

or a salt thereof

or a solvate or hydrate thereof.

25

In a further aspect the present invention relates to processes for preparing a compound of general formula (I) and to new intermediate compounds in these processes.

30 A further aspect of the invention relates to a salt of the compounds of general formula (I) according to this invention, in particular to a pharmaceutically acceptable salt thereof.

In a further aspect this invention relates to a pharmaceutical composition, comprising one or more compounds of general formula (I) or one or more pharmaceutically acceptable salts thereof according to the invention, optionally together with one or more inert carriers and/or diluents.

5

In a further aspect this invention relates to a method for treating diseases or conditions which are mediated by inhibiting the activity of acetyl-CoA carboxylase(s) in a patient in need thereof characterized in that a compound of general formula (I) or a pharmaceutically acceptable salt thereof is administered to the patient.

10

According to another aspect of the invention, there is provided a method for treating a metabolic disease or disorder in a patient in need thereof characterized in that a compound of general formula (I) or a pharmaceutically acceptable salt thereof is administered to the patient.

15

According to another aspect of the invention, there is provided a method for treating a cardiovascular disease or disorder in a patient in need thereof characterized in that a compound of general formula (I) or a pharmaceutically acceptable salt thereof is administered to the patient.

20

According to another aspect of the invention, there is provided a method for treating a neurodegenerative disease or disorder or for treating a disease or disorder of the central nervous system in a patient in need thereof characterized in that a compound of general formula (I) or a pharmaceutically acceptable salt thereof is administered to 25 the patient.

According to another aspect of the invention, there is provided a method for treating a cancer, a malignant disorder or a neoplasia in a patient in need thereof characterized in that a compound of general formula (I) or a pharmaceutically acceptable salt thereof is administered to the patient.

30

According to another aspect of the invention, there is provided the use of a compound of the general formula (I) or a physiologically acceptable salt thereof for

the manufacture of a medicament for a therapeutic method as described hereinbefore and hereinafter.

According to another aspect of the invention, there is provided a compound of the
5 general formula (I) or a physiologically acceptable salt thereof for a therapeutic
method as described hereinbefore and hereinafter.

In a further aspect this invention relates to a method for treating a disease or
condition mediated by the inhibition of acetyl-CoA carboxylase(s) in a patient that
10 includes the step of administering to the patient in need of such treatment a
therapeutically effective amount of a compound of the general formula (I) in
combination with a therapeutically effective amount of one or more additional
therapeutic agents.

15 In a further aspect this invention relates to a use of a compound of the general
formula (I) in combination with one or more additional therapeutic agents for the
treatment or prevention of diseases or conditions which are mediated by the
inhibition of acetyl-CoA carboxylase(s).

20 In a further aspect this invention relates to a pharmaceutical composition which
comprises a compound according to general formula (I) or a pharmaceutically
acceptable salt thereof and one or more additional therapeutic agents, optionally
together with one or more inert carriers and/or diluents.

25 Other aspects of the invention become apparent to the one skilled in the art from the
specification and the experimental part as described hereinbefore and hereinafter.

Detailed Description

30 Unless otherwise stated, the groups, residues, and substituents, particularly Ar¹, Ar²,
X, R, T, R¹, R², R^c, R^{N1}, R^{N2}, and L are defined as above and hereinafter. If residues,
substituents, or groups occur several times in a compound, as for example R^c, R^{N1},
R^{N2} or R², they may have the same or different meanings. Some preferred meanings
of individual groups and substituents of the compounds according to the invention will

be given hereinafter. Any and each of these definitions may be combined with each other.

Ar¹:

5 **Ar¹-G1 :**

The group Ar¹ is preferably selected from the group Ar¹-G1 as defined hereinbefore and hereinafter.

Ar¹-G2:

10 In one embodiment the group Ar¹ is selected from the group Ar¹-G2 consisting of: phenyl, naphthyl, dihydronaphthyl, tetrahydronaphthyl, naphthyl and a 5- or 6-membered monocyclic or 9- or 10-membered bicyclic ring system containing 1, 2 or 3 heteroatoms selected from N, O, S, or S(O)_r with r = 1 or 2, wherein at least one of the heteroatoms is part of an aromatic ring, and

15

wherein all of the before mentioned groups may be optionally substituted with one or more substituents R¹,

20 particularly wherein all of the before mentioned groups may be optionally substituted with a substituent R¹ and optionally one or more substituents R², and

wherein two substituents R¹ linked to adjacent C-atoms of Ar¹ may be connected with each other and together form a C₃₋₅-alkylene bridging group in which 1, 2 or 3 CH₂-groups may be replaced independently of each other by O, C(=O), S, S(=O), S(=O)2, 25 NH or N(Ci₁₋₄-alkyl)-, wherein the alkylene bridging group may optionally be substituted by one or two F atoms or one or two Ci-3-alkyl groups.

Ar¹-G3:

In another embodiment the group Ar¹ is selected from the group Ar¹-G3 consisting of: 30 phenyl, naphthyl, pyridyl, 2H-pyridin-2-onyl, pyrimidinyl, pyridazinyl, pyrazinyl, furanyl, thienyl, pyrrolyl, imidazolyl, triazolyl, oxazolyl, isoxazolyl, pyrazolyl, thiazolyl, quinolinyl, isoquinolinyl, quinoxalinyl, indolyl, benzofuranyl, benzoimidazolyl, benzoxazolyl, benzoisoxazolyl, benzthiazolyl, benzotriazolyl, oxazolopyrimidinyl, 2,3-

dihydrobenzofuranyl, benzo[1,3]dioxolyl, 2,3-dihydro-benzo[1,4]dioxinyl, 2,3,4,5-tetrahydro-benzo[b]oxepinyl, and 3,4-dihydro-2H-benzo[b][1,4]dioxepinyl,

5 wherein the before mentioned bicyclic groups preferably are linked to the azetidine ring of the core structure of the formula (I) via an aromatic or heteroaromatic ring of the bicyclic group, and

wherein all of the before mentioned mono- and bicyclic groups may be optionally substituted with one or more substituents R¹,

10 particularly wherein all of the before mentioned mono- or bicyclic groups may be optionally substituted with a substituent R¹ and optionally one or more substituents R².

15 **Ar¹-G4:**

In another embodiment the group Ar¹ is selected from the group Ar¹-G4 consisting of: phenyl, pyridinyl, pyrimidinyl, benzoxazolyl, benzoisoxazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinoxalinyl and oxazolopyrimidinyl,

20 wherein the above-mentioned phenyl, pyridinyl and pyrimidinyl are each substituted with one to three groups independently selected from R¹, and/or

25 wherein two adjacent carbon atoms of a phenyl group may be linked to each other via a -O-CH₂-O-, -O-CF₂-O-, -O-CH₂-CH₂-O-, -O-CH₂-CH₂-CH₂-CH₂-O-CH₂-CH₂-CH₂-C(CH₃)₂-O-, -O-CH₂-CH₂-CH₂-O- or -O-CH₂-C(CH₃)₂-CH₂-O-bridge, and

30 wherein the above-mentioned benzoxazolyl, benzoisoxazolyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinoxalinyl and oxazolopyrimidinyl groups are each optionally substituted with Cl, CH₃ or phenyl.

R¹ may be any of the subgroups of R¹ as defined below. However, with respect to Ar¹-G4, R¹ is preferably a group selected from the group R¹-G4 as defined below.

Ar¹-G5:

In another embodiment the group Ar¹ is selected from the group Ar¹-G5 consisting of: phenyl, pyridinyl and pyrimidinyl,

5 wherein the above-mentioned phenyl and pyridinyl groups are each substituted with one to three groups independently selected from R¹, and

wherein the above-mentioned pyrimidinyl groups is substituted with one or two R¹, and

10 wherein two adjacent carbon atoms of a phenyl group may be linked to each other via a -O-CH₂-CH₂-O-, -O-CH₂-CH₂-C(CH₃)₂-,-O-CH₂-CH₂-CH₂-O- , or -O-CH₂-C(CH₃)₂-CH₂-O- bridge.

15 R¹ may be any of the subgroups of R¹ as defined below. However, with respect to Ar¹-G5, R¹ is preferably a group selected from the group R¹-G5 as defined below.


Ar¹-G6:

In another embodiment the group Ar¹ is selected from the group Ar¹-G6 consisting of: phenyl and pyridinyl, which are each optionally substituted with one to three substituents independently selected from the group consisting of:

F, Cl, Br, CN, CF₃, Ci₄ alkyl, phenyl, -CO-O-Ci₄-alkyl, -O-phenyl-O-pyrimidinyl, -O-Ci₄ alkyl, -O-C₃₋₇ cycloalkyl and -O-CH₂-C₃₋₅ cycloalkyl, wherein the cycloalkyl groups may optionally be substituted with one CN or one or two F.


Preferred substituents of the phenyl group are:

30 F, Cl, Br, CN, CF₃, Ci₄ alkyl, -O-Ci₄ alkyl, -O-C₃₋₅ cycloalkyl, -O-CH₂-cyclopropyl and



Preferred substituents of the pyridinyl group are:

F, Cl, Br, CN, CF₃, phenyl, -CO-O-CH₂CH₃, -O-phenyl.-O-pyrimidinyl, C₁₋₃ alkyl, -O-Ci₄ alkyl, -O-CH₂-cyclopropyl and



5

Ar¹-G6a:

In another embodiment the group Ar¹ is selected from the group Ar¹-G6a consisting of:


10 phenyl, which is optionally substituted with one to three substituents independently selected from the group consisting of:

F, Cl, Br, CN, CF₃, C₁₋₄ alkyl, -O-CM alkyl, -O-C₃₋₇ cycloalkyl and -O-CH₂-C₃₋₅ cycloalkyl, wherein the cycloalkyl groups may optionally be substituted with

15 one or two F.

Preferred substituents of the phenyl group are:

F, Cl, Br, CN, CF₃, C₁₋₄ alkyl, -O-CM alkyl, -O-C₃₋₅ cycloalkyl, -O-CH₂-cyclopropyl and



20

R¹:

R¹-G1 :

The group R¹ is preferably selected from the group R¹-G1 as defined hereinbefore
25 and hereinafter,

R¹-G2:

In another embodiment the group R¹ is selected from the group R¹-G2 consisting of:

H, F, Cl, Br, I, CN, OH, -NO₂, Ci₁₋₆-alkyl, C₂₋₆-alkenyl, C₃₋₇-cycloalkyl, C₃₋₇-cycloalkyl-Ci₁₋₃-alkyl, Ci₁₋₆-alkyl-O-, C₃₋₇-cycloalkyl-O-, C₃₋₇-cycloalkyl-Ci₁₋₃-alkyl-O-, Ci₁₋₆-alkyl-S-, Ci₁₋₆-alkyl-S(=O)₂-, C₃₋₇-cycloalkyl-S-, C₃₋₇-cycloalkyl-Ci₁₋₃-alkyl-S-, Ci₁₋₄-alkyl-C(=O)-, C₃₋₇-cycloalkyl-C(=O)-, R^{N1}R^{N2}N-, R^{N1}R^{N2}N-C₂₋₃-alkyl-O-, R^{N1}R^{N2}N-C(=O)-, R^{N1}R^{N2}N-
5 S(=O)₂-, Ci₁₋₄-alkyl-C(=O)-NR^{N1}-, Ci₁₋₄-alkyl-S(=O)₂-NR^{N1}-, Ci₁₋₄-alkyl-C(=O)-NR^{N1}-Ci₁₋₃-alkyl-, HO-C(=O)-, Ci₁₋₆-alkyl-O-C(=O)-, heterocydyl, heterocycl-O-, heterocycl-Ci₁₋₃-alkyl, heterocycl-Ci₁₋₃-alkyl-O-, heterocycl-C(=O)-, phenyl, phenyl-Ci₁₋₃-alkyl, phenyl-O-, phenyl-Ci₁₋₃-alkyl-O-, heteroaryl, heteroaryl-Ci₁₋₃-alkyl, heteroaryl-O- and heteroaryl-Ci₁₋₃-alkyl-O-,
10

10

wherein in each cydoalkyi and heterocydyl a CH₂-group may optionally be replaced by -C(=O)-, and

15

wherein each cydoalkyi and heterocydyl may be optionally substituted with one or more Ci₁₋₄-alkyl, which may be optionally substituted with one or more substituents R^c, and

20

wherein each heterocydyl may be optionally substituted with aryl or heteroaryl, and

25

wherein each phenyl and heteroaryl group may be optionally substituted with one or more substituents R², and

30

wherein each heterocydyl group is selected from the group consisting of: piperidinyl, piperazinyl, morpholinyl, pyrrolidinyl, tetrahydrofuranyl and azetidinyl, and

35

wherein each heteroaryl group is selected from the group consisting of: pyridinyl, pyrimidinyl, pyrazinyl, furanyl, thienyl, pyridazinyl, pyrrolyl, pyrazolyl, imidazolyl, oxazolyl, isoxazolyl, thiazolyl, indolyl, and benzimidazolyl.

R¹-G3:

In another embodiment the group R¹ is selected from the group R¹-G3 consisting of:

- 16 -

H, F, Cl, Br, CN, OH, Cl_{-4} -alkyl, C_{2-4} -alkenyl, C_{3-7} -cycloalkyl, C_{3-7} -cycloalkyl- CH_2 , Cl_{-4} -alkyl-O-, C_{3-7} -cycloalkyl-O-, C_{3-7} -cycloalkyl- Cl_{-3} -alkyl-O-, Cl_{-4} -alkyl-S(=O)₂, $\text{R}^{\text{N}1}\text{R}^{\text{N}2}\text{N}$, HO-C(=O)-, Cl_{-4} -alkyl-O-C(=O)-, phenyl, phenyl-O-, phenyl- CH_2 -O-, pyridinyl,

5

wherein each alkyl, cydoalkyi and heterocydyl may be optionally substituted with one to three substituents independently selected from the group consisting of F and CN, and

10 wherein each phenyl and pyridinyl group may be optionally substituted with one F or -OCH₃.

R¹-G4:


In another embodiment the group R¹ is selected from the group R¹-G4 consisting of:
 15 F, Cl, Br, CN, Cl_{-4} alkyl, -O- Cl_{-4} alkyl, C_{3-7} cydoalkyi, -O- C_{3-7} cydoalkyi, -O-(Cl_{-2} alkyl)-
 C_{3-7} cydoalkyi, -NH- C_{3-7} cydoalkyi, -NH(Cl_{-4} alkyl), -CO-O(Cl_{-4} alkyl), phenyl, -O-
 phenyl, pyrimidinyl and pyridinyl,

20 wherein the alkyl and cydoalkyi groups may optionally be substituted with one to three F and/or one CN, and

wherein the phenyl and pyridinyl groups may optionally be substituted with one F or OCH₃.

R¹-G5:


In another embodiment the group R¹ is selected from the group R¹-G5 consisting of:
 F, Cl, CN, CF₃, Cl_{-4} alkyl, -O- Cl_{-4} alkyl, cyclopropyl, -O- C_{3-5} cydoalkyi, -O- CH_2 -
 cyclopropyl, and



R¹-G5a:


In another embodiment the group R¹ is selected from the group R¹-G5a consisting of:

F, Cl, Br, CN, CF_3 , phenyl, -CO-O-CH₂CH₃, -O-phenyl-O-pyrimidinyl, $\text{C}_1\text{-}_3$ alkyl, -O-C₁₋₄ alkyl, -O-CH₂-cyclopropyl and



5 **R¹-G5b:**

In another embodiment the group R¹ is selected from the group R¹-G5b consisting of: Cl, Br, CF_3 , $\text{C}_1\text{-}_4$ alkyl, -O-C₁₋₄ alkyl, -CO-O-CH₂CH₃, cyclopropyl, phenyl, pyridinyl, -O-CH₂-cyclopropyl, and



10 wherein the phenyl and pyridinyl may be substituted by F or -O-CH₃.


R¹-G5c:

In another embodiment the group R¹ is selected from the group R¹-G5c consisting of: -NH-cyclobutyl, -O-cyclobutyl and -NH(CH₂CH₃).

15

R¹-G5d:

In another embodiment the group R¹ is selected from the group R¹-G5d consisting of: CF_3 , -O-(C₁₋₂-alkyl), -O-(C₃₋₄-cycloalkyl), -O-CH₂-cyclopropyl and



20

R^c

R^c-G1 :

The group R^c is preferably selected from the group R^c-G1 as defined hereinbefore and hereinafter.

25

R^c-G2:

In one embodiment the group R^c is selected from the group R^c-G2 consisting of:

F, **Cl**, Br, **CN**, OH, **Ci₃-alkyl-O-**, **C₃₋₆-cycloalkyl-O-**, H₂N-, **(Ci₃-alkyl)NH-**, **(Ci₃-alkyl)₂N-**, **Ci₃-alkyl-C(=O)-**, **Ci₃-alkyl-S(=O)₂-**, HO-C(=O)- and **Ci₃-alkyl-O-C(=O)-**, wherein each alkyl may be optionally substituted with one or more F-atoms and/or may be substituted with OH.

5

R^c-G3:

In another embodiment the group R^c is selected from the group R^c-G3 consisting of: F, **Cl**, **CN**, OH and **Ci₃-alkyl-O-**.

10 **R^c-G4:**

In another embodiment the group R^c is selected from the group R^c-G4 consisting of: F, **CN**, OH and **-O-CH₃-**.

R^{N1}15 **R^{N1}-G1 :**

The group R^{N1} is preferably selected from the group R^{N1}-G1 as defined hereinbefore and hereinafter.

With regard to an alkenyl or alkynyl group, for example R^{N1}, attached to the **N**-atom 20 of an amino-group it is to be understood that the double or triple bond is preferably not conjugated with the **N**-atom.

R^{N1}-G2:

In one embodiment, the group R^{N1} is selected from the group R^{N1}-G2 consisting of: 25 **H**, **Ci₄-alkyl**, **C₃₋₆-cycloalkyl**, **C₃₋₆-cycloalkyl-CH₂-**, heterocydyl, heterocycl¹-CH₂-², phenyl, **phenyl-CH₂-**, pyridyl, **pyridyl-CH₂-**, pyrazolyl-CH₂- and oxazolyl-**Ci₃-alkyl**;

wherein heterocydyl is defined as hereinbefore and hereinafter, or each heterocydyl 30 is preferably selected from tetrahydrofuran, tetrahydropyran, pyrrolidinyl and piperidinyl; and

wherein each cycloalkyl and heterocydyl may be optionally substituted with one or two **Ci₄-alkyl**; and

wherein each alkyl, cydoalkyl and heterocydyl may be optionally substituted with one to three F; and

5 wherein each alkyl, cydoalkyl and heterocydyl may be optionally substituted with a substituent selected from OH, $\text{C}_{i-3}\text{-alkyl-O-}$, $\text{H}_2\text{N-}$, HO-C(=O)- and $\text{C}_{i-4}\text{-alkyl-O-C(=O)-}$; and

wherein each phenyl, pyridyl, pyrazolyl and oxazolyl may be optionally substituted with one to three substituents R^2 .

10

 $\text{R}^{N1}\text{-G3:}$

In another embodiment the group R^{N1} is selected from the group $\text{R}^{N1}\text{-G3}$ consisting of H, $\text{C}_{i-4}\text{-alkyl}$, c3-6-cycloalkyl- $\text{C}_{i-3}\text{-alkyl-}$ and c3-6-cycloalkyl.

15

 R^{N2} **$\text{R}^{N2}\text{-G1 :}$**

The group R^{N2} is preferably selected from the group $\text{R}^{N2}\text{-G1}$ as defined hereinbefore and hereinafter.

20

 $\text{R}^{N2}\text{-G2:}$

In another embodiment the group R^{N2} is selected from the group $\text{R}^{N2}\text{-G2}$ consisting of H and $\text{C}_{i-4}\text{-alkyl}$.

 $\text{R}^{N2}\text{-G3:}$

25 In another embodiment the group R^{N2} is selected from the group $\text{R}^{N2}\text{-G3}$ consisting of H and CH_3 .

 $\text{Ar}^2:$ **$\text{Ai}^\wedge\text{-G1 :}$**

30 The group Ar^2 is preferably selected from the group $\text{Ar}^2\text{-G1}$ as defined hereinbefore and hereinafter.

 $\text{Ar}^2\text{-G2:}$

- 20 -

In another embodiment the group Ar² is selected from the group Ar²-G2 consisting of: phenylene, pyridylene, pyrimidinylene, pyridazinylene, pyrazinylene, furanylene, thienylene, pyrrolylene, imidazolylene, triazolylene, oxazolylene, isoxazolylene, pyrazolylen e and thiazolylene, wherein all of the before mentioned groups may be 5 optionally substituted with one or more substituents R².

Ar²-G3:

In another embodiment the group Ar² is selected from the group Ar²-G3 consisting of phenylene and pyridylene, wherein all of the before mentioned groups may be 10 optionally substituted with one to three substituents R².

Ai¹-GSa:

In another embodiment the group Ar² is selected from the group Ar²-G3a consisting of phenylene and pyridylene.

15

Ar²-G4:


In another embodiment the group Ar² is selected from the group Ar²-G4 consisting of: phenylene, which may be optionally substituted with one or two substituents R².

20 **Ar²-G4a:**

In another embodiment the group Ar² is selected from the group Ar²-G4a consisting of phenylene.

Ar²-G5:

25 In another embodiment the group Ar² is selected from the group Ar²-G5 consisting of:



R²:

R²-G1 :

30 The group R² is preferably selected from the group R²-G1 as defined hereinbefore and hereinafter.

R²-G2:

In another embodiment the group R² is selected from the group R²-G2 consisting of: F, Cl, Br, CN, OH, Ci₁₋₃-alkyl-, Ci₁₋₃-alkyl-O-, Ci₁₋₃-alkyl-S-, H₂N-, (Ci₁₋₃-alkyl)NH-, (Ci₁₋₃-alkyl)₂N-, Ci₁₋₃-alkyl-C(=O)-NH- and heterocyclyl;

5

wherein each alkyl may be optionally substituted with one or more F-atoms and/or a substituent selected from OH, Ci₁₋₃-alkyl-O- and CN; and

wherein heterocyclyl is defined as hereinbefore and hereinafter, or heterocyclyl
10 preferably denotes a C₃₋₆-cycloalkyl ring wherein one or two -CH₂-groups are replaced by a group selected from -O-, -NH-, -N(CH₃)-; and

wherein two substituents R² attached to an aryl or heteroaryl group may be linked to each other and form a C2-5-alkylene bridging group in which 1 or 2 -CH₂-groups may
15 be replaced by a group independently of each other selected from O, NH and N(CH₃)-, wherein the C2-5-alkylene bridging group is optionally be substituted by 1 or 2 CH₃- groups.

R²-G3:

20 In another embodiment the group R² is selected from the group R²-G3 consisting of: F, Cl, CN, OH, Ci₁₋₃-alkyl-, Ci₁₋₃-alkyl-O-, Ci₁₋₃-alkyl-S-, H₂N-, (Ci₁₋₃-alkyl)NH-, (Ci₁₋₃-alkyl)₂N- and Ci₁₋₃-alkyl-C(=O)-NH-;

wherein each alkyl may be optionally substituted with one or more F-atoms and/or a
25 substituent selected from OH, CH₃-O- and CN; and

wherein two substituents R² attached to adjacent C-atoms of an aryl or heteroaryl group may be linked to each other and form a -CH₂-CH₂-O-, -O-CH₂-CH₂-O- or -O-CH₂-O- bridging group which is optionally substituted by 1 or 2 CH₃- groups.

30

R²-G4:

In another embodiment the group R² is selected from the group R²-G4 consisting of: F, Cl, Ci₁₋₂-alkyl-, Ci₁₋₂-alkyl-O-, Ci₁₋₃-alkyl-C(=O)-NH and HO-CH₂-C(=O)-NH-.

R²-G4a:

In another embodiment the group R² is selected from the group R²-G4a consisting of: F and -OCH₃.

5 **R²-G4b:**

In another embodiment the group R² is selected from the group R²-G4b consisting of: CH₃, CH₃-C(=O)-NH- and HO-CH₂-C(=O)-NH-.

L:10 **L-G1 :**

The group L is preferably selected from the group L-G1 as defined hereinbefore and hereinafter.

L-G2:

15 In one embodiment the group L is selected from the group L-G2 consisting of: a linear Ci-3-alkylene group that is optionally substituted with one or two CH₃.

L-G3:

In another embodiment the group L is selected from the group L-G3 consisting of: 20 a linear Ci-2-alkylene group that is optionally substituted with one CH₃.

L-G4:

In another embodiment the group L is selected from the group L-G4 consisting of: -CH(CH₃)- and -CH(CH₃)-(CH₂)-.

25

L-G5:

In another embodiment the group L is selected from the group L-G5 consisting of: -CH(CH₃)- and -CH(CH₃)-(CH₂)-,
wherein the -CH(CH₃)- moiety of the -CH(CH₃)-(CH₂)- group is attached to the
30 N atom in formula (I) and the -CH₂-* moiety of the -CH(CH₃)-(CH₂)- group is attached to the Ar² group.

L-G6:

In another embodiment the group L is selected from the group L-G6 consisting of

-CH(CH₃)-.

L-G6a:




According an embodiment L-G6a the group L is , embracing and



5 , wherein the right-hand side of each of the before-mentiones moieties is attached to the N atom in formula (I) and the left-hand side of each of the before-mentiones moieties is attached to the Ar² group.


A preferred example of the group L-G6a is



10

wherein the right-hand side of each of the before-mentiones moieties is attached to the N atom in formula (I) and the left-hand side of each of the before-mentiones moieties is attached to the Ar² group.

15 Another preferred example of the group L-G6a is



wherein the right-hand side of each of the before-mentiones moieties is attached to the N atom in formula (I) and the left-hand side of each of the before-mentiones moieties is attached to the Ar² group.

20

L-G7:

In another embodiment the group L is selected from the group L-G7 consisting of:


-CH(CH₃)-(CH₂)-,

wherein the -CH(CH₃)- moiety of the -CH(CH₃)-(CH₂)- group is attached to the

25 N atom in formula (I) and the -CH₂- moiety of the -CH(CH₃)-(CH₂)- group is attached to the Ar² group.

L-G7a:

In another embodiment the group L is selected from the group L-G7a consisting of:




wherein the $-\text{CH}(\text{CH}_3)-$ moiety of the above group is attached to the N atom in formula (I) and the $-\text{CH}_2-$ moiety of the above group is attached to the Ar^2 group.

5

L-G7b:

In another embodiment the group L is selected from the group L-G7b consisting of:



10

wherein the $-\text{CH}(\text{CH}_3)-$ moiety of the above group is attached to the N atom in formula (I) and the $-\text{CH}_2-$ moiety of the above group is attached to the Ar^2 group.

X:

15 **X-G1 :**

The group X is preferably selected from the group X-G1 as defined hereinbefore and hereinafter.

X-G2:

20 In another embodiment the group X is selected from the group X-G2 consisting of O.

X-G3:

In another embodiment the group X is selected from the group X-G3 consisting of S.

25 **R:**

R-G1 :

The group R is preferably selected from the group $\text{R}^N\text{-G1}$ as defined hereinbefore and hereinafter.

30 **R-G2:**

In another embodiment the group R is selected from the group R^N-G2 consisting of H and methyl.

R-G3:

5 In another embodiment the group R is selected from the group R^N-G3 consisting of H.

T:**T-G1:**

10 The group T is preferably selected from the group T-G1 as defined hereinbefore and hereinafter.

T-G2:

In another embodiment the group T is selected from the group T-G2 consisting of:

15 Ci-6-alkyl, C2-6-alkenyl, C2-6-alkynyl, C3-7-cycloalkyl, C3-7-cycloalkenyl, C₃₋₇-cycloalkyl-Ci-3-alkyl, Ci-6-alkyl-O-, C3-7-cycloalkyl-O-, C3-7-cycloalkyl-Ci-3-alkyl-O-, Ci-6-alkyl-S-, Cs-7-cycloalkyl-S-, C₃₋₇-cycloalkyl-Ci-3-alkyl-S-, Ci-4-alkyl-C(=O)-, Ci-4-alkyl-S(=O)₂-, R^{T1}R^{T2}-N-, R^{T1}R^{T2}-N-Ci-3-alkyl-, R^{T1}R^{T2}-N-CO-, Ci-4-alkyl-C(=O)-R^{T2}N-Ci-3-alkyl, heterocyclyl, phenyl and heteroaryl;

20

wherein in each cycloalkyl and heterocyclyl a CH₂-group may optionally be replaced by -C(=O)-; and

25 wherein each cycloalkyl and heterocyclyl may be optionally substituted with one or more Ci-4-alkyl, which may be optionally substituted with one or more substituents R^c; and

wherein each alkyl, cycloalkyl and heterocyclyl may be optionally substituted with one or more substituents R^c; and

30

wherein each alkyl may optionally be substituted with a heteroaryl group; and

wherein heterocyclyl is defined as hereinbefore and hereinafter; preferably heterocyclyl is azetidinyl, pyrrolidinyl, piperidinyl, morpholinyl, oxetanyl, tetrahydrofuranyl or tetrahydropyranyl; and

5 wherein each heteroaryl is selected from the group consisting of phenyl, pyridyl, pyrimidinyl, pyridazinyl, pyrazinyl, furanyl, thienyl, pyrrolyl, imidazolyl, pyrazolyl, triazolyl, oxazolyl, isoxazolyl, isothiazolyl and thiazolyl; and

wherein each aryl and heteroaryl group may be optionally substituted with one or
10 more substituents \mathbf{R}^2 .

T-G3:

In another embodiment the group T is selected from the group T-G3 consisting of:


\mathbf{C}_{1-4} -alkyl, \mathbf{C}_{2-4} -alkenyl, c3-6-cycloalkyl, c3-7-cycloalkenyl, c3-5-cycloalkyl- \mathbf{C}_{1-3} -alkyl-,
15 heteroaryl- \mathbf{C}_{1-3} -s-alkyl-, \mathbf{C}_{1-3} -alkyl-O-, $\mathbf{R}^{T1}\mathbf{R}^{T2}$ -N-, \mathbf{C}_{1-3} -alkyl-C(=O)- \mathbf{R}^{T2} N- \mathbf{C}_{1-3} -alkyl-,
heterocyclyl, phenyl and heteroaryl,

wherein in each heterocyclyl, a \mathbf{CH}_2 -group may optionally be replaced by -C(=O)-;
and

20 wherein each cycloalkyl and heterocyclyl may be optionally substituted with one or
more \mathbf{C}_{1-3} -alkyl, which may be optionally substituted with one or more substituents
 \mathbf{R}^c ; and

25 wherein each alkyl, cycloalkyl and heterocyclyl may be optionally substituted with one
or more F; and

wherein heterocyclyl is defined as hereinbefore and hereinafter, preferably
heterocyclyl is selected from the group consisting of



wherein heteroaryl is selected from the group consisting of pyridyl, pyrimidinyl,

5 pyridazinyl, pyrazinyl, furanyl, thienyl, pyrrolyl, imidazolyl, pyrazolyl, triazolyl, oxazolyl, isoxazolyl, thiazolyl and isothiazolyl; and

wherein each phenyl and heteroaryl group may be optionally substituted with one or more substituents R^2 .

10

T-G4:

In another embodiment the group T is selected from the group T-G4 consisting of:


Ci_{-4} -alkyl, C_{2-4} -alkenyl, C_{3-5} -cycloalkyl, C_{5-6} -cycloalkenyl, C_{3-5} -cycloalkyl- CH_2 -, Ci_{-3} -alkyl-O-, $R^{T1}R^{T2}$ -N-, Ci_{-3} -alkyl-C(=O)-R T2 N- Ci_{-3} -alkyl, heterocyclyl, phenyl and

15 heteroaryl,

wherein R^{T1} is H, CH_3 or CH_2CH_3 ; and R^{T2} is H or CH_3 ; and

wherein each alkyl, cycloalkyl and heterocyclyl may be optionally substituted with one 20 F, CN, CH_3 , CF_3 , cyclopropyl, -NH $_2$, -N(CH_3) $_2$, pyridinyl, OH or -OCH $_3$; and

wherein heterocyclyl is selected from the group consisting of:



25 wherein heteroaryl is selected from the group consisting of pyridyl, pyrimidinyl, pyridazinyl, pyrazinyl, furanyl, thienyl, pyrrolyl, imidazolyl, pyrazolyl, oxazolyl, isoxazolyl, thiazolyl and isothiazolyl; and

wherein each phenyl and heteroaryl group may be optionally substituted with one or two substituents independently selected from CH_3 , $-\text{NH}-\text{CO}-\text{CH}_3$, $-\text{NH}-\text{CO}-\text{CH}_2-\text{OH}$.


T-G5:

5 In another embodiment the group T is selected from the group T-G5 consisting of:
 $\text{C}_{1-4}\text{-alkyl}$, $\text{C}_{2-4}\text{-alkenyl}$, $\text{C}_{3-5}\text{-cycloalkyl}$, $\text{C}_{5-6}\text{-cycloalkenyl}$, cyclopropyl- CH_2 -, CH_3O -, $\text{R}^{\text{T}1}\text{R}^{\text{T}2}\text{-N-}$, $\text{CH}_3\text{-C(=O)-NH-Ci-3-alkyl}$, heterocyclyl, phenyl and heteroaryl,


wherein $\text{R}^{\text{T}1}$ is H, CH_3 or CH_2CH_3 ; and $\text{R}^{\text{T}2}$ is H or CH_3 ; and

10 wherein each alkyl, cycloalkyl and heterocyclyl may be optionally substituted with one F, CN, CH_3 , CF_3 , cyclopropyl, $-\text{N}(\text{CH}_3)_2$, pyridinyl, OH or $-\text{OCH}_3$; and

wherein heterocyclyl is selected from the group consisting of:



and



15 ; and

wherein heteroaryl is selected from the group consisting of pyridyl, pyrimidinyl, pyridazinyl, furanyl, pyrazolyl, oxazolyl, isoxazolyl, thiazolyl and isothiazolyl; and

20 wherein each phenyl and heteroaryl group may be optionally substituted with one or two substituents independently selected from CH_3 , $-\text{NH}-\text{CO}-\text{CH}_3$, $-\text{NH}-\text{CO}-\text{CH}_2-\text{OH}$.

T-G5a:

In another embodiment the group T is selected from the group T-G5a consisting of:

25 CH_3 , cyclopropyl and $-\text{N}(\text{CH}_3)_2$,

wherein the CH_3 group may optionally be substituted with one F, CN, CH_3 , CF_3 , cyclopropyl, $-\text{N}(\text{CH}_3)_2$, pyridinyl, OH, or $-\text{OCH}_3$.

30 **T-G5b:**

In another embodiment the group T is selected from the group T-G5a consisting of:

CH_3 and $-\text{N}(\text{CH}_3)_2$,

wherein the CH_3 group may optionally be substituted with one F, CN, CH_3 , CF_3 , cyclopropyl, $-\text{N}(\text{CH}_3)_2$, pyridinyl, OH, or $-\text{OCH}_3$.

5 **T-G6:**

In another embodiment the group T is selected from the group T-G6 consisting of: CH_3 optionally substituted with one F, CN, CH_3 , CF_3 , cyclopropyl, OH or $-\text{OCH}_3$.

T-G6a:

10 In another embodiment the group T is selected from the group T-G6a consisting of: CH_3 .

R^{T1}

$\mathbf{R}^{\mathbf{T}1}\text{-G1}$:

15 The group $\mathbf{R}^{\mathbf{T}1}$ is preferably selected from the group $\mathbf{R}^{\mathbf{T}1}\text{-G1}$ consisting of H, **Ci-6**-alkyl, **C₃₋₆-cycloalkyl**, **C₃₋₆-cycloalkyl-Ci₁₋₃-alkyl**, phenyl and pyridyl, wherein each cycloalkyl may be optionally substituted with one or more **Ci₁₋₄-alkyl**, and

20 wherein each alkyl and cycloalkyl may be optionally substituted with one or more substituents selected from F, and

wherein each alkyl and cycloalkyl may be optionally substituted with a substituent selected from OH, **Ci₁₋₃-alkyl-O-**, **Ci₁₋₃-alkyl-C(=O)-**, **HO-C(=O)-** and **Ci₁₋₄-alkyl-O-C(=O)-**; and

25 wherein each phenyl and pyridyl group may be optionally substituted with one or more substituents \mathbf{R}^2 .

$\mathbf{R}^{\mathbf{T}1}\text{-G2}$:

30 In another embodiment the group $\mathbf{R}^{\mathbf{T}1}$ is selected from the group $\mathbf{R}^{\mathbf{T}1}\text{-G2}$ consisting of:

H, **Ci₁₋₄-alkyl**, phenyl and pyridinyl.

$\mathbf{R}^{\mathbf{T}1}\text{-G3}$:

In another embodiment the group $\mathbf{R}^{\mathbf{T}1}$ is selected from the group $\mathbf{R}^{\mathbf{T}1}\text{-G3}$ consisting of

H, methyl and ethyl.

R^{T2}

5 **R^{T2-G1} :**

The group R^{T2} is preferably selected from the group R^{T2-G1} consisting of H and C₁₋₄-alkyl.

R^{T2-G2}:

10 In another embodiment the group R^{T2} is selected from the group R^{T2-G2} consisting of H and methyl.


15 Examples of preferred subgeneric embodiments according to the present invention are set forth in the following table, wherein each substituent group of each embodiment is defined according to the definitions set forth hereinbefore and wherein all other substituents of the formula (I) are defined according to the definitions set forth hereinbefore:







Embodi ment	Ar ¹ and R ¹	Ar ²	L	X	R	T
E-1	Ar ¹ -G1, R ¹ -G1	Ar ² -G1	L-G1	X-G1	R-G1	T-G1
E-2	Ar ¹ -G2, R ¹ -G1	Ar ² -G2	L-G1	X-G1	R-G1	T-G2
E-3	Ar ¹ -G2, R ¹ -G2	Ar ² -G3	L-G1	X-G1	R-G1	T-G3
E-4	Ar ¹ -G3, R ¹ -G2	Ar ² -G3a	L-G1	X-G1	R-G2	T-G3
E-5	Ar ¹ -G4, R ¹ -G3	Ar ² -G3	L-G1	X-G1	R-G2	T-G3
E-6	Ar ¹ -G4, R ¹ -G3	Ar ² -G3a	L-G2	X-G1	R-G2	T-G3
E-7	Ar ¹ -G4, R ¹ -G3	Ar ² -G4	L-G3	X-G2	R-G3	T-G4
E-8	Ar ¹ -G4, R ¹ -G3	Ar ² -G4	L-G3	X-G2	R-G3	T-G5
E-9	Ar ¹ -G4, R ¹ -G3	Ar ² -G4	L-G3	X-G2	R-G3	T-G5a







Embodi ment	Ar ¹ and R ¹	Ar ²	L	X	R	T
E-10	Ar ¹ -G4, R ¹ -G3	Ar ² -G4	L-G3	X-G2	R-G3	T-G6
E-11	Ar ¹ -G4, R ¹ -G3	Ar ² -G4	L-G3	X-G2	R-G3	T-G6a
E-12	Ar ¹ -G4, R ¹ -G4	Ar ² -G4a	L-G4	X-G2	R-G3	T-G5
E-13	Ar ¹ -G4, R ¹ -G4	Ar ² -G4a	L-G4	X-G2	R-G3	T-G5a
E-14	Ar ¹ -G4, R ¹ -G4	Ar ² -G4a	L-G4	X-G2	R-G3	T-G6
E-15	Ar ¹ -G4, R ¹ -G4	Ar ² -G4a	L-G4	X-G2	R-G3	T-G6a
E-16	Ar ¹ -G5, R ¹ -G4	Ar ² -G4	L-G3	X-G2	R-G2	T-G5
E-17	Ar ¹ -G5, R ¹ -G4	Ar ² -G4a	L-G5	X-G2	R-G2	T-G5
E-18	Ar ¹ -G5, R ¹ -G4	Ar ² -G4a	L-G6	X-G2	R-G2	T-G6a
E-19	Ar ¹ -G5, R ¹ -G4	Ar ² -G5	L-G3	X-G2	R-G2	T-G4
E-20	Ar ¹ -G5, R ¹ -G4	Ar ² -G5	L-G4	X-G2	R-G2	T-G5
E-21	Ar ¹ -G5, R ¹ -G4	Ar ² -G5	L-G6	X-G2	R-G2	T-G5
E-22	Ar ¹ -G5, R ¹ -G4	Ar ² -G5	L-G5	X-G2	R-G2	T-G5
E-23	Ar ¹ -G6	Ar ² -G4	L-G2	X-G2	R-G2	T-G3
E-24	Ar ¹ -G6	Ar ² -G4	L-G3	X-G2	R-G2	T-G4
E-25	Ar ¹ -G6	Ar ² -G4	L-G5	X-G2	R-G2	T-G4
E-26	Ar ¹ -G6	Ar ² -G4a	L-G2	X-G2	R-G2	T-G3
E-27	Ar ¹ -G6	Ar ² -G4a	L-G3	X-G2	R-G2	T-G4
E-28	Ar ¹ -G6	Ar ² -G4a	L-G3	X-G2	R-G2	T-G5
E-29	Ar ¹ -G6	Ar ² -G4a	L-G5	X-G2	R-G2	T-G5
E-30	Ar ¹ -G6	Ar ² -G5	L-G2	X-G2	R-G2	T-G3
E-31	Ar ¹ -G6	Ar ² -G5	L-G3	X-G2	R-G2	T-G4
E-32	Ar ¹ -G6	Ar ² -G5	L-G3	X-G2	R-G2	T-G5
E-33	Ar ¹ -G6	Ar ² -G5	L-G5	X-G2	R-G2	T-G5

Embodiment	Ar ¹ and R ¹	Ar ²	L	X	R	T
E-34	Ar ¹ -G6a	Ar ² -G4	L-G2	X-G2	R-G2	T-G3
E-35	Ar ¹ -G6a	Ar ² -G4a	L-G3	X-G2	R-G2	T-G4
E-36	Ar ¹ -G6a	Ar ² -G4a	L-G5	X-G2	R-G2	T-G4
E-37	Ar ¹ -G6a	Ar ² -G5	L-G3	X-G2	R-G2	T-G4
E-38	Ar ¹ -G6a	Ar ² -G5	L-G5	X-G2	R-G2	T-G4
E-39	Ar ¹ -G6a	Ar ² -G5	L-G6	X-G2	R-G2	T-G4

The following preferred embodiments of compounds of the formula (I) are described using generic formulas (1.1) to (1.8b), wherein any tautomers and stereoisomers, 5 solvates, hydrates and salts thereof, in particular the pharmaceutically acceptable salts thereof, are encompassed.



(I.2b)	
(I.2c)	
(I.3)	
(I.3a)	
(I.3b)	
(I.4)	

(I.4a)	
(I.4b)	
(I.5)	
(I.5a)	
(I.5b)	
(I.6)	

(I.6a)	
(I.6b)	
(I.7)	
(I.7a)	
(I.7b)	
(I.8)	

(I.8a)	
(I.8b)	

wherein in each of the above formulas (1.1) to (1.8b) the groups Ar¹, R¹, L and T are defined as hereinbefore and hereinafter; and

5 n is 0, 1, 2 or 3.

Preferred embodiments of the above formulas (1.1) to (1.8b) according to the present invention are set forth in the following table, wherein each group Ar¹, R¹, L and T of each embodiment is defined according to the definitions set forth hereinbefore and 10 wherein all other substituents of the formula (I) are defined according to the definitions set forth hereinbefore and n is 0, 1, 2 or 3:

Embodiment	Formula	Ar ¹	R ¹	n	L	T
E-A	(1.1)	Ar ¹ -G4	R ¹ -G3	1, 2 or 3	L-G2	T-G3
E-Aa	(1.1)	Ar ¹ -G4	R ¹ -G4	1 or 2	L-G3	T-G5
E-Ab	(1.1)	Ar ¹ -G5	R ¹ -G3	1, 2 or 3	L-G3	T-G4
E-B	(I.2)	-	R ¹ -G3	1, 2 or 3	L-G3	T-G3
E-Ba	(I.2)	-	R ¹ -G3	1 or 2	L-G5	T-G5
E-Bb	(I.2)	-	R ¹ -G4	1, 2 or 3	L-G3	T-G3
E-Bc	(I.2)	-	R ¹ -G4	1, 2 or 3	L-G3	T-G5
E-Bd	(I.2)	-	R ¹ -G4	1 or 2	L-G3	T-G4

Embodiment	Formula	Ar ¹	R ¹	n	L	T
E-Be	(1.2)	-	R ¹ -G4	1 or 2	L-G3	T-G5
E-Bf	(1.2)	-	R ¹ -G4	1 or 2	L-G3	T-G5a
E-Bg	(1.2)	-	R ¹ -G4	1 or 2	L-G3	T-G6
E-Bh	(1.2)	-	R ¹ -G4	1	L-G3	T-G4
E-Bi	(1.2)	-	R ¹ -G5	1	L-G3	T-G5
E-Bj	(1.2)	-	R ¹ -G5	1	L-G3	T-G5a
E-Bk	(1.2)	-	R ¹ -G5	1	L-G3	T-G6
E-C	(1.2a)	-	R ¹ -G3	1, 2 or 3	-	T-G4
E-Ca	(1.2a)	-	R ¹ -G3	1, 2 or 3	-	T-G5
E-Cb	(1.2a)	-	R ¹ -G4	1, 2 or 3	-	T-G4
E-Cc	(1.2a)	-	R ¹ -G4	1 or 2	-	T-G5
E-Cd	(1.2a)	-	R ¹ -G5	1 or 2	-	T-G4
E-D	(1.2a)	-	R ¹ -G5	1 or 2	-	T-G5
E-Da	(1.2b)	-	R ¹ -G5	1, 2 or 3	-	T-G4
E-Db	(1.2b)	-	R ¹ -G5	1 or 2	-	T-G5
E-E	(1.3)	-	R ¹ -G3	1 or 2	L-G3	T-G3
E-Ea	(1.3)	-	R ¹ -G4	1 or 2	L-G3	T-G4
E-Eb	(1.3)	-	R ¹ -G5a	1	L-G3	T-G5
E-F	(1.3a)	-	R ¹ -G3	1 or 2	-	T-G4
E-Fa	(1.3a)	-	R ¹ -G4	1 or 2	-	T-G4
E-Fb	(1.3a)	-	R ¹ -G4	1 or 2	-	T-G5
E-Fc	(1.3a)	-	R ¹ -G5a	1 or 2	-	T-G4
E-Fd	(1.3a)	-	R ¹ -G5a	1 or 2	-	T-G5
E-G	(1.3b)	-	R ¹ -G3	1	-	T-G4
E-Ga	(1.3b)	-	R ¹ -G4	1	-	T-G4
E-Gb	(1.3b)	-	R ¹ -G4	1	-	T-G5
E-Gc	(1.3b)	-	R ¹ -G5a	1	-	T-G4
E-Gd	(1.3b)	-	R ¹ -G5a	1	-	T-G5
E-H	(1.4)	-	R ¹ -G3	1 or 2	L-G3	T-G3
E-Ha	(1.4)	-	R ¹ -G4	1 or 2	L-G3	T-G4
E-Hb	(1.4)	-	R ¹ -G5a	1	L-G3	T-G5
E-I	(1.4a)	-	R ¹ -G3	1 or 2	-	T-G4
E-Ia	(1.4a)	-	R ¹ -G4	1 or 2	-	T-G4

Embodiment	Formula	Ar ¹	R ¹	n	L	T
E-lb	(1.4a)	-	R ¹ -G4	1 or 2	-	T-G5
E-lc	(1.4a)	-	R ¹ -G5a	1 or 2	-	T-G4
E-ld	(1.4a)	-	R ¹ -G5a	1 or 2	-	T-G5
E-J	(1.4b)	-	R ¹ -G3	1	-	T-G4
E-Ja	(1.4b)	-	R ¹ -G4	1	-	T-G4
E-Jb	(1.4b)	-	R ¹ -G4	1	-	T-G5
E-Jc	(1.4b)	-	R ¹ -G5a	1	-	T-G4
E-Jd	(1.4b)	-	R ¹ -G5a	1	-	T-G5
E-K	(1.5)	-	R ¹ -G3	1 or 2	L-G3	T-G3
E-Ka	(1.5)	-	R ¹ -G4	1 or 2	L-G3	T-G4
E-Kb	(1.5)	-	R ¹ -G5a	1	L-G3	T-G5
E-L	(1.5a)	-	R ¹ -G3	1 or 2	-	T-G4
E-La	(1.5a)	-	R ¹ -G4	1 or 2	-	T-G4
E-Lb	(1.5a)	-	R ¹ -G4	1 or 2	-	T-G5
E-Lc	(1.5a)	-	R ¹ -G5a	1 or 2	-	T-G4
E-Ld	(1.5a)	-	R ¹ -G5a	1 or 2	-	T-G5
E-M	(1.5b)	-	R ¹ -G3	1	-	T-G4
E-Ma	(1.5b)	-	R ¹ -G4	1	-	T-G4
E-Mb	(1.5b)	-	R ¹ -G4	1	-	T-G5
E-Mc	(1.5b)	-	R ¹ -G5a	1	-	T-G4
E-Md	(1.5b)	-	R ¹ -G5a	1	-	T-G5
E-N	(1.6)	-	R ¹ -G3	1 or 2	L-G3	T-G3
E-Na	(1.6)	-	R ¹ -G4	1	L-G3	T-G4
E-Nb	(1.6)	-	R ¹ -G5b	1	L-G3	T-G5
E-O	(1.6a)	-	R ¹ -G3	1 or 2	-	T-G4
E-Oa	(1.6a)	-	R ¹ -G4	1	-	T-G4
E-Ob	(1.6a)	-	R ¹ -G4	1	-	T-G5
E-Oc	(1.6a)	-	R ¹ -G5b	1	-	T-G4
E-Od	(1.6a)	-	R ¹ -G5b	1	-	T-G5
E-P	(1.6b)	-	R ¹ -G3	1	-	T-G4
E-Pa	(1.6b)	-	R ¹ -G4	1	-	T-G4
E-Pb	(1.6b)	-	R ¹ -G4	1	-	T-G5
E-Pc	(1.6b)	-	R ¹ -G5b	1	-	T-G4

Embodiment	Formula	Ar ¹	R ¹	n	L	T
E-Pd	(1.6b)	-	R ¹ -G5b	1	-	T-G5
E-Q	(1.7)	-	R ¹ -G3	1 or 2	L-G3	T-G3
E-Qa	(1.7)	-	R ¹ -G4	1	L-G3	T-G4
E-Qb	(1.7)	-	R ¹ -G5c	1	L-G3	T-G5
E-R	(1.7a)	-	R ¹ -G3	1 or 2	-	T-G4
E-Ra	(1.7a)	-	R ¹ -G4	1	-	T-G4
E-Rb	(1.7a)	-	R ¹ -G4	1	-	T-G5
E-Rc	(1.7a)	-	R ¹ -G5c	1	-	T-G4
E-Rd	(1.7a)	-	R ¹ -G5c	1	-	T-G5
E-S	(1.7b)	-	R ¹ -G3	1	-	T-G4
E-Sa	(1.7b)	-	R ¹ -G4	1	-	T-G4
E-Sb	(1.7b)	-	R ¹ -G4	1	-	T-G5
E-Sc	(1.7b)	-	R ¹ -G5c	1	-	T-G4
E-Sd	(1.7b)	-	R ¹ -G5c	1	-	T-G5
E-T	(1.8)	-	R ¹ -G3	1 or 2	L-G3	T-G3
E-Ta	(1.8)	-	R ¹ -G4	1	L-G3	T-G4
E-Tb	(1.8)	-	R ¹ -G5d	1	L-G3	T-G5
E-U	(1.8a)	-	R ¹ -G3	1 or 2	-	T-G4
E-Ua	(1.8a)	-	R ¹ -G4	1	-	T-G4
E-Ub	(1.8a)	-	R ¹ -G4	1	-	T-G5
E-Uc	(1.8a)	-	R ¹ -G5d	1	-	T-G4
E-Ud	(1.8a)	-	R ¹ -G5d	1	-	T-G5
E-V	(1.8b)	-	R ¹ -G3	1	-	T-G4
E-Va	(1.8b)	-	R ¹ -G4	1	-	T-G4
E-Vb	(1.8b)	-	R ¹ -G4	1	-	T-G5
E-Vc	(1.8b)	-	R ¹ -G5d	1	-	T-G4
E-Vd	(1.8b)	-	R ¹ -G5d	1	-	T-G5

including any tautomers and stereoisomers, solvates, hydrates and salts thereof, in particular the pharmaceutically acceptable salts thereof.

5 Preferred are those compounds of formula (I), wherein

- 40 -

Ar¹ is selected from the group consisting of phenyl, pyridinyl and pyrimidinyl,


wherein the above-mentioned phenyl and pyridinyl groups are each substituted with one to three groups independently selected from R¹, and

5

wherein the above-mentioned pyrimidinyl groups is substituted with one or two R¹, and

10 wherein two adjacent carbon atoms of a phenyl group may be linked to each other via a -O-CH₂-CH₂-O-, -O-CH₂-CH₂-C(CH₃)₂-, -O-CH₂-CH₂-CH₂-O-, or -O-CH₂-C(CH₃)₂-CH₂-O- bridge;

15 R¹ is selected from the group consisting of F, Cl, Br, CN, CF₃, Ci₄ alkyl, -O-Ci₄ alkyl, cyclopropyl, -O-C₃₋₅ cycloalkyl, -O-CH₂- cyclopropyl, and



15

Ar² is phenylene;

L is a linear Ci-2-alkylene group that is optionally substituted with one CH₃;

X is O;

20 R is H; and

25 T is selected from the group consisting of: Ci₄-alkyl, C₂₋₄-alkenyl, C₃₋₅-cycloalkyl, C₅₋₆-cycloalkenyl, cyclopropyl-CH₂-, CH₃O-, R^{T1}R^{T2}-N-, CH₃-C(=O)-NH-Ci₃-alkyl, heterocyclyl, phenyl and heteroaryl,


25

wherein R^{T1} is H, CH₃ or CH₂CH₃; and

R^{T2} is H or CH₃; and

30 wherein each alkyl, cycloalkyl and heterocyclyl may be optionally substituted with one F, CN, CH₃, CF₃, cyclopropyl, -N(CH₃)₂, pyridinyl, OH or -OCH₃; and

wherein heterocyclyl is selected from the group consisting of:



wherein heteroaryl is selected from the group consisting of pyridyl,

5 pyrimidinyl, pyridazinyl, furanyl, pyrazolyl, oxazolyl, isoxazolyl, thiazolyl and isothiazolyl; and


wherein each phenyl and heteroaryl group may be optionally substituted with one or two substituents independently selected from CH_3 , $-\text{NH}-\text{CO}-$


10 CH_3 , $-\text{NH}-\text{CO}-\text{CH}_2-\text{OH}$;

including any tautomers and stereoisomers thereof,
or a salt thereof.

15 Particularly preferred compounds, including their tautomers and stereoisomers, the salts thereof, or any solvates or hydrates thereof, are described in the experimental section hereinafter.

Specifically mentioned are the following compounds of the invention:






and their pharmaceutically acceptable salts.


The compounds according to the invention may be obtained using methods of synthesis which are known to the one skilled in the art and described in the literature of organic synthesis. Preferably the compounds are obtained analogously to the methods of preparation explained more fully hereinafter, in particular as described in the experimental section.

Synthesis Schemes


Compounds of general formula (I) may be prepared by palladium-mediated Buchwald reactions or copper-mediated Ullmann reactions of aryl halogenides or aryl triflates (II) with azetidines (III) wherein Z is a leaving group which for example denotes Cl, 5 Br, I or OTf (triflate).



Compounds of general formula (I) may alternatively be prepared by nucleophilic aromatic substitution reactions (SnAr) of aryl/heteroaryl halogenides or aryl/heteroaryl 10 triflates (II) with azetidines (III), wherein Z is a leaving group which for example denotes Cl, Br, I, S(=O)CH₃ or triflate.




Compounds of general formula (I) may be prepared by amide coupling reactions of 15 amines (IV) with carboxylic acids (V-I) mediated by coupling reagents such as eg 2-(1H-benzotriazol-1-yl)-1,1,3,3-tetramethyluronium tetrafluoroborat (TBTU), N-hydroxybenzotriazole (HOBT), 2-(1H-7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate (HATU) or N,N-carbonyldiimidazole (CDI).




Alternatively, compounds of general formula (I) may be prepared by amide coupling 20 reactions of amines (IV) with carboxylic acids chlorides (V-II) or carboxylic acid anhydrides (V-III).

- 44 -



Compounds of general formula (VI) may be prepared by nucleophilic aromatic substitution reactions (SnAr) of pyrimidines (VI I) with nucleophiles HR^1 (VI II), wherein
 5 Z is a leaving group which for example denotes Cl, Br, I, $S(=O)CH_3$ or triflate and wherein HR^1 is a nucleophile, such as for example an alcohol or an amine and wherein the reaction may be performed with other regioisomers of pyrimidine or other heteroaryls also. Alcohols may be deprotonated by an appropriate base before used as nucleophiles.


10




15

Compounds of general formula (IX) may be prepared by alkylation reactions of aromatic alcohols (X) with electrophiles (XI) wherein Z is a leaving group which for example denotes Cl, Br, I, mesylate, tosylate or triflate and R^3 is Ci_6 -alkyl, $C3_6$ -alkenyl, $C3_6$ -alkynyl, $C3_{10}$ -carbocyclyl, $C3_{10}$ -carbocyclyl- Ci_3 -alkyl, $R^{N1}R^{N2}N-C2_3$ -alkyl, heterocyclyl, heterocyclyl- Ci_3 -alkyl, aryl, aryl- Ci_3 -alkyl, heteroaryl or heteroaryl- Ci_3 -alkyl, wherein in each carbocyclyl and heterocyclyl a CH_2 -group may optionally be replaced by $-C(=O)-$ or $-C(=CR^{N2})-$; wherein each carbocyclyl and heterocyclyl may be optionally substituted with one or more Ci_4 -alkyl, which may be optionally substituted with one or more substituents R^c ; wherein each alkyl, carbocyclyl and heterocyclyl may be optionally substituted with one or more substituents R^c ; wherein each heterocyclyl may be optionally substituted with aryl or heteroaryl; and wherein each aryl and heteroaryl group may be optionally substituted with one or more substituents R^2 .

- 45 -




Compounds of general formula (XII) may be prepared by palladium-catalyzed Suzuki coupling reactions of aryl/heteroaryl halogenides or aryl/heteroaryl triflates (XIII) with 5 boronic acids (XIV) or trimethylboroxine wherein Z is a leaving group which for example denotes Cl, Br, I or triflate and R⁵ is C₂₋₆-alkenyl, C₂₋₆-alkynyl, aryl or heteroaryl, wherein each aryl and heteroaryl group may be optionally substituted with one or more substituents R².




10

Compounds of general formula (XV) may be prepared by urea forming reactions such as reaction of amines (IV) with amines (XVI) after reaction with reagents such as N,N-carbonylditriazole (CDT) or N,N-carbonyldiimidazole (CDI).



15

Alternatively, compounds of general formula (XV) may be prepared by urea forming reactions such as reaction of amines (IV) with carbamoyl chlorides (XVII) or isocyanates (XVIII).




20


Compounds of general formula (XX), wherein R⁸ is Ci-6-alkyl, C3-*io*-carbocyclyl or C3-*io*-carbocyclyl-Ci-3-alkyl, may be prepared by urethane forming reactions such as

- 46 -


reaction of amines (IV) with alcohols (IXX) after reaction with reagents such as CDT or CDI. Alcohols may be used in their deprotonated form.




5 Alternatively, compounds of general formula (XX) may be prepared by urethane forming reactions such as reaction of amines (IV) with chloro formates (XXI).




Compounds of general formula (I) may be prepared by alkylation reactions of aromatic alcohols (XXIII) with electrophiles (XXII) wherein Z is a leaving group such as for example Cl, Br, I, mesylate, tosylate or triflate



Alternatively compounds of general formula (I) may be prepared by a Mitsunobu reaction of aromatic alcohols (XXIII) with alcohols (XXIV).



Compounds of general formula XXII, wherein X is S, may be prepared by analogous reactions as described before for X = O, except for the Mitsunobu reaction.



Terms and definitions

Terms not specifically defined herein should be given the meanings that would be given to them by one of skill in the art in light of the disclosure and the context. As used in the specification, however, unless specified to the contrary, the following 5 terms have the meaning indicated and the following conventions are adhered to.

The terms "compound(s) according to this invention", "compound(s) of formula (I)", "compound(s) of the invention" and the like denote the compounds of the formula (I) according to the present invention including their tautomers, stereoisomers and 10 mixtures thereof and the salts thereof, in particular the pharmaceutically acceptable salts thereof, and the solvates and hydrates of such compounds, including the solvates and hydrates of such tautomers, stereoisomers and salts thereof.

The terms "treatment" and "treating" embraces both preventative, i.e. prophylactic, or 15 therapeutic, i.e. curative and/or palliative, treatment. Thus the terms "treatment" and "treating" comprise therapeutic treatment of patients having already developed said condition, in particular in manifest form. Therapeutic treatment may be symptomatic treatment in order to relieve the symptoms of the specific indication or causal treatment in order to reverse or partially reverse the conditions of the indication or to 20 stop or slow down progression of the disease. Thus the compositions and methods of the present invention may be used for instance as therapeutic treatment over a period of time as well as for chronic therapy. In addition the terms "treatment" and "treating" comprise prophylactic treatment, i.e. a treatment of patients at risk to develop a condition mentioned hereinbefore, thus reducing said risk.

25

When this invention refers to patients requiring treatment, it relates primarily to treatment in mammals, in particular humans.

The term "therapeutically effective amount" means an amount of a compound of the 30 present invention that (i) treats or prevents the particular disease or condition, (ii) attenuates, ameliorates, or eliminates one or more symptoms of the particular disease or condition, or (iii) prevents or delays the onset of one or more symptoms of the particular disease or condition described herein.

The terms "modulated" or "modulating", or "modulate(s)", as used herein, unless otherwise indicated, refers to the inhibition of acetyl-CoA carboxylase(s) (ACC)(s) with one or more compounds of the present invention.

5 The terms "mediated" or "mediating" or "mediate", as used herein, unless otherwise indicated, refers to the (i) treatment, including prevention the particular disease or condition, (ii) attenuation, amelioration, or elimination of one or more symptoms of the particular disease or condition, or (iii) prevention or delay of the onset of one or more symptoms of the particular disease or condition described herein.

10

The term "substituted" as used herein, means that any one or more hydrogens on the designated atom, radical or moiety is replaced with a selection from the indicated group, provided that the atom's normal valence is not exceeded, and that the substitution results in an acceptably stable compound.

15

In the groups, radicals, or moieties defined below, the number of carbon atoms is often specified preceding the group, for example, **Ci-6-alkyl** means an alkyl group or radical having 1 to 6 carbon atoms. In general, for groups comprising two or more subgroups, the last named subgroup is the radical attachment point, for example, the 20 substituent "aryl-**Ci-3-alkyl**-" means an aryl group which is bound to a **Ci-3-alkyl**- group, the latter of which is bound to the core or to the group to which the substituent is attached.


In case a compound of the present invention is depicted in form of a chemical name 25 and as a formula in case of any discrepancy the formula shall prevail.

An asterisk is may be used in sub-formulas to indicate the bond which is connected to the core molecule as defined.

30 The numeration of the atoms of a substituent starts with the atom which is closest to the core or to the group to which the substituent is attached.


For example, the term "3-carboxypropyl-group" represents the following substituent:

- 49 -



wherein the carboxy group is attached to the third carbon atom of the propyl group.

The terms "1-methylpropyl-", "2,2-dimethylpropyl-" or "cyclopropylmethyl-" group represent the following groups:



The asterisk may be used in sub-formulas to indicate the bond which is connected to the core molecule as defined.

10 In a definition of a group the term "wherein each X, Y and Z group is optionally substituted with" and the like denotes that each group X, each group Y and each group Z either each as a separate group or each as part of a composed group may be substituted as defined. For example a definition " R^{ex} denotes H, **Ci**-3-alkyl, c3-6-cycloalkyl, c3-6-cycloalkyl -**Ci**-3-alkyl or **Ci**-3-alkyl-O-, wherein each alkyl group is

15 optionally substituted with one or more L^{ex} ." or the like means that in each of the beforementioned groups which comprise the term alkyl, i.e. in each of the groups **Ci**-3-alkyl, c3-6-cycloalkyl -**Ci**-3-alkyl and **Ci**-3-alkyl-O-, the alkyl moiety may be substituted with L^{ex} as defined.

20 In the following the term bicyclic includes spirocyclic.

Unless specifically indicated, throughout the specification and the appended claims, a given chemical formula or name shall encompass tautomers and all stereo, optical and geometrical isomers (e.g. enantiomers, diastereomers, E/Z isomers etc ..) and

25 racemates thereof as well as mixtures in different proportions of the separate enantiomers, mixtures of diastereomers, or mixtures of any of the foregoing forms where such isomers and enantiomers exist, as well as salts, including pharmaceutically acceptable salts thereof and solvates thereof such as for instance hydrates including solvates of the free compounds or solvates of a salt of the

30 compound.

The phrase "pharmaceutically acceptable" is employed herein to refer to those compounds, materials, compositions, and/or dosage forms which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of 5 human beings and animals without excessive toxicity, irritation, allergic response, or other problem or complication, and commensurate with a reasonable benefit/risk ratio.

As used herein, "pharmaceutically acceptable salts" refer to derivatives of the 10 disclosed compounds wherein the parent compound is modified by making acid or base salts thereof. Examples of pharmaceutically acceptable salts include, but are not limited to, mineral or organic acid salts of basic residues such as amines; alkali or organic salts of acidic residues such as carboxylic acids; and the like.

The pharmaceutically acceptable salts of the present invention can be synthesized 15 from the parent compound which contains a basic or acidic moiety by conventional chemical methods.

Salts of other acids than those mentioned above which for example are useful for purifying or isolating the compounds of the present invention are also part of the invention.

20

The term halogen generally denotes fluorine, chlorine, bromine and iodine.

The term " Cl_{-n} -alkyl", wherein n is an integer from 1 to n, either alone or in combination with another radical denotes an acyclic, saturated, branched or linear 25 hydrocarbon radical with 1 to n C atoms. For example the term Cl_{-5} -alkyl embraces the radicals $\text{H}_3\text{C}-$, $\text{H}_3\text{C}-\text{CH}_2-$, $\text{H}_3\text{C}-\text{CH}_2-\text{CH}_2-$, $\text{H}_3\text{C}-\text{CH}(\text{CH}_3)-$, $\text{H}_3\text{C}-\text{CH}_2-\text{CH}_2-\text{CH}_2-$, $\text{H}_3\text{C}-\text{CH}_2-\text{CH}(\text{CH}_3)-$, $\text{H}_3\text{C}-\text{CH}(\text{CH}_3)-\text{CH}_2-$, $\text{H}_3\text{C}-\text{C}(\text{CH}_3)_2-$, $\text{H}_3\text{C}-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{CH}_2-$, $\text{H}_3\text{C}-\text{CH}_2-\text{CH}_2-\text{CH}(\text{CH}_3)-$, $\text{H}_3\text{C}-\text{CH}_2-\text{CH}(\text{CH}_3)-\text{CH}_2-$, $\text{H}_3\text{C}-\text{CH}(\text{CH}_3)-\text{CH}_2-\text{CH}_2-$, $\text{H}_3\text{C}-\text{CH}_2-\text{C}(\text{CH}_3)_2-$, $\text{H}_3\text{C}-\text{C}(\text{CH}_3)_2-\text{CH}_2-$, $\text{H}_3\text{C}-\text{CH}(\text{CH}_3)-\text{CH}(\text{CH}_3)-$ and $\text{H}_3\text{C}-\text{CH}_2-$ 30 $\text{CH}(\text{CH}_2\text{CH}_3)-$.

The term " Cl_{-n} -alkylene" wherein n is an integer 1 to n, either alone or in combination with another radical, denotes an acyclic, straight or branched chain divalent alkyl radical containing from 1 to n carbon atoms. For example the term Cl_{-4} -alkylene

includes $-(CH_2)-$, $-(CH_2-CH_2)-$, $-(CH(CH_3))-$, $-(CH_2-CH_2-CH_2)-$, $-(C(CH_3)_2)-$, $-(CH(CH_2CH_3))-$, $-(CH(CH_3)-CH_2)-$, $-(CH_2-CH(CH_3))-$, $-(CH_2-CH_2-CH_2-CH_2)-$, $-(CH_2-CH_2-CH(CH_3))-$, $-(CH(CH_3)-CH_2-CH_2)-$, $-(CH_2-CH(CH_3)-CH_2)-$, $-(CH_2-C(CH_3)_2)-$, $-(C(CH_3)_2-CH_2)-$, $-(CH(CH_3)-CH(CH_3))-$, $-(CH_2-CH(CH_2CH_3))-$, $-(CH(CH_2CH_3)-CH_2)-$, $-(CH(CH_2CH_2CH_3))-$, $-(CHCH(CH_3)_2)-$ and $-C(CH_3)(CH_2CH_3)-$.

5

The term " C_{2-n} -alkenyl", is used for a group as defined in the definition for " C_{1-n} -alkyl" with at least two carbon atoms, if at least two of those carbon atoms of said group are bonded to each other by a double bond. For example the term C_{2-3} -alkenyl

10 includes $-CH=CH_2$, $-CH=CH-CH_3$, $-CH_2-CH=CH_2$.

The term " C_{2-n} -alkenylene" is used for a group as defined in the definition for " C_{1-n} -alkylene" with at least two carbon atoms, if at least two of those carbon atoms of said group are bonded to each other by a double bond. For example the term C_{2-3} -alkenylene includes $-CH=CH-$, $-CH=CH-CH_2-$, $-CH_2-CH=CH-$.

15

The term " C_{2-n} -alkynyl", is used for a group as defined in the definition for " C_{1-n} -alkyl" with at least two carbon atoms, if at least two of those carbon atoms of said group are bonded to each other by a triple bond. For example the term C_{2-3} -alkynyl

20 includes $-C\equiv CH$, $-C\equiv C-CH_3$, $-CH_2-C\equiv CH$.

The term " C_{2-n} -alkynylene" is used for a group as defined in the definition for " C_{1-n} -alkylene" with at least two carbon atoms, if at least two of those carbon atoms of said group are bonded to each other by a triple bond. For example the term C_{2-3} -alkynylene includes $-C\equiv C-$, $-C\equiv C-CH_2-$, $-CH_2-C\equiv C-$.

25

The term " C_{3-n} -carbocyclyl" as used either alone or in combination with another radical, denotes a monocyclic, bicyclic or tricyclic, saturated or unsaturated hydrocarbon radical with 3 to n C atoms. The hydrocarbon radical is preferably nonaromatic. Preferably the 3 to n C atoms form one or two rings. In case of a bicyclic or tricyclic ring system the rings may be attached to each other via a single bond or may be fused or may form a spirocyclic or bridged ring system. For example the term $C_{3-}io$ -carbocyclyl includes $C_{3-}io$ -cylcoalkyl, $C_{3-}io$ -cycloalkenyl,

30

octahydropentalenyl, octahydroindenyl, decahydronaphthyl, indanyl, tetrahydronaphthyl. Most preferably the term C_{3-n} -carbocyclyl denotes C_{3-n} -cycloalkyl, in particular C_{3-7} -cycloalkyl.

5 The term " C_{3-n} -cycloalkyl", wherein n is an integer 4 to n, either alone or in combination with another radical denotes a cyclic, saturated, unbranched hydrocarbon radical with 3 to n C atoms. The cyclic group may be mono-, bi-, tri- or spirocyclic, most preferably monocyclic. Examples of such cycloalkyl groups include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl, cyclononyl, 10 cyclododecyl, bicyclo[3.2.1.]octyl, spiro[4.5]decyl, norpinyl, norbonyl, norcaryl, adamantlyl, etc.


The term " C_{3-n} -cycloalkenyl", wherein n is an integer 3 to n, either alone or in combination with another radical, denotes a cyclic, unsaturated but nonaromatic, 15 unbranched hydrocarbon radical with 3 to n C atoms, at least two of which are bonded to each other by a double bond. For example the term C_{3-7} -cycloalkenyl includes cyclopropenyl, cyclobutenyl, cyclopentenyl, cyclopentadienyl, cyclohexenyl, cyclohexadienyl, cycloheptenyl, cycloheptadienyl and cycloheptatrienyl.


20 The term "aryl" as used herein, either alone or in combination with another radical, denotes a carbocyclic aromatic monocyclic group containing 6 carbon atoms which may be further fused to a second 5- or 6-membered carbocyclic group which may be aromatic, saturated or unsaturated. Aryl includes, but is not limited to, phenyl, indanyl, indenyl, naphthyl, anthracenyl, phenanthrenyl, tetrahydronaphthyl and 25 dihydronaphthyl. More preferably the term "aryl" as used herein, either alone or in combination with another radical, denotes phenyl or naphthyl, most preferably phenyl.


The term "heterocyclyl" means a saturated or unsaturated mono-, bi-, tri- or 30 spirocyclic, preferably mono- or bicyclic-ring system containing one or more heteroatoms selected from N, O or $S(O)_r$ with r=0, 1 or 2, which in addition may have a carbonyl group. More preferably the term "heterocyclyl" as used herein, either alone or in combination with another radical, means a saturated or unsaturated, even more preferably a saturated mono- or bicyclic-ring system containing 1, 2, 3 or 4

heteroatoms selected from N, O or S(O)_r with r=0, 1 or 2 which in addition may have a carbonyl group. The term "heterocydyl" is intended to include all the possible isomeric forms. Examples of such groups include aziridinyl, oxiranyl, azetidinyl, oxetanyl, pyrrolidinyl, tetrahydrofuranyl, piperidinyl, tetrahydropyranyl, azepanyl, 5 piperazinyl, morpholinyl, tetrahydrofuranonyl, tetrahydropyranonyl, pyrrolidinonyl, piperidinonyl, piperazinonyl, morpholinonyl.

Thus, the term "heterocydyl" includes the following exemplary structures which are not depicted as radicals as each form may be attached through a covalent bond to 10 any atom so long as appropriate valences are maintained:








The term "heteroaryl" means a mono- or polycyclic, preferably mono- or bicyclic-ring system containing one or more heteroatoms selected from N, O or S(O)_r with r=0, 1 or 2 wherein at least one of the heteroatoms is part of an aromatic ring, and wherein said ring system may have a carbonyl group. More preferably the term "heteroaryl"


5 as used herein, either alone or in combination with another radical, means a mono- or bicyclic-ring system containing 1, 2, 3 or 4 heteroatoms selected from N, O or S(O)_r with r=0, 1 or 2 wherein at least one of the heteroatoms is part of an aromatic ring, and wherein said ring system may have a carbonyl group. The term "heteroaryl" is intended to include all the possible isomeric forms.

10

Thus, the term "heteroaryl" includes the following exemplary structures which are not depicted as radicals as each form may be attached through a covalent bond to any atom so long as appropriate valences are maintained:



15



Many of the terms given above may be used repeatedly in the definition of a formula or group and in each case have one of the meanings given above, independently of one another.

5

Pharmacological Activity

The activity of the compounds of the invention may be demonstrated using the following ACC2 assay:

10 **Spectrophotometric 384 well assay**

Malonyl CoA formation by acetyl CoA carboxylases is stoichiometrically linked to the consumption of ATP. ACC2 activity is measured in a NADH-linked kinetic method measuring ADP generated during the ACC reaction using a coupled lactate dehydrogenase / pyruvate kinase reaction.

15 For biological testing, a human ACC2 construct which lacks the 128 amino acids at the N-terminus for increased solubility (nt 385-6966 in Genbank entry AJ575592) is cloned. The protein is then expressed in insect cells using a baculoviral expression system. Protein purification is performed by anion exchange.

20 All compounds are dissolved in dimethyl sulfoxide (DMSO) to a concentration of 10 mM.

Assay reactions are then carried out in 384-well plates, with hACC2 in an appropriate dilution and at final assay concentrations (f.c.) of 100 mM Tris (pH 7.5), 10 mM

trisodium citrate, 25 mM KHCO₃, 10 mM MgCl₂, 0.5 mg/ml BSA, 3.75 mM reduced L-glutathione, 15 U/ml lactate dehydrogenase, 0.5 mM phosphoenolpyruvate, 15 U/ml pyruvate kinase, compounds at different concentrations at final DMSO concentrations of 1%.

5 The enzymatic reaction is then started by addition of a mixture of NADH, acetyl Coenzyme A (both 200 μ M f.c.) and ATP (500 μ M f.c.). The decrease of the optical density (slope S) is then determined at 25°C at a wavelength of 340 nm over 15 minutes in a spectrophotometric reader.

Each assay microtiter plate contains wells with vehicle instead of compound as 10 controls for the non-inhibited enzyme (100% CTL; 'HIGH') and wells without acetyl-CoA as controls for non-specific NADH degradation (0% CTL; 'LOW').

The slope S is used for calculation of %CTL= $(S(\text{compound})-S(\text{'LOW'}))/(S(\text{'HIGH'})-S(\text{'LOW'})) * 100$. Compounds will give values between 100%CTL (no inhibition) and 0%CTL (complete inhibition).

15 For IC₅₀ value determination, the sample slope in the presence of the test compound after subtraction of the low controls (S(compound)-S('LOW')) are used.

An IC₅₀ value is derived from the compound slopes at different dosages after subtraction of the low controls (S(compound)-S('LOW')) by non-linear regression curve fitting (equation $y = (A + ((B-A)/(1 + ((C/x)^D))))$).

20

The compounds of general formula (I) according to the invention for example have IC₅₀ values below 15000 nM, particularly below 1000 nM, preferably below 300 nM.

In the following table the activity expressed as IC₅₀ (μ M) of compounds according to 25 the invention is presented wherein the IC₅₀ values are determined in the ACC2 assay as described hereinbefore. The term "Ex." refers to the example numbers according to the following experimental section.

Ex.	IC ₅₀ [μ M]	Ex.	IC ₅₀ [μ M]	Ex.	IC ₅₀ [μ M]	Ex.	IC ₅₀ [μ M]
1.1	0.18	1.51	0.27	4.18	1.80	8.16	0.70
1.2	0.24	1.52	0.09	4.19	0.20	8.17	2.79
1.3	0.53	1.53	0.76	4.20	0.77	8.18	0.74

1.4	0.1 3	1.54	0.04	4.21	0.41	8.1 9	0.69
1.5	0.20	1.55	0.1 1	4.22	0.45	8.20	0.48
1.6	0.08	1.56	0.1 5	4.23	1.18	8.21	0.30
1.7	1.25	1.57	0.74	4.24	0.39	8.22	1.87
1.8	0.55	1.58	1.39	4.25	0.75	8.23	5.79
1.9	0.56	1.59	0.30	4.26	13.34	8.24	3.94
1.10	0.67	1.60	0.65	4.27	0.57	8.25	0.53
1.11	2.21	1.61	2.54	4.28	0.36	8.26	2.78
1.12	5.34	1.62	2.91	4.29	1.17	8.27	0.76
1.13	0.07	2.1	0.88	4.30	1.38	8.28	0.1 7
1.14	0.07	2.2	0.06	4.31	0.1 9	8.29	0.41
1.15	0.1 8	2.3	0.24	4.32	0.1 0	8.30	0.67
1.16	0.45	2.4	6.14	4.33	1.73	8.31	1.09
1.17	1.89	2.5	0.1 7	4.34	0.94	8.32	0.55
1.18	0.14	2.6	0.09	4.35	1.11	8.33	0.1 9
1.19	0.09	2.7	2.06	4.36	1.14	8.34	0.81
1.20	0.49	2.8	4.1 0	4.37	0.29	8.35	3.1 6
1.21	0.1 9	2.9	10.63	4.38	3.05	8.36	3.04
1.22	1.18	2.1 0	0.14	4.39	0.57	8.37	2.06
1.23	2.81	2.1 1	2.55	4.39	0.60	8.38	1.26
1.24	0.1 2	2.1 2	4.50	4.40	1.05	8.39	5.20
1.25	0.58	3.1	0.21	4.41	3.48	8.40	0.35
1.26	0.74	3.2	1.12	4.42	0.53	8.41	0.51
1.27	1.75	3.3	0.32	4.43	0.53	8.42	1.69
1.28	1.62	3.4	1.47	5.1	0.1 7	8.43	2.34
1.29	0.1 9	3.5	0.20	5.2	2.1 8	8.44	0.93
1.30	0.67	3.6	3.88	6.1	0.09	8.45	2.31
1.31	0.1 9	3.7	0.47	6.2	1.74	8.46	1.19
1.32	0.1 9	3.8	0.24	6.3	0.1 5	8.47	1.91
1.33	0.20	3.9	3.82	7.1	1.44	9	1.10
1.34	0.34	4.1	0.46	7.2	1.25	10.1	0.34
1.35	0.06	4.2	2.04	7.3	0.78	10.2	1.17

1.36	2.83	4.3	0.48	8.1	7.70	10.3	0.67
1.37	2.27	4.4	3.99	8.2	2.37	10.4	0.44
1.38	2.31	4.5	2.45	8.3	1.09	10.5	0.48
1.39	0.35	4.6	6.23	8.4	0.43	10.6	0.42
1.40	1.03	4.7	3.47	8.5	0.16	10.7	0.23
1.41	1.23	4.8	0.24	8.6	1.77	11.1	6.01
1.42	1.05	4.9	0.72	8.7	0.05	11.2	10.60
1.43	1.29	4.10	0.37	8.8	0.43	11.3	11.75
1.44	0.25	4.11	3.62	8.9	1.15	11.4	2.39
1.45	0.13	4.12	1.06	8.10	0.37	11.5	3.20
1.46	0.17	4.13	0.25	8.11	2.84	11.6	8.28
1.47	0.40	4.14	6.11	8.12	1.00	12.1	4.12
1.48	1.02	4.15	0.69	8.13	0.45	12.2	0.70
1.49	1.13	4.16	0.47	8.14	2.25	13	0.10
1.50	0.89	4.17	0.29	8.15	1.74	14	2.04

In view of their ability to inhibit acetyl-CoA carboxylase(s), the compounds of general formula (I) according to the invention and the corresponding salts thereof are theoretically suitable for the treatment, including preventative treatment of all those 5 diseases or conditions which may be affected or which are mediated by the inhibition of acetyl-CoA carboxylase(s), in particular ACC2, activity.

Accordingly, the present invention relates to a compound of general formula (I) as a medicament.

10

Furthermore, the present invention relates to the use of a compound of general formula (I) for the treatment and/or prevention of diseases or conditions which are mediated by the inhibition of acetyl-CoA carboxylase(s), in particular ACC2, in a patient, preferably in a human.

15

In yet another aspect the present invention relates a method for treating, including preventing a disease or condition mediated by the inhibition of acetyl-CoA carboxylase(s) in a mammal that includes the step of administering to a patient,

preferably a human, in need of such treatment a therapeutically effective amount of a compound of the present invention, or a pharmaceutical composition thereof.

5 Diseases and conditions mediated by inhibitors of acetyl-CoA carboxylases embrace metabolic and/or cardiovascular and/or neurodegenerative diseases or conditions.

According to one aspect the compounds of the present invention are particularly suitable for treating diabetes mellitus, in particular Type 2 diabetes, Type 1 diabetes, and diabetes-related diseases, such as ishyperglycemia, metabolic syndrome, 10 impaired glucose tolerance, diabetic neuropathy, diabetic nephropathy, diabetic retinopathy, dyslipidemia, hypertension, hyperinsulinemia, and insulin resistance syndrome, hepatic insulin resistance, including complications such as macro- and microvascular disorders, including thromboses, hypercoagulable and prothrombotic states (arterial and venous), high blood pressure, coronary artery disease and heart 15 failure, increased abdominal girth, hypercoagulability, hyperuricemia, micro-albuminemia.

According to another aspect the compounds of the present invention are particularly suitable for treating overweight, obesity, including visceral (abdominal) obesity, 20 nonalcoholic fatty liver disease (NAFLD) and obesity related disorders, such as for example weight gain or weight maintenance

Obesity and overweight are generally defined by body mass index (BMI), which is correlated with total body fat and estimates the relative risk of disease. BMI 25 is calculated by weight in kilograms divided by height in meters squared (kg/m^2). Overweight is typically defined as a BMI of $25\text{-}29.9\text{ kg}/\text{m}^2$, and obesity is typically defined as a BMI of $30\text{ kg}/\text{m}^2$ or greater.

According to another aspect the compounds of the present invention are particularly 30 suitable for treating, including preventing, or delaying the progression or onset of diabetes or diabetes-related disorders including Type 1 (insulin-dependent diabetes mellitus, also referred to as "IDDM") and Type 2 (noninsulin-dependent diabetes mellitus, also referred to as "NIDDM") diabetes, impaired glucose tolerance, insulin resistance, hyperglycemia, pancreatic beta cell degeneration and diabetic

complications (such as macro- and microvascular disorders, atherosclerosis, coronary heart disease, stroke, peripheral vascular disease, nephropathy, hypertension, neuropathy, and retinopathy).

5 In addition the compounds of the present invention are suitable for treating dyslipidemias in general and more specifically elevated lipid concentrations in the blood and in tissues, dysregulation of LDL, HDL and VLDL, in particular high plasma triglyceride concentrations, high postprandial plasma triglyceride concentrations, low HDL cholesterol concentration, low apoA lipoprotein concentrations, high LDL

10 cholesterol concentrations, high apoB lipoprotein concentrations, including atherosclerosis, coronary heart disease, cerebrovascular disorders, diabetes mellitus, metabolic syndrome, obesity, insulin resistance and/or cardiovascular disorders.

15 ACC inhibition may lead to a centrally stimulating effect on food intake. Therefore compounds of the present invention may be suitable for treating eating disorders such as anorexia nervosa.

20 In addition the compounds of the present invention may provide neuroprotective effects in patients with Parkinson's disease, Alzheimer's disease, hypoxia, ischemia, amyotrophic lateral sclerosis or glioma and may improve cognitive scores in Alzheimer's diseases patients.

Further diseases and conditions mediated by inhibitors of acetyl-CoA carboxylases embrace but are not limited to:

25 A. disorders of fatty acid metabolism and glucose utilization disorders; disorders in which insulin resistance is involved;

B. hepatic disorders and conditions related thereto, including:

30 fatty liver, hepatic steatosis, non-alcoholic hepatitis, non-alcoholic steatohepatitis (NASH), alcoholic hepatitis, acute fatty liver, fatty liver of pregnancy, drug-induced hepatitis, iron storage diseases, hepatic fibrosis, hepatic cirrhosis, hepatoma, viral hepatitis;

C. skin disorders and conditions and those associated with polyunsaturated fatty acids, such as

- eczema, acne, sebaceous gland diseases, psoriasis, keloid scar formation or prevention, other diseases related to mucous membrane fatty acid composition;

5

D. primary hypertriglyceridemia or secondary hypertriglyceridemias following familial histiocytic reticulosis, lipoprotein lipase deficiency, hyperlipoproteinemias, apolipoprotein deficiency (e.g. apoCII or apoE deficiency);

10

E. diseases or conditions related to neoplastic cellular proliferation, for example benign or malignant tumors, cancer, neoplasias, metastases, carcinogenesis;

F. diseases or conditions related to neurological, psychiatric or immune disorders 15 or conditions;

G. other diseases or conditions in which inflammatory reactions, cell differentiation and/or other ACC-mediated aspects may for example be involved are:

- atherosclerosis such as, for example (but not restricted thereto), coronary sclerosis including angina pectoris or myocardial infarction, stroke, ischemic, stroke and transient ischemic attack (TIA),
- peripheral occlusive disease,
- vascular restenosis or reocclusion,
- chronic inflammatory bowel diseases such as, for example, Crohn's disease 20 and ulcerative colitis,
- pancreatitis,
- sinusitis,
- retinopathy, ischemic retinopathy,
- adipose cell tumors,

25

- lipomatous carcinomas such as, for example, liposarcomas,
- solid tumors and neoplasms such as, for example (but not restricted thereto), carcinomas of the gastrointestinal tract, of the liver, of the biliary tract and of the pancreas, endocrine tumors, carcinomas of the lungs, of the kidneys and the urinary tract, of the genital tract, prostate carcinomas, breast cancer (in

particular breast cancer with BRCA1 mutations), etc.,

- tumors in which ACC is up regulated,
- acute and chronic myeloproliferative disorders and lymphomas, angiogenesis
- neurodegenerative disorders including Alzheimer's disease, multiple sclerosis,

5 Parkinson's disease, epilepsy,

- erythema-squamous dermatoses such as, for example, psoriasis,
- acne vulgaris,
- other skin disorders and dermatological conditions which are modulated by PPAR,

10 - eczemas and neurodermatitis,

- dermatitis such as, for example, seborrheic dermatitis or photodermatitis,
- keratitis and keratoses such as, for example, seborrheic keratoses, senile keratoses, actinic keratoses, photo-induced keratoses or keratosis follicularis,
- keloids and keloid prophylaxis,

15 - bacterial infections,

- fungal infections,
- warts, including condylomata or condylomata acuminata
- human papilloma viral (HPV) infections such as, for example, venereal papillomata, viral warts such as, for example, molluscum contagiosum,

20 leukoplakia,

- papular dermatoses such as, for example, lichen planus,
- skin cancer such as, for example, basal-cell carcinomas, melanomas or cutaneous T-cell lymphomas,
- localized benign epidermal tumors such as, for example, keratoderma,

25 epidermal naevi,

- chilblains;
- high blood pressure,
- polycystic ovary syndrome (PCOS),
- asthma,

30 - cystic fibrosis,

- osteoarthritis,
- lupus erythematosus (LE) or inflammatory rheumatic disorders such as, for example rheumatoid arthritis,
- vasculitis,

- wasting (cachexia),
- gout,
- ischemia/reperfusion syndrome,
- acute respiratory distress syndrome (ARDS)

5 - viral diseases and infections

- lipodystrophy and lipodystrophic conditions, also for treating adverse drug effect;
- myopathies and lipid myopathies (such as carnitine palmitoyltransferase I or II deficiency);

10

H. formation of muscles and a lean body or muscle mass formation.

The dose range of the compounds of general formula (I) applicable per day is usually from 0.001 to 10 mg per kg body weight, for example from 0.01 to 8 mg per kg body weight of the patient. Each dosage unit may conveniently contain from 0.1 to 1000 mg, for example 0.5 to 500 mg.

The actual therapeutically effective amount or therapeutic dosage will of course depend on factors known by those skilled in the art such as age and weight of the patient, route of administration and severity of disease. In any case the combination will be administered at dosages and in a manner which allows a therapeutically effective amount to be delivered based upon patient's unique condition.

Pharmaceutical Compositions

25 Suitable preparations for administering the compounds of formula (I) will be apparent to those with ordinary skill in the art and include for example tablets, pills, capsules, suppositories, lozenges, troches, solutions, syrups, elixirs, sachets, injectables, inhalatives and powders etc. The content of the pharmaceutically active compound(s) is advantageously in the range from 0.1 to 90 wt.-%, for example from 1 to 70 wt.-%
30 of the composition as a whole.

Suitable tablets may be obtained, for example, by mixing one or more compounds according to formula (I) with known excipients, for example inert diluents, carriers,

disintegrants, adjuvants, surfactants, binders and/or lubricants. The tablets may also consist of several layers.

5 **Combination Therapy**

The compounds of the invention may further be combined with one or more, preferably one additional therapeutic agent. According to one embodiment the additional therapeutic agent is selected from the group of therapeutic agents useful in the treatment of diseases or conditions associated with metabolic diseases or 10 conditions such as for example diabetes mellitus, obesity, diabetic complications, hypertension, hyperlipidemia.

Therefore a compound of the invention may be combined with one or more additional therapeutic agents selected from the group consisting of anti-obesity agents 15 (including appetite suppressants), agents which lower blood glucose, anti-diabetic agents, agents for treating dyslipidemias, such as lipid lowering agents, anti-hypertensive agents, antiatherosclerotic agents, anti-inflammatory active ingredients, agents for the treatment of malignant tumors, antithrombotic agents, agents for the treatment of heart failure and agents for the treatment of complications caused by 20 diabetes or associated with diabetes.

Suitable anti-obesity agents include 11beta-hydroxy steroid dehydrogenase-1 (11beta-HSD type 1) inhibitors, stearoyl-CoA desaturase-1 (SCD-1) inhibitor, MCR-4 agonists, cholecystokinin-A (CCK-A) agonists, monoamine reuptake inhibitors, 25 sympathomimetic agents, beta3 adrenergic agonists, dopamine agonists, melanocyte-stimulating hormone analogs, 5HT2c agonists, melanin concentrating hormone antagonists, leptin (the OB protein), leptin analogs, leptin agonists, galanin antagonists, lipase inhibitors, anorectic agents, neuropeptide-Y antagonists (e.g., NPY Y5 antagonists), PY₃-36 (including analogs thereof), thyromimetic agents, 30 dehydroepiandrosterone or an analog thereof, glucocorticoid agonists or antagonists, orexin antagonists, glucagon-like peptide-1 agonists, ciliary neurotrophic factors, human agouti-related protein (AGRP) inhibitors, ghrelin antagonists, histamine 3 antagonists or inverse agonists, neuromedin U agonists, MTP/ApoB inhibitors (e.g., gut-selective MTP inhibitors), opioid antagonist, orexin antagonist, and the like.

Preferred anti-obesity agents for use in the combination aspects of the present invention include gut-selective MTP inhibitors CCK_A agonists, 5HT_{2c} agonists, MCR4 agonist, lipase inhibitor, opioid antagonists, oleoyl-estrone, obineptide, 5 pramlintide (Symlin®), tesofensine (NS2330), leptin, liraglutide, bromocriptine, orlistat, exenatide (Byetta®), AOD-9604 (CAS No. 221231-10-3) and sibutramine.

Suitable anti-diabetic agents include a sodium-glucose co-transporter (SGLT) inhibitor, a phosphodiesterase (PDE) 10 inhibitor, a diacylglycerol acyltransferase (DGAT) 1 or 2 inhibitor, a sulfonylurea (e.g., acetohexamide, chlorpropamide, diabinese, glibenclamide, glipizide, glyburide, glimepiride, gliclazide, glipentide, gliquidone, glisolamide, tolazamide, and tolbutamide), a meglitinide, an alpha-amylase inhibitor (e.g., tendamistat, trestatin and AL-3688), an alpha-glucosidase hydrolase inhibitor (e.g., acarbose), an alpha-glucosidase inhibitor (e.g., adiposine, 10 camiglibose, emiglitate, miglitol, voglibose, pradimicin-Q, and salbostatin), a PPAR_Y agonist (e.g., balaglitazone, ciglitazone, darglitazone, englitazone, isaglitazone, pioglitazone, rosiglitazone and troglitazone), a PPAR alpha/ gamma agonist (e.g., CLX-0940, GW-1536, GW-20 1929, GW-2433, KRP-297, L-796449, LR-90, MK-0767 and SB-21 9994), a biguanide (e.g., metformin), a GLP-1 derivative, a glucagon-like 15 peptide 1 (GLP-1) agonist (e.g., Byetta™, exendin-3 and exendin-4), a protein tyrosine phosphatase-1 B (PTP-1 B) inhibitor (e.g., trodusquemine, hyrtiosal extract), SIRT-1 inhibitor (e.g. reservatrol), a dipeptidyl peptidase IV (DPP-IV) inhibitor (e.g., sitagliptin, vildagliptin, alogliptin, linagliptin and saxagliptin), an insulin 20 secretagogue, a fatty acid oxidation inhibitor, an A2 antagonist, a c-jun amino-terminal kinase (JNK) inhibitor, an insulin, an insulin derivative, fast acting insulins, inhalable insulins, oral insulins, an insulin mimetic, a glycogen phosphorylase inhibitor, a VPAC2 receptor agonist and a glucokinase activator. Preferred anti-diabetic agents are metformin, a glucagon-like peptide 1 (GLP-1) agonist (e.g., ByettaTM) and DPP-IV inhibitors (e.g. sitagliptin, vildagliptin, alogliptin, linagliptin and 25 saxagliptin).

Preferably, compounds of the present invention and/or pharmaceutical compositions comprising a compound of the present invention optionally in combination with one or

more additional therapeutic agents are administered in conjunction with exercise and/or a diet.

Therefore, in another aspect, this invention relates to the use of a compound
5 according to the invention in combination with one or more additional therapeutic agents described hereinbefore and hereinafter for the treatment or prevention of diseases or conditions which may be affected or which are mediated by the inhibition of acetyl-CoA carboxylase(s), in particular ACC2, in particular diseases or conditions as described hereinbefore and hereinafter.

10

In yet another aspect the present invention relates a method for treating, including preventing a disease or condition mediated by the inhibition of acetyl-CoA carboxylase(s) in a patient that includes the step of administering to the patient, preferably a human, in need of such treatment a therapeutically effective amount of a
15 compound of the present invention in combination with a therapeutically effective amount of one or more additional therapeutic agents described in hereinbefore and hereinafter,

The use of the compound according to the invention in combination with the
20 additional therapeutic agent may take place simultaneously or at staggered times.

The compound according to the invention and the one or more additional therapeutic agents may both be present together in one formulation, for example a tablet or capsule, or separately in two identical or different formulations, for example as a so-
25 called kit-of-parts.

Consequently, in another aspect, this invention relates to a pharmaceutical composition which comprises a compound according to the invention and one or more additional therapeutic agents described hereinbefore and hereinafter, optionally
30 together with one or more inert carriers and/or diluents.

Further aspects of the invention include the use of a compound according to the invention or a salt thereof as a crop protection agent to combat and/or prevent fungal infestations, or to control other pests such as weeds, insects, or acarids that are

harmful to crops. Another aspect of the invention relates to the use of a compound according to the invention or a salt thereof for controlling and/or preventing plant pathogenic microorganisms, for example plant pathogenic fungi. Therefore one aspect of the invention is a compound according to the formula (I) or a salt thereof for 5 use as a fungicide, insecticide, acaricide and/or herbicide. Another aspect of the invention relates to an agricultural composition comprising a compound of the present invention together with one or more suitable carriers. Another aspect of the invention relates to an agricultural composition comprising a compound of the present invention in combination with at least one additional fungicide and/or 10 systemically acquired resistance inducer together with one or more suitable carriers.

Examples

The Examples that follow are intended to illustrate the present invention without 15 restricting it. The terms "ambient temperature" and "room temperature" are used interchangeably and designate a temperature of about 20 °C.

Preliminary remarks:

As a rule, $^1\text{H-NMR}$ and/or mass spectra have been obtained for the compounds 20 prepared. The R_f values are determined using Merck silica gel 60 F_{254} plates and UV light at 254 nm.

Abbreviations:

ACN	acetonitrile
AcOH	acetic acid
aq.	aqueous
BOC	tert-butoxy-carbonyl-
CDI	N,N-carbonyldiimidazole
CDT	N,N-carbonyldi triazole
CyH	cyclohexane
DBAD	di-tert-butyl azodicarboxylate
DCM	dichloromethane
DIAD	diisopropyl azodicarboxylate
DIBAIH	diisobutyl aluminium hydride


DIPE	diisopropyl ether
DIPEA	N,N-diisopropylethylamine
DMAP	4-dimethylaminopyridine
DMF	N,N-dimethylformamide
DMSO	dimethyl sulfoxide
EI-MS	electron induced mass spectrometry
ESI-MS	electrospray ionisation mass spectrometry
EtOAc	ethyl acetate
EtOH	ethanol
Ex	example
FA	formic acid
GC	gas chromatography
HATU	2-(1 H-7-azabenzotriazol-1-yl)-1 ,1 ,3,3-tetramethyluronium hexafluorophosphate
HOEt	N-hydroxybenzotriazole
HPLC	high performance liquid chromatography
LAH	lithium aluminium hydride
MeOH	methanol
m.p.	melting point
MS	mass spectrum
MsCl	methanesulfonyl chloride
n.d.	not determined
NMP	N-methyl-2-pyrrolidone
Pd/C	palladium on activated carbon
PE	petroleum ether
PG	protecting group
R _f	retention factor
r.t.	room temperature (about 20°C)
R _t	retention time
sat.	saturated
TBME	i _{ie} /f-butyl methyl ether
TEA	triethylamine
TF / TFA	trifluoroacetic acid
THF	tetrahydrofuran
TBAF	tetrabutylammonium fluoride

TBTU	2-(1 H-benzotriazol-1-yl)-1,1,3,3-tetramethyluronium tetrafluoroborat
TLC	thin layer chromatography
TMS	trimethylsilyl
Ts	4-toluenesulfonyl
X-Phos	2-dicyclohexylphosphino-2',4',6'-triisopropylbiphenyl

Preparation of Starting Compounds

5 Example I

1-(4-Hydroxyphenyl)propane-2-amine




10 A mixture of 7.00 g (46.6 mmol) 4-hydroxyphenylacetone and 200 mL **7N** ammonia in MeOH is charged with 0.70 g Raney nickel. The mixture is stirred in an hydrogen atmosphere (50 psi) at r.t. over night. After complete reaction, the mixture is filtrated, the solvent is removed in vacuo and the crude product is purified by HPLC (MeOH/H₂O/NH₃).

15 C₉H₁₃NO (M= 151.2 g/mol)
ESI-MS: 152 [M+H]⁺
R_t (HPLC): 0.95 min (method A)

Example II

20 (S)-4-(1-Amino-ethyl)-phenol hydrobromid




To 11.0 g (72.7 mmol) (S)-1-(4-methoxy-phenyl)-ethylamine is added carefully 30 mL HBr (30% in HOAc). The mixture is stirred at 100 °C for 4 h. After cooling down to r.t., the solvent is removed in vacuo and the residue is dried in vacuo. The resulting product is used without further purification.

5 $C_6H_{11}NO \cdot HBr$ (M= 218.1 g/mol)
 ESI-MS: 121 [M+H-NH₃]⁺
 Rt (HPLC): 0.50 min (method A)

Example III

10 Example III.1 (general route)
(S)-N-(1 -(4-Bromophenyl)ethyl)acetamide



15 To 200 g (1.00 mol) (S)-1 -(4-bromophenyl)ethylamine in 800 mL DCM are slowly added 94.5 mL (1.00 mol) acetic anhydride while cooling the mixture to 20-30 °C. Then the cooling is removed and the reaction mixture is stirred at r.t. over night. Afterwards the mixture is consecutively washed with water, sat. aq. NaHCO₃ solution, water, diluted aq. citric acid solution and again water. The org. layer is dried with
 20 MgSO₄ and the solvent is removed in vacuo. The crude product is used without further purification.

$C_9H_{11}BrNO$ (M= 242.1 g/mol)
 ESI-MS: 242/244 [M+H]⁺
 Rt (HPLC): 1.67 min (method A)


25

The following compounds are prepared analogously to example 11.1. For the examples 11.2-3 acetic acid is used as solvent and after complete reaction the solvent is removed in vacuo and the residue is partitioned between EtOAc and water.

Ex.	Starting material	Product structure	Mass spec result	HPLC retention time (method)
III.1			242/244 [M+H] ⁺	1.67 (A)
III.2			242 [M+H] ⁺	2.47 (B)
III.3			194 [M+H] ⁺	1.80 (B)

Example IV

5 N-(1 -(4-iodophenyl)ethyl)acetamide



To 8.70 g (35.9 mmol) N-(1 -(4-bromophenyl)ethyl)acetamide (III.2) in 100 mL 1,4-dioxane are added 0.69 g (3.62 mmol) Cul, 0.97 mL (9.11 mmol) N,N'-dimethyl-10 ethylenediamine and 10.8 g (71.9 mmol) NaI. The reaction mixture is stirred at 120°C for 3 d. The mixture is allowed to cool to r.t. and half of the solvent is removed in vacuo. EtOAc and diluted aq. ammonia solution are added and the layers are separated. The aq. layer is once more extracted with EtOAc. The organic layers are combined, dried with Na₂SO₄ and the solvent is removed in vacuo. The crude product is triturated with diethylether and dried in vacuo.


C₁₀H₁₂INO (M= 289.1 g/mol)

ESI-MS: 290 [M+H]⁺

Rt (HPLC): 2.95 min (method B)

Example V(S)-N-(1-(4-Hydroxyphenyl)ethyl)acetamide


5



19.5 g (71.5 mmol) of example II are added to 50 mL THF and charged with 40 mL (287 mmol) TEA. 5.59 mL (78.7 mmol) acetyl chloride are added carefully and the mixture is stirred at r.t. for 2 h. 100 mL water are added and the reaction mixture is slightly acidified with citric acid. The aqueous phase is extracted four times with EtOAc. The org. phases are combined, dried with MgSO_4 and the solvent is removed in vacuo. The crude product is dissolved in 40 mL MeOH and charged with 30 mL aq. 4 N NaOH-solution and stirred at r.t for 1.5 h. Then 34 mL aq. 4 N HCl solution are added and the aqueous phase is extracted three times with ethyl acetate. The org. phases are combined, dried with MgSO_4 and the solvent is removed in vacuo. The crude product is purified by HPLC (MeOH/H₂O/NH₃).

C₁₀H₁₃NO₂ (M= 179.2 g/mol)
 ESI-MS: 180 [M+H]⁺
 Rt (HPLC): 0.52 min (method A)

20

Example VIN-(1-(4-Hydroxyphenyl)ethyl)acetamide

a) 1.00 g (3.46 mmol) of example IV.3 and 1.02 g (10.4 mmol) KOAc are added to 10 mL DMSO. After degassing the mixture 1.32 g (5.19 mmol) bis(pinakolato)diboron and 141 mg (0.17 mmol) $\text{PdCl}_2(\text{dppf})^*\text{CH}_2\text{Cl}_2$ are added and the mixture is stirred at 80 °C for 1.5 h. 80 mL EtOAc are added and the resulting mixture is washed with water (3x). The org. phase is dried with MgSO_4 and the solvent is removed in vacuo. The crude product is purified by column chromatography (silica gel, DCM/MeOH 9/1).

- 75 -

$\text{C}_6\text{H}_{24}\text{BNO}_3$ (M= 289.2 g/mol)

ESI-MS: 290 [M+H]⁺

R_t (HPLC): 1.81 min (method A)

5 b) 0.90 g (3.11 mmol) of the above mentioned product is added to 20 mL THF and cooled down to 0 °C. 0.78 mL (3.11 mmol) **4N** aq. NaOH solution and 0.41 mL (4.67 mmol) **H2O2** (35% in water) are added and the resulting mixture is stirred for 3 h at constant temperature. The mixture is separated between water and EtOAc and the aq. phase is extracted with EtOAc (3x). The org. phases are combined and the solvent is removed in vacuo. The crude product is purified by HPLC (MeOH/H₂O/NH₃). The resulting product is triturated with diethylether.

10 C10H13NO2 (M= 179.2 g/mol)


ESI-MS: 180 [M+H]⁺

R_t (HPLC): 1.16 min (method E)

15

Example VII

(S)-3-(1-(4-hydroxyphenyl)ethyl)-1,1-dimethylurea



20 7.20 g (33.0 mmol) of example II are added to 80 mL DCM and charged with 18.5 mL (132 mmol) TEA. After cooling down to 0 °C 5.88 g (36.3 mmol) CDI are added, the cooling is removed and stirring is continued for 20 min. 4.46 g (99.0 mmol) dimethylamine are added and the mixture is stirred at r.t. over night. 1 N aq. citic acid is added, the phases are separated and the aq. phase is extracted one more time with DCM.


25 The org. phases are combined, dried with MgSO₄ and the solvent is removed in vacuo. The crude product is purified by column chromatography (silica gel, DCM/MeOH 95/5 → 90/10).

C₁₁H₁₆N₂O₂ (M= 208.3 g/mol)

ESI-MS: 209 [M+H]⁺

30 R_t (HPLC): 0.94 min (method A)

Example VIII

(S)-N-(1-(4-hydroxyphenyl)ethyl)cyclopropanecarboxannide


7.20 g (33.0 mmol) of example II are added to 80 mL DCM and charged with 18.5 mL
 5 (132 mmol) TEA. After cooling down to 0 °C 3.59 mL (39.6 mmol) cyclopropane-
 carbonyl chloride are carefully added. Afterwards the cooling is removed and the
 reaction mixture is stirred at r.t. over night. The reaction mixture is further diluted with
 DCM and washed with aq. citric acid. The aq. phase is extracted one more time with
 DCM. The org. phases are combined, dried with MgSO_4 and the solvent is removed
 10 in vacuo. The crude product is purified by column chromatography (silica gel,
 DCM/MeOH 95/5 → 90/1 0).

C₁₂H₁₅NO₂ (M= 205.3 g/mol)

ESI-MS: 206 [M+H]⁺

Rt (HPLC): 0.81 min (method A)

15


Example IX(S)-tert-Butyl 1-(4-hydroxyphenyl)ethylcarbamate

20 5.00 g (22.9 mmol) of example II are added to 80 mL DCM and charged with 34.4 mL
 (68.8 mmol) of a 2N aq. Na_2CO_3 solution. 5.25 g (24.1 mmol) Boc_2O in 20 mL DCM
 are added and the mixture is stirred at r.t. for 2 h. 100 mL water are added and the
 resulting mixture is extracted three times with DCM. The org. phases are combined,
 washed with water, dried with MgSO_4 and the solvent is removed in vacuo. The
 25 crude product is purified by column chromatography (silica gel, DCM/MeOH 95/5 →
 80/20).

C₁₃H₁₉NO₃ (M= 237.3 g/mol)

ESI-MS: 238 [M+H]⁺

Rt (HPLC): 1.58 min (method A)

Example X(S)-Benzyl 1-(4-hydroxyphenyl)ethylcarbannate

5


7.20 g (33.0 mmol) of example II are added to a mixture of 10 mL THF and 10 mL water. 19.4 g (231 mmol) NaHCO₃ are slowly added so that the pH > 7. 5.18 mL (36.3 mmol) benzyl chloroformate is added slowly and the resulting mixture is stirred at r.t. for 3 h. 100 mL water are added and the mixture is slightly acidified with 10% aq.

10 citric acid. The resulting mixture is extracted two times with EtOAc. The org. phases are combined, dried with MgSO₄ and the solvent is removed in vacuo. The crude product is purified by column chromatography (silica gel, DCM/MeOH 95/5 → 80/20).
 C₁₃H₁₉NO₃ (M= 271.3 g/mol)
 ESI-MS: 272 [M+H]⁺

15 R_t (HPLC): 1.69 min (method A)

Example XIExample XI.1 (general route)(S)-tert-Butyl 3-(4-(1-acetamidoethyl)phenoxy)azetidine-1-carboxylate

20



25 6.00 g (33.5 mmol) of example I are added to 120 mL DMF and charged with 8.41 g (33.5 mmol) 1-Boc-3-methanesulfonyloxyazetidine and 21.8 g (67.0 mmol) CS₂CO₃. The reaction mixture is stirred at 80 °C over night. The reaction is quenched by the addition of water. The resulting mixture is extracted two times with EtOAc. The org. phases are combined, washed with diluted aq. NaHCO₃ solution (3x), dried with

MgSO_4 and the solvent is removed in vacuo. The crude product is purified by HPLC (MeOH/H₂O/NH₃).

$\text{C}_{18}\text{H}_{26}\text{N}_2\text{O}_4$ (M= 334.4 g/mol)


ESI-MS: 335 [M+H]⁺

5 Rt (HPLC): 1.88 min (method A)

The following compounds are prepared analogously to example XI.1.


For the example XI.6 1-Boc-3-iodoazetidine is used.

Ex.	Starting material	Product structure	Mass spec result	HPLC retention time (method)
XI.1	V		335 [M+H] ⁺	1.88 (A)
XI.2	VI		335 [M+H] ⁺	1.87 (A)
XI.3	III.4		349 [M+H] ⁺	1.93 (A)
XI.4	VIII		361 [M+H] ⁺	2.19 (A)
XI.5	VII		364 [M+H] ⁺	2.12 (A)
XI.6	X		444 [M+NH4] ⁺	2.25 (A)

Example XII(S)-tert-Butyl 1-(4-(1-benzhydrylazetidin-3-yloxy)phenyl)ethylcarbamate

5 7.48 g (31.5 mmol) of example I are added to 150 mL DMF and charged with 10.0 g (33.5 mmol) 1-diphenylmethyl-3-methanesulfonyloxyazetidine and 20.5 g (63.0 mmol) Cs_2CO_3 . The reaction mixture is stirred at 80 °C for 3 h, filtered and the solvent is removed in vacuo. The crude product is purified by column chromatography (PE/EtOAc 80/20 \rightarrow 50/50).

10 $\text{C}_{29}\text{H}_{34}\text{N}_2\text{O}_3$ (M= 458.6 g/mol)
ESI-MS: 459 $[\text{M}+\text{H}]^+$
Rt (HPLC): 2.32 min (method A)

Example XIIIExample XIII.1 (general route)(S)-N-(1-(4-(Azetidin-3-yloxy)phenyl)ethyl)acetamide

20 To 1.20 g (3.59 mmol) of example XI.1 in 10 mL MeOH are added 8.61 mL (10.8 mmol) HCl (c = 1.3 mol/l in MeOH). The reaction mixture is stirred at r.t. over night. The solvent is removed in vacuo and the residue is purified by HPLC (MeOH/H₂O/NH₃).
 $\text{C}_{13}\text{H}_{18}\text{N}_2\text{O}_2$ (M= 234.3 g/mol)

25 ESI-MS: 235 $[\text{M}+\text{H}]^+$
Rt (HPLC): 1.06 min (method A)


The following compounds are prepared analogously to example XI1.1.

For the examples XIII.3 and XIV.4 a gentle warming (-35-40 °C) is applied.

For the example XIII.5 aq. NaOH (1N) is added to the residue and the resulting precipitate is filtered, washed and dried.


5 For the example XIII.6 the residue is not purified by HPLC.

Ex.	Starting material	Product structure	Mass spec result	HPLC retention time (method)
XIII.1	XI.1		235 [M+H] ⁺	1.06 (A)
XIII.2	XI.2		235 [M+H] ⁺	1.06 (A)
XIII.3	XI.4		261 [M+H] ⁺	1.23 (A)
XIII.4	XI.5		264 [M+H] ⁺	1.17 (A)
XIII.5	XI.6		327 [M+H] ⁺	1.72 (A)
XIII.6	XI.1		235 [M+H] ⁺	1.06 (A)

Example XIVN-(1 -(4-(Azetidin-3-yloxy)phenyl)propan-2-yl)acetamide


5 To 0.90 g (2.58 mmol) of example XI.3 in 15 mL DCM is added 1 mL TFA. The reaction mixture is stirred at r.t. for 3 h and afterwards neutralised with sat. aq. NaHCO₃ solution. The aq. Phase is saturated with K₂CO₃ and extracted with EtOAc and MeOH (3x). The org. phases are combined, dried with MgSO₄ and the solvent is removed in vacuo.

10 Ci₄H₂₀N₂O₂ (M= 248.3 g/mol)
 ESI-MS: 249 [M+H]⁺
 Rt (HPLC): 1.23 min (method A)

Example XV(S)-tert-Butyl 1-(4-(azetidin-3-yloxy)phenyl)ethylcarbamate

7.20 g (15.7 mmol) of example XII are added to 50 mL EtOH and charged with 1.00 g 20 Pd/C (10%). The reaction mixture is hydrogenated at r.t. for 24 h at a hydrogen pressure of 3 bar. The reaction mixture is filtered and the solvent is removed in vacuo. The crude product is triturated with TBME.

Ci₆H₂₄N₂O₃ (M= 292.4 g/mol)
 ESI-MS: 293 [M+H]⁺
 25 R_t (HPLC): 1.72 min (method A)

Example XVIExample XVI.1 (general route)1-Bromo-4-cyclopropylmethoxybenzene

5.0 g (28.9 mmol) 4-bromophenol, 3.93 g (43.4 mmol) (chloromethyl)cyclopropane and 7.99 g (57.8 mmol) K_2CO_3 are added to 10 mL DMF and stirred at 80°C over night. Afterwards the reaction mixture is diluted with water and extracted with DCM. The organic layer is dried with $MgSO_4$ and the solvent is removed in vacuo.

10 C₁₀H₁₁BrO (M= 227.1 g/mol)

EI-MS: 226/228 [M]⁺

Rt (HPLC): 8.16 min (method J)

The following compounds are prepared analogously to example XVI.1

























15 For example XVI.2 the reaction temperature is 120 °C.


For example XVI.1 7 the reaction temperature is 140 °C and the product is purified by column chromatography (silica gel, PE/EtOAc 8/2).

For example XVI.1 9 KOtBu is used as base and the reaction mixture is stirred at 100 °C for 6 h.

Ex.	Starting material	Starting material	Product structure	Mass spec result	HPLC retention time (method)
XVI.1				226/228 [M] ⁺	8.16 (J)
XVI.2				226/228 [M] ⁺	4.90 (Q)

XVI.3				n.d.	9.01 (J)
XVI.4				n.d.	8.53 (J)
XVI.5				n.d.	4.81 (R)
XVI.6				244/246 [M] ⁺	2.08 (A)
XVI.7				219 [M-C ₄ H ₇] ⁻	2.17 (A)
XVI.8				231/233 [M+H] ⁺	2.00 (A)
XVI.9				257/259 [M+H] ⁺	2.18 (A)
XVI.10				259/261 [M+H] ⁺	2.21 (A)
XVI.11				257/259 [M+H] ⁺	2.11 (A)
XVI.12				244/246 [M] ⁺	2.12 (A)
XVI.13				245/247 [M+H] ⁺	1.30 (N)

XVI.14				257/259 [M+NH ₄] ⁺	2.27 (A)
XVI.15				252/254 [M+H] ⁺	2.00 (A)
XVI.16				251/253 [M] ⁺	2.07 (A)
XVI.17				n.d.	2.06 (A)
XVI.18				n.d.	TLC: $R_f = 0.9$ PE/EtOAc 8/2
XVI.19				287/289 [M+H] ⁺	1.57(P)
XVI.20				n.d.	1.31 (N)
XVI.21				n.d.	1.27 (N)

Example XVIIExample XVII.1 (general route)1-Bromo-4-(2-bromoethoxy)benzene

5

a) 55.0 g (318 mmol) 4-bromophenol and 14.1 g (352 mmol) NaOH are added to 110 mL water. 41.1 mL (477 mmol) dibromoethane are added slowly and the reaction mixture is stirred for 16 h under reflux. Afterwards the reaction mixture is extracted with DCM and the solvent is removed in vacuo. The crude product is purified by

10 column chromatography (silica gel, CyH/EtOAc 4/1).

b) 52.0 g (186 mmol) of 1-bromo-4-(2-bromoethoxy)benzene is added to 300 mL THF and cooled down to 0 °C. Within 30 min 25.0 g (223 mmol) KOtBu are added to this mixture in several portions. Cooling is removed and the reaction mixture is stirred at r.t. over night. The reaction is quenched by the addition of water. The resulting mixture is extracted with EtOAc (2x). The org. phases are combined, washed with sat. aq. NaCl solution, dried with MgSO₄ and the solvent is removed in vacuo. The resulting product is used without further purification.

c) 39.0 g (176 mmol) of 1-bromo-4-vinyloxybenzene and 32.4 mL (441 mmol) chloroiodomethane are added to 500 mL dichloroethane and cooled down to 0 °C. 20 During 1 h 200 mL (200 mmol) diethylzinc solution (c = 1 mol/l in hexane) are added and stirring is continued for 2 h at 0 °C. The reaction is quenched by the addition of 200 mL of a sat. aq. NH₄Cl solution and extracted with TBME (2x). The org. phases are combined, washed with sat. aq. NaCl solution, dried with MgSO₄ and the solvent is removed in vacuo. The crude product is purified by column chromatography (silica 25 gel, PE).

C₉H₉BrO (M= 213.1 g/mol)

EI-MS: 212/214 [M]⁺

R_f (TLC): 0.4 (silica gel, PE)

The following compounds are prepared analogously to example XVI_{1.1}.


For the examples XVI_{1.2} - XVI_{1.4} the phenolate in step a) is preformed by reacting the appropriate phenol with NaOH in a MeOH/water (1/1) mixture at r.t. for 1 h. Then the solvent is removed in vacuo and the resulting sodium salt is reacted with

5 dibromoethane (5 eq.) at 100 °C for 24 h. The reaction mixture is quenched by the addition of water and extracted with DCM.

For example XVII.5 step a: To the diphenol and dibromoethane (8 eq.) in acetone is added CS₂CO₃ (5 eq.) and the reaction mixture is stirred at 90 °C for 45 h. The reaction mixture is quenched by the addition of water and extracted with EtOAc.

10 For example XVII.6 the phenol is added to THF, deprotonated with NaH at 0 °C and after the addition of dibromoethane the reaction mixture is refluxed for 16 h followed by an aq. work up.


Ex.	Starting material	Product structure	Mass spec result	HPLC retention time (method)
XVII. 1	^{HO} XX.		212/214 [M] ⁺	TLC: R_f = 0.4 (silica gel, PE)
XVI _{1.2}			n.d.	5.89 (H)
XVI _{1.3}			n.d.	5.30 (I)
XVI _{1.4}			n.d.	7.34 (I)
XVI _{1.5}			n.d.	8.50 (J)
XVI _{1.6}			230/232 [M] ⁺	TLC: R_f = 0.54 (silica gel, hexane/EtOAc 9/1)

Example XVIII5-Bromo-2-cyclobutoxy-pyrimidine

5 2.42 ml (31.0 mmol) cyclobutanol and 3.00 g (15.5 mmol) 5-bromo-2-chloropyrimidine are added to 40 ml 1,4-dioxane and cooled down to 0 °C. Then the reaction mixture is charged with 1.86 g (46.5 mmol) NaH. After removing of the cooling bath the reaction mixture is stirred at r.t. for 1 h. The reaction is quenched by the addition of water and sat. aq. NaHCO₃ solution. The 1,4-dioxane is removed in

10 vacuo and the aq. residue is extracted with DCM. The org. phases are combined, washed with water and dried with MgSO₄. The solvent is removed under reduced pressure. The crude product is used without further purification.


15 C₉H₁₀BrN₂O (M= 229.1 g/mol)
ESI-MS: 229/231 [M+H]⁺
R_t (HPLC): 1.93 (method A)

Example XIX5-Bromo-N-cyclobutylpyrimidin-2-amine

20 In a sealed tube 2.00 g (10.3 mmol) 5-bromo-2-chloropyrimidine, 1.15 ml (13.4 mmol) cyclobutylamin and 2.70 ml (15.5 mmol) DIPEA are added to 12 mL ACN. The reaction mixture is stirred at 50 °C over night, then diluted with EtOAc and washed with water (2x). The org. phase is dried with MgSO₄ and the solvent is

25 removed in vacuo.

C₈H₁₀BrN₃ (M= 228.1 g/mol)
ESI-MS: 228/230 [M+H]⁺
R_t (HPLC): 1.87 (method A)

Example XX5-Bromo-N-sec-butylpyrimidin-2-amine

In a sealed tube 2.00 g (10.3 mmol) 5-bromo-2-chloropyrimidine and 3.13 ml (31.0 mmol) sec-butylamin are added to 12 mL ACN. The reaction mixture is stirred at 120 °C for 2 h, then diluted with EtOAc and washed with water (2x). The org. phase is dried with MgSO₄ and the solvent is removed in vacuo.


C₈H₁₂BrN₃ (M= 230.1 g/mol)

ESI-MS: 230/232 [M+H]⁺

10 R_t (HPLC): 1.93 (method A)

Example XXIExample XXI.1 (general route)2-Chloro-5-cyclopropylpyrimidine

15




3.60 g (18.6 mmol) 5-bromo-2-chloropyrimidine, 4.80 g (55.8 mmol) cyclopropyl boronic acid, 13.8 g (65.1 mmol) K₃PO₄, 568 mg (2.03 mmol) tneydohexylphosphine and 4 mL water are added to 80 mL toluene. The mixture is degassed thoroughly and charged with 627 mg (2.79 mmol) Pd(OAc)₂. After degassing again the mixture is stirred at 100 °C over night, filtered and the solvent is removed in vacuo. The crude product is purified by column chromatography (silica gel, PE/EtOAc 90/1 0 → 70/30).

C₇H₇ClN₂ (M= 154.6 g/mol)

EI-MS: 155 [M+H]⁺

25 R_t (HPLC): 1.34 (method A)

Example XXIIExample XXII.1 (general route)5-Chloro-2-phenyloxazolo[5,4-d]pyrimidine

5

7.82 g (64.0 mmol) benzoic acid and 20 mL POCl_3 are stirred together at 100 °C for 30 min. Within 45 min 7.00 g (42.7 mmol) 2,6-dichloro-5-aminopyrimidine are added by several portions. The reaction mixture is stirred for additional 2 h at the same temperature. After that the mixture is carefully added to an ice cold aq. NaOH solution.

10 The resulting precipitate is filtered, washed with water and dried.

$\text{C}_{12}\text{H}_6\text{ClN}_3\text{O}$ ($M= 231.6$ g/mol)

EI-MS: 232 $[\text{M}+\text{H}]^+$


Rt (HPLC): 2.22 (method S)

15 The following compounds are prepared analogously to example XXII.1.

For the example XXII.2 the solvent of reaction mixture is removed in vacuo and the residue is triturated first with diethylether, then dissolved in DCM, filtered and the solvent is removed in vacuo again. The residue is one more time triturated with diethylether.

Ex.	Starting material	Product structure	Mass spec result	HPLC retention time (method)
XXV.1			232 $[\text{M}+\text{H}]^+$	2.22 (S)
XXV.2			238 $[\text{M}+\text{H}]^+$	1.43 (S)

20

Example XXIII5-(3-Bromophenoxy)-2-nnethylpentan-2-ol

5 50.2 mL (150 mmol) MeMgBr (3 M in diethylether) are cooled down to 0 °C. 24.0 g (75 mmol) of the intermediate XIX.20 are added slowly and after 1h additional 50 mL THF are added and stirring is continued over night. The reaction mixture is poured onto a ice /aq. HCl mixture and extracted with EtOAc. The org. phases are combined, washed with a sat. aq. NaHCO₃ solution and dried with Na₂SO₄. The solvent is

10 removed in vacuo and the residue is purified by column chromatography (silica gel, CyH/EtOAc 80/20).

C₁₂H₁₇BrO₂ (M= 273.2 g/mol)

ESI-MS: 273/275 [M+H]⁺

Rt (HPLC): 1.51 (method P)

15


Example XXIV8-Bromo-5,5-dimethyl-2,3,4,5-tetrahydrobenzo[b]oxepine

20 1.09 g (4.00 mmol) of intermediate XXIII are added to 1 l DCM and cooled down to 0 °C. 4.26 mL cone. H₂SO₄ are dropped into the mixture and stirring is continued for 1 h. 300 mL water are added and the phases are separated. The org. phase is washed with sat. aq. NaCl solution, dried with Na₂SO₄ and the solvent is removed in vacuo.

25 C₁₂H₁₅BrO (M= 255.2 g/mol)

EI-MS: 254 [M]⁺


Rt (HPLC): 1.75 (method P)

Example XXV1-(6-Chloropyridin-3-yl)cyclopropanecarbonitrile

To 32 ml of an aq. NaOH solution (50%) and 32 ml ACN 50.2 ml (150 mmol) are added 6.60 g (21.2 mmol) $BnBu_3NBr$ and 3.20 g (21.0 mmol) of 6-chloropyridin-3-yl-acetonitrile. Then 2.17 ml (25.2 mmol) 1,2-dibromoethane are added and the mixture is stirred at r.t. for 20 h. Afterwards the mixture is extracted with EtOAc (2x), the org. phases are combined, washed with water and sat. aq. NaCl solution, dried with $MgSO_4$ and the solvent is removed in vacuo. The residue is purified by column chromatography (silica gel, CyH/EtOAc 95/5 \rightarrow 50/50).

C₉H₇CIN₂ (M = 178.6 g/mol)

ESI-MS: 179 [M+H]⁺

Example XXVI3-(4-Chlorophenyl)-6-iodopyridazine

20 ml ACN are added to 0.50 g (1.11 mmol) 3-(4-chlorophenyl)-6-chloropyridazine and 3.33 g (22.2 mmol) NaI before 189 μ l fuming cone. HCl are added and the mixture is stirred at 80 °C for 4 h. The mixture is alkalised with aq. ammonia solution (32%) diluted with water and extracted with EtOAc. The org. phases are combined, dried with Na_2SO_4 and the solvent is removed in vacuo. The resulting crude product is triturated with TBME.


C₁₀H₆ClIIN₂ (M= 316.5 g/mol)

ESI-MS: 317 [M+H]⁺

R_t (HPLC): 3.15 (method B)

Example XXVIIExample XXVII.1 (general route)2-Bromo-5-ethoxy-pyridine

5



2.51 g (22.99 mmol) 1-bromoethane and 1.99 g (14.37 mmol) K_2CO_3 are added to a mixture of 1.0 g (5.75 mmol) 2-bromo-5-hydroxypyridine in 100 mL ACN. The mixture is stirred at reflux over night. Then the reaction mixture is poured onto water and extracted with TBME. The organic layer is dried with Na_2SO_4 and the solvent is removed under vacuo. The residue is purified by column chromatography (silica gel, PE/EtOAc 1/1).

C_7H_8BrNO (M = 202.1 g/mol)


ESI-MS: 202 [M+H]⁺

Rt (HPLC): 1.70 min (method F)

15

The following compounds are prepared analogously to example XXVII.1

Ex.	Starting material	Product structure	Mass spec result	HPLC retention time (method)
XXVII.1			202 [M+H] ⁺	1.70 (F)
XXVII.2			216 [M+H] ⁺	1.70 (F)

Example XXVIIIExample XXVIII.1 (general route)5-(4-Iodo-pyridin-2-yloxy)-pyrimidine

To 259 mg (26.9 mmol) pyrimidin-5-ol (J. Chem. Soc. 1956, 2033) in 200 mL DMF are added 108 mg (26.9 mmol) sodium hydride (60% dispersion in mineral oil). The mixture is stirred for 20 min at r.t.. After that time, 500 mg (22.4 mmol) 2-fluoro-4-iodopyridine are added and the mixture is stirred for 12 h at 80°C. Subsequently the 10 mixture is poured into water and extracted with ethyl acetate (3x). The combined organic layers are washed with brine. After drying over sodium sulphate, the solvent is removed in vacuo and the residue is purified by column chromatography (silica gel; heptane/EtOAc, 100/0 → 40/60).

15 $C_9H_6IN_3O$ (M = 299.1 g/mol)

ESI-MS: 300 $[M+H]^+$

Rt (HPLC) : 2.87 min (method C)

The following compounds are prepared analogously to Example XXVIII.1
For the example XXVI 1.2 the reaction mixture is stirred at 50 °C for 4 h.
20 For the example XXVIII.3 THF is used as solvent and the reaction mixture is stirred at r.t. over night. For the example XXVI 1.4 6 eq. of the alcohol and 4 eq. of KOtBu as base are used and the reaction mixture is stirred at r.t. for 3 h.


Ex.	Starting material	Starting material	Product structure	Mass spec result	HPLC retention time (method)
XXVIII.1				300 $[M+H]^+$	2.87 (C)
XXVIII.2				249/251 $[M+H]^+$	1.12 (K)

XXVIII.3				228/230 [M+H] ⁺	1.14 (K)
XXVIII.4				296 [M] ⁺	GC: 4.56 (AA)
XXVIII.5				276 [M] ⁺	GC: 4.31 (AA)

Example XXIX and XXX

4-iodo-2-propoxy-pyridine (XXIX) and 2-chloro-4-propoxypyridine (XXX)

5



0.58 g (25.06 mmol) sodium are carefully added to 40 ml n-propanol by several portions. The mixture is stirred until the metal is dissolved completely (ca. 45 min). Then 6.00 g (25.1 mmol) 2-chloro-4-iodo-pyridine are slowly added to the mixture.


10 The mixture is stirred at reflux for 3 h. The reaction is quenched by the addition of water. The solvent is removed in vacuo and to the residue are added 20 ml DMF/MeOH. The mixture is filtrated and the filtrate is purified by HPLC (MeOH/H₂O/NH₃).

4-iodo-2-propoxy-pyridine (XXIX):

15 C₈H₁₀INO (M= 263.1 g/mol)
ESI-MS: 264 [M+H]⁺
Rt (HPLC): 2.22 min (method F)

2-Chloro-4-propoxypyridine (XXX):

20 C₈H₁₀CINO (M= 171.6 g/mol)
ESI-MS: 172 [M+H]⁺
Rt (HPLC): 1.85 min (method F)

Example XXXI4-ido-2-isopropoxy-pyridine


5 To 2.30 g (10.4 mmol) 4-ido-1 H-pyridin-2-one in 130 mL DCM are added 0.88 mL (11.4 mmol) 2-propanol and 3.00 (11.4 mmol) triphenylphosphine. After cooling down to 0 °C 2.23 mL (11.4 mmol) DIAD are added dropwise. After 5 min the cooling is removed and the mixture is stirred at r.t. for 2 h. The reaction mixture is washed with water and brine, dried with Na_2SO_4 and the solvent is removed in vacuo. The crude

10 product is purified by column chromatography (silica gel, heptane/EtOAc 90/1 0).

$\text{C}_8\text{H}_10\text{INO}$ (M= 263.1 g/mol)

ESI-MS: 264 [M+H]⁺

Rt(HPLC): 3.72 (method C)

15 Example XXXIIExample XXXII.1 (general route)7-Bromo-8-fluoro-3,4-dihydro-2H-benzo-[1,4]-dioxepine

20 a) To 1.00 g (7.57 mmol) 4-fluorocatechol in 50 mL DCM are added 0.58 mL (11.4 mmol) bromine in 10 mL DCM. The reaction mixture is stirred at r.t. for 3 h. Then the solvent is removed in vacuo and the crude product is purified by column chromatography (silica gel, DCM/MeOH 9/1).

25 b) To 1.50 g (7.25 mmol) of 4-bromo-5-fluorobenzene-1,2-diol in 25 mL DMF are added 5.90 g (18.1 mmol) Cs_2CO_3 and 0.89 mL (8.70 mmol) 1,3-dibromopropane. The reaction mixture is stirred at 120 °C over night. The solvent is removed in vacuo



and to the residue is added water. After extracting several times with EtOAc the org. phases are combined, washed with sat. aq. NaCl solution and dried with MgSO_4 . The solvent is removed in vacuo and the resulting residue is purified by column chromatography (silica gel, PE/EtOAc 9/1 \rightarrow 7/3).

5 $\text{C}_9\text{H}_8\text{BrFO}_2$ (M= 247.1 g/mol)

EI-MS: 246/248 $[\text{M}]^+$

R_f (TLC): 0.50 (silica gel, PE/EtOAc 4/1)


The following compounds are prepared analogously to example XXXI 1.1

Ex.	Starting material	Structure	Mass spec result	TLC R_f -value (silica gel)
XXXII.1	4-fluorocatechol		246/248 $[\text{M}]^+$	0.50 (PE/EtOAc 4/1)
XXXII.2	4-fluorocatechol		232/234 $[\text{M}^*]^+$	0.59 (PE/EtOAc 4/1)

10

Example XXXIII

(S)-N-(1-(4-(1-(3-Hydroxyphenyl)azetidin-3-yloxy)phenyl)ethyl)acetamide




To 0.80 g (2.96 mmol) of the amine XIII. 1 in 10 mL dioxane are added 0.31 mL (2.96

15 mmol) 3-bromophenol, 1.42 g (14.8 mmol) NaOtBu 135 mg (0.15 mmol) Pd_2dba_3 and and 176 mg (0.59 mmol) 2-(di-tert-butylphosphino)biphenyl. The mixture is degassed thoroughly and stirred at 80 °C for 2 h. Some water is added and the mixture is purified by HPLC (MeOH/H₂O/NH₃).

C₁₉H₂₂N₂O₃ (M= 326.4 g/mol)

20 ESI-MS: 327 $[\text{M}+\text{H}]^+$

Rt(HPLC): 0.86 (method K)

Example XXXIV(S)-1 -(4-(1 -(5-Bromopyrimidin-2-yl)azetidin-3-yloxy)phenyl)ethylamine

5

a) 600 mg (2.05 mmol) of the azetidine XV and 317 mg (2.05 mmol) of the pyrimidine XXI.1 are added to 15 mL DMSO and 0.53 mL (3.08 mmol) DIPEA. The mixture is stirred at 35 °C over night and afterwards purified by HPLC (MeOH/H₂O/NH₃).

10

b) 550 mg (1.34 mmol) of (S)-tert-butyl 1-(4-(1-(5-bromopyrimidin-2-yl)azetidin-3-yloxy)phenyl)ethylcarbamate are added to 3 mL MeOH and 3.22 mL (4.02 mmol) of a methanolic HCl solution (c = 1.3 mol/l). The reaction mixture is stirred at r.t. over night. The solvent is removed in vacuo and the residue is purified by HPLC (MeOH/H₂O/NH₃).

15


C₁₈H₂₂N₄O (M= 310.4 g/mol)

ESI-MS: 311 [M+H]⁺

Rt (HPLC): 2.01 min (method A)

Example XXXV

20

(S)-tert-Butyl 1-(4-(1-(4-ethoxyphenyl)azetidin-3-yloxy)phenyl)ethylcarbamate

To 360 mg (1.10 mmol) of amine XIII.6 in 10 mL dioxane are added 222 mg (1.10 mmol) 4-bromophenetole, 437 mg (4.41 mmol) NaOtBu, 101 mg (0.11 mmol) Pd₂dba₃ and 132 mg (0.44 mmol) 2-(di-tert-butylphosphino)biphenyl. The mixture is degassed thoroughly and stirred at 80 °C for 45 min in a microwave oven. A small

25

amount of water and MeOH are added, the mixture is filtered and afterwards purified by HPLC (MeOH/H₂O/NH₃).


C₂₄H₃₂N₂O₄ (M= 412.5 g/mol)

ESI-MS: 413 [M+H]⁺

5 R_t (HPLC): 2.36 min (method A)

Example XXXVI

(S)-1-(4-(1-(4-Ethoxyphenyl)azetidin-3-yloxy)phenyl)ethylamine



10 270 mg (0.66 mmol) of the example XXXV are added to 2 mL MeOH and 1.05 mL (1.31 mmol) of a methanolic HCl solution (c = 1.3 mol/l). The reaction mixture is stirred at r.t. for 2 h. The solvent is removed in vacuo and to the residue is added diluted aq. NaOH solution. The aq. mixture is extracted with DCM. The org. phase is washed with water, dried with MgSO₄ and the solvent is removed in vacuo.


15 C₁₉H₂₄N₂O₂ (M= 312.4 g/mol)

ESI-MS: 313 [M+H]⁺

R_t (HPLC): 0.89 min (method M)

Example XXXVII

(S)-Benzyl 1-(4-(1-(2-methoxy-4-propoxyphenyl)azetidin-3-yloxy)phenyl)-ethylcarbamate



To 3.70 g (8.16 mmol) of amine XIII.5 (used as its HCl salt) in 16 mL toluene and 3 mL

tert-butanol are added 2.00 g (8.16 mmol) of the aryl bromide XVI. 12, 6.64 g (20.4 mmol) Cs_2CO_3 , 91.6 mg (0.41 mmol) PdOAc_2 and 194 mg (0.41 mmol) X-Phos. The mixture is degassed thoroughly and stirred at 120 °C over night. EtOAc and a small amount of water are added and the resulting mixture is filtered, the solvent is

5 removed in vacuo and the residue is purified by HPLC (acetone/ $\text{H}_2\text{O}/\text{NH}_3$).


$\text{C}_{29}\text{H}_{34}\text{N}_2\text{O}_5$ (M= 490.6 g/mol)

ESI-MS: 491 [M+H]⁺

R_t (HPLC): 2.38 min (method G)

10 Example XXXVIII

(S)-1-(4-(1-(2-Methoxy-4-propoxyphenyl)azetidin-3-yloxy)phenyl)ethylamine




1.70 g (3.47 mmol) of example XXXVII is added to 50 mL MeOH and charged with

15 100 mg Pd/C (10%). The reaction mixture is hydrogenated at r.t. at a hydrogen pressure of 2 bar. Then the reaction mixture is filtered, the solvent is removed in vacuo and the residue is purified by HPLC (acetone/ $\text{H}_2\text{O}/\text{NH}_3$).

$\text{C}_{21}\text{H}_{28}\text{N}_2\text{O}_3$ (M= 356.5 g/mol)

ESI-MS: 357 [M+H]⁺

20 R_t (HPLC): 2.20 min (method A)

Example XXXIXExample XXXIX.1 (general route)(S)-N-(1-(4-(1-(2-Chloropyrimidin-4-yl)azetidin-3-yloxy)phenyl)ethyl)acetamide

5 1.00 g (3.69 mmol) of the azetidine XIII.6 and 605 mg (4.06 mmol) of 2,4-dichloropyrimidine are added to 30 mL DMSO and 0.57 mL (4.06 mmol) TEA. The mixture is stirred at r.t. over night, the solvent is removed in vacuo and the residue is purified by HPLC (MeOH/H₂O/NH₃).

C₁₇H₁₉ClN₄O₂ (M = 346.8 g/mol)


10 ESI-MS: 347 [M+H]⁺

Rt (HPLC): 0.84 min (method K)

The following compounds are prepared analogously to example XXXIX.1.1

For the example XXXIX.2 the reaction mixture is stirred at r.t. for 3 h.

Ex.	Starting material (s)	Structure	Mass spec result	HPLC retention time (method)
XXXIX.1	XIII.6		347 [M+H] ⁺	0.84 (K)
XXXIX.2	XIII.6		347 [M+H] ⁺	0.87 (K)

Example XLExample XL.1 (general route)3-Bromo-6-ethoxy-pyridine-2-carbonitrile

5 a) To 4.50 g (22.3 mmol) of 3-bromo-6-ethoxy-pyridine in 100 mL DCM are added 20.5 g (89.1 mmol) mCPBA by several portions. The reaction mixture is stirred at r.t. over night, charged with MgSO_4 and filtered over 100 g of basic aluminium oxide. The solvent is removed in vacuo before DCM and an aq. NaHCO_3 solution are added and the layers are separated. The aq. layer is saturated with NaCl and extracted

10 several times with DCM. The org. layers are combined and the solvent is removed in vacuo. The residue is filtered through a plug of silica gel.

b) 1.00 g (4.59 mmol) 5-bromo-2-ethoxy-pyridine 1-oxide, 2.46 mL (18.3 mmol) trimethylsilyl cyanide and 1.92 mL (13.8 mmol) TEA in 10 mL ACN are stirred together at 100 °C for 24 h. Then the volatile components are removed in vacuo and the residue is dissolved in DCM, washed with aq. NaHCO_3 solution, dried with MgSO_4 and filtered through a plug of silica gel. Then the solvent is removed in vacuo.

15

$\text{C}_8\text{H}_7\text{BrN}_2\text{O}$ ($M= 227.1$ g/mol),

ESI-MS: 227/229 $[\text{M}+\text{H}]^+$





R_f (TLC): 0.85 (silica gel, PE/EtOAc 7/3)


20

The following compounds are prepared analogously to example XL.1.

For the examples XL.2 and XL.3 the intermediates from step a are purified by HPLC after the filtration over basic aluminium oxide.

Example	Starting material	Product structure	Mass spec result	R_f (HPLC) [min] (method)
XL.1			227/229 $[\text{M}+\text{H}]^+$	TLC: $R_f = 0.85$ (silica gel, PE/EtOAc 7/3)





XL.2			241/243 [M+H] ⁺	1.09 (K)
XL.3			275/277 [M+Na] ⁺	1.08 (K)


Example XLIExample XLI.1 (general route)5 2-Chloro-5-((2,2-difluorocyclopropyl)methoxy)pyrimidine

295 mg (1.72 mmol) 1-bromomethyl-2,2-difluorocyclopropane and 318 mg (2.30 mmol) K₂CO₃ are added to a mixture of 150 mg (1.15 mmol) 2-chloro-5-hydroxypyrimidine in 2 mL DMF. The mixture is stirred at 80 °C over night. Then the 10 reaction mixture is quenched by the addition of water and extracted with DCM. The organic layer is dried with Na₂SO₄ and the solvent is removed in vacuo. The residue is purified by HPLC (MeOH/H₂O/NH₃).

C₈H₇ClF₂N₂O (M= 220.6 g/mol)
ESI-MS: 221 [M+H]⁺
15 R_t (HPLC): 1.21 min (method V)


The following compounds are prepared analogously to example XLI.1.

Example	Starting material	Product structure	Mass spec result	R _t (HPLC) [min] (method)
XLI.1			221 [M+H] ⁺	1.21 (V)
XLI.2			173 [M+H] ⁺	1.47 (A)

Example XLII4-Bromo-2-ethoxy-5-fluoropyridine

5 2.08 mL (26.0 mmol) Ethyliodide and 1.08 g (3.91 mmol) Ag_2CO_3 are added to a mixture of 500 mg (2.160 mmol) 4-bromo-5-fluoro-pyridin-2-ol in 10 mL DCM. The mixture is stirred at r.t. over night. Then the reaction mixture is quenched by the addition of water and DCM. After filtration the org. layer is separated, dried with Na_2SO_4 and the solvent is removed in vacuo.

10 $\text{C}_7\text{H}_7\text{BrFNO}$ (M = 220.0 g/mol)
 ESI-MS: 220 [M+H]⁺
 Rt (HPLC) : 1.27 min (method X)

Example XLIII5-tert-Butyl-2-chloropyrimidine

15 a) 25.0 g (0.21 mol) tert-Butylmalononitrile and 150 mL toluene are cooled down to -60 °C and 400 mL DIBALH (c = 1.5 mol/l in toluene) are carefully added. The reaction mixture is warmed to r.t. and stirred for 4 h. The reaction is quenched by the addition of aq. HCl solution (c = 1 mol/l). The resulting mixture is extracted with EtOAc, the org. layers are combined, dried with Na_2SO_4 and the solvent is removed in vacuo. The crude product is used without further purification.

20 b) 25 mL Hydrochloric acid (c = 10 mol/l) are added to a mixture of 5.62 g (0.09 mol) urea and 200 mL ethanol. The mixture is stirred for 10 min and then treated with 10.0 g of the above mentioned crude product. The resulting mixture is heated at reflux for 40 h, then concentrated and dried with toluene. The crude product is used without further purification.

c) 200 mL POCl_3 and 14 g (0.09 mol) 5-tert-butylpyrimidin-2(1 H)-one are heated in a sealed glass ampoule for 3 h at 160 °C. The mixture is cooled and the excess POCl_3 is removed in vacuo and the residue is treated with water and basified to pH 5 10 with aq. NaOH solution (c = 3 mol/l). The mixture is extract with DCM, dried with Na_2SO_4 and the solvent is removed in vacuo. The crude product is purified by flash column chromatography (silica gel, DCM/MeOH 98/2).

$\text{C}_8\text{H}_{11}\text{ClN}_2$ (M= 170.6 g/mol)

ESI-MS: 171 [M]⁺


10 R_f (HPLC): 6.34 (method W)

Example XLIV

Example XLIV.1 (general route)

A-[1 -(4-Bromo-3-fluoro-phenyl)-ethyl]1-acetamide

15



a) 11.1 g (91.0 mmol) 2-Methyl-2-propanesulfinamide are added to 16.5 g (76.0 mmol) 4-bromo-3-fluoro-benzaldehyde and 34.6 g (152 mmol) titanium(IV) ethoxide in 100 mL THF. Stirring is continued for 1.5 h at 50°C. After that time, the mixture is 20 poured into brine. The mixture is filtered over celite and the filtrate is transferred to a separation funnel. The organic layer is separated, dried over magnesium sulphate and the solvent is removed by evaporation. The residue is purified by column chromatography (silica gel; gradient hexane/EtOAc 1:0 → 3:2) to yield the desired product.

25 $\text{C}_9\text{H}_{11}\text{BrFNOS}$ (M = 306.2 g/mol),

ESI-MS: 306/308 [M+H]⁺

R_f (TLC): 0.30 (silica gel, hexane/EtOAc 4:1)

b) Under inert gas atmosphere 37.2 mL (112 mmol) of a 3N solution of methyl 30 magnesium bromide in THF are added dropwise to 17.1 g (55.8 mmol) 2-methyl-

propane-2-sulfonic acid 4-bromo-3-fluoro-benzylideneannide (IX.1) in 170 mL THF at -78°C. The cooling bath is removed and stirring is continued for 2 h. After that time, the mixture is poured into sat. NH₄Cl-solution (300 mL) and extracted with EtOAc. The organic layer is separated, washed with brine and dried over sodium sulphate.

5 The solvent is removed by evaporation to yield the desired product.

C₁₂H₁₇BrFNOS (M = 322.2 g/mol)

ESI-MS: 322/324 [M+H]⁺

c) 20.0 mL 4N HCl in dioxane are added to 17.2 g (53.4 mmol) 2-methyl-propane-2-sulfonic acid [1-(4-bromo-3-fluoro-phenyl)-ethyl]-amide (X.1) in 150 mL MeOH. 10 Stirring is continued for 1 h. After that time, the mixture is concentrated and the residue is triturated from diethylether (150 mL). The precipitate is filtered off, washed with diethylether and suspended in 250 mL DCM. 8.64 mL (107 mmol) pyridine and 5.29 mL (56.1 mmol) acetic anhydride are added and the mixture is stirred for 12 h at 15 rt. After that time, the mixture is transferred to a separation funnel, washed with 1N HCl (200 mL) and sat. aq. NaHCO₃-solution (200 mL) and dried over sodium sulphate. The solvent is removed by evaporation to yield the desired product.


C₁₀H₁₁BrFNO (M = 260.1 g/mol)

ESI-MS: 260/262 [M+H]⁺

20 R_t (HPLC): 2.60 min (method Y)

Example XLV

N-[1-(4-Hydroxy-3-fluoro-phenyl)-ethyl]-acetamide



25 Under inert gas atmosphere 200 mg (0.77 mmol) of example III are added to a 1/1 mixture of dioxane and water. Then 240 mg (4.20 mmol) powdered KOH and 58.7 mg (0.12 mmol) 2-di-*t*-butylphosphino-3,4,5,6-tetramethyl-2',4',6'-trisopropyl-1,1'-biphenyl are added and the mixture is degassed before 36.6 mg (0.04 mmol) Pd₂(dba)₃ are added and the mixture is stirred at 140 °C for 10 min in a microwave 30 oven. After cooling down, the reaction mixture is diluted with EtOAc, washed with an

aq. HCl solution ($c = 4$ mol/L) and a sat. aq. NaCl solution, dried over $MgSO_4$, filtered and the solvent is removed in vacuo. The resulting crude product is purified by HPLC (ACN/ H_2O /TFA).


C₁₀H₁₂FNO₂ (M = 197.2 g/mol)

5 ESI-MS: 198 [M+H]⁺

R_t (HPLC): 0.55 min (method AA)

Example XLVI

(S)-N-(1-(5-Hydroxypyridin-2-yl)ethyl)acetamide



To 1.00 g (5.73 mmol) (S)-6-(1-aminoethyl)pyridin-3-ol in 10 mL THF and 3.99 mL TEA (28.6 mmol) are slowly added 494 mg (6.30 mmol) acetyl chloride and stirring is continued for 2 h. The reaction is quenched by the addition of water and extracted several times with DCM. The org. layers are combined, dried over $MgSO_4$, filtered and the solvent is removed in vacuo. The crude product is purified by column chromatography (silica gel, DCM/MeOH/NH₃ 90/9/1).

15 C₉H₁₂N₂O₂ (M= 180.2 g/mol)


ESI-MS: 181 [M+H]⁺

R_t (HPLC): 0.15 min (method Z)

20

Example XLVII

1-(4-(Cyclopropylmethoxy)phenyl)azetidin-3-ol



25 a) 6.50 g (26.0 mmol) 3-(tert-butyldimethylsilyloxy)azetidine, 7.13 g (26.0 mmol) of 1-(cyclopropylmethoxy)-4-iodobenzene (WO2012/028676), 10.0 g (104 mmol) NaOtBu and 1.54 g (2.08 mmol) chloro(2-dicyclohexylphosphino-2',4',6'-triisopropyl-1,1'-biphenyl)(2-(2-aminoethyl)-phenyl)-palladium (II) are added to 65 mL dioxane and stirred at 75 °C for 2.5 h under inert gas atmosphere. Water is added and the

resulting mixture is extracted with EtOAc. The org. layers are combined, dried over MgSO_4 , filtered and the solvent is removed in vacuo. The resulting crude product is purified by HPLC (ACN/ $\text{H}_2\text{O}/\text{NH}_3$).

C₁₉H₃₁N₂O₂S₁ (M= 333.5 g/mol)

5 ESI-MS: 334 [M+H]⁺

R_t (HPLC): 1.46 min (method K)

b) To 3.65 g (10.9 mmol) of the above mentioned product in 60 mL THF are added 3.36 g (12.0 mmol) tetrabutylammoniumfluoride monohydrate and the mixture is

10 stirred at r.t. over night. 150 mL of a half saturated aq. NaHCO₃ solution is added and the resulting mixture is extracted with TBME (3x). The org. layers are combined, washed with diluted aq. NaHCO₃ solution, dried over MgSO_4 , filtered and the solvent is removed in vacuo. The resulting crude product is purified by HPLC (MeOH/ $\text{H}_2\text{O}/\text{NH}_3$).


15 C₁₃H₁₇NO₂ (M= 219.3 g/mol)

ESI-MS: 220 [M+H]⁺

R_t (HPLC): 0.76 min (method AB)

Example XLVIII

20 1-(4-(Cyclopropylmethoxy)phenyl)azetidin-3-yl methanesulfonate



To 1.50 g (6.84 mmol) of the above mentioned product in 10 mL DCM are added 2.38 mL (17.1 mmol) TEA and 635 μL (8.21 mmol) methanesulfonyl chloride and the mixture is stirred at r.t. for 1 h. Water is added and the mixture is extracted with DCM.

25 The org. layers are combined, washed with brine, dried over MgSO_4 , filtered and the solvent is removed in vacuo. The resulting crude product is triturated with diethylether.

C₁₄H₁₉NO₄S (M= 297.4 g/mol)

ESI-MS: 298 [M+H]⁺


30 R_t (HPLC): 0.86 min (method AB)

Preparation of Final Compounds

Example 1

5 Example 1.1 (general route)

N-(1-(4-(1 -(4-tert-Butylphenyl)azetidin-3-yloxy)phenyl)ethyl)acetamide



To 120 mg (0.51 mmol) of amine XIII.2 in 2.5 ml 1,4-dioxane are added 100 μ l (0.56 mmol) 1-bromo-4-tert-butyl-benzene, 203 mg (2.05 mmol) NaOtBu, 46.9 mg (0.05 mmol) Pd₂dba3 and 61.1 mg (0.21 mmol) 2-(di-tert-butylphosphino)biphenyl. The mixture is degassed thoroughly and stirred at 45 °C over night. A small amount of water and MeOH is added, the mixture is filtered and afterwards purified by HPLC (MeOH/H₂O/NH₃).

C₂₃H₃₀N₂O₂ (M= 366.5 g/mol)

15 ESI-MS: 367 [M+H]⁺

Rt (HPLC): 1.32 min (method L)









The following compounds are prepared analogously to example 1.1:

For the examples 1.26 - 1.31 and 1.53 - 1.54 the amount of amine in dioxane/DMF is added to the appropriate arylbromide and the reaction mixture is stirred at 45 °C for 3 d.

For the examples 1.32-1.35 the reaction temperature is 50 °C.

For the examples 1.36-1.43 and 1.55 the reactions are done in a microwave oven at 80 °C for 45 min.









25 For the examples 1.44-1.52 the reaction mixtures are stirred at 80-90 °C for 1-2 h.

Ex.	Starting material(s)	Structure	Mass spec result	HPLC retention time (method)
1.1	XIII.2		367 [M+H] ⁺	1.32 (L)
1.2	XIII.2		369 [M+H] ⁺	1.83 (A)
1.3	XIII.2 + XVII.1		367 [M+H] ⁺	2.00 (A)
1.4	XIII.2 + XVI.2		381 [M+H] ⁺	2.09 (A)
1.5	XIII.2		355 [M+H] ⁺	1.95 (A)
1.6	XIII.2 + XVI.3		395 [M+H] ⁺	2.16 (A)
1.7	XIII.2		356 [M+H] ⁺	1.62 (A)
1.8	XIII.2		391/393 [M+H] ⁺	1.87 (A)

1.9	XIII.2		370 [M+H] ⁺	2.00 (A)
1.10	XIII.2 + XVIII		383 [M+H] ⁺	1.91 (A)
1.11	XIV		381 [M+H] ⁺	2.36 (F)
1.12	XIV		369 [M+H] ⁺	2.10 (F)
1.13	XIII.1		369 [M+H] ⁺	2.09 (A)
1.14	XIII.1 + XVI.1		381 [M+H] ⁺	2.06 (A)
1.15	XIII.1 + XIX		382 [M+H] ⁺	1.81 (A)
1.16	XIII.1		383 [M+H] ⁺	1.86 (A)
1.17	XIII.1 + XVI.8		385 [M+H] ⁺	1.98 (A)

1.18	XIII.1 + XVI.18		411 [M+H] ⁺	2.11 (A)
1.19	XIII.1		355 [M+H] ⁺	1.96 (A)
1.20	XIII.1 + XVI.9		411 [M+H] ⁺	2.13 (A)
1.21	XIII.1 + XVI.12		399 [M+H] ⁺	2.10 (A)
1.22	XIII.1		366 [M+H] ⁺	1.89 (A)
1.23	XIII.1		404 [M+H] ⁺	2.07 (A)
1.24	XIII.1		369 [M+H] ⁺	1.84 (A)
1.25	XIII.1 + XVI.14		394 [M+H] ⁺	2.12 (A)
1.26	XIII.1		391 [M+H] ⁺	2.12 (A)

1.35	XIII.4		Chiral 395 [M+H] ⁺	1.11 (M)
1.36	XIII.1 + XVI.11		Chiral 411 [M+H] ⁺	1.17 (K)
1.37	XIII.1 + XVI.13		Chiral 399 [M+H] ⁺	1.05 (K)
1.38	XIII.1 + XVI.10		Chiral 413 [M+H] ⁺	1.13 (K)
1.39	XIII.4 + XVII.1		Chiral 393 [M+H] ⁺	1.15 (K)
1.40	XIII.4 + XVI.1		Chiral 407 [M+H] ⁺	1.20 (K)
1.41	XIII.4		Chiral 395 [M+H] ⁺	1.07 (K)
1.42	XIII.5		Chiral 398 [M+H] ⁺	1.23 (K)
1.43	XIII.6 + XVII.3		Chiral 397 [M+H] ⁺	1.13 (K)

1.44	XIII.6 + XVII.2		381 [M+H] ⁺	1.16 (K)
1.45	XIII.6 + XVII.4		403 [M+H] ⁺	1.23 (K)
1.46	XIII.6 + XVI.16		406 [M+H] ⁺	1.19 (K)
1.47	XIII.6 + XVI.15		406 [M+H] ⁺	1.16 (K)
1.48	XIII.6 + XVI.4		369 [M+H] ⁺	1.20 (K)
1.49	XIII.6 + XVI.5		383 [M+H] ⁺	1.30 (K)
1.50	XIII.6		383 [M+H] ⁺	1.17 (K)
1.51	XIII.1 + XXVII.1		356 [M+H] ⁺	1.82 (A)


1.52	XIII.1 + XXVII.2		370 [M+H] ⁺	1.94 (A)
1.53	XIII.6 + XXXII		401 [M+H] ⁺	1.07 (K)
1.54	XIII.6 + XVI.20		435 [M+H] ⁺	1.29 (N)
1.55	XIII.6 + XVII.6		385 [M+H] ⁺	1.30 (N)
1.56	XIII.6		379 [M+H] ⁺	1.33 (N)
1.57	XIII.6 + XXXII.2		387 [M+H] ⁺	1.10 (U)
1.58	XIII.6 + XL.1		381 [M+H] ⁺	1.23 (N)
1.59	XIII.6 + XL.2		395 [M+H] ⁺	1.29 (N)
1.60	XIII.6 + XL.3		407 [M+H] ⁺	1.29 (N)

1.61	XIII.6 + XVI.21		447 [M+H] ⁺	1.27 (N)
1.62	XIII.6 + XLII		374 [M+H] ⁺	1.24 (N)

Example 2

Example 2.1 (general route)

5 (S)-N-(1-(4-(1-(6-(cyclopropylmethoxy)pyridin-3-yl)azetidin-3-yl)phenyl)ethyl)acetamide



To 80 mg (0.30 mmol) of amine XIII.6 in 0.5 mL toluene and 0.2 mL tert-butanol are added 67.4 mg (0.30 mmol) 5-bromo-2-(cyclopropylmethoxy)pyridine (XXVIII. 3), 0.29 g (0.89 mmol) Cs₂CO₃, 3.35 mg (0.02 mmol) PdOAc₂ and 7.12 mg (0.02 mmol) X-
10 Phos. The mixture is degassed thoroughly and stirred at 120 °C over night. A small amount of water and MeOH are added and the mixture is purified by HPLC (MeOH/H₂O/NH₃).

C₂₂H₂₇N₃O₃ (M = 381.5 g/mol)

15 ESI-MS: 382 [M+H]⁺

Rt (HPLC): 2.05 min (method A)

The following compounds are prepared analogously to example 2.1:


Ex.	Starting material(s)	Structure	Mass spec result	HPLC retention time (method)
2.1	XIII.6 + XXVIII.3		382 [M+H] ⁺	2.05 (A)
2.2	XIII.1 + XVI.6		399 [M+H] ⁺	2.20 (A)
2.3	XIII.6 + XVI.7		373 [M+H] ⁺	2.11 (A)
2.4	XIII.1 + XVII.5		423 [M+H] ⁺	1.33 (N)
2.5	XIII.1 + XXVIII.5		383 [M+H] ⁺	2.22.(A)
2.6	XIII.1 + XXVIII.4		403 [M+H] ⁺	2.29.(A)
2.7	XIII.1		370 [M+H] ⁺	2.02 (A)
2.8	XIII.1 + XXVIII.2		404 [M+H] ⁺	2.04 (A)

2.9	XIII.1 + XXVIII.1		406 [M+H] ⁺	1.79 (A)
2.10	XIII.3		381 [M+H] ⁺	2.11 (A)
2.11	XIII.6 + XVI.17		399 [M+H] ⁺	1.28 (L)
2.12	XIII.6 + XXX		370 [M+H] ⁺	1.07 (A)

Example 3

Example 3.1 (general route)

5 (S)-N-(1-(4-(1-(5-bromopyrimidin-2-yl)azetidin-3-yloxy)phenyl)ethyl)acetamide



0.69 g (3.59 mmol) 5-bromo-2-chloro-pyrimidine and 0.70 g (2.99 mmol) of the amine XIII.1 are added to 10 mL DMSO and 0.78 mL (4.49 mmol) DIPEA. The mixture is stirred at 35 °C over night and afterwards purified by HPLC (MeOH/H₂O/NH₃).

10 Ci₇H₁₉BrN₄O₂ (M= 391.3 g/mol)

ESI-MS: 391/393 [M+H]⁺

R_t (HPLC): 1.80 min (method A)

15 The following compounds are prepared analogously to example 3.1 :







For the example 3.2 the reaction mixture is stirred for 4 d at 45 °C.




For the examples 3.3-3.4 and 3.8 the reaction mixture is stirred for 3 d at 35 °C.

For the example 3.6 the reaction mixture is stirred at 50 °C over night. Then some water and MeOH is added and the resulting precipitate is filtered, washed with water and dried in vacuo.

For the example 3.7 the reaction is done in ACN instead of DMSO and the reaction 5 mixture is stirred at 70 °C for 3 h.

For the examples 3.9 - 3.11 the reaction is done in ACN instead of DMSO and the reaction mixture is stirred at 60 °C over night.


Ex.	Starting material(s)	Structure	Mass spec result	HPLC retention time (method)
3.1	XIII.1		390/392 [M+H] ⁺	1.80 (A)
3.2	XIII.1		343 [M+H] ⁺	1.63 (A)
3.3	XIII.2		355 [M+H] ⁺	1.89 (A)
3.4	XIII.2		419 [M+H] ⁺	1.97 (A)
3.5	XIII.3 + XXI.1		379 [M+H] ⁺	2.15 (A)
3.6	XIII.3		416/418 [M+H] ⁺	2.05 (A)

3.7	XIII.3	 Chiral	419/421 [M+H] ⁺	2.01 (A)
3.8	XIII.3	 Chiral	409 [M+H] ⁺	1.12 (K)
3.9	XIII.6	 Chiral	395 [M+H] ⁺	2.00 (A)

Example 4

Example 4.1 (general route)

5 (S)-N-(1 -(4-(1-(5-Chloropyridin-2-yl)azetidin-3-yloxy)phenyl)ethyl)acetamide



To 14.8 mg (0.10 mmol) 2,5-dichloro-pyridine is added 23.4 mg (0.1 mmol) of the amine XIII.1 in 1.45 mL NMP and 0.05 mL (0.30 mmol) DIPEA. The mixture is stirred at 130 °C over night. Afterwards the solvent is removed in vacuo and the residue is purified by HPLC (MeOH/H₂O/TFA).

C18H20ClN3O2 (M= 345.8 g/mol)






ESI-MS: 346 [M+H]⁺










R_t (HPLC): 1.51 min (method E)

15

The following compounds are prepared analogously to example 4.1;

For the example 4.26 DMSO is used as solvent and the reaction mixture is stirred at 150 °C over night.

4.9	XIII.1		Chiral	403 [M+Na] ⁺	1.73 (E)
4.10	XIII.1		Chiral	398 [M+H] ⁺	2.01 (E)
4.11	XIII.1		Chiral	356 [M+H] ⁺	1.93 (E)
4.12	XIII.1		Chiral	377 [M+H] ⁺	1.10 (E)
4.13	XIII.1		Chiral	368 [M+H] ⁺	1.68 (E)
4.14	XIII.1		Chiral	381 [M+H] ⁺	1.69 (E)
4.15	XIII.1		Chiral	396 [M+H] ⁺	1.30 (E)
4.16	XIII.1		Chiral	381 [M+H] ⁺	1.87 (E)

4.17	XIII.1		389/391 [M+H] ⁺	1.56 (E)
4.18	XIII.1		389/391 [M+H] ⁺	1.12 (E)
4.19	XIII.1		380 [M+H] ⁺	1.76 (E)
4.20	XIII.1		380 [M+H] ⁺	1.72 (E)
4.21	XIII.1		380 [M+H] ⁺	1.90 (E)
4.22	XIII.1		392 [M+H] ⁺	1.25 (E)
4.23	XIII.1 + XXV.1		430 [M+H] ⁺	2.03 (E)
4.24	XIII.1		347 [M+H] ⁺	1.81 (E)
4.25	XIII.1		352 [M+H] ⁺	1.84 (E)


4.42	XIII.6 + XLI		419 [M+H] ⁺	1.42 (V)
4.43	XIII.6		398 [M+H] ⁺	1.14 (K)

* The appropriate 2-chloropyridyl can be prepared according to J. Med.Chem 1980, 23, 92 - 95.

Example 5

5 Example 5.1 (general route)

(S)-N-(1 -(4-(1-(5-Cyclopropylpyrimidin-2-yl)azetidin-3-yloxy)phenyl)ethyl)acetamide



To 0.1 g (0.26 mmol) of the arylbromide 3.1 , 65.9 mg (0.77 mmol) cyclopropyl
10 boronic acid, 0.19 g (0.90 mmol) K_3PO_4 , 11.5 mg (0.05 mmol) $PdOAc_2$ and 7.17 mg
(0.03 mmol) tricyclohexylphosphine are added 3 mL toluene and 150 μ l water. The
resulting mixture is degassed and stirred at 100 °C over night. Then the reaction
mixture is filtered and the solvent is removed in vacuo. The residue is purified by
HPLC (acetone/ H_2O/NH_3).

15 $C_{20}H_{24}N_4O_2$ (M= 352.4 g/mol)

ESI-MS: 353 [M+H]⁺

R_t (HPLC): 1.81 min (method A)

The following compounds are prepared analogously to example 5.1 :


20 For the example 5.2 3 eq of trimethylboroxine are used for the cross coupling.

Ex.	Starting material	Structure	Mass spec result	HPLC retention time (method)
5.1	3.1		353 [M+H] ⁺	1.81 (A)
5.2	3.1		327 [M+H] ⁺	1.64 (A)


Example 6

Example 6.1 (general route)

5 (S)-N-(1 -(4-(1-(5-Cyclopropylpyrimidin-2-yl)azetidin-3-yloxy)phenyl)ethyl)acetamide



To 50.0 g (0.13 mmol) of the arylbromide 3.1 and 19.7 mg (0.14 mmol) 4-fluoro-phenyl boronic acid in 1.5 mL dioxane and 1.5 mL methanol are added 0.13 mL (0.26 mmol) of an aq. Na₂CO₃ solution (c = 2 mol/l) and 3.58 mg (0.005 mmol) 10 Pd(PPh₃)₂Cl₂. The resulting mixture is stirred at 95 °C for 3 h. Then water and EtOAc are added and the layers are separated. The org. layer is filtered and the solvent is removed in vacuo. The residue is purified by HPLC (acetone/H₂O/NH₃).



15 ESI-MS: 407 [M+H]⁺

Rt (HPLC): 1.99 min (method A)

The following compounds are prepared analogously to example 6.1 :


Ex.	Starting material	Structure	Mass spec result	HPLC retention time (method)
6.1	3.1		407 [M+H] ⁺	1.99 (A)
6.2	3.1		420 [M+H] ⁺	1.87 (A)
6.3	1.8		353 [M+H] ⁺	1.81 (A)

5 Example 7

Example 7.1 (general route)

(S)-N-(1 -(4-(1-(3-(Cyclopropylmethoxy)phenyl)azetidin-3-yloxy)phenyl)ethyl)-acetamide

10



15

To 50.0 mg (0.15 mmol) of the phenol XXXIII and 22.0 μ l (0.23 mmol) (bromomethyl)-cyclopropane in 1 mL DMF are added 52.9 mg (0.38 mmol) K₂CO₃. The resulting mixture is stirred at 80 °C for 4 h. Then a small amount of water and methanol are added and the mixture is directly purified by HPLC (ACN/H₂O/FA).

C₂₃H₂₈N₂O₃ (M = 380.5 g/mol)

ESI-MS: 381 [M+H]⁺

Rt (HPLC): 1.18 min (method K)


The following compounds are prepared analogously to example 7.1 :

Ex.	Starting material	Structure	Mass spec result	HPLC retention time (method)
7.1	XXXIII		381 [M+H] ⁺	1.18 (K)
7.2	XXXIII		417 [M+H] ⁺	1.14 (K)
7.3	XXXIII		395 [M+H] ⁺	1.26 (K)

5

Example 8Example 8.1 (general route)(S)-N-(1-(4-(1-(2-Methoxy-4-propoxyphenyl)azetidin-3-yloxy)phenyl)ethyl)cyclopropanecarboxamide

10



15.5 mg (0.18 mmol) cyclopropanecarboxylic acid, 46.9 μ l (0.18 mmol) DIPEA and 75.1 mg (0.27 mmol) HATU are added to 5 mL THF and stirred for 5 min. Then

15 80.0 mg (0.18 mmol) of the amine XXXVIII in 2 mL THF are added and the resulting

mixture is stirred at r.t. for 5 h. Afterwards the mixture is directly purified by HPLC (acetone/H₂O/NH₃).

C₂₅H₃₂N₂O₄ (M= 424.5 g/mol)








ESI-MS: 425 [M+H]⁺









5 R_t (HPLC): 2.23 min (method A)








The following compounds are prepared analogously to example 8.1 :








For the examples 8.6 - 8.48 TBTU and DMF are used instead of HATU and THF.








Ex.	Starting material	Structure	Mass spec result	HPLC retention time (method)
8.1	XXXVIII		425 [M+H] ⁺	2.23 (A)
8.2	XXXVIII		417 [M+H] ⁺	2.20 (A)
8.3	XXXVIII		451 [M+H] ⁺	2.16 (A)
8.4	XXXVIII		468 [M+H] ⁺	2.22 (A)
8.5	XXXIV		405 [M+H] ⁺	1.94 (A)







8.6	XXXIV	 Chiral	378 [M+H] ⁺	1.92 (A)
8.7	XXXVI	 Chiral	407 [M+H] ⁺	1.05 (M)
8.8	XXXVI	 Chiral	380 [M+H] ⁺	1.04 (M)
8.9	XXXVI	 Chiral	395 [M+H] ⁺	0.63 (T)
8.10	XXXVI	 Chiral	497 [M+H] ⁺	0.61 (T)
8.11	XXXVI	 Chiral	474 [M+H] ⁺	0.62 (T)
8.12	XXXVI	 Chiral	406 [M+H] ⁺	0.62 (T)

8.13	XXXVI		381 [M+H] ⁺	0.62 (T)
8.14	XXXVI		419 [M+H] ⁺	0.61 (T)
8.15	XXXVI		395 [M+H] ⁺	0.62 (T)
8.16	XXXVI		383 [M+H] ⁺	0.62 (T)
8.17	XXXVI		371 [M+H] ⁺	0.61 (T)
8.18	XXXVI		423 [M+H] ⁺	0.62 (T)
8.19	XXXVI		413 [M+H] ⁺	0.62 (T)
8.20	XXXVI		395 [M+H] ⁺	0.62 (T)

8.21	XXXVI	 Chiral	449 [M+H] ⁺	0.64 (T)
8.22	XXXVI	 Chiral	452 [M+H] ⁺	0.62 (T)
8.23	XXXVI	 Chiral	398 [M+H] ⁺	0.61 (T)
8.24	XXXVI	 Chiral	432 [M+H] ⁺	0.60 (T)
8.25	XXXVI	 Chiral	475 [M+H] ⁺	0.61 (T)
8.26	XXXVI	 Chiral	399 [M+H] ⁺	0.61 (T)
8.27	XXXVI	 Chiral	409 [M+H] ⁺	0.63 (T)


8.28	XXXVI	 Chiral	495 [M+H] ⁺	0.62 (T)
8.29	XXXVI	 Chiral	369 [M+H] ⁺	0.61 (T)
8.30	XXXVI	 Chiral	373 [M+H] ⁺	0.61 (T)
8.31	XXXVI	 Chiral	408 [M+H] ⁺	0.62 (T)
8.32	XXXVI	 Chiral	481 [M+H] ⁺	0.62 (T)
8.33	XXXVI	 Chiral	408 [M+H] ⁺	0.61 (T)
8.34	XXXVI	 Chiral	424 [M+H] ⁺	0.62 (T)

8.35	XXXVI	 Chiral	481 [M+H] ⁺	0.62 (T)
8.36	XXXVI	 Chiral	413 [M+H] ⁺	0.63 (T)
8.37	XXXVI	 Chiral	421 [M+H] ⁺	0.63 (T)
8.38	XXXVI	 Chiral	418 [M+H] ⁺	0.60 (T)
8.39	XXXVI	 Chiral	424 [M+H] ⁺	0.61 (T)
8.40	XXXVI	 Chiral	407 [M+H] ⁺	0.63 (T)
8.41	XXXVI	 Chiral	424 [M+H] ⁺	0.62 (T)

8.42	XXXVI		Chiral	383 [M+H] ⁺	0.62 (T)
8.43	XXXVI		Chiral	440 [M+H] ⁺	0.61 (T)
8.44	XXXVI		Chiral	425 [M+H] ⁺	0.62 (T)
8.45	XXXVI		Chiral	418 [M+H] ⁺	0.60 (T)
8.46	XXXVI		Chiral	411 [M+H] ⁺	0.61 (T)
8.47	XXXVI		Chiral	385 [M+H] ⁺	0.61 (T)

Example 9

(S)-N-(1-(4-(1-(5-Cyclopropylpyrimidin-2-yl)azetidin-3-yl)oxy)phenyl)ethyl-
5 propionamide



60.0 mg (0.19 mmol) of the amine XXXIV are added to 1 mL DCM and charged with 0.11 mL (0.77 mmol) TEA and 0.02 mL (0.23 mmol) propionyl chloride. The resulting mixture is stirred at r.t. for 3 h. Afterwards a small amount of MeOH is added and the mixture is directly purified by HPLC (MeOH/H₂O/NH₃).

5 C₂₁H₂₆N₄O₂ (M= 366.5 g/mol)


ESI-MS: 367 [M+H]⁺

Rt (HPLC): 2.00 min (method A)

Example 10

10 Example 10.1 (general route)

(S)-N-(1-(4-(1-(2-Methoxy-4-propoxyphenyl)azetidin-3-yloxy)phenyl)ethyl)cyclopropanecarboxamide



15

65 mg (0.21 mmol) amine XXXIV and 0.06 mL (0.42 mmol) TEA are added to 1 mL DCM and charged with 36.1 mg (0.22 mmol) CDT. The reaction mixture is stirred at r.t. for 15 min. Then 28.3 mg (0.63 mmol) dimethylamine are added and stirring is continued for 3 h. Some MeOH is added and the mixture is directly purified by HPLC

20 (MeOH/H₂O/NH₃).

C₂₁H₂₇N₅O₂ (M= 381.5 g/mol)

ESI-MS: 382 [M+H]⁺


Rt (HPLC): 2.01 min (method A)

25 The following compounds are prepared analogously to example 10.1:


For the example 10.5 the CDT is added at 0 °C and stirring is continued at 0-5 °C for 15 min before the mixture is warmed to r.t., stirred for additional 5 min and then charged with the dimethylamine.

For the examples 10.6 - 10.7 a mixture of 1.8 eq NaOMe in 1 mL MeOH is added 30 after the CDT activation.

Ex.	Starting material	Structure	Mass spec result	HPLC retention time (method)
10.1	XXXIV		382 [M+H] ⁺	2.01 (A)
10.2	XXXIV		382 [M+H] ⁺	2.00 (A)
10.3	XXXVI		384 [M+H] ⁺	2.11 (A)
10.4	XXXVI		384 [M+H] ⁺	2.11 (A)
10.5	XXXVI		428 [M+H] ⁺	2.20 (A)
10.6	XXXIV		369 [M+H] ⁺	2.05 (A)
10.7	XXXVI		371 [M+H] ⁺	1.14 (M)

Example 11Example 11.1 (general route)(S)-N-(1-(4-(1-(6-Propoxypyrimidin-4-yl)azetidin-3-yloxy)phenyl)ethyl)acetamide

5 To 20.8 mg (0.35 mmol) 1-propanol in 1 mL THF is added 9.1 3 mg (0.38 mmol) NaH. The mixture is stirred at r.t. for 10 min and afterwards charged with 60.0 mg (0.17 mmol) of the chloropyrimidine XXXIX.2. The resulting mixture is stirred at 80 °C over night. Then the mixture is filtered and directly purified by HPLC (MeOH/H₂O/NH₃).



10 ESI-MS: 371 [M+H]⁺

Rt (HPLC): 1.04 min (method K)

The following compounds are prepared analogously to example 11.1:


Ex.	Starting material	Structure	Mass spec result	HPLC retention time (method)
11.1	XXXIX.2		371 [M+H] ⁺	1.04 (K)
11.2	XXXIX.2		371 [M+H] ⁺	1.02 (K)
11.3	XXXIX.2		357 [M+H] ⁺	0.95 (K)

11.4	XXXIX.2		Chiral	383 [M+H] ⁺	1.03 (K)
11.5	XXXIX.2		Chiral	383 [M+H] ⁺	1.04 (K)
11.6	XXXIX.1		Chiral	371 [M+H] ⁺	1.01 (K)

Example 12

Example 12.1 (general route)

5 (S)-N-(1-(4-(1-(6-((Cyclopropylmethyl)(methyl)amino)pyrimidin-4-yl)azetidin-3-yl)oxy)phenyl)ethyl)acetamide



10 60.0 mg (0.17 mmol) of the chloropyrimidine XXXIX.2, 31.6 mg (0.26 mmol) 1-cyclopropyl-N-methylmethanamine and 150 μ l (0.87 mmol) DIPEA are added to 1.5 mL NMP and stirred at 130 °C for 5 h. Then the mixture is filtered and directly purified by HPLC (MeOH/H₂O/NH₃).

C₂₂H₂₉N₅O₂ (M= 395.5 g/mol)

15 ESI-MS: 396 [M+H]⁺


Rt (HPLC): 1.04 min (method K)

The following compounds are prepared analogously to example 12.1 :

Ex.	Starting material	Structure	Mass spec result	HPLC retention time (method)
12.1	XXXIX.2		396 [M+H] ⁺	1.04 (K)
12.2	XXXIX.1		396 [M+H] ⁺	1.13 (K)

Example 13

5 N-(1-(4-(1-(4-(cyclopropylmethoxy)phenyl)azetidin-3-yl)-3-fluorophenyl)ethyl)acetamide



To 22.4 mg (0.11 mmol) of the phenole XLV in 2.5 ml THF are added 24.1 mg (0.11 mmol) of intermediate XLVII, 28.9 mg (0.11 mmol) triphenylphosphine and 25.3 mg (0.11 mmol) DBAD. The resulting mixture is stirred at 60 °C over night. To the mixture, 28.9 mg (0.11 mmol) triphenylphosphine and 25.3 g mg (0.11 mmol) DBAD were added and the mixture is stirred for further 5h at 60 °C. The solvent is evaporated, the residue is dissolved in 1 ml DMF and the mixture is purified by HPLC (ACN/H₂O/NH₃).


15 C₂₃H₂₇FN₂O₃ (M= 398.5 g/mol)

ESI-MS: 399 [M+H]⁺

Rt (HPLC): 0.54 (method AC)

Example 14

N-(1-{5-[1-(4-Cyclopropylmethoxy-phenyl)-azetidin-3-yloxy]-pyridin-2-yl}-ethyl)-acetamide



To 23.4 mg (0.13 mmol) of the phenole XLVI are added 29.7 mg (0.10 mmol) of the mesylate XLVIII in 2 mL DMF and 81.5 mg (0.25 mmol) Cs_2CO_3 . The resulting mixture is stirred at 80 °C over night. Then 1 mL water is added and mixture is directly purified by HPLC (ACN/H₂O/NH₃).

10 C₂₂H₂₇N₃O₃ (M= 381.5 g/mol)

ESI-MS: 382 [M+H]⁺

Rt (HPLC): 0.72 (method AD)

15 **Analytic methods**

1 HPLC

Method A

time (min)	Vol% water (incl. 0.2% NH ₄ OH)	Vol% methanol (incl. 3% water)
0.0	95	5
0.2	95	5
2.2	5	95
2.3	5	95
2.4	0	100
2.6	0	100

Analytical column: XBridge C 18 (Waters); 2.5 μm ; 3.0 x 30 mm;

column temperature: 40°C; flow: 1.3 mL/min;

Method B

time (min)	Vol% water (incl. 0.1% FA)	Vol% acetonitrile (incl. 0.1% FA)
0.0	95	5
0.1	95	5
3.1	2	98
4.5	2	98
5.0	95	5

Analytical column: X-terra™ MS C18 (Waters); 2.5 µm; 4.6 x 30 mm; column temperature: r.t.; flow: 1.0 mL/min; detection 210-420 nm.

Method C

time (min)	Vol% water (incl. 0.1% FA)	Vol% methanol (incl. 0.1% FA)
0.0	90	10
2.5	2	98
6.0	2	98

Analytical column: XBridge C18 (Waters); 3.5 µm; 2.1 x 50 mm; column temperature: 35°C; flow: 0.8 mL/min; detection 220-320 nm.

Method D

time (min)	Vol% water (incl. 0.2% FA)	Vol% methanol (incl. 3% water)
0.0	95	5
0.2	95	5
2.2	5	95
2.3	5	95
2.4	0	100
2.6	0	100

Analytical column: Zorbax StableBond C18 (Agilent); 1.8 µm; 3.0 x 30 mm; column temperature: 40°C; flow: 1.3 mL/min;

Method E

time (min)	Vol% water (incl. 0.1% TFA)	Vol% methanol
0.0	80	20
1.7	0	100
2.5	0	100
2.6	80	20

Analytical column: Sunfire C 18 (Waters); 3.5 μ m; 4.6 x 50 mm;

column temperature: 60°C; flow: 2 mL/min;

Method F

time (min)	Vol% water (incl. 0.2% NH ₄ OH)	Vol% methanol (incl. 3% water)
0.0	95	5
0.2	95	5
2.2	5	95
2.3	5	95
2.4	0	100
2.6	0	100

Analytical column: Gemini C 18 (Phenomenex); 2.5 μ m; 3.0 x 30 mm;

column temperature: 40°C; flow: 1.3 mL/min

Method G

time (min)	Vol% water (incl. 0.2% NH ₄ OH)	Vol% methanol (incl. 3% water)
0.0	95	5
0.2	95	5
2.2	5	95
2.3	5	95
2.4	0	100
2.6	0	100

Analytical column: XBridge C 18 (Waters); 2.5 μ m; 3.0 x 30 mm;

column temperature: 60°C; flow: 1.3 mL/min;

Method H

time (min)	Vol% water (incl. 0.05% TFA)	Vol% ACN
0.0	40	60
6	10	90
15	10	90
15.1	40	60

Analytical column: XBridge C18 (Waters) 3.5 μ m; 4.6 x 50 mm;
column temperature: r.t.;

Method I

time (min)	Vol% water (incl. 0.01 M NH4OAc)	Vol% ACN
0.0	50	50
6	10	90
15	10	90
15.1	40	60

Analytical column: Eclipse-XDB-C18 (Agilent), 5.0 μ m; 4.6 x 150 mm; column temperature: r.t.; flow: 1.0 ml/min.

Method J

time (min)	Vol% water (incl. 0.01 M NH4OAc)	Vol% ACN
0.0	70	30
8	10	90
15	10	90
15.1	70	30

Analytical column: XBridge C8 (Waters) 5.0 μ m; 4.6 x 150 mm;
column temperature: r.t.; flow: 1 ml/min.

Method K

time (min)	Vol% water (incl. 0.2% NH ₄ OH)	Vol% MeOH
0.0	95	5
0.05	95	5
1.40	0	100
1.80	0	100

Analytical column: XBridge C18 (Waters) 2.5 μ m; 3.0 x 30 mm;

column temperature: 60 °C; flow: 2.2 ml/min.

Method L

time (min)	Vol% water (incl. 0.2% NH ₄ OH)	Vol% MeOH
0.0	95	5
1.0	10	90
2.2	10	90
2.3	0	100
2.5	0	100

Analytical column: XBridge C18 (Waters) 2.5 μ m; 3.0 x 30 mm;

column temperature: 40 °C; flow: 1.3 ml/min.

Method M

time (min)	Vol% water (incl. 0.2% TFA)	Vol% MeOH
0.00	95	5
0.05	95	5
1.40	0	100
1.80	0	100

Analytical column: Sunfire C18 (Waters) 2.5 μ m; 3.0 x 30 mm;

column temperature: 60 °C; flow: 2.2 ml/min.

Method N

time (min)	Vol% water (incl. 0.1% NH ₄ OH)	Vol% MeOH	Flow [ml/min]
0.0	95	5	2.2
0.30	95	5	2.2
1.50	0	100	2.2
1.55	0	100	2.9
1.70	0	100	2.9

Analytical column: XBridge C 18 (Waters) 2.5 μ m; 3.0 x 30 mm;

column temperature: 60 °C.

Method O

time (min)	Vol% water (incl. 0.1% NH ₄ OH)	Vol% MeOH
0.0	95	5
0.15	95	5
1.70	0	100
2.10	0	100

Analytical column: XBridge C 18 (Agilent) 3.5 μ m; 4.6 x 30 mm;

column temperature: 60 °C; flow: 4 ml/min

Method P

time (min)	Vol% water (incl. 0.15% TFA)	Vol% MeOH
0.00	95	5
0.25	95	5
1.90	0	100
2.05	0	100

Analytical column: Microsorb C 18 (Waters) 5 μ m; 4.6 x 20 mm;

column temperature: r.t.; flow: 5.2 ml/min.

Method Q

time (min)	Vol% water (incl. 0.05% TFA)	Vol% ACN
0.0	70	30
8	10	90
15	10	90
15.1	70	30

Analytical column: XBridge C18 (Waters) 3.5 μm ; 4.6 x 150 mm;
column temperature: r.t.; flow: 1 ml/min.

Method R

time (min)	Vol% water (incl. 0.1% FA)	Vol% ACN
0.0	40	60
4	10	90
15	10	90
15.1	10	60

Analytical column: Symmetry C 18 (Waters) 3.5 μm ; 4.6 x 75 mm;
column temperature: r.t.; flow: 1 ml/min.

Method S

time (min)	Vol% water (incl. 0.1% TFA)	Vol% ACN (incl. 0.1% TFA)
0.0.	95	5
2.0.	0	100
2.49	0	100
2.50	95	5

Analytical column: Sunfire C18 (Waters) 3.5 μm ; 4.6 x 50 mm;
column temperature: 40 °C; flow: 1.5 ml/min.

Method T

time (min)	Vol% water (incl. 0.1% TFA)	Vol% MeOH	Flow [ml/min]
0.0	95	5	1.5
0.7	1	99	1.5
0.8	1	99	1.5
0.81	95	5	1.5

Analytical column: Ascentis Express C 18 (Supeico) 2.7 μ m; 2.1 x

50 mm; column temperature: 60 °C;

Method U

time (min)	Vol% water (incl. 0.1% FA)	Vol% MeOH	Flow [ml/min]
0.0	95	5	2.2
0.05	95	5	2.2
1.40	0	100	2.2
1.80	0	100	2.2

Analytical column: Sunfire C 18 (Waters) 2.5 μ m; 3 x 30 mm; column

temperature: 60 °C;

Method V

time (min)	Vol% water (incl. 0.1% TFA)	Vol% MeOH (incl. 0.1% TFA)	Flow [ml/min]
0.0	95	5	4.0
0.15	95	5	4.0
1.70	0	100	4.0
2.25	0	100	4.0

Analytical column: Sunfire C 18 (Waters) 3.5 μ m; 4.6 x 30 mm;

column temperature: 60 °C.

Method W

time (min)	Vol% water (incl. 0.1% FA)	Vol% MeOH	Flow [ml/min]
0.0	40	60	1.0
6	10	90	1.0
16	10	90	1.0
18	40	60	1.0
20	40	60	1.0

Analytical column: Kromasil C 18 100 (Akzo Nobel) 5.0 μ m; 4.6 x

250 mm; column temperature: r.t.;

Method X

time (min)	Vol% water (incl. 0.1% TFA)	Vol% MeOH	Flow [ml/min]
0.0	95	5	2.2
0.30	95	5	2.2
1.50	0	100	2.2
1.55	0	100	2.9
1.65	0	100	2.9

Analytical column: Sunfire C 18 (Waters) 2.5 μ m; 3 x 30 mm; column

temperature: 60 °C.

Method Y

time (min)	Vol% water (incl. 0.1% FA)	Vol% ACN (incl. 0.1% FA)
0.0	95	5
3.50	2	98
6.00	2	98

Analytical column: X-Bridge C 18 (Waters) 3.5 pm; 2.1 x 50 mm;

column temperature: 35 °C; flow: 0.8 ml/min.

Method Z

time (min)	Vol% water (incl. 0.1% TFA)	Vol% MeOH
0.0	95	5
0.05	95	5
1.40	0	100
1.80	0	100

Analytical column: StableBond C 18 (Waters) 1.8 μm ; 3.0 x 30 mm; column temperature: 60 °C; flow: 2.2 ml/min.

Method AA

time (min)	Vol% water (incl. 0.1% TFA)	Vol% ACN	Flow [mL/min]
0.00	97	3	2.2
0.20	97	3	2.2
1.20	0	100	2.2
1.25	0	100	3
1.40	0	100	3

Analytical column: Sunfire C 18 (Waters) 2.5 μm ; 3.0 x 30 mm; column temperature: 60 °C.

Method AB

time (min)	Vol% water (incl. 0.1% NH ₄ OH)	Vol% MeOH	Flow [ml/min]
0.0	97	3	2.2
0.20	97	3	2.2
1.20	0	100	2.2
1.25	0	100	3.0
1.40	0	100	3.0

Analytical column: XBridge C 18 (Waters) 2.5 μm ; 3.0 x 30 mm; column temperature: 60 °C.

Method AC

time (min)	Vol% water (incl. 0.1% TFA)	Vol% ACN (incl. 0.08% TFA)
0.0	95.0	5.0
0.75	0.0	100.0
0.85	0.0	100.0

Analytical column: Sunfire C 18 (Waters) 2.5 μm ; 2.1 x 50 mm; column

temperature: 60 °C; flow: 1.5 ml/min.

Method AD

time (min)	Vol% water (incl. 0.1% NH ₄ OH)	Vol% ACN	Flow [ml/min]
0.0	98	3	2.0
1.20	0	100	2.0
1.40	0	100	2.0

Analytical column: XBridge C 18 (Waters) 2.5 μm ; 3.0 x 30 mm;

column temperature: 60 °C.

2 GC

5

Method AAA


Analytical column: SLB-5MS 15 m, ID 100 μm , df 0.10 μm .

Average velocity 45 cm/s, carrier gas : He, split ratio: 300 :1, injector temp: 250 °C, injection volume: 1 μL .

Initial temp: 60 °C, initial time: 1.0 min, solvent delay: 0.6 min, rate: 50 °C/min, final temp: 250 °C, final time: 1.0 min.

Patent Claims:

1. A compound of general formula I



5

wherein

Ar¹ is selected from the group Ar¹-G1 consisting of:

10 6- to 10-membered aryl and 5- to 10-membered heteroaryl, which may each
be substituted with one or more substituents R¹,

15 wherein two substituents R¹ linked to adjacent C-atoms of Ar¹ together
may form a C₃₋₅-alkylene bridge in which 1, 2 or 3 C₂H₂-groups may be
replaced independently of each other by O, C(=O), S, S(=O), S(=O)₂, NH
or N(Ci-4-alkyl)-, wherein the alkylene bridge may optionally be substituted
by one or two F atoms or one or two Ci-3-alkyl groups; and

R¹ is selected from the group R¹-G1 consisting of:

20 H, F, Cl, Br, I, CN, OH, -NO₂, Ci₁₋₆-alkyl, C₂₋₆-alkenyl, C₂₋₆-alkynyl, C₃₋₁₀-
carbocyclyl, C₃₋₁₀-carbocyclyl-Ci₃-alkyl, Ci₆-alkyl-O-, C₃₋₆-alkenyl-O-, C₃₋₆-
alkynyl-O-, C₃₋₁₀-carbocyclyl-O-, C₃₋₁₀-carbocyclyl-Ci₃-alkyl-O-, Ci₆-alkyl-S-,
Ci₆-alkyl-S(=O)-, Ci₆-alkyl-S(=O)₂-, C₃₋₁₀-carbocyclyl-S-, C₃₋₁₀-carbocyclyl-C₁₋₃-
alkyl-S-, Ci₄-alkyl-C(=O)-, C₃₋₁₀-carbocyclyl-C(=O)-, R^{N1}R^{N2}N-, R^{N1}R^{N2}N-
25 C₂₋₃-alkyl-O-, R^{N1}R^{N2}N-C(=O)-, R^{N1}R^{N2}N-S(=O)₂-, Ci₆-alkyl-C(=O)-N R^{N1}-, Ci₆-
alkyl-S(=O)₂-N R^{N1}-, Ci₆-alkyl-C(=O)-N R^{N1}-Ci₃-alkyl-, HO-C(=O)-, Ci₆-
alkyl-O-C(=O)-, heterocyclyl, heterocyclyl-O-, heterocyclyl-Ci₃-alkyl,
heterocyclyl-Ci₃-alkyl-O-, heterocyclyl-C(=O)-, aryl, aryl-Ci₃-alkyl, aryl-O-,
aryl-Ci₃-alkyl-O-, heteroaryl, heteroaryl-Ci₃-alkyl, heteroaryl-O- and
30 heteroaryl-Ci₃-alkyl-O-,

wherein in each carbocydyl and heterocyclyl a CH₂-group may optionally be replaced by -C(=O)- or -C(=CR^{N2}₂)-, and

5 wherein each carbocydyl and heterocyclyl may be optionally substituted with one or more Ci₄-alkyl, which may be optionally substituted with one or more substituents R^c, and

10 wherein each alkyl, carbocydyl and heterocyclyl may be optionally substituted with one or more substituents R^c, and

wherein each heterocyclyl may be optionally substituted with aryl or heteroaryl, and

15 wherein each aryl and heteroaryl group may be optionally substituted with one or more substituents R²,

R^c is selected from the group R^c-G1 consisting of:

20 F, Cl, Br, CN, OH, Ci₄-alkyl -O-, C₃₋₇-cycloalkyl -O-, C₃₋₇-cycloalkyl -Ci₃-alkyl-
O-, H₂N-, (Ci₄-alkyl)NH-, (Ci₄-alkyl)₂N-, Ci₄-alkyl -C(=O)-, Ci₄-alkyl -S(=O)₂-,
HO-C(=O)- and Ci₄-alkyl -O-C(=O)-, wherein each alkyl or cycloalkyl may be
optionally substituted with one or more substituents selected from F and OH,
and

25 R^{N1} is selected from the group R^{N1}-G1 consisting of:
H, Ci-6-alkyl, C3-alkyl-carbocydyl, C3-alkyl-carbocydyl-Ci-3-alkyl, C3-alkenyl, C3-alkynyl,
heterocyclyl, heterocyclyl -Ci-3-alkyl, aryl, aryl -Ci-3-alkyl, heteroaryl
and heteroaryl -Ci-3-alkyl,

30 wherein each carbocydyl and heterocyclyl may be optionally substituted with one or more Ci₄-alkyl, and

wherein in each carbocydyl and heterocyclyl a CH₂-group may optionally be replaced by -C(=O)-, and

wherein each alkyl, carbocyclyl and heterocyclyl may be optionally substituted with one or more substituents R^c , and

5 wherein each aryl and heteroaryl group may be optionally substituted with one or more substituents R^2 ,

R^{N2} is selected from the group R^{N2} -G1 consisting of: H and Ci-6-alkyl, and

10 Ar^2 is selected from the group Ar^2 -G1 consisting of:
phenylene and a 5- or 6-membered monocyclic heteroaryl group containing 1, 2 or 3 heteroatoms selected from N, O, or S, wherein all of the before mentioned groups may be optionally substituted with one or more substituents R^2 , and

15 R^2 is selected from the group R^2 -G1 consisting of:
F, Cl, Br, CN, OH, Ci₄-alkyl-, Ci₄-alkyl-O-, Ci₄-alkyl-S-, Ci₄-alkyl-S(=O)₂-, H₂N-, (Ci₄-alkyl)NH-, (Ci₄-alkyl)₂N-, Ci₄-alkyl-C(=O)-NH- and heterocyclyl,

20 wherein each alkyl may be optionally substituted with one or more F-atoms and/ or with a substituent selected from OH, Ci-3-alkyl-O- and CN; and

wherein two substituents R^2 attached to an aryl or heteroaryl group may be linked to each other and form a C2-5-alkylene bridging group in which 1 or 2 -
25 CH_2 -groups may be replaced by a group independently of each other selected from O, S, NH and N(Ci₄-alkyl)-, wherein the C2-5-alkylene bridging group is optionally be substituted by 1 or 2 Ci-3-alkyl groups;

30 L is selected from the group L-G1 consisting of:
a Ci₄-alkylene group that is optionally substituted by one or two CH_3 ;

X is selected from the group X-G1 consisting of: O and S;

R is selected from the group R-G1 consisting of: H and Ci-3-alkyl; and

T is selected from the group T-G1 consisting of:

Ci-6-alkyl, C2-6-alkenyl, C2-6-alkynyl, C3-*io*-carbocyclyl, C3-*io*-carbocyclyl-Ci-3-alkyl, Ci-6-alkyl-O-, C3-*io*-carbocyclyl-O-, C3-*io*-carbocyclyl-Ci-3-alkyl-O-, Ci-6-alkyl-S-, C3-*io*-carbocyclyl-S-, C3-*io*-carbocyclyl-Ci-3-alkyl-S-, Ci₄-alkyl-C(=O)-, Ci-4-alkyl-S(=O)₂⁻, R^{N1}R^{N2}-N-, R^{N1}R^{N2}-N-Ci₃-alkyl-, R^{N1}R^{N2}-N-CO-, Ci₄-alkyl-C(=O)-R^{N2}N-Ci-3-alkyl, heterocyclyl, aryl and heteroaryl,

wherein in each carbocyclyl and heterocyclyl a CH₂-group may optionally be replaced by -C(=O)-, and

wherein each carbocyclyl and heterocyclyl may be optionally substituted with one or more Ci₄-alkyl, which may be optionally substituted with one or more substituents R^c, and

wherein each alkyl, carbocyclyl and heterocyclyl may be optionally substituted with one or more substituents R^c, and

wherein each alkyl may optionally be substituted with a heteroaryl group; and

wherein each aryl and heteroaryl group may be optionally substituted with one or more substituents R²;

including any tautomers and stereoisomers thereof,

or a salt thereof.

2. A compound according to claim 1 wherein

X is O, and

30 R is H or CH₃.

3. A compound according to claim 1 or 2 wherein Ar¹ is selected from a group consisting of:

phenyl, pyridinyl, pyrimidinyl, benzoxazolyl, benzoisoxalyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinoxalinyl and oxazolopyhmidinyl,

wherein the above-mentioned phenyl, pyridinyl and pyrimidinyl are each
5 substituted with one to three groups independently selected from R¹, and/or

wherein two adjacent carbon atoms of a phenyl group may be linked to each other via a -O-CH₂-O-, -O-CF₂-O-, -O-CH₂-CH₂-O-, -O-CH₂-CH₂-CH₂-CH₂-, -O-CH₂-CH₂-CH₂-C(CH₃)₂-, -O-CH₂-CH₂-CH₂-O- or -O-CH₂-C(CH₃)₂-CH₂-O-
10 bridge, and

wherein the above-mentioned benzoxazolyl, benzoisoxalyl, benzthiazolyl, quinolinyl, isoquinolinyl, quinoxalinyl and oxazolopyhmidinyl groups are each optionally substituted with Cl, CH₃ or phenyl,

15

wherein R¹ is defined according to claim 1.

20

4. A compound according to any one of claims 1 to 3 wherein R¹ is selected from the group consisting of:

H, F, Cl, Br, CN, OH, Ci₁₋₄-alkyl, C₂₋₄-alkenyl, C₃₋₇-cycloalkyl, C₃₋₇-cycloalkyl-CH₂-, Ci₁₋₄-alkyl-O-, C₃₋₇-cycloalkyl-O-, C₃₋₇-cycloalkyl-Ci₁₋₃-alkyl-O-, Ci₁₋₄-alkyl-S(=O)₂, R^{N1}R^{N2}N-, HO-C(=O)-, Ci₁₋₄-alkyl-O-C(=O)-, phenyl, phenyl-O-, phenyl-CH₂-O-, pyridinyl,

25

wherein each alkyl, cydoalkyl and heterocydyl may be optionally substituted with one to three substituents independently selected from the group consisting of F and CN, and

30

wherein each phenyl and pyridinyl group may be optionally substituted with one F or -OCH₃.

5. A compound according to one or more of the previous claims wherein Ar² is selected from the group consisting of: phenylene and pyridylene, wherein all of the before mentioned groups may be optionally substituted with one to three substituents R²,

5

wherein R² is F, Cl, Ci₁₋₂-alkyl-, Ci₁₋₂-alkyl-O-, Ci₁₋₃-alkyl-C(=O)-NH or HO-CH₂-C(=O)-NH-.

10 6. A compound according to one or more of the previous claims wherein L is a linear Ci-2-alkylene group that is optionally substituted with one CH₃.

15 7. A compound according to one or more of the previous claims wherein T is selected from the group consisting of:


Ci₁₋₄-alkyl, C₂₋₄-alkenyl, C₃₋₆-cycloalkyl, C₃₋₇-cycloalkenyl, C₃₋₅-cycloalkyl-Ci₁₋₃-alkyl-, heteroaryl-Ci₁₋₃-alkyl-, Ci₁₋₃-alkyl-O-, R^{T1}R^{T2}-N-, Ci₁₋₃-alkyl-C(=O)-R^{T2}N-Ci₁₋₃-alkyl-, heterocydyl, phenyl and heteroaryl,

20 wherein in each heterocydyl, a CH₂-group may optionally be replaced by -C(=O)-; and

25 wherein each cydoalkyl and heterocydyl may be optionally substituted with one or more Ci₁₋₃-alkyl, which may be optionally substituted with one or more substituents R^c; and

wherein each alkyl, cydoalkyl and heterocydyl may be optionally substituted with one or more F; and

30 wherein heterocydyl is defined as hereinbefore and hereinafter, preferably heterocydyl is selected from the group consisting of



wherein heteroaryl is selected from the group consisting of pyridyl, pyrimidinyl,

5 pyridazinyl, pyrazinyl, furanyl, thieryl, pyrrolyl, imidazolyl, pyrazolyl, triazolyl, oxazolyl, isoxazolyl, thiazolyl and isothiazolyl; and

wherein each phenyl and heteroaryl group may be optionally substituted with one or more substituents selected from the group consisting of CH_3 , $\text{CH}_3\text{-C}(=\text{O})\text{-NH-}$ and

10 $\text{HO-CH}_2\text{-C}(=\text{O})\text{-NH-};$ and

wherein R^c is selected from the group consisting of F, Cl, CN, OH and Ci-3-alkyl-; and

15 wherein R^{t1} is selected from the group consisting of H, methyl and ethyl; and

wherein R^{t2} is H or methyl.

20 8. A compound according to claim 1, wherein


Ar^1 is selected from the group consisting of phenyl, pyridinyl and pyrimidinyl,

wherein the above-mentioned phenyl and pyridinyl groups are each substituted
25 with one to three groups independently selected from R^1 , and

wherein the above-mentioned pyrimidinyl groups is substituted with one or two R^1 , and

wherein two adjacent carbon atoms of a phenyl group may be linked to each other via a -O-CH₂-CH₂-O-, -O-CH₂-CH₂-CH₂-C(CH₃)₂-, -O-CH₂-CH₂-CH₂-CH₂-O-, or -O-CH₂-C(CH₃)₂-CH₂-O- bridge;

5 R¹ is selected from the group consisting of F, Cl, Br, CN, CF₃, C₁₋₄ alkyl, -O-C₁₋₄ alkyl, cyclopropyl, -O-C₃₋₅ cycloalkyl, -O-CH₂- cyclopropyl, and



Ar² is phenylene;

10 L is a linear C₁₋₂-alkylene group that is optionally substituted with one CH₃;

X is O;

R is H; and


15 T is selected from the group consisting of: C₁₋₄-alkyl, C₂₋₄-alkenyl, C₃₋₅-cycloalkyl, C₅₋₆-cycloalkenyl, cyclopropyl-CH₂-, CH₃O-, R^{T1}R^{T2}-N-, CH₃-C(=O)-NH-C₁₋₃-alkyl, heterocyclyl, phenyl and heteroaryl,

wherein R^{T1} is H, CH₃ or CH₂CH₃; and

R^{T2} is H or CH₃; and

20 wherein each alkyl, cycloalkyl and heterocyclyl may be optionally substituted with one F, CN, CH₃, CF₃, cyclopropyl, -N(CH₃)₂, pyridinyl, OH or -OCH₃; and

25 wherein heterocyclyl is selected from the group consisting of:



30 wherein heteroaryl is selected from the group consisting of pyridyl, pyrimidinyl, pyridazinyl, furanyl, pyrazolyl, oxazolyl, isoxazolyl, thiazolyl and isothiazolyl; and


wherein each phenyl and heteroaryl group may be optionally substituted with one or two substituents independently selected from CH_3 , $-\text{NH}-\text{CO}-\text{CH}_3$, $-\text{NH}-\text{CO}-\text{CH}_2-\text{OH}$;


5

including any tautomers and stereoisomers thereof,
or a salt thereof.

9. A compound according to claim 1, which is:

10





or a pharmaceutically acceptable salt thereof.

5 10. A pharmaceutically acceptable salt of a compound according to one or more of the claims 1 to 9.

11. A pharmaceutical composition comprising one or more compounds according to one or more of the claims 1 to 9 or one or more pharmaceutically acceptable salts 10 thereof, optionally together with one or more inert carriers and/or diluents.

12. A method for treating diseases or conditions which are mediated by inhibiting the activity of acetyl-CoA carboxylase(s) in a patient in need thereof characterized in that a compound according to one or more of the claims 1 to 9 or a pharmaceutically acceptable salt thereof is administered to the patient.

15 13. A method according to claim 12, wherein the disease or condition mediated by inhibiting the activity of acetyl-CoA carboxylase(s) is obesity, diabetes or dyslipidemia.

14. A compound according to any one of claim 1 to 9 or a pharmaceutically acceptable salts thereof for use in a method of treating obesity, diabetes or dyslipidemia.

5 15. Use of a compound according to any one of claim 1 to 9 or a pharmaceutically acceptable salts thereof for the preparation of a medicament for treating obesity, diabetes or dyslipidemia.

10 16. A pharmaceutical composition comprising one or more compounds according to one or more of the claims 1 to 9 or one or more pharmaceutically acceptable salts thereof and one or more additional therapeutic agents, optionally together with one or more inert carriers and/or diluents.

INTERNATIONAL SEARCH REPORT

International application No

PCT/ EP2012/075996

A. CLASSIFICATION OF SUBJECT MATTER					
I NV . C07D401/04	C07D403/04	C07D405/04	C07D407/04	C07D413/04	
A61 K3 1/397	A61 K3 1/4439	A61 K3 1/506	C07D40 1/ 12	C07D401/ 14	
C07D403/ 12	C07D403/ 14	C07D405/ 12	C07D417/O4	C07D417/ 12	

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

C07D A61 K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EPO- Internal , CHEM ABS Data , WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 2010/ 1272 12 A1 (FOREST LAB HOLDINGS LTD; SARMA PAKALA KUMARA SAVITHRU [IN]; ACHARYA VI) 4 November 2010 (2010 - 11-04) the whole document -----	1- 16
A	JP 2008 17962 1 A (TAISHO PHARMA CO LTD) 7 August 2008 (2008-08-07) the whole document -----	1- 16
A	JP 2010 0430 19 A (TAISHO PHARMA CO LTD) 25 February 2010 (2010-02 -25) the whole document -----	1- 16
A	WO 03/072 197 A1 (PFI ZER PROD INC [US]; PERRY DAVID AUSTEN [US]; HARWOOD HAROLD JAMES JR) 4 September 2003 (2003 -09 -04) the whole document ----- -/-	1- 16

Further documents are listed in the continuation of Box C.

See patent family annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

23 January 2013

Date of mailing of the international search report

31/01/2013

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040,
Fax: (+31-70) 340-3016

Authorized officer

Sotoca Usina, E

INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2012/075996

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	JP 2006 131559 A (TAKEDA CHEMICAL INDUSTRIES LTD) 25 May 2006 (2006-05-25) the whole document -----	1-16
A	US 2011/263562 AI (YAMASHITA TOHRU [JP] ET AL) 27 October 2011 (2011-10-27) the whole document -----	1-16

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/EP2012/075996

Patent document cited in search report		Publication date	Patent family member(s)			Publication date
wo 2010127212	AI	04-11-2010	US 2010280067	AI	04-11-2010	
			wo 2010127212	AI	04-11-2010	
JP 2008179621	A	07-08-2008	NONE			
JP 2010043019	A	25-02-2010	NONE			
wo 03072197	AI	04-09-2003	AT 303178	T	15-09-2005	
			AU 2003248354	AI	09-09-2003	
			CN 1642599	A	20-07-2005	
			CO 5601002	A2	31-01-2006	
			DE 60301491	DI	06-10-2005	
			DE 60301491	T2	18-05-2006	
			EP 1478437	AI	24-11-2004	
			ES 2246481	T3	16-02-2006	
			HR P20040716	A2	28-02-2005	
			IS 7391	A	09-08-2004	
			MA 27178	AI	03-01-2005	
			NZ 534582	A	31-03-2006	
			US 2003187254	AI	02-10-2003	
			wo 03072197	AI	04-09-2003	
			ZA 200406332	A	28-09-2005	
JP 2006131559	A	25-05-2006	NONE			
US 2011263562	AI	27-10-2011	AR 080969	AI	23-05-2012	
			AU 2011246004	AI	29-11-2012	
			CA 2797767	AI	03-11-2011	
			SG 185759	AI	28-12-2012	
			TW 201206911	A	16-02-2012	
			US 2011263562	AI	27-10-2011	
			UY 33354	A	01-12-2011	
			wo 2011136385	AI	03-11-2011	