




(72) HOUTHOFF, HENDRIK JAN, NL

(72) REEDIJK, JAN, NL

(72) VOLKERS, HERMAN H., NL

(72) HEETEBRIJ, ROBERT JOCHEM, NL


(71) KREATECH BIOTECHNOLOGY B.V., NL

(51) Int.Cl.⁶ C07F 15/00, G01N 33/543, C07H 23/00

(30) 1997/04/10 (97201066.4) EP

(54) **COMPOSE A BASE DE TRANS-PLATINE ET NECESSAIRE DE
DIAGNOSTIC**

(54) **TRANS-PLATINUM COMPOUND, AND DIAGNOSTIC KIT**



(57) La présente invention concerne un composé à base de trans-platine, utile dans un procédé de marquage d'une molécule bio-organique et correspondant à la formule (I) dans laquelle A représente une fraction réactive de fixation sur un marqueur, D représente une fraction réactive de fixation sur une molécule bio-organique et X1 et X2 représentent des fractions inertes, semblables ou différentes.

(57) The present invention is concerned with a trans-platinum based compound for use in a method for labelling a bio-organic molecule, having formula (I), wherein A represents a reactive moiety for attachment to a label, D represents a reactive moiety for attachment to a bio-organic molecule, and X1 and X2 represent the same or different inert moieties.



**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07F 15/00, C07H 23/00, G01N 33/543	A1	(11) International Publication Number: WO 98/45304 (43) International Publication Date: 15 October 1998 (15.10.98)
(21) International Application Number: PCT/NL98/00206 (22) International Filing Date: 9 April 1998 (09.04.98) (30) Priority Data: 97201066.4 10 April 1997 (10.04.97) EP (34) Countries for which the regional or international application was filed: NL et al. (71) Applicant (for all designated States except US): KREATECH BIOTECHNOLOGY B.V. [NL/NL]; Keienbergweg 3, Postbus 12756, NL-1100 AT Amsterdam (NL). (72) Inventors; and (75) Inventors/Applicants (for US only): HOUTHOF, Hendrik, Jan [NL/NL]; Schellingswouderdijk 125, NL-1023 NB Amsterdam (NL). REEDIJK, Jan [NL/NL]; Anthoni Duycklaan 4, NL-2334 CD Leiden (NL). VOLKERS, Herman, H. [NL/NL]; Noordervesting 3, NL-1141 SL Monnickendam (NL). HEETEBRIJ, Robert, Jochem [NL/NL]; Zadelstraat 2, NL-3511 LT Utrecht (NL). (74) Agent: SMULDERS, Th., A., H., J.; Vereenigde Octrooibureaux, Nieuwe Parklaan 97, NL-2587 BN The Hague (NL).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
(54) Title: TRANS-PLATINUM COMPOUND, AND DIAGNOSTIC KIT (I) (57) Abstract The present invention is concerned with a trans-platinum based compound for use in a method for labelling a bio-organic molecule, having formula (I), wherein A represents a reactive moiety for attachment to a label, D represents a reactive moiety for attachment to a bio-organic molecule, and X1 and X2 represent the same or different inert moieties.		

TRANS-PLATINUM COMPOUND, AND DIAGNOSTIC KIT

The invention relates to a trans-platinum based compound, to a diagnostic kit comprising said compound, and to a method for labeling a bio-organic molecule wherein use is made of said compound.

5 Platinum (coordination) compounds have been considered interesting molecules for a very long time. For a review of these compounds and their uses we refer to Reedijk et al. (Structure and Bonding, 67, pp. 53-89, 1987). Especially cis-platinum has received a lot of
10 attention as an anti-tumor drug. This anti-tumor reactivity of platinum compounds originates from their having at least two reactive groups (preferable cis-oriented towards each other), which make it possible to cross-link DNA molecules, thereby inhibiting the
15 replication of these DNA molecules.

 A different use of cis-platinum compounds has been described in European patent application no. 95201197.1. Herein a method is disclosed for linking bio-organic molecules and labels through cis-platinum compounds, of
20 which two coordination sites are occupied by two ends of a stabilizing bridge, such as an ethylene diamine group.

 These known cis-platinum compounds are suitable for linking labels to several kinds of bio-organic molecules. Examples are (poly)-peptides, polynucleic acids and
25 nucleotides.

 The labeling of polynucleic acids, such as DNA molecules, is desirable for applications in fields such as recombinant DNA technology. In many cases, however, one wishes not to label the nucleic acid macromolecule,
30 but to label a certain nucleotide and enzymatically build this into a polynucleic acid. This way, the location of the label on the resulting polynucleic acid can be

influenced, which is not possible when a label is attached to the macromolecule.


It has been found, however, that nucleotides linked to a label by cis-platinum linkers are not satisfactorily
5 built in into DNA molecules by DNA polymerase, if at all. Therefore, in order to use a cis-platinum linker for obtaining a labeled DNA molecule one has to label the macromolecule.

The available alternatives for labeling a
10 nucleotide, which can be incorporated into a polynucleic acid are the more conventional methods of labeling. However, these conventional methods also have a major disadvantage as they are not suitable for labeling any nucleotide. In some cases, for instance when only a few
15 residues of a certain nucleotide are present in a certain polynucleic acid, or when the terminating nucleotide residue of a polynucleic acid is to be labeled, it is desired to be able to label any nucleotide.

An example of such a conventional method has been
20 described by Dale et al., Biochemistry, **14**, (1975), 2447-2457, which method involves direct covalent mercuriation as a labeling technique. Dale et al. report that cytosine and uracil may be mercurated at their C5-position under mild conditions. However, they also report that for
25 adenine, thymine and guanine bases negative results were obtained.

Thus, there is a need for a universal linking system, which is excellent for linking labels and bio-organic molecules, including all different nucleotides,
30 and which also makes it possible to enzymatically build in any nucleotide labeled through said linking system in a polynucleic acid in an efficient manner.

The present invention provides such a universal linking system. It has been found that a trans-platinum
35 based compound, having the formula:



wherein A represents a reactive moiety for attachment to a label, D represents a reactive moiety for attachment to a bio-organic molecule, and X1 and X2
5 represent the same or different inert moieties, meets the above requirements for use in a method for labeling a bio-organic molecule.

Surprisingly, linking labels to several kinds of bio-organic molecules by using the compound of the
10 invention is at least as efficient as the known methods. Further, it has been found that nucleotides linked to a label by the compound of the invention may be built in into polynucleotides very efficiently.

Nucleotides can assume two conformations: the syn-
15 conformation and the anti-conformation. Definitions of these conformations can be found in Saenger, "Principles of Nucleic Acid Structure", Springer-Verlag Inc., 1984. It is believed that for a nucleotide to be satisfactorily built in into e.g. a DNA molecule it has to be in an
20 anti-configuration. In US-A-4,711,955 it has been reported that altering of the normal anti-conformation of a nucleoside must be avoided, because otherwise the nucleotide derivatives are unacceptable as polymerase substrates. Thus nucleotides having a syn-conformation
25 will not be enzymatically incorporated into a polynucleotide in an efficient manner, if at all.

Besides this, it is known from the above-mentioned publication by Saenger that the presence of a bulky group or substituent at the N7 position of a nucleotide, i.e.
30 the position to which a metal coordinates most easily, induces a nucleotide to have a syn-conformation. Thus, it would be expected that a nucleotide having a platinum compound coordinated to its N7 position would be in a syn-orientation. Accordingly, it would be expected to be
35 not possible to incorporate said nucleotide into a polynucleotide with good results.

It is therefore very surprising that nucleotides linked to a label by a trans-platinum based compound in accordance with the invention may be built in into polynucleotides in a highly efficient manner.

5 The moieties X1 and X2 in formula (I) are inert. This means, that these moieties are ligands that remain bonded to the platinum core during the labeling reactions, i.e. they are stable under physiological conditions. They may be different or the same and may be
10 any group that shows no interfering interactions between them and the label, bio-organic molecule or any other compound to be used in the application of the labeling techniques.

In these moieties X1 and X2 resides another
15 advantage of the invention compared to the cis-platinum linkers. In order to obtain a stable cis-platinum compound it is imperative that said compound comprises a stabilizing bridge. For the trans-platinum compounds such a stabilizing bridge is not necessary. Accordingly, the
20 possible variations in structure for the trans-platinum compounds of the invention are more numerous. Thus, according to the invention the skilled person has many instruments at his disposal for obtaining the desired properties of the compound.

25 In a preferred embodiment, X1 and X2 are the same or different non-leaving ligands. This means, that they are groups which have a very small leaving group character. Most preferred examples thereof are NH_3 , NH_2R , NHRR' , and $\text{NRR}'\text{R}''$ groups, wherein R, R' and R'' either represent an
30 alkyl group having from 1 to 6 carbon atoms, as these groups form very stable bonds to the platinum core.

Particularly preferred is the embodiment wherein X1 and X2 are both an $\text{N}(\text{CH}_3)_3$ group. The presence of the methyl groups has a positive effect in that they prevent
35 conformational modifications of the substrate due to for instance hydrogen bonding of a label or bio-organic molecule to the inert moieties.

The reactive moieties A and D are chosen such that they are easily replaced by or substituted with a label, respectively a bio-organic molecule. It is preferred that A and D are chosen from the group of acetate, NO_3^- , HCO_3^- , CO_3^{2-} , SO_3^- , Cl^- , I^- , and other halogens. These groups are good leaving groups and are therefore easily replaced.

In a highly preferred embodiment, a spacer, such as an oligolysine or a polylysine, is used. In accordance with this embodiment, A and/or D in formula (I) comprises a spacer, which spacer comprises a chain having at least four atoms, which chain comprises an electron donating moiety on one end and a reactive moiety on the other end, wherein the chain is attached to the platinum through the electron donating moiety. To the spacer, a label or a bio-organic molecule may be attached by a reaction with the end of the chain which is the most distant end seen from the platinum.

The use of a spacer has the advantage that the distance between the label and the bio-organic molecule can be optimized, so that steric factors cannot be an obstacle to an efficient labeling reaction.

The electron donating moiety of the spacer is preferably an amine group or a thiolate anion, because it has been found that these groups are capable of forming very strong bonds to platinum.

In a further preferred embodiment, the chain of the spacer comprises at least one heteroatom, preferably an oxygen atom, as this positively effects the hydrophilicity of the chain, which is an important aspect in physiological systems.

It is also possible to use a spacer that comprises a platinum atom. In this embodiment, the invention provides a bis-platinum compound for linking labels and bio-organic molecules. The second platinum can either have a cis or a trans orientation.

The spacer comprises preferably no more than 20 carbon atoms in the chain, which is preferably a non-

branched chain, thus causing no steric hindrance. The reasons for this will be clear.

A highly preferred spacer is 1,8-diamino-3,6-dioxaoctane, herein referred to as Dadoo. Dadoo is a very flexible spacer having two primary amine groups and a size that makes it very suitable for use as a spacer according to the invention.

The labels that may be linked to a bio-organic molecule by the compound of the invention are not critical. In principle all labels which can be attached to a bio-organic molecule and are employed to date can be used. These labels may be radioactive labels, enzymes (which need reaction with a substrate to be detected), specific binding pairs components such as avidin, streptavidin or biotin, biocytin, iminobiotin, colloidal dye substances, fluorochromes (rhodamin, etc.) including Cy[®]-colourants, reducing substances (eosin, erythrosin, etc.), (coloured) latex sols, digoxigenin, metals (ruthenium), metal sols or other particulate sols (selenium, carbon and the like), dansyl lysin, Infra Red Dyes, coumarines (amino methyl coumarine), antibodies, protein A, protein G, etc. The invention has most benefits with bulkier labels such as biotin, avidin, streptavidin, and digoxigenin.

Cy[®]-colourants are preferred labels to be linked to bio-organic molecules by the compound of the invention. These labels can very suitably be used for a technique referred to as multi-colour labeling. The reason for this is that different colourants of this kind, while having different colours, are very much alike in chemical structure.

Almost every bio-organic molecule which contains an accessible sulfur or nitrogen atom can be linked to a label by the present compounds. Very suitable compounds to be labeled are proteins, peptides, DNA molecules, RNA molecules, and (oligo)-nucleotides.

The platinum binds very easily to the N7 position of nucleotides. This way DNA or RNA molecules, be it single

stranded or otherwise can be easily detected. It also allows for the production of probes for hybridization techniques wherein unlabeled DNA/RNA molecules hybridize to the labeled probe. The platinum compounds do hardly
5 interfere with the hybridization, if at all.

The platinum compounds of the invention are also very suitable for attaching bio-organic molecules to solid surfaces such as nitrocellulose, nylon filters or microtiter plates.


10 Nucleotides modified by using a linker according to the invention and oligo- and polynucleotides into which the nucleotides have been built in, or oligo- and polynucleotides that have been directly modified using these novel platinum compounds may be used as probes in
15 biomedical research, clinical diagnostics and recombinant DNA technology.

The wide variety of utilities are based upon the ability of the platinum compounds to form stable complexes with polypeptides which in turn can be detected
20 either by means of detectable moieties which are attached to or which interact with the polypeptide. Some uses include detecting and identifying nucleic acid containing etiological agents, e.g. bacteria and viruses; screening bacteria for antibiotic resistance; screening animals for
25 genetic disorders in relation to pharmaceutical effects; diagnosing genetic disorders, e.g. trisomy 21, sickle cell anemia: chromosomal karyotyping; and identifying tumor cells.

The invention also encompasses a diagnostic kit for
30 detecting, determining and/or localizing biological substances of interest, comprising the platinum compound of the invention, optionally together with other suitable means for detection. The platinum compound in the kit may comprise one or more of a spacer, a label and a bio-
35 organic molecule. Of course, the invention also encompasses a kit, wherein these components are present separately, i.e. in unbound form.

In a different embodiment, the invention is directed to a method for labeling bio-organic molecules wherein use is made of the compounds disclosed herein-above.

In a method for labeling a bio-organic molecule in accordance with this embodiment, a reactive moiety of a
5 trans-platinum based compound, having the formula:



wherein A, D, X1 and X2 have the above meanings, is reacted with a label; and the other reactive moiety of
10 said trans-platinum based compound is reacted with a bio-organic molecule or vice versa.

In a preferred method for labeling a bio-organic molecule, the label and/or bio-organic molecule are connected to the platinum through a spacer, which spacer
15 comprises a chain having at least four atoms, which chain comprises an electron donating moiety on one end and a reactive moiety on the other end, wherein the chain is attached to the platinum through the electron donating moiety. This spacer may be any of the spacers disclosed
20 herein-above.

Of course, the spacer(s), the label, the bio-organic molecule and the trans-platinum may be attached to each other in any order. For instance, the spacer(s) may first be attached to the platinum followed by reacting the
25 obtained compound with label and bio-organic molecule, but it is also possible first to attach the spacer(s) to the label and/or bio-organic molecule before the reaction with the platinum compound.

In a highly preferred embodiment, the invention
30 provides a method as described above wherein the bio-organic molecule is a nucleotide.

The invention will be elucidated by the following, non-restrictive examples.

EXAMPLE 1: Preparation of trans-diamminedichloro-platinum(II) (trans- $[\text{Pt}(\text{NH}_3)_2\text{Cl}_2]$)

1 g of potassium tetrachloroplatinate was dissolved in 20
5 ml Milli-Q and 0.6 ml of concentrated HCl was added. The
mixture was heated until boiling and 2.5 ml of a 25%
solution of NH_4OH was added dropwise with swirling. The
resulting solution was evaporated to about 10 ml. To the
10 resulting pale yellow residue, 100 ml of 6N HCl was
added, and the mixture was evaporated as above to a
volume of 15-20 ml. The trans-diamminedichloro-
platinum(II) was deposited as a fine yellow powder. After
cooling in ice, the mixture was filtered, washed with
three 10 ml portions of ice-cold water, ethanol and
15 ether, and air-dried. The trans-diamminedichloro-
platinum(II) was crystallized from a minimum amount of
boiling 0.1 N HCl. The product was characterized by
infrared spectroscopy.

20

EXAMPLE 2: Preparation of trans-platinum diammine-DigDadoo chloride (trans- $[\text{Pt}(\text{NH}_3)_2(\text{DigDadoo-NH}_2)\text{Cl}](\text{NO}_3)$)

30 mg of trans- $[\text{Pt}(\text{NH}_3)_2\text{Cl}_2]$ (0.1 mmol) was suspended in
25 25 ml of N,N'-dimethylformamide and 17 mg (0.1 mmol) of
 AgNO_3 was added to the mixture which was reacted for 16
hours at room temperature shielded from the light. The
precipitated AgCl was filtered off and 3.4 ml was taken
from the filtrate (3.9 mg (0.014 mmol) of the trans-
30 $[\text{Pt}(\text{NH}_3)_2(\text{Cl})(\text{NO}_3)]$) and incubated with 8 mg of DigDadoo
(digoxigenin-1,8-diamino-3,6-dioxaoctane, 0.014 mmol,
purchased from Boehringer Mannheim) for 48 hours at 40°C,
shielded from the light. The N,N'-dimethylformamide was
removed *in vacuo* and the remainder was stirred in CH_2Cl_2
35 whereupon a yellowish powder precipitated. The
precipitated powder was filtered off, washed with CH_2Cl_2
and dried. The product was characterized by high
resolution $^1\text{H-NMR}$.

EXAMPLE 3: Preparation of trans-platinum diammine-DigDadoo-2'-deoxyGuanosine-5'-triphosphate (trans-[Pt(NH₃)₂(DigDadoo-NH₂)(dGTP-N7)]²⁺)

5 12 mg trans-[Pt(NH₃)₂(DigDadoo-NH₂)Cl](NO₃) (0.014 mmol) was dissolved in 2 ml of Milli-Q and 3 mg (0.006 mmol) of 2'-deoxyguanosine-5'-triphosphate was added. The pH of the mixture was adjusted to ~6 with 0.1 N NaOH and the reaction was performed for 24 hours at 50°C shielded from
10 the light. The mixture was lyophilized and the residue was applied to a preparative anion exchange chromatography column (Q-Sepharose 26/10, Pharmacia BioTech) and a gradient starting with 100% Milli-Q to 100% NH₄HCO₃ was applied. The appropriate fraction was collected,
15 concentrated and lyophilized to remove NH₄HCO₃. The product was analyzed by high resolution ¹H and ³¹P-NMR and by analytical FPLC-methods, anion exchange (MonoQ 5/5, Pharmacia BioTech) and reversed phase (PepRPC 5/5, Pharmacia BioTech). The product was converted into its
20 tetralithium salt by passing it over a Dowex cation exchange column.

EXAMPLE 4: Preparation of trans-platinum diammine-Cy5[®]-
25 chloride (trans-[Pt(NH₃)₂(Cy5[®]-NH₂)Cl](NO₃))

7.9 mg Cy5 Monofunctional reactive dye (0.01 mmol) was dissolved in 5 ml of dry N,N'-dimethylformamide and reacted with an excess of ethylenediamine for 4 hours at
30 room temperature in an inert atmosphere (argon). The N,N'-dimethylformamide and excess ethylenediamine were removed in vacuo. The residue was redissolved in 10 ml of N,N'-dimethylformamide and 3 mg (0.009 mmol) trans-[Pt(NH₃)₂(Cl)(NO₃)] was added. The mixture was stirred
35 for 16 hours at 50°C, shielded from the light. The N,N'-dimethylformamide was removed in vacuo and the residue was stirred in CH₂Cl₂, whereupon the product precipitated.

The precipitate was filtered off and dried in vacuo. The product was characterized by high resolution ^1H -NMR.

5 EXAMPLE 5: Enzymatic incorporation of trans-
[Pt(NH₃)₂(DigDadoo-NH₂)(dGTP)] into pBR 322 DNA

The enzymatic incorporation was performed by incubating the following mixture overnight at 37°C:

10

Denaturated DNA (1 µg)	8.0 µl
dTTP (5.0 mM)	0.8 µl
dCTP (5.0 mM)	0.8 µl
dATP (5.0 mM)	0.8 µl
15 dGTP (1.0 mM)	2.6 µl
trans-[Pt(NH ₃) ₂ (DigDadoo-NH ₂)(dGTP)] (1 mM)	1.4 µl
High Prime Mixture*	4.0 µl
Sterile demineralized water	<u>1.6 µl</u>
Total	20 µl

20

* The High Prime Mixture contained Klenow polymerase and hexanucleotides in an optimized buffer.

The incubation was stopped by adding 2 µl of 0.2 M EDTA
25 and the volume of the mixture was adjusted to 40 µl by adding sterile demineralized water. A filter hybridization was performed with a probe concentration of 25 ng/ml, using dentarued homologous DNA dotted on filterstrips with the following dilution range: 100pg, 30
30 pg, 3 pg, 1 pg, 0.3 pg, 0.1 pg, 0.03 pg, 0.01 pg and 0 pg. Detection was carried out with anti-Dig-AP (Fab fragments from an anti-digoxigening antibody from sheep conjugated with alkaline phosphatase (AP)) (1:10,000) and CDP-Star (chemiluminescent substrate). As a result 0.3 pg
35 hybridization was observed.

12

EXAMPLE 6: trans-[Pt(NH₃)₂(DigDadoo-NH₂)(dGTP)] Oligo
tailing using Terminal Transferase

The tailing reaction was performed by incubating the
5 following mixture for 15 minutes at 37°C:

Reaction buffer:	1M potassium cacodylate,	4 µl
	0.125 M Tris-HCl, 1.25 mg/m;	
	Bovine serum albumine pH 6.6	
10	(25°C)	
	CoCl ₂ solution (25 mM)	4 µl
	oligo (20 pmol/µl)	5 µl
	trans-[Pt(NH ₃) ₂ (DigDadoo-NH ₂)(dGTP)]/dATP*	1 µl
	Terminal Transferase 50U/µl	1 µl
15	Sterile demineralized water	5 µl
	Total	20 µl

* Mixture of 9 µl of trans-[Pt(NH₃)₂(DigDadoo-
NH₂)(dGTP)] (1 mM) and 1 µl dATP (100 mM)

20

The incubation was stopped by adding 2 µl of a mixture of
1 µl glycogen and 200 µl 0.2 M EDTA having a pH of 8.0.
Subsequently, a purification was performed by
precipitation from ethanol and a hybridization analogous
25 to the filter hybridization described in Example 5 was
performed. As a result 10 pg hybridization was observed.

EXAMPLE 7: Normal DNA labeling using trans-
30 [Pt(NH₃)₂(DigDadoo-NH₂)Cl](NO₃)

From a standard QC procedure, testing a range of
different ratios, the following protocol was performed.
A mixture of 2 µl DNA (2 µg), 2 µl trans-
35 [Pt(NH₃)₂(DigDadoo-NH₂)Cl](NO₃) (0.1 mg/ml) and 16 µl
sterile demineralized water was incubated for 30 minutes
at 85°C. The incubation was stopped by adding 5 µl 1%

sodium diethyldithiocarbamic acid. In a filter hybridization, which was performed analogous to the hybridization described in Example 5, 3 pg hybridization was observed.

5

EXAMPLE 8: Ratio labeling of DNA by mixing painting probes that have been labeled independently

10 Identical painting probes were labeled in independent reactions using Fluorescein (Flu) and Rhodamine (Rho). After labeling, independent probes were mixed using a ratio of 1:9, 1:5, 1:1, 5:1 and 9:1, respectively, and tested spot blot and by filter hybridization. Detection
15 was performed by alkaline phosphatase conjugated antibodies raised against Flu and Rho, respectively. After detection, ratios of labeled painting probes appeared to be identical when compared with the ratio of input marker molecule.

20

EXAMPLE 9: Ratio labeling of DNA using a defined mixed input of trans-[Pt(NH₃)₂(Dadoo-NH₂)Cl]NO₃ marker complexes in one reaction


25

Probes were labeled using similar ratios as described in Example 8, with the exception that the probe DNA was incubated with a mixture of trans-[Pt(NH₃)₂(Dadoo-NH₂)Cl]NO₃ marker (Flu and Rho) complexes in one reaction
30 tube. After detection by conjugated antibodies, it appeared that distribution of label occurred equimolar and that different marker molecules did not interfere with each other, nor did they affect each other negatively. All results were confirmed by microscopial
35 analysis

P22321PC00

New Claims

1. A method for labeling a bio-organic molecule, wherein a reactive moiety of a trans-platinum based compound, having the formula:



- 5 wherein A and D represent the same or different reactive moieties, and X1 and X2 represent the same or different inert moieties, is reacted with a label, and wherein the other reactive moiety of said trans-platinum based compound is reacted with a bio-organic molecule, in any order.
- 10 2. A method according to claim 1, wherein X1 and X2 are the same or different non-leaving ligands.
3. A method according to any of the preceding claims, wherein X1 and X2 are chosen from the group of NH_3 , NH_2R , $NHRR'$, $NRR'R''$ groups, wherein R, R' and R'' represent an alkyl
- 15 group having from 1 to 6 carbon atoms.
4. A method according to any of the preceding claims, wherein X1 and X2 both represent an $N(CH_3)_3$ group.
5. A method according to any of the preceding claims, wherein A and/or D are chosen from the group of acetate,
- 20 NO_3^- , HCO_3^- , CO_3^{2-} , SO_3^- , Cl^- , I^- , and other halogens.
6. A method according to any of the preceding claims, wherein the label and/or the bio-organic molecule are connected to the platinum through a spacer, which spacer comprises a chain having at least four atoms, which chain
- 25 comprises an electron donating moiety on one end and a reactive moiety on the other end, wherein one or both of the reactive moieties of the trans-platinum based compound are reacted with the electron donating moiety of said spacer(s)

and wherein the reactive moiety of said spacer(s) are reacted with the label and/or bio-organic molecule, in any order.

7. A method according to claim 6, wherein the electron donating moiety is an amine group or a thiolate anion.

5 8. A method according to claim 6 or 7, wherein the chain further comprises at least one heteroatom.

9. A method according to any of the claims 6-8, wherein the spacer comprises no more than 20 carbon atoms in the chain and wherein the chain is essentially non-branched.


10 10. A method according to claim 9, wherein A and/or D is 1,8-diamino-3,6-dioxaoctane.

11. A method according to any of the preceding claims, wherein the bio-organic molecule is chosen from the group of proteins, peptides, DNA molecules, RNA molecules, and
15 (oligo)-nucleotides.

12. A method according to claim 11, wherein the bio-organic molecule is a nucleotide.

13. A method according to any of the preceding claims, wherein the label is chosen from the group biotin, avidin,
20 streptavidin, digoxigenin, and fluorochromes including Cy[®]-colourants.

14. A trans-platinum based compound for use in a method for labeling a bio-organic molecule according to any of the preceding claims, having the formula:



25

wherein A represents a reactive moiety for attachment to a label, D represents a reactive moiety for attachment to a bio-organic molecule, and X1 and X2 represent the same or different inert moieties.

30 15. A compound according to claim 14, having a label attached thereto.

16. A compound according to claim 14 or 15 having a bio-organic molecule attached thereto.

17. A compound according to claim 16, wherein the bio-organic molecule is a nucleotide.

18. A diagnostic kit for detecting, determining and/or localizing biological substances of interest, comprising a
5 compound according to any of the claims 14-16, optionally together with other suitable means for detection.

AMENDED SHEET