

(19) AUSTRALIAN PATENT OFFICE

(54) Title
4-phenylpiperidine derivatives as renin inhibitors

(51)⁶ International Patent Classification(s)
C07D 211/42 8BMEP **C07D**
(2006.01) 401/10
C07D 401/10 20060101ALI2005100
(2006.01) 8BMEP **C07D**
C07D 405/04 405/04
(2006.01) 20060101ALI2005100
C07D 409/10 8BMEP **C07D**
(2006.01) 409/10
C07D 211/42 20060101ALI2005100
20060101AFI2005100 8BMEP
PCT/EP2004/013410

(21) Application No: 2004293177

(22) Application Date: 2004 .11 .25

(87) WIPO No: W005/051911

(30) Priority Data

(31) Number	(32) Date	(33) Country
60/525,375	2003 .11 .26	US

(43) Publication Date : 2005 .06 .09

(71) Applicant(s)
Novartis AG

(72) Inventor(s)
Sellner, Holger; Ostermann, Nils; Maibaum, Jurgen Klaus; Ehrhardt, Claus; Sedrani, Richard ; Breitenstein, Werner

(74) Agent/Attorney
Davies Collison Cave, Level 15 1 Nicholson Street, Melbourne, VIC, 3000

(56) Related Art
WO 1997/009311

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
9 June 2005 (09.06.2005)

PCT

(10) International Publication Number
WO 2005/051911 A1

(51) International Patent Classification⁷: **C07D 211/42**,
405/04, 401/10, 409/10, A61K 31/4418, 31/443, 31/4436,
31/4439, 31/444

Grossmannstrasse 17, 79541 Lörrach (DE). **OSTER-
MANN, Nils** [DE/DE]; Mattenleestrasse 28, 79650
Schopfheim (DE).

(21) International Application Number:
PCT/EP2004/013410

(74) Agent: **GRUBB, Philip**; Novartis AG, Corporate Intellec-
tual Property, CH-4002 Basel (CH).

(22) International Filing Date:
25 November 2004 (25.11.2004)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/525,375 26 November 2003 (26.11.2003) US

(71) Applicant (for all designated States except AT, US): **NO-
VARTIS AG** [CH/CH]; Lichtstrasse 35, CH-4056 Basel
(CH).

(71) Applicant (for AT only): **NOVARTIS PHARMA GMBH**
[AT/AT]; Brunner Strasse 59, A-1230 Vienna (AT).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **SEDRANI, Richard**
[LU/CH]; Herrengrabenweg 15, CH-4054 Basel (CH).
MAIBAUM, Jürgen, Klaus [DE/DE]; Lütemannsweg
14/1, 79576 Weil-Haltingen (DE). **BREITENSTEIN,**
Werner [CH/CH]; St. Galler-Ring 179, CH-4054 Basel
(CH). **SELLNER, Holger** [DE/CH]; Keltenstr. 37,
CH-4106 Therwil (CH). **EHRHARDT, Claus** [DE/DE];

(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AI, AM,
AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CI, CN,
CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI,
GB, GD, GF, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE,
KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD,
MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG,
PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SI, SY, TJ, TM,
TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM,
ZW.


(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM,
ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI,
FR, GB, GR, HU, IE, IS, IT, LU, MC, NL, PL, PT, RO, SE,
SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: 4-PHENYLPIPERIDINE DERIVATIVES AS RENIN INHIBITORS




(57) Abstract: Compounds of the present invention having the formula (I) exhibit inhibitory activity on the natural enzyme renin. Thus, compounds of formula (I) may be employed for the treatment of hypertension, atherosclerosis, unstable coronary syndrome, congestive heart failure, cardiac hypertrophy, cardiac fibrosis, cardiomyopathy postinfarction, unstable coronary syndrome, diastolic dysfunction, chronic kidney disease, hepatic fibrosis, complications resulting from diabetes, such as nephropathy, vasculopathy and neuropathy, diseases of the coronary vessels, restenosis following angioplasty, raised intra-ocular pressure, glaucoma, abnormal vascular growth, hyperaldosteronism, cognitive impairment, alzheimers, dementia, anxiety states and cognitive disorders.

WO 2005/051911 A1

4-PHENYLPYPERIDINE DERIVATIVES AS RENIN INHIBITORS

The present invention provides compounds of the formula



wherein

R_1 is $-\text{CH}_2\text{-X}$, $-\text{O-X}$ or $-\text{S(O)}_{0-2}\text{-X}$; or

R_1 is $-\text{NR}_8\text{-X}$, $-\text{NR}_8\text{C(O)-X}$ or $-\text{NR}_8\text{S(O)}_2\text{-X}$ in which

R_8 is hydrogen or lower alkyl; and

X is $-(\text{CH}_2)_m\text{-(CR}_9\text{R}_{10})_p\text{-(CH}_2)_n\text{-Z-(CH}_2)_q\text{-W}$ in which

m , n and q are independently zero or an integer from 1 to 5;

p is zero or 1;

R_9 and R_{10} are independently hydrogen, hydroxy, halogen, lower alkyl, lower alkoxy or cycloalkyl; or

R_9 and R_{10} combined are alkylene which together with the carbon atom to which they are attached form a 3- to 6-membered ring;

Z is a bond; or

Z is O , S(O)_{0-2} , or $-\text{NR}_{11}-$ in which

R_{11} is hydrogen or lower alkyl, provided that R_1 is $-\text{CH}_2\text{-X}$ when m , n and p are all zero;

W is aryl or heterocyclyl;

R_2 is hydrogen, halogen, cyano, hydroxy or lower alkoxy;

L is a bond; or

L is $-(\text{CH}_2)_s\text{-O-(CH}_2)_v\text{-}$ in which

s and v are independently zero or an integer from 1 to 3; or

- 2 -

L is $-C(O)-$, $-C(O)O-$, $-OC(O)-$, $-OC(O)NR_{12}-$, $-NR_{12}-$, $-NR_{13}C(O)-$, $-NR_{13}C(O)O-$ or $-NR_{13}C(O)NR_{12}-$ in which

R_{12} and R_{13} are independently hydrogen or lower alkyl;

R_3 is hydrogen, hydroxy, halogen or cyano provided that L is a bond; or

R_3 is optionally substituted lower alkyl, aralkyl, heteroaralkyl, aryl or heterocyclyl; or

R_3 and R_{12} combined are alkylene which together with the nitrogen atom to which they are attached form a 5- to 6-membered ring;

R_4 is hydrogen, optionally substituted lower alkyl or aryl;

R_5 and R_6 are independently hydrogen, halogen, hydroxy, trifluoromethyl, optionally substituted lower alkyl, lower alkoxy or cycloalkyl; or

R_5 and R_6 combined together with the carbon atoms to which they are attached form a fused 5- to 6-membered aromatic or heteroaromatic ring provided that R_5 and R_6 are attached to carbon atoms adjacent to each other; or

R_5 and R_6 combined are alkylene which together with the carbon atoms to which they are attached form a fused 5- to 7-membered ring provided that R_5 and R_6 are attached to carbon atoms adjacent to each other; or

C- R_6 and C- R_5 may be replaced with nitrogen;

R_7 is hydrogen, halogen, hydroxy, trifluoromethyl, optionally substituted lower alkyl, lower alkoxy, cycloalkyl, alkanoyl, alkoxyalkoxy, alkanoyloxy, amino, alkylamino, dialkylamino, acylamino, carbamoyl, thiol, alkylthio, alkylthiono, sulfonyl, sulfonamido, sulfamoyl, nitro, cyano, carboxy, alkoxycarbonyl, aryl, alkenyl, alkynyl, aralkoxy, heterocyclyl including indolyl, imidazolyl, furyl, thienyl, thiazolyl, pyrrolidyl, pyridyl, pyrimidyl, piperidyl, morpholinyl and tetrazolyl; or

R_7 and R_8 combined are O, $S(O)_{0-2}$, $-NR_{14}-$, $-(CH_2)_{1-2}-$, $-O-CH_2-$, $-CH_2-O-$, $-S(O)_{0-2}-CH_2-$, $-CH_2-S(O)_{0-2}-$, $-NR_{14}-CH_2-$, $-CH_2-NR_{14}-$, $-S(O)_{0-2}-NR_{14}-$ or $-NR_{14}-S(O)_{0-2}-$ in which

R_{14} is hydrogen or lower alkyl, provided R_8 is located at the 2' position; or

C- R_7 may be replaced with nitrogen;

Y is $-(CH_2)_r-$, $-O-(CH_2)_r-$, $-(CH_2)_r-O-$, $-S_{0-2}-(CH_2)_r-$ or $-(CH_2)_r-S_{0-2}-$ in which

r is zero or an integer from 1 to 3;

Q combined with the atoms to which it is attached form a 5- to 6-membered monocyclic aromatic or heteroaromatic ring; or

Q combined with the atoms to which it is attached form a 7- to 12-membered bicyclic aromatic or heterocyclic ring;

or a pharmaceutically acceptable salt thereof.

The compounds of the present invention exhibit inhibitory activity on the natural enzyme renin. Thus, compounds of formula (I) may be employed for the treatment of hypertension, atherosclerosis, unstable coronary syndrome, congestive heart failure, cardiac hypertrophy, cardiac fibrosis, cardiomyopathy postinfarction, unstable coronary syndrome, diastolic dysfunction, chronic kidney disease, hepatic fibrosis, complications resulting from diabetes, such as nephropathy, vasculopathy and neuropathy, diseases of the coronary vessels, restenosis following angioplasty, raised intra-ocular pressure, glaucoma, abnormal vascular growth, hyperaldosteronism, cognitive impairment, alzheimers, dementia, anxiety states and cognitive disorders.

Listed below are definitions of various terms used to describe the compounds of the present invention. These definitions apply to the terms as they are used throughout the specification unless they are otherwise limited in specific instances either individually or as part of a larger group.

The term "optionally substituted alkyl" refers to unsubstituted or substituted straight- or branched-chain hydrocarbon groups having 1-20 carbon atoms, preferably 1-7 carbon atoms. Exemplary unsubstituted alkyl groups include methyl, ethyl, propyl, isopropyl, *n*-butyl, *t*-butyl, isobutyl, pentyl, hexyl, isohexyl, heptyl, 4,4-dimethylpentyl, octyl and the like. Substituted alkyl groups include, but are not limited to, alkyl groups substituted by one or more of the following groups: halo, hydroxy, cycloalkyl, alkanoyl, alkoxy, alkyloxyalkoxy, alkanoyloxy, amino, alkylamino, dialkylamino, acylamino, carbamoyl, thiol, alkylthio, alkylthiono, sulfonyl, sulfonamido, sulfamoyl, nitro, cyano, carboxy, alkoxycarbonyl, aryl, alkenyl, alkynyl, aralkoxy, guanidino, heterocyclyl including indolyl, imidazolyl, furyl, thienyl, thiazolyl, pyrrolidyl, pyridyl, pyrimidyl, piperidyl, morpholinyl and the like.

The term "lower alkyl" refers to those optionally substituted alkyl groups as described above having 1-7, preferably 1-4 carbon atoms.

The term "halogen" or "halo" refers to fluorine, chlorine, bromine and iodine.

The term "alkenyl" refers to any of the above alkyl groups having at least two carbon atoms and further containing a carbon to carbon double bond at the point of attachment. Groups having 2-4 carbon atoms are preferred.

The term "alkynyl" refers to any of the above alkyl groups having at least two carbon atoms and further containing a carbon to carbon triple bond at the point of attachment. Groups having 2-4 carbon atoms are preferred.

The term "alkylene" refers to a straight-chain bridge of 2-5 carbon atoms connected by single bonds, e.g., $-(CH_2)_x-$, wherein x is 2 to 5, which may be interrupted with one or more heteroatoms selected from O, $S(O)_{0.2}$ or $-NR-$, wherein R may be hydrogen, alkyl, cycloalkyl, aryl, acyl, carbamoyl, sulfonyl, sulfamoyl, alkoxycarbonyl, aryloxy carbonyl or aralkoxycarbonyl, or the alkylene may be substituted with one or more substituents selected from alkyl, cycloalkyl, oxo, halogen, hydroxy, carboxy, alkoxy, alkoxycarbonyl and the like.

The term "cycloalkyl" refers to optionally substituted monocyclic, bicyclic or tricyclic hydrocarbon groups of 3-12 carbon atoms, each of which may be substituted by one or more substituents, such as alkyl, halo, oxo, hydroxy, alkoxy, alkanoyl, acylamino, carbamoyl, alkylamino, dialkylamino, thiol, alkylthio, nitro, cyano, carboxy, alkoxycarbonyl, sulfonyl, sulfonamido, sulfamoyl, heterocyclyl and the like.

Exemplary monocyclic hydrocarbon groups include, but are not limited to, cyclopropyl, cyclobutyl, cyclopentyl, cyclopentenyl, cyclohexyl and cyclohexenyl and the like.

Exemplary bicyclic hydrocarbon groups include bornyl, indyl, hexahydroindyl, tetrahydronaphthyl, decahydronaphthyl, bicyclo[2.1.1]hexyl, bicyclo[2.2.1]heptyl, bicyclo[2.2.1]heptenyl, 6,6-dimethylbicyclo[3.1.1]heptyl, 2,6,6-trimethylbicyclo[3.1.1]heptyl, bicyclo[2.2.2]octyl and the like.

Exemplary tricyclic hydrocarbon groups include adamantyl and the like.

The term "alkoxy" refers to alkyl-O-.

The term "alkanoyl" refers to alkyl-C(O)-.

The term "alkanoyloxy" refers to alkyl-C(O)-O-.

The terms "alkylamino" and "dialkylamino" refer to alkyl-NH- and (alkyl)₂N-, respectively.

The term "alkanoylamino" refers to alkyl-C(O)-NH-.

The term "alkylthio" refers to alkyl-S-.

The term "alkylaminothiocarbonyl" refers to alkyl-NHC(S)-.

The term "trialkylsilyl" refers to (alkyl)₃Si-.

The term "trialkylsilyloxy" refers to (alkyl)₃SiO-.

The term "alkylthiono" refers to alkyl-S(O)-.

The term "alkylsulfonyl" refers to alkyl-S(O)₂-.

The term "alkoxycarbonyl" refers to alkyl-O-C(O)-.

The term "alkoxycarbonyloxy" refers to alkyl-O-C(O)O-.

The term "carbamoyl" refers to H₂NC(O)-, alkyl-NHC(O)-, (alkyl)₂NC(O)-, aryl-NHC(O)-, alkyl(aryl)-NC(O)-, heteroaryl-NHC(O)-, alkyl(heteroaryl)-NC(O)-, aralkyl-NHC(O)-, alkyl(aralkyl)-NC(O)- and the like.

The term "sulfamoyl" refers to H₂NS(O)₂-, alkyl-NHS(O)₂-, (alkyl)₂NS(O)₂-, aryl-NHS(O)₂-, alkyl(aryl)-NS(O)₂-, (aryl)₂NS(O)₂-, heteroaryl-NHS(O)₂-, aralkyl-NHS(O)₂-, heteroaralkyl-NHS(O)₂- and the like.

The term "sulfonamido" refers to alkyl-S(O)₂-NH-, aryl-S(O)₂-NH-, aralkyl-S(O)₂-NH-, heteroaryl-S(O)₂-NH-, heteroaralkyl-S(O)₂-NH-, alkyl-S(O)₂-N(alkyl)-, aryl-S(O)₂-N(alkyl)-, aralkyl-S(O)₂-N(alkyl)-, heteroaryl-S(O)₂-N(alkyl)-, heteroaralkyl-S(O)₂-N(alkyl)- and the like.

The term "sulfonyl" refers to alkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, aralkylsulfonyl, heteroaralkylsulfonyl and the like.

The term "optionally substituted amino" refers to a primary or secondary amino group which may optionally be substituted by a substituent, such as acyl, sulfonyl, alkoxycarbonyl, cycloalkoxycarbonyl, aryloxycarbonyl, heteroaryloxycarbonyl, aralkoxycarbonyl, heteroaralkoxycarbonyl, carbamoyl and the like.

The term "aryl" or "aromatic ring" refers to monocyclic or bicyclic aromatic hydrocarbon groups having 6-12 carbon atoms in the ring portion, such as phenyl, biphenyl, naphthyl and tetrahydronaphthyl, each of which may optionally be substituted by 1-4 substituents, such as alkyl, trifluoromethyl, cycloalkyl, halo, hydroxy, alkoxy, acyl, alkanoyloxy, aryloxy, optionally substituted amino, thiol, alkylthio, arylthio, nitro, cyano, carboxy, alkoxy carbonyl, carbamoyl, alkylthiono, sulfonyl, sulfonamido, heterocyclyl and the like.

The term "monocyclic aryl" refers to optionally substituted phenyl as described under aryl.

The term "aralkyl" refers to an aryl group bonded directly through an alkyl group, such as benzyl.

The term "aralkanoyl" refers to aralkyl-C(O)-.

The term "aralkylthio" refers to aralkyl-S-.

The term "aralkoxy" refers to an aryl group bonded directly through an alkoxy group.

The term "arylsulfonyl" refers to aryl-S(O)₂-.

The term "arylthio" refers to aryl-S-.

The term "aroyl" refers to aryl-C(O)-.

The term "aroyloxy" refers to aryl-C(O)-O-.

The term "aroylamino" refers to aryl-C(O)-NH-.

The term "aryloxycarbonyl" refers to aryl-O-C(O)-.

The term "heterocyclyl", "heterocyclo" or "heterocyclic ring" refers to an optionally substituted, fully saturated or unsaturated, aromatic or nonaromatic cyclic group, e.g., which is a 4- to 7-membered monocyclic, 7- to 12-membered bicyclic or 10- to 15-membered tricyclic ring system, which has at least one heteroatom in at least one carbon atom-containing ring. Each ring of the heterocyclic group containing a heteroatom may have 1, 2 or 3 heteroatoms selected from nitrogen atoms, oxygen atoms and sulfur atoms, where the nitrogen and sulfur heteroatoms may also optionally be oxidized. The heterocyclic group may be attached at a heteroatom or a carbon atom.

Exemplary monocyclic heterocyclic groups include pyrrolidinyl, pyrrolyl, pyrazolyl, oxetanyl, pyrazolinyl, imidazolyl, imidazolyl, imidazolidinyl, triazolyl, oxazolyl, oxazolidinyl, isoxazolyl, isoxazolyl, thiazolyl, thiadiazolyl, thiazolidinyl, isothiazolyl, isothiazolidinyl, furyl, tetrahydrofuryl, thienyl, oxadiazolyl, piperidinyl, piperazinyl, 2-oxopiperazinyl, 2-oxopiperidinyl, 2-oxopyrrolidinyl, 2-oxoazepinyl, azepinyl, 4-piperidonyl, pyridyl, pyrazinyl, pyrimidinyl, pyridazinyl, tetrahydropyranyl, morpholinyl, thiamorpholinyl, thiamorpholinyl sulfoxide, thiamorpholinyl sulfone, 1,3-dioxolane and tetrahydro-1,1-dioxothieryl and the like.

Exemplary bicyclic heterocyclic groups include indolyl, dihydroidolyl, benzothiazolyl, benzoxaziny, benzoxazolyl, benzothienyl, benzothiazinyl, quinuclidinyl, quinolinyl, tetrahydroquinolinyl, decahydroquinolinyl, isoquinolinyl, tetrahydroisoquinolinyl, decahydroisoquinolinyl, benzimidazolyl, benzopyranyl, indoliziny, benzofuryl, chromonyl, coumarinyl, benzopyranyl, cinnolyl, quinoxaliny, indazolyl, pyrrolopyridyl, furopyridinyl (such as furo[2,3-c]pyridinyl, furo[3,2-b]-pyridinyl or furo[2,3-b]pyridinyl), dihydroisoindolyl, 1,3-dioxo-1,3-dihydroisoindol-2-yl, dihydroquinazolinyl (such as 3,4-dihydro-4-oxo-quinazolinyl), phthalazinyl and the like.

Exemplary tricyclic heterocyclic groups include carbazolyl, dibenzoazepinyl, dithienoazepinyl, benzindolyl, phenanthrolinyl, acridinyl, phenanthridinyl, phenoxazinyl, phenothiazinyl, xanthenyl, carbolinyl and the like.

The term "heterocyclyl", "heterocyclo" or "heterocyclic ring" includes substituted heterocyclic groups. Substituted heterocyclic groups refer to heterocyclic groups substituted with 1, 2 or 3 substituents selected from the group consisting of the following:

- (a) alkyl;
- (b) hydroxy (or protected hydroxy);
- (c) halo;
- (d) oxo, i.e., =O;
- (e) optionally substituted amino, alkylamino or dialkylamino;
- (f) alkoxy;
- (g) cycloalkyl;
- (h) carboxy;
- (i) heterocycloxy;
- (j) alkoxycarbonyl, such as unsubstituted lower alkoxycarbonyl;
- (k) mercapto;

- (l) nitro;
- (m) cyano;
- (n) sulfamoyl or sulfonamido;
- (o) aryl;
- (p) alkanoyloxy;
- (q) aroyloxy;
- (r) arylthio;
- (s) aryloxy;
- (t) alkylthio;
- (u) formyl;
- (v) carbamoyl; and
- (w) aralkyl.

The term "heterocyclooxy" denotes a heterocyclic group bonded through an oxygen bridge.

The term "heteroaryl" or "heteroaromatic ring" refers to an aromatic heterocycle, e.g., monocyclic or bicyclic aryl, such as pyrrolyl, pyrazolyl, imidazolyl, triazolyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, furyl, thienyl, pyridyl, pyrazinyl, pyrimidinyl, pyridazinyl, indolyl, benzothiazolyl, benzoxazolyl, benzothieryl, quinoliny, isoquinoliny, benzimidazolyl, benzofuryl and the like, optionally substituted by, e.g., lower alkyl, lower alkoxy or halo.

The term "heteroarylsulfonyl" refers to heteroaryl-S(O)₂-.

The term "heteroaroyle" refers to heteroaryl-C(O)-.

The term "heteroaroylelamino" refers to heteroaryl-C(O)NH-.

The term "heteroaralkyl" refers to a heteroaryl group bonded through an alkyl group.

The term "heteroaralkanoyl" refers to heteroaralkyl-C(O)-.

The term "heteroaralkanoylamino" refers to heteroaralkyl-C(O)NH-.

The term "acyl" refers to alkanoyl, aroyl, heteroaryl, aralkanoyl, heteroaralkanoyl and the like.

The term "acylamino" refers to alkanoylamino, aroylamino, heteroaroylelamino, aralkanoylamino, heteroaralkanoylamino and the like.

Pharmaceutically acceptable salts of any compound of the present invention refer to salts formed with acids, namely acid addition salts, such as of mineral acids, organic carboxylic and organic sulfonic acids, e.g., hydrochloric acid, maleic acid and methanesulfonic acid.

Similarly salts formed with bases, namely cationic salts, such as alkali and alkaline earth metal salts, such as sodium, lithium, potassium, calcium, magnesium, as well as ammonium salts, such as ammonium, trimethylammonium, diethylammonium, and tris(hydroxymethyl)methylammonium salts and salts with amino acids, are possible provided an acidic group constitutes part of the structure.

As described herein above, the present invention provides piperidine derivatives of formula (I), pharmaceutical compositions containing them, methods for preparing said compounds, and methods of treating renin mediated conditions by administration of a therapeutically effective amount of a compound of the present invention, or a pharmaceutical composition thereof.

The groups of compounds mentioned below are not to be regarded as exclusive, rather, e.g., in order to replace general definitions with more specific definitions, parts of those groups of compounds can be interchanged or exchanged for the definitions given above, or omitted, as appropriate.

Preferred are compounds of formula (I) wherein

R_1 is $-\text{CH}_2\text{-X}$, $-\text{O-X}$ or $-\text{S(O)}_{0-2}\text{-X}$; or

R_1 is $-\text{NR}_8\text{-X}$, $-\text{NR}_8\text{C(O)-X}$ or $-\text{NR}_8\text{S(O)}_2\text{-X}$ in which

R_8 is hydrogen or lower alkyl; and

X is $-(\text{CH}_2)_m\text{-(CR}_9\text{R}_{10})_p\text{-(CH}_2)_n\text{-Z-(CH}_2)_q\text{-W}$ in which

m and n are independently zero or an integer from 1 to 5;

p is zero or 1;

q is zero;

R_9 and R_{10} are independently hydrogen, hydroxy, halogen, lower alkyl, lower alkoxy or cycloalkyl; or

R_9 and R_{10} combined are alkylene which together with the carbon atom to which they are attached form a 3- to 6-membered ring;

Z is a bond; or

Z is O, S(O)₀₋₂, or -NR₁₁- in which

R₁₁ is hydrogen or lower alkyl, provided that R₁ is -CH₂-X when m, n and p are all zero;

W is aryl or heterocyclyl;

R₂ is hydrogen, halogen, cyano, hydroxy or lower alkoxy;

L is a bond; or

L is -(CH₂)_s-O-(CH₂)_v- in which

s and v are zero; or

L is -C(O)-, -C(O)O-, -OC(O)-, -OC(O)NR₁₂-, -NR₁₂-, -NR₁₃C(O)-, -NR₁₃C(O)O- or -NR₁₃C(O)NR₁₂- in which

R₁₂ and R₁₃ are independently hydrogen or lower alkyl;

R₃ is hydrogen, halogen or cyano provided that L is a bond; or

R₃ is optionally substituted lower alkyl, aralkyl, heteroaralkyl, aryl or heterocyclyl; or

R₃ and R₁₂ combined are alkylene which together with the nitrogen atom to which they are attached form a 5- to 6-membered ring;

R₄ is hydrogen, optionally substituted lower alkyl or aryl;

R₅ and R₆ are independently hydrogen, halogen, hydroxy, trifluoromethyl, optionally substituted lower alkyl, lower alkoxy or cycloalkyl; or

R₅ and R₆ combined together with the carbon atoms to which they are attached form a fused 5- to 6-membered aromatic or heteroaromatic ring provided that R₅ and R₆ are attached to carbon atoms adjacent to each other; or

R₅ and R₆ combined are alkylene which together with the carbon atoms to which they are attached form a fused 5- to 7-membered ring provided that R₅ and R₆ are attached to carbon atoms adjacent to each other; or

R₇ is hydrogen, halogen, hydroxy, trifluoromethyl, optionally substituted, lower alkyl, lower alkoxy or cycloalkyl; or

R₇ and R₈ combined are O, S(O)₀₋₂, -NR₁₄-, -(CH₂)₁₋₂-, -O-CH₂-, -CH₂-O-, -S(O)₀₋₂-CH₂-, -CH₂-S(O)₀₋₂-, -NR₁₄-CH₂-, -CH₂-NR₁₄-, -S(O)₀₋₂-NR₁₄- or -NR₁₄-S(O)₀₋₂- in which

R₁₄ is hydrogen or lower alkyl, provided R₈ is located at the 2' position;

- 11 -

Y is $-(CH_2)_r-$ in which


r is zero;

Q combined with the carbon atoms to which it is attached form a 5- to 6-membered monocyclic aromatic or heteroaromatic ring; or

Q combined with the carbon atoms to which it is attached form a 9- to 10-membered bicyclic aromatic or heterocyclic ring;

or a pharmaceutically acceptable salt thereof.

Further preferred are compounds of formula (I) having the formula




wherein

R₁, R₂, L, R₃, R₄, R₅, R₆, R₇ and Q have meanings as defined herein above;

or a pharmaceutically acceptable salt thereof.

Further preferred are also the compounds of formula (I) having the formula



wherein

R₁, R₂, L, R₃, R₄, R₅, R₆, R₇ and Q have meanings as defined herein above;

or a pharmaceutically acceptable salt thereof.

Preferred are compounds of formula (IB), designated as the A group, wherein

R_1 is $-\text{CH}_2\text{-X}$, $-\text{O-X}$ or $-\text{S-X}$; or

R_1 is $-\text{NR}_6\text{-X}$, $-\text{NR}_6\text{C(O)-X}$ or $-\text{NR}_6\text{S(O)}_2\text{-X}$ in which

R_6 is hydrogen or lower alkyl; and

X is $-(\text{CH}_2)_m\text{-(CR}_9\text{R}_{10})_p\text{-(CH}_2)_n\text{-Z-W}$ in which

m and n are independently zero or an integer of 1 or 2;

p is zero or 1;

R_9 and R_{10} are independently hydrogen, hydroxy, halogen, lower alkyl, lower alkoxy or cycloalkyl; or

R_9 and R_{10} combined are alkylene which together with the carbon atom to which they are attached form a 3- to 6-membered ring;

Z is a bond; or

Z is O , S(O)_{0-2} , or $-\text{NR}_{11}-$ in which

R_{11} is hydrogen or lower alkyl, provided that R_1 is $-\text{CH}_2\text{-X}$ when m , n and p are all zero;

W is aryl or heterocyclyl;

R_2 is hydrogen, halogen or hydroxy;

L is a bond;

R_3 is hydrogen or halogen;

R_4 is hydrogen, optionally substituted lower alkyl or aryl;

R_5 and R_8 are independently hydrogen, halogen, hydroxy, trifluoromethyl, optionally substituted lower alkyl, lower alkoxy or cycloalkyl;

R_7 is hydrogen, halogen, hydroxy, trifluoromethyl, optionally substituted lower alkyl, lower alkoxy or cycloalkyl; or

R_7 and R_8 combined are O , S(O)_{0-2} , $-\text{NR}_{14}-$, $-(\text{CH}_2)_{1-2}-$, $-\text{O-CH}_2-$, $-\text{CH}_2\text{-O-}$, $-\text{S(O)}_{0-2}\text{-CH}_2-$, $-\text{CH}_2\text{-S(O)}_{0-2}-$, $-\text{NR}_{14}\text{-CH}_2-$, $-\text{CH}_2\text{-NR}_{14}-$, $-\text{S(O)}_{0-2}\text{-NR}_{14}-$ or $-\text{NR}_{14}\text{-S(O)}_{0-2}-$ in which

R_{14} is hydrogen or lower alkyl, provided R_8 is located at the 2'-position;

Q combined with the atoms to which it is attached form a 5- to 6-membered monocyclic aromatic or heteroaromatic ring; or

Q combined with the atoms to which it is attached form a 9- to 10-membered bicyclic aromatic or heterocyclic ring;

or a pharmaceutically acceptable salt thereof.

Preferred are the compounds in the A group, designated as the B group, wherein

R_1 is $-\text{CH}_2\text{-X}$, $-\text{O-X}$ or $-\text{S-X}$; or

R_1 is $-\text{NR}_8\text{-X}$, $-\text{NR}_8\text{C(O)-X}$ or $-\text{NR}_8\text{S(O)}_2\text{-X}$ in which

R_8 is hydrogen or lower alkyl; and

X is $-(\text{CH}_2)_m\text{-(CR}_9\text{R}_{10})_p\text{-(CH}_2)_n\text{-Z-W}$ in which

m and n are independently zero or an integer of 1 or 2;

p is zero or 1;

R_9 and R_{10} are independently hydrogen or lower alkyl; or

Z is a bond; or

Z is O, S(O)_{0-2} , or $-\text{NR}_{11}-$ in which

R_{11} is hydrogen or lower alkyl, provided that R_1 is $-\text{CH}_2\text{-X}$ when m, n and p are all zero;

W is aryl or heterocyclyl;

R_2 is hydrogen, halogen or hydroxy;

L is a bond;

R_3 is hydrogen or halogen;

R_4 is hydrogen;

R_5 and R_6 are independently hydrogen, halogen, hydroxy, trifluoromethyl, optionally substituted lower alkyl, lower alkoxy or cycloalkyl;

R_7 is hydrogen, halogen, hydroxy, trifluoromethyl, optionally substituted lower alkyl, lower alkoxy or cycloalkyl; or

R_7 and R_8 combined are O, S(O)_{0-2} , $-\text{NR}_{14}-$, $-(\text{CH}_2)_{1-2}-$, $-\text{O-CH}_2-$, $-\text{CH}_2\text{-O-}$, $-\text{S(O)}_{0-2}\text{-CH}_2-$, $-\text{CH}_2\text{-S(O)}_{0-2}-$, $-\text{NR}_{14}\text{-CH}_2-$, $-\text{CH}_2\text{-NR}_{14}-$, $-\text{S(O)}_{0-2}\text{-NR}_{14}-$ or $-\text{NR}_{14}\text{-S(O)}_{0-2}-$ in which

R_{14} is hydrogen or lower alkyl, provided R_6 is located at the 2'-position;

Q combined with the atoms to which it is attached form a 5- to 6-membered monocyclic aromatic or heteroaromatic ring; or

Q combined with the atoms to which it is attached form a 9- to 10-membered bicyclic aromatic or heterocyclic ring;

or a pharmaceutically acceptable salt thereof.

Preferred are the compounds in the B group wherein

R_1 is $-NR_8-X$, $-NR_8C(O)-X$ or $-NR_8S(O)_2-X$ in which

R_8 is hydrogen or lower alkyl;

or a pharmaceutically acceptable salt thereof.

Preferred are also the compounds in the B group wherein

Q combined with the carbon atoms to which it is attached form a pyridyl or pyrimidinyl ring;


or a pharmaceutically acceptable salt thereof.

Preferred are also the compounds in the B group wherein

Q combined with the carbon atoms to which it is attached form a thienyl, furyl, pyrrolyl or indolyl ring;

or a pharmaceutically acceptable salt thereof.

Preferred are also the compounds in the A group, designated as the C group, having the formula



wherein

R_1 is $-CH_2-X$, $-O-X$ or $-S-X$; or

- 15 -

R_1 is $-NR_8-X$, $-NR_8C(O)-X$ or $-NR_8S(O)_2-X$ in which

R_8 is hydrogen or lower alkyl; and

X is $-(CH_2)_m-(CR_9R_{10})_p-(CH_2)_n-Z-W$ in which

m , n and p are independently zero or 1;

R_9 is hydrogen;

R_{10} is hydrogen or lower alkyl;

Z is a bond; or

Z is O , $S(O)_{0-2}$, or $-NR_{11}-$ in which

R_{11} is hydrogen or lower alkyl, provided that R_1 is $-CH_2-X$ when m , n and p are all zero;

W is aryl or heterocyclyl;

R_2 is hydrogen;

R_3 is hydrogen or halogen;

R_5 and R_6 are independently hydrogen, halogen, hydroxy, trifluoromethyl, optionally substituted lower alkyl, lower alkoxy or cycloalkyl;

R_7 is hydrogen, halogen, hydroxy, trifluoromethyl, optionally substituted lower alkyl, lower alkoxy or cycloalkyl; or

R_7 and R_6 combined are O , $S(O)_{0-2}$, $-NR_{14}-$, $-(CH_2)_{1-2}-$, $-O-CH_2-$, $-CH_2-O-$, $-S(O)_{0-2}-CH_2-$, $-CH_2-S(O)_{0-2}-$, $-NR_{14}-CH_2-$, $-CH_2-NR_{14}-$, $-S(O)_{0-2}-NR_{14}-$ or $-NR_{14}-S(O)_{0-2}-$ in which

R_{14} is hydrogen or lower alkyl, provided R_6 is located at the 2'-position;

R_{15} is hydrogen, halogen, hydroxy, trifluoromethyl, optionally substituted lower alkyl, lower alkoxy or cycloalkyl;

or a pharmaceutically acceptable salt thereof.

Preferred are the compounds in the C group wherein

R_1 is $-O-X$ or $-S-X$; and

X is $-(CH_2)_m-(CR_9R_{10})_p-(CH_2)_n-Z-W$ in which

m is 1;

n and p are zero;

Z is a bond;

W is aryl or heterocyclyl;

R₃ is hydrogen or halogen;

R₅ is hydrogen, halogen, hydroxy, trifluoromethyl, optionally substituted lower alkyl, lower alkoxy or cycloalkyl;

R₆ is hydrogen;

R₇ is hydrogen;

R₁₅ is hydrogen, halogen, hydroxy, trifluoromethyl, optionally substituted lower alkyl, lower alkoxy or cycloalkyl;

or a pharmaceutically acceptable salt thereof.

Further preferred are the compounds in the C group wherein

R₃ is hydrogen;

or a pharmaceutically acceptable salt thereof.

More preferred are the compounds in the C group wherein

W is monocyclic aryl;

or a pharmaceutically acceptable salt thereof.


Preferred are also the compounds in the A group, designated as the D group, wherein

p is 1;

R₉ and R₁₀ combined are alkylene which together with the carbon atom to which they are attached form a 3- to 6-membered ring;

or a pharmaceutically acceptable salt thereof.

Preferred are the compounds in the D group having the formula



wherein

R_1 is $-\text{CH}_2\text{-X}$, $-\text{O-X}$ or $-\text{S-X}$; or

R_1 is $-\text{NR}_8\text{-X}$, $-\text{NR}_8\text{C(O)-X}$ or $-\text{NR}_8\text{S(O)}_2\text{-X}$ in which

R_8 is hydrogen or lower alkyl; and

X is $-(\text{CH}_2)_m\text{-CR}_9\text{R}_{10}\text{-(CH}_2)_n\text{-Z-W}$ in which

m and n are 1;

Z is a bond; or

Z is O , S(O)_{0-2} , or $-\text{NR}_{11}-$ in which

R_{11} is hydrogen or lower alkyl, provided that R_1 is $-\text{CH}_2\text{-X}$ when m , n and p are all zero;

W is aryl or heterocyclyl;

R_2 is hydrogen;

R_3 is hydrogen or halogen;

R_5 and R_6 are independently hydrogen, halogen, hydroxy, trifluoromethyl, optionally substituted lower alkyl, lower alkoxy or cycloalkyl;

R_7 is hydrogen, halogen, hydroxy, trifluoromethyl, optionally substituted lower alkyl, lower alkoxy or cycloalkyl; or

R_7 and R_6 combined are O , S(O)_{0-2} , $-\text{NR}_{14}-$, $-(\text{CH}_2)_{1-2}-$, $-\text{O-CH}_2-$, $-\text{CH}_2\text{-O-}$, $-\text{S(O)}_{0-2}\text{-CH}_2-$, $-\text{CH}_2\text{-S(O)}_{0-2}-$, $-\text{NR}_{14}\text{-CH}_2-$, $-\text{CH}_2\text{-NR}_{14}-$, $-\text{S(O)}_{0-2}\text{-NR}_{14}-$ or $-\text{NR}_{14}\text{-S(O)}_{0-2}-$ in which

R_{14} is hydrogen or lower alkyl, provided R_8 is located at the 2'-position;

R_{15} is hydrogen, halogen, hydroxy, trifluoromethyl, optionally substituted lower alkyl, lower alkoxy or cycloalkyl;

or a pharmaceutically acceptable salt thereof.

Further preferred are the compounds in the D group wherein

R_4 is $-\text{O-X}$ or $-\text{S-X}$; and

X is $-\text{CH}_2\text{-CR}_9\text{R}_{10}\text{-CH}_2\text{-Z-W}$ in which

Z is a bond;

W is aryl;

R₃ is hydrogen;

R₅ is hydrogen, halogen, hydroxy, trifluoromethyl, optionally substituted lower alkyl, lower alkoxy or cycloalkyl;

R₆ is hydrogen;

R₇ is hydrogen;

R₁₅ is hydrogen, halogen, hydroxy, trifluoromethyl, optionally substituted lower alkyl, lower alkoxy or cycloalkyl;

or a pharmaceutically acceptable salt thereof.

The compounds of the present invention possess two or more asymmetric centers depending on the choice of the substituents. The preferred absolute configuration at the C-3 and C-4 asymmetric centers is maintained throughout the specification and the appended claims as indicated herein above. However, any possible diastereoisomers, enantiomers and geometric isomers, and mixtures thereof, e.g., racemates, are encompassed by the instant invention.

Particular embodiments of the invention are:

(3R*,4R*)-4-Biphenyl-3-yl-3-(naphthalen-2-ylmethoxy)-piperidine hydrochloride;
 (3R*,4R*)-4-Biphenyl-3-yl-3-(biphenyl-4-ylmethoxy)-piperidine hydrochloride;
 (3R*,4R*)-3-Benzoyloxy-4-biphenyl-3-yl-piperidine hydrochloride;
 (3R*,4R*)-4-Biphenyl-3-yl-3-(4-bromo-benzoyloxy)-piperidine hydrochloride;
 (3R*,4R*)-4-Biphenyl-3-yl-3-(4-trifluoromethoxy-benzoyloxy)-piperidine hydrochloride;
 (3R*,4R*)-4-Biphenyl-3-yl-3-(3-phenoxy-benzoyloxy)-piperidine hydrochloride;
 (3R*,4R*)-4-Dibenzofuran-4-yl-3-(naphthalen-2-ylmethoxy)-piperidine hydrochloride;
 (3R*,4R*)-3-(Biphenyl-4-ylmethoxy)-4-dibenzofuran-4-yl-piperidine hydrochloride;
 (3R*,4R*)-3-Benzoyloxy-4-dibenzofuran-4-yl-piperidine hydrochloride;
 (3R*,4R*)-4-Dibenzofuran-4-yl-3-(3-phenoxy-benzoyloxy)-piperidine hydrochloride;
 [3-((3R*,4R*)-4-Dibenzofuran-4-yl-piperidin-3-ylloxymethyl)-phenyl]-phenyl-methanone hydrochloride;
 (3R*,4R*)-4-(4'-Methoxy-biphenyl-3-yl)-3-(naphthalen-2-ylmethoxy)-piperidine hydrochloride;
 3'-[(3R*,4R*)-3-(Naphthalen-2-ylmethoxy)-piperidin-4-yl]-biphenyl-3-ol hydrochloride;

(3R*,4R*)-4-[3'-(2-Methoxy-ethoxy)-biphenyl-3-yl]-3-(naphthalen-2-ylmethoxy)-piperidine hydrochloride;
N-{3'-[(3R*,4R*)-3-(Naphthalen-2-ylmethoxy)-piperidin-4-yl]-biphenyl-3-yl}-acetamide hydrochloride;
1-{3'-[(3R*,4R*)-3-(Naphthalen-2-ylmethoxy)-piperidin-4-yl]-biphenyl-4-yl}-ethanone hydrochloride;
1-{3'-[(3R*,4R*)-3-(Naphthalen-2-ylmethoxy)-piperidin-4-yl]-biphenyl-3-yl}-ethanone hydrochloride;
3'-[(3R*,4R*)-3-(Naphthalen-2-ylmethoxy)-piperidin-4-yl]-biphenyl-3-carboxylic acid ethyl ester hydrochloride;
(3R*,4R*)-4-{3'-Chloro-biphenyl-3-yl}-3-(naphthalen-2-ylmethoxy)-piperidine hydrochloride;
(3R*,4R*)-3-(Naphthalen-2-ylmethoxy)-4-{3'-trifluoromethyl-biphenyl-3-yl}-piperidine hydrochloride;
(3R*,4R*)-4-{4'-Chloro-biphenyl-3-yl}-3-(naphthalen-2-ylmethoxy)-piperidine hydrochloride;
3'-[(3R*,4R*)-3-(Naphthalen-2-ylmethoxy)-piperidin-4-yl]-biphenyl-4-ol hydrochloride;
Dimethyl-{3'-[(3R*,4R*)-3-(Naphthalen-2-ylmethoxy)-piperidin-4-yl]biphenyl-4-yl}-amine hydrochloride;
3'-[(3R*,4R*)-3-(Naphthalen-2-ylmethoxy)-piperidin-4-yl]-biphenyl-3-carboxylic acid hydrochloride;
3'-[(3R*,4R*)-3-(Naphthalen-2-ylmethoxy)-piperidin-4-yl]-biphenyl-4-carboxylic acid methylamide hydrochloride;
4-{3'-[(3R*,4R*)-3-(Naphthalen-2-ylmethoxy)-piperidin-4-yl]-phenyl}-pyridine hydrochloride;
(3R*,4R*)-3-(Naphthalen-2-ylmethoxy)-4-(3-thiophen-3-yl-phenyl)-piperidine hydrochloride;
3-{3'-[(3R*,4R*)-3-(Naphthalen-2-ylmethoxy)-piperidin-4-yl]-phenyl}-pyridine hydrochloride;
2-{3'-[(3R*,4R*)-3-(Naphthalen-2-ylmethoxy)-piperidin-4-yl]-phenyl}-1H-indole hydrochloride;
5-{3'-[(3R*,4R*)-3-(Naphthalen-2-ylmethoxy)-piperidin-4-yl]-phenyl}-pyrimidine hydrochloride;
(3R,4R)-4-Biphenyl-3-yl-3-[2-(3-methoxy-propoxy)-4-methyl-benzyloxy]-piperidine;
(3R,4R)-4-Biphenyl-3-yl-3-[2-(3-methoxy-propoxy)-benzyloxy]-piperidine trifluoroacetic acid;
(3R,4R)-4-Biphenyl-3-yl-3-[4-methoxy-3-(3-methoxy-propoxy)-benzyloxy]-piperidine;
(3R,4R)-4-Biphenyl-3-yl-3-[3-(4-methoxy-butyl)-benzyloxy]-piperidine;
(3R,4R)-4-Biphenyl-3-yl-3-[3-(3-methoxy-propyl)-benzyloxy]-piperidine;


(3R,4R)-4-Biphenyl-3-yl-3-[3-(2-methoxy-ethyl)-benzyloxy]-piperidine;
 (3R,4R)-4-Biphenyl-3-yl-3-[3-(3-methoxy-propyl)-4-methyl-benzyloxy]-piperidine;
 (3R,4R)-4-Biphenyl-3-yl-3-[3-(4-methoxy-butyl)-4-methyl-benzyloxy]-piperidine;
 (3R,4R)-4-Biphenyl-3-yl-3-[2-(3-methoxy-propyl)-benzyloxy]-piperidine trifluoroacetic acid;
 (3R,4R)-3-(1-Benzyl-cyclopropylmethoxy)-4-biphenyl-3-yl-piperidine trifluoroacetic acid; and
 (3R,4R)-4-Biphenyl-3-yl-3-[1-[2-(3-methoxy-propyl)-benzyl]-cyclopropylmethoxy]-piperidine hydrochloride;

or a pharmaceutically acceptable salt thereof.


Compounds of formula (I) may be prepared from known starting materials according to methods well known in the art, e.g., using methods disclosed in International PCT Patent Application No. WO 97/09311, or as described herein in the illustrative Examples, or modifications thereof.

For example, compounds of formula (IC) having formula (IC') wherein R_5 , R_6 , R_7 , R_{15} have meanings as defined herein above, and R_1 is -O-X in which X has a meaning as defined herein above, may be prepared as outlined in Scheme 1.

Scheme 1:



Compounds of formula (IIa) wherein Pg represents a protecting group such as a lower alkoxy carbonyl, e.g., *t*-butoxycarbonyl or ethoxycarbonyl, may be converted to compounds of formula (III) under conditions of Suzuki or Stille coupling, e.g., a compound of formula (IIa) may be reacted with a boronic acid of the formula



wherein R_5 , R_6 , R_7 and R_{15} have meanings as defined above, in the presence of a palladium catalyst such as tetrakis(triphenylphosphine)palladium(0) and a base such as sodium or potassium carbonate in an appropriate solvent, e.g., water, dimethoxyethane, acetonitrile, methanol, ethanol or THF, or a mixture of solvents thereof. If desired, the Suzuki reaction may be conducted in the presence of an additive such as lithium chloride. Boronic acids of formula (VI) are known, or if they are novel, they may be prepared using methods known in the art.

A resulting compound of formula (III) may then be hydroxylated under appropriate reaction conditions, e.g., under conditions of hydroboration, to afford a compound of formula (IV) wherein Pg, R_5 , R_6 , R_7 and R_{15} have meanings as defined above.

A resulting compound of formula (IV) is obtained as a racemic mixture according to the reaction sequence as outlined in Scheme 1. Racemic compounds, e.g., those of formula (IV) and (IC'), may be separated into their pure enantiomers using conventional methods known in the art.

Alternatively, compounds of formula (IV) wherein Pg, R_5 , R_6 , R_7 and R_{15} have meanings as defined above may be obtained in high enantioselectivity from compounds of formula (III) by a reaction sequence involving asymmetric dihydroxylation followed by catalytic reduction, as described by Rich et al. in *Organic Letters*, **2001**, 3, 2317-2320, or as described in International PCT Patent Application No. WO 00/63173.

Compounds of formula (IV) wherein Pg, R_5 , R_6 , R_7 and R_{15} have meanings as defined above may then be treated, e.g., with an alkylating agent of the formula





wherein X has a meaning as defined herein, and Lg₁ represents a leaving group such as iodide, bromide, chloride, trifluoromethylsulfonate, mesylate or tosylate, preferably bromide, in the presence of a base such as sodium hydride in an organic solvent, such as tetrahydrofuran (THF), *N,N*-dimethylformamide (DMF) or *N,N*-dimethylacetamide (DMA), to afford compounds of formula (V). Alkylating agents of formula (VII) are known, or if they are novel, they may be prepared as illustrated herein in the Examples, or using methods well known in the art.

Finally, compounds of formula (V) wherein Pg, X, R₅, R₆, R₇ and R₁₅ have meanings as defined above may be converted to compounds of formula (IC') wherein R₅, R₆, R₇ R₁₅ have meanings as defined herein above, and R₁ is -O-X in which X has a meaning as defined herein above, or a pharmaceutically acceptable salt thereof, by removal of the protecting group, e.g., when Pg is *t*-butoxycarbonyl by treatment with an acid such as trifluoroacetic acid (TFA) or hydrochloric acid (HCl), or when Pg is ethoxycarbonyl by treatment with a base such as aqueous sodium or potassium hydroxide.


Alternatively, compounds of formula (IC') wherein R₅, R₆, R₇ R₁₅ have meanings as defined herein above, and R₁ is -O-X in which X has a meaning as defined herein above, may be obtained from compounds of formula (IIb) following the reaction sequence as outlined in Scheme 2.

Scheme 2:





As illustrated in Scheme 2, compounds of formula (IIb) wherein Pg represents a protecting group as described herein above, and R_{16} and R_{17} are lower alkyl, or R_{16} and R_{17} combined are alkylene which together with the boron and the oxygen atoms form a 5- or 6-membered ring, preferably, R_{16} and R_{17} combined are 1,1,2,2-tetramethylethylene, may be converted to compounds of formula (VIII) under conditions of Suzuki coupling, e.g., a compound of formula (IIb) may be treated with a compound of the formula




wherein R_5 and R_6 have meanings as defined above, in the presence of a palladium catalyst such as tetrakis(triphenylphosphine)palladium(0) and a base such as sodium or potassium carbonate in an appropriate solvent, e.g., water, dimethoxyethane, acetonitrile, methanol, ethanol or THF, or a mixture of solvents thereof. If desired, the Suzuki reaction may be conducted in the presence of an additive such as lithium chloride. Compounds of formulae (IIb) and (XI) are known, or if they are novel, they may be prepared using to methods known in the art.

A resulting compound of formula (VIII) may then be converted to a compound of formula (IX) wherein Pg, R_5 and R_6 have meanings as defined above, as described herein for compounds of formula (IV).

Compounds of formula (IX) wherein Pg, R_5 and R_6 have meanings as defined herein above, may then be treated, e.g., with an alkylating agent of formula (VII), to afford compounds of formula (X) wherein Pg, X, R_5 and R_6 have meanings as defined herein above, under reaction conditions as described herein for compounds of formula (V).


Finally, compounds of formula (X) wherein Pg, X, R₅ and R₆ have meanings as defined herein above may be converted to compounds of formula (IC') wherein X, R₅, R₆, R₇ and R₁₅ have meanings as defined herein above, or a pharmaceutically acceptable salt thereof, by first coupling a compound of formula (X) with a boronic acid of the formula




wherein R₇ and R₁₅ have meanings as defined herein above, followed by removal of the protecting group under appropriate reaction conditions, e.g., as illustrated herein above.


In addition, compounds of formula (IC') wherein R₅, R₆, R₇ and R₁₅ have meanings as defined herein above, and R₁ is -S(O)₀₋₂-X, -NR₈-X, -NR₈C(O)-X or -NR₈S(O)₂-X in which R₈ and X have meanings as defined herein above, may be prepared from compounds of formula (IV) using methods well known in the art.

Scheme 3:





(XIV)



(XV)

For example, as illustrated in Scheme 3, a compound of formula (IV) wherein Pg, R₅, R₆, R₇ and R₁₅ have meanings as defined herein above, may be converted to a compound of formula (XIV) or (XV), e.g., by first converting the hydroxyl group to a leaving group (Lg₂) such as bromide or iodide to afford a compound of formula (XIII) wherein Pg, R₅, R₆, R₇ and R₁₅ have meanings as defined above and Lg₂ represents a leaving group as described herein above. Subsequent reaction with a nucleophile, such as thiolacetic acid or sodium azide, followed by hydrolysis or reduction, respectively, then affords a compound of formula (XIV) or a compound of formula (XV) wherein Pg, R₅, R₆, R₇ and R₁₅ have meanings as defined herein above. Compounds of formulae (XIV) or (XV) may be converted to compounds of formula (IC') wherein R₅, R₆, R₇ and R₁₅ have meanings as defined herein above, and R₁ is -S(O)₀₋₂-X, -NR₈-X, -NR₈C(O)-X or -NR₈S(O)₂-X in which R₈ and X have meanings as defined herein above, using methods described herein or using methods well known in the art.

The processes described herein above may be conducted under inert atmosphere, preferably under nitrogen atmosphere.

As exemplified herein above, in starting compounds and intermediates which are converted to the compounds of the invention in a manner described herein, functional groups present, such as amino, thiol, carboxyl and hydroxy groups, are optionally protected by conventional protecting groups that are common in preparative organic chemistry. Protected amino, thiol, carboxyl and hydroxyl groups are those that can be converted under mild conditions into free amino thiol, carboxyl and hydroxyl groups without the molecular framework being destroyed or other undesired side reactions taking place.

The purpose of introducing protecting groups is to protect the functional groups from undesired reactions with reaction components under the conditions used for carrying out a desired chemical transformation. The need and choice of protecting groups for a particular reaction is known to those skilled in the art and depends on the nature of the functional group to be protected (hydroxyl group, amino group, etc.), the structure and stability of the molecule of which the substituent is a part and the reaction conditions.

Well-known protecting groups that meet these conditions and their introduction and removal are described, e.g., in McOmie, "Protective Groups in Organic Chemistry", Plenum Press, London, NY (1973); and Greene and Wuts, "Protective Groups in Organic Synthesis", John Wiley and Sons, Inc., NY (1999).

The above-mentioned reactions are carried out according to standard methods, in the presence or absence of diluent, preferably, such as are inert to the reagents and are solvents thereof, of catalysts, condensing or said other agents, respectively and/or inert atmospheres, at low temperatures, room temperature (RT) or elevated temperatures, preferably at or near the boiling point of the solvents used, and at atmospheric or super-atmospheric pressure. The preferred solvents, catalysts and reaction conditions are set forth in the appended illustrative Examples.

The invention further includes any variant of the present processes, in which an intermediate product obtainable at any stage thereof is used as starting material and the remaining steps are carried out, or in which the starting materials are formed *in situ* under the reaction conditions, or in which the reaction components are used in the form of their salts or optically pure antipodes.

Compounds of the invention and intermediates can also be converted into each other according to methods generally known *per se*.

The invention also relates to any novel starting materials, intermediates and processes for their manufacture.

Depending on the choice of starting materials and methods, the new compounds may be in the form of one of the possible isomers or mixtures thereof, for example, as substantially pure geometric (*cis* or *trans*) isomers, diastereomers, optical isomers (antipodes), racemates or mixtures thereof. The aforesaid possible isomers or mixtures thereof are within the purview of this invention.

Any resulting mixtures of isomers can be separated on the basis of the physicochemical differences of the constituents, into the pure geometric or optical isomers, diastereomers, racemates, for example, by chromatography and/or fractional crystallization.

Any resulting racemates of final products or intermediates can be resolved into the optical antipodes by known methods, e.g., by separation of the diastereomeric salts thereof, obtained with an optically active acid or base, and liberating the optically active acidic or basic compound. In particular, the piperidinyl moiety may be employed to resolve the compounds of the present invention into their optical antipodes, e.g., by fractional crystallization of a salt formed with an optically active acid, e.g., tartaric acid, dibenzoyl tartaric acid, diacetyl tartaric acid, di-O,O'-p-toluoyl tartaric acid, mandelic acid, malic acid or camphor-10-sulfonic acid. Racemic compounds, e.g., those of formula (IV), (IX), (XIV), (XV) and (IC'), may also be resolved by forming covalent diastereomeric derivatives, e.g. esters or amides, by condensing an alcohol or a thiol, e.g., those of formula (IV), (IX) or (XIV), or an amine, e.g., those of formula (XV) and (IC'), with a chiral, enantiomerically pure carboxylic acid. The resulting diastereomeric esters or amides can then be separated by crystallization or by chromatography and subsequently hydrolyzed to yield enantiomerically pure compounds of formula (IV), (IX), (XIV), (XV) and (IC'), respectively. Racemic products can also be resolved by chiral chromatography, e.g., high pressure liquid chromatography (HPLC) using a chiral adsorbent.

Finally, compounds of the invention are either obtained in the free form, as a salt thereof, or as prodrug derivatives thereof.

Compounds of the invention having basic groups, in particular, the piperidinyl moiety, can be converted into acid addition salts, especially pharmaceutically acceptable salts. These are formed, for example, with inorganic acids, such as mineral acids, for example, sulfuric acid, a phosphoric or hydrohalic acid, or with organic carboxylic acids, such as (C₁-C₄)-alkanecarboxylic acids which, for example, are unsubstituted or substituted by halogen, for example, acetic acid, such as saturated or unsaturated dicarboxylic acids, for example, oxalic, succinic, maleic or fumaric acid, such as hydroxycarboxylic acids, for example glycolic, lactic, malic, tartaric or citric acid, such as amino acids, for example, aspartic or glutamic acid, or with organic sulfonic acids, such as (C₁-C₄)-alkylsulfonic acids, for example, methanesulfonic acid; or arylsulfonic acids which are unsubstituted or substituted (for example by halogen). Preferred are salts formed with hydrochloric acid, methanesulfonic acid and maleic acid.

Compounds of the instant invention which contain acidic groups may be converted into salts with pharmaceutically acceptable bases. Such salts include alkali metal salts, like sodium, lithium and potassium salts; alkaline earth metal salts, like calcium and magnesium salts; ammonium salts with organic bases, e.g., trimethylamine salts, diethylamine salts, tris(hydroxymethyl)methylamine salts, dicyclohexylamine salts and N-methyl-D-glucamine salts; salts with amino acids like arginine, lysine and the like. Salts may be formed using conventional methods, advantageously in the presence of an ethereal or alcoholic solvent, such as a lower alkanol. From the solutions of the latter, the salts may be precipitated with ethers, e.g., diethyl ether. Resulting salts may be converted into the free compounds by treatment with acids. These or other salts can also be used for purification of the compounds obtained.

Prodrug derivatives of any compound of the invention are derivatives of said compounds which following administration release the parent compound *in vivo* via some chemical or physiological process, e.g., a prodrug on being brought to the physiological pH or through enzyme action is converted to the parent compound. Exemplary prodrug derivatives are, e.g., esters of free carboxylic acids and S-acyl and O-acyl derivatives of thiols, alcohols or phenols, wherein acyl has a meaning as defined herein. Preferred are pharmaceutically acceptable ester derivatives convertible by solvolysis under physiological conditions to the parent carboxylic acid, e.g., lower alkyl esters, cycloalkyl esters, lower alkenyl esters, benzyl esters, mono- or di-substituted lower alkyl esters, such as the ω -(amino, mono- or di-lower alkylamino, carboxy, lower alkoxy carbonyl)-lower alkyl esters, the α -(lower alkanoyloxy, lower alkoxy carbonyl or di-lower alkylaminocarbonyl)-lower alkyl esters, such as the pivaloyloxymethyl ester and the like conventionally used in the art.

In view of the close relationship between the free compounds, the prodrug derivatives and the compounds in the form of their salts, whenever a compound is referred to in this context, a prodrug derivative and a corresponding salt is also intended, provided such is possible or appropriate under the circumstances.

The compounds, including their salts, can also be obtained in the form of their hydrates, or include other solvents used for their crystallization.

As described above, the compounds of the present invention are inhibitors of renin activity and, thus, may be employed for the treatment of hypertension, atherosclerosis, unstable coronary syndrome, congestive heart failure, cardiac hypertrophy, cardiac fibrosis,

cardiomyopathy postinfarction, unstable coronary syndrome, diastolic dysfunction, chronic kidney disease, hepatic fibrosis, complications resulting from diabetes, such as nephropathy, vasculopathy and neuropathy, diseases of the coronary vessels, restenosis following angioplasty, raised intra-ocular pressure, glaucoma, abnormal vascular growth, hyperaldosteronism, cognitive impairment, alzheimers, dementia, anxiety states and cognitive disorders.

The present invention further provides pharmaceutical compositions comprising a therapeutically effective amount of a pharmacologically active compound of the instant invention, alone or in combination with one or more pharmaceutically acceptable carriers.

The pharmaceutical compositions according to the present invention are those suitable for enteral, such as oral or rectal, transdermal and parenteral administration to mammals, including man, to inhibit renin activity, and for the treatment of conditions associated with renin activity. Such conditions include hypertension, atherosclerosis, unstable coronary syndrome, congestive heart failure, cardiac hypertrophy, cardiac fibrosis, cardiomyopathy postinfarction, unstable coronary syndrome, diastolic dysfunction, chronic kidney disease, hepatic fibrosis, complications resulting from diabetes, such as nephropathy, vasculopathy and neuropathy, diseases of the coronary vessels, restenosis following angioplasty, raised intra-ocular pressure, glaucoma, abnormal vascular growth, hyperaldosteronism, cognitive impairment, alzheimers, dementia, anxiety states and cognitive disorders.

Thus, the pharmacologically active compounds of the invention may be employed in the manufacture of pharmaceutical compositions comprising an effective amount thereof in conjunction or admixture with excipients or carriers suitable for either enteral or parenteral application. Preferred are tablets and gelatin capsules comprising the active ingredient together with:

- a) diluents, e.g., lactose, dextrose, sucrose, mannitol, sorbitol, cellulose and/or glycine;
- b) lubricants, e.g., silica, talcum, stearic acid, its magnesium or calcium salt and/or polyethyleneglycol; for tablets also
- c) binders, e.g., magnesium aluminum silicate, starch paste, gelatin, tragacanth, methylcellulose, sodium carboxymethylcellulose and or polyvinylpyrrolidone; if desired
- d) disintegrants, e.g., starches, agar, alginic acid or its sodium salt, or effervescent mixtures; and/or
- e) absorbants, colorants, flavors and sweeteners.

Injectable compositions are preferably aqueous isotonic solutions or suspensions, and suppositories are advantageously prepared from fatty emulsions or suspensions.

Said compositions may be sterilized and/or contain adjuvants, such as preserving, stabilizing, wetting or emulsifying agents, solution promoters, salts for regulating the osmotic pressure and/or buffers. In addition, they may also contain other therapeutically valuable substances. Said compositions are prepared according to conventional mixing, granulating or coating methods, respectively, and contain about 0.1-75%, preferably about 1-50%, of the active ingredient.

Suitable formulations for transdermal application include a therapeutically effective amount of a compound of the invention with carrier. Advantageous carriers include absorbable pharmacologically acceptable solvents to assist passage through the skin of the host. Characteristically, transdermal devices are in the form of a bandage comprising a backing member, a reservoir containing the compound optionally with carriers, optionally a rate controlling barrier to deliver the compound of the skin of the host at a controlled and pre-determined rate over a prolonged period of time, and means to secure the device to the skin.

Accordingly, the present invention provides pharmaceutical compositions as described above for the treatment of conditions mediated by renin activity, preferably, hypertension, atherosclerosis, unstable coronary syndrome, congestive heart failure, cardiac hypertrophy, cardiac fibrosis, cardiomyopathy postinfarction, unstable coronary syndrome, diastolic dysfunction, chronic kidney disease, hepatic fibrosis, complications resulting from diabetes, such as nephropathy, vasculopathy and neuropathy, diseases of the coronary vessels, restenosis following angioplasty, raised intra-ocular pressure, glaucoma, abnormal vascular growth, hyperaldosteronism, cognitive impairment, alzheimers, dementia, anxiety states and cognitive disorders.

The pharmaceutical compositions may contain a therapeutically effective amount of a compound of the invention as defined above, either alone or in a combination with another therapeutic agent, e.g., each at an effective therapeutic dose as reported in the art. Such therapeutic agents include:

a) antidiabetic agents such as insulin, insulin derivatives and mimetics; insulin secretagogues such as the sulfonylureas, e.g., Glipizide, glyburide and Amaryl; insulinotropic sulfonylurea receptor ligands such as meglitinides, e.g., nateglinide and repaglinide; peroxisome proliferator-activated receptor (PPAR) ligands; protein tyrosine phosphatase-1B

(PTP-1B) inhibitors such as PTP-112; GSK3 (glycogen synthase kinase-3) inhibitors such as SB-517955, SB-4195052, SB-216763, NN-57-05441 and NN-57-05445; RXR ligands such as GW-0791 and AGN-194204; sodium-dependent glucose cotransporter inhibitors such as T-1095; glycogen phosphorylase A inhibitors such as BAY R3401; biguanides such as metformin; alpha-glucosidase inhibitors such as acarbose; GLP-1 (glucagon like peptide-1), GLP-1 analogs such as Exendin-4 and GLP-1 mimetics; and DPPIV (dipeptidyl peptidase IV) inhibitors such as LAF237;

b) hypolipidemic agents such as 3-hydroxy-3-methyl-glutaryl coenzyme A (HMG-CoA) reductase inhibitors, e.g., lovastatin, pitavastatin, simvastatin, pravastatin, cerivastatin, mevastatin, velostatin, fluvastatin, dalvastatin, atorvastatin, rosuvastatin and rivastatin; squalene synthase inhibitors; FXR (farnesoid X receptor) and LXR (liver X receptor) ligands; cholestyramine; fibrates; nicotinic acid and aspirin;

c) anti-obesity agents such as orlistat; and

d) anti-hypertensive agents, e.g., loop diuretics such as ethacrynic acid, furosemide and torsemide; angiotensin converting enzyme (ACE) inhibitors such as benazepril, captopril, enalapril, fosinopril, lisinopril, moexipril, perinodopril, quinapril, ramipril andtrandolapril; inhibitors of the Na-K-ATPase membrane pump such as digoxin; neutralendopeptidase (NEP) inhibitors; ACE/NEP inhibitors such as omapatrilat, sampatrilat and fasidotril; angiotensin II antagonists such as candesartan, eprosartan, irbesartan, losartan, telmisartan and valsartan, in particular valsartan; β -adrenergic receptor blockers such as acebutolol, atenolol, betaxolol, bisoprolol, metoprolol, nadolol, propranolol, sotalol and timolol; inotropic agents such as digoxin, dobutamine and milrinone; calcium channel blockers such as amlodipine, bepridil, diltiazem, felodipine, nifedipine, nimodipine, nisoldipine and verapamil; aldosterone receptor antagonists; and aldosterone synthase inhibitors.

Other specific anti-diabetic compounds are described by Patel Mona in *Expert Opin Investig Drugs*, 2003, 12(4), 623-633, in the figures 1 to 7, which are herein incorporated by reference. A compound of the present invention may be administered either simultaneously, before or after the other active ingredient, either separately by the same or different route of administration or together in the same pharmaceutical formulation.

The structure of the therapeutic agents identified by code numbers, generic or trade names may be taken from the actual edition of the standard compendium "The Merck Index" or from databases, e.g., Patents International (e.g. IMS World Publications). The corresponding content thereof is hereby incorporated by reference.

Accordingly, the present invention provides pharmaceutical compositions comprising a therapeutically effective amount of a compound of the invention in combination with a therapeutically effective amount of another therapeutic agent, preferably selected from anti-diabetics, hypolipidemic agents, anti-obesity agents or anti-hypertensive agents, most preferably from antidiabetics, anti-hypertensive agents or hypolipidemic agents as described above.

The present invention further relates to pharmaceutical compositions as described above for use as a medicament.

The present invention further relates to use of pharmaceutical compositions or combinations as described above for the preparation of a medicament for the treatment of conditions mediated by renin activity, preferably, hypertension, atherosclerosis, unstable coronary syndrome, congestive heart failure, cardiac hypertrophy, cardiac fibrosis, cardiomyopathy postinfarction, unstable coronary syndrome, diastolic dysfunction, chronic kidney disease, hepatic fibrosis, complications resulting from diabetes, such as nephropathy, vasculopathy and neuropathy, diseases of the coronary vessels, restenosis following angioplasty, raised intra-ocular pressure, glaucoma, abnormal vascular growth, hyperaldosteronism, cognitive impairment, alzheimers, dementia, anxiety states and cognitive disorders.

Thus, the present invention also relates to a compound of formula (I) for use as a medicament, to the use of a compound of formula (I) for the preparation of a pharmaceutical composition for the prevention and/or treatment of conditions mediated by renin activity, and to a pharmaceutical composition for use in conditions mediated by renin activity comprising a compound of formula (I), or a pharmaceutically acceptable salt thereof, in association with a pharmaceutically acceptable diluent or carrier therefor.

The present invention further provides a method for the prevention and/or treatment of conditions mediated by renin activity, which comprises administering a therapeutically effective amount of a compound of the present invention.

A unit dosage for a mammal of about 50-70 kg may contain between about 1 mg and 1000 mg, advantageously between about 5-600 mg of the active ingredient. The therapeutically effective dosage of active compound is dependent on the species of warm-blooded animal (mammal), the body weight, age and individual condition, on the form of administration, and on the compound involved.

In accordance with the foregoing the present invention also provides a therapeutic combination, e.g., a kit, kit of parts, e.g., for use in any method as defined herein, comprising a compound of formula (I), or a pharmaceutically acceptable salt thereof, to be used concomitantly or in sequence with at least one pharmaceutical composition comprising at least another therapeutic agent, preferably selected from anti-diabetic agents, hypolipidemic agents, anti-obesity agents or anti-hypertensive agents. The kit may comprise instructions for its administration.

Similarly, the present invention provides a kit of parts comprising: (i) a pharmaceutical composition of the invention; and (ii) a pharmaceutical composition comprising a compound selected from an anti-diabetic, a hypolipidemic agent, an anti-obesity agent, an anti-hypertensive agent, or a pharmaceutically acceptable salt thereof, in the form of two separate units of the components (i) to (ii).

Likewise, the present invention provides a method as defined above comprising co-administration, e.g., concomitantly or in sequence, of a therapeutically effective amount of a compound of formula (I), or a pharmaceutically acceptable salt thereof, and a second drug substance, said second drug substance being an anti-diabetic, a hypolipidemic agent, an anti-obesity agent or an anti-hypertensive agent, e.g., as indicated above.

Preferably, a compound of the invention is administered to a mammal in need thereof.

Preferably, a compound of the invention is used for the treatment of a disease which responds to modulation of renin activity.

Preferably, the condition associated with renin activity is selected from hypertension, atherosclerosis, unstable coronary syndrome, congestive heart failure, cardiac hypertrophy, cardiac fibrosis, cardiomyopathy postinfarction, unstable coronary syndrome, diastolic dysfunction, chronic kidney disease, hepatic fibrosis, complications resulting from diabetes, such as nephropathy, vasculopathy and neuropathy, diseases of the coronary vessels, restenosis following angioplasty, raised intra-ocular pressure, glaucoma, abnormal vascular growth, hyperaldosteronism, cognitive impairment, alzheimers, dementia, anxiety states and cognitive disorders.

Finally, the present invention provides a method or use which comprises administering a compound of formula (I) in combination with a therapeutically effective amount of an anti-diabetic agent, a hypolipidemic agent, an anti-obesity agent or an anti-hypertensive agent.

Ultimately, the present invention provides a method or use which comprises administering a compound of formula (I) in the form of a pharmaceutical composition as described herein.

As used throughout the specification and in the claims, the term "treatment" embraces all the different forms or modes of treatment as known to those of the pertinent art and in particular includes preventive, curative, delay of onset and/or progression, and palliative treatment.

The above-cited properties are demonstrable *in vitro* and *in vivo* tests using advantageously mammals, e.g., mice, rats, rabbits, dogs, monkeys or isolated organs, tissues and preparations thereof. Said compounds can be applied *in vitro* in the form of solutions, e.g., preferably aqueous solutions, and *in vivo* either enterally, parenterally, advantageously intravenously, e.g., as a suspension or in aqueous solution. The dosage *in vitro* may range between about 10^{-3} molar and 10^{-10} molar concentrations. A therapeutically effective amount *in vivo* may range depending on the route of administration, between about 0.001 and 500 mg/kg, preferably between about 0.1 and 100 mg/kg.

As described above, the compounds of the present invention have enzyme-inhibiting properties. In particular, they inhibit the action of the natural enzyme renin. Renin passes from the kidneys into the blood where it effects the cleavage of angiotensinogen, releasing the decapeptide angiotensin I which is then cleaved in the lungs, the kidneys and other organs to form the octapeptide angiotensin II. The octapeptide increases blood pressure both directly by arterial vasoconstriction and indirectly by liberating from the adrenal glands the sodium-ion-retaining hormone aldosterone, accompanied by an increase in extracellular fluid volume which increase can be attributed to the action of angiotensin II. Inhibitors of the enzymatic activity of renin lead to a reduction in the formation of angiotensin I, and consequently a smaller amount of angiotensin II is produced. The reduced concentration of that active peptide hormone is the direct cause of the hypotensive effect of renin inhibitors.

The action of renin inhibitors may be demonstrated *inter alia* experimentally by means of *in vitro* tests, the reduction in the formation of angiotensin I being measured in various systems (human plasma, purified human renin together with synthetic or natural renin substrate).

Inter alia the following *in vitro* tests may be used:

An extract of human renin from the kidney (0.5 mGU [milli-Goldblatt units]/mL) is incubated for one h at 37 °C and pH 7.2 in 1M aqueous 2-N-(tris-hydroxymethylmethyl)amino-ethanesulfonic acid buffer solution with 23 µg/mL of synthetic renin substrate, the tetradecapeptide H-Asp-Arg-Val-Tyr-Ile-His-ProPhe-His-Leu-Leu-Val-Tyr-Ser-OH. The amount of angiotensin I formed is determined by radioimmunoassay. Each of the inhibitors according to the invention is added to the incubation mixture at different concentrations. The IC₅₀ is defined as the concentration of a particular inhibitor that reduces the formation of angiotensin I by 50%.

Recombinant human renin (expressed in Chinese Hamster Ovary cells and purified using standard methods) at 4 nM concentration is incubated with test compound at various concentrations for 1 h at RT in 0.1 M Tris-HCl buffer, pH 7.4, containing 0.05 M NaCl, 0.5 mM EDTA and 0.05 % CHAPS. Synthetic peptide substrate Arg-Glu(EDANS)-Ile-His-Pro-Phe-His-Leu-Val-Ile-His-Thr-Lys(DABCYL)-Arg₉ is added to a final concentration of 2 µM and increase in fluorescence is recorded at an excitation wave-length of 340 nm and at an emission wave-length of 485 nm in a microplate spectro-fluorimeter. IC₅₀ values are calculated from percentage of inhibition of renin activity as a function of test compound concentration (Fluorescence Resonance Energy Transfer, FRET, assay).

Recombinant human renin (expressed in Chinese Hamster Ovary cells and purified using standard methods) at 1 nM concentration is incubated with test compound at various concentrations for 1.5 h at 37°C in 0.1 M Tris/HCl pH 7.4 containing 0.05 M NaCl, 0.5 mM EDTA and 0.025% (w/v) CHAPS. Synthetic peptide substrate Ac-Ile-His-Pro-Phe-His-Leu-Val-Ile-His-Asn-Lys-[DY-505-X5] is added to a final concentration of 5 µM. The enzyme reaction is stopped by adding 6 µL of 1.0% TFA. The product of the reaction is separated by HPLC and quantified by spectrophotometric measurement at 505 nm wave-length. IC₅₀ values are calculated from percentage of inhibition of renin activity as a function of test compound concentration.

Recombinant human renin (expressed in Chinese Hamster Ovary cells and purified using standard methods) at 3.3 nM concentration, 125I-NVP-AJ1891-NX-1 (0.27 µCi/mL) and streptavidin-SPA (0.67 mg/mL) beads are incubated with test compound at various concentrations for 2.0 h at RT in 0.1 M Tris/HCl pH 7.4 containing 0.5M NaCl and 0.5% (w/v) Brij35. At the end of the incubation time, the plates are centrifuged (55g, 60 seconds) and

counted in a Wallac MicroBeta reader. IC_{50} values are calculated from percentage of displacement of radioligand binding to renin as a function of test compound concentration.

In animals deficient in salt, renin inhibitors bring about a reduction in blood pressure. Human renin may differ from the renin of other species. In order to test inhibitors of human renin, primates, e.g., marmosets (*Callithrix jacchus*) may be used, because human renin and primate renin are substantially homologous in the enzymatically active region. *Inter alia* the following *in vivo* tests may be used:

The test compounds are tested on normotensive marmosets of both sexes having a body weight of approximately 350 g that are conscious, allowed to move freely and in their normal cages. The blood pressure and heart rate are measured via a catheter in the descending aorta and recorded radiometrically. The endogenous release of renin is stimulated by the combination of a 1-week low-salt diet and a single intramuscular injection of furosemide (5-(aminosulfonyl)-4-chloro-2-[(2-furanylmethyl)amino]benzoic acid) (5 mg/kg). 16 h after the injection of furosemide the test compounds are administered either directly into the femoral artery using an injection cannula or, in the form of a suspension or solution, via an oesophageal tube into the stomach, and their action on the blood pressure and heart rate are evaluated. In the *in vivo* test described, the compounds of the present invention have hypotensive action at doses of from approximately 0.003 to approximately 1 mg/kg i.v. and at doses of from approximately 0.3 to approximately 100 mg/kg p.o.

Alternatively, renin inhibitors may be tested on male normotensive marmosets weighing 250 to 500 g that are conscious, allowed to move freely and in their normal cages. The blood pressure, and heart rate are measured via a catheter placed in the descending aorta and recorded radiometrically. Electrocardiogram are obtained by placing electrodes of transmitter in lead II. The endogenous release of renin is stimulated by two intramuscular injection of furosemide (5-(aminosulfonyl)-4-chloro-2-[(2-furanylmethyl)amino]benzoic acid) (10 mg/kg) 43 and 19 hours prior compound application. Test compounds are administered either directly into the femoral artery using an injection cannula or, in the form of a suspension or solution, via an oesophageal tube into the stomach, and their action on the blood pressure, heart rate and ECG are evaluated. In the *in vivo* test described, compounds of the present invention have hypotensive action at doses of from approximately 0.003 to approximately 0.3 mg/kg i.v. and at doses of from approximately 0.31 to approximately 30 mg/kg p.o.

The compounds of the present invention also have the property of regulating, especially reducing, intra-ocular pressure.

The extent of the reduction in intra-ocular pressure after administration of a pharmaceutical active ingredient of formula (I) according to the present invention can be determined, for example, in animals, for example rabbits or monkeys. Two typical experimental procedures that illustrate the present invention, but are not limited to in any way, are described hereinafter.

The *in vivo* test on a rabbit of the "Fauve de Bourgogne" type to determine the intra-ocular-pressure-reducing activity of topically applied compositions can be designed, for example, as follows: The intra-ocular pressure (IOP) is measured using an aplanation tonometer both before the experiment and at regular intervals of time. After a local anaesthetic has been administered, the suitably formulated test compound is applied topically in a precisely defined concentration (e.g. 0.000001-5% by weight) to one eye of the animal in question. The contralateral eye is treated, for example, with physiological saline. The measured values thus obtained are evaluated statistically.

The *in vivo* tests on monkeys of the species *Macaca Fascicularis* to determine the intra-ocular-pressure-reducing activity of topically applied compositions can be carried out, e.g., as follows: The suitably formulated test compound is applied in a precisely defined concentration (e.g. 0.000001-5% by weight) to one eye of each monkey. The other eye of the monkey is treated correspondingly, for example with physiological saline. Before the start of the test the animals are anaesthetised with intramuscular injections of, for example, ketamine. At regular intervals of time, the intra-ocular pressure (IOP) is measured. The test is carried out and evaluated in accordance with the rules of "good laboratory practice" (GLP).


Illustrative of the invention, the compound of Example 1 demonstrates inhibition of renin activity with an IC_{50} value of about 70 nM in the FRET assay.

The following Examples are intended to illustrate the invention and are not to be construed as being limitations thereon. If not mentioned otherwise, all evaporations are performed under reduced pressure, preferably between about 10 and 100 mmHg (= 20-133 mbar). The structure of final products, intermediates and starting materials is confirmed by standard analytical methods, e.g., microanalysis, melting point (m.p.) and spectroscopic

characteristics, e.g., MS, LC/MS, IR, NMR. In general, abbreviations used are those conventional in the art.

Example 1

(3R*,4R*)-4-Biphenyl-3-yl-3-(naphthalen-2-ylmethoxy)-piperidine hydrochloride



A. 4-Biphenyl-3-yl-3,6-dihydro-2H-pyridine-1-carboxylic acid *t*-butyl ester

A stirred mixture of 4-trifluoromethanesulfonyloxy-3,6-dihydro-2H-pyridine-1-carboxylic acid *t*-butyl ester (3.98 g, 12 mmol), 3-biphenylboronic acid (2.77 g, 14 mmol), LiCl (1.53 g, 36 mmol), dimethoxyethane (35 mL), tetrakis-(triphenylphosphin)-palladium (0.81 g, 0.7 mmol) and Na₂CO₃ (2M solution, 14 mL, 28 mmol) is heated at reflux under argon for 14 h. The mixture is cooled to RT, dimethoxyethane is removed *in vacuo* and the residue is diluted with aqueous 2N Na₂CO₃ solution containing a few mL of concentrated NH₄OH. The aqueous layer is extracted three times with CH₂Cl₂. The combined organic extracts are dried (Na₂SO₄) and evaporated *in vacuo*. Flash chromatography of the dark residue (SiO₂, hexane/ethyl acetate) affords 4-biphenyl-3-yl-3,6-dihydro-2H-pyridine-1-carboxylic acid *t*-butyl ester as a colourless oil: MS (after treatment with trifluoroacetic acid) 236.3 [M+H-C₅H₈O₂]⁺; R_f 0.40 (hexane/ethyl acetate 8:2).

B. (3R*,4R*)-4-Biphenyl-3-yl-3-hydroxy-piperidine-1-carboxylic acid *t*-butyl ester

To a stirred mixture of the title A compound, 4-biphenyl-3-yl-3,6-dihydro-2H-pyridine-1-carboxylic acid *t*-butyl ester (2.57 g, 7.66 mmol) and 6 mL of dry THF is added dropwise under argon at 5°C a 1M borane-tetrahydrofuran complex solution (11 mL, 11 mmol) within 15 min. The reaction mixture is stirred for 10 min. at 5°C and for 1 h at RT. NaBO₃ (3.38 g, 22 mmol) is then added in several portions followed by H₂O (10 mL). After 2 h at RT, the reaction mixture is extracted twice with ethyl acetate. The combined organic extracts are dried (Na₂SO₄) and evaporated *in vacuo*. Flash chromatography of the residue (SiO₂, hexane/ethyl acetate) affords (3R*,4R*)-4-biphenyl-3-yl-3-hydroxy-piperidine-1-carboxylic

acid *t*-butyl ester as a white amorphous solid: MS (after treatment with trifluoroacetic acid) 254.3 $[M+H-C_5H_6O_2]^+$; R_f 0.10 (hexane/ethyl acetate 8:2).

C. (3R*,4R*)-4-Biphenyl-3-yl-3-(naphthalen-2-ylmethoxy)-piperidine-1-carboxylic acid *t*-butyl ester


A suspension of the title B compound, (3R*,4R*)-4-biphenyl-3-yl-3-hydroxy-piperidine-1-carboxylic acid *t*-butyl ester (318 mg, 0.9 mmol) and sodium hydride (60% dispersion in mineral oil, 72 mg, 1.8 mmol) in 8 mL of dry DMA is shaken for 10 min. at 50°C. After cooling to RT, 2-(bromomethyl)naphthalene (298 mg, 1.35 mmol) is added and the mixture shaken for 16 h at 50°C. After addition of H₂O, the aqueous layer is extracted twice with ethyl acetate. The combined organic extracts are dried (Na₂SO₄) and evaporated *in vacuo*. Flash chromatography of the residue (SiO₂, hexane/ethyl acetate) affords (3R*,4R*)-4-biphenyl-3-yl-3-(naphthalen-2-ylmethoxy)-piperidine-1-carboxylic acid tert-butyl ester as a yellowish viscous oil: MS (LC/MS) 494.2 $[M+H]^+$; R_f 0.30 (hexane/ethyl acetate 8:2).

D. (3R*,4R*)-4-Biphenyl-3-yl-3-(naphthalen-2-ylmethoxy)-piperidine hydrochloride


A mixture of the title C compound, (3R*,4R*)-4-biphenyl-3-yl-3-(naphthalen-2-ylmethoxy)-piperidine-1-carboxylic acid *t*-butyl ester (54 mg, 0.11 mmol) and HCl (5M in 2-propanol, 1 mL, 5 mmol) is stirred for several h at RT (until complete consumption of the starting material). The reaction mixture is evaporated *in vacuo* to afford (3R*,4R*)-4-biphenyl-3-yl-3-(naphthalen-2-ylmethoxy)-piperidine hydrochloride as a slightly beige foam: MS (LC/MS) 394.1 $[M+H]^+$; R_f 0.23 (DCM/MeOH 9:1).

Example 2


(3R*,4R*)-4-Biphenyl-3-yl-3-(biphenyl-4-ylmethoxy)-piperidine hydrochloride




The title compound is prepared analogously as described in Example 1: MS 420.6 $[M+H]^+$; retention time 2.77 min (HPLC, Nucleosil C18; 30→100% CH₃CN in H₂O within 3 min).

Example 3**(3R*,4R*)-3-Benzoyloxy-4-biphenyl-3-yl-piperidine hydrochloride**

The title compound is prepared analogously as described in Example 1: MS (LC/MS) 344.2 [M+H]⁺; R_f 0.20 (DCM/MeOH 9:1).

Example 4**(3R*,4R*)-4-Biphenyl-3-yl-3-(4-bromo-benzyloxy)-piperidine hydrochloride**


The title compound is prepared analogously as described in Example 1: MS (LC/MS) 422.1/424.1 [M+H]⁺; R_f 0.22 (DCM/MeOH 9:1).

Example 5**(3R*,4R*)-4-Biphenyl-3-yl-3-(4-trifluoromethoxy-benzyloxy)-piperidine hydrochloride**

The title compound is prepared analogously as described in Example 1: MS (LC/MS) 428.2 $[M+H]^+$; R_f 0.19 (DCM/MeOH 9:1).

Example 6


(3R*,4R*)-4-Biphenyl-3-yl-3-(3-phenoxy-benzyloxy)-piperidine hydrochloride




The title compound is prepared analogously as described in Example 1: MS (LC/MS) 436.2 $[M+H]^+$; R_f 0.26 (DCM/MeOH 9:1).

Example 7


(3R*,4R*)-4-Dibenzofuran-4-yl-3-(naphthalen-2-ylmethoxy)-piperidine hydrochloride



A mixture of (3R*,4R*)-4-dibenzofuran-4-yl-3-(naphthalen-2-ylmethoxy)-piperidine-1-carboxylic acid *t*-butyl ester (prepared analogously as described in Example 1; 191 mg, 0.38 mmol) and HCl (5 M in 2-propanol, 2 mL, 10 mmol) is stirred for 1 h at RT. The solvent is removed *in vacuo* and the residue purified by preparative HPLC (CH₃CN/H₂O). A saturated aqueous NaHCO₃ solution is added to the combined pure fractions, CH₃CN is evaporated *in vacuo*, and the remaining aqueous layer extracted twice with DCM. The combined organic phases are dried (Na₂SO₄) and evaporated. The resulting gum is treated with HCl (4M in 1,4-dioxane, 1 mL) and stirred for 30 min at RT. Evaporation of the solvent *in vacuo* yields (3R*,4R*)-4-dibenzofuran-4-yl-3-(naphthalen-2-ylmethoxy)-piperidine hydrochloride as a white solid: MS 408.6 $[M+H]^+$; R_f 0.20 (DCM/MeOH 9:1).

Example 8**(3R*,4R*)-3-(Biphenyl-4-ylmethoxy)-4-dibenzofuran-4-yl-piperidine hydrochloride**


A mixture of (3R*,4R*)-3-(biphenyl-4-ylmethoxy)-4-dibenzofuran-4-yl-piperidine-1-carboxylic acid *t*-butyl ester (prepared analogously as described in Example 1; 218 mg, 0.41 mmol) and HCl (5M in 2-propanol, 2 mL, 10 mmol) is stirred for 1 h at RT. The solvent is removed *in vacuo*, the residue triturated with hot MeOH and the suspension filtered. The filter cake is washed with MeOH and ether and dried at 60°C *in vacuo* to afford (3R*,4R*)-3-(biphenyl-4-ylmethoxy)-4-dibenzofuran-4-yl-piperidine hydrochloride: MS 434.6 [M+H]⁺; R_f 0.23 (DCM/MeOH 9:1).

Example 9**(3R*,4R*)-3-Benzylloxy-4-dibenzofuran-4-yl-piperidine hydrochloride**

The title compound is prepared analogously as described in Example 8: MS 358.4 [M+H]⁺; R_f 0.19 (DCM/MeOH 9:1).

Example 10**(3R*,4R*)-4-Dibenzofuran-4-yl-3-(3-phenoxy-benzylloxy)-piperidine hydrochloride**


- 43 -



The title compound is prepared analogously as described in Example 8: MS 450.7 $[M+H]^+$; R_f 0.23 (DCM/MeOH 9:1).

Example 11


[3-((3R*,4R*)-4-Dibenzofuran-4-yl-piperidin-3-yloxymethyl)-phenyl]-phenyl-methanone hydrochloride



The title compound is prepared analogously as described in Example 8: MS 462.6 $[M+H]^+$; R_f 0.21 (DCM/MeOH 9:1).

Example 12

(3R*,4R*)-4-(4'-Methoxy-biphenyl-3-yl)-3-(naphthalen-2-ylmethoxy)-piperidine hydrochloride



A. (3-Hydroxy-phenyl)-3,6-dihydro-2H-pyridine-1-carboxylic acid *t*-butyl ester

The title compound is prepared analogously as described for the title A compound in Example 1 using 3-hydroxyphenylboronic acid: MS 274.3 [M-H]⁺; retention time 6.23 min (HPLC, Nucleosil C18; 5→100% CH₃CN in H₂O within 8 min).

B. (3R*,4R*)-3-Hydroxy-4-(4'-methoxy-biphenyl-3-yl)-piperidine-1-carboxylic acid *t*-butyl ester

A mixture of (3-hydroxy-phenyl)-3,6-dihydro-2H-pyridine-1-carboxylic acid *t*-butyl ester (1.0 g, 3.63 mmol) and triethylamine (1.02 mL, 7.26 mmol) in 100 mL CH₂Cl₂ is stirred for 10 min. at RT. After the addition of N-phenyltrifluoromethanesulfonimide (1.56 g, 4.36 mmol) and K₂CO₃, the reaction mixture is heated at reflux for 5 h. The mixture is cooled to RT, filtered and the solvent removed *in vacuo*. Filtration of the residue (SiO₂, hexane/ethyl acetate) affords crude 4-(3-trifluoromethanesulfonyloxy-phenyl)-3,6-dihydro-2H-pyridine-1-carboxylic acid *t*-butyl ester as a colourless oil. The crude product is dissolved in 12 mL dimethoxyethane. After the addition of 4-methoxyphenylboronic acid (669 mg, 4.4 mmol), LiCl (466 mg, 11 mmol), tetrakis-(triphenylphosphin)-palladium (0.25 g, 0.22 mmol) and Na₂CO₃ (2M solution, 4.4 mL, 8.8 mmol) the mixture is heated at reflux under argon for 12 h. The mixture is cooled to RT, dimethoxyethane is removed *in vacuo* and the residue is diluted with ethyl acetate. The organic layer is washed with aqueous 2M Na₂CO₃ sol., dried (Na₂SO₄) and evaporated *in vacuo*. Flash chromatography of the brown residue (SiO₂, hexane/ethyl acetate) affords 4-(4'-methoxy-biphenyl-3-yl)-3,6-dihydro-2H-pyridine-1-carboxylic acid *t*-butyl ester as a yellow oil.

To a stirred mixture of this compound (858 mg, 2.35 mmol) in 6 mL of dry THF is added dropwise under argon at 5°C a 1M borane-tetrahydrofuran complex solution (3.84 mL, 3.84 mmol) within 10 min. The reaction mixture is stirred for 10 min. at 5°C and for 1 h at RT. NaBO₃ (1.18 g, 7.68 mmol) is then added in several portions followed by H₂O (2.25 mL). After 4 h at RT, the reaction mixture is diluted with brine and extracted twice with ethyl acetate. The combined organic extracts are dried (Na₂SO₄) and evaporated *in vacuo*. Flash chromatography of the residue (SiO₂, hexane/ethyl acetate) affords (3R*,4R*)-3-hydroxy-4-(4'-methoxy-biphenyl-3-yl)-piperidine-1-carboxylic acid *t*-butyl ester as a white amorphous solid: MS 384.4 [M+H]⁺; retention time 7.01 min (HPLC, Nucleosil C18; 5→100% CH₃CN in H₂O within 8 min).

C. (3R*,4R*)-4-(4'-Methoxy-biphenyl-3-yl)-3-(naphthalen-2-ylmethoxy)-piperidine-1-carboxylic acid *t*-butyl ester


The title compound is prepared analogously as described for the title C compound in Example 1 from the title B compound, (3R*,4R*)-3-hydroxy-4-(4'-methoxy-biphenyl-3-yl)-piperidine-1-carboxylic acid *t*-butyl ester: MS 524.5 [M+H]⁺; retention time 8.83 min (HPLC, Nucleosil C18; 5→100% CH₃CN in H₂O within 8 min, then 100% CH₃CN for 2 min).

D. (3R*,4R*)-4-(4'-Methoxy-biphenyl-3-yl)-3-(naphthalen-2-ylmethoxy)-piperidine hydrochloride

The title compound is prepared analogously as described for the title D compound in Example 1 from (3R*,4R*)-4-(4'-methoxy-biphenyl-3-yl)-3-(naphthalen-2-ylmethoxy)-piperidine-1-carboxylic acid *t*-butyl ester : MS 424.5 [M+H]⁺; R_f 0.24 (DCM/MeOH 9:1).

Example 13

3'-[(3R*,4R*)-3-(Naphthalen-2-ylmethoxy)-piperidin-4-yl]-biphenyl-3-ol hydrochloride



A. 4-(3-Bromo-phenyl)-3,6-dihydro-2H-pyridine-1-carboxylic acid *t*-butyl ester

A stirred mixture of 4-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-3,6-dihydro-2H-pyridine-1-carboxylic acid *tert*-butyl ester (prepared as described in Tetrahedron Letters **2000**, 3705-3708; 6.52 g, 21.1 mmol), 1-bromo-3-iodobenzene (4.05 mL, 31.8 mmol), bis-(triphenylphosphine)-palladium(II) chloride (892 mg, 1.27 mmol), K₂CO₃ (8.7 g, 63 mmol) and 1,1'-bis(diphenylphosphino)ferrocene (704 mg, 1.27 mmol) in 120 mL dimethylformamide is heated at 80°C under argon for 14 h. The mixture is cooled to RT, dimethylformamide is removed *in vacuo*. After addition of brine, the aqueous layer is extracted three times with CH₂Cl₂. The combined organic extracts are dried (Na₂SO₄) and evaporated *in vacuo*. Flash chromatography of the residue (SiO₂, hexane/ethyl acetate) affords 4-(3-bromo-phenyl)-3,6-dihydro-2H-pyridine-1-carboxylic acid *t*-butyl ester as a colourless viscous resin: MS 338.2 / 340.2 [M+H]⁺; retention time 8.01 min (HPLC, Nucleosil C18; 5→100% CH₃CN in H₂O within 8 min, then 100% CH₃CN for 2 min).

B. (3R*,4R*)-4-(3-Bromo-phenyl)-3-hydroxy-piperidine-1-carboxylic acid *t*-butyl ester

The title compound is prepared analogously as described for the title B compound in Example 1 from the title A compound, 4-(3-bromo-phenyl)-3,6-dihydro-2H-pyridine-1-carboxylic acid *t*-butyl ester: MS 356.2 / 358.1 [M+H]⁺; retention time 6.59 min (HPLC, Nucleosil C18; 5→100% CH₃CN in H₂O within 8 min).

C. (3R*,4R*)-4-(3-Bromo-phenyl)-3-(naphthalen-2-ylmethoxy)-piperidine-1-carboxylic acid *t*-butyl ester

The title compound is prepared analogously as described for the title C compound in Example 1 from the title B compound, (3R*,4R*)-4-(3-bromo-phenyl)-3-hydroxy-piperidine-1-carboxylic acid *t*-butyl ester: MS 496.3 / 498.2 [M+H]⁺; retention time 8.87 min (HPLC, Nucleosil C18; 5→100% CH₃CN in H₂O within 8 min, then 100% CH₃CN for 2 min).

D. (3R*,4R*)-4-(3'-Hydroxy-biphenyl-3-yl)-3-(naphthalen-2-ylmethoxy)-piperidine-1-carboxylic acid *t*-butyl ester

A stirred mixture of (3R*,4R*)-4-(3-bromo-phenyl)-3-(naphthalen-2-ylmethoxy)-piperidine-1-carboxylic acid *t*-butyl ester (150 mg, 0.3 mmol), 3-hydroxyphenylboronic acid (48 mg, 0.35 mmol), dimethoxyethane (3 mL), H₂O (1 mL), tetrakis-(triphenylphosphin)-palladium (21 mg, 0.02 mmol) and Na₂CO₃ (47 mg, 0.45 mmol) is heated at reflux under argon for 16 h. The mixture is cooled to RT, dimethoxyethane is removed *in vacuo* and the residue is diluted with aqueous 2N Na₂CO₃ solution containing a few mL of concentrated NH₄OH. The aqueous layer is extracted three times with CH₂Cl₂. The combined organic extracts are dried (Na₂SO₄) and evaporated *in vacuo*. Flash chromatography of the dark residue (SiO₂, hexane/ethyl acetate) affords (3R*,4R*)-4-(3'-hydroxy-biphenyl-3-yl)-3-(naphthalen-2-ylmethoxy)-piperidine-1-carboxylic acid *t*-butyl ester as a yellowish resin: MS 508.4 [M-H]⁻; retention time 8.01 min (HPLC, Nucleosil C18; 5→100% CH₃CN in H₂O within 8 min, then 100% CH₃CN for 2 min).


E. 3'-[(3R*,4R*)-3-(Naphthalen-2-ylmethoxy)-piperidin-4-yl]-biphenyl-3-ol hydrochloride

A mixture of (3R*,4R*)-4-(3'-hydroxy-biphenyl-3-yl)-3-(naphthalen-2-ylmethoxy)-piperidine-1-carboxylic acid *t*-butyl ester (96.5 mg, 0.19 mmol) and HCl (5M in 2-propanol, 2 mL, 10 mmol) is stirred for 1 h at RT. The reaction mixture is evaporated *in vacuo* and the residue purified using preparative HPLC (H₂O / CH₃CN). The combined pure fractions are treated with solid K₂CO₃ and CH₃CN is removed *in vacuo*. The aqueous layer is extracted twice with

CH_2Cl_2 . The combined organic extracts are dried (Na_2SO_4) and evaporated *in vacuo*. The residue is stirred with HCl (4M in dioxane, 1 mL, 8 mmol) for 30 min. at RT. Evaporation of the solvent affords 3'-[(3R*,4R*)-3-(naphthalen-2-ylmethoxy)-piperidin-4-yl]-biphenyl-3-ol hydrochloride as a colourless white solid: MS 410.5 $[\text{M}+\text{H}]^+$; R_f 0.11 (DCM/MeOH 9:1).

Example 14


(3R*,4R*)-4-[3'-(2-Methoxy-ethoxy)-biphenyl-3-yl]-3-(naphthalen-2-ylmethoxy)-piperidine hydrochloride



A mixture of the title D compound in Example 13, (3R*,4R*)-4-(3'-hydroxy-biphenyl-3-yl)-3-(naphthalen-2-ylmethoxy)-piperidine-1-carboxylic acid t-butyl ester (80 mg, 0.16 mmol), 2-chloroethyl methyl ether (57 μL , 0.62 mmol), K_2CO_3 (29 mg, 0.21 mmol), KI (16.6 mg, 0.1 mmol) and dimethylacetamide (1.5 mL) is stirred under argon for 16 h at 75°C. After the addition of a second portion of 2-chloroethyl methyl ether (35 μL , 0.38 mmol) and K_2CO_3 (44 mg, 0.32 mmol) stirring is continued for 18 h at 75°C. The mixture is filtered and the filtrate evaporated *in vacuo*. HCl (5M in 2-propanol, 1 mL, 5 mmol) is added to the crude residue and the mixture is stirred for 45 min. at RT. The reaction mixture is evaporated *in vacuo* and the residue purified using preparative HPLC (H_2O / CH_3CN). The combined pure fractions are treated with solid K_2CO_3 and CH_3CN is removed *in vacuo*. The aqueous layer is extracted twice with CH_2Cl_2 . The combined organic extracts are dried (Na_2SO_4) and evaporated *in vacuo*. The residue is stirred with HCl (5M in 2-propanol, 1 mL, 5 mmol) for 30 min. at RT. Evaporation of the solvent affords (3R*,4R*)-4-[3'-(2-methoxy-ethoxy)-biphenyl-3-yl]-3-(naphthalen-2-ylmethoxy)-piperidine hydrochloride as a brownish amorphous solid: MS (LC/MS) 468.4 $[\text{M}+\text{H}]^+$; retention time 5.85 min (HPLC, Nucleosil C18; 5→100% CH_3CN in H_2O within 8 min).

Example 15


N-{3'-[[3R*,4R*]-3-(Naphthalen-2-ylmethoxy)-piperidin-4-yl]-biphenyl-3-yl}-acetamide hydrochloride



The title compound is prepared analogously as described in Example 13 using 3-acetamidobenzeneboronic acid: MS 451.5 [M+H]⁺; R_f 0.11 (DCM/MeOH 9:1).

Example 16

1-{3'-[[3R*,4R*]-3-(Naphthalen-2-ylmethoxy)-piperidin-4-yl]-biphenyl-4-yl}-ethanone hydrochloride




The title compound is prepared analogously as described in Example 13 using 4-acetylphenylboronic acid: MS 436.6 [M+H]⁺; R_f 0.24 (DCM/MeOH 9:1).

Example 17

1-{3'-[[3R*,4R*]-3-(Naphthalen-2-ylmethoxy)-piperidin-4-yl]-biphenyl-3-yl}-ethanone hydrochloride


- 49 -



The title compound is prepared analogously as described in Example 13 using 3-acetylphenylboronic acid: MS 436.6 $[M+H]^+$; R_f 0.24 (DCM/MeOH 9:1).

Example 18


3'-[(3R*,4R*)-3-(Naphthalen-2-ylmethoxy)-piperidin-4-yl]-biphenyl-3-carboxylic acid ethyl ester hydrochloride



The title compound is prepared analogously as described in Example 13 using 3-ethoxycarbonylphenylboronic acid: MS 466.5 $[M+H]^+$; R_f 0.28 (DCM/MeOH 9:1).

Example 19


(3R*,4R*)-4-(3'-Chloro-biphenyl-3-yl)-3-(naphthalen-2-ylmethoxy)-piperidine hydrochloride



The title compound is prepared analogously as described in Example 13 using 3-chlorophenylboronic acid: MS 428.4 [M+H]⁺; R_f 0.27 (DCM/MeOH 9:1).

Example 20


(3R*,4R*)-3-(Naphthalen-2-ylmethoxy)-4-(3'-trifluoromethyl-biphenyl-3-yl)-piperidine hydrochloride



The title compound is prepared analogously as described in Example 13 using 3-(trifluoromethyl)phenylboronic acid: MS 462.5 [M+H]⁺; R_f 0.25 (DCM/MeOH 9:1).

Example 21

(3R*,4R*)-4-(4'-Chloro-biphenyl-3-yl)-3-(naphthalen-2-ylmethoxy)-piperidine hydrochloride




The title compound is prepared analogously as described in Example 13 using 4-chlorophenylboronic acid: MS 428.4 [M+H]⁺; R_f 0.25 (DCM/MeOH 9:1).

Example 22

3'-[(3R*,4R*)-3-(Naphthalen-2-ylmethoxy)-piperidin-4-yl]-biphenyl-4-ol hydrochloride


- 51 -



The title compound is prepared analogously as described in Example 13 using 4-(*t*-butyldimethylsilyloxy)phenylboronic acid: MS 410.6 $[M+H]^+$; R_f 0.12 (DCM/MeOH 9:1).

Example 23

Dimethyl-3'-[(3R*,4R*)-3-(Naphthalen-2-ylmethoxy)-piperidin-4-yl]biphenyl-4-yl]-amine hydrochloride




The title compound is prepared analogously as described in Example 13 using 4-(N,N-dimethylamino)phenylboronic acid: MS 437.7 $[M+H]^+$; R_f 0.25 (DCM/MeOH 9:1).


Example 24

3'-[(3R*,4R*)-3-(Naphthalen-2-ylmethoxy)-piperidin-4-yl]-biphenyl-3-carboxylic acid hydrochloride

- 52 -




A stirred mixture of (3R*,4R*)-4-(3-bromo-phenyl)-3-(naphthalen-2-ylmethoxy)-piperidine-1-carboxylic acid *t*-butyl ester (300 mg, 0.6 mmol), 3-ethoxycarbonylphenyl boronic acid (136.8 mg, 0.71 mmol), dimethoxyethane (3 mL), H₂O (1 mL), tetrakis-(triphenylphosphin)-palladium (46.5 mg, 0.04 mmol) and Na₂CO₃ (96 mg, 0.91 mmol) is heated at reflux under argon for 16 h. The mixture is cooled to RT, dimethoxyethane is removed *in vacuo* and the residue is diluted with aqueous 2N Na₂CO₃ solution containing a few mL of concentrated NH₄OH. The aqueous layer is extracted three times with CH₂Cl₂. The combined organic extracts are dried (Na₂SO₄) and evaporated *in vacuo*. Flash chromatography of the dark residue (SiO₂, hexane/ethyl acetate) affords (3R*,4R*)-4-(3'-ethoxycarbonyl-biphenyl-3-yl)-3-(naphthalen-2-ylmethoxy)-piperidine-1-carboxylic acid *t*-butyl ester as a yellowish resin. The ester (138.7 mg, 0.245 mmol) is dissolved in MeOH (1 mL) and THF (1 mL), treated with aqueous NaOH (2 M, 0.5 mL, 1 mmol) and the mixture is stirred for 14 h at RT. The solvents are removed *in vacuo*, the residue is diluted with H₂O and the pH adjusted to 5 by the addition of 1N HCl. The aqueous phase is extracted twice with ethyl acetate. The combined organic extracts are dried (Na₂SO₄) and evaporated *in vacuo* to afford (3R*,4R*)-4-(3'-carboxy-biphenyl-3-yl)-3-(naphthalen-2-ylmethoxy)-piperidine-1-carboxylic acid *t*-butyl ester as a white amorphous solid. A mixture of this acid (110.1 mg, 0.20 mmol) and HCl (5M in 2-propanol, 2 mL, 10 mmol) is stirred for 1 h at RT. The reaction mixture is evaporated *in vacuo* and the residue purified using preparative HPLC (H₂O / CH₃CN). The combined pure fractions are treated with solid K₂CO₃ and CH₃CN is removed *in vacuo*. The aqueous layer is extracted twice with CH₂Cl₂. The combined organic extracts are dried (Na₂SO₄) and evaporated *in vacuo*. The residue is stirred with HCl (5M in 2-propanol, 1 mL, 5 mmol) for 2 h at RT. Evaporation of the solvent affords 3'-[(3R*,4R*)-3-(naphthalen-2-ylmethoxy)-piperidin-4-yl]-biphenyl-3-carboxylic acid hydrochloride as a white solid: MS 438.6 [M+H]⁺; retention time 5.23 min (HPLC, Nucleosil C18; 5→100% CH₃CN in H₂O within 8 min).

Example 25**3'-[(3R*,4R*)-3-(Naphthalen-2-ylmethoxy)-piperidin-4-yl]-biphenyl-4-carboxylic acid methylamide hydrochloride**

A stirred mixture of (3R*,4R*)-4-(3-bromo-phenyl)-3-(naphthalen-2-ylmethoxy)-piperidine-1-carboxylic acid *t*-butyl ester (173.8 mg, 0.35 mmol), 4-ethoxycarbonylphenyl boronic acid (135.8 mg, 0.70 mmol), dimethoxyethane (4.5 mL), H₂O (1.5 mL), tetrakis-(triphenylphosphin)-palladium (46.2 mg, 0.04 mmol) and Na₂CO₃ (111.3 mg, 1.05 mmol) is heated at reflux under argon for 14 h. The mixture is cooled to RT, dimethoxyethane is removed *in vacuo* and the residue is diluted with aqueous 2N Na₂CO₃ solution containing a few mL of concentrated NH₄OH. The aqueous layer is extracted three times with CH₂Cl₂. The combined organic extracts are dried (Na₂SO₄) and evaporated *in vacuo*. Flash chromatography of the dark residue (SiO₂, hexane/ethyl acetate) affords (3R*,4R*)-4-(4'-ethoxycarbonyl-biphenyl-3-yl)-3-(naphthalen-2-ylmethoxy)-piperidine-1-carboxylic acid *t*-butyl ester as a yellowish resin. A stirred mixture of this ester and CH₃NH₂ (8M in EtOH, 5 mL, 40 mmol) is heated at 80°C for 24 h in a pressure bottle. The reaction mixture is evaporated *in vacuo* to afford (3R*,4R*)-4-(4'-methylcarbamoyl-biphenyl-3-yl)-3-(naphthalen-2-ylmethoxy)-piperidine-1-carboxylic acid *t*-butyl ester as a viscous yellowish oil. A mixture of this compound (62.4 mg, 0.11 mmol) and HCl (5M in 2-propanol, 3 mL, 15 mmol) is stirred for 1 h at RT. The reaction mixture is evaporated *in vacuo* to afford 3'-[(3R*,4R*)-3-(naphthalen-2-ylmethoxy)-piperidin-4-yl]-biphenyl-4-carboxylic acid methylamide hydrochloride as a white solid. MS 451.7 [M+H]⁺; R_f 0.13 (DCM/MeOH 9:1).

Example 26**4-{3'-[(3R*,4R*)-3-(Naphthalen-2-ylmethoxy)-piperidin-4-yl]-phenyl}-pyridine hydrochloride**


- 54 -



The title compound is prepared analogously as described in Example 13 using pyridine-4-boronic acid: MS 395.6 $[M+H]^+$; retention time 3.76 min (HPLC, Nucleosil C18; 5→100% CH_3CN in H_2O within 8 min).

Example 27

(3R*,4R*)-3-(Naphthalen-2-ylmethoxy)-4-(3-thiophen-3-yl-phenyl)-piperidine hydrochloride




The title compound is prepared analogously as described in Example 13 using thiophene-3-boronic acid: MS 400.6 $[M+H]^+$; retention time 5.88 min (HPLC, Nucleosil C18; 5→100% CH_3CN in H_2O within 8 min).

Example 28

3-{3-[(3R*,4R*)-3-(Naphthalen-2-ylmethoxy)-piperidin-4-yl]-phenyl}-pyridine hydrochloride


- 55 -



The title compound is prepared analogously as described in Example 13 using pyridine-3-boronic acid: MS 395.5 $[M+H]^+$; retention time 3.87 min (HPLC, Nucleosil C18; 5→100% CH_3CN in H_2O within 8 min).

Example 29

2-{3-[(3R*,4R*)-3-(Naphthalen-2-ylmethoxy)-piperidin-4-yl]-phenyl}-1H-indole hydrochloride




The title compound is prepared analogously as described in Example 13 using 1-(*t*-butoxycarbonyl)indole-2-boronic acid: MS 433.6 $[M+H]^+$; retention time 6.09 min (HPLC, Nucleosil C18; 5→100% CH_3CN in H_2O within 8 min).

Example 30

5-{3-[(3R*,4R*)-3-(Naphthalen-2-ylmethoxy)-piperidin-4-yl]-phenyl}-pyrimidine hydrochloride


- 56 -



The title compound is prepared analogously as described in Example 13 using pyrimidine-5-boronic acid: MS 396.6 $[M+H]^+$; retention time 4.47 min (HPLC, Nucleosil C18; 5→100% CH_3CN in H_2O within 8 min).

Example 31

(3R,4R)-4-Biphenyl-3-yl-3-[2-(3-methoxy-propoxy)-4-methyl-benzyloxy]-piperidine



A. 4-Biphenyl-3-yl-3,6-dihydro-2H-pyridine-1-carboxylic acid ethyl ester

The title compound is prepared analogously as described for the title A compound in Example 1.

B. (3S,4S)-4-Biphenyl-3-yl-3,4-dihydroxy-piperidine-1-carboxylic acid ethyl ester

To a stirred solution of AD-mix- α (20.13 g, 18.3 mmol) in $tBuOH$ (120 mL) and H_2O (120 mL) is added methanesulfonamid (1.92 g, 19.6 mmol). The reaction mixture is cooled to $0^\circ C$ followed by addition of a solution of the title A compound, 4-biphenyl-3-yl-3,6-dihydro-2H-pyridine-1-carboxylic acid ethyl ester (4.1 g, 13.07 mmol) in $tBuOH$ (10 mL) and H_2O (10 mL). The reaction is stirred at $0^\circ C$ for 30 min and allowed to stir over the week end at room temperature. To the reaction mixture is added 18 g of Na_2SO_3 followed by stirring for 1 h before the addition of methylene chloride. The layers are separated and the aqueous one extracted 3 times with methylene chloride. The combined organic extracts are washed with a

2N sodium hydroxide solution, dried over anhydrous sodium sulfate and concentrated under reduced pressure. Purification by flash column chromatography ($\text{CH}_2\text{Cl}_2/\text{MeOH}$: 95/5) affords the desired product of (3S,4S)-4-biphenyl-3-yl-3,4-dihydroxy-piperidine-1-carboxylic acid ethyl ester. TLC, Rf ($\text{CH}_2\text{Cl}_2/\text{MeOH}$: 95/5) = 0.69. MS 339.9 [M-H]. Rt (HPLC, Nucleosil C18, 10:90-100:0 $\text{CH}_3\text{CN}/\text{H}_2\text{O}$ + 0.1% TFA within 5 min, then 100% CH_3CN + 0.1% TFA): 5.26 min.

C. (3R,4R)-4-Biphenyl-3-yl-3-hydroxy-piperidine-1-carboxylic acid ethyl ester


To a vigorously stirred solution of the title B compound, (3S,4S)-4-biphenyl-3-yl-3,4-dihydroxy-piperidine-1-carboxylic acid ethyl ester (4.2 g, 12.05 mmol) in EtOH (250 mL) is added 6.5 g Raney-Nickel. The stirred suspension is heated to reflux for 4 h, then allowed to cool to room temperature, filtered over a celite pad and rinsed with EtOH. The solvent is removed under reduced pressure and purification by flash column chromatography (hexane/EtOAc 4/1 to 1/2) affords the desired product: (3R,4R)-4-Biphenyl-3-yl-3-hydroxy-piperidine-1-carboxylic acid ethyl ester. TLC, Rf (Hexane/EtOAc 2/1) = 0.65. MS 326.1 [M+H]. Rt (HPLC, Nucleosil C18, 10:90-100:0 $\text{CH}_3\text{CN}/\text{H}_2\text{O}$ + 0.1% TFA within 5 min, then 100% CH_3CN + 0.1% TFA): 5.62 min.

D. (3R,4R)-4-Biphenyl-3-yl-3-[2-(3-methoxy-propoxy)-4-methyl-benzyloxy]-piperidine-1-carboxylic acid ethyl ester

A suspension of the title C compound, (3R,4R)-4-biphenyl-3-yl-3-hydroxy-piperidine-1-carboxylic acid ethyl ester (150 mg, 0.45 mmol) and sodium hydride (60% dispersion in mineral oil, 30 mg, 0.68 mmol) in 10 mL of dry DMF is stirred for 10 min at 60°C under argon. After cooling to room temperature, 1-bromomethyl-2-(3-methoxy-propoxy)-4-methyl-benzene (252 mg, 0.9 mmol) is added and the mixture further stirred for 3 h at 60°C. After addition of H_2O , the aqueous layer is extracted twice with ethyl acetate and the combined organic extracts are dried over anhydrous sodium sulfate and concentrated under reduced pressure. The residue is purified by preparative HPLC (Macherey-Nagel 250/4 nucleosil 100-10 C18 column, 20:80 CH_3CN + 0.1% TFA over 11 min, then 100% CH_3CN + 0.1% TFA for 5.5 min, 40 mL/min). To the combined fractions containing the desired product is added 1M aqueous sodium carbonate solution and the acetonitrile is removed under reduced pressure. The resulting aqueous suspension is extracted twice with ethyl acetate. The combined organic layers are dried over anhydrous sodium sulfate, filtered and concentrated under reduced pressure to afford the title compound. MS 535.2 [M+18]; Rt (HPLC, Nucleosil C18, 10:90-100:0 $\text{CH}_3\text{CN}/\text{H}_2\text{O}$ + 0.1% TFA within 5 min, then 100% CH_3CN + 0.1% TFA): 7.29 min.

E. (3R,4R)-4-Biphenyl-3-yl-3-[2-(3-methoxy-propoxy)-4-methyl-benzyloxy]-piperidine

To a solution of the title D compound, (3R,4R)-4-biphenyl-3-yl-3-[2-(3-methoxy-propoxy)-4-methyl-benzyloxy]-piperidine-1-carboxylic acid ethyl ester (50 mg, 0.95 mmol) in 2 mL of absolute ethanol is added 2 mL of a 10% aqueous solution of sodium hydroxide. The resulting mixture is heated at 170°C for 45 min in a microwave. The solvent is removed under reduced pressure and the resulting aqueous suspension is extracted twice with methylene chloride. The combined organic layers are dried over anhydrous sodium sulfate, filtered and concentrated under reduced pressure. The residue is purified by preparative HPLC (Macherey-Nagel 250/4 nucleosil 100-10 C18 column, 20:80 CH₃CN + 0.1% TFA over 11 min, then 100% CH₃CN + 0.1% TFA for 5.5 min, 40 mL/min). To the combined fractions containing the desired product is added 1M aqueous sodium carbonate solution and the acetonitrile is removed under reduced pressure. The resulting aqueous suspension is extracted twice with ethyl acetate. The combined organic layers are dried over anhydrous sodium sulfate, filtered and concentrated under reduced pressure affording the title compound as a colorless oil. A solution of the compound in acetonitrile is lyophilized. MS 446.1 [M+H]. Rt (HPLC, Nucleosil C18, 10:90-100:0 CH₃CN/H₂O + 0.1% TFA within 5 min, then 100% CH₃CN + 0.1% TFA): 5.48 min.


Example 32**(3R,4R)-4-Biphenyl-3-yl-3-[2-(3-methoxy-propoxy)-benzyloxy]-piperidine trifluoroacetic acid**

The title compound is prepared analogously as described in Example 31, except that the product is purified by HPLC, and the combined fractions containing the desired product are concentrated under reduced pressure and a solution of the compound in acetonitrile is lyophilized to afford (3R,4R)-4-biphenyl-3-yl-3-[2-(3-methoxy-propoxy)-benzyloxy]-piperidine

trifluoroacetic acid: MS 432.2 [M+H]; Rt (HPLC, Nucleosil C18, 10:90-100:0 CH₃CN/H₂O + 0.1% TFA within 5 min, then 100% CH₃CN + 0.1% TFA): 5.53 min.

Example 33


(3R,4R)-4-Biphenyl-3-yl-3-[4-methoxy-3-(3-methoxy-propoxy)-benzyloxy]-piperidine



The title compound is prepared analogously as described in Example 31. MS 462.0 [M+H].
Rt (HPLC, Nucleosil C18, 10:90-100:0 CH₃CN/H₂O + 0.1% TFA within 5 min, then 100% CH₃CN + 0.1% TFA): 5.13 min.

Example 34


(3R,4R)-4-Biphenyl-3-yl-3-[3-(4-methoxy-butyl)-benzyloxy]-piperidine



The title compound is prepared analogously as described in Example 31. MS 429.9 [M+H].
Rt (HPLC, Nucleosil C18, 10:90-100:0 CH₃CN/H₂O + 0.1% TFA within 5 min, then 100% CH₃CN + 0.1% TFA): 5.64 min.

Example 35


(3R,4R)-4-Biphenyl-3-yl-3-[3-(3-methoxy-propyl)-benzyloxy]-piperidine



The title compound is prepared analogously as described in Example 31. MS 416.1 [M+H].
Rt (HPLC, Nucleosil C18, 10:90-100:0 CH₃CN/H₂O + 0.1% TFA within 5 min, then 100% CH₃CN + 0.1% TFA): 5.44 min.

Example 36


(3R,4R)-4-Biphenyl-3-yl-3-[3-(2-methoxy-ethyl)-benzyloxy]-piperidine



The title compound is prepared analogously as described in Example 31. MS 402.0 [M+H].
Rt (HPLC, Nucleosil C18, 10:90-100:0 CH₃CN/H₂O + 0.1% TFA within 5 min, then 100% CH₃CN + 0.1% TFA): 5.24 min.

Example 37


(3R,4R)-4-Biphenyl-3-yl-3-[3-(3-methoxy-propyl)-4-methyl-benzyloxy]-piperidine



The title compound is prepared analogously as described in Example 31. MS 429.9 [M+H].
Rt (HPLC, Nucleosil C18, 10:90-100:0 CH₃CN/H₂O + 0.1% TFA within 5 min, then 100% CH₃CN + 0.1% TFA) 5.56 min.

Example 38


(3R,4R)-4-Biphenyl-3-yl-3-[3-(4-methoxy-butyl)-4-methyl-benzyloxy]-piperidine




The title compound is prepared analogously as described in Example 31. MS 444.0 [M+H].
Rt (HPLC, Nucleosil C18, 10:90-100:0 CH₃CN/H₂O + 0.1% TFA within 5 min, then 100% CH₃CN + 0.1% TFA): 5.65 min.

Example 39


(3R,4R)-4-Biphenyl-3-yl-3-[2-(3-methoxy-propyl)-benzyloxy]-piperidine trifluoroacetic acid



The title compound is prepared analogously as described in Example 31, except that the product is purified by HPLC, and the combined fractions containing the desired product are concentrated under reduced pressure and a solution of the compound in acetonitrile is lyophilized to afford (3R,4R)-4-biphenyl-3-yl-3-[2-(3-methoxy-propyl)-benzyloxy]-piperidine trifluoroacetic acid: MS 416.2 [M+H]; Rt (HPLC, Nucleosil C18, 10:90-100:0 CH₃CN/H₂O + 0.1% TFA within 5 min, then 100% CH₃CN + 0.1% TFA): 5.56 min.


Example 40**(3R,4R)-3-(1-Benzyl-cyclopropylmethoxy)-4-biphenyl-3-yl-piperidine trifluoroacetic acid**

The title compound is prepared analogously as described in Example 31, except that the product is purified by HPLC, and the combined fractions containing the desired product are concentrated under reduced pressure and a solution of the compound in acetonitrile is lyophilized to afford: (3R,4R)-3-(1-benzyl-cyclopropylmethoxy)-4-biphenyl-3-yl-piperidine trifluoroacetic acid: MS 397.8 [M+H]; Rt (HPLC, Nucleosil C18, 10:90-100:0 CH₃CN/H₂O + 0.1% TFA within 5 min, then 100% CH₃CN + 0.1% TFA): 5.67 min.

Example 41**(3R,4R)-4-Biphenyl-3-yl-3-{1-[2-(3-methoxy-propyl)-benzyl]-cyclopropylmethoxy}-piperidine hydrochloride**

The free base of the title compound is prepared analogously as described in Example 31. The compound is then treated with HCl (4M in 1,4-dioxane, 2 eq.) and stirred for 30 min at RT. Evaporation of the solvent *in vacuum* yields (3R,4R)-4-biphenyl-3-yl-3-{1-[2-(3-methoxy-propyl)-benzyl]-cyclopropylmethoxy}-piperidine hydrochloride: MS 470.1 [M+H]; Rt (HPLC, Nucleosil C18, 10:90-100:0 CH₃CN/H₂O + 0.1% TFA within 5 min, then 100% CH₃CN + 0.1% TFA): 5.88 min.

Example 42

1-Bromomethyl-2-(3-methoxy-propoxy)-4-methyl-benzene:**A. Toluene-4-sulfonic acid 3-methoxy-propyl ester**

A solution of 3-methoxy-1-propanol (6 g, 65.24 mmol), triethylamine (27.5 mL, 195.7 mmol) and tosylchloride (18.85 g, 97.9 mmol) in methylene chloride (50 mL) is stirred for 3 h at room temperature under nitrogen. The mixture is poured into H₂O, and the aqueous layer extracted twice with methylene chloride. The combined organic extracts are dried over anhydrous sodium sulfate and concentrated under reduced pressure. The residue is purified by flash column chromatography on silica gel (hexane/EtOAc 4/1) to afford the title compound as an orange oil: MS 261.9 [M+18]; Rt (HPLC, Nucleosil C18, 10:90-100:0 CH₃CN/H₂O + 0.1% TFA within 5 min, then 100% CH₃CN + 0.1% TFA): 5.21 min.

B. 2-(3-Methoxy-propoxy)-4-methyl-benzaldehyde

A solution of 2-hydroxy-4-methylbenzaldehyde (3 g, 21.6 mmol), potassium carbonate (14.9 g, 107.8 mmol) and the title A compound, toluene-4-sulfonic acid 3-methoxy-propyl ester (8.07 g, 32.4 mmol) in DMF (50 mL) is stirred at 50°C for 3 h. The solvent is concentrated under reduced pressure, H₂O is added, and the aqueous layer extracted twice with ethyl acetate. The combined organic extracts are dried over anhydrous sodium sulfate and concentrated under reduced pressure to afford the title compound together with 30% of remaining tosylate: Rt (HPLC, Nucleosil C18, 10:90-100:0 CH₃CN/H₂O + 0.1% TFA within 5 min, then 100% CH₃CN + 0.1% TFA): 5.25, 5.39 min. The mixture is further used as is in the next step.

C. [2-(3-Methoxy-propoxy)-4-methyl-phenyl]-methanol

To a solution of the title B compound, 2-(3-methoxy-propoxy)-4-methyl-benzaldehyde (3.7 g, 17.41 mmol) in 30 mL of methanol is added portion wise sodium borohydride (1.03 g, 26.12 mmol). The reaction mixture is stirred for 30 min at room temperature. Sodium borohydride (1.03 g, 26.12 mmol) is added and the mixture further stirred overnight to complete the reaction. After addition of H₂O, the aqueous layer is extracted twice with ethyl acetate. The combined organic extracts are dried over anhydrous sodium sulfate and concentrated under


reduced pressure. The residue is purified by flash column chromatography on silica gel (hexane/EtOAc 4/1 to 1/2) to afford the title compound as a colorless oil: TLC, R_f (hexane/AcOEt 2/1) = 0.2. Rt (HPLC, Nucleosil C18, 10:90-100:0 CH₃CN/H₂O + 0.1% TFA within 5 min, then 100% CH₃CN + 0.1% TFA): 4.88 min.

D. 1-Bromomethyl-2-(3-methoxy-propoxy)-benzene

To a solution of the title C compound, [2-(3-methoxy-propoxy)-4-methyl-phenyl]-methanol (300 mg, 1.4 mmol) in 5 mL of chloroform is added trimethylbromosilane (0.28 mL, 2.1 mmol), with stirring at room temperature. After 2 h the solvent is evaporated off and the crude residue is purified by flash column chromatography on silica gel (hexane/EtOAc 1/1) to afford the title compound as a colorless oil (0.4 g). Rt (HPLC, Nucleosil C18, 10:90-100:0 CH₃CN/H₂O + 0.1% TFA within 5 min, then 100% CH₃CN + 0.1% TFA): 4.70 min.

Example 43

1-Bromomethyl-2-(3-methoxy-propoxy)-benzene



A. 2-(3-Methoxy-propoxy)-benzaldehyde


A solution of 2-hydroxy-benzaldehyde (1.5 g, 12.16 mmol), potassium carbonate (8.4 g, 60.8 mmol) and toluene-4-sulfonic acid 3-methoxy-propyl ester (14.85 g, 60.8 mmol) in DMF (30 mL) is stirred at 50°C for 3 h. The solvent is concentrated under reduced pressure, H₂O is added, and the aqueous layer extracted twice with ethyl acetate. The combined organic extracts are dried over anhydrous sodium sulfate and concentrated under reduced pressure. The residue is purified by flash column chromatography on silica gel (hexane/EtOAc 4/1 to 1/2) to afford the title compound together with 50% of remaining tosylate: MS 195.0 [M+H]. Rt (HPLC, Nucleosil C18, 10:90-100:0 CH₃CN/H₂O + 0.1% TFA within 5 min, then 100% CH₃CN + 0.1% TFA): 5.08 min. The mixture is further used as is in the next step.

B. [2-(3-Methoxy-propoxy)-phenyl]-methanol

The title compound is prepared analogously as described for the title C compound in Example 42: TLC, R_f (hexane/AcOEt 2/1) = 0.46. Rt (HPLC, Nucleosil C18, 10:90-100:0 CH₃CN/H₂O + 0.1% TFA within 5 min, then 100% CH₃CN + 0.1% TFA): 4.58 min.

C. 1-Bromomethyl-2-(3-methoxy-propoxy)-benzene

The title compound is prepared analogously as described for the title D compound in Example 42: MS 275.7, 277.8 [M+18]. Rt (HPLC, Nucleosil C18, 10:90-100:0 CH₃CN/H₂O + 0.1% TFA within 5 min, then 100% CH₃CN + 0.1% TFA): 5.84 min.

Example 44**4-Bromomethyl-1-methoxy-2-(3-methoxy-propoxy)-benzene****A. 4-Methoxy-3-(3-methoxy-propoxy)-benzaldehyde**

To a solution of 3-hydroxy-4-methoxy-benzaldehyde (3.04 g, 20 mmol) and 3-methoxy-propanol (1.80 g, 20 mmol) in THF (150 mL) is added triphenylphosphine (5.24 g, 20 mmol) at room temperature under nitrogen atmosphere. To the stirring mixture is added diethyl azodicarboxylate (3.11 mL, 20 mmol) over 10 min at room temperature, and the resulting solution is further stirred over 20 h. THF is removed under reduced pressure and the remaining residue is purified by flash chromatography on silica gel (hexane/EtOAc 2/1) to afford the title compound: TLC, Rf (hexane/EtOAc 2/1) = 0.2.


B. [4-Methoxy-3-(3-methoxy-propoxy)-phenyl]-methanol

To a solution of the title A compound, 4-methoxy-3-(3-methoxy-propoxy)-benzaldehyde (2.02 g, 8.9 mmol) in 60 mL of THF is added portion wise sodium borohydride (0.74 g, 18.73 mmol) at 0°C. The reaction mixture is stirred for 6 hours at room temperature. After addition of H₂O, the aqueous layer is extracted twice with ethyl acetate. The combined organic extracts are dried over anhydrous sodium sulfate and concentrated under reduced pressure to afford the title compound as a colorless oil: TLC, Rf (hexane/AcOEt 1/1) = 0.34.

C. 4-Bromomethyl-1-methoxy-2-(3-methoxy-propoxy)-benzene

The title compound is prepared analogously as described for the title D compound in Example 42: TLC, Rf (hexane/AcOEt 1/1) = 0.57. Rt (HPLC, Nucleosil C18, 10:90-100:0 CH₃CN/H₂O + 0.1% TFA within 5 min, then 100% CH₃CN + 0.1% TFA): 4.89 min.

Example 45**1-Bromomethyl-3-(4-methoxy-butyl)-benzene**



A. (3-Methoxy-propyl)-triphenyl-phosphonium bromide

A solution of PPh_3 (42.8 g, 163.2 mmol) and 1-bromo-3-methoxypropane (25 g, 163.3 mmol) in toluene (70 mL) is heated at 150°C in an autoclave for 44 h. After completion of the reaction, the mixture is filtered and the precipitate washed with toluene and dried under high vacuum for 4 h affording the title compound as a white powder: Rt (HPLC, Nucleosil C18, 10:90-100:0 $\text{CH}_3\text{CN}/\text{H}_2\text{O}$ + 0.1% TFA within 5 min, then 100% CH_3CN + 0.1% TFA): 5.06 min.

B. 1-Bromo-3-(4-methoxy-but-1-enyl)-benzene

To a stirred solution of NaHMDS (9.02 g, 49.2 mmol) in THF (50 mL) under nitrogen atmosphere is added drop wise at 0°C a THF solution (50 mL) of the title A compound, (3-methoxy-propyl)-triphenyl-phosphonium bromide (20.4 g, 49.2 mmol). The resulting mixture is stirred for 1 h at 0°C before the addition of a THF solution (50 mL) of m-bromobenzaldehyde (7 g, 37.8 mmol). The reaction mixture is further stirred for 2 h at room temperature and poured into a saturated NH_4Cl aqueous solution, the aqueous layer is extracted twice with EtOAc. The combined organic extracts are dried over anhydrous magnesium sulfate and concentrated under reduced pressure. The residue is taken up into ether and the triphenylphosphine oxide precipitate is filtered off through a pad of celite. The filtrate is concentrate and the residual material purified by flash column chromatography on silica gel (hexane/EtOAc 95/5) to afford the title compound (as a mixture Z and E stereoisomers) as a yellow oil: Rt (HPLC, Nucleosil C18, 10:90-100:0 $\text{CH}_3\text{CN}/\text{H}_2\text{O}$ + 0.1% TFA within 5 min, then 100% CH_3CN + 0.1% TFA): 6.30 min.

C. 1-Bromo-3-(4-methoxy-butyl)-benzene

A suspension of the title B compound, 1-bromo-3-(4-methoxy-but-1-enyl)-benzene (1.8 g, 7.46 mmol) and Pd/c 5% (0.36 g) in THF (20 mL) is shaken under an hydrogen atmosphere. After completion of the reaction, the mixture is filtered through a pad of celite, the solvent is evaporated under reduced pressure and the residue purified by flash chromatography on silica gel (hexane/EtOAc 98/2) to give the title compound as a yellow oil: MS 260.1, 261.9

[M+18]. Rt (HPLC, Nucleosil C18, 10:90-100:0 CH₃CN/H₂O + 0.1% TFA within 5 min, then 100% CH₃CN + 0.1% TFA): 6.45 min.

D. 3-(4-Methoxy-butyl)-benzaldehyde

To a stirred solution of the title C compound, 1-bromo-3-(4-methoxy-butyl)-benzene (11 g, 45.24 mmol) in 200 mL of THF is added drop wise *n*-butyl lithium (31.1 mL, 49.76 mmol, 1.6 M solution in hexane) over 30 min at -72°C. The reaction mixture is further stirred 5 min at -72°C before the addition of a THF solution (50 mL) of DMF (7.67 mL, 99.52 mmol) over 45 min. The reaction mixture is further stirred for 15 min at -72°C, 1 h at room and poured into aqueous 2 M HCl solution. The aqueous layer is extracted twice with ether and the combined organic extracts are dried over anhydrous sodium sulfate and concentrated under reduced pressure. The residue is purified by flash column chromatography on silica gel (hexane/EtOAc 9/1) to afford the title compound as a yellow oil: MS 210.0 [M+18]; Rt (HPLC, Nucleosil C18, 10:90-100:0 CH₃CN/H₂O + 0.1% TFA within 5 min, then 100% CH₃CN + 0.1% TFA): 5.35 min.

E. [3-(4-Methoxy-butyl)-phenyl]-methanol


To a solution of the title D compound, 3-(4-methoxy-butyl)-benzaldehyde (0.1 g, 0.52 mmol) and MeOH (0.063 mL, 1.56 mmol) in 2 mL of THF is added portion wise sodium borohydride (20 mg, 0.52 mmol). The reaction mixture is stirred for 1 hour at room and poured into aqueous 1 M HCl. The aqueous layer is extracted twice with ethyl acetate and the combined organic extracts are dried over anhydrous sodium sulfate and concentrated under reduced pressure. The residue is purified by flash column chromatography on silica gel (hexane/EtOAc 3/1) to afford the title compound as a colorless oil: TLC, Rf (hexane/AcOEt 2/1) = 0.34. MS 212.1 [M+18].

F. 1-Bromomethyl-3-(4-methoxy-butyl)-benzene

The title compound is prepared analogously as described for the title D compound in Example 42: TLC, Rf (hexane/AcOEt 1/1) = 0.8. MS 273.9, 276.0 [M+18]. Rt (HPLC, Nucleosil C18, 10:90-100:0 CH₃CN/H₂O + 0.1% TFA within 5 min, then 100% CH₃CN + 0.1% TFA): 6.10 min.

Example 46

1-Bromomethyl-3-(3-methoxy-propyl)-benzene

**A. (2-Methoxy-ethyl)-triphenyl-phosphonium bromide**

The title compound is prepared analogously as described for the title C compound in Example 45: Rt (HPLC, Nucleosil C18, 10:90-100:0 CH₃CN/H₂O + 0.1% TFA within 5 min, then 100% CH₃CN + 0.1% TFA): 5.09 min.

B. 1-Bromo-3-(3-methoxy-propenyl)-benzene

The title compound is prepared analogously as described for the title B compound in Example 45: Rt (HPLC, Nucleosil C18, 10:90-100:0 CH₃CN/H₂O + 0.1% TFA within 5 min, then 100% CH₃CN + 0.1% TFA): 5.97 min.

C. 1-Bromo-3-(3-methoxy-propyl)-benzene

The title compound is prepared analogously as described for the title C compound in Example 45: MS 226.9, 228.9 [M+H].

D. 3-(3-Methoxy-propyl)-benzaldehyde

The title compound is prepared analogously as described for the title D compound in Example 45: TLC, R_f (hexane/AcOEt 5/1) = 0.37. MS 195.9 [M+18].


E. [3-(3-Methoxy-propyl)-phenyl]-methanol

The title compound is prepared analogously as described for the title E compound in Example 45: MS 198 [M+18]; TLC, R_f (hexane/AcOEt 2/1) = 0.33.

F. 1-Bromomethyl-3-(3-methoxy-propyl)-benzene

The title compound is prepared analogously as described for the title F compound in Example 45: TLC, R_f (hexane/AcOEt 1/1) = 0.85; MS 261.9, 262.8 [M+18]; Rt (HPLC, Nucleosil C18, 10:90-100:0 CH₃CN/H₂O + 0.1% TFA within 5 min, then 100% CH₃CN + 0.1% TFA): 5.85 min.

Example 47**1-bromomethyl-3-(2-methoxy-ethyl)-benzene**

**A. Methoxymethyl-triphenyl-phosphonium bromide**

The title compound is prepared analogously as described for the title A compound in Example 45: Rt (HPLC, Nucleosil C18, 10:90-100:0 CH₃CN/H₂O + 0.1% TFA within 5 min, then 100% CH₃CN + 0.1% TFA): 4.83 min.

B. 1-Bromo-3-(2-methoxy-vinyl)-benzene

The title compound is prepared analogously as described for the title B compound in Example 45: TLC, R_f (hexane/AcOEt 9/1) = 0.65. Rt (HPLC, Nucleosil C18, 10:90-100:0 CH₃CN/H₂O + 0.1% TFA within 5 min, then 100% CH₃CN + 0.1% TFA): 6.01 and 6.09 min.

C. 1-Bromo-3-(2-methoxy-ethyl)-benzene

The title compound is prepared analogously as described for the title C compound in Example 45: TLC, R_f (hexane/AcOEt 9/1) = 0.6. Rt (HPLC, Nucleosil C18, 10:90-100:0 CH₃CN/H₂O + 0.1% TFA within 5 min, then 100% CH₃CN + 0.1% TFA): 5.75 min.

D. 3-(2-Methoxy-ethyl)-benzaldehyde

The title compound is prepared analogously as described for the title D compound in Example 45: TLC, R_f (hexane/AcOEt 5/1) = 0.37. MS 182.1 [M+18].


E. [3-(2-Methoxy-ethyl)-phenyl]-methanol

The title compound is prepared analogously as described for the title E compound in Example 45: MS 183.9 [M+18]. TLC, R_f (hexane/AcOEt 2/1) = 0.22.

F. 1-Bromomethyl-3-(2-methoxy-ethyl)-benzene

The title compound is prepared analogously as described for the title F compound in Example 45: MS 246.0, 247.8 [M+18]; Rt (HPLC, Nucleosil C18, 10:90-100:0 CH₃CN/H₂O + 0.1% TFA within 5 min, then 100% CH₃CN + 0.1% TFA): 5.52 min.

Example 48**4-Bromomethyl-2-(3-methoxy-propyl)-1-methyl-benzene**

**A. 5-Bromo-2-methyl-benzaldehyde**

To a stirred solution of aluminum chloride (57.93 g, 434.46 mmol) in methylene chloride (200 mL) is added at 0°C over 45 min o-methylbenzaldehyde (30 g, 249.69 mmol). To the resulting brown mixture is further added a methylene chloride solution (200 mL) of bromine (12.71 mL, 249.69 mmol) over 3 h and the reaction mixture is allowed to reach room temperature overnight. The reaction mixture is then poured into ice (600 g) and the aqueous layer is extracted twice with methylene chloride. The combined organic extracts are washed with a saturated aqueous solution of NaHCO₃, dried over anhydrous magnesium sulfate and concentrated under reduced pressure. The residue is taken up into hexane and the precipitate is filtered, washed with hexane (this protocol is repeated twice) and dried under high vacuum to afford the title compound as a mixture of two regioisomers: 5-bromo-2-methyl-benzaldehyde and 3-bromo-2-methyl-benzaldehyde in a 80/20 ratio: MS 197.0, 199.0 [M-H]. Rt (HPLC, Nucleosil C18, 10:90-100:0 CH₃CN/H₂O + 0.1% TFA within 5 min, then 100% CH₃CN + 0.1% TFA): 5.52 min.

B. 4-Bromo-2-(3-methoxy-propenyl)-1-methyl-benzene

The title compound is prepared analogously as described for the title B compound in Example 45 using (2-methoxy-ethyl)-triphenyl-phosphonium bromide (prepared in Example 46) and the title A compound, 5-bromo-2-methyl-benzaldehyde (containing 20 % of 3-bromo-2-methyl-benzaldehyde) (6 g, 30.14 mmol): TLC, R_f (hexane/AcOEt 9/1) = 0.48. Rt (HPLC, Nucleosil C18, 10:90-100:0 CH₃CN/H₂O + 0.1% TFA within 5 min, then 100% CH₃CN + 0.1% TFA): 6.26, 6.34 min.

C. 4-Bromo-2-(3-methoxy-propyl)-1-methyl-benzene

The title compound is prepared analogously as described for the title C compound in Example 45: MS 243.0, 245.0 [M-H]. TLC, R_f (hexane/AcOEt 9/1) = 0.63.

D. 3-(3-Methoxy-propyl)-4-methyl-benzaldehyde

The title compound is prepared analogously as described for the title D compound in Example 45: MS 192.9 [M+H], 210.0 [M+18].

E. [3-(3-Methoxy-propyl)-4-methyl-phenyl]-methanol


The title compound is prepared analogously as described for the title E compound in Example 45: TLC, R_f (hexane/AcOEt 2/1) = 0.38; MS 212.2 [M+18].

F. 4-Bromomethyl-2-(3-methoxy-propyl)-1-methyl-benzene

The title compound is prepared analogously as described for the title F compound in Example 45: MS 273.9, 276 [M+18]. Rt (HPLC, Nucleosil C18, 10:90-100:0 CH₃CN/H₂O + 0.1% TFA within 5 min, then 100% CH₃CN + 0.1% TFA): 6.09 min.

Example 49

4-Bromomethyl-2-(4-methoxy-butyl)-1-methyl-benzene



A. 4-Bromo-2-(4-methoxy-but-1-enyl)-1-methyl-benzene

The title compound is prepared analogously as described for the title B compound in Example 45 using (3-methoxy-propyl)-triphenyl-phosphonium bromide and 5-bromo-2-methyl-benzaldehyde (prepared in Example 48; containing 20 % of 3-bromo-2-methyl-benzaldehyde): Rt (HPLC, Nucleosil C18, 10:90-100:0 CH₃CN/H₂O + 0.1% TFA within 5 min, then 100% CH₃CN + 0.1% TFA): 6.60 min.

B. 4-Bromo-2-(4-methoxy-butyl)-1-methyl-benzene

The title compound is prepared analogously as described for the title C compound in Example 45: Rt (HPLC, Nucleosil C18, 10:90-100:0 CH₃CN/H₂O + 0.1% TFA within 5 min, then 100% CH₃CN + 0.1% TFA): 6.72 min.

C. 3-(4-Methoxy-butyl)-4-methyl-benzaldehyde


The title compound is prepared analogously as described for the title D compound in Example 45: TLC, R_f (hexane/AcOEt 9/1) = 0.5; MS 224.1 [M+18].

D. [3-(4-Methoxy-butyl)-4-methyl-phenyl]-methanol

The title compound is prepared analogously as described for the title E compound in Example 45: MS 225.9 [M+18]; TLC, R_f (hexane/AcOEt 2/1) = 0.4. Rt (HPLC, Nucleosil C18, 10:90-100:0 CH₃CN/H₂O + 0.1% TFA within 5 min, then 100% CH₃CN + 0.1% TFA): 5.04 min.

E. 4-Bromomethyl-2-(4-methoxy-butyl)-1-methyl-benzene

The title compound is prepared analogously as described for the title F compound in Example 45: TLC, R_f (hexane/AcOEt 1/1) = 0.85; MS 288.0, 289.8 [M+18]; Rt (HPLC, Nucleosil C18, 10:90-100:0 CH₃CN/H₂O + 0.1% TFA within 5 min, then 100% CH₃CN + 0.1% TFA): 6.35 min.

Example 50**1-Bromomethyl-2-(3-methoxy-propyl)-benzene****A. 3-(2-Hydroxymethyl-phenyl)-propan-1-ol** (Malandra, J. L.; Trahanovski, W. S. *JOC* **1995**, 60, 261-263)

In a solution of 1,2-dihydronaphtalene (6.5 g, 50 mmol) in methylene chloride (200 mL) ozone is bubbled at -75°C until a blue color developed. Excess of ozone is then removed by bubbling nitrogen through the solution until the blue color has dissipated. The reaction mixture is allowed to reach room temperature and the solvent partially removed under *vacuum*. The resulting ozonide is dissolved in THF (50 mL) and this solution is added drop wise at 0°C to a solution of LiAlH₄ (55 mL, 55 mmol, 1 M in THF) in THF (125 mL). The reaction mixture is further stirred for 4 h at room temperature, and wet Na₂SO₄ is added to the reaction mixture until the evolution of H₂ ceased. The mixture is filtered off and the white solid washed with EtOAc. The filtrate is dried over anhydrous magnesium sulfate and concentrated under reduced pressure. The residue is purified by flash column chromatography on silica gel (hexane/EtOAc 4/1 to EtOAc/MeOH 95/5) to afford the title compound: TLC, R_f (AcOEt) = 0.40. MS 183.9 [M+18].

B. [2-(3-Methoxy-propyl)-phenyl]-methanol

To a suspension of the title A compound, 3-(2-hydroxymethyl-phenyl)-propan-1-ol (14.9 g, 90 mmol) and proton-sponge (25 g, 1.3 eq.) in methylene chloride (500 mL) is added at 0°C under nitrogen trimethyloxonium-tetrafluoroborate (17.3 g, 117 mmol). The reaction mixture is further stirred for 5 h at room temperature before filtration. The filtrate is poured into water and the aqueous layer extracted twice with methylene chloride. The combined organic


extracts are dried over Na_2SO_4 , filtered and concentrated under reduced pressure. The residue is purified by flash column chromatography on silica gel (hexane/EtOAc 7/3 to 1/1 and to AcOEt/MeOH 9/1) to afford the title compound: TLC, Rf (hexane/AcOEt 4/1) = 0.57. MS 198.0 [M+18].

C. 1-Bromomethyl-2-(3-methoxy-propyl)-benzene

The title compound is prepared analogously as described for the title D compound in Example 42: TLC, Rf (hexane/AcOEt 95/5) = 0.4. MS 259.8, 261.9 [M+18].

Example 51

(1-Bromomethyl-cyclopropylmethyl)-benzene



A. Cyclopropanecarboxylic acid *tert*-butyl ester

To a solution of *tert*-butanol (9.4 mL, 100 mmol) in 50 mL of anhydrous THF is added under nitrogen during several minutes *n*-butyl lithium (69 mL, 110 mmol, 1.6 M solution in hexane). After 30 min, the resulting solution is treated by the drop wise addition of a THF solution (40 mL) of cyclopropanecarbonyl chloride (10 mL, 110 mmol). The reaction mixture is further stirred at reflux for 1 h, cooled to 0°C by an ice bath, and slowly hydrolyzed by the addition of water. The aqueous layer is extracted twice with ether and the combined organic extracts are dried over anhydrous sodium sulfate and concentrated under reduced pressure. The residue is purified by vacuum distillation (bp = 70°C, 76 mmHg) to afford the title compound as a yellow oil.

B. 1-Benzyl-cyclopropanecarboxylic acid *tert*-butyl ester

To a stirring solution of LDA (11 mL, 22 mmol, 2 M in THF) in THF (40 mL) under nitrogen is added drop wise at -75°C a THF solution (10 mL) of the title A compound, cyclopropanecarboxylic acid *tert*-butyl ester (2.85 g, 20 mmol). The resulting mixture is stirred for 5 h at -75°C before the addition of a THF solution (10 mL) of benzylbromide (3.8 mL, 32 mmol). The reaction mixture is allowed to reach room temperature overnight and poured into a saturated NH_4Cl aqueous solution. The aqueous layer is extracted twice with EtOAc and the combined organic extracts are dried over anhydrous magnesium sulfate and concentrated under reduced pressure. The residue is purified by flash column

chromatography on silica gel (hexane/EtOAc 97/3) to afford the title compound: TLC, Rf (hexane/AcOEt 95/5) = 0.47; MS 249.9 [M+18].

C. (1-Benzyl-cyclopropyl)-methanol


To a stirring solution of the title B compound, 1-benzyl-cyclopropanecarboxylic acid *tert*-butyl ester (0.232 g, 1 mmol) in THF (8 mL) under nitrogen is added drop wise at -50°C LiAlH₄ (3 mL, 3 mmol, 1 m in THF). The resulting mixture is further stirred for 1 h at -50°C, then slowly allowed to reach room temperature and poured into a saturated NaHCO₃ aqueous solution. The aqueous layer is extracted twice with EtOAc, and the combined organic extracts are dried over anhydrous magnesium sulfate and concentrated under reduced pressure. The residue is purified by flash column chromatography on silica gel (hexane/EtOAc 4/1 to 1/1) to afford the title compound: TLC, Rf (hexane/AcOEt 1/1) = 0.52. MS 180.0 [M+18]. Rt (HPLC, Nucleosil C18, 10:90-100:0 CH₃CN/H₂O + 0.1% TFA within 5 min, then 100% CH₃CN + 0.1% TFA): 4.95 min.

D. (1-Bromomethyl-cyclopropylmethyl)-benzene

To a solution of the title C compound, (1-benzyl-cyclopropyl)-methanol (0.8 g, 4.93 mmol), pyridine (994 μ L, 12.3 mmol) and CBr₄ (1.8 g, 5.42 mmol) in methylene chloride (25 mL) is added triphenylphosphine (1.8 g, 6.9 mmol), with stirring at 0°C. After completion of the reaction, the mixture is poured into a saturated aqueous solution of NaHCO₃. The aqueous layer is extracted twice with methylene chloride and the combined organic extracts are dried over anhydrous magnesium sulfate and concentrated under reduced pressure. The residue is taken up into ether and the triphenylphosphine oxide precipitate is filtered off. The filtrate is concentrated under reduced pressure and purified by flash column chromatography on silica gel (hexane/EtOAc 9/1) to afford the title compound: TLC, Rf (hexane/AcOEt 99/1) = 0.54. Rt (HPLC, Nucleosil C18, 10:90-100:0 CH₃CN/H₂O + 0.1% TFA within 5 min, then 100% CH₃CN + 0.1% TFA): 6.41 min.

Example 52

1-(1-Bromomethyl-cyclopropylmethyl)-2-(3-methoxy-propyl)-benzene



A. 1-[2-(3-Methoxy-propyl)-benzyl]-cyclopropanecarboxylic acid *tert*-butyl ester

The title compound is prepared analogously as described for the title B compound in Example 51: TLC, R_f (hexane/AcOEt 95/5) = 0.2. MS 322.2 [M+18].

B. {1-[2-(3-Methoxy-propyl)-benzyl]-cyclopropyl}-methanol

The title compound is prepared analogously as described for the title C compound in Example 51: TLC, R_f (hexane/AcOEt 1/1) = 0.42.

C. 1-(1-Bromomethyl-cyclopropylmethyl)-2-(3-methoxy-propyl)-benzene


The title compound is prepared analogously as described for the title D compound in Example 51: TLC, R_f (hexane/AcOEt 95/5) = 0.52.

Throughout this specification and the claims which follow, unless the context requires otherwise, the word "comprise", and variations such as "comprises" and "comprising", will be understood to imply the inclusion of a stated integer or step or group of integers or steps but not the exclusion of any other integer or step or group of integers or steps.

The reference in this specification to any prior publication (or information derived from it), or to any matter which is known, is not, and should not be taken as an acknowledgment or admission or any form of suggestion that that prior publication (or information derived from it) or known matter forms part of the common general knowledge in the field of endeavour to which this specification relates.

THE CLAIMS DEFINING THE INVENTION ARE AS FOLLOWS:

1. A compound of the formula



wherein

R₁ is -CH₂-X, -O-X or -S(O)₀₋₂-X; or

R₁ is -NR₈-X, -NR₈C(O)-X or -NR₈S(O)₂-X in which

R₈ is hydrogen or C₁₋₇ alkyl; and

X is -(CH₂)_m-(CR₉R₁₀)_p-(CH₂)_n-Z-(CH₂)_q-W in which

m, n and q are independently zero or an integer from 1 to 5;

p is zero or 1;

R₉ and R₁₀ are independently hydrogen, hydroxy, halogen, C₁₋₇ alkyl, C₁₋₇ alkoxy or cycloalkyl; or

R₉ and R₁₀ combined are alkylene which together with the carbon atom to which they are attached form a 3- to 6-membered ring;

Z is a bond; or

Z is O, S(O)₀₋₂, or -NR₁₁- in which

R₁₁ is hydrogen or C₁₋₇ alkyl, provided that R₁ is -CH₂-X when m, n and p are all zero;

W is aryl or heterocyclyl;

R₂ is hydrogen, halogen, cyano, hydroxy or C₁₋₇ alkoxy;

L is a bond; or

- 77 -

L is $-(CH_2)_s-O-(CH_2)_v-$ in which

s and v are independently zero or an integer from 1 to 3; or

L is $-C(O)-$, $-C(O)O-$, $-OC(O)-$, $-OC(O)NR_{12}-$, $-NR_{12}-$, $-NR_{13}C(O)-$, $-NR_{13}C(O)O-$ or $-NR_{13}C(O)NR_{12}-$ in which

R_{12} and R_{13} are independently hydrogen or C_{1-7} alkyl;

R_3 is hydrogen, hydroxy, halogen or cyano provided that L is a bond; or

R_3 is optionally substituted C_{1-7} alkyl, aralkyl, heteroaralkyl, aryl or heterocyclyl; or

R_3 and R_{12} combined are alkylene which together with the nitrogen atom to which they are attached form a 5- to 6-membered ring;

R_4 is hydrogen, optionally substituted C_{1-7} alkyl or aryl;

R_5 and R_6 are independently hydrogen, halogen, hydroxy, trifluoromethyl, optionally substituted C_{1-7} alkyl, C_{1-7} alkoxy or cycloalkyl; or

R_5 and R_6 combined together with the carbon atoms to which they are attached form a fused 5- to 6-membered aromatic or heteroaromatic ring provided that R_5 and R_6 are attached to carbon atoms adjacent to each other; or

R_5 and R_6 combined are alkylene which together with the carbon atoms to which they are attached form a fused 5- to 7-membered ring provided that R_5 and R_6 are attached to carbon atoms adjacent to each other; or

C- R_5 and C- R_6 may be replaced with nitrogen;

R_7 is hydrogen, halogen, hydroxy, trifluoromethyl, optionally substituted C_{1-7} alkyl, C_{1-7} alkoxy, cycloalkyl, alkanoyl, alkoxyalkoxy, alkanoyloxy, amino, alkylamino, dialkylamino, acylamino, carbamoyl, thiol, alkylthio, alkylthiono, sulfonyl, sulfonamido, sulfamoyl, nitro, cyano, carboxy, alkoxy carbonyl, aryl, alkenyl, alkynyl, aralkoxy, heterocyclyl including indolyl, imidazolyl, furyl, thienyl, thiazolyl, pyrrolidyl, pyridyl, pyrimidyl, piperidyl, morpholinyl and tetrazolyl; or

R_7 and R_6 combined are O, $S(O)_{0-2}$, $-NR_{14}-$, $-(CH_2)_{1-2}-$, $-O-CH_2-$, $-CH_2-O-$, $-S(O)_{0-2}-CH_2-$, $-CH_2-S(O)_{0-2}-$, $-NR_{14}-CH_2-$, $-CH_2-NR_{14}-$, $-S(O)_{0-2}-NR_{14}-$ or $-NR_{14}-S(O)_{0-2}-$ in which

R_{14} is hydrogen or C_{1-7} alkyl, provided R_6 is located at the 2' position; or

C- R_7 may be replaced with nitrogen;

- 78 -

Y is $-(CH_2)_r-$, $-O-(CH_2)_r-$, $-(CH_2)_r-O-$, $-S_{0-2}-(CH_2)_r-$ or $-(CH_2)_r-S_{0-2}-$ in which
r is zero or an integer from 1 to 3;

Q combined with the atoms to which it is attached form a 5- to 6-membered monocyclic aromatic or heteroaromatic ring; or

Q combined with the atoms to which it is attached form a 7- to 12-membered bicyclic aromatic or heterocyclic ring;

or a pharmaceutically acceptable salt thereof.

2. A compound according to Claim 1 wherein

R_1 is $-CH_2-X$, $-O-X$ or $-S(O)_{0-2}-X$; or

R_1 is $-NR_8-X$, $-NR_8C(O)-X$ or $-NR_8S(O)_2-X$ in which

R_8 is hydrogen or C_{1-7} alkyl; and

X is $-(CH_2)_m-(CR_9R_{10})_p-(CH_2)_n-Z-(CH_2)_q-W$ in which

m and n are independently zero or an integer from 1 to 5;

p is zero or 1;

q is zero;

R_9 and R_{10} are independently hydrogen, hydroxy, halogen, C_{1-7} alkyl, C_{1-7} alkoxy or cycloalkyl; or

R_9 and R_{10} combined are alkylene which together with the carbon atom to which they are attached form a 3- to 6-membered ring;

Z is a bond; or

Z is O, $S(O)_{0-2}$, or $-NR_{11}-$ in which

R_{11} is hydrogen or C_{1-7} alkyl, provided that R_1 is $-CH_2-X$ when m, n and p are all zero;

W is aryl or heterocyclyl;

R_2 is hydrogen, halogen, cyano, hydroxy or C_{1-7} alkoxy;

L is a bond; or

L is $-(CH_2)_s-O-(CH_2)_v-$ in which

s and v are zero; or

- 79 -

L is $-\text{C}(\text{O})-$, $-\text{C}(\text{O})\text{O}-$, $-\text{OC}(\text{O})-$, $-\text{OC}(\text{O})\text{NR}_{12}-$, $-\text{NR}_{12}-$, $-\text{NR}_{13}\text{C}(\text{O})-$, $-\text{NR}_{13}\text{C}(\text{O})\text{O}-$ or $-\text{NR}_{13}\text{C}(\text{O})\text{NR}_{12}-$ in which

R_{12} and R_{13} are independently hydrogen or C_{1-7} alkyl;

R_3 is hydrogen, halogen or cyano provided that L is a bond; or

R_3 is optionally substituted C_{1-7} alkyl, aralkyl, heteroaralkyl, aryl or heterocyclyl; or

R_3 and R_{12} combined are alkylene which together with the nitrogen atom to which they are attached form a 5- to 6-membered ring;

R_4 is hydrogen, optionally substituted C_{1-7} alkyl or aryl;

R_5 and R_6 are independently hydrogen, halogen, hydroxy, trifluoromethyl, optionally substituted C_{1-7} alkyl, C_{1-7} alkoxy or cycloalkyl; or

R_5 and R_6 combined together with the carbon atoms to which they are attached form a fused 5- to 6-membered aromatic or heteroaromatic ring provided that R_5 and R_6 are attached to carbon atoms adjacent to each other; or

R_5 and R_6 combined are alkylene which together with the carbon atoms to which they are attached form a fused 5- to 7-membered ring provided that R_5 and R_6 are attached to carbon atoms adjacent to each other; or

R_7 is hydrogen, halogen, hydroxy, trifluoromethyl, optionally substituted, C_{1-7} alkyl, C_{1-7} alkoxy or cycloalkyl; or

R_7 and R_6 combined are O , $\text{S}(\text{O})_{0-2}$, $-\text{NR}_{14}-$, $-(\text{CH}_2)_{1-2}-$, $-\text{O}-\text{CH}_2-$, $-\text{CH}_2-\text{O}-$, $-\text{S}(\text{O})_{0-2}-\text{CH}_2-$, $-\text{CH}_2-\text{S}(\text{O})_{0-2}-$, $-\text{NR}_{14}-\text{CH}_2-$, $-\text{CH}_2-\text{NR}_{14}-$, $-\text{S}(\text{O})_{0-2}-\text{NR}_{14}-$ or $-\text{NR}_{14}-\text{S}(\text{O})_{0-2}-$ in which

R_{14} is hydrogen or C_{1-7} alkyl, provided R_6 is located at the 2' position;

Y is $-(\text{CH}_2)_r-$ in which


r is zero;

Q combined with the carbon atoms to which it is attached form a 5- to 6-membered monocyclic aromatic or heteroaromatic ring; or

Q combined with the carbon atoms to which it is attached form a 9- to 10-membered bicyclic aromatic or heterocyclic ring;

or a pharmaceutically acceptable salt thereof.

3. A compound according to Claim 2 of the formula




wherein

R_1 , R_2 , L , R_3 , R_4 , R_5 , R_6 , R_7 and Q have meanings as defined in Claim 2;

or a pharmaceutically acceptable salt thereof.

4. A compound according to Claim 2 of the formula



wherein

R_1 , R_2 , L , R_3 , R_4 , R_5 , R_6 , R_7 and Q have meanings as defined in Claim 2;

or a pharmaceutically acceptable salt thereof.

5. A compound according to Claim 4 wherein

R_1 is $-\text{CH}_2-\text{X}$, $-\text{O}-\text{X}$ or $-\text{S}-\text{X}$; or

R_1 is $-\text{NR}_8-\text{X}$, $-\text{NR}_8\text{C}(\text{O})-\text{X}$ or $-\text{NR}_8\text{S}(\text{O})_2-\text{X}$ in which

R_8 is hydrogen or C_{1-7} alkyl; and

X is $-(\text{CH}_2)_m-(\text{CR}_9\text{R}_{10})_p-(\text{CH}_2)_n-\text{Z}-\text{W}$ in which

m and n are independently zero or an integer of 1 or 2;

p is zero or 1;

- 81 -

R_9 and R_{10} are independently hydrogen, hydroxy, halogen, C_{1-7} alkyl, C_{1-7} alkoxy or cycloalkyl; or

R_9 and R_{10} combined are alkylene which together with the carbon atom to which they are attached form a 3- to 6-membered ring;

Z is a bond; or

Z is O, $S(O)_{0-2}$, or $-NR_{11}-$ in which

R_{11} is hydrogen or C_{1-7} alkyl, provided that R_1 is $-CH_2-X$ when m, n and p are all zero;

W is aryl or heterocyclyl;

R_2 is hydrogen, halogen or hydroxy;

L is a bond;

R_3 is hydrogen or halogen;

R_4 is hydrogen, optionally substituted C_{1-7} alkyl or aryl;

R_5 and R_6 are independently hydrogen, halogen, hydroxy, trifluoromethyl, optionally substituted C_{1-7} alkyl, C_{1-7} alkoxy or cycloalkyl;

R_7 is hydrogen, halogen, hydroxy, trifluoromethyl, optionally substituted C_{1-7} alkyl, C_{1-7} alkoxy or cycloalkyl; or

R_7 and R_6 combined are O, $S(O)_{0-2}$, $-NR_{14}-$, $-(CH_2)_{1-2}-$, $-O-CH_2-$, $-CH_2-O-$, $-S(O)_{0-2}-CH_2-$, $-CH_2-S(O)_{0-2}-$, $-NR_{14}-CH_2-$, $-CH_2-NR_{14}-$, $-S(O)_{0-2}-NR_{14}-$ or $-NR_{14}-S(O)_{0-2}-$ in which

R_{14} is hydrogen or C_{1-7} alkyl, provided R_6 is located at the 2'-position;

Q combined with the atoms to which it is attached form a 5- to 6-membered monocyclic aromatic or heteroaromatic ring; or

Q combined with the atoms to which it is attached form a 9- to 10-membered bicyclic aromatic or heterocyclic ring;

or a pharmaceutically acceptable salt thereof.

6. A compound according to Claim 5 wherein

R_1 is $-CH_2-X$, $-O-X$ or $-S-X$; or

R_1 is $-NR_8-X$, $-NR_8C(O)-X$ or $-NR_8S(O)_2-X$ in which

- 82 -

R_8 is hydrogen or C_{1-7} alkyl; and

X is $-(CH_2)_m-(CR_9R_{10})_p-(CH_2)_n-Z-W$ in which

m and n are independently zero or an integer of 1 or 2;

p is zero or 1;

R_9 and R_{10} are independently hydrogen or C_{1-7} alkyl; or

Z is a bond; or

Z is O, $S(O)_{0-2}$, or $-NR_{11}-$ in which

R_{11} is hydrogen or C_{1-7} alkyl, provided that R_1 is $-CH_2-X$ when m, n and p are all zero;

W is aryl or heterocyclyl;

R_2 is hydrogen, halogen or hydroxy;

L is a bond;

R_3 is hydrogen or halogen;

R_4 is hydrogen;

R_5 and R_6 are independently hydrogen, halogen, hydroxy, trifluoromethyl, optionally substituted C_{1-7} alkyl, C_{1-7} alkoxy or cycloalkyl;

R_7 is hydrogen, halogen, hydroxy, trifluoromethyl, optionally substituted C_{1-7} alkyl, C_{1-7} alkoxy or cycloalkyl; or

R_7 and R_6 combined are O, $S(O)_{0-2}$, $-NR_{14}-$, $-(CH_2)_{1-2}-$, $-O-CH_2-$, $-CH_2-O-$, $-S(O)_{0-2}-CH_2-$, $-CH_2-S(O)_{0-2}-$, $-NR_{14}-CH_2-$, $-CH_2-NR_{14}-$, $-S(O)_{0-2}-NR_{14}-$ or $-NR_{14}-S(O)_{0-2}-$ in which

R_{14} is hydrogen or C_{1-7} alkyl, provided R_6 is located at the 2'-position;

Q combined with the atoms to which it is attached form a 5- to 6-membered monocyclic aromatic or heteroaromatic ring; or

Q combined with the atoms to which it is attached form a 9- to 10-membered bicyclic aromatic or heterocyclic ring;

or a pharmaceutically acceptable salt thereof.

7. A compound according to Claim 6 wherein

- 83 -

R_1 is $-NR_8-X$, $-NR_8C(O)-X$ or $-NR_8S(O)_2-X$ in which

R_8 is hydrogen or C_{1-7} alkyl;

or a pharmaceutically acceptable salt thereof.

8. A compound according to Claim 6 wherein

Q combined with the carbon atoms to which it is attached form a pyridyl or pyrimidinyl ring;


or a pharmaceutically acceptable salt thereof.

9. A compound according to Claim 6 wherein

Q combined with the carbon atoms to which it is attached form a thienyl, furyl, pyrrolyl or indolyl ring;

or a pharmaceutically acceptable salt thereof.

10. A compound according to Claim 5 of the formula



wherein

R_1 is $-CH_2-X$, $-O-X$ or $-S-X$; or

R_1 is $-NR_8-X$, $-NR_8C(O)-X$ or $-NR_8S(O)_2-X$ in which

R_8 is hydrogen or C_{1-7} alkyl; and

X is $-(CH_2)_m-(CR_9R_{10})_p-(CH_2)_n-Z-W$ in which

m, n and p are independently zero or 1;

R_9 is hydrogen;

R_{10} is hydrogen or C_{1-7} alkyl;

- 84 -

Z is a bond; or

Z is O, S(O)₀₋₂, or -NR₁₁- in which

R₁₁ is hydrogen or C₁₋₇ alkyl, provided that R₁ is -CH₂-X when m, n and p are all zero;

W is aryl or heterocyclyl;

R₂ is hydrogen;

R₃ is hydrogen or halogen;

R₅ and R₆ are independently hydrogen, halogen, hydroxy, trifluoromethyl, optionally substituted C₁₋₇ alkyl, C₁₋₇ alkoxy or cycloalkyl;

R₇ is hydrogen, halogen, hydroxy, trifluoromethyl, optionally substituted C₁₋₇ alkyl, C₁₋₇ alkoxy or cycloalkyl; or

R₇ and R₈ combined are O, S(O)₀₋₂, -NR₁₄-, -(CH₂)₁₋₂-, -O-CH₂-, -CH₂-O-, -S(O)₀₋₂-CH₂-, -CH₂-S(O)₀₋₂-, -NR₁₄-CH₂-, -CH₂-NR₁₄-, -S(O)₀₋₂-NR₁₄- or -NR₁₄-S(O)₀₋₂- in which

R₁₄ is hydrogen or C₁₋₇ alkyl, provided R₆ is located at the 2'-position;

R₁₅ is hydrogen, halogen, hydroxy, trifluoromethyl, optionally substituted C₁₋₇ alkyl, C₁₋₇ alkoxy or cycloalkyl;

or a pharmaceutically acceptable salt thereof.

11. A compound according to Claim 10 wherein

R₁ is -O-X or -S-X; and

X is -(CH₂)_m-(CR₉R₁₀)_p-(CH₂)_n-Z-W in which

m is 1;

n and p are zero;

Z is a bond;

W is aryl or heterocyclyl;

R₃ is hydrogen or halogen;

R₅ is hydrogen, halogen, hydroxy, trifluoromethyl, optionally substituted C₁₋₇ alkyl, C₁₋₇ alkoxy or cycloalkyl;

- 85 -

R_6 is hydrogen;

R_7 is hydrogen;

R_{15} is hydrogen, halogen, hydroxy, trifluoromethyl, optionally substituted C_{1-7} alkyl, C_{1-7} alkoxy or cycloalkyl;

or a pharmaceutically acceptable salt thereof.

12. A compound according to Claim 11 wherein

R_3 is hydrogen;

or a pharmaceutically acceptable salt thereof.

13. A compound according to Claim 12 wherein

W is monocyclic aryl;

or a pharmaceutically acceptable salt thereof.


14. A compound according to Claim 5 wherein

p is 1;

R_9 and R_{10} combined are alkylene which together with the carbon atom to which they are attached form a 3- to 6-membered ring;

or a pharmaceutically acceptable salt thereof.

15. A compound according to Claim 14 of the formula



wherein

R_1 is $-CH_2-X$, $-O-X$ or $-S-X$; or

R_1 is $-NR_8-X$, $-NR_8C(O)-X$ or $-NR_8S(O)_2-X$ in which

- 86 -

R_8 is hydrogen or C_{1-7} alkyl; and

X is $-(CH_2)_m-CR_9R_{10}-(CH_2)_n-Z-W$ in which

m and n are 1;

Z is a bond; or

Z is O, $S(O)_{0-2}$, or $-NR_{11}-$ in which

R_{11} is hydrogen or C_{1-7} alkyl, provided that R_1 is $-CH_2-X$ when m, n and p are all zero;

W is aryl or heterocyclyl;

R_2 is hydrogen;

R_3 is hydrogen or halogen;

R_5 and R_6 are independently hydrogen, halogen, hydroxy, trifluoromethyl, optionally substituted C_{1-7} alkyl, C_{1-7} alkoxy or cycloalkyl;

R_7 is hydrogen, halogen, hydroxy, trifluoromethyl, optionally substituted C_{1-7} alkyl, C_{1-7} alkoxy or cycloalkyl; or

R_7 and R_6 combined are O, $S(O)_{0-2}$, $-NR_{14}-$, $-(CH_2)_{1-2}-$, $-O-CH_2-$, $-CH_2-O-$, $-S(O)_{0-2}-CH_2-$, $-CH_2-S(O)_{0-2}-$, $-NR_{14}-CH_2-$, $-CH_2-NR_{14}-$, $-S(O)_{0-2}-NR_{14}-$ or $-NR_{14}-S(O)_{0-2}-$ in which

R_{14} is hydrogen or C_{1-7} alkyl, provided R_6 is located at the 2'-position;

R_{15} is hydrogen, halogen, hydroxy, trifluoromethyl, optionally substituted C_{1-7} alkyl, C_{1-7} alkoxy or cycloalkyl;

or a pharmaceutically acceptable salt thereof.

16. A compound of claim 15 wherein

R_1 is $-O-X$ or $-S-X$; and

X is $-CH_2-CR_9R_{10}-CH_2-Z-W$ in which

Z is a bond;

W is aryl;

R_3 is hydrogen;

R_5 is hydrogen, halogen, hydroxy, trifluoromethyl, optionally substituted C_{1-7} alkyl, C_{1-7} alkoxy or cycloalkyl;

- 87 -

R₈ is hydrogen;

R₇ is hydrogen;

R₁₅ is hydrogen, halogen, hydroxy, trifluoromethyl, optionally substituted C₁₋₇ alkyl, C₁₋₇ alkoxy or cycloalkyl;

or a pharmaceutically acceptable salt thereof.

17. A method for the inhibition of renin activity in mammals which method comprises administering to a mammal in need thereof a therapeutically effective amount of a compound of any one of Claims 1 to 16.

18. A method for the prevention and/or treatment of conditions associated with renin activity in mammals which method comprises administering to a mammal in need thereof a therapeutically effective amount of a compound of any one of Claims 1 to 16.

19. A method according to Claim 18, which method comprises administering said compound in combination with a therapeutical effective amount of an anti-diabetic agent, a hypolipidemic agent, an anti-obesity agent or an anti-hypertensive agent.

20. A method for the treatment of hypertension, atherosclerosis, unstable coronary syndrome, congestive heart failure, cardiac hypertrophy, cardiac fibrosis, cardiomyopathy postinfarction, unstable coronary syndrome, diastolic dysfunction, chronic kidney disease, hepatic fibrosis, complications resulting from diabetes, such as nephropathy, vasculopathy and neuropathy, diseases of the coronary vessels, restenosis following angioplasty, raised intra-ocular pressure, glaucoma, abnormal vascular growth, hyperaldosteronism, cognitive impairment, alzheimers, dementia, anxiety states and cognitive disorders, which method comprises administering to a mammal in need thereof a therapeutical effective amount of a compound of any one of Claims 1 to 16.

21. A pharmaceutical composition comprising a therapeutically effective amount of a compound of any one of Claims 1 to 16 in combination with one or more pharmaceutical acceptable carriers.

22. A pharmaceutical composition comprising a therapeutically effective amount of a compound of any one of Claims 1 to 16 in combination with a therapeutically effective

amount of an anti-diabetic agent, a hypolipidemic agent, an anti-obesity agent or an anti-hypertensive agent.

23. A pharmaceutical composition according to Claim 21 or Claim 22 for the treatment of hypertension, atherosclerosis, unstable coronary syndrome, congestive heart failure, cardiac hypertrophy, cardiac fibrosis, cardiomyopathy postinfarction, unstable coronary syndrome, diastolic dysfunction, chronic kidney disease, hepatic fibrosis, complications resulting from diabetes, such as nephropathy, vasculopathy and neuropathy, diseases of the coronary vessels, restenosis following angioplasty, raised intra-ocular pressure, glaucoma, abnormal vascular growth, hyperaldosteronism, cognitive impairment, alzheimers, dementia, anxiety states and cognitive disorders.

24. A pharmaceutical composition according to Claim 21 or Claim 22, for use as medicament.

25. Use of a pharmaceutical composition according to Claim 21 or Claim 22, for the preparation of a medicament for the treatment of conditions associated with renin activity.

26. Use of a compound according to any one of Claims 1 to 16, for the preparation of a pharmaceutical composition for the treatment of conditions associated with renin activity.

27. Use according to Claim 25 or Claim 26, wherein the condition associated with renin activity is selected from hypertension, atherosclerosis, unstable coronary syndrome, congestive heart failure, cardiac hypertrophy, cardiac fibrosis, cardiomyopathy postinfarction, unstable coronary syndrome, diastolic dysfunction, chronic kidney disease, hepatic fibrosis, complications resulting from diabetes, such as nephropathy, vasculopathy and neuropathy, diseases of the coronary vessels, restenosis following angioplasty, raised intra-ocular pressure, glaucoma, abnormal vascular growth, hyperaldosteronism, cognitive impairment, alzheimers, dementia, anxiety states and cognitive disorders.

28. A compound according to any one of Claims 1 to 16, for use as a medicament.

29. Compound according to Claim 1 substantially as hereinbefore described with reference to any one of the Examples.