

(12) STANDARD PATENT

(11) Application No. AU 2003229961 B2

(19) AUSTRALIAN PATENT OFFICE

(54) Title
Immunoglobulin-domain containing cell surface recognition molecules

(51) International Patent Classification(s)
C07K 14/705 (2006.01) **C12N 15/12** (2006.01)

(21) Application No: **2003229961** (22) Date of Filing: **2003.04.30**

(87) WIPO No: **WO03/093316**

(30) Priority Data

(31) Number
0209884.6 (32) Date
2002.04.30 (33) Country
GB

(43) Publication Date: **2003.11.17**
(43) Publication Journal Date: **2004.01.08**
(44) Accepted Journal Date: **2009.02.19**

(71) Applicant(s)
Ares Trading S.A.

(72) Inventor(s)
Phelps, Christopher Benjamin;Fagan, Richard Joseph;Davids, Andrew Robert;Power, Christine

(74) Agent / Attorney
Griffith Hack, Level 3 509 St Kilda Road, Melbourne, VIC, 3004

(56) Related Art
WO 2001/054477
WO 2002/034783
Edwards, J. B. D. M. EMBL Accession No: BD511416

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization International Bureau



(43) International Publication Date
13 November 2003 (13.11.2003)

PCT

(10) International Publication Number
WO 2003/093316 A3

(51) International Patent Classification⁷: **C07K 14/705**, C12N 15/63, A01K 67/027, C12N 15/12, 5/10

(21) International Application Number: PCT/GB2003/001851

(22) International Filing Date: 30 April 2003 (30.04.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data: 0209884.6 30 April 2002 (30.04.2002) GB

(71) Applicant (for all designated States except US): **ARES TRADING S.A. [CH/CH]**; Chateau de Vaumarcus, CH-2028 Vaumarcus (CH).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **DAVIDS, Andrew, Robert [GB/GB]**; Inpharmatica, 60 Charlotte Street, London W1T 2NU (GB). **FAGAN, Richard, Joseph [US/GB]**; Inpharmatica, 60 Charlotte Street, London W1T 2NU (GB). **PHELPS, Christopher, Benjamin [GB/GB]**; Inpharmatica, 60 Charlotte Street, London W1T 2NU (GB). **POWER, Christine [GB/FR]**; 10, rue des Jonquilles, F-01710 Thoiry (FR).

(74) Agents: **GOODFELLOW, Hugh, Robin et al.**; Carpmaels & Ransford, 43 Bloomsbury Square, London WC1A 2RA (GB).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GII, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:
— with international search report
— before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

(88) Date of publication of the international search report: 12 February 2004

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

WO 2003/093316 A3

(54) Title: IMMUNOGLOBULIN-DOMAIN CONTAINING CELL SURFACE RECOGNITION MOLECULES

(57) Abstract: This invention relates to a novel protein (INSP052), herein identified as an immunoglobulin domain-containing cell surface recognition molecule and to the use of this protein and nucleic acid sequences from the encoding gene in the diagnosis, prevention and treatment of disease.

21 Jan 2009

2003229961

IMMUNOGLOBULIN-DOMAIN CONTAINING CELL SURFACE RECOGNITION MOLECULES

This invention relates to novel proteins (termed INSP052 and INSP055), herein identified as immunoglobulin domain-containing cell surface recognition molecules and to the use of these proteins and nucleic acid sequences from the encoding genes in the diagnosis, prevention and treatment of disease.

All publications, patents and patent applications cited herein are incorporated in full by reference.

It is to be understood that, if any prior art publication is referred to herein, such reference does not constitute an admission that the publication forms a part of the common general knowledge in the art, in Australia or any other country.

In the claims which follow and in the preceding description of the invention, except where the context requires otherwise due to express language or necessary implication, the word "comprise" or variations such as "comprises" or "comprising" is used in an inclusive sense, i.e. to specify the presence of the stated features but not to preclude the presence or addition of further features in various embodiments of the invention.

BACKGROUND

The process of drug discovery is presently undergoing a fundamental revolution as the era of functional genomics comes of age. The term "functional genomics" applies to an approach utilising bioinformatics tools to ascribe function to protein sequences of interest. Such tools are becoming increasingly necessary as the speed of generation of sequence data is rapidly outpacing the ability of research laboratories to assign functions to these protein sequences.

As bioinformatics tools increase in potency and in accuracy, these tools are rapidly replacing the conventional techniques of biochemical characterisation. Indeed, the advanced bioinformatics tools used in identifying the present invention are now capable of outputting results in which a high degree of confidence can be placed.

Various institutions and commercial organisations are examining sequence data as they become available and significant discoveries are being made on an on-going basis. However, there remains a continuing need to identify and characterise further genes and the polypeptides that they encode, as targets for research and drug discovery.

21 Jan 2009

20003229961

Recently, a remarkable tool for the evaluation of sequences of unknown function has been developed by the Applicant for the present invention. This tool is a database system, termed the Biopendium search database, that is the subject of WO 01/69507. This database system consists of an integrated data resource created using proprietary technology and containing information generated from an all-by-all comparison of all available protein or nucleic acid sequences.

The aim behind the integration of these sequence data from separate data resources is to combine as much data as possible, relating both to the sequences themselves and to

information relevant to each sequence, into one integrated resource. All the available data relating to each sequence, including data on the three-dimensional structure of the encoded protein, if this is available, are integrated together to make best use of the information that is known about each sequence and thus to allow the most educated predictions to be made 5 from comparisons of these sequences. The annotation that is generated in the database and which accompanies each sequence entry imparts a biologically relevant context to the sequence information.

This data resource has made possible the accurate prediction of protein function from sequence alone. Using conventional technology, this is only possible for proteins that 10 exhibit a high degree of sequence identity (above about 20%-30% identity) to other proteins in the same functional family. Accurate predictions are not possible for proteins that exhibit a very low degree of sequence homology to other related proteins of known function.

SIGNAL PEPTIDE-CONTAINING PROTEINS

15 The ability of cells to make and secrete extracellular proteins is central to many biological processes. Enzymes, growth factors, extracellular matrix proteins and signaling molecules are all secreted by cells. This is through fusion of a secretory vesicle with the plasma membrane. In most cases, but not all, proteins are directed to the endoplasmic reticulum and into secretory vesicles by a signal peptide. Signal peptides are cis-acting sequences 20 that affect the transport of polypeptide chains from the cytoplasm to a membrane bound compartment such as a secretory vesicle. Polypeptides that are targeted to the secretory vesicles are either secreted into the extracellular matrix or are retained in the plasma membrane. The polypeptides that are retained in the plasma membrane will have one or more transmembrane domains. Examples of signal peptide containing proteins that play a 25 central role in the functioning of a cell are cytokines, hormones, extracellular matrix proteins, adhesion molecules, receptors, proteases, and growth and differentiation factors.

IMMUNOGLOBULIN DOMAIN-CONTAINING CELL SURFACE RECOGNITION MOLECULES

30 Immunoglobulin domain-containing cell surface recognition molecules have been shown to play a role in diverse physiological functions, many of which can play a role in disease processes. Alteration of their activity is a means to alter the disease phenotype and as such identification of novel immunoglobulin domain-containing cell surface recognition

molecules is highly relevant as they may play a role in many diseases, particularly inflammatory disease, oncology, and cardiovascular disease. Immunoglobulin domain-containing cell surface recognition molecules are involved in a range of biological processes, including: embryogenesis (Martin-Bermudo, M.D. *et al.*, *Development*. 2000 127(12):2607-15; Chen, L.M., *et al.*, *J Neurosci*. 2000 20(10):3776-84; Zweegman, S., *et al.*, *Exp Hematol*. 2000 28(4):401-10; Darribere, T., *et al.*, *Biol Cell*. 2000 92(1):5-25), maintenance of tissue integrity (Eckes, B., *et al.*, *J Cell Sci*. 2000 113(Pt 13):2455-2462; Buckwalter, J.A., *et al.*, *Instr Course Lect*. 2000 49:481-9; Frenette, P.S., *et al.*, *J Exp Med*. 2000 191(8):1413-22; Delmas, V., *et al.*, *Dev Biol*. 1999 216(2):491-506; Humphries, M.J., *et al.*, *Trends Pharmacol Sci*. 2000 21(1):29-32; Miosge, N., *et al.*, *Lab Invest*. 1999 79(12):1591-9; Nagaoka T, *et al.*, *Am J Pathol* 2000 Jul 157:1 237-47; Nwariaku FE, *et al.*, *J Trauma* 1995 39(2): 285-8; Zhu X, *et al.*, *Zhonghua Zheng Xing Shao Shang Wai Ke Za Zhi* 1999 15(1): 53-5), leukocyte extravasation/inflammation (Lim, L.H., *et al.*, *Am J Respir Cell Mol Biol*. 2000 22(6):693-701; Johnston, B., *et al.*, *Microcirculation*. 2000 7(2):109-18; Mertens, A.V., *et al.*, *Clin Exp Allergy*. 1993 23(10):868-73; Chcialowski, A., *et al.*, *Pol Merkuriusz Lek*. 2000 7(43):13-7; Rojas, A.I., *et al.*, *Crit Rev Oral Biol Med*. 1999 10(3):337-58; Marinova-Mutafchieva, L., *et al.*, *Arthritis Rheum*. 2000 43(3):638-44; Vijayan, K.V., *et al.*, *J Clin Invest*. 2000 105(6):793-802; Currie, A.J., *et al.*, *J Immunol*. 2000 164(7):3878-86; Rowin, M.E., *et al.*, *Inflammation*. 2000 24(2):157-73; Johnston, B., *et al.*, *J Immunol*. 2000 164(6):3337-44; Gerst, J.L., *et al.*, *J Neurosci Res*. 2000 59(5):680-4; Kagawa, T.F., *et al.*, *Proc Natl Acad Sci U S A*. 2000 97(5):2235-40; Hillan, K.J., *et al.*, *Liver*. 1999 19(6):509-18; Panes, J., 1999 22(10):514-24; Arao, T., *et al.*, *J Clin Endocrinol Metab*. 2000 85(1):382-9; Souza, H.S., *et al.*, *Gut*. 1999 45(6):856-63; Grunstein, M.M., *et al.*, *Am J Physiol Lung Cell Mol Physiol*. 2000 278(6):L1154-63; Mertens, A.V., *et al.*, *Clin Exp Allergy*. 1993 23(10):868-73; Berends, C., *et al.*, *Clin Exp Allergy*. 1993 23(11):926-33; Fernvik, E., *et al.*, *Inflammation*. 2000 24(1):73-87; Bocchino, V., *et al.*, *J Allergy Clin Immunol*. 2000 105(1 Pt 1):65-70; Jones SC, *et al.*, *Gut* 1995 36(5):724-30; Liu CM, *et al.*, *Ann Allergy Asthma Immunol* 1998 81(2):176-80; McMurray RW, *Semin Arthritis Rheum* 1996 25(4):215-33; Takahashi H, *et al.*, *Eur J Immunol* 1992 22(11): 2879-85; Carlos T, *et al.*, *J Heart Lung Transplant* 1992 11(6): 1103-8; Fabrega E, *et al.*, *Transplantation* 2000 69(4): 569-73; Zohrens G, *et al.*, *Hepatology* 1993 18(4): 798-802; Montefort S, *et al.*, *Am J Respir Crit Care Med* 1994 149(5): 1149-52), oncogenesis (Orr, F.W., *et al.*, *Cancer*. 2000 88(S12):2912-2918; Zeller, W., *et al.*, *J Hematother Stem Cell Res*. 1999 8(5):539-46; Okada, T., *et al.*, *Clin*

Exp Metastasis. 1999 17(7):623-9; Mateo, V., *et al.*, Nat Med. 1999 5(11):1277-84; Yamaguchi, K., *et al.*, J Exp Clin Cancer Res. 2000 19(1):113-20; Maeshima, Y., *et al.*, J Biol Chem. 2000 275(28):21340-8; Van Waes, C., *et al.*, Int J Oncol. 2000 16(6):1189-95; Damiano, J.S., *et al.*, Leuk Lymphoma. 2000 38(1-2):71-81; Seftor, R.E., *et al.*, Cancer 5 Metastasis Rev. 1999 18(3):359-75; Shaw, L.M., J Mammary Gland Biol Neoplasia. 1999 4(4):367-76; Weyant, M.J., *et al.*, Clin Cancer Res. 2000 6(3):949-56), angiogenesis (Koch AE, *et al* Nature 1995 376 (6540): 517-9; Wagener C & Ergun S. Exp Cell Res 2000 261(1): 19-24; Ergun S, *et al.* Mol Cell 2000 5(2): 311-20), bone resorption (Hartman GD, & Duggan ME. Expert Opin Investig Drugs 2000 9(6): 1281-91; Tanaka Y, *et al.* J Bone Miner Res 10 1995 10(10): 1462-9; Lark MW, *et al.* J Pharmacol Exp Ther 1999 291(2): 612-7; Raynal C, *et al.* Endocrinology 1996 137(6):2347-54; Ilvesaro JM, *et al.* Exp Cell Res 1998 242(1): 75-83), neurological dysfunction (Ossege LM, *et al.* Int Immunopharmacol 2001 1:1085-100; Bitsch A, *et al*, Stroke 1998 29:2129-35; Iadecola C & Alexander M. Curr Opin Neurol 2001 14:89-94; Becker K, *et al* Stroke 2001 32(1): 206-11; Relton JK, *et al* Stroke 2001 32(1): 199-15 205; Hamada Y, *et al* J Neurochem 1996 66:1525-31), thrombogenesis (Wang, Y.G., *et al.*, J Physiol (Lond). 2000 526(Pt 1):57-68; Matsuno, H., *et al.*, Nippon Yakurigaku Zasshi. 2000 115(3):143-50; Eliceiri, B.P., *et al.*, Cancer J Sci Am. 2000 6(Suppl 3):S245-9; von Beckerath, N., *et al.*, Blood. 2000 95(11):3297-301; Topol, E.J., *et al.*, Am Heart J. 2000 139(6):927-33; Kroll, H., *et al.*, Thromb Haemost. 2000 83(3):392-6), and invasion/adherence 20 of bacterial pathogens to the host cell (Dersch P, *et al.* EMBO J 1999 18(5): 1199-1213).

The detailed characterisation of the structure and function of several immunoglobulin-domain containing cell surface recognition molecule families has led to active programs by a number of pharmaceutical companies to develop modulators for use in the treatment of diseases involving inflammation, oncology, neurology, immunology and cardiovascular 25 function. Immunoglobulin domain containing cell surface recognition molecules are involved in virtually every aspect of biology from embryogenesis to apoptosis. They are essential to the structural integrity and homeostatic functioning of most tissues. It is therefore not surprising that defects in immunoglobulin domain containing cell surface recognition molecules cause disease and that many diseases involve modulation of immunoglobulin 30 domain containing cell surface recognition molecule function. The members of this family are described below in Table 1.

The Immunoglobulin domain containing cell surface recognition molecule family in fact contains several distinct families. Of these families, some are of particular pharmaceutical interest due to small molecule tractability. They include:

1. The immunoglobulin adhesion molecules represent the counter receptors for the integrins and includes the intracellular adhesion molecules (ICAMs) and vascular cell adhesion molecules (VCAMs). Members are composed of variable numbers of globular, immunoglobulin-like, extracellular domains. Some members of the family, for example, PECAM-1 (CD31) and NCAM, mediate homotypic adhesion. Some members of the family, for example ICAM-1 and VCAM-1, mediate adhesion via interactions with integrins.
2. Cell surface growth factor receptors. Growth factors are extracellular and in order to exert a biological effect they interact with specific, high affinity receptors located on the plasma membranes of target cells. The molecular characterisation of a variety of different growth factor receptors revealed that they fall into defined families; the tyrosine kinase receptors, G-protein associated seven transmembrane receptors, and the serine/threonine kinase receptors. The tyrosine kinase receptors are characterised by an extracellular domain, a transmembrane domain, and an intracellular domain which possess tyrosine kinase activity. VEGFR, PDGFR, FGFR, CSF-1R and c-KIT are examples of tyrosine kinase growth factor receptors which also contain immunoglobulin domains in the extracellular portion. Dys-regulation of growth factor function results in many different disease phenotypes, including, but not exclusive to oncology (Bartucci M *et al.*, (2001) *Cancer Res.* Sep 15;61(18):6747-54, Dias S *et al.*, (2001) *Proc Natl Acad Sci U S A.* Sep 11;98(19):10857-62, Djavan B *et al.*, (2001) *World J Urol.* 19(4):225-33), inflammation (Fiocchi C. (2001) *J Clin Invest.* Aug;108(4):523-6, Hodge S *et al.*, (2001) *Respirology.* Sep;6(3):205-211, Fenwick SA *et al.*, (2001) *J Anat.* Sep;199(Pt 3):231-40), neurological (Cooper JD *et al.*, (2001) *Proc Natl Acad Sci U S A* 98(18):10439-44, Fahnstock M *et al.*, (2001) *Mol Cell Neurosci* 18(2):210-20), and metabolism (Vickers MH *et al.*, (2001) *Endocrinology.* 142(9):3964-73).

30 **Table 1: Immunoglobulin domain-containing cell surface recognition molecules**

<i>Receptor</i>	<i>Ligand</i>	<i>Distribution</i>
-----------------	---------------	---------------------

ICAM-1 5 Ig domains	LFA-1 (CD11a/CD18) Mac-1 (CD11b/CD18), CD43	Widespread, endothelial cells, fibroblasts, epithelium, monocytes, lymphocytes, dendritic cells, chondrocytes.
ICAM-2 2 Ig domains	LFA-1 (CD11b)	endothelial cells (high): lymphocytes, monocytes, basophils, platelets (low).
ICAM-3 5 Ig domains	LFA-1 (α d/CD18)	Lymphocytes, monocytes, neutrophils, eosinophils, basophils.
VCAM-1 6 or 7 Ig domains	α 4 β 1, α 4 β 7	Endothelial cells, monocytes, fibroblasts, dendritic cells, bone marrow stromal cells, myoblasts.
LFA-3 6 Ig domains	CD2	Endothelial cells, leukocytes, epithelial cells
PECAM-1 (CD31)	CD31, heparin	Endothelial cells (at EC-EC junctions), T cell subsets, platelets, neutrophils, eosinophils, monocytes, smooth muscle cells, bone marrow stem cells.
NCAM	NCAM, heparin SO ₄	Neural cells, muscle
MAdCAM-1 4 Ig domains	α 4 β 7, L-selectin	Peyer's patch, mesenteric lymph nodes, mucosal endothelial cells, spleen.
CD2	CD58, CD59, CD48	T lymphocytes
VEGFR	VEGF	Widespread, retina, umbilical vein, adrenal, neuronal precursor cells
FGFR	FGF	Widespread, brain, colon, ovary
KIT	Stem Cell Factor, MGF	Widespread, foetus, melanocytes, gall bladder, cerebellum, gastric epithelium (low)
PDGFR	PDGF	Widespread, breast, placenta, fibroblast, liver, ovary, skin, heart
CSF-1R	CSF	Widespread, placenta, liver, multiple sclerosis lesions, spleen, lung, breast.

Immunoglobulin domain-containing cell surface recognition molecules have thus been shown to play a role in diverse physiological functions, many of which can play a role in disease processes. Alteration of their activity is a means to alter the disease phenotype and

5 as such identification of novel Immunoglobulin domain-containing cell surface recognition molecules is highly relevant as they may play a role in many diseases, particularly

immunology, inflammatory disease, oncology, cardiovascular disease, central nervous system disorders and infection.

THE INVENTION

The invention is based on the discovery that the INSP052 and INSP055 proteins function 5 as immunoglobulin domain-containing cell surface recognition molecules. Examples of immunoglobulin domain-containing cell surface recognition molecules are listed in Table 1.

In one embodiment of the first aspect of the invention, there is provided a polypeptide which:

- 10 (i) comprises or consists of the amino acid sequence as recited in SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO: 6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:16 or the extracellular domain of INSP052;
- (ii) is a fragment thereof having the activity of a polypeptide according to (i), or having an antigenic determinant in common with a polypeptide according to (i); or
- 15 (iii) is a functional equivalent of (i) or (ii).

By "the activity of a polypeptide according to (i)", we refer to immunoglobulin domain-containing cell surface recognition molecule activity. By immunoglobulin domain-containing cell surface recognition molecule activity we refer to polypeptides that comprise amino acid sequence or structural features that can be identified as conserved 20 features within the immunoglobulin domain-containing cell surface recognition molecule family.

The polypeptide having the sequence recited in SEQ ID NO:2 is referred to hereafter as "the INSP052 exon 1 polypeptide". The polypeptide having the sequence recited in SEQ ID NO:4 is referred to hereafter as "the INSP052 exon 2 polypeptide". The polypeptide 25 having the sequence recited in SEQ ID NO:6 is referred to hereafter as "the INSP052 exon 3 polypeptide". The polypeptide having the sequence recited in SEQ ID NO:8 is referred to hereafter as "the INSP052 exon 4 polypeptide". The polypeptide having the sequence recited in SEQ ID NO:10 is referred to hereafter as "the INSP052 exon 5 polypeptide". The polypeptide having the sequence recited in SEQ ID NO:12 is referred to hereafter as "the 30 INSP052 exon 6 polypeptide". The polypeptide having the sequence recited in SEQ ID NO:14 is referred to hereafter as "the INSP052 exon 7 polypeptide". Combining SEQ ID

NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12 and SEQ ID NO:14 produces the sequence recited in SEQ ID NO:16. The polypeptide having the sequence recited in SEQ ID NO:16 is referred to hereafter as the INSP052 polypeptide.

The term “INSP052 exon polypeptides” as used herein includes polypeptides comprising 5 or consisting of the INSP052 exon 1 polypeptide, the INSP052 exon 2 polypeptide, the INSP052 exon 3 polypeptide, the INSP052 exon 4 polypeptide, the INSP052 exon 5 polypeptide, the INSP052 exon 6 polypeptide, the INSP052 exon 7 polypeptide, the INSP052 polypeptide and the extracellular domain of INSP052.

In one embodiment, the polypeptide according to this embodiment consists of the amino 10 acid sequence recited in SEQ ID NO:16 or is a fragment of or functional equivalent thereof. In another embodiment, the polypeptide consists of the amino acid sequence recited in SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO: 6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, or SEQ ID NO:14, or a variant thereof.

In one embodiment of the first aspect of the invention there is provided a polypeptide 15 which:

- i) comprises or consists of the amino acid sequence of the extracellular domain of INSP052;
- ii) is a fragment thereof having the activity of a polypeptide according to (i), or having an antigenic determinant in common with a polypeptide according to (i); or
- 20 iii) is a functional equivalent of (i) or (ii).

The extracellular domain of INSP052 corresponds to amino acids 1-240 (see the Examples section). See also Figure 7 for the extracellular domain of INSP052.

In a second embodiment of the first aspect of the invention, there is provided a polypeptide which:

- 25 (i) comprises or consists of the amino acid sequence as recited in SEQ ID NO:18,
- (ii) is a fragment thereof having the activity of a polypeptide of (i), or having an antigenic determinant in common with a polypeptide of (i); or
- (iii) is a functional equivalent of (i) or (ii).

By “the activity of a polypeptide according to (i)”, we refer to immunoglobulin domain-30 containing cell surface recognition molecule activity.

Preferably, the polypeptide according to this embodiment consists of the amino acid sequence recited in SEQ ID NO:18 or is a fragment of or functional equivalent thereof.

The polypeptide having the sequence recited in SEQ ID NO:18 is referred to hereafter as "the INSP055 polypeptide".

5 In a second aspect, the invention provides a purified nucleic acid molecule which encodes a polypeptide of the first aspect of the invention.

Preferably, the purified nucleic acid molecule comprises or consists of the nucleic acid sequence as recited in SEQ ID NO:1 (encoding the INSP052 A exon 1 polypeptide), SEQ ID NO:3 (encoding the INSP052 exon 2 polypeptide), SEQ ID NO:5 (encoding the

10 INSP052 exon 3 polypeptide), SEQ ID NO:7 (encoding the INSP052 exon 4 polypeptide), SEQ ID NO:9 (encoding the INSP052 exon 5 polypeptide), SEQ ID NO:11 (encoding the INSP052 exon 6 polypeptide), SEQ ID NO:13 (encoding the INSP052 exon 7 polypeptide), SEQ ID NO:15 (encoding the INSP052 polypeptide), or SEQ ID NO:17 (encoding the INSP055 polypeptide), or is a redundant equivalent or fragment of any one

15 of these sequences.

Combining the sequences recited in SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:11 and SEQ ID NO:13 produces the sequence recited in SEQ ID NO:15.

In one embodiment of the second aspect of the invention there is provided a nucleic acid
20 molecule which encodes a polypeptide which comprises or consists of the extracellular domain of INSP052. Preferably, the nucleic acid molecule comprises or consists of the nucleic acid sequence set forth in Figure 7 or the coding portion of the nucleic acid sequence set forth in Figure 7.

In a third aspect, the invention provides a purified nucleic acid molecule which hybridizes
25 under high stringency conditions with a nucleic acid molecule of the second aspect of the invention.

In a fourth aspect, the invention provides a vector, such as an expression vector, that contains a nucleic acid molecule of the second or third aspect of the invention.

In a fifth aspect, the invention provides a host cell transformed with a vector of the fourth
30 aspect of the invention.

In a sixth aspect, the invention provides a ligand which binds specifically to, and which

preferably inhibits the activity of a polypeptide of the first aspect of the invention.

By "the activity of a polypeptide of the invention" and similar expressions, we refer to activity characteristic of immunoglobulin domain-containing cell surface recognition molecules.

5 In a seventh aspect, the invention provides a compound that is effective to alter the expression of a natural gene which encodes a polypeptide of the first aspect of the invention or to regulate the activity of a polypeptide of the first aspect of the invention.

A compound of the seventh aspect of the invention may either increase (agonise) or decrease (antagonise) the level of expression of the gene or the activity of the polypeptide.

10 Importantly, the identification of the function of the INSP052 and INSP055 polypeptides allows for the design of screening methods capable of identifying compounds that are effective in the treatment and/or diagnosis of disease. Ligands and compounds according to the sixth and seventh aspects of the invention may be identified using such methods. These methods are included as aspects of the present invention.

15 In an eighth aspect, the invention provides a polypeptide of the first aspect of the invention, or a nucleic acid molecule of the second or third aspect of the invention, or a vector of the fourth aspect of the invention, or a host cell of the fifth aspect, or a ligand of the sixth aspect of the invention, or a compound of the seventh aspect of the invention, for use in therapy or diagnosis. These molecules may also be used in the manufacture of a

20 medicament for the treatment of diseases including, but not limited to, cell proliferative disorders, autoimmune/inflammatory disorders, cardiovascular disorders, neurological and psychiatric disorders, developmental disorders, genetic disorders, metabolic disorders, infections and other pathological conditions. These diseases preferably include neoplasm, cancer, brain tumour, glioma, bone tumor, lung tumor, breast tumour, prostate tumour,

25 colon tumour, hemangioma, myeloproliferative disorder, leukemia, hematological disease, neutropenia, thrombocytopenia, angiogenesis disorders, dermatological disease, ageing, wounds, burns, fibrosis, cardiovascular disease, restenosis, heart disease, peripheral vascular disease, coronary artery disease, oedema, thromboembolism, dysmenorrhea, endometriosis, pre-eclampsia, lung disease, COPD, asthma bone disease, renal disease,

30 glomerulonephritis, liver disease, Crohn's disease, gastritis, ulcerative colitis, ulcer, immune disorder, autoimmune disease, arthritis, rheumatoid arthritis, psoriasis, epidermolysis bullosa, systemic lupus erythematosus, ankylosing spondylitis, Lyme

disease, multiple sclerosis, neurodegeneration, stroke, brain/spinal cord injury, Alzheimer's disease, Parkinson's disease, motor neurone disease, neuromuscular disease, HIV, AIDS, cytomegalovirus infection, fungal infection, ocular disorder, macular degeneration, glaucoma, diabetic retinopathy, ocular hypertension and other conditions in which 5 immunoglobulin domain containing cell surface recognition molecules are implicated. These moieties of the first, second, third, fourth, sixth or seventh aspect of the invention may also be used in the manufacture of a medicament for the treatment of such diseases.

In a ninth aspect, the invention provides a method of diagnosing a disease in a patient, comprising assessing the level of expression of a natural gene encoding a polypeptide of 10 the first aspect of the invention or the activity of a polypeptide of the first aspect of the invention in tissue from said patient and comparing said level of expression or activity to a control level, wherein a level that is different to said control level is indicative of disease. Such a method will preferably be carried out *in vitro*. Similar methods may be used for monitoring the therapeutic treatment of disease in a patient, wherein altering the level of 15 expression or activity of a polypeptide or nucleic acid molecule over the period of time towards a control level is indicative of regression of disease.

A preferred method for detecting polypeptides of the first aspect of the invention comprises the steps of: (a) contacting a ligand, such as an antibody, of the sixth aspect of the invention with a biological sample under conditions suitable for the formation of a 20 ligand-polypeptide complex; and (b) detecting said complex.

A number of different methods according to the ninth aspect of the invention exist, as the skilled reader will be aware, such as methods of nucleic acid hybridization with short probes, point mutation analysis, polymerase chain reaction (PCR) amplification and methods using antibodies to detect aberrant protein levels. Similar methods may be used on 25 a short or long term basis to allow therapeutic treatment of a disease to be monitored in a patient. The invention also provides kits that are useful in these methods for diagnosing disease.

Preferably, the disease diagnosed by a method of the ninth aspect of the invention is a disease in which immunoglobulin domain-containing cell surface recognition molecules 30 are implicated, as described above.

In a tenth aspect, the invention provides for the use of the polypeptides of the first aspect of the invention as immunoglobulin domain-containing cell surface recognition molecules.

The importance of the Ig domain in cell surface receptors is described in Lokker NA et al., "Functional importance of platelet-derived growth factor (PDGF) receptor extracellular immunoglobulin-like domains. Identification of PDGF binding site and neutralizing monoclonal antibodies," *J Biol Chem* 1997 Dec 26;272(52):33037-44.

- 5 The invention also provides for the use of a nucleic acid molecule according to the second or third aspects of the invention to express a protein that possesses immunoglobulin domain-containing cell surface recognition molecule activity. The invention also provides a method for effecting immunoglobulin domain-containing cell surface recognition molecule activity, said method utilising a polypeptide of the first aspect of the invention.
- 10 In an eleventh aspect, the invention provides a pharmaceutical composition comprising a polypeptide of the first aspect of the invention, or a nucleic acid molecule of the second or third aspect of the invention, or a vector of the fourth aspect of the invention, or a host cell of the fifth aspect of the invention, or a ligand of the sixth aspect of the invention, or a compound of the seventh aspect of the invention, in conjunction with a pharmaceutically-
15 acceptable carrier.

In a twelfth aspect, the present invention provides a polypeptide of the first aspect of the invention, or a nucleic acid molecule of the second or third aspect of the invention, or a vector of the fourth aspect of the invention, or a host cell of the fifth aspect of the invention, or a ligand of the sixth aspect of the invention, or a compound of the seventh aspect of the invention, for use in therapy or diagnosis. These molecules may also be used in the manufacture of a medicament for the treatment of diseases including, but not limited to, cell proliferative disorders, autoimmune/inflammatory disorders, cardiovascular disorders, neurological and psychiatric disorders, developmental disorders, genetic disorders, metabolic disorders, infections and other pathological conditions. These diseases
20 preferably include neoplasm, cancer, brain tumour, glioma, bone tumor, lung tumor, breast tumour, prostate tumour, colon tumour, hemangioma, myeloproliferative disorder, leukemia, hematological disease, neutropenia, thrombocytopenia, angiogenesis disorders, dermatological disease, ageing, wounds, burns, fibrosis, cardiovascular disease, restenosis, heart disease, peripheral vascular disease, coronary artery disease, oedema,
25 thromboembolism, dysmenorrhea, endometriosis, pre-eclampsia, lung disease, COPD, asthma, bone disease, renal disease, glomerulonephritis, liver disease, Crohn's disease, gastritis, ulcerative colitis, ulcer, immune disorder, autoimmune disease, arthritis, rheumatoid arthritis, psoriasis, epidermolysis bullosa, systemic lupus erythematosus,

5 ankylosing spondylitis, Lyme disease, multiple sclerosis, neurodegeneration, stroke, brain/spinal cord injury, Alzheimer's disease, Parkinson's disease, motor neurone disease, neuromuscular disease, HIV, AIDS, cytomegalovirus infection, fungal infection, ocular disorder, macular degeneration, glaucoma, diabetic retinopathy, ocular hypertension and other conditions in which immunoglobulin domain containing cell recognition molecules are implicated.

In a thirteenth aspect, the invention provides a method of treating a disease in a patient comprising administering to the patient a polypeptide of the first aspect of the invention, or a nucleic acid molecule of the second or third aspect of the invention, or a vector of the 10 fourth aspect of the invention, or a host cell of the fifth aspect of the invention, or a ligand of the sixth aspect of the invention, or a compound of the seventh aspect of the invention.

For diseases in which the expression of a natural gene encoding a polypeptide of the first aspect of the invention, or in which the activity of a polypeptide of the first aspect of the invention, is lower in a diseased patient when compared to the level of expression or 15 activity in a healthy patient, the polypeptide, nucleic acid molecule, ligand or compound administered to the patient should be an agonist. Conversely, for diseases in which the expression of the natural gene or activity of the polypeptide is higher in a diseased patient when compared to the level of expression or activity in a healthy patient, the polypeptide, nucleic acid molecule, ligand or compound administered to the patient should be an 20 antagonist. Examples of such antagonists include antisense nucleic acid molecules, ribozymes and ligands, such as antibodies.

In a fourteenth aspect, the invention provides transgenic or knockout non-human animals that have been transformed to express higher, lower or absent levels of a polypeptide of the first aspect of the invention. Such transgenic animals are very useful models for the study 25 of disease and may also be used in screening regimes for the identification of compounds that are effective in the treatment or diagnosis of such a disease.

Preferably, the disease is a disease in which immunoglobulin domain-containing cell surface recognition molecules are implicated, as described above.

It should be appreciated that the scope of protection sought for the polypeptides and 30 nucleic acids of the present invention does not extend to nucleic acids or polypeptides present in their natural source. Rather, the polypeptides and nucleic acids claimed by the present invention may be regarded as being "isolated" or "purified". The terms "isolated"

and "purified" as used herein refer to a nucleic acid or polypeptide separated from at least one other component (e. g., nucleic acid or polypeptide) present with the nucleic acid or polypeptide in its natural source. Thus, for example, a polypeptide contained in a tissue extract would constitute an "isolated" or "purified" polypeptide, as would a polypeptide 5 synthetically or recombinantly produced. In one embodiment, the nucleic acid or polypeptide is found in the presence of (if anything) only a solvent, buffer, ion, or other component normally present in a solution of the same.

It should be noted that the terms "isolated" and "purified" do not denote the method by which 10 the polypeptide or nucleic acid is obtained or the level of purity of the preparation. Thus, such isolated or purified species may be produced recombinantly, isolated directly from the cell or tissue of interest or produced synthetically based on the determined sequences.

A summary of standard techniques and procedures which may be employed in order to utilise the invention is given below. It will be understood that this invention is not limited to the particular methodology, protocols, cell lines, vectors and reagents described. It is 15 also to be understood that the terminology used herein is for the purpose of describing particular embodiments only and it is not intended that this terminology should limit the scope of the present invention. The extent of the invention is limited only by the terms of the appended claims.

Standard abbreviations for nucleotides and amino acids are used in this specification.

20 The practice of the present invention will employ, unless otherwise indicated, conventional techniques of molecular biology, microbiology, recombinant DNA technology and immunology, which are within the skill of those working in the art.

Such techniques are explained fully in the literature. Examples of particularly suitable texts for consultation include the following: Sambrook Molecular Cloning; A Laboratory 25 Manual, Second Edition (1989); DNA Cloning, Volumes I and II (D.N. Glover ed. 1985); Oligonucleotide Synthesis (M.J. Gait ed. 1984); Nucleic Acid Hybridization (B.D. Hames & S.J. Higgins eds. 1984); Transcription and Translation (B.D. Hames & S.J. Higgins eds. 1984); Animal Cell Culture (R.I. Freshney ed. 1986); Immobilized Cells and Enzymes (IRL Press, 1986); B. Perbal, A Practical Guide to Molecular Cloning (1984); the Methods 30 in Enzymology series (Academic Press, Inc.), especially volumes 154 & 155; Gene Transfer Vectors for Mammalian Cells (J.H. Miller and M.P. Calos eds. 1987, Cold Spring

Harbor Laboratory); Immunochemical Methods in Cell and Molecular Biology (Mayer and Walker, eds. 1987, Academic Press, London); Scopes, (1987) Protein Purification: Principles and Practice, Second Edition (Springer Verlag, N.Y.); and Handbook of Experimental Immunology, Volumes I-IV (D.M. Weir and C. C. Blackwell eds. 1986).

5 As used herein, the term "polypeptide" includes any peptide or protein comprising two or more amino acids joined to each other by peptide bonds or modified peptide bonds, i.e. peptide isosteres. This term refers both to short chains (peptides and oligopeptides) and to longer chains (proteins).

The polypeptide of the present invention may be in the form of a mature protein or may be

10 a pre-, pro- or prepro- protein that can be activated by cleavage of the pre-, pro- or prepro- portion to produce an active mature polypeptide. In such polypeptides, the pre-, pro- or prepro- sequence may be a leader or secretory sequence or may be a sequence that is employed for purification of the mature polypeptide sequence.

The polypeptide of the first aspect of the invention may form part of a fusion protein. For

15 example, it is often advantageous to include one or more additional amino acid sequences which may contain secretory or leader sequences, pro-sequences, sequences which aid in purification, or sequences that confer higher protein stability, for example during recombinant production. Alternatively or additionally, the mature polypeptide may be fused with another compound, such as a compound to increase the half-life of the
20 polypeptide (for example, polyethylene glycol).

Polypeptides may contain amino acids other than the 20 gene-encoded amino acids, modified either by natural processes, such as by post-translational processing or by chemical modification techniques which are well known in the art. Among the known modifications which may commonly be present in polypeptides of the present invention are

25 glycosylation, lipid attachment, sulphation, gamma-carboxylation, for instance of glutamic acid residues, hydroxylation and ADP-ribosylation. Other potential modifications include acetylation, acylation, amidation, covalent attachment of flavin, covalent attachment of a haeme moiety, covalent attachment of a nucleotide or nucleotide derivative, covalent attachment of a lipid derivative, covalent attachment of phosphatidylinositol, cross-linking,
30 cyclization, disulphide bond formation, demethylation, formation of covalent cross-links, formation of cysteine, formation of pyroglutamate, formylation, GPI anchor formation, iodination, methylation, myristoylation, oxidation, proteolytic processing, phosphorylation,

prenylation, racemization, selenoylation, transfer-RNA mediated addition of amino acids to proteins such as arginylation, and ubiquitination.

Modifications can occur anywhere in a polypeptide, including the peptide backbone, the amino acid side-chains and the amino or carboxyl termini. In fact, blockage of the amino 5 or carboxyl terminus in a polypeptide, or both, by a covalent modification is common in naturally-occurring and synthetic polypeptides and such modifications may be present in polypeptides of the present invention.

The modifications that occur in a polypeptide often will be a function of how the polypeptide is made. For polypeptides that are made recombinantly, the nature and extent 10 of the modifications in large part will be determined by the post-translational modification capacity of the particular host cell and the modification signals that are present in the amino acid sequence of the polypeptide in question. For instance, glycosylation patterns vary between different types of host cell.

The polypeptides of the present invention can be prepared in any suitable manner. Such 15 polypeptides include isolated naturally-occurring polypeptides (for example purified from cell culture), recombinantly-produced polypeptides (including fusion proteins), synthetically-produced polypeptides or polypeptides that are produced by a combination of these methods.

The functionally-equivalent polypeptides of the first aspect of the invention may be 20 polypeptides that are homologous to the INSP052 and INSP055 polypeptides. Two polypeptides are said to be "homologous", as the term is used herein, if the sequence of one of the polypeptides has a high enough degree of identity or similarity to the sequence of the other polypeptide. "Identity" indicates that at any particular position in the aligned sequences, the amino acid residue is identical between the sequences. "Similarity" 25 indicates that, at any particular position in the aligned sequences, the amino acid residue is of a similar type between the sequences. Degrees of identity and similarity can be readily calculated (Computational Molecular Biology, Lesk, A.M., ed., Oxford University Press, New York, 1988; Biocomputing. Informatics and Genome Projects, Smith, D.W., ed., Academic Press, New York, 1993; Computer Analysis of Sequence Data, Part 1, Griffin, 30 A.M., and Griffin, H.G., eds., Humana Press, New Jersey, 1994; Sequence Analysis in Molecular Biology, von Heijne, G., Academic Press, 1987; and Sequence Analysis Primer, Gribskov, M. and Devereux, J., eds., M Stockton Press, New York, 1991).

Homologous polypeptides therefore include natural biological variants (for example, allelic variants or geographical variations within the species from which the polypeptides are derived) and mutants (such as mutants containing amino acid substitutions, insertions or deletions) of the INSP052 and INSP055 polypeptides. Such mutants may include 5 polypeptides in which one or more of the amino acid residues are substituted with a conserved or non-conserved amino acid residue (preferably a conserved amino acid residue) and such substituted amino acid residue may or may not be one encoded by the genetic code. Typical such substitutions are among Ala, Val, Leu and Ile; among Ser and Thr; among the acidic residues Asp and Glu; among Asn and Gln; among the basic 10 residues Lys and Arg; or among the aromatic residues Phe and Tyr. Particularly preferred are variants in which several, i.e. between 5 and 10, 1 and 5, 1 and 3, 1 and 2 or just 1 amino acids are substituted, deleted or added in any combination. Especially preferred are silent substitutions, additions and deletions, which do not alter the properties and activities 15 of the protein. Also especially preferred in this regard are conservative substitutions. Such mutants also include polypeptides in which one or more of the amino acid residues includes a substituent group.

Typically, greater than 30% identity between two polypeptides is considered to be an indication of functional equivalence. Preferably, functionally equivalent polypeptides of the first aspect of the invention have a degree of sequence identity with the INSP052 and 20 INSP055 polypeptides, or with active fragments thereof, of greater than 80%. More preferred polypeptides have degrees of identity of greater than 85%, 90%, 95%, 98% or 99%, respectively.

The functionally-equivalent polypeptides of the first aspect of the invention may also be polypeptides which have been identified using one or more techniques of structural 25 alignment. For example, the Inpharmatica Genome ThreaderTM technology that forms one aspect of the search tools used to generate the Biopendium search database may be used (see co-pending International Patent Application No. PCT/GB01/01105, published as WO 01/69507) to identify polypeptides of presently-unknown function which, while having low sequence identity as compared to the INSP052 and INSP055 polypeptides, are 30 predicted to be immunoglobulin domain-containing cell surface recognition molecules, said method utilising a polypeptide of the first aspect of the invention, by virtue of sharing significant structural homology with the INSP052 and INSP055 polypeptide sequences. By "significant structural homology" is meant that the Inpharmatica Genome ThreaderTM

predicts two proteins to share structural homology with a certainty of at least 10% and more preferably, at least 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90% and above.

The polypeptides of the first aspect of the invention also include fragments of the INSP052 and INSP055 polypeptides and fragments of the functional equivalents of the INSP052 and 5 INSP055 polypeptides, provided that those fragments retain immunoglobulin domain-containing cell surface recognition molecule activity or have an antigenic determinant in common with the INSP052 and INSP055 polypeptides.

As used herein, the term "fragment" refers to a polypeptide having an amino acid sequence that is the same as part, but not all, of the amino acid sequence of the INSP052 and 10 INSP055 polypeptides or one of its functional equivalents. The fragments should comprise at least n consecutive amino acids from the sequence and, depending on the particular sequence, n preferably is 7 or more (for example, 8, 10, 12, 14, 16, 18, 20 or more). Small fragments may form an antigenic determinant.

Such fragments may be "free-standing", i.e. not part of or fused to other amino acids or 15 polypeptides, or they may be comprised within a larger polypeptide of which they form a part or region. When comprised within a larger polypeptide, the fragment of the invention most preferably forms a single continuous region. For instance, certain preferred embodiments relate to a fragment having a pre - and/or pro- polypeptide region fused to the amino terminus of the fragment and/or an additional region fused to the carboxyl 20 terminus of the fragment. However, several fragments may be comprised within a single larger polypeptide.

The polypeptides of the present invention or their immunogenic fragments (comprising at least one antigenic determinant) can be used to generate ligands, such as polyclonal or 25 monoclonal antibodies, that are immunospecific for the polypeptides. Such antibodies may be employed to isolate or to identify clones expressing the polypeptides of the invention or to purify the polypeptides by affinity chromatography. The antibodies may also be employed as diagnostic or therapeutic aids, amongst other applications, as will be apparent to the skilled reader.

The term "immunospecific" means that the antibodies have substantially greater affinity for 30 the polypeptides of the invention than their affinity for other related polypeptides in the prior art. As used herein, the term "antibody" refers to intact molecules as well as to fragments thereof, such as Fab, F(ab')2 and Fv, which are capable of binding to the

antigenic determinant in question. Such antibodies thus bind to the polypeptides of the first aspect of the invention.

By "substantially greater affinity" we mean that there is a measurable increase in the affinity for a polypeptide of the invention as compared with the affinity for known

5 immunoglobulin domain-containing cell surface recognition molecules.

Preferably, the affinity is at least 1.5-fold, 2-fold, 5-fold 10-fold, 100-fold, 10^3 -fold, 10^4 -fold, 10^5 -fold or 10^6 -fold greater for a polypeptide of the invention than for known immunoglobulin domain-containing cell surface recognition molecules.

If polyclonal antibodies are desired, a selected mammal, such as a mouse, rabbit, goat or 10 horse, may be immunised with a polypeptide of the first aspect of the invention. The polypeptide used to immunise the animal can be derived by recombinant DNA technology or can be synthesized chemically. If desired, the polypeptide can be conjugated to a carrier protein. Commonly used carriers to which the polypeptides may be chemically coupled include bovine serum albumin, thyroglobulin and keyhole limpet haemocyanin. The 15 coupled polypeptide is then used to immunise the animal. Serum from the immunised animal is collected and treated according to known procedures, for example by immunoaffinity chromatography.

Monoclonal antibodies to the polypeptides of the first aspect of the invention can also be readily produced by one skilled in the art. The general methodology for making 20 monoclonal antibodies using hybridoma technology is well known (see, for example, Kohler, G. and Milstein, C., *Nature* 256: 495-497 (1975); Kozbor *et al.*, *Immunology Today* 4: 72 (1983); Cole *et al.*, 77-96 in *Monoclonal Antibodies and Cancer Therapy*, Alan R. Liss, Inc. (1985).

Panels of monoclonal antibodies produced against the polypeptides of the first aspect of 25 the invention can be screened for various properties, i.e., for isotype, epitope, affinity, etc. Monoclonal antibodies are particularly useful in purification of the individual polypeptides against which they are directed. Alternatively, genes encoding the monoclonal antibodies of interest may be isolated from hybridomas, for instance by PCR techniques known in the art, and cloned and expressed in appropriate vectors.

30 Chimeric antibodies, in which non-human variable regions are joined or fused to human constant regions (see, for example, Liu *et al.*, *Proc. Natl. Acad. Sci. USA*, 84, 3439 (1987)), may also be of use.

The antibody may be modified to make it less immunogenic in an individual, for example by humanisation (see Jones *et al.*, *Nature*, 321, 522 (1986); Verhoeyen *et al.*, *Science*, 239, 1534 (1988); Kabat *et al.*, *J. Immunol.*, 147, 1709 (1991); Queen *et al.*, *Proc. Natl Acad. Sci. USA*, 86, 10029 (1989); Gorman *et al.*, *Proc. Natl Acad. Sci. USA*, 88, 34181 (1991); 5 and Hodgson *et al.*, *Bio/Technology*, 9, 421 (1991)). The term "humanised antibody", as used herein, refers to antibody molecules in which the CDR amino acids and selected other amino acids in the variable domains of the heavy and/or light chains of a non-human donor antibody have been substituted in place of the equivalent amino acids in a human antibody. The humanised antibody thus closely resembles a human antibody but has the binding 10 ability of the donor antibody.

In a further alternative, the antibody may be a "bispecific" antibody, that is an antibody having two different antigen binding domains, each domain being directed against a different epitope.

Phage display technology may be utilised to select genes which encode antibodies with 15 binding activities towards the polypeptides of the invention either from repertoires of PCR amplified V-genes of lymphocytes from humans screened for possessing the relevant antibodies, or from naive libraries (McCafferty, J. *et al.*, (1990), *Nature* 348, 552-554; Marks, J. *et al.*, (1992) *Biotechnology* 10, 779-783). The affinity of these antibodies can also be improved by chain shuffling (Clackson, T. *et al.*, (1991) *Nature* 352, 624-628).

20 Antibodies generated by the above techniques, whether polyclonal or monoclonal, have additional utility in that they may be employed as reagents in immunoassays, radioimmunoassays (RIA) or enzyme-linked immunosorbent assays (ELISA). In these applications, the antibodies can be labelled with an analytically-detectable reagent such as a radioisotope, a fluorescent molecule or an enzyme.

25 Preferred nucleic acid molecules of the second and third aspects of the invention are those which encode the polypeptide sequences recited in SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14 and/or SEQ ID NO:16, or SEQ ID NO:18, and functionally equivalent polypeptides. These nucleic acid molecules may be used in the methods and applications described herein. The nucleic acid 30 molecules of the invention preferably comprise at least n consecutive nucleotides from the sequences disclosed herein where, depending on the particular sequence, n is 10 or more (for example, 12, 14, 15, 18, 20, 25, 30, 35, 40 or more).

The nucleic acid molecules of the invention also include sequences that are complementary to nucleic acid molecules described above (for example, for antisense or probing purposes).

Nucleic acid molecules of the present invention may be in the form of RNA, such as

5 mRNA, or in the form of DNA, including, for instance cDNA, synthetic DNA or genomic DNA. Such nucleic acid molecules may be obtained by cloning, by chemical synthetic techniques or by a combination thereof. The nucleic acid molecules can be prepared, for example, by chemical synthesis using techniques such as solid phase phosphoramidite chemical synthesis, from genomic or cDNA libraries or by separation from an organism.

10 RNA molecules may generally be generated by the *in vitro* or *in vivo* transcription of DNA sequences.

The nucleic acid molecules may be double-stranded or single-stranded. Single-stranded DNA may be the coding strand, also known as the sense strand, or it may be the non-coding strand, also referred to as the anti-sense strand.

15 The term "nucleic acid molecule" also includes analogues of DNA and RNA, such as those containing modified backbones, and peptide nucleic acids (PNA). The term "PNA", as used herein, refers to an antisense molecule or an anti-gene agent which comprises an oligonucleotide of at least five nucleotides in length linked to a peptide backbone of amino acid residues, which preferably ends in lysine. The terminal lysine confers solubility to the

20 composition. PNAs may be pegylated to extend their lifespan in a cell, where they preferentially bind complementary single stranded DNA and RNA and stop transcript elongation (Nielsen, P.E. *et al.* (1993) *Anticancer Drug Des.* 8:53-63).

These molecules also may have a different sequence which, as a result of the degeneracy of the genetic code, encode a polypeptide of SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6,

25 SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14 and/or SEQ ID NO:16, or SEQ ID NO:18. Such molecules may include, but are not limited to, the coding sequence for the mature polypeptide by itself; the coding sequence for the mature polypeptide and additional coding sequences, such as those encoding a leader or secretory sequence, such as a pro-, pre- or prepro- polypeptide sequence; the coding sequence of the mature

30 polypeptide, with or without the aforementioned additional coding sequences, together with further additional, non-coding sequences, including non-coding 5' and 3' sequences, such as the transcribed, non-translated sequences that play a role in transcription (including

termination signals), ribosome binding and mRNA stability. The nucleic acid molecules may also include additional sequences which encode additional amino acids, such as those which provide additional functionalities.

The nucleic acid molecules of the second and third aspects of the invention may also 5 encode the fragments or the functional equivalents of the polypeptides and fragments of the first aspect of the invention. Such a nucleic acid molecule may be a naturally-occurring variant such as a naturally-occurring allelic variant, or the molecule may be a variant that is not known to occur naturally. Such non-naturally occurring variants of the nucleic acid molecule may be made by mutagenesis techniques, including those applied to nucleic acid 10 molecules, cells or organisms.

Among variants in this regard are variants that differ from the aforementioned nucleic acid molecules by nucleotide substitutions, deletions or insertions. The substitutions, deletions or insertions may involve one or more nucleotides. The variants may be altered in coding or non-coding regions or both. Alterations in the coding regions may produce conservative 15 or non-conservative amino acid substitutions, deletions or insertions.

The nucleic acid molecules of the invention can also be engineered, using methods generally known in the art, for a variety of reasons, including modifying the cloning, processing, and/or expression of the gene product (the polypeptide). DNA shuffling by random fragmentation and PCR reassembly of gene fragments and synthetic 20 oligonucleotides are included as techniques which may be used to engineer the nucleotide sequences. Site-directed mutagenesis may be used to insert new restriction sites, alter glycosylation patterns, change codon preference, produce splice variants, introduce mutations and so forth.

Nucleic acid molecules which encode a polypeptide of the first aspect of the invention may 25 be ligated to a heterologous sequence so that the combined nucleic acid molecule encodes a fusion protein. Such combined nucleic acid molecules are included within the second or third aspects of the invention. For example, to screen peptide libraries for inhibitors of the activity of the polypeptide, it may be useful to express, using such a combined nucleic acid molecule, a fusion protein that can be recognised by a commercially-available antibody. A 30 fusion protein may also be engineered to contain a cleavage site located between the sequence of the polypeptide of the invention and the sequence of a heterologous protein so that the polypeptide may be cleaved and purified away from the heterologous protein.

The nucleic acid molecules of the invention also include antisense molecules that are partially complementary to nucleic acid molecules encoding polypeptides of the present invention and that therefore hybridize to the encoding nucleic acid molecules (hybridization). Such antisense molecules, such as oligonucleotides, can be designed to

5 recognise, specifically bind to and prevent transcription of a target nucleic acid encoding a polypeptide of the invention, as will be known by those of ordinary skill in the art (see, for example, Cohen, J.S., Trends in Pharm. Sci., 10, 435 (1989), Okano, J. Neurochem. 56, 560 (1991); O'Connor, J. Neurochem 56, 560 (1991); Lee *et al.*, Nucleic Acids Res 6, 3073 (1979); Cooney *et al.*, Science 241, 456 (1988); Dervan *et al.*, Science 251, 1360 (1991)).

10 The term "hybridization" as used herein refers to the association of two nucleic acid molecules with one another by hydrogen bonding. Typically, one molecule will be fixed to a solid support and the other will be free in solution. Then, the two molecules may be placed in contact with one another under conditions that favour hydrogen bonding. Factors that affect this bonding include: the type and volume of solvent; reaction temperature; time

15 of hybridization; agitation; agents to block the non-specific attachment of the liquid phase molecule to the solid support (Denhardt's reagent or BLOTO); the concentration of the molecules; use of compounds to increase the rate of association of molecules (dextran sulphate or polyethylene glycol); and the stringency of the washing conditions following hybridization (see Sambrook *et al.* [*supra*]).

20 The inhibition of hybridization of a completely complementary molecule to a target molecule may be examined using a hybridization assay, as known in the art (see, for example, Sambrook *et al* [*supra*]). A substantially homologous molecule will then compete for and inhibit the binding of a completely homologous molecule to the target molecule under various conditions of stringency, as taught in Wahl, G.M. and S.L. Berger (1987;

25 Methods Enzymol. 152:399-407) and Kimmel, A.R. (1987; Methods Enzymol. 152:507-511).

"Stringency" refers to conditions in a hybridization reaction that favour the association of very similar molecules over association of molecules that differ. High stringency hybridisation conditions are defined as overnight incubation at 42°C in a solution

30 comprising 50% formamide, 5XSSC (150mM NaCl, 15mM trisodium citrate), 50mM sodium phosphate (pH7.6), 5x Denhardts solution, 10% dextran sulphate, and 20 microgram/ml denatured, sheared salmon sperm DNA, followed by washing the filters in

0.1X SSC at approximately 65°C. Low stringency conditions involve the hybridisation reaction being carried out at 35°C (see Sambrook *et al.* [*supra*]). Preferably, the conditions used for hybridization are those of high stringency.

Preferred embodiments of this aspect of the invention are nucleic acid molecules that are at least 70% identical over their entire length to a nucleic acid molecule encoding the INSP052 or INSP055 polypeptides (SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:13 and/or SEQ ID NO:15, or SEQ ID NO:17) and nucleic acid molecules that are substantially complementary to such nucleic acid molecules. Preferably, a nucleic acid molecule according to this aspect of the invention comprises a region that is at least 80% identical over its entire length to the coding sequence for SEQ ID NO:2 given in SEQ ID NO:1, the coding sequence for SEQ ID NO:4 given in SEQ ID NO:3, the coding sequences for SEQ ID NO:6 given in SEQ ID NO:5, the coding sequence for SEQ ID NO:8 given in SEQ ID NO:7, the coding sequence for SEQ ID NO:10 given in SEQ ID NO:9, the coding sequence for SEQ ID NO:12 given in SEQ ID NO:11, the coding sequence for SEQ ID NO:14 given in SEQ ID NO:13, the coding sequence for SEQ ID NO:16 given in SEQ ID NO:15, the coding sequence for SEQ ID NO:18 given in SEQ ID NO:17 or is a nucleic acid molecule that is complementary thereto. In this regard, nucleic acid molecules at least 90%, preferably at least 95%, more preferably at least 98% or 99% identical over their entire length to the same are particularly preferred. Preferred embodiments in this respect are nucleic acid molecules that encode polypeptides which retain substantially the same biological function or activity as the INSP052 and INSP055 polypeptides.

The invention also provides a process for detecting a nucleic acid molecule of the invention, comprising the steps of: (a) contacting a nucleic probe according to the invention with a biological sample under hybridizing conditions to form duplexes; and (b) detecting any such duplexes that are formed.

As discussed additionally below in connection with assays that may be utilised according to the invention, a nucleic acid molecule as described above may be used as a hybridization probe for RNA, cDNA or genomic DNA, in order to isolate full-length cDNAs and genomic clones encoding the INSP052 and INSP055 polypeptides and to isolate cDNA and genomic clones of homologous or orthologous genes that have a high sequence similarity to the gene encoding this polypeptide.

In this regard, the following techniques, among others known in the art, may be utilised and are discussed below for purposes of illustration. Methods for DNA sequencing and analysis are well known and are generally available in the art and may, indeed, be used to practice many of the embodiments of the invention discussed herein. Such methods may 5 employ such enzymes as the Klenow fragment of DNA polymerase I, Sequenase (US Biochemical Corp, Cleveland, OH), Taq polymerase (Perkin Elmer), thermostable T7 polymerase (Amersham, Chicago, IL), or combinations of polymerases and proof-reading exonucleases such as those found in the ELONGASE Amplification System marketed by Gibco/BRL (Gaithersburg, MD). Preferably, the sequencing process may be automated 10 using machines such as the Hamilton Micro Lab 2200 (Hamilton, Reno, NV), the Peltier Thermal Cycler (PTC200; MJ Research, Watertown, MA) and the ABI Catalyst and 373 and 377 DNA Sequencers (Perkin Elmer).

One method for isolating a nucleic acid molecule encoding a polypeptide with an equivalent function to that of the INSP052 and INSP055 polypeptides is to probe a 15 genomic or cDNA library with a natural or artificially-designed probe using standard procedures that are recognised in the art (see, for example, "Current Protocols in Molecular Biology", Ausubel *et al.* (eds). Greene Publishing Association and John Wiley Interscience, New York, 1989,1992). Probes comprising at least 15, preferably at least 30, and more preferably at least 50, contiguous bases that correspond to, or are complementary 20 to, nucleic acid sequences from the appropriate encoding gene (SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:11, SEQ ID NO:13, SEQ ID NO:15 or SEQ ID NO:17) are particularly useful probes. Such probes may be labelled with an analytically-detectable reagent to facilitate their identification. Useful reagents include, but are not limited to, radioisotopes, fluorescent dyes and enzymes that are 25 capable of catalysing the formation of a detectable product. Using these probes, the ordinarily skilled artisan will be capable of isolating complementary copies of genomic DNA, cDNA or RNA polynucleotides encoding proteins of interest from human, mammalian or other animal sources and screening such sources for related sequences, for example, for additional members of the family, type and/or subtype.

30 In many cases, isolated cDNA sequences will be incomplete, in that the region encoding the polypeptide will be cut short, normally at the 5' end. Several methods are available to obtain full length cDNAs, or to extend short cDNAs. Such sequences may be extended utilising a partial nucleotide sequence and employing various methods known in the art to

detect upstream sequences such as promoters and regulatory elements. For example, one method which may be employed is based on the method of Rapid Amplification of cDNA Ends (RACE; see, for example, Frohman *et al.*, PNAS USA 85, 8998-9002, 1988). Recent modifications of this technique, exemplified by the MarathonTM technology (Clontech Laboratories Inc.), for example, have significantly simplified the search for longer cDNAs. A slightly different technique, termed "restriction-site" PCR, uses universal primers to retrieve unknown nucleic acid sequence adjacent a known locus (Sarkar, G. (1993) PCR Methods Applic. 2:318-322). Inverse PCR may also be used to amplify or to extend sequences using divergent primers based on a known region (Triglia, T. *et al.* (1988) 5 Nucleic Acids Res. 16:8186). Another method which may be used is capture PCR which involves PCR amplification of DNA fragments adjacent a known sequence in human and yeast artificial chromosome DNA (Lagerstrom, M. *et al.* (1991) PCR Methods Applic., 1, 111-119). Another method which may be used to retrieve unknown sequences is that of Parker, J.D. *et al.* (1991); Nucleic Acids Res. 19:3055-3060). Additionally, one may use 10 PCR, nested primers, and PromoterFinderTM libraries to walk genomic DNA (Clontech, Palo Alto, CA). This process avoids the need to screen libraries and is useful in finding 15 intron/exon junctions.

When screening for full-length cDNAs, it is preferable to use libraries that have been size-selected to include larger cDNAs. Also, random-primed libraries are preferable, in that 20 they will contain more sequences that contain the 5' regions of genes. Use of a randomly primed library may be especially preferable for situations in which an oligo d(T) library does not yield a full-length cDNA. Genomic libraries may be useful for extension of sequence into 5' non-transcribed regulatory regions.

In one embodiment of the invention, the nucleic acid molecules of the present invention 25 may be used for chromosome localisation. In this technique, a nucleic acid molecule is specifically targeted to, and can hybridize with, a particular location on an individual human chromosome. The mapping of relevant sequences to chromosomes according to the present invention is an important step in the confirmatory correlation of those sequences with the gene-associated disease. Once a sequence has been mapped to a precise 30 chromosomal location, the physical position of the sequence on the chromosome can be correlated with genetic map data. Such data are found in, for example, V. McKusick, Mendelian Inheritance in Man (available on-line through Johns Hopkins University Welch Medical Library). The relationships between genes and diseases that have been mapped to

the same chromosomal region are then identified through linkage analysis (coinheritance of physically adjacent genes). This provides valuable information to investigators searching for disease genes using positional cloning or other gene discovery techniques. Once the disease or syndrome has been crudely localised by genetic linkage to a particular 5 genomic region, any sequences mapping to that area may represent associated or regulatory genes for further investigation. The nucleic acid molecule may also be used to detect differences in the chromosomal location due to translocation, inversion, etc. among normal, carrier, or affected individuals.

The nucleic acid molecules of the present invention are also valuable for tissue 10 localisation. Such techniques allow the determination of expression patterns of the polypeptide in tissues by detection of the mRNAs that encode them. These techniques include *in situ* hybridization techniques and nucleotide amplification techniques, such as PCR. Results from these studies provide an indication of the normal functions of the polypeptide in the organism. In addition, comparative studies of the normal expression 15 pattern of mRNAs with that of mRNAs encoded by a mutant gene provide valuable insights into the role of mutant polypeptides in disease. Such inappropriate expression may be of a temporal, spatial or quantitative nature.

Gene silencing approaches may also be undertaken to down-regulate endogenous expression of a gene encoding a polypeptide of the invention. RNA interference (RNAi) 20 (Elbashir, SM *et al.*, *Nature* 2001, 411, 494-498) is one method of sequence specific post-transcriptional gene silencing that may be employed. Short dsRNA oligonucleotides are synthesised *in vitro* and introduced into a cell. The sequence specific binding of these dsRNA oligonucleotides triggers the degradation of target mRNA, reducing or ablating target protein expression.

25 Efficacy of the gene silencing approaches assessed above may be assessed through the measurement of polypeptide expression (for example, by Western blotting), and at the RNA level using TaqMan-based methodologies.

The vectors of the present invention comprise nucleic acid molecules of the invention and may be cloning or expression vectors. The host cells of the invention, which may be 30 transformed, transfected or transduced with the vectors of the invention may be prokaryotic or eukaryotic.

The polypeptides of the invention may be prepared in recombinant form by expression of

their encoding nucleic acid molecules in vectors contained within a host cell. Such expression methods are well known to those of skill in the art and many are described in detail by Sambrook *et al* (*supra*) and Fernandez & Hoeffler (1998, eds. "Gene expression systems. Using nature for the art of expression". Academic Press, San Diego, London, 5 Boston, New York, Sydney, Tokyo, Toronto).

Generally, any system or vector that is suitable to maintain, propagate or express nucleic acid molecules to produce a polypeptide in the required host may be used. The appropriate nucleotide sequence may be inserted into an expression system by any of a variety of well-known and routine techniques, such as, for example, those described in Sambrook *et al.*, 10 (*supra*). Generally, the encoding gene can be placed under the control of a control element such as a promoter, ribosome binding site (for bacterial expression) and, optionally, an operator, so that the DNA sequence encoding the desired polypeptide is transcribed into RNA in the transformed host cell.

Examples of suitable expression systems include, for example, chromosomal, episomal and 15 virus-derived systems, including, for example, vectors derived from: bacterial plasmids, bacteriophage, transposons, yeast episomes, insertion elements, yeast chromosomal elements, viruses such as baculoviruses, papova viruses such as SV40, vaccinia viruses, adenoviruses, fowl pox viruses, pseudorabies viruses and retroviruses, or combinations thereof, such as those derived from plasmid and bacteriophage genetic elements, including 20 cosmids and phagemids. Human artificial chromosomes (HACs) may also be employed to deliver larger fragments of DNA than can be contained and expressed in a plasmid.

Particularly suitable expression systems include microorganisms such as bacteria transformed with recombinant bacteriophage, plasmid or cosmid DNA expression vectors; yeast transformed with yeast expression vectors; insect cell systems infected with virus 25 expression vectors (for example, baculovirus); plant cell systems transformed with virus expression vectors (for example, cauliflower mosaic virus, CaMV; tobacco mosaic virus, TMV) or with bacterial expression vectors (for example, Ti or pBR322 plasmids); or animal cell systems. Cell-free translation systems can also be employed to produce the polypeptides of the invention.

30 Introduction of nucleic acid molecules encoding a polypeptide of the present invention into host cells can be effected by methods described in many standard laboratory manuals, such as Davis *et al.*, Basic Methods in Molecular Biology (1986) and Sambrook *et al.*,*[supra]*.

Particularly suitable methods include calcium phosphate transfection, DEAE-dextran mediated transfection, transvection, microinjection, cationic lipid-mediated transfection, electroporation, transduction, scrape loading, ballistic introduction or infection (see Sambrook *et al.*, 1989 [*supra*]; Ausubel *et al.*, 1991 [*supra*]; Spector, Goldman & 5 Leinwald, 1998). In eukaryotic cells, expression systems may either be transient (for example, episomal) or permanent (chromosomal integration) according to the needs of the system.

The encoding nucleic acid molecule may or may not include a sequence encoding a control sequence, such as a signal peptide or leader sequence, as desired, for example, for secretion 10 of the translated polypeptide into the lumen of the endoplasmic reticulum, into the periplasmic space or into the extracellular environment. These signals may be endogenous to the polypeptide or they may be heterologous signals. Leader sequences can be removed by the bacterial host in post-translational processing.

In addition to control sequences, it may be desirable to add regulatory sequences that allow 15 for regulation of the expression of the polypeptide relative to the growth of the host cell. Examples of regulatory sequences are those which cause the expression of a gene to be increased or decreased in response to a chemical or physical stimulus, including the presence of a regulatory compound or to various temperature or metabolic conditions. Regulatory sequences are those non-translated regions of the vector, such as enhancers, 20 promoters and 5' and 3' untranslated regions. These interact with host cellular proteins to carry out transcription and translation. Such regulatory sequences may vary in their strength and specificity. Depending on the vector system and host utilised, any number of suitable transcription and translation elements, including constitutive and inducible promoters, may be used. For example, when cloning in bacterial systems, inducible 25 promoters such as the hybrid lacZ promoter of the Bluescript phagemid (Stratagene, LaJolla, CA) or pSportTM plasmid (Gibco BRL) and the like may be used. The baculovirus polyhedrin promoter may be used in insect cells. Promoters or enhancers derived from the genomes of plant cells (for example, heat shock, RUBISCO and storage protein genes) or from plant viruses (for example, viral promoters or leader sequences) may be cloned into 30 the vector. In mammalian cell systems, promoters from mammalian genes or from mammalian viruses are preferable. If it is necessary to generate a cell line that contains multiple copies of the sequence, vectors based on SV40 or EBV may be used with an appropriate selectable marker.

An expression vector is constructed so that the particular nucleic acid coding sequence is located in the vector with the appropriate regulatory sequences, the positioning and orientation of the coding sequence with respect to the regulatory sequences being such that the coding sequence is transcribed under the "control" of the regulatory sequences, i.e.,

- 5 RNA polymerase which binds to the DNA molecule at the control sequences transcribes the coding sequence. In some cases it may be necessary to modify the sequence so that it may be attached to the control sequences with the appropriate orientation; i.e., to maintain the reading frame.

The control sequences and other regulatory sequences may be ligated to the nucleic acid
10 coding sequence prior to insertion into a vector. Alternatively, the coding sequence can be cloned directly into an expression vector that already contains the control sequences and an appropriate restriction site.

For long-term, high-yield production of a recombinant polypeptide, stable expression is preferred. For example, cell lines which stably express the polypeptide of interest may be
15 transformed using expression vectors which may contain viral origins of replication and/or endogenous expression elements and a selectable marker gene on the same or on a separate vector. Following the introduction of the vector, cells may be allowed to grow for 1-2 days in an enriched media before they are switched to selective media. The purpose of the selectable marker is to confer resistance to selection, and its presence allows growth and
20 recovery of cells that successfully express the introduced sequences. Resistant clones of stably transformed cells may be proliferated using tissue culture techniques appropriate to the cell type.

Mammalian cell lines available as hosts for expression are known in the art and include many immortalised cell lines available from the American Type Culture Collection
25 (ATCC) including, but not limited to, Chinese hamster ovary (CHO), HeLa, baby hamster kidney (BHK), monkey kidney (COS), C127, 3T3, BHK, HEK 293, Bowes melanoma and human hepatocellular carcinoma (for example Hep G2) cells and a number of other cell lines.

In the baculovirus system, the materials for baculovirus/insect cell expression systems are
30 commercially available in kit form from, inter alia, Invitrogen, San Diego CA (the "MaxBac" kit). These techniques are generally known to those skilled in the art and are described fully in Summers and Smith, Texas Agricultural Experiment Station Bulletin No.

1555 (1987). Particularly suitable host cells for use in this system include insect cells such as *Drosophila S2* and *Spodoptera Sf9* cells.

There are many plant cell culture and whole plant genetic expression systems known in the art. Examples of suitable plant cellular genetic expression systems include those described 5 in US 5,693,506; US 5,659,122; and US 5,608,143. Additional examples of genetic expression in plant cell culture has been described by Zenk, *Phytochemistry* 30, 3861-3863 (1991).

In particular, all plants from which protoplasts can be isolated and cultured to give whole regenerated plants can be utilised, so that whole plants are recovered which contain the 10 transferred gene. Practically all plants can be regenerated from cultured cells or tissues, including but not limited to all major species of sugar cane, sugar beet, cotton, fruit and other trees, legumes and vegetables.

Examples of particularly preferred bacterial host cells include streptococci, staphylococci, *E. coli*, *Streptomyces* and *Bacillus subtilis* cells.

15 Examples of particularly suitable host cells for fungal expression include yeast cells (for example, *S. cerevisiae*) and *Aspergillus* cells.

Any number of selection systems are known in the art that may be used to recover transformed cell lines. Examples include the herpes simplex virus thymidine kinase (Wigler, M. *et al.* (1977) *Cell* 11:223-32) and adenine phosphoribosyltransferase (Lowy, I. 20 *et al.* (1980) *Cell* 22:817-23) genes that can be employed in tk- or apt \pm cells, respectively.

Also, antimetabolite, antibiotic or herbicide resistance can be used as the basis for selection; for example, dihydrofolate reductase (DHFR) that confers resistance to methotrexate (Wigler, M. *et al.* (1980) *Proc. Natl. Acad. Sci.* 77:3567-70); npt, which confers resistance to the aminoglycosides neomycin and G-418 (Colbere-Garapin, F. *et al* 25 (1981) *J. Mol. Biol.* 150:1-14) and als or pat, which confer resistance to chlorsulfuron and phosphinotricin acetyltransferase, respectively. Additional selectable genes have been described, examples of which will be clear to those of skill in the art.

Although the presence or absence of marker gene expression suggests that the gene of interest is also present, its presence and expression may need to be confirmed. For 30 example, if the relevant sequence is inserted within a marker gene sequence, transformed cells containing the appropriate sequences can be identified by the absence of marker gene

function. Alternatively, a marker gene can be placed in tandem with a sequence encoding a polypeptide of the invention under the control of a single promoter. Expression of the marker gene in response to induction or selection usually indicates expression of the tandem gene as well.

5 Alternatively, host cells that contain a nucleic acid sequence encoding a polypeptide of the invention and which express said polypeptide may be identified by a variety of procedures known to those of skill in the art. These procedures include, but are not limited to, DNA-DNA or DNA-RNA hybridizations and protein bioassays, for example, fluorescence activated cell sorting (FACS) or immunoassay techniques (such as the enzyme-linked 10 immunosorbent assay [ELISA] and radioimmunoassay [RIA]), that include membrane, solution, or chip based technologies for the detection and/or quantification of nucleic acid or protein (see Hampton, R. *et al.* (1990) Serological Methods, a Laboratory Manual, APS Press, St Paul, MN) and Maddox, D.E. *et al.* (1983) *J. Exp. Med.*, 158, 1211-1216).

A wide variety of labels and conjugation techniques are known by those skilled in the art 15 and may be used in various nucleic acid and amino acid assays. Means for producing labelled hybridization or PCR probes for detecting sequences related to nucleic acid molecules encoding polypeptides of the present invention include oligolabelling, nick translation, end-labelling or PCR amplification using a labelled polynucleotide. Alternatively, the sequences encoding the polypeptide of the invention may be cloned into 20 a vector for the production of an mRNA probe. Such vectors are known in the art, are commercially available, and may be used to synthesise RNA probes *in vitro* by addition of an appropriate RNA polymerase such as T7, T3 or SP6 and labelled nucleotides. These procedures may be conducted using a variety of commercially available kits (Pharmacia & Upjohn, (Kalamazoo, MI); Promega (Madison WI); and U.S. Biochemical Corp., 25 Cleveland, OH)).

Suitable reporter molecules or labels, which may be used for ease of detection, include radionuclides, enzymes and fluorescent, chemiluminescent or chromogenic agents as well as substrates, cofactors, inhibitors, magnetic particles, and the like.

Nucleic acid molecules according to the present invention may also be used to create 30 transgenic animals, particularly rodent animals. Such transgenic animals form a further aspect of the present invention. This may be done locally by modification of somatic cells, or by germ line therapy to incorporate heritable modifications. Such transgenic animals

may be particularly useful in the generation of animal models for drug molecules effective as modulators of the polypeptides of the present invention.

The polypeptide can be recovered and purified from recombinant cell cultures by well-known methods including ammonium sulphate or ethanol precipitation, acid extraction, 5 anion or cation exchange chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography, hydroxylapatite chromatography and lectin chromatography. High performance liquid chromatography is particularly useful for purification. Well known techniques for refolding proteins may be employed to regenerate an active conformation when the polypeptide is denatured during 10 isolation and or purification.

Specialised vector constructions may also be used to facilitate purification of proteins, as desired, by joining sequences encoding the polypeptides of the invention to a nucleotide sequence encoding a polypeptide domain that will facilitate purification of soluble proteins. Examples of such purification-facilitating domains include metal chelating 15 peptides such as histidine-tryptophan modules that allow purification on immobilised metals, protein A domains that allow purification on immobilised immunoglobulin, and the domain utilised in the FLAGS extension/affinity purification system (Immunex Corp., Seattle, WA). The inclusion of cleavable linker sequences such as those specific for Factor XA or enterokinase (Invitrogen, San Diego, CA) between the purification domain and the 20 polypeptide of the invention may be used to facilitate purification. One such expression vector provides for expression of a fusion protein containing the polypeptide of the invention fused to several histidine residues preceding a thioredoxin or an enterokinase cleavage site. The histidine residues facilitate purification by IMAC (immobilised metal ion affinity chromatography as described in Porath, J. *et al.* (1992), *Prot. Exp. Purif.* 3: 25 263-281) while the thioredoxin or enterokinase cleavage site provides a means for purifying the polypeptide from the fusion protein. A discussion of vectors which contain fusion proteins is provided in Kroll, D.J. *et al.* (1993; *DNA Cell Biol.* 12:441-453).

If the polypeptide is to be expressed for use in screening assays, generally it is preferred that it be produced at the surface of the host cell in which it is expressed. In this event, the 30 host cells may be harvested prior to use in the screening assay, for example using techniques such as fluorescence activated cell sorting (FACS) or immunoaffinity techniques. If the polypeptide is secreted into the medium, the medium can be recovered in

order to recover and purify the expressed polypeptide. If polypeptide is produced intracellularly, the cells must first be lysed before the polypeptide is recovered.

The polypeptide of the invention can be used to screen libraries of compounds in any of a variety of drug screening techniques. Such compounds may activate (agonise) or inhibit 5 (antagonise) the level of expression of the gene or the activity of the polypeptide of the invention and form a further aspect of the present invention. Preferred compounds are effective to alter the expression of a natural gene which encodes a polypeptide of the first aspect of the invention or to regulate the activity of a polypeptide of the first aspect of the invention.

10 Agonist or antagonist compounds may be isolated from, for example, cells, cell-free preparations, chemical libraries or natural product mixtures. These agonists or antagonists may be natural or modified substrates, ligands, enzymes, receptors or structural or functional mimetics. For a suitable review of such screening techniques, see Coligan *et al.*, Current Protocols in Immunology 1(2):Chapter 5 (1991).

15 Compounds that are most likely to be good antagonists are molecules that bind to the polypeptide of the invention without inducing the biological effects of the polypeptide upon binding to it. Potential antagonists include small organic molecules, peptides, polypeptides and antibodies that bind to the polypeptide of the invention and thereby inhibit or extinguish its activity. In this fashion, binding of the polypeptide to normal 20 cellular binding molecules may be inhibited, such that the normal biological activity of the polypeptide is prevented.

The polypeptide of the invention that is employed in such a screening technique may be free in solution, affixed to a solid support, borne on a cell surface or located intracellularly. In general, such screening procedures may involve using appropriate cells or cell 25 membranes that express the polypeptide that are contacted with a test compound to observe binding, or stimulation or inhibition of a functional response. The functional response of the cells contacted with the test compound is then compared with control cells that were not contacted with the test compound. Such an assay may assess whether the test compound results in a signal generated by activation of the polypeptide, using an 30 appropriate detection system. Inhibitors of activation are generally assayed in the presence of a known agonist and the effect on activation by the agonist in the presence of the test compound is observed.

A preferred method for identifying an agonist or antagonist compound of a polypeptide of the present invention comprises:

- (a) contacting a cell expressing on the surface thereof the polypeptide according to the first aspect of the invention, the polypeptide being associated with a second component capable of providing a detectable signal in response to the binding of a compound to the polypeptide, with a compound to be screened under conditions to permit binding to the polypeptide; and
- (b) determining whether the compound binds to and activates or inhibits the polypeptide by measuring the level of a signal generated from the interaction of the compound with the polypeptide.

A further preferred method for identifying an agonist or antagonist of a polypeptide of the invention comprises:

- (a) contacting a cell expressing on the surface thereof the polypeptide, the polypeptide being associated with a second component capable of providing a detectable signal in response to the binding of a compound to the polypeptide, with a compound to be screened under conditions to permit binding to the polypeptide; and
- (b) determining whether the compound binds to and activates or inhibits the polypeptide by comparing the level of a signal generated from the interaction of the compound with the polypeptide with the level of a signal in the absence of the compound.

- 20 In further preferred embodiments, the general methods that are described above may further comprise conducting the identification of agonist or antagonist in the presence of labelled or unlabelled ligand for the polypeptide.

In another embodiment of the method for identifying agonist or antagonist of a polypeptide of the present invention comprises:

- 25 determining the inhibition of binding of a ligand to cells which have a polypeptide of the invention on the surface thereof, or to cell membranes containing such a polypeptide, in the presence of a candidate compound under conditions to permit binding to the polypeptide, and determining the amount of ligand bound to the polypeptide. A compound capable of causing reduction of binding of a ligand is considered to be an agonist or antagonist.
- 30 Preferably the ligand is labelled.

More particularly, a method of screening for a polypeptide antagonist or agonist compound

comprises the steps of:

- (a) incubating a labelled ligand with a whole cell expressing a polypeptide according to the invention on the cell surface, or a cell membrane containing a polypeptide of the invention;
- (b) measuring the amount of labelled ligand bound to the whole cell or the cell membrane;
- 5 (c) adding a candidate compound to a mixture of labelled ligand and the whole cell or the cell membrane of step (a) and allowing the mixture to attain equilibrium;
- (d) measuring the amount of labelled ligand bound to the whole cell or the cell membrane after step (c); and
- (e) comparing the difference in the labelled ligand bound in step (b) and (d), such that the

10 compound which causes the reduction in binding in step (d) is considered to be an agonist or antagonist.

In certain of the embodiments described above, simple binding assays may be used, in which the adherence of a test compound to a surface bearing the polypeptide is detected by means of a label directly or indirectly associated with the test compound or in an assay

15 involving competition with a labelled competitor. In another embodiment, competitive drug screening assays may be used, in which neutralising antibodies that are capable of binding the polypeptide specifically compete with a test compound for binding. In this manner, the antibodies can be used to detect the presence of any test compound that possesses specific binding affinity for the polypeptide.

20 Persons skilled in the art will be able to devise assays for identifying modulators of a polypeptide of the invention. Of interest in this regard is Lokker NA et al J Biol Chem 1997 Dec 26;272(52):33037-44 which reports an example of an assay to identify antagonists (in this case neutralizing antibodies).

Assays may also be designed to detect the effect of added test compounds on the

25 production of mRNA encoding the polypeptide in cells. For example, an ELISA may be constructed that measures secreted or cell-associated levels of polypeptide using monoclonal or polyclonal antibodies by standard methods known in the art, and this can be used to search for compounds that may inhibit or enhance the production of the polypeptide from suitably manipulated cells or tissues. The formation of binding

30 complexes between the polypeptide and the compound being tested may then be measured.

Assay methods that are also included within the terms of the present invention are those that involve the use of the genes and polypeptides of the invention in overexpression or ablation assays. Such assays involve the manipulation of levels of these genes/polypeptides in cells and assessment of the impact of this manipulation event on the physiology of the 5 manipulated cells. For example, such experiments reveal details of signalling and metabolic pathways in which the particular genes/polypeptides are implicated, generate information regarding the identities of polypeptides with which the studied polypeptides interact and provide clues as to methods by which related genes and proteins are regulated.

Another technique for drug screening which may be used provides for high throughput 10 screening of compounds having suitable binding affinity to the polypeptide of interest (see International patent application WO84/03564). In this method, large numbers of different small test compounds are synthesised on a solid substrate, which may then be reacted with the polypeptide of the invention and washed. One way of immobilising the polypeptide is to use non-neutralising antibodies. Bound polypeptide may then be detected using methods 15 that are well known in the art. Purified polypeptide can also be coated directly onto plates for use in the aforementioned drug screening techniques.

The polypeptide of the invention may be used to identify membrane-bound or soluble receptors, through standard receptor binding techniques that are known in the art, such as ligand binding and crosslinking assays in which the polypeptide is labelled with a 20 radioactive isotope, is chemically modified, or is fused to a peptide sequence that facilitates its detection or purification, and incubated with a source of the putative receptor (for example, a composition of cells, cell membranes, cell supernatants, tissue extracts, or bodily fluids). The efficacy of binding may be measured using biophysical techniques such as surface plasmon resonance and spectroscopy. Binding assays may be used for the 25 purification and cloning of the receptor, but may also identify agonists and antagonists of the polypeptide, that compete with the binding of the polypeptide to its receptor. Standard methods for conducting screening assays are well understood in the art.

The invention also includes a screening kit useful in the methods for identifying agonists, antagonists, ligands, receptors, substrates, enzymes, that are described above.

30 The invention includes the agonists, antagonists, ligands, receptors, substrates and enzymes, and other compounds which modulate the activity or antigenicity of the polypeptide of the invention discovered by the methods that are described above.

The invention also provides pharmaceutical compositions comprising a polypeptide, nucleic acid, ligand or compound of the invention in combination with a suitable pharmaceutical carrier. These compositions may be suitable as therapeutic or diagnostic reagents, as vaccines, or as other immunogenic compositions, as outlined in detail below.

- 5 According to the terminology used herein, a composition containing a polypeptide, nucleic acid, ligand or compound [X] is "substantially free of" impurities [herein, Y] when at least 85% by weight of the total X+Y in the composition is X. Preferably, X comprises at least about 90% by weight of the total of X+Y in the composition, more preferably at least about 95%, 98% or even 99% by weight.
- 10 The pharmaceutical compositions should preferably comprise a therapeutically effective amount of the polypeptide, nucleic acid molecule, ligand, or compound of the invention. The term "therapeutically effective amount" as used herein refers to an amount of a therapeutic agent needed to treat, ameliorate, or prevent a targeted disease or condition, or to exhibit a detectable therapeutic or preventative effect. For any compound, the 15 therapeutically effective dose can be estimated initially either in cell culture assays, for example, of neoplastic cells, or in animal models, usually mice, rabbits, dogs, or pigs. The animal model may also be used to determine the appropriate concentration range and route of administration. Such information can then be used to determine useful doses and routes for administration in humans.
- 20 The precise effective amount for a human subject will depend upon the severity of the disease state, general health of the subject, age, weight, and gender of the subject, diet, time and frequency of administration, drug combination(s), reaction sensitivities, and tolerance/response to therapy. This amount can be determined by routine experimentation and is within the judgement of the clinician. Generally, an effective dose will be from 0.01 25 mg/kg to 50 mg/kg, preferably 0.05 mg/kg to 10 mg/kg. Compositions may be administered individually to a patient or may be administered in combination with other agents, drugs or hormones.

A pharmaceutical composition may also contain a pharmaceutically acceptable carrier, for administration of a therapeutic agent. Such carriers include antibodies and other 30 polypeptides, genes and other therapeutic agents such as liposomes, provided that the carrier does not itself induce the production of antibodies harmful to the individual receiving the composition, and which may be administered without undue toxicity.

Suitable carriers may be large, slowly metabolised macromolecules such as proteins, polysaccharides, polylactic acids, polyglycolic acids, polymeric amino acids, amino acid copolymers and inactive virus particles.

5 Pharmaceutically acceptable salts can be used therein, for example, mineral acid salts such as hydrochlorides, hydrobromides, phosphates, sulphates, and the like; and the salts of organic acids such as acetates, propionates, malonates, benzoates, and the like. A thorough discussion of pharmaceutically acceptable carriers is available in Remington's Pharmaceutical Sciences (Mack Pub. Co., N.J. 1991).

Pharmaceutically acceptable carriers in therapeutic compositions may additionally contain

10 liquids such as water, saline, glycerol and ethanol. Additionally, auxiliary substances, such as wetting or emulsifying agents, pH buffering substances, and the like, may be present in such compositions. Such carriers enable the pharmaceutical compositions to be formulated as tablets, pills, dragees, capsules, liquids, gels, syrups, slurries, suspensions, and the like, for ingestion by the patient.

15 Once formulated, the compositions of the invention can be administered directly to the subject. The subjects to be treated can be animals; in particular, human subjects can be treated.

The pharmaceutical compositions utilised in this invention may be administered by any number of routes including, but not limited to, oral, intravenous, intramuscular, intra-
20 arterial, intramedullary, intrathecal, intraventricular, transdermal or transcutaneous applications (for example, see WO98/20734), subcutaneous, intraperitoneal, intranasal, enteral, topical, sublingual, intravaginal or rectal means. Gene guns or hyposprays may also be used to administer the pharmaceutical compositions of the invention. Typically, the therapeutic compositions may be prepared as injectables, either as liquid solutions or
25 suspensions; solid forms suitable for solution in, or suspension in, liquid vehicles prior to injection may also be prepared.

Direct delivery of the compositions will generally be accomplished by injection, subcutaneously, intraperitoneally, intravenously or intramuscularly, or delivered to the interstitial space of a tissue. The compositions can also be administered into a lesion.

30 Dosage treatment may be a single dose schedule or a multiple dose schedule.

If the activity of the polypeptide of the invention is in excess in a particular disease state, several approaches are available. One approach comprises administering to a subject an

inhibitor compound (antagonist) as described above, along with a pharmaceutically acceptable carrier in an amount effective to inhibit the function of the polypeptide, such as by blocking the binding of ligands, substrates, enzymes, receptors, or by inhibiting a second signal, and thereby alleviating the abnormal condition. Preferably, such antagonists 5 are antibodies. Most preferably, such antibodies are chimeric and/or humanised to minimise their immunogenicity, as described previously.

In another approach, soluble forms of the polypeptide that retain binding affinity for the ligand, substrate, enzyme, receptor, in question, may be administered. Typically, the polypeptide may be administered in the form of fragments that retain the relevant portions.

- 10 In an alternative approach, expression of the gene encoding the polypeptide can be inhibited using expression blocking techniques, such as the use of antisense nucleic acid molecules (as described above), either internally generated or separately administered. Modifications of gene expression can be obtained by designing complementary sequences or antisense molecules (DNA, RNA, or PNA) to the control, 5' or regulatory regions
- 15 (signal sequence, promoters, enhancers and introns) of the gene encoding the polypeptide. Similarly, inhibition can be achieved using "triple helix" base-pairing methodology. Triple helix pairing is useful because it causes inhibition of the ability of the double helix to open sufficiently for the binding of polymerases, transcription factors, or regulatory molecules. Recent therapeutic advances using triplex DNA have been described in the literature (Gee,
- 20 J.E. *et al.* (1994) In: Huber, B.E. and B.I. Carr, Molecular and Immunologic Approaches, Futura Publishing Co., Mt. Kisco, NY). The complementary sequence or antisense molecule may also be designed to block translation of mRNA by preventing the transcript from binding to ribosomes. Such oligonucleotides may be administered or may be generated *in situ* from expression *in vivo*.
- 25 In addition, expression of the polypeptide of the invention may be prevented by using ribozymes specific to its encoding mRNA sequence. Ribozymes are catalytically active RNAs that can be natural or synthetic (see for example Usman, N, *et al.*, Curr. Opin. Struct. Biol (1996) 6(4), 527-33). Synthetic ribozymes can be designed to specifically cleave mRNAs at selected positions thereby preventing translation of the mRNAs into
- 30 functional polypeptide. Ribozymes may be synthesised with a natural ribose phosphate backbone and natural bases, as normally found in RNA molecules. Alternatively the ribozymes may be synthesised with non-natural backbones, for example, 2'-O-methyl

RNA, to provide protection from ribonuclease degradation and may contain modified bases.

RNA molecules may be modified to increase intracellular stability and half-life. Possible modifications include, but are not limited to, the addition of flanking sequences at the 5' 5 and/or 3' ends of the molecule or the use of phosphorothioate or 2' O-methyl rather than phosphodiesterase linkages within the backbone of the molecule. This concept is inherent in the production of PNAs and can be extended in all of these molecules by the inclusion of non-traditional bases such as inosine, queosine and butosine, as well as acetyl-, methyl-, thio- and similarly modified forms of adenine, cytidine, guanine, thymine and uridine 10 which are not as easily recognised by endogenous endonucleases.

For treating abnormal conditions related to an under-expression of the polypeptide of the invention and its activity, several approaches are also available. One approach comprises administering to a subject a therapeutically effective amount of a compound that activates the polypeptide, i.e., an agonist as described above, to alleviate the abnormal condition. 15 Alternatively, a therapeutic amount of the polypeptide in combination with a suitable pharmaceutical carrier may be administered to restore the relevant physiological balance of polypeptide.

Gene therapy may be employed to effect the endogenous production of the polypeptide by the relevant cells in the subject. Gene therapy is used to treat permanently the inappropriate 20 production of the polypeptide by replacing a defective gene with a corrected therapeutic gene.

Gene therapy of the present invention can occur *in vivo* or *ex vivo*. *Ex vivo* gene therapy requires the isolation and purification of patient cells, the introduction of a therapeutic gene and introduction of the genetically altered cells back into the patient. In contrast, *in vivo* 25 gene therapy does not require isolation and purification of a patient's cells.

The therapeutic gene is typically "packaged" for administration to a patient. Gene delivery vehicles may be non-viral, such as liposomes, or replication-deficient viruses, such as adenovirus as described by Berkner, K.L., in *Curr. Top. Microbiol. Immunol.*, 158, 39-66 (1992) or adeno-associated virus (AAV) vectors as described by Muzyczka, N., in *Curr. 30 Top. Microbiol. Immunol.*, 158, 97-129 (1992) and U.S. Patent No. 5,252,479. For example, a nucleic acid molecule encoding a polypeptide of the invention may be engineered for expression in a replication-defective retroviral vector. This expression

construct may then be isolated and introduced into a packaging cell transduced with a retroviral plasmid vector containing RNA encoding the polypeptide, such that the packaging cell now produces infectious viral particles containing the gene of interest. These producer cells may be administered to a subject for engineering cells *in vivo* and 5 expression of the polypeptide *in vivo* (see Chapter 20, Gene Therapy and other Molecular Genetic-based Therapeutic Approaches, (and references cited therein) in Human Molecular Genetics (1996), T Strachan and A P Read, BIOS Scientific Publishers Ltd).

Another approach is the administration of "naked DNA" in which the therapeutic gene is directly injected into the bloodstream or muscle tissue.

10 In situations in which the polypeptides or nucleic acid molecules of the invention are disease-causing agents, the invention provides that they can be used in vaccines to raise antibodies against the disease causing agent.

Vaccines according to the invention may either be prophylactic (ie. to prevent infection) or therapeutic (ie. to treat disease after infection). Such vaccines comprise immunising 15 antigen(s), immunogen(s), polypeptide(s), protein(s) or nucleic acid, usually in combination with pharmaceutically-acceptable carriers as described above, which include any carrier that does not itself induce the production of antibodies harmful to the individual receiving the composition. Additionally, these carriers may function as immunostimulating agents ("adjuvants"). Furthermore, the antigen or immunogen may be conjugated to a 20 bacterial toxoid, such as a toxoid from diphtheria, tetanus, cholera, H. pylori, and other pathogens.

Since polypeptides may be broken down in the stomach, vaccines comprising polypeptides are preferably administered parenterally (for instance, subcutaneous, intramuscular, intravenous, or intradermal injection). Formulations suitable for parenteral administration 25 include aqueous and non-aqueous sterile injection solutions which may contain anti-oxidants, buffers, bacteriostats and solutes which render the formulation isotonic with the blood of the recipient, and aqueous and non-aqueous sterile suspensions which may include suspending agents or thickening agents.

The vaccine formulations of the invention may be presented in unit-dose or multi-dose 30 containers. For example, sealed ampoules and vials and may be stored in a freeze-dried condition requiring only the addition of the sterile liquid carrier immediately prior to use. The dosage will depend on the specific activity of the vaccine and can be readily

determined by routine experimentation.

Genetic delivery of antibodies that bind to polypeptides according to the invention may also be effected, for example, as described in International patent application WO98/55607.

5 The technology referred to as jet injection (see, for example, www.powderject.com) may also be useful in the formulation of vaccine compositions.

A number of suitable methods for vaccination and vaccine delivery systems are described in International patent application WO00/29428.

This invention also relates to the use of nucleic acid molecules according to the present 10 invention as diagnostic reagents. Detection of a mutated form of the gene characterised by the nucleic acid molecules of the invention which is associated with a dysfunction will provide a diagnostic tool that can add to, or define, a diagnosis of a disease, or susceptibility to a disease, which results from under-expression, over-expression or altered spatial or temporal expression of the gene. Individuals carrying mutations in the gene may 15 be detected at the DNA level by a variety of techniques.

Nucleic acid molecules for diagnosis may be obtained from a subject's cells, such as from 20 blood, urine, saliva, tissue biopsy or autopsy material. The genomic DNA may be used directly for detection or may be amplified enzymatically by using PCR, ligase chain reaction (LCR), strand displacement amplification (SDA), or other amplification techniques (see Saiki *et al.*, *Nature*, 324, 163-166 (1986); Bej, *et al.*, *Crit. Rev. Biochem. Molec. Biol.*, 26, 301-334 (1991); Birkenmeyer *et al.*, *J. Virol. Meth.*, 35, 117-126 (1991); Van Brunt, *J. Bio/Technology*, 8, 291-294 (1990)) prior to analysis.

In one embodiment, this aspect of the invention provides a method of diagnosing a disease 25 in a patient, comprising assessing the level of expression of a natural gene encoding a polypeptide according to the invention and comparing said level of expression to a control level, wherein a level that is different to said control level is indicative of disease. The method may comprise the steps of:

a)contacting a sample of tissue from the patient with a nucleic acid probe under stringent conditions that allow the formation of a hybrid complex between a nucleic acid 30 molecule of the invention and the probe;

b)contacting a control sample with said probe under the same conditions used in step a);

c)and detecting the presence of hybrid complexes in said samples; wherein detection of levels of the hybrid complex in the patient sample that differ from levels of the hybrid complex in the control sample is indicative of disease.

A further aspect of the invention comprises a diagnostic method comprising the steps of:

- 5 a)obtaining a tissue sample from a patient being tested for disease;
- b)isolating a nucleic acid molecule according to the invention from said tissue sample; and
- c)diagnosing the patient for disease by detecting the presence of a mutation in the nucleic acid molecule which is associated with disease.

To aid the detection of nucleic acid molecules in the above-described methods, an 10 amplification step, for example using PCR, may be included.

Deletions and insertions can be detected by a change in the size of the amplified product in comparison to the normal genotype. Point mutations can be identified by hybridizing amplified DNA to labelled RNA of the invention or alternatively, labelled antisense DNA sequences of the invention. Perfectly-matched sequences can be distinguished from 15 mismatched duplexes by RNase digestion or by assessing differences in melting temperatures. The presence or absence of the mutation in the patient may be detected by contacting DNA with a nucleic acid probe that hybridises to the DNA under stringent conditions to form a hybrid double-stranded molecule, the hybrid double-stranded molecule having an unhybridised portion of the nucleic acid probe strand at any portion 20 corresponding to a mutation associated with disease; and detecting the presence or absence of an unhybridised portion of the probe strand as an indication of the presence or absence of a disease-associated mutation in the corresponding portion of the DNA strand.

Such diagnostics are particularly useful for prenatal and even neonatal testing.

Point mutations and other sequence differences between the reference gene and "mutant" 25 genes can be identified by other well-known techniques, such as direct DNA sequencing or single-strand conformational polymorphism, (see Orita *et al.*, Genomics, 5, 874-879 (1989)). For example, a sequencing primer may be used with double-stranded PCR product or a single-stranded template molecule generated by a modified PCR. The sequence determination is performed by conventional procedures with radiolabelled nucleotides or 30 by automatic sequencing procedures with fluorescent-tags. Cloned DNA segments may also be used as probes to detect specific DNA segments. The sensitivity of this method is

greatly enhanced when combined with PCR. Further, point mutations and other sequence variations, such as polymorphisms, can be detected as described above, for example, through the use of allele-specific oligonucleotides for PCR amplification of sequences that differ by single nucleotides.

- 5 DNA sequence differences may also be detected by alterations in the electrophoretic mobility of DNA fragments in gels, with or without denaturing agents, or by direct DNA sequencing (for example, Myers *et al.*, *Science* (1985) 230:1242). Sequence changes at specific locations may also be revealed by nuclease protection assays, such as RNase and S1 protection or the chemical cleavage method (see Cotton *et al.*, *Proc. Natl. Acad. Sci.*
- 10 USA (1985) 85: 4397-4401).

In addition to conventional gel electrophoresis and DNA sequencing, mutations such as microdeletions, aneuploidies, translocations, inversions, can also be detected by *in situ* analysis (see, for example, Keller *et al.*, *DNA Probes*, 2nd Ed., Stockton Press, New York, N.Y., USA (1993)), that is, DNA or RNA sequences in cells can be analysed for mutations without need for their isolation and/or immobilisation onto a membrane. Fluorescence *in situ* hybridization (FISH) is presently the most commonly applied method and numerous reviews of FISH have appeared (see, for example, Trachuck *et al.*, *Science*, 250, 559-562 (1990), and Trask *et al.*, *Trends, Genet.*, 7, 149-154 (1991)).

In another embodiment of the invention, an array of oligonucleotide probes comprising a nucleic acid molecule according to the invention can be constructed to conduct efficient screening of genetic variants, mutations and polymorphisms. Array technology methods are well known and have general applicability and can be used to address a variety of questions in molecular genetics including gene expression, genetic linkage, and genetic variability (see for example: M.Chee *et al.*, *Science* (1996), Vol 274, pp 610-613).

- 25 In one embodiment, the array is prepared and used according to the methods described in PCT application WO95/11995 (Chee *et al.*); Lockhart, D. J. *et al.* (1996) *Nat. Biotech.* 14: 1675-1680); and Schena, M. *et al.* (1996) *Proc. Natl. Acad. Sci.* 93: 10614-10619. Oligonucleotide pairs may range from two to over one million. The oligomers are synthesized at designated areas on a substrate using a light-directed chemical process. The
- 30 substrate may be paper, nylon or other type of membrane, filter, chip, glass slide or any other suitable solid support. In another aspect, an oligonucleotide may be synthesized on the surface of the substrate by using a chemical coupling procedure and an ink jet

application apparatus, as described in PCT application W095/251116 (Baldeschweiler *et al*). In another aspect, a "gridded" array analogous to a dot (or slot) blot may be used to arrange and link cDNA fragments or oligonucleotides to the surface of a substrate using a vacuum system, thermal, UV, mechanical or chemical bonding procedures. An array, such

5 as those described above, may be produced by hand or by using available devices (slot blot or dot blot apparatus), materials (any suitable solid support), and machines (including robotic instruments), and may contain 8, 24, 96, 384, 1536 or 6144 oligonucleotides, or any other number between two and over one million which lends itself to the efficient use of commercially-available instrumentation.

10 In addition to the methods discussed above, diseases may be diagnosed by methods comprising determining, from a sample derived from a subject, an abnormally decreased or increased level of polypeptide or mRNA. Decreased or increased expression can be measured at the RNA level using any of the methods well known in the art for the quantitation of polynucleotides, such as, for example, nucleic acid amplification, for

15 instance PCR, RT-PCR, RNase protection, Northern blotting and other hybridization methods.

Assay techniques that can be used to determine levels of a polypeptide of the present invention in a sample derived from a host are well-known to those of skill in the art and are discussed in some detail above (including radioimmunoassays, competitive-binding assays,

20 Western Blot analysis and ELISA assays). This aspect of the invention provides a diagnostic method which comprises the steps of: (a) contacting a ligand as described above with a biological sample under conditions suitable for the formation of a ligand-polypeptide complex; and (b) detecting said complex.

Protocols such as ELISA, RIA, and FACS for measuring polypeptide levels may

25 additionally provide a basis for diagnosing altered or abnormal levels of polypeptide expression. Normal or standard values for polypeptide expression are established by combining body fluids or cell extracts taken from normal mammalian subjects, preferably humans, with antibody to the polypeptide under conditions suitable for complex formation. The amount of standard complex formation may be quantified by various methods, such as

30 by photometric means.

Antibodies which specifically bind to a polypeptide of the invention may be used for the diagnosis of conditions or diseases characterised by expression of the polypeptide, or in

assays to monitor patients being treated with the polypeptides, nucleic acid molecules, ligands and other compounds of the invention. Antibodies useful for diagnostic purposes may be prepared in the same manner as those described above for therapeutics. Diagnostic assays for the polypeptide include methods that utilise the antibody and a label to detect 5 the polypeptide in human body fluids or extracts of cells or tissues. The antibodies may be used with or without modification, and may be labelled by joining them, either covalently or non-covalently, with a reporter molecule. A wide variety of reporter molecules known in the art may be used, several of which are described above.

Quantities of polypeptide expressed in subject, control and disease samples from biopsied 10 tissues are compared with the standard values. Deviation between standard and subject values establishes the parameters for diagnosing disease. Diagnostic assays may be used to distinguish between absence, presence, and excess expression of polypeptide and to monitor regulation of polypeptide levels during therapeutic intervention. Such assays may also be used to evaluate the efficacy of a particular therapeutic treatment regimen in animal 15 studies, in clinical trials or in monitoring the treatment of an individual patient.

A diagnostic kit of the present invention may comprise:

(a) a nucleic acid molecule of the present invention;
(b) a polypeptide of the present invention; or
(c) a ligand of the present invention.

20 In one aspect of the invention, a diagnostic kit may comprise a first container containing a nucleic acid probe that hybridises under stringent conditions with a nucleic acid molecule according to the invention; a second container containing primers useful for amplifying the nucleic acid molecule; and instructions for using the probe and primers for facilitating the diagnosis of disease. The kit may further comprise a third container holding an agent for 25 digesting unhybridised RNA.

In an alternative aspect of the invention, a diagnostic kit may comprise an array of nucleic acid molecules, at least one of which may be a nucleic acid molecule according to the invention.

To detect polypeptide according to the invention, a diagnostic kit may comprise one or 30 more antibodies that bind to a polypeptide according to the invention; and a reagent useful for the detection of a binding reaction between the antibody and the polypeptide.

Such kits will be of use in diagnosing a disease or susceptibility to disease, including, but not limited to, diseases including, but not limited to, cell proliferative disorders, autoimmune/inflammatory disorders, cardiovascular disorders, neurological and psychiatric disorders, developmental disorders, genetic disorders, metabolic disorders, 5 infections and other pathological conditions. These diseases preferably include neoplasm, cancer, brain tumour, glioma, bone tumor, lung tumor, breast tumour, prostate tumour, colon tumour, hemangioma, myeloproliferative disorder, leukemia, hematological disease, neutropenia, thrombocytopenia, angiogenesis disorders, dermatological disease, ageing, wounds, burns, fibrosis, cardiovascular disease, restenosis, heart disease, peripheral 10 vascular disease, coronary artery disease, oedema, thromboembolism, dysmenorrhea, endometriosis, pre-eclampsia, lung disease, COPD, asthma bone disease, renal disease, glomerulonephritis, liver disease, Crohn's disease, gastritis, ulcerative colitis, ulcer, immune disorder, autoimmune disease, arthritis, rheumatoid arthritis, psoriasis, epidermolysis bullosa, systemic lupus erythematosus, ankylosing spondylitis, Lyme 15 disease, multiple sclerosis, neurodegeneration, stroke, brain/spinal cord injury, Alzheimer's disease, Parkinson's disease, motor neurone disease, neuromuscular disease, HIV, AIDS, cytomegalovirus infection, fungal infection, ocular disorder, macular degeneration, glaucoma, diabetic retinopathy, ocular hypertension and other conditions in which immunoglobulin domain containing cell recognition molecules are implicated.

20 Various aspects and embodiments of the present invention will now be described in more detail by way of example, with particular reference to the INSP052 and INSP055 polypeptides.

It will be appreciated that modification of detail may be made without departing from the scope of the invention.

25 **Brief description of the Figures**

Figure 1: Results from BLAST against NCBI non-redundant database using full-length INSP052 polypeptide sequence.

Figure 2: Alignment generated by BLAST between the full-length INSP052 polypeptide sequence and the closest related sequence, biliary glycoprotein H (mouse).

30 **Figure 3:** Results from BLAST against NCBI non-redundant database using full-length INSP055 polypeptide sequence.

Figure 4: Alignment generated by BLAST between the full-length INSP055 polypeptide sequence and the closest related sequence, biliary glycoprotein H (mouse).

Figure 5: Predicted nucleotide sequence of INSP052 with translation. underlined sequence denotes predicted signal peptide. **[Boxed sequence]** denotes predicted 5 transmembrane domain.

Figure 6: INSP052 coding exon organization in genomic DNA. Bottom = INSP052.cDNA, 1251 bp. Top = chr11.genomic_DNA. Sequence encoding the putative extracellular domain is underlined. Start and Stop codons are in bold type.

Figure 7: Nucleotide sequence and translation of cloned INSP052 extracellular 10 domain

Figure 8: Map of pENTR-INSP052-EC-6HIS

Figure 9: Map of pEAK12d-INSP052-EC-6HIS

EXAMPLES

Example 1 INSP052 and INSP055

15 The polypeptide sequence derived from combining SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12 and SEQ ID NO:14 and SEQ ID NO:16 which represents the translation of consecutive exons from INSP052 is derived from human genomic sequence. The polynucleotide and polypeptide sequences SEQ ID NO 17 and SEQ ID 18 representing INSP055 are polynucleotide and polypeptide sequences of the mouse 20 orthologue of INSP052 respectively. The existence of a mouse orthologue supports the gene model for the human sequence INSP052.

INSP052 and INSP055 polypeptide sequences represented by SEQ ID NO 16 and SEQ ID NO 18, respectively, are predicted to contain signal peptide sequences and a transmembrane spanning domain.

25 The polypeptide sequence derived from combining SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12 and SEQ ID NO:14 and SEQ ID NO:16 which represents the translation of consecutive exons from INSP052, was used as a BLAST query against the NCBI non-redundant Sequence database. The top ten matches are shown in Figure 1, all of which are immunoglobulin domain containing proteins.

30 Figure 2 shows the alignment of the INSP052 query sequence to the sequence of the highest matching known protein, biliary glycoprotein H (mouse).

The polypeptide sequence INSP055, was used as a BLAST query against the NCBI non-redundant Sequence database. The top ten matches are shown in Figure 3. Figure 4 shows the alignment of the INSP055 query sequence to the sequence of the highest matching known protein, biliary glycoprotein H (mouse).

5 Expressed sequence tags (ESTs) representing the INSP052 and INSP055 transcripts in human and mouse originate from the following cDNA libraries: brain, including cerebellum, cortex, hippocampus, hypothalamus, medulla oblongata; inner ear and breast. Transcripts are also represented by ESTs from oligodendrogloma, glioblastoma and multiple sclerosis lesions. This suggests that INSP052 can be cloned from the above tissues
10 10 and may be associated with diseases of the above tissues. Accordingly, the polypeptides, antibodies and other moieties described herein may have utility in the treating a disease in one of the above tissues.

Example 2 Cloning of the INSP052 extracellular domain by exon assembly

The INSP052 full length prediction encodes a type I membrane protein of 416 amino acids, 15 related to the VEGF/PDGF receptors, belonging to the immunoglobulin superfamily. The predicted nucleotide sequence, starting from the initiating ATG codon to the poly A tail is 2025 nucleotides long (figure 5). The coding sequence (cds) spans 7 exons (figure 6). A putative signal sequence (encoding amino acids 1-33) is located in exon 1. The sequence encoding the predicted transmembrane (TM) domain (amino acids 241 to 263) is located at 20 the exon 3-4 boundary.

The extracellular (EC) domain encoding amino acids 1-240 was cloned by exon assembly from genomic DNA. An overview of the exon assembly method is summarized below:

25 – Individual exons 1, 2 and 3 were amplified from genomic DNA by PCR. The reverse primer for exon 3 also contained an 11 base overlap with the 5' sequence of exon 4.

– Gel-purified exons were mixed and a 2nd PCR reaction was performed to amplify the re-assembled DNA.

– The full length PCR product corresponding to the INSP052 EC domain was gel-purified and subcloned sequentially into pDONR 201 (Gateway entry vector) and 30 pEAK12d (expression vector) using the Invitrogen GatewayTM methodology.

–
1. PCR amplification of exons encoding the extracellular domain of INSP052 from

genomic DNA.

PCR primers were designed to amplify exons 1, 2 and 3 individually (table 1). The forward primer for exon 1 (INSP052-B1P-exon1F) also contains the partial sequence of the Gateway attB1 site (5' GCAGGGCTTC) and a Kozak sequence (5' GCCACC). The reverse 5 primer for exon 1 (INSP052 -exon1R) has an overlap of 18 bases with exon 2 at its 5' end. The forward primer for exon 2 (INSP052 -exon2F) has an 18 bp overlap with exon 1 at its 5' end. The reverse primer for exon 2 (INSP052 -exon2R) has an overlap of 18 bases with exon 3 at its 5' end. The forward primer for exon 3 (INSP052 -exon3F) contains an 17 bp overlap with exon 2 at its 5' end. The reverse primer for exon 3 (INSP052 -exon3R) 10 has an overlap of 11 bases with exon 4 at its 5' end.

For amplification of INSP052 exon 1, the PCR reaction was performed in a final volume of 50 μ l and contained 1.5 μ l of human genomic DNA (0.1 μ g/ μ l, Novagen cat. no. 69237), 2 μ l of 5 mM dNTPs (Amersham Pharmacia Biotech), 6 μ l of INSP052-B1P-exon1F (10 μ M), 6 μ l of INSP052 -exon1R , 5 μ l of 10X Pwo buffer and 0.5 μ l of Pwo polymerase (5 15 U/ μ l) (Roche, cat. no. 1 644 955). The PCR conditions were 94 0 C for 2 min; 35 cycles of 94 0 C for 30 s, 60 0 C for 30 s and 72 0 C for 1 min; an additional elongation cycle of 72 0 C for 5 min; and a holding cycle of 4 0 C. Reaction products were loaded onto a 1.5 % agarose 20 gel (1X TAE) and PCR products of the correct size (118 bp) were gel- purified using a Qiaquick Gel Extraction Kit (Qiagen cat. no. 28704) and eluted in 50 μ l of elution buffer (Qiagen).

Exon 2 was amplified using the same reaction conditions with primers INSP052 -exon2F and INSP052 -exon2R. PCR products of 378 bp were gel purified as above.

Exon 3 was amplified using the same reaction conditions with primers INSP052 -exon3F and INSP052 -exon3R. PCR products of 321 bp were gel purified as above.

25 2. Assembly of extracellular domain-encoding exons of INSP052

Exons 1, 2 and 3-4 were re-assembled in a PCR reaction containing 5 μ l of each gel purified exon, 2 μ l of 5 mM dNTPs, 6 μ l of INSP052-B1P-exon1F (10 μ M), 6 μ l of INSP052-5HIS-R (10 μ M), 5 μ l of 10X Pfu buffer, 14.5 μ l H₂O and 0.5 μ l Pfu polymerase (3 U/ μ l; Promega cat. no. M774B). The reaction conditions were: 94 0 C, 4 30 min; 10 cycles of 94 0 C for 30 s, 48 0 C for 30 s and 70 0 C for 2 min; 25 cycles of 94 0 C for 30 s, 52 0 C, for 30 s and 70 0 C for 2 min ; an additional elongation step of 70 0 C for 10 min; and a holding cycle at 4 0 C. Reaction products were analysed on a 1.5 % agarose gel (1X TAE). PCR products of the correct size (750 bp) were gel purified using a Qiaquick

Gel Extraction Kit (Qiagen cat. no. 28704) and eluted in 50 µl of elution buffer (Qiagen). The resultant product (INSP052 EC ORF) contains the ORF of the INSP052 EC domain flanked at the 5' end by an attB1 site and Kozak sequence, and at the 3' end by a 5HIS tag encoding sequence.

5 3. Subcloning of the INSP052 EC domain ORF into pDONR201

AttB1 and attB2 recombination sites were added to the 5' and 3' end of the full length INSP052 EC domain sequence in a PCR reaction containing 2 µl of gel purified INSP052 EC ORF, 2 µl of 5 mM dNTPs (Amersham Pharmacia Biotech), 6 µl of GCP-Forward (10 µM), 6 µl of GCP-Reverse (10 µM), 5 µl of 10X Vent buffer and 0.5 µl of Vent DNA polymerase (2 U/µl) (New England Biolabs, cat. no. M0254S) in a final volume of 50 µl. The PCR conditions were 94 °C for 2 min; 30 cycles of 94 °C for 30 sec; 55 °C for 30 sec and 72 °C for 1 min; an additional elongation step of 72 °C for 3 min and a holding cycle of 4 °C. Reaction products were analysed on a 1.5 % agarose gel (1X TAE) and PCR products of the correct size (808 bp) were gel purified using a Qiaquick Gel Extraction Kit (Qiagen cat. no. 28704) and eluted in 50 µl of elution buffer (Qiagen). The purified PCR product (Gateway-modified INSP052 EC domain) was then transferred to pDONR201 using BP clonase as follows: 5 µl of Gateway-modified INSP052 EC domain was incubated with 1.5 µl pDONR201 (0.1 µg/µl), 2 µl BP buffer and 1.5 µl of BP clonase enzyme mix (Invitrogen) at RT for 1 h. The reaction was stopped by addition of proteinase K (2 µg) and incubated at 37 °C for a further 10 min. An aliquot of this reaction (1 µl) was transformed into 20 µl of *E. coli* DH10B cells (diluted 1/5 in H₂O) by electroporation using a Biorad Gene Pulser. Electroporated cells were diluted by addition of 1 ml of SOC medium and incubated for 1 h at 37 °C. Transformants were plated on LB-kanamycin plates and incubated overnight at 37 °C. Plasmid mini prep DNA was isolated from 1-10 resultant colonies using a Qiaprep Turbo 9600 robotic system (Qiagen) and subjected to DNA sequencing with pENTR-F1 and pENTR-R1 sequencing primers using the BigDyeTerminator system (Applied Biosystems cat. no. 4390246) according to the manufacturer's instructions. Sequencing reactions were purified using Dye-Ex columns (Qiagen) or Montage SEQ 96 cleanup plates (Millipore cat. no. LSKS09624) then analyzed on an Applied Biosystems 3700 sequencer.

4. Subcloning of the INSP052 EC domain ORF to expression vector pEAK12d

Plasmid eluate (1.5 µl) from a pDONR201 clone containing the correct sequence of the INSP052 EC domain (plasmid ID # 13497) was then used in a recombination reaction

containing 1.5 μ l pEAK12d (0.1 μ g/ μ l), 2 μ l LR buffer and 1.5 μ l of LR clonase (Invitrogen) in a final volume of 10 μ l. The mixture was incubated at RT for 1 h, stopped by addition of proteinase K (2 μ g) and incubated at 37°C for a further 10 min. An aliquot of this reaction (1 μ l) was used to transform *E. coli* DH10B cells by electroporation as described above. Electroporated cells were diluted by addition of 1 ml of SOC medium and incubated for 1 h at 37 °C. Transformants were plated on LB-ampicillin plates and incubated overnight at 37 °C. Mini prep DNA was prepared from 4 colonies using a Qiaprep Turbo 9600 robotic system (Qiagen) and eluted in 50 μ l of elution buffer. Two μ l of each miniprep was then subjected to PCR in a total reaction volume of 50 μ l containing 2 μ l of 5mM dNTPs, 6 μ l of 10 μ M pEAK12-F, 6 μ l of 10 μ M pEAK12-R, 5 μ l of 10X AmpliTaq™ buffer and 0.5 μ l AmpliTaq™ (Applied Biosystems cat. no. N808-0155). The cycling conditions were as follows: 94 °C for 2 min; 30 cycles of 94 °C for 30 sec, 55°C for 30 sec, and 72 °C for 1 min; 1 cycle, 72 °C for 3 min. Samples were then maintained at 4 °C (holding cycle) before further analysis.

Plasmid mini prep DNA was isolated from colonies which gave the expected PCR product size (1074 bp) was then subjected to DNA sequencing with pEAK12-F and pEAK12-R sequencing primers.

CsCl gradient purified maxi-prep DNA of plasmid pEAK12d-INSP052EC-6HIS (plasmid ID # 13495) was prepared from a 500 ml culture of a sequence verified clone (Sambrook J. et al., in Molecular Cloning, a Laboratory Manual, 2nd edition, 1989, Cold Spring Harbor Laboratory Press), resuspended at a concentration of 1 μ g/ μ l in sterile water and stored at -20 C.

Table 2: Primers for INSP052 EC domain cloning and sequencing

Primer	Sequence (5'-3')
GCP Forward	G GGG ACA AGT TTG TAC AAA AAA GCA GGC TTC <u>GCC</u> <u>ACC</u>
GCP Reverse	GGG GAC CAC TTT GTA CAA GAA AGC TGG GTT TCA ATG GTG ATG GTG ATG GTG
INSP052-B1P- exon1F	GCA GGC TTC <u>GCC</u> <u>ACC</u> ATG AAG AGA GAA AGG GGA GCC CTG TC
INSP052- exon1R	TCA CCC CCT CCA GGG GGT <u>CTG</u> <u>TCT</u> GGA TCA GAA GAA

INSP052- exon2F	<u>TTC</u> <u>TTC</u> TGA TCC AGA CAG ACC CCC TGG AGG GGG TGA
INSP052- exon2R	GTG GCC TCG AAA TGG GCA <u>CAT</u> <u>CTA</u> <u>CAG</u> <u>TAA</u> GGT <u>TGA</u>
INSP052- exon3F	<u>CAA</u> <u>CCT</u> TAC TGT AGA <u>TGT</u> GCC CAT TTC GAG GCC ACA
INSP052- exon3R	<u>GGA</u> <u>GCT</u> TCT TCT GTA TAC GGT GAT CTT GAC AG
INSP052-5HIS-R	<i>GTG ATG GTG ATG GTG</i> GGA GCT TCT TCT GTA TAC GG
pEAK12-F	GCC AGC TTG GCA CTT GAT GT
pEAK12-R	GAT GGA GGT GGA CGT GTC AG
pENTR-F1	TCG CGT TAA CGC TAG CAT GGA TCT C
pENTR-R1	GTA ACA TCA GAG ATT TTG AGA CAC

Underlined sequence = Kozak sequence

Bold = Stop codon

Italic sequence = His tag

Double underlined = overlap with adjacent exon

5

Example 3: Expression in mammalian cells of the cloned, His-tagged INSP052-6His-V1 (plasmid No. 13495)

Human Embryonic Kidney 293 cells expressing the Epstein-Barr virus Nuclear Antigen (HEK293-EBNA, Invitrogen) were maintained in suspension in Ex-cell VPRO serum-free medium (seed stock, maintenance medium, JRH). Sixteen to 20 hours prior to transfection (Day-1), cells were seeded in 2x T225 flasks (50 ml per flask in DMEM / F12 (1:1) containing 2% FBS seeding medium (JRH) at a density of 2×10^5 cells/ ml). The next day (transfection day 0) the transfection took place by using the JetPEI™ reagent (2 μ l/ μ g of plasmid DNA, PolyPlus-transfection). For each flask, 113 μ g of cDNA (plasmid No. 13495) was co-transfected with 2.3 μ g of GFP (fluorescent reporter gene). The transfection mix was then added to the 2xT225 flasks and incubated at 37°C (5%CO₂) for 6 days. In order to increase our chances to get more material, we repeated this procedure into two extra flasks such as to generate 200 ml total. Confirmation of positive transfection was done by qualitative fluorescence examination at day 1 and day 6 (Axiovert 10 Zeiss).

15 On day 6 (harvest day), supernatants (200ml) from the four flasks were pooled and centrifuged (4°C, 400g) and placed into a pot bearing a unique identifier.

20 One aliquot (500ul) was kept for QC of the 6His-tagged protein (internal bioprocessing

One aliquot (500ul) was kept for QC of the 6His-tagged protein (internal bioprocessing

QC).

Purification process

The 200 ml culture medium sample containing the recombinant protein with a C-terminal 6His tag was diluted to a final volume of 200 ml with cold buffer A (50 mM NaH₂PO₄; 5 600 mM NaCl; 8.7 % (w/v) glycerol, pH 7.5). The sample was filtered through a 0.22 μ m sterile filter (Millipore, 500 ml filter unit) and kept at 4°C in a 250 ml sterile square media bottle (Nalgene).

The purification was performed at 4°C on the VISION workstation (Applied Biosystems) connected to an automatic sample loader (Labomatic). The purification procedure was 10 composed of two sequential steps, metal affinity chromatography on a Poros 20 MC (Applied Biosystems) column charged with Ni ions (4.6 x 50 mm, 0.83 ml), followed by gel filtration on a Sephadex G-25 medium (Amersham Pharmacia) column (1,0 x 10 cm).

For the first chromatography step the metal affinity column was regenerated with 30 column volumes of EDTA solution (100 mM EDTA; 1 M NaCl; pH 8.0), recharged with 15 Ni ions through washing with 15 column volumes of a 100 mM NiSO₄ solution, washed with 10 column volumes of buffer A, followed by 7 column volumes of buffer B (50 mM NaH₂PO₄; 600 mM NaCl; 8.7 % (w/v) glycerol, 400 mM; imidazole, pH 7.5), and finally equilibrated with 15 column volumes of buffer A containing 15 mM imidazole. The sample was transferred, by the Labomatic sample loader, into a 200 ml sample loop and 20 subsequently charged onto the Ni metal affinity column at a flow rate of 10 ml/min. The column was washed with 12 column volumes of buffer A, followed by 28 column volumes of buffer A containing 20 mM imidazole. During the 20 mM imidazole wash loosely attached contaminating proteins were elution of the column. The recombinant His-tagged protein was finally eluted with 10 column volumes of buffer B at a flow rate of 2 ml/min, 25 and the eluted protein was collected in a 1.6 ml fraction.

For the second chromatography step, the Sephadex G-25 gel-filtration column was regenerated with 2 ml of buffer D (1.137 M NaCl; 2.7 mM KCl; 1.5 mM KH₂PO₄; 8 mM Na₂HPO₄; pH 7.2), and subsequently equilibrated with 4 column volumes of buffer C (137 mM NaCl; 2.7 mM KCl; 1.5 mM KH₂PO₄; 8 mM Na₂HPO₄; 20 % (w/v) glycerol; pH 7.4). 30 The peak fraction eluted from the Ni-column was automatically through the integrated sample loader on the VISION loaded onto the Sephadex G-25 column and the protein was eluted with buffer C at a flow rate of 2 ml/min. The desalting sample was recovered in a 2.2

ml fraction. The fraction was filtered through a 0.22 um sterile centrifugation filter (Millipore), frozen and stored at -80C. An aliquot of the sample was analyzed on SDS-PAGE (4-12 % NuPAGE gel; Novex) by coomassie staining and Western blot with anti-His antibodies.

5 Coomassie staining. The NuPAGE gel was stained in a 0.1 % coomassie blue R250 staining solution (30 % methanol, 10 % acetic acid) at room temperature for 1 h and subsequently destained in 20 % methanol, 7.5 % acetic acid until the background was clear and the protein bands clearly visible.

Western blot. Following the electrophoresis the proteins were electrotransferred from the 10 gel to a nitrocellulose membrane at 290 mA for 1 hour at 4°C. The membrane was blocked with 5 % milk powder in buffer E (137 mM NaCl; 2.7 mM KCl; 1.5 mM KH₂PO₄; 8 mM Na₂HPO₄; 0.1 % Tween 20, pH 7.4) for 1 h at room temperature, and subsequently incubated with a mixture of 2 rabbit polyclonal anti-His antibodies (G-18 and H-15, 0.2ug/ml each; Santa Cruz) in 2.5 % milk powder in buffer E overnight at 4°C. After 15 further 1 hour incubation at room temperature, the membrane was washed with buffer E (3 x 10 min), and then incubated with a secondary HRP-conjugated anti-rabbit antibody (DAKO, HRP 0399) diluted 1/3000 in buffer E containing 2.5 % milk powder for 2 hours at room temperature. After washing with buffer E (3 x 10 minutes), the membrane was developed with the ECL kit (Amersham Pharmacia) for 1 min. The membrane was 20 subsequently exposed to a Hyperfilm (Amersham Pharmacia), the film developed and the western blot image visually analyzed.

Protein assay. The protein concentration was determined using the BCA protein assay kit (Pierce) with bovine serum albumin as standard. 890 µg purified protein was recovered from the 200 ml culture medium.

CLAIMS

1. A polypeptide, which polypeptide comprises or consists of the amino acid sequence as recited in SEQ ID NO:16 (INSP052).
2. A polypeptide which is a functional equivalent of the polypeptide according to claim 1, which has greater than 80% sequence identity with SEQ ID NO:16 and has immunoglobulin domain-containing cell surface recognition molecule activity.
3. A polypeptide which is a fragment or a functional equivalent of a polypeptide according to claim 1, which has greater than 80% sequence identity with the amino acid sequence recited in SEQ ID NO:16.
- 10 4. A purified nucleic acid molecule which encodes a polypeptide according to any one of the preceding claims.
5. A purified nucleic acid molecule according to claim 4, which comprises the nucleic acid sequence as recited in SEQ ID NO:15 or is a redundant equivalent or fragment thereof.
- 15 6. A purified nucleic acid molecule according to claim 5 which consists of the nucleic acid sequence as recited in SEQ ID NO:15 or is a redundant equivalent or fragment thereof.
7. A purified nucleic acid molecule which hybridizes under high stringency conditions with a nucleic acid molecule according to any one of claims 4 to 6.
- 20 8. A vector comprising a nucleic acid molecule as recited in any one of claims 4 to 7.
9. A host cell transformed with a vector according to claim 8.
10. An antibody ligand which binds specifically to a polypeptide according to any one of claims 1 to 3.
11. An antibody ligand according to claim 10, which inhibits the activity of a 25 polypeptide according to any one of claims 1 to 3.
12. A method of diagnosing a disease in a patient, comprising assessing the level of expression of a natural gene encoding a polypeptide according to any one of claims 1 to 3, or assessing the activity of a polypeptide according to any one of claims 1 to 3, in tissue from said patient and comparing said level of expression or activity to a control level, 30 wherein a level that is different to said control level is indicative of disease.

13. A method according to claim 12 that is carried out *in vitro*.
14. A method according to claim 12 or claim 13, which comprises the steps of: (a) contacting an antibody ligand according to claim 10 or claim 11 with a biological sample under conditions suitable for the formation of a ligand-polypeptide complex; and (b) 5 detecting said complex.
15. A method according to claim 12 or claim 13, comprising the steps of:
 - a) contacting a sample of tissue from the patient with a nucleic acid probe under stringent conditions that allow the formation of a hybrid complex between a nucleic acid molecule according to any one of claims 4 to 7 and the probe;
 - 10 b) contacting a control sample with said probe under the same conditions used in step a); and
 - c) detecting the presence of hybrid complexes in said samples; wherein detection of levels of the hybrid complex in the patient sample that differ from levels of the hybrid complex in the control sample is indicative of disease.
- 15 16. A method according to claim 12 or claim 13, comprising:
 - a) contacting a sample of nucleic acid from tissue of the patient with a nucleic acid primer under stringent conditions that allow the formation of a hybrid complex between a nucleic acid molecule according to any one of claims 4 to 7 and the primer;
 - 20 b) contacting a control sample with said primer under the same conditions used in step a); and
 - c) amplifying the sampled nucleic acid; and
 - d) detecting the level of amplified nucleic acid from both patient and control samples; wherein detection of levels of the amplified nucleic acid in the patient 25 sample that differ significantly from levels of the amplified nucleic acid in the control sample is indicative of disease.
17. A method according to claim 12 or claim 13 comprising:
 - a) obtaining a tissue sample from a patient being tested for disease;
 - b) isolating a nucleic acid molecule according to any one of claims 4 to 7 from said

tissue sample; and

c) diagnosing the patient for disease by detecting the presence of a mutation which is associated with disease in the nucleic acid molecule as an indication of the disease.

5 18. A method according to claim 17, further comprising amplifying the nucleic acid molecule to form an amplified product and detecting the presence or absence of a mutation in the amplified product.

19. A method according to claim 17 or claim 18, wherein the presence or absence of the mutation in the patient is detected by contacting said nucleic acid molecule with a 10 nucleic acid probe that hybridises to said nucleic acid molecule under stringent conditions to form a hybrid double-stranded molecule, the hybrid double-stranded molecule having an unhybridised portion of the nucleic acid probe strand at any portion corresponding to a mutation associated with disease; and detecting the presence or absence of an unhybridised portion of the probe strand as an indication of the presence or absence of a disease- 15 associated mutation.

20. A method according to any one of claims 12 to 19, wherein said disease is a cell proliferative disorder, an autoimmune/inflammatory disorder, a cardiovascular disorder, a neurological disorder, a psychiatric disorder, a developmental disorder, a genetic disorder, a metabolic disorder, an infection, a liver disease or other pathological condition.

20 21. Use of a polypeptide according to any one of claims 1 to 3 as an immunoglobulin domain-containing cell surface recognition molecule.

22. A pharmaceutical composition comprising a polypeptide according to any one of claims 1 to 3, a nucleic acid molecule according to any one of claims 4 to 7, a vector according to claim 8, a host cell according to claim 9, or an antibody ligand according to 25 claim 10 or claim 11.

23. A vaccine composition comprising a polypeptide according to any one of claims 1 to 3 or a nucleic acid according to any one of claims 4 to 7.

24. Use of a polypeptide according to any one of claims 1 to 3, a nucleic acid molecule according to any one of claims 4 to 7, a vector according to claim 8, a host cell according 30 to claim 9, an antibody ligand according to claim 10 or claim 11, or a pharmaceutical composition according to claim 22, in the manufacture of a medicament for the treatment

of a disease selected from the group consisting of: cell proliferative disorders, autoimmune/inflammatory disorders, cardiovascular disorders, neurological and psychiatric disorders, developmental disorders, genetic disorders, metabolic disorders, infections, and other pathological conditions.

5 25. A method of treating a disease in a patient, comprising administering to the patient a polypeptide according to any one of claims 1 to 3, a nucleic acid molecule according to any one of claims 4 to 7, a vector according to claim 8, a host cell according to claim 9, an antibody ligand according to claim 10 or claim 11, or a pharmaceutical composition according to claim 22.

10 26. A method according to claim 25, wherein, for diseases in which the expression of the natural gene or the activity of the polypeptide is lower in a diseased patient when compared to the level of expression or activity in a healthy patient, the polypeptide, nucleic acid molecule, vector, antibody ligand, compound or composition administered to the patient is an agonist.

15 27. A method according to claim 25, wherein, for diseases in which the expression of the natural gene or activity of the polypeptide is higher in a diseased patient when compared to the level of expression or activity in a healthy patient, the polypeptide, nucleic acid molecule, vector, antibody ligand, compound or composition administered to the patient is an antagonist.

20 28. A method of monitoring the therapeutic treatment of disease in a patient, comprising monitoring over a period of time the level of expression or activity of a polypeptide according to any one of claims 1 to 3, or the level of expression of a nucleic acid molecule according to any one of claims 4 to 7 in tissue from said patient, wherein altering said level of expression or activity over the period of time towards a control level
25 is indicative of regression of said disease.

29. A method for the identification of a compound that is effective in the treatment and/or diagnosis of disease, comprising contacting a polypeptide according to any one of claims 1 to 3, or a nucleic acid molecule according to any one of claims 4 to 7 with one or more compounds suspected of possessing binding affinity for said polypeptide or nucleic acid molecule, and selecting a compound that binds specifically to said nucleic acid molecule or polypeptide.

30. A kit useful for diagnosing disease comprising a first container containing a nucleic acid probe that hybridises under stringent conditions with a nucleic acid molecule according to any one of claims 4 to 7; a second container containing primers useful for amplifying said nucleic acid molecule; and instructions for using the probe and primers for 5 facilitating the diagnosis of disease.

31. A kit according to claim 30, further comprising a third container holding an agent for digesting unhybridised RNA.

32. A kit comprising an array of nucleic acid molecules, at least one of which is a nucleic acid molecule according to any one of claims 4 to 7.

10 33. A kit comprising one or more antibodies that bind to a polypeptide as recited in any one of claims 1 to 3, and a reagent useful for the detection of a binding reaction between said antibody and said polypeptide.

34. A transgenic or knockout non-human animal that has been transformed to express higher, lower or absent levels of a polypeptide according to any one of claims 1 to 3.

15 35. A method for screening for a compound effective to treat disease, by contacting a non-human transgenic animal according to claim 34 with a candidate compound and determining the effect of the compound on the disease of the animal.

36. A polypeptide according to any one of claims 1 to 3, a nucleic acid molecule according to any one of claims 4 to 7, a vector according to claim 8, a host cell according 20 to claim 9, an antibody ligand according to claim 10 or 11, a method according to claim 12, 25, 28, 29 or 35, a pharmaceutical composition according to claim 22, a vaccine according to claim 23, a use according to claim 24, a kit according to any one of claims 30 to 33, or a transgenic knockout non-human animal according to claim 34, substantially as hereinbefore described with reference to the accompanying examples and/or figures.

1/15

FIG. 1

BLASTP 2.2.1 [Jul-12-2001]

Reference: Altschul, Stephen F., Thomas L. Madden, Alejandro A. Schaffer, Jinghui Zhang, Zheng Zhang, Webb Miller, and David J. Lipman (1997), "Gapped BLAST and PSI-BLAST: a new generation of protein database search programs", Nucleic Acids Res. 25:3389-3402.

Query= genscan2

(416 letters)

Database: ncbi-nr

897,014 sequences; 280,886,335 total letters

Searching.....done

	Score	E
Sequences producing significant alignments:	(bits)	Value
pir JC1512 biliary glycoprotein H - mouse	81	2e-14
pir JC1506 biliary glycoprotein B - mouse	81	2e-14
pir A39037 carcinoembryonic antigen mmCGM2 precursor - mouse >g...	79	9e-14
ref NP_036056.1 (NM_011926) CEA-related cell adhesion molecule ...	79	9e-14
pir JC1509 biliary glycoprotein E - mouse	73	5e-12
ref NP_001758.1 (NM_001767) CD2 antigen (p50), sheep red blood ...	73	6e-12
ref NP_113943.1 (NM_031755) carcinoembryonic antigen-related ce...	72	8e-12
pir RWHUC2 T-cell surface glycoprotein CD2 precursor - human >g...	72	8e-12
gb AAA51946.1 (MI6336) CD2 surface antigen [Homo sapiens]	72	8e-12
ref NP_291021.1 (NM_033543) hypothetical protein R29124_1 [Homo...	72	1e-11
pir JC1507 biliary glycoprotein C - mouse	71	2e-11
emb CAA47697.1 (X67280) biliary glycoprotein [Mus musculus]	71	2e-11
pir S34338 biliary glycoprotein F - mouse >gi 312586 emb CAA476...	71	2e-11
pir JC1511 biliary glycoprotein G - mouse	71	2e-11

FIG. 1 (contd.)

3/15

FIG. 2

>pir||JC1512 biliary glycoprotein H - mouse

Length = 341

Score = 80.9 bits (198), Expect = 2e-14

Identities = 54/168 (32%), Positives = 86/168 (51%), Gaps = 9/168 (5%)

Query: 73 RDKPVTVVQSIGTEVIGTLR---PDYRDRIRLFENGSLLLSDLQLADEGTYEVEISITD 128

+ PV+ I +V GT + P + R ++ NGSLL+ + + D G Y +E+ TD

Sbjct: 69 KGNPVSTNAEIVHQVTGTNKTTPAHSGRETVYSNGSLLIQRVTVKDTGVYTIEM--TD 126

Query: 129 DTFTG-EKTINLTVDVPISRPQVLVASTTVLELSEAFTLNCSHENGTKPSYTWLKDGPL 187

+ F E T+ V P+++P + V +TTV EL ++ TL C N + WL ++ L

Sbjct: 127 ENFRRTEATVQFHVHQPVTPQPSLQVTNTTVKEL-DSVLTCL-SNDIGANIQWLFNSQL 184

Query: 188 LNDSRMLLSPDQKVLTITRVLMEDDDLYSCMVNPISQGRSLPVKITV 235

RM LS + +L I + ED Y C + NP+S RS +K+ +

Sbjct: 185 QLTERMTLSQNNISILRIDPIKREDAGEYQCEISNPVSVKRSNSIKLDI 232

4/15

FIG. 3

Sequences producing significant alignments:	Score	E
	(bits)	Value
gi 483306 pir JC1506 biliary glycoprotein B - mouse	79	1e-13
gi 111207 pir A39037 carcinoembryonic antigen mmCGM2 precu...	77	3e-13
gi 483312 pir JC1512 biliary glycoprotein H - mouse	77	4e-13
gi 13937381 ref NP_036056.1 (NM_011926) CEA-related cell a...	75	1e-12
gi 228710 prf 1809184A pregnancy-specific glycoprotein (Ra...	70	5e-11
gi 483307 pir JC1507 biliary glycoprotein C - mouse	70	6e-11
gi 16117775 ref NP_291021.1 (NM_033543) hypothetical prote...	69	8e-11
gi 483309 pir JC1509 biliary glycoprotein E - mouse	69	9e-11
gi 312582 emb CAA47695.1 (X67278) biliary glycoprotein (Mu...	69	1e-10
gi 483311 pir JC1511 biliary glycoprotein G - mouse	68	2e-10

5/15

FIG. 4

>gi|483306|pir||JC1506 biliary glycoprotein B - mouse

Length = 278

Score = 78.6 bits (192), Expect = 1e-13

Identities = 54/168 (32%), Positives = 86/168 (51%), Gaps = 9/168 (5%)

Query: 73 RDKPVTVVQSIGTEVIGTLR----PDYRDRIRLFENGSLLSDLQLADEGTYEVEISITD 128

+ PV+ I +V GT + P + R ++ NGSL+ + + D G Y +E +TD

Sbjct: 69 KGNPVSTNAEIVHQVIGTNKTTGPAHSGRETVYSNGSLLIQRVTVKDTGVYTIE--MTD 126

Query: 129 DTF-TGEKTINLTVDVPISRPQVLVASTTVLELSEAFTLNCSHENGTKPSYTWLKDGPL 187

+ F E T+ V P+++P + V +TTV EL ++ TL C N + WL + + L

Sbjct: 127 ENFRRTEATVQFHVHQPVTLQPSLQVTNTTVKEL-DSVTLTCL-SNDIGANIQWLSQL 184

Query: 188 LNDSRMLLSPDQKVLTITRVLMEDODLYSCVVENPISQVRSLPVKITV 235

RM LS + +L I + ED Y C + NP+S RS +K+ +

Sbjct: 185 QLTERMTLSQNNISILRIDPIKREDAGEYQCEISNPVSVKRSNSIKLDI 232

6/15

FIG. 5

1 ATGAAGAGAG AAAGGGGAGC CCTGTCCAGA GCCTCCAGGG CCCTGCGCCT TGCTCCTTT
m k r e r g a l s r a s r a l r l a p f

61 GTCTACCTTC TTCTGATCCA GACAGACCCC CTGGAGGGGG TGAACATCAC CAGCCCCGTG
v y l l l i q t d p l e g v n i t s p v

121 CGCCTGATCC ATGGCACCGT GGGGAAGTCG GCTCTGCTTT CTGTGCAGTA CAGCAGTACC
r l i h g t v g k s a l l s v q y s s t

181 AGCAGCGACA GGCTGTAGT GAAGTGGCAG CTGAAGCGGG ACAAGCCAGT GACCGTGGTG
s s d r p v v k w q l k r d k p v t v v

241 CAGTCCATTG GCACAGAGGT CATCGGCACC CTGCGGCCTG ACTATCGAGA CCGTATCCGA
q s i g t e v i g t l r p d y r d r i r

301 CTCTTGAAA ATGGCTCCCT GCTTCTCAGC GACCTGCAGC TGGCCGATGA GGGCACCTAT
l f e n g s l l l s d l q l a d e g t y

361 GAGGTCGAGA TCTCCATCAC CGACGACACC TTCACTGGGG AGAAGACCAT CAACCTTACT
e v e i s i t d d t f t g e k t i n l t

421 GTAGATGTGC CCATTCGAG GCCACAGGTG TTGGTGGCTT CAACCACTGT GCTGGAGCTC
v d v p i s r p q v l v a s t t v l e l

481 AGCGAGGCCT TCACCTTGAA CTGCTCACAT GAGAATGGCA CCAAGCCCAG CTACACCTGG
s e a f t l n c s h e n g t k p s y t w

541 CTGAAGGATG GCAAGCCCCT CCTCAATGAC TCGAGAATGC TCCGTCCCC CGACCAAAAG
l k d g k p l l n d s r m l l s p d q k

601 GTGCTCACCA TCACCCGCGT GCTCATGGAG GATGACGACC TGTACAGCTG CATGGTGGAG
v l t i t r v l m e d d d l y s c m v e

661 AACCCCATCA GCCAGGGCCG CAGCCTGCCCT GTCAAGATCA CCGTATACAG AAGAAGCTCC
n p i s q g r s l p v k i t v y r r s s

721 CTTTACATCA TCTTGTCTAC AGGAGGCATC TTCCCTCCTTG TGACCTTGGT GACAGTCTGT
[l y i i l s t g g i f l l v t l v t v c]

781 GCCTGCTGGA AACCCCTCAA AAGGAAACAG AAGAAGCTAG AAAAGCAAAA CTCCCTGGAA
[a c w k p s k r k q k k l e k q n s l e]

7/15

841 TACATGGATC AGAATGATGA CCGCCTGAAA CCAGAAGCAG ACACCCCTCCC TCGAAGTGGT
y m d q n d d r l k p e a d t l p r s g

901 GAGCAGGAAC GGAAGAACCC CATGGCACTC TATATCCTGA AGGACAAGGA CTCCCCGGAG
e q e r k n p m a l y i l k d k d s p e

961 ACCGAGGAGA ACCCGGCCCG GGAGCCTCGA AGCGCGACGG AGCCCGGCCGCC GCCCGGCTAC
t e e n p a p e p r s a t e p g p p g y

1021 TCCGTGTCTC CCGCCGTGCC CGGCCGCTCG CCAGGGCTGC CCATCCGCTC TGCCCGCCGC
s v s p a v p g r s p g l p i r s a r r

1081 TACCCGCGCT CCCCAGCGCG CTCCCCAGCC ACCGGCCGGA CACACTCGTC GCCGGCCAGG
y p r s p a r s p a t g r t h s s p p r

1141 GCCCCGAGCT CGCCCGGCCCG CTCGGCGAGC GCCTCGCGCA CACTGGGAC TGCGGGCGTG
a p s s p g r s r s a s r t l r t a g v

1201 CACATAATCC GCGAGCAAGA CGAGGCCGGC CCGGTGGAGA TCAGGCCCTG AGCCGCCTCG
h i i r e q d e a g p v e i s a

1261 GGATCCCCCTG AGAGGCGCCC GCGGTCTGCG GCCAGTGGCC CGGGGGAAAG CTGGGGCTGG
1321 GAAGCCCGGG CGCGCGCGC TGGGGACGAG GGGAGGTCCC GGGGGGGCGC TGGTGTCTCG
1381 GGTGTGAACG TGTATGAGCA TGCGCAGACG GAGGCAGGTG CGCGGAGGCG GCAGTGTGTA
1441 TATGGTGAAA CCGGGTCGCA TTTGCTTCCG GTTTACTGGC TGTGTCTCA CTTGGTATAG
1501 GTTGTGCCCT CTTAGGACCA CATAGATTAT TACATTTCTG GCCCAATACC CAAAAGGGTT
1561 TTATGGAAAC TAACATCAGT AACCTAACCC CCGTGACTAT CCTGTGCTCT TCCTAGGGAG
1621 CTGTGTTGTT TCCCACCCAC CACCCCTCCC TCTGAACAAA TGCCTGAGTG CTGGGGCACT
1681 TTTTTTTTTT TTTTTTTTTT GCAAGTTCA ATTAGAGAGG CCACTTTCCC
1741 AGAATCCACA GCTGCACTAA GCTAAGGAGA AGCCAGATGC CGGTTACTGG GTGTGCAGGG
1801 GCTGTTCTGA GCTGGGGGGA TCATTGTGAA GGCCTTCTTC CCTGGGCACC TGGTACCTGG
1861 GGACCTACAA GGTGGTGAGG GAAGGGTACG AGTACATTCC TTTTCCCTCT GACCTGGGCG
1921 CTAGCAAGGG CAAAGAACCC GAGCCTGCCA GCTTGGCCTC CTCCACAGC CTCCCTCGGA
1981 GGCATGCCAT GCCAAGCACT CTTCTGTCT CTGTTCATGA ATAAA

FIG. 5(contd.)

FIG. 6

gi 115886 **ATGAAGAGAGAAAGGGGAGCCCTGTCCAGAGCCTCCAGGGCCCTGCGCCT** 115935

1. ATGAAGAGAGAAAGGGGAGCCCTGTCCAGAGCCTCCAGGGCCCTGCGCCT 50
 INSP052-BIP-exon2F →
 INSP052-exon2F →

gi 115936 **TGCTCCTTTGTCTACCTCTTCTGATCCAGACAGgtagg.....cac** 115970

51 TGCTCCTTTGTCTACCTCTTCTGATCCAGACAG..... 85
 ← INSP052-exon1R cont
 INSP052-exon2F cont →

gi 115970 **agACCCCTGGAGGGGGTGAACATCACCAAGCCCCGTGCGCCTGATCCATG** 126870

>>|||||.....|||||.....|||||.....|||||.....|||||.....|||||.....
 85 . ACCCCTGGAGGGGGTGAACATCACCAAGCCCCGTGCGCCTGATCCATG 133
 ← INSP052-exon1R

gi 126871 **GCACCGTGGGAAGTCGGCTCTGCTTCTGTGCAGTACAGCAGTACCGAC** 126920

|||||.....|||||.....|||||.....|||||.....|||||.....|||||.....
 134 GCACCGTGGGAAGTCGGCTCTGCTTCTGTGCAGTACAGCAGTACCGAC 183

gi 126921 **AGCGACAGGCCTGTAGTGAAGTGGCAGCTGAAGCGGGACAAGCCAGTGAC** 126970

|||||.....|||||.....|||||.....|||||.....|||||.....|||||.....
 184 AGCGACAGGCCTGTAGTGAAGTGGCAGCTGAAGCGGGACAAGCCAGTGAC 233

gi 126971 **CGTGGTGCAGTCCATTGGCACAGAGGTATCGCACCCCTGCGGCCTGACT** 127020

|||||.....|||||.....|||||.....|||||.....|||||.....|||||.....
 234 CGTGGTGCAGTCCATTGGCACAGAGGTATCGCACCCCTGCGGCCTGACT 283

gi 127021 **ATCGAGACCGTATCCGACTCTTGAAAATGGCTCCCTGCTTCTCAGCGAC** 127070

|||||.....|||||.....|||||.....|||||.....|||||.....|||||.....
 284 ATCGAGACCGTATCCGACTCTTGAAAATGGCTCCCTGCTTCTCAGCGAC 333

gi 127071 **CTGCAGCTGGCCGATGAGGGCACCTATGAGGTGAGATCTCCATCACCGA** 127120

|||||.....|||||.....|||||.....|||||.....|||||.....|||||.....
 334 CTGCAGCTGGCCGATGAGGGCACCTATGAGGTGAGATCTCCATCACCGA 383

9/15

FIG. 6(contd.)

INSP052-exon3F

gi 127121 CGACACCTTCACTGGGGAGAAGACCATCAACCTTACTGTAGATGgtaaa. 127164
 384 CGACACCTTCACTGGGGAGAAGACCATCAACCTTACTGTAGATG..... 427

INSP052-exon3F cont INSP052-exon2R cont

gi 127164ctcagTGCCCATTCGAGGCCACAGGTGTTGGCTTCAACCACT 127922
 717 >>>|||||.....|||||.....|||||.....|||||.....|||||.....

427TGCCCATTCGAGGCCACAGGTGTTGGCTTCAACCACT 468
 INSP052-exon2R

gi 127923 GTGCTGGAGCTCAGCGAGGCCTCACCTGAACTGCTCACATGAGAATGG 127972
 469 GTGCTGGAGCTCAGCGAGGCCTCACCTGAACTGCTCACATGAGAATGG 518

gi 127973 CACCAAGCCCAGCTACACCTGGCTGAAGGATGGCAAGCCCCCTCCTCAATG 128022
 519 CACCAAGCCCAGCTACACCTGGCTGAAGGATGGCAAGCCCCCTCCTCAATG 568

gi 128023 ACTCGAGAATGCTCCTGTCCCCGACCAAAAGGTGCTCACCATCACCGC 128072
 569 ACTCGAGAATGCTCCTGTCCCCGACCAAAAGGTGCTCACCATCACCGC 618

gi 128073 GTGCTCATGGAGGATGACGACCTGTACAGCTGCATGGTGGAGAACCCAT 128122
 619 GTGCTCATGGAGGATGACGACCTGTACAGCTGCATGGTGGAGAACCCAT 668

gi 128123 CAGCCAGGGCCGCAGCCTGCCTGTCAAGATCACCGTATACAGttag.... 128163
 669 CAGCCAGGGCCGCAGCCTGCCTGTCAAGATCACCGTATACA..... 709
 INSP052-exon3R cont

gi 128163 .cctagGAAGAAGCTCCCTTACATCATCTTGTCTACAGGAGGCATCTC 128502
 >>>|||||.....|||||.....|||||.....|||||.....|||||.....>>> 295
 709GAAGAAGCTCCCTTACATCATCTTGTCTACAGGAGGCATCTC 753
 INSP052-exon3R

gi 128503 CTCCTTGTGACCTTGGTGACAGTCTGTGCCTGCTGGAAACCTCCAAAAG 128552
 754 CTCCTTGTGACCTTGGTGACAGTCTGTGCCTGCTGGAAACCTCCAAAAG 803

10/15

FIG. 6(contd.)

```

gi 128552 gtctg.....cacagGAAACAGAAGAAGCTAGAAAAGCAAAACTCCCTGG 129108
      >>>> 521 >>>>|||||||||||||||||||||||||||||||||||||||||||
803 .....GAAACAGAAGAAGCTAGAAAAGCAAAACTCCCTGG 838

gi 129109 AATACATGGATCAGAATGATGACCGCCTGAAACCAGAAGGttag.....t 129147
      |||||||||||||||||||||||||||||||||||||||||||||>>>> 286 >
839 AATACATGGATCAGAATGATGACCGCCTGAAACCAGAAG..... 877

gi 129147 gcagCAGACACCCCTCCCTCGAAGTGGTGAGCAGGAACGGAAGAACCCAT 129479
      >>>>|||||||||||||||||||||||||||||||||||||||||||||||||||
877 ....CAGACACCCCTCCCTCGAAGTGGTGAGCAGGAACGGAAGAACCCAT 923

gi 129480 GGCACTCTATATCCTGAAGGACAAGgttag.....tgtagGACTCCCCGG 130461
      |||||||||||||||||||||>>>> 947 >>>>|||||||||||
924 GGCACTCTATATCCTGAAGGACAAG.....GACTCCCCGG 958

gi 130462 AGACCGAGGAGAACCCGGCCCCGGAGCCTCGAAGCGCGACGGAGCCGGC 130511
      |||||||||||||||||||||||||||||||||||||||||||||||||||||
959 AGACCGAGGAGAACCCGGCCCCGGAGCCTCGAAGCGCGACGGAGCCGGC 1008

gi 130512 CCGCCCGGCTACTCCGTGTCTCCGCCGTGCCCGGCCGCTCGCCGGGGCT 130561
      |||||||||||||||||||||||||||||||||||||||||||||||||
1009 CCGCCCGGCTACTCCGTGTCTCCGCCGTGCCCGGCCGCTCGCCGGGGCT 1058

gi 130562 GCCCATCCGCTCTGCCGCCGCTACCGCGCTCCCCAGCGCGCTCCCCAG 130611
      |||||||||||||||||||||||||||||||||||||||||||||||||
1059 GCCCATCCGCTCTGCCGCCGCTACCGCGCTCCCCAGCGCGCTCCCCAG 1108

gi 130612 CCACCGGCCGGACACACTCGTCGCCGCCAGGGCCCCGAGCTCGCCCGGC 130661
      |||||||||||||||||||||||||||||||||||||||||||||
1109 CCACCGGCCGGACACACTCGTCGCCGCCAGGGCCCCGAGCTCGCCCGGC 1158

gi 130662 CGCTCGCGCAGCGCCTCGCGCACACTGCCGACTGCCGGCTGCACATAAT 130711
      |||||||||||||||||||||||||||||||||||||||||||||
1159 CGCTCGCGCAGCGCCTCGCGCACACTGCCGACTGCCGGCTGCACATAAT 1208

gi 130712 CCGCGAGCAAGACGAGGCCGGCCGGTGGAGATCAGCGCCTGA 130754
      |||||||||||||||||||||||||||||||||||||||||
1209 CCGCGAGCAAGACGAGGCCGGCCGGTGGAGATCAGCGCCTGA 1251

```

11/15

FIG. 7

1 ACAAGTTGT ACAAAAAAGC AGGCTTCGCC ACCATGAAGA GAGAAAGGGG AGCCCTGTCC
m k r e r g a l s

61 AGAGCCTCCA GGGCCCTGCG CCTTGCTCCT TTTGTCTACC TTCTTCTGAT CCAGACAGAC
r a s r a l r l a p f v y l l l i q t d

121 CCCCTGGAGG GGGTGAACAT CACCAGCCCC GTGCGCCTGA TCCATGGCAC CGTGGGAAAG
p l e g v n i t s p v r l i h g t v g k

181 TCGGCTCTGC TTTCTGTGCA GTACAGCAGT ACCAGCAGCG ACAGGCCTGT AGTGAAGTGG
s a l l s v q y s s t s s d r p v v k w

241 CAGCTGAAGC GGGACAAGCC AGTGACCGTG GTGCAGTCCA TTGGCACAGA GGTACATCGGC
q l k r d k p v t v v q s i g t e v i g

301 ACCCTGCGGC CTGACTATCG AGACCGTATC CGACTCTTG AAAATGGCTC CCTGCTTCTC
t l r p d y r d r i r l f e n g s l l l

361 AGCGACCTGC AGCTGGCCGA TGAGGGCACC TATGAGGTAG AGATCTCCAT CACCGACGAC
s d l q l a d e g t y e v e i s i t d d

421 ACCTTCACTG GGGAGAAGAC CATCACCTT ACTGTAGATG TGCCCATTTC GAGGCCACAG
t f t g e k t i n l t v d v p i s r p q

481 GTGTTGGTGG CTTCAACCAC TGTGCTGGAG CTCAGCGAGG CCTTCACCTT GAACTGCTCA
v l v a s t t v l e l s e a f t l n c s

541 CATGAGAATG GCACCAAGCC CAGCTACACC TGGCTGAAGG ATGGCAAGCC CCTCCTCAAT
h e n g t k p s y t w l k d g k p l l n

601 GACTCGAGAA TGCTCCTGTC CCCCCGACCAA AAGGTGCTCA CCATCACCCG CGTGCTCATG
d s r m l l s p d q k v l t i t r v l m

661 GAGGATGACG ACCTGTACAG CTGCATGGTG GAGAACCCCA TCAGCCAGGG CCGCAGCCTG
e d d d l y s c m v e n p i s q g r s l

721 CCTGTCAAGA TCACCGTATA CAGAAGAAGC TCCCCACCAC ACCATCACCA TTGAAACCCA
p v k i t v y r r s s h h h h h h h

781 GCTTTCTTGT ACAAAAGTGGT

12/15

FIG. 8

Molecule: pENTR-INSPO52- EC-6HIS, 3005 bps DNA Circular
File Name: pENTR-INSPO52-6HIS.cm5, dated 21 Feb 2003

Description: Ligation of Cons-6His.SEQ into pENTR-attL1-attL2


Molecule Features:

Type	Start	End	Name	Description
MARKER	21			pENTR-F1 primer
MARKER	110		C attL1	
GENE	136	873	INSPO52-EC-6HIS	
MARKER	888		attL2	
MARKER	1001		C	pENTR-R1 primer
GENE	1100	1909	KanR	
REGION	2030	2669	ori	

13/15

FIG. 8(contd.)

SUBSTITUTE SHEET (RULE 26)



14/15

FIG. 9

Molecule: pEAK12d-INSP052-EC-6HIS, 7687 bps DNA Circular
File Name: pEAK12d-INSP052-6HIS.cm5, dated 21 Feb 2003

Description: Ligation of Cons-6His.SEQ into pEAK12d-attB1-attB2


Molecule Features:

Type	Start	End	Name	Description
REGION	2	595		pmb-ori
GENE	596	1519	AmpR	
REGION	1690	2795	EF-1a	
MARKER	2703			pEAK12F primer
REGION	2855	2887	attB1	
GENE	2888	3625	INSP052-EC-6HIS (aal-240)	
REGION	3629	3654	attB2	
MARKER	3656	C		pEAK12R primer
REGION	3661	4089		poly A/splice
GENE	4708	4090	C	PUROMYCIN resistance
REGION	4932	4709	C tK	tK promoter
REGION	5427	4933	C Ori P	
GENE	7479	5427	C EBNA-1	
REGION	7480	7679	sv40	

15/15

SUBSTITUTE SHEET (RULE 26)

FIG. 9 (contd.)



Sequence Listing

Note: for amino acids encoded by exon-exon junctions, the amino acid will be assigned to the more 5' exon.

SEQ ID NO 1: (INSP052 Nucleotide sequence exon1)

5 1 ATGAAGAGAG AAAGGGGAGC CCTGTCCAGA GCCTCCAGGG CCCTGCGCCT TGCTCCTTT
61 GTCTACCTTC TTCTGATCCA GACAG

SEQ ID NO 2: (INSP052 polypeptide sequence of Exon 1)

10 1 MKRERGALSR ASRALRLAPF VYLLLIQTD
SEQ ID NO 3: (INSP052 Nucleotide sequence exon2)

15 1 ACCCCCTGGA GGGGGTGAAC ATCACCAAGCC CCGTGCCT GATCCATGGC ACCGTGGGGA
61 AGTCGGCTCT GCTTCTGTG CAGTACAGCA GTACCAGCAG CGACAGGCCT GTAGTGAAGT
121 GGCAGCTGAA GCGGGACAAG CCAGTGACCG TGTTGCAGTC CATTGGCACA GAGGTCATCG
181 GCACCCCTGCG GCCTGACTAT CGAGACCGTA TCCGACTCTT TGAAAATGGC TCCCTGCTTC
241 TCAGCGACCT GCAGCTGGCC GATGAGGGCA CCTATGAGGT CGAGATCTCC ATCACCGACG
301 ACACCTTCAC TGGGGAGAAG ACCATCAACC TTACTGTAGA TG

SEQ ID NO 4: (INSP052 Protein Sequence of Exon 2)

20 1 PLEGVNITSP VRLIHGTVGK SALLSVQYSS TSSDRPVVKW QLKRDKPVTV VQSIGTEVIG
61 TLRPDYRDRI RLFENGSLLL SDLQLADEGT YEVEISITDD TFTGEKTINL TVDV

SEQ ID NO 5: (INSP052 Nucleotide sequence Exon3)

25 1 TGCCCATTTC GAGGCCACAG GTGTTGGTGG CTTCAACCAC TGTGCTGGAG CTCAGCGAGG
61 CCTTCACCTT GAACTGCTCA CATGAGAATG GCACCAAGCC CAGCTACACC TGGCTGAAGG
121 ATGGCAAGCC CCTCCTCAAT GACTCGAGAA TGCTCCTGTC CCCCCACCA AAGGTGCTCA
181 CCATCACCCG CGTGCTCATG GAGGATGACG ACCTGTACAG CTGCATGGTG GAGAACCCCA
241 TCAGCCAGGG CCGCAGCCTG CCTGTCAAGA TCACCGTATA CA

30 SEQ ID NO 7: (INSP052 Polypeptide sequence of Exon 3)

1 PISRPQVLVA STTVLELSEA FTLNCSHENG TKPSYTWLKD GKPLLNDSRM LLSPDQKVLT
61 ITRVLMEDDD LYSCMVENPI SQGRSLPVKI TVYR

SEQ ID NO 7: (INSP052 Nucleotide Sequence Exon 4)

1 GAAGAAGCTC CCTTTACATC ATCTTGTCTA CAGGAGGCAT CTTCCCTCCTT GTGACCTTGG
61 TGACAGTCTG TGCCTGCTGG AAACCCCTCCA AAAG

5

SEQ ID NO 8: (INSP052 Polypeptide sequence of Exon 4)

1 RSSLYIILST GGIFLLVTLV TVCACWKPSK R

SEQ ID NO 9: (INSP052 Nucleotide Sequence Exon 5)

10 1 GAAACAGAAG AAGCTAGAAA AGCAAAACTC CCTGGAATAC ATGGATCAGA ATGATGACCG
61 CCTGAAACCA GAAG

SEQ ID NO 10: (INSP052 Polypeptide Sequence Exon 5)

1 KQKKLEKQNS LEYMDQNDDR LKPEA

15

SEQ ID NO 11: (INSP052 Nucleotide Sequence Exon 6)

1 CAGACACCCT CCCTCGAAGT GGTGAGCAGG AACGGAAGAA CCCCATGGCA CTCTATATCC
61 TGAAGGACAA G

20 SEQ ID NO 12: (INSP052 Polypeptide Sequence Exon 6)

1 DTLPRSGEQE RKNPMALYIL KDK

SEQ ID NO 13: (INSP052 Nucleotide Sequence Exon 7)

1 GACTCCCCGG AGACCGAGGA GAACCCGGCC CCGGAGCCTC GAAGCGCGAC GGAGCCCGGC
25 61 CCGCCCGGCT ACTCCGTGTC TCCCGCCGTG CCCGGCCGCT CGCCGGGGCT GCCCATCCGC
121 TCTGCCCGCC GCTACCCGCG CTCCCCAGCG CGCTCCCCAG CCACCGGCCG GACACACTCG
181 TCGCCGCCCA GGGCCCCGAG CTGCCCCGGC CGCTCGCGCA GCGCCTCGCC CACACTGCCG
241 ACTGCGGGCG TGCACATAAT CCGCGAGCAA GACGAGGCCG GCCCGGTGGA GATCAGCGCC
301 TGA

30

SEQ ID NO 14: (INSP052 Polypeptide sequence for exon 7)

1 DSPEEENPA PEPRSATEPG PPGYSVSPA V PGRSPGLPIR SARRYPRSPA RSPATGRTHS
61 SPPRAPSSPG RSRSASRTL R TAGVHITREQ DEAGPVEISA

SEQ ID NO :15 (INSP052 Combined Nucleotide sequence exons 1,2,3,4,5,6 and 7)

1 ATGAAGAGAG AAAGGGGAGC CCTGTCCAGA GCCTCCAGGG CCCTGCGCCT TGCTCCTTT
61 GTCTACCTTC TTCTGATCCA GACAGACCCC CTGGAGGGGG TGAACATCAC CAGCCCCGTG
5 121 CGCCTGATCC ATGGCACCGT GGGGAAGTCG GCTCTGCTTT CTGTGCAGTA CAGCAGTACC
181 AGCAGCGACA GGCCTGTAGT GAAGTGGCAG CTGAAGCGGG ACAAGCCAGT GACCGTGGTG
241 CAGTCCATTG GCACAGAGGT CATCGGCACC CTGCGGCCTG ACTATCGAGA CCGTATCCGA
301 CTCTTGAAA ATGGCTCCCT GCTTCTCAGC GACCTGCAGC TGGCCGATGA GGGCACCTAT
361 GAGGTCGAGA TCTCCATCAC CGACGACACC TTCACTGGGG AGAAGACCAT CAACCTTACT
10 421 GTAGATGTGC CCATTCGAG GCCACAGGTG TTGGTGGCTT CAACCACTGT GCTGGAGCTC
481 AGCGAGGCCT TCACCTTGAA CTGCTCACAT GAGAATGGCA CCAAGCCAG CTACACCTGG
541 CTGAAGGATG GCAAGCCCT CCTCAATGAC TCGAGAATGC TCCTGTCCCC CGACCAAAAG
601 GTGCTCACCA TCACCCCGT GCTCATGGAG GATGACGACC TGTACAGCTG CATGGTGGAG
661 AACCCCATCA GCCAGGGCCG CAGCCTGCCT GTCAAGATCA CCGTATACAG AAGAAGCTCC
15 721 CTTTACATCA TCTTGTCTAC AGGAGGCATC TTCTCCTTG TGACCTTGGT GACAGTCTGT
781 GCCTGCTGGA AACCCCTCAA AAGGAAACAG AAGAAGCTAG AAAAGCAAA CTCCCTGGAA
841 TACATGGATC AGAATGATGA CCGCCTGAAA CCAGAAGCAG ACACCCTCCC TCGAAGTGGT
901 GAGCAGGAAC GGAAGAACCC CATGGCACTC TATATCCTGA AGGACAAGGA CTCCCCGGAG
961 ACCGAGGAGA ACCCGGCCCG GGAGCCTCGA AGCGCGACGG AGCCCGGCCG GCCCGGCTAC
20 1021 TCCGTGTCTC CCGCCGTGCC CGGCCGCTCG CGGGGCTGC CCATCCGCTC TGCCCCGCC
1081 TACCCGCGCT CCCCAGCGC CTCCCCAGCC ACCGGCCGGA CACACTCGTC GCCGCCAGG
1141 GCCCCGAGCT CGCCCGGCCG CTCGCGCAGC GCCTCGCGCA CACTGCGGAC TGCGGCGTG
1201 CACATAATCC GCGAGCAAGA CGAGGCCGGC CCCGTGGAGA TCAGCGCCTG A

25 SEQ ID NO:16 (INSP052 Combined polypeptide sequence for exons 1,2,3,4,5,6 and 7.)

1 MKRERGALSR ASRALRLAPF VYLLIQTDP LEGVNITSPV RLIHGTVGKS ALLSVQYSST
61 SSDRPVVWKQ LKRDKPVTVV QSIGTEVIGT LRPDYRDRIR LFENGSLLLS DLQLADEGTY
121 EVEISITDDT FTGEKTIINT VDVPISRPQV LVASTTVLEL SEAFTLNC SH ENGTKPSYTW
181 LKDGPPLLND SRMLLSPDQK VLTITRVLME DDDLYSCMVE NPISQGRSLP VKITVYRRSS
30 241 LYIILSTGGI FLLVTLTV C ACWKPSKRQ KKLEKQNSLE YMDQNDDRLK PEADTLPRSG
301 EGERKNPMAL YILKDKDSPE TEENPAPEPR SATEPGPPGY SVSPAVPGRS PGLPIRSARR
361 YPRSPARSPA TGRTHSSPPR APSSPGRSRS ASRTLRTAGV HIIREQDEAG PVEISA

SEQ ID NO:17 (INSP055 Mouse virtual cDNA)

1 ATGAAGAGAG AAAGGGGAGC CCTGTCAAGA GCCTCCAGGG CTCTGCGCCT CTCTCCTTT
61 GTCTACCTGC TTCTCATCCA GCCAGTCCCC CTGGAGGGGG TGAACATCAC CAGCCCAGTA
121 CGTCTGATCC ACGGCACAGT GGGGAAGTCG GCCCTGCTTT CCGTGCAGTA CAGTAGCACC
5 181 AGCAGCGACA AGCCCCTGGT GAAGTGGCAG CTGAAGCGTG ACAAGCCAGT GACCGTGGTG
241 CAGTCTATAG GCACAGAGGT CATTGGCACT CTGCGGCCTG ACTATCGAGA CCGTATCCGG
301 CTCTTGAAA ATGGCTCCTT GCTTCTCAGC GACCTGCAGC TGGCGGATGA GGGAACCTAT
361 GAAGTGGAGA TTTCCATCAC TGACGACACC TTCACCGGGG AGAACACCAT CAACCTCACC
421 GTGGATGTGC CCATTTCAAG GCCGCAGGTA TTAGTGGCTT CAACCACTGT GCTGGAGCTC
10 481 AGTGAGGCCT TCACCCCTCAA CTGCTCCCAT GAGAATGGCA CCAAGCCTAG CTACACGTGG
541 CTGAAGGATG GCAAACCCCT CCTCAATGAC TCCCGAATGC TCCTGTCCTT TGACCAAAAG
601 GTGCTCACCA TCACCCGAGT ACTCATGGAA GATGACGACC TGTACAGCTG TGTGGTGGAG
661 AACCCCATCA GCCAGGTCCG CAGCCTGCCT GTCAAGATCA CTGTGTATAG AAGAAGCTCC
721 CTCTATATCA TCTTGTCTAC AGGAGGCATC TTCCTCCTTG TGACCCCTGGT GACAGTTTGT
15 781 GCCTGCTGGA AACCCCTAAA AAAGTCTAGG AAGAAGAGGA AGTTGGAGAA GCAAACCTCC
841 TTGGAATACA TGGATCAGAA TGATGACCGC CTAAAATCAG AAGCAGATAC CCTACCCCGA
901 AGTGGAGAAC AGGAGCGGAA GAACCCAATG GCACTCTATA TCCTGAAGGA TAAGGATTCC
961 TCAGAGCCAG ATGAAAACCC TGCTACAGAG CCACGGAGCA CCACAGAACCGGGTCC
1021 GGCTACTCCG TGTCGCCGCC CGTCCCCGGC CGCTCTCCGG GGCTTCCCAT CCGCTCAGCC
20 1081 CGCCGCTACC CGCGCTCCCC AGCACGTTCC CCTGCCACTG GCCGGACGCA CACGTCGCCA
1141 CGCGGGGCC CGAGCTGCC AGGCCGCTCG CGCAGCTCTT CGCGCTCACT GCGGACTGCA
1201 GGCGTGCAGA GAATCCGGGA GCAGGACGAG TCAGGGCAGG TGGAGATCAG TGCCTGA

SEQ ID NO:18 (INSP055 Mouse Predicted Protein)

25 1 MKRERGALSR ASRALRLSPF VYLLIQPVP LEGVNITSPV RLIHGTVGKS ALLSVQYSST
61 SSDKPVVKWQ LKRDKPVTVV QSIGTEVIGT LRPDYRDRIR LFENGSLLLS DLQIADEGTY
121 EVEISITDDT FTGEKTIINLT VDVPISRPQV LVASTTVLEL SEAFTLNC SH ENGTPSYTW
181 LKDGPPLLND SRMLLSPDQK VLTITRVLME DDDLYSCVVE NPISQVRSLP VKITVYRRSS
241 LYIILSTGGI FLLVTLVTVC ACWKPSKKSR KKRKLEKQNS LEYMDQNDDR LKSEADTLPR
30 301 SGEQERKNPM ALYILKDKDS SEPDENPATE PRSTTEPGPP GYSVSPPVPG RSPGLPIRSA
361 RRYPRSPARS PATGRHTSP PRAPSSPGRS RSSSRSRLRTA GVQRIREQDE SGQVEISA