

(43) International Publication Date
30 January 2014 (30.01.2014)(51) International Patent Classification:
A01N 43/40 (2006.01)(21) International Application Number:
PCT/US2013/051299(22) International Filing Date:
19 July 2013 (19.07.2013)


(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
61/675,067 24 July 2012 (24.07.2012) US
13/840,346 15 March 2013 (15.03.2013) US(71) Applicant: DOW AGROSCIENCES LLC [US/US];
9330 Zionsville Road, Indianapolis, IN 46268 (US).(72) Inventors: YERKES, Carla; 4571 South State Road 47,
Crawfordsville, IN 47933 (US). MANN, Richard; 1828
South 550 East, Franklin, IN 46250 (US). SCHMITZER,
Paul; 7690 North Whittier Place, Indianapolis, IN 46250
(US). SATCHIVI, Norbert; 3100 Cherub Court, Carmel,
IN 46074 (US).(74) Agent: BICKLEY, Patrick, M.; Faegre Baker Daniels
LLP, 300 North Meridian Street, Suite 2700, Indianapolis,
IN 46204 (US).(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY,
BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM,
DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT,
HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KN, KP, KR,
KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME,
MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ,
OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC,
SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN,
TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ,
UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ,
TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK,
EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV,
MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM,
TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW,
KM, ML, MR, NE, SN, TD, TG).**Published:**

- with international search report (Art. 21(3))
- before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments (Rule 48.2(h))

(54) Title: HERBICIDAL COMPOSITIONS COMPRISING 4-AMINO-3-CHLORO-5-FLUORO-6-(4-CHLORO-2-FLUORO-3-METHOXYPHENYL) PYRIDINE-2-CARBOXYLIC ACID



(57) Abstract: Provided herein are synergistic herbicidal compositions containing (a) a compound of formula (I): 4-amino-3-chloro-5-fluoro-6-(4-chloro-2-fluoro-3-methoxyphenyl)pyridine-2-carboxylic acid or a derivative thereof, or an agriculturally acceptable salt or ester thereof and (b) a protoporphyrinogen oxidase inhibitor, e.g., acifluorfen, azafeniden, bifenox, carfentrazone-ethyl-ethyl, flumioxazin, ipfencarbazone, oxadiargyl, oxadiazon, oxyfluorfen, pentoxazone, pyraclonil, pyraflufenethyl and saflufenacil, or a salt or ester thereof. The compositions and methods provided herein provide control of undesirable vegetation, e.g., in direct-seeded, water-seeded and transplanted rice, cereals, wheat, barley, oats, rye, sorghum, corn or maize, sugarcane, sunflower, oilseed rape, canola, sugar beet, soybean, cotton, pineapple, pastures, grasslands, rangelands, fallowland, turf, tree and vine orchards, aquatics, plantation crops, vegetables, industrial vegetation management (IVM) and rights-of-way (ROW).

HERBICIDAL COMPOSITIONS COMPRISING 4-AMINO-3-CHLORO-5-FLUORO-6-(4-CHLORO-2-FLUORO-3-METHOXYPHENYL) PYRIDINE-2-CARBOXYLIC ACID OR A DERIVATIVE THEREOF AND A PROTOPORPHYRINOGEN OXIDASE INHIBITOR

Priority Claim

5 This application claims the benefit of United States provisional patent application number 61/675,067 filed on July 24, 2012, and United States patent application serial number 13/840,346, filed March 15, 2013, the disclosure of each of which is incorporated herein by reference in its entirety.

Field


10 Provided herein are herbicidal compositions comprising (a) 4-amino-3-chloro-5-fluoro-6-(4-chloro-2-fluoro-3-methoxyphenyl)pyridine-2-carboxylic acid or an agriculturally acceptable ester or salt thereof and (b) a protoporphyrinogen oxidase inhibitor. Provided herein are also methods of controlling undesirable vegetation comprising applying (a) 4-amino-3-chloro-5-fluoro-6-(4-chloro-2-fluoro-3-methoxyphenyl)pyridine-2-carboxylic acid
15 or an agriculturally acceptable ester or salt thereof and (b) a protoporphyrinogen oxidase inhibitor.

Background

 The protection of crops from weeds and other vegetation which inhibit crop growth is a constantly recurring problem in agriculture. To help combat this problem, researchers in
20 the field of synthetic chemistry have produced an extensive variety of chemicals and chemical formulations effective in the control of such unwanted growth. Chemical herbicides of many types have been disclosed in the literature and a large number are in commercial use. However, there remains a need for compositions and methods that are effective in controlling undesirable vegetation.

25 Summary

 Provided herein are herbicidal compositions comprising a herbicidally effective amount of (a) a compound of the formula (I)




or an agriculturally acceptable salt or ester of thereof, and (b) a protoporphyrinogen oxidase inhibitor. The compositions may also contain an agriculturally acceptable adjuvant or carrier. Provided herein are also methods of controlling undesirable vegetation comprising applying

5 (a) a compound of formula (I) or an agriculturally acceptable ester or salt thereof and (b) a protoporphyrinogen oxidase inhibitor, *e.g.*, acifluorfen, azafeniden, bifenox, carfentrazone-ethyl, flumioxazin, ipfencarbazone, oxadiargyl, oxadiazon, oxyfluorfen, pentoxazone, pyraclonil, pyraflufen and saflufenacil, or an agriculturally acceptable salt or ester thereof.

Several embodiments are recited below. In the embodiments, the ratio of compound

10 (a) to compound (b) can be expressed in units of weight to weight (g to g), gae/ha to gae/ha or gae/ha to gai/ha.

1. A synergistic herbicidal composition comprising a herbicidally effective amount of (a) a compound of the formula (I)



15 or an agriculturally acceptable salt or ester thereof and (b) a protoporphyrinogen oxidase inhibitor.

2. The composition of embodiment 1, wherein (a) is a C₁₋₄ alkyl or benzyl ester of compound (I).
3. The composition of embodiment 2, wherein (a) is a benzyl ester of compound (I).
- 20 4. The composition of embodiment 1, wherein (a) is the compound of formula (I), which is the carboxylic acid.

5. The composition of any of embodiments 1-4, wherein (b) is acifluorfen, azafeniden, bifenox, carfentrazone-ethyl, flumioxazin, ipfencarbazone, oxadiargyl, oxadiazon, oxyfluorfen, pentoxazone, pyraclonil, pyraflufen-ethyl and saflufenacil, or agriculturally acceptable salt, carboxylic acid, carboxylate salt, or ester thereof.
- 5 6. The composition of embodiment 1, wherein (a) is the compound of formula (I) or an agriculturally acceptable benzyl ester and (b) is acifluorfen sodium.
7. The composition of embodiment 1, wherein (a) is the compound of formula (I) or an agriculturally acceptable benzyl ester and (b) is azafeniden.
8. The composition of embodiment 1, wherein (a) is the compound of formula (I) or an
10 agriculturally acceptable benzyl ester and (b) is bifenox.
9. The composition of embodiment 1, wherein (a) is the compound of formula (I) or an agriculturally acceptable benzyl ester and (b) is carfentrazone-ethyl.
10. The composition of embodiment 1, wherein (a) is the compound of formula (I) or an agriculturally acceptable benzyl ester and (b) is ipfencarbazone.
- 15 11. The composition of embodiment 1, wherein (a) is the compound of formula (I) or an agriculturally acceptable benzyl ester and (b) is flumioxazin.
12. The composition of embodiment 1, wherein (a) is the compound of formula (I) or an agriculturally acceptable benzyl ester and (b) is oxadiargyl.
13. The composition of embodiment 1, wherein (a) is the compound of formula (I) or an
20 agriculturally acceptable benzyl ester and (b) is oxadiazon.
14. The composition of embodiment 1, wherein (a) is the compound of formula (I) or an agriculturally acceptable benzyl ester and (b) is oxyfluorfen.
15. The composition of embodiment 1, wherein (a) is the compound of formula (I) or an agriculturally acceptable benzyl ester and (b) is pentoxazone.
- 25 16. The composition of embodiment 1, wherein (a) is the compound of formula (I) or an agriculturally acceptable benzyl ester and (b) is pyraclonil.
17. The composition of embodiment 1, wherein (a) is the compound of formula (I) or an agriculturally acceptable benzyl ester and (b) is pyraflufen-ethyl.
18. The composition of embodiment 1, wherein (a) is the compound of formula (I) or an
30 agriculturally acceptable benzyl ester and (b) is saflufenacil.
19. The composition of any of embodiments 1-18, further comprising a herbicide safener.

20. The composition of embodiment 5 wherein the weight ratio of the compound of formula (I) or agriculturally acceptable salt or ester thereof to acifluorfen or agriculturally acceptable salt thereof is from about 1:210 to about 9:1.
21. The composition of embodiment 5 wherein the weight ratio of the compound of formula (I) or agriculturally acceptable salt or ester thereof to azafeniden or agriculturally acceptable salt thereof is from about 1:220 to about 10:1.
22. The composition of embodiment 5 wherein the weight ratio of the compound of formula (I) or agriculturally acceptable salt or ester thereof to bifenox or agriculturally acceptable carboxylic acid, carboxylate salt, or ester thereof is from about 1:500 to about 3:1.
23. The composition of embodiment 5, wherein the weight ratio of compound of formula (I) or agriculturally acceptable salt or ester thereof to carfentrazone-ethyl or agriculturally acceptable salt or ester thereof is from about 1:60 to about 86:1.
24. The composition of embodiment 5, wherein the weight ratio of the compound of formula (I) or agriculturally acceptable salt or ester thereof to ipfencarbazone or agriculturally acceptable salt thereof is from about 1:250 to about 5:1.
25. The composition of embodiment 5, wherein the weight ratio of the compound of formula (I) or agriculturally acceptable salt or ester thereof to flumioxazin or agriculturally acceptable salt thereof is from about 1:300 to about 17:1.
26. The composition of embodiment 5, wherein the weight ratio of the compound of formula (I) or agriculturally acceptable salt or ester thereof to oxadiargyl or agriculturally acceptable salt thereof is from about 1:1100 to about 6:1.
27. The composition of embodiment 5 wherein the weight ratio of the compound of formula (I) or agriculturally acceptable salt or ester thereof to oxadiazon or agriculturally acceptable salt thereof is from about 1:2240 to about 10:1.
28. The composition of embodiment 5 wherein the weight ratio of the compound of formula (I) or agriculturally acceptable salt or ester thereof to oxyfluorfen is from about 1:1120 to about 3:1.
29. The composition of embodiment 5 wherein the weight ratio of the compound of formula (I) or agriculturally acceptable salt or ester thereof to pentoxazone or agriculturally acceptable salt thereof is from about 1:225 to about 6:1.

30. The composition of embodiment 5 wherein the weight ratio of the compound of formula (I) or agriculturally acceptable salt or ester thereof to pyraclonil or agriculturally acceptable salt thereof is from about 1:180 to about 9:1.

31. The composition of embodiment 5 wherein the weight ratio of the compound of formula (I) or agriculturally acceptable salt or ester thereof to pyraflufen-ethyl or agriculturally acceptable carboxylic acid, carboxylate salt, or ester thereof is from about 1:6 to about 150:1.


32. The composition of any of embodiments 5 wherein the weight ratio of the compound of formula (I) or agriculturally acceptable salt or ester thereof to saflufenacil or agriculturally acceptable salt thereof is from about 1:100 to about 60:1.

33. The composition of any of embodiments 1-32, further comprising an agriculturally acceptable adjuvant or carrier.

34. The composition of any of embodiments 1-32 which is synergistic as determined by the Colby equation.

35. A method of controlling undesirable vegetation which comprises contacting the vegetation or the locus thereof with or applying to the soil or water to prevent the emergence or growth of vegetation the composition of any of embodiments 1-34.

36. A method of controlling undesirable vegetation which comprises contacting the vegetation or the locus thereof with or applying to the soil or water to prevent the emergence or growth of vegetation a herbicidally effective amount of (a) a compound of the formula (I)



or an agriculturally acceptable salt or ester thereof and (b) a protoporphyrinogen oxidase inhibitor.

37. The method of embodiment 35 or 36, wherein the method is practiced in at least one member of the group consisting of direct-seeded, water-seeded and transplanted rice, cereals, wheat, barley, oats, rye, sorghum, corn, maize, sugarcane, sunflower, oilseed rape, canola, sugar beet, soybean, cotton, pineapple, pastures, grasslands, rangelands, fallowland, turf, tree

and vine orchards, aquatics, plantation crops, vegetables, industrial vegetation management (IVM) or rights-of-way (ROW).

38. The method of embodiment 35 or 36, wherein the undesirable vegetation is immature.

39. The method of embodiment 35 or 36, wherein the (a) and (b) are applied to water.

5 40. The method of embodiment 39, wherein the water is part of a flooded rice paddy.

41. The method of embodiment 35 or 36, wherein the (a) and (b) are applied pre-emergently to the weed or the crop.

42. The method of embodiment 35 or 36, wherein the (a) and (b) are applied post-emergently to the weed or the crop.

10 43. The method of of embodiment 35 or 36, wherein the undesirable vegetation is controlled in glyphosate-, 5-enolpyruvylshikimate-3-phosphate (EPSP) synthase inhibitor-, glufosinate-, glutamine synthetase inhibitor-, dicamba-, phenoxy auxin-, pyridyloxy auxin-, synthetic auxin-, auxin transport inhibitor-, aryloxyphenoxypropionate-, cyclohexanedione-, phenylpyrazoline-, acetyl CoA carboxylase (ACCase) inhibitor-, imidazolinone-,
15 sulfonyleurea-, pyrimidinylthiobenzoate-, triazolopyrimidine-, sulfonyleaminocarbonyl triazolinone-, acetolactate synthase (ALS) or acetohydroxy acid synthase (AHAS) inhibitor-, 4-hydroxyphenyl-pyruvate dioxygenase (HPPD) inhibitor-, phytoene desaturase inhibitor-, carotenoid biosynthesis inhibitor-, protoporphyrinogen oxidase (PPO) inhibitor-, cellulose biosynthesis inhibitor-, mitosis inhibitor-, microtubule
20 inhibitor-, very long chain fatty acid inhibitor-, fatty acid and lipid biosynthesis inhibitor-, photosystem I inhibitor-, photosystem II inhibitor-, triazine-, or bromoxynil- tolerant crops.

44. The method of embodiment 43, wherein the tolerant crop possesses multiple or stacked traits conferring tolerance to multiple herbicides or multiple modes of action.

45. The method of embodiment 35 or 36, wherein the undesirable vegetation comprises a
25 herbicide resistant or tolerant weed.

46. The method of embodiment 45, wherein the resistant or tolerant weed is a biotype with resistance or tolerance to multiple herbicides, multiple chemical classes, multiple herbicide modes-of-action or via multiple resistance mechanisms.

47. The method of embodiment 45, wherein the resistant or tolerant weed is a biotype
30 resistant or tolerant to acetolactate synthase (ALS) or acetohydroxy acid synthase (AHAS) inhibitors, photosystem II inhibitors, acetyl CoA carboxylase (ACCase) inhibitors, synthetic auxins, auxin transport inhibitors, photosystem I inhibitors, 5-enolpyruvylshikimate-3-

phosphate (EPSP) synthase inhibitors, microtubule assembly inhibitors, fatty acid and lipid synthesis inhibitors, protoporphyrinogen oxidase (PPO) inhibitors, carotenoid biosynthesis inhibitors, very long chain fatty acid (VLCFA) inhibitors, phytoene desaturase (PDS) inhibitors, glutamine synthetase inhibitors, 4-hydroxyphenyl-pyruvate-dioxygenase (HPPD) inhibitors, mitosis inhibitors, cellulose biosynthesis inhibitors, herbicides with multiple modes-of-action, quinclorac, arylaminopropionic acids, difenzoquat, endothall, or organoarsenicals.

48. The composition of any of embodiments 1-5, wherein (b) is acifluorfen, azafeniden, bifenox, carfentrazone-ethyl, flumioxazin, ipfencarbazone, oxadiargyl, oxadiazon, oxyfluorfen, pentoxazone, pyraclonil, pyraflufen-ethyl and saflufenacil, or agriculturally acceptable salt, carboxylic acid, carboxylate salt, or ester thereof.

49. The composition of embodiment 1, wherein (a) is the compound of formula (I) or an agriculturally acceptable benzyl ester and (b) is acifluorfen sodium.

50. The composition of embodiment 1, wherein (a) is the compound of formula (I) or an agriculturally acceptable benzyl ester and (b) is azafeniden.

51. The composition of embodiment 1, wherein (a) is the compound of formula (I) or an agriculturally acceptable benzyl ester and (b) is bifenox.

52. The composition of embodiment 1, wherein (a) is the compound of formula (I) or an agriculturally acceptable benzyl ester and (b) is carfentrazone-ethyl.

53. The composition of embodiment 1, wherein (a) is the compound of formula (I) or an agriculturally acceptable benzyl ester and (b) is ipfencarbazone.

54. The composition of embodiment 1, wherein (a) is the compound of formula (I) or an agriculturally acceptable benzyl ester and (b) is flumioxazin.

55. The composition of embodiment 1, wherein (a) is the compound of formula (I) or an agriculturally acceptable benzyl ester and (b) is oxadiargyl.

56. The composition of embodiment 1, wherein (a) is the compound of formula (I) or an agriculturally acceptable benzyl ester and (b) is oxadiazon.

57. The composition of embodiment 1, wherein (a) is the compound of formula (I) or an agriculturally acceptable benzyl ester and (b) is oxyfluorfen.

58. The composition of embodiment 1, wherein (a) is the compound of formula (I) or an agriculturally acceptable benzyl ester and (b) is pentoxazone.

59. The composition of embodiment 1, wherein (a) is the compound of formula (I) or an agriculturally acceptable benzyl ester and (b) is pyraclonil.
60. The composition of embodiment 1, wherein (a) is the compound of formula (I) or an agriculturally acceptable benzyl ester and (b) is pyraflufen-ethyl.
- 5 61. The composition of embodiment 1, wherein (a) is the compound of formula (I) or an agriculturally acceptable benzyl ester and (b) is saflufenacil.
62. The composition of embodiment 48 wherein the weight ratio of the compound of formula (I) or agriculturally acceptable salt or ester thereof to acifluorfen, acifluorfen sodium, or agriculturally acceptable salt thereof is from about 1:210 to about 9:1.
- 10 63. The composition of embodiment 48 wherein the weight ratio of the compound of formula (I) or agriculturally acceptable salt or ester thereof to azafeniden is from about 1:220 to about 10:1.
64. The composition of embodiment 48 wherein the weight ratio of the compound of formula (I) or agriculturally acceptable salt or ester thereof to bifenox or agriculturally acceptable carboxylic acid, carboxylate salt, or ester thereof is from about 1:500 to about 3:1.
- 15 65. The composition of embodiment 48, wherein the weight ratio of compound of formula (I) or agriculturally acceptable salt or ester thereof to carfentrazone-ethyl or agriculturally acceptable salt or ester thereof is from about 1:60 to about 86:1.
66. The composition of embodiment 48 wherein the weight ratio of the compound of formula (I) or agriculturally acceptable salt or ester thereof to ipfencarbazone or agriculturally acceptable salt thereof is from about 1:250 to about 5:1.
- 20 67. The composition of embodiment 48 wherein the weight ratio of the compound of formula (I) or agriculturally acceptable salt or ester thereof to flumioxazin or agriculturally acceptable salt thereof is from about 1:300 to about 17:1.
- 25 68. The composition of embodiment 48 wherein the weight ratio of the compound of formula (I) or agriculturally acceptable salt or ester thereof to oxadiargyl or agriculturally acceptable salt thereof is from about 1:1100 to about 6:1.
69. The composition of embodiment 48 wherein the weight ratio of the compound of formula (I) or agriculturally acceptable salt or ester thereof to oxadiazon or agriculturally acceptable salt thereof is from about 1:2240 to about 10:1.
- 30

70. The composition of embodiment 48 wherein the weight ratio of the compound of formula (I) or agriculturally acceptable salt or ester thereof to oxyfluorfen is from about 1:1120 to about 3:1.

71. The composition of embodiment 48 wherein the weight ratio of the compound of formula (I) or agriculturally acceptable salt or ester thereof to pentoxazone or agriculturally acceptable salt thereof is from about 1:225 to about 6:1.

72. The composition of embodiment 48 wherein the weight ratio of the compound of formula (I) or agriculturally acceptable salt or ester thereof to pyraclonil or agriculturally acceptable salt thereof is from about 1:180 to about 9:1.

73. The composition of embodiment 48 wherein the weight ratio of the compound of formula (I) or agriculturally acceptable salt or ester thereof to pyraflufen-ethyl or agriculturally acceptable carboxylic acid, carboxylate salt, or ester thereof is from about 1:6 to about 150:1.

74. The composition of embodiment 48, wherein the weight ratio of the compound of formula (I) or agriculturally acceptable salt or ester thereof to saflufenacil or agriculturally acceptable salt thereof is from about 1:100 to about 60:1.

75. The composition of any of embodiments 1-34 or 48-74, further comprising an agriculturally acceptable adjuvant or carrier.

76. The composition of any of embodiments 1-34 or 48-74, further comprising a herbicide safener.

77. The composition of any of embodiments 1-34 or 48-74, wherein the ratio of compound (a) and compound (b) are present in proportions that result in synergistic herbicidal activity as determined by the Colby equation.


78. A method of controlling undesirable vegetation which comprises contacting the vegetation or the locus thereof with or applying to the soil or water to prevent the emergence or growth of vegetation the composition of any of embodiments 1-34 or 48-74.

79. A method according to any of embodiments 35-47, wherein the undesirable vegetation is immature.

Detailed Description

DEFINITIONS


As used herein, the compound of formula (I) has the following structure:



The compound of formula (I) can be identified by the name 4-amino-3-chloro-6-(4-chloro-2-fluoro-3-methoxyphenyl)-5-fluoropyridine-2-carboxylic acid and has been described in U.S. Patent 7,314,849 (B2), which is incorporated herein by reference in its entirety. Exemplary uses of the compound of the formula (I) include controlling undesirable vegetation, including grass, broadleaf and sedge weeds, in multiple non-crop and cropping situations.


Without being limited to any theory, protoporphyrinogen oxidase inhibitors are believed to interfere with chlorophyll, lipid and protein production, causing plant cells to dry and disintegrate. Exemplary protoporphyrinogen oxidase inhibitors include, but are not limited to, acifluorfen, azafeniden, bifenox, carfentrazone-ethyl, flumioxazin, ipfencarbazone, oxadiargyl, oxadiazon, oxyfluorfen, pentoxazone, pyraclonil, pyraflufen-ethyl and saflufenacil and derivatives thereof.

As used herein, acifluorfen is 5-[2-chloro-4-(trifluoromethyl)phenoxy]-2-nitrobenzoic acid. Acifluorfen possesses the following structure:




The herbicidal activity of acifluorfen is exemplified in Tomlin, C., ed. A World Compendium The Pesticide Manual. 15th ed. Alton: BCPC Publications, 2009 (hereafter "*The Pesticide Manual*, Fifteenth Edition, 2009."). An exemplary form of acifluorfen includes its sodium salt. Exemplary uses of acifluorfen include its use for post-emergence control of broadleaf weeds in soybeans and rice.

As used herein, azafenidin is 2-[2,4-dichloro-5-(2-propynyloxy)phenyl]-5,6,7,8-tetrahydro-1,2,4-triazolo[4,3-*a*]pyridin-3(2*H*)-one and possesses the following structure:




Its herbicidal activity is exemplified in *The Pesticide Manual*, Fifteenth Edition, 2009. Exemplary uses of azafenidin include as a pre-emergence, and, in mixture, post-emergence, herbicide in citrus, grapes, olives, sugar cane and other perennial crops.

5 As used herein, bifenox is methyl 5-(2,4-dichlorophenoxy)-2-nitrobenzoate. Bifenox possesses the following structure:




The herbicidal activity of bifenox is exemplified in *The Pesticide Manual*, Fifteenth Edition, 2009. Exemplary uses of bifenox include its use for control of barnyardgrass and
10 sedges, *e.g.*, in transplanted rice.

As used herein, carfentrazone-ethyl is ethyl α ,2-dichloro-5-[4-(difluoromethyl)-4,5-dihydro-3-methyl-5-oxo-1*H*-1,2,4-triazol-1-yl]-4-fluorobenzenepropanoate. Carfentrazone-ethyl possesses the following structure:




15 The herbicidal activity of carfentrazone-ethyl is exemplified in *The Pesticide Manual*, Fifteenth Edition, 2009. Exemplary uses of carfentrazone-ethyl include its use for post-emergence control of a wide range of broadleaf weeds, *e.g.*, in cereals.

As used herein, flumioxazin is 2-[7-fluoro-3,4-dihydro-3-oxo-4-(2-propynyl)-2*H*-1,4-benzoxazin-6-yl]-4,5,6,7-tetrahydro-1*H*-isoindole-1,3(2*H*)-dione. Flumioxazin possesses the
20 following structure:




The herbicidal activity of flumioxazin is exemplified in *The Pesticide Manual*, Fifteenth Edition, 2009. Exemplary uses of flumioxazin include its use for pre-emergence control of broad-leaved weeds and grasses, *e.g.*, in soybeans and tree and vine crops.

- 5 As used herein, oxadiargyl is 3-[2,4-dichloro-5-(2-propynyloxy)phenyl]-5-(1,1-dimethylethyl)-1,3,4-oxadiazol-2(3*H*)-one. Oxadiargyl possesses the following structure:




- 10 Its herbicidal activity is exemplified in *The Pesticide Manual*, Fifteenth Edition, 2009. Exemplary uses of oxadiargyl include its use for pre-emergence and early post-emergence control of broad-leaved weeds, grasses and annual sedges, *e.g.*, in rice, sunflowers, potatoes, vegetables, sugar cane, fruit trees, and citrus.

As used herein, oxadiazon is 3-[2,4-dichloro-5-(1-methylethoxy)phenyl]-5-(1,1-dimethylethyl)-1,3,4-oxadiazol-2(3*H*)-one. Oxadiazon possesses the following structure:




- 15 Its herbicidal activity is exemplified in *The Pesticide Manual*, Fifteenth Edition, 2009. Exemplary uses of oxadiazon include its use for control of mono- and dicotyledonous weeds, *e.g.*, in rice, and pre-emergence control of bindweed, annual broad-leaved weeds and grass; and post-emergence control of bindweed and annual broad-leaved weeds, *e.g.*, in carnations, gladioli, roses, fruit trees and bushes, vines, ornamental trees and shrubs, hops, cotton, rice, soya beans, sunflower, onions and turf.

20 As used herein, oxyfluorfen is 2-chloro-1-(3-ethoxy-4-nitrophenoxy)-4-(trifluoromethyl)benzene. It possesses the following structure:




Its herbicidal activity is exemplified in *The Pesticide Manual*, Fifteenth Edition, 2009. Exemplary uses of oxyfluorfen include its use for pre- and post-emergence control of annual broadleaved and grass weeds, *e.g.*, in tropical and subtropical crops.

- 5 As used herein, pentoxazone, is 3-[4-chloro-5-(cyclopentyloxy)-2-fluorophenyl]-5-(1-methylethylidene)-2,4-oxazolidinedione and possesses the following structure:




Its herbicidal activity is exemplified in *The Pesticide Manual*, Fifteenth Edition, 2009. Exemplary uses of pentoxazone include its use for pre- and post-emergence control of *Echinochloa* spp. and *Monchoria vaginalis*, *e.g.*, in paddy rice, before, during or after transplantation.

As used herein, pyraclonil is 1-(3-chloro-4,5,6,7-tetrahydropyrazolo[1,5-*a*]pyridin-2-yl)-5-(methyl-2-propynylamino)-1*H*-pyrazole-4-carbonitrile and possesses the following structure:




Exemplary uses of pyraclonil include its use for weed control in rice.

As used herein, pyraflufen-ethyl is ethyl 2-chloro-5-(4-chloro-5-difluoromethoxy-1-methylpyrazol-3-yl)-4-fluorophenoxy]acetate. Pyraflufen-ethyl possesses the following structure:



The herbicidal activity of pyraflufen-ethyl is exemplified in *The Pesticide Manual*, Fifteenth Edition, 2009. Exemplary uses of pyraflufen-ethyl include its use for post-emergence control of broadleaf weeds, *e.g.*, in cereals.

- 5 As used herein, saflufenacil is 2-chloro-5-[3,6-dihydro-3-methyl-2,6-dioxo-4-(trifluoromethyl)-1(2*H*)-pyrimidinyl]-4-fluoro-*N*-[[methyl(1-methylethyl)amino]sulfonyl]benzamide and saflufenacil possesses the following structure:



Its herbicidal activity is exemplified in *The Pesticide Manual*, Fifteenth Edition, 2009.

- 10 Exemplary uses of saflufenacil include its use for foliar and residual control of broadleaf weeds.

As used herein, herbicide means a compound, *e.g.*, active ingredient that kills, controls or otherwise adversely modifies the growth of plants.

- 15 As used herein, a herbicidally effective or vegetation controlling amount is an amount of active ingredient which causes an adversely modifying effect to the vegetation *e.g.*, causing deviations from natural development, killing, effecting regulation, causing desiccation, causing retardation, and the like.

As used herein, controlling undesirable vegetation means preventing, reducing, killing, or otherwise adversely modifying the development of plants and vegetation.


- 20 Described herein are methods of controlling undesirable vegetation through the application of certain herbicide combinations or compositions. Methods of application include, but are not limited to applications to the vegetation or locus thereof, *e.g.*, application to the area adjacent to the vegetation, as well as preemergence, postemergence, foliar (broadcast, directed, banded, spot, mechanical, over-the-top, or rescue), and in-water applications (emerged and
25 submerged vegetation, broadcast, spot, mechanical, water-injected, granular broadcast,

granular spot, shaker bottle, or stream spray) via hand, backpack, machine, tractor, or aerial (airplane and helicopter) application methods.

As used herein, plants and vegetation include, but are not limited to, germinant seeds, emerging seedlings, plants emerging from vegetative propagules, immature vegetation, and established vegetation.

As used herein, agriculturally acceptable salts and esters refer to salts and esters that exhibit herbicidal activity, or that are or can be converted in plants, water, or soil to the referenced herbicide. Exemplary agriculturally acceptable esters are those that are or can be hydrolyzed, oxidized, metabolized, or otherwise converted, *e.g.*, in plants, water, or soil, to the corresponding carboxylic acid which, depending on the pH, may be in the dissociated or undissociated form.

Exemplary salts include those derived from alkali or alkaline earth metals and those derived from ammonia and amines. Exemplary cations include sodium, potassium, magnesium, and aminium cations of the formula:




wherein R^1 , R^2 , R^3 and R^4 each, independently represents hydrogen or C_1 - C_{12} alkyl, C_3 - C_{12} alkenyl or C_3 - C_{12} alkynyl, each of which is optionally substituted by one or more hydroxy, C_1 - C_4 alkoxy, C_1 - C_4 alkylthio or phenyl groups, provided that R^1 , R^2 , R^3 and R^4 are sterically compatible. Additionally, any two of R^1 , R^2 , R^3 and R^4 together may represent an aliphatic difunctional moiety containing one to twelve carbon atoms and up to two oxygen or sulfur atoms. Salts can be prepared by treatment with a metal hydroxide, such as sodium hydroxide, with an amine, such as ammonia, trimethylamine, diethanolamine, 2-methylthiopropylamine, bisallylamine, 2-butoxyethylamine, morpholine, cyclododecylamine, or benzylamine or with a tetraalkylammonium hydroxide, such as tetramethylammonium hydroxide or choline hydroxide.

Exemplary esters include those derived from C_1 - C_{12} alkyl, C_3 - C_{12} alkenyl, C_3 - C_{12} alkynyl or C_7 - C_{10} aryl-substituted alkyl alcohols, such as methyl alcohol, isopropyl alcohol, 1-butanol, 2-ethylhexanol, butoxyethanol, methoxypropanol, allyl alcohol, propargyl alcohol, cyclohexanol or unsubstituted or substituted benzyl alcohols. Benzyl alcohols may be substituted with from 1-3 substituents independently selected from halogen, C_1 - C_4 alkyl or C_1 - C_4 alkoxy. Esters can be prepared by coupling of the acids with the alcohol using any number of suitable activating agents such as those used for peptide couplings such as

dicyclohexylcarbodiimide (DCC) or carbonyl diimidazole (CDI); by reacting the acids with alkylating agents such as alkylhalides or alkylsulfonates in the presence of a base such as triethylamine or lithium carbonate; by reacting the corresponding acid chloride of an acid with an appropriate alcohol; by reacting the corresponding acid with an appropriate alcohol in the presence of an acid catalyst or by transesterification.

COMPOSITIONS AND METHODS

Provided herein are herbicidal compositions comprising a herbicidally effective amount of (a) a compound of the formula (I)



or an agriculturally acceptable salt or ester of thereof, and (b) a protoporphyrinogen oxidase inhibitor. In some embodiments, the protoporphyrinogen oxidase inhibitor is acifluorfen, azafeniden, bifenox, carfentrazone-ethyl, flumioxazin, ipfencarbazone, oxadiargyl, oxadiazon, oxyfluorfen, pentoxazone, pyraclonil, pyraflufen-ethyl and saflufenacil, or agriculturally acceptable salts or esters thereof.

Provided herein are also methods of controlling undesirable vegetation comprising contacting the vegetation or the locus thereof, *i.e.*, area adjacent to the plant, with or applying to the soil or water to prevent the emergence or growth of vegetation a herbicidally effective amount of (a) the compound of formula (I) and (b) a protoporphyrinogen oxidase inhibitor. In certain embodiments, the methods employ the compositions described herein.

Furthermore, in some embodiments, the combination of compound (I) or agriculturally acceptable salt or ester thereof and a protoporphyrinogen oxidase inhibitor, or an agriculturally acceptable salt or ester thereof exhibit synergism, *e.g.*, the herbicidal active ingredients are more effective in combination than when applied individually. Synergism has been defined as “an interaction of two or more factors such that the effect when combined is greater than the predicted effect based on the response of each factor applied separately.”

Senseman, S., ed. Herbicide Handbook. 9th ed. Lawrence: Weed Science Society of America, 2007. In certain embodiments, the compositions exhibit synergy as determined by

the Colby's equation. Colby, S.R. 1967. Calculation of the synergistic and antagonistic response of herbicide combinations. Weeds 15:20-22.

In certain embodiments of the compositions and methods described herein, the compound of formula (I), *i.e.*, the carboxylic acid, is employed. In certain embodiments, a
5 carboxylate salt of the compound of formula (I) is employed. In certain embodiments, an aralkyl or alkyl ester is employed. In certain embodiments, a benzyl, substituted benzyl, or C₁₋₄ alkyl, *e.g.*, n-butyl ester is employed. In certain embodiments, the benzyl ester is employed.

In some embodiments, the compound of formula (I) or salt or ester thereof and
10 acifluorfen, azafeniden, bifenox, carfentrazone-ethyl, flumioxazin, ipfencarbazone, oxadiargyl, oxadiazon, oxyfluorfen, pentoxazone, pyraclonil, pyraflufen-ethyl and saflufenacil, or an agriculturally acceptable salt or ester thereof are formulated in one composition, tank mixed, applied simultaneously, or applied sequentially.

Herbicidal activity is exhibited by the compounds when they are applied directly to
15 the plant or to the locus of the plant at any stage of growth. The effect observed depends upon the plant species to be controlled, the stage of growth of the plant, the application parameters of dilution and spray drop size, the particle size of solid components, the environmental conditions at the time of use, the specific compound employed, the specific adjuvants and carriers employed, the soil type, and the like, as well as the amount of chemical
20 applied. These and other factors can be adjusted to promote non-selective or selective herbicidal action. In some embodiments, the compositions described herein are applied as a post-emergence application, pre-emergence application, or in-water application to flooded paddy rice or water bodies (*e.g.*, ponds, lakes and streams), to relatively immature undesirable vegetation to achieve the maximum control of weeds.

In some embodiments, the compositions and methods provided herein are utilized to
25 control weeds in crops, including but not limited to direct-seeded, water-seeded and transplanted rice, cereals, wheat, barley, oats, rye, sorghum, corn/maize, sugarcane, sunflower, oilseed rape, canola, sugar beet, soybean, cotton, pineapple, pastures, grasslands, rangelands, fallowland, turf, tree and vine orchards, aquatics, plantation crops, vegetables,
30 industrial vegetation management (IVM) and rights-of-way (ROW).

In certain embodiments, the compositions and methods provided herein are utilized to control weeds in rice. In certain embodiments, the rice is direct-seeded, water-seeded, or transplanted rice.

The compositions and methods described herein may be used to control undesirable
5 vegetation in glyphosate-tolerant-, 5-enolpyruvylshikimate-3-phosphate (EPSP) synthase
inhibitor-tolerant-, glufosinate-tolerant-, glutamine synthetase inhibitor-tolerant-, dicamba-
tolerant-, phenoxy auxin-tolerant-, pyridyloxy auxin-tolerant-, auxin-tolerant-, auxin transport
inhibitor-tolerant-, aryloxyphenoxypropionate-tolerant-, acetyl CoA carboxylase (ACCase)
inhibitor-tolerant-, sulfonyleurea-tolerant-, pyrimidinylthiobenzoate-tolerant-,
10 triazolopyrimidine-tolerant-, sulfonylaminocarbonyltriazolinone-tolerant-, imidazolinone-
tolerant-, acetolactate synthase (ALS) or acetohydroxy acid synthase (AHAS) inhibitor-
tolerant-, 4-hydroxyphenyl-pyruvate dioxygenase (HPPD) inhibitor -tolerant-, phytoene
desaturase inhibitor-tolerant-, carotenoid biosynthesis inhibitor-tolerant-, protoporphyrinogen
oxidase (PPO) inhibitor -tolerant-, cellulose biosynthesis inhibitor-tolerant-, mitosis inhibitor-
15 tolerant-, microtubule inhibitor-tolerant-, very long chain fatty acid inhibitor-tolerant-, fatty
acid and lipid biosynthesis inhibitor-tolerant-, photosystem I inhibitor-tolerant-, photosystem
II inhibitor-tolerant-, triazine-tolerant-, and bromoxynil-tolerant- crops (such as, but not
limited to, soybean, cotton, canola/oilseed rape, rice, cereals, corn, sorghum, sunflower, sugar
beet, sugarcane, turf, etc.), for example, in conjunction with glyphosate, EPSP synthase
20 inhibitors, glufosinate, glutamine synthase inhibitors, dicamba, phenoxy auxins, pyridyloxy
auxins, synthetic auxins, auxin transport inhibitors, aryloxyphenoxypropionates,
cyclohexanediones, phenylpyrazolines, ACCase inhibitors, imidazolinones, sulfonyleureas,
pyrimidinylthiobenzoates, triazolopyrimidines, sulfonylaminocarbonyltriazolinones, ALS or
AHAS inhibitors, HPPD inhibitors, phytoene desaturase inhibitors, carotenoid biosynthesis
25 inhibitors, PPO inhibitors, cellulose biosynthesis inhibitors, mitosis inhibitors, microtubule
inhibitors, very long chain fatty acid inhibitors, fatty acid and lipid biosynthesis inhibitors,
photosystem I inhibitors, photosystem II inhibitors, triazines, and bromoxynil. The
compositions and methods may be used in controlling undesirable vegetation in crops
possessing multiple or stacked traits conferring tolerance to multiple chemistries and/or
30 inhibitors of multiple modes of action. In some embodiments, the compound of formula (I)
or salt or ester thereof and complementary herbicide or salt or ester thereof are used in
combination with herbicides that are selective for the crop being treated and which

complement the spectrum of weeds controlled by these compounds at the application rate employed. In some embodiments, the compositions described herein and other complementary herbicides are applied at the same time, either as a combination formulation, as a tank mix, or sequentially.

5 The compositions and methods may be used in controlling undesirable vegetation in crops possessing agronomic stress tolerance (including but not limited to drought, cold, heat, salt, water, nutrient, fertility, pH), pest tolerance (including but not limited to insects, fungi and pathogens) and crop improvement traits (including but not limited to yield; protein, carbohydrate, or oil content; protein, carbohydrate, or oil composition; plant stature and plant
10 architecture).

The compositions and methods provided herein are utilized to control undesirable vegetation. Undesirable vegetation includes, but is not limited to, undesirable vegetation that occurs in rice, cereals, wheat, barley, oats, rye, sorghum, corn/maize, sugarcane, sunflower, oilseed rape, canola, sugar beet, soybean, cotton, pineapple, pastures, grasslands, rangelands,
15 fallowland, turf, tree and vine orchards, aquatics, plantation crops, vegetables, industrial vegetation management (IVM) and rights of way (ROW).

In some embodiments, the methods provided herein are utilized to control undesirable vegetation in rice. In certain embodiments, the undesirable vegetation is *Brachiaria platyphylla* (Groseb.) Nash or *Urochloa platyphylla* (Nash) R.D. Webster (broadleaf
20 signalgrass, BRAPP), *Digitaria sanguinalis* (L.) Scop. (large crabgrass, DIGSA), *Echinochloa* species (ECHSS), *Echinochloa crus-galli* (L.) P. Beauv. (barnyardgrass, ECHCG), *Echinochloa crus-pavonis* (Kunth) Schult. (gulf cockspur, ECHCV), *Echinochloa colonum* (L.) LINK (junglerice, ECHCO), *Echinochloa oryzoides* (Ard.) Fritsch (early watergrass, ECHOR), *Echinochloa oryzicola* (Vasinger) Vasinger (late watergrass, ECHPH),
25 *Echinochloa phyllopogon* (Stapf) Koso-Pol. (rice barnyardgrass, ECHPH), *Echinochloa polystachya* (Kunth) Hitchc. (creeping river grass, ECHPO), *Ischaemum rugosum* Salisb. (saramollagrass, ISCRU), *Leptochloa chinensis* (L.) Nees (Chinese sprangletop, LEFCH), *Leptochloa fascicularis* (Lam.) Gray (bearded sprangletop, LEFFA), *Leptochloa panicoides* (Presl.) Hitchc. (Amazon sprangletop, LEFPA), *Oryza* species (red and weedy rice, ORYSS),
30 *Panicum dichotomiflorum* (L.) Michx. (fall panicum, PANDI), *Paspalum dilatatum* Poir. (dallisgrass, PASDI), *Rottboellia cochinchinensis* (Lour.) W.D. Clayton (itchgrass, ROOEX), *Cyperus* species (CYPSS), *Cyperus difformis* L. (smallflower flatsedge, CYPDI), *Cyperus*

dubius Rottb. (MAPDU), *Cyperus esculentus* L. (yellow nutsedge, CYPES), *Cyperus iria* L. (rice flatsedge, CYPPIR), *Cyperus rotundus* L. (purple nutsedge, CYPRO), *Cyperus serotinus* Rottb./C.B. Clarke (tidalmarsh flatsedge, CYPSE), *Eleocharis* species (ELOSS), *Fimbristylis miliacea* (L.) Vahl (globe fringerush, FIMMI), *Schoenoplectus* species (SCPSS),
 5 *Schoenoplectus juncooides* Roxb. (Japanese bulrush, SCPJU), *Bolboschoenus maritimus* (L.) Palla or *Schoenoplectus maritimus* L. Lye (sea clubrush, SCPMA), *Schoenoplectus mucronatus* L. (ricefield bulrush, SCPMU), *Aeschynomene* species, (jointvetch, AESSS), *Alternanthera philoxeroides* (Mart.) Griseb. (alligatorweed, ALRPH), *Alisma plantago-aquatica* L. (common waterplantain, ALSPA), *Amaranthus* species, (pigweeds and
 10 amaranths, AMASS), *Ammannia coccinea* Rottb. (redstem, AMMCO), *Commelina benghalensis* L. (Benghal dayflower, COMBE), *Eclipta alba* (L.) Hassk. (American false daisy, ECLAL), *Heteranthera limosa* (SW.) Willd./Vahl (ducksalad, HETLI), *Heteranthera reniformis* R. & P. (roundleaf mudplantain, HETRE), *Ipomoea* species (morningglories, IPOSS), *Ipomoea hederacea* (L.) Jacq. (ivyleaf morningglory, IPOHE), *Lindernia dubia* (L.)
 15 Pennell (low false pimpernel, LIDDU), *Ludwigia* species (LUDSS), *Ludwigia linifolia* Poir. (southeastern primrose-willow, LUDLI), *Ludwigia octovalvis* (Jacq.) Raven (longfruited primrose-willow, LUDOC), *Monochoria korsakowii* Regel & Maack (monochoria, MOOKA), *Monochoria vaginalis* (Burm. F.) C. Presl ex Kuhth, (monochoria, MOOVA), *Murdannia nudiflora* (L.) Brennan (doveweed, MUDNU), *Polygonum pensylvanicum* L.,
 20 (Pennsylvania smartweed, POLPY), *Polygonum persicaria* L. (ladysthumb, POLPE), *Polygonum hydropiperoides* Michx. (POLHP, mild smartweed), *Rotala indica* (Willd.) Koehne (Indian toothcup, ROTIN), *Sagittaria* species, (arrowhead, SAGSS), *Sesbania exaltata* (Raf.) Cory/Rydb. Ex Hill (hemp sesbania, SEBEX), or *Sphenoclea zeylanica* Gaertn. (gooseweed, SPDZE).

25 In some embodiments, the methods provided herein are utilized to control undesirable vegetation in cereals. In certain embodiments, the undesirable vegetation is *Alopecurus myosuroides* Huds. (blackgrass, ALOMY), *Apera spica-venti* (L.) Beauv. (windgrass, APESV), *Avena fatua* L. (wild oat, AVEFA), *Bromus tectorum* L. (downy brome, BROTE), *Lolium multiflorum* Lam. (Italian ryegrass, LOLMU), *Phalaris minor* Retz. (littleseed
 30 canarygrass, PHAMI), *Poa annua* L. (annual bluegrass, POANN), *Setaria pumila* (Poir.) Roemer & J.A. Schultes (yellow foxtail, SETLU), *Setaria viridis* (L.) Beauv. (green foxtail, SETVI), *Amaranthus retroflexus* L. (redroot pigweed, AMARE), *Brassica* species (BRSSS),

Chenopodium album L. (common lambsquarters, CHEAL), *Cirsium arvense* (L.) Scop. (Canada thistle, CIRAR), *Galium aparine* L. (catchweed bedstraw, GALAP), *Kochia scoparia* (L.) Schrad. (kochia, KCHSC), *Lamium purpureum* L. (purple deadnettle, LAMPU), *Matricaria recutita* L. (wild chamomile, MATCH), *Matricaria matricarioides* (Less.) Porter (pineappleweed, MATMT), *Papaver rhoeas* L. (common poppy, PAPRH), *Polygonum convolvulus* L. (wild buckwheat, POLCO), *Salsola tragus* L. (Russian thistle, SASKR), *Sinapis* species (SINSS), *Sinapis arvensis* L. (wild mustard, SINAR), *Stellaria media* (L.) Vill. (common chickweed, STEME), *Veronica persica* Poir. (Persian speedwell, VERPE), *Viola arvensis* Murr. (field violet, VIOAR), or *Viola tricolor* L. (wild violet, VIOTR).

In some embodiments, the methods provided herein are utilized to control undesirable vegetation in range and pasture, fallowland, IVM and ROW. In certain embodiments, the undesirable vegetation is *Ambrosia artemisiifolia* L. (common ragweed, AMBEL), *Cassia obtusifolia* (sickle pod, CASOB), *Centaurea maculosa* auct. non Lam. (spotted knapweed, CENMA), *Cirsium arvense* (L.) Scop. (Canada thistle, CIRAR), *Convolvulus arvensis* L. (field bindweed, CONAR), *Daucus carota* L. (wild carrot, DAUCA), *Euphorbia esula* L. (leafy spurge, EPHES), *Lactuca serriola* L./Torn. (prickly lettuce, LACSE), *Plantago lanceolata* L. (buckhorn plantain, PLALA), *Rumex obtusifolius* L. (broadleaf dock, RUMOB), *Sida spinosa* L. (prickly sida, SIDSP), *Sinapis arvensis* L. (wild mustard, SINAR), *Sonchus arvensis* L. (perennial sowthistle, SONAR), *Solidago* species (goldenrod, SOOSS), *Taraxacum officinale* G.H. Weber ex Wiggers (dandelion, TAROF), *Trifolium repens* L. (white clover, TRFRE), or *Urtica dioica* L. (common nettle, URTDI).

In some embodiments, the methods provided herein are utilized to control undesirable vegetation found in row crops, tree and vine crops, and perennial crops. In certain embodiments, the undesirable vegetation is *Alopecurus myosuroides* Huds. (blackgrass, ALOMY), *Avena fatua* L. (wild oat, AVEFA), *Brachiaria decumbens* Stapf. or *Urochloa decumbens* (Stapf) R.D. Webster (Surinam grass, BRADC), *Brachiaria brizantha* (Hochst. ex A. Rich.) Stapf. or *Urochloa brizantha* (Hochst. ex A. Rich.) R.D. (beard grass, BRABR), *Brachiaria platyphylla* (Groseb.) Nash or *Urochloa platyphylla* (Nash) R.D. Webster (broadleaf signalgrass, BRAPP), *Brachiaria plantaginea* (Link) Hitchc. or *Urochloa plantaginea* (Link) R.D. Webster (alexandergrass, BRAPL), *Cenchrus echinatus* L. (southern sandbur, CENEC), *Digitaria horizontalis* Willd. (Jamaican crabgrass, DIGHO), *Digitaria*

- insularis* (L.) Mez ex Ekman (sourgrass, TRCIN), *Digitaria sanguinalis* (L.) Scop. (large crabgrass, DIGSA), *Echinochloa crus-galli* (L.) P. Beauv. (barnyardgrass, ECHCG), *Echinochloa colonum* (L.) Link (junglerice, ECHCO), *Eleusine indica* (L.) Gaertn. (goosegrass, ELEIN), *Lolium multiflorum* Lam. (Italian ryegrass, LOLMU), *Panicum*
- 5 *dichotomiflorum* Michx. (fall panicum, PANDI), *Panicum miliaceum* L. (wild-proso millet, PANMI), *Setaria faberi* Herrm. (giant foxtail, SETFA), *Setaria viridis* (L.) Beauv. (green foxtail, SETVI), *Sorghum halepense* (L.) Pers. (Johnsongrass, SORHA), *Sorghum bicolor* (L.) Moench ssp. *Arundinaceum* (shattercane, SORVU), *Cyperus esculentus* L. (yellow nutsedge, CYPES), *Cyperus rotundus* L. (purple nutsedge, CYPRO), *Abutilon theophrasti*
- 10 Medik. (velvetleaf, ABUTH), *Amaranthus* species (pigweeds and amaranths, AMASS), *Ambrosia artemisiifolia* L. (common ragweed, AMBEL), *Ambrosia psilostachya* DC. (western ragweed, AMBPS), *Ambrosia trifida* L. (giant ragweed, AMBTR), *Anoda cristata* (L.) Schlecht. (spurred anoda, ANVCR), *Asclepias syriaca* L. (common milkweed, ASCSY), *Bidens pilosa* L. (hairy beggarticks, BIDPI), *Borreria* species (BOISS), *Borreria alata*
- 15 (Aubl.) DC. or *Spermacoce alata* Aubl. (broadleaf buttonweed, BOILF), *Spermacoce latifolia* (broadleaved button weed, BOILF), *Chenopodium album* L. (common lambsquarters, CHEAL), *Cirsium arvense* (L.) Scop. (Canada thistle, CIRAR), *Commelina benghalensis* L. (tropical spiderwort, COMBE), *Datura stramonium* L. (jimsonweed, DATST), *Daucus carota* L. (wild carrot, DAUCA), *Euphorbia heterophylla* L. (wild poinsettia, EPHHL),
- 20 *Euphorbia hirta* L. or *Chamaesyce hirta* (L.) Millsp. (garden spurge, EPHHI), *Euphorbia dentata* Michx. (toothed spurge, EPHDE), *Erigeron bonariensis* L. or *Conyza bonariensis* (L.) Cronq. (hairy fleabane, ERIBO), *Erigeron canadensis* L. or *Conyza canadensis* (L.) Cronq. (Canadian fleabane, ERICA), *Conyza sumatrensis* (Retz.) E. H. Walker (tall fleabane, ERIFL), *Helianthus annuus* L. (common sunflower, HELAN), *Jacquemontia tamnifolia* (L.)
- 25 Griseb. (smallflower morningglory, IAQTA), *Ipomoea hederacea* (L.) Jacq. (ivyleaf morningglory, IPOHE), *Ipomoea lacunosa* L. (white morningglory, IPOLA), *Lactuca serriola* L./Torn. (prickly lettuce, LACSE), *Portulaca oleracea* L. (common purslane, POROL), *Richardia* species (pusley, RCHSS), *Sida* species (sida, SIDSS), *Sida spinosa* L. (prickly sida, SIDSP), *Sinapis arvensis* L. (wild mustard, SINAR), *Solanum ptychanthum*
- 30 Dunal (eastern black nightshade, SOLPT), *Tridax procumbens* L. (coat buttons, TRQPR), or *Xanthium strumarium* L. (common cocklebur, XANST).

In some embodiments, the methods provided herein are utilized to control undesirable vegetation in turf. In certain embodiments, the undesirable vegetation is *Bellis perennis* L. (English daisy, BELPE), *Cyperus esculentus* L. (yellow nutsedge, CYPES), *Cyperus* species (CYPSS), *Digitaria sanguinalis* (L.) Scop. (large crabgrass, DIGSA), *Diodia virginiana* L. (Virginia buttonweed, DIQVI), *Euphorbia* species (spurge, EPHSS), *Glechoma hederacea* L. (ground ivy, GLEHE), *Hydrocotyle umbellata* L. (dollarweed, HYDUM), *Kyllinga* species (kyllinga, KYLSS), *Lamium amplexicaule* L. (henbit, LAMAM), *Murdannia nudiflora* (L.) Brenan (doveweed, MUDNU), *Oxalis* species (woodsorrel, OXASS), *Plantago major* L. (broadleaf plantain, PLAMA), *Plantago lanceolata* L. (buckhorn/narrowleaf plantain, PLALA), *Phyllanthus urinaria* L. (chamberbitter, PYLTE), *Rumex obtusifolius* L. (broadleaf dock, RUMOB), *Stachys floridana* Shuttlew. (Florida betony, STAFL), *Stellaria media* (L.) Vill. (common chickweed, STEME), *Taraxacum officinale* G.H. Weber ex Wiggers (dandelion, TAROF), *Trifolium repens* L. (white clover, TRFRE), or *Viola* species (wild violet, VIOSS).

In some embodiments, the compositions and methods provided herein are utilized to control undesirable vegetation consisting of grass, broadleaf and sedge weeds. In certain embodiments, the compositions and methods provided herein are utilized to control undesirable grass, broadleaf and sedge vegetation including but not limited to *Brassica*, *Brachiaria* or *Urochloa*, *Bolboschoenus*, *Cyperus*, *Digitaria*, *Echinochloa*, *Fimbristylis*, *Ipomoea*, *Ischaemum*, *Leptochloa*, *Papaver*, *Setaria* or *Xanthium*.

In some embodiments, the combination of compound (I) or agriculturally acceptable ester or salt thereof and a protoporphyrinogen oxidase inhibitor are used to control, including but not limited to oilseed rape (*Brassica napus* L.), large crabgrass (*Digitaria sanguinalis*), barnyardgrass (*Echinochloa crus-galli*), jungle rice (*Echinochloa colona*), early watergrass (*Echinochloa oryzoides*), ivyleaf morningglory (*Ipomoea hederacea*), Chinese sprangletop (*Leptochloa chinensis*), smallflower umbrella sedge (*Cyperus difformis*), yellow nutsedge (*Cyperus esculentus*), rice flatsedge (*Cyperus iria*), purple nutsedge (*Cyperus rotundus*), globe fringerush (*Fimbristylis miliacea*), Japanese bulrush (*Schoenoplectus juncooides*), sea clubrush (*Bolboschoenus maritimus* or *Schoenoplectus maritimus*), broadleaf signalgrass (*Brachiaria platyphylla* or *Urochloa platyphylla*), saramollagrass (*Ischaemum rugosum* Salis.), green foxtail (*Setaria viridis*), common poppy (*Papaver rhoeas*), or common cocklebur (*Xanthium strumarium*).

The compounds of formula I or agriculturally acceptable salt or ester thereof may be used to control herbicide resistant or tolerant weeds. The methods employing the combination of a compound of formula I or agriculturally acceptable salt or ester thereof and the compositions described herein may also be employed to control herbicide resistant or tolerant weeds. Exemplary resistant or tolerant weeds include, but are not limited to, biotypes resistant or tolerant to acetolactate synthase (ALS) or acetohydroxy acid synthase (AHAS) inhibitors (*e.g.*, imidazolinones, sulfonylureas, pyrimidinylthiobenzoates, triazolopyrimidines, sulfonylaminocarbonyltriazolinones), photosystem II inhibitors (*e.g.*, phenylcarbamates, pyridazinones, triazines, triazinones, uracils, amides, ureas, benzothiadiazinones, nitriles, phenylpyridazines), acetyl CoA carboxylase (ACCase) inhibitors (*e.g.*, aryloxyphenoxypropionates, cyclohexanediones, phenylpyrazolines), synthetic auxins (*e.g.*, benzoic acids, phenoxy-carboxylic acids, pyridine carboxylic acids, quinoline carboxylic acids), auxin transport inhibitors (*e.g.*, phthalamates, semicarbazones), photosystem I inhibitors (*e.g.*, bipyridyliums), 5-enolpyruvylshikimate-3-phosphate (EPSP) synthase inhibitors (*e.g.*, glyphosate), glutamine synthetase inhibitors (*e.g.*, glufosinate, bialafos), microtubule assembly inhibitors (*e.g.*, benzamides, benzoic acids, dinitroanilines, phosphoramidates, pyridines), mitosis inhibitors (*e.g.*, carbamates), very long chain fatty acid (VLCFA) inhibitors (*e.g.*, acetamides, chloroacetamides, oxyacetamides, tetrazolinones), fatty acid and lipid synthesis inhibitors (*e.g.*, phosphorodithioates, thiocarbamates, benzofuranes, chlorocarbonic acids), protoporphyrinogen oxidase (PPO) inhibitors (*e.g.*, diphenylethers, N-phenylphthalimides, oxadiazoles, oxazolidinediones, phenylpyrazoles, pyrimidindiones, thiadiazoles, triazolinones), carotenoid biosynthesis inhibitors (*e.g.*, clomazone, amitrole, aclonifen), phytoene desaturase (PDS) inhibitors (*e.g.*, amides, anilidex, furanones, phenoxybutan-amides, pyridiazinones, pyridines), 4-hydroxyphenyl-pyruvate-dioxygenase (HPPD) inhibitors (*e.g.*, callistemones, isoxazoles, pyrazoles, triketones), cellulose biosynthesis inhibitors (*e.g.*, nitriles, benzamides, quinclorac, triazolocarboxamides), herbicides with multiple modes-of-action such as quinclorac, and unclassified herbicides such as arylaminopropionic acids, difenzoquat, endothall, and organoarsenicals. Exemplary resistant or tolerant weeds include, but are not limited to, biotypes with resistance or tolerance to multiple herbicides, biotypes with resistance or tolerance to multiple chemical classes, biotypes with resistance or tolerance to multiple

herbicide modes-of-action, and biotypes with multiple resistance or tolerance mechanisms (*e.g.*, target site resistance or metabolic resistance).

In certain embodiments of the compositions and methods described herein, the compound of formula (I) or salt or ester thereof is used in combination with acifluorfen.

5 With regard to the compositions, in some embodiments, the weight ratio of the compound of formula (I) or salt or ester thereof to acifluorfen or an agriculturally acceptable salt thereof is within the range of from about 1:210 to about 9:1. In certain embodiments, the weight ratio of the compound of formula (I) or salt or ester thereof to acifluorfen is within the range of from about 1:191 to about 2:1. In certain embodiments, the compositions comprise the
10 compound of formula (I) or its benzyl or n-butyl ester and acifluorfen sodium. In one embodiment, the composition comprises the benzyl ester of the compound of formula (I) and acifluorfen, wherein the weight ratio of the benzyl ester of the compound of formula (I) to acifluorfen is about 1:17.5 to about 1:35. With respect to the methods, in certain
15 embodiments, the methods comprise contacting the undesirable vegetation or locus thereof or applying to the soil or water to prevent the emergence or growth of vegetation a composition described herein. In some embodiments, the composition is applied at an application rate of from about 37 grams active ingredient per hectare (g ai/ha) to about 720 g ai/ha based on the total amount of active ingredients in the composition. In certain embodiments, the composition is applied at an application rate of from about 38 g ai/ha to about 470 g ai/ha
20 based on the total amount of active ingredients in the composition. In some embodiments, the methods comprise contacting the undesirable vegetation or locus thereof or applying to the soil or water to prevent the emergence or growth of vegetation with a compound of formula (I) or salt or ester thereof and acifluorfen, *e.g.*, sequentially or simultaneously. In some embodiments, the acifluorfen is applied at a rate from about 35 g ai/ha to about 420 g
25 ai/ha and the compound of formula (I) or salt or ester thereof is applied at a rate from about 2 grams acid equivalent per hectare (g ae/ha) to about 300 g ae/ha. In certain embodiments, the methods utilize the compound of formula (I) or its benzyl or n-butyl ester and acifluorfen sodium. In one embodiment, the methods utilize the benzyl ester of the compound of formula (I) and acifluorfen, wherein the benzyl ester of the compound of formula (I) is
30 applied at a rate of from about 8 g acid equivalent per hectare (g ae/ha), and acifluorfen is applied at a rate of about 140 g ai/ha to about 280 g ai/ha. In certain embodiments, the

methods and compositions utilizing the compound of formula (I) or salt or ester thereof in combination with acifluorfen are used to control ECHCO.

In certain embodiments of the compositions and methods described herein, the compound of formula (I) or salt or ester thereof is used in combination with azafeniden. In some embodiments, the weight ratio of the compound of formula (I) or salt or ester thereof to azafeniden is within the range of from about 1:220 to about 10:1. In certain embodiments, the weight ratio of the compound of formula (I) or salt or ester thereof to azafeniden is within the range of from about 1:109 to about 2:1. In certain embodiments, the compositions comprise the compound of formula (I) or its benzyl or n-butyl ester and azafeniden. With respect to the methods, in certain embodiments, the methods comprise contacting the undesirable vegetation or locus thereof or applying to the soil or water to prevent the emergence or growth of vegetation a composition described herein. In some embodiments, the composition is applied at an application rate of from about 32 grams active ingredient per hectare (g ai/ha) to about 540 g ai/ha based on the total amount of active ingredients in the composition. In certain embodiments, the composition is applied at an application rate of from about 33 g ai/ha to about 290 g ai/ha based on the total amount of active ingredients in the composition. In some embodiments, the methods comprise contacting the undesirable vegetation or locus thereof or applying to the soil or water to prevent the emergence or growth of vegetation with a compound of formula (I) or salt or ester thereof and azafeniden, *e.g.*, sequentially or simultaneously. In some embodiments, the azafeniden is applied at a rate from about 30 g ai/ha to about 240 g ai/ha and the compound of formula (I) or salt or ester thereof is applied at a rate from about 2 grams acid equivalent per hectare (g ae/ha) to about 300 g ae/ha. In certain embodiments, the methods utilize the compound of formula (I) or its benzyl or n-butyl ester and azafeniden.

In certain embodiments of the compositions and methods described herein, the compound of formula (I) or salt or ester thereof is used in combination with bifenox. With regard to the compositions, in some embodiments, the weight ratio of the compound of formula (I) or salt or ester thereof to bifenox is within the range of from about 1:500 to about 3:1. In certain embodiments, the weight ratio of the compound of formula (I) or salt or ester thereof to bifenox is within the range of from about 1:455 to about 1:2.5. In certain embodiments, the compositions comprise the compound of formula (I) or its benzyl or n-butyl ester and bifenox. In one embodiment, the composition comprises the benzyl ester of

the compound of formula (I) and bifenox, wherein the weight ratio of the benzyl ester of the compound of formula (I) to bifenox is about 1:5 to about 1:90. In one embodiment, the composition comprises the benzyl ester of the compound of formula (I) and bifenox, wherein the weight ratio of the benzyl ester of the compound of formula (I) to bifenox is about 1:9 to about 1:90. With respect to the methods, in certain embodiments, the methods comprise contacting the undesirable vegetation or locus thereof or applying to the soil or water to prevent the emergence or growth of vegetation a composition described herein. In some embodiments, the composition is applied at an application rate of from about 127 grams active ingredient per hectare (g ai/ha) to about 1300 g ai/ha based on the total amount of active ingredients in the composition. In certain embodiments, the composition is applied at an application rate of from about 128 g ai/ha to about 1050 g ai/ha based on the total amount of active ingredients in the composition. In some embodiments, the methods comprise contacting the undesirable vegetation or locus thereof or applying to the soil or water to prevent the emergence or growth of vegetation with a compound of formula (I) or salt or ester thereof and bifenox, *e.g.*, sequentially or simultaneously. In some embodiments, the bifenox is applied at a rate from about 125 g ai/ha to about 1000 g ai/ha and the compound of formula (I) or salt or ester thereof is applied at a rate from about 2 grams acid equivalent per hectare (g ae/ha) to about 300 g ae/ha. In certain embodiments, the methods utilize the compound of formula (I) or its benzyl or n-butyl ester and bifenox. In one embodiment, the methods utilize the benzyl ester of the compound of formula (I) and bifenox, wherein the benzyl ester of the compound of formula (I) is applied at a rate of from about 8 g acid equivalent per hectare (g ae/ha) to about 32 g ae/ha, and bifenox is applied at a rate of about 180 g ai/ha to about 720 g ai/ha. In one embodiment, the methods utilize the benzyl ester of the compound of formula (I) and bifenox, wherein the benzyl ester of the compound of formula (I) is applied at a rate of from about 96 g acid equivalent per hectare (g ae/ha), and bifenox is applied at a rate of about 180 g ai/ha to about 720 g ai/ha. In certain embodiments, the methods and compositions utilizing the compound of formula (I) or salt or ester thereof in combination with bifenox are used to control BRSNW, CYPIR, ECHOR, IPOHE, PAPRH, or SCPMA.

In certain embodiments of the compositions and methods described herein, the compound of formula (I) or salt or ester thereof is used in combination with carfentrazone-ethyl or salt or ester thereof. With regard to the compositions, in some embodiments, the

weight ratio of the compound of formula (I) or salt or ester thereof to carfentrazone-ethyl or carboxylic acid or carboxylate salt thereof or other ester is within the range of from about 1:60 to about 86:1. In certain embodiments, the weight ratio of the compound of formula (I) or salt or ester thereof to carfentrazone-ethyl or carboxylic acid or carboxylate salt thereof or other ester is within the range of from about 1:49 to about 49:1. In certain embodiments, the weight ratio of the compound of formula (I) or salt or ester thereof to carfentrazone-ethyl or carboxylic acid or carboxylate salt thereof or other ester is within the range of from about 1:0.2 to about 1:10. In certain embodiments, the weight ratio of the compound of formula (I) or salt or ester thereof to carfentrazone-ethyl or carboxylic acid or carboxylate salt thereof or other ester is within the range of from about 1:0.4 to about 1:5. In certain embodiments, the compositions provided herein comprise the compound of formula (I) or its benzyl or *n*-butyl ester and carfentrazone-ethyl. In one embodiment, the composition comprises the compound of formula (I) and carfentrazone-ethyl, wherein the weight ratio of the compound of formula (I) to carfentrazone-ethyl is about 1:0.4 to about 1:5. In one embodiment, the composition comprises the benzyl ester of the compound of formula (I) and carfentrazone-ethyl, wherein the weight ratio of the benzyl ester of the compound of formula (I) to carfentrazone-ethyl is about 1:0.4 to about 1:3.2. In one embodiment, the composition comprises the *n*-butyl ester of the compound of formula (I) and carfentrazone-ethyl, wherein the weight ratio of the *n*-butyl ester of the compound of formula (I) to carfentrazone-ethyl is about 1:1.75. With respect to the methods, in certain embodiments, the methods comprise contacting the undesirable vegetation or locus thereof or applying to the soil or water to prevent the emergence or growth of vegetation a composition described herein. The application rate will depend upon the particular type of weed to be controlled, the degree of control required, and the timing and method of application. In some embodiments, the composition is applied at an application rate of from about 5.5 grams active ingredient per hectare (g ai/ha) to about 420 g ai/ha based on the total amount of active ingredients in the composition. In certain embodiments, the composition is applied at an application rate of from about 8 g ai/ha to about 257 g ai/ha based on the total amount of active ingredients in the composition. In some embodiments, the methods comprise contacting the undesirable vegetation or locus thereof or applying to the soil or water to prevent the emergence or growth of vegetation with a compound of formula (I) or salt or ester thereof and carfentrazone-ethyl or carboxylic acid or carboxylate salt thereof or other ester, *e.g.*, sequentially or simultaneously. In some

embodiments, the carfentrazone-ethyl or carboxylic acid or carboxylate salt thereof or other ester is applied at a rate from about 3.5 g ai/ha to about 120 g ai/ha and the compound of formula (I) or salt or ester thereof is applied at a rate from about 2 grams acid equivalent per hectare (g ae/ha) to about 300 g ae/ha. In some embodiments, the carfentrazone-ethyl or carboxylic acid or carboxylate salt thereof or other ester is applied at a rate from about 3 g ai/ha to about 430 g ai/ha and the compound of formula (I) or salt or ester thereof is applied at a rate from about 2 g acid equivalent per hectare (g ae/ha) to about 84 g ae/ha. In some embodiments, the carfentrazone-ethyl or carboxylic acid or carboxylate salt thereof or other ester is applied at a rate from about 7 g ai/ha to about 215 g ai/ha and the compound of formula (I) or salt or ester thereof is applied at a rate from about 4.38 g acid equivalent per hectare (g ae/ha) to about 42.4 g ae/ha. In certain embodiments, the methods utilize the compound of formula (I), or its benzyl or *n*-butyl ester and carfentrazone-ethyl. In one embodiment, the methods utilize the compound of formula (I) and carfentrazone-ethyl, wherein the compound of formula (I) is applied at a rate of from about 4.38 g acid equivalent per hectare (g ae/ha) to about 42.4 g ae/ha, and carfentrazone-ethyl is applied at a rate of about 7 g ai/ha to about 215 g ai/ha. In one embodiment, the methods utilize the benzyl ester of the compound of formula (I) and carfentrazone-ethyl, wherein the benzyl ester of the compound of formula (I) is applied at a rate of from about 8.75 g acid equivalent per hectare (g ae/ha) to about 35 g ae/ha, and carfentrazone-ethyl is applied at a rate of about 3.5 g ai/ha to about 28 g ai/ha. In one embodiment, the methods utilize the *n*-butyl ester of the compound of formula (I) and carfentrazone-ethyl, wherein the *n*-butyl ester of the compound of formula (I) is applied at a rate of about 16 g acid equivalent per hectare (g ae/ha), and carfentrazone-ethyl is applied at a rate of about 28 g ai/ha. In certain embodiments, the methods and compositions utilizing the compound of formula (I) or salt or ester thereof in combination with carfentrazone-ethyl or carboxylic acid or carboxylate salt thereof or other ester are used to control CYPES, CYPIR, DIGSA, ECHCG, ECHOR or LEFCH.

In certain embodiments of the compositions and methods described herein, the compound of formula (I) or salt or ester thereof is used in combination with flumioxazin. With regard to the compositions, in some embodiments, the weight ratio of the compound of formula (I) or salt or ester thereof to flumioxazin is within the range of from about 1:300 to about 17:1. In certain embodiments, the weight ratio of the compound of formula (I) or salt or ester thereof to flumioxazin is within the range of from about 1:272 to about 3:1. In certain

embodiments, the compositions comprise the compound of formula (I) or its benzyl or n-butyl ester and flumioxazin. In one embodiment, the composition comprises the benzyl ester of the compound of formula (I) and flumioxazin, wherein the weight ratio of the benzyl ester of the compound of formula (I) to flumioxazin is about 1:9 to about 1:35. In one embodiment,

5 the composition comprises the benzyl ester of the compound of formula (I) and flumioxazin, wherein the weight ratio of the benzyl ester of the compound of formula (I) to flumioxazin is about 1:3 to about 1:27. With respect to the methods, in certain embodiments, the methods comprise contacting the undesirable vegetation or locus thereof or applying to the soil or water to prevent the emergence or growth of vegetation a composition described herein. In

10 some embodiments, the composition is applied at an application rate of from about 19 grams active ingredient per hectare (g ai/ha) to about 900 g ai/ha based on the total amount of active ingredients in the composition. In certain embodiments, the composition is applied at an application rate of from about 20 g ai/ha to about 650 g ai/ha based on the total amount of active ingredients in the composition. In some embodiments, the methods comprise

15 contacting the undesirable vegetation or locus thereof or applying to the soil or water to prevent the emergence or growth of vegetation with a compound of formula (I) or salt or ester thereof and flumioxazin, *e.g.*, sequentially or simultaneously. In some embodiments, the flumioxazin is applied at a rate from about 17.5 g ai/ha to about 600 g ai/ha and the compound of formula (I) or salt or ester thereof is applied at a rate from about 2 grams acid

20 equivalent per hectare (g ae/ha) to about 300 g ae/ha. In certain embodiments, the methods utilize the compound of formula (I) or its benzyl or n-butyl ester and flumioxazin. In one embodiment, the methods utilize the benzyl ester of the compound of formula (I) and flumioxazin, wherein the benzyl ester of the compound of formula (I) is applied at a rate of from about 8 g acid equivalent per hectare (g ae/ha) to about 16 g ai/ha, and flumioxazin is

25 applied at a rate of about 140 g ai/ha to about 280 g ai/ha. In one embodiment, the methods utilize the benzyl ester of the compound of formula (I) and flumioxazin, wherein the benzyl ester of the compound of formula (I) is applied at a rate of from about 16 g acid equivalent per hectare (g ae/ha) to about 32 g ae/ha, and bifenox is applied at a rate of about 105 g ai/ha to about 420 g ai/ha. In certain embodiments, the methods and compositions utilizing the

30 compound of formula (I) or salt or ester thereof in combination with flumioxazin are used to control ECHCO, IPOHE, SETVI, or XANST.

In certain embodiments of the compositions and methods described herein, the compound of formula (I) or salt or ester thereof is used in combination with ipfencarbazone. With regard to the compositions, in some embodiments, the weight ratio of the compound of formula (I) or salt or ester thereof to ipfencarbazone is within the range of from about 1:250 to about 5:1. In certain embodiments, the weight ratio of the compound of formula (I) or salt or ester thereof to ipfencarbazone is within the range of from about 1:227 to about 1:1. In certain embodiments, the compositions comprise the compound of formula (I) or its benzyl or n-butyl ester and ipfencarbazone. With respect to the methods, in certain embodiments, the methods comprise contacting the undesirable vegetation or locus thereof or applying to the soil or water to prevent the emergence or growth of vegetation a composition described herein. In some embodiments, the composition is applied at an application rate of from about 65 grams active ingredient per hectare (g ai/ha) to about 800 g ai/ha based on the total amount of active ingredients in the composition. In certain embodiments, the composition is applied at an application rate of from about 66 g ai/ha to about 550 g ai/ha based on the total amount of active ingredients in the composition. In some embodiments, the methods comprise contacting the undesirable vegetation or locus thereof or applying to the soil or water to prevent the emergence or growth of vegetation with a compound of formula (I) or salt or ester thereof and ipfencarbazone, *e.g.*, sequentially or simultaneously. In some embodiments, the ipfencarbazone is applied at a rate from about 62.5 g ai/ha to about 500 g ai/ha and the compound of formula (I) or salt or ester thereof is applied at a rate from about 2 grams acid equivalent per hectare (g ae/ha) to about 300 g ae/ha. In certain embodiments, the methods utilize the compound of formula (I) or its benzyl or n-butyl ester and ipfencarbazone.

In certain embodiments of the compositions and methods described herein, the compound of formula (I) or salt or ester thereof is used in combination with oxadiargyl. With regard to the compositions, in some embodiments, the weight ratio of the compound of formula (I) or salt or ester thereof to oxadiargyl is within the range of from about 1:1100 to about 6:1. In certain embodiments, the weight ratio of the compound of formula (I) or salt or ester thereof to oxadiargyl is within the range of from about 1:509 to about 4:1. In certain embodiments, the compositions comprise the compound of formula (I) or its benzyl or n-butyl ester and oxadiargyl. In one embodiment, the composition comprises the compound of formula (I) and oxadiargyl, wherein the weight ratio of the compound of formula (I) to

oxadiargyl is about 1:3 to about 1:11. In one embodiment, the composition comprises the benzyl ester of the compound of formula (I) and oxadiargyl, wherein the weight ratio of the benzyl ester of the compound of formula (I) to oxadiargyl is about 1: 5 to about 1:11. With respect to the methods, in certain embodiments, the methods comprise contacting the

5 undesirable vegetation or locus thereof or applying to the soil or water to prevent the emergence or growth of vegetation a composition described herein. In some embodiments, the composition is applied at an application rate of from about 15 grams active ingredient per hectare (g ai/ha) to about 2500 g ai/ha based on the total amount of active ingredients in the composition. In certain embodiments, the composition is applied at an application rate of

10 from about 16 g ai/ha to about 1170 g ai/ha based on the total amount of active ingredients in the composition. In some embodiments, the methods comprise contacting the undesirable vegetation or locus thereof or applying to the soil or water to prevent the emergence or growth of vegetation with a compound of formula (I) or salt or ester thereof and oxadiargyl, e.g., sequentially or simultaneously. In some embodiments, the oxadiargyl is applied at a rate

15 from about 50 g ai/ha to about 2200 g ai/ha and the compound of formula (I) of salt or ester thereof is applied at a rate from about 2 grams acid equivalent per hectare (g ae/ha) to about 300 g ae/ha. In certain embodiments, the methods utilize the compound of formula (I) or its benzyl or n-butyl ester and oxadiargyl. In one embodiment, the methods utilize the compound of formula (I) and oxadiargyl, wherein of the compound of formula (I) is applied

20 at a rate of from about 4.38 g acid equivalent per hectare (g ae/ha) to about 17.5 g ae/ha, and oxadiargyl is applied at a rate of about 50 g ai/ha. In one embodiment, the methods utilize the benzyl ester of the compound of formula (I) and oxadiargyl, wherein the benzyl ester of the compound of formula (I) is applied at a rate of from about 4.38 g acid equivalent per hectare (g ae/ha) to about 8.75 g ae/ha, and oxadiargyl is applied at a rate of about 50 g ai/ha.

25 In certain embodiments, the methods and compositions utilizing the compound of formula (I) or salt or ester thereof in combination with oxadiargyl are used to control IPOHE.

In certain embodiments of the compositions and methods described herein, the compound of formula (I) or salt or ester thereof is used in combination with oxadiazon. With regard to the compositions, in some embodiments, the weight ratio of the compound of

30 formula (I) or salt or ester thereof to oxadiazon is within the range of from about 1:2240 to about 10:1. In certain embodiments, the weight ratio of the compound of formula (I) or salt or ester thereof to oxadiazon is within the range of from about 1:57 to about 2:1. In certain

embodiments, the weight ratio of the compound of formula (I) or salt or ester thereof to oxadiazon is within the range of from about 1:0.9 to about 1:57. In certain embodiments, the weight ratio of the compound of formula (I) or salt or ester thereof to oxadiazon is within the range of from about 1:1.8 to about 1:28.5. In certain embodiments, the compositions provided

5 herein comprise the compound of formula (I) or its benzyl or *n*-butyl ester and oxadiazon. In one embodiment, the composition comprises the compound of formula (I) and oxadiazon, wherein the weight ratio of the compound of formula (I) to oxadiazon is about 1:1.8 to about 1:14.3. In one embodiment, the composition comprises the benzyl ester of the compound of formula (I) and oxadiazon, wherein the weight ratio of the benzyl ester of the compound of

10 formula (I) to oxadiazon is about 1:1.8 to about 1:70. In one embodiment, the composition comprises the *n*-butyl ester of the compound of formula (I) and oxadiazon, wherein the weight ratio of the *n*-butyl ester of the compound of formula (I) to oxadiazon is about 1:3.6 to about 1:14.3. With respect to the methods, in certain embodiments, the methods comprise contacting the undesirable vegetation or locus thereof or applying to the soil or water to

15 prevent the emergence or growth of vegetation a composition described herein. In some embodiments, the composition is applied at an application rate of from about 33 grams active ingredient per hectare (g ai/ha) to about 4780 g ai/ha based on the total amount of active ingredients in the composition. In certain embodiments, the composition is applied at an application rate of from about 35 g ai/ha to about 320 g ai/ha based on the total amount of

20 active ingredients in the composition. In some embodiments, the methods comprise contacting the undesirable vegetation or locus thereof or applying to the soil or water to prevent the emergence or growth of vegetation with a compound of formula (I) or salt or ester thereof and oxadiazon, *e.g.*, sequentially or simultaneously. In some embodiments, the oxadiazon is applied at a rate from about 31 g ai/ha to about 4480 g ai/ha and the compound

25 of formula (I) of salt or ester thereof is applied at a rate from about 2 grams acid equivalent per hectare (g ae/ha) to about 300 g ae/ha. In some embodiments, the oxadiazon is applied at a rate from about 15 g ai/ha to about 500 g ai/ha and the compound of formula (I) of salt or ester thereof is applied at a rate from about 2 g acid equivalent per hectare (g ae/ha) to about 140 g ae/ha. In some embodiments, the oxadiazon is applied at a rate from about 31 g ai/ha

30 to about 250 g ai/ha and the compound of formula (I) of salt or ester thereof is applied at a rate from about 4.38 g acid equivalent per hectare (g ae/ha) to about 70 g ae/ha. In certain embodiments, the methods utilize the compound of formula (I), or its benzyl or *n*-butyl ester

and oxadiazon. In one embodiment, the methods utilize the compound of formula (I) and oxadiazon, wherein the compound of formula (I) is applied at a rate of from about 8.75 g acid equivalent per hectare (g ae/ha) to about 35 g ae/ha, and oxadiazon is applied at a rate of about 31 g ai/ha to about 250 g ai/ha. In one embodiment, the methods utilize the benzyl ester of the compound of formula (I) and oxadiazon, wherein the benzyl ester of the compound of formula (I) is applied at a rate of from about 4.38 g acid equivalent per hectare (g ae/ha) to about 17.5 g ae/ha, and oxadiazon is applied at a rate of about 31 g ai/ha to about 250 g ai/ha. In one embodiment, the methods utilize the *n*-butyl ester of the compound of formula (I) and oxadiazon, wherein the *n*-butyl ester of the compound of formula (I) is applied at a rate of about 17.5 g acid equivalent per hectare (g ae/ha) to about 70 g ae/ha, and oxadiazon is applied at a rate of about 250 g ai/ha. In certain embodiments, the methods and compositions utilizing the compound of formula (I) or salt or ester thereof in combination with oxadiazon are used to control ECHOR or IPOHE.

In certain embodiments of the compositions and methods described herein, the compound of formula (I) or salt or ester thereof is used in combination with oxyfluorfen. With regard to the compositions, in some embodiments, the weight ratio of the compound of formula (I) or salt or ester thereof to oxyfluorfen is within the range of from about 1:1120 to about 3:1. In certain embodiments, the weight ratio of the compound of formula (I) or salt or ester thereof to oxyfluorfen is within the range of from 1:82 to about 1:2. In certain embodiments, the weight ratio of the compound of formula (I) or salt or ester thereof to oxyfluorfen is within the range of from about 1:2 to about 1:164. In certain embodiments, the weight ratio of the compound of formula (I) or salt or ester thereof to oxyfluorfen is within the range of from about 1:4.2 to about 1:82. In certain embodiments, the compositions provided herein comprise the compound of formula (I) or its benzyl or *n*-butyl ester and oxyfluorfen. In one embodiment, the composition comprises the compound of formula (I) and oxyfluorfen, wherein the weight ratio of the compound of formula (I) to oxyfluorfen is about 1:4.2 to about 1:68. In one embodiment, the composition comprises the benzyl ester of the compound of formula (I) and oxyfluorfen, wherein the weight ratio of the benzyl ester of the compound of formula (I) to oxyfluorfen is about 1:5 to about 1:82. With respect to the methods, in certain embodiments, the methods comprise contacting the undesirable vegetation or locus thereof or applying to the soil or water to prevent the emergence or growth of vegetation a composition described herein. In some embodiments, the composition

is applied at an application rate of from about 92 grams active ingredient per hectare (g ai/ha) to about 2540 g ai/ha based on the total amount of active ingredients in the composition. In certain embodiments, the composition is applied at an application rate of from about 94 g ai/ha to about 402 g ai/ha based on the total amount of active ingredients in the composition.

5 In some embodiments, the methods comprise contacting the undesirable vegetation or locus thereof or applying to the soil or water to prevent the emergence or growth of vegetation with a compound of formula (I) or salt or ester thereof and oxyfluorfen, *e.g.*, sequentially or simultaneously. In some embodiments, the oxyfluorfen is applied at a rate from about 100 g ai/ha to about 2240 g ai/ha and the compound of formula (I) or salt or ester thereof is applied
10 at a rate from about 2 grams acid equivalent per hectare (g ae/ha) to about 300 g ae/ha. In some embodiments, the oxyfluorfen is applied at a rate from about 45 g ai/ha to about 720 g ai/ha and the compound of formula (I) or salt or ester thereof is applied at a rate from about 2 g acid equivalent per hectare (g ae/ha) to about 85 g ae/ha. In some embodiments, the oxyfluorfen is applied at a rate from about 90 g ai/ha to about 360 g ai/ha and the compound
15 of formula (I) or salt or ester thereof is applied at a rate from about 4.38 g acid equivalent per hectare (g ae/ha) to about 42.4 g ae/ha. In certain embodiments, the methods utilize the compound of formula (I), or its benzyl or *n*-butyl ester and oxyfluorfen. In one embodiment, the methods utilize the compound of formula (I) and oxyfluorfen, wherein the compound of formula (I) is applied at a rate of from about 4.38 g acid equivalent per hectare (g ae/ha) to
20 about 64 g ae/ha, and oxyfluorfen is applied at a rate of about 90 g ai/ha to about 360 g ai/ha. In one embodiment, the methods utilize the benzyl ester of the compound of formula (I) and oxyfluorfen, wherein the benzyl ester of the compound of formula (I) is applied at a rate of from about 4.38 g acid equivalent per hectare (g ae/ha) to about 32 g ae/ha, and oxyfluorfen is applied at a rate of about 90 g ai/ha to about 360 g ai/ha. In certain embodiments, the
25 methods and compositions utilizing the compound of formula (I) or salt or ester thereof in combination with oxyfluorfen are used to control CYPIR, CYPRO, DIGSA, ECHCG, ECHCO, IPOHE, or SCPJU.

In certain embodiments of the compositions and methods described herein, the compound of formula (I) or salt or ester thereof is used in combination with pentoxazone or
30 salt thereof. With regard to the compositions, in some embodiments, the weight ratio of the compound of formula (I) or salt or ester thereof to pentoxazone is within the range of from about 1:225 to about 6:1. In certain embodiments, the weight ratio of the compound of

formula (I) or salt or ester thereof to pentoxazone is within the range of from 1:26 to about 1:2. In certain embodiments, the weight ratio of the compound of formula (I) or salt or ester thereof to pentoxazone is within the range of from about 1:0.8 to about 1:52. In certain embodiments, the weight ratio of the compound of formula (I) or salt or ester thereof to pentoxazone is within the range of from about 1:1.6 to about 1:26. In certain embodiments, the compositions provided herein comprise the compound of formula (I) or its benzyl ester and pentoxazone. In one embodiment, the composition comprises the compound of formula (I) and pentoxazone, wherein the weight ratio of the compound of formula (I) to pentoxazone is about 1:1.6 to about 1:13. In one embodiment, the composition comprises the benzyl ester of the compound of formula (I) and pentoxazone, wherein the weight ratio of the benzyl ester of the compound of formula (I) to pentoxazone is about 1:6 to about 1:26. With respect to the methods, in certain embodiments, the methods comprise contacting the undesirable vegetation or locus thereof or applying to the soil or water to prevent the emergence or growth of vegetation a composition described herein. In some embodiments, the composition is applied at an application rate of from about 58 grams active ingredient per hectare (g ai/ha) to about 750 g ai/ha based on the total amount of active ingredients in the composition. In certain embodiments, the composition is applied at an application rate of from about 61 g ai/ha to about 148 g ai/ha based on the total amount of active ingredients in the composition. In some embodiments, the methods comprise contacting the undesirable vegetation or locus thereof or applying to the soil or water to prevent the emergence or growth of vegetation with a compound of formula (I) or salt or ester thereof and pentoxazone, *e.g.*, sequentially or simultaneously. In some embodiments, the pentoxazone is applied at a rate from about 56 g ai/ha to about 450 g ai/ha and the compound of formula (I) or salt or ester thereof is applied at a rate from about 2 grams acid equivalent per hectare (g ae/ha) to about 300 g ae/ha. In some embodiments, the pentoxazone is applied at a rate from about 25 g ai/ha to about 230 g ai/ha and the compound of formula (I) or salt or ester thereof is applied at a rate from about 2 g acid equivalent per hectare (g ae/ha) to about 70 g ae/ha. In some embodiments, the pentoxazone is applied at a rate from about 56 g ai/ha to about 112 g ai/ha and the compound of formula (I) or salt or ester thereof is applied at a rate from about 4.38 g acid equivalent per hectare (g ae/ha) to about 35 g ae/ha. In certain embodiments, the methods utilize the compound of formula (I), or its benzyl ester and pentoxazone. In one embodiment, the methods utilize the compound of formula (I) and pentoxazone, wherein the compound of

formula (I) is applied at a rate of from about 8.75 g acid equivalent per hectare (g ae/ha) to about 35 g ae/ha, and pentoxazone is applied at a rate of about 56 g ai/ha to about 113 g ai/ha. In one embodiment, the methods utilize the benzyl ester of the compound of formula (I) and pentoxazone, wherein the benzyl ester of the compound of formula (I) is applied at a rate of
5 from about 4.38 g acid equivalent per hectare (g ae/ha) to about 8.75 g ae/ha, and pentoxazone is applied at a rate of about 56 g ai/ha to about 113 g ai/ha. In certain embodiments, the methods and compositions utilizing the compound of formula (I) or salt or ester thereof in combination with pentoxazone are used to control CYPPO, ECHCG, FIMMI or SCPJU.

10 In certain embodiments of the compositions and methods described herein, the compound of formula (I) or salt or ester thereof is used in combination with pyraclonil or salt thereof. With regard to the compositions, in some embodiments, the weight ratio of the compound of formula (I) or salt or ester thereof to pyraclonil or salt thereof is within the range of from about 1:180 to about 9:1. In certain embodiments, the weight ratio of the
15 compound of formula (I) or salt or ester thereof to pyraclonil or salt thereof is within the range of from about 1:62 to about 1:1. In certain embodiments, the weight ratio of the compound of formula (I) or salt or ester thereof to pyraclonil or salt thereof is within the range of from about 1:4.4 to about 2:1. In certain embodiments, the weight ratio of the compound of formula (I) or salt or ester thereof to pyraclonil or salt thereof is within the
20 range of from about 1:34 to about 2:1. In certain embodiments, the compositions provided herein comprise the compound of formula (I) or its benzyl or *n*-butyl ester and pyraclonil. In one embodiment, the composition comprises the compound of formula (I) and pyraclonil, wherein the weight ratio of the compound of formula (I) to pyraclonil is about 1:31 to about 1:3. In one embodiment, the composition comprises the benzyl ester of the compound of
25 formula (I) and pyraclonil, wherein the weight ratio of the benzyl ester of the compound of formula (I) to pyraclonil is about 1:2 to about 1:34. With respect to the methods, in certain embodiments, the methods comprise contacting the undesirable vegetation or locus thereof or applying to the soil or water to prevent the emergence or growth of vegetation a composition described herein. In some embodiments, the composition is applied at an application rate of
30 from about 36 grams active ingredient per hectare (g ai/ha) to about 660 g ai/ha based on the total amount of active ingredients in the composition. In certain embodiments, the composition is applied at an application rate of from about 41 g ai/ha to about 305 g ai/ha

based on the total amount of active ingredients in the composition. In some embodiments, the methods comprise contacting the undesirable vegetation or locus thereof or applying to the soil or water to prevent the emergence or growth of vegetation with a compound of formula (I) or salt or ester thereof and pyraclonil or salt thereof, *e.g.*, sequentially or simultaneously. In some embodiments, the pyraclonil is applied at a rate from about 34 g ai/ha to about 360 g ai/ha and the compound of formula (I) or salt or ester thereof is applied at a rate from about 2 grams acid equivalent per hectare (g ae/ha) to about 300 g ae/ha. In some embodiments, the pyraclonil or salt thereof is applied at a rate from about 4 g ai/ha to about 140 g ai/ha and the compound of formula (I) or salt or ester thereof is applied at a rate from about 2 g acid equivalent per hectare (g ae/ha) to about 45 g ae/ha. In some embodiments, the pyraclonil or salt thereof is applied at a rate from about 33.75 g ai/ha to about 270 g ai/ha and the compound of formula (I) or salt or ester thereof is applied at a rate from about 4.38 g acid equivalent per hectare (g ae/ha) to about 35 g ae/ha. In certain embodiments, the methods utilize the compound of formula (I), or its benzyl or *n*-butyl ester and pyraclonil. In one embodiment, the methods utilize the compound of formula (I) and pyraclonil, wherein the compound of formula (I) is applied at a rate of from about 4.38 g acid equivalent per hectare (g ae/ha) to about 35 g ae/ha, and pyraclonil is applied at a rate of about 33.75 g ai/ha to about 270 g ai/ha. In one embodiment, the methods utilize the benzyl ester of the compound of formula (I) and pyraclonil, wherein the benzyl ester of the compound of formula (I) is applied at a rate of from about 4.38 g acid equivalent per hectare (g ae/ha) to about 35 g ae/ha, and pyraclonil is applied at a rate of about 33.75 g ai/ha to about 270 g ai/ha. In certain embodiments, the methods and compositions utilizing the compound of formula (I) or salt or ester thereof in combination with pyraclonil are used to control BRAPP, CYPRO, ECHCG, ECHCO, ECHOR IPOHE or ISCRU.

In certain embodiments of the compositions and methods described herein, the compound of formula (I) or salt or ester thereof is used in combination with pyraflufen-ethyl or carboxylic acid or carboxylate salt thereof or other ester. With regard to the compositions, in some embodiments, the weight ratio of the compound of formula (I) or salt or ester thereof to pyraflufen-ethyl or carboxylic acid or carboxylate salt thereof or other ester is within the range of from about 1:6 to about 150:1. In certain embodiments, the weight ratio of the compound of formula (I) or salt or ester thereof to pyraflufen-ethyl or carboxylic acid or carboxylate salt thereof or other ester is within the range of from about 1:5.5 to about 25:1. In

certain embodiments, the compositions comprise the compound of formula (I) or its benzyl or n-butyl ester and pyraflufen-ethyl. In one embodiment, the composition comprises the benzyl ester of the compound of formula (I) and pyraflufen-ethyl, wherein the weight ratio of the benzyl ester of the compound of formula (I) to pyraflufen-ethyl is about 1:3 to about 1.5:1.

5 With respect to the methods, in certain embodiments, the methods comprise contacting the undesirable vegetation or locus thereof or applying to the soil or water to prevent the emergence or growth of vegetation a composition described herein. In some embodiments, the composition is applied at an application rate of from about 4 grams active ingredient per hectare (g ai/ha) to about 312 g ai/ha based on the total amount of active ingredients in the
10 composition. In certain embodiments, the composition is applied at an application rate of from about 5 g ai/ha to about 62 g ai/ha based on the total amount of active ingredients in the composition. In some embodiments, the methods comprise contacting the undesirable vegetation or locus thereof or applying to the soil or water to prevent the emergence or growth of vegetation with a compound of formula (I) or salt or ester thereof and pyraflufen-ethyl or carboxylic acid or carboxylate salt thereof or other ester, *e.g.*, sequentially or
15 simultaneously. In some embodiments, the pyraflufen-ethyl or carboxylic acid or carboxylate salt thereof or other ester is applied at a rate from about 2 g ai/ha to about 12 g ai/ha and the compound of formula (I) or salt or ester thereof is applied at a rate from about 2 grams acid equivalent per hectare (g ae/ha) to about 300 g ae/ha. In certain embodiments, the
20 methods utilize the compound of formula (I) or its benzyl or n-butyl ester and pyraflufen-ethyl. In one embodiment, the methods utilize the benzyl ester of the compound of formula (I) and pyraflufen-ethyl, wherein the benzyl ester of the compound of formula (I) is applied at a rate of from about 8 g acid equivalent per hectare (g ae/ha) to about 16 g ae/ha, and pyraflufen-ethyl is applied at a rate of about 2.6 g ai/ha to about 10.6 g ai/ha. In certain
25 embodiments, the methods and compositions utilizing the compound of formula (I) or salt or ester thereof in combination with pyraflufen-ethyl are used to control CYP1R or ECHCG.

In certain embodiments of the compositions and methods described herein, the compound of formula (I) or salt or ester thereof is used in combination with saflufenacil or salt thereof. With regard to the compositions, in some embodiments, the weight ratio of the
30 compound of formula (I) or salt or ester thereof to saflufenacil or salt or ester thereof is within the range of from about 1:100 to about 60:1. In certain embodiments, the weight ratio of the compound of formula (I) or salt or ester thereof to saflufenacil or salt or ester thereof is

within the range of from about 1:9 to about 12:1. In certain embodiments, the weight ratio of the compound of formula (I) or salt or ester thereof to saflufenacil or salt thereof is within the range of from about 1:0.1 to about 1:10. In certain embodiments, the weight ratio of the compound of formula (I) or salt or ester thereof to saflufenacil or salt thereof is within the range of from about 1:0.3 to about 1:5. In certain embodiments, the compositions provided herein comprise the compound of formula (I) or its benzyl or *n*-butyl ester and saflufenacil. In one embodiment, the composition comprises the compound of formula (I) and saflufenacil, wherein the weight ratio of the compound of formula (I) to saflufenacil is about 1:0.3 to about 1:5. In one embodiment, the composition comprises the benzyl ester of the compound of formula (I) and saflufenacil, wherein the weight ratio of the benzyl ester of the compound of formula (I) to saflufenacil is about 1:0.3 to about 1:5. In one embodiment, the composition comprises the *n*-butyl ester of the compound of formula (I) and saflufenacil, wherein the weight ratio of the *n*-butyl ester of the compound of formula (I) to saflufenacil is about 1:1.1. With respect to the methods, in certain embodiments, the methods comprise contacting the undesirable vegetation or locus thereof or applying to the soil or water to prevent the emergence or growth of vegetation a composition described herein. In some embodiments, the composition is applied at an application rate of from about 7 grams active ingredient per hectare (g ai/ha) to about 500 g ai/ha based on the total amount of active ingredients in the composition. In certain embodiments, the composition is applied at an application rate of from about 9 g ai/ha to about 110 g ai/ha based on the total amount of active ingredients in the composition. In some embodiments, the methods comprise contacting the undesirable vegetation or locus thereof or applying to the soil or water to prevent the emergence or growth of vegetation with a compound of formula (I) or salt or ester thereof and saflufenacil or salt thereof, *e.g.*, sequentially or simultaneously. In some embodiments, the saflufenacil or salt or ester thereof is applied at a rate from about 5 g ai/ha to about 200 g ai/ha and the compound of formula (I) of salt or ester thereof is applied at a rate from about 2 grams acid equivalent per hectare (g ae/ha) to about 300 g ae/ha. In some embodiments, the saflufenacil or salt thereof is applied at a rate from about 2 g ai/ha to about 80 g ai/ha and the compound of formula (I) of salt or ester thereof is applied at a rate from about 2 g acid equivalent per hectare (g ae/ha) to about 140 g ae/ha. In some embodiments, the saflufenacil or salt thereof is applied at a rate from about 5 g ai/ha to about 40 g ai/ha and the compound of formula (I) of salt or ester thereof is applied at a rate from about 4.38 g acid equivalent per hectare (g

ae/ha) to about 70 g ae/ha. In certain embodiments, the methods utilize the compound of formula (I), or its benzyl or *n*-butyl ester and saflufenacil. In one embodiment, the methods utilize the compound of formula (I) and saflufenacil, wherein the compound of formula (I) is applied at a rate of from about 4.38 g acid equivalent per hectare (g ae/ha) to about 70 g
5 ae/ha, and saflufenacil is applied at a rate of about 5 g ai/ha to about 40 g ai/ha. In one embodiment, the methods utilize the benzyl ester of the compound of formula (I) and saflufenacil, wherein the benzyl ester of the compound of formula (I) is applied at a rate of from about 4.38 g acid equivalent per hectare (g ae/ha) to about 17.5 g ae/ha, and saflufenacil is applied at a rate of about 5 g ai/ha to about 20 g ai/ha. In one embodiment, the methods
10 utilize the *n*-butyl ester of the compound of formula (I) and saflufenacil, wherein the *n*-butyl ester of the compound of formula (I) is applied at a rate of about 35 g acid equivalent per hectare (g ae/ha), and saflufenacil is applied at a rate of about 40 g ai/ha. In certain embodiments, the methods and compositions utilizing the compound of formula (I) or salt or ester thereof in combination with saflufenacil or salt thereof are used to control CYPUR,
15 CYPUR, ECHCG, ECHCO, ECHOR, ISCRU or LEFCH.

The components of the mixtures described herein can be applied either separately or as part of a multipart herbicidal system.

The mixtures described herein can be applied in conjunction with one or more other herbicides to control a wider variety of undesirable vegetation. When used in conjunction
20 with other herbicides, the composition can be formulated with the other herbicide or herbicides, tank mixed with the other herbicide or herbicides or applied sequentially with the other herbicide or herbicides. Some of the herbicides that can be employed in conjunction with the compositions and methods described herein include, but are not limited to: 4-CPA; 4-CPB; 4-CPP; 2,4-D; 2,4-D choline salt, 2,4-D esters and amines, 2,4-DB; 3,4-DA; 3,4-DB;
25 2,4-DEB; 2,4-DEP; 3,4-DP; 2,3,6-TBA; 2,4,5-T; 2,4,5-TB; acetochlor, acifluorfen, acrolein, alachlor, allidochlor, alloxymid, allyl alcohol, alorac, ametrifone, ametryn, amibuzin, amicarbazone, amidosulfuron, aminocyclopyrachlor, aminopyralid, amiprofos-methyl, amitrole, ammonium sulfamate, anilofos, anisuron, asulam, atraton, atrazine, azimsulfuron, aziprotryne, barban, BCPC, beflubutamid, benazolin, bencarbazone, benfluralin, benfuresate,
30 bensulfuron-methyl, bensulide, benthicarb, bentazon-sodium, benzadox, benzfendazole, benzipram, benzobicyclon, benzofenap, benzoxyprop, benzthiazuron, bialaphos, bicyclopyrone, bilanafos, bispyribac-sodium, borax, bromacil, bromobonil, bromobutide,

bromofenoxim, bromoxynil, brompyrazon, butachlor, butafenacil, butamifos, butenachlor,
 buthidazole, buthiuron, butralin, butroxydim, buturon, butylate, cacodylic acid, cafenstrole,
 calcium chlorate, calcium cyanamide, cambendichlor, carbasulam, carbetamide, carboxazole
 chlorprocarb, CDEA, CEPC, chlomethoxyfen, chloramben, chloranocryl, chlorazifop,
 5 chlorazine, chlorbromuron, chlorbufam, chloreturon, chlorfenac, chlorfenprop,
 chlorflurazole, chlorflurenol, chloridazon, chlorimuron, chlornitrofen, chloropon,
 chlorotoluron, chloroxuron, chloroxynil, chlorpropham, chlorsulfuron, chlorthal,
 chlorthiamid, cinidon-ethyl, cinmethylin, cinosulfuron, cisanilide, clethodim, clidinate,
 clodinafop-propargyl, clofop, clomazone, clomeprop, cloprop, cloproxydim, clopyralid,
 10 cloransulam-methyl, CMA, copper sulfate, CPMF, CPPC, credazine, cresol, cumyluron,
 cyanatryn, cyanazine, cycloate, cyclopyrimorate, cyclosulfamuron, cycloxydim, cycluron,
 cyhalofop-butyl, cyperquat, cyprazine, cyprazole, cypromid, daimuron, dalapon, dazomet,
 delachlor, desmedipham, desmetryn, di-allate, dicamba, dichlobenil, dichloralurea,
 dichlormate, dichlorprop, dichlorprop-P, diclofop-methyl, diclosulam, diethamquat, diethatyl,
 15 difenopenten, difenoxuron, difenzoquat, diflufenican, diflufenzopyr, dimefuron,
 dimepiperate, dimethachlor, dimethametryn, dimethenamid, dimethenamid-P, dimexano,
 dimidazon, dinitramine, dinofenate, dinoprop, dinosam, dinoseb, dinoterb, diphenamid,
 dipropetryn, diquat, disul, dithiopyr, diuron, DMPA, DNOC, DSMA, EBEP, eglinazone,
 endothal, epronaz, EPTC, erbon, esprocarb, ethalfluralin, ethbenzamide, ethametsulfuron,
 20 ethidimuron, ethiolate, ethobenzamid, etobenzamid, ethofumesate, ethoxyfen,
 ethoxysulfuron, etinofen, etnipromid, etobenzanid, EXD, fenasulam, fenoprop, fenoxaprop,
 fenoxaprop-P-ethyl, fenoxaprop-P-ethyl + isoxadifen-ethyl, fenoxasulfone, fenteracol,
 fenthiaprop, fentrazamide, fenuron, ferrous sulfate, flamprop, flamprop-M, flazasulfuron,
 florasulam, fluazifop, fluazifop-P-butyl, fluazolate, flucarbazone, flucetosulfuron,
 25 fluchloralin, flumipropyn, fluometuron, fluorodifen, fluoroglycofen, fluoromidine,
 fluoronitrofen, fluothiuron, flupoxam, flupropacil, flupropanate, flupyrsulfuron, fluridone,
 flurochloridone, fluroxypyr, fluroxypyr-meptyl, flurtamone, fluthiacet, fomesafen,
 foramsulfuron, fosamine, fumiclorac, furyloxyfen, glufosinate, glufosinate-ammonium,
 glufosinate-P-ammonium, glyphosate, halauxifen, halauxifen-methyl, halosafen,
 30 halosulfuron-methyl, haloxydine, haloxyfop-methyl, haloxyfop-P-methyl, hexachloroacetone,
 hexaflurate, hexazinone, imazamethabenz, imazamox, imazapic, imazapyr, imazaquin,
 imazosulfuron, imazethapyr, indanofan, indaziflam, iodobonil, iodomethane, iodosulfuron,

iodosulfuron-ethyl-sodium, iofensulfuron, ioxynil, ipazine, iprymidam, isocarbamid, isocil,
 isomethiozin, isonoruron, isopolinate, isopropalin, isoproturon, isouron, isoxaben,
 isoxachlortole, isoxaflutole, isoxapyrifop, karbutilate, ketospiradox, lactofen, lenacil, linuron,
 MAA, MAMA, MCPA esters and amines, MCPA-thioethyl, MCPB, mecoprop, mecoprop-P,
 5 medinoterb, mefenacet, mefluidide, mesoprazine, mesosulfuron, mesotrione, metam,
 metamifop, metamitron, metazachlor, metazosulfuron, metflurazon, methabenzthiazuron,
 methalpropalin, methazole, methiobencarb, methiozolin, methiuron, methometon,
 methoprottryne, methyl bromide, methyl isothiocyanate, methyldymron, metobenzuron,
 metobromuron, metolachlor, metosulam, metoxuron, metribuzin, metsulfuron, metsulfuron-
 10 methyl, molinate, monalide, monisouron, monochloroacetic acid, monolinuron, monuron,
 morfamquat, MSMA, naproanilide, napropamide, naptalam, neburon, nicosulfuron,
 nipyraclufen, nitralin, nitrofen, nitrofluorfen, norflurazon, noruron, OCH, orbencarb, *ortho*-
 dichlorobenzene, orthosulfamuron, oryzalin, oxapyrazon, oxasulfuron, oxaziclomefone,
 paraflufen-ethyl, parafluron, paraquat, pebulate, pelargonic acid, pendimethalin, penoxsulam,
 15 pentachlorophenol, pentanochlor, perfluidone, pethoxamid, phenisopham, phenmedipham,
 phenmedipham-ethyl, phenobenzuron, phenylmercury acetate, picloram, picolinafen,
 pinoxaden, piperophos, potassium arsenite, potassium azide, potassium cyanate, pretilachlor,
 primisulfuron-methyl, procyazine, prodiamine, profluzol, profluralin, profoxydim,
 proglinazine, prohexadione-calcium, prometon, prometryn, pronamide, propachlor, propanil,
 20 propaquizafop, propazine, propham, propisochlor, propoxycarbazone, propyrisulfuron,
 propyzamide, prosulfalin, prosulfocarb, prosulfuron, proxan, prynachlor, pydanon,
 pyrasulfotole, pyrazogyl, pyrazolynate, pyrazosulfuron-ethyl, pyrazoxyfen, pyribenzoxim,
 pyributicarb, pyriclor, pyridafol, pyridate, pyriftalid, pyriminobac, pyrimisulfan, pyritiobac-
 sodium, pyroxasulfone, pyroxsulam, quinclorac, quinmerac, quinochloramine, quinonamid,
 25 quizalofop, quizalofop-P-ethyl, rhodethanil, rimsulfuron, S-metolachlor, sebuthylazine,
 secbumeton, sethoxydim, siduron, simazine, simeton, simetryn, SMA, sodium arsenite,
 sodium azide, sodium chlorate, sulcotrione, sulfallate, sulfentrazone, sulfometuron, sulfosate,
 sulfosulfuron, sulfuric acid, sulglycapin, swep, SYN-523, TCA, tebutam, tebuthiuron,
 tefuryltrione, tembotrione, tepraloxymid, terbacil, terbucarb, terbutachlor, terbumeton,
 30 terbuthylazine, terbutryn, tetrafluron, thenylchlor, thiazafluron, thiazopyr, thidiazimin,
 thidiazuron, thienencarbazone-methyl, thifensulfuron, thifensulfuron-methyl, thiobencarb,
 tiocarbamil, tioclorim, topramezone, tralkoxydim, triafamone, tri-allate, triasulfuron,

triaziflam, tribenuron, tribenuron-methyl, tricamba, triclopyr choline salt, triclopyr esters and salts, tridiphane, trietazine, trifloxysulfuron, trifluralin, triflusulfuron, trifop, trifopsime, trihydroxytriazine, trimeturon, tripropindan, tritac tritosulfuron, vernolate, xylachlor and salts, esters, optically active isomers and mixtures thereof.

5 The compositions and methods described herein, can further be used in conjunction with glyphosate, 5-enolpyruvylshikimate-3-phosphate (EPSP) synthase inhibitors, glufosinate, glutamine synthetase inhibitors, dicamba, phenoxy auxins, pyridyloxy auxins, synthetic auxins, auxin transport inhibitors, aryloxyphenoxypropionates, cyclohexanediones, phenylpyrazolines, acetyl CoA carboxylase (ACCase) inhibitors, imidazolinones,
10 sulfonyleureas, pyrimidinylthiobenzoates, triazolopyrimidines, sulfonyleaminocarbonyltriazolinones, acetolactate synthase (ALS) or acetohydroxy acid synthase (AHAS) inhibitors, 4-hydroxyphenyl-pyruvate dioxygenase (HPPD) inhibitors, phytoene desaturase inhibitors, carotenoid biosynthesis inhibitors, protoporphyrinogen oxidase (PPO) inhibitors, cellulose biosynthesis inhibitors, mitosis inhibitors, microtubule
15 inhibitors, very long chain fatty acid inhibitors, fatty acid and lipid biosynthesis inhibitors, photosystem I inhibitors, photosystem II inhibitors, triazines, and bromoxynil on glyphosate-tolerant, EPSP synthase inhibitor-tolerant, glufosinate-tolerant, glutamine synthetase inhibitor-tolerant, dicamba-tolerant, phenoxy auxin-tolerant, pyridyloxy auxin-tolerant, auxin-tolerant, auxin transport inhibitor-tolerant, aryloxyphenoxypropionate-tolerant,
20 cyclohexanedione-tolerant, phenylpyrazoline-tolerant, ACCase-tolerant, imidazolinone-tolerant, sulfonyleurea-tolerant, pyrimidinylthiobenzoate-tolerant, triazolopyrimidine-tolerant, sulfonyleaminocarbonyltriazolinone-tolerant, ALS- or AHAS-tolerant, HPPD-tolerant, phytoene desaturase inhibitor-tolerant, carotenoid biosynthesis inhibitor tolerant, PPO-tolerant, cellulose biosynthesis inhibitor-tolerant, mitosis inhibitor-tolerant, microtubule
25 inhibitor-tolerant, very long chain fatty acid inhibitor-tolerant, fatty acid and lipid biosynthesis inhibitor-tolerant, photosystem I inhibitor-tolerant, photosystem II inhibitor-tolerant, triazine-tolerant, and bromoxynil-tolerant crops, and crops possessing multiple or stacked traits conferring tolerance to multiple chemistries and/or multiple modes of action via single and/or multiple resistance mechanisms. In some embodiments, the compound of
30 formula (I) or salt or ester thereof and complementary herbicide or salt or ester thereof are used in combination with herbicides that are selective for the crop being treated and which complement the spectrum of weeds controlled by these compounds at the application rate

employed. In some embodiments, the compositions described herein and other complementary herbicides are applied at the same time, either as a combination formulation, as a tank mix, or as a sequential application.

In some embodiments, the compositions described herein are employed in combination with one or more herbicide safeners, such as AD-67 (MON 4660), benoxacor, benthicarb, brassinolide, cloquintocet (mexyl), cyometrinil, daimuron, dichlormid, dicyclonon, dimepiperate, disulfoton, fenchlorazole-ethyl, fenclorim, flurazole, fluxofenim, furilazole, harpin proteins, isoxadifen-ethyl, jiecaowan, jiecaoxi, mefenpyr-diethyl, mephenate, naphthalic anhydride (NA), oxabetrinil, R29148 and *N*-phenyl-sulfonylbenzoic acid amides, to enhance their selectivity. In some embodiments, the safeners are employed in rice, cereal, corn, or maize settings. In some embodiments, the safener is cloquintocet or an ester or salt thereof. In certain embodiments, cloquintocet is utilized to antagonize harmful effects of the compositions on rice and cereals. In some embodiments, the safener is cloquintocet (mexyl).

In some embodiments, the compositions described herein are employed in combination with one or more plant growth regulators, such as 2,3,5-tri-iodobenzoic acid, IAA, IBA, naphthaleneacetamide, α -naphthaleneacetic acids, benzyladenine, 4-hydroxyphenethyl alcohol, kinetin, zeatin, endothal, ethephon, pentachlorophenol, thidiazuron, tribufos, aviglycine, gibberellins, gibberellic acid, abscisic acid, ancymidol, fosamine, glyphosine, isopyrimol, jasmonic acid, maleic hydrazide, mepiquat, 2,3,5-tri-iodobenzoic acid, morphactins, dichlorflurenol, flurprimidol, mefluidide, paclobutrazol, tetcyclacis, uniconazole, brassinolide, brassinolide-ethyl, cycloheximide, ethylene, methasulfocarb, prohexadione, triapenthenol and trinexapac.

In some embodiments, the plant growth regulators are employed in one or more crops or settings, such as rice, cereal crops, corn, maize, broadleaf crops, oilseed rape/canola, turf, pineapple, sugarcane, sunflower, pastures, grasslands, rangelands, fallowland, turf, tree and vine orchards, plantation crops, vegetables, and non-crop (ornamentals) settings. In some embodiments, the plant growth regulator is mixed with the compound of formula (I), or mixed with the compound of formula (I) and a protoporphyrinogen oxidase to cause a preferentially advantageous effect on plants.

In some embodiments, compositions provided herein further comprise at least one agriculturally acceptable adjuvant or carrier. Suitable adjuvants or carriers should not be

phytotoxic to valuable crops, particularly at the concentrations employed in applying the compositions for selective weed control in the presence of crops, and should not react chemically with herbicidal components or other composition ingredients. Such mixtures can be designed for application directly to weeds or their locus or can be concentrates or formulations that are normally diluted with additional carriers and adjuvants before application. They can be solids, such as, for example, dusts, granules, water-dispersible granules, or wettable powders, or liquids, such as, for example, emulsifiable concentrates, solutions, emulsions or suspensions. They can also be provided as a pre-mix or tank mixed.

Suitable agricultural adjuvants and carriers include, but are not limited to, crop oil concentrate; nonylphenol ethoxylate; benzylcocoalkyldimethyl quaternary ammonium salt; blend of petroleum hydrocarbon, alkyl esters, organic acid, and anionic surfactant; C₉-C₁₁ alkylpolyglycoside; phosphated alcohol ethoxylate; natural primary alcohol (C₁₂-C₁₆) ethoxylate; di-*sec*-butylphenol EO-PO block copolymer; polysiloxane-methyl cap; nonylphenol ethoxylate + urea ammonium nitrate; emulsified methylated seed oil; tridecyl alcohol (synthetic) ethoxylate (8EO); tallow amine ethoxylate (15 EO); PEG(400) dioleate-99.

Liquid carriers that can be employed include water and organic solvents. The organic solvents include, but are not limited to, petroleum fractions or hydrocarbons such as mineral oil, aromatic solvents, paraffinic oils, and the like; vegetable oils such as soybean oil, rapeseed oil, olive oil, castor oil, sunflower seed oil, coconut oil, corn oil, cottonseed oil, linseed oil, palm oil, peanut oil, safflower oil, sesame oil, tung oil and the like; esters of the above vegetable oils; esters of monoalcohols or dihydric, trihydric, or other lower polyalcohols (4-6 hydroxy containing), such as 2-ethyl hexyl stearate, *n*-butyl oleate, isopropyl myristate, propylene glycol dioleate, di-octyl succinate, di-butyl adipate, di-octyl phthalate and the like; esters of mono, di and polycarboxylic acids and the like. Specific organic solvents include, but are not limited to toluene, xylene, petroleum naphtha, crop oil, acetone, methyl ethyl ketone, cyclohexanone, trichloroethylene, perchloroethylene, ethyl acetate, amyl acetate, butyl acetate, propylene glycol monomethyl ether and diethylene glycol monomethyl ether, methyl alcohol, ethyl alcohol, isopropyl alcohol, amyl alcohol, ethylene glycol, propylene glycol, glycerine, *N*-methyl-2-pyrrolidinone, *N,N*-dimethyl alkylamides, dimethyl sulfoxide, liquid fertilizers and the like. In certain embodiments, Water is the carrier for the dilution of concentrates.

Suitable solid carriers include but are not limited to talc, pyrophyllite clay, silica, attapulgus clay, kaolin clay, kieselguhr, chalk, diatomaceous earth, lime, calcium carbonate, bentonite clay, Fuller's earth, cottonseed hulls, wheat flour, soybean flour, pumice, wood flour, walnut shell flour, lignin, cellulose, and the like.

5 In some embodiments, the compositions described herein further comprise one or more surface-active agents. In some embodiments, such surface-active agents are employed in both solid and liquid compositions, and in certain embodiments those designed to be diluted with carrier before application. The surface-active agents can be anionic, cationic or nonionic in character and can be employed as emulsifying agents, wetting agents, suspending
10 agents, or for other purposes. Surfactants which may also be used in the present formulations are described, *inter alia*, in "McCutcheon's Detergents and Emulsifiers Annual," MC Publishing Corp., Ridgewood, New Jersey, 1998 and in "Encyclopedia of Surfactants," Vol. I-III, Chemical Publishing Co., New York, 1980-81. Surface-active agents include, but are not limited to salts of alkyl sulfates, such as diethanolammonium lauryl sulfate;
15 alkylarylsulfonate salts, such as calcium dodecylbenzenesulfonate; alkylphenol-alkylene oxide addition products, such as nonylphenol-C₁₈ ethoxylate; alcohol-alkylene oxide addition products, such as tridecyl alcohol-C₁₆ ethoxylate; soaps, such as sodium stearate; alkyl-naphthalene-sulfonate salts, such as sodium dibutylnaphthalenesulfonate; dialkyl esters of sulfosuccinate salts, such as sodium di(2-ethylhexyl) sulfosuccinate; sorbitol esters, such as
20 sorbitol oleate; quaternary amines, such as lauryl trimethylammonium chloride; polyethylene glycol esters of fatty acids, such as polyethylene glycol stearate; block copolymers of ethylene oxide and propylene oxide; salts of mono and dialkyl phosphate esters; vegetable or seed oils such as soybean oil, rapeseed/canola oil, olive oil, castor oil, sunflower seed oil, coconut oil, corn oil, cottonseed oil, linseed oil, palm oil, peanut oil, safflower oil, sesame oil,
25 tung oil and the like; and esters of the above vegetable oils, and in certain embodiments, methyl esters.

In some embodiments, these materials, such as vegetable or seed oils and their esters, can be used interchangeably as an agricultural adjuvant, as a liquid carrier or as a surface active agent.

30 Other exemplary additives for use in the compositions provided herein include but are not limited to compatibilizing agents, antifoam agents, sequestering agents, neutralizing agents and buffers, corrosion inhibitors, dyes, odorants, spreading agents, penetration aids,

sticking agents, dispersing agents, thickening agents, freezing point depressants, antimicrobial agents, and the like. The compositions may also contain other compatible components, for example, other herbicides, plant growth regulants, fungicides, insecticides, and the like and can be formulated with liquid fertilizers or solid, particulate fertilizer carriers such as ammonium nitrate, urea and the like.

In some embodiments, the concentration of the active ingredients in the compositions described herein is from about 0.0005 to 98 percent by weight. In some embodiments, the concentration is from about 0.0006 to 90 percent by weight. In compositions designed to be employed as concentrates, the active ingredients, in certain embodiments, are present in a concentration from about 0.1 to 98 weight percent, and in certain embodiment's about 0.5 to 90 weight percent. Such compositions are, in certain embodiments, diluted with an inert carrier, such as water, before application. The diluted compositions usually applied to weeds or the locus of weeds contain, in certain embodiments, about 0.0006 to 3.0 weight percent active ingredient and in certain embodiments contain about 0.01 to 1.0 weight percent.

The present compositions can be applied to weeds or their locus by the use of conventional ground or aerial dusters, sprayers, and granule applicators, by addition to irrigation or paddy water, and by other conventional means known to those skilled in the art.

The described embodiments and following examples are for illustrative purposes and are not intended to limit the scope of the claims. Other modifications, uses, or combinations with respect to the compositions described herein will be apparent to a person of ordinary skill in the art without departing from the spirit and scope of the claimed subject matter.

Examples

Results in Examples I, II, III and IV are greenhouse trial results.

Example I. Evaluation of Postemergence Foliar-Applied Herbicidal Mixtures for Weed

Control in Direct Seeded Rice

Seeds or nutlets of the desired test plant species were planted in a soil matrix prepared by mixing a loam or sandy loam soil (e.g., 28.6 percent silt, 18.8 percent clay, and 52.6 percent sand, with a pH of about 5.8 and an organic matter content of about 1.8 percent) and calcareous grit in an 80 to 20 ratio. The soil matrix was contained in plastic pots with a volume of 1 quart and a surface area of 83.6 square centimeters (cm²). When required to ensure good germination and healthy plants, a fungicide treatment and/or other chemical or physical treatment was applied. The plants were grown for 8-22 days in a greenhouse with

an approximate 14 h photoperiod which was maintained at about 29° C during the day and 26° C during the night. Nutrients (Peters Excel[®] 15-5-15 5-Ca 2-Mg and iron chelate) were applied in the irrigation solution as needed and water was added on a regular basis.


Supplemental lighting was provided with overhead metal halide 1000-Watt lamps as

5 necessary. The plants were employed for testing when they reached the first through fourth true leaf stage.


Treatments consisted of the acid or esters of 4-amino-3-chloro-5-fluoro-6-(4-chloro-2-fluoro-3-methoxy-phenyl)pyridine-2-carboxylic acid (Compound A), each formulated as an SC (suspension concentrate), and various herbicidal components alone and in combination.

10 Forms of compound A were applied on an acid equivalent basis.


Forms of compound A (compound of formula I) tested include:



Compound A Acid



Compound A *n*-Butyl Ester



Compound A Benzyl Ester

- 5 Other herbicidal components were applied on an active ingredient basis and included protoporphyrinogen oxidase (PPO)-inhibiting herbicides carfentrazone-ethyl formulated as Aim[®] EC, oxyfluorfen formulated as GoalTender[®] SC or technical material, pyraclonil formulated as Pyraclon[®], oxadiazon formulated as Ronstar[®] 50W, and saflufenacil formulated as Sharpen[®], acifluorfen sodium salt formulated as Blazer, bifenox formulated as Fox or
10 technical grade material, flumioxazin formulated as Payload, and pyraflufen-ethyl (technical grade material).

Treatment requirements were calculated based upon the rates being tested, the concentration of active ingredient or acid equivalent in the formulation, and a 12 mL application volume at a rate of 187 L/ha.

- 15 For treatments comprised of formulated compounds, measured amounts of compounds were placed individually in 25 mL glass vials and diluted in a volume of 1.25% (v/v) Agri-Dex[®] crop oil concentrated to obtain 12X stock solutions. If a test compound did not dissolve readily, the mixture was warmed and/or sonicated. Application solutions were prepared by adding an appropriate amount of each stock solution (*e.g.*, 1 mL) and diluted to
20 the appropriate final concentrations with the addition of 10 mL of an aqueous mixture of 1.25% (v/v) crop oil concentrate so that the final spray solutions contained 1.25+/-0.05% (v/v) crop oil concentrate.

- For treatments comprised of technical compounds, weighed amounts can be placed individually in 25 mL glass vials and dissolved in a volume of 97:3 v/v acetone/DMSO to
25 obtain 12X stock solutions. If a test compound does not dissolve readily, the mixture can be warmed and/or sonicated. Application solutions can be prepared by adding an appropriate amount of each stock solution (*e.g.*, 1 mL) and diluted to the appropriate final concentrations

with the addition of 10 mL of an aqueous mixture of 1.5% (v/v) crop oil concentrate so that the final spray solutions contain 1.25% (v/v) crop oil concentrate. When technical materials are used, the concentrated stock solutions can be added to the spray solutions so that the final acetone and DMSO concentrations of the application solutions are 16.2% and 0.5%,
5 respectively.

For treatments comprised of formulated and technical compounds, weighed amounts of the technical materials were placed individually in 25 mL glass vials and dissolved in a volume of 97:3 v/v acetone/DMSO to obtain 12X stock solutions, and measured amounts of the formulated compounds were placed individually in 25 mL glass vials and diluted in a
10 volume of 1.5% (v/v) crop oil concentrate or water to obtain 12X stock solutions. If a test compound did not dissolve readily, the mixture was warmed and/or sonicated. Application solutions were prepared by adding an appropriate amount of each stock solution (*e.g.*, 1 mL) and diluted to the appropriate final concentrations with the addition of an appropriate amount of an aqueous mixture of 1.5% (v/v) crop oil concentrate so that the final spray solutions
15 contained 1.25% (v/v) crop oil concentrate. When required, additional water and/or 97:3 v/v acetone/DMSO was added to individual application solutions so that the final acetone and DMSO concentrations of the application solutions being compared were 8.1% and 0.25%, respectively.

All stock solutions and applications solutions were visually inspected for compound
20 compatibility prior to application. Spray solutions were applied to the plant material with an overhead Mandel track sprayer equipped with a 8002E nozzles calibrated to deliver 187 L/ha over an application area of 0.503 m² at a spray height of 18 to 20 inches (46 to 50 cm) above average plant canopy height. Control plants were sprayed in the same manner with the solvent blank.

25 The treated plants and control plants were placed in a greenhouse as described above and watered by sub-irrigation to prevent wash-off of the test compounds. After approximately 3 weeks, the condition of the test plants as compared with that of the untreated plants was determined visually and scored on a scale of 0 to 100 percent where 0 corresponds to no injury or growth inhibition and 100 corresponds to complete kill.

30 Colby's equation was used to determine the herbicidal effects expected from the mixtures (Colby, S.R. 1967. Calculation of the synergistic and antagonistic response of herbicide combinations. Weeds 15:20-22.) .

The following equation was used to calculate the expected activity of mixtures containing two active ingredients, A and B:

$$\text{Expected} = A + B - (A \times B/100)$$

A = observed efficacy of active ingredient A at the same concentration as used in the mixture.

B = observed efficacy of active ingredient B at the same concentration as used in the mixture.

The compounds tested, application rates employed, plant species tested, and results are given in Tables 1-21.

10 Table 1. Synergistic Activity of Foliar-Applied Compound A Acid and Carfentrazone-ethyl Herbicidal Compositions on Weed Control in a Rice Cropping System.

Compound A Acid	Carfentrazone-ethyl	Visual Weed Control (%) - 21 DAA			
		DIGSA		ECHCG	
g ae/ha	g ai/ha	Obs	Exp	Obs	Exp
4.38	0	33	-	33	-
8.75	0	40	-	55	-
17.5	0	43	-	93	-
0	14	0	-	10	-
4.38	14	50	33	50	39
8.75	14	35	40	70	60
17.5	14	60	43	95	93

Compound A Acid	Carfentrazone-ethyl	Visual Weed Control (%) - 21 DAA	
		CYPES	
g ae/ha	g ai/ha	Obs	Exp
4.38	0	18	-
8.75	0	80	-
0	7	0	-
0	14	0	-
4.38	7	40	18
8.75	7	95	80
4.38	14	50	18
8.75	14	90	80

Compound A Acid	Carfentrazone-ethyl	Visual Weed Control (%) - 19 DAA	
		LEFCH	
g ae/ha	g ai/ha	Obs	Exp
19.4	0	5	-
0	28	10	-
19.4	28	38	15

Table 2. Synergistic Activity of Foliar-Applied Compound A Benzyl Ester and Carfentrazone-ethyl Herbicidal Compositions on Weed Control in a Rice Cropping System.

Compound A Benzyl Ester	Carfentrazone-ethyl	Visual Weed Control (%) - 21 DAA			
		ECHCG		CYPUR	
g ae/ha	g ai/ha	Obs	Exp	Obs	Exp
8.75	0	55	-	85	-
0	3.5	0	-	0	-
0	7	0	-	0	-
0	14	10	-	0	-
8.75	3.5	75	55	100	85
8.75	7	90	55	100	85
8.75	14	80	60	85	85

Table 3. Synergistic Activity of Foliar-Applied Compound A n-Butyl Ester and

5 Carfentrazone-ethyl Herbicidal Compositions on Weed Control in a Rice Cropping System.

Compound A n-Butyl Ester	Carfentrazone-ethyl	Visual Weed Control (%) - 19 DAA	
		LEFCH	
g ae/ha	g ai/ha	Obs	Exp
16	0	10	-
0	28	10	-
16	28	35	19

Table 4. Synergistic Activity of Foliar-Applied Compound A Acid and Oxyfluorfen (technical) Herbicidal Compositions on Weed Control in a Rice Cropping System.

Compound A Acid	Oxyfluorfen	Visual Weed Control (%) - 21 DAA
		DIGSA

g ae/ha	g ai/ha	Obs	Exp
5.3	0	20	-
10.6	0	30	-
21.2	0	35	-
0	90	70	-
0	180	50	-
0	360	85	-
5.3	90	60	76
10.6	90	99	79
21.2	90	99	81
5.3	180	100	60
10.6	180	95	65
21.2	180	100	68
5.3	360	100	88
10.6	360	100	90
21.2	360	100	90

Table 5. Synergistic Activity of Foliar-Applied Compound A Benzyl Ester and Oxyfluorfen (technical) Herbicidal Compositions on Weed Control in a Rice Cropping System.

Compound A Benzyl Ester	Oxyfluorfen	Visual Weed Control (%) - 21 DAA	
		DIGSA	
g ae/ha	g ai/ha	Obs	Exp
4.38	0	10	-
8.75	0	45	-
17.5	0	70	-
0	180	50	-
4.38	180	100	55
8.75	180	99	73
17.5	180	95	85

Compound A Benzyl Ester	Oxyfluorfen	Visual Weed Control (%) - 21 DAA	
		CYPIR	
g ae/ha	g ai/ha	Obs	Exp
4.38	0	30	-
0	90	0	-
0	180	30	-
0	360	40	-
4.38	90	95	30

4.38	180	100	51
4.38	360	100	58

Table 6. Synergistic Activity of Foliar-Applied Compound A Acid and Oxyfluorfen (SC) Herbicidal Compositions on Weed Control in a Rice Cropping System.

Compound A Acid	Oxyfluorfen	Visual Weed Control (%) - 23 DAA	
		ECHCG	
g ae/ha	g ai/ha	Obs	Exp
4.38	0	45	-
8.75	0	55	-
17.5	0	80	-
0	180	30	-
4.38	180	80	62
8.75	180	85	69
17.5	180	80	86

Compound A Acid	Oxyfluorfen	Visual Weed Control (%) - 23 DAA	
		ECHCO	
g ae/ha	g ai/ha	Obs	Exp
4.38	0	15	-
8.75	0	60	-
0	90	50	-
4.38	90	95	58
8.75	90	100	80

5

Table 7. Synergistic Activity of Foliar-Applied Compound A Benzyl Ester and Oxyfluorfen (SC) Herbicidal Compositions on Weed Control in a Rice Cropping System.

Compound A Benzyl Ester	Oxyfluorfen	Visual Weed Control (%) - 23 DAA	
		DIGSA	
g ae/ha	g ai/ha	Obs	Exp
4.38	0	0	-
8.75	0	0	-
17.5	0	15	-
0	180	60	-
4.38	180	65	60
8.75	180	100	60

17.5	180	100	66
------	-----	-----	----

Compound A Benzyl Ester	Oxyfluorfen	Visual Weed Control (%) - 23 DAA	
		ECHCG	
g ae/ha	g ai/ha	Obs	Exp
4.38	0	40	-
0	90	30	-
0	180	30	-
0	360	95	-
4.38	90	80	58
4.38	180	100	58
4.38	360	100	97

Compound A Benzyl Ester	Oxyfluorfen	Visual Weed Control (%) - 23 DAA	
		IPOHE	
g ae/ha	g ai/ha	Obs	Exp
4.38	0	10	-
8.75	0	10	-
17.5	0	40	-
0	90	85	-
0	180	85	-
4.38	90	100	87
8.75	90	100	87
17.5	90	100	91
4.38	180	99	87
8.75	180	100	87
17.5	180	100	91

Table 8. Synergistic Activity of Foliar-Applied Compound A Acid and Pyraclonil Herbicidal Compositions on Weed Control in a Rice Cropping System.

Compound A Acid	Pyraclonil	Visual Weed Control (%) - 20 DAA	
		BRAPP	
g ae/ha	g ai/ha	Obs	Exp
4.38	0	10	-
0	33.75	0	-
0	67.5	0	-

0	135	0	-
4.38	33.75	45	10
4.38	67.5	50	10
4.38	135	55	10

Compound A Acid	Pyraclonil	Visual Weed Control (%) - 20 DAA			
		ECHCG		ECHCO	
g ac/ha	g ai/ha	Obs	Exp	Obs	Exp
4.38	0	25	-	50	-
8.75	0	80	-	75	-
0	33.75	0	-	0	-
0	67.5	0	-	15	-
0	135	0	-	20	-
4.38	33.75	65	25	75	50
8.75	33.75	90	80	95	75
4.38	67.5	85	25	70	58
8.75	67.5	95	80	80	79
4.38	135	90	25	70	60
8.75	135	95	80	90	80

Compound A Acid	Pyraclonil	Visual Weed Control (%) - 21 DAA	
		ECHCG	
g ac/ha	g ai/ha	Obs	Exp
17.5	0	88	-
35	0	85	-
0	270	0	-
17.5	270	90	88
35	270	100	85

Compound A Acid	Pyraclonil	Visual Weed Control (%) - 21 DAA	
		IPOHE	
g ac/ha	g ai/ha	Obs	Exp
35	0	45	-
0	270	45	-
35	270	88	70

Table 9. Synergistic Activity of Foliar-Applied Compound A Benzyl Ester and Pyraclonil Herbicidal Compositions on Weed Control in a Rice Cropping System.

Compound A Benzyl Ester	Pyraclonil	Visual Weed Control (%) - 20 DAA	
		ECHCO	
g ae/ha	g ai/ha	Obs	Exp
4.38	0	40	-
0	33.75	0	-
0	67.5	15	-
0	135	20	-
4.38	33.75	50	40
4.38	67.5	60	49
4.38	135	75	52

Compound A Benzyl Ester	Pyraclonil	Visual Weed Control (%) - 21 DAA	
		IPOHE	
g ae/ha	g ai/ha	Obs	Exp
17.5	0	10	-
35	0	50	-
0	270	45	-
17.5	270	70	51
35	270	93	73

Compound A Benzyl Ester	Pyraclonil	Visual Weed Control (%) - 21 DAA	
		ISCRU	
g ae/ha	g ai/ha	Obs	Exp
8	0	20	-
16	0	0	-
0	67.5	0	-
0	135	0	-
0	270	0	-
8	67.5	50	20
16	67.5	30	0
8	135	70	20
16	135	100	0
8	270	95	20

16	270	60	0
----	-----	----	---

Table 10. Synergistic Activity of Foliar-Applied Compound A Acid and Oxadiazon Herbicidal Compositions on Weed Control in a Rice Cropping System.

Compound A Acid	Oxadiazon	Visual Weed Control (%) - 20 DAA	
		IPOHE	
g ae/ha	g ai/ha	Obs	Exp
8.75	0	10	-
0	31.25	15	-
0	62.5	20	-
0	125	30	-
8.75	31.25	20	24
8.75	62.5	50	28
8.75	125	50	37

Compound A Acid	Oxadiazon	Visual Weed Control (%) - 20 DAA	
		IPOHE	
g ae/ha	g ai/ha	Obs	Exp
17.5	0	30	-
35	0	45	-
0	250	5	-
17.5	250	63	34
35	250	75	48

5

Table 11. Synergistic Activity of Foliar-Applied Compound A Benzyl Ester and Oxadiazon Herbicidal Compositions on Weed Control in a Rice Cropping System.

Compound A Benzyl Ester	Oxadiazon	Visual Weed Control (%) - 20 DAA	
		IPOHE	
g ae/ha	g ai/ha	Obs	Exp
4.38	0	10	-
8.75	0	10	-
17.5	0	15	-
0	31.25	15	-
0	62.5	20	-

0	125	30	-
4.38	31.25	40	24
8.75	31.25	40	24
17.5	31.25	60	28
4.38	62.5	60	28
8.75	62.5	30	28
17.5	62.5	30	32
4.38	125	50	37
8.75	125	55	37
17.5	125	50	41

Compound A Benzyl Ester	Oxadiazon	Visual Weed Control (%) - 21 DAA	
		IPOHE	
g ae/ha	g ai/ha	Obs	Exp
17.5	0	10	-
0	250	5	-
17.5	250	45	15

Table 12. Synergistic Activity of Foliar-Applied Compound A n-Butyl Ester and Oxadiazon Herbicidal Compositions on Weed Control in a Rice Cropping System.

Compound A n-Butyl Ester	Oxadiazon	Visual Weed Control (%) - 21 DAA	
		IPOHE	
g ae/ha	g ai/ha	Obs	Exp
17.5	0	40	-
0	250	5	-
17.5	250	60	43

5

Table 13. Synergistic Activity of Foliar-Applied Compound A Acid and Saflufenacil Herbicidal Compositions on Weed Control in a Rice Cropping System.

Compound A Acid	Saflufenacil	Visual Weed Control (%) - 20 DAA	
		LEFCH	
g ae/ha	g ai/ha	Obs	Exp
4.38	0	0	-
8.75	0	30	-
17.5	0	30	-

0	5	0	-
4.38	5	30	0
8.75	5	30	30
17.5	5	45	30

Compound A Acid	Saflufenacil	Visual Weed Control (%) - 20 DAA			
		ECHCO		CYPIR	
g ae/ha	g ai/ha	Obs	Exp	Obs	Exp
4.38	0	40	-	30	-
0	5	0	-	0	-
0	10	40	-	0	-
0	20	40	-	100	-
4.38	5	45	40	100	30
4.38	10	70	64	70	30
4.38	20	80	64	100	100

Table 14. Synergistic Activity of Foliar-Applied Compound A Benzyl Ester and Saflufenacil Herbicidal Compositions on Weed Control in a Rice Cropping System.

Compound A Benzyl Ester	Saflufenacil	Visual Weed Control (%) - 20 DAA	
		ECHCO	
g ae/ha	g ai/ha	Obs	Exp
4.38	0	60	-
8.75	0	80	-
0	10	40	-
0	20	40	-
4.38	10	40	76
8.75	10	99	88
4.38	20	90	76
8.75	20	90	88

5

Compound A Benzyl Ester	Saflufenacil	Visual Weed Control (%) - 20 DAA	
		LEFCH	
g ae/ha	g ai/ha	Obs	Exp
8.75	0	30	-
17.5	0	50	-
0	5	0	-

0	10	0	-
8.75	5	80	30
17.5	5	40	50
8.75	10	20	30
17.5	10	70	50

Compound A Benzyl Ester	Saflufenacil	Visual Weed Control (%) - 20 DAA	
		CYPIR	
g ae/ha	g ai/ha	Obs	Exp
4.38	0	70	-
0	5	0	-
0	10	0	-
0	20	100	-
4.38	5	100	70
4.38	10	100	70
4.38	20	100	100

Compound A Benzyl Ester	Saflufenacil	Visual Weed Control (%) - 21 DAA	
		ISCRU	
g ae/ha	g ai/ha	Obs	Exp
8	0	20	-
16	0	0	-
0	10	0	-
0	20	0	-
8	10	100	20
16	10	70	0
8	20	70	20
16	20	70	0

Table 15. Synergistic Activity of Foliar-Applied Compound A Acid and Oxadiargyl

5 Herbicidal Compositions on Weed Control in a Rice Cropping System.

Compound A Acid	Oxadiargyl	Visual Weed Control (%) - 22 DAA	
		IPOHE	
g ae/ha	g ai/ha	Obs	Exp
4.38	0	15	-

8.75	0	35	-
17.5	0	50	-
0	50	25	-
4.38	50	85	36
8.75	50	70	51
17.5	50	70	63

Table 16. Synergistic Activity of Foliar-Applied Compound A Benzyl Ester and Oxadiargyl Herbicidal Compositions on Weed Control in a Rice Cropping System.

Compound A Benzyl Ester	Oxadiargyl	Visual Weed Control (%) - 22 DAA	
		IPOHE	
g ae/ha	g ai/ha	Obs	Exp
4.38	0	10	-
8.75	0	35	-
0	50	25	-
4.38	50	85	33
8.75	50	75	51

- 5 Table 17. Synergistic Activity of Foliar-Applied Compound A Benzyl Ester and Acifluorfen Sodium Herbicidal Compositions on Weed Control in a Rice Cropping System.

Compound A Benzyl Ester	Acifluorfen sodium	Visual Weed Control (%) - 21 DAA	
		ECHCO	
g ae/ha	g ai/ha	Obs	Exp
8	0	40	-
0	140	0	-
0	280	25	-
8	140	60	40
8	280	65	55

Table 18. Synergistic Activity of Foliar-Applied Compound A Benzyl Ester and Bifenox (technical) Herbicidal Compositions on Weed Control in a Rice Cropping System.

Compound A Benzyl Ester	Bifenox	Visual Weed Control (%) - 22 DAA	
		IPOHE	
g ae/ha	g ai/ha	Obs	Exp

8	0	10	-
16	0	40	-
32	0	45	-
0	360	40	-
0	720	65	-
8	360	70	46
16	360	80	64
32	360	70	67
8	720	95	69
16	720	80	79
32	720	100	81

Compound A Benzyl Ester	Bifenox	Visual Weed Control (%) - 22 DAA	
		CYPUR	
g ae/ha	g ai/ha	Obs	Exp
8	0	10	-
16	0	60	-
0	180	0	-
0	720	0	-
8	180	30	10
16	180	100	60
8	720	30	10
16	720	70	60

Table 19. Synergistic Activity of Foliar-Applied Compound A Benzyl Ester and Bifenox (Fox[®]) Herbicidal Compositions on Weed Control in a Rice Cropping System.

Compound A Benzyl Ester	Bifenox	Visual Weed Control (%) - 21 DAA	
		IPOHE	
g ae/ha	g ai/ha	Obs	Exp
16	0	10	-
32	0	45	-
0	180	55	-
16	180	95	60
32	180	100	75

Compound A Benzyl Ester	Bifenox	Visual Weed Control (%) - 21 DAA	
		CYPiR	
g ae/ha	g ai/ha	Obs	Exp
8	0	30	-
0	180	0	-
0	360	0	-
0	720	0	-
8	180	30	30
8	360	85	30
8	720	50	30

Table 20. Synergistic Activity of Foliar-Applied Compound A Benzyl Ester and Flumioxazin Herbicidal Compositions on Weed Control in a Rice Cropping System.

Compound A Benzyl Ester	Flumioxazin	Visual Weed Control (%) - 20 DAA	
		ECHCO	
g ae/ha	g ai/ha	Obs	Exp
8	0	40	-
16	0	60	-
0	140	60	-
0	280	70	-
8	140	90	76
16	140	90	84
8	280	90	82
16	280	100	88

5 Table 21. Synergistic Activity of Foliar-Applied Compound A Benzyl Ester and Pyraflufen-ethyl Herbicidal Compositions on Weed Control in a Rice Cropping System.

Compound A Benzyl Ester	Pyraflufen- ethyl	Visual Weed Control (%) - 21 DAA	
		ECHCG	
g ae/ha	g ai/ha	Obs	Exp
8	0	45	-
0	5.3	15	-
0	10.6	15	-
8	5.3	65	53
8	10.6	80	53

Compound A Benzyl Ester	Pyraflufen- ethyl	Visual Weed Control (%) - 21 DAA	
		CYPİR	
g ae/ha	g ai/ha	Obs	Exp
8	0	35	-
0	2.65	0	-
0	5.3	0	-
0	10.6	20	-
8	2.65	50	35
8	5.3	70	35
8	10.6	60	48

BRAPP	<i>Brachiaria platyphylla</i> (Griseb.) Nash	signalgrass, broadleaf
CYPDI	<i>Cyperus difformis</i> L.	sedge, smallflower umbrella
5 CYPES	<i>Cyperus esculentus</i> L.	nutsedge, yellow
CYPİR	<i>Cyperus iria</i> L.	flatsedge, rice
DIGSA	<i>Digitaria sanguinalis</i> (L.) Scop.	crabgrass, large
ECHCG	<i>Echinochloa crusgalli</i> (L.) Beauv.	barnyardgrass
ECHCO	<i>Echinochloa colona</i> (L.) Link	junglerice
10 IPOHE	<i>Ipomoea hederacea</i> Jacq.	morningglory, ivyleaf
ISCRU	<i>Ischaemum rugosum</i> Salisb.	saramollagrass
LEFCH	<i>Leptochloa chinensis</i> (L.) Nees	sprangletop, Chinese
SCPJU	<i>Schoenoplectus juncooides</i> (Roxb.) Palla	bulrush, Japanese

g ae/ha = grams acid equivalent per hectare

15 g ai/ha = grams active ingredient per hectare

Obs = observed value

Exp = expected value as calculated by Colby's equation

DAA = days after application

Example II. Evaluation of In-Water Applied Herbicidal Mixtures for Weed Control in

20 Transplanted Paddy Rice

Weed seeds or nutlets of the desired test plant species were planted in puddled soil (mud) prepared by mixing a shredded, non-sterilized mineral soil (50.5 percent silt, 25.5 percent clay, and 24 percent sand, with a pH of about 7.6 and an organic matter content of

about 2.9 percent) and water at a 1:1 volumetric ratio. The prepared mud was dispensed in 365 mL aliquots into 16-ounce (oz.) non-perforated plastic pots with a surface area of 86.59 square centimeters (cm²) leaving a headspace of 3 centimeters (cm) in each pot. Mud was allowed to dry overnight prior to planting or transplanting. Rice seeds were planted in Sun

5 Gro MetroMix[®] 306 planting mixture, which typically has a pH of 6.0 to 6.8 and an organic matter content of about 30 percent, in plastic plug trays. Seedlings at the second or third leaf stage of growth were transplanted into 840 mL of mud contained in 32-oz. non-perforated plastic pots with a surface area of 86.59 cm² 4 days prior to herbicide application. The paddy was created by filling the headspace of the pots with 2.5 to 3 cm of water. When required to


10 ensure good germination and healthy plants, a fungicide treatment and/or other chemical or physical treatment was applied. The plants were grown for 4-22 days in a greenhouse with an approximate 14 h photoperiod which was maintained at about 29°C during the day and 26°C during the night. Nutrients were added as Osmocote[®] (19:6:12, N:P:K + minor nutrients) at 2 g per 16-oz. pot and 4 g per 32-oz. pot. Water was added on a regular basis to

15 maintain the paddy flood, and supplemental lighting was provided with overhead metal halide 1000-Watt lamps as necessary. The plants were employed for testing when they reached the first through fourth true leaf stage.


Treatments consisted of the acid or esters of 4-amino-3-chloro-5-fluoro-6-(4-chloro-2-fluoro-3-methoxy-phenyl)pyridine-2-carboxylic acid (compound A) each formulated as an

20 SC and various herbicidal components alone and in combination. Forms of compound A were applied on an acid equivalent basis.


Forms of compound A (compound of formula I) tested include:



Compound A Acid



Compound A *n*-Butyl Ester



Compound A Benzyl Ester

5

Other herbicidal components were applied on an active ingredient basis and included protoporphyrinogen oxidase (PPO)-inhibiting herbicides carfentrazone-ethyl formulated as Aim[®] EC, oxyfluorfen formulated as an SC or technical material, pyraclonil formulated as Pyraclon[®], oxadiazon formulated as Ronstar[®] 50W, saflufenacil formulated as Sharpen[®], pentoxazone (technical grade material), bifenox formulated as Fox[®] or technical grade material and oxadiargyl (technical grade material).

10

Treatment requirements for each compound or herbicidal component were calculated based upon the rates being tested, the concentration of active ingredient or acid equivalent in the formulation, an application volume of 2 mL per component per pot, and an application area of 86.59 cm² per pot.

15

For formulated compounds, a measured amount was placed in an individual 100 or 200 mL glass vial and was dissolved in a volume of 1.25% (v/v) Agri-Dex[®] crop oil concentrate to obtain application solutions. If the test compound did not dissolve readily, the mixture was warmed and/or sonicated.

20

For technical grade compounds, a weighed amount was placed in an individual 100 to 200 mL glass vial and was dissolved in a volume of acetone to obtain concentrated stock solutions. If the test compound did not dissolve readily, the mixture was warmed and/or sonicated. The concentrated stock solutions obtained were diluted with an equivalent volume of an aqueous mixture containing 2.5% (v/v) crop oil concentrate so that the final application solutions contained 1.25% (v/v) crop oil concentrate.

Applications were made by injecting with a pipetter appropriate amounts of the application solutions, individually and sequentially, into the aqueous layer of the paddy. Control plants were treated in the same manner with the solvent blank. Applications were made so that all treated plant material received the same concentrations of acetone and crop oil concentrate.

The treated plants and control plants were placed in a greenhouse as described above and water was added as needed to maintain a paddy flood. After approximately 3 weeks the condition of the test plants as compared with that of the untreated plants was determined visually and scored on a scale of 0 to 100 percent where 0 corresponds to no injury or growth inhibition and 100 corresponds to complete kill.

Colby's equation was used to determine the herbicidal effects expected from the mixtures (Colby, S.R. 1967. Calculation of the synergistic and antagonistic response of herbicide combinations. Weeds 15:20-22.) .

The following equation was used to calculate the expected activity of mixtures containing two active ingredients, A and B:

$$\text{Expected} = A + B - (A \times B/100)$$

A = observed efficacy of active ingredient A at the same concentration as used in the mixture.

B = observed efficacy of active ingredient B at the same concentration as used in the mixture.

Some of the compounds tested, application rates employed, plant species tested, and results are given in Tables 22-38.

Table 22. Synergistic Activity of In-Water Applications of Compound A Acid and Carfentrazone-ethyl Herbicidal Compositions on Weed Control in a Rice Cropping System.

Compound A Acid	Carfentrazone-ethyl	Visual Weed Control (%) - 21 DAA
-----------------	---------------------	----------------------------------

		ECHCG	
g ae/ha	g ai/ha	Obs	Exp
8.75	0	0	-
17.5	0	0	-
35	0	40	-
0	14	0	-
0	28	20	-
8.75	14	0	0
17.5	14	20	0
35	14	95	40
8.75	28	30	20
17.5	28	50	20
35	28	95	52

Compound A Acid	Carfentrazone-ethyl	Visual Weed Control (%) - 19 DAA	
		ECHOR	
g ae/ha	g ai/ha	Obs	Exp
42.4	0	15	-
0	215	65	-
42.4	215	100	70

Table 23. Synergistic Activity of In-Water Applications of Compound A Benzyl Ester and Carfentrazone-ethyl Herbicidal Compositions on Weed Control in a Rice Cropping System.

Compound A Benzyl Ester	Carfentrazone-ethyl	Visual Weed Control (%) - 21 DAA	
		LEFCH	
g ae/ha	g ai/ha	Obs	Exp
8.75	0	20	-
17.5	0	30	-
35	0	55	-
0	28	10	-
8.75	28	40	28
17.5	28	40	37
35	28	65	60

Table 24. Synergistic Activity of In-Water Applications of Compound A Acid and Oxyfluorfen (technical) Herbicidal Compositions on Weed Control in a Rice Cropping System.

Compound A Acid	Oxyfluorfen	Visual Weed Control (%) - 22 DAA	
		CYPRO	
g ae/ha	g ai/ha	Obs	Exp
10.6	0	0	-
21.2	0	25	-
42.4	0	95	-
0	180	50	-
10.6	180	90	50
21.2	180	90	63
42.4	180	100	98

Compound A Acid	Oxyfluorfen	Visual Weed Control (%) - 22 DAA	
		SCPJU	
g ae/ha	g ai/ha	Obs	Exp
8.75	0	60	-
17.5	0	70	-
35	0	95	-
0	180	0	-
0	360	0	-
8.75	180	55	60
17.5	180	90	70
35	180	100	95
8.75	360	75	60
17.5	360	80	70
35	360	100	95

5

Table 25. Synergistic Activity of In-Water Applications of Compound A Acid and Oxyfluorfen (SC) Herbicidal Compositions on Weed Control in a Rice Cropping System.

Compound A Acid	Oxyfluorfen	Visual Weed Control (%) - 22 DAA	
		CYPRO	
g ae/ha	g ai/ha	Obs	Exp

8.75	0	0	-
17.5	0	20	-
35	0	50	-
0	180	30	-
0	360	20	-
8.75	180	50	30
17.5	180	100	44
35	180	95	65
8.75	360	85	20
17.5	360	85	36
35	360	50	60

Compound A Acid	Oxyfluorfen	Visual Weed Control (%) - 21 DAA	
		CYPRO	
g ae/ha	g ai/ha	Obs	Exp
16	0	0	-
32	0	20	-
64	0	85	-
0	135	0	-
0	270	0	-
16	135	20	0
32	135	20	20
64	135	95	85
16	270	40	0
32	270	60	20
64	270	100	85

Table 26. Synergistic Activity of In-Water Applications of Compound A Benzyl Ester and Oxyfluorfen (SC) Herbicidal Compositions on Weed Control in a Rice Cropping System.

Compound A Benzyl Ester	Oxyfluorfen	Visual Weed Control (%) - 21 DAA	
		CYPRO	
g ae/ha	g ai/ha	Obs	Exp
32	0	90	-
0	135	0	-
0	270	0	-
32	135	95	90
32	270	100	90

Table 27. Synergistic Activity of In-Water Applications of Compound A Acid and Pyraclonil Herbicidal Compositions on Weed Control in a Rice Cropping System.

Compound A Acid	Pyraclonil	Visual Weed Control (%) - 20 DAA	
		ECHCG	
g ae/ha	g ai/ha	Obs	Exp
8.75	0	40	-
17.5	0	30	-
0	45	0	-
8.75	45	100	40
17.5	45	100	30

Table 28. Synergistic Activity of In-Water Applications of Compound A Benzyl Ester and Pyraclonil Herbicidal Compositions on Weed Control in a Rice Cropping System.

Compound A Benzyl Ester	Pyraclonil	Visual Weed Control (%) - 20 DAA	
		ECHOR	
g ae/ha	g ai/ha	Obs	Exp
4.38	0	10	-
8.75	0	20	-
17.5	0	30	-
0	45	40	-
4.38	45	45	46
8.75	45	80	52
17.5	45	100	58

Compound A Benzyl Ester	Pyraclonil	Visual Weed Control (%) - 20 DAA	
		CYPRO	
g ae/ha	g ai/ha	Obs	Exp
4.38	0	20	-
8.75	0	20	-
0	45	0	-
0	90	0	-
4.38	45	20	20
8.75	45	90	20
4.38	90	85	20
8.75	90	85	20

Table 29. Synergistic Activity of In-Water Applications of Compound A Acid and Oxadiazon Herbicidal Compositions on Weed Control in a Rice Cropping System.

Compound A Acid	Oxadiazon	Visual Weed Control (%) - 21 DAA	
		ECHOR	
g ae/ha	g ai/ha	Obs	Exp
8.75	0	10	-
17.5	0	20	-
35	0	35	-
0	62.5	15	-
0	125	20	-
8.75	62.5	40	24
17.5	62.5	40	32
35	62.5	75	45
8.75	125	40	28
17.5	125	50	36
35	125	60	48

5

Compound A Acid	Oxadiazon	Visual Weed Control (%) - 21 DAA	
		ECHOR	
g ae/ha	g ai/ha	Obs	Exp
35	0	10	-
70	0	28	-
0	250	18	-
35	250	40	26
70	250	85	40

Table 30. Synergistic Activity of In-Water Applications of Compound A Benzyl Ester and Oxadiazon Herbicidal Compositions on Weed Control in a Rice Cropping System.

Compound A Benzyl Ester	Oxadiazon	Visual Weed Control (%) - 21 DAA	
		ECHOR	
g ae/ha	g ai/ha	Obs	Exp
8.75	0	25	-

17.5	0	60	-
0	62.5	15	-
8.75	62.5	70	36
17.5	62.5	75	66

Table 31. Synergistic Activity of In-Water Applications of Compound A n-Butyl Ester and Oxadiazon Herbicidal Compositions on Weed Control in a Rice Cropping System.

Compound A n-Butyl Ester	Oxadiazon	Visual Weed Control (%) - 21 DAA	
		ECHOR	
g ae/ha	g ai/ha	Obs	Exp
35	0	25	-
70	0	38	-
0	250	18	-
35	250	43	38
70	250	70	48

5 Table 32. Synergistic Activity of In-Water Applications of Compound A Acid and Saflufenacil Herbicidal Compositions on Weed Control in a Rice Cropping System.

Compound A Acid	Saflufenacil	Visual Weed Control (%) - 20 DAA			
		ECHCG		LEFCH	
g ae/ha	g ai/ha	Obs	Exp	Obs	Exp
17.5	0	50	-	0	-
35	0	95	-	30	-
0	10	0	-	0	-
0	20	10	-	0	-
17.5	10	85	50	30	0
35	10	100	95	30	30
17.5	20	70	55	30	0
35	20	100	96	70	30

Compound A Acid	Saflufenacil	Visual Weed Control (%) - 20 DAA	
		CYPRO	
g ae/ha	g ai/ha	Obs	Exp
8.75	0	0	-
17.5	0	20	-

35	0	80	-
0	10	0	-
8.75	10	20	0
17.5	10	30	20
35	10	95	80

Compound A Acid	Saflufenacil	Visual Weed Control (%) - 21 DAA	
		LEFCH	
g ac/ha	g ai/ha	Obs	Exp
35	0	5	-
70	0	23	-
0	40	5	-
35	40	58	33
70	40	55	38

Table 33. Synergistic Activity of In-Water Applications of Compound A Benzyl Ester and Saflufenacil Herbicidal Compositions on Weed Control in a Rice Cropping System.

Compound A Benzyl Ester	Saflufenacil	Visual Weed Control (%) - 20 DAA	
		ECHOR	
g ac/ha	g ai/ha	Obs	Exp
4.38	0	10	-
8.75	0	15	-
0	10	0	-
0	20	0	-
4.38	10	0	10
8.75	10	30	15
4.38	20	40	10
8.75	20	45	15

5

Compound A Benzyl Ester	Saflufenacil	Visual Weed Control (%) - 20 DAA	
		LEFCH	
g ac/ha	g ai/ha	Obs	Exp
4.38	0	0	
8.75	0	20	-
17.5	0	30	-

0	10	0	-
0	20	0	-
4.38	10	50	0
8.75	10	40	20
17.5	10	100	30
4.38	20	100	0
8.75	20	60	20
17.5	20	100	30

Table 34. Synergistic Activity of In-Water Applications of Compound A n-Butyl Ester and Saflufenacil Herbicidal Compositions on Weed Control in a Rice Cropping System.

Compound A n-Butyl Ester	Saflufenacil	Visual Weed Control (%) - 21 DAA	
		LEFCH	
g ae/ha	g ai/ha	Obs	Exp
35	0	8	-
0	40	5	-
35	40	28	12

Table 35. Synergistic Activity of In-Water Applications of Compound A Acid and

5 Pentoxazone Herbicidal Compositions on Weed Control in a Rice Cropping System.

Compound A Acid	Pentoxazone	Visual Weed Control (%) - 21 DAA	
		ECHCG	
g ae/ha	g ai/ha	Obs	Exp
8.75	0	15	-
17.5	0	20	-
35	0	95	-
0	56.25	0	-
8.75	56.25	30	15
17.5	56.25	40	20
35	56.25	99	95

Compound A Acid	Pentoxazone	Visual Weed Control (%) - 21 DAA	
		CYPRO	
g ae/ha	g ai/ha	Obs	Exp
35	0	70	-
0	56.25	0	-

0	112.5	0	-
35	56.25	85	70
35	112.5	100	70

Compound A Acid	Pentoxazone	Visual Weed Control (%) - 21 DAA	
		FIMMI	
g ae/ha	g ai/ha	Obs	Exp
8.75	0	0	-
0	56.25	25	-
0	112.5	20	-
8.75	56.25	50	25
8.75	112.5	99	20

Compound A Acid	Pentoxazone	Visual Weed Control (%) - 21 DAA	
		SCPJU	
g ae/ha	g ai/ha	Obs	Exp
8.75	0	40	-
17.5	0	60	-
35	0	95	-
0	56.25	0	-
0	112.5	0	-
8.75	56.25	40	40
17.5	56.25	95	60
35	56.25	100	95
8.75	112.5	60	40
17.5	112.5	90	60
35	112.5	100	95

Table 36. Synergistic Activity of In-Water Applications of Compound A Benzyl Ester and Pentoxazone Herbicidal Compositions on Weed Control in a Rice Cropping System.

Compound A Benzyl Ester	Pentoxazone	Visual Weed Control (%) - 21 DAA	
		CYPRO	
g ae/ha	g ai/ha	Obs	Exp
4.38	0	30	-
8.75	0	60	-
0	56.25	0	-

0	112.5	0	-
4.38	56.25	80	30
8.75	56.25	75	60
4.38	112.5	50	30
8.75	112.5	80	60

Table 37. Synergistic Activity of In-Water Applications of Compound A Benzyl Ester and Bifenox (technical) Herbicidal Compositions on Weed Control in a Rice Cropping System.

Compound A Benzyl Ester	Bifenox	Visual Weed Control (%) - 21 DAA	
		ECHOR	
g ae/ha	g ai/ha	Obs	Exp
8	0	20	-
16	0	45	-
32	0	60	-
0	720	40	-
8	720	85	52
16	720	100	67
32	720	100	76

Table 38. Synergistic Activity of In-Water Applications of Compound A Benzyl Ester and Bifenox (Fox[®]) Herbicidal Compositions on Weed Control in a Rice Cropping System.

Compound A Benzyl Ester	Bifenox	Visual Weed Control (%) - 21 DAA	
		SCPMA	
g ae/ha	g ai/ha	Obs	Exp
48	0	0	-
96	0	0	-
0	180	40	-
48	180	0	40
96	180	60	40

	CYPRO	<i>Cyperus rotundus</i> L.	nutsedge, purple
	ECHCG	<i>Echinochloa crusgalli</i> (L.) Beauv.	barnyardgrass
	ECHOR	<i>Echinochloa oryzoides</i> (Ard.) Fritsch	watergrass, early
10	FIMMI	<i>Fimbristylis miliacea</i> (L.) Vahl	fringerush, globe
	LEFCH	<i>Leptochloa chinensis</i> (L.) Nees	sprangletop, Chinese
	SCPJU	<i>Schoenoplectus juncooides</i> (Roxb.) Palla	bulrush, Japanese

SCPMA *Schoenoplectus maritimus* (L.) Lye clubrush, sea

g ae/ha = grams acid equivalent per hectare

g ai/ha = grams active ingredient per hectare

Obs = observed value

5 Exp = expected value as calculated by Colby's equation


DAA = days after application

Example III. Evaluation of Postemergence Herbicidal Activity of Mixtures in Cereal Crops in the Greenhouse.

10 Seeds of the desired test plant species were planted in Sun Gro MetroMix[®] 306
planting mixture, which typically has a pH of 6.0 to 6.8 and an organic matter content of
about 30 percent, in plastic pots with a surface area of 103.2 square centimeters (cm²). When
required to ensure good germination and healthy plants, a fungicide treatment and/or other
chemical or physical treatment was applied. The plants were grown for 7-36 days in a
15 greenhouse with an approximate 14 hour photoperiod which was maintained at about 18° C
during the day and about 17° C during the night. Nutrients and water were added on a
regular basis and supplemental lighting was provided with overhead metal halide 1000-Watt
lamps as necessary. The plants were employed for testing when they reached the second or
third true leaf stage.

20 Treatments consisted of the benzyl ester of 4-amino-3-chloro-5-fluoro-6-(4-chloro-2-
fluoro-3-methoxy-phenyl)pyridine-2-carboxylic acid (Compound A), formulated as an SC, a
second cereal herbicide alone and then both in combination.

Forms of compound A (compound of formula I) tested include:



Compound A Benzyl Ester

Other herbicidal components were applied on an active ingredient basis and included protoporphyrinogen oxidase-inhibiting herbicides.

Measured aliquots of benzyl ester of 4-amino-3-chloro-5-fluoro-6-(4-chloro-2-fluoro-3-methoxy-phenyl)pyridine-2-carboxylic acid (Compound A) were placed in 25 milliliter (mL) glass vials and diluted in a volume of 1.25% (v/v) Agri-Dex[®] crop oil concentrate to obtain stock solutions. Compound requirements are based upon a 12 mL application volume at a rate of 187 liters per hectare (L/ha). Spray solutions of the second cereal herbicide and experimental compound mixtures were prepared by adding the stock solutions to the appropriate amount of dilution solution to form 12 mL spray solution with active ingredients in two- and three-way combinations. Formulated compounds were applied to the plant material with an overhead Mandel track sprayer equipped with an 8002E nozzle calibrated to deliver 187 L/ha over an application area of 0.503 square meters (m²) at a spray height of 18 inches (46 cm) above average plant canopy. Control plants were sprayed in the same manner with the solvent blank.

The treated plants and control plants were placed in a greenhouse as described above and watered by sub-irrigation to prevent wash-off of the test compounds. After 20-22 days, the condition of the test plants as compared with that of the control plants was determined visually and scored on a scale of 0 to 100 percent where 0 corresponds to no injury and 100 corresponds to complete kill.

Colby's equation was used to determine the herbicidal effects expected from the mixtures (Colby, S.R. 1967. Calculation of the synergistic and antagonistic response of herbicide combinations. Weeds 15:20-22.) .

The following equation was used to calculate the expected activity of mixtures containing two active ingredients, A and B:

$$\text{Expected} = A + B - (A \times B/100)$$

A = observed efficacy of active ingredient A at the same concentration as used in the mixture.

B = observed efficacy of active ingredient B at the same concentration as used in the mixture.

The compounds tested, application rates employed, plant species tested, and results are given in Table 39.

Table 39. Synergistic Activity of Foliar-Applied Compound A Benzyl Ester and Bifenox Herbicidal Compositions on Weed Control in a Cereals Cropping System.

Compound A Benzyl Ester	Bifenox	Visual Weed Control (%) - 21 DAA			
		PAPRH		BRSNW	
g ai/ha	g ai/ha	Obs	Exp	Obs	Exp
5	0	28	-	13	-
10	0	35	-	13	-
0	180	5	-	13	-
0	360	8	-	18	-
0	780	8	-	37	-
5	180	33	32	25	24
5	360	35	33	33	28
5	780	70	34	67	45
10	780	68	40	72	45

BRSNW *Brassica napus* L. rape, oilseed (winter)

5 PAPRH *Papaver rhoeas* L. poppy, common

g ac/ha = grams acid equivalent per hectare

g ai/ha = grams active ingredient per hectare

Obs = observed value

Exp = expected value as calculated by Colby's equation

10 DAA = days after application

Example IV. Evaluation of Pre-emergence Soil-Applied Herbicidal Mixtures for Weed Control


15 Seeds or nutlets of the desired test plant species were planted in a soil matrix prepared by mixing a loam soil (32 percent silt, 23 percent clay, and 45 percent sand, with a pH of about 6.5 and an organic matter content of about 1.9 percent) and calcareous grit in an 80 to 20 ratio. The soil matrix was contained in plastic pots with a volume of 1 quart and a surface area of 83.6 square centimeters (cm²).

20 Treatments consisted of the benzyl ester of 4-amino-3-chloro-5-fluoro-6-(4-chloro-2-fluoro-3-methoxy-phenyl)pyridine-2-carboxylic acid (compound A) formulated as an SC

(suspension concentrate) and various herbicidal components alone and in combination.

Forms of compound A were applied on an acid equivalent basis.

Forms of compound A (compound of formula I) tested include:



Compound A Benzyl Ester

Other herbicidal components were applied on an acid equivalent or active ingredient basis and included the protoporphyrinogen oxidase (PPO)-inhibiting herbicides, flumioxazin formulated as Payload[®] and oxadiazon formulated as Ronstar[®] 50W.

Treatment requirements were calculated based upon the rates being tested, the concentration of active ingredient or acid equivalent in the formulation, and a 12 mL application volume at a rate of 187 L/ha.

For treatments comprised of formulated compounds, measured amounts of compounds were placed individually in 25 mL glass vials and diluted in a volume of 1.25% (v/v) Agri-Dex[®] crop oil concentrate (COC) to obtain 12X stock solutions. If a test compound did not dissolve readily, the mixture was warmed and/or sonicated. Application solutions were prepared by adding an appropriate amount of each stock solution (e.g., 1 mL) and diluted to the appropriate final concentrations with the addition of 10 mL of an aqueous mixture of 1.25% (v/v) COC so that the final spray solutions contained 1.25% (v/v) COC.

For treatments comprised of technical compounds, weighed amounts can be placed individually in 25 mL glass vials and dissolved in a volume of 97:3 (v/v) acetone/DMSO to obtain 12X stock solutions. If a test compound does not dissolve readily, the mixture can be warmed and/or sonicated. Application solutions can be prepared by adding an appropriate amount of each stock solution (e.g., 1 mL) and diluted to the appropriate final concentrations with the addition of 10 mL of an aqueous mixture of 1.5% (v/v) COC so that the final spray solutions contain 1.25% (v/v) COC. When technical materials are used, the concentrated

stock solutions can be added to the spray solutions so that the final acetone and DMSO concentrations of the application solutions are 16.2% and 0.5%, respectively.

For treatments comprised of formulated and technical compounds, weighed amounts of the technical materials can be placed individually in 25 mL glass vials and dissolved in a volume of 97:3 (v/v) acetone/DMSO to obtain 12X stock solutions, and measured amounts of the formulated compounds can be placed individually in 25 mL glass vials and diluted in a volume of 1.5% (v/v) COC or water to obtain 12X stock solutions. If a test compound does not dissolve readily, the mixture can be warmed and/or sonicated. Application solutions can be prepared by adding an appropriate amount of each stock solution (*e.g.*, 1 mL) and diluted to the appropriate final concentrations with the addition of an appropriate amount of an aqueous mixture of 1.5% (v/v) COC so that the final spray solutions contain 1.25% (v/v) COC. As required, additional water and/or 97:3 (v/v) acetone/DMSO can be added to individual application solutions so that the final acetone and DMSO concentrations of the application solutions being compared are 8.1% and 0.25%, respectively.

All stock solutions and applications solutions were visually inspected for compound compatibility prior to application. Spray solutions were applied to the soil with an overhead Mandel track sprayer equipped with a 8002E nozzles calibrated to deliver 187 L/ha over an application area of 0.503 m² at a spray height of 18 inches (46 cm) above average pot height. Control pots were sprayed in the same manner with the solvent blank.

The treated and control pots were placed in a greenhouse and top watered as needed. When required to ensure good germination and healthy plants, a fungicide treatment and/or other chemical or physical treatment was applied. The pots were maintained in a greenhouse with an approximate 14 hr photoperiod which was maintained at about 29° C during the day and 26° C during the night. Nutrients (Peters[®] Excel 15-5-15 5-Ca 2-Mg) were applied in the irrigation solution as needed and water was added on a regular basis. Supplemental lighting was provided with overhead metal halide 1000-Watt lamps as necessary. After approximately 4 weeks, the condition of the test plants as compared with that of the untreated plants was determined visually and scored on a scale of 0 to 100 percent where 0 corresponds to no injury or growth inhibition and 100 corresponds to complete kill.

Colby's equation was used to determine the herbicidal effects expected from the mixtures (Colby, S.R. 1967. Calculation of the synergistic and antagonistic response of herbicide combinations. Weeds 15:20-22.) .

The following equation was used to calculate the expected activity of mixtures containing two active ingredients, A and B:

$$\text{Expected} = A + B - (A \times B/100)$$

A = observed efficacy of active ingredient A at the same concentration as used in the mixture.

B = observed efficacy of active ingredient B at the same concentration as used in the mixture.

Some of the compounds tested, application rates employed, plant species tested, and results are given in Tables 40-41.

Table 40. Synergistic Activity of Soil-applied, Pre-emergence Applications of Compound A Benzyl Ester and Flumioxazin Herbicidal Compositions on Weed Control.

Compound A Benzyl Ester	Flumioxazin	Visual Weed Control (%) - 28 DAA	
		IPOHE	
g ae/ha	g ai/ha	Obs	Exp
16	0	13	-
32	0	23	-
0	105	70	-
0	210	70	-
0	420	90	-
16	105	100	74
32	105	80	77
16	210	85	74
32	210	100	77
16	420	100	91
32	420	100	92

Compound A Benzyl Ester	Flumioxazin	Visual Weed Control (%) - 29 DAA	
		XANST	
g ae/ha	g ai/ha	Obs	Exp
16	0	0	-
32	0	5	-
0	105	18	-
0	210	48	-
16	105	38	18

32	105	30	22
16	210	40	48
32	210	65	50

Compound A Benzyl Ester	Fumioxazin	Visual Weed Control (%) - 28 DAA	
		SETVI	
g ae/ha	g ai/ha	Obs	Exp
16	0	10	-
32	0	10	-
0	105	73	-
16	105	100	75
32	105	80	75

Table 41. Synergistic Activity of Soil-applied, Pre-emergence Applications of Compound A Benzyl Ester and Oxadiazon Herbicidal Compositions on Weed Control.

Compound A Benzyl Ester	Oxadiazon	Visual Weed Control (%) - 32 DAA	
		IPOHE	
g ae/ha	g ai/ha	Obs	Exp
8	0	65	-
16	0	83	-
0	560	50	-
0	1120	100	-
8	560	100	83
16	560	100	91
32	560	100	100

5

IPOHE *Ipomoea hederacea* (L.) Jacq.

morningglory, ivyleaf

SETVI *Setaria viridis* (L.) Beauv.

foxtail, green

XANST *Xanthium strumarium* L.

cocklebur, common

g ae/ha = grams acid equivalent per hectare

10 g ai/ha = grams active ingredient per hectare


Obs = observed value

Exp = expected value as calculated by Colby's equation


DAA = days after application

WHAT IS CLAIMED IS:

1. A synergistic herbicidal composition comprising a herbicidally effective amount of (a) a compound of the formula (I)



- 5 or an agriculturally acceptable salt or ester thereof and (b) a protoporphyrinogen oxidase inhibitor.
2. The composition of claim 1, wherein (a) is the compound of formula (I), a C₁₋₄ alkyl ester of the compound of formula (I), or a benzyl ester of the compound of formula (I).
3. The composition of any of claims 1-2, wherein (b) is at least one compound, or an agriculturally acceptable salt, carboxylic acid, carboxylate salt, or ester thereof, selected from the group consisting of: acifluorfen, azafeniden, bifenox, carfentrazone-ethyl, flumioxazin, ipfencarbazone, oxadiargyl, oxadiazon, oxyfluorfen, pentoxazone, pyraclonil, pyraflufen-ethyl, and saflufenacil.
4. The composition of any of claims 1-3, further comprising an agriculturally acceptable adjuvant or carrier.
5. The composition of any of claims 1-4, further comprising a herbicide safener.
6. A method of controlling undesirable vegetation which comprises contacting the vegetation or the locus thereof with or applying to the soil or water to prevent the emergence or growth of vegetation a herbicidally effective amount of (a) a compound of the formula (I)



- or an agriculturally acceptable salt or ester thereof and (b) a protoporphyrinogen oxidase inhibitor.

- 88

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US13/51299

A. CLASSIFICATION OF SUBJECT MATTER

IPC(8) - A01N 43/40 (2013.01)

USPC - 504/244

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC(8): A01N 43/40, 43/54, 57/12, 57/16, 25/00, 25/32, 63/00; A01P 13/00; C07D 213/78, 213/79, 213/72, 211/70, 401/04 (2013.01)

USPC: 504/244, 260, 105, 103, 116.1, 118; 546/304, 310, 314, 326

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

MicroPatent (US-G, US-A, EP-A, EP-B, WO, JP-bib, DE-C,B, DE-A, DE-T, DE-U, GB-A, FR-A); ProQuest; Google; Google Scholar; Scirus; PubMed; PubChem; pyridine; fluoropyridine; picolinate; herbicide; protoporphyrinogen oxidase inhibitor; acifluorfen; bifenox; oxadiazon; synergism

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 2010/0137137 A1 (ROSINGER, CH et al.) 3 June 2010; paragraphs [0005]-[0015], [0039]-[0041], [0445], [0557], [0565], [0568], [0604]-[0605] and [0610]	6-7, 8/6-7
Y	WO 2009/029518 A2 (SATCHIVI, N et al.) 5 March 2009; lines 5-21, page 2; line 10, page 2 to line 2, page 3; lines 3-8, page 7; line 10, page 20 to line 7, page 24	1-2, 3/1-2
A	US 2012/0115727 A1 (SATCHIVI, N et al.) 10 May 2012; abstract	1-2, 3/1-2, 6-7, 8/6-7

☐ Further documents are listed in the continuation of Box C.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

21 November 2013 (21.11.2013)

Date of mailing of the international search report

06 DEC 2013

Name and mailing address of the ISA/US

Mail Stop PCT, Attn: ISA/US, Commissioner for Patents
P.O. Box 1450, Alexandria, Virginia 22313-1450

Facsimile No. 571-273-3201

Authorized officer:

Shane Thomas

PCT Helpdesk: 571-272-4300
PCT OSP: 571-272-7774

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US13/51299

Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☒ Claims Nos.: 4-5 and 9-13
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying additional fees, this Authority did not invite payment of additional fees.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.
- ☐ The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.
- ☐ No protest accompanied the payment of additional search fees.