



(86) Date de dépôt PCT/PCT Filing Date: 2016/08/26
(87) Date publication PCT/PCT Publication Date: 2017/03/09
(45) Date de délivrance/Issue Date: 2024/01/02
(85) Entrée phase nationale/National Entry: 2018/02/27
(86) N° demande PCT/PCT Application No.: EP 2016/070257
(87) N° publication PCT/PCT Publication No.: 2017/036993
(30) Priorité/Priority: 2015/08/28 (GB1515391.9)

(51) Cl.Int./Int.Cl. *A23L 2/00* (2006.01),
A61K 31/366 (2006.01), *A61K 38/01* (2006.01),
A61P 21/00 (2006.01), *C07D 311/80* (2006.01)
(72) Inventeurs/Inventors:
ANDREUX, PENELOPE, CH;
RINSCH, CHRISTOPHER, CH;
BLANCO-BOSE, WILLIAM, CH
(73) Propriétaire/Owner:
AMAZENTIS SA, CH
(74) Agent: BORDEN LADNER GERVAIS LLP

(54) Titre : COMPOSITIONS COMPRENANT UN COMPOSE UROLITHINE
(54) Title: COMPOSITIONS COMPRISING AN UROLITHIN COMPOUND

(57) Abrégé/Abstract:

The invention provides compositions comprising a source of protein and a urolithin. The invention also provides uses and methods associated with, or making use of the compositions, such as a medicament, dietary supplement, functional food or medical food and in the treatment and/or prophylaxis of a muscle-related pathological condition. The invention also provides kits comprising urolithin and protein.



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(10) International Publication Number
WO 2017/036993 A1

(43) International Publication Date
9 March 2017 (09.03.2017)

(51) International Patent Classification:

A23L 2/00 (2006.01) *C07D 311/80* (2006.01)
A61K 38/01 (2006.01) *A61P 21/00* (2006.01)
A61K 31/366 (2006.01)

(21) International Application Number:

PCT/EP2016/070257

(22) International Filing Date:

26 August 2016 (26.08.2016)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

1515391.9 28 August 2015 (28.08.2015) GB

(71) Applicant: **AMAZENTIS SA** [CH/CH]; Parc Scientifique EPFL, PSE-C, 1015 Lausanne (CH).

(72) Inventors: **ANDREUX, Penelope**; 3 Av d'Evian, 1006 Lausanne (CH). **RINSCH, Christopher**; 60 Rue De Lausanne, 1110 Morges (CH). **BLANCO-BOSE, William**; Chemin Chantemerle 13t, 1010 Lausanne (CH).

(74) Agent: **BRADY, Paul Andrew**; 20 Red Lion Street, London Greater London WC1R 4PQ (GB).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))

(54) Title: COMPOSITIONS COMPRISING AN UROLITHIN COMPOUND

(57) Abstract: The invention provides compositions comprising a source of protein and a urolithin. The invention also provides uses and methods associated with, or making use of the compositions, such as a medicament, dietary supplement, functional food or medical food and in the treatment and/or prophylaxis of a muscle-related pathological condition. The invention also provides kits comprising urolithin and protein.



WO 2017/036993 A1

COMPOSITIONS COMPRISING AN UROLITHIN COMPOUND

The current invention relates to nutritional and medical formulations of urolithins and protein, in particular Urolithin A and protein.

Background

- 5 Low muscle mass or poor muscle performance are characteristics of many diseases and conditions.

Muscle-related pathological conditions include myopathies, neuromuscular diseases, such as Duchenne muscular dystrophy, acute sarcopenia, for example muscle atrophy and/or cachexia, for example associated with burns, bed rest, limb immobilization, or major
10 thoracic, abdominal, neck and/or orthopedic surgery. Age-related muscle-loss is an especially prevalent condition. Cachexia due to prolonged immobilization or other diseases, for example cancer, are other conditions that are often characterised by poor muscle performance.

Good muscle performance is important for effective living at all stages of life in healthy
15 individuals as well as in those individuals suffering from a disease, especially the elderly. Improved muscle performance is of particular interest to athletes. For example an increase in muscular contraction strength, increase in amplitude of muscle contraction, or shortening of muscle reaction time between stimulation and contraction are all of benefit to individuals, especially athletes.

- 20 In severe cases of muscular atrophy, an anabolic steroid such as methandrostenolone is administered to patients to assist in the curative process. Such drugs can have numerous side-effects so their long-term use is preferably avoided.


Since the absence of muscle-building amino acids can contribute to muscle wasting, amino acid therapy is beneficial for regenerating damaged or atrophied muscle tissue, and for
25 building muscle mass. The branched-chain amino acids or BCAAs (leucine, isoleucine, and valine) are important to this process, in addition to lysine and other amino acids.

Protein-containing nutritional products are available that are specifically formulated to increase muscle mass, improve muscle performance and/or reduce muscle wasting in elderly or hospitalised individuals. In general, it is necessary to exercise regularly for the desired
30 improvements to be achieved.

Summary of the invention

The present invention provides a composition comprising:

- a) a source of protein; and
- b) a compound of formula (I) or a salt thereof:



(I)

wherein:

A, B, C and D are each independently selected from H and OH;

W, X and Y are each independently selected from H and OH; and

10 Z is selected from H and OH.

Compounds of Formula (I) are members of the Urolithin family; in particular, the compound of Formula (I) is Urolithin A. Administration of a composition of protein and Urolithin A has been found to have a surprisingly beneficial and enhanced effect compared with administration of protein on its own. The composition of the invention is thus useful in the treatment diseases and conditions characterised by low muscle mass or poor muscle performance, and in the enhancement of muscle growth and/or muscle performance. It is also useful in maintaining muscle function.

In general, it is necessary for a subject to exercise in order for amino acid/protein treatments to have an optimal effect. There is evidence that the compositions of the invention are effective in enhancement of muscle function in the absence of exercise, or with less exercise being necessary.

The invention further provides a composition of the invention for use in the treatment of muscle-related pathological conditions. The invention also provides a method of treating a muscle-related pathological condition in a subject comprising administering to the subject an effective amount of a composition of the invention. The invention provides a composition of the invention for use in enhancing muscle performance. The invention also provides a method of enhancing muscle performance by administering to a subject an effective amount of a composition of the invention.

In a treatment of the invention, it is not essential for the Urolithin and protein to be administered simultaneously as part of a single composition. The invention also provides a method of treating a muscle-related pathological condition in a subject, or enhancing muscle function in a subject comprising administering to the subject an effective amount of a urolithin (for example Urolithin A) and an effective amount of protein. The urolithin and the protein can be administered at the same time or separated by a time interval. The invention further provides a kit comprising urolithin and protein for use in such a method. The amount of each of the urolithin and the protein that is administered is as described herein below for unit doses of compositions of the invention.

The invention further provides a compound of Formula (I) or a salt thereof having D_{50} size in the range 0.5 to 50 μm and a D_{90} size in the range 5 to 100 μm . Preferably, the compound or salt has a D_{90} size in the range 8.2 to 16.0 μm , a D_{50} size in the range 2.8 to 5.5 μm and a D_{10} size in the range 0.5 to 1.0 μm . It has been found that the compounds of the invention with those particle sizes have good dispersion and dissolution properties, and enhanced bioavailability.

Summary of the Figures

Figure 1 shows the results of a muscle endurance experiment in which mice were fed with either a high protein diet alone or a high protein diet containing Urolithin A.

Figure 2 shows the results of experiments in which rats were fed with formulations comprising Urolithin A at a variety of different particle sizes and the level of the Urolithin A in the blood was assessed.

Detailed description

As described above, the invention provides a composition comprising a source of protein and a urolithin.

The protein in the composition of the invention is preferably purified protein; that is to say that it has been isolated from the native food ingredient that it was created in. The protein content, as assessed by weight %, in a composition of the invention is preferably higher than in most unprocessed food ingredients. The protein typically makes up at least 15% w/w of the composition of the invention. The protein preferably makes up 20% w/w or more of the composition, for example 25% w/w or more by weight, for example 30% w/w or more by weight, for example 40% w/w or more of the composition, for example 50% w/w or more of the composition, for example 55% w/w or more of the composition, for example 60% w/w or more of the composition. For example the protein may make up 20-99% w/w of the composition, for example 20-90% w/w of the composition, for example, 30-80% /w of the composition, for example 40-80% /w of the composition, for example 50-80% /w of the composition, for example 40-70% /w of the composition.

The weight ratio between the protein component and the urolithin is generally in the range 3:1 to 5000:1, for example 3:1 to 1000:1, for example 5:1 to 1000:1; for example 3:1 to 500:1, for example 3:1 to 250:1, for example 5:1 to 500:1; for example 5:1 to 200:1; for example 5:1 to 100:1; for example 5:1 to 50:1; for example 5:1 to 20:1, for example 6:1 to 15:1. Alternatively, the ratio can be for example in the range of 10:1 to 500:1, for example in the range 10:1 to 100:1, for example 20:1 to 50:1, for example 30:1 to 40:1. In some embodiments, the ratio can be in the range 50:1 to 5000:1; for example 70:1 to 2000:1, for example 100:1 to 500:1. The composition of the invention can be provided as a single serving supplement to a subject's general diet (for example as a bar or a drink); alternatively, it can be provided as a part or the whole of every meal. The weight ratio between the protein component and the urolithin is generally higher if the composition is provided as a part or the whole of every meal than if it is provided as a single serving supplement to a subject's general diet.

The urolithin typically makes up from 0.05 to 10% w/w of the composition of the invention, for example 0.1 to 10% w/w. For example the urolithin may make up 0.2-8% w/w of the composition, for example 0.2-5% w/w of the composition, for example, 0.3-3% /w of the composition, for example 0.5-2% w/w of the composition. In some embodiments, the

urolithin may make up 0.05-5% w/w of the composition, for example 0.1-2% w/w of the composition, for example, 0.2-1% /w of the composition. In some embodiments, the urolithin may make up from 0.05 to 80% w/w of the composition, for example from 0.1 to 75 %w/w of the composition.

5 *Sources of protein:*

Non-limiting examples of suitable protein or sources thereof for use in the compositions include hydrolyzed, partially hydrolyzed or non-hydrolyzed proteins or protein sources. They may be derived from any known or otherwise suitable source such as milk (e.g., casein, whey), animal (e.g., meat, fish), cereal (e.g., rice, corn) or vegetable (e.g., soy, pea) sources.

- 10 Combinations of sources or types of proteins may be used. Non- limiting examples of proteins or sources thereof include intact pea protein, intact pea protein isolates, intact pea protein concentrates, milk protein isolates, milk protein concentrates, casein protein isolates, casein protein concentrates, whey protein concentrates, whey protein isolates, sodium or calcium caseinates, whole cow's milk, partially or completely defatted milk, yoghurt, soy
- 15 protein isolates and soy protein concentrates, and combinations thereof. Combinations of sources or types of proteins may be used.

Preferred proteins include pea protein, whey protein, soy protein and casein. Casein proteins may, for example, be or include sodium caseinate and calcium caseinate.

- Mixtures of sources of proteins may have particular benefits. For example, two or more of
- 20 pea protein, whey protein, soy protein and casein may be present. For example sodium caseinate, milk protein concentrate, and soy protein isolate may all be present. Combinations of casein protein and whey protein are especially preferred, for example containing 60 to 90% w/w whey protein and 10 to 40% w/w casein protein, for example 65 to 80% w/w whey protein and 20 to 35% w/w casein protein, for example 70% w/w whey protein and 30% w/w
- 25 casein protein.

The source of protein may be provided by individual amino acids or polypeptide made up of many amino acids, or mixtures of those.

- For many muscle growth, muscle maintenance and/or muscle enhancement treatments, it is beneficial for certain particular amino acids to be provided. For example, L-arginine, L-
- 30 glutamine, lysine and the branched-chain amino acids (ie leucine, isoleucine, and valine; in

particular leucine and isoleucine) are considered important. In certain embodiments, the source of protein contains a high proportion of those amino acids.

These particular amino acids may be provided as the source of protein, or they may be additional to a main source of protein. Thus, the source of protein in the composition of the invention may include one or more branched-chain amino acids (leucine, isoleucine, and valine). The source of protein in the composition of the invention may include one or both of L-arginine and L-glutamine. The source of protein in the composition of the invention may include lysine.

In a preferred composition, the protein component comprises whey protein and/or casein protein together with one or more individual amino acids, for example one or more of (or all of) leucine, isoleucine and L-arginine.


Urolithins:

Urolithins are metabolites produced by the action of mammalian, including human, gut microbiota on ellagitannins and ellagic acid. Ellagitannins and ellagic acid are compounds commonly found in foods such as pomegranates, nuts and berries. Ellagitannins are minimally absorbed in the gut themselves. Urolithins are a class of compounds with the representative structure (I) shown above. The structures of some particularly common urolithins are described in Table 1 below, with reference to structure (I).

	Substituent of structure (I)					
	A	B	C	D	W, X and Y	Z
Urolithin A	H	H	H	OH	H	OH
Urolithin B	H	H	H	H	H	OH
Urolithin C	H	H	OH	OH	H	OH
Urolithin D	OH	H	OH	OH	H	OH
Urolithin E	OH	OH	H	OH	H	OH
Isourolithin A	H	H	OH	H	H	OH
Isourolithin B	H	H	OH	H	H	H
Urolithin M-5	OH	OH	OH	OH	H	OH
Urolithin M-6	H	OH	OH	OH	H	OH
Urolithin M-7	H	OH	H	OH	H	OH

In practice, for commercial scale products, it is convenient to synthesise the urolithins. Routes of synthesis are described, for example, in WO2014/004902.

Particularly suitable compounds for use in compositions of the invention are the naturally-occurring urolithins. Thus, Z is preferably OH and W, X and Y are preferably all H. When W, X and Y are all H, and A, and B are both H, and C, D and Z are all OH, then the compound is Urolithin C. When W, X and Y are all H, and A, B and C are all H, and D and Z are both OH, then the compound is Urolithin A. Preferably, the Urolithin used in a formulation of the invention is Urolithin A or Urolithin C. Most preferably, the Urolithin used in a formulation of the invention is Urolithin A.



Urolithin A

Preferably, urolithin for use in compositions of the invention is micronized. It has been found by the current inventors that micronized urolithin can be dissolved or suspended more rapidly and more effectively than unmicronised urolithin.

The invention further provides micronized urolithin. The invention further provides a composition contained micronized urolithin. Micronized urolithin preferably has a D_{50} size of under $100\text{ }\mu\text{m}$ – that is to say that 50% of the urolithin by mass has a particle diameter size of under $100\text{ }\mu\text{m}$. More preferably, the urolithin has a D_{50} size of under $75\text{ }\mu\text{m}$, for example under $50\text{ }\mu\text{m}$, for example under $25\text{ }\mu\text{m}$, for example under $20\text{ }\mu\text{m}$, for example under $10\text{ }\mu\text{m}$. More preferably, the urolithin has a D_{50} in the range $0.5\text{--}50\text{ }\mu\text{m}$, for example 0.5 to $20\text{ }\mu\text{m}$, for example 0.5 to $10\text{ }\mu\text{m}$, for example 1 to $10\text{ }\mu\text{m}$, for example 2.8 to $5.5\text{ }\mu\text{m}$. Preferably, the urolithin has a D_{90} size of under $100\text{ }\mu\text{m}$ - that is to say that 90% of the urolithin by mass has

a particle diameter size of under 100 μm . More preferably, the urolithin has a D_{90} size of under 75 μm , for example under 50 μm , for example under 25 μm , for example under 20 μm , for example under 15 μm . The urolithin preferably has a D_{90} in the range 5 to 100 μm , for example 5 to 50 μm , for example 5 to 20 μm , for example 8.2 to 16.0 μm . Preferably, the urolithin has a D_{10} in the range 0.5 – 1.0 μm . Preferably, the urolithin has a D_{90} in the range 8.2 to 16.0 μm , a D_{50} in the range 2.8 to 5.5 μm and a D_{10} in the range 0.5 to 1.0 μm .

Micronisation can be achieved by methods established in the art, for example compressive force milling, hammermilling, universal or pin milling, or jet milling (for example spiral jet milling or fluidised-bed jet milling) may be used. Jet milling is especially suitable.

Forms of compositions:

The compositions of the invention can take any suitable physical form. They may be in the form of a solid (for example a tablet or a bar), a semi-solid (for example a softgel, capsule (for example a hard capsule) or dragee), a powder or a liquid (including emulsions). The compositions of the invention may be pharmaceutical compositions. The compositions of the invention may be nutritional compositions.

Tablet form compositions may be of any suitable type, and they may contain excipients conventional in the art. The excipients can, for example, provide a desired hardness, shelf-life and flavour such that the composition has an acceptable taste, an attractive appearance and good storage stability. A bar may be of any suitable type and it may contain ingredients conventionally used for the preparation of snack bars.

Semi-solid forms may likewise contain excipients conventional in the art. The excipients can, for example, provide a desired hardness, shelf-life and flavour such that the composition has an acceptable taste, an attractive appearance and good storage stability.

Powders are commonly used for the supply of nutritional and medical compositions.

Powders have the advantage that multiple doses can be provided in a simple container, and doses of various sizes can be used from the same supplied container. Powders generally have good storage properties. Powder compositions may also contain excipients conventional in the art. The excipients can, for example, provide a shelf-life, flavour and moisture resistance such that the composition has an acceptable taste, an attractive appearance and good storage stability. Protein powders are widely available on the market. The current invention may take the form of a kit comprising a protein powder composition together with a separate solid

or liquid composition containing urolithin. A solid or liquid composition containing urolithin (for example a tablet or a drink, or other form described herein) may be provided with instructions for use together with a protein powder. The current invention may take the form of a protein powder mixed with urolithin.

- 5 Liquid compositions may be in the form of a medicine, in the form of a drink. Liquid formulations may be solutions, emulsions, slurries or other semi-liquids. Excipients in a liquid composition can, for example, provide a shelf-life, visual appearance, flavour and mouthfeel such that the composition has an acceptable taste, an attractive appearance and good storage stability.
- 10 Compositions may be in the form of a yoghurt. For example, Greek- and Icelandic-style yoghurts are known to commonly have an especially high protein content which makes them especially suitable for use in formulations of the invention. Yoghurts for use in compositions of the invention may contain, for example, from 2 to 15g of protein per 100g. Particularly preferred are yoghurts with a high protein content, for example from 6 to 15g per 100g, for
15 example from 7 to 15g per 100g, for example from 8 to 15g per 100g. The yogurts for use in compositions of the invention may be either spoonable or drinkable in format. Optionally, supplemental protein may also be added to a yoghurt formulation to increase the protein content of the formulation. Yoghurts of the invention may contain live cultures, such as *S. thermophilus*, *L. bulgaricus*. and/or *L. acidophilus*. *L. lactis*.
- 20 *Additional components in compositions of the invention:*
The composition according to the invention may contain additional components beyond the urolithin and the protein. The additional components may be compounds that provide health benefits, for example selected from vitamins, minerals, polyunsaturated fatty acids, functional amino acids and other compounds.
- 25 Amongst vitamins, there may specifically be mentioned Vitamin A, Vitamin C, Vitamin D, Vitamin B12 and Vitamin K2. As used herein, "vitamin D" refers, to any of known form of vitamin D, and specifically includes vitamin D2 (ergocalciferol), vitamin D3 (cholecalciferol), vitamin D precursors, metabolites and another analogues, and combinations thereof, as well as the various active and inactive forms of vitamin D. For example, vitamin
30 D3 may be provided in its unhydroxylated inactive form as cholecalciferol, or may be provided in its hydroxylated active form as calcitriol.

Creatine has been described as having beneficial effects in the treatment of muscle disorders. It can be included in compositions of the invention. β -hydroxyl- β -methylbutyrate (HMB) has been described as having beneficial effects in the treatment of muscle disorders. It can be included in compositions of the invention.

- 5 Amongst minerals, there may specifically be mentioned calcium salts (for example calcium phosphate), selenium, and iron salts.

Polyunsaturated fatty acids are fatty acids that contain more than one double bond in the backbone. This class includes many important compounds, such as essential fatty acids, e.g., omega-3 and omega-6 fatty acids. Long chain polyunsaturated fatty acids are suitable, and preferably those having at least 20 carbon atoms in the molecule. Such long chain omega-3 fatty acids include cis-11, 14, 17-eicosatrienoic acid (ETE) C20:3, cis-8, 11, 14, 17-eicosatetraenoic acid (ETA) C20:4, cis-5,8, 11, 14, 17-eicosapentaenoic acid (EPA) C20:5, cis- 7, 10, 13, 16, 19-docosapentaenoic acid (DPA, Clupanodonic acid) C22:5, cis- 4, 7, 10, 13, 16, 19-docosaheptaenoic acid (DHA) C22:6, cis-9, 12, 15, 18,21- tetracosapentaenoic acid C24:5; cis-6,9, 12, 15, 18,21-tetracosahexaenoic acid (Nisinic acid) C24:6. Long chain omega-6 fatty acids having at least 20 carbon atoms include cis-11, 14-eicosadienoic acid C20:2, cis-8, 11, 14-eicosatrienoic acid (Dihomo-gamma-linolenic acid) (DGLA) C20:3, cis-5,8, 11, 14-eicosatetraenoic acid (Arachidonic acid) (AA) C20:4, cis-13, 16-docosadienoic acid C22:2, cis- 7, 10, 13, 16-docosatetraenoic acid (Adrenic acid) C22:4, cis-4, 7, 10, 13, 16-docosapentaenoic acid (Osbond acid) C22:5. The composition according to the invention preferably contains EPA, DHA or a combination of them, for example in an amount from 10 to 1,000 mg; for example in an amount from 25 to 250mg.

For many muscle growth and/or muscle enhancement treatments, it is beneficial for certain particular amino acids to be provided. For example, L-arginine, L-glutamine, lysine and the branched-chain amino acids are considered important. These amino acids are sometimes known as “functional amino acids”. The composition of the invention may include one or more branched-chain amino acids (leucine, isoleucine, and valine). The composition of the invention may include one or both of L-arginine and L-glutamine. The composition of the invention may include lysine.

- 30 Pharmaceutical compositions of the invention may include additional pharmaceutically active compounds.

A composition of the invention may include one or more agents that are useful for mitochondrial biogenesis or the treatment of mitochondrial disorders. Such compounds include, without limitation, resveratrol, pyrroloquinoline quinone, genistein, hydroxytyrosol, quercetin, L-carnitine, alpha-lipoic acid, and folinic acid (e.g., as leucovorin).

- 5 Additional compounds may further (or alternatively) be included in a composition of the invention, including for example tomatidine, ursolic acid, curcumin, capsaicin, menthol, troamine salicylate and methylsalicylate.

In some exemplary embodiments, the compositions of the present disclosure may comprise, in addition to protein and urolithin, one or more additional macronutrients, typically fat or
10 carbohydrate, or both fat and carbohydrate.

Any suitable source of fat or oil of the type commonly used in the preparation of foodstuffs and pharmaceuticals may be used in compositions of the invention. Non-limiting examples of suitable sources of fats for use in the compositions described herein also include polyunsaturated fatty acids such as docosahexaenoic acid (DHA), arachidonic acid (ARA),
15 eicosapentaenoic acid (EPA) and combinations thereof.

Non-limiting examples of suitable carbohydrates or sources thereof for use in the compositions described herein may include maltodextrin, hydrolyzed or modified starch or cornstarch, glucose polymers, corn syrup, corn syrup solids, rice-derived carbohydrates, glucose, fructose, lactose, high fructose corn syrup, tapioca dextrin, isomaltulose, sucromalt,
20 maltitol powder, glycerin, fructooligosaccharides, soy fiber, corn fiber, guar gum, konjac flour, polydextrose, honey, sugar alcohols (e.g., maltitol, erythritol, sorbitol), and combinations thereof. Maltodextrin, sucrose and fructose are especially preferred.

The total concentrations or amounts of the protein, fat, carbohydrates and other components vary depending upon the nutritional needs of the intended user.

- 25 Additional components in a composition of the invention may be compounds that do not provide health benefits to the subject, but instead improve the composition in some other way, for example its taste, texture or shelflife as mentioned above. The composition of the invention may thus further contain one or more compounds selected from emulsifiers, colorants, preservatives, gums, setting agents, thickeners, sweeteners and flavourings.

Suitable emulsifiers, colorants, preservatives, gums, setting agents and thickeners are well known in the art of manufacture of emulsions and other semi-liquids. For example preservatives, such as benzoic acid, sorbic acid, phosphoric acid, lactic acid, acetic acid, hydrochloric acid and the soluble salts thereof may be used.

5 A sweetener may be especially beneficial in a composition of the invention. High potency non-nutritive carbohydrate sweetening agents may be used, for example selected from aspartame, sucrose, potassium acesulfame, saccharin, cyclamates, Stevia, thaumatin and mixtures thereof. Aspartame is especially suitable.

A flavouring may be especially beneficial in a composition of the invention. In a liquid or
10 semi-liquid composition, fruit flavour can be provided by inclusion of a fruit sauce or puree. Typical flavorings include strawberry, raspberry, blueberry, apricot, pomegranate, peach, pineapple, lemon, orange and apple. Generally, fruit flavorings include fruit extract, fruit preserve or fruit puree, with any of a combination of sweeteners, starch, stabilizer, natural and/or artificial flavors, colorings, preservatives, water and citric acid or other suitable acid to
15 control the pH.

Dosing

The effective amount of the composition to be taken will vary depending upon the manner of administration, the age, body weight, and general health of the subject. Factors such as the
20 disease state, age, and weight of the subject may be important, and dosage regimens may be adjusted to provide the optimum response.

Conventional protein compositions are generally taken at a level of 5 to 30g per serving, and a subject generally takes one, two or three servings per day. The daily dose of protein from the composition is thus generally in the range of 5 to 90g, for example 5 to 80g, for example
25 5 to 70g, for example 5 to 60g, for example 5 to 50g, for example 10 to 40g, for example 20 to 40g.

For an elderly subject, recommended protein intake and exercise for optimal muscle function have been described, for example in: 'Recommendations from the ESPEN Expert Group': *Clinical Nutrition*, 33 (2014) 929e936. It is recommended: (a) for healthy older people, the
30 diet should provide at least 1.0 -1.2 g protein/kg body weight/day; (b) for older people who are malnourished or at risk of malnutrition because they have acute or chronic illness, the diet

should provide 1.2 - 1.5 g protein/kg body weight/day, with even higher intake for individuals with severe illness or injury.

A unit dose composition of the invention preferably contains 5 to 90g, for example 5 to 80g, for example 5 to 70g, for example 5 to 60g of protein, for example 5 to 40g of protein, for example 10 to 40g of protein for example 10 to 30g of protein. A unit dose may be in the form of a snack bar; a snack bar of weight 25 to 150g, for example 40 to 100g may contain the necessary amount of protein (such as 10 to 40g of protein, or another amount mentioned above). A unit dose composition may alternatively be in the form of a drink, for example provided in a container (for example a pouch or a bottle) of a volume suitable for a single dose (for example 50 to 500ml, for example 100 to 300ml). A drink of 100 to 300ml may contain the necessary amount of protein. A unit dose composition may alternatively be in the form of a powder to be reconstituted into a drink, for example a suitable quantity of powder for a single dose (for example 20gr to 60gr of powder, containing 10gr to 40gr of protein). A reconstituted drink of 100 to 500ml may contain the necessary amount of protein.

Daily intake of the urolithin (for example Urolithin A) component is typically in the range of 10mg to 5g per day, for example 20mg to 2500mg per day, for example 20mg to 500mg per day, for example, 10mg to 100mg per day, for example 50mg to 1500mg per day, for example 250mg to 1500mg per day, for example 50mg to 1000mg per day, for example 20mg to 250mg per day, for example 250mg to 1000mg per day, for example 500mg to 1000mg per day, for example 750mg to 1000mg per day. In one embodiment, the composition is taken in an amount to provide a dosage of urolithin in the range from about 0.2 mg/kg/day to greater than about 100 mg/kg/day. For example, the dosage of urolithin may be 0.2 to 100, 0.2 to 50, 0.2 to 25, 0.2 to 10, 0.2 to 7.5, 0.2 to 5, 0.25 to 100, 0.25 to 25, 0.25 to 25, 0.25 to 10, 0.25 to 7.5, 0.25 to 5, 0.5 to 50, 0.5 to 25, 0.5 to 20, 0.5 to 15, 0.5 to 10, 0.5 to 7.5, 0.5 to 5, 0.75 to 50, 0.75 to 25, 0.75 to 20, 0.75 to 15, 0.75 to 10, 0.75 to 7.5, 0.75 to 5, 1.0 to 50, 1 to 25, 1 to 20, 1 to 15, 1 to 10, 1 to 7.5, 1 to 5, 2 to 50, 2 to 25, 2 to 20, 2 to 15, 2 to 10, 2 to 7.5, or 2 to 5 mg/kg/day.

A unit dose composition of the invention preferably contains 10mg to 5g of urolithin, for example 20mg to 2500mg, for example 50mg to 1500mg, for example 250mg to 1500mg, for example 50mg to 1000mg, for example 50mg to 250mg, for example 250mg to 1000mg. A unit dose may be in the form of a snack bar; a snack bar of weight 25 to 150g (for example 40 to 100g) may contain the necessary amount of urolithin. A unit dose composition may

alternatively be in the form of a drink, for example provided in a container (for example a pouch) of a volume suitable for a single dose (for example 100 to 300ml). A drink of 50 to 500ml (for example 100 to 300ml) may contain the necessary amount of urolithin. A drink providing the composition of the invention may contain the urolithin at a concentration of 0.1 to 50mg per ml, for example 0.5 to 10mg per ml, for example 1 to 5mg per ml.

The weight ratio between the protein component and the urolithin is generally in the range 1:10 to 5000:1; for example 1:10 to 1:1; for example 1:2.5 to 5000:1; for example 1:2.5 to 1200:1; for example 1:1 to 1200:1; for example 1:1 to 600:1; for example 2:1 to 5000:1; for example 2:1 to 12000:1; for example 3:1 to 5000:1; for example 3:1 to 1200:1; for example 3:1 to 1000:1; for example 3:1 to 500:1; for example 3:1 to 250:1; for example 5:1 to 1200:1; for example 5:1 to 600:1; for example 5:1 to 500:1; for example 5:1 to 400:1; for example 5:1 to 200:1; for example 5:1 to 100:1; for example 5:1 to 80:1; for example 5:1 to 50:1; for example 5:1 to 20:1, for example 6:1 to 15:1. Alternatively, the ratio can be for example 10:1 to 500:1, for example 10:1 to 100:1, for example 20:1 to 50:1, for example 30:1 to 40:1. In some embodiments, the ratio can be in the range 50:1 to 5000:1; for example 70:1 to 2000:1, for example 100:1 to 500:1.

The urolithin typically makes up from 0.05 to 10% w/w of the composition of the invention, 0.1 to 10% w/w of the composition of the invention. For example the urolithin may make up 0.2-8% w/w of the composition, for example 0.2-5% w/w of the composition, for example, 0.3-3% /w of the composition, for example 0.5-2% /w of the composition. In some embodiments, the urolithin may make up 0.05-5% w/w of the composition, for example 0.1-2% w/w of the composition, for example, 0.2-1% /w of the composition.

A composition of the invention may thus contain 5 to 80g of protein and 10mg to 5g of urolithin; for example 5 to 60g of protein and 10mg to 5g of urolithin; for example 5 to 40g of protein and 20mg to 2500mg; for example 5 to 40g of protein and 50mg to 1000mg of urolithin; for example 10 to 30g of protein and 50mg to 1000mg of urolithin; for example 10 to 30g of protein and 50mg to 500mg of urolithin; for example 10 to 30g of protein and 100mg to 500mg of urolithin. The compositions preferably further contain carbohydrates, vitamins and minerals. Representative compositions are shown in Tables 1, 2 and 3:

Table 1: Representative powder composition:

Composition	Per 100 g
-------------	-----------

Protein	10-80 g
Carbohydrates	20-40 g
Fat	0-20 g
Polyunsaturated Fatty Acids	0-5g
Fiber	0-5 g
Vitamins	0-100% of respective RDAs
Minerals	0-100% of respective RDAs
Folic Acid	0-1000 ug
Niacin	0-100 mg
Creatine	0-20g
Urolithin A	0.025-5g

In a bulk powder, the protein preferably makes up 20% or more of the powder by weight, for example 20% or more by weight, for example 30% or more by weight, for example 40% or more by weight of the powder. For example the protein may make up 20-90% by weight of the powder, for example 30-80% by weight of the powder, for example 40-80% by weight of the powder, for example 40-70%, for example 60-80% by weight, for example 60-70% by weight of the powder.

Bulk powder is generally provided with instructions informing the subject how much of the powder to use for one serving. For example the bulk powder may be supplied in a container accompanied by a scoop of the necessary size to enable the correct amount of powder to be measured out. Powder may be taken neat, mixed with food, or added to water to make a drink.

Table 2: Representative drink composition:

Composition	Per 100 mL
Protein (for example 100% Hydrolyzed Whey)	5-15 g
Carbohydrates	1-20 g
Fat	0-9.5g
Polyunsaturated Fatty Acids	0-2.5 g
Fiber	0-2 g
Vitamins	0-100% of respective RDAs
Minerals	0-100% of respective RDAs
Folic Acid	0-500 ug
Niacin	0-20 mg
Urolithin A	10-2000 mg

Table 3: Representative bar composition:

Composition	Per bar of 35g
Protein	6.7 g
Carbohydrates	10-20g
Fat	0-15 g
Polyunsaturated Fatty Acids	2-6 g
Fiber	0-5 g
Niacin	0-100 mg
Vitamins	0-100% of respective RDAs
Minerals	0-100% of respective RDAs
L-Carnitine	0-500 mg
Urolithin A	10-2000 mg

5 Table 4: Representative yoghurt composition:

Composition	Per 100g Yoghurt
Protein	2-15 g
Carbohydrates	3-20g
Fat	0-12 g
Calcium	5-20% RDA
Fiber	0-4 g
Vitamins	0-100% of respective RDAs
Minerals	0-100% of respective RDAs
Live Cultures	<i>S. thermophilus</i> , <i>L. bulgaricus</i> , <i>L. acidophilus</i> , <i>L. lactis</i>
Urolithin A	10-2000 mg

The composition of the invention can be taken as a single treatment or, more commonly, as a series of treatments. In one example, a subject takes a dose before or after exercise. For a subject who is not able to exercise, a dose of the composition may, for example, be taken
10 once, twice or three times per day, or one, two, three, four, five or six times per week. It will also be appreciated that the effective dosage of the compound may increase or decrease over the course of a particular treatment.

Treatments:

The compositions of the invention find use in improving muscle performance, improving
15 muscle function, preventing a decline in muscle function, increasing muscle mass and/or

reducing muscle wasting. The improvement in muscle performance, improving or maintaining muscle function, the increase in muscle mass and/or reduction in muscle wasting may be as part of a medical treatment, or it may be for personal preference (“lifestyle”) or cosmetic reasons. The compositions of the invention can be for use as a medicament. The compositions can be used as a dietary supplement, as a functional food, functional beverage, or as a medical food.

Most muscle enhancement regimes require the subject to exercise as well as taking a muscle enhancement composition. There is evidence that the combination of protein and urolithin of the invention is effective in enhancing muscle growth, muscle strength, muscle endurance and muscle function in the absence of exercise, or with less exercise being necessary.

The compositions find use in the treatment of both diseases and disease states. The compositions find use in the management normal physiological function in healthy individuals of conditions characterised by poor physical performance, impaired endurance capacity, and impaired muscle function. Compositions of the invention may improve physical performance in individuals with a disease, including young and elderly individuals. Compositions of the invention may improve physical performance, for example, short-term performance or long-term performance in healthy individuals, including athletes, non-athletic individuals, sedentary individuals and the elderly. This improvement of performance may be measured by the time spent to walk or run a certain distance (for example, an improved performance during the 6 minute walk test (MWT)), an improved time to run a certain distance, an improved IPAQ score on the international physical activity questionnaire, an increased number of chair-stands in a certain time, or another test designed to measure physical performance.

Compositions of the invention further provide for the improvement of endurance capacity.

The endurance capacity refers to the time to fatigue when exercising at a constant workload, generally at an intensity $<80\%$ VO_2max . Compositions of the invention may improve endurance capacity in individuals with a disease, including young and elderly individuals.

Compositions of the invention may improve endurance capacity in healthy individuals, including athletes, non-athletic individuals, sedentary individuals and the elderly. The

invention provides for a method of increasing the time to fatigue while performing a specific activity, for example, fitness training, walking, running, swimming, or cycling. This improvement of endurance capacity may be assessed with objective measurements (for

example, speed, oxygen consumption or heart rate) or it can be self-reported measurements (for example, using a validated questionnaire).

The invention further provides a composition to improve, maintain or reduce the loss of muscle function. Compositions of the invention may improve, maintain or reduce the loss of muscle function in individuals with a disease, including young and elderly individuals.

Compositions of the invention may improve, maintain or reduce the loss of muscle function in healthy individuals, including athletes, non-athletic individuals, sedentary individuals and the elderly. For example, compositions of the invention may improve, maintain or reduce the loss of muscle function in frail or pre-frail individuals. For example, compositions of the invention may increase muscle strength as evidenced by the improvement of performing a physical activity, such as an exercise, for example, increased ability to lift weights or increased hand grip strength. Also, compositions of the invention may improve muscle structure, for example by increasing or maintaining muscle mass in conditions of normal muscle function, declining muscle function or impaired muscle function.

This invention further provides a composition to improve the physical performance or endurance capacity as perceived by the individual. For example, by the reduction of in perceived exertion or effort during exercise or an activity as determined using a self-reported questionnaire.

Medical treatments:

The composition of the invention can be for use as a medicament. The compositions of the invention find use in the treatment of muscle-related pathological conditions. Accordingly, the invention provides a composition of the invention for use in the treatment of a muscle-related pathological condition. The invention also provides a method of treating a muscle-related pathological condition in a subject comprising administering to the subject an effective amount of a composition of the invention. Muscle-related pathological conditions include both conditions impacting generally healthy individuals as well as pathological conditions. Such muscle conditions found in healthy people or people affected by a disease include musculoskeletal diseases or disorders; cachexia; muscle wasting; myopathies; age-related decline in muscle function; pre-frailty; frailty; neuromuscular diseases, such as Duchenne muscular dystrophy and other dystrophies; sarcopenia, for example, acute sarcopenia; muscle atrophy and/or cachexia, for example muscle atrophy and/or cachexia

associated with burns, bed rest, limb immobilization, or major thoracic, abdominal, and/or orthopedic surgery; and muscle degenerative disease.

Examples of age-related disease conditions that may be treated with compositions of the invention include sarcopenia and muscle wasting.

- 5 It has been reported in WO2014/111580 that Urolithin B (but not Urolithin A) increased the mean diameter of myotubes *in vitro*. The effect was not seen with Urolithin A.

Muscle Performance:

- The composition of the invention is useful in enhancing muscle performance. The invention thus provides a composition of the invention for use in enhancing muscle performance. The
10 invention also provides a method of enhancing muscle performance by administering to a subject an effective amount of a composition of the invention. Administration can be self-administration.

- The enhanced muscle performance may be one or more improved muscle function, reduced decline in muscle function, improved muscle strength, improved muscle endurance and
15 improved muscle recovery.

The composition of the invention can thus be used in a method of improving physical endurance (e.g., ability to perform a physical task such as exercise, physical labor, sports activities), inhibiting or retarding physical fatigue, enhancing working capacity and endurance, and reducing muscle fatigue.

- 20 Improved muscle function can be particularly beneficial in elderly subjects with reduced muscle function as a result of an age-related condition. For example, a subject who may benefit from improved muscle function may experience a decline in muscle function which then leads to pre-frailty and frailty. Such subjects may not necessarily experience muscle wastage in addition to their decline in muscle function. Some subjects do experience both
25 muscle wasting and a decline in muscle function, for example subjects with sarcopenia. The composition of the invention may be used in enhancing muscle performance by administering a composition of the invention to a subject who is frail or pre-frail.

- Muscle performance may be sports performance, which is to say the ability of an athlete's muscles to perform when participating in sports activities. Enhanced sports performance,
30 strength, speed, and endurance are measured by an increase in muscular contraction strength,

increase in amplitude of muscle contraction, or shortening of muscle reaction time between stimulation and contraction. Athlete refers to an individual who participates in sports at any level and who seeks to achieve an improved level of strength, speed, or endurance in their performance, such as, for example, body builders, bicyclists, long distance runners, and short distance runners. Enhanced sports performance is manifested by the ability to overcome muscle fatigue, ability to maintain activity for longer periods of time, and have a more effective workout.


Examples

The following Examples illustrate the invention.

Compounds

Urolithin A was prepared as follows:

Urolithin A (**4**) was prepared in two steps starting from bromide **1** and resorcinol **2**. The pure compound was obtained as a pale yellow powder.



Step 1:

A mixture of 2-bromo-5-methoxybenzoic acid **1** (27.6 g; 119 mmol; 1.0 eq.), resorcinol **2** (26.3 g; 239 mmol; 2.0 eq.) and sodium hydroxide (10.5 g; 263 mmol; 2.2 eq.) in water (120 mL) was heated under reflux for 1 hour. A 5 % aqueous solution of copper sulphate (3.88 g of $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$ in 50 mL water; 15.5 mmol; 0.1 eq.) was then added and the mixture was refluxed for an additional 30 minutes. The mixture was allowed to cool to room temperature and the solid was filtered on a Büchner filter. The residue was washed with cold water to give a pale red solid which was triturated in hot MeOH. The suspension was left overnight at 4 °C. The resultant precipitate was filtered and washed with cold MeOH to yield the title compound **3** as a pale brown solid.

Step 2:

To a suspension of **3** (10.0 g; 41 mmol; 1.0 eq.) in dry dichloromethane (100 mL) was added dropwise at 0 °C a 1 M solution of boron tribromide in dry dichloromethane (11.93 mL of

pure BBr₃ in 110 mL of anhydrous dichloromethane; 124 mmol; 3.0 eq.). The mixture was left at 0 °C for 1 hour and was then allowed to warm up to room temperature. The solution was stirred at that temperature for 17 hours. Then ice was added thoroughly to the mixture. The yellow precipitate was filtered and washed with cold water to give a yellow solid which was heated to reflux in acetic acid for 3 hours. The hot solution was filtered quickly and the precipitate was washed with acetic acid, then with diethyl ether to yield the title compound **4** as a yellow solid. ¹H and ¹³C NMR were in accordance with the structure of **4**.

Example 1: Experimental trial on muscle function with Optimal Protein with or without Urolithin A supplementation

a) Optimal Protein Diet containing Urolithin A

To mimic age related muscle loss in humans, an experimental animal model was setup in old mice. The control diet designed to be used in the experimental study was a high protein diet that would be analogous to a clinically recommended high protein diet in elderly humans at risk of muscle loss. The diet was formulated to improve the performance of old animals that consume them. The high protein diet has a better balance of essential nutrients (amino acids, vitamins and minerals) and is supplemented with 200gr casein per kg of diet, instead of 140gr of casein in the regular maintenance diet (Reeves PG, J. Nutr. 1997, 127(5) 838S-841S).

Casein proteins are nutritionally complete, highly digestible proteins with high amino scores that contain all amino acids in amounts sufficient to support maintenance of all body proteins.

The components of the experimental high protein diet are shown in Table 5 below. Urolithin A was admixed to the high protein diet at a level of 0.57g per kg of diet. Both the diets were isocaloric.

Table 5: Composition of feeds:

Diet	High Protein Diet		High Protein Diet + Urolithin A	
	gm%	Kcal%	gm%	Kcal%
Protein	20	20	20	20
Carbohydrate	64	64	64	64
Fat	7	16	7	16
Total		100		100
Kcal/gm	4.0		4.0	
Ingredient	Gm	Kcal	Gm	Kcal
Casein protein	200	800	200	800

L-Cystine	3	12	3	12
Corn Starch	397.486	1590	397.486	1590
Maltodextrin 10	132	528	132	528
Sucrose	100	400	100	400
Cellulose, BW200	50	0	50	0
Soybean Oil	70	630	70	630
t-butylhydroquinone	0.014	0	0.014	0
Mineral Mix S 100022G	35	0	35	0
Vitamin Mix V10037	10	40	10	40
Choline Bitartrate	2.5	0	2.5	0
Urolithin A	0	0	0.57	0
FD&C Red Dye #40	0	0	0.05	0
Total	1000	4000	1000.62	4000

b) Protein in combination with urolithin A in diet is better compared with protein alone

22 month old C57BL/6J old mice that would be equivalent to a 65yr to 75yr old elderly human were treated with either the high protein diet alone or a high protein diet containing Urolithin A. Both the diets contained in addition to the casein protein, essential nutrient for optimal muscle function such as L-cysteine and vitamins and mineral mix. The level of Urolithin A in the diet results in a dose of 50 mg/kg/day delivered to mice.

Before and following 6 weeks of treatment with Urolithin A and protein combination, mice were tested for their endurance on treadmill. Endurance test was performed during the day.

- 10 Mice had access to food and water before and after the exercise. The test was performed using treadmill placed in front of a motivational grid (Panlab, Barcelona, Spain). Mice were acclimatized to the treadmill for 5 minutes set on to 0.3mA. The exercise started with 12 min running at a speed of 15 cm/second. Then speed increased by 3 cm/second every 12 minutes. The animal was removed from the treadmill when touching the back grid more than 5 times
- 15 per minute during two consecutive minutes. The total distance and running time after 6 weeks of treatment was normalized over the basal values. Neither group of mice were subjected to an exercise regimen prior to the endurance test. Mice were maintained in standard animal housing throughout the course of the study.

The results of the test are shown in Figure 1. It was strikingly observed that the mice fed with Urolithin A in addition to the high protein diet displayed an increase of 42% in running distance and 28% in running time as compared to mice receiving only a protein diet. This improvement in running endurance occurs in the absence of an exercise regimen. Statistical significance was determined using a Fischer t test. ***:p<0.001.

These results demonstrate the ability of Urolithin A combined with an optimal protein diet to increase muscle function in aged animals. It is anticipated that similar improvements would be observed with urolithin A in elderly humans where as standard of care today, only an optimal protein diet is recommended.

Example 2: Powder formula composition targeting healthy aging and age-related muscle loss containing high protein and urolithin A

Table 6:

Composition	Per 100 g (single serving)
Protein	51.5 g
Whey	(45 g)
Leucine	(3.5 g)
Isoleucine	(1.5 g)
L-Arginine	(1.5 g)
Carbohydrates	28 g
Fat	11.5 g
Polyunsaturated Fatty Acids	2.5 g
Fiber	2.0 g
Vitamin A	50 ug
Vitamin D3	50 ug
Vitamin E	20 mg
Vitamin C	100 mg
Vitamin B6	2 mg
Vitamin B12	10 ug
Folic Acid	500 ug
Niacin	50 mg
Zinc	5 mg
Calcium	100 mg
Selenium	40 ug
Iron	20 mg
Magnesium	100 mg

Creatine	1.5 g
Urolithin A	1.0 g

The composition with the nutrient profile shown in Table 6 is given to a subject to counteract age related muscle loss.

5 Example 3: An enteral nutrition liquid composition targeting immobilized subject in intensive care or hospital settings containing protein and urolithin A

Table 7:

Composition	Per 100 mL
Protein	10.5 g
100% Hydrolyzed Whey	
Carbohydrates	15.4 g
Fat	9.9 g
Polyunsaturated Fatty Acids	3.8 g
Fiber	0 g
Vitamin A	170 ug
Vitamin D3	2 ug
Vitamin E	3 mg
Vitamin C	20 mg
Vitamin B6	0.3 mg
Vitamin B12	1 ug
Folic Acid	50 ug
Niacin	5 mg
Zinc	1.5 mg
Calcium	100 mg
Selenium	10 ug
Iron	2 mg
Magnesium	40 mg
Urolithin A	100 mg

The drink composition with the nutrient profile shown in Table 7 is given to an immobilised subject in intensive care or a hospital setting.

10

Example 4: A cereal bar composition targeting an active athlete for optimal muscle function during endurance training containing protein and urolithin A

Table 8:

Composition	Per bar of 35g
Energy	600 kcal
Protein	6.7 g
Carbohydrates	17.2 g
Fat	7.6 g
Polyunsaturated Fatty Acids	3.8 g
Fiber	1.6 g
Niacin	25 mg
Zinc	2.5 mg
Calcium	180 mg
Sodium	25 mg
Potassium	60 mg
Magnesium	80 mg
L-Carnitine	200 mg
Urolithin A	250 mg

5

The bar composition with the nutrient profile shown in Table 8 is given to an active athlete for optimal muscle function during endurance.

Example 5: A yogurt composition

10 Table 9:

Nutrition value:	per 100 g	
Fat	0,2 g	
Carbohydrates	3,7 g	
Protein	9,8 g	
Vitamin B2	0,18 mg	13 % of RDA
Calcium	95 mg	12 % of RDA
Phosphorus	170 mg	24 % of RDA
Live Active Cultures		
Urolithin A	250mg, 500mg or 1000 mg	

Example 6: Example 3: Effects of particle size on urolithin A bioavailability

The particle size of urolithin A was reduced in a controlled manner using an MC50 Spiral Jetmill, using filtered nitrogen, with a feed rate of 240 g/hr, a Venturi pressure of 12 bar, and a mill pressure of 12 bar. Different particle size distributions of urolithin A were determined on a Malvern Particle Size analyzer (Malvern Instruments, UK). Three samples were evaluated in depth. Sample #1 of urolithin A had a particle size distribution of D₁₀ of 1.03 µm, a D₅₀ of 53.4 µm and a D₉₀ of 365 µm. Sample #2 of urolithin A had a particle size of D₁₀ of 0.272 µm, a D₅₀ of 2.17 µm and a D₉₀ of 6.84 µm. Sample #3 of urolithin A had a particle size of D₁₀ of 0.597 µm, a D₅₀ of 5.67 µm and a D₉₀ of 40.1 µm

Table 10: Particle size determinations for urolithin A before and after micronization

Urolithin A	D ₁₀	D ₅₀	D ₉₀
Sample #1	1.03 µm	53.4 µm	365 µm
Sample #2	0.272 µm	2.17 µm	6.84 µm
Sample #3	0.597 µm	5.67 µm	40.1 µm

To demonstrate the impact of particular particle size distributions on bioavailability, male Sprague-Dawley rats were fasted overnight and then administered by oral gavage with either sample #1, sample #2, or sample #3 of urolithin A suspended in 15% DMSO, 85% (0.5% methyl cellulose, 0.25% Tween 80 in Water), at a dose corresponding to 25 mg/kg/day.

Table 11: Design of the bioavailability study.

Test Article	Animals N=	Dose (mg/kg)	Dosing Solution Conc. (mg/mL)	Dosing Volume (mL/kg)	Vehicle	Plasma Sampling Time Points
Urolithin A Sample #1	3	25	5	5	15% DMSO 85% (0.5% methyl cellulose/0.25% Tween 80 in Water)	Pre-dose, 0.25, 0.5, 1, 2, 4, 8, 12, 24 and 48 hours
Urolithin A Sample #2	3	25	5	5	15% DMSO 85% (0.5% methyl cellulose/0.25% Tween 80 in Water)	Pre-dose, 0.25, 0.5, 1, 2, 4, 8, 12, 24 and 48 hours

Urolithin A Sample #3	3	25	5	5	15% DMSO 85% (0.5% methyl cellulose/0.25% Tween 80 in Water)	Pre-dose, 0.25, 0.5, 1, 2, 4, 8, 12, 24 and 48 hours
--------------------------	---	----	---	---	---	--

The solutions for gavage were prepared as follows: 35mg of urolithin A powder was diluted in 7 ml of 15% DMSO, 0.5% methyl cellulose/0.25% Tween 80 in water, resulting in a 5 mg/ml fine suspension. The DMSO was obtained from BDH, methyl cellulose was obtained from Sigma, and Tween 80 was obtained from Sigma-Aldrich.

Blood was collected from the rats by jugular vein cannulation at different time points, and urolithin A was quantified in the plasma to determine its pharmacokinetic profile following oral gavage of Sample #1, Sample #2, and Sample #3. The study was repeated on three rats for each sample. Figure 3 shows the resulting pharmacokinetic profiles of urolithin sample #1 and sample #2. The data collected is summarized below in Table 12.

Table 12: Results of bioavailability study for urolithin A with different particle sizes

Parameter	Sample #1	Sample #2	Sample #3
C _{max} (ng/ml)	17.1	37	17.4
Relative C _{max} (% of non-micronized C _{max})	100%	216%	102%
T _{max} (h)	0.25	0.33	0.50
AUC	75.1	96.2	94.8
Relative bioavailability (% AUC of non-micronized urolithin A)	100%	128%	126%

Urolithin A showed an increase in bioavailability when particle size was reduced below a D₉₀ of 50µm. For Sample #2 there was a relative increase in the C_{max} of 116% (2.16-fold increase) and an increase in the AUC of 28%, in comparison to Sample #1. For Sample #3 there was a relative increase in the AUC of 26%.

These results demonstrate that urolithin A preparations having a particle size of D₉₀ < 50µm consistently shows a higher bioavailability when compared to urolithin A preparations having a particle size of D₉₀ > 300µm. Additionally, it was observed that urolithin A preparations

having particle size of $D_{90} < 20\mu\text{m}$ were especially advantageous and enabled achieving dramatically higher peak urolithin A blood levels as the C_{max} achieved more than doubled.

CLAIMS

1. A composition comprising:
 - a) a source of protein and
 - b) urolithin A;wherein the source of protein is the range 5 to 90g and the urolithin A is in the range 20mg to 2500mg.
2. The composition as claimed in claim 1 wherein the urolithin A has a D₅₀ size in the range 0.5 to 50 µm and a D₉₀ size in the range 5 to 100 µm.
3. The composition as claimed in claim 1 or claim 2 wherein the protein makes up at least 15% w/w of the composition.
4. The composition as claimed in claim 1 or claim 2 wherein the protein makes up 20-99% w/w of the composition.
5. The composition as claimed in claim 4 wherein the protein makes up 30-80% w/w of the composition.
6. The composition as claimed in any one of claims 1 to 5 wherein the protein makes up 5 to 40g of the composition.
7. The composition as claimed in any one of claims 1 to 5 wherein the composition is a unit dose composition and the protein makes up 10 to 40g of the unit dose composition.
8. The composition as claimed in any one of claims 1 to 5 wherein the the composition is a unit dose composition and protein makes up 10-30g of the unit dose composition.
9. A composition comprising:

- a) a source of protein and
- b) urolithin A;

wherein the composition is a yogurt and the protein makes up 2 to 15g protein per 100g and the urolithin A is in the range 10mg to 2000mg.

10. The composition as claimed in any one of claims 1 to 9 wherein the urolithin A makes up 0.05 to 10% w/w of the composition.

11. The composition as claimed in any one of claims 1 to 9 wherein the urolithin A makes up 0.2 to 5% w/w of the composition.

12. The composition as claimed in any one of claims 1 to 11 wherein the weight ratio between the protein component and the urolithin A is in the range 2:1 to 1000:1.

13. The composition of any one of claims 1 to 12 wherein the urolithin A is in the range 250mg to 1500mg.

14. The composition as claimed in claim 13 wherein the urolithin A is in the range 250mg to 1000mg.

15. The composition as claimed in any one of claims 1 to 14 wherein the protein is hydrolyzed, partially hydrolyzed or non-hydrolyzed protein.

16. The composition as claimed in claim 15 wherein the protein is derived from a source selected from milk, animal, cereal or vegetable sources.

17. The composition as claimed in claim 16 wherein the milk source is casein or whey.

18. The composition as claimed in claim 16 wherein the animal source is meat or fish.

19. The composition as claimed in claim 16 wherein the cereal source is rice or corn.
20. The composition as claimed in claim 16 wherein the vegetable source is soy or pea.
21. A composition as claimed in any one of claims 16 to 20 wherein the protein is selected from intact pea protein, intact pea protein isolates, intact pea protein concentrates, milk protein isolates, milk protein concentrates, casein protein isolates, casein protein concentrates, whey protein concentrates, whey protein isolates, sodium or calcium caseinates, whole cow's milk, partially or completely defatted milk, soy protein isolates and soy protein concentrates, and combinations thereof.
22. The composition as claimed in any one of claims 1 to 8, wherein the composition is in the form of a yoghurt.
23. The composition as claimed in any one of claims 1 to 8 wherein the composition is in the form of a solid, a semi-solid, capsule, a powder or a liquid.
24. The composition as claimed in any one of claims 1 to 23 wherein the composition is a powder and the protein makes up 40-80% by weight of the powder.
25. The composition as claimed in claim 24 wherein the composition is in the form of a solid, or a capsule.
26. The composition as claimed in claim 23 wherein the composition is in the form of a solid is a tablet or a bar.
27. The composition as claimed in claim 23 wherein the composition is in the form of a semi-solid is a softgel, capsule, or dragee.

28. The composition as claimed in claim 27 wherein the composition is in the form of a capsule and is a hard capsule.
29. The composition as claimed in claim 23 wherein the composition is in the form of a liquid or an emulsion.
30. The composition as claimed in any one of claims 1 to 29 for use as a medicament, dietary supplement, functional food, functional beverage or medical food.
31. The composition for use as claimed in claim 30 for use as a medicament.
32. The composition for use as claimed in claim 31 for use as a medicament for the treatment of a muscle-related pathological condition.
33. The composition for use as claimed in claim 32 wherein the muscle-related pathological condition is selected from musculoskeletal diseases or disorders; muscle wasting; myopathies; neuromuscular diseases, sarcopenia; and muscle atrophy or cachexia, or a combination of atrophy and cachexia.
34. The composition for use as claimed in claim 32 wherein the muscle-related pathological condition is neuromuscular diseases wherein the neuromuscular diseases are selected from Duchenne muscular dystrophy and other dystrophies.
35. The composition for use as claimed in claim 33 wherein the sarcopenia is acute sarcopenia.
36. The composition for use as claimed in claim 33 wherein the muscle atrophy or cachexia or a combination of atrophy and cachexia is selected from muscle atrophy or cachexia associated with burns, bed rest, limb immobilization, or major thoracic, abdominal, and/or orthopedic surgery.

37. Use of the composition as claimed in any one of claims 1 to 29 as a medicament, dietary supplement, functional food, functional beverage or medical food.
38. The use as claimed in claim 37 wherein the use is as a medicament.
39. The use as claimed in claim 38 wherein the medicament is for the treatment of a muscle-related pathological condition.
40. The use as claimed in claim 39 wherein the muscle-related pathological condition is selected from musculoskeletal diseases or disorders; muscle wasting; myopathies; neuromuscular diseases, sarcopenia; and muscle atrophy or cachexia, or a combination of atrophy and cachexia.
41. The use as claimed in claim 39 wherein the muscle-related pathological condition is neuromuscular diseases wherein the neuromuscular diseases are selected from Duchenne muscular dystrophy and other dystrophies.
42. The use as claimed in claim 40 wherein the sarcopenia is acute sarcopenia.
43. The use as claimed in claim 40 wherein the muscle atrophy or cachexia or a combination of atrophy and cachexia is selected from muscle atrophy or cachexia associated with burns, bed rest, limb immobilization, or major thoracic, abdominal, and/or orthopedic surgery.
44. Use of an effective amount of a composition as claimed in any one of claims 1 to 29 for enhancing muscle performance, wherein the composition is for administration to a subject.

45. The use as claimed in claim 44 wherein the enhanced muscle performance is selected from one or more of improved muscle function, reduced decline in muscle function, improved muscle strength, improved muscle endurance and improved muscle recovery.
46. The use as claimed in claim 44 or claim 45, wherein the subject suffers age-related decline in muscle function, age-related sarcopenia, age-related muscle wasting, physical fatigue, muscle fatigue, and/or is frail or pre-frail.
47. The use as claimed in claim 46 wherein the subject suffers physical fatigue or muscle fatigue.
48. The use of claim 46 or claim 47, wherein the subject is frail or pre-frail.
49. Use of an effective amount of a composition as claimed in any one of claims 1 to 29 for improving, maintaining or reducing the loss of muscle function, wherein the composition is for administration to a subject.
50. The use as claimed in claim 49 wherein the loss in muscle function is in healthy individuals, including athletes, non-athletic individuals, sedentary individuals and the elderly.
51. Use of an effective amount of a composition as claimed in any one of claims 1 to 29 for improving endurance capacity, wherein the composition is for administration to a subject.
52. The use as claimed in claim 51 wherein the improvement in endurance capacity is in healthy individuals, including athletes, non-athletic individuals, sedentary individuals and the elderly.
53. Use of an effective amount of a composition as claimed in any one of claims 1 to 29 for enhancing physical performance, wherein the composition is for administration to a subject.

54. The use as claimed in claim 53 wherein the physical performance is in a healthy individual or in the elderly.
55. Use of an effective amount of a composition as claimed in any one of claims 1 to 29 for increasing muscle strength, increasing or maintaining muscle mass or muscle recovery, wherein the composition is for administration to a subject.
56. Use of an effective amount of a composition as claimed in any one of claims 1 to 29 for improving physical endurance, wherein the composition is for administration to a subject.
57. Use of an effective amount of a composition as claimed in any one of claims 1 to 29 for inhibiting or retarding physical fatigue, enhancing working capacity and endurance and reducing muscle fatigue, wherein the composition is for administration to a subject.
58. Use of an effective amount of a composition as claimed in any one of claims 1 to 29 for enhancing sports performance, wherein the composition is for administration to a subject.
59. A composition as defined in claim 1 for use in the treatment of a muscle-related pathological condition in a subject, or enhancing muscle function in a subject, wherein an effective amount of urolithin A and an effective amount of a source of protein are for administration to the subject, wherein the administration of the urolithin A and the administration of the protein are at the same time or separated by a time interval.
60. Use of a composition as defined in claim 1 in the treatment of a muscle-related pathological condition in a subject, or enhancing muscle function in a subject, wherein an effective amount of urolithin A and an effective amount of a source of protein are for administration to the subject, wherein the administration of the urolithin A and the administration of the protein are at the same time or separated by a time interval.

61. A kit comprising the compound for use of claim 59 or claim 60.
62. An effective amount of a composition as claimed in any one of claims 1 to 29 for use in enhancing muscle performance, wherein the composition is for administration to a subject.
63. The composition for use as claimed in claim 62 wherein the enhanced muscle performance is selected from one or more of improved muscle function, reduced decline in muscle function, improved muscle strength, improved muscle endurance and improved muscle recovery.
64. The composition for use as claimed in claim 62 or claim 63, wherein the subject suffers age-related decline in muscle function, age-related sarcopenia, age-related muscle wasting, physical fatigue, muscle fatigue, and/or is frail or pre-frail.
65. The composition for use as claimed in claim 64 wherein the subject suffers physical fatigue or muscle fatigue.
66. The composition for use of claim 64 or claim 65, wherein the subject is frail or pre-frail.
67. An effective amount of a composition as claimed in any one of claims 1 to 29 for use in improving, maintaining or reducing the loss of muscle function, wherein the composition is for administration to a subject.
68. The composition for use as claimed in claim 67 wherein the loss in muscle function is in healthy individuals, including athletes, non-athletic individuals, sedentary individuals and the elderly.
69. An effective amount of a composition as claimed in any one of claims 1 to 29 for use in improving endurance capacity, wherein the composition is for administration to a subject.

70. The composition for use as claimed in claim 69 wherein the improvement in endurance capacity is in healthy individuals, including athletes, non-athletic individuals, sedentary individuals and the elderly.

71. An effective amount of a composition as claimed in any one of claims 1 to 29 for use in enhancing physical performance, wherein the composition is for administration to a subject.

72. The composition for use as claimed in claim 71 wherein the physical performance is in a healthy individual or in the elderly.


73. An effective amount of a composition as claimed in any one of claims 1 to 29 for use in increasing muscle strength, increasing or maintaining muscle mass or muscle recovery, wherein the composition is for administration to a subject.

74. An effective amount of a composition as claimed in any one of claims 1 to 29 for use in improving physical endurance, wherein the composition is for administration to a subject.

75. An effective amount of a composition as claimed in any one of claims 1 to 29 for use in inhibiting or retarding physical fatigue, enhancing working capacity and endurance and reducing muscle fatigue, wherein the composition is for administration to a subject.

76. An effective amount of a composition as claimed in any one of claims 1 to 29 for use in enhancing sports performance, wherein the composition is for administration to a subject.

Figure 1:



5 Figure 2:

