



(51) International Patent Classification:

C07H 21/04 (2006.01) C12N 15/86 (2006.01)

A61K 35/30 (2015.01)

(21) International Application Number:

PCT/US2020/015183

(22) International Filing Date:

27 January 2020 (27.01.2020)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/801,483 05 February 2019 (05.02.2019) US

62/823,281 25 March 2019 (25.03.2019) US

62/916,477 17 October 2019 (17.10.2019) US

(71) Applicants: **THE BROAD INSTITUTE, INC.** [US/US]; 415 Main Street, Cambridge, Massachusetts 02142 (US). **PRESIDENT AND FELLOWS OF HARVARD COLLEGE** [US/US]; 17 Quincy Street, Cambridge, Massachusetts 02138 (US). **NEW YORK UNIVERSITY** [US/US]; 70 Washington Square South, New York, New York 10012 (US).

(72) Inventors: **DIMIDSCHSTEIN, Jordane**; c/o The Broad Institute, Inc., 415 Main Street, Cambridge, Massachusetts 02138 (US). **FISHELL, Gordon**; c/o President and Fellows of Harvard College, 17 Quincy Street, Cambridge, Massachusetts 02138 (US). **DEVINSKY, Orrin**; 4 East 70th Street, 9A, New York, New York 10021 (US).

(74) Agent: **HUNTER-ENSOR, Melissa**; c/o Greenberg Traurig, LLP, One International Place, Boston, Massachusetts 0210 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA,

SC, SD, SE, SG, SK, SL, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, WS, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

- with international search report (Art. 21(3))
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))

(54) Title: INTERNEURON-SPECIFIC THERAPEUTICS FOR NORMALIZING NEURONAL CELL EXCITABILITY AND TREATING DRAVET SYNDROME

(57) Abstract: Provided are therapeutic virus vectors, particularly, recombinant adeno-associated virus (rAAV) vectors, designed to contain an enhancer sequence that specifically restricts expression of an effector gene (e.g., an SCN1A-encoding polynucleotide, Gq-DREADD-encoding polynucleotide, or PSAM-encoding polynucleotide) contained in the vector to PV-expressing GABAergic interneuron or to neuron cell populations in the brain. The rAAV vectors, compositions and methods thereof are useful for treating subjects afflicted with neuropathologies, seizures, pharmacologically-intractable forms of epilepsy including Dravet syndrome (DS), a form of infantile epilepsy associated with severe seizures, cognitive impairment and premature death, as the cause of DS involves loss of function of a sodium channel encoded by the SCN1A gene. The described vectors restore expression of effector genes to the appropriate interneuron or neuron cell populations with specificity and sensitivity, advantageously to address the root cause of the disease by restoring the excitation-inhibition balance by means of gene-therapy (with SCN1A) or pharmacogenetics.



INTERNEURON-SPECIFIC THERAPEUTICS FOR NORMALIZING NEURONAL CELL EXCITABILITY AND TREATING DRAVET SYNDROME

5

CROSS REFERENCE TO RELATED APPLICATIONS

This application is an International PCT application which claims priority to and benefit of U.S. Provisional Application No. 62/801,483, filed on February 5, 2019, U.S. Provisional Application No. 62/823,281, filed on March 25, 2019, and
10 U.S. Provisional Application No. 62/916,477, filed on October 17, 2019, the contents of each of which are incorporated by reference herein in its entirety.

STATEMENT REGARDING FEDERALLY SPONSORED RESEARCH

The invention was made with government support under grant number MH111529 awarded by the National Institutes of Health. The government has certain
15 rights in the invention.

BACKGROUND

A delicate balance exists between excitation and inhibition that must be carefully maintained for proper functioning of brain circuits and the activities of the neuronal cells that function within these circuits. An alteration, defect, or disruption
20 in the balance of excitation versus inhibition in the brain circuitry was shown to result in a number of neurological, neurogenetic, or neurodegenerative diseases and disorders. Moreover, the lack of proper cortical interneuron function was linked to neurodevelopment and neurological diseases and disorders.

Abnormal or aberrant interneuron function and activity may be a consequence
25 of a deviation from the course of interneuron development (e.g., aberrant fate specification during embryonic development due to genetic mutation) or acute insult (e.g., stroke, concussion). Aberrant GABAergic neurotransmission and alterations in inhibitory cortical circuits may cause and induce the clinical features and symptoms, e.g., seizures and epilepsy, that afflict patients having serious neurological diseases
30 and disorders, such as Dravet syndrome (DS), a pharmaco-resistant form of infantile epilepsy associated with cognitive impairment and premature death.

A dearth of therapeutic compositions and methods capable of modulating the activity of GABAergic interneurons, or other cortical neurons, with specificity and sensitivity severely impedes the ability of the medical community to alleviate the seizures in a wide variety of cases of epilepsy, particularly, in patients suffering from focal seizures and DS. Such compositions and methods are urgently needed to combat and treat the severe symptoms of these devastating conditions, as well as other neuropsychiatric diseases. The products, compositions and methods described herein are provided to address and meet these needs.

SUMMARY OF THE DISCLOSURE AND EMBODIMENTS

Featured herein are viral vectors, particularly, recombinant adeno-associated virus (rAAV) vectors, virus particles, and compositions and methods thereof. The rAAV vectors contain (are molecularly engineered to contain) at least one transgene (e.g., an effector gene such as the hM3Dq modified muscarinic receptor (Gq-DREADD), pharmacologically selective actuator molecule (PSAM), or a therapeutic gene such as *SCN1A*) and a specific regulatory polynucleotide sequence that restricts expression of the transgene to interneuron (IN) cells, particularly fast-spiking parvalbumin-expressing GABAergic interneurons (called PV-interneurons (PV INs) herein), or neuron cells of the brain cortex. In an embodiment, the specific regulatory polynucleotide sequence is derived from an enhancer sequence in the vicinity of the gene *SCN1A* and restricts expression of the transgene carried by the rAAV to fast-spiking parvalbumin-expressing GABAergic interneuron populations in brain. In an embodiment, the therapeutic gene is *SCN1A*. In a particular embodiment, the vector specifically transduces interneuron cells that are deficient or defective in the expression of the *SCN1A* gene that encodes the sodium chloride channel Nav1.1 in interneuron cells, in particular, cortical interneuron cells, and normalizes the excitability of the *SCN1A*-deficient or defective interneurons, thereby alleviating seizures and seizure symptoms in subjects suffering from Dravet syndrome (DS).

In an aspect, a suitable viral vector, e.g., a lentiviral vector or, in particular, a recombinant adeno-associated virus (rAAV) vector, is used to restrict expression of a transgene in GABA-ergic PV-expressing interneurons, or a pyramidal (PYR) neuron, or a vaso-intestinal peptide (VIP)-expressing cortical interneuron, in a mammal, and comprises an enhancer element polynucleotide (also called a regulatory element

herein) as described herein. In an embodiment, the enhancer element is provided in *cis*. In an embodiment, the regulatory element is S5E1, S5E2, S5E3, S5E4, S5E5, S5E6, S5E7, S5E8, S5E9., or S5E10, particularly human E1-E10, as described herein. In an embodiment, the enhancer element is human E11-E35 as described herein. In an embodiment, the enhancer element is S5E1 (E1). In an embodiment, the enhancer element is S5E2 (E2). In an embodiment, the enhancer element is S5E3 (E3). In an embodiment, the enhancer element is S5E4 (E4). In an embodiment, the enhancer element is E5. In an embodiment, the enhancer element is E6. In an embodiment, the enhancer element is E11. In an embodiment, the enhancer element is E14. In an embodiment, the enhancer element is E22. In an embodiment, the enhancer element is E29.

In an aspect, the viral vector or rAAV vector comprising the enhancer drives the expression of a copy of *SCN1A* in a transduced PV-expressing interneuron cell for the treatment and therapy of DS. In other embodiments, the vector or rAAV vector comprising the enhancer drives the expression of effector genes such as Gq-DREADD receptor or such as a pharmacologically selective actuator molecule (PSAM), an orthogonal ligand-gated ion channel, (and its pharmacologically selective effector molecule (PSEMs)) for chemogenetic modulation of PV-interneuron activity for the treatment of all forms of epilepsy, including focal and pharmacologically intractable epilepsy and for the treatment of DS.

In an aspect, a viral vector comprising a transgene polynucleotide sequence and an enhancer polynucleotide sequence that specifically restricts expression of the transgene in parvalbumin (PV)-expressing interneuron cells of the brain is provided.

In an aspect, a viral vector comprising an enhancer polynucleotide sequence specifically associated with *SCN1A* gene expression and a transgene polynucleotide sequence, wherein the enhancer sequence restricts expression of the transgene in PV-expressing interneuron cells of the brain is provided.

In an aspect, a suitable viral vector, e.g., a lentiviral vector or, in particular, a recombinant adeno-associated virus (rAAV) vector, is used to restrict expression of a transgene in GABA-ergic, vaso-intestinal peptide-expressing cortical interneuron cells (VIP cINs) of the brain in a mammal, in which an enhancer element as described

herein provided in *cis*. In an embodiment, the enhancer element is S5E6 as described herein.

In an aspect, a suitable viral vector, e.g., a lentiviral vector or, in particular, a recombinant adeno-associated virus (rAAV) vector, is used to restrict expression of a transgene in both GABA-ergic interneurons and glutamatergic pyramidal neurons in the brain in a mammal, in which an enhancer element as described herein provided in *cis*. In an embodiment, the pyramidal neurons are in cortical layer 5 of the brain in a mammal. In an embodiment, the enhancer element that restricts expression to pyramidal neurons is S5E5 as described herein.

In embodiments of the viral vector of the above-delineated aspects, the transgene is a reporter gene, a Designer receptor exclusively activated by designer drug (DREADD)-encoding gene, a pharmacologically selective actuator molecule (PSAM)-encoding gene, or a therapeutic gene, e.g., *SCN1A*. In an embodiment, the transgene is an *SCN1A* gene. In an embodiment, the transgene is a DREADD-encoding polynucleotide. In an embodiment, the DREADD-encoding polynucleotide is a Gq-DREADD-encoding gene that is activated by the chemogen clozapine-N4-oxide (CNO). In an embodiment, the transgene is a pharmacologically selective actuator molecule (PSAM)-encoding gene. In an embodiment, the expressed PSAM specifically interacts with a PSEM ligand. In an embodiment, the viral vector is recombinant adeno-associated virus (rAAV) vector.

In another aspect, a recombinant adeno-associated virus (rAAV) vector comprising an *SCN1A* transgene polynucleotide sequence, or a functional portion thereof, and an enhancer polynucleotide sequence that specifically restricts expression of the *SCN1A* transgene in interneuron cells of the brain is provided.

In embodiments of the viral vector or the rAAV vector of the above-delineated aspects, an Nav1.1 sodium channel encoded by the *SCN1A* transgene is functionally expressed in interneuron cells or neuron cells following transduction of the interneuron or neuron cells by the viral vector or rAAV vector. In embodiments of the viral vector or the rAAV vector of the above-delineated aspects, an Nav1.1 sodium channel encoded by the *SCN1A* transgene is functionally expressed in both GABA-ergic interneurons and glutamatergic pyramidal neurons following transduction of the interneuron or neuron cells by the viral vector or rAAV vector. In

an embodiment, the interneuron cells are GABAergic interneuron cells. In an embodiment, the interneuron cells are GABAergic interneuron cells within the brain telencephalon. In an embodiment, the GABAergic interneuron cells express parvalbumin (PV). In an embodiment, the neuron cells are pyramidal neuron cells, e.g., glutamatergic pyramidal cells in the brain cortex. In an embodiment of any one of the above-delineated aspects, the enhancer polynucleotide sequence comprises the polynucleotide sequence of the mouse enhancer element E1, E2, E3, E4, E5, E6, E7, E8, E9, or E10 (SEQ ID NOs: 5-14, respectively), or an ortholog, such as a human ortholog, thereof. In an embodiment, the enhancer polynucleotide sequence comprises the polynucleotide sequence of human enhancer element E1, E2, E3, E4, E5, E6, E7, E8, E9, or E10 (SEQ ID NOs: 15-24, respectively). In an embodiment, the viral vector or rAAV vector comprises an enhancer polynucleotide sequence comprising a nucleotide sequence which contains one or more regions of about 100 bp or longer having at least 75% or greater sequence identity to a polynucleotide sequence of a human enhancer element E1, E2, E3, E4, E5, E6, E7, E8, E9, or E10 (SEQ ID NOs: 15-24, respectively). In another embodiment, the viral vector or rAAV vector comprises an enhancer polynucleotide sequence comprising a nucleotide sequence which contains one or more regions of about 100 bp or longer having at least 75% or greater sequence identity to a polynucleotide sequence of human enhancer element E2 (SEQ ID NO: 16). In another embodiment, the viral vector or rAAV vector comprises an enhancer polynucleotide sequence comprising a nucleotide sequence which contains one or more regions of about 100 bp or longer having at least 75% or greater sequence identity to a polynucleotide sequence of human enhancer element E5 (SEQ ID NO: 19). In another embodiment, the viral vector or rAAV vector comprises an enhancer polynucleotide sequence comprising a nucleotide sequence which contains one or more regions of about 100 bp or longer having at least 75% or greater sequence identity to a polynucleotide sequence of human enhancer element E6 (SEQ ID: 20). In an embodiment of the above-delineated aspects, the viral vector or rAAV vector comprises an enhancer polynucleotide sequence comprising the polynucleotide sequence of human enhancer element E2 (SEQ ID NO: 16). In other embodiments of the above-delineated aspects, the viral vector or rAAV vector comprises an enhancer polynucleotide sequence comprising

the polynucleotide sequence of human enhancer element E5 (SEQ ID NO: 19) or an enhancer polynucleotide sequence comprising the polynucleotide sequence of human enhancer element E6 (SEQ ID NO: 20). In other embodiments of the above-delineated aspects, the viral vector or rAAV vector comprises any one (or one or more) of an enhancer polynucleotide sequence comprising the polynucleotide sequence of human enhancer element E11 (SEQ ID NO: 25) to E35 (SEQ ID NO: 49). In an embodiment, the capacity of the vector to package polynucleotide sequences of greater than about 4.7 kb comprises reassembly of multiple rAAV vectors by homologous recombination or by splicing mediated by acceptor sites. In an embodiment, the vector delivers the *SCN1A* gene to *SCN1A*-expressing GABAergic interneuron or glutamatergic pyramidal neuron cells in the brain, and wherein the *SCN1A* gene is functionally expressed, thereby restoring normal levels of *SCN1A* in the interneuron and neuron cells following administration of the vector to a subject. In an embodiment, the subject is a human patient. In an embodiment, the human patient is an infant suffering from Dravet syndrome (DS).

In another aspect, a viral particle or virus-like particle comprising the viral vector or rAAV vector of any of the above-delineated aspects is provided.

In another aspect, a cell comprising the viral vector or rAAV vector of any of the above-delineated aspects is provided. In an embodiment, the cell comprises the viral particle as delineated above.

In another aspect is provided a pharmaceutical composition comprising the viral vector or rAAV vector of any of the above-delineated aspects, and a pharmaceutically acceptable vehicle, carrier, or diluent.

In another aspect is provided a pharmaceutical composition comprising the viral particle of any of the above-delineated aspects, and a pharmaceutically acceptable vehicle, carrier, or diluent. In an embodiment of the above-delineated aspects, the pharmaceutical composition is in liquid dosage form.

In an aspect, a method of restoring normal levels of *SCN1A* expression in GABAergic interneuron cells in which *SCN1A* expression levels are deficient or defective is provided, in which the method comprises contacting the cells with an effective amount of the viral or rAAV vector of any of the above-delineated aspects,

or a viral particle or a pharmaceutical composition thereof, to restore normal levels of *SCN1A* expression in the GABAergic interneuron cells.

In an aspect, a method of treating infantile epilepsy and/or seizures in an infant who has or is at risk of having epilepsy, seizures, or Dravet syndrome (DS) is

5 provided, in which the method comprises administering to the infant a therapeutically effective amount of the viral or rAAV vector of any of the above-delineated aspects, the viral particle of any of the above-delineated aspects, or a pharmaceutical composition of any of the above-delineated aspects, to treat seizures, epilepsy, or DS in the subject.

10 In an aspect, a method of treating Dravet syndrome (DS) in a subject who has or is at risk of having DS is provided, in which the method comprises administering to the subject a therapeutically effective amount of the viral or rAAV vector of any of the above-delineated aspects, or a viral particle or a pharmaceutical composition thereof, to treat DS in the subject.

15 In an aspect, a method of inhibiting or preventing seizures and/or epilepsy in a subject having or at risk of having seizures and/or epilepsy is provided, the method comprising systemically administering to the subject a recombinant adeno-associated virus (rAAV) vector comprising an *SCN1A* transgene polynucleotide sequence, or a functional portion thereof, an enhancer polynucleotide sequence that specifically
20 restricts expression of the *SCN1A* transgene in interneuron cells of the cerebral cortex of the subject, and a capsid that enhances transduction of the vector into interneuron cells.

In an embodiment of the methods in any of the above-delineated aspects, the infant or the subject is a human patient. In an embodiment of the methods in any of
25 the above-delineated aspects, the enhancer polynucleotide sequence in the viral vector or rAAV vector is selected from human enhancer elements E1, E2, E3, E4, E5, E6, E7, E8, E9, or E10, or E11-E35 (SEQ ID NOs: 25-49, respectively). In an embodiment, the viral vector or rAAV vector comprises an enhancer polynucleotide sequence comprising a nucleotide sequence which contains one or more regions of
30 about 100 bp or longer having at least 75% or greater sequence identity to a polynucleotide sequence of a human enhancer element E1, E2, E3, E4, E5, E6, E7, E8, E9, or E10 (SEQ ID NOs: 15-24, respectively) or E11-E35 (SEQ ID NOs: 25-49,

respectively). In an embodiment, the enhancer polynucleotide sequence is the human E2 enhancer polynucleotide sequence or the enhancer polynucleotide sequence contains one or more regions of about 100 bp or longer having at least 75% or greater sequence identity to a polynucleotide sequence of a human enhancer element E1, E2, E3, E4, E5, E6, E7, E8, E9, or E10 (SEQ ID NOs: 15-24, respectively) or E11-E35 (SEQ ID NOs: 25-49, respectively). In an embodiment, the enhancer polynucleotide sequence is the human E5 enhancer polynucleotide sequence. In an embodiment, the enhancer polynucleotide sequence is the human E6 enhancer polynucleotide sequence. In a certain embodiment, the enhancer polynucleotide sequence contains one or more regions of about 100 bp or longer having at least 75% or greater sequence identity to a polynucleotide sequence of human enhancer element E2 (SEQ ID NO: 16). In other embodiments, the enhancer polynucleotide sequence contains one or more regions of about 100 bp or longer having at least 75% or greater sequence identity to a polynucleotide sequence of human enhancer element E5 (SEQ ID NO: 19) or to a polynucleotide sequence of human enhancer element E6 (SEQ ID NO: 20).

In an aspect, a method of delivering a transgene for restricted expression in an interneuronal cell or neuronal cell that expresses an *SCN1A* gene to inhibit or prevent seizures and/or epilepsy in a subject in need thereof is provided, in which the method comprises contacting the cell with a recombinant adeno-associated virus (rAAV) vector comprising an *SCN1A* transgene polynucleotide sequence, or a functional portion thereof, and an enhancer polynucleotide sequence that specifically restricts expression of the *SCN1A* transgene in interneuron or neuron cells of the cerebral cortex of the subject, thereby inhibiting or preventing seizures and/or epilepsy in the subject.

In an embodiment of the methods in any of the above-delineated aspects, the rAAV vector, viral particle, virus-like particle, or pharmaceutical composition is administered systemically. In an embodiment of the methods in any of the above-delineated aspects, the rAAV vector, viral particle, virus-like particle, or pharmaceutical composition is administered parenterally or intravenously. In an embodiment of the methods in any of the above-delineated aspects, the rAAV vector, viral particle, or pharmaceutical composition is administered intracerebrally. In an embodiment of the methods in any of the above-delineated aspects, the rAAV vector,

viral particle, or pharmaceutical composition is administered as a prophylactic. In an embodiment of the methods in any of the above-delineated aspects, the method further comprises administering an adjunct anti-epileptic treatment to the infant or subject.

5 In another aspect, a viral vector comprising a transgene polynucleotide sequence and an enhancer polynucleotide sequence that specifically restricts expression of the transgene in vaso-intestinal peptide-expressing cortical interneuron cells (VIP cINs) of the brain is provided. In another aspect, a viral vector is provided, in which the vector comprises an enhancer polynucleotide sequence specifically
10 associated with *SCN1A* gene expression and a transgene polynucleotide sequence, wherein the enhancer sequence restricts expression of the transgene in vaso-intestinal peptide-expressing cortical interneuron cells (VIP cINs) of the brain. In an embodiment, the enhancer polynucleotide sequence comprises a nucleotide sequence which contains one or more regions of about 100 bp or longer having at least 75% or
15 greater sequence identity to a polynucleotide sequence of human enhancer element E6 (SEQ ID NO: 20). In a particular embodiment, the enhancer polynucleotide sequence is human enhancer element E6 (SEQ ID NO: 20). In an embodiment, the viral vector is recombinant adeno-associated virus (rAAV) vector. In an embodiment, the transgene is the *SCN1A* gene.

20 In another aspect, a viral vector comprising a transgene polynucleotide sequence and an enhancer polynucleotide sequence that specifically restricts expression of the transgene in pyramidal neurons of the brain is provided. In another aspect, a viral vector is provided, in which the vector comprises an enhancer polynucleotide sequence specifically associated with *SCN1A* gene expression and a
25 transgene polynucleotide sequence, wherein the enhancer sequence restricts expression of the transgene in pyramidal neurons of the brain. In an embodiment, the enhancer polynucleotide sequence comprises a nucleotide sequence which contains one or more regions of about 100 bp or longer having at least 75% or greater sequence identity to a polynucleotide sequence of human enhancer element E5 (SEQ ID NO:
30 19). In a particular embodiment, the enhancer polynucleotide sequence is human enhancer element E5 (SEQ ID NO: 19). In another particular embodiment, the enhancer sequence restricts expression of the transgene in pyramidal neurons in

cortical layer 5 of the brain. In an embodiment, the viral vector is recombinant adeno-associated virus (rAAV) vector. In an embodiment, the transgene is the *SCN1A* gene.

In an aspect, a viral vector that comprises an enhancer polynucleotide sequence selected from SEQ ID NOs: 15-24, or a functional portion thereof, is provided, wherein the vector specifically targets neuronal cells expressing *SCN1A*. In an embodiment, the neuronal cells are parvalbumin cortical interneurons (PV cINs), pyramidal (PYR) neurons, or vaso-intestinal peptide cortical interneurons (VIP cIN).

In an aspect, a viral vector that comprises an enhancer polynucleotide sequence selected from SEQ ID NOs: 25-27, or a functional portion thereof, is provided, wherein the vector specifically targets cells expressing *Pvalb*.

In an aspect, a viral vector that comprises an enhancer polynucleotide sequence selected from SEQ ID NOs: 28-31, or a functional portion thereof, is provided, wherein the vector specifically targets cells expressing *Acan*.

In an aspect, a viral vector that comprises an enhancer polynucleotide sequence selected from SEQ ID NOs: 32-39, or a functional portion thereof, is provided, wherein the vector specifically targets cells expressing *Tmem132c*.

In an aspect, a viral vector that comprises an enhancer polynucleotide sequence selected from SEQ ID NO: 40 or SEQ ID NO: 41, or a functional portion thereof, is provided, wherein the vector specifically targets cells expressing *Lrrc38*.

In an aspect, a viral vector that comprises an enhancer polynucleotide sequence selected from SEQ ID NO: 42 or SEQ ID NO: 43, or a functional portion thereof, is provided, wherein the vector specifically targets cells expressing *Inpp5j*.

In an aspect, a viral vector that comprises an enhancer polynucleotide sequence selected from SEQ ID NOs: 44-47, or a functional portion thereof, is provided, wherein the vector specifically targets cells expressing *Mef2c*.

In an aspect, a viral vector that comprises an enhancer polynucleotide sequence selected from SEQ ID NO: 48 or SEQ ID NO: 49, or a functional portion thereof, is provided, wherein the vector specifically targets cells expressing *Pthlh*.

In an aspect, a viral vector that comprises an enhancer polynucleotide sequence selected from SEQ ID NOS: 15-49, or a functional portion thereof, is provided, wherein the vector specifically targets cells PV-expressing cells.

In an embodiment of the viral vector of any of the above-delineated aspects, the target cells are PV-expressing neuronal cells. In an embodiment of the viral vector of any of the above-delineated aspects, the viral vector is a lentiviral vector or a recombinant adeno-associated virus (rAAV) vector.

5 In an aspect is provided a cell comprising the viral vector of any of the above-delineated aspects and embodiments.

In an aspect is provided a viral particle or virus-like particle comprising the viral vector of any of the above-delineated aspects and embodiments. In an aspect is provided a cell comprising the viral particle or virus-like particle comprising the viral
10 vector of any of the above-delineated aspects and embodiments.

In an aspect is provided a pharmaceutical composition comprising the viral vector, or the viral particle or virus-like particle, of any of the above-delineated aspects and embodiments and a pharmaceutically acceptable vehicle, carrier, or diluent.

15 In another aspect, a method of restricting expression of a transgene in a neuronal cell of a subject is provided, in which the method comprises administering to the subject a delivery vector comprising at least one enhancer element polynucleotide comprising a sequence of SEQ ID NO: 15-49 and a transgene polynucleotide, wherein the transgene is specifically expressed in the neuronal cell. In an embodiment of the
20 method, the transgene is *SCN1A*. In an embodiment, the neuronal cell is a cortical interneuron expressing parvalbumin (PV cIN). In an embodiment, the enhancer element polynucleotide comprises a sequence set forth in SEQ ID NOS: 15-18 or SEQ ID NOS: 21-24.

In another embodiment of the above method, the neuronal cell is a pyramidal
25 (PYR) cell. In an embodiment, the enhancer element polynucleotide comprises the sequence set forth in SEQ ID NO: 19.

In another embodiment of the above method, the neuronal cell is a cortical interneuron expressing the vaso-intestinal peptide (VIP cIN). In an embodiment, the enhancer element polynucleotide comprises the sequence set forth in SEQ ID NO: 20.

30 In an embodiment of the above-delineated method and its embodiments, the delivery vector is a lentiviral vector or rAAV. In an embodiment of the method, the delivery vector is administered to the brain. In an embodiment of the method, the

delivery vector is administered locally or systemically. In an embodiment, the subject is a mammal. In an embodiment, the subject is human.

In another aspect, a viral vector comprising a human enhancer polynucleotide sequence selected from SEQ ID NOS: 15-49 is provided. In an embodiment, the viral
5 vector is a recombinant adeno-associated virus (rAAV) vector. In embodiments, a viral particle or virus-like particle comprises the above-delineated viral vector. In another embodiment, a cell comprises the above-delineated viral vector. In an embodiment, a cell comprises the above-delineated viral particle or the virus-like particle. In an embodiment, a pharmaceutical composition comprises the above-
10 delineated viral vector, or the above-delineated viral particle or virus-like particle, and a pharmaceutically acceptable vehicle, carrier, or diluent.

Definitions

Unless defined otherwise, all technical and scientific terms used herein have the meaning commonly understood by a person skilled in the art to which the
15 described aspects and embodiments belong. The following references provide one of skill with a general definition of many of the terms used in the described embodiments: Singleton et al., Dictionary of Microbiology and Molecular Biology (2nd ed. 1994); The Cambridge Dictionary of Science and Technology (Walker ed., 1988); The Glossary of Genetics, 5th Ed., R. Rieger et al. (eds.), Springer Verlag
20 (1991); and Hale & Marham, The Harper Collins Dictionary of Biology (1991). As used herein, the following terms have the meanings ascribed to them below, unless specified otherwise.

By “administering” is meant giving, supplying, dispensing a composition, agent, therapeutic product, e.g., a virus vector (rAAV) harboring a transgene (e.g., an
25 effector or a therapeutic gene), and the like to a subject, or applying or bringing the composition and the like into contact with the subject. Administering or administration may be accomplished by any of a number of routes, such as, for example, without limitation, parenteral or systemic, intravenous (IV), (injection), subcutaneous, intrathecal, intracranial, intramuscular, dermal, intradermal, inhalation,
30 rectal, intravaginal, topical, oral, subcutaneous, intramuscular, or intraocular. In embodiments, administration is systemic, such as by inoculation, injection, or intravenous injection.

By “agent” is meant a peptide, polypeptide, nucleic acid molecule, or small molecule chemical compound, antibody, or a fragment thereof.

By "alteration" is meant a change (increase or decrease) in the expression levels or activity of a gene or polypeptide as detected by standard art known methods
5 such as those described herein. As used herein, an alteration includes a 10% change in expression levels, a 25% change, a 40% change, or a 50% or greater change in expression levels. "

By “ameliorate” and “amelioration” is meant decrease, suppress, attenuate, diminish, arrest, or stabilize the development or progression of a disease.

10 By "analog" or “derivative” is meant a molecule that is not identical, but has analogous functional or structural features. For example, a polypeptide analog retains the biological activity of a corresponding naturally-occurring polypeptide, while having certain biochemical modifications that enhance the analog's function relative to a naturally occurring polypeptide. Such biochemical modifications could increase
15 the analog's protease resistance, membrane permeability, or half-life, without altering, for example, polynucleotide binding activity. In another example, a polynucleotide analog retains the biological activity of a corresponding naturally-occurring polynucleotide while having certain modifications that enhance the analog's function relative to a naturally occurring polynucleotide. Such modifications could increase
20 the polynucleotide's affinity for DNA, half-life, and/or nuclease resistance, an analog may include an unnatural nucleotide or amino acid.

As used herein, the term “at risk” as it applies to a neurological or neurogenetic disease, disorder, or pathology, such as seizures or epilepsy, refers to patients or individuals who have a family history or genetic risk factor genes for a
25 neurological or neurogenetic disease, disorder, or pathology.

As used herein, the term "carrier" refers to a diluent, adjuvant, excipient, or vehicle with which a composition or pharmaceutical composition, e.g., comprising a polynucleotide, viral vector, or viral particle) can be administered. Pharmaceutical and pharmaceutically acceptable carriers include sterile liquids, such as water and
30 oils, including those of petroleum, animal, vegetable or synthetic origin, such as peanut oil, soybean oil, mineral oil, sesame oil, and the like. Water or aqueous saline solutions and aqueous dextrose and glycerol solutions may be employed as

carriers, particularly for injectable solutions. Carriers may also include solid dosage forms, including, but not limited to, one or more of a binder (for compressed pills), a glidant, an encapsulating agent, a flavorant, and a colorant. Suitable pharmaceutical carriers are described in "Remington's Pharmaceutical Sciences" by E.W. Martin.

5 As used herein, "comprises," "comprising," "containing" and "having" and the like can have the meaning ascribed to them in U.S. Patent law and can mean "includes," "including," and the like; "consisting essentially of" or "consists essentially" likewise has the meaning ascribed in U.S. Patent law and the term is open-ended, allowing for the presence of more than that which is recited so long as
10 basic or novel characteristics of that which is recited are not changed by the presence of more than that which is recited, but excludes prior art embodiments.

 "DREADD" is an acronym for "designer receptor exclusively activated by a designer drug," which is a modified G protein coupled receptor (GPCR) that may be administered or specifically introduced into a subject, or cells thereof, e.g., PV-
15 expressing interneurons, by use of a viral vector (which contains a polynucleotide sequence encoding the DREADD) or through genetic breeding. DREADDs are known as chemical genetic or "chemogenetic" molecules allow for a precise level of temporal control over the excitation and inhibition of neurons. Following expression of the DREADD, it may be activated by a specific ligand (or agonist), which may be
20 administered by intravenous injection or orally. The DREADD and its ligand are designed to be orthogonal, i.e., they bind specifically to each other and do not cross-react. By way of nonlimiting example, five different classes of DREADDs are available for use: hM3Dq raises calcium levels in a cell, causing burst firing; hM4Di lowers cAMP and the activation of a particular potassium channel, causing neuronal
25 silencing, and also inhibits presynaptic neurotransmitter release; GsD enhances cAMP, causing modulation signaling; and Rq(R165L) enhances arrestin signaling, a specific pathway that has been linked to the mechanisms of psychoactive drugs; and κ -opioid receptor DREADD or KORD, which reduces or inhibits excitation of neurons and also inhibits presynaptic neurotransmitter release. (See, e.g., Kelly Rae
30 Chi, 2015, *The Scientist*; and S.M. Sternson and B.L. Roth, 2014, *Ann Rev Neuroscience*, 37:387-407).

Orthogonal ligand-gated ion channels, called pharmacologically selective actuator molecules (PSAMs) and pharmacologically selective effector molecules (PSEMs) are other types of chemogenetic molecules that are used as optogenetic agents and in optogenetic methods, in a manner similar to the use of DREADDs.

- 5 Each PSAM is exclusively activated by a PSEM cognate synthetic agonist. By way of example, three specific PSAM/PSEM tools have been designed, each with different ion conductance properties for controlling neuronal excitability. (See, e.g., Shapiro, M.G. et al., 2012, *ACS Chem. Neurosci.*, 3(8):619-629). These include the cation-selective activator, PSAM^{Q79G,Q139G}-5HT3HC/PSEM^{22S}, the anion-selective silencer, 10 PSAM^{L141F,Y115F}-GlyR/PSEM^{89S}, and a third Ca²⁺-selective channel, PSAM^{Q79G,L141S}-nAChR V13/T/PSEM^{9S}. (See, *Ibid.*, and Magnus, C.J. et al., 2011, *Science*, 333(6047):1292-1296). Both DREADDs and PSAMs-PSEMs allow control over neuronal activity, in a temporal manner, from minutes to hours. (See, e.g., Kelly Rae Chi, 2015, *The Scientist*; and S.M. Sternson and B.L. Roth, 2014, *Ann Rev* 15 *Neuroscience*, 37:387-407). By way of example, different PSAMs have been used with various ion channels and PSEMs to control neurons, e.g., E/I balance in neurons. Such PSAM-PSEM pairings include, without limitation, PSAM^{L141F, Y115F}- 5HT3 HC, which is activated by the ligand PSEM^{89S}, allowing cations to flow into the cell and boost excitability; PSAM^{L141F, Y115F} - GlyR, which is activated by the ligand 20 PSEM^{89S}, silencing neurons; and PSAM^{Q79G, L141S}-nAChR V13, which is activated by the ligand PSEM^{9S}, enhancing calcium signaling. Because there are two different PSEM ligands, PSAMs-PSEMs can also be combined in the same animal (subject).

“Detect” refers to identifying the presence, absence or amount of a molecule, compound, or agent to be detected.

- 25 By “disease” is meant any condition or disorder that adversely affects, damages or interferes with the normal function of a cell, tissue, organ, or part of the body, such as the brain, including the cerebral cortex of the brain and brain tissues. In one embodiment, the disease is a seizure or epilepsy. In another embodiment, the disease is Dravet syndrome.

- 30 By “effective amount” is meant the amount of a required to ameliorate the symptoms of a disease relative to an untreated patient. The effective amount of active compound(s) used to practice the described methods for therapeutic treatment of a

disease varies depending upon the manner of administration, the age, body weight, and general health of the subject. Ultimately, the attending physician, clinician, or veterinarian will decide the appropriate amount and dosage regimen. Such amount is referred to as an "effective" amount. In one embodiment, an effective amount is the amount of an rAAV vector comprising a specific enhancer sequence (e.g., an *SCN1A*-specific enhancer, such as E1-E10, as described herein) and one or more transgene sequences (e.g., *SCN1A*) inserted therein that is required to reduce, ameliorate, abate, inhibit, or stabilize a symptom of a neurological disease or disorder, such as seizures, epilepsy, Dravet syndrome (DS), or the severity thereof. In another embodiment, an effective amount is the amount of an rAAV vector comprising a specific enhancer sequence (e.g., an *SCN1A*-specific enhancer, e.g., E1-E10, as described herein) and one or more transgene sequences (e.g., *SCN1A*) inserted therein required to cause specific inhibitory activity of an interneuron cell, such as a GABAergic interneuron cell or a PV-expressing, GABAergic interneuron cell. In an embodiment, the enhancer is E2, as described herein, which restricts expression of a transgene, e.g., *SCN1A* or effectors like Gq-DREADD or PSAM for chemogenetic modulation of PV-interneuron activity, to PV-interneuron cells.

As used herein, the term "endogenous" describes a molecule (e.g., a polypeptide, peptide, nucleic acid, or cofactor) that is found naturally in a particular organism (e.g., a human) or in a particular location within an organism (e.g., an organ, a tissue, or a cell, such as a human cell).

As used herein, the term "exogenous" refers to a molecule (e.g., a polypeptide, peptide nucleic acid, or cofactor) that is not found naturally or endogenously in a particular organism (e.g., a human) or in a particular location within an organism (e.g., an organ, a tissue, or a cell, such as a human cell). Exogenous materials include those that are provided from an external source to an organism or to cultured matter extracted therefrom.

A "regulatory element," "regulatory sequence," "enhancer," "enhancer element" or "enhancer sequence" refers to a nucleic acid or polynucleotide sequence or a region of a nucleic acid or polynucleotide sequence, e.g., DNA or RNA, of about 50-2500 nucleotides, that contains one or more binding sites that are recognized and bound by one or more binding protein(s), e.g., transcription factor(s). In general, the

binding proteins function as activators to increase the likelihood that transcription of a particular target gene will occur. Enhancers can activate transcription independent of their location, distance or orientation with respect to the promoters of genes. For example, enhancer sequences may be located upstream of a gene, downstream of a gene, within the coding region of a gene, or up to one million base pairs away from the gene. Typically, the binding of a DNA binding protein(s) or transcription factor(s) to an enhancer changes or alters the conformation of the DNA, thereby allowing interactions to occur between or among the transcription factor(s) bound to the DNA.

Enhancers have been described as clusters of DNA sequences capable of binding combinations of transcription factors that then interact with components of the mediator complex or TFIID to help recruit RNA polymerase II (RNAPII). To accomplish this, enhancer-bound transcription factors loop out the intervening sequences and contact the promoter region of a gene, thus allowing enhancers to act in a distance-independent fashion. In addition, activation of eukaryotic genes requires de-compaction of the chromatin fiber, which is carried out by enhancer-bound transcription factors that can recruit histone modifying enzymes or ATP-dependent chromatin remodeling complexes to alter chromatin structure and increase the accessibility of the DNA to other proteins. (For a review of enhancer function, *see, e.g., Ong, C.-T. and Corces, V.G., 2011, Nat. Rev. Genetics, 12(4):283-293*).

As described herein, ten enhancers in the vicinity of the *SCN1A* gene were screened for the ability to restrict expression of a transgene, namely, *SCN1A*, to PV-expressing interneuron cells (PV-cells), most of which express the *SCN1A* gene. The isolated enhancer sequences, called S5E1 (E1) – S5E10 (E10) herein, were discovered to have the ability to restrict expression of *SCN1A* to GABA-ergic interneurons. By way of example, the E2 enhancer (S5E2) was demonstrated to target and restrict expression of a transgene to PV-expressing interneurons, which express *SCN1A*. It will be appreciated that a large fraction of cells that express *SCN1A* are not PV-expressing interneurons. In an embodiment, the enhancers as described herein allow the restriction of expression of a transgene, e.g., *SCN1A* or another effector gene, e.g., Gq-DREADD or PSAM, in PV-interneurons, rather than in all *SCN1A*-expressing neurons. By way of further example, the isolated E5 enhancer (S5E5) was

demonstrated to target and restrict expression of a transgene to glutamatergic pyramidal neurons in the brain. In embodiments, such an enhancer is E1, E2, E3, E4, E5, E6, E7, E8, E9, or E10, as described herein. In an embodiment, an enhancer element is isolated from its naturally occurring environment. Such an enhancer
5 element is used in a vector, e.g., a viral vector, for delivery to a cell, tissue, or region of the body, such as the brain.

By "fragment" is meant a portion of a polypeptide or nucleic acid molecule. This portion contains at least 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, or 90% of the entire length of the reference nucleic acid molecule or polypeptide. A fragment
10 may contain 10, 20, 30, 40, 50, 60, 70, 80, 90, or 100, 200, 300, 400, 500, 600, 700, 800, 900, or 1000 nucleotides or amino acids.

By "functionally expressed" is meant that a gene or transgene contained in or inserted into the polynucleotide of an rAAV or rAAV vector as described herein is expressed in an infected or transduced cell and produces its encoded product, which is
15 functional and/or active in the cell. In an embodiment, the cell is an interneuron cell. In an embodiment, the cell is a GABAergic interneuron cell. In an embodiment, the cell is a GABAergic interneuron cell that expresses parvalbumin (PV). In an embodiment, the cell is a neuron, in particular, a glutamatergic pyramidal interneuron cell. In an embodiment, the transgene is a detectable reporter gene, such as d-Tomato,
20 ChR2, GFP, RFP, and the like. In an embodiment, the transgene is a Designer receptor exclusively activated by designer drugs (DREADD) or Gq-DREADD. In an embodiment, the transgene is PSAM. In an embodiment, the transgene is *SCN1A* which encodes the sodium channel Nav1.1.

"Hybridization" means hydrogen bonding, which may be Watson-Crick,
25 Hoogsteen or reversed Hoogsteen hydrogen bonding, between complementary nucleobases. For example, adenine and thymine are complementary nucleobases that pair through the formation of hydrogen bonds.

The term "interneuron" refers to a neuron (nerve cell), or local circuit neuron in the central nervous system (CNS) that relays impulses between sensory neurons
30 and motor neurons. In general, neurons are specialized cells that function primarily in the transmission of nerve impulses. Neurons have cellular processes, such as dendrites and axons. Dendrites, which are shorter processes in the cell body of a

neuron, receive inputs from other neurons and conduct signals to the cell body.

Axons are longer, single processes of the cell soma and relay signals toward the tip of the neuron (called the synaptic terminal). The three, main types of neurons include sensory neurons, interneurons (of the CNS), and motor neurons. In the human brain, there are about 100 billion interneurons, which receive impulses from the sensory neurons. Interneurons interpret the information received from other neurons and relay impulses to motor neurons for an appropriate response in a function called 'integration.'

The terms "isolated," "purified," or "biologically pure" refer to material that is free to varying degrees from components which normally accompany or are associated with it as found in its native state. "Isolate" denotes a degree of separation from original source or surroundings. "Purify" denotes a degree of separation that is higher than isolation. A "purified" or "biologically pure" protein or polynucleotide is sufficiently free of other materials such that any impurities do not materially affect the biological properties of the protein or polynucleotide, or cause other adverse consequences. That is, a polynucleotide (nucleic acid), polypeptide, or peptide is purified if it is substantially free of cellular material, viral material, or culture medium when produced by recombinant DNA techniques, or chemical precursors or other chemicals when chemically synthesized. Purity and homogeneity are typically determined using analytical chemistry techniques, for example, polyacrylamide gel electrophoresis or high-performance liquid chromatography. The term "purified" can denote that a nucleic acid, protein, or peptide gives rise to essentially one band in an electrophoretic gel. For a protein that can be subjected to modifications, for example, phosphorylation or glycosylation, different modifications may give rise to different isolated proteins, which can be separately purified.

By "isolated polynucleotide" is meant a nucleic acid (e.g., a DNA) that is free of the genes which flank the gene in the naturally-occurring genome of the organism from which a nucleic acid molecule, such as a nucleic acid molecule described herein, is derived. The term therefore includes, for example, a recombinant DNA that is incorporated into a vector; into an autonomously replicating plasmid or virus; or into the genomic DNA of a prokaryote or eukaryote; or that exists as a separate molecule (for example, a cDNA or a genomic or cDNA fragment produced by PCR or

restriction endonuclease digestion) independent of other sequences. In addition, the term includes an RNA molecule that is transcribed from a DNA molecule, as well as a recombinant DNA that is part of a hybrid gene encoding additional polypeptide sequence.

5 By an "isolated polypeptide" is meant a polypeptide that has been separated from components that naturally accompany it. Typically, a polypeptide is isolated when it is at least 60%, by weight, free from the proteins and naturally-occurring organic molecules with which it is naturally associated. Preferably, the preparation is at least 75%, or at least 85%, or at least 90%, or at least 99%, by weight, a desired
10 polypeptide. An isolated polypeptide may be obtained, for example, by extraction from a natural source, by expression of a recombinant nucleic acid encoding such a polypeptide; or by chemically synthesizing the protein. Purity can be measured by any appropriate method, for example, column chromatography, polyacrylamide gel electrophoresis, or by HPLC analysis.

15 By "marker" is meant any protein or polynucleotide having an alteration in expression, level or activity that is associated with a disease or disorder. In one embodiment, a marker is an *SCN1A* polynucleotide or *SCN1A* polypeptide.

The term "mutation," as used herein, refers to a substitution of a nucleotide base or amino acid residue within a sequence, *e.g.*, a nucleic acid or amino acid
20 sequence, respectively, with another residue, or a deletion or insertion of one or more residues within a sequence. Mutations are typically described herein by identifying the original residue followed by the position of the residue within the sequence and by the identity of the newly substituted residue. Various methods for making the amino acid substitutions (mutations) provided herein are well known in the art, and are
25 provided by, for example, Green and Sambrook, *Molecular Cloning: A Laboratory Manual* (4th ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (2012)).

As used herein, "obtaining" as in "obtaining an agent" includes synthesizing, purchasing, or otherwise acquiring the agent.

30 By "polynucleotide" is meant a nucleic acid molecule, *e.g.*, a double-stranded (ds) DNA polynucleotide, a single-stranded (ss) DNA polynucleotide, a dsRNA polynucleotide, or a ssRNA polynucleotide, that encodes one or more polypeptides.

The term encompasses positive-sense (i.e., protein-coding) DNA polynucleotides, which are capable of being transcribed to form an RNA transcript, which can be subsequently translated to produce a polypeptide following one or more optional RNA processing events (e.g., intron excision by RNA splicing, or ligation of a 5' cap or a 3' polyadenyl tail). The term additionally encompasses positive-sense RNA polynucleotides, capable of being directly translated to produce a polypeptide following one or more optional RNA processing events. As used herein, a polynucleotide may be contained within a viral vector, such as a recombinant adeno-associated viral vector (rAAV).

The terms “nucleic acid” and “nucleic acid molecule,” as used herein, refer to a compound comprising a nucleobase and an acidic moiety, e.g., a nucleoside, a nucleotide, or a polymer of nucleotides. Typically, polymeric nucleic acids, e.g., nucleic acid molecules comprising three or more nucleotides are linear molecules, in which adjacent nucleotides are linked to each other via a phosphodiester linkage. In some embodiments, “nucleic acid” refers to individual nucleic acid residues (e.g. nucleotides and/or nucleosides). In some embodiments, “nucleic acid” refers to an oligonucleotide chain comprising three or more individual nucleotide residues. As used herein, the terms “oligonucleotide” and “polynucleotide” can be used interchangeably to refer to a polymer of nucleotides (e.g., a string of at least three nucleotides). In some embodiments, “nucleic acid” encompasses RNA as well as single and/or double-stranded DNA. Nucleic acids may be naturally occurring, for example, in the context of a genome, a transcript, an mRNA, tRNA, rRNA, siRNA, snRNA, a plasmid, cosmid, chromosome, chromatid, or other naturally occurring nucleic acid molecule. On the other hand, a nucleic acid molecule may be a non-naturally occurring molecule, e.g., a recombinant DNA or RNA, an artificial chromosome, an engineered genome, or fragment thereof, or a synthetic DNA, RNA, DNA/RNA hybrid, or including non-naturally occurring nucleotides or nucleosides. Furthermore, the terms “nucleic acid,” “DNA,” “RNA,” and/or similar terms include nucleic acid analogs, e.g., analogs having other than a phosphodiester backbone. Nucleic acids can be purified from natural sources, produced using recombinant expression systems and optionally purified, chemically synthesized, *etc.* Where appropriate, e.g., in the case of chemically synthesized molecules, nucleic acids can

comprise nucleoside analogs such as analogs having chemically modified bases or sugars, and backbone modifications. A nucleic acid sequence is presented in the 5' to 3' direction unless otherwise indicated. In some embodiments, a nucleic acid is or comprises natural nucleosides (e.g., adenosine, thymidine, guanosine, cytidine, uridine, deoxyadenosine, deoxythymidine, deoxyguanosine, and deoxycytidine); nucleoside analogs (e.g., 2-aminoadenosine, 2-thiothymidine, inosine, pyrrolo-pyrimidine, 3-methyl adenosine, 5-methylcytidine, 2-aminoadenosine, C5-bromouridine, C5-fluorouridine, C5-iodouridine, C5-propynyl-uridine, C5-propynyl-cytidine, C5-methylcytidine, 2-aminoadenosine, 7-deazaadenosine, 7-deazaguanosine, 8-oxoadenosine, 8-oxoguanosine, O(6)-methylguanine, and 2-thiocytidine); chemically modified bases; biologically modified bases (e.g., methylated bases); intercalated bases; modified sugars (2'-e.g., fluororibose, ribose, 2'-deoxyribose, arabinose, and hexose); and/or modified phosphate groups (e.g., phosphorothioates and 5'-N-phosphoramidite linkages).

As used herein, the term "pharmaceutically acceptable" refers to molecular entities, biological products and compositions that are physiologically tolerable and do not typically produce an allergic or other adverse reaction, such as gastric upset, dizziness and the like, when administered to a patient (e.g., a human patient).

As used herein, the terms "prevent," "preventing," "prevention," "prophylactic treatment" and the like refer to reducing the probability of developing a disorder or condition in a subject, who does not have, but who is at risk of, susceptible to, or predisposed to, developing a disorder or condition.

As used herein, the term "pseudotyped" refers to a viral vector that contains one or more foreign viral structural proteins, e.g., envelope glycoproteins. A pseudotyped virus may be one in which the envelope glycoproteins of an enveloped virus or the capsid proteins of a non-enveloped virus originate from a virus that differs from the source of the original virus genome and the genome replication apparatus. (D.A. Sanders, 2002, *Curr. Opin. Biotechnol.*, 13:437-442). The foreign viral envelope proteins of a pseudotyped virus can be utilized to alter host tropism or to increase or decrease the stability of the virus particles. Examples of pseudotyped viral vectors include a virus that contains one or more envelope glycoproteins that do not naturally occur on the exterior of the wild-type virus. Pseudotyped viral vectors can

infect cells and express and produce proteins or molecules encoded by polynucleotides, e.g., reporter or effector proteins or molecules, contained within the viral vectors, e.g., the sodium channel Nav1.1 encoded by the *SCN1A* gene.

The term "recombinant" as used herein in the context of proteins or nucleic acids refers to proteins or nucleic acids that do not occur in nature (or in a naturally occurring protein or nucleic acid sequence), but are the product of human engineering, often or typically utilizing molecular biological or molecular genetic tools and techniques practiced by the skilled practitioner in the art. For example, in some embodiments, a recombinant protein or nucleic acid molecule comprises an amino acid or nucleotide sequence that comprises at least one, at least two, at least three, at least four, at least five, at least six, at least seven, or at least eight mutations as compared to any naturally occurring sequence.

By "reduces" is meant a negative alteration of at least 5%, 10%, 25%, 50%, 75%, or 100%.

By "reference" is meant a standard or control condition. A "reference sequence" is a defined sequence used as a basis for sequence comparison. A reference sequence may be a subset of or the entirety of a specified sequence, for example, a segment of a full-length cDNA or gene sequence, or the complete cDNA or gene sequence. For polypeptides, the length of the reference polypeptide sequence will generally be at least about 16 amino acids, at least about 20 amino acids, at least about 25 amino acids, or about 35 amino acids, about 50 amino acids, or about 100 amino acids. For nucleic acids, the length of the reference nucleic acid sequence will generally be at least about 50 nucleotides, at least about 60 nucleotides, at least about 75 nucleotides, or about 100 nucleotides, or about 300 nucleotides, or any integer thereabouts or therebetween.

By "specifically binds" is meant a nucleic acid molecule, polypeptide, or complex thereof (e.g., a binding protein such as a transcription factor and its cognate nucleic acid binding region), or a compound, or molecule that recognizes and binds a given polypeptide and/or nucleic acid molecule, but which does not substantially recognize and bind other molecules in a sample, for example, a biological sample.

By "subject" is meant a mammal, including, but not limited to, a human or non-human mammal, such as a non-human primate, e.g., a marmoset, or a non-human

mammal, such as a bovine, equine, canine, ovine, or feline mammal, or a sheep, goat, llama, camel, or a rodent (rat, mouse), ferret, gerbil, hamster, or zebrafish. A subject is typically a patient, such as a human patient, who receives treatment for a particular disease or condition as described herein (e.g., a neuropsychiatric, neurological, or neurogenetic disease, disorder, or pathology, such as seizures, epilepsy, or DS). Examples of subjects and patients include mammals, such as humans, receiving treatment for such diseases or conditions or who are at risk of having such diseases or conditions.

Ranges provided herein are understood to be shorthand for all of the values within the range. For example, a range of 1 to 50 is understood to include any number, combination of numbers, or sub-range from the group consisting 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, or 50, inclusive of the first and last values.

As used herein, the term “therapeutically effective amount” refers to a quantity of a therapeutic agent that is sufficient to treat, abate, reduce, diagnose, prevent, and/or delay the onset of one or more symptoms of a disease, disorder, and/or condition upon administration to a patient in need of treatment. In some cases, a therapeutically effective amount may also refer to a quantity of a therapeutic agent that is administered prophylactically (e.g., in advance of the development of full-blown disease) to a subject who is at risk of developing a disease or the symptoms thereof, such as a neurological, neurodegenerative, or neurogenetic disease or disorder. In an embodiment, the disorder is Dravet syndrome (DS).

As used herein, the terms “treat,” “treating,” “treatment,” and the like refer to reducing or ameliorating a disorder and/or symptoms associated therewith. It will be appreciated that, although not precluded, treating a disorder or condition does not require that the disorder, condition or symptoms associated therewith be completely eliminated. “Treat” or “treatment” may refer to therapeutic treatment, in which the object is to prevent or slow down (lessen or reduce) an undesired physiological change or disorder. Beneficial or desired clinical results include, but are not limited to, alleviation of symptoms, diminishment of extent of disease, stabilized (i.e., not worsening) state of disease, delay or slowing of disease progression, amelioration or

palliation of the disease state, and remission (whether partial or total), whether detectable or undetectable. Those in need of treatment include those already with the condition or disorder, as well as those prone to have the condition or disorder or those in whom the condition or disorder is to be prevented.

5 As used herein, the terms “prevent,” “preventing,” “prevention,” “prophylactic treatment” and the like, refer to inhibiting or blocking a disease state, or the full development of a disease in a subject, or reducing the probability of developing a disease, disorder or condition in a subject, who does not have, but is at risk of developing, or is susceptible to developing, a disease, disorder, or condition.

10 As used herein, the term “vector” refers to a nucleic acid (e.g., a DNA vector, such as a plasmid), a RNA vector, virus or other suitable replicon (e.g., viral vector). A “vector” further refers to a nucleic acid (polynucleotide) molecule into which foreign nucleic acid can be inserted without disrupting the ability of the vector to be expressed in, replicate in, and/or integrate into a host cell. A variety of vectors have
15 been developed for the delivery of polynucleotides encoding exogenous proteins into a prokaryotic or eukaryotic cell. A vector may contain a polynucleotide sequence that includes gene of interest (e.g., a transgene, such as a therapeutic gene, a reporter gene, or, more specifically, an *SCN1A* gene encoding an Nav1.1 sodium channel) as well as, for example, additional sequence elements capable of regulating transcription,
20 translation, and/or the integration of these polynucleotide sequences into the genome of a cell. A vector may contain regulatory sequences, such as a promoter, e.g., a subgenomic promoter, region and an enhancer region, which direct gene transcription. A vector may contain polynucleotide sequences (enhancer sequences) that enhance the rate of translation of these genes or improve the stability or nuclear export of the
25 mRNA that results from gene transcription. These sequence elements may include, e.g., 5' and 3' untranslated regions, an internal ribosomal entry site (IRES), and/or a polyadenylation signal site in order to direct efficient transcription of a gene carried on the expression vector. Vectors, such as viral vectors or the rAAV vectors described herein, may also be referred to as expression vectors.

30 “Transduction” refers to a process by which DNA or polynucleotide, e.g., one or more transgenes, contained in a virus or virus vector is introduced or transferred into a cell by the virus or virus vector, wherein the DNA or polynucleotide is

expressed. In an embodiment, the DNA or polynucleotide transduced into a cell by a virus vector, such as an rAAV vector as described herein, is stably expressed in the cell. In some cases, a virus or virus vector is said to infect a cell.

As used herein, the term “vehicle” refers to a solvent, diluent, or carrier
5 component of a pharmaceutical composition.

By “virus particle” (also called a virion) is meant a virus (infectious agent) that exists as an independent particle comprising the core viral genome or genetic material (RNA or DNA); a protein coat, called the capsid, which surrounds the genetic material and protects it; and, in some cases, an envelope of lipids surrounding
10 the capsid. A virus particle may refer to the form of a virus before it infects a cell and becomes intracellular, or to the form of the virus that infects a cell.

By “virus-like particles (VLPs)” is meant virus particles made up of one of more viral structural proteins, but lacking the viral genome. Because VLPs lack a viral genome, they are non-infectious and yield safer and potentially more-economical
15 vaccines and vaccine products. In addition, VLPs can often be produced by heterologous expression and can be easily purified. Most VLPs comprise at least a viral core protein that drives budding and release of particles from a host cell.

By “substantially identical” is meant a polypeptide or nucleic acid molecule exhibiting at least 50% identity to a reference amino acid sequence (for example, any
20 one of the amino acid sequences described herein) or nucleic acid sequence (for example, any one of the nucleic acid sequences described herein). Preferably, such a sequence is at least 60%, preferably at least 70%, more preferably 80% or 85%, and most preferably 90%, 95% or even 99% identical at the amino acid level or nucleic acid to the sequence used for comparison, for example, over a specified comparison
25 window. Optimal alignment may be conducted using the homology alignment algorithm of Needleman and Wunsch, 1970, *J. Mol. Biol.*, 48:443. An indication that two peptide or polypeptide sequences are substantially identical is that one peptide or polypeptide is immunologically reactive with specific antibodies raised against the second peptide or polypeptide, although such cross-reactivity is not required for two
30 polypeptides to be deemed substantially identical. Thus, a peptide or polypeptide is substantially identical to a second peptide or polypeptide, for example, where the two differ only by a conservative substitution. Peptides or polypeptides that are

"substantially similar" share sequences as noted above except that residue positions which are not identical may differ by conservative amino acid changes. Conservative substitutions typically include, but are not limited to, substitutions within the following groups: glycine and alanine; valine, isoleucine, and leucine; aspartic acid and glutamic acid; asparagine and glutamine; serine and threonine; lysine and arginine; and phenylalanine and tyrosine, and others as known to the skilled person in the art.

Sequence identity is typically measured using sequence analysis software (for example, Sequence Analysis Software Package of the Genetics Computer Group, University of Wisconsin Biotechnology Center, 1710 University Avenue, Madison, Wis. 53705, BLAST, BESTFIT, GAP, or PILEUP/PRETTYBOX programs). Such software matches identical or similar sequences by assigning degrees of homology to various substitutions, deletions, and/or other modifications. Conservative substitutions typically include substitutions within the following groups: glycine, alanine; valine, isoleucine, leucine; aspartic acid, glutamic acid, asparagine, glutamine; serine, threonine; lysine, arginine; and phenylalanine, tyrosine. In an exemplary approach to determining the degree of identity, a BLAST program may be used, with a probability score between e^{-3} and e^{-100} indicating a closely related sequence.

By "substantially identical" is generally meant a polypeptide or nucleic acid molecule exhibiting at least 50% identity to a reference amino acid sequence (for example, any one of the amino acid sequences described herein) or nucleic acid sequence (for example, any one of the nucleic acid sequences described herein). In embodiments, such a sequence is at least 60%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, or greater, or at least 99% identical at the amino acid level or nucleic acid to the sequence used for comparison.

Polynucleotides or viral nucleic acid molecules useful in the methods and compositions as described herein include any nucleic acid molecule that encodes a polypeptide, or a fragment thereof, or that encodes the components of viral vectors described herein. The polynucleotides or viral nucleic acid molecules may encode polypeptide products harbored by the viral vectors, such as recombinant adeno-associated virus (rAAV) and the like, as well as a peptide or fragment thereof. Such

nucleic acid molecules need not be 100% identical with an endogenous sequence or a viral vector nucleic acid sequence, but will typically exhibit substantial identity.

Polynucleotides having substantial identity to an endogenous sequence or to a viral vector sequence are typically capable of hybridizing with at least one strand of a double-stranded nucleic acid molecule or to a viral vector nucleic acid molecule.

Nucleic acid molecules useful in the described methods include any nucleic acid molecule that encodes a polypeptide as described herein, or a fragment thereof. By "hybridize" is meant pairing of the nucleic acid molecules to form a double-stranded molecule between complementary polynucleotide sequences (e.g., a gene or nucleic acid sequence described herein), or portions thereof, under various conditions of stringency. (See, e.g., Wahl, G. M. and S. L. Berger (1987) *Methods Enzymol.* 152:399; Kimmel, A. R. (1987) *Methods Enzymol.* 152:507).

For example, stringent salt concentration will ordinarily be less than about 750 mM NaCl and 75 mM trisodium citrate, preferably less than about 500 mM NaCl and 50 mM trisodium citrate, and more preferably less than about 250 mM NaCl and 25 mM trisodium citrate. Low stringency hybridization can be obtained in the absence of organic solvent, e.g., formamide, while high stringency hybridization can be obtained in the presence of at least about 35% formamide, and more preferably at least about 50% formamide. Stringent temperature conditions will ordinarily include temperatures of at least about 30°C, more preferably of at least about 37°C, and most preferably of at least about 42°C. Varying additional parameters, such as hybridization time, the concentration of detergent, e.g., sodium dodecyl sulfate (SDS), and the inclusion or exclusion of carrier DNA, are well known to those skilled in the art. Various levels of stringency are accomplished by combining these various conditions as needed. In one embodiment, hybridization will occur at 30°C in 750 mM NaCl, 75 mM trisodium citrate, and 1% SDS. In a more preferred embodiment, hybridization will occur at 37°C in 500 mM NaCl, 50 mM trisodium citrate, 1% SDS, 35% formamide, and 100 µg/ml denatured salmon sperm DNA (ssDNA). In another embodiment, hybridization will occur at 42°C in 250 mM NaCl, 25 mM trisodium citrate, 1% SDS, 50% formamide, and 200 µg/ml ssDNA. Useful variations on these conditions will be readily apparent to those skilled in the art.

For most applications, washing steps that follow hybridization will also vary in stringency. Wash stringency conditions can be defined by salt concentration and by temperature. As above, wash stringency can be increased by decreasing salt concentration or by increasing temperature. For example, stringent salt concentration for the wash steps will preferably be less than about 30 mM NaCl and 3 mM trisodium citrate, and most preferably less than about 15 mM NaCl and 1.5 mM trisodium citrate. Stringent temperature conditions for the wash steps will ordinarily include a temperature of at least about 25°C, more preferably of at least about 42°C, and even more preferably of at least about 68° C. In an embodiment, wash steps will occur at 25°C in 30 mM NaCl, 3 mM trisodium citrate, and 0.1% SDS. In another embodiment, wash steps will occur at 42 C in 15 mM NaCl, 1.5 mM trisodium citrate, and 0.1% SDS. In yet another embodiment, wash steps will occur at 68° C in 15 mM NaCl, 1.5 mM trisodium citrate, and 0.1% SDS. Additional variations of these conditions will be readily apparent to those skilled in the art. Hybridization techniques are well known to those skilled in the art and are described, for example, in Benton and Davis (*Science*, 196:180, 1977); Grunstein and Hogness (*Proc. Natl. Acad. Sci., USA*, 72:3961, 1975); Ausubel et al. (*Current Protocols in Molecular Biology*, Wiley Interscience, New York, 2001); Berger and Kimmel (*Guide to Molecular Cloning Techniques*, 1987, Academic Press, New York); and Sambrook et al., *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, New York.

Nucleic acids that do not hybridize to each other under stringent conditions are still substantially identical if the polypeptides that they encode are substantially identical. This occurs, for example, when a copy of a nucleic acid is created using the maximum codon degeneracy permitted by the genetic code. In such cases, the nucleic acids typically hybridize under moderately stringent hybridization conditions. Nonlimiting examples of "moderately stringent hybridization conditions" include a hybridization in a buffer of 40% formamide, 1 M NaCl, 1% SDS at 37 C, and a wash in 1 x SSC at 45 C. A positive hybridization is at least twice background. Those of ordinary skill will readily recognize that alternative hybridization and wash conditions can be utilized to provide conditions of similar stringency.

By "ortholog" is meant any polypeptide or nucleic acid molecule of an organism that is highly related to a reference protein or nucleic acid sequence from another organism. The degree of relatedness may be expressed as the probability that a reference protein would identify a sequence, for example, in a blast search. The probability that a reference sequence would identify a random sequence as an ortholog is extremely low, less than e^{-10} , e^{-20} , e^{-30} , e^{-40} , e^{-50} , e^{-75} , e^{-100} . The skilled artisan understands that an ortholog is likely to be functionally related to the reference protein or nucleic acid sequence. In other words, the ortholog and its reference molecule would be expected to fulfill similar, if not equivalent, functional roles in their respective organisms, e.g., mouse and human orthologs.

It is not required that an ortholog, when aligned with a reference sequence, have a particular degree of amino acid sequence identity to the reference sequence. A protein ortholog might share significant amino acid sequence identity over the entire length of the protein, for example, or, alternatively, might share significant amino acid sequence identity over only a single functionally important domain of the protein. Such functionally important domains may be defined by genetic mutations or by structure-function assays. Orthologs may be identified using methods practiced in the art. The functional role of an ortholog may be assayed using methods well known to the skilled artisan. For example, function might be assayed *in vivo* or *in vitro* using a biochemical, immunological, or enzymatic assay; or transformation rescue. Alternatively, bioassays may be carried out in tissue culture; function may also be assayed by gene inactivation (e.g., by RNAi, siRNA, or gene knockout), or gene over-expression, as well as by other methods.

Ranges as provided herein are understood to be shorthand for all of the values within the range. For example, a range of 1 to 50 is understood to include any number, combination of numbers, or sub-range from the group 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, or 50, inclusive of the first and last values.

By *SCN1A* is meant a polypeptide or protein (the sodium channel Nav1.1) or fragment thereof having at least about or equal to 85%, or at least about or equal to 90%, 95%, 98%, 99%, or greater, amino acid sequence identity to the amino acid

sequence of the canonical amino acid sequence of *SCN1A*, Human Isoform 1,
 OmniProt Identifier No. P35498-1 (Length 2,009 amino acids; Mass (Da): 228,972);
 RefSeq Nos. NP_001159435.1; NP_001189364.1; NP_001340877. The polypeptide
 (protein) sequence of human *SCN1A* is as follows:

5	10	20	30	40	50
	MEQTVLVPPG	PDSFNFFTRE	SLAAIERRIA	EEKAKNPKPD	KKDDDENGPK
	60	70	80	90	100
	PNSDLEAGKN	LPFIYGDIPP	EMVSEPLEDL	DPYYINKKTF	IVLNKGKAI F
	110	120	130	140	150
10	RFSATSALYI	LTPFNPLRKI	AIKILVHSLF	SMLIMCTILT	NCVFMTMSNP
	160	170	180	190	200
	PDWTKNVEYT	FTGIYTFESL	IKIIARGFCL	EDFTFLRDPW	NWLDFTVITF
	210	220	230	240	250
	AYVTEFVDLG	NVSALRTFRV	LRALKTISVI	PGLKTIVGAL	IQSVKKLSDV
15	260	270	280	290	300
	MILTVFCLSV	FALIGLQLFM	GNLRNKCIQW	PPTNASLEEH	SIEKNITVNY
	310	320	330	340	350
	NGTLINETVF	EFDWKSIIQD	SRYHYFLEGF	LDALLCGNSS	DAGQCPEGYM
	360	370	380	390	400
20	CVKAGRNPNY	GYTSFDTFSW	AFLSLFRLMT	QDFWENLYQL	TLRAAGKTYM
	410	420	430	440	450
	IFFVLVIFLG	SFYLINLILA	VVAMAYEEQN	QATLEEAQK	EAEFQQMIEQ
	460	470	480	490	500
	LKKQQEAAQQ	AATATASEHS	REPSAAGRLS	DSSSEASKLS	SKSAKERRNR
25	510	520	530	540	550
	RKKRKQKEQS	GGEKDEDEF	QKSESEDSIR	RKGFRFSIEG	NRLTYEKRY S
	560	570	580	590	600
	SPHQSLLSIR	GSLFSPRRNS	RTSLFSFRGR	AKDVGSEND F	ADDEHSTFED
	610	620	630	640	650
30	NESRRDSL FV	PRRHGERRNS	NLSQTSRSSR	MLAVFPANGK	MHSTVDCNGV
	660	670	680	690	700
	VSLVGGPSVP	TSPVGQLLPE	VIIDKPATDD	NGTTTETEMR	KRRSSSFHVS
	710	720	730	740	750
	MDFLEDPSQR	GRAMSIASIL	TNTVEELEES	RQKCPPCWYK	FSNIFLIWDC
35	760	770	780	790	800
	SPYWLKVKHV	VNLVVMDPFV	DLAITICIVL	NTLFMAMEHY	PMTDHFNNVL
	810	820	830	840	850
	TVGNLVFTGI	FTAEMFLKII	AMDPYYYFQE	GWNIFDGFIV	TLSLVELGLA
	860	870	880	890	900
40	NVEGLSVLRS	FRLLRVFKLA	KSWPTLNMLI	KIIGNSVGAL	GNLTLVLAI I
	910	920	930	940	950
	VFIFAVVGMQ	LFGKSYKDCV	CKIASDCQLP	RWHMNDFFHS	FLIVFRVLCG
	960	970	980	990	1000
	EWIETMWDCM	EVAGQAMCLT	VFMMVMVIGN	LVVLNLFLAL	LLSSFSADNL
45	1010	1020	1030	1040	1050
	AATDDDNE MN	NLQIAVDRMH	KGVAIVKRKI	YEFIQQSFIR	KQKILDEIKP
	1060	1070	1080	1090	1100

	LDDLNNKKDS	CMSNHTAEIG	KDL DY LKDVN	GTTSGIGTGS	SVEKYIIDES
	1110	1120	1130	1140	1150
	DYMSFINNPS	LTVTVPPIAVG	ESDFENLNTE	DFSSESDLEE	SKEKLNESSS
	1160	1170	1180	1190	1200
5	SSEGSTVDIG	APVEEQPVVE	PEETLEPEAC	FTEGCVQRFK	CCQINVEEGR
	1210	1220	1230	1240	1250
	GKQWWNLRT	CFRIVEHNWF	ETFIVFMILL	SSGALAFEDI	YIDQRKTIKT
	1260	1270	1280	1290	1300
	MLEYADKVFT	YIFILEMLLK	WVAYGYQTYF	TNAWCWLDFL	IVDVSLVSLT
10	1310	1320	1330	1340	1350
	ANALGYSELG	AIKSLRTLRA	LRPLRALS RF	EGMRVVVNAL	LGAIPSIMNV
	1360	1370	1380	1390	1400
	LLVCLIFWLI	FSIMGVNLFA	GKFYHCINTT	TGDRFDIEDV	NNHTDCLKLI
	1410	1420	1430	1440	1450
15	ERNETARWKN	VKVNFDNVGF	GYLSLLQVAT	FKGWMDIMYA	AVDSRNVELQ
	1460	1470	1480	1490	1500
	PKYEESLYMY	LYFVIFII FG	SFFTLNLF IG	VIIDNFNQOK	KKFGGQDI FM
	1510	1520	1530	1540	1550
	TEEQKKYYNA	MKKLGSKKPQ	KPIPRPGNKF	QGMVFDFVTR	QVFDISIMIL
20	1560	1570	1580	1590	1600
	ICLNMTMMV	ETDDQSEYVT	TILSRINLVF	IVLFTGECVL	KLISLRHYFF
	1610	1620	1630	1640	1650
	TIGWNIFDFV	VVILSIVGMF	LAELIEKYFV	SPTLFRVIRL	ARIGRILRLI
	1660	1670	1680	1690	1700
25	KGAKGIRTLL	FALMMSLPAL	FNIGLLLFLV	MFIYAI FGMS	NFAYVKREVG
	1710	1720	1730	1740	1750
	IDDMFNFETF	GNSMICLFQI	TTSAGWDGLL	APILNSKPPD	CDPNKVNPGS
	1760	1770	1780	1790	1800
	SVKGDGCGNS	VGIFFFVSYI	IISFLVVNM	YIAVILENFS	VATEESAEP L
30	1810	1820	1830	1840	1850
	SEDDFEMFYE	VWEKFDPDAT	QFMFEKLSQ	FAAALEPPLN	LPQPNKLQLI
	1860	1870	1880	1890	1900
	AMDLP MVSGD	RIHCLDILFA	FTKRVLGESG	EMDALRIQME	ERFMASNPSK
	1910	1920	1930	1940	1950
35	VSYQPITTTL	KRKQEEVSAV	IIQRAYRRHL	LKRTVKQASF	TYNKNKIKGG
	1960	1970	1980	1990	2000
	ANLLIKEDMI	IDRINENSIT	EKTDLTMTA	ACPPSYDRVT	KPIVEKHEQE
	GKDEKAKGK	(SEQ ID NO: 1).			

The Nav1.1 sodium channel is encoded by a human *SCN1A* polynucleotide sequence or fragment thereof having at least about or equal to 85%, or at least about or equal to 90%, 95%, 98%, 99%, or greater, sequence identity to the *SCN1A* polynucleotide sequence under Accession No. NCBI CCDS 54413.1 (RefSeq Nos. NM_001165963.2; NM_001202435.2; NM_001353948.1) as set forth below.

(Genome information from Genome Reference Consortium GRCh38.p12. GenBank

assembly accession: GCA_000001405.27 (latest); RefSeq assembly accession:

GCF_000001405.38 (latest))

SCN1A Nucleotide Sequence (6030 nt):

```

5  atggagcaaacagtgcttgtaccaccaggacctgacagcttcaacttcttcaccagagaatctcttgcg
   gctattgaaagacgcattgcagaagaaaaggcaaagaatcccaaaccagacaaaaaagatgacgacgaa
   aatggcccaaagccaaatagtgacttggagctggaaagaaccttccatttatttatggagacattcct
10  ccagagatgggtgtcagagccccctggaggacctggaccctactatatcaataagaaaacttttatagta
   ttgaataaaggggaaggccatcttccgggttcagtgccacctctgccctgtacatttttaactcccttcaat
   cctcttaggaaaatagctattaagatttttggtacattcattattcagcatgctaattatgtgcactatt
   ttgacaaaactgtgtgtttatgacaatgagtaaccctcctgattggacaaagaatgtagaatacaccttc
   acaggaatatatacttttgaatcacttataaaaaattattgcaaggggatttctgtttagaagattttact
   ttccctcgggatccatggaactggctcgatttactgtcattacatttgcgtacgtcacagagtttgtg
15  gacctgggcaatgtctcggcattgagaacattcagagttctccgagcattgaagacgatttcagtcatt
   ccaggcctgaaaaccattgtgggagccctgatccagtcctgtgaagaagctctcagatgtaatgatcctg
   actgtgttctgtctgagcgtatttgcctaaattgggctgcagctgttcatgggcaacctgaggaataaa
   tgtatacaatggcctcccaccaatgcttccttggagggaacatagtatagaaaagaatataactgtgaat
   tataatggtagcacttataaatgaaactgtcttggagtttgactggaagtcatatattcaagattcaaga
   tatcattatttctcctggagggttttttagatgcactactatgtggaaaatagctctgatgcaggccaatgt
20  ccagagggatataatgtgtgtgaaagctggtagaaaatcccaattatggctacacaagctttgataccttc
   agttgggctttttgtccttgtttcgactaatgactcaggacttctgggaaaatctttatcaactgaca
   ttacgtgctgctgggaaaacgtacatgataatttttgtattggtcatttttcttgggctcattctaccta
   ataaatttgatcctggctgtggtggccatggcctacgaggaacagaatcaggccaccttggagaagca
   gaacagaaagaggccgaatttcagcagatgattgaacagcttaaaaagcaacaggaggcagctcagcag
25  gcagcaacggcaactgcctcagaacattccagagagcccagtcagcaggcaggctctcagacagctca
   tctgaagcctctaagttgagttccaagagtgtcaaggaaagaagaaatcggaggaagaaaagaaaacag
   aaagagcagtcctgggtggggaagagaaagatgaggatgaattccaaaaatctgaatctgaggacagcatc
   aggaggaaagggttttctccttccattgaagggaaccgattgacatatgaaaagaggtaactcctcccca
   caccagtctttgttgagcatccgtggctccctattttcaccaaggcgaaatagcagaacaagccttttc
30  agcttttagagggcgagcaaaaggatgtgggatctgagaacgacttcgcagatgatgagcacagcaccttt
   gaggataacgagagccgtagagattccttgtttgtgccccgacgacacggagagagacgcaacagcaac
   ctgagtcagaccagtaggtcatcccggatgctggcagtggttccagcgaatgggaagatgcacagcact
   gtggattgcaatgggtgtggtttccttgggtgggaccttcagttcctacatcgctgttggacagctt
   ctgccagaggtgataatagataagccagctactgtacaaatggaacaaccactgaaactgaaatgaga
35  aagagaagggtcaagttctttccacggttccatggactttctagaagatccttcccaaaggcaacgagca
   atgagtatagccagcatttcaacaaatacagtagaagaacttgaagaatccaggcagaaatgccaccc
   tgttgggtataaaattttccaacatattcttaactctgggactgttctccatatttggttaaaagtgaacat
   gttgtcaacctggttgtgatggaccatttgttgacctggccatcaccatctgtattgtcttaataact
   cttttcatggccatggagcactatccaatgacggaccatttcaataatgtgcttacagtaggaaacttg
40  gttttcactgggatctttacagcagaaatgtttctgaaaattattgcatggatccttactattatttc
   caagaaggctggaatatctttgacgggttttattgtgacgcttagcctggtagaacttggaactcgccaat
   gtggaaggattatctgttctccgttcatttccgattgtcgcaggttttcaagttggcaaaatcttggcca
   acgttaaatatgtctaataaagatcatcggcaattccgtgggggctctgggaaatttaaccctcgtcttg
   gccatcatcgtcttcatTTTTTgcccgtggctcggcatgcagctctttggtaaaaagctacaaagattgtgtc
45  tgcaagatcgccagtgattgtcaactcccacgctggcacatgaatgacttcttccactccttctgatt
   gtgttccgcgtgctgtgtggggagtgatagagaccatgtgggactgtatggagggttctggtcaagcc
   atgtgccttactgtcttcatgatggtcatggtgattggaaacctagtggtcctgaatctctttctggcc
   ttgcttctgagctcatttagtgacagacaaccttgcagccactgatgatgataatgaaatgaataatctc
   caaattgctgtggataggtatgcacaaaggagtagcttatgtgaaaagaaaaatatgaatttattcaa
50  cagtccttcattaggaaacaaaagatttttagatgaaattaaaccacttgatgatctaaacaacaagaaa
   gacagttgtatgtccaatcatacagcagaaattgggaaagatcttgactatcttaagatgtaaatgga
   actaagaagtggtataggaaactggcagcagtggttgaaaaatacattattgatgaaagtgttacatgtca
   ttcataaacaacccccagtccttactgtgactgtaccaattgctgtaggagaatctgactttgaaaattta
   aacacggaagacttttagtagtgaatcggtatctggaagaaagcaaaagagaaactgaatgaaagcagtagc
55  tcatcagaaggtagcactgtggacatcggcgacacctgtagaagaacagcccgtagtggaacctgaagaa
   actcttgaaccagaagcttgtttcactgaaggctgtgtacaaaagattcaagtggtgtcaaatcaatgtg
   gaagaaggcagaggaaaacaatggtggaacctgagaaggacgtgtttccgaatagttgaacataactgg

```

tttgagaccttcattgttttcatgattctccttagtagtggtgctctggcatttgaagatatatatatt
 gatcagcgaaagacgattaagacgatgttggaatatgctgacaaggttttcacttacattttcattctg
 gaaatgcttctaaaatgggtggcatatggctatcaaacatatattcaccaatgcctgggtgttggtggac
 ttcttaattgttgatgtttcattggtcagtttaacagcaaatgccttgggttactcagaacttgagacc
 5 atcaaactctcaggaactaagagctctgagacctctaagagccttatctcgatttgaagggatgagg
 gtgggttgtaatgcccttttaggagcaattccatccatcatgaatgtgcttctggtttgtcttatattc
 tggctaattttcagcatcatgggcgtaaatttggttgctggcaaattctaccactgtattaacaccaca
 actggtgacaggtttgacatcgaagacgtgaataatcatactgattgcctaaaaactaatagaagaaat
 gagactgctcgatggaaaaatgtgaaagtaaactttgataatgtaggatttgggtatctctctttgctt
 10 caagttgccacattcaaaggatggatggatataatgtatgcagcagttgattccagaaatgtggaactc
 cagcctaagtatgaagaaagtctgtacatgtatctttactttgttattttcatcatctttgggtccttc
 ttacacttgaacctgtttattggtgtcatcatagataatttcaaccagcagaaaaagaagtttgagggt
 caagacatctttatgacagaagaacagaagaataactataatgcaatgaaaaaattaggatcgaaaaaa
 ccgcaaaagcctatacctcgaccaggaaacaaatttcaaggaatggtctttgacttcgtaaccagacaa
 15 gtttttgacataagcatcatgattctcatctgtcttaacatggtcacaatgatggtggaaacagatgac
 cagagtgaatatgtgactaccattttgtcacgcatcaatctggtgttcattgtgtctatttactggagag
 tgtgtactgaaactcatctctctacgccattattattttaccatttggtggaatatttttgattttgtg
 gttgtcattctctccattgtaggtatgtttcttgccgagctgatagaaaagtatttctgtctccctacc
 ctgttccgagtgtaccgtcttgctaggattggccgaatcctacgtctgatcaaaggagcaaaggggatc
 20 cgcaagctgctctttgctttgatgatgtcccttctcgttggtttaacatcggcctcctactcttctta
 gtcatgttcatctacgccatctttgggatgtccaaactttgcctatgttaagaggggaagttgggatcgat
 gacatgttcaactttgagacctttggcaacagcatgatctgcctattccaaattacaacctctgtggc
 tgggatggattgttagcaccattctcaacagtaagccaccgactgtgaccctaataaaagttaaccct
 ggaagctcagttaagggagactgtgggaaccatctgttggaattttctttttgtcagttacatcatc
 25 atatccttctcgtgtgtgtgaacatgtacatcgcggtcatctctggagaacttcagtggtgtactgaa
 gaaagtgcagagcctctgagtgaggatgactttgagatgttctatgaggtttgggagaagttgatccc
 gatgcaactcagttcatggaatttgaaaaattatctcagtttgcagctgcgcttgaaaccgcctctcaat
 ctgccacaaccaaaacaaactccagctcattgccatggatttgcctatgggtgagtggtgaccggatccac
 tgtcttgatatcttattttgcttttcaaaagcggttctaggagagagtggagagatggatgctctacga
 30 atacagatggaagagcgattcatggcttccaatccttccaagggtctcctatcagccaatcactactact
 ttaaaacgaaaaacaagaggaagtatctgctgtcattattcagcgtgcttacagacgccaccttttaag
 cgaactgtaaaaacaagcttctcttacgtacaataaaaaacaaaatcaaaggtggggctaattctcttata
 aaagaagacatgataattgacagaataaatgaaaactctattacagaaaaaactgatctgaccatgtcc
 actgcagcttgtccaccttctatgaccgggtgacaaagccaattgtggaaaaacatgagcaagaaggc
 35 aaagatgaaaaagccaaagggaataa (SEQ ID NO: 2).

The sodium channel Nav1.1 encoded by the *SCN1A* gene is expressed in
 multiple distinct neuronal populations in the cortex. These include 3 non-overlapping
 neuronal populations: fast-spiking cortical interneurons expressing parvalbumin (PV
 cINs), dis-inhibitory cortical interneurons expressing the vaso-intestinal peptide (VIP
 40 cINs) and layer 5 pyramidal neurons.

The amino acid sequence of the unmodified human muscarinic acetylcholine
 receptor M3 is provided under NCBI Reference Sequence NP_000731.1 as set forth
 below. Also encompassed herein is a polypeptide or protein or functional fragment
 thereof having at least about or equal to 85%, or at least about or equal to 90%, 95%,
 45 98%, 99%, or greater, amino acid sequence identity to the following amino acid
 sequence:

1 MTLHNNSTTS PLFPNISSSW IHSPSDAGLP PGTVTHFGSY NVSRAAGNFS SPDGTTDDPL
 61 GGHTVWQVVF IAFLTGILAL VTIIGNILVI VSFKNKQLK TVNNYFLLSL ACADLIIGVI
 121 SMNLFTTYII MNRWALGNLA CDLWLAIIDYV ASNASVMNLL VISFDRYFSI TRPLTYRAKR

181 TTKRAGVMIG LAWVISFVLW APAILEWQYF VGKRTVPPGE CFIQFLSEPT ITFGTAIAAF
 241 YMPVTIMTIL YWRIYKETEK RTKELAGLQA SGTEAETENF VHPTGSSRSC SSYELQQQSM
 301 KRSNRKYGR CHFWFTTKSW KPSSEQMDQD HSSSDSWNNN DAAASLENSA SSDEEDIGSE
 361 TRAIYSIVLK LPGHSTILNS TKLPSSDNLQ VPEEELGMVD LERKADKLQA QKSVDGGSF
 5 421 PKSFSKLPIQ LESAVDTAKT SDVNSSVGKS TATLPLSFKE ATLAKEFALK TRSQITKRKR
 481 MSLVKEKKAA QTLSAILLAF IITWTPYNIM VLVNTFCDSC IPKTFWNLYG WLCYINSTVN
 541 PVCYALCNKT FRTTFKMLLL CQCDKKKRRK QQYQQRQSVI FHKRAPEQAL
 (SEQ ID NO: 3).

The amino acid sequence of the human Gq-DREADD (hM3Dq) excitatory
 10 receptor is derived from the amino-acid sequence of the unmodified human
 muscarinic acetylcholine receptor M3 set forth above. In the Gq-DREADD (hM3Dq)
 receptor amino acid sequence (590 aa), the tyrosine in position 149 is replaced by a
 cysteine, and the arginine in position 239 is replaced by a glycine (US Publication No.
 2018/0078658), as shown below:

15 Met Thr Leu His Asn Asn Ser Thr Thr Ser Pro Leu Phe Pro Asn Ile Ser Ser Ser Trp
 Ile His Ser Pro Ser Asp Ala Gly Leu Pro Pro Gly Thr Val Thr His Phe Gly Ser Tyr
 Asn Val Ser Arg Ala Ala Gly Asn Phe Ser Ser Pro Asp Gly Thr Thr Asp Asp Pro Leu
 Gly Gly His Thr Val Trp Gln Val Val Phe Ile Ala Phe Leu Thr Gly Ile Leu Ala Leu
 Val Thr Ile Ile Gly Asn Ile Leu Val Ile Val Ser Phe Lys Val Asn Lys Gln Leu Lys Thr
 20 Val Asn Asn Tyr Phe Leu Leu Ser Leu Ala Cys Ala Asp Leu Ile Ile Gly Val Ile Ser
 Met Asn Leu Phe Thr Thr Tyr Ile Ile Met Asn Arg Trp Ala Leu Gly Asn Leu Ala Cys
 Asp Leu Trp Leu Ala Ile Asp Cys Val Ala Ser Asn Ala Ser Val Met Asn Leu Leu Val
 Ile Ser Phe Asp Arg Tyr Phe Ser Ile Thr Arg Pro Leu Thr Tyr Arg Ala Lys Arg Thr
 Thr Lys Arg Ala Gly Val Met Ile Gly Leu Ala Trp Val Ile Ser Phe Val Leu Trp Ala
 25 Pro Ala Ile Leu Phe Trp Gln Tyr Phe Val Gly Lys Arg Thr Val Pro Pro Gly Glu Cys
 Phe Ile Gln Phe Leu Ser Glu Pro Thr Ile Thr Phe Gly Thr Ala Ile Ala Gly Phe Tyr
 Met Pro Val Thr Ile Met Thr Ile Leu Tyr Trp Arg Ile Tyr Lys Glu Thr Glu Lys Arg
 Thr Lys Glu Leu Ala Gly Leu Gln Ala Ser Gly Thr Glu Ala Glu Thr Glu Asn Phe Val
 His Pro Thr Gly Ser Ser Arg Ser Cys Ser Ser Tyr Glu Leu Gln Gln Gln Ser Met Lys
 30 Arg Ser Asn Arg Arg Lys Tyr Gly Arg Cys His Phe Trp Phe Thr Thr Lys Ser Trp Lys
 Pro Ser Ser Glu Gln Met Asp Gln Asp His Ser Ser Ser Asp Ser Trp Asn Asn Asn Asp
 Ala Ala Ala Ser Leu Glu Asn Ser Ala Ser Ser Asp Glu Glu Asp Ile Gly Ser Glu Thr
 Arg Ala Ile Tyr Ser Ile Val Leu Lys Leu Pro Gly His Ser Thr Ile Leu Asn Ser Thr Lys
 Leu Pro Ser Ser Asp Asn Leu Gln Val Pro Glu Glu Glu Leu Gly Met Val Asp Leu
 35 Glu Arg Lys Ala Asp Lys Leu Gln Ala Gln Lys Ser Val Asp Asp Gly Gly Ser Phe Pro

Lys Ser Phe Ser Lys Leu Pro Ile Gln Leu Glu Ser Ala Val Asp Thr Ala Lys Thr Ser
 Asp Val Asn Ser Ser Val Gly Lys Ser Thr Ala Thr Leu Pro Leu Ser Phe Lys Glu Ala
 Thr Leu Ala Lys Arg Phe Ala Leu Lys Thr Arg Ser Gln Ile Thr Lys Arg Lys Arg Met
 Ser Leu Val Lys Glu Lys Lys Ala Ala Gln Thr Leu Ser Ala Ile Leu Leu Ala Phe Ile
 5 Ile Thr Trp Thr Pro Tyr Asn Ile Met Val Leu Val Asn Thr Phe Cys Asp Ser Cys Ile
 Pro Lys Thr Phe Trp Asn Leu Gly Tyr Trp Leu Cys Tyr Ile Asn Ser Thr Val Asn Pro
 Val Cys Tyr Ala Leu Cys Asn Lys Thr Phe Arg Thr Thr Phe Lys Met Leu Leu Leu
 Cys Gln Cys Asp Lys Lys Lys Arg Arg Lys Gln Gln Tyr Gln Gln Arg Gln Ser Val Ile
 Phe His Lys Arg Ala Pro Glu Gln Ala Leu (SEQ ID NO: 4).

10 Unless specifically stated or obvious from context, as used herein, the term
 "or" is understood to be inclusive. Unless specifically stated or obvious from context,
 as used herein, the terms "a", "an", and "the" are understood to be singular or plural.

As used herein, the term "about" or "approximately" means within an
 acceptable error range for the type of value described and the method used to measure
 15 the value. For example, these terms can signify within 20%, more preferably within
 10%, and most preferably still within 5% of a given value or range. More
 specifically, "about" can be understood as within 20%, 10%, 9%, 8%, 7%, 6%, 5%,
 4%, 3%, 2%, 1%, 0.5%, 0.1%, 0.05%, or 0.01% of the stated value or range.
 Alternatively, especially in biological systems, the term "about" means within one log
 20 unit (i.e., one order of magnitude), preferably within a factor of two of a given value.
 Unless specifically stated or obvious from context, as used herein, the term "about" is
 understood as within a range of normal tolerance in the art, for example within 2
 standard deviations of the mean. Unless otherwise clear from context, all numerical
 values provided herein are modified by the term about.

25 The recitation of a listing of chemical groups or component groups in any
 definition of a variable herein includes definitions of that variable as any single group
 or combination of listed groups. The recitation of an embodiment for a variable or
 aspect herein includes that embodiment as any single embodiment or in combination
 with any other embodiments or portions thereof as described in the disclosure.

30 Any compositions or methods provided herein can be combined with one or
 more of any of the other compositions and methods provided herein.

BRIEF DESCRIPTION OF THE DRAWINGS

FIGS. 1A-1, 1A-2, 1A-3, 1B-1, 1B-2, 1C and 1D present tabular data and information related to the discovery and identification of specific enhancer (regulatory) sequences, called “E1-E35” herein. Shown are enhancers specific for *SCN1A*-restricted gene expression, such as E1-E10, in GABAergic interneurons, such as PV-expressing interneurons, as well as enhancers that target other genes presented in the tables. **FIG. 1A-1** presents tabular data depicting gene, target (e.g., neuronal cell type), specificity, position (e.g., intergenic or intronic), chromosome location and genome sequence start and stop site characteristics of thirty-five (35) enhancer elements, called E1-E35, in the mouse genome. Similarly, **FIGS. 1A-2 and 1A-3** present tabular data depicting the gene, target (e.g., neuronal cell type), specificity, position (e.g., intergenic or intronic), chromosome location and genome sequence start and stop site characteristics of these thirty-five (35) E1-E35 enhancer elements in the human genome. By way of example, enhancer (regulatory) elements E1-E10 (also called S5E1-S5E10 herein) were identified in the vicinity of the human *SCN1A* gene, in the mouse genome (**FIG. 1A-1**) and in the human genome (**FIGS. 1A-2 and 1A-3**). In **FIGS. 1A-1 to 1A-3**, the polynucleotide sequences of the mouse and human enhancer elements described herein have the start and stop sites in the mouse and human genomes as set forth in the tables (as well as in the tables of **FIGS. 15A-1, 15A-2, 16A-1 and 16A-2**); the mouse and human enhancer sequences are provided via the web-accessible genome information listed in the tables of **FIGS. 1A-1 to 1A-3**. **FIGS. 1B-1 and 1B-2** present images showing E1-E10 enhancer element-restricted reporter gene expression in PV-expressing interneurons in cortical layers of brain. The images show the results of immunohistochemical (IHC) staining analysis for dTomato in brain sections following systemic *in vivo* injection of the pAAV-S5-E2-dTomato vector into an animal (mouse), allowing for detection of specific cells transduced by the vector. **FIGS. 1C and 1D** show graphs depicting the quantification of the degree of the specificity (**FIG. 1C**) and sensitivity (**FIG. 1D**) of expression of a reporter gene in PV-expressing interneurons in the cortex. The expression of the reporter gene is controlled by the E1-E10 enhancer elements contained in rAAV vectors. The specificity was quantified as the proportion of cells expressing the viral reporter dTomato co-expressing the PV-interneuron marker PV assessed by

immunohistochemistry on brain sections following systemic *in vivo* injection of the pAAV-S5-E2-dTomato vector into an animal (mouse). The sensitivity was quantified as the proportion of cells expressing the PV-interneuron marker PV that co-expressed in the viral reporter dTomato as assessed by immunohistochemistry on brain sections following systemic *in vivo* injection of the pAAV-S5-E2-dTomato vector into an animal (mouse). Bar graphs represent mean +/- standard error of the mean (s.e.m.).

FIGS. 2A and 2B present images showing localization of reporter gene expression, using rAAV vectors containing the E2 enhancer element sequence and reporter transgene (e.g., d-Tomato) or an effector gene (e.g. Gq-DREADD) across brain structures including the cortex. **FIG. 2A** presents an image showing the results of immunohistochemical (IHC) staining analysis for the dTomato reporter in brain sections (sagittal sections in the top portion of the figure; coronal sections in the lower portions of the figure) following systemic *in vivo* injection of the pAAV-S5-E2-dTomato vector into an animal (mouse), allowing for detection of specific cells transduced by the vector. **FIG. 2B** presents images showing the results of immunohistochemical (IHC) staining analysis for the dTomato reporter expressed in brain sections following systemic *in vivo* injection of the pAAV-S5-E2-dTomato vector, or into an animal (mouse), allowing for detection of specific cells expressing PV. Reporter gene expression from the pAAV-S5-E2-dTomato vector is visualized in brain sections (in red in the left panel of **FIG. 2B**). Reporter gene expression from the pAAV-S5-E2-Gq-DREADD-dTomato is visualized in green for Gq-DREADD and red for dTomato in the right panel of **FIG. 2B**). Detection of specific PV-expressing cells transduced by the vector is visualized (in the left panel of **FIG. 2B**, and in the right panel of **FIG. 2B**).

FIGS. 3A-3F show schematics, plots, graphs and confocal microscope images related to the identification of SCN1A enhancers. **FIG. 3A** provides a schematic representation of the scATAC-seq pipeline. Interneurons were collected from the visual cortex of adult *Dlx6a*^{Cre}::*Sun1-eGFP* mice. **FIG. 3B** shows a plot of the 3500 nuclei in UMAP space. The clusters obtained from the SnapATAC pipeline were lumped into the four cardinal classes of interneurons. **FIG. 3C** presents a Venn diagram showing the numbers of unique and shared peaks across the four interneuron populations, PV, SST, VIP and ID2. **FIG. 3D** shows a schematic representation of

the enhancer selection method at the *SCN1A* locus, as described in the Methods herein (Example 8). **FIGS. 3E and 3F** illustrate results obtained following the systemic injection of adult mice with the indicated rAAV-E[x]-dTomato vector containing an enhancer element as described and the analysis 3 weeks post-injection.

- 5 Immunohistochemical (IHC) evaluation for the reporter and indicated markers in the S1 cortex was used to assess the strength of expression of the reporter (**FIG. 3E**, upper panel) and the specificity of expression of the viral reporter for the indicated markers (all other panels). Representative fluorescent images of the indicated viral reporter in the somatosensory cortex (**FIG. 3F**, left panels). Dashed lines represent
10 the limits of anatomical structures. Scale bars represent 100 μ m. On the graphs, the dots represent individual measurements and the lines represent average $\pm$ s.e.m.

- FIGS. 4A-4E** presents images, graphs and recording traces related to viral targeting of PV cortical interneurons (PV cINs) in mice. Adult mice were injected systemically (**FIGS. 4A-4B** images) or locally (**FIG. 4D**) with rAAV-E2-dTomato
15 expressing the reporter dTomato under the control of the E2 regulatory element and analyzed 3 weeks post-injection by immunohistochemistry (IHC) or ISH for both the reporter and the PV marker. **FIG. 4C** shows a slice recoding of the intrinsic properties of virally labeled neurons. **FIG. 4D** (right panel) presents a graph illustrating the specificity of expression shown as the proportion of cells expressing
20 the reporter that co-express the PV relative to the strength of expression of the reporter. **FIG. 4E** presents images resulting from experiments in which mice were injected locally with rAAV-E2-dTomato expressing the reporter dTomato under the control of the E2 regulatory element and analyzed at the indicated developmental stages for the reporter and the indicated markers. Scale bars represent 250 μ m (**FIG.**
25 **4A**) and 50 μ m (**FIGS. 4B, 4D, 4E**). In the graphs, the dots represent individual measurements, and the lines represent average $\pm$ s.e.m.

- FIGS. 5A-5E** present images, current clamp recording traces and graphs related to viral monitoring and manipulation of PV cortical interneurons (PC cINs) in mice. Mice were injected locally in the somatosensory (S1) cortex with rAAVs (**FIG.**
30 **5A** – P10 injection with rAAV-E2-SYP-dTomato; **FIG. 5B** – P14 injection with rAAV-E2-GCaMP6f; **FIGS. 5D and 5E** – Adult injection with rAAV-E2-C1V1-eYFP), or systemically (**FIG. 5C** – Adult injection with rAAV-E2-PSAM4-5HT3-

LC-GFP). **FIG. 5A** presents representative images of the co-localization between the SYP-dTomato reporter and the synaptic marker Syt2 one-week post-injection and corresponding quantification. **FIG. 5B** shows results of Ca^{2+} imaging upon whisker stimulation performed 2-3 weeks post-injection. In the right panel, the success rate was calculated as the proportions of $\Delta F/F$ peaks above threshold in response to whisker stimulation. **FIG. 5C** shows the results of current clamp recording performed on brain sections 4 weeks after injection. The traces show a representative cellular response at the indicated currents at both baseline and after bath application of varenicline. **FIG. 5D** shows the results of current clamp recording performed on brain sections 1 week after injection. Cells expressing the viral reporter were exposed to 2 seconds of constant laser stimulation (550 nm) while the voltage was recorded over 3 seconds. Neighboring pyramidal cells that did not express the viral reporter were also recorded from during laser stimulation. **FIG. 5E** illustrates *in vivo* single-unit analysis of neuronal activity and shows Raster plots of virally infected neurons upon laser stimulation and corresponding population quantification data. The left panels show fast-spiking cells and the right panels show regular spiking excitatory cells. Notably, due to the mosaic nature of local viral injection, individual cell responses were bimodal. This likely reflects whether or not particular cells were infected. Scale bars represent 5 μm . The middle bars at the top of the “Trial” versus “Time” graphs represent laser stimulation. In the graphs, dots represent individual measurements and the lines represent average $\pm$ s.e.m.

FIGS. 6A and 6B present drawings, graphs, images and recording traces related to viral targeting and manipulation PV cortical interneurons (PV cINs) in primates, including humans. **FIG. 6A:** Animals from indicated species were locally (rat and macaque) or systemically (marmoset) injected with rAAV-E2-C1V1-eYFP (macaque) or rAAV-E2-dTomato (rat and marmoset) and analyzed 2-8 weeks post-injection. The specificity of expression is shown as the proportion of virally labeled cells co-expressing PV. **FIG. 6B:** Human brain tissue obtained from surgical resection was exposed to either rAAV-E2-dTomato **(i-iii)** or rAAV-E2-C1V1-eYFP **(iv)** and maintained in culture for 7-14 days. The upper right panel shows the proportion of fast-spiking neurons among the virally-labeled cells assessed by electrophysiological recordings of intrinsic properties. **(iv)** Electrophysiology current

clamp recording of virally labeled cells upon laser stimulation. Scale bars represent 25 μm . The bar at the top of the “Direct photoactivation (PV)” trace represents laser stimulation and the arrowheads point at neurons co-expressing PV and the viral reporter. On the graphs, dots represent individual measurements and the lines

5 represent average $\pm$ s.e.m.

FIG. 7 depicts fluorescent images of sagittal sections from adult mice that were injected systemically with the indicated rAAV-E[x]-dTom viral reporter vector and analyzed 3 weeks post-injection with IHC for the viral reporter. Scale bar represents 500 μm .

10 **FIGS. 8A-8D** present images and graphs of results following systemic injection of adult mice with rAAV-E2-dTomato. **FIG. 8A** relates to slice recording of the intrinsic properties of virally labeled neurons. The left panel shows a representative cell expressing the viral reporter. The triangular trace in the middle, top represents the recording pipet. The quantifications show the indicated parameters.

15 The darker gray dots in the “Identity” graph represent cells with stereotypical fast-spiking (FS) properties. **FIG. 8B** shows representative slice recording traces of positive and negative fast-spiking cells (FS and nFS, respectively). Scale bars represent 20 μm . On the graphs, dots represent individual measurements and the lines represent average $\pm$ s.e.m. **FIGS. 8C and 8D** show results following systemic

20 injection of adult mice with rAAV-E2-dTomato and analysis 3 weeks post-injection. **FIG. 8C**: Coronal and sagittal sections were analyzed with IHC for the viral reporter and PV and the specificity to PV was reported across brain regions. **FIG. 8D**: The native viral expression was analyzed from the indicated organs. Scale bars represent 100 μm (**FIG. 8C**) and 250 μm (**FIG. 8D**). On the graphs, dots represent individual

25 measurements and the lines represent average $\pm$ s.e.m.

FIGS. 9A-9C present images, recording trace data and graphs. Mice were injected systemically (**FIG. 9A**: P14 injection with rAAV-E2- GCaMP6f) and locally (**FIG. 9B**: rAAV-E2-C1V1-eYFP; **FIG. 9C**: rAAV-E2- GqDREADD) in the somatosensory cortex. **FIG. 9A**: Mice were analyzed 1-week post-injection. The left

30 panel shows widefield images of two representative peaks shown by the pound sign in the middle panels. The right panel shows a fluorescent image taken after GCaMP recordings. **FIG. 9B**: Slice electrophysiology current clamp recording were

performed 1-week post-injection. Cells expressing the viral reporter were targeted with either 10Hz or 40Hz laser stimulation (550 nm) while the voltage was recorded over 3 seconds. **FIG. 9C:** Slice electrophysiology current clamp recordings were performed 1-week post- injection. The voltage was recorded before and after bath application of CNO. Scale bars represent 500 μ m. The “+CNO” bars represent laser stimulation. On the graphs, dots represent individual measurements.

FIGS. 10A and 10B present stained images and data plots related to studies in which human brain tissue obtained from surgical resection was exposed to either AAV-E2-dTomato and maintained in culture for 7-14 days. **FIG. 10A:**

Representative image of the dendrites of virally labeled cells filled with Biocytin during the recording session. **FIG. 10B:** Slice recording of the intrinsic properties of virally labeled neurons. The quantifications show the indicated parameters. The darker, rightmost dots in the “Identity” graph represent cells with stereotypical fast-spiking (FS) properties. Scale bar represent 100 μ m. In the graphs, the dots represent individual measurements and the lines represent average $\pm$ s.e.m.

FIG. 11 provides a table showing quantifications of cells expressing markers/reporters. As described in Example 7, quantifications were performed using a minimum of two independent biological replicates, and the specific numbers of cells and conditions are indicated for each individual quantification in the table.

FIG. 12 presents UMAP plots of 3500 neuronal nuclei collected from 4 *Dlx6a^{Cre}::Sun1*- GFP mice reflecting promoter accessibility of the indicated canonical interneuron markers.

FIGS. 13A and 13B present slices, images and graphs related to the identification of viral enhancers with regional specificity. **FIG. 13A:** Adult mice were injected systemically with the indicated rAAV vector containing an enhancer element polynucleotide sequence and a detectable reporter or marker (e.g., GFP) polynucleotide, i.e., rAAV-E[x]-eGFP, and analyzed 3 weeks post-injection. Immunohistochemistry (IHC) for the reporter and indicated markers in the S1 cortex was used to assess the density of neuronal cell-bodies expressing the viral reporter (left panels) and the specificity of expression of the viral reporter for the indicated markers (right panels). For the E29 virus, no cell bodies are observed in the thalamus, with the exception of the thalamic reticular nucleus (TRN). **FIG. 13B:** An adult

macaque was injected in V1 with rAAV-E22-eGFP and analyzed 8 weeks post-injection with IHC for the reporter and indicated markers. Scale bars represent 100 μm (a), 50 μm (b, left) and 10 μm (b, right). On the graphs, dots represent individual measurements and the lines represent average $\pm$ s.e.m.

5 **FIG. 14** presents images and a graph related to studies in which adult mice were injected with the indicated modified rAAV-E2-dTomato construct and analyzed 3 weeks post-injection with IHC for the viral reporter and PV. The corresponding specificity is shown in the graph at the right. Scale bars represent 2 μm . On the graphs, dots represent individual measurements and the lines represent average $\pm$ s.e.m.

10 **FIGS. 15A-1 and 15A-2** present a table containing the specifications for all tested enhancers, including their associated gene, target population, specificity for target population, location, presence of ATAC peaks, and conservation with the human sequence.

15 **FIGS. 16A-1 and 16A-2** present a table that compiles various parameters related to each of the tested enhancers, including enhancer name (E1-E35), gene, target, % specificity, murine chromosome location (Mouse_mm10_Chr), enhancer sequence start site in murine genome (Mouse_mm10_Start), enhancer sequence stop site in murine genome (Mouse_mm10_Stop), size (base pairs (bp)), human
20 chromosome location (Human_hg38_Chr), enhancer sequence start site in human genome (Human_hg38_Start), enhancer sequence stop site in human genome (Human_hg38_Stop), and Percentage of conservation between the mouse and human enhancer sequences. In the tables presented in FIGS. 15A-1 and 15A-2, and FIGS. 16-A1 and 16-A2, as well as in FIGS. 1A1 to 1A3, the total number of base pairs (bp)
25 shown for the polynucleotide sequence of each of the listed enhancers E1-E35 reflects that the numerical value of the first base pair (bp) counted in the sequence is valued at zero (0), as would be appreciated by one skilled in the art. Notwithstanding, the total number of bp comprising the polynucleotide sequence of each enhancer (E1-E35) is obtainable simply by counting the number of total bp in the sequence, based on the
30 tabularized data presented in the figures described herein.

DETAILED DESCRIPTION OF THE EMBODIMENTS

The embodiments featured and described herein relate to strategies, methods and products developed to identify multiple new enhancers, (E1-E35), for use with viral vectors, such as recombinant adeno-associated virus (rAAV) vectors, for example, to target functionally distinct neuronal subtypes, particularly, within the cerebral cortex. Investigation of the regulatory landscape of the disease gene *SCN1A* led to the identification of enhancers that target the breadth of its expression, including, by way of nonlimiting example, two enhancers that are selective for parvalbumin (PV) and vasoactive intestinal peptide (VIP) cortical interneurons. The functional utility of these regulatory elements was demonstrated, and it was found that the PV-specific enhancers allowed for the selective targeting and manipulation of these neurons across species, from mice to humans. Moreover, the selection method as described herein is generalizable to other genes and characterized certain PV-specific enhancers, such as, for example, E11, E14, E22 and E29, which have a high degree of specificity for distinct regions of the brain. Recombinant viral vectors, e.g., rAAV vectors, harboring the enhancer sequences provide viral tools for use in cell-type specific circuit manipulation and in therapeutic interventions to treat and ameliorate neuropathological or neuropsychiatric diseases, conditions and pathologies.

Specific viral-based therapeutic products, compositions, methods and approaches for treating or ameliorating neurological, neurodevelopmental, neurogenetic, or neuropsychiatric diseases, disorders, and pathologies are described herein. As described, virus vectors and vehicles for gene delivery are designed and produced to contain a specific enhancer sequence (enhancer) and a polynucleotide sequence of a gene of interest, such as an effector gene (e.g., a transgene or reporter gene), which is specifically and functionally expressed in specific interneuron or neuron cell populations following transduction of the interneuron or neuron cells by the virus vector or vehicle. In an embodiment, a virus vector or vehicle is provided which comprises the polynucleotide of a specific enhancer sequence (enhancer), which is specifically and functionally expressed in specific interneuron or neuron cell populations following transduction of the interneuron or neuron cells by the virus vector or vehicle. In an embodiment, the enhancer harbored by the virus is capable of

restricting the expression of the transgene to certain interneuron cells or neuronal cells. In embodiments, expression of the transgene is restricted to expression in cells that are deficient for that gene. In an embodiment, the expression of the transgene is specifically modulated in the interneuron cell or other neuronal cell. In other
5 embodiments, the transgene is an effector gene or a therapeutic gene. In
embodiments, the enhancer element restricts expression of a gene to one or more neuronal cell types, including a parvalbumin (PV)-expressing cortical interneuron cell (PV-cIN cell), which is a fast-spiking cortical interneuron; a dis-inhibitory cortical interneuron cell expressing vaso-intestinal peptide (VIP), (VIP cIN cell); and a
10 pyramidal (PYR) neuron, in particular, a pyramidal neuron of cortical layer 5 of the brain.

In an embodiment, the virus vector contains a specific enhancer sequence and a transgene (effector gene) associated with a neurological, neurodevelopmental or neurogenetic disease, disorder, or condition, and the enhancer is capable of restricting
15 the expression of the transgene to an interneuron cell population that has loss-of-function for the gene, is deficient for the gene, or that expresses a mutant, variant, or defective form of the gene associated with the neurological or neurogenetic disease, disorder, and pathology. In a particular embodiment, the enhancer sequence inserted in the virus vector polynucleotide is identified as one having specificity for regulating
20 the expression of the *SCN1A* gene, which encodes the Nav1.1 sodium channel, and restricting expression to *SCN1A*-expressing cells, in particular, GABAergic interneuron cells. Loss of function of the *SCN1A* gene is the most prevalent cause of the debilitating disease Dravet syndrome (DS), which is a pharmaco-resistant form of infantile epilepsy associated with cognitive impairment and premature death. In
25 certain embodiments, the specific expression of the transgene (effector gene) in interneurons may be determined by the detection of markers that are specific for interneuron cells, e.g., without limitation, GABA GAD67, or PV interneuron cell markers. In an embodiment, the virus vector or vehicle is an adeno-associated virus (AAV) or a recombinant AAV (rAAV). The terms “AAV” and “rAAV” are used
30 interchangeably herein.

The term “transgene” is used herein to refer to a gene (or genes) of interest (an effector gene) contained in the rAAV vector or vehicle as described herein and is

specifically expressed and functional in a certain cell types or populations as described herein, especially by virtue of the enhancer sequence also contained in the rAAV vector, which restricts the expression of the gene to a defined population of cells, e.g., PV-expressing or *SCN1A*-expressing interneurons or subtypes thereof. In some cases, the gene of interest (effector gene) is a normal form of a gene that is expressed in the cell type transduced by rAAV and whose encoded product functions to provide a normal or normally-functioning product in the cell, such as a cell in which there is a loss of function of the same gene as the transgene. In some cases, the transgene or effector gene may be a reporter gene, e.g., green fluorescent protein (GFP) or red fluorescent protein (RFP) that provides a detectable signal following transduction of a cell by the rAAV vector. In some cases, the transgene or effector gene may be both a reporter and a gene that encodes a product whose expression and activity provide for normal cell function. The latter type of gene may be considered to be a therapeutic gene. In a particular embodiment, the rAAV contains an *SCN1A*-specific enhancer sequence and an *SCN1A* transgene.

The rAAV vectors and methods described herein are based, at least in part, on the discovery and demonstration that a specific enhancer can restrict the expression of a transgene carried by the virus vector, such as a gene associated with a neurological disease, disorder, or pathology, or a reporter gene, to interneuron cells (“interneurons”) in the brain where the gene is expressed and the encoded gene (transgene) product is functional. In an embodiment, such an expressed, functional gene offsets, replaces, or substitutes for, the abnormal, aberrant, or lack of function of a gene encoding a product involved in the normal functioning of an interneuron cell.

In an embodiment, a suitable viral vector, e.g., a lentiviral vector or, in particular, a recombinant adeno-associated virus (rAAV) vector, is used to restrict expression of a transgene in GABA-ergic PV-expressing interneurons a mammal, in which an enhancer element as described herein provided in *cis*. In embodiments, the enhancer element is one of S5E1 (E1), S5E2 (E2), S5E3 (E3), S5E4 (E4), S5E6 (E6), S5E7 (E7), S5E8 (E8), S5E9 (E9), S5E10 (E10). In embodiments, the enhancer element is E2, which is capable of restricting the expression of a viral reporter to parvalbumin (PV)-expressing cortical interneurons (PV cINs), E6, which is selective for VIP interneurons; or E5, which labels interneuron populations across all cortical

layers, yet is especially selective for pyramidal neurons in layer 5 of the brain cortex, in particular, glutamatergic pyramidal neurons, as described herein. In a particular embodiment, the enhancer element is E2. In another particular embodiment, the enhancer element is E5. In yet another particular embodiment, the enhancer element is E6.

In an embodiment, the viral vector or rAAV vector comprising the enhancer drives the expression of a copy of *SCN1A* in a transduced PV-expressing interneuron cell for the treatment and therapy of seizures, all forms of epilepsy, or DS. In other embodiments, the vector or rAAV vector comprising the enhancer drives the expression of effectors like Gq-DREADD or PSAM for chemogenetic modulation of PV-interneuron activity for the treatment of all forms of seizures, epilepsy, including focal and pharmacologically intractable epilepsy, and also for the treatment of DS and the symptoms thereof.

In general, a viral vector or rAAV vector comprises a polynucleotide comprising an enhancer sequence selected from S5E1-S5E10 as described herein, and a transgene sequence, such as, a polynucleotide sequence encoding an *SCN1A* gene, a polynucleotide sequence encoding hM3Dq modified muscarinic receptor (Gq-DREADD) receptor, or a polynucleotide sequence encoding PSAM. In an embodiment, the polynucleotide comprises an enhancer sequence selected from E2, E5, or E6 as described herein. In certain embodiments, methods are provided for therapeutic and prophylactic treatments for seizures and epilepsy, and more specifically, Dravet syndrome, in an individual (e.g., a human patient) in need thereof.

In an embodiment, a method is provided in which an individual or subject in need, e.g., a patient afflicted with seizures, epilepsy, or DS, is administered a viral vector, such as a recombinant adeno-associated virus (rAAV) vector, comprising an enhancer sequence as described herein, such as E2, E5, or E6, and a transgene polynucleotide sequence encoding, for example, an *SCN1A*-encoding polynucleotide sequence, a hM3Dq modified muscarinic receptor (Gq-DREADD)-encoding polynucleotide sequence, or a PSAM-encoding polynucleotide sequence, such that *SCN1A*, Gq-DREADD, or PSAM, respectively, is expressed in interneurons of the individual or subject, especially in PV-expressing interneurons. Thus, a method is provided for converting interneurons, especially, PV-expressing interneurons, in an

individual or subject in need, that do not express SCN1A, Gq-DREADD, or PSAM to interneurons that do express SCN1A, Gq-DREADD, or PSAM, respectively. As such, the expression of the genes and encoded proteins is linked to the presence of the enhancer element (E1-E10) as described herein that is also provided as a component of the rAAV vector genome. In an embodiment, the enhancer element is E2, E5, or E6. In an embodiment, an individual or subject in need, e.g., a patient afflicted with seizures, epilepsy, or DS, is administered a viral vector, such as a recombinant adeno-associated virus (rAAV) vector, comprising an enhancer sequence as described herein, such as E2, and a transgene polynucleotide sequence encoding *SCN1A*.

In an embodiment, a prophylactic or therapeutic treatment method is provided for prophylaxis and/or therapy for seizures, epilepsy, or DS, which comprises introducing into an individual or subject in need a viral vector or an rAAV vector which comprises an enhancer sequence (E1-E10) as described herein, and a sequence encoding an SCN1A-encoding polynucleotide sequence such that the severity of the seizures, epilepsy, or DS symptoms experienced by the individual or subject is reduced, or the seizures, epilepsy, or DS symptoms are treated or prevented. In an embodiment, the enhancer element is E2, E5, or E6. In an embodiment, the individual or subject in need is experiencing a seizure (e.g., an epileptic seizure) or a symptom of DS at the time of administering the vector. Following administration of the vector to the individual or subject, the severity of the seizures, epilepsy, or DS symptoms is reduced, or the seizures, epilepsy, or DS symptoms are treated or prevented.

In an embodiment, a prophylactic or therapeutic treatment method is provided for prophylaxis and/or therapy for seizures, epilepsy, or DS, which comprises introducing into an individual a viral vector or an rAAV vector which comprises an enhancer sequence (E1-E10) as described herein, and a sequence encoding an hM3Dq modified muscarinic receptor (Gq-DREADD)-encoding polynucleotide sequence, and subsequently administering to the individual an effective amount of an agonist of the Gq-DREADD such that the severity of the seizures, epilepsy, or DS symptoms is reduced, or the seizures, epilepsy, or DS symptoms are treated or prevented. In an embodiment, the enhancer element is E2, E5, or E6. In an embodiment, the individual or subject in need is experiencing a seizure (e.g., and epileptic seizure) at

the time of administering the agonist of the Gq-DREADD receptor. Following administration of the agonist, the severity of the seizure is reduced. In embodiments, Gq-DREADD receptor agonist is clozapine-N4-oxide (CNO) or another suitable Gq-DREADD receptor agonist as known and used in the art.

5 In embodiments of the therapeutic and prophylactic methods described herein, the individual or subject is experiencing, or is at risk for developing, a partial seizure or a generalized seizure. In other embodiments the individual or subject has, is suspected of having, or has been diagnosed with epilepsy of any form, including, without limitation, pharmaco-resistant epilepsy. In accordance with the described
10 methods, seizures, epilepsy, or DS symptoms are inhibited, blocked, reduced, abated, or prevented.

In an embodiment, a composition comprising a viral vector or rAAV vector is administered to a subject in need thereof. In an embodiment, the administration of a composition comprising a vector (or the vector itself) comprising an enhancer
15 element, e.g., E1-E10, as described herein and a polynucleotide encoding *SCN1A* facilitates conversion of interneurons or PV-expressing interneurons of an individual or subject that do not express *SCN1A* into *SCN1A*-expressing interneurons or PV-expressing interneurons in the brain. In another embodiment, the administration of a composition comprising a vector (or the vector itself) comprising an enhancer
20 element, e.g., E1-E10, as described herein and a polynucleotide encoding Gq-DREADD receptor facilitates conversion of interneurons or PV-expressing interneurons of an individual or subject that do not express Gq-DREADD receptor into Gq-DREADD receptor-expressing interneurons or PV-expressing interneurons in the brain, thereby resulting in interneurons or PV-expressing interneurons that are
25 responsive to a Gq-DREADD agonist. In another embodiment, the administration of a composition comprising a vector (or the vector itself) comprising an enhancer element, e.g., E1-E10, as described herein and a polynucleotide encoding a PSAM facilitates conversion of interneurons or PV-expressing interneurons of an individual or subject that do not express PSAM into PSAM-expressing interneurons or PV-
30 expressing interneurons in the brain. In an embodiment, the vectors, compositions and methods as described herein are used in the prophylactic or therapeutic treatment

of partial and/or generalized seizures. In an embodiment, the enhancer element is E2, E5, or E6.

In an embodiment, the vectors, compositions and methods as described herein are used in the prophylactic or therapeutic treatment of various forms of epilepsy, including, without limitation, pharmaco-resistant epilepsy and/or may constitute a replacement of a pharmacological treatment. In embodiments, the vectors, compositions and methods as described herein are used in the prophylactic or therapeutic treatment of one or more seizure disorders, which include, but are not limited to, epilepsy, including, localization-related epilepsies, generalized epilepsies, epilepsies with both generalized and/or local seizures, and the like, seizures associated with Lennox-Gastaut syndrome, seizures as a complication of a disease or condition (such as seizures associated with encephalopathy, phenylketonuria, juvenile Gaucher's disease, Unvericht-Lundborg's progressive myoclonic epilepsy, stroke, head trauma, stress, hormonal changes, drug use or withdrawal, alcohol use or withdrawal, sleep deprivation, fever, infection, brain cancer, and the like, or chemically-induced seizure disorders.

In embodiments, the vectors or rAAV vectors, compositions and methods as described herein are used in the prophylactic or therapeutic treatment of an individual or subject in need, e.g., one who has experienced, and/or is at risk of experiencing a seizure, and thus may be diagnosed with or be suspected of having any seizure disorder. In an embodiment, administration of a viral vector or rAAV vector comprising an enhancer element as described herein and a transgene may occur at a time prior to the onset of the seizure, e.g., epileptic seizure, or DS symptom, for example, days, weeks, months, or years prior to administration. By way of example, those in the art have demonstrated that rAAV driven expression can last for at least six years in a non-human primate model (Rivera, V.M. et al., 2005, *Blood*, 105:1424-1430).

In an embodiment, the rAAV vector, which comprises an *SCN1A*-specific enhancer sequence, also comprises capsid proteins that enhance the targeting ability of the virus vector and allow the vector to specifically transduce interneuron cells, such as GABAergic interneuron cells, and/or specific subpopulations of GABAergic interneuron cells, particularly in the cerebral cortex of the brain. rAAV vectors that

transduce GABAergic interneurons and rAAV vectors that comprise capsid proteins which increase the likelihood that the virus will specifically transduce GABAergic interneurons, in particular, the subpopulation of GABAergic interneurons that also express parvalbumin (PV), called PV-expressing interneurons, (also called PV-expressing cortical interneurons) are highly suitable for use in the compositions and methods described herein. In another embodiment, the rAAV vector containing an *SCN1A*-specific enhancer sequence, e.g., E5, also comprises capsid proteins that enhance the targeting ability of the virus vector and allow the vector to specifically transduce pyramidal neurons, e.g., glutamatergic pyramidal neuron cells of the brain cortex.

In an embodiment, the transgene (effector gene) inserted into the virus vector is one whose function (or loss of function) has been found to be causally associated with a neurological disease characterized by the deleterious symptoms of seizures or epilepsy, such as infantile febrile epilepsy, or Dravet syndrome (DS). The enhancer sequence in the vector restricts expression of the transgene to interneurons or subtypes thereof, or neurons, such as pyramidal neurons, and specifically modulates, e.g., increases or enhances, the expression of a normal, functional version of this gene in an interneuron cell. In an embodiment, the interneuron cell is a GABAergic interneuron cell. In an embodiment, the interneuron GABAergic cell is a PV-expressing interneuron cell. In an embodiment, the neuron cell is a pyramidal neuron cell. In an embodiment, the pyramidal neuron cell is a glutamatergic pyramidal neuron.

In a particular embodiment, the AAV vectors, vector-based compositions, and delivery and treatment methods provided herein are useful for treating a patient who is afflicted with Dravet syndrome (DS), and the serious symptoms thereof, such as epilepsy and accompanying seizures. In an embodiment, the patient is a human patient, in particular, an infant or young child afflicted with DS. As described further *infra*, Dravet syndrome (DS) is a form of infantile epilepsy that is associated with many serious symptoms, including cognitive impairment and life-threatening seizures. The loss of function of the sodium channel Nav1.1 encoded by the *SCN1A* gene is the most prevalent cause for DS. Previous studies using mouse models of DS suggest that it is the loss of *SCN1A* gene function in GABAergic interneurons that is the

primary defect underlying the seizures that represent the most deleterious symptom in this syndrome. There is currently no reliable treatment to eliminate or reduce seizures in DS patients. Therefore, the viral products, compositions and methods as described herein provide a much needed and highly beneficial treatment for patients afflicted
5 with DS.

Accordingly, in a particular embodiment, the transgene or effector gene contained in the AAV vector or vehicle is *SCN1A* and the enhancer is a nucleic acid sequence (e.g., a *cis*-acting control element in the AAV vector) that restricts the expression of the *SCN1A* gene to *SCN1A*-expressing interneurons and is specific for
10 modulating the expression of the *SCN1A* gene in interneuron cells, e.g., GABAergic interneurons, or PV-expressing, GABAergic interneurons. In another particular embodiment, the transgene or effector gene contained in the AAV vector or vehicle is *SCN1A* and the enhancer is a nucleic acid sequence (e.g., a *cis*-acting control element in the AAV vector) that restricts the expression of the *SCN1A* gene to *SCN1A*-
15 expressing pyramidal neurons and is specific for modulating the expression of the *SCN1A* gene in pyramidal neuron cells, e.g., glutamatergic pyramidal neurons, in the brain cortex, e.g., cortical layer 5 of the brain.

Methods utilizing an AAV vector, which is designed and molecularly engineered to harbor a specific enhancer that restricts that expression of a normal
20 *SCN1A* effector gene encoding the Nav1.1 sodium channel to interneuron cells, involve administering a therapeutically effective amount of the viral vector, a viral particle, or a pharmaceutical composition comprising the viral vector or particle to a subject (e.g., a human infant having DS), in particular, to transduce interneuron cells in the subject with the vector harboring an *SCN1A*-specific enhancer sequence and an
25 *SCN1A* gene, express the gene in the interneuron cells and provide a functional response, e.g., the provision of a functional Nav1.1 sodium channel or an increase in function of the sodium channel, in interneuron cells of the subject following administration. The functional expression of *SCN1A* in the transduced interneuronal cells normalizes the excitability of *SCN1A*-deficient interneuron cell populations, such
30 as GABAergic interneurons and PV-expressing, GABAergic interneurons. Such a result restores the delicate E/I balance in regions of the brain.

To successfully and specifically express genes contained in AAV as a form of therapy for DS, an approach was developed in which the regulatory landscape of the *SCN1A* gene was explored to identify enhancer polynucleotide sequences capable of restricting expression specifically to the neuronal cell population that is deficient for this effector gene. (FIGS. 3A-3D). In an embodiment, the enhancer sequence is a *cis*-acting element that modulates, e.g., increases, enhances, augments, or otherwise improves, expression of the *SCN1A* gene, particularly in an interneuron cell, such as a GABAergic interneuron cell or a PV-expressing GABAergic interneuron cell, particularly in interneurons in which there is a loss of function of the *SCN1A* gene. In an embodiment, the enhancer sequence is a *cis*-acting element that modulates, e.g., increases, enhances, augments, or otherwise improves, expression of the *SCN1A* gene, particularly in a pyramidal neuron, such as a glutamatergic pyramidal neuron cell. The terms “enhancer” and “enhancer element” are used interchangeably herein. In some cases herein, the term “enhancer element” is referred to as a “regulatory element.”

In an embodiment, the enhancer polynucleotide sequence that specifically regulates the expression of the *SCN1A* gene in an interneuron cell is about 25-50, 50-100, 100-150, 150-200, 200-250, 250-300, 300-350, 350-400, 400-450, 450-500, 500-550, 550-600, 600-650, 650-700, 700-750, 750-800, 800-850, 850-900, 900-950, 950-1000, 1050, 1100, 1150, 1200, 1250, 1300, 1350, 1400, 1450, 1500, 1550, 1600, 1650, 1700, 1650, 1800, 1850, 1900, 1950, 2000, 2050, or 2500 nucleotides (base pairs (bp)), or longer, e.g., greater than 2500 nucleotides (bp) in length, including all larger and smaller values in between these aforementioned bp lengths. In some embodiments, PV-specific enhancer sequence suitable for use is 261 bp, 521 bp, 547 bp, 606 bp, 618 bp, 663 bp, 832 bp, 1280 bp, 1644 bp, or 2430 bp. In other embodiments, PV-specific enhancer sequence suitable for use is 267 bp, 586 bp, 636 bp, 665 bp, 844 bp, 849 bp, 894 bp, 1636 bp, 1766 bp, or 5124 bp. The enhancer sequence having specificity for modulating (e.g., enhancing) expression of the *SCN1A* gene in an interneuron cell may be derived from an intronic or intergenic sequence of genomic polynucleotide, e.g., DNA or RNA. (FIGS. 1A-1 to 1A-3).

In an embodiment, an *SCN1A*-specific enhancer sequence comprises a nucleotide sequence which contains one or more regions of 50-500 bp or longer, 50-

250 bp or longer, 100-200 bp or longer, or 100 bp or longer, having at least 70% or greater, at least 75% or greater, at least 80% or greater, at least 85% or greater, at least 90% or greater, or at least 95% or greater sequence identity to the following mouse polynucleotide (DNA) sequence (E1), also called “S5E1,” located on chromosome 2,
 5 at start/stop positions 66256056/ 66257335, shown in FIG. 1A-1, or a human ortholog thereof.

10 caaagtggacagaggggaggggaggggatgcgaggggaggggaggggaatgatcgcgaaaggcttccaa
 accttgctccttggtttttcaccattttctgaaatatatgctgagtgcactatgggaagaccattttcata
 atctataataactcctccttttaaaaggactttcgtaaccgcttgcaaaggtgagtgcgggtagaggac
 15 attagctcgctaagtccttagaaatcacacttgagactaagcaggctttcccaggagaagtcctaaagc
 caacataagcaggagggctgggggctggcgttaaccgcaaggcagtggttgagccctcgggatcatccc
 ggcgggggcgcgagcatctccgccaaggccgaggtctcaccatcagctgccgagccaccctgtacc
 tcgcagtcactcgccctgcccacgccccgcgcccgcctcaccttcagccccctgggagtcctatggcc
 gccggctacccggagggtgcccaccgctgcccgcgcaggggttaggggttcagaccacttcccgggccc
 20 ctcaaaccctaagggacgcggcgtgcagcacgaggggctggcccgatctccattggttgctggcgtga
 gggggcgaggcttagttgtaggactaggaaggagggggccaccggagcaggcgaggagggaaaccccgag
 ggaggaccgcgagggcgactggggctggaatccgctgagcattgagtgtgctgccgagttgtgggctaga
 ggaggggaggtccagcctggaaacggcgagggaggggattgggtggagcaagagatatgagattaaa
 gaataaagatgatgaagcagcaaataggagggagagccatgccgcttttcatccctgcaaacaaaggcc
 25 gactccattttctcagcattttttgtggaagccgatttgcgcaatgcggcttagtacttgaccagggaa
 aatgatttacctgacacgtgtagtaatcggtgtctgggccacaaggtggcgagaaaaatcacagttcgg
 caaaaaacctgaaacctggcttgggcttgttctaaatcttttcaggcgctgctgtaattttgctattcg
 agtgcttattaaactgctccgcccagatttccacccccaaagtcttatttataaaatatgtggttacctct
 tttagattttctattttcttaagtgtttgctgtagtttgatctaaactgtccctcaaagacacacgtgct
 30 gaatgttc ccagccccgtgtgctgttgggagtggtgga (SEQ ID NO: 5).

In an embodiment, an *SCN1A*-specific enhancer sequence comprises a nucleotide sequence which contains one or more regions of 50-500 bp or longer, 50-250 bp or longer, 100-200 bp or longer, or 100 bp or longer, having at least 70% or greater, at least 75% or greater, at least 80% or greater, at least 85% or greater, at least
 35 90% or greater, or at least 95% or greater sequence identity to the following mouse polynucleotide (DNA) sequence (E2), also called “S5E2,” located on chromosome 2, at start/stop positions 66364036/66364653, shown in FIG. 1A-1, or a human ortholog thereof.

35 aatctaacatggctgctatagcttactgactagaagttaagtgcacacttcctaaaagaaggctttgac
 acaagccacttcagttccctcctcattttctgtccccattcctctctctgtagaattctgagatttca
 attcagttttatacagaaaccacattactgtaagccctacaaagttaggcaatatagctatatggagt
 caagtaatgtaggttatttttttcccaatggtgctggtgaagggtggcaattatgtagctatacttagca
 gactgaggaaattctgctagagtcagcatttgtctcttcattgctatgaaacagtaatggaaaaataaa
 40 caaaaaacaaaggcaaacactatgcataattccctcagatcatattaacatgtgatgttggagtaaat
 gttataacccccatttttgaaatacttaccttaattaactatgatttcccttaaaataatgcagtatattac
 aatctatatgaaagcactatatgggacacatggtatgatggaacagtgacccaagagacaccaagaac

attcctgtctgtggcagtccttttctctatacagaggcatttagtctcaattgctcagagttat
(SEQ ID NO: 6).

In an embodiment, an *SCN1A*-specific enhancer sequence comprises a nucleotide sequence which contains one or more regions of 50-500 bp or longer, 50-250 bp or longer, 100-200 bp or longer, or 100 bp or longer, having at least 70% or greater, at least 75% or greater, at least 80% or greater, at least 85% or greater, at least 90% or greater, or at least 95% or greater sequence identity to the following mouse polynucleotide (DNA) sequence (E3), also called "S5E3," located on chromosome 2, at start/stop positions 66383190/66384021, shown in FIG. 1A-1, or a human ortholog thereof.

ataaaatatttttcttaaaatttacattttaaagccatccaacttaccaaagtgatttcaaaccaca
attacattttatctcaactaccagcattttatatttagccaggatctacatgagacacatatcatgatgtg
ctatgtacatcttgtttatacagtggttatattgataacaaatgtcatcaatataacattgaattaatctt
ccataattttatggggaaaaaaggagcagccttactgaagggcaaagttatacaacagctttacagaagc
tgcatgcgagtgagtgacacagaaaagagaccgcccgttttgaaaactcacagaactagcctcacgggtttg
cttgctttcatctgacacagaaaagagaccgcccgttttgaaaactcacagaactagcctcacgggtttg
tgagtgccattgagcgctggctgccaagaacgggtgtttaactcgagaaatcattgaacaagtttagaaa
ataaagatgcttatgacaatttcaaacttgaaaggtctccaaagaaggactgagatattggtgagaggag
taaagagaatcctggtgcattttatctcatgcttccttctgttcgaagatcattttgaggtttataaaag
gtgggggtgatccaaaaatctccaggctgagagtcctggctgaggctgtgaactgggctgcagagaaagg
gccacgcctccctcctctgctcgcattactcagcagccttttctgcatgtggctggctgcagacaatcta
aacccttccgctgtgcgtcccccccttatactgttctgccaaaaggaaggcagagaggaaatcagctacg
gggc (SEQ ID NO: 7).

In an embodiment, an *SCN1A*-specific enhancer sequence comprises a nucleotide sequence which contains one or more regions of 50-500 bp or longer, 50-250 bp or longer, 100-200 bp or longer, or 100 bp or longer, having at least 70% or greater, at least 75% or greater, at least 80% or greater, at least 85% or greater, at least 90% or greater, or at least 95% or greater sequence identity to the following mouse polynucleotide (DNA) sequence (E4), also called "S5E4," located on chromosome 2, at start/stop positions 66387764/66388024, shown in FIG. 1A-1, or a human ortholog thereof.

tctgacagagcaagtcttgacctgcttaacattatgttatgctagtcatttttaaagttagtctttat
cccatagaaggtcagttttttacattattatataatcttttgacagaataacaaataacattctgaat
gtctcatttctaaatacaaaacatcttagtataaaattatgcattgttttaagtcttggaagtaggtc

cacatgtagaaaacaaagtacgtatgataaaaaatatcaaaattgtatattcag (SEQ ID NO: 8).

In an embodiment, an *SCN1A*-specific enhancer sequence comprises a nucleotide sequence which contains one or more regions of 50-500 bp or longer, 50-250 bp or longer, 100-200 bp or longer, or 100 bp or longer, having at least 70% or greater, at least 75% or greater, at least 80% or greater, at least 85% or greater, at least 90% or greater, or at least 95% or greater sequence identity to the following mouse polynucleotide (DNA) sequence (E5), also called "S5E5," located on chromosome 2, at start/stop positions 66392447/66393109, shown in FIG. 1A-1, or a human ortholog thereof.

aatgttttgatatttaggagaaaattgcaaaacaaaatgatgacagtgtttgaaagtgtttgatcagtg
ccaagcatcactttatgtacttggaacatgacttgaggccttaagctgtgatttgcaaatgtagatt
ggaatcaagatctttatagatgaggaagcaaaaatcagaagacaaaataacattatcaacttgatctca
tgtgcagccagggctgaactgcaaatgctgatttgccccagctctgggctcctcaaactcgttccttgga
tcttattagttggaactttatctctgctcgtggcaggggtgcctgggaccatgtttataaatatctgctg
aatgaagaataagtgagtcaatcgaaccagaactcactttggttagttaatttcattcgtggatattat
ggagagcagaagaagaattccagagacacgatttggtcaaaactctctaagaaaatgatgacactata
tattgatgaaaatgaatgttcttgttcttgcctttatgtattttcttgtccccccactccccatctgct
agggctctcattacagcatagttcttgaatatcccagggttgacctgaagttacaatatattcttgattta
gatggcagacattgggaatatgttgcactcttaaaatttaata (SEQ ID NO: 9).

In an embodiment, an *SCN1A*-specific enhancer sequence comprises a nucleotide sequence which contains one or more regions of 50-500 bp or longer, 50-250 bp or longer, 100-200 bp or longer, or 100 bp or longer, having at least 70% or greater, at least 75% or greater, at least 80% or greater, at least 85% or greater, at least 90% or greater, or at least 95% or greater sequence identity to the following mouse polynucleotide (DNA) sequence (E6), also called "S5E6," located on chromosome 2, at start/stop positions 66401767/66402372, shown in FIG. 1A-1, or a human ortholog thereof.

ttgtcactttgttactctacagtgttgccctggagttcgatacttcattactctatagtggggtgaagaa
gttcacacttcttatttttcatttccctccctcaatgatttcttcagagctagctcttaccagctagaaat
tcttcaaacgacactcgtgccttcccttcacacaggttgaaactatttgtctctaattgccctaaagtactg
gtgttcaatcttccaggcacttccaatgatctgaaatctgacctgcttaggtcagctggctctgagatt
atgggtattctagtcctcaaaccaacctgttggtcgttggtttgtaccaaacacactgacttacatag
ctcaaaataccactggccttttaaaatggcatatcacattccaggggaggatcaaaactgctggctgg
tgatatttgtcaagtctctcaaagttgcactttccaggattttcaattcactgaattcttagacagaca

tgtttatgtgaaagaattctttatatatTTTTTctcctcttttgagtgggcaaagtaaaatcttgacctc
 tgggttccttattttatttgactctctgtagtattttaaatcttaaaattttcct (SEQ ID NO:
 10).

In an embodiment, an *SCN1A*-specific enhancer sequence comprises a
 5 nucleotide sequence which contains one or more regions of 50-500 bp or longer, 50-
 250 bp or longer, 100-200 bp or longer, or 100 bp or longer, having at least 70% or
 greater, at least 75% or greater, at least 80% or greater, at least 85% or greater, at least
 90% or greater, or at least 95% or greater sequence identity to the following mouse
 polynucleotide (DNA) sequence (E7), also called "S5E7," located on chromosome 2,
 10 at start/stop positions 66407834/66410263, shown in FIG. 1A-1, or a human ortholog
 thereof.

gatactgtataattaattaggcctccaatcatgccttcccagcctccacggatggagaaaccctctccg
 ccatgccttaaagaggaattgctgtaataaatgagtcctctgatagcaaatttctcagcaagggggaat
 cgcgtaaatggagacatagattgacagcaaagtccaatgtgtatttttaccagaacgaactctccgg
 15 ttcaagcctttgaaagagacatttgaaaacaaaaacaaacaatgtaatggagcgaggaaaaaagccac
 agaagtgagtggcagggagtttaaaagagcagatgccactgccaggtctatgggacataaccagccact
 tgtgctgggtcttggcagtttataatgctacctcatcttctccgcgaaattgttttcccgtaaatctct
 gtggccatccattcctgtctacacattatgttcctaaaaatagacaccatctaaaaatcacttcaaggag
 ctttgtggaggaaggcctaaattgcaacactcctccagcgaagatagatgcagtggttgatggcattac
 20 cagtcggtagccaggaaggggagtttgtgaggagtttttccaccacagttaatctgtttctggaaggaa
 agggagtgctcagacttcccagggaggcacaacgtgtgtggaagctctcatttgcacccccggcctg
 tcaggtattgcagcaaaagggagaggtgagctaccctggctctccttgggcaggagggacagaatcagg
 aagcatcaacctcagcatggaattttcctattcctggttggcatcctcctcttgggatgatttacagcg
 cgggttggagaaacacgctctgccactccactagcgcaccagatagacagtgacagcctgcagatccat
 25 acccgaggagaagccacatttctacgtgtgatagcaacagcgtttggcaatttgcgactttgctactg
 cagcttagaaaaatatttagtcacatgcacatctgaacagaaaagacaccagccttgactcagtcatttc
 cgtcagacacacgaaagaaaaagcgtctctgctcacaagcttatttggactgctttgttgaaaggaggg
 gcggcagacactttgtagatgtggcaagagggcctttatatccagacctcaaacaggtaggagagaagga
 agccaggagaggttaaggaagggcggtggaagcctcacagccacctcgaagaaaaacagtttttttggc
 30 ctgttcagaaagcaagaggttccacagtggttttgtgtcaatggagcacatctgcagtatcatttgccgt
 tgggtgacctctgtctaattaaaagtaagtcagtccttcccacccggcattgtctgaaacccgggactct
 ttatcacttttgctaaagtccatttgcaagtgtagttaaggaagagtcaggggggaaacagcatctgtcc
 cttctggtcctggggaggaggcactcctttccaagagtcaagcctctgccaaagaagctgcctccct
 gcaatgctaggatccaggagcagccccgctgccttcttgcctcctctgtgaggtctaatttttgcacat
 35 tctttaggagcgatatgacctctattcacagccatcgaatccagttccaaagcaccaatgacagagggg
 gcttcaagacaagaccttgcctaggaggtgagggcaagcaaggcaagagctggcccgatgccaggtt
 attttaggccaagaatctcatccttctatcaaaatgctgaactgcaaaacgaacctgatttcagttca
 tgggaaggttgagaggaggaggagggggaggaggaggaggaagagaggaggaggaggaggaggaggag
 gagggaggaggaggaggaggaggaggttggccgaattcacatgcaaaaatagacttcttctgttctgc
 40 cccaactcttatttccgtgggctcttctcccccaaggatttaccaggtaagaattcaccaccaaagaaga
 tcacaatgagataatcagatggcttacctgataaaaaggaaaattatccatctgcagtgaggagcaaca
 tctccccacgacgagtcgccaccttccggttgcaacgattcagattccttcttgcaaaagggtgaccaagt
 gcttcacaagggtgcagcctcataggggcagaacacacgtacacaaacacacgcacacacacacacac
 atgcaccagagacctctgcagtatcctctcggttcatcctcgccctcactctatgggtaccttaatacaaa
 45 tcagcaaatagcttgttttaaaaaaaaaaagaaagaaaaaaaagcggagacagcacctaacttacagtg
 ccatctagtggctacatcgtaaataggttctcacagcctggatttctgtgttctttctcaaccgcttcc
 ttctgggttccttttt (SEQ ID NO: 11).

In an embodiment, an *SCN1A*-specific enhancer sequence comprises a nucleotide sequence which contains one or more regions of 50-500 bp or longer, 50-250 bp or longer, 100-200 bp or longer, or 100 bp or longer, having at least 70% or greater, at least 75% or greater, at least 80% or greater, at least 85% or greater, at least 90% or greater, or at least 95% or greater sequence identity to the following mouse polynucleotide (DNA) sequence (E8), also called “S5E8,” located on chromosome 2, at start/stop positions 66439814/66441457, shown in FIG. 1A-1, or a human ortholog thereof.

attgatctccaacttttttaaatccctctgtcatttttaaatgaggtgcactcggttgtgtcatcatctcg
 10 gtttttaattgtgtgaaaatttccctgccaatctcacaccgctgcggcaacctcaccttgctacttgccct
 gcaagttctgcagtggtgcccgttctgagtatgcccgttttaactagttcttgagcaggacacaaagcgag
 atagtctgatagaaccagttctcctctggttttaacctttactcttagatgagttaagggtcacatcaaa
 ccagggctcagcccgcagatctcctaagcacagcccctcctgacccaatgcagtttaaccaacctcat
 15 tcagcgttagtatcaaatgacactggagctgctgcagtatgcatcccagagactaagtaggcaggattta
 ttatcagcagaagtcccctaactaccagggttattcaagctccgttcttgctcaciaaacaggcgccggga
 agacacagtgagcagactcagagctcattttacaagacaagcgaattctcagtttagagacaagggcagc
 gcggcagcgaactgcagtaaatcttttcacgctcacagcaacatctaacaatgctctcctgcaacgcct
 cagatcaaacgaatcctacttggttttaacatcaaatcaacaccataaaaaaggcttcattagcaaagt
 20 tcaatttaggatgtttttaatcgtgtcttaattctagaaccagtgcgagactttccatgcttattcaag
 catgctgacagaattggaacctcttagaattgcctacctgcacctatcagcctggctgacaggagcccg
 ccaaaggattaaaaaaacaaaacccaaaacataaaatcatgcaaaaaaatatttacccccgaagat
 gtatgtagttaaagctcagcttccctgcagcctcgatagcccctgaagtgttaatctgaagaaacagttc
 catgagtttccacaggccggtagtgagtctcctacacttgacctagacagacttacataatgaagcatc
 25 agtgctggggagcttgacgatgtcatcaccagcaagagtaagaagtattggcagcagcaagcaggcgg
 gcaggctgagatcttgcatggaaatcatgaaccaggctcttgcttttctgtttttgaaacgttttggaagg
 agagttatgaatagcccagaaataggtctcattttgtgggtaggaagaatgaccagaagcatgaaagct
 aaatctcctggcaagtgcaggggacctctcttgagtggtgcagtaaacccgaggggacgacttctcctg
 ctgtcaactcctgaaccatcacatctggagtgaaggaagggctggtgaagccttgtaataaatgcaaa
 30 ggatgctgctgagagctttggtctgcctttaactcattgtggtgagtagaggggatgtggcagtatgca
 atgagagttgggtgtgtaggttgctttgcagagtaataaccccccccccccccaatctgtgaagtgtca
 atacttttagacacattttaataaacaagatgatagtaaaattactcttctccatcaaattgagactgtg
 ctgggttaaactgttttaaatgcattttaactcctgatgttcatccaagtaataagag (SEQ ID
 NO: 12).

In an embodiment, an *SCN1A*-specific enhancer sequence comprises a nucleotide sequence which contains one or more regions of 50-500 bp or longer, 50-250 bp or longer, 100-200 bp or longer, or 100 bp or longer, having at least 70% or greater, at least 75% or greater, at least 80% or greater, at least 85% or greater, at least 90% or greater, or at least 95% or greater sequence identity to the following mouse polynucleotide (DNA) sequence (E9), also called “S5E9,” located on chromosome 2, at start/stop positions 66441748/66442268, shown in FIG. 1A-1, or a human ortholog thereof.

atctcaagtgtatgtaacatgagctacagtctttaaaccctacaaacagtacatccagtctcctaccatg
 attctgagtgatgatgatttcatatgagcacaagatgacatcatactatttagttatatgtaaaatcatg

gtcttacatggggttggtggacaaaaccatctagttttggagggtgacagaaatagagaggacgccatgcac
 tacttaaaaaataatcgcagccttcttttcttagctagggagtttgctgctatgagccacattaagacca
 ggggtgaggagatgagacgatacaggggcatgaaagaacacggtgatctacttttctcctgttaattaacg
 agtaaggaaatagacattaaaagaagttaaattgtgtctgagccaacgtaggtgaggtttcccccaaatt
 5 cacttggtagttttgctactgcagtatagtaaatacttgttttcatttgttttttttttttttgtttttt
 ttgtttttttgtttttttgtctttttgttttttttttt (SEQ ID NO: 13).

In an embodiment, an *SCN1A*-specific enhancer sequence comprises a nucleotide sequence which contains one or more regions of 50-500 bp or longer, 50-250 bp or longer, 100-200 bp or longer, or 100 bp or longer, having at least 70% or greater, at least 75% or greater, at least 80% or greater, at least 85% or greater, at least 90% or greater, or at least 95% or greater sequence identity to the following mouse polynucleotide (DNA) sequence (E10), also called "S5E10," located on chromosome 2, at start/stop positions 66450594/66451140, shown in FIG. 1A-1, or a human ortholog thereof.

15 tattgcaaaaggaaggaatgagacagtttatgcagagctaagggtttgtgcgttattatgattaatcac
 aaggacagctgccaaagcttccatcatgacaatattctctgggagaattcatcaggttctactgtctatt
 aatttctgttgatgtatcttatctggcatcttcaatgacagaggacacttgtagttttttttttttaag
 tgaagggttaaaagacaaagttcattaaagaaatgatttatatatgacatttaagaactagcaatgtcat
 tgcttcaagaaaattatgagaatttagtcttgtaggagtttacaccatgtccttgaagtgtctaatta
 20 tgtgacttgatagttttacttagtacatatcgattaggctgtatctattatttatcaagaaattatgga
 aggaggcaatgtggcataggcatacacattctgattttaaaataatcctgcttttaccattaactcctt
 ctcagataattctgaatacatatcttgctctatgaatctgtgtaatcatggaaaaagaaaaaatc
 (SEQ ID NO: 14).

In an aspect, the human sequences (human ortholog sequences) for the ten above-described murine enhancer sequences were determined based on alignment of the mouse sequence to the human genomic sequence of *SCN1A*, including 100 kb both upstream and downstream. Accordingly, human ortholog sequences that are highly conserved between mouse and human sequences were identified.

In an embodiment, an *SCN1A*-specific enhancer sequence comprises a nucleotide sequence which contains one or more regions of 50-500 bp or longer, 50-250 bp or longer, 100-200 bp or longer, or 100 bp or longer, having at least 70% or greater, at least 75% or greater, at least 80% or greater, at least 85% or greater, at least 90% or greater, or at least 95% or greater sequence identity to the following human polynucleotide (DNA) sequence, called E1 or S5E1 herein, located in the human

genome sequence at human_hg38 start 165953030/human_hg38 stop 165954796

(FIGS. 1A-2 and 1A-3):

5 tctaattggacatacagtaacccttcataaatatatttgcgtgaacgagtgattcagtgaaacaaatgaataga
 gaagaccaacatccgaaaagttatattttatatttcaagcctcatgtctttaactgttttatatcagccttt
 ctttaagttgaccgtcattaatatttgcgtgaatgaatgagtcagtgataaacagagaagaccatcacctt
 aaaataacgacccctccacttttaagtcttacgtctttaatgggtttcatataatctttctgcgtcttt
 tttactgtccagtggtgggagctgacactagtttgccttaagtccttaaaaaatcgaccccgaggcgag
 tgtcataggttaacccaagctttcctagtaaacatgatacaaaaagtaaacacacacacagcatggggac
 10 cagcaattcagaaacaccgagcgggcggtgcccagacctgggcttccccagcaggggcccgcggagac
 cgccgtgagcagaggctgcaggccccaccccgcaaccgcagcagccggggcaccgcaggggaaacagcgg
 cctagcgaagccaccgagctccctccgcgcccccgggccaaaaggccgcaaaggaaactccgcccggcc
 gcccgctcacccgctcacccgctcacccgctcaccttcaattcctgcgagtcctatggctgccccgaggc
 cgggcccgcggggctctggggattgtctcgccgcagcctaaaggaaagcgcagaaattcagctcccttagc
 ctccccgagcgtctagcgcggggggcccgagcgggagggggcggggtcgcgccgcgattggctgtcgga
 15 gggagaggcgggctgtgtggcggggatcgtgtgtgaatggagcagggggcggggggacccggagggtga
 gggctgcgagggggcgccgggaggggtccgggctgggaaaaggccctccgcccgcgagagtgacagctgaaa
 aggaggtcacactgggaaacggctgtctgaggacagtggtggcgggccgaggaaatggaattcagga
 ataaaggaaacggagtgatgaagaagggaagtctgtttcctgtcactggttgtaaaggaaagacaccatt
 ttctgcacgtttgtctggaggcggattcccgagtgccggtctcagcaaggctctgcccgcgcccggaaa
 20 aagcgggtcaactttcacgtgggcaagttgttttacggccacaagggtggcgagaaaaaaaatcacac
 gttcttaacagaaatacgggtgcgcttgggcccgtctttgcaggcgttgctgcaatctttgttagaatgt
 gtgttcaattagcccttttttaccagccccgataataagaggggacaaataaattaaacttccagaaaat
 tagtgtcttgttttcaatgataactactgattttaaactgagaataaaaatgaatcccaatgcaattttt
 atgtttgcaccccataggaactcaatcagtcacacatagattttcttaagtcaggaaattaaatgga
 25 aatataatagactaagattttctatttctgttaataataatatttaaaatagtgataaggtctgagat
 ttaagtgatctttgcagaatctttcacgtggattccaaattttgatcctagtgtaattatcttacttt
 agttgacatgatacgtagttgccttttccagattttaaggttttctaaggagtttataaacattgacttt
 tcccccatgccaataggttatgtaaggacagtctt (SEQ ID NO: 15).

In an embodiment, an *SCN1A*-specific enhancer sequence comprises a
 30 nucleotide sequence which contains one or more regions of 50-500 bp or longer, 50-
 250 bp or longer, 100-200 bp or longer, or 100 bp or longer, having at least 70% or
 greater, at least 75% or greater, at least 80% or greater, at least 85% or greater, at least
 90% or greater, or at least 95% or greater sequence identity to the following human
 polynucleotide (DNA) sequence, called E2 or S5E2 herein, located in the human
 35 genome sequence at human_hg38 start 166084035/human_hg38 stop 166084884
 (FIGS. 1A-2 and 1A-3):

agtggtgggcctcccagggtgttttagctagcaatgagagaggcactgcctatatccaagttgtatatgg
 cagggttttgcacaaagtggtgattactcgaagagaaaagcctaattggccagtcatttcatcttccccctttct
 cgatgttcatcttttctctcccagctctcctttattctcaattttcttttttttttttttttttttgcag
 40 cctccatctcacttccgttgctgtcctctccccaccccttcccactctggactgtgcctctcctttgta
 gacacttcaagtcattctatttcatcaaaaaccatgggtctagaagtaacttaattgtaaacccacaaa
 gatggagacagaatgaatgccattcttcttgcgtctctcagacaaatgcaggtcatttttgcctatgg
 tgctggtaaaagccaggagttatgtagctataagtagcagccagaggaaatagtgctgagtcagcaatt
 gtcttttttattgctgtggggcaataatgggagaaaaaatcaggcttgggtacaattccctttgaaggaaa
 45 aagatgccaacactagcatttttaacacaaaatgctggttgggggttgggaggaaggatgcttacattcc

ttcttttgaaatatctacttttgataaccattttggtaaaataatgcagtgttttcagtgtgcaaaccct
 ttcaggactcatgggtgtatggcagacgcacctgacagcaataatttaagggtagacctgagaatgactc
 tgtggtctaaaaagaatgtgtgtttggaagtctgaggtaagaaatctggctggaagtggccaacctgga
 aatttgctccttattattaagg (SEQ ID NO: 16).

- 5 In an embodiment, an *SCN1A*-specific enhancer sequence comprises a nucleotide sequence which contains one or more regions of 50-500 bp or longer, 50-250 bp or longer, 100-200 bp or longer, or 100 bp or longer, having at least 70% or greater, at least 75% or greater, at least 80% or greater, at least 85% or greater, at least 90% or greater, or at least 95% or greater sequence identity to the following human
- 10 polynucleotide (DNA) sequence, called E3 or S5E3 herein, located in the human genome sequence at human_hg38 start 166090876/human_hg38 stop 166091720 (FIGS. 1A-2 and 1A-3):

atagtgc aaagt ttaaat ttcatt tttccta agattt gttttaaa ataacac gatttacc caagt gattt
 caaaccaca attacatt ctgtttaa attacta atatttt attgcat cacaat ctgcat gaaacag atgt
 15 caggatata atgaactaac ctgcattgt atttttt atttttt gtctcct gtggcata acgattt catagga
 aagagaact acacagct gactgact gatggggaa agttacaca atggatag ctttgcag caacatacta
 atgcggtag ggagatgt ctgcagag aggctagaa ataaaaa tcat tttctt ccggagc agcactg cttgct
 gtccggtg agacaaaaa agagattt cctttttt cctttt cttttttt tga aaactc acataac attaat
 tctgtta agcactgg atacacg gaagggt gtttacct tagaaaaa tcat ttagca attttt tagaaact ag
 20 acatatag caatttttaaat cttttta actatcta atgaccaa agcagagg gtcctcaca agaggg attt
 agatgct actgaatt gaataaa agaaaat atggata catttatt gtatgc cttattc agtttgagg ttca
 ttttgag ttttagaa atagggat ataaaaa catcagg ggttaaat agcatgg gtaaagg acatgaacaa
 gctgcag agaagagg ctgactgc ctgctata tttgcagg cattactc agcactt tttcttaa accgatac
 atcttgct ggctgc ataagca agacaa gaccctt tttccct atggctc aggaaggc agagaagt caactt
 25 cagccttg aaaaaagg ca (SEQ ID NO: 17).

- In an embodiment, an *SCN1A*-specific enhancer sequence comprises a nucleotide sequence which contains one or more regions of 50-500 bp or longer, 50-250 bp or longer, 100-200 bp or longer, or 100 bp or longer, having at least 70% or greater, at least 75% or greater, at least 80% or greater, at least 85% or greater, at least
- 30 90% or greater, or at least 95% or greater sequence identity to the following human polynucleotide (DNA) sequence, called E4 or S5E4 herein, located in the human genome sequence at human_hg38 start 166094366/human_hg38 stop 166094633 (FIGS. 1A-2 and 1A-3):

tgccagacagaacaagtttttagtgtagttgatagtaagttgtgcccagaatattaaattgagtcaaatt
 35 tat tttccacataa agtcacag ttttatatgt cattatata atctcttggcagaaataaggaataacat

(SEQ ID NO: 18).

catgtaaaaattaatatgatcttttagtcacttagaaaaaataaccataaagaacactaatagtgttttaa
agcatctacccagtgccaagaactgcattatgtattggtgaacataacttttagactttaccatacaacg
tgaaaatatatattattatcactatttttacagatgaagcaataaaaagtcagaaaaaatgtagctaatta
aagtgatactgtgtatagctagagcagtgatagctagagctgatttgtctgactctagccctagtttc
ttccattatatcaatttctctggaatgtatctctgttcatggcatagtgccctgacactatgcttatta
atatcttttgaataaaaagaaccactgagtgatttgaataaaaactaaatttagttagttaattttattg
gtggtatatagagatagtaggaaaaataattgaaaagagacataaacagatttgccaatactttctaag
aaaaattatggaactagagtttagtcaaaatgaatgctttcattgttagaattcaactttaatctttgc
agaatacaaaacaaagacccattttctagaagaagtaacaggggaagagagagtaagaaaagagataatgat
gaacattgtctaattgttacagcataatctagtaaggtaagaacagagagagttcattgacttaccaac
atagttgtccctaatacacctctgtgaacctagagtgctacgatataataatgattgtggtggtttaaaa
agtaaatgggctgggcatggtggctcacatctgtaatcccacactttggaaggctgaggcaggtgta
ttgcttgagctcacaaagctcgacaccagcctgggcaacatggcaaaaccccgtctctacaaaaaata

In an embodiment, an *SCN1A*-specific enhancer sequence comprises a nucleotide sequence which contains one or more regions of 50-500 bp or longer, 50-250 bp or longer, 100-200 bp or longer, or 100 bp or longer, having at least 70% or greater, at least 75% or greater, at least 80% or greater, at least 85% or greater, at least 90% or greater, or at least 95% or greater sequence identity to the following human polynucleotide (DNA) sequence, called E6 or S5E6 herein, located in the human genome sequence at human_hg38 start 166118214/human_hg38 stop 166118879 (FIGS. 1A-2 and 1A-3):

62

gcgcaggctggagtgacgtggctagattcttcttgagtatgctcaaacttcctttttggaatgtcttcc
 aaaggcactcttgcccttcatttgtacaagttgattgaccctttaaaaggccttaaatattattgtgacac
 ctcacagactcctcaaatacctgaaacctgaaatgctgaggcccagggtggcactgaaatgatgggtatt
 ctagacctgacaccggactgttttctccttggtttgtcccaacacactgacatacatagcccaaaata
 5 ctactggcctttttaagtggcatatcacattccagggttaatatcaaaactgctgcttggttagcatttgt
 gaagtctcaaagtaactctttccaggattttcaaataccactgaatttcttagattgaaatatgtatgtg
 acagaattctcttagcttttcttctctatgaatatgtaattggaaaactctgagatccggtttctcatc
 ttatttggatttttctttaatcttaaaattatgaatatgttgcct

(SEQ ID NO: 20).

- 10 In an embodiment, an *SCN1A*-specific enhancer sequence comprises a
 nucleotide sequence which contains one or more regions of 50-500 bp or longer, 50-
 250 bp or longer, 100-200 bp or longer, or 100 bp or longer, having at least 70% or
 greater, at least 75% or greater, at least 80% or greater, at least 85% or greater, at least
 90% or greater, or at least 95% or greater sequence identity to the following human
 15 polynucleotide (DNA) sequence, called E7 or S5E7 herein, located in the human
 genome sequence at human_hg38 start 165892760/human_hg38 stop 165897884
 (FIGS. 1A-2 and 1A-3):

tggaacaaacgcacaaacggttgatggataacgggtgatgacttacacaacaatgcgaatgcatttaatgcc
 actgaacagtagacacttaaaatggttaagatgatgaattttgtgttatatatgtttcaccacaatacaa
 20 aatattcttttaaaaagacttttggaataactgtatctacttaattacaggatgtcaaaactaatacagg
 ctgatagtagtatttctgtcccttgacacacaatcttgggtccagagattttgttcaccacaccttttag
 catcactaaaaagggcacaataagaatatggtttcagaaaaagacaattcaaatatttggtcttgcctt
 tagctatgtgaattcaatcaaattactcaaatcttcttgagtccaatatacttattttcttaaaatagga
 ttataatattgactgtaggagtgctacagaaaataaaggcatgaaaaatatttataaaattacaaatgtta
 25 ttaataatattttatacttccaaaaatgttgacaagaaatagagtaactaccccataataaagccacagc
 atctggaagctatattggattaagcaagaactaaagggttaaaatctcgattaaatttttttgcatga
 tactgctagtagtattatcaacattgggaaggcaatttcttgaatatcttatactattgaaatgtatt
 cattattagttcaagttataattaccagtgacagattaaattacattcacttgcctttggttaacctg
 acatttgacagaaggcaatttctgcacttaagaaatgtattaaaaactaaaatgtatattaccttcta
 30 aaaaacttagctgggtccatctttattgatgaatagtaggaagatatcaaaatagttatagggtgatgag
 atgtggcaagcatgcagtgctatggtatttacaagcacaggattcttaactttgcctggaggag
 ttgggaattttcacataggagttgacctttgagcagcctcaaggataggaggaagatcctactagacgg
 acaaaggcattccaagtagcagaaggcatgcgccaaggaggaagcagagaacagtggtggggagtgttgg
 taactttgatattattataaagcggaggaagaaggataaagaaatataaatggccaataattttgcggccat
 35 attattattaaaaataatgctatgatttttagactttatcctgaagcactaacttaatttttaagcaaagg
 gtaggtttttgatttttagaactgatatgctagtcctatgatgacctggagcagccagaacctagaagct
 ggaagatgagttgggaatctgcactagttcagatgagaggtgataagggctttcattagagcagtggtg
 taggatacagacagactggatgtgttagctagctatcaagcaaacagagctgaggagacatgttaaccaa
 ttagtatgaaggaaggggaaagctcaaggcgatctggagattctgagagagaaaaggggcaatctgtc
 40 tgagagcagtaattagatctagaagaggaatttttcaactacttaaatagggtcaaatttgtatggtac
 atttctgaaataagctaaaaatagagccttaattctaaagtacaagatgagttactgaggataaccaataa
 tgtacacataaaatgaacggagatgcatgttttagagtaattccaacaaaatagatctgtggataagta
 tgtaaggtagtagtaagaataaagcatacaacacaagattaaaaactcttaagattaaaaatcacat
 agacaataaaaaatttacttaaaattttgggtgttttttgagaccaagctctcactctgtcaccgaggct
 45 ggagtgacgtggtgtgatcttggctcacggcaacctccacctcccagggttcaagcgattctcctgctgt
 gtttttaattgacttgggtgttttacagtcattcactgatccattcaaccaataaacatctatgttgcca
 catccatgtgtgtggtgcatcttctgatattgggaatatggtgtgatccctgaactcgaggagctaca

gtgtaataaagaacacaaatatgcacataaatatTTTTtagtaaaatactataagtaataaagacgtatg
 gacaaagtaaccgaggattagggttaccaactcatcccagtttgctgtgactttctgttagcatggga
 agtccctgcatccaggaaaaccctttgctccgaggcaaatctaggatgggttggtcatgctaccagcacc
 acataagacattttttaatgtaggtggtagtgaggatgcagatttactttttattttcccccaaga
 5 gggatataatttttagattatgtgtttggaaagagaaaaggataaggaagctaaagttcatttttaggcaa
 aaggaaaaaacccaagcaaaagacttggaagcatgggtgtatgtcatttggtgttgctggagtacaagat
 ccttatattgcatttataaaaaattgctttttatatttggtttacaaacagtcctaaagcagccagtctact
 aagccaatttttttgggaaaaaggctgctgccaagcaacagaaacttacgtgaaacaaaaccacaaga
 10 cacatgaagacttcttcaaactcttagaaaactataatgtgtgagattcttcaaactcttagaaaactata
 atacttttaatgactttaaataattcacagtggaagaagtctgttttttaaagaaataaagttagatcat
 tgtctcaaagggaaagactgtgaaatgggaacagcttgagatagaatgaatatattatgtatattactt
 ttaaattggtagtttagagaagaggaataagagaaaaacagtggtggaacacaaggtagaaatggcagggaa
 aaaaacgatacaggcctgaagaaaataaaagtaggtttgggcaatgtgggtggcaagatgagcccatat
 tttggaccagagtgagtggaaagaggtgagatgggtcaagtagttcacagcattcttttaataaacatct
 15 gagtatactctggaatagacagggcaaaaaacaaatgaaattgcctgtggttagtccccatattaaataa
 atttaattttatttaattaaaaaactcaaagagtaaaaaataaaagagaaaagtgttgatattactgtaa
 aaaacagaccatattttctccttttcaatttttgctgctcgctgggataattttatttttaatacaaaact
 gatgttctctaaattcaaacatctttttattaaagcggttactagaggtagcctgcagagcatatctag
 ttctttgagttgcccctgcttgaaggattgccacctctgtgtctctgaggctctgagcacagatgcctag
 20 ggcacagcctgagatgaagggtgggtgggtttagaagaactgaaacacagctctaggacttctctgcc
 atttcaaactcttttttagtaacacaggttagtaacatacagtcattgtatttagtcaaacagttcccacctc
 tttctttttctctttctctcctccctccatccctgcttccctttcttctctttctcctcactcttct
 ttccatctagtattttgttgagtactaaactaggccagatattcttctaaattctggaacacagcagtga
 aggacaacgtttctgggtcacttgatgcttccattctatgattttgtagttttttgctttttgtttcacaag
 25 ggctacacaaaaccccaaaaatcctaaagccaaacccccaaaactcaactgaacagaaaaacaggataaag
 ggacagagagcaagaagtgggtgctacttgatgtaggccagtgaggaggggcatttctaaggaagaagca
 gcaccagagcaaaaggcacaagaaagaaatgaggcagcagccagatagacctgtgggagcagcatattccg
 ggaagtagaaacagcaaaacaaaaggctcaaggctgtaattggcttgggcttttgcttacctagtttct
 gttgtccctttcttcttttactatcagaattctgattttattctacagggatttagattcagctaaaaga
 30 taacattttcccaccttctttttgcagccacggaggtgatactactaaagggttttttgtttgtttgtttg
 ttttgtttttaataaaacgatgattttgctgggtgggaagcccttttgccagctctgcaccacaccttc
 ccacctgggcccctggaacatcaatgccagcactgcgggtggtcatcttgaaccatgaggtgatgtcag
 gcaagtccagggtaacaacgtagaacaagagagaaggctctggtttcctgatgatactgtggaaccg
 ccacaccagtcctcacagtgtagttttcatttcttttacagagagaaaaactaaaaaccgtagtttt
 35 agcctagaattttcaggggtatctctgccattttcaatgaaagtattttctaaattcttcataggctagg
 gatggaacacagatgagttatgacgacgctgcaataatctatgtggaagatggcaatgccttagacca
 ggggtggcactaacagaggtggtaaaaagtgtggtcattctagggtatactttgaatgtagcactaacaag
 gatttgctgatagactggaggtgatatatgagagaaagatgagacaaagggttaactgtgaggtctgggc
 cggcacaacagtgagcagtgatgccagtcactgaggtgagaggtgggggtggagcaaacttagaggcgga
 40 gggaaagtccaggtgttctattttggacatgttaagcttgagttactcctagacatctgagtggaatg
 caaagaggcagaggtgtacatgagtaagggtgcagataaatgtttaggacacatctgcacatacatgg
 gatataaagccacgcacctggacaaagtcacctaggggtgaaatataaaaaagaaaggagttcagga
 actgaagctacgagtggttctaataatttaaagctcatcacagagaagagaactccaaaaaaggaaactcaa
 acagcaggcaatggggcaaaagaagactgagtgagtttaggatctcagagacaaatgaagaaagtctt
 45 tcattgtggaagggaataa (SEQ ID NO: 21).

In an embodiment, an *SCN1A*-specific enhancer sequence comprises a
 nucleotide sequence which contains one or more regions of 50-500 bp or longer, 50-
 250 bp or longer, 100-200 bp or longer, or 100 bp or longer, having at least 70% or
 greater, at least 75% or greater, at least 80% or greater, at least 85% or greater, at least
 50 90% or greater, or at least 95% or greater sequence identity to the following human
 polynucleotide (DNA) sequence, called E8 or S5E8 herein, located in the human

genome sequence at human_hg38 start 166148156/human_hg38 stop 166149792

(FIGS. 1A-2 and 1A-3):

tttgtcttcaactttttaaatatccatctatTTTTtagattagatgccatctgttgtattatcatatct
 ggttaaatcttattaaaattcctggccaatatatcactctgctgcggaatcttacctgctacctctc
 5 taggcttttctgcagcattcaataaagagcctgctatTTTtaactaattgctgaggcaggacacagtgtg
 agagtctaaaacagaatcagcgctatatTTTtctactTTTaccatagagtgaacaactgttcatatc
 aaacctggaatcatcccttcaagtctcctaacaacagcacagtttgagccaatgcagttaatccaccct
 ccttcagtgtctagtgtcgaatggcgctTTTgctgcagttacaccttgaaactaagtaggcaggattca
 tttttgttgaagtcacctaactgccagtttattcttatacgatcacaacaatcacaacagaagacaa
 10 atacaggcatatatataactcacttacaagggaagcaaatTTgcagccagagacaagggcaacgtaaca
 gccagaactgcagtaaatctctTTgagctaatagccacctctaataatgctctcctacaacacctcca
 atcaaataaattgtattcagagtttaaatatcaaatcaacattcattccttgatggttacagatgatgt
 ccgataagcaaatttgaatttatgatttatttactccttaagtgtctcaaagccaggattactggaaga
 ctactatgcttattcaagcatgctgcagaaactgtaatctcagtaatttctccctgcacctctcagca
 15 tgactgcagggcatctgccaaactgtagtacataaaactgctgaaacatgcaaaaatatttaccctcaa
 aagatgtagtaaaagctcagcttctccagcttccataaccctgaagtgttaatctggaggaacagtt
 ccatgagtttccacaggccagcagtggtgtctcctacacttgacctagacagccttacataacgaagcac
 cagtgtctggggagctctctgaatgtcatcaccagcaagagcaagaagtattggcagcagcaggcagcca
 ggcaggctgggagtttgcatggaaatcatgaagtctttcttgtcttctcctcttttgaaatatTTTggaag
 20 gcgagttaagaatagctcagaaactgggtctcattctTTTTgtgggaagaatgaccagaagcataaaagc
 taagtcttccagcaagtgcagacacctctTTTgggtgtttgcagtaaacctaacaagaatgaattgcta
 tcagtaaaagtctgtaccatgacacctaaaaggaaggaaatggtaaaagcaaagtaataactcaaagaca
 gtcaccgagggctTTTgtccacctTTtaactcattgtgggtgagtagagggaatatgtatatatgtaatg
 agattattattgggggtgtgagttgtTTTgcaggatggtaactaaaggatttgtaaagagtgtttatgt
 25 tcttaaaagatcttgtttgtgagcaaggtaataggatatcaaattcagaacgtgggtgggttaaactgggt
 tgtatttaatgtatttcaacttctgaaaatcttatgcaactaataagaaa (SEQ ID NO: 22).

In an embodiment, an *SCN1A*-specific enhancer sequence comprises a
 nucleotide sequence which contains one or more regions of 50-500 bp or longer, 50-
 250 bp or longer, 100-200 bp or longer, or 100 bp or longer, having at least 70% or
 30 greater, at least 75% or greater, at least 80% or greater, at least 85% or greater, at least
 90% or greater, or at least 95% or greater sequence identity to the following human
 polynucleotide (DNA) sequence, called E9 or S5E9 herein, located in the human
 genome sequence at human_hg38 start 166150066/human_hg38 stop 166150702
 (FIGS. 1A-2 and 1A-3):

35 tcccaaatgtgtgcaatgcaagttatgctTTTTaaaagctagaaataataaaaccagtcttctatttctga
 ttttgagtatgggtgatatagttattttaataaaaagatgacattatattgtttaattatatataaaattg
 tggtttgatatgagttgttggttaatatgtataattcctgaggtaacagaaatagagaaggaaccacac

atcatttataaaataatcttaatgttctgctcttagctgggaaacctatctgctaataatgcatacactaag
tagagttagggaaataagagaatttagatctatgagggaaacacagtgatctaattccaacccattactta
actcataaggaaactgaggtagaaagaagtttagatgatatgcctgacatagaggaagaggtgagtga
aatggttttcctgacactaacttggtattttgtcagctatactgcaatgaataattgtcttttgatact
5 ggagtaaaggcttgatgtacagtgattttttatatcatatacaaatgacagaaaaaaaaaagtggagtagt
actaaatatctgcttttagcagtagtctgatttttgaaaaacaagttctgtactgattggaatgagaaa
ctttcttcagttattt (SEQ ID NO: 23).

In an embodiment, an *SCN1A*-specific enhancer sequence comprises a
nucleotide sequence which contains one or more regions of 50-500 bp or longer, 50-
10 250 bp or longer, 100-200 bp or longer, or 100 bp or longer, having at least 70% or
greater, at least 75% or greater, at least 80% or greater, at least 85% or greater, at least
90% or greater, or at least 95% or greater sequence identity to the following human
polynucleotide (DNA) sequence, called E10 or S5E10 herein, located in the human
genome sequence at human_hg38 start 166160023/human_hg38 stop 166160609
15 (FIGS. 1A-2 and 1A-3):

tatcccagaaagaaggaaatgggtcagtttatctggagttaagcatttgtgtattatcatgattaatcac
aggaacagtttgccaagctttcattataaaaaatattctccaggagaattcatcaagttccattgcctatt
aatctctgtccatgcattttatgttggcatcttcaatgacagaggacacttttaaaaaaagaaatgaag
acaaaagaaaaagttcattagagaaataatgtatgtgtgatatttaaaaaattaagccacatcatcatcc
20 taagaaaactacgagacttttagtttttagtataaaacttgcagtggtgttcttgagattttctaaatataagg
cttaacattttttccttaatacacatcgatttggcatctcatatgtattatttatcaggaaattataaa
acaaagtaaaatgatgttttactaaaatgcacatcatttttagatatgggatttttaaaacttgatttat
aatactacttttaccatgaaatactcttttgttgatgaccttgagtacatttcccatctgtgaatctg
tgtaatcatgtacaaaaataaatgagacaaaacct (SEQ ID NO: 24).

25 In an embodiment, an *SCN1A*-specific enhancer sequence comprises a
nucleotide sequence which contains one or more regions of about 100 bp or longer
having at least 75% or greater sequence identity to a human polynucleotide (DNA)
sequence of the above-described E1 (S5E1) to E10 (S5E10) enhancer element
sequences (e.g., SEQ ID NOs: 15-24). In another embodiment, an *SCN1A*-specific
30 enhancer sequence comprises a nucleotide sequence which contains one or more
regions of about 100 bp or longer having at least 75% or greater sequence identity to a
human polynucleotide (DNA) sequence of the above-described E2 (S5E2) enhancer
element sequence (e.g., SEQ ID NO: 16).

In another embodiment, an enhancer sequence as described herein comprises a
35 nucleotide sequence which contains one or more regions of 50-500 bp or longer, 50-

250 bp or longer, 100-200 bp or longer, or 100 bp or longer, having at least 70% or greater, at least 75% or greater, at least 80% or greater, at least 85% or greater, at least 90% or greater, or at least 95% or greater sequence identity to the following human polynucleotide (DNA) sequence, called E11 herein, located in the human genome
 5 sequence at human_hg38 start 36816984 /human_hg38 stop 36817612 (FIGS. 1A-2 and 1A-3):

tcagcaagtctgtcatcgacatcctgcaactgtttgagcgggcagagcaagtgcgaaaagattaaaaag
 tgctttttctcatcatttctgctcatatgaccagcgctgcagtgtctgcgcgcccggcgccagcccgccgg
 gcctggcatggcgccaggggcccggactctgagcgcagcgggagcggctcagtccagccgcgcgctga
 10 gcagcgcgcggccgcccggcaagaaggcgcgcggacctgctaccactcctgcaccgccaggccaggggtcc
 gcgggatcccaggggctgcggccagggcacgaggggaaggggccacctctgggatttagggggcactggc
 gtcaccagctgggtctggaaagtccacctgccgtcaaggacacgcaggaggtgcgcgctctcagatctg
 ggaaccttgccgatgtcctgccgcgtgggggaagatcctgaaccttcagcggccagcctgcacctcag
 gacctcctagggcctgctccctttctctctccactcctacctcagcctctgctctggtctgtcctggat
 15 gcaaatattatgctgcaaaatctgagcgcgtgaggtcctgaaacctgacccacccgacgcagggaggaggt
 ggcagggga (SEQ ID NO: 25)

In another embodiment, an enhancer sequence as described herein comprises a nucleotide sequence which contains one or more regions of 50-500 bp or longer, 50-250 bp or longer, 100-200 bp or longer, or 100 bp or longer, having at least 70% or greater, at least 75% or greater, at least 80% or greater, at least 85% or greater, at least 90% or greater, or at least 95% or greater sequence identity to the following human polynucleotide (DNA) sequence, called E12 herein, located in the human genome
 20 sequence at human_hg38 start 36817484 /human_hg38 stop 36817720 (FIGS. 1A-2 and 1A-3):

Tccctttctctctccactcctacctcagcctctgctctggtctgtcctggatgcaaatattatgctgcaa
 aatctgagcgcgtgaggtcctgaaacctgacccacccgacgcagggaggaggtggcagggacagggacag
 ggacagggcaggagctgctggggcccacttcgggtgccccatcccacatctggccagggatgcataattct
 25 aaaacctgatttgatgttttacttttattt (SEQ ID NO: 26)

In another embodiment, an enhancer sequence as described herein comprises a nucleotide sequence which contains one or more regions of 50-500 bp or longer, 50-250 bp or longer, 100-200 bp or longer, or 100 bp or longer, having at least 70% or greater, at least 75% or greater, at least 80% or greater, at least 85% or greater, at least 90% or greater, or at least 95% or greater sequence identity to the following human polynucleotide (DNA) sequence, called E13 herein, located in the human genome
 30

sequence at human_hg38 start 36818134 /human_hg38 stop 36818727 (FIGS. 1A-2 and 1A-3):

Cctgggctgcactaagtcccagtggtgaccttgggttggtgaccttctctgggcttctgtctccttctgat
 gtgttgatgacgtcagtggtcccatgtagtgggacctggggactgcaacttaaggatttggcaggtagg
 5 cagggccttgggctgtggtggccctgggtgggtggggaccagggagagcagctgtccagctgcccagtaa
 ctcaagttccctgacatcgctgtcaacattgtctcctgcagctcagccctggatgggtgccccttcttg
 aaaccttaggatacctctgctggctccagctgccccctccctgtgagtcagctccttcaagccacagcc
 cgccagatggcttccaaggcaccaggtgacgctcctgacctgatgcctctcagctccaggacttccc
 aggacccctcagctgcccctggaccctgctgctactgcccgtcacctctgcaccttgtccccagctgggct
 10 gctgactcagatatgccaggtcctatgctatcatttcaactcccaggtcagctcactccaggagcct
 agttggagaatggatttccccagctgaaggacgcttcagcta (SEQ ID NO: 27)

In another embodiment, an enhancer sequence as described herein comprises a nucleotide sequence which contains one or more regions of 50-500 bp or longer, 50-250 bp or longer, 100-200 bp or longer, or 100 bp or longer, having at least 70% or
 15 greater, at least 75% or greater, at least 80% or greater, at least 85% or greater, at least 90% or greater, or at least 95% or greater sequence identity to the following human polynucleotide (DNA) sequence, called E14 herein, located in the human genome sequence at human_hg38 start 88802240 /human_hg38 stop 88802877 (FIGS. 1A-2 and 1A-3):

gcacttcagttccttttcatcaaggaactgattaaagtgtggtctattatctctgtagtggggagtccc
 gaaagattccacttcttctcttcttcttgcccaatgagaggggtcaggaagcttccaccaccatcctgactg
 tggccaccacatcctggtgtgaagcaccaccagcttccctccacaagaactctgaaggctaccagccagc
 ttgagtcctccgaagggtgctgtggtcaccagcagcctgtgtagcagagagcacaagacctgggactc
 gaactcagtcccaccttcaagagggatctccaccactttctgagcctcagttttcacatctttggttag
 25 gggcaggtgggaagttgccatatttaccttggttgggtcctttggaaaaattaaatgaaatgttaatgta
 tgttacacaacgggtcataggagggttcaaaagctgctagttctttttctccttttctgaaaacgg
 tattaagagtactgtagagacactggagaattcctgcttcatatgagatgtccttgggtctctcccggtg
 gaatttgaagacccagagactcgcagctccccgtgagcctccccgctgacgccttacctccccctctcc
 accatccccctgccatcc (SEQ ID NO: 28)

In another embodiment, an enhancer sequence as described herein comprises a nucleotide sequence which contains one or more regions of 50-500 bp or longer, 50-250 bp or longer, 100-200 bp or longer, or 100 bp or longer, having at least 70% or
 30 greater, at least 75% or greater, at least 80% or greater, at least 85% or greater, at least 90% or greater, or at least 95% or greater sequence identity to the following human polynucleotide (DNA) sequence, called E15 herein, located in the human genome

sequence at human_hg38 start 88803290 /human_hg38 stop 88803678 (FIGS. 1A-2 and 1A-3):

tgcccttccccctccccctgtcgccgccccctcggtccctgggggtgggggtttccctttgcgctcgcccc
 tcccgcccccaacctcacgggccctccccctcccccgccgtccctatgtatgtgtcacagcgcccat
 5 gcccggcccgcccgcccaacctccccgcccgtccagagggggctcgagagctgaggacgcgcgcag
 cgctgctcaaggtctctctctctcagcaccctcgccggccggcgctgtgacgcgggtgccaggggtctccg
 ggcacctttcagtggtccattccctcagccagccaggactccgcaaccagcagttgccgctgcggccac
 agccccgaggggacctgcggacaggacgcggcgaggaggagggt (SEQ ID NO: 29)

In another embodiment, an enhancer sequence as described herein comprises a
 10 nucleotide sequence which contains one or more regions of 50-500 bp or longer, 50-
 250 bp or longer, 100-200 bp or longer, or 100 bp or longer, having at least 70% or
 greater, at least 75% or greater, at least 80% or greater, at least 85% or greater, at least
 90% or greater, or at least 95% or greater sequence identity to the following human
 polynucleotide (DNA) sequence, called E16 herein, located in the human genome
 15 sequence at human_hg38 start 88807290 /human_hg38 stop 88807962 (FIGS. 1A-2
 and 1A-3):

gccttgggaaaggggatcaggagcacatcttgggaacaggagccttccctctcctgctgtcaacggcctg
 gcctcgtggccatgcttcgtgtcctgatggtagccgcactgccgccccgataacttaaaggaagccccct
 tcaatgggatgaagggcccggtttctgtgacagctactttgggagtggcagctcccccttccccagatcc
 20 aacaggacacagccatccctgcgagggagtctgggccccctggacgcagttgaagaagggccactgggag
 gccggcagaacaaggggagggctggagaaggagcgggagtgcaggcgagaggaggaccagagagggggga
 atttgtagagaaaacggtaaaacggtttcttttttcaaaagttgaatccaggggcaagaacggaaacggt
 ggagtttacttttaaaaactcagagcctccttataagtggtagtggtgtgtgtgtgtgtgtgtgtgtgt
 tgtgtgtgtgtgtgtgcatgctggcagggcgggccctgcgggctggccaggcctgtgagaggtgacttctc
 25 tccctctcaatggccttacagtgtatcctaaagaagctttttgttaaaactcatgaataggtggatttgg
 ggtgtgtgatgctggccatcacgatttgatatttggttacctttgagggtta
 (SEQ ID NO: 30)

In another embodiment, an enhancer sequence as described herein comprises a
 nucleotide sequence which contains one or more regions of 50-500 bp or longer, 50-
 30 250 bp or longer, 100-200 bp or longer, or 100 bp or longer, having at least 70% or
 greater, at least 75% or greater, at least 80% or greater, at least 85% or greater, at least
 90% or greater, or at least 95% or greater sequence identity to the following human
 polynucleotide (DNA) sequence, called E17 herein, located in the human genome
 sequence at human_hg38 start 88833390 /human_hg38 stop 88833984 (FIGS. 1A-2
 35 and 1A-3):

TtcatTTtcagggtccccgtcccccttccctctgcctcacctatgctgtcgccctgtcagcacctaattcag
 cttcccatggagaaaggcctccctgttgacagggccgtgctgggactcagggctgccaaagtcagtctt
 ctgcataaaaggctcagtgagtcctggagacacttggaagccagacagaatggaatttctccctatTTt
 tctttacagctgagaaaaacacacacaaacacaagagcatatTTtattgcgatatttctatcccaaagttt
 5 gtctTTtaaaaaaaaaaaaaaagaaataaaattagtttctctgcccactccaccctatccccctacccccac
 tctcctccccacctctTTttcattccttcccatTTtctggtttaggccagggagagaaatcaagccgtcc
 aagccccacagagcactcctacacccccggacaatgtccagcttgttcagagagtgaggggaagaaacac
 agctccgaaaacatacacacaacctcccaatgaaggggtgttctgaggggaagaacaggcgggccttgtgt
 ctgaacacgaatccctaagggtctgagggaagagaggagccgga (SEQ ID NO: 31)

10 In another embodiment, an enhancer sequence as described herein comprises a
 nucleotide sequence which contains one or more regions of 50-500 bp or longer, 50-
 250 bp or longer, 100-200 bp or longer, or 100 bp or longer, having at least 70% or
 greater, at least 75% or greater, at least 80% or greater, at least 85% or greater, at least
 90% or greater, or at least 95% or greater sequence identity to the following human
 15 polynucleotide (DNA) sequence, called E18 herein, located in the human genome
 sequence at human_hg38 start 128377753 /human_hg38 stop 128378783 (FIGS. 1A-2
 and 1A-3):

ggctcttaagtaggaaaaacacatgggtgtgatttgggttggtgaatagtcgacttgtagatctgcccact
 cgtcaacattgctctgacatcagattttctgaagagcagtggtgggctccttccccagggctggccgagt
 20 tttgaggggaactgcaggttctgatgtttccaaactctgtatctctgccctcgtcatttccatggacaag
 ttatttgtgctggtgtgagatccagaatcggtcctgctacgtgacaatgcctggagcacggagccagga
 acccaagctgcctcaccagcgtgttaggggtgttactgtgcccgttttagaagatcacttggaggtgca
 aaaatagagcagttttttttttgttttttttttttttgacatggagttctcgctctgtcgcccagggtggag
 tgcagtgggcgcaatctcggtcactgcaagctccgcctccccgggttcacgccatttctcctgacctagcc
 25 tccccagtagctgggactacaggagcctgccaccatgcccggttaattttttttgtatttttttagtagag
 acgggggttgaccatgttagccaggatgggtctcgatctcctgacttcatgatccgcccgcctcggcctc
 ccaaagtgtctgggattacaggcgtgagccaccacgcctggccaagagcagatttttttaaaaaaattaa
 gtacctctattcatttgcaccttcaactaccagtgaggagatcaaaatttcttagagcaaatgcattcg
 atgccactcacagatttctgacaggagagcacaaatttcagggaacgcctacatcaaagcactaattggcac
 30 ttttacagtgcttttctccgcacgtgagccttgtggtggaaggagctgtcatagtaatgcgtattcct
 ccatgcccagtgagtaggggtgacggtcaattcacagttcactaggcacaaaagatgacggggctctcct
 ctgctcgggacagcaagaagggttgaggtgatacggtttgtcggtgtccccacccaaatctcatct
 (SEQ ID NO: 32)

35 In another embodiment, an enhancer sequence as described herein comprises a
 nucleotide sequence which contains one or more regions of 50-500 bp or longer, 50-
 250 bp or longer, 100-200 bp or longer, or 100 bp or longer, having at least 70% or

greater, at least 75% or greater, at least 80% or greater, at least 85% or greater, at least 90% or greater, or at least 95% or greater sequence identity to the following human polynucleotide (DNA) sequence, called E19 herein, located in the human genome sequence at human_hg38 start 128289803 /human_hg38 stop 128290279 (FIGS. 1A-2 and 1A-3):

```
catgaaagtcactggttttacataactaattgcattttcccaaaacatcaagctaaatacataactgata
gcatgtttaaaggctcctatgtttcacctcaaattatctcatatttcacatgaggcaaaccctgtcctga
ggccctgatgaagatgggcaggcagatctgatatcagctgcttttcttctccttgatggaacctctggat
tgcggtgccctatcctataatgtgaaaaagggtctccagaaaagggtggaggaattactttctgaaattc
tgaagggtggtatccaaagggtgcagaaaggaacattatcttctaccatataaaaaccagtagggcggtg
gatgctgggacactgtatgagtccattctcactactgccataaagaagtacctgagactgggtatcttat
aaaggaaagagggttggttgactcacagttctgcaggcttaatagaaagcataactgggagggc
```

(SEQ ID NO: 33)

In another embodiment, an enhancer sequence as described herein comprises a nucleotide sequence which contains one or more regions of 50-500 bp or longer, 50-250 bp or longer, 100-200 bp or longer, or 100 bp or longer, having at least 70% or greater, at least 75% or greater, at least 80% or greater, at least 85% or greater, at least 90% or greater, or at least 95% or greater sequence identity to the following human polynucleotide (DNA) sequence, called E20 herein, located in the human genome sequence at human_hg38 start 128323153 /human_hg38 stop 128323718 (FIGS. 1A-2 and 1A-3):

```
aaaaatgaagaatttatatgccaaactccaggagaaatatttcagtgaggctcctggcttggtgcaaggat
atggaaccagagaccaagaattctatcagttaaaagcagcttagtttcttgagcctggactgatggggga
cgtggaagacaaagtgtcagggtccatcagtggaagattggccttgagccactgtacacagaatggagag
cccactggcctaaaaggagagattgtcaggcgtgacgaagcaggaatttttagccgaagaatattcacia
attacaggccaagaggggaagtggggacgttcgtcttctctcatagccttgctcgttgggggaccagct
gtcctttattgttaatagaaaaatcaatatagcaagaggcgaatctttgctgtgataacattggctcct
ttcaccaggcggtgtggaattagattactgatagatgcacctctgtcgctccccagggtccagatagaa
tctatgggcttttgccaataagcacggtaacagagtggtggatcaggaaccagcgggtggccaatggcagt
ggagaaaatgtaat
```

(SEQ ID NO: 34)

In another embodiment, an enhancer sequence as described herein comprises a nucleotide sequence which contains one or more regions of 50-500 bp or longer, 50-250 bp or longer, 100-200 bp or longer, or 100 bp or longer, having at least 70% or greater, at least 75% or greater, at least 80% or greater, at least 85% or greater, at least 90% or greater, or at least 95% or greater sequence identity to the following human

polynucleotide (DNA) sequence, called E21 herein, located in the human genome sequence at human_hg38 start 128332503 /human_hg38 stop 128332974 (FIGS. 1A-2 and 1A-3):

atagaccatataattctcacatgtcaagggttttaagccaaagccctcaggcaccacttctgattttctt
 5 gaaggatcaaaaataaaaagggttgcaaccctcacagccgtaggctcctgcagcaactctttggtgcacct
 gtcacctgatacctgggaggaggctctgagtcctatgctgtggaagggtgctggccttcatggatgggc
 ctccccctggtgtgtccactgtggacctgagtggtgtcccagagccctggctctcccttcttttccctct
 agaaagggaatcgcatgttcccaatcatctctgagattatctttattcttcaaggagttgcagtggct
 cttgccaagtgccctgggtcttgacatctgcctagtggccctgtagagacctccacctccagacag
 10 ctcagaatttgctaaagaaaatgtgaattttggagtcagggttagaaaatatgacat

(SEQ ID NO: 35)

In another embodiment, an enhancer sequence as described herein comprises a nucleotide sequence which contains one or more regions of 50-500 bp or longer, 50-250 bp or longer, 100-200 bp or longer, or 100 bp or longer, having at least 70% or
 15 greater, at least 75% or greater, at least 80% or greater, at least 85% or greater, at least 90% or greater, or at least 95% or greater sequence identity to the following human polynucleotide (DNA) sequence, called E22 herein, located in the human genome sequence at human_hg38 start 128336003 /human_hg38 stop 128336491 (FIGS. 1A-2 and 1A-3):

aagaaaagttttatttttgcctctgtagtattgggggtttaagtgatcacggtaattttccattatcattt
 20 tgtgttttaataataacaacaggctttattgtgaaaatatttgtctaataattgggcagtaaatgttta
 agtgattttggtttaattactattacagtcatactattacagtgcatataataagaattcttcttgagtt
 tgttcattagatgggaagaggctgcatttttaaaaaatatatgcatgcctataatactacatttaata
 tgtgcgtatataaagagatgctttcttatttatatacatggtcattatagagctttgtgagaaatagaa
 25 ttttctctgtgcaatctgtactctgggaggggttatttggctgacactgtatgccatttccctaacagaa
 tgtctctagttaagtaatcatatgatgaagacatcccagctgggactctatatattaagccaagttacta
 tttcta (SEQ ID NO: 36)

In another embodiment, an enhancer sequence as described herein comprises a nucleotide sequence which contains one or more regions of 50-500 bp or longer, 50-250 bp or longer, 100-200 bp or longer, or 100 bp or longer, having at least 70% or
 30 greater, at least 75% or greater, at least 80% or greater, at least 85% or greater, at least 90% or greater, or at least 95% or greater sequence identity to the following human polynucleotide (DNA) sequence, called E23 herein, located in the human genome sequence at human_hg38 start 128365603 /human_hg38 stop 1283366181 (FIGS. 1A-2 and 1A-3):

ctaaaaatgcctcctcgctctgatttttagccgtggttgttgagtagccggttcagcaggagctgtga
 ttccattgagctctcaaaccataaaatgcaaatctccgaggatggctcctctccctgccccacag
 ttgtgctccgaatagtgtctgagtttcattttttacaaggggcctttaaaaactcctgggcccccttgaaa
 actcccagccccctttgtccagatggggatggaggtggccaggctgccccgttgattgtgtgccgagga
 5 gccctccccgggaaggctgtgattttatacgcgcaggccttgtcacggggtgaaaggaagggccacttttt
 cttttgatccaatgttaggtttgaaagccaccactgctgtaaactcagctggatccgcggggcctga
 ttaaacacattgcccgtttgttgccgagatggtgtttcggaaaggcgtgtgaatgcacttcctttgc
 ggggctcacacagacaagatgtgtgttgcaaggatgagggcgcctgctcggcctccagcccagggccggg
 aaggggagaagggtgctgtgcgtcgctgc (SEQ ID NO: 37)

- 10 In another embodiment, an enhancer sequence as described herein comprises a
 nucleotide sequence which contains one or more regions of 50-500 bp or longer, 50-
 250 bp or longer, 100-200 bp or longer, or 100 bp or longer, having at least 70% or
 greater, at least 75% or greater, at least 80% or greater, at least 85% or greater, at least
 90% or greater, or at least 95% or greater sequence identity to the following human
 15 polynucleotide (DNA) sequence, called E24 herein, located in the human genome
 sequence at human_hg38 start 128375853 /human_hg38 stop 128376606 (FIGS. 1A-2
 and 1A-3):

Ttgaggcaagagcgcagggtggcatatccagggtggccactgggtctggagtgtcagtagcagggcagat
 ttagaagggtgactttgcatacctaggcaaggccagctcatgcgggatgtcggagcccatgggaagcacc
 20 ttgcgtttgaggctgcctgcgggtgggaagcttcagagtttcaagcggggctttgctatgggtttgttct
 gctttcccgttttcccttttgaggaggccttacagagatagtgatgactttgcagctgttaatcatcag
 gaagctgtaatcactaagaatgtttgaaatcatcagtttaaggattttttagaaggaagtaaaccaaagaa
 atactgcagtagcctgccctaattatttcctgggcttaaaagtaaccagggtgcattggagagattat
 tttcttctctgatttatgaaggctctcagggtccaaattttgaaaactgctgatcgaatttgttcttgatg
 25 ttgtcatagaaatctgaaactttcctacttgtctgagagtgaatcttttgattattcactcaagggt
 ttgatagggtttaaaaaaaggccttcgggacatctcttgttataaagtgtcaacttttagatatcaagaga
 atcatgatatatatttattactacaaaagagaaaataagcaactgaaaaactcatgaacttgaagcatgaa
 gcaaaccccttaagttctaggggtttcaagatgtggatgccaacatgtgatgacattttaaaaga
 (SEQ ID NO: 38)

- 30 In another embodiment, an enhancer sequence as described herein comprises a
 nucleotide sequence which contains one or more regions of 50-500 bp or longer, 50-
 250 bp or longer, 100-200 bp or longer, or 100 bp or longer, having at least 70% or
 greater, at least 75% or greater, at least 80% or greater, at least 85% or greater, at least
 90% or greater, or at least 95% or greater sequence identity to the following human
 35 polynucleotide (DNA) sequence, called E25 herein, located in the human genome

sequence at human_hg38 start 128408553 /human_hg38 stop 128408930 (FIGS. 1A-2 and 1A-3):

ggcttttgggtttttacaaaatattacaagttgcctaaatagtccgtgtttaaggacatagagccagagc
tctttctggaatgtcatacctcggcagggccttttgtgcatgttttaagctgattctgaaattaggggg
5 ttaaaatggaagcgccgagccatccctaaagagagggagggcgaatgtgcccttgttgctggtgacccca
gaacaaggcctctgggctgagaacaggagagaatgttatttctttgaaaagccatcttgacaatccaag
tccgtttggctgcagcaccaaaggcagctttgatctgctcgccagtgtccctgccgggaaaaggattag
ggccttccagaggacagcagagccaggctgcc (SEQ ID NO: 39)

In another embodiment, an enhancer sequence as described herein comprises a
10 nucleotide sequence which contains one or more regions of 50-500 bp or longer, 50-
250 bp or longer, 100-200 bp or longer, or 100 bp or longer, having at least 70% or
greater, at least 75% or greater, at least 80% or greater, at least 85% or greater, at least
90% or greater, or at least 95% or greater sequence identity to the following human
polynucleotide (DNA) sequence, called E26 herein, located in the human genome
15 sequence at human_hg38 start 13388723 /human_hg38 stop 13390212 (FIGS. 1A-2
and 1A-3):

gcccaggctggagtgtggtggcaaatctcagataactgaaacctctgcttcccaggctcaagccatcc
tcccacctctgtctgcagagtagctgagactataggcatgtgccacaatgctcagataattacttaaca
ttctagtagagtctagtagacatgggctatcactatgttgccctggctggtctggaactcctgggctca
20 agtgattgttctgccttggcttcccaaagtgttgggattacggctgtaagccgcatgcttggcttcgc
tttacaattttttttttttttttttttgagacagagtcttactctgccaccaggctggagtgtagtggc
tagattttggctcactgcaaactctggcccttgggttaagagattctcctgcctcagcttcccaagtag
ctgggattacaggcatggacaaccatacctggctaataattttgtattagcagagacggtatttcaccgt
gtcggccgggctggtctcgaactcccgacctcatgatccgcctacctcgggctcccaaagtgtgggat
25 tacaggcatgagccaccgtgcttggccaagaagacattttgttttctcaaaaaagtggagatctgagct
tcaaagatccttggtaacacttcccagtgctatcagtgtagtggtgcagtggctaataattcatggacc
ctataggagggatcttgctgctcttttagagggtgggacacactcttcttggtagcagaagggcagaac
tatgcctctgtggccacttattgcagaatggaattggagtaaactgagggccctttcacacatgctaga
gaactgactttggccctaggagaagtgggggttgaggggattggcctgagaaacttgcttttctactg
30 gattgtcctctagagtttttcactggagatttgtcagaatgagcctccagtcccatccagactcctgg
agctggcaggccagagcctgctgaggaaccagttcttgaccatcttcatcctggacgagctgccaggg
aggtcttccctctgatgttcatggaggcctccagcatgagacattttgagggcctgaagctgatggtgc
aggcctggcccttctcctccgcctcctctgggatccctgatgaagacacctcatctggagaccttgcaag
ctgtgctgaagggacttgatacactgctggcccagaagcttcgccccagggtgaggtgactcaggtggcc
35 tgggtgggaagggtccaggcatccagggaaggacagctggctcaggaggagtgggtgggggtggggagct
aggggtggctcagaggcttctgatggtgcccattgagagaccttgaccattgccagatcctctggaaaag

gactgctcaccatacaggggtccactgaggaacaggaacctgcttcctcccagtggaaggtaaaggttc
tagaagtgagaaccaggcagaatccaagggggagcgggatg (SEQ ID NO: 40)

In another embodiment, an enhancer sequence as described herein comprises a nucleotide sequence which contains one or more regions of 50-500 bp or longer, 50-
5 250 bp or longer, 100-200 bp or longer, or 100 bp or longer, having at least 70% or greater, at least 75% or greater, at least 80% or greater, at least 85% or greater, at least 90% or greater, or at least 95% or greater sequence identity to the following human polynucleotide (DNA) sequence, called E27 herein, located in the human genome sequence at human_hg38 start 13469123 /human_hg38 stop 13470861 (FIGS. 1A-2
10 and 1A-3):

ttgggttccaatggaactacagagagcaatgactactggtcctgaatgtgggtataaagttttacccta
aaagtaccttgtatTTTTTTTccaaggccaaaatatacaactcatttgcacccctggaaaggatatgttta
tttaaaaaaaaaagattgcatTTTgcaaacagtagagaacactgctctTTTTTatttaaaaaaatctttac
catggaaaaacaataaagtttgcgtgtgtgtttattggctctggggacttaaagaacaaagataccttgt
15 ggattgcagacaaatgaaacccacagaggtttgctttgggtaggtttcaccatacaggtgtttcaatag
accactttgcaaataataaattacttaacactcaaggccgctagaggccactaaaaaggagtttatggc
caaggcacagggctgggtggctggctcagtgagcgggtggcaggatattaatgagactcagagcctggacg
tgctctggatccagttaaatgtaatagagttggaaaaccacctgccccagccactgacggcacctagg
attcatgcctgtaactttgaccatctgagcctgtagggacattggggaggagggggagggtgagaggag
20 gcagtggcaacagcagcatggatgttccatgcaaaccttctctgtcaccagggaaagcagtcctgagca
catgaatttcttagcctctcttccaggatgaagcctagtttgaaccagcactgccaggttgaagtgtta
ctgcatcctgcagccagagccagggcatgtggccaccccccttggtccctgctgtggtacccagagtcac
ttggaacatgtgtgaagccaggatgaggggtgcataaccctccagatgctgatattctaaatatattacaa
gtcacaataacagagaaaactgtTTTTTTTTTgttattgttgttgttttgagaaggagtctcgctct
25 gtgcgccaggctggagtgcagtagcgtgatctcggtcctcctcctcccggttcacgccattct
cctgcctcagcctcccaagtagctgggattataggcatgcaccaccacgcctagcaaatTTTgtatTTT
tagtagagatggggTTTTTctccatgttgggtcaggctggtctcaactcctgatctcaggtgatctgccac
ctcagcctcccaaagtgtgtgggattacaggtgtgagccaccgcacccggccaacaaaagtaccttctta
atgacttcgaagactaggtttaaatggtaaattattaaattcttggaaatctgccacagaatatggcat
30 tgtggggacagctgagctgattgaaacctgctccctTTTcttccactccagctccatcctgcacct
taggggtctatgcacacctgtgtggacatccaccctcacatccaacctctattcacattccccaccac
catcctgtgtggccactcagcctgctctaaagcagggatgctgggaagatgccacatccaagcttggga
atcgTTTTTgccagaaattgggggccctaagtacccaaaaaatgttctagaaggggacatgttctggat
ggccatggactccttgcctccctgggggaagagcacagctggaggaggactggagcaaggccccctaaagc
35 actggaccaagataatgcccctcttggccagggtccaagggctgtactagtgggtaccgctgtcatcac
agcattcattactg (SEQ ID NO: 41)

In another embodiment, an enhancer sequence as described herein comprises a nucleotide sequence which contains one or more regions of 50-500 bp or longer, 50-250 bp or longer, 100-200 bp or longer, or 100 bp or longer, having at least 70% or greater, at least 75% or greater, at least 80% or greater, at least 85% or greater, at least 90% or greater, or at least 95% or greater sequence identity to the following human polynucleotide (DNA) sequence, called E28 herein, located in the human genome sequence at human_hg38 start 31124894 /human_hg38 stop 31125629 (FIGS. 1A-2 and 1A-3):

cctcggaaaccaaggttggtctctggcacctgtagggccacgggcagctatgtcagcttcctcgggaaggac
 cgaggctgggtctctggcatctccccgaccaatcctgggtccactgtgtacccctgaagggcagaaaacag
 ctactgcccaccgcagctccagcctggccccaacatctgtgggcccagctgggtgatgtctgcctcagctg
 gaccaaagcctccccagcgaccacaggtcagttctgggtccgacgtccctggggctgggtgatgcctg
 cctcagcaggggccaagatctccccagtcaccctggggcccaatctggccccaacctccagagaccaga
 agcaggagccacctgcctccgtgggacccaagccaacactggcagcctctggcctgagcctggcctgg
 cttctgaggagcagccccagaaactccctccaccccttccccgggtgccagtcagttctgtctccaa
 ctcaggaacaggccctgggtccagcatccacggcatcaggcgcagcctctgtgggacagacatcagcta
 gaaagaggggatgccccagccccctagacctctccctgcttctgaggggcatctccagcctccagctcaga
 catctggtcctacaggtccccaccctgcatccaaacctcccagaccctcggtctccccctccttcc
 gagcccggtc tgaggccctccacagcagccctgaggatcctgtttt (SEQ ID NO: 42)

In another embodiment, an enhancer sequence as described herein comprises a nucleotide sequence which contains one or more regions of 50-500 bp or longer, 50-250 bp or longer, 100-200 bp or longer, or 100 bp or longer, having at least 70% or greater, at least 75% or greater, at least 80% or greater, at least 85% or greater, at least 90% or greater, or at least 95% or greater sequence identity to the following human polynucleotide (DNA) sequence, called E29 herein, located in the human genome sequence at human_hg38 start 31132544 /human_hg38 stop 31133831 (FIGS. 1A-2 and 1A-3):

agcctggccaacatgggtgaaacgctgcctctactaaaaatacaaaaattagccaggcgtgatggcggtta
 cctgtggtctcagctactggggaggctgagacaggagaatcacttgaaccggggaggtggaggttcag
 tgagccgagattgcaccactgcactccagcctgggtgacagagcgagactccatctcagaaaaagaaaa
 aaaaaaaaaagagtctctgagtttacagatgagggccctggcattcagagaggctgaggaactcaccca
 gcctgtcaacggcagaaccagagccaaatccaggatttgctagcttcaaagctatgttctcactcactc
 cctaaggaggctgtgggcagaaggaacctgggctgggaggcagcacagggttggtatttatactaga
 cctgttctgcctcagtttcccagctctgtaaagtggccctttgtctcaggcaattttgtgctaagacccaa
 gagccttaagtgtgtgggatactagagggtctccctgatgtggccccctgccccctgccttgccctggac
 agtttgccttcagggaacgacatgccactgggtgcggctggaggtggcagatgagtgggtgccccgagc

aggcgggtggtgaggtaccgcatggaaacagtgttcgcccgcagctcctgggactggatcggcttatacc
 ggggtgagagggggcagtggtggtcagcgactcaggggaagaaaggggcctggaggagcagctgaacagcat
 ggtggggtcactggcttgtccagatcttgatgccacactgggagactgctgggatcagacattataggg
 tcacaacactgattccacaacactgatccccaggtgggtttccgccattgcaaggactatgtggctta
 5 tgtctggggccaaacatgaagatgtggatgggaatacctaccaggtacttaaaaggagtgggagagtcag
 ggcaagtccttgttgcctttgggacctcagaactcaccttgggggctctcaggtggcctccctgacccc
 caacttaggcttatacccttgggcctaccaggttaacattcagtgaggaatcactgcccgaaggccatgg
 agacttcactcctgggctactatagtcacaaccacagcatcctcatcggcactgaacccttcaggt
 aagtaggccagactgctgggtgggggtgcctaaagacttttgtcaaagtccacagcctctacattctg
 10 ctcccttgagttcagacaaataacctgacctcccaagatctgccaac (SEQ ID NO: 43)

In another embodiment, an enhancer sequence as described herein comprises a
 nucleotide sequence which contains one or more regions of 50-500 bp or longer, 50-
 250 bp or longer, 100-200 bp or longer, or 100 bp or longer, having at least 70% or
 greater, at least 75% or greater, at least 80% or greater, at least 85% or greater, at least
 15 90% or greater, or at least 95% or greater sequence identity to the following human
 polynucleotide (DNA) sequence, called E30 herein, located in the human genome
 sequence at human_hg38 start 88655733 /human_hg38 stop 88657379 (FIGS. 1A-2
 and 1A-3):

acatcttaaacagtccttttaatgttatgaatttgatttttcaagaaatacatgtcatttattttcaaaa
 20 cgaaatgatagctatactttctagagtctatcaatagtattttaaaataagatactcataactttcfaat
 actgcttttactagtcactcgtcattaaatgtaactgtatattcaagagcttttctaataatagcct
 ttaattaaacgaaggactgtagagggtttctgttgccctttgaagttcttaattattacttgtatcca
 gcattttatggtacacttaagggttaaatataatcattttaaataacattgaagagaaatatgaagactt
 ttgcccatttttaattaaatctctgaatttcagtatgttgaataataaacatatgttttgatttttttt
 25 catggccgaatggcaaatgctcactatattaaacaacaaaaaagaaatggttagctttttatgggact
 aatcgctaagcagatgcatgtaaagagctattttctatgcatggcttccaaaagtgctaattaaatag
 ttggtattcaaggctatgctcgtcattgttttagtgacacacaaatccagcgatgtgtgcccagcagaca
 ttttaagttgaatgttttctcctctacggtctttgtcatgaaatggtggcaccatgatgagaacactag
 tgaagcaaaacattgaaatatgctttaataatgttttaacatgtagtgacactagcctagttttcta
 30 atgaatttttaatttctgttttcttataagggtgatatgagttatcgctgatgcatattaaatcatata
 catgagtcatttttctctaaatttgcataaaatggctaaatgctaatagcaccaaaggagcttactatat
 gtggtacagcaaatattcccttgaagattttctgcaatcaatctcctgtattttcattagcaaccagata
 aggtgtggtctgcagaataaaaaaagaaagtgtgtagctcatgaacttatgaggcttcagatgatttc
 tacgtggtgatttagagtggttctgcaattagaatttatgtaggtaaaacacacatgtgcttccttttaa
 35 aggacagtgcaacaaaagtctgaatacagccttgcaattgttaaacaaatgaaaaggcaccattcaat
 tattgtgatttttttacatctataattaaatgaaggaaagccatacttttaatttagtatcatttgatt
 ggcataacccttactgaaattttacaatttccctactatgtttataaaaagaacttttaaaaataacat

gtgtgaaatattttgtttgctaactgttcccatTTTTccttgtcaaataatggtgaagaattttctggac
 taatgtttaacattttaaaaatgttttttctatcatcaaatactcttactgaactgacattaggatcata
 tgctttataaaaaattgcattagggtaacagtattattgggcaaaccagagatgtttacttgaaggata
 aacttgctgcttactcactccactcatcaacccttttctcgtctcctacagttccaccatctggaatat
 5 tttttaaccagt aaagaaaaaattggggaaggggatggctattttaaaaataaatgcttt

(SEQ ID NO: 44)

In another embodiment, an enhancer sequence as described herein comprises a
 nucleotide sequence which contains one or more regions of 50-500 bp or longer, 50-
 250 bp or longer, 100-200 bp or longer, or 100 bp or longer, having at least 70% or
 10 greater, at least 75% or greater, at least 80% or greater, at least 85% or greater, at least
 90% or greater, or at least 95% or greater sequence identity to the following human
 polynucleotide (DNA) sequence, called E31 herein, located in the human genome
 sequence at human_hg38 start 88872683 /human_hg38 stop 88872997 (FIGS. 1A-2
 and 1A-3):

15 tattatcctagtaaataccttaaaaaactttaagaggtgggtgattttataattcccatTTTtacagatcc
 tgggtactggggctttctgggtcattaaaacacctgcctaaaaccactaatcagtaaattgggaggctggct
 tttgaaccagtttttgggtcgttgttcttaatacattattctttattgtttatggacatgtttgtctaata
 gcataatatgtagaatcaaagaaatgatattaagtgtggaaatggagtctccaaactctttatgcttgt
 ttaaacgatcttctctctcgcgagagtgtatcttcatcctt (SEQ ID NO: 45)

20 In another embodiment, an enhancer sequence as described herein comprises a
 nucleotide sequence which contains one or more regions of 50-500 bp or longer, 50-
 250 bp or longer, 100-200 bp or longer, or 100 bp or longer, having at least 70% or
 greater, at least 75% or greater, at least 80% or greater, at least 85% or greater, at least
 90% or greater, or at least 95% or greater sequence identity to the following human
 25 polynucleotide (DNA) sequence, called E32 herein, located in the human genome
 sequence at human_hg38 start 88745133 /human_hg38 stop 88745535 (FIGS. 1A-2
 and 1A-3):

tttcccttactcagctaacaacattttaccaagtatctgctgtgtgctaacgcttaggtgttaaactgg
 gcatacaaactgaatgagaaagagtttgcctccacagagctgagcgtcctagagagatgtgccagatg
 30 ttgcaatcataatgcaatgagaaatgtaattgttggtacaggctactatgtaagcacaggaaagaggtgc
 ataacttgtctgttagagtcaggaaaggcttttctcaaatggctgaactgaattctgtgggatgacaaa
 gagtgctcaatagcatgaagcagaagaaggaaaggcatgctaggattgcataggaagagtaagcgcc
 gtgacattgccaaagtggcgccacagtgtagcaattaagagcacacactgaggccgggt

(SEQ ID NO: 46)

In another embodiment, an enhancer sequence as described herein comprises a nucleotide sequence which contains one or more regions of 50-500 bp or longer, 50-250 bp or longer, 100-200 bp or longer, or 100 bp or longer, having at least 70% or greater, at least 75% or greater, at least 80% or greater, at least 85% or greater, at least 90% or greater, or at least 95% or greater sequence identity to the following human polynucleotide (DNA) sequence, called E33 herein, located in the human genome sequence at human_hg38 start 88799783 /human_hg38 stop 88801354 (FIGS. 1A-2 and 1A-3):

ataaaaatatcaggtaaaatataatctctcatctctcatcttctctccatctccctatgtccctttccctc
tctctctctctctctctctctcacacacacacacacacacacacacacacacacacacacagagaga
gagacacacatgttcttctcctctaaaaagaaaaaccaataatcctctactgagacagttgtgaatcaaag
gtttcttctgacaggagttacatccatctctgaatttcctagagagcagcaaagggccttgtgttttatt
ccccttccacacttaatcactggactgtggggccagactgaatgagtagctcattagaatcactgagtt
cactgaggggatgagagattccttcctggctgggtgctaagtgatactcccataaggattttgtggta
caaaacgtgctggatatggaggtaacctgtctgggagtccctgtcactccaaggatcacttggaatgctc
tggaaaaaacacatgacctggctgaatgagttctgtttgaattgtttagcctacaccttcatttcagcagc
ttatactgcattaatgaggttattgttcctttgccgtccaattgttcccaagctgattttttgcatata
tgttttacatccttaacaagaatgcctgtctcctgctgttttcagagtctcttccacagtgctgagcatg
agtggagcttgctaaatcattgctaaatgaagcaatgggctgtaagcatgtcctgtgggatctgcatct
tcagatcatcctgaagtactcaacaaccacatcttcttccaggaacagagcccaacataaaactggtagg
gtttgctgtcttagacagctaagagaacgaggagtggagctagtgaacaagcagtggaagggggcagttc
cttaatgccatccgaactgaatttcaacagtctgacaagctagcgttttgggtaaaatcccagtatac
ttgtcacagagttaagtaaaaatggacttccttcaaagggaagtgccttttaatacaataaactgtttttgtt
tttttaaccaatggattaaaaaatttaacacatttactaaatctggcataatttatatattgtatctaaac
agatattcaagctgcattataatataatcataaaaaaactgatctcagtcgtctgtgtaagcctttgtg
agtctttgtgccattgttggagtagtgctaattatcaagcaaagacatgataattacgcagagcctttt
tgctaaaagaaggaatctttttcaacacccacgcactgcacaatttcctatgaccctgtagcactactc
tggtatggcccagaaaatttgtattttctgtgtaaaaggctggaaattatatttttctatctctcccgata
ccttttcttcttgtgagtaaaactgttttttagagggttaaggaagaggggtaatgggccaaatgggaaat
ataaaaactaatgacccttcatgaaacatattatgctcctcattaaacttattcagttaaatgtattcca
ttaaattaaaaataaataaggatttaaaattttaccaggagcagtaaacaaatctt

(SEQ ID NO: 47)

In another embodiment, an enhancer sequence as described herein comprises a nucleotide sequence which contains one or more regions of 50-500 bp or longer, 50-250 bp or longer, 100-200 bp or longer, or 100 bp or longer, having at least 70% or greater, at least 75% or greater, at least 80% or greater, at least 85% or greater, at least

90% or greater, or at least 95% or greater sequence identity to the following human polynucleotide (DNA) sequence, called E34 herein, located in the human genome sequence at human_hg38 start 27969472 /human_hg38 stop 27969690 (FIGS. 1A-2 and 1A-3):

5 t g c t g a a c c a g t t c t c c g c t g c a t c g t t c c g c t c g c g c t c g g g a c c t g c a a c a g a a g g g a a t g g g a c c
g a g t g t c a g t c t g g a c t c t c c a t c t c c c c g c a c t a c t c c g c t c c c c c t t t t t a g c c c g c t c t c a a a a g
c c t c t t c a a c a t c a a g g g c a t c t c c c a a g t t g a a a g a a a a a a a t t t t c t c t g g a g c c t c t c a g c a c t t
a c t t a t t t t a g c a (SEQ ID NO: 48)

In another embodiment, an enhancer sequence as described herein comprises a nucleotide sequence which contains one or more regions of 50-500 bp or longer, 50-250 bp or longer, 100-200 bp or longer, or 100 bp or longer, having at least 70% or greater, at least 75% or greater, at least 80% or greater, at least 85% or greater, at least 90% or greater, or at least 95% or greater sequence identity to the following human polynucleotide (DNA) sequence, called E35 herein, located in the human genome sequence at human_hg38 start 27973822 /human_hg38 stop 27974489 (FIGS. 1A-2 and 1A-3):

[illegible]

In an embodiment, an enhancer sequence as described herein comprises a nucleotide sequence which contains one or more regions of about 100 bp or longer having at least 75% or greater sequence identity to a human polynucleotide (DNA) sequence of the above-described E11 to E35 enhancer element sequences (e.g., SEQ ID NOs: 25-49). In another embodiment, an enhancer sequence comprises a nucleotide sequence which contains one or more regions of about 100 bp or longer having at least 75% or greater sequence identity to a human polynucleotide (DNA) sequence of the above-described E1 (SEQ ID NO: 15), E2 (SEQ ID NO: 16), E5

(SEQ ID NO: 19), E6 (SEQ ID NO: 20), E11 (SEQ ID NO: 25), E14 (SEQ ID NO: 28), E22 (SEQ ID NO: 36), or E29 (SEQ ID NO: 43) enhancer element sequences, for example.

Genetic Epilepsy with Febrile Seizures plus (GEFS+) and Dravet syndrome

5 Genetic epilepsy with febrile seizures plus (GEFS+) is a rare condition that constitutes a spectrum of seizure disorders of varying severity. GEFS+ is usually diagnosed in families whose members have a combination of febrile seizures, which are triggered by a high fever and recurrent seizures (epilepsy) of other types, including seizures that are not related to fevers (afebrile seizures). The additional
10 seizure types, called generalized seizures, usually involve both sides of the brain; however, seizures that involve only one side of the brain (partial seizures) occur in some affected individuals. The most common types of seizures in individuals with GEFS+ include myoclonic seizures that cause involuntary muscle twitches; atonic seizures that involve sudden episodes of weak muscle tone; and absence seizures that
15 cause loss of consciousness for short periods that appear as staring spells. While GEFS+ is usually diagnosed in families, it can occur in individuals with no history of the condition in their family.

 The most common and mildest feature of the GEFS+ spectrum is simple febrile seizures, which begin in infancy and typically stop by the age of five. When
20 the febrile seizures continue after age five, or when other types of seizures develop, the condition is called febrile seizures plus (FS+), which typically cease in early adolescence.

 Dravet syndrome (DS), also known as severe myoclonic epilepsy of infancy (SMEI) or early infantile epileptic encephalopathy-6 (EIEE6), is a condition
25 frequently considered to be part of the GEFS+ spectrum and is the most severe disorder in this group of disorders. The term Dravet syndrome is preferably used, because not all affected individuals exhibit myoclonic epilepsy. Affected infants typically have prolonged seizures that last several minutes (status epilepticus) and are triggered by fever. Other types of seizures, including afebrile seizures, begin in early
30 childhood. These seizure types can include myoclonic or absence seizures. In Dravet syndrome, these seizures are difficult to control with medication, and they can worsen over time. A decline in brain function is also common in Dravet syndrome. Children

affected with Dravet syndrome usually develop normally in the first year of life, but then development stalls; some affected children lose previously acquired skills and suffer developmental regression. Many children afflicted with Dravet syndrome have difficulty coordinating movements (ataxia) and have intellectual disabilities.

5 Causes of GEFS+

Mutations in several genes, including some that have not been identified, can cause GEFS+. The most commonly associated gene is *SCN1A*. More than 80% of Dravet syndrome cases and about 10% of other GEFS+ cases are caused by changes in this gene. Mutations in other genes have been found in only a small number of
10 affected individuals or families. The *SCN1A* gene and other genes associated with GEFS+ encode subunits of ion channels that transport positively charged ions into cells. The transport of these ions helps generate and transmit electrical signals between neurons (nerve cells). Mutations in the *SCN1A* gene have a variety of effects on sodium channels. Many genetic mutations that cause or are associated with Dravet
15 syndrome reduce the number of functional channels in each cell. Mutations that cause the milder GEFS+ disorders likely alter the channel's structure. All of these genetic changes affect the ability of the channels to transport sodium ions into neurons. Some deleterious mutations are thought to reduce channel activity while others may increase it. Changes in GABAergic receptor subunit genes impair the channel's function,
20 causing uncontrolled signaling between neurons, which likely leads to seizures. Without wishing or intending to be bound by theory, some studies have reported that certain *SCN1A* gene mutations cause constant stimulation of signaling between neurons. Such overstimulation of certain neurons in the brain triggers the abnormal brain activity associated with seizures.

25 While it is not known if all *SCN1A* gene mutations have the same effect, genome-wide association studies have demonstrated that loss of function of the voltage-gated sodium channel Nav1.1 encoded by the *SCN1A* gene is the most prevalent cause for Dravet syndrome. Previous studies using mouse models of Dravet syndrome suggest that it is the loss of *SCN1A* gene function in GABAergic
30 interneurons that is the primary defect underlying the seizures that represent the most deleterious symptom of this syndrome.

In animal studies, it was found that *SCN1A* $-/-$ mice developed severe ataxia and seizures and died on postnatal day 15. *SCN1A* $+/-$ mice had spontaneous seizures and sporadic deaths beginning after postnatal day 21, with a notable dependence on genetic background. Loss of *SCN1A* did not change voltage-dependent activation or inactivation of sodium channels in hippocampal neurons. However, the sodium current density was substantially reduced in inhibitory interneurons of *SCN1A* $-/-$ and $+/-$ mice. (Yu, F.H. et al., 2006, *Nat Neurosci*, 9(9):1142-1149; Yu et al., 2007, *Nat Neurosci*, 10(1):134). The studies suggested that reduced sodium currents in GABAergic inhibitory interneurons resulting from heterozygous *SCN1A* mutations may cause the hyperexcitability that leads to epilepsy in patients with SMEI.

GABAergic cortical interneurons

GABAergic interneurons, which release the neurotransmitter gamma-aminobutyric acid (GABA) are inhibitory neurons of the central nervous system and are essential for regulating and maintaining neural circuitry and activity. (Kelsom, C. and Lu, W., 2013, *Cell Biosci.*, 3:19). GABAergic interneurons of the mammalian cerebral cortex comprise several different cortical interneuron subtypes that may be categorized and classified by their expressed protein markers.

Interneurons play a key role in the wiring and neural circuitry of the developing nervous system of both invertebrate and vertebrate organisms. In general, an interneuron is a specialized type of neuron (nerve cell) whose primary role is to form a connection between other types of neurons. Interneurons, which are neither motor neurons nor sensory neurons, differ from projection neurons in that projection neurons send their signals to more distant locations, such as the brain or the spinal cord. Critically, interneurons function to modulate neural circuitry and circuit activity. A large majority of interneurons of the central nervous system are of the inhibitory type. In contrast to excitatory neurons, inhibitory cortical interneurons typically release the neurotransmitters gamma-aminobutyric acid (GABA) and glycine. Cortical interneurons are localized in the cerebral cortex, which is defined as a sheet of outer neural tissue that functions to cover the cerebrum and cerebellum structures in the brain. (*Id.*)

GABAergic interneurons include numerous interneuron subtypes that may be categorized by the surface markers they express. Four major cortical interneuron

subtypes are parvalbumin (PV)-expressing interneurons, somatostatin (SST)-expressing interneurons (which constitute a heterogeneous population), and ionotropic serotonin receptor 5HT3a (5HT3aR)-expressing interneurons. These three subtypes together account for approximately 100% of the neocortical GABAergic interneuron population in mice. Although these interneurons home to their respective layers of the cerebral cortex, they are generated in various subpallial locations and they subsequently migrate to the cerebral cortex.

Cortical circuit function is maintained by the balance between excitatory inputs and inhibitory inputs. A disruption of the balance of neural circuits is likely to contribute to the emergence of neurological, neurodevelopmental, or neuropsychiatric disorders such as, without limitation, epilepsy, autism spectrum disorders, and intellectual disabilities.

The role of GABAergic cortical interneurons

GABAergic neurons play an inhibitory role and synaptically release the neurotransmitter GABA to regulate the firing rate of target neurons. Neurotransmitter release typically acts through postsynaptic GABA_A ionotropic receptors in order to trigger a neuronal signaling pathway. Interneuron role/function is typically categorized into three components: (1) afferent input, (2) intrinsic properties of the interneuron, and (3) targets of the interneuron. In general, interneurons receive input from various sources, including pyramidal cells, as well as cells from other cortical and subcortical regions. (Kelsom, C. and Lu, W., 2013, *Cell Biosci.*, 3:19). With regard to output, cortical interneurons engage in feed-forward and feedback inhibition. Regardless of the mode of output, the cortical interneuron network is further complicated by the fact that a single cortical interneuron is capable of making multiple connections with its excitatory neuronal target(s).

Cortical interneuron subtypes

It is estimated that there are over 20 different subtypes of GABAergic interneurons in the cerebral cortex. The subtypes are also distinguished from each other based upon the calcium-binding proteins they express, which serve as markers. Based on studies performed in both mouse and rat brain tissue, the calcium-binding protein, parvalbumin (PV), and the neuropeptide somatostatin (SST), are key markers

found to define the most predominant interneuron subtypes within the cerebral cortex. Of particular note, the PV-expressing interneuron population is independent from the SST-expressing population, in that expression of these markers does not overlap. In addition to PV- and SST-positive GABAergic interneurons, which together comprise approximately 70% of the total GABAergic cortical interneuron population, another subgroup of interneurons that express 5HT3aR were found to comprise approximately 30% of all interneurons. These three interneuron subpopulations account for nearly or equal to 100% of all GABAergic cortical interneurons, yet each of these populations, especially the 5HT3aR-expressing population, is heterogeneous and expresses other proteins or neuropeptides that contribute to their characterization. (Kelsom, C. and Lu, W., 2013, *Cell Biosci.*, 3:19).

Parvalbumin (PV)-expressing interneurons

PV-expressing interneuron represent approximately 40% of the GABAergic cortical interneuron population. This population of interneurons possesses a fast-spiking pattern, and fire sustained high-frequency trains of brief action potentials. These interneurons also possess the lowest input resistance and the fastest membrane time constant of all interneurons.

Two types of PV-interneurons comprise the PV interneuron group: basket cells and chandelier cells. Basket cells are interneurons that make synapses at the soma and proximal dendrite of target neurons, and usually have multipolar morphology. Several studies have shown that fast-spiking basket neurons are the dominant inhibitory system in the neocortex, where they mediate the fast inhibition of target neurons, among many other functions. Such fast-spiking basket neurons likely play a large role in regulating the delicate balance between excitatory and inhibitory inputs in the cerebral cortex. Unlike basket neurons, the chandelier cell subgroup of PV-expressing interneurons targets the axon initial segment of pyramidal neurons. Both basket cells and chandelier cells are fast-spiking, but they differ in electrophysiological properties. In contrast to other interneurons, chandelier cells may be excitatory rather than inhibitory due to their depolarizing effects on membrane potential. (Kelsom, C. and Lu, W., 2013, *Cell Biosci.*, 3:19).

Another group of PV-expressing cells that is independent from chandelier and basket neurons in the neocortex, e.g., mouse neurocortex, are called multipolar

bursting cells, which differ from chandelier and basket cells in both electrophysiology and connectivity. Multipolar bursting neurons possess synapses with pyramidal cells (or other multipolar bursting cells) that demonstrate a paired-pulse facilitation; in contrast, chandelier and basket cells are usually strongly depressing. (Kelsom, C. and
5 Lu, W., 2013, *Cell Biosci.*, 3:19).

Somatostatin (SST)-expressing interneurons

SST-expressing interneurons constitute the second-largest interneuron group in the mouse neocortex and represent approximately 30% of the total cortical interneuron population. SST GABAergic interneurons represent a heterogeneous
10 population of cortical interneurons. SST-positive interneurons are called Martinotti cells and possess ascending axons that arborize layer I of the cerebral cortex and establish synapses onto the dendritic tufts of pyramidal neurons. Martinotti cells are also found throughout cortical layers II-VI, but are most abundant in layer V. In contrast to PV-positive interneurons, excitatory inputs onto Martinotti cells are
15 strongly facilitating. Additional subpopulations of SST-expressing cortical interneurons show differences in firing properties, expression of molecular markers and connectivity of different neurons within this population. (Kelsom, C. and Lu, W., 2013, *Cell Biosci.*, 3:19).

5HT3aR-expressing interneurons

20 The third population of GABAergic cortical interneurons is designated as the 5HT3aR interneuron group, which accounts for approximately 30% of the GABAergic cortical interneuron population. Based on mouse studies, this population of GABAergic interneurons in the cortex express the 5HTa3 receptor, but do not express either PV or SST.

25 5HT3aR interneurons represent a heterogeneous population. Within the 5HT3aR interneuron group are several subsets of interneurons that also express other protein or neuropeptide markers, including vasoactive intestinal peptide (VIP). VIP-expressing interneurons are localized in cortical layers II and III. The VIP-expressing interneurons do not express PV or SST, but do express the 5HTa3 receptor, accounting
30 for approximately 40% of the 5HT3aR population. VIP interneurons generally make synapses onto dendrites; some have been observed to target other interneurons.

Compared with other cortical interneurons, VIP interneurons possess a very high input resistance and are among the most excitable of interneurons.

60% of cortical interneurons in the 5HT3aR-expressing population do not express VIP. Of this VIP-negative 5HT3aR group, nearly 80% express the interneuron marker reelin. In this latter category of cortical interneurons, the neurogliaform cell population, called spiderweb cells, express neuropeptide Y (NPY), and exhibit multiple dendrites radiating from a round soma. Neurogliaform interneurons can form synaptic connections with each other as well as with other interneuron types, in contrast to other types of interneurons that can only make synapses onto homologous neurons. Thus, neurogliaform cells play an important role in regulating neural circuitry and function by activating slow GABA_A and GABA_B receptors in order to provoke long-lasting inhibitory postsynaptic potentials onto pyramidal neurons and other interneurons.

Pyramidal neurons

Pyramidal neurons, also known as pyramidal cells, are neurons with a pyramidal shaped cell body (soma), which ranges from 20-120 μm in diameter, and two distinct dendritic trees. The basal dendrites emerge from the base and the apical dendrites from the apex of the pyramidal cell body. Like most neurons, pyramidal neurons have multiple dendrites and a single axon, but both dendrites and axons branch extensively. The dendrites of pyramidal neurons are usually regarded as input structures, receiving synaptic contacts from other neurons, while the axon serves as its output to other neurons. Pyramidal neuron dendrites can also release retrograde signaling molecules (e.g. endocannabinoids), so communication is somewhat bidirectional. The extensive branching of the dendrites and the axon allows a single neuron to communicate with thousands of other neurons in a network. (Spruston, N., 2009, *Scholarpedia*, 4(5):6130).

Pyramidal neurons are found in forebrain structures, such as the cerebral cortex, hippocampus, and amygdala, but not in the olfactory bulbs, striatum, midbrain, hindbrain, or spinal cord of mammals, as well as birds, fish and reptiles. Pyramidal neurons are the most populous members of the excitatory family of neurons, e.g., neurons that release the neurotransmitter glutamate, in the brain areas that they inhabit, such as brain cortical structures. Their abundance suggests that they

play critical roles in the functioning of the nervous system, as well as in cognitive processing. Pyramidal neurons comprise about two-thirds of all neurons in the mammalian cerebral cortex, where they function to transform synaptic inputs into a patterned output of action potentials. Pyramidal neurons receive synaptic inputs from
5 tens of thousands of excitatory synapses and several thousand inhibitory synapses. Most of the excitatory inputs use glutamate as the neurotransmitter, e.g., glutamatergic pyramidal neurons, while inhibitory inputs use GABA.

While the nature of a stimulus can determine the type of output generated by a pyramidal neuron (e.g., single spike vs. burst), the intrinsic neuronal excitability is
10 another important determinant of how the neuron responds to an input. Typically, neurons are classified according to how they respond to current injection, which may vary from one type of pyramidal neuron to the next. Most pyramidal neurons respond to continuous depolarizing current injection with a train of spikes that exhibits spike-frequency adaptation (accommodation). Many pyramidal neurons respond with one
15 or more bursts of action potentials. The nature of this response is largely determined by the types of voltage-gated ion channels expressed in the neuron, but the structure of the dendritic tree is also important (Mainen, Z.F. et al., 1996, *Nature*, 382:363-366; Spruston, N., 2008, *Nature Reviews Neuroscience*, 9:206-221; Spruston, 2009, *Scholarpedia*, 4(5):6130).

20 **Adeno-associated Virus (AAV)**

AAV is a small (25 nm), nonenveloped virus that contains a linear single-stranded DNA genome packaged into the viral capsid. It belongs to the family *Parvoviridae* and is of the genus *Dependovirus*, because productive infection by AAV occurs only in the presence of either an adenovirus or herpesvirus helper virus. In the
25 absence of helper virus, AAV (serotype 2) can establish latency after transduction into a cell by specific but rare integration into chromosome 19q13.4. Accordingly, AAV is the only mammalian DNA virus known to be capable of site-specific integration. (Daya, S. and Berns, K.I., 2008, *Clin. Microbiol. Rev.*, 21(4):583-593).

There are two stages to the AAV life cycle after successful infection: a lytic
30 stage and a lysogenic stage. In the presence of adenovirus or herpesvirus helper virus, the lytic stage persists. During this period, AAV undergoes productive infection characterized by genome replication, viral gene expression, and virion production.

The adenoviral genes that provide helper functions for AAV gene expression include E1a, E1b, E2a, E4, and VA RNA. While adenovirus and herpesvirus provide different sets of genes for helper function, they both regulate cellular gene expression and provide a permissive intracellular milieu for a productive AAV infection.

- 5 Herpesvirus aids in AAV gene expression by providing viral DNA polymerase and helicase as well as the early functions necessary for HSV transcription.

In the absence of adenovirus or herpesvirus, AAV replication is limited; viral gene expression is repressed; and the AAV genome can establish latency by integrating into a 4-kb region on chromosome 19 (q13.4), called AAVS1. The AAVS1 locus is near several muscle-specific genes, TNNT1 and TNNI3. The AAVS1 region itself is an upstream part of the gene MBS85 whose product has been shown to be involved in actin organization. Tissue culture experiments suggest that the AAVS1 locus is a safe integration site.

15 **Recombinant AAV (rAAV) as a vector for gene delivery and therapeutic treatment**

- AAVs are well suited for use as vectors and vehicles for gene transfer to the nervous system, as they enable gene expression and knockdown, gene editing, circuit modulation, *in vivo* imaging, disease model development, and the assessment of therapeutic candidates for the treatment of neurological diseases. AAVs provide safe, long-term expression in the nervous system. Most of the foregoing applications rely on local AAV injections into the adult brain to bypass the blood-brain barrier (BBB) and to temporally and spatially restrict transgene expression.

- AAV vectors have been highly successful in fulfilling all of the features desired for a delivery vehicle, such as the ability to attach to and enter the target cell, successful transfer to the nucleus, the ability to be expressed in the nucleus for a sustained period of time, and a general lack of pathogenicity and toxicity. Recombinant AAV (rAAV) is advantageous as a delivery vector, particularly for delivery to interneurons in brain tissue, as it is focally injectable; it exhibits stable expression over time; and it is both non-pathogenic and non-integrative into the genome of the cell into which it is transduced. Twelve human serotypes of AAV (AAV serotype 1 (AAV-1) to AAV-12) and more than 100 serotypes from nonhuman primates have been reported to date. (Daya, S. and Berns, K.I., 2008, *Clin. Microbiol.*

Rev., 21(4):583-593). In addition, rAAV has been approved by the FDA for use as a vector in at least 38 protocols for several different human clinical trials. AAV's lack of pathogenicity, persistence and its many available serotypes have increased the potential of the virus as a delivery vehicle for a gene therapy application in
5 accordance with the described compositions and methods.

Recombinant AAV (rAAV) vectors have been constructed that do not encode the replication (Rep) proteins and that lack the *cis*-active, 38 base pair integration efficiency element (IEE), which is required for frequent site-specific integration. The inverted terminal repeats (ITRs) are retained because they are the *cis* signals required
10 for packaging. Thus, current recombinant AAV (rAAV) vectors persist primarily as extrachromosomal elements.

Recombinant AAV (rAAV) vectors for gene therapy have been based mostly on the AAV-2 serotype. AAV-2-based rAAV vectors can transduce muscle, liver, brain, retina, and lungs, requiring several weeks for optimal expression. The
15 efficiency of rAAV transduction is dependent on the efficiency at each step of AAV infection, i.e., virus binding, entry, trafficking, nuclear entry, uncoating, and second-strand synthesis.

Several novel AAV vector technologies have been developed to either increase the genome capacity for AAV or enhance gene expression. *Trans*-splicing
20 AAV vectors have been used to increase the capacity of the vector for harboring heterologous polynucleotides by taking advantage of AAV's ability to form head-to-tail concatemers via recombination in the ITRs. In this approach, the transgene cassette is split between two rAAV vectors containing adequately placed splice donor and acceptor sites. Transcription from recombined AAV molecules, followed by the
25 correct splicing of the mRNA transcript, results in a functional gene product. While somewhat less efficient than rAAV vectors, *trans*-splicing AAV vectors permit delivery of therapeutic genes up to 9 kb in size and have been successfully used for gene expression in the retina, lung and muscle.

Polynucleotides encoding rAAVs as described herein comprise an *SCN1A*
30 enhancer polynucleotide sequence. Because of its nature as an enhancer, the orientation of the enhancer polynucleotide sequence, i.e., 5'-3' or 3'-5', is not material to its function. Accordingly, the enhancer sequences (e.g., the E1-E10, e.g., E2, a PV-

specific enhancer sequence, or E5 or E6, as described herein) may be used in a reverse orientation and may be used as reverse-complementary sequences. A “PV-specific enhancer” refers to the enhancer sequences described herein that target and restrict expression of a transgene in PV-expressing cortical interneurons (PV-cINs) as described herein.

Moreover, the enhancer need not be specifically spaced relative to other sequences, such as the *SCN1A* coding sequence. In addition, the rAAV polynucleotides may include additional elements, for example, a sequence encoding a reporter or a detectable marker, such as a fluorescent protein, or an element such as a Woodchuck Hepatitis Virus Posttranscriptional Regulatory Element (WPRE), which may increase RNA stability and protein yield. An rAAV polynucleotide may also comprise a promoter to drive transcription of one or more polynucleotides (genes) which are inserted between inverted terminal repeats (ITRs). A polyadenylation signal, such as bovine growth hormone polyadenylation signal and/or SV40 polyomavirus simian virus 40 polyadenylation signal, may be included as elements in the rAAV polynucleotide. The rAAV polynucleotide can comprise a minimal promoter, e.g., a human beta-globin minimal promoter (ph β g) and a chimeric intron sequence (Hermeming et al., 2004, *J Virol Methods*, 122(1):73-77). Without wishing to be bound by theory, ITRs may aid in concatamer formation in the nucleus after the single-stranded, AAV vector DNA is converted into double stranded (ds) DNA by host cell DNA polymerase complexes. Thus, the administration of the described rAAVs may form episomal concatemers in the nucleus of interneuron cells into which they are transduced. In non-dividing cells, such as adult interneurons, concatemers may remain intact in these cells for the lifetime of the interneurons. Advantageously, integration of rAAV polynucleotides into host chromosomes is likely to be negligible or absent and will not alter or affect the expression or regulation of any other human gene.

Recombinant AAV vectors can be made using standard and practiced techniques in the art and employing commercially available reagents. It will be appreciated by the skilled practitioner that rAAV vectors that been used in several clinical trials that have yielded promising results. By way of example, rAAV based therapy received marketing approval by the European Union in 2012, as reported by

Kotterman, M.A. et al., 2014, *Nat. Rev. Genet.*, 15:445-451. In some embodiments, plasmid vectors may encode all or some of the well-known replication (rep), capsid (cap) and adeno-helper components. The rep component comprises four overlapping genes encoding Rep proteins required for the AAV life cycle (e.g., Rep78, Rep68, 5 Rep52 and Rep40). The cap component comprises overlapping nucleotide sequences of capsid proteins VP1, VP2 and VP3, which interact together to form a capsid of an icosahedral symmetry. A second plasmid that encodes helper components and provides helper function for the AAV vector may also be co-transfected into cells. The helper components comprise the adenoviral genes E2A, E4orf6, and VA RNAs 10 for viral replication.

In an embodiment, a method of making rAAVs for the products, compositions, and uses described herein involves culturing cells that comprise an rAAV polynucleotide expression vector as described; culturing the cells to allow for expression of the polynucleotides to produce the rAAVs within the cell, and 15 separating or isolating the rAAVs from cells in the cell culture and/or from the cell culture medium. Such methods are known and practiced by those having skill in the art. The rAAVs can be purified from the cells and cell culture medium to any desired degree of purity using conventional techniques.

In an embodiment, the rAAV vector contains an *SCN1A*-restricted enhancer 20 polynucleotide sequence and a chemogenetic DREADD ('Designer receptor exclusively activated by designer drug')-encoding sequence, e.g., a Gq-DREADD receptor (Hu, J. et al., 2016, *J Biol Chem*, 291:7809-7820), or a. The amino acid sequence of the Gq-DREADD receptor has been reported by Armbruster et al. (2007, *Proc Natl Acad Sci USA*, 104:5163-5168). The amino acid sequence of the Gq- 25 DREADD receptor is a derivative of the amino-acid sequence of the human muscarinic acetylcholine receptor, M3, in which the tyrosine in position 149 is replaced by a cysteine, and the arginine in position 239 is replaced by a glycine. The unmodified human sequence is provided under NCBI accession no. NP 000731.1. In an embodiment, the polynucleotide sequence that encodes the Gq-DREADD receptor 30 in the rAAV vector can be modified, for example, by including optimized codons for expression of the Gq-DREADD receptor in human interneurons.

In an embodiment, the rAAV vector contains an *SCN1A*-restricted enhancer polynucleotide sequence and a chemogenetic PSAM-encoding sequence.

Recombinant AAV vectors, which have a genome of small size (about 5 kb), can be engineered to package and contain larger genomes (transgenes), e.g., those that are greater than 4.7 kb. By way of example, two approaches developed to package larger amounts of genetic material (genes, polynucleotides, nucleic acid) include split AAV vectors and fragment AAV (fAAV) genome reassembly (Hirsch, M.L. et al., 2010, *Mol Ther* 18(1):6-8; Hirsch, M.L. et al., 2016, *Methods Mol Biol*, 1382:21-39). Split rAAV vector applications were developed to take advantage of the fact that rAAV genomes naturally concatemerize in the cell post-transduction and are substrates for enhanced homologous recombination (HR) (Hirsch, M.L. et al., 2016, *Methods Mol Biol*, 1382:21-39). This approach comprises "splitting" a large transgene into two separate vectors and upon co-transduction, intracellular large gene reconstruction via vector genome concatemerization occurs via HR or nonhomologous end joining (NHEJ). In general, three strategies exist within the split rAAV approaches: overlapping, trans-splicing, and hybrid trans-splicing.

Fragment AAV (fAAV) as an approach for AAV-mediated large gene delivery was developed based on reports that attempted encapsidation of transgenic cassettes exceeding the packaging capacity of the AAV capsid resulted in the packaging of heterogeneous single-strand genome fragments (<5 kb) of both polarities. After transduction by multiple fAAV particles, the genome fragments can undergo opposite strand annealing, followed by host-mediated DNA synthesis to reconstruct the intended oversized genome within the cell. (Hirsch, M.L. et al., 2016, *Methods Mol Biol*, 1382:21-39).

An advantage and benefit of the vectors, compositions and methods described herein is the identification and use of sufficiently small enhancer elements (*cis*-acting elements) that are capable of specifically restricting gene expression to a defined population of cells, e.g., interneuron cells. In an embodiment, the enhancer element is at least one of the E1-E10 enhancer sequences as described herein, which are *SCN1A*-specific and restrict gene expression, e.g., the *SCN1A* gene, to interneuron cells such as GABAergic interneurons and PV-expressing GABAergic interneurons, or pyramidal neurons, such as glutamatergic pyramidal neurons. The genes (transgenes)

delivered by the rAAV vectors described herein are active and functional in the specific cells in which they are expressed, i.e., the products that they encode are produced, and are functionally expressed by the cells. By way of specific example, an rAAV vector as described herein which is engineered to contain an enhancer sequence that specifically restricts expression of a transgene, e.g., a reporter gene or *SCN1A*, to a GABAergic interneuron cell or a GABAergic, PV-expressing, cortical interneuron cell, transduces these specific cell types, and the encoded reporter protein, or Nav1.1 sodium channel in the case of *SCN1A*, is functionally expressed in the specific cell type. By way of another specific example, an rAAV vector as described herein is engineered to contain an enhancer sequence that specifically restricts expression of a transgene, e.g., a reporter gene or *SCN1A*, to pyramidal cell, such as a glutamatergic pyramidal cell in the brain cortex.

As another advantage, the described *SCN1A*-specific enhancer control elements E1-E10 are of a size/length (kb), e.g., less than approximately 2 kb, to allow for their insertion in a rAAV vector along with other effector element polynucleotide sequences, e.g., reporter polynucleotides, DREADDs, transgenes. By way of example, given the obligate minimal size of reporter elements (e.g., Enhanced green fluorescent protein (EGFP), orange fluorescent protein (dTomato)), alone or in combination with effector or reporter elements, (e.g. Channelrhodopsin (ChR2), DREADDs), which average about 700bp to 2kb, respectively, a maximum of ~2kb in packaging capacity remains for the insertion of a cis-acting DNA control element such as an enhancer sequence into an rAAV vector. The *SCN1A*-restrictive enhancer sequences identified and described herein are capable of restricting expression to a defined population of cells, e.g., interneurons or GABAergic interneurons, or pyramidal neuron cells, and are sufficiently small elements to allow for additional nucleic acid sequences, reporter elements and transgenes, to also be cloned into the AAV vector.

Cell-specific AAV capsids

The rational design of AAV vectors that display selective tissue/organ targeting has broadened the applications of AAV as vector/vehicle for gene therapy. Both direct and indirect targeting approaches have been used to enhance AAV vector cell targeting specificity and retargeting. By way of example, in direct targeting,

AAV vector targeting to certain cell types is mediated by small peptides or ligands that have been directly inserted into the viral capsid sequence. This approach has been successfully employed to target endothelial cells. Direct targeting requires detailed knowledge of the capsid structure such that peptides or ligands are positioned at sites that are exposed to the capsid surface; the insertion does not significantly affect capsid structure and assembly; and the native tropism is ablated to maximize targeting to a specific cell type. In indirect targeting, AAV vector targeting is mediated by an associating molecule that interacts with both the viral surface and the specific cell surface receptor. Such associating molecules for AAV vectors may include bispecific antibodies and biotin. The advantages of indirect targeting are that different adaptors can be coupled to the capsid without resulting in significant changes in the capsid structure, and the native tropism can be easily ablated. A disadvantage of using adaptors for targeting involves a potential for decreased stability of the capsid-adaptor complex *in vivo*.

In addition, AAV vectors may be produced that comprise capsids that allow for the increased transduction of cells and gene transfer to the central nervous system and the brain via the vasculature. (Chan, K.Y. et al., 2017, *Nat. Neurosci.*, 20(8):1172-1179). Such vectors facilitate robust transduction of neuronal cells, including interneurons. When used with enhancers and cell-type specific promoters, such AAVs provide targeted gene expression in neuronal cells of the nervous system.

For applications that do not require high expression levels per cell, the amount of virus used, i.e., the viral dose, could be lowered. Lowering the viral load used for systemic gene delivery can reduce cost and production burden and minimize a potential risk for adverse reactions to viral components.

Delivery of recombinant adeno-associated viral vectors and treatment approaches

In general, the delivery of an effector gene to treat a neurological disease at the genetic level, e.g., by modifying or correcting gene expression, such as by gene therapy, may be achieved using appropriate and effective vectors, such as viral or virus vectors, e.g., AAV or rAAV. The use of a rAAV vector provides efficient delivery of therapeutic genes to a cell where the genes are expressed. While other methods and approaches for delivering genes to cells involve, for example, the use of

purified DNA under hydrodynamic pressure, a shotgun approach using DNA adhering to gold particles, or lipid-DNA complexes, such methods and approaches frequently do not provide efficient gene delivery and result in gene expression that is lower than that required for therapeutic efficacy. Moreover, such methods are not applicable to human use. Viruses, on the other hand, represent natural vectors for the delivery and expression of exogenous genes in host cells *in vivo*.

An advantage associated with the use of rAAV as a viral vector is that rAAV transgene expression typically persists for years or for a life time, as has been demonstrated in animal models. This stands in contrast to non-rAAV viral vectors, which often lead to an initial burst of transgene expression that commonly disappears after a relatively short time, e.g., weeks.

To achieve enhanced therapy or treatment, the dose of rAAV vector that is required for a therapeutic response may be reduced, e.g., by using certain rAAV serotypes. Alternatively, the surface of the rAAV vector capsid may be altered to include specific ligands for attachment to target tissues and cells as described above. Another approach takes into consideration the trafficking of the virus particle from the endocytosomal vesicle to the nucleus. (Zhao, W. et al., 2007, *Gene Ther.*, 14:545-550; Daya, S. and Berns, K.I., 2008, *Clin. Microbiol. Rev.*, 21(4):583-593). Typically, the virus particle-to-infectivity ratio of rAAV vector preparations ranges from 10:1 to 100:1. The high ratios reflect incomplete or empty vector particles, as well as trafficking from the endocytosomal vesicle to the nucleus. During trafficking, the vector particle may become ubiquitinated and directed to a proteasome for degradation, rather than to the nucleus where the transgene may be expressed. It was found that ubiquitination and direction to the proteasome require phosphorylation of tyrosine residues on the surface of the rAAV vector capsid. When the seven tyrosine residues on the surface of the AAV-2 capsid were replaced phenylalanine residues, the multiplicity of infection (MOI) required for the detection of transgene expression was greatly reduced both in cell culture and in several mouse models of transduction of cells in the liver and eye. Consequently, the ability to increase transgene expression to therapeutic levels in the treatment of diseases may be enhanced.

One or more treatment approaches to gain control over seizures are embraced by the therapeutic products, compositions and methods described herein involving state-of-the-art gene therapy or pharmaco-genetic approaches. Such approaches may likely lead to the development of a clinically relevant therapies to alleviate the seizure symptoms of DS.

For direct delivery to the brain, rAAV vectors may be administered by open neurosurgical procedure or by focal injection in order to bypass the blood-brain barrier, to temporally and spatially restrict transgene expression, and to target specific areas of the brain, e.g., interneuron cells and brain tissue comprising these cells.

Systemic rAAV delivery (by intravenous injection) provides a non-invasive alternative for broad gene delivery to the nervous system; however, the high viral load required and relatively low transduction efficiency have limited wide adoption of this method. Several groups have developed rAAV capsids that enhance gene transfer to the CNS and certain tissues and cell populations after intravenous delivery. By way of example, AAV-AS capsid¹⁸ utilizes a polyalanine N-terminal extension to the AAV9.4719 VP2 capsid protein to provide higher neuronal transduction, particularly in the striatum. The AAV-BR1 capsid²⁰, based on AAV2, may be useful for more efficient and selective transduction of brain endothelial cells. Another AAV capsid, AAV-PHP.B, comprises a capsid that transduces the majority of neurons and astrocytes across many regions of the adult mouse brain and spinal cord after intravenous injection. In an embodiment, rAAV comprises a capsid which specifically transduced interneurons, including PV interneurons, in the cerebral cortex (brain).

Other modes of rAAV vector administration may include lipid-mediated vector delivery, hydrodynamic delivery, and a gene gun. In a particular embodiment, the rAAV vectors comprise a capsid that increases the likelihood of directly infecting or transducing interneuron cells, such as GABAergic interneuron cells and GABAergic, PV-expressing interneuron cells, or pyramidal cells, e.g., glutamatergic pyramidal cells, and brain tissue comprising these cells.

The virus vectors and compositions thereof as described herein may be used in the treatment of neurological, neurodevelopmental and neurodegenerative diseases and disorders, particularly, for the treatment of DS, which includes epilepsy and its

attendant, often severe, seizure symptoms. A characteristic that distinguishes categories of seizures is whether the seizure activity is partial (e.g., focal) or generalized. In an embodiment, virus vectors and compositions thereof as described herein are used to treat partial and/or generalized seizures. Partial seizures are typically considered to be those in which the seizure activity is restricted to discrete areas of the cerebral cortex. As will be appreciated by the skilled practitioner, a seizure is characterized as a simple-partial seizure if consciousness is fully preserved during the course of the seizure. If consciousness is impaired, then the seizure is characterized as a complex-partial seizure. Complex-partial seizures also include those that initiate as partial seizures and subsequently extend through the cortex; as such, these types of seizures are considered to be partial seizures with secondary generalization.

Generalized seizures encompass distant regions of the brain simultaneously in a bilaterally symmetric manner and can include sudden, brief lapses of consciousness, such as in the case of absence or petit mal seizures, without loss of postural control. Atypical absence seizures usually include a longer period of lapse of consciousness and more gradual onset and termination. Generalized tonic-clonic or grand mal seizures, considered as the main type of generalized seizures, typically have an abrupt onset without warning. The initial phase of the seizure usually involves tonic contraction of muscles, impaired respiration, a marked enhancement of sympathetic tone leading to increased heart rate, blood pressure and pupil size. After approximately 10-20 seconds, the tonic phase of the seizure typically evolves into a clonic phase, which is produced by periods of muscle relaxation superimposed on the tonic muscle contraction. The periods of relaxation progressively increase until the end of the ictal phase, which usually lasts no more than one minute. The postictal phase is characterized by unresponsiveness, muscular flaccidity, and excessive salivation that can cause stridorous breathing and partial airway obstruction.

Atonic seizures are characterized by sudden loss of postural muscle tone lasting approximately 1-2 seconds. While consciousness is briefly impaired, there is usually no postictal confusion. Myoclonic seizures are characterized by a sudden and brief muscle contraction that may involve one part of the body or the entire body. Without limitation, the rAAV products, compositions and methods of use thereof as

described herein embrace the prophylactic and/or therapeutic treatment of the above-described seizures, including the seizures afflicting those with DS. In an embodiment, the rAAV products, compositions and methods of use thereof as described herein are used for the prophylactic and/or therapeutic treatment of epilepsy associated with a loss of function or impairment of function of the sodium channel Nav1.1 encoded by the *SCN1A* gene. In a particular embodiment, the rAAV products, compositions and methods of use thereof as described herein are used for the prophylactic and/or therapeutic treatment of Dravet syndrome (DS). In another embodiment, the rAAV products, compositions and methods of use thereof as described herein are used for the prophylactic and/or therapeutic treatment of pharmaco-resistant epilepsy, which refers to an epileptic condition that is uncontrolled, despite use of two or more drugs that are suitable for treating this type of epilepsy and that have been administered at maximum tolerated doses (MTDs). In embodiments, a pharmaco-resistant epilepsy embraces a condition in which seizures have failed to be eliminated by previous anti-epileptic drug treatments or treatment combinations.

Pharmacogenetic approaches

Pharmacogenetic approaches are contemplated for use with the virus vectors, rAAV vectors, compositions thereof, and methods described herein. Such approaches deliver either Gq-DREADD receptor or PSAM into PV-interneurons specifically using a viral vector, such as a rAAV vector comprising an enhancer element (e.g., E1-E10) as described herein and a polynucleotide encoding a Gq-DREADD receptor or PSAM. The targeted PV-neurons, either in a specific region upon focal injection or throughout the cortex upon systemic injection, as dictated by the type of pathology being treated, stably express the receptor (Gq-DREADD or PSAM). Thereafter, an individual (patient) is administered the drug that activates the receptor (e.g. CNO or PSEM, respectively). This approach results in a controlled alteration of the excitability of the PV-interneurons expressing the receptor and allows for a dose-dependent and time-dependent modulation of the excitation/inhibition (E/I) balance in neurons (interneurons and PV-expressing interneurons), resulting in a normalization of brain activity.

Pharmaceutical Compositions

Provided also are pharmaceutical compositions or formulations for treating subjects who are afflicted with, or who are at risk of developing, a neurological or neurogenetic disease, disorder, or pathology such as DS. In an embodiment, the pharmaceutical composition includes an AAV vector or virus particle, such as one containing an *SCN1A*-specific enhancer sequence, as described herein (as active agent) and a pharmaceutically acceptable carrier, excipient, or diluent. When formulated in a pharmaceutical composition, an rAAV vector as therapeutic compound or product can be admixed with a pharmaceutically acceptable carrier, diluent, or excipient.

The therapeutic agent(s) may be contained in any appropriate amount in any suitable carrier substance, and is/are generally present in an amount of 1-95% by weight of the total weight of the composition. The composition may be provided in a dosage form that is suitable for a parenteral (e.g., subcutaneous, intravenous, intramuscular, or intraperitoneal) administration route, such that the agent, such as a viral vector described herein, is systemically delivered. In an embodiment, systemic injection of an rAAV vector as described herein allows for the characterization of specificity of expression across brain regions, particularly when a reporter product is also encoded by the vector. The pharmaceutical compositions may be formulated according to conventional pharmaceutical practice (see, e.g., Remington: The Science and Practice of Pharmacy (20th ed.), ed. A. R. Gennaro, Lippincott Williams & Wilkins, 2000 and Encyclopedia of Pharmaceutical Technology, eds. J. Swarbrick and J. C. Boylan, 1988-1999, Marcel Dekker, New York).

Pharmaceutical compositions may be formulated to release the active agent substantially immediately upon administration or at any predetermined time or time after administration. The latter types of compositions are generally known as controlled release formulations, which include (i) formulations that create a substantially constant concentration of the agent within the body over an extended period of time; (ii) formulations that after a predetermined lag time create a substantially constant concentration of the drug within the body over an extended period of time; (iii) formulations that sustain action during a predetermined time period by maintaining a relatively constant, effective level in the body with

concomitant minimization of undesirable side effects associated with fluctuations in the plasma level of the active substance (sawtooth kinetic pattern); (iv) formulations that localize action by, e.g., spatial placement of a controlled release composition adjacent to or in contact with a target site or location, e.g., in a region of a tissue or organ; (v) formulations that allow for convenient dosing, such that doses are administered, for example, once every one, two, or several weeks; and (vi) formulations that target a specific tissue or cell type using carriers, chemical derivatives, or specifically designed vectors (e.g., comprising a certain capsid composition) to deliver the therapeutic agent, e.g., to interneurons or PV-expressing GABAergic interneurons, or pyramidal neurons, e.g., glutamatergic pyramidal neurons. For some applications, controlled release formulations obviate the need for frequent dosing during the day to sustain the plasma level of the administered agent at a therapeutic level.

Methods by which to obtain controlled release in which the rate of release outweighs the rate of metabolism of the agent in question are not meant to be limiting. By way of example, controlled release is obtained by appropriate selection of various formulation parameters and ingredients, including, e.g., various types of controlled release compositions and coatings. Thus, the therapeutic agent is formulated with appropriate excipients into a pharmaceutical composition that, upon administration, releases the agent in a controlled manner. Examples include single or multiple unit tablet or capsule compositions, oil solutions, suspensions, emulsions, microcapsules, microspheres, molecular complexes, nanoparticles, patches, and liposomes.

The administration of a composition comprising a combination of agents for the treatment of a neurological disease or disorder such as DS may be by any suitable means that results in a concentration of the therapeutic that, combined with other components, is effective in ameliorating, abating, reducing, decreasing, or stabilizing seizures in a subject. The composition may be administered systemically, for example, formulated in a pharmaceutically-acceptable buffer such as physiological saline. In an embodiment, systemic injection of an rAAV vector as described herein allows for the characterization of specificity of expression across brain regions, particularly when a reporter product is also encoded by the vector.

Routes of administration include, for example, intracranial, parenteral, subcutaneous (s.c.), intravenous (i.v.), intraperitoneal (i.p.), intramuscular (i.m.), or intradermal administration, e.g., by injection, that optimally provide continuous, sustained levels of the agent in the patient. The amount of the therapeutic agent to be administered varies depending upon the manner of administration, the age, physical condition and body weight of the patient, and with the clinical symptoms of the neurological disease or disorder, such as DS. Generally, amounts will be in the range of those used for other viral vector-based agents employed in the treatment of neurological diseases and disorders, particularly in the brain, although in certain instances lower amounts are needed if the agent exhibits increased specificity. A composition is administered at a dosage that shows a therapeutic effect, such as, for example, ameliorating, abating, reducing, decreasing, or stabilizing seizures in a patient, as determined by methods known to one skilled in the art.

The pharmaceutical composition may be administered parenterally by injection, infusion or implantation (subcutaneous, intravenous, intramuscular, intraperitoneal, intracranial, or the like) in dosage forms, formulations, or via suitable delivery devices or implants containing conventional, non-toxic pharmaceutically acceptable carriers and adjuvants. The formulation and preparation of such compositions are well known to those skilled in the art of pharmaceutical formulation, and can be found, for example, in Remington: The Science and Practice of Pharmacy, *supra*. In particular embodiments, administration is systemic and parenteral, such as by injection or intravenous delivery.

Compositions for parenteral delivery and administration may be provided in unit dosage forms (e.g., in single-dose ampules), or in vials containing several doses and in which a suitable preservative may be added (see below). The composition may be in the form of a solution, a suspension, an emulsion, an infusion device, or a delivery device for implantation, or it may be presented as a dry powder to be reconstituted with water or another suitable vehicle before use. Apart from the active agent (e.g., viral vector or particle comprising enhancer sequences and polynucleotides encoding an effector gene and associated regulatory sequences, as described herein), the composition may include suitable parenterally acceptable carriers and/or excipients. The active therapeutic agent(s) may be incorporated into

microspheres, microcapsules, nanoparticles, liposomes, or the like for controlled release. Furthermore, the composition may include suspending, solubilizing, stabilizing, pH-adjusting agents, tonicity adjusting agents, and/or dispersing agents.

In some embodiments, the composition comprising the active therapeutic(s) (i.e., viral vector or particle described herein) is formulated for intravenous delivery. As noted above, the pharmaceutical compositions according to the described embodiments may be in the form suitable for sterile injection. To prepare such a composition, the suitable therapeutic(s) are dissolved or suspended in a parenterally acceptable liquid vehicle. Acceptable vehicles and solvents that may be employed include water, water adjusted to a suitable pH by addition of an appropriate amount of hydrochloric acid, sodium hydroxide or a suitable buffer, 1,3-butanediol, Ringer's solution, isotonic sodium chloride solution and dextrose solution. The aqueous formulation may also contain one or more preservatives (e.g., methyl, ethyl or n-propyl p-hydroxybenzoate). In cases where one of the agents is only sparingly or slightly soluble in water, a dissolution enhancing or solubilizing agent can be added, or the solvent may include 10-60% w/w of propylene glycol or the like.

Methods of administration and delivery

Administration of a viral vector or pharmaceutical composition as described herein to a subject, e.g., a patient or infant patient having DS. In embodiments, the viral vector, viral particle, or pharmaceutical composition may be delivered to a cell (a target cell such as an interneuron or a brain layer comprising interneurons) in any manner such that the viral vector, particle or composition is functional and active to express the sequences contained in the vector or virus particle. Illustratively, rAAV comprising an *SCN1A*-specific enhancer and an effector gene (e.g. *SCN1A*) polynucleotide sequence may be delivered to interneuron cells or tissue comprising interneuron cells to provide for targeted expression of *SCN1A* in the interneurons. Thus, viral vectors or viral particles are delivered to a cell by contacting the cell with a composition comprising the viral vectors, or viral particles and by heterologous expression of the polynucleotides harbored by the viral vector or viral particles in the cell. The polynucleotides harbored by the rAAV vector must be delivered to the cells of a subject in a form in which they can be taken up so that therapeutically effective levels of the encoded products can be produced.

Transducing rAAV vectors are used for the delivery and expression of genes encoding desired proteins, polypeptides, or peptides to cells, especially because of their high efficiency of infection and stable integration and expression (see, e.g., Cayouette et al., *Human Gene Therapy*, 8:423-430, 1997; Kido et al., *Current Eye Research*, 15:833-844, 1996; Bloomer et al., *Journal of Virology*, 71:6641-6649, 1997; Naldini et al., *Science*, 272:263-267, 1996; and Miyoshi et al., *Proc. Natl. Acad. Sci. U.S.A.*, 94:10319, 1997). By way of example, rAAV is engineered to contain a polynucleotide encoding an *SCN1A*-specific enhancer nucleic acid sequence as described herein that preferentially directs gene expression in specific interneuron cell types and is used to direct and restrict the expression of a gene, e.g., *SCN1A*, in GABAergic interneuron target cells or in pyramidal target cells, such as glutamatergic pyramidal cells. In an embodiment, expression of the gene can be driven from any suitable promoter, such as a promoter specific for the target cells. In an embodiment, the rAAV vector is administered systemically. In an embodiment, systemic injection of an rAAV vector as described herein allows for the characterization of specificity of expression across brain regions, particularly, for example, when a reporter product is also encoded by the vector.

Gene transfer can also be achieved using *in vitro* transfection methods. Such methods include the use of calcium phosphate, DEAE dextran, electroporation, and protoplast fusion. Liposomes can also be potentially beneficial for delivery of DNA into a cell.

Treatment methods and protocols

Provided are methods of administering a therapeutic agent to a subject in need, such as a subject having, undergoing, having experienced, and/or at risk of experiencing a neurological disease or disorder, more particularly, a seizure, epilepsy, or DS, and who also may be diagnosed with, or be suspected of having, or having symptoms of, a seizure disorder, or who is identified as being in need of such treatment, in which an effective amount of a viral vector or viral particle as described herein, or a composition described herein, is administered to the subject to produce a therapeutic effect. According to the described methods, a therapeutic effect includes, without limitation, that amount of rAAV that is introduced into a sufficient number of interneurons so as to inhibit, reduce, or ameliorate one or more symptoms of the

neurological disease or disorder, e.g., a seizure or epilepsy, or to prevent one or more symptoms subsequent to the administration of the rAAV vector product or composition to the subject. The amount of rAAV that is administered may be determined by the skilled practitioner in the art, such as a medical or clinical practitioner, and, as appreciated by one skilled in the art, is based on factors such as the size of the epileptic focus, the titer of the virus preparation and from data acquired in non-human primates (e.g., Colle, M.-A. et al., 2010, *Hum. Mol. Genet.*, 19:147-158). By way of example, from 10^{10} to 10^{12} rAAV particles may be used to transduce rAAV vectors or particles thereof to a therapeutically relevant number of interneurons. Identifying a subject in need of such treatment can be in the judgment of a subject or a health care professional and can be subjective (e.g. opinion) or objective (e.g. measurable by a test or diagnostic method).

The therapeutic methods (which include prophylactic treatment) in general comprise administration of a therapeutically effective amount of the agents described herein, such as an rAAV vector, a viral particle, or composition containing the aforementioned agents, to a subject (e.g., animal, human) in need thereof, including a mammal, particularly a human. Such treatment will be suitably administered to subjects, particularly humans or infant humans, suffering from, having, susceptible to, or at risk for a neurological disease or disorder, such as seizures and/or epilepsy, or DS. Determination of those subjects "at risk" can be made by any objective or subjective determination by a diagnostic test or opinion of a subject or health care provider (e.g., genetic test, enzyme or protein marker or biomarker, family history, and the like).

Viral vectors and pharmaceutical compositions as described can be used therapeutically to treat patients suffering from neurological or neurodegenerative diseases or disorders, e.g., seizures, epilepsy, or DS, or prophylactically to provide advanced treatment or protection to patients at risk for certain neurological or neurodegenerative diseases or disorders, such as a prophylactic vaccination to reduce, diminish, abate, or ward off a seizure, epilepsy, one or more symptoms of DS, or the severity thereof. A prophylactically effective amount of the rAAV vectors as described herein are not intended to be limiting herein, and may range between about 10^2 TU (transducing units) per kilogram body weight of the recipient and about 10^{20}

TU kilogram body weight of the recipient, or any TUs in between those values.

Mouse models of seizures and DS can be used to optimize dosages and regimens.

The therapeutic vectors as described herein may be administered to a subject in need thereof in an effective amount to normalize the excitability of SCN1A-

5 deficient interneurons and alleviate seizures and seizure symptoms of Dravet syndrome (DS). The vectors and methods described herein may be of therapeutic value for an individual, e.g., a human infant, child or adult, who experiences or is at risk for experiencing one or more seizures and/or DS. In an embodiment, an rAAV or a composition comprising an rAAVs as described herein is administered to an
10 individual whose interneurons do not express or exhibit loss of function or expression, at the time of administration, of the *SCN1A* gene encoding the Nav1.1 sodium channel, which is dependent on an *SCN1A*-specific enhancer, such as E1-E10 described herein, for expression. In an embodiment, the expression of SCN1a in interneuron cells transduced by the described rAAV vectors containing an *SCN1A*-
15 restricting enhancer sequence normalizes the excitability of interneurons deficient in, or having abnormal expression of, *SCN1A*. In an embodiment, a composition comprising an rAAV vector as described herein is administered to an individual whose interneurons no longer express the *SCN1A* gene. In an embodiment, a composition comprising an rAAV vector as described herein is administered to an
20 individual who is at least one month old. In embodiments, the individual is at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17 or 18 years of age.

Subjects, e.g., mammalian subjects, and human patients to whom the rAAV vectors as described herein are administered may also benefit from adjunct or additional treatments or therapeutic compounds or drugs, such as anti-seizure
25 modalities, including but not necessarily limited to, use with other anti-epileptic therapeutic agents, and/or surgical techniques. as are well known to those having skill in the art. By way of example, anti-epileptic drugs (AEDs) that may be used in conjunction with the therapeutic products and compositions described herein include, without limitation, Acetazolamide, Brivaracetam, Carbamazepine,
30 Clobazam, Clonazepam; Eslicarbazepine acetate, Ethosuximide, Gabapentin, Lacosamide, Lamotrigine, Levetiracetam, Oxcarbazepine, Perampanel, Phenobarbital, Phenytoin, Pregabalin, Primidone, Rufinamide, Sodium valproate, Stiripentol,

Tiagabine, Topiramate, Valproic acid, (available as Convulex, Epilim Chrono, Epilim Chronosphere), Vigabatrin and Zonisamide.

Kits

Also provided are kits for preventing or treating a neurological or neuropsychiatric disease, condition, or pathology, e.g., seizures and/or epilepsy, as well as the symptoms of Dravet syndrome (DS), in a subject in need thereof. In one embodiment, the kit provides a therapeutic or prophylactic composition containing an effective amount of a rAAV vector or viral particle as described herein, which comprises an enhancer polynucleotide sequence specific for the *SCN1A* gene that restricts the expression of an *SCN1A* gene, e.g., contained in the virus vector, to interneuron cells, including GABAergic interneuron cells in the brain (i.e., in the telecephalon), or to pyramidal cells, such as glutamatergic pyramidal cells in the brain cortex, or to VIP cells. In an embodiment, the *SCN1A*-specific enhancer is an E1, E2, E3, E4, E5, E6, E7, E8, E9, or E10 human enhancer sequence as described herein. In an embodiment, the *SCN1A*-specific enhancer is an E2 human enhancer polynucleotide sequence. In an embodiment, the *SCN1A*-specific enhancer is an E5 human enhancer polynucleotide sequence. In an embodiment, the *SCN1A*-specific enhancer is an E6 human enhancer polynucleotide sequence.

In another embodiment, the kit provides a therapeutic or prophylactic composition containing an effective amount of a rAAV vector or viral particle as described herein, which comprises an E11-E35 enhancer polynucleotide sequence, in particular a human E11-E35 sequence, specific for a gene expressed in a neuron or interneuron cell, especially a PV-expressing neuron.

In some embodiments, the kit comprises a sterile container which contains the therapeutic or prophylactic composition; such containers can be boxes, ampoules, bottles, vials, tubes, bags, pouches, blister-packs, or other suitable container forms known in the art. The containers can be made of plastic, glass, laminated paper, metal foil, or other materials suitable for holding medicaments.

A composition comprising an rAAV vector comprising at least an *SCN1A*-specific enhancer polynucleotide sequence as described herein is provided together with instructions for administering the composition to a subject having or at risk of developing a seizure, epilepsy, or DS. In an embodiment, the rAAV vector comprises

an *SCN1A* transgene for expression in interneuron cells including GABAergic interneurons and PV-expressing interneurons, or in pyramidal cells, such as glutamatergic pyramidal cells. The instructions will generally include information about the use of the composition for the treatment or prevention of the seizure, epilepsy, or DS. In other embodiments, the instructions include at least one of the following: description of the therapeutic agent (rAAV comprising *SCN1A*-specific enhancer polynucleotide sequence, etc.); dosage schedule and administration for treatment or prevention of ischemia or symptoms thereof; precautions; warnings; indications; counter-indications; overdose information; adverse reactions; animal pharmacology; clinical studies; and/or references. The instructions may be printed directly on the container (when present), or as a label applied to the container, or as a separate sheet, pamphlet, card, or folder supplied in or with the container.

Further embodiments and advantages thereof

Understanding and developing methods of treating neurological disorders stems from the complexity of the neuronal types involved. The products and methods of the embodiments described herein were developed to deconvolve the cellular actions of a disease gene, or a disease-associated gene. As such, the *SCN1A* locus was systematically dissected, thereby resulting in the identification of 10 different enhancer elements (enhancers E1-E10), in particular, human enhancer elements and the sequences thereof, that were found to be distributed across the intronic and intergenic region of the *SCN1A* gene (**FIG 3D**). By creating AAVs whose expression was dependent on each of these enhancers, at least three enhancers that recapitulated the global pattern of *SCN1A* gene expression were identified, e.g., E2 (for PV-specific expression), E6 (for VIP-specific expression) and E5 (for expression related to pyramidal layer 5. The other seven elements (e.g., E1, E3, E4, E7, E8, E9 and E10) were all highly specific for GAD1, labeled an assortment of interneuron subpopulations and may recruit distinct combinations of subtypes. In a particular embodiment, the E2 enhancer element was identified as being selective for a certain cIN subtype, namely, the PV-expressing fast spiking cells. As loss of expression of *SCN1A* is especially associated with PV cIN dysfunction, the E2 enhancer proved to be particularly adept at selectively targeting this cell population, not only in rodents but also within various primates, including humans. In addition, the E2 enhancer was

determined to be useful for investigating aspects of PV cIN function, including, without limitation, connectivity, monitoring excitability, and manipulating PV cIN activity with optogenetics. The demonstration of the utility of the E2 enhancer in a range of species highlights the breadth of basic and clinical applications that are provided by this approach. Other uses provided by the E2 enhancer include, by way of example, broader exploration of circuits (e.g. creating starter cells for monosynaptic tracing using recombinant-virus, such as rabies), cell type-specific gene loss of function (e.g CRISPR) and targeted drug screening. In addition, use of the E2 enhancer provides an agent for investigating species specific differences in the numbers, distribution or physiological properties of PV cINs. Generalized to other cell types, the approach is advantageous for investigating a range of species, most notably, both primates and humans.

As described herein, the strategy of systematically examining enhancers at a specific disease locus, such as the *SCN1A* gene locus, successfully identified key regulatory elements for each of the cell types that expresses this gene, thus, highlighting the benefits of the approach. It both clarifies the regulatory landscape controlling the expression of the *SCN1A* gene, as well as providing a tool kit for the manipulation of the distinct subpopulations of cells that express it.

Many of the SNPs associated with the *SCN1A* locus map to intron 1. In particular embodiments and as described herein, the three enhancers that were identified as having high specificity for *SCN1A*-expressing populations, namely, E2, E5 and E6, were located within this region. Without wishing to be limited by theory, the identified SNPs may represent mutations in these enhancers that affect the expression of *SCN1A*. It has been reported that GTEx data show multiple eQTLs within these enhancers that are associated with alterations in *SCN1A* expression in humans (Auget, F. et al., 2017, *Nature*, 550:204-213). E2 is especially noted, as conditional removal of *SCN1A* from forebrain interneurons has been shown to recapitulate the seizure phenotype in mice. As *SCN1A* expression is largely restricted to the PV-expressing subpopulation of interneurons, mutations in the E2 enhancer may be a direct cause of Dravet syndrome.

One of the great impediments to examining the early dynamics of circuit maturation has been the inaccessibility to specific cell types without the use of

transgenic animals. Young PV cINs have been particularly problematic to target even with complex genetic strategies. In view of the abundance of PV-cINs (they represent 40% of all inhibitory cINs), as well as their implication in neurodevelopmental disorders, accessing these cells prior to the onset of PV expression is a priority in the field. The specificity of the E2 enhancer at these developmental stages and the use of viral injection provide agents and tools to understand the normal development of neuronal cell types such as PV cINs and their role in neurological or neuropsychiatric diseases. In an embodiment, the E2 enhancer provides an agent for studying the normal development of PV-cINs and their role in disease. In addition, the E2 enhancer, as well as other enhancer elements provided herein, may serve to target specific cells and are advantageous for the treatment of diseases, e.g., neuronal diseases, including Dravet syndrome.

In other embodiments, the enhancers identified and described herein provide access to particular cell populations with distinct clinical relevance. By way of example, these enhancers be used to alleviate the debilitating aspects of Dravet syndrome, e.g., either through gene therapy or via modulation of neuronal activity, e.g., via optogenetic or chemogenetic approaches. (*See, e.g.,* Walker, M.C. et al., 2019, *Neuropharmacology*, 107751. doi: 10.1016/j.neuropharm.2019.107751. Review. PMID: 31494141). As described and demonstrated herein, local and systemic injections can be used for effective viral delivery to the brain, thus providing delivery and administration methods for clinical interventions. By way of example, local injections (e.g., of recombinant virus carrying an enhancer element and target polynucleotide) may be employed to alleviate focal epilepsy, prefrontal cortex dysfunction or hippocampal memory disorders. Systemic administration or delivery of virus may be employed in contexts where global interventions are necessary, for example, to correct generalized seizures or for psychiatric and neurodegenerative disorders. As provided by the embodiments described and exemplified herein, the rigorous identification of regulatory elements allows for accessing specific cell types. Such elements are advantageous for use in both experimental and therapeutic procedures and methods.

The practice of the described embodiments employs, unless otherwise indicated, conventional techniques of molecular biology (including recombinant techniques), microbiology, cell biology, biochemistry and immunology, which are well within the purview of the skilled artisan. Such techniques are explained fully in the literature, such as, “Molecular Cloning: A Laboratory Manual”, second edition (Sambrook, 1989); “Oligonucleotide Synthesis” (Gait, 1984); “Animal Cell Culture” (Freshney, 1987); “Methods in Enzymology” “Handbook of Experimental Immunology” (Weir, 1996); “Gene Transfer Vectors for Mammalian Cells” (Miller and Calos, 1987); “Current Protocols in Molecular Biology” (Ausubel, 1987); “PCR: The Polymerase Chain Reaction”, (Mullis, 1994); “Current Protocols in Immunology” (Coligan, 1991). These techniques are applicable to the production of the polynucleotides, viral vectors and viral particles and, as such, may be considered in making and practicing the embodiments described herein. Particularly useful techniques for particular embodiments will be discussed in the sections that follow.

The following examples are put forth so as to provide those of ordinary skill in the art with a complete disclosure and description of how to make and use the products, compositions and therapeutic methods as described herein, and are not intended to limit the scope of what is described and exemplified herein.

EXAMPLES

20 **EXAMPLE 1 – Identification of cis-regulatory sequences (PV-interneuron-specific enhancer sequences) that restrict expression of reporter and effector genes to PV-expressing cortical interneuron cell populations**

SCN1A, the gene that encodes the Nav1.1 sodium channel, is expressed in multiple distinct neuronal populations in the cortex. These include three, non-overlapping neuronal populations: fast-spiking cortical interneurons expressing parvalbumin (PV cINs), dis-inhibitory cortical interneurons expressing the vaso-intestinal peptide (VIP cINs) and layer 5 pyramidal neurons. In a particular embodiment, *SCN1A* is expressed in PV-expressing cortical interneurons. *SCN1A* is of particular interest, as its loss of function is associated with Dravet syndrome, an early-onset and intractable form of epileptic encephalopathy characterized by the early onset of seizures. More specifically, haploinsufficiency or pathogenic variants of *SCN1A* cause Dravet Syndrome.

An integrative method to systematically identify candidate enhancers within the *SCN1A* locus was developed and devised as a genetic strategy to target the distinct cortical populations expressing this gene. Regulatory sequences were selected based on the following three criteria. First, because it has been posited that the proximity of the enhancer to the transcriptional start site (TSS) of a gene scales directly to the level of expression, the intergenic and intronic regions of *SCN1A* closest to its TSS were examined to identify enhancers capable of driving functional levels of transgenes. Second, because the location of active enhancers within a given cell type correlates with chromatin accessibility, interneurons from the visual cortex were collected using *Dlx6a*^{cre};Sun1-eGFP transgenic mice to assess the chromatin landscape of the cellular populations expressing *SCN1A*. The location of active enhancers within a given cell type correlates with chromatin accessibility and DNA hypomethylation.

After the nuclei were isolated, single-cell ATAC-seq profiling (see, e.g., Buenrostro, J.D. et al., *Nature*, 523:486–90 (2015) and Cusanovich, D.A. et al., *Science*, 348: 910–4 (2015)) was performed using the SnapATAC analysis pipeline (described in the Methods *infra*) to identify differentially accessible chromatin regions in each of the four major classes of cortical interneurons, including PV cINs, VIP cINs and pyramidal neurons, which express the highest levels of *SCN1A*. (**FIGS. 3A-3C and FIG. 12**). Third, because regulatory elements are subject to positive selection pressure, sequences showing the highest conservation across mammalian species, including humans, were identified. Thus, to determine and isolate enhancers with therapeutic potential, ten selectively accessible intronic and intergenic regions near the TSS of *SCN1A* that are highly conserved through evolution were evaluated, e.g., E1-E10 as described herein. (**FIG. 3D and FIGS. 15A-1, 15A-2, 16A-1 and 16A-2**).

To examine the ability of the candidate enhancers to target the neuronal populations that express *SCN1A*, each enhancer sequence was inserted into an rAAV-backbone containing a minimal promoter upstream of a red fluorescent reporter (rAAV-E[x]-dTomato). From these constructs, rAAVs were then produced with the PHPeB capsid (Chan, K.Y. et al., *Nat. Neurosci.*, 20:1172–1179 (2017)) and were systemically injected into adult mice. After 3 weeks, all viruses showed strong and sparse expression within the cortex, as well as across multiple brain regions. Except for E5, the vast majority of virally-labelled cells expressed the pan-interneuron

marker Gad1. However, the degree of co-localization for PV within cortical neurons varied and ranged from over 90% for E2 to below 5% for E6, with all remaining enhancers displaying intermediate levels of PV specificity (**FIG. 3E and FIG. 7**). Thereafter, the identity and layer distribution of the neuronal populations captured by the E2, E5 and E6 enhancers were further examined. Consistent with their layer distribution, co-localization analysis with various markers revealed that the E2 regulatory element restricted the expression of the viral reporter to PV cINs, while E6 was selective for VIP interneurons. By contrast, the E5 regulatory element, while sparsely labeling interneurons across all layers, had a notable enrichment for pyramidal neurons in layer 5 (**FIG. 3F**). Thus, a significant fraction of the cortical expression profile of SCN1A was mirrored by the collective expression of three enhancers. These regulatory elements therefore account for largely non-overlapping expression in populations of interneurons and neurons with distinct functions and developmental origins. The viral tools developed as described herein provide a means for dissecting neuronal subtypes and can be advantageously used to study their normal function, as well as abnormalities in diseased cortex.

In a particular aspect, the S5E2 (E2) enhancer element sequence was incorporated into a recombinant AAV (rAAV) vector, which comprised a minimal basal promoter and a reporter transgene (e.g., d-Tomato) or an effector gene (e.g. Gq-DREADD), to generate the rAAV vector called pAAV-S5-E2-dTomato. The ability of the E2 enhancer to restrict expression of the reporter gene (transgene) to PV-expressing interneurons in brain was assessed by injecting the E2 enhancer-containing rAAV vector systemically into animals (mice) and analyzing the co-localization between the expressed reporter across brain structures including the cortex. An image showing the results of immunohistochemical (IHC) staining analysis for the dTomato reporter in brain sections is shown in **FIG. 2A** (sagittal sections in the top portion of the figure; coronal sections in the lower portions of the figure) following systemic *in vivo* injection of the pAAV-S5-E2-dTomato vector into an animal (mouse), allowing for detection of specific cells transduced by the vector. Images showing the results of immunohistochemical (IHC) staining analysis for the dTomato reporter expressed in brain sections following systemic *in vivo* injection of the pAAV-S5-E2-dTomato vector, or into an animal (mouse), allowing for detection of specific cells expressing

PV are shown in **FIG. 2B**. Reporter gene expression from the pAAV-S5-E2-dTomato vector is visualized in brain sections (**FIG. 2B**, left panel, red). Reporter gene expression from the pAAV-S5-E2-Gq-DREADD-dTomato is visualized for Gq-DREADD (green) and for dTomato (red) (**FIG. 2B**, right panel). Detection of specific PV-expressing cells transduced by the vector is also visualized (**FIG. 2B**, left panel, green; **FIG. 2B**, right panel, blue).

Identification of candidate enhancers

Using the above enhancer sequence selection approach, ten candidate enhancer sequences proximal to the *SCN1A* gene transcriptional start site were discovered in the mouse genome. These enhancer sequences, called S5E1 (E1), S5E2 (E2), S5E3 (E3), S5E4 (E4), S5E5 (E5), S5E6 (E6), S5E7 (E7), S5E8 (E8), S5E9 (E9) and S5E10 (E10) herein, were identified in the vicinity of the *SCN1A* gene (**FIG. 1A-1**). The human polynucleotide sequences corresponding to the E1-E10 enhancer sequences (SEQ ID NOs: 15-24) are also provided and described herein, as well as additional human enhancer polynucleotide sequences E11-E35 (SEQ ID NOs: 25-49) as described herein. (**FIGS. 1A-2 and 1A-3**).

The human (human ortholog) sequences for the E1-E10 enhancers were determined based on alignment of the mouse sequences to the human genomic sequence of *SCN1A*, including 100 kb both upstream and downstream, leading to the identification of human ortholog sequences that are highly conserved between the two species (**FIGS. 1A-1 to 1A-3, 16A-1, 16A-2**). As will be appreciated by the skilled practitioner, enhancer regulatory elements comprise a series of transcription binding sites that are relatively well conserved across species, but are interspersed with spacer sequences that are not contiguously conserved across species. Accordingly, in an embodiment, a *SCN1A* enhancer element can constitute a nucleotide sequence containing any regions of more than 100 bp that have at least 75% or greater sequence identity with a human polynucleotide (DNA) enhancer sequence as described herein, namely, E1-E10. In an embodiment, a *SCN1A* enhancer element constitutes a nucleotide sequence containing any regions of more than 100 bp that has at least 75% or greater sequence identity with the human E2 (S5E2) polynucleotide (DNA) enhancer sequence. For such enhancer sequences, the size of the nucleic acid sequence is not limiting, so long as the sequence contains any regions of more than

100 bp that have at least 75% or greater sequence identity with a human polynucleotide (DNA) E1-E10 or E11-E35 enhancer sequence as described herein. The data related to each of the identified enhancer sequences (35 enhancer sequences) described herein is provided in the tables shown in **FIGS. 1A-1 to 1A-3, 15A-1, 15A-2, 16A-1 and 16A-2.**

E1-E10 enhancer element-restricted reporter gene expression in PV-expressing interneurons in cortical layers of the mouse brain is shown in **FIGS. 1B-1 and 1B-2**, which present immunohistochemical (IHC) staining analysis for dTomato in brain sections following systemic *in vivo* injection of the pAAV-S5-E2-dTomato vector into an animal (mouse). Quantification of the degree of the specificity (**FIG. 1C**) and sensitivity (**FIG. 1D**) of expression of the reporter gene in PV-expressing interneurons in the cortex is demonstrated graphically. The expression of the reporter gene is controlled by the E1-E10 enhancer elements contained in rAAV vectors. The specificity was quantified as the proportion of cells expressing the viral reporter dTomato co-expressing the PV-interneuron marker PV assessed by immunohistochemistry on brain sections following systemic *in vivo* injection of the pAAV-S5-E2-dTomato vector into an animal (mouse). The sensitivity was quantified as the proportion of cells expressing the PV-interneuron marker PV that co-expressed in the viral reporter dTomato as assessed by immunohistochemistry on brain sections following systemic *in vivo* injection of the pAAV-S5-E2-dTomato vector into an animal (mouse). Bar graphs represent mean +/- standard error of the mean.

EXAMPLE 2 –Viral targeting of PV cortical interneurons (PV cINs) in mice

The 90% specificity of the E2 regulatory element for PV cINs provides a means for targeting fast-spiking neurons (e.g., basket and chandelier cells), which collectively constitute 40% of all cortical (GABAergic) interneurons. These neurons exert a strong level of inhibition over local networks, and their dysfunction has been directly implicated in neurological and neuropsychiatric disorders including Dravet syndrome, focal epilepsy, Autism Spectrum Disorder (ASD) and schizophrenia. As such, gaining control over their activity is of particular interest for both fundamental research and clinical applications. Thus, the E2 regulatory element was investigated and characterized in order to develop an agent having broad utility, e.g., as a viral tool or a therapeutic agent.

Adult mice systemically-injected with rAAV-E2-dTomato showed detectable expression of the viral reporter after one week and reached a high and stable level of expression after 3 weeks. Immunohistochemistry and *in situ* hybridization were performed and consistently showed that ~90% of virally labeled cells were PV INs in the cortex (i.e., PV-expressing cortical interneurons). Conversely, on average, 75% of PV cINs expressed the viral reporter, with a maximum sensitivity reaching 93% (FIGS. 4A and 4B). This is indicative of the capability of E2 to target all PV cINs without bias for layers or subtypes. Consistent with the specificity for PV cINs, slice recordings from mice showed that the neurons expressing the viral reporter exhibited electrophysiological properties characteristic of fast-spiking PV cINs both within the primary somatosensory cortex (S1) and the pre-frontal cortex (PFC), (FIG. 4C and FIGS. 8A and 8B).

Although the viral reporter was predominantly confined to the brain cortex, some positive cells were observed in other brain regions closely corresponding to areas of *SCN1A* expression. E2 maintained high specificity for PV-expressing neurons within the primary visual cortex (V1) and cingulate cortex, subiculum, hippocampal CA1, substantia nigra pars reticulata (FIG. 8C). Of note, virtually no viral reporter expression was observed outside of the brain, with the exception of a few cells observed in the liver (which is expected upon systemic delivery of any AAVs) and in the lungs (where *SCN1A* is expressed at a low level (FIG. 8D). These results show that, despite systemic delivery, the vector containing E2 can be used to selectively target PV-expressing neurons in various brain regions, with insignificant off-target expression outside of the central nervous system.

Many experimental paradigms and clinical applications may require local rather than systemic injection. To be useful in these contexts, viral expression must retain a high level of specificity for PV cINs. Stereotactically guided injections typically lead to a higher number of viral particles per cell compared to systemic delivery, which may result in off-target expression. To test whether increasing the viral load altered the specificity, the same volume of rAAV-E2-dTomato was locally injected at various titers into the cortex of adult mice and reporter expression was assessed within PV cINs after one week (FIG. 4D). The results showed that while

higher titers had increased levels of reporter expression, no significant alteration of specificity was observed.

Despite the prevalence of PV-expressing interneurons in the mature cortex, targeting these PV cINs at early postnatal stages has been hampered by the relatively late expression of parvalbumin (approximately 15 days after birth, i.e., P15) and the lack of other early markers for this population. The involvement of PV cINs in developmental disorders highlights the need to target and manipulate this cell population during cortical circuit assembly. Complex genetic strategies offer only a partial solution to achieving this in mice (i.e., Lhx6-Cre, Sst-Flp and Cre and Flp-dependent reporter); however, these strategies do not offer the means to easily manipulate these neurons prior to the second postnatal week.

To test whether the E2 enhancer targeted fast-spiking cINs before the onset of expression of parvalbumin, its activity was examined at various postnatal stages. To this end, the analysis was tiled across the early postnatal period, through a series of stereotactically-guided injections of rAAV-E2-dTomato (**FIG. 4E**). The selectivity of the reporter was assessed upon the onset of parvalbumin expression at P15. This assessment revealed that greater than 50% selectivity was obtained for PV cINs upon injection at P1, increasing to 67% by P7 injection, and to over 80% after P10 injection. This approach was further used to label PV cINs prior to P15. To identify fast spiking cINs in this context, Lhx6-Cre/Intact transgenic mice, in which GFP is expressed in medial ganglionic eminence (MGE)-derived interneurons (both PV cINs and SST cINs), were used. By co-staining for SST, the PV cINs could be distinguished as GFP-positive/SST-negative. 72% and 78% specificity was obtained for PV with a P4-P7 or a P7-P10 time course, respectively. Thus, this approach provides a means to study such neurons during circuit maturation using a single viral injection.

EXAMPLE 3 – Viral monitoring and manipulation of PV cortical interneurons in mice

Having demonstrated the fidelity of E2 expression for PV cINs with differing modes of injection and across developmental stages as described in Example 2, the utility of this vector was assessed for studying connectivity (using a presynaptic reporter) and activity (using imaging, coupled with a genetically encoded calcium-

reporter). When E2 was used to drive a synaptophysin-tdTomato fusion gene (*see, e.g.,* Madisen, L. et al., 2012, *Nat Neurosci*, 15(5):793-802), reporter expression was restricted pre-synaptically to PV cINs, with terminals peri-somatically located onto pyramidal neurons (**FIG. 5A**). When this vector was used to drive GCaMP6f expression (Chen, T.W. et al., *Nature*, 499: 295–300 (2013)), it was demonstrated that PV cINs were recruited upon whisker stimulation (**FIGS. 5B and FIG. 9A**). Together these results demonstrated that E2 provides an effective means to monitor various aspects of PV cIN biology.

Further studies were conducted to examine whether E2 was sufficient to elicit functional changes in activity using chemo- or optogenetic approaches. E2 was used to direct the expression of the chemogenetic receptor PSAM4-5HT3-LC (Magnus, C.J. et al., 2019, *Science*, 364(6436) in adult animals. It was observed that PV cINs in brain sections collected from these animals, when exposed to the actuator varenicline, could be induced to fire when current clamped below threshold (**FIG. 5C**). Similar results were obtained using the chemogenetic receptor Gq-DREADD (Armbruster, B.N. et al., *PNAS USA*, 104:5163–5168 (2007), (**FIG. 9C**). Finally, both constant and high frequency laser stimulation of PV cINs expressing the red-shifted opsin C1V1 in brain slices resulted in firing time-locked to the stimulus (**FIGS. 5D and 5E and FIGS. 9A and 9B**). Demonstrating that engagement of these neurons resulted in concomitant local inhibition, pyramidal neuron activity in the vicinity of virally labeled PV cINs was consistently interrupted by laser stimulation. Notably, this effect was abolished by treatment with picrotoxin (**FIG. 5D and FIG. 9B**). Having demonstrated the efficacy of the method *ex vivo*, the ability to alter excitatory networks *in vivo* by opto-genetically stimulating PV cINs was examined. Three weeks following local injection of AAV-E2-C1V1 into the primary visual cortex of adult animals, single unit recordings within the infected region were performed both at baseline and upon laser stimulation. The identity of recorded neurons was distinguished based upon their spike width and maximal firing frequency. Reliably, inhibitory interneuron firing rates were increased by laser stimulation, while excitatory neuronal firing was silenced (**FIG. 5E**). Together, these results demonstrated that E2 can functionally engage PV cINs and elicit network inhibition using chemo- or optogenetics approaches both *ex vivo* and *in vivo*.

EXAMPLE 4 – Viral monitoring and manipulation of PV cortical interneurons in primates, including humans

The sequence of the E2 enhancer is highly conserved across mammalian species, including humans, thus suggesting a conserved role in gene regulation.

5 Studies were conducted to establish whether the E2 regulatory element could be used to target PV cINs across mammalian species. Using systemic injections (in marmoset) or focal injections (in rat and macaque) of virus vector harboring E2 (E2 virus), it was demonstrated that PV cINs were targeted with approximately 90% specificity (**FIG. 6A**). It has been reported that human brain tissue obtained during
10 surgical resection can be cultured for prolonged periods (Eugène, E. et al., 2014, *J. Neurosci Methods*, 235:234-244). Taking advantage of the resilience of human brain to remain healthy *ex vivo*, freshly resected subiculum or medial temporal cortex were exposed to E2 virus. Over the two-week culture period, the progressive appearance of fluorescently labeled cells was observed. In regions where PV staining reflected the
15 expected distribution of these cells, virally labeled cells were PV-positive (**FIG. 6B** (i); see methods for details). In addition, the majority of cells within both the cortex and the subiculum showed the characteristic hallmarks of PV INs as indicated by multiple criteria, including morphology, maximal firing rate when evoked through direct depolarization, or optogenetic light stimulation (**FIG. 6B** (ii-iv) and **FIGS. 10A**
20 **and 10B**).

Notably, the human E2 enhancer showed the same degree of specificity for PV cINs upon injection in mice, further demonstrating that non-coding regions of the genome characterized by a high degree of sequence conservation are likely to retain their functional properties across species. Finally, truncation of both the 5' and 3'
25 ends of the human E2 enhancer resulted in a drastic reduction of specificity, suggesting that the functional boundaries of the E2 enhancer have been optimally identified (**FIG. 14**). Together, these results indicate that the E2 vector provides an effective tool for targeting and manipulating PV cINs across mammals, including humans.

EXAMPLE 5 – Identification of viral enhancers with regional specificity

To demonstrate that the enhancer selection method described herein was generalizable, 25 additional enhancer/regulatory element candidates (E11-E35 herein

were identified in the vicinity of seven genes whose expression was enriched in PV cINs across species (**FIGS. 1A-1 to 1A-3; FIGS. 15A-1 and 15A-2; FIGS. 16A-1 and 16A-2**). (see, Methods, *infra*). Systemic injection of AAVs containing these sequences revealed that four of them displayed greater than 90% selectivity for PV cINs. Notably, among these enhancers, the relatively few virally-labeled neurons that did not express PV were positive for the pan-interneuron marker Gad1. In **FIGS. 1A-1 to 1A-3**, *Pvalb* (UniProtKB – P20472) refers to the gene that encodes calcium-binding parvalbumin alpha protein; *ACAN* (NCBI Gene ID: 176; UniProt P16112) refers to the gene that encodes the aggrecan core protein (also called cartilage-specific proteoglycan core protein), which may be involved in the disease spondyloepimetaphyseal dysplasia; *Tmem132c* (NCBI Gene ID: 92293) refers to the gene that encodes the transmembrane protein 132c, a type of protein that spans a biological membrane of a cell or organelle; *Lrrc38* (UniProtKB – Q5VT99) refers to the leucine rich repeat containing 38 gene, which shows relatively high expression in adrenal and prostate tissues; *Inpp5j* (UniProtKB – Q15735) refers to the gene that encodes phosphatidylinositol 4,5-bisphosphate 5-phosphatase A, which may be involved in the modulation of the function of inositol and phosphatidylinositol phosphate binding proteins in membrane ruffles; *Mef2c* (UniProtKB – Q06413) refers to the gene that encodes myocyte-specific enhancer factor 2C, a transcription factor in the Mef2 family, involved in cardiac morphogenesis and myogenesis and vascular development, as well as in neurogenesis and in the development of cortical architecture. In humans, mutations in the *Mef2c* gene result in autosomal dominant mental retardation 20 (MRD20), which is characterised by severe psychomotor impairment, periodic tremor as well as abnormal EEG and epilepsy. *Pth1h* (NCBI Gene ID: 5744) refers to the gene that encodes parathyroid hormone-like peptide, which is secreted by cancer cells, e.g., breast, lung, ovarian, pancreatic, prostate, liver, or colorectal cancer cells, causing humoral hypercalcemia of malignancy by activating the type 1 PTH/PTHrP receptor in kidney and bone. Similar to *SCN1A*, the above-noted genes are highly enriched in PV-interneurons compared with all other cells in the brain. As such, these genes were selected as candidates for targeting by enhancer elements, and the enhancers as described were identified and located in the vicinity of the coding sequences of these genes.

In particular, four PV-specific regulatory elements, namely, E11 (SEQ ID NO: 25, human), E14 (SEQ ID NO: 28, human), E22 (SEQ ID NO: 36, human) and E29 (SEQ ID NO: 43, human) were identified as having highly selective expression within specific brain regions. (FIGS. 13A and 13B). Each of these four enhancers was specific for distinct but overlapping subsets of the PV-expressing neurons. Specifically, while E11 and E14 showed a bias for targeting PV cINs in the upper layers of the cortex, the E22 enhancer showed restricted expression almost exclusively to the cortex, with only a few neurons showing low levels of expression elsewhere. By contrast, the E29 enhancer showed the most global expression, as it targeted the entire population of PV-expressing neurons throughout the central nervous system. All of these enhancers exhibit a high degree of sequence conservation and were selected from genes whose expression profile is similar across species. To directly test that the cross-species similarity among the enhancers results in similar functionality across species, AAV-E22-dTomato was locally injected in V1 of a macaque. This showed that, in a manner similar to that of mouse, the expression of the viral reporter was restricted to PV cINs. The combination of regional selectivity and conservation of expression across species provides a utility for these viral agents in targeted therapies to correct abnormal brain function in different mammalian species.

EXAMPLE 6 –*SCN1A* expression is restored to normal levels by delivering a functional copy of the *SCN1A* gene within the *SCN1A*-expressing population in a mouse model of DS.

To restore *SCN1A* gene expression to normal levels by delivering a functional copy of the *SCN1A* gene to an *SCN1A*-expressing interneuron cell population, e.g., in a mouse model of DS, the ‘limited nucleic acid (DNA) payload’ (i.e., the size of exogenous nucleic acid (DNA), e.g., a transgene and associated nucleic acid sequences, that is contained or carried within the rAAV vector) is increased using one or more approaches that result in an rAAV vector that can accommodate the size of the *SCN1A* gene. As noted *supra*, the AAV DNA is on the order of 4.7-5 kb, while genes desired for insertion within an rAAV vector and delivery by the vector are often twice that size or larger. The delivery of larger genes using rAAVs has been demonstrated in other contexts using multiple vectors that re-assemble by homologous recombination or by splicing mediated by acceptor-sites. (See, e.g.,

Hirsch, M.L. et al., 2016, *Methods Mol Biol*, 1382:21-39). Both of these approaches are available to overcome the packaging limits of rAAV.

As both DS animal models and human patients have demonstrated, the requirement for *SCN1A* is dose-dependent. Therefore, the level of expression of rAAV-driven *SCN1A* is appropriately titrated as known and practiced in the art to match, or to match as closely as possible, the normal endogenous level of *SCN1A* expression. Several methods can be used to precisely modulate the levels of *SCN1A* gene expression. Various strategies are used to modulate the levels of *SCN1A* expression, using amelioration of seizures as a direct readout of the effectiveness of the treatment.

EXAMPLE 7 – A pharmacogenetic approach to selectively normalize the excitability of an *SCN1A* -deficient neuronal population in a mouse model of DS

As an alternative to direct gene therapy using rAAV vectors harboring specific enhancer and gene nucleic acid sequences for delivery and restricted expression in interneuron cells, pharmacogenetic methods may be employed to directly correct neuronal activity within the *SCN1A* neuronal populations. To this end, a chemogenetic approach involving ‘designer receptors’ may be used to modulate interneuron activity. Designer receptors exclusively activated by designer drugs (DREADDs) are modified human muscarinic receptors. In addition, PSAM-PSEM chemogenetic agents are suitable for use.

Using Gq-DREADD, a receptor exclusively activated by clozapine-N4-oxide, (CNO), a pharmacologically inert and orally bioavailable drug, excitability/inhibitory balance (E/I balance) may be corrected in a mouse model of DS (DS mice). In brief, the Gq-DREADD receptor is expressed in *SCN1A*-deficient interneuron cells using an rAAV vector harboring an *SCN1A*-specific enhancer, e.g., E1-E10, as described *supra* and the *SCN1A* gene. Based on other studies using Gq-DREADD, the receptor is expected to be functional and located at the membrane of the transduced/infected cells. In addition, the rAAV vector containing a *SCN1A*-specific regulatory element, e.g., E1-E10 as described herein, should drive the expression of the G1-DREADD receptor exclusively within interneurons, such as GABAergic interneurons and PV-expressing, GABAergic interneurons. The functionality of the Gq-DREADD within infected cells may be assessed. Upon bath application of CNO, all interneurons

expressing Gq-DREADD are expected to show membrane potential depolarization within less than a minute, consistent with the expression of a functional receptor). Moreover, voltage clamp recordings of a pyramidal cell in the vicinity of interneurons that express Gq-DREADD are expected to show an increase in inhibitory post-synaptic currents (IPSCs) upon application of clozapine-*N*-oxide (CNO). Such experiments demonstrate that rAAV comprising an SCN1a-specific E1-E10 enhancer sequence and a Gq-DREADD-encoding polynucleotide allows specific, functional and restricted expression of Gq-DREADD and that CNO-treatment effectively and selectively increases the activity of interneurons, thereby providing a localized and marked increase in inhibitory activity by the interneurons within neighboring excitatory neurons.

If the lack of *SCN1A* (loss of function of *SCN1A*) should impair the ability of the DREADD to increase the cells' excitability, DREADD can be delivered to all interneurons by using a pan-interneuron enhancer, such as, for example, the distinct and different *Dlx* enhancer, as described by Dimidschstein, J. et al. (2016, *Nature Neuroscience*, 19(12):1743-1749) to circumvent the impairment by increasing the activity of other types of interneurons that are not affected by the loss of function of *SCN1A*.

EXAMPLE 8 – Materials and Methods of the above-described Examples

scATAC-seq library preparation and sequencing. Male hemizygous *Dlx6a*-Cre mice (Jax stock #008199) were crossed with female homozygous INTACT mice (flox-Sun1-eGFP, Jax stock #021039) to yield *Dlx6a*-Cre::INTACT offspring for scATAC-seq experiments. Brains from P28 *Dlx6a*Cre::INTACT mice were harvested, sectioned coronally on a mouse brain slicer (Zivic Instruments), and regions of interest were dissected in ice-cold artificial cerebrospinal fluid (ACSF). Tissue was then transferred to a dounce homogenizer containing Lysis Buffer (10 mM Tris-HCl, 10 mM NaCl, 3 mM MgCl₂, 0.01% Tween-20, and 0.01% IGEPAL CA-630, 0.001% Digitonin). Tissue was homogenized with 10 strokes of pestle A, 10 strokes of pestle B, and incubated for 5 minutes on ice before being filtered through a 30 µm filter and centrifuged at 500xg for 10 minutes at 4°C. The pellet was resuspended in 1% BSA for sorting for GFP⁺ nuclei on a Sony SH800S cell sorter. Nuclei were sorted into Diluted Nuclei Buffer (10X Genomics). Single-cell ATAC-seq libraries were

prepared using the Chromium Single Cell ATAC Solution (10X Genomics). Libraries were sequenced using a Nova-Seq S2 100 cycle kit (Illumina). (FIGS. 3A-3C)

scATAC analysis. Raw sequencing data were passed through the Cell Ranger ATAC pipeline (10X Genomics). The fragments files were then used to generate snap files for analysis using the snapATAC package (<https://doi.org/10.1101/615179>). Cells were clustered using graph-based clustering (k=15, 24 principle components). Gene activity scores were generated as described in the snapATAC package and used to determine clusters corresponding to interneuron cardinal classes. For each cardinal class, bigwig files were generated and peaks were called using macs2 for input into the Integrated Genome Browser and enhancer selection. Peaks across cardinal classes were compared using bedtools Jaccard.

Enhancer selection. All enhancers presented herein (S5E1-E10 and E11-E35) were selected based on the co-presence of ATACseq data (for DNA accessibility) and conservation across species (using UCSC genome browser vertebrate conservation track). The genomic coordinates for mice and their human orthologs are presented in FIGS. 1A-1 to 1A-3.

For **Selection**, candidate regulatory elements were manually curated from a list of elements generated by intersecting the “context” region (*SCN1A* intergenic region + intron1) with both the “ATACseq peak union” file and the “Phastcons 60-way” file – see below. **Accessibility.** ATAC-seq data (Mo et al., 2015, *Neuron*, 86:1369-1384) were downloaded on the GEO repository and discretized as peaks using MACS2 ran with default parameters (<https://github.com/taoliu/MACS>). Using a custom R script, a file containing the union of all peaks across datasets was generated and used for enhancer selection as described below. The final selection relied on the inspection of the peaks for individual cell-types rather than on the union of all peaks. **Methylation.** Mouse mCH levels for non-overlapping 100kb bins across the entire genome for mouse (Luo et al., 2018, *Nat Commun*, 9(1):3824) were downloaded from Brainome portal (<http://brainome.org>). These data were used as a confirmation for the positioning of the candidates selected using the ATAC-seq dataset described above. **Conservation.** The “phascons 60-way” track was downloaded from the UCSC portal (<https://genome.ucsc.edu>) in BED file format and

filtered using a custom R script to remove any element smaller than 10bp and fuse any element separated by less than 50bp using Bedtools / Intersect.

rAAV cloning and viral production. All viral constructs were generated using standard cloning methods and protocols in molecular biology. The plasmid pAAV-mDlx-GFP (Addgene#83900; Addgene, Watertown, MA), (Dimidschstein, J. et al., 2016, *Nat. Neuroscience*, 19(12):1743-1749) was used to create a standard backbone containing the elements necessary for the production of AAVs (internal terminal repeats, minimal promoter, woodchuck posttranscriptional response element).

The enhancer sequences (necessary for restricting expression to specific types of neurons) were synthesized *de novo* by Genewiz (Cambridge, MA) and the reporters and effectors were amplified by PCR. In particular, the enhancer sequences were amplified by PCR from mouse genomic DNA using the following primers: E1: caaagtggacagagggagg (SEQ ID NO: 50) and gtgctgtgggagtggtgga (1280 bp), (SEQ ID NO: 51); E2: aatctaactggctgtata (SEQ ID NO: 52) and caattgctcagagtatttt (618 bp), (SEQ ID NO: 53); E3: ataaaatttttttcttaa (SEQ ID NO: 54) and gaggaatcagctacggggc (832 bp), (SEQ ID NO: 55); E4: tctgacagagcaagtcttga (SEQ ID NO: 56) and tatcaaaattgtatattcag (261 bp), (SEQ ID NO: 57); E5: aatgtttgatatttaggag (SEQ ID NO: 58) and ttgactcttaaaatttaata (663 bp), (SEQ ID NO: 59); E6: ttgtcactttgttactctac (SEQ ID NO: 60) and ttaaatcttaaaatttctt (606 bp), (SEQ ID NO: 61); E7: gatactgtataattaattag (SEQ ID NO: 62) and ctctctctgtgttcctttt (2430 bp), (SEQ ID NO: 63); E8: attgatctccaacttttaa (SEQ ID NO: 64) and gttcatccaagtaataagag (1644 bp), (SEQ ID NO: 65); E9: atctcaagtgtatgaacat (SEQ ID NO: 66) and gtctttttgttttttttt (521 bp), (SEQ ID NO: 67); E10: tattgcaaaaggaaggaatg (SEQ ID NO: 68) and tcatggaaaaagaaaaaatc (547 bp), (SEQ ID NO: 69). The enhancers, reporters and effectors were cloned using the Gibson Cloning Assembly Kit (NEB-E5510S) following standard procedures. Specifically, for AAV-E1:10-dTomato, the dTomato coding sequence was amplified from the plasmid Addgene # 83897; for AAV-E2-SYP-dTomato, the Synaptophysin-tdTomato coding sequence was amplified from the plasmid Addgene # 34881; for AAV-E2-GCaMP6f, the GCaMP6f coding sequence was amplified from the plasmid Addgene # 83899; for AAV-E2-C1V1-eYFP, the C1V1-eYFP coding sequence was amplified from the plasmid Addgene # 35499.

Final plasmids were assembled using the Gibson Assembly® Cloning Kit (NEB-E5510S), (New England BioLabs, Ipswich, MA), following the manufacturer's instructions and standard protocol. The rAAVs were produced using standard production methods. Polyethylenimine (PEI) was used for transfection (*see, e.g.,* Longo, P.A. et al., 2013, *Methods Enzymol.*, 529:227-240) and OptiPrep™ density gradient (Sigma-Aldrich, St. Louis, MO) was used for viral particle purification and isolation. Serotype 1 was used to produce the AAVs for local injections in mice and rats. Serotype 9 was used for systemic injection in marmosets and serotype PHPeB was used for both local injection in macaques and systemic injections in mice. Viral titer was estimated by qPCR with primers annealing via the WPRE sequence that is common to all constructs. All batches produced were in the range of 10^{10} to 10^{12} viral genomes per ml. In particular, Woodchuck Hepatitis Virus (WHP) Posttranscriptional Regulatory Element (WPRE) is a DNA sequence, which, when transcribed, creates a tertiary structure enhancing expression. WPRE, a tripartite regulatory element with gamma, alpha, and beta components, is commonly used in molecular biology to increase expression of genes delivered by viral vectors, e.g., rAAV-dTomato. (*see, e.g.,* Choi, J.-H. et al., 2014, *Mol. Brain*, 7:17). All rAAV batches produced were in the range of 10^{10} to 10^{12} viral genomes per ml.

Animals. Mice: Female C57BL/6J mice (*Mus musculus*; 10 weeks old) were obtained from Jackson Labs (Bar Harbor, ME - stock# 000664). **Rats.** Sprague Dawley rats (adult 150-- 250 gm) were obtained from Charles River labs, Kingston, NY. **Marmosets.** One female common marmoset (*Callithrix jacchus*, 6.0 years old) was obtained from the colony at Massachusetts Institute of Technology. **Macaques.** One male macaque (*Macaca mulatta*, 15.0 years old) was obtained from the California National Primate Research Center at the University of California, Davis. All animals were maintained in a 12 light/12 dark cycle with a maximum of five animals per cage for mice and one animal per cage for rats. Marmosets and macaques were socially housed. All animal maintenance and experimental procedures were performed according to the guidelines established by the Institutional Animal Care and Use Committee at the Broad Institute of MIT and Harvard (mice), McGovern research institute at MIT (rats and marmosets) and Salk Institute for Biological studies (Macaques) and adhered to the standards of the National Institutes of Health.

Local and systemic viral injections. Mouse local S1. Local injection in adult mice were performed by stereotactically guided injections in the somatosensory cortex with the following coordinates: 1.0 mm posterior, 2.9 mm lateral, 0.7/0.45 mm ventral relative to Bregma with 150 nL of virus. **Mouse systemic.** For systemic injection in
5 adult mice, approximately 10^{11} viral particles were injected in the retro-orbital sinus per animal. Post-operative monitoring was performed for five days post injection.

Rat local in V1. Local injection in adult rats was performed by stereotactically guided injections in the primary visual cortex with the following coordinates: 5.4 mm posterior, 4.2 mm lateral, 2.0 mm ventral relative to bregma with 670 nL of virus.

10 **Marmoset systemic injection.** For systemic injection in adult marmosets, approximately 10^{12} viral particles in ~ 0.7 ml of sterile PBS were injected into the saphenous vein, followed by another infusion with ~ 0.5 ml of saline. After the final infusion, pressure was applied to the injection site to ensure hemostasis. The animal was returned to its home cage and monitored closely for normal behavior post

15 anesthesia. The animal was euthanized 51 days after viral injection. **Macaque local in V1.** Local injection in an adult macaque was performed by a stereotactically guided injection in the left primary visual cortex with the following coordinates: 13 mm posterior, 19 mm lateral, 23 mm superior relative to the center of the inter-aural line (based on the animal's MRI). A total of volume of 333 nL was injected at 4
20 depths (i.e., 1.8, 1.3, 0.8 and 0.3 mm from the cortical surface).

Surgery. For stereotactically guided viral injection, animals were anesthetized under isoflurane (1–3% in oxygen) and placed in a stereotactic head frame on a temperature-controlled heating pad. A craniotomy and a durotomy were performed above the brain region of interest. The animals were injected with 50–500 nl of the indicated
25 virus (rAAV) at a rate of 10–25 nl/minute using a sharp glass pipette (25–35 mm in diameter), which was left in place for 5–15 minutes after the injection to minimize backflow. The craniotomy site was covered with sterile bone wax, the surgical opening was closed with Vetbond, and the animals were returned to their home cages for at least 1 week. The injection sites were defined by the following coordinates:
30 somatosensory cortex S1: 1.0 mm posterior, 3.0 mm lateral, 0.7 / 0.4 mm ventral relative to bregma; hippocampus CA1: 1.6 mm posterior, 1.8 mm lateral, 1.2 mm

ventral relative to bregma; striatum: 0.5 mm posterior, 2.0 mm lateral, 3.2 mm ventral relative to bregma.

For retro-orbital vein injection, animals were anesthetized under isoflurane (1–3% in oxygen) and placed on a temperature-controlled heating pad. Intravenous (IV) injections were performed in the retro-orbital plexus. More specifically, the animal (mouse) was placed in a funnel-shaped nose cone connected to a non-rebreathing apparatus (Surgivet, Dublin, OH) and the needle was inserted, bevel down, at the medial canthus, into the retroorbital sinus. Up to 150 μ L of supernatant containing replication-defective rAAV vectors were injected into the tail vein or retro-orbital plexus. Following injection, the eye was held shut for a minimum of 30 seconds to ensure homeostasis.

Electrophysiological recordings in mice:

Slice preparation for 2 to 6-weeks-old mice. Virally injected mice were anesthetized with isoflurane. Upon loss of reflexes, mice were transcardially perfused with ice-cold oxygenated ACSF containing the following (in mM): 87 NaCl, 75 sucrose, 2.5 KCl, 1.25 NaH_2PO_4 , 26 NaHCO_3 , 10 glucose, 1 CaCl_2 and 2 MgCl_2 . Mice were then decapitated and 300- μ m thick coronal slices were sectioned using a Leica VT-1200-S vibratome and incubated in a holding chamber at 32–35 $^\circ\text{C}$ for 15–30 min followed by continued incubation at room temperature 20–23.5 $^\circ\text{C}$ (68–74 $^\circ\text{F}$) for at least 45–60 minutes before physiological recordings. Slices containing the injection site were transferred in a recording chamber submerged with oxygenated ACSF containing the following (in mM): 125 NaCl, 2.5 KCl, 1.25 NaH_2PO_4 , 26 NaHCO_3 , 10 glucose, 2 CaCl_2 and 1 MgCl_2 (pH = 7.4, bubbled with 95% O_2 and 5% CO_2).

Slice preparation for 6-weeks-old and older mice. Acute coronal brain slices were prepared as follows: Mice were anesthetized with Avertin solution (20 mg/ml, 0.5 mg/g body weight) and transcardially perfused with 15 to 20 ml of ice-cold carbogenated (95% O_2 , 5% CO_2) cutting solution containing the following: 194 mM sucrose, 30 mM NaCl, 4.5 mM KCl, 1.2 mM NaH_2PO_4 , 0.2 mM CaCl_2 , 2 mM MgCl_2 , 26 mM NaHCO_3 , and 10 mM D-(+)-glucose (with osmolarity of 340–350 mOsm). The brains were then rapidly removed and placed in ice-cold cutting solution for slice preparation. Coronal slices (300 μ m) were prepared and then incubated at 32 $^\circ\text{C}$ with carbogenated artificial cerebral spinal fluid (aCSF) for 10 to

15 minutes. The slices were then incubated at room temperature for at least 1 hour in a CSF that contained the following: 119 mM NaCl, 2.3 mM KCl, 1.0 mM NaH₂PO₄, 26 mM NaHCO₃, 11 mM glucose, 1.3 mM MgSO₄, and 2.5 mM CaCl₂ (pH 7.4, with osmolality of 295–305 mOsm) at room temperature for at least 1 hour. **Current**

5 **clamp.** For interneuron recording, 10 μ M CNQX, 25 μ M AP-5 and 10 μ M SR-95531 were also added to block AMPA, NMDA and GABA_A receptors, respectively, to measure the cell-intrinsic effect of optogenetic and chemogenetic stimulation.

Whole-cell current-clamp recordings were obtained from visually-identified cells expressing the viral reporter using borosilicate pipettes (3–5 M Ω) containing (in mM):

10 130 K-gluconate, 6.3 KCl, 0.5 EGTA, 10 HEPES, 4 Mg-ATP, 0.3 Na-GTP and 0.3% biocytin (pH adjusted to 7.3 with KOH). Upon break-in, series resistance (typically 15–25 M Ω) was compensated and only stable recordings (<20% change) were included. Data were acquired using a MultiClamp 700B amplifier (Molecular Devices), sampled at 20 kHz and filtered at 10 kHz. All cells were held at -60 mV
15 with a DC current, and current-step protocols were applied to obtain firing patterns and to extract basic sub-threshold and supra-threshold electrophysiological properties.

Voltage clamp. Cells not expressing the viral reporter were selected according to their pyramidal-cell-shaped soma under IR-DIC visualization and recorded with pipettes containing (in mM): 130 Cs-gluconate, 0.5 EGTA, 7 KCl, 10 HEPES, 4 Mg-

20 ATP, 0.3 Na-GTP, 5 phosphocreatine, 5 QX-314 and 0.3% biocytin (pH adjusted to 7.3 with CsOH). Cells were held continuously at 0 mV for baseline and optogenetic or chemogenetic stimulation. For both current and voltage clamp recording, a baseline of at least 2 minutes was recorded before stimulation. Small pulses (-20 pA or -5 mV, 100 ms at 0.2 Hz or 0.5 Hz) were applied throughout the baseline and
25 CNO application to monitor series resistance changes. Data were analyzed offline using Clampfit 10.2 software (Molecular Devices).

***In vivo* calcium imaging.** Approximately 100 nL of AAV-E2-GCaMP6 virus was injected into the barrel cortex of animals at postnatal day 10. At P27-P34, craniotomies were implanted over the injection site and widefield calcium imaging
30 was performed after recovery from the craniotomy procedure. Briefly, anesthetized (1.5% isoflurane) mice were imaged at 3-4Hz with 4x magnification (Thorlabs CCD camera – 1501M-USB, Thorlabs LED stimulation – DC4104), while air puffs (100-

200ms duration, Picospritzer III) at specific intervals (5-20s) were directed at contralateral whiskers. Multiple recordings were performed, and afterward, the mouse was perfused for histological analysis. Recordings were analyzed in ImageJ by calculating the F/F (change in fluorescence/average fluorescence) for each recording and synched whisker stimulation. A threshold of (5%) F/F was set for both stimulated and spontaneous calcium signal response.

Electrophysiological recordings in humans.

Tissue preparation, culture protocol and inoculation of virus. Four participants (2 male/2 female; age range 22-57 years) underwent a surgical procedure in which brain tissue (temporal lobe and hippocampus) was resected for the treatment of drug resistant epilepsy. In all cases, each participant had previously undergone an initial surgery for placement of subdural and/or depth electrodes for intracranial monitoring to identify the location of seizure onset. The NINDS Institutional Review Board (IRB) approved the research protocol (ClinicalTrials.gov Identifier NCT01273129), and informed consent was obtained from the participants for experimental use of the resected tissue. 300 μ m slices from both hippocampus and temporal lobe were obtained (Leica 1200S Vibratome; Leica Microsystems, Bannockburn, IL) in ice-cold oxygenated sucrose based cutting solution (100mM sucrose, 80mM NaCl, 3.5mM KCl, 24mM NaHCO₃, 1.25mM NaH₂PO₄, 4.5mM MgCl₂, 0.5mM CaCl₂, and 10mM glucose, saturated with 95% O₂ and 5% CO₂) within 30 minutes following neurosurgical resection. Slices were then incubated in the sucrose cutting solution at 33°C for 30 minutes and allowed to cool to room temperature for 15-30 minutes. The slices were transferred to culture medium (Eugène et al., 2014) and placed in an incubator (5% CO₂) at 35°C, for 15 minutes of equilibration. Each individual slice was then transferred onto a 30 mm Millicell Cell Culture Insert (Millipore; Cat No. PICM0RG50) for interface culture and incubated as above. After 12 hours, the culture medium was changed and 1-2 μ l of pAAV_S5E2-dTomato with or without pAAV_S5E2_C1V1-eYFP was directly pipetted onto each slice and placed back into the incubator. For hippocampal slices, the virus was targeted to the subiculum subfield. Culture medium was routinely changed every 2-3 days until electrophysiological analyses. **Electrophysiological recordings.**

Electrophysiological recordings from cultured human slices were performed between

7 to 14 days after viral inoculation. Cultured human slices were transferred to a recording chamber perfused with extracellular solution (130 mM NaCl, 3.5 mM KCl, 24 mM NaHCO₃, 1.25 mM NaH₂PO₄-H₂O, 10 mM glucose, 2.5 mM CaCl₂ and 1.5 mM MgCl₂ saturated with 95% O₂/5% CO₂ (pH 7.4; 300-310 mOsm) at a rate of 3 - 4 ml/min at 33°C. Whole cell patch clamp recordings from pAAV_S5E2-dTomato or pAAV_S5E2_C1V1-eYFP infected neurons were performed with an intracellular solution of the following composition: 130 mM K-gluconate, 10 mM HEPES, 0.6 mM EGTA, 2 mM MgCl₂, 2 mM Na₂ATP, 0.3 mM NaGTP and 0.5% biocytin (pH adjusted to 7.4; osmolality adjusted to 285 -300 mOsm). In some recordings, 130mM K-gluconate was replaced by 90 mM K-gluconate/40 KCl. Intrinsic membrane and firing properties were assayed essentially as described previously (Tricoire, L. et al., 2011, *J. Neurosci*, 31(30):10948-70). 550nm light stimulated optogenetic activation of C1V1 was delivered to the slices via the 40X water immersion objective using a CoolLED pE-4000 Illumination system (Andover, UK). **Biocytin reconstruction and immunocytochemistry.** After electrophysiological recording, slices were drop-fixed in 4% paraformaldehyde in 0.1M PB overnight. Slices were washed in 0.1M PB (3 X 15 minutes) and permeabilized/blocked in 0.5% Triton X-100/10% goat serum in 0.1M PB for at least 2 hours at room temperature. For combined biocytin recovery and immunocytochemistry, an initial incubation (4°C for 40 hours) in primary antibodies diluted at 1:1000 was performed (rabbit anti-PV, Abcam Cat No: ab11427; guinea-pig anti-RFP, SYSY, Cat No: 390005). Slices were washed in 0.1M PB at room temperature 4 x 30 minutes and incubated in secondary antibodies (1:1000 for goat anti guinea-pig Alex-fluor 555, Thermofisher Cat No. A21435; 1:500 for goat anti-rabbit Alex-fluor 647, Thermofisher Cat No. A32733 and 1:1000 Streptavidin Alexa Fluor™ 488; Thermofisher S1123) overnight at 4°C. After a final wash procedure (4 x 30 minutes) the slices were mounted on microscope slides with Prolong Gold antifade (Thermofisher; Cat No. P36930) for subsequent confocal microscopy analysis.

Immunohistochemistry (IHC). Animals injected with the virus were euthanized with Euthasol (Virbac, USA) and transcardially perfused with 4% paraformaldehyde (PFA). The brains were placed in 4% PFA overnight, and then were sectioned at 50–60 µm (in particular, 50 µm) using a Leica VTS1000 vibrosector. Floating brain

sections were permeabilized with 0.1% Triton X-100 and phosphate buffered saline (PBS) for 30 minutes, washed three times with PBS, and incubated in blocking buffer (5% normal donkey serum in PBS) for 30 minutes. The sections were then incubated overnight in blocking buffer with the indicated combinations of the following primary antibodies at 4°C: chicken anti-GFP at 1:1,000 (Abcam USA, ab13970); rabbit anti-DsRed at 1:1000 (Clontech USA 632496); goat anti-PV at 1:1,000 (Swant USA, PVG-213); guinea-pig anti-PV at 1:1,000 (Swant USA, GP-72); rabbit anti-SST at 1:2000 (Peninsula USA, T-4103.0050); mouse anti-Synaptotagmin-2 at 1:250 (ZFIN USA, #ZDB-ATB-081002-25). The sections were then washed three times with PBS incubated with Alexa Fluor-conjugated secondary antibodies at 1:1000 (Invitrogen, USA), counterstained with DAPI (Sigma, USA) and mounted on glass slides using Fluoromount-G (Sigma, USA). Images of brain regions were acquired using a Zeiss LSM800 confocal microscope or a Zeiss Axioimager A1 epifluorescence microscope. The staining of PV IHC within human brain tissues was highly variable; therefore, estimates of viral specificity were made within regions of cortex and subiculum where staining density reflected the known distribution and density of these cells. In view of the variability for human brain tissue, accurate quantification was not obtained by this method.

***In situ* hybridization.** The *in-situ* hybridization probes (Gad1; product #400951, Pvalb; product #421931, VIP; product #415961) used in the studies described herein were designed by Advanced Cell Diagnostics (Newark, CA, USA). The reagents in the RNAscope® Multiplex Fluorescent Reagent Kit v2 (product # 323100), RNAscope® Probe Diluent (product #300041), HYBEZ™ oven (product #321710/321720), humidity control tray (product # 310012), and HYBEZ Humidifying Paper (product #310025) were also from Advanced Cell Diagnostics. TSA Plus Fluorescein, TSA Plus Cyanine 3, and TSA Plus Cyanine 5 from PerkinElmer (#NEL741, #NEL744, and #NEL745). Brain tissue was processed as mentioned in the immunohistochemistry section *supra*. Brain sections were washed one time in PBS followed by three washes in 0.1% Triton X-100 and PBS, mounted on Superfrost Plus glass slides (Fisher Scientific, 12-550-15) and baked at 60°C in the HYBEZ oven for 25 minutes. The slides were then submerged in 4% PFA for 30 minutes then washed 3 times in H₂O. RNAscope H₂O₂ was applied to each section

for 5 minutes at room temperature. The slides were then washed 3 times in H₂O before being submerged in pre-warmed 90°C H₂O for 15 seconds, followed by pre-warmed 90°C RNAscope Target Retrieval for 15 minutes. Slides were washed 3 times in H₂O before RNAscope Protease III was applied onto each section and then
5 incubated for 15 minutes at 40°C in the HYBEZ oven. Slides were washed 3 times in H₂O and then were incubated with probe solution diluted to 1:50 with probe diluent for 2 hours at 40°C in HYBEZ oven. Next, the sections were washed three times in RNAscope wash buffer followed by fluorescence amplification. Of note, probes against the RNA of the reporter revealed a non-specific staining that was likely
10 attributed to the viral DNA. To reveal the viral reporter, the RNAscope protocol was performed with an IHC amplification of the dTomato. The sections were incubated in blocking solution (0.3% Triton X-100 plus 5% normal horse serum in PBS) for 30 minutes. Following this, sections were incubated in antibody solution (0.1% Triton X-100 plus 5% normal horse serum in PBS) with rabbit anti-DsRed at 1:250
15 (Clontech USA 632496) at 4°C overnight. The sections were then washed three times with PBS, incubated with Alexa Fluor-conjugated secondary antibodies at 1:500 (Invitrogen, USA), counterstained with DAPI (Sigma, USA) and mounted on glass slides using Fluoromount-G (Sigma, USA).

Quantifications and statistics. For strength of expression, fluorescence images were
20 taken at a standardized magnification and exposure time. The average pixel intensity of the cell bodies of each cell expressing the viral reporter was recorded and reported as an average over all cells per enhancer. For quantification of co-localization, cells expressing the indicated reporter were counted using only the corresponding color channel, and then, among these cells, the number of cells co-expressing the marker of
25 interest was counted. A cell was considered to be positive for a given marker if the corresponding signal was above background fluorescence. The ratio of cells co-expressing both markers over the total number of cells expressing only the reporter was then calculated, reported herein as mean $\pm$ s.e.m (represented as bar plots in figures herein, for example). Quantifications were performed using a minimum of
30 two independent biological replicates (the specific number of cells, animals and conditions are indicated for each individual quantification in the table presented in FIG. 11, and/or described in the figure legends. Several sections from the same

animal were used when indicated. Data collection and analysis were not performed blind to the conditions of the experiments, but experimenters from different research groups performed the quantifications. No statistical methods were used to predetermine sample sizes, but the sample sizes described were similar to those reported in previous publications.

Example 9 – Viral manipulation of functionally distinct neurons from mice to humans

Described herein are methods and approaches for understanding and treating neuronal and neuropsychiatric diseases by targeting and manipulating specific neuronal cell populations and subtypes. Gaining access to these cell populations in non-human primates and humans has become paramount. While AAVs may be useful for gene delivery in the nervous system, they have a limited genomic payload and are not intrinsically selective for particular neuronal populations. Described herein is the identification of regulatory elements capable of restricting viral expression to broad neuronal classes. To focus the selection of the enhancers as described herein, the regulatory landscape of *SCN1A*, a gene expressed in distinct neuronal populations and whose disruption is associated with severe epilepsy, was specifically examined.

Combining single-cell ATAC-seq data with sequence conservation across species, ten candidate regulatory sequences were identified in the vicinity of the *SCN1A* gene. By investigating each of these elements for its ability to direct viral expression, three enhancers (E2, E5, E6) that collectively targeted the breadth of neuronal populations expressing *SCN1A* were identified. Among these, a particular short regulatory sequence (E2 herein) was found to be capable of restricting viral expression to parvalbumin-expressing cortical interneurons (PV cINs). To fully assess the utility of this element beyond reporter expression, the enhancer element was validated in a variety of contexts, including synaptic tagging, calcium imaging, as well as opto- and chemo-genic approaches, both *ex vivo* and *in vivo*. Moreover, this enhancer element allowed for the selective targeting of PV cINs both during development and across species, including rodents, non-human primates and humans. Demonstrating that this approach provided a generalizable strategy for enhancer discovery, twenty-five additional regulatory elements were selected in the vicinity of

seven genes enriched in PV INs (**FIGS. 15A-1, 15A-2, 16A-1 and 16A-2**). From these, an additional four PV-specific regulatory elements (E11, E14, E22 and E29) were identified, each of which had remarkably selective expression within specific brain regions. Together, the utility of a variety of functionally-tested tools that can be utilized across animal models was demonstrated. Such “viral reagents,” comprising viral delivery vectors harboring a polynucleotide encoding one or more enhancer elements as described herein, as well as one or more target polynucleotide, can be employed to interrogate how functionally distinct neuronal cell-types are affected in the context of neurological, neurodevelopmental and neurodegenerative disease in non-human primates. Ultimately, enhancer-containing viral vectors can serve as agents that therapeutically normalize pathological neuronal activity or gene expression in specific neuronal cell populations.

The enhancers identified and described herein provide access to neuronal populations with particular clinical relevance. These enhancers may be leveraged to alleviate debilitating aspects of Dravet syndrome, for example, by the use of gene therapy or by modulation of neuronal activity. As described in the Examples *supra*, local and systemic injections were utilized for effective viral vector delivery to the brain. With local injections, neurological conditions and pathologies such as focal epilepsy, prefrontal cortex dysfunction or hippocampal memory disorders may be treated or ameliorated. Alternatively, the systemic introduction of virus vectors could be used in contexts where global interventions are necessary, for example, to correct generalized seizures, or for psychiatric and neurodegenerative disorders. The regulatory elements described herein provide for specifically accessing specific cell types for therapeutic contexts.

Indeed, the method and approach for enhancer selection as described herein is advantageous as it is generalizable to other genes. Without intending to be limiting, a subset of seven, representative enhancers (e.g., E1, E5, E6, E11, E14, E22, E29 herein) were identified and demonstrated to have unique specificity for both distinct neuronal populations and regions of the central nervous system. Even with application of stringent criteria (>90% selectivity for the target population), the described enhancer selection method has a high (>20%) success rate. Moreover, as predicted by the high degree of sequence conservation, the representative subset of

enhancers proved equally selective and effective across species, including humans. As such, the described methods provide a reliable means to identify systematically cell-type specific enhancers that are functional across species.

Other Embodiments

5 From the foregoing description, it will be apparent that variations and modifications may be made to the embodiments described herein to adopt them to various usages and conditions. Such embodiments are also within the scope of the following claims.

10 The recitation of a listing of elements in any definition of a variable herein includes definitions of that variable as any single element or combination (or subcombination) of listed elements. The recitation of an embodiment herein includes that embodiment as any single embodiment or in combination with any other embodiments or portions thereof, such as described in one or more sections herein. All patents and publications mentioned in this specification are herein incorporated by
15 reference to the same extent as if each independent patent and publication was specifically and individually indicated to be incorporated by reference.

What is claimed is:

1. A viral vector comprising a transgene polynucleotide sequence and an enhancer polynucleotide sequence that specifically restricts expression of the transgene in parvalbumin (PV)-expressing interneuron cells of the brain.
2. A viral vector comprising an enhancer polynucleotide sequence specifically associated with *SCN1A* gene expression and a transgene polynucleotide sequence, wherein the enhancer sequence restricts expression of the transgene in PV-expressing interneuron cells of the brain.
3. The viral vector of claim 1 or claim 2, wherein the transgene is a reporter gene, a Designer receptor exclusively activated by designer drug (DREADD)-encoding gene, pharmacologically selective actuator molecule (PSAM)-encoding therapeutic gene, or a therapeutic gene.
4. The viral vector of any one of claims 1-3, wherein the transgene is an *SCN1A* gene.
5. The viral vector of any one of claims 1-3, wherein the transgene is a DREADD-encoding gene.
6. The viral vector of claim 5, wherein the DREADD-encoding gene is a Gq-DREADD-encoding gene that is activated by the chemogen clozapine-N4-oxide (CNO).
7. The viral vector of claim 3, wherein the transgene is a pharmacologically selective actuator molecule (PSAM)-encoding therapeutic gene.
8. The viral vector of any one of claims 1-7, wherein the viral vector is recombinant adeno-associated virus (rAAV) vector.
9. A recombinant adeno-associated virus (rAAV) vector comprising an *SCN1A* transgene polynucleotide sequence, or a functional portion thereof, and an enhancer polynucleotide sequence that specifically restricts expression of the *SCN1A* transgene in interneuron or neuron cells of the brain.

10. The rAAV vector of claim 4 or claim 9, wherein an Nav1.1 sodium channel encoded by the *SCN1A* transgene is functionally expressed in interneuron cells following transduction of the interneuron cells by the rAAV vector.
11. The viral or rAAV vector of any one of claims 1-10, wherein the interneuron
5 cells are GABAergic interneuron cells.
12. The rAAV vector of claim 9, wherein the interneuron cells are GABAergic interneuron cells within the brain telencephalon.
13. The rAAV vector of claim 11, wherein the GABAergic interneuron cells express parvalbumin (PV).
- 10 14. The rAAV vector of any one of claims 9-11, wherein the GABAergic interneuron cells express vaso-intestinal peptide (VIP).
15. The rAAV vector of any one of claims 9-11, wherein neuron cells are pyramidal (PYR) neurons of the brain cortex.
- 15 16. The viral or rAAV vector of any one of claims 1-13, wherein the enhancer polynucleotide sequence comprises a nucleotide sequence which contains one or more regions of about 100 bp or longer having at least 75% or greater sequence identity to a polynucleotide sequence of a human enhancer element E1, E2, E3, E4, E7, E8, E9, or E10 (SEQ ID NOs: 15-18 or 21-24, respectively).
- 20 17. The viral or rAAV vector of any one of claims 1-13, wherein the enhancer polynucleotide sequence comprises the polynucleotide sequence of human enhancer element E1, E2, E3, E4, E7, E8, E9, or E10 (SEQ ID NOs: 15-18 or 21-24, respectively).
- 25 18. The viral or rAAV vector of claim 13, wherein the enhancer polynucleotide sequence comprises a nucleotide sequence which contains one or more regions of about 100 bp or longer having at least 75% or greater sequence identity to a polynucleotide sequence of human enhancer element E2 (SEQ ID NO: 16).
19. The rAAV vector of claim 14 or claim 15, wherein the enhancer polynucleotide sequence comprises a nucleotide sequence which contains one or more regions of about 100 bp or longer having at least 75% or greater sequence identity to a

polynucleotide sequence of human enhancer element E6 (SEQ ID NO: 20) or human enhancer element E5 (SEQ ID NO: 19).

20. The rAAV vector of claim 19, wherein the enhancer polynucleotide sequence comprises the polynucleotide sequence of human enhancer element E6 (SEQ ID NO: 20) or the polynucleotide sequence of human enhancer element E5 (SEQ ID NO: 19).

21. The viral or rAAV vector of any one of claims 1-20, wherein the capacity of the vector to package polynucleotide sequences of greater than about 4.7 kb comprises reassembly of multiple rAAV vectors by homologous recombination or by splicing mediated by acceptor sites.

22. The viral or rAAV vector of any one of claims 4 or 9-20, wherein the vector delivers the *SCN1A* gene to *SCN1A*-expressing GABAergic interneuron cells in the brain, and wherein the *SCN1A* gene is functionally expressed, thereby restoring normal levels of *SCN1A* in the interneuron cells following administration of the vector to a subject.

23. The viral or rAAV vector of claim 22, wherein the subject is a human patient.

24. The viral or rAAV vector of claim 23, wherein the human patient is an infant suffering from Dravet syndrome (DS).

25. A viral particle or virus-like particle comprising the viral or rAAV vector of any one of claims 1-24.

26. A cell comprising the viral or rAAV vector of any one of claims 1-24.

27. A cell comprising the viral particle or virus-like particle of claim 25.

28. A pharmaceutical composition comprising the viral or rAAV vector of any one of claims 1-24, and a pharmaceutically acceptable vehicle, carrier, or diluent.

29. A pharmaceutical composition comprising the viral particle or virus-like particle of claim 25, and a pharmaceutically acceptable vehicle, carrier, or diluent.

30. The pharmaceutical composition of claim 28 or claim 29, wherein the pharmaceutical composition is in liquid dosage form.

31. A method of restoring normal levels of *SCN1A* expression in GABAergic interneuron cells or neuron cells in which *SCN1A* expression levels are deficient or

defective, the method comprising contacting the cells with an effective amount of the viral or rAAV vector of any one of claims 4 or 9-22, a viral particle, or a pharmaceutical composition thereof, to restore normal levels of *SCN1A* expression in the GABAergic interneuron cells or neuron cells.

5 32. A method of treating infantile epilepsy and/or seizures in an infant who has or is at risk of having epilepsy, seizures, or Dravet syndrome (DS), the method comprising administering to the infant a therapeutically effective amount of the viral or rAAV vector of any one of claims 1-24, the viral particle or virus-like particle of claim 25, or a pharmaceutical composition of any one of claims 28-30 to treat
10 seizures, epilepsy, or DS in the subject.

33. A method of treating Dravet syndrome (DS) in a subject who has or is at risk of having DS, the method comprising administering to the subject a therapeutically effective amount of the viral or rAAV vector of any one of claims 4 or 9-22, a viral particle or a pharmaceutical composition thereof, to treat DS in the subject.

15 34. A method of inhibiting or preventing seizures and/or epilepsy in a subject having or at risk of having seizures and/or epilepsy, the method comprising systemically administering to the subject a recombinant adeno-associated virus (rAAV) vector comprising an *SCN1A* transgene polynucleotide sequence, or a functional portion thereof, an enhancer polynucleotide sequence that specifically
20 restricts expression of the *SCN1A* transgene in interneuron cells or neuron cells of the cerebral cortex of the subject, and a capsid that enhances transduction of the vector into interneuron cells.

35. The method of any one of claims 32-34, wherein the infant or the subject is a human patient.

25 36. The method of any one of claims 31-35, wherein the enhancer polynucleotide sequence in the rAAV vector comprises one or more regions of about 100 bp or longer having at least 75% or greater sequence identity to a polynucleotide sequence of a human enhancer element E1, E2, E3, E4, E5, E6, E7, E8, E9, or E10 (SEQ ID NOs: 15-24, respectively).

37. The method of claim 36, wherein the enhancer polynucleotide sequence in the rAAV vector comprises one or more regions of about 100 bp or longer having at least 75% or greater sequence identity to a polynucleotide sequence of human enhancer element E2 (SEQ ID NO: 16).

5 38. The method of claim 37, wherein the enhancer polynucleotide sequence is the human enhancer element E2 polynucleotide sequence of SEQ ID NO: 16.

39. The method of claim 36, wherein the enhancer polynucleotide sequence in the rAAV vector comprises one or more regions of about 100 bp or longer having at least 75% or greater sequence identity to a polynucleotide sequence of human enhancer
10 element E6 (SEQ ID NO: 20) or to a polynucleotide sequence of human enhancer element E5 (SEQ ID NO: 19).

40. The method of claim 39, wherein the enhancer polynucleotide sequence is the human enhancer element E6 polynucleotide sequence of SEQ ID NO: 20 or the human enhancer element E5 polynucleotide sequence of SEQ ID NO: 19.

15 41. A method of delivering a transgene for restricted expression in an interneuron cell or neuron cell that expresses an *SCN1A* gene to inhibit or prevent seizures and/or epilepsy in a subject in need thereof, the method comprising: contacting the cell with a recombinant adeno-associated virus (rAAV) vector comprising an *SCN1A* transgene polynucleotide sequence, or a functional portion thereof, and an enhancer
20 polynucleotide sequence that specifically restricts expression of the *SCN1A* transgene in interneuron cells or neuron cells of the cerebral cortex of the subject, thereby inhibiting or preventing seizures and/or epilepsy in the subject.

42. The method of any one of claims 31, or 34-41, wherein the interneuron cells are selected from PV-expressing cortical interneurons (PV-cINs), dis-inhibitory
25 cortical interneurons expressing the vaso-intestinal peptide (VIP cINs), or pyramidal neurons.

43. The method of claim 41 or claim 42, wherein the enhancer polynucleotide sequence in the rAAV vector comprises one or more regions of about 100 bp or longer having at least 75% or greater sequence identity to a polynucleotide sequence

of a human enhancer element E1, E2, E3, E4, E5, E6, E7, E8, E9, or E10 (SEQ ID NOs: 15-24, respectively).

44. The method of claim 41 or claim 42, wherein the enhancer polynucleotide sequence in the rAAV vector comprises one or more regions of about 100 bp or
5 longer having at least 75% or greater sequence identity to a polynucleotide sequence of human enhancer element E2 (SEQ ID NO: 16), to a polynucleotide sequence of human enhancer element E6 (SEQ ID NO: 20), or to a polynucleotide sequence of human enhancer element E5 (SEQ ID NO: 19).

45. The method of claim 41 or claim 42, wherein the enhancer polynucleotide
10 sequence in the rAAV vector is selected from human enhancer elements E1, E2, E3, E4, E5, E6, E7, E8, E9, or E10 (SEQ ID NOs: 15-24, respectively).

46. The method of claim 45, wherein the enhancer polynucleotide sequence is the human enhancer element E2 polynucleotide sequence of SEQ ID NO: 16, the human enhancer element E6 (SEQ ID NO: 20), or the human enhancer element E5 (SEQ ID
15 NO: 19).

47. The method of any one of claims 32-46, wherein the rAAV vector, viral particle, virus-like particle, or pharmaceutical composition is administered systemically.

48. The method of any one of claims 32-46, wherein the rAAV vector, viral
20 particle, or pharmaceutical composition is administered parenterally.

49. The method of claim 47 or claim 48, wherein the rAAV vector, viral particle, virus-like particle, or pharmaceutical composition is administered intravenously.

50. The method of any one of claims 32-46, wherein the rAAV vector, viral particle, virus-like particle, or pharmaceutical composition is administered
25 intracerebrally.

51. The method of any one of claims 32-50, wherein the rAAV vector, viral particle, virus-like particle, or pharmaceutical composition is administered as a prophylactic.

52. The method according to any one of claims 32-51, further comprising
30 administering an adjunct anti-epileptic treatment to the infant or subject.

53. A viral vector comprising a transgene polynucleotide sequence and an enhancer polynucleotide sequence that specifically restricts expression of the transgene in vaso-intestinal peptide-expressing cortical interneuron cells (VIP cINs) of the brain.

5 54. A viral vector comprising an enhancer polynucleotide sequence specifically associated with *SCN1A* gene expression and a transgene polynucleotide sequence, wherein the enhancer sequence restricts expression of the transgene in vaso-intestinal peptide-expressing cortical interneuron cells (VIP cINs) of the brain.

55. The viral vector of claim 53 or claim 54, wherein the enhancer polynucleotide
10 sequence comprises a nucleotide sequence which contains one or more regions of about 100 bp or longer having at least 75% or greater sequence identity to a polynucleotide sequence of human enhancer element E6 (SEQ ID NO: 20).

56. The viral vector of any one of claims 53-55, wherein the enhancer polynucleotide sequence is human enhancer element E6 (SEQ ID NO: 20).

15 57. A viral vector comprising a transgene polynucleotide sequence and an enhancer polynucleotide sequence that specifically restricts expression of the transgene in pyramidal neurons of the brain.

58. A viral vector comprising an enhancer polynucleotide sequence specifically associated with *SCN1A* gene expression and a transgene polynucleotide sequence,
20 wherein the enhancer sequence restricts expression of the transgene in pyramidal neurons of the brain.

59. The viral vector of claim 57 or claim 58, wherein the enhancer polynucleotide sequence comprises a nucleotide sequence which contains one or more regions of about 100 bp or longer having at least 75% or greater sequence identity to a
25 polynucleotide sequence of human enhancer element E5 (SEQ ID NO: 19).

60. The viral vector of any one of claims 57-59, wherein the enhancer polynucleotide sequence is human enhancer element E5 (SEQ ID NO: 19).

61. The viral vector of any one of claims 58-60, wherein the enhancer sequence restricts expression of the transgene in glutamatergic pyramidal neurons of the brain.

62. The viral vector of any one of claims 58-61, wherein the enhancer sequence restricts expression of the transgene in pyramidal neurons in cortical layer 5 of the brain.

63. The viral vector of any one of claims 53-62, wherein the viral vector is a
5 lentiviral vector or a recombinant adeno-associated virus (rAAV) vector.

64. The viral vector of any one of claims 53-63, wherein the transgene is an *SCN1A* gene.

65. A viral vector comprising an enhancer polynucleotide sequence selected from
SEQ ID NOs: 15-24, or a functional portion thereof, wherein the vector specifically
10 targets neuronal cells expressing *SCN1A*.

66. The viral vector of claim 65, wherein the neuronal cells are parvalbumin cortical interneurons (PV cINs), pyramidal (PYR) neurons, or vaso-intestinal peptide cortical interneurons (VIP cIN).

67. A viral vector comprising an enhancer polynucleotide sequence selected from
15 SEQ ID NOs: 25-27, or a functional portion thereof, wherein the vector specifically targets cells expressing *Pvalb*.

68. A viral vector comprising an enhancer polynucleotide sequence selected from SEQ ID NOs: 28-31, or a functional portion thereof, wherein the vector specifically targets cells expressing *Acan*.

69. A viral vector comprising an enhancer polynucleotide sequence selected from
20 SEQ ID NOs: 32-39, or a functional portion thereof, wherein the vector specifically targets cells expressing *Tmem132c*.

70. A viral vector comprising an enhancer polynucleotide sequence selected from
SEQ ID NO: 40 or SEQ ID NO: 41, or a functional portion thereof, wherein the
25 vector specifically targets cells expressing *Lrrc38*.

71. A viral vector comprising an enhancer polynucleotide sequence selected from
SEQ ID NO: 42 or SEQ ID NO: 43, or a functional portion thereof, wherein the
vector specifically targets cells expressing *Inpp5j*.

72. A viral vector comprising an enhancer polynucleotide sequence selected from SEQ ID NOs: 44-47, or a functional portion thereof, wherein the vector specifically targets cells expressing *Mef2c*.
73. A viral vector comprising an enhancer polynucleotide sequence selected from
5 SEQ ID NO: 48 or SEQ ID NO: 49, or a functional portion thereof, wherein the vector specifically targets cells expressing *Pthlh*.
74. A viral vector comprising an enhancer polynucleotide sequence selected from SEQ ID NOS: 15-49, or a functional portion thereof, wherein the vector specifically targets cells PV-expressing cells.
- 10 75. The viral vector of any one of claims 65, or 67-74, wherein the target cells are PV-expressing neuronal cells.
76. The viral vector of any one of claims 65-75, which is a recombinant adeno-associated virus (rAAV) vector.
77. A viral particle or virus-like particle comprising the viral vector of any one of
15 claims 65-76.
78. A cell comprising the viral vector of any one of claims 65-76.
79. A cell comprising the viral particle or virus-like particle of claim 77.
80. A pharmaceutical composition comprising the viral vector of any one of claims 65-76, or the viral particle or virus-like particle of claim 77, and a
20 pharmaceutically acceptable vehicle, carrier, or diluent.
81. A method of restricting expression of a transgene in a neuronal cell of a subject, the method comprising administering to the subject a delivery vector comprising at least one enhancer element polynucleotide comprising a sequence of SEQ ID NO: 15-49 and a transgene polynucleotide, wherein the transgene is
25 specifically expressed in the neuronal cell.
82. The method of claim 81, wherein the transgene is *SCN1A*.
83. The method of claim 82, wherein the neuronal cell is a cortical interneuron expressing parvalbumin (PV cIN).

84. The method of claim 82 or claim 83, wherein the enhancer element polynucleotide comprises a sequence set forth in SEQ ID NOS: 15-18 or SEQ ID NOS: 21-24.
85. The method of claim 82, wherein the neuronal cell is a pyramidal (PYR) cell.
- 5 86. The method of claim 85, wherein the enhancer element polynucleotide comprises the sequence set forth in SEQ ID NO: 19.
87. The method of claim 82, wherein the neuronal cell is a cortical interneuron expressing the vaso-intestinal peptide (VIP cIN).
88. The method of claim 87, wherein the enhancer element polynucleotide
10 comprises the sequence set forth in SEQ ID NO: 20.
89. The method of any one of claims 81-88, wherein the delivery vector is a lentiviral vector or rAAV.
90. The method of claim 89, wherein the delivery vector is administered to the brain.
- 15 91. The method of claim 90, wherein the delivery vector is administered locally or systemically.
92. The method of any one of claims 81-91, wherein the subject is a mammal.
93. The method of claim 92, wherein the subject is a human.
94. A viral vector comprising a human enhancer polynucleotide sequence selected
20 from SEQ ID NOS: 15-49.
95. The viral vector of claim 94, which is a recombinant adeno-associated virus (rAAV) vector.
96. A viral particle or virus-like particle comprising the viral vector of claim 94 or claim 95.
- 25 97. A cell comprising the viral vector of claim 94 or claim 95.
98. A cell comprising the viral particle or the virus-like particle of claim 96.

99. A pharmaceutical composition comprising the viral vector of claim 94 or claim 95, or the viral particle or virus-like particle of claim 96, and a pharmaceutically acceptable vehicle, carrier, or diluent.

1/46

Enhancer	Gene	Target	Specificity	Position	Mouse_mr	Mouse_mm10
E1	Scn1a	PV	0.22	intergenic	chr2	66256056
E2	Scn1a	PV	0.9	intronic	chr2	66364036
E3	Scn1a	PV	0.57	intronic	chr2	66383190
E4	Scn1a	PV	0.14	intronic	chr2	66387764
E5	Scn1a	PYR	0.2	intronic	chr2	66392447
E6	Scn1a	VIP	0.88	intronic	chr2	66401767
E7	Scn1a	PV	0.33	intronic	chr2	66407834
E8	Scn1a	PV	0.61	intronic	chr2	66439814
E9	Scn1a	PV	0.47	intergenic	chr2	66441748
E10	Scn1a	PV	0.23	intergenic	chr2	66450594
E11	Pvalb	PV	0.9	intronic	chr15	78204152
E12	Pvalb	PV	0.59	intronic	chr15	78204583
E13	Pvalb	PV	0.67	intronic	chr15	78205234
E14	Acan	PV	0.94	intergenic	chr7	79052127
E15	Acan	PV	0.79	intergenic	chr7	79053118
E16	Acan	PV	0.58	intronic	chr7	79056553
E17	Acan	PV	0.54	intronic	chr7	79079999
E18	Tmem132c	PV	0.57	intronic	chr5	127243448
E19	Tmem132c	PV	0.57	intronic	chr5	127257256
E20	Tmem132c	PV	0.66	intronic	chr5	127290515
E21	Tmem132c	PV	0.71	intronic	chr5	127300767
E22	Tmem132c	PV	0.94	intronic	chr5	127305150
E23	Tmem132c	PV	0.64	intronic	chr5	127323924
E24	Tmem132c	PV	0.82	intronic	chr5	127331966
E25	Tmem132c	PV	0.73	intronic	chr5	127355818
E26	Lrrc38	PV	0.72	intergenic	chr4	143348892
E27	Lrrc38	PV	0.66	intronic	chr4	143361408
E28	Inpp5j	PV	0.83	intergenic	chr11	3504821
E29	Inpp5j	PV	0.94	intergenic	chr11	3509025
E30	Mef2c	PV	0.77	intergenic	chr13	83503268
E31	Mef2c	PV	0.63	intronic	chr13	83507235
E32	Mef2c	PV	0.7	intronic	chr13	83515122
E33	Mef2c	PV	0.82	intronic	chr13	83518268
E34	Pthlh	PV	0.48	intronic	chr6	147263395
E35	Pthlh	PV	0.86	intergenic	chr6	147266874

FIG. 1A-1

2/46

Mouse_mm10__	Size (bp)	mouse_mm10__DAS__Link
66257335	1279	http://genome.ucsc.edu/cgi-bin/das/mm10/dna?segment=chr2:66256056,66257335
66364653	617	http://genome.ucsc.edu/cgi-bin/das/mm10/dna?segment=chr2:66364036,66364653
66384021	831	http://genome.ucsc.edu/cgi-bin/das/mm10/dna?segment=chr2:66383190,66384021
66388024	260	http://genome.ucsc.edu/cgi-bin/das/mm10/dna?segment=chr2:66387764,66388024
66393109	662	http://genome.ucsc.edu/cgi-bin/das/mm10/dna?segment=chr2:66392447,66393109
66402372	605	http://genome.ucsc.edu/cgi-bin/das/mm10/dna?segment=chr2:66401767,66402372
66410263	2429	http://genome.ucsc.edu/cgi-bin/das/mm10/dna?segment=chr2:66407834,66410263
66441457	1643	http://genome.ucsc.edu/cgi-bin/das/mm10/dna?segment=chr2:66439814,66441457
66442268	520	http://genome.ucsc.edu/cgi-bin/das/mm10/dna?segment=chr2:66441748,66442268
66451140	546	http://genome.ucsc.edu/cgi-bin/das/mm10/dna?segment=chr2:66450594,66451140
78204655	503	http://genome.ucsc.edu/cgi-bin/das/mm10/dna?segment=chr15:78204152,78204655
78204784	201	http://genome.ucsc.edu/cgi-bin/das/mm10/dna?segment=chr15:78204583,78204784
78205766	532	http://genome.ucsc.edu/cgi-bin/das/mm10/dna?segment=chr15:78205234,78205766
79052622	495	http://genome.ucsc.edu/cgi-bin/das/mm10/dna?segment=chr7:79052127,79052622
79053435	317	http://genome.ucsc.edu/cgi-bin/das/mm10/dna?segment=chr7:79053118,79053435
79057054	501	http://genome.ucsc.edu/cgi-bin/das/mm10/dna?segment=chr7:79056553,79057054
79080472	473	http://genome.ucsc.edu/cgi-bin/das/mm10/dna?segment=chr7:79079999,79080472
127244121	673	http://genome.ucsc.edu/cgi-bin/das/mm10/dna?segment=chr5:127243448,127244121
127257594	338	http://genome.ucsc.edu/cgi-bin/das/mm10/dna?segment=chr5:127257256,127257594
127291016	501	http://genome.ucsc.edu/cgi-bin/das/mm10/dna?segment=chr5:127290515,127291016
127301107	340	http://genome.ucsc.edu/cgi-bin/das/mm10/dna?segment=chr5:127300767,127301107
127305592	442	http://genome.ucsc.edu/cgi-bin/das/mm10/dna?segment=chr5:127305150,127305592
127324468	544	http://genome.ucsc.edu/cgi-bin/das/mm10/dna?segment=chr5:127323924,127324468
127332522	556	http://genome.ucsc.edu/cgi-bin/das/mm10/dna?segment=chr5:127331966,127332522
127356133	315	http://genome.ucsc.edu/cgi-bin/das/mm10/dna?segment=chr5:127355818,127356133
143349749	857	http://genome.ucsc.edu/cgi-bin/das/mm10/dna?segment=chr4:143348892,143349749
143362362	954	http://genome.ucsc.edu/cgi-bin/das/mm10/dna?segment=chr4:143361408,143362362
3505244	423	http://genome.ucsc.edu/cgi-bin/das/mm10/dna?segment=chr11:3504821,3505244
3509652	627	http://genome.ucsc.edu/cgi-bin/das/mm10/dna?segment=chr11:3509025,3509652
83504033	765	http://genome.ucsc.edu/cgi-bin/das/mm10/dna?segment=chr13:83503268,83504033
83507457	222	http://genome.ucsc.edu/cgi-bin/das/mm10/dna?segment=chr13:83507235,83507457
83515409	287	http://genome.ucsc.edu/cgi-bin/das/mm10/dna?segment=chr13:83515122,83515409
83519179	911	http://genome.ucsc.edu/cgi-bin/das/mm10/dna?segment=chr13:83518268,83519179
147263584	189	http://genome.ucsc.edu/cgi-bin/das/mm10/dna?segment=chr6:147263395,147263584
147267390	516	http://genome.ucsc.edu/cgi-bin/das/mm10/dna?segment=chr6:147266874,147267390

FIG. 1A-1 (cont.)

3/46

Enhancer	Gene	Target	Specificity	Position	Human_hg38_chr	Human_hg38_start	Human_hg38_stop
E1	Scn1a	PV	22%	intergenic	chr2	165953030	165954796
E2	Scn1a	PV	90%	intronic	chr2	166084035	166084884
E3	Scn1a	PV	57%	intronic	chr2	166090876	166091720
E4	Scn1a	PV	14%	intronic	chr2	166094366	166094633
E5	Scn1a	PYR	20%	intronic	chr2	166103693	166104587
E6	Scn1a	VIP	88%	intronic	chr2	166118214	166118879
E7	Scn1a	PV	33%	intronic	chr2	165892760	165897884
E8	Scn1a	PV	61%	intronic	chr2	166148156	166149792
E9	Scn1a	PV	47%	intergenic	chr2	166150066	166150702
E10	Scn1a	PV	23%	intergenic	chr2	166160023	166160609
E11	Pvalb	PV	90%	intronic	chr22	36816984	36817612
E12	Pvalb	PV	59%	intronic	chr22	36817484	36817720
E13	Pvalb	PV	67%	intronic	chr22	36818134	36818727
E14	Acan	PV	94%	intergenic	chr15	88802240	88802877
E15	Acan	PV	79%	intergenic	chr15	88803290	88803678
E16	Acan	PV	58%	intronic	chr15	88807290	88807962
E17	Acan	PV	54%	intronic	chr15	88833390	88833984
E18	Tmem132c	PV	57%	intronic	chr12	128377753	128378783
E19	Tmem132c	PV	57%	intronic	chr12	128289803	128290279

FIG. 1A-2

4/46		% Conservation between mouse and human
Human_hg38_size_(bp)	human_hg38_DAS_link	
1766	http://genome.ucsc.edu/cgi-bin/das/hg38/dna?segment=chr2:165953030,165954796	69%
849	http://genome.ucsc.edu/cgi-bin/das/hg38/dna?segment=chr2:166084035,166084884	71%
844	http://genome.ucsc.edu/cgi-bin/das/hg38/dna?segment=chr2:166090876,166091720	67%
267	http://genome.ucsc.edu/cgi-bin/das/hg38/dna?segment=chr2:166094366,166094633	78%
894	http://genome.ucsc.edu/cgi-bin/das/hg38/dna?segment=chr2:166103693,166104587	72%
665	http://genome.ucsc.edu/cgi-bin/das/hg38/dna?segment=chr2:166118214,166118879	72%
5124	http://genome.ucsc.edu/cgi-bin/das/hg38/dna?segment=chr2:165892760,165897884	74%
1636	http://genome.ucsc.edu/cgi-bin/das/hg38/dna?segment=chr2:166148156,166149792	75%
636	http://genome.ucsc.edu/cgi-bin/das/hg38/dna?segment=chr2:166150066,166150702	72%
586	http://genome.ucsc.edu/cgi-bin/das/hg38/dna?segment=chr2:166160023,166160609	75%
628	http://genome.ucsc.edu/cgi-bin/das/hg38/dna?segment=chr22:36816984,36817612	78%
236	http://genome.ucsc.edu/cgi-bin/das/hg38/dna?segment=chr22:36817484,36817720	74%
593	http://genome.ucsc.edu/cgi-bin/das/hg38/dna?segment=chr22:36818134,36818727	73%
637	http://genome.ucsc.edu/cgi-bin/das/hg38/dna?segment=chr15:88802240,88802877	72%
388	http://genome.ucsc.edu/cgi-bin/das/hg38/dna?segment=chr15:88803290,88803678	84%
672	http://genome.ucsc.edu/cgi-bin/das/hg38/dna?segment=chr15:88807290,88807962	82%
594	http://genome.ucsc.edu/cgi-bin/das/hg38/dna?segment=chr15:88833390,88833984	86%
1030	http://genome.ucsc.edu/cgi-bin/das/hg38/dna?segment=chr12:128377753,128378783	70%
476	http://genome.ucsc.edu/cgi-bin/das/hg38/dna?segment=chr12:128289803,128290279	78%

FIG. 1A-2 (cont.)

5/46

E20	Tmem132c	PV	66%	intronic	chr12	128323153	128323718
E21	Tmem132c	PV	71%	intronic	chr12	128332503	128332974
E22	Tmem132c	PV	94%	intronic	chr12	128336003	128336491
E23	Tmem132c	PV	64%	intronic	chr12	128365603	128366181
E24	Tmem132c	PV	82%	intronic	chr12	128375853	128376606
E25	Tmem132c	PV	73%	intronic	chr12	128408553	128408930
E26	Lrrc38	PV	72%	intergenic	chr1	13388723	13390212
E27	Lrrc38	PV	66%	intronic	chr1	13469123	13470861
E28	Inpp5j	PV	83%	intergenic	chr22	31124894	31125629
E29	Inpp5j	PV	94%	intergenic	chr22	31132544	31133831
E30	Mef2c	PV	77%	intergenic	chr5	88655733	88657379
E31	Mef2c	PV	63%	intronic	chr5	88872683	88872997
E32	Mef2c	PV	70%	intronic	chr5	88745133	88745535
E33	Mef2c	PV	82%	intronic	chr5	88799783	88801354
E34	Pthlh	PV	48%	intronic	chr12	27969472	27969690
E35	Pthlh	PV	86%	intergenic	chr12	27973822	27974489

FIG. 1A-3

6/46

565	http://genome.ucsc.edu/cgi-bin/das/hg38/dna?segment=chr12:128323153,128323718	77%
471	http://genome.ucsc.edu/cgi-bin/das/hg38/dna?segment=chr12:128332503,128332974	74%
488	http://genome.ucsc.edu/cgi-bin/das/hg38/dna?segment=chr12:128336003,128336491	75%
578	http://genome.ucsc.edu/cgi-bin/das/hg38/dna?segment=chr12:128365603,128366181	85%
753	http://genome.ucsc.edu/cgi-bin/das/hg38/dna?segment=chr12:128375853,128376606	74%
377	http://genome.ucsc.edu/cgi-bin/das/hg38/dna?segment=chr12:128408553,128408930	77%
1489	http://genome.ucsc.edu/cgi-bin/das/hg38/dna?segment=chr1:13388723,13390212	70%
1738	http://genome.ucsc.edu/cgi-bin/das/hg38/dna?segment=chr1:13469123,13470861	71%
735	http://genome.ucsc.edu/cgi-bin/das/hg38/dna?segment=chr22:31124894,31125629	77%
1287	http://genome.ucsc.edu/cgi-bin/das/hg38/dna?segment=chr22:31132544,31133831	74%
1646	http://genome.ucsc.edu/cgi-bin/das/hg38/dna?segment=chr5:88655733,88657379	-
314	http://genome.ucsc.edu/cgi-bin/das/hg38/dna?segment=chr5:88872683,88872997	68%
402	http://genome.ucsc.edu/cgi-bin/das/hg38/dna?segment=chr5:88745133,88745535	70%
1571	http://genome.ucsc.edu/cgi-bin/das/hg38/dna?segment=chr5:88799783,88801354	-
218	http://genome.ucsc.edu/cgi-bin/das/hg38/dna?segment=chr12:27969472,27969690	68%
667	http://genome.ucsc.edu/cgi-bin/das/hg38/dna?segment=chr12:27973822,27974489	69%

FIG. 1A-3 (cont.)

7/46




FIG. 1B-1

8/46




FIG. 1B-2




FIG. 1C




FIG. 1D

11/46





FIG. 2A

12/46

AAV-S5E2-dTomato-Gq-DREADD + PV IHC



AAV-S5E2-dTomato + PV IHC







FIG. 2B

13/46

**FIG. 3A****FIG. 3B****FIG. 3C**






FIG. 3D

15/46

**FIG. 3E**

16/46

**FIG. 3F**

17/46




FIG. 4A




FIG. 4B




FIG. 4C




FIG. 4D







FIG. 4E

18/46

**FIG. 5A****FIG. 5B****FIG. 5C**




FIG. 5D




FIG. 5E




FIG. 6A




FIG. 6B

22/46






FIG. 7



24/46




FIG. 8C




FIG. 8D




FIG. 9A




FIG. 9B






FIG. 9C

27/46

**FIG. 10A****FIG. 10B**

28/46

FIG.	Panel	Enhancer	Marker	Layer	No.	Mean	s.e.m
3	B	E1	Reporter	1-6	3	17.1	4.7
3	B	E2	Reporter	1-6	3	78.7	11.4
3	B	E3	Reporter	1-6	3	49.1	5.5
3	B	E4	Reporter	1-6	3	16	2.6
3	B	E5	Reporter	1-6	3	38.3	17
3	B	E6	Reporter	1-6	3	8.1	0.2
3	B	E7	Reporter	1-6	3	10.4	1.4
3	B	E8	Reporter	1-6	3	41.9	9.2
3	B	E9	Reporter	1-6	3	28.2	5.6
3	B	E10	Reporter	1-6	3	22.3	7.2
3	B	Dlx	Reporter	1-6	3	100.1	7.9
3	B	E1	Reporter / Gad1	1-6	2	95.1	4.3
3	B	E2	Reporter / Gad1	1-6	2	97.8	0
3	B	E3	Reporter / Gad1	1-6	2	95	2.8
3	B	E4	Reporter / Gad1	1-6	2	97.6	2.4
3	B	E5	Reporter / Gad1	1-6	2	63.7	1.5

FIG. 11

29/46

FIG.	Panel	Enhancer	Marker	Layer	No.	Mean	s.e.m
3	B	E6	Reporter / Gad1	1-6	2	98.7	1.3
3	B	E7	Reporter / Gad1	1-6	2	89.5	4
3	B	E8	Reporter / Gad1	1-6	2	85.4	0.5
3	B	E9	Reporter / Gad1	1-6	2	98.6	1.4
3	B	E10	Reporter / Gad1	1-6	2	96.7	2.4
3	B	Dlx	Reporter / Gad1	1-6	4	93.4	1.3
3	B	E1	Reporter / PV	1-6	6	50.3	5.6
3	B	E2	Reporter / PV	1-6	6	74	2.5
3	B	E3	Reporter / PV	1-6	4	45.5	2.1
3	B	E4	Reporter / PV	1-6	6	30.7	3.8
3	B	E5	Reporter / PV	1-6	6	22.8	2.4
3	B	E6	Reporter / PV	1-6	2	16	2
3	B	E7	Reporter / PV	1-6	6	50.8	2.1
3	B	E8	Reporter / PV	1-6	4	47.5	3.4
3	B	E9	Reporter / PV	1-6	6	52.8	6.3
3	B	E10	Reporter / PV	1-6	4	29	2.7

FIG. 11 (cont.)

30/46

FIG.	Panel	Enhancer	Marker	Layer	No.	Mean	s.e.m
3	C	E2	Reporter / Gad1	1	3	0	0
3	C	E2	Reporter / Gad1	2/3	3	94.7	0.9
3	C	E2	Reporter / Gad1	4	3	100	0
3	C	E2	Reporter / Gad1	5	3	97.1	1
3	C	E2	Reporter / Gad1	6	3	97.2	2.8
3	C	E2	Reporter / PV	1	3	0	0
3	C	E2	Reporter / PV	2/3	3	86.5	2.8
3	C	E2	Reporter / PV	4	3	93.5	1.7
3	C	E2	Reporter / PV	5	3	87.3	2.1
3	C	E2	Reporter / PV	6	3	87.3	2.1
3	C	E2	Reporter / SST	1	3	0	0
3	C	E2	Reporter / SST	2/3	3	0	0
3	C	E2	Reporter / SST	4	3	6.1	0.8
3	C	E2	Reporter / SST	5	3	3.3	1
3	C	E2	Reporter / SST	6	3	7.6	1
3	C	E2	Reporter / VIP	1	3	na	na

FIG. 11 (cont.)

31/46

FIG.	Panel	Enhancer	Marker	Layer	No.	Mean	s.e.m
3	C	E2	Reporter / VIP	2/3	3	na	na
3	C	E2	Reporter / VIP	4	3	na	na
3	C	E2	Reporter / VIP	5	3	na	na
3	C	E2	Reporter / VIP	6	3	na	na
3	C	E5	Reporter / Gad1	1	3	91.7	8.3
3	C	E5	Reporter / Gad1	2/3	3	98.9	1.1
3	C	E5	Reporter / Gad1	4	3	95.7	0
3	C	E5	Reporter / Gad1	5	3	35	1.7
3	C	E5	Reporter / Gad1	6	3	94.7	5.3
3	C	E5	Reporter / PV	1	3	0	0
3	C	E5	Reporter / PV	2/3	3	36.3	7.7
3	C	E5	Reporter / PV	4	3	15.8	3.8
3	C	E5	Reporter / PV	5	3	17.8	4.5
3	C	E5	Reporter / PV	6	3	12.9	2.1
3	C	E5	Reporter / SST	1	3	0	0
3	C	E5	Reporter / SST	2/3	3	7.1	4.3

FIG. 11 (cont.)

32/46

FIG.	Panel	Enhancer	Marker	Layer	No.	Mean	s.e.m
3	C	E5	Reporter / SST	4	3	22.4	10.3
3	C	E5	Reporter / SST	5	3	13.1	4.3
3	C	E5	Reporter / SST	6	3	12.3	12.3
3	C	E5	Reporter / VIP	1	3	na	na
3	C	E5	Reporter / VIP	2/3	3	na	na
3	C	E5	Reporter / VIP	4	3	na	na
3	C	E5	Reporter / VIP	5	3	na	na
3	C	E5	Reporter / VIP	6	3	na	na
3	C	E6	Reporter / Gad1	1	3	0	0
3	C	E6	Reporter / Gad1	2/3	3	95.5	4.5
3	C	E6	Reporter / Gad1	4	3	100	0
3	C	E6	Reporter / Gad1	5	3	98.3	1.7
3	C	E6	Reporter / Gad1	6	3	100	0
3	C	E6	Reporter / PV	1	3	0	0
3	C	E6	Reporter / PV	2/3	3	5.4	1.4
3	C	E6	Reporter / PV	4	3	3.4	0.6

FIG. 11 (cont.)

33/46

FIG.	Panel	Enhancer	Marker	Layer	No.	Mean	s.e.m
3	C	E6	Reporter / PV	5	3	2.3	0.6
3	C	E6	Reporter / PV	6	3	1.8	0.3
3	C	E6	Reporter / SST	1	3	0	0
3	C	E6	Reporter / SST	2/3	3	0	0
3	C	E6	Reporter / SST	4	3	7.5	4.6
3	C	E6	Reporter / SST	5	3	5.9	2.6
3	C	E6	Reporter / SST	6	3	2.9	2.9
3	C	E6	Reporter / VIP	1	3	0	0
3	C	E6	Reporter / VIP	2/3	3	92.5	2.6
3	C	E6	Reporter / VIP	4	3	76.3	8.8
3	C	E6	Reporter / VIP	5	3	76.7	3.8
3	C	E6	Reporter / VIP	6	3	96.2	3.8
3	C	E2	Density Gad1+	1	3	0.3	0.5
3	C	E2	Density Gad1+	2/3	3	12.5	1
3	C	E2	Density Gad1+	4	3	30.9	3.1
3	C	E2	Density Gad1+	5	3	35	1.4

FIG. 11 (cont.)

34/46

FIG.	Panel	Enhancer	Marker	Layer	No.	Mean	s.e.m
3	C	E2	Density Gad1+	6	3	18.8	4.8
3	C	E2	Density Gad1-	1	3	0	0
3	C	E2	Density Gad1-	2/3	3	0.7	0.1
3	C	E2	Density Gad1-	4	3	0	0
3	C	E2	Density Gad1-	5	3	1.1	0.6
3	C	E2	Density Gad1-	6	3	0.7	0.9
3	C	E2	Density Gad1+	1	3	2.2	0.9
3	C	E5	Density Gad1+	2/3	3	22.2	1.5
3	C	E5	Density Gad1+	4	3	12.2	0.5
3	C	E5	Density Gad1+	5	3	18.6	0.3
3	C	E5	Density Gad1+	6	3	8.4	1.9
3	C	E5	Density Gad1-	1	3	0.3	0.4
3	C	E5	Density Gad1-	2/3	3	0.3	0.4
3	C	E5	Density Gad1-	4	3	0.6	0
3	C	E5	Density Gad1-	5	3	34.7	3
3	C	E5	Density Gad1-	6	3	0.6	0.8

FIG. 11 (cont.)

35/46

FIG.	Panel	Enhancer	Marker	Layer	No.	Mean	s.e.m
3	C	E6	Density Gad1+	1	3	0	0
3	C	E6	Density Gad1+	2/3	3	34.7	7.2
3	C	E6	Density Gad1+	4	3	36.8	1.3
3	C	E6	Density Gad1+	5	3	13.7	6.6
3	C	E6	Density Gad1+	6	3	13.6	1
3	C	E6	Density Gad1-	1	3	0	0
3	C	E6	Density Gad1-	2/3	3	0	0
3	C	E6	Density Gad1-	4	3	0.6	0.9
3	C	E6	Density Gad1-	5	3	0	0
3	C	E6	Density Gad1-	6	3	0.6	0.9
4	B	E2	Reporter / PV	1-6	4	91.6	0.9
4	B	E2	Reporter / PV	1-6	8	90.8	1.1
4	B	E2	PV / reporter	1-6	19	75.7	2.9
4	D	E2	Reporter at .5	1-6	3	10	1.1
4	D	E2	Reporter at 2	1-6	3	33	10.6
4	D	E2	Reporter at 5	1-6	3	96.5	8.4

FIG. 11 (cont.)

36/46

FIG.	Panel	Enhancer	Marker	Layer	No.	Mean	s.e.m
4	D	E2	Reporter / PV at .5	1-6	3	86.4	0.5
4	D	E2	Reporter / PV at 2	1-6	3	86.6	0.6
4	D	E2	Reporter / PV at 5	1-6	3	91.3	0.8
4	E	E2	Reporter / PV 1-15	1-6	4	56.7	2.3
4	E	E2	Reporter / PV 7-15	1-6	5	67.2	1.5
4	E	E2	Reporter / PV 10-15	1-6	3	81.7	1.1
4	E	E2	Reporter / PV 4-7	1-6	2	60.9	1.9
4	E	E2	Reporter / PV 7-10	1-6	5	78.1	1.8
5	A	E2	Reporter / PV	1-6	4	85.3	0.6
5	A	E2	PV / reporter	1-6	4	87.7	1.2
5	B	E2	Delta	1-6	14	8.2	0.8
5	A	E2	Rat	1-6	1	93	na


FIG. 11 (cont.)

37/46

FIG.	Panel	Enhancer	Marker	Layer	No.	Mean	s.e.m
5	A	E2	Marmoset	1-6	4	91.8	3.1
5	A	E2	Macaque	1-6	4	87.3	0.5
8	A	E2	IR	na	10	119.1	11.8
8	A	E2	AHP	na	10	15	0.9
8	B	E2	Reporter / PV V1	1-6	2	88.1	2.3
8	B	E2	Reporter / PV CG	na	2	80.2	3.7
8	B	E2	Reporter / PV SUB	na	3	91.1	2.1
8	B	E2	Reporter / PV CA1	na	6	80.5	1.9
8	B	E2	Reporter / PV STR	na	3	54.8	5.8
8	B	E2	Reporter / PV BLA	na	3	38.1	4.1
8	B	E2	Reporter / PV SNR	na	2	84.2	1.1
10	A	E2	IR	na	44	56.2	6.7
10	A	E2	AHP	na	45	28.8	0.8

FIG. 11 (cont.)

38/46

**FIG. 12**




FIG. 13A

40/46




FIG. 13A (cont.)

41/46




FIG. 13B




FIG. 14

RE	Gene	Target	Specificity	Position	Chr	Start	Stop	Size (bp)	scATAC PV	scATAC SST	scATAC VIP	scATAC ID2	bATAC Ex	Ms/Hm
E1	Scn1a	PV	22%	intergenic	chr2	66256056	66257335	1279	1	1	1	1	1	69%
E2	Scn1a	PV	90%	intronic	chr2	66364036	66364653	617	1	0	0	1	0	71%
E3	Scn1a	PV	57%	intronic	chr2	66383190	66384021	831	1	1	1	1	1	67%
E4	Scn1a	PV	14%	intronic	chr2	66387764	66388024	260	0	0	0	0	0	78%
E5	Scn1a	PYR	20%	intronic	chr2	66392447	66393109	662	1	1	1	1	1	72%
E6	Scn1a	VIP	88%	intronic	chr2	66401767	66402372	605	0	0	0	0	0	72%
E7	Scn1a	PV	33%	intronic	chr2	66407834	66410263	2429	1	1	1	1	1	74%
E8	Scn1a	PV	61%	intronic	chr2	66439814	66441457	1643	1	1	1	1	1	75%
E9	Scn1a	PV	47%	intergenic	chr2	66441748	66442268	520	1	0	0	0	1	72%
E10	Scn1a	PV	23%	intergenic	chr2	66450594	66451140	546	1	1	1	1	1	75%
E11	Pvalb	PV	90%	intronic	chr15	78204152	78204655	503	1	1	1	1	0	78%
E12	Pvalb	PV	59%	intronic	chr15	78204583	78204784	201	1	1	0	0	0	74%
E13	Pvalb	PV	67%	intronic	chr15	78205234	78205766	532	0	1	0	0	0	73%
E14	Acan	PV	94%	intergenic	chr7	79052127	79052622	495	1	0	0	0	0	72%
E15	Acan	PV	79%	intergenic	chr7	79053118	79053435	317	1	1	1	1	1	84%
E16	Acan	PV	58%	intronic	chr7	79056553	79057054	501	0	0	0	0	0	82%
E17	Acan	PV	54%	intronic	chr7	79079999	79080472	473	1	0	0	0	0	86%

43/46

FIG. 15A-1

RE	Gene	Target	Specificity	Position	Chr	Start	Stop	Size (bp)	scATAC PV	scATAC SST	scATAC VIP	scATAC ID2	bATAC Ex	Ms/Hm
E18	Tmem132c	PV	57%	intronic	chr5	127243448	127244121	673	1	1	0	0	0	70%
E19	Tmem132c	PV	57%	intronic	chr5	127257256	127257594	338	0	1	0	0	0	78%
E20	Tmem132c	PV	66%	intronic	chr5	127290515	127291016	501	1	0	0	0	0	77%
E21	Tmem132c	PV	71%	intronic	chr5	127300767	127301107	340	0	0	0	0	0	74%
E22	Tmem132c	PV	94%	intronic	chr5	127305150	127305592	442	1	0	0	0	0	75%
E23	Tmem132c	PV	64%	intronic	chr5	127323924	127324468	544	1	1	1	1	0	85%
E24	Tmem132c	PV	82%	intronic	chr5	127331966	127332522	556	1	1	1	1	0	74%
E25	Tmem132c	PV	73%	intronic	chr5	127355818	127356133	315	1	0	0	0	0	77%
E26	Lrrc38	PV	72%	intergenic	chr4	143348892	143349749	857	1	1	1	0	0	70%
E27	Lrrc38	PV	66%	intronic	chr4	143361408	143362362	954	1	0	1	0	0	71%
E28	Inpp5j	PV	83%	intergenic	chr11	3504821	3505244	423	1	1	1	1	1	77%
E29	Inpp5j	PV	94%	intergenic	chr11	3509025	3509652	627	1	1	0	0	0	74%
E30	Mef2c	PV	77%	intergenic	chr13	83503268	83504033	765	0	0	0	0	0	-
E31	Mef2c	PV	63%	intronic	chr13	83507235	83507457	222	0	0	0	0	0	68%
E32	Mef2c	PV	70%	intronic	chr13	83515122	83515409	287	0	0	0	0	0	70%
E33	Mef2c	PV	82%	intronic	chr13	83518268	83519179	911	1	1	1	1	1	-
E34	Pthlh	PV	48%	intronic	chr6	147263395	147263584	189	1	1	1	1	1	68%
E35	Pthlh	PV	86%	intergenic	chr6	147266874	147267390	516	0	0	0	0	0	69%

44/46

FIG. 15A-2

45/46

RE	Gene	Target	Specificity	Position	Mouse_ mm10_Chr	Mouse_ mm10_Start	Mouse_ mm10_Stop	Size (bp)	Human_ hg38_Chr	Human_ hg38_Start	Human_ hg38_Stop	Human_hg38 _size_(bp)	Percentage of conservation Mouse vs Human
E1	Scn1a	PV	22%	intergenic	chr2	66256056	66257335	1279	chr2	165953030	165954796	1766	69%
E2	Scn1a	PV	90%	intronic	chr2	66364036	66364653	617	chr2	166084035	166084884	849	71%
E3	Scn1a	PV	57%	intronic	chr2	66383190	66384021	831	chr2	166090876	166091720	844	67%
E4	Scn1a	PV	14%	intronic	chr2	66387764	66388024	260	chr2	166094366	166094633	267	78%
E5	Scn1a	PVR	20%	intronic	chr2	66392447	66393109	662	chr2	166103693	166104587	894	72%
E6	Scn1a	VIP	88%	intronic	chr2	66401767	66402372	605	chr2	166118214	166118879	665	72%
E7	Scn1a	PV	33%	intronic	chr2	66407834	66410263	2429	chr2	165892760	165897884	5124	74%
E8	Scn1a	PV	61%	intronic	chr2	66439814	66441457	1643	chr2	166148156	166149792	1636	75%
E9	Scn1a	PV	47%	intergenic	chr2	66441748	66442268	520	chr2	166150066	166150702	636	72%
E10	Scn1a	PV	23%	intergenic	chr2	66450594	66451140	546	chr2	166160023	166160609	586	75%
E11	Pvalb	PV	90%	intronic	chr15	78204152	78204655	503	chr22	36816984	36817612	628	78%
E12	Pvalb	PV	59%	intronic	chr15	78204583	78204784	201	chr22	36817484	36817720	236	74%
E13	Pvalb	PV	67%	intronic	chr15	78205234	78205766	532	chr22	36818134	36818727	593	73%
E14	Acan	PV	94%	intergenic	chr7	79052127	79052622	495	chr15	88802240	88802877	637	72%
E15	Acan	PV	79%	intergenic	chr7	79053118	79053435	317	chr15	88803290	88803678	388	84%
E16	Acan	PV	58%	intronic	chr7	79056553	79057054	501	chr15	88807290	88807962	672	82%
E17	Acan	PV	54%	intronic	chr7	79079999	79080472	473	chr15	88833390	88833984	594	86%
E18	Tmem132c	PV	57%	intronic	chr5	127243448	127244121	673	chr12	128377753	128378783	1030	70%

FIG. 16A-1

46/46

RE	Gene	Target	Specificity	Position	Mouse_ mm10_Chr	Mouse_ mm10_Start	Mouse_ mm10_Stop	Size (bp)	Human_ hg38_Chr	Human_ hg38_Start	Human_ hg38_Stop	Human_hg38 _size_(bp)	Percentage of conservation Mouse vs Human
E19	Tmem132c	PV	57%	intronic	chr5	127257256	127257594	338	chr12	128289803	128290279	476	78%
E20	Tmem132c	PV	66%	intronic	chr5	127290515	127291016	501	chr12	128323153	128323718	565	77%
E21	Tmem132c	PV	71%	intronic	chr5	127300767	127301107	340	chr12	128332503	128332974	471	74%
E22	Tmem132c	PV	94%	intronic	chr5	127305150	127305592	442	chr12	128336003	128336491	488	75%
E23	Tmem132c	PV	64%	intronic	chr5	127323924	127324468	544	chr12	128365603	128366181	578	85%
E24	Tmem132c	PV	82%	intronic	chr5	127331966	127332522	556	chr12	128375853	128376606	753	74%
E25	Tmem132c	PV	73%	intronic	chr5	127355818	127356133	315	chr12	128408553	128408930	377	77%
E26	Lrrc38	PV	72%	intergenic	chr4	143348892	143349749	857	chr1	13388723	13390212	1489	70%
E27	Lrrc38	PV	66%	intronic	chr4	143361408	143362362	954	chr1	13469123	13470861	1738	71%
E28	Inpp5j	PV	83%	intergenic	chr11	3504821	3505244	423	chr22	31124894	31125629	735	77%
E29	Inpp5j	PV	94%	intergenic	chr11	3509025	3509652	627	chr22	31132544	31133831	1287	74%
E30	Mei2c	PV	77%	intergenic	chr13	83503268	83504033	765	chr5	88655733	88657379	1646	-
E31	Mei2c	PV	63%	intronic	chr13	83507235	83507457	222	chr5	88872683	88872997	314	68%
E32	Mei2c	PV	70%	intronic	chr13	83515122	83515409	287	chr5	88745133	88745535	402	70%
E33	Mei2c	PV	82%	intronic	chr13	83518268	83519179	911	chr5	88799783	88801354	1571	-
E34	Pthlh	PV	48%	intronic	chr6	147263395	147263584	189	chr12	27969472	27969690	218	68%
E35	Pthlh	PV	86%	intergenic	chr6	147266874	147267390	516	chr12	27973822	27974489	667	69%

FIG. 16A-2

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 20/15183

A. CLASSIFICATION OF SUBJECT MATTER

IPC - C07H 21/04, A61K 35/30, C12N 15/86 (2020.01)

CPC - A61K 48/005, C12N 15/861, A61K 48/00, C12N 15/85, G01N 33/5058

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

See Search History document

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

See Search History document

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

See Search History document

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 2013/0347137 A1 (DEISSEROTH et al.) 26 December 2013 (26.12.2013) para [0147], [0132], [0152], [0048], [0054]	1, 3/1
X ----- Y	US 2019/0024119 A1 (ENCODED THERAPEUTICS, INC.) 24 January 2019 (24.01.2019) para [0105], [0051], [0215-0216], [0169], [0213], [0134], [0005], [0009]	2, 3/2, 9 ----- 74, 75/74, 94-98
Y	GenBank accession number AC010127.12 "Homo sapiens BAC clone RP11-218 from 2, complete sequence", 8 April 2005 [online]. [Retrieved 04 May 2020]. Retrieved from the internet <URL: https://www.ncbi.nlm.nih.gov/nuccore/AC010127.12 > whole doc.	74, 75/74, 94-98

☐ Further documents are listed in the continuation of Box C.☐ See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"D" document cited by the applicant in the international application

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

4 May 2020

Date of mailing of the international search report

17 JUL 2020

Name and mailing address of the ISA/US

Mail Stop PCT, Attn: ISA/US, Commissioner for Patents
P.O. Box 1450, Alexandria, Virginia 22313-1450

Facsimile No. 571-273-8300

Authorized officer

Lee Young

Telephone No. PCT Helpdesk: 571-272-4300

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 20/15183

Box No. I Nucleotide and/or amino acid sequence(s) (Continuation of item 1.c of the first sheet)

1. With regard to any nucleotide and/or amino acid sequence disclosed in the international application, the international search was carried out on the basis of a sequence listing:

- a. ☐ forming part of the international application as filed:
 ☐ in the form of an Annex C/ST.25 text file.
 ☐ on paper or in the form of an image file.
- b. ☐ furnished together with the international application under PCT Rule 13ter.1(a) for the purposes of international search only in the form of an Annex C/ST.25 text file.
- c. ☒ furnished subsequent to the international filing date for the purposes of international search only:
 ☒ in the form of an Annex C/ST.25 text file (Rule 13ter.1(a)).
 ☐ on paper or in the form of an image file (Rule 13ter.1(b) and Administrative Instructions, Section 713).

2. ☒ In addition, in the case that more than one version or copy of a sequence listing has been filed or furnished, the required statements that the information in the subsequent or additional copies is identical to that forming part of the application as filed or does not go beyond the application as filed, as appropriate, were furnished.

3. Additional comments:

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 20/15183

Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

3. ☒ Claims Nos.: 4-6, 8, 10-11, 13-33, 35-40, 42-52, 56, 60-64, 76-80, 89-93 and 99
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:
This application contains the following inventions or groups of inventions which are not so linked as to form a single general inventive concept under PCT Rule 13.1. In order for all inventions to be searched, the appropriate additional search fees must be paid.

-----Please see Supplemental Sheet-----

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying additional fees, this Authority did not invite payment of additional fees.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:
1-3, 9, 74, 75(in part), 94-98, limited to target PV-expressing interneuron cells, transgene is a reporter gene, and SEQ ID NO: 15

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.
- ☐ The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT
Information on patent family members

International application No.

PCT/US 20/15183

Continuation of Box No. III Observations where unity of invention is lacking:

Group I+: Claims 1-3, 7, 9, 12, 53-55, 57-59, 65-75 and 94-98 directed to a viral vector comprising an enhancer polynucleotide. The viral vector will be searched to the extent that the vector specifically targets parvalbumin (PV)-expressing interneuron cells of the brain (claims 1, 2, 66, 74), the transgene is a reporter gene (claim 3), and the enhancer polynucleotide sequence is SEQ ID NO: 15 (claim 65, 74, 94). It is believed that claims 1-3, 65-66, 74, 75(in part), 94-98, limited to a viral vector that targets PV-expressing interneuron cells, the transgene is a reporter gene, and the enhancer polynucleotide sequence is SEQ ID NO: 15, encompass this first named invention, and thus these claims will be searched without fee to the extent that the vector encompasses said sequences. [Note that claim 9 is not included in the first named invention because it requires the transgene to be SCN1A instead of a reporter gene.] Additional viral vectors will be searched upon the payment of additional fees. Applicants must specify the claims that encompass any additionally elected viral vectors. Applicants must further indicate, if applicable, the claims which encompass the first named invention, if different than what was indicated above for this group. Failure to clearly identify how any paid additional invention fees are to be applied to the "+" group(s) will result in only the first claimed invention to be searched. An exemplary election would be a viral vector that specifically targets vaso-intestinal peptide-expressing cortical interneuron cells (VIP cINs) of the brain (claims 53, 54, 66), and the enhancer polynucleotide sequence is SEQ ID NO: 20 (claim 55, 65, 94) (i.e. claims 53-55, 65-66, 94-98).

Group II: Claims 34 and 41, directed to a method of inhibiting or preventing seizures and/or epilepsy in a subject by systemically administering to the subject a recombinant adeno-associated virus (rAAV) vector comprising an SCN1A transgene polynucleotide sequence.

Group III+: Claims 81-88, directed to a method of restricting transgene expression to a neuronal cell, comprising administering to a subject a delivery vector polynucleotide, comprising at least one enhancer element and a SCN1A transgene that is specifically expressed in the neuronal cell. Group III+ will be searched upon payment of additional fees. The neuronal expression method will be searched, for example, to the extent that the neuronal cell is a cortical interneuron expressing parvalbumin (PvIN) and the enhancer element polynucleotide comprises the sequence set forth in SEQ ID NO: 15. It is believed that claims 81-84 limited to a neuronal expression method comprising administering a vector with an enhancer element comprising the sequence set forth in SEQ ID NO: 15. Applicants must specify the claims that encompass any additionally elected neuronal expression methods. Failure to clearly identify how any paid additional invention fees are to be applied to the "+" group(s) will result in only the first claimed invention to be searched. An exemplary election would be a neuronal expression method wherein the neuronal cell is a pyramidal (PYR) cell and the enhancer element polynucleotide comprises the sequence set forth in SEQ ID NO: 19 (Claims 81-82 and 85-86).

The inventions listed as Groups I+, II and III+ do not relate to a single general inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical features for the following reasons:

Special Technical Features

No technical features are shared between the viral vector polynucleotide sequences of Group I+ and, accordingly, these groups lack unity a priori.

No technical features are shared between the viral vector polynucleotide sequences of Group III+ and, accordingly, these groups lack unity a priori.

Additionally, even if the inventions listed as Group I+ or Group III+ were considered to share technical features, these shared technical features are previously disclosed by the prior art, as further discussed below.

Groups II and III+ do not require an isolated viral vector composition, as required by group I+.

Groups I+ and III+ do not require a method of inhibiting or preventing seizures and/or epilepsy in a subject, as required by group II.

Group I+ and II do not require a method of restricting expression of a transgene to: a cortical interneuron expressing parvalbumin, a pyramidal cell, or a cortical interneuron expressing the vaso-intestinal peptide of a subject, as required by group III+.

Common Technical Features

The inventions of Groups I+, II and III+ share the technical feature of a viral vector comprising a transgene polynucleotide sequence and an enhancer polynucleotide sequence that specifically restricts expression of the transgene to specific neurons in the brain.

However, this shared technical feature does not represent a contribution over prior art, because the shared technical feature is anticipated by US 2019/0024119 A1 to Encoded Therapeutics Inc. (hereinafter "Encoded"). Encoded teaches said viral vector having regulatory elements restricting expression to PV neurons (para [0105] "In some cases, the expression cassette is part of a vector or a plasmid, e.g., a viral vector or rAAV vector. ... In some cases, the expression cassette comprises one or more PV cell selective regulatory elements ... operably linked to a transgene, wherein the transgene is SCN1A", para [0141] "Parvalbumin ... is expressed in ... GABAergic interneurons ... PV cells are generally considered GABAergic cells."; para [0036] "a regulatory element, which can comprise promoter and/or enhancer sequences, is operatively linked to a coding region if the regulatory element helps initiate transcription of the coding sequence. ").

-----Continued on next page-----

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 20/15183

-----Continued from previous page-----

The inventions of Groups I+, II and III+ also share the technical feature of a recombinant adeno-associated virus (rAAV) vector comprising a SCN1A transgene polynucleotide sequence, however, Encoded teaches said viral vector (para [0105] - "In some cases, the expression cassette is part of a vector or a plasmid, e.g., a viral vector or rAAV vector. ... In some cases, the expression cassette comprises one or more PV cell selective regulatory elements ... operably linked to a transgene, wherein the transgene is SCN1A").

The inventions of Groups I+ and III+ share the technical feature of a vector specifically that targets neuronal cells expressing SCN1A that are parvalbumin cortical interneurons (PV cINs), pyramidal (PYR) neurons, or vaso-intestinal peptide cortical interneurons (VIP cIN).

However, this shared technical feature does not represent a contribution over prior art, because the shared technical feature is made obvious by Encoded in view of US 2014/0309181 A1 to Curna Inc (hereinafter 'Curna Inc'). Encoded teaches expressing SCN1A that are PV cINs and VIP cIN (para [0105] "In some cases, the expression cassette is part of a vector or a plasmid, e.g., a viral vector or rAAV vector. ... In some cases, the expression cassette comprises one or more PV cell selective regulatory elements ... operably linked to a transgene, wherein the transgene is SCN1A", para [0006] "a nucleic acid cassette comprising one or more regulatory elements operably linked to a transgene that results in selective expression in any target cell type, e.g., PV neurons in the CNS ... or non-PV cells in the CNS. para [0148] "One way of selectively expressing a transgene within a subpopulation of cells in the brain is to use a viral vector comprising a transgene operably linked to a cell-type selective regulatory element, or a regulatory element that is selective (or has selective activity) in the subpopulation of cells in the brain, e.g., PV cells", para [0007] "In some cases, the regulatory elements contain less than 600 bp of contiguous sequence from within 10 kb of the transcription start site of ... VIP"). Although Encoded does not expressly teach cells that are PYR, Curna teaches upregulating SCN1A in PYR neurons (para [0192] "The purpose of this experiment is to ensure that the SCN1A protein up-regulated by the small compounds increases the amplitude of the sodium current in the hippocampal GABAergic interneurons, where it is shown to be affected in Dravet syndrome."). Since Encoded teaches treating Dravet's syndrome by over-expression of SCN1A in PV and VIN (above) for treatment of Dravet syndrome (para [0005] "compositions for treating a neurological condition or disorder associated with a genetic defect in the CNS ... In some instances, the neurological condition or disease is Dravet syndrome") it would have been obvious to an artisan of ordinary skill to prepare SCN1A expression vectors for PV, VIP cIN or PYR neurons in a patient, such as to treat Dravet's syndrome.

As the technical features were known in the art at the time of the invention, they cannot be considered special technical features that would otherwise unify the groups.

Groups I+, II and III+ therefore lack unity under PCT Rule 13 because they do not share the same or corresponding special technical feature.

Item 4, continued: claims 4-6, 8, 10-11, 13-33, 35-40, 42-52, 56, 60-64, 76-80, 89-93 and 99 are not not drafted in accordance with the second and third sentences of Rule 6.4(a) regarding multiply dependent claims.