



Office de la Propriété

Intellectuelle
du Canada

Un organisme
d'Industrie Canada

Canadian
Intellectual Property
Office

An agency of
Industry Canada

CA 2195856 C 2008/09/30

(11)(21) **2 195 856**

(12) **BREVET CANADIEN
CANADIAN PATENT**

(13) **C**


(86) Date de dépôt PCT/PCT Filing Date: 1995/07/31
(87) Date publication PCT/PCT Publication Date: 1996/02/15
(45) Date de délivrance/Issue Date: 2008/09/30
(85) Entrée phase nationale/National Entry: 1997/01/23
(86) N° demande PCT/PCT Application No.: US 1995/009702
(87) N° publication PCT/PCT Publication No.: 1996/004281
(30) Priorité/Priority: 1994/08/01 (US283,953)

(51) Cl.Int./Int.Cl. *C07D 473/18* (2006.01),
A61K 31/52 (2006.01), *A61P 35/00* (2006.01),
C07D 473/06 (2006.01), *C07D 473/40* (2006.01),
C07D 239/46 (2006.01), *C07D 239/48* (2006.01),
C07D 239/50 (2006.01)

(72) Inventeurs/Inventors:
MOSCHEL, ROBERT C., US;
PEGG, ANTHONY E., US;
DOLAN, M. EILEEN, US;
CHAE, MI-YOUNG, US

(73) Propriétaires/Owners:

(54) Titre : O⁶-BENZYLGUANINES SUBSTITUEES ET UTILISATION CONNEXE
(54) Title: SUBSTITUTED O⁶-BENZYLGUANINES AND USE THEREOF



(57) Abrégé/Abstract:

The present invention provides AGT inactivating compounds such as substituted O⁶-benzylguanines of formula (I) wherein 7- or 9-substituted 8-aza-O⁶-benzylguanines, 7,8-disubstituted O⁶-benzylguanines, 7,9-disubstituted O⁶-benzylguanines, 4(6)-substituted 2-amino-5-nitro-6(4)-benzyloxypyrimidines, and 4(6)-substituted 2-amino-5-nitroso-6(4)-benzyloxypyrimidines, as well as pharmaceutical compositions comprising such compounds along with a pharmaceutically acceptable carrier. The present invention further provides a method of enhancing the chemotherapeutic treatment of tumor cells in a mammal with an antineoplastic

(73) Propriétaires(suite)/Owners(continued):

THE UNITED STATES OF AMERICA, REPRESENTED BY THE SECRETARY, DEPARTMENT OF HEALTH AND HUMAN SERVICES, US; ARCH DEVELOPMENT CORPORATION, US; PENN STATE RESEARCH FOUNDATION (THE), US

(74) Agent: OGILVY RENAULT LLP/S.E.N.C.R.L.,S.R.L.

(57) Abrégé(suite)/Abstract(continued):

alkylating agent which causes cytotoxic lesion at the O⁶-position of guanine, by administering to a mammal an effective amount of one of the aforesaid compounds, 2,4-diamino-6-benzyloxy-s-triazine, 5-substituted 2,4-diamino-6-benzyloxyprymidines, or 8-aza-O⁶-benzylguanine, and administering to the mammal an effective amount of an antineoplastic alkylating agent which causes cytotoxic lesions of the O⁶-position of guanine.




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ :		A1	(11) International Publication Number: WO 96/04281
C07D 473/18, 487/04, 473/06, 473/40, 239/46, 239/48, 239/50, 251/52, A61K 31/52, 31/505 // (C07D 487/04, 249:00, 239:00)			(43) International Publication Date: 15 February 1996 (15.02.96)
(21) International Application Number:	PCT/US95/09702		(72) Inventors; and
(22) International Filing Date:	31 July 1995 (31.07.95)		(75) Inventors/Applicants (for US only): MOSCHEL, Robert, C. [US/US]; 8204 Glendale Drive, Frederick, MD 21702 (US). PEGG, Anthony, E. [US/US]; 9 Brownstone Drive, Hershey, PA 17033 (US). DOLAN, M., Eileen [US/US]; 821 S. East Avenue, Oak Park, IL 60304 (US). CHAE, Mi-Young [KR/US]; Apartment 203, 1418H Taney Avenue, Frederick, MD 21702 (US).
(30) Priority Data:	283,953	1 August 1994 (01.08.94)	US
(60) Parent Application or Grant			(74) Agents: GAGALA, Bruce, M. et al.; Leydig, Voit & Mayer, Ltd., Suite 4900, Two Prudential Plaza, 180 North Stetson, Chicago, IL 60601-6780 (US).
(63) Related by Continuation	US	08/283,953 (CIP)	2195856
	Filed on	1 August 1994 (01.08.94)	
(71) Applicants (for all designated States except US): THE UNITED STATES OF AMERICA, represented by THE SECRETARY, DEPARTMENT OF HEALTH AND HUMAN SERVICESNational Institutes of Health [US/US]; Office of Technology Transfer, Box OTT, Bethesda, MD 20892-9902 (US). ARCH DEVELOPMENT CORPORATION [US/US]; The University of Chicago, 1101 East 58th Street, Chicago, IL 60637 (US). THE PENN STATE RESEARCH FOUNDATION, INC. [US/US]; 113 Technology Center, University Park, IL 16802 (US).	(81) Designated States: AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LT, LU, LV, MD, MG, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TT, UA, UG, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD, SZ, UG).		
Published With international search report.			

(54) Title: SUBSTITUTED O⁶-BENZYLGUANINES AND 6(4)-BENZYLOXYPYRIMIDINES

(57) Abstract

The present invention provides AGT inactivating compounds such as substituted O⁶-benzylguanines of formula (I) wherein 7- or 9-substituted 8-aza-O⁶-benzylguanines, 7,8-disubstituted O⁶-benzylguanines, 7,9-disubstituted O⁶-benzylguanines, 4(6)-substituted 2-amino-5-nitro-6(4)-benzyloxyprymidines, and 4(6)-substituted 2-amino-5-nitro-6(4)-benzyloxyprymidines, as well as pharmaceutical compositions comprising such compounds along with a pharmaceutically acceptable carrier. The present invention further provides a method of enhancing the chemotherapeutic treatment of tumor cells in a mammal with an antineoplastic alkylating agent which causes cytotoxic lesion at the O⁶-position of guanine, by administering to a mammal an effective amount of one of the aforesaid compounds, 2,4-diamino-6-benzyloxy-s-triazine, 5-substituted 2,4-diamino-6-benzyloxyprymidines, or 8-aza-O⁶-benzylguanine, and administering to the mammal an effective amount of an antineoplastic alkylating agent which causes cytotoxic lesions of the O⁶-position of guanine.



SUBSTITUTED O⁶-BENZYLGUANINES AND USE THEREOFTECHNICAL FIELD OF THE INVENTION

The present invention relates to substituted O⁶-benzylguanines, O⁶-benzyl-8-azaguanines, and 6(4)-benzyloxypyrimidines, pharmaceutical compositions comprising such compounds, and methods of using such compounds. The subject compounds are particularly useful in inactivating the human DNA repair protein O⁶-alkylguanine-DNA alkyltransferase.

10

BACKGROUND OF THE INVENTION

The inactivation of the human DNA repair protein O⁶-alkylguanine-DNA alkyltransferase (AGT) by O⁶-benzylguanine leads to a dramatic enhancement in the cytotoxic response of human tumor cells and tumor xenografts to chemotherapeutic drugs whose mechanism of action involves modification of DNA guanine residues at the O⁶-position (Dolan et al., *Proc. Natl. Acad. Sci. U.S.A.*, 87, 5368-5372 (1990); Dolan et al., *Cancer Res.*, 51, 3367-3372 (1991); Dolan et al., *Cancer Commun.*, 2, 371-377 (1990); Mitchell et al., *Cancer Res.*, 52, 1171-1175 (1992); Friedman et al., *J. Natl. Cancer Inst.*, 84, 1926-1931 (1992); Felker et al., *Cancer Chem. Pharmacol.*, 32, 471-476 (1993); Dolan et al., *Cancer Chem. Pharmacol.*, 32, 221-225 (1993); Dolan et al., *Biochem. Pharmacol.*, 46, 285-290 (1993)). The AGT inactivating activity of a large number of O⁶-benzylguanine analogs have been compared with the aim of obtaining information about the types of substituent groups and the sites at which they could be attached to O⁶-benzylguanine without significantly lowering its AGT-inactivating activity (Moschel et al., *J. Med. Chem.*, 35, 4486-4491 (1992); Chae et al., *J. Med. Chem.*, 37, 342-347 (1994)). While these studies led to the production of a variety of analogs that were as potent or somewhat less potent than O⁶-benzylguanine, none of the analogs were better than O⁶-benzylguanine.


Thus, there remains a need for additional compounds which are capable of enhancing the chemotherapeutic treatment of tumor cells in a mammal with an antineoplastic alkylating agent which causes cytotoxic lesions at the O^6 -position of guanine. The present invention provides such compounds and associated pharmaceutical compositions and treatment methods. These and other objects and advantages of the present invention, as well as additional inventive features, will be apparent from the description of the invention provided herein.

BRIEF SUMMARY OF THE INVENTION

The present invention provides 7- and 8-substituted O^6 -benzylguanine derivatives, 7,8-disubstituted O^6 -benzylguanine derivatives, 7,9-disubstituted O^6 -benzylguanine derivatives, 8-aza- O^6 -benzylguanine derivatives, and 4(6)-substituted 2-amino-5-nitro-6(4)-benzyloxypyrimidine and 2-amino-5-nitroso-6(4)-benzyloxypyrimidine derivatives which have been found to be effective AGT inactivators, as well as pharmaceutical compositions comprising such derivatives along with a pharmaceutically acceptable carrier. The present invention further provides a method of enhancing the chemotherapeutic treatment of tumor cells in a mammal with an antineoplastic alkylating agent which causes cytotoxic lesions at the O^6 -position of guanine, by administering to a mammal an effective amount of one of the aforesaid derivatives, 2,4-diamino-6-benzyloxy-s-triazine, 5-substituted 2,4-diamino-6-benzyloxypyrimidines, or 8-aza- O^6 -benzylguanine, and administering to the mammal an effective amount of an antineoplastic alkylating agent which causes cytotoxic lesions at the O^6 -position of guanine.

DESCRIPTION OF THE PREFERRED EMBODIMENTS

The present invention provides a compound of the formula I



5


wherein R₁ is a substituent selected from the group consisting of amino, hydroxy, C₁-C₄ alkylamino, C₁-C₄ dialkylamino, and C₁-C₄ acylamino (although, as explained in further detail below, other substituents can be placed at this 2-position), R₂ is a substituent selected from the group consisting of hydrogen, C₁-C₄ alkyl, C₁-C₄ aminoalkyl, C₁-C₄ hydroxyalkyl, C₁-C₄ alkylaminoalkyl, C₁-C₄ dialkylaminoalkyl, C₁-C₄ cyanoalkyl, C₁-C₄ carbamoylalkyl, C₁-C₄ pivaloylalkyl, C₁-C₆ alkylcarbonyloxy C₁-C₄ alkyl, carbo C₁-C₄ alkoxyalkyl, ribose, 2'-deoxyribose, the conjugate acid form of a C₁-C₄ carboxyalkyl, and the carboxylate anion of a C₁-C₄ carboxyalkyl as the sodium salt (although, as explained in further detail below, other substituents can be placed at this N⁹-position), and R₃ is a substituent selected from the group consisting of hydrogen, halo, C₁-C₄ alkyl, C₁-C₄ hydroxyalkyl, thiol, C₁-C₄ alkylthio, trifluoromethylthio, C₁-C₄ thioacyl, hydroxy, C₁-C₄ alkoxy, trifluoromethoxy, methanesulfonyloxy, trifluoromethanesulfonyloxy, C₁-C₄ acyloxy, amino, C₁-C₄ aminoalkyl, C₁-C₄ alkylamino, C₁-C₄ dialkylamino, trifluoromethylamino, ditrifluoromethylamino, aminomethanesulfonyl, C₁-C₄ aminoacyl, aminotrifluoromethylcarbonyl, formylamino, nitro, nitroso, C₁-C₄ alkyldiazo, C₅-C₆ aryldiazo, trifluoromethyl, C₁-C₄ haloalkyl, halomethyl, C₁-C₄ cyanoalkyl, cyanomethyl, cyano,

C_1-C_4 alkyloxycarbonyl, C_1-C_4 alkylcarbonyl, phenyl, phenylcarbonyl, formyl, C_1-C_4 alkoxyethyl, phenoxyethyl, C_2-C_4 vinyl, C_2-C_4 ethynyl, and SO_nR' wherein n is 0, 1, 2, or 3 and R' is hydrogen, C_1-C_4 alkyl, amino, or phenyl, with the proviso that R_1 is not amino when both R_2 and R_3 are hydrogen, and with the proviso that R_1 is not amino or methylamino when R_2 is ribose or 2'-deoxyribose and R_3 is hydrogen. It is to be understood that the substituents are defined herein such that the group farthest from the point of attachment of the substituent is named first. By way of illustration, C_1-C_6 alkylcarbonyloxy C_1-C_4 alkyl includes pivaloyloxymethyl.

Suitable compounds of the above formula include those compounds wherein R_1 is selected from the group consisting of amino, hydroxy, C_1-C_4 alkylamino, C_1-C_4 dialkylamino, and C_1-C_4 alkylcarbonylamino, R_2 is selected from the group consisting of hydrogen, C_1-C_4 alkyl, and C_1-C_6 alkylcarbonyloxy C_1-C_4 alkyl, and R_3 is selected from the group consisting of amino, halo, C_1-C_4 alkyl, hydroxy, and trifluoromethyl. Other suitable compounds include those wherein R_1 is selected from the group consisting of amino, hydroxy, methylamino, dimethylamino, and acetylamino, R_2 is selected from the group consisting of hydrogen, methyl, and pivaloyloxymethyl, and R_3 is selected from the group consisting of amino, bromo, methyl, hydroxy, and trifluoromethyl. Examples of suitable compounds include 8-amino- O^6 -benzylguanine, 8-methyl- O^6 -benzylguanine, 8-hydroxy- O^6 -benzylguanine, 8-bromo- O^6 -benzylguanine, 8-trifluoromethyl- O^6 -benzylguanine, O^6 -benzylxanthine, O^6 -benzyluric acid, N^2 -acetyl- O^6 -benzyl-8-oxoguanine, O^6 -benzyl- N^2 -methylguanine, O^6 -benzyl- N^2,N^2 -dimethylguanine, O^6 -benzyl-8-trifluoromethyl-9-methylguanine, O^6 -benzyl-8-bromo-9-methylguanine, and O^6 -benzyl-8-bromo-9-(pivaloyloxymethyl)guanine.

4a

In one aspect of the present invention there is a pharmaceutical composition comprising a pharmaceutically acceptable carrier and a compound of the formula




wherein R_1 is a substituent selected from the group consisting of amino, hydroxy, C_1-C_4 alkylamino, C_1-C_4 dialkylamino, and C_1-C_4 alkylcarbonylamino, R_2 is a substituent selected from the group consisting of hydrogen, C_1-C_4 alkyl, C_1-C_4 aminoalkyl, C_1-C_4 hydroxyalkyl, C_1-C_4 alkylamino C_1-C_4 alkyl, C_1-C_4 dialkylamino alkyl, C_1-C_4 cyanoalkyl, C_1-C_4 carbamoylalkyl, C_1-C_4 pivaloylalkyl, C_1-C_6 alkylcarbonyloxy C_1-C_4 alkyl, C_1-C_4 alkoxycarbonylalkyl, 2'-deoxyribose, the conjugate acid form of a C_1-C_4 carboxyalkyl, and the carboxylate anion of a C_1-C_4 carboxyalkyl as the sodium salt, and R_3 is a substituent selected from the group consisting of halo, C_1-C_4 alkyl, C_1-C_4 hydroxyalkyl, mercapto, C_1-C_4 alkylthio, trifluoromethylthio, C_1-C_4 alkylthiocarbonyl, hydroxy, C_1-C_4 alkoxy, trifluoromethoxy, methanesulfonyloxy, trifluoromethanesulfonyloxy, C_1-C_4 alkylcarbonyloxy, amino, C_1-C_4 aminoalkyl, C_1-C_4 alkylamino, C_1-C_4 dialkylamino, trifluoromethylamino, ditrifluoromethylamino, aminomethanesulfonyl, amino C_1-C_4 alkylcarbonyl, aminotrifluoromethylcarbonyl, formylamino, nitro, nitroso, C_1-C_4 alkyldiazo, C_5-C_6 aryldiazo, trifluoromethyl, C_1-C_4

4b

haloalkyl, C₁-C₄ cyanoalkyl, cyano, C₁-C₄ alkyloxycarbonyl, C₁-C₄ alkylcarbonyl, phenyl, phenylcarbonyl, formyl, C₁-C₄ alkoxyethyl, phenoxyethyl, C₂-C₄ vinyl, C₂-C₄ ethynyl, and SO_nR' wherein n is 0, 1, 2, or 3 and R' is hydrogen, C₁-C₄ alkyl, amino, or phenyl.


In another aspect of the present invention there is a use of a compound of the formula



wherein R₁ is a substituent selected from the group consisting of amino, hydroxy, C₁-C₄ alkylamino, C₁-C₄ dialkylamino, and C₁-C₄ alkylcarbonyl, R₂ is a substituent selected from the group consisting of hydrogen, C₁-C₄ alkyl, C₁-C₄ aminoalkyl, C₁-C₄ hydroxyalkyl, C₁-C₄ alkylamino C₁-C₄ alkyl, C₁-C₄ dialkylamino alkyl, C₁-C₄ cyanoalkyl, C₁-C₄ carbamoylalkyl, C₁-C₄ pivaloylalkyl, C₁-C₆ alkylcarbonyloxy C₁-C₄ alkyl, C₁-C₄ alkoxy carbonylalkyl, 2'-deoxyribose, the conjugate acid form of a C₁-C₄ carboxyalkyl, and the carboxylate anion of a C₁-C₄ carboxyalkyl as the sodium salt, and R₃ is a substituent selected from the group consisting of halo, C₁-C₄ alkyl, C₁-C₄ hydroxyalkyl, mercapto, C₁-C₄ alkylthio, trifluoromethylthio, C₁-C₄ alkylthiocarbonyl, hydroxy, C₁-C₄ alkoxy, trifluoromethoxy, methanesulfonyloxy, trifluoromethanesulfonyloxy, C₁-C₄ alkylcarbonyloxy, amino, C₁-C₄ aminoalkyl, C₁-C₄ alkylamino, C₁-C₄ dialkylamino,

trifluoromethylamino, ditrifluoromethylamino, aminomethanesulfonyl, amino C₁-C₄ alkylcarbonyl, aminotrifluoromethylcarbonyl, formylamino, nitro, nitroso, C₁-C₄ alkyldiazo, C₅-C₆ aryldiazo, trifluoromethyl, C₁-C₄ haloalkyl, C₁-C₄ cyanoalkyl, cyano, C₁-C₄ alkyloxycarbonyl, C₁-C₄ alkylcarbonyl, phenyl, phenylcarbonyl, formyl, C₁-C₄ alkoxyethyl, phenoxyethyl, C₂-C₄ vinyl, C₂-C₄ ethynyl, and SO_nR' wherein n is 0, 1, 2, or 3 and R' is hydrogen, C₁-C₄ alkyl, amino, or phenyl, in the manufacture of a medicament for enhancing the chemotherapeutic treatment of tumor cells in a mammal with an antineoplastic alkylating agent that causes cytotoxic lesions at the O⁶-position of guanine.

In yet another aspect of the present invention there is a product containing compound of the formula



wherein R₁ is a substituent selected from the group consisting of amino, hydroxy, C₁-C₄ alkylamino, C₁-C₄ dialkylamino, and C₁-C₄ alkylcarbonyl, R₂ is a substituent selected from the group consisting of hydrogen, C₁-C₄ alkyl, C₁-C₄ aminoalkyl, C₁-C₄ hydroxyalkyl, C₁-C₄ alkylamino C₁-C₄ alkyl, C₁-C₄ dialkylamino alkyl, C₁-C₄ cyanoalkyl, C₁-C₄ carbamoylalkyl, C₁-C₄ pivaloylalkyl, C₁-C₆ alkylcarbonyloxy C₁-C₄ alkyl, C₁-C₄ alkoxy carbonylalkyl, 2'-deoxyribose, the conjugate acid form of a C₁-C₄ carboxyalkyl, and the carboxylate anion of a C₁-C₄

4d

carboxyalkyl as the sodium salt, and R₃ is a substituent selected from the group consisting of halo, C₁-C₄ alkyl, C₁-C₄ hydroxyalkyl, mercapto, C₁-C₄ alkylthio, trifluoromethylthio, C₁-C₄ alkylthiocarbonyl, hydroxy, C₁-C₄ alkoxy, trifluoromethoxy, methanesulfonyloxy, trifluoromethanesulfonyloxy, C₁-C₄ alkylcarbonyloxy, amino, C₁-C₄ aminoalkyl, C₁-C₄ alkylamino, C₁-C₄ dialkylamino, trifluoromethylamino, ditrifluoromethylamino, aminomethanesulfonyl, amino C₁-C₄ alkylcarbonyl, aminotrifluoromethylcarbonyl, formylamino, nitro, nitroso, C₁-C₄ alkyldiazo, C₅-C₆ aryl diazo, trifluoromethyl, C₁-C₄ haloalkyl, C₁-C₄ cyanoalkyl, cyano, C₁-C₄ alkyloxycarbonyl, C₁-C₄ alkylcarbonyl, phenyl, phenylcarbonyl, formyl, C₁-C₄ alkoxyethyl, phenoxyethyl, C₂-C₄ vinyl, C₂-C₄ ethynyl, and SO_nR' wherein n is 0, 1, 2, or 3 and R' is hydrogen, C₁-C₄ alkyl, amino, or phenyl, and an antineoplastic alkylating agent which causes cytotoxic lesions at the O⁶-position of guanine, for simultaneous, separate or sequential use for enhancing chemotherapeutic treatment of tumor cells in a mammal.

The following terms are understood as equivalent:


C₁₋₄ acylamino - C₁₋₄ alkylcarbonylamino;

thiol - mercapto;

C₁₋₄ thioacyl - C₁₋₄ alkylthiocarbonyl; and

C₁₋₄ acyloxy - C₁₋₄ alkylcarbonyloxy.

The present invention also provides a compound of the formula II




(II)

5


wherein R₁ is NO₂ or NO, and R₂ is a substituent selected from the group consisting of hydrogen, halo, C₁-C₄ alkyl, C₁-C₄ hydroxyalkyl, thiol, C₁-C₄ alkylthio, trifluoromethylthio, C₁-C₄ thioacyl, hydroxy, C₁-C₄ alkyloxy, trifluoromethoxy, methanesulfonyloxy, trifluoromethanesulfonyloxy, C₁-C₄ acyloxy, C₁-C₄ aminoalkyl, C₁-C₄ alkylamino, C₁-C₄ dialkylamino, trifluoromethylamino, ditrifluoromethylamino, aminomethanesulfonyl, amino C₁-C₄ alkylcarbonyl, aminotrifluoromethylcarbonyl, formylamino, nitro, nitroso, C₁-C₄ alkyldiazo, C₅-C₆ aryldiazo, trifluoromethyl, C₁-C₄ haloalkyl, cyanomethyl, C₁-C₄ cyanoalkyl, cyano, C₁-C₄ alkyloxycarbonyl, C₁-C₄ alkylcarbonyl, phenyl, phenylcarbonyl, formyl, C₁-C₄ alkoxyethyl, phenoxyethyl, C₂-C₄ vinyl, C₂-C₄ ethynyl, and SO_nR' wherein n is 0, 1, 2, or 3 and R' is hydrogen, C₁-C₄ alkyl, amino, or phenyl. Suitable compounds include those compounds wherein R₁ is NO₂ and R₂ is hydrogen or a C₁-C₄ alkyl. Examples of suitable compounds include 2-amino-4-benzyl-5-nitropyrimidine and 2-amino-4-benzyl-6-methyl-5-nitropyrimidine.

The present invention further provides a compound of the formula III




wherein R is selected from the group consisting of C₁-C₄ alkyl, C₁-C₄ alkyloxycarbonyl C₁-C₄ alkyl, carboxy C₁-C₄ alkyl, cyano C₁-C₄ alkyl, aminocarbonyl C₁-C₄ alkyl, hydroxy C₁-C₄ alkyl, and C₁-C₄ alkyloxy C₁-C₄ alkyl. Suitable compounds of the above formula include those wherein R is selected from the group consisting of C₁-C₄ alkyl and C₁-C₆ alkylcarbonyloxy C₁-C₄ alkyl. Examples of suitable compounds include 8-aza-*O*⁶-benzyl-9-methylguanine and 8-aza-*O*⁶-benzyl-9-(pivaloyloxymethyl)guanine.

The present invention further provides a compound of the formula IV



wherein R is selected from the group consisting of C₁-C₄ alkyl, C₁-C₆ alkylcarbonyloxy C₁-C₄ alkyl, C₁-C₄ alkyloxycarbonyl C₁-C₄ alkyl, carboxy C₁-C₄ alkyl, cyano C₁-C₄ alkyl, aminocarbonyl C₁-C₄ alkyl, hydroxy C₁-C₄ alkyl, and C₁-C₄ alkyloxy C₁-C₄ alkyl. Suitable compounds include those wherein said R is C₁-C₆ alkylcarbonyloxy C₁-C₄ alkyl. An example of a suitable compound is 8-aza-*O*⁶-benzyl-7-(pivaloyloxymethyl)guanine.

The present invention further provides a compound of the formula V



5

wherein R₁ is selected from the group consisting of hydrogen, halo, C₁-C₄ alkyl, halo C₁-C₄ alkyl, C₁-C₆ alkylcarbonyloxy C₁-C₄ alkyl, C₁-C₄ alkyloxycarbonyl C₁-C₄ alkyl, carboxy C₁-C₄ alkyl, cyano C₁-C₄ alkyl, aminocarbonyl C₁-C₄ alkyl, hydroxy C₁-C₄ alkyl, and C₁-C₄ alkyloxy C₁-C₄ alkyl, and R₂ is selected from the group consisting of C₁-C₄ alkyl, halo C₁-C₄ alkyl, C₁-C₆ alkylcarbonyloxy C₁-C₄ alkyl, C₁-C₄ alkyloxycarbonyl C₁-C₄ alkyl, carboxy C₁-C₄ alkyl, cyano C₁-C₄ alkyl, aminocarbonyl C₁-C₄ alkyl, hydroxy C₁-C₄ alkyl, and C₁-C₄ alkyloxy C₁-C₄ alkyl, with the proviso that when R₁ is hydrogen, R₂ is selected from the group consisting of halo C₁-C₄ alkyl, C₁-C₄ alkyloxy C₁-C₄ alkyl, C₁-C₆ alkylcarbonyloxy C₁-C₄ alkyl, C₃-C₄ alkyloxycarbonyl C₁-C₄ alkyl, carboxy C₂-C₄ alkyl, cyano C₂-C₄ alkyl, aminocarbonyl C₂-C₄ alkyl, and hydroxy C₁-C₃ alkyl. Suitable compounds include those wherein R₁ is hydrogen or halo, and R₂ is C₁-C₆ alkylcarbonyloxy C₁-C₄ alkyl. Examples of suitable compounds include O⁶-benzyl-8-bromo-7-(pivaloyloxymethyl)guanine and O⁶-benzyl-7-(pivaloyloxymethyl)guanine.

The present invention additionally provides treatment methods, which are generally administered via pharmaceutical compositions comprising one or more of the O⁶-substituted compounds of the present invention. In particular, the present invention provides a method of enhancing the chemotherapeutic treatment of tumor cells in a mammal with an antineoplastic alkylating agent that


causes cytotoxic lesions at the O^6 -position of guanine, which method comprises administering to a mammal an effective amount of one or more of the aforescribed present inventive compounds of formulas I-V, and

5 administering to the mammal an effective amount of an antineoplastic alkylating agent that causes cytotoxic lesions at the O^6 -position of guanine. The present invention also includes the method of enhancing the chemotherapeutic treatment of tumor cells in a mammal with

10 an antineoplastic alkylating agent that causes cytotoxic lesions at the O^6 -position of guanine, which method comprises (i) administering to a mammal an effective amount of

(a) 8-aza- O^6 -benzylguanine


15



(b) a compound of the formula VI

(VI)

20



25

wherein R is a substituent selected from the group consisting of hydrogen, halo, C_1 - C_4 alkyl, C_1 - C_4 hydroxyalkyl, thiol, C_1 - C_4 alkylthio, trifluoromethylthio, C_1 - C_4 thioacyl, hydroxy, C_1 - C_4 alkoxy, trifluoromethoxy, methanesulfonyloxy, trifluoromethanesulfonyloxy, C_1 - C_4 acyloxy, amino, C_1 - C_4 aminoalkyl, C_1 - C_4 alkylamino, C_1 - C_4


dialkylamino, trifluoromethylamino, ditrifluoromethylamino, aminomethanesulfonyl, amino C₁-C₄ alkylcarbonyl, aminotrifluoromethylcarbonyl, formylamino, nitro, nitroso, C₁-C₄ alkyldiazo, C₅-C₆ aryldiazo, trifluoromethyl, C₁-C₄ haloalkyl, halomethyl, cyanomethyl, C₁-C₄ cyanoalkyl, cyano, C₁-C₄ alkyloxycarbonyl, C₁-C₄ alkylcarbonyl, phenyl, phenylcarbonyl, formyl, C₁-C₄ alkoxymethyl, phenoxyethyl, C₂-C₄ vinyl, C₂-C₄ ethynyl, and SO_nR' wherein n is 0, 1, 2, or 3 and R' is hydrogen, C₁-C₄ alkyl, amino, or phenyl, or

10 (c) 2,4-diamino-6-benzyl-s-triazine,
and (ii) administering to the mammal an effective amount of an antineoplastic alkylating agent which causes cytotoxic lesions at the O⁶-position of guanine.

Several O⁶-substituted compounds were tested for their ability to inactivate the human DNA repair protein, O⁶-alkylguanine-DNA alkyltransferase (AGT, alkyltransferase).


Two classes of compounds were identified as being significantly better than O⁶-benzylguanine (the prototype low-molecular-weight inactivator) in inactivating AGT in human HT29 colon tumor cell extracts. These were 8-substituted O⁶-benzylguanines bearing electron-withdrawing groups at the 8-position and 5-substituted 2,4-diamino-6-benzyloxypyrimidines bearing electron-withdrawing groups at the 5-position. The latter derivatives were also more effective than O⁶-benzylguanine in inactivating AGT in intact HT29 colon tumor cells. Both types of compounds were as effective or more effective than O⁶-benzylguanine in enhancing cell killing by 1,3-bis(2-chloroethyl)-1-nitrosourea (BCNU) of colon, breast and prostate cancer cells grown in culture. Provided 8-substituted O⁶-benzylguanine derivatives bearing electron-withdrawing substituents at the 8-position and 5-substituted 2,4-diamino-6-benzyloxypyrimidines bearing electron-withdrawing substituents at the 5-position do not exhibit undesirable toxicity, they should be superior to O⁶-benzylguanine as chemotherapeutic adjuvants for enhancing the effectiveness of antitumor drugs whose mechanism of action involves

modification of the σ^6 -position of DNA guanine residues. The specific compounds surveyed for AGT inactivating activity are illustrated below.




1a-e

- 1a, R=NH₂
- 1b, R=CH₃
- 1c, R=OH
- 1d, R=Br
- 1e, R=CF₃




2




3a-f

- 3a, R₁=H; R₂=NO₂
- 3b, R₁=NH₂; R₂=H
- 3c, R₁=NH₂; R₂=NH₂
- 3d, R₁=NH₂; R₂=NO
- 3e, R₁=NH₂; R₂=NO₂
- 3f, R₁=NH₂; R₂=Br




4a-f

- 4a, R₁=HO; R₂=H
- 4b, R₁=HO; R₂=OH
- 4c, R₁=F; R₂=H
- 4d, R₁=CH₃CONH; R₂=OH
- 4e, R₁=CH₃NH; R₂=H
- 4f, R₁=(CH₃)₂N; R₂=H




5a,b


- 5a, R = H
- 5b, R = CH₃





6





7



8a, R = CH₃
8b, R = CH₂OCOC(CH₃)₃



10a, R = CH₃
10b, R = CH₂OCOC(CH₃)₃



Preparations of the 8-substituted O^6 -benzylguanine derivatives 8-amino- O^6 -benzylguanine (1a) and O^6 -benzyl-8-methylguanine (1b) were accomplished by treating 2,8-diamino-6-chloropurine and 2-amino-6-chloro-8-methylpurine, respectively, with sodium benzyloxide in benzyl alcohol. O^6 -Benzyl-8-oxoguanine (O^6 -benzyl-7,8-dihydro-8-oxoguanine, 1c) was prepared by reacting 1,1'-carbonyldiimidazole with 2,4,5-triamino-6-benzyloxypyrimidine (Pfleiderer et al., *Chem. Ber.*, 94, 12-18 (1961)). For convenience, the compound is illustrated in the 8-hydroxy tautomeric form although it most probably exists in solution in the 8-keto form with a hydrogen attached to the 7-nitrogen atom. O^6 -Benzyl-8-bromoguanine (1d) was prepared by bromination of O^6 -benzylguanine. O^6 -Benzyl-8-trifluoromethylguanine (1e) was prepared by reacting 2-amino-6-chloro-8-trifluoromethylpurine with sodium benzyloxide in benzyl alcohol. 8-Aza- O^6 -benzylguanine (2) was prepared through nitrous acid treatment of 2,4,5-triamino-6-benzyloxypyrimidine. Compound 2 had been prepared previously by another route (Shealy et al., *J. Org. Chem.*, 27, 4518-4523 (1962)).

With respect to the pyrimidine derivatives (3a-f), 4-amino-6-benzyloxy-5-nitropyrimidine (3a) was prepared by treating 4-amino-6-chloro-5-nitropyrimidine (Boon et al., *J. Chem. Soc.*, 96-102 (1951)) with sodium benzyloxide in benzyl alcohol. Derivatives 3b-d were prepared by the method of Pfleiderer et al. (*Chem. Ber.*, 94, 12-18 (1961)).

2,4-Diamino-6-benzyloxy-5-nitropyrimidine (3e) and 2,4-diamino-6-benzyloxy-5-bromopyrimidine (3f) were prepared previously by Kosary et al. (*Acta Pharm. Hung.*, 49, 241-247 (1989)).

The purines, O^6 -benzylxanthine (4a) and O^6 -benzyluric acid (4b) were prepared by nitrous acid deamination of O^6 -benzylguanine and O^6 -benzyl-8-oxoguanine, respectively. N^2 -Acetyl- O^6 -benzyl-8-oxoguanine (N^2 -acetyl- O^6 -benzyl-7,8-dihydro-8-oxoguanine) (4d) was prepared through acetylation

of O^6 -benzyl-8-oxoguanine (1c). O^6 -Benzyl-2-fluorohypoxanthine (4c) was prepared previously by Robins and Robins (J. Org. Chem., 34, 2160-2163 (1969)). This material was treated with methylamine and dimethylamine to 5 produce O^6 -benzyl- N^2 -methylguanine (4e) and O^6 -benzyl- N^2,N^2 -dimethylguanine (4f), respectively.

Compounds 5a (2-amino-4-benzyloxy-5-nitropyrimidine) and 5b (2-amino-4-benzyloxy-6-methyl-5-nitropyrimidine) were prepared by treating 2-amino-4-chloro-5-nitropyrimidine and 2-amino-4-chloro-6-methyl-5-nitropyrimidine (Boon et al., J. Chem. Soc., 96-102 (1951)), respectively, with sodium benzyloxide in benzyl alcohol. Compound 6 (2,4-diamino-6-benzyloxy-s-triazine) was prepared previously under similar conditions 10 (Wakabayashi et al., Nippon Dojo-Hiryogaku Zasshi, 41, 193-200 (1970)). O^6 -Benzyl-8-trifluoromethyl-9-methylguanine (7) was prepared by treating the anion of 1e with methyl iodide in *N,N*-dimethylformamide.

Compound 8a was prepared by methylating the sodium 20 salt of 8-aza- O^6 -benzylguanine using methyl iodide as the methylating agent. Compounds 8b and 9 were prepared by the reaction of the sodium salt of 8-aza- O^6 -benzylguanine and chloromethyl pivalate. Compound 10a was prepared by direct bromination of O^6 -benzyl-9-methylguanine. Compounds 10b and 25 11 were prepared by the reaction of the sodium salt of O^6 -benzyl-8-bromoguanine and chloromethyl pivalate. Compound 12 was prepared by the reaction of the sodium salt of O^6 -benzylguanine and chloromethyl pivalate.

The ability of these compounds to inactivate the AGT 30 protein in HT29 human colon tumor cell extracts and in intact HT29 cells is summarized in Tables 1 and 2. The data represent the dose of compound required to produce 50% inactivation in cell-free extracts upon incubation for 30 min or in cells upon incubation for 4 hr.

Table 1. AGT-Inactivating Activity of 6-Benzylxopurine, 6(4)-Benzylxopyrimidine, and 6-Benzylx-o-*triazine* Derivatives

Compound	ED ₅₀ (mM) ^a	
	In HT29 cell-free extract	In HT29 cells
2,4-diamino-6-benzylx-5-nitrosopyrimidine (3d)	0.06	0.02
2,4-diamino-6-benzylx-5-nitropyrimidine (3e)	0.06	0.02
8-aza- <i>O</i> ⁶ -benzylguanine (2)	0.07	0.06
<i>O</i> ⁶ -benzyl-8-bromoguanine (1d)	0.08	0.06
<i>O</i> ⁶ -benzylguanine	0.2	0.05
<i>O</i> ⁶ -benzyl-8-methyl-guanine (1b)	0.3	0.1
<i>O</i> ⁶ -benzyl-8-oxoguanine (1c)	0.3	0.15
<i>O</i> ⁶ -benzyl-8-trifluoromethylguanine (1e)	0.4	0.25
2,4,5-triamino-6-benzylxopyrimidine (3c)	0.4	0.3
2-amino-4-benzylx-6-methyl-5-nitropyrimidine (5b)	0.4	0.06
2-amino-4-benzylx-5-nitropyrimidine (5a)	0.4	0.05
8-amino- <i>O</i> ⁶ -benzylguanine (1a)	0.7	2
2,4-diamino-6-benzylx-5-bromopyrimidine (3f)	2	0.8
2,4-diamino-6-benzylx-o- <i>triazine</i> (6)	4	1.0
2,4-diamino-6-benzylxopyrimidine (3b)	15	5
<i>O</i> ⁶ -benzyluric acid (4b)	25	45
4-amino-6-benzylx-5-nitropyrimidine (3a)	28	8
<i>O</i> ⁶ -benzyl-2-fluorohypoxanthine (4c)	48	12
<i>O</i> ⁶ -benzylxanthine (4a)	60	35
<i>N</i> ² -acetyl- <i>O</i> ⁶ -benzyl-8-oxoguanine (4d)	65	11
<i>O</i> ⁶ -benzyl- <i>N</i> ² -methylguanine (4e)	160	60
<i>O</i> ⁶ -benzyl- <i>N</i> ² , <i>N</i> ² -dimethylguanine (4f)	200	110

^a The effective dose required to produce 50% inactivation in cell-free extracts upon incubation for 30 min or in cells upon incubation for 4 hr. The values for *O*⁶-benzylguanine are from Moschel et al., *J. Med. Chem.*, 35, 4486-4491 (1992).

Within these series of compounds, O^6 -benzyl- N^2 -methyl- and O^6 -benzyl- N^2,N^2 -dimethylguanine (4e and 4f) were the least active agents exhibiting ED₅₀ values for inactivation of AGT in HT29 cell extracts of 160 and 200 mM, respectively. For comparison, the ED₅₀ value exhibited by O^6 -benzylguanine was 0.2 mM (Table 1). The other 2- and/or 8-substituted 6-benzylxopurines, N^2 -acetyl- O^6 -benzyl-8-oxoguanine (4d), O^6 -benzylxanthine (4a), O^6 -benzyl-2-fluorohypoxanthine (4c) and O^6 -benzyluric acid (4b), together with the substituted pyrimidines 4-amino-6-benzylxy-5-nitropyrimidine (3a) and 2,4-diamino-6-benzylxy-5-nitropyrimidine (3b), comprised a group of increasingly more active AGT inactivating agents exhibiting intermediate ED₅₀ values in the range of 65 to 15 mM. 2,4-Diamino-6-benzylxy-s-triazine (6) and 2,4-diamino-6-benzylxy-5-bromopyrimidine (3f) were considerably more active than 3b indicating that electron-withdrawing groups at the 5-position of a 2,4-diamino-6-benzylxy-5-nitropyrimidine derivative are positive contributors to efficient AGT inactivation. This is further emphasized by the very high activity exhibited by 2,4-diamino-6-benzylxy-5-nitrosopyrimidine (3d) and 2,4-diamino-6-benzylxy-5-nitropyrimidine (3e), which contain strongly electron-withdrawing nitroso and nitro substituents, respectively. These two derivatives are the most active AGT inactivators tested to date. The observation that 2-amino-4-benzylxy-5-nitropyrimidine (5a) is much more active than 3a indicates that a 2-amino group is critical for high activity for a 6(4)-benzylxy-5-nitropyrimidine derivative. An additional alkyl group at the 4(6)-position (e.g., as in 5b) does not enhance activity significantly over that for 5a although an amino group at the 4(6)-position significantly enhances activity. Thus, AGT inactivating activity increases substantially over the series 5a=5b<3d=3e. With these considerations in mind the activity of 2,4,5-triamino-6-benzylxy-5-nitropyrimidine (3c) seems exceptional and the reasons for its relatively high activity are unclear at present. It is also

significant that pyrimidines 5a and 5b are quite active in cells, which is not totally predicted by their corresponding activity in HT29 cell extracts.

All the O^6 -benzylguanine analogs 1a-d were much more active than the purines in the series 4a-f and the activity differences among 1a-d also reflect enhancements due to introduction of electron withdrawing groups. Thus, activity increased in the series 8-amino- O^6 -benzylguanine (1a) < O^6 -benzyl-8-oxoguanine (1c) < O^6 -benzyl-8-methylguanine (1b) < O^6 -benzyl-8-bromoguanine (1d) <8-aza- O^6 -benzylguanine (2). Indeed, derivatives 1d and 2 were essentially as active as pyrimidines 3d and 3e in cell-free extracts although 1d and 2 were somewhat less active in cells than expected from their activity in cell-free extracts.

The compounds listed in Table 2 also had AGT-inactivating activity in cell free extracts and in cells. The activity of 7, 8a, and 10a in cells is significantly higher than their activity in cell-free extracts. Thus the ratio of ED₅₀ values in cell-free extracts/intact cells is 1.6, 1.6, 1.1, respectively, for derivatives 1d, 1e, and 2 (Table 1). This ratio increases to 7.2, 6.3, and 6.3, respectively, for the corresponding methylated derivatives 10a, 7, and 8a. It is believed that the higher activity of the methylated derivatives in the cells is due to the fact that these compounds do not possess readily dissociable hydrogens in the imidazole portion of the purine ring system and therefore they can readily enter the cells as neutral molecules.

Table 2. AGT-Inactivating Activity of 7,8- and 8,9-Disubstituted O^6 -Benzylguanine Derivatives and Related Compounds

Compound	ED ₅₀ value (mM) ^a	
	In HT29 cell-free extract	In HT29 cells
O^6 -benzyl-8-trifluoromethyl-9-methylguanine (7)	2.5	0.4
8-aza- O^6 -benzyl-9-methylguanine (8a)	0.5	0.08
8-aza- O^6 -benzyl-9-(pivaloyloxymethyl)guanine (8b)	0.28 ^b	0.23
8-aza- O^6 -benzyl-7-(pivaloyloxymethyl)guanine (9)	0.11 ^c	0.16
O^6 -benzyl-8-bromo-9-methylguanine (10a)	1.9	0.25
O^6 -benzyl-8-bromo-9-(pivaloyloxymethyl)guanine (10b)	0.08 ^d	0.05
O^6 -benzyl-7-(pivaloyloxymethyl)guanine (12)	9 ^e	0.3
O^6 -benzyl-9-(pivaloyloxymethyl)guanine	3.1 ^{f,g}	0.3
O^6 -benzyl-9-methylguanine	2.6 ^g	0.4

^a The dose required to produce 50% inactivation in cell-free extracts upon incubation for 30 min. or in cells upon incubation for 4 hr. ^bED₅₀=4 with purified human AGT. ^cED₅₀=2 with purified human AGT. ^dED₅₀ > 100 with purified human AGT. ^eED₅₀ >> 100 with purified human AGT. ^fED₅₀ = 95 with purified human AGT. ^gData from Chae et al., *J. Med. Chem.*, 37, 342-347 (1994).

The ability of increasing concentrations of 1a-d, 2, 5 and 3c-e to enhance the killing of human HT29 colon cancer cells, DU-145 prostate cancer cells, and MCF-71 breast cancer cells by BCNU (40 mM) is shown in Tables 3, 4, and 5, respectively. The data reflect the number of cell colonies that result following exposure to AGT inactivator 10 alone or AGT inactivator 2 hr before exposure to BCNU as described in Dolan et al. (*Proc. Natl. Acad. Sci., U.S.A.*, 87, 5368-5372 (1990)). Data for O^6 -benzylguanine are included for comparison. As indicated, at 10 mM

concentrations, all the 8-substituted purines with the exception of 1a were as effective as O^6 -benzylguanine in enhancing the cytotoxicity of BCNU (40 mM); such treatment killed essentially all the tumor cells. Treatment of the 5 cells with the modified 8-substituted O^6 -benzylguanine alone or BCNU alone had no significant effect on cell colony number. The comparatively low activity of 1a in all but the breast cancer cells may reflect its poor transport into other tumor cell types or its rapid metabolic conversion to 10 an ineffective AGT inactivator. Its ineffective enhancement of BCNU cytotoxicity parallels its relatively poor AGT inactivating ability in colon tumor cells (Table 1).

For the pyrimidines tested, 2,4,5-triamino-6-benzylxypyrimidine (3c) was as effective as the 8-substituted O^6 -benzylguanine derivatives and O^6 -benzylguanine itself in enhancing BCNU toxicity although the nitroso- and nitropyrimidine derivatives (3d and 3e) were similarly effective at a 4-fold lower dose.

20

Table 3. Killing of HT-29 Colon Cancer Cells by BCNU Combined with AGT Inactivators

Inactivator	Inactivator Concentration (mM)	BCNU (mM)	Colony Formation per 1000 cells
None		None	435±63
None		40	442±34
O^6 -benzylguanine	10	None	431±33
	10	40	13±6
	2.5	40	38±15
	1	40	277±25
8-aza- O^6 -benzylguanine (2)	10	None	537±48
	10	40	2±1
	1	40	423±42
O^6 -benzyl-8-bromoguanine (1d)	10	None	401±22
	10	40	1±0
	1	40	299±30

O^6 -benzyl-8-oxoguanine (1c)	10 10 1	None 40 40	401±22 <1 221±15
O^6 -benzyl-8- methylguanine (1b)	10 10 1	None 40 40	513±76 <1 230±51
O^6 -benzyl-8-aminoguanine (1a)	10 10 1	None 40 40	504±30 430±41 475±26
2,4,5-triamino-6- benzyloxyprymidine (3c)	10 10 1	None 40 40	453±59 3±1 487±32
2,4-diamino-6-benzyloxy- 5-nitrosopyrimidine (3d)	2.5 2.5 1	None 40 40	528±64 <1 19±4
2,4-diamino-6-benzyloxy- 5-nitropyrimidine (3e)	2.5 2.5 1	None 40 40	438±25 <1 45±4

Table 4. Killing of DU-145 Prostate Cancer Cells by BCNU
Combined with AGT Inactivators

Inactivator	Inactivator Concentration (mM)	BCNU (mM)	Colony Formation per 1000 cells
None		None	453±81
None		40	394±76
O^6 -benzylguanine	10 10 1	None 40 40	462±68 28±5 299±18
8-aza- O^6 -benzylguanine (2)	10 10 1	None 40 40	452±72 28±5 248±21
O^6 -benzyl-8- bromoguanine (1d)	10 10 1	None 40 40	493±90 16±3 267±39
O^6 -benzyl-8-oxoguanine (1c)	10 10 1	None 40 40	379±34 34±3 329±43
O^6 -benzyl-8-	10 10	None 40	357±43 50±7

methylguanine (1b)	1	40	306±157
O^6 -benzyl-8-aminoguanine (1a)	10	None	380±36
	10	40	435±70
	1	40	295±45
2,4,5-triamino-6-benzyloxypyrimidine (3c)	10	None	429±101
	10	40	57±7
	1	40	378±60
2,4-diamino-6-benzyloxy-5-nitrosopyrimidine (3d)	2.5	None	403±35
	2.5	40	7±3
	1	40	25±4
	0.25	40	192±17
2,4-diamino-6-benzyloxy-5-nitropyrimidine (3e)	2.5	None	407±80
	2.5	40	9±2
	1	40	59±6
	0.25	40	129±26

Table 5. Killing of MCF-71 Breast Cancer Cells by BCNU Combined with AGT Inactivators

Inactivator	Inactivator Concentration (mM)	BCNU (mM)	Colony Formation per 1000 cells
None		None	426±78
None		40	364±60
O^6 -benzylguanine	10	None	455±63
	10	40	4±2
	2.5	40	12±6
8-aza- O^6 -benzylguanine (2)	10	None	483±27
	10	40	2±1
O^6 -benzyl-8-bromoguanine (1d)	10	None	380±109
	10	40	3±1
	2.5	40	4±3
O^6 -benzyl-8-oxoguanine (1c)	10	None	522±78
	10	40	4±2
O^6 -benzyl-8-methylguanine (1b)	10	None	376±76
	10	40	2±1
O^6 -benzyl-8-aminoguanine (1a)	10	None	432±36
	10	40	95±8

2,4,5-triamino-6-	10	None	448±55
benzyloxypyrimidine (3c)	10	40	12±4
2,4-diamino-6-benzyloxy-	2.5	None	447±87
5-nitrosopyrimidine (3d)	2.5	40	2±1
2,4-diamino-6-benzyloxy-	2.5	None	314±49
5-nitropyrimidine (3e)	2.5	40	2±1

Although the human alkyltransferase is very sensitive to inactivation by O^6 -benzylguanine and the various compounds described above, a number of mutants have been generated that are resistant to O^6 -benzylguanine (Crone and Pegg, *Cancer Res.*, 53, 4750-4753 (1993)). This resistance is probably caused by a reduction in the space surrounding the active site of the alkyltransferase, which limits the access to O^6 -benzylguanine. These mutants are produced by single base changes in the alkyltransferase DNA-coding sequence causing changes in one or two amino acids in the alkyltransferase (Crone and Pegg, *Cancer Res.*, 53, 4750-4753 (1993)). Thus, as indicated in Table 6, changing the proline residue at position 140 to alanine (protein P140A) or the glycine residue at position 156 to an alanine (protein G156A) causes a 20-fold and a 240-fold increase in resistance to O^6 -benzylguanine, respectively. The alkyltransferase containing an arginine in place of a proline at residue 138 together with an arginine in place of a proline at residue 140 (protein P138A/P140A) is 88-fold more resistant to inactivation by O^6 -benzylguanine. It is possible that such resistant mutants will arise or be selected for in tumors under the selective pressure generated by treatment with O^6 -benzylguanine plus an alkylating agent. More potent inhibitors and/or those of a smaller size that are better able to fit into the space of the active site of the mutant alkyltransferase can be used to advantage to overcome this resistance.

Table 6. Inhibition of Mutant Alkyltransferase Proteins by O^6 -Benzylguanine or 2,4-Diamino-6-benzyloxy-5-nitrosopyrimidine

Protein	ED ₅₀ value (mM) ^a	
	O^6 -benzylguanine	2,4-diamino-6-benzyloxy-5-nitrosopyrimidine
Control	0.25	0.05
P140A	5	0.1
P138A/P140A	22	0.3
G156A	60	1

^aThe concentration needed to inactivate 50% of the activity in 30 minutes.

As shown in Table 6, 2,4-diamino-6-benzyloxy-5-nitrosopyrimidine (3d) was 50 to 60 times better at inactivating the mutant alkyltransferases than O^6 -benzylguanine. Doses of 2,4-diamino-6-benzyloxy-5-nitrosopyrimidine leading to intracellular concentrations greater than 5 mM will therefore be effective at inactivating such resistant alkyltransferases. Concentrations greater than 200 mM of O^6 -benzylguanine would be needed to get such inactivation, and these are much more than can be achieved with this compound in current formulations. However, 8-substituted O^6 -benzylguanine derivatives that are significantly more potent than O^6 -benzylguanine may be useful in inactivating mutant alkyltransferases provided their required intracellular concentrations can be achieved. These data for mutant alkyltransferase inactivation and the data presented earlier indicate that pyrimidine derivatives bearing electron-withdrawing groups at the 5-position as well as substituted O^6 -benzylguanine derivatives bearing electron-withdrawing groups at the 8-position are superior to O^6 -benzylguanine for use as adjuvants in chemotherapy with agents whose mechanism of action, like that of BCNU,

involves modification of the O⁶-position of DNA guanine residues.

Other 8-substituted O⁶-benzylguanine derivatives bearing electron-withdrawing 8-substituents (e.g., NO₂) are 5 readily available. For example, O⁶-benzyl-8-nitroguanine could be prepared by treatment of 8-nitroguanine (Jones and Robins, *J. Am. Chem. Soc.*, 82, 3773-3779 (1960)) with phosphorus oxychloride to produce 2-amino-6-chloro-8-nitropurine which when treated with sodium benzyloxide in 10 benzyl alcohol would produce the desired O⁶-benzyl-8-nitroguanine.

Additional 2,4-diamino-6-benzyloxypyrimidine derivatives bearing electron-withdrawing groups other than halogen or nitro groups (e.g., formyl or cyano groups) 15 could also be readily prepared. 2,4-Diamino-5-formyl-6-hydroxypyrimidine, a known compound (Delia and Otteman, *Heterocycles*, 20, 1805-1809 (1983)), can be treated with phosphorus oxychloride to produce a 2,4-diamino-6-chloro-5-formylpyrimidine intermediate, which on treatment with 20 sodium benzyloxide in benzyl alcohol produces 2,4-diamino-6-benzyloxy-5-formylpyrimidine. Treatment of the formyl pyrimidine with hydroxylamine affords 2,4-diamino-6-benzyloxy-5-cyanopyrimidine. The preparation of a large number of 5-substituted 6(4)-benzyloxypyrimidines or 8-25 substituted O⁶-benzylguanine derivatives is possible for those skilled in the art of synthesis of heterocyclic aromatic compounds (D.J. Brown, "The Pyrimidines," in *The Chemistry of Heterocyclic Compounds*, Vol. 16, A. Weissberger, Ed., Wiley Interscience, New York, 1962; D.J. 30 Brown, "The Pyrimidines," Supplement I, in *The Chemistry of Heterocyclic Compounds*, Vol. 16, A. Weissberger and E.C. Taylor, Eds., Wiley Interscience, New York, 1970; J.H. Lister, "Fused Pyrimidines Part II Purines," in *The Chemistry of Heterocyclic Compounds*, Vol. 24 Part II, A. 35 Weissberger and E.C. Taylor, Eds., Wiley Interscience, New York, 1971).

Because many 9-substituted O^6 -benzylguanine derivatives exhibit excellent AGT inactivation properties (Moschel et al., *J. Med. Chem.*, 35, 4486-4491 (1992); Chae et al., *J. Med. Chem.*, 37, 342-347 (1994)), 8,9-disubstituted analogs are expected to be similarly active. These can be readily prepared by reacting the anion of 8-substituted O^6 -benzylguanines (e.g., 1a-e) or the anion of 8-aza- O^6 -benzylguanine (2) with any of the range of compounds already described (Moschel et al., *J. Med. Chem.*, 35, 4486-4491 (1992); Chae et al., *J. Med. Chem.*, 37, 342-347 (1994)) to produce a mixture of isomeric 7,8- and 8,9-disubstituted O^6 -benzylguanine derivatives. The desired 8,9-disubstituted derivative can be isolated and purified by silica gel column chromatography as already described (Moschel et al., *J. Med. Chem.*, 35, 4486-4491 (1992); Chae et al., *J. Med. Chem.*, 37, 342-347 (1994)). Compound 7 was prepared by treating the anion of compound 1e with methyl iodide in *N,N*-dimethylformamide. Compounds 8-12 were prepared using similar procedures.

The O^6 -substituted compounds of the present invention can be administered in any suitable manner to a mammal for the purpose of enhancing the chemotherapeutic treatment of a particular cancer. Although more than one route can be used to administer a particular compound, a particular route can provide a more immediate and more effective reaction than another route. Accordingly, the described methods provided herein are merely exemplary and are in no way limiting.

Generally, the O^6 -substituted compounds of the present invention as described above will be administered in a pharmaceutical composition to an individual afflicted with a cancer. Those undergoing or about to undergo chemotherapy can be treated with the O^6 -substituted compounds separately or in conjunction with other treatments, as appropriate. In therapeutic applications, compositions are administered to a patient in an amount sufficient to elicit an effective depression of AGT

activity thereby potentiating the cytotoxicity of the aforescribed chemotherapeutic treatment. An amount adequate to accomplish this is defined as a "therapeutically effective dose," which is also an "AGT inactivating effective amount." Amounts effective for a therapeutic or prophylactic use will depend on, e.g., the stage and severity of the disease being treated, the age, weight, and general state of health of the patient, and the judgment of the prescribing physician. The size of the dose will also be determined by the O^6 -substituted compound selected, method of administration, timing and frequency of administration as well as the existence, nature, and extent of any adverse side-effects that might accompany the administration of a particular O^6 -substituted compound and the desired physiological effect. It will be appreciated by one of skill in the art that various disease states may require prolonged treatment involving multiple administrations, perhaps using a series of different AGT inactivators and/or chemotherapeutic agents in each or various rounds of administration.

Suitable chemotherapeutic agents usefully administered in coordination with the O^6 -substituted compounds of the present invention include alkylating agents, such as chloroethylating and methylating agents. Such agents may be administered using conventional techniques such as those described in Wasserman et al., Cancer, 36, pp. 1258-1268 (1975), and Physicians' Desk Reference, 48th ed., Edward R. Barnhart publisher (1994). For example, 1,3-bis(2-chloroethyl)-1-nitrosourea (carmustine or BCNU, Bristol-Myers, Evansville, IN) may be administered intravenously at a dosage of from about 150 to 200 mg/m² every six weeks. Another alkylating agent, 1-(2-chloroethyl)-3-cyclohexyl-1-nitrosourea (lomustine or CCNU, Bristol-Myers), may be administered orally at a dosage of about 130 mg/m² every six weeks. Other alkylating agents may be administered in appropriate dosages via appropriate routes of administration known to skilled medical practitioners.

Suitable doses and dosage regimens can be determined by conventional range-finding techniques known to those of ordinary skill in the art. Generally, treatment is initiated with smaller dosages that are less than the optimum dose of the compound. Thereafter, the dosage is increased by small increments until the optimum effect under the circumstances is reached. The present inventive method typically will involve the administration of about 0.1 mg to about 50 mg of one or more of the compounds described above per kg body weight of the individual. For a 70 kg patient, dosages of from about 10 mg to about 200 mg of O^6 -substituted compound would be more commonly used, possibly followed by further lesser dosages from about 1 mg to about 1 mg of O^6 -substituted compound over weeks to months, depending on a patient's physiological response, as determined by measuring cancer-specific antigens or other measurable parameters related to the tumor load of a patient.

It must be kept in mind that the compounds and compositions of the present invention generally are employed in serious disease states, that is, life-threatening or potentially life-threatening situations. In such cases, in view of the minimization of extraneous substances and the relative nontoxic nature of the O^6 -substituted compounds, it is possible and may be felt desirable by the treating physician to administer substantial excesses of these O^6 -substituted compounds.

Single or multiple administrations of the compounds can be carried out with dose levels and pattern being selected by the treating physician. In any event, the pharmaceutical formulations should provide a quantity of AGT-inactivating compounds of the invention sufficient to effectively enhance the cytotoxic impact of the chemotherapy.

The pharmaceutical compositions for therapeutic treatment are intended for parenteral, topical, oral or local administration and generally comprise a

pharmaceutically acceptable carrier and an amount of the active ingredient sufficient to reduce, and preferably prevent, the activity of the AGT protein. The carrier may be any of those conventionally used and is limited only by 5 chemico-physical considerations, such as solubility and lack of reactivity with the compound, and by the route of administration.

Examples of pharmaceutically acceptable acid addition salts for use in the present inventive pharmaceutical 10 composition include those derived from mineral acids, such as hydrochloric, hydrobromic, phosphoric, metaphosphoric, nitric and sulfuric acids, and organic acids, such as tartaric, acetic, citric, malic, lactic, fumaric, benzoic, glycolic, gluconic, succinic, *p*-toluenesulphonic acids, and 15 arylsulphonic, for example.

The pharmaceutically acceptable excipients described herein, for example, vehicles, adjuvants, carriers or diluents, are well-known to those who are skilled in the art and are readily available to the public. It is 20 preferred that the pharmaceutically acceptable carrier be one that is chemically inert to the active compounds and one that has no detrimental side effects or toxicity under the conditions of use. Such pharmaceutically acceptable excipients preferably include saline (e.g., 0.9% saline), 25 Cremophor EL (which is a derivative of castor oil and ethylene oxide available from Sigma Chemical Co., St. Louis, MO) (e.g., 5% Cremophor EL/5% ethanol/90% saline, 10% Cremophor EL/90% saline, or 50% Cremophor EL/50% ethanol), propylene glycol (e.g., 40% propylene glycol/10% 30 ethanol/50% water), polyethylene glycol (e.g., 40% PEG 400/60% saline), and alcohol (e.g., 40% *t*-butanol/60% water). The most preferred pharmaceutical excipient for use in conjunction with the present invention is polyethylene glycol, such as PEG 400, and particularly a 35 composition comprising 40% PEG 400 and 60% water or saline.

The choice of excipient will be determined in part by the particular O^6 -substituted compound chosen, as well as by

the particular method used to administer the composition. Accordingly, there is a wide variety of suitable formulations of the pharmaceutical composition of the present invention.

5 The following formulations for oral, aerosol, parenteral, subcutaneous, intravenous, intraarterial, intramuscular, interperitoneal, rectal, and vaginal administration are merely exemplary and are in no way limiting.

10 The pharmaceutical compositions can be administered parenterally, e.g., intravenously, intraarterially, subcutaneously, intradermally, or intramuscularly. Thus, the invention provides compositions for parenteral administration that comprise a solution of the O^6 -

15 substituted compound dissolved or suspended in an acceptable carrier suitable for parenteral administration, including aqueous and non-aqueous, isotonic sterile injection solutions.

Overall, the requirements for effective pharmaceutical

20 carriers for parenteral compositions are well known to those of ordinary skill in the art. See Pharmaceutics and Pharmacy Practice, J.B. Lippincott Company, Philadelphia, PA, Bunker and Chalmers, eds., pages 238-250 (1982), and ASHP Handbook on Injectable Drugs, Toissel, 4th ed., pages

25 622-630 (1986). Such solutions can contain anti-oxidants, buffers, bacteriostats, and solutes that render the formulation isotonic with the blood of the intended recipient, and aqueous and non-aqueous sterile suspensions that can include suspending agents, solubilizers,

30 thickening agents, stabilizers, and preservatives. The compound may be administered in a physiologically acceptable diluent in a pharmaceutical carrier, such as a sterile liquid or mixture of liquids, including water, saline, aqueous dextrose and related sugar solutions, an

35 alcohol, such as ethanol, isopropanol, or hexadecyl alcohol, glycols, such as propylene glycol or polyethylene glycol, dimethylsulfoxide, glycerol ketals, such as 2,2-

dimethyl-1,3-dioxolane-4-methanol, ethers, such as poly(ethyleneglycol) 400, an oil, a fatty acid, a fatty acid ester or glyceride, or an acetylated fatty acid glyceride with or without the addition of a

5 pharmaceutically acceptable surfactant, such as a soap or a detergent, suspending agent, such as pectin, carbomers, methylcellulose, hydroxypropylmethylcellulose, or carboxymethylcellulose, or emulsifying agents and other pharmaceutical adjuvants.

10 Oils useful in parenteral formulations include petroleum, animal, vegetable, or synthetic oils. Specific examples of oils useful in such formulations include peanut, soybean, sesame, cottonseed, corn, olive, petrolatum, and mineral. Suitable fatty acids for use in

15 parenteral formulations include oleic acid, stearic acid, and isostearic acid. Ethyl oleate and isopropyl myristate are examples of suitable fatty acid esters.

Suitable soaps for use in parenteral formulations include fatty alkali metal, ammonium, and triethanolamine

20 salts, and suitable detergents include (a) cationic detergents such as, for example, dimethyl dialkyl ammonium halides, and alkyl pyridinium halides, (b) anionic detergents such as, for example, alkyl, aryl, and olefin sulfonates, alkyl, olefin, ether, and monoglyceride

25 sulfates, and sulfosuccinates, (c) nonionic detergents such as, for example, fatty amine oxides, fatty acid alkanolamides, and polyoxyethylenepolypropylene copolymers, (d) amphoteric detergents such as, for example, alkyl-*b*-aminopropionates, and 2-alkyl-imidazoline quaternary

30 ammonium salts, and (e) mixtures thereof.

The parenteral formulations typically will contain from about 0.5% to about 25% by weight of the active ingredient in solution. Preservatives and buffers may be used. In order to minimize or eliminate irritation at the

35 site of injection, such compositions may contain one or more nonionic surfactants having a hydrophile-lipophile balance (HLB) of from about 12 to about 17. The quantity

of surfactant in such formulations will typically range from about 5% to about 15% by weight. Suitable surfactants include polyethylene sorbitan fatty acid esters, such as sorbitan monooleate and the high molecular weight adducts 5 of ethylene oxide with a hydrophobic base, formed by the condensation of propylene oxide with propylene glycol. The parenteral formulations can be presented in unit-dose or multi-dose sealed containers, such as ampules and vials, and can be stored in a freeze-dried (lyophilized) condition 10 requiring only the addition of the sterile liquid excipient, for example, water, for injections, immediately prior to use. Extemporaneous injection solutions and suspensions can be prepared from sterile powders, granules, and tablets of the kind previously described.

15 Topical formulations, including those that are useful for transdermal drug release, are well-known to those of skill in the art and are suitable in the context of the present invention for application to skin.

Formulations suitable for oral administration require 20 extra considerations considering the peptidyl and/or carbohydrate nature of some of the O^6 -substituted compounds of the present invention and the likely breakdown thereof if such compounds are administered orally without protecting them from the digestive secretions of the 25 gastrointestinal tract. Such a formulation can consist of (a) liquid solutions, such as an effective amount of the compound dissolved in diluents, such as water, saline, or orange juice; (b) capsules, sachets, tablets, lozenges, and troches, each containing a predetermined amount of the 30 active ingredient, as solids or granules; (c) powders; (d) suspensions in an appropriate liquid; and (e) suitable emulsions. Liquid formulations may include diluents, such as water and alcohols, for example, ethanol, benzyl alcohol, and the polyethylene alcohols, either with or 35 without the addition of a pharmaceutically acceptable surfactant, suspending agent, or emulsifying agent.

Capsule forms can be of the ordinary hard- or soft-shelled

gelatin type containing, for example, surfactants, lubricants, and inert fillers, such as lactose, sucrose, calcium phosphate, and corn starch. Tablet forms can include one or more of lactose, sucrose, mannitol, corn 5 starch, potato starch, alginic acid, microcrystalline cellulose, acacia, gelatin, guar gum, colloidal silicon dioxide, croscarmellose sodium, talc, magnesium stearate, calcium stearate, zinc stearate, stearic acid, and other excipients, colorants, diluents, buffering agents, 10 disintegrating agents, moistening agents, preservatives, flavoring agents, and pharmacologically compatible excipients. Lozenge forms can comprise the active ingredient in a flavor, usually sucrose and acacia or tragacanth, as well as pastilles comprising the active 15 ingredient in an inert base, such as gelatin and glycerin, or sucrose and acacia, emulsions, gels, and the like containing, in addition to the active ingredient, such excipients as are known in the art.

The O^6 -substituted compounds of the present 20 invention, alone or in combination with other suitable components, can be made into aerosol formulations to be administered via inhalation. The compounds are preferably supplied in finely divided form along with a surfactant and propellant. Typical percentages of active compound are 25 0.01%-20% by weight, preferably 1%-10%. The surfactant must, of course, be nontoxic, and preferably soluble in the propellant. Representative of such surfactants are the esters or partial esters of fatty acids containing from 6 to 22 carbon atoms, such as caproic, octanoic, lauric, 30 palmitic, stearic, linoleic, linolenic, olesteric and oleic acids with an aliphatic polyhydric alcohol or its cyclic anhydride. Mixed esters, such as mixed or natural glycerides may be employed. The surfactant may constitute 0.1%-20% by weight of the composition, preferably 0.25-5%. 35 The balance of the composition is ordinarily propellant. A carrier can also be included as desired, e.g., lecithin for intranasal delivery. These aerosol formulations can be

placed into acceptable pressurized propellants, such as dichlorodifluoromethane, propane, nitrogen, and the like. They also may be formulated as pharmaceuticals for non-pressured preparations, such as in a nebulizer or an 5 atomizer. Such spray formulations may be used to spray mucosa.

Additionally, the compounds and polymers useful in the present inventive methods may be made into suppositories by mixing with a variety of bases, such as emulsifying bases 10 or water-soluble bases. Formulations suitable for vaginal administration may be presented as pessaries, tampons, creams, gels, pastes, foams, or spray formulas containing, in addition to the active ingredient, such carriers as are known in the art to be appropriate.

15 The concentration of the O^6 -substituted compounds of the present invention in the pharmaceutical formulations can vary widely, i.e., from less than about 1%, usually at or at least about 10%, to as much as 20% to 50% or more by weight, and will be selected primarily by fluid volumes, 20 viscosities, etc., in accordance with the particular mode of administration selected.

Thus, a typical pharmaceutical composition for intravenous infusion could be made up to contain 250 ml of sterile Ringer's solution, and 100 mg of the O^6 -substituted 25 compound. Actual methods for preparing parenterally administrable compounds will be known or apparent to those skilled in the art and are described in more detail in, for example, Remington's Pharmaceutical Science (17th ed., Mack Publishing Company, Easton, PA, 1985).

30 It will be appreciated by one of ordinary skill in the art that, in addition to the aforescribed pharmaceutical compositions, the O^6 -substituted compounds of the present inventive method may be formulated as inclusion complexes, such as cyclodextrin inclusion complexes, or liposomes. 35 Liposomes serve to target the compounds to a particular tissue, such as lymphoid tissue or cancerous hepatic cells. Liposomes can also be used to increase the half-life of

the O^6 -substituted compound. Liposomes useful in the present invention include emulsions, foams, micelles, insoluble monolayers, liquid crystals, phospholipid dispersions, lamellar layers and the like. In these 5 preparations, the O^6 -substituted compound to be delivered is incorporated as part of a liposome, alone or in conjunction with a suitable chemotherapeutic agent. Thus, liposomes filled with a desired O^6 -substituted compound of the invention can be directed to the site of a specific tissue 10 type, hepatic cells, for example, where the liposomes then deliver the selected chemotherapeutic-enhancement compositions. Liposomes for use in the invention are formed from standard vesicle-forming lipids, which generally include neutral and negatively charged 15 phospholipids and a sterol, such as cholesterol. The selection of lipids is generally guided by consideration of, for example, liposome size and stability of the liposomes in the blood stream. A variety of methods are available for preparing liposomes, as described in, for 20 example, Szoka et al., *Ann. Rev. Biophys. Bioeng.*, 9, 467 (1980), and U.S. Patents 4,235,871, 4,501,728, 4,837,028, and 5,019,369. For targeting to the cells of a particular tissue type, a ligand to be incorporated into the liposome can include, for example, antibodies or fragments thereof 25 specific for cell surface determinants of the targeted tissue type. A liposome suspension containing an O^6 -substituted compound may be administered intravenously, locally, topically, etc. in a dose that varies according to the mode of administration, the O^6 -substituted compound 30 being delivered, the stage of disease being treated, etc.

While the efficacy of the O^6 -substituted compounds of the present invention has been demonstrated with respect to particular types of cancerous cells, e.g., colon, prostate, and breast cancer cells, the present invention has 35 applicability to the treatment of any type of cancer capable of being treated with an antineoplastic alkylating agent which causes cytotoxic lesions at the O^6 -position of

guanine. Such cancers include, for example, colon tumors, prostate tumors, brain tumors, lymphomas, leukemias, breast tumors, ovarian tumors, lung tumors, Wilms' tumor, rhabdomyosarcoma, multiple myeloma, stomach tumors, soft-tissue sarcomas, Hodgkin's disease, and non-Hodgkin's lymphomas.

Similarly, in view of the mode of action of the O^6 -substituted compounds of the present invention, such compounds can be used in conjunction with any type of antineoplastic alkylating agent which causes cytotoxic lesions at the O^6 -position of guanine. Such antineoplastic alkylating agents include, for example, chloroethylating agents (e.g. chloroethylnitrosoureas and chloroethyltriazines) and monofunctional alkylating agents such as Streptozotocin, Procarbazine, Dacarbazine, and Temozolomide.

Among the chloroethylating agents, the most frequently used chemotherapeutic drugs are 1-(2-chloroethyl)-3-cyclohexyl-1-nitrosourea (CCNU, lomustine), 1,3-bis(2-chloroethyl)-1-nitrosourea (BCNU, carmustine), 1-(2-chloroethyl)-3-(4-methylcyclohexyl)-1-nitrosourea (MeCCNU, semustine), and 1-(2-chloroethyl)-3-(4-amino-2-methyl-5-pyrimidinyl)methyl-1-nitrosourea (ACNU). These agents have been used clinically against tumors of the central nervous system, multiple myeloma, melanoma, lymphoma, gastrointestinal tumors, and other solid tumors (Colvin and Chabner, *Alkylating Agents*. In: *Cancer Chemotherapy: Principles and Practice*, Chabner and Collins, eds., Lippincott, Philadelphia, pp. 276-313 (1990); McCormick and McElhinney, *Eur. J. Cancer*, 26, 207-221 (1990)). Chloroethylating agents currently under development with fewer side effects are 1-(2-chloroethyl)-3-(2-hydroxyethyl)-1-nitrosourea (HECNU), 2-chloroethyl-methylsulfonylmethanesulfonate (Clomesone), and 1-[N-(2-chloroethyl)-N-nitrosoureido]ethylphosphonic acid diethyl ester (Fotemustine) (Colvin and Chabner, *Alkylating Agents*. In: *Cancer Chemotherapy: Principles and Practice*, Chabner

and Collins, eds., Lippincott, Philadelphia, pp. 276-313 (1990); McCormick and McElhinney, *Eur. J. Cancer*, 26, 207-221 (1990)). Methylating chemotherapeutic agents include Streptozotocin (2-deoxy-2-(3-methyl-3-nitrosoureido)-D-5 glucopyranose), Procarbazine (N-(1-methylethyl)-4-[(2-methylhydrazino)methyl]benzamide), Dacarbazine or DTIC (5-(3,3-dimethyl-1-triazenyl)-1H-imidazole-4-carboxamide), and Temozolomide (8-carbamoyl-3-methylimidazo[5,1-d]-1,2,3,5-tetrazine-4-(3H)-one). Temozolomide is active against 10 malignant melanomas, brain tumors, and mycosis fungoides. Streptozotocin is effective against pancreatic tumors. Procarbazine is used to treat Hodgkin's disease and brain tumors, and DTIC is used in treatment of melanoma and lymphomas (Colvin and Cabner, *Alkylating Agents. In: Cancer Chemotherapy: Principles and Practice*, Chabner and Collins, eds., Lippincott, Philadelphia, pp. 276-313 (1990); Longo, *Semin. Concol.*, 17, 716-735 (1990)).

The examples set forth below describe the syntheses of the aforescribed compounds. As regards the methods and 20 materials set forth in these examples, ¹H-NMR spectra were recorded on a VarianTM VXR 500S spectrometer equipped with Sun 2/110 data stations or a VarianTM XL 200 instrument interfaced to an Advanced data system. Samples were dissolved in DMSO-d₆ with Me₄Si as an internal standard. EI 25 mass spectra were obtained on a reversed geometry VG MicromassTM ZAB-2F spectrometer interfaced to a VG 2035 data system. Elemental analyses were performed by Galbraith Laboratories, Inc., Knoxville, TN.

Most of the reagents and solvents were from Aldrich 30 Chemical Co., Inc., Milwaukee, WI. 8-Aza-*O*⁶-benzylguanine (2) (Shealy et al., *J. Org. Chem.*, 27, 4518-4523 (1962)), 2,4-diamino-6-benzyloxypyrimidine (3b) (Pfleiderer and Lohrmann, *Chem. Ber.*, 94, 12-18 (1961)), 2,4,5-triamino-6-benzyloxypyrimidine (3c) (Pfleiderer and Lohrmann, *Chem. Ber.*, 94, 12-18 (1961)), 2,4-diamino-6-benzyloxy-5-35 nitrosopyrimidine (3d) (Pfleiderer and Lohrmann, *Chem. Ber.*, 94, 12-18 (1961)), 2,4-diamino-6-benzyloxy-5-

nitropyrimidine (3e) (Kosary et al., *Acta Pharm. Hung.*, 49, 241-247 (1989)), 2,4-diamino-6-benzyloxy-5-bromopyrimidine (3f) (Kosary et al., *Acta Pharm. Hung.*, 49, 241-247 (1989)), 4-amino-6-benzyloxy-5-nitropyrimidine (3a) and O^6 -benzyl-2-fluorohypoxanthine (4c) (Robins and Robins, *J. Org. Chem.*, 34, 2160-2163 (1969)) were prepared previously. Alternative synthetic methods are provided below for some of these compounds together with spectroscopic data not provided previously. AGT inactivation studies were carried out as described in Moschel et al., *J. Med. Chem.*, 35, 4486-4491 (1992). Cell killing experiments involving various AGT inactivators in combination with BCNU were carried out as in Dolan, et al. (*Proc. Natl. Acad. Sci. U.S.A.*, 87, 5368-5372 (1990)). Cells were treated for 2 h with AGT inactivator prior to exposure to BCNU.

Example 1: 2,8-Diamino-6-chloropurine

A suspension of 8-aminoguanine (Fischer, *Z. Physiol. Chem.*, 60, 69 (1909); Beaman et al., in Zorbach and Tipson, *Synthetic Procedures in Nucleic Acid Chemistry*, Vol. 1, pp 41 - 43, John Wiley & Sons, New York, 1968) (3.0 g, 18.1 mmol) in phosphorus oxychloride (90 mL) and *N,N*-diethylaniline (3 mL) was refluxed for 30 min and the excess phosphorus oxychloride was evaporated under reduced pressure. Ice (20 g) was added slowly to the resulting solution and the pH was adjusted to 6 with a concentrated aqueous sodium hydroxide solution. A yellow solid formed and was collected by filtration, washed with water, and dried to give a green solid. Crystallization from water with charcoal treatment produced 2,8-diamino-6-chloropurine as a white solid: yield, 2.11 g (63%); mp >275 °C dec.; ^1H NMR δ 6.09 (s, 2 H, NH₂, exchange with D₂O), 6.71 (s, 2 H, NH₂, exchange with D₂O); MS (EI) calcd. *m/z* for C₅H₅N₆³⁵Cl 184.0264, found 184.0266; calcd. *m/z* C₅H₅N₆³⁷Cl 186.0235, found 186.0237.

Example 2: 8-Amino- O^6 -benzylguanine (1a)

2,8-Diamino-6-chloropurine (0.9 g, 4.9 mmol) was added to the solution of sodium (0.22 g, 10 mmol) in benzyl alcohol (9.0 mL). The solution was heated in a 130 °C oil bath for 5 h, and was poured into water (100 mL) with constant stirring for 10 min. Undissolved solid was removed by filtration and the filtrate was neutralized with glacial acetic acid. The solution was mixed with methanol (100 mL), and half of the aqueous methanol solution was loaded on a 3 x 80 cm SephadexTM LH-20 column eluted with methanol/water (1:1) at 1 mL/min. Column eluent was continuously monitored at 280 nm and fractions (10 mL) were collected. The remainder of the reaction mixture in MeOH/H₂O was chromatographed separately under identical conditions. The desired product eluted in fractions 100 - 130. Evaporation of solvent from the pooled fractions 100-130 from both chromatographic runs afforded analytically pure 1a: yield, 0.26 g (21%); mp 269 - 271 °C dec.; UV (pH 1) λ_{max} 241 nm ($\epsilon = 0.699 \times 10^4$), 300 (1.109×10^4); (pH 6.9) 250 (sh) (0.447×10^4), 292 (1.027×10^4); (pH 13) 255 (sh) (0.355×10^4), 295 (0.932×10^4); ¹H NMR δ 5.41 (s, 2 H, ArCH₂), 5.70 (s, 2 H, NH₂, exchange with D₂O), 6.18 (s, 2 H, NH₂, exchange with D₂O), 7.25-7.55 (m, 5 H, ArH), 11.1 (br s, 1 H, NH, exchanges with D₂O); MS (EI) calcd. m/z for C₁₂H₁₂N₆O 256.1072, found 256.1059; Anal. (C₁₂H₁₂N₆O) C, H, N.

Example 3: 2-Amino-6-chloro-8-methylpurine

A suspension of 8-methylguanine (Daves et al., *J. Am. Chem. Soc.*, 82, 2633 - 2640 (1960)) (1.0 g, 6.1 mmol) in phosphorous oxychloride (30 mL) and *N,N*-diethylaniline (1 mL) was refluxed for 3 h. The excess phosphorous oxychloride was evaporated under reduced pressure. The resulting brown oil was dissolved in ice-water and was neutralized with a concentrated aqueous NaOH solution. After evaporation of the solvent, the solid residue was suspended in 70 mL of H₂O. Undissolved solid was filtered

off, and the filtrate was loaded on a 3 x 80 cm Sephadex LH-20 column eluted with methanol/water (1:1) at 1 mL/min. Column eluent was continuously monitored at 280 nm and fractions (10 mL) were collected. Evaporation of pooled 5 fractions 50-60 produced 2-amino-6-chloro-8-methylpurine as a crude solid. Crystallization from ethanol/water with charcoal treatment afforded 2-amino-6-chloro-8-methylpurine as a white solid: yield, 0.57 g (51%); mp > 265 °C dec.; ¹H NMR d 2.39 (s, 3 H, CH₃), 6.62 (s, 2 H, NH₂, exchange with 10 D₂O), 12.56 (s, 1 H, NH, exchanges with D₂O); MS (EI) calcd. m/z for C₆H₆N₅³⁵Cl 183.0312, found 183.0309; calcd. m/z for C₆H₆N₅³⁷Cl 185.0283, found 185.0286.

15 Example 4: O⁶-Benzyl-8-methylguanine (1b)

Sodium (0.1 g, 4.4 mmol) was stirred in 4.1 mL of benzyl alcohol until all sodium had reacted. 2-Amino-6-chloro-8-methylpurine (0.41 g, 2.2 mmol) was added, and the reaction mixture was heated in a 130 °C oil bath for 5 h. 20 After cooling to room temperature 40 mL of ether was added to remove excess benzyl alcohol. The sticky precipitate that formed was collected by filtration and was dissolved in water (50 mL). The pH of the yellow solution was adjusted to 5 - 6 with glacial acetic acid. The solution 25 was mixed with methanol (50 mL) and was loaded on a 3 x 80 cm SephadexTM LH-20 column eluted with methanol/water (1:1) at 1 mL/min. Column eluent was continuously monitored at 280 nm and fractions (10 mL) were collected. Evaporation of pooled fractions 78-93 afforded analytically pure 1b: 30 yield, 0.25 g (44%); mp 214 - 216 °C; UV (pH 1) λ_{max} 238 nm (sh) (e = 0.648 x 10⁴), 290 (1.136 x 10⁴); (pH 6.9) 242 (0.758 x 10⁴), 284 (0.897 x 10⁴); (pH 13) 240 (sh) (0.495 x 10⁴), 286 (0.932 x 10⁴); ¹H NMR d 2.33 (s, 3 H, CH₃), 5.46 (s, 2 H, ArCH₂), 6.17 (s, 2 H, NH₂, exchange with D₂O), 7.34- 35 7.51 (m, 5 H, ArH), 12.18 (br s, 1 H, NH, exchanges with D₂O); MS (EI) calcd. m/z for C₁₃H₁₃N₅O 255.1120, found 255.1125; Anal. (C₁₃H₁₃N₅O.1/4 H₂O) C, H, N.

Example 5: O^6 -Benzyl-8-oxoguanine (1c)

2,4,5-Triamino-6-benzyloxypyrimidine (Pfleiderer et al., *Chem. Ber.*, 94, 12 - 18 (1961)) (1.85 g, 8 mmol) and 1,1'-carbonyldiimidazole (1.30 g, 8 mmol) were dissolved in anhydrous *N,N*-dimethylformamide (5 mL) under argon. The solution was stirred at room temperature overnight and was mixed with water (200 mL) to precipitate a white solid. The solid was collected by filtration, and dissolved in 250 mL of aqueous 2 N NaOH solution. Undissolved material was removed by filtration, and the filtrate was neutralized with glacial acetic acid to precipitate a white solid. The solid was collected by filtration, was washed with water, and was recrystallized from 50% aqueous ethanol to afford analytically pure 1c: yield, 1.63 g (79 %); mp 256 - 257 °C dec.; UV (pH 1) λ_{max} 243 nm ($\epsilon = 0.717 \times 10^4$), 306 (1.499 $\times 10^4$); (pH 6.9) 243 (0.915 $\times 10^4$), 290 (1.108 $\times 10^4$); (pH 13) 249 (sh) (0.443 $\times 10^4$), 293 (1.368 $\times 10^4$); ^1H NMR δ 5.41 (s, 2 H, ArCH_2), 6.13 (s, 2 H, NH_2 , exchange with D_2O), 7.33 - 7.51 (m, 5 H, ArH), 10.46 (s, 1 H, exchanges with D_2O), 11.04 (s, 1 H, exchanges with D_2O); MS (EI) Calcd. *m/z* for $\text{C}_{12}\text{H}_{11}\text{N}_5\text{O}_2$: 257.0912. Found: 257.0914. Anal. ($\text{C}_{12}\text{H}_{11}\text{N}_5\text{O}_2 \cdot 1/2 \text{H}_2\text{O}$) C, N, H.

25

Example 6: O^6 -Benzyl-8-bromoguanine (1d)

Bromine (0.26 mL, 5.1 mmol) was added slowly to the solution of O^6 -benzylguanine (1.205 g, 5.0 mmol) in anhydrous DMF (10 mL) under argon. The resulting deep green solution was stirred at room temperature overnight. The solution was mixed with water (70 mL) to precipitate crude product. This product was collected by filtration and was dissolved in 50% aqueous methanol (100 mL). The solution was loaded on a 3 x 80 cm SephadexTM LH-20 column eluted with methanol/water (1:1) at 1 mL/min. Column eluent was continuously monitored at 280 nm and fractions (10 mL) were collected. The desired product eluted in

fractions 110 - 190. Evaporation of solvent from the pooled fractions 110 - 190 afforded 1d as a pale yellow solid. Crystallization from ethanol/water (1:1) produced analytically pure 1d: yield, 0.166 g (10%); mp 135 - 137
5 °C dec.; UV (pH 1) λ_{max} 236 nm (sh) ($\epsilon = 0.517 \times 10^4$), 294 (1.429 $\times 10^4$); (pH 6.9) 244 (0.666 $\times 10^4$), 287 (1.043 $\times 10^4$); (pH 13) 245 (sh) (0.544 $\times 10^4$), 289 (1.030 $\times 10^4$); ^1H NMR d 5.45 (s, 2 H, ArCH₂), 6.35 (s, 2 H, NH₂, exchange with D₂O), 7.34 - 7.52 (m, 5 H, ArH), 13.08 (b s, 1 H, NH, 10 exchanges with D₂O); MS (EI) calcd. m/z for C₁₂H₁₀N₅O⁷⁹Br 319.0068, found 319.0069; calcd. m/z for C₁₂H₁₀N₅O⁸¹Br 321.0048, found 321.0048; Anal. (C₁₂H₁₀N₅OBr₃/2 H₂O) C, H, N, Br.

15 Example 7: 8-Aza- O^6 -benzylguanine (2)

Glacial acetic acid (1 mL) was added into the mixture of 2,4,5-triamino-6-benzyloxypyrimidine (0.231 g, 1.0 mmol) and sodium nitrite (0.069 g, 1.0 mmol) in acetone (5 mL).
20 The resulting mixture was stirred at room temperature for 2 h. The solution was poured in water (100 mL) with stirring to precipitate a crude solid. The solid was collected by filtration and air dried. Crystallization from ethanol/water (1:1) with charcoal treatment produced 2 as a
25 white solid: yield, 105 mg (43%); mp 191 - 192 °C (192 - 193 °C; Shealy et. al., J. Org. Chem., 27, 4518 - 4523 (1962)); ^1H NMR d 5.56 (s, 2 H, ArCH₂), 7.00 (s, 2 H, NH₂, exchange with D₂O), 7.41 - 7.58 (m, 5 H, ArH); MS (EI) calcd. m/z for C₁₁H₁₀N₆O 242.0916, found 242.0924.

30

Example 8: 4-Amino-6-benzyloxy-5-nitropyrimidine (3a)

4-Amino-6-chloro-5-nitropyrimidine (Boon et al., J. Chem. Soc., 96-102 (1951)) (1.5 g, 8.6 mmol) was added to a
35 solution of sodium (0.23 g, 9.9 mmol) in benzyl alcohol (14 mL). The solution was heated in a 130 °C oil bath for 3.5 h, and was poured into benzene (50 mL). A yellow solid was collected by filtration and washed with benzene.

Crystallization from benzene/ether afforded an analytically pure sample of 3a: yield, 0.71 g (34%); mp 149 - 150 °C; UV (pH 1) λ_{max} 284 nm ($\epsilon = 0.368 \times 10^4$), 333 (0.488×10^4); (pH 6.9) 284 (0.329×10^4), 336 (0.470×10^4); (pH 13) 290 (0.344 $\times 10^4$), 333 (0.494×10^4); ^1H NMR d 5.50 (s, 2 H, ArCH₂), 7.33 - 7.49 (m, 5 H, ArH), 8.12 - 8.24 (br d, 2 H, NH_a and NH_b, exchange with D₂O), 8.24 (s, 1 H, H-2); MS (EI) calcd. m/z for C₁₁H₁₀N₄O₃ 246.0752, found 246.0751; Anal. (C₁₁H₁₀N₄O₃) C, H, N.

10

Example 9: 2,4-Diamino-6-benzylxy-5-nitropyrimidine (3e)

2,4-Diamino-6-chloro-5-nitropyrimidine (O'Brien et. al., *J. Med. Chem.*, 9, 573 - 575 (1966)) (1.0 g, 5.28 mmol) was added to a solution of sodium (0.14 g, 6.08 mmol) in benzyl alcohol (9 mL). The solution was heated in a 160 °C oil bath for 3.5 h and the solvent was evaporated under reduced pressure to provide a yellow solid. This solid was washed with water, and air dried. Crystallization from benzene/ether gave a pale yellow filamentous solid: yield, 0.69 g (50%); mp 194 - 195 °C (171 °C; Kosary et. al., *Acta. Pharm. Hung.*, 49, 241 - 247 (1989)); UV (pH 1) λ_{max} 236 nm (sh) ($\epsilon = 1.452 \times 10^4$), 264 (0.522×10^4), 321 (1.294 $\times 10^4$); (pH 6.9) 242 (sh) (0.965×10^4), 337 (1.493×10^4); (pH 13) 242 (sh) (0.952×10^4), 338 (1.479×10^4); ^1H NMR d 5.43 (s, 2 H, ArCH₂), 7.26 (br s, 2 H, NH₂, exchange with D₂O), 7.33 - 7.51 (m, 5 H, ArH), 7.93 (br s, 2 H, NH₂, exchange with D₂O); MS (EI) calcd. m/z for C₁₁H₁₁N₅O₃ 261.0861, found 261.0866; Anal. (C₁₁H₁₁N₅O₃).

30

Example 10: O⁶-Benzylxanthine (4a)

A suspension of O⁶-benzylguanine (0.83 g, 3.4 mmol) in acetone (15 mL) was poured into a solution of sodium nitrite (5 g) in 15 mL of H₂O. Acetic acid (8 mL) was added to the suspension with stirring. Minimum amounts of acetone were added as necessary to dissolve any suspended solid. The resulting pale yellow-green solution was

stirred for 3 h. A pale green precipitate that formed was collected by filtration and washed with water (200 mL). Recrystallization of the air-dried solid from ethanol/water (1:1) afforded analytically pure 4a: yield, 0.43 g (52%);
5 mp 145 - 147 °C dec.; UV (pH 1) λ_{max} 270 nm ($\epsilon = 0.749 \times 10^4$); (pH 6.9) 286 (1.143×10^4); (pH 13) 290 (0.914×10^4);
 ^1H NMR d 5.49 (s, 2 H, ArCH₂), 7.36-7.54 (m, 5 H, ArH), 8.02
(s, 1 H, H-8), 11.8 (br s, 1 H, NH, exchanges with D₂O),
13.2 (br s, 1 H, NH, exchanges with D₂O); MS (EI) calcd. m/z
10 for C₁₂H₁₀N₄O₂ 242.0803, found 242.0828; Anal. (C₁₂H₁₀N₄O₂ · H₂O) C, H, N.

Example 11: O⁶-Benzyluric acid (4b)

15 Sodium nitrite (1.5 g, 43 mmol) dissolved in water (5 mL) was added to a suspension of O⁶-benzyl-8-oxoguanine (1c) (0.257 g, 1.0 mmol) in acetone (5 mL). Glacial acetic acid (3 mL) was added to the suspension with stirring. After stirring for 3 h at room temperature a bright yellow precipitate formed. The suspension was mixed with water (150 mL) and undissolved solid was filtered off. Saturated aqueous sodium carbonate solution was added to the filtrate to adjust the pH to approximately 5. A yellow precipitate (130 mg) was collected and washed with water. This solid
20 was crystallized from 50% aqueous ethanol to give an analytically pure sample of 4b: yield, 75 mg (29%); mp >230 °C; UV (pH 1) λ_{max} 236 nm (sh) ($\epsilon = 0.972 \times 10^4$), 299 (1.427×10^4); (pH 6.9) 240 (sh) (0.821×10^4), 304 (2.134×10^4); (pH 13) 245 (sh) (0.846×10^4), 297 (1.861×10^4);
 ^1H NMR d 5.43 (s, 2 H, ArCH₂), 7.35-7.51 (m, 5 H, ArH), 10.76
(s, 1 H, NH, exchanges with D₂O), 11.23 (s, 1 H, NH,
30 exchanges with D₂O), 11.39 (s, 1 H, NH, exchanges with D₂O); MS (EI) calcd. m/z for C₁₂H₁₀N₄O₃ 258.0752, found 258.0753; Anal. (C₁₂H₁₀N₄O₃ · 5/2 H₂O) C, H, N.

Example 12: Diacetyl- O^6 -benzyl-8-oxoguanine

Acetic anhydride (2 mL) was added to the suspension of O^6 -benzyl-8-oxoguanine (1c) (0.257 g, 1.0 mmol) in dry toluene (10 mL). The suspension was vigorously refluxed for 24 hr, and was cooled to room temperature. After storing at 4 °C for 4 hr, the resulting precipitate was collected by filtration, washed with benzene and air dried to give an analytically pure sample of a diacetylated product: yield, 0.287 g (84%); mp 272 - 274 °C dec.; UV (100% MeOH) λ_{max} 275 nm ($\epsilon = 1.313 \times 10^4$); (pH 1) 275 (1.143 $\times 10^4$); (pH 6.9) 238 (0.995 $\times 10^4$), 276 (1.115 $\times 10^4$); (pH 13) 285 (2.138 $\times 10^4$); ^1H NMR d 2.18 (s, 3 H, CH_3), 2.57 (s, 3 H, CH_3), 5.51 (s, 2 H, ArCH_2), 7.30 - 7.57 (m, 5 H, ArH), 10.41 (s, 1 H, exchanges with D_2O), 12.30 (s, 1 H, exchanges with D_2O); MS (EI) Calcd. m/z for $\text{C}_{16}\text{H}_{15}\text{N}_5\text{O}_4$: 341.1123. Found: 341.1130. Anal. ($\text{C}_{16}\text{H}_{15}\text{N}_5\text{O}_4$) C, N, H.

Example 13: N^2 -Acetyl- O^6 -benzyl-8-oxoguanine (4d)

Diacetyl- O^6 -benzyl-8-oxoguanine (85 mg, 0.25 mmol) was dissolved in methanol (10 mL) and ammonium hydroxide (28%, 5 mL) and was allowed stand for 1 hr. The clear solution became cloudy and a precipitate formed on standing. The precipitate was collected by filtration, washed with water, and dried to give an analytically pure sample of 4d: yield, 48 mg (65%); mp 335 - 337 °C dec.; UV (pH 1) λ_{max} 276 nm ($\epsilon = 1.723 \times 10^4$), 303 (sh) (0.679 $\times 10^4$); (pH 6.9) 276 (1.379 $\times 10^4$); (pH 13) 284 (1.683 $\times 10^4$); ^1H NMR d 2.15 (s, 3 H, CH_3), 5.49 (s, 2 H, ArCH_2), 7.30 - 7.55 (m, 5 H, ArH), 10.21 (s, 1 H, exchanges with D_2O), 10.99 (s, 1 H, exchanges with D_2O), 11.60 (s, 1 H, exchanges with D_2O); MS (EI) Calcd. m/z for $\text{C}_{14}\text{H}_{13}\text{N}_5\text{O}_3$: 299.1018. Found: 299.1023. Anal. ($\text{C}_{14}\text{H}_{13}\text{N}_5\text{O}_3$) C, N, H.

35

Example 14: O^6 -Benzyl-2-fluorohypoxanthine (4c)

O^6 -Benzylguanine (1.21 g, 5 mmol) was added to 100 mL of 48% fluoboric acid at -20 °C. Sodium nitrite (1.23 g,

35 mmole) was dissolved in water (5 mL) and 2.5 mL of this sodium nitrite solution was added slowly to the cold fluoboric acid solution. The resulting mixture was stirred for 1 h at or below -15 °C. Additional fluoboric acid (25 mL) was added followed by an additional 2.5 mL of the aqueous sodium nitrite solution. After stirring for an additional 1 h below -15 °C, fluoboric acid (25 mL) was again added and stirring was continued for 1 h. The resulting solution was neutralized with saturated aqueous sodium carbonate solution at -20 °C and was allowed to warm to room temperature. A white precipitate that formed was collected by filtration and was washed with water and dried under vacuum to afford crude 4c: yield, 0.52 g, 43%. An analytical sample was prepared by chromatography on a 15 Sephadex™ LH-20 column (3 x 80 cm) eluted with methanol/water (1:1) at 1 mL/min. The desired 4c eluted in fractions 66 - 77: mp 182 - 183 °C (184 - 185 °C; Robins and Robins, *J. Org. Chem.*, 34, 2160-2163 (1969)); UV (pH 1) λ_{max} 256 nm ($\epsilon = 1.117 \times 10^4$); (pH 6.9) 257 (1.078×10^4); (pH 13) 264 (1.063×10^4); ^1H NMR d 5.60 (s, 2 H, ArCH_2), 7.37-7.57 (m, 5 H, ArH), 8.40 (s, 1 H, H-8), 13.60 (s, 1 H, NH, exchanges with D_2O), ^{19}F NMR d 23.54 downfield from trifluoroacetic acid standard; MS (EI) calcd. *m/z* for $\text{C}_{12}\text{H}_9\text{FN}_4\text{O}$ 244.0760, found 244.0756; Anal. ($\text{C}_{12}\text{H}_9\text{FN}_4\text{O}$ 0.2/3 H_2O) 25 C, H, N.

Example 15: O^6 -Benzyl- N^2 -methylguanine (4e)

Fluoboric acid (48%, 30 mL) was cooled to -20 °C in an 30 dry ice-acetone bath. O^6 -Benzylguanine (0.362 g, 1.5 mmol) was added with stirring. Sodium nitrite (0.369 g, 10.5 mmol) was dissolved in water (1 mL) and 0.5 mL of this solution was added slowly to the cold fluoboric acid solution. The resulting solution was stirred at or below 35 -15 °C for 1 h. More fluoboric acid (5 mL) was then added followed by 0.5 mL of the sodium nitrite solution. After stirring for 1 h at or below -15 °C, fluoboric acid (5 mL)

was again added and stirring was continued for an additional 1 h. Methylamine (40% in water, 60 mL) was then added at -20 °C, and the resulting basic solution was stirred at room temperature for 2 days. The solvent was 5 evaporated under reduced pressure to produce a white solid. The solid was suspended in 50 mL of H₂O with stirring for 10 min. Undissolved material was collected by filtration and washed with water. This solid was dissolved in 40 mL methanol/water (1:1) to which was added 1.2 mL of 28% 10 aqueous ammonia solution. The solution was loaded on a 3 x 80 cm SephadexTM LH-20 column eluted with MeOH/H₂O/NH₄OH (30:70:3) at 1 mL/min. Column eluent was continuously monitored at 280 nm and fractions (10 mL) were collected. Evaporation of the pooled fractions 106-127 gave an 15 analytically pure sample of 4e: yield, 85 mg (22%); mp 189 - 190 °C; UV (pH 1) λ_{max} 238 nm (sh) ($\epsilon = 0.665 \times 10^4$), 297 (0.904 $\times 10^4$); (pH 6.9) 246 (0.898 $\times 10^4$), 290 (0.676 $\times 10^4$); (pH 13) 240 (sh) (0.615 $\times 10^4$), 294 (0.674 $\times 10^4$); ¹H NMR d 2.30 (d, 3 H, CH₃), 5.50 (s, 2 H, ArCH₂), 6.75 (m, 1 20 H, MeNH, exchanges with D₂O), 7.31-7.53 (m, 5 H, ArH), 7.82 (s, 1 H, H-8), 12.53 (s, 1 H, NH, exchanges with D₂O); MS (EI) calcd. m/z for C₁₃H₁₃N₅O 255.1120, found 255.1107; Anal. (C₁₃H₁₃N₅O.1/2 H₂O) C, H, N.

25 Example 16: O⁶-Benzyl-N²,N²-dimethylguanine (4f)

Fluoboric acid (48%, 40 mL) was cooled to -20 °C in an dry ice-acetone bath. O⁶-Benzylguanine (0.482 g, 2.0 mmol) was added with stirring. Sodium nitrite (0.492 g, 14.0 30 mmol) was dissolved in water (2 mL) and 1 mL of this solution was added slowly to the cold fluoboric acid solution. The resulting solution was stirred at or below -15 °C for 1 h. More fluoboric acid (10 mL) was added followed by the addition of 1 mL of the sodium nitrite 35 solution. After stirring for 1 h at or below -15 °C, additional fluoboric acid (10 mL) was added with stirring for 1 h. Dimethylamine (40% in water, 60 mL) was then

added to the solution at -20 °C, and the resulting mixture was allowed to warm to room temperature. The suspension became a clear solution and a precipitate formed within 10 min. After standing overnight at room temperature the precipitate was collected by filtration and was washed with water. The solid was crystallized from 50% aqueous ethanol to give an analytically pure sample of 4f: yield, 0.25 g (46%); mp 220 - 221 °C dec.; UV (pH 1) λ_{max} 248 nm (sh) (ϵ = 0.512 $\times 10^4$), 303 (0.908 $\times 10^4$); (pH 6.9) 251 (1.152 $\times 10^4$), 10 299 (0.686 $\times 10^4$); (pH 13) 248 (sh) (0.766 $\times 10^4$), 299 (0.710 $\times 10^4$); ^1H NMR d 3.12 (s, 6 H, CH_3), 5.54 (s, 2 H, ArCH_2), 7.36-7.51 (m, 5 H, ArH), 7.84 (s, 1 H, H-8), 12.56 (s, 1 H, NH, exchanges with D_2O); MS (EI) calcd. m/z for $\text{C}_{14}\text{H}_{15}\text{N}_5\text{O}$ 269.1276, found 269.1254; Anal. ($\text{C}_{14}\text{H}_{15}\text{N}_5\text{O}$) C, H, N.

15

Example 17: 2,4-Diamino-6-benzyloxy-5-bromopyrimidine (3f)

2,4-Diamino-5-bromo-6-chloropyrimidine (Phillips et. al., *J. Org. Chem.*, 29, 1488 - 1490 (1963)) (2.3 g, 10 20 mmol) was added to a solution of sodium (0.29 g, 12.5 mmol) in benzyl alcohol (10 mL) under argon. The solution was heated in a 130 °C oil bath for 3 h and the benzyl alcohol was evaporated under reduced pressure to give a white solid. This solid was washed with water, and air dried. 25 Crystallization from 50% aqueous ethanol gave white crystalline needles of 3f: yield, 2.32 g (76%); mp 165 - 166 °C (lit. 136 °C; Kosary et. al., *Acta Pharm. Hung.*, 49, 241 - 247 (1989)); UV (pH 1) λ_{max} 236 nm (ϵ = 0.873 $\times 10^4$), 291 (1.388 $\times 10^4$); (pH 6.9) 236 (0.850 $\times 10^4$), 277 (0.835 \times 30 10 4); (pH 13) 234 (0.869 $\times 10^4$), 277 (0.827 $\times 10^4$); ^1H NMR d 5.30 (s, 2 H, ArCH_2), 6.15 (s, 2 H, NH_2 , exchange with D_2O), 6.32 (s, 2 H, NH_2 , exchange with D_2O), 7.31 - 7.45 (m, 5 H, ArH); MS (EI) calcd. m/z for $\text{C}_{11}\text{H}_{11}\text{N}_4\text{O}^{79}\text{Br}$ 294.0115, found 294.0127; calcd. m/z for $\text{C}_{11}\text{H}_{11}\text{N}_4\text{O}^{81}\text{Br}$ 296.0094, found 35 296.0083; Anal. ($\text{C}_{11}\text{H}_{11}\text{N}_4\text{OBr}$) C, H, N.

Example 18: 2-Amino-4-chloro-5-nitropyrimidine

A suspension of 2-amino-4-hydroxy-5-nitropyrimidine (5.0 g, 32.1 mmol) in phosphorous oxychloride (100 mL) was 5 refluxed overnight, and the excess phosphorous oxychloride was evaporated under reduced pressure. The residue was mixed with ice (100 g) in an ice-bath, and the mixture was neutralized with concentrated aqueous sodium carbonate solution. A yellow precipitate was collected by filtration 10 and washed with water: yield, 1.39 g (25%); mp 191 - 194 °C dec.; ^1H NMR δ 8.45 (br s, 2 H, NH_2 , exchange with D_2O), 9.03 (s, 1 H, H-6); MS (EI) calcd. m/z for $\text{C}_4\text{H}_3\text{N}_4\text{O}_2^{35}\text{Cl}$ 173.9944, found 173.9934; calcd. m/z for $\text{C}_4\text{H}_3\text{N}_4\text{O}_2^{37}\text{Cl}$ 175.9915, found 175.9916.

15

Example 19: 2-Amino-4-benzyloxy-5-nitropyrimidine (5a)

2-Amino-4-chloro-5-nitropyrimidine (0.70 g, 4.0 mmol) was added to a solution of sodium (0.12 g, 5.2 mmol) in 20 benzyl alcohol (8 mL) under argon. The solution was heated in a 130 °C oil bath for 3 h, and approximately half of the benzyl alcohol was evaporated under reduced pressure. The residue was poured into water (50 mL) with constant stirring for 10 min. After neutralization with glacial 25 acetic acid, a brown precipitate formed which was collected by filtration and washed with water. This solid was crystallized from benzene to give 5a as a golden crystalline solid: yield, 126 mg (13%); mp 164 - 167 °C; UV (pH 1) λ_{max} 262 nm ($\epsilon = 0.879 \times 10^4$), 295 (sh) (0.571 x 10⁴); (pH 6.9) 235 (sh) (0.448 x 10⁴), 273 (0.360 x 10⁴), 30 326 (1.085 x 10⁴); (pH 13) 273 (0.404 x 10⁴), 327 (1.055 x 10⁴); ^1H NMR δ 5.51 (s, 2 H, ArCH_2), 7.35 - 7.54 (m, 5 H, ArH), 8.05 (d, 2 H, NH_2 , exchange with D_2O), 8.92 (s, 1 H, H-6); MS (EI) calcd. m/z for $\text{C}_{11}\text{H}_{10}\text{N}_4\text{O}_3$ 246.0752, found 35 246.0758; Anal. ($\text{C}_{11}\text{H}_{10}\text{N}_4\text{O}_3$) C, H, N.

Example 20: 2-Amino-4-benzyloxy-6-methyl-5-nitropyrimidine (5b)

2-Amino-4-chloro-6-methyl-5-nitropyrimidine (Boon et al., *J. Chem. Soc.*, 96 - 102 (1951)) (1.24 g, 6.58 mmol) was added to a solution of sodium (0.21 g, 9.13 mmol) in benzyl alcohol (14 mL) under argon. The solution was heated in a 135 °C oil bath for 3.5 h, and was poured into water (70 mL) with constant stirring for 10 min. After neutralization with glacial acetic acid, a yellow precipitate formed which was collected by filtration and washed with water. This solid was crystallized from benzene to give 5b as a bright yellow crystalline solid: yield, 0.57 g (33%); mp 159 - 160 °C; UV (pH 1) λ_{max} 268 nm (e = 0.783 x 10⁴), 345 (sh) (0.104 x 10⁴); (pH 6.9) 282 (0.564 x 10⁴), 345 (sh) (0.338 x 10⁴); (pH 13) 282 (0.549 x 10⁴), 345 (sh) (0.332 x 10⁴); ¹H NMR d 2.35 (s, 3 H, CH₃), 5.44 (s, 2 H, ArCH₂), 7.34 - 7.46 (m, 5 H, ArH), 7.64 (b s, 2 H, NH₂, exchange with D₂O); MS (EI) calcd. m/z for C₁₂H₁₂N₄O₃ 260.0908, found 260.0913; Anal. (C₁₂H₁₂N₄O₃) C, H, N.

Example 21: 2,4-Diamino-6-benzyloxy-s-triazine (6)

2,4-Diamino-6-chloro-s-triazine (2.25 g, 15.0 mmol) was added to a solution of sodium (0.43 g, 18.8 mmol) in benzyl alcohol (30 mL) under argon. The suspension was heated in a 130 °C oil bath for 3.5 h. The excess benzyl alcohol was removed under vacuum and the resulting solid was collected with the aid of benzene, and washed with water (100 mL): yield, 1.83 g (56%); mp 184 - 185 °C (lit. 186 - 188 °C; Wakabayashi et al., *Nippon Dojo-Hiryogaku Zasshi*, 41, 193 - 200 (1970)); UV (pH 1) λ_{max} 233 nm (sh) (e = 0.589 x 10⁴); (pH 6.9) 238 (sh) (0.111 x 10⁴); (pH 13) 240 (sh) (0.073 x 10⁴); ¹H NMR d 5.25 (s, 2 H, ArCH₂), 6.63 (s, 4 H, NH₂, exchange with D₂O), 7.30 - 7.42 (m, 5 H, ArH); MS (EI) calcd. m/z for C₁₀H₁₁N₅O 217.0963, found 217.0955.

Example 22: 2-Amino-6-chloro-8-trifluoromethylpurine

A suspension of 8-trifluoromethylguanine (Pfleiderer and Shanshal, *Liebigs Ann. Chem.*, 726, 201 - 215 (1969)) (2.0 g, 9.1 mmol) in phosphorous oxychloride (20 mL) was refluxed for 3 h. Excess phosphorous oxychloride was evaporated under reduced pressure. The resulting residue was mixed with ice-water (100 g), and the pH was adjusted to 3 - 4 with a concentrated aqueous NaOH solution. The resulting solution was mixed with MeOH (100 mL) and approximately half (i.e., 100 mL) of the aqueous methanol solution was loaded on a 3 x 80 cm SephadexTM LH-20 column eluted with methanol/water (1:1) at 1 mL/min. Column eluent was continuously monitored at 280 nm and fractions (10 mL) were collected. The remainder of the reaction mixture in MeOH/H₂O was chromatographed separately under identical conditions. The desired product eluted in fractions 73 - 85. Evaporation of solvent from the pooled fractions 73 - 85 from both chromatographic runs afforded analytically pure 2-amino-6-chloro-8-trifluoromethylpurine: yield, 0.94 g (43%); mp >225 °C dec.; UV (pH 1) λ_{max} 245 nm ($\epsilon = 0.501 \times 10^4$), 314 (0.746×10^4); (pH 6.9) 270 (0.265 $\times 10^4$), 315 (0.612×10^4); (pH 13) 272 (0.269 $\times 10^4$), 314 (0.612×10^4); ¹H NMR d 7.19 (s, 2 H, NH₂, exchange with D₂O), 14.25 (br s, 1 H, NH, exchanges with D₂O); MS (EI) calcd. m/z for C₆H₃N₅F₃³⁵Cl 237.0029, found 237.0011; calcd. m/z for C₆H₃N₅F₃³⁷Cl 239.0000, found 238.9987; Anal. (C₆H₃N₅F₃C) C, H, N, F, Cl.

30 Example 23: O⁶-Benzyl-8-trifluoromethylguanine (1e)

Sodium (0.10 g, 4.3 mmol) was stirred in 5 mL of benzyl alcohol until all had reacted. 2-Amino-6-chloro-8-trifluoromethylpurine (0.475 g, 2.0 mmol) was added, and the reaction mixture was heated in a 135 °C oil bath for 3.5 h. The benzyl alcohol was removed by vacuum distillation yielding a brown oil. The oil was dissolved in water (50 mL) and was acidified with glacial acetic acid

to produce a pale yellow precipitate. The precipitate was collected by filtration and washed with water. The crude product was loaded on a 2.5 x 35 cm silica gel column (DavisilTM grade 633, 200 - 425 mesh, 60 Å). Elution was carried out with 5% EtOH in CHCl₃ to provide analytically pure *O*⁶-benzyl-8-trifluoromethylguanine (1e): yield, 0.42 g (67%); mp 214 - 216 °C dec.; UV (pH 1) λ_{max} 291 nm (ϵ = 1.229 $\times 10^4$); (pH 6.9) 244 (0.470 $\times 10^4$), 289 (1.023 $\times 10^4$); (pH 13) 247 (sh) (0.393 $\times 10^4$), 290 (0.923 $\times 10^4$); ¹H NMR δ 5.51 (s, 2 H, ArCH₂), 6.82 (s, 2 H, NH₂, exchange with D₂O), 7.38 - 7.55 (m, 5 H, ArH), 13.75 (br s, 1 H, NH, exchanges with D₂O); MS (EI) calcd. *m/z* for C₁₃H₁₀N₅OF₃ 309.0837, found 309.0827; Anal. (C₁₃H₁₀N₅OF₃) C, H, N, F.

15 Example 24: *O*⁶-Benzyl-8-trifluoromethyl-9-methylguanine (7)

To *O*⁶-benzyl-8-trifluoromethylguanine (1e) (200 mg, 0.65 mmol) under argon was added 0.66 mL of a 1.0 M solution of sodium ethoxide in ethanol. The solution was stirred for 10 min and the ethanol was removed under vacuum. The remaining solid was dissolved in anhydrous DMF (1.5 mL), and methyl iodide (49 μL, 0.78 mmol) was added to the solution. This solution was stirred at room temperature for 1 h, and 1.5 mL additional DMF was added. The solution was stirred at room temperature overnight. The solvent was evaporated under reduced pressure. The crude solid was loaded on a 2.5 x 35 cm silica gel column (Davisil grade 633, 200 - 425 mesh, 60 Å). Elution was carried out with chloroform/hexane (3:1) to provide analytically pure *O*⁶-benzyl-8-trifluoromethyl-9-methylguanine (7): yield, 95 mg (45%); mp 86 - 89 °C; UV (pH 1) λ_{max} 244 nm (ϵ = 0.581 $\times 10^4$), 286 (1.274 $\times 10^4$); (pH 6.9) 252 (0.608 $\times 10^4$), 288 (1.022 $\times 10^4$); (pH 13) 252 (0.618 $\times 10^4$), 288 (1.038 $\times 10^4$); ¹H NMR δ 3.70 (s, 3 H, CH₃), 5.51 (s, 2 H, ArCH₂), 6.91 (s, 2 H, NH₂, exchange with D₂O), 7.38 - 7.54 (m, 5 H, ArH); MS (EI) calcd. *m/z* for

$C_{14}H_{12}N_5OF_3$ 323.0994, found 323.0978; Anal. ($C_{14}H_{12}N_5OF_3$) C, H, N, F.

Example 25: 8-Aza- O^6 -benzyl-9-methylguanine (8a)

8-Aza- O^6 -benzylguanine (0.484 g, 2.0 mmol) was mixed with 4 mL of 0.5 M sodium ethoxide in ethanol and stirred for 30 min. The ethanol was evaporated under reduced pressure. The residue was dissolved in anhydrous DMF (6 mL), and methyl iodide (0.15 mL, 2.4 mmol) was added. The clear solution became cloudy within 10 min, and the resulting mixture was stirred overnight at room temperature. DMF was evaporated under reduced pressure to give a brown solid. The solid was dissolved in chloroform and loaded on a silica gel column (DavisilTM grade 633, 200-425 mesh, 60 Å). Product 8a was eluted with chloroform; yield, 138 mg (27%); mp 178-179 °C; UV (pH 1) λ_{max} 243 nm (sh) ($\epsilon = 0.556 \times 10^4$), 284 (1.112×10^4); (pH 6.9) 243 (0.553×10^4), 290 (0.998×10^4); (pH 13) 242 (0.549×10^4), 290 (1.010×10^4); ¹H NMR δ 3.96 (s, 3 H, CH_3), 5.57 (s, 2 H, $ArCH_2$), 7.18 (s, 2 H, NH_2 , exchange with D_2O), 7.38-7.57 (m, 5 H, ArH); MS (EI) calcd m/z for $C_{12}H_{12}N_6O$ 256.1072, found 256.1086; Anal. ($C_{12}H_{12}N_6O$) C, H, N.

Example 26: 8-Aza- O^6 -benzyl-9-(pivaloyloxymethyl)guanine (8b) and 8-Aza- O^6 -benzyl-7-(pivaloyloxymethyl)guanine (9)

8-Aza- O^6 -benzylguanine (0.484 g, 2.0 mmol) was mixed with 4 mL of 0.5 M sodium ethoxide in ethanol and stirred for 30 min. The ethanol was evaporated under reduced pressure. The residue was dissolved in anhydrous DMF (6 mL), and chloromethyl pivalate (0.3 mL, 2.1 mmol) was added. The clear solution was stirred for 8 h at room temperature. DMF was evaporated under reduced pressure to give a brown solid. The solid was dissolved in chloroform and loaded on a silica gel column (DavisilTM grade 633, 200-425 mesh, 60 Å). The 9-isomer (8b) was eluted from the column with $CHCl_3$:hexane (4:1) while the 7-isomer (9) was

subsequently eluted with CHCl_3 . 8-Aza- O^6 -benzyl-9-(pivaloyloxymethyl)guanine (**8b**): yield, 405 mg (57%); mp 119-120 °C; UV (pH 1) λ_{max} 246 nm ($\epsilon = 0.494 \times 10^4$), 286 (0.878×10^4); (pH 6.9) 247 (0.472×10^4), 288 (0.819×10^4); (pH 13) (decomposes to 8-aza- O^6 -benzylguanine); ^1H NMR δ 1.10 (s, 9 H, $\text{C}(\text{CH}_3)_3$), 5.50 (s, 2 H, ArCH_2), 6.31 (s, 2 H, CH_2), 7.38 (s, 2 H, NH_2 , exchange with D_2O), 7.40-7.54 (m, 5 H, ArH); MS (EI) calcd m/z for $\text{C}_{17}\text{H}_{20}\text{N}_6\text{O}_3$ 356.1596, found 356.1578; Anal. ($\text{C}_{17}\text{H}_{20}\text{N}_6\text{O}_3 \cdot 1/5\text{H}_2\text{O}$) C, H, N. 8-Aza- O^6 -benzyl-7-(pivaloyloxymethyl)guanine (**9**): yield, 103 mg (15%); mp 153-154 °C; UV (pH 1) λ_{max} 244 nm ($\epsilon = 0.820 \times 10^4$), 294 (1.249×10^4); (pH 6.9) 250 (sh) (0.296×10^4), 313 (0.503×10^4); (pH 13) (decomposes to 8-aza- O^6 -benzylguanine); ^1H NMR δ 1.12 (s, 9 H, $\text{C}(\text{CH}_3)_3$), 5.56 (s, 2 H, ArCH_2), 6.40 (s, 2 H, CH_2), 7.04 (s, 2 H, NH_2 , exchange with D_2O), 7.4-7.58 (m, 5 H, ArH); MS (EI) calcd m/z for $\text{C}_{17}\text{H}_{20}\text{N}_6\text{O}_3$ 356.1596, found 356.1602; Anal. ($\text{C}_{17}\text{H}_{20}\text{N}_6\text{O}_3$).

Example 27: O^6 -Benzyl-8-bromo-9-methylguanine (**10a**)

O^6 -Benzyl-9-methylguanine (0.252 g, 1.0 mmol) and sodium bicarbonate (0.084 g, 1.0 mmol) were dissolved in anhydrous DMF (2 mL) under argon. Bromine (52 mL, 1.0 mmol) was added to the solution and the resulting mixture was stirred overnight at room temperature. The solvent was evaporated under reduced pressure. The residue was dissolved in chloroform, and loaded on a silica gel column (DavisilTM grade 633, 200 - 425 mesh, 60 Å). Product **10a** was eluted with chloroform; yield, 180 mg (52%); mp 150-152 °C; UV (pH 1) λ_{max} 248 nm ($\epsilon = 0.753 \times 10^4$), 292 (1.398×10^4); (pH 6.9) 251 (0.919×10^4), 287 (1.306×10^4); (pH 13) 251 (0.906×10^4), 287 (1.296×10^4); ^1H NMR δ 3.53 (s, 3 H, CH_3), 5.47 (s, 2 H, ArCH_2), 6.61 (s, 2 H, NH_2 , exchange with D_2O), 7.35-7.52 (m, 5 H, ArH); MS (EI) calcd m/z for $\text{C}_{13}\text{H}_{12}\text{N}_5\text{O}^{79}\text{Br}$ 333.0225, found 333.0228; calcd m/z for $\text{C}_{13}\text{H}_{12}\text{N}_5\text{O}^{81}\text{Br}$ 335.0205, found 335.0188; Anal. ($\text{C}_{13}\text{H}_{12}\text{N}_5\text{OBr}$) C, H, N, Br.

Example 28: O^6 -Benzyl-8-bromo-9-(pivaloyloxymethyl)guanine (10b) and O^6 -benzyl-8-bromo-7-(pivaloyloxymethyl)guanine (11)

5 O^6 -Benzyl-8-bromoguanine (Chae et al., *J. Med. Chem.*, 38, 342-347 (1995)) (0.48 g, 1.5 mmol) was mixed with 1.5 mL of a 1.0 M solution of sodium ethoxide in ethanol and was stirred for 20 min. The ethanol was removed under 10 reduced pressure and the solid residue was dissolved in DMF (5 mL). Chloromethylpivalate (0.24 mL, 1.65 mmol) was then added and the solution was stirred overnight. The DMF was removed under reduced pressure. The residue was dissolved in chloroform and was loaded on a silica gel column 15 (Davisil grade 633, 200-425 mesh, 60A) eluted with chloroform. The 9-isomer (10b) eluted earlier than the 7-isomer with chloroform and 10b was recovered in pure form under these conditions. O^6 -Benzyl-8-bromo-9-(pivaloyloxymethyl)guanine (10b): yield, 150 mg (23%); mp 20 217-218°C; UV (pH 1) λ_{max} 250 nm ($\epsilon = 0.944 \times 10^4$), 291 (1.166 $\times 10^4$); (pH 6.9) 266 (0.916 $\times 10^4$), 295 (0.916 $\times 10^4$); (pH 13) decomposes to O^6 -benzyl-8-bromoguanine; ^1H NMR 25 d 1.13 (s, 9H, $\text{C}(\text{CH}_3)_3$), 5.48 (s, 2H, ArCH_2), 5.93 (s, 2H, CH_2), 6.80 (s, 2H, NH_2 , exchange with D_2O), 7.35-7.52 (m, 5H, ArH). MS (EI) calcd m/z for $\text{C}_{18}\text{H}_{20}\text{N}_5\text{O}_3^{79}\text{Br}$ 433.0750, found 433.0725; calcd m/z for $\text{C}_{18}\text{H}_{20}\text{N}_5\text{O}_3^{81}\text{Br}$ 435.0729, found 435.0672; Anal. ($\text{C}_{18}\text{H}_{20}\text{N}_5\text{O}_3\text{Br}$) C, H, N, Br. The recovered 7-isomer (11) was rechromatographed on a silica gel column 30 (DavisilTM grade 633, 200-425 mesh, 60 A) which was eluted first with CHCl_3 /hexane (1:1) followed by CHCl_3 to recover O^6 -benzyl-8-bromo-7-(pivaloyloxymethyl)guanine (11).


Example 29: O^6 -Benzyl-7-(pivaloyloxymethyl)guanine (12)

35 O^6 -Benzylguanine (2.41 g, 10 mmol) was mixed with 10 mL of a 1.0 M solution of sodium ethoxide in ethanol and was stirred for 30 min. The ethanol was evaporated under reduced pressure. The residue was dissolved in anhydrous DMF (30 mL), and chloromethyl-pivalate (Aldrich) (1.5 mL,

10.4 mmol) was added. The clear solution was stirred overnight at room temperature. DMF was evaporated under reduced pressure to give a pale peach-colored solid. The solid was dissolved in chloroform/ethanol (9:1) and loaded 5 on a silica gel column (Davisil grade 633, 200 - 425 mesh, 60 Å). The column was eluted with chloroform/ethanol (9:1) to elute the 9-isomer (Chae et al., *J. Med. Chem.*, 37, 342-347 (1994)) followed by the 7-isomer. The 7-isomer (12) was further purified by silica gel column 10 chromatography (Davisil grade 633, 200 - 425 mesh, 60 Å) using chloroform/ethanol (98:2) as eluent: yield, 36 mg (1%); mp 166-168 °C dec; UV (pH 1) λ_{max} 240 nm (sh) (ϵ = 0.656 $\times 10^4$), 290 (1.164 $\times 10^4$); (pH 6.9) 240 (sh) (0.635 $\times 10^4$), 293 (0.528 $\times 10^4$); (pH 13) decomposes to O^6 - 15 benzylguanine; ^1H NMR δ 0.98 (s, 9 H, $\text{C}(\text{CH}_3)_3$), 5.51 (s, 2 H, ArCH_2), 6.07 (s, 2 H, CH_2), 6.32 (s, 2 H, NH_2 , exchange with D_2O), 7.36-7.58 (m, 5 H, ArH), 8.25 (s, 1 H, H-8); MS (EI) calcd m/z for $\text{C}_{18}\text{H}_{21}\text{N}_5\text{O}_3$ 355.1644, found 355.1626.

CLAIMS:

1. A compound of the formula



wherein R₁ is a substituent selected from the group consisting of amino, hydroxy, C₁-C₄ alkylamino, C₁-C₄ dialkylamino, and C₁-C₄ alkylcarbonylamino, R₂ is a substituent selected from the group consisting of hydrogen, C₁-C₄ alkyl, C₁-C₄ aminoalkyl, C₁-C₄ hydroxyalkyl, C₁-C₄ alkylamino C₁-C₄ alkyl, C₁-C₄ dialkylamino alkyl, C₁-C₄ cyanoalkyl, C₁-C₄ carbamoylalkyl, C₁-C₄ pivaloylalkyl, C₁-C₆ alkylcarbonyloxy C₁-C₄ alkyl, C₁-C₄ alkoxy carbonylalkyl, 2'-deoxyribose, the conjugate acid form of a C₁-C₄ carboxyalkyl, and the carboxylate anion of a C₁-C₄ carboxyalkyl as the sodium salt, and R₃ is a substituent selected from the group consisting of halo, C₁-C₄ alkyl, C₁-C₄ hydroxyalkyl, mercapto, C₁-C₄ alkylthio, trifluoromethylthio, C₁-C₄ alkylthiocarbonyl, hydroxy, C₁-C₄ alkoxy, trifluoromethoxy, methanesulfonyloxy, trifluoromethanesulfonyloxy, C₁-C₄ alkylcarbonyloxy, amino, C₁-C₄ aminoalkyl, C₁-C₄ alkylamino, C₁-C₄ dialkylamino, trifluoromethylamino, ditrifluoromethylamino, aminomethanesulfonyl, amino C₁-C₄ alkylcarbonyl, aminotrifluoromethylcarbonyl, formylamino, nitro, nitroso, C₁-C₄ alkyldiazo, C₅-C₆ aryldiazo, trifluoromethyl, C₁-C₄ haloalkyl, C₁-C₄ cyanoalkyl, cyano, C₁-C₄ alkoxy carbonyl,

C_1 - C_4 alkylcarbonyl, phenyl, phenylcarbonyl, formyl, C_1 - C_4 alkoxyethyl, phenoxyethyl, C_2 - C_4 vinyl, C_2 - C_4 ethynyl, and SO_nR' wherein n is 0, 1, 2, or 3 and R' is hydrogen, C_1 - C_4 alkyl, amino, or phenyl.

2. The compound of claim 1, wherein R_1 is selected from the group consisting of amino, hydroxy, C_1 - C_4 alkylamino, C_1 - C_4 dialkylamino, and C_1 - C_4 alkylcarbonylamino, R_2 is selected from the group consisting of hydrogen, C_1 - C_4 alkyl, and C_1 - C_6 alkylcarbonyloxy C_1 - C_4 alkyl, and R_3 is selected from the group consisting of amino, halo, C_1 - C_4 alkyl, hydroxy, and trifluoromethyl.


3. The compound of claim 2, wherein R_1 is selected from the group consisting of amino, hydroxy, methylamino, dimethylamino, and acetylamino, R_2 is selected from the group consisting of hydrogen, methyl, and pivaloyloxymethyl, and R_3 is selected from the group consisting of amino, bromo, methyl, hydroxy, and trifluoromethyl.

4. The compound of claim 3, wherein said compound is selected from the group consisting of 8-amino- O^6 -benzylguanine, 8-methyl- O^6 -benzylguanine, 8-hydroxy- O^6 -benzylguanine, 8-bromo- O^6 -benzylguanine, 8-trifluoromethyl- O^6 -benzylguanine, O^6 -benzylxanthine, O^6 -benzyluric acid, N^2 -acetyl- O^6 -benzyl-8-oxoguanine, O^6 -benzyl-8-trifluoromethyl-9-methylguanine, O^6 -benzyl-8-bromo-9-methylguanine, and O^6 -benzyl-8-bromo-9-(pivaloyloxymethyl)-guanine.

5. The compound of claim 1, where R_1 is amino.

6. The compound of claim 1, wherein R_1 is hydroxy, C_1 - C_4 alkylamino, C_1 - C_4 dialkylamino, or C_1 - C_4 alkylcarbonylamino.


7. A compound of the formula



wherein R₁ is selected from the group consisting of hydrogen, halo, C₁-C₄ alkyl, halo C₁-C₄ alkyl, C₁-C₆ alkylcarbonyloxy C₁-C₄ alkyl, C₁-C₄ alkyloxycarbonyl C₁-C₄ alkyl, carboxy C₁-C₄ alkyl, cyano C₁-C₄ alkyl, aminocarbonyl C₁-C₄ alkyl, hydroxy C₁-C₄ alkyl, and C₁-C₄ alkyloxy C₁-C₄ alkyl, and R₂ is selected from the group consisting of C₁-C₄ alkyl, halo C₁-C₄ alkyl, C₁-C₆ alkylcarbonyloxy C₁-C₄ alkyl, C₁-C₄ alkyloxycarbonyl C₁-C₄ alkyl, carboxy C₁-C₄ alkyl, cyano C₁-C₄ alkyl, aminocarbonyl C₁-C₄ alkyl, hydroxy C₁-C₄ alkyl, and C₁-C₄ alkyloxy C₁-C₄ alkyl, with the proviso that when R₁ is hydrogen, R₂ is selected from the group consisting of halo C₁-C₄ alkyl, C₁-C₄ alkyloxy C₁-C₄ alkyl, C₁-C₆ alkylcarbonyloxy C₁-C₄ alkyl, carboxy C₂-C₄ alkyl, cyano C₂-C₄ alkyl, aminocarbonyl C₂-C₄ alkyl, and hydroxy C₁-C₃ alkyl.


8. The compound of claim 7, wherein said compound is selected from the group consisting of O⁶-benzyl-8-bromo-7-(pivaloyloxymethyl)guanine and O⁶-benzyl-7-(pivaloyloxymethyl)guanine.

9. A compound of the formula



wherein R_1 is hydroxy, R_2 is a substituent selected from the group consisting of hydrogen, $\text{C}_1\text{-C}_4$ alkyl, $\text{C}_1\text{-C}_4$ aminoalkyl, $\text{C}_1\text{-C}_4$ hydroxyalkyl, $\text{C}_1\text{-C}_4$ alkylamino $\text{C}_1\text{-C}_4$ alkyl, $\text{C}_1\text{-C}_4$ dialkylamino alkyl, $\text{C}_1\text{-C}_4$ cyanoalkyl, $\text{C}_1\text{-C}_4$ carbamoylalkyl, $\text{C}_1\text{-C}_4$ pivaloylalkyl, $\text{C}_1\text{-C}_6$ alkylcarbonyloxy $\text{C}_1\text{-C}_4$ alkyl, $\text{C}_1\text{-C}_4$ alkoxy carbonylalkyl, 2'-deoxyribose, the conjugate acid form of a $\text{C}_1\text{-C}_4$ carboxyalkyl, and the carboxylate anion of a $\text{C}_1\text{-C}_4$ carboxyalkyl as the sodium salt, and R_3 is a substituent selected from the group consisting of hydrogen, halo, $\text{C}_1\text{-C}_4$ alkyl, $\text{C}_1\text{-C}_4$ hydroxyalkyl, mercapto, $\text{C}_1\text{-C}_4$ alkylthio, trifluoromethylthio, $\text{C}_1\text{-C}_4$ alkylthiocarbonyl, hydroxy, $\text{C}_1\text{-C}_4$ alkoxy, trifluoromethoxy, methanesulfonyloxy, trifluoromethanesulfonyloxy, $\text{C}_1\text{-C}_4$ alkylcarbonyloxy, amino, $\text{C}_1\text{-C}_4$ aminoalkyl, $\text{C}_1\text{-C}_4$ alkylamino, $\text{C}_1\text{-C}_4$ dialkylamino, trifluoromethylamino, ditrifluoromethylamino, aminomethanesulfonyl, amino $\text{C}_1\text{-C}_4$ alkylcarbonyl, aminotrifluoromethylcarbonyl, formylamino, nitro, nitroso, $\text{C}_1\text{-C}_4$ alkyldiazo, $\text{C}_5\text{-C}_6$ aryldiazo, trifluoromethyl, $\text{C}_1\text{-C}_4$ haloalkyl, $\text{C}_1\text{-C}_4$ cyanoalkyl, cyano, $\text{C}_1\text{-C}_4$ alkyloxycarbonyl, $\text{C}_1\text{-C}_4$ alkylcarbonyl, phenyl, phenylcarbonyl, formyl, $\text{C}_1\text{-C}_4$ alkoxy methyl, phenoxy methyl, $\text{C}_2\text{-C}_4$ vinyl, $\text{C}_2\text{-C}_4$ ethynyl, and $\text{SO}_n\text{R}'$ wherein n is 0, 1, 2, or 3 and R' is hydrogen, $\text{C}_1\text{-C}_4$ alkyl, amino, or phenyl.

10. A compound of the formula



wherein R₁ is a substituent selected from the group consisting of amino, hydroxy, C₁-C₄ alkylamino, C₁-C₄ alkylcarbonylamino, and C₁-C₄ dialkylamino, R₂ is a substituent selected from the group consisting of C₁-C₄ aminoalkyl, C₁-C₄ alkylamino C₁-C₄ alkyl, C₁-C₄ dialkylamino alkyl, C₁-C₄ carbamoylalkyl, C₁-C₄ pivaloylalkyl, C₁-C₆ alkylcarbonyloxy C₁-C₄ alkyl, C₁-C₄ alkoxy carbonylalkyl, and 2'-deoxyribose, and R₃ is a substituent selected from the group consisting of hydrogen, halo, C₁-C₄ alkyl, C₁-C₄ hydroxyalkyl, mercapto, C₁-C₄ alkylthio, trifluoromethylthio, C₁-C₄ alkylthiocarbonyl, hydroxy, C₁-C₄ alkoxy, trifluoromethoxy, methanesulfonyloxy, trifluoromethanesulfonyloxy, C₁-C₄ alkylcarbonyloxy, amino, C₁-C₄ aminoalkyl, C₁-C₄ alkylamino, C₁-C₄ dialkylamino, trifluoromethylamino, ditrifluoromethylamino, aminomethanesulfonyl, amino C₁-C₄ alkylcarbonyl, aminotrifluoromethylcarbonyl, formylamino, nitro, nitroso, C₁-C₄ alkyldiazo, C₅-C₆ aryldiazo, trifluoromethyl, C₁-C₄ haloalkyl, C₁-C₄ cyanoalkyl, cyano, C₁-C₄ alkyloxycarbonyl, C₁-C₄ alkylcarbonyl, phenyl, phenylcarbonyl, formyl, alkoxy methyl, phenoxy methyl, C₂-C₄ vinyl, C₂-C₄ ethynyl, and SO_nR' wherein n is 0, 1, 2, or 3 and R' is hydrogen, C₁-C₄ alkyl, amino, or phenyl, with the provisos that R₁ is not

methylamino when R_2 is ribose or 2'-deoxyribose and R_3 is hydrogen and that when R_1 is amino and R_3 is hydrogen, R_2 is not alkylcarbonyloxyalkyl, alkoxy carbonylalkyl, or carbamoylalkyl.

11. A pharmaceutical composition comprising a pharmaceutically acceptable carrier and at least one compound of any one of claims 1 to 10.

12. The pharmaceutical composition of claim 11, wherein said pharmaceutically acceptable carrier comprises polyethylene glycol.


13. Use of a compound of any one of claims 1 to 10 in the manufacture of a medicament for enhancing the chemotherapeutic treatment of tumor cells in a mammal with an antineoplastic alkylating agent that causes cytotoxic lesions at the O^6 -position of guanine.

14. A product containing a compound of any one of claims 1 to 10 and an antineoplastic alkylating agent which causes cytotoxic lesions at the O^6 -position of guanine, for simultaneous, separate or sequential use for enhancing chemotherapeutic treatment of tumor cells in a mammal.

15. The pharmaceutical composition of claim 11, for enhancing the chemotherapeutic treatment of tumor cells in a mammal with an antineoplastic alkylating agent that causes cytotoxic lesions at the O^6 -position of guanine.

16. A method of inhibiting the reaction of O^6 -alkylguanine-DNA alkyltransferase in vitro with an alkylated DNA comprising reacting the O^6 -alkylguanine-DNA alkyltransferase with a compound of any one of claims 1 to 10.

17. An enhancing chemotherapeutic pharmaceutical composition for treatment of tumor cells in a mammal with an antineoplastic alkylating agent which causes cytotoxic lesions at the O^6 -position of guanine comprising a therapeutically effective amount of a compound defined in any one of claims 1 to 10 in association with a pharmaceutically acceptable carrier.



(I)