



HU000034251T2

(19) **HU****MAGYARORSZÁG****Szellemi Tulajdon Nemzeti Hivatala**(11) Lajstromszám: **E 034 251**(13) **T2**

EURÓPAI SZABADALOM SZÖVEGÉNEK FORDÍTÁSA

(21) Magyar ügyszám: **E 10 782040**(22) A bejelentés napja: **2010. 11. 01.**(51) Int. Cl.: **A61K 39/085** (2006.01)**A61K 39/385** (2006.01)

(96) Az európai bejelentés bejelentési száma:

EP 20100782040

(86) A nemzetközi (PCT) bejelentési szám:

PCT/IB 10/054934

(97) Az európai bejelentés közzétételi adatai:

EP 2493498 A1 **2011. 05. 05.**

(87) A nemzetközi közzétételi szám:

WO 11051917

(97) Az európai szabadalom megadásának meghirdetési adatai:

EP 2493498 B1 **2017. 03. 22.**

(30) Elsőbbségi adatok:

256905 P**2009. 10. 30.****US**

(73) Jogosult(ak):

**GlaxoSmithKline Biologicals SA, 1330
Rixensart (BE)**

(72) Feltaláló(k):

COSTANTINO, Paolo, I-53100 Siena (IT)**ROMANO, Maria Rosaria, I-53100 Siena (IT)****BERTI, Francesco, I-53100 Siena (IT)**

(74) Képviselő:

**Danubia Szabadalmi és Jogi Iroda Kft.,
Budapest**

(54)

Staphylococcus aureusból származó 5. típusú és 8. típusú kapszuláris szacharidok tisztítása

Az európai szabadalom ellen, megadásának az Európai Szabadalmi Közlönyben való meghirdetésétől számított kilenc hónapon belül, felszólalást lehet benyújtani az Európai Szabadalmi Hivatalnál. (Európai Szabadalmi Egyezmény 99. cikk(1))

A fordítást a szabadalmas az 1995. évi XXXIII. törvény 84/H. §-a szerint nyújtotta be. A fordítás tartalmi helyességét a Szellemi Tulajdon Nemzeti Hivatala nem vizsgálta.



(11)

EP 2 493 498 B1

(12)

EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention
of the grant of the patent:
22.03.2017 Bulletin 2017/12

(51) Int Cl.:
A61K 39/085 ^(2006.01) **A61K 39/385** ^(2006.01)

(21) Application number: **10782040.9**

(86) International application number:
PCT/IB2010/054934

(22) Date of filing: **01.11.2010**

(87) International publication number:
WO 2011/051917 (05.05.2011 Gazette 2011/18)

(54) **PURIFICATION OF STAPHYLOCOCCUS AUREUS TYPE 5 AND TYPE 8 CAPSULAR
SACCHARIDES**

REINIGUNG VON KAPSELFÖRMIGEN POLYSACCHARIDEN TYP 5 UND TYP 8 AUS
STAPHYLOCOCCUS AUREUS

PURIFICATION DE SACCHARIDES CAPSULAIRES DE STAPHYLOCOCCUS AUREUS DE TYPE
5 ET DE TYPE 8

(84) Designated Contracting States:
**AL AT BE BG CH CY CZ DE DK EE ES FI FR GB
GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO
PL PT RO RS SE SI SK SM TR**

(30) Priority: **30.10.2009 US 256905 P**

(43) Date of publication of application:
05.09.2012 Bulletin 2012/36

(73) Proprietor: **GlaxoSmithKline Biologicals SA
1330 Rixensart (BE)**

(72) Inventors:
• **COSTANTINO, Paolo**
I-53100 Siena (IT)
• **ROMANO, Maria Rosaria**
I-53100 Siena (IT)
• **BERTI, Francesco**
I-53100 Siena (IT)

(74) Representative: **Sampson, Catherine**
GlaxoSmithKline
Global Patents CN925.1
980 Great West Road
Brentford Middlesex TW8 9GS (GB)

(56) References cited:
WO-A1-99/42130 WO-A1-2005/000346

- **FATTOM A ET AL:** "Synthesis and immunologic properties in mice of vaccines composed of Staphylococcus aureus type 5 and type 8 capsular polysaccharides conjugated to Pseudomonas aeruginosa exotoxin A.", INFECTION AND IMMUNITY JUL 1990 LNKD-PUBMED:2114365, vol. 58, no. 7, July 1990 (1990-07), pages 2367-2374, XP002620550, ISSN: 0019-9567 cited in the application
- **GILBERT F B ET AL:** "Purification of type 5 capsular polysaccharide from Staphylococcus aureus by a simple efficient method", JOURNAL OF MICROBIOLOGICAL METHODS, ELSEVIER, AMSTERDAM, NL, vol. 20, no. 1, 1 June 1994 (1994-06-01), pages 39-46, XP023698993, ISSN: 0167-7012, DOI: DOI:10.1016/0167-7012(94)90062-0 [retrieved on 1994-06-01] cited in the application
- **CESCUTTI P ET AL:** "Determination of the size and degree of acetyl substitution of oligosaccharides from Neisseria meningitidis group A by ionspray mass spectrometry.", BIOCHEMICAL AND BIOPHYSICAL RESEARCH COMMUNICATIONS 16 JUL 1996 LNKD-PUBMED:8702408, vol. 224, no. 2, 16 July 1996 (1996-07-16), pages 444-450, XP002620551, ISSN: 0006-291X

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

EP 2 493 498 B1

- JENNINGS H: "Further approaches for optimizing polysaccharide-protein conjugate vaccines for prevention of invasive bacterial disease.", THE JOURNAL OF INFECTIOUS DISEASES JUN 1992 LNKD- PUBMED:1588152, vol. 165 Suppl 1, June 1992 (1992-06), pages S156-S159, XP009144149, ISSN: 0022-1899
- LEFEBER D J ET AL: "Isolation of oligosaccharides from a partial-acid hydrolysate of pneumococcal type 3 polysaccharide for use in conjugate vaccines", CARBOHYDRATE RESEARCH, PERGAMON, GB, vol. 337, no. 9, 30 April 2002 (2002-04-30), pages 819-825, XP004355355, ISSN: 0008-6215, DOI: DOI:10.1016/S0008-6215(02)00059-9
- DENG L ET AL: "CHARACTERIZATION OF THE LINKAGE BETWEEN THE TYPE III CAPSULAR POLYSACCHARIDE AND THE BACTERIAL CELL WALL OF GROUP B STREPTOCOCCUS", JOURNAL OF BIOLOGICAL CHEMISTRY.(MICROFILMS), AMERICAN SOCIETY OF BIOLOGICAL CHEMISTS, BALTIMORE, MD, US, vol. 275, no. 11, 17 March 2000 (2000-03-17), pages 7497-7504, XP008068869,
- LEE J C ET AL: "EFFECTS OF IN VITRO AND IN VIVO GROWTH CONDITIONS ON EXPRESSION OF TYPE 8 CAPSULAR POLYSACCHARIDE BY STAPHYLOCOCCUS AUREUS", INFECTION AND IMMUNITY, AMERICAN SOCIETY FOR MICROBIOLOGY, US, vol. 61, no. 5, 1 May 1993 (1993-05-01), pages 1853-1858, XP009051250, ISSN: 0019-9567
- FOURNIER J-M ET AL: "PURIFICATION AND CHARACTERIZATION OF STAPHYLOCOCCUS AUREUS TYPE 8 CAPSULAR POLYSACCHARIDE", INFECTION AND IMMUNITY, AMERICAN SOCIETY FOR MICROBIOLOGY, US, vol. 45, no. 1, 1 July 1984 (1984-07-01), pages 87-93, XP009051281, ISSN: 0019-9567
- Henning H.-G., W. Jugelt, G. Sauer: "Praktische Chemie", 1982, VEB Verlag Volk und Gesundheit, Berlin page 515, 540,

Description**TECHNICAL FIELD**

5 **[0001]** This invention is in the field of purifying bacterial capsular polysaccharides, particularly those of *Staphylococcus aureus* type 5 and type 8, and particularly for use in the preparation of vaccines.

BACKGROUND ART

10 **[0002]** The capsular saccharides of bacteria have been used for many years in vaccines against capsulated bacteria. As saccharides are T-independent antigens, however, they are poorly immunogenic. Conjugation to a carrier can convert T-independent antigens into T-dependent antigens, thereby enhancing memory responses and allowing protective immunity to develop. The most effective saccharide vaccines are therefore based on glycoconjugates, and the prototype conjugate vaccine was against *Haemophilus influenzae* type b ('Hib') [e.g. see chapter 14 of ref. 96].

15 **[0003]** Another bacterium for which conjugate vaccines have been described is *Staphylococcus aureus* (*S.aureus*). Various polysaccharides have been isolated from *S.aureus* for use in glycoconjugates. Two polysaccharides of particular interest are the type 5 and type 8 capsular polysaccharides. Approximately 60% of human *S.aureus* strains are type 8 and approximately 30% are type 5. Much of the work on type 5 and type 8 conjugates has been performed by Fattom *et al.*, and is described in documents such as references 1 to 9.

20 **[0004]** The starting point for polysaccharide-based vaccines is the polysaccharide itself, and this is generally purified from the target bacterium. Fattom *et al.* have developed a complex process for purification of the type 5 and type 8 capsular polysaccharides that is described in detail in reference 1, and involves the following key steps after bacterial culture: suspension of bacterial cells in buffer, treatment with lysostaphin, treatment with DNase and RNase, centrifugation, dialysis against buffer, treatment with protease, further dialysis, filtration, addition of ethanol to 25% with calcium chloride to precipitate contaminants; further addition of ethanol to 75% to precipitate the polysaccharide; collection and drying of the precipitate; anion exchange chromatography; dialysis; lyophilisation; size exclusion chromatography; dialysis and final lyophilisation.

25 **[0005]** The Fattom process involves the use of lysostaphin to lyse the bacterial cell walls and thereby release capsular polysaccharide. However, this step is time-consuming and makes the process difficult to scale-up to an industrial setting. It also increases the overall cost and complexity of the process. Other researchers have attempted to omit this step and develop a simpler, more efficient method of purifying the polysaccharide. For example, reference [10] describes an alternative process that involves autoclaving *S.aureus* cells, ultrafiltration of the polysaccharide-containing supernatant, concentration, lyophilisation, treatment with sodium metaperiodate, further ultrafiltration, diafiltration, high performance size exclusion liquid chromatography, dialysis and freeze-drying. The authors suggest that this method provides a good yield and is suitable for large scale production of polysaccharide. In this method, the lysostaphin treatment is replaced by autoclaving to release capsular polysaccharide. The method was further developed in reference [11]. An important step in these alternative methods is the treatment with sodium metaperiodate. This step is carried out to remove teichoic acid contamination of the capsular polysaccharide. However, once again this step increases the duration, complexity and overall cost of the process. Reference [12] describes a similar process that again involves autoclaving to release capsular polysaccharide and treatment with sodium metaperiodate to remove teichoic acid. In contrast, most other groups use processes that retain lysostaphin treatment (see, for example, references 13, 14, 15, 16, 17 and 18), sometimes including treatment with sodium metaperiodate (e.g. in references 13 and 14).

35 **[0006]** The above methods are complex and may leave contamination in the resultant polysaccharide. There is thus a need for further and improved processes for purifying *S.aureus* type 5 and type 8 capsular polysaccharides, and particularly for less complex processes that result in less contamination.

DISCLOSURE OF THE INVENTION

40 **[0007]** The invention is based on a purification process in which the polysaccharide is initially released from the bacterial cells by treatment with an acid. This step removes the need for lysostaphin treatment and can be used as an alternative to autoclaving, as in the above processes. The inventors have found that the process results in a purified polysaccharide with low teichoic acid contamination. This means that it is not necessary to treat the polysaccharide with sodium metaperiodate. The purified polysaccharide also has low peptidoglycan contamination, making it particularly suitable for medical uses. The inventors' process can be quick and simple because laborious steps in previous processes are not necessary.

45 **[0008]** We disclose a method for releasing capsular polysaccharide from *S.aureus* type 5 or type 8 cells, comprising the step of treating the cells with acid. Further disclosed is a process for purifying capsular polysaccharide from *S.aureus* type 5 or type 8 cells comprising this method. Other processing steps may be included in the process, such as enzymatic

treatment, e.g. to remove nucleic acid, protein and/or peptidoglycan contaminants; diafiltration, e.g. to remove low molecular weight contaminants; anion exchange chromatography, e.g. to remove residual protein; and concentration.

[0009] Accordingly, we disclose a process for purifying *S.aureus* type 5 or type 8 capsular polysaccharide, comprising the step of releasing the polysaccharide from *S.aureus* type 5 or type 8 cells by treating the cells with acid. Similarly, we disclose, in a process for purifying *S.aureus* type 5 or type 8 capsular polysaccharide, the improvement consisting of the use of acid treatment of *S.aureus* type 5 or type 8 cells to release the polysaccharide from the cells. Release by acid treatment removes the need for lysostaphin treatment or autoclaving to release the polysaccharide.

[0010] We also disclose a process for purifying *S.aureus* type 5 or type 8 capsular polysaccharide, wherein the process does not involve a step of lysostaphin treatment. Similarly, we disclose a process for purifying *S.aureus* type 5 or type 8 capsular polysaccharide, wherein the process does not involve a step of sodium metaperiodate treatment. Typically, the process does not involve one or both of these steps.

[0011] We disclose a process for purifying *S.aureus* type 5 or type 8 capsular polysaccharide, wherein the process provides a composition comprising the polysaccharide and a level of peptidoglycan contamination that is less than 5% (e.g. $\leq 4\%$, $\leq 3\%$, $\leq 2\%$, $\leq 1\%$, etc.) by weight peptidoglycan relative to the total weight of the polysaccharide. Typically, the composition comprises less than 4%, particularly less than 3%, by weight peptidoglycan. The inventors have found that levels of about 2% or even about 1% can be obtained using these methods. The inventors have found that compositions with this level of peptidoglycan are useful in vaccine manufacture. In contrast, reference 17 teaches that levels above 5% should be used for this purpose. The level of peptidoglycan contamination may be measured using the methods described herein.

[0012] Similarly, we disclose a process for purifying *S.aureus* type 5 or type 8 capsular polysaccharide, wherein the process provides a composition comprising the polysaccharide and a level of protein contamination that is less than 5% (e.g. $\leq 4\%$, $\leq 3\%$, $\leq 2\%$, $\leq 1\%$, $\leq 0.5\%$, etc.) by weight protein relative to the total weight of the polysaccharide. Typically, the composition comprises less than 3%, particularly about 2.4%, by weight protein. The level of protein contamination may be measured using a MicroBCA assay (Pierce).

[0013] Disclosed is a process for purifying *S.aureus* type 5 or type 8 capsular polysaccharide, wherein the process provides a composition comprising the polysaccharide and a level of nucleic acid contamination that is less than 1% (e.g. $\leq 0.75\%$, $\leq 0.50\%$, $\leq 0.25\%$, $\leq 0.10\%$, $\leq 0.01\%$, etc.) by weight nucleic acid relative to the total weight of the polysaccharide. Typically, the composition comprises less than 0.25%, particularly about 0.09%, by weight nucleic acid. The level of nucleic acid contamination may be measured by absorption at 260nm in a spectrophotometer.

[0014] A process for purifying *S.aureus* type 5 or type 8 capsular polysaccharide, wherein (a) the level of peptidoglycan acid contamination is less than 5% (as described above); (b) the level of protein contamination is less than 5% (as described above); (c) the level of nucleic acid contamination that is less than 1% (as described above), is disclosed.

[0015] A composition comprising a *S.aureus* type 5 or type 8 capsular polysaccharide, obtainable by any of the processes of the invention, is also disclosed.

[0016] We disclose a composition comprising *S.aureus* type 5 or type 8 capsular polysaccharide, wherein the composition comprises a level of peptidoglycan contamination that is less than 5% (e.g. $\leq 4\%$, $\leq 3\%$, $\leq 2\%$, $\leq 1\%$, etc.) by weight peptidoglycan relative to the total weight of the polysaccharide. Typically, the composition comprises less than 3%, particularly less than 2%, by weight peptidoglycan. Compositions with levels of about 2% or even about 1% are specifically disclosed.

[0017] Similarly, we disclose a composition comprising *S.aureus* type 5 or type 8 capsular polysaccharide, wherein the composition comprises a level of protein contamination that is less than 5% (e.g. $\leq 4\%$, $\leq 3\%$, $\leq 2\%$, $\leq 1\%$, $\leq 0.5\%$, etc.) by weight protein relative to the total weight of the polysaccharide. Typically, the composition comprises less than 3%, particularly about 2.4%, by weight protein.

[0018] We disclose a composition comprising *S.aureus* type 5 or type 8 capsular polysaccharide, wherein the composition comprises a level of nucleic acid contamination that is less than 1% (e.g. $\leq 0.75\%$, $\leq 0.50\%$, $\leq 0.25\%$, $\leq 0.10\%$, $\leq 0.01\%$, etc.) by weight nucleic acid relative to the total weight of the polysaccharide. Typically, the composition comprises less than 0.25%, particularly about 0.09%, by weight nucleic acid.

[0019] We disclose a composition comprising *S.aureus* type 5 or type 8 capsular polysaccharide, wherein a) a level of peptidoglycan acid contamination is less than 5% (as described above); (b) the level of protein contamination is less than 5% (as described above); (c) the level of nucleic acid contamination that is less than 1% (as described above).

[0020] The invention provides a method for making an immunogenic composition comprising an *S. aureus* type 5 capsular polysaccharide conjugated to a carrier molecule and an *S. aureus* type 8 capsular polysaccharide conjugated to a carrier molecule, said method comprising:

- (a) a process for purifying capsular polysaccharide from *S. aureus* type 5 cells comprising a step of releasing the capsular polysaccharide from the cells by acid treatment of the cells, wherein the cells are in the form of a wet cell paste or are suspended in aqueous medium, and the molecular mass of the purified polysaccharide is between 2-3500kDa;

- (b) conjugation of the capsular polysaccharide to a carrier molecule to make a conjugate; and
- (c) mixing the conjugate with an *S. aureus* type 8 capsular polysaccharide conjugated to a carrier molecule.

[0021] The invention further provides a method for making an immunogenic composition comprising an *S. aureus* type 8 capsular polysaccharide conjugated to a carrier molecule and an *S. aureus* type 5 capsular polysaccharide conjugated to a carrier molecule, said method comprising:

- (a) a process for purifying capsular polysaccharide from *S. aureus* type 8 cells comprising a step of releasing the capsular polysaccharide from the cells by acid treatment of the cells, wherein the cells are in the form of a wet cell paste or are suspended in an aqueous medium, and the molecular mass of the purified polysaccharide is between 2-3500 kDa;
- (b) conjugation of the capsular polysaccharide to a carrier molecule to make a conjugate; and
- (c) mixing the conjugate with an *S. aureus* type 5 capsular polysaccharide conjugated to a carrier molecule.

The capsular polysaccharide

[0022] The invention is based on the capsular polysaccharides of *S. aureus* type 5 and type 8. The structures of type 5 and type 8 capsular polysaccharides were described in references 19 and 20 as:

Type 5 → 4)-β-D-ManNAcA(3OAc)-(1 → 4)-α-L-FucNAc(1 → 3)-β-D-FucNAc-(1 →
Type 8 → 3)-β-D-ManNAcA(4OAc)-(1 → 3)-α-L-FucNAc(1 → 3)-β-D-FucNAc-(1 →.

[0023] Recent NMR spectroscopy data [21] has led to a revision of these structures to:

Type 5 → 4)-β-D-ManNAcA-(1 → 4)-α-L-FucNAc(3OAc)-(1 → 3)-β-D-FucNAc-(1 →
Type 8 → 3)-β-D-ManNAcA(4OAc)-(1 → 3)-α-L-FucNAc(1 → 3)-α-D-FucNAc(1 →.

[0024] After release from the *S. aureus* type 5 or type 8 cells, the polysaccharide may be chemically modified relative to the capsular polysaccharide as found in nature. For example, the polysaccharide may be de-O-acetylated (partially or fully), de-N-acetylated (partially or fully), N-propionated (partially or fully), etc. De-acetylation may occur before, during or after other processing steps, but typically occurs before any conjugation step. Depending on the particular polysaccharide, de-acetylation may or may not affect immunogenicity e.g. the NeisVac-C™ vaccine uses a de-O-acetylated polysaccharide, whereas Menjugate™ is acetylated, but both vaccines are effective. The effect of de-acetylation etc. can be assessed by routine assays. For example, the relevance of O-acetylation on *S. aureus* type 5 or type 8 capsular polysaccharides is discussed in reference 6. The native polysaccharides are said in this document to have 75% O-acetylation. These polysaccharides induced antibodies to both the polysaccharide backbone and O-acetyl groups. Polysaccharides with 0% O-acetylation still elicited antibodies to the polysaccharide backbone. Both types of antibody were opsonic against *S. aureus* strains that varied in their O-acetyl content. Accordingly, the type 5 or type 8 capsular polysaccharides used in the present invention may have between 0 and 100% O-acetylation. For example, the degree of O-acetylation of the type 5 capsular polysaccharide may be 10-100%, 10-100%, 20-100%, 30-100%, 40-100%, 50-100%, 60-100%, 70-100%, 80-100%, 90-100%, 50-90%, 60-90%, 70-90% or 80-90%. Alternatively, 0% O-acetylated type 5 capsular polysaccharide may be used. Similarly, the degree of O-acetylation of the type 8 capsular polysaccharide may be 10-100%, 10-100%, 20-100%, 30-100%, 40-100%, 50-100%, 60-100%, 70-100%, 80-100%, 90-100%, 50-90%, 60-90%, 70-90% or 80-90%. Alternatively, 0% O-acetylated type 8 capsular polysaccharide may be used. In one embodiment, the degree of O-acetylation of the type 5 and type 8 capsular polysaccharides may be 10-100%, 20-100%, 30-100%, 40-100%, 50-100%, 60-100%, 70-100%, 80-100%, 90-100%, 50-90%, 60-90%, 70-90% or 80-90%. In other embodiments, 0% O-acetylated type 5 and type 8 capsular polysaccharides are used. The degree of N-acetylation of the type 5 capsular polysaccharide used in the invention may be 0-100%, 50-100%, 75-100%, 80-100%, 90-100%, or 95-100%. Typically, the degree of N-acetylation of the type 5 capsular polysaccharide is 100%. Similarly, the degree of N-acetylation of the type 8 capsular polysaccharide used in the invention may be 0-100%, 50-100%, 75-100%, 80-100%, 90-100%, or 95-100%. Typically, the degree of N-acetylation of the type 8 capsular polysaccharide is 100%. In one embodiment, the degree of N-acetylation of the type 5 and type 8 capsular polysaccharides may be 0-100%, 50-100%, 75-100%, 80-100%, 90-100%, or 95-100%. Typically, the degree of N-acetylation of the type 5 and type 8 capsular polysaccharides are 100%.

[0025] The degree of O-acetylation of the polysaccharide can be determined by any method known in the art, for example, by proton NMR (e.g. as described in references 22, 23, 24 or 25). A further method is described in reference 26. Similar methods may be used to determine the degree of N-acetylation of the polysaccharide. O-acetyl groups may

be removed by hydrolysis, for example by treatment with a base such as anhydrous hydrazine [27] or NaOH [6]. Similar methods may be used to remove N-acetyl groups. To maintain high levels of O-acetylation on type 5 and/or 8 capsular polysaccharides, treatments that lead to hydrolysis of the O-acetyl groups are minimised, e.g. treatments at extremes of pH.

Starting material

[0026] The process of the invention starts with *S.aureus* type 5 or type 8 cells. Typically, the cells are grown by fermentation prior to release of capsular polysaccharide. Suitable methods of cultivating *S.aureus* type 5 or type 8 cells are well known to the skilled person and are disclosed, for example, in references 1 to 21 and the references cited therein. After cell growth, the cells are usually deactivated. A suitable method for deactivation is treatment with phenol:ethanol, e.g. as described in reference 1.

[0027] The cells may be centrifuged prior to release of capsular polysaccharide. The process may therefore start with the cells in the form of a wet cell paste. Typically, however, the cells are resuspended in an aqueous medium that is suitable for the next step in the process, e.g. in a buffer or in distilled water. The cells may be washed with this medium prior to re-suspension. In another embodiment, the cells may be treated in suspension in their original culture medium. Alternatively, the cells are treated in a dried form.

Acid treatment

[0028] In the method of the invention and methods disclosed herein, *S.aureus* type 5 or type 8 cells are treated with acid. This step results in release of capsular polysaccharide from the cells. In contrast, previous methods have used lysostaphin treatment or autoclaving to release the polysaccharide. The acid treatment of the invention and as disclosed herein is preferably carried out using a mild acid, e.g. acetic acid, to minimise damage to the polysaccharide. The skilled person would be capable of identifying suitable acids and conditions (e.g. of concentration, temperature and/or time) for release of the polysaccharide. For example, the inventors have found that treatment of cells suspended at about 0.5mg/ml in distilled water with 1% acetic acid (v/v) at 100°C for 2 hours is suitable. Treatment with other acids, e.g. trifluoroacetic or other organic acids, may also be suitable.

[0029] The efficacy of different acid treatments may be tested using routine methods. For example, after acid treatment, the cells may be isolated and treated using known methods of *S.aureus* type 5 or type 8 capsular polysaccharide release (e.g. the lysostaphin-based method of reference 1) to see if additional capsular polysaccharide can be released. If additional capsular polysaccharide is released, then the acid treatment conditions may be altered so that a greater proportion of the capsular saccharide is released during acid treatment. In this way, it is possible to optimise the acid treatment conditions so that an optimal amount of capsular saccharide is released. For example, the inventors have found that after treatment of cells suspended at about 0.5mg/ml in distilled water with 1% acetic acid (v/v) at 100°C for 2 hours, very little additional capsular saccharide is releasable from the cells by subsequent lysostaphin treatment.

[0030] The inventors have found that after acid treatment, the degree of O-acetylation of the type 5 capsular polysaccharide may be between 60-100%. In particular, the degree of O-acetylation may be between the 65-95%, particularly 70-90%. Typically, the degree of O-acetylation is between 75-85%, e.g. about 80%. Similar values may be obtained for the type 8 capsular saccharide. If desired, the degree of O-acetylation of the capsular saccharide may then be altered by further processing steps as discussed above.

[0031] After acid treatment, the reaction mixture is typically neutralised. This may be achieved by the addition of a base, e.g. NaOH. The cells may be centrifuged and the polysaccharide-containing supernatant collected for storage and/or additional processing.

Enzymatic treatment

[0032] The polysaccharide obtained after acid treatment may be impure and contaminated with bacterial nucleic acids and proteins. These contaminants may be removed by enzymatic treatment. For example, RNA may be removed by treatment with RNase, DNA with DNase and protein with protease (e.g. pronase). The skilled person would be capable of identifying suitable enzymes and conditions for removal of the contaminants. For example, the inventors have found that treatment of polysaccharide-containing supernatant with 50µg/ml each of DNase and RNase at 37°C for 6-8 hours is suitable. Other suitable conditions are disclosed in the literature, e.g. in reference 1.

[0033] The polysaccharide obtained after acid treatment may also or alternatively be contaminated with peptidoglycan. This contaminant may also be removed by enzymatic treatment. The inventors have found that treatment with mutanolysin is effective at removing peptidoglycan contamination. The skilled person would be capable of identifying suitable conditions for removal of the peptidoglycan with mutanolysin. For example, the inventors have found that treatment of polysaccharide-containing supernatant with 180U/ml each of mutanolysin at 37°C for 16 hours is suitable. After treatment,

the suspension may be clarified by centrifugation and the polysaccharide-containing supernatant collected for storage and/or additional processing.

Diafiltration

[0034] The process of the invention may involve a step of diafiltration. This step is typically performed after the acid treatment and/or enzymatic treatment discussed above. The inventors have found that a diafiltration step, particularly by tangential flow filtration, is particularly effective for removing impurities from the polysaccharide. The impurities are typically low molecular weight contaminants like teichoic and/or peptidoglycan fragments. The tangential flow filtration is suitably carried out against 1M NaCl (e.g. against about 10 volumes) and then NaPi 10mM pH 7.2 buffer (e.g. against another 10 volumes). The filtration membrane should thus be one that allows passage of small molecular weight contaminants while retaining the capsular polysaccharide. A cut-off in the range 10kDa-30kDa is typical. The inventors have found that tangential flow filtration using a 30kDa cut-off membrane is particularly suitable for large-scale processes.

[0035] At least 5 cycles of tangential flow diafiltration are usually performed e.g. 6, 7, 8, 9, 10, 11 or more.

[0036] The polysaccharide-containing retentate from the diafiltration is collected for storage and/or additional processing.

Anion exchange chromatography

[0037] The polysaccharide may be further purified by a step of anion exchange chromatography. The inventors have found that anion exchange chromatography is particularly effective at removing residual protein and nucleic acid contamination, while maintaining a good yield of the polysaccharide.

[0038] The anion exchange chromatography step may be performed after the acid treatment, enzymatic treatment and/or diafiltration steps discussed above.

[0039] The anion exchange chromatography may be carried out using any suitable anionic exchange matrix. Commonly used anion exchange matrices are resins such as Q-resins (based on quaternary amines) and DEAE resins (based on diethylaminoethane). The inventors have found that DEAE-resins (e.g. a DEAE-Sepharose™ Fast Flow resin (GE Healthcare)) are particularly suitable, although other resins may be used.

[0040] Appropriate starting buffers and mobile phase buffers for the anion exchange chromatography can also be determined by routine experiments without undue burden. Typical buffers for use in anion exchange chromatography include N-methyl piperazine, piperazine, L-histidine, bis-Tris, bis-Tris propane, triethanolamine, Tris, N-methyl-diethanolamine, diethanolamine, 1,3-diaminopropane, ethanolamine, piperidine, sodium chloride and phosphate buffers. The inventors have found that phosphate buffers, e.g. a sodium phosphate buffer, are suitable as the starting buffer for the anion exchange chromatography. The buffer may be at any suitable concentration. For example, 10 mM sodium phosphate has been found to be suitable. Material bound to the anionic exchange resin may be eluted with a suitable buffer. The inventors have found that a gradient of NaCl 1M is suitable.

[0041] Eluate fractions containing polysaccharide may be determined by measuring UV absorption at 215 nm. Fractions containing polysaccharide, usually combined together, are collected for storage and/or additional processing.

[0042] The anion exchange chromatography step may be repeated, e.g. 1, 2, 3, 4 or 5 times. Typically the anion exchange chromatography step is carried out once.

Gel filtration

[0043] The process of the invention may involve one or more step(s) of gel filtration. This gel filtration is used to select polysaccharide molecules of a particular length and to further reduce contamination, particularly by proteins. However, the inventors have found that contrary to previous methods like those of references 1 to 9, a gel filtration step is not required to obtain polysaccharide of high purity. Accordingly, this step may be omitted from the processes of the invention. The omission of this step is advantageous because it simplifies the process and reduces the overall cost.

[0044] When present, the gel filtration step(s) may be performed after the acid treatment, enzymatic treatment, diafiltration and/or anion exchange chromatography steps discussed above. Typically, any gel filtration step(s) are carried out after the anion exchange chromatography step discussed above.

[0045] The gel filtration step(s) may be carried out using any suitable gel filtration matrix. Commonly used gel filtration matrices are based on dextran gels, agarose gels, polyacrylamide gels, polyacryloylmorpholine gels, and polystyrene gels etc. Cross-linked dextran gels and mixed polyacrylamide/agarose gels may also be used. The inventors have found that dextran gels (e.g. a Sephacryl™ S300 gel (GE Healthcare)) are particularly suitable, although other gels may be used.

[0046] Appropriate mobile phase buffers for the gel filtration can be determined by routine experiments without undue burden. Typical buffers for use in gel filtration include N-methyl piperazine, piperazine, L-histidine, bis-Tris, bis-Tris propane, triethanolamine, Tris, N-methyl-diethanolamine, diethanolamine, 1,3-diaminopropane, ethanolamine, piperid-

ine, sodium chloride and phosphate buffers. For example, sodium chloride buffers may be suitable. The buffer may be at any suitable concentration. For example, 50 mM sodium chloride may be used for the mobile phase.

[0047] Eluate fractions containing polysaccharide may be determined by measuring UV absorption at 215 nm. Fractions containing polysaccharide, usually combined together, are collected for storage and/or additional processing.

Concentration

[0048] In addition to, or instead of, the one or more step(s) of gel filtration, the process of the invention may involve one or more steps of concentrating the polysaccharide. This concentration is useful for obtaining a sample of the correct concentration for any subsequent conjugation of the polysaccharide to a carrier molecule, as described below. However, the inventors have found that this concentration step is not required to obtain polysaccharide of high purity. Accordingly, this step may be omitted from the processes of the invention.

[0049] When present, the concentration step(s) may be performed after the acid treatment, enzymatic treatment, diafiltration, anion exchange chromatography and/or gel filtration steps discussed above. Typically, any concentration step(s) are carried out after the anion exchange chromatography step discussed above. If used in addition to the gel filtration step(s) discussed above, the concentration step(s) may be carried out before or after the gel filtration step(s) discussed above. However, typically, concentration step(s) are used instead of gel filtration step(s).

[0050] The concentration step(s) may be carried out by any suitable method. For example, the inventors have found that the concentration step(s) may be diafiltration step(s) as described above, for example tangential flow filtration using a 30kDa cut-off membrane. For example, a Hydrosart™ (Sartorius) 30kDa cut-off membrane (with a 200 cm² membrane area) may be used.

[0051] The concentrated polysaccharide sample is collected for storage and/or additional processing.

Further treatment of the capsular polysaccharide

[0052] After purification, the polysaccharide may be further treated to remove contaminants. This is particularly important in situations where even minor contamination is not acceptable (e.g. for human vaccine production).

[0053] The molecular mass of the purified *S.aureus* type 5 or type 8 capsular polysaccharide can be measured by gel filtration relative to pullulan standards, such as those available from Polymer Standard Service [28]. Typically, the purified polysaccharide is a mixture of polysaccharides with masses within a range of values. For the type 5 capsular polysaccharide, the molecular mass of the purified polysaccharide is between 2-3500 kDa, e.g. between 10-2000 kDa, particularly between 20-1000 kDa and more particularly between 100-600 kDa. Similarly, for the type 8 capsular polysaccharide, the molecular mass of the purified polysaccharide is between 2-3500 kDa, e.g. between 10-2000 kDa, particularly between 20-1000 kDa and more particularly between 100-600 kDa.

[0054] The purified polysaccharide may be depolymerised to form an oligosaccharide. Oligosaccharides may be preferred for use in vaccines. Depolymerisation to oligosaccharide may occur before or after any of the steps mentioned above. Typically, depolymerisation takes place after the anion exchange chromatography described above. If the polysaccharide is concentrated after this chromatography, then depolymerisation typically takes place after this concentration. Where the composition of the invention includes a depolymerised polysaccharide, it is preferred that depolymerisation precedes any conjugation. Full-length polysaccharides may be depolymerised to give shorter fragments for use in the invention by various methods. Preferably, the method described in reference 29 is used. Alternatively, other methods for depolymerisation of the polysaccharide may be used. For example, the polysaccharide may be heated or subjected to microfluidisation [30] or sonic radiation [3]. Alternatively, depolymerisation by oxidation-reduction [31] or ozonolysis [32] may be used.

[0055] Oligosaccharides can be identified by chromatography, e.g. size exclusion chromatography. The products may be sized in order to remove short-length oligosaccharides. This can be achieved in various ways, such as gel filtration. Specific molecular masses can be measured by gel filtration relative to pullulan standards, such as those available from Polymer Standard Service [33].

[0056] If N-acetyl groups in the native capsular polysaccharide have been de-N-acetylated then the processes of the invention may include a step of re-N-acetylation. Controlled re-N-acetylation can conveniently be performed using a reagent such as acetic anhydride (CH₃CO)₂O e.g. in 5% ammonium bicarbonate [34].

[0057] Further rounds of filtration, e.g. sterile filtration, can also be performed.

[0058] These additional steps can generally be performed at room temperature.

Storage

[0059] The *S.aureus* type 5 or type 8 capsular polysaccharide preparation may be lyophilised, e.g. by freeze-drying under vacuum, or frozen in solution (e.g. as the eluate from the final concentration step, if included) for storage at any

stage during the purification process. Accordingly, it is not necessary for the preparation to be transferred immediately from one step of the process to another. For example, if the polysaccharide preparation is to be purified by diafiltration, then it may be lyophilised or frozen in solution prior to this purification. Similarly, the polysaccharide may be lyophilised or frozen in solution prior to the anion exchange chromatography step. If the polysaccharide preparation is to be purified by gel filtration, then it may be lyophilised or frozen in solution prior to this step. Similarly, if the polysaccharide preparation is to be concentrated, then it may be lyophilised or frozen in solution prior to this step. The lyophilised preparation is reconstituted in an appropriate solution prior to further treatment. Similarly, the frozen solution is defrosted prior to further treatment.

[0060] The purified polysaccharide obtained by the process of the invention may be processed for storage in any suitable way. For example, the polysaccharide may be lyophilised as described above. Alternatively, the polysaccharide may be stored in aqueous solution, typically at low temperature, e.g. at -20°C. Conveniently, the polysaccharide may be stored as the eluate from the anion exchange chromatography, gel filtration or concentration steps.

Conjugation

[0061] The final purified capsular polysaccharide of the invention can be used as an antigen without further modification e.g. for use in *in vitro* diagnostic assays, for use in immunisation, etc.

[0062] For immunisation purposes, however, it is preferred to conjugate the polysaccharide to a carrier molecule, such as a protein. In general, covalent conjugation of polysaccharides to carriers enhances the immunogenicity of polysaccharides as it converts them from T-independent antigens to T-dependent antigens, thus allowing priming for immunological memory. Conjugation is particularly useful for paediatric vaccines [e.g. ref. 35] and is a well known technique [e.g. reviewed in refs. 36 to 44]. Thus the processes of the invention may include the further step of conjugating the purified polysaccharide to a carrier molecule.

[0063] Conjugation of *S.aureus* type 5 and type 8 capsular polysaccharides has been widely reported e.g. see references 1 to 9. The typical process used in the literature for conjugation involves thiolation of a purified polysaccharide using cystamine. The reaction relies on the presence of carboxylate groups in the capsular polysaccharide. These groups react with cystamine in the presence of a carbodiimide, e.g. EDAC. The derivatised polysaccharide is then conjugated to a carrier protein such as the *Pseudomonas aeruginosa* endotoxin A (ETA), typically via a linker [2]. Conjugate vaccines prepared in this manner have been shown to be safe and immunogenic in humans [5]. Other researchers have carried out conjugation of purified type 5 and type 8 capsular polysaccharides by reductive amination [45 and 12]; glutaraldehyde coupling [45]; or reaction of hydroxyl groups on the polysaccharides with cyanating agents like CDAP [46] or cyanuric trichloride [11]. Preferably, the process described in reference 29 is used.

[0064] Preferred carrier proteins are bacterial toxins, such as diphtheria or tetanus toxins, or toxoids or mutants thereof. The inventors have found that the CRM197 diphtheria toxin mutant [47] is suitable. *Pseudomonas aeruginosa* exotoxin A (ETA) and its non-toxic mutant recombinant exoprotein A (rEPA) have been used as carrier proteins for *S.aureus* type 5 or type 8 capsular polysaccharides ([1] and [2]). *S.aureus* α -haemolysin (α -toxin) ([45] and [48]), ovalbumin [11] and human serum albumin [12] have also been used. These carriers may be used in the present invention.

[0065] Other suitable carrier proteins include the *N.meningitidis* outer membrane protein complex [49], synthetic peptides [50,51], heat shock proteins [52,53], pertussis proteins [54,55], cytokines [56], lymphokines [56], hormones [56], growth factors [56], human serum albumin (typically recombinant), artificial proteins comprising multiple human CD4⁺ T cell epitopes from various pathogen-derived antigens [57] such as N19 [58], protein D from *H.influenzae* [59-61], pneumococcal surface protein PspA [62], pneumolysin [63] or its non-toxic derivatives [64], iron-uptake proteins [65], toxin A or B from *C.difficile* [66], a GBS protein [67], a GAS protein [68] etc.

[0066] Other suitable carrier proteins include *S.aureus* protein antigens, for example the *S.aureus* protein antigens set out below.

[0067] Attachment to the carrier is preferably via a -NH₂ group e.g. in the side chain of a lysine residue in a carrier protein, or of an arginine residue. Attachment may also be via a -SH group e.g. in the side chain of a cysteine residue.

[0068] It is possible to use more than one carrier protein e.g. to reduce the risk of carrier suppression. Thus different carrier proteins can be used for the type 5 and type 8 capsular polysaccharides, e.g. type 5 polysaccharide might be conjugated to CRM197 while type 8 polysaccharide might be conjugated to rEPA. It is also possible to use more than one carrier protein for a particular polysaccharide antigen e.g. type 5 polysaccharide might be in two groups, with one group conjugated to CRM197 and the other conjugated to rEPA. Typically, however, the same carrier protein is used for all polysaccharides.

[0069] A single carrier protein might carry more than one polysaccharide antigen [69,70]. For example, a single carrier protein might have conjugated to it type 5 and type 8 capsular polysaccharides. To achieve this goal, different polysaccharides can be mixed prior to the conjugation process. Typically, however, there are separate conjugates for each polysaccharide, with the different polysaccharides being mixed after conjugation. The separate conjugates may be based on the same carrier.

[0070] Conjugates with a polysaccharide:protein ratio (w/w) of between 1:20 (*i.e.* excess protein) and 20:1 (*i.e.* excess polysaccharide) are typically used. Ratios of 1:10 to 1:1 are preferred, particularly ratios between 1:5 and 1:2 and, most preferably, about 1:3. In contrast, type 5 and type 8 capsular polysaccharide conjugates used in the literature tend to have higher ratios, *e.g.* between 0.73 and 1.08 in references 1, 2 and 3. In particular embodiments of the invention, the polysaccharide:protein ratio (w/w) for type 5 capsular polysaccharide conjugate is between 1:10 and 1:2; and/or the polysaccharide:protein ratio (w/w) for type 8 capsular polysaccharide conjugate is between 1:5 and 7:10.

[0071] Conjugates may be used in conjunction with free carrier [71]. When a given carrier protein is present in both free and conjugated form in a composition of the invention, the unconjugated form is preferably no more than 5% of the total amount of the carrier protein in the composition as a whole, and more preferably present at less than 2% by weight.

[0072] After conjugation, free and conjugated polysaccharides can be separated. There are many suitable methods, including hydrophobic chromatography, tangential ultrafiltration, diafiltration *etc.* [see also refs. 72 & 73, *etc.*].

Combinations of conjugates and other antigens

[0073] The methods of the invention are for making an immunogenic composition comprising an *S. aureus* type 5 capsular polysaccharide conjugated to a carrier molecule and an *S. aureus* type 8 capsular polysaccharide conjugated to a carrier molecule.

[0074] Polysaccharides prepared by the methods as disclosed herein (in particular after conjugation as described above) can be mixed *e.g.* with each other and/or with other antigens. The processes disclosed herein may include the further step of mixing the polysaccharide with one or more further antigens. We disclose a composition comprising a polysaccharide prepared by the method as disclosed and one or more further antigens. The composition is typically an immunogenic composition.

[0075] The further antigen(s) may comprise further polysaccharides prepared by the disclosed methods, and so a composition comprising more than one polysaccharide as disclosed is also disclosed. In particular, the present invention provides a composition comprising a type 5 capsular polysaccharide of the invention and a type 8 capsular polysaccharide of the invention. We disclose that, the further antigen(s) may be type 5 or type 8 capsular polysaccharides prepared by methods other than those disclosed herein, *e.g.* the methods of references 1 to 18 above. Accordingly, we disclose a composition comprising a type 5 capsular polysaccharide and a type 8 capsular polysaccharide, wherein one of the polysaccharides (the type 5 polysaccharide or the type 8 polysaccharide) is a polysaccharide as disclosed and the other polysaccharide is not a polysaccharide as disclosed.

[0076] Where multiple different *S. aureus* conjugates are mixed then these may include different types of conjugate from the same *S. aureus* serotype and/or conjugates from different *S. aureus* serotypes. For example, the conjugates may be from *S. aureus* type 5 and type 8. The composition will be produced by preparing separate conjugates (*e.g.* a different conjugate for each serotype) and then combining the conjugates.

[0077] Further antigen(s) may comprise other *S. aureus* antigens, including the saccharide and protein antigens set out below.

[0078] The further antigen(s) may comprise antigens from non-*S. aureus* pathogens. Thus the disclosed compositions and the composition of the invention may further comprise one or more non-*S. aureus* antigens, including additional bacterial, viral or parasitic antigens. These may be selected from the following:

- a protein antigen from *N. meningitidis* serogroup B, such as those in refs. 74 to 80, with protein '287' (see below) and derivatives (*e.g.* 'ΔG287') being particularly preferred.
- an outer-membrane vesicle (OMV) preparation from *N. meningitidis* serogroup B, such as those disclosed in refs. 81, 82, 83, 84 *etc.*
- a saccharide antigen from *N. meningitidis* serogroup A, C, W135 and/or Y, such as the oligosaccharide disclosed in ref. 85 from serogroup C or the oligosaccharides of ref. 86.
- a saccharide antigen from *Streptococcus pneumoniae* [*e.g.* refs. 87-89; chapters 22 & 23 of ref. 96].
- an antigen from hepatitis A virus, such as inactivated virus [*e.g.* 90, 91; chapter 15 of ref. 96].
- an antigen from hepatitis B virus, such as the surface and/or core antigens [*e.g.* 91,92; chapter 16 of ref. 96].
- an antigen from hepatitis C virus [*e.g.* 93].
- an antigen from *Bordetella pertussis*, such as pertussis holotoxin (PT) and filamentous haemagglutinin (FHA) from *B. pertussis*, optionally also in combination with pertactin and/or agglutinogens 2 and 3 [*e.g.* refs. 94 & 95; chapter 21 of ref. 96].
- a diphtheria antigen, such as a diphtheria toxoid [*e.g.* chapter 13 of ref. 96].
- a tetanus antigen, such as a tetanus toxoid [*e.g.* chapter 27 of ref. 96].
- a saccharide antigen from *Haemophilus influenzae* B [*e.g.* chapter 14 of ref. 96]
- an antigen from *N. gonorrhoeae* [*e.g.* 74, 75, 76].
- an antigen from *Chlamydia pneumoniae* [*e.g.* 97, 98, 99, 100, 101, 102, 103].

- an antigen from *Chlamydia trachomatis* [e.g. 104].
- an antigen from *Porphyromonas gingivalis* [e.g. 105].
- polio antigen(s) [e.g. 106, 107; chapter 24 of ref. 96] such as IPV.
- rabies antigen(s) [e.g. 108] such as lyophilised inactivated virus [e.g. 109, RabAvert™].
- measles, mumps and/or rubella antigens [e.g. chapters 19, 20 and 26 of ref. 96].
- influenza antigen(s) [e.g. chapters 17 & 18 of ref. 96], such as the haemagglutinin and/or neuraminidase surface proteins.
- an antigen from *Moraxella catarrhalis* [e.g. 110].
- an antigen from *Streptococcus pyogenes* (group A streptococcus) [e.g. 111, 112, 113].
- an antigen from *Streptococcus agalactiae* (group B streptococcus) [e.g. 68, 114-116].
- an antigen from *S. epidermidis* [e.g. type I, II and/or III capsular polysaccharide obtainable from strains ATCC-31432, SE-360 and SE-10 as described in refs. 117, 118 and 119].

[0079] Where a saccharide or carbohydrate antigen is used, it is preferably conjugated to a carrier in order to enhance immunogenicity. Conjugation of *H. influenzae* B, meningococcal and pneumococcal saccharide antigens is well known.

[0080] Toxic protein antigens may be detoxified where necessary (e.g. detoxification of pertussis toxin by chemical and/or genetic means [95]).

[0081] Where a diphtheria antigen is included in the composition it is preferred also to include tetanus antigen and pertussis antigens. Similarly, where a tetanus antigen is included it is preferred also to include diphtheria and pertussis antigens. Similarly, where a pertussis antigen is included it is preferred also to include diphtheria and tetanus antigens.

[0082] Antigens may be adsorbed to an aluminium salt.

[0083] One type of preferred composition includes further antigens that affect the immunocompromised, and so the *S. aureus* polysaccharides of the invention can be combined with one or more antigens from the following *non-S. aureus* pathogens: *Streptococcus agalactiae*, *Staphylococcus epidermis*, influenza virus, *Enterococcus faecalis*, *Pseudomonas aeruginosa*, *Legionella pneumophila*, *Listeria monocytogenes*, *Neisseria meningitidis*, and parainfluenza virus.

[0084] Another type of preferred composition includes further antigens from bacteria associated with nosocomial infections, and so the *S. aureus* polysaccharides of the invention can be combined with one or more antigens from the following *non-S. aureus* pathogens: *Clostridium difficile*, *Pseudomonas aeruginosa*, *Candida albicans*, and extraintestinal pathogenic *Escherichia coli*.

[0085] Antigens in the composition will typically be present at a concentration of at least 1 µg/ml each. In general, the concentration of any given antigen will be sufficient to elicit an immune response against that antigen.

[0086] As an alternative to using proteins antigens in the composition of the invention, nucleic acid encoding the antigen may be used [e.g. refs. 120 to 128]. Protein components of the compositions of the invention may thus be replaced by nucleic acid (preferably DNA e.g. in the form of a plasmid) that encodes the protein.

[0087] In practical terms, there may be an upper limit to the number of antigens included in compositions of the invention. The number of antigens (including *S. aureus* antigens) in a composition of the invention may be less than 20, less than 19, less than 18, less than 17, less than 16, less than 15, less than 14, less than 13, less than 12, less than 11, less than 10, less than 9, less than 8, less than 7, less than 6, less than 5, less than 4, or less than 3. The number of *S. aureus* antigens in a composition of the invention may be less than 6, less than 5, or less than 4.

Pharmaceutical compositions and methods

[0088] We disclose processes for preparing pharmaceutical compositions, comprising the steps of mixing (a) a polysaccharide disclosed herein (optionally in the form of a conjugate) with (b) a pharmaceutically acceptable carrier. The processes of the invention are for making immunogenic compositions, which may comprise a pharmaceutically acceptable carrier. Typical 'pharmaceutically acceptable carriers' include any carrier that does not itself induce the production of antibodies harmful to the individual receiving the composition. Suitable carriers are typically large, slowly metabolised macromolecules such as proteins, polysaccharides, polylactic acids, polyglycolic acids, polymeric amino acids, amino acid copolymers, lactose, and lipid aggregates (such as oil droplets or liposomes). Such carriers are well known to those of ordinary skill in the art. The vaccines may also contain diluents, such as water, saline, glycerol, etc. Additionally, auxiliary substances, such as wetting or emulsifying agents, pH buffering substances, and the like, may be present. Sterile pyrogen-free, phosphate-buffered physiologic saline is a typical carrier. A thorough discussion of pharmaceutically acceptable excipients is available in reference 129.

[0089] Compositions of the invention and as disclosed herein may be in aqueous form (i.e. solutions or suspensions) or in a dried form (e.g. lyophilised). If a dried vaccine is used then it will be reconstituted into a liquid medium prior to injection. Lyophilisation of conjugate vaccines is known in the art e.g. the Menjugate™ product is presented in lyophilised form, whereas NeisVac-C™ and Meningitec™ are presented in aqueous form. To stabilise conjugates during lyophilisation, it may be typical to include a sugar alcohol (e.g. mannitol) or a disaccharide (e.g. sucrose or trehalose) e.g. at

between 1mg/ml and 30mg/ml (e.g. about 25 mg/ml) in the composition.

[0090] The pharmaceutical compositions may be packaged into vials or into syringes. The syringes may be supplied with or without needles. A syringe will include a single dose of the composition, whereas a vial may include a single dose or multiple doses.

[0091] Aqueous compositions of polysaccharides of the invention or as disclosed herein are suitable for reconstituting other vaccines from a lyophilised form. Where a composition of the invention or as disclosed herein is to be used for such extemporaneous reconstitution, a process for reconstituting such a lyophilised vaccine, comprising the step of mixing the lyophilised material with an aqueous composition of the invention is provided or disclosed. The reconstituted material can be used for injection.

S.aureus antigens

[0092] As mentioned above, one or more further *S.aureus* antigens can be included in compositions of the invention or as disclosed. The antigens may be protein or saccharide antigens. *S.aureus* protein antigens may be used as carrier proteins for conjugates of the invention or as disclosed, carrier proteins for other conjugates, or as unconjugated protein antigens. *S.aureus* saccharide antigens may be used as the saccharides for other conjugates or as unconjugated saccharide antigens.

[0093] Suitable *S.aureus* saccharide antigens include the exopolysaccharide of *S.aureus*, which is a poly-N-acetylglucosamine (PNAG). This polysaccharide is present in both *S.aureus* and *S.epidermidis* and can be isolated from either source [130,131]. For example, PNAG may be isolated from *S.aureus* strain MN8m [132]. The saccharide antigen may be a polysaccharide having the size that arises during purification of the exopolysaccharide from bacteria, or it may be an polysaccharide achieved by fragmentation of such a polysaccharide e.g. size can vary from over 400kDa to between 75 and 400kDa, or between 10 and 75kDa, or up to 30 repeat units. The saccharide antigen can have various degrees of N-acetylation and, as described in reference 133, the PNAG may be less than 40% N-acetylated (e.g. less than 35, 30, 20, 15, 10 or 5% N-acetylated; deacetylated PNAG is also known as dPNAG). Deacetylated epitopes of PNAG can elicit antibodies that are capable of mediating opsonic killing. The preparation of dPNAG is described in reference 134. The PNAG may or may not be O-succinylated e.g. it may be O-succinylated on fewer less than 25, 20, 15, 10, 5, 2, 1 or 0.1% of residues. The PNAG may be conjugated to a carrier molecule as described above or alternatively unconjugated.

[0094] Another suitable *S.aureus* saccharide antigen is the type 336 antigen, which is a β -linked hexosamine with no O-acetylation [135,136]. The type 336 antigen is cross-reactive with antibodies raised against the 336 strain (ATCC 55804). The type 336 antigen may be conjugated to a carrier molecule as described above or alternatively unconjugated.

[0095] Suitable *S.aureus* protein antigens include the following *S.aureus* antigens (or antigens comprising immunogenic fragment(s) thereof) [e.g. see references 137-144]: AhpC, AhpF, Autolysin amidase, Autolysin glucosaminidase, Collagen binding protein CAN, EbhB, GehD lipase, Heparin binding protein HBP (17kDa), Laminin receptor, MAP, MntC (also known as SitC), MRPII, Npase, ORF0594, ORF0657n, ORF0826, PBP4, RAP (RNA III activating protein), Sai-1, SasK, SBI, SdrG, SdrH, SSP-1, SSP-2 and Vitronectin-binding protein.

[0096] Further suitable *S.aureus* protein antigens include a clfA antigen; a clfB antigen; a sdrE2 antigen; a sdrC antigen; a sasF antigen, a emp antigen; a sdrD antigen; a spa antigen; a esaC antigen; a esxA antigen; a esxB antigen; a sta006 antigen; a isdC antigen; a Hla antigen; a sta011 antigen; a isdA antigen; a isdB antigen; and a sta073 antigen, as described below. One or more (i.e. 1, 2, 3, 4, 5, 6 or more) of these antigens may be present in a composition of the invention. Of these antigens, the use of one or more (i.e. 1, 2, 3, 4, 5, 6 or more) of a esxA antigen; a esxB antigen; a sta006 antigen; a Hla antigen; a sta011 antigen; and/or a sta073 antigen is specifically envisaged.

[0097] For example, a composition of the invention or as disclosed may comprise one of the following combinations of *S.aureus* protein antigens:

(1) A esxA antigen, a esxB antigen, a sta006 antigen and a Hla antigen. The esxA and esxB antigens can usefully be combined as a hybrid polypeptide, as discussed below, e.g. a EsxAB hybrid with a esxB antigen downstream of a esxA antigen. The Hla antigen may be a detoxified mutant e.g. including a H35L mutation.

(2) A esxA antigen, a esxB antigen, a sta006 antigen and a sta011 antigen. The esxA and esxB antigens may be combined as a hybrid polypeptide, as discussed below, e.g. an EsxAB hybrid.

(3) A esxA antigen, a esxB antigen and a sta011 antigen. The esxA and esxB antigens can usefully be combined as a hybrid polypeptide, as discussed below, e.g. a EsxAB hybrid.

(4) A esxA antigen, a esxB antigen, a Hla antigen, a sta006 antigen and a sta011 antigen. The esxA and esxB antigens may be combined as a hybrid polypeptide, as discussed below, e.g. an EsxAB hybrid. The Hla antigen may be a detoxified mutant e.g. including a H35L mutation.

(5) A esxA antigen, a esxB antigen and a Hla antigen. The esxA and esxB antigens can usefully be combined as a hybrid polypeptide, as discussed below, e.g. a EsxAB hybrid. The Hla antigen may be a detoxified mutant e.g. including a H35L mutation.

(6) A Hla antigen, a sta006 antigen and a sta011 antigen. The Hla antigen may be a detoxified mutant e.g. including a H35L mutation.

(7) A esxA antigen and a esxB antigen. The esxA and esxB antigens can usefully be combined as a hybrid polypeptide, as discussed below, e.g. an EsxAB hybrid.

(8) A esxA antigen, a esxB antigen and a sta006 antigen. The esxA and esxB antigens can usefully be combined as a hybrid polypeptide, as discussed below, e.g. a EsxAB hybrid.

(9) A esxA antigen, a esxB antigen, a sta011 antigen and a sta073 antigen. The esxA and esxB antigens may be combined as a hybrid polypeptide, as discussed below, e.g. an EsxAB hybrid.

(10) A sta006 antigen and a sta011 antigen.

[0098] Further *Staphylococcus aureus* antigens are disclosed in reference 145.

clfA

[0099] The 'clfA' antigen is annotated as 'clumping factor A'. In the NCTC 8325 strain clfA is SAOUHSC_00812 and has amino acid sequence SEQ ID NO: 1 (GI:88194572). In the Newman strain it is nwmn_0756 (GI:151220968).

[0100] Useful clfA antigens can elicit an antibody (e.g. when administered to a human) that recognises SEQ ID NO: 1 and/or may comprise an amino acid sequence: (a) having 50% or more identity (e.g. 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5% or more) to SEQ ID NO: 1; and/or (b) comprising a fragment of at least 'n' consecutive amino acids of SEQ ID NO: 1, wherein 'n' is 7 or more (e.g. 8, 10, 12, 14, 16, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 250 or more). These clfA proteins include variants of SEQ ID NO: 1. Preferred fragments of (b) comprise an epitope from SEQ ID NO: 1. Other preferred fragments lack one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the C-terminus and/or one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the N-terminus of SEQ ID NO: 1 while retaining at least one epitope of SEQ ID NO: 1. The final 368 C-terminal amino acids of SEQ ID NO: 1 can usefully be omitted. The first 39 N-terminal amino acids of SEQ ID NO: 1 can usefully be omitted. Other fragments omit one or more protein domains.

[0101] SEQ ID NO: 2 is a useful fragment of SEQ ID NO: 1 ('ClfA₄₀₋₅₅₉'). This fragment omits the long repetitive region towards the C-terminal of SEQ ID NO: 1.

clfB

[0102] The 'clfB' antigen is annotated as 'clumping factor B'. In the NCTC 8325 strain clfB is SAOUHSC_02963 and has amino acid sequence SEQ ID NO: 3 (GI:88196585). In the Newman strain it is nwmn_2529 (GI:151222741).

[0103] Useful clfB antigens can elicit an antibody (e.g. when administered to a human) that recognises SEQ ID NO: 3 and/or may comprise an amino acid sequence: (a) having 50% or more identity (e.g. 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5% or more) to SEQ ID NO: 3; and/or (b) comprising a fragment of at least 'n' consecutive amino acids of SEQ ID NO: 3, wherein 'n' is 7 or more (e.g. 8, 10, 12, 14, 16, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 250 or more). These clfB proteins include variants of SEQ ID NO: 3. Preferred fragments of (b) comprise an epitope from SEQ ID NO: 3. Other preferred fragments lack one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the C-terminus and/or one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the N-terminus of SEQ ID NO: 3 while retaining at least one epitope of SEQ ID NO: 3. The final 40 C-terminal amino acids of SEQ ID NO: 3 can usefully be omitted. The first 44 N-terminal amino acids of SEQ ID NO: 3 can usefully be omitted. Other fragments omit one or more protein domains. ClfB is naturally a long protein and so the use of fragments is helpful e.g. for purification, handling, fusion, expression, etc.

[0104] SEQ ID NO: 4 is a useful fragment of SEQ ID NO: 3 ('ClfB₄₅₋₅₅₂'). This fragment includes the most exposed domain of ClfB and is more easily used at an industrial scale. It also reduces the antigen's similarity with human proteins. Other useful fragments, based on a 3-domain model of ClfB, include: ClfB₄₅₋₃₆₀ (also known as CLfB-N12; SEQ ID NO: 5); ClfB₂₁₂₋₅₄₂ (also known as CLfB-N23; SEQ ID NO: 6); and ClfB₃₆₀₋₅₄₂ (also known as CLfB-N3; SEQ ID NO: 7).

sdrE2

[0105] The 'sdrE2' antigen is annotated as 'Ser-Asp rich fibrinogen/bone sialoprotein-binding protein SdrE'. In the Newman strain sdrE2 is NWMN_0525 and has amino acid sequence SEQ ID NO: 8 (GI:151220737).

[0106] Useful sdrE2 antigens can elicit an antibody (e.g. when administered to a human) that recognises SEQ ID NO: 8 and/or may comprise an amino acid sequence: (a) having 50% or more identity (e.g. 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5% or more) to SEQ ID NO: 8; and/or (b) comprising a fragment of at least 'n' consecutive amino acids of SEQ ID NO: 8, wherein 'n' is 7 or more (e.g. 8, 10, 12, 14, 16, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 250 or more). These sdrE2 proteins include variants of SEQ ID

NO: 8. Preferred fragments of (b) comprise an epitope from SEQ ID NO: 8. Other preferred fragments lack one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the C-terminus and/or one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the N-terminus of SEQ ID NO: 8 while retaining at least one epitope of SEQ ID NO: 8. The final 38 C-terminal amino acids of SEQ ID NO: 8 can usefully be omitted. The first 52 N-terminal amino acids of SEQ ID NO: 8 can usefully be omitted. Other fragments omit one or more protein domains. SdrE2 is naturally a long protein and so the use of fragments is very helpful e.g. for purification, handling, fusion, expression, etc.

[0107] SEQ ID NO: 9 is a useful fragment of SEQ ID NO: 8 ('SdrE₅₃₋₆₃₂'). This fragment includes the most exposed domain of SdrE2 and is more easily used at an industrial scale. It also reduces the antigen's similarity with human proteins.

sdrC

[0108] The 'sdrC' antigen is annotated as 'sdrC protein'. In the NCTC 8325 strain sdrC is SAOUHSC_00544 and has amino acid sequence SEQ ID NO: 10 (GI:88194324).

[0109] Useful sdrC antigens can elicit an antibody (e.g. when administered to a human) that recognises SEQ ID NO: 10 and/or may comprise an amino acid sequence: (a) having 50% or more identity (e.g. 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5% or more) to SEQ ID NO: 10; and/or (b) comprising a fragment of at least 'n' consecutive amino acids of SEQ ID NO: 10, wherein 'n' is 7 or more (e.g. 8, 10, 12, 14, 16, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 250 or more). These sdrC proteins include variants of SEQ ID NO: 10. Preferred fragments of (b) comprise an epitope from SEQ ID NO: 10. Other preferred fragments lack one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the C-terminus and/or one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the N-terminus of SEQ ID NO: 10 while retaining at least one epitope of SEQ ID NO: 10. The final 38 C-terminal amino acids of SEQ ID NO: 10 can usefully be omitted. The first 50 N-terminal amino acids of SEQ ID NO: 10 can usefully be omitted. Other fragments omit one or more protein domains. SdrC is naturally a long protein and so the use of fragments is helpful e.g. for purification, handling, fusion, expression, etc.

[0110] SEQ ID NO: 11 is a useful fragment of SEQ ID NO: 10 ('SdrC₅₁₋₅₁₈'). This fragment includes the most exposed domain of SdrC and is more easily used at an industrial scale. It also reduces the antigen's similarity with human proteins.

sasF

[0111] The 'sasF' antigen is annotated as 'sasF protein'. In the NCTC 8325 strain sasF is SAOUHSC_02982 and has amino acid sequence SEQ ID NO: 12 (GI:88196601).

[0112] Useful sasF antigens can elicit an antibody (e.g. when administered to a human) that recognises SEQ ID NO: 12 and/or may comprise an amino acid sequence: (a) having 50% or more identity (e.g. 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5% or more) to SEQ ID NO: 12; and/or (b) comprising a fragment of at least 'n' consecutive amino acids of SEQ ID NO: 12, wherein 'n' is 7 or more (e.g. 8, 10, 12, 14, 16, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 250 or more). These sasF proteins include variants of SEQ ID NO: 12. Preferred fragments of (b) comprise an epitope from SEQ ID NO: 12. Other preferred fragments lack one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the C-terminus and/or one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the N-terminus of SEQ ID NO: 12 while retaining at least one epitope of SEQ ID NO: 12. The final 39 C-terminal amino acids of SEQ ID NO: 12 can usefully be omitted. The first 37 N-terminal amino acids of SEQ ID NO: 12 can usefully be omitted. Other fragments omit one or more protein domains.

emp

[0113] The 'emp' antigen is annotated as 'extracellular matrix and plasma binding protein'. In the NCTC 8325 strain emp is SAOUHSC_00816 and has amino acid sequence SEQ ID NO: 13 (GI:88194575). In the Newman strain it is nwmn_0758 (GI:151220970).

[0114] Useful emp antigens can elicit an antibody (e.g. when administered to a human) that recognises SEQ ID NO: 13 and/or may comprise an amino acid sequence: (a) having 50% or more identity (e.g. 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5% or more) to SEQ ID NO: 13; and/or (b) comprising a fragment of at least 'n' consecutive amino acids of SEQ ID NO: 13, wherein 'n' is 7 or more (e.g. 8, 10, 12, 14, 16, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 250 or more). These emp proteins include variants of SEQ ID NO: 13. Preferred fragments of (b) comprise an epitope from SEQ ID NO: 13. Other preferred fragments lack one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the C-terminus and/or one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the N-terminus of SEQ ID NO: 13 while retaining at least one epitope of SEQ ID NO: 13. The first 26 N-terminal amino acids of SEQ ID NO: 13 can usefully be omitted. Other fragments omit one or more protein domains.

[0115] SEQ ID NOs: 14, 15, 16 and 17 are useful fragments of SEQ ID NO: 13 ('Emp₃₅₋₃₄₀', 'Emp₂₇₋₃₃₄', 'Emp₃₅₋₃₃₄' and 'Emp₂₇₋₁₄₇', respectively).

sdrD

[0116] The 'sdrD' antigen is annotated as 'sdrD protein'. In the NCTC 8325 strain sdrD is SAOUHSC_00545 and has amino acid sequence SEQ ID NO: 18 (GI:88194325).

[0117] Useful sdrD antigens can elicit an antibody (e.g. when administered to a human) that recognises SEQ ID NO: 18 and/or may comprise an amino acid sequence: (a) having 50% or more identity (e.g. 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5% or more) to SEQ ID NO: 18; and/or (b) comprising a fragment of at least 'n' consecutive amino acids of SEQ ID NO: 18, wherein 'n' is 7 or more (e.g. 8, 10, 12, 14, 16, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 250 or more). These sdrD proteins include variants of SEQ ID NO: 18. Preferred fragments of (b) comprise an epitope from SEQ ID NO: 18. Other preferred fragments lack one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the C-terminus and/or one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the N-terminus of SEQ ID NO: 18 while retaining at least one epitope of SEQ ID NO: 18. The final 38 C-terminal amino acids of SEQ ID NO: 18 can usefully be omitted. The first 52 N-terminal amino acids of SEQ ID NO: 18 can usefully be omitted. Other fragments omit one or more protein domains. SdrD is naturally a long protein and so the use of fragments is very helpful e.g. for purification, handling, fusion, expression, etc.

[0118] SEQ ID NO: 19 is a useful fragment of SEQ ID NO: 18 ('SdrD₅₃₋₅₉₂'). This fragment includes the most exposed domain of SdrD and is more easily used at an industrial scale. It also reduces the antigen's similarity with human proteins. A nother useful fragment, with the same C-terminus residue, is SdrD₃₉₄₋₅₉₂ (also known as SdrD-N3; SEQ ID NO: 20).

spa

[0119] The 'spa' antigen is annotated as 'protein A' or 'SpA'. In the NCTC 8325 strain spa is SAOUHSC_00069 and has amino acid sequence SEQ ID NO: 21 (GI:88193885). In the Newman strain it is nwmn_0055 (GI:151220267). All *S.aureus* strains express the structural gene for spa, a well characterized virulence factor whose cell wall-anchored surface protein product has five highly homologous immunoglobulin binding domains designated E, D, A, B, and C [146]. These domains display ~80% identity at the amino acid level, are 56 to 61 residues in length, and are organized as tandem repeats [147]. SpA is synthesized as a precursor protein with an N-terminal signal peptide and a C-terminal sorting signal [148,149]. Cell wall-anchored spa is displayed in great abundance on the staphylococcal surface [150,151]. Each of its immunoglobulin binding domains is composed of antiparallel α -helices that assemble into a three helix bundle and can bind the Fc domain of immunoglobulin G (IgG) [152,153], the VH3 heavy chain (Fab) of IgM (i.e. the B cell receptor) [154], the von Willebrand factor at its A1 domain [155] and/or the TNF- α receptor I (TNFRI) [156], which is displayed on surfaces of airway epithelia.

[0120] Useful spa antigens can elicit an antibody (e.g. when administered to a human) that recognises SEQ ID NO: 21 and/or may comprise an amino acid sequence: (a) having 50% or more identity (e.g. 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5% or more) to SEQ ID NO: 21; and/or (b) comprising a fragment of at least 'n' consecutive amino acids of SEQ ID NO: 21, wherein 'n' is 7 or more (e.g. 8, 10, 12, 14, 16, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 250 or more). These spa proteins include variants of SEQ ID NO: 21. Preferred fragments of (b) comprise an epitope from SEQ ID NO: 21. Other preferred fragments lack one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the C-terminus and/or one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the N-terminus of SEQ ID NO: 21 while retaining at least one epitope of SEQ ID NO: 21. The final 35 C-terminal amino acids of SEQ ID NO: 21 can usefully be omitted. The first 36 N-terminal amino acids of SEQ ID NO: 21 can usefully be omitted. Other fragments omit one or more protein domains. Reference 157 suggests that individual IgG-binding domains might be useful immunogens, alone or in combination.

[0121] SEQ ID NO: 22 is a useful fragment of SEQ ID NO: 21 ('Spa₃₇₋₃₂₅'). This fragment contains all the five SpA Ig-binding domains and includes the most exposed domain of SpA. It also reduces the antigen's similarity with human proteins. Other useful fragments may omit 1, 2, 3 or 4 of the natural A, B, C, D and/or E domains. As reported in reference 157, other useful fragments may include only 1, 2, 3 or 4 of the natural A, B, C, D and/or E domains e.g. comprise only the SpA(A) domain but not B to E, or comprise only the SpA(D) domain but not A, B, C or E, etc. Thus a spa antigen useful with the invention may include 1, 2, 3, 4 or 5 IgG-binding domains, but ideally has 4 or fewer. If an antigen includes only one type of spa domain (e.g. only the SpA(A) or SpA(D) domain), it may include more than one copy of this domain e.g. multiple SpA(D) domains in a single polypeptide chain. An individual domain within the antigen may be mutated at 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more amino acids relative to SEQ ID NO: 21 (e.g. see ref. 157, disclosing mutations at residues 3 and/or 24 of domain D, at residue 46 and/or 53 of domain A, etc.). Such mutants should not remove the antigen's ability to elicit an antibody that recognises SEQ ID NO: 21, but may remove the antigen's binding to IgG. In

certain aspects a spa antigen includes a substitution at (a) one or more amino acid substitution in an IgG Fc binding sub-domain of SpA domain A, B, C, D and/or E that disrupts or decreases binding to IgG Fc, and (b) one or more amino acid substitution in a V_H3 binding sub-domain of SpA domain A, B, C, D, and/or E that disrupts or decreases binding to V_H3. In certain embodiments, a variant SpA comprises at least or at most 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more variant SpA domain D peptides.

esaC

[0122] The 'esaC' antigen is annotated as 'esaC'. In the NCTC 8325 strain esaC is SAOUHSC_00264 and has amino acid sequence SEQ ID NO: 23 (GI:88194069).

[0123] Useful esaC antigens can elicit an antibody (e.g. when administered to a human) that recognises SEQ ID NO: 23 and/or may comprise an amino acid sequence: (a) having 50% or more identity (e.g. 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5% or more) to SEQ ID NO: 23; and/or (b) comprising a fragment of at least 'n' consecutive amino acids of SEQ ID NO: 23, wherein 'n' is 7 or more (e.g. 8, 10, 12, 14, 16, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100 or more). These esaC proteins include variants of SEQ ID NO: 23. Preferred fragments of (b) comprise an epitope from SEQ ID NO: 23. Other preferred fragments lack one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the C-terminus and/or one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the N-terminus of SEQ ID NO: 23 while retaining at least one epitope of SEQ ID NO: 23. Other fragments omit one or more protein domains.

esxA

[0124] The 'esxA' antigen is annotated as 'protein'. In the NCTC 8325 strain esxA is SAOUHSC_00257 and has amino acid sequence SEQ ID NO: 24 (GI:88194063).

[0125] Useful esxA antigens can elicit an antibody (e.g. when administered to a human) that recognises SEQ ID NO: 24 and/or may comprise an amino acid sequence: (a) having 50% or more identity (e.g. 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5% or more) to SEQ ID NO: 24; and/or (b) comprising a fragment of at least 'n' consecutive amino acids of SEQ ID NO: 24, wherein 'n' is 7 or more (e.g. 8, 10, 12, 14, 16, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90 or more). These esxA proteins include variants of SEQ ID NO: 24. Preferred fragments of (b) comprise an epitope from SEQ ID NO: 24. Other preferred fragments lack one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the C-terminus and/or one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the N-terminus of SEQ ID NO: 24 while retaining at least one epitope of SEQ ID NO: 24. Other fragments omit one or more protein domains.

esxB

[0126] The 'esxB' antigen is annotated as 'esxB'. In the NCTC 8325 strain esxB is SAOUHSC_00265 and has amino acid sequence SEQ ID NO: 25 (GI:88194070).

[0127] Useful esxB antigens can elicit an antibody (e.g. when administered to a human) that recognises SEQ ID NO: 25 and/or may comprise an amino acid sequence: (a) having 50% or more identity (e.g. 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5% or more) to SEQ ID NO: 25; and/or (b) comprising a fragment of at least 'n' consecutive amino acids of SEQ ID NO: 25, wherein 'n' is 7 or more (e.g. 8, 10, 12, 14, 16, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100 or more). These esxB proteins include variants of SEQ ID NO: 25. Preferred fragments of (b) comprise an epitope from SEQ ID NO: 25. Other preferred fragments lack one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the C-terminus and/or one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the N-terminus of SEQ ID NO: 25 while retaining at least one epitope of SEQ ID NO: 25. Other fragments omit one or more protein domains.

sta006

[0128] The 'sta006' antigen is annotated as 'ferrichrome-binding protein', and has also been referred to as 'FhuD2' in the literature [158]. In the NCTC 8325 strain sta006 is SAOUHSC_02554 and has amino acid sequence SEQ ID NO: 26 (GI:88196199). In the Newman strain it is nwmn_2185 (GI:151222397).

[0129] Useful sta006 antigens can elicit an antibody (e.g. when administered to a human) that recognises SEQ ID NO: 26 and/or may comprise an amino acid sequence: (a) having 50% or more identity (e.g. 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5% or more) to SEQ ID NO: 26; and/or (b) comprising a fragment of at least 'n' consecutive amino acids of SEQ ID NO: 26, wherein 'n' is 7 or more (e.g. 8, 10, 12, 14, 16, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 250 or more). These sta006 proteins include variants

of SEQ ID NO: 26. Preferred fragments of (b) comprise an epitope from SEQ ID NO: 26. Other preferred fragments lack one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the C-terminus and/or one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the N-terminus of SEQ ID NO: 26 while retaining at least one epitope of SEQ ID NO: 26. The first 17 N-terminal amino acids of SEQ ID NO: 26 can usefully be omitted. Other fragments omit one or more protein domains. Mutant forms of sta006 are reported in reference 159. A sta006 antigen may be lipidated e.g. with an acylated N-terminus cysteine.

isdC

[0130] The 'isdC' antigen is annotated as 'protein'. In the NCTC 8325 strain isdC is SAOUHSC_01082 and has amino acid sequence SEQ ID NO: 27 (GI:88194830).

[0131] Useful isdC antigens can elicit an antibody (e.g. when administered to a human) that recognises SEQ ID NO: 27 and/or may comprise an amino acid sequence: (a) having 50% or more identity (e.g. 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5% or more) to SEQ ID NO: 27; and/or (b) comprising a fragment of at least 'n' consecutive amino acids of SEQ ID NO: 27, wherein 'n' is 7 or more (e.g. 8, 10, 12, 14, 16, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200 or more). These isdC proteins include variants of SEQ ID NO: 27. Preferred fragments of (b) comprise an epitope from SEQ ID NO: 27. Other preferred fragments lack one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the C-terminus and/or one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the N-terminus of SEQ ID NO: 27 while retaining at least one epitope of SEQ ID NO: 27. The final 39 C-terminal amino acids of SEQ ID NO: 27 can usefully be omitted. The first 28 N-terminal amino acids of SEQ ID NO: 27 can usefully be omitted. Other fragments omit one or more protein domains. Useful fragments of IsdB are disclosed in reference 165.

[0132] Reference 160 discloses antigens which usefully include epitopes from both IsdB and IsdH.

Hla

[0133] The 'Hla' antigen is the 'alpha-hemolysin precursor' also known as 'alpha toxin' or simply 'hemolysin'. In the NCTC 8325 strain Hla is SAOUHSC_01121 and has amino acid sequence SEQ ID NO: 28 (GI:88194865). In the Newman strain it is nwmn_1073 (GI:151221285). Hla is an important virulence determinant produced by most strains of *S.aureus*, having pore-forming and haemolytic activity. Anti-Hla antibodies can neutralise the detrimental effects of the toxin in animal models, and Hla is particularly useful for protecting against pneumonia.

[0134] Useful Hla antigens can elicit an antibody (e.g. when administered to a human) that recognises SEQ ID NO: 28 and/or may comprise an amino acid sequence: (a) having 50% or more identity (e.g. 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5% or more) to SEQ ID NO: 28; and/or (b) comprising a fragment of at least 'n' consecutive amino acids of SEQ ID NO: 28, wherein 'n' is 7 or more (e.g. 8, 10, 12, 14, 16, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 250 or more). These Hla proteins include variants of SEQ ID NO: 28. Preferred fragments of (b) comprise an epitope from SEQ ID NO: 28. Other preferred fragments lack one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the C-terminus and/or one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the N-terminus of SEQ ID NO: 28 while retaining at least one epitope of SEQ ID NO: 28. The first 26 N-terminal amino acids of SEQ ID NO: 28 can usefully be omitted. Truncation at the C-terminus can also be used e.g. leaving only 50 amino acids (residues 27-76 of SEQ ID NO: 28) [161]. Other fragments omit one or more protein domains.

[0135] Hla's toxicity can be avoided in compositions of the invention by chemical inactivation (e.g. using formaldehyde, glutaraldehyde or other cross-linking reagents). Instead, however, it is preferred to use mutant forms of Hla which remove its toxic activity while retaining its immunogenicity. Such detoxified mutants are already known in the art. One useful Hla antigen has a mutation at residue 61 of SEQ ID NO: 28, which is residue 35 of the mature antigen (i.e. after omitting the first 26 N-terminal amino acids). Thus residue 61 may not be histidine, and may instead be e.g. Ile, Val or preferably Leu. A His-Arg mutation at this position can also be used. For example, SEQ ID NO: 29 is the mature mutant Hla-H35L sequence and a useful Hla antigen comprises SEQ ID NO: 29. Another useful mutation replaces a long loop with a short sequence e.g. to replace the 39mer at residues 136-174 of SEQ ID NO: 28 with a tetramer such as PSGS (SEQ ID NO: 30), as in SEQ ID NO: 31 (which also includes the H35L mutation) and SEQ ID NO: 32 (which does not include the H35L mutation).

[0136] Further useful Hla antigens are disclosed in references 162 and 163.

[0137] SEQ ID NOs: 33, 34 & 35 are three useful fragments of SEQ ID NO: 28 ('Hla₂₇₋₇₆', 'Hla₂₇₋₈₉' and 'Hla₂₇₋₇₉', respectively). SEQ ID NOs: 36, 37 and 38 are the corresponding fragments from SEQ ID NO: 29.

sta011

[0138] The 'sta011' antigen is annotated as 'lipoprotein'. In the NCTC 8325 strain sta011 is SAOUHSC_00052 and has amino acid sequence SEQ ID NO: 39 (GI:88193872).

[0139] Useful sta011 antigens can elicit an antibody (e.g. when administered to a human) that recognises SEQ ID NO: 39 and/or may comprise an amino acid sequence: (a) having 50% or more identity (e.g. 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5% or more) to SEQ ID NO: 39; and/or (b) comprising a fragment of at least 'n' consecutive amino acids of SEQ ID NO: 39, wherein 'n' is 7 or more (e.g. 8, 10, 12, 14, 16, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 250 or more). These sta011 proteins include variants of SEQ ID NO: 39. Preferred fragments of (b) comprise an epitope from SEQ ID NO: 39. Other preferred fragments lack one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the C-terminus and/or one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the N-terminus of SEQ ID NO: 39 while retaining at least one epitope of SEQ ID NO: 39. The first 23 N-terminal amino acids of SEQ ID NO: 39 can usefully be omitted. Other fragments omit one or more protein domains. A sta006 antigen may be lipidated e.g. with an acylated N-terminus cysteine.

[0140] Variant forms of SEQ ID NO: 39 which may be used for preparing sta011 antigens include, but are not limited to, SEQ ID NOs: 40, 41 and 42 with various Ile/Val/Leu substitutions.

isdA

[0141] The 'isdA' antigen is annotated as 'IsdA protein'. In the NCTC 8325 strain isdA is SAOUHSC_01081 and has amino acid sequence SEQ ID NO: 43 (GI:88194829). In the Newman strain it is nwmn_1041 (GI:151221253).

[0142] Useful isdA antigens can elicit an antibody (e.g. when administered to a human) that recognises SEQ ID NO: 43 and/or may comprise an amino acid sequence: (a) having 50% or more identity (e.g. 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5% or more) to SEQ ID NO: 43; and/or (b) comprising a fragment of at least 'n' consecutive amino acids of SEQ ID NO: 43, wherein 'n' is 7 or more (e.g. 8, 10, 12, 14, 16, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 250 or more). These isdA proteins include variants of SEQ ID NO: 43. Preferred fragments of (b) comprise an epitope from SEQ ID NO: 43. Other preferred fragments lack one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the C-terminus and/or one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the N-terminus of SEQ ID NO: 43 while retaining at least one epitope of SEQ ID NO: 43. The final 38 C-terminal amino acids of SEQ ID NO: 43 can usefully be omitted. The first 46 N-terminal amino acids of SEQ ID NO: 43 can usefully be omitted. Truncation to exclude the C-terminal 38mer of SEQ ID NO: 43 (beginning with the LPKTG motif) is also useful. Other fragments omit one or more protein domains.

[0143] SEQ ID NO: 44 is a useful fragment of SEQ ID NO: 43 (amino acids 40-184 of SEQ ID NO: 43; 'IsdA₄₀₋₁₈₄') which includes the natural protein's heme binding site and includes the antigen's most exposed domain. It also reduces the antigen's similarity with human proteins. Other useful fragments are disclosed in references 164 and 165.

[0144] IsdA does not adsorb well to aluminium hydroxide adjuvants, so IsdA present in a composition may be unadsorbed or may be adsorbed to an alternative adjuvant e.g. to an aluminium phosphate.

isdB

[0145] The 'isdB' antigen is annotated as 'neurofilament protein isdB'. In the NCTC 8325 strain isdB is SAOUHSC_01079 and has amino acid sequence SEQ ID NO: 45 (GI:88194828). IsdB has been proposed for use as a vaccine antigen on its own [166], but this may not prevent pneumonia.

[0146] Useful isdB antigens can elicit an antibody (e.g. when administered to a human) that recognises SEQ ID NO: 45 and/or may comprise an amino acid sequence: (a) having 50% or more identity (e.g. 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5% or more) to SEQ ID NO: 45; and/or (b) comprising a fragment of at least 'n' consecutive amino acids of SEQ ID NO: 45, wherein 'n' is 7 or more (e.g. 8, 10, 12, 14, 16, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 250 or more). These isdB proteins include variants of SEQ ID NO: 45. Preferred fragments of (b) comprise an epitope from SEQ ID NO: 45. Other preferred fragments lack one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the C-terminus and/or one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the N-terminus of SEQ ID NO: 45 while retaining at least one epitope of SEQ ID NO: 45. The final 36 C-terminal amino acids of SEQ ID NO: 45 can usefully be omitted. The first 40 N-terminal amino acids of SEQ ID NO: 45 can usefully be omitted. Other fragments omit one or more protein domains. Useful fragments of IsdB are disclosed in references 165 and 167 e.g. lacking 37 internal amino acids of SEQ ID NO: 45.

[0147] In some embodiments, compositions of the invention do not include an isdB antigen.

sta073

[0148] The 'sta073' antigen is annotated as 'bifunctional autolysin precursor'. In the NCTC 8325 strain sta073 is SAOUHSC_00994 and has amino acid sequence SEQ ID NO: 46 (GI:88194750). In the Newman strain it is nwmn_0922 (GI:151221134). Proteomic analysis has revealed that this protein is secreted or surface-exposed.

[0149] Useful sta073 antigens can elicit an antibody (e.g. when administered to a human) that recognises SEQ ID NO: 46 and/or may comprise an amino acid sequence: (a) having 50% or more identity (e.g. 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5% or more) to SEQ ID NO: 46; and/or (b) comprising a fragment of at least 'n' consecutive amino acids of SEQ ID NO: 46, wherein 'n' is 7 or more (e.g. 8, 10, 12, 14, 16, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 250 or more). These sta073 proteins include variants of SEQ ID NO: 46. Preferred fragments of (b) comprise an epitope from SEQ ID NO: 46. Other preferred fragments lack one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the C-terminus and/or one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the N-terminus of SEQ ID NO: 46 while retaining at least one epitope of SEQ ID NO: 46. The first 24 N-terminal amino acids of SEQ ID NO: 46 can usefully be omitted. Other fragments omit one or more protein domains.

[0150] Sta073 does not adsorb well to aluminium hydroxide adjuvants, so Sta073 present in a composition may be unadsorbed or may be adsorbed to an alternative adjuvant e.g. to an aluminium phosphate.

Hybrid polypeptides

[0151] *S.aureus* protein antigens used in the invention or as disclosed may be present in the composition as individual separate polypeptides. Where more than one antigen is used, however, they do not have to be present as separate polypeptides. Instead, at least two (e.g. 2, 3, 4, 5, or more) antigens can be expressed as a single polypeptide chain (a 'hybrid' polypeptide). Hybrid polypeptides offer two main advantages: first, a polypeptide that may be unstable or poorly expressed on its own can be assisted by adding a suitable hybrid partner that overcomes the problem; second, commercial manufacture is simplified as only one expression and purification need be employed in order to produce two polypeptides which are both antigenically useful.

[0152] The hybrid polypeptide may comprise two or more polypeptide sequences from each of the antigens listed above, or two or more variants of the same antigen in the cases in which the sequence has partial variability across strains.

[0153] Hybrids consisting of amino acid sequences from two, three, four, five, six, seven, eight, nine, or ten antigens are useful. In particular, hybrids consisting of amino acid sequences from two, three, four, or five antigens are preferred, such as two or three antigens.

[0154] Different hybrid polypeptides may be mixed together in a single formulation. Hybrids may be combined with non-hybrid antigens selected from the first, second or third antigen groups. Within such combinations, an antigen may be present in more than one hybrid polypeptide and/or as a non-hybrid polypeptide. It is preferred, however, that an antigen is present either as a hybrid or as a non-hybrid, but not as both.

[0155] Hybrid polypeptides can be represented by the formula $\text{NH}_2\text{-A}\{-\text{X-L}\}_n\text{-B-COOH}$, wherein: X is an amino acid sequence of a *S.aureus* antigen, as described above; L is an optional linker amino acid sequence; A is an optional N-terminal amino acid sequence; B is an optional C-terminal amino acid sequence; *n* is an integer of 2 or more (e.g. 2, 3, 4, 5, 6, etc.). Usually *n* is 2 or 3.

[0156] If a -X- moiety has a leader peptide sequence in its wild-type form, this may be included or omitted in the hybrid protein. In some embodiments, the leader peptides will be deleted except for that of the -X-moiety located at the N-terminus of the hybrid protein i.e. the leader peptide of X_1 will be retained, but the leader peptides of $\text{X}_2 \dots \text{X}_n$ will be omitted. This is equivalent to deleting all leader peptides and using the leader peptide of X_1 as moiety -A-.

[0157] For each *n* instances of $\{-\text{X-L}\}$, linker amino acid sequence -L- may be present or absent. For instance, when *n*=2 the hybrid may be $\text{NH}_2\text{-X}_1\text{-L}_1\text{-X}_2\text{-L}_2\text{-COOH}$, $\text{NH}_2\text{-X}_1\text{-X}_2\text{-COOH}$, $\text{NH}_2\text{-X}_1\text{-L}_1\text{-X}_2\text{-COOH}$, $\text{NH}_2\text{-X}_1\text{-X}_2\text{-L}_2\text{-COOH}$, etc. Linker amino acid sequence(s) -L- will typically be short (e.g. 20 or fewer amino acids i.e. 20, 19, 18, 17, 16, 15, 14, 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, 3, 2, 1). Examples comprise short peptide sequences which facilitate cloning, poly-glycine linkers (i.e. comprising Gly_n where *n* = 2, 3, 4, 5, 6, 7, 8, 9, 10 or more), and histidine tags (i.e. His_n where *n* = 3, 4, 5, 6, 7, 8, 9, 10 or more). Other suitable linker amino acid sequences will be apparent to those skilled in the art. A useful linker is GSGGGG (SEQ ID NO: 47) or GSGSGGGG (SEQ ID NO: 48), with the Gly-Ser dipeptide being formed from a *Bam*HI restriction site, thus aiding cloning and manipulation, and the $(\text{Gly})_4$ tetrapeptide being a typical poly-glycine linker. Other suitable linkers, particularly for use as the final L_n are ASGGGS (SEQ ID NO: 49 e.g. encoded by SEQ ID NO: 50) or a Leu-Glu dipeptide.

[0158] -A- is an optional N-terminal amino acid sequence. This will typically be short (e.g. 40 or fewer amino acids i.e. 40, 39, 38, 37, 36, 35, 34, 33, 32, 31, 30, 29, 28, 27, 26, 25, 24, 23, 22, 21, 20, 19, 18, 17, 16, 15, 14, 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, 3, 2, 1). Examples include leader sequences to direct protein trafficking, or short peptide sequences which facilitate cloning or purification (e.g. histidine tags i.e. His_n where *n* = 3, 4, 5, 6, 7, 8, 9, 10 or more). Other suitable N-

terminal amino acid sequences will be apparent to those skilled in the art. If X₁ lacks its own N-terminus methionine, -A- is preferably an oligopeptide (e.g. with 1, 2, 3, 4, 5, 6, 7 or 8 amino acids) which provides a N-terminus methionine e.g. Met-Ala-Ser, or a single Met residue.

[0159] -B- is an optional C-terminal amino acid sequence. This will typically be short (e.g. 40 or fewer amino acids *i.e.* 39, 38, 37, 36, 35, 34, 33, 32, 31, 30, 29, 28, 27, 26, 25, 24, 23, 22, 21, 20, 19, 18, 17, 16, 15, 14, 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, 3, 2, 1). Examples include sequences to direct protein trafficking, short peptide sequences which facilitate cloning or purification (e.g. comprising histidine tags *i.e.* His_n where *n* = 3, 4, 5, 6, 7, 8, 9, 10 or more, such as SEQ ID NO: 51), or sequences which enhance protein stability. Other suitable C-terminal amino acid sequences will be apparent to those skilled in the art.

[0160] One hybrid polypeptide of the invention may include both EsxA and EsxB antigens. These may be in either order, N- to C- terminus. SEQ ID NOs: 52 ('EsxAB'; encoded by SEQ ID NO: 53) and 54 ('EsxBA') are examples of such hybrids, both having hexapeptide linkers ASGGGS (SEQ ID NO: 49).

General

[0161] The practice of the present invention will employ, unless otherwise indicated, conventional methods of chemistry, biochemistry, molecular biology, immunology and pharmacology, within the skill of the art. Such techniques are explained fully in the literature. See, e.g., references 168-175, *etc.*

[0162] "GI" numbering is used above. A GI number, or "GenInfo Identifier", is a series of digits assigned consecutively to each sequence record processed by NCBI when sequences are added to its databases. The GI number bears no resemblance to the accession number of the sequence record. When a sequence is updated (e.g. for correction, or to add more annotation or information) then it receives a new GI number. Thus the sequence associated with a given GI number is never changed.

[0163] References to a percentage sequence identity between two amino acid sequences means that, when aligned, that percentage of amino acids are the same in comparing the two sequences. This alignment and the percent homology or sequence identity can be determined using software programs known in the art, for example those described in section 7.7.18 of ref. 176. A preferred alignment is determined by the Smith-Waterman homology search algorithm using an affine gap search with a gap open penalty of 12 and a gap extension penalty of 2, BLOSUM matrix of 62. The Smith-Waterman homology search algorithm is disclosed in ref. 177.

[0164] Where the invention concerns an "epitope", this epitope may be a B-cell epitope and/or a T-cell epitope. Such epitopes can be identified empirically (e.g. using PEPSCAN [178,179] or similar methods), or they can be predicted (e.g. using the Jameson-Wolf antigenic index [180], matrix-based approaches [181], MAPITOPE [182], TEPITOPE [183,184], neural networks [185], OptiMer & EpiMer [186, 187], ADEPT [188], Tsites [189], hydrophilicity [190], antigenic index [191] or the methods disclosed in references 192-196, *etc.*). Epitopes are the parts of an antigen that are recognised by and bind to the antigen binding sites of antibodies or T-cell receptors, and they may also be referred to as "antigenic determinants".

[0165] Where an antigen "domain" is omitted, this may involve omission of a signal peptide, of a cytoplasmic domain, of a transmembrane domain, of an extracellular domain, *etc.*

[0166] The term "comprising", encompasses "including" as well as "consisting" e.g. a composition "comprising" X may consist exclusively of X or may include something additional e.g. X + Y.

[0167] The term "about" in relation to a numerical value *x* means, for example, $x \pm 10\%$.

[0168] The word "substantially" does not exclude "completely" e.g. a composition which is "substantially free" from Y may be completely free from Y. Where necessary, the word "substantially" may be omitted from the definition of the invention.

[0169] Where the invention provides a process involving multiple sequential steps, the invention can also provide a process involving less than the total number of steps. The different steps can be performed at very different times by different people in different places (e.g. in different countries).

[0170] It will be appreciated that sugar rings can exist in open and closed form and that, whilst closed forms are shown in structural formulae herein, open forms are also encompassed by the invention. Similarly, it will be appreciated that sugars can exist in pyranose and furanose forms and that, whilst pyranose forms are shown in structural formulae herein, furanose forms are also encompassed. Different anomeric forms of sugars are also encompassed.

BRIEF DESCRIPTION OF DRAWINGS

[0171]

Figure 1 illustrates a process for purifying *S.aureus* type 5 and type 8 capsular polysaccharides based on the method of reference 13.

Figure 2 shows a DEAE Sepharose chromatogram of capsular polysaccharide and a ^1H NMR spectrum of capsular polysaccharide-containing fractions (fractions 68-80) prepared according to the method of Figure 1.

Figure 3 shows a S300 Sephacryl chromatogram of capsular polysaccharide and a ^1H NMR spectrum of capsular polysaccharide-containing fractions (fractions 22-44) prepared according to the method of Figure 1.

Figure 4 illustrates an exemplary process of the invention for purifying *S.aureus* type 5 and type 8 capsular polysaccharides.

Figure 5 shows a DEAE Sepharose chromatogram of capsular polysaccharide prepared according to a method of the invention.

Figure 6 shows a ^1H NMR spectrum for purified *S.aureus* type 5 capsular polysaccharide.

Figure 7 shows the chemical structure of the peptidoglycan of *S.aureus* based on references 197, 198, 199 and 200. The repeat unit is highlighted.

MODES FOR CARRYING OUT THE INVENTION

A. Purification of *S.aureus* type 5 capsular polysaccharide (comparative example)

[0172] *S.aureus* type 5 capsular polysaccharide was purified according to the scheme illustrated in Figure 1, based on the method of reference 13. The conditions and rationale for the various steps of this method are described in Table 1:

Table 1

Step	Conditions	Rationale
Bacterial growth on plates		
Bacterial pellet centrifugation		Harvest of cells
Reaction with Lysostaphin	100 $\mu\text{g/ml}$ of Lysostaphin overnight at 37°C	Cell wall lysis and release of capsular polysaccharide
Reaction with DNase/RNase	50 $\mu\text{g/ml}$ of DNase and RNase at 37°C for 6-8hrs	Nucleic acid hydrolysis
Reaction with NaIO_4	0.05M NaIO_4 for 5hrs at RT in the dark	Teichoic acid hydrolysis
Diafiltration 30kDa	Washing with NaCl 1M and H_2O	Low molecular weight species removal
Anion exchange chromatography (DEAE SepharoseFF resin)	NaCl 1M gradient	Separation according to charge (protein removal)
Gel filtration (Sephacryl S300)	NaPi 10 mM pH 7.2 and NaCl 10 mM	Separation according to molecular weight

Bacterial pellet centrifugation and enzymatic reactions (Lysostaphin and RNase/DNase)

[0173] *S.aureus* was grown in solid medium to provide a bacterial suspension of 600-800 ml. The wet cell pellet, harvested by centrifugation at 8000 rpm, had a mass of around 30-50 g. The harvested pellet was washed three times with 50mM Tris-2mM MgSO_4 pH7.5 and then suspended at 0.25-0.5g per ml in 50mM Tris-2mM MgSO_4 pH7.5 and treated with 0.1-0.13mg/ml of lysostaphin (Sigma-Aldrich). The reaction mixture was incubated at 37°C for 16hrs (ON) with mild stirring. 0.05 mg/ml of DNase/RNase (Sigma-Aldrich) was added to the suspension and incubated for 5-7hrs at 37°C . The suspension was then clarified by centrifugation.

Reaction with NaIO_4

[0174] The material was incubated with 50mM NaIO_4 (Sigma-Aldrich) in the dark for 5-7hrs. NaIO_4 was then removed

EP 2 493 498 B1

by the addition of excess glycerol for 30 minutes with stirring in the light.

30 kDa tangential flow filtration

[0175] Tangential flow filtration was carried out as indicated in Table 2:

Table 2

Membrane type	Sartorius Hydrosart™ 30kDa
Surface area	0.1m ²
P _{in} /P _{out}	0.4/0.0 bar
Permeate flow rate	80 ml/min
Diafiltration volumes	10 volumes of NaCl 1M followed by 10 volumes of distilled water
Product recovery	Retentate volume + two washings with distilled water equal to the dead volume of the system (with completely open retentate and closed permeate)

[0176] The tangential flow filtration was performed in a Sartorius™ holder for 0.1m² cassettes using a WatsonMarlon™ peristaltic pump. Afterwards, the membrane was washed with NaOH 1M and stored in NaOH 0.1M at +2-8°C.

DEAE Sepharose Fast Flow chromatography

[0177] Residual protein, nucleic acid and other impurities were removed by anion exchange chromatography carried out in accordance with Table 3:

Table 3

Resin	DEAE Sepharose™ Fast Flow resin (G&E Healthcare)
Column dimension	Ø = 5cm; h = 7.5cm; V = 150ml
Equilibration	10mM NaPi buffer pH7.2 q.b. to reach 1.8-2.0mS/cm eluate conductivity
Load	Retentate from 30K UF buffered to 10mM NaPi buffer pH7.2
Elution	20 column volumes of 10mM NaPi buffer pH7.2
Stripping	20 column volumes of NaCl 1M

[0178] The chromatography was performed using an Akta™ system (G&E Healthcare) and the capsular polysaccharide was detected by measuring UV absorption at 215nm. The capsular polysaccharide solution was first added to 100mM NaPi buffer pH7.2 to obtain a final buffer concentration of 10mM NaPi pH7.2. The DEAE resin was pre-equilibrated with 100mM NaPi buffer pH7.2 to pH7.2 and then equilibrated with 10m NaPi buffer pH7.2 to achieve the indicated conductivity (10mM NaPi buffer pH7.2 conductivity). The resultant fractions were analyzed by NMR and those containing capsular polysaccharide pooled together (Figure 2).

S300 Sephacryl chromatography

[0179] The polysaccharide was further purified by gel-filtration chromatography carried out in accordance with Table 4:

Table 4

Resin	S300 Sephacryl™ resin (G&E Healthcare)
Column dimension	Ø = 2.6cm; h = 95cm; V = 500ml
Equilibration	50mMNaCl buffer q.b. to reach 6.3-6.5mS/cm eluate conductivity
Load	12-14ml
Elution	50mMNaCl buffer

[0180] The chromatography was performed on an Akta™ system (G&E Healthcare) and the capsular polysaccharide was detected by measuring UV absorption at 215nm. The resultant fractions were analyzed by NMR and those containing capsular polysaccharide pooled together (Figure 3).

B. Purification of *S.aureus* type 5 and type 8 capsular polysaccharides (example)

[0181] *S.aureus* type 5 and type 8 capsular polysaccharides were purified according to the scheme illustrated in Figure 4. The conditions and rationale for the various steps of this method are described in Table 5:

Table 5

Step	Conditions	Rationale
Bacterial growth on plates		
Bacterial pellet centrifugation		Harvest of cells
Reaction with AcOH 1%	2hrs at 100°C	Cell wall lysis and release of capsular polysaccharide
Reaction with mutanolysin	180U/ml of mutanolysin at 37°C over-night	Further removal of peptidoglycan
Reaction with DNase/RNase	50 µg/ml of DNase and RNase at 37°C for 6-8hrs	Nucleic acid hydrolysis
Diafiltration 30kDa	Washing with NaCl 1M and H ₂ O	Low molecular weight species removal
Anion exchange chromatography (DEAE SepharoseFF resin)	NaCl 1M gradient	Separation according to charge (protein removal)

Bacterial pellet centrifugation and acid and enzymatic reactions (acetic acid, RNase/DNase and mutanolysin)

[0182] *S.aureus* was grown in solid medium to provide a bacterial suspension of 600-800 ml. The wet cell pellet, harvested by centrifugation at 8000 rpm, had a mass of around 30-50 g. The harvested pellet was washed three times with 50mM Tris-2mM MgSO₄ pH 7.5 and then suspended at 0.5-0.6g per ml in distilled water and stirred vigorously while the temperature was raised to 100°C. Acetic acid was then added to a final concentration of 1% and the mixture kept at 100°C for 2hrs. The mixture was neutralised with NaOH 1M and centrifuged at 8000 rpm.

[0183] The supernatant was decanted from the pellet and combined with 0.05 mg/ml of DNase/RNase (Sigma-Aldrich). The mixture was then incubated for 5-7hrs at 37°C and afterwards clarified by centrifugation.

[0184] 180U/ml of mutanolysin (Sigma-Aldrich) was then added to the suspension and the mixture incubated over-night (for 16hrs) at 37°C with mild stirring. The suspension was then clarified again by centrifugation

30 kDa tangential flow filtration

[0185] Tangential flow filtration was carried out as indicated in Table 6:

Table 6

Membrane type	Sartorius Hydrosart™ 30kDa
Surface area	0.2m ²
P _{in} /P _{out}	0.7/0.0 bar
Permeate flow rate	11 ml/min
Diafiltration volumes	10 volumes of NaCl 1M followed by 10 volumes of NaPi 10mM pH 7.2 buffer
Product recovery	Retentate volume + two washings with distilled water equal to the dead volume of the system (with completely open retentate and closed permeate)

[0186] The tangential flow filtration was performed in a Sartorius™ holder for 0.2m² cassettes using a WatsonMarlon™

peristaltic pump. Afterwards, the membrane was washed with NaOH 1M and stored in NaOH 0.1 M at +2-8°C.

DEAE Sepharose Fast Flow chromatography

[0187] Residual protein, nucleic acid and other impurities were removed by anion exchange chromatography carried out in accordance with Table 7:

Table 7

Resin	DEAE Sepharose™ Fast Flow resin (G&E Healthcare)
Column dimension	Ø = 5cm; h = 7.5cm; V = 150ml
Equilibration	10mM NaPi buffer pH7.2 q.b. to reach 1.8-2.0mS/cm eluate conductivity
Load	Retentate from 30K UF
Elution	20 column volumes of 10mM NaPi buffer pH7.2
<u>Stripping</u>	20 column volumes of NaCl 1M

[0188] The chromatography was performed using an Akta™ system (G&E Healthcare) and the capsular polysaccharide was detected by measuring UV absorption at 215nm. The capsular polysaccharide solution was first added to 100mM NaPi buffer pH7.2 to obtain a final buffer concentration of 10mM NaPi pH7.2. The DEAE resin was pre-equilibrated with 100mM NaPi buffer pH7.2 to pH7.2 and then equilibrated with 10mM NaPi buffer pH7.2 to achieve the indicated conductivity (10mM NaPi buffer pH7.2 conductivity). The resultant fractions were analyzed by NMR and those containing capsular polysaccharide pooled together (Figure 5).

30 kDa tangential flow filtration

[0189] Tangential flow filtration was carried out to remove NaCl left over from the anion exchange chromatography and to concentrate the purified polysaccharides. The filtration was carried out as indicated in Table 8:

Table 8

Membrane type	Sartorius Hydrosart™ 30kDa
Surface area	0.2m ²
P _{in} /P _{out}	0.7/0.0 bar
Permeate flow rate	11 ml/min
Diafiltration volumes	10 volumes of distilled water
Product recovery	Retentate volume + two washings with distilled water equal to the dead volume of the system (with completely open retentate and closed permeate)

[0190] The tangential flow filtration was performed in a Sartorius™ holder for 0.2m² cassettes using a WatsonMarlon™ peristaltic pump. Afterwards, the membrane was washed with NaOH 1M and stored in NaOH 0.1M at +2-8°C. The purified polysaccharide was analysed by NMR (e.g. Figure 6 for the type 5 capsular polysaccharide).

C. Determination of peptidoglycan contamination in purified polysaccharide

[0191] The peptidoglycan (Figure 7) content of purified type 5 polysaccharide obtained according to the methods in sections A and B above was determined by amino acid analysis using HPAEC-PAD according to the Dionex AAA-Direct™ system (AminoPac™ PA10 AAA-Direct™, Dionex) in accordance with the manufacturer's instructions. Briefly, 20µL of 100µM norleucine was added to 200µL of polysaccharide at 250µg/mL in water in a 400°C treated glass tube and dried using a Speedvac system. The norleucine serves as an internal standard. Samples were hydrolyzed *in vacuo* using the vapor of boiling hydrochloric acid/phenol in order to yield free amino acids from residual protein and peptidoglycan contamination. Separation of free amino acids was performed on an AminoPac™ PA10 column (2x250mm) equipped with an AminoPac™ PA10 guard column (2x50mm) using a gradient condition for amino acids and carbohydrates according to the manufacturer's recommendations. These gradient conditions are summarized in Table 9:

Table 9

Time (min)	%E1	%E2	%E3	Curve	Comments
Initiation	84	16	0		Autosampler fills the sample loop
0.0	84	16	0		Valve from Load to Inject
2.0	84	16	0		Begin hydroxide gradient
12.1	68	32	0	8	
16.0	68	32	0		Begin acetate gradient
24.0	36	24	40	8	
40.0	36	24	40		
40.1	20	80	0	5	Column wash with hydroxide
42.1	20	80	0		
42.2	84	16	0	5	Equilibrate to starting conditions
65.0	84	16	0		
<i>Eluent E1: Deionized Water; Eluent E2: 0.250 M Sodium Hydroxide; Eluent E3: 1.0 M Sodium Acetate and Flow=0.25mL/min</i>					

[0192] Detection was performed using a AAA-Direct waveform potential (Table 10).

Table 10

Time (sec)	Potential (V) vs. Ag/AgCl	Potential (V) vs. pH	Integration
0.000	-0.20	+0.13	
0.040	-0.20	+0.13	
0.050	0.00	+0.33	
0.210	0.00	+0.33	Begin
0.220	+0.22	+0.55	
0.460	+0.22	+0.55	
0.470	0.00	+0.33	
0.560	0.00	+0.33	End
0.570	-0.20	-1.67	
0.580	-0.20	-1.67	
0.590	+0.60	+0.93	
0.600	-0.20	+0.13	

[0193] The quantification was performed using a non-hydrolyzed 17 amino acid standard solution (Fluka P/N 09428) in the range 2.5-50 μ M. Standard samples were analyzed with and without norleucine, at the same sample concentration. The ratio of the norleucine peak area in the sample divided by the average norleucine peak area in the standards was used as a correction factor for possible amino acid loss in the hydrolysis step. A BSA sample was used as control sample.

Peptidoglycan content estimation

[0194] Peptidoglycan content was estimated using two different methods. The first method (method 1) was based on the method used in reference 17, which involves a summation of the lysine, alanine, glycine and glutamate content. In the second method (method 2), a conversion factor is calculated for each amino acid according to the following formula:

$$\frac{(\text{molecular mass of amino acid}) \times (\text{number of residues in the peptidoglycan structure})}{(\text{molecular mass of the repeating unit of peptidoglycan})}$$

[0195] The molecular mass of the repeating unit of peptidoglycan is 1233.27 Da (Figure 7). The peptidoglycan content was then calculated as the average peptidoglycan concentration obtained by calculating the ratio of the amino acid concentration and the conversion factor.

[0196] The peptidoglycan content of the purified type 5 capsular polysaccharide after anionic exchange chromatography is given in Table 11:

Table 11

Measurement method	Details of calculation	% Peptidoglycan	
		Measurement 1	Measurement 2
1	Calculated according to reference 17 as sum of Lys-Ala-Gly-Glx concentration	2.04	0.74
1	Calculated according to reference 17 as sum of all amino acids detectable except for Lys-Ala-Gly-Glx	0.48	0.85
2	Calculated using Ala and Gly concentration divided by PG conversion factor (Ala=0.2167, Gly=0.3043)	0.88	0.81

[0197] The method of the invention provides a very low content of peptidoglycan in the purified polysaccharide.

D. Conjugation and immunogenicity of purified polysaccharides

[0198] Purified type 5 polysaccharides obtained from the methods in sections A and B above were conjugated to CRM197 according to the method of reference 29. Total saccharide in the conjugate was determined by HPAEC-PAD analysis and protein content by MicroBCA assay (Table 12).

TABLE 12

Purification method	Lot	Protein (μg/ml)	Saccharide (μg/ml)	Saccharide/protein (w/w)
A	1	51.52	1.72	0.03
A	2	161.80	17.10	0.11
A	3	34.42	4.22	0.12
B	4	444.0	139.0	0.31
B	5	40.56	12.70	0.31

[0199] The conjugates prepared using polysaccharides purified by the method of the invention (lots 4 and 5) had higher polysaccharide:protein ratios.

[0200] The immunogenicity of lot 5 was tested in a mouse lethal model of *S.aureus* infection. Briefly, CD1 mice were immunised by intraperitoneal injection with a 2 μg dose of antigen in an injection volume of 200 μl. Immunisations were carried out in groups of twelve mice according to the following scheme, prior to challenge by intraperitoneal injection of a bacterial suspension of 5x10⁸ CFU type 5 *S.aureus*. Cultures of *S.aureus* were centrifuged, washed twice and diluted in PBS before challenge. Further dilutions were needed for the desired inoculum, which was experimentally verified by agar plating and colony formation. Animals were monitored for 14 days and lethal disease recorded.

Group 1 - PBS plus alum

Group 2 - Type 5 capsular polysaccharide-CRM conjugate (Lot 5) plus alum

EP 2 493 498 B1

Group 4 - Type 5 capsular polysaccharide-CRM conjugate (Lot 5) plus EsxAB, Sta006 and Sta011 proteins and alum

Group 5 - Type 5 capsular polysaccharide-CRM conjugate (Lot 5) plus HlaH35L, Sta006 and Sta011 proteins and alum

[0201] Survival data is presented in Table 13:

TABLE 13

Group	Time (days)													
	1	2	3	4	5	6	7	8	9	10	11	12	13	14
1	100	25	17	17	17	17	17	17	17	17	8	0	0	0
2	100	50	50	50	50	50	50	50	50	42	42	42	42	42
4	100	67	67	67	67	67	67	67	67	67	67	67	67	67
5	100	100	100	100	100	100	83	83	75	75	75	75	75	75

[0202] The conjugates prepared using polysaccharides purified by the method of the invention gave a high level of survival. Survival was enhanced by addition of *S. aureus* protein antigens.

REFERENCES

[0203]

- [1] Fattom et al. (1990) Infect Immun. 58(7):2367-74.
- [2] Fattom et al. (1992) Infect Immun. 60(2):584-9.
- [3] Fattom et al. (1993) Infect Immun. 61(3):1023-32.
- [4] Fattom et al. (1996) Infect Immun. 64(5):1659-65.
- [5] Welch et al. (1996) J Am Soc Nephrol. 7(2):247-53.
- [6] Fattom et al. (1998) Infect Immun. 66(10):4588-92.
- [7] Fattom et al. (1993) Vaccine 17(2):126-33.
- [8] Fattom et al. (2002) N Engl J Med. 346(7):491-6.
- [9] Robbins et al. (2005) Ann N Y Acad Sci. 754:68-82.
- [10] Gilbert et al. (1994) J. Microb. Meth. 20:39-46.
- [11] Gilbert et al. (1994) Vaccine. 12(4):369-74.
- [12] Tollersrud et al. (2001) Vaccine. 19(28-29):3896-903.
- [13] Lee et al. (1993) Infect Immun 61:1853-8.
- [14] WO2004/080490.
- [15] WO2006/032475.
- [16] WO2006/032500.
- [17] WO2006/065553.
- [18] WO2006/114500.
- [19] Moreau et al. (1990) Carbohydrate Res. 339(5):285-91
- [20] Fournier et al. (1984) Infect. Immun. 45(1):87-93.
- [21] Jones (2005) Carbohydrate Res. 340(6):1097-106.
- [22] Lemercinier and Jones (1996) Carbohydrate Res. 296:83-96.
- [23] Jones and Lemercinier (2002) J Pharm Biomed Anal. 30(4):1233-47.
- [24] WO05/033148
- [25] WO 00/56357
- [26] Hestrin (1949) J. Biol. Chem. 180:249-261.
- [27] Konadu et al. (1994) Infect. Immun. 62:5048-5054.
- [28] www.polymer.de
- [29] US patent application 61/247,518, 'CONJUGATION OF STAPHYLOCOCCUS AUREUS TYPE 5 AND TYPE 8 CAPSULAR POLYSACCHARIDES' (NOVARTIS AG). Assignee reference no. 53594-US-PSP and PCT application no. PCT/IB2010/002565 (NOVARTIS AG).
- [30] WO2007/113222

- [31] US patent 6,045,805
- [32] US patents 6,027,733 & 6,274,144.
- [33] www.polymer.de
- [34] Wessels et al. (1989) *Infect Immun* 57:1089-94.
- 5 [35] Ramsay et al. (2001) *Lancet* 357(9251):195-196.
- [36] Lindberg (1999) *Vaccine* 17 Suppl 2:S28-36.
- [37] Buttery & Moxon (2000) *J R Coll Physicians Lond* 34:163-68.
- [38] Ahmad & Chapnick (1999) *Infect Dis Clin North Am* 13:113-33, vii.
- [39] Goldblatt (1998) *J. Med. Microbiol.* 47:563-7.
- 10 [40] European patent 0477508.
- [41] US patent 5,306,492.
- [42] WO98/42721.
- [43] Dick et al. in *Conjugate Vaccines* (eds. Cruse et al.) Karger, Basel, 1989, 10:48-114.
- [44] Hermanson *Bioconjugate Techniques*, Academic Press, San Diego (1996) ISBN: 0123423368.
- 15 [45] Reynaud-Rondier et al. (1991) *FEMS Microbiology Immunology* 76:193-200.
- [46] WO03/061558.
- [47] Research Disclosure, 453077 (Jan 2002)
- [48] Herbelin et al. (1997) *J Dairy Sci.* 80(9):2025-34.
- [49] EP-A-0372501.
- 20 [50] EP-A-0378881.
- [51] EP-A-0427347.
- [52] WO93/17712.
- [53] WO94/03208.
- [54] WO98/58668.
- 25 [55] EP-A-0471177.
- [56] WO91/01146
- [57] Falugi et al. (2001) *Eur J Immunol* 31:3816-3824.
- [58] Baraldo et al. (2004) *Infect Immun* 72(8):4884-7.
- [59] EP-A-0594610.
- 30 [60] Ruan et al. (1990) *J Immunol* 145:3379-3384.
- [61] WO00/56360.
- [62] WO02/091998.
- [63] Kuo et al. (1995) *Infect Immun* 63:2706-13.
- [64] Michon et al. (1998) *Vaccine*. 16:1732-41.
- 35 [65] WO01/72337
- [66] WO00/61761.
- [67] WO2004/041157.
- [68] WO02/34771.
- [69] WO99/42130.
- 40 [70] WO2004/011027.
- [71] WO96/40242.
- [72] Lei et al. (2000) *Dev Biol (Basel)* 103:259-264.
- [73] WO00/38711; US patent 6,146,902.
- [74] WO99/24578.
- 45 [75] WO99/36544.
- [76] WO99/57280.
- [77] WO00/22430.
- [78] Tettelin et al. (2000) *Science* 287:1809-1815.
- [79] WO96/29412.
- 50 [80] Pizza et al. (2000) *Science* 287:1816-1820.
- [81] WO01/52885.
- [82] Bjune et al. (1991) *Lancet* 338(8775):1093-1096.
- [83] Fukasawa et al. (1999) *Vaccine* 17:2951-2958.
- [84] Rosenqvist et al. (1998) *Dev. Biol. Stand* 92:323-333..
- 55 [85] Costantino et al. (1992) *Vaccine* 10:691-698.
- [86] WO03/007985.
- [87] Watson (2000) *Pediatr Infect Dis J* 19:331-332.
- [88] Rubin (2000) *Pediatr Clin North Am* 47:269-285, v.

- [89] Jedrzejewski (2001) *Microbiol Mol Biol Rev* 65:187-207.
- [90] Bell (2000) *Pediatr Infect Dis J* 19:1187-1188.
- [91] Iwarson (1995) *APMIS* 103:321-326.
- [92] Gerlich et al. (1990) *Vaccine* 8 Suppl:S63-68 & 79-80.
- 5 [93] Hsu et al. (1999) *Clin Liver Dis* 3:901-915.
- [94] Gustafsson et al. (1996) *N. Engl. J. Med.* 334:349-355.
- [95] Rappuoli et al. (1991) *TIBTECH* 9:232-238.
- [96] Vaccines (2004) eds. Plotkin & Orenstein. ISBN 0-7216-9688-0.
- [97] WO02/02606.
- 10 [98] Kalman et al. (1999) *Nature Genetics* 21:385-389.
- [99] Read et al. (2000) *Nucleic Acids Res* 28:1397-406.
- [100] Shirai et al. (2000) *J. Infect. Dis.* 181(Suppl 3):S524-S527.
- [101] WO99/27105.
- [102] WO00/27994.
- 15 [103] WO00/37494.
- [104] WO99/28475.
- [105] Ross et al. (2001) *Vaccine* 19:4135-4142.
- [106] Sutter et al. (2000) *Pediatr Clin North Am* 47:287-308.
- [107] Zimmerman & Spann (1999) *Am Fam Physician* 59:113-118, 125-126.
- 20 [108] Dreesen (1997) *Vaccine* 15 Suppl:S2-6.
- [109] *MMWR Morb Mortal Wkly Rep* 1998 Jan 16;47(1):12, 19.
- [110] McMichael (2000) *Vaccine* 19 Suppl 1:S101-107.
- [111] WO02/34771.
- [112] Dale (1999) *Infect Dis Clin North Am* 13:227-43, viii.
- 25 [113] Ferretti et al. (2001) *PNAS USA* 98: 4658-4663.
- [114] WO03/093306.
- [115] WO2004/018646.
- [116] WO2004/041157.
- [117] Ichiman and Yoshida (1981) *J. Appl. Bacteriol.* 51:229.
- 30 [118] US4197290
- [119] Ichiman et al. (1991) *J. Appl. Bacteriol.* 71:176.
- [120] Robinson & Torres (1997) *Seminars in Immunology* 9:271-283.
- [121] Donnelly et al. (1997) *Annu Rev Immunol* 15:617-648.
- [122] Scott-Taylor & Dalgleish (2000) *Expert Opin Investig Drugs* 9:471-480.
- 35 [123] Apostolopoulos & Plebanski (2000) *Curr Opin Mol Ther* 2:441-447.
- [124] Ilan (1999) *Curr Opin Mol Ther* 1:116-120.
- [125] Dubensky et al. (2000) *Mol Med* 6:723-732.
- [126] Robinson & Pertmer (2000) *Adv Virus Res* 55:1-74.
- [127] Donnelly et al. (2000) *Am J Respir Crit Care Med* 162(4 Pt 2):S190-193.
- 40 [128] Davis (1999) *Mt. Sinai J. Med* 66:84-90.
- [129] Gennaro (2000) Remington: The Science and Practice of Pharmacy. 20th edition, ISBN: 0683306472.
- [130] Joyce et al. (2003) *Carbohydrate Research* 338 :903.
- [131] Maira-Litran et al. (2002) *Infect. Immun.* 70:4433.
- [132] WO2004/043407.
- 45 [133] WO2007/113224.
- [134] WO2004/043405
- [135] WO98/10788.
- [136] WO2007/053176.
- [137] WO2007/113222.
- 50 [138] WO2005/009379.
- [139] WO2009/029132.
- [140] WO2008/079315.
- [141] WO2005/086663.
- [142] WO2005/115113.
- 55 [143] WO2006/033918.
- [144] WO2006/078680.
- [145] Kuroda et al. (2001) *Lancet* 357(9264):1225-1240; see also pages 1218-1219.
- [146] Sjodahl (1977) *J. Biochem.* 73:343-351.

- [147] Uhlen et al. (1984) *J. Biol. Chem.* 259:1695-1702 & 13628 (Corr.).
- [148] Schneewind et al. (1992) *Cell* 70:267-281.
- [149] DeDent et al. (2008) *EMBO J.* 27:2656-2668.
- [150] Sjoquist et al. (1972) *Eur. J. Biochem.* 30:190-194.
- 5 [151] DeDent et al. (2007) *J. Bacteriol.* 189:4473-4484.
- [152] Deisenhofer et al., (1978) *Hoppe-Seyh Zeitsch. Physiol. Chem.* 359:975-985.
- [153] Deisenhofer (1981) *Biochemistry* 20:2361-2370.
- [154] Graille et al. (2000) *Proc. Nat. Acad. Sci. USA* 97:5399-5404.
- [155] O'Seaghdha et al. (2006) *FEBS J.* 273:4831-41.
- 10 [156] Gomez et al. (2006) *J. Biol. Chem.* 281:20190-20196.
- [157] WO2007/071692.
- [158] Sebulsky & Heinrichs (2001) *J Bacteriol* 183:4994-5000.
- [159] Sebulsky et al. (2003) *J Biol Chem* 278:49890-900.
- [160] WO2005/009378.
- 15 [161] Rable & Wardenburg (2009) *Infect Immun* 77:2712-8.
- [162] WO2007/145689.
- [163] WO2009/029831.
- [164] WO2005/079315.
- [165] WO2008/152447.
- 20 [166] Kuklin et al. (2006) *Infect Immun.* 74(4):2215-23.
- [167] WO2005/009379.
- [168] Gennaro (2000) *Remington: The Science and Practice of Pharmacy*. 20th edition, ISBN: 0683306472.
- [169] *Methods In Enzymology* (S. Colowick and N. Kaplan, eds., Academic Press, Inc.)
- [170] *Handbook of Experimental Immunology*, Vols. I-IV (D.M. Weir and C.C. Blackwell, eds, 1986, Blackwell Scientific Publications)
- 25 [171] Sambrook et al. (2001) *Molecular Cloning: A Laboratory Manual*, 3rd edition (Cold Spring Harbor Laboratory Press).
- [172] *Handbook of Surface and Colloidal Chemistry* (Birdi, K.S. ed., CRC Press, 1997)
- [173] Ausubel et al. (eds) (2002) *Short protocols in molecular biology*, 5th edition (Current Protocols).
- 30 [174] *Molecular Biology Techniques: An Intensive Laboratory Course*, (Ream et al., eds., 1998, Academic Press)
- [175] *PCR (Introduction to Biotechniques Series)*, 2nd ed. (Newton & Graham eds., 1997, Springer Verlag)
- [176] *Current Protocols in Molecular Biology* (F.M. Ausubel et al., eds., 1987) Supplement 30
- [177] Smith & Waterman (1981) *Adv. Appl. Math.* 2: 482-489.
- [178] Geysen et al. (1984) *PNAS USA* 81:3998-4002.
- 35 [179] Carter (1994) *Methods Mol Biol* 36:207-23.
- [180] Jameson, BA et al. 1988, *CABIOS* 4(1):181-186.
- [181] Raddrizzani & Hammer (2000) *Brief Bioinform* 1(2):179-89.
- [182] Bublil et al. (2007) *Proteins* 68(1):294-304.
- [183] De Lalla et al. (1999) *J. Immunol.* 163:1725-29.
- 40 [184] Kwok et al. (2001) *Trends Immunol* 22:583-88.
- [185] Brusica et al. (1998) *Bioinformatics* 14(2):121-30
- [186] Meister et al. (1995) *Vaccine* 13(6):581-91.
- [187] Roberts et al. (1996) *AIDS Res Hum Retroviruses* 12(7):593-610.
- [188] Maksyutov & Zagrebelnaya (1993) *ComputAppl Biosci* 9(3):291-7.
- 45 [189] Feller & de la Cruz (1991) *Nature* 349(6311):720-1.
- [190] Hopp (1993) *Peptide Research* 6:183-190.
- [191] Welling et al. (1985) *FEBS Lett.* 188:215-218.
- [192] Davenport et al. (1995) *Immunogenetics* 42:392-297.
- [193] Tsurui & Takahashi (2007) *J Pharmacol Sci.* 105(4):299-316.
- 50 [194] Tong et al. (2007) *Brief Bioinform.* 8(2):96-108.
- [195] Schirle et al. (2001) *J Immunol Methods.* 257(1-2):1-16.
- [196] Chen et al. (2007) *Amino Acids* 33(3):423-8.
- [197] Kim et al. (2008) *Biochemistry* 47(12):3822-3831.
- [198] Patti et al. (2008) *Biochemistry* 47(32):8378-8385.
- 55 [199] Kim and Schaefer (2008) *Biochemistry* 47(38):10155-10161.
- [200] Biswas (2006) PhD Thesis: Characterization of *Staphylococcus aureus* peptidoglycan hydrolases and isolation of defined peptidoglycan structures der Eberhard Karls Universität Tübingen

SEQUENCE LISTING

[0204]

5 <110> NOVARTIS AG

 <120> PURIFICATION OF STAPHYLOCOCCUS AUREUS TYPE 5 AND TYPE 8 CAPSULAR SACCHARIDES

 <130> P055840WO

10 <140> PCT/IB2010/
 <141> 2010-11-01

 <150> US-61/256,905

15 <151> 2009-10-30

 <160> 54

 <170> SeqWin2010, version 1.0

20 <210> 1
 <211> 927
 <212> PRT
 <213> Staphylococcus aureus

25 <400> 1

30

35

40

45

50

55

EP 2 493 498 B1

	Met	Asn	Met	Lys	Lys	Lys	Glu	Lys	His	Ala	Ile	Arg	Lys	Lys	Ser	Ile	
	1				5					10					15		
5	Gly	Val	Ala	Ser	Val	Leu	Val	Gly	Thr	Leu	Ile	Gly	Phe	Gly	Leu	Leu	
				20				25					30				
	Ser	Ser	Lys	Glu	Ala	Asp	Ala	Ser	Glu	Asn	Ser	Val	Thr	Gln	Ser	Asp	
			35					40					45				
10	Ser	Ala	Ser	Asn	Glu	Ser	Lys	Ser	Asn	Asp	Ser	Ser	Ser	Val	Ser	Ala	
		50					55					60					
	Ala	Pro	Lys	Thr	Asp	Asp	Thr	Asn	Val	Ser	Asp	Thr	Lys	Thr	Ser	Ser	
	65					70					75					80	
15	Asn	Thr	Asn	Asn	Gly	Glu	Thr	Ser	Val	Ala	Gln	Asn	Pro	Ala	Gln	Gln	
					85					90					95		
	Glu	Thr	Thr	Gln	Ser	Ser	Ser	Thr	Asn	Ala	Thr	Thr	Glu	Glu	Thr	Pro	
				100					105					110			
20	Val	Thr	Gly	Glu	Ala	Thr	Thr	Thr	Thr	Thr	Asn	Gln	Ala	Asn	Thr	Pro	
			115					120					125				
	Ala	Thr	Thr	Gln	Ser	Ser	Asn	Thr	Asn	Ala	Glu	Glu	Leu	Val	Asn	Gln	
		130					135					140					
25	Thr	Ser	Asn	Glu	Thr	Thr	Ser	Asn	Asp	Thr	Asn	Thr	Val	Ser	Ser	Val	
	145					150					155					160	
30	Asn	Ser	Pro	Gln	Asn	Ser	Thr	Asn	Ala	Glu	Asn	Val	Ser	Thr	Thr	Gln	
				165						170					175		
	Asp	Thr	Ser	Thr	Glu	Ala	Thr	Pro	Ser	Asn	Asn	Glu	Ser	Ala	Pro	Gln	
				180					185					190			
35																	
40																	
45																	
50																	
55																	

EP 2 493 498 B1

	Ser	Thr	Asp	Ala	Ser	Asn	Lys	Asp	Val	Val	Asn	Gln	Ala	Val	Asn	Thr	
			195					200					205				
5	Ser	Ala	Pro	Arg	Met	Arg	Ala	Phe	Ser	Leu	Ala	Ala	Val	Ala	Ala	Asp	
		210					215					220					
	Ala	Pro	Val	Ala	Gly	Thr	Asp	Ile	Thr	Asn	Gln	Leu	Thr	Asn	Val	Thr	
		225				230					235					240	
10	Val	Gly	Ile	Asp	Ser	Gly	Thr	Thr	Val	Tyr	Pro	His	Gln	Ala	Gly	Tyr	
					245					250					255		
	Val	Lys	Leu	Asn	Tyr	Gly	Phe	Ser	Val	Pro	Asn	Ser	Ala	Val	Lys	Gly	
				260					265					270			
15	Asp	Thr	Phe	Lys	Ile	Thr	Val	Pro	Lys	Glu	Leu	Asn	Leu	Asn	Gly	Val	
			275					280					285				
	Thr	Ser	Thr	Ala	Lys	Val	Pro	Pro	Ile	Met	Ala	Gly	Asp	Gln	Val	Leu	
		290					295					300					
20	Ala	Asn	Gly	Val	Ile	Asp	Ser	Asp	Gly	Asn	Val	Ile	Tyr	Thr	Phe	Thr	
		305				310					315					320	
	Asp	Tyr	Val	Asn	Thr	Lys	Asp	Asp	Val	Lys	Ala	Thr	Leu	Thr	Met	Pro	
25					325					330					335		
	Ala	Tyr	Ile	Asp	Pro	Glu	Asn	Val	Lys	Lys	Thr	Gly	Asn	Val	Thr	Leu	
				340					345					350			
30	Ala	Thr	Gly	Ile	Gly	Ser	Thr	Thr	Ala	Asn	Lys	Thr	Val	Leu	Val	Asp	
			355					360					365				
	Tyr	Glu	Lys	Tyr	Gly	Lys	Phe	Tyr	Asn	Leu	Ser	Ile	Lys	Gly	Thr	Ile	
		370					375					380					
35	Asp	Gln	Ile	Asp	Lys	Thr	Asn	Asn	Thr	Tyr	Arg	Gln	Thr	Ile	Tyr	Val	
		385				390					395					400	
	Asn	Pro	Ser	Gly	Asp	Asn	Val	Ile	Ala	Pro	Val	Leu	Thr	Gly	Asn	Leu	
				405						410					415		
40	Lys	Pro	Asn	Thr	Asp	Ser	Asn	Ala	Leu	Ile	Asp	Gln	Gln	Asn	Thr	Ser	
				420					425					430			
	Ile	Lys	Val	Tyr	Lys	Val	Asp	Asn	Ala	Ala	Asp	Leu	Ser	Glu	Ser	Tyr	
			435				440						445				
45	Phe	Val	Asn	Pro	Glu	Asn	Phe	Glu	Asp	Val	Thr	Asn	Ser	Val	Asn	Ile	
		450					455					460					
	Thr	Phe	Pro	Asn	Pro	Asn	Gln	Tyr	Lys	Val	Glu	Phe	Asn	Thr	Pro	Asp	
		465				470					475					480	
50	Asp	Gln	Ile	Thr	Thr	Pro	Tyr	Ile	Val	Val	Val	Asn	Gly	His	Ile	Asp	
					485					490					495		
	Pro	Asn	Ser	Lys	Gly	Asp	Leu	Ala	Leu	Arg	Ser	Thr	Leu	Tyr	Gly	Tyr	
55				500					505					510			
	Asn	Ser	Asn	Ile	Ile	Trp	Arg	Ser	Met	Ser	Trp	Asp	Asn	Glu	Val	Ala	

EP 2 493 498 B1

	515						520					525				
5	Phe	Asn	Asn	Gly	Ser	Gly	Ser	Gly	Asp	Gly	Ile	Asp	Lys	Pro	Val	Val
	530						535					540				
	Pro	Glu	Gln	Pro	Asp	Glu	Pro	Gly	Glu	Ile	Glu	Pro	Ile	Pro	Glu	Asp
	545					550					555					560
10	Ser	Asp	Ser	Asp	Pro	Gly	Ser	Asp	Ser	Gly	Ser	Asp	Ser	Asn	Ser	Asp
					565					570					575	
	Ser	Gly	Ser	Asp	Ser	Gly	Ser	Asp	Ser	Thr	Ser	Asp	Ser	Gly	Ser	Asp
				580					585					590		
15	Ser	Ala	Ser	Asp	Ser	Asp	Ser	Ala	Ser	Asp	Ser	Asp	Ser	Ala	Ser	Asp
			595					600					605			
	Ser	Asp	Ser	Ala	Ser	Asp	Ser	Asp	Ser	Ala	Ser	Asp	Ser	Asp	Ser	Asp
	610						615					620				
20	Asn	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp
	625					630					635					640
	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp
					645					650					655	
25	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp
				660					665					670		
	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp
				675				680					685			
30	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp
	690						695					700				
	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp
	705					710					715					720
35	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp
					725					730					735	
	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp
				740					745					750		
40	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Ala
				755				760					765			
45	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp
	770						775					780				
	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp
	785					790					795					800
50	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Glu	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp
					805					810					815	
	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Ala
				820					825					830		
55	Ser	Asp	Ser	Asp	Ser	Gly	Ser	Asp	Ser	Asp	Ser	Ser	Ser	Asp	Ser	Asp
					835			840						845		

EP 2 493 498 B1

	Ser	Glu	Ser	Asp	Ser	Asn	Ser	Asp	Ser	Glu	Ser	Val	Ser	Asn	Asn	Asn
	850						855					860				
5	Val	Val	Pro	Pro	Asn	Ser	Pro	Lys	Asn	Gly	Thr	Asn	Ala	Ser	Asn	Lys
	865					870					875					880
	Asn	Glu	Ala	Lys	Asp	Ser	Lys	Glu	Pro	Leu	Pro	Asp	Thr	Gly	Ser	Glu
					885					890					895	
10	Asp	Glu	Ala	Asn	Thr	Ser	Leu	Ile	Trp	Gly	Leu	Leu	Ala	Ser	Ile	Gly
				900					905					910		
	Ser	Leu	Leu	Leu	Phe	Arg	Arg	Lys	Lys	Glu	Asn	Lys	Asp	Lys	Lys	
15		915						920					925			
	<210> 2															
	<211> 520															
	<212> PRT															
20	<213> Staphylococcus aureus															
	<400> 2															
25																
30																
35																
40																
45																
50																
55																

EP 2 493 498 B1

	Ser	Glu	Asn	Ser	Val	Thr	Gln	Ser	Asp	Ser	Ala	Ser	Asn	Glu	Ser	Lys
	1				5					10				15		
5	Ser	Asn	Asp	Ser	Ser	Ser	Val	Ser	Ala	Ala	Pro	Lys	Thr	Asp	Asp	Thr
				20					25					30		
	Asn	Val	Ser	Asp	Thr	Lys	Thr	Ser	Ser	Asn	Thr	Asn	Asn	Gly	Glu	Thr
			35					40					45			
10	Ser	Val	Ala	Gln	Asn	Pro	Ala	Gln	Gln	Glu	Thr	Thr	Gln	Ser	Ser	Ser
		50					55					60				
	Thr	Asn	Ala	Thr	Thr	Glu	Glu	Thr	Pro	Val	Thr	Gly	Glu	Ala	Thr	Thr
	65					70					75					80
15	Thr	Thr	Thr	Asn	Gln	Ala	Asn	Thr	Pro	Ala	Thr	Thr	Gln	Ser	Ser	Asn
				85						90					95	
	Thr	Asn	Ala	Glu	Glu	Leu	Val	Asn	Gln	Thr	Ser	Asn	Glu	Thr	Thr	Ser
20				100					105				110			
	Asn	Asp	Thr	Asn	Thr	Val	Ser	Ser	Val	Asn	Ser	Pro	Gln	Asn	Ser	Thr
			115					120					125			
25	Asn	Ala	Glu	Asn	Val	Ser	Thr	Thr	Gln	Asp	Thr	Ser	Thr	Glu	Ala	Thr
		130					135					140				
	Pro	Ser	Asn	Asn	Glu	Ser	Ala	Pro	Gln	Ser	Thr	Asp	Ala	Ser	Asn	Lys
	145				150						155					160
30	Asp	Val	Val	Asn	Gln	Ala	Val	Asn	Thr	Ser	Ala	Pro	Arg	Met	Arg	Ala
				165						170					175	
	Phe	Ser	Leu	Ala	Ala	Val	Ala	Ala	Asp	Ala	Pro	Val	Ala	Gly	Thr	Asp
35				180					185					190		
	Ile	Thr	Asn	Gln	Leu	Thr	Asn	Val	Thr	Val	Gly	Ile	Asp	Ser	Gly	Thr
			195					200					205			

40

45

50

55

EP 2 493 498 B1

	Thr	Val	Tyr	Pro	His	Gln	Ala	Gly	Tyr	Val	Lys	Leu	Asn	Tyr	Gly	Phe	
	210						215					220					
5	Ser	Val	Pro	Asn	Ser	Ala	Val	Lys	Gly	Asp	Thr	Phe	Lys	Ile	Thr	Val	
	225					230					235					240	
	Pro	Lys	Glu	Leu	Asn	Leu	Asn	Gly	Val	Thr	Ser	Thr	Ala	Lys	Val	Pro	
					245					250					255		
10	Pro	Ile	Met	Ala	Gly	Asp	Gln	Val	Leu	Ala	Asn	Gly	Val	Ile	Asp	Ser	
				260					265					270			
	Asp	Gly	Asn	Val	Ile	Tyr	Thr	Phe	Thr	Asp	Tyr	Val	Asn	Thr	Lys	Asp	
			275					280					285				
15	Asp	Val	Lys	Ala	Thr	Leu	Thr	Met	Pro	Ala	Tyr	Ile	Asp	Pro	Glu	Asn	
	290						295					300					
	Val	Lys	Lys	Thr	Gly	Asn	Val	Thr	Leu	Ala	Thr	Gly	Ile	Gly	Ser	Thr	
20	305					310					315					320	
	Thr	Ala	Asn	Lys	Thr	Val	Leu	Val	Asp	Tyr	Glu	Lys	Tyr	Gly	Lys	Phe	
					325					330					335		
25	Tyr	Asn	Leu	Ser	Ile	Lys	Gly	Thr	Ile	Asp	Gln	Ile	Asp	Lys	Thr	Asn	
				340					345					350			
	Asn	Thr	Tyr	Arg	Gln	Thr	Ile	Tyr	Val	Asn	Pro	Ser	Gly	Asp	Asn	Val	
			355					360					365				
30	Ile	Ala	Pro	Val	Leu	Thr	Gly	Asn	Leu	Lys	Pro	Asn	Thr	Asp	Ser	Asn	
	370						375					380					
	Ala	Leu	Ile	Asp	Gln	Gln	Asn	Thr	Ser	Ile	Lys	Val	Tyr	Lys	Val	Asp	
	385					390					395					400	
35	Asn	Ala	Ala	Asp	Leu	Ser	Glu	Ser	Tyr	Phe	Val	Asn	Pro	Glu	Asn	Phe	
				405						410					415		
	Glu	Asp	Val	Thr	Asn	Ser	Val	Asn	Ile	Thr	Phe	Pro	Asn	Pro	Asn	Gln	
40				420					425					430			
	Tyr	Lys	Val	Glu	Phe	Asn	Thr	Pro	Asp	Asp	Gln	Ile	Thr	Thr	Pro	Tyr	
			435					440					445				
45	Ile	Val	Val	Val	Asn	Gly	His	Ile	Asp	Pro	Asn	Ser	Lys	Gly	Asp	Leu	
	450						455					460					
	Ala	Leu	Arg	Ser	Thr	Leu	Tyr	Gly	Tyr	Asn	Ser	Asn	Ile	Ile	Trp	Arg	
	465					470					475					480	
50	Ser	Met	Ser	Trp	Asp	Asn	Glu	Val	Ala	Phe	Asn	Asn	Gly	Ser	Gly	Ser	
				485						490					495		
	Gly	Asp	Gly	Ile	Asp	Lys	Pro	Val	Val	Pro	Glu	Gln	Pro	Asp	Glu	Pro	
				500					505					510			
55	Gly	Glu	Ile	Glu	Pro	Ile	Pro	Glu									
			515					520									

<210> 3
<211> 877
<212> PRT
<213> Staphylococcus aureus

5

<400> 3

10

15

20

25

30

35

40

45

50

55

EP 2 493 498 B1

	Met	Lys	Lys	Arg	Ile	Asp	Tyr	Leu	Ser	Asn	Lys	Gln	Asn	Lys	Tyr	Ser	
	1				5					10					15		
5	Ile	Arg	Arg	Phe	Thr	Val	Gly	Thr	Thr	Ser	Val	Ile	Val	Gly	Ala	Thr	
				20					25					30			
	Ile	Leu	Phe	Gly	Ile	Gly	Asn	His	Gln	Ala	Gln	Ala	Ser	Glu	Gln	Ser	
			35					40					45				
10	Asn	Asp	Thr	Thr	Gln	Ser	Ser	Lys	Asn	Asn	Ala	Ser	Ala	Asp	Ser	Glu	
		50					55					60					
	Lys	Asn	Asn	Met	Ile	Glu	Thr	Pro	Gln	Leu	Asn	Thr	Thr	Ala	Asn	Asp	
	65					70					75					80	
15	Thr	Ser	Asp	Ile	Ser	Ala	Asn	Thr	Asn	Ser	Ala	Asn	Val	Asp	Ser	Thr	
				85						90					95		
	Thr	Lys	Pro	Met	Ser	Thr	Gln	Thr	Ser	Asn	Thr	Thr	Thr	Thr	Glu	Pro	
20				100						105					110		
	Ala	Ser	Thr	Asn	Glu	Thr	Pro	Gln	Pro	Thr	Ala	Ile	Lys	Asn	Gln	Ala	
			115					120					125				
25	Thr	Ala	Ala	Lys	Met	Gln	Asp	Gln	Thr	Val	Pro	Gln	Glu	Ala	Asn	Ser	
	130						135					140					
	Gln	Val	Asp	Asn	Lys	Thr	Thr	Asn	Asp	Ala	Asn	Ser	Ile	Ala	Thr	Asn	
	145					150					155					160	
30	Ser	Glu	Leu	Lys	Asn	Ser	Gln	Thr	Leu	Asp	Leu	Pro	Gln	Ser	Ser	Pro	
				165						170					175		
	Gln	Thr	Ile	Ser	Asn	Ala	Gln	Gly	Thr	Ser	Lys	Pro	Ser	Val	Arg	Thr	
			180						185					190			
35	Arg	Ala	Val	Arg	Ser	Leu	Ala	Val	Ala	Glu	Pro	Val	Val	Asn	Ala	Ala	
			195					200					205				
	Asp	Ala	Lys	Gly	Thr	Asn	Val	Asn	Asp	Lys	Val	Thr	Ala	Ser	Asn	Phe	
40		210				215						220					
	Lys	Leu	Glu	Lys	Thr	Thr	Phe	Asp	Pro	Asn	Gln	Ser	Gly	Asn	Thr	Phe	
	225					230					235					240	
45	Met	Ala	Ala	Asn	Phe	Thr	Val	Thr	Asp	Lys	Val	Lys	Ser	Gly	Asp	Tyr	
				245						250					255		
	Phe	Thr	Ala	Lys	Leu	Pro	Asp	Ser	Leu	Thr	Gly	Asn	Gly	Asp	Val	Asp	
				260					265					270			
50	Tyr	Ser	Asn	Ser	Asn	Asn	Thr	Met	Pro	Ile	Ala	Asp	Ile	Lys	Ser	Thr	
			275					280					285				
	Asn	Gly	Asp	Val	Val	Ala	Lys	Ala	Thr	Tyr	Asp	Ile	Leu	Thr	Lys	Thr	
55		290					295					300					

EP 2 493 498 B1

	Tyr	Thr	Phe	Val	Phe	Thr	Asp	Tyr	Val	Asn	Asn	Lys	Glu	Asn	Ile	Asn	
	305					310					315					320	
5	Gly	Gln	Phe	Ser	Leu	Pro	Leu	Phe	Thr	Asp	Arg	Ala	Lys	Ala	Pro	Lys	
					325					330					335		
	Ser	Gly	Thr	Tyr	Asp	Ala	Asn	Ile	Asn	Ile	Ala	Asp	Glu	Met	Phe	Asn	
				340					345					350			
10	Asn	Lys	Ile	Thr	Tyr	Asn	Tyr	Ser	Ser	Pro	Ile	Ala	Gly	Ile	Asp	Lys	
			355					360					365				
	Pro	Asn	Gly	Ala	Asn	Ile	Ser	Ser	Gln	Ile	Ile	Gly	Val	Asp	Thr	Ala	
		370					375					380					
15	Ser	Gly	Gln	Asn	Thr	Tyr	Lys	Gln	Thr	Val	Phe	Val	Asn	Pro	Lys	Gln	
	385					390					395					400	
	Arg	Val	Leu	Gly	Asn	Thr	Trp	Val	Tyr	Ile	Lys	Gly	Tyr	Gln	Asp	Lys	
20					405					410					415		
	Ile	Glu	Glu	Ser	Ser	Gly	Lys	Val	Ser	Ala	Thr	Asp	Thr	Lys	Leu	Arg	
				420					425					430			
	Ile	Phe	Glu	Val	Asn	Asp	Thr	Ser	Lys	Leu	Ser	Asp	Ser	Tyr	Tyr	Ala	
25			435					440					445				
	Asp	Pro	Asn	Asp	Ser	Asn	Leu	Lys	Glu	Val	Thr	Asp	Gln	Phe	Lys	Asn	
		450					455					460					
30	Arg	Ile	Tyr	Tyr	Glu	His	Pro	Asn	Val	Ala	Ser	Ile	Lys	Phe	Gly	Asp	
	465					470					475					480	
	Ile	Thr	Lys	Thr	Tyr	Val	Val	Leu	Val	Glu	Gly	His	Tyr	Asp	Asn	Thr	
					485					490					495		
35	Gly	Lys	Asn	Leu	Lys	Thr	Gln	Val	Ile	Gln	Glu	Asn	Val	Asp	Pro	Val	
				500					505					510			
	Thr	Asn	Arg	Asp	Tyr	Ser	Ile	Phe	Gly	Trp	Asn	Asn	Glu	Asn	Val	Val	
40			515					520					525				
	Arg	Tyr	Gly	Gly	Gly	Ser	Ala	Asp	Gly	Asp	Ser	Ala	Val	Asn	Pro	Lys	
		530					535					540					
	Asp	Pro	Thr	Pro	Gly	Pro	Pro	Val	Asp	Pro	Glu	Pro	Ser	Pro	Asp	Pro	
45						550					555					560	
	Glu	Pro	Glu	Pro	Thr	Pro	Asp	Pro	Glu	Pro	Ser	Pro	Asp	Pro	Glu	Pro	
					565					570					575		
50	Glu	Pro	Ser	Pro	Asp	Pro	Asp	Pro	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	
				580					585					590			
	Gly	Ser	Asp	Ser	Asp	Ser	Gly	Ser	Asp	Ser	Asp	Ser	Glu	Ser	Asp	Ser	
			595					600					605				
55	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Glu	Ser	
		610					615					620					

EP 2 493 498 B1

[illegible]

<210> 4

<211> 508

<212> PRT

<213> Staphylococcus aureus

<400> 4

EP 2 493 498 B1

Ser Glu Gln Ser Asn Asp Thr Thr Gln Ser Ser Lys Asn Asn Ala Ser
 1 5 10 15

5 Ala Asp Ser Glu Lys Asn Asn Met Ile Glu Thr Pro Gln Leu Asn Thr
 20 25 30

Thr Ala Asn Asp Thr Ser Asp Ile Ser Ala Asn Thr Asn Ser Ala Asn

10

15

20

25

30

35

40

45

50

55

EP 2 493 498 B1

55

EP 2 493 498 B1

	Tyr	Gln	Asp	Lys	Ile	Glu	Glu	Ser	Ser	Gly	Lys	Val	Ser	Ala	Thr	Asp	
	370						375					380					
5	Thr	Lys	Leu	Arg	Ile	Phe	Glu	Val	Asn	Asp	Thr	Ser	Lys	Leu	Ser	Asp	
	385					390					395					400	
	Ser	Tyr	Tyr	Ala	Asp	Pro	Asn	Asp	Ser	Asn	Leu	Lys	Glu	Val	Thr	Asp	
					405					410					415		
10	Gln	Phe	Lys	Asn	Arg	Ile	Tyr	Tyr	Glu	His	Pro	Asn	Val	Ala	Ser	Ile	
				420					425					430			
	Lys	Phe	Gly	Asp	Ile	Thr	Lys	Thr	Tyr	Val	Val	Leu	Val	Glu	Gly	His	
			435					440					445				
15	Tyr	Asp	Asn	Thr	Gly	Lys	Asn	Leu	Lys	Thr	Gln	Val	Ile	Gln	Glu	Asn	
	450						455					460					
	Val	Asp	Pro	Val	Thr	Asn	Arg	Asp	Tyr	Ser	Ile	Phe	Gly	Trp	Asn	Asn	
20	465					470					475					480	
	Glu	Asn	Val	Val	Arg	Tyr	Gly	Gly	Gly	Ser	Ala	Asp	Gly	Asp	Ser	Ala	
					485					490					495		
25	Val	Asn	Pro	Lys	Asp	Pro	Thr	Pro	Gly	Pro	Pro	Val					
				500					505								
	<210> 5																
	<211> 316																
	<212> PRT																
30	<213> Staphylococcus aureus																
	<400> 5																
35	Ser	Glu	Gln	Ser	Asn	Asp	Thr	Thr	Gln	Ser	Ser	Lys	Asn	Asn	Ala	Ser	
	1				5					10					15		
	Ala	Asp	Ser	Glu	Lys	Asn	Asn	Met	Ile	Glu	Thr	Pro	Gln	Leu	Asn	Thr	
				20					25					30			
40	Thr	Ala	Asn	Asp	Thr	Ser	Asp	Ile	Ser	Ala	Asn	Thr	Asn	Ser	Ala	Asn	
			35					40					45				
	Val	Asp	Ser	Thr	Thr	Lys	Pro	Met	Ser	Thr	Gln	Thr	Ser	Asn	Thr	Thr	
	50						55					60					
45	Thr	Thr	Glu	Pro	Ala	Ser	Thr	Asn	Glu	Thr	Pro	Gln	Pro	Thr	Ala	Ile	
	65					70					75					80	
	Lys	Asn	Gln	Ala	Thr	Ala	Ala	Lys	Met	Gln	Asp	Gln	Thr	Val	Pro	Gln	
				85						90					95		
50	Glu	Ala	Asn	Ser	Gln	Val	Asp	Asn	Lys	Thr	Thr	Asn	Asp	Ala	Asn	Ser	
				100					105					110			
55	Ile	Ala	Thr	Asn	Ser	Glu	Leu	Lys	Asn	Ser	Gln	Thr	Leu	Asp	Leu	Pro	
			115					120					125				
	Gln	Ser	Ser	Pro	Gln	Thr	Ile	Ser	Asn	Ala	Gln	Gly	Thr	Ser	Lys	Pro	
	130						135					140					

EP 2 493 498 B1

	Ser	Val	Arg	Thr	Arg	Ala	Val	Arg	Ser	Leu	Ala	Val	Ala	Glu	Pro	Val	
	145					150					155					160	
5	Val	Asn	Ala	Ala	Asp	Ala	Lys	Gly	Thr	Asn	Val	Asn	Asp	Lys	Val	Thr	
					165					170					175		
	Ala	Ser	Asn	Phe	Lys	Leu	Glu	Lys	Thr	Thr	Phe	Asp	Pro	Asn	Gln	Ser	
				180						185					190		
10	Gly	Asn	Thr	Phe	Met	Ala	Ala	Asn	Phe	Thr	Val	Thr	Asp	Lys	Val	Lys	
			195					200					205				
	Ser	Gly	Asp	Tyr	Phe	Thr	Ala	Lys	Leu	Pro	Asp	Ser	Leu	Thr	Gly	Asn	
	210						215					220					
15	Gly	Asp	Val	Asp	Tyr	Ser	Asn	Ser	Asn	Asn	Thr	Met	Pro	Ile	Ala	Asp	
	225					230					235					240	
	Ile	Lys	Ser	Thr	Asn	Gly	Asp	Val	Val	Ala	Lys	Ala	Thr	Tyr	Asp	Ile	
20					245					250					255		
	Leu	Thr	Lys	Thr	Tyr	Thr	Phe	Val	Phe	Thr	Asp	Tyr	Val	Asn	Asn	Lys	
				260					265					270			
25	Glu	Asn	Ile	Asn	Gly	Gln	Phe	Ser	Leu	Pro	Leu	Phe	Thr	Asp	Arg	Ala	
			275				280					285					
	Lys	Ala	Pro	Lys	Ser	Gly	Thr	Tyr	Asp	Ala	Asn	Ile	Asn	Ile	Ala	Asp	
	290					295						300					
30	Glu	Met	Phe	Asn	Asn	Lys	Ile	Thr	Tyr	Asn	Tyr	Ser					
	305					310					315						

<210> 6

<211> 331

35 <212> PRT

<213> Staphylococcus aureus

<400> 6

40

45

50

55

EP 2 493 498 B1

	Gly	Thr	Asn	Val	Asn	Asp	Lys	Val	Thr	Ala	Ser	Asn	Phe	Lys	Leu	Glu
	1				5					10					15	
5	Lys	Thr	Thr	Phe	Asp	Pro	Asn	Gln	Ser	Gly	Asn	Thr	Phe	Met	Ala	Ala
				20				25						30		
	Asn	Phe	Thr	Val	Thr	Asp	Lys	Val	Lys	Ser	Gly	Asp	Tyr	Phe	Thr	Ala
			35					40					45			
10	Lys	Leu	Pro	Asp	Ser	Leu	Thr	Gly	Asn	Gly	Asp	Val	Asp	Tyr	Ser	Asn
		50					55					60				
	Ser	Asn	Asn	Thr	Met	Pro	Ile	Ala	Asp	Ile	Lys	Ser	Thr	Asn	Gly	Asp
15	65					70					75					80
	Val	Val	Ala	Lys	Ala	Thr	Tyr	Asp	Ile	Leu	Thr	Lys	Thr	Tyr	Thr	Phe
					85					90					95	
	Val	Phe	Thr	Asp	Tyr	Val	Asn	Asn	Lys	Glu	Asn	Ile	Asn	Gly	Gln	Phe
20				100					105					110		
	Ser	Leu	Pro	Leu	Phe	Thr	Asp	Arg	Ala	Lys	Ala	Pro	Lys	Ser	Gly	Thr

25

30

35

40

45

50

55

EP 2 493 498 B1

	115	120	125
5	Tyr Asp Ala Asn Ile Asn Ile Ala Asp Glu Met Phe Asn Asn Lys Ile 130 135 140		
	Thr Tyr Asn Tyr Ser Ser Pro Ile Ala Gly Ile Asp Lys Pro Asn Gly 145 150 155 160		
10	Ala Asn Ile Ser Ser Gln Ile Ile Gly Val Asp Thr Ala Ser Gly Gln 165 170 175		
	Asn Thr Tyr Lys Gln Thr Val Phe Val Asn Pro Lys Gln Arg Val Leu 180 185 190		
15	Gly Asn Thr Trp Val Tyr Ile Lys Gly Tyr Gln Asp Lys Ile Glu Glu 195 200 205		
	Ser Ser Gly Lys Val Ser Ala Thr Asp Thr Lys Leu Arg Ile Phe Glu 210 215 220		
20	Val Asn Asp Thr Ser Lys Leu Ser Asp Ser Tyr Tyr Ala Asp Pro Asn 225 230 235 240		
	Asp Ser Asn Leu Lys Glu Val Thr Asp Gln Phe Lys Asn Arg Ile Tyr 245 250 255		
25	Tyr Glu His Pro Asn Val Ala Ser Ile Lys Phe Gly Asp Ile Thr Lys 260 265 270		
	Thr Tyr Val Val Leu Val Glu Gly His Tyr Asp Asn Thr Gly Lys Asn 275 280 285		
30	Leu Lys Thr Gln Val Ile Gln Glu Asn Val Asp Pro Val Thr Asn Arg 290 295 300		
	Asp Tyr Ser Ile Phe Gly Trp Asn Asn Glu Asn Val Val Arg Tyr Gly 305 310 315 320		
35	Gly Gly Ser Ala Asp Gly Asp Ser Ala Val Asn 325 330		

<210> 7
 <211> 183
 <212> PRT
 <213> Staphylococcus aureus

<400> 7

EP 2 493 498 B1

	Ser	Ser	Pro	Ile	Ala	Gly	Ile	Asp	Lys	Pro	Asn	Gly	Ala	Asn	Ile	Ser
	1				5					10					15	
5	Ser	Gln	Ile	Ile	Gly	Val	Asp	Thr	Ala	Ser	Gly	Gln	Asn	Thr	Tyr	Lys
				20					25					30		
	Gln	Thr	Val	Phe	Val	Asn	Pro	Lys	Gln	Arg	Val	Leu	Gly	Asn	Thr	Trp
			35					40					45			
10	Val	Tyr	Ile	Lys	Gly	Tyr	Gln	Asp	Lys	Ile	Glu	Glu	Ser	Ser	Gly	Lys
		50					55					60				
	Val	Ser	Ala	Thr	Asp	Thr	Lys	Leu	Arg	Ile	Phe	Glu	Val	Asn	Asp	Thr
15	65					70					75					80
	Ser	Lys	Leu	Ser	Asp	Ser	Tyr	Tyr	Ala	Asp	Pro	Asn	Asp	Ser	Asn	Leu
				85						90					95	
20	Lys	Glu	Val	Thr	Asp	Gln	Phe	Lys	Asn	Arg	Ile	Tyr	Tyr	Glu	His	Pro
				100					105					110		
	Asn	Val	Ala	Ser	Ile	Lys	Phe	Gly	Asp	Ile	Thr	Lys	Thr	Tyr	Val	Val
			115					120					125			
25	Leu	Val	Glu	Gly	His	Tyr	Asp	Asn	Thr	Gly	Lys	Asn	Leu	Lys	Thr	Gln
		130					135					140				
	Val	Ile	Gln	Glu	Asn	Val	Asp	Pro	Val	Thr	Asn	Arg	Asp	Tyr	Ser	Ile
30	145				150						155					160
	Phe	Gly	Trp	Asn	Asn	Glu	Asn	Val	Val	Arg	Tyr	Gly	Gly	Gly	Ser	Ala
				165						170					175	
	Asp	Gly	Asp	Ser	Ala	Val	Asn									
35				180												

<210> 8

<211> 1166

<212> PRT

40 <213> Staphylococcus aureus

<400> 8

45

50

55

EP 2 493 498 B1

	Met	Ile	Asn	Arg	Asp	Asn	Lys	Lys	Ala	Ile	Thr	Lys	Lys	Gly	Met	Ile
	1				5					10					15	
5	Ser	Asn	Arg	Leu	Asn	Lys	Phe	Ser	Ile	Arg	Lys	Tyr	Thr	Val	Gly	Thr
				20					25					30		
	Ala	Ser	Ile	Leu	Val	Gly	Thr	Thr	Leu	Ile	Phe	Gly	Leu	Gly	Asn	Gln
			35					40					45			
10	Glu	Ala	Lys	Ala	Ala	Glu	Asn	Thr	Ser	Thr	Glu	Asn	Ala	Lys	Gln	Asp
		50					55					60				
	Asp	Ala	Thr	Thr	Ser	Asp	Asn	Lys	Glu	Val	Val	Ser	Glu	Thr	Glu	Asn
	65					70					75					80
15	Asn	Ser	Thr	Thr	Glu	Asn	Asn	Ser	Thr	Asn	Pro	Ile	Lys	Lys	Glu	Thr
					85					90					95	
	Asn	Thr	Asp	Ser	Gln	Pro	Glu	Ala	Lys	Lys	Glu	Ser	Thr	Ser	Ser	Ser
20				100					105					110		
	Thr	Gln	Lys	Gln	Gln	Asn	Asn	Val	Thr	Ala	Thr	Thr	Glu	Thr	Lys	Pro
			115					120					125			
	Gln	Asn	Ile	Glu	Lys	Glu	Asn	Val	Lys	Pro	Ser	Thr	Asp	Lys	Thr	Ala
25		130					135					140				
	Thr	Glu	Asp	Thr	Ser	Val	Ile	Leu	Glu	Glu	Lys	Lys	Ala	Pro	Asn	Asn
	145					150					155					160
30	Thr	Asn	Asn	Asp	Val	Thr	Thr	Lys	Pro	Ser	Thr	Ser	Glu	Pro	Ser	Thr
					165					170					175	

35

40

45

50

55

EP 2 493 498 B1

	Ser	Glu	Ile	Gln	Thr	Lys	Pro	Thr	Thr	Pro	Gln	Glu	Ser	Thr	Asn	Ile	
				180					185					190			
5	Glu	Asn	Ser	Gln	Pro	Gln	Pro	Thr	Pro	Ser	Lys	Val	Asp	Asn	Gln	Val	
			195					200					205				
	Thr	Asp	Ala	Thr	Asn	Pro	Lys	Glu	Pro	Val	Asn	Val	Ser	Lys	Glu	Glu	
		210					215					220					
10	Leu	Lys	Asn	Asn	Pro	Glu	Lys	Leu	Lys	Glu	Leu	Val	Arg	Asn	Asp	Ser	
	225					230					235					240	
	Asn	Thr	Asp	His	Ser	Thr	Lys	Pro	Val	Ala	Thr	Ala	Pro	Thr	Ser	Val	
				245						250					255		
15	Ala	Pro	Lys	Arg	Val	Asn	Ala	Lys	Met	Arg	Phe	Ala	Val	Ala	Gln	Pro	
				260					265					270			
	Ala	Ala	Val	Ala	Ser	Asn	Asn	Val	Asn	Asp	Leu	Ile	Lys	Val	Thr	Lys	
			275					280					285				
20	Gln	Thr	Ile	Lys	Val	Gly	Asp	Gly	Lys	Asp	Asn	Val	Ala	Ala	Ala	His	
		290					295					300					
	Asp	Gly	Lys	Asp	Ile	Glu	Tyr	Asp	Thr	Glu	Phe	Thr	Ile	Asp	Asn	Lys	
25	305					310					315					320	
	Val	Lys	Lys	Gly	Asp	Thr	Met	Thr	Ile	Asn	Tyr	Asp	Lys	Asn	Val	Ile	
				325					330						335		
30	Pro	Ser	Asp	Leu	Thr	Asp	Lys	Asn	Asp	Pro	Ile	Asp	Ile	Thr	Asp	Pro	
				340					345					350			
	Ser	Gly	Glu	Val	Ile	Ala	Lys	Gly	Thr	Phe	Asp	Lys	Ala	Thr	Lys	Gln	
			355					360					365				
35	Ile	Thr	Tyr	Thr	Phe	Thr	Asp	Tyr	Val	Asp	Lys	Tyr	Glu	Asp	Ile	Lys	
		370					375					380					
	Ser	Arg	Leu	Thr	Leu	Tyr	Ser	Tyr	Ile	Asp	Lys	Lys	Thr	Val	Pro	Asn	
	385					390					395					400	
40	Glu	Thr	Ser	Leu	Asn	Leu	Thr	Phe	Ala	Thr	Ala	Gly	Lys	Glu	Thr	Ser	
				405					410						415		
	Gln	Asn	Val	Thr	Val	Asp	Tyr	Gln	Asp	Pro	Met	Val	His	Gly	Asp	Ser	
				420					425					430			
45	Asn	Ile	Gln	Ser	Ile	Phe	Thr	Lys	Leu	Asp	Glu	Asp	Lys	Gln	Thr	Ile	
			435					440					445				
	Glu	Gln	Gln	Ile	Tyr	Val	Asn	Pro	Leu	Lys	Lys	Ser	Ala	Thr	Asn	Thr	
		450					455					460					
50	Lys	Val	Asp	Ile	Ala	Gly	Ser	Gln	Val	Asp	Asp	Tyr	Gly	Asn	Ile	Lys	
	465					470					475					480	
	Leu	Gly	Asn	Gly	Ser	Thr	Ile	Ile	Asp	Gln	Asn	Thr	Glu	Ile	Lys	Val	
55				485						490					495		
	Tyr	Lys	Val	Asn	Ser	Asp	Gln	Gln	Leu	Pro	Gln	Ser	Asn	Arg	Ile	Tyr	

EP 2 493 498 B1

	500						505						510					
	Asp	Phe	Ser	Gln	Tyr	Glu	Asp	Val	Thr	Ser	Gln	Phe	Asp	Asn	Lys	Lys		
			515					520					525					
5																		
	Ser	Phe	Ser	Asn	Asn	Val	Ala	Thr	Leu	Asp	Phe	Gly	Asp	Ile	Asn	Ser		
		530					535					540						
10	Ala	Tyr	Ile	Ile	Lys	Val	Val	Ser	Lys	Tyr	Thr	Pro	Thr	Ser	Asp	Gly		
	545					550					555					560		
	Glu	Leu	Asp	Ile	Ala	Gln	Gly	Thr	Ser	Met	Arg	Thr	Thr	Asp	Lys	Tyr		
					565					570					575			
15	Gly	Tyr	Tyr	Asn	Tyr	Ala	Gly	Tyr	Ser	Asn	Phe	Ile	Val	Thr	Ser	Asn		
				580					585					590				
	Asp	Thr	Gly	Gly	Gly	Asp	Gly	Thr	Val	Lys	Pro	Glu	Glu	Lys	Leu	Tyr		
			595					600					605					
20	Lys	Ile	Gly	Asp	Tyr	Val	Trp	Glu	Asp	Val	Asp	Lys	Asp	Gly	Val	Gln		
		610					615					620						
	Gly	Thr	Asp	Ser	Lys	Glu	Lys	Pro	Met	Ala	Asn	Val	Leu	Val	Thr	Leu		
	625					630					635					640		
25	Thr	Tyr	Pro	Asp	Gly	Thr	Thr	Lys	Ser	Val	Arg	Thr	Asp	Ala	Asn	Gly		
					645					650					655			
	His	Tyr	Glu	Phe	Gly	Gly	Leu	Lys	Asp	Gly	Glu	Thr	Tyr	Thr	Val	Lys		
				660					665					670				
30	Phe	Glu	Thr	Pro	Thr	Gly	Tyr	Leu	Pro	Thr	Lys	Val	Asn	Gly	Thr	Thr		
			675					680					685					
	Asp	Gly	Glu	Lys	Asp	Ser	Asn	Gly	Ser	Ser	Val	Thr	Val	Lys	Ile	Asn		
		690					695					700						
35																		
	Gly	Lys	Asp	Asp	Met	Ser	Leu	Asp	Thr	Gly	Phe	Tyr	Lys	Glu	Pro	Lys		
	705					710					715					720		
	Tyr	Asn	Leu	Gly	Asp	Tyr	Val	Trp	Glu	Asp	Thr	Asn	Lys	Asp	Gly	Ile		
				725						730					735			
40																		
	Gln	Asp	Ala	Asn	Glu	Pro	Gly	Ile	Lys	Asp	Val	Lys	Val	Thr	Leu	Lys		
				740					745					750				
	Asp	Ser	Thr	Gly	Lys	Val	Ile	Gly	Thr	Thr	Thr	Thr	Asp	Ala	Ser	Gly		
			755					760					765					
	Lys	Tyr	Lys	Phe	Thr	Asp	Leu	Asp	Asn	Gly	Asn	Tyr	Thr	Val	Glu	Phe		
		770					775					780						
50	Glu	Thr	Pro	Ala	Gly	Tyr	Thr	Pro	Thr	Val	Lys	Asn	Thr	Thr	Ala	Asp		
	785					790					795					800		
	Asp	Lys	Asp	Ser	Asn	Gly	Leu	Thr	Thr	Thr	Gly	Val	Ile	Lys	Asp	Ala		
					805					810					815			
55	Asp	Asn	Met	Thr	Leu	Asp	Arg	Gly	Phe	Tyr	Lys	Thr	Pro	Lys	Tyr	Ser		
				820					825					830				

EP 2 493 498 B1

	Leu	Gly	Asp	Tyr	Val	Trp	Tyr	Asp	Ser	Asn	Lys	Asp	Gly	Lys	Gln	Asp	
			835					840					845				
5	Ser	Thr	Glu	Lys	Gly	Ile	Lys	Asp	Val	Thr	Val	Thr	Leu	Gln	Asn	Glu	
			850				855					860					
	Lys	Gly	Glu	Val	Ile	Gly	Thr	Thr	Lys	Thr	Asp	Glu	Asn	Gly	Lys	Tyr	
			865			870					875					880	
10	Arg	Phe	Asp	Asn	Leu	Asp	Ser	Gly	Lys	Tyr	Lys	Val	Ile	Phe	Glu	Lys	
					885					890						895	
	Pro	Ala	Gly	Leu	Thr	Gln	Thr	Val	Thr	Asn	Thr	Thr	Glu	Asp	Asp	Lys	
				900						905				910			
15	Asp	Ala	Asp	Gly	Gly	Glu	Val	Asp	Val	Thr	Ile	Thr	Asp	His	Asp	Asp	
			915					920					925				
	Phe	Thr	Leu	Asp	Asn	Gly	Tyr	Phe	Glu	Glu	Asp	Thr	Ser	Asp	Ser	Asp	
20			930				935					940					
	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	
						950					955					960	
25	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	
					965					970						975	
	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	
					980				985					990			
30	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	
								1000						1005			
	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	
							1010					1020					
35	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	
						1030					1035					1040	
	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	
40						1045				1050						1055	
	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	
						1060				1065				1070			
45	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	
								1080					1085				
	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	
								1090					1100				
50	Ser	Asp	Ala	Gly	Lys	His	Thr	Pro	Val	Lys	Pro	Met	Ser	Thr	Thr	Lys	
						1110					1115					1120	
	Asp	His	His	Asn	Lys	Ala	Lys	Ala	Leu	Pro	Glu	Thr	Gly	Ser	Glu	Asn	
					1125					1130					1135		
55	Asn	Gly	Ser	Asn	Asn	Ala	Thr	Leu	Phe	Gly	Gly	Leu	Phe	Ala	Ala	Leu	
					1140				1145					1150			

EP 2 493 498 B1

Gly Ser Leu Leu Leu Phe Gly Arg Arg Lys Lys Gln Asn Lys
1155 1160 1165

5

<210> 9
<211> 580
<212> PRT
<213> Staphylococcus aureus

10

<400> 9

15

20

25

30

35

40

45

50

55

EP 2 493 498 B1

	Ala	Glu	Asn	Thr	Ser	Thr	Glu	Asn	Ala	Lys	Gln	Asp	Asp	Ala	Thr	Thr	
	1				5					10					15		
5	Ser	Asp	Asn	Lys	Glu	Val	Val	Ser	Glu	Thr	Glu	Asn	Asn	Ser	Thr	Thr	
				20					25					30			
	Glu	Asn	Asn	Ser	Thr	Asn	Pro	Ile	Lys	Lys	Glu	Thr	Asn	Thr	Asp	Ser	
				35				40					45				
10	Gln	Pro	Glu	Ala	Lys	Lys	Glu	Ser	Thr	Ser	Ser	Ser	Thr	Gln	Lys	Gln	
		50					55					60					
	Gln	Asn	Asn	Val	Thr	Ala	Thr	Thr	Glu	Thr	Lys	Pro	Gln	Asn	Ile	Glu	
	65					70					75					80	
15	Lys	Glu	Asn	Val	Lys	Pro	Ser	Thr	Asp	Lys	Thr	Ala	Thr	Glu	Asp	Thr	
					85					90					95		
	Ser	Val	Ile	Leu	Glu	Glu	Lys	Lys	Ala	Pro	Asn	Asn	Thr	Asn	Asn	Asp	
				100					105					110			
20	Val	Thr	Thr	Lys	Pro	Ser	Thr	Ser	Glu	Pro	Ser	Thr	Ser	Glu	Ile	Gln	
			115					120					125				
	Thr	Lys	Pro	Thr	Thr	Pro	Gln	Glu	Ser	Thr	Asn	Ile	Glu	Asn	Ser	Gln	
25		130					135					140					
	Pro	Gln	Pro	Thr	Pro	Ser	Lys	Val	Asp	Asn	Gln	Val	Thr	Asp	Ala	Thr	
	145					150					155					160	
30	Asn	Pro	Lys	Glu	Pro	Val	Asn	Val	Ser	Lys	Glu	Glu	Leu	Lys	Asn	Asn	
					165					170					175		
	Pro	Glu	Lys	Leu	Lys	Glu	Leu	Val	Arg	Asn	Asp	Ser	Asn	Thr	Asp	His	
				180					185					190			
35	Ser	Thr	Lys	Pro	Val	Ala	Thr	Ala	Pro	Thr	Ser	Val	Ala	Pro	Lys	Arg	
			195					200					205				
	Val	Asn	Ala	Lys	Met	Arg	Phe	Ala	Val	Ala	Gln	Pro	Ala	Ala	Val	Ala	
40		210					215					220					
	Ser	Asn	Asn	Val	Asn	Asp	Leu	Ile	Lys	Val	Thr	Lys	Gln	Thr	Ile	Lys	
	225					230					235					240	
	Val	Gly	Asp	Gly	Lys	Asp	Asn	Val	Ala	Ala	Ala	His	Asp	Gly	Lys	Asp	
45					245					250					255		
	Ile	Glu	Tyr	Asp	Thr	Glu	Phe	Thr	Ile	Asp	Asn	Lys	Val	Lys	Lys	Gly	
				260					265					270			
50	Asp	Thr	Met	Thr	Ile	Asn	Tyr	Asp	Lys	Asn	Val	Ile	Pro	Ser	Asp	Leu	
55																	

EP 2 493 498 B1

	275					280					285					
5	Thr	Asp	Lys	Asn	Asp	Pro	Ile	Asp	Ile	Thr	Asp	Pro	Ser	Gly	Glu	Val
	290						295					300				
	Ile	Ala	Lys	Gly	Thr	Phe	Asp	Lys	Ala	Thr	Lys	Gln	Ile	Thr	Tyr	Thr
	305					310					315					320
10	Phe	Thr	Asp	Tyr	Val	Asp	Lys	Tyr	Glu	Asp	Ile	Lys	Ser	Arg	Leu	Thr
					325					330					335	
	Leu	Tyr	Ser	Tyr	Ile	Asp	Lys	Lys	Thr	Val	Pro	Asn	Glu	Thr	Ser	Leu
				340					345					350		
15	Asn	Leu	Thr	Phe	Ala	Thr	Ala	Gly	Lys	Glu	Thr	Ser	Gln	Asn	Val	Thr
			355					360					365			
	Val	Asp	Tyr	Gln	Asp	Pro	Met	Val	His	Gly	Asp	Ser	Asn	Ile	Gln	Ser
	370						375					380				
20	Ile	Phe	Thr	Lys	Leu	Asp	Glu	Asp	Lys	Gln	Thr	Ile	Glu	Gln	Gln	Ile
	385					390					395					400
	Tyr	Val	Asn	Pro	Leu	Lys	Lys	Ser	Ala	Thr	Asn	Thr	Lys	Val	Asp	Ile
25					405					410					415	
	Ala	Gly	Ser	Gln	Val	Asp	Asp	Tyr	Gly	Asn	Ile	Lys	Leu	Gly	Asn	Gly
				420					425					430		
30	Ser	Thr	Ile	Ile	Asp	Gln	Asn	Thr	Glu	Ile	Lys	Val	Tyr	Lys	Val	Asn
			435					440					445			
	Ser	Asp	Gln	Gln	Leu	Pro	Gln	Ser	Asn	Arg	Ile	Tyr	Asp	Phe	Ser	Gln
	450						455					460				
35	Tyr	Glu	Asp	Val	Thr	Ser	Gln	Phe	Asp	Asn	Lys	Lys	Ser	Phe	Ser	Asn
	465					470					475					480
	Asn	Val	Ala	Thr	Leu	Asp	Phe	Gly	Asp	Ile	Asn	Ser	Ala	Tyr	Ile	Ile
					485					490					495	
40	Lys	Val	Val	Ser	Lys	Tyr	Thr	Pro	Thr	Ser	Asp	Gly	Glu	Leu	Asp	Ile
				500					505					510		
	Ala	Gln	Gly	Thr	Ser	Met	Arg	Thr	Thr	Asp	Lys	Tyr	Gly	Tyr	Tyr	Asn
45			515					520					525			
	Tyr	Ala	Gly	Tyr	Ser	Asn	Phe	Ile	Val	Thr	Ser	Asn	Asp	Thr	Gly	Gly
	530						535					540				
50	Gly	Asp	Gly	Thr	Val	Lys	Pro	Glu	Glu	Lys	Leu	Tyr	Lys	Ile	Gly	Asp
	545					550					555					560
	Tyr	Val	Trp	Glu	Asp	Val	Asp	Lys	Asp	Gly	Val	Gln	Gly	Thr	Asp	Ser
				565						570					575	
55	Lys	Glu	Lys	Pro												
				580												

<210> 10

<211> 995
<212> PRT
<213> Staphylococcus aureus

5 <400> 10

10

15

20

25

30

35

40

45

50

55

EP 2 493 498 B1

	Met	Asn	Asn	Lys	Lys	Thr	Ala	Thr	Asn	Arg	Lys	Gly	Met	Ile	Pro	Asn	
	1				5					10					15		
5	Arg	Leu	Asn	Lys	Phe	Ser	Ile	Arg	Lys	Tyr	Ser	Val	Gly	Thr	Ala	Ser	
				20					25					30			
	Ile	Leu	Val	Gly	Thr	Thr	Leu	Ile	Phe	Gly	Leu	Ser	Gly	His	Glu	Ala	
			35					40					45				
10	Lys	Ala	Ala	Glu	His	Thr	Asn	Gly	Glu	Leu	Asn	Gln	Ser	Lys	Asn	Glu	
		50					55					60					
	Thr	Thr	Ala	Pro	Ser	Glu	Asn	Lys	Thr	Thr	Lys	Lys	Val	Asp	Ser	Arg	
	65					70					75					80	
15	Gln	Leu	Lys	Asp	Asn	Thr	Gln	Thr	Ala	Thr	Ala	Asp	Gln	Pro	Lys	Val	
					85					90					95		
	Thr	Met	Ser	Asp	Ser	Ala	Thr	Val	Lys	Glu	Thr	Ser	Ser	Asn	Met	Gln	
20				100					105					110			
	Ser	Pro	Gln	Asn	Ala	Thr	Ala	Asn	Gln	Ser	Thr	Thr	Lys	Thr	Ser	Asn	
			115					120					125				
25	Val	Thr	Thr	Asn	Asp	Lys	Ser	Ser	Thr	Thr	Tyr	Ser	Asn	Glu	Thr	Asp	
		130				135						140					
	Lys	Ser	Asn	Leu	Thr	Gln	Ala	Lys	Asp	Val	Ser	Thr	Thr	Pro	Lys	Thr	
	145					150				155						160	
30	Thr	Thr	Ile	Lys	Pro	Arg	Thr	Leu	Asn	Arg	Met	Ala	Val	Asn	Thr	Val	
					165					170					175		
	Ala	Ala	Pro	Gln	Gln	Gly	Thr	Asn	Val	Asn	Asp	Lys	Val	His	Phe	Ser	
35				180					185					190			
	Asn	Ile	Asp	Ile	Ala	Ile	Asp	Lys	Gly	His	Val	Asn	Gln	Thr	Thr	Gly	
			195					200					205				
	Lys	Thr	Glu	Phe	Trp	Ala	Thr	Ser	Ser	Asp	Val	Leu	Lys	Leu	Lys	Ala	
40		210				215						220					
	Asn	Tyr	Thr	Ile	Asp	Asp	Ser	Val	Lys	Glu	Gly	Asp	Thr	Phe	Thr	Phe	
	225				230						235					240	
45	Lys	Tyr	Gly	Gln	Tyr	Phe	Arg	Pro	Gly	Ser	Val	Arg	Leu	Pro	Ser	Gln	
				245						250					255		
	Thr	Gln	Asn	Leu	Tyr	Asn	Ala	Gln	Gly	Asn	Ile	Ile	Ala	Lys	Gly	Ile	
				260					265					270			
50	Tyr	Asp	Ser	Thr	Thr	Asn	Thr	Thr	Thr	Tyr	Thr	Phe	Thr	Asn	Tyr	Val	
			275					280					285				
	Asp	Gln	Tyr	Thr	Asn	Val	Arg	Gly	Ser	Phe	Glu	Gln	Val	Ala	Phe	Ala	
55		290					295					300					

EP 2 493 498 B1

	Lys	Arg	Lys	Asn	Ala	Thr	Thr	Asp	Lys	Thr	Ala	Tyr	Lys	Met	Glu	Val	
	305					310					315					320	
5	Thr	Leu	Gly	Asn	Asp	Thr	Tyr	Ser	Glu	Glu	Ile	Ile	Val	Asp	Tyr	Gly	
				325						330					335		
	Asn	Lys	Lys	Ala	Gln	Pro	Leu	Ile	Ser	Ser	Thr	Asn	Tyr	Ile	Asn	Asn	
				340					345					350			
10	Glu	Asp	Leu	Ser	Arg	Asn	Met	Thr	Ala	Tyr	Val	Asn	Gln	Pro	Lys	Asn	
			355					360					365				
	Thr	Tyr	Thr	Lys	Gln	Thr	Phe	Val	Thr	Asn	Leu	Thr	Gly	Tyr	Lys	Phe	
	370						375						380				
15	Asn	Pro	Asn	Ala	Lys	Asn	Phe	Lys	Ile	Tyr	Glu	Val	Thr	Asp	Gln	Asn	
	385					390					395					400	
	Gln	Phe	Val	Asp	Ser	Phe	Thr	Pro	Asp	Thr	Ser	Lys	Leu	Lys	Asp	Val	
				405						410					415		
20	Thr	Asp	Gln	Phe	Asp	Val	Ile	Tyr	Ser	Asn	Asp	Asn	Lys	Thr	Ala	Thr	
			420						425					430			
	Val	Asp	Leu	Met	Lys	Gly	Gln	Thr	Ser	Ser	Asn	Lys	Gln	Tyr	Ile	Ile	
25			435					440					445				
	Gln	Gln	Val	Ala	Tyr	Pro	Asp	Asn	Ser	Ser	Thr	Asp	Asn	Gly	Lys	Ile	
		450					455					460					
30	Asp	Tyr	Thr	Leu	Asp	Thr	Asp	Lys	Thr	Lys	Tyr	Ser	Trp	Ser	Asn	Ser	
	465					470					475					480	
	Tyr	Ser	Asn	Val	Asn	Gly	Ser	Ser	Thr	Ala	Asn	Gly	Asp	Gln	Lys	Lys	
				485						490					495		
35	Tyr	Asn	Leu	Gly	Asp	Tyr	Val	Trp	Glu	Asp	Thr	Asn	Lys	Asp	Gly	Lys	
			500						505					510			
	Gln	Asp	Ala	Asn	Glu	Lys	Gly	Ile	Lys	Gly	Val	Tyr	Val	Ile	Leu	Lys	
			515					520					525				
40	Asp	Ser	Asn	Gly	Lys	Glu	Leu	Asp	Arg	Thr	Thr	Thr	Asp	Glu	Asn	Gly	
		530					535						540				
	Lys	Tyr	Gln	Phe	Thr	Gly	Leu	Ser	Asn	Gly	Thr	Tyr	Ser	Val	Glu	Phe	
	545					550					555					560	
45	Ser	Thr	Pro	Ala	Gly	Tyr	Thr	Pro	Thr	Thr	Ala	Asn	Val	Gly	Thr	Asp	
				565						570					575		
	Asp	Ala	Val	Asp	Ser	Asp	Gly	Leu	Thr	Thr	Thr	Gly	Val	Ile	Lys	Asp	
				580				585						590			
50	Ala	Asp	Asn	Met	Thr	Leu	Asp	Ser	Gly	Phe	Tyr	Lys	Thr	Pro	Lys	Tyr	
			595					600					605				
	Ser	Leu	Gly	Asp	Tyr	Val	Trp	Tyr	Asp	Ser	Asn	Lys	Asp	Gly	Lys	Gln	
55		610					615					620					
	Asp	Ser	Thr	Glu	Lys	Gly	Ile	Lys	Gly	Val	Lys	Val	Thr	Leu	Gln	Asn	

EP 2 493 498 B1

	625				630					635				640		
	Glu	Lys	Gly	Glu	Val	Ile	Gly	Thr	Thr	Glu	Thr	Asp	Glu	Asn	Gly	Lys
					645					650					655	
5																
	Tyr	Arg	Phe	Asp	Asn	Leu	Asp	Ser	Gly	Lys	Tyr	Lys	Val	Ile	Phe	Glu
				660					665					670		
10																
	Lys	Pro	Ala	Gly	Leu	Thr	Gln	Thr	Gly	Thr	Asn	Thr	Thr	Glu	Asp	Asp
			675					680					685			
	Lys	Asp	Ala	Asp	Gly	Gly	Glu	Val	Asp	Val	Thr	Ile	Thr	Asp	His	Asp
		690					695					700				
15																
	Asp	Phe	Thr	Leu	Asp	Asn	Gly	Tyr	Tyr	Glu	Glu	Glu	Thr	Ser	Asp	Ser
	705					710					715					720
	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser
				725						730					735	
20																
	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser
				740					745					750		
	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser
			755				760					765				
25																
	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser
		770					775					780				
	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser
	785					790				795						800
30																
	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser
				805					810					815		
	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser
				820					825					830		
35																
	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser
			835				840					845				
	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Asn	Asp	Ser	Asp	Ser	Asp	Ser
40							855					860				
	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser
	865					870				875						880
45																
	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser
				885					890					895		
	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser
				900					905					910		
50																
	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Asn	Asp	Ser	Asp	Ser
			915				920						925			
	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ala	Gly	Lys	His	Thr	Pro	Ala	Lys	Pro
		930					935					940				
55																
	Met	Ser	Thr	Val	Lys	Asp	Gln	His	Lys	Thr	Ala	Lys	Ala	Leu	Pro	Glu
	945					950					955					960

EP 2 493 498 B1

Thr Gly Ser Glu Asn Asn Asn Ser Asn Asn Gly Thr Leu Phe Gly Gly
965 970 975

5 Leu Phe Ala Ala Leu Gly Ser Leu Leu Leu Phe Gly Arg Arg Lys Lys
980 985 990

Gln Asn Lys
995

10 <210> 11
<211> 468
<212> PRT
<213> Staphylococcus aureus

15 <400> 11

20

25

30

35

40

45

50

55

EP 2 493 498 B1

	Ala	Glu	His	Thr	Asn	Gly	Glu	Leu	Asn	Gln	Ser	Lys	Asn	Glu	Thr	Thr	
	1				5					10					15		
5	Ala	Pro	Ser	Glu	Asn	Lys	Thr	Thr	Lys	Lys	Val	Asp	Ser	Arg	Gln	Leu	
				20					25					30			
	Lys	Asp	Asn	Thr	Gln	Thr	Ala	Thr	Ala	Asp	Gln	Pro	Lys	Val	Thr	Met	
			35				40						45				
10	Ser	Asp	Ser	Ala	Thr	Val	Lys	Glu	Thr	Ser	Ser	Asn	Met	Gln	Ser	Pro	
	50						55					60					
	Gln	Asn	Ala	Thr	Ala	Asn	Gln	Ser	Thr	Thr	Lys	Thr	Ser	Asn	Val	Thr	
15	65					70					75					80	
	Thr	Asn	Asp	Lys	Ser	Ser	Thr	Thr	Tyr	Ser	Asn	Glu	Thr	Asp	Lys	Ser	
				85						90					95		
	Asn	Leu	Thr	Gln	Ala	Lys	Asp	Val	Ser	Thr	Thr	Pro	Lys	Thr	Thr	Thr	
20				100					105					110			
	Ile	Lys	Pro	Arg	Thr	Leu	Asn	Arg	Met	Ala	Val	Asn	Thr	Val	Ala	Ala	
			115					120					125				
25	Pro	Gln	Gln	Gly	Thr	Asn	Val	Asn	Asp	Lys	Val	His	Phe	Ser	Asn	Ile	
	130						135					140					
	Asp	Ile	Ala	Ile	Asp	Lys	Gly	His	Val	Asn	Gln	Thr	Thr	Gly	Lys	Thr	
	145					150					155					160	
30	Glu	Phe	Trp	Ala	Thr	Ser	Ser	Asp	Val	Leu	Lys	Leu	Lys	Ala	Asn	Tyr	
				165						170					175		
	Thr	Ile	Asp	Asp	Ser	Val	Lys	Glu	Gly	Asp	Thr	Phe	Thr	Phe	Lys	Tyr	
35				180					185					190			
	Gly	Gln	Tyr	Phe	Arg	Pro	Gly	Ser	Val	Arg	Leu	Pro	Ser	Gln	Thr	Gln	
			195					200					205				
	Asn	Leu	Tyr	Asn	Ala	Gln	Gly	Asn	Ile	Ile	Ala	Lys	Gly	Ile	Tyr	Asp	
40	210						215					220					
	Ser	Thr	Thr	Asn	Thr	Thr	Thr	Tyr	Thr	Phe	Thr	Asn	Tyr	Val	Asp	Gln	
	225					230					235					240	

EP 2 493 498 B1

	Tyr	Thr	Asn	Val	Arg	Gly	Ser	Phe	Glu	Gln	Val	Ala	Phe	Ala	Lys	Arg	
					245					250					255		
5	Lys	Asn	Ala	Thr	Thr	Asp	Lys	Thr	Ala	Tyr	Lys	Met	Glu	Val	Thr	Leu	
				260					265					270			
	Gly	Asn	Asp	Thr	Tyr	Ser	Glu	Glu	Ile	Ile	Val	Asp	Tyr	Gly	Asn	Lys	
			275					280					285				
10	Lys	Ala	Gln	Pro	Leu	Ile	Ser	Ser	Thr	Asn	Tyr	Ile	Asn	Asn	Glu	Asp	
		290					295					300					
	Leu	Ser	Arg	Asn	Met	Thr	Ala	Tyr	Val	Asn	Gln	Pro	Lys	Asn	Thr	Tyr	
15	305					310					315					320	
	Thr	Lys	Gln	Thr	Phe	Val	Thr	Asn	Leu	Thr	Gly	Tyr	Lys	Phe	Asn	Pro	
					325					330					335		
	Asn	Ala	Lys	Asn	Phe	Lys	Ile	Tyr	Glu	Val	Thr	Asp	Gln	Asn	Gln	Phe	
20				340					345					350			
	Val	Asp	Ser	Phe	Thr	Pro	Asp	Thr	Ser	Lys	Leu	Lys	Asp	Val	Thr	Asp	
			355					360					365				
25	Gln	Phe	Asp	Val	Ile	Tyr	Ser	Asn	Asp	Asn	Lys	Thr	Ala	Thr	Val	Asp	
		370					375					380					
	Leu	Met	Lys	Gly	Gln	Thr	Ser	Ser	Asn	Lys	Gln	Tyr	Ile	Ile	Gln	Gln	
	385					390					395					400	
30	Val	Ala	Tyr	Pro	Asp	Asn	Ser	Ser	Thr	Asp	Asn	Gly	Lys	Ile	Asp	Tyr	
					405					410					415		
	Thr	Leu	Asp	Thr	Asp	Lys	Thr	Lys	Tyr	Ser	Trp	Ser	Asn	Ser	Tyr	Ser	
35				420					425					430			
	Asn	Val	Asn	Gly	Ser	Ser	Thr	Ala	Asn	Gly	Asp	Gln	Lys	Lys	Tyr	Asn	
			435					440					445				
	Leu	Gly	Asp	Tyr	Val	Trp	Glu	Asp	Thr	Asn	Lys	Asp	Gly	Lys	Gln	Asp	
40		450					455					460					
	Ala	Asn	Glu	Lys													
	465																
45	<210>	12															
	<211>	635															
	<212>	PRT															
	<213>	Staphylococcus aureus															
50	<400>	12															
55																	

EP 2 493 498 B1

	Met	Ala	Lys	Tyr	Arg	Gly	Lys	Pro	Phe	Gln	Leu	Tyr	Val	Lys	Leu	Ser
	1				5					10					15	
5	Cys	Ser	Thr	Met	Met	Ala	Thr	Ser	Ile	Ile	Leu	Thr	Asn	Ile	Leu	Pro
				20					25					30		
	Tyr	Asp	Ala	Gln	Ala	Ala	Ser	Glu	Lys	Asp	Thr	Glu	Ile	Thr	Lys	Glu
			35					40					45			
10	Ile	Leu	Ser	Lys	Gln	Asp	Leu	Leu	Asp	Lys	Val	Asp	Lys	Ala	Ile	Arg
15																
20																
25																
30																
35																
40																
45																
50																
55																

EP 2 493 498 B1

	50		55		60												
	Gln	Ile	Glu	Gln	Leu	Lys	Gln	Leu	Ser	Ala	Ser	Ser	Lys	Glu	His	Tyr	
	65					70					75					80	
5																	
	Lys	Ala	Gln	Leu	Asn	Glu	Ala	Lys	Thr	Ala	Ser	Gln	Ile	Asp	Glu	Ile	
				85						90					95		
	Ile	Lys	Arg	Ala	Asn	Glu	Leu	Asp	Ser	Lys	Asp	Asn	Lys	Ser	Ser	His	
10				100					105					110			
	Thr	Glu	Met	Asn	Gly	Gln	Ser	Asp	Ile	Asp	Ser	Lys	Leu	Asp	Gln	Leu	
			115					120					125				
	Leu	Lys	Asp	Leu	Asn	Glu	Val	Ser	Ser	Asn	Val	Asp	Arg	Gly	Gln	Gln	
15		130					135					140					
	Ser	Gly	Glu	Asp	Asp	Leu	Asn	Ala	Met	Lys	Asn	Asp	Met	Ser	Gln	Thr	
	145					150					155					160	
	Ala	Thr	Thr	Lys	His	Gly	Glu	Lys	Asp	Asp	Lys	Asn	Asp	Glu	Ala	Met	
20					165					170					175		
	Val	Asn	Lys	Ala	Leu	Glu	Asp	Leu	Asp	His	Leu	Asn	Gln	Gln	Ile	His	
				180					185					190			
	Lys	Ser	Lys	Asp	Ala	Ser	Lys	Asp	Thr	Ser	Glu	Asp	Pro	Ala	Val	Ser	
25			195					200					205				
	Thr	Thr	Asp	Asn	Asn	His	Glu	Val	Ala	Lys	Thr	Pro	Asn	Asn	Asp	Gly	
		210					215					220					
	Ser	Gly	His	Val	Val	Leu	Asn	Lys	Phe	Leu	Ser	Asn	Glu	Glu	Asn	Gln	
30		225				230					235					240	
	Ser	His	Ser	Asn	Arg	Leu	Thr	Asp	Lys	Leu	Gln	Gly	Ser	Asp	Lys	Ile	
				245						250					255		
35																	
	Asn	His	Ala	Met	Ile	Glu	Lys	Leu	Ala	Lys	Ser	Asn	Ala	Ser	Thr	Gln	
			260						265					270			
	His	Tyr	Thr	Tyr	His	Lys	Leu	Asn	Thr	Leu	Gln	Ser	Leu	Asp	Gln	Arg	
40			275					280					285				
	Ile	Ala	Asn	Thr	Gln	Leu	Pro	Lys	Asn	Gln	Lys	Ser	Asp	Leu	Met	Ser	
		290					295					300					
	Glu	Val	Asn	Lys	Thr	Lys	Glu	Arg	Ile	Lys	Ser	Gln	Arg	Asn	Ile	Ile	
45		305				310					315					320	
	Leu	Glu	Glu	Leu	Ala	Arg	Thr	Asp	Asp	Lys	Lys	Tyr	Ala	Thr	Gln	Ser	
				325						330					335		
	Ile	Leu	Glu	Ser	Ile	Phe	Asn	Lys	Asp	Glu	Ala	Val	Lys	Ile	Leu	Lys	
50				340					345					350			
	Asp	Ile	Arg	Val	Asp	Gly	Lys	Thr	Asp	Gln	Gln	Ile	Ala	Asp	Gln	Ile	
			355					360					365				
	Thr	Arg	His	Ile	Asp	Gln	Leu	Ser	Leu	Thr	Thr	Ser	Asp	Asp	Leu	Leu	
55		370					375					380					

EP 2 493 498 B1

	Thr	Ser	Leu	Ile	Asp	Gln	Ser	Gln	Asp	Lys	Ser	Leu	Leu	Ile	Ser	Gln	
	385					390					395					400	
5	Ile	Leu	Gln	Thr	Lys	Leu	Gly	Lys	Ala	Glu	Ala	Asp	Lys	Leu	Ala	Lys	
					405					410					415		
	Asp	Trp	Thr	Asn	Lys	Gly	Leu	Ser	Asn	Arg	Gln	Ile	Val	Asp	Gln	Leu	
				420					425					430			
10	Lys	Lys	His	Phe	Ala	Ser	Thr	Gly	Asp	Thr	Ser	Ser	Asp	Asp	Ile	Leu	
			435					440					445				
	Lys	Ala	Ile	Leu	Asn	Asn	Ala	Lys	Asp	Lys	Lys	Gln	Ala	Ile	Glu	Thr	
		450					455					460					
15	Ile	Leu	Ala	Thr	Arg	Ile	Glu	Arg	Gln	Lys	Ala	Lys	Leu	Leu	Ala	Asp	
	465					470					475					480	
	Leu	Ile	Thr	Lys	Ile	Glu	Thr	Asp	Gln	Asn	Lys	Ile	Phe	Asn	Leu	Val	
20					485					490					495		
	Lys	Ser	Ala	Leu	Asn	Gly	Lys	Ala	Asp	Asp	Leu	Leu	Asn	Leu	Gln	Lys	
				500					505					510			
25	Arg	Leu	Asn	Gln	Thr	Lys	Lys	Asp	Ile	Asp	Tyr	Ile	Leu	Ser	Pro	Ile	
			515					520					525				
	Val	Asn	Arg	Pro	Ser	Leu	Leu	Asp	Arg	Leu	Asn	Lys	Asn	Gly	Lys	Thr	
		530					535					540					
30	Thr	Asp	Leu	Asn	Lys	Leu	Ala	Asn	Leu	Met	Asn	Gln	Gly	Ser	Asp	Leu	
	545					550					555					560	
	Leu	Asp	Ser	Ile	Pro	Asp	Ile	Pro	Thr	Pro	Lys	Pro	Glu	Lys	Thr	Leu	
				565						570					575		
35	Thr	Leu	Gly	Lys	Gly	Asn	Gly	Leu	Leu	Ser	Gly	Leu	Leu	Asn	Ala	Asp	
				580				585						590			
	Gly	Asn	Val	Ser	Leu	Pro	Lys	Ala	Gly	Glu	Thr	Ile	Lys	Glu	His	Trp	
40			595					600					605				
	Leu	Pro	Ile	Ser	Val	Ile	Val	Gly	Ala	Met	Gly	Val	Leu	Met	Ile	Trp	
		610					615					620					
45	Leu	Ser	Arg	Arg	Asn	Lys	Leu	Lys	Asn	Lys	Ala						
	625					630					635						
	<210>	13															
	<211>	340															
	<212>	PRT															
50	<213>	Staphylococcus aureus															
	<400>	13															
55	Met	Lys	Lys	Lys	Leu	Leu	Val	Leu	Thr	Met	Ser	Thr	Leu	Phe	Ala	Thr	
	1				5					10					15		
	Gln	Ile	Met	Asn	Ser	Asn	His	Ala	Lys	Ala	Ser	Val	Thr	Glu	Ser	Val	
				20					25					30			

EP 2 493 498 B1

	Asp	Lys	Lys	Phe	Val	Val	Pro	Glu	Ser	Gly	Ile	Asn	Lys	Ile	Ile	Pro
			35					40					45			
5	Ala	Tyr	Asp	Glu	Phe	Lys	Asn	Ser	Pro	Lys	Val	Asn	Val	Ser	Asn	Leu
		50					55					60				
	Thr	Asp	Asn	Lys	Asn	Phe	Val	Ala	Ser	Glu	Asp	Lys	Leu	Asn	Lys	Ile
	65					70					75					80
10	Ala	Asp	Ser	Ser	Ala	Ala	Ser	Lys	Ile	Val	Asp	Lys	Asn	Phe	Val	Val
					85					90					95	
	Pro	Glu	Ser	Lys	Leu	Gly	Asn	Ile	Val	Pro	Glu	Tyr	Lys	Glu	Ile	Asn
				100					105					110		
15	Asn	Arg	Val	Asn	Val	Ala	Thr	Asn	Asn	Pro	Ala	Ser	Gln	Gln	Val	Asp
			115					120					125			
	Lys	His	Phe	Val	Ala	Lys	Gly	Pro	Glu	Val	Asn	Arg	Phe	Ile	Thr	Gln
20		130					135					140				
	Asn	Lys	Val	Asn	His	His	Phe	Ile	Thr	Thr	Gln	Thr	His	Tyr	Lys	Lys
	145					150					155					160
	Val	Ile	Thr	Ser	Tyr	Lys	Ser	Thr	His	Val	His	Lys	His	Val	Asn	His
25					165					170					175	
	Ala	Lys	Asp	Ser	Ile	Asn	Lys	His	Phe	Ile	Val	Lys	Pro	Ser	Glu	Ser
				180					185					190		
30	Pro	Arg	Tyr	Thr	His	Pro	Ser	Gln	Ser	Leu	Ile	Ile	Lys	His	His	Phe
			195					200					205			
	Ala	Val	Pro	Gly	Tyr	His	Ala	His	Lys	Phe	Val	Thr	Pro	Gly	His	Ala
		210					215					220				
35	Ser	Ile	Lys	Ile	Asn	His	Phe	Cys	Val	Val	Pro	Gln	Ile	Asn	Ser	Phe
	225					230					235					240
	Lys	Val	Ile	Pro	Pro	Tyr	Gly	His	Asn	Ser	His	Arg	Met	His	Val	Pro
40					245					250					255	
	Ser	Phe	Gln	Asn	Asn	Thr	Thr	Ala	Thr	His	Gln	Asn	Ala	Lys	Val	Asn
				260					265					270		
45	Lys	Ala	Tyr	Asp	Tyr	Lys	Tyr	Phe	Tyr	Ser	Tyr	Lys	Val	Val	Lys	Gly
			275					280					285			
	Val	Lys	Lys	Tyr	Phe	Ser	Phe	Ser	Gln	Ser	Asn	Gly	Tyr	Lys	Ile	Gly
		290					295					300				
50	Lys	Pro	Ser	Leu	Asn	Ile	Lys	Asn	Val	Asn	Tyr	Gln	Tyr	Ala	Val	Pro
	305					310					315					320
	Ser	Tyr	Ser	Pro	Thr	His	Tyr	Val	Pro	Glu	Phe	Lys	Gly	Ser	Leu	Pro
					325					330					335	
55	Ala	Pro	Arg	Val												
				340												

<210> 14
<211> 306
<212> PRT
<213> Staphylococcus aureus

5

<400> 14

10

15

20

25

30

35

40

45

50

55

EP 2 493 498 B1

	Lys	Phe	Val	Val	Pro	Glu	Ser	Gly	Ile	Asn	Lys	Ile	Ile	Pro	Ala	Tyr	
	1				5					10					15		
5	Asp	Glu	Phe	Lys	Asn	Ser	Pro	Lys	Val	Asn	Val	Ser	Asn	Leu	Thr	Asp	
				20					25					30			
	Asn	Lys	Asn	Phe	Val	Ala	Ser	Glu	Asp	Lys	Leu	Asn	Lys	Ile	Ala	Asp	
			35					40					45				
10	Ser	Ser	Ala	Ala	Ser	Lys	Ile	Val	Asp	Lys	Asn	Phe	Val	Val	Pro	Glu	
		50					55					60					
	Ser	Lys	Leu	Gly	Asn	Ile	Val	Pro	Glu	Tyr	Lys	Glu	Ile	Asn	Asn	Arg	
	65					70					75					80	
15	Val	Asn	Val	Ala	Thr	Asn	Asn	Pro	Ala	Ser	Gln	Gln	Val	Asp	Lys	His	
					85					90					95		
	Phe	Val	Ala	Lys	Gly	Pro	Glu	Val	Asn	Arg	Phe	Ile	Thr	Gln	Asn	Lys	
20				100					105					110			
	Val	Asn	His	His	Phe	Ile	Thr	Thr	Gln	Thr	His	Tyr	Lys	Lys	Val	Ile	
			115					120					125				
25	Thr	Ser	Tyr	Lys	Ser	Thr	His	Val	His	Lys	His	Val	Asn	His	Ala	Lys	
		130					135					140					
	Asp	Ser	Ile	Asn	Lys	His	Phe	Ile	Val	Lys	Pro	Ser	Glu	Ser	Pro	Arg	
	145					150					155					160	
30	Tyr	Thr	His	Pro	Ser	Gln	Ser	Leu	Ile	Ile	Lys	His	His	Phe	Ala	Val	
					165					170					175		
	Pro	Gly	Tyr	His	Ala	His	Lys	Phe	Val	Thr	Pro	Gly	His	Ala	Ser	Ile	
				180					185					190			
35	Lys	Ile	Asn	His	Phe	Cys	Val	Val	Pro	Gln	Ile	Asn	Ser	Phe	Lys	Val	
			195					200					205				
	Ile	Pro	Pro	Tyr	Gly	His	Asn	Ser	His	Arg	Met	His	Val	Pro	Ser	Phe	
40		210					215					220					
	Gln	Asn	Asn	Thr	Thr	Ala	Thr	His	Gln	Asn	Ala	Lys	Val	Asn	Lys	Ala	
	225					230					235					240	
45	Tyr	Asp	Tyr	Lys	Tyr	Phe	Tyr	Ser	Tyr	Lys	Val	Val	Lys	Gly	Val	Lys	
					245					250					255		
	Lys	Tyr	Phe	Ser	Phe	Ser	Gln	Ser	Asn	Gly	Tyr	Lys	Ile	Gly	Lys	Pro	
				260					265					270			
50	Ser	Leu	Asn	Ile	Lys	Asn	Val	Asn	Tyr	Gln	Tyr	Ala	Val	Pro	Ser	Tyr	
			275					280					285				
	Ser	Pro	Thr	His	Tyr	Val	Pro	Glu	Phe	Lys	Gly	Ser	Leu	Pro	Ala	Pro	
55		290					295					300					

Arg Val
305

5

<210> 15
<211> 308
<212> PRT
<213> Staphylococcus aureus

10

<400> 15

15

20

25

30

35

40

45

50

55

EP 2 493 498 B1

	Ser	Val	Thr	Glu	Ser	Val	Asp	Lys	Lys	Phe	Val	Val	Pro	Glu	Ser	Gly	
	1				5					10					15		
5	Ile	Asn	Lys	Ile	Ile	Pro	Ala	Tyr	Asp	Glu	Phe	Lys	Asn	Ser	Pro	Lys	
				20					25					30			
	Val	Asn	Val	Ser	Asn	Leu	Thr	Asp	Asn	Lys	Asn	Phe	Val	Ala	Ser	Glu	
			35					40					45				
10	Asp	Lys	Leu	Asn	Lys	Ile	Ala	Asp	Ser	Ser	Ala	Ala	Ser	Lys	Ile	Val	
		50					55					60					
	Asp	Lys	Asn	Phe	Val	Val	Pro	Glu	Ser	Lys	Leu	Gly	Asn	Ile	Val	Pro	
	65					70					75					80	
15	Glu	Tyr	Lys	Glu	Ile	Asn	Asn	Arg	Val	Asn	Val	Ala	Thr	Asn	Asn	Pro	
					85					90					95		
	Ala	Ser	Gln	Gln	Val	Asp	Lys	His	Phe	Val	Ala	Lys	Gly	Pro	Glu	Val	
20				100					105					110			
	Asn	Arg	Phe	Ile	Thr	Gln	Asn	Lys	Val	Asn	His	His	Phe	Ile	Thr	Thr	
			115					120					125				
	Gln	Thr	His	Tyr	Lys	Lys	Val	Ile	Thr	Ser	Tyr	Lys	Ser	Thr	His	Val	
25		130					135					140					
	His	Lys	His	Val	Asn	His	Ala	Lys	Asp	Ser	Ile	Asn	Lys	His	Phe	Ile	
	145					150					155					160	
30	Val	Lys	Pro	Ser	Glu	Ser	Pro	Arg	Tyr	Thr	His	Pro	Ser	Gln	Ser	Leu	
					165					170					175		
	Ile	Ile	Lys	His	His	Phe	Ala	Val	Pro	Gly	Tyr	His	Ala	His	Lys	Phe	
				180					185					190			
35	Val	Thr	Pro	Gly	His	Ala	Ser	Ile	Lys	Ile	Asn	His	Phe	Cys	Val	Val	
			195					200					205				
	Pro	Gln	Ile	Asn	Ser	Phe	Lys	Val	Ile	Pro	Pro	Tyr	Gly	His	Asn	Ser	
40		210					215					220					
	His	Arg	Met	His	Val	Pro	Ser	Phe	Gln	Asn	Asn	Thr	Thr	Ala	Thr	His	
	225					230					235					240	
	Gln	Asn	Ala	Lys	Val	Asn	Lys	Ala	Tyr	Asp	Tyr	Lys	Tyr	Phe	Tyr	Ser	
45					245					250					255		
	Tyr	Lys	Val	Val	Lys	Gly	Val	Lys	Lys	Tyr	Phe	Ser	Phe	Ser	Gln	Ser	
				260					265					270			
50	Asn	Gly	Tyr	Lys	Ile	Gly	Lys	Pro	Ser	Leu	Asn	Ile	Lys	Asn	Val	Asn	
			275					280					285				
	Tyr	Gln	Tyr	Ala	Val	Pro	Ser	Tyr	Ser	Pro	Thr	His	Tyr	Val	Pro	Glu	
55		290					295					300					
	Phe	Lys	Gly	Ser													
	305																

EP 2 493 498 B1

<210> 16
 <211> 300
 <212> PRT
 <213> Staphylococcus aureus

5

<400> 16

10

Lys Phe Val Val Pro Glu Ser Gly Ile Asn Lys Ile Ile Pro Ala Tyr
 1 5 10 15

Asp Glu Phe Lys Asn Ser Pro Lys Val Asn Val Ser Asn Leu Thr Asp
 20 25 30

15

Asn Lys Asn Phe Val Ala Ser Glu Asp Lys Leu Asn Lys Ile Ala Asp
 35 40 45

Ser Ser Ala Ala Ser Lys Ile Val Asp Lys Asn Phe Val Val Pro Glu
 50 55 60

20

Ser Lys Leu Gly Asn Ile Val Pro Glu Tyr Lys Glu Ile Asn Asn Arg
 65 70 75 80

Val Asn Val Ala Thr Asn Asn Pro Ala Ser Gln Gln Val Asp Lys His
 85 90 95

25

Phe Val Ala Lys Gly Pro Glu Val Asn Arg Phe Ile Thr Gln Asn Lys
 100 105 110

Val Asn His His Phe Ile Thr Thr Gln Thr His Tyr Lys Lys Val Ile
 115 120 125

30

Thr Ser Tyr Lys Ser Thr His Val His Lys His Val Asn His Ala Lys
 130 135 140

35

Asp Ser Ile Asn Lys His Phe Ile Val Lys Pro Ser Glu Ser Pro Arg
 145 150 155 160

Tyr Thr His Pro Ser Gln Ser Leu Ile Ile Lys His His Phe Ala Val
 165 170 175

40

Pro Gly Tyr His Ala His Lys Phe Val Thr Pro Gly His Ala Ser Ile
 180 185 190

Lys Ile Asn His Phe Cys Val Val Pro Gln Ile Asn Ser Phe Lys Val
 195 200 205

45

Ile Pro Pro Tyr Gly His Asn Ser His Arg Met His Val Pro Ser Phe
 210 215 220

50

Gln Asn Asn Thr Thr Ala Thr His Gln Asn Ala Lys Val Asn Lys Ala
 225 230 235 240

Tyr Asp Tyr Lys Tyr Phe Tyr Ser Tyr Lys Val Val Lys Gly Val Lys

55

EP 2 493 498 B1

				245					250					255			
	Lys	Tyr	Phe	Ser	Phe	Ser	Gln	Ser	Asn	Gly	Tyr	Lys	Ile	Gly	Lys	Pro	
5				260					265					270			
	Ser	Leu	Asn	Ile	Lys	Asn	Val	Asn	Tyr	Gln	Tyr	Ala	Val	Pro	Ser	Tyr	
			275					280					285				
	Ser	Pro	Thr	His	Tyr	Val	Pro	Glu	Phe	Lys	Gly	Ser					
10		290					295					300					

<210> 17

<211> 121

<212> PRT

15 <213> Staphylococcus aureus

<400> 17

20	Ser	Val	Thr	Glu	Ser	Val	Asp	Lys	Lys	Phe	Val	Val	Pro	Glu	Ser	Gly	
	1				5					10					15		
	Ile	Asn	Lys	Ile	Ile	Pro	Ala	Tyr	Asp	Glu	Phe	Lys	Asn	Ser	Pro	Lys	
				20					25					30			
25	Val	Asn	Val	Ser	Asn	Leu	Thr	Asp	Asn	Lys	Asn	Phe	Val	Ala	Ser	Glu	
			35					40					45				
	Asp	Lys	Leu	Asn	Lys	Ile	Ala	Asp	Ser	Ser	Ala	Ala	Ser	Lys	Ile	Val	
30		50					55					60					
	Asp	Lys	Asn	Phe	Val	Val	Pro	Glu	Ser	Lys	Leu	Gly	Asn	Ile	Val	Pro	
		65				70					75				80		
	Glu	Tyr	Lys	Glu	Ile	Asn	Asn	Arg	Val	Asn	Val	Ala	Thr	Asn	Asn	Pro	
35					85					90					95		
	Ala	Ser	Gln	Gln	Val	Asp	Lys	His	Phe	Val	Ala	Lys	Gly	Pro	Glu	Val	
				100					105					110			
40	Asn	Arg	Phe	Ile	Thr	Gln	Asn	Lys	Val								
			115					120									

<210> 18

<211> 1349

45 <212> PRT

<213> Staphylococcus aureus

<400> 18

50

55

EP 2 493 498 B1

	Met	Leu	Asn	Arg	Glu	Asn	Lys	Thr	Ala	Ile	Thr	Arg	Lys	Gly	Met	Val
	1				5					10					15	
5	Ser	Asn	Arg	Leu	Asn	Lys	Phe	Ser	Ile	Arg	Lys	Tyr	Thr	Val	Gly	Thr
				20					25					30		
	Ala	Ser	Ile	Leu	Val	Gly	Thr	Thr	Leu	Ile	Phe	Gly	Leu	Gly	Asn	Gln
			35					40					45			
10	Glu	Ala	Lys	Ala	Ala	Glu	Ser	Thr	Asn	Lys	Glu	Leu	Asn	Glu	Ala	Thr
		50					55					60				
	Thr	Ser	Ala	Ser	Asp	Asn	Gln	Ser	Ser	Asp	Lys	Val	Asp	Met	Gln	Gln
15	65					70					75					80
20																
25																
30																
35																
40																
45																
50																
55																

EP 2 493 498 B1

	Leu	Asn	Gln	Glu	Asp	Asn	Thr	Lys	Asn	Asp	Asn	Gln	Lys	Glu	Met	Val	
					85					90					95		
5	Ser	Ser	Gln	Gly	Asn	Glu	Thr	Thr	Ser	Asn	Gly	Asn	Lys	Leu	Ile	Glu	
				100					105					110			
	Lys	Glu	Ser	Val	Gln	Ser	Thr	Thr	Gly	Asn	Lys	Val	Glu	Val	Ser	Thr	
			115					120					125				
10	Ala	Lys	Ser	Asp	Glu	Gln	Ala	Ser	Pro	Lys	Ser	Thr	Asn	Glu	Asp	Leu	
		130					135						140				
	Asn	Thr	Lys	Gln	Thr	Ile	Ser	Asn	Gln	Glu	Ala	Leu	Gln	Pro	Asp	Leu	
	145					150					155					160	
15	Gln	Glu	Asn	Lys	Ser	Val	Val	Asn	Val	Gln	Pro	Thr	Asn	Glu	Glu	Asn	
					165					170					175		
	Lys	Lys	Val	Asp	Ala	Lys	Thr	Glu	Ser	Thr	Thr	Leu	Asn	Val	Lys	Ser	
20				180					185					190			
	Asp	Ala	Ile	Lys	Ser	Asn	Asp	Glu	Thr	Leu	Val	Asp	Asn	Asn	Ser	Asn	
			195					200					205				
	Ser	Asn	Asn	Glu	Asn	Asn	Ala	Asp	Ile	Ile	Leu	Pro	Lys	Ser	Thr	Ala	
25		210					215					220					
	Pro	Lys	Arg	Leu	Asn	Thr	Arg	Met	Arg	Ile	Ala	Ala	Val	Gln	Pro	Ser	
	225					230					235					240	
30	Ser	Thr	Glu	Ala	Lys	Asn	Val	Asn	Asp	Leu	Ile	Thr	Ser	Asn	Thr	Thr	
					245					250					255		
	Leu	Thr	Val	Val	Asp	Ala	Asp	Lys	Asn	Asn	Lys	Ile	Val	Pro	Ala	Gln	
				260					265					270			
35	Asp	Tyr	Leu	Ser	Leu	Lys	Ser	Gln	Ile	Thr	Val	Asp	Asp	Lys	Val	Lys	
		275						280					285				
	Ser	Gly	Asp	Tyr	Phe	Thr	Ile	Lys	Tyr	Ser	Asp	Thr	Val	Gln	Val	Tyr	
40		290					295					300					
	Gly	Leu	Asn	Pro	Glu	Asp	Ile	Lys	Asn	Ile	Gly	Asp	Ile	Lys	Asp	Pro	
	305					310					315					320	
	Asn	Asn	Gly	Glu	Thr	Ile	Ala	Thr	Ala	Lys	His	Asp	Thr	Ala	Asn	Asn	
45					325					330					335		
	Leu	Ile	Thr	Tyr	Thr	Phe	Thr	Asp	Tyr	Val	Asp	Arg	Phe	Asn	Ser	Val	
				340					345					350			
50	Gln	Met	Gly	Ile	Asn	Tyr	Ser	Ile	Tyr	Met	Asp	Ala	Asp	Thr	Ile	Pro	
			355					360					365				
	Val	Ser	Lys	Asn	Asp	Val	Glu	Phe	Asn	Val	Thr	Ile	Gly	Asn	Thr	Thr	
		370					375					380					
55	Thr	Lys	Thr	Thr	Ala	Asn	Ile	Gln	Tyr	Pro	Asp	Tyr	Val	Val	Asn	Glu	
	385					390					395					400	

EP 2 493 498 B1

	Lys	Asn	Ser	Ile	Gly	Ser	Ala	Phe	Thr	Glu	Thr	Val	Ser	His	Val	Gly	
					405					410					415		
5	Asn	Lys	Glu	Asn	Pro	Gly	Tyr	Tyr	Lys	Gln	Thr	Ile	Tyr	Val	Asn	Pro	
				420					425					430			
	Ser	Glu	Asn	Ser	Leu	Thr	Asn	Ala	Lys	Leu	Lys	Val	Gln	Ala	Tyr	His	
			435					440					445				
10	Ser	Ser	Tyr	Pro	Asn	Asn	Ile	Gly	Gln	Ile	Asn	Lys	Asp	Val	Thr	Asp	
		450					455					460					
	Ile	Lys	Ile	Tyr	Gln	Val	Pro	Lys	Gly	Tyr	Thr	Leu	Asn	Lys	Gly	Tyr	
	465					470					475					480	
15	Asp	Val	Asn	Thr	Lys	Glu	Leu	Thr	Asp	Val	Thr	Asn	Gln	Tyr	Leu	Gln	
					485					490					495		
	Lys	Ile	Thr	Tyr	Gly	Asp	Asn	Asn	Ser	Ala	Val	Ile	Asp	Phe	Gly	Asn	
				500					505					510			
20	Ala	Asp	Ser	Ala	Tyr	Val	Val	Met	Val	Asn	Thr	Lys	Phe	Gln	Tyr	Thr	
			515					520					525				
	Asn	Ser	Glu	Ser	Pro	Thr	Leu	Val	Gln	Met	Ala	Thr	Leu	Ser	Ser	Thr	
25		530					535					540					
	Gly	Asn	Lys	Ser	Val	Ser	Thr	Gly	Asn	Ala	Leu	Gly	Phe	Thr	Asn	Asn	
	545					550				555						560	
	Gln	Ser	Gly	Gly	Ala	Gly	Gln	Glu	Val	Tyr	Lys	Ile	Gly	Asn	Tyr	Val	
30					565					570					575		
	Trp	Glu	Asp	Thr	Asn	Lys	Asn	Gly	Val	Gln	Glu	Leu	Gly	Glu	Lys	Gly	
				580				585					590				
	Val	Gly	Asn	Val	Thr	Val	Thr	Val	Phe	Asp	Asn	Asn	Thr	Asn	Thr	Lys	
35			595					600					605				
	Val	Gly	Glu	Ala	Val	Thr	Lys	Glu	Asp	Gly	Ser	Tyr	Leu	Ile	Pro	Asn	
		610					615					620					
40	Leu	Pro	Asn	Gly	Asp	Tyr	Arg	Val	Glu	Phe	Ser	Asn	Leu	Pro	Lys	Gly	
	625					630				635						640	
	Tyr	Glu	Val	Thr	Pro	Ser	Lys	Gln	Gly	Asn	Asn	Glu	Glu	Leu	Asp	Ser	
					645					650					655		
45	Asn	Gly	Leu	Ser	Ser	Val	Ile	Thr	Val	Asn	Gly	Lys	Asp	Asn	Leu	Ser	
				660					665					670			
	Ala	Asp	Leu	Gly	Ile	Tyr	Lys	Pro	Lys	Tyr	Asn	Leu	Gly	Asp	Tyr	Val	
			675					680					685				
50	Trp	Glu	Asp	Thr	Asn	Lys	Asn	Gly	Ile	Gln	Asp	Gln	Asp	Glu	Lys	Gly	
		690					695					700					
	Ile	Ser	Gly	Val	Thr	Val	Thr	Leu	Lys	Asp	Glu	Asn	Gly	Asn	Val	Leu	
55	705					710				715						720	
	Lys	Thr	Val	Thr	Thr	Asp	Ala	Asp	Gly	Lys	Tyr	Lys	Phe	Thr	Asp	Leu	

EP 2 493 498 B1

	725							730					735				
5	Asp	Asn	Gly	Asn	Tyr	Lys	Val	Glu	Phe	Thr	Thr	Pro	Glu	Gly	Tyr	Thr	
				740					745					750			
	Pro	Thr	Thr	Val	Thr	Ser	Gly	Ser	Asp	Ile	Glu	Lys	Asp	Ser	Asn	Gly	
				755				760					765				
10	Leu	Thr	Thr	Thr	Gly	Val	Ile	Asn	Gly	Ala	Asp	Asn	Met	Thr	Leu	Asp	
		770					775					780					
	Ser	Gly	Phe	Tyr	Lys	Thr	Pro	Lys	Tyr	Asn	Leu	Gly	Asn	Tyr	Val	Trp	
	785					790					795					800	
15	Glu	Asp	Thr	Asn	Lys	Asp	Gly	Lys	Gln	Asp	Ser	Thr	Glu	Lys	Gly	Ile	
					805					810					815		
	Ser	Gly	Val	Thr	Val	Thr	Leu	Lys	Asn	Glu	Asn	Gly	Glu	Val	Leu	Gln	
				820					825					830			
20	Thr	Thr	Lys	Thr	Asp	Lys	Asp	Gly	Lys	Tyr	Gln	Phe	Thr	Gly	Leu	Glu	
			835					840					845				
	Asn	Gly	Thr	Tyr	Lys	Val	Glu	Phe	Glu	Thr	Pro	Ser	Gly	Tyr	Thr	Pro	
		850					855					860					
25	Thr	Gln	Val	Gly	Ser	Gly	Thr	Asp	Glu	Gly	Ile	Asp	Ser	Asn	Gly	Thr	
	865					870					875					880	
	Ser	Thr	Thr	Gly	Val	Ile	Lys	Asp	Lys	Asp	Asn	Asp	Thr	Ile	Asp	Ser	
				885					890						895		
30	Gly	Phe	Tyr	Lys	Pro	Thr	Tyr	Asn	Leu	Gly	Asp	Tyr	Val	Trp	Glu	Asp	
				900					905					910			
	Thr	Asn	Lys	Asn	Gly	Val	Gln	Asp	Lys	Asp	Glu	Lys	Gly	Ile	Ser	Gly	
			915					920					925				
35	Val	Thr	Val	Thr	Leu	Lys	Asp	Glu	Asn	Asp	Lys	Val	Leu	Lys	Thr	Val	
		930					935					940					
40	Thr	Thr	Asp	Glu	Asn	Gly	Lys	Tyr	Gln	Phe	Thr	Asp	Leu	Asn	Asn	Gly	
	945					950					955					960	
	Thr	Tyr	Lys	Val	Glu	Phe	Glu	Thr	Pro	Ser	Gly	Tyr	Thr	Pro	Thr	Ser	
				965						970					975		
45	Val	Thr	Ser	Gly	Asn	Asp	Thr	Glu	Lys	Asp	Ser	Asn	Gly	Leu	Thr	Thr	
				980					985					990			
	Thr	Gly	Val	Ile	Lys	Asp	Ala	Asp	Asn	Met	Thr	Leu	Asp	Ser	Gly	Phe	
			995					1000					1005				
50	Tyr	Lys	Thr	Pro	Lys	Tyr	Ser	Leu	Gly	Asp	Tyr	Val	Trp	Tyr	Asp	Ser	
		1010					1015					1020					
	Asn	Lys	Asp	Gly	Lys	Gln	Asp	Ser	Thr	Glu	Lys	Gly	Ile	Lys	Asp	Val	
	1025					1030						1035				1040	
55	Lys	Val	Thr	Leu	Leu	Asn	Glu	Lys	Gly	Glu	Val	Ile	Gly	Thr	Thr	Lys	
					1045					1050					1055		

EP 2 493 498 B1

	Thr	Asp	Glu	Asn	Gly	Lys	Tyr	Cys	Phe	Asp	Asn	Leu	Asp	Ser	Gly	Lys	
				1060					1065					1070			
5	Tyr	Lys	Val	Ile	Phe	Glu	Lys	Pro	Ala	Gly	Leu	Thr	Gln	Thr	Val	Thr	
			1075					1080					1085				
	Asn	Thr	Thr	Glu	Asp	Asp	Lys	Asp	Ala	Asp	Gly	Gly	Glu	Val	Asp	Val	
		1090					1095					1100					
10	Thr	Ile	Thr	Asp	His	Asp	Asp	Phe	Thr	Leu	Asp	Asn	Gly	Tyr	Phe	Glu	
	1105					1110					1115					1120	
	Glu	Asp	Thr	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	
				1125					1130						1135		
15	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	
			1140					1145						1150			
	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	
20			1155					1160					1165				
	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	
		1170					1175					1180					
25	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	
	1185					1190				1195						1200	
	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	
				1205					1210					1215			
30	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	
			1220						1225					1230			
	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	
			1235					1240					1245				
35	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	
		1250					1255					1260					
	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	
40		1265				1270				1275						1280	
	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ser	Asp	Ala	Gly	Lys	His	Thr	Pro	Val	
				1285					1290						1295		
45	Lys	Pro	Met	Ser	Thr	Thr	Lys	Asp	His	His	Asn	Lys	Ala	Lys	Ala	Leu	
			1300					1305						1310			
	Pro	Glu	Thr	Gly	Ser	Glu	Asn	Asn	Gly	Ser	Asn	Asn	Ala	Thr	Leu	Phe	
			1315					1320					1325				
50	Gly	Gly	Leu	Phe	Ala	Ala	Leu	Gly	Ser	Leu	Leu	Leu	Phe	Gly	Arg	Arg	
		1330					1335					1340					
	Lys	Lys	Gln	Asn	Lys												
55		1345															
	<210> 19																
	<211> 540																
	<212> PRT																

<213> Staphylococcus aureus

<400> 19

5

10

15

20

25

30

35

40

45

50

55

EP 2 493 498 B1

	Ala	Glu	Ser	Thr	Asn	Lys	Glu	Leu	Asn	Glu	Ala	Thr	Thr	Ser	Ala	Ser	
	1				5					10					15		
5	Asp	Asn	Gln	Ser	Ser	Asp	Lys	Val	Asp	Met	Gln	Gln	Leu	Asn	Gln	Glu	
				20					25					30			
	Asp	Asn	Thr	Lys	Asn	Asp	Asn	Gln	Lys	Glu	Met	Val	Ser	Ser	Gln	Gly	
			35					40					45				
10	Asn	Glu	Thr	Thr	Ser	Asn	Gly	Asn	Lys	Leu	Ile	Glu	Lys	Glu	Ser	Val	
		50					55					60					
	Gln	Ser	Thr	Thr	Gly	Asn	Lys	Val	Glu	Val	Ser	Thr	Ala	Lys	Ser	Asp	
	65					70					75					80	
15	Glu	Gln	Ala	Ser	Pro	Lys	Ser	Thr	Asn	Glu	Asp	Leu	Asn	Thr	Lys	Gln	
					85					90					95		
	Thr	Ile	Ser	Asn	Gln	Glu	Ala	Leu	Gln	Pro	Asp	Leu	Gln	Glu	Asn	Lys	
20				100					105					110			
	Ser	Val	Val	Asn	Val	Gln	Pro	Thr	Asn	Glu	Glu	Asn	Lys	Lys	Val	Asp	
			115					120					125				
	Ala	Lys	Thr	Glu	Ser	Thr	Thr	Leu	Asn	Val	Lys	Ser	Asp	Ala	Ile	Lys	
25		130					135					140					
	Ser	Asn	Asp	Glu	Thr	Leu	Val	Asp	Asn	Asn	Ser	Asn	Ser	Asn	Asn	Glu	
	145					150					155					160	
30	Asn	Asn	Ala	Asp	Ile	Ile	Leu	Pro	Lys	Ser	Thr	Ala	Pro	Lys	Arg	Leu	
				165						170					175		
	Asn	Thr	Arg	Met	Arg	Ile	Ala	Ala	Val	Gln	Pro	Ser	Ser	Thr	Glu	Ala	
				180					185					190			
35	Lys	Asn	Val	Asn	Asp	Leu	Ile	Thr	Ser	Asn	Thr	Thr	Leu	Thr	Val	Val	
			195					200					205				
	Asp	Ala	Asp	Lys	Asn	Asn	Lys	Ile	Val	Pro	Ala	Gln	Asp	Tyr	Leu	Ser	
40		210					215					220					
	Leu	Lys	Ser	Gln	Ile	Thr	Val	Asp	Asp	Lys	Val	Lys	Ser	Gly	Asp	Tyr	
	225				230						235					240	
	Phe	Thr	Ile	Lys	Tyr	Ser	Asp	Thr	Val	Gln	Val	Tyr	Gly	Leu	Asn	Pro	
45				245						250					255		
	Glu	Asp	Ile	Lys	Asn	Ile	Gly	Asp	Ile	Lys	Asp	Pro	Asn	Asn	Gly	Glu	
				260					265					270			
50	Thr	Ile	Ala	Thr	Ala	Lys	His	Asp	Thr	Ala	Asn	Asn	Leu	Ile	Thr	Tyr	
			275					280					285				
	Thr	Phe	Thr	Asp	Tyr	Val	Asp	Arg	Phe	Asn	Ser	Val	Gln	Met	Gly	Ile	
		290					295					300					
55	Asn	Tyr	Ser	Ile	Tyr	Met	Asp	Ala	Asp	Thr	Ile	Pro	Val	Ser	Lys	Asn	

EP 2 493 498 B1

	305		310		315		320
5	Asp Val Glu Phe	Asn Val Thr Ile Gly	Asn Thr Thr Thr Lys Thr Thr				
		325	330			335	
	Ala Asn Ile Gln Tyr Pro Asp Tyr Val	Val Asn Glu Lys Asn Ser Ile					
		340	345			350	
10	Gly Ser Ala Phe Thr Glu Thr Val Ser His Val Gly	Asn Lys Glu Asn					
		355	360			365	
	Pro Gly Tyr Tyr Lys Gln Thr Ile Tyr Val Asn Pro Ser Glu Asn Ser						
		370	375			380	
15	Leu Thr Asn Ala Lys Leu Lys Val Gln Ala Tyr His Ser Ser Tyr Pro						
		385	390			395	400
	Asn Asn Ile Gly Gln Ile Asn Lys Asp Val Thr Asp Ile Lys Ile Tyr						
		405	410			415	
20	Gln Val Pro Lys Gly Tyr Thr Leu Asn Lys Gly Tyr Asp Val Asn Thr						
		420	425			430	
	Lys Glu Leu Thr Asp Val Thr Asn Gln Tyr Leu Gln Lys Ile Thr Tyr						
25		435	440			445	
	Gly Asp Asn Asn Ser Ala Val Ile Asp Phe Gly Asn Ala Asp Ser Ala						
		450	455			460	
30	Tyr Val Val Met Val Asn Thr Lys Phe Gln Tyr Thr Asn Ser Glu Ser						
		465	470			475	480
	Pro Thr Leu Val Gln Met Ala Thr Leu Ser Ser Thr Gly Asn Lys Ser						
		485	490			495	
35	Val Ser Thr Gly Asn Ala Leu Gly Phe Thr Asn Asn Gln Ser Gly Gly						
		500	505			510	
	Ala Gly Gln Glu Val Tyr Lys Ile Gly Asn Tyr Val Trp Glu Asp Thr						
		515	520			525	
40	Asn Lys Asn Gly Val Gln Glu Leu Gly Glu Lys Gly						
		530	535			540	

<210> 20

<211> 199

<212> PRT

<213> Staphylococcus aureus

<400> 20

EP 2 493 498 B1

	Pro	Asp	Tyr	Val	Val	Asn	Glu	Lys	Asn	Ser	Ile	Gly	Ser	Ala	Phe	Thr	
	1				5					10					15		
5	Glu	Thr	Val	Ser	His	Val	Gly	Asn	Lys	Glu	Asn	Pro	Gly	Tyr	Tyr	Lys	
				20					25					30			
	Gln	Thr	Ile	Tyr	Val	Asn	Pro	Ser	Glu	Asn	Ser	Leu	Thr	Asn	Ala	Lys	
			35					40					45				
10	Leu	Lys	Val	Gln	Ala	Tyr	His	Ser	Ser	Tyr	Pro	Asn	Asn	Ile	Gly	Gln	
		50					55					60					
	Ile	Asn	Lys	Asp	Val	Thr	Asp	Ile	Lys	Ile	Tyr	Gln	Val	Pro	Lys	Gly	
15	65					70					75					80	
	Tyr	Thr	Leu	Asn	Lys	Gly	Tyr	Asp	Val	Asn	Thr	Lys	Glu	Leu	Thr	Asp	
				85						90					95		
	Val	Thr	Asn	Gln	Tyr	Leu	Gln	Lys	Ile	Thr	Tyr	Gly	Asp	Asn	Asn	Ser	
20				100					105					110			
	Ala	Val	Ile	Asp	Phe	Gly	Asn	Ala	Asp	Ser	Ala	Tyr	Val	Val	Met	Val	
			115					120					125				
25	Asn	Thr	Lys	Phe	Gln	Tyr	Thr	Asn	Ser	Glu	Ser	Pro	Thr	Leu	Val	Gln	
		130					135					140					
	Met	Ala	Thr	Leu	Ser	Ser	Thr	Gly	Asn	Lys	Ser	Val	Ser	Thr	Gly	Asn	
	145					150					155					160	
30	Ala	Leu	Gly	Phe	Thr	Asn	Asn	Gln	Ser	Gly	Gly	Ala	Gly	Gln	Glu	Val	
					165					170					175		
	Tyr	Lys	Ile	Gly	Asn	Tyr	Val	Trp	Glu	Asp	Thr	Asn	Lys	Asn	Gly	Val	
35				180					185					190			
	Gln	Glu	Leu	Gly	Glu	Lys	Gly										
				195													

<210> 21

40 <211> 516

<212> PRT

<213> Staphylococcus aureus

45 <400> 21

50

55

EP 2 493 498 B1

	Met	Lys	Lys	Lys	Asn	Ile	Tyr	Ser	Ile	Arg	Lys	Leu	Gly	Val	Gly	Ile
	1				5					10					15	
5	Ala	Ser	Val	Thr	Leu	Gly	Thr	Leu	Leu	Ile	Ser	Gly	Gly	Val	Thr	Pro
				20					25					30		
	Ala	Ala	Asn	Ala	Ala	Gln	His	Asp	Glu	Ala	Gln	Gln	Asn	Ala	Phe	Tyr
			35					40					45			
10	Gln	Val	Leu	Asn	Met	Pro	Asn	Leu	Asn	Ala	Asp	Gln	Arg	Asn	Gly	Phe
		50					55					60				
	Ile	Gln	Ser	Leu	Lys	Asp	Asp	Pro	Ser	Gln	Ser	Ala	Asn	Val	Leu	Gly
	65					70					75				80	
15	Glu	Ala	Gln	Lys	Leu	Asn	Asp	Ser	Gln	Ala	Pro	Lys	Ala	Asp	Ala	Gln
					85					90					95	
	Gln	Asn	Asn	Phe	Asn	Lys	Asp	Gln	Gln	Ser	Ala	Phe	Tyr	Glu	Ile	Leu
20				100					105					110		
	Asn	Met	Pro	Asn	Leu	Asn	Glu	Ala	Gln	Arg	Asn	Gly	Phe	Ile	Gln	Ser
			115					120					125			
25	Leu	Lys	Asp	Asp	Pro	Ser	Gln	Ser	Thr	Asn	Val	Leu	Gly	Glu	Ala	Lys
		130					135					140				

30

35

40

45

50

55

EP 2 493 498 B1

	Lys	Leu	Asn	Glu	Ser	Gln	Ala	Pro	Lys	Ala	Asp	Asn	Asn	Phe	Asn	Lys	
	145					150					155					160	
5	Glu	Gln	Gln	Asn	Ala	Phe	Tyr	Glu	Ile	Leu	Asn	Met	Pro	Asn	Leu	Asn	
					165					170					175		
	Glu	Glu	Gln	Arg	Asn	Gly	Phe	Ile	Gln	Ser	Leu	Lys	Asp	Asp	Pro	Ser	
				180					185					190			
10	Gln	Ser	Ala	Asn	Leu	Leu	Ser	Glu	Ala	Lys	Lys	Leu	Asn	Glu	Ser	Gln	
			195					200					205				
	Ala	Pro	Lys	Ala	Asp	Asn	Lys	Phe	Asn	Lys	Glu	Gln	Gln	Asn	Ala	Phe	
	210						215					220					
15	Tyr	Glu	Ile	Leu	His	Leu	Pro	Asn	Leu	Asn	Glu	Glu	Gln	Arg	Asn	Gly	
	225					230					235					240	
	Phe	Ile	Gln	Ser	Leu	Lys	Asp	Asp	Pro	Ser	Gln	Ser	Ala	Asn	Leu	Leu	
					245					250					255		
20	Ala	Glu	Ala	Lys	Lys	Leu	Asn	Asp	Ala	Gln	Ala	Pro	Lys	Ala	Asp	Asn	
				260					265					270			
	Lys	Phe	Asn	Lys	Glu	Gln	Gln	Asn	Ala	Phe	Tyr	Glu	Ile	Leu	His	Leu	
25			275					280					285				
	Pro	Asn	Leu	Thr	Glu	Glu	Gln	Arg	Asn	Gly	Phe	Ile	Gln	Ser	Leu	Lys	
		290					295					300					
30	Asp	Asp	Pro	Ser	Val	Ser	Lys	Glu	Ile	Leu	Ala	Glu	Ala	Lys	Lys	Leu	
	305					310					315					320	
	Asn	Asp	Ala	Gln	Ala	Pro	Lys	Glu	Glu	Asp	Asn	Asn	Lys	Pro	Gly	Lys	
				325						330					335		
35	Glu	Asp	Asn	Asn	Lys	Pro	Gly	Lys	Glu	Asp	Asn	Asn	Lys	Pro	Gly	Lys	
			340						345					350			
	Glu	Asp	Asn	Asn	Lys	Pro	Gly	Lys	Glu	Asp	Asn	Asn	Lys	Pro	Gly	Lys	
			355					360					365				
40	Glu	Asp	Gly	Asn	Lys	Pro	Gly	Lys	Glu	Asp	Asn	Lys	Lys	Pro	Gly	Lys	
	370						375					380					
	Glu	Asp	Gly	Asn	Lys	Pro	Gly	Lys	Glu	Asp	Asn	Lys	Lys	Pro	Gly	Lys	
	385					390					395					400	
45	Glu	Asp	Gly	Asn	Lys	Pro	Gly	Lys	Glu	Asp	Gly	Asn	Lys	Pro	Gly	Lys	
				405						410					415		
	Glu	Asp	Gly	Asn	Gly	Val	His	Val	Val	Lys	Pro	Gly	Asp	Thr	Val	Asn	
				420				425						430			
50	Asp	Ile	Ala	Lys	Ala	Asn	Gly	Thr	Thr	Ala	Asp	Lys	Ile	Ala	Ala	Asp	
		435					440						445				
	Asn	Lys	Leu	Ala	Asp	Lys	Asn	Met	Ile	Lys	Pro	Gly	Gln	Glu	Leu	Val	
55		450					455					460					
	Val	Asp	Lys	Lys	Gln	Pro	Ala	Asn	His	Ala	Asp	Ala	Asn	Lys	Ala	Gln	

EP 2 493 498 B1

	465		470		475		480									
	Ala	Leu	Pro	Glu	Thr	Gly	Glu	Glu	Asn	Pro	Phe	Ile	Gly	Thr	Thr	Val
5					485				490						495	
	Phe	Gly	Gly	Leu	Ser	Leu	Ala	Leu	Gly	Ala	Ala	Leu	Leu	Ala	Gly	Arg
				500					505					510		
10	Arg	Arg	Glu	Leu												
			515													
	<210> 22															
	<211> 289															
	<212> PRT															
15	<213> Staphylococcus aureus															
	<400> 22															
20																
25																
30																
35																
40																
45																
50																
55																

EP 2 493 498 B1

	Ala	Gln	His	Asp	Glu	Ala	Gln	Gln	Asn	Ala	Phe	Tyr	Gln	Val	Leu	Asn	
	1				5					10					15		
5	Met	Pro	Asn	Leu	Asn	Ala	Asp	Gln	Arg	Asn	Gly	Phe	Ile	Gln	Ser	Leu	
				20					25					30			
	Lys	Asp	Asp	Pro	Ser	Gln	Ser	Ala	Asn	Val	Leu	Gly	Glu	Ala	Gln	Lys	
			35					40					45				
10	Leu	Asn	Asp	Ser	Gln	Ala	Pro	Lys	Ala	Asp	Ala	Gln	Gln	Asn	Asn	Phe	
		50					55					60					
	Asn	Lys	Asp	Gln	Gln	Ser	Ala	Phe	Tyr	Glu	Ile	Leu	Asn	Met	Pro	Asn	
15	65					70					75					80	
	Leu	Asn	Glu	Ala	Gln	Arg	Asn	Gly	Phe	Ile	Gln	Ser	Leu	Lys	Asp	Asp	
					85					90					95		
	Pro	Ser	Gln	Ser	Thr	Asn	Val	Leu	Gly	Glu	Ala	Lys	Lys	Leu	Asn	Glu	
20				100					105					110			
	Ser	Gln	Ala	Pro	Lys	Ala	Asp	Asn	Asn	Phe	Asn	Lys	Glu	Gln	Gln	Asn	
			115					120					125				
25	Ala	Phe	Tyr	Glu	Ile	Leu	Asn	Met	Pro	Asn	Leu	Asn	Glu	Glu	Gln	Arg	
		130					135					140					
	Asn	Gly	Phe	Ile	Gln	Ser	Leu	Lys	Asp	Asp	Pro	Ser	Gln	Ser	Ala	Asn	
	145					150					155					160	
30	Leu	Leu	Ser	Glu	Ala	Lys	Lys	Leu	Asn	Glu	Ser	Gln	Ala	Pro	Lys	Ala	
				165						170					175		
	Asp	Asn	Lys	Phe	Asn	Lys	Glu	Gln	Gln	Asn	Ala	Phe	Tyr	Glu	Ile	Leu	
35				180					185					190			
	His	Leu	Pro	Asn	Leu	Asn	Glu	Glu	Gln	Arg	Asn	Gly	Phe	Ile	Gln	Ser	
			195					200					205				
	Leu	Lys	Asp	Asp	Pro	Ser	Gln	Ser	Ala	Asn	Leu	Leu	Ala	Glu	Ala	Lys	
40		210					215					220					
	Lys	Leu	Asn	Asp	Ala	Gln	Ala	Pro	Lys	Ala	Asp	Asn	Lys	Phe	Asn	Lys	
	225					230					235					240	
45	Glu	Gln	Gln	Asn	Ala	Phe	Tyr	Glu	Ile	Leu	His	Leu	Pro	Asn	Leu	Thr	
					245					250					255		
	Glu	Glu	Gln	Arg	Asn	Gly	Phe	Ile	Gln	Ser	Leu	Lys	Asp	Asp	Pro	Ser	
50				260					265					270			
	Val	Ser	Lys	Glu	Ile	Leu	Ala	Glu	Ala	Lys	Lys	Leu	Asn	Asp	Ala	Gln	
			275					280					285				
55	Ala																

<210> 23
 <211> 130
 <212> PRT

EP 2 493 498 B1

<213> Staphylococcus aureus

<400> 23

```

5      Met Asn Phe Asn Asp Ile Glu Thr Met Val Lys Ser Lys Phe Lys Asp
      1              5              10              15

      Ile Lys Lys His Ala Glu Glu Ile Ala His Glu Ile Glu Val Arg Ser
      20              25              30

10     Gly Tyr Leu Arg Lys Ala Glu Gln Tyr Lys Arg Leu Glu Phe Asn Leu
      35              40              45

      Ser Phe Ala Leu Asp Asp Ile Glu Ser Thr Ala Lys Asp Val Gln Thr
15     50              55              60

      Ala Lys Ser Ser Ala Asn Lys Asp Ser Val Thr Val Lys Gly Lys Ala
      65              70              75              80

20     Pro Asn Thr Leu Tyr Ile Glu Lys Arg Asn Leu Met Lys Gln Lys Leu
      85              90              95

      Glu Met Leu Gly Glu Asp Ile Asp Lys Asn Lys Glu Ser Leu Gln Lys
      100             105             110

25     Ala Lys Glu Ile Ala Gly Glu Lys Ala Ser Glu Tyr Phe Asn Lys Ala
      115             120             125

      Met Asn
      130

```

30

<210> 24

<211> 97

<212> PRT

<213> Staphylococcus aureus

35

<400> 24

40

```

      Met Ala Met Ile Lys Met Ser Pro Glu Glu Ile Arg Ala Lys Ser Gln
      1              5              10              15

      Ser Tyr Gly Gln Gly Ser Asp Gln Ile Arg Gln Ile Leu Ser Asp Leu
      20              25              30

45     Thr Arg Ala Gln Gly Glu Ile Ala Ala Asn Trp Glu Gly Gln Ala Phe
      35              40              45

      Ser Arg Phe Glu Glu Gln Phe Gln Gln Leu Ser Pro Lys Val Glu Lys
      50              55              60

50     Phe Ala Gln Leu Leu Glu Glu Ile Lys Gln Gln Leu Asn Ser Thr Ala
      65              70              75              80

      Asp Ala Val Gln Glu Gln Asp Gln Gln Leu Ser Asn Asn Phe Gly Leu
      85              90              95

55     Gln

```

55

<210> 25

EP 2 493 498 B1

<211> 104
 <212> PRT
 <213> Staphylococcus aureus

5 <400> 25

	Met	Gly	Gly	Tyr	Lys	Gly	Ile	Lys	Ala	Asp	Gly	Gly	Lys	Val	Asp	Gln	
	1				5					10					15		
10	Ala	Lys	Gln	Leu	Ala	Ala	Lys	Thr	Ala	Lys	Asp	Ile	Glu	Ala	Cys	Gln	
				20					25					30			
	Lys	Gln	Thr	Gln	Gln	Leu	Ala	Glu	Tyr	Ile	Glu	Gly	Ser	Asp	Trp	Glu	
15				35				40					45				
	Gly	Gln	Phe	Ala	Asn	Lys	Val	Lys	Asp	Val	Leu	Leu	Ile	Met	Ala	Lys	
		50					55					60					
	Phe	Gln	Glu	Glu	Leu	Val	Gln	Pro	Met	Ala	Asp	His	Gln	Lys	Ala	Ile	
20	65					70					75					80	
	Asp	Asn	Leu	Ser	Gln	Asn	Leu	Ala	Lys	Tyr	Asp	Thr	Leu	Ser	Ile	Lys	
					85					90					95		
25	Gln	Gly	Leu	Asp	Arg	Val	Asn	Pro									
				100													

<210> 26
 <211> 302
 <212> PRT
 <213> Staphylococcus aureus

30 <400> 26

35	Met	Lys	Lys	Leu	Leu	Leu	Pro	Leu	Ile	Ile	Met	Leu	Leu	Val	Leu	Ala	
	1				5					10					15		
	Ala	Cys	Gly	Asn	Gln	Gly	Glu	Lys	Asn	Asn	Lys	Ala	Glu	Thr	Lys	Ser	
40				20					25					30			
	Tyr	Lys	Met	Asp	Asp	Gly	Lys	Thr	Val	Asp	Ile	Pro	Lys	Asp	Pro	Lys	
				35				40					45				
	Arg	Ile	Ala	Val	Val	Ala	Pro	Thr	Tyr	Ala	Gly	Gly	Leu	Lys	Lys	Leu	
45		50					55					60					
	Gly	Ala	Asn	Ile	Val	Ala	Val	Asn	Gln	Gln	Val	Asp	Gln	Ser	Lys	Val	
	65					70					75					80	
50	Leu	Lys	Asp	Lys	Phe	Lys	Gly	Val	Thr	Lys	Ile	Gly	Asp	Gly	Asp	Val	

55

EP 2 493 498 B1

	85	90	95
5	Glu Lys Val Ala Lys Glu Lys Pro Asp Leu Ile Ile Val Tyr Ser Thr 100 105 110		
	Asp Lys Asp Ile Lys Lys Tyr Gln Lys Val Ala Pro Thr Val Val Val 115 120 125		
10	Asp Tyr Asn Lys His Lys Tyr Leu Glu Gln Gln Glu Met Leu Gly Lys 130 135 140		
	Ile Val Gly Lys Glu Asp Lys Val Lys Ala Trp Lys Lys Asp Trp Glu 145 150 155 160		
15	Glu Thr Thr Ala Lys Asp Gly Lys Glu Ile Lys Lys Ala Ile Gly Gln 165 170 175		
	Asp Ala Thr Val Ser Leu Phe Asp Glu Phe Asp Lys Lys Leu Tyr Thr 180 185 190		
20	Tyr Gly Asp Asn Trp Gly Arg Gly Gly Glu Val Leu Tyr Gln Ala Phe 195 200 205		
	Gly Leu Lys Met Gln Pro Glu Gln Gln Lys Leu Thr Ala Lys Ala Gly 210 215 220		
25	Trp Ala Glu Val Lys Gln Glu Glu Ile Glu Lys Tyr Ala Gly Asp Tyr 225 230 235 240		
	Ile Val Ser Thr Ser Glu Gly Lys Pro Thr Pro Gly Tyr Glu Ser Thr 245 250 255		
30	Asn Met Trp Lys Asn Leu Lys Ala Thr Lys Glu Gly His Ile Val Lys 260 265 270		
	Val Asp Ala Gly Thr Tyr Trp Tyr Asn Asp Pro Tyr Thr Leu Asp Phe 275 280 285		
35	Met Arg Lys Asp Leu Lys Glu Lys Leu Ile Lys Ala Ala Lys 290 295 300		

<210> 27

<211> 227

<212> PRT

<213> Staphylococcus aureus

<400> 27

EP 2 493 498 B1

	Met	Lys	Asn	Ile	Leu	Lys	Val	Phe	Asn	Thr	Thr	Ile	Leu	Ala	Leu	Ile	
	1				5					10					15		
5	Ile	Ile	Ile	Ala	Thr	Phe	Ser	Asn	Ser	Ala	Asn	Ala	Ala	Asp	Ser	Gly	
				20				25					30				
	Thr	Leu	Asn	Tyr	Glu	Val	Tyr	Lys	Tyr	Asn	Thr	Asn	Asp	Thr	Ser	Ile	
			35					40					45				
10	Ala	Asn	Asp	Tyr	Phe	Asn	Lys	Pro	Ala	Lys	Tyr	Ile	Lys	Lys	Asn	Gly	
		50					55					60					
	Lys	Leu	Tyr	Val	Gln	Ile	Thr	Val	Asn	His	Ser	His	Trp	Ile	Thr	Gly	
	65				70						75					80	
15																	
	Met	Ser	Ile	Glu	Gly	His	Lys	Glu	Asn	Ile	Ile	Ser	Lys	Asn	Thr	Ala	
					85					90					95		
20	Lys	Asp	Glu	Arg	Thr	Ser	Glu	Phe	Glu	Val	Ser	Lys	Leu	Asn	Gly	Lys	
				100					105					110			
	Ile	Asp	Gly	Lys	Ile	Asp	Val	Tyr	Ile	Asp	Glu	Lys	Val	Asn	Gly	Lys	
			115					120					125				
25	Pro	Phe	Lys	Tyr	Asp	His	His	Tyr	Asn	Ile	Thr	Tyr	Lys	Phe	Asn	Gly	
		130					135					140					
	Pro	Thr	Asp	Val	Ala	Gly	Ala	Asn	Ala	Pro	Gly	Lys	Asp	Asp	Lys	Asn	
	145					150					155					160	
30																	
	Ser	Ala	Ser	Gly	Ser	Asp	Lys	Gly	Ser	Asp	Gly	Thr	Thr	Thr	Gly	Gln	
					165					170					175		
	Ser	Glu	Ser	Asn	Ser	Ser	Asn	Lys	Asp	Lys	Val	Glu	Asn	Pro	Gln	Thr	
35				180					185					190			
	Asn	Ala	Gly	Thr	Pro	Ala	Tyr	Ile	Tyr	Ala	Ile	Pro	Val	Ala	Ser	Leu	
			195					200					205				
40	Ala	Leu	Leu	Ile	Ala	Ile	Thr	Leu	Phe	Val	Arg	Lys	Lys	Ser	Lys	Gly	
		210					215					220					
	Asn	Val	Glu														
	225																
45	<210>	28															
	<211>	319															
	<212>	PRT															
	<213>	Staphylococcus aureus															
50	<400>	28															
55																	

EP 2 493 498 B1

	Met	Lys	Thr	Arg	Ile	Val	Ser	Ser	Val	Thr	Thr	Thr	Leu	Leu	Leu	Gly	
	1				5					10					15		
5	Ser	Ile	Leu	Met	Asn	Pro	Val	Ala	Asn	Ala	Ala	Asp	Ser	Asp	Ile	Asn	
				20					25					30			
	Ile	Lys	Thr	Gly	Thr	Thr	Asp	Ile	Gly	Ser	Asn	Thr	Thr	Val	Lys	Thr	
				35				40					45				
10	Gly	Asp	Leu	Val	Thr	Tyr	Asp	Lys	Glu	Asn	Gly	Met	His	Lys	Lys	Val	
		50					55					60					
	Phe	Tyr	Ser	Phe	Ile	Asp	Asp	Lys	Asn	His	Asn	Lys	Lys	Leu	Leu	Val	
15	65					70					75					80	
	Ile	Arg	Thr	Lys	Gly	Thr	Ile	Ala	Gly	Gln	Tyr	Arg	Val	Tyr	Ser	Glu	
					85					90					95		
	Glu	Gly	Ala	Asn	Lys	Ser	Gly	Leu	Ala	Trp	Pro	Ser	Ala	Phe	Lys	Val	
20				100					105					110			
	Gln	Leu	Gln	Leu	Pro	Asp	Asn	Glu	Val	Ala	Gln	Ile	Ser	Asp	Tyr	Tyr	
			115					120					125				
25	Pro	Arg	Asn	Ser	Ile	Asp	Thr	Lys	Glu	Tyr	Met	Ser	Thr	Leu	Thr	Tyr	
		130					135					140					
	Gly	Phe	Asn	Gly	Asn	Val	Thr	Gly	Asp	Asp	Thr	Gly	Lys	Ile	Gly	Gly	
30	145					150					155					160	
	Leu	Ile	Gly	Ala	Asn	Val	Ser	Ile	Gly	His	Thr	Leu	Lys	Tyr	Val	Gln	
					165					170					175		
	Pro	Asp	Phe	Lys	Thr	Ile	Leu	Glu	Ser	Pro	Thr	Asp	Lys	Lys	Val	Gly	
35				180					185					190			
	Trp	Lys	Val	Ile	Phe	Asn	Asn	Met	Val	Asn	Gln	Asn	Trp	Gly	Pro	Tyr	
			195					200					205				
40	Asp	Arg	Asp	Ser	Trp	Asn	Pro	Val	Tyr	Gly	Asn	Gln	Leu	Phe	Met	Lys	
		210					215					220					
	Thr	Arg	Asn	Gly	Ser	Met	Lys	Ala	Ala	Asp	Asn	Phe	Leu	Asp	Pro	Asn	
45	225					230					235					240	
	Lys	Ala	Ser	Ser	Leu	Leu	Ser	Ser	Gly	Phe	Ser	Pro	Asp	Phe	Ala	Thr	
					245					250					255		
	Val	Ile	Thr	Met	Asp	Arg	Lys	Ala	Ser	Lys	Gln	Gln	Thr	Asn	Ile	Asp	
50				260					265					270			
	Val	Ile	Tyr	Glu	Arg	Val	Arg	Asp	Asp	Tyr	Gln	Leu	His	Trp	Thr	Ser	
			275					280					285				
55	Thr	Asn	Trp	Lys	Gly	Thr	Asn	Thr	Lys	Asp	Lys	Trp	Ile	Asp	Arg	Ser	
		290					295					300					
	Ser	Glu	Arg	Tyr	Lys	Ile	Asp	Trp	Glu	Lys	Glu	Glu	Met	Thr	Asn		
	305					310					315						

EP 2 493 498 B1

<210> 29
 <211> 293
 <212> PRT
 <213> Staphylococcus aureus

5

<400> 29

10

Ala Asp Ser Asp Ile Asn Ile Lys Thr Gly Thr Thr Asp Ile Gly Ser
 1 5 10 15

Asn Thr Thr Val Lys Thr Gly Asp Leu Val Thr Tyr Asp Lys Glu Asn
 20 25 30

15

Gly Met Leu Lys Lys Val Phe Tyr Ser Phe Ile Asp Asp Lys Asn His
 35 40 45

Asn Lys Lys Leu Leu Val Ile Arg Thr Lys Gly Thr Ile Ala Gly Gln
 50 55 60

20

Tyr Arg Val Tyr Ser Glu Glu Gly Ala Asn Lys Ser Gly Leu Ala Trp
 65 70 75 80

Pro Ser Ala Phe Lys Val Gln Leu Gln Leu Pro Asp Asn Glu Val Ala
 85 90 95

25

Gln Ile Ser Asp Tyr Tyr Pro Arg Asn Ser Ile Asp Thr Lys Glu Tyr

30

35

40

45

50

55

EP 2 493 498 B1

	100	105	110
5	Met Ser Thr Leu Thr Tyr Gly Phe Asn Gly Asn Val Thr Gly Asp Asp 115 120 125		
	Thr Gly Lys Ile Gly Gly Leu Ile Gly Ala Asn Val Ser Ile Gly His 130 135 140		
10	Thr Leu Lys Tyr Val Gln Pro Asp Phe Lys Thr Ile Leu Glu Ser Pro 145 150 155 160		
	Thr Asp Lys Lys Val Gly Trp Lys Val Ile Phe Asn Asn Met Val Asn 165 170 175		
15	Gln Asn Trp Gly Pro Tyr Asp Arg Asp Ser Trp Asn Pro Val Tyr Gly 180 185 190		
	Asn Gln Leu Phe Met Lys Thr Arg Asn Gly Ser Met Lys Ala Ala Asp 195 200 205		
20	Asn Phe Leu Asp Pro Asn Lys Ala Ser Ser Leu Leu Ser Ser Gly Phe 210 215 220		
	Ser Pro Asp Phe Ala Thr Val Ile Thr Met Asp Arg Lys Ala Ser Lys 225 230 235 240		
25	Gln Gln Thr Asn Ile Asp Val Ile Tyr Glu Arg Val Arg Asp Asp Tyr 245 250 255		
	Gln Leu His Trp Thr Ser Thr Asn Trp Lys Gly Thr Asn Thr Lys Asp 260 265 270		
30	Lys Trp Ile Asp Arg Ser Ser Glu Arg Tyr Lys Ile Asp Trp Glu Lys 275 280 285		
35	Glu Glu Met Thr Asn 290		

<210> 30
 <211> 4
 <212> PRT
 <213> Artificial Sequence

 <220>
 <223> Tetrapeptide loop replacement
 <400> 30

Pro Ser Gly Ser
 1

<210> 31
 <211> 258
 <212> PRT
 <213> Staphylococcus aureus

 <400> 31

EP 2 493 498 B1

	Ala	Asp	Ser	Asp	Ile	Asn	Ile	Lys	Thr	Gly	Thr	Thr	Asp	Ile	Gly	Ser	
	1				5					10					15		
5	Asn	Thr	Thr	Val	Lys	Thr	Gly	Asp	Leu	Val	Thr	Tyr	Asp	Lys	Glu	Asn	
				20					25					30			
	Gly	Met	Leu	Lys	Lys	Val	Phe	Tyr	Ser	Phe	Ile	Asp	Asp	Lys	Asn	His	
10			35					40					45				
	Asn	Lys	Lys	Leu	Leu	Val	Ile	Arg	Thr	Lys	Gly	Thr	Ile	Ala	Gly	Gln	
		50					55					60					
15	Tyr	Arg	Val	Tyr	Ser	Glu	Glu	Gly	Ala	Asn	Lys	Ser	Gly	Leu	Ala	Trp	
	65					70					75					80	
	Pro	Ser	Ala	Phe	Lys	Val	Gln	Leu	Gln	Leu	Pro	Asp	Asn	Glu	Val	Ala	
					85					90					95		
20	Gln	Ile	Ser	Asp	Tyr	Tyr	Pro	Arg	Asn	Ser	Ile	Asp	Thr	Pro	Ser	Gly	
				100					105					110			
	Ser	Val	Gln	Pro	Asp	Phe	Lys	Thr	Ile	Leu	Glu	Ser	Pro	Thr	Asp	Lys	
25			115					120					125				
	Lys	Val	Gly	Trp	Lys	Val	Ile	Phe	Asn	Asn	Met	Val	Asn	Gln	Asn	Trp	
		130					135					140					
30	Gly	Pro	Tyr	Asp	Arg	Asp	Ser	Trp	Asn	Pro	Val	Tyr	Gly	Asn	Gln	Leu	
	145				150						155					160	
	Phe	Met	Lys	Thr	Arg	Asn	Gly	Ser	Met	Lys	Ala	Ala	Asp	Asn	Phe	Leu	
					165					170					175		
35	Asp	Pro	Asn	Lys	Ala	Ser	Ser	Leu	Leu	Ser	Ser	Gly	Phe	Ser	Pro	Asp	
				180					185					190			
	Phe	Ala	Thr	Val	Ile	Thr	Met	Asp	Arg	Lys	Ala	Ser	Lys	Gln	Gln	Thr	
			195					200					205				
40	Asn	Ile	Asp	Val	Ile	Tyr	Glu	Arg	Val	Arg	Asp	Asp	Tyr	Gln	Leu	His	
		210					215					220					
	Trp	Thr	Ser	Thr	Asn	Trp	Lys	Gly	Thr	Asn	Thr	Lys	Asp	Lys	Trp	Ile	
45		225				230					235					240	
	Asp	Arg	Ser	Ser	Glu	Arg	Tyr	Lys	Ile	Asp	Trp	Glu	Lys	Glu	Glu	Met	
					245					250					255		
50	Thr	Asn															
	<210> 32																
	<211> 258																
	<212> PRT																
	<213> Staphylococcus aureus																
55	<400> 32																

EP 2 493 498 B1

	Ala	Asp	Ser	Asp	Ile	Asn	Ile	Lys	Thr	Gly	Thr	Thr	Asp	Ile	Gly	Ser	
	1				5					10					15		
5	Asn	Thr	Thr	Val	Lys	Thr	Gly	Asp	Leu	Val	Thr	Tyr	Asp	Lys	Glu	Asn	
				20					25					30			
	Gly	Met	His	Lys	Lys	Val	Phe	Tyr	Ser	Phe	Ile	Asp	Asp	Lys	Asn	His	
			35					40					45				
10	Asn	Lys	Lys	Leu	Leu	Val	Ile	Arg	Thr	Lys	Gly	Thr	Ile	Ala	Gly	Gln	
		50					55					60					
	Tyr	Arg	Val	Tyr	Ser	Glu	Glu	Gly	Ala	Asn	Lys	Ser	Gly	Leu	Ala	Trp	
15	65					70					75					80	
	Pro	Ser	Ala	Phe	Lys	Val	Gln	Leu	Gln	Leu	Pro	Asp	Asn	Glu	Val	Ala	
					85					90					95		
	Gln	Ile	Ser	Asp	Tyr	Tyr	Pro	Arg	Asn	Ser	Ile	Asp	Thr	Pro	Ser	Gly	
20				100					105					110			
	Ser	Val	Gln	Pro	Asp	Phe	Lys	Thr	Ile	Leu	Glu	Ser	Pro	Thr	Asp	Lys	
			115					120					125				
25	Lys	Val	Gly	Trp	Lys	Val	Ile	Phe	Asn	Asn	Met	Val	Asn	Gln	Asn	Trp	
		130					135					140					
	Gly	Pro	Tyr	Asp	Arg	Asp	Ser	Trp	Asn	Pro	Val	Tyr	Gly	Asn	Gln	Leu	
	145					150					155					160	
30	Phe	Met	Lys	Thr	Arg	Asn	Gly	Ser	Met	Lys	Ala	Ala	Asp	Asn	Phe	Leu	
					165					170					175		
	Asp	Pro	Asn	Lys	Ala	Ser	Ser	Leu	Leu	Ser	Ser	Gly	Phe	Ser	Pro	Asp	
35				180					185					190			
	Phe	Ala	Thr	Val	Ile	Thr	Met	Asp	Arg	Lys	Ala	Ser	Lys	Gln	Gln	Thr	
			195					200					205				
	Asn	Ile	Asp	Val	Ile	Tyr	Glu	Arg	Val	Arg	Asp	Asp	Tyr	Gln	Leu	His	
40		210					215					220					
	Trp	Thr	Ser	Thr	Asn	Trp	Lys	Gly	Thr	Asn	Thr	Lys	Asp	Lys	Trp	Ile	
	225					230					235					240	
45	Asp	Arg	Ser	Ser	Glu	Arg	Tyr	Lys	Ile	Asp	Trp	Glu	Lys	Glu	Glu	Met	
					245					250					255		
	Thr	Asn															
50	<210>	33															
	<211>	50															
	<212>	PRT															
	<213>	Staphylococcus aureus															
55	<400>	33															

EP 2 493 498 B1

	Ala	Asp	Ser	Asp	Ile	Asn	Ile	Lys	Thr	Gly	Thr	Thr	Asp	Ile	Gly	Ser
	1				5					10					15	
5	Asn	Thr	Thr	Val	Lys	Thr	Gly	Asp	Leu	Val	Thr	Tyr	Asp	Lys	Glu	Asn
				20					25					30		
	Gly	Met	His	Lys	Lys	Val	Phe	Tyr	Ser	Phe	Ile	Asp	Asp	Lys	Asn	His
			35					40					45			
10	Asn	Lys														
		50														
	<210> 34															
	<211> 63															
15	<212> PRT															
	<213> Staphylococcus aureus															
	<400> 34															
20	Ala	Asp	Ser	Asp	Ile	Asn	Ile	Lys	Thr	Gly	Thr	Thr	Asp	Ile	Gly	Ser
	1				5					10					15	
	Asn	Thr	Thr	Val	Lys	Thr	Gly	Asp	Leu	Val	Thr	Tyr	Asp	Lys	Glu	Asn
25				20					25					30		
	Gly	Met	His	Lys	Lys	Val	Phe	Tyr	Ser	Phe	Ile	Asp	Asp	Lys	Asn	His
			35					40					45			
30	Asn	Lys	Lys	Leu	Leu	Val	Ile	Arg	Thr	Lys	Gly	Thr	Ile	Ala	Gly	
		50					55					60				
	<210> 35															
	<211> 53															
	<212> PRT															
35	<213> Staphylococcus aureus															
	<400> 35															
40	Ala	Asp	Ser	Asp	Ile	Asn	Ile	Lys	Thr	Gly	Thr	Thr	Asp	Ile	Gly	Ser
	1				5					10					15	
	Asn	Thr	Thr	Val	Lys	Thr	Gly	Asp	Leu	Val	Thr	Tyr	Asp	Lys	Glu	Asn
				20					25					30		
45	Gly	Met	His	Lys	Lys	Val	Phe	Tyr	Ser	Phe	Ile	Asp	Asp	Lys	Asn	His
			35					40					45			
	Asn	Lys	Lys	Leu	Leu											
50		50														
	<210> 36															
	<211> 50															
	<212> PRT															
55	<213> Staphylococcus aureus															
	<400> 36															

EP 2 493 498 B1

Ala Asp Ser Asp Ile Asn Ile Lys Thr Gly Thr Thr Asp Ile Gly Ser
 1 5 10 15
 5 Asn Thr Thr Val Lys Thr Gly Asp Leu Val Thr Tyr Asp Lys Glu Asn
 20 25 30
 Gly Met Leu Lys Lys Val Phe Tyr Ser Phe Ile Asp Asp Lys Asn His
 35 40 45
 10 Asn Lys
 50
 <210> 37
 <211> 63
 15 <212> PRT
 <213> Staphylococcus aureus
 <400> 37
 20
 Ala Asp Ser Asp Ile Asn Ile Lys Thr Gly Thr Thr Asp Ile Gly Ser
 1 5 10 15
 25 Asn Thr Thr Val Lys Thr Gly Asp Leu Val Thr Tyr Asp Lys Glu Asn
 20 25 30
 Gly Met Leu Lys Lys Val Phe Tyr Ser Phe Ile Asp Asp Lys Asn His
 35 40 45
 30 Asn Lys Lys Leu Leu Val Ile Arg Thr Lys Gly Thr Ile Ala Gly
 50 55 60
 <210> 38
 35 <211> 53
 <212> PRT
 <213> Staphylococcus aureus
 <400> 38
 40
 Ala Asp Ser Asp Ile Asn Ile Lys Thr Gly Thr Thr Asp Ile Gly Ser
 1 5 10 15
 45 Asn Thr Thr Val Lys Thr Gly Asp Leu Val Thr Tyr Asp Lys Glu Asn
 20 25 30
 Gly Met Leu Lys Lys Val Phe Tyr Ser Phe Ile Asp Asp Lys Asn His
 35 40 45
 50 Asn Lys Lys Leu Leu
 50
 <210> 39
 <211> 256
 55 <212> PRT
 <213> Staphylococcus aureus
 <400> 39

EP 2 493 498 B1

	Met	Met	Lys	Arg	Leu	Asn	Lys	Leu	Val	Leu	Gly	Ile	Ile	Phe	Leu	Phe	
	1				5					10					15		
5	Leu	Val	Ile	Ser	Ile	Thr	Ala	Gly	Cys	Gly	Ile	Gly	Lys	Glu	Ala	Glu	
				20					25					30			
	Val	Lys	Lys	Ser	Phe	Glu	Lys	Thr	Leu	Ser	Met	Tyr	Pro	Ile	Lys	Asn	
			35					40					45				
10	Leu	Glu	Asp	Leu	Tyr	Asp	Lys	Glu	Gly	Tyr	Arg	Asp	Asp	Gln	Phe	Asp	
		50					55					60					
	Lys	Asn	Asp	Lys	Gly	Thr	Trp	Ile	Ile	Asn	Ser	Glu	Met	Val	Ile	Gln	
	65					70					75					80	
15	Pro	Asn	Asn	Glu	Asp	Met	Val	Ala	Lys	Gly	Met	Val	Leu	Tyr	Met	Asn	
				85						90					95		
	Arg	Asn	Thr	Lys	Thr	Thr	Asn	Gly	Tyr	Tyr	Tyr	Val	Asp	Val	Thr	Lys	
20				100					105					110			
	Asp	Glu	Asp	Glu	Gly	Lys	Pro	His	Asp	Asn	Glu	Lys	Arg	Tyr	Pro	Val	
			115					120					125				
25	Lys	Met	Val	Asp	Asn	Lys	Ile	Ile	Pro	Thr	Lys	Glu	Ile	Lys	Asp	Glu	
		130					135					140					
	Lys	Ile	Lys	Lys	Glu	Ile	Glu	Asn	Phe	Lys	Phe	Phe	Val	Gln	Tyr	Gly	
	145					150					155					160	
30	Asp	Phe	Lys	Asn	Leu	Lys	Asn	Tyr	Lys	Asp	Gly	Asp	Ile	Ser	Tyr	Asn	
					165					170					175		
	Pro	Glu	Val	Pro	Ser	Tyr	Ser	Ala	Lys	Tyr	Gln	Leu	Thr	Asn	Asp	Asp	
35				180					185					190			
	Tyr	Asn	Val	Lys	Gln	Leu	Arg	Lys	Arg	Tyr	Asp	Ile	Pro	Thr	Ser	Lys	
			195					200					205				
40	Ala	Pro	Lys	Leu	Leu	Leu	Lys	Gly	Ser	Gly	Asn	Leu	Lys	Gly	Ser	Ser	
		210					215					220					
	Val	Gly	Tyr	Lys	Asp	Ile	Glu	Phe	Thr	Phe	Val	Glu	Lys	Lys	Glu	Glu	
	225					230					235					240	
45	Asn	Ile	Tyr	Phe	Ser	Asp	Ser	Leu	Asp	Tyr	Lys	Lys	Ser	Gly	Asp	Val	
					245					250					255		

<210> 40

<211> 256

50 <212> PRT

<213> Staphylococcus aureus

<400> 40

55

EP 2 493 498 B1

	Met	Met	Lys	Arg	Leu	Asn	Lys	Leu	Val	Leu	Gly	Ile	Ile	Phe	Leu	Phe
	1				5					10					15	
5	Leu	Val	Ile	Ser	Ile	Thr	Ala	Gly	Cys	Gly	Ile	Gly	Lys	Glu	Ala	Glu
				20					25					30		
	Val	Lys	Lys	Ser	Phe	Glu	Lys	Thr	Leu	Ser	Met	Tyr	Pro	Ile	Lys	Asn
			35					40					45			
10	Leu	Glu	Asp	Leu	Tyr	Asp	Lys	Glu	Gly	Tyr	Arg	Asp	Asp	Gln	Phe	Asp
		50					55					60				
	Lys	Asn	Asp	Lys	Gly	Thr	Trp	Ile	Ile	Asn	Ser	Glu	Met	Val	Ile	Gln
	65					70					75					80
15	Pro	Asn	Asn	Glu	Asp	Met	Val	Ala	Lys	Gly	Met	Val	Leu	Tyr	Met	Asn
					85					90					95	
	Arg	Asn	Thr	Lys	Thr	Thr	Asn	Gly	Tyr	Tyr	Tyr	Val	Asp	Val	Thr	Lys
20				100					105					110		
	Asp	Glu	Asp	Glu	Gly	Lys	Pro	His	Asp	Asn	Glu	Lys	Arg	Tyr	Pro	Val
			115					120					125			
25	Lys	Met	Val	Asp	Asn	Lys	Ile	Ile	Pro	Thr	Lys	Glu	Ile	Lys	Asp	Glu
		130					135					140				
	Lys	Leu	Lys	Lys	Glu	Ile	Glu	Asn	Phe	Lys	Phe	Phe	Val	Gln	Tyr	Gly
	145					150					155					160
30	Asp	Phe	Lys	Asn	Ile	Lys	Asn	Tyr	Lys	Asp	Gly	Asp	Ile	Ser	Tyr	Asn
					165					170					175	
	Pro	Glu	Val	Pro	Ser	Tyr	Ser	Ala	Lys	Tyr	Gln	Leu	Thr	Asn	Asp	Asp
35																
				180					185					190		
	Tyr	Asn	Val	Lys	Gln	Leu	Arg	Lys	Arg	Tyr	Asp	Ile	Pro	Thr	Ser	Lys
			195					200					205			
40	Ala	Pro	Lys	Leu	Leu	Leu	Lys	Gly	Ser	Gly	Asn	Leu	Lys	Gly	Ser	Ser
		210					215					220				
	Val	Gly	Tyr	Lys	Asp	Ile	Glu	Phe	Thr	Phe	Val	Glu	Lys	Lys	Glu	Glu
	225					230					235					240
45	Asn	Ile	Tyr	Phe	Ser	Asp	Ser	Leu	Asp	Tyr	Lys	Lys	Ser	Gly	Asp	Val
					245					250					255	
50	<210>	41														
	<211>	256														
	<212>	PRT														
	<213>	Staphylococcus aureus														
55	<400>	41														

EP 2 493 498 B1

	Met	Met	Lys	Arg	Leu	Asn	Lys	Leu	Val	Leu	Gly	Ile	Ile	Phe	Leu	Phe	
	1				5					10					15		
5	Leu	Val	Ile	Ser	Ile	Thr	Ala	Gly	Cys	Gly	Ile	Gly	Lys	Glu	Ala	Glu	
				20					25					30			
	Val	Lys	Lys	Ser	Phe	Glu	Lys	Thr	Leu	Ser	Met	Tyr	Pro	Ile	Lys	Asn	
			35					40					45				
10	Leu	Glu	Asp	Leu	Tyr	Asp	Lys	Glu	Gly	Tyr	Arg	Asp	Asp	Gln	Phe	Asp	
		50					55					60					
	Lys	Asn	Asp	Lys	Gly	Thr	Trp	Ile	Ile	Asn	Ser	Glu	Met	Val	Ile	Gln	
	65					70					75					80	
15	Pro	Asn	Asn	Glu	Asp	Met	Val	Ala	Lys	Gly	Met	Val	Leu	Tyr	Met	Asn	
				85						90					95		
	Arg	Asn	Thr	Lys	Thr	Thr	Asn	Gly	Tyr	Tyr	Tyr	Val	Asp	Val	Thr	Lys	
20				100					105					110			
	Asp	Glu	Asp	Glu	Gly	Lys	Pro	His	Asp	Asn	Glu	Lys	Arg	Tyr	Pro	Val	
			115					120					125				
25	Lys	Met	Val	Asp	Asn	Lys	Ile	Ile	Pro	Thr	Lys	Glu	Ile	Lys	Asp	Glu	
		130					135					140					
	Lys	Val	Lys	Lys	Glu	Ile	Glu	Asn	Phe	Lys	Phe	Phe	Val	Gln	Tyr	Gly	
	145					150					155					160	
30	Asp	Phe	Lys	Asn	Ile	Lys	Asn	Tyr	Lys	Asp	Gly	Asp	Ile	Ser	Tyr	Asn	
				165						170					175		
	Pro	Glu	Val	Pro	Ser	Tyr	Ser	Ala	Lys	Tyr	Gln	Leu	Thr	Asn	Asp	Asp	
				180					185					190			
35	Tyr	Asn	Val	Lys	Gln	Leu	Arg	Lys	Arg	Tyr	Asp	Ile	Pro	Thr	Ser	Lys	
			195					200					205				
	Ala	Pro	Lys	Leu	Leu	Leu	Lys	Gly	Ser	Gly	Asn	Leu	Lys	Gly	Ser	Ser	
40		210					215					220					
	Val	Gly	Tyr	Lys	Asp	Ile	Glu	Phe	Thr	Phe	Val	Glu	Lys	Lys	Glu	Glu	
	225					230					235				240		
45	Asn	Ile	Tyr	Phe	Ser	Asp	Ser	Leu	Asp	Tyr	Lys	Lys	Ser	Gly	Asp	Val	
				245						250					255		

<210> 42

<211> 256

50 <212> PRT

<213> Staphylococcus aureus

<400> 42

55

EP 2 493 498 B1

	Met	Met	Lys	Arg	Leu	Asn	Lys	Leu	Val	Leu	Gly	Ile	Ile	Phe	Leu	Phe	
	1				5					10					15		
5	Leu	Val	Ile	Ser	Ile	Thr	Ala	Gly	Cys	Gly	Ile	Gly	Lys	Glu	Ala	Glu	
				20					25					30			
	Val	Lys	Lys	Ser	Phe	Glu	Lys	Thr	Leu	Ser	Met	Tyr	Pro	Ile	Lys	Asn	
			35					40					45				
10	Leu	Glu	Asp	Leu	Tyr	Asp	Lys	Glu	Gly	Tyr	Arg	Asp	Asp	Gln	Phe	Asp	
		50					55					60					
	Lys	Asn	Asp	Lys	Gly	Thr	Trp	Ile	Ile	Asn	Ser	Glu	Met	Val	Ile	Gln	
	65					70					75					80	
15	Pro	Asn	Asn	Glu	Asp	Met	Val	Ala	Lys	Gly	Met	Val	Leu	Tyr	Met	Asn	
				85						90					95		
	Arg	Asn	Thr	Lys	Thr	Thr	Asn	Gly	Tyr	Tyr	Tyr	Val	Asp	Val	Thr	Lys	
20				100					105					110			
	Asp	Glu	Asp	Glu	Gly	Lys	Pro	His	Asp	Asn	Glu	Lys	Arg	Tyr	Pro	Val	
			115					120					125				
	Lys	Met	Val	Asp	Asn	Lys	Ile	Ile	Pro	Thr	Lys	Glu	Ile	Lys	Asp	Glu	
25		130					135					140					
	Lys	Leu	Lys	Lys	Glu	Ile	Glu	Asn	Phe	Lys	Phe	Phe	Val	Gln	Tyr	Gly	
	145					150					155					160	
30	Asp	Phe	Lys	Asn	Val	Lys	Asn	Tyr	Lys	Asp	Gly	Asp	Ile	Ser	Tyr	Asn	
				165						170					175		
	Pro	Glu	Val	Pro	Ser	Tyr	Ser	Ala	Lys	Tyr	Gln	Leu	Thr	Asn	Asp	Asp	
				180					185					190			
35	Tyr	Asn	Val	Lys	Gln	Leu	Arg	Lys	Arg	Tyr	Asp	Ile	Pro	Thr	Ser	Lys	
			195					200					205				
	Ala	Pro	Lys	Leu	Leu	Leu	Lys	Gly	Ser	Gly	Asn	Leu	Lys	Gly	Ser	Ser	
40		210					215					220					
	Val	Gly	Tyr	Lys	Asp	Ile	Glu	Phe	Thr	Phe	Val	Glu	Lys	Lys	Glu	Glu	
	225					230					235					240	
	Asn	Ile	Tyr	Phe	Ser	Asp	Ser	Leu	Asp	Tyr	Lys	Lys	Ser	Gly	Asp	Val	
45				245						250					255		

<210> 43

<211> 350

<212> PRT

50 <213> Staphylococcus aureus

<400> 43

55

EP 2 493 498 B1

	Met	Thr	Lys	His	Tyr	Leu	Asn	Ser	Lys	Tyr	Gln	Ser	Glu	Gln	Arg	Ser	
	1				5					10					15		
5	Ser	Ala	Met	Lys	Lys	Ile	Thr	Met	Gly	Thr	Ala	Ser	Ile	Ile	Leu	Gly	
				20					25					30			
	Ser	Leu	Val	Tyr	Ile	Gly	Ala	Asp	Ser	Gln	Gln	Val	Asn	Ala	Ala	Thr	
			35					40					45				
10	Glu	Ala	Thr	Asn	Ala	Thr	Asn	Asn	Gln	Ser	Thr	Gln	Val	Ser	Gln	Ala	
		50					55					60					
	Thr	Ser	Gln	Pro	Ile	Asn	Phe	Gln	Val	Gln	Lys	Asp	Gly	Ser	Ser	Glu	
	65				70						75					80	
15	Lys	Ser	His	Met	Asp	Asp	Tyr	Met	Gln	His	Pro	Gly	Lys	Val	Ile	Lys	
				85						90					95		
	Gln	Asn	Asn	Lys	Tyr	Tyr	Phe	Gln	Thr	Val	Leu	Asn	Asn	Ala	Ser	Phe	
20				100					105					110			
	Trp	Lys	Glu	Tyr	Lys	Phe	Tyr	Asn	Ala	Asn	Asn	Gln	Glu	Leu	Ala	Thr	
			115					120					125				
25	Thr	Val	Val	Asn	Asp	Asn	Lys	Lys	Ala	Asp	Thr	Arg	Thr	Ile	Asn	Val	
		130					135					140					
	Ala	Val	Glu	Pro	Gly	Tyr	Lys	Ser	Leu	Thr	Thr	Lys	Val	His	Ile	Val	
	145					150					155					160	
30	Val	Pro	Gln	Ile	Asn	Tyr	Asn	His	Arg	Tyr	Thr	Thr	His	Leu	Glu	Phe	
				165						170					175		
	Glu	Lys	Ala	Ile	Pro	Thr	Leu	Ala	Asp	Ala	Ala	Lys	Pro	Asn	Asn	Val	
				180					185					190			
35	Lys	Pro	Val	Gln	Pro	Lys	Pro	Ala	Gln	Pro	Lys	Thr	Pro	Thr	Glu	Gln	
			195					200					205				
	Thr	Lys	Pro	Val	Gln	Pro	Lys	Val	Glu	Lys	Val	Lys	Pro	Thr	Val	Thr	
40		210					215					220					
	Thr	Thr	Ser	Lys	Val	Glu	Asp	Asn	His	Ser	Thr	Lys	Val	Val	Ser	Thr	
	225				230						235					240	
45	Asp	Thr	Thr	Lys	Asp	Gln	Thr	Lys	Thr	Gln	Thr	Ala	His	Thr	Val	Lys	
				245						250					255		
	Thr	Ala	Gln	Thr	Ala	Gln	Glu	Gln	Asn	Lys	Val	Gln	Thr	Pro	Val	Lys	
				260					265					270			
50	Asp	Val	Ala	Thr	Ala	Lys	Ser	Glu	Ser	Asn	Asn	Gln	Ala	Val	Ser	Asp	
			275					280					285				
	Asn	Lys	Ser	Gln	Gln	Thr	Asn	Lys	Val	Thr	Lys	His	Asn	Glu	Thr	Pro	

55

EP 2 493 498 B1

	290		295		300												
5	Lys	Gln	Ala	Ser	Lys	Ala	Lys	Glu	Leu	Pro	Lys	Thr	Gly	Leu	Thr	Ser	
	305					310					315					320	
	Val	Asp	Asn	Phe	Ile	Ser	Thr	Val	Ala	Phe	Ala	Thr	Leu	Ala	Leu	Leu	
					325					330					335		
10	Gly	Ser	Leu	Ser	Leu	Leu	Leu	Phe	Lys	Arg	Lys	Glu	Ser	Lys			
				340					345					350			
	<210> 44																
	<211> 145																
	<212> PRT																
15	<213> Staphylococcus aureus																
	<400> 44																
20	Asp	Ser	Gln	Gln	Val	Asn	Ala	Ala	Thr	Glu	Ala	Thr	Asn	Ala	Thr	Asn	
	1				5					10					15		
	Asn	Gln	Ser	Thr	Gln	Val	Ser	Gln	Ala	Thr	Ser	Gln	Pro	Ile	Asn	Phe	
				20					25					30			
25	Gln	Val	Gln	Lys	Asp	Gly	Ser	Ser	Glu	Lys	Ser	His	Met	Asp	Asp	Tyr	
			35					40					45				
	Met	Gln	His	Pro	Gly	Lys	Val	Ile	Lys	Gln	Asn	Asn	Lys	Tyr	Tyr	Phe	
	50					55						60					
30	Gln	Thr	Val	Leu	Asn	Asn	Ala	Ser	Phe	Trp	Lys	Glu	Tyr	Lys	Phe	Tyr	
	65				70						75					80	
	Asn	Ala	Asn	Asn	Gln	Glu	Leu	Ala	Thr	Thr	Val	Val	Asn	Asp	Asn	Lys	
35					85					90					95		
	Lys	Ala	Asp	Thr	Arg	Thr	Ile	Asn	Val	Ala	Val	Glu	Pro	Gly	Tyr	Lys	
				100					105					110			
40	Ser	Leu	Thr	Thr	Lys	Val	His	Ile	Val	Val	Pro	Gln	Ile	Asn	Tyr	Asn	
			115					120					125				
	His	Arg	Tyr	Thr	Thr	His	Leu	Glu	Phe	Glu	Lys	Ala	Ile	Pro	Thr	Leu	
	130						135					140					
45	Ala																
	145																
	<210> 45																
	<211> 645																
50	<212> PRT																
	<213> Staphylococcus aureus																
	<400> 45																
55																	

EP 2 493 498 B1

Met Asn Lys Gln Gln Lys Glu Phe Lys Ser Phe Tyr Ser Ile Arg Lys
1 5 10 15

5 Ser Ser Leu Gly Val Ala Ser Val Ala Ile Ser Thr Leu Leu Leu Leu
20 25 30

Met Ser Asn Gly Glu Ala Gln Ala Ala Glu Glu Thr Gly Gly Thr
35 40 45

10

15

20

25

30

35

40

45

50

55

EP 2 493 498 B1

	Asn	Thr	Glu	Ala	Gln	Pro	Lys	Thr	Glu	Ala	Val	Ala	Ser	Pro	Thr	Thr	
	50						55					60					
5	Thr	Ser	Glu	Lys	Ala	Pro	Glu	Thr	Lys	Pro	Val	Ala	Asn	Ala	Val	Ser	
	65					70					75					80	
	Val	Ser	Asn	Lys	Glu	Val	Glu	Ala	Pro	Thr	Ser	Glu	Thr	Lys	Glu	Ala	
				85						90					95		
10	Lys	Glu	Val	Lys	Glu	Val	Lys	Ala	Pro	Lys	Glu	Thr	Lys	Glu	Val	Lys	
				100					105					110			
	Pro	Ala	Ala	Lys	Ala	Thr	Asn	Asn	Thr	Tyr	Pro	Ile	Leu	Asn	Gln	Glu	
			115					120					125				
15	Leu	Arg	Glu	Ala	Ile	Lys	Asn	Pro	Ala	Ile	Lys	Asp	Lys	Asp	His	Ser	
	130						135					140					
	Ala	Pro	Asn	Ser	Arg	Pro	Ile	Asp	Phe	Glu	Met	Lys	Lys	Lys	Asp	Gly	
20	145					150					155					160	
	Thr	Gln	Gln	Phe	Tyr	His	Tyr	Ala	Ser	Ser	Val	Lys	Pro	Ala	Arg	Val	
				165						170					175		
	Ile	Phe	Thr	Asp	Ser	Lys	Pro	Glu	Ile	Glu	Leu	Gly	Leu	Gln	Ser	Gly	
25				180					185					190			
	Gln	Phe	Trp	Arg	Lys	Phe	Glu	Val	Tyr	Glu	Gly	Asp	Lys	Lys	Leu	Pro	
			195					200					205				
30	Ile	Lys	Leu	Val	Ser	Tyr	Asp	Thr	Val	Lys	Asp	Tyr	Ala	Tyr	Ile	Arg	
	210						215					220					
	Phe	Ser	Val	Ser	Asn	Gly	Thr	Lys	Ala	Val	Lys	Ile	Val	Ser	Ser	Thr	
	225					230					235					240	
35	His	Phe	Asn	Asn	Lys	Glu	Glu	Lys	Tyr	Asp	Tyr	Thr	Leu	Met	Glu	Phe	
					245					250					255		
	Ala	Gln	Pro	Ile	Tyr	Asn	Ser	Ala	Asp	Lys	Phe	Lys	Thr	Glu	Glu	Asp	
40				260					265					270			
	Tyr	Lys	Ala	Glu	Lys	Leu	Leu	Ala	Pro	Tyr	Lys	Lys	Ala	Lys	Thr	Leu	
			275					280					285				
	Glu	Arg	Gln	Val	Tyr	Glu	Leu	Asn	Lys	Ile	Gln	Asp	Lys	Leu	Pro	Glu	
45							295					300					
	Lys	Leu	Lys	Ala	Glu	Tyr	Lys	Lys	Lys	Leu	Glu	Asp	Thr	Lys	Lys	Ala	
	305					310					315					320	
50	Leu	Asp	Glu	Gln	Val	Lys	Ser	Ala	Ile	Thr	Glu	Phe	Gln	Asn	Val	Gln	
					325					330					335		
	Pro	Thr	Asn	Glu	Lys	Met	Thr	Asp	Leu	Gln	Asp	Thr	Lys	Tyr	Val	Val	
				340					345					350			
55	Tyr	Glu	Ser	Val	Glu	Asn	Asn	Glu	Ser	Met	Met	Asp	Thr	Phe	Val	Lys	
			355					360					365				

EP 2 493 498 B1

	His	Pro	Ile	Lys	Thr	Gly	Met	Leu	Asn	Gly	Lys	Lys	Tyr	Met	Val	Met	
	370						375						380				
5	Glu	Thr	Thr	Asn	Asp	Asp	Tyr	Trp	Lys	Asp	Phe	Met	Val	Glu	Gly	Gln	
	385				390					395						400	
	Arg	Val	Arg	Thr	Ile	Ser	Lys	Asp	Ala	Lys	Asn	Asn	Thr	Arg	Thr	Ile	
				405						410					415		
10	Ile	Phe	Pro	Tyr	Val	Glu	Gly	Lys	Thr	Leu	Tyr	Asp	Ala	Ile	Val	Lys	
			420					425						430			
	Val	His	Val	Lys	Thr	Ile	Asp	Tyr	Asp	Gly	Gln	Tyr	His	Val	Arg	Ile	
			435				440						445				
15	Val	Asp	Lys	Glu	Ala	Phe	Thr	Lys	Ala	Asn	Thr	Asp	Lys	Ser	Asn	Lys	
	450						455					460					
	Lys	Glu	Gln	Gln	Asp	Asn	Ser	Ala	Lys	Lys	Glu	Ala	Thr	Pro	Ala	Thr	
20	465				470						475					480	
	Pro	Ser	Lys	Pro	Thr	Pro	Ser	Pro	Val	Glu	Lys	Glu	Ser	Gln	Lys	Gln	
					485					490					495		
25	Asp	Ser	Gln	Lys	Asp	Asp	Asn	Lys	Gln	Leu	Pro	Ser	Val	Glu	Lys	Glu	
				500				505						510			
	Asn	Asp	Ala	Ser	Ser	Glu	Ser	Gly	Lys	Asp	Lys	Thr	Pro	Ala	Thr	Lys	
			515					520					525				
30	Pro	Thr	Lys	Gly	Glu	Val	Glu	Ser	Ser	Ser	Thr	Thr	Pro	Thr	Lys	Val	
	530					535						540					
	Val	Ser	Thr	Thr	Gln	Asn	Val	Ala	Lys	Pro	Thr	Thr	Ala	Ser	Ser	Lys	
	545				550						555					560	
35	Thr	Thr	Lys	Asp	Val	Val	Gln	Thr	Ser	Ala	Gly	Ser	Ser	Glu	Ala	Lys	
				565						570					575		
	Asp	Ser	Ala	Pro	Leu	Gln	Lys	Ala	Asn	Ile	Lys	Asn	Thr	Asn	Asp	Gly	
40				580				585						590			
	His	Thr	Gln	Ser	Gln	Asn	Asn	Lys	Asn	Thr	Gln	Glu	Asn	Lys	Ala	Lys	
			595					600					605				
45	Ser	Leu	Pro	Gln	Thr	Gly	Glu	Glu	Ser	Asn	Lys	Asp	Met	Thr	Leu	Pro	
	610					615						620					
	Leu	Met	Ala	Leu	Leu	Ala	Leu	Ser	Ser	Ile	Val	Ala	Phe	Val	Leu	Pro	
	625				630						635					640	
50	Arg	Lys	Arg	Lys	Asn												
					645												

<210> 46

<211> 1256

55 <212> PRT

<213> Staphylococcus aureus

<400> 46

EP 2 493 498 B1

Met Ala Lys Lys Phe Asn Tyr Lys Leu Pro Ser Met Val Ala Leu Thr

5

10

15

20

25

30

35

40

45

50

55

EP 2 493 498 B1

	1			5					10					15		
	Leu	Val	Gly	Ser	Ala	Val	Thr	Ala	His	Gln	Val	Gln	Ala	Ala	Glu	Thr
				20					25					30		
5	Thr	Gln	Asp	Gln	Thr	Thr	Asn	Lys	Asn	Val	Leu	Asp	Ser	Asn	Lys	Val
			35					40					45			
	Lys	Ala	Thr	Thr	Glu	Gln	Ala	Lys	Ala	Glu	Val	Lys	Asn	Pro	Thr	Gln
10		50					55					60				
	Asn	Ile	Ser	Gly	Thr	Gln	Val	Tyr	Gln	Asp	Pro	Ala	Ile	Val	Gln	Pro
	65					70					75					80
	Lys	Thr	Ala	Asn	Asn	Lys	Thr	Gly	Asn	Ala	Gln	Val	Ser	Gln	Lys	Val
15					85					90					95	
	Asp	Thr	Ala	Gln	Val	Asn	Gly	Asp	Thr	Arg	Ala	Asn	Gln	Ser	Ala	Thr
				100					105					110		
	Thr	Asn	Asn	Thr	Gln	Pro	Val	Ala	Lys	Ser	Thr	Ser	Thr	Thr	Ala	Pro
20				115					120					125		
	Lys	Thr	Asn	Thr	Asn	Val	Thr	Asn	Ala	Gly	Tyr	Ser	Leu	Val	Asp	Asp
		130					135					140				
25	Glu	Asp	Asp	Asn	Ser	Glu	Asn	Gln	Ile	Asn	Pro	Glu	Leu	Ile	Lys	Ser
	145					150					155					160
	Ala	Ala	Lys	Pro	Ala	Ala	Leu	Glu	Thr	Gln	Tyr	Lys	Thr	Ala	Ala	Pro
					165					170					175	
30	Lys	Ala	Ala	Thr	Thr	Ser	Ala	Pro	Lys	Ala	Lys	Thr	Glu	Ala	Thr	Pro
				180					185					190		
	Lys	Val	Thr	Thr	Phe	Ser	Ala	Ser	Ala	Gln	Pro	Arg	Ser	Val	Ala	Ala
35			195					200					205			
	Thr	Pro	Lys	Thr	Ser	Leu	Pro	Lys	Tyr	Lys	Pro	Gln	Val	Asn	Ser	Ser
		210					215					220				
	Ile	Asn	Asp	Tyr	Ile	Cys	Lys	Asn	Asn	Leu	Lys	Ala	Pro	Lys	Ile	Glu
40	225					230					235					240
	Glu	Asp	Tyr	Thr	Ser	Tyr	Phe	Pro	Lys	Tyr	Ala	Tyr	Arg	Asn	Gly	Val
					245					250					255	
	Gly	Arg	Pro	Glu	Gly	Ile	Val	Val	His	Asp	Thr	Ala	Asn	Asp	Arg	Ser
45				260					265					270		
	Thr	Ile	Asn	Gly	Glu	Ile	Ser	Tyr	Met	Lys	Asn	Asn	Tyr	Gln	Asn	Ala
			275					280					285			
	Phe	Val	His	Ala	Phe	Val	Asp	Gly	Asp	Arg	Ile	Ile	Glu	Thr	Ala	Pro
50		290					295					300				
	Thr	Asp	Tyr	Leu	Ser	Trp	Gly	Val	Gly	Ala	Val	Gly	Asn	Pro	Arg	Phe
	305					310					315					320
55	Ile	Asn	Val	Glu	Ile	Val	His	Thr	His	Asp	Tyr	Ala	Ser	Phe	Ala	Arg
				325						330					335	

EP 2 493 498 B1

	Ser	Met	Asn	Asn	Tyr	Ala	Asp	Tyr	Ala	Ala	Thr	Gln	Leu	Gln	Tyr	Tyr	
				340					345					350			
5	Gly	Leu	Lys	Pro	Asp	Ser	Ala	Glu	Tyr	Asp	Gly	Asn	Gly	Thr	Val	Trp	
			355					360					365				
	Thr	His	Tyr	Ala	Val	Ser	Lys	Tyr	Leu	Gly	Gly	Thr	Asp	His	Ala	Asp	
		370					375					380					
10	Pro	His	Gly	Tyr	Leu	Arg	Ser	His	Asn	Tyr	Ser	Tyr	Asp	Gln	Leu	Tyr	
		385				390					395					400	
	Asp	Leu	Ile	Asn	Glu	Lys	Tyr	Leu	Ile	Lys	Met	Gly	Lys	Val	Ala	Pro	
					405					410					415		
15	Trp	Gly	Thr	Gln	Ser	Thr	Thr	Thr	Pro	Thr	Thr	Pro	Ser	Lys	Pro	Thr	
				420					425					430			
	Thr	Pro	Ser	Lys	Pro	Ser	Thr	Gly	Lys	Leu	Thr	Val	Ala	Ala	Asn	Asn	
20			435					440					445				
	Gly	Val	Ala	Gln	Ile	Lys	Pro	Thr	Asn	Ser	Gly	Leu	Tyr	Thr	Thr	Val	
		450					455					460					
25	Tyr	Asp	Lys	Thr	Gly	Lys	Ala	Thr	Asn	Glu	Val	Gln	Lys	Thr	Phe	Ala	
		465				470					475					480	
	Val	Ser	Lys	Thr	Ala	Thr	Leu	Gly	Asn	Gln	Lys	Phe	Tyr	Leu	Val	Gln	
					485					490					495		
30	Asp	Tyr	Asn	Ser	Gly	Asn	Lys	Phe	Gly	Trp	Val	Lys	Glu	Gly	Asp	Val	
				500					505					510			
	Val	Tyr	Asn	Thr	Ala	Lys	Ser	Pro	Val	Asn	Val	Asn	Gln	Ser	Tyr	Ser	
			515					520					525				
35	Ile	Lys	Pro	Gly	Thr	Lys	Leu	Tyr	Thr	Val	Pro	Trp	Gly	Thr	Ser	Lys	
		530					535					540					
	Gln	Val	Ala	Gly	Ser	Val	Ser	Gly	Ser	Gly	Asn	Gln	Thr	Phe	Lys	Ala	
40		545				550					555					560	
	Ser	Lys	Gln	Gln	Gln	Ile	Asp	Lys	Ser	Ile	Tyr	Leu	Tyr	Gly	Ser	Val	
					565					570					575		
45	Asn	Gly	Lys	Ser	Gly	Trp	Val	Ser	Lys	Ala	Tyr	Leu	Val	Asp	Thr	Ala	
				580					585					590			
	Lys	Pro	Thr	Pro	Thr	Pro	Thr	Pro	Lys	Pro	Ser	Thr	Pro	Thr	Thr	Asn	
			595					600					605				
50	Asn	Lys	Leu	Thr	Val	Ser	Ser	Leu	Asn	Gly	Val	Ala	Gln	Ile	Asn	Ala	
		610					615					620					
	Lys	Asn	Asn	Gly	Leu	Phe	Thr	Thr	Val	Tyr	Asp	Lys	Thr	Gly	Lys	Pro	
		625				630					635					640	
55	Thr	Lys	Glu	Val	Gln	Lys	Thr	Phe	Ala	Val	Thr	Lys	Glu	Ala	Ser	Leu	
					645					650					655		

EP 2 493 498 B1

	Gly	Gly	Asn	Lys	Phe	Tyr	Leu	Val	Lys	Asp	Tyr	Asn	Ser	Pro	Thr	Leu	
				660					665					670			
5	Ile	Gly	Trp	Val	Lys	Gln	Gly	Asp	Val	Ile	Tyr	Asn	Asn	Ala	Lys	Ser	
			675					680					685				
	Pro	Val	Asn	Val	Met	Gln	Thr	Tyr	Thr	Val	Lys	Pro	Gly	Thr	Lys	Leu	
		690					695					700					
10	Tyr	Ser	Val	Pro	Trp	Gly	Thr	Tyr	Lys	Gln	Glu	Ala	Gly	Ala	Val	Ser	
	705					710					715					720	
	Gly	Thr	Gly	Asn	Gln	Thr	Phe	Lys	Ala	Thr	Lys	Gln	Gln	Gln	Ile	Asp	
				725						730					735		
15	Lys	Ser	Ile	Tyr	Leu	Phe	Gly	Thr	Val	Asn	Gly	Lys	Ser	Gly	Trp	Val	
				740					745					750			
	Ser	Lys	Ala	Tyr	Leu	Ala	Val	Pro	Ala	Ala	Pro	Lys	Lys	Ala	Val	Ala	
			755					760					765				
20	Gln	Pro	Lys	Thr	Ala	Val	Lys	Ala	Tyr	Thr	Val	Thr	Lys	Pro	Gln	Thr	
		770					775					780					
	Thr	Gln	Thr	Val	Ser	Lys	Ile	Ala	Gln	Val	Lys	Pro	Asn	Asn	Thr	Gly	
25	785					790					795					800	
	Ile	Arg	Ala	Ser	Val	Tyr	Glu	Lys	Thr	Ala	Lys	Asn	Gly	Ala	Lys	Tyr	
				805						810					815		
	Ala	Asp	Arg	Thr	Phe	Tyr	Val	Thr	Lys	Glu	Arg	Ala	His	Gly	Asn	Glu	
30				820					825					830			
	Thr	Tyr	Val	Leu	Leu	Asn	Asn	Thr	Ser	His	Asn	Ile	Pro	Leu	Gly	Trp	
			835					840					845				
	Phe	Asn	Val	Lys	Asp	Leu	Asn	Val	Gln	Asn	Leu	Gly	Lys	Glu	Val	Lys	
35		850					855					860					
	Thr	Thr	Gln	Lys	Tyr	Thr	Val	Asn	Lys	Ser	Asn	Asn	Gly	Leu	Ser	Met	
	865					870					875					880	
40	Val	Pro	Trp	Gly	Thr	Lys	Asn	Gln	Val	Ile	Leu	Thr	Gly	Asn	Asn	Ile	
				885						890					895		
	Ala	Gln	Gly	Thr	Phe	Asn	Ala	Thr	Lys	Gln	Val	Ser	Val	Gly	Lys	Asp	
				900					905					910			
45	Val	Tyr	Leu	Tyr	Gly	Thr	Ile	Asn	Asn	Arg	Thr	Gly	Trp	Val	Asn	Ala	
			915					920					925				
	Lys	Asp	Leu	Thr	Ala	Pro	Thr	Ala	Val	Lys	Pro	Thr	Thr	Ser	Ala	Ala	
		930					935					940					
50	Lys	Asp	Tyr	Asn	Tyr	Thr	Tyr	Val	Ile	Lys	Asn	Gly	Asn	Gly	Tyr	Tyr	
	945					950					955					960	
	Tyr	Val	Thr	Pro	Asn	Ser	Asp	Thr	Ala	Lys	Tyr	Ser	Leu	Lys	Ala	Phe	
55				965						970					975		
	Asn	Glu	Gln	Pro	Phe	Ala	Val	Val	Lys	Glu	Gln	Val	Ile	Asn	Gly	Gln	

EP 2 493 498 B1

	980	985	990
5	Thr Trp Tyr Tyr Gly Lys Leu Ser Asn Gly Lys Leu Ala Trp Ile Lys 995 1000 1005		
	Ser Thr Asp Leu Ala Lys Glu Leu Ile Lys Tyr Asn Gln Thr Gly Met 1010 1015 1020		
10	Thr Leu Asn Gln Val Ala Gln Ile Gln Ala Gly Leu Gln Tyr Lys Pro 1025 1030 1035 1040		
	Gln Val Gln Arg Val Pro Gly Lys Trp Thr Asp Ala Lys Phe Asn Asp 1045 1050 1055		
15	Val Lys His Ala Met Asp Thr Lys Arg Leu Ala Gln Asp Pro Ala Leu 1060 1065 1070		
	Lys Tyr Gln Phe Leu Arg Leu Asp Gln Pro Gln Asn Ile Ser Ile Asp 1075 1080 1085		
20	Lys Ile Asn Gln Phe Leu Lys Gly Lys Gly Val Leu Glu Asn Gln Gly 1090 1095 1100		
	Ala Ala Phe Asn Lys Ala Ala Gln Met Tyr Gly Ile Asn Glu Val Tyr 1105 1110 1115 1120		
25	Leu Ile Ser His Ala Leu Leu Glu Thr Gly Asn Gly Thr Ser Gln Leu 1125 1130 1135		
	Ala Lys Gly Ala Asp Val Val Asn Asn Lys Val Val Thr Asn Ser Asn 1140 1145 1150		
30	Thr Lys Tyr His Asn Val Phe Gly Ile Ala Ala Tyr Asp Asn Asp Pro 1155 1160 1165		
	Leu Arg Glu Gly Ile Lys Tyr Ala Lys Gln Ala Gly Trp Asp Thr Val 1170 1175 1180		
	Ser Lys Ala Ile Val Gly Gly Ala Lys Phe Ile Gly Asn Ser Tyr Val 1185 1190 1195 1200		
40	Lys Ala Gly Gln Asn Thr Leu Tyr Lys Met Arg Trp Asn Pro Ala His 1205 1210 1215		
	Pro Gly Thr His Gln Tyr Ala Thr Asp Val Asp Trp Ala Asn Ile Asn 1220 1225 1230		
45	Ala Lys Ile Ile Lys Gly Tyr Tyr Asp Lys Ile Gly Glu Val Gly Lys 1235 1240 1245		
	Tyr Phe Asp Ile Pro Gln Tyr Lys 1250 1255		

<210> 47

<211> 6

<212> PRT

55 <213> Artificial Sequence

<220>

<223> linker

EP 2 493 498 B1

<400> 47

5 Gly Ser Gly Gly Gly Gly
1 5

<210> 48

<211> 8

<212> PRT

10 <213> Artificial Sequence

<220>

<223> linker

15 <400> 48

20 Gly Ser Gly Ser Gly Gly Gly Gly
1 5

<210> 49

<211> 6

<212> PRT

25 <213> Artificial Sequence

<220>

<223> linker

30 <400> 49

35 Ala Ser Gly Gly Gly Ser
1 5

<210> 50

<211> 18

<212> DNA

<213> Artificial Sequence

40 <220>

<223> Sequence encoding SEQ ID NO: 49

<400> 50

45 gctagcgggtg gcggatcc 18

<210> 51

<211> 6

<212> PRT

50 <213> Artificial Sequence

<220>

<223> hexahistidine tag

55 <400> 51

His His His His His His
1 5

EP 2 493 498 B1

<210> 52
 <211> 207
 <212> PRT
 <213> Staphylococcus aureus

5

<400> 52

10

Met Ala Met Ile Lys Met Ser Pro Glu Glu Ile Arg Ala Lys Ser Gln
 1 5 10 15

Ser Tyr Gly Gln Gly Ser Asp Gln Ile Arg Gln Ile Leu Ser Asp Leu

15

20

25

30

Thr Arg Ala Gln Gly Glu Ile Ala Ala Asn Trp Glu Gly Gln Ala Phe
 35 40 45

20

Ser Arg Phe Glu Glu Gln Phe Gln Gln Leu Ser Pro Lys Val Glu Lys
 50 55 60

Phe Ala Gln Leu Leu Glu Glu Ile Lys Gln Gln Leu Asn Ser Thr Ala
 65 70 75 80

25

Asp Ala Val Gln Glu Gln Asp Gln Gln Leu Ser Asn Asn Phe Gly Leu
 85 90 95

Gln Ala Ser Gly Gly Gly Ser Met Gly Gly Tyr Lys Gly Ile Lys Ala
 100 105 110

30

Asp Gly Gly Lys Val Asp Gln Ala Lys Gln Leu Ala Ala Lys Thr Ala
 115 120 125

35

Lys Asp Ile Glu Ala Cys Gln Lys Gln Thr Gln Gln Leu Ala Glu Tyr
 130 135 140

Ile Glu Gly Ser Asp Trp Glu Gly Gln Phe Ala Asn Lys Val Lys Asp
 145 150 155 160

40

Val Leu Leu Ile Met Ala Lys Phe Gln Glu Glu Leu Val Gln Pro Met
 165 170 175

Ala Asp His Gln Lys Ala Ile Asp Asn Leu Ser Gln Asn Leu Ala Lys
 180 185 190

45

Tyr Asp Thr Leu Ser Ile Lys Gln Gly Leu Asp Arg Val Asn Pro
 195 200 205

<210> 53
 <211> 618
 <212> DNA
 <213> Staphylococcus aureus

50

<400> 53

55

EP 2 493 498 B1

```

atggcaatga ttaagatgag tccagaggaa atcagagcaa aatcgcaatc ttacgggcaa 60
ggttcagacc aaatccgtca aattttatct gatttaacac gtgcacaagg tgaaattgca 120
gcgaactggg aaggtcaagc tttcagccgt ttcgaagagc aattccaaca acttagtcct 180
aaagtagaaa aatttgcaca attattagaa gaaattaaac aacaattgaa tagcactgct 240
5 gatgccgttc aagaacaaga ccaacaactt tctaataatt tcggtttgca agctagcggg 300
ggcggatccg gtggatataa aggtattaaa gcagatgggtg gcaagggttg tcaagcgaaa 360
caattagcgg caaaaacagc taaagatatt gaagcatgtc aaaagcaaac gcaacagctc 420
gctgagtata tcgaaggtag tgattgggaa ggacagttcg ccaataagggt gaaagatgtg 480
ttactcatta tggcaaagtt tcaagaagaa ttagtacaac cgatgggtga ccatcaaaaa 540
10 gcaattgata acttaagtca aaatctagcg aaatacgata cattatcaat taagcaaggg 600
cttgataggg tgaaccca 618

```

<210> 54

<211> 207

15 <212> PRT

<213> Staphylococcus aureus

<400> 54

```

20 Met Gly Gly Tyr Lys Gly Ile Lys Ala Asp Gly Gly Lys Val Asp Gln
   1           5           10           15

25 Ala Lys Gln Leu Ala Ala Lys Thr Ala Lys Asp Ile Glu Ala Cys Gln
   20           25           30

30 Lys Gln Thr Gln Gln Leu Ala Glu Tyr Ile Glu Gly Ser Asp Trp Glu
   35           40           45

35 Gly Gln Phe Ala Asn Lys Val Lys Asp Val Leu Leu Ile Met Ala Lys
   50           55           60

   Phe Gln Glu Glu Leu Val Gln Pro Met Ala Asp His Gln Lys Ala Ile
   65           70           75           80

   Asp Asn Leu Ser Gln Asn Leu Ala Lys Tyr Asp Thr Leu Ser Ile Lys
   85           90           95

40 Gln Gly Leu Asp Arg Val Asn Pro Ala Ser Gly Gly Gly Ser Met Ala
   100          105          110

   Met Ile Lys Met Ser Pro Glu Glu Ile Arg Ala Lys Ser Gln Ser Tyr
   115          120          125

45 Gly Gln Gly Ser Asp Gln Ile Arg Gln Ile Leu Ser Asp Leu Thr Arg
   130          135          140

   Ala Gln Gly Glu Ile Ala Ala Asn Trp Glu Gly Gln Ala Phe Ser Arg
   145          150          155          160

50 Phe Glu Glu Gln Phe Gln Gln Leu Ser Pro Lys Val Glu Lys Phe Ala
   165          170          175

   Gln Leu Leu Glu Glu Ile Lys Gln Gln Leu Asn Ser Thr Ala Asp Ala
   180          185          190

   Val Gln Glu Gln Asp Gln Gln Leu Ser Asn Asn Phe Gly Leu Gln
   195          200          205

```

Claims

1. A method for making an immunogenic composition comprising an *S. aureus* type 5 capsular polysaccharide conjugated to a carrier molecule and an *S. aureus* type 8 capsular polysaccharide conjugated to a carrier molecule, said method comprising:
 - (a) a process for purifying capsular polysaccharide from *S. aureus* type 5 cells comprising a step of releasing the capsular polysaccharide from the cells by acid treatment of the cells, wherein the cells are in the form of a wet cell paste or are suspended in an aqueous medium, and the molecular mass of the purified polysaccharide is between 2-3500 kDa;
 - (b) conjugation of the capsular polysaccharide to a carrier molecule to make a conjugate; and
 - (c) mixing the conjugate with an *S. aureus* type 8 capsular polysaccharide conjugated to a carrier molecule.
2. A method for making an immunogenic composition comprising an *S. aureus* type 8 capsular polysaccharide conjugated to a carrier molecule and an *S. aureus* type 5 capsular polysaccharide conjugated to a carrier molecule, said method comprising:
 - (a) a process for purifying capsular polysaccharide from *S. aureus* type 8 cells comprising a step of releasing the capsular polysaccharide from the cells by acid treatment of the cells, wherein the cells are in the form of a wet cell paste or are suspended in an aqueous medium, and the molecular mass of the purified polysaccharide is between 2-3500 kDa;
 - (b) conjugation of the capsular polysaccharide to a carrier molecule to make a conjugate; and
 - (c) mixing the conjugate with an *S. aureus* type 5 capsular polysaccharide conjugated to a carrier molecule.
3. The method of claim 1 or claim 2, wherein (i) the acid treatment is carried out using acetic acid and/or (ii) the method further comprises a step of neutralisation.
4. The method according to any preceding claim, wherein the method further comprises a step of centrifugation of the cells and collection of the polysaccharide-containing supernatant.
5. The method according to any one of claims 1-4, wherein the process in step (a) further comprises a step of treatment of the capsular polysaccharide with DNase and/or RNase.
6. The method according to any one of claims 1-5, wherein the process in step (a) further comprises a step of treatment of the capsular polysaccharide with mutanolysin.
7. The method according to any one of claims 1-6, wherein the process in step (a) further comprises:
 - (a) a step of diafiltration;
 - (b) a step of anion exchange chromatography;
 - (c) a step of gel filtration; and/or
 - (d) a step of concentration of the polysaccharide.
8. The method according to any one of claims 1-7, wherein the process in step (a) further comprises:
 - (a) a step of depolymerisation of the purified polysaccharide to form an oligosaccharide; and/or
 - (b) a step of sterile filtration.
9. The method according to any one of claims 1-8, wherein the process in step (a) provides a composition comprising:
 - (a) the polysaccharide and a level of peptidoglycan contamination that is less than 5% by weight peptidoglycan relative to the total weight of the polysaccharide; and/or
 - (b) the polysaccharide and a level of protein contamination that is less than 5% by weight protein relative to the total weight of the polysaccharide.
10. The method according to any one of claims 1-9, wherein the process in step (a) provides a composition comprising the polysaccharide and a level of nucleic acid contamination that is less than 1% by weight nucleic acid relative to the total weight of the polysaccharide.

11. The method according to any one of claims of claims 1-10 further comprising including in the composition one or more *S. aureus* protein antigen(s) selected from the group consisting of a clfA antigen; a clfB antigen; a sdrE2 antigen; a sdrC antigen; a sasF antigen; a emp antigen; a sdrD antigen; a spa antigen; a esaC antigen; a esxA antigen; a esxB antigen; a sta006 antigen; a isdC antigen; a Hla antigen; a sta011 antigen; a isdA antigen; a isdB antigen; and a sta073 antigen.

Patentansprüche

1. Verfahren zum Herstellen einer immunogenen Zusammensetzung, umfassend ein *S. aureus* Typ 5-Kapselpolysaccharid, das an ein Trägermolekül konjugiert ist, und ein *S. aureus* Typ 8-Kapselpolysaccharid, das an ein Trägermolekül konjugiert ist, wobei das Verfahren umfasst:

- (a) einen Prozess zum Aufreinigen von Kapselpolysaccharid von *S. aureus* Typ 5-Zellen, umfassend einen Schritt des Freisetzens des Kapselpolysaccharids aus den Zellen durch Säurebehandlung der Zellen, wobei die Zellen in der Form einer nassen Zellpaste sind oder in einem wässrigen Medium suspendiert sind, und die molekulare Masse des aufgereinigten Polysaccharids zwischen 2-3500 kDa ist;
- (b) Konjugation des Kapselpolysaccharids an ein Trägermolekül, um ein Konjugat zu erzeugen; und
- (c) Mischen des Konjugats mit einem *S. aureus* Typ 8-Kapselpolysaccharid, das an ein Trägermolekül konjugiert ist.

2. Verfahren zum Herstellen einer immunogenen Zusammensetzung, umfassend ein *S. aureus* Typ 8-Kapselpolysaccharid, das an ein Trägermolekül konjugiert ist, und ein *S. aureus* Typ 5-Kapselpolysaccharid, das an ein Trägermolekül konjugiert ist, wobei das Verfahren umfasst:

- (a) einen Prozess zum Aufreinigen von Kapselpolysaccharid von *S. aureus* Typ 8-Zellen, umfassend einen Schritt des Freisetzens des Kapselpolysaccharids aus den Zellen durch Säurebehandlung der Zellen, wobei die Zellen in der Form einer nassen Zellpaste sind oder in einem wässrigen Medium suspendiert sind, und die molekulare Masse des aufgereinigten Polysaccharids zwischen 2-3500 kDa ist;
- (b) Konjugation des Kapselpolysaccharids an ein Trägermolekül, um ein Konjugat zu erzeugen; und
- (c) Mischen des Konjugats mit einem *S. aureus* Typ 5-Kapselpolysaccharid, das an ein Trägermolekül konjugiert ist.

3. Verfahren gemäß Anspruch 1 oder Anspruch 2, wobei (i) die Säurebehandlung unter Verwendung von Essigsäure durchgeführt wird, und/oder (ii) das Verfahren weiterhin einen Schritt der Neutralisation umfasst.

4. Verfahren gemäß irgendeinem der vorangegangenen Ansprüche, wobei das Verfahren weiterhin einen Schritt der Zentrifugation der Zellen und des Gewinnens des Polysaccharid-enthaltenden Überstands umfasst.

5. Verfahren gemäß irgendeinem der Ansprüche 1-4, wobei der Prozess in Schritt (a) weiterhin einen Schritt der Behandlung des Kapselpolysaccharids mit DNase und/oder RNase umfasst.

6. Verfahren gemäß irgendeinem der Ansprüche 1-5, wobei der Prozess in Schritt (a) weiterhin einen Schritt der Behandlung des Kapselpolysaccharids mit Mutanolysin umfasst.

7. Verfahren gemäß irgendeinem der Ansprüche 1-6, wobei der Prozess in Schritt (a) weiterhin umfasst:

- (a) einen Schritt der Diafiltration;
- (b) einen Schritt der Anionenaustauschchromatographie;
- (c) einen Schritt der Gelfiltration; und/oder
- (d) einen Schritt der Konzentration des Polysaccharids.

8. Verfahren gemäß irgendeinem der Ansprüche 1-7, wobei der Prozess in Schritt (a) weiterhin umfasst:

- (a) einen Schritt der Depolymerisation des aufgereinigten Polysaccharids, um ein Oligosaccharid zu bilden; und/oder
- (b) einen Schritt der Sterilfiltration.

9. Verfahren gemäß irgendeinem der Ansprüche 1-8, wobei der Prozess in Schritt (a) eine Zusammensetzung bereitstellt, umfassend:

- (a) das Polysaccharid und einen Gehalt an Peptidoglycankontamination, der weniger als 5 Gewichts-% Peptidoglycan relativ zu dem Gesamtgewicht des Polysaccharids beträgt; und/oder
- (b) das Polysaccharid und einen Gehalt an Proteinkontamination, der weniger als 5 Gewichts-% Protein relativ zu dem Gesamtgewicht des Polysaccharids beträgt.

10. Verfahren gemäß irgendeinem der Ansprüche 1-9, wobei der Prozess in Schritt (a) eine Zusammensetzung bereitstellt, umfassend das Polysaccharid und einen Gehalt an Nukleinsäurekontamination, der weniger als 1 Gewichts-% Nukleinsäure relativ zu dem Gesamtgewicht des Polysaccharids beträgt.

11. Verfahren gemäß irgendeinem der Ansprüche 1-10, weiterhin umfassend das Einschließen eines oder mehrerer *S. aureus*-Proteinantigen(e) in der Zusammensetzung, die ausgewählt sind aus der Gruppe bestehend aus einem clfA-Antigen; einem clfB-Antigen; einem sdrE2-Antigen; einem sdrC-Antigen; einem sasF-Antigen; einem emp-Antigen; einem sdrD-Antigen; einem spa-Antigen; einem esaC-Antigen; einem esxA-Antigen; einem esxB-Antigen; einem sta006-Antigen; einem isdC-Antigen; einem Hla-Antigen; einem sta011-Antigen; einem isdA-Antigen; einem isdB-Antigen; und einem sta073-Antigen.

Revendications

1. Procédé de fabrication d'une composition immunogène comprenant un polysaccharide capsulaire de *S. aureus* de type 5 conjugué à une molécule de support et un polysaccharide capsulaire de *S. aureus* de type 8 conjugué à une molécule de support, ledit procédé comprenant :

- (a) un procédé de purification de polysaccharide capsulaire à partir de cellules de *S. aureus* de type 5 comprenant une étape de libération du polysaccharide capsulaire à partir des cellules par traitement à l'acide des cellules, dans lequel les cellules sont sous la forme d'une pâte cellulaire humide ou sont en suspension dans un milieu aqueux, et la masse moléculaire du polysaccharide purifié se situe entre 2 et 3500 kDa ;
- (b) la conjugaison du polysaccharide capsulaire à une molécule de support pour fabriquer un conjugué ; et
- (c) le mélange du conjugué avec un polysaccharide capsulaire de *S. aureus* de type 8 conjugué à une molécule de support.

2. Procédé de fabrication d'une composition immunogène comprenant un polysaccharide capsulaire de *S. aureus* de type 8 conjugué à une molécule de support et un polysaccharide capsulaire de *S. aureus* de type 5 conjugué à une molécule de support, ledit procédé comprenant :

- (a) un procédé de purification de polysaccharide capsulaire à partir de cellules de *S. aureus* de type 8 comprenant une étape de libération du polysaccharide capsulaire à partir des cellules par traitement à l'acide des cellules, dans lequel les cellules sont sous la forme d'une pâte cellulaire humide ou sont en suspension dans un milieu aqueux, et la masse moléculaire du polysaccharide purifié se situe entre 2 et 3500 kDa ;
- (b) la conjugaison du polysaccharide capsulaire à une molécule de support pour fabriquer un conjugué ; et
- (c) le mélange du conjugué avec un polysaccharide capsulaire de *S. aureus* de type 5 conjugué à une molécule de support.

3. Procédé selon la revendication 1 ou la revendication 2, dans lequel (i) le traitement à l'acide est réalisé en utilisant de l'acide acétique et/ou (ii) le procédé comprend en outre une étape de neutralisation.

4. Procédé selon l'une quelconque des revendications précédentes, le procédé comprenant en outre une étape de centrifugation des cellules et de recueil du surnageant contenant le polysaccharide.

5. Procédé selon l'une quelconque des revendications 1 à 4, le procédé dans l'étape (a) comprenant en outre une étape de traitement du polysaccharide capsulaire avec de la DNase et/ou de la RNase.

6. Procédé selon l'une quelconque des revendications 1 à 5, le procédé dans l'étape (a) comprenant en outre une étape de traitement du polysaccharide capsulaire avec de la mutanolysine.

7. Procédé selon l'une quelconque des revendications 1 à 6, le procédé dans l'étape (a) comprenant en outre :

- (a) une étape de diafiltration ;
- (b) une étape de chromatographie à échange d'anions ;
- (c) une étape de filtration sur gel ; et/ou
- (d) une étape de concentration du polysaccharide.

8. Procédé selon l'une quelconque des revendications 1 à 7, le procédé dans l'étape (a) comprenant en outre :

- (a) une étape de dépolymérisation du polysaccharide purifié pour former un oligosaccharide ; et/ou
- (b) une étape de stérilisation par filtration.

9. Procédé selon l'une quelconque des revendications 1 à 8, le procédé dans l'étape (a) fournissant une composition comprenant :

- (a) le polysaccharide et un taux de contamination par des peptidoglycanes qui est inférieur à 5 % en poids de peptidoglycanes par rapport au poids total du polysaccharide ; et/ou
- (b) le polysaccharide et un taux de contamination par des protéines qui est inférieur à 5 % en poids de protéines par rapport au poids total du polysaccharide.

10. Procédé selon l'une quelconque des revendications 1 à 9, le procédé dans l'étape (a) fournissant une composition comprenant le polysaccharide et un taux de contamination par un acide nucléique qui est inférieur à 1 % en poids d'acide nucléique par rapport au poids total du polysaccharide.

11. Procédé selon l'une quelconque des revendications 1 à 10, comprenant en outre l'incorporation dans la composition d'un ou de plusieurs antigènes protéiniques de *S. aureus* choisis dans le groupe constitué d'un antigène clfA ; un antigène clfB ; un antigène sdrE2 ; un antigène sdrC ; un antigène sasF ; un antigène cmp ; un antigène sdrD ; un antigène spa ; un antigène esaC ; un antigène esxA ; un antigène esxB ; un antigène sta006 ; un antigène isdC ; un antigène Hla ; un antigène sta011 ; un antigène isdA ; un antigène isdB ; et un antigène sta073.

FIGURE 1




FIGURE 2





FIGURE 3





FIGURE 4




FIGURE 5




FIGURE 6





FIGURE 7

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description


- WO 2004080490 A [0203]
- WO 2006032475 A [0203]
- WO 2006032500 A [0203]
- WO 2006065553 A [0203]
- WO 2006114500 A [0203]
- WO 05033148 A [0203]
- WO 0056357 A [0203]
- US 61247518 B [0203]
- US IB201000256 W [0203]
- WO 2007113222 A [0203]
- US 6045805 A [0203]
- US 6027733 A [0203]
- US 6274144 B [0203]
- EP 0477508 A [0203]
- US 5306492 A [0203]
- WO 9842721 A [0203]
- WO 03061558 A [0203]
- EP 0372501 A [0203]
- EP 0378881 A [0203]
- EP 0427347 A [0203]
- WO 9317712 A [0203]
- WO 9403208 A [0203]
- WO 9858668 A [0203]
- EP 0471177 A [0203]
- WO 9101146 A [0203]
- EP 0594610 A [0203]
- WO 0056360 A [0203]
- WO 02091998 A [0203]
- WO 0172337 A [0203]
- WO 0061761 A [0203]
- WO 2004041157 A [0203]
- WO 0234771 A [0203]
- WO 9942130 A [0203]
- WO 2004011027 A [0203]
- WO 9640242 A [0203]
- WO 0038711 A [0203]
- US 6146902 A [0203]
- WO 9924578 A [0203]
- WO 9936544 A [0203]
- WO 9957280 A [0203]
- WO 0022430 A [0203]
- WO 9629412 A [0203]
- WO 0152885 A [0203]
- WO 03007985 A [0203]
- WO 0202606 A [0203]
- WO 9927105 A [0203]
- WO 0027994 A [0203]
- WO 0037494 A [0203]
- WO 9928475 A [0203]
- WO 03093306 A [0203]
- WO 2004018646 A [0203]
- US 4197290 A [0203]
- WO 2004043407 A [0203]
- WO 2007113224 A [0203]
- WO 2004043405 A [0203]
- WO 9810788 A [0203]
- WO 2007053176 A [0203]
- WO 2005009379 A [0203]
- WO 2009029132 A [0203]
- WO 2008079315 A [0203]
- WO 2005086663 A [0203]
- WO 2005115113 A [0203]
- WO 2006033918 A [0203]
- WO 2006078680 A [0203]
- WO 2007071692 A [0203]
- WO 2005009378 A [0203]
- WO 2007145689 A [0203]
- WO 2009029831 A [0203]
- WO 2005079315 A [0203]
- WO 2008152447 A [0203]
- US IB2010 W [0204]
- US 61256905 B [0204]

Non-patent literature cited in the description

- **FATTOM et al.** *Infect Immun*, 1990, vol. 58 (7), 2367-74 [0203]
- **FATTOM et al.** *Infect Immun*, 1992, vol. 60 (2), 584-9 [0203]
- **FATTOM et al.** *Infect Immun*, 1993, vol. 61 (3), 1023-32 [0203]
- **FATTOM et al.** *Infect Immun*, 1996, vol. 64 (5), 1659-65 [0203]
- **WELCH et al.** *J Am Soc Nephrol*, 1996, vol. 7 (2), 247-53 [0203]
- **FATTOM et al.** *Infect Immun*, 1998, vol. 66 (10), 4588-92 [0203]
- **FATTOM et al.** *Vaccine*, 1993, vol. 17 (2), 126-33 [0203]
- **FATTOM et al.** *N Engl J Med*, 2002, vol. 346 (7), 491-6 [0203]

- **ROBBINS et al.** *Ann N Y Acad Sci*, 2005, vol. 754, 68-82 [0203]
- **GILBERT et al.** *J. Microb. Meth.*, 1994, vol. 20, 39-46 [0203]
- **GILBERT et al.** *Vaccine*, 1994, vol. 12 (4), 369-74 [0203]
- **TOLLERSRUD et al.** *Vaccine*, 2001, vol. 19 (28-29), 3896-903 [0203]
- **LEE et al.** *Infect Immun*, 1993, vol. 61, 1853-8 [0203]
- **MOREAU et al.** *Carbohydrate Res*, 1990, vol. 339 (5), 285-91 [0203]
- **FOURNIER et al.** *Infect. Immun.*, 1984, vol. 45 (1), 87-93 [0203]
- **JONES.** *Carbohydrate Res.*, 2005, vol. 340 (6), 1097-106 [0203]
- **LEMERCINIER ; JONES.** *Carbohydrate Res.*, 1996, vol. 296, 83-96 [0203]
- **JONES ; LEMERCINIER.** *J Pharm Biomed Anal.*, 2002, vol. 30 (4), 1233-47 [0203]
- **HESTRIN.** *J. Biol. Chem.*, 1949, vol. 180, 249-261 [0203]
- **KONADU et al.** *Infect. Immun.*, 1994, vol. 62, 5048-5054 [0203]
- **WESSELS et al.** *Infect Immun*, 1989, vol. 57, 1089-94 [0203]
- **RAMSAY et al.** *Lancet*, 2001, vol. 357 (9251), 195-196 [0203]
- **LINDBERG.** *Vaccine*, 1999, vol. 17 (2), 28-36 [0203]
- **BUTTERY ; MOXON.** *J R Coll Physicians Lond*, 2000, vol. 34, 163-68 [0203]
- **AHMAD ; CHAPNICK.** *Infect Dis Clin North Am*, 1999, vol. 13, 113-33 [0203]
- **GOLDBLATT.** *J. Med. Microbiol.*, 1998, vol. 47, 563-7 [0203]
- **DICK et al.** *Conjugate Vaccines.* Karger, 1989, vol. 10, 48-114 [0203]
- **HERMANSON.** *Bioconjugate Techniques.* Academic Press, 1996 [0203]
- **REYNAUD-RONDIER et al.** *FEMS Microbiology Immunology*, 1991, vol. 76, 193-200 [0203]
- *Research Disclosure*, January 2002 [0203]
- **HERBELIN et al.** *J Dairy Sci.*, 1997, vol. 80 (9), 2025-34 [0203]
- **FALUGI et al.** *Eur J Immunol*, 2001, vol. 31, 3816-3824 [0203]
- **BARALDO et al.** *Infect Immun*, 2004, vol. 72 (8), 4884-7 [0203]
- **RUAN et al.** *J Immunol*, 1990, vol. 145, 3379-3384 [0203]
- **KUO et al.** *Infect Immun*, 1995, vol. 63, 2706-13 [0203]
- **MICHON et al.** *Vaccine*, 1998, vol. 16, 1732-41 [0203]
- **LEI et al.** *Dev Biol*, 2000, vol. 103, 259-264 [0203]
- **TETTELIN et al.** *Science*, 2000, vol. 287, 1809-1815 [0203]
- **PIZZA et al.** *Science*, vol. 287, 1816-1820 [0203]
- **BJUNE et al.** *Lancet*, 1991, vol. 338 (8775), 1093-1096 [0203]
- **FUKASAWA et al.** *Vaccine*, 1999, vol. 17, 2951-2958 [0203]
- **ROSENQVIST et al.** *Dev. Biol. Stand*, 1998, vol. 92, 323-333 [0203]
- **COSTANTINO et al.** *Vaccine*, 1992, vol. 10, 691-698 [0203]
- **WATSON.** *Pediatr Infect Dis J*, 2000, vol. 19, 331-332 [0203]
- **RUBIN.** *Pediatr Clin North Am*, 2000, vol. 47, 269-285 [0203]
- **JEDRZEJAS.** *Microbiol Mol Biol Rev*, 2001, vol. 65, 187-207 [0203]
- **BELL.** *Pediatr Infect Dis J*, 2000, vol. 19, 1187-1188 [0203]
- **IWARSON.** *APMIS*, 1995, vol. 103, 321-326 [0203]
- **GERLICH et al.** *Vaccine*, 1990, vol. 8, S63-68, 79-80 [0203]
- **HSU et al.** *Clin Liver Dis*, 1999, vol. 3, 901-915 [0203]
- **GUSTAFSSON et al.** *N. Engl. J. Med.*, 1996, vol. 334, 349-355 [0203]
- **RAPPUOLI et al.** *TIBTECH*, 1991, vol. 9, 232-238 [0203]
- *Vaccines.* 2004 [0203]
- **KALMAN et al.** *Nature Genetics*, 1999, vol. 21, 385-389 [0203]
- **READ et al.** *Nucleic Acids Res*, 2000, vol. 28, 1397-406 [0203]
- **SHIRAI et al.** *J. Infect. Dis.*, 2000, vol. 181 (3), S524-S527 [0203]
- **ROSS et al.** *Vaccine*, 2001, vol. 19, 4135-4142 [0203]
- **SUTTER et al.** *Pediatr Clin North Am*, 2000, vol. 47, 287-308 [0203]
- **ZIMMERMAN ; SPANN.** *Am Fam Physician*, 1999, vol. 59, 113-118, 125-126 [0203]
- **DREESEN.** *Vaccine*, 1997, vol. 15, S2-6 [0203]
- *MMWR Morb Mortal Wkly Rep*, 16 January 1998, vol. 47 (1), 12, , 19 [0203]
- **MCMICHAEL.** *Vaccine*, 2000, vol. 19 (1), 101-107 [0203]
- **DALE.** *Infect Dis Clin North Am*, 1999, vol. 13, 227-43 [0203]
- **FERRETTI et al.** *PNAS USA*, 2001, vol. 98, 4658-4663 [0203]
- **LCHIMAN ; YOSHIDA.** *J. Appl. Bacteriol.*, 1981, vol. 51, 229 [0203]
- **LCHIMAN et al.** *J. Appl. Bacteriol.*, 1991, vol. 71, 176 [0203]
- **ROBINSON ; TORRES.** *Seminars in Immunology*, 1997, vol. 9, 271-283 [0203]
- **DONNELLY et al.** *Annu Rev Immunol*, 1997, vol. 15, 617-648 [0203]
- **SCOTT-TAYLOR ; DALGLEISH.** *Expert Opin Investig Drugs*, 2000, vol. 9, 471-480 [0203]
- **APOSTOLOPOULOS ; PLEBANSKI.** *Curr Opin Mol Ther*, 2000, vol. 2, 441-447 [0203]

- **ILAN.** *Curr Opin Mol Ther*, 1999, vol. 1, 116-120 [0203]
- **DUBENSKY et al.** *Mol Med*, 2000, vol. 6, 723-732 [0203]
- **ROBINSON ; PERTMER.** *Adv Virus Res*, 2000, vol. 55, 1-74 [0203]
- **DONNELLY et al.** *Am J Respir Crit Care Med*, 2000, vol. 162 (4), S190-193 [0203]
- **DAVIS.** *Mt. Sinai J. Med*, 1999, vol. 66, 84-90 [0203]
- **GENNARO.** *Remington: The Science and Practice of Pharmacy*. 2000 [0203]
- **JOYCE et al.** *Carbohydrate Research*, 2003, vol. 338, 903 [0203]
- **MAIRA-LITRAN et al.** *Infect. Immun.*, 2002, vol. 70, 4433 [0203]
- **KURODA et al.** *Lancet*, 2001, vol. 357 (9264), 1225-1240 [0203]
- **SJODAHL.** *J. Biochem.*, 1977, vol. 73, 343-351 [0203]
- **UHLEN et al.** *J. Biol. Chem.*, 1984, vol. 259, 1695-1702, 13628 [0203]
- **SCHNEEWIND et al.** *Cell*, 1992, vol. 70, 267-281 [0203]
- **DEDENT et al.** *EMBO J.*, 2008, vol. 27, 2656-2668 [0203]
- **SJOQUIST et al.** *Eur. J. Biochem.*, 1972, vol. 30, 190-194 [0203]
- **DEDENT et al.** *J. Bacteriol.*, 2007, vol. 189, 4473-4484 [0203]
- **DEISENHOFER et al.** *Hoppe-Seyh Zeitsch. Physiol. Chem.*, 1978, vol. 359, 975-985 [0203]
- **DEISENHOFER.** *Biochemistry*, 1981, vol. 20, 2361-2370 [0203]
- **GRAILLE et al.** *Proc. Nat. Acad. Sci. USA*, 2000, vol. 97, 5399-5404 [0203]
- **O'SEAGHDHA et al.** *FEBS J.*, 2006, vol. 273, 4831-41 [0203]
- **GOMEZ et al.** *J. Biol. Chem.*, 2006, vol. 281, 20190-20196 [0203]
- **SEBULSKY ; HEINRICHS.** *J Bacteriol*, 2001, vol. 183, 4994-5000 [0203]
- **SEBULSKY et al.** *J Biol Chem*, 2003, vol. 278, 49890-900 [0203]
- **RABLE ; WARDENBURG.** *Infect Immun*, 2009, vol. 77, 2712-8 [0203]
- **KUKLIN et al.** *Infect Immun.*, 2006, vol. 74 (4), 2215-23 [0203]
- **GENNARO.** *Remington: The Science and Practice of Pharmacy*, 2000 [0203]
- *Methods In Enzymology*. Academic Press, Inc, [0203]
- *Handbook of Experimental Immunology*. Scientific Publications, 1986, vol. I-IV [0203]
- **SAMBROOK et al.** *Molecular Cloning: A Laboratory Manual*. Cold Spring Harbor Laboratory Press, 2001 [0203]
- *Handbook of Surface and Colloidal Chemistry*. CRC Press, 1997 [0203]
- *Short protocols in molecular biology*. 2002 [0203]
- *Molecular Biology Techniques: An Intensive Laboratory Course*. Academic Press, 1998 [0203]
- *PCR (Introduction to Biotechniques Series)*. Springer Verlag, 1997 [0203]
- *Current Protocols in Molecular Biology*. 1987 [0203]
- **SMITH ; WATERMAN.** *Adv. Appl. Math.*, 1981, vol. 2, 482-489 [0203]
- **GEYSEN et al.** *PNAS USA*, 1984, vol. 81, 3998-4002 [0203]
- **CARTER.** *Methods Mol Biol*, 1994, vol. 36, 207-23 [0203]
- **JAMESON, BA et al.** *CABIOS*, 1988, vol. 4 (1), 181-186 [0203]
- **RADDRIZZANI ; HAMMER.** *Brief Bioinform*, 2000, vol. 1 (2), 179-89 [0203]
- **BUBLIL et al.** *Proteins*, 2007, vol. 68 (1), 294-304 [0203]
- **DELALLA et al.** *J. Immunol.*, 1999, vol. 163, 1725-29 [0203]
- **KWOK et al.** *Trends Immunol*, 2001, vol. 22, 583-88 [0203]
- **BRUSIC et al.** *Bioinformatics*, 1998, vol. 14 (2), 121-30 [0203]
- **MEISTER et al.** *Vaccine*, 1995, vol. 13 (6), 581-91 [0203]
- **ROBERTS et al.** *AIDS Res Hum Retroviruses*, 1996, vol. 12 (7), 593-610 [0203]
- **MAKSYUTOV ; ZAGREBELNAYA.** *ComputAppl Biosci*, 1993, vol. 9 (3), 291-7 [0203]
- **FELLER ; DE LA CRUZ.** *Nature*, 1991, vol. 349 (6311), 720-1 [0203]
- **HOPP.** *Peptide Research*, 1993, vol. 6, 183-190 [0203]
- **WELLING et al.** *FEBS Lett.*, 1985, vol. 188, 215-218 [0203]
- **DAVENPORT et al.** *Immunogenetics*, 1995, vol. 42, 392-297 [0203]
- **TSURUI ; TAKAHASHI.** *J Pharmacol Sci*, 2007, vol. 105 (4), 299-316 [0203]
- **TONG et al.** *Brief Bioinform*, 2007, vol. 8 (2), 96-108 [0203]
- **SCHIRLE et al.** *J Immunol Methods*, 2001, vol. 257 (1-2), 1-16 [0203]
- **CHEN et al.** *Amino Acids*, 2007, vol. 33 (3), 423-8 [0203]
- **KIM et al.** *Biochemistry*, 2008, vol. 47 (12), 3822-3831 [0203]
- **PATTI et al.** *Biochemistry*, 2008, vol. 47 (32), 8378-8385 [0203]
- **KIM ; SCHAEFER.** *Biochemistry*, 2008, vol. 47 (38), 10155-10161 [0203]
- **BISWAS.** PhD Thesis: Characterization of *Staphylococcus aureus* peptidoglycan hydrolases and isolation of defined peptidoglycan structures. Eberhard Karls Universität Tübingen, 2006 [0203]



SZTNH-100026122

SZABADALMI IGÉNYPONTOK

1. Eljárás egy hordozómolekulához konjugált, *S. aureus*ból származó 5. típusú kapszuláris poliszacharidot és egy hordozómolekulához konjugált, *S. aureus*ból származó 8. típusú kapszuláris poliszacharidot tartalmazó immunogén készítmény előállítására, mely eljárás tartalmazza a következőket:

(a) eljárás kapszuláris poliszacharid tisztítására 5. típusú *S. aureus* sejtekből, mely tartalmaz egy lépést, melyben a kapszuláris poliszacharidot felszabadítjuk a sejtekből a sejtek savas kezelése útján, ahol a sejtek nedves sejtmassza formájában vannak vagy vizes közegben szuszpendálva vannak, és a tisztított poliszacharid molekulatömege 2-3500 kDa;

(b) a kapszuláris poliszacharid konjugálása egy hordozómolekulához konjugátum előállítása céljából; és

(c) a konjugátum elkeverése egy hordozómolekulához konjugált, *S. aureus*ból származó 8. típusú kapszuláris poliszachariddal.

2. Eljárás egy hordozómolekulához konjugált, *S. aureus*ból származó 8. típusú kapszuláris poliszacharidot és egy hordozómolekulához konjugált, *S. aureus*ból származó 5. típusú kapszuláris poliszacharidot tartalmazó immunogén készítmény előállítására, mely eljárás tartalmazza a következőket:

(a) eljárás kapszuláris poliszacharid tisztítására 8. típusú *S. aureus* sejtekből, mely tartalmaz egy lépést, melyben a kapszuláris poliszacharidot felszabadítjuk a sejtekből a sejtek savas kezelése útján, ahol a sejtek nedves sejtmassza formájában vannak vagy vizes közegben szuszpendálva vannak, és a tisztított poliszacharid molekulatömege 2-3500 kDa;

(b) a kapszuláris poliszacharid konjugálása egy hordozómolekulához konjugátum előállítása céljából; és

(c) a konjugátum elkeverése egy hordozómolekulához konjugált, *S. aureus*ból származó 5. típusú kapszuláris poliszachariddal.

3. Az 1. vagy 2. igénypont szerinti eljárás, ahol (i) a savas kezelést ecetsav alkalmazásával végezzük és/vagy (ii) az eljárás tartalmaz továbbá egy semlegesítési lépést.

4. Az előző igénypontok bármelyike szerinti eljárás, ahol az eljárás tartalmaz továbbá egy lépést a sejtek centrifugálására és a poliszacharid-tartalmú felülúszó begyűjtésére.

5. Az 1-4. igénypontok bármelyike szerinti eljárás, ahol az eljárás az (a) lépésben tartalmaz továbbá egy lépést a kapszuláris poliszacharid DNÁzzal és/vagy RNÁzzal történő kezelésére.
6. Az 1-5. igénypontok bármelyike szerinti eljárás, ahol az eljárás az (a) lépésben tartalmaz továbbá egy lépést a kapszuláris poliszacharid mutanolizinnel történő kezelésére.
7. Az 1-6. igénypontok bármelyike szerinti eljárás, ahol az eljárás az (a) lépésben tartalmazza továbbá a következőket:
- (a) egy diaszűrési lépés;
 - (b) egy anioncserélő kromatográfiás lépés;
 - (c) egy gélszűrési lépés; és/vagy
 - (d) egy lépés a poliszacharid töményítésére.
8. Az 1-7. igénypontok bármelyike szerinti eljárás, ahol az eljárás az (a) lépésben tartalmazza továbbá a következőket:
- (a) egy lépés a tisztított poliszacharid depolimerizálására oligoszacharid előállítása céljából; és/vagy
 - (d) egy sterilszűrési lépés.
9. Az 1-8. igénypontok bármelyike szerinti eljárás, ahol az eljárás az (a) lépésben egy következőket tartalmazó készítményt eredményez:
- (a) a poliszacharidot és peptidoglikán szennyeződést, melynek mennyisége a poliszacharid össztömegére vonatkoztatva kevesebb, mint 5 tömeg% peptidoglikán; és/vagy
 - (d) a poliszacharidot és fehérje szennyeződést, melynek mennyisége a poliszacharid össztömegére vonatkoztatva kevesebb, mint 5 tömeg% fehérje.
10. Az 1-9. igénypontok bármelyike szerinti eljárás, ahol az eljárás az (a) lépésben olyan készítményt eredményez, amely a poliszacharidot és nukleinsav szennyeződést tartalmaz, melynek mennyisége a poliszacharid össztömegére vonatkoztatva kevesebb, mint 1 tömeg% nukleinsav.
11. Az 1-10. igénypontok bármelyike szerinti eljárás, amely tartalmazza továbbá a következő csoportból választott egy vagy több *S. aureus* fehérje antigén belefoglalását a készítménybe: egy clfA antigén; egy clfB antigén; egy sdrE2 antigén; egy sdrC antigén; egy sasF antigén; egy emp antigén; egy sdrD antigén; egy spa antigén; egy esaC antigén; egy esxA antigén; egy esxB antigén; egy sta006 antigén; egy isdC antigén; egy Hla antigén; egy sta011 antigén; egy isdA antigén; egy isdB antigén; és egy sta073 antigén.