



US 20030204867A1

(19) **United States**

(12) **Patent Application Publication** (10) **Pub. No.: US 2003/0204867 A1**
(43) **Pub. Date: Oct. 30, 2003**

(54) **DNA SEQUENCES FOR AMMONIUM
TRANSPORTER, PLASMIDS, BACTERIA,
YEASTS, PLANT CELLS AND PLANTS
CONTAINING THE TRANSPORTER**

(76) Inventors: **Wolf-Bernd Frommer**, Berlin (DE);
Olaf Ninnemann, Berlin (DE)

Correspondence Address:

FROMMER LAWRENCE & HAUG
745 FIFTH AVENUE- 10TH FL.
NEW YORK, NY 10151 (US)

(21) Appl. No.: **10/449,148**

(22) Filed: **May 30, 2003**

Related U.S. Application Data

(62) Division of application No. 08/635,967, filed on Apr.
29, 1996, filed as 371 of international application No.
PCT/EP94/03499, filed on Oct. 24, 1994.

(30) **Foreign Application Priority Data**

Oct. 28, 1993 (DE)..... P4337597.9

Publication Classification

(51) **Int. Cl.⁷** **C12Q 1/68**; C07H 21/04;
C12P 21/02; C12N 5/04; C12N 1/21;
C12N 1/18; A01H 1/00; C12N 15/82
(52) **U.S. Cl.** **800/278**; 435/6; 435/69.1;
435/320.1; 435/254.2; 435/252.3;
530/370; 536/23.6

(57) **ABSTRACT**

There are described DNA sequences, that contain the coding region of ammonium transporters, which after introduction in suitable vectors are introduced into the plant genome and lead to the formation of mRNA which makes possible either the formation of new ammonium transporters, or prevents the formation of endogenous ammonium transporters in transgenic plants. There are further described plasmids, bacteria, yeast strains, plant cells and transgenic plants, which contain the sequences of ammonium transporters as a constituent of a recombinant DNA, and also a process for the identification and isolation of DNA sequences, coding for ammonium transporters.

FIG. 1





FIG. 2



DNA SEQUENCES FOR AMMONIUM TRANSPORTER, PLASMIDS, BACTERIA, YEASTS, PLANT CELLS AND PLANTS CONTAINING THE TRANSPORTER

[0001] The present invention relates to DNA sequences that contain the coding region of ammonium transporters, whose introduction into a plant genome modifies the uptake and transfer of nitrogen compounds in transgenic plants, as well as plasmids, bacteria, yeasts, plant cells and plants containing these DNA sequences, and also a process for the identification and isolation of DNA sequences that code for the ammonium transporter.

[0002] The supplying of growing plants with nitrogen compounds is a limiting factor in biomass production and is thus a limit on the yield of agricultural production. For this reason, nitrogen compounds, often in the form of mineral fertilisers, are added in agricultural biomass production.

[0003] An only partial uptake of the added nitrogen compounds by the plants makes it on the one hand necessary that the nitrogen fertiliser produced with high energy input is used in an excess, and on the other hand it leads only to a partial uptake so that the nitrogen compounds are washed into the ground water which can lead to considerable ecological problems.

[0004] There is thus a great interest in plants which are capable of taking up large amounts of nitrogen as well as in the provision of the possibility of modifying the nitrogen uptake in plants.

[0005] For many plants there is provided information that the uptake of nitrogen is essentially in the form of nitrate salts. In strongly acid soils or in soils which follow intensive cultivation or have a strong tannin content, the nitrate formation (nitrification) is however strongly reduced and the uptake of nitrogen in the form of ammonium becomes the most important mechanism for the uptake of nitrogen compounds (Raven & Smith 1976 New Phytol 76: 415-431). Plants which are well adapted to acid soils, appear in part to favour ammonium rather than nitrate uptake and can tolerate ammonium ion concentrations that would be toxic for other plants. Examples of these plants are sugar cane, *Betula verucosa* or *Lolium rigidum* (Foy et al 1978, Ann Rev Plant Physiol 29: 511-566). The toxicity of the ammonium follows from a displacement of the ion balance: the uptake of the positively charged ammonium ion leads to an acidification of the cytoplasm, provided that no cations are secreted in counter-exchange. In an uptake of ammonium via a transport system whose uptake mechanism is based on an ammonium ion proton antiport, the ionic imbalance is not a problem.

[0006] There is thus a great interest in transport systems which work by the mechanism of an ammonium ion proton antiport and/or in systems which could be converted through techniques of protein engineering into ammonium ion proton antiports.

[0007] In spite of extensive efforts, it has not been possible up until now to isolate transport systems with whose help plants are protected against an ammonium ion loss caused by membrane diffusion (retrieval system).

[0008] By the term ammonium is also to be understood methylamine which is analogous to ammonium.

[0009] An active uptake system has been investigated in the fungus *Aspergillus nidulans* (e.g. Arst et al., 1973, Mol Gen Genet 121: 239-245), as well as in *Penicillium chrysogenum* (Hackette et al., J Biol Chem 245: 4241-4250). In these studies methylamine was used as the ammonium analogue. For *Aspergillus nidulans*, five genetic loci were established, which take part in the transport of methylamine (Pateman et al., 1973, J Bacteriol 114: 943-950). In biochemical experiments concerning the methylammonium transport in *Penicillium chrysogenum* and *Saccharomyces cerevisiae*, it has been shown that the transport is temperature and pH dependent, and that the pH optimum is 6.0 to 6.5 and the transport efficiency steadily rises up to a temperature of 35° C. (Roon et al., 1975, J Bacteriol 122: 502-509). The methylamine and/or ammonium transport in *Saccharomyces cerevisiae* is dependent on the supply of easily usable chemical energy, e.g. in the form of glucose. The transport system consists of at least three independent transporters, which differ in transport capacity and affinity for the substrate: besides a high affinity transporter with low capacity (K_m value=250 μ M, maximum speed V_{max} =20 nmol/min per mg cells (dry weight)) there is a low affinity system with high capacity (K_m =2 mM, V_{max} =50 nmol/min per mg cells) and a low affinity system with medium capacity (K_m =20 mM, V_{max} =33 nmol/min per mg cells) (Dubois & Grenson, 1979, Mol Gen Genet 175: 67-76). In the presence of glutamine or asparagine in the surrounding medium, the ammonium uptake is reduced by 60 to 70%, whilst other amines hardly have any influence (Roon et al., 1975, J Bacteriol 122: 502-509).

[0010] Nothing is known about the molecular nature of the ammonium transporter of the above-mentioned or different fungi and other organisms, such as for example bacteria. Equally nothing is known about systems, with whose help fungi or bacteria protect against ammonium ion loss by membrane diffusion (retrieval systems) at the molecular level. Further genes, which code for ammonium transporters are not known.

[0011] Various evidence suggests that the ammonium transport in *Saccharomyces cerevisiae* is accomplished by at least two functionally different transport systems (Dubois & Grenson, 1979, Mol Gen Genet 175: 67-76):

[0012] 1. Kinetic analyses of the methylamine-uptake by *Saccharomyces* show an abrupt transition between apparent linear sections, whereby both functions are inhibitable by ammonium.

[0013] 2. Both functions can be separately excluded by mutation. The resulting mutations mep-1 and mep-2 are genetically independent.

[0014] 3. The mutants mep-1 and mep-2 can each grow in media with ammonium as the only nitrogen source, whilst a double mutant mep-1/mep-2 shows hardly any growth under these conditions (data taken from Dubois & Grenson, 1979).

[0015] A clarification of the ammonium transport processes in plants, that leads to similar detailed information such as for yeast, is not available and because of the difficulty of the molecular biological analysis of mutations is scarcely possible in a corresponding manner.

[0016] The object of the present invention is to provide, coding for DNA-sequences of ammonium transporters,

which cause a change in the uptake and transfer of nitrogen compounds in transgenic plants.

[0017] The object of the present invention is further to provide, DNA-constructs, such as plasmids, with which the ammonium transport in transgenic plants can be modified by introduction of the corresponding construct (plasmids) into the plant genome which leads upon transcription to the formation of a new ammonium transporter molecule in the transgenic plant and/or the suppression of the formation the plant's own ammonium transporter molecules.

[0018] There have now surprisingly been found DNA sequences, that contain the coding region of a plant ammonium transporter, whereby the information contained in the nucleotide sequence when integrated in a plant genome

[0019] a) makes possible under control of a promoter in sense orientation, the expression of a translatable mRNA, which leads to the synthesis of an ammonium transporter in transgenic plants, or

[0020] b) makes possible under control of a promoter in anti-sense orientation, the expression of a non-translatable mRNA, which prevents the synthesis of an endogenous ammonium transporter in transgenic plants.

[0021] A further aspect of the invention are therefore DNA sequences, which contain the coding region of a plant ammonium transporter.

[0022] In an analogous way the ammonium transporter can also be used to modify animal cells.

[0023] The identification of DNA sequences, which code for a plant ammonium transporter can take place by a process, in which DNA sequences are identified and isolated, which are able to complement specific mutations in the yeast *Saccharomyces cerevisiae*, especially mutations, which have the result that the corresponding strains cannot grow any further in media which contain ammonium as the only nitrogen source. Such a strain can be transformed with a plant cDNA-library and transformands can be selected which can grow in media which contain ammonium as the only nitrogen source.

[0024] A further aspect of the invention is a process for the identification and isolation of DNA sequences from plants that code for ammonium transporters, which includes the following steps:

[0025] a) transformation of a yeast strain, which cannot grow in media which contain ammonium as the only nitrogen source, with a plant cDNA-library using suitable expression vectors,

[0026] b) selection and propagation of transformands, which after expression of plant cDNA-sequences, can grow in media which contain ammonium as the only nitrogen source, and

[0027] c) isolation of the expression vectors, which carry a plant cDNA insert from the selected transformand.

[0028] The yeast strain in process step a) is preferably one which cannot take up ammonium from the medium because of mutations in the transport systems for the ammonium uptake. Preferred is the double mutant mep1/mep2 (strain 26972c), described by Dubois & Gresson (1979, Mol. Gen. Genet. 175:67-76), and with which two uptake systems for ammonium are interrupted following mutation.

[0029] A further aspect of the invention are DNA sequences from plants, which are obtainable using the above described process and which code for a protein with the biological activity of an ammonium transporter.

[0030] Whether it is possible using such complementation process to identify DNA sequences which code for plant ammonium transporters depends on various factors. First the expression plasmid suitable for use in yeast must contain cDNA fragments which code for the plant ammonium transporter, that means the mRNA fundamental for the cDNA synthesis must result from tissues which express the ammonium transporter. Since nothing is known about plant ammonium transporters, tissues which code for the ammonium transporter are therefore also not known.

[0031] A further prerequisite for the success of the complementation strategy is that ammonium transport systems existing in plants can be functionally expressed in yeast, since only in this case is a complementation of the deficiency liberated by the mutation possible. Since nothing is known about plant ammonium transporters, it is also not known whether a functional expression in yeast is possible.

[0032] It has now further been surprisingly found that by expression of a cDNA library, for example from leaf tissue of *Arabidopsis thaliana*, by means of expression plasmids suitable for use in yeast which contain the promoter of phosphoglycerate kinase from yeast, the complementation of the double mutation mep-1/mep-2 is possible, if the expression plasmids contain specified plant cDNA fragments. These cDNA fragments code for plant ammonium transporters.

[0033] The identification of plant ammonium transporters is described here using *Arabidopsis thaliana* as an example, but it is not however limited to this plant species.

[0034] A cDNA fragment that codes for a plant ammonium transporter containing for example the following sequence. (Seq. ID No.1):

TTCTTCTCTAAACTCTCAAC

20

ATG TCT TGC TCG GCC ACC GAT CTC GCC GTC CTG TTG CGT CCT AAT.65
Met Ser Cys Ser Ala Thr Asp Leu Ala Val Leu Leu Gly Pro Asn

GCC ACG GCG GCG GCC AAC TAC ATA TGT GGC CAG CTA GGC GAC GTC 110
Ala Thr Ala Ala Ala Asn Tyr Ile Cys Gly Gln Leu Gly Asp Val

-continued

AAC AAC AAA TTC ATC GAC ACC GCT TTC GCT ATA GAC AAC ACT TAC 155
Asn Asn Lys Phe Ile Asp Thr Ala Phe Ala Ile Asp Asn Thr Tyr

CTC CTC TTC TCC GCC TAC CTT GTC TTC TCT ATG CAG CTT GGC TTC 200
Leu Leu Phe Ser Ala Tyr Leu Val Phe Ser Met Gln Leu Gly Phe

GCT ATG CTC TGT GCC CGT TCC GTG AGA GCC AAG AAT ACT ATG AAC 245
Ala Met Leu Cys Ala Gly Ser Val Arg Ala Lys Asn Thr Met Asn

ATC ATG CTT ACC AAC GTC CTT GAC GCT GCA GCC GGT GGT CTC TTC 290
Ile Met Leu Thr Asn Val Leu Asp Ala Ala Ala Gly Gly Leu Phe

TAT TAT CTG TTT GGC TAC GCC TTT GCC TTT GGA TCT CCG TCC AAT 335
Tyr Tyr Leu Phe Gly Tyr Ala Phe Ala Phe Gly Ser Pro Ser Asn

GGT TTC ATC GGT AAA CAC TAC TTT GGT CTC AAA GAC ATC CCC ACG 380
Gly Phe Ile Gly Lys His Tyr Phe Gly Leu Lys Asp Ile Pro Thr

GCC TCT GCT GAC TAC TCC AAC TTT CTC TAC CAA TGG GCC TTT GCA 425
Ala Ser Ala Asp Tyr Ser Asn Phe Leu Tyr Gln Trp Ala Phe Ala

ATC GCT GCG GCT GGA ATC ACA ACT GGC TCC ATC GCT GAA CGG ACA 470
Ile Ala Ala Ala Gly Ile Thr Ser Gly Ser Ile Ala Glu Arg Thr

CAG TTC GTG GCT TAC CTA ATC TAT TCC TCT TTC TTA ACC GGG TTT 515
Gln Phe Val Ala Tyr Leu Ile Tyr Ser Ser Phe Leu Thr Gly Phe

GTT TAC CCG GTC GTC TCT CAC TCG TTC TGG TCA GTT CAT GGA TGG 560
Val Tyr Pro Val Val Ser His Trp Phe Trp Ser Val Asp Gly Trp

GCC AGC CCG TTC CGT ACC GAT GGA CAT TTG CTT TTC AGC ACC GGA 605
Ala Ser Pro Phe Arg Thr Asp Gly Asp Leu Leu Phe Ser Thr Gly

GCG ATA GAT TTC GCT GGG TCC GGT GTT GTT CAT ATG GTC GGA GGT 650
Ala Ile Asp Phe Ala Gly Ser Gly Val Val His Met Val Gly Gly

ATC GCT GGA CTC TGG GGT GCG CTC ATC GAA GGT CCA CGA CTT GGC 695
Ile Ala Gly Leu Trp Gly Ala Leu Ile Glu Gly Pro Arg Leu Gly

CGG TTC GAT AAC GGA GGC CGT GCC ATC GCT CTT CGT GGC CAC TCG 740
Arg Phe Asp Asn Gly Gly Arg Ala Ile Ala Leu Arg Gly His Ser

GCG TCA CTT GTT GTC CTT GGA ACA TTC CTC CTC TGG TTT GGA TGG 785
Ala Ser Leu Val Val Leu Gly Thr Phe Leu Leu Trp Phe Gly Trp

TAC GGA TTT AAC CCC GGT TCC TTC AAC AAG ATC CTA GTC ACG TAC 830
Tyr Gly Phe Asn Pro Gly Ser Phe Asn Lys Ile Leu Val Thr Tyr

GAG ACA GGC ACA TAC AAC GGC CAG TGG AGC GCG GTC GGA CGG ACA 875
Glu Thr Gly Thr Tyr Asn Gly Gln Trp Ser Ala Val Gly Arg Thr

GCT GTC ACA ACA ACG TTA GCT GGC TGC ACC GCG GCG CTG ACA ACC 920
Ala Val Thr Thr Thr Leu Ala Gly Cys Thr Ala Ala Leu Thr Thr

CTA TTT CCC AAA CGT CTA CTC TCG GGA CAT TGG AAC GTC ACT GAT 965
Leu Phe Gly Lys Arg Leu Leu Ser Gly His Trp Asn Val Thr Asp

GTA TGC AAC GGC CTC CTC GGA GGG TTT GCA GCC ATA ACT GGT GGC 1010
Val Cys Asn Gly Leu Leu Gly Gly Phe Ala Ala Ile Thr Gly Gly

TGC TCT GTC GTT GAG CCA TGG GCT GCG ATC ATC TGC GGG TTC GTG 1055
Cys Ser Val Val Glu Pro Trp Ala Ala Ile Ile Cys Gly Phe Val

GCG GCC CTA GTC CTC CTC GGA TGC AAC AAG CTC GCT GAG AAG CTC 1100
Ala Ala Leu Val Leu Leu Gly Cys Asn Lys Leu Ala Glu Lys Leu

AAA TAC GAC GAC CCT CTT GAG GCA GCA CAA CTA CAC GGT GGT TGC 1145
Lys Tyr Asp Asp Pro Leu Glu Ala Ala Gln Leu His Gly Gly Cys

GGT GCG TGG GGA CTA ATA TTC ACG GCT CTC TTC GCT CAA GAA AAG 1190
Gly Ala Trp Gly Leu Ile Phe Thr Ala Leu Phe Ala Gln Glu Lys

TAC TTG AAC CAG ATT TAC GGC AAC AAA CCC GGA AGG CCA CAC GGT 1235
Tyr Leu Asn Gln Ile Tyr Gly Asn Lys Pro Gly Arg Pro His Gly

TTG TTT ATG GGC GGT GGA GGA AAA CTA CTT GGA GCT CAG CTG ATT 1280

-continued

Leu Phe Met Gly Gly Gly Lys Leu Leu Gly Ala Gln Leu Ile

CAG ATC ATT GTG ATC ACG GGT TGG GTA AGT GCG ACC ATG GGG ACA 1325
Gln Ile Ile Val Ile Thr Gly Trp Val Ser Ala Thr Met Gly Thr

CTT TTC TTC ATC CTC AAG AAA ATG AAA TTG TTG CCG ATA TCG TCC 1370
Leu Phe Phe Ile Leu Lys Lys Met Lys Leu Leu Arg Ile Ser Ser

GAG GAT GAG ATG GCC GGT ATG GAT ATG ACC AGG CAC GGT GGT TTT 1415
Glu Asp Glu Met Ala Gly Met Asp Met Thr Arg His Gly Gly Phe

GCT TAT ATG TAC TTT GAT CAT GAT GAG TCT CAC AAA GCC ATT CAG 1460
Ala Tyr Met Tyr Phe Asp Asp Glu Ser His Lys Ala Ile Gln

CTT AGG AGA GTT GAG CCA CGA TCT CCT TCT CCT TCT GGT GCT AAT 1505
Leu Arg Arg Val Glu Pro Arg Ser Pro Ser Pro Ser Gly Ala Asn

ACT ACA CCT ACT CCG GTT TGATTGGAT TTTTACTTTT ATTCTCTATT 1553
Thr Thr Pro Thr Pro Val

TTCTAGAGTA TTATTTTAAA TGATGTTTTG TGATACTTAA ATATTGTTTT 1603
GGATATTTTT TTCGCATTTC AGTAATGTTT TAGATGTACA GTTTCATCGG 1653

GTTGTGATGA TAATATCTAT GTGGTCATTT GTGTTCTCTT TCGAGTTAAA 1703
AAAAAAAAAA AAAAAAAAAA AAAAAAAAAA AAAAAAAAAA AAAAA 1748

[0035] The DNA sequences of the invention, identified using the transformed yeast strain, such as e.g. the sequence Seq. ID No. 1, can be introduced into plasmids and thereby be combined with regulatory elements for expression in eukaryotic cells (see Example 4). These regulatory elements are on the one hand transcription promoters, and on the other hand transcription terminators. With the plasmids, eukaryotic cells can be transformed, with the aim of expression of a translatable mRNA which makes possible the synthesis of an ammonium transporter in the cells (sense orientation) or with the aim of expression of a non-translatable RNA, which prevents synthesis of an endogenous ammonium transporter in the cells (anti-sense orientation).

[0036] These plasmids are also an aspect of the invention.

[0037] Preferred plasmids are plasmids p35S-MEP-a and p35S-a-MEP-a, which have been deposited at the Deutsche Sammlung von Mikroorganismen (DSM).

[0038] A still further aspect of the present invention are yeasts, which contain the DNA sequences of the invention.

[0039] The yeast strains used for the identification of a plant methylamine or ammonium transporter can be used for studies of the properties of the transporter as well as its substrate. The DNA sequences of a plant ammonium transporter present in plasmids can, in accordance with results of these studies, be subjected to a mutagenesis or sequence modification by recombination, in order to change the properties of the transporter. By changing the specificity of the transport system, the transport of new compounds is possible, which opens up interesting applications (see below). In addition, by suitable change of the transporter, the transport mechanism can be modified. One can envisage, especially but not exclusively, a change, that modifies the cotransport properties of the transporter, for example with the effect that this results in an ammonium ion-proton-antiport, which prevents a toxicity of ammonium ions taken up for plants as a result of acidification of the cytoplasm. In this way, the yeast strain of the invention, which contains the cDNA sequence of a plant ammonium transporter in an expression plasmid, can be used for a mutation selection system.

[0040] By expression of an RNA corresponding to the DNA sequences of the invention of plant ammonium transporters in transgenic plants it is possible to have a change of the plant nitrogen metabolism, whose economic significance is obvious. Nitrogen is the nutrient mainly responsible for limiting growth. The viability of seedlings as well as germination capacity of seeds is directly dependent on the nitrogen content of storage tissue. The formation of high value food materials with a high protein content is dependent on a sufficient nitrogen supply.

[0041] The change of nitrogen uptake, for example by addition of a new uptake system of nitrogen compounds such as ammonium, can thus lead to an increase in yield of transgenic crops, especially under nitrogen limitation. In this way, transgenic plants can be cultivated in high yields under low input conditions.

[0042] The possibility of suppressing the take-up of ammonium in the transgenic plants can however also be desirable under certain conditions. For example one can envisage the cultivation of transgenic plants on acid soils which are not suitable for growth. As a result of suppressing nitrification ammonium ion concentrations on such soils could be present which would be toxic for certain plants, because, the ammonium take up from the plants in the cells can no longer be fully metabolised and acts as a cell poison. The suppression of the take-up of ammonium by suppressing the biosynthesis of the ammonium transporter can thus alter transgenic plants to the extent that cultivation on acid soils becomes possible.

[0043] A further aspect of the present invention are transgenic plants, in which the DNA sequences of the invention are introduced as a constituent of a recombinant DNA molecule, in which this recombinant DNA molecule is stably integrated into the genome, and in whose cells based on the presence of these sequences there is achieved either a synthesis of an additional plant ammonium transporter whereby these cells can take up larger amounts of ammonium in comparison with untransformed plants or there is achieved an inhibition of the synthesis of endogenous

ammonium transporters whereby such cells show a reduced uptake of ammonium in comparison with untransformed cells.

[0044] Transgenic crops are, for example tobacco, potatoes, sugar beet, soya beans, peas, beans or maize.

[0045] The genetic modification of dicotyledonous and monocotyledonous plants can be carried out by currently known processes, (see for example Gasser, C. S., Fraley, R. T., 1989, Science 244: 1293-1299; Potrykus, 1991, Ann Rev Plant Mol Biol Plant Physiol 42: 205-225). For expression in plants the coding sequences must be coupled with the transcriptional regulatory elements. Such elements called promoters, are known (EP 375091).

[0046] Further, the coding regions must be provided with transcription termination signals with which they can be correctly transcribed. Such elements are also described (see Gielen et al., 1989, EMBO J 8: 23-29). The transcriptional start region can be both native and/or homologous as well as foreign and/or heterologous to the host plant. If desired, termination regions are interchangeable with one another. The DNA sequence of the transcription starting and termination regions can be prepared synthetically or obtained naturally, or obtained from a mixture of synthetic and natural DNA constituents. For introduction of foreign genes in higher plants a large number of cloning vectors are available that include a replication signal for *E. coli* and a marker which allows a selection of the transformed cells. Examples of such vectors are pBR 322, pUC-Series, M13 mp-Series, pACYC 184 etc. Depending on the method of introduction of the desired gene in the plants, other DNA sequences may be suitable. Should the Ti- or Ri-plasmid be used, e.g. for the transformation of the plant cell, then at least the right boundary, often however both the right and left boundary of the Ti- and Ri-Plasmid T-DNA, is attached, as a flanking region, to the gene being introduced. The use of T-DNA for the transformation of plants cells has been intensively researched and is well described in EP 120 516; Hoekama, In: The Binary Plant Vector System, Offsetdrukkerij Kanters B. V. Alblasserdam, (1985), Chapter V; Fraley, et al., Crit. Rev. Plant Sci., 4:1-46 and An et al. (1985) EMBO J. 4: 277-287. Once the introduced DNA is integrated in the genome, it is as a rule stable there and remains also in the offspring of the original transformed cells. It normally contains a selection marker, which induces resistance in the transformed plant cells against a biocide or antibiotic such as kanamycin, G 418, bleomycin, hygromycin or phosphinotricin etc. The individual marker employed should therefore allow the selection of transformed cells from cells, which lack the introduced DNA.

[0047] For the introduction of DNA into a plant host cell, besides transformation using *Agrobacteria*, there are many other techniques available. These techniques include the fusion of protoplasts, microinjection of DNA and electroporation, as well as ballistic methods and virus infection. From the transformed plant material, whole plants can be regenerated in a suitable medium, which contains antibiotics or biocides for the selection. The resulting plants can then be tested for the presence of introduced DNA.

[0048] A further aspect of the present invention are transformed plant cells, in which the DNA sequences of the invention are introduced as a constituent of a recombinant DNA molecule, in which this recombinant DNA molecule is

stably integrated into the genome, and in whose cells based on the presence of these sequences there is achieved either a synthesis of an additional plant ammonium transporter whereby these cells can take up larger amounts of ammonium in comparison with untransformed plants or there is achieved an inhibition of the synthesis of endogenous ammonium transporters whereby such cells show a reduced uptake of ammonium in comparison with untransformed cells.

[0049] No special demands are placed on the plasmids in injection and electroporation. Simple plasmids, such as e.g. pUC-derivatives can be used. Should however whole plants be regenerated from such transformed cells the presence of a selectable marker gene is necessary. The transformed cells grow within the plants in the usual manner (see also McCormick et al. (1986) Plant Cell Reports 5: 81-84). These plants can be grown normally and crossed with plants, that possess the same transformed genes or different. The resulting hybrid individuals have the corresponding phenotypical properties.

[0050] The introduction of DNA plant ammonium transporters for changing the uptake of nitrogen compounds is described here using *Arabidopsis thaliana* and tobacco as examples. The use is not however limited to this plant species.

[0051] The DNA sequences of the invention can also be introduced in plasmids and thereby combined with steering elements for an expression in prokaryotic cells. The formation of a translatable RNA sequence of a eukaryotic ammonium transporter from bacteria leads, in spite of the considerable differences in the membrane structures of prokaryotes and eukaryotes, means in addition surprisingly, that prokaryotes can now express a functional eukaryotic ammonium transporter with its substrate specificity. This makes possible the production of bacterial strains, which as for the yeast strain used for identifying the ammonium transporter could be used for studies of the properties of the transporter as well as its substrate, which opens up interesting applications.

[0052] The invention also relates to bacteria, that contain the plasmids of the invention.

[0053] The DNA sequences of the invention can also be introduced in plasmids which allow mutagenesis or a sequence modification through recombination of DNA sequences using standard microbiological processes. In this way the specificity of the ammonium transporter can be modified.

[0054] Modified ammonium transporters can be used for example for transformation of agricultural transgenic plants, whereby both transport of pesticides and also plant growth regulators to plants can be envisaged.

[0055] By using standard processes (see Sambrook et al., 1989, Molecular Cloning: A Laboratory Manual, 2. Edn., Cold Spring Harbor Laboratory Press, NY, USA) base exchanges can be carried out or natural or synthetic sequences can be added. For fusing DNA fragments with one another adaptors or linkers can be added to the fragments. Further, manipulations can be carried out which prepare suitable restriction cleavage sites or remove the excess DNA or restriction cleavage sites. Where insertions, deletions or substitutions such as for example transitions and transversions are to be carried out, in vitro mutagenesis,

primer repair, restrictions or ligations can be used. For methods of analysis, in general a sequence analysis, restriction analysis and other biochemical molecular biological methods can be used. After each manipulation, the DNA sequence, used, can be cleaved and fused with another DNA sequence. Each plasmid sequence can be cloned in the same or different plasmids.

[0056] The invention also relates to derivatives or parts of plasmids on which the DNA sequences of the invention are localised.

[0057] Derivatives or parts of the DNA sequences and plasmids of the invention can also be used for the transformation of prokaryotic and eukaryotic cells.

[0058] Further, the DNA sequences of the invention can be used according to standard processes for the isolation of homologous sequences from the genome of plants of various species, which also code for ammonium transporter molecules. With these sequences, constructs for the transformation of plant cells can be prepared which modify the transport processes in transgenic plants.

[0059] By the terms "homology" or "homologous sequences" are to be understood, a sequence identity of 60% to 80%, preferably 80% to 95% and especially 95% to 100%.

[0060] In order to specify related DNA sequences, gene libraries must first be prepared, which are representative for the content in genes of a plant species or for the expression of genes in a plant species. The former are genomic libraries, whilst the latter are cDNA libraries. From these, related sequences can be isolated using the DNA sequences of the invention as probes. Once the related gene has been identified and isolated, a determination of the sequence and an analysis of the properties of the proteins coded from this sequence is possible.

[0061] DNA sequences of ammonium transporters obtained in this way are also part of the invention and could be used as described above.

[0062] The use of the DNA sequences as described above is also part of the invention.

[0063] A further aspect of the invention are DNA sequences from plants, which hybridise with DNA sequences of the invention and code for a protein that possesses the biological activity of an ammonium transporter. The term "hybridisation" means in this connection, a hybridisation under conventional hybridisation conditions, preferably under stringent conditions, such as described for example by Sambrook et al. (1989, Molecular Cloning, A Laboratory Manual, 2. Edn. Cold Spring Harbor Laboratory Press, Cold Spring Harbour, N.Y.). An important biological activity of an ammonium transporter is the capability of transporting ammonium or analogues thereof through biological membranes. This activity can be measured by uptake of ammonium or analogues thereof by cells, which express the particular ammonium transporter, as described for example in example 3 of the present invention.

[0064] Deposits

[0065] The following plasmids were deposited at the Deutschen Sammlung von Mikroorganismen (DSM) in Braunschweig, Germany on the 26.10.1993 (deposit number):

Plasmid	p35S-MEP-a	(DSM 8651)
Plasmid	p35S-a-MEP-a	(DSM 8652)

DESCRIPTION OF THE FIGURES

[0066] FIG. 1 shows the plasmid p35S-MEP-a, which is a derivative of the plasmids pBIN19 (Bevan, M., 1984, Nucl Acids Res 12: 8711-8721). It comprises:

[0067] A=Fragment from the genome of the cauliflower mosaic virus that carries the 35S promoter (nt 6909-7437). The promoter fragment was prepared as an EcoRI/KpnI fragment from plasmid pDH51 (Pietrzak et al., Nucl. Acid Res. 14, 5857-5868).

[0068] B=NotI/NotI fragment of the cDNA with the coding region of the ammonium transporter of *Arabidopsis thaliana* in sense orientation to the fragment A. The arrow marked in fragment B indicates the reading direction of the cDNA.

[0069] C=Polyadenylation signal of the gene 3 of the T-DNA of the plasmid pTiACH5 (Gielen et al., EMBO J 3: 835-846), nucleotides 11749 to 11939, which was isolated as a PvuII/HindIII fragment from plasmid pAGV40 (Herrera-Estrella et al., Nature 303, 209-213) and, after the addition of SphI linker onto the PvuII cleavage site, was cloned between the SphI and HindIII cleavage sites of the polylinker from pBIN19.

[0070] FIG. 2 shows the plasmid p35S-a-MEP-a, which is a derivative of the plasmids pBIN19 (Bevan, M., 1984, Nucl Acids Res 12: 8711-8721). It comprises:

[0071] A=Fragment from the genome of the cauliflower mosaic virus that carries the 35S promoter (nt 6909-7437). The promoter fragment was prepared as an EcoRI/KpnI fragment from plasmid pDH51 (Pietrzak et al., Nucl. Acid Res. 14, 5857-5868).

[0072] B=NotI/NotI fragment of the cDNA with the coding region of the ammonium transporter of *Arabidopsis thaliana* in anti-sense orientation to the fragment A. The arrow marked in fragment B indicates the reading direction of the cDNA.

[0073] C=Polyadenylation signal of the gene 3 of the T-DNA of the plasmid pTiACH5 (Gielen et al., EMBO J 3: 835-846), nucleotides 11749 to 11939, which was isolated as a PvuII/HindIII fragment from plasmid pAGV40 (Herrera-Estrella et al., Nature 303, 209-213) and, after the addition of SphI linker onto the PvuII cleavage site, was cloned between the SphI and HindIII cleavage sites of the polylinker from pBIN19.

[0074] The following examples illustrate the invention without limiting it in which there is described the cloning and identification as well as the function of a plant ammonium transporter and its use for the genetic modification of plants with aim of changing the uptake and transfer of nitrogen compounds. In an analogous manner to the Examples, other plants especially crop plants such as potato, sugar beet, maize, etc, can be modified.

[0075] In Examples 1 to 4 the following standard processes and special techniques were used.

[0076] 1. Cloning Process

[0077] For cloning in *E. coli*, a derivative of the vector pACYC that contains the polylinker from pBluescriptSK was used.

[0078] For the transformation of yeasts, the vector pFL61 (Minet & Lacroute, 1990, Curr Genet 18: 287-291) was used.

[0079] For the plant transformation the gene constructs were cloned in the binary vector pBinAR, a derivative of pBIN19 (Bevan, 1984, Nucl Acids Res 12: 8711-8721), was cloned (see Example 4).

[0080] 2. Bacterial and Yeast Strains

[0081] For the pACYC vector as well as for pBinAR constructs, *E. coli* strain DH5 α was used.

[0082] As starting strain for the production of the cDNA library in yeast, the yeast strain 26972c (Dubois & Gresson, 1979, Mol Gen Gnete 175: 67-76) with the mutations mep-1/mep-2 and ura3 was used.

[0083] The transformation of the plasmid in potato plants was carried out using *Agrobacterium tumefaciens* strain LBA4404 (Bevan (1984) Nucl. Acids Res 12: 8711-8720).

[0084] 3. Transformation of *Agrobacterium tumefaciens*

[0085] The transfer of the DNA in the *Agrobacteria* was carried out by direct transformation by the method of Höfgen & Willmitzer (1988, Nucleic Acids Res 16: 9877). The plasmid DNA of the transformed *Agrobacterium* was isolated in accordance with the method of Birnboim and Doly (1979) (Nucl Acids Res 7: 1513-1523) and was analysed by gel electrophoresis after suitable restriction cleavage.

[0086] 4. Plant Transformation

[0087] Tobacco

[0088] 10 ml of an *Agrobacterium tumefaciens* overnight culture grown under selection was sedimented and resuspended in the same volume of antibiotic free medium. In a sterile petri dish, leaf discs of sterile plants (approximately 1 cm²), the central vein of which had been removed, were immersed in this bacterial suspension. The leaf discs were then placed in a closely packed arrangement in petri dishes containing MS medium (Murashige et al. (1962) Physiologia Plantarum 15, 473-497) with 2% sucrose and 0.8% bacto agar. After two days incubation in the dark at 25° C., they were transferred onto MS medium containing 500 mg/l Claforan, 50 mg/l kanamycin, 1 mg/l benzylaminopurine (BAP), 0.2 mg/l of naphthylacetic acid (NAA) and 0.8% bacto agar. Growing shoots were transferred onto hormone-free MS medium with 250 mg/l Claforan and 50 mg/l Kanamycin.

[0089] *Arabidopsis thaliana*

[0090] Cotyledons from a seven day old sterile culture of *Arabidopsis thaliana* grown from seeds, were pre-incubated on CI-Medium for two days with a layer of tobacco suspension culture (as feeder layer) and then for 5 minutes in a suspension with *Agrobacterium tumefaciens* (preparation of suspension see under tobacco transformation) and then incubated for two days on CI-Medium with feeder layer in low light conditions. After this cocultivation with Agrobak-

teria the explantates were laid out on SIM I-medium, that was weekly renewed. After callous-formation the explantates were set transplanted to SIM II-medium and the cultivated under selection with kanamycin. Small shoots were separated from the callous material and transplanted to RI-Medium. After separation of seeds, these could be analysed.

[0091] The individual media were as follows:

Cl-medium	SIM I-medium	SIM II-medium	RI-Medium
Murashige-Skoog medium with 2% sucrose, 8 g/l agar, 1 mg/l 2,4-dichloro-phenoxy-acetic acid (2,4-D), 0.2 mg/l kinetin	Murashige-Skoog medium with 2% sucrose, 8 g/l agar, 1 mg/l 6-benzyl-aminopurine, 0.4 mg/l naphthyl-acetic acid, 0.5 g/l Claforan, 100 mg/l kanamycin	Murashige-Skoog medium with 2% sucrose, 8 g/l agar, 7 mg/l N ⁶ -[2-iso-pentenyl]-adenine, 0.05 mg/l indolyl-acetic acid, 0.5 mg/l Clatoran, 100 mg/l kanamycin	Murashige-Skoog medium with 1% sucrose, 8 g/l agar, 1 mg/l indolyl-butyric acid

EXAMPLE 1

[0092] Cloning of the cDNA of a Plant Methylamine or Ammonium Transporter

[0093] For complementation of the yeast strain 26972c (Dubois & Gresson, 1979, Mol Gen Genet 175: 67-76), there was used a cDNA of young germ lines from *Arabidopsis thaliana* (two leaf stage) in the yeast expression vector pFL61 (Minet & Lacroute, 1990, Curr Genet 18: 287-291) which had been made available by Minet (Minet et al., 1992, Plant J 2: 417-422). Around 1 μ g of the vector with the cDNA-insert was transformed in the yeast strain 26972c by the method of Dohmen et al. (1991, Yeast 7: 691-692). Yeast transformants, which could grow in minimal medium with 1 mM NH₄Cl as the sole nitrogen source were propagated. From the lines, plasmid-DNA was prepared by standard methods. This DNA was immediately transformed in strain 26972c. In this way a plasmid pFL61-MEP-a was obtained that can complement the mep-1/mep-2 double mutation. This plasmid has an insertion of size 1.75 kbp.

[0094] The yeast strain 26972c::pFL61-MEP-a obtain by transformation of 26972c with plasmid pFL61-MEP-a can be used for uptake studies with methylamine or ammonium (see example 3). By genetic modification of the coding region of the ammonium transporter gene MEP-a by standard methods (cf. Sambrook et al., 1989, Molecular cloning: A laboratory manual, 2. Edn., Cold Spring Harbor Laboratory Press, NY, USA) the specificity or characteristic of the transport mechanism of which can be modified. The strain 26972c::pFL61-MEP-a is suitable directly for testing for inhibitors or promoters of the ammonium transport, using the ammonium transport system of the invention (see Example 3).The cDNA insertion of the plasmids pFL61-MEP-a can be used for the identification of similar DNA sequences from plants of other species or from other organisms, such as bacteria or animal systems. For this hybridi-

sation techniques are usable. Likewise using the cDNA sequence, a construct for the expression of the gene product as a fusion protein in bacteria can be prepared by standard processes. Using the fusion protein, an antibody, which identifies similar proteins in other organisms, can be prepared.

EXAMPLE 2

[0095] Sequence Analyses of the cDNA Insert of the Plasmid pFL61-MEP-a

[0096] From a yeast line 26972c::pFL61-MEP-a, obtained from example 1, the plasmid pFL61-MEP-a was isolated and its cDNA insert prepared as a NotI fragment. The fragment was cloned in a modified pACYC vector (Chang & Cohen, 1978, J Bacteriol 134. 1141-1156), which contained, between a HindII (nt 3211) and a filled HindIII (nt 1523) cutting position, the polylinker from pBluescript as BssHII fragment. Through the modification the tetracycline resistance is lost. The cDNA fragment was cloned in the thus obtained vector pACH-H as NotI fragment in the NotI cutting position. Using synthetic oligonucleotides, the insert was sequenced by the method of Sanger et al. (1977, Proc Natl Acad Sci USA 74: 5463-5467). The sequence is given in Seq ID No 1.

EXAMPLE 3

[0097] Uptake Studies With ¹⁴C-labelled Methylamine Into the Yeast Line 26972c::pFL61-MEP-a

[0098] The yeast lines 26972c::pFL61, 26972c::pFL61-MEP-a and their starting lines S1278b (Dubois & Grenson, 1979, Mol Gen Genet 175: 67-76, MEP-1/MEP-2 ura3) were grown in liquid medium with 1.7 g yeast nitrogen base without ammonium and without amino acids (Difco), 20 g agarose, 1% glucose (w/v), 0.5 mg/l proline until the culture reached the logarithmic phase. After centrifuging of each 25 ml of the culture the cells were washed with 10 mM phosphate pH 7 or pH 4 and taken up in 1.5 ml 10 mM phosphate buffer. Ten minutes before starting the transport measurements, it was made up to an end concentration of 10 mM glucose. 100 µml of the suspension was added to a solution of 10 mM phosphate pH 4 or 7, 100 µM unlabelled methylamine and 5 µCi ¹⁴C-labelled methylamine (0.1 µCi/ 1.9 µmol) (end concentrations). The uptake of the methylamine was measured after 10, 60, 120 and 180 seconds. For this, to 50 µl of starting material was taken and in a volume of 1 ml H₂O+3 mM unlabelled methylamine absorbed on a glass fibre filter. After washing with 10 ml H₂O+3 mM unlabelled methylamine the amount of ¹⁴C-labelled methylamine taken up was measured by scintillation (Table Ia and Ib). The uptake of the labelled methylamine was compared for the case of coincubation with 0.5 mM NH₄Cl (Table II) and 0.5 mM KCl (Table III), as well as for the case of coincubation with the uncouplers, dinitrophenol (DNP) and carbonyl cyanide m-chlorophenylhydrazone (CCCP) (Table IV). The values of a typical experiment are given in Tables I to IV:

TABLE 1

Uptake of ¹⁴C-methylamine in yeast strains after incubation with 10 mM glucose at different pH values (at each time-point for the MEP-a transformants and the zero controls (plasmid pFL61 without cDNA insertion) values from two independent series of measurements are given. Measurements in cpm (counts per minute)

Time point [sec]	Σ1278b	26972c::pFL61	26972c::pFL61-MEP-a		
pH = 4					
10	289.00	220.20	117.20	204.80	214.80
60	1151.20	200.60	138.60	283.80	345.60
120	19605.46	284.80	197.40	434.40	594.60
180	32651.43	317.60	285.60	479.60	842.20
pH = 7					
10	469.60	202.20	./.	302.80	352.20
60	1022.80	291.80	./.	1247.80	1292.20
120	6176.36	524.80	./.	3152.50	3478.28
180	16826.67	797.80	./.	8490.83	6099.39

[0099]

TABLE II

Uptake of ¹⁴C-methylamine in yeast strains after incubation with 10 mM glucose in presence of 0.5 mM NH₄Cl. Measurements in cpm. pH = 7

Time point [sec]	1278b	26972c::pFL61	26972c::pFL61-MEP-a	
10	356.60	271.60	268.60	
60	885.60	795.80	577.60	
120	1237.80	1388.20	978.00	
180	1597.40	1773.00	1226.60	

[0100]

TABLE III

Uptake of ¹⁴C-methylamine in yeast strains after incubation with 10 mM glucose in presence of 0.5 mM KCl, unless otherwise noted. Measurements in cpm. pH = 7

Time point [sec]	Σ1278b	26972c::pFL61	26972c::pFL61-MEP-a	26972c::pFL61-MEP-a without KCl-competition
10	391.20	127.60	384.00	205.00
60	2054.90	224.00	1276.60	467.60
120	8455.00	374.80	3128.44	1541.00
180	14377.14	567.40	5056.00	2416.39

[0101]

TABLE IV

Uptake of ¹⁴C-methylamine in yeast strains after incubation with 10 mM glucose in presence of 0.1 mM dinitrophenol (DNP) or 10 µM carbonyl cyanide m-chlorophenylhydrazone. Measurements in cpm (counts per minute). pH = 7

Time point [sec]	Σ1278b +DNP	Σ1278b +CCCP	26972c::pFL61-MEP-a +DNP	26972c::pFL61-MEP-a +CCCP	26972c::pFL6-MEP-a
10	167.20	173.20	378.00	484.60	373.60
60	386.00	316.80	1349.40	1407.00	1214.20

TABLE IV-continued

Uptake of ¹⁴ C-methylamine in yeast strains after incubation with 10 mM glucose in presence of 0.1 mM dinitrophenol (DNP) or 10 μM carbonyl cyanide m-chlorophenylhydraZone. Measurements in cpm (counts per minute). pH = 7					
Time point [sec]	Σ1278b +DNP	Σ1278b +CCCP	26972c::pFL61-MEP-a +DNP	26972c::pFL61-MEP-a +CCCP	26972c::pFL6-MEP-a
120	706.80	460.40	2794.44	2597.92	3673.45
180	906.00	608.00	2794.44	3620.36	6135.15

EXAMPLE 4

[0102] Transformation of Plants With a construct for Over-Expression of the Coding Region of the Ammonium Transporter

[0103] From the plasmid pFL61-MEP-a, that contains as insert the cDNA for the methylamine or ammonium transporter from *Arabidopsis thaliana*, the insert was isolated as NotI fragment and cloned after filling in the overhanging ends in the SmaI cutting position of pBinAR. The cDNA has the designation “B” in the plasmid map (FIG. 1). According to whether or not B is introduced in sense orientation to the 35S Promoter of pBinAR, the resulting plasmid has the

designation p35S-MEP-a or p35S-a-MEP-a. The plasmid pBinAR is a derivative of pBIN19 (Bevan, 1984, Nucl Acids Res 12: 8711-8720). Between its EcoRI and KpnI cutting positions, a fragment from the genome of the cauliflower mosaic virus, that carries the 35S promoter (nt 6909-7437), was introduced. The promoter fragment prepared as EcoRI/KpnI fragment from the plasmid pDH51 (Pietrzak et al., Nucl Acids Res 14: 5857-5868). In the plasmid map the promoter fragment has the designation “A”. Between the SphI and the HindIII cutting position of pBinAR, the polyadenylation signal of the gene 3 of the T-DNA of the plasmid pTiACH5 (Gielen et al., EMBO J 3:835-846) is inserted. Also a PvuII/HindIII fragment (nt 11749-11939) from the plasmid pAGV 40 (Herrera-Estrella et al., 1983, Nature 303: 209-2139) was supplied at the PvuII cutting position with a SphI linker. The polyadenylation signal has the designation “C” in the plasmid map.

[0104] After transformation of Agrobacteria with the plasmids p35S-MEP-a and p35S-a-MEP-a, these were used for infection of leaf segments of tobacco and *Arabidopsis thaliana*.

[0105] Ten independently obtained transformands for both constructs, in which the presence of the intact, non-rearranged chimeric gene was demonstrated using Southern blot analysis, were tested for changes in nitrogen content.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(iii) NUMBER OF SEQUENCES: 2

(2) INFORMATION FOR SEQ ID NO: 1:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 1748 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: cDNA
- (vi) ORIGINAL SOURCE:
 - (A) ORGANISM: *Arabidopsis thaliana*
 - (C) INDIVIDUAL ISOLATE: Ammonium transporter
- (vii) IMMEDIATE SOURCE:
 - (A) LIBRARY: cDNA library in plasmid pFL61
- (viii) POSITION IN GENOME:
 - (B) MAP POSITION: from 21 to 1523 coding region
- (ix) FEATURE:
 - (A) NAME/KEY: CDS
 - (B) LOCATION: 21..1526
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

TTCTTCTCTA	AACTCTCAAC	ATG TCT TGC TCG GCC ACC GAT CTC GCC GTC	50
	Met Ser Cys Ser Ala Thr Asp Leu Ala Val		
	1	5	10
CTG TTG GGT CCT AAT GCC ACG GCG GCG GCC AAC TAC ATA TGT GGC CAG			98

-continued

Leu	Leu	Gly	Pro	Asn	Ala	Thr	Ala	Ala	Ala	Asn	Tyr	Ile	Cys	Gly	Gln	
				15						20				25		
CTA	GGC	GAC	GTC	AAC	AAC	AAA	TTC	ATC	GAC	ACC	GCT	TTC	GCT	ATA	GAC	146
Leu	Gly	Asp	Val	Asn	Asn	Lys	Phe	Ile	Asp	Thr	Ala	Phe	Ala	Ile	Asp	
			30					35					40			
AAC	ACT	TAC	CTC	CTC	TTC	TCC	GCC	TAC	CTT	GTC	TTC	TCT	ATG	CAG	CTT	194
Asn	Thr	Tyr	Leu	Leu	Phe	Ser	Ala	Tyr	Leu	Val	Phe	Ser	Met	Gln	Leu	
		45				50						55				
GGC	TTC	GCT	ATG	CTC	TGT	GCC	GGT	TCC	GTG	AGA	GCC	AAG	AAT	ACT	ATG	242
Gly	Phe	Ala	Met	Leu	Cys	Ala	Gly	Ser	Val	Arg	Ala	Lys	Asn	Thr	Met	
	60					65					70					
AAC	ATC	ATG	CTT	ACC	AAC	GTC	CTT	GAC	GCT	GCA	GCC	GGT	GGT	CTC	TTC	290
Asn	Ile	Met	Leu	Thr	Asn	Val	Leu	Asp	Ala	Ala	Ala	Gly	Gly	Leu	Phe	
	75				80				85						90	
TAT	TAT	CTG	TTT	GGC	TAC	GCC	TTT	GCC	TTT	GGA	TCT	CCG	TCC	AAT	GGT	338
Tyr	Tyr	Leu	Phe	Gly	Tyr	Ala	Phe	Ala	Phe	Gly	Ser	Pro	Ser	Asn	Gly	
			95					100						105		
TTC	ATC	GGT	AAA	CAC	TAC	TTT	GGT	CTC	AAA	GAC	ATC	CCC	ACG	GCC	TCT	386
Phe	Ile	Gly	Lys	His	Tyr	Phe	Gly	Leu	Lys	Asp	Ile	Pro	Thr	Ala	Ser	
			110				115						120			
GCT	GAC	TAC	TCC	AAC	TTT	CTC	TAC	CAA	TGG	GCC	TTT	GCA	ATC	GCT	GCG	434
Ala	Asp	Tyr	Ser	Asn	Phe	Leu	Tyr	Gln	Trp	Ala	Phe	Ala	Ile	Ala	Ala	
	125						130					135				
GCT	GGA	ATC	ACA	AGT	GGC	TCG	ATC	GCT	GAA	CGG	ACA	CAG	TTC	GTG	GCT	482
Ala	Gly	Ile	Thr	Ser	Gly	Ser	Ile	Ala	Glu	Arg	Thr	Gln	Phe	Val	Ala	
	140					145					150					
TAC	CTA	ATC	TAT	TCC	TCT	TTC	TTA	ACC	GGG	TTT	GTT	TAC	CCG	GTC	GTC	530
Tyr	Leu	Ile	Tyr	Ser	Ser	Phe	Leu	Thr	Gly	Phe	Val	Tyr	Pro	Val	Val	
	155				160				165					170		
TCT	CAC	TGG	TTC	TGG	TCA	GTT	GAT	GGA	TGG	GCC	AGC	CCG	TTC	CGT	ACC	578
Ser	His	Trp	Phe	Trp	Ser	Val	Asp	Gly	Trp	Ala	Ser	Pro	Phe	Arg	Thr	
			175					180						185		
GAT	GGA	GAT	TTG	CTT	TTC	AGC	ACC	GGA	GCG	ATA	GAT	TTC	GCT	GGG	TCC	626
Asp	Gly	Asp	Leu	Phe	Ser	Thr	Thr	Gly	Ala	Ile	Asp	Phe	Ala	Gly	Ser	
		190						195				200				
GGT	GTT	GTT	CAT	ATG	GTC	GGA	GGT	ATC	GCT	GGA	CTC	TGG	GGT	GCG	CTC	674
Gly	Val	Val	His	Met	Val	Gly	Gly	Ile	Ala	Gly	Leu	Trp	Gly	Ala	Leu	
	205					210						215				
ATC	GAA	GGT	CCA	CGA	CTT	GGC	CGG	TTC	GAT	AAC	GGA	GGC	CGT	GCC	ATC	722
Ile	Glu	Gly	Pro	Arg	Leu	Gly	Arg	Phe	Asp	Asn	Gly	Gly	Arg	Ala	Ile	
	220					225				230						
GCT	CTT	CGT	GGC	CAC	TCG	GCG	TCA	CTT	GTT	GTC	CTT	GGA	ACA	TTC	CTC	770
Ala	Leu	Arg	Gly	His	Ser	Ala	Ser	Leu	Val	Val	Leu	Gly	Thr	Phe	Leu	
	235				240					245				250		
CTC	TGG	TTT	GGA	TGG	TAC	GGA	TTT	AAC	CCC	GGT	TCC	TTC	AAC	AAG	ATC	818
Leu	Trp	Phe	Gly	Trp	Tyr	Gly	Phe	Asn	Pro	Gly	Ser	Phe	Asn	Lys	Ile	
			255					260						265		
CTA	GTC	ACG	TAC	GAG	ACA	GGC	ACA	TAC	AAC	GGC	CAG	TGG	AGC	GCG	GTC	866
Leu	Val	Thr	Tyr	Glu	Thr	Gly	Thr	Tyr	Asn	Gly	Gln	Trp	Ser	Ala	Val	
		270						275					280			
GGA	CGG	ACA	GCT	GTC	ACA	ACA	ACG	TTA	GCT	GGC	TGC	ACC	GCG	GCG	CTG	914
Gly	Arg	Thr	Ala	Val	Thr	Thr	Thr	Leu	Ala	Gly	Cys	Thr	Ala	Ala	Leu	
	285						290					295				
ACA	ACC	CTA	TTT	GGG	AAA	CGT	CTA	CTC	TCG	GGA	CAT	TGG	AAC	GTC	ACT	962
Thr	Thr	Leu	Phe	Gly	Lys	Arg	Leu	Leu	Ser	Gly	His	Trp	Asn	Val	Thr	
	300					305					310					
GAT	GTA	TGC	AAC	GGC	CTC	CTC	GGA	GGG	TTT	GCA	GCC	ATA	ACT	GGT	GGC	1010

-continued

Asp	Val	Cys	Asn	Gly	Leu	Leu	Gly	Gly	Phe	Ala	Ala	Ile	Thr	Gly	Gly	
315					320				325						330	
TGC	TCT	GTC	GTT	GAG	CCA	TGG	GCT	GCG	ATC	ATC	TGC	GGG	TTC	GTG	GCG	1058
Cys	Ser	Val	Val	Glu	Pro	Trp	Ala	Ala	Ile	Ile	Cys	Gly	Phe	Val	Ala	
				335					340					345		
GCC	CTA	GTC	CTC	CTC	GGA	TGC	AAC	AAG	CTC	GCT	GAG	AAG	CTC	AAA	TAC	1106
Ala	Leu	Val	Leu	Leu	Gly	Cys	Asn	Lys	Leu	Ala	Glu	Lys	Leu	Lys	Tyr	
				350				355					360			
GAC	GAC	CCT	CTT	GAG	GCA	GCA	CAA	CTA	CAC	GGT	GGT	TGC	GGT	GCG	TGG	1154
Asp	Asp	Pro	Leu	Glu	Ala	Ala	Gln	Leu	His	Gly	Gly	Cys	Gly	Ala	Trp	
		365					370					375				
GGA	CTA	ATA	TTC	ACG	GCT	CTC	TTC	GCT	CAA	GAA	AAG	TAC	TTG	AAC	CAG	1202
Gly	Leu	Ile	Phe	Thr	Ala	Leu	Phe	Ala	Gln	Glu	Lys	Tyr	Leu	Asn	Gln	
	380					385					390					
ATT	TAC	GGC	AAC	AAA	CCC	GGA	AGG	CCA	CAC	GGT	TTG	TTT	ATG	GGC	GGT	1250
Ile	Tyr	Gly	Asn	Lys	Pro	Gly	Arg	Pro	His	Gly	Leu	Phe	Met	Gly	Gly	
395					400					405					410	
GGA	GGA	AAA	CTA	CTT	GGA	GCT	CAG	CTG	ATT	CAG	ATC	ATT	GTG	ATC	ACG	1298
Gly	Gly	Lys	Leu	Leu	Gly	Ala	Gln	Leu	Ile	Gln	Ile	Ile	Val	Ile	Thr	
			415					420					425			
GGT	TGG	GTA	AGT	GCG	ACC	ATG	GGG	ACA	CTT	TTC	TTC	ATC	CTC	AAG	AAA	1346
Gly	Trp	Val	Ser	Ala	Thr	Met	Gly	Thr	Leu	Phe	Phe	Ile	Leu	Lys	Lys	
			430					435					440			
ATG	AAA	TTG	TTG	CGG	ATA	TCG	TCC	GAG	GAT	GAG	ATG	GCC	GGT	ATG	GAT	1394
Met	Lys	Leu	Leu	Arg	Ile	Ser	Ser	Glu	Asp	Glu	Met	Ala	Gly	Met	Asp	
		445					450					455				
ATG	ACC	AGG	CAC	GGT	GGT	TTT	GCT	TAT	ATG	TAC	TTT	GAT	GAT	GAT	GAG	1442
Met	Thr	Arg	His	Gly	Gly	Phe	Ala	Tyr	Met	Tyr	Phe	Asp	Asp	Asp	Glu	
		460				465					470					
TCT	CAC	AAA	GCC	ATT	CAG	CTT	AGG	AGA	GTT	GAG	CCA	CGA	TCT	CCT	TCT	1490
Ser	His	Lys	Ala	Ile	Gln	Leu	Arg	Arg	Val	Glu	Pro	Arg	Ser	Pro	Ser	
475					480					485					490	
CCT	TCT	GGT	GCT	AAT	ACT	ACA	CCT	ACT	CCG	GTT	TGA	TTTGGATTTT				1536
Pro	Ser	Gly	Ala	Asn	Thr	Thr	Pro	Thr	Pro	Val	*					
			495							500						
TACTTTTATT	CTCTATTTTC	TAGAGTATTA	TTTTAAATGA	TGTTTTGTGA	TACTTAAATA											1596
TTGTTTTGGA	TATTTTTTTTG	GCATTTCACT	AATGTTTTAG	ATGTACAGTT	TCATGGGGTT											1656
GTGATGATAA	TATCTATGTG	GTCATTTGTG	TTCTCTTTGG	AGTTAAAAAA	AAAAAAAAAA											1716
AAAAAAAAAA	AAAAAAAAAA	AAAAAAAAAA	AA													1748

(2) INFORMATION FOR SEQ ID NO: 2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 501 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

Met Ser Cys Ser Ala Thr Asp Leu Ala Val Leu Leu Gly Pro Asn Ala
 1 5 10 15

Thr Ala Ala Ala Asn Tyr Ile Cys Gly Gln Leu Gly Asp Val Asn Asn
 20 25 30

Lys Phe Ile Asp Thr Ala Phe Ala Ile Asp Asn Thr Tyr Leu Leu Phe
 35 40 45

-continued

Ser	Ala	Tyr	Leu	Val	Phe	Ser	Met	Gln	Leu	Gly	Phe	Ala	Met	Leu	Cys
50						55					60				
Ala	Gly	Ser	Val	Arg	Ala	Lys	Asn	Thr	Met	Asn	Ile	Met	Leu	Thr	Asn
65					70					75					80
Val	Leu	Asp	Ala	Ala	Ala	Gly	Gly	Leu	Phe	Tyr	Tyr	Leu	Phe	Gly	Tyr
			85						90					95	
Ala	Phe	Ala	Phe	Gly	Ser	Pro	Ser	Asn	Gly	Phe	Ile	Gly	Lys	His	Tyr
		100						105					110		
Phe	Gly	Leu	Lys	Asp	Ile	Pro	Thr	Ala	Ser	Ala	Asp	Tyr	Ser	Asn	Phe
	115						120					125			
Leu	Tyr	Gln	Trp	Ala	Phe	Ala	Ile	Ala	Ala	Ala	Gly	Ile	Thr	Ser	Gly
	130				135						140				
Ser	Ile	Ala	Glu	Arg	Thr	Gln	Phe	Val	Ala	Tyr	Leu	Ile	Tyr	Ser	Ser
145					150					155					160
Phe	Leu	Thr	Gly	Phe	Val	Tyr	Pro	Val	Val	Ser	His	Trp	Phe	Trp	Ser
			165						170					175	
Val	Asp	Gly	Trp	Ala	Ser	Pro	Phe	Arg	Thr	Asp	Gly	Asp	Leu	Leu	Phe
		180						185					190		
Ser	Thr	Gly	Ala	Ile	Asp	Phe	Ala	Gly	Ser	Gly	Val	Val	His	Met	Val
	195						200					205			
Gly	Gly	Ile	Ala	Gly	Leu	Trp	Gly	Ala	Leu	Ile	Glu	Gly	Pro	Arg	Leu
	210				215						220				
Gly	Arg	Phe	Asp	Asn	Gly	Gly	Arg	Ala	Ile	Ala	Leu	Arg	Gly	His	Ser
225				230						235					240
Ala	Ser	Leu	Val	Val	Leu	Gly	Thr	Phe	Leu	Leu	Trp	Phe	Gly	Trp	Tyr
			245						250					255	
Gly	Phe	Asn	Pro	Gly	Ser	Phe	Asn	Lys	Ile	Leu	Val	Thr	Tyr	Glu	Thr
		260						265					270		
Gly	Thr	Tyr	Asn	Gly	Gln	Trp	Ser	Ala	Val	Gly	Arg	Thr	Ala	Val	Thr
	275						280					285			
Thr	Thr	Leu	Ala	Gly	Cys	Thr	Ala	Ala	Leu	Thr	Thr	Leu	Phe	Gly	Lys
	290				295						300				
Arg	Leu	Leu	Ser	Gly	His	Trp	Asn	Val	Thr	Asp	Val	Cys	Asn	Gly	Leu
305					310					315					320
Leu	Gly	Gly	Phe	Ala	Ala	Ile	Thr	Gly	Gly	Cys	Ser	Val	Val	Glu	Pro
			325						330					335	
Trp	Ala	Ala	Ile	Ile	Cys	Gly	Phe	Val	Ala	Ala	Leu	Val	Leu	Leu	Gly
		340						345					350		
Cys	Asn	Lys	Leu	Ala	Glu	Lys	Leu	Lys	Tyr	Asp	Asp	Pro	Leu	Glu	Ala
	355						360					365			
Ala	Gln	Leu	His	Gly	Gly	Cys	Gly	Ala	Trp	Gly	Leu	Ile	Phe	Thr	Ala
	370					375					380				
Leu	Phe	Ala	Gln	Glu	Lys	Tyr	Leu	Asn	Gln	Ile	Tyr	Gly	Asn	Lys	Pro
385					390					395					400
Gly	Arg	Pro	His	Gly	Leu	Phe	Met	Gly	Gly	Gly	Gly	Lys	Leu	Leu	Gly
			405						410					415	
Ala	Gln	Leu	Ile	Gln	Ile	Ile	Val	Ile	Thr	Gly	Trp	Val	Ser	Ala	Thr
		420						425					430		
Met	Gly	Thr	Leu	Phe	Phe	Ile	Leu	Lys	Lys	Met	Lys	Leu	Leu	Arg	Ile
	435						440					445			
Ser	Ser	Glu	Asp	Glu	Met	Ala	Gly	Met	Asp	Met	Thr	Arg	His	Gly	Gly

-continued

450					455					460					
Phe	Ala	Tyr	Met	Tyr	Phe	Asp	Asp	Asp	Glu	Ser	His	Lys	Ala	Ile	Gln
465					470					475					480
Leu	Arg	Arg	Val	Glu	Pro	Arg	Ser	Pro	Ser	Pro	Ser	Gly	Ala	Asn	Thr
				485					490					495	
Thr	Pro	Thr	Pro	Val											
			500												

1. DNA sequences that contain the coding region of a plant ammonium transporter, characterised in that the information contained in the nucleotide sequence when integrated in a plant genome makes possible under control of a promoter in sense orientation, the expression of a translatable mRNA, which leads to the synthesis of an ammonium transporter in transgenic plants.

2. DNA sequences, that contain the coding region of a plant ammonium transporter, characterised in that the information contained in the nucleotide sequence when integrated in a plant genome makes possible under control of a promoter in anti-sense orientation, the expression of a non-translatable mRNA, which prevents the synthesis of an endogenous ammonium transporter in transgenic plants.

3. A DNA sequence according to claim 1 or 2, characterised in that, it contains the following nucleotide sequence (Seq-ID No 1):

TTGTTCTGTAAACTCTCAAC																				20
ATG	TCT	TGC	TGG	GCC	ACC	GAT	CTC	GCC	GTC	CTG	TTG	GGT	CCT	AAT	65					
Net	Ser	Gys	Ser	Ala	Thr	Asp	Leu	Ala	Val	Leu	Leu	Gly	Pro	Asn						
GCC	ACG	GCG	GCG	GCC	AAC	TAC	ATA	TGT	GGC	CAG	CTA	GGC	GAC	GTC	110					
Ala	Thr	Ala	Ala	Ala	Asn	Tyr	Ile	Cys	Gly	Gln	Leu	Gly	Asp	Val						
AAC	AAC	AAA	TTC	ATC	GAC	ACC	GCT	TTC	GCT	ATA	GAC	AAC	ACT	TAC	155					
Asn	Asn	Lys	Phe	Ile	Asp	Thr	Ala	Phe	Ala	Ile	Asp	Asn	Thr	Tyr						
CTC	CTC	TTC	TCC	GCC	TAC	CTT	GTC	TTC	TCT	ATG	CAG	CTT	GGC	TTC	200					
Leu	Leu	Phe	Ser	Ala	Tyr	Leu	Val	Phe	Ser	Met	GIn	Leu	Gly	Phe						
GCT	ATG	CTC	TGT	GCC	CGT	TCC	GTG	AGA	GCC	AAG	AAT	ACT	ATG	AAC	245					
Ala	Met	Leu	Cys	Ala	Gly	Ser	Val	Arg	Ala	Lys	Asn	Thr	Met	Asn						
ATC	ATG	CTT	ACC	AAC	GTC	CTT	GAC	GCT	GCA	GCC	GGT	GGT	CTC	TTC	290					
Ile	Met	Leu	Thr	Asn	Val	Leu	Asp	Ala	Ala	Ala	Gly	Gly	Leu	Phe						
TAT	TAT	CTG	TTT	GGC	TAC	GCC	TTT	GCC	TTT	GGA	TCT	CCG	TCC	AAT	335					
Tyr	Tyr	Leu	Phe	Gly	Tyr	Ala	Phe	Ala	Phe	Gly	Ser	Pro	Ser	Asn						
GGT	TTC	ATC	GGT	AAA	CAC	TAC	TTT	GGT	CTC	AAA	GAC	ATC	CCC	ACG	380					
Gly	Phe	Ile	Gly	Lys	His	Tyr	Phe	Gly	Leu	Lys	Asp	Ile	Pro	Thr						
GCC	TCT	GCT	GAC	TAC	TCC	AAC	TTT	CTC	TAC	CAA	TGG	GCC	TTT	GCA	425					
Ala	Ser	Ala	Asp	Tyr	Ser	Asn	Phe	Leu	Tyr	Gln	Trp	Ala	Phe	Ala						
ATC	GCT	GCG	GCT	GGA	ATC	ACA	AGT	GGC	TCG	ATC	GCT	GAA	CGG	ACA	470					
Ile	Ala	Ala	Ala	Gly	Ile	Thr	Ser	Gly	Ser	Ile	Ala	Glu	Arg	Thr						
CAG	TTC	GTG	GCT	TAC	CTA	ATC	TAT	TCC	TCT	TTC	TTA	ACC	GGG	TTT	515					
Gln	Phe	Val	Ala	Tyr	Leu	Ile	Tyr	Ser	Ser	Phe	Leu	Thr	Gly	Phe						
GTT	TAC	CCG	GTC	GTC	TCT	CAC	TGG	TTC	TGG	TCA	GTT	GAT	GGA	TGG	560					
Val	Tyr	Pro	Val	Val	Ser	His	Trp	Phe	Trp	Ser	Val	Asp	Gly	Trp						
GCC	AGC	CCG	TTC	CGT	ACC	GAT	GGA	GAT	TTG	CTT	TTC	AGC	ACC	GGA	605					
Ala	Ser	Pro	Phe	Arg	Thr	Asp	Gly	Asp	Leu	Leu	Phe	Ser	Thr	Gly						
GCG	ATA	GAT	TTC	GCT	GGG	TCC	GGT	GTT	GTT	CAT	ATG	GTC	GGA	GGT	650					
Ala	Ile	Asp	Phe	Ala	Gly	Ser	Gly	Val	Val	His	Met	Val	Gly	Gly						
ATC	GCT	GGA	CTC	TGG	GGT	GCG	CTC	ATC	GAA	GGT	CCA	CGA	CTT	GGC	695					
Ile	Ala	Gly	Leu	Trp	Gly	Ala	Leu	Ile	Glu	Gly	Pro	Arg	Leu	Gly						
CGG	TTC	GAT	AAC	GGA	GGC	CGT	GCC	ATC	GCT	CTT	CGT	GGC	CAC	TCG	740					

-continued

Arg Phe Asp Asn Gly Gly Arg Ala Ile Ala Leu Arg Gly His Ser

GCG TCA CTT GTT GTC CTT GGA ACA TTC CTC CTC TGG TTT GGA TGG 785
Ala Ser Leu Val Val Leu Gly Thr Phe Leu Leu Trp Phe Gly Trp

TAC GGA TTT AAC CCC GGT TCC TTC AAC AAG ATC CTA GTC ACG TAC 830
Tyr Gly Phe Asn Pro Gly Ser Phe Asn Lys Ile Leu Val Thr Tyr

GAG ACA GGC ACA TAC AAC GGC CAG TGG AGC GCG GTC GGA CGG ACA 875
Glu Thr Gly Thr Tyr Asn Gly Gln Trp Ser Ala Val Gly Arg Thr

GCT GTC ACA ACA ACG TTA GCT GGC TGC ACC GCG GCG CTG ACA ACC 920
Ala Val Thr Thr Leu Leu Ala Gly Cys Thr Ala Ala Leu Thr Thr

CTA TTT GGG AAA CGT CTA CTC TCG GGA CAT TGG AAC GTC ACT GAT 965
Leu Phe Gly Lys Arg Leu Leu Ser Gly His Trp Asn Val Thr Asp

GTA TGC AAC GGC CTC CTC GGA GGG TTT GCA GCC ATA ACT GGT GGC 1010
Val Cys Asn Gly Leu Leu Gly Gly Phe Ala Ala Ile Thr Gly Gly

TGC TCT GTC GTT GAG CCA TGG GCT GCG ATC ATC TGC GGG TTC GTG 1055
Cys Ser Val Val Glu Pro Trp Ala Ala Ile Ile Cys Gly Phe Val

GCG GCC CTA GTC CTC CTC GGA TGC AAC AAG CTC GCT GAG AAG CTC 1100
Ala Ala Leu Val Leu Leu Gly Cys Asn Lys Leu Ala Glu Lys Leu

AAA TAC GAC GAC CCT CTT GAG GCA GCA CAA CTA CAC GGT GGT TGC 1145
Lys Tyr Asp Asp Pro Leu Glu Ala Ala Gln Leu His Gly Gly Cys

GCT GCG TGG GGA CTA ATA TTC ACG GCT CTC TTC GCT CAA GAA AAG 1190
Gly Ala Trp Gly Leu Ile Phe Thr Ala Leu Phe Ala Gln Glu Lys

TAC TTG AAC CAG ATT TAC GGC AAC AAA CCC GGA AGG CCA CAC GGT
1235
Tyr Leu Asn Gln Ile Tyr Gly Asn Lys Pro Gly Arg Pro His Gly

TTG TTT ATG GGC GGT GGA GGA AAA CTA CTT GGA GCT CAG CTG ATT 1280
Leu Phe Met Gly Gly Gly Gly Lys Leu Leu Gly Ala Gln Leu Ile

CAG ATC ATT GTG ATC ACG GGT TGG GTA ACT GCG ACC ATG GGG ACA 1325
Gln Ile Ile Val Ile Thr Gly Trp Val Ser Ala Thr Met Gly Thr

CTT TTC TTC ATC CTC AAG AAA ATG AAA TTG TTG CGG ATA TCG TCC 1370
Leu Phe Phe Ile Leu Lys Lys Met Lys Leu Leu Arg Ile Ser Ser

GAG GAT GAG ATG GCC GGT ATG GAT ATG ACC AGG CAC GGT GGT TTT 1415
Glu Asp Glu Met Ala Gly Met Asp Met Thr Arg His Gly Gly Phe

GCT TAT ATG TAC TTT GAT GAT CAT GAG TCT CAC AAA GCC ATT CAG 1460
Ala Tyr Met Tyr Phe Asp Asp Asp Glu Ser His Lys Ala Ile Gln

CTT AGG AGA GTT GAG CCA CGA TCT CCT TCT CCT TCT GGT GCT AAT 1505
Leu Arg Arg Val Glu Pro Arg Ser Pro Ser Pro Ser Gly Ala Asn

ACT ACA CCT ACT CCG GTT TGATTTGGAT TTTTACTTTT ATTCTCTATT 1553
Thr Thr Pro Thr Pro Val

TTCTAGAGTA TTATTTTAAA TGATGTTTTG TGATACTTAA ATATTGTTTT 1603
GGATATTTTT TTGGCATTTC AGTAATGTTT TAGATGTACA GTTTCATGGG 1653

GTTGTGATGA TAATATCTAT GTGGTCATTT GTGTTCTCTT TGGAGTTAAA 1703
AAAAAAAAAA AAAAAAAAAA AAAAAAAAAA AAAAAAAAAA AAAAA 1748

4. A process for the identification and isolation of DNA sequences from plants that code for ammonium transporters, characterised in that it includes the following steps:

- transformation of a yeast strain, which cannot grow in media which contain ammonium as the only nitrogen source, with a plant cDNA-library using suitable expression vectors,
- selection and propagation of transformands, which after expression of plant cDNA-sequences, can grow in media which contain ammonium as the only nitrogen source, and

c) isolation of the expression vectors, which carry a plant cDNA insert from the selected transformand.

5. DNA sequences from plants, that code for a protein with the biological activity of an ammonium transporter, obtainable by a process according to claim 4.

6. Plasmids, containing DNA sequences according to any one of claims 1-3.

7. Plasmid p35S-MEP-a (DSM 8651).

8. Plasmid p35S-a-MEP-a (DSM 8652).

9. Use of the plasmid according to any one of claims 6-8 or derivatives or parts thereof for the transformation of pro- and eukaryotic cells.

10. Transgenic plants, in which the DNA sequences of the invention are introduced as a constituent of a recombinant DNA molecule, in which this recombinant DNA molecule is stably integrated into the genome, and in whose cells based on the presence of these sequences there is achieved either a synthesis of an additional plant ammonium transporter whereby these cells can take up larger amounts of ammonium in comparison with untransformed plants or there is achieved an inhibition of the synthesis of endogenous ammonium transporters whereby such cells show a reduced uptake of ammonium in comparison with untransformed cells

11. Yeasts, containing DNA sequences according to any one of claims 1-3.

12. Bacteria, containing DNA sequences according to any one of claims 1-3.

13. Plant cells, in which the DNA sequences of the invention are introduced as a constituent of a recombinant DNA molecule, in which this recombinant DNA molecule is stably integrated into the genome, and in whose cells based on the presence of these sequences there is achieved either a synthesis of an additional plant ammonium transporter whereby these cells can take up larger amounts of ammonium in comparison with untransformed plants or there is achieved an inhibition of the synthesis of endogenous ammonium transporters whereby such cells show a reduced uptake of ammonium in comparison with untransformed cells.

14. Use of plant cells according to claim 13 for the regeneration of whole plants.

15. Use of the DNA sequences according to any one of claims 1-3, for the preparation of derivatives with changed specificity of the ammonium transporter.

16. Use of the DNA sequences according to any one of claims 1-3, for the isolation of homologous sequences from the genome of bacteria, fungi and transgenic plants.

17. Use of the DNA sequences according to claims 1, 3 or 5, in combination with a transcription promoter in sense orientation, for the expression of a translatable mRNA, which makes possible the synthesis of an ammonium transporter in pro- and eukaryotic cells.

18. Use of the DNA sequences according to claims 2, 3 or 5, in combination with a transcription promoter in anti-sense orientation, for the expression of a non-translatable mRNA, which prevents the synthesis of an endogenous ammonium transporter in pro- and eukaryotic cells.

19. Use of the DNA sequences according to any one of claims 1-3 or 5 for the preparation of transgenic plants with changed nitrogen metabolism.

20. Use of the DNA sequences according to any one of claims 1, 3 or 5 for the preparation of transgenic crop plants, such as tobacco, potatoes, beets and maize with increased yield.

21. Use of the DNA sequences according to any one of claims 2, 3 or 5 for the preparation of transgenic crop plants, such as tobacco, potatoes, beets and maize under low input conditions.

22. Use of the DNA sequences according to any one of claims 2, 3 or 5 for the preparation of transgenic crop plants, which also grow on acid soils.

* * * * *