



(86) Date de dépôt PCT/PCT Filing Date: 2003/12/23
(87) Date publication PCT/PCT Publication Date: 2004/07/15
(45) Date de délivrance/Issue Date: 2012/07/10
(85) Entrée phase nationale/National Entry: 2005/05/05
(86) N° demande PCT/PCT Application No.: US 2003/041657
(87) N° publication PCT/PCT Publication No.: 2004/058704
(30) Priorité/Priority: 2002/12/23 (US60/436,145)


(51) Cl.Int./Int.Cl. *C07D 239/95* (2006.01),
A61K 31/517 (2006.01), *A61K 31/519* (2006.01),
A61K 31/549 (2006.01), *A61P 25/00* (2006.01),
C07D 285/24 (2006.01), *C07D 401/12* (2006.01),
C07D 471/04 (2006.01), *C07D 487/04* (2006.01),
C07D 495/04 (2006.01)

(72) Inventeurs/Inventors:
MCNAUGHTON-SMITH, GRANT ANDREW, US;
THOMAS, JAMES BARNWELL, JR., US;
AMATO, GEORGE SALVATORE, US

(73) Propriétaire/Owner:
ICAGEN, INC., US

(74) Agent: FETHERSTONHAUGH & CO.

(54) Titre : QUINAZOLINONES COMME MODULATEURS DE CANAUX POTASSIQUES
(54) Title: QUINAZOLINONES AS POTASSIUM CHANNEL MODULATORS



(57) Abrégé/Abstract:

Compounds, compositions and methods are provided which are useful in the treatment of diseases through the modulation of potassium ion flux through voltage-dependent potassium channels. More particularly, the invention provides quinazolinone, compositions and methods that are useful in the treatment of central or peripheral nervous system disorders (e.g., migraine, ataxia, Parkinson's disease, bipolar disorders, trigeminal neuralgia, spasticity, mood disorders, brain tumors, psychotic disorders, myokymia, seizures, epilepsy, hearing and vision loss, Alzheimer's disease, age-related memory loss, learning deficiencies, anxiety and motor neuron diseases) and as neuroprotective agents (e.g., to prevent stroke and the like) by modulating potassium channels associated with the onset or recurrence of the indicated conditions.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
15 July 2004 (15.07.2004)

PCT

(10) International Publication Number
WO 2004/058704 A2

(51) International Patent Classification⁷: **C07D**

(21) International Application Number:
PCT/US2003/041657

(22) International Filing Date:
23 December 2003 (23.12.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/436,145 23 December 2002 (23.12.2002) US

(71) Applicant (for all designated States except US): **ICA-GEN, INC.** [US/US]; 4222 Emperor Boulevard, Suite 350, Durham, NC 27703 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **MCNAUGHTON-SMITH, Grant, Andrew** [GB/US]; 143 Walton's Creek

Road, Morrisville, NC 27560 (US). **THOMAS, James, Barnwell, Jr.** [US/US]; 3620 Rabbit Hill Lane, Efland, NC 27243 (US). **AMATO, George, Salvatore** [US/US]; 101 Ashley Drive, Cary, NC 27512 (US).

(74) Agents: **MANN, Jeffrey, S.** et al.; Townsend and Townsend and Crew LLP, Two Embarcadero Center, 8th Floor, San Francisco, CA 94111 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),

[Continued on next page]

(54) Title: QUINAZOLINONES AS POTASSIUM CHANNEL MODULATORS

#	MOLSTRUCTURE	MW	Mass obs	#	MOLSTRUCTURE	MW	Mass obs
1		389	390	13		428	429
2		426	427	14		342	343
3		442	443	15		411	412
4		356	357	16		462	463
5		426	427	17		437	438
6		409	410	18		371	372
7		476	477	19		388	389
8		418	419	20		371	372
9		478	479	21		421	422
10		375	376	22		444	445
11		411	412	23		426	427
12		375	376	24		331	332

(57) Abstract: Compounds, compositions and methods are provided which are useful in the treatment of diseases through the modulation of potassium ion flux through voltage-dependent potassium channels. More particularly, the invention provides quinazolinone, compositions and methods that are useful in the treatment of central or peripheral nervous system disorders (*e.g.*, migraine, ataxia, Parkinson's disease, bipolar disorders, trigeminal neuralgia, spasticity, mood disorders, brain tumors, psychotic disorders, myokymia, seizures, epilepsy, hearing and vision loss, Alzheimer's disease, age-related memory loss, learning deficiencies, anxiety and motor neuron diseases) and as neuroprotective agents (*e.g.*, to prevent stroke and the like) by modulating potassium channels associated with the onset or recurrence of the indicated conditions.

WO 2004/058704 A2

WO 2004/058704 A2

Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,
ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE,
SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA,
GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

- *as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii)) for the following designations AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW, ARIPO patent (BW, GH, GM, KE, LS, MW, MZ, SD,*

SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG)

- *as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii)) for all designations*
- *of inventorship (Rule 4.17(iv)) for US only*

Published:

- *without international search report and to be republished upon receipt of that report*

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

QUINAZOLINONES AS POTASSIUM CHANNEL MODULATORS

[0001]

5

FIELD OF THE INVENTION

[0002] This invention relates to the use of certain quinazolinones as potassium channel modulators and to the treatment of diseases in which a potassium channel is implicated.

10 Additionally, this invention relates to novel compounds that are useful as potassium channel modulators.

BACKGROUND OF THE INVENTION

[0003] Ion channels are cellular proteins that regulate the flow of ions, including calcium, potassium, sodium and chloride, into and out of cells. These channels are
15 present in all human cells and affect such processes as nerve transmission, muscle contraction and cellular secretion. Among the ion channels, potassium channels are the most ubiquitous and diverse, being found in a variety of animal cells such as nervous, muscular, glandular, immune, reproductive, and epithelial tissue. These channels allow the flow of potassium in and/or out of the cell under certain conditions. For example, the
20 outward flow of potassium ions upon opening of these channels makes the interior of the cell more negative, counteracting depolarizing voltages applied to the cell. These channels are regulated, *e.g.*, by calcium sensitivity, voltage-gating, second messengers, extracellular ligands, and ATP-sensitivity.

[0004] Potassium channels are associated with a number of physiological processes,
25 including regulation of heartbeat, dilation of arteries, release of insulin, excitability of nerve cells, and regulation of renal electrolyte transport. Potassium channels are made by alpha subunits that fall into at least 8 families, based on predicted structural and functional similarities (Wei *et al.*, *Neuropharmacology* 35(7): 805-829 (1997)). Three of these families (Kv, eag-related, and KQT) share a common motif of six transmembrane

domains and are primarily gated by voltage. Two other families, CNG and SK/IK, also contain this motif but are gated by cyclic nucleotides and calcium, respectively. The three other families of potassium channel alpha subunits have distinct patterns of transmembrane domains. Slo family potassium channels, or BK channels have seven
5 transmembrane domains (Meera *et al.*, *Proc. Natl. Acad. Sci. U.S.A.* **94**(25): 14066-71 (1997)) and are gated by both voltage and calcium or pH (Schreiber *et al.*, *J. Biol. Chem.* **273**: 3509-16 (1998)). Another family, the inward rectifier potassium channels (Kir), belongs to a structural family containing two transmembrane domains, and an eighth functionally diverse family (TP, or "two-pore") contains two tandem repeats of this
10 inward rectifier motif.

[0005] Potassium channels are typically formed by four alpha subunits, and can be homomeric (made of identical alpha subunits) or heteromeric (made of two or more distinct types of alpha subunits). In addition, potassium channels made from Kv, KQT and Slo or BK subunits have often been found to contain additional, structurally distinct
15 auxiliary, or beta, subunits. These subunits do not form potassium channels themselves, but instead they act as auxiliary subunits to modify the functional properties of channels formed by alpha subunits. For example, the Kv beta subunits are cytoplasmic and are known to increase the surface expression of Kv channels and/or modify inactivation kinetics of the channel (Heinemann *et al.*, *J. Physiol.* **493**: 625-633 (1996); Shi *et al.*,
20 *Neuron* **16**(4): 843-852 (1996)). In another example, the KQT family beta subunit, minK, primarily changes activation kinetics (Sanguinetti *et al.*, *Nature* **384**: 80-83 (1996)).

[0006] Slo or BK potassium channels are large conductance potassium channels found in a wide variety of tissues, both in the central nervous system and periphery. They play a key role in the regulation of processes such as neuronal integration, muscular
25 contraction and hormone secretion. They may also be involved in processes such as lymphocyte differentiation and cell proliferation, spermatocyte differentiation and sperm motility. Three alpha subunits of the Slo family have been cloned, *i.e.*, Slo1, Slo2, and Slo3 (Butler *et al.*, *Science* **261**: 221-224 (1993); Schreiber *et al.*, *J. Biol. Chem.*, **273**:
3509-16 (1998); and Joiner *et al.*, *Nature Neurosci.* **1**: 462-469 (1998)). These Slo family
30 members have been shown to be voltage and/or calcium gated, and/or regulated by intracellular pH.

[0007] Certain members of the Kv family of potassium channels were recently renamed (see, Biervert, *et al.*, *Science* **279**: 403-406 (1998)). KvLQT1 was re-named KCNQ1, and the KvLQT1-related channels (KvLR1 and KvLR2) were renamed KCNQ2 and KCNQ3, respectively. More recently, additional members of the KCNQ subfamily were identified.

5 For example, KCNQ4 was identified as a channel expressed in sensory outer hair cells (Kubisch, *et al.*, *Cell* **96**(3): 437-446 (1999)). KCNQ5 (Kananura *et al.*, *Neuroreport* **11**(9):2063 (2000)), KCNQ 2/3 (Main *et al.*, *Mol. Pharmacol.* **58**: 253-62 (2000), and KCNQ 3/5 (Wickenden *et al.*, *Br. J. Pharma* **132**: 381 (2001)) have also recently been described.


10 [0008] KCNQ2 and KCNQ3 have been shown to be nervous system-specific potassium channels associated with benign familial neonatal convulsions ("BFNC"), a class of idiopathic generalized epilepsy (see, Leppert, *et al.*, *Nature* **337**: 647-648 (1989)). These channels have been linked to M-current channels (see, Wang, *et al.*, *Science* **282**: 1890-1893 (1998)). The discovery and characterization of these channels and currents provides
15 useful insights into how these voltage dependent (Kv) potassium channels function in different environments, and how they respond to various activation mechanisms. Such information has now led to the identification of modulators of KCNQ2 and KCNQ3 potassium channels or the M-current, and the use of such modulators as therapeutic agents. The modulators are the subject of the present invention.

20

SUMMARY OF THE INVENTION

[0009] The present invention provides quinazolinones and pharmaceutically acceptable salts thereof ("compounds of the invention"), which are useful in the treatment of diseases through the modulation of potassium ion flux through voltage-dependent potassium channels.

25 [0010] In one aspect, the present invention provides compounds of the formula:



(I).

[0011] In Formula I, the symbol A represents a ring structure, e.g., a five- or six-membered substituted or unsubstituted aryl, five- and six-membered substituted or

unsubstituted heteroaryl, substituted or unsubstituted C₄-C₈ cycloalkyl, and substituted or unsubstituted 5-8 membered heterocyclyl ring system. X represents a group such as CO, CS or SO₂.

[0012] W is a member selected from N and CR³. The symbol R³ represents H, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, substituted or unsubstituted C₃-C₈ cycloalkyl, substituted or unsubstituted 5-7 membered heterocyclyl, or substituted or unsubstituted C₁-C₈ alkyl.

[0013] Z represents a bond, -CH₂-, -CHF-, -CF₂-, -CH=CH- or -N(R⁴)(CR^{4a}R^{4b})_s-, in which R⁴ represents H or a substituted or unsubstituted C₁-C₅ alkyl group. The symbols R^{4a} and R^{4b} represent groups that are independently selected from H, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, substituted or unsubstituted C₃-C₈ cycloalkyl, substituted or unsubstituted 5-7 membered heterocyclyl, or substituted or unsubstituted C₁-C₈ alkyl. R¹ is substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, substituted or unsubstituted C₃-C₈ cycloalkyl, substituted or unsubstituted 5-7 membered heterocyclyl, or substituted or unsubstituted C₁-C₈ alkyl. The index "s" is an integer from 1 to 3. Moreover, when "s" is greater than 1, each R^{4a} and R^{4b} are independently selected.

[0014] The symbol Y represents S(O)_n, in which the index "n" is an integer from 0-2. R² is CF₃, substituted or unsubstituted C₁-C₈ alkyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, substituted or unsubstituted C₃-C₈ cycloalkyl, or substituted or unsubstituted 3-7-membered heterocyclyl group.


[0015] In another aspect, the present invention provides pharmaceutical compositions comprising a pharmaceutically acceptable excipient and a compound of the formula provided above.

[0016] In yet another aspect, the present invention provides a method for increasing flow through voltage dependent potassium channels in a cell, comprising contacting the cell with a compound of the formula provided above in an amount sufficient to open the potassium channels.

[0017] In still another aspect, the present invention provides a method for treating a central or peripheral nervous system disorder or condition through the modulation of a voltage-dependent potassium channel, the method comprising administering to a subject

in need of such treatment an effective amount of a compound of the formula provided above.

[0017a] In another aspect of the invention, the present invention provides a compound having the formula:



in which A is a member selected from the group consisting of five- and six-membered substituted or unsubstituted aryl, five- and six-membered substituted or unsubstituted heteroaryl, substituted or unsubstituted C₄-C₈ cycloalkyl, and substituted or unsubstituted 5-8 membered heterocyclyl; X is a member selected from the group consisting of CO, CS and SO₂; Z is a member selected from the group consisting of -CH₂-, -CHF-, -CF₂-, -CH=CH- and -N(R⁴)(CR^{4a}R^{4b})_s- wherein R⁴ is a member selected from the group consisting of H and a substituted or unsubstituted C₁-C₅ alkyl group; R^{4a} and R^{4b} are members independently selected from the group consisting of H, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, substituted or unsubstituted C₃-C₈ cycloalkyl, substituted or unsubstituted 5-7 membered heterocyclyl, and substituted or unsubstituted C₁-C₈ alkyl; and s is an integer from 1 to 3; Y is S(O)_n, wherein n is an integer from 0-2; R¹ is a member selected from the group consisting of substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, substituted or unsubstituted C₃-C₈ cycloalkyl, substituted or unsubstituted 5-7 membered heterocyclyl, and substituted or unsubstituted C₁-C₈ alkyl; and R² is a member selected from the group consisting of CF₃, substituted or unsubstituted C₁-C₈ alkyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, substituted or unsubstituted C₃-C₈ cycloalkyl, and substituted or unsubstituted 3-7-membered heterocyclyl; provided that when A is phenyl, Z is -CH₂- and R¹ is phenyl, substituted phenyl or heteroaryl, then R² is other than a benzyl, substituted benzyl, alkylheteroaryl, alkylheterocyclyl or cyanomethyl group.

[0017b] The substituents for the alkyl, cycloalkyl, heterocyclyl and heteroalkyl groups are selected from the group consisting of -OR', =O, =NR', =N-OR', -NR'R'', -SR', -halogen, -SiR'R''R''', -OC(O)R', -C(O)R', -CO₂R', -CONR'R'', -OC(O)NR'R'', -NRC(O)R', -NR-

C(O)NR'R'', -NRC(O₂)R', -NR-C(NR'R'')=NR''', -S(O)R', -S(O)₂R', -S(O)₂NR'R'', -
 NRSO₂R', -CN and -NO₂ in a number ranging from zero to (2m'+1), where m' is the total
 number of carbon atoms in such group and R, R', R'' and R''' are each independentl selected
 from the group consisting of hydrogen, unsubstituted heteroalkyl, unsubstituted aryl,
 5 unsubstituted alkyl, alkoxy or thioalkoxy groups, or arylalkyl groups.

[0017c] The substituents for the aryl and heteroaryl groups are selected from the group
 consisting of -halogen, -OR', =O, =NR', =N-OR', -NR'R'', -SR', -SiR'R''R''', -OC(O)R', -
 C(O)R', -CO₂R', -CONR'R'', -OC(O)NR'R'', -NRC(O)R', -NR-C(O)NR'R'', -NRC(O)₂R',
 10 -NR-C(NR'R'')=NR''', -S(O)R', -S(O)₂R', -S(O)₂NR'R'', -NRSO₂R', -CN, -NO₂, -R', -N₃,
 -CH(Ph)₂, fluoro (C₁-C₄)alkoxy and fluoro (C₁-C₄)alkyl in a number ranging from zero to
 the total number of open valences on the aromatic ring system, where R, R', R'' and R''' are
 each independently selected from the group consisting of hydrogen, substituted or
 unsubstituted C₁-C₈ alkyl and heteroalkyl, substituted or unsubstituted C₃-C₈ cycloalkyl,
 15 unsubstituted aryl and heteroaryl, (unsubstituted aryl)-(C₁-C₄) alkyl, and (unsubstituted
 aryl)oxy-(C₁-C₄)alkyl, and where R' and R'', together with the nitrogen atom to which they
 are attached, are optionally joined to form a 5- to 7-membered ring.





[0018] Other objects and advantages of the present invention will be apparent from the
 20 detail description that follows.





BRIEF DESCRIPTION OF THE CHEMICAL STRUCTURES





[0019] Figures 1A-1G below display structures of representative compounds of the
 invention.





25





FIGURE 1A

#	MOLSTRUCTURE	MW	Mass Obs
1		389	390
2		426	427
3		442	443
4		356	357

410	409		6
419	418		8
427	477		
426	476		5
			7

9		478	479
11		411	412
10		375	376
12		375	376

343	342		14
463	462		16
429	428		13
412	411		15

17		437	438		18		371	372
19		388	389		20		371	372





















445	444		332
22	24		331
422	427		
421	426		23
21	23		





FIGURE 1B

#	MOLSTRUCTURE	MW	Mass Obs
26		429	430
28		361	362
25		438	439
27		411	412

318	317		390
	389		
30	32		
361	425		
360	424		
29	31		

33		371	372		34		321	322
35		320	321		36		277	278

37		325	326		38		454	455
39		472	473	40	40		343	344

41		361	362		42		343	344
43		339	340		44		353	354













45		375	376		46		389	390
47		357	358		48		371	372





FIGURE 1C

#	MOLSTRUCTURE	MW	Mass Obs
49		383	384
50		375	376
51		393	394
52		407	408

53	421	422	54	361	362
					
55	375	376	56	375	376
					

57	379	380		58		379	380
59	389	390		60		361	362

61	361	362		62		411	412
63	411	412		64		343	344

65		357	358		66		361	362
67		375	376		68		343	344





















69		357	358	
70		418	419	
71		389	390	
72		343	344	





FIGURE 1D



#	MOLSTRUCTURE	MW	Mass Obs	#	MOLSTRUCTURE	MW	Mass Obs
73		393	394	74		379	380
75		371	372	76		375	376

77		389	390	
79		401	402	
		78		
		383	384	
		416	417	
		80		

81		427	428		82		409	410
83		427	428		84		455	456

85		361	362		86		357	358	
87		437	438		88		409	410	

89		433	434		90		345	346
91		407	408		92		391	392

93	373	374	94		391	392
95	371	372	96		441	442

[0109] Unless otherwise specified, all solvents (HPLC grade) and reagents were purchased from suppliers and used without further purification. Analytical thin layer chromatography (TLC) was performed on Whatman Inc. 60 silica gel plates (0.25 mm thickness). Compounds were visualized under UV lamp (254 nm) or by developing with KMnO₄/KOH, ninhydrin or Hanessian's solution. Flash chromatography was done using silica gel from Selectro Scientific (particle size 32-63). ¹H NMR, ¹⁹F NMR and ¹³C NMR spectra were recorded on a Varian 300 machine at 300 MHz, 282 MHz and 75.7 MHz, respectively. Melting points were recorded on an Electrothermal IA9100 apparatus and were uncorrected.

10

EXAMPLE 1

1.1 General procedure for preparation of isatoic anhydrides (II)

[0110] Anhydrous pyridine (2 eq) was added to a solution of 2-aminobenzoic acid derivative (IA) (1 eq) in dry methylene chloride and acetonitrile (1:1, 40 mL/g of 2-aminobenzoic acid) at room temperature. Solid triphosgene (1/3 eq) was then added in one portion and the resulting mixture was heated at 50 °C for 2 h. The resulting solid was collected by filtration and dried in vacuo. The crude isatoic anhydrides (IIA) were typically obtained in 50 – 80% yields. Though contaminated with some pyridinium hydrochloride, the anhydrides were used in the next step without further purification.

1.2 General procedure for the preparation of acylhydrazides (IV)

20 [0111] A mixture of isatoic anhydride (IIA, 1 eq) and appropriate phenylacetylhydrazide (IIIA, 1.1 eq) were heated in glacial AcOH (2 mL/mmol) at 50 °C for 2-6 h. The resulting solution was cooled and water was added while shaking. The white precipitate was collected by filtration, washed with water and dried *in vacuo* at 50 °C for 4 h. The desired products (IV) were obtained as white solids in high purity (typically >90%) and moderate yields (typically 45-60%).

25

1.3 Representative compounds





[0112] 1.3a 2-Amino-4-fluoro-benzoic acid N'-phenylacetyl-hydrazide



Obtained as a white powder. ¹H NMR (300 MHz, d-6 DMSO) δ 3.05(2H, s), 6.30 (1H, dt, J = 8.5, 2.4 Hz), 6.45 (1H, dd, J = 12.0, 2.6 Hz), 6.72 (2H, brs), 7.18-7.34 (5H, m), 7.57 (1H, dd, J = 8.9, 6.8 Hz), 10.02 (2H, brs).





30





FIGURE 1E

#	MOLSTRUCTURE	MW	Mass Obs	#	MOLSTRUCTURE	MW	Mass Obs
97		455	456	98		427	428
99		437	438	100		371	372

101		423	424		102		423	424	
103		340	341		104		385	386	

105	425	426		106		393	394
107	393	394		108		407	408

109		371	372					
111		401	402	112		421	422	
110								
390		389						

113		393	394		114		357	358
115		371	372		116		375	376




















117	389	390			
118	329		330		
119	343	344	120	357	358
118	329		330		
120	357		358		





FIGURE 1F

#	MOLSTRUCTURE	MW	Mass Obs	#	MOLSTRUCTURE	MW	Mass Obs
121		354	355	122		343	344
123		425	426	124		407	408

428	427		126		428	427		125
390	389		128		376	375		127

129		455	456		130		455	456
131		437	438		132		361	362

133		437	438		
135		409	410		
134		455	456		
136		423	424		

137		383	384		
139		455	456		
138					
428		427			
438		437			


















141		441	442		142		411	412
143		357	358		144		358	359

FIGURE 1G

#	MOLSTRUCTURE	MW	Mass Obs
145		389	390
146		407	408
147		427	428
148		355	356

149	437	438		150		372	373
							
151	361	362		152		372	373
							

153		426	427		154		372	373
155		426	427		156		369	370




157		403	404
-----	---	-----	-----

[0020] Figures 2A and 2B below are activity data for selected compounds of the invention.

FIG. 2A

Structure #	NG-108 EC50 (μ M)
40	0.24
44	0.17
45	0.19
46	0.21
47	0.38
48	0.09
61	0.23
50	0.37
51	0.19
52	0.07
53	0.1
54	0.25
55	0.31
56	0.45
57	0.23
59	0.24
60	0.38
73	0.19
74	0.81
62	0.07
63	0.47
64	0.63
65	0.85
66	0.48
67	1
68	0.46

FIG. 2B

NG-108 EC50 = 0.36 μ MNG-108 EC50 = 0.9 μ MNG-108 EC50 = 0.69 μ M

DETAILED DESCRIPTION OF THE INVENTION AND THE PREFERRED EMBODIMENTS

Abbreviations and Definitions:

[0021] The abbreviations used herein have their conventional meaning within the
5 chemical and biological arts. For example: CHO, Chinese hamster ovary; EBSS, Earl's
Balanced Salt Solution; KCNQ, potassium channel Q; KCNQ2, potassium channel Q2;
Et₃N, triethylamine; MeOH, methanol; and DMSO, dimethylsulfoxide.

[0022] "Compound of the invention," as used herein refers to a compound according to
Formulae I or II or a combination thereof, and a pharmaceutically acceptable salt of a
10 compound according to Formulae I or II or a combination thereof.

[0023] "Modulating," as used herein, refers to the ability of a compound of the
invention to activate and/or inhibit a potassium channel, preferably, a KCNQ potassium
channel.

[0024] "Opening" and "activating" are used interchangeably herein to refer to the
15 partial or full activation of a KCNQ channel by a compound of the invention, which leads
to an increase in ion flux either into or out of a cell in which a KCNQ channel is found.

[0025] Where substituent groups are specified by their conventional chemical formulae,
written from left to right, they equally encompass the chemically identical substituents
which would result from writing the structure from right to left, e.g., -CH₂O- is intended
20 to also recite -OCH₂-; -NHS(O)₂- is also intended to represent. -S(O)₂HN-, etc.

[0026] The term "alkyl," by itself or as part of another substituent, means, unless
otherwise stated, a straight or branched chain, or cyclic hydrocarbon radical, or
combination thereof, which may be fully saturated, mono- or polyunsaturated and can
include di- and multivalent radicals, having the number of carbon atoms designated (*i.e.*
25 C₁-C₁₀ means one to ten carbons). Examples of saturated hydrocarbon radicals include,
but are not limited to, groups such as methyl, ethyl, n-propyl, isopropyl, n-butyl, t-butyl,
isobutyl, sec-butyl, cyclohexyl, (cyclohexyl)methyl, cyclopropylmethyl, homologs and
isomers of, for example, n-pentyl, n-hexyl, n-heptyl, n-octyl, and the like. An unsaturated
alkyl group is one having one or more double bonds or triple bonds. Examples of
30 unsaturated alkyl groups include, but are not limited to, vinyl, 2-propenyl, crotyl, 2-

isopentenyl, 2-(butadienyl), 2,4-pentadienyl, 3-(1,4-pentadienyl), ethynyl, 1- and 3-propynyl, 3-butyryl, and the higher homologs and isomers. The term “alkyl,” unless otherwise noted, is also meant to include those derivatives of alkyl defined in more detail below, such as “heteroalkyl.” Alkyl groups which are limited to hydrocarbon groups are
 5 termed “homoalkyl”.

[0027] The term “alkylene” by itself or as part of another substituent means a divalent radical derived from an alkane, as exemplified, but not limited, by $-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2-$, and further includes those groups described below as “heteroalkylene.” Typically, an alkyl (or alkylene) group will have from 1 to 24 carbon atoms, with those groups having
 10 10 or fewer carbon atoms being preferred in the present invention. A “lower alkyl” or “lower alkylene” is a shorter chain alkyl or alkylene group, generally having eight or fewer carbon atoms.

[0028] The terms “alkoxy,” “alkylamino” and “alkylthio” (or thioalkoxy) are used in their conventional sense, and refer to those alkyl groups attached to the remainder of the
 15 molecule via an oxygen atom, an amino group, or a sulfur atom, respectively.

[0029] The term “heteroalkyl,” by itself or in combination with another term, means, unless otherwise stated, a stable straight or branched chain, or cyclic hydrocarbon radical, or combinations thereof, consisting of the stated number of carbon atoms and at least one heteroatom selected from the group consisting of O, N, Si and S, and wherein the nitrogen
 20 and sulfur atoms may optionally be oxidized and the nitrogen heteroatom may optionally be quaternized. The heteroatom(s) O, N and S and Si may be placed at any interior position of the heteroalkyl group or at the position at which the alkyl group is attached to the remainder of the molecule. Examples include, but are not limited to, $-\text{CH}_2\text{CH}_2\text{OCH}_3$, $-\text{CH}_2\text{CH}_2\text{NHCH}_3$, $-\text{CH}_2\text{CH}_2\text{N}(\text{CH}_3)\text{CH}_3$, $-\text{CH}_2\text{SCH}_2\text{CH}_3$, $-\text{CH}_2\text{CH}_2\text{S}(\text{O})\text{CH}_3$, $-\text{CH}_2\text{CH}_2\text{S}(\text{O})_2\text{CH}_3$, $-\text{CH}=\text{CHOCH}_3$, $-\text{Si}(\text{CH}_3)_3$, $-\text{CH}_2\text{CH}=\text{NOCH}_3$, and $-\text{CH}=\text{CHN}(\text{CH}_3)\text{CH}_3$. Up to two heteroatoms may be consecutive, such as, for example, $-\text{CH}_2\text{NH}\text{OCH}_3$ and $-\text{CH}_2\text{OSi}(\text{CH}_3)_3$. Similarly, the term “heteroalkylene” by itself or
 25 as part of another substituent means a divalent radical derived from heteroalkyl, as exemplified, but not limited by, $-\text{CH}_2\text{CH}_2\text{SCH}_2\text{CH}_2-$ and $-\text{CH}_2\text{SCH}_2\text{CH}_2\text{NHCH}_2-$.
 30 For heteroalkylene groups, heteroatoms can also occupy either or both of the chain termini (*e.g.*, alkyleneoxy, alkylenedioxy, alkyleneamino, alkylenediamino, and the like). Still further, for alkylene and heteroalkylene linking groups, no orientation of the linking

group is implied by the direction in which the formula of the linking group is written. For example, the formula $-C(O)_2R'$ represents both $-C(O)_2R'$ and $-R'C(O)_2-$.

[0030] The terms “cycloalkyl” and “heterocycloalkyl”, by themselves or in combination with other terms, represent, unless otherwise stated, cyclic versions of “alkyl” and “heteroalkyl”, respectively. Additionally, for heterocycloalkyl, a heteroatom can occupy the position at which the heterocycle is attached to the remainder of the molecule. Examples of cycloalkyl include, but are not limited to, cyclopentyl, cyclohexyl, 1-cyclohexenyl, 3-cyclohexenyl, cycloheptyl, and the like. Examples of heterocycloalkyl include, but are not limited to, 1-(1,2,5,6-tetrahydropyridyl), 1-piperidinyl, 2-piperidinyl, 3-piperidinyl, 4-morpholinyl, 3-morpholinyl, tetrahydrofuran-2-yl, tetrahydrofuran-3-yl, tetrahydrothien-2-yl, tetrahydrothien-3-yl, 1-piperazinyl, 2-piperazinyl, and the like.

[0031] The terms “halo” or “halogen,” by themselves or as part of another substituent, mean, unless otherwise stated, a fluorine, chlorine, bromine, or iodine atom. Additionally, terms such as “haloalkyl,” are meant to include monohaloalkyl and polyhaloalkyl. For example, the term “halo(C₁-C₄)alkyl” is meant to include, but not be limited to, trifluoromethyl, 2,2,2-trifluoroethyl, 4-chlorobutyl, 3-bromopropyl, and the like.

[0032] The term “aryl” means, unless otherwise stated, a polyunsaturated, aromatic, hydrocarbon substituent which can be a single ring or multiple rings (preferably from 1 to 3 rings) which are fused together or linked covalently. The term “heteroaryl” refers to aryl groups (or rings) that contain from one to four heteroatoms selected from N, O, and S, wherein the nitrogen and sulfur atoms are optionally oxidized, and the nitrogen atom(s) are optionally quaternized. A heteroaryl group can be attached to the remainder of the molecule through a heteroatom. Non-limiting examples of aryl and heteroaryl groups include phenyl, 1-naphthyl, 2-naphthyl, 4-biphenyl, 1-pyrrolyl, 2-pyrrolyl, 3-pyrrolyl, 3-pyrazolyl, 2-imidazolyl, 4-imidazolyl, pyrazinyl, 2-oxazolyl, 4-oxazolyl, 2-phenyl-4-oxazolyl, 5-oxazolyl, 3-isoxazolyl, 4-isoxazolyl, 5-isoxazolyl, 2-thiazolyl, 4-thiazolyl, 5-thiazolyl, 2-furyl, 3-furyl, 2-thienyl, 3-thienyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, 2-pyrimidyl, 4-pyrimidyl, 5-benzothiazolyl, purinyl, 2-benzimidazolyl, 5-indolyl, 1-isoquinolyl, 5-isoquinolyl, 2-quinoxalyl, 5-quinoxalyl, 3-quinolyl, and 6-quinolyl.

Substituents for each of the above noted aryl and heteroaryl ring systems are selected from the group of acceptable substituents described below.

[0033] For brevity, the term “aryl” when used in combination with other terms (*e.g.*, aryloxy, arylthioxy, arylalkyl) includes both aryl and heteroaryl rings as defined above.

5 Thus, the term “arylalkyl” is meant to include those radicals in which an aryl group is attached to an alkyl group (*e.g.*, benzyl, phenethyl, pyridylmethyl and the like) including those alkyl groups in which a carbon atom (*e.g.*, a methylene group) has been replaced by, for example, an oxygen atom (*e.g.*, phenoxymethyl, 2-pyridyloxymethyl, 3-(1-naphthyloxy)propyl, and the like).

10 [0034] Each of the above terms (*e.g.*, “alkyl,” “heteroalkyl,” “aryl” and “heteroaryl”) include both substituted and unsubstituted forms of the indicated radical. Preferred substituents for each type of radical are provided below.

[0035] Substituents for the alkyl and heteroalkyl radicals (including those groups often referred to as alkylene, alkenyl, heteroalkylene, heteroalkenyl, alkynyl, cycloalkyl, 15 heterocycloalkyl, cycloalkenyl, and heterocycloalkenyl) can be one or more of a variety of groups selected from, but not limited to: -OR', =O, =NR', =N-OR', -NR'R'', -SR', -halogen, -SiR'R''R''', -OC(O)R', -C(O)R', -CO₂R', -CONR'R'', -OC(O)NR'R'', -NR''C(O)R', -NR'-C(O)NR''R''', -NR''C(O)₂R', -NR-C(NR'R'')=NR''', -S(O)R', -S(O)₂R', -S(O)₂NR'R'', -NRSO₂R', -CN and -NO₂ in a number ranging from zero to 20 (2m'+1), where m' is the total number of carbon atoms in such radical. R', R'', R''' and R'''' each preferably independently refer to hydrogen, substituted or unsubstituted heteroalkyl, substituted or unsubstituted aryl, *e.g.*, aryl substituted with 1-3 halogens, substituted or unsubstituted alkyl, alkoxy or thioalkoxy groups, or arylalkyl groups.

When a compound of the invention includes more than one R group, for example, each of 25 the R groups is independently selected as are each R', R'', R''' and R'''' groups when more than one of these groups is present. When R' and R'' are attached to the same nitrogen atom, they can be combined with the nitrogen atom to form a 5-, 6-, or 7-membered ring. For example, -NR'R'' is meant to include, but not be limited to, 1-pyrrolidinyl and 4-morpholinyl. From the above discussion of substituents, one of skill in the art will 30 understand that the term “alkyl” is meant to include groups including carbon atoms bound to groups other than hydrogen groups, such as haloalkyl (*e.g.*, -CF₃ and -CH₂CF₃) and

acyl (*e.g.*, -C(O)CH₃, -C(O)CF₃, -C(O)CH₂OCH₃, and the like). These substituents are referred to herein generically as "alkyl group substituents."

[0036] Similar to the substituents described for the alkyl radical, substituents for the aryl and heteroaryl groups, referred to generically herein as "aryl group substituents," are varied and are selected from, for example: halogen, -OR', =O, =NR', =N-OR', -NR'R'', -SR', -halogen, -SiR'R''R''', -OC(O)R', -C(O)R', -CO₂R', -CONR'R'', -OC(O)NR'R'', -NR''C(O)R', -NR'-C(O)NR''R''', -NR''C(O)₂R', -NR-C(NR'R'')=NR''', -S(O)R', -S(O)₂R', -S(O)₂NR'R'', -NRSO₂R', -CN and -NO₂, -R', -N₃, -CH(Ph)₂, fluoro(C₁-C₄)alkoxy, and fluoro(C₁-C₄)alkyl, in a number ranging from zero to the total number of open valences on the aromatic ring system; and where R', R'', R''' and R'''' are preferably independently selected from hydrogen, (C₁-C₈)alkyl and heteroalkyl, unsubstituted aryl and heteroaryl, (unsubstituted aryl)-(C₁-C₄)alkyl, and (unsubstituted aryl)oxy-(C₁-C₄)alkyl. When a compound of the invention includes more than one R group, for example, each of the R groups is independently selected as are each R', R'', R''' and R'''' groups when more than one of these groups is present.

[0037] Two of the substituents on adjacent atoms of the aryl or heteroaryl ring may optionally be replaced with a substituent of the formula -T-C(O)-(CRR')_q-U-, wherein T and U are independently -NR-, -O-, -CRR'- or a single bond, and q is an integer of from 0 to 3. Alternatively, two of the substituents on adjacent atoms of the aryl or heteroaryl ring may optionally be replaced with a substituent of the formula -A-(CH₂)_r-B-, wherein A and B are independently -CRR'-, -O-, -NR-, -S-, -S(O)-, -S(O)₂-, -S(O)₂NR'- or a single bond, and r is an integer of from 1 to 4. One of the single bonds of the new ring so formed may optionally be replaced with a double bond. Alternatively, two of the substituents on adjacent atoms of the aryl or heteroaryl ring may optionally be replaced with a substituent of the formula -(CRR')_s-X-(CR''R''')_d-, where s and d are independently integers of from 0 to 3, and X is -O-, -NR'-, -S-, -S(O)-, -S(O)₂-, or -S(O)₂NR'-. The substituents R, R', R'' and R''' are preferably independently selected from hydrogen or substituted or unsubstituted (C₁-C₆)alkyl.

[0038] As used herein, the term "heteroatom" is meant to include oxygen (O), nitrogen (N), sulfur (S) and silicon (Si).

[0039] The term “pharmaceutically acceptable salts” is meant to include salts of the active compounds which are prepared with relatively nontoxic acids or bases, depending on the particular substituents found on the compounds described herein. When compounds of the present invention contain relatively acidic functionalities, base addition salts can be obtained by contacting the neutral form of such compounds with a sufficient amount of the desired base, either neat or in a suitable inert solvent. Examples of pharmaceutically acceptable base addition salts include sodium, potassium, calcium, ammonium, organic amino, or magnesium salt, or a similar salt. When compounds of the present invention contain relatively basic functionalities, acid addition salts can be obtained by contacting the neutral form of such compounds with a sufficient amount of the desired acid, either neat or in a suitable inert solvent. Examples of pharmaceutically acceptable acid addition salts include those derived from inorganic acids like hydrochloric, hydrobromic, nitric, carbonic, monohydrogencarbonic, phosphoric, monohydrogenphosphoric, dihydrogenphosphoric, sulfuric, monohydrogensulfuric, hydriodic, or phosphorous acids and the like, as well as the salts derived from relatively nontoxic organic acids like acetic, propionic, isobutyric, maleic, malonic, benzoic, succinic, suberic, fumaric, lactic, mandelic, phthalic, benzenesulfonic, p-tolylsulfonic, citric, tartaric, methanesulfonic, and the like. Also included are salts of amino acids such as arginate and the like, and salts of organic acids like glucuronic or galactunoric acids and the like (*see*, for example, Berge *et al.*, “Pharmaceutical Salts”, *Journal of Pharmaceutical Science*, 1977, 66, 1-19). Certain specific compounds of the present invention contain both basic and acidic functionalities that allow the compounds to be converted into either base or acid addition salts.

[0040] The neutral forms of the compounds are preferably regenerated by contacting the salt with a base or acid and isolating the parent compound in the conventional manner. The parent form of the compound differs from the various salt forms in certain physical properties, such as solubility in polar solvents, but otherwise the salts are equivalent to the parent form of the compound for the purposes of the present invention.

[0041] In addition to salt forms, the present invention provides compounds, which are in a prodrug form. Prodrugs of the compounds described herein are those compounds that readily undergo chemical changes under physiological conditions to provide the compounds of the present invention. Additionally, prodrugs can be converted to the

compounds of the present invention by chemical or biochemical methods in an *ex vivo* environment. For example, prodrugs can be slowly converted to the compounds of the present invention when placed in a transdermal patch reservoir with a suitable enzyme or chemical reagent.

5 [0042] Certain compounds of the present invention can exist in unsolvated forms as well as solvated forms, including hydrated forms. In general, the solvated forms are equivalent to unsolvated forms and are encompassed within the scope of the present invention. Certain compounds of the present invention may exist in multiple crystalline or amorphous forms. In general, all physical forms are equivalent for the uses
10 contemplated by the present invention and are intended to be within the scope of the present invention.

[0043] Certain compounds of the present invention possess asymmetric carbon atoms (optical centers) or double bonds; the racemates, diastereomers, geometric isomers and individual isomers are encompassed within the scope of the present invention.

15 [0044] The compounds of the present invention may also contain unnatural proportions of atomic isotopes at one or more of the atoms that constitute such compounds. For example, the compounds may be radiolabeled with radioactive isotopes, such as for example tritium (^3H), iodine-125 (^{125}I) or carbon-14 (^{14}C). All isotopic variations of the compounds of the present invention, whether radioactive or not, are intended to be
20 encompassed within the scope of the present invention.

Introduction

[0045] The present invention provides compounds which, *inter alia*, are useful in the treatment of diseases through the modulation of potassium ion flux through voltage-dependent potassium channels. More particularly, the invention provides compounds,
25 compositions and methods that are useful in the treatment of central or peripheral nervous system disorders (e.g., migraine, ataxia, Parkinson's disease, bipolar disorders, trigeminal neuralgia, spasticity, mood disorders, brain tumors, psychotic disorders, myokymia, seizures, epilepsy, hearing and vision loss, Alzheimer's disease, age-related memory loss, learning deficiencies, anxiety and motor neuron diseases), and as neuroprotective agents
30 (e.g., to prevent stroke, retinal degeneration and the like). Compounds of the invention have use as agents for treating convulsive states, for example that following grand mal,

petit mal, psychomotor epilepsy or focal seizure. The compounds of the invention are also useful in treating disease states such as gastroesophageal reflux disorder and gastrointestinal hypomotility disorders.

[0046] Moreover, compounds of the invention are useful in the treatment of pain, for example, neuropathic pain, diabetic pain, inflammatory pain, cancer pain, migraine pain, and musculoskeletal pain. The compounds are also useful to treat conditions, which may themselves be the origin of pain, for example, inflammatory conditions, including arthritic conditions (*e.g.*, rheumatoid arthritis, rheumatoid spondylitis, osteoarthritis and gouty arthritis) and non-articular inflammatory conditions (*e.g.*, herniated, ruptured and prolapsed disc syndrome, bursitis, tendonitis, tenosynovitis, fibromyalgia syndrome, and other conditions associated with ligamentous sprain and regional musculoskeletal strain). Particularly preferred compounds of the invention are less ulcerogenic than other anti-inflammatory agents (*e.g.*, ibuprofen, naproxen and aspirin). Furthermore, the compounds of the invention are useful in treating conditions and pain associated with abnormally raised skeletal muscle tone.

[0047] The compounds of the invention are also of use in treating anxiety (*e.g.* anxiety disorders). Anxiety disorders are defined in the Diagnostic and Statistical Manual of Mental Disorders (Third Edition-revised 1987, published by the American Psychiatric Association, Washington, D.C., *see*, pages 235 to 253), as psychiatric conditions having symptoms of anxiety and avoidance behavior as characteristic features. Included amongst such disorders are generalized anxiety disorder, simple phobia and panic disorder.

[0048] Anxiety also occurs as a symptom associated with other psychiatric disorders, for example, obsessive compulsive disorder, post-traumatic stress disorder, schizophrenia, mood disorders and major depressive disorders, and with organic clinical conditions including, but not limited to, Parkinson's disease, multiple sclerosis, and other physically incapacitating disorders.

[0049] In view of the above-noted discovery, the present invention provides compounds, compositions, and methods for increasing ion flux in voltage-dependent potassium channels, particularly those channels responsible for the M-current. As used herein, the term "M-current," "channels responsible for the M-current" and the like, refers to a slowly activating, non-inactivating, slowly deactivating voltage-gated K⁺ channel.


M-current is active at voltages close to the threshold for action potential generation in a wide variety of neuronal cells, and thus, is an important regulator of neuronal excitability.

[0050] Recently, members of the voltage-dependent potassium channel family were shown to be directly involved in diseases of the central or peripheral nervous system. The quinazolinones provided herein are now shown to act as potassium channel modulators, particularly openers, for KCNQ2 and KCNQ3, KCNQ4, and KCNQ5 as well as the heteromultimer channels such as KCNQ2/3, KCNQ3/5 or the M-current.

Description of the Embodiments

I. MODULATORS OF VOLTAGE-DEPENDENT POTASSIUM CHANNELS

[0051] The present invention provides a novel class of potassium ion channel modulators, particularly effective at modulating KCNQ, according to Formula I:



[0052] In Formula I, the symbol A represents a ring structure, e.g., a five- or six-membered substituted or unsubstituted aryl, five- and six-membered substituted or unsubstituted heteroaryl, substituted or unsubstituted C₄-C₈ cycloalkyl, and substituted or unsubstituted 5-8 membered heterocyclyl ring system. X represents a group such as CO, CS or SO₂.

[0053] Z represents a bond, -CH₂-, -CHF-, -CF₂-, -CH=CH- or -N(R⁴)(CR^{4a}R^{4b})_s-, in which R⁴ represents H or a substituted or unsubstituted C₁-C₅ alkyl group. The symbols R^{4a} and R^{4b} represent groups that are independently selected from H, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, substituted or unsubstituted C₃-C₈ cycloalkyl, substituted or unsubstituted 5-7 membered heterocyclyl, or substituted or unsubstituted C₁-C₈ alkyl. R¹ is substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, substituted or unsubstituted C₃-C₈ cycloalkyl, substituted or unsubstituted 5-7 membered heterocyclyl, or substituted or unsubstituted C₁-C₈ alkyl. The index "s" is an integer from 1 to 3. Moreover, when "s" is greater than 1, each R^{4a} and R^{4b} are independently selected.

[0054] W is selected from N and CR^3 , wherein R^3 is H, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, substituted or unsubstituted C_3 - C_8 cycloalkyl, substituted or unsubstituted 5-7 membered heterocyclyl, or substituted or unsubstituted C_1 - C_8 alkyl.

5 [0055] The symbol Y represents $S(O)_n$, in which the index "n" is an integer from 0-2. R^2 is CF_3 , substituted or unsubstituted C_1 - C_8 alkyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, substituted or unsubstituted C_3 - C_8 cycloalkyl, or substituted or unsubstituted 3-7-membered heterocyclyl.


[0056] In an exemplary embodiment according to Formula I, the invention provides a
10 compound in which, when A is phenyl, Z is a bond, $-CH_2-$ or $-NH-$, and R^1 is phenyl, substituted phenyl or heteroaryl, then R^2 is other than a benzyl, substituted benzyl, alkylheteroaryl, alkylheterocyclyl or cyanomethyl group.

[0057] In another exemplary embodiment, A is substituted or unsubstituted aryl, e.g., substituted or unsubstituted phenyl, or and substituted or unsubstituted heteroaryl. When
15 A is substituted phenyl, it is generally substituted one, two or more times with an "aryl substituent" as defined herein. Exemplary "aryl substituents" include halogen, nitrile, substituted or unsubstituted C_1 - C_4 alkyl, SCF_3 , trifluoromethyl and trifluoromethoxy.

[0058] As discussed above, in selected compounds of the invention, R^1 is a substituted or unsubstituted aryl, e.g., substituted or unsubstituted phenyl, or substituted or
20 unsubstituted heteroaryl. When R^1 is substituted phenyl, it is generally substituted with one, two or more aryl group substituents, as defined herein, such as substituted halogen, CF_3 and OCF_3 .

[0059] In a selected embodiment, R^2 is a substituted or unsubstituted C_1 - C_6 saturated acyclic alkyl group. In yet another embodiment, R^2 is a C_1 - C_4 saturated acyclic alkyl
25 group.


[0060] In a further exemplary embodiment, the invention provides compounds having the formula:



(II)

in which the substituents are generally identical to those discussed in the context of Formula I. Additionally, R^5 and R^6 are independently selected aryl group substituents, as defined herein, such as H, halo, CF_3 , CF_3O , NO_2 , CN, $S(O)_mR^7$, $COOR^8$, $CONR^9R^{10}$, $SO_2NR^{11}R^{12}$, $S(O)_mCF_3$, substituted or unsubstituted C_1 - C_6 alkyl, or substituted or unsubstituted C_3 - C_7 cycloalkyl. The symbols R^7 and R^8 represent groups that are independently substituted or unsubstituted C_1 - C_5 alkyl, and substituted or unsubstituted C_3 - C_7 cycloalkyl. The index "m" is an integer from 0 to 2. R^9 , R^{10} , R^{11} and R^{12} are independently H, substituted or unsubstituted C_1 - C_5 alkyl, or substituted or unsubstituted C_3 - C_7 cycloalkyl. R^9 and R^{10} or R^{11} and R^{12} , together with the nitrogen atom to which they are attached, are optionally joined to form a 5- to 7-membered ring.

In yet another exemplary embodiment, the invention provides compounds according to Formula III:



(III)

in which the substituents are generally identical to those discussed in the context of Formulae I and II.


[0061] Also within the scope of the present invention are compounds that are poly- or multi-valent species, including, for example, species such as dimers, trimers, tetramers and higher homologs of the compounds of the invention or reactive analogues thereof. The poly- and multi-valent species can be assembled from a single species or more than one species of the invention. For example, a dimeric construct can be "homo-dimeric" or "heterodimeric." Moreover, poly- and multi-valent constructs in which a compound of the invention, or a reactive analogue thereof, is attached to an oligomeric or polymeric framework (*e.g.*, polylysine, dextran, hydroxyethyl starch and the like) are within the scope of the present invention. The framework is preferably polyfunctional (*i.e.* having an array of reactive sites for attaching compounds of the invention). Moreover, the

framework can be derivatized with a single species of the invention or more than one species of the invention.

[0062] Moreover, the present invention includes compounds within the motif set forth in Formula I, which are functionalized to afford compounds having a water-solubility that is enhanced relative to analogous compounds that are not similarly functionalized. Methods of enhancing the water-solubility of organic compounds are known in the art. Such methods include, but are not limited to, functionalizing an organic nucleus with a permanently charged moiety, e.g., quaternary ammonium, or a group that is charged at a physiologically relevant pH, e.g. carboxylic acid, amine. Other methods include, appending to the organic nucleus hydroxyl- or amine-containing groups, e.g. alcohols, polyols, polyethers, and the like. Representative examples include, but are not limited to, polylysine, polyethyleneimine, poly(ethyleneglycol) and poly(propyleneglycol). Suitable functionalization chemistries and strategies for these compounds are known in the art. See, for example, Dunn, R.L., *et al.*, Eds. POLYMERIC DRUGS AND DRUG DELIVERY SYSTEMS, ACS Symposium Series Vol. 469, American Chemical Society, Washington, D.C. 1991.


Preparation of Potassium Channel Modulators

[0063] Compounds of the present invention can be prepared using readily available starting materials or known intermediates. However, some intermediates produced during the synthesis are themselves novel, as indicated. Briefly, the synthesis of the compounds of the invention is performed starting with an appropriately substituted ortho-nitro or -amino carboxylic acid. The acid has the general formula:




(IA): B=NH₂; (VII): B=NO₂

[0064] By reaction involving a phenylacetylhydrazide (Z¹ = CH₂) (IIIA)



(IIIA)


or an analog that, depending on the identity of Z^1 in the desired product, will contain an appropriate group in place of the methylene group (and Y^1 represents hydrogen or one or more various "aryl group substituents" as defined herein), there is formed an intermediate (IV) or (VIII):



(IV): $B = NH_2$; (VIII): $B = NO_2$

[0065] Nitro groups in intermediates (VIII) are reduced to amino groups to form the corresponding amino intermediates (IV) (and X^1 represents hydrogen or one or more various "aryl group substituents" as defined herein).

- 10 **[0066]** The amino compounds (IV) are reacted with a suitable reactant, such as O-ethylxanthic acid, potassium salt, forming intermediate (V)



(V)


which is then alkylated, yielding the desired product, a compound of Formula I in which R^2 is -S-.

- 15 **[0067]** Compounds in which R^2 is $S(O)$ or $S(O)_2$ are prepared by specific oxidation of these compounds using, for example, m-chloroperbenzoic acid or ozone as oxidizing agents.

- [0068]** Thus, to produce compounds in which the pyrimidine-containing ring is a quinazoline ring, an ortho-amino or ortho-nitrobenzoic acid is converted by use of a phenylacetylhydrazide or analogous compound IIIA to intermediate (IV) or (VIII)
- 20

[0069] Intermediates (IV), (V), and (VIII) above are novel and form an aspect of this invention.

[0070] The process is further illustrated by Scheme 1



Scheme 1

[0071] Methods for preparing dimers, trimers and higher homologs of small organic molecules, such as those of the present invention, as well as methods of functionalizing a polyfunctional framework molecule are well known to those of skill in the art. For example, an aromatic amine of the invention is converted to the corresponding isothiocyanate by the action of thiophosgene. The resulting isothiocyanate is coupled to an amine of the invention, thereby forming either a homo- or heterodimeric species. Alternatively, the isothiocyanate is coupled with an amine-containing backbone, such as polylysine, thereby forming a conjugate between a polyvalent framework and a compound of the invention. If it is desired to prepare a heterofunctionalized polyvalent species, the polylysine is underlabeled with the first isothiocyanate and subsequently

labeled with one or more different isothiocyanates. Alternatively, a mixture of isothiocyanates is added to the backbone. Purification proceeds by, for example, size exclusion chromatography, dialysis, nanofiltration and the like.

II. ASSAYS FOR MODULATORS OF KCNQ CHANNELS

5 [0072] Assays for determining the ability of a compound of the invention to modulate, e.g., open, a potassium ion channel are generally known in the art. One of skill in the art is able to determine an appropriate assay for investigating the activity of a selected compound of the invention towards a particular ion channel. For simplicity, portions of the following discussion focuses on KCNQ2 as a representative example, however, the
10 discussion is equally applicable to other KCNQ potassium ion channels.

[0073] KCNQ monomers as well as KCNQ alleles and polymorphic variants are subunits of potassium channels. The activity of a potassium channel comprising KCNQ subunits can be assessed using a variety of *in vitro* and *in vivo* assays, e.g., measuring current, measuring membrane potential, measuring ion flux, e.g., potassium or rubidium,
15 measuring potassium concentration, measuring second messengers and transcription levels, using potassium-dependent yeast growth assays, and using e.g., voltage-sensitive dyes, radioactive tracers, and patch-clamp electrophysiology.

[0074] Furthermore, such assays can be used to test for inhibitors and activators of channels comprising KCNQ. Such modulators of a potassium channel are useful for
20 treating various disorders involving potassium channels, including but not limited to, for example, central and peripheral nervous system disorders (e.g., migraine, ataxia, Parkinson's disease, bipolar disorders, trigeminal neuralgia, spasticity, mood disorders, brain tumors, psychotic disorders, myokymia, seizures, epilepsy, hearing and vision loss, Alzheimer's disease, age-related memory loss, learning deficiencies, anxiety and motor
25 neuron diseases, and can also be used as neuroprotective agents (e.g., to prevent stroke and the like). Such modulators are also useful for investigation of the channel diversity provided by KCNQ and the regulation/modulation of potassium channel activity provided by KCNQ.

[0075] Modulators of the potassium channels are tested using biologically active
30 KCNQ, either recombinant or naturally occurring, or by using native cells, like cells from the nervous system expressing the M-current. KCNQ can be isolated, co-expressed or

expressed in a cell, or expressed in a membrane derived from a cell. In such assays, KCNQ2 is expressed alone to form a homomeric potassium channel or is co-expressed with a second subunit (*e.g.*, another KCNQ family member, preferably KCNQ3) so as to form a heteromeric potassium channel. Modulation is tested using one of the *in vitro* or *in vivo* assays described above. Samples or assays that are treated with a potential potassium channel inhibitor or activator are compared to control samples without the test compound, to examine the extent of modulation. Control samples (untreated with activators or inhibitors) are assigned a relative potassium channel activity value of 100. Activation of channels comprising KCNQ2 is achieved when the potassium channel activity value relative to the control is 130%, more preferably 150%, more preferably 170% higher. Compounds that increase the flux of ions will cause a detectable increase in the ion current density by increasing the probability of a channel comprising KCNQ2 being open, by decreasing the probability of it being closed, by increasing conductance through the channel, and increasing the number or expression of channels.

[0076] The activity of the compounds of the invention can also be represented by EC₅₀. Preferred compounds of the invention have an EC₅₀ in a potassium ion channel assay of from about 1 nM to about 10 μ M, preferably from about 1 nM to about 1 μ M, and more preferably from about 1 nM to about 500 nM.

[0077] Changes in ion flux may be assessed by determining changes in polarization (*i.e.*, electrical potential) of the cell or membrane expressing an exemplary potassium channel such as KCNQ2, KCNQ2/3 or the M-current. A preferred means to determine changes in cellular polarization is by measuring changes in current or voltage with the voltage-clamp and patch-clamp techniques, using the “cell-attached” mode, the “inside-out” mode, the “outside-out” mode, the “perforated cell” mode, the “one or two electrode” mode, or the “whole cell” mode (*see, e.g.*, Ackerman *et al.*, *New Engl. J. Med.* **336**: 1575-1595 (1997)). Whole cell currents are conveniently determined using the standard methodology (*see, e.g.*, Hamil *et al.*, *Pflugers. Archiv.* **391**: 85 (1981)). Other known assays include: radiolabeled rubidium flux assays and fluorescence assays using voltage-sensitive dyes (*see, e.g.*, Vestergaard-Bogind *et al.*, *J. Membrane Biol.* **88**: 67-75 (1988); Daniel *et al.*, *J. Pharmacol. Meth.* **25**: 185-193 (1991); Holevinsky *et al.*, *J. Membrane Biology* **137**: 59-70 (1994)). Assays for compounds capable of inhibiting or increasing potassium flux through the channel proteins comprising KCNQ2 or

heteromultimers of KCNQ subunits can be performed by application of the compounds to a bath solution in contact with and comprising cells having a channel of the present invention (*see, e.g.,* Blatz *et al.*, *Nature* **323**: 718-720 (1986); Park, *J. Physiol.* **481**: 555-570 (1994)). Generally, the compounds to be tested are present in the range from about 1 pM to about 100 mM, preferably from about 1 pM to about 1 μ M .

[0078] The effects of the test compounds upon the function of the channels can be measured by changes in the electrical currents or ionic flux or by the consequences of changes in currents and flux. Changes in electrical current or ionic flux are measured by either increases or decreases in flux of ions such as potassium or rubidium ions. The cations can be measured in a variety of standard ways. They can be measured directly by concentration changes of the ions or indirectly by membrane potential or by radio-labeling of the ions. Consequences of the test compound on ion flux can be quite varied. Accordingly, any suitable physiological change can be used to assess the influence of a test compound on the channels of this invention. The effects of a test compound can be measured by a toxin-binding assay. When the functional consequences are determined using intact cells or animals, one can also measure a variety of effects such as transmitter release (*e.g.,* dopamine), hormone release (*e.g.,* insulin), transcriptional changes to both known and uncharacterized genetic markers (*e.g.,* northern blots), cell volume changes (*e.g.,* in red blood cells), immunoresponses (*e.g.,* T cell activation), changes in cell metabolism such as cell growth or pH changes, and changes in intracellular second messengers such as Ca^{2+} , or cyclic nucleotides.

[0079] KCNQ2 orthologs will generally confer substantially similar properties on a channel comprising such KCNQ2, as described above. In a preferred embodiment, the cell placed in contact with a compound that is suspected to be a KCNQ2 homolog is assayed for increasing or decreasing ion flux in a eukaryotic cell, *e.g.,* an oocyte of *Xenopus* (*e.g., Xenopus laevis*) or a mammalian cell such as a CHO or HeLa cell. Channels that are affected by compounds in ways similar to KCNQ2 are considered homologs or orthologs of KCNQ2.

III. PHARMACEUTICAL COMPOSITIONS OF POTASSIUM CHANNEL MODULATORS

[0080] In another aspect, the present invention provides pharmaceutical compositions comprising a pharmaceutically acceptable excipient and a compound of Formula I provided above.

Formulation of the Compounds (Compositions)

5 [0081] The compounds of the present invention can be prepared and administered in a wide variety of oral, parenteral and topical dosage forms. Thus, the compounds of the present invention can be administered by injection, that is, intravenously, intramuscularly, intracutaneously, subcutaneously, intraduodenally, or intraperitoneally. Also, the compounds described herein can be administered by inhalation, for example, intranasally.
10 Additionally, the compounds of the present invention can be administered transdermally. Accordingly, the present invention also provides pharmaceutical compositions comprising a pharmaceutically acceptable carrier or excipient and either a compound of Formula I, or a pharmaceutically acceptable salt thereof.

[0082] For preparing pharmaceutical compositions from the compounds of the present
15 invention, pharmaceutically acceptable carriers can be either solid or liquid. Solid form preparations include powders, tablets, pills, capsules, cachets, suppositories, and dispersible granules. A solid carrier can be one or more substances, which may also act as diluents, flavoring agents, binders, preservatives, tablet disintegrating agents, or an encapsulating material.

20 [0083] In powders, the carrier is a finely divided solid, which is in a mixture with the finely divided active component. In tablets, the active component is mixed with the carrier having the necessary binding properties in suitable proportions and compacted in the shape and size desired.

[0084] The powders and tablets preferably contain from 5% or 10% to 70% of the
25 active compound. Suitable carriers are magnesium carbonate, magnesium stearate, talc, sugar, lactose, pectin, dextrin, starch, gelatin, tragacanth, methylcellulose, sodium carboxymethylcellulose, a low melting wax, cocoa butter, and the like. The term “preparation” is intended to include the formulation of the active compound with encapsulating material as a carrier providing a capsule in which the active component
30 with or without other carriers, is surrounded by a carrier, which is thus in association with

it. Similarly, cachets and lozenges are included. Tablets, powders, capsules, pills, cachets, and lozenges can be used as solid dosage forms suitable for oral administration.

[0085] For preparing suppositories, a low melting wax, such as a mixture of fatty acid glycerides or cocoa butter, is first melted and the active component is dispersed
5 homogeneously therein, as by stirring. The molten homogeneous mixture is then poured into convenient sized molds, allowed to cool, and thereby to solidify.

[0086] Liquid form preparations include solutions, suspensions, and emulsions, for example, water or water/propylene glycol solutions. For parenteral injection, liquid preparations can be formulated in solution in aqueous polyethylene glycol solution.

10 [0087] Aqueous solutions suitable for oral use can be prepared by dissolving the active component in water and adding suitable colorants, flavors, stabilizers, and thickening agents as desired. Aqueous suspensions suitable for oral use can be made by dispersing the finely divided active component in water with viscous material, such as natural or synthetic gums, resins, methylcellulose, sodium carboxymethylcellulose, and other well-
15 known suspending agents.

[0088] Also included are solid form preparations, which are intended to be converted, shortly before use, to liquid form preparations for oral administration. Such liquid forms include solutions, suspensions, and emulsions. These preparations may contain, in addition to the active component, colorants, flavors, stabilizers, buffers, artificial and
20 natural sweeteners, dispersants, thickeners, solubilizing agents, and the like.

[0089] The pharmaceutical preparation is preferably in unit dosage form. In such form the preparation is subdivided into unit doses containing appropriate quantities of the active component. The unit dosage form can be a packaged preparation, the package containing discrete quantities of preparation, such as packeted tablets, capsules, and
25 powders in vials or ampoules. Also, the unit dosage form can be a capsule, tablet, cachet, or lozenge itself, or it can be the appropriate number of any of these in packaged form.

[0090] The quantity of active component in a unit dose preparation may be varied or adjusted from 0.1 mg to 10000 mg, more typically 1.0 mg to 1000 mg, most typically 10 mg to 500 mg, according to the particular application and the potency of the active
30 component. The composition can, if desired, also contain other compatible therapeutic agents.

Effective Dosages

- [0091] Pharmaceutical compositions provided by the present invention include compositions wherein the active ingredient is contained in a therapeutically effective amount, *i.e.*, in an amount effective to achieve its intended purpose. The actual amount effective for a particular application will depend, *inter alia*, on the condition being treated. For example, when administered in methods to treat pain or anxiety, such compositions will contain an amount of active ingredient effective to achieve a clinically relevant degree of reduction in the condition being treated. Similarly, when the pharmaceutical composition is used to treat or prevent a central or peripheral nervous system disorder, e.g., Parkinson's disease a therapeutically effective amount will reduce one or more symptoms characteristic of the diseases (e.g., tremors) to below a predetermined pressure threshold. Determination of a therapeutically effective amount of a compound of the invention is well within the capabilities of those skilled in the art, especially in light of the detailed disclosure herein.
- 15 [0092] For any compound described herein, the therapeutically effective amount can be initially determined from cell culture assays. Target plasma concentrations will be those concentrations of active compound(s) that are capable of modulating, e.g., activating or opening the KCNQ channel. In preferred embodiments, the KCNQ channel activity is altered by at least 30%. Target plasma concentrations of active compound(s) that are
- 20 capable of inducing at least about 50%, 70%, or even 90% or higher alteration of the KCNQ channel potassium flux are presently preferred. The percentage of alteration of the KCNQ channel in the patient can be monitored to assess the appropriateness of the plasma drug concentration achieved, and the dosage can be adjusted upwards or downwards to achieve the desired percentage of alteration.
- 25 [0093] As is well known in the art, therapeutically effective amounts for use in humans can also be determined from animal models. For example, a dose for humans can be formulated to achieve a circulating concentration that has been found to be effective in animals. A particularly useful animal model for predicting anticonvulsant dosages is the maximal electroshock assay (Fischer RS, *Brain Res. Rev.* **14**: 245-278 (1989)). The
- 30 dosage in humans can be adjusted by monitoring KCNQ channel activation and adjusting the dosage upwards or downwards, as described above.

[0094] A therapeutically effective dose can also be determined from human data for compounds which are known to exhibit similar pharmacological activities, such as retigabine (Rudnfeldt *et al.*, *Neuroscience Lett.* **282**: 73-76 (2000)).

5 [0095] Adjusting the dose to achieve maximal efficacy in humans based on the methods described above and other methods as are well-known in the art is well within the capabilities of the ordinarily skilled artisan.

[0096] By way of example, when a compound of the invention is used in the prophylaxis and/or treatment of an exemplary disease such as epilepsy, a circulating concentration of administered compound of about 0.001 μM to 20 μM is considered to be
10 effective, with about 0.01 μM to 5 μM being preferred.

[0097] Patient doses for oral administration of the compounds described herein, which is the preferred mode of administration for prophylaxis and for treatment of an exemplary disease such as epilepsy, typically range from about 1 mg/day to about 10,000 mg/day, more typically from about 10 mg/day to about 1,000 mg/day, and most typically from
15 about 1 mg/day to about 500 mg/day. Stated in terms of patient body weight, typical dosages range from about 0.01 to about 150 mg/kg/day, more typically from about 0.1 to about 15 mg/kg/day, and most typically from about 0.5 to about 10 mg/kg/day.

[0098] For other modes of administration, dosage amount and interval can be adjusted individually to provide plasma levels of the administered compound effective for the
20 particular clinical indication being treated. For example, if acute epileptic seizures are the most dominant clinical manifestation, in one embodiment, a compound according to the invention can be administered in relatively high concentrations multiple times per day. Alternatively, if the patient exhibits only periodic epileptic seizures on an infrequent, periodic or irregular basis, in one embodiment, it may be more desirable to administer a
25 compound of the invention at minimal effective concentrations and to use a less frequent administration regimen. This will provide a therapeutic regimen that is commensurate with the severity of the individual's disease.

[0099] Utilizing the teachings provided herein, an effective prophylactic or therapeutic treatment regimen can be planned which does not cause substantial toxicity and yet is
30 entirely effective to treat the clinical symptoms demonstrated by the particular patient. This planning should involve the careful choice of active compound by considering

factors such as compound potency, relative bioavailability, patient body weight, presence and severity of adverse side effects, preferred mode of administration and the toxicity profile of the selected agent.

Compound Toxicity

5 [0100] The ratio between toxicity and therapeutic effect for a particular compound is its therapeutic index and can be expressed as the ratio between LD₅₀ (the amount of compound lethal in 50% of the population) and ED₅₀ (the amount of compound effective in 50% of the population). Compounds that exhibit high therapeutic indices are preferred. Therapeutic index data obtained from cell culture assays and/or animal studies can be
10 used in formulating a range of dosages for use in humans. The dosage of such compounds preferably lies within a range of plasma concentrations that include the ED₅₀ with little or no toxicity. The dosage may vary within this range depending upon the dosage form employed and the route of administration utilized. See, *e.g.* Fingl *et al.*, *In: THE PHARMACOLOGICAL BASIS OF THERAPEUTICS*, Ch.1, p.1, 1975. The exact
15 formulation, route of administration and dosage can be chosen by the individual physician in view of the patient's condition and the particular method in which the compound is used.

IV. METHODS FOR INCREASING ION FLOW IN VOLTAGE-DEPENDENT POTASSIUM CHANNELS

20 [0101] In yet another aspect, the present invention provides methods for increasing ion flow through voltage dependent potassium channels in a cell. The method includes contacting a cell containing the target ion channels with an amount of a compound of the invention sufficient to enhance the activity of a potassium channel.

[0102] The methods provided in this aspect of the invention are useful for the diagnosis
25 of conditions that can be treated by modulating ion flux through voltage-dependent potassium channels, or for determining if a patient will be responsive to therapeutic agents, which act by opening potassium channels. In particular, a patient's cell sample can be obtained and contacted with a compound of the invention and the ion flux can be measured relative to a cell's ion flux in the absence of a compound of the invention. An
30 increase in ion flux will typically indicate that the patient will be responsive to a therapeutic regimen of ion channel openers.

V. METHODS FOR TREATING CONDITIONS MEDIATED BY VOLTAGE-DEPENDENT POTASSIUM CHANNELS

[0103] In still another aspect, the present invention provides a method for the treatment of a central or peripheral nervous system disorder or condition through modulation of a voltage-dependent potassium channel. In this method, a subject in need of such treatment is administered an effective amount of a compound having the formula provided above.

[0104] The compounds provided herein are useful as potassium channel modulators and find therapeutic utility via modulation of voltage-dependent potassium channels in the treatment of diseases or conditions. The potassium channels targets for the compounds of the invention are described herein as voltage-dependent potassium channels such as the KCNQ potassium channels. As noted above, these channels may include homomultimers and heteromultimers of KCNQ2, KCNQ3, KCNQ4, and KCNQ5. A heteromultimer of two proteins, *e.g.*, KCNQ2 and KCNQ3 is referred to as, for example, KCNQ2/3, KCNQ3/5, etc. The conditions that can be treated with the compounds and compositions of the present invention may include, but are not limited to, central or peripheral nervous system disorders (*e.g.*, migraine, ataxia, Parkinson's disease, bipolar disorders, trigeminal neuralgia, spasticity, mood disorders, brain tumors, psychotic disorders, myokymia, seizures, epilepsy, hearing and vision loss, Alzheimer's disease, age-related memory loss, learning deficiencies, anxiety, and motor neuron diseases). The compounds and compositions of the present invention may also serve as neuroprotective agents (*e.g.*, to prevent stroke, retinal degeneration and the like). In a preferred embodiment, the condition or disorder to be treated is epilepsy or seizures. In another preferred embodiment, the condition or disorder is hearing loss.

[0105] In therapeutic use for the treatment of epilepsy or other neurological conditions, the compounds utilized in the pharmaceutical method of the invention are administered at the initial dosage of about 0.001 mg/kg to about 1000 mg/kg daily. A daily dose range of about 0.1 mg/kg to about 100 mg/kg is more typical. The dosages, however, may be varied depending upon the requirements of the patient, the severity of the condition being treated, and the compound being employed. Determination of the proper dosage for a particular situation is within the skill of the practitioner. Generally, treatment is initiated with smaller dosages, which are less than the optimum dose of the compound. Thereafter, the dosage is increased by small increments until the optimum effect under

circumstances is reached. For convenience, the total daily dosage may be divided and administered in portions during the day, if desired.

EXAMPLES

[0106] The following examples are offered to illustrate, but not to limit the claimed invention.

[0107] In the examples below, unless otherwise stated, temperatures are given in degrees Celsius ($^{\circ}\text{C}$); operations were carried out at room or ambient temperature (typically a range of from about 18-25 $^{\circ}\text{C}$; evaporation of solvent was carried out using a rotary evaporator under reduced pressure (typically, 4.5-30 mmHg) with a bath temperature of up to 60 $^{\circ}\text{C}$; the course of reactions was typically followed by TLC and reaction times are provided for illustration only; melting points are uncorrected; products exhibited satisfactory ^1H -NMR and/or microanalytical data; yields are provided for illustration only; and the following conventional abbreviations are also used: mp (melting point), L (liter(s)), mL (milliliters), mmol (millimoles), g (grams), mg (milligrams), min (minutes), and h (hours).

EXAMPLES

[0108] In the description below, unless otherwise stated, temperatures are given in degrees Celsius ($^{\circ}\text{C}$). The course of reactions was typically followed by TLC and reaction times are provided for illustration only. Melting points are uncorrected; products exhibited satisfactory ^1H -NMR and/or microanalytical data; yields are provided for illustration only; and the following conventional abbreviations are also used: mp (melting point), L (liter(s)), mL (milliliters), eq (equivalents), mmol (millimoles), g (grams), mg (milligrams), min (minutes), and h (hours).

[0109] Unless otherwise specified, all solvents (HPLC grade) and reagents were purchased from suppliers and used without further purification. Analytical thin layer chromatography (TLC) was performed on Whatman Inc. 60 silica gel plates (0.25 mm thickness). Compounds were visualized under UV lamp (254 nm) or by developing with KMnO_4/KOH , ninhydrin or Hanessian's solution. Flash chromatography was done using silica gel from Selectro Scientific (particle size 32-63). ^1H NMR, ^{19}F NMR and ^{13}C NMR spectra were recorded on a Varian 300 machine at 300 MHz, 282 MHz and 75.7 MHz,

respectively. Melting points were recorded on an Electrothermal IA9100 apparatus and were uncorrected.

EXAMPLE 1

1.1 General procedure for preparation of isatoic anhydrides (II)

- 5 [0110] Anhydrous pyridine (2 eq) was added to a solution of 2-aminobenzoic acid derivative (IA) (1 eq) in dry methylene chloride and acetonitrile (1:1, 40 mL/g of 2-aminobenzoic acid) at room temperature. Solid triphosgene (1/3 eq) was then added in one portion and the resulting mixture was heated at 50 °C for 2 h. The resulting solid was collected by filtration and dried in vacuo. The crude isatoic anhydrides (IIA) were
10 typically obtained in 50 – 80% yields. Though contaminated with some pyridinium hydrochloride, the anhydrides were used in the next step without further purification.

1.2 General procedure for the preparation of acylhydrazides (IV)

- [0111] A mixture of isatoic anhydride (IIA, 1 eq) and appropriate phenylacetylhydrazide (IIIA, 1.1 eq) were heated in glacial AcOH (2 mL/mmol) at 50 °C
15 for 2-6 h. The resulting solution was cooled and water was added while shaking. The white precipitate was collected by filtration, washed with water and dried *in vacuo* at 50 °C for 4 h. The desired products (IV) were obtained as white solids in high purity (typically >90%) and moderate yields (typically 45-60%).

1.3 Representative compounds

- 20 [0112] *1.3a 2-Amino-4-fluoro-benzoic acid N'-phenylacetyl-hydrazide*
Obtained as a white powder. ¹H NMR (300 MHz, d-6 DMSO) δ 3.05(2H, s), 6.30 (1H, dt, *J* = 8.5, 2.4 Hz), 6.45 (1H, dd, *J* = 12.0, 2.6 Hz), 6.72 (2H, brs), 7.18-7.34 (5H, m), 7.57 (1H, dd, *J* = 8.9, 6.8 Hz), 10.02 (2H, brs).
- 25 [0113] *1.3b 2-Amino-4-fluoro-benzoic acid N'-[2-(3-fluoro-phenyl)-acetyl]-hydrazide*.
Obtained as a white powder. ¹H NMR (300 MHz, d-6 DMSO) δ 3.54(2H, s), 6.30 (1H, dt, *J* = 8.7, 2.6 Hz), 6.46 (1H, dd, *J* = 11.8, 2.4 Hz), 6.72 (2H, brs), 7.02-7.19 (3H, m), 7.34 (1H, dt, *J* = 14.4, 7.8 Hz), 7.57 (1H, dd, *J* = 8.7, 6.8 Hz), 10.05 (2H, brs).

EXAMPLE 2

2.1 *General procedure for the preparation of thioureas (V)*

[0114] A mixture of an acylhydrazide (IV, 1 eq) and KSCSOEt (1.5-2 eq) in ethanol (10 mL/mmol) was heated at 70 °C overnight. The resulting yellow mixture was cooled and neutralized with 1N aqueous HCl. Water was added (10-20 mL/mmol) and the mixture was vortexed for 10 sec. The resulting white solid was collected by filtration, washed with water and dried in vacuo at 50 °C for 6-10 h. The desired product (V) was obtained as a white solid in high purity (typically >90%) and high yield (typically >80%).

2.2 *Representative compounds*

10 [0115] 2.2a *N*-(5-Fluoro-2-mercapto-4-oxo-4H-quinazolin-3-yl)-2-phenyl-acetamide
¹H NMR (300 MHz, d-6 DMSO) δ 3.65 (2H, s), 7.09-7.17 (3H, m), 7.20 (1H, d, *J* = 8.4 Hz), 7.34-7.41 (2H, m), 7.76 (1H, dt, *J* = 8.2, 5.4 Hz), 11.11 (1H, brs), 13.21 (1H, brs).

15 [0116] 2.2b *N*-(7-Fluoro-2-mercapto-4-oxo-4H-quinazolin-3-yl)-2-(3-fluoro-phenyl)-acetamide
¹H NMR (300 MHz, d-6 DMSO) δ 3.70 (2H, s), 7.02-7.24 (5H, m), 7.36 (1H, dt, *J* = 8.0, 7.6 Hz), 8.02 (1H, dd, *J* = 8.9, 5.9 Hz), 11.27 (1H, brs), 13.21 (1H, brs).

20 [0117] 2.2c *N*-(7-Fluoro-2-mercapto-4-oxo-4H-quinazolin-3-yl)-2-phenyl-acetamide
¹H NMR (300 MHz, d-6 DMSO) δ 3.65 (2H, s), 7.12 (1H, dd, *J* = 9.7, 2.3 Hz), 7.17-7.39 (6H, m), 8.02 (1H, dd, *J* = 8.7, 8.9 Hz), 11.16 (1H, brs), 13.20 (1H, brs).

EXAMPLE 3

3.1 *General procedure for preparation of sulfides (VI)*

[0118] Aqueous sodium hydroxide (3N, 1.1 eq) was added to the appropriate sulfide intermediate (V, 1 eq) in methanol or ethanol at room temperature. To the resulting homogeneous solution was added the appropriate alkylating agent (1.1 eq) in one portion. The reaction was then shaken at 20-60 °C. When the reaction was judged to be complete (by LCMS), water was added to the reaction mixture (75 mL/g of intermediate) and the pH was adjusted to 7 with 6N HCl. The alcoholic solvent was removed under reduced pressure and the resulting solids were collected by filtration, washed with water and dried in vacuo affording the desired final products (VI) in 40-90% yields.

EXAMPLE 4*4.1 General procedure for preparation of nitrobenzoic-N'-phenylacetyl hydrazides (VIII)*

[0119] The appropriate nitrobenzoic acid derivative (VII) (1 eq) was stirred in dry methylene chloride (100 mL/g of acid) at room temperature and to this was added two drops of N,N-dimethylformamide (DMF). Neat oxalyl chloride (2 eq) was then added to the mixture dropwise at such a rate as to control gas evolution. After stirring for 2 h, the volatiles were removed by rotary evaporation and the remaining material was re-dissolved in dry methylene chloride (100 mL/g of acid). Pyridine (2 eq) and the appropriate hydrazide derivative (IIIA) (1 eq) were added consecutively and the mixture was allowed to stir at room temperature until the reaction was judged to be complete by HPLC analysis, whereupon the reaction mixture was poured into water. The organic layer was removed and the water layer was extracted three times with ethyl acetate. The combined organic layers were dried (Na₂SO₄) and concentrated to provide the desired nitrobenzoic-N'-phenylacetyl hydrazides (VIII) in high purity (typically >95%) and yields ranging from 40 to 90%.

EXAMPLE 5*5.1 General procedure for preparation of 2-aminobenzoic acid N'-phenylacetyl-hydrazides (IV) from nitrobenzoic-N'-phenylacetyl hydrazides (VIII)*

[0120] 1 atmosphere of hydrogen was applied to a mixture of nitrobenzoic-N'-phenylacetyl hydrazide (VIII) in methanol (100 mL/g of hydrazide) and 10% palladium on activated carbon (100 mg/g of hydrazide). The reaction mixture was stirred at room temperature for 1-10 h. The resulting mixture was filtered through celiteTM/silica gel and concentrated under reduced pressure. The desired 2-aminobenzoic acid N'-phenylacetyl-hydrazides (IV) were, in general, used directly in the next step without any further purification. In those instances when the desired products were contaminated with significant impurities, the hydrazides were purified by silica gel chromatography using EtOAc/hexanes.

EXAMPLE 6

[0121] Example 6 sets forth the characterization of a number of representative compounds of the invention, using the general procedures above. The compounds were characterized using a combination of melting point, ¹H NMR and mass spectrometry. The results of the characterization are presented below. The structures for the compounds

set forth below are provided in **FIG. 1**. Numbers in parentheses refer to compound numbers in **FIG. 1**.

[0122] 6.1 *2-(3-Fluoro-phenyl)-N-(2-methylsulfanyl-4-oxo-4H-quinazolin-3-yl)-acetamide (72)*

5 Obtained as a white powder in 49% yield and 98% purity. ¹H NMR (300 MHz, CDCl₃) δ 2.51(3H, s), 3.82 (2H, AB, *J* = 16.0, 2.4 Hz), 7.01(1H, td, *J* = 8.4, 1.7 Hz), 7.14 (1H, d, *J* = 9.4Hz), 7.19 (1H, d, *J* = 7.7Hz), 7.56 (1H, d, *J* = 8.0Hz), 7.35 (2H, m), 7.70 (1H, td, *J* = 1.6Hz), 8.15 (2H, m).

10 **[0123]** 6.2 *N-(2-Ethylsulfanyl-4-oxo-4H-quinazolin-3-yl)-2-(4-fluoro-phenyl)-acetamide (65)*

Obtained as a white powder in 35% yield and >95% purity. ¹H NMR (300 MHz, CDCl₃) δ 1.37 (3H, t, *J* = 7.3Hz), 3.12 (2H, m), 3.80 (2H, AB, *J* = 16.0, 3.4Hz), 7.20 (2H, m), 7.35 (3H, m), 7.53 (1H, d, *J* = 8.2Hz), 7.69 (1H, t, *J* = 8Hz), 8.13 (1H, d, *J* = 8.0Hz).

15 **[0124]** 6.3 *N-(7-Fluoro-2-methylsulfanyl-4-oxo-4H-quinazolin-3-yl)-2-phenyl-acetamide (40)*

Obtained as a white solid in 61% yield and 98% purity. ¹H NMR (300 MHz, CDCl₃) δ 2.50 (3H, s), 3.84 (2H, dd, *J* = 18.1, 16.7Hz), 7.07 (1H, dt, *J* = 8.5, 2.4Hz), 7.20 (1H, dd, *J* = 9.7, 2.4Hz), 7.30–7.46 (4H, m), 7.77 (1H, s), 8.14 (1H, dd, *J* = 8.9, 6.1Hz).

20 **[0125]** 6.4 *N-(7-Fluoro-2-isopropylsulfanyl-4-oxo-4H-quinazolin-3-yl)-2-(3-fluoro-phenyl)-acetamide (71)*

Obtained as a tan solid in 27% yield and 97% purity. ¹H NMR (300 MHz, CDCl₃) δ 1.35 (6H, dd, *J* = 12.6, 6.8Hz), 3.55 (1H, d, *J* = 14.6Hz), 3.70 (1H, d, *J* = 15.0Hz), 3.82 (1H, pentet, *J* = 7.1Hz), 6.75 (1H, t, *J* = 6.4Hz), 6.89 (1H, t, *J* = 8.7Hz), 6.96-7.08 (2H, m), 7.18 (1H, q, *J* = 7.1Hz), 7.81 (1H, t, *J* = 6.1Hz).

30 **[0126]** 6.5 *N-(2-Ethylsulfanyl-7-fluoro-4-oxo-4H-quinazolin-3-yl)-2-(4-fluoro-phenyl)-acetamide (50)*

Obtained as a white solid in 86% yield and 97% purity. ¹H NMR (300 MHz, CDCl₃) δ 1.37 (3H, t, *J* = 7.5Hz), 3.01-3.19 (2H, m), 3.79 (2H, s), 7.03-7.12 (3H, m), 7.19 (1H, dd, *J* = 9.7, 2.4 Hz), 7.38 (1H, dd, *J* = 8.5, 5.4Hz), 7.96 (1H, s), 8.13 (1H, dd, *J* = 8.9, 6.2 Hz).

35

[0127] 6.6 *N*-(2-Ethylsulfanyl-5-fluoro-4-oxo-4*H*-quinazolin-3-yl)-2-phenyl-acetamide (69)

Obtained as a white solid in 78% yield and 100% purity. ¹H NMR (300 MHz, CDCl₃) δ 1.37 (3H, t, *J* = 7.5 Hz), 3.08-3.18 (2H, m), 3.13 (2H, dd, *J* = 12.0, 7.4 Hz), 6.96 (1H, t, *J* = 8.7 Hz), 7.30-7.46 (5H, m), 7.55-7.63 (1H, m).

[0128] 6.7 *N*-(7-Fluoro-2-isopropylsulfanyl-4-oxo-4*H*-quinazolin-3-yl)-2-phenyl-acetamide (48)

Obtained as a white solid in 56% yield and 99% purity. ¹H NMR (300 MHz, CDCl₃) δ 1.40 (6H, dd, *J* = 7.1, 1.8 Hz), 3.82 (2H, dd, *J* = 18.6, 16.3 Hz), 3.92 (1H, septet, *J* = 6.9 Hz), 7.05 (1H, dt, *J* = 8.7, 2.4 Hz), 7.17 (1H, dd, *J* = 9.7, 2.4 Hz), 7.28-7.44 (4H, m), 7.86 (1H, s), 8.13 (1H, dd, *J* = 8.7, 6.0 Hz).

[0129] 6.8 *N*-(5-Fluoro-2-methylsulfanyl-4-oxo-4*H*-quinazolin-3-yl)-2-(4-fluoro-phenyl)-acetamide (66)

Obtained as a white solid in 70% yield and 100% purity. ¹H NMR (300 MHz, CD₃OD) δ 2.52 (3H, s), 3.77 (2H, s), 7.08 (3H, m), 7.41 (3H, m), 7.74 (1H, m).

[0130] 6.9 *N*-(2-Methylsulfanyl-4-oxo-7-trifluoromethyl-4*H*-quinazolin-3-yl)-2-phenyl-acetamide (51)

Obtained as a white solid in 50% yield and 98% purity. ¹H NMR (300 MHz, CDCl₃) δ 2.53 (3H, s), 3.86 (2H, s), 7.30-7.47 (5H, m), 7.56 (1H, d, *J* = 8.4 Hz), 7.76 (1H, s), 7.84 (1H, s), 8.24 (1H, d, *J* = 8.3 Hz).

[0131] 6.10 *N*-(2-Ethylsulfanyl-4-oxo-7-trifluoromethyl-4*H*-quinazolin-3-yl)-2-phenyl-acetamide (52)

Obtained as a white solid in 23% yield and 98% purity. ¹H NMR (300 MHz, CDCl₃) δ 1.40 (3H, t, *J* = 7.5 Hz), 3.10-3.20 (2H, m), 3.87 (3H, s), 7.33-7.48 (5H, m), 7.56 (1H, d, *J* = 8.5 Hz), 7.60 (1H, s), 7.84 (1H, s), 8.25 (1H, d, *J* = 8.0 Hz).

[0132] 6.11 *N*-(2-Isopropylsulfanyl-4-oxo-7-trifluoromethyl-4*H*-quinazolin-3-yl)-2-phenyl-acetamide (53)

Obtained as a white solid in 23% yield and 98% purity. ¹H NMR (300 MHz, CDCl₃) δ 1.43 (6H, dd, *J* = 6.8, 4.6 Hz), 3.86 (2H, s), 3.97 (1H, pentet, *J* = 6.8 Hz), 7.34-7.48 (4H, m), 7.52-7.58 (2H, m), 7.82 (1H, s), 8.25 (1H, d, *J* = 8.4 Hz).

[0133] 6.12 2-(3-Fluoro-phenyl)-N-(2-methylsulfanyl-4-oxo-7-trifluoromethyl-4H-quinazolin-3-yl)-acetamide (63)

Obtained as an off-white solid in 100% yield and 98% purity. ¹H NMR (300 MHz, CDCl₃) δ 2.54 (3H, s), 3.85 (3H, s), 7.05 (1H, dt, *J* = 8.5, 2.4Hz), 7.15 (1H, d, *J* = 9.4Hz),
 5 7.21 (1H, d, *J* = 7.6Hz), 7.38 (1H, dt, *J* = 7.8, 6.1Hz), 7.57 (1H, d, *J* = 7.5Hz), 7.82 (1H, s), 7.86 (1H, s), 8.26 (1H, d, *J* = 8.2Hz).

[0134] 6.13 2-(4-Fluoro-phenyl)-N-(2-methylsulfanyl-4-oxo-7-trifluoromethyl-4H-quinazolin-3-yl)-acetamide (62)

10 Obtained as a tan solid in 95% yield and 98% purity. ¹H NMR (300 MHz, CDCl₃) δ 2.54 (3H, s), 3.82 (2H, s), 7.39 (2H, m), 7.57(1H, d, *J* = 8.4Hz), 7.85 (1H, s), 7.90 (2H, t, *J* = 8.5Hz), 8.25 (1H, d, *J* = 8.2Hz).

[0135] 6.14 N-[4-Oxo-2-(3,3,3-trifluoro-propylsulfanyl)-5,6,7,8-tetrahydro-4H-quinazolin-3-yl]-2-phenyl-acetamide (142)

15 ¹H NMR (300 MHz, d-6 DMSO) δ 1.55-1.78 (5H, m), 2.26-2.39 (2H, m), 2.50-2.65 (3H, m), 3.05-3.22 (2H, m), 3.67 (2H, s), 7.23-7.32 (5H, m), 11.25 (1H, brs).

[0136] 6.15 N-(7-Methyl-2-methylsulfanyl-4-oxo-4H-thieno[3,2-d]pyrimidin-3-yl)-2-phenyl-acetamide (90)

20 ¹H NMR (300 MHz, d-6 DMSO) δ 2.30 (3H, s), 2.48 (3H, s), 3.72 (3H, s), 7.23-7.35 (5H, m), 7.86 (1H, s), 11.39 (1H, brs).

[0137] 6.16 N-(7-Methanesulfonyl-2-methylsulfanyl-4-oxo-4H-quinazolin-3-yl)-2-phenyl-acetamide (157)

25 ¹H NMR (300 MHz, d-6 DMSO) δ 2.52 (3H, s), 2.33 (3H, s), 3.74 (3H, s), 7.24-7.35 (5H, m), 7.92 (1H, dd, *J* = 8.4, 1.6 Hz), 8.07 (1H, d, *J* = 1.4 Hz), 8.28 (1H, d, *J* = 8.4 Hz), 11.50 (1H, brs).

[0138] 6.17 2-(3-Fluoro-phenyl)-N-(2-isopropylsulfanyl-7-methyl-4-oxo-4H-thieno[3,2-d]pyrimidin-3-yl)-acetamide (94)

30 ¹H NMR (300 MHz, d-6 DMSO) δ 1.35 (3H, d, *J* = 6.9 Hz), 1.38 (3H, d, *J* = 6.8 Hz), 2.29 (3H, s), 3.74 (3H, s), 3.83 (1H, sept, *J* = 7.0 Hz), 7.08 (1H, d, *J* = 8.9, 2.3 Hz), 7.17-7.20 (2H, m), 7.33-7.41 (1H, m), 7.87 (1H, s), 11.37 (1H, brs).

EXAMPLE 7

[0139] This example illustrates a screening protocol for evaluating compounds of the present invention for the ability to open voltage-gated potassium channels.

5 [0140] NG108-15 cells, a mouse neuroblastoma, rat glioma hybrid cell line, functionally express M-currents (Robbins *et al.*, *J. Physiol.* **451**: 159-85 (1992). NG108-15 M-currents are likely comprised, at least in part, of KCNQ2, KCNQ3 and KCNQ5, since these genes are reportedly robustly expressed in differentiated NG108-15 cells (Selyanko *et al.*, *J. Neurosci.* **19**(18): 7742-56 (1999); Schroeder *et al.*, *J. Biol. Chem.* **275**(31): 24089-95 (2000)) and KCNQ3 dominant-negative constructs reduce M-current density in
10 these cells (Selyanko *et al.*, *J. Neurosci.* **22**(5): RC212 (2002).

[0141] NG108-15 were maintained in DMEM (high glucose) supplemented with 10% fetal bovine serum, 0.05 mM pyridoxine, 0.1 mM hypoxanthine, 400 nM aminopterin, 16 mM thymidine, 50 μgml^{-1} gentamycin and 10 mM HEPES, in an incubator at 37°C with a humidified atmosphere of 5 % CO₂. Cells were plated in 96 well plates differentiated by
15 addition of 10 μM PGE1 and 50 μM isomethylbutylxanthine to the growth media prior to study.

[0142] Differentiated NG108-15 cells were loaded with voltage-sensitive dye by incubation in Earls Balanced Salt Solution (EBSS) containing 5 mM DiBAC for 1h. Following loading, drug solution containing 5 mM DiBAC was added to each well.
20 Changes in fluorescence were measured every 30 s for 25 min. The maximum change in fluorescence was measured and expressed as a percentage of the maximum response obtained in the presence of a positive control agent.


[0143] FIG. 2 includes results of assays of compounds of the invention by the above procedure.

25

[0144] It is understood that the examples and embodiments described herein are for illustrative purposes only and that various modifications or changes in light thereof will be suggested to persons skilled in the art and are to be included within the spirit and purview of this application and scope of the appended claims.

WHAT IS CLAIMED IS:

1. A compound having the formula:



5 in which

A is a member selected from the group consisting of five- and six-membered substituted or unsubstituted aryl, five- and six-membered substituted or unsubstituted heteroaryl, substituted or unsubstituted C₄-C₈ cycloalkyl, and substituted or unsubstituted 5-8 membered heterocyclyl;

10 X is a member selected from the group consisting of CO, CS and SO₂;

Z is a member selected from the group consisting of -CH₂-, -CHF-, -CF₂-, -CH=CH- and -N(R⁴)(CR^{4a}R^{4b})_s-

wherein R⁴ is a member selected from the group consisting of H and a substituted or unsubstituted C₁-C₅ alkyl group;

15 R^{4a} and R^{4b} are members independently selected from the group consisting of H, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, substituted or unsubstituted C₃-C₈ cycloalkyl, substituted or unsubstituted 5-7 membered heterocyclyl, and substituted or unsubstituted C₁-C₈ alkyl;

20 s is an integer from 1 to 3;

Y is S(O)_n, wherein n is an integer from 0-2;

R¹ is a member selected from the group consisting of substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, substituted or unsubstituted C₃-C₈ cycloalkyl, substituted or unsubstituted 5-7 membered heterocyclyl, and substituted or unsubstituted C₁-C₈ alkyl; and

25

R² is a member selected from the group consisting of CF₃, substituted or unsubstituted C₁-C₈ alkyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, substituted or unsubstituted C₃-C₈ cycloalkyl, and substituted or unsubstituted 3-7-membered heterocyclyl;

provided that when A is phenyl, Z is -CH₂- and R¹ is phenyl, substituted phenyl or heteroaryl, then R² is other than a benzyl, substituted benzyl, alkylheteroaryl, alkylheterocyclyl or cyanomethyl group;

wherein:

- 5 the substituents for the alkyl, cycloalkyl, heterocyclyl and heteroalkyl groups are selected from the group consisting of -OR', =O, =NR', =N-OR', -NR'R'', -SR', -halogen, -SiR'R''R''', -OC(O)R', -C(O)R', -CO₂R', -CONR'R'', -OC(O)NR'R'', -NRC(O)R', -NR-C(O)NR'R'', -NRC(O₂)R', -NR-C(NR'R'')=NR''', -S(O)R', -S(O)₂R', -S(O)₂NR'R'', -NRSO₂R', -CN and -
- 10 NO₂ in a number ranging from zero to (2m'+1), where m' is the total number of carbon atoms in such group and R, R', R'' and R''' are each independently selected from the group consisting of hydrogen, unsubstituted heteroalkyl, unsubstituted aryl, unsubstituted alkyl, alkoxy or thioalkoxy groups, or arylalkyl groups; and
- 15 the substituents for the aryl and heteroaryl groups are selected from the group consisting of -halogen, -OR', =O, =NR', =N-OR', -NR'R'', -SR', -SiR'R''R''', -OC(O)R', -C(O)R', -CO₂R', -CONR'R'', -OC(O)NR'R'', -NRC(O)R', -NR-C(O)NR'R'', -NRC(O)₂R', -NR-C(NR'R'')=NR''', -S(O)R', -S(O)₂R', -S(O)₂NR'R'', -NRSO₂R', -CN, -NO₂, -R', -N₃, -CH(Ph)₂, fluoro
- 20 (C₁-C₄)alkoxy and fluoro (C₁-C₄)alkyl in a number ranging from zero to the total number of open valences on the aromatic ring system, where R, R', R'' and R''' are each independently selected from the group consisting of hydrogen, substituted or unsubstituted C₁-C₈ alkyl and heteroalkyl, substituted or unsubstituted C₃-C₈ cycloalkyl, unsubstituted aryl and
- 25 heteroaryl, (unsubstituted aryl)-(C₁-C₄) alkyl, and (unsubstituted aryl)oxy-(C₁-C₄)alkyl, and where R' and R'', together with the nitrogen atom to which they are attached, are optionally joined to form a 5- to 7-membered ring.

2. A compound according to claim 1 in which A is a member selected from the group consisting of substituted or unsubstituted aryl, and substituted or unsubstituted heteroaryl.

3. A compound according to claim 2 in which A is substituted or unsubstituted phenyl.

4. A compound according to claim 3 in which A is phenyl substituted by one or two groups selected from the group consisting of halogen, nitrile, substituted or unsubstituted C₁-C₄ alkyl, SCF₃, trifluoromethyl and trifluoromethoxy.

5

5. A compound according to claim 1 in which X is CO.

6. A compound according to claim 1 in which Z is a member selected from the group consisting of -CH₂- and -CH=CH-.

10

7. A compound according to claim 1 in which R¹ is a member selected from the group consisting of substituted or unsubstituted aryl, and substituted or unsubstituted heteroaryl.

15

8. A compound according to claim 1 in which R¹ is substituted or unsubstituted phenyl.

9. A compound according to claim 8 in which R¹ is a member selected from the group consisting of phenyl, and phenyl substituted with one or more of halogen, CF₃ and OCF₃.


20

10. A compound according to claim 1 in which R² is a substituted or unsubstituted C₁-C₆ saturated acyclic alkyl group.

25

11. A compound according to claim 10 in which R² is a C₁-C₄ saturated acyclic alkyl group.

12. The compound according to claim 1 having the formula:



30 in which

X is a member selected from the group consisting of CO, CS and SO₂;

Z is a member selected from the group consisting of -CH₂-, -CHF-, -CF₂- and -CH=CH- and -N(R⁴)(CR^{4a}R^{4b})_s-

wherein R^4 is a member selected from the group consisting of H and a substituted or unsubstituted C_1 - C_5 alkyl group;

R^{4a} and R^{4b} are members independently selected from the group consisting of H, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, substituted or unsubstituted C_3 - C_8 cycloalkyl, substituted or unsubstituted 5-7 membered heterocyclyl, and substituted or unsubstituted C_1 - C_8 alkyl;

s is an integer from 1 to 3;

Y is S;

R^1 is a member selected from the group consisting of substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, substituted or unsubstituted C_3 - C_8 cycloalkyl, substituted or unsubstituted 5-7 membered heterocyclyl, and substituted or unsubstituted C_1 - C_8 alkyl;

R^2 is a member selected from the group consisting of CF_3 , substituted or unsubstituted C_1 - C_8 alkyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, substituted or unsubstituted C_3 - C_8 cycloalkyl, substituted or unsubstituted 3-7 membered saturated heterocyclyl;

R^5 and R^6 are independently selected from the group consisting of H, halo, CF_3 , CF_3O , NO_2 , CN, $S(O)_mR^7$, $COOR^8$, $CONR^9R^{10}$, $SO_2NR^{11}R^{12}$, $S(O)_mCF_3$, substituted or unsubstituted C_1 - C_6 alkyl, and substituted or unsubstituted C_3 - C_7 cycloalkyl

wherein R^7 and R^8 are independently selected from the group consisting of substituted or unsubstituted C_1 - C_5 alkyl, and substituted or unsubstituted C_3 - C_7 cycloalkyl;

m is an integer from 0 to 2; and

R^9 , R^{10} , R^{11} and R^{12} are independently selected from the group consisting of H, substituted or unsubstituted C_1 - C_5 alkyl, substituted or unsubstituted C_3 - C_7 cycloalkyl, and R^9 and R^{10} or R^{11} and R^{12} , together with the nitrogen atom to which they are attached, are optionally joined to form a 5- to 7-membered ring.

13. The compound according to claim 12 wherein

X is a member selected from the group consisting of CO and SO_2 ;

Z is CH₂; and


R¹ is substituted or unsubstituted phenyl.

14. The compound according to claim 12 wherein

R² is a member selected from the group consisting of substituted or unsubstituted C₁-C₄ alkyl, substituted or unsubstituted C₃-C₆ cycloalkyl and substituted or unsubstituted C₃-C₆ heterocyclyl; and

R⁵ and R⁶ are members independently selected from the group consisting of H, halo, CF₃, OCF₃, substituted or unsubstituted C₁-C₅ alkyl, SCF₃ and CN.

15. A compound according to claim 1 having the formula:



in which

X is a member selected from the group consisting of CO, CS and SO₂;

Z is a member selected from the group consisting of -CH₂-, -CHF-, -CF₂- and -CH=CH- and -N(R⁴)(CR^{4a}R^{4b})_s-

wherein R⁴ is a member selected from the group consisting of H and a substituted or unsubstituted C₁-C₅ alkyl group;

R^{4a} and R^{4b} are members independently selected from the group consisting of H, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, substituted or unsubstituted C₃-C₈ cycloalkyl, substituted or unsubstituted 5-7 membered heterocyclyl, and substituted or unsubstituted C₁-C₈ alkyl;

s is an integer from 1 to 3;

R¹ is a member selected from the group consisting of substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, substituted or unsubstituted C₃-C₈ cycloalkyl, substituted or unsubstituted 5-7 membered heterocyclyl, substituted or unsubstituted C₁-C₈ alkyl;

Y is S;

R^2 is a member selected from the group consisting of CF_3 , substituted or unsubstituted C_1 - C_8 alkyl, substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, substituted or unsubstituted C_3 - C_8 cycloalkyl, substituted or unsubstituted 3-7 membered saturated heterocyclyl;

5 R^5 and R^6 are independently selected from the group consisting of halo, CF_3 , CF_3O , NO_2 , CN , $S(O)_mR^7COOR^8$, $CONR^9R^{10}$, $SO_2NR^{11}R^{12}$, $S(O)_mCF_3$, substituted or unsubstituted C_1 - C_6 alkyl, and substituted or unsubstituted C_3 - C_7 cycloalkyl wherein R^7 and R^8 are independently selected from the group consisting of substituted or unsubstituted C_1 - C_5 alkyl, and substituted or unsubstituted C_3 -
10 C_7 cycloalkyl;

m is an integer from 0 to 2; and

R^9 , R^{10} , R^{11} and R^{12} are independently selected from the group consisting of H, substituted or unsubstituted C_1 - C_5 alkyl, substituted or unsubstituted C_3 - C_7 cycloalkyl, and R^9 and R^{10} or R^{11} and R^{12} , together with the nitrogen atom to
15 which they are attached, are optionally joined to form a 5-to 7-membered ring.

16. The compound according to claim 15 wherein

X is a member selected from the group consisting of CO and SO_2 ;

20 Z is CH_2 ; and







R^1 is substituted or unsubstituted phenyl.







17. The compound according to claim 16 wherein







R^2 is a member selected from the group consisting of substituted or unsubstituted C_1 -
25 C_4 alkyl, substituted or unsubstituted C_3 - C_6 cycloalkyl and substituted or unsubstituted C_3 - C_7 heterocyclyl;







and R^5 and R^6 are members independently selected from the group consisting of halo, CF_3 , OCF_3 , substituted or unsubstituted C_1 - C_5 alkyl, SCF_3 and CN.







30 18. A compound of claim 1, wherein the compound has a structure selected from the following formulae:







1		2	
3		4	
5		6	







7		8	
9		10	
11		12	







13		14	
15		16	
17		18	







19		20	
21		22	
23		24	







25		26	
27		28	
29		30	







31	 <chem>COc1cccc(c1)SC2=C3N4C(=O)N(C2)C(=O)N4CC5CCCC5</chem>	32	 <chem>CCOC1=CC=C(C=C1)SC2=C3N4C(=O)N(C2)C(=O)N4CC5CCCC5</chem>
33	 <chem>CCOC1=CC=C(C=C1)SC2=C3N4C(=O)N(C2)C(=O)N4CC5CCCC5</chem>	34	 <chem>CCOC1=CC=C(C=C1)SC2=C3N4C(=O)N(C2)C(=O)N4CC5CCCC5</chem>
35	 <chem>NC(=O)CCSC2=C3N4C(=O)N(C2)C(=O)N4CC5CCCC5</chem>	36	 <chem>CCSC2=C3N4C(=O)N(C2)C(=O)N4CC5CCCC5</chem>

37		38	
39		40	
41		42	

43		44	
45		46	
47		48	







49	 <chem>CCCC(=O)N1C(=O)N2C(=N1)SCC3=CC=CC=C3N2C3=CC=CC=C3</chem>	50	 <chem>CCCC(=O)N1C(=O)N2C(=N1)SCC3=CC=CC=C3N2C4=CC=C(C=C4)F</chem>
51	 <chem>CCCC(=O)N1C(=O)N2C(=N1)SC3=CC=CC=C3N2C4=CC=C(C=C4)C(F)(F)F</chem>	52	 <chem>CCCC(=O)N1C(=O)N2C(=N1)SC3=CC=CC=C3N2C4=CC=C(C=C4)C(F)(F)F</chem>
53	 <chem>CCCC(=O)N1C(=O)N2C(=N1)SC(C)C3=CC=CC=C3N2C4=CC=C(C=C4)C(F)(F)F</chem>	54	 <chem>CCCC(=O)N1C(=O)N2C(=N1)SC3=CC=CC=C3N2C4=CC=C(C=C4)F</chem>







55	 <chem>CCSC1=NC2=CC=C(C=C2C(=O)N1N(C(=O)CC3=CC=C(C=C3)F)C)C=C3C=CC(=C3)F</chem>	56	 <chem>CCSC1=NC2=CC=C(C=C2C(=O)N1N(C(=O)CC3=CC=C(C=C3)F)C)C=C3C=CC(=C3)F</chem>
57	 <chem>CS1=NC2=CC=C(C=C2C(=O)N1N(C(=O)CC3=CC(=CC=C3)F)C)C=C3C=CC(=C3)F</chem>	58	 <chem>CS1=NC2=CC=C(C=C2C(=O)N1N(C(=O)CC3=CC=C(C=C3)F)C)C=C3C=CC(=C3)F</chem>
59	 <chem>CC(C)SC1=NC2=CC=C(C=C2C(=O)N1N(C(=O)CC3=CC=C(C=C3)F)C)C=C3C=CC(=C3)F</chem>	60	 <chem>CS1=NC2=CC=C(C=C2C(=O)N1N(C(=O)CC3=CC=C(C=C3)F)C)C=C3C=CC(=C3)F</chem>







61	 <chem>CC1=NC2=C(N1C(=O)c3ccc(F)cc3)N(CS2)C(=O)CCc4ccc(F)cc4</chem>	62	 <chem>CC1=NC2=C(N1C(=O)c3ccc(C(F)(F)F)cc3)N(CS2)C(=O)CCc4ccc(F)cc4</chem>
63	 <chem>CC1=NC2=C(N1C(=O)c3ccc(C(F)(F)F)cc3)N(CS2)C(=O)CCc4ccc(F)cc4</chem>	64	 <chem>CC1=NC2=C(N1C(=O)c3ccccc3)N(CS2)C(=O)CCc4ccc(F)cc4</chem>
65	 <chem>CC1=NC2=C(N1C(=O)c3ccccc3)N(CSC2)C(=O)CCc4ccc(F)cc4</chem>	66	 <chem>CC1=NC2=C(N1C(=O)c3cc(F)ccc3)N(CS2)C(=O)CCc4ccc(F)cc4</chem>

67	<chem>CCSC1=NC2=CC=CC=C2C(=O)N1NCC(=O)Cc3ccc(F)cc3</chem>	68	<chem>CS1=NC2=CC=CC=C2C(=O)N1NCC(=O)Cc3ccccc3</chem>
69	<chem>CCSC1=NC2=CC=CC=C2C(=O)N1NCC(=O)Cc3ccccc3</chem>	70	<chem>CCSC1=NC2=CC=C(C(=O)S(=O)(=O)C)C(=O)N1NCC(=O)Cc3ccccc3</chem>
71	<chem>CC(C)SC1=NC2=CC(F)=CC=C2C(=O)N1NCC(=O)Cc3ccc(F)cc3</chem>	72	<chem>CS1=NC2=CC=CC=C2C(=O)N1NCC(=O)Cc3ccc(F)cc3</chem>







73	 <chem>CCSC1=NC2=CC=C(C=C2C(=O)N1N(C(=O)CC3=CC=C(C=C3)F)F)C4=CC=C(C=C4)F</chem>	74	 <chem>CSC1=NC2=CC=C(C=C2C(=O)N1N(C(=O)CC3=CC=C(C=C3)F)F)C4=CC=C(C=C4)F</chem>
75	 <chem>CC(C)SC1=NC2=CC=CC=C2C(=O)N1N(C(=O)CC3=CC=C(C=C3)F)C4=CC=CC=C4</chem>	76	 <chem>CCSC1=NC2=CC=C(C=C2C(=O)N1N(C(=O)CC3=CC=C(C=C3)F)F)C4=CC=CC=C4F</chem>
77	 <chem>CC(C)SC1=NC2=CC=C(C=C2C(=O)N1N(C(=O)CC3=CC=C(C=C3)F)F)C4=CC=CC=C4F</chem>	78	 <chem>C1CC1SC2=NC3=CC=C(C=C3C(=O)N2N(C(=O)CC4=CC=CC=C4)F)C5=CC=C(C=C5)F</chem>

79		80	
81		82	
83		84	







85		86	
87		88	
89		90	







91	 <chem>CC(C)S=C1N=C2C(=O)N(C1)C(=O)CCc3ccc(F)cc3C2(F)F</chem>	92	 <chem>CC(C)S=C1N=C2C(=O)N(C1)C(=O)CCc3ccc(F)cc3C2=C3SC=C(C)C3</chem>
93	 <chem>CC(C)S=C1N=C2C(=O)N(C1)C(=O)CCc3ccccc3C2=C3SC=C(C)C3</chem>	94	 <chem>CC(C)S=C1N=C2C(=O)N(C1)C(=O)CCc3ccc(F)cc3C2=C3SC=C(C)C3</chem>
95	 <chem>CC(C)S=C1N=C2C(=O)N(C1)C(=O)CCc3ccc(F)cc3C2=NC3=CC=CC=C3</chem>	96	 <chem>CCSC1=NC2=CC=C(C=C2)C(=O)N(C1)C(=O)CCc3ccc(OC(F)F)cc3C4=CC=CC=C4F</chem>







97	Chemical structure 97: A quinazolinone derivative. The quinazolinone core has a fluorine atom at position 6. The 2-position is substituted with an isopropylthio group (-S-CH(CH₃)₂). The 4-position is substituted with a 4-(trifluoromethoxy)benzyl group (-CH₂-C₆H₄-OCF₃).	98	Chemical structure 98: A quinazolinone derivative. The quinazolinone core has a fluorine atom at position 6. The 2-position is substituted with a methylthio group (-SCH₃). The 4-position is substituted with a 4-(trifluoromethoxy)benzyl group (-CH₂-C₆H₄-OCF₃).
99	Chemical structure 99: A quinazolinone derivative. The quinazolinone core has a fluorine atom at position 6. The 2-position is substituted with an isopropylthio group (-S-CH(CH₃)₂). The 4-position is substituted with a 4-(trifluoromethoxy)benzyl group (-CH₂-C₆H₄-OCF₃).	100	Chemical structure 100: A quinazolinone derivative. The quinazolinone core has a fluorine atom at position 6. The 2-position is substituted with an ethylthio group (-SCH₂CH₃). The 4-position is substituted with a benzyl group (-CH₂-C₆H₅).
101	Chemical structure 101: A quinazolinone derivative. The quinazolinone core has a fluorine atom at position 6. The 2-position is substituted with an ethylthio group (-SCH₂CH₃). The 4-position is substituted with a 4-(trifluoromethoxy)benzyl group (-CH₂-C₆H₄-OCF₃).	102	Chemical structure 102: A quinazolinone derivative. The quinazolinone core has a fluorine atom at position 6. The 2-position is substituted with an ethylthio group (-SCH₂CH₃). The 4-position is substituted with a 4-(trifluoromethoxy)benzyl group (-CH₂-C₆H₄-OCF₃).







103		104	
105		106	
107		108	







109	<chem>CC(C)S=C1N=C2C(=O)N(C1)C(=O)C2c3ccccc3F</chem>	110	<chem>CC(C)S=C1N=C2C(=O)N(C1)C(=O)C2c3ccc(F)cc3</chem>
111	<chem>CCOCCSC=C1N=C2C(=O)N(C1)C(=O)C2c3ccccc3F</chem>	112	<chem>CC(C)S=C1N=C2C(=O)N(C1)C(=O)C2c3cc(F)(F)Fcc3</chem>
113	<chem>CCSC=C1N=C2C(=O)N(C1)C(=O)C2c3cc(F)(F)Fcc3</chem>	114	<chem>CCSC=C1N=C2C(=O)N(C1)C(=O)C2c3ccccc3F</chem>







115	 <chem>CC(C)SC1=NC2=CC=CC=C2C(=O)N1N(C(=O)CCc3ccccc3)C4=CC=CC=C4F</chem>	116	 <chem>CCSC1=NC2=CC=CC=C2C(=O)N1N(C(=O)CCc3ccc(F)cc3)C4=CC=CC=C4F</chem>
117	 <chem>CC(C)SC1=NC2=CC=CC=C2C(=O)N1N(C(=O)CCc3ccc(F)cc3)C4=CC=CC=C4F</chem>	118	 <chem>CCSC1=NC2=CC=CC=C2C(=O)N1N(C(=O)CCc3ccccc3)C4=CC=CC=C4N(C)C5=CC=CC=C5</chem>
119	 <chem>CCSC1=NC2=CC=CC=C2C(=O)N1N(C(=O)CCc3ccccc3)C4=CC=CC=C4N(C)C5=CC=CC=C5</chem>	120	 <chem>CC(C)SC1=NC2=CC=CC=C2C(=O)N1N(C(=O)CCc3ccccc3)C4=CC=CC=C4N(C)C5=CC=CC=C5</chem>







121	 <chem>CC(C)S=C1N=CN2C(=O)N(C1=CC=CC2)C(=O)CCc3ccccc3</chem>	122	 <chem>CCSC=C1N=CN2C(=O)N(C1=CCCCC2)C(=O)CCc3ccccc3</chem>
123	 <chem>CCSC=C1N=CN2C(=O)N(C1=CC=C(C2)C(F)(F)F)C(=O)CCc3ccc(F)cc3</chem>	124	 <chem>CCSC=C1N=CN2C(=O)N(C1=CC=C(C2)C(F)(F)F)C(=O)CCc3ccccc3</chem>
125	 <chem>CS=C1N=CN2C(=O)N(C1=CC=C(C2)OC(F)(F)F)C(=O)CCc3ccc(F)cc3</chem>	126	 <chem>CS=C1N=CN2C(=O)N(C1=CC=C(C2)OC(F)(F)F)C(=O)CCc3ccc(F)cc3</chem>

127	 <chem>CCSC1=NC2=CC=CC=C2C(=O)N1N3C(=O)CCc4ccc(F)cc43</chem>	128	 <chem>CC(C)SC1=NC2=CC=CC=C2C(=O)N1N3C(=O)CCc4ccc(F)cc43</chem>
129	 <chem>CC(C)SC1=NC2=CC(OC(F)(F)F)=CC=C2C(=O)N1N3C(=O)CCc4ccc(F)cc43</chem>	130	 <chem>CC(C)SC1=NC2=CC(OC(F)(F)F)=CC=C2C(=O)N1N3C(=O)CCc4ccc(F)cc43</chem>
131	 <chem>CC(C)SC1=NC2=CC(OC(F)(F)F)=CC=C2C(=O)N1N3C(=O)CCc4ccccc43</chem>	132	 <chem>CSC1=NC2=CC(F)=CC(F)=C2C(=O)N1N3C(=O)CCc4ccccc43</chem>


133	 <chem>CC(C)S=C1N=C2C(=O)N(C1)N(C2)C(=O)CCc3ccccc3Oc4ccccc4F(F)F</chem>	134	 <chem>CC(C)S=C1N=C2C(=O)N(C1)N(C2)C(=O)CCc3ccc(F)cc3Oc4ccccc4F(F)F</chem>
135	 <chem>CS=C1N=C2C(=O)N(C1)N(C2)C(=O)CCc3ccccc3Oc4ccccc4F(F)F</chem>	136	 <chem>CCSC=C1N=C2C(=O)N(C1)N(C2)C(=O)CCc3ccccc3Oc4ccccc4F(F)F</chem>
137	 <chem>CC(C)S=C1N=C2C(=O)N(C1)N(C2)C(=O)CCc3ccccc3Oc4ccccc4OC</chem>	138	 <chem>CS=C1N=C2C(=O)N(C1)N(C2)C(=O)CCc3ccc(F)cc3Oc4ccccc4F(F)F</chem>

139		140	
141		142	
143		144	

145	 <chem>CC(C)S[C@H]1C(=O)N2C(=N1)C(F)=C(F)C=C2N(C2=O)N(C2=O)CC3=CC=CC=C3</chem>	146	 <chem>CC(F)(F)FSC1=NC2=CC=CC=C2C(=O)N1N(C1=O)CC3=CC=CC=C3</chem>
147	 <chem>CC1=C(C)SC=C1[C@H]2C(=O)N3C(=N2)SC(C3=O)N(C3=O)CC4=CC=CC=C4</chem>	148	 <chem>CO[C@H]1C=CC=C1C2=CC(=O)N3C(=N2)SC(C3=O)N(C3=O)CC4=CC=CC=C4</chem>
149	 <chem>CC(F)(F)FSC1=NC2=CC(OC)=CC=C2C(=O)N1N(C1=O)CC3=CC=CC=C3</chem>	150	 <chem>CC(C)S[C@H]1C(=O)N2C(=N1)C(F)=CC=C2N(C2=O)N(C2=O)CC3=CC=CC=C3</chem>

151		152	
153		154	
155		156	


157



19. The compound of claim 18, wherein the compound is selected from compounds 38, 40, 41, 42, 44, 45, 46, 47, 48, 50, 51, 52, 53, 54, 55, 56, 57, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 73 or 74.


5

20. A compound of claim 1, wherein the compound has a structure selected from the following formulae:



10

21. A compound of claim 1, having the formula:



22. A composition comprising a pharmaceutically acceptable excipient and a compound according to any one of claims 1-17.

23. A composition for increasing ion flow in a voltage-dependent potassium channel, said composition comprising a pharmaceutically acceptable excipient and a compound according to any one of claims 1-17.

5

24. A method of increasing ion flow through voltage-dependent potassium channels in a cell, said method comprising contacting said cell *in vitro* with a potassium channel-opening amount of a compound according to any one of claims 1-17.

10

25. The method according to claim 24, wherein said voltage-dependent potassium channel is responsible for the M-current.

15

26. A method of increasing ion flow through KCNQ voltage-dependent potassium channels in a cell, said method comprising contacting said cell *in vitro* with a potassium channel-opening amount of a compound according to any one of claims 1-17.

20

27. Use of a compound according to any one of claims 1-17 for the preparation of a medicament for modulation of a KCNQ voltage-dependent potassium channel in a subject.

28. Use of a compound according to any one of claims 1-17 for modulation of a KCNQ voltage-dependent potassium channel in a subject.

25

29. A compound according to any one of claims 1-17 for use in modulation of a KCNQ voltage-dependent potassium channel in a subject.

30

30. Use of a compound according to any one of claims 1-17 for the preparation of a medicament for treating a central or peripheral nervous system disorder or condition in a subject.

31. Use of a compound according to any one of claims 1-17 for treating a central or peripheral nervous system disorder or condition in a subject.

32. A compound according to any one of claims 1-17 for use in treating a central or peripheral nervous system disorder or condition in a subject.

33. The use according to claim 30 or 31 wherein said disorder or condition is selected from the group consisting of migraine, ataxia, Parkinson's disease, bipolar disorders, spasticity, mood disorders, brain tumors, psychotic disorders, myokymia, seizures, epilepsy, hearing loss, vision loss, Alzheimer's disease, age-related memory loss, learning deficiencies, motor neuron diseases, trigeminal neuralgia, retinal degeneration, and stroke.

34. The compound according to claim 32 wherein said disorder or condition is selected from the group consisting of migraine, ataxia, Parkinson's disease, bipolar disorders, spasticity, mood disorders, brain tumors, psychotic disorders, myokymia, seizures, epilepsy, hearing loss, vision loss, Alzheimer's disease, age-related memory loss, learning deficiencies, motor neuron diseases, trigeminal neuralgia, retinal degeneration and stroke.

35. Use of a compound according to any one of claims 1-17 for the preparation of a medicament for modulation of a voltage-dependent potassium channel in a subject.

36. Use of a compound according to any one of claims 1-17 for modulation of a voltage-dependent potassium channel in a subject.

37. A compound according to any one of claims 1-17 for modulation of a voltage-dependent potassium channel in a subject.

38. Use of a compound according to any one of claims 1-17 for the preparation of a medicament for treating epilepsy, retinal degeneration, pain, anxiety, or bipolar disorder in a subject.

39. Use of a compound according to any one of claims 1-17 for treating epilepsy, retinal degeneration, pain, anxiety or bipolar disorder in a subject.

40. A compound according to any one of claims 1-17 for use in treating epilepsy, retinal degeneration, pain, anxiety or bipolar disorder in a subject.

5 41. The use according to claim 38 or 39 wherein said pain is a member selected from the group consisting of neuropathic pain, diabetic pain, inflammatory pain, cancer pain, migraine pain, or musculoskeletal pain.

10 42. The compound according to claim 40 wherein said pain is neuropathic pain, diabetic pain, inflammatory pain, cancer pain, migraine pain, or musculoskeletal pain.

43. The use according to claim 30 or 31 wherein said condition or disorder is epilepsy or seizures.

15 44. The compound according to claim 32 wherein said condition or disorder is epilepsy or seizures.

45. The use according to claim 30 or 31 wherein said condition or disorder is hearing loss.

20 46. The use according to claim 33 wherein said condition or disorder is hearing loss.

25 47. The use according to claim 30 or 31 wherein said condition or disorder is pain or anxiety.

48. The compound according to claim 32 wherein said condition or disorder is pain or anxiety.

30 49. The use according to claim 30 or 31 wherein said condition or disorder is retinal degeneration.

50. The compound according to claim 32 wherein said condition or disorder is retinal degeneration.

51. A packaged preparation for use in modulation of a voltage-dependent potassium channel in a subject, the packaged preparation comprising a compound according to any one of claims 1-17 and instructions for use.

5

52. A packaged preparation for use in modulation of a KCNQ voltage-dependent potassium channel in a subject, the packaged preparation comprising a compound according to any one of claims 1-17 and instructions for use.

10

53. A packaged preparation for use in treating a central or peripheral nervous system disorder or condition in a subject, the packaged preparation comprising a compound according to any one of claims 1-17 and instructions for use.

54. The packaged preparation of claim 53 wherein said disorder or condition is selected from the group consisting of migraine, ataxia, Parkinson's disease, bipolar disorders, spasticity, mood disorders, brain tumors, psychotic disorders, myokymia, seizures, epilepsy, hearing loss, vision loss, Alzheimer's disease, age-related memory loss, learning deficiencies, motor neuron diseases, trigeminal neuralgia, retinal degeneration, and stroke.

20

55. The packaged preparation of claim 53 wherein said condition or disorder is epilepsy or seizures.

56. The packaged preparation of claim 53 wherein said condition or disorder is hearing loss.

25

57. The packaged preparation of claim 53 wherein said condition or disorder is pain or anxiety.

30

58. The packaged preparation of claim 53 wherein said condition or disorder is retinal degeneration.

59. A packaged preparation for use in treating epilepsy, retinal degeneration, pain, anxiety, or bipolar disorder in a subject, the commercial package comprising a compound according to any one of claims 1-17 and instructions for use.

5 60. The packaged preparation of claim 59 wherein said pain is neuropathic pain, diabetic pain, inflammatory pain, cancer pain, migraine pain, or musculoskeletal pain.

