

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
13 July 2006 (13.07.2006)

PCT

(10) International Publication Number
WO 2006/074199 A1

(51) International Patent Classification:

A61P 9/00 (2006.01) C07K 19/00 (2006.01)
 A61P 9/10 (2006.01) C07K 14/505 (2006.01)
 A61K 38/17 (2006.01) C07K 14/56 (2006.01)
 A61K 38/18 (2006.01) C07K 14/565 (2006.01)
 A61K 38/21 (2006.01) C12N 9/64 (2006.01)
 A61K 38/48 (2006.01)

(21) International Application Number:

PCT/US2006/000140

(22) International Filing Date: 4 January 2006 (04.01.2006)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

11/029,003 5 January 2005 (05.01.2005) US

(71) Applicant (for all designated States except US): SYNTONIX PHARMACEUTICALS, INC. [US/US]; 9 Fourth Avenue, Waltham, Massachusetts 02461 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): PETERS, Robert, T. [US/US]; 51 Newfield Street, West Roxbury, Massachusetts 02132 (US). MEZO, Adam, R. [CA/US]; 15 Montclair Avenue, Waltham, Massachusetts 02451 (US). RIVERA, Daniel, S. [US/US]; 100 Providence Place,

#451, Providence, Rhode Island 02903 (US). BITONTI, Alan, J. [US/US]; 32 Carlton Drive, Acton, Massachusetts 01720 (US).


(74) Agent: MCDONELL, Leslie, A.; Finnegan, Henderson, Farabow, Garrett & Dunner, L.L.P., 901 New York Avenue, Washington, DC 20001-4413 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

[Continued on next page]

(54) Title: IMMUNOGLOBULIN CHIMERIC MONOMER-DIMER HYBRIDS



(57) Abstract: The invention relates to a chimeric monomer-dimer hybrid protein wherein said protein comprises a first and a second polypeptide chain, said first polypeptide chain comprising at least a portion of an immunoglobulin constant region and a biologically active molecule, and said second polypeptide chain comprising at least a portion of an immunoglobulin constant region without the biologically active molecule of the first chain. The invention also relates to methods of using and methods of making the chimeric monomer-dimer hybrid protein of the invention.



Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

IMMUNOGLOBULIN CHIMERIC MONOMER-DIMER HYBRIDS

RELATED APPLICATIONS

[001] This application claims priority to U.S. Application No. 11/029,003, filed on January 5, 2005, and is incorporated herein by reference.

FIELD OF THE INVENTION

[002] The invention relates generally to therapeutic chimeric proteins, comprised of two polypeptide chains, wherein the first chain is comprised of a therapeutic biologically active molecule and the second chain is not comprised of the therapeutic biologically active molecule of the first chain. More specifically, the invention relates to chimeric proteins, comprised of two polypeptide chains, wherein both chains are comprised of at least a portion of an immunoglobulin constant region wherein the first chain is modified to further comprise a biologically active molecule, and the second chain is not so modified. The invention, thus relates to a chimeric protein that is a monomer-dimer hybrid, *i.e.*, a chimeric protein having a dimeric aspect and a monomeric aspect, wherein the dimeric aspect relates to the fact that it is comprised of two polypeptide chains each comprised of a portion of an immunoglobulin constant region, and wherein the monomeric aspect relates to the fact that only one of the two chains is comprised of a therapeutic biologically active molecule. Figure 1 illustrates one example of a monomer-dimer hybrid wherein the biologically active molecule is erythropoietin (EPO) and the portion of an immunoglobulin constant region is an IgG Fc region.

BACKGROUND OF THE INVENTION

[003] Immunoglobulins are comprised of four polypeptide chains, two heavy chains and two light chains, which associate via disulfide bonds to form tetramers. Each chain is further comprised of one variable region and one constant region. The variable regions mediate antigen recognition and binding, while the constant regions, particularly the heavy chain constant regions, mediate a variety of effector functions, *e.g.*, complement binding and Fc receptor binding (see, *e.g.*, U.S. Patent Nos.: 6,086,875; 5,624,821; 5,116,964).

[004] The constant region is further comprised of domains denoted CH (constant heavy) domains (CH1, CH2, etc.). Depending on the isotype, (*i.e.* IgG,

IgM, IgA IgD, IgE) the constant region can be comprised of three or four CH domains. Some isotypes (e.g. IgG) constant regions also contain a hinge region Janeway et al. 2001, *Immunobiology*, Garland Publishing, N.Y., N.Y.

[005] The creation of chimeric proteins comprised of immunoglobulin constant regions linked to a protein of interest, or fragment thereof, has been described (see, e.g., U.S. Patent Nos. 5,480,981 and 5,808,029; Gascoigne et al. 1987, *Proc. Natl. Acad. Sci. USA* 84:2936; Capon et al. 1989, *Nature* 337:525; Traunecker et al. 1989, *Nature* 339:68; Zettmeissl et al. 1990, *DNA Cell Biol. USA* 9:347; Byrn et al. 1990, *Nature* 344:667; Watson et al. 1990, *J. Cell. Biol.* 110:2221; Watson et al. 1991, *Nature* 349:164; Aruffo et al. 1990, *Cell* 61:1303; Linsley et al. 1991, *J. Exp. Med.* 173:721; Linsley et al. 1991, *J. Exp. Med.* 174:561; Stamenkovic et al., 1991, *Cell* 66:1133; Ashkenazi et al. 1991, *Proc. Natl. Acad. Sci. USA* 88:10535; Lesslauer et al. 1991, *Eur. J. Immunol.* 27:2883; Peppel et al. 1991, *J. Exp. Med.* 174:1483; Bennett et al. 1991, *J. Biol. Chem.* 266:23060; Kurschner et al. 1992, *J. Biol. Chem.* 267:9354; Chalupny et al. 1992, *Proc. Natl. Acad. Sci. USA* 89:10360; Ridgway and Gorman, 1991, *J. Cell. Biol.* 115, Abstract No. 1448; Zheng et al. 1995, *J. Immun.* 154:5590). These molecules usually possess both the biological activity associated with the linked molecule of interest as well as the effector function, or some other desired characteristic associated with the immunoglobulin constant region (e.g. biological stability, cellular secretion).

[006] The Fc portion of an immunoglobulin constant region, depending on the immunoglobulin isotype can include the CH2, CH3, and CH4 domains, as well as the hinge region. Chimeric proteins comprising an Fc portion of an immunoglobulin bestow several desirable properties on a chimeric protein including increased stability, increased serum half life (see Capon et al. 1989, *Nature* 337:525) as well as binding to Fc receptors such as the neonatal Fc receptor (FcRn) (U.S. Patent Nos. 6,086,875, 6,485,726, 6,030,613; WO 03/077834; US2003-0235536A1).

[007] FcRn is active in adult epithelial tissue and expressed in the lumen of the intestines, pulmonary airways, nasal surfaces, vaginal surfaces, colon and rectal surfaces (U.S. Patent No. 6,485,726). Chimeric proteins comprised of FcRn binding partners (e.g. IgG, Fc fragments) can be effectively shuttled across

epithelial barriers by FcRn, thus providing a non-invasive means to systemically administer a desired therapeutic molecule. Additionally, chimeric proteins comprising an FcRn binding partner are endocytosed by cells expressing the FcRn. But instead of being marked for degradation, these chimeric proteins are recycled out into circulation again, thus increasing the in vivo half life of these proteins.

[008] Portions of immunoglobulin constant regions, *e.g.*, FcRn binding partners typically associate, via disulfide bonds and other non-specific interactions, with one another to form dimers and higher order multimers. The instant invention is based in part upon the surprising discovery that transcytosis of chimeric proteins comprised of FcRn binding partners appears to be limited by the molecular weight of the chimeric protein, with higher molecular weight species being transported less efficiently.

[009] Chimeric proteins comprised of biologically active molecules, once administered, typically will interact with a target molecule or cell. The instant invention is further based in part upon the surprising discovery that monomer-dimer hybrids, with one biologically active molecule, but two portions of an immunoglobulin constant region, *e.g.*, two FcRn binding partners, function and can be transported more effectively than homodimers, also referred to herein simply as "dimers" or higher order multimers with two or more copies of the biologically active molecule. This is due in part to the fact that chimeric proteins, comprised of two or more biologically active molecules, which exist as dimers and higher order multimers, can be sterically hindered from interacting with their target molecule or cell, due to the presence of the two or more biologically active molecules in close proximity to one another and that the biologically active molecule can have a high affinity for itself.

[010] Accordingly one aspect of the invention provides chimeric proteins comprised of a biologically active molecule that is transported across the epithelium barrier. An additional aspect of the invention provides chimeric proteins comprised of at least one biologically active molecule that is able to interact with its target molecule or cell with little or no steric hindrance or self aggregation.

[011] The aspects of the invention provide for chimeric proteins comprising a first and second polypeptide chain, the first chain comprising at least a portion of immunoglobulin constant region, wherein the portion of an immunoglobulin constant region has been modified to include a biologically active molecule and the second chain comprising at least a portion of immunoglobulin constant region, wherein the portion of an immunoglobulin constant region has not been so modified to include the biologically active molecule of the first chain.

SUMMARY OF THE INVENTION

[012] The invention relates to a chimeric protein comprising one biologically active molecule and two molecules of at least a portion of an immunoglobulin constant region. The chimeric protein is capable of interacting with a target molecule or cell with less steric hindrance compared to a chimeric protein comprised of at least two biologically active molecules and at least a portion of two immunoglobulin constant regions. The invention also relates to a chimeric protein comprising at least one biologically active molecule and two molecules of at least a portion of an immunoglobulin constant region that is transported across an epithelium barrier more efficiently than a corresponding homodimer, *i.e.*, wherein both chains are linked to the same biologically active molecule. The invention, thus relates to a chimeric protein comprising a first and a second polypeptide chain linked together, wherein said first chain comprises a biologically active molecule and at least a portion of an immunoglobulin constant region, and said second chain comprises at least a portion of an immunoglobulin constant region, but no immunoglobulin variable region and without any biologically active molecule attached.

[013] The invention relates to a chimeric protein comprising a first and a second polypeptide chain linked together, wherein said first chain comprises a biologically active molecule and at least a portion of an immunoglobulin constant region, and said second chain comprises at least a portion of an immunoglobulin constant region without an immunoglobulin variable region or any biologically active molecule and wherein said second chain is not covalently bonded to any molecule having a molecular weight greater than 1 kD, 2 kD, 5 kD, 10 kD, or 20 kD. In one embodiment, the second chain is not covalently bonded to any molecule having a molecular weight greater than 0-2 kD. In one embodiment, the

second chain is not covalently bonded to any molecule having a molecular weight greater than 5-10 kD. In one embodiment, the second chain is not covalently bonded to any molecule having a molecular weight greater than 15-20 kD.

[014] The invention relates to a chimeric protein comprising a first and a second polypeptide chain linked together, wherein said first chain comprises a biologically active molecule and at least a portion of an immunoglobulin constant region, and said second chain comprises at least a portion of an immunoglobulin constant region not covalently linked to any other molecule except the portion of an immunoglobulin of said first polypeptide chain.

[015] The invention relates to a chimeric protein comprising a first and a second polypeptide chain linked together, wherein said first chain comprises a biologically active molecule and at least a portion of an immunoglobulin constant region, and said second chain consists of at least a portion of an immunoglobulin constant region and optionally an affinity tag.

[016] The invention relates to a chimeric protein comprising a first and a second polypeptide chain linked together, wherein said first chain comprises a biologically active molecule and at least a portion of an immunoglobulin constant region, and said second chain consists essentially of at least a portion of an immunoglobulin constant region and optionally an affinity tag.

[017] The invention relates to a chimeric protein comprising a first and a second polypeptide chain linked together, wherein said first chain comprises a biologically active molecule and at least a portion of an immunoglobulin constant region, and said second chain comprises at least a portion of an immunoglobulin constant region without an immunoglobulin variable region or any biologically active molecule and optionally a molecule with a molecular weight less than 10 kD, 5 kD, 2 kD or 1 kD. In one embodiment, the second chain comprises a molecule less than 15-20 kD. In one embodiment, the second chain comprises a molecule less than 5-10 kD. In one embodiment, the second chain comprises a molecule less than 1-2 kD.

[018] The invention relates to a chimeric protein comprising a first and second polypeptide chain, wherein said first chain comprises a biologically active molecule, at least a portion of an immunoglobulin constant region, and at least a first domain, said first domain having at least one specific binding partner, and

wherein said second chain comprises at least a portion of an immunoglobulin constant region, and at least a second domain, wherein said second domain is a specific binding partner of said first domain, without any immunoglobulin variable region or a biologically active molecule.

[019] The invention relates to a method of making a chimeric protein comprising a first and second polypeptide chain, wherein the first polypeptide chain and the second polypeptide chain are not the same, said method comprising transfecting a cell with a first DNA construct comprising a DNA molecule encoding a first polypeptide chain comprising a biologically active molecule and at least a portion of an immunoglobulin constant region and optionally a linker, and a second DNA construct comprising a DNA molecule encoding a second polypeptide chain comprising at least a portion of an immunoglobulin constant region without any biologically active molecule or an immunoglobulin variable region, and optionally a linker, culturing the cells under conditions such that the polypeptide chain encoded by the first DNA construct is expressed and the polypeptide chain encoded by the second DNA construct is expressed and isolating monomer-dimer hybrids comprised of the polypeptide chain encoded by the first DNA construct and the polypeptide chain encoded by the second DNA construct.

[020] The invention relates to a method of making a chimeric protein comprising a first and second polypeptide chain, wherein the first polypeptide chain and the second polypeptide chain are not the same, and wherein said first polypeptide chain comprises a biologically active molecule, at least a portion of an immunoglobulin constant region, and at least a first domain, said first domain, having at least one specific binding partner, and wherein said second polypeptide chain comprises at least a portion of an immunoglobulin constant region and a second domain, wherein said second domain, is a specific binding partner of said first domain, without any biologically active molecule or an immunoglobulin variable region, said method comprising transfecting a cell with a first DNA construct comprising a DNA molecule encoding said first polypeptide chain and a second DNA construct comprising a DNA molecule encoding, said second polypeptide chain, culturing the cells under conditions such that the polypeptide chain encoded by the first DNA construct is expressed and the polypeptide chain

encoded by the second DNA construct is expressed and isolating monomer-dimer hybrids comprised of the polypeptide chain encoded by the first DNA construct and polypeptide chain encoded by the second DNA construct.

[021] The invention relates to a method of making a chimeric protein of the invention said method comprising transfecting a cell with a first DNA construct comprising a DNA molecule encoding a first polypeptide chain comprising a biologically active molecule and at least a portion of an immunoglobulin constant region and optionally a linker, culturing the cell under conditions such that the polypeptide chain encoded by the first DNA construct is expressed, isolating the polypeptide chain encoded by the first DNA construct and transfecting a cell with a second DNA construct comprising a DNA molecule encoding a second polypeptide chain comprising at least a portion of an immunoglobulin constant region without any biologically active molecule or immunoglobulin variable region, culturing the cell under conditions such that the polypeptide chain encoded by the second DNA construct is expressed, isolating the polypeptide chain, encoded by the second DNA construct, combining the polypeptide chain, encoded by the first DNA construct and the polypeptide chain encoded by the second DNA construct under conditions such that monomer-dimer hybrids comprising the polypeptide chain encoded by the first DNA construct and the polypeptide chain encoded by the second DNA construct form, and isolating said monomer-dimer hybrids.

[022] The invention relates to a method of making a chimeric protein comprising a first and second polypeptide chain, wherein the first polypeptide chain and the second polypeptide chain are not the same, said method comprising transfecting a cell with a DNA construct comprising a DNA molecule encoding a polypeptide chain comprising at least a portion of an immunoglobulin constant region, culturing the cells under conditions such that the polypeptide chain encoded by the DNA construct is expressed with an N terminal cysteine such that dimers of the polypeptide chain form and isolating dimers comprised of two copies of the polypeptide chain encoded by the DNA construct and chemically reacting the isolated dimers with a biologically active molecule, wherein said biologically active molecule has a C terminus thioester, under conditions such that the biologically active molecule reacts predominantly with only one polypeptide chain of the dimer thereby forming a monomer-dimer hybrid.

[023] The invention relates to a method of making a chimeric protein comprising a first and second polypeptide chain, wherein the first polypeptide chain and the second polypeptide chain are not the same, said method comprising transfecting a cell with a DNA construct comprising a DNA molecule encoding a polypeptide chain comprising at least a portion of an immunoglobulin constant region, culturing the cells under conditions such that the polypeptide chain encoded by the DNA construct is expressed with an N terminal cysteine such that dimers of the polypeptide chains form, and isolating dimers comprised of two copies of the polypeptide chain encoded by the DNA construct, and chemically reacting the isolated dimers with a biologically active molecule, wherein said biologically active molecule has a C terminus thioester, such that the biologically active molecule is linked to each chain of the dimer, denaturing the dimer comprised of the portion of the immunoglobulin linked to the biologically active molecule such that monomeric chains form, combining the monomeric chains with a polypeptide chain comprising at least a portion of an immunoglobulin constant region without a biologically active molecule linked to it, such that monomer-dimer hybrids form, and isolating the monomer-dimer hybrids.

[024] The invention relates to a method of making a chimeric protein comprising a first and second polypeptide chain, wherein the first polypeptide chain and the second polypeptide chain are not the same, said method comprising transfecting a cell with a DNA construct comprising a DNA molecule encoding a polypeptide chain comprising at least a portion of an immunoglobulin constant region, culturing the cells under conditions such that the polypeptide chain encoded by the DNA construct is expressed as a mixture of two polypeptide chains, wherein the mixture comprises a polypeptide with an N terminal cysteine, and a polypeptide with a cysteine in close proximity to the N terminus, isolating dimers comprised of the mixture of polypeptide chains encoded by the DNA construct and chemically reacting the isolated dimers with a biologically active molecule, wherein said biologically active molecule has an active thioester, such that at least some monomer-dimer hybrid forms and isolating the monomer-dimer hybrid from said mixture.

[025] The invention relates to a method of treating a disease or condition comprising administering a chimeric protein of the invention thereby treating the disease or condition.

[026] Additional objects and advantages of the invention will be set forth in part in the description which follows, and in part will be obvious from the description, or may be learned by practice of the invention. The objects and advantages of the invention will be realized and attained by means of the elements and combinations particularly pointed out in the appended claims.

[027] It is to be understood that both the foregoing general description and the following detailed description are exemplary and explanatory only and are not restrictive of the invention, as claimed.

BRIEF DESCRIPTION OF THE DRAWINGS

[028] Figure 1 is a schematic diagram comparing the structure of an EPO-Fc homodimer, or dimer, and the structure of an Epo-FC monomer-dimer hybrid.

[029] Figure 2a is the amino acid sequence of the chimeric protein Factor VII-Fc. Included in the sequence is the signal peptide (underlined), which is cleaved by the cell and the propeptide (bold), which is recognized by the vitamin K-dependent γ carboxylase which modifies the Factor VII to achieve full activity. The sequence is subsequently cleaved by PACE to yield Factor VII-Fc.

[030] Figure 2b is the amino acid sequence of the chimeric protein Factor IX-Fc. Included in the sequence is the signal peptide (underlined) which is cleaved by the cell and the propeptide (bold) which is recognized by the vitamin K-dependent γ carboxylase which modifies the Factor IX to achieve full activity. The sequence is subsequently cleaved by PACE to yield Factor IX-Fc.

[031] Figure 2c is the amino acid sequence of the chimeric protein IFN α -Fc. Included in the sequence is the signal peptide (underlined), which is cleaved by the cell resulting in the mature IFN α -Fc.

[032] Figure 2d is the amino acid sequence of the chimeric protein IFN α -Fc Δ linker. Included in the sequence is the signal peptide (underlined) which is cleaved by the cell resulting in the mature IFN α -Fc Δ linker.

[033] Figure 2e is the amino acid sequence of the chimeric protein Flag-Fc. Included in the sequence is the signal peptide (underlined), which is cleaved by the cell resulting in the mature Flag-Fc.

[034] Figure 2f is the amino acid sequence of the chimeric protein Epo-CCA-Fc. Included in the sequence is the signal peptide (underlined), which is cleaved by the cell resulting in the mature Epo-CCA-Fc. Also shown in **bold** is the acidic coiled coil domain.

[035] Figure 2g is the amino acid sequence of the chimeric protein CCB-Fc. Included in the sequence is the signal peptide (underlined), which is cleaved by the cell resulting in the mature CCB-Fc. Also shown in **bold** is the basic coiled coil domain.

[036] Figure 2h is the amino acid sequence of the chimeric protein Cys-Fc. Included in the sequence is the signal peptide (underlined), which is cleaved by the cell resulting in the mature Cys-Fc. When this sequence is produced in CHO cells a percentage of the molecules are incorrectly cleaved by the signal peptidase such that two extra amino acids are left on the N terminus, thus preventing the linkage of a biologically active molecule with a C terminal thioester (e.g., via native ligation). When these improperly cleaved species dimerize with the properly cleaved Cys-Fc and are subsequently reacted with biologically active molecules with C terminal thioesters, monomer-dimer hybrids form.

[037] Figure 2i is the amino acid sequence of the chimeric protein IFN α -GS15-Fc. Included in the sequence is the signal peptide (underlined) which is cleaved by the cell resulting in the mature IFN α -GS15-Fc.

[038] Figure 2j is the amino acid sequence of the chimeric protein Epo-Fc. Included in the sequence is the signal peptide (underlined), which is cleaved by the cell resulting in the mature Epo-Fc. Also shown in **bold** is the 8 amino acid linker.

[039] Figure 3a is the nucleic acid sequence of the chimeric protein Factor VII-Fc. Included in the sequence is the signal peptide (underlined) and the propeptide (**bold**) which is recognized by the vitamin K-dependent γ carboxylase which modifies the Factor VII to achieve full activity. The translated sequence is subsequently cleaved by PACE to yield mature Factor VII-Fc.

[040] Figure 3b is the nucleic acid sequence of the chimeric protein Factor IX-Fc. Included in the sequence is the signal peptide (underlined) and the propeptide (**bold**) which is recognized by the vitamin K-dependent γ carboxylase which modifies the Factor IX to achieve full activity. The translated sequence is subsequently cleaved by PACE to yield mature Factor IX-Fc.

[041] Figure 3c is the nucleic acid sequence of the chimeric protein IFN α -Fc. Included in the sequence is the signal peptide (underlined), which is cleaved by the cell after translation resulting in the mature IFN α -Fc.

[042] Figure 3d is the nucleic acid sequence of the chimeric protein IFN α -Fc Δ linker. Included in the sequence is the signal peptide (underlined) which is cleaved by the cell after translation resulting in the mature IFN α -Fc Δ linker.

[043] Figure 3e is the amino acid sequence of the chimeric protein Flag-Fc. Included in the sequence is the signal peptide (underlined), which is cleaved by the cell after translation resulting in the mature Flag-Fc.

[044] Figure 3f is the nucleic acid sequence of the chimeric protein Epo-CCA-Fc. Included in the sequence is the signal peptide (underlined), which is cleaved by the cell after translation resulting in the mature Epo-CCA-Fc. Also shown in bold is the acidic coiled coil domain.

[045] Figure 3g is the nucleic acid sequence of the chimeric protein CCB-Fc. Included in the sequence is the signal peptide (underlined), which is cleaved by the cell after translation resulting in the mature CCB-Fc. Also shown in bold is the basic coiled coil domain.

[046] Figure 3h is the nucleic acid sequence of the chimeric protein Cys-Fc. Included in the sequence is the signal peptide (underlined), which is cleaved by the cell after translation resulting in the mature Cys-Fc.

[047] Figure 3i is the nucleic acid sequence of the chimeric protein IFN α -GS15-Fc. Included in the sequence is the signal peptide (underlined) which is cleaved by the cell after translation resulting in the mature IFN α -GS15-Fc.

[048] Figure 3j is the nucleic acid sequence of the chimeric protein Epo-Fc. Included in the sequence is the signal peptide (underlined), which is cleaved by the cell after translation resulting in the mature Epo-Fc. Also shown in bold is a nucleic acid sequence encoding the 8 amino acid linker.

[049] Figure 4 demonstrates ways to form monomer-dimer hybrids through native ligation.

[050] Figure 5a shows the amino acid sequence of Fc MESNA (SEQ ID NO:4).

[051] Figure 5b shows the DNA sequence of Fc MESNA (SEQ ID NO:5).

[052] Figure 6 compares antiviral activity of IFN α homo-dimer (*i.e.* comprised of 2 IFN α molecules) with an IFN α monomer-dimer hybrid (*i.e.* comprised of 1 IFN α molecule).

[053] Figure 7 is a comparison of clotting activity of a chimeric monomer-dimer hybrid Factor VIIa-Fc (one Factor VII molecule) and a chimeric homodimer Factor VIIa-Fc (two Factor VII molecules).

[054] Figure 8 compares oral dosing in neonatal rats of a chimeric monomer-dimer hybrid Factor VIIa-Fc (one Factor VII molecule) and a chimeric homodimer Factor VIIa-Fc (two Factor VII molecules).

[055] Figure 9 compares oral dosing in neonatal rats of a chimeric monomer-dimer hybrid Factor IX-Fc (one Factor IX molecule) with a chimeric homodimer.

[056] Figure 10 is a time course study comparing a chimeric monomer-dimer hybrid Factor IX-Fc (one Factor IX molecule) administered orally to neonatal rats with an orally administered chimeric homodimer.

[057] Figure 11 demonstrates pharmacokinetics of Epo-Fc dimer compared to Epo-Fc monomer-dimer hybrid in cynomolgus monkeys after a single pulmonary dose.

[058] Figure 12 compares serum concentration in monkeys of subcutaneously administered Epo-Fc monomer-dimer hybrid with subcutaneously administered Aranesp[®] (darbepoetin alfa).

[059] Figure 13 compares serum concentration in monkeys of intravenously administered Epo-Fc monomer-dimer hybrid with intravenously administered Aranesp[®] (darbepoetin alfa) and Epogen[®] (epoetin alfa).

[060] Figure 14 shows a trace from a Mimetic Red 2[™] column (ProMetic LifeSciences, Inc., Wayne, NJ) and an SDS-PAGE of fractions from the column containing EpoFc monomer-dimer hybrid, EpoFc dimer, and Fc. EpoFc

monomer-dimer hybrid is found in fractions 11, 12, 13, and 14. EpoFc dimer is found in fraction 18. Fc is found in fractions 1/2.

[061] Figure 15 shows the pharmacokinetics of IFN β Fc with an 8 amino acid linker in cynomolgus monkeys after a single pulmonary dose.

[062] Figure 16 shows neopterin stimulation in response to the IFN β -Fc homodimer and the IFN β -Fc N297A monomer-dimer hybrid in cynomolgus monkeys.

[063] Figure 17a shows the nucleotide sequence of interferon β -Fc; Figure 17b shows the amino acid sequence of interferon β -Fc.

[064] Figure 18 shows the amino acid sequence of T20(a); T21(b) and T1249(c).

DESCRIPTION OF THE EMBODIMENTS

A. Definitions

[065] **Affinity tag**, as used herein, means a molecule attached to a second molecule of interest, capable of interacting with a specific binding partner for the purpose of isolating or identifying said second molecule of interest.

[066] **Analogs of** chimeric proteins of the invention, or proteins or peptides substantially identical to the chimeric proteins of the invention, as used herein, means that a relevant amino acid sequence of a protein or a peptide is at least 70%, 75%, 80%, 85%, 90%, 95%, 97%, 98%, 99%, or 100% identical to a given sequence. By way of example, such sequences may be variants derived from various species, or they may be derived from the given sequence by truncation, deletion, amino acid substitution or addition. Percent identity between two amino acid sequences is determined by standard alignment algorithms such as, for example, Basic Local Alignment Tool (BLAST) described in Altschul et al. 1990, *J. Mol. Biol.*, 215:403-410, the algorithm of Needleman et al. 1970, *J. Mol. Biol.*, 48:444-453; the algorithm of Meyers et al. 1988, *Comput. Appl. Biosci.*, 4:11-17; or Tatusova et al. 1999, *FEMS Microbiol. Lett.*, 174:247-250, etc. Such algorithms are incorporated into the BLASTN, BLASTP and "BLAST 2 Sequences" programs (see www.ncbi.nlm.nih.gov/BLAST). When utilizing such programs, the default parameters can be used. For example, for nucleotide sequences the following settings can be used for "BLAST 2 Sequences": program BLASTN, reward for match 2, penalty for mismatch -2, open gap and extension

gap penalties 5 and 2 respectively, gap x_dropoff 50, expect 10, word size 11, filter ON. For amino acid sequences the following settings can be used for "BLAST 2 Sequences": program BLASTP, matrix BLOSUM62, open gap and extension gap penalties 11 and 1 respectively, gap x_dropoff 50, expect 10, word size 3, filter ON.

[067] **Bioavailability**, as used herein, means the extent and rate at which a substance is absorbed into a living system or is made available at the site of physiological activity.

[068] **Biologically active molecule**, as used herein, means a non-immunoglobulin molecule or fragment thereof, capable of treating a disease or condition or localizing or targeting a molecule to a site of a disease or condition in the body by performing a function or an action, or stimulating or responding to a function, an action or a reaction, in a biological context (e.g. in an organism, a cell, or an in vitro model thereof). Biologically active molecules may comprise at least one of polypeptides, nucleic acids, small molecules such as small organic or inorganic molecules.

[069] A **chimeric protein**, as used herein, refers to any protein comprised of a first amino acid sequence derived from a first source, bonded, covalently or non-covalently, to a second amino acid sequence derived from a second source, wherein the first and second source are not the same. A first source and a second source that are not the same can include two different biological entities, or two different proteins from the same biological entity, or a biological entity and a non-biological entity. A chimeric protein can include for example, a protein derived from at least 2 different biological sources. A biological source can include any non-synthetically produced nucleic acid or amino acid sequence (e.g. a genomic or cDNA sequence, a plasmid or viral vector, a native virion or a mutant or analog, as further described herein, of any of the above). A synthetic source can include a protein or nucleic acid sequence produced chemically and not by a biological system (e.g. solid phase synthesis of amino acid sequences). A chimeric protein can also include a protein derived from at least 2 different synthetic sources or a protein derived from at least one biological source and at least one synthetic source. A chimeric protein may also comprise a first amino acid sequence derived from a first source, covalently or non-covalently linked to a

nucleic acid, derived from any source or a small organic or inorganic molecule derived from any source. The chimeric protein may comprise a linker molecule between the first and second amino acid sequence or between the first amino acid sequence and the nucleic acid, or between the first amino acid sequence and the small organic or inorganic molecule.

[070] **Clotting factor**, as used herein, means any molecule, or analog thereof, naturally occurring or recombinantly produced which prevents or decreases the duration of a bleeding episode in a subject with a hemostatic disorder. In other words, it means any molecule having clotting activity.

[071] **Clotting activity**, as used herein, means the ability to participate in a cascade of biochemical reactions that culminates in the formation of a fibrin clot and/or reduces the severity, duration or frequency of hemorrhage or bleeding episode.

[072] **Dimer** as used herein refers to a chimeric protein comprising a first and second polypeptide chain, wherein the first and second chains both comprise a biologically active molecule, and at least a portion of an immunoglobulin constant region. A homodimer refers to a dimer where both biologically active molecules are the same.

[073] **Dimerically linked monomer-dimer hybrid** refers to a chimeric protein comprised of at least a portion of an immunoglobulin constant region, e.g. an Fc fragment of an immunoglobulin, a biologically active molecule and a linker which links the two together such that one biologically active molecule is bound to 2 polypeptide chains, each comprising a portion of an immunoglobulin constant region. Figure 4 shows an example of a dimerically linked monomer-dimer hybrid.

[074] **DNA construct**, as used herein, means a DNA molecule, or a clone of such a molecule, either single- or double-stranded that has been modified through human intervention to contain segments of DNA combined in a manner that as a whole would not otherwise exist in nature. DNA constructs contain the information necessary to direct the expression of polypeptides of interest. DNA constructs can include promoters, enhancers and transcription terminators. DNA constructs containing the information necessary to direct the secretion of a polypeptide will also contain at least one secretory signal sequence.

[075] **Domain**, as used herein, means a region of a polypeptide (including proteins as that term is defined) having some distinctive physical feature or role including for example an independently folded structure composed of one section of a polypeptide chain. A domain may contain the sequence of the distinctive physical feature of the polypeptide or it may contain a fragment of the physical feature which retains its binding characteristics (*i.e.*, it can bind to a second domain). A domain may be associated with another domain. In other words, a first domain may naturally bind to a second domain.

[076] A **fragment**, as used herein, refers to a peptide or polypeptide comprising an amino acid sequence of at least 2 contiguous amino acid residues, of at least 5 contiguous amino acid residues, of at least 10 contiguous amino acid residues, of at least 15 contiguous amino acid residues, of at least 20 contiguous amino acid residues, of at least 25 contiguous amino acid residues, of at least 40 contiguous amino acid residues, of at least 50 contiguous amino acid residues, of at least 100 contiguous amino acid residues, or of at least 200 contiguous amino acid residues or any deletion or truncation of a protein, peptide, or polypeptide.

[077] **Hemostasis**, as used herein, means the stoppage of bleeding or hemorrhage; or the stoppage of blood flow through a blood vessel or body part.

[078] **Hemostatic disorder**, as used herein, means a genetically inherited or acquired condition characterized by a tendency to hemorrhage, either spontaneously or as a result of trauma, due to an impaired ability or inability to form a fibrin clot.

[079] **Linked**, as used herein, refers to a first nucleic acid sequence covalently joined to a second nucleic acid sequence. The first nucleic acid sequence can be directly joined or juxtaposed to the second nucleic acid sequence or alternatively an intervening sequence can covalently join the first sequence to the second sequence. Linked as used herein can also refer to a first amino acid sequence covalently, or non-covalently, joined to a second amino acid sequence. The first amino acid sequence can be directly joined or juxtaposed to the second amino acid sequence or alternatively an intervening sequence can covalently join the first amino acid sequence to the second amino acid sequence.

[080] **Operatively linked**, as used herein, means a first nucleic acid sequence linked to a second nucleic acid sequence such that both sequences are capable of being expressed as a biologically active protein or peptide.

[081] **Polypeptide**, as used herein, refers to a polymer of amino acids and does not refer to a specific length of the product; thus, peptides, oligopeptides, and proteins are included within the definition of polypeptide. This term does not exclude post-expression modifications of the polypeptide, for example, glycosylation, acetylation, phosphorylation, pegylation, addition of a lipid moiety, or the addition of any organic or inorganic molecule. Included within the definition, are for example, polypeptides containing one or more analogs of an amino acid (including, for example, unnatural amino acids) and polypeptides with substituted linkages, as well as other modifications known in the art, both naturally occurring and non-naturally occurring.

[082] **High stringency**, as used herein, includes conditions readily determined by the skilled artisan based on, for example, the length of the DNA. Generally, such conditions are defined in Sambrook et al. *Molecular Cloning: A Laboratory Manual*, 2 ed. Vol. 1, pp. 1.101-104, Cold Spring Harbor Laboratory Press (1989), and include use of a prewashing solution for the nitrocellulose filters 5X SSC, 0.5% SDS, 1.0 mM EDTA (PH 8.0), hybridization conditions of 50% formamide, 6X SSC at 42°C (or other similar hybridization solution, such as Stark's solution, in 50% formamide at 42°C, and with washing at approximately 68°C, 0.2X SSC, 0.1% SDS. The skilled artisan will recognize that the temperature and wash solution salt concentration can be adjusted as necessary according to factors such as the length of the probe.

[083] **Moderate stringency**, as used herein, include conditions that can be readily determined by those having ordinary skill in the art based on, for example, the length of the DNA. The basic conditions are set forth by Sambrook et al. *Molecular Cloning: A Laboratory Manual*, 2d ed. Vol. 1, pp. 1.101-104, Cold Spring Harbor Laboratory Press (1989), and include use of a prewashing solution for the nitrocellulose filters 5X SSC, 0.5% SDS, 1.0 mM EDTA (pH 8.0), hybridization conditions of 50% formamide, 6X SSC at 42°C (or other similar hybridization solution, such as Stark's solution, in 50% formamide at 42°C), and washing conditions of 60°C, 0.5X SSC, 0.1% SDS.

[084] A **small inorganic molecule**, as used herein means a molecule containing no carbon atoms and being no larger than 50 kD.

[085] A **small organic molecule**, as used herein means a molecule containing at least one carbon atom and being no larger than 50 kD.

[086] **Treat, treatment, treating**, as used herein means, any of the following: the reduction in severity of a disease or condition; the reduction in the duration of a disease course; the amelioration of one or more symptoms associated with a disease or condition; the provision of beneficial effects to a subject with a disease or condition, without necessarily curing the disease or condition, the prophylaxis of one or more symptoms associated with a disease or condition.

B. Improvements Offered by Certain Embodiments of the Invention

[087] The invention provides for chimeric proteins (monomer-dimer hybrids) comprising a first and a second polypeptide chain, wherein said first chain comprises a biologically active molecule and at least a portion of an immunoglobulin constant region, and said second chain comprises at least a portion of an immunoglobulin constant region without any biologically active molecule or variable region of an immunoglobulin. Figure 1 contrasts traditional fusion protein dimers with one example of the monomer-dimer hybrid of the invention. In this example, the biologically active molecule is EPO and the portion of an immunoglobulin is IgG Fc region.

[088] Like other chimeric proteins comprised of at least a portion of an immunoglobulin constant region, the invention provides for chimeric proteins which afford enhanced stability and increased bioavailability of the chimeric protein compared to the biologically active molecule alone. Additionally, however, because only one of the two chains comprises the biologically active molecule, the chimeric protein has a lower molecular weight than a chimeric protein wherein all chains comprise a biologically active molecule and while not wishing to be bound by any theory, this may result in the chimeric protein being more readily transcytosed across the epithelium barrier, e.g., by binding to the FcRn receptor thereby increasing the half-life of the chimeric protein. In one embodiment, the invention thus provides for an improved non-invasive method (e.g. via any mucosal surface, such as, orally, buccally, sublingually, nasally, rectally, vaginally,

or via pulmonary or ocular route) of administering a therapeutic chimeric protein of the invention. The invention thus provides methods of attaining therapeutic levels of the chimeric proteins of the invention using less frequent and lower doses compared to previously described chimeric proteins (e.g. chimeric proteins comprised of at least a portion of an immunoglobulin constant region and a biologically active molecule, wherein all chains of the chimeric protein comprise a biologically active molecule).

[089] In another embodiment, the invention provides an invasive method, e.g., subcutaneously, intravenously, of administering a therapeutic chimeric protein of the invention. Invasive administration of the therapeutic chimeric protein of the invention provides for an increased half life of the therapeutic chimeric protein which results in using less frequent and lower doses compared to previously described chimeric proteins (e.g. chimeric proteins comprised of at least a portion of an immunoglobulin constant region and a biologically active molecule, wherein all chains of the chimeric protein comprise a biologically active molecule).

[090] Yet another advantage of a chimeric protein wherein only one of the chains comprises a biologically active molecule is the enhanced accessibility of the biologically active molecule for its target cell or molecule resulting from decreased steric hindrance, decreased hydrophobic interactions, decreased ionic interactions, or decreased molecular weight compared to a chimeric protein wherein all chains are comprised of a biologically active molecule.

C. Chimeric Proteins

[091] The invention relates to chimeric proteins comprising one biologically active molecule, at least a portion of an immunoglobulin constant region, and optionally at least one linker. The portion of an immunoglobulin will have both an N, or an amino terminus, and a C, or carboxy terminus. The chimeric protein may have the biologically active molecule linked to the N terminus of the portion of an immunoglobulin. Alternatively, the biologically active molecule may be linked to the C terminus of the portion of an immunoglobulin. In one embodiment, the linkage is a covalent bond. In another embodiment, the linkage is a non-covalent bond.

[092] The chimeric protein can optionally comprise at least one linker; thus, the biologically active molecule does not have to be directly linked to the portion of an immunoglobulin constant region. The linker can intervene in between the biologically active molecule and the portion of an immunoglobulin constant region. The linker can be linked to the N terminus of the portion of an immunoglobulin constant region, or the C terminus of the portion of an immunoglobulin constant region. If the biologically active molecule is comprised of at least one amino acid the biologically active molecule will have an N terminus and a C terminus and the linker can be linked to the N terminus of the biologically active molecule, or the C terminus the biologically active molecule.

[093] The invention relates to a chimeric protein of the formula $X-L_a-F:F$ or $F:F-L_a-X$, wherein X is a biologically active molecule, L is an optional linker, F is at least a portion of an immunoglobulin constant region and, a is any integer or zero. The invention also relates to a chimeric protein of the formula $T_a-X-L_a-F:F$ or $T_a-F:F-L_a-X$, wherein X is a biologically active molecule, L is an optional linker, F is at least a portion of an immunoglobulin constant region, a is any integer or zero, T is a second linker or alternatively a tag that can be used to facilitate purification of the chimeric protein, e.g., a FLAG tag, a histidine tag, a GST tag, a maltose binding protein tag and (:) represents a chemical association, e.g. at least one non-peptide bond. In certain embodiments, the chemical association, i.e., (:) is a covalent bond. In other embodiments, the chemical association, i.e., (:) is a non-covalent interaction, e.g., an ionic interaction, a hydrophobic interaction, a hydrophilic interaction, a Van der Waals interaction, a hydrogen bond. It will be understood by the skilled artisan that when a equals zero X will be directly linked to F. Thus, for example, a may be 0, 1, 2, 3, 4, 5, or more than 5.

[094] In one embodiment, the chimeric protein of the invention comprises the amino acid sequence of figure 2a (SEQ ID NO:6). In one embodiment, the chimeric protein of the invention comprises the amino acid sequence of figure 2b (SEQ ID NO:8). In one embodiment, the chimeric protein of the invention comprises the amino acid sequence of figure 2c (SEQ ID NO:10). In one embodiment, the chimeric protein of the invention comprises the amino acid sequence of figure 2d (SEQ ID NO:12). In one embodiment, the chimeric protein of the invention comprises the amino acid sequence of figure 2e (SEQ ID NO:14).

In one embodiment, the chimeric protein of the invention comprises the amino acid sequence of figure 2f (SEQ ID NO:16). In one embodiment, the chimeric protein of the invention comprises the amino acid sequence of figure 2g (SEQ ID NO:18). In one embodiment, the chimeric protein of the invention comprises the amino acid sequence of figure 2h (SEQ ID NO:20). In one embodiment, the chimeric protein of the invention comprises the amino acid sequence of figure 2i (SEQ ID NO:22). In one embodiment, the chimeric protein of the invention comprises the amino acid sequence of figure 2j (SEQ ID NO:24). In one embodiment, the chimeric protein of the invention comprises the amino acid sequence of figure 17b (SEQ ID NO:27).

1. Chimeric Protein Variants

[095] Derivatives of the chimeric proteins of the invention, antibodies against the chimeric proteins of the invention and antibodies against binding partners of the chimeric proteins of the invention are all contemplated, and can be made by altering their amino acids sequences by substitutions, additions, and/or deletions/truncations or by introducing chemical modification that result in functionally equivalent molecules. It will be understood by one of ordinary skill in the art that certain amino acids in a sequence of any protein may be substituted for other amino acids without adversely affecting the activity of the protein.

[096] Various changes may be made in the amino acid sequences of the chimeric proteins of the invention or DNA sequences encoding therefore without appreciable loss of their biological activity, function, or utility. Derivatives, analogs, or mutants resulting from such changes and the use of such derivatives is within the scope of the present invention. In a specific embodiment, the derivative is functionally active, *i.e.*, capable of exhibiting one or more activities associated with the chimeric proteins of the invention, *e.g.*, FcRn binding, viral inhibition, hemostasis, production of red blood cells. Many assays capable of testing the activity of a chimeric protein comprising a biologically active molecule are known in the art. Where the biologically active molecule is an HIV inhibitor, activity can be tested by measuring reverse transcriptase activity using known methods (see, *e.g.*, Barre-Sinoussi et al. 1983, *Science* 220:868; Gallo et al. 1984, *Science* 224:500). Alternatively, activity can be measured by measuring fusogenic activity (see, *e.g.*, Nussbaum et al. 1994, *J. Virol.* 68(9):5411). Where

the biological activity is hemostasis, a StaCLot FVIIa-rTF assay can be performed to assess activity of Factor VIIa derivatives (Johannessen et al. 2000, *Blood Coagulation and Fibrinolysis* 11:S159).

[097] Substitutes for an amino acid within the sequence may be selected from other members of the class to which the amino acid belongs (see Table 1). Furthermore, various amino acids are commonly substituted with neutral amino acids, e.g., alanine, leucine, isoleucine, valine, proline, phenylalanine, tryptophan, and methionine (see, e.g., MacLennan et al. 1998, *Acta Physiol. Scand. Suppl.* 643:55-67; Sasaki et al. 1998, *Adv. Biophys.* 35:1-24).

TABLE 1

Original Residues	Exemplary Substitutions	Typical Substitutions
Ala (A)	Val, Leu, Ile	Val
Arg (R)	Lys, Gln, Asn	Lys
Asn (N)	Gln	Gln
Asp (D)	Glu	Glu
Cys (C)	Ser, Ala	Ser
Gln (Q)	Asn	Asn
Gly (G)	Pro, Ala	Ala
His (H)	Asn, Gln, Lys, Arg	Arg
Ile (I)	Leu, Val, Met, Ala, Phe, Norleucine	Leu
Leu (L)	Norleucine, Ile, Val, Met, Ala, Phe	Ile
Lys (K)	Arg, 1,4-Diamino-butyrlic Acid, Gln, Asn	Arg
Met (M)	Leu, Phe, Ile	Leu
Phe (F)	Leu, Val, Ile, Ala, Tyr	Leu
Pro (P)	Ala	Gly
Ser (S)	Thr, Ala, Cys	Thr
Thr (T)	Ser	Ser
Trp (W)	Tyr, Phe	Tyr
Tyr (Y)	Trp, Phe, Thr, Ser	Phe
Val (V)	Ile, Met, Leu, Phe, Ala, Norleucine	Leu

2. Biologically Active Molecules

[0098] The invention contemplates the use of any biologically active molecule as the therapeutic molecule of the invention. The biologically active molecule can be a polypeptide. The biologically active molecule can be a single amino acid. The biologically active molecule can include a modified polypeptide.

[0099] The biologically active molecule can include a lipid molecule (e.g. a steroid or cholesterol, a fatty acid, a triacylglycerol, glycerophospholipid, or sphingolipid). The biologically active molecule can include a sugar molecule (e.g. glucose, sucrose, mannose). The biologically active molecule can include a nucleic acid molecule (e.g. DNA, RNA). The biologically active molecule can include a small organic molecule or a small inorganic molecule.

a. Cytokines and Growth Factors

[0100] In one embodiment, the biologically active molecule is a growth factor, hormone or cytokine or analog or fragment thereof. The biologically active molecule can be any agent capable of inducing cell growth and proliferation. In a specific embodiment, the biologically active molecule is any agent which can induce erythrocytes to proliferate. Thus, one example of a biologically active molecule contemplated by the invention is EPO. The biologically active molecule can also include, but is not limited to, RANTES, MIP1 α , MIP1 β , IL-2, IL-3, GM-CSF, growth hormone, tumor necrosis factor (e.g. TNF α or β).

[0101] The biologically active molecule can include interferon α , whether synthetically or recombinantly produced, including but not limited to, any one of the about twenty-five structurally related subtypes, as for example interferon- α 2a, now commercially available for clinical use (ROFERON®, Roche) and interferon- α 2b also approved for clinical use (INTRON®, Schering) as well as genetically engineered versions of various subtypes, including, but not limited to, commercially available consensus interferon α (INFERGEN®, Intermune, developed by Amgen) and consensus human leukocyte interferon see, e.g., U.S. Patent Nos.: 4,695,623; 4,897,471, interferon β , epidermal growth factor, gonadotropin releasing hormone (GnRH), leuprolide, follicle stimulating hormone, progesterone, estrogen, or testosterone.

[0102] A list of cytokines and growth factors which may be used in the chimeric protein of the invention has been previously described (see, e.g., U.S.

Patent Nos. 6,086,875, 6,485,726, 6,030,613; WO 03/077834; US2003-0235536A1).

b. Antiviral Agents

[0103] In one embodiment, the biologically active molecule is an antiviral agent, including fragments and analogs thereof. An antiviral agent can include any molecule that inhibits or prevents viral replication, or inhibits or prevents viral entry into a cell, or inhibits or prevents viral egress from a cell. In one embodiment, the antiviral agent is a fusion inhibitor. In one embodiment, the antiviral agent is a cytokine which inhibits viral replication. In another embodiment, the antiviral agent is interferon α .

[0104] The viral fusion inhibitor for use in the chimeric protein can be any molecule which decreases or prevents viral penetration of a cellular membrane of a target cell. The viral fusion inhibitor can be any molecule that decreases or prevents the formation of syncytia between at least two susceptible cells. The viral fusion inhibitor can be any molecule that decreases or prevents the joining of a lipid bilayer membrane of a eukaryotic cell and a lipid bilayer of an enveloped virus. Examples of enveloped virus include, but are not limited to HIV-1, HIV-2, SIV, influenza, parainfluenza, Epstein-Barr virus, CMV, herpes simplex 1, herpes simplex 2 and respiratory syncytia virus.

[0105] The viral fusion inhibitor can be any molecule that decreases or prevents viral fusion including, but not limited to, a polypeptide, a small organic molecule or a small inorganic molecule. In one embodiment, the fusion inhibitor is a polypeptide. In one embodiment, the viral fusion inhibitor is a polypeptide of 3-36 amino acids. In another embodiment, the viral fusion inhibitor is a polypeptide of 3-50 amino acids, 10-65 amino acids, 10-75 amino acids. The polypeptide can be comprised of a naturally occurring amino acid sequence (e.g. a fragment of gp41) including analogs and mutants thereof or the polypeptide can be comprised of an amino acid sequence not found in nature, so long as the polypeptide exhibits viral fusion inhibitory activity.

[0106] In one embodiment, the viral fusion inhibitor is a polypeptide, identified as being a viral fusion inhibitor using at least one computer algorithm, e.g., ALLMOTI5, 107x178x4 and PLZIP (see, e.g., U.S. Patent Nos.: 6,013,263;

6,015,881; 6,017,536; 6,020,459; 6,060,065; 6,068,973; 6,093,799; and 6,228,983).

[0107] In one embodiment, the viral fusion inhibitor is an HIV fusion inhibitor. In one embodiment, HIV is HIV-1. In another embodiment, HIV is HIV-2. In one embodiment, the HIV fusion inhibitor is a polypeptide comprised of a fragment of the gp41 envelope protein of HIV-1. The HIV fusion inhibitor can comprise, e.g., T20 (SEQ ID NO:1) or an analog thereof, T21 (SEQ ID NO:2) or an analog thereof, T1249 (SEQ ID NO:3) or an analog thereof, N_{CCG}gp41 (Louis et al. 2001, *J. Biol. Chem.* 276:(31)29485) or an analog thereof, or 5 helix (Root et al. 2001, *Science* 291:884) or an analog thereof.

[0108] Assays known in the art can be used to test for viral fusion inhibiting activity of a polypeptide, a small organic molecule, or a small inorganic molecule. These assays include a reverse transcriptase assay, a p24 assay, or syncytia formation assay (see, e.g., U.S. Patent No. 5,464,933).

[0109] A list of antiviral agents which may be used in the chimeric protein of the invention has been previously described (see, e.g., U.S. Patent Nos. 6,086,875, 6,485,726, 6,030,613; WO 03/077834; US2003-0235536A1).

c. Hemostatic Agents

[0110] In one embodiment, the biologically active molecule is a clotting factor or other agent that promotes hemostasis, including fragments and analogs thereof. The clotting factor can include any molecule that has clotting activity or activates a molecule with clotting activity. The clotting factor can be comprised of a polypeptide. The clotting factor can be, as an example, but not limited to Factor VIII, Factor IX, Factor XI, Factor XII, fibrinogen, prothrombin, Factor V, Factor VII, Factor X, Factor XIII or von Willebrand Factor. In one embodiment, the clotting factor is Factor VII or Factor VIIa. The clotting factor can be a factor that participates in the extrinsic pathway. The clotting factor can be a factor that participates in the intrinsic pathway. Alternatively, the clotting factor can be a factor that participates in both the extrinsic and intrinsic pathway.

[0111] The clotting factor can be a human clotting factor or a non-human clotting factor, e.g., derived from a non-human primate, a pig or any mammal. The clotting factor can be chimeric clotting factor, e.g., the clotting factor can comprise a portion of a human clotting factor and a portion of a porcine clotting

factor or a portion of a first non-human clotting factor and a portion of a second non-human clotting factor.

[0112] The clotting factor can be an activated clotting factor. Alternatively, the clotting factor can be an inactive form of a clotting factor, *e.g.*, a zymogen. The inactive clotting factor can undergo activation subsequent to being linked to at least a portion of an immunoglobulin constant region. The inactive clotting factor can be activated subsequent to administration to a subject. Alternatively, the inactive clotting factor can be activated prior to administration.

[0113] In certain embodiments an endopeptidase, *e.g.*, paired basic amino acid cleaving enzyme (PACE), or any PACE family member, such as PCSK1-9, including truncated versions thereof, or its yeast equivalent Kex2 from *S. cerevisiae* and truncated versions of Kex2 (Kex2 1-675) (*see, e.g.*, U.S. Patent Nos. 5,077,204; 5,162,220; 5,234,830; 5,885,821; 6,329,176) may be used to cleave a propeptide to form the mature chimeric protein of the invention (*e.g.* factor VII, factor IX).

d. Other Proteinaceous Biologically Active Molecules

[0114] In one embodiment, the biologically active molecule is a receptor or a fragment or analog thereof. The receptor can be expressed on a cell surface, or alternatively the receptor can be expressed on the interior of the cell. The receptor can be a viral receptor, *e.g.*, CD4, CCR5, CXCR4, CD21, CD46. The biologically active molecule can be a bacterial receptor. The biologically active molecule can be an extra-cellular matrix protein or fragment or analog thereof, important in bacterial colonization and infection (*see, e.g.*, U.S. Patent Nos.: 5,648,240; 5,189,015; 5,175,096) or a bacterial surface protein important in adhesion and infection (*see, e.g.*, U.S. Patent No. 5,648,240). The biologically active molecule can be a growth factor, hormone or cytokine receptor, or a fragment or analog thereof, *e.g.*, TNF α receptor, the erythropoietin receptor, CD25, CD122, or CD132.

[0115] A list of other proteinaceous molecules which may be used in the chimeric protein of the invention has been previously described (*see, e.g.*, U.S. Patent Nos. 6,086,875; 6,485,726; 6,030,613; WO 03/077834; US2003-0235536A1).

e. Nucleic Acids

[0116] In one embodiment, the biologically active molecule is a nucleic acid, e.g., DNA, RNA. In one specific embodiment, the biologically active molecule is a nucleic acid that can be used in RNA interference (RNAi). The nucleic acid molecule can be as an example, but not as a limitation, an anti-sense molecule or a ribozyme or an aptamer.

[0117] Antisense RNA and DNA molecules act to directly block the translation of mRNA by hybridizing to targeted mRNA and preventing protein translation. Antisense approaches involve the design of oligonucleotides that are complementary to a target gene mRNA. The antisense oligonucleotides will bind to the complementary target gene mRNA transcripts and prevent translation. Absolute complementarity, is not required.

[0118] A sequence "complementary" to a portion of an RNA, as referred to herein, means a sequence having sufficient complementarity to be able to hybridize with the RNA, forming a stable duplex; in the case of double-stranded antisense nucleic acids, a single strand of the duplex DNA may thus be tested, or triplex formation may be assayed. The ability to hybridize will depend on both the degree of complementarity and the length of the antisense nucleic acid. Generally, the longer the hybridizing nucleic acid, the more base mismatches with an RNA it may contain and still form a stable duplex (or triplex, as the case may be). One skilled in the art can ascertain a tolerable degree of mismatch by use of standard procedures to determine the melting point of the hybridized complex.

[0119] Antisense nucleic acids should be at least six nucleotides in length, and are preferably oligonucleotides ranging from 6 to about 50 nucleotides in length. In specific aspects, the oligonucleotide is at least 10 nucleotides, at least 17 nucleotides, at least 25 nucleotides or at least 50 nucleotides.

[0120] The oligonucleotides can be DNA or RNA or chimeric mixtures or derivatives or modified versions thereof, single-stranded or double-stranded. The oligonucleotide can be modified at the base moiety, sugar moiety, or phosphate backbone, for example, to improve stability of the molecule, hybridization, etc. The oligonucleotide may include other appended groups such as polypeptides (e.g. for targeting host cell receptors in vivo), or agents facilitating transport across the cell membrane (see, e.g., Letsinger et al. 1989, *Proc. Natl. Acad. Sci. USA*

86:6553; Lemaitre et al. 1987, *Proc. Natl. Acad. Sci. USA* 84:648; WO 88/09810,) or the blood-brain barrier (see, e.g., WO 89/10134), hybridization-triggered cleavage agents (see, e.g., Krol et al. 1988, *BioTechniques* 6:958) or intercalating agents (see, e.g., Zon 1988, *Pharm. Res.* 5:539). To this end, the oligonucleotide may be conjugated to another molecule, e.g., a polypeptide, hybridization triggered cross-linking agent, transport agent, or hybridization-triggered cleavage agent.

[0121] Ribozyme molecules designed to catalytically cleave target gene mRNA transcripts can also be used to prevent translation of target gene mRNA and, therefore, expression of target gene product. (See, e.g., WO 90/11364; Sarver et al. 1990, *Science* 247, 1222-1225).

[0122] Ribozymes are enzymatic RNA molecules capable of catalyzing the specific cleavage of RNA. (See Rossi 1994, *Current Biology* 4:469). The mechanism of ribozyme action involves sequence specific hybridization of the ribozyme molecule to complementary target RNA, followed by an endonucleolytic cleavage event. The composition of ribozyme molecules must include one or more sequences complementary to the target gene mRNA, and must include the well known catalytic sequence responsible for mRNA cleavage. For this sequence, see, e.g., U.S. Pat. No. 5,093,246.

[0123] In one embodiment, ribozymes that cleave mRNA at site specific recognition sequences can be used to destroy target gene mRNAs. In another embodiment, the use of hammerhead ribozymes is contemplated. Hammerhead ribozymes cleave mRNAs at locations dictated by flanking regions that form complementary base pairs with the target mRNA. The sole requirement is that the target mRNA have the following sequence of two bases: 5'-UG-3'. The construction and production of hammerhead ribozymes is well known in the art and is described more fully in Myers 1995, *Molecular Biology and Biotechnology: A Comprehensive Desk Reference*, VCH Publishers, New York, and in Haseloff and Gerlach 1988, *Nature*, 334:585.

f. Small Molecules

[0124] The invention also contemplates the use of any therapeutic small molecule or drug as the biologically active molecule in the chimeric protein of the invention. A list of small molecules and drugs which may be used in the chimeric

protein of the invention has been previously described (see, e.g., U.S. Patent Nos. 6,086,875; 6,485,726; 6,030,613; WO 03/077834; US2003-0235536A1).

3. Immunoglobulins

[0125] The chimeric proteins of the invention comprise at least a portion of an immunoglobulin constant region. Immunoglobulins are comprised of four protein chains that associate covalently—two heavy chains and two light chains. Each chain is further comprised of one variable region and one constant region. Depending upon the immunoglobulin isotype, the heavy chain constant region is comprised of 3 or 4 constant region domains (e.g. CH1, CH2, CH3, CH4). Some isotypes are further comprised of a hinge region.

[0126] The portion of an immunoglobulin constant region can be obtained from any mammal. The portion of an immunoglobulin constant region can include a portion of a human immunoglobulin constant region, a non-human primate immunoglobulin constant region, a bovine immunoglobulin constant region, a porcine immunoglobulin constant region, a murine immunoglobulin constant region, an ovine immunoglobulin constant region or a rat immunoglobulin constant region.

[0127] The portion of an immunoglobulin constant region can be produced recombinantly or synthetically. The immunoglobulin can be isolated from a cDNA library. The portion of an immunoglobulin constant region can be isolated from a phage library (See, e.g., McCafferty *et al.* 1990, *Nature* 348:552, Kang *et al.* 1991, *Proc. Natl. Acad. Sci. USA* 88:4363; EP 0 589 877 B1). The portion of an immunoglobulin constant region can be obtained by gene shuffling of known sequences (Mark *et al.* 1992, *Bio/Technol.* 10:779). The portion of an immunoglobulin constant region can be isolated by in vivo recombination (Waterhouse *et al.* 1993, *Nucl. Acid Res.* 21:2265). The immunoglobulin can be a humanized immunoglobulin (U.S. Patent No. 5,585,089, Jones *et al.* 1986, *Nature* 332:323).

[0128] The portion of an immunoglobulin constant region can include a portion of an IgG, an IgA, an IgM, an IgD, or an IgE. In one embodiment, the immunoglobulin is an IgG. In another embodiment, the immunoglobulin is IgG1. In another embodiment, the immunoglobulin is IgG2.

[0129] The portion of an immunoglobulin constant region can include the entire heavy chain constant region, or a fragment or analog thereof. In one embodiment, a heavy chain constant region can comprise a CH1 domain, a CH2 domain, a CH3 domain, and/or a hinge region. In another embodiment, a heavy chain constant region can comprise a CH1 domain, a CH2 domain, a CH3 domain, and/or a CH4 domain.

[0130] The portion of an immunoglobulin constant region can include an Fc fragment. An Fc fragment can be comprised of the CH2 and CH3 domains of an immunoglobulin and the hinge region of the immunoglobulin. The Fc fragment can be the Fc fragment of an IgG1, an IgG2, an IgG3 or an IgG4. In one specific embodiment, the portion of an immunoglobulin constant region is an Fc fragment of an IgG1. In another embodiment, the portion of an immunoglobulin constant region is an Fc fragment of an IgG2.

[0131] In another embodiment, the portion of an immunoglobulin constant region is an Fc neonatal receptor (FcRn) binding partner. An FcRn binding partner is any molecule that can be specifically bound by the FcRn receptor with consequent active transport by the FcRn receptor of the FcRn binding partner. Specifically bound refers to two molecules forming a complex that is relatively stable under physiologic conditions. Specific binding is characterized by a high affinity and a low to moderate capacity as distinguished from nonspecific binding which usually has a low affinity with a moderate to high capacity. Typically, binding is considered specific when the affinity constant K_A is higher than 10^6 M^{-1} , or more preferably higher than 10^8 M^{-1} . If necessary, non-specific binding can be reduced without substantially affecting specific binding by varying the binding conditions. The appropriate binding conditions such as concentration of the molecules, ionic strength of the solution, temperature, time allowed for binding, concentration of a blocking agent (e.g. serum albumin, milk casein), etc., may be optimized by a skilled artisan using routine techniques.

[0132] The FcRn receptor has been isolated from several mammalian species including humans. The sequences of the human FcRn, monkey FcRn rat FcRn, and mouse FcRn are known (Story et al. 1994, *J. Exp. Med.* 180:2377). The FcRn receptor binds IgG (but not other immunoglobulin classes such as IgA, IgM, IgD, and IgE) at relatively low pH, actively transports the IgG transcellularly in

a luminal to serosal direction, and then releases the IgG at relatively higher pH found in the interstitial fluids. It is expressed in adult epithelial tissue (U.S. Patent Nos. 6,485,726, 6,030,613, 6,086,875; WO 03/077834; US2003-0235536A1) including lung and intestinal epithelium (Israel et al. 1997, *Immunology* 92:69) renal proximal tubular epithelium (Kobayashi et al. 2002, *Am. J. Physiol. Renal Physiol.* 282:F358) as well as nasal epithelium, vaginal surfaces, and biliary tree surfaces.

[0133] FcRn binding partners of the present invention encompass any molecule that can be specifically bound by the FcRn receptor including whole IgG, the Fc fragment of IgG, and other fragments that include the complete binding region of the FcRn receptor. The region of the Fc portion of IgG that binds to the FcRn receptor has been described based on X-ray crystallography (Burmeister et al. 1994, *Nature* 372:379). The major contact area of the Fc with the FcRn is near the junction of the CH2 and CH3 domains. Fc-FcRn contacts are all within a single Ig heavy chain. The FcRn binding partners include whole IgG, the Fc fragment of IgG, and other fragments of IgG that include the complete binding region of FcRn. The major contact sites include amino acid residues 248, 250-257, 272, 285, 288, 290-291, 308-311, and 314 of the CH2 domain and amino acid residues 385-387, 428, and 433-436 of the CH3 domain. References made to amino acid numbering of immunoglobulins or immunoglobulin fragments, or regions, are all based on Kabat et al. 1991, *Sequences of Proteins of Immunological Interest*, U.S. Department of Public Health, Bethesda, MD.

[0134] The Fc region of IgG can be modified according to well recognized procedures such as site directed mutagenesis and the like to yield modified IgG or Fc fragments or portions thereof that will be bound by FcRn. Such modifications include modifications remote from the FcRn contact sites as well as modifications within the contact sites that preserve or even enhance binding to the FcRn. For example, the following single amino acid residues in human IgG1 Fc (Fcγ1) can be substituted without significant loss of Fc binding affinity for FcRn: P238A, S239A, K246A, K248A, D249A, M252A, T256A, E258A, T260A, D265A, S267A, H268A, E269A, D270A, E272A, L274A, N276A, Y278A, D280A, V282A, E283A, H285A, N286A, T289A, K290A, R292A, E293A, E294A, Q295A, Y296F, N297A, S298A, Y300F, R301A, V303A, V305A, T307A, L309A, Q311A, D312A, N315A,

K317A, E318A, K320A, K322A, S324A, K326A, A327Q, P329A, A330Q, P331A, E333A, K334A, T335A, S337A, K338A, K340A, Q342A, R344A, E345A, Q347A, R355A, E356A, M358A, T359A, K360A, N361A, Q362A, Y373A, S375A, D376A, A378Q, E380A, E382A, S383A, N384A, Q386A, E388A, N389A, N390A, Y391F, K392A, L398A, S400A, D401A, D413A, K414A, R416A, Q418A, Q419A, N421A, V422A, S424A, E430A, N434A, T437A, Q438A, K439A, S440A, S444A, and K447A, where for example P238A represents wildtype proline substituted by alanine at position number 238. As an example, one specific embodiment, incorporates the N297A mutation, removing a highly conserved N-glycosylation site. In addition to alanine other amino acids may be substituted for the wildtype amino acids at the positions specified above. Mutations may be introduced singly into Fc giving rise to more than one hundred FcRn binding partners distinct from native Fc. Additionally, combinations of two, three, or more of these individual mutations may be introduced together, giving rise to hundreds more FcRn binding partners. Moreover, one of the FcRn binding partners of the monomer-dimer hybrid may be mutated and the other FcRn binding partner not mutated at all, or they both may be mutated but with different mutations. Any of the mutations described herein, including N297A, may be used to modify Fc, regardless of the biologically active molecule (e.g., EPO, IFN, Factor IX, T20).

[0135] Certain of the above mutations may confer new functionality upon the FcRn binding partner. For example, one embodiment incorporates N297A, removing a highly conserved N-glycosylation site. The effect of this mutation is to reduce immunogenicity, thereby enhancing circulating half life of the FcRn binding partner, and to render the FcRn binding partner incapable of binding to FcγRI, FcγRIIA, FcγRIIB, and FcγRIIIA, without compromising affinity for FcRn (Routledge et al. 1995, *Transplantation* 60:847; Friend et al. 1999, *Transplantation* 68:1632; Shields et al. 1995, *J. Biol. Chem.* 276:6591). As a further example of new functionality arising from mutations described above affinity for FcRn may be increased beyond that of wild type in some instances. This increased affinity may reflect an increased "on" rate, a decreased "off" rate or both an increased "on" rate and a decreased "off" rate. Mutations believed to impart an increased affinity for FcRn include T256A, T307A, E380A, and N434A (Shields et al. 2001, *J. Biol. Chem.* 276:6591).

[0136] Additionally, at least three human Fc gamma receptors appear to recognize a binding site on IgG within the lower hinge region, generally amino acids 234-237. Therefore, another example of new functionality and potential decreased immunogenicity may arise from mutations of this region, as for example by replacing amino acids 233-236 of human IgG1 "ELLG" to the corresponding sequence from IgG2 "PVA" (with one amino acid deletion). It has been shown that FcγRI, FcγRII, and FcγRIII, which mediate various effector functions will not bind to IgG1 when such mutations have been introduced. Ward and Ghetie 1995, *Therapeutic Immunology* 2:77 and Armour et al.1999, *Eur. J. Immunol.* 29:2613.

[0137] In one embodiment, the FcRn binding partner is a polypeptide including the sequence PKNSSMISNTP (SEQ ID NO:26) and optionally further including a sequence selected from HQSLGTQ (SEQ ID NO:27), HQNLSDGK (SEQ ID NO:28), HQNISDGK (SEQ ID NO:29), or VISSHLGQ (SEQ ID NO:30) (U.S. Patent No. 5,739,277).

[0138] Two FcRn receptors can bind a single Fc molecule. Crystallographic data suggest that each FcRn molecule binds a single polypeptide of the Fc homodimer. In one embodiment, linking the FcRn binding partner, e.g., an Fc fragment of an IgG, to a biologically active molecule provides a means of delivering the biologically active molecule orally, buccally, sublingually, rectally, vaginally, as an aerosol administered nasally or via a pulmonary route, or via an ocular route. In another embodiment, the chimeric protein can be administered invasively, e.g., subcutaneously, intravenously.

[0139] The skilled artisan will understand that portions of an immunoglobulin constant region for use in the chimeric protein of the invention can include mutants or analogs thereof, or can include chemically modified immunoglobulin constant regions (e.g. pegylated), or fragments thereof (see, e.g., Aslam and Dent 1998, *Bioconjugation: Protein Coupling Techniques For the Biomedical Sciences Macmilan Reference*, London). In one instance, a mutant can provide for enhanced binding of an FcRn binding partner for the FcRn. Also contemplated for use in the chimeric protein of the invention are peptide mimetics of at least a portion of an immunoglobulin constant region, e.g., a peptide mimetic of an Fc fragment or a peptide mimetic of an FcRn binding partner. In one

embodiment, the peptide mimetic is identified using phage display or via chemical library screening (see, e.g., McCafferty *et al.* 1990, *Nature* 348:552, Kang *et al.* 1991, *Proc. Natl. Acad. Sci. USA* 88:4363; EP 0 589 877 B1).

4. Optional Linkers

[0140] The chimeric protein of the invention can optionally comprise at least one linker molecule. The linker can be comprised of any organic molecule. In one embodiment, the linker is polyethylene glycol (PEG). In another embodiment, the linker is comprised of amino acids. The linker can comprise 1-5 amino acids, 1-10 amino acids, 1-20 amino acids, 10-50 amino acids, 50-100 amino acids, 100-200 amino acids. In one embodiment, the linker is the eight amino acid linker EFAGAAAV (SEQ ID NO:31). Any of the linkers described herein may be used in the chimeric protein of the invention, e.g., a monomer-dimer hybrid, including EFAGAAAV, regardless of the biologically active molecule (e.g. EPO, IFN, Factor IX).

[0141] The linker can comprise the sequence G_n . The linker can comprise the sequence $(GA)_n$ (SEQ ID NO:32). The linker can comprise the sequence $(GGG)_n$ (SEQ ID NO:33). The linker can comprise the sequence $(GGG)_n(GGGG)_n$ (SEQ ID NO:34). In these instances, n may be an integer from 1-10, i.e., 1, 2, 3, 4, 5, 6, 7, 8, 9, 10. Examples of linkers include, but are not limited to, GGG (SEQ ID NO:35), SGGSGGS (SEQ ID NO:36), GGS GGSGGS GGSGGG (SEQ ID NO:37), GGS GGSGGGGGSGGGGS (SEQ ID NO:38), GGS GGSGGS GGSGGS GGSGGS (SEQ ID NO:39). The linker does not eliminate or diminish the biological activity of the chimeric protein. Optionally, the linker enhances the biological activity of the chimeric protein, e.g., by further diminishing the effects of steric hindrance and making the biologically active molecule more accessible to its target binding site.

[0142] In one specific embodiment, the linker for interferon α is 15-25 amino acids long. In another specific embodiment, the linker for interferon α is 15-20 amino acids long. In another specific embodiment, the linker for interferon α is 10-25 amino acids long. In another specific embodiment, the linker for interferon α is 15 amino acids long. In one embodiment, the linker for interferon α is $(GGGGS)_n$ (SEQ ID NO:40) where G represents glycine, S represents serine and n is an integer from 1-10. In a specific embodiment, n is 3.

[0143] The linker may also incorporate a moiety capable of being cleaved either chemically (e.g. hydrolysis of an ester bond), enzymatically (i.e. incorporation of a protease cleavage sequence) or photolytically (e.g., a chromophore such as 3-amino-3-(2-nitrophenyl) propionic acid (ANP)) in order to release the biologically active molecule from the Fc protein.

5. Chimeric Protein Dimerization Using Specific Binding Partners

[0144] In one embodiment, the chimeric protein of the invention comprises a first polypeptide chain comprising at least a first domain, said first domain having at least one specific binding partner, and a second polypeptide chain comprising at least a second domain, wherein said second domain, is a specific binding partner of said first domain. The chimeric protein thus comprises a polypeptide capable of dimerizing with another polypeptide due to the interaction of the first domain and the second domain. Methods of dimerizing antibodies using heterologous domains are known in the art (U.S. Patent Nos.: 5,807,706 and 5,910,573; Kostelny et al. 1992, *J. Immunol.* 148(5):1547).

[0145] Dimerization can occur by formation of a covalent bond, or alternatively a non-covalent bond, e.g., hydrophobic interaction, Van der Waal's forces, interdigitation of amphiphilic peptides such as, but not limited to, alpha helices, charge-charge interactions of amino acids bearing opposite charges, such as, but not limited to, lysine and aspartic acid, arginine and glutamic acid. In one embodiment, the domain is a helix bundle comprising a helix, a turn and another helix. In another embodiment, the domain is a leucine zipper comprising a peptide having several repeating amino acids in which every seventh amino acid is a leucine residue. In one embodiment, the specific binding partners are fos/jun. (see Branden et al. 1991, *Introduction To Protein Structure*, Garland Publishing, New York).

[0146] In another embodiment, binding is mediated by a chemical linkage (see, e.g., Brennan et al. 1985, *Science* 229:81). In this embodiment, intact immunoglobulins, or chimeric proteins comprised of at least a portion of an immunoglobulin constant region are cleaved to generate heavy chain fragments. These fragments are reduced in the presence of the dithiol complexing agent sodium arsenite to stabilize vicinal dithiols and prevent intermolecular disulfide formation. The fragments generated are then converted to thionitrobenzoate

(TNB) derivatives. One of the TNB derivatives is then reconverted to the heavy chain fragment thiol by reduction with mercaptoethylamine and is then mixed with an equimolar amount of the other TNB derivative to form a chimeric dimer.

D. Nucleic Acids

[0147] The invention relates to a first nucleic acid construct and a second nucleic acid construct each comprising a nucleic acid sequence encoding at least a portion of the chimeric protein of the invention. In one embodiment, the first nucleic acid construct comprises a nucleic acid sequence encoding a portion of an immunoglobulin constant region operatively linked to a second DNA sequence encoding a biologically active molecule, and said second DNA construct comprises a DNA sequence encoding an immunoglobulin constant region without the second DNA sequence encoding a biologically active molecule.

[0148] The biologically active molecule can include, for example, but not as a limitation, a viral fusion inhibitor, a clotting factor, a growth factor or hormone, or a receptor, or analog, or fragment of any of the preceding. The nucleic acid sequences can also include additional sequences or elements known in the art (e.g., promoters, enhancers, poly A sequences, affinity tags). In one embodiment, the nucleic acid sequence of the second construct can optionally include a nucleic acid sequence encoding a linker placed between the nucleic acid sequence encoding the biologically active molecule and the portion of the immunoglobulin constant region. The nucleic acid sequence of the second DNA construct can optionally include a linker sequence placed before or after the nucleic acid sequence encoding the biologically active molecule and/or the portion of the immunoglobulin constant region.

[0149] In one embodiment, the nucleic acid construct is comprised of DNA. In another embodiment, the nucleic acid construct is comprised of RNA. The nucleic acid construct can be a vector, e.g., a viral vector or a plasmid. Examples of viral vectors include, but are not limited to adeno virus vector, an adeno associated virus vector or a murine leukemia virus vector. Examples of plasmids include but are not limited to pUC, pGEM and pGEX.

[0150] In one embodiment, the nucleic acid construct comprises the nucleic acid sequence of figure 3a (SEQ ID NO:7). In one embodiment, the nucleic acid construct comprises the nucleic acid sequence of figure 3b (SEQ ID NO:9). In

one embodiment, the nucleic acid construct comprises the nucleic acid sequence of figure 3c (SEQ ID NO:11). In one embodiment, the nucleic acid construct comprises the nucleic acid sequence of figure 3d (SEQ ID NO:13). In one embodiment, the nucleic acid construct comprises the nucleic acid sequence of figure 3e (SEQ ID NO:15). In one embodiment, the nucleic acid construct comprises the nucleic acid sequence of figure 3f (SEQ ID NO:17). In one embodiment, the nucleic acid construct comprises the nucleic acid sequence of figure 3g (SEQ ID NO:19). In one embodiment, the nucleic acid construct comprises the nucleic acid sequence of figure 3h (SEQ ID NO:21). In one embodiment, the nucleic acid construct comprises the nucleic acid sequence of figure 3i (SEQ ID NO:23). In one embodiment, the nucleic acid construct comprises the nucleic acid sequence of figure 3j (SEQ ID NO:25). In one embodiment, the nucleic acid construct comprises the nucleic acid sequence of figure 17a (SEQ ID NO:27).

[0151] Due to the known degeneracy of the genetic code, wherein more than one codon can encode the same amino acid, a DNA sequence can vary from that shown in SEQ ID NOS:7, 9, 11, 13, 15, 17, 19, 21, 23, 25 or 27 and still encode a polypeptide having the corresponding amino acid sequence of SEQ ID NOS:6, 8, 10, 12, 14, 16, 18, 20, 22, 24 or 26 respectively. Such variant DNA sequences can result from silent mutations (*e.g.* occurring during PCR amplification), or can be the product of deliberate mutagenesis of a native sequence. The invention thus provides isolated DNA sequences encoding polypeptides of the invention, chosen from: (a) DNA comprising the nucleotide sequence of SEQ ID NOS:7, 9, 11, 13, 15, 17, 19, 21, 23, 25 or 27; (b) DNA encoding the polypeptides of SEQ ID NOS:6, 8, 10, 12, 14, 16, 18, 20, 22, 24 or 26; (c) DNA capable of hybridization to a DNA of (a) or (b) under conditions of moderate stringency and which encodes polypeptides of the invention; (d) DNA capable of hybridization to a DNA of (a) or (b) under conditions of high stringency and which encodes polypeptides of the invention, and (e) DNA which is degenerate as a result of the genetic code to a DNA defined in (a), (b), (c), or (d) and which encode polypeptides of the invention. Of course, polypeptides encoded by such DNA sequences are encompassed by the invention.

[0152] In another embodiment, the nucleic acid molecules comprising a sequence encoding the chimeric protein of the invention can also comprise nucleotide sequences that are at least 80% identical to a native sequence. Also contemplated are embodiments in which a nucleic acid molecules comprising a sequence encoding the chimeric protein of the invention comprises a sequence that is at least 90% identical, at least 95% identical, at least 98% identical, at least 99% identical, or at least 99.9% identical to a native sequence. A native sequence can include any DNA sequence not altered by the human hand. The percent identity may be determined by visual inspection and mathematical calculation. Alternatively, the percent identity of two nucleic acid sequences can be determined by comparing sequence information using the GAP computer program, version 6.0 described by Devereux et al. 1984, *Nucl. Acids Res.* 12:387, and available from the University of Wisconsin Genetics Computer Group (UWGCG). The preferred default parameters for the GAP program include: (1) a unary comparison matrix (containing a value of 1 for identities and 0 for non identities) for nucleotides, and the weighted comparison matrix of Gribskov and Burgess 1986, *Nucl. Acids Res.* 14:6745, as described by Schwartz and Dayhoff, eds. 1979, *Atlas of Protein Sequence and Structure*, National Biomedical Research Foundation, pp. 353-358; (2) a penalty of 3.0 for each gap and an additional 0.10 penalty for each symbol in each gap; and (3) no penalty for end gaps. Other programs used by one skilled in the art of sequence comparison may also be used.

E. Synthesis of Chimeric Proteins

[0153] Chimeric proteins comprising at least a portion of an immunoglobulin constant region and a biologically active molecule can be synthesized using techniques well known in the art. For example, the chimeric proteins of the invention can be synthesized recombinantly in cells (see, e.g., Sambrook et al. 1989, *Molecular Cloning A Laboratory Manual*, Cold Spring Harbor Laboratory, N.Y. and Ausubel et al. 1989, *Current Protocols in Molecular Biology*, Greene Publishing Associates and Wiley Interscience, N.Y.). Alternatively, the chimeric proteins of the invention can be synthesized using known synthetic methods such as solid phase synthesis. Synthetic techniques are well known in the art (see, e.g., Merrifield, 1973, *Chemical Polypeptides*, (Katsoyannis and Panayotis eds.)

pp. 335-61; Merrifield 1963, *J. Am. Chem. Soc.* 85:2149; Davis et al. 1985, *Biochem. Intl.* 10:394; Finn et al. 1976; *The Proteins* (3d ed.) 2:105; Erikson et al. 1976, *The Proteins* (3d ed.) 2:257; U.S. Patent No. 3,941,763. Alternatively, the chimeric proteins of the invention can be synthesized using a combination of recombinant and synthetic methods. In certain applications, it may be beneficial to use either a recombinant method or a combination of recombinant and synthetic methods.

[0154] Nucleic acids encoding a biologically active molecule can be readily synthesized using recombinant techniques well known in the art. Alternatively, the peptides themselves can be chemically synthesized. Nucleic acids of the invention may be synthesized by standard methods known in the art, e.g., by use of an automated DNA synthesizer (such as are commercially available from Biosearch, Applied Biosystems, etc.). As examples, phosphorothioate oligonucleotides may be synthesized by the method of Stein et al. 1988, *Nucl. Acids Res.* 16:3209, methylphosphonate oligonucleotides can be prepared by use of controlled pore glass polymer supports as described in Sarin et al. 1988, *Proc. Natl. Acad. Sci. USA* 85:7448. Additional methods of nucleic acid synthesis are known in the art. (see, e.g., U.S. Patent Nos. 6,015,881; 6,281,331; 6,469,136).

[0155] DNA sequences encoding immunoglobulin constant regions, or fragments thereof, may be cloned from a variety of genomic or cDNA libraries known in the art. The techniques for isolating such DNA sequences using probe-based methods are conventional techniques and are well known to those skilled in the art. Probes for isolating such DNA sequences may be based on published DNA sequences (see, for example, Hieter et al. 1980, *Cell* 22:197-207). The polymerase chain reaction (PCR) method disclosed by Mullis et al. (U.S. Patent No. 4,683,195) and Mullis (U.S. Patent No. 4,683,202) may be used. The choice of library and selection of probes for the isolation of such DNA sequences is within the level of ordinary skill in the art. Alternatively, DNA sequences encoding immunoglobulins or fragments thereof can be obtained from vectors known in the art to contain immunoglobulins or fragments thereof.

[0156] For recombinant production, a first polynucleotide sequence encoding a portion of the chimeric protein of the invention (e.g. a portion of an immunoglobulin constant region) and a second polynucleotide sequence encoding

a portion of the chimeric protein of the invention (e.g. a portion of an immunoglobulin constant region and a biologically active molecule) are inserted into appropriate expression vehicles, *i.e.* vectors which contains the necessary elements for the transcription and translation of the inserted coding sequence, or in the case of an RNA viral vector, the necessary elements for replication and translation. The nucleic acids encoding the chimeric protein are inserted into the vector in proper reading frame.

[0157] The expression vehicles are then transfected or co-transfected into a suitable target cell, which will express the polypeptides. Transfection techniques known in the art include, but are not limited to, calcium phosphate precipitation (Wigler et al. 1978, *Cell* 14:725) and electroporation (Neumann et al. 1982, *EMBO, J.* 1:841), and liposome based reagents. A variety of host-expression vector systems may be utilized to express the chimeric proteins described herein including both prokaryotic or eukaryotic cells. These include, but are not limited to, microorganisms such as bacteria (e.g. *E. coli*) transformed with recombinant bacteriophage DNA or plasmid DNA expression vectors containing an appropriate coding sequence; yeast or filamentous fungi transformed with recombinant yeast or fungi expression vectors containing an appropriate coding sequence; insect cell systems infected with recombinant virus expression vectors (e.g. baculovirus) containing an appropriate coding sequence; plant cell systems infected with recombinant virus expression vectors (e.g. cauliflower mosaic virus or tobacco mosaic virus) or transformed with recombinant plasmid expression vectors (e.g. Ti plasmid) containing an appropriate coding sequence; or animal cell systems, including mammalian cells (e.g. CHO, Cos, HeLa cells).

[0158] When the chimeric protein of the invention is recombinantly synthesized in a prokaryotic cell it may be desirable to refold the chimeric protein. The chimeric protein produced by this method can be refolded to a biologically active conformation using conditions known in the art, e.g., denaturing under reducing conditions and then dialyzed slowly into PBS.

[0159] Depending on the expression system used, the expressed chimeric protein is then isolated by procedures well-established in the art (e.g. affinity chromatography, size exclusion chromatography, ion exchange chromatography).

[0160] The expression vectors can encode for tags that permit for easy purification of the recombinantly produced chimeric protein. Examples include, but are not limited to vector pUR278 (Ruther et al. 1983, *EMBO J.* 2:1791) in which the chimeric protein described herein coding sequences may be ligated into the vector in frame with the lac z coding region so that a hybrid protein is produced; pGEX vectors may be used to express chimeric proteins of the invention with a glutathione S-transferase (GST) tag. These proteins are usually soluble and can easily be purified from cells by adsorption to glutathione-agarose beads followed by elution in the presence of free glutathione. The vectors include cleavage sites (thrombin or Factor Xa protease or PreScission Protease™ (Pharmacia, Peapack, N.J.)) for easy removal of the tag after purification.

[0161] To increase efficiency of production, the polynucleotides can be designed to encode multiple units of the chimeric protein of the invention separated by enzymatic cleavage sites. The resulting polypeptide can be cleaved (e.g. by treatment with the appropriate enzyme) in order to recover the polypeptide units. This can increase the yield of polypeptides driven by a single promoter. When used in appropriate viral expression systems, the translation of each polypeptide encoded by the mRNA is directed internally in the transcript; e.g., by an internal ribosome entry site, IRES. Thus, the polycistronic construct directs the transcription of a single, large polycistronic mRNA which, in turn, directs the translation of multiple, individual polypeptides. This approach eliminates the production and enzymatic processing of polyproteins and may significantly increase yield of polypeptide driven by a single promoter.

[0162] Vectors used in transformation will usually contain a selectable marker used to identify transformants. In bacterial systems, this can include an antibiotic resistance gene such as ampicillin or kanamycin. Selectable markers for use in cultured mammalian cells include genes that confer resistance to drugs, such as neomycin, hygromycin, and methotrexate. The selectable marker may be an amplifiable selectable marker. One amplifiable selectable marker is the DHFR gene. Another amplifiable marker is the DHFR cDNA (Simonsen and Levinson 1983, *Proc. Natl. Acad. Sci. USA* 80:2495). Selectable markers are reviewed by Thilly (*Mammalian Cell Technology*, Butterworth Publishers, Stoneham, MA) and the choice of selectable markers is well within the level of ordinary skill in the art.

[0163] Selectable markers may be introduced into the cell on a separate plasmid at the same time as the gene of interest, or they may be introduced on the same plasmid. If on the same plasmid, the selectable marker and the gene of interest may be under the control of different promoters or the same promoter, the latter arrangement producing a dicistronic message. Constructs of this type are known in the art (for example, U.S. Pat. No. 4,713,339).

[0164] The expression elements of the expression systems vary in their strength and specificities. Depending on the host/vector system utilized, any of a number of suitable transcription and translation elements, including constitutive and inducible promoters, may be used in the expression vector. For example, when cloning in bacterial systems, inducible promoters such as pL of bacteriophage λ , plac, ptrp, ptac (ptrp-lac hybrid promoter) and the like may be used; when cloning in insect cell systems, promoters such as the baculovirus polyhedron promoter may be used; when cloning in plant cell systems, promoters derived from the genome of plant cells (e.g. heat shock promoters; the promoter for the small subunit of RUBISCO; the promoter for the chlorophyll a/b binding protein) or from plant viruses (e.g. the 35S RNA promoter of CaMV; the coat protein promoter of TMV) may be used; when cloning in mammalian cell systems, promoters derived from the genome of mammalian cells (e.g. metallothionein promoter) or from mammalian viruses (e.g. the adenovirus late promoter; the vaccinia virus 7.5 K promoter) may be used; when generating cell lines that contain multiple copies of expression product, SV40-, BPV- and EBV-based vectors may be used with an appropriate selectable marker.

[0165] In cases where plant expression vectors are used, the expression of sequences encoding linear or non-cyclized forms of the chimeric proteins of the invention may be driven by any of a number of promoters. For example, viral promoters such as the 35S RNA and 19S RNA promoters of CaMV (Brisson et al. 1984, *Nature* 310:511-514), or the coat protein promoter of TMV (Takamatsu et al. 1987, *EMBO J.* 6:307-311) may be used; alternatively, plant promoters such as the small subunit of RUBISCO (Coruzzi et al. 1984, *EMBO J.* 3:1671-1680; Broglie et al. 1984, *Science* 224:838-843) or heat shock promoters, e.g., soybean hsp17.5-E or hsp17.3-B (Gurley et al. 1986, *Mol. Cell. Biol.* 6:559-565) may be used. These constructs can be introduced into plant cells using Ti plasmids, Ri

plasmids, plant virus vectors, direct DNA transformation, microinjection, electroporation, etc. For reviews of such techniques see, e.g., Weissbach & Weissbach 1988, *Methods for Plant Molecular Biology*, Academic Press, NY, Section VIII, pp. 421-463; and Grierson & Corey 1988, *Plant Molecular Biology*, 2d Ed., Blackie, London, Ch. 7-9.

[0166] In one insect expression system that may be used to produce the chimeric proteins of the invention, *Autographa californica* nuclear polyhidrosis virus (AcNPV) is used as a vector to express the foreign genes. The virus grows in *Spodoptera frugiperda* cells. A coding sequence may be cloned into non-essential regions (for example, the polyhedron gene) of the virus and placed under control of an AcNPV promoter (for example, the polyhedron promoter). Successful insertion of a coding sequence will result in inactivation of the polyhedron gene and production of non-occluded recombinant virus (*i.e.* virus lacking the proteinaceous coat coded for by the polyhedron gene). These recombinant viruses are then used to infect *Spodoptera frugiperda* cells in which the inserted gene is expressed. (see, e.g., Smith et al. 1983, *J. Virol.* 46:584; U.S. Patent No. 4,215,051). Further examples of this expression system may be found in Ausubel et al., eds. 1989, *Current Protocols in Molecular Biology*, Vol. 2, Greene Publish. Assoc. & Wiley Interscience.

[0167] Another system which can be used to express the chimeric proteins of the invention is the glutamine synthetase gene expression system, also referred to as the "GS expression system" (Lonza Biologics PLC, Berkshire UK). This expression system is described in detail in U.S. Patent No. 5,981,216.

[0168] In mammalian host cells, a number of viral based expression systems may be utilized. In cases where an adenovirus is used as an expression vector, a coding sequence may be ligated to an adenovirus transcription/translation control complex, e.g., the late promoter and tripartite leader sequence. This chimeric gene may then be inserted in the adenovirus genome by in vitro or in vivo recombination. Insertion in a non-essential region of the viral genome (e.g. region E1 or E3) will result in a recombinant virus that is viable and capable of expressing peptide in infected hosts (see, e.g., Logan & Shenk 1984, *Proc. Natl. Acad. Sci. USA* 81:3655). Alternatively, the vaccinia 7.5 K promoter may be used (see, e.g., Mackett et al. 1982, *Proc. Natl. Acad. Sci.*

USA 79:7415; Mackett et al. 1984, *J. Virol.* 49:857; Panicali et al. 1982, *Proc. Natl. Acad. Sci. USA* 79:4927).

[0169] In cases where an adenovirus is used as an expression vector, a coding sequence may be ligated to an adenovirus transcription/translation control complex, e.g., the late promoter and tripartite leader sequence. This chimeric gene may then be inserted in the adenovirus genome by in vitro or in vivo recombination. Insertion in a non-essential region of the viral genome (e.g. region E1 or E3) will result in a recombinant virus that is viable and capable of expressing peptide in infected hosts (see, e.g., Logan & Shenk 1984, *Proc. Natl. Acad. Sci. USA* 81:3655). Alternatively, the vaccinia 7.5 K promoter may be used (see, e.g., Mackett et al. 1982, *Proc. Natl. Acad. Sci. USA* 79:7415; Mackett et al. 1984, *J. Virol.* 49:857; Panicali et al. 1982, *Proc. Natl. Acad. Sci. USA* 79:4927).

[0170] Host cells containing DNA constructs of the chimeric protein are grown in an appropriate growth medium. As used herein, the term "appropriate growth medium" means a medium containing nutrients required for the growth of cells. Nutrients required for cell growth may include a carbon source, a nitrogen source, essential amino acids, vitamins, minerals and growth factors. Optionally the media can contain bovine calf serum or fetal calf serum. In one embodiment, the media contains substantially no IgG. The growth medium will generally select for cells containing the DNA construct by, for example, drug selection or deficiency in an essential nutrient which is complemented by the selectable marker on the DNA construct or co-transfected with the DNA construct. Cultured mammalian cells are generally grown in commercially available serum-containing or serum-free media (e.g. MEM, DMEM). Selection of a medium appropriate for the particular cell line used is within the level of ordinary skill in the art.

[0171] The recombinantly produced chimeric protein of the invention can be isolated from the culture media. The culture medium from appropriately grown transformed or transfected host cells is separated from the cell material, and the presence of chimeric proteins is demonstrated. One method of detecting the chimeric proteins, for example, is by the binding of the chimeric proteins or portions of the chimeric proteins to a specific antibody recognizing the chimeric protein of the invention. An anti-chimeric protein antibody may be a monoclonal or polyclonal antibody raised against the chimeric protein in question. For

example, the chimeric protein contains at least a portion of an immunoglobulin constant region. Antibodies recognizing the constant region of many immunoglobulins are known in the art and are commercially available. An antibody can be used to perform an ELISA or a western blot to detect the presence of the chimeric protein of the invention.

[0172] The chimeric protein of the invention can be synthesized in a transgenic animal, such as a rodent, cow, pig, sheep, or goat. The term "transgenic animals" refers to non-human animals that have incorporated a foreign gene into their genome. Because this gene is present in germline tissues, it is passed from parent to offspring. Exogenous genes are introduced into single-celled embryos (Brinster et al. 1985, *Proc. Natl. Acad. Sci. USA* 82:4438). Methods of producing transgenic animals are known in the art, including transgenics that produce immunoglobulin molecules (Wagner et al. 1981, *Proc. Natl. Acad. Sci. USA* 78:6376; McKnight et al. 1983, *Cell* 34:335; Brinster et al. 1983, *Nature* 306:332; Ritchie et al. 1984, *Nature* 312:517; Baldassarre et al. 2003, *Theriogenology* 59:831; Robl et al. 2003, *Theriogenology* 59:107; Malassagne et al. 2003, *Xenotransplantation* 10(3):267).

[0173] The chimeric protein of the invention can also be produced by a combination of synthetic chemistry and recombinant techniques. For example, the portion of an immunoglobulin constant region can be expressed recombinantly as described above. The biologically active molecule, can be produced using known chemical synthesis techniques (e.g. solid phase synthesis):

[0174] The portion of an immunoglobulin constant region can be ligated to the biologically active molecule using appropriate ligation chemistry and then combined with a portion of an immunoglobulin constant region that has not been ligated to a biologically active molecule to form the chimeric protein of the invention. In one embodiment, the portion of an immunoglobulin constant region is an Fc fragment. The Fc fragment can be recombinantly produced to form Cys-Fc and reacted with a biologically active molecule expressing a thioester to make a monomer-dimer hybrid. In another embodiment, an Fc-thioester is made and reacted with a biologically active molecule expressing an N terminus Cysteine (Figure 4).

[0175] In one embodiment, the portion of an immunoglobulin constant region ligated to the biologically active molecule will form homodimers. The homodimers can be disrupted by exposing the homodimers to denaturing and reducing conditions (e.g. beta-mercaptoethanol and 8M urea) and then subsequently combined with a portion of an immunoglobulin constant region not linked to a biologically active molecule to form monomer-dimer hybrids. The monomer-dimer hybrids are then renatured and refolded by dialyzing into PBS and isolated, e.g., by size exclusion or affinity chromatography.

[0176] In another embodiment, the portion of an immunoglobulin constant region will form homodimers before being linked to a biologically active molecule. In this embodiment, reaction conditions for linking the biologically active molecule to the homodimer can be adjusted such that linkage of the biologically active molecule to only one chain of the homodimer is favored (e.g. by adjusting the molar equivalents of each reactant).

[0177] The biologically active molecule can be chemically synthesized with an N terminal cysteine. The sequence encoding a portion of an immunoglobulin constant region can be sub-cloned into a vector encoding intein linked to a chitin binding domain (New England Biolabs, Beverly, MA). The intein can be linked to the C terminus of the portion of an immunoglobulin constant region. In one embodiment, the portion of the immunoglobulin with the intein linked to its C terminus can be expressed in a prokaryotic cell. In another embodiment, the portion of the immunoglobulin with the intein linked to its C terminus can be expressed in a eukaryotic cell. The portion of immunoglobulin constant region linked to intein can be reacted with MESNA. In one embodiment, the portion of an immunoglobulin constant region linked to intein is bound to a column, e.g., a chitin column and then eluted with MESNA. The biologically active molecule and portion of an immunoglobulin can be reacted together such that nucleophilic rearrangement occurs and the biologically active molecule is covalently linked to the portion of an immunoglobulin via an amide bond. (Dawsen et al. 2000, *Annu. Rev. Biochem.* 69:923). The chimeric protein synthesized this way can optionally include a linker peptide between the portion of an immunoglobulin and the biologically active molecule. The linker can for example be synthesized on the N terminus of the biologically active molecule. Linkers can include peptides and/or

organic molecules (e.g. polyethylene glycol and/or short amino acid sequences). This combined recombinant and chemical synthesis allows for the rapid screening of biologically active molecules and linkers to optimize desired properties of the chimeric protein of the invention, e.g., viral inhibition, hemostasis, production of red blood cells, biological half-life, stability, binding to serum proteins or some other property of the chimeric protein. The method also allows for the incorporation of non-natural amino acids into the chimeric protein of the invention which may be useful for optimizing a desired property of the chimeric protein of the invention. If desired, the chimeric protein produced by this method can be refolded to a biologically active conformation using conditions known in the art, e.g., reducing conditions and then dialyzed slowly into PBS.

[0178] Alternatively, the N-terminal cysteine can be on the portion of an immunoglobulin constant region, e.g., an Fc fragment. An Fc fragment can be generated with an N- terminal cysteine by taking advantage of the fact that a native Fc has a cysteine at position 226 (see Kabat et al. 1991, *Sequences of Proteins of Immunological Interest*, U.S. Department of Public Health, Bethesda, MD).

[0179] To expose a terminal cysteine, an Fc fragment can be recombinantly expressed. In one embodiment, the Fc fragment is expressed in a prokaryotic cell, e.g., E.coli. The sequence encoding the Fc portion beginning with Cys 226 (EU numbering) can be placed immediately following a sequence encoding a signal peptide, e.g., OmpA, PhoA, STII. The prokaryotic cell can be osmotically shocked to release the recombinant Fc fragment. In another embodiment, the Fc fragment is produced in a eukaryotic cell, e.g., a CHO cell, a BHK cell. The sequence encoding the Fc portion fragment can be placed directly following a sequence encoding a signal peptide, e.g., mouse Igk light chain or MHC class I Kb signal sequence, such that when the recombinant chimeric protein is synthesized by a eukaryotic cell, the signal sequence will be cleaved, leaving an N terminal cysteine which can then be isolated and chemically reacted with a molecule bearing a thioester (e.g. a C terminal thioester if the molecule is comprised of amino acids).

[0180] The N terminal cysteine on an Fc fragment can also be generated using an enzyme that cleaves its substrate at its N terminus, e.g., Factor X^a,

enterokinase, and the product isolated and reacted with a molecule with a thioester.

[0181] The recombinantly expressed Fc fragment can be used to make homodimers or monomer-dimer hybrids.

[0182] In a specific embodiment, an Fc fragment is expressed with the human α interferon signal peptide adjacent to the Cys at position 226. When a construct encoding this polypeptide is expressed in CHO cells, the CHO cells cleave the signal peptide at two distinct positions (at Cys 226 and at Val within the signal peptide 2 amino acids upstream in the N terminus direction). This generates a mixture of two species of Fc fragments (one with an N-terminal Val and one with an N-terminal Cys). This in turn results in a mixture of dimeric species (homodimers with terminal Val, homodimers with terminal Cys and heterodimers where one chain has a terminal Cys and the other chain has a terminal Val). The Fc fragments can be reacted with a biologically active molecule having a C terminal thioester and the resulting monomer-dimer hybrid can be isolated from the mixture (e.g. by size exclusion chromatography). It is contemplated that when other signal peptide sequences are used for expression of Fc fragments in CHO cells a mixture of species of Fc fragments with at least two different N termini will be generated.

[0183] In another embodiment, a recombinantly produced Cys-Fc can form a homodimer. The homodimer can be reacted with peptide that has a branched linker on the C terminus, wherein the branched linker has two C terminal thioesters that can be reacted with the Cys-Fc. In another embodiment, the biologically active molecule has a single non-terminal thioester that can be reacted with Cys-Fc. Alternatively, the branched linker can have two C terminal cysteines that can be reacted with an Fc thioester. In another embodiment, the branched linker has two functional groups that can be reacted with the Fc thioester, e.g., 2-mercaptoamine. The biologically active molecule may be comprised of amino acids. The biologically active molecule may include a small organic molecule or a small inorganic molecule.

F. Methods of Using Chimeric Proteins

[0184] The chimeric proteins of the invention have many uses as will be recognized by one skilled in the art, including, but not limited to methods of

treating a subject with a disease or condition. The disease or condition can include, but is not limited to, a viral infection, a hemostatic disorder, anemia, cancer, leukemia, an inflammatory condition or an autoimmune disease (e.g. arthritis, psoriasis, lupus erythematosus, multiple sclerosis), or a bacterial infection (see, e.g., U.S. Patent Nos. 6,086,875, 6,030,613, 6,485,726; WO 03/077834; US2003-0235536A1).

1. Methods of Treating Cardiovascular Diseases or Conditions

[0185] In one embodiment, the invention relates to a method of treating a subject having a cardiovascular disease or condition, comprising administering a therapeutically effective amount of at least one chimeric protein, wherein the chimeric protein comprises a first and a second polypeptide chain, wherein the first chain comprises at least a portion of an immunoglobulin constant region and at least one agent capable of promoting mitosis, promoting angiogenesis and/or inhibiting apoptosis, such as erythropoietin, and the second polypeptide chain comprises at least a portion of an immunoglobulin without said agent. The cardiovascular disease or condition can include, but is not limited to, stroke, brain ischemia, cerebral ischemia, myocardial ischemia, cardiac ischemia, ischemic heart disease, myocardial infarction, coronary artery disease, acute coronary syndrome, atherosclerosis, chronic heart failure, congestive heart failure, or a reperfusion injury.

[0186] In one embodiment, administering a therapeutically effective amount of at least one chimeric protein of the invention to a subject having a cardiovascular disease or condition promotes vascularization, organ growth, mitogenesis, angiogenesis, endothelial cell proliferation, endothelial cell differentiation, vascular repair, cardiovascular regeneration, or infarct reduction. In one embodiment, administering a therapeutically effective amount of at least one chimeric protein of the invention inhibits apoptosis, limits the sequelae of cerebral ischemia, reduces ventricular mass, ameliorates exercise-related cardiac ischemia, enhances exercise capacity, or protects neurons from ischemic damage.

2. Methods of Treating a Subject with a Red Blood Cell Deficiency

[0187] The invention relates to a method of treating a subject having a deficiency of red blood cells, e.g., anemia, comprising administering a

therapeutically effective amount of at least one chimeric protein, wherein the chimeric protein comprises a first and a second polypeptide chain, wherein the first chain comprises at least a portion of an immunoglobulin constant region and at least one agent capable of inducing proliferation of red blood cells, *e.g.*, EPO, and the second polypeptide chain comprises at least a portion of an immunoglobulin without the agent capable of inducing red blood cell proliferation of the first chain.

3. Methods of Treating a Subject with a Viral Infection

[0188] The invention relates to a method of treating a subject having a viral infection or exposed to a virus comprising administering a therapeutically effective amount of at least one chimeric protein, wherein the chimeric protein comprises a first and a second polypeptide chain, wherein the first chain comprises at least a portion of an immunoglobulin constant region and at least one antiviral agent, *e.g.*, a fusion inhibitor or interferon α and the second polypeptide chain comprises at least a portion of an immunoglobulin without the antiviral agent of the first chain. In one embodiment, the subject is infected with a virus which can be treated with IFN α , *e.g.*, hepatitis C virus. In one embodiment, the subject is infected with HIV, such as HIV-1 or HIV-2.

[0189] In one embodiment, the chimeric protein of the invention inhibits viral replication. In one embodiment, the chimeric protein of the invention prevents or inhibits viral entry into target cells, thereby stopping, preventing, or limiting the spread of a viral infection in a subject and decreasing the viral burden in an infected subject. By linking a portion of an immunoglobulin to a viral fusion inhibitor the invention provides a chimeric protein with viral fusion inhibitory activity with greater stability and greater bioavailability compared to viral fusion inhibitors alone, *e.g.*, T20, T21, T1249. Thus, in one embodiment, the viral fusion inhibitor decreases or prevents HIV infection of a target cell, *e.g.*, HIV-1.

a. Conditions That May Be Treated

[0190] The chimeric protein of the invention can be used to inhibit or prevent the infection of a target cell by a hepatitis virus, *e.g.*, hepatitis virus C. The chimeric protein may comprise an anti-viral agent which inhibits viral replication.

[0191] In one embodiment, the chimeric protein of the invention comprises a fusion inhibitor. The chimeric protein of the invention can be used to inhibit or prevent the infection of any target cell by any virus (see, e.g., U.S. Patent Nos. 6,086,875, 6,030,613, 6,485,726; WO 03/077834; US2003-0235536A1). In one embodiment, the virus is an enveloped virus such as, but not limited to HIV, SIV, measles, influenza, Epstein-Barr virus, respiratory syncytia virus, or parainfluenza virus. In another embodiment, the virus is a non-enveloped virus such as rhino virus or polio virus

[0192] The chimeric protein of the invention can be used to treat a subject already infected with a virus. The subject can be acutely infected with a virus. Alternatively, the subject can be chronically infected with a virus. The chimeric protein of the invention can also be used to prophylactically treat a subject at risk for contracting a viral infection, e.g., a subject known or believed to in close contact with a virus or subject believed to be infected or carrying a virus. The chimeric protein of the invention can be used to treat a subject who may have been exposed to a virus, but who has not yet been positively diagnosed.

[0193] In one embodiment, the invention relates to a method of treating a subject infected with HCV comprising administering to the subject a therapeutically effective amount of a chimeric protein, wherein the chimeric protein comprises an Fc fragment of an IgG and a cytokine, e.g., IFN α .

[0194] In one embodiment, the invention relates to a method of treating a subject infected with HIV comprising administering to the subject a therapeutically effective amount of a chimeric protein wherein the chimeric protein comprises an Fc fragment of an IgG and the viral fusion inhibitor comprises T20.

4. Methods of Treating a Subject Having a Hemostatic Disorder

[0195] The invention relates to a method of treating a subject having a hemostatic disorder comprising administering a therapeutically effective amount of at least one chimeric protein, wherein the chimeric protein comprises a first and a second chain, wherein the first chain comprises at least one clotting factor and at least a portion of an immunoglobulin constant region, and the second chain comprises at least a portion of an immunoglobulin constant region.

[0196] The chimeric protein of the invention treats or prevents a hemostatic disorder by promoting the formation of a fibrin clot. The chimeric protein of the

invention can activate any member of a coagulation cascade. The clotting factor can be a participant in the extrinsic pathway, the intrinsic pathway or both. In one embodiment, the clotting factor is Factor VII or Factor VIIa. Factor VIIa can activate Factor X which interacts with Factor Va to cleave prothrombin to thrombin, which in turn cleaves fibrinogen to fibrin. In another embodiment, the clotting factor is Factor IX or Factor IXa. In yet another embodiment, the clotting factor is Factor VIII or Factor VIIIa. In yet another embodiment, the clotting factor is von Willebrand Factor, Factor XI, Factor XII, Factor V, Factor X or Factor XIII.

a. Conditions That May Be Treated

[0197] The chimeric protein of the invention can be used to treat any hemostatic disorder. The hemostatic disorders that may be treated by administration of the chimeric protein of the invention include, but are not limited to, hemophilia A, hemophilia B, von Willebrand's disease, Factor XI deficiency (PTA deficiency), Factor XII deficiency, as well as deficiencies or structural abnormalities in fibrinogen, prothrombin, Factor V, Factor VII, Factor X, or Factor XIII.

[0198] In one embodiment, the hemostatic disorder is an inherited disorder. In one embodiment, the subject has hemophilia A, and the chimeric protein comprises Factor VIII or Factor VIIIa. In another embodiment, the subject has hemophilia A and the chimeric protein comprises Factor VII or Factor VIIa. In another embodiment, the subject has hemophilia B and the chimeric protein comprises Factor IX or Factor IXa. In another embodiment, the subject has hemophilia B and the chimeric protein comprises Factor VII or Factor VIIa. In another embodiment, the subject has inhibitory antibodies to Factor VIII or Factor VIIIa and the chimeric protein comprises Factor VII or Factor VIIa. In yet another embodiment, the subject has inhibitory antibodies against Factor IX or Factor IXa and the chimeric protein comprises Factor VII or Factor VIIa.

[0199] The chimeric protein of the invention can be used to prophylactically treat a subject with a hemostatic disorder. The chimeric protein of the invention can be used to treat an acute bleeding episode in a subject with a hemostatic disorder

[0200] In one embodiment, the hemostatic disorder is the result of a deficiency in a clotting factor, e.g., Factor IX, Factor VIII. In another embodiment,

the hemostatic disorder can be the result of a defective clotting factor, e.g., von Willebrand's Factor.

[0201] In another embodiment, the hemostatic disorder can be an acquired disorder. The acquired disorder can result from an underlying secondary disease or condition. The unrelated condition can be, as an example, but not as a limitation, cancer, an autoimmune disease, or pregnancy. The acquired disorder can result from old age or from medication to treat an underlying secondary disorder (e.g. cancer chemotherapy).

5. Methods of Treating a Subject In Need of a General Hemostatic Agent

[0202] The invention also relates to methods of treating a subject that does not have a hemostatic disorder or a secondary disease or condition resulting in acquisition of a hemostatic disorder. The invention thus relates to a method of treating a subject in need of a general hemostatic agent comprising administering a therapeutically effective amount of at least one chimeric protein, wherein the chimeric protein comprises a first and a second polypeptide chain wherein the first polypeptide chain comprises at least a portion of an immunoglobulin constant region and at least one clotting factor and the second chain comprises at least a portion of an immunoglobulin constant region without the clotting factor of the first polypeptide chain.

a. Conditions That May Be Treated

[0203] In one embodiment, the subject in need of a general hemostatic agent is undergoing, or is about to undergo, surgery. The chimeric protein of the invention can be administered prior to or after surgery as a prophylactic. The chimeric protein of the invention can be administered during or after surgery to control an acute bleeding episode. The surgery can include, but is not limited to, liver transplantation, liver resection, or stem cell transplantation.

[0204] The chimeric protein of the invention can be used to treat a subject having an acute bleeding episode who does not have a hemostatic disorder. The acute bleeding episode can result from severe trauma, e.g., surgery, an automobile accident, wound, laceration gun shot, or any other traumatic event resulting in uncontrolled bleeding.

6. Treatment Modalities

[0205] The chimeric protein of the invention can be administered intravenously, subcutaneously, intra-muscularly, or via any mucosal surface, e.g., orally, sublingually, buccally, sublingually, nasally, rectally, vaginally or via pulmonary route. The chimeric protein can be implanted within or linked to a biopolymer solid support that allows for the slow release of the chimeric protein to the desired site.

[0206] The dose of the chimeric protein of the invention will vary depending on the subject and upon the particular route of administration used. Dosages can range from 0.1 to 100,000 µg/kg body weight. In one embodiment, the dosing range is 0.1-1,000 µg/kg. The protein can be administered continuously or at specific timed intervals. In vitro assays may be employed to determine optimal dose ranges and/or schedules for administration. Many in vitro assays that measure viral infectivity are known in the art. For example, a reverse transcriptase assay, or an rt PCR assay or branched DNA assay can be used to measure HIV concentrations. A StaClot assay can be used to measure clotting activity. Additionally, effective doses may be extrapolated from dose-response curves obtained from animal models.

[0207] The invention also relates to a pharmaceutical composition comprising a viral fusion inhibitor, at least a portion of an immunoglobulin and a pharmaceutically acceptable carrier or excipient. Examples of suitable pharmaceutical carriers are described in *Remington's Pharmaceutical Sciences* by E.W. Martin. Examples of excipients can include starch, glucose, lactose, sucrose, gelatin, malt, rice, flour, chalk, silica gel, sodium stearate, glycerol monostearate, talc, sodium chloride, dried skim milk, glycerol, propylene, glycol, water, ethanol, and the like. The composition can also contain pH buffering reagents, and wetting or emulsifying agents.

[0208] For oral administration, the pharmaceutical composition can take the form of tablets or capsules prepared by conventional means. The composition can also be prepared as a liquid for example a syrup or a suspension. The liquid can include suspending agents (e.g. sorbitol syrup, cellulose derivatives or hydrogenated edible fats), emulsifying agents (lecithin or acacia), non-aqueous vehicles (e.g. almond oil, oily esters, ethyl alcohol, or fractionated vegetable oils),

and preservatives (e.g. methyl or propyl -p-hydroxybenzoates or sorbic acid). The preparations can also include flavoring, coloring and sweetening agents. Alternatively, the composition can be presented as a dry product for constitution with water or another suitable vehicle.

[0209] For buccal and sublingual administration the composition may take the form of tablets, lozenges or fast dissolving films according to conventional protocols.

[0210] For administration by inhalation, the compounds for use according to the present invention are conveniently delivered in the form of an aerosol spray from a pressurized pack or nebulizer (e.g. in PBS), with a suitable propellant, e.g., dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoromethane, carbon dioxide or other suitable gas. In the case of a pressurized aerosol the dosage unit can be determined by providing a valve to deliver a metered amount. Capsules and cartridges of, e.g., gelatin for use in an inhaler or insufflator can be formulated containing a powder mix of the compound and a suitable powder base such as lactose or starch.

[0211] The pharmaceutical composition can be formulated for parenteral administration (*i.e.* intravenous or intramuscular) by bolus injection. Formulations for injection can be presented in unit dosage form, e.g., in ampoules or in multidose containers with an added preservative. The compositions can take such forms as suspensions, solutions, or emulsions in oily or aqueous vehicles, and contain formulatory agents such as suspending, stabilizing and/or dispersing agents. Alternatively, the active ingredient can be in powder form for constitution with a suitable vehicle, e.g., pyrogen free water.

[0212] The pharmaceutical composition can also be formulated for rectal administration as a suppository or retention enema, e.g., containing conventional suppository bases such as cocoa butter or other glycerides.

7. Combination Therapy

[0213] The chimeric protein of the invention can be used to treat a subject with a disease or condition in combination with at least one other known agent to treat said disease or condition.

[0214] In one embodiment, the invention relates to a method of treating a subject infected with HIV comprising administering a therapeutically effective

amount of at least one chimeric protein comprising a first and a second chain, wherein the first chain comprises an HIV fusion inhibitor and at least a portion of an immunoglobulin constant region and the second chain comprises at least a portion of an immunoglobulin without an HIV fusion inhibitor of the first chain, in combination with at least one other anti-HIV agent. Said other anti-HIV agent can be any therapeutic with demonstrated anti-HIV activity. Said other anti-HIV agent can include, as an example, but not as a limitation, a protease inhibitor (e.g. Amprenavir[®], Crixivan[®], Ritonivir[®]), a reverse transcriptase nucleoside analog (e.g. AZT, DDI, D4T, 3TC, Ziagen[®]), a nonnucleoside analog reverse transcriptase inhibitor (e.g. Sustiva[®]), another HIV fusion inhibitor, a neutralizing antibody specific to HIV, an antibody specific to CD4, a CD4 mimic, e.g., CD4-IgG2 fusion protein (U.S. Patent Application 09/912,824) or an antibody specific to CCR5, or CXCR4, or a specific binding partner of CCR5, or CXCR4.

[0215] In another embodiment, the invention relates to a method of treating a subject with a hemostatic disorder comprising administering a therapeutically effective amount of at least one chimeric protein comprising a first and a second chain, wherein the first chain comprises at least one clotting factor and at least a portion of an immunoglobulin constant region and the second chain comprises at least a portion of an immunoglobulin constant region without the clotting factor of the first chain, in combination with at least one other clotting factor or agent that promotes hemostasis. Said other clotting factor or agent that promotes hemostasis can be any therapeutic with demonstrated clotting activity. As an example, but not as a limitation, the clotting factor or hemostatic agent can include Factor V, Factor VII, Factor VIII, Factor IX, Factor X, Factor XI, Factor XII, Factor XIII, prothrombin, or fibrinogen or activated forms of any of the preceding. The clotting factor of hemostatic agent can also include anti-fibrinolytic drugs, e.g., epsilon-amino-caproic acid, tranexamic acid.

8. Methods of Inhibiting Viral Fusion With a Target Cell

[0216] The invention also relates to an in vitro method of inhibiting HIV fusion with a mammalian cell comprising combining the mammalian cell with at least one chimeric protein, wherein the chimeric protein comprises a first and a second chain, wherein the first chain comprises at least a portion of an immunoglobulin constant region and an HIV inhibitor and the second chain

comprises at least a portion of an immunoglobulin constant region without the HIV inhibitor of the first chain. The mammalian cell can include any cell or cell line susceptible to infection by HIV including but not limited to primary human CD4⁺ T cells or macrophages, MOLT-4 cells, CEM cells, AA5 cells or HeLa cells which express CD4 on the cell surface.

G. Methods of Isolating Chimeric Proteins

[0217] Typically, when chimeric proteins of the invention are produced they are contained in a mixture of other molecules such as other proteins or protein fragments. The invention thus provides for methods of isolating any of the chimeric proteins described supra from a mixture containing the chimeric proteins. It has been determined that the chimeric proteins of the invention bind to dye ligands under suitable conditions and that altering those conditions subsequent to binding can disrupt the bond between the dye ligand and the chimeric protein, thereby providing a method of isolating the chimeric protein. In some embodiments the mixture may comprise a monomer-dimer hybrid, a dimer and at least a portion of an immunoglobulin constant region, e.g., an Fc. Thus, in one embodiment, the invention provides a method of isolating a monomer-dimer hybrid. In another embodiment, the invention provides a method of isolating a dimer.

[0218] Accordingly, in one embodiment, the invention provides a method of isolating a monomer-dimer hybrid from a mixture, where the mixture comprises

a) the monomer-dimer hybrid comprising a first and second polypeptide chain, wherein the first chain comprises a biologically active molecule, and at least a portion of an immunoglobulin constant region and wherein the second chain comprises at least a portion of an immunoglobulin constant region without a biologically active molecule or immunoglobulin variable region;

b) a dimer comprising a first and second polypeptide chain, wherein the first and second chains both comprise a biologically active molecule, and at least a portion of an immunoglobulin constant region; and

c) a portion of an immunoglobulin constant region; said method comprising

- 1) contacting the mixture with a dye ligand linked to a solid support under suitable conditions such that both the monomer-dimer hybrid and the dimer bind to the dye ligand;

- 2) removing the unbound portion of an immunoglobulin constant region;
- 3) altering the suitable conditions of 1) such that the binding between the monomer-dimer hybrid and the dye ligand linked to the solid support is disrupted;
- 4) isolating the monomer-dimer hybrid.

In some embodiments, prior to contacting the mixture with a dye ligand, the mixture may be contacted with a chromatographic substance such as protein A sepharose or the like. The mixture is eluted from the chromatographic substance using an appropriate elution buffer (e.g. a low pH buffer) and the eluate containing the mixture is then contacted with the dye ligand.

[0219] Suitable conditions for contacting the mixture with the dye ligand may include a buffer to maintain the mixture at an appropriate pH. An appropriate pH may include a pH of from, 3-10, 4-9, 5-8. In one embodiment, the appropriate pH is 8.0. Any buffering agent known in the art may be used so long as it maintains the pH in the appropriate range, e.g., tris, HEPES, PIPES, MOPS. Suitable conditions may also include a wash buffer to elute unbound species from the dye ligand. The wash buffer may be any buffer which does not disrupt binding of a bound species. For example, the wash buffer can be the same buffer used in the contacting step.

[0220] Once the chimeric protein is bound to the dye ligand, the chimeric protein is isolated by altering the suitable conditions. Altering the suitable conditions may include the addition of a salt to the buffer. Any salt may be used, e.g., NaCl, KCl. The salt should be added at a concentration that is high enough to disrupt the binding between the dye ligand and the desired species, e.g., a monomer-dimer hybrid.

[0221] In some embodiments where the mixture is comprised of an Fc, a monomer-dimer hybrid, and a dimer, it has been found that the Fc does not bind to the dye ligand and thus elutes with the flow through. The dimer binds more tightly to the dye ligand than the monomer-dimer hybrid. Thus a higher concentration of salt is required to disrupt the bond (e.g. elute) between the dimer and the dye ligand compared to the salt concentration required to disrupt the bond between the dye ligand and the monomer-dimer hybrid.

[0222] In some embodiments NaCl may be used to isolate the monomer-dimer hybrid from the mixture. In some embodiments the appropriate concentration of salt which disrupts the bond between the dye ligand and the monomer-dimer hybrid is from 200-700 mM, 300-600 mM, 400-500 mM. In one embodiment, the concentration of NaCl required to disrupt the binding between the dye ligand the monomer-dimer hybrid is 400 mM.

[0223] NaCl may also be used to isolate the dimer from the mixture. Typically, the monomer-dimer hybrid is isolated from the mixture before the dimer. The dimer is isolated by adding an appropriate concentration of salt to the buffer, thereby disrupting the binding between the dye ligand and the dimer. In some embodiments the appropriate concentration of salt which disrupts the bond between the dye ligand and the dimer is from 800 mM to 2 M, 900 mM to 1.5 M, 950 mM to 1.2 M. In one specific embodiment, 1 M NaCl is used to disrupt the binding between the dye ligand and the dimer.

[0224] The dye ligand may be a bio-mimetic. A bio-mimetic is a human-made substance, device, or system that imitates nature. Thus in some embodiments the dye ligand imitates a molecule's naturally occurring ligand. The dye ligand may be chosen from Mimetic Red 1TM, Mimetic Red 2TM, Mimetic Orange 1TM, Mimetic Orange 2TM, Mimetic Orange 3TM, Mimetic Yellow 1TM, Mimetic Yellow 2TM, Mimetic Green 1TM, Mimetic Blue 1TM, and Mimetic Blue 2TM (Prometic Biosciences (USA) Inc., Wayne, NJ). In one specific embodiment, the dye ligand is Mimetic Red 2TM (Prometic Biosciences (USA) Inc., Wayne, NJ). In certain embodiments the dye ligand is linked to a solid support, e.g., from Mimetic Red 1A6XLTM, Mimetic Red 2 A6XLTM, Mimetic Orange 1 A6XLTM, Mimetic Orange 2 A6XLTM, Mimetic Orange 3 A6XLTM, Mimetic Yellow 1 A6XLTM, Mimetic Yellow 2 A6XLTM, Mimetic Green 1 A6XLTM, Mimetic Blue 1 A6XLTM, and Mimetic Blue 2 A6XLTM (Prometic Biosciences (USA) Inc., Wayne, NJ).

[0225] The dye ligand may be linked to a solid support. The solid support may be any solid support known in the art (see, e.g., www.seperationsNOW.com). Examples of solid supports may include a bead, a gel, a membrane, a nanoparticle, or a microsphere. The solid support may comprise any material which can be linked to a dye ligand (e.g. agarose, polystyrene, sepharose, sephadex). Solid supports may comprise any synthetic organic polymer such as

polyacrylic, vinyl polymers, acrylate, polymethacrylate, and polyacrylamide. Solid supports may also comprise a carbohydrate polymer, e.g., agarose, cellulose, or dextran. Solid supports may comprise inorganic oxides, such as silica, zirconia, titania, ceria, alumina, magnesia (i.e., magnesium oxide), or calcium oxide. Solid supports may also comprise combinations of some of the above-mentioned supports including, but not limited to, dextran-acrylamide.

Examples

Example 1: Molecular Weight Affects FcRn Mediated Transcytosis

[0226] Chimeric proteins comprised of various proteins of interest and IgG Fc were recombinantly produced (Sambrook et al. *Molecular Cloning: A Laboratory Manual*, 2 ed., Cold Spring Harbor Laboratory Press, (1989)) or in the case of contactin-Fc, MAB- β -gal, (a complex of a monoclonal antibody bound to β -gal) (Biodesign International, Saco, ME) and MAB-GH (a complex of monoclonal antibody and growth hormone)(Research Diagnostics, Inc. Flanders, NJ) were purchased commercially. Briefly, the genes encoding the protein of interest were cloned by PCR, and then sub-cloned into an Fc fusion expression plasmid. The plasmids were transfected into DG44 CHO cells and stable transfectants were selected and amplified with methotrexate. The chimeric protein homodimers were purified over a protein A column. The proteins tested included interferon α , growth hormone, erythropoietin, follicle stimulating hormone, Factor IX, beta-galactosidase, contactin, and Factor VIII. Linking the proteins to immunoglobulin portions, including the FcRn receptor binding partner, or using commercially available whole antibody (including the FcRn binding region)-antigen complexes permitted the investigation of transcytosis as a function of molecular weight (see U.S. Patent No. 6,030,613). The chimeric proteins were administered to rats orally and serum levels were measured 2-4 hours post administration using an ELISA for recombinantly produced chimeric proteins and both a western blot and ELISA for commercially obtained antibody complexes and chimeric proteins. Additionally, all of the commercially obtained proteins or complexes as well as Factor VIII-Fc, Factor IX-Fc and Epo-Fc controls were iodinated using IODO beads (Pierce, Pittsburgh, PA). The results indicated serum levels of Fc and monoclonal antibody chimeric proteins orally administered to rats are directly

related to the size of the protein. The apparent cutoff point for orally administered Fc chimeric proteins is between 200-285 kD. (Table 2).

TABLE 2

Protein	Size (kD)	Transcytosis
IFN α -Fc	92	++++
GH-Fc	96	+++
Epo-Fc	120	+++
FSH-Fc	170	+++
MAB:GH	172-194	+++
FIX-Fc	200	+
MAB: β Gal	285-420	-
Contactin-Fc	300	-
FVIII Δ -Fc	380	-

Example 2: Cloning of pcDNA 3.1-Flag-Fc

[0227] The sequence for the FLAG peptide (Asp-Tyr-Lys-Asp-Asp-Asp-Asp-Lys), a common affinity tag used to identify or purify proteins, was cloned into the pcDNA 3.1-Fc plasmid, which contains the mouse Igk signal sequence followed by the Fc fragment of human IgG1 (amino acids 221-447, EU numbering). The construct was created by overlapping PCR using the following primers:

FlagFc-F1: 5'- GCTGGCTAGCCACCATGGA -3'(SEQ ID NO:41)

FlagFc-R1: 5'- CTTGTCATCGTCGTCCTTGTAGTCGTCA
CCAGTGGAACCTGGAAC -3' (SEQ ID NO:42)

FlagFc-F2: 5'- GACTACAAGG ACGACGATGA CAAGGACAAA ACTCACACAT
GCCCACCGTG CCCAGCTCCG GAACTCC -3' (SEQ ID NO:43)

FlagFc-R2: 5'- TAGTGGATCCTCATTTACCCG -3' (SEQ ID NO:44)

[0228] The pcDNA 3.1-Fc template was then added to two separate PCR reactions containing 50 pmol each of the primer pairs FlagFc-F1/R1 or FlagFc-

F2/R2 in a 50 µl reaction using Pfu Ultra DNA polymerase (Stratagene, CA) according to manufacturer's standard protocol in a MJ Thermocycler using the following cycles: 95°C 2 minutes; 30 cycles of (95°C 30 seconds, 52°C 30 seconds, 72°C 45 seconds), followed by 72°C for 10 minutes. The products of these two reactions were then mixed in another PCR reaction (2 µl each) with 50 pmol of FlagFc-F1 and FlagFc-R2 primers in a 50 µl reaction using Pfu Ultra DNA polymerase (Stratagene, CA) according to manufacturer's standard protocol in a MJ Thermocycler using the following cycles: 95°C 2 minutes; 30 cycles of (95°C 30 seconds, 52°C 30 seconds, 72°C 45 seconds), followed by 72°C for 10 minutes. The resulting fragment was gel purified, digested and inserted into the pcDNA 3.1-Fc plasmid NheI-Bam HI. The resulting plasmid contains the mouse Igk signal sequence producing the FlagFc protein.

Example 3: Cloning of -Factor VII-Fc construct

[0229] The coding sequence for Factor VII, was obtained by RT-PCR from human fetal liver RNA (Clontech, Palo Alto, CA). The cloned region is comprised of the cDNA sequence from bp 36 to bp 1430 terminating just before the stop codon. A SbfI site was introduced on the N-terminus. A BspEI site was introduced on the C-terminus. The construct was cloned by PCR using the primers:

Downstream: 5' GCTACCTGCAGGCCACCATGGTCTCCCAGGCCCTCAGG
3'(SEQ ID NO:45)

Upstream : 5' CAGTTCCGGAGCTGGGCACGGCGGGCACGTGTGAGTTT
TGTCGGGAAAT GG 3' (SEQ ID NO:46)

and the following conditions: 95°C for 5 minutes followed by 30 cycles of 95°C for 30 seconds, 55°C for 30 seconds, 72°C for 1 minute and 45 seconds, and a final extension cycle of 72°C for 10 minutes.

[0230] The fragment was digested SbfI - BspE I and inserted into pED.dC-Fc a plasmid encoding for the Fc fragment of an IgG1.

Example 4: Cloning of Factor IX-Fc construct

[0231] The human Factor IX coding sequence, including the prepropeptide sequence, was obtained by RT-PCR amplification from adult human liver RNA using the following primers:

natFIX-F: 5'-TTACTGCAGAAGGTTATGCAGCGCGTGAACATG- 3'(SEQ ID NO:47)

F9-R: 5'-TTTTTCGAATTCAGTGAGCTTTGTTTTTTCCTTAATCC- 3'(SEQ ID NO:48)

[0232] 20 ng of adult human liver RNA (Clontech, Palo Alto, CA) and 25 pmol each primer were added to a RT-PCR reaction using the SuperScript.™ One-Step RT-PCR with PLATINUM® Taq system (Invitrogen, Carlsbad, CA) according to manufacturers protocol. Reaction was carried out in a MJ Thermocycler using the following cycles: 50°C 30 minutes; 94°C 2 minutes; 35 cycles of (94°C 30 seconds, 58°C 30 seconds, 72°C 1 minute), and a final 72°C 10 minutes. The fragment was gel purified using Qiagen Gel Extraction Kit (Qiagen, Valencia, CA), and digested with PstI-EcoRI, gel purified, and cloned into the corresponding digest of the pED.dC.XFc plasmid.

Example 5: Cloning of PACE construct

[0233] The coding sequence for human PACE (paired basic amino acid cleaving enzyme), an endoprotease, was obtained by RT-PCR. The following primers were used:

PACE-F1: 5'- GGTAAGCTTGCCATGGAGCTGAGGCCCTGGTTGC -3'(SEQ ID NO:49)

PACE-R1: 5'- GTTTTCAATCTCTAGGACCCACTCGCC -3'(SEQ ID NO:50)

PACE-F2: 5'- GCCAGGCCACATGACTACTCCGC -3'(SEQ ID NO:51)

PACE-R2: 5'- GGTGAATTCTCACTCAGGCAGGTGTGAGGGCAGC -3'(SEQ ID NO:52)

[0234] The PACE-F1 primer adds a HindIII site to the 5' end of the PACE sequence beginning with 3 nucleotides before the start codon, while the PACE-R2 primer adds a stop codon after amino acid 715, which occurs at the end of the extracellular domain of PACE, as well as adding an EcoRI site to the 3' end of the stop codon. The PACE-R1 and -F2 primers anneal on the 3' and 5' sides of an internal BamHI site, respectively. Two RT-PCR reactions were then set up using 25 pmol each of the primer pairs of PACE-F1/R1 or PACE-F2/R2 with 20 ng of adult human liver RNA (Clontech; Palo Alto, CA) in a 50 μ l RT-PCR reaction using the SuperScript.™ One-Step RT-PCR with PLATINUM® Taq system (Invitrogen, Carlsbad, CA) according to manufacturers protocol. The reaction was carried out in a MJ Thermocycler using the following cycles: 50°C 30 minutes; 94°C 2 minutes; 30 cycles of (94°C 30 seconds, 58°C 30 seconds, 72°C 2 minutes), followed by 72°C 10 minutes. These fragments were each ligated into the vector pGEM T-Easy (Promega, Madison, WI) and sequenced fully. The F2-R2 fragment was then subcloned into pcDNA6 V5/His (Invitrogen, Carlsbad, CA) using the BamHI/EcoRI sites, and then the F1-R1 fragment was cloned into this construct using the HindIII/BamHI sites. The final plasmid, pcDNA6-PACE, produces a soluble form of PACE (amino acids 1-715), as the transmembrane region has been deleted. The sequence of PACE in pcDNA6-PACE is essentially as described in Harrison et al. 1998, *Seminars in Hematology* 35:4.

Example 6: Cloning of IFN α -Fc eight amino acid linker construct

[0235] The human interferon α 2b (hIFN α) coding sequence, including the signal sequence, was obtained by PCR from human genomic DNA using the following primers:

IFN α -Sig-F: 5'-GCTACTGCAGCCACCATGGCCTTGACCTTTGCTTTAC-3'(SEQ ID NO:53)

IFN α -EcoR-R: 5'-CGTTGAATTCTTCCTTACTTCTTAACTTTCTTGC-3'(SEQ ID NO:54)

[0236] Genomic DNA was prepared from 373MG human astrocytoma cell line, according to standard methods (Sambrook et al. 1989, *Molecular Cloning: A Laboratory Manual*, 2d ed., Cold Spring Harbor Laboratory Press). Briefly,

approximately 2×10^5 cells were pelleted by centrifugation, resuspended in 100 μ l phosphate buffered saline pH 7.4, then mixed with an equal volume of lysis buffer (100 mM Tris pH 8.0/ 200 mM NaCl / 2% SDS / 5 mM EDTA). Proteinase K was added to a final concentration of 100 μ g/ml, and the sample was digested at 37°C for 4 hours with occasional gentle mixing. The sample was then extracted twice with phenol:chloroform, the DNA precipitated by adding sodium acetate pH 7.0 to 100 mM and an equal volume of isopropanol, and pelleted by centrifugation for 10 min at room temperature. The supernatant was removed and the pellet was washed once with cold 70% ethanol and allowed to air dry before resuspending in TE (10 mM Tris pH 8.0 / 1 mM EDTA).

[0237] 100 ng of this genomic DNA was then used in a 25 μ l PCR reaction with 25 pmol of each primer using Expand High Fidelity System (Boehringer Mannheim, Indianapolis, IN) according to manufacturer's standard protocol in a MJ Thermocycler using the following cycles: 94°C 2 minutes; 30 cycles of (94°C 30 seconds, 50°C 30 seconds, 72°C 45 seconds), and finally 72°C 10 minutes. The expected sized band (~550 bp) was gel purified with a Gel Extraction kit (Qiagen, Valencia, CA), digested with PstI/EcoRI, gel purified again, and cloned into the PstI/EcoRI site of pED.dC.XFc, which contains an 8 amino acid linker (EFAGAAAV) followed by the Fc region of human IgG1.

Example 7: Cloning of IFN α Fc Δ linker construct

[0238] 1 μ g of purified pED.dC.native human IFN α Fc DNA, from Example 6, was then used as a template in a 25 μ l PCR reaction with 25 pmol of each primer IFNa-Sig-F and the following primer:

hIFNaNoLinkFc-R: 5'CAGTTCCGGAGCTGGGCACGGCGGG

CACGTGTGAGTTTTGTCTTCCTTACTTCTTAACTTTTTGCAAGTTTG- 3'(SEQ ID NO:55)

[0239] The PCR reaction was carried out using Expand High Fidelity System (Boehringer Mannheim, Indianapolis, IN) according to the manufacturer's standard protocol in a RapidCycler thermocycler (Idaho Technology, Salt Lake City, UT), denaturing at 94°C for 2 minutes followed by 18 cycles of 95°C for 15 seconds, 55°C for 0 seconds, and 72°C for 1 minute with a slope of 6, followed by

72°C extension for 10 minutes. A PCR product of the correct size (~525 bp) was gel purified using a Gel Extraction kit (Qiagen; Valencia, CA), digested with the PstI and BspEI restriction enzymes, gel purified, and subcloned into the corresponding sites of a modified pED.dC.XFc, where amino acids 231-233 of the Fc region were altered using the degeneracy of the genetic code to incorporate a BspEI site while maintaining the wild type amino acid sequence.

Example 8: Cloning of IFN α Fc GS15 linker construct

[0240] A new backbone vector was created using the Fc found in the Δ linker construct (containing BspEI and RsrII sites in the 5' end using the degeneracy of the genetic code to maintain the amino acid sequence), using this DNA as a template for a PCR reaction with the following primers:

5' B2xGGGGS: 5' gtcaggatccggcgggtggagggagcgacaaaactcacacgtgccc
3'(SEQ ID NO:56)

3' GGGGS: 5' tgacgcggccgctcattacccggagacaggg 3'(SEQ ID NO:57)

[0241] A PCR reaction was carried out with 25 pmol of each primer using Pfu Turbo enzyme (Stratagene, La Jolla, CA) according to manufacturer's standard protocol in a MJ Thermocycler using the following method: 95°C 2 minutes; 30 cycles of (95°C 30 seconds, 54°C 30 seconds, 72°C 2 minutes), 72°C 10 minutes. The expected sized band (~730 bp) was gel purified with a Gel Extraction kit (Qiagen, Valencia CA), digested BamHI/NotI; gel purified again, and cloned into the BamHI/NotI digested vector of pcDNA6 ID, a version of pcDNA6 with the IRES sequence and dhfr gene inserted into NotI/XbaI site.

[0242] 500 ng of purified pED.dC.native human IFN α Fc DNA was then used as a template in a 25 μ l PCR reaction with the following primers:

5' IFN α for GGGGS: 5' ccgctagcctgcaggccaccatggccttgacc 3'(SEQ ID NO:58)

3' IFN α for GGGGS: 5' ccggatccgccgccaccttcttactacgtaaac 3'(SEQ ID NO:59)

[0243] A PCR reaction was carried out with 25 pmol of each primer using Expand High Fidelity System (Boehringer Mannheim, Indianapolis, IN) according to manufacturer's standard protocol in a MJ Thermocycler using the following cycles: 95°C 2 minutes; 14 cycles of (94°C 30 seconds, 48°C 30 seconds, 72°C 1 minute), 72°C 10 minutes. The expected sized band (~600 bp) was gel purified with a Gel Extraction kit (Qiagen, Valencia CA), digested NheI/BamHI, gel purified again, and cloned into the NheI/BamHI site of the pcDNA6 ID/Fc vector, above, to create an IFN α Fc fusion with a 10 amino acid Gly/Ser linker (2xGGGGS), pcDNA6 ID/IFN α -GS10-Fc.

[0244] A PCR reaction was then performed using 500 ng of this pcDNA6 ID/IFN α -GS10-Fc with the following primers

5' B3XGGGGS:5'(SEQ ID NO:60)

gtcaggatccggtggaggcggtccggcggtggaggagcgacaaaactcacacgtgccc
3'(SEQ ID NO:61)

fcclv-R: 5' atagaagcctttgaccaggc 3'(SEQ ID NO:62)

[0245] A PCR reaction was carried out with 25 pmol of each primer using Expand High Fidelity System (Boehringer Mannheim, Indianapolis, IN) according to manufacturer's standard protocol in a MJ Thermocycler using the following cycles: 95°C 2 minutes; 14 cycles of (94°C 30 seconds, 48°C 30 seconds, 72°C 1 minute), 72°C 10 minutes. The expected sized band (504 bp) was gel purified with a Gel Extraction kit (Qiagen, Valencia CA), digested BamHI/BspEI, the 68 bp band was gel purified, and cloned into the BamHI/BspEI site of the pcDNA6 ID/IFN α -GS10-Fc vector, above, to create an IFN α Fc fusion with a 15 amino acid Gly/Ser linker (3xGGGGS), pcDNA6 ID/IFN α -GS15-Fc.

Example 9: Cloning of a Basic Peptide Construct

[0246] The hinge region of the human IgG1 Fc fragment from amino acid 221-229 (EU numbering) was replaced with a basic peptide (CCB). Four overlapping oligos were used (IDT, Coralville, IA):

1. CCB-Fc Sense 1:

5' GCC GGC GAA TTC GGT GGT GAG TAC CAG GCC CTG AAG AAG AAG
GTG GCC CAG CTG AAG GCC AAG AAC CAG GCC CTG AAG AAG AAG
3'(SEQ ID NO:63)

2. CCB-Fc Sense 2:

5' GTG GCC CAG CTG AAG CAC AAG GGC GGC GGC CCC GCC CCA GAG
CTC CTG GGC GGA CCG A 3'(SEQ ID NO:64)

3. CCB-Fc Anti-Sense 1:

5' CGG TCC GCC CAG GAG CTC TGG GGC GGG GCC GCC GCC CTT GTG
CTT CAG CTG GGC CAC CTT CTT CTT CAG GGC CTG GTT CTT G 3'(SEQ ID
NO:65)

4. CCB-Fc Anti-Sense 2:

5' GCC TTC AGC TGG GCC ACC TTC TTC TTC AGG GCC TGG TAC TCA CCA
CCG AAT TCG CCG GCA 3'(SEQ ID NO:66)

[0247] The oligos were reconstituted to a concentration of 50 μ M with dH₂O. 5 μ l of each oligo were annealed to each other by combining in a thin walled PCR tube with 2.2 μ l of restriction buffer #2 (*i.e.* final concentration of 10 mM Tris HCl pH 7.9, 10 mM MgCl₂, 50 mM Na Cl, 1 mM dithiothreitol) (New England Biolabs, Beverly, MA) and heated to 95°C for 30 seconds and then allowed to anneal by cooling slowly for 2 hours to 25°C. 5 pmol of the now annealed oligos were ligated into a pGEM T-Easy vector as directed in the kit manual. (Promega, Madison WI). The ligation mixture was added to 50 μ l of DH5 α competent *E. coli* cells (Invitrogen, Carlsbad, CA) on ice for 2 minutes, incubated at 37°C for 5 minutes, incubated on ice for 2 minutes, and then plated on LB+100 μ g/L ampicillin agar plates and placed at 37°C for 14 hours. Individual

bacterial colonies were picked and placed in 5 ml of LB+100 µg/L ampicillin and allowed to grow for 14 hours. The tubes were spun down at 2000xg, 4°C for 15 minutes and the vector DNA was isolated using Qiagen miniprep kit (Qiagen, Valencia, CA) as indicated in the kit manual. 2 µg of DNA was digested with NgoM IV-Rsr-II. The fragment was gel purified by the Qiaquick method as instructed in the kit manual (Qiagen, Valencia, CA) and ligated to pED.dcEpoFc with NgoM IV/Rsr II. The ligation was transformed into DH5α competent *E. coli* cells and the DNA prepared as described for the pGEM T-Easy vector.

Example 10: Cloning of the erythropoietin-acidic peptide Fc construct

[0248] The hinge region of the human IgG1 Fc fragment in EPO-Fc from amino acid 221-229 (EU numbering) was replaced with an acidic peptide (CCA). Four overlapping oligos were used (IDT, Coralville, IA):

1. Epo-CCA-Fc Sense 1:

5' CCG GTG ACA GGG AAT TCG GTG GTG AGT ACC AGG CCC TGG AGA
AGG AGG TGG CCC AGC TGG AG 3'(SEQ ID NO:67)

2. Epo-CCA-Fc Sense 2:

5' GCC GAG AAC CAG GCC CTG GAG AAG GAG GTG GCC CAG CTG GAG
CAC GAG GGT GGT GGT CCC GCT CCA GAG CTG CTG GGC GGA CA
3'(SEQ ID NO:68)

3. Epo-CCA-Fc Anti-Sense 1:

5' GTC CGC CCA GCA GCT CTG GAG CGG GAC CAC CAC CCT CGT GCT
CCA GCT GGG CCA C 3'(SEQ ID NO:69)

4. Epo-CCA-Fc Anti-Sense 2:

5' CTC CTT CTC CAG GGC CTG GTT CTC GGC CTC CAG CTG GGC CAC
 CTC CTT CTC CAG GGC CTG GTA CTC ACC ACC GAA TTC CCT GTC ACC
 GGA 3'(SEQ ID NO:70)

[0249] The oligos were reconstituted to a concentration of 50 μ M with dH₂O. 5 μ l of each oligo were annealed to each other by combining in a thin walled PCR tube with 2.2 μ l of restriction buffer No. 2 (New England Biolabs, Beverly, MA) and heated to 95°C for 30 seconds and then allowed to cool slowly for 2 hours to 25°C. 5 pmol of the now annealed oligos were ligated into a pGEM T-Easy vector as directed in the kit manual. (Promega, Madison, WI). The ligation mixture was added to 50 μ l of DH5 α competent *E. coli* cells (Invitrogen, Carlsbad, CA) on ice for 2 minutes, incubated at 37°C 5 minutes, incubated on ice for 2 minutes, and then plated on LB+100 μ g/L ampicillin agar plates and placed at 37°C for 14 hours. Individual bacterial colonies were picked and placed in 5 ml of LB+100 μ g/L ampicillin and allowed to grow for 14 hours. The tubes were spun down at 2000xg, 4°C for 15 minutes and the vector DNA was prepared using Qiagen miniprep kit (Qiagen, Valencia, CA) as indicated in the kit manual. 2 μ g of DNA was digested with Age I-Rsr-II. The fragment was gel purified by the Qiaquick method as instructed in the kit manual (Qiagen, Valencia, CA) and ligated into pED.Epo Fc.1 Age I-Rsr II. The ligation was transformed into DH5 α competent *E. coli* cells and DNA prepped as described above.

Example 11: Cloning of Cys-Fc construct

[0250] Using PCR and standard molecular biology techniques (Sambrook et al. 1989, *Molecular Cloning: A Laboratory Manual*, 2nd ed., Cold Spring Harbor Laboratory Press), a mammalian expression construct was generated such that the coding sequence for the human IFN α signal peptide was directly abutted against the coding sequence of Fc beginning at the first cysteine residue (Cys 226, EU Numbering). Upon signal peptidase cleavage and secretion from mammalian cells, an Fc protein with an N-terminal cysteine residue was thus generated. Briefly, the primers

IFNa-Sig-F (IFNa-Sig-F: 5'-GCTACTGCAGCCACCATGGCCTTGACCTT
 TGCTTTAC-3')(SEQ ID NO:71) and Cys-Fc-R
 (5'-CAGTTCCGGAGCTGGGACACGGCGGA

GAGCCACAGAGCAGCTTG-3') (SEQ ID NO:72) were used in a PCR reaction to create a fragment linking the IFN α signal sequence with the N terminus of Fc, beginning with Cys 226. 500 ng of pED.dC.native hIFN α Δ linker was added to 25 pmol of each primer in a PCR reaction with Expand High Fidelity System (Boehringer Mannheim, Indianapolis, IN) according to manufacturer's standard protocol. The reaction was carried out in a MJ Thermocycler using the following cycles: 94°C 2 minutes; 30 cycles of (94°C 30 seconds, 50°C 30 seconds, 72°C 45 seconds), and finally 72°C 10 minutes. The expected sized band (~112 bp) was gel purified with a Gel Extraction kit (Qiagen, Valencia CA), digested with the PstI and BspEI restriction enzymes, gel purified, and subcloned into the corresponding sites pED.dC.native hIFN α Δ linker to generate pED.dC.Cys-Fc (Figure 5).

Example 12: Protein Expression and Preparation of Fc-MESNA

[0251] The coding sequence for Fc (the constant region of human IgG1) was obtained by PCR amplification from an Fc-containing plasmid using standard conditions and reagents, following the manufacturer's recommended procedure to subclone the Fc coding sequence *NdeI/SapI*. Briefly, the primers 5'-GTGGTCATA TGGGCATTGAAGGCAGAGGCGCCGCTGCGGTCTG - 3' (SEQ ID NO:73) and 5' - GGTGGTTGC TCTTCCGCAAAAACCCGGAGACAGGGAGAGACTCTTCTGCG - 3' (SEQ ID NO:74) were used to amplify the Fc sequence from 500 ng of the plasmid pED.dC.Epo-Fc using Expand High Fidelity System (Boehringer Mannheim, Basel Switzerland) in a RapidCycler thermocycler (Idaho Technology Salt Lake City, Utah), denaturing at 95°C for 2 minutes followed by 18 cycles of 95°C for 0 sec, 55°C for 0 sec, and 72°C for 1 minute with a slope of 4, followed by 72°C extension for 10 minutes. The PCR product was subcloned into an intermediate cloning vector and sequenced fully, and then subcloned using the *NdeI* and *SapI* sites in the pTWIN1 vector following standard procedures. Sambrook, J., Fritsch, E.F. and Maniatis, T. 1989, *Molecular Cloning: A Laboratory Manual*, 2nd ed.; Cold Spring Harbor, New York: Cold Spring Harbor Laboratory Press. This plasmid was then transformed into BL21(DE3) pLysS cells using standard methods. Id. A 1 liter culture of cells was grown to an absorbance reading of 0.8 AU at 37°C, induced with 1 mM isopropyl beta-D-1-thiogalactopyranoside, and grown

overnight at 25°C. Cells were pelleted by centrifugation, lysed in 20 mM Tris 8.8/1% NP40/0.1 mM phenylmethanesulfonyl fluoride/ 1 µg/ml Benzonase (Novagen Madison, WI), and bound to chitin beads (New England Biolabs; Beverly, MA) overnight at 4°C. Beads were then washed with several column volumes of 20 mM Tris 8.5/ 500 mM NaCl/ 1 mM EDTA, and then stored at -80°C. Purified Fc-MESNA was generated by eluting the protein from the beads in 20 mM Tris 8.5/ 500 mM NaCl / 1 mM EDTA / 500 mM 2-mercapto ethane sulfonic acid (MESNA), and the eluate was used directly in the coupling reaction, below.

Example 13: Factor VII-Fc monomer-dimer hybrid expression and purification

[0252] CHO DG-44 cells expressing Factor VII-Fc were established. CHO DG-44 cells were grown at 37°C, 5% CO₂, in MEM Alpha plus nucleoside and ribonucleosides and supplemented with 5% heat-inactivated fetal bovine serum until transfection.

[0253] DG44 cells were plated in 100 mm tissue culture petri dishes and grown to a confluency of 50%- 60%. A total of 10 µg of DNA was used to transfect one 100 mm dish: 7.5 µg of pED.dC.FVII-Fc + 1.5 µg pcDNA3/Flag-Fc + 1 µg of pcDNA6-PACE. The cells were transfected as described in the Superfect transfection reagent manual (Qiagen, Valencia, CA). The media was removed from transfection after 48 hours and replaced with MEM Alpha without nucleosides plus 5% dialyzed fetal bovine serum and 10 µg/ml of Blasticidin (Invitrogen, Carlsbad, CA) and 0.2 mg/ml geneticin (Invitrogen, Carlsbad, CA). After 10 days, the cells were released from the plate with 0.25% trypsin and transferred into T25 tissue culture flasks, and the selection was continued for 10-14 days until the cells began to grow well as stable cell lines were established. Protein expression was subsequently amplified by the addition 25 nM methotrexate.

[0254] Approximately 2×10^7 cells were used to inoculate 300 ml of growth medium in a 1700 cm² roller bottle (Corning, Corning, NY) supplemented with 5 µg/ml of vitamin K₃ (menadione sodium bisulfite) (Sigma, St Louis, MO). The roller bottles were incubated in a 5% CO₂ at 37°C for 72 hours. Then the growth medium was exchanged with 300 ml serum-free production medium (DMEM/F12 with 5 µg/ml bovine insulin and 10 µg/ml Gentamicin) supplemented

with 5 µg/L of vitamin K₃. The production medium (conditioned medium) was collected every day for 10 days and stored at 4°C. Fresh production medium was added to the roller bottles after each collection and the bottles were returned to the incubator. Pooled media was first clarified using a Sartoclean glass fiber filter (3.0 µm + 0.2 µm) (Sartorius Corp. Gottingen, Germany) followed by an Acropack 500 filter (0.8 µm + 0.2 µm) (Pall Corp., East Hills, NY). The clarified media was then concentrated approximately 20-fold using Pellicon Biomax tangential flow filtration cassettes (10 kDa MWCO) (Millipore Corp., Billerica, MA).

[0255] Fc chimeras were then captured from the concentrated media by passage over a Protein A Sepharose 4 Fast Flow Column (AP Biotech, Piscataway, NJ). A 5 x 5 cm (100 ml) column was loaded with ≤5 mg Fc protein per ml column volume at a linear flow rate of 100 cm/hour to achieve a residence time of ≥ 3 minutes. The column was then washed with >5 column volumes of 1X DPBS to remove non-specifically bound proteins. The bound proteins were eluted with 100 mM Glycine pH 3.0. Elution fractions containing the protein peak were then neutralized by adding 1 part 1 M Tris-HCL, pH 8 to 10 parts elute fraction.

[0256] To remove FLAG-Fc homodimers (that is, chimeric Fc dimers with FLAG peptide expressed as fusions with both Fc molecules) from the preparation, the Protein A Sepharose 4 Fast Flow pool was passed over a Unosphere S cation-exchange column (BioRad Corp., Richmond, CA). Under the operating conditions for the column, the FLAG-Fc monomer-dimer hybrid is uncharged (FLAG-Fc theoretical pI=6.19) and flows through the column while the hFVII-Fc constructs are positively charged, and thus bind to the column and elute at higher ionic strength. The Protein A Sepharose 4 Fast Flow pool was first dialyzed into 20 mM MES, 20 mM NaCl, pH 6.1. The dialyzed material was then loaded onto a 1.1 x 11 cm (9.9 ml) column at 150 cm/hour. During the wash and elution, the flow rate was increased to 500 cm/hour. The column was washed sequentially with 8 column volumes of 20 mM MES, 20 mM NaCl, pH 6.1 and 8 column volumes of 20 mM MES, 40 mM NaCl, pH 6.1. The bound protein was eluted with 20 mM MES, 750 mM NaCl, pH 6.1. Elution fractions containing the protein peak were pooled and sterile filtered through a 0.2 µm filter disc prior to storage at -80°C.

[0257] An anti-FLAG MAB affinity column was used to separate chimeric Fc dimers with hFVII fused to both Fc molecules from those with one FLAG peptide and one hFVII fusion. The Unosphere S Eluate pool was diluted 1:1 with 20 mM Tris, 50 mM NaCl, 5 mM CaCl₂, pH 8 and loaded onto a 1.6 x 5 cm M2 anti-FLAG sepharose column (Sigma Corp., St. Louis, MO) at a linear flow rate of 60 cm/hour. Loading was targeted to < 2.5 mg monomer-dimer hybrid /ml column volume. After loading the column was washed with 5 column volumes 20 mM Tris, 50 mM NaCl, 5 mM CaCl₂, pH 8.0, monomer-dimer hybrids were then eluted with 100 mM Glycine, pH 3.0. Elution fractions containing the protein peak were then neutralized by adding 1 part 1 M Tris-HCl, pH 8 to 10 parts eluate fraction. Pools were stored at -80°C.

Example 14: Factor IX-Fc homodimer and monomer-dimer hybrid expression and purification

[0258] CHO DG-44 cells expressing Factor IX-Fc were established. DG44 cells were plated in 100 mm tissue culture petri dishes and grown to a confluency of 50%- 60%. A total of 10 µg of DNA was used to transfect one 100 mm dish: for the homodimer transfection, 8 µg of pED.dC.Factor IX-Fc + 2 µg of pcDNA6-PACE was used; for the monomer-dimer hybrid transfection, 8 µg of pED.dC.Factor IX-Fc + 1 µg of pcDNA3-FlagFc +1 µg pcDNA6-PACE was used. The cells were transfected as described in the Superfect transfection reagent manual (Qiagen, Valencia, CA). The media was removed from transfection after 48 hours and replaced with MEM Alpha without nucleosides plus 5% dialyzed fetal bovine serum and 10 µg/ml of Blasticidin (Invitrogen, Carlsbad, CA) for both transfections, while the monomer-dimer hybrid transfection was also supplemented with 0.2 mg/ml geneticin (Invitrogen, Carlsbad, CA). After 3 days, the cells were released from the plate with 0.25% trypsin and transferred into T25 tissue culture flasks, and the selection was continued for 10-14 days until the cells began to grow well as stable cell lines were established. Protein expression was subsequently amplified by the addition 10 nM or 100 nM methotrexate for the homodimer or monomer-dimer hybrid, respectively.

[0259] For both cell lines, approximately 2×10^7 cells were used to inoculate 300 ml of growth medium in a 1700 cm² roller bottle (Corning, Corning,

NY), supplemented with 5 µg/L of vitamin K₃ (menadione sodium bisulfite) (Sigma, St. Louis, MO). The roller bottles were incubated in a 5% CO₂ at 37°C for approximately 72 hours. The growth medium was exchanged with 300 ml serum-free production medium (DMEM/F12 with 5 µg/ml bovine insulin and 10 µg/ml Gentamicin), supplemented with 5 µg/L of vitamin K₃. The production medium (conditioned medium) was collected everyday for 10 days and stored at 4°C. Fresh production medium was added to the roller bottles after each collection and the bottles were returned to the incubator. Prior to chromatography, the medium was clarified using a SuporCap-100 (0.8/0.2 µm) filter (Pall Gelman Sciences, Ann Arbor, MI). All of the following steps were performed at 4°C. The clarified medium was applied to Protein A Sepharose, washed with 5 column volumes of 1X PBS (10 mM phosphate, pH 7.4, 2.7 mM KCl, and 137 mM NaCl), eluted with 0.1 M glycine, pH 2.7, and then neutralized with 1/10 volume of 1 M Tris-HCl, pH 9.0. The protein was then dialyzed into PBS.

[0260] The monomer-dimer hybrid transfection protein sample was subject to further purification, as it contained a mixture of FIX-Fc:FIX-Fc homodimer, FIX-Fc:Flag-Fc monomer-dimer hybrid, and Flag-Fc:Flag-Fc homodimer. Material was concentrated and applied to a 2.6 cm x 60 cm (318 ml) Superdex 200 Prep Grade column at a flow rate of 4 ml/minute (36 cm/hour) and then eluted with 3 column volumes of 1X PBS. Fractions corresponding to two peaks on the UV detector were collected and analyzed by SDS-PAGE. Fractions from the first peak contained either FIX-Fc:FIX-Fc homodimer or FIX-Fc:FlagFc monomer-dimer hybrid, while the second peak contained FlagFc:FlagFc homodimer. All fractions containing the monomer-dimer hybrid but no FlagFc homodimer were pooled and applied directly to a 1.6 x 5 cm M2 anti-FLAG sepharose column (Sigma Corp., St. Louis, MO) at a linear flow rate of 60 cm/hour. After loading, the column was washed with 5 column volumes PBS. Monomer-dimer hybrids were then eluted with 100 mM Glycine, pH 3.0. Elution fractions containing the protein peak were then neutralized by adding 1/10 volume of 1 M Tris-HCl, and analyzed by reducing and nonreducing SDS-PAGE. Fractions were dialyzed into PBS, concentrated to 1-5 mg/ml, and stored at -80°C.

Example 15: IFN α homodimer and monomer-dimer hybrid expression and purification

[0261] CHO DG-44 cells expressing hIFN α were established. DG44 cells were plated in 100 mm tissue culture petri dishes and grown to a confluency of 50%-60%. A total of 10 μ g of DNA was used to transfect one 100 mm dish: for the homodimer transfection, 10 μ g of the hIFN α Fc constructs; for the monomer-dimer hybrid transfection, 8 μ g of the hIFN α Fc constructs + 2 μ g of pcDNA3-FlagFc. The cells were transfected as described in the Superfect transfection reagent manual (Qiagen, Valencia, CA). The media was removed from transfection after 48 hours and replaced with MEM Alpha without nucleosides plus 5% dialyzed fetal bovine serum, while the monomer-dimer hybrid transfection was also supplemented with 0.2 mg/ml geneticin (Invitrogen, Carlsbad, CA). After 3 days, the cells were released from the plate with 0.25% trypsin and transferred into T25 tissue culture flasks, and the selection was continued for 10-14 days until the cells began to grow well and stable cell lines were established. Protein expression was subsequently amplified by the addition methotrexate: ranging from 10 to 50 nM.

[0262] For all cell lines, approximately 2×10^7 cells were used to inoculate 300 ml of growth medium in a 1700 cm² roller bottle (Corning, Corning, NY). The roller bottles were incubated in a 5% CO₂ at 37°C for approximately 72 hours. Then the growth medium was exchanged with 300 ml serum-free production medium (DMEM/F12 with 5 μ g/ml bovine insulin and 10 μ g/ml Gentamicin). The production medium (conditioned medium) was collected every day for 10 days and stored at 4°C. Fresh production medium was added to the roller bottles after each collection and the bottles were returned to the incubator. Prior to chromatography, the medium was clarified using a SuporCap-100 (0.8/0.2 μ m) filter from Pall Gelman Sciences (Ann Arbor, MI). All of the following steps were performed at 4°C. The clarified medium was applied to Protein A Sepharose, washed with 5 column volumes of 1X PBS (10 mM phosphate, pH 7.4, 2.7 mM KCl, and 137 mM NaCl), eluted with 0.1 M glycine, pH 2.7, and then neutralized with 1/10 volume of 1 M Tris-HCl, pH 9.0. The protein was then dialyzed into PBS.

[0263] The monomer-dimer hybrid transfection protein samples were then subject to further purification, as it contained a mixture of IFN α Fc:IFN α Fc homodimer, IFN α Fc:FlagFc monomer-dimer hybrid, and FlagFc:FlagFc homodimer (or Δ linker or GS15 linker). Material was concentrated and applied to a 2.6 cm x 60 cm (318 ml) Superdex 200 Prep Grade column at a flow rate of 4 ml/min (36 cm/hr) and then eluted with 3 column volumes of 1X PBS. Fractions corresponding to two peaks on the UV detector were collected and analyzed by SDS-PAGE. Fractions from the first peak contained either IFN α Fc:IFN α Fc homodimer or IFN α Fc:FlagFc monomer-dimer hybrid, while the second peak contained FlagFc:FlagFc homodimer. All fractions containing the monomer-dimer hybrid, but no FlagFc homodimer, were pooled and applied directly to a 1.6 x 5 cm M2 anti-FLAG sepharose column (Sigma Corp., St. Louis, MO) at a linear flow rate of 60 cm/hour. After loading the column was washed with 5 column volumes PBS monomer-dimer hybrids were then eluted with 100 mM Glycine, pH 3.0. Elution fractions containing the protein peak were then neutralized by adding 1/10 volume of 1 M Tris-HCl, and analyzed by reducing and nonreducing SDS-PAGE. Fractions were dialyzed into PBS, concentrated to 1-5 mg/ml, and stored at -80°C.

Example 16: Coiled coil protein expression and purification

[0264] The plasmids, pED.dC Epo-CCA-Fc and pED.dC CCB-Fc will be transfected either alone or together at a 1:1 ratio into CHO DG44 cells. The cells will be transfected as described in the Superfect transfection reagent manual (Qiagen, Valencia, CA). The media will be removed after 48 hours and replaced with MEM Alpha w/o nucleosides plus 5% dialyzed fetal bovine serum. Purification will be done by affinity chromatography over a protein A column according to methods known in the art. Alternatively, purification can be achieved using size exclusion chromatography.

Example 17: Cys-Fc expression and purification

[0265] CHO DG-44 cells expressing Cys-Fc were established. The pED.dC.Cys-Fc expression plasmid, which contains the mouse dihydrofolate reductase (dhfr) gene, was transfected into CHO DG44 (dhfr deficient) cells using Superfect reagent (Qiagen; Valencia, CA) according to manufacturer's protocol, followed by selection for stable transfectants in α MEM (without nucleosides)

tissue culture media supplemented with 5% dialyzed FBS and penicillin/streptomycin antibiotics (Invitrogen; Carlsbad, CA) for 10 days. The resulting pool of stably transfected cells were then amplified with 50 nM methotrexate to increase expression. Approximately 2×10^7 cells were used to inoculate 300 ml of growth medium in a 1700 cm² roller bottle (Corning, Corning, NY). The roller bottles were incubated in a 5% CO₂ at 37°C for approximately 72 hours. The growth medium was exchanged with 300 ml serum-free production medium (DMEM/F12 with 5 µg/ml bovine insulin and 10 µg/ml Gentamicin). The production medium (conditioned medium) was collected every day for 10 days and stored at 4°C. Fresh production medium was added to the roller bottles after each collection and the bottles were returned to the incubator. Prior to chromatography, the medium was clarified using a SuporCap-100 (0.8/0.2 µm) filter from Pall Gelman Sciences (Ann Arbor, MI). All of the following steps were performed at 4°C. The clarified medium was applied to Protein A Sepharose, washed with 5 column volumes of 1X PBS (10 mM phosphate, pH 7.4, 2.7 mM KCl, and 137 mM NaCl), eluted with 0.1 M glycine, pH 2.7, and then neutralized with 1/10 volume of 1 M Tris-HCl, pH 9.0. Protein was dialyzed into PBS and used directly in conjugation reactions.

Example 18: Coupling of T20-thioesters to Cys-Fc

[0266] Cys-Fc (4 mg, 3.2 mg/ml final concentration) and either T20-thioester or T20-PEG-thioester (2 mg, approximately 5 molar equivalents) were incubated for 16 hours at room temperature in 0.1 M Tris 8/ 10 mM MESNA. Analysis by SDS-PAGE (Tris-Gly gel) using reducing sample buffer indicated the presence of a new band approximately 5 kDa larger than the Fc control (>40-50% conversion to the conjugate). Previous N-terminal sequencing of Cys-Fc and unreacted Cys-Fc indicated that the signal peptide is incorrectly processed in a fraction of the molecules, leaving a mixture of (Cys)-Fc, which will react through native ligation with peptide-thioesters, and (Val)-(Gly)-(Cys)-Fc, which will not. As the reaction conditions are insufficient to disrupt the dimerization of the Cys-Fc molecules, this reaction generated a mixture of T20-Cys-Fc:T20-Cys-Fc homodimers, T20-Cys-Fc: Fc monomer-dimer hybrids, and Cys-Fc:Cys-Fc Fc-dimers. This protein was purified using size exclusion chromatography as

indicated above to separate the three species. The result was confirmed by SDS-PAGE analysis under nonreducing conditions.

Example 19: Antiviral assay for IFN α activity

[0267] Antiviral activity (IU/ml) of IFN α fusion proteins was determined using a CPE (cytopathic effect) assay. A549 cells were plated in a 96 well tissue culture plate in growth media (RPMI 1640 supplemented with 10% fetal bovine serum (FBS) and 2 mM L-glutamine) for 2 hours at 37°C, 5% CO₂. IFN α standards and IFN α fusion proteins were diluted in growth media and added to cells in triplicate for 20 hours at 37°C, 5% CO₂. Following incubation, all media was removed from wells, encephalomyocarditis virus (EMC) virus was diluted in growth media and added (3000 pfu/well) to each well with the exception of control wells. Plates were incubated at 37°C, 5% CO₂ for 28 hours. Living cells were fixed with 10% cold trichloroacetic acid (TCA) and then stained with Sulforhodamine B (SRB) according to published protocols (Rubinstein et al. 1990, *J. Natl. Cancer Inst.* 82, 1113). The SRB dye was solubilized with 10 mM Tris pH 10.5 and read on a spectrophotometer at 490 nm. Samples were analyzed by comparing activities to a known standard curve World Health Organization IFN α 2b International Standard ranging from 5 to 0.011 IU/ml. The results are presented below in Table 3 and Figure 6 and demonstrate increased antiviral activity of monomer-dimer hybrids.

TABLE 3: INTERFERON ANTIVIRAL ASSAY
HOMODIMER V. MONOMER-DIMER HYBRID

Protein	Antiviral Activity (IU/nmol)	Std dev
IFN α Fc 8aa linker homodimer	0.45×10^5	0.29×10^5
IFN α Fc 8aa linker:FlagFc monomer-dimer hybrid	4.5×10^5	1.2×10^5
IFN α Fc Δ linker homodimer	0.22×10^5	0.07×10^5
IFN α Fc Δ delta linker: FlagFc monomer-dimer hybrid	2.4×10^5	0.0005×10^5
IFN α Fc GS15 linker homodimer	2.3×10^5	1.0×10^5
IFN α Fc GS15 linker monomer-dimer hybrid	5.3×10^5	0.15×10^5

Example 20: FVIIa Clotting Activity Analysis

[0268] The StaClot FVIIa-rTF assay kit was purchased from Diagnostica Stago (Parsippany, NJ) and modified as described in Johannessen et al. 2000, *Blood Coagulation and Fibrinolysis* 11:S159. A standard curve was preformed with the FVIIa World Health Organization standard 89/688. The assay was used to compare clotting activity of monomer-dimer hybrids compared to homodimers. The results showed the monomer-dimer hybrid had four times the clotting activity compared to the homodimer (Figure 7).

Example 21: FVIIa-Fc Oral dosing in day 10 rats

[0269] 25 gram day 9 newborn Sprague Dawley rats were purchased from Charles River (Wilmington, MA) and allowed to acclimate for 24 hours. The rats were dosed orally with FVIIaFc homodimer, monomer-dimer hybrid or a 50:50 mix of the two. A volume of 200 μ l of a FVIIaFc solution for a dose of 1 mg/kg was administered. The solution was composed of a Tris-HCl buffer pH 7.4 with 5 mg/ml soybean trypsin inhibitor. The rats were euthanized with CO₂ at several time points, and 200 μ l of blood was drawn by cardiac puncture. Plasma was

obtained by the addition of a 3.8% sodium citrate solution and centrifugation at room temperature at a speed of 1268xg. The plasma samples were either assayed fresh or frozen at 20°C. Orally dosed monomer-dimer hybrid resulted in significantly higher maximum (C_{max}) serum concentrations compared to homodimeric Factor VII (Figure 8).

Example 22: Factor IX-Fc Oral dosing of neonatal rats

[0270] Ten-day old neonatal Sprague-Dawley rats were dosed p.o. with 200 μ l of FIX-Fc homodimer or FIX-Fc: FlagFc monomer-dimer hybrid at approximately equimolar doses of 10 nmol/kg in 0.1 M sodium phosphate buffer, pH 6.5 containing 5 mg/ml soybean trypsin inhibitor and 0.9% NaCl. At 1, 2, 4, 8, 24, 48, and 72 hours post injection, animals were euthanized with CO₂, blood was drawn via cardiac puncture and plasma was obtained by the addition of a 3.8% sodium citrate solution and centrifugation at room temperature at a speed of 1268xg. Samples were then sedimented by centrifugation, serum collected and frozen at -20°C until analysis of the fusion proteins by ELISA.

Example 23: Factor IX-Fc ELISA

[0271] A 96-well Immulon 4HBX ELISA plate (Thermo LabSystems, Vantaa, Finland) was coated with 100 μ l/well of goat anti-Factor IX IgG (Affinity Biologicals, Ancaster, Canada) diluted 1:100 in 50 mM carbonate buffer, pH 9.6. The plates were incubated at ambient temperature for 2 hours or overnight at 4°C sealed with plastic film. The wells were washed 4 times with PBST, 300 μ l/well using the TECAN plate washer. The wells were blocked with PBST + 6% BSA, 200 μ l/well, and incubated 90 minutes at ambient temperature. The wells were washed 4 times with PBST, 300 μ l/well using the TECAN plate washer. Standards and blood samples from rats described in Example 18 were added to the wells, (100 μ l/well), and incubated 90 minutes at ambient temperature. Samples and standards were diluted in HBET buffer (HBET: 5.95 g HEPES, 1.46 g NaCl, 0.93 g Na₂EDTA, 2.5 g Bovine Serum Albumin, 0.25 ml Tween-20, bring up to 250 ml with dH₂O, adjust pH to 7.2). Standard curve range was from 200 ng/ml to 0.78 ng/ml with 2 fold dilutions in between. Wells were washed 4 times with PBST, 300 μ l/well using the TECAN plate washer. 100 μ l/well of conjugated goat anti-human IgG-Fc-HARP antibody (Pierce, Rockford, IL) diluted in HBET

1:25,000 was added to each well. The plates were incubated 90 minutes at ambient temperature. The wells were washed 4 times with PBST, 300 µl/well using the TECAN plate washer. The plates were developed with 100 µl/well of tetramethylbenzidine peroxidase substrate (TMB) (Pierce, Rockford, IL) was added according to the manufacturer's instructions. The plates were incubated 5 minutes at ambient temperature in the dark or until color developed. The reaction was stopped with 100 µl/well of 2 M sulfuric acid. Absorbance was read at 450 nm on SpectraMax plusplate reader (Molecular Devices, Sunnyvale, CA).

Analysis of blood drawn at 4 hours indicated more than a 10 fold difference in serum concentration between Factor IX-Fc monomer-dimer hybrids compared to Factor IX Fc homodimers (Figure 9). The results indicated Factor IX-Fc monomer-dimer hybrid levels were consistently higher than Factor IX-Fc homodimers (Figure 10).

Example 24: Cloning of Epo-Fc

[0272] The mature Epo coding region was obtained by PCR amplification from a plasmid encoding the mature erythropoietin coding sequence, originally obtained by RT-PCR from Hep G2 mRNA, and primers hepoxba-F and hepoeco-R, indicated below. Primer hepoxba-F contains an *Xba*I site, while primer hepoeco-R contains an *Eco*RI site. PCR was carried out in the Idaho Technology RapidCycler using Vent polymerase, denaturing at 95°C for 15 seconds, followed by 28 cycles with a slope of 6.0 of 95°C for 0 seconds, 55°C for 0 seconds, and 72°C for 1 minute 20 seconds, followed by 3 minute extension at 72°C. An approximately 514 bp product was gel purified, digested with *Xba*I and *Eco*RI, gel purified again and directionally subcloned into an *Xba*I/*Eco*RI-digested, gel purified pED.dC.XFc vector, mentioned above. This construct was named pED.dC.EpoFc.

[0273] The Epo sequence, containing both the endogenous signal peptide and the mature sequence, was obtained by PCR amplification using an adult kidney QUICK-clone cDNA preparation as the template and primers Epo+Pep-Sbf-F and Epo+Pep-Sbf-R, described below. The primer Epo+Pep-Sbf-F contains an *Sbf*I site upstream of the start codon, while the primer Epo+Pep-Sbf-R anneals downstream of the endogenous *Sbf*I site in the Epo sequence. The PCR reaction was carried out in the PTC-200 MJ Thermocycler using Expand polymerase,

denaturing at 94°C for 2 minutes, followed by 32 cycles of 94°C for 30 seconds, 57°C for 30 seconds, and 72°C for 45 seconds, followed by a 10 minute extension at 72°C. An approximately 603 bp product was gel isolated and subcloned into the pGEM-T Easy vector. The correct coding sequence was excised by *Sbf*I digestion, gel purified, and cloned into the *Pst*I-digested, shrimp alkaline phosphatase (SAP)-treated, gel purified pED.dC.EpoFc plasmid. The plasmid with the insert in the correct orientation was initially determined by *Kpn*I digestion. A *Xmn*I and *Pvu*II digestion of this construct was compared with pED.dC.EpoFc and confirmed to be in the correct orientation. The sequence was determined and the construct was named pED.dC.natEpoFc.

PCR Primers:

hepoxba-F (EPO-F): 5'-AATCTAGAGCCCCACCGCCTCATCTGTGAC-3'(SEQ ID NO:75)

hepoeco-R (EPO-R) 5'-TTGAATTCTCTGTCCCCTGTCCTGCAGGCC-3'(SEQ ID NO:76)

Epo+Pep-Sbf-F: 5'-GTACCTGCAGGCGGAGATGGGGGTGCA-3'(SEQ ID NO:77)

Epo+Pep-Sbf-R: 5'-CCTGGTCATCTGTCCCCTGTCC-3'(SEQ ID NO:78)

Example 25: Cloning of Epo-Fc

[0274] An alternative method of cloning EPO-Fc is described herein. Primers were first designed to amplify the full length Epo coding sequence, including the native signal sequence, as follows:

Epo-F: 5'-GTCCAACCTG CAGGAAGCTTG CCGCCACCAT GGGAGTGCAC GAATGTCCTG CCTGG- 3'(SEQ ID NO:79)

Epo-R: 5'-GCCGAATTCA GTTTTGTGCA CCGCAGCGG CGCCGGCGAA CTCTCTGTCC CCTGTTCTGC AGGCCTCC- 3'(SEQ ID NO:80)

[0275] The forward primer incorporates an SbfI and HindIII site upstream of a Kozak sequence, while the reverse primer removes the internal SbfI site, and adds an 8 amino acid linker to the 3' end of the coding sequence (EFAGAAAV) (SEQ ID NO:81) as well as SalI and EcoRI restriction sites. The Epo coding sequence was then amplified from a kidney cDNA library (BD Biosciences Clontech, Palo Alto, CA) using 25 pmol of these primers in a 25 µl PCR reaction using Expand High Fidelity System (Boehringer Mannheim, Indianapolis, IN) according to manufacturer's standard protocol in a MJ Thermocycler using the following cycles: 94°C 2 minutes; 30 cycles of (94°C 30 seconds, 58°C 30 seconds, 72°C 45 seconds), followed by 72°C for 10 minutes. The expected sized band (641 bp) was gel purified with a Gel Extraction kit (Qiagen, Valencia, CA) and ligated into the intermediate cloning vector pGEM T-Easy (Promega, Madison, WI). DNA was transformed into DH5α cells (Invitrogen, Carlsbad, CA) and miniprep cultures grown and purified with a Plasmid Miniprep Kit (Qiagen, Valencia, CA) both according to manufacturer's standard protocols. Once the sequence was confirmed, this insert was digested out with SbfI/EcoRI restriction enzymes, gel purified, and cloned into the PstI/EcoRI sites of the mammalian expression vector pED.dC in a similar manner.

[0276] Primers were designed to amplify the coding sequence for the constant region of human IgG1 (the Fc region, EU numbering 221-447) as follows:
Fc-F: 5'-GCTGCGGTCG ACAAACACTCA CACATGCCCA CCGTGCCCAG
CTCCGGAACCT CCTGGGCGGA CCGTCAGTC- 3'(SEQ ID NO:82)

Fc-R 5'-ATTGGAATTC TCATTACCC GGAGACAGGG AGAGGC- 3'(SEQ ID NO:83)

The forward primer incorporates a SalI site at the linker-Fc junction, as well as introducing BspEI and RsrII sites into the Fc region without affecting the coding sequence, while the reverse primer adds an EcoRI site after the stop codon. The Fc coding sequence was then amplified from a leukocyte cDNA library (BD Biosciences Clontech, Palo Alto, CA) using 25 pmol of these primers in a 25 µl PCR reaction using Expand High Fidelity System (Boehringer Mannheim, Indianapolis, IN) according to manufacturer's standard protocol in a MJ Thermocycler using the following cycles: 94°C 2 minutes; 30 cycles of (94°C 30

seconds, 58°C 30 seconds, 72°C 45 seconds), followed by 72°C for 10 minutes. The expected sized band (696 bp) was gel purified with a Gel Extraction kit (Qiagen, Valencia, CA) and ligated into the intermediate cloning vector pGEM T-Easy (Promega, Madison, WI). DNA was transformed into DH5 α cells (Invitrogen, Carlsbad, CA) and miniprep cultures grown and purified with a Plasmid Miniprep Kit (Qiagen, Valencia, CA), both according to manufacturer's standard protocols. Once the sequence was confirmed, this insert was digested out with Sal/EcoRI restriction enzymes, gel purified, and cloned into the Sall/EcoRI sites of the plasmid pED.dC.Epo (above) in a similar manner, to generate the mammalian expression plasmid pED.dC.EpoFc. In another experiment this plasmid was also digested with RsrII/XmaI, and the corresponding fragment from pSYN-Fc-002, which contains the Asn 297 Ala mutation (EU numbering) was cloned in to create pED.dC.EPO-Fc N297A (pSYN-EPO-004). Expression in mammalian cells was as described in Example 26. The amino acid sequence of EpoFc with an eight amino acid linker is provided in figure 2j. During the process of this alternative cloning method, although the exact EpoFc amino acid sequence was preserved (figure 2J), a number of non-coding changes were made at the nucleotide level (figure 3J). These are G6A (G at nucleotide 6 changed to A) (eliminate possible secondary structure in primer), G567A (removes endogenous SbfI site from Epo), A582G (removes EcoRI site from linker), A636T and T639G (adds unique BspEI site to Fc), and G651C (adds unique RsrII site to Fc). The nucleotide sequence in figure 3J is from the construct made in Example 25, which incorporates these differences from the sequence of the construct from Example 24.

Example 26: EPO-Fc Homodimer And Monomer-dimer Hybrid Expression And Purification

[0277] DG44 cells were plated in 100 mm tissue culture petri dishes and grown to a confluency of 50%-60%. A total of 10 μ g of DNA was used to transfect one 100 mm dish: for the homodimer transfection, 10 μ g of pED.dC.EPO-Fc; for the monomer-dimer hybrid transfection, 8 μ g of pED.dC.EPO-Fc + 2 μ g of pcDNA3-FlagFc. The constructs used were cloned as described in Example 24. The cloning method described in Example 25 could also be used to obtain constructs for use in this example. The cells were transfected as described in the Superfect transfection reagent manual (Qiagen, Valencia, CA). Alternatively,

pED.dC.EPO-Fc was cotransfected with pSYN-Fc-016 to make an untagged monomer. The media was removed from transfection after 48 hours and replaced with MEM Alpha without nucleosides plus 5% dialyzed fetal bovine serum for both transfections, while the monomer-dimer hybrid transfection was also supplemented with 0.2 mg/ml geneticin (Invitrogen, Carlsbad, CA). After 3 days, the cells were released from the plate with 0.25% trypsin and transferred into T25 tissue culture flasks, and the selection was continued for 10-14 days until the cells began to grow well as stable cell lines were established. Protein expression was subsequently amplified by the addition methotrexate.

[0278] For both cell lines, approximately 2×10^7 cells were used to inoculate 300 ml of growth medium in a 1700 cm² roller bottle (Corning, Corning, NY). The roller bottles were incubated in a 5% CO₂ at 37°C for approximately 72 hours. The growth medium was exchanged with 300 ml serum-free production medium (DMEM/F12 with 5 µg/ml bovine insulin and 10 µg/ml Gentamicin). The production medium (conditioned medium) was collected every day for 10 days and stored at 4°C. Fresh production medium was added to the roller bottles after each collection and the bottles were returned to the incubator. Prior to chromatography, the medium was clarified using a SuporCap-100 (0.8/0.2 µm) filter from Pall Gelman Sciences (Ann Arbor, MI). All of the following steps were performed at 4°C. The clarified medium was applied to Protein A Sepharose, washed with 5 column volumes of 1X PBS (10 mM phosphate, pH 7.4, 2.7 mM KCl, and 137 mM NaCl), eluted with 0.1 M glycine, pH 2.7, and then neutralized with 1/10 volume of 1 M Tris-HCl, pH 9.0. Protein was then dialyzed into PBS.

[0279] The monomer-dimer hybrid transfection protein sample was subject to further purification, as it contained a mixture of EPO-Fc:EPO-Fc homodimer, EPO-Fc:Flag-Fc monomer-dimer hybrid, and Flag-Fc:Flag-Fc homodimer. Material was concentrated and applied to a 2.6 cm x 60 cm (318 ml) Superdex 200 Prep Grade column at a flow rate of 4 ml/min (36 cm/hour) and then eluted with 3 column volumes of 1X PBS. Fractions corresponding to two peaks on the UV detector were collected and analyzed by SDS-PAGE. Fractions from the first peak contained either EPO-Fc:EPO-Fc homodimer or EPO-Fc:FlagFc monomer-dimer hybrid, while the second peak contained FlagFc:FlagFc homodimer. All fractions containing the monomer-dimer hybrid but

no FlagFc homodimer were pooled and applied directly to a 1.6 x 5 cm M2 anti-FLAG sepharose column (Sigma Corp.) at a linear flow rate of 60 cm/hour. After loading the column was washed with 5 column volumes PBS. Monomer-dimer hybrids were then eluted with 100 mM Glycine, pH 3.0. Elution fractions containing the protein peak were then neutralized by adding 1/10 volume of 1 M Tris-HCl, and analyzed by reducing and nonreducing SDS-PAGE. Fractions were dialyzed into PBS, concentrated to 1-5 mg/ml, and stored at -80°C.

[0280] Alternatively, fractions from first peak of the Superdex 200 were analyzed by SDS-PAGE, and only fractions containing a majority of EpoFc monomer-dimer hybrid, with a minority of EpoFc homodimer, were pooled. This pool, enriched for the monomer-dimer hybrid, was then reappplied to a Superdex 200 column, and fractions containing only EpoFc monomer-dimer hybrid were then pooled, dialyzed and stored as purified protein. Note that this alternate purification method could be used to purify non-tagged monomer-dimer hybrids as well.

Example 27: Administration of EpoFc Dimer and Monomer-Dimer Hybrid With an Eight Amino Acid Linker to Cynomolgus Monkeys

[0281] For pulmonary administration, aerosols of either EpoFc dimer or EpoFc monomer-dimer hybrid proteins (both with the 8 amino acid linker) in PBS, pH 7.4 were created with the Aeroneb Pro™ (AeroGen, Mountain View, CA) nebulizer, in-line with a Bird Mark 7A respirator, and administered to anesthetized naïve cynomolgus monkeys through endotracheal tubes (approximating normal tidal breathing). Both proteins were also administered to naïve cynomolgus monkeys by intravenous injection. Samples were taken at various time points, and the amount of Epo-containing protein in the resulting plasma was quantitated using the Quantikine IVD Human Epo Immunoassay (R&D Systems, Minneapolis, MN). Pharmacokinetic parameters were calculated using the software WinNonLin. Table 4 presents the bioavailability results of cynomolgus monkeys treated with EpoFc monomer-dimer hybrid or EpoFc dimer.

**TABLE 4: ADMINISTRATION OF EPOFC MONOMER-DIMER
HYBRID AND EPOFC DIMER TO MONKEYS**

Protein	Monkey #	Route	Approx. Deposited Dose ¹ (µg/kg)	C _{max} (ng/ml)	C _{max} (fmol/ml)	t _{1/2} (hr)	t _{1/2} avg (hr)
EpoFc monomer-dimer hybrid	CO6181	pulm	20	72.3	1014	23.6	25.2
	CO6214	pulm	20	50.1	703	23.5	
	CO7300	pulm	20	120	1684	36.2	
	CO7332	pulm	20	100	1403	17.5	
	CO7285	IV	25	749	10508	21.3	22.6
	CO7288	IV	25	566	7941	23	
	CO7343	IV	25	551	1014	23.5	
EpoFc dimer	DD026	pulm	15	10.7	120	11.5	22.1
	DD062	pulm	15	21.8	244	27.3	
	DD046	pulm	15	6.4	72	21.8	
	DD015	pulm	15	12.8	143	20.9	
	DD038	pulm	35	27	302	29	
	F4921	IV	150	3701	41454	15.1	14.6
	96Z002	IV	150	3680	41219	15.3	
	1261CQ	IV	150	2726	30533	23.6	
	127-107	IV	150	4230	47379	15.0	
	118-22	IV	150	4500	50403	8.7	
	126-60	IV	150	3531	39550	9.8	

¹ Based on 15% deposition fraction of nebulized dose as determined by gamma scintigraphy

[0282] The percent bioavailability (F) was calculated for the pulmonary doses using the following equation:

$$F = (\text{AUC pulmonary} / \text{Dose pulmonary}) / (\text{AUC IV} / \text{Dose IV}) * 100$$

**TABLE 5: CALCULATION OF PERCENT BIOAVAILABILITY FOR EPOFc
MONOMER-DIMER HYBRID V. DIMER AFTER PULMONARY
ADMINISTRATION TO NAÏVE CYNOMOLGUS MONKEYS**

Protein	Monkey #	Approx. Dose ¹ (deposited)	AUC ng•hr/mL	Bioavailability ² (F)	Average Bioavailability
EpoFc monomer- dimer hybrid	CO6181	20 µg/kg	3810	25.2%	34.9%
	CO6214	20 µg/kg	3072	20.3%	
	CO7300	20 µg/kg	9525	63.0%	
	CO7332	20 µg/kg	4708	31.1%	
EpoFc dimer	DD026	15 µg/kg	361	5.1%	10.0 %
	DD062	15 µg/kg	1392	19.6%	
	DD046	15 µg/kg	267	3.8%	
	DD015	15 µg/kg	647	9.1%	
	DD038	35 µg/kg	2062	12.4%	

¹ Based on 15% deposition fraction of nebulized dose as determined by gamma scintigraphy

² Mean AUC for IV EpoFc monomer-dimer hybrid = 18,913 ng•hr/mL (n=3 monkeys), dosed at 25 µg/kg. Mean AUC for IV EpoFc dimer = 70, 967 ng•hr/mL (n=6 monkeys), dosed at 150 µg/kg

[0283] The pharmacokinetics of EpoFc with an 8 amino acid linker administered to cynomolgus monkeys is presented in figure 11. The figure compares the EpoFc dimer with the EpoFc monomer-dimer hybrid in monkeys after administration of a single pulmonary dose. Based on a molar comparison significantly higher serum levels were obtained in monkeys treated with the monomer-dimer hybrid compared to the dimer.

Example 28: Subcutaneous Administration of EPOFc Monomer-dimer Hybrid

[0284] To compare serum concentrations of known erythropoietin agents with EPOFc monomer-dimer hybrids, both EPOFc monomer-dimer hybrid and Aranesp® (darbepoetin alfa), which is not a chimeric fusion protein, were administered subcutaneously to different monkeys and the serum concentration of both was measured over time.

[0285] Cynomolgus monkeys (n = 3 per group) were injected subcutaneously with 0.025 mg/kg EpoFc monomer-dimer hybrid. Blood samples were collected predose and at times up to 144 hours post dose. Serum was prepared from the blood and stored frozen until analysis by ELISA (Human Epo Quantikine Immunoassay) (R & D Systems, Minneapolis, MN). Pharmacokinetic parameters were determined using WinNonLin[®] software (Pharsight, Mountainview, CA).

[0286] The results indicated the serum concentrations of both EPOFc monomer-dimer hybrid and Aranesp[®] (darbepoetin alfa) were equivalent over time, even though the administered molar dose of Aranesp[®] (darbepoetin alfa) was slightly larger (Table 6) (figure 12).

TABLE 6

	Route	Dose ($\mu\text{g/kg}$)	Dose (nmol/kg)	Cmax (ng/mL)	AUC (ng•hr•mL ⁻¹)	T _{1/2} (hr)	% Bioavailability (F)
EpoFc Monomer -dimer hybrid	Subcutaneous	25	0.3	133 $\pm$ 34	10,745 $\pm$ 3,144	26 $\pm$ 5	57 $\pm$ 17
Aranesp [®]	Subcutaneous	20	0.54	83 $\pm$ 11	5390 $\pm$ 747	22 $\pm$ 2	53 $\pm$ 8

Example 29: Intravenous Administration of EPOFc Monomer-dimer Hybrid

[0287] To compare serum concentrations of known erythropoietin agents with EPOFc monomer-dimer hybrids, EPOFc monomer-dimer hybrid, Aranesp[®] (darbepoetin alfa), and Epogen[®] (epoetin alfa), neither of which is a chimeric fusion protein, were administered intravenously to different monkeys and the serum concentration of both was measured over time.

[0288] Cynomolgus monkeys (n = 3 per group) were injected intravenously with 0.025 mg/kg EpoFc monomer-dimer hybrid. Blood samples were collected predose and at times up to 144 hours post dose. Serum was prepared from the blood and stored frozen until analysis by ELISA (Human Epo Quantikine Immunoassay) (R & D Systems, Minneapolis, MN). Pharmacokinetic

parameters were determined using WinNonLin[®] software (Pharsight, Mountainview, CA).

[0289] The results indicated the serum concentration versus time (AUC) of EPOFc monomer-dimer hybrid was greater than the concentrations of either Epogen[®] (epoetin alfa) or Aranesp[®] (darbepoetin alfa), even though the monkeys received larger molar doses of both Epogen[®] (epoetin alfa) and Aranesp[®] (darbepoetin alfa) (Table 7) (Figure 13).

TABLE 7

	Route	Dose ($\mu\text{g/kg}$)	Dose (nmol/kg)	Cmax (ng/mL)	AUC ($\text{ng}\cdot\text{hr}\cdot\text{mL}^{-1}$)	T _{1/2} (hr)
EpoFc Monomer- dimer hybrid	Intravenous	25	0.3	622 $\pm$ 110	18,913 $\pm$ 3,022	23 $\pm$ 1
Aranesp [®]	Intravenous	20	0.54	521 $\pm$ 8	10,219 $\pm$ 298	20 $\pm$ 1
Epogen	Intravenous	20	0.66	514 $\pm$ 172	3936 $\pm$ 636	6.3 $\pm$ 0.6

Example 30: Alternative Purification of EpoFc Monomer-dimer Hybrid

[0290] Yet another alternative for purifying EPO-Fc is described herein. A mixture containing Fc, EpoFc monomer-dimer hybrid, and EpoFc dimer was applied to a Protein A Sepharose column (Amersham, Uppsala, Sweden). The mixture was eluted according to the manufacturer's instructions. The Protein A Sepharose eluate, containing the mixture was buffer exchanged into 50 mM Tris-Cl (pH 8.0). The protein mixture was loaded onto an 8 mL Mimetic Red 2 XL column (ProMetic Life Sciences, Inc., Wayne, NJ) that had been equilibrated in 50 mM Tris-Cl (pH 8.0). The column was then washed with 50 mM Tris-Cl (pH 8.0); 50 mM NaCl. This step removed the majority of the Fc. EpoFc monomer-dimer hybrid was specifically eluted from the column with 50 mM Tris-Cl (pH 8.0); 400 mM NaCl. EpoFc dimer can be eluted and the column regenerated with 5 column volumes of 1 M NaOH. Eluted fractions from the column were analyzed by SDS-PAGE (Figure 14).

Example 31: Cloning of Igk signal sequence - Fc construct for making untagged Fc alone.

[0291] The coding sequence for the constant region of IgG1 (EU # 221-447; the Fc region) was obtained by PCR amplification from a leukocyte cDNA library (Clontech, CA) using the following primers:

rcFc-F 5'- GCTGCGGTGACAAAACCTCACACATGCCCACCGTGCCCAGCTCCGGAACCTCCTGGGCGGACCGTCAGTC -3' (SEQ ID NO: 84)

rcFc-R 5'- ATTGGAATTCTCATTTACCCGGAGACAGGGAGAGGC -3' (SEQ ID NO: 85)

[0292] The forward primer adds three amino acids (AAV) and a Sall cloning site before the beginning of the Fc region, and also incorporates a BspEI restriction site at amino acids 231-233 and an RsrII restriction site at amino acids 236-238 using the degeneracy of the genetic code to preserve the correct amino acid sequence (EU numbering). The reverse primer adds an EcoRI cloning site after the stop codon of the Fc. A 25 µl PCR reaction was carried out with 25 pmol of each primer using Expand High Fidelity System (Boehringer Mannheim, Indianapolis, IN) according to the manufacturer's standard protocol in a MJ Thermocycler using the following cycles: 94°C 2 minutes; 30 cycles of (94°C 30 seconds, 58°C 30 seconds, 72°C 45 seconds), 72°C 10 minutes. The expected sized band (~696 bp) was gel purified with a Gel Extraction kit (Qiagen, Valencia CA), and cloned into pGEM T-Easy (Promega, Madison, WI) to produce an intermediate plasmid pSYN-Fc-001 (pGEM T-Easy/Fc).

[0293] The mouse Igk signal sequence was added to the Fc CDS using the following primers:

rc-Igk sig seq-F: 5'-TTTAAGCTTGCCGCCACCATGGAGACAGACACACTCC TGCTATGGGTACTGCTGCTCTGGGTTCCAGGTTCCACTGGTGACAAAACCTCACACATGCCCACCG -3' (SEQ ID NO: 86)

Fc-noXma-GS-R: 5'- GGTCAGCTCATCGCGGGATGGG -3' (SEQ ID NO: 87)

Fc-noXma-GS-F: 5'- CCCATCCCGCGATGAGCTGACC -3' (SEQ ID NO: 88)

[0294] The rc-Igk signal sequence-F primer adds a HindIII restriction site to the 5' end of the molecule, followed by a Kozak sequence (GCCGCCACC) (SEQ ID NO: 89) followed by the signal sequence from the mouse Igk light chain, directly abutted to the beginning of the Fc sequence (EU# 221). The Fc-noXma-

GS-F and -R primers remove the internal XmaI site from the Fc coding sequence, using the degeneracy of the genetic code to preserve the correct amino acid sequence. Two 25 µl PCR reactions were carried out with 25 pmol of either rc-Igk signal sequence-F and Fc-noXma-GS-R or Fc-noXma-GS-F and rcFc-R using Expand High Fidelity System (Boehringer Mannheim, Indianapolis, IN) according to the manufacturer's standard protocol in a MJ Thermocycler. The first reaction was carried out with 500 ng of leukocyte cDNA library (BD Biosciences Clontech, Palo Alto, CA) as a template using the following cycles: 94°C 2 minutes; 30 cycles of (94°C 30 seconds, 55°C 30 seconds, 72°C 45 seconds), 72°C 10 minutes. The second reaction was carried out with 500 ng of pSYN-Fc-001 as a template (above) using the following cycles: 94°C 2 minutes; 16 cycles of (94°C 30 seconds, 58°C 30 seconds, 72°C 45 seconds), 72°C 10 minutes. The expected sized bands (~495 and 299 bp, respectively) were gel purified with a Gel Extraction kit (Qiagen, Valencia CA), then combined in a PCR reaction with 25 pmol of rc-Igk signal sequence-F and rcFc-R primers and run as before, annealing at 58°C and continuing for 16 cycles. The expected sized band (~772 bp) was gel purified with a Gel Extraction kit (Qiagen, Valencia CA) and cloned into pGEM T-Easy (Promega, Madison, WI) to produce an intermediate plasmid pSYN-Fc-007 (pGEM T-Easy/Igk sig seq-Fc). The entire Igk signal sequence-Fc cassette was then subcloned using the HindIII and EcoRI sites into either the pEE6.4 (Lonza, Slough, UK) or pcDNA3.1 (Invitrogen, Carlsbad, CA) mammalian expression vector, depending on the system to be used, to generate pSYN-Fc-009 (pEE6.4/Igk sig seq-Fc) and pSYN-Fc-015 (pcDNA3/Igk sig seq-Fc).

Example 32: Cloning of Igk signal sequence - Fc N297A construct for making untagged Fc N297A alone.

[0295] In order to mutate Asn 297 (EU numbering) of the Fc to an Ala residue, the following primers were used:

N297A-F 5'- GAGCAGTACGCTAGCACGTACCG -3' (SEQ ID NO: 90)

N297A-R 5'- GGTACGTGCTAGCGTACTGCTCC -3' (SEQ ID NO: 91)

[0296] Two PCR reactions were carried out with 25 pmol of either rc-Igk signal sequence-F and N297A-R or N297A-F and rcFc-R using Expand High Fidelity System (Boehringer Mannheim, Indianapolis, IN) according to the manufacturer's standard protocol in a MJ Thermocycler. Both reactions were

carried out using 500 ng of pSYN-Fc-007 as a template using the following cycles: 94°C 2 minutes; 16 cycles of (94°C 30 seconds, 48°C 30 seconds, 72°C 45 seconds), 72°C 10 minutes. The expected sized bands (~319 and 475 bp, respectively) were gel purified with a Gel Extraction kit (Qiagen, Valencia CA), then combined in a PCR reaction with 25 pmol of rc-Igk signal sequence-F and rcFc-R primers and run as before, annealing at 58°C and continuing for 16 cycles. The expected sized band (~772 bp) was gel purified with a Gel Extraction kit (Qiagen, Valencia CA) and cloned into pGEM T-Easy (Promega, Madison, WI) to produce an intermediate plasmid pSYN-Fc-008 (pGEM T-Easy/Igk sig seq-Fc N297A). The entire Igk signal sequence-Fc alone cassette was then subcloned using the HindIII and EcoRI sites into either the pEE6.4 (Lonza, Slough, UK) or pcDNA3.1 (Invitrogen, Carlsbad, CA) mammalian expression vector, depending on the system to be used, to generate pSYN-Fc-010 (pEE6.4/Igk sig seq-Fc N297A) and pSYN-Fc-016 (pcDNA3/Igk sig seq-Fc N297A).

[0297] These same N297A primers were also used with rcFc-F and rcFc-R primers and pSYN-Fc-001 as a template in a PCR reaction followed by subcloning as indicated above to generate pSYN-Fc-002 (pGEM T Easy/Fc N297A).

Example 33: Cloning of EpoFc and Fc into single plasmid for double gene vectors for making EpoFc wildtype or N297A monomer-dimer hybrids, and expression.

[0298] An alternative to transfecting the EpoFc and Fc constructs on separate plasmids is to clone them into a single plasmid, also called a double gene vector, such as used in the Lonza Biologics (Slough, UK) system. The RsrII/EcoRI fragment from pSYN-Fc-002 was subcloned into the corresponding sites in pEE12.4 (Lonza Biologics, Slough, UK) according to standard procedures to generate pSYN-Fc-006 (pEE12.4/Fc N297A fragment). The pSYN-EPO-004 plasmid was used as a template for a PCR reaction using Epo-F primer from Example 25 and the following primer:

EpoRsr-R: 5'- CTGACGGTCCGCCCAGGAGTTCCG
GAGCTGGGCACGGTGGGCATG TGTGAGTTTTGTGACCGCAGCGG -3'
(SEQ ID NO: 91)

[0299] A PCR reaction was carried out using Expand High Fidelity System (Boehringer Mannheim, Indianapolis, IN) according to the manufacturer's standard protocol in a MJ Thermocycler as indicated above, for 16 cycles with 55°C annealing temperature. The expected sized band (~689 bp) was gel purified with a Gel Extraction kit (Qiagen, Valencia CA) and cloned into pSYN-Fc-006 using the HindIII/RsrII restriction sites, to generate pSYN-EPO-005 (pEE12.4/EpoFc N297A). The double gene vector for the EpoFc N297A monomer-dimer hybrid was then constructed by cloning the NotI/BamHI fragment from pSYN-Fc-010 into the corresponding sites in pSYN-EPO-005 to generate pSYN-EPO-008 (pEE12.4-6.4/EpoFc N297A/Fc N297A).

[0300] The wild type construct was also made by subcloning the wild type Fc sequence from pSYN-Fc-001 into pSYN-EPO-005 using the RsrII and EcoRI sites, to generate pSYN-EPO-006 (pEE12.4/EpoFc). The double gene vector for the EpoFc monomer-dimer hybrid was then constructed by cloning the NotI/BamHI fragment from pSYN-Fc-009 into the corresponding sites in pSYN-EPO-006 to generate pSYN-EPO-007 (pEE12.4-6.4/EpoFc /Fc).

[0301] Each plasmid was transfected into CHOK1SV cells and positive clones identified and adapted to serum-free suspension, as indicated in the Lonza Biologics Manual for Standard Operating procedures (Lonza Biologics, Slough, UK), and purified as indicated for the other monomer-dimer constructs.

Example 34: Cloning of human IFN β Fc, IFN β -Fc N297A with eight amino acid linkers and Ig κ -Fc-6His constructs

[0302] 10 ng of a human genomic DNA library from Clontech (BD Biosciences Clontech, Palo Alto, CA) was used as a template to isolate human IFN β with its native signal sequence using the following primers:

IFN β -F H3/SbfI:

5'- CTAGCCTGCAGGAAGCTTGCCGCCACCATGACCA
ACAAGTGTCTCCTC -3' (SEQ ID NO: 92)

IFN β -R (EFAG) SalI:

5'TTTGTGCGACCGCAGCGGCCGGCGAACTCGTTTCGG
AGGTAACCTGTAAG -3' (SEQ ID NO: 93)

[0303] The reverse primer was also used to create an eight amino acid linker sequence (EFAGAAAV) (SEQ ID NO: 94) on the 3' end of the human IFN β sequence. The PCR reaction was carried out using the Expand High Fidelity System (Boehringer Mannheim, Indianapolis, IN) according to the manufacturer's standard protocol in a Rapid Cyclor thermocycler (Idaho Technology, Salt Lake City, UT). A PCR product of the correct size (~607 bp) was gel purified using a Gel Extraction kit (Qiagen; Valencia, CA), cloned into TA cloning vector (Promega, Madison, WI) and sequenced. This construct was named pSYN-IFN β -002. pSYN-IFN β -002 was digested with SbfI and Sall and cloned into pSP72 (Promega) at PstI and Sall sites to give pSYN-IFN β -005.

[0304] Purified pSYN-Fc-001 (0.6 μ g) was digested with Sall and EcoRI and cloned into the corresponding sites of pSYN-IFN β -005 to create the plasmid pSYN-IFN β -006 which contains human IFN β linked to human Fc through an eight amino acid linker sequence. pSYN-IFN β -006 was then digested with SbfI and EcoRI and the full-length IFN β -Fc sequence cloned into the PstI and EcoRI sites of pEDdC.sig to create plasmid pSYN-IFN β -008.

[0305] pSYN-Fc-002 containing the human Fc DNA with a single amino acid change from asparagine to alanine at position 297 (N297A; EU numbering) was digested with BspEI and XmaI to isolate a DNA fragment of ~365 bp containing the N297A mutation. This DNA fragment was cloned into the corresponding sites in pSYN-IFN β -008 to create plasmid pSYN-IFN β -009 that contains the IFN β -Fc sequence with an eight amino acid linker and an N297A mutation in Fc in the expression vector, pED.dC.

[0306] Cloning of Igk signal sequence-Fc N297A – 6His. The following primers were used to add a 6xHis tag to the C terminus of the Fc N297A coding sequence:

Fc GS-F: 5'- GGCAAGCTTGCCGCCACCATGGAGACAGACACACTCC -3' (SEQ ID NO: 95)

Fc.6His-R: 5'- TCAGTGGTGATGGTGATGATGTTTACCCGGAGACAGGGAG -3' (SEQ ID NO: 96)

Fc.6His-F: 5'- GGTAACATCATCACCATCACCCTGAGAATTCC AATATCACTAGTGAATTCCG -3' (SEQ ID NO: 97)

Sp6+T-R: 5'- GCTATTTAGGTGACACTATAGAATACTCAAGC -3' (SEQ ID NO: 98)

[0307] Two PCR reactions were carried out with 50 pmol of either Fc GS-F and Fc.6His-R or Fc.6His-F and Sp6+T-R using the Expand High Fidelity System (Boehringer Mannheim, Indianapolis, IN) according to the manufacturer's standard protocol in a MJ Thermocycler. Both reactions were carried out using 500 ng of pSYN-Fc-008 as a template in a 50 µl reaction, using standard cycling conditions. The expected sized bands (~780 and 138 bp, respectively) were gel purified with a Gel Extraction kit (Qiagen, Valencia CA), then combined in a 50 µl PCR reaction with 50 pmol of Fc GS-F and Sp6+T-R primers and run as before, using standard cycling conditions. The expected sized band (~891 bp) was gel purified with a Gel Extraction kit (Qiagen, Valencia CA) and cloned into pcDNA6 V5-His B using the HindIII and EcoRI sites to generate pSYN-Fc-014 (pcDNA6/Igk sig seq-Fc N297A-6 His).

Example 35: Expression and purification of IFNβFc, IFNβ-Fc N297A homodimer and IFNβ-Fc N297A monomer-dimer hybrid

[0308] CHO DG44 cells were plated in 100 mm tissue culture dishes and grown to a confluency of 50%-60%. A total of 10 µg of DNA was used to transfect a single 100 mm dish. For the homodimer transfection, 10 µg of the pSYN-FNβ-008 or pSYN-IFNβ-009 construct was used; for the monomer-dimer hybrid transfection, 8 µg of the pSYN-IFNβ-009 + 2 µg of pSYN-Fc-014 construct was used. The cells were transfected using Superfect transfection reagents (Qiagen, Valencia, CA) according to the manufacturer's instructions. 48 to 72 hours post-transfection, growth medium was removed and cells were released from the plates with 0.25% trypsin and transferred to T75 tissue culture flasks in selection medium (MEM Alpha without nucleosides plus 5% dialyzed fetal bovine serum). The selection medium for the monomer-dimer hybrid transfection was supplemented with 5 µg/ml Blastidin (Invitrogen, Carlsbad, CA). Selection was continued for 10-14 days until the cells began to grow well and stable cell lines were established. Protein expression was subsequently amplified by the addition methotrexate: ranging from 10 to 50 nM.

[0309] For all cell lines, approximately 2×10^7 cells were used to inoculate 300 ml of growth medium in a 1700 cm² roller bottle (Corning, Corning, NY). The

roller bottles were incubated in a 5% CO₂ incubator at 37°C for approximately 72 hours. The growth medium was then exchanged with 300 ml serum-free production medium (DMEM/F12 with 5 µg/ml human insulin). The production medium (conditioned medium) was collected every day for 10 days and stored at 4°C. Fresh production medium was added to the roller bottles after each collection and the bottles were returned to the incubator. Prior to chromatography, the medium was clarified using a SuporCap-100 (0.8/0.2 µm) filter from Pall Gelman Sciences (Ann Arbor, MI). All of the following steps were performed at 4°C. The clarified medium was applied to Protein A Sepharose, washed with 5 column volumes of 1X PBS (10 mM phosphate, pH 7.4, 2.7 mM KCl, and 137 mM NaCl), eluted with 0.1 M glycine, pH 2.7, and then neutralized with 1/10 volume of 1 M Tris-HCl pH 8.0, 5 M NaCl. The homodimer proteins were further purified over a Superdex 200 Prep Grade sizing column run and eluted in 50 mM sodium phosphate pH 7.5, 500 mM NaCl, 10% glycerol.

[0310] The monomer-dimer hybrid protein was subject to further purification since it contained a mixture of IFNβFc N297A:IFNβFc N297A homodimer, IFNβFc N297A: Fc N297A His monomer-dimer hybrid, and Fc N297A His: Fc N297A His homodimer. Material was applied to a Nickel chelating column in 50 mM sodium phosphate pH 7.5, 500 mM NaCl. After loading, the column was washed with 50 mM imidazole in 50 mM sodium phosphate pH 7.5, 500 mM NaCl and protein was eluted with a gradient of 50 – 500 mM imidazole in 50 mM sodium phosphate pH 7.5, 500 mM NaCl. Fractions corresponding to elution peaks on a UV detector were collected and analyzed by SDS-PAGE. Fractions from the first peak contained IFNβFc N297A: Fc N297A His monomer-dimer hybrid, while the second peak contained Fc N297A His: Fc N297A His homodimer. All fractions containing the monomer-dimer hybrid, but no Fc homodimer, were pooled and applied directly to a Superdex 200 Prep Grade sizing column, run and eluted in 50 mM sodium phosphate pH 7.5, 500 mM NaCl, 10% glycerol. Fractions containing IFNβ-Fc N297A:Fc N297A His monomer-dimer hybrids were pooled and stored at -80°C.

Example 36: Antiviral assay for IFN β activity

[0311] Antiviral activity (IU/ml) of IFN β fusion proteins was determined using a CPE (cytopathic effect) assay. A549 cells were plated in a 96 well tissue culture plate in growth media (RPMI 1640 supplemented with 10% fetal bovine serum (FBS) and 2 mM L-glutamine) for 2 hours at 37°C, 5% CO₂. IFN β standards and IFN β fusion proteins were diluted in growth media and added to cells in triplicate for 20 hours at 37°C, 5% CO₂. Following incubation, all media was removed from wells, encephalomyocarditis virus (EMCV) was diluted in growth media and added (3000 pfu/well) to each well with the exception of control wells. Plates were incubated at 37°C, 5% CO₂ for 28 hours. Living cells were fixed with 10% cold trichloroacetic acid (TCA) and then stained with Sulforhodamine B (SRB) according to published protocols (Rubinstein et al. 1990, *J. Natl. Cancer Inst.* 82, 1113). The SRB dye was solubilized with 10 mM Tris pH 10.5 and read on a spectrophotometer at 490 nm. Samples were analyzed by comparing activities to a known standard curve ranging from 10 to 0.199 IU/ml. The results are presented below in Table 8 and demonstrate increased antiviral activity of monomer-dimer hybrids.

TABLE 8: INTERFERON BETA ANTIVIRAL ASSAY
HOMODIMER V. MONOMER-DIMER HYBRID

Protein	Antiviral Activity (IU/nmol)	Std dev
IFN β -Fc 8aa linker homodimer	4.5×10^5	0.72×10^5
IFN β Fc N297A 8aa linker homodimer	3.21×10^5	0.48×10^5
IFN β Fc N297A 8aa linker: Fc His monomer-dimer hybrid	12.2×10^5	2×10^5

Example 37: Administration of IFN β Fc Homodimer and Monomer-Dimer Hybrid With an Eight Amino Acid Linker to Cynomolgus Monkeys

[0312] For pulmonary administration, aerosols of either IFN β Fc homodimer or IFN β Fc N297A monomer-dimer hybrid proteins (both with the 8

amino acid linker) in PBS, pH 7.4, 0.25% HSA were created with the Aeronex Pro™ (AeroGen, Mountain View, CA) nebulizer, in-line with a Bird Mark 7A respirator, and administered to anesthetized naïve cynomolgus monkeys through endotracheal tubes (approximating normal tidal breathing). Blood samples were taken at various time points, and the amount of IFN β -containing protein in the resulting serum was quantitated using a human IFN β Immunoassay (Biosource International, Camarillo, CA). Pharmacokinetic parameters were calculated using the software WinNonLin. Table 9 presents the results of cynomolgus monkeys treated with IFN β Fc N297A monomer-dimer hybrid or IFN β Fc homodimer.

TABLE 9: ADMINISTRATION OF IFN β FC N297A MONOMER-DIMER HYBRID AND IFN β FC HOMODIMER TO MONKEYS

Protein	Monkey #	Route	Approx. Deposited Dose ¹ (μ g/kg)	C _{max} (ng/ml)	AUC (hr*ng/ml)	t _{1/2} (hr)	t _{1/2} avg (hr)
IFN β Fc N297A monomer-dimer hybrid	CO7308	pulm	20	23.3	987.9	27.6	27.1
	CO7336	pulm	20	22.4	970.6	25.6	
	CO7312	pulm	20	21.2	1002.7	28.0	
IFN β Fc homodimer	CO7326	pulm	20	2.6	94.6	11.1	11.4
	CO7338	pulm	20	5.0	150.6	11.7	

¹ Based on 15% deposition fraction of nebulized dose as determined by gamma scintigraphy

[0313] The pharmacokinetics of IFN β Fc with an 8 amino acid linker administered to cynomolgus monkeys is presented in figure 15. The figure compares the IFN β Fc homodimer with the IFN β Fc N297A monomer-dimer hybrid in monkeys after administration of a single pulmonary dose. Significantly higher serum levels were obtained in monkeys treated with the monomer-dimer hybrid compared to the homodimer.

[0314] Serum samples were also analyzed for neopterin levels (a biomarker of IFN β activity) using a neopterin immunoassay (MP Biomedicals,

Orangeburg, NY). The results for this analysis are shown in figure 16. The figure compares neopterin stimulation in response to the IFN β -Fc homodimer and the IFN β -Fc N297A monomer-dimer hybrid. It can be seen that significantly higher neopterin levels were detected in monkeys treated with IFN β -Fc N297A monomer-dimer hybrid as compared to the IFN β -Fc homodimer.

[0315] All numbers expressing quantities of ingredients, reaction conditions, and so forth used in the specification and claims are to be understood as being modified in all instances by the term "about." Accordingly, unless indicated to the contrary, the numerical parameters set forth in the specification and attached claims are approximations that may vary depending upon the desired properties sought to be obtained by the present invention. At the very least, and not as an attempt to limit the application of the doctrine of equivalents to the scope of the claims, each numerical parameter should be construed in light of the number of significant digits and ordinary rounding approaches.

[0316] All references cited herein are incorporated herein by reference in their entirety and for all purposes to the same extent as if each individual publication or patent or patent application was specifically and individually indicated to be incorporated by reference in its entirety for all purposes. To the extent publications and patents or patent applications incorporated by reference contradict the disclosure contained in the specification, the specification is intended to supercede and/or take precedence over any such contradictory material.

[0317] Many modifications and variations of this invention can be made without departing from its spirit and scope, as will be apparent to those skilled in the art. The specific embodiments described herein are offered by way of example only and are not meant to be limiting in any way. It is intended that the specification and examples be considered as exemplary only, with a true scope and spirit of the invention being indicated by the following claims.

Claims:

1. A method of treating a subject with a cardiovascular disease or condition comprising administering a chimeric protein to the subject, such that said cardiovascular disease or condition is treated, wherein said chimeric protein comprises a first and second polypeptide chain,
 - a) said first chain comprising an FcRn binding partner, and a biologically active molecule and
 - b) said second chain comprising an FcRn binding partner without a biologically active molecule or a variable region of an immunoglobulin.
2. The method of claim 1, wherein the cardiovascular disease or condition is chosen from stroke, brain ischemia, cerebral ischemia, myocardial ischemia, cardiac ischemia, ischemic heart disease, myocardial infarction, coronary artery disease, acute coronary syndrome, atherosclerosis, chronic heart failure, congestive heart failure, and reperfusion injury.
3. The method of claim 1, wherein the biologically active molecule is erythropoietin.




FIG. 1

FACTOR VII-Fc AMINO ACID SEQUENCE (SIGNAL PEPTIDE UNDERLINED, PROPEPTIDE IN BOLD)

```

1  MVSOALRLLC LLLGLOGCLA AVFVTQEEAH GVLHRRRRAN AFLEELRPGS
51  LERECKEEQC SFEEAREIFK DAERTKLEWI SYSDGDQCAS SPCQNGGSCK
101 DQLQSYICFC LPAFEGRNCE THKDDQLICV NENGGEQYC SDHTGTRSC
151 RCHEGYSLLA DGVSCTPTVE YPCGKIPILE KRNASKPQGR IVGGKVCPKG
201 ECPWQVLLLV NGAQLCGGTL INTIWVVSAA HCFDKIKNWR NLI AVLGEHD
251 LSEHDGDEQS RRVAQVIIPS TYVPGTTNHD IALLRLHQPV VLTDHVPLC
301 LPERTFSERT LAFVRFSLSV GWGQLLD RGA TALELMVLNV PRLMTQDCLQ
351 QSRKVG DSPN ITEYMFCAGY SDGSKDCKG DSGGPHATHY RGTWYLTGIV
401 SWGQGCATVG HFGVYTRVSQ YIEWLQKLMR SEPRPGVLLR APFPDKTHTC
451 PPCPAPELLG GPSVFLFPPK PKDTLMISRT PEVTCVVVDV SHEDPEVKFN
501 WYVDGVEVHN AKTKPREEQY NSTYRVVSVL TVLHQDWLNG KEYKCKVSNK
551 ALPAPIEKTI SKAKGQPREP QVYTLPPSRD ELTKNQVSLT CLVKGFYPSD
601 IAVEWESNGQ PENNYKTPP VLDS DGSFFL YSKLTVDKSR WQQGNVFSCS
651 VMHEALHNHY TQKSLSLSPG K

```

FIG. 2A

FACTOR IX-Fc AMINO ACID SEQUENCE (SIGNAL PEPTIDE UNDERLINED, PROPEPTIDE IN BOLD)

```

1  MORVNMIMAE SPGLITICLL GILLSAECTV FLDHENANKI LNRPKRYNSG
51  KLEEFVQGNL ERECMECKCS FEEAREVFEN TERTTEFWKQ YVDGDQCESN
101 PCLNGGSCKD DINSYECWCP FGFEGKNCEL DVTCKNIKNGR CEQFCKNSAD
151 NKVVCSCTEG YRLAENQKSC EPAVPFPCGR VSVSQTSLT RAETVFPDVD
201 YVNSTEAETI LDNITQSTQS FNDFTRVVGG EDAKPGQFPW QVVLNGKVDA
251 FCGGSIVNEK WIVTAAHCVE TGVKITVVAG EHNIEETEHT EQKRN VIRII
301 PHHNYNAAIN KYNHDIALLE LDEPLVLNSY VTPICIADKE YTNIFLKFGS
351 GYVSGWGRVF HKGRSALVLQ YLRVPLVDRA TCLRSTKFTI YNNMFCAGFH
401 EGGRDSCQGD SGGPHVTEVE GTSFLTGIIS WGEECAMKGK YGIYTKVSRY
451 VNWIKEKTKL TEFAGAAVD KTHTCPPCPA PELLGGPSVF LFPPKPKDTL
501 MISRTPEVTC VVVDVSHEDP EVKFNWYVDG VEVHNAKTKP REEQYNSTYR
551 VVSVLTVLHQ DWLNGKEYKC KVSNAKALPAP IEKTISKAKG QPREPQVYTL
601 PPSRDELTKN QVSLTCLVKG FYPSDIAVEW ESNGQPENNY KTPPVLDSD
651 GSFFLYSKLT VDKSRWQQGN VFSCSVMHEA LHNHYTQKSL SLSPGK

```

FIG. 2B

IFN α -Fc AMINO ACID SEQUENCE (8 AMINO ACID LINKER) (SIGNAL SEQUENCE UNDERLINED)

```

1  MALTFALLVA LLVLSCKSSC SVGCDLPQTH SLGSRRTLML LAQMRRISLF
51 SCLKDRHDFG FPQEEFGNQF QKAETIPVLH EMIQQIFNLF STKDSSAAWD
101 ETLLDKFYTE LYQQLNDLEA CVIQGVGVTE TPLMKEDSIL AVRKYFQRIT
151 LYLKEKKYSP CAWEVVRAEI MRSFSLSTNL QESLRSKEEF AGAAAVDKTH
201 TCPPCPAPEL LGGPSVFLFP PKPKDTLMIS RTPEVTCVVV DVSHEDPEVK
251 FNWYVDGVEV HNAKTKPREE QYNSTYRVVS VLTVLHQDWL NGKEYKCKVS
301 NKALPAPIEK TISKAKGQPR EPQVYTLPPS RDELTKNQVS LTCLVKGFYP
351 SDIAVEWESN GOPENNYKTT PPVLDSGGSF FLYSKLTVDK SRWQQGNVFS
401 CSVMHEALHN HYTQKSLSL PGK

```

FIG. 2C

IFN α -Fc Δ LINKER AMINO ACID SEQUENCE (SIGNAL SEQUENCE UNDERLINED)

```

1  MALTFALLVA LLVLSCKSSC SVGCDLPQTH SLGSRRTLML LAQMRRISLF
51 SCLKDRHDFG FPQEEFGNQF QKAETIPVLH EMIQQIFNLF STKDSSAAWD
101 ETLLDKFYTE LYQQLNDLEA CVIQGVGVTE TPLMKEDSIL AVRKYFQRIT
151 LYLKEKKYSP CAWEVVRAEI MRSFSLSTNL QESLRSKEDK THTCPPCPAP
201 ELLGGPSVFL FPPKPKDTLM ISRTPEVTCV VVDVSHEDPE VKFNWYVDGV
251 EVHNAKTKPR EEQYNSTYRV VSVLTVLHQD WLNGKEYKCK VSNKALPAPI
301 EKTISKAKGQ PREPQVYTL PPSRDELTKNQ VSLTCLVKGF YPSDIAVEWE
351 SNGQPENNYK TTPPVLDSDG SFFLYSKLTV DKSRWQQGNV FSCSVMHEAL
401 HNHYTQKSLS LSPGK

```

FIG. 2D

FLAGFc AMINO ACID SEQUENCE (SIGNAL SEQUENCE UNDERLINED)

```

1  METDTLLLWV LLLWVPGSTG DDYKDDDDKD KTHTCPPCPA PELLGGPSVF
51 LFPPKPKDTL MISRTPEVTC VVDVSHEDP EVKFNWYVDG VEVHNAKTKP
101 REEQYNSTYR VSVLTVLHQD DWLNGKEYKC KVSNAKPAP IEKTISKAKG
151 QPREPQVYTL PPSRDELTKN QVSLTCLVKG FYPSDIAVEW ESNGQPENNY
201 KTTTPVLDSG GSFFLYSKLT VDKSRWQQGN VFSCSVMHEA LHNHYTQKSL
251 SLSPGK

```

FIG. 2E

Epo-CCA-Fc AMINO ACID SEQUENCE (K^b SIGNAL SEQUENCE
UNDERLINED, ACIDIC COILED COIL IN BOLD)

```

1  MVPCTLLLLL AAALAPTOTR AGSRAPPRLI CDSRVLQRYL LEAEAENIT
51 TGCAEHCSLN ENITVPDTKV NFIYAWKRMEV GQQAVEVWQG LALLSEAVLR
101 GQALLVNSSQ PWEPLQLHVD KAVSGLRSLT TLLRALGAQK EAI SPPDAAS
151 AAPLRTITAD TFRKLFVYS NFLRGKLY TGEACRTGDR EFGGEYQALE
201 KEVAQLEAEN QALEKEVAQL EHEGGGPAPE LLGGPSVFLF PPKPKDTLMI
251 SRTPEVTCVV VDVSHEDPEV KFNWYVDGVE VHNATKPRE EQYNSTYRVV
301 SVLTVLHQDW LNGKEYKCKV SNKALPAPIE KTISKAKGQP REPQVYTLPP
351 SRDELTKNQV SLTCLVKGFY PSDIAVEWES NGQPENNYKT TPPVLDSGDS
401 FFLYSKLTVD KSRWQQGNVF SCSVMHEALH NHYTQKSLSL SPGK

```

FIG. 2F

CCB-Fc AMINO ACID SEQUENCE (K^b SIGNAL SEQUENCE UNDERLINED,
BASIC COILED COIL IN BOLD)

```

1  MVPCTLLLLL AAALAPTOTR AGEFGGEYQA LKKKVAQLKA KNQALKKKVA
51 QLKHKGGGPA PELLGGPSVF LFPPKPKDTL MISRTPEVTC VVDVSHEDP
101 EVKFNWYVDG VEVHNATKP REEQYNSTYR VSVLTVLHQ DWLNGKEYKC
151 KVSNAKALPAP IEKTISKAKG QPREPQVYTL PPSRDELTKN QVSLTCLVKG
201 FYPSDIAVEW ESNQOPENNY KTPPVLDSD GSFFLYSKLT VDKSRWQQGN
251 VFSCSVMHEA LHNHYTQKSL SLSPGK

```

FIG. 2G

CysFc AMINO ACID SEQUENCE (hIFNa SIGNAL SEQUENCE UNDERLINED)

```

1  MALTFALLVA LLVLSCSSC SVGCPPCPAP ELLGGPSVFL FPPKPKDTLM
51 ISRTPEVTCV VVDVSHEDPE VKFNWYVDGV EVHNATKPR EEQYNSTYRV
101 VSVLTVLHQD WLNGKEYKCK VSNKALPAPI EKTISKAKGQ PREPQVYTL
151 PSRDELTKNQ VSLTCLVKGF YPSDIAVEWE SNGQPENNYK TPPVLDSG
201 SFFLYSKLTV DKSrwQQGNV FSCSVMHEAL HNHYTQKSL LSPGK

```

FIG. 2H

IFN α GS15 Fc PROTEIN SEQUENCE (SIGNAL SEQUENCE UNDERLINED):

```
1  MALTFALLVA LLVLSCKSSC SVGCDLPQTH SLGSRRTLML LAQMRRISLF
51 SCLKDRHDFG FPQEEFGNQF QKAETIPVLH EMIQQIFNLF STKDSSAAWD
101 ETLLDKFYTE LYQQLNDLEA CVIQGVGVTE TPLMKEDSIL AVRKYFQRIT
151 LYLKEKKYSP CAWEVVRAEI MRSFSLSTNL QESLRSKEGG GSGGGGGSGG
201 GGSDKTHTCP PCPAPELLGG PSVFLFPPKP KDTLMIS RTP EVTCVVVDVS
251 HEDPEVKFNW YVDGVEVHNA KTKPREEQYN STYRVVSVLT VLHQDWLNGK
301 EYKCKVSNKA LPAPIEKTIS KAKGQPREPQ VYTLPPSRDE LTKNQVSLTC
351 LVKGFYPSDI AVEWESNGQP ENNYKTTTPV LDSDGSFFLY SKLTVDKSRW
401 QQGNVFSCSV MHEALHNHYT QKSLSLSPGK
```

FIG. 2I

EpoFc AMINO ACID SEQUENCE (SIGNAL SEQUENCE UNDERLINED,
LINKER IN BOLD)

```
1  MGVHECPAWL WLLLSLLSLP LGLPVLGAPP RLICDSRVLE RYLLEAKEAE
51 NITTGCAEHC SLNENITVPD TKVNFYAWKR MEVGQQAQEV WQGLALLSEA
101 VLRGQALLVN SSQPWEPLQL HVDKAVSGLR SLTTLLRALG AQKEAISPDP
151 AASAAPLRTI TADTFRKLFR VYSNFLRGKL KLYTGEACRT GDREFAGAAA
201 VDKTHTCPPC PAPELLGGPS VFLFPPKPKD TLMISRTPEV TCVVVDVSHE
251 DPEVKFNWYV DGVEVHNAKT KPREEQYNST YRVVSVLTVL HQDWLNGKEY
301 KCKVSNKALP APIEKTISKA KGQPREPQVY TLPPSRDEL TKNQVSLTCLV
351 KGFYPSDIAV EWESNGQPEN NYKTTTPVLD SDGSFFLYSK LTVDKSRWQQ
401 GNVFSCSVMH EALHNHYTQK SLSLSPGK
```

FIG. 2J

FACTOR VII-FC NUCLEOTIDE SEQUENCE (SIGNAL PEPTIDE UNDERLINED, PROPEPTIDE IN BOLD)

atggtctcccaggccctcaggctcctctgccttctgcttgggcttcagggtgcctggctgcag
tcttcgtaaccaggaggaagccacggcgctcctgcaccggcgccggcgcgccaacgcgttcct
ggaggagctgcggccgggctccctggagagggagtgcaaggaggagcagtgtccttcgaggag
gcccgggagatcttcaaggacgcggagaggacgaagctgttctggatttcttacagtgatggg
accagtgtgcctcaagtccatgccagaatgggggctcctgcaaggaccagctccagtcctatat
ctgcttctgcctccctgccttcgagggccggaactgtgagacgcacaaggatgaccagctgac
tgtgtgaacgagaacggcggtgtgagcagtactgcagtaccacacgggcaccaagcgctcct
gtcgggtgccacgaggggtactctctgctggcagacggggtgtcctgcacaccacagttgaata
tccatgtggaaaaatacctattctagaaaaaagaaatgccagcaaaccccaaggccgaattgtg
gggggcaagggtgtgccccaaaggggagtgccatggcaggtcctgttggttgatggagctc
agttgtgtggggggaccctgatcaacaccatctgggtggtctccgcggccactgtttcgaaa
aatcaagaactggaggaacctgatcgcggtgctgggcgagcacgacctcagcgagcacgacggg
gatgagcagagccggcgggtggcgcaggtcatcatcccagcacgtacgtcccgggcaccacca
accacgacatcgcgctgctccgctgcaccagcccggtggtcctcactgacctgtggtgcccct
ctgcctgcccgaacggacgttctctgagaggacgctggccttcgtgcgcttctcattggtcagc
ggctggggccagctgctggaccgtggcgccacggccctggagctcatgggtcctcaacgtgcccc
ggctgatgaccaggactgctgcagcagtcacggaagggtgggagactcccaaatatcacgga
gtacatgttctgtgccggctactcggatggcagcaaggactcctgcaagggggcacgtggaggc
ccacatgccaccaactaccggggcacgtggtacctgacgggcatcgtcagctggggccagggt
gcgcaaccgtggggccactttgggggttacaccaggggtctcccagtacatcgagtggctgcaaaa
gctcatgcgctcagagccacgcccaggagtcctcctgcgagccccatttcccgaaaaactcac
acgtgcccgcgctgcccagctccggaactgctgggcggacgctcagtcttctcttccccccaa
aaccaaggacacctcatgatctcccggaccctgaggtcacatgcgtggtggtggacgtgag
ccacgaagacctgaggtcaagttcaactggtacgtggacggcggtggaggtgcataatgccaag
acaaagccgcgggaggagcagtacaacagcacgtaccgtgtggtcagcgtcctcacgctcctgc
accaggactggctgaatggcaaggagtacaagtgaagggtctccaacaaagccctcccagcccc
catcgagaaaacctctccaaagccaaagggcagccccgagaaccacaggtgtacacctgccc
ccatcccgggatgagctgaccaagaaccaggtcagcctgacctgcctgggtcaaaggcttctatc
ccagcgacatcgccgtggagtgaggagcaatgggcagccggagaacaactacaagaccacgcc
tcccggtgttgactccgacggctccttcttctctacagcaagctcacggtggacaagagcagg
tggcagcaggggaacgtcttctcatgctccgtgatgcatgaggctctgcacaaccactacacgc
agaagagcctctccctgtctccgggtaaatga

FIG. 3A

FACTOR IX-FC NUCLEOTIDE SEQUENCE (SIGNAL PEPTIDE UNDERLINED, PROPEPTIDE IN BOLD)

atgcagcgcgtgaacatgatcatggcagaatcaccaggcctcatcaccatctgccttttaggat
atctactcagtgctgaat**gtcacg**tttttcttgatcatgaaaacgccaacaaaattctgaatcg
gccaaagaggtataattcaggtaaattggaagagtttgttcaagggaaccttgagagagaatgt
atggaagaaaagtgtagttttgaagaagcagagaagttttgaaaacactgaaagaacaactg
aattttggaagcagtatgttgatggagatcagtggtgagtc caatccatgtttaaatggcggcag
ttgcaaggatgacattaattcctatgaatgttggtgtccctttggatttgaaggaaagaactgt
gaattagatgtaacatgtaacattaagaatggcagatgcgagcagttttgtaaaaatagtgtg
ataacaaggtggtttgctcctgtactgagggatatcgacttgcagaaaaccagaagtcctgtga
accagcagtgccatttccatgtggaagagtttctgtttcaciaaacttctaagctcacccgtgct
gagactgtttttcctgatgtggactatgtaaattctactgaagctgaaaccattttggataaca
tcaactcaagcacccaatcatttaatagacttcactcgggtgttggtggagaagatgccaaacc
aggtcaattcccttggcaggttgtttgaatggtaaagttgatgcattctgtggaggctctatc
gttaatgaaaaatggattgtaactgctgccactgtgttgaaactggtgttaaaattacagttg
tcgcaggtgaacataatattgaggagacagaacatacagagcaaaagcgaaatgtgattcgaat
tattcctcaccacaactacaatgcagctattaataagtacaaccatgacattgcccttctggaa
ctggacgaacccttagtgctaaacagctacgttacacctatttgcaattgctgacaaggaataca
cgaacatcttcctcaaatttggatctggctatgtaagtggctggggaagagtcctccacaaagg
gagatcagcttttagttcttcagtaccttagagttccacttgtgaccgagccacatgtcttcga
tctacaaagttcaccatctataacaacatgttctgtgctggcttccatgaaggaggtagagatt
catgtcaaggagatagtgggggaccccatgttactgaagtgggaaggaccagtttcttaactgg
aattattagctggggtgaagagtggtgcaatgaaaggcaaataatggaatatataccaaggtatcc
cggatgtcaactggattaaggaaaaaacaagctcactgaattcgccggcgccgctgcggtcg
acaaaactcacacgtgcccgccgtgccagctccggaactgctggcgcgaccgtcagtcctcct
cttccccccaaaacccaaggacaccctcatgatctcccggaccctgaggtcacatgcgtgggtg
gtggacgtgagccacgaagaccctgaggtcaagttcaactggtacgtggacggcgtggaggtgc
ataatgccaagacaaagccgcgaggagcagtaacacagcagcgtaccgtgtggtcagcgtcct
cacgcctcctgcaccaggactggctgaatggcaaggagtacaagtgaaggtctccaacaaagcc
ctcccagccccatcgagaaaaccatctccaaagccaaaggcgagccccgagaaccacaggtgt
acacctgcccccatcccgggatgagctgaccaagaaccaggtcagcctgacctgcctggtcaa
aggcttctatcccagcgacatcgccgtggagtgaggagcaatgggcagccggagaacaactac
aagaccacgcctcccgtgttgactccgacggctccttcttctctacagcaagctcacgcgtgg
acaagagcaggtggcagcaggggaacgtcttctcatgctccgtgatgcatgaggctctgcacaa
ccactacacgcagagaagagcctctccctgtctccgggtaaatga

FIG. 3B

IFN α -Fc NUCLEOTIDE SEQUENCE (8 AMINO ACID LINKER)

atggccttgacctttgctttactggtggccctcctggtgctcagctgcaagtcaagctgctctg
tggcctgtgatctgcctcaaaccacagcctgggtagcaggaggaccttgatgctcctggcaca
gatgaggagaatctctcttttctcctgcttgaaggacagacatgactttggatttccccaggag
gagtttggcaaccagttccaaaaggctgaaaccatccctgtcctccatgagatgatccagcaga
tcttcaatctcttcagcacaaaggactcatctgctgcttgggatgagaccctcctagacaaatt
ctacactgaactctaccagcagctgaatgacctggaagcctgtgtgatacagggggtgggggtg
acagagactcccctgatgaaggaggactccattctggctgtgaggaaatacttccaaagaatca
ctctctatctgaaagagaagaaatacagcccttgtgcctgggaggttgtcagagcagaaatcat
gagatctttttctttgtcaacaaacttgcaagaaagtttaagaagtaaggaagaattcgccggc
gccgtgcggtcgacaaaactcacacatgcccaccgtgccagcacctgaactcctggggggac
cgctcagtcttctcttccccccaaaacccaaggacaccctcatgatctcccggaaccttgaggt
cacatgcgtggtggtggacgtgagccacgaagaccctgaggtcaagttcaactggtacgtggac
ggcgtggaggtgcataatgccaagacaaagccgaggaggagcagtacaacagcacgtaccgtc
tggtcagcgtcctcaccgtcctgcaccaggactggctgaatggcaaggagtacaagtgaaggt
ctccaacaaagccctcccagccccatcgagaaaaccatctccaaagccaaagggcagccccga
gaaccacaggtgtacacctgcccccatcccgggatgagctgaccaagaaccaggtcagcctga
cctgcctggtcaaaggcttctatcccagcgacatcgccgtggagtgggagagcaatgggcagcc
ggagaacaactacaagaccacgcctcccgtggtggactccgacggctccttcttctctacagc
aagctcaccgtggacaagagcaggtggcagcaggggaacgtcttctcatgctccgtgatgcatg
aggctctgcacaaccactacacgcagaagagcctctccctgtctccgggtaaataga

FIG. 3C

IFN α -Fc Δ LINKER NUCLEOTIDE SEQUENCE

atggccttgacctttgctttactggtggccctcctggtgctcagctgcaagtcaagctgctctg
tgggctgtgatctgcctcaaaccacagcctgggtagcaggaggaccttgatgctcctggcaca
gatgaggagaatctctcttttctcctgcttgaaggacagacatgactttggatttccccaggag
gagtttggcaaccagttccaaaaggctgaaaccatccctgtcctccatgagatgatccagcaga
tcttcaatctcttcagcacaaaggactcatctgctgcttgggatgagaccctcctagacaaatt
ctacactgaactctaccagcagctgaatgacctggaagcctgtgtgatacagggggtgggggtg
acagagactcccctgatgaaggaggactccattctggctgtgaggaaatacttccaaagaatca
ctctctatctgaaagagaagaaatacagcccttgtgcctgggagggtgtcagagcagaaatcat
gagatctttttctttgtcaacaaacttgcaagaaagtttaagaagtaaggaagacaaaactcac
acgtgcccgccgtgccagctccggaactgctgggcggaccgtcagttcttctcttcccccaa
aacccaaggacaccctcatgatctcccgaccctgaggtcacatgcgtggtggtggacgtgag
ccacgaagaccctgaggtcaagttcaactggtacgtggacggcgtggaggtgcataatgccaag
acaaagccgcgggaggagcagtacaacagcacgtaccgtgtggtcagcgtcctcacctgctgc
accaggactggctgaatggcaaggagtacaagtgaaggtctccaacaaagccctcccagcccc
catcgagaaaaccatctccaaagccaaagggcagccccgagaaccacaggtgtacaccctgccc
ccatcccgggatgagctgaccaagaaccaggtcagcctgacctgcctggtcaaaggcttctatc
ccagcgacatcgccgtggagtgaggagcaatgggcagccggagaaactacaagaccacgcc
tcccggtgttgactccgacggctccttcttctctacagcaagctcacctggacaagagcagg
tggcagcaggggaacgtcttctcatgctccgtgatgcatgaggctctgcacaaccactacacgc
agaagagcctctcctgtctccgggtaaatga

FIG. 3D

FLAGFc NUCLEOTIDE SEQUENCE

atggagacagacacactcctgctatgggtactgctgctctgggttccaggttccactggtgacg
actacaaggacgacgatgacaaggacaaaactcacacatgccaccgtgccagctccggaact
cctgggggggaccgtcagttcttcttccccccaaaacccaaggacaccctcatgatctcccg
accctgaggtcacatgcgtggtggtggacgtgagccacgaagaccctgaggtcaagttcaact
ggtacgtggacggcgtggaggtgcataatgccaagacaaagccgcgggaggagcagtacaacag
cacgtaccgtgtggtcagcgtcctcacctgctgaccaggactggctgaatggcaaggagtac
aagtgaaggtctccaacaaagccctcccagcccccatcgagaaaaccatctccaaagccaaag
ggcagccccgagaaccacaggtgtacaccctgcccccatcccgggatgagctgaccaagaacca
ggtcagcctgacctgcctggtcaaaggcttctatcccagcgacatcgccgtggagtgaggagc
aatgggcagccggagaacaactacaagaccacgcctcccgtgttgactccgacggctccttct
tctctacagcaagctcacctggacaagagcaggtggcagcaggggaacgtcttctcatgctc
cgtgatgcatgaggctctgcacaaccactacacgcagaagagcctctcctgtctccgggtaaa
tga

FIG. 3E

Epo-CCA-Fc NUCLEOTIDE SEQUENCE (K^b SIGNAL SEQUENCE UNDERLINED, ACIDIC COILED COIL IN BOLD)

atggtaccgtgcacgctgctcctgctgttgccggccgcccctggctccgactcagacccgcgccg
gctctagagccccaccacgcctcatctgtgacagccgagtcctgcagaggtacctcttgaggc
caaggagggcgagaatatcacgacgggctgtgctgaacactgcagcttgaatgagaatatcact
gtcccagacaccaaagttaattttctatgcctggaagaggatggaggtcgggagcaggccgtag
aagtctggcagggcctggccctgctgtcggaagctgtcctgcggggccaggccctgttggtcaa
ctcttcccagccgtgggagccccctgcagctgcatgtggataaagccgtcagtgcccttcgcagc
ctcaccactctgcttcgggctctgggagcccagaaggaagccatctcccctccagatgcggcct
cagctgctccactccgaacaatcactgctgacactttccgcaaactcttccgagtctactccaa
tttctccggggaaagctgaagctgtacacaggggaggcctgcaggaccggtgacagggaattc
ggtggtgagtaccaggccctggagaaggaggtggcccagctggaggccgagaaccaggccctgg
agaaggaggtggcccagctgggacacgaggggtggtggtcccgacccgagctgctgggcggacc
gtcagctcttctcttccccccaaaaccaaggacaccctcatgatctcccgacccctgaggtc
acatgcgtggtggtggacgtgagccacgaagaccctgaggtcaagttcaactggtacgtggacg
gcgtggaggtgcataatgccaagacaaagccgcgggaggagcagtacaacagcacgtaccgtgt
ggtcagcgtcctcacgctcctgcaccaggactggctgaatggcaaggagtacaagtgaaggctc
tccaacaaagccctcccagcccccatcgagaaaaccatctccaaagccaaagggcagccccgag
aaccacaggtgtacaccctgcccccatcccgggatgagctgaccaagaaccaggtcagcctgac
ctgcctggtcaaaggcttctatcccagcgacatcgccgtggagtgggagagcaatgggcagccg
gagaacaactacaagaccacgcctcccggtgttgactccgacggctccttcttctctacagca
agctcacccgtggacaagagcaggtggcagcaggggaacgtcttctcatgctccgtgatgcatga
ggctctgcacaaccactacacgcagaagagcctctcctgtctccgggtaaata

FIG. 3F

CCB-Fc NUCLEOTIDE SEQUENCE (SIGNAL SEQUENCE UNDERLINED, BASIC COILED COIL IN BOLD)

atggtaccgtgcacgctgctcctgctgttgccggccgcccctggctccgactcagacccgcgccg
gcgaattc**ggtggtgagtaccaggccctgaagaagaagggtggcccagctgaaggccaagaacca**
ggccctgaagaagaagggtggcccagctgaagcacaaggcgccgccccgccccagagctcctg
ggcggaccgtcagttcttcttccccccaaaaccaaggacaccctcatgatctcccgaccc
ctgaggtcacatgcgtggtggtggacgtgagccacgaagaccctgaggtcaagttcaactggta
cgtggacggcgtggaggtgcataatgccaagacaaagccgcgggaggagcagtacaacagcacg
taccgtgtggtcagcgtcctcacgctcctgcaccaggactggctgaatggcaaggagtacaagt
gcaaggcttccaacaaagccctcccagcccccatcgagaaaaccatctccaaagccaaagggca
gccccgagaaccacaggtgtacaccctgcccccatcccgggatgagctgaccaagaaccaggtc
agcctgacctgcctggtcaaaggcttctatcccagcgacatcgccgtggagtgggagagcaatg
ggcagccggagaacaactacaagaccacgcctcccggtgttgactccgacggctccttcttctc
ctacagcaagctcacccgtggacaagagcaggtggcagcaggggaacgtcttctcatgctccgtg
atgcatgaggctctgcacaaccactacacgcagaagagcctctcctgtctccgggtaaata

FIG. 3G

CysFc NUCLEOTIDE SEQUENCE (hIFN α SIGNAL SEQUENCE UNDERLINED)

atggccttgacctttgctttactggtggccctcctggtgctcagctgcaagtcaagctgctctg
tgggctgcccgcctgcccagctccggaactgctggcgacgctcagctcttcttcccccc
aaaacccaaggacaccctcatgatctcccgacccctgaggtcacatgctggtggtggacgtg
agccacgaagaccctgaggtcaagttcaactggtacgtggacggcgtggaggtgcataatgcc
agacaaagccgcgaggagcagtacaacagcacgtaccgtgtggtcagcgtcctcaccgtcct
gcaccaggactggctgaatggcaaggagtacaagtgaaggctccaacaagccctcccagcc
cccatcgagaaaaccatctccaaagccaaagggcagccccgagaaccacaggtgtacaccctgc
ccccatcccgggatgagctgaccaagaaccaggtcagcctgacctgctggtcaaaggcttcta
tcccagcgacatcgccgtggagtgggagagcaatggcgagccggagacaactacaagaccacg
cctcccggtgttgactccgacggctccttcttctctacagcaagctcaccgtggacaagagca
ggtggcagcaggggaacgtcttctcatgctccgtgatgcatgaggctctgcacaaccctacac
gcagaagagcctctccctgtctccgggtaaatga

FIG. 3HIFN α GS15 Fc NUCLEOTIDE SEQUENCE (SIGNAL SEQUENCE UNDERLINED):

atggccttgacctttgctttactggtggccctcctggtgctcagctgcaagtcaagctgctctg
tgggctgtgatctgcctcaaaccacagcctgggtagcaggaggaccttgatgctcctggcaca
gatgaggagaatctctcttttctcctgcttgaaggacagacatgactttggatttccccaggag
gagtttggaaccagttccaaaaggctgaaaccatccctgtcctccatgagatgatccagcaga
tcttcaatctcttcagcacaaggactcatctgctgcttgggatgagaccctcctagacaaatt
ctacactgaactctaccagcagctgaatgacctggaggcctgtgtgatacaggggggtgggggtg
acagagactcccctgatgaaggaggactccattctggctgtgaggaaatacttccaaagaatca
ctctctatctgaaagagaagaaatacagcccttgctgctgggaggttgctcagagcagaaatcat
gagatcttttctttgtcaacaaacttgcaagaaagtttacgtagtaaggaaggtggcgggcgga
tccggtggaggcggtccggcggtggaggagcgacaaaactcacacgtgcccgcctgcccag
ctccggaactgctggcgacgctcagctcttcttccccccaaaacccaaggacaccctcat
gatctcccgacccctgaggtcacatgctggtggtggacgtgagccacgaagaccctgaggtc
aagttcaactggtacgtggacggcgtggaggtgcataatgccaaagacaaagccgcgggaggagc
agtacaacagcacgtaccgtgtggtcagcgtcctcaccgtcctgcaccaggactggctgaatgg
caaggagtacaagtgaaggctccaacaagccctcccagccccatcgagaaaaccatctcc
aaagecaaagggcagccccgagaaccacaggtgtacaccctgcccccatcccgggatgagctga
ccaagaaccaggtcagcctgacctgctggtcaaaggcttctatcccagcgacatcgccgtgga
gtgggagagcaatggcgagccggagacaactacaagaccacgcctcccgtgttgactccgac
ggctccttcttctctacagcaagctcaccgtggacaagagcaggtggcagcaggggaacgtct
tctcatgctccgtgatgcatgaggctctgcacaaccctacacgcagaagagcctctccctgtc
tccgggtaaatga

FIG. 3I


EpoFc NUCLEOTIDE SEQUENCE (SIGNAL SEQUENCE UNDERLINED, LINKER IN BOLD)

atgggagtgacgaatgtcctgcctggctgtggcttctcctgtccctgctgtcgcctccctctgg
gcctcccagtcctgggcgccccaccacgcctcatctgtgacagccgagtcctggagaggtacct
cttgaggccaaggaggccgagaatatcacgacgggctgtgctgaacactgcagcttgaatgag
aatatcactgtcccagacaccaaagttaatttctatgcctggaagaggatggaggtcgggcagc
aggccgtagaagtctggcagggcctggccctgctgtcggaagctgtcctgcggggccaggccct
gttgggtcaactcttcccagccgtgggagcccctgcagctgcatgtggataaagccgtcagtggc
cttcgcagcctcaccactctgcttcgggctctgggagcccagaaggaagccatctcccctccag
atgcggcctcagctgtccactccgaacaatcactgtgacactttccgcaaactcttccgagt
ctactccaatttctcggggaaagctgaagctgtacacaggggaggcctgcagaacaggggac
agag**agttcgcggcgccgctgcggtc**gacaaaactcacacatgccaccgtgccagctccgg
aactcctgggcggaccgtcagtccttctcttccccccaaaacccaaggacaccctcatgatctc
ccggacccctgaggtcacatgcgtggtggtggacgtgagccacgaagaccctgaggtcaagttc
aactggtacgtggacggcgtggaggtgcataatgccaagacaaagccgcgggaggagcagtaca
acagcacgtaccgtgtggtcagcgtcctcaccgtcctgcaccaggactggtgaatggcaagga
gtacaagtgcaaggtctccaacaaagccctcccagccccatcgagaaaaccatctccaaagcc
aaagggcagccccgagaaccacaggtgtacaccctgccccatcccgggatgagctgaccaaga
accaggtcagcctgacctgcctggtcaaaggcttctatcccagcgacatcgccgtggagtgagg
gagcaatgggcagccggagaacaactacaagaccacgcctcccgtgttgactccgacggctcc
ttcttctctacagcaagctcaccgtggacaagagcaggtggcagcaggggaacgtcttctcat
gctccgtgatgcatgaggctctgcacaaccactacacgcagaagagcctctccctgtctccggg
taaata

FIG. 3J

FIG. 4

VARIOUS WAYS TO FORM MONOMER/DIMER HYBRIDS THROUGH NATIVE LIGATION
NOTE THAT PEPTIDE COULD BE SUBSTITUTED WITH ANY OTHER MOLECULE, DNA, ETC.



Amino acid sequence of Fc-MESNA (produced in pTWIN1 vector from NEB; when Fc-Intein-CED is eluted from chitin beads with MESNA, produces the following protein with a C-terminal thioester on the final Phe residue)

1	MGIEGRGAAA	VDTSHTCPPC	PAPELLGGPS	VFLFPPKPKD	TLMISRTPEV
51	TCVVVDVSHE	DPEVKFNWYV	DGVEVHNAKT	KPREEQYNST	YRVVSVLTVL
101	HQDWLNGKEY	KCKVSNKALP	APIEKTISKA	KGQPREPQVY	TLPPSRDELT
151	KNQVSLTCLV	KGFYPSDIAV	EWESNGQPEN	NYKTTTPVLD	SDGSFFLYSK
201	LTVDKSRWQQ	GNVFSCSVMH	EALHNHYTQK	SLSLSPGF	

FIG. 5A

Nucleotide sequence of Fc CDS in pTWIN1 (the final F residue, ttt, directly abuts the Mxe GyrA intein CDS in pTWIN1)

atgggcattgaaggcagaggcgccgctgcggtcgatactagtcacacatgccaccgtgcccag
cacctgaactccggggggaccgtcagctcttctcttccccccaaaacccaaggacaccctcat
gatctcccggaccctgaggtcacatgcgtggtggtggacgtgagccacgaagaccctgaggtc
aagttcaactggtacgtggacggcgtggaggtgcataatgccaagacaaagccgcgggaggagc
agtacaacagcacgtaccgtgtggtcagcgtcctcaccgtcctgcaccaggactggctgaatgg
caaggagtacaagtgaaggtctccaacaaagccctcccagcccccatcgagaaaaccatctcc
aaagccaaagggcagccccgagaaccacaggtgtacaccctgcccccatcccgggatgagctga
ccaagaaccaggtcagcctgacctgcctggtcaaaggcttctatcccagcgacatcgccgtgga
gtgggagagcaatgggcagccggagaacaactacaagaccacgcctcccgtgttgactccgac
ggctccttcttctctacagcaagctcaccgtggacaagagcaggtggcagcaggggaacgtct
tctcatgctccgtgatgcatgaggctctgcacaaccactacacgcagaagagtctctccctgtc
tccgggtttt

FIG. 5B




FIG. 6




FIG. 7




FIG. 8

C_{max} AFTER SINGLE ORAL DOSE OF FACTOR IX-Fc HOMODIMER
V. FACTOR IX-Fc:FLAG-Fc MONOMER-DIMER HYBRID
INTO 10 DAY OLD SPRAGUE DAWLEY RATS




FIG. 9




FIG. 10




FIG. 11





FIG. 12

**FIG. 13**




FIG. 14

PHARMACOKINETICS OF IFN β -008 (IFN β -Fc WT DIMER) AND
IFN β -009/Fc-014 (IFN β -Fc N297A MONOMER) IN CYNOMOLGUS
MONKEY SERUM AFTER A SINGLE PULMONARY DOSE OF 20 μ g/kg




FIG. 15




FIG. 16

IFN β -Fc NUCLEOTIDE SEQUENCE (SIGNAL PEPTIDE UNDERLINED)

atgaccaacaagtgtctcctccaaattgctctcctggttgcttctccactacagctctttcca
 tgagctacaacttgcttggttcctacaaagaagcagcaattttcagtgtcagaagctcctgtg
 gcaattgaatgggaggcttgaatattgcctcaaggacaggatgaactttgacatccctgaggag
 attaagcagctgcagcagttccagaaggaggacgccgcattgaccatctatgagatgctccaga
 acatcttttgctattttcagacaagattcatctagcactggctggaatgagactattgttgagaa
 cctcctggctaattgtctatcatcagataaaccatctgaagacagtcctggaagaaaaactggag
 aaagaagatttcaccaggggaaaactcatgagcagctctgcacctgaaaagatattatgggagga
 ttctgcattacctgaaggccaaggagtacagtcactgtgcctggaccatagtcagagtggaaat
 cctaaggaacttttacttcattaacagacttacaggttacctccgaaacgagttcgccggcgcc
 gctgcggtcgacaaaactcacacatgccaccgtgccagctccggaactcctgggaggaccgt
 cagtcttcctcttccccccaaaaccaaggacacctcatgatctccggaccctgaggtcac
 atgcgtggtggaggcgtgagccacgaagacctgaggtcaagttcaactggtacgtggacggc
 gtggagggtgcataatgccaaagacaaagccgcgggaggagcagtacaacagcacgtaccgtgtgg
 tcagcgtcctcacgctcctgcaccaggactggctgaatggcaaggagtacaagtgaaggtctc
 caacaaagccctcccagcccccatcgagaaaaccatctccaaagccaaagggcagccccgagaa
 ccacaggtgtacacctgcccccatcccggtgagctgaccaagaaccaggtcagcctgacct
 gcctggtcaaaggcttctatcccagcgacatcgccgtggagtgggagagcaatgggcagccgga
 gaacaactacaagaccagcctcccggtgttgactccgacggctccttcttctctacagcaag
 ctaccgtggacaagagcaggtggcagcaggggaacgtcttctcatgctccgtgatgcatgagg
 ctctgcacaaccactacacgcagaagagcctctcctgtctccgggtaaataga

FIG. 17AIFN β -Fc AMINO ACID SEQUENCE (SIGNAL SEQUENCE UNDERLINED, LINKER SEQUENCE IN BOLD, N297 IN BOLD UNDERLINED).

1 MTNKCLLOIA LLLCFSTTAL SMSYNLLGFL QRSSNFQCQK LLWQLNGRLE
 51 YCLKDRMNFD IPEEIKQLQQ FQKEDAALTI YEMLQNI FAI FRQDSSSTGW
 101 NETIVENLLA NVYHQINHLK TVLEEKLEKE DFTRGKLMSS LHLKRYYGRI
 151 LHYLKAKEYS HCAWTIVRVE ILRNFYFINR LTGYLRNEFA **GAAAVDKTHT**
 201 CPPCPAPELL GGPSVFLFPP KPKDTLMISR TPEVTCVVVD VSHEDPEVKF
 251 NWYVDGVEVH NAKTKPREEQ **YN**STYRVVSV LTVLHQDWLN GKEYKCKVSN
 301 KALPAPIEKT ISKAKGQPRE PQVYTLPPSR DELTKNQVSL TCLVKGFYPS
 351 DIAVEWESNG QPENNYK TTP PVLDS DGSFF LYSKLTVDKS RWQQGNVFSC
 401 SVMHEALHNH YTQKSLSLSP GK

FIG. 17B

YTSLIHSLIEESQNQQEKNEQELLELDKWASLWNWF (SEQ ID NO: 1).

FIG. 18A

NNLRAIEAQQHLLQLTVWGIKQLQARILAVERYLKDQ (SEQ ID NO: 2)

FIG. 18B

WQEWQKITALLEQAQIQQEKNEYELQKLDKWASLWEWF (SEQ ID NO: 3)

FIG. 18C

SEQUENCE LISTING

<110> SYNTONIX PHARMACEUTICALS, INC.
PETERS, ROBERT T.
MEZO, ADAM R.
RIVERA, DANIEL S.
BITONTI, ALAN J.

<120> IMMUNOGLOBULIN CHIMERIC MONOMER-DIMER HYBRIDS

<130> 08945.0007-01304

<140> Not Yet Assigned

<141> Herewith

<150> 11/029,003

<151> 2005-01-05

<160> 91

<170> PatentIn Ver. 3.2

<210> 1

<211> 36

<212> PRT

<213> Human immunodeficiency virus

<400> 1

Tyr	Thr	Ser	Leu	Ile	His	Ser	Leu	Ile	Glu	Glu	Ser	Gln	Asn	Gln	Gln
1				5					10					15	

Glu	Lys	Asn	Glu	Gln	Glu	Leu	Leu	Glu	Leu	Asp	Lys	Trp	Ala	Ser	Leu
		20						25					30		

Trp	Asn	Trp	Phe
		35	

<210> 2

<211> 37

<212> PRT

<213> Human immunodeficiency virus

<400> 2

Asn	Asn	Leu	Arg	Ala	Ile	Glu	Ala	Gln	Gln	His	Leu	Leu	Gln	Leu	Thr
1				5					10					15	

Val	Trp	Gly	Ile	Lys	Gln	Leu	Gln	Ala	Arg	Ile	Leu	Ala	Val	Glu	Arg
		20						25					30		

Tyr	Leu	Lys	Asp	Gln
			35	

<210> 3

<211> 39

<212> PRT

<213> Human immunodeficiency virus

<400> 3

Trp Gln Glu Trp Glu Gln Lys Ile Thr Ala Leu Leu Glu Gln Ala Gln
1 5 10 15

Ile Gln Gln Glu Lys Asn Glu Tyr Glu Leu Gln Lys Leu Asp Lys Trp
20 25 30

Ala Ser Leu Trp Glu Trp Phe
35

<210> 4

<211> 238

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
construct

<400> 4

Met Gly Ile Glu Gly Arg Gly Ala Ala Ala Val Asp Thr Ser His Thr
1 5 10 15

Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser Val Phe
20 25 30

Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro
35 40 45

Glu Val Thr Cys Val Val Val Asp Val Ser His Glu Asp Pro Glu Val
50 55 60

Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr
65 70 75 80

Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser Val
85 90 95

Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys
100 105 110

Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser
115 120 125

Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro
130 135 140

Ser Arg Asp Glu Leu Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val
145 150 155 160

Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly
165 170 175

Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp

3/41

180	185	190
Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp		
195	200	205
Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu His		
210	215	220
Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Phe		
225	230	235

<210> 5

<211> 714

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic construct

<400> 5

```

atggggcattg aaggcagagg cgccgctgcg gtcgatacta gtcacacatg cccaccgtgc 60
ccagcacctg aactcctggg gggaccgtca gtcttctctt tcccccaaa acccaaggac 120
accctcatga tctcccgac ccctgaggtc acatgcgtgg tgggtggacgt gagccacgaa 180
gaccctgagg tcaagttcaa ctggtacgtg gacggcgtgg aggtgcataa tgccaagaca 240
aagccgcggg aggagcagta caacagcacg taccgtgtgg tcagcgtcct caccgtcctg 300
caccaggact ggctgaatgg caaggagtac aagtgcaagg tctccaacaa agccctccca 360
gcccccatcg agaaaaccat ctccaaagcc aaagggcagc cccgagaacc acaggtgtac 420
accctgcccc catcccgga tgagctgacc aagaaccagg tcagcctgac ctgcctggtc 480
aaaggcttct atcccagcga catcgccgtg gagtgggaga gcaatgggca gccggagAAC 540
aactacaaga ccacgcctcc cgtgttgac tccgacggct ccttcttct ctacagcaag 600
ctcaccgtgg acaagagcag gtggcagcag gggaacgtct tctcatgctc cgtgatgcat 660
gaggctctgc acaaccacta cagcagaag agtctctccc tgtctccggg tttt 714

```

<210> 6

<211> 671

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic construct

<400> 6

Met Val Ser Gln Ala Leu Arg Leu Leu Cys Leu Leu Leu Gly Leu Gln
1 5 10 15
Gly Cys Leu Ala Ala Val Phe Val Thr Gln Glu Glu Ala His Gly Val
20 25 30
Leu His Arg Arg Arg Arg Ala Asn Ala Phe Leu Glu Glu Leu Arg Pro
35 40 45
Gly Ser Leu Glu Arg Glu Cys Lys Glu Glu Gln Cys Ser Phe Glu Glu
50 55 60

Ala Arg Glu Ile Phe Lys Asp Ala Glu Arg Thr Lys Leu Phe Trp Ile
 65 70 75 80
 Ser Tyr Ser Asp Gly Asp Gln Cys Ala Ser Ser Pro Cys Gln Asn Gly
 85 90 95
 Gly Ser Cys Lys Asp Gln Leu Gln Ser Tyr Ile Cys Phe Cys Leu Pro
 100 105 110
 Ala Phe Glu Gly Arg Asn Cys Glu Thr His Lys Asp Asp Gln Leu Ile
 115 120 125
 Cys Val Asn Glu Asn Gly Gly Cys Glu Gln Tyr Cys Ser Asp His Thr
 130 135 140
 Gly Thr Lys Arg Ser Cys Arg Cys His Glu Gly Tyr Ser Leu Leu Ala
 145 150 155 160
 Asp Gly Val Ser Cys Thr Pro Thr Val Glu Tyr Pro Cys Gly Lys Ile
 165 170 175
 Pro Ile Leu Glu Lys Arg Asn Ala Ser Lys Pro Gln Gly Arg Ile Val
 180 185 190
 Gly Gly Lys Val Cys Pro Lys Gly Glu Cys Pro Trp Gln Val Leu Leu
 195 200 205
 Leu Val Asn Gly Ala Gln Leu Cys Gly Gly Thr Leu Ile Asn Thr Ile
 210 215 220
 Trp Val Val Ser Ala Ala His Cys Phe Asp Lys Ile Lys Asn Trp Arg
 225 230 235 240
 Asn Leu Ile Ala Val Leu Gly Glu His Asp Leu Ser Glu His Asp Gly
 245 250 255
 Asp Glu Gln Ser Arg Arg Val Ala Gln Val Ile Ile Pro Ser Thr Tyr
 260 265 270
 Val Pro Gly Thr Thr Asn His Asp Ile Ala Leu Leu Arg Leu His Gln
 275 280 285
 Pro Val Val Leu Thr Asp His Val Val Pro Leu Cys Leu Pro Glu Arg
 290 295 300
 Thr Phe Ser Glu Arg Thr Leu Ala Phe Val Arg Phe Ser Leu Val Ser
 305 310 315 320
 Gly Trp Gly Gln Leu Leu Asp Arg Gly Ala Thr Ala Leu Glu Leu Met
 325 330 335
 Val Leu Asn Val Pro Arg Leu Met Thr Gln Asp Cys Leu Gln Gln Ser
 340 345 350
 Arg Lys Val Gly Asp Ser Pro Asn Ile Thr Glu Tyr Met Phe Cys Ala
 355 360 365

Gly Tyr Ser Asp Gly Ser Lys Asp Ser Cys Lys Gly Asp Ser Gly Gly
 370 375 380
 Pro His Ala Thr His Tyr Arg Gly Thr Trp Tyr Leu Thr Gly Ile Val
 385 390 395 400
 Ser Trp Gly Gln Gly Cys Ala Thr Val Gly His Phe Gly Val Tyr Thr
 405 410 415
 Arg Val Ser Gln Tyr Ile Glu Trp Leu Gln Lys Leu Met Arg Ser Glu
 420 425 430
 Pro Arg Pro Gly Val Leu Leu Arg Ala Pro Phe Pro Asp Lys Thr His
 435 440 445
 Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser Val
 450 455 460
 Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr
 465 470 475 480
 Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu Asp Pro Glu
 485 490 495
 Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala Lys
 500 505 510
 Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser
 515 520 525
 Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys
 530 535 540
 Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile
 545 550 555 560
 Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro
 565 570 575
 Pro Ser Arg Asp Glu Leu Thr Lys Asn Gln Val Ser Leu Thr Cys Leu
 580 585 590
 Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn
 595 600 605
 Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser
 610 615 620
 Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg
 625 630 635 640
 Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu
 645 650 655
 His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys

660

665

670

<210> 7
<211> 2016
<212> DNA
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
construct

<400> 7

```
atggtctctccc aggccctcag gctcctctgc cttctgcttg ggcttcaggg ctgcctggct 60
gcagtcttcg taaccagga ggaagccac ggcgtcctgc accggcgccg gcgcgccaac 120
gcgttcctgg aggagctgcy gccgggctcc ctggagaggg agtgcaagga ggagcagtgc 180
tccttcgagg agggccggga gatcttcaag gacgcggaga ggacgaagct gttctggatt 240
tcttacagt atggggacca gtgtgcctca agtccatgcc agaatggggg ctctgcaag 300
gaccagctcc agtcctatat ctgcttctgc ctccctgcct tcgagggccg gaactgtgag 360
acgcacaagg atgaccagct gatctgtgtg aacgagaacg gcggctgtga gcagtactgc 420
agtgaccaca cgggcaccaa gcgctcctgt cggtgccaac aggggtactc tctgctggca 480
gacgggggtgt cctgcacacc cacagttgaa tatccatgtg gaaaaatacc tattctagaa 540
aaaagaaatg ccagcaaacc ccaaggccga attgtggggg gcaaggtgtg ccccaaaggg 600
gagtgtccat ggaggtcct gttgttggtg aatggagctc agttgtgtgg ggggaccctg 660
atcaacacca tctgggtggt ctccgcggcc cactgtttcg acaaaatcaa gaactggagg 720
aacctgatcg cgtgtgtgg cgagcacgac ctacgcgagc acgacgggga tgagcagagc 780
cggcggtgg cgaggtcat catcccagc acgtacgtcc cgggcaccac caaccagac 840
atcgcgctgc tccgctgca ccagccgtg gtctcactg accatgtggt gcccctctgc 900
ctgcccgaac ggagcttctc tgagaggagc ctggccttcg tgcgttctc attggtcagc 960
ggctggggcc agctgtgga ccgtggcgcc acggccctgg agctcatggt cctcaacgtg 1020
ccccggctga tgaccagga ctgcctgcag cagtcaagga aggtgggaga ctcccaaat 1080
atcacggagt acatgttctg tgccggctac tggatggca gcaaggactc ctgcaagggg 1140
gacagtggag gccacatgc caccactac cggggcacgt ggtacctgac gggcatcgtc 1200
agctggggcc agggctgcgc aaccgtgggc cactttgggg tgtacaccag ggtctcccag 1260
tacatcgagt ggctgcaaaa gctcatgcgc tcagagccac gccaggagt cctcctgcga 1320
gccccatttc ccgacaaaac tcacacgtgc ccgcctgccc cagctccgga actgctgggc 1380
ggaccgtcag tcttcctctt cccccaaaa cccaaggaca ccctcatgat ctcccgacc 1440
cctgaggtca catgcgtggt ggtggacgtg agccacgaag accctgaggt caagttcaac 1500
tggtacgtgg acggcgtgga ggtgcataat gccaaagaca agccgcggga ggagcagtac 1560
aacgacagc accgtgtggt cagcgtcctc accgtcctgc accaggactg gctgaatggc 1620
aaggagtaca agtgcaaggt ctccaacaaa gccctcccag ccccatcga gaaaaccatc 1680
tccaaagcca aagggcagcc ccgagaacca caggtgtaca ccctgcccc atcccgat 1740
gagctgacca agaaccaggt cagcctgacc tgctggtca aaggcttcta tcccagcagc 1800
atcgccgtgg agtgggagag caatgggcag ccggagaaca actacaagac cagcctccc 1860
gtgttgact ccgacggctc cttcttctc tacagcaagc tcaccgtgga caagagcagg 1920
tggcagcagg ggaacgtctt ctcatgctcc gtgatgcatg aggtcttgca caaccactac 1980
acgcagaaga gcctctccct gtctccgggt aatatga 2016
```

<210> 8
<211> 696
<212> PRT
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
construct

<400> 8

```

Met Gln Arg Val Asn Met Ile Met Ala Glu Ser Pro Gly Leu Ile Thr
 1           5           10           15

Ile Cys Leu Leu Gly Tyr Leu Leu Ser Ala Glu Cys Thr Val Phe Leu
      20           25           30

Asp His Glu Asn Ala Asn Lys Ile Leu Asn Arg Pro Lys Arg Tyr Asn
      35           40           45

Ser Gly Lys Leu Glu Glu Phe Val Gln Gly Asn Leu Glu Arg Glu Cys
      50           55           60

Met Glu Glu Lys Cys Ser Phe Glu Glu Ala Arg Glu Val Phe Glu Asn
      65           70           75           80

Thr Glu Arg Thr Thr Glu Phe Trp Lys Gln Tyr Val Asp Gly Asp Gln
      85           90           95

Cys Glu Ser Asn Pro Cys Leu Asn Gly Gly Ser Cys Lys Asp Asp Ile
      100          105          110

Asn Ser Tyr Glu Cys Trp Cys Pro Phe Gly Phe Glu Gly Lys Asn Cys
      115          120          125

Glu Leu Asp Val Thr Cys Asn Ile Lys Asn Gly Arg Cys Glu Gln Phe
      130          135          140

Cys Lys Asn Ser Ala Asp Asn Lys Val Val Cys Ser Cys Thr Glu Gly
      145          150          155          160

Tyr Arg Leu Ala Glu Asn Gln Lys Ser Cys Glu Pro Ala Val Pro Phe
      165          170          175

Pro Cys Gly Arg Val Ser Val Ser Gln Thr Ser Lys Leu Thr Arg Ala
      180          185          190

Glu Thr Val Phe Pro Asp Val Asp Tyr Val Asn Ser Thr Glu Ala Glu
      195          200          205

Thr Ile Leu Asp Asn Ile Thr Gln Ser Thr Gln Ser Phe Asn Asp Phe
      210          215          220

Thr Arg Val Val Gly Gly Glu Asp Ala Lys Pro Gly Gln Phe Pro Trp
      225          230          235          240

Gln Val Val Leu Asn Gly Lys Val Asp Ala Phe Cys Gly Gly Ser Ile
      245          250          255

Val Asn Glu Lys Trp Ile Val Thr Ala Ala His Cys Val Glu Thr Gly
      260          265          270

Val Lys Ile Thr Val Val Ala Gly Glu His Asn Ile Glu Glu Thr Glu
      275          280          285

His Thr Glu Gln Lys Arg Asn Val Ile Arg Ile Ile Pro His His Asn

```

290	295	300
Tyr Asn Ala Ala Ile Asn Lys Tyr Asn His Asp Ile Ala Leu Leu Glu 305 310 315 320		
Leu Asp Glu Pro Leu Val Leu Asn Ser Tyr Val Thr Pro Ile Cys Ile 325 330 335		
Ala Asp Lys Glu Tyr Thr Asn Ile Phe Leu Lys Phe Gly Ser Gly Tyr 340 345 350		
Val Ser Gly Trp Gly Arg Val Phe His Lys Gly Arg Ser Ala Leu Val 355 360 365		
Leu Gln Tyr Leu Arg Val Pro Leu Val Asp Arg Ala Thr Cys Leu Arg 370 375 380		
Ser Thr Lys Phe Thr Ile Tyr Asn Asn Met Phe Cys Ala Gly Phe His 385 390 395 400		
Glu Gly Gly Arg Asp Ser Cys Gln Gly Asp Ser Gly Gly Pro His Val 405 410 415		
Thr Glu Val Glu Gly Thr Ser Phe Leu Thr Gly Ile Ile Ser Trp Gly 420 425 430		
Glu Glu Cys Ala Met Lys Gly Lys Tyr Gly Ile Tyr Thr Lys Val Ser 435 440 445		
Arg Tyr Val Asn Trp Ile Lys Glu Lys Thr Lys Leu Thr Glu Phe Ala 450 455 460		
Gly Ala Ala Ala Val Asp Lys Thr His Thr Cys Pro Pro Cys Pro Ala 465 470 475 480		
Pro Glu Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro 485 490 495		
Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val 500 505 510		
Val Asp Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val 515 520 525		
Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln 530 535 540		
Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln 545 550 555 560		
Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala 565 570 575		
Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro 580 585 590		
Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Asp Glu Leu Thr		

595 600 605

Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser
610 615 620

Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr
625 630 635 640

Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr
645 650 655

Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe
660 665 670

Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys
675 680 685

Ser Leu Ser Leu Ser Pro Gly Lys
690 695

<210> 9

<211> 2091

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
construct

<400> 9 -

atgcagcgcg tgaacatgat catggcagaa tcaccaggcc tcatcaccat ctgcctttta 60
ggatatctac tcagtgtgta atgtacagtt tttcttgatc atgaaaacgc caacaaaatt 120
ctgaatcggc caaagaggta taattcaggt aaattggaag agtttgttca agggaaacct 180
gagagagaat gtatggaaga aaagtgtagt tttgaagaag cagagaagt ttttgaaaac 240
actgaaagaa caactgaatt ttggaagcag tatgttgatg gagatcagtg tgagtccaat 300
ccatgtttta atggcggcag ttgcaaggat gacattaatt cctatgaatg ttggtgtccc 360
tttggttttg aaggaaagaa ctgtgaatta gatgtaacat gtaacattaa gaatggcaga 420
tgcgagcagt tttgtaaaaa tagtgctgat aacaagggtg tttgctcctg tactgaggga 480
tatcgacttg cagaaaacca gaagtcctgt gaaccagcag tgccatttcc atgtggaaga 540
gtttctgttt cacaaacttc taagctcacc cgtgctgaga ctgtttttcc tgatgtggac 600
tatgtaaatt ctactgaagc tgaaccatt ttggataaca tctactcaaag caccatca 660
tttaatgact tctactcgggt tgttggtgga gaagatgcca aaccagggtca attcccttgg 720
caggttggtt tgaatggtaa agttgatgca ttctgtggag gctctatcgt taatgaaaaa 780
tggattgtaa ctgctgccc aatgtgtgaa actggtgtta aaattacagt tgtcgcagg 840
gaacataata ttgaggagac agaacataca gagcaaaagc gaaatgtgat tcgaattatt 900
cctcaccaca actacaatgc agctattaat aagtacaacc atgacattgc cttctgga 960
ctggacgaac ccttagtgct aaacagctac gttacaccta tttgcattgc tgacaaggaa 1020
tacacgaaca tcttcctcaa atttggatct ggctatgtaa gtggctgggg aagagtcctc 1080
cacaaagga gatcagcttt agttcttcag taccttagag ttccacttgt tgaccgagcc 1140
acatgtcttc gatctacaaa gtccaccatc tataacaaca tgttctgtgc tggcttccat 1200
gaaggaggta gagattcatg tcaaggagat agtgggggac cccatgttac tgaagtggaa 1260
gggaccagtt tcttaactgg aattattagc tggggtgaag agtgtgcaat gaaaggcaaa 1320
tatggaatat ataccaaggt atcccggat gtcaactgga ttaaggaaaa acaaagctc 1380
actgaattcg ccggcgccgc tgcggtcgac aaaactcaca catgccacc gtgcccagca 1440
cctgaactcc tggggggacc gtcagtcttc ctcttcccc caaaacccaa ggacaccctc 1500
atgatctccc ggaccctga ggtcacatgc gtggtggtgg acgtgagcca cgaagacctt 1560

```

gaggtcaagt tcaactggta cgtggacggc gtggaggtgc ataatgcaa gacaaagccg 1620
cgggaggagc agtacaacag cacgtaccgt gtggtcagcg tcctcaccgt cctgcaccag 1680
gactggctga atggcaagga gtacaagtgc aaggtctcca acaaagccct ccagccccc 1740
atcgagaaaa ccatctccaa agccaaaggg cagccccgag aaccacaggt gtacaccctg 1800
cccccatccc gggatgagct gaccaagaac caggtcagcc tgacctgcct ggtcaaaggc 1860
ttctatccca gcgacatcgc cgtggagtgg gagagcaatg ggcagccgga gaacaactac 1920
aagaccacgc ctcccgtgtt ggactccgac ggtccttctt tcctctacag caagctcacc 1980
gtggacaaga gcaggtggca gcaggggaac gtcttctcat gctccgtgat gcatgagggt 2040
ctgcacaacc actacacgca gaagagcctc tccctgtctc cgggtaaatg a 2091

```

<210> 10

<211> 423

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
construct

<400> 10

```

Met Ala Leu Thr Phe Ala Leu Leu Val Ala Leu Leu Val Leu Ser Cys
  1              5              10             15

Lys Ser Ser Cys Ser Val Gly Cys Asp Leu Pro Gln Thr His Ser Leu
      20              25              30

Gly Ser Arg Arg Thr Leu Met Leu Leu Ala Gln Met Arg Arg Ile Ser
      35              40              45

Leu Phe Ser Cys Leu Lys Asp Arg His Asp Phe Gly Phe Pro Gln Glu
      50              55              60

Glu Phe Gly Asn Gln Phe Gln Lys Ala Glu Thr Ile Pro Val Leu His
      65              70              75             80

Glu Met Ile Gln Gln Ile Phe Asn Leu Phe Ser Thr Lys Asp Ser Ser
      85              90              95

Ala Ala Trp Asp Glu Thr Leu Leu Asp Lys Phe Tyr Thr Glu Leu Tyr
      100             105             110

Gln Gln Leu Asn Asp Leu Glu Ala Cys Val Ile Gln Gly Val Gly Val
      115             120             125

Thr Glu Thr Pro Leu Met Lys Glu Asp Ser Ile Leu Ala Val Arg Lys
      130             135             140

Tyr Phe Gln Arg Ile Thr Leu Tyr Leu Lys Glu Lys Lys Tyr Ser Pro
      145             150             155             160

Cys Ala Trp Glu Val Val Arg Ala Glu Ile Met Arg Ser Phe Ser Leu
      165             170             175

Ser Thr Asn Leu Gln Glu Ser Leu Arg Ser Lys Glu Glu Phe Ala Gly
      180             185             190

```

11/41

Ala Ala Ala Val Asp Lys Thr His Thr Cys Pro Pro Cys Pro Ala Pro
 195 200 205
 Glu Leu Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys
 210 215 220
 Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val
 225 230 235 240
 Asp Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp
 245 250 255
 Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr
 260 265 270
 Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp
 275 280 285
 Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu
 290 295 300
 Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg
 305 310 315 320
 Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Asp Glu Leu Thr Lys
 325 330 335
 Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp
 340 345 350
 Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys
 355 360 365
 Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser
 370 375 380
 Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser
 385 390 395 400
 Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser
 405 410 415
 Leu Ser Leu Ser Pro Gly Lys
 420

<210> 11

<211> 1272

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
 construct

<400> 11

atggccttga cctttgcttt actggtggcc ctctggtgc tcagctgcaa gtcaagctgc 60

```

tctgtgggct gtgatctgcc tcaaaccac agcctgggta gcaggaggac cttgatgctc 120
ctggcacaga tgaggagaat ctctcttttc tcctgcttga aggacagaca tgacttttga 180
tttccccagg aggagtttgg caaccagttc caaaaggctg aaaccatccc tgtcctccat 240
gagatgatcc agcagatctt caatctcttc agcacaagg actcatctgc tgccttgggat 300
gagaccctcc tagacaaatt ctacactgaa ctctaccagc agctgaatga cctggaagcc 360
tgtgtgatac aggggggtggg ggtgacagag actcccctga tgaaggagga ctccattctg 420
gctgtgagga aatacttcca aagaatcact ctctatctga aagagaagaa atacagccct 480
tgtgcctggg aggttgtcag agcagaaatc atgagatctt tttctttgtc aacaaacttg 540
caagaaagtt taagaagtaa ggaagaattc gccggcgccg ctgcggtcga caaaactcac 600
acatgcccac cgtgcccagc acctgaactc ctggggggac cgtcagtctt cctcttcccc 660
ccaaaaccca aggacaccct catgatctcc cggaccctcg aggtcacatg cgtggtggtg 720
gacgtgagcc acgaagaccc tgaggtcaag ttcaactggt acgtggacgg cgtggagggtg 780
cataatgcca agacaaagcc gcgggaggag cagtacaaca gcacgtaccg tgtggtcagc 840
gtcctcaccc tcctgcacca ggactggctg aatggcaagg agtacaagtg caaggtctcc 900
aacaaagccc tcccagcccc catcgagaaa accatctcca aagccaaagg gcagccccga 960
gaaccacagg tgtacaccct gccccatcc cgggatgagc tgaccaagaa ccaggtcagc 1020
ctgacctgcc tgggtcaaagg cttctatccc agcgacatcg ccgtggagtg ggagagcaat 1080
gggcagccgg agaacaacta caagaccacg cctcccgtgt tggactccga cggctccttc 1140
ttcctctaca gcaagctcac cgtggacaag agcaggtggc agcaggggaa cgtcttctca 1200
tgctccgtga tgcattgaggc tctgcacaac cactacacgc agaagagcct ctccctgtct 1260
ccgggtaaat ga 1272

```

<210> 12

<211> 415

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic construct

<400> 12

```

Met Ala Leu Thr Phe Ala Leu Leu Val Ala Leu Leu Val Leu Ser Cys
  1                      5                      10                      15

```

```

Lys Ser Ser Cys Ser Val Gly Cys Asp Leu Pro Gln Thr His Ser Leu
          20                      25                      30

```

```

Gly Ser Arg Arg Thr Leu Met Leu Leu Ala Gln Met Arg Arg Ile Ser
          35                      40                      45

```

```

Leu Phe Ser Cys Leu Lys Asp Arg His Asp Phe Gly Phe Pro Gln Glu
          50                      55                      60

```

```

Glu Phe Gly Asn Gln Phe Gln Lys Ala Glu Thr Ile Pro Val Leu His
          65                      70                      75                      80

```

```

Glu Met Ile Gln Gln Ile Phe Asn Leu Phe Ser Thr Lys Asp Ser Ser
          85                      90                      95

```

```

Ala Ala Trp Asp Glu Thr Leu Leu Asp Lys Phe Tyr Thr Glu Leu Tyr
          100                      105                      110

```

```

Gln Gln Leu Asn Asp Leu Glu Ala Cys Val Ile Gln Gly Val Gly Val
          115                      120                      125

```

Thr Glu Thr Pro Leu Met Lys Glu Asp Ser Ile Leu Ala Val Arg Lys
 130 135 140
 Tyr Phe Gln Arg Ile Thr Leu Tyr Leu Lys Glu Lys Lys Tyr Ser Pro
 145 150 155 160
 Cys Ala Trp Glu Val Val Arg Ala Glu Ile Met Arg Ser Phe Ser Leu
 165 170 175
 Ser Thr Asn Leu Gln Glu Ser Leu Arg Ser Lys Glu Asp Lys Thr His
 180 185 190
 Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser Val
 195 200 205
 Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr
 210 215 220
 Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu Asp Pro Glu
 225 230 235 240
 Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala Lys
 245 250 255
 Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser
 260 265 270
 Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys
 275 280 285
 Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile
 290 295 300
 Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro
 305 310 315 320
 Pro Ser Arg Asp Glu Leu Thr Lys Asn Gln Val Ser Leu Thr Cys Leu
 325 330 335
 Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn
 340 345 350
 Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser
 355 360 365
 Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg
 370 375 380
 Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu
 385 390 395 400
 His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys
 405 410 415

<210> 13

<211> 1248

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic construct

<400> 13

```

atggccttga cctttgcttt actggtggcc ctctggtgct tcagctgcaa gtcaagctgc 60
tctgtgggct gtgatctgcc tcaaaccac agcctgggta gcaggaggac cttgatgctc 120
ctggcacaga tgaggagaat ctctcttttc tcctgcttga aggacagaca tgactttgga 180
tttccccagg aggagtttgg caaccagttc caaaaggctg aaaccatccc tgtcctccat 240
gagatgatcc agcagatctt caatctcttc agcacaaagg actcatctgc tgcttgggat 300
gagaccctcc tagacaaatt ctacactgaa ctctaccagc agctgaatga cctggaagcc 360
tgtgtgatac agggggtggg ggtgacagag actccccctga tgaaggagga ctccattctg 420
gctgtgagga aatacttcca aagaatcact ctctatctga aagagaagaa atacagccct 480
tgtgcctggg aggttgtcag agcagaaatc atgagatctt tttctttgtc aacaaacttg 540
caagaaagtt taagaagtaa ggaagacaaa actcacacgt gcccgccgtg cccagctccg 600
gaactgctgg gcggaccgtc agtcttcctc tccccccaa aaccaagga caccctcatg 660
atctcccga cccctgaggt cacatgcgtg gtggtggacg tgagccacga agaccctgag 720
gtcaagttca actggtacgt ggacggcgtg gaggtgcata atgccaagac aaagccgcgg 780
gaggagcagt acaacagcac gtaccgtgtg gtcagcgtcc tcaccgtcct gcaccaggac 840
tggctgaatg gcaaggagta caagtgcagg gtctccaaca aagccctccc agcccccatc 900
gagaaaacca tctccaaagc caaagggcag ccccgagaac cacaggtgta caccctgccc 960
ccatcccggg atgagctgac caagaaccag gtcagcctga cctgcctggt caaaggcttc 1020
tatcccagcg acatcgccgt ggagtgggag agcaatgggc agccggagaa caactacaag 1080
accacgcctc cgtgtttgga ctccgacggc tccttcttcc tctacagcaa gctcaccgtg 1140
gacaagagca ggtggcagca ggggaacgtc ttctcatgct ccgtgatgca tgaggctctg 1200
cacaaccact acacgcagaa gagcctctcc ctgtctccgg gtaaataga 1248

```

<210> 14

<211> 256

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic construct

<400> 14

```

Met Glu Thr Asp Thr Leu Leu Leu Trp Val Leu Leu Leu Trp Val Pro
  1                      5                      10                      15

Gly Ser Thr Gly Asp Asp Tyr Lys Asp Asp Asp Asp Lys Asp Lys Thr
          20                      25                      30

His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser
          35                      40                      45

Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg
          50                      55                      60

Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu Asp Pro
          65                      70                      75                      80

Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala

```

	85		90		95
Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val	100		105		110
Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr	115		120		125
Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr	130		135		140
Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu	145		150		155
Pro Pro Ser Arg Asp Glu Leu Thr Lys Asn Gln Val Ser Leu Thr Cys	165		170		175
Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser	180		185		190
Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp	195		200		205
Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser	210		215		220
Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala	225		230		235
Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys	245		250		255

<210> 15

<211> 771

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic construct

<400> 15

```

atggagacag acacactcct gctatgggta ctgctgctct gggttccagg ttccactggg 60
gacgactaca aggacgacga tgacaaggac aaaactcaca catgcccacc gtgcccagct 120
ccggaactcc tgggggggacc gtcagtcttc ctcttcccc caaaacccaa ggacaccctc 180
atgatctccc ggacccctga ggtcacatgc gtgggtgggtg acgtgagcca cgaagaccct 240
gaggtcaagt tcaactggta cgtggacggc gtggaggtgc ataatgccaa gacaaagccg 300
cgggaggagc agtacaacag caggtaccgt gtggtcagcg tcctcaccgt cctgcaccag 360
gactggctga atggcaagga gtacaagtgc aaggtctcca acaaagccct ccagccccc 420
atcgagaaaa ccattctcaa agccaaaggg cagccccgag aaccacaggt gtacaccctg 480
cccccatccc gggatgagct gaccaagaac caggtcagcc tgacctgcct ggtcaaaggc 540
ttctatccca gcgacatcgc cgtggagtgg gagagcaatg ggcagccgga gaacaactac 600
aagaccacgc ctcccgtgtt ggactccgac ggctccttct tcctctacag caagctcacc 660

```

gtggacaaga gcaggtggca gcaggggaac gtcttctcat gctccgtgat gcatgaggct 720
 ctgcacaacc actacacgca gaagagcctc tccctgtctc cgggtaaatg a 771

<210> 16

<211> 444

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
 construct

<400> 16

Met	Val	Pro	Cys	Thr	Leu	Leu	Leu	Leu	Leu	Ala	Ala	Ala	Leu	Ala	Pro
1				5					10					15	
Thr	Gln	Thr	Arg	Ala	Gly	Ser	Arg	Ala	Pro	Pro	Arg	Leu	Ile	Cys	Asp
			20					25					30		
Ser	Arg	Val	Leu	Gln	Arg	Tyr	Leu	Leu	Glu	Ala	Lys	Glu	Ala	Glu	Asn
		35					40					45			
Ile	Thr	Thr	Gly	Cys	Ala	Glu	His	Cys	Ser	Leu	Asn	Glu	Asn	Ile	Thr
	50					55					60				
Val	Pro	Asp	Thr	Lys	Val	Asn	Phe	Tyr	Ala	Trp	Lys	Arg	Met	Glu	Val
65					70					75					80
Gly	Gln	Gln	Ala	Val	Glu	Val	Trp	Gln	Gly	Leu	Ala	Leu	Leu	Ser	Glu
				85				90						95	
Ala	Val	Leu	Arg	Gly	Gln	Ala	Leu	Leu	Val	Asn	Ser	Ser	Gln	Pro	Trp
			100					105					110		
Glu	Pro	Leu	Gln	Leu	His	Val	Asp	Lys	Ala	Val	Ser	Gly	Leu	Arg	Ser
		115					120					125			
Leu	Thr	Thr	Leu	Leu	Arg	Ala	Leu	Gly	Ala	Gln	Lys	Glu	Ala	Ile	Ser
	130					135					140				
Pro	Pro	Asp	Ala	Ala	Ser	Ala	Ala	Pro	Leu	Arg	Thr	Ile	Thr	Ala	Asp
145					150					155					160
Thr	Phe	Arg	Lys	Leu	Phe	Arg	Val	Tyr	Ser	Asn	Phe	Leu	Arg	Gly	Lys
			165						170					175	
Leu	Lys	Leu	Tyr	Thr	Gly	Glu	Ala	Cys	Arg	Thr	Gly	Asp	Arg	Glu	Phe
			180					185					190		
Gly	Gly	Glu	Tyr	Gln	Ala	Leu	Glu	Lys	Glu	Val	Ala	Gln	Leu	Glu	Ala
		195					200					205			
Glu	Asn	Gln	Ala	Leu	Glu	Lys	Glu	Val	Ala	Gln	Leu	Glu	His	Glu	Gly
	210					215					220				
Gly	Gly	Pro	Ala	Pro	Glu	Leu	Leu	Gly	Gly	Pro	Ser	Val	Phe	Leu	Phe

atggtaccgt	gcacgctgct	cctgctgttg	gcggccgccc	tggctccgac	tcagaccgcg	60
gccggtctcta	gagccccacc	acgcctcatc	tgtgacagcc	gagtcctgca	gaggtaacctc	120
ttggaggcca	aggaggccga	gaatatcacg	acgggctgtg	ctgaacactg	cagcttgaat	180
gagaatatca	ctgtcccaga	caccaaagtt	aatttctatg	cctggaagag	gatggaagtc	240

```

gggcagcagg ccgtagaagt ctggcagggc ctggccctgc tgtcgggaagc tgtcctgcgg 300
ggccaggccc tgttgggtcaa ctcttcccag ccgtggggagc ccctgcagct gcatgtggat 360
aaagccgtca gtggccttcg cagcctcacc actctgtctt gggctctggg agcccagaag 420
gaagccatct cccctccaga tgcggcctca gctgtctccac tccgaacaat cactgctgac 480
actttccgca aactcttccg agtctactcc aatttctctc ggggaaagct gaagctgtac 540
acaggggagg cctgcaggac cggtgacagg gaattcgggt gtgagtacca ggccctggag 600
aaggagggtg cccagctgga ggccgagaac caggccctgg agaaggaggt ggccagctg 660
gagcacgagg gtggtgggtcc cgcacccgag ctgctggggc gaccgtcagt cttcctcttc 720
ccccaaaac ccaaggacac cctcatgata tcccggaccc ctgaggtcac atgctggtg 780
gtggacgtga gccacgaaga ccctgagggtc aagttcaact ggtacgtgga cggcgtggag 840
gtgcataatg ccaagacaaa gccgcgggag gagcagtaca acagcacgta ccgtgtgggtc 900
agcgtcctca ccgtcctgca ccaggactgg ctgaatggca aggagtacaa gtgcaagggtc 960
tccaacaaag cctcccagc ccccatcgag aaaaccatct ccaaagccaa agggcagccc 1020
cgagaaccac aggtgtacac cctgccccca tcccgggatg agctgaccaa gaaccagggtc 1080
agcctgacct gcctgggtcaa aggcttctat ccagcgaca tcgccgtgga gtgggagagc 1140
aatgggcagc cggagaacaa ctacaagacc acgcctcccg tgttggtact cgacgggtcc 1200
ttcttctct acagcaagct caccgtggac aagagcaggt ggcagcaggg gaacgtcttc 1260
tcatgctccg tgatgcatga ggctctgcac aaccactaca cgcagaagag cctctccctg 1320
tctccgggta aatga 1335

```

<210> 18

<211> 276

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic construct

<400> 18

```

Met Val Pro Cys Thr Leu Leu Leu Leu Leu Ala Ala Ala Leu Ala Pro
  1              5              10              15

```

```

Thr Gln Thr Arg Ala Gly Glu Phe Gly Gly Glu Tyr Gln Ala Leu Lys
      20              25              30

```

```

Lys Lys Val Ala Gln Leu Lys Ala Lys Asn Gln Ala Leu Lys Lys Lys
      35              40              45

```

```

Val Ala Gln Leu Lys His Lys Gly Gly Gly Pro Ala Pro Glu Leu Leu
      50              55              60

```

```

Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu
      65              70              75              80

```

```

Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser
      85              90              95

```

```

His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu
      100              105              110

```

```

Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr
      115              120              125

```

```

Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn
      130              135              140

```

Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro
 145 150 155 160
 Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln
 165 170 175
 Val Tyr Thr Leu Pro Pro Ser Arg Asp Glu Leu Thr Lys Asn Gln Val
 180 185 190
 Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val
 195 200 205
 Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro
 210 215 220
 Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr
 225 230 235 240
 Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val
 245 250 255
 Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu
 260 265 270
 Ser Pro Gly Lys
 275

<210> 19

<211> 831

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic construct

<400> 19

```

atggtaccgt gcacgtgtgt cctgctgttg gggccgccc tggctccgac tcagacccgc 60
gccggcgaat tcggtggtga gtaccaggcc ctgaagaaga aggtggcca gctgaaggcc 120
aagaaccagg ccctgaagaa gaaggtggcc cagctgaagc acaaggcg cgccccgc 180
ccagagctcc tgggcggacc gtcagtcttc ctcttcccc caaaaccac ggacaccctc 240
atgatctccc ggaccctga ggtcacatgc gtggtggtgg acgtgagcca cgaagaccct 300
gaggtcaagt tcaactggta cgtggacggc gtggaggtgc ataatgcaa gacaaagccg 360
cgggaggagc agtacaacag cacgtaccgt gtggtcagcg tcctcaccgt cctgcaccag 420
gactggctga atggcaagga gtacaagtgc aaggtctcca acaaagccct cccagcccc 480
atcgagaaaa ccattctcaa agccaaaggg cagccccgag aaccacaggt gtacaccctg 540
ccccatccc gggatgagct gaccaagaac caggtcagcc tgacctgcct ggtcaaaggc 600
ttctatccca gcgacatcgc cgtggagtg gagagcaatg ggcagccgga gaacaactac 660
aagaccacgc ctcccgtgtt ggactccgac ggctccttct tcctctacag caagctcacc 720
gtggacaaga gcaggtggca gcaggggaac gtcttctcat gtcctgtgat gcatgagggt 780
ctgcacaacc actacacgca gaagagcctc tcctgtctc cgggtaaatg a 831

```

<210> 20

<211> 245

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic construct

<400> 20

Met	Ala	Leu	Thr	Phe	Ala	Leu	Leu	Val	Ala	Leu	Leu	Val	Leu	Ser	Cys
1				5					10					15	

Lys	Ser	Ser	Cys	Ser	Val	Gly	Cys	Pro	Pro	Cys	Pro	Ala	Pro	Glu	Leu
			20					25					30		

Leu	Gly	Gly	Pro	Ser	Val	Phe	Leu	Phe	Pro	Pro	Lys	Pro	Lys	Asp	Thr
		35					40					45			

Leu	Met	Ile	Ser	Arg	Thr	Pro	Glu	Val	Thr	Cys	Val	Val	Val	Asp	Val
	50					55					60				

Ser	His	Glu	Asp	Pro	Glu	Val	Lys	Phe	Asn	Trp	Tyr	Val	Asp	Gly	Val
65					70					75					80

Glu	Val	His	Asn	Ala	Lys	Thr	Lys	Pro	Arg	Glu	Glu	Gln	Tyr	Asn	Ser
				85					90					95	

Thr	Tyr	Arg	Val	Val	Ser	Val	Leu	Thr	Val	Leu	His	Gln	Asp	Trp	Leu
			100					105					110		

Asn	Gly	Lys	Glu	Tyr	Lys	Cys	Lys	Val	Ser	Asn	Lys	Ala	Leu	Pro	Ala
		115					120					125			

Pro	Ile	Glu	Lys	Thr	Ile	Ser	Lys	Ala	Lys	Gly	Gln	Pro	Arg	Glu	Pro
	130					135					140				

Gln	Val	Tyr	Thr	Leu	Pro	Pro	Ser	Arg	Asp	Glu	Leu	Thr	Lys	Asn	Gln
145					150					155					160

Val	Ser	Leu	Thr	Cys	Leu	Val	Lys	Gly	Phe	Tyr	Pro	Ser	Asp	Ile	Ala
				165					170					175	

Val	Glu	Trp	Glu	Ser	Asn	Gly	Gln	Pro	Glu	Asn	Asn	Tyr	Lys	Thr	Thr
		180						185					190		

Pro	Pro	Val	Leu	Asp	Ser	Asp	Gly	Ser	Phe	Phe	Leu	Tyr	Ser	Lys	Leu
		195					200					205			

Thr	Val	Asp	Lys	Ser	Arg	Trp	Gln	Gln	Gly	Asn	Val	Phe	Ser	Cys	Ser
	210					215					220				

Val	Met	His	Glu	Ala	Leu	His	Asn	His	Tyr	Thr	Gln	Lys	Ser	Leu	Ser
225					230					235					240

Leu	Ser	Pro	Gly	Lys
				245

<210> 21
 <211> 738
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 construct

<400> 21
 atggccttgga ccttttgcttt actggtggcc ctcttggtgc tcagctgcaa gtcaagctgc 60
 tctgtgggct gcccgccgtg cccagctccg gaactgctgg gcggaccgtc agtcttcctc 120
 ttcccccaaa aacccaagga caccctcatg atctcccgga cccctgaggt cacatgcgtg 180
 gtggtggacg tgagccacga agaccctgag gtcaagttca actggtacgt ggacggcgtg 240
 gaggtgcata atgccaagac aaagccgcgg gaggagcagt acaacagcac gtaccgtgtg 300
 gtcagcgtcc tcaccgtcct gcaccaggac tggctgaatg gcaaggagta caagtgaag 360
 gtctccaaca aagccctccc agcccccatc gagaaaacca tctccaaagc caaagggcag 420
 ccccgagaac cacagggtgta caccctgccc ccatcccgga atgagctgac caagaaccag 480
 gtcagcctga cctgcctggt caaaggcttc tatccagcg acatcgccgt ggagtgggag 540
 agcaatgggc agccggagaa caactacaag accacgcctc ccgtgttgga ctccgacggc 600
 tccttcttcc tctacagcaa gctcaccgtg gacaagagca ggtggcagca ggggaacgtc 660
 ttctcatgct ccgtgatgca tgaggctctg cacaaccact acacgcagaa gagcctctcc 720
 ctgtctccgg gtaaata 738

<210> 22
 <211> 430
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 construct

<400> 22
 Met Ala Leu Thr Phe Ala Leu Leu Val Ala Leu Leu Val Leu Ser Cys
 1 5 10 15
 Lys Ser Ser Cys Ser Val Gly Cys Asp Leu Pro Gln Thr His Ser Leu
 20 25 30
 Gly Ser Arg Arg Thr Leu Met Leu Leu Ala Gln Met Arg Arg Ile Ser
 35 40 45
 Leu Phe Ser Cys Leu Lys Asp Arg His Asp Phe Gly Phe Pro Gln Glu
 50 55 60
 Glu Phe Gly Asn Gln Phe Gln Lys Ala Glu Thr Ile Pro Val Leu His
 65 70 75 80
 Glu Met Ile Gln Gln Ile Phe Asn Leu Phe Ser Thr Lys Asp Ser Ser
 85 90 95
 Ala Ala Trp Asp Glu Thr Leu Leu Asp Lys Phe Tyr Thr Glu Leu Tyr
 100 105 110

Gln Gln Leu Asn Asp Leu Glu Ala Cys Val Ile Gln Gly Val Gly Val
 115 120 125
 Thr Glu Thr Pro Leu Met Lys Glu Asp Ser Ile Leu Ala Val Arg Lys
 130 135 140
 Tyr Phe Gln Arg Ile Thr Leu Tyr Leu Lys Glu Lys Lys Tyr Ser Pro
 145 150 155 160
 Cys Ala Trp Glu Val Val Arg Ala Glu Ile Met Arg Ser Phe Ser Leu
 165 170 175
 Ser Thr Asn Leu Gln Glu Ser Leu Arg Ser Lys Glu Gly Gly Gly Gly
 180 185 190
 Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Asp Lys Thr His Thr
 195 200 205
 Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser Val Phe
 210 215 220
 Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro
 225 230 235 240
 Glu Val Thr Cys Val Val Val Asp Val Ser His Glu Asp Pro Glu Val
 245 250 255
 Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala Lys Thr
 260 265 270
 Lys Pro Arg Glu Glu Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser Val
 275 280 285
 Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys
 290 295 300
 Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser
 305 310 315 320
 Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro
 325 330 335
 Ser Arg Asp Glu Leu Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val
 340 345 350
 Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly
 355 360 365
 Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp
 370 375 380
 Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp
 385 390 395 400
 Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu His
 405 410 415

Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys
 420 425 430

<210> 23
 <211> 1291
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 construct

<400> 23
 atggccttga cctttgcttt actggtggcc ctctggtgc tcagctgcaa gtcaagctgc 60
 tctgtgggct gtgatctgcc tcaaaccac agcctgggta gcaggaggac cttgatgctc 120
 ctggcacaga tgaggagaat ctctcttttc tctgtcttga aggacagaca tgacttttga 180
 tttccccagg aggagtttgg caaccagttc caaaaggctg aaaccatccc tgtcctccat 240
 gagatgatcc agcagatott caatctcttc agcacaaagg actcatctgc tgcttgggat 300
 gagaccctcc tagacaaatt ctacactgaa ctctaccagc agctgaatga cctggaggcc 360
 tgtgtgatac aggggggtggg ggtgacagag actcccctga tgaaggagga ctccattctg 420
 gctgtgagga aatacttcca aagaatcact ctctatctga aagagaagaa atacagccct 480
 tgtgcctggg aggttgtcag agcagaaatc atgagatctt tttctttgtc aacaaacttg 540
 caagaaagtt tacgtagtaa ggaaggtggc ggcggatccg gtggaggcgg gtccggcggt 600
 ggaggggagcg aaaaaactca cacgtgcccg ccgtgcccag ctccggaact gctgggcgga 660
 ccgtcagttt cctcttcccc caaaaaccca aggacaccct catgatctcc cggacccctg 720
 aggtcacatg cgtgggtggtg gacgtgagcc acgaagacc tgaggtcaag ttcaactggt 780
 acgtggacgg cgtggagggtg cataatgcca agacaaagcc gcgggaggag cagtacaaca 840
 gcacgtaccg tgtggtcagc gtcctcaccg tcctgcacca ggactggctg aatggcaaag 900
 agtacaagtg caaggtctcc aacaaagccc tcccagcccc catcgagaaa accatctcca 960
 aagcaaaggg cagccccgag aaccacaggt gtacaccctg ccccatccc gggatgagct 1020
 gaccaagaac caggtcagcc tgacctgcct ggtcaaaggc ttctatccca gcgacatcgc 1080
 cgtggagtgg gagagcaatg ggcagccgga gaacaactac aagaccacgc ctcccgtgtt 1140
 ggactccgac ggctccttct tcctctacag caagctcacc gtggacaaga gcagggtggca 1200
 gcaggggaac gtcttctcat gctccgtgat gcatgaggct ctgcacaacc actacacgca 1260
 gaagagcctc tccctgtctc cgggtaaagt a 1291

<210> 24
 <211> 428
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence: Synthetic
 construct

<400> 24
 Met Gly Val His Glu Cys Pro Ala Trp Leu Trp Leu Leu Leu Ser Leu
 1 5 10 15
 Leu Ser Leu Pro Leu Gly Leu Pro Val Leu Gly Ala Pro Pro Arg Leu
 20 25 30
 Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu Leu Glu Ala Lys Glu

35					40					45					
Ala	Glu	Asn	Ile	Thr	Thr	Gly	Cys	Ala	Glu	His	Cys	Ser	Leu	Asn	Glu
	50					55					60				
Asn	Ile	Thr	Val	Pro	Asp	Thr	Lys	Val	Asn	Phe	Tyr	Ala	Trp	Lys	Arg
	65					70					75				80
Met	Glu	Val	Gly	Gln	Gln	Ala	Val	Glu	Val	Trp	Gln	Gly	Leu	Ala	Leu
				85					90					95	
Leu	Ser	Glu	Ala	Val	Leu	Arg	Gly	Gln	Ala	Leu	Leu	Val	Asn	Ser	Ser
			100					105					110		
Gln	Pro	Trp	Glu	Pro	Leu	Gln	Leu	His	Val	Asp	Lys	Ala	Val	Ser	Gly
			115					120					125		
Leu	Arg	Ser	Leu	Thr	Thr	Leu	Leu	Arg	Ala	Leu	Gly	Ala	Gln	Lys	Glu
	130					135					140				
Ala	Ile	Ser	Pro	Pro	Asp	Ala	Ala	Ser	Ala	Ala	Pro	Leu	Arg	Thr	Ile
	145					150					155				160
Thr	Ala	Asp	Thr	Phe	Arg	Lys	Leu	Phe	Arg	Val	Tyr	Ser	Asn	Phe	Leu
				165					170					175	
Arg	Gly	Lys	Leu	Lys	Leu	Tyr	Thr	Gly	Glu	Ala	Cys	Arg	Thr	Gly	Asp
			180					185					190		
Arg	Glu	Phe	Ala	Gly	Ala	Ala	Ala	Val	Asp	Lys	Thr	His	Thr	Cys	Pro
	195						200					205			
Pro	Cys	Pro	Ala	Pro	Glu	Leu	Leu	Gly	Gly	Pro	Ser	Val	Phe	Leu	Phe
	210					215					220				
Pro	Pro	Lys	Pro	Lys	Asp	Thr	Leu	Met	Ile	Ser	Arg	Thr	Pro	Glu	Val
	225					230					235			240	
Thr	Cys	Val	Val	Val	Asp	Val	Ser	His	Glu	Asp	Pro	Glu	Val	Lys	Phe
				245					250					255	
Asn	Trp	Tyr	Val	Asp	Gly	Val	Glu	Val	His	Asn	Ala	Lys	Thr	Lys	Pro
			260				265						270		
Arg	Glu	Glu	Gln	Tyr	Asn	Ser	Thr	Tyr	Arg	Val	Val	Ser	Val	Leu	Thr
	275						280					285			
Val	Leu	His	Gln	Asp	Trp	Leu	Asn	Gly	Lys	Glu	Tyr	Lys	Cys	Lys	Val
	290					295					300				
Ser	Asn	Lys	Ala	Leu	Pro	Ala	Pro	Ile	Glu	Lys	Thr	Ile	Ser	Lys	Ala
	305					310					315				320
Lys	Gly	Gln	Pro	Arg	Glu	Pro	Gln	Val	Tyr	Thr	Leu	Pro	Pro	Ser	Arg
				325					330					335	
Asp	Glu	Leu	Thr	Lys	Asn	Gln	Val	Ser	Leu	Thr	Cys	Leu	Val	Lys	Gly

340	345	350
Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro		
355	360	365
Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser		
370	375	380
Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln		
385	390	395
Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His		
405	410	415
Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys		
420	425	

<210> 25

<211> 1287

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic construct

<400> 25

```

atgggagtgac acgaatgtcc tgcctggctg tggcttctcc tgtccctgct gtcgctccct 60
ctgggcctcc cagtctctggg cgccccacca cgctcatct gtgacagccg agtcctggag 120
aggtacctct tggaggccaa ggaggccgag aatatcacga cgggctgtgc tgaacactgc 180
agcttgaatg agaatatcac tgtcccagac accaaagtta atttctatgc ctggaagagg 240
atggaggtcg ggcagcaggc cgtagaagtc tggcagggcc tggccctgct gtcggaagct 300
gtcctgctgg gccaggccct gttggtcaac tcttcccagc cgtgggagcc cctgcagctg 360
catgtggata aagccgtcag tggccttcgc agcctcacca ctctgcttcg ggctctggga 420
gccagaagg aagccatctc cctccagat gcggcctcag ctgctccact ccgaacaatc 480
actgctgaca ctttccgcaa actcttcga gtctactcca atttctccg gggaaagctg 540
aagctgtaca caggggaggg ctgcagaaca ggggacagag agttcgccgg cgccgctgcg 600
gtcgacaaaa ctcacacatg cccaccgtgc ccagctccgg aactcctggg cggaccgtca 660
gtcttctct tcccccaaaa acccaaggac accctcatga tctcccggac ccctgaggtc 720
acatgcgtgg tgggtggacgt gagccacgaa gaccctgagg tcaagttcaa ctggtacgtg 780
gacggcgtgg aggtgcataa tgccaagaca aagccgcggg aggagcagta caacagcacg 840
taccgtgtgg tcagcgtcct caccgtcctg caccaggact ggctgaatgg caaggagtac 900
aagtgaagg tctccaacaa agccctcca gcccctatcg agaaaacat ctccaaagcc 960
aaagggcagc cccgagaacc acaggtgtac accctgcccc catcccggga tgagctgacc 1020
aagaaccagg tcagcctgac ctgcctggtc aaaggcttct atcccagcga catcgccgtg 1080
gagtgggaga gcaatgggca gccggagaac aactacaaga ccacgcctcc cgtgttggac 1140
tccgacggct ccttcttct ctacagcaag ctcaccgtgg acaagagcag gtggcagcag 1200
gggaacgtct tctcatgctc cgtgatgcat gaggctctgc acaaccacta cagcagaag 1260
agcctctccc tgtctccggg taaatga
1287

```

<210> 26

<211> 11

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 26

Pro Lys Asn Ser Ser Met Ile Ser Asn Thr Pro
1 5 10

<210> 27

<211> 7

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 27

His Gln Ser Leu Gly Thr Gln
1 5

<210> 28

<211> 8

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 28

His Gln Asn Leu Ser Asp Gly Lys
1 5

<210> 29

<211> 8

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 29

His Gln Asn Ile Ser Asp Gly Lys
1 5

<210> 30

<211> 8

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide

<400> 30

Val Ile Ser Ser His Leu Gly Gln
1 5

<210> 31

<211> 8

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: linker peptide

<400> 31

Glu Phe Ala Gly Ala Ala Ala Val
1 5

<210> 32

<211> 20

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: linker peptide

<220>

<223> this sequence may encompass 1-10 Gly-Ala repeats

<400> 32

Gly Ala Gly Ala Gly Ala Gly Ala Gly Ala Gly Ala Gly Ala Gly Ala
1 5 10 15

Gly Ala Gly Ala
20

<210> 33

<211> 30

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: linker peptide

<220>

<223> this sequence may encompass 1-10 Gly-Gly-Ser repeats

<400> 33

Gly Gly Ser Gly Gly Ser Gly Gly Ser Gly Gly Ser Gly Gly Ser Gly
1 5 10 15

Gly Ser Gly Gly Ser Gly Gly Ser Gly Gly Ser Gly Gly Ser
20 25 30

<210> 34
<211> 80
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: linker peptide

<220>
<221> MISC_FEATURE
<222> (1)..(30)
<223> this region may encompass 1-10 Gly-Gly-Ser repeats

<220>
<221> MISC_FEATURE
<222> (31)..(80)
<223> this region may encompass 1-10 Gly-Gly-Gly-Gly-Ser repeats

<400> 34
Gly Gly Ser Gly Gly Ser Gly Gly Ser Gly Gly Ser Gly Gly Ser Gly
1 5 10 15
Gly Ser Gly Gly Ser Gly Gly Ser Gly Gly Ser Gly Gly Ser Gly Gly
20 25 30
Gly Gly Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Gly Gly Gly
35 40 45
Gly Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly
50 55 60
Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser
65 70 75 80

<210> 35
<211> 3
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: linker peptide

<400> 35
Gly Gly Gly
1

<210> 36
<211> 7
<212> PRT
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: linker peptide

<400> 36

Ser Gly Gly Ser Gly Gly Ser
1 5

<210> 37

<211> 15

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: linker peptide

<400> 37

Gly Gly Ser Gly Gly Ser Gly Gly Ser Gly Gly Ser Gly Gly Gly
1 5 10 15

<210> 38

<211> 16

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: linker peptide

<400> 38

Gly Gly Ser Gly Gly Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser
1 5 10 15

<210> 39

<211> 18

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: linker peptide

<400> 39

Gly Gly Ser Gly Gly Ser Gly Gly Ser Gly Gly Ser Gly Gly Ser Gly
1 5 10 15

Gly Ser

<210> 40

<211> 50

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: linker peptide

<220>

<223> this sequence may encompass 1-10 Gly-Gly-Gly-Gly-Ser repeats

<400> 40

Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Gly
1 5 10 15

Gly Gly Gly Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Gly Gly
20 25 30

Gly Gly Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Gly Gly Gly
35 40 45

Gly Ser
50

<210> 41

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer

<400> 41

gctggctagc caccatgga 19

<210> 42

<211> 45

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer

<400> 42

cttgtcatcg tcgtccttgt agtcgtcacc agtggaaacct ggaac 45

<210> 43

<211> 67

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer

<400> 43

gactacaagg acgacgatga caaggacaaa actcacacat gccaccgtg cccagctccg 60
gaactcc 67

<210> 44

<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer

<400> 44
tagtggatcc tcatttaccc g 21

<210> 45
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer

<400> 45
gctacctgca ggccaccatg gtctcccagg ccctcagg 38

<210> 46
<211> 51
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer

<400> 46
cagttccgga gctgggcacg gcgggcacgt gtgagttttg tcgggaaatg g 51

<210> 47
<211> 33
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer

<400> 47
ttactgcaga aggttatgca gcgcgtgaac atg 33

<210> 48
<211> 38
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer

<400> 48
tttttcgaat tcagtgagct ttgttttttc cttaatcc 38

<210> 49
<211> 34
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer

<400> 49
ggtaagcttg ccatggagct gaggcctgg ttgc 34

<210> 50
<211> 27
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer

<400> 50
gttttcaatc tctaggaccc actcgcc 27

<210> 51
<211> 23
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer

<400> 51
gccaggccac atgactactc cgc 23

<210> 52
<211> 34
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer

<400> 52
ggtgaattct cactcaggca ggtgtgaggg cagc 34

<210> 53
<211> 37
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer

<400> 53
gctactgcag ccaccatggc cttgaccttt gctttac 37

<210> 54
<211> 35
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer

<400> 54
cggtgaattc ttccttactt cttaaacttt cttgc 35

<210> 55
<211> 73
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer

<400> 55
cagttccgga gctgggcacg gcgggcacgt gtgagttttg tcttccttac ttcttaaact 60
ttttgcaagt ttg 73

<210> 56
<211> 45
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer

<400> 56
gtcaggatcc ggcggtggag ggagcgacaa aactcacacg tgccc 45

<210> 57
<211> 32
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer

<400> 57
tgacgcggcc gctcatttac ccggagacag gg 32

<210> 58
<211> 32
<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer

<400> 58

ccgctagcct gcaggccacc atggccttga cc

32

<210> 59

<211> 34

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer

<400> 59

ccggatccgc cgccaccttc cttactacgt aaac

34

<210> 60

<211> 15

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Linker peptide

<400> 60

Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser
1 5 10 15

<210> 61

<211> 60

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer

<400> 61

gtcaggatcc ggtggaggcg ggtccggcgg tggagggagc gacaaaactc acacgtgccc 60

<210> 62

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer

<400> 62

atagaagcct ttgaccaggc

20

<210> 63
<211> 84
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 63
gccggcggaat tcggtggtga gtaccaggcc ctgaagaaga aggtggccca gctgaaggcc 60
aagaaccagg ccctgaagaa gaag 84

<210> 64
<211> 58
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 64
gtggcccagc tgaagcaciaa gggcggcggc cccgcccag agctcctggg cggaccga 58

<210> 65
<211> 82
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 65
cggtcgccc aggagctctg gggcggggcc gccgcccttg tgcttcagct gggccacctt 60
cttcttcagg gcctggttct tg 82

<210> 66
<211> 60
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 66
gccttcagct gggccacctt cttcttcagg gcctggtact caccaccgaa ttcgccggca 60

<210> 67
<211> 62

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 67

ccggtgacag ggaattcggg ggtgagtacc aggccctgga gaaggaggtg gccagctgg 60
ag 62

<210> 68

<211> 83

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 68

gccgagaacc aggccctgga gaaggaggtg gccagctgg agcacgaggg tgggtggccc 60
gctccagagc tgctgggcgg aca 83

<210> 69

<211> 55

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 69

gtccgcccag cagctctgga gcgggaccac caccctcgtg ctccagctgg gccac 55

<210> 70

<211> 90

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
oligonucleotide

<400> 70

ctccttctcc agggcctggg tctcggcctc cagctggggc acctccttct ccagggcctg 60
gtactcacca ccgaattccc tgtcaccgga 90

<210> 71

<211> 37

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer

<400> 71

gctactgcag ccaccatggc cttgaccttt gctttac

37

<210> 72

<211> 44

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer

<400> 72

cagttccgga gctgggcacg gcggagagcc cacagagcag cttg

44

<210> 73

<211> 42

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer

<400> 73

gtggtcatat gggcattgaa ggcagaggcg ccgctgcggt cg

42

<210> 74

<211> 50

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer

<400> 74

ggtggttgct cttccgcaaa aaccggaga caggagaga ctcttctgcg

50

<210> 75

<211> 32

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Primer

<400> 75

aatctagagc cccaccacgc ctcattctgtg ac

32

<210> 76
<211> 30
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer

<400> 76
ttgaattctc tgtccctgt cctgcaggcc 30

<210> 77
<211> 27
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer

<400> 77
gtacctgcag gcggagatgg gggtgca 27

<210> 78
<211> 22
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer

<400> 78
cctggtcac tgtccctgt cc 22

<210> 79
<211> 56
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer

<400> 79
gtccaacctg caggaagctt gccgccacca tgggagtgca cgaatgtcct gcctgg 56

<210> 80
<211> 67
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer

<400> 80
gccgaattca gttttgtcga ccgcagcggc gccggcgaac tctctgtccc ctgttctgca 60
ggcctcc 67

<210> 81
<211> 8
<212> PRT
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Linker peptide

<400> 81
Glu Phe Ala Gly Ala Ala Val
1 5

<210> 82
<211> 69
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer

<400> 82
gctgcggtcg acaaaactca cacatgccca ccgtgccag ctccggaact cctgggcgga 60
ccgtcagtc 69

<210> 83
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence: Primer

<400> 83
attggaattc tcatttacct ggagacaggg agaggg 36

<210> 84
<211> 4
<212> PRT
<213> Homo sapiens

<400> 84
Glu Leu Leu Gly
1

<210> 85
<211> 10
<212> PRT
<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Linker peptide

<220>

<223> this sequence may encompass 1-10 Gly repeats

<400> 85

Gly Gly Gly Gly Gly Gly Gly Gly Gly Gly
1 5 10

<210> 86

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Linker peptide

<400> 86

Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser
1 5 10

<210> 87

<211> 7

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
peptide fragment

<400> 87

Ser Val Gly Cys Pro Pro Cys
1 5

<210> 88

<211> 6

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic
peptide fragment

<400> 88

Val Gly Cys Pro Pro Cys
1 5

<210> 89

<211> 7

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide fragment

<400> 89

Ser Thr Gly Cys Pro Pro Cys
1 5

<210> 90

<211> 4

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic peptide fragment

<400> 90

Cys Pro Pro Cys
1

<210> 91

<211> 8

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Synthetic FLAG peptide

<400> 91

Asp Tyr Lys Asp Asp Asp Asp Lys
1 5

INTERNATIONAL SEARCH REPORT

International application No
PCT/US2006/000140

A. CLASSIFICATION OF SUBJECT MATTER

INV. A61P9/00 A61P9/10 A61K38/17 A61K38/18 A61K38/21
A61K38/48
ADD. C07K19/00 C07K14/505 C07K14/56 C07K14/565 C12N9/64

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

A61P A61K C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, BIOSIS, EMBASE, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 2004/101740 A (SYNTONIX PHARMACEUTICALS, INC; RIVERA, DANIEL, S; PETERS, ROBERT, T; B) 25 November 2004 (2004-11-25) figures 6-8 paragraph [0030] examples 5-8	1-3
Y	WO 2004/003176 A (THE KENNETH S. WARREN INSTITUTE, INC; H. LUNDBECK A/S) 8 January 2004 (2004-01-08) pages 112-113; examples 6.7-6.8	1-3
Y	WO 01/91780 A (ORTHO-MCNEIL PHARMACEUTICAL, INC) 6 December 2001 (2001-12-06) example 5	1-3
	----- -/-	

☒ Further documents are listed in the continuation of Box C.

☒ See patent family annex.

* Special categories of cited documents :

A document defining the general state of the art which is not considered to be of particular relevance

E earlier document but published on or after the international filing date

L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

O document referring to an oral disclosure, use, exhibition or other means

P document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

G document member of the same patent family

Date of the actual completion of the international search

22 May 2006

Date of mailing of the international search report

02/06/2006

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Bumb, P

INTERNATIONAL SEARCH REPORT

International application No

PCT/US2006/000140

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 2004/006962 A (NOVO NORDISK A/S) 22 January 2004 (2004-01-22) examples 1-3 claims 1-24,33,34 -----	1,2
Y	WO 02/089828 A (INSTITUTE OF MOLECULAR AND CELL BIOLOGY; TAN, YIN, HWEE; SCHELLEKENS,) 14 November 2002 (2002-11-14) pages 11-14 claims 1-16 -----	1,2
P,Y	WO 2005/001025 A (SYNTONIX, INC; PETERS, ROBERT, T; MEZO, ADAM, R; RIVERA, DANIEL, S; BI) 6 January 2005 (2005-01-06) examples 1-37 -----	1-3

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US2006/000140

Box II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 1-3 (with respect to industrial applicability)
because they relate to subject matter not required to be searched by this Authority, namely:

Although claims 1-3 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/US2006/000140

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 2004101740	A	25-11-2004	AU 2004239244 A1	25-11-2004
			CA 2522859 A1	25-11-2004
			EP 1624891 A2	15-02-2006
WO 2004003176	A	08-01-2004	AU 2003251770 A1	19-01-2004
			CA 2491567 A1	08-01-2004
			EP 1552298 A2	13-07-2005
			JP 2006507228 T	02-03-2006
			MX PA05000063 A	08-04-2005
WO 0191780	A	06-12-2001	AU 7490401 A	11-12-2001
			BR 0111182 A	25-02-2004
			CA 2410453 A1	06-12-2001
			CN 1452492 A	29-10-2003
			EP 1296702 A1	02-04-2003
			JP 2003534384 T	18-11-2003
			MX PA02011727 A	24-10-2003
			NZ 522924 A	24-12-2004
			ZA 200210304 A	19-03-2004
WO 2004006962	A	22-01-2004	AU 2003242507 A1	02-02-2004
			EP 1523504 A2	20-04-2005
WO 02089828	A	14-11-2002	AU 2002310922 A1	18-11-2002
WO 2005001025	A	06-01-2005	AU 2004252422 A1	06-01-2005
			CA 2522590 A1	06-01-2005
			EP 1654378 A2	10-05-2006
			EP 1625209 A2	15-02-2006
			WO 2004101739 A2	25-11-2004