




(86) Date de dépôt PCT/PCT Filing Date: 2004/09/07
(87) Date publication PCT/PCT Publication Date: 2005/03/24
(85) Entrée phase nationale/National Entry: 2006/03/03
(86) N° demande PCT/PCT Application No.: US 2004/029206
(87) N° publication PCT/PCT Publication No.: 2005/026137
(30) Priorité/Priority: 2003/09/06 (US60/500,444)

(51) Cl.Int./Int.Cl. *C07D 277/82* (2006.01),
C07D 417/12 (2006.01), *C07D 417/06* (2006.01),
C07D 417/04 (2006.01), *C07D 277/46* (2006.01),
C07D 263/48 (2006.01), *A61P 43/00* (2006.01),
A61K 31/426 (2006.01)
(71) Demandeur/Applicant:
VERTEX PHARMACEUTICALS INCORPORATED, US
(72) Inventeurs/Inventors:
MILLER, MARK T., US;
HADIDA RUAH, SARA S., US;
SINGH, ASHAVINI K., US;
CLEVELAND, THOMAS, US;
MAKINGS, LEWIS R., US;
...

(54) Titre : MODULATEURS DE TRANSPORTEURS A CASSETTE DE LIAISON A L'ATP
(54) Title: MODULATORS OF ATP-BINDING CASSETTE TRANSPORTERS

(57) **Abrégé/Abstract:**

The present invention relates to modulators of ATP-Binding Cassette ("ABC") transporters or fragments thereof, including CF Transmembrane Regulator ("CFTR"), compositions thereof, and methods therewith. The present invention also relates to methods of treating ABC transporter mediated diseases using such modulators.



(21) **2 537 841**

(13) **A1**

(72) Inventeurs(suite)/Inventors(continued): HAMILTON, MATTHEW, US; GROOTENHUIS, PETER D.J., US

(74) Agent: SMART & BIGGAR

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
24 March 2005 (24.03.2005)

PCT

(10) International Publication Number
WO 2005/026137 A3

(51) International Patent Classification⁷: **C07D 277/82**,
277/46, 417/12, 263/48, 417/04, 417/06, A61K 31/426,
A61P 43/00

(74) Agents: **HALEY, James F., Jr.** et al.; c/o Fish & Neave,
1251 Avenue of the Americas, New York, NY 10020 (US).

(21) International Application Number:
PCT/US2004/029206

(81) Designated States (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(22) International Filing Date:
7 September 2004 (07.09.2004)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/500,444 6 September 2003 (06.09.2003) US

(84) Designated States (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **MILLER, Mark T.** [US/US]; 5075 La Jolla Blvd., San Diego, CA 92109 (US). **HADIDA RUAH, Sara S.** [ES/US]; 3717 Nobel Drive #1401, San Diego, CA 92122 (US). **SINGH, Ashavini K.** [IN/US]; 13597 Tiverton Road, San Diego, CA 92130 (US). **CLEVELAND, Thomas** [US/US]; 4431 Vision Drive #7, San Diego, CA 92121 (US). **MAKINGS, Lewis R.** [US/US]; 337 Cerro Street, Encinitas, CA 92024 (US). **HAMILTON, Matthew** [US/US]; 3005 Ridgecliff Lane, Spring Valley, CA 91977 (US). **GROOTENHUIS, Peter D.J.** [NL/US]; 4801 Riding Ridge Road, San Diego, CA 92122 (US).

Published:


- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

(88) Date of publication of the international search report:
21 July 2005

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: MODULATORS OF ATP-BINDING CASSETTE TRANSPORTERS

(57) Abstract: The present invention relates to modulators of ATP-Binding Cassette ("ABC") transporters or fragments thereof, including CF Transmembrane Regulator ("CFTR"), compositions thereof, and methods therewith. The present invention also relates to methods of treating ABC transporter mediated diseases using such modulators.



WO 2005/026137 A3

MODULATORS OF ATP-BINDING CASSETTE TRANSPORTERSTECHNICAL FIELD OF THE INVENTION

[0001] The present invention relates to modulators of ATP-Binding Cassette ("ABC") transporters or fragments thereof, including CF Transmembrane Regulator ("CFTR"), compositions thereof, and methods therewith. The present invention also relates to methods of treating ABC transporter mediated diseases using such modulators.

BACKGROUND OF THE INVENTION

[0002] ABC transporters are a group of membrane transporter proteins that play a major role in the transport and protection of cells against a wide variety of pharmacological agents, potentially toxic drugs, and xenobiotics. ABC transporters are homologous membrane proteins that bind and use cellular adenosine triphosphate (ATP) for their specific activities. Some of these transporters were discovered as multidrug resistance proteins (like the MDR1-P glycoprotein, or the multidrug resistance protein, MRP1), defending malignant cancer cells against chemotherapeutic agents. Up until the present time, 48 Human ABC Transporters have been identified, and these have been arranged into 7 families based on their sequence identity and function.

[0003] ABC transporters play a variety of important physiological roles within the body, as well as providing a defense against harmful compounds from the environment.

Moreover they represent important potential drug targets both in their own right, as well as, because in many cases therapeutic drugs are also transported out of the target cell by these molecules.

[0004] One of the members of the ABC transporter family, namely, CFTR, is believed be the chloride channel responsible for cAMP-mediated chloride secretion in epithelial cells, and to play a key role in the secretion of chloride and maintenance of normal electrolyte transport throughout the body. CFTR is a protein of approximately 1480 amino acids made up of two repeated elements, each comprising six transmembrane segments and a nucleotide-binding domain. The two repeats are separated by a large, polar, regulatory (R)-domain containing multiple potential phosphorylation sites.

[0005] The gene associated with CFTR has been identified and sequenced (See Gregory, R. J. et al. (1990) Nature 347:382-386; Rich, D. P. et al. (1990) Nature 347:358-362), (Riordan, J. R. et al. (1989) Science 245:1066-1073). A defect in this gene leads to cystic fibrosis (hereinafter "CF"), the most common fatal genetic disease in humans affecting approximately one in every 2,500 infants born in the United States. Within the general United States population, up to 10 million people carry a single copy of the defective gene without apparent ill effects. In contrast, individuals with two copies of the CF associated gene suffer from the chronic effects of CF, including chronic lung destruction and death.

[0006] In patients with Cystic fibrosis, expression of the CF associated gene in airway cells, leads to reduced cellular apical chloride conductance causing an imbalance in ion and fluid transport. It is widely believed that this leads to the abnormal mucus secretion

in pancreatic ductules and in the airways that ultimately results in the pulmonary infections and epithelial cell damage typically associated with disease progression in CF. In addition to respiratory problems, CF patients typically suffer from gastrointestinal problems, and pancreatic insufficiency. Males are almost uniformly infertile and fertility is decreased in females. In contrast to the severe effects of two copies of the CF associated gene, individuals with a single copy of the CF associated gene exhibit increased resistance to cholera and to dehydration resulting from diarrhea - perhaps explaining the relatively high frequency of the CF gene within the population.

[0007] Sequence analysis of the CFTR gene of CF chromosomes has revealed a variety of disease causing mutations (Cutting, G. R. et al. (1990) Nature 346:366-369; Dean, M. et al. (1990) Cell 61:863:870; and Kerem, B-S. et al. (1989) Science 245:1073-1080; Kerem, B-S et al. (1990) Proc. Natl. Acad. Sci. USA 87:8447-8451). At present, more than 1000 mutations in the CF gene have been identified (<http://www.genet.sickkids.on.ca/cftr/>), but population studies have indicated that the most common CF mutation, a deletion of the 3 nucleotides that encode phenylalanine at position 508 of the CFTR amino acid sequence, is associated with approximately 70% of the cases of cystic fibrosis. The mutated CFTR protein is referred to as $\Delta F508$.

[0008] It is believed that the deletion of residue 508 in $\Delta F508$ -CFTR prevents the nascent protein from folding correctly, resulting in the inability of this mutant protein to exit the endoplasmic reticulum (hereinafter "ER"), and traffic to the plasma membrane. As a result, insufficient amounts of the mature protein are present at the plasma membrane and chloride transport

within epithelial tissues is significantly reduced (Quinton, P. M. (1990), FASEB J. 4: 2709-2727). Hence, the cellular phenomenon of defective ER processing of other protein/s like CFTR, by the ER machinery, has been shown to be the underlying basis for a wide range of isolated and inherited diseases. The two ways that the ER machinery can malfunction is either by loss of coupling to ER export of the proteins leading to degradation, or by the ER accumulation of these defective/misfolded proteins [Aridor M, et al., Nature Med., 5(7), pp 745- 751 (1999); Shastry, B.S., et al., Neurochem. International, 43, pp 1-7 (2003); Rutishauser, J., et al., Swiss Med Wkly, 132, pp 211-222 (2002); Morello, JP et al., TIPS, 21, pp. 466- 469 (2000); Bross P., et al., Human Mut., 14, pp. 186-198 (1999)]. Studies have shown, however, that $\Delta F508$ -CFTR, when presented at the plasma membrane is functional as a cAMP-responsive Cl^- channel (Dalemans et al. (1991), Nature Lond. 354: 526-528; Denning et al., supra.; Pasyk and Foskett (1995), J. Cell. Biochem. 270: 12347-50).

[0009] Although CFTR transports a variety of molecules in addition to anions, it is clear that this role (the transport of anions) represents one element in an important mechanism of transporting ions and water across the epithelium. The other elements include the epithelial Na^+ channel, ENaC, $Na^+/2Cl^-/K^+$ co-transporter, Na^+-K^+ -ATPase pump and the basolateral membrane K^+ channels, that are responsible for the uptake of chloride into the cell.

[0010] These elements work together to achieve directional transport across the epithelium via their selective expression and localization within the cell. Chloride absorption takes place by the coordinated activity of ENaC and CFTR present on the apical membrane

and the Na^+ - K^+ -ATPase pump and Cl^- channels expressed on the basolateral surface of the cell. Secondary active transport of chloride from the luminal side leads to the accumulation of intracellular chloride, which can then passively leave the cell via Cl^- channels, resulting in a vectorial transport. Arrangement of $\text{Na}^+/\text{2Cl}^-/\text{K}^+$ co-transporter, Na^+ - K^+ -ATPase pump and the basolateral membrane K^+ channels on the basolateral surface and CFTR on the luminal side coordinate the secretion of chloride via CFTR on the luminal side. Because water is probably never actively transported itself, its flow across epithelia depends on tiny transepithelial osmotic gradients generated by the bulk flow of sodium and chloride.

[0011] In addition to CF, modulation of CFTR activity may be beneficial for other diseases not directly caused by mutations in CFTR, such as secretory diseases and other protein folding diseases mediated by CFTR. These include, but are not limited to, chronic obstructive pulmonary disease (hereinafter "COPD"), dry eye disease, and Sjögren's Syndrome.

[0012] COPD is characterized by airflow limitation that is progressive and not fully reversible. The airflow limitation is due to mucus hypersecretion, emphysema, and bronchiolitis. Activators of mutant or wild-type CFTR offer a potential treatment of mucus hypersecretion and impaired mucociliary clearance that is common in COPD. Specifically, increasing anion secretion across CFTR may facilitate fluid transport into the airway surface liquid to hydrate the mucus and optimized periciliary fluid viscosity. This would lead to enhanced mucociliary clearance and a reduction in the symptoms associated with COPD. Dry eye disease is characterized by a decrease in tear aqueous production

and abnormal tear film lipid, protein and mucin profiles. There are many causes of dry eye, some of which include age, Lasik eye surgery, arthritis, medications, chemical/thermal burns, allergies, and diseases, such as Cystic Fibrosis and Sjögren's syndrome. Increasing anion secretion via CFTR would enhance fluid transport from the corneal endothelial cells and secretory glands surrounding the eye to increase corneal hydration. This would help to alleviate the symptoms associated with dry eye disease. Sjögren's syndrome is an autoimmune disease in which the immune system attacks moisture-producing glands throughout the body, including the eye, mouth, skin, respiratory tissue, liver, vagina, and gut. Symptoms, include, dry eye, mouth, and vagina, as well as lung disease. The disease is also associated with rheumatoid arthritis, systemic lupus, systemic sclerosis, and polymyositis/dermatomyositis. Defective protein trafficking is believed to cause the disease, for which treatment options are limited. Modulators of CFTR activity may hydrate the various organs afflicted by the disease and help to elevate the associated symptoms.

[0013] As discussed above, it is believed that the deletion of residue 508 in $\Delta F508$ -CFTR prevents the nascent protein from folding correctly, resulting in the inability of this mutant protein to exit the ER, and traffic to the plasma membrane. As a result, insufficient amounts of the mature protein are present at the plasma membrane and chloride transport within epithelial tissues is significantly reduced. In fact, this cellular phenomenon of defective ER processing of ABC transporters by the ER machinery, has been shown to be the underlying basis not only for CF disease, but for a wide range of other isolated and inherited diseases.

The two ways that the ER machinery can malfunction is either by loss of coupling to ER export of the proteins leading to degradation, or by the ER accumulation of these defective/misfolded proteins [Aridor M, et al., Nature Med., 5(7), pp 745- 751 (1999); Shastri, B.S., et al., Neurochem. International, 43, pp 1-7 (2003); Rutishauser, J., et al., Swiss Med Wkly, 132, pp 211-222 (2002); Morello, JP et al., TIPS, 21, pp. 466- 469 (2000); Bross P., et al., Human Mut., 14, pp. 186-198 (1999)].

[0014] The diseases associated with the first class of ER malfunction are CF (due to misfolded $\Delta F508$ -CFTR), hereditary emphysema (due to $\alpha 1$ -antitrypsin; non Piz variants), hereditary hemochromatosis, coagulation-fibrinolysis deficiencies, such as protein C deficiency, Type 1 hereditary angioedema, lipid processing deficiencies, such as familial hypercholesterolemia, Type 1 chylomicronemia, abetalipoproteinemia, lysosomal storage diseases, such as I-cell disease/pseudo-Hurler, mucopolysaccharidoses (due to lysosomal processing enzymes), Sandhof/Tay-Sachs (due to β -hexosaminidase), Crigler-Najjar type II (due to UDP-glucuronyl-sialyltransferase), polyendocrinopathy/ hyperinsulemia, Diabetes mellitus (due to insulin receptor), Laron dwarfism (due to growth hormone receptor), myeloperoxidase deficiency, primary hypoparathyroidism (due to preproparathyroid hormone), melanoma (due to tyrosinase). The diseases associated with the latter class of ER malfunction are glycanosis CDG type 1, Hereditary emphysema (due to $\alpha 1$ -Antitrypsin (PiZ variant), Congenital hyperthyroidism, osteogenesis imperfecta (due to Type I, II, IV procollagen), hereditary hypofibrinogenemia (due to fibrinogen), ACT deficiency (due to $\alpha 1$ -Antichymotrypsin), Diabetes

insipidus (DI), neurophyseal DI (due to Vasopressin hormone/V2-receptor), neprogenic DI (due to aquaporin II), Charcot-Marie Tooth syndrome (due to Peripheral myelin protein 22), Perlizaeus-Merzbacher disease, neurodegenerative diseases such as Alzheimer's disease (due to β APP and presenilins), Parkinson's disease, amyotrophic lateral sclerosis, progressive supranuclear palsy, Pick's disease, several polyglutamine neurological disorders such as Huntington, spinocerebellar ataxia type I, spinal and bulbar muscular atrophy, dentatorubal pallidoluysian, and myotonic dystrophy, as well as Spongiform encephalopathies, such as hereditary Creutzfeldt-Jakob disease (due to Prion protein processing defect), Fabry disease (due to lysosomal α -galactosidase A) and Strausler-Scheinker syndrome (due to Prp processing defect).

[0015] In CF, chloride transport mediated by the CFTR is reduced resulting in the abnormal mucus secretion that characterizes the disease. By contrast in secretory diarrheas epithelial water transport is dramatically increased as a result of secretagogue activated chloride transport. The mechanism involves elevation of cAMP and stimulation of CFTR.

[0016] Although there are numerous causes of diarrhea, the major consequences of diarrheal diseases, resulting from excessive chloride transport are common to all, and include dehydration, acidosis, death and impaired growth.

[0017] Acute and chronic diarrheas represent a major medical problem in many areas of the world. Diarrhea is both a significant factor in malnutrition and the leading cause of death (5,000,000 deaths/year) in children less than five years old.

[0018] Secretory diarrheas are also a dangerous condition in patients of acquired immunodeficiency syndrome (AIDS) and chronic inflammatory bowel disease (IBD). Sixteen million travelers to developing countries from industrialized nations every year develop diarrhea, with the severity and number of cases of diarrhea varying depending on the country and area of travel.

[0019] Diarrhea in barn animals and pets such as cows, pigs and horses, sheep, goats, cats and dogs, also known as scours, is a major cause of death in these animals. Diarrhea can result from any major transition, such as weaning or physical movement, as well as in response to a variety of bacterial or viral infections and generally occurs within the first few hours of the animal's life.

[0020] The most common diarrheal causing bacteria is enterotoxogenic E-coli (ETEC) having the K99 pilus antigen. Common viral causes of diarrhea include rotavirus and coronavirus. Other infectious agents include cryptosporidium, giardia lamblia, and salmonella, among others.

[0021] Symptoms of rotaviral infection include excretion of watery feces, dehydration and weakness. Coronavirus causes a more severe illness in the newborn animals, and has a higher mortality rate than rotaviral infection. Often, however, a young animal may be infected with more than one virus or with a combination of viral and bacterial microorganisms at one time. This dramatically increases the severity of the disease.

[0022] Accordingly, there is a need for modulators of an ABC transporter activity, and compositions thereof, that can be used to modulate the activity of the ABC transporter in the cell membrane of a mammal.

[0023] There is a need for methods of treating ABC transporter mediated diseases using such modulators of ABC transporter activity.

[0024] There is a need for methods of modulating an ABC transporter activity in an ex vivo cell membrane of a mammal.

[0025] There is a need for modulators of CFTR activity that can be used to modulate the activity of CFTR in the cell membrane of a mammal.



[0026] There is a need for methods of treating CFTR-mediated diseases using such modulators of CFTR activity.

[0027] There is a need for methods of modulating CFTR activity in an ex vivo cell membrane of a mammal.

[0028] There is a need for correctors that enhance the density of CFTR in the plasma membrane by facilitating the migration of the CFTR thereto.

SUMMARY OF THE INVENTION

[0029] The present invention provides a method of modulating ABC transporter activity, comprising the step of contacting said ABC transporter with a compound of formula I or formula I':



or a pharmaceutically acceptable salt thereof;
wherein:

Y' is O, S, or NR;

p is 0-2;


X is a bond, O, S, S(O), S(O)₂, CF₂, CH₂, -CHOR-, -C(O)-, -O-C(O)-, -C(O)-O-, -C(O)-NR-, -NR-C(O)-, -NR-C(O)-O-, -O-C(O)-NR-, -NR-C(O)-NR-, or NR;

R is H, R², or R⁶;

A is aliphatic, aryl, heteroaryl, heterocyclic, or cycloalkyl;

C is a phenyl or 5-8 membered cycloaliphatic ring;

Q is selected from:



each B is independently selected from 3-7 membered monocyclic or 8-14 membered bicyclic or tricyclic, saturated, unsaturated or aromatic ring containing 0-4 heteroatoms in each ring, wherein each said heteroatom is independently selected from N, NH, S, or O;

wherein each A, B, and C is independently and optionally substituted with up to 4 substituents independently selected from R¹, R², R³, R⁴, or R⁵;

R^L is -OR^A, -SR^A, or -N(R^{AB})₂;

each R^A is independently hydrogen, C1-C6 aliphatic, or a 3-7 membered carbocyclic or heterocyclic ring, saturated or unsaturated ring, having up to 3 heteroatoms selected from O, N, or S, wherein each R^A is optionally substituted with up to 3 substituents independently selected from R¹, R⁴ or R⁷,

each R^{AB} is independently hydrogen or C1-C6 aliphatic optionally substituted with up to 3 substituents independently selected from R¹, R⁴ or R⁷;

wherein up to two methylene units in R^A or R^{AB} are optionally replaced with $-CO-$, $-CS-$, $-COCO-$, $-CONR-$, $-CO_2-$, $-OCO-$, $-NRCO_2-$, $-O-$, $-NRCONR-$, $-OCONR-$, $-NRCO-$, $-S-$, $-SO-$, $-SO_2-$, $-NR-$, $-SO_2NR-$, $NRSO_2-$, or $-NRSO_2NR$; or

two R^{AB} , taken together with the nitrogen atom, is a 3-7 membered heterocyclic or heteroaryl ring containing up to 4 heteroatoms selected from O, N, or S, wherein said ring is optionally substituted with up to 2 substituents selected from oxo or $(C_{1-4}aliphatic)_p-Y$;

R^M is C1-C6 aliphatic, optionally substituted with up to two substituents selected from R^1 , R^2 , R^3 , or R^4 ;

each of X_1 and X_2 is independently selected from O, S, or NR;

R^N is C1-C6 aliphatic or phenyl, wherein R^N is optionally substituted with up to two substituents selected from R^1 , R^2 , R^3 , or R^4 ;

R^P is C1-C6 aliphatic, optionally substituted with up to two substituents selected from R^1 , R^2 , R^3 , or R^4 ;

R^Q is C1-C6 aliphatic or aryl, wherein R^Q is optionally substituted with up to two substituents selected from R^1 , R^2 , R^3 , or R^4 ;

R^1 is oxo, R^6 or $((C_{1-4}aliphatic)_n-Y$;

n is 0 or 1;

Y is halo, CN, NO_2 , CF_3 , OCF_3 , OH, SR^6 , $S(O)R^6$, SO_2R^6 , NH_2 , NHR^6 , $N(R^6)_2$, NR^6R^8 , $N(R^8)^2$, COOH, $COOR^6$ or OR^6 ; or

two R^1 on adjacent ring atoms, taken together, form 1,2-methylenedioxy or 1,2-ethylenedioxy;

R^2 is aliphatic, wherein each R^2 optionally comprises up to 2 substituents independently selected from R^1 , R^4 , or R^5 ;

R^3 is a cycloaliphatic, aryl, heterocyclic, or heteroaryl ring optionally comprising up to 3 substituents, independently selected from R^1 , R^2 , R^4 or R^5 ;

R^4 is OR^5 , OR^6 , $OC(O)R^6$, $OC(O)R^5$, $OC(O)OR^6$, $OC(O)OR^5$, $OC(O)N(R^6)_2$, $OC(O)N(R^5)_2$, $OC(O)N(R^6R^5)$, SR^6 , SR^5 , $S(O)R^6$, $S(O)R^5$, SO_2R^6 , SO_2R^5 , $SO_2N(R^6)_2$, $SO_2N(R^5)_2$, $SO_2NR^5R^6$, SO_3R^6 , SO_3R^5 , $C(O)R^5$, $C(O)OR^5$, $C(O)R^6$, $C(O)OR^6$, $C(O)N(R^6)_2$, $C(O)N(R^5)_2$, $C(O)N(R^5R^6)$, $C(O)N(OR^6)R^6$, $C(O)N(OR^5)R^6$, $C(O)N(OR^6)R^5$, $C(O)N(OR^5)R^5$, $C(NOR^6)R^6$, $C(NOR^6)R^5$, $C(NOR^5)R^6$, $C(NOR^5)R^5$, $N(R^6)_2$, $N(R^5)_2$, $N(R^5R^6)$, $NR^5C(O)R^5$, $NR^6C(O)R^6$, $NR^6C(O)R^5$, $NR^5C(O)R^6$, $NR^6C(O)OR^6$, $NR^5C(O)OR^6$, $NR^6C(O)OR^5$, $NR^5C(O)OR^5$, $NR^6C(O)N(R^6)_2$, $NR^6C(O)NR^5R^6$, $NR^6C(O)N(R^5)_2$, $NR^5C(O)N(R^6)_2$, $NR^5C(O)NR^5R^6$, $NR^5C(O)N(R^5)_2$, $NR^6SO_2R^6$, $NR^6SO_2R^5$, $NR^5SO_2R^5$, $NR^5SO_2R^6$, $NR^6SO_2N(R^6)_2$, $NR^6SO_2NR^5R^6$, $NR^6SO_2N(R^5)_2$, $NR^5SO_2N(R^6)_2$, $NR^5SO_2NR^5R^6$, $NR^5SO_2N(R^5)_2$, $N(OR^6)R^6$, $N(OR^6)R^5$, $N(OR^5)R^5$, or $N(OR^5)R^6$;

R^5 is a cycloaliphatic, aryl, heterocyclic, or heteroaryl ring, optionally comprising up to 3 R^1 substituents;

R^6 is H or aliphatic, wherein R^6 optionally comprises a R^7 substituent;

R^7 is a cycloaliphatic, aryl, heterocyclic, or heteroaryl ring, and each R^7 optionally comprises up to 2 substituents independently chosen from H, (C_1-C_6) -straight or branched alkyl, (C_2-C_6) straight or branched alkenyl or alkynyl, 1,2-methylenedioxy, 1,2-ethylenedioxy, or $(CH_2)_n-Z$;

Z is selected from halo, CN, NO₂, CF₃, OCF₃, OH, S-aliphatic, S(O)-aliphatic, SO₂-aliphatic, NH₂, NH-aliphatic, N(aliphatic)₂, N(aliphatic)R⁸, NHR⁸, N(R⁸)₂, COOH, C(O)O(-aliphatic), or O-aliphatic; and

R⁸ is an amino-capping group.

[0030] The present invention also provides compositions comprising compounds of formula (I) and formula (I'), and methods of treating ABC transporter mediated diseases using compounds of formula (I) and formula (I').

DETAILED DESCRIPTION OF THE INVENTION

[0031] As used herein, the following definitions shall apply unless otherwise indicated.

[0032] The term "ABC-transporter" as used herein means an ABC-transporter protein or a fragment thereof comprising at least one binding domain, wherein said protein or fragment thereof is present *in vivo* or *in vitro*. The term "binding domain" as used herein means a domain on the ABC-transporter that can bind to a modulator. See, e.g., Hwang, T. C. et al., J. Gen. Physiol. (1998): 111(3), 477-90.

[0033] The term "CFTR" as used herein means cystic fibrosis transmembrane regulator or a mutation thereof capable of its regulator activity, in part or full, including, but not limited to, ΔF508 CFTR and G551D CFTR (see, e.g., <http://www.genet.sickkids.on.ca/cftr/>, for CFTR mutations).

[0034] The term "modulating" as used herein means increasing or decreasing by a measurable amount.

[0035] The phrase "optionally substituted" is used interchangeably with the phrase "substituted or unsubstituted."

[0036] The term "aliphatic" or "aliphatic group", as used herein, means a straight-chain or branched, hydrocarbon chain that is completely saturated (alkyl) or is unsaturated (alkenyl or alkynyl). Unless otherwise specified, an aliphatic group has 1 to 12 carbon atoms, preferably, 1-6 carbon atoms, and more preferably, 1-4 carbon atoms. Up to three, and preferably two, $-CH_2-$ in said aliphatic may be replaced with O, S, or -NR.

[0037] The term "alkylidene" as used herein means a straight-chain or branched hydrocarbon chain that is completely saturated or unsaturated, and is connected to the rest of the molecule through covalent bonds. Exemplary alkylidene groups include methylene, ethylene, or propylene. Unless otherwise specified, an alkylidene group has 1 to 12 carbon atoms, preferably, 1-6 carbon atoms, and more preferably, 1-4 carbon atoms.

[0038] The term "cycloaliphatic" means a saturated or partially unsaturated monocyclic or bicyclic hydrocarbon ring that has up to two points of attachment to the rest of the molecule. Unless otherwise specified, preferred cycloaliphatic rings are 3-8 membered monocyclic rings, more preferably 3-6, and even more preferably, 3, 5, or 6. Also preferred, unless otherwise specified, are 8-12 membered bicyclic hydrocarbon rings, more preferably 10 membered bicyclic hydrocarbon rings.

[0039] The term "heteroatom," unless otherwise specified, means nitrogen, oxygen, or sulfur and includes any oxidized form of nitrogen and sulfur, and the quaternized form of any basic nitrogen. Also the term "nitrogen" includes a substitutable nitrogen of a heterocyclic ring. As an example, in a saturated or partially unsaturated ring having 0-3 heteroatoms selected from oxygen, sulfur or nitrogen, the nitrogen

may be N (as in 3,4-dihydro-2*H*-pyrrolyl), NH (as in pyrrolidinyl) or as in N-substituted pyrrolidinyl.

[0040] The term "unsaturated", as used herein, means a double bond or a triple bond. Each such bond constitutes one unit of unsaturation.



[0041] The term "aryl" used alone or as part of a larger moiety as in "aralkyl", "aralkoxy", or "aryloxyalkyl", refers to monocyclic, bicyclic and tricyclic aromatic carbocyclic ring systems. Unless otherwise specified, preferred aryl rings have a total of five to fourteen ring members, wherein at least one ring, if bicyclic or tricyclic, in the system is aromatic and wherein each ring in the system contains up to 6 ring members. The term "aryl" may be used interchangeably with the term "aryl ring". Phenyl is an example of aryl.

[0042] The term "heterocycle", "heterocyclyl", or "heterocyclic" as used herein means non-aromatic, monocyclic, bicyclic or tricyclic ring systems wherein one or more ring members is a heteroatom. Unless otherwise specified, each ring in the system preferably contains contains 3 to 7 ring members with preferably 1-3 heteroatoms.

[0043] The term "heteroaryl", used alone or as part of a larger moiety as in "heteroaralkyl" or "heteroarylalkoxy", refers to monocyclic, bicyclic and tricyclic ring systems, wherein at least one ring in the system is aromatic, and at least one ring in the system contains one or more heteroatoms. Unless otherwise specified, such ring systems preferably have a total of 5 to 15 ring members, wherein each ring in the system preferably contains 3 to 7 ring members, with preferably 1-3 heteroatoms. The term "heteroaryl" may be used interchangeably with the term "heteroaryl ring" or the term "heteroaromatic".

[0044] A combination of substituents or variables is permissible only if such a combination results in a stable or chemically feasible compound. A stable compound or chemically feasible compound is one that is not substantially altered when kept at a temperature of 40°C or less, in the absence of moisture or other chemically reactive conditions, for at least a week.

[0045] The present invention provides a method of modulating ABC transporter activity, comprising the step of contacting said ABC transporter with a compound of formula I or formula I':



or a pharmaceutically acceptable salt thereof;
wherein:

Y' is O, S, or NR;

p is 0-2;


X is a bond, O, S, S(O), S(O)₂, CF₂, CH₂, -CHOR-, -C(O)-, -O-C(O)-, -C(O)-O-, -C(O)-NR-, -NR-C(O)-, -NR-C(O)-O-, -O-C(O)-NR-, -NR-C(O)-NR-, or NR;

R is H, R², or R⁶;

A is aliphatic, aryl, heteroaryl, heterocyclic, or cycloalkyl;

C is a phenyl or 5-8 membered cycloaliphatic ring;

Q is selected from:



each B is independently selected from 3-7 membered monocyclic or 8-14 membered bicyclic or tricyclic, saturated, unsaturated or aromatic ring containing 0-4 heteroatoms in each ring, wherein each said heteroatom is independently selected from N, NH, S, or O;

wherein each A, B, and C is independently and optionally substituted with up to 4 substituents independently selected from R^1 , R^2 , R^3 , R^4 , or R^5 ;

R^{L} is ---OR^{A} , ---SR^{A} , or $\text{---N(R}^{\text{AB}}\text{)}_2$;

each R^{A} is independently hydrogen, C1-C6 aliphatic, or a 3-7 membered carbocyclic or heterocyclic ring, saturated or unsaturated ring, having up to 3 heteroatoms selected from O, N, or S, wherein each R^{A} is optionally substituted with up to 3 substituents independently selected from R^1 , R^4 or R^7 ,

each R^{AB} is independently hydrogen or C1-C6 aliphatic optionally substituted with up to 3 substituents independently selected from R^1 , R^4 or R^7 ;

wherein up to two methylene units in R^{A} or R^{AB} are optionally replaced with ---CO--- , ---CS--- , ---COCO--- , ---CONR--- , $\text{---CO}_2\text{---}$, ---OCO--- , $\text{---NRCO}_2\text{---}$, ---O--- , ---NRCONR--- , ---OCONR--- , ---NRCO--- , ---S--- , ---SO--- , $\text{---SO}_2\text{---}$, ---NR--- , $\text{---SO}_2\text{NR---}$, $\text{NRSO}_2\text{---}$, or $\text{---NRSO}_2\text{NR---}$; or

two R^{AB} , taken together with the nitrogen atom, is a 3-7 membered heterocyclic or heteroaryl ring containing up to 4 heteroatoms selected from O, N, or S, wherein

said ring is optionally substituted with up to 2 substituents selected from oxo or (C₁₋₄aliphatic)_p-Y;

R^M is C1-C6 aliphatic, optionally substituted with up to two substituents selected from R¹, R², R³, or R⁴;

each of X₁ and X₂ is independently selected from O, S, or NR;

R^N is C1-C6 aliphatic or phenyl, wherein R^N is optionally substituted with up to two substituents selected from R¹, R², R³, or R⁴;

R^P is C1-C6 aliphatic, optionally substituted with up to two substituents selected from R¹, R², R³, or R⁴;

R^Q is C1-C6 aliphatic or aryl, wherein R^Q is optionally substituted with up to two substituents selected from R¹, R², R³, or R⁴;

R¹ is oxo, R⁶ or ((C1-C4)aliphatic)_n-Y;

n is 0 or 1;

Y is halo, CN, NO₂, CF₃, OCF₃, OH, SR⁶, S(O)R⁶, SO₂R⁶, NH₂, NHR⁶, N(R⁶)₂, NR⁶R⁸, N(R⁸)₂, COOH, COOR⁶ or OR⁶; or

two R¹ on adjacent ring atoms, taken together, form 1,2-methylenedioxy or 1,2-ethylenedioxy;

R² is aliphatic, wherein each R² optionally comprises up to 2 substituents independently selected from R¹, R⁴, or R⁵;

R³ is a cycloaliphatic, aryl, heterocyclic, or heteroaryl ring optionally comprising up to 3 substituents, independently selected from R¹, R², R⁴ or R⁵;

R⁴ is OR⁵, OR⁶, OC(O)R⁶, OC(O)R⁵, OC(O)OR⁶, OC(O)OR⁵, OC(O)N(R⁶)₂, OC(O)N(R⁵)₂, OC(O)N(R⁶R⁵), SR⁶, SR⁵, S(O)R⁶, S(O)R⁵, SO₂R⁶, SO₂R⁵, SO₂N(R⁶)₂, SO₂N(R⁵)₂,

$\text{SO}_2\text{NR}^5\text{R}^6$, SO_3R^6 , SO_3R^5 , $\text{C}(\text{O})\text{R}^5$, $\text{C}(\text{O})\text{OR}^5$, $\text{C}(\text{O})\text{R}^6$, $\text{C}(\text{O})\text{OR}^6$,
 $\text{C}(\text{O})\text{N}(\text{R}^6)_2$, $\text{C}(\text{O})\text{N}(\text{R}^5)_2$, $\text{C}(\text{O})\text{N}(\text{R}^5\text{R}^6)$, $\text{C}(\text{O})\text{N}(\text{OR}^6)\text{R}^6$,
 $\text{C}(\text{O})\text{N}(\text{OR}^5)\text{R}^6$, $\text{C}(\text{O})\text{N}(\text{OR}^6)\text{R}^5$, $\text{C}(\text{O})\text{N}(\text{OR}^5)\text{R}^5$, $\text{C}(\text{NOR}^6)\text{R}^6$,
 $\text{C}(\text{NOR}^6)\text{R}^5$, $\text{C}(\text{NOR}^5)\text{R}^6$, $\text{C}(\text{NOR}^5)\text{R}^5$, $\text{N}(\text{R}^6)_2$, $\text{N}(\text{R}^5)_2$, $\text{N}(\text{R}^5\text{R}^6)$,
 $\text{NR}^5\text{C}(\text{O})\text{R}^5$, $\text{NR}^6\text{C}(\text{O})\text{R}^6$, $\text{NR}^6\text{C}(\text{O})\text{R}^5$, $\text{NR}^5\text{C}(\text{O})\text{R}^6$, $\text{NR}^6\text{C}(\text{O})\text{OR}^6$,
 $\text{NR}^5\text{C}(\text{O})\text{OR}^6$, $\text{NR}^6\text{C}(\text{O})\text{OR}^5$, $\text{NR}^5\text{C}(\text{O})\text{OR}^5$, $\text{NR}^6\text{C}(\text{O})\text{N}(\text{R}^6)_2$,
 $\text{NR}^6\text{C}(\text{O})\text{NR}^5\text{R}^6$, $\text{NR}^6\text{C}(\text{O})\text{N}(\text{R}^5)_2$, $\text{NR}^5\text{C}(\text{O})\text{N}(\text{R}^6)_2$, $\text{NR}^5\text{C}(\text{O})\text{NR}^5\text{R}^6$,
 $\text{NR}^5\text{C}(\text{O})\text{N}(\text{R}^5)_2$, $\text{NR}^6\text{SO}_2\text{R}^6$, $\text{NR}^6\text{SO}_2\text{R}^5$, $\text{NR}^5\text{SO}_2\text{R}^5$, $\text{NR}^5\text{SO}_2\text{R}^6$,
 $\text{NR}^6\text{SO}_2\text{N}(\text{R}^6)_2$, $\text{NR}^6\text{SO}_2\text{NR}^5\text{R}^6$, $\text{NR}^6\text{SO}_2\text{N}(\text{R}^5)_2$, $\text{NR}^5\text{SO}_2\text{N}(\text{R}^6)_2$,
 $\text{NR}^5\text{SO}_2\text{NR}^5\text{R}^6$, $\text{NR}^5\text{SO}_2\text{N}(\text{R}^5)_2$, $\text{N}(\text{OR}^6)\text{R}^6$, $\text{N}(\text{OR}^6)\text{R}^5$, $\text{N}(\text{OR}^5)\text{R}^5$,
 or $\text{N}(\text{OR}^5)\text{R}^6$;

R^5 is a cycloaliphatic, aryl, heterocyclic, or heteroaryl ring, optionally comprising up to 3 R^1 substituents;

R^6 is H or aliphatic, wherein R^6 optionally comprises a R^7 substituent;

R^7 is a cycloaliphatic, aryl, heterocyclic, or heteroaryl ring, and each R^7 optionally comprises up to 2 substituents independently chosen from H, $(\text{C}_1\text{-C}_6)$ -straight or branched alkyl, $(\text{C}_2\text{-C}_6)$ straight or branched alkenyl or alkynyl, 1,2-methylenedioxy, 1,2-ethylenedioxy, or $(\text{CH}_2)_n\text{-Z}$;

Z is selected from halo, CN, NO_2 , CF_3 , OCF_3 , OH, S-aliphatic, S(O)-aliphatic, SO_2 -aliphatic, NH_2 , NH-aliphatic, $\text{N}(\text{aliphatic})_2$, $\text{N}(\text{aliphatic})\text{R}^8$, NHR^8 , $\text{N}(\text{R}^8)_2$, COOH, $\text{C}(\text{O})\text{O}(-\text{aliphatic})$, or O-aliphatic; and

R^8 is an amino-capping group.

[0046] The term "amino capping group" refers to a suitable chemical group that may be attached to a

nitrogen atom. The term "capping" refers to when the designated amino group is attached to a suitable chemical group (e.g., protecting group). Examples of suitable amino capping groups are described in T.W. Greene et al., Protective Groups in Organic Synthesis, 3d. Ed., John Wiley and Sons (1999); L. Fieser and M. Fieser, Fieser and Fieser's Reagents for Organic Synthesis, John Wiley and Sons (1994); L. Paquette, ed. Encyclopedia of Reagents for Organic Synthesis, John Wiley and Sons (1995) and are exemplified in certain of the specific compounds used in this invention.

[0047] According to a preferred embodiment, Y' is S.

[0048] According to one embodiment, Y' is O.

[0049] According to another embodiment, Y' is NR.

In one embodiment, R is H. Or, R is R². Or, R is R⁶.

[0050] According to another embodiment, p is 1. Or, p is 1 and X is attached to the carbon adjacent to Y' atom. Or, p is 1 and X is attached to the carbon adjacent to the ring nitrogen atom.

[0051] According to another embodiment, p is 2.

[0052] According to another embodiment, X is a bond, O, S, CH₂, CF₂, CHOR, C(O)NR, C(O)O-, NRC(O), or NR. In certain embodiments, X is a bond, O, or CH₂. In other embodiments, X is CH₂.

[0053] According to another embodiment, A is an optionally substituted C3-C7 cycloaliphatic ring. Preferably, A is an optionally substituted cyclopropyl, cyclopentyl, or cyclohexyl.


[0054] According to another embodiment, A is optionally substituted (C1-C10)aliphatic. In certain embodiments, A is optionally substituted methyl, ethyl, propyl, butyl, pentyl, or hexyl.














[0055] According to another embodiment, A is optionally substituted C6-C10 aryl ring. In one embodiment, A is optionally substituted phenyl or naphthyl.

[0056] According to another embodiment, A is optionally substituted C5-C12 heteroaryl ring. In certain embodiments, A is selected from optionally substituted triazinyl, pyrazinyl, pyrimidinyl, pyridazinyl, pyridinyl, thiadiazolyl, triazolyl, oxadiazolyl, isothiazolyl, pyrazolyl, imidazolyl, thiazolyl, oxazolyl, pyrrolyl, thienyl, furanyl, indoliziny, indolyl, isoindolyl, benzofuranyl, benzo[b]thienyl, 1H-indazolyl, benzimidazolyl, benzthiazolyl, purinyl, quinolinyl, isoquinolinyl, cinnolinyl, phthazinyl, quinazolinyl, quinoxalinyl, 1,8-naphthyridinyl, pteridinyl, acridinyl, phenazinyl, phenothiazinyl, or phenoxazinyl.


[0057] According to another embodiment, A is optionally substituted C3-C12 heterocyclic ring. In certain embodiments, A is selected from optionally substituted aziridine, oxirane, thirane, pyrrolidyl, tetrahydrofuranyl, tetrahydrothienyl, dioxolanyl, pyrrolinyl, pyranyl, pyrazolinyl, pyrazolidinyl, piperidinyl, 1,4-dioxanyl, morpholinyl, 1,4-dithianyl, thiomorpholinyl, piperazinyl, 3H-indolyl, or indolinyl.

[0058] In some embodiments, A, X, and the ring attached thereto, taken together, is selected from:



**v****vi****vii****viii****ix****x****xi****xii****xiii****xiv****xv****xvi****xvii**

or

**xviii**

wherein:

 R^{Ph} is independently R^1 , R^2 , or R^3 ; and r is 0-3. X_5 is CH_2 , $\text{C}(\text{O})$, or CHOR ; X_6 is O or NR^2 ; and R^{Ak} is C1-C6 aliphatic, optionally substituted with R^1 , R^2 , or R^3 .

[0059] In another embodiment, A, X, and the ring attached thereto, taken together, is selected from any of the rings i to xviii, wherein the sulfur atom in each of the thiazole ring is replaced with an oxygen atom (to provide the corresponding oxazole).

[0060] According to another embodiment, each B is independently selected from optionally substituted C6-C10 aryl. In certain embodiments, each B is an optionally substituted phenyl or naphthyl. Or, each B is an unsubstituted phenyl.

[0061] According to another embodiment, each B is independently selected from optionally substituted C5-C12 heteroaryl. In certain embodiments, each B is independently an optionally substituted C5-C7 heteroaryl.

[0062] According to another embodiment, each B is independently selected from triazinyl, pyrazinyl, pyrimidinyl, pyridazinyl, pyridinyl, thiadiazolyl, triazolyl, oxadiazolyl, isothiazolyl, pyrazolyl, imidazolyl, thiazolyl, oxazolyl, pyrrolyl, thienyl, furanyl, indolizinyll, indolyl, isoindolyl, benzofuranyl, benzo[b]thienyl, 1H-indazolyl, benzimidazolyl, benzthiazolyl, purinyl, quinolinyl, isoquinolinyl, cinnolinyl, phthazinyl, quinazolinyl, quinoxalinyl, 1,8-naphthyridinyl, pteridinyl, carbazolyl, acridinyl, phenazinyl, phenothiazinyl, phenoxazinyl, indenyl, naphthyl, azulinyll, or anthracenyl.

[0063] According to another embodiment, R¹ is 1,2-methylene dioxy, or 1,2-ethylenedioxy.

[0064] According to another embodiment, R¹ is R⁶, wherein R⁶ is straight chain or branched (C1-C6)alkyl or (C2-C6) alkenyl or alkynyl, optionally substituted with R⁷.

[0065] According to another embodiment, R¹ is (C1-C4 aliphatic)_n-Y, wherein n is 0 or 1, and Y is halo, CN,

NO_2 , CF_3 , OCF_3 , OH , SR^6 , S(O)R^6 , SO_2R^6 , NH_2 , NHR^6 , $\text{N(R}^6)_2$, NR^6R^8 , COOH , COOR^6 or OR^6 ;

[0066] According to another embodiment, R^1 is selected from halo, CF_3 , NH_2 , NH(C1-C4 alkyl) , NHC(O)CH_3 , OH , O(C1-C4 alkyl) , OPh , O-benzyl , S-(C1-C4 alkyl) , C1-C4 aliphatic , CN , methylenedioxy, ethylenedioxy, $\text{SO}_2\text{NH(C1-C4 alkyl)}$, or $\text{SO}_2\text{N(C1-C4 alkyl)}_2$.

[0067] According to another embodiment, R^1 is selected from methyl, n-propyl, i-propyl, t-butyl, halo, CF_3 , NH_2 , $\text{NH(CH}_3)$, NHC(O)CH_3 , OH , OCH_3 , OPh , O-benzyl , $\text{S-(C}_2\text{H}_5)$, S-CH_3 , NO_2 , CN , methylenedioxy, $\text{SO}_2\text{NH(n-propyl)}$, or $\text{SO}_2\text{N(n-propyl)}_2$.

[0068] According to one embodiment, R^2 is a straight chain or branched (C1-C6)alkyl or (C2-C6) alkenyl or alkynyl, optionally substituted with R^1 , R^4 , or R^5 . In certain embodiments, R^2 is a straight chain or branched (C1-C4)alkyl or (C2-C4) alkenyl or alkynyl, optionally substituted with R^1 , R^4 , or R^5 .

[0069] According to another embodiment, R^3 is optionally substituted phenyl, naphthyl, C5-C10 heteroaryl or C3-C7 heterocyclyl. In certain embodiments, R^3 is an optionally substituted phenyl, C5-C6 heteroaryl, or C3-C6 heterocyclyl.

[0070] According to one embodiment, R^4 is selected from OR^5 , OR^6 , SR^5 , SR^6 , NR^5COR^5 , NR^5COR^6 , NR^6COR^5 , or NR^6COR^6 .

[0071] According to one embodiment, R^5 is C5-C6 cycloalkyl, C6 or C10 aryl, C5-C10 heteroaryl or C3-C7 heterocyclyl, optionally substituted with up to 2 R^1 . In certain embodiments, R^5 is an optionally substituted cyclohexyl, phenyl, C5-C6 heteroaryl, or C3-C6 heterocyclyl.

[0072] According to one embodiment, R^6 is H.


[0073] According to another embodiment, R^6 is a straight chain or branched (C1-C6)alkyl or (C2-C6 alkenyl) or alkynyl, optionally substituted with R^7 .

[0074] According to another embodiment, R^6 is a straight chain or branched (C1-C6)alkyl or (C2-C6 alkenyl) or alkynyl.


[0075] According to one embodiment, R^7 is C5-C6 cycloalkyl, phenyl, naphthyl, C5-C10 heteroaryl or C3-C7 heterocyclyl, optionally substituted with straight chain or branched (C1-C6)alkyl or (C2-C6 alkenyl) or alkynyl. Or, R^7 is C5-C6 cycloalkyl, phenyl, naphthyl, C5-C10 heteroaryl or C3-C7 heterocyclyl, optionally substituted with 1-2-methylenedioxy, 1,2-ethylenedioxy, or $(CH_2)_n-Z$. In certain embodiments, R^7 is an optionally substituted cyclohexyl, phenyl, C5-C6 heteroaryl, or C3-C6 heterocyclyl.

[0076] According to a preferred embodiment, R^8 is acetyl, arylsulfonyl or alkylsulfonyl.

[0077] In another embodiment, Q in compounds of formula I is selected from:



[0078] In one embodiment, the present invention provides compounds having formula I-a:



[0079] In certain embodiments, p is 1. Or, p is 2.

[0080] In certain embodiments, X is a bond, O, CH_2 , $CHOH$, $C(O)$, or $C(O)O$. Or, X is a bond, O, or CH_2 .


[0081] In certain embodiments, p is 1 and X is a bond. In one embodiment, p is 1, and X-A is attached to the carbon adjacent to the ring nitrogen atom. Or, p is 1, and X-A is attached to the carbon adjacent to the Y'.

[0082] In certain embodiments, p is 1, X is CH₂, CHOH, or C(O), and A is an optionally substituted phenyl.

[0083] In certain embodiments, X is a bond, and A is an optionally substituted phenyl.

[0084] In certain embodiments, p is 2, each X is a bond, and each A is an optionally substituted phenyl.

[0085] In certain embodiments, each B is independently and optionally substituted ring selected from:



wherein X₃ is O, S, or NR.

[0086] Preferred substituents on B include C1-C4 alkyl, -O-C1-C4 alkyl, CN, halo, COOH, -C(O)NH₂, -C(O)O(C1-C4 alkyl), -C(O)NH(C1-C4 alkyl), -C(O)N(C1-C4 alkyl)₂, or phenyl optionally substituted with up to two substituents selected from C1-C4 alkyl, -O-C1-C4 alkyl, CN, halo, COOH, -C(O)NH₂, -C(O)O(C1-C4 alkyl), -C(O)NH(C1-C4 alkyl), -C(O)N(C1-C4 alkyl)₂.

[0087] In certain embodiments, each B is independently an optionally substituted ring selected from ring **i**, **iii**,

iv, v, or vi. Or, wherein B is an optionally substituted ring vii.

[0088] In some embodiments, B is independently ring x optionally substituted with up to two substituents selected from R^1 or phenyl optionally substituted with up to two R^1 . Preferably, B is phenyl optionally substituted with up to two substituents selected from C1-C4 alkyl, -O-C1-C4 alkyl, CN, halo, COOH, -C(O)NH₂, -C(O)O(C1-C4 alkyl), -C(O)NH(C1-C4 alkyl), -C(O)N(C1-C4 alkyl)₂, or phenyl optionally substituted with up to two substituents selected from C1-C4 alkyl, -O-C1-C4 alkyl, CN, halo, COOH, -C(O)NH₂, -C(O)O(C1-C4 alkyl), -C(O)NH(C1-C4 alkyl), -C(O)N(C1-C4 alkyl)₂.

[0089] Or, each B is an optionally substituted ring selected from ring xi, xii, xiii, or xiv.

[0090] In certain embodiments, p is 1. Or, p is 2.

[0091] In some embodiments, R is hydrogen.

[0092] In some embodiments, Y' is S. Or, Y' is O.

[0093] In certain embodiments, X is a bond and A is optionally substituted phenyl. In certain embodiments, A is attached to the carbon atom adjacent to the nitrogen ring atom.

[0094] In certain embodiments, A is phenyl optionally substituted with up to two substituents selected from C1-C4 alkyl, C1-C4 alkoxy, cyano, halo, N-pyrrolidinyl, N-piperidinyl, or methylenedioxy.

[0095] In certain embodiments, A is phenyl, methoxyphenyl, dimethoxyphenyl, cyanophenyl, N-pyrrolidinylphenyl, methylenedioxyphenyl, halophenyl, methylphenyl, or dimethylphenyl.

[0096] In certain embodiments, A is phenyl, 3-methoxyphenyl, 2-methoxyphenyl, 4-chlorophenyl, 4-cyanophenyl, 4-(N-pyrrolidinyl)phenyl, 4-tolyl, 3,4-methylenedioxyphenyl, 3-chlorophenyl, 2,4-


dimethoxyphenyl, 2-chlorophenyl, 4-bromophenyl, or 2,4-dimethylphenyl.

[0097] In certain other embodiments, A is C3-C10 cycloaliphatic ring. Exemplary ring include cyclopropyl, cyclopentyl, cyclohexyl, cycloheptyl, or adamantyl. In certain embodiments, A is cyclohexyl or adamantyl.

[0098] In certain embodiments the compounds of the present invention have one of more of the following features:

- a) R is hydrogen;
- b) Y' is S;
- c) A is phenyl optionally substituted with up to two substituents selected from C1-C4 alkyl, C1-C4 alkoxy, cyano, halo, N-pyrrolidinyl, N-piperidinyl, or methylenedioxy; and
- d) B is phenyl optionally substituted with up to two substituents selected from C1-C4 alkyl, -O-C1-C4 alkyl, CN, halo, COOH, -C(O)NH₂, -C(O)O(C1-C4 alkyl), -C(O)NH(C1-C4 alkyl), -C(O)N(C1-C4 alkyl)₂, or phenyl optionally substituted with up to two substituents selected from C1-C4 alkyl, -O-C1-C4 alkyl, CN, halo, COOH, -C(O)NH₂, -C(O)O(C1-C4 alkyl), -C(O)NH(C1-C4 alkyl), -C(O)N(C1-C4 alkyl)₂.

[0099] In one embodiment, the present invention provides compounds having formula **I-b**:



I-b.

[00100] In certain embodiments, p is 1 and X is a bond. In one embodiment, p is 1, and X-A is attached to the carbon adjacent to the ring nitrogen atom. Or, p is 1, and X-A is attached to the carbon adjacent to the Y'.

[00101] In certain embodiments, p is 1, X is CH₂, CHOH, or C(O), preferably CH₂, and A is an optionally substituted phenyl.


[00102] In certain embodiments, X is a bond, and A is an optionally substituted phenyl.

[00103] In certain embodiments, p is 2, each X is a bond, and each A is an optionally substituted phenyl.

[00104] In certain embodiments, said C1-C6 aliphatic is C1-C4 straight or branched alkylidene. Exemplary alkylidenes include -CH₂-, -CH(Me)-, -C(Me)₂-, -CH(Et)-, -C(Et)₂-, or -CH₂-CH(Me)-.

[00105] In certain embodiments, B is selected from optionally substituted C3-C8 cycloalkyl, phenyl, piperidyl, or pyrrolidinyl. Preferably, B is phenyl, cyclopentyl, cyclohexyl, or piperidyl, optionally substituted with up to two R¹ substituents.

[00106] In some embodiments, said (C1-C6 aliphatic)-B in formula I-b is selected from:



wherein:

Ak is C1-C6 straight or branched alkylidene;

X₄ is CH₂, O or S;

Ar' is phenyl optionally substituted with up to two R¹; and

B is optionally substituted with up to two R¹.


[00107] In some embodiments, Ak is selected from CH₂, CH(CH₃), C(CH₃)₂, CH(Et), C(Et)₂, CH(n-propyl), CH(i-Pr), CH(n-butyl), CH(but-2-yl), or CH(t-butyl).

[00108] In some embodiments, Ar' is phenyl optionally substituted with halo, C1-C4 alkyl, or O-(C1-C4 alkyl).

[00109] In some embodiments, X_4 is S. Or, X_4 is O.

[00110] In some embodiments, R is hydrogen.

[00111] In one embodiment, the present invention provides compounds of formula I-f:



wherein:

Y' is O or S;

X_1 is O, S, or NR;

R^M is C1-C6 aliphatic or phenyl, wherein R^M is optionally substituted with up to two substituents independently selected from R^1 , R^2 , or R^3 ;

R^N is C1-C6 aliphatic or a 3-7 membered monocyclic, saturated, unsaturated or aromatic ring containing 0-4 heteroatoms in each ring, wherein each said heteroatom is independently selected from N, NH, S, or O;

wherein R^N is optionally substituted with up to 4 substituents independently selected from R^1 , R^2 , R^3 , R^4 , or R^5 .

[00112] In certain embodiments, p is 1 and X is a bond. In one embodiment, p is 1, and X-A is attached to the carbon adjacent to the ring nitrogen atom. Or, p is 1, and X-A is attached to the carbon adjacent to the Y' .

[00113] In certain embodiments, p is 1, X is CH_2 , CHOH, or C(O), preferably CH_2 , and A is an optionally substituted phenyl.

[00114] In certain embodiments, X is a bond, and A is an optionally substituted phenyl.

[00115] In certain embodiments, p is 2, each X is a bond, and each A is an optionally substituted phenyl.

[00116] In some embodiments, X_1 is NH or N(C1-C4 alkyl). Or, X_1 is O.

[00117] In some embodiments, R^M is optionally substituted phenyl.


[00118] In some embodiments, R^M is C1-C6 alkyl, optionally substituted with phenyl. In some embodiments, R^M is C1-C4 alkyl.

[00119] In some embodiments, R^N is optionally substituted C3-C7 cycloaliphatic, phenyl, or benzyl.

[00120] In some embodiments, R^N is C1-C6 aliphatic.

[00121] In some embodiments, R is hydrogen.

[00122] In one embodiment, the present invention provides a compound of formula I-g:



wherein:

Y' is O or S;

R^P is C1-C8 aliphatic optionally substituted with up to two substituents independently selected from R^1 , R^2 , or R^3 .

[00123] In certain embodiments, p is 1 and X is a bond. In one embodiment, p is 1, and X-A is attached to the carbon adjacent to the ring nitrogen atom. Or, p is 1, and X-A is attached to the carbon adjacent to the Y' .

[00124] In certain embodiments, p is 1, X is CH_2 , CHOH, or C(O), preferably CH_2 , and A is an optionally substituted phenyl.

[00125] In certain embodiments, X is a bond, and A is an optionally substituted phenyl.

[00126] In certain embodiments, p is 2, each X is a bond, and each A is an optionally substituted phenyl.

[00127] In some embodiments, R^p is C1-C4 alkyl, optionally substituted with up to two R^1 .

[00128] In some embodiments, R^p is selected from ethyl, n-propyl, i-propyl, n-butyl, but-2-yl, or t-butyl, isoamyl, optionally substituted with halo, CN, COOH, or CONH₂.

[00129] In some embodiments, R is hydrogen.


[00130] In certain embodiments, p is 1. Or, p is 2.

[00131] In certain embodiments, p is 2, and each A is optionally substituted phenyl. Or, p is 2, and each A is phenyl.

[00132] In certain embodiments, compounds of formula 1-g have one or more of the following features:

- a) Y' is S;
- b) R is hydrogen;
- c) p is 2 and each A is phenyl;
- d) R^p is isoamyl, t-butyl, ethyl, isopropyl, n-propyl, 1-carboxy-prop-3-yl, or 1-carboxy-2-methyl-prop-3-yl.

[00133] In one embodiment, the present invention provides compounds of formula I-h:



wherein:

Y' is O or S;

X₂ is O, S, or NR;

R^Q is C1-C6 aliphatic or phenyl, optionally substituted with up to two substituents independently selected from R¹, R², or R³.

[00134] In certain embodiments, p is 1 and X is a bond. In one embodiment, p is 1, and X-A is attached to

the carbon adjacent to the ring nitrogen atom. Or, p is 1, and X-A is attached to the carbon adjacent to the Y'.

[00135] In certain embodiments, p is 1, X is CH₂, CHOH, or C(O), preferably CH₂, and A is an optionally substituted phenyl.

[00136] In certain embodiments, X is a bond, and A is an optionally substituted phenyl.


[00137] In certain embodiments, p is 2, each X is a bond, and each A is an optionally substituted phenyl.

[00138] In some embodiments, X₂ is S. Or, X₂ is O.

[00139] In some embodiments, R⁰ is C1-C4 alkyl, optionally substituted with up to three R¹.

[00140] In some embodiments, R⁰ is phenyl optionally substituted with C1-C4 alkyl, or R¹.

[00141] According to another preferred embodiment, the methods of the present invention employ a compound having formula (IA):




wherein:

Y' is O, S, or NR;

X is a bond, CH₂, CHOR, C(O)O, C(O), NR, or O;

A is aliphatic, aryl, heteroaryl, heterocyclic, or cycloaliphatic;

Q is selected from:



each B is independently selected from 3-7 membered monocyclic or 8-14 membered bicyclic or tricyclic,

saturated, unsaturated or aromatic ring containing 0-4 heteroatoms in each ring, wherein each said heteroatom is independently selected from N, NH, S, or O;

R is H, R², or R⁶;

wherein each A and B is independently and optionally substituted with up to 4 substituents independently selected from R¹, R², R³, R⁴, or R⁵; and

R¹, R², R³, R⁴, or R⁵ are as defined above for formula (I).

[00142] According to one embodiment of formula (IA), Y' is S.

[00143] According to another embodiment of formula (IA), Y' is O.

[00144] According to another embodiment of formula (IA), Y' is NR.

[00145] In one embodiment, X is a bond, CH₂, NR, or O;

[00146] According to another embodiment of formula (IA), X is CH₂. According to another embodiment of formula (IA), X is CF₂. According to yet another embodiment of formula (IA), X is a bond. According to yet another embodiment of formula (IA), X is O. According to yet another embodiment of formula (IA), X is NR.

[00147] According to another embodiment of formula (IA), A is phenyl or a 5-6 membered heteroaryl, preferably phenyl, wherein A is optionally substituted with up to 3 substituents selected from R¹, R², R³, or R⁴.

[00148] According to one embodiment of formula (IA), Q is B. Alternatively, Q is -(C1-C6)- aliphatic-B. Or, Q is CH(B)₂. According to another embodiment, Q is C(B)₃. Preferably, B is phenyl.


[00149] In one embodiment, X is a bond, -CHOR-, or -C(O)-,

[00150] Exemplary compounds of formula (IA) are those wherein:

- (i) X is a bond, CH₂, or O;
- (i) A is optionally substituted phenyl;
- (iii) Q is diphenylmethyl.

[00151] Exemplary compounds of formula (IA) useful in the methods of the present invention are as shown below in Table 1.

[00152] According to another embodiment, the methods of the present invention employ a compound having formula (IB):




wherein:

Y' is O, S, or NR;

X is a bond, CH₂, CHOR, C(O), C(O)O, NR, or O;

A is aliphatic, aryl, heteroaryl, heterocyclic, or cycloaliphatic;

Q is selected from:



each B is independently selected from 3-7 membered monocyclic or 8-14 membered bicyclic or tricyclic, saturated, unsaturated or aromatic ring containing 0-4 heteroatoms in each ring, wherein each said heteroatom is independently selected from N, NH, S, or O;

R is H, R², or R⁶;

wherein each A and B is independently and optionally substituted with up to 4 substituents independently selected from R^1 , R^2 , R^3 , R^4 , or R^5 ; and

R^1 , R^2 , R^3 , R^4 , or R^5 are as defined above for formula (I).

[00153] According to one embodiment of formula (IB), Y' is S.

[00154] According to another embodiment of formula (IB), Y' is O.

[00155] According to another embodiment of formula (IB), Y' is NR.

[00156] According to one embodiment of formula (IB), X is a bond, CH_2 , NR, or O, Or, X is a bond or CH_2 . According to another embodiment of formula (IB), X is CF_2 . According to yet another embodiment of formula (IB), X is a bond. According to yet another embodiment of formula (IB), X is O. According to yet another embodiment of formula (IB), X is NR.


[00157] According to another embodiment of formula (IB), A is phenyl or a 5-6 membered heteroaryl, preferably phenyl, wherein A is optionally substituted with up to 3 substituents selected from R^1 , R^2 , R^3 , or R^4 .

[00158] According to another embodiment of formula (IB), Q is B. Alternatively, Q is $-(C1-C6)-$ aliphatic-B. Or, Q is $CH(B)_2$. According to another embodiment, Q is $C(B)_3$. Preferably, B is phenyl.

[00159] Preferred compounds of formula IB are those wherein:

- (i) X is a bond, CH_2 , or O;
- (ii) A is optionally substituted phenyl;
- (iii) Q is diphenylmethyl.

[00160] According to another embodiment, the methods of the present invention employ a compound having formula (IC):




wherein:

Y' is O, S, or NR;

each X is independently a bond, CH₂, -CHOR-, -C(O)O-, -C(O)-, NR, or O;

A is aliphatic, aryl, heteroaryl, heterocyclic, or cycloaliphatic;

Q is selected from:



each B is independently selected from 3-7 membered monocyclic or 8-14 membered bicyclic or tricyclic, saturated, unsaturated or aromatic ring containing 0-4 heteroatoms in each ring, wherein each said heteroatom is independently selected from N, NH, S, or O;

R is H, R², or R⁶;

wherein each A and B is independently and optionally substituted with up to 4 substituents independently selected from R¹, R², R³, R⁴, or R⁵; and

R¹, R², R³, R⁴, or R⁵ are as defined above for formula (I).

[00161] According to one embodiment of formula (IC), Y' is S.

[00162] According to another embodiment of formula (IC), Y' is O.

[00163] According to another embodiment of formula (IC), Y' is NR.

[00164] According to another embodiment of formula (IC), X is a bond or CH₂. According to another embodiment of formula (IC), X is CH₂. According to yet another embodiment of formula (IC), X is a bond. Or, X is -CHOR-. Or, X is -C(O)-. According to yet another embodiment of formula (IC), X is O. According to yet another embodiment of formula (IC), X is NR.


[00165] According to another embodiment of formula (IC), A is phenyl or a 5-6 membered heteroaryl, preferably phenyl, wherein A is optionally substituted with up to 3 substituents selected from R¹, R², R³, or R⁴.

[00166] According to another embodiment of formula (IC), Q is B. Alternatively, Q is -(C1-C6)- aliphatic-B. Or, Q is CH(B)₂. According to another embodiment, Q is C(B)₃. Preferably, B is phenyl.

[00167] Exemplary compounds of formula (IC) are those wherein:

- (i) each X is a bond, -CHOR-, or -C(O)-;
- (ii) each A is optionally substituted phenyl, (C1-C6)aliphatic, or CF₃;
- (iii) Q is optionally substituted phenyl, (C1-C6)aliphatic, or diphenylmethyl.

[00168] According to another embodiment, the present invention provides a compound having formula (II):



or a pharmaceutically acceptable salt thereof;
wherein:

X_1 is a bond, O, S, CHOR, C(O), C(O)O, CF_2 , CH_2 , or NR;

R is H or R^2

A_1 is (C2-C10) aliphatic, aryl, heteroaryl, heterocyclic, or cycloaliphatic;

each B_1 is independently selected from 3-7 membered monocyclic, saturated, unsaturated or aromatic ring containing 0-4 heteroatoms selected from N, NH, S, or O;

wherein each A_1 is optionally substituted with up to 4 substituents independently selected from R^1 , R^2 , R^3 , R^4 , or R^5 ;

R^1 is oxo, R^6 or ((C1-C4)aliphatic) $_n$ -Y;

n is 0 or 1;

Y is halo, CN, NO_2 , CF_3 , OCF_3 , OH, SR^6 , $S(O)R^6$, SO_2R^6 , NH_2 , NHR^6 , $N(R^6)_2$, NR^6R^8 , $N(R^8)_2$, COOH, $COOR^6$ or OR^6 ; or

two R^1 on adjacent ring atoms, taken together, form 1,2-methylenedioxy or 1,2-ethylenedioxy;

R^2 is aliphatic, wherein each R^2 optionally comprises up to 2 substituents independently selected from R^1 , R^4 , or R^5 ;

R^3 is a cycloaliphatic, aryl, heterocyclic, or heteroaryl ring optionally comprising up to 3 substituents, independently selected from R^1 , R^2 , R^4 or R^5 ;

R^4 is OR^5 , OR^6 , $OC(O)R^6$, $OC(O)R^5$, $OC(O)OR^6$, $OC(O)OR^5$, $OC(O)N(R^6)_2$, $OC(O)N(R^5)_2$, $OC(O)N(R^6R^5)$, SR^6 , SR^5 , $S(O)R^6$, $S(O)R^5$, SO_2R^6 , SO_2R^5 , $SO_2N(R^6)_2$, $SO_2N(R^5)_2$, $SO_2NR^5R^6$, SO_3R^6 , SO_3R^5 , $C(O)R^5$, $C(O)OR^5$, $C(O)R^6$, $C(O)OR^6$, $C(O)N(R^6)_2$, $C(O)N(R^5)_2$, $C(O)N(R^5R^6)$, $C(O)N(OR^6)R^6$, $C(O)N(OR^5)R^6$, $C(O)N(OR^6)R^5$, $C(O)N(OR^5)R^5$, $C(NOR^6)R^6$, $C(NOR^6)R^5$, $C(NOR^5)R^6$, $C(NOR^5)R^5$, $N(R^6)_2$, $N(R^5)_2$, $N(R^5R^6)$, $NR^5C(O)R^5$, $NR^6C(O)R^6$, $NR^6C(O)R^5$, $NR^5C(O)R^6$, $NR^6C(O)OR^6$, $NR^5C(O)OR^6$, $NR^6C(O)OR^5$, $NR^5C(O)OR^5$, $NR^6C(O)N(R^6)_2$, $NR^6C(O)NR^5R^6$, $NR^6C(O)N(R^5)_2$, $NR^5C(O)N(R^6)_2$, $NR^5C(O)NR^5R^6$, $NR^5C(O)N(R^5)_2$, $NR^6SO_2R^6$, $NR^6SO_2R^5$, $NR^5SO_2R^5$, $NR^5SO_2R^6$, $NR^6SO_2N(R^6)_2$, $NR^6SO_2NR^5R^6$, $NR^6SO_2N(R^5)_2$, $NR^5SO_2N(R^6)_2$, $NR^5SO_2NR^5R^6$, $NR^5SO_2N(R^5)_2$, $N(OR^6)R^6$, $N(OR^6)R^5$, $N(OR^5)R^5$, or $N(OR^5)R^6$;

R^5 is a cycloaliphatic, aryl, heterocyclic, or heteroaryl ring, optionally comprising up to 3 R^1 substituents;

R^6 is H or aliphatic, wherein R^6 optionally comprises a R^7 substituent;

R^7 is a cycloaliphatic, aryl, heterocyclic, or heteroaryl ring, and each R^7 optionally comprises up to 2 substituents independently chosen from H, $(C_1-C_6)-$

straight or branched alkyl, (C₂-C₆) straight or branched alkenyl or alkynyl, 1,2-methylenedioxy, 1,2-ethylenedioxy, or (CH₂)_n-Z;

Z is selected from halo, CN, NO₂, CF₃, OCF₃, OH, S-aliphatic, S(O)-aliphatic, SO₂-aliphatic, NH₂, NH-aliphatic, N(aliphatic)₂, N(aliphatic)R⁸, NHR⁸, N(R⁸)₂, COOH, C(O)O(-aliphatic, or O-aliphatic; and

R⁸ is an amino capping group.

provided that:

- (i) when both B₁ are simultaneously phenyl and X₁ is CH₂, then A is not 4-fluoro-phenyl, 4-phenyl-piperidyl, phenyl, 2,4-dichloro-phenyl, 4-methoxy-phenyl, 3,4-dichloro-phenyl, 2,5-dichloro-phenyl, 4-nitro-phenyl, 4-bromo-phenyl, 4-methyl-phenyl, 2-chloro-phenyl, 1-naphthyl, 3-trifluoromethyl-phenyl, 2,3-dichlorophenyl, N-morpholinyl, 4-chloro-phenyl, 3-chloro-phenyl, or 3-nitro-phenyl;
- (ii) when X₁ is a bond or CH₂, one B₁ is a substituted phenyl and the other B₁ is cycloaliphatic, then A₁ is not (C₂-C₈)aliphatic; and
- (iii) when X₁ is a bond, then A₁ is not an optionally substituted 6-membered heteroaryl ring with 1-3 nitrogen ring atoms.

[00169] According to one embodiment of formula (II), X₁ is a bond, O, S, CF₂, CH₂, or NR. X₁ is CH₂, CF₂, or O. Or X₁ is CH₂ or O. In certain embodiments, X₁ is CH₂.

[00170] According to another embodiment, A₁ is an optionally substituted C₃-C₇ cycloaliphatic ring. In certain embodiments, A₁ is an optionally substituted cyclopropyl, cyclopentyl, or cyclohexyl.

[00171] According to another embodiment, A₁ is optionally substituted (C₁-C₁₀)aliphatic. In certain

embodiments, A₁ is optionally substituted methyl, ethyl, propyl, butyl, pentyl, or hexyl.

[00172] According to another embodiment, A₁ is optionally substituted C6-C10 aryl ring. In certain embodiments, A₁ is optionally substituted phenyl or naphthyl.

[00173] According to another embodiment, A₁ is optionally substituted C5-C12 heteroaryl ring. In certain embodiments, A is selected from optionally substituted triazinyl, pyrazinyl, pyrimidinyl, pyridazinyl, pyridinyl, thiadiazolyl, triazolyl, oxadiazolyl, isothiazolyl, pyrazolyl, imidazolyl, thiazolyl, oxazolyl, pyrrolyl, thienyl, furanyl, indoliziny, indolyl, isoindolyl, benzofuranyl, benzo[b]thienyl, 1H-indazolyl, benzimidazolyl, benzthiazolyl, purinyl, quinolinyl, isoquinolinyl, cinnolinyl, phthazinyl, quinazolinyl, quinoxalinyl, 1,8-naphthyridinyl, pteridinyl, acridinyl, phenazinyl, phenothiazinyl, or phenoxazinyl.

[00174] According to another embodiment, A₁ is optionally substituted C3-C12 heterocyclic ring. In certain embodiments, A₁ is selected from optionally substituted aziridine, oxirane, thirane, pyrrolidyl, tetrahydrofuranyl, tetrahydrothienyl, dioxolanyl, pyrrolinyl, pyranyl, pyrazolinyl, pyrazolidinyl, piperidinyl, 1,4-dioxanyl, morpholinyl, 1,4-dithianyl, thiomorpholinyl, piperazinyl, 3H-indolyl, or indolinyl.

[00175] According to another embodiment of formula (II), each B₁ is independently selected from optionally substituted C6-C10 aryl. In certain embodiments, each B₁ is independently an optionally substituted phenyl or naphthyl. Or, each B₁ is an unsubstituted phenyl.

[00176] According to another embodiment, each B₁ is independently selected from optionally substituted C5-C12

heteroaryl ring. In certain embodiments, each B₁ is independently and optionally substituted C5-C7 heteroaryl ring. Or, each B₁ is independently selected from optionally substituted triazinyl, pyrazinyl, pyrimidinyl, pyridazinyl, pyridinyl, thiadiazolyl, triazolyl, oxadiazolyl, isothiazolyl, pyrazolyl, imidazolyl, thiazolyl, oxazolyl, pyrrolyl, thienyl, furanyl, indolizinyll, indolyl, isoindolyl, benzofuranyl, benzo[b]thienyl, 1H-indazolyl, benzimidazolyl, benzthiazolyl, purinyl, quinolinyl, isoquinolinyl, cinnolinyl, phthazinyl, quinazolinyl, quinoxalinyl, 1,8-naphthyridinyl, pteridinyl, carbazolyl, acridinyl, phenazinyl, phenothiazinyl, phenoxazinyl, indenyl, naphthyl, azulinyll, or anthracenyl.

[00177] According to another embodiment, each B₁ is independently an optionally substituted 3-12 membered heterocyclic ring having up to 4 heteroatoms selected from O, S, or NR. In certain embodiments, each B₁ is independently selected from optionally substituted aziridine, oxirane, thiirane, pyrrolidyl, tetrahydrofuranyl, tetrahydrothienyl, dioxolanyl, pyrrolinyl, pyranyl, pyrazolinyl, pyrazolidinyl, piperidinyl, 1,4-dioxanyl, morpholinyl, 1,4-dithianyl, thiomorpholinyl, piperazinyl, 3H-indolyl, or indolinyl.


[00178] Embodiments of R¹, R², R³, R⁴, R⁵, R⁶, R⁷, R⁸, and Z in compound of formula (II) are as described above for compound of formula (I).

[00179] In certain embodiments, one B₁ is phenyl with up to two R¹ substituents, and the other B₁ is selected from pyrazolyl, pyrrolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, or piperazinyl, optionally substituted with up to two R¹ substituents.

[00180] In certain embodiments, one B₁ is phenyl, and the other B₁ is selected from 1,2-pyrazol-1-yl, 1-

piperidinyl, 2-carboethoxy-1-piperidinyl, 4-morpholinyl, 3-carboethoxy-1-piperidinyl, 3-methyl-1-piperidinyl, 2-methyl-1-pyrrolidinyl, 3-hydroxymethyl-1-piperidinyl, 4-carboethoxy-1-piperidinyl, 4-methyl-1-piperidinyl, 1-pyrrolidinyl, 4-(pyrimidin-2-yl)-1-piperazinyl, or 4-hydroxy-piperidinyl.

[00181] According to another embodiment, the present invention provides a compound of formula IIA:



IIA

wherein:

Y' is O or S;

B is a 3-8 membered, saturated, monocyclic, ring having 0-4 heteroatoms selected from O, S, or N; and

ring G and B are optionally substituted with up to four substituents independently selected from R¹, R², R³, R⁴, or R⁵;

provided that when Y' is S, and:

a) when B is cyclohexyl, tetrahydrofuran-2-yl, or cyclopropyl, and ring G has 1-3 halo substituents, then ring G has at least one additional substituent other than halo; and

b) when B is tetrahydrofuran-2-yl, then ring G is not phenyl, trifluoromethylphenyl, methoxyphenyl, or tolyl;

c) when B is cyclohexyl, then ring G is not phenyl or trifluoromethylphenyl.

[00182] According to one embodiment, B is tetrahydrofuranyl, piperidyl, morpholinyl, or thiomorpholinyl.

[00183] According to another embodiment, B is C3-C8 saturated, carbocyclic, monocyclic ring. Exemplary rings


include cyclopropyl, cyclopentyl, cyclohexyl, or cycloheptyl.

[00184] According to another embodiment, ring G is phenyl optionally substituted with R^1 . Preferably, ring G is optionally substituted with up to two substituents selected from halo, cyano, C1-C4 alkyl, or O-(C1-C4 alkyl).

[00185] In one embodiment, compounds of formula IIA have one or more of the following features:

- a) Y' is S;
- b) ring G is halo-substituted phenyl;
- c) B is phenyl optionally substituted with halo, cyano, C1-C4 alkyl, or O-(C1-C4 alkyl).

[00186] According to another embodiment, the present invention provides a compound of formula IIB:



IIB;

wherein:

Y' is S or O;

R is H, R^2 , or R^6 ;

R^B is C1-C6 aliphatic or a 3-7 membered monocyclic or 8-14 membered bicyclic or tricyclic, saturated, unsaturated or aromatic ring containing 0-4 heteroatoms in each ring, wherein each said heteroatom is independently selected from N, NH, S, or O; wherein each of ring G, ring H, and R^B is independently and optionally substituted with up to 4 substituents selected from R^1 , R^2 , R^3 , R^4 , or R^5 ; provided that when Y is S, and:

- a) when R^B is hydrogen, and ring G and ring H both have 1-3 halo substituents, then at least one of ring G and ring H has an additional substituent other than halo;

b) when R^B is hydrogen and ring H is unsubstituted phenyl, then ring G is not phenyl or phenyl substituted with methyl, CF_3 , -OMe, NO_2 , or 1-3 halo;

c) when R^B is hydrogen and ring H is phenyl with methyl, 1-2 methoxy or 1-2 halo substituents, then ring G is not phenyl substituted with CF_3 or 1-2 halo;

d) when R^B is methyl and ring H phenyl substituted with butyl, then ring G is not phenyl substituted with methyl, or 1-2 halo; and

e) when R^B and ring H are both unsubstituted phenyl, then ring G is not unsubstituted phenyl, or phenyl substituted with methyl, CF_3 , OMe, NO_2 , or 1-2 halo.

[00187] In certain embodiments, ring G is phenyl optionally substituted with up to two R^1 . Exemplary R^1 includes C1-C4 alkyl, O-(C1-C4 alkyl), halo, or cyano.

[00188] In one embodiment, R^B is C1-6 aliphatic optionally substituted with up to 4 substituents selected from R^1 , R^2 , R^3 , R^4 , or R^5 . In certain embodiments, R^B is C1-C4 alkyl optionally substituted with up to 2 substituents selected from R^1 . Exemplary R^B include methyl, ethyl, n-propyl, isopropyl, sec-butyl, n-butyl, or t-butyl.


[00189] In other embodiments, R^B is a 3-7 membered monocyclic saturated, unsaturated or aromatic ring containing 0-4 heteroatoms optionally substituted with up to 4 substituents selected from R^1 , R^2 , R^3 , R^4 , or R^5 . In certain embodiments, R^B is a 3-7 membered monocyclic saturated, carbocyclic ring optionally substituted with up to 2 substituents selected from R^1 . Exemplary rings include cyclopropyl, cyclopentyl, cyclohexyl, or cycloheptyl.

[00190] In other embodiments, R^B is a 3-7 membered monocyclic saturated, unsaturated or aromatic ring

containing 1-3 heteroatoms optionally substituted with up to 4 substituents selected from R^1 , R^2 , R^3 , R^4 , or R^5 . In certain embodiments, R^B is a 3-7 membered monocyclic saturated ring containing 1-3 heteroatoms optionally substituted with up to 2 substituents selected from R^1 . Exemplary rings include piperidinyl, morpholinyl, or thiomorpholinyl.

[00191] In other embodiments, R^B is a 8-14 membered bicyclic or tricyclic, saturated, unsaturated or aromatic ring containing 0-4 heteroatoms in each ring, wherein each said heteroatom is independently selected from N, NH, S, or O, optionally substituted with up to 4 substituents selected from R^1 , R^2 , R^3 , R^4 , or R^5 .

[00192] According to another embodiment, the present invention provides compounds of formula IIC:



wherein:

Y' is O or S;

X_7 is O, S, or NR' ;

R' is hydrogen, R^2 , or R^6 ;

R is hydrogen, R^2 , or R^6 ;

R^B is C1-6 aliphatic or a 3-7 membered monocyclic or 8-14 membered bicyclic or tricyclic, saturated, unsaturated or aromatic ring containing 0-4 heteroatoms in each ring, wherein each said heteroatom is independently selected from N, NH, S, or O;

R^C is C1-C6 aliphatic;

wherein each of ring G, R^B , and R^C is independently and optionally substituted with up to 4 substituents selected from R^1 , R^2 , R^3 , R^4 , or R^5 , as defined above.

[00193] According to one embodiment, X_7 is O. Or, X_7 is S. Or, X_7 is NR' .

[00194] According to one embodiments, R^C is C1-C6 alkyl, optionally substituted with up to two substituents selected from R^1 , R^2 , R^3 , R^4 , or R^5 . Or, R^C is C1-C6 alkyl. Exemplary R^C includes methyl, ethyl, isopropyl, n-propyl, n-butyl, sec-butyl, or t-butyl.


[00195] According to another embodiment, R^B is phenyl optionally substituted with up to two R^1 substituents. Or, R^B is phenyl.

[00196] According to another embodiment, R^B is C1-C6 alkyl. Exemplary R^B includes methyl, ethyl, isopropyl, n-propyl, n-butyl, sec-butyl, or t-butyl.

[00197] In one embodiment, compound of formula IIC includes one or more of the following features:

- a) ring G is benzyl optionally substituted with one R^1 substituent, preferably halo;
- b) Y' is S and R is hydrogen;
- c) R^C is C1-C4 alkyl;
- d) X_7 is NH or NR' wherein R' is C1-C4 alkyl; and
- e) R^B is C1-C4 alkyl.

[00198] According to another embodiment, the present invention provides a compound of formula IID:



IID;

wherein:

- Y' is O or S;
- R is hydrogen or R^2 ;
- X_8 is O, S, or NR' ;
- R' is hydrogen, R^2 , or R^6 ;

R^{BB} is C1-6 aliphatic or a 3-7 membered monocyclic or 8-14 membered bicyclic or tricyclic, saturated, unsaturated or aromatic ring containing 0-4 heteroatoms in each ring, wherein each said heteroatom is independently selected from N, S, or O;

wherein each of ring G and R^{BB} is independently and optionally substituted with up to 4 substituents selected from R^1 , R^2 , R^3 , R^4 , or R^5 , as defined above.

[00199] In certain embodiments, X_8 is O. Or, X_8 is S. Or, X_8 is NR' .

[00200] According to another embodiment, R^{BB} is phenyl optionally substituted with up to two R^1 substituents. Or, R^{BB} is phenyl.

[00201] According to another embodiment, R^{BB} is C1-C6 alkyl. Exemplary R^{BB} includes methyl, ethyl, isopropyl, n-propyl, n-butyl, sec-butyl, or t-butyl.


[00202] According to another embodiment, R^{BB} is optionally substituted C3-C8 cycloalkyl, e.g., cyclopropyl, cyclopentyl, or cyclohexyl.

[00203] According to another embodiment, R^{BB} is optionally substituted benzyl.

[00204] In one embodiment, compounds of formula IID have one or more of the following features:

- a) Y' is S and R is hydrogen;
- b) each ring G is unsubstituted phenyl;
- d) X_8 is NR' , and R' is hydrogen or C1-C4 alkyl; and
- e) R^{BB} is C1-C4 alkyl, benzyl, cyclopentyl, or cyclohexyl.

[00205] According to another preferred embodiment, the present invention provides a compound having formula (III):



or a pharmaceutically acceptable salt thereof;
wherein:

X_1 is a bond, O, S, CHOR, C(O), C(O)O, CF₂, CH₂, or NR;

R is H or R²

A₁ is (C2-C10) aliphatic, aryl, heteroaryl, heterocyclic, or cycloaliphatic;

each B₁ is independently selected from 3-7 membered monocyclic, saturated, unsaturated or aromatic ring containing 0-4 heteroatoms selected from N, NH, S, or O;

wherein each A₁ is optionally substituted with up to 4 substituents independently selected from R¹, R², R³, R⁴, or R⁵;

R¹ is R⁶ or ((C1-C4)aliphatic)_n-Y;

n is 0 or 1;

Y is halo, CN, NO₂, CF₃, OCF₃, OH, SR⁶, S(O)R⁶, SO₂R⁶, NH₂, NHR⁶, N(R⁶)₂, NR⁶R⁸, COOH, COOR⁶ or OR⁶; or

two R¹ on adjacent ring atoms, taken together, form 1,2-methylenedioxy or 1,2-ethylenedioxy;

R² is aliphatic, wherein each R² optionally comprises up to 2 substituents independently selected from R¹, R⁴, or R⁵;

R³ is a cycloaliphatic, aryl, heterocyclic, or heteroaryl ring optionally comprising up to 3

substituents, independently selected from R^1 , R^2 , R^4 or R^5 ;

R^4 is OR^5 , OR^6 , $OC(O)R^6$, $OC(O)R^5$, $OC(O)OR^6$, $OC(O)OR^5$, $OC(O)N(R^6)_2$, $OC(O)N(R^5)_2$, $OC(O)N(R^6R^5)$, SR^6 , SR^5 , $S(O)R^6$, $S(O)R^5$, SO_2R^6 , SO_2R^5 , $SO_2N(R^6)_2$, $SO_2N(R^5)_2$, $SO_2NR^5R^6$, SO_3R^6 , SO_3R^5 , $C(O)R^5$, $C(O)OR^5$, $C(O)R^6$, $C(O)OR^6$, $C(O)N(R^6)_2$, $C(O)N(R^5)_2$, $C(O)N(R^5R^6)$, $C(O)N(OR^6)R^6$, $C(O)N(OR^5)R^6$, $C(O)N(OR^6)R^5$, $C(O)N(OR^5)R^5$, $C(NOR^6)R^6$, $C(NOR^6)R^5$, $C(NOR^5)R^6$, $C(NOR^5)R^5$, $N(R^6)_2$, $N(R^5)_2$, $N(R^5R^6)$, $NR^5C(O)R^5$, $NR^6C(O)R^6$, $NR^6C(O)R^5$, $NR^6C(O)OR^6$, $NR^5C(O)OR^6$, $NR^6C(O)OR^5$, $NR^5C(O)OR^5$, $NR^6C(O)N(R^6)_2$, $NR^6C(O)NR^5R^6$, $NR^6C(O)N(R^5)_2$, $NR^5C(O)N(R^6)_2$, $NR^5C(O)NR^5R^6$, $NR^5C(O)N(R^5)_2$, $NR^6SO_2R^6$, $NR^6SO_2R^5$, $NR^5SO_2R^5$, $NR^5SO_2R^6$, $NR^6SO_2N(R^6)_2$, $NR^6SO_2NR^5R^6$, $NR^6SO_2N(R^5)_2$, $NR^5SO_2NR^5R^6$, $NR^5SO_2N(R^5)_2$, $N(OR^6)R^6$, $N(OR^6)R^5$, $N(OR^5)R^5$, or $N(OR^5)R^6$;

R^5 is a cycloaliphatic, aryl, heterocyclic, or heteroaryl ring, optionally comprising up to 3 R^1 substituents;

R^6 is H or aliphatic, wherein R^6 optionally comprises a R^7 substituent;

R^7 is a cycloaliphatic, aryl, heterocyclic, or heteroaryl ring, and each R^7 optionally comprises up to 2 substituents independently chosen from H, (C_1-C_6) -straight or branched alkyl, (C_2-C_6) straight or branched alkenyl or alkynyl, 1,2-methylenedioxy, 1,2-ethylenedioxy, or $(CH_2)_n-Z$;


Z is selected from halo, CN, NO_2 , CF_3 , OCF_3 , OH, S-aliphatic, S(O)-aliphatic, SO_2 -aliphatic, NH_2 , N-

aliphatic, $N(\text{aliphatic})_2$, $N(\text{aliphatic})R^8$, COOH , $\text{C}(\text{O})\text{O}(-\text{aliphatic})$, or O-aliphatic ; and

R^8 is an amino capping group;

provided that:

(i) when X_1 is a bond, one B_1 is phenyl and the other B_1 is N -piperidyl, then A is not:



(ii) when X_1 is a bond, then A_1 is not an optionally substituted 6-membered heteroaryl ring with 1-3 nitrogen ring atoms.

[00206] In certain embodiments, According to another embodiment of formula (III), X_1 is CH_2 , CF_2 , or O . In another embodiment, X_1 is a bond, O , S , CF_2 , CH_2 , or NR . Or, X_1 is CH_2 or O . Or, X_1 is CH_2 .

[00207] According to another embodiment of formula (III), A_1 is an optionally substituted C3-C7 cycloaliphatic ring. In certain embodiments,, A_1 is an optionally substituted cyclopropyl, cyclopentyl, or cyclohexyl.

[00208] According to another embodiment of formula (III), A_1 is optionally substituted (C1-C10)aliphatic. In certain embodiments, A_1 is optionally substituted methyl, ethyl, propyl, butyl, pentyl, or hexyl.

[00209] According to another embodiment of formula (III), A_1 is optionally substituted C6-C10 aryl ring. In certain embodiments, A_1 is optionally substituted phenyl or naphthyl.

[00210] According to another embodiment of formula (III), A₁ is optionally substituted C5-C12 heteroaryl ring. In certain embodiments, A is selected from optionally substituted triazinyl, pyrazinyl, pyrimidinyl, pyridazinyl, pyridinyl, thiadiazolyl, triazolyl, oxadiazolyl, isothiazolyl, pyrazolyl, imidazolyl, thiazolyl, oxazolyl, pyrrolyl, thienyl, furanyl, indoliziny, indolyl, isoindolyl, benzofuranyl, benzo[b]thienyl, 1H-indazolyl, benzimidazolyl, benzthiazolyl, purinyl, quinolinyl, isoquinolinyl, cinnolinyl, phthazinyl, quinazolinyl, quinoxalinyl, 1,8-naphthyridinyl, pteridinyl, acridinyl, phenazinyl, phenothiazinyl, or phenoxazinyl.

[00211] According to another embodiment of formula (III), A₁ is optionally substituted C3-C12 heterocyclic ring. In certain embodiments, A₁ is selected from optionally substituted aziridine, oxirane, thiirane, pyrrolidyl, tetrahydrofuranyl, tetrahydrothienyl, dioxolanyl, pyrrolinyl, pyranyl, pyrazolinyl, pyrazolidinyl, piperidinyl, 1,4-dioxanyl, morpholinyl, 1,4-dithianyl, thiomorpholinyl, piperazinyl, 3H-indolyl, or indolinyl.

[00212] According to another embodiment of formula (III), each B₁ is independently selected from optionally substituted C6-C10 aryl. In certain embodiments, each B₁ is independently an optionally substituted phenyl or naphthyl. Or, each B₁ is an unsubstituted phenyl.


[00213] According to another embodiment of formula (III), each B₁ is independently selected from optionally substituted C5-C12 heteroaryl. In certain embodiments, each B₁ is independently and optionally substituted C5-C7 heteroaryl. Or, each B₁ is independently selected from optionally substituted triazinyl, pyrazinyl, pyrimidinyl, pyridazinyl, pyridinyl, thiadiazolyl, triazolyl,

oxadiazolyl, isothiazolyl, pyrazolyl, imidazolyl, thiazolyl, oxazolyl, pyrrolyl, thienyl, furanyl, indoliziny, indolyl, isoindolyl, benzofuranyl, benzo[b]thienyl, 1H-indazolyl, benzimidazolyl, benzthiazolyl, purinyl, quinolinyl, isoquinolinyl, cinnolinyl, phthazinyl, quinazolinyl, quinoxalinyl, 1,8-naphthyridinyl, pteridinyl, carbazolyl, acridinyl, phenazinyl, phenothiazinyl, phenoxazinyl, indenyl, naphthyl, azuliny, or anthracenyl.

[00214] According to another embodiment of formula (III), each B₁ is independently an optionally substituted 3-12 membered heterocyclic ring having up to 4 heteroatoms selected from O, S, or NR. In certain embodiments, each B₁ is independently selected from optionally substituted aziridine, oxirane, thiirane, pyrrolidyl, tetrahydrofuranyl, tetrahydrothienyl, dioxolanyl, pyrrolinyl, pyranyl, pyrazolinyl, pyrazolidinyl, piperidinyl, 1,4-dioxanyl, morpholinyl, 1,4-dithianyl, thiomorpholinyl, piperazinyl, 3H-indolyl, or indolinyl.

[00215] Exemplary embodiments of R¹, R², R³, R⁴, R⁵, R⁶, R⁷, R⁸, and Z in compounds of formula (III) are as described above for compound of formula (I).


[00216] According to another embodiment, the present invention provides compounds of formula IV:



IV;

wherein:

B is selected from:



wherein:

Y is O or S; and

X₃ is O or S.


[00217] According to one embodiment, X₃ is O. Or, X₃ is S.

[00218] According to one embodiment, B is structure (i) above. Or, B is structure (ii) above.

[00219] According to another embodiment, R is hydrogen.

[00220] According to another embodiment, Y' is S. Or, Y' is O.


[00221] According to another embodiment, the present invention provides compounds of formula V:



wherein:

Y' is O or S;

B is selected from:



wherein:

Ak is C1-C6 alkylidene;

X₄ is CH₂, O or S;

Ar' is phenyl optionally substituted with up to two R¹; and

B is optionally substituted with up to two R¹.

[00222] In some embodiments, Ak is selected from CH_2 , $\text{CH}(\text{CH}_3)$, $\text{C}(\text{CH}_3)_2$, $\text{CH}(\text{Et})$, $\text{C}(\text{Et})_2$, $\text{CH}(\text{n-propyl})$, $\text{CH}(\text{i-Pr})$, $\text{CH}(\text{n-butyl})$, $\text{CH}(\text{but-2-yl})$, or $\text{CH}(\text{t-butyl})$.

[00223] In some embodiments, Ar' is phenyl optionally substituted with halo, C1-C4 alkyl, or O-(C1-C4 alkyl).

[00224] In some embodiments, X_4 is CH_2 . X_4 is S. Or, X_4 is O.


[00225] In some embodiments, R is hydrogen.

[00226] In certain embodiments, p is 2, X is a bond, and each A is optionally substituted phenyl.

[00227] In other embodiments, the compounds have one or more of the following features:

- a) Y' is S;
- b) R is hydrogen;
- c) p is 2, X is a bond, and each A is phenyl;
- d) B is ring (iii) above, wherein Ak is $\text{CH}(\text{CH}_3)$ and Ar' is phenyl optionally substituted with halo, C1-C4 alkyl, or O-(C1-C4 alkyl).

[00228] According to another embodiment, the present invention provides a compound of formula VI:



VI;

Y' is O or S;

R is hydrogen or R^2 ;

B is phenyl, 3-7 membered, monocyclic, saturated, carbocyclic ring, or 3-10 membered saturated or unsaturated, monocyclic or bicyclic heterocyclic ring having up to 4 heteroatoms selected from O, S, or N, or

5-10 membered monocyclic or bicyclic heteroaryl ring having up to 4 heteroatoms selected from O, S, or N;

wherein each ring G_1 , G_2 , and B is independently substituted with up to 4 substituents selected from R^1 , R^2 , R^3 , R^4 , or R^5 ;

R^1 is oxo, R^6 or $((C1-C4)aliphatic)_n-Y$;

n is 0 or 1;

Y is halo, CN, NO_2 , CF_3 , OCF_3 , OH, SR^6 , $S(O)R^6$, SO_2R^6 , NH_2 , NHR^6 , $N(R^6)_2$, NR^6R^8 , $N(R^8)^2$, COOH, COOR⁶ or OR⁶; or

two R^1 on adjacent ring atoms, taken together, form 1,2-methylenedioxy or 1,2-ethylenedioxy;

R^2 is aliphatic, wherein each R^2 optionally comprises up to 2 substituents independently selected from R^1 , R^4 , or R^5 ;

R^3 is a cycloaliphatic, aryl, heterocyclic, or heteroaryl ring optionally comprising up to 3 substituents, independently selected from R^1 , R^2 , R^4 or R^5 ;

R^4 is OR^5 , OR^6 , $OC(O)R^6$, $OC(O)R^5$, $OC(O)OR^6$, $OC(O)OR^5$, $OC(O)N(R^6)_2$, $OC(O)N(R^5)_2$, $OC(O)N(R^6R^5)$, SR^6 , SR^5 , $S(O)R^6$, $S(O)R^5$, SO_2R^6 , SO_2R^5 , $SO_2N(R^6)_2$, $SO_2N(R^5)_2$, $SO_2NR^5R^6$, SO_3R^6 , SO_3R^5 , $C(O)R^5$, $C(O)OR^5$, $C(O)R^6$, $C(O)OR^6$, $C(O)N(R^6)_2$, $C(O)N(R^5)_2$, $C(O)N(R^5R^6)$, $C(O)N(OR^6)R^6$, $C(O)N(OR^5)R^6$, $C(O)N(OR^6)R^5$, $C(O)N(OR^5)R^5$, $C(NOR^6)R^6$, $C(NOR^6)R^5$, $C(NOR^5)R^6$, $C(NOR^5)R^5$, $N(R^6)_2$, $N(R^5)_2$, $N(R^5R^6)$, $NR^5C(O)R^5$, $NR^6C(O)R^6$, $NR^6C(O)R^5$, $NR^5C(O)R^6$, $NR^6C(O)OR^6$, $NR^5C(O)OR^6$, $NR^6C(O)OR^5$, $NR^5C(O)OR^5$, $NR^6C(O)N(R^6)_2$, $NR^6C(O)NR^5R^6$, $NR^6C(O)N(R^5)_2$, $NR^5C(O)N(R^6)_2$, $NR^5C(O)NR^5R^6$,

NR⁵C(O)N(R⁵)₂, NR⁶SO₂R⁶, NR⁶SO₂R⁵, NR⁵SO₂R⁵, NR⁵SO₂R⁶, NR⁶SO₂N(R⁶)₂, NR⁶SO₂NR⁵R⁶, NR⁶SO₂N(R⁵)₂, NR⁵SO₂N(R⁶)₂, NR⁵SO₂NR⁵R⁶, NR⁵SO₂N(R⁵)₂, N(OR⁶)R⁶, N(OR⁶)R⁵, N(OR⁵)R⁵, or N(OR⁵)R⁶;

R⁵ is a cycloaliphatic, aryl, heterocyclic, or heteroaryl ring, optionally comprising up to 3 R¹ substituents;

R⁶ is H or aliphatic, wherein R⁶ optionally comprises a R⁷ substituent;

R⁷ is a cycloaliphatic, aryl, heterocyclic, or heteroaryl ring, and each R⁷ optionally comprises up to 2 substituents independently chosen from H, (C₁-C₆)-straight or branched alkyl, (C₂-C₆) straight or branched alkenyl or alkynyl, 1,2-methylenedioxy, 1,2-ethylenedioxy, or (CH₂)_n-Z;

Z is selected from halo, CN, NO₂, CF₃, OCF₃, OH, S-aliphatic, S(O)-aliphatic, SO₂-aliphatic, NH₂, NH-aliphatic, N(aliphatic)₂, N(aliphatic)R⁸, NHR⁸, N(R⁸)₂, COOH, C(O)O(-aliphatic, or O-aliphatic; and

R⁸ is an amino capping group.

[00229] In certain embodiments of formula VI, when R is hydrogen, then the following compounds are excluded:

- B is not quinolin-2-yl or 1,2-dihydro-2-oxo-quinolin-4-yl;
- when G₁ and G₂ both are phenyl, and Y' is S, then B is not 1,4-benzodioxin-2-yl, cyclopropyl, cyclohexyl, thien-2-yl, 1H-thieno[2,3-c]pyrazol-1-phenyl-3-methyl-5-yl, 5-methyl-thien-3-yl, 2,5-dichloro-thien-3-yl, 2-phenyl-quinolin-4-yl, furan-2-yl, thien-5-(4,5-diphenyl-2-thiazolyl-carboxamide)-2-yl, benzo[b]thiophen-2-yl,


pyridin-2-(4,5-diphenyl-2-thiazolyl-carboxamide)-6-yl, 5-nitro-thien-2-yl, 3-chloro-benzo[b]thiophen-2-yl, 4H-1-benzopyran-3-yl or 2H-1-benzopyran-3,4-dihydro-3-oxo-4-yl, 4H-1-benzopyran-3-yl or 2H-1-benzopyran-3,4-dihydro-3-oxo-4-yl;

c) when G_1 and G_2 both are phenyl, and Y' is O, then B is not 1,2-dihydro-2-oxo-quinolin-4-yl or 3,4-dihydro-3-phenyl-phthalazin-1-yl or thien-2-yl;

d) the following compounds are excluded:

Y'	G_1	G_2	B
S	Ph	4-Me-Ph	2-Cl-thien-5-yl or 2,5-dichloro-thien-3-yl
S	Ph or 3,4-dimethylphenyl	3,4-dimethylphenyl or Ph	pyridin-3-yl
S	3,4-dimethylphenyl	Ph	pyridin-4-yl
S	4-Cl-Ph	Ph	thien-2-yl
S	3,4-dimethylphenyl	Ph	pyridin-4-yl
S	Ph	4-Me-phenyl	thien-2-yl or benzothiazol-2-yl
S	4-NO ₂ -Ph or 4-Me-Ph	Ph or 4-OMe-Ph or 2,4-dimethylphenyl	thien-2-yl or furan-2-yl
S	4-OMe-Ph or 2,4-dimethylphenyl	4-NO ₂ -Ph	furan-2-yl or thien-2-yl
S	4-OMe-Ph or Ph	4-OMe-Ph	furan-2-yl or thien-2-yl
S	Ph	4-Me-Ph	furan-2-yl
S	Ph	4-Me-Ph	5-nitro-thien-2-yl
S	4-Me-Ph	4-Me-Ph	2-chloro-pyridin-3-yl
S	Ph	4-Me-Ph	3-chloro-benzo[b]thiophen-2-yl
S	4-Me-Ph or Ph	4-NO ₂ -Ph	thien-2-yl or furan-2-yl
S	2,4-dimethylphenyl	4-NO ₂ -Ph	pyridin-3-yl or thien-2-yl
S	4-Cl-Ph	4-NO ₂ -Ph	thien-2-yl
S	Ph or 2-OMe-Ph or 3-OMe-Ph or 4-OMe-Ph	2,4-dimethoxyphenyl	1H-indol-2-yl.

d) when Y' is S, G₁ and G₂ are both phenyl, then B is not



[00230] In one embodiment, G₁ and G₂ are both phenyl. Or, each is independently and optionally substituted with up to two substituents selected from halo, or C1-C4 alkyl.

[00231] In certain embodiments of formula VI, B is phenyl optionally substituted with up to two substituents selected from halo, C1-C4 alkyl, O-(C1-C4 alkyl), COOH, COO(C1-C4 alkyl), or cyano.


[00232] In other embodiments, B is 3,4-dichlorophenyl, 4-chlorophenyl, 4-methoxyphenyl, 4-methylphenyl, 2,6-difluorophenyl, 2-methylphenyl, 3-methylphenyl, phenyl-2-carboxylic acid, 2-chlorophenyl, 4-cyanophenyl, or 3-methoxyphenyl.

[00233] In another embodiment, B is 3-7 membered, monocyclic, saturated, carbocyclic ring. Exemplary rings include cyclopropyl, cyclopentyl, cyclohexyl, or cycloheptyl.

[00234] In another embodiment, B is a 3-10 membered saturated or unsaturated, monocyclic or bicyclic heterocyclic ring having up to 4 heteroatoms selected from O, S, or N. Exemplary rings include tetrahydrofuranyl, thienyl, or pyrrolyl.

[00235] In another embodiment, B is a 5-10 membered monocyclic or bicyclic heteroaryl ring having up to 4 heteroatoms selected from O, S, or N.

[00236] According to another embodiment, the present invention provides a compound of formula VII:



VII;

wherein:

Y' is O or S;

R is hydrogen or R²;

R^{Ak} is C1-C6 aliphatic, optionally substituted with up to 3 substituents independently selected from R¹, R², or R³;

B is phenyl, 3-7 membered, monocyclic, saturated, carbocyclic ring, or 3-10 membered saturated or unsaturated, monocyclic or bicyclic heterocyclic ring having up to 4 heteroatoms selected from O, S, or N, or 5-10 membered monocyclic or bicyclic heteroaryl ring having up to 4 heteroatoms selected from O, S, or N;

wherein each ring G₁, G₂, and B is independently substituted with up to 4 substituents selected from R¹, R², R³, R⁴, or R⁵;

R¹ is oxo, R⁶ or ((C1-C4)aliphatic)_n-Y;

n is 0 or 1;

Y is halo, CN, NO₂, CF₃, OCF₃, OH, SR⁶, S(O)R⁶, SO₂R⁶, NH₂, NHR⁶, N(R⁶)₂, NR⁶R⁸, N(R⁸)₂, COOH, COOR⁶ or OR⁶; or

two R¹ on adjacent ring atoms, taken together, form 1,2-methylenedioxy or 1,2-ethylenedioxy;

R² is aliphatic, wherein each R² optionally comprises up to 2 substituents independently selected from R¹, R⁴, or R⁵;

R³ is a cycloaliphatic, aryl, heterocyclic, or heteroaryl ring optionally comprising up to 3

substituents, independently selected from R^1 , R^2 , R^4 or R^5 ;

R^4 is OR^5 , OR^6 , $OC(O)R^6$, $OC(O)R^5$, $OC(O)OR^6$, $OC(O)OR^5$, $OC(O)N(R^6)_2$, $OC(O)N(R^5)_2$, $OC(O)N(R^6R^5)$, SR^6 , SR^5 , $S(O)R^6$, $S(O)R^5$, SO_2R^6 , SO_2R^5 , $SO_2N(R^6)_2$, $SO_2N(R^5)_2$, $SO_2NR^5R^6$, SO_3R^6 , SO_3R^5 , $C(O)R^5$, $C(O)OR^5$, $C(O)R^6$, $C(O)OR^6$, $C(O)N(R^6)_2$, $C(O)N(R^5)_2$, $C(O)N(R^5R^6)$, $C(O)N(OR^6)R^6$, $C(O)N(OR^5)R^6$, $C(O)N(OR^6)R^5$, $C(O)N(OR^5)R^5$, $C(NOR^6)R^6$, $C(NOR^6)R^5$, $C(NOR^5)R^6$, $C(NOR^5)R^5$, $N(R^6)_2$, $N(R^5)_2$, $N(R^5R^6)$, $NR^5C(O)R^5$, $NR^6C(O)R^6$, $NR^6C(O)R^5$, $NR^5C(O)R^6$, $NR^6C(O)OR^6$, $NR^5C(O)OR^6$, $NR^6C(O)OR^5$, $NR^5C(O)OR^5$, $NR^6C(O)N(R^6)_2$, $NR^6C(O)NR^5R^6$, $NR^6C(O)N(R^5)_2$, $NR^5C(O)N(R^6)_2$, $NR^5C(O)NR^5R^6$, $NR^5C(O)N(R^5)_2$, $NR^6SO_2R^6$, $NR^6SO_2R^5$, $NR^5SO_2R^5$, $NR^5SO_2R^6$, $NR^6SO_2N(R^6)_2$, $NR^6SO_2NR^5R^6$, $NR^6SO_2N(R^5)_2$, $NR^5SO_2N(R^6)_2$, $NR^5SO_2NR^5R^6$, $NR^5SO_2N(R^5)_2$, $N(OR^6)R^6$, $N(OR^6)R^5$, $N(OR^5)R^5$, or $N(OR^5)R^6$;

R^5 is a cycloaliphatic, aryl, heterocyclic, or heteroaryl ring, optionally comprising up to 3 R^1 substituents;

R^6 is H or aliphatic, wherein R^6 optionally comprises a R^7 substituent;

R^7 is a cycloaliphatic, aryl, heterocyclic, or heteroaryl ring, and each R^7 optionally comprises up to 2 substituents independently chosen from H, (C_1-C_6) -straight or branched alkyl, (C_2-C_6) straight or branched alkenyl or alkynyl, 1,2-methylenedioxy, 1,2-ethylenedioxy, or $(CH_2)_n-Z$;

Z is selected from halo, CN, NO_2 , CF_3 , OCF_3 , OH, S-aliphatic, $S(O)$ -aliphatic, SO_2 -aliphatic, NH_2 , NH-

aliphatic, $N(\text{aliphatic})_2$, $N(\text{aliphatic})R^8$, NHR^8 , $N(R^8)_2$, COOH , $\text{C(O)O}(-\text{aliphatic}, \text{ or } \text{O-aliphatic})$; and

R^8 is an amino-capping group.

[00237] In one embodiment of formula VII, when each of G_1 , G_2 , and B is unsubstituted phenyl, and R is hydrogen, then B is not 3,4,5-trimethoxyphenyl.

[00238] In one embodiment, R^{Ak} is C1-C6 alkyl. Exemplary R^{Ak} include methyl, ethyl, isopropyl, n-propyl, sec-butyl, n-butyl, or t-butyl.


[00239] In one embodiment, B is optionally substituted phenyl.

[00240] In another embodiment, B is 3-7 membered, monocyclic, saturated, carbocyclic ring. Exemplary rings include cyclopropyl, cyclopentyl, cyclohexyl, or cycloheptyl.

[00241] In another embodiment, B is a 3-10 membered saturated or unsaturated, monocyclic or bicyclic heterocyclic ring having up to 4 heteroatoms selected from O, S, or N. Exemplary rings include tetrahydrofuranyl, tetrahydrothiophenyl, pyrrolidinyl, or piperazinyl.

[00242] In another embodiment, B is a 5-10 membered monocyclic or bicyclic heteroaryl ring having up to 4 heteroatoms selected from O, S, or N.

[00243] According to another embodiment, the present invention provides a compound having formula I':





or a pharmaceutically acceptable salt thereof;
wherein:

Y' is O, S, or NR; R is H, R², or R⁶;

C is a phenyl or 5-8 membered cycloaliphatic ring;

Q is selected from:



each B is independently selected from 3-7 membered monocyclic or 8-14 membered bicyclic or tricyclic, saturated, unsaturated or aromatic ring containing 0-4 heteroatoms in each ring, wherein each said heteroatom is independently selected from N, NH, S, or O;

wherein each A, B, and C is independently and optionally substituted with up to 4 substituents independently selected from R¹, R², R³, R⁴, or R⁵;

R^L is -OR^A, -SR^A, or -N(R^{AB})₂;

each R^A is independently hydrogen, C1-C6 aliphatic, or a 3-7 membered carbocyclic or heterocyclic ring, saturated or unsaturated ring, having up to 3 heteroatoms selected from O, N, or S, wherein each R^A is optionally substituted with up to 3 substituents independently selected from R¹, R⁴ or R⁷,

each R^{AB} is independently hydrogen or C1-C6 aliphatic optionally substituted with up to 3 substituents independently selected from R¹, R⁴ or R⁷;

wherein up to two methylene units in R^A or R^{AB} are optionally replaced with -CO-, -CS-, -COCO-, -CONR-, -CO₂-

, -OCO-, -NRCO₂-, -O-, -NRCONR-, -OCONR-, -NRCO-, -S-, -SO, -SO₂-, -NR-, -SO₂NR-, NRSO₂-, or -NRSO₂NR; or

two R^{AB}, taken together with the nitrogen atom, is a 3-7 membered heterocyclic or heteroaryl ring containing up to 4 heteroatoms selected from O, N, or S, wherein said ring is optionally substituted with up to 2 substituents selected from oxo or (C₁₋₄aliphatic)_p-Y;

R^M is C1-C6 aliphatic, optionally substituted with up to two substituents selected from R¹, R², R³, or R⁴;

each of X₁ and X₂ is independently selected from O, S, or NR;

R^N is C1-C6 aliphatic or phenyl, wherein R^N is optionally substituted with up to two substituents selected from R¹, R², R³, or R⁴;

R^P is C1-C6 aliphatic, optionally substituted with up to two substituents selected from R¹, R², R³, or R⁴;

R^Q is C1-C6 aliphatic or aryl, wherein R^Q is optionally substituted with up to two substituents selected from R¹, R², R³, or R⁴;

R¹ is oxo, R⁶ or ((C1-C4)aliphatic)_n-Y;

n is 0 or 1;

Y is halo, CN, NO₂, CF₃, OCF₃, OH, SR⁶, S(O)R⁶, SO₂R⁶, NH₂, NHR⁶, N(R⁶)₂, NR⁶R⁸, N(R⁸)₂, COOH, COOR⁶ or OR⁶; or

two R¹ on adjacent ring atoms, taken together, form 1,2-methylenedioxy or 1,2-ethylenedioxy;

R² is aliphatic, wherein each R² optionally comprises up to 2 substituents independently selected from R¹, R⁴, or R⁵;

R³ is a cycloaliphatic, aryl, heterocyclic, or heteroaryl ring optionally comprising up to 3

substituents, independently selected from R^1 , R^2 , R^4 or R^5 ;

R^4 is OR^5 , OR^6 , $OC(O)R^6$, $OC(O)R^5$, $OC(O)OR^6$, $OC(O)OR^5$, $OC(O)N(R^6)_2$, $OC(O)N(R^5)_2$, $OC(O)N(R^6R^5)$, SR^6 , SR^5 , $S(O)R^6$, $S(O)R^5$, SO_2R^6 , SO_2R^5 , $SO_2N(R^6)_2$, $SO_2N(R^5)_2$, $SO_2NR^5R^6$, SO_3R^6 , SO_3R^5 , $C(O)R^5$, $C(O)OR^5$, $C(O)R^6$, $C(O)OR^6$, $C(O)N(R^6)_2$, $C(O)N(R^5)_2$, $C(O)N(R^5R^6)$, $C(O)N(OR^6)R^6$, $C(O)N(OR^5)R^6$, $C(O)N(OR^6)R^5$, $C(O)N(OR^5)R^5$, $C(NOR^6)R^6$, $C(NOR^6)R^5$, $C(NOR^5)R^6$, $C(NOR^5)R^5$, $N(R^6)_2$, $N(R^5)_2$, $N(R^5R^6)$, $NR^5C(O)R^5$, $NR^6C(O)R^6$, $NR^6C(O)R^5$, $NR^5C(O)R^6$, $NR^6C(O)OR^6$, $NR^5C(O)OR^6$, $NR^6C(O)OR^5$, $NR^5C(O)OR^5$, $NR^6C(O)N(R^6)_2$, $NR^6C(O)NR^5R^6$, $NR^6C(O)N(R^5)_2$, $NR^5C(O)N(R^6)_2$, $NR^5C(O)NR^5R^6$, $NR^5C(O)N(R^5)_2$, $NR^6SO_2R^6$, $NR^6SO_2R^5$, $NR^5SO_2R^5$, $NR^5SO_2R^6$, $NR^6SO_2N(R^6)_2$, $NR^6SO_2NR^5R^6$, $NR^6SO_2N(R^5)_2$, $NR^5SO_2N(R^6)_2$, $NR^5SO_2NR^5R^6$, $NR^5SO_2N(R^5)_2$, $N(OR^6)R^6$, $N(OR^6)R^5$, $N(OR^5)R^5$, or $N(OR^5)R^6$;

R^5 is a cycloaliphatic, aryl, heterocyclic, or heteroaryl ring, optionally comprising up to 3 R^1 substituents;

R^6 is H or aliphatic, wherein R^6 optionally comprises a R^7 substituent;

R^7 is a cycloaliphatic, aryl, heterocyclic, or heteroaryl ring, and each R^7 optionally comprises up to 2 substituents independently chosen from H, (C_1-C_6) -straight or branched alkyl, (C_2-C_6) straight or branched alkenyl or alkynyl, 1,2-methylenedioxy, 1,2-ethylenedioxy, or $(CH_2)_n-Z$;

Z is selected from halo, CN, NO_2 , CF_3 , OCF_3 , OH, S-aliphatic, S(O)-aliphatic, SO_2 -aliphatic, NH_2 , NH-

aliphatic, $N(\text{aliphatic})_2$, $N(\text{aliphatic})R^8$, NHR^8 , $N(R^8)_2$, COOH , $\text{C(O)O}(-\text{aliphatic}, \text{ or } \text{O-aliphatic}; \text{ and}$

R^8 is an amino-capping group.

[00244] In one embodiment, Y' is O. Or, Y' is S.

[00245] In another embodiment, R is hydrogen.

[00246] In one embodiment, Q in formula I' is B (structure (a)). Preferred B include optionally substituted phenyl, or C3-C8 cycloaliphatic. In one embodiment, B in formula I' is phenyl or C3-C8 cycloalkyl optionally substituted with up to two substituents selected from R^1 , R^2 , or phenyl optionally substituted with up to two substituents selected from R^1 or R^2 .

[00247] In another embodiment, Q in formula I' is - (C1-C6 aliphatic)-B (structure (b)).

[00248] In certain embodiments, said C1-C6 aliphatic is C1-C4 straight or branched alkylidene. Exemplary alkylidenes include $-\text{CH}_2-$, $-\text{CH}(\text{Me})-$, $-\text{C}(\text{Me})_2-$, $-\text{CH}(\text{Et})-$, $-\text{C}(\text{Et})_2-$, or $-\text{CH}_2-\text{CH}(\text{Me})-$.

[00249] In certain embodiments, B is selected from optionally substituted C3-C8 cycloalkyl, phenyl, piperidyl, or pyrrolidinyl. Preferably, B is phenyl, cyclopentyl, cyclohexyl, or piperidyl, optionally substituted with up to two R^1 or R^2 substituents.

[00250] In one embodiment, ring C is phenyl optionally substituted with up to two R^1 . Or ring C is cyclohexenyl optionally substituted with up to two R^1 .

[00251] According to a preferred embodiment, R^1 is 1,2-methylene dioxy, or 1,2-ethylenedioxy.

[00252] According to another preferred embodiment, R^1 is R^6 , wherein R^6 is straight chain or branched (C1-C6)alkyl or (C2-C6) alkenyl or alkynyl, optionally substituted with R^7 .

[00253] According to another preferred embodiment, R^1 is (C1-C4 aliphatic)_n-Y, wherein n is 0 or 1, and Y is halo, CN, NO₂, CF₃, OCF₃, OH, SR⁶, S(O)R⁶, SO₂R⁶, NH₂, NHR⁶, N(R⁶)₂, NR⁶R⁸, COOH, COOR⁶ or OR⁶;

[00254] According to another preferred embodiment, R^1 is selected from halo, CF₃, NH₂, NH(C1-C4 alkyl), NHC(O)CH₃, OH, O(C1-C4 alkyl), OPh, O-benzyl, S-(C1-C4 alkyl), C1-C4 aliphatic, CN, methylenedioxy, ethylenedioxy, SO₂NH(C1-C4 alkyl), or SO₂N(C1-C4 alkyl)₂.

[00255] According to another more preferred embodiment, R^1 is selected from methyl, n-propyl, i-propyl, t-butyl, halo, CF₃, NH₂, NH(CH₃), NHC(O)CH₃, OH, OCH₃, OPh, O-benzyl, S-(C₂H₅), S-CH₃, NO₂, CN, methylenedioxy, SO₂NH(n-propyl), or SO₂N(n-propyl)₂.



[00256] According to a preferred embodiment, R^2 is a straight chain or branched (C1-C6)alkyl or (C2-C6) alkenyl or alkynyl, optionally substituted with R^1 , R^4 , or R^5 . More preferably, R^2 is a straight chain or branched (C1-C4)alkyl or (C2-C4) alkenyl or alkynyl, optionally substituted with R^1 , R^4 , or R^5 .



[00257] In formula I', other embodiments of B, R^L , R^M , R^N , R^P , R^Q , X₁, X₂, and R are as described above for formula I.



[00258] Exemplary compounds of the present invention are shown below in Table 1.



[0001]



Table 1



Cmpd #	Compound
1	 <chem>CCOC1=CC=C2C(=C1)S=C(NC(=O)C3CCCCC3)N2</chem>
2	 <chem>CC1=CC(=C(C)C=C1C2=CN=C(S2)NC(=O)C3=CC=CC=C3</chem>



Cmpd #	Compound
3	 <chem>CC1(C)SC(C1)C(=O)CC2=CC=C(C)C=C2C3=NC=SC3</chem>
4	 <chem>COc1ccccc1C2=NC=SC2C(=O)CC3=CC=C(C)C=C3</chem>



Cmpd #	Compound
5	 <chem>COc1ccccc1c2nc3c(s2)nc(=O)c4ccc(C)cc43</chem>
6	 <chem>COc1ccccc1c2nc3c(s2)nc(=O)c4ccc(OC)cc43</chem>



Cmpd #	Compound
7	 <chem>COc1ccc(cc1)C2=NC(=SNC2=O)c3cc(OC)cc(c3)</chem>
8	 <chem>O=C(Nc1nc2c(c1)CCCC2)s3cc(Br)ccc3</chem>



Cmpd #	Compound
9	 <chem>NC(=O)c1cc(Br)ccc1Nc2cc(sc2C3=CC=CS3)</chem>
10	 <chem>CC1=CC=C(C=C1)C2=CC=C(C=C2)C(=O)Nc3nc4ccccc4s3</chem>



Cmpd #	Compound
11	 <chem>CCCC1=CC=C(C=C1)C(=O)Nc2nc3ccccc3s2</chem>
12	 <chem>CCCC1=CC=C(C=C1)C(=O)Nc2nc3ccccc3s2</chem>



Cmpd #	Compound
13	 <chem>COc1ccc(cc1)-c2ccc(cc2)C(=O)Nc3ccsc3</chem>
14	 <chem>COc1ccc(cc1)-c2ccc(cc2)C(=O)Nc3cc4ccccc4s3</chem>



Cmpd #	Compound
15	 <chem>COc1ccc(cc1)NC(=O)c2ccc(cc2)-c3ccc(OC)cc3</chem>
16	 <chem>CCCc1ccc(cc1)C(=O)Nc2ccsc2</chem>



Cmpd #	Compound
17	 <chem>O=C1C2CCC3C1CC2C3NC4=NC(=C(C=C4)C5=CC=CC=C5)S=C4</chem>
18	 <chem>O=C1NC(=C2C(=N1)SC(C2)C3=CC=CC=C3)C4=CC=CC=C4</chem>



Cmpd #	Compound
19	 <chem>Clc1ccc(cc1)Cc2sc(NC(=O)c3cc(Cl)ccc3)n2</chem>
20	 <chem>O=C(O)CSNC(=O)Nc1nc(C2=CC=CC=C2)c(C3=CC=CC=C3)s1</chem>



Cmpd #	Compound
21	 <chem>O=C(Nc1cc(C2=CC=CC=C2)sc1)C3=CC=CC=C3</chem>
22	 <chem>O=C(NC1CC1)c2nc(Cc3cc(Cl)cc(Cl)c3)sc2</chem>



Cmpd #	Compound
23	 <chem>Nc1nc(s1)Cc2ccc(Cl)cc2</chem>
24	 <chem>CC1=CC=C2C(=C1)S=C(NC(C)Cc3ccccc3)N2</chem>



Cmpd #	Compound
25	 <chem>CC1=CC=CC=C1C(=O)NCSCC2=CC=CC=C2CSCC3=CC=CC=C3</chem>
26	 <chem>Clc1ccc(cc1)CC2=CN=C(S2)CC(=O)NCC3=CC4=CC=CC=C4C5=CC=CC=C35</chem>



Cmpd #	Compound
27	 <chem>COc1ccc(cc1)Cc2sc(NC(=O)C(c3ccccc3)c4ccccc4)n2</chem>
28	 <chem>Clc1ccc(cc1)Cc2sc(NC(=O)c3ccccc3)n2</chem>



Cmpd #	Compound
29	 <chem>Clc1ccccc1CC1=CN(C(=O)C2CCCCC2)S1</chem>
30	 <chem>Clc1cc(Cl)ccc1CC1=CN(C(=O)C2CCCCC2)S1</chem>



Cmpd #	Compound
31	 <chem>O=C(Nc1ccsc1Cc2ccc(Cl)cc2)Cc3ccccc3</chem>
32	 <chem>O=C(Nc1ccsc1Cc2ccccc2)Nc3ccccc3</chem>



Cmpd #	Compound
33	 <chem>Clc1ccc(cc1)CC2=CN=C(S2)NC(=O)c3cc(Cl)ccc3</chem>
34	 <chem>C[C@H](C)[C@@H](C(=O)Nc1nc(C2=CC=CC=N2)s1)c3cc(F)c(F)cc3</chem>

Cmpd #	Compound
35	 <chem>c1ccc(cc1)-c2cc(sc2NC(=O)C(c3ccccc3)C(C)C)</chem>
36	 <chem>Clc1ccc(cc1)Cc2cc(sc2NC(=O)CC#N)</chem>



Cmpd #	Compound
37	 <chem>Clc1cccc(c1)Cc2nc(s2)NC(=O)C(c3ccccc3)c4ccccc4</chem>
38	 <chem>Clc1cccc(c1)Cc2nc(s2)NC(=O)C(c3ccccc3)c4ccccc4</chem>



Cmpd #	Compound
39	 <chem>Cc1ccc(cc1)C(=O)Nc2nc(c(s2)c3ccc(C)cc3)c4ccccc4</chem>
40	 <chem>Cc1ccc2c(c1)sc(c2)n3c(c4ccccc43)C(=O)N[C@@H](C)c5ccccc5</chem>



Cmpd #	Compound
41	 Chemical structure of compound 41: 2-(4-chlorobenzoyl)-4-methyl-1,3-benzoxazole. The structure consists of a benzoxazole ring with a methyl group (CH₃) at the 4-position and a 2-(4-chlorobenzoyl) group at the 2-position. The benzoyl group is attached to the nitrogen atom of the benzoxazole ring. The benzene ring of the benzoyl group has a chlorine atom (Cl) at the para position.
42	 Chemical structure of compound 42: 2-(4-bromobenzoyl)-4-methyl-1,3-benzoxazole. The structure is similar to compound 41, but the benzoyl group is attached to the nitrogen atom of the benzoxazole ring, and the benzene ring of the benzoyl group has a bromine atom (Br) at the para position.



Cmpd #	Compound
43	 <chem>Clc1ccc(cc1)C(=O)Nc2sc(cc2-c3ccc(Br)cc3)C(=O)N</chem>
44	 <chem>Cc1ccc(cc1)C(=O)Nc2sc(C)c(c2-c3ccccc3)C(=O)N</chem>



Cmpd #	Compound
45	 <chem>N#Cc1ccc(cc1)C(=O)Nc2nc(s2)c3ccccc3c4ccc(C)cc4</chem>
46	 <chem>c1ccc(cc1)CC(=O)Nc2nc(s2)c3cccc4ccccc34</chem>



Cmpd #	Compound
47	 <chem>Cc1ccc(cc1)C(=O)Nc2nc(cc2-c3ccc(C)cc3)S</chem>
48	 <chem>c1ccccc1C(=O)Nc2nc(cc2-c3ccc(C)cc3)S</chem>



Cmpd #	Compound
49	 <chem>COc1ccc(cc1)C(=O)Nc2cc(sc2-c3ccccc3)</chem>
50	 <chem>CCC(=O)Nc1cc(sc1-c2ccccc2-c3ccccc3)</chem>



Cmpd #	Compound
51	 <chem>COc1ccccc1C(=O)Nc2nc(c3ccccc3c2c4ccccc4)C5=CC=CC=C5</chem>
52	 <chem>Cc1ccc(cc1)C(=O)Nc2nc(c3ccccc3c2c4ccccc4)C5=CC=CC=C5</chem>



Cmpd #	Compound
53	 <chem>Nc1nc(s1)C2=CC=CC=C2C(=O)c3ccc(Cl)cc3</chem>
54	 <chem>Cc1ccc(cc1)Cc2nc(s2)C(=O)c3ccc(Cl)cc3</chem>



Cmpd #	Compound
55	 <chem>NC(=O)C1CCCCC1c2nc(C3=CC=CC=C3)cs2</chem>
56	 <chem>Clc1cc(Cl)ccc1Cc2nc(C(=O)N(C(=O)C(c3ccccc3)(c4ccccc4)C5=CC=CC=C5)c6ccccc6)cs2</chem>



Cmpd #	Compound
57	
58	



Cmpd #	Compound
59	 <chem>O=C(O)c1cccc2c1c(=O)nc(Nc3nc(c(c3)c4ccccc4)c5ccccc5)c2</chem>
60	 <chem>CC(O)CC(=O)Nc1nc(c(c1)c2ccccc2)c3ccccc3</chem>



Cmpd #	Compound
61	 <chem>O=C(O)CCCC(=O)Nc1nc(c(s1)c2ccccc2)c3ccccc3</chem>
62	 <chem>Fc1ccccc1C(=O)Nc2nc(c(s2)c3ccccc3)c4ccccc4</chem>



Cmpd #	Compound
63	 <chem>CCc1ccc(OCC(=O)Nc2nc3ccccc3s2)cc1</chem>
64	 <chem>CC1=C(C)SC(=NC1C)NC(=O)COc2ccccc2C(C)C</chem>



Cmpd #	Compound
65	 <chem>CC1(CCN(C1)CC(=O)Nc2nc(Cc3ccccc3c4ccccc42)cs2</chem>
66	 <chem>CC1(C)N(C(=N1)NC(=O)c2ccc(C)cc2)S1</chem>



Cmpd #	Compound
67	 <chem>Cc1ccc(cc1)C(=O)Nc2sc(C3=CC=CC=C3)nc2</chem>
68	 <chem>c1ccc(cc1)C(=O)Nc2sc(C3=CC=CC=C3)c(C4=CC=CC=C4)n2</chem>



Cmpd #	Compound
69	 <chem>O=C(Nc1nc(C2=CC=C(C=C2)OC)c(C3=CC=CC=C3)s1)c4ccccc4</chem>
70	 <chem>Cc1ccc(cc1)C(=O)Nc2nc(C3=CC=CC=C3)c(C4=CC=CC=C4)s2</chem>



Cmpd #	Compound
71	 <chem>CC1=CC=C(C=C1)C(=O)Nc2nc(c3ccccc3)c4ccccc24</chem>
72	 <chem>COc1ccc(cc1)C(=O)Nc2nc(c3ccccc3)c4ccccc24</chem>



Cmpd #	Compound
73	 <chem>Clc1cc(Cl)ccc1C(=O)Nc2nc(c(c2)c3ccccc3)c4ccccc4</chem>
74	 <chem>Clc1ccc(cc1)C(=O)Nc2nc(c(c2)c3ccccc3)c4ccccc4</chem>



Cmpd #	Compound
75	 <chem>CC1=CC=C(C(=O)Nc2nc(C3=CC=CC=C3)cs2)C=C1</chem>
76	 <chem>COc1cccc(C(=O)Nc2nc(C3=CC=CC=C3)cs2)c1</chem>



Cmpd #	Compound
77	 <chem>Nc1nc(s1)C(=O)c2ccccc2Cl</chem>
78	 <chem>COc1ccc(cc1)C(=O)Nc2nc(s2)C3=CC=CC=C3</chem>



Cmpd #	Compound
79	 Chemical structure of compound 79: 1-(4-methoxyphenyl)-2-((2-methoxyphenyl)amino)thiazole. The structure consists of a thiazole ring substituted at the 1-position with a 4-methoxyphenyl group and at the 2-position with a (2-methoxyphenyl)amino group.
80	 Chemical structure of compound 80: 1-(4-methoxyphenyl)-2-((4-methoxyphenyl)amino)thiazole. The structure consists of a thiazole ring substituted at the 1-position with a 4-methoxyphenyl group and at the 2-position with a (4-methoxyphenyl)amino group.



Cmpd #	Compound
81	 <chem>COc1ccc(cc1)-c2cc(sc2-c3ccccc3)C(=O)N</chem>
82	 <chem>COc1ccc(cc1)-c2cc(sc2-c3ccc(OC)cc3)C(=O)N</chem>



Cmpd #	Compound
83	 Chemical structure of compound 83: 4-(4-methoxyphenyl)-2-((4-methylphenyl)amino)thiazole. The structure consists of a thiazole ring substituted at the 2-position with a (4-methylphenyl)amino group and at the 4-position with a 4-methoxyphenyl group. The thiazole ring is connected to a benzene ring at the 4-position, which has a methoxy group (-OCH₃) at the para position. The thiazole ring is also connected to an amino group (-NH-) at the 2-position, which is further connected to a benzene ring at the para position, which has a methyl group (-CH₃) at the para position.
84	 Chemical structure of compound 84: 4-(4-methoxyphenyl)-2-((4-methoxyphenyl)amino)thiazole. The structure consists of a thiazole ring substituted at the 2-position with a (4-methoxyphenyl)amino group and at the 4-position with a 4-methoxyphenyl group. The thiazole ring is connected to a benzene ring at the 4-position, which has a methoxy group (-OCH₃) at the para position. The thiazole ring is also connected to an amino group (-NH-) at the 2-position, which is further connected to a benzene ring at the para position, which has a methoxy group (-OCH₃) at the para position.



Cmpd #	Compound
85	 <chem>CC(C)C(=O)Nc1cc(sc1-c2ccccc2)</chem>
86	 <chem>CC(C)CC(=O)Nc1cc(sc1-c2ccccc2)</chem>



Cmpd #	Compound
87	 <chem>CC(C)(C)C(=O)Nc1nc(C2=CC=CC=C2)cs1</chem>
88	 <chem>CC1=CC=C(C=C1)C(=O)Nc2nc(C3=CC=CC=C3)cs2</chem>



Cmpd #	Compound
89	 <chem>O=C(Nc1ccsc1Cc2ccccc2)c3c4ccccc4Oc5ccccc35</chem>
90	 <chem>O=C(Nc1ccsc1Cc2cc(Cl)ccc2)c3c4ccccc4Oc5ccccc35</chem>



Cmpd #	Compound
91	 <chem>Clc1ccccc1Cc2nc(NC(=O)C(c3ccccc3)C4CCCCC4)s2</chem>
92	 <chem>c1ccccc1Cc2nc3c(ncn3)NC(=O)C(c4ccccc4)C5CCCCC5</chem>



Cmpd #	Compound
93	 <chem>O=C(Nc1cc(Cc2ccccc2)sc1)c3c4ccccc4c5ccccc35</chem>
95	 <chem>O=C(Nc1cc(Cc2cc(Cl)cc(Cl)c2)sc1)C(c3ccc(Cl)cc3)c4ccc(Cl)cc4</chem>



Cmpd #	Compound
96	 <chem>CC(C(=O)Nc1nc(Cc2ccccc2)s1)c3ccccc3</chem>
97	 <chem>CC(C(=O)Nc1nc(Cc2cc(Cl)ccc2)s1)c3ccccc3</chem>



Cmpd #	Compound
98	 <chem>CC(C)C(C(=O)Nc1ccsc1Cc2ccccc2)c3ccccc3</chem>
99	 <chem>CC(C)C(C(=O)Nc1nc(Cc2cc(Cl)ccc2)sc1)c3ccccc3</chem>



Cmpd #	Compound
100	 <chem>CC(C)C(C(=O)Nc1nc(Cc2ccccc2)oc1)C(c3ccccc3)C</chem>
101	 <chem>CC(=O)Nc1nc(Cc2ccccc2)sc1C(C(=O)Nc3ccccc3)C4CCCC4</chem>



Cmpd #	Compound
102	 <chem>O=C(C1CCCC1)Nc2nc(Cc3ccccc3Cl)cs2c4ccccc4</chem>
103	 <chem>CC(C)C(C1=CC=CC=C1)C(=O)Nc2nc(Cc3ccccc3Cl)cs2</chem>



Cmpd #	Compound
104	 <chem>Nc1nc(c(s1)c2ccccc2)c3cc(Cl)ccc3</chem>
105	 <chem>COc1cccc(Nc2nc(c(s2)c3ccccc3)c4ccccc4)c1</chem>



Cmpd #	Compound
106	 <chem>Clc1ccc(cc1)C(=O)Nc2nc(c(s2)C3=CC=CC=C3)C4=CC=CC=C4</chem>
107	 <chem>c1ccc(cc1)C(=O)Nc2nc(c(s2)C3=CC=CC=C3)C4=CC=CC=C4</chem>



Cmpd #	Compound
108	 <chem>CCCC(=O)Nc1nc(c2ccccc12)C3=CC=CC=C3</chem>
109	 <chem>CC(C)C(=O)Nc1nc(c2ccccc12)C3=CC=CC=C3</chem>



Cmpd #	Compound
110	 <chem>CC(C)C(=O)Nc1nc(c(s1)c2ccccc2)c3ccccc3</chem>
111	 <chem>C1CC1C(=O)Nc2nc(c(s2)c3ccccc3)c4ccccc4</chem>



Cmpd #	Compound
112	 <chem>CC(C)(N)C(=O)Nc1sc(C2=CC=CC=C2)c(C3=CC=CC=C3)c1</chem>
113	 <chem>CC(C)(N)C(=O)Nc1sc(C2=CC=CC=C2)c(C3=CC=CC=C3)c1</chem>



Cmpd #	Compound
114	 <chem>c1ccccc1c2nc(s2)C(=N)NC(=O)C3CCCCC3</chem>
115	 <chem>c1ccccc1c2nc(s2)C(=N)NC(=O)C3OCCC3</chem>



Cmpd #	Compound
116	 <chem>O=C(O[C@H]1CCOC1)Nc2nc(c(s2)C1=CC=CC=C1)C2=CC=CC=C2</chem>
117	 <chem>O=C(O[C@H]1CCCCC1)Nc2nc(c(s2)C1=CC=C(C=C1)Cl)C2=CC=CC=C2</chem>



Cmpd #	Compound
118	 <chem>O=C(Nc1nc(Cc2ccc(Cl)cc2)sc1)C(c3ccc(Cl)cc3)Cc4ccc(Cl)cc4</chem>
119	 <chem>CC(C1=CC=CC=C1)C(=O)Nc2nc(Cc3ccc(Cl)cc3)sc2C(=O)Cc4ccccc4</chem>



Cmpd #	Compound
120	 <chem>CC(C1=CC=CC=C1)C(=O)Nc2nc(Cc3ccc(Cl)cc3)sc2</chem>
121	 <chem>CC(C)C(C1=CC=CC=C1)C(=O)Nc2nc(Cc3ccc(Cl)cc3)sc2</chem>



Cmpd #	Compound
122	 <chem>O=C(Nc1nc(Cc2ccc(Cl)cc2)sc1)C(c3ccccc3)CC4CCCC4</chem>
123	 <chem>O=C(Nc1nc(Cc2ccc(Cl)cc2)sc1)C(c3ccccc3)C(C)C4=CC=CC=C4</chem>



Cmpd #	Compound
124	 <chem>O=C(Nc1ccsc1Cc2ccc(Cl)cc2)C(c3ccccc3)Cc4ccccc4</chem>
125	 <chem>O=C(Nc1ccsc1Cc2ccc(Cl)cc2)C(c3ccc(Cl)cc3)Cc4ccc(Cl)cc4</chem>



Cmpd #	Compound
126	 <chem>Clc1ccc(cc1)Cc2sc(NC(=O)c3c4ccccc4CC3)c[nH]2</chem>
127	 <chem>CC(C1=CC=CC=C1)C(=O)Nc2nc(Cc3ccc(Cl)cc3)sc2</chem>



Cmpd #	Compound
128	 <chem>CC(C(=O)Nc1nc(Cc2ccc(Cl)cc2)s1)c3ccccc3</chem>
129	 <chem>CC(C)C(=O)Nc1nc(Cc2ccc(Cl)cc2)s1</chem>



Cmpd #	Compound
130	 <chem>CC(C)(c1ccccc1)C(=O)Nc2cc(sc2Cc3ccc(Cl)cc3)</chem>
131	 <chem>O=C1Nc2cc(sc2Cc3ccc(Cl)cc3)C(=O)N1c4c5ccccc5oc6ccccc46</chem>



Cmpd #	Compound
132	 <chem>COc1ccc(cc1)Cc2cc(sc2)NC(=O)Cc3ccccc3</chem>
133	 <chem>COc1ccc(cc1)Cc2cc(sc2)NC(=O)C3c4ccccc4c5ccccc35</chem>



Cmpd #	Compound
134	 <chem>COc1ccc(cc1)Cc2cc(sc2)NC(=O)C(c3ccc(Cl)cc3)Cc4ccc(Cl)cc4</chem>
135	 <chem>COc1cccc(c1)Cc2cc(sc2)NC(=O)C(C)c3ccccc3</chem>



Cmpd #	Compound
136	 <chem>CC(C(=O)Nc1nc(Cc2cc(OC)ccc2)s1)c3ccccc3</chem>
137	 <chem>CC(C)(C(=O)Nc1nc(Cc2cc(OC)ccc2)s1)c3ccccc3</chem>



Cmpd #	Compound
138	 Chemical structure of compound 138: 2-(3-methoxybenzyl)-4-((S)-1-phenylpropan-2-ylamino)thiazole. The structure features a thiazole ring substituted at the 2-position with a 3-methoxybenzyl group and at the 4-position with an (S)-1-phenylpropan-2-ylamino group.
139	 Chemical structure of compound 139: 2-(benzylthio)-4-((S)-1-phenylcyclohexylamino)thiazole. The structure features a thiazole ring substituted at the 2-position with a benzylthio group and at the 4-position with an (S)-1-phenylcyclohexylamino group.



Cmpd #	Compound
140	 <chem>COc1ccc(cc1)CC2=CN=C(S2)NC(=O)C3c4ccccc4Oc5ccccc35</chem>
141	 <chem>c1ccc(cc1)CC2=CN=C(O2)NC(=O)c3ccccc3</chem>



Cmpd #	Compound
142	
143	



Cmpd #	Compound
145	 <chem>O=C1C2=CC=CC=C2O1C(=O)Nc3cc(Cl)ccc3</chem>
146	 <chem>CC(C1=CC=CC=C1)C(=O)Nc2cc(Cl)ccc2</chem>



Cmpd #	Compound
147	 <chem>CC(C)C(C(=O)Nc1cc(Cc2ccc(Cl)cc2)oc1)c3ccccc3</chem>
148	 <chem>C1CCC(CC1)C(C(=O)Nc2cc(Cc3ccc(Cl)cc3)oc2)c4ccccc4</chem>



Cmpd #	Compound
149	 <chem>CC(=O)Nc1nc(Cc2ccc(Cl)cc2)oc1Cc3ccccc3</chem>
150	 <chem>CC(=O)Nc1nc(Cc2ccc(Cl)cc2)oc1Cc3ccccc3</chem>



Cmpd #	Compound
151	 <chem>Clc1ccc(cc1)COc2nc(O=C(c3ccccc3)C4CCCCC4)nn2</chem>
152	 <chem>Clc1ccc(cc1)COc2nc(NC(=O)C3Cc4ccccc43)nn2</chem>



Cmpd #	Compound
153	
154	



Cmpd #	Compound
155	
156	



Cmpd #	Compound
157	
158	



Cmpd #	Compound
159	
160	



Cmpd #	Compound
161	 Chemical structure of compound 161: N-(4-cyanophenyl)-2-(benzoylamino)thiazole. The structure consists of a thiazole ring substituted at the 2-position with a benzoylamino group (a benzene ring attached to a carbonyl group, which is attached to the nitrogen of the thiazole ring) and at the 4-position with a 4-cyanophenyl group (a benzene ring attached to a cyano group).
162	 Chemical structure of compound 162: N-(3,5-dimethoxyphenyl)-2-(benzoylamino)thiazole. The structure consists of a thiazole ring substituted at the 2-position with a benzoylamino group (a benzene ring attached to a carbonyl group, which is attached to the nitrogen of the thiazole ring) and at the 4-position with a 3,5-dimethoxyphenyl group (a benzene ring with methoxy groups at the 3 and 5 positions, attached to the thiazole ring).



Cmpd #	Compound
163	
164	



Cmpd #	Compound
165	 <chem>O=C(Nc1nc(C2=CC=C(C=C2)Cl)sc1C3=CC=CC=C3)C4=CC=CC=C4</chem>
166	 <chem>O=C(NC(=O)C(C)C5=CC=CC=C5)c1nc(C2=CC=CC=C2)sc1</chem>



Cmpd #	Compound
167	 <chem>c1ccc(cc1)-c2nc(s2)NC(=O)C(c3ccccc3)C</chem>
168	 <chem>c1ccc(cc1)-c2nc(s2)NC(=O)C(c3ccccc3)C(C)C</chem>



Cmpd #	Compound
169	
170	



Cmpd #	Compound
171	 <chem>COC(=O)c1ccccc1C(=O)Nc2nc(c3ccccc3s2)c4ccccc4</chem>
172	 <chem>Cc1ccccc1C(=O)Nc2nc(c3ccccc3s2)c4ccccc4</chem>



Cmpd #	Compound
173	 <chem>CC(C1=CC=CC=C1)C(=O)Nc2nc(c3ccccc3s2)c4ccccc4</chem>
174	 <chem>CC(C)C(C1=CC=CC=C1)C(=O)Nc2nc(c3ccccc3s2)c4ccccc4</chem>



Cmpd #	Compound
175	 <chem>CC1(Cc2ccccc2)C(=O)Nc3nc(cc3C4=CC=CC=C4C5=CC=CC=C5)C6=CC=CC=C6</chem>
176	 <chem>CCCC(C)C(Cc1ccccc1)C(=O)Nc2nc(cc2C3=CC=CC=C3C4=CC=CC=C4)S5=CC=CC=C5</chem>



Cmpd #	Compound
177	 <chem>c1ccc(cc1)c2c(c(c(s2)N1C(=O)C(c1ccccc1)C2CCCCC2)N)C3=CC=CC=C3</chem>
178	 <chem>COC(=O)c1ccccc1C(=O)Nc2nc(c(s2)C3=CC=CC=C3)C4=CC=CC=C4</chem>



Cmpd #	Compound
179	 <chem>COc1ccc(cc1)Cc2cc3nc(NC(=O)C(c4ccccc4)c5ccccc5)oc3cc2</chem>
180	 <chem>COc1ccc(cc1)Cc2cc3nc(NC(=O)C(c4ccccc4)c5ccccc5)oc3cc2</chem>



Cmpd #	Compound
181	 <chem>COc1ccc(cc1Cc2oc(C(=O)Nc3c4ccccc4c3)c2)</chem>
182	 <chem>COc1ccc(cc1Cc2oc(C(=O)C(C)C3=CC=CC=C3)c2)</chem>



Cmpd #	Compound
183	 Chemical structure of compound 183: 1-(4-methoxybenzyl)-4-phenyl-1H-imidazole-5-carboxamide, 1-ethyl-. The structure shows a benzimidazole core with a 4-methoxybenzyl group at position 1 and a 1-ethyl-4-phenylcarbamoyl group at position 5. <chem>CCOC1=CC=C(C=C1)CC2=CN=C(NC(=O)C(C)C3=CC=CC=C3)O2</chem>
184	 Chemical structure of compound 184: 1-(4-methoxybenzyl)-4-phenyl-1H-imidazole-5-carboxamide, 1-isopropyl-. The structure shows a benzimidazole core with a 4-methoxybenzyl group at position 1 and a 1-isopropyl-4-phenylcarbamoyl group at position 5. <chem>CCOC1=CC=C(C=C1)CC2=CN=C(NC(=O)C(C)C(C)C3=CC=CC=C3)O2</chem>



Cmpd #	Compound
185	 Chemical structure of compound 185: 1-(4-methoxybenzyl)-2-(benzoylamino)oxazole. The structure consists of a benzoylamino group (a benzene ring attached to a carbonyl group, which is attached to an NH group) connected to the 2-position of an oxazole ring. The 1-position of the oxazole ring is connected to a methoxybenzyl group (a benzene ring with a methoxy group at the para position, attached to a CH2 group).
186	 Chemical structure of compound 186: 1-(4-chlorobenzyl)-2-(1-cyclohexyl-2-phenylethan-1-yl)oxazole. The structure consists of a 1-cyclohexyl-2-phenylethan-1-yl group (a cyclohexane ring attached to a CH group, which is attached to a CH2 group, which is attached to a phenyl ring) connected to the 2-position of an oxazole ring. The 1-position of the oxazole ring is connected to a 4-chlorobenzyl group (a benzene ring with a chlorine atom at the para position, attached to a CH2 group).



Cmpd #	Compound
187	 Chemical structure of compound 187: A 1,3,4-oxadiazole ring is substituted at the 2-position with a 3-chlorobenzyl group and at the 5-position with a carbonyl group. The carbonyl group is part of an amide linkage to a 1,1-diphenylethyl group, where one phenyl ring is substituted with a chlorine atom at the para position. <chem>Clc1ccccc1Cc2nc([N-]H)c([N+])c2C(=O)C(c3ccccc3Cl)c4ccccc4</chem>
188	 Chemical structure of compound 188: A 1,3,4-oxadiazole ring is substituted at the 2-position with a 3-chlorobenzyl group and at the 5-position with a carbonyl group. The carbonyl group is part of an amide linkage to a bicyclo[4.1.0]hept-2-en-2-yl group. <chem>Clc1ccccc1Cc2nc([N-]H)c([N+])c2C(=O)Nc3C=Cc4ccccc34</chem>



Cmpd #	Compound
189	 <chem>CC(C(=O)Nc1nc(Cc2ccccc2Cl)oc1)c3ccccc3</chem>
190	 <chem>CCC(C(=O)Nc1nc(Cc2ccccc2Cl)oc1)c3ccccc3</chem>



Cmpd #	Compound
191	 <chem>CC(C)C(C(=O)Nc1ccoc1Cc2ccccc2Cl)c3ccccc3</chem>
192	 <chem>C1CCCC1C(C(=O)Nc2ccoc2Cc3ccccc3Cl)c4ccccc4</chem>



Cmpd #	Compound
193	 <chem>CC(C)C(C(=O)Nc1nc(Cc2ccccc2Cl)co1)c3ccccc3</chem>
194	 <chem>Nc1ccccc1C(=O)Nc2nc(Cc3ccccc3Cl)co2</chem>



Cmpd #	Compound
195	 <chem>Nc1ccccc1C(=O)Nc2nc(Cc3ccccc3Cl)oc2</chem>
196	 <chem>COc1ccc(cc1)N2CCCN2C(=O)Nc3sc(cc3c4ccccc4)C5=CC=CC=C5</chem>



Cmpd #	Compound
197	
198	



Cmpd #	Compound
199	 Chemical structure of compound 199: A 4-bromophenyl group is connected via a sulfonyl group (SO₂) to a piperazine ring. The piperazine ring is further connected to a 1-(4-phenylthiazol-2-yl)propan-1-one moiety. The thiazole ring is substituted with a phenyl group at the 4-position.
201	 Chemical structure of compound 201: A 4-chlorophenyl group is connected via a methylene group to a thiazole ring. The thiazole ring is substituted with a 1-phenylpropan-1-one moiety. The thiazole ring is also substituted with a phenyl group at the 4-position.



Cmpd #	Compound
202	 <chem>CC([C@@H](C(=O)c1ccccc1)Nc2nc(s2)Cc3cc(Cl)ccc3)c4ccccc4</chem>
203	 <chem>CC([C@H](C(=O)c1ccccc1)Nc2nc(s2)Cc3ccc(Cl)cc3)c4ccccc4</chem>



Cmpd #	Compound
204	 <chem>CC([C@@H](C(=O)Nc1cc(s1)Cc2ccc(Cl)cc2)c3ccccc3)C(=O)N</chem>
205	 <chem>CC([C@@H](C(=O)Nc1cc(s1)Cc2ccc(Cl)cc2)c3ccccc3)C(=O)N</chem>



Cmpd #	Compound
206	 Chemical structure of compound 206: 2-(4-chlorobenzyl)-4-(3-methylbutanoylamino)thiazole. The structure consists of a thiazole ring substituted at the 2-position with a 4-chlorobenzyl group and at the 4-position with a 3-methylbutanoylamino group. <chem>CC(C)CC(=O)Nc1nc(Cc2ccc(Cl)cc2)s1</chem>
207	 Chemical structure of compound 207: 2-(4-chlorobenzyl)-4-(cyclopropylamino)thiazole. The structure consists of a thiazole ring substituted at the 2-position with a 4-chlorobenzyl group and at the 4-position with a cyclopropylamino group. <chem>C1CC1C(=O)Nc2nc(Cc3ccc(Cl)cc3)s2</chem>



Cmpd #	Compound
208	 <chem>CCCC(=O)Nc1nc(Cc2ccccc2Cl)s1</chem>
209	 <chem>C1CCC(CC1)C(=O)Nc2nc(Cc3ccccc3Cl)s2</chem>



Cmpd #	Compound
210	 <chem>CCCC(=O)Nc1nc(Cc2ccc(Cl)cc2)s1</chem>
211	 <chem>CC(NC(=O)Nc1nc(Cc2ccc(Cl)cc2)s1)Cc3ccccc3</chem>



Cmpd #	Compound
212	 <chem>CCN(CC)C(=O)Nc1nc(Cc2ccccc2Cl)s1Cc3ccccc3</chem>
213	 <chem>CCN(CC)C(=O)Nc1nc(Cc2ccccc2Cl)s1Cc3ccccc3</chem>



Cmpd #	Compound
214	 <chem>Clc1ccc(cc1)Cc2cc[nH]c2NC(=O)C(c3ccccc3)N4CCOCC4</chem>
215	 <chem>Clc1ccc(cc1)Cc2cc[nH]c2NC(=O)C(c3ccccc3)N4CCCCC4</chem>



Cmpd #	Compound
216	 <chem>CCOC(=O)C(N)N1C=NC(SC1Cc2ccc(Cl)cc2)=N</chem>
217	 <chem>CCNC(=O)N1C=NC(SC1Cc2ccc(Cl)cc2)=N</chem>



Cmpd #	Compound
218	 <chem>CC(C)C(=O)Nc1nc(Cc2ccccc2Cl)s1</chem>
219	 <chem>CCOC(C)C(=O)Nc1nc(Cc2ccccc2Cl)s1</chem>



Cmpd #	Compound
220	 <chem>CCOC(C)C(=O)Nc1nc(Cc2ccc(Cl)cc2)sc1</chem>
221	 <chem>CC(C)OC(=O)Nc1nc(Cc2ccc(Cl)cc2)sc1</chem>



Cmpd #	Compound
222	 <chem>CCN(C)C(=O)Nc1cc(Cc2ccccc2Cl)cs1</chem>
223	 <chem>CCN1CCOCC1C(=O)Nc2cc(Cc3ccccc3Cl)cs2</chem>



Cmpd #	Compound
224	 <chem>CCSCC1CCN1C(=N)Cc2cc(Cl)ccc2</chem>
225	 <chem>CC1CCN1C(=N)Cc2cc(Cl)ccc2</chem>



Cmpd #	Compound
226	 <chem>CC1=CC=C(C=C1)C(=O)Nc2nc(s2)c3ccccc3</chem>
227	 <chem>COc1cccc(c1)c2nc(s2)c3cc(ccc3C(=O)N)Cl</chem>



Cmpd #	Compound
228	 <chem>CC1=CC=C(C=C1)C(=O)Nc2nc(C3=CC=C(C=C3)Cl)cs2</chem>
229	 <chem>ClC1=CC=C(C=C1)C(=O)Nc2nc(C3=CC=C(C=C3)Cl)cs2</chem>



Cmpd #	Compound
230	 <chem>CC1=CC=C(C=C1)C(=O)Nc2nc(c3cc(Cl)ccc3s2)C(=O)N</chem>
231	 <chem>CC1=CC=C(C=C1)C(=O)Nc2nc(c3ccc(OC)cc3s2)C(=O)N</chem>



Cmpd #	Compound
232	 <chem>COc1ccc(cc1)-c2cc(sc2)NC(=O)c3ccc(Cl)cc3</chem>
233	 <chem>Clc1ccc(cc1)C(=O)Nc2cc(sc2)c3ccccc3</chem>



Cmpd #	Compound
234	 <chem>CCOC(=O)C(c1ccccc1)NC2=NC=C(Cc3ccccc3Cl)S2</chem>
235	 <chem>C1CCNC1C(=O)NC2=NC=C(Cc3ccccc3Cl)S2</chem>



Cmpd #	Compound
237	 Chemical structure of compound 237: 1-(4-(3-methoxybenzyl)-1,3,4-thiazol-5-yl)propan-1-one, 2-phenyl-2-methyl-. The structure features a thiazole ring substituted at the 4-position with a (3-methoxyphenyl)methyl group and at the 5-position with a 2-phenyl-2-methylpropanamido group.
238	 Chemical structure of compound 238: 1-(4-(3-chlorobenzyl)-1,3,4-thiazol-5-yl)propan-1-one, 2-phenyl-2-methyl-. The structure features a thiazole ring substituted at the 4-position with a (3-chlorophenyl)methyl group and at the 5-position with a 2-phenyl-2-methylpropanamido group.



Cmpd #	Compound
239	 <chem>CCN(Cc1ccccc1)C(=O)Nc2nc(Cc3ccccc3Cl)s2</chem>
240	 <chem>CCN(C1CCCCC1)(Cc2ccccc2)C(=O)Nc3nc(Cc4ccccc4Cl)s3</chem>



Cmpd #	Compound
241	 <chem>O=C(Nc1cc[nH]1C2=CC=CC=C2)C(N3C=CC=C3)Cc4ccc(Cl)cc4</chem>
242	 <chem>CC1=CC=C(C=C1)N(C1=CC=CC=C1)C(=O)Nc2cc[nH]2C3=CC=CC=C3</chem>



Cmpd #	Compound
243	 <chem>CC1CCN(CC1)C(=O)C(NC2=CN=C(C2)Cc3ccccc3Cl)Cc4ccccc4</chem>
244	 <chem>OC1CCN(CC1)C(=O)C(NC2=CN=C(C2)Cc3ccccc3Cl)Cc4ccccc4</chem>



Cmpd #	Compound
245	 <chem>CCOC(=O)C1CCN(C1)C(C(=O)Nc2cc(sc2Cc3ccccc3Cl)c4ccccc4)C5=CC=CC=C5</chem>
246	 <chem>CC1CCN(C1)C(C(=O)Nc2cc(sc2Cc3ccccc3Cl)c4ccccc4)C5=CC=CC=C5</chem>



Cmpd #	Compound
247	 Chemical structure of compound 247: A thiazole ring substituted with a 4-chlorobenzyl group at the 4-position and an N-(1-((4-ethoxybenzoyl)amino)-4-phenylpiperidin-4-yl) group at the 2-position. <chem>CCOC(=O)C1CCN(C1)C(=O)Nc2nc(Cc3ccc(Cl)cc3)cs2</chem>
248	 Chemical structure of compound 248: A thiazole ring substituted with a 4-chlorobenzyl group at the 4-position and an N-(1-hydroxy-4-phenylpiperidin-4-yl) group at the 2-position. <chem>O[C@@H]1CCN(C1)C(=O)Nc2nc(Cc3ccc(Cl)cc3)cs2</chem>



Cmpd #	Compound
249	 <chem>Clc1ccccc1Cc2cc(sc2NC(=O)C(c3ccccc3)N4CCN(C5=NC=CC=N5)CC4)</chem>
250	 <chem>Clc1ccccc1Cc2cc(sc2NC(=O)N(C3=CC=CC=C3)NC4CCCC4)</chem>



Cmpd #	Compound
251	 <chem>CC(NC(=O)Nc1nc(Cc2ccc(Cl)cc2)cs1)CCc3ccccc3</chem>
252	 <chem>CC(O)C1c2ccccc2N1c3nc(Cc4ccc(Cl)cc4)cs3</chem>



Cmpd #	Compound
253	 <chem>CCOC(=O)C1CCCCN1C(C(=O)Nc2nc(Cc3ccccc3Cl)s2)c4ccccc4</chem>
254	 <chem>CN(C)Cc1ccccc1C(C(=O)Nc2nc(Cc3ccccc3Cl)s2)c4ccccc4</chem>



Cmpd #	Compound
255	 <chem>CC1=CC=C(C=C1)C(=O)Nc2nc(Cc3cc(Cl)ccc3)cs2</chem>
256	 <chem>C1=CC=C(C=C1)C(=O)Nc2nc(Cc3cc(C#N)ccc3)cs2</chem>



Cmpd #	Compound
257	
258	



Cmpd #	Compound
259	 <chem>CC(C)C(C(=O)Nc1nc(Cc2ccc(C#N)cc2)s1)c3ccccc3</chem>
260	 <chem>C1CCC(CC1)C(C(=O)Nc2nc(Cc3ccc(C#N)cc3)s2)c4ccccc4</chem>



Cmpd #	Compound
261	 <chem>CC(C)(C)C(=O)Nc1nc(Cc2ccc(C#N)cc2)s1</chem>
262	 <chem>CCC(C(=O)Nc1nc(C(=O)C2=CC=CC=C2Cl)s1)C3=CC=CC=C3</chem>



Cmpd #	Compound
263	 <chem>CC(C1=CC=CC=C1)C(=O)NC2=CN=C(C3=CC=CC=C3Cl)C2O</chem>
264	 <chem>CC(C1=CC=CC=C1)C(=O)NC2=NC3=CC=C(C=C3S2)S(=O)(=O)C</chem>



Cmpd #	Compound
265	 <chem>CCOC1=CC=C(C=C1)c2nc(C(=O)C(c3ccccc3)CC)cs2</chem>
266	 <chem>COC1=CC=C(C=C1)c2nc(C(=O)C(c3ccccc3)CC)cs2</chem>



Cmpd #	Compound
267	 Chemical structure of compound 267: 2-((1-ethyl-1-phenylethyl)carbamoyl)-5,6-dimethyl-1,2,4-benzothiazole. The structure features a benzothiazole ring system with methyl groups at the 5 and 6 positions. The 2-position of the benzothiazole ring is substituted with a carbonyl group, which is further substituted with a 1-ethyl-1-phenylethyl group.
268	 Chemical structure of compound 268: 2-((1-ethyl-1-phenylethyl)carbamoyl)-4-methyl-1,2,4-benzothiazole. The structure features a benzothiazole ring system with a methyl group at the 4-position. The 2-position of the benzothiazole ring is substituted with a carbonyl group, which is further substituted with a 1-ethyl-1-phenylethyl group.



Cmpd #	Compound
269	
270	



Cmpd #	Compound
271	 <chem>CC(C1=CC=CC=C1)C(=O)Nc2nc(Cc3ccc(Cl)cc3)sc2</chem>
272	 <chem>CC(C1=CC=CC=C1)C(=O)Nc2nc(Cc3ccc(Cl)cc3)sc2</chem>



Cmpd #	Compound
273	 Chemical structure of compound 273: A thiazole ring substituted with a 4-methoxybenzyl group at position 2, an N-(2-methoxyphenyl)propanamide group at position 4, and an N-(4-methoxyphenyl)propanamide group at position 5. <chem>COc1ccc(cc1)C(=O)Nc2sc(cc2Cc3ccccc3OC)C(=O)Nc4ccc(OC)cc4</chem>
274	 Chemical structure of compound 274: A thiazole ring substituted with a 2-methoxybenzyl group at position 4, an N-(1-phenylpropan-2-yl)propanamide group at position 2, and an N-(1-phenylpropan-2-yl)propanamide group at position 5. <chem>COc1ccccc1Cc2ccn(c2)C(=O)Nc3ccccc3C(C)C(=O)Nc4ccccc4C(C)C(=O)N</chem>



Cmpd #	Compound
275	 Chemical structure of compound 275: A thiazole ring substituted with a 4-methoxybenzyl group at position 4 and an N-(1-phenyl-1-(dimethylamino)ethyl)carbamoyl group at position 2. <chem>COc1ccc(cc1)Cc2cc(C(=O)N(Cc3ccccc3)C(C)(C)N)nn2</chem>
276	 Chemical structure of compound 276: A thiazole ring substituted with a 3-cyanobenzyl group at position 4 and an N-(1-phenylpropan-2-yl)carbamoyl group at position 2. <chem>N#Cc1ccccc1Cc2cc(C(=O)N(Cc3ccccc3)CC)nn2</chem>



Cmpd #	Compound
277	
278	


Cmpd #	Compound
279	 <chem>CC1=CN=C(S1)NC(=O)C(c2ccccc2)C(c3ccccc3)</chem>
280	 <chem>CCOC(=O)c1c[nH]c(s1)NC(=O)C(c2ccccc2)C(c3ccccc3)</chem>

Cmpd #	Compound
281	 <chem>CCOC(=O)C=Cc1nc(s1)C(=O)NC(C(=O)c2ccccc2)c3ccccc3</chem>
282	 <chem>COC(=O)c1ccncc1Cc2nc(s2)C(=O)NC(C(=O)c3ccccc3)c4ccccc4</chem>

Cmpd #	Compound
283	 <chem>COc1ccc(cc1)Cc2cc(sc2)Nc3ccccc3CC</chem>
284	 <chem>COc1ccc(cc1)Cc2cc(sc2)Nc3ccccc3CC</chem>

Cmpd #	Compound
285	 <chem>CC(C1=CC=CC=C1)C(=O)NNC2=NC=C(S2)CC3=CC=C(OC)C(F)=C3</chem>
286	 <chem>CC(C1=CC=CC=C1)C(=O)NNC2=NC=C(S2)CC3=CC=C(OC)C=C3</chem>

Cmpd #	Compound
287	 <chem>CCOC(=O)C1=C(C(=N1)C(=O)N(C2=CC=CC=C2)C3=CC=CC=C3)CC(C)C</chem>
288	 <chem>CCOC(=O)C1=C(C(=N1)C(=O)N(C2=CC=CC=C2)C3=CC=CC=C3)CC(C)C</chem>

Cmpd #	Compound
289	 <chem>CC(C(=O)Nc1nc2c(s1)cccc2C(=O)c3ccccc3)C(c4ccccc4)C</chem>

[00260] According to an alternative embodiment, preferred compounds of the present invention are those that measurably increase the activity of an ABC-transporter or of a fragment thereof, and preferably CFTR activity.

[00261] According to another alternative embodiment, preferred compounds of the present invention are those that measurably decrease the activity of an ABC-transporter or of a fragment thereof.

[00262] One of skill in the art would be well aware of techniques and assays useful in measuring the increase or decrease of activity of an ABC-transporter or of a fragment thereof.

[00263] According to an alternative preferred embodiment, the present invention provides a method of modulating CFTR activity in a cell membrane of a mammal in need thereof, comprising the step of administering to said mammal a composition comprising a compound of the present invention as defined above.

[00264] The preferred embodiments of compound of formula (I) useful in potentiating the activity of CFTR include the preferred embodiments of the present invention described above.

[00265] According to an alternative embodiment, the present invention provides a method of increasing the number of functional ABC transporters in a membrane of a cell, comprising the step of contacting said cell with a compound of the present invention. The term "functional ABC transporter" as used herein means an ABC transporter that is capable of transport activity.

[00266] According to a preferred embodiment, said functional ABC transporter is CFTR.

[00267] The preferred embodiments of compounds of formula (I) useful in increasing the number of functional

ABC transporters include preferred embodiments of the compounds of the present invention as described above.


[00268] Unless otherwise stated, structures depicted herein are also meant to include all stereochemical forms of the structure; i.e., the R and S configurations for each asymmetric center. Therefore, single stereochemical isomers as well as enantiomeric and diastereomeric mixtures of the present compounds are within the scope of the invention. Unless otherwise stated, structures depicted herein are also meant to include compounds that differ only in the presence of one or more isotopically enriched atoms. For example, compounds having the present structures except for the replacement of a hydrogen by a deuterium or tritium, or the replacement of a carbon by a ^{13}C - or ^{14}C -enriched carbon are within the scope of this invention. Such compounds are useful, for example, as analytical tools or probes in biological assays.

[00269] The present invention includes within its scope pharmaceutically acceptable prodrugs of the compounds of the present invention. A "pharmaceutically acceptable prodrug" means any pharmaceutically acceptable salt, ester, salt of an ester, or other derivative of a compound of the present invention which, upon administration to a recipient, is capable of providing (directly or indirectly) a compound of this invention or an active metabolite or residue thereof. Preferred prodrugs are those that increase the bioavailability of the compounds of this invention when such compounds are administered to a mammal or which enhance delivery of the parent compound to a biological compartment relative to the parent species.

[00270] The compounds of the present invention may be readily prepared using methods known in the art. One such synthetic route is illustrated in Scheme 1 below:

Scheme 1


Step 1:




[00271] **2-Aminothiazoles.** The appropriate bromoketone was dissolved in a minimum of ethanol or methanol with 1.1 equivalents of thiourea. The reaction mixture was stirred overnight at room temperature, evaporated to dryness, and then dissolved in either dichloromethane or ethyl acetate. The reaction mixture was then extracted with 1M sodium hydroxide followed by a saturated aqueous sodium chloride solution. The organic layer was then separated, dried over sodium sulfate, and evaporated to dryness to yield the desired 2-aminothiazole.

Step 2:


(a)




(b)



(c)



(d)



[00272] **Amides.** (a) If the appropriate acid chloride was commercially available, it was added to one equivalent of the appropriate amine in minimum of pyridine. The reaction mixture was allowed to stir overnight at room temperature. The reaction mixture was then filtered and evaporated to dryness. The crude

product was purified by reverse phase preparative liquid chromatography.

[00273] (b) Alternatively, the acid chloride was added to one equivalent of the appropriate amine in minimum of 1,4-dioxane containing an excess of triethylamine. The reaction mixture was then either allowed to stir overnight at room temperature or subjected to microwave irradiation for 5 minutes at 200°C. The crude product was then filtered, evaporated to dryness, dissolved in a minimum of dimethylsulfoxide and then purified by reverse phase preparative liquid chromatography.

[00274] (c) If the appropriate acid chloride was not commercially available the appropriate carboxylic acid was added to a solution containing one equivalent of the appropriate amine in a minimum of pyridine. O-(7-Azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (HATU, 1.4 eq.) is added, and the reaction is stirred overnight. The crude product was purified by reverse phase preparative liquid chromatography.

[00275] (d) If the appropriate acid chloride was not commercially available the appropriate carboxylic acid was added to a solution containing one equivalent of the appropriate amine in a minimum of N,N-dimethylformamide with an excess of triethylamine. O-(7-Azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (HATU, 1.4 eq.) is added, and the reaction is stirred overnight. The crude product was purified by reverse phase preparative liquid chromatography.

[00276] One of skill in the art will recognize that the above two synthetic routes are generic and can be

readily exploited for any embodiment of compound formula (I).

[00277] The term "pharmaceutically acceptable carrier, adjuvant, or vehicle" refers to a non-toxic carrier, adjuvant, or vehicle that does not destroy the pharmacological activity of the compound with which it is formulated. Pharmaceutically acceptable carriers, adjuvants or vehicles that may be used in the compositions of this invention include, but are not limited to, ion exchangers, alumina, aluminum stearate, lecithin, serum proteins, such as human serum albumin, buffer substances such as phosphates, glycine, sorbic acid, potassium sorbate, partial glyceride mixtures of saturated vegetable fatty acids, water, salts or electrolytes, such as protamine sulfate, disodium hydrogen phosphate, potassium hydrogen phosphate, sodium chloride, zinc salts, colloidal silica, magnesium trisilicate, polyvinyl pyrrolidone, cellulose-based substances, polyethylene glycol, sodium carboxymethylcellulose, polyacrylates, waxes, polyethylene-polyoxypropylene-block polymers, polyethylene glycol and wool fat.

[00278] Pharmaceutically acceptable salts of the compounds of this invention include those derived from pharmaceutically acceptable inorganic and organic acids and bases. Examples of suitable acid salts include acetate, adipate, alginate, aspartate, benzoate, benzenesulfonate, bisulfate, butyrate, citrate, camphorate, camphorsulfonate, cyclopentanepropionate, digluconate, dodecylsulfate, ethanesulfonate, formate, fumarate, glucoheptanoate, glycerophosphate, glycolate, hemisulfate, heptanoate, hexanoate, hydrochloride, hydrobromide, hydroiodide, 2-hydroxyethanesulfonate, lactate, maleate, malonate, methanesulfonate, 2-

naphthalenesulfonate, nicotinate, nitrate, oxalate, palmoate, pectinate, persulfate, 3-phenylpropionate, phosphate, picrate, pivalate, propionate, salicylate, succinate, sulfate, tartrate, thiocyanate, tosylate and undecanoate. Other acids, such as oxalic, while not in themselves pharmaceutically acceptable, may be employed in the preparation of salts useful as intermediates in obtaining the compounds of the invention and their pharmaceutically acceptable acid addition salts.

[00279] Salts derived from appropriate bases include alkali metal (e.g., sodium and potassium), alkaline earth metal (e.g., magnesium), ammonium and $N^+(C_{1-4} \text{ alkyl})_4$ salts. This invention also envisions the quaternization of any basic nitrogen-containing groups of the compounds disclosed herein. Water or oil-soluble or dispersible products may be obtained by such quaternization.

[00280] The compositions of the present invention may be administered orally, parenterally, by inhalation spray, topically, rectally, nasally, buccally, vaginally or via an implanted reservoir. The term "parenteral" as used herein includes subcutaneous, intravenous, intramuscular, intra-articular, intra-synovial, intrasternal, intrathecal, intrahepatic, intralesional and intracranial injection or infusion techniques. Preferably, the compositions are administered orally, intraperitoneally or intravenously. Sterile injectable forms of the compositions of this invention may be aqueous or oleaginous suspension. These suspensions may be formulated according to techniques known in the art using suitable dispersing or wetting agents and suspending agents. The sterile injectable preparation may also be a sterile injectable solution or suspension in a non-toxic parenterally-acceptable diluent or solvent, for example as a solution in 1,3-butanediol.

Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium.

[00281] For this purpose, any bland fixed oil may be employed including synthetic mono- or di-glycerides. Fatty acids, such as oleic acid and its glyceride derivatives are useful in the preparation of injectables, as are natural pharmaceutically-acceptable oils, such as olive oil or castor oil, especially in their polyoxyethylated versions. These oil solutions or suspensions may also contain a long-chain alcohol diluent or dispersant, such as carboxymethyl cellulose or similar dispersing agents that are commonly used in the formulation of pharmaceutically acceptable dosage forms including emulsions and suspensions. Other commonly used surfactants, such as Tweens, Spans and other emulsifying agents or bioavailability enhancers which are commonly used in the manufacture of pharmaceutically acceptable solid, liquid, or other dosage forms may also be used for the purposes of formulation.

[00282] The pharmaceutically acceptable compositions of this invention may be orally administered in any orally acceptable dosage form including, but not limited to, capsules, tablets, aqueous suspensions or solutions. In the case of tablets for oral use, carriers commonly used include lactose and corn starch. Lubricating agents, such as magnesium stearate, are also typically added. For oral administration in a capsule form, useful diluents include lactose and dried cornstarch. When aqueous suspensions are required for oral use, the active ingredient is combined with emulsifying and suspending

agents. If desired, certain sweetening, flavoring or coloring agents may also be added.

[00283] Alternatively, the pharmaceutically acceptable compositions of this invention may be administered in the form of suppositories for rectal administration. These can be prepared by mixing the agent with a suitable non-irritating excipient that is solid at room temperature but liquid at rectal temperature and therefore will melt in the rectum to release the drug. Such materials include cocoa butter, beeswax and polyethylene glycols.

[00284] The pharmaceutically acceptable compositions of this invention may also be administered topically, especially when the target of treatment includes areas or organs readily accessible by topical application, including diseases of the eye, the skin, or the lower intestinal tract. Suitable topical formulations are readily prepared for each of these areas or organs.

[00285] Topical application for the lower intestinal tract can be effected in a rectal suppository formulation (see above) or in a suitable enema formulation. Topically-transdermal patches may also be used.

[00286] For topical applications, the pharmaceutically acceptable compositions may be formulated in a suitable ointment containing the active component suspended or dissolved in one or more carriers. Carriers for topical administration of the compounds of this invention include, but are not limited to, mineral oil, liquid petrolatum, white petrolatum, propylene glycol, polyoxyethylene, polyoxypropylene compound, emulsifying wax and water. Alternatively, the pharmaceutically acceptable compositions can be formulated in a suitable lotion or cream containing the active components suspended or dissolved in one or more

pharmaceutically acceptable carriers. Suitable carriers include, but are not limited to, mineral oil, sorbitan monostearate, polysorbate 60, cetyl esters wax, cetearyl alcohol, 2-octyldodecanol, benzyl alcohol and water.

[00287] For ophthalmic use, the pharmaceutically acceptable compositions may be formulated as micronized suspensions in isotonic, pH adjusted sterile saline, or, preferably, as solutions in isotonic, pH adjusted sterile saline, either with or without a preservative such as benzylalkonium chloride. Alternatively, for ophthalmic uses, the pharmaceutically acceptable compositions may be formulated in an ointment such as petrolatum.

[00288] The pharmaceutically acceptable compositions of this invention may also be administered by nasal aerosol or inhalation. Such compositions are prepared according to techniques well-known in the art of pharmaceutical formulation and may be prepared as solutions in saline, employing benzyl alcohol or other suitable preservatives, absorption promoters to enhance bioavailability, fluorocarbons, and/or other conventional solubilizing or dispersing agents.

[00289] Most preferably, the pharmaceutically acceptable compositions of this invention are formulated for oral administration.

[00290] The amount of the compounds of the present invention that may be combined with the carrier materials to produce a composition in a single dosage form will vary depending upon the host treated, the particular mode of administration. Preferably, the compositions should be formulated so that a dosage of between 0.01-100 mg/kg body weight/day of the modulator can be administered to a patient receiving these compositions.

[00291] It should also be understood that a specific dosage and treatment regimen for any particular patient

will depend upon a variety of factors, including the activity of the specific compound employed, the age, body weight, general health, sex, diet, time of administration, rate of excretion, drug combination, and the judgment of the treating physician and the severity of the particular disease being treated. The amount of a compound of the present invention in the composition will also depend upon the particular compound in the composition.

[00292] Depending upon the particular condition, or disease, to be treated or prevented, additional therapeutic agents, which are normally administered to treat or prevent that condition, may also be present in the compositions of this invention. As used herein, additional therapeutic agents that are normally administered to treat or prevent a particular disease, or condition, are known as "appropriate for the disease, or condition, being treated."

[00293] According to an alternative embodiment, the present invention provides a method of treating a ABC transporter mediated disease in a mammal, comprising the step of administering to said mammal a composition comprising a compound of the present invention, or a preferred embodiment thereof as set forth above.

[00294] According to a preferred embodiment, the ABC transporter mediated disease is selected from immunodeficiency disorder, inflammatory disease, allergic disease, autoimmune disease, destructive bone disorder, proliferative disorder, infectious disease or viral disease.

[00295] In certain preferred embodiments, the present invention provides a method of treating cystic fibrosis, hereditary emphysema, hereditary hemochromatosis, coagulation-fibrinolysis deficiencies,

such as protein C deficiency, Type-1 hereditary angioedema, lipid processing deficiencies, such as Familial hypercholesterolemia, Type-1 chylomicronemia, abetalipoproteinemia, lysosomal storage diseases, such as I-cell disease/Pseudo-Hurler, mucopolysaccharidoses, Sandhof/Tay-Sachs, Crigler-Najjar type II, polyendocrinopathy/hyperinsulemia, Diabetes mellitus, Laron dwarfism, myeloperoxidase deficiency, primary hypoparathyroidism, melanoma, glycanosis CDG type 1, hereditary emphysema, congenital hyperthyroidism, osteogenesis imperfecta, hereditary hypofibrinogenemia, ACT deficiency, Diabetes insipidus (DI), neurophyseal DI, neprogenic DI, Charcot-Marie Tooth syndrome, Perlizaeus-Merzbacher disease, neurodegenerative diseases such as Alzheimer's disease, Parkinson's disease, amyotrophic lateral sclerosis, progressive supranuclear palsy, Pick's disease, several polyglutamine neurological disorders asuch as Huntington, spinocerebellar ataxia type I, Spinal and bulbar muscular atrophy, dentatorubal pallidoluysian, and myotonic dystrophy, as well as spongiform encephalopathies, such as hereditary Creutzfeldt-Jakob disease (due to prion protein processing defect), Fabry disease, Straussler-Scheinker disease, secretory diarrhea, polycystic kidney disease, chronic obstructive pulmonary disease (COPD), dry eye disease, and Sjögren's Syndrome, comprising the step of administering to said mammal an effective amount of a composition comprising a compound of the present invention, or a preferred embodiment thereof as set forth above.

[00296] According to an alternative preferred embodiment, the present invention provides a method of treating cystic fibrosis comprising the step of administering to said mammal a composition comprising the

step of administering to said mammal an effective amount of a composition comprising a compound of the present invention.

[00297] According to the invention an "effective amount" of the compound or pharmaceutically acceptable composition is that amount effective for treating or lessening the severity of any one of the above diseases.

[00298] The compounds and compositions, according to the method of the present invention, may be administered using any amount and any route administration effective for treating or lessening the severity of any one of the above diseases.

[00299] In one embodiment, the compounds of the present invention are useful in treating cystic fibrosis.

[00300] According to a more preferred embodiment, the disease so treated is selected from Tangier's disease, stargardt disease 1, age related macular dystrophy 2, retinitis pigmentosa, bare lymphocyte syndrome, PFIC-3, anemia, progressive intrahepatic cholestasis-2, Dublin-Johnson syndrome, Pseudoxanthoma elasticum, cystic fibrosis, familial persistent hyperinsulinemic hypoglycemia of infancy, adrenoleukodystrophy, sitosterolemia, chronic obstructive pulmonary disease, asthma, disseminated bronchiectasis, chronic pancreatitis, male infertility, emphysema, or pneumonia.

[00301] According to another more preferred embodiment, the ABC transporter mediated disease is secretory diarrhea, or polycystic kidney disease in a mammal.

[00302] According to an alternative preferred embodiment, the present invention provides a method of treating cystic fibrosis or secretory diarrhea comprising the step of administering to said mammal a composition

comprising the step of administering to said mammal a composition comprising a compound of the present invention, or a preferred embodiment thereof as set forth above. Most preferably, said disease is cystic fibrosis.

[00303] According to another embodiment, the present invention provides a method of modulating activity of an anion channel *in vitro* or *in vivo*, comprising the step of contacting said channel with a compound of the present invention. Preferably, said anion channel is a chloride channel or a bicarbonate channel. More preferably, said anion channel is a chloride channel.

[00304] According to yet another embodiment, the present invention provides a method of treating an anion channel mediated disease in a mammal, comprising the step of administering to said mammal a composition comprising a compound according to the present invention.

[00305] According to another embodiment, the present invention provides a pharmaceutical composition comprising:

- (i) a compound of the present invention as described above;
- (ii) a pharmaceutically acceptable carrier; and
- (iii) an additional agent selected from a mucolytic agent, bronchodilator, an antibiotic, an anti-infective agent, an anti-inflammatory agent, a CFTRmodulator, or a nutritional agent.

[00306] Preferred embodiments of compounds the present invention in the above pharmaceutical composition are those as described above.

[00307] According to another embodiment, the present invention provides a kit for use in measuring the activity of a ABC transporter or a fragment thereof in a biological sample *in vitro* or *in vivo*, comprising:

(i) a composition comprising a compound of the present invention; and

(ii) instructions for:

a) contacting the composition with the biological sample;

b) measuring activity of said ABC transporter or a fragment thereof.

[00308] According to a preferred embodiment, the kit is useful in measuring the activity of CFTR.

[00309] According to another preferred embodiment, the activity of the ABC transporter is measured by measuring the transmembrane voltage potential.

[00310] Means for measuring the voltage potential across a membrane in the biological sample may employ any of the known methods in the art, such as optical membrane potential assay or other electrophysiological methods.

[00311] The optical membrane potential assay utilized voltage-sensitive FRET sensors described by Gonzalez and Tsien (See, Gonzalez, J. E. and R. Y. Tsien (1995) "Voltage sensing by fluorescence resonance energy transfer in single cells" Biophys J **69**(4): 1272-80, and Gonzalez, J. E. and R. Y. Tsien (1997) "Improved indicators of cell membrane potential that use fluorescence resonance energy transfer" Chem Biol **4**(4): 269-77) in combination with instrumentation for measuring fluorescence changes such as the Voltage/Ion Probe Reader (VIPR) (See, Gonzalez, J. E., K. Oades, et al. (1999) "Cell-based assays and instrumentation for screening ion-channel targets" Drug Discov Today **4**(9): 431-439).


[00312] These voltage sensitive assays are based on the change in fluorescence resonant energy transfer (FRET) between the membrane-soluble, voltage-sensitive dye, DiSBAC₂(3), and a fluorescent phospholipid, CC2-DMPE, which is attached to the outer leaflet of the plasma

membrane and acts as a FRET donor. Changes in membrane potential (V_m) cause the negatively charged DiSBAC₂(3) to redistribute across the plasma membrane and the amount of energy transfer from CC2-DMPE changes accordingly. The changes in fluorescence emission can be monitored using VIPRTM II, which is an integrated liquid handler and fluorescent detector designed to conduct cell-based screens in 96- or 384-well microtiter plates.

[00313] Preferred ABC transporters in the kit of the present invention include CFTR.


[00314] In order that the invention described herein may be more fully understood, the following examples are set forth. It should be understood that these examples are for illustrative purposes only and are not to be construed as limiting this invention in any manner.

EXAMPLE 1




General Procedure: The appropriate bromoketone was dissolved in a minimum of ethanol or methanol with 1.1 equivalents of thiourea. The reaction mixture was stirred overnight at room temperature, evaporated to dryness, and then dissolved in either dichloromethane or ethyl acetate. The reaction mixture was then extracted with 1 M sodium hydroxide followed by a saturated aqueous solution of sodium chloride. The organic layer was then separated, dried over sodium sulfate, and evaporated to dryness to yield the desired 2-aminothiazole.

Example 2



General Procedure: One equivalent of the appropriate carboxylic acid and one equivalent of the appropriate amine were dissolved in *N,N*-dimethylformamide (DMF) containing triethylamine (3 equivalents). *O*-(7-Azabenzotriazol-1-yl)-*N,N,N',N'*-tetramethyluronium hexafluorophosphate (HATU) was added and the solution was allowed to stir. The crude product was purified by reverse-phase preparative liquid chromatography to yield the pure product.


Example 3



General Procedure: The appropriate acid chloride was added to one equivalent of the appropriate amine in minimum of pyridine. The reaction mixture was allowed to stir overnight at room temperature. The reaction mixture was then filtered and evaporated to dryness. The crude product was purified by reverse-phase preparative liquid chromatography. Alternatively, the acid chloride was added to one equivalent of the appropriate amine in minimum of 1,4-dioxane containing an excess of triethylamine. The reaction mixture was then either allowed to stir overnight at room temperature or subjected to microwave irradiation for 5 minutes at 200 °C. The crude product was then filtered, evaporated to

dryness, dissolved in a minimum of dimethylsulfoxide and then purified by reverse-phase preparative liquid chromatography.

Example 4




Hal = Cl, Br, I

wherein W is a group as described in the compounds of the present invention.

General Procedure: One equivalent of the halide was dissolved in a minimum of the alcohol. The reaction vessel was sealed and then subjected to microwave irradiation for 15 minutes at 125 °C. The crude mixture was evaporated to dryness, dissolved in 1 mL of dimethylsulfoxide and purified by reverse-phase preparative liquid chromatography.

Example 5




Hal = Cl, Br, I

wherein W is a group as described in the compounds of the present invention.

General Procedure: One equivalent of the halide was dissolved in a minimum of *N,N*-dimethylformamide (DMF) containing 20 equivalents of amine. The reaction vessel was sealed and then subjected to microwave irradiation for 5 minutes at 80 °C. The crude mixture was evaporated to dryness and purified by reverse-phase preparative liquid chromatography.

1)



a) 4-Fluoro-1-methoxy-2-nitro-benzene A solution of methyl iodide (127.8 g, 0.9004 mol) in acetonitrile (100 mL) was slowly added to a solution of 5-fluoro-2-nitrophenol (94.2 g, 0.600 mol) and potassium carbonate (207 g, 1.50 mol) in acetonitrile (450 mL). The mixture was heated to reflux for 15 hours. The mixture was allowed to cool, filtered, and washed twice with dichloromethane (100 mL). The combined filtrate was evaporated to dryness to give the desired product (96 g, 0.56 mol, 93 %), which was used directly in the next step.

b) 5-Fluoro-2-methoxy-phenylamine A solution of 4-fluoro-1-methoxy-2-nitro-benzene (85 g, 0.50 mol) in methanol (300 mL) containing palladium on carbon (10 %, 8 g) was stirred for 15 hours under an atmosphere of hydrogen. The catalyst was filtered and the filtrate was evaporated to dryness to give the crude product (61.5 g, 0.436 mol, 87 %), which was used directly in the next step.

c) 2-Chloro-3-(5-fluoro-2-methoxy-phenyl)-propionaldehyde A solution of sodium nitrite (36 g, 0.51


mole) in water (50 mL) was slowly added to a solution of 5-fluoro-2-methoxy-phenylamine (61.5 g, 0.48 mol) in hydrochloric acid (20 % aqueous solution, 115 mL) at 0 °C. After stirring for 10 minutes, a cooled (0 °C) solution of acrolein (50 mL, 0.75 mol) in acetone (100 mL) containing calcium oxide (0.56 g, 0.010 mol) was added slowly to the reaction mixture. This was then followed by a solution of cuprous chloride (5 g, 0.05 mol) in acetone (100 mL) containing hydrochloric acid (20 % aqueous solution, 10 mL). The mixture was stirred at 0 to 30 °C for 3 hours, and then extracted three times with dichloromethane (300 mL). The combined organic layers were washed with a saturated aqueous solution of sodium bicarbonate, a saturated aqueous solution of sodium chloride, dried over sodium sulfate, filtered, and evaporated to dryness to give a black viscous oil. The crude product was passed through a short silica column to give the crude product, which was used directly in the next step.

d) 5-(5-Fluoro-2-methoxy-benzyl)-thiazol-2-ylamine

A mixture of 2-chloro-3-(5-fluoro-2-methoxy-phenyl)-propionaldehyde (crude from above) and thiourea (29.2 g, 0.384 mol) in ethanol (400 mL) was heated to reflux for 15 hours. The solvent was removed and the residue was diluted with dichloromethane (300 mL), sodium hydroxide (10 % aqueous solution, 150 mL) and water (200 mL). The aqueous phase was extracted twice with dichloromethane (150 mL). The combined organic layer was washed with water, a saturated aqueous solution of sodium chloride, dried over sodium sulfate, and evaporated to dryness. The crude product was recrystallized from a mixture of ethyl acetate and hexanes to give the pure product (6.5 g, 0.027 mol, 6.2 % from 5-fluoro-2-methoxy-phenylamine). ESI-MS m/z calc. 238.1, found 239.2 ($M+1$)⁺ ¹H NMR(CDCl₃):

δ 6.90-6.81 (m, 2 H), 6.79-6.76 (m, 2 H), 4.75 (br, 2 H), 3.91 (s, 2 H), 3.82 (s, 3 H).

2)



a) 2-Methoxy-3-nitro-pyridine A suspension of sodium methoxide (40.5 g, 0.750 mol) in 200 mL of methanol was slowly added to a solution of 2-chloro-3-nitro-pyridine (79.3 g, 0.500 mol) in 800 mL of methanol at 0 °C. The reaction mixture was stirred for 4 hours and then poured into 1000 g of ice. The resulting precipitate was filtered, washed with water, and dried to give 2-methoxy-3-nitro-pyridine (70. g, 0.45 mmol, 90 %) as a white solid.


b) 2-Methoxy-pyridin-3-ylamine A solution of 2-methoxy-3-nitro-pyridine (70. g, 0.45 mol) in methanol (700 mL) containing palladium on carbon (7 g, 10 %) was stirred under an atmosphere of hydrogen for 15 hours. The catalyst was filtered and washed with methanol. The filtrate was evaporated to dryness to give crude 2-methoxy-pyridin-3-ylamine (48 g, 0.39 mol, 87 %), which was used directly in the next step.

c) 2-Chloro-3-pyridin-3-yl-propionaldehyde A solution of sodium nitrite (20. g, 0.28 mol) in water (100 mL) was slowly added to a solution of 2-methoxy-pyridin-3-ylamine (36 g, 0.25 mol) in hydrochloric acid (20 % aqueous solution, 60 mL) at 0 °C. After stirring

for 10 minutes, a cooled (0 °C) solution of acrolein (25 mL, 0.37 mol) in acetone (25 mL) containing calcium oxide (5 g, 0.09 mol) was slowly added to the reaction mixture. This was then followed by a solution of cuprous chloride (2.5 g, 0.025 mol) in acetone (25 mL) containing hydrochloric acid (20 % aqueous solution, 5 mL). The mixture was stirred at 0 to 30 °C for 3 hours, and then extracted three times with dichloromethane (150mL). The combined organic layers were washed with a saturated aqueous solution of sodium bicarbonate, a saturated aqueous solution of sodium chloride, dried over sodium sulfate, filtered, and evaporated to dryness to give a black viscous oil. The crude product was passed through a short silica column to give a crude product, which was used directly in the next step.

d) 5-(2-Methoxy-pyridin-3-ylmethyl)-thiazol-2-ylamine A mixture of 2-chloro-3-pyridin-3-yl-propionaldehyde (crude from above) and thiourea (14.8 g, 0.194 mol) in ethanol (200 mL) was heated to reflux overnight. The solvent was removed and the residue was diluted with dichloromethane (1.2 L) and then washed with sodium hydroxide (10 % aqueous solution, 400 mL) and water (200 mL). The organic layer was extracted three times with hydrochloric acid (5 % aqueous solution, 400 mL) and the combined aqueous layer was brought to between pH 9 and 10 with sodium hydroxide (10 % aqueous solution). The resulting precipitate was filtered to give the crude product, which was recrystallized, from ethyl acetate and hexanes to give the pure product (5.1 g, 0.023 mol, 5.6 % from 2-methoxy-pyridin-3-ylamine). ESI-MS m/z calc. 221.1, found 222.2 (M+1)⁺.

3)




a) **2-Bromo-1-(chloro-phenyl)-ethanone** Bromine (3.8 mL, 65 mmol) was added dropwise to a solution of 1-(2-chloro-phenyl)-ethanone (10. g, 65 mmol) in acetic acid (75 mL) at 0°C. The mixture was then warmed to room temperature and stirred overnight. The mixture was evaporated to dryness and used in the next step without further purification.

N'-[5-(2-Chloro-benzoyl)-thiazol-2-yl]-*N,N*-dimethyl-formamidine. A mixture of thiourea (4.95 g, 65.0 mmol) and dimethoxymethyl-dimethyl-amine (23.2 g, 195 mmol) in methanol (80 mL) was heated under reflux for 30 minutes. After allowing the mixture to cool, triethylamine (19.8 g, 195 mmol) and a solution of 2-bromo-1-(chloro-phenyl)-ethanone (crude from last step) in methanol (50 mL) were added. The mixture was heated to reflux for 4 hours. The solvent was removed and the residue was used directly in the next procedure.

b) **(2-Amino-thiazol-5-yl)-(2-chloro-phenyl)-methanone** The crude *N'*-[5-(2-chloro-benzoyl)-thiazol-2-yl]-*N,N*-dimethyl-formamidine was dissolved in 10% aqueous hydrochloric acid (150 mL) and heated to 70°C for 4 hours. The precipitate was filtered, washed with ether, and then suspended in a 10% aqueous sodium carbonate solution (250mL). The suspension was stirred for 1 hour and the precipitate was filtered, washed with ether, and dried in air to give (2-amino-thiazol-5-yl)-(2-chloro-phenyl)-methanone as a brown solid (8.5 g, 36 mmol, 55% from 1-(2-chloro-phenyl)-ethanone). ESI-MS *m/z* calc.

238.0, found 239.3 (M+1)⁺ ¹H NMR (DMSO): δ : 7.252 (s, 1 H), 7.420-7.553 (m, 4 H), 8.345 (s, 2 H).

4)




a) 2-Chloro-3-(2-chloro-phenyl)-propionaldehyde To a solution of 2-chloroaniline (12.7 g, 100. mmol) in hydrochloric acid (20 % aqueous solution, 40 mL) was added dropwise a solution of sodium nitrite (7.5 g, 110 mmol) in water (20 mL) at 0 to 5 °C. After stirring for 10 minutes, a cooled (0 °C) solution of acrolein (15 g, 270 mmol) in acetone (100 mL) containing calcium oxide (2.0 g, 36 mmol) was added gradually, and then followed by a solution of cuprous chloride (1 g, 10 mmol) in acetone (10 mL) containing hydrochloric acid (20 % aqueous solution, 2 mL). The mixture was stirred at 0 to 30 °C for 3 hours and then extracted three times with dichloromethane (100mL). The combined organic layers were washed with a saturated aqueous solution of sodium bicarbonate followed by a saturated aqueous solution of sodium chloride. The organic layer was separated, dried over sodium sulfate, and evaporated to dryness to give a black viscous oil. The crude product was passed through a short silica gel column to give 12 g of crude product, which was used directly in the next step.

b) 5-(2-Chloro-benzyl)-thiazol-2-ylamine A mixture of 2-chloro-3-(2-chloro-phenyl)-propionaldehyde (12 g, crude from above) and urea (6.0 g, 0.10 mol) in ethanol (120 mL) was heated to reflux overnight. The solvent was evaporated to dryness. The residue was diluted with


dichloromethane (120 mL) and then washed with sodium hydroxide (10% aqueous solution, 50 mL) and water (30 mL). The organic layer was extracted three times with hydrochloric acid (5% aqueous solution, 120 mL). The combined aqueous layer was adjusted with a 10 % aqueous solution of sodium hydroxide to between pH 9 and 10 and then extracted three times with dichloromethane (150 mL). The organic layers were combined, dried over sodium sulfate, evaporated to dryness, and purified by silica gel column chromatography to yield a yellow solid. (5.2 g, 0.023 mol, 23% from 2-chloroaniline). ESI-MS m/z calc. 224.0, found 225.2 $(M+1)^+$ ^1H NMR ($CDCl_3$) δ 4.07 (s, 2H), 4.90 (bs, 2H), 6.80 (s, 1H), 7.37-7.15 (m, 4H).

6) 5-(2-methoxy-benzyl)-thiazol-2-ylamine



5-(2-methoxy-benzyl)-thiazol-2-ylamine was prepared in a manner analogous to that of 5-(2-chloro-benzyl)-thiazol-2-ylamine. ESI-MS m/z calc. 220.1, found 221.2 $(M+1)^+$ ^1H NMR ($CDCl_3$) δ 7.26-7.19 (m, 1H), 7.15 (d, J = 6.8 Hz, 1H), 6.90-6.85 (m, 2H), 6.79 (s, 1H), 4.77 (bs, 2H), 3.93 (s, 2H), 3.84 (s, 3H).


7) 5-(3-Chloro-benzyl)-thiazol-2-ylamine



5-(3-Chloro-benzyl)-thiazol-2-ylamine was prepared in a manner analogous to that of 5-(2-chloro-benzyl)-thiazol-2-ylamine. ESI-MS m/z calc. 224.0, found 225.2


(M+1)⁺ ¹H NMR (CDCl₃) δ 7.26-7.21 (m, 3H), 7.10 (d, *J* = 6.8 Hz, 1H), 6.81 (s, 1H), 4.82 (bs, 2H), 3.93 (s, 2H).

8) 5-(4-Chloro-benzyl)-thiazol-2-ylamine




5-(4-Chloro-benzyl)-thiazol-2-ylamine was prepared in a manner analogous to that of 5-(2-chloro-benzyl)-thiazol-2-ylamine. ESI-MS *m/z* calc. 224.0, found 225.2 (M+1)⁺ ¹H NMR (CDCl₃) δ 7.26 (d, *J* = 8.4 Hz, 2H), 7.14 (d, *J* = 8.4 Hz, 2H), 6.79 (s, 1H), 4.85 (bs, 2H), 3.92 (s, 2H).

9) 5-(2-Cyano-benzyl)-thiazol-2-ylamine



5-(2-Cyano-benzyl)-thiazol-2-ylamine was prepared in a manner analogous to that of 5-(2-chloro-benzyl)-thiazol-2-ylamine (12 g, 56 mmol, 11 % from 2-cyanoaniline). ESI-MS *m/z* calc. 215.05, found 216.16 (M+1)⁺ ¹H NMR (CDCl₃): δ 7.64 (d, 1H), 7.54 (t, 1H), 7.34 (m, 2H), 6.87 (s, 1H), 4.89 (br, 2H), 4.19 (s, 2H).

10)




a) 2-Chloro-3-(2-methoxy-phenyl)-propionaldehyde. A solution of 2-methoxyaniline (24.6 g, 0.200 mol) in

hydrochloric acid (20 % aqueous solution, 80 mL) was slowly added to a solution of sodium nitrite (15 g, 0.22 mol) in water (40 mL) at 0 to 5 °C. After stirring for 10 minutes, a cooled (0 °C) solution of acrolein (30 g, 0.56 mol) in acetone (200 mL) containing calcium oxide (4.0 g, 72 mmol) was slowly added, followed by a solution of cuprous chloride (2.0 g, 20 mmol) in acetone (20 mL) containing hydrochloric acid (20 % aqueous solution, 4 mL). The mixture was stirred at 0 to 30 °C for 3 hours, and then extracted with three 150 mL portions of dichloromethane. The combined organic layers were washed with a saturated aqueous solution of sodium bicarbonate, a saturated aqueous solution of sodium chloride, dried over sodium sulfate, filtered, and concentrated to give a black viscous oil. The crude product was passed through a short silica column to give 10 g of crude product, which was used directly in the next procedure.

b) 5-(2-methoxy-benzyl)-oxazol-2-ylamine. A mixture of 2-chloro-3-(2-methoxyphenyl)-propionaldehyde (10 g, crude from above) and urea (9.6 g, 0.16 mol) was dissolved in ethanol (250 mL) and then heated to reflux overnight. The solvent was evaporated to dryness. The residue was diluted with dichloromethane (250 mL) and then washed with sodium hydroxide (10 % aqueous solution, 100 mL) and water (50 mL). The organic layer was extracted three times with hydrochloric acid (5 % aqueous solution, 250 mL). The combined aqueous layers were adjusted to pH 9 to 10 with a 10 % aqueous solution of sodium hydroxide and then extracted three times with dichloromethane (300mL). The organic layer was separated, dried over sodium sulfate, and evaporated to dryness. The crude product was purified by silica gel column chromatography to yield the yellow-red solid product.


(0.72 g, 0.35 % from 2-methoxyaniline). ESI-MS m/z calc. 204.1, found 205.1 $(M+1)^+$ ^1H NMR(CDCl_3) δ 7.26-7.20 (m, 1H), 7.14 (d, $J = 7.2\text{Hz}$, 1H), 6.91-6.86 (m, 2H), 6.35 (s, 1H), 4.49 (bs, 2H), 3.85 (s, 2H), 3.82 (s, 3H).

11)




5-(2-Chloro-benzyl)-oxazol-2-ylamine 5-(2-Chloro-benzyl)-oxazol-2-ylamine was prepared in a manner analogous to that of the preparation of 5-(2-methoxy-benzyl)-oxazol-2-ylamine to yield the product as a yellow solid. (3.5 g, 8.4% from 2-chloroaniline). ESI-MS m/z calc. 208.0, found 209.1 $(M+1)^+$ ^1H NMR (CDCl_3) δ 7.37-7.18 (m, 4H), 6.40 (s, 1H), 4.66 (bs, 2H), 3.97 (s, 2H).

12)




5-(3-Chloro-benzyl)-oxazol-2-ylamine 5-(3-Chloro-benzyl)-oxazol-2-ylamine was prepared in a manner analogous to that of the preparation of 5-(2-methoxy-benzyl)-oxazol-2-ylamine to yield the product as a yellow solid (1.2 g, 2.9 % from 3-chloroaniline). ESI-MS m/z calc. 208.0, found 209.2 ^1H NMR(CDCl_3) δ 7.26-7.22 (m, 3H), 7.10 (d, $J = 6.0\text{Hz}$, 1H), 6.44 (s, 1H), 4.73 (bs, 2H), 3.82 (s, 2H).

13)



5-(4-Chloro-benzyl)-oxazol-2-ylamine 5-(4-Chloro-benzyl)-oxazol-2-ylamine was prepared in a manner analogous to that of the preparation of 5-(2-methoxy-benzyl)-oxazol-2-ylamine to yield the product as a yellow solid (1.6 g, yield 3.86 % from 4-chloroaniline). ESI-MS m/z calc. 208.0, found 209.1 ^1H NMR(CDCl_3) δ 7.27 (d, J = 8.4Hz, 2H), 7.17 (d, J = 8.0 Hz, 2H), 6.38 (s, 1H), 4.66 (bs, 2H), 3.81(s, 2H).

14)




a) 2-Bromo-3-phenylpropionaldehyde A solution of bromine (15.2 g, 95.1 mmol) in 30 mL of dichloromethane was added to a solution of 3-phenyl-propionaldehyde (13.4 g, 100 mmol) in dichloromethane (150 mL) at 0 °C over 20 minutes. The reaction mixture was allowed to stir for 2 hours and then a saturated aqueous solution of sodium bicarbonate (100 mL) was added to the mixture. The organic layer was separated and the aqueous layer was washed with dichloromethane (50 mL). The combined organic layers were washed with water, a saturated aqueous solution of sodium chloride, and then evaporated to dryness to give an orange oil (14.2 g), which was used directly in the next step.

b) 5-Benzyl-oxazol-2-ylamine A mixture of 2-bromo-3-phenylpropionaldehyde (14.2 g, crude from above) and urea (7.2 g, 0.12 mol) were heated to reflux for 15 hours in 200 mL of ethanol. The solvent was evaporated to dryness

and the residue was diluted with dichloromethane (250 mL) and then washed with sodium hydroxide (10 % aqueous solution, 100 mL) and water (50 mL). The organic layer was extracted three times with hydrochloric acid (5 % aqueous solution, 250mL). The combined aqueous layers were adjusted to between pH 9 to 10 with a 10 % aqueous solution of sodium hydroxide and then extracted three times with dichloromethane (300mL). The organic layer was dried over sodium sulfate, evaporated to dryness, and purified by silica gel column chromatography to give a pale yellow solid. (1.6 g, 9.2 mmol, 9.2% from 3-phenyl-propionaldehyde). ESI-MS m/z calc. 174.1, found 175.1 ^1H NMR(CDCl_3) δ 7.32-7.22 (m, 5H), 6.39 (s, 1H), 4.72 (bs, 2H), 3.84 (s, 2H).


15)



a) 2-Bromo-3-phenylpropionaldehyde A solution of bromine (15.2 g, 95.1 mmol) in 30 mL of dichloromethane was added to a solution of 3-phenyl-propionaldehyde (13.4 g, 100 mmol) in dichloromethane (150 mL) at 0 °C over 20 minutes. The reaction mixture was allowed to stir for 2 hours and then a saturated aqueous solution of sodium bicarbonate (100 mL) was added to the mixture. The organic layer was separated and the aqueous layer was washed with dichloromethane (50 mL). The combined organic layers were washed with water, a saturated aqueous solution of sodium chloride, and then evaporated to dryness to give an orange oil (14.2 g), which was used directly in the next step.

b) 5-benzyl-thiazol-2-ylamine A mixture of 2-bromo-3-phenylpropionaldehyde (14.2 g, crude from above) and urea (7.2 g, 0.12 mol) were heated to reflux for 15 hours in 200 mL of ethanol. The solvent was evaporated to dryness and the residue was diluted with dichloromethane (250 mL) and then washed with sodium hydroxide (10 % aqueous solution, 100 mL) and water (50 mL). The organic layer was extracted three times with hydrochloric acid (5 % aqueous solution, 250 mL). The combined aqueous layers were brought to pH 9 to 10 with a 10 % aqueous solution of sodium hydroxide and then extracted three times with dichloromethane (300 mL). The organic layer was dried over sodium sulfate, evaporated to dryness, and purified by silica gel column chromatography to give a pale yellow solid. (5.2 g, 27 mmol, 27 % from 3-phenylpropionaldehyde). ESI-MS m/z calc. 190.1, found 191.2 ^1H NMR (CDCl_3) δ 7.32-7.21 (m, 5H), 6.79 (s, 1H), 4.91 (bs, 2H), 3.95 (s, 2H).

16)



a) 2-Chloro-3-(2-chloro-phenyl)-propionaldehyde. A solution of sodium nitrite (15 g, 0.22 mol) in water (40 mL) was slowly added to a solution of 2-chloroaniline (25.5 g, 0.200 mol) in hydrochloric acid (20 % aqueous solution, 100 mL) at 0 to 5 °C. The mixture was stirred for ten minutes and then poured into a cooled (0 °C) solution of acrolein (30. g, 0.56 mol) in acetone (200 mL) containing calcium oxide (4.0 g, 72 mmol), followed by a solution of cuprous chloride (2.0 g, 20 mmol) in acetone (20 mL) containing hydrochloric acid (20 %


aqueous solution, 4 mL). The mixture was stirred for 3 hours at room temperature, and then extracted three times with dichloromethane (150 mL). The combined organic layers were washed with a solution of saturated aqueous sodium chloride, dried over sodium sulfate, filtered, and evaporated to dryness to give a black viscous oil that was used directly in the next procedure.

b) [2-(2-Chloro-phenyl)-1-methyliminomethyl-ethyl]-methyl-amine. A solution of methylamine in methanol (27 %, 69 g) was slowly added to a solution of 2-chloro-3-(2-chloro-phenyl)-propionaldehyde in dichloromethane (20 mL). The reaction mixture was allowed to stir for 12 hours and then used immediately in the next procedure.

c) 5-(2-Chloro-benzyl)-1-methyl-1H-imidazol-2-ylamine. A solution of cyanamide in water (50 %, 150 mL) was added to a boiling solution of [2-(2-chloro-phenyl)-1-methyliminomethyl-ethyl]-methyl-amine in methanol and dichloromethane. The pH was brought to 4.5 by the continual addition of an aqueous solution of sulfuric acid (9 M). The mixture was refluxed for 2 hours, allowed to cool to room temperature, and adjusted to pH 9 through the addition of powdered sodium bicarbonate. The mixture was extracted three times with dichloromethane (200 mL) and the combined organic layers were extracted three times with hydrochloric acid (20 % aqueous solution, 150 mL). The aqueous solution was adjusted to pH 10 with sodium hydroxide (10 % aqueous solution) and extracted three times with dichloromethane (150 mL). The combined organic layers were washed with a saturated aqueous solution of sodium chloride, dried over sodium sulfate, filtered, and evaporated to dryness to give a black solid which was purified by column chromatography to yield the

product (5.0 g, 0.23 mmol, 11 % from 2-chloroaniline) as a brown solid. ESI-MS m/z calc. 221.1, found 222.3 ($M+1$)⁺ ¹H NMR (CDCl₃): δ 7.30-7.37 (m, 1 H), 7.15-7.18 (m, 2 H), 7.03-7.06 (m, 1 H), 6.43 (s, 1 H), 3.94 (s, 2H), 3.80 (br, 2H), 3.15 (s, 3 H).

17)




a) 2-Bromo-3-oxo-hexanoic acid ethyl ester 3-Oxo-hexanoic acid ethyl ester (4.0 mL, 25 mmol) and magnesium perchlorate (1.7 g, 7.6 mmol) were placed in 500 mL of ethyl acetate and allowed to stir for 5 minutes. *N*-Bromosuccinimide (4.7 g, 26 mmol) was added and the reaction mixture was allowed to stir for 15 minutes, at which time thin-layer chromatography (10 % ethyl acetate in hexanes, silica gel, 254 nm irradiation) indicated the reaction was complete. The reaction mixture was diluted with 500 mL of ethyl ether and washed three times with an equal volume of saturated aqueous sodium chloride. The organic layer was then dried over sodium sulfate and evaporated to dryness. This material was used in the next step without further purification.

b) Amino-4-propyl-thiazole-5-carboxylic acid ethyl ester 2-Bromo-3-oxo-hexanoic acid ethyl ester (5.9 g, 25 mmol), was dissolved in 60 mL of ethanol containing triethylamine (4.2 mL, 30 mmol) and thiourea (1.9 g, 25


mmol). The colorless solution was protected from light and allowed to stir for 16 hours. The resulting red suspension was evaporated to dryness and dissolved in a minimum of dichloromethane. This solution was washed three times with an equal volume of a saturated aqueous solution of sodium bicarbonate, followed by a saturated aqueous solution of sodium chloride. The organic layer was separated and filtered to remove a fine red precipitate which remained suspended in the organic phase. The solvent was removed and then the solid was dissolved in a minimum of 50 / 50 (v / v) ethyl acetate and 1 N aqueous solution of hydrochloric acid. The layers were separated and the aqueous layer was washed with an equal volume of ethyl acetate. After discarding the organic layers, the aqueous layer was then placed in an ice bath with an equal volume of ethyl acetate. Sodium hydroxide (1N) was then slowly added with vigorous swirling until the aqueous phase was basic. The layers were separated and the aqueous layer was washed two additional times with ethyl acetate. The combined organic layers were washed three times with an equal volume of a solution of saturated aqueous sodium bicarbonate followed by a solution of saturated aqueous sodium chloride. The organic layer was then dried over sodium sulfate and evaporated to dryness to yield a pale yellow solid (1.8 g, 8.4 mmol, 34 %) ESI-MS m/z calc. 214.1, found 215.3 $(M+1)^+$ Retention time 1.90 minutes.

18) 4-(2-Methoxy-phenyl)-thiazol-2-ylamine.



2-Bromo-1-(2-methoxy-phenyl)-ethanone (0.6388 g, 2.789 mmol) and thiourea (0.2289 g, 3.007 mmol) were dissolved in a 20 mL of ethanol. The reaction mixture was allowed to stir overnight at room temperature. The ethanol was evaporated to dryness and the crude product was dissolved in a minimum of dichloromethane. The crude product was then extracted twice with 1M sodium hydroxide and once with a saturated aqueous solution of sodium chloride. The organic layer was then dried over sodium sulfate, filtered, and evaporated to dryness to yield the pure product (0.529 g, 2.56 mmol, 92.0 %). ESI-MS m/z calc. 206.3, found 207.1 ($M+1$)⁺ Retention time 1.86 minutes. ¹H NMR (400 MHz, CD₃CN) δ 3.91 (s, 3H), 5.54 (s, 2H), 6.97-7.02 (m, 1H), 7.03-7.06 (m, 1H), 7.23 (s, 1H), 7.24-7.29 (m, 1H), 8.06 (dd, J = 7.8, 1.8 Hz, 1H).


19) 4-(3-Methoxy-phenyl)-thiazol-2-ylamine.



2-Bromo-1-(3-methoxy-phenyl)-ethanone (0.7291 g, 3.183 mmol) and thiourea (0.2665 g, 3.501 mmol) were dissolved in a 20 mL of ethanol. The reaction mixture was allowed to stir overnight at room temperature. The ethanol was evaporated to dryness and the crude product was dissolved in a minimum of dichloromethane. The crude product was then extracted twice with 1M sodium hydroxide and once with a saturated aqueous solution of sodium chloride. The organic layer was then dried over sodium sulfate, filtered, and evaporated to dryness to yield the product (0.619 g, 3.00 mmol, 94.3 %). ESI-MS m/z calc. 206.3, found 207.0 ($M+1$)⁺ Retention time 1.86 minutes. ¹H


NMR (400 MHz, CD₃CN) δ 3.84 (s, 3H), 5.67 (s, 2H), 6.85-6.91 (m, 2H), 7.31 (t, J = 7.9 Hz, 1H), 7.36-7.43 (m, 2H).

20) 4-Phenyl-thiazol-2-ylamine



2-Bromo-1-phenyl-ethanone (19.9 g, 0.100 mol) and thiourea (7.9 g, 0.10 mol) were mixed in a 150 mL of methanol. The reaction mixture was warmed to dissolve the reagents and then allowed to stir overnight at room temperature. The methanol was evaporated to dryness and the crude product was dissolved in a minimum of ethyl acetate. The crude product was then extracted twice with 1M sodium hydroxide and once with a saturated aqueous solution of sodium chloride. The organic layer was then dried over sodium sulfate, filtered, and evaporated to dryness to yield the pure product (16.8 g, 0.0953 mol, 95.3 %) of product. ESI-MS m/z calc. 176.0, found 177.2 ($M+1$)⁺ Retention time 1.41 minutes. ¹H NMR (400 MHz, CD₃CN) δ 5.73 (s, 2H), 6.87 (s, 1H), 7.28-7.34 (m, 1H), 7.36-7.43 (m, 2H), 7.79-7.85 (m, 2H).

21)




a) 2-Pyridin-3-yl-butylbutyric acid ethyl ester Pyridin-3-yl-acetic acid ethyl ester (22.5g, 0.136 mol) in tetrahydrofuran (50 mL) was slowly added to a suspension

of sodium hydride (60% in mineral oil, 6 g, 0.4 mol) in tetrahydrofuran (50 mL) at 0 °C. A solution of ethyl iodide (23.4g, 0.150 mol) in tetrahydrofuran (50 mL) was added to and the reaction mixture was allowed to stir overnight at room temperature. The mixture was poured into ice and the aqueous layer was extracted three times with ethyl acetate. The organic layer was washed with a saturated aqueous solution of sodium chloride, dried over sodium sulfate, and evaporated to dryness. The crude residue was purified by preparative reverse phase liquid chromatography (13.5g, 0.0699 mol, 51.3 %). ESI-MS m/z calc. 193.1, found 194.2 $(M+1)^+$ ^1H NMR ($CDCl_3$) δ : 9.42-9.39 (m, 2 H), 8.58-8.55 (m, 1 H), 8.17-8.13 (m, 1 H), 5.09-4.97 (m, 2 H), 4.35 (t, 1 H, $J = 7.6$ Hz), 3.04-2.95 (m, 1 H), 2.74-2.65 (m, 1 H), 2.10 (t, 3 H, $J=7.2$ Hz), 1.79 (t, 3 H, $J = 7.6$ Hz).


b) 2-Pyridin-3-yl-butyric acid A mixture of 2-Pyridin-3-yl-butyric acid ethyl ester (8 g, 0.04 mol) and a 20 % aqueous solution of hydrochloric acid (50 mL) was heated to reflux for 3 hours. The solvent evaporated to dryness to yield the desired product (5 g, 0.02 mol, 50 %), ESI-MS m/z calc. 165.1, found 166.5 $(M+1)^+$ ^1H NMR ($DMSO-d_6$): δ 8.86 (s, 1 H), 8.80 (d, 1 H, $J = 5.6$ Hz), 8.47 (d, 1 H, $J = 8.0$ Hz), 7.97 (dd, 1 H, $J = 5.6, 8.0$ Hz), 3.83 (t, 1 H, $J = 8.0$ Hz), 2.07-2.04 (m, 1 H), 1.82-1.80 (m, 1 H), 0.80 (t, 3 H, $J = 7.6$ Hz).

22) N-{5-[(2-Chloro-phenyl)-hydroxy-methyl]-thiazol-2-yl}-2-phenyl-butyramide




N-[5-(2-Chloro-benzoyl)-thiazol-2-yl]-2-phenylbutyramide (102 mg, 0.265 mmol) was suspended in 1 mL of anhydrous methanol. Sodium borohydride (30.3 mg, 0.801 mmol) was slowly added and the resulting pale yellow solution was allowed to stir for 1 hour at room temperature. After stirring for one hour a second aliquot of sodium borohydride (30.3 mg, 0.801 mmol) was added. The reaction mixture was allowed to stir for an additional hour and then the crude product was evaporated to dryness and then dissolved in a minimum of ethyl acetate. The organic layer was washed three times with an equal volume of 1N hydrochloric acid, saturated aqueous sodium bicarbonate, and saturated aqueous sodium chloride. The organic layer was then dried over sodium sulfate, filtered, and evaporated to dryness. The crude product was further purified by reverse-phase preparative liquid chromatography to yield the pure product (46 mg, 0.12 mmol, 45 %) ESI-MS *m/z* calc. 386.1, found 387.3 (M+1)⁺ Retention time 3.83 minutes.

23) *N*-[5-(2-Chloro-benzyl)-thiazol-2-yl]-2,2-diphenylacetamide




5-(2-Chloro-benzyl)-thiazol-2-ylamine (45 mg, 0.20 mmol) and diphenyl-acetic acid (42 mg, 0.20 mmol) were dissolved in *N,N*-dimethylformamide (1 mL) containing triethylamine (84.1 μ L, 0.600 mmol). *O*-(7-Azabenzotriazol-1-yl)-*N,N,N',N'*-tetramethyluronium hexafluorophosphate (84 mg, 0.22 mmol) was added and the solution was allowed to stir for 16 hours. The crude product was purified by reverse-phase preparative liquid chromatography (10.3 mg, 0.0246 mmol, 12 %). ESI-MS *m/z* calc. 418.1, found 419.2 (*M*+1)⁺ Retention time 3.85 minutes. ¹H NMR (400 MHz, DMSO-*d*₆) δ 4.19 (s, 2H), 5.30 (s, 1H), 7.23-7.47 (m, 15H).

24) *N*-(7-Oxo-4,5,6,7-tetrahydro-benzothiazol-2-yl)-2-phenyl-butyramide




2-Amino-5,6-dihydro-4*H*-benzothiazol-7-one (34 mg, 0.20 mmol) and 2-phenyl-butyric acid (33 mg, 0.20 mmol) were dissolved in *N,N*-dimethylformamide (1 mL) containing triethylamine (84.1 μ L, 0.600 mmol). *O*-(7-Azabenzotriazol-1-yl)-*N,N,N',N'*-tetramethyluronium hexafluorophosphate (84 mg, 0.22 mmol) was added and the solution was allowed to stir for 16 hours. The crude product was purified by reverse-phase preparative liquid chromatography (31 mg, 0.099 mmol, 49 %). ESI-MS *m/z* calc. 314.1, found 315.3 (*M*+1)⁺ Retention time 2.90 minutes.

25) (2-Diphenylacetyl-amino-thiazol-4-yl)-acetic acid methyl ester



(2-Amino-thiazol-4-yl)-acetic acid methyl ester (45 mg, 0.20 mmol) and diphenyl-acetic acid (37 mg, 0.20 mmol) were dissolved in *N,N*-dimethylformamide (2 mL) containing triethylamine (84.1 μ L, 0.600 mmol). *O*-(7-Azabenzotriazol-1-yl)-*N,N,N',N'*-tetramethyluronium hexafluorophosphate (84 mg, 0.22 mmol) was added and the solution was allowed to stir for 16 hours. The crude product was purified by reverse-phase preparative liquid chromatography (29.4 mg, 0.0802 mmol, 40. %). ESI-MS *m/z* calc. 380.1, found 381.3 (*M*+1)⁺ Retention time 3.28 minutes.


26) 2-Diphenylacetyl-amino-thiazole-4-carboxylic acid ethyl ester



2-Amino-thiazole-4-carboxylic acid methyl ester (32 mg, 0.20 mmol) and diphenyl-acetic acid (42 mg, 0.20 mmol) were dissolved in *N,N*-dimethylformamide (1 mL) containing triethylamine (84.1 μ L, 0.600 mmol). *O*-(7-Azabenzotriazol-1-yl)-*N,N,N',N'*-tetramethyluronium hexafluorophosphate (84 mg, 0.22 mmol) was added and the solution was allowed to stir for 16 hours. The crude


product was purified by reverse-phase preparative liquid chromatography (19 mg, 0.052 mmol, 26 %). ESI-MS m/z calc. 366.1, found 367.1 ($M+1$)⁺ Retention time 3.34 minutes.

27) *N*-[5-(2-Methoxy-pyridin-3-ylmethyl)-thiazol-2-yl]-2-phenyl-butamide



5-(2-Methoxy-pyridin-3-ylmethyl)-thiazol-2-ylamine (44 mg, 0.20 mmol) and 2-phenyl-butyric acid (33 mg, 0.20 mmol) were dissolved in acetonitrile (1 mL) containing triethylamine (84.1 μ L, 0.600 mmol). O-(7-Azabenzotriazol-1-yl)-*N,N,N',N'*-tetramethyluronium hexafluorophosphate (84 mg, 0.22 mmol) was added and the solution was allowed to stir for 16 hours. The crude product was purified by reverse-phase preparative liquid chromatography (15 mg, 0.041 mmol, 21 %). ESI-MS m/z calc. 367.1, found 368.1 ($M+1$)⁺ Retention time 3.24 minutes.


28) 2-Bromo-*N*-[5-(2-chloro-benzyl)-thiazol-2-yl]-2-phenyl-acetamide



5-(2-Chloro-benzyl)-thiazol-2-ylamine (450 mg, 2.0 mmol) and bromo-phenyl-acetic acid (430 mg, 2.0 mmol) were dissolved in acetonitrile (20 mL) containing


triethylamine (280 μ L, 2.0 mmol). O-(7-Azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (836 mg, 2.2 mmol) was added and the solution was allowed to stir for 16 hours. The crude product was purified by silica gel chromatography using a gradient of 10-30 % ethyl acetate in hexanes to yield a pale yellow solid (633 mg, 1.50 mmol, 75.0 %). ESI-MS m/z calc. 420.0, found 421.2 (M+1)⁺ Retention time 3.63 minutes.

29) 2-Bromo-N-[5-(2-chloro-benzyl)-thiazol-2-yl]-2-phenyl-acetamide




5-(2-Chloro-benzyl)-thiazol-2-ylamine (0.484 g, 2.16 mmol) and 2-bromo-butyrac acid (0.360 g, 2.16 mmol) were dissolved in acetonitrile (20 mL) containing triethylamine (302 μ L, 2.16 mmol). O-(7-Azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (1.15 g, 3.02 mmol) was added and the solution was allowed to stir for 16 hours. The crude product was purified by silica gel chromatography. ESI-MS m/z calc. 372.0, found 373.2 (M+1)⁺ Retention time 3.41 minutes.

30) N-[5-(4-Chloro-benzyl)-oxazol-2-yl]-2-cyclopentyl-2-phenyl-acetamide




5-(4-Chloro-benzyl)-oxazol-2-ylamine (42 mg, 0.20 mmol) and cyclohexyl-phenyl-acetic acid (44 mg, 0.20 mmol) were dissolved in *N,N*-dimethylformamide (1 mL) containing triethylamine (84.1 μ L, 0.600 mmol). *O*-(7-Azabenzotriazol-1-yl)-*N,N,N',N'*-tetramethyluronium hexafluorophosphate (84 mg, 0.22 mmol) was added and the solution was allowed to stir for 16 hours. The crude product was purified by reverse-phase preparative liquid chromatography (9.6 mg, 0.023 mmol, 12 %). ESI-MS *m/z* calc. 408.9, found 409.4 (*M*+1)⁺ Retention time 3.76 minutes.

31) *N*-[5-(2-Chloro-benzyl)-1-methyl-1*H*-imidazol-2-yl]-2-phenyl-butyramide




5-(2-Chloro-benzyl)-1-methyl-1*H*-imidazol-2-ylamine (44 mg, 0.20 mmol) and 2-phenyl-butyric acid (33 mg, 0.20 mmol) were dissolved in *N,N*-dimethylformamide (1 mL) containing triethylamine (84.1 μ L, 0.600 mmol). *O*-(7-Azabenzotriazol-1-yl)-*N,N,N',N'*-tetramethyluronium hexafluorophosphate (84 mg, 0.22 mmol) was added and the solution was allowed to stir for 16 hours. The crude product was purified by reverse-phase preparative liquid chromatography (13 mg, 0.035 mmol, 18 %). ESI-MS *m/z* calc. 367.2, found 368.1 (*M*+1)⁺ Retention time 2.42 minutes.

32) *N*-(6-Ethoxy-benzothiazol-2-yl)-2-phenyl-butyramide




6-Ethoxy-benzothiazol-2-ylamine (39 mg, 0.20 mmol) and 2-phenyl-butyrlic acid (33 mg, 0.20 mmol) were dissolved in *N,N*-dimethylformamide (1 mL) containing triethylamine (84.1 μ L, 0.600 mmol). *O*-(7-Azabenzotriazol-1-yl)-*N,N,N',N'*-tetramethyluronium hexafluorophosphate (84 mg, 0.22 mmol) was added and the solution was allowed to stir for 16 hours. The crude product was purified by reverse-phase preparative liquid chromatography (17 mg, 0.050 mmol, 25 %). ESI-MS *m/z* calc. 340.1, found 340.9 (*M*+1)⁺ Retention time 3.55 minutes.

33) 4-Methyl-*N*-(4-phenyl-thiazol-2-yl)-benzamide




4-Phenyl-thiazol-2-ylamine (35.2 mg, 0.200 mmol) and 4-methyl-benzoyl chloride (30.9 mg, 0.200 mmol) were dissolved in 1 mL of pyridine. The reaction mixture was stirred at room temperature overnight and then purified by reverse-phase preparative liquid chromatography (20.6 mg, 0.0635 mmol, 31.8 %). ESI-MS *m/z* calc. 294.1, found 295.2 (*M*+1)⁺ Retention time 3.55 minutes.

34) 4-Methyl-*N*-(4-phenyl-thiazol-2-yl)-benza




mide 5-Phenyl-thiazol-2-ylamine (35.2 mg, 0.200 mmol) and 4-methyl-benzoyl chloride (30.9 mg, 0.200 mmol) were dissolved in 1 mL of pyridine. The reaction mixture was stirred at room temperature overnight and then purified by reverse-phase preparative liquid chromatography (10.4 mg, 0.0353 mmol, 17.7 %). ESI-MS m/z calc. 294.1, found 295.4 ($M+1$)⁺ Retention time 3.40 minutes.

35) *N*-[4-(2-Methoxy-phenyl)-thiazol-2-yl]-4-methyl-benzamide




4-(2-Methoxy-phenyl)-thiazol-2-ylamine (41.3 mg, 0.200 mmol) and 4-methyl-benzoyl chloride (30.9 mg, 0.200 mmol) were dissolved in 1 mL of pyridine. The reaction mixture was stirred at room temperature overnight and then purified by reverse-phase preparative liquid chromatography (7.32 mg, 0.0226 mmol, 11.3 %). ESI-MS m/z calc. 324.1, found 325.2 ($M+1$)⁺ Retention time 3.75 minutes.

36) *N*-[4-(2-Methoxy-phenyl)-thiazol-2-yl]-benzamide




4-(2-Methoxy-phenyl)-thiazol-2-ylamine (41.3 mg, 0.200 mmol) and benzoyl chloride (28.1 mg, 0.200 mmol) were dissolved in 1 mL of pyridine. The reaction mixture was stirred at room temperature overnight and then purified by reverse-phase preparative liquid chromatography (11.4 mg, 0.0367 mmol, 18.4 %). ESI-MS m/z calc. 310.1, found 311.2 ($M+1$)⁺ Retention time 3.55 minutes.

37) N-(4,5-Diphenyl-thiazol-2-yl)-benzamide




4,5-Diphenyl-thiazol-2-ylamine (50.5 mg, 0.200 mmol) and benzoyl chloride (28.1 mg, 0.200 mmol) were dissolved in 1,4-dioxane (2 mL) containing triethylamine (84.1 μ L, 0.600 mmol). The reaction mixture was subjected to microwave irradiation for 5 minutes at 200 °C. The crude product was filtered, evaporated to dryness, dissolved in 1 mL of dimethylsulfoxide and purified by reverse-phase preparative liquid chromatography (7.21 mg, 0.0202 mmol, 10.1 %). ESI-MS m/z calc. 356.1, found 357.2 ($M+1$)⁺ Retention time 3.95 minutes.

38) N-(4,5-Diphenyl-thiazol-2-yl)-4-methyl-benzamide



4,5-Diphenyl-thiazol-2-ylamine (50.5 mg, 0.200 mmol) and 4-methyl-benzoyl chloride (30.9 mg, 0.200 mmol) were dissolved in 1,4-dioxane (2 mL) containing triethylamine (84.1 μ L, 0.600 mmol). The reaction mixture was subjected to microwave irradiation for 5 minutes at 200 °C. The crude product was filtered, evaporated to dryness, dissolved in 1 mL of dimethylsulfoxide and purified by reverse-phase preparative liquid chromatography (18.6 mg, 0.0502 mmol, 25.1 %). ESI-MS *m/z* calc. 370.1, found 371.2 (M+1)⁺ Retention time 4.13 minutes.


39) *N*-(4,5-Diphenyl-thiazol-2-yl)-4-methoxy-benzamide



4,5-Diphenyl-thiazol-2-ylamine (50.5 mg, 0.200 mmol) and 4-methoxy-benzoyl chloride (34.1 mg, 0.200 mmol) were dissolved in 1,4-dioxane (2 mL) containing triethylamine (84.1 μ L, 0.600 mmol). The reaction mixture was stirred overnight at room temperature. The crude product was filtered, evaporated to dryness, dissolved in 1 mL of dimethylsulfoxide and purified by reverse-phase preparative liquid chromatography (12.5 mg, 0.0323 mmol,


16.2 %). ESI-MS m/z calc. 386.1, found 387.2 ($M+1$)⁺
Retention time 3.95 minutes.

40) 4-Methyl-N-(4-*p*-tolyl-thiazol-2-yl)-benzamide



4-*p*-Tolyl-thiazol-2-ylamine (38.1 mg, 0.200 mmol) and 4-methyl-benzoyl chloride (30.9 mg, 0.200 mmol) were dissolved in 1,4-dioxane (2 mL) containing triethylamine (84.1 μ L, 0.600 mmol). The reaction mixture was stirred overnight at room temperature. The crude product was filtered, evaporated to dryness, dissolved in 1 mL of dimethylsulfoxide and purified by reverse-phase preparative liquid chromatography (13.5 mg, 0.0438, 21.9 %). ESI-MS m/z calc. 308.1, found 309.0 ($M+1$)⁺ Retention time 3.72 minutes.


41) 4-Methyl-N-(5-methyl-thiazol-2-yl)-benzamide



5-Methyl-thiazol-2-ylamine (22.8 mg, 0.200 mmol) and 4-methyl-benzoyl chloride (30.9 mg, 0.200 mmol) were dissolved in 1,4-dioxane (2 mL) containing triethylamine (84.1 μ L, 0.600 mmol). The reaction mixture was stirred overnight at room temperature. The crude product was filtered, evaporated to dryness, dissolved in 1 mL of dimethylsulfoxide and purified by reverse-phase preparative liquid chromatography (9.29 mg, 0.0400, 20.0


%). ESI-MS m/z calc. 232.1, found 233.2 ($M+1$)⁺ Retention time 2.65 minutes.

42) *N*-[5-(2-Chloro-benzyl)-thiazol-2-yl]-2-methoxy-2-phenyl-acetamide



2-Bromo-*N*-[5-(2-chloro-benzyl)-thiazol-2-yl]-2-phenyl-acetamide (42 mg, 0.10 mmol) was dissolved in 5 mL of methanol. The reaction vessel was sealed and then subjected to microwave irradiation for 15 minutes at 125 °C. The crude mixture was evaporated to dryness, dissolved in 1 mL of dimethylsulfoxide and purified by reverse-phase preparative liquid chromatography (16 mg, 0.043, 43 %). ESI-MS m/z calc. 372.1, found 373.2 ($M+1$)⁺ Retention time 3.43 minutes. ¹H NMR (400 MHz, CD₃CN) δ 3.42 (s, 3H), 4.23 (s, 2H), 4.94 (s, 1H), 7.21-7.53 (m, 10H)


43) *N*-[5-(2-Chloro-benzyl)-thiazol-2-yl]-2-methylamino-2-phenyl-acetamide



2-Bromo-*N*-[5-(2-chloro-benzyl)-thiazol-2-yl]-2-phenyl-acetamide (42 mg, 0.10 mmol) was dissolved in 0.5 mL of *N,N*-dimethylformamide containing 1.0 mL of methylamine (2.0 M in tetrahydrofuran, 2.0 mmol). The reaction vessel was sealed and then subjected to


microwave irradiation for 5 minutes at 80 °C. The crude mixture was evaporated to dryness, dissolved in 1 mL of dimethylsulfoxide and purified by reverse-phase preparative liquid chromatography (29 mg, 0.078, 78 %). ESI-MS m/z calc. 371.1, found 372.2 (M+1)⁺ Retention time 2.33 minutes. ¹H NMR (400 MHz, MeOD) δ 2.66 (s, 3H), 4.23 (s, 2H), 5.09 (s, 1H), 7.13-7.56 (m, 10H).

44) N-[5-(2-Chloro-benzyl)-thiazol-2-yl]-2-morpholin-4-yl-2-phenyl-acetamide



2-Bromo-N-[5-(2-chloro-benzyl)-thiazol-2-yl]-2-phenyl-acetamide (42 mg, 0.10 mmol) was dissolved in 0.5 mL of *N,N*-dimethylformamide containing morpholine (174 mg, 2.00 mmol). The reaction vessel was sealed and then subjected to microwave irradiation for 5 minutes at 80 °C. The crude mixture was evaporated to dryness and purified by reverse-phase preparative liquid chromatography (31 mg, 0.072, 72 %). ESI-MS m/z calc. 427.1, found 428.0 (M+1)⁺ Retention time 2.40 minutes.


45) N-[5-(2-Chloro-benzyl)-thiazol-2-yl]-2-dimethylamino-2-phenyl-acetamide



2-Bromo-N-[5-(2-chloro-benzyl)-thiazol-2-yl]-2-phenyl-acetamide (42 mg, 0.10 mmol) was dissolved in 0.5


mL of *N,N*-dimethylformamide containing dimethylamine (90.2 mg, 2.00 mmol). The reaction vessel was sealed and then subjected to microwave irradiation for 5 minutes at 80 °C. The crude mixture was evaporated to dryness and purified by reverse-phase preparative liquid chromatography (13 mg, 0.034, 34 %). ESI-MS *m/z* calc. 385.1, found 386.0 (M+1)⁺ Retention time 2.40 minutes.

46) *N*-[5-(2-Chloro-benzyl)-thiazol-2-yl]-2-dimethylamino-2-phenyl-acetamide



2-Bromo-*N*-[5-(2-chloro-benzyl)-thiazol-2-yl]-butyramide (37 mg, 0.10 mmol) was dissolved in 0.5 mL of *N,N*-dimethylformamide containing aniline (186 mg, 2.00 mmol). The reaction vessel was sealed and then subjected to microwave irradiation for 5 minutes at 80 °C. The crude mixture was evaporated to dryness and purified by reverse-phase preparative liquid chromatography (24 mg, 0.062, 62 %). ESI-MS *m/z* calc. 385.1, found 386.1 (M+1)⁺ Retention time 3.48 minutes.


47) *N*-[5-(2-Chloro-benzyl)-thiazol-2-yl]-2-ethylamino-butylamide



2-Bromo-*N*-[5-(2-chloro-benzyl)-thiazol-2-yl]-butyramide (37 mg, 0.10 mmol) was dissolved in 0.5 mL of *N,N*-dimethylformamide containing ethylamine (2 M in


tetrahydrofuran, 1.00 mL, 2.00 mmol). The reaction vessel was sealed and then subjected to microwave irradiation for 5 minutes at 80 °C. The crude mixture was evaporated to dryness and purified by reverse-phase preparative liquid chromatography (19 mg, 0.056, 56 %). ESI-MS m/z calc. 337.1, found 338.0 (M+1)⁺ Retention time 2.08 minutes.

48) N-[5-(2-Chloro-benzyl)-thiazol-2-yl]-2-piperidin-1-yl-butyramide



2-Bromo-N-[5-(2-chloro-benzyl)-thiazol-2-yl]-butyramide (37 mg, 0.10 mmol) was dissolved in 0.5 mL of *N,N*-dimethylformamide containing piperidine (170 mg, 2.0 mmol). The reaction vessel was sealed and then subjected to microwave irradiation for 5 minutes at 80 °C. The crude mixture was evaporated to dryness and purified by reverse-phase preparative liquid chromatography (31 mg, 0.082, 82 %). ESI-MS m/z calc. 377.1, found 378.2 (M+1)⁺ Retention time 2.21 minutes.

49) N-[5-(2-Chloro-benzyl)-thiazol-2-yl]-2-diethylamino-butyramide



2-Bromo-N-[5-(2-chloro-benzyl)-thiazol-2-yl]-butyramide (37 mg, 0.10 mmol) was dissolved in 0.5 mL of

N,N-dimethylformamide containing diethylamine (146 mg, 2.00 mmol). The reaction vessel was sealed and then subjected to microwave irradiation for 5 minutes at 80 °C. The crude mixture was evaporated to dryness and purified by reverse-phase preparative liquid chromatography (14 mg, 0.038, 38 %). ESI-MS *m/z* calc. 365.1, found 366.2 (M+1)⁺ Retention time 2.18 minutes.

Analytical data for exemplary compounds of the present invention is recited below in Table 2 below.

Table 2

Cmpd #	LC/MS	LC/RT
2	309.00	3.00
4	325.00	3.63
5	325.20	3.53
6	341.00	3.42
7	341.00	3.47
18	281.20	3.20
21	280.80	3.35
23	329.00	3.38
24	310.90	3.58
25	385.20	3.67
28	329.00	3.37
29	335.00	3.50
31	343.20	3.33
32	309.00	3.13
34	374.00	1.83
35	338.00	1.27
37	419.20	3.92
38	419.20	3.85
39	385.30	4.24
47	309.00	3.72
48	295.20	2.83
49	311.00	3.41
50	309.20	2.01
51	387.40	2.96
52	371.20	4.13
53	315.00	3.63
55	287.00	3.53
58	324.20	2.05
66	247.00	2.73
67	295.20	3.52
68	357.20	3.95
69	311.00	3.43
70	371.20	4.11
71	371.20	3.94
72	387.20	3.95
73	425.20	4.41
74	391.20	4.18
75	295.00	3.45
76	311.00	3.52

Cmpd #	LC/MS	LC/RT
77	314.80	3.38
78	311.20	3.37
79	341.00	3.53
80	340.80	3.49
81	311.20	3.35
82	341.00	3.42
83	325.00	3.53
84	340.80	3.38
85	247.00	3.03
86	261.00	3.27
87	261.00	3.30
88	295.00	3.55
89	399.40	3.63
90	433.40	3.79
91	425.20	4.24
92	375.40	3.57
93	383.20	3.57
94	383.40	3.42
95	487.00	4.27
96	337.20	3.53
97	371.00	3.72
98	365.00	3.85
99	399.40	4.02
100	349.20	3.28
101	377.40	3.89
102	411.40	4.07
103	385.20	3.85
104	391.20	2.70
105	387.20	2.78
106	391.20	3.03
107	371.20	2.55
108	323.20	2.30
109	323.20	2.33
110	337.20	2.58
111	321.20	1.98
112	337.00	3.27
113	349.20	3.33
114	363.20	3.57
115	351.00	2.88

Cmpd #	LC/MS	LC/RT
116	351.00	2.51
117	425.20	4.25
118	489.00	4.32
119	357.20	3.53
120	371.20	3.72
121	399.40	4.02
122	411.40	4.09
123	385.20	3.87
124	419.20	3.84
125	487.20	4.27
126	355.20	3.43
127	357.20	3.52
128	371.00	3.70
129	399.20	4.00
130	411.20	4.09
131	433.40	3.75
132	421.20	4.05
133	413.40	3.57
134	483.20	4.10
135	353.00	3.32
136	367.20	3.50
137	395.20	3.84
138	381.20	3.67
139	391.20	4.09
140	429.40	3.62
141	279.00	2.66
142	409.40	3.76
143	403.40	3.42
144	417.20	3.58
145	339.00	3.02
146	355.00	3.22
147	383.40	3.52
148	395.00	3.57
149	327.40	2.91
150	313.20	2.91
151	409.40	3.72
152	339.00	3.00
153	341.00	3.03
154	355.00	3.20
155	383.20	3.52
156	395.00	3.57
157	327.40	2.90
158	313.00	2.90
159	315.00	2.81
160	315.00	2.96
161	306.20	2.48
162	341.00	2.65
163	350.20	2.53
164	325.20	2.51
165	391.20	3.68
166	309.20	2.80
167	323.20	3.05
168	337.20	3.22
169	363.20	3.53
170	351.20	3.43
171	339.00	2.68

Cmpd #	LC/MS	LC/RT
172	385.20	3.45
173	399.20	3.70
174	413.40	3.87
175	439.40	4.15
176	427.40	4.05
177	453.40	4.30
178	415.20	3.40
179	405.40	3.55
180	399.20	3.27
181	335.40	2.83
182	337.00	2.86
183	351.20	3.03
184	379.40	3.37
185	309.20	2.73
186	409.40	3.70
187	471.20	3.85
188	339.20	2.96
189	341.20	3.02
190	355.20	3.18
191	383.20	3.50
192	395.00	3.55
193	369.20	3.32
194	327.40	2.86
195	313.20	2.85
196	563.40	2.65
197	567.20	3.03
198	547.40	2.83
199	537.40	2.51
200	396.20	3.38
201	-	3.83
202	-	99.00
204	371.00	3.67
205	-	99.00
206	309.20	3.25
207	293.00	2.96
208	281.20	2.86
209	321.20	3.35
210	295.20	3.07
211	386.00	2.24
212	414.20	2.31
213	386.20	2.24
214	428.00	2.40
215	426.00	2.33
216	373.20	3.48
217	372.20	2.33
218	324.40	2.03
219	325.20	3.15
220	338.00	2.08
221	366.20	2.18
222	338.00	2.03
223	380.00	2.11
224	396.00	2.33
225	378.20	2.21
226	295.40	3.40
227	345.00	3.62
228	329.00	3.70

Cmpd #	LC/MS	LC/RT
229	349.00	3.79
230	329.00	3.65
231	325.20	3.48
232	344.80	3.60
233	315.00	2.91
234	387.20	3.64
235	409.40	3.44
236	383.20	5.11
237	366.70	3.48
238	370.90	3.67
239	400.30	2.46
240	454.30	2.70
241	412.30	2.48
242	426.10	2.55
243	440.10	2.61
244	456.30	2.41
245	498.30	2.76
246	440.10	2.63
247	498.10	2.70
248	442.30	2.45
249	505.30	2.78
250	426.30	2.63
251	386.10	3.48
252	387.10	3.62
253	498.30	3.66
254	462.00	3.02
255	371.90	3.48
256	416.30	4.54
257	348.10	4.14
258	362.10	4.22
259	390.30	4.42

Cmpd #	LC/MS	LC/RT
260	402.50	4.44
261	376.30	4.32
262	385.00	3.80
263	387.00	3.83
264	375.10	3.07
265	340.90	3.55
266	327.30	3.38
267	325.10	3.70
268	311.10	3.67
269	415.30	3.38
270	401.30	3.10
271	371.10	3.00
272	400.10	1.62
273	397.30	2.93
274	367.10	2.85
275	396.10	1.41
276	362.10	2.56
277	391.10	0.71
278	368.10	2.42
279	309.10	3.14
280	367.10	3.34
281	381.30	3.28
282	416.50	3.41
283	368.10	3.24
284	367.00	3.51
285	385.00	3.60
286	367.00	3.49
287	409.30	3.93
288	361.10	3.84
289	315.30	2.90

EXAMPLE 6

Assays for Detecting and Measuring $\Delta F508$ -CFTR Correction and Potentiator Properties of Compounds

A) Membrane potential optical methods for assaying $\Delta F508$ -CFTR modulation properties of compounds

The optical membrane potential assay utilized voltage-sensitive FRET sensors described by Gonzalez and Tsien (See, Gonzalez, J. E. and R. Y. Tsien (1995) "Voltage sensing by fluorescence resonance energy transfer in single cells" Biophys J 69(4): 1272-80, and Gonzalez, J. E. and R. Y. Tsien (1997) "Improved indicators of cell membrane potential that use fluorescence resonance energy transfer" Chem Biol 4(4): 269-77) in combination with

instrumentation for measuring fluorescence changes such as the Voltage/Ion Probe Reader (VIPR) (See, Gonzalez, J. E., K. Oades, et al. (1999) "Cell-based assays and instrumentation for screening ion-channel targets" Drug Discov Today 4(9): 431-439).

These voltage sensitive assays are based on the change in fluorescence resonant energy transfer (FRET) between the membrane-soluble, voltage-sensitive dye, DiSBAC₂(3), and a fluorescent phospholipid, CC2-DMPE, which is attached to the outer leaflet of the plasma membrane and acts as a FRET donor. Changes in membrane potential (V_m) cause the negatively charged DiSBAC₂(3) to redistribute across the plasma membrane and the amount of energy transfer from CC2-DMPE changes accordingly. The changes in fluorescence emission were monitored using VIPRTM II, which is an integrated liquid handler and fluorescent detector designed to conduct cell-based screens in 96- or 384-well microtiter plates.

Identification of Correction Compounds

To identify small molecules that correct the trafficking defect associated with $\Delta F508$ -CFTR; a single-addition HTS assay format was developed. The cells were incubated in serum-free medium for 16 hrs at 37 °C in the presence or absence (negative control) of test compound. As a positive control, cells plated in 384-well plates were incubated for 16 hrs at 27 °C to "temperature-correct" $\Delta F508$ -CFTR. The cells were subsequently rinsed 3X with Krebs Ringers solution and loaded with the voltage-sensitive dyes. To activate $\Delta F508$ -CFTR, 10 mM forskolin and the CFTR potentiator, genistein (20 mM), were added along with Cl⁻-free medium to each well. The addition of Cl⁻-free medium promoted Cl⁻ efflux in

response to $\Delta F508$ -CFTR activation and the resulting membrane depolarization was optically monitored using the FRET-based voltage-sensor dyes.

Identification of Potentiator Compounds

To identify potentiators of $\Delta F508$ -CFTR, a double-addition HTS assay format was developed. During the first addition, a Cl^- -free medium with or without test compound was added to each well. After 22 sec, a second addition of Cl^- -free medium containing 2 - 10 μM forskolin was added to activate $\Delta F508$ -CFTR. The extracellular Cl^- concentration following both additions was 28 mM, which promoted Cl^- efflux in response to $\Delta F508$ -CFTR activation and the resulting membrane depolarization was optically monitored using the FRET-based voltage-sensor dyes.

Solutions

Bath Solution #1: (in mM) NaCl 160, KCl 4.5, CaCl_2 2,
 MgCl_2 1, HEPES 10, pH 7.4 with NaOH.

Chloride-free bath solution: Chloride salts in Bath
 Solution #1 are substituted with
 gluconate salts.

CC2-DMPE: Prepared as a 10 mM stock solution in
 DMSO and stored at -20°C .

DiSBAC₂(3): Prepared as a 10 mM stock in DMSO and
 stored at -20°C .

Cell Culture

NIH3T3 mouse fibroblasts stably expressing $\Delta F508$ -CFTR are used for optical measurements of membrane potential. The cells are maintained at 37°C in 5% CO_2 and 90 % humidity in Dulbecco's modified Eagle's medium supplemented with 2 mM glutamine, 10 % fetal bovine serum, 1 X NEAA, β -ME, 1 X pen/strep, and 25 mM HEPES in

175 cm² culture flasks. For all optical assays, the cells were seeded at 30,000/well in 384-well matrigel-coated plates and cultured for 2 hrs at 37 °C before culturing at 27 °C for 24 hrs. for the potentiator assay. For the correction assays, the cells are cultured at 27 °C or 37 °C with and without compounds for 16 - 24 hours.

B) Electrophysiological Assays for assaying $\Delta F508$ -CFTR modulation properties of compounds

1. Ussing Chamber Assay

Ussing chamber experiments were performed on polarized epithelial cells expressing $\Delta F508$ -CFTR to further characterize the $\Delta F508$ -CFTR modulators identified in the optical assays. FRT ^{$\Delta F508$ -CFTR} epithelial cells grown on Costar Snapwell cell culture inserts were mounted in an Ussing chamber (Physiologic Instruments, Inc., San Diego, CA), and the monolayers were continuously short-circuited using a Voltage-clamp System (Department of Bioengineering, University of Iowa, IA, and, Physiologic Instruments, Inc., San Diego, CA). Transepithelial resistance was measured by applying a 2-mV pulse. Under these conditions, the FRT epithelia demonstrated resistances of 4 K Ω / cm² or more. The solutions were maintained at 27 °C and bubbled with air. The electrode offset potential and fluid resistance were corrected using a cell-free insert. Under these conditions, the current reflects the flow of Cl⁻ through $\Delta F508$ -CFTR expressed in the apical membrane. The I_{sc} was digitally acquired using an MP100A-CE interface and AcqKnowledge software (v3.2.6; BIOPAC Systems, Santa Barbara, CA).

Identification of Correction Compounds

Typical protocol utilized a basolateral to apical membrane Cl^- concentration gradient. To set up this gradient, normal ringer was used on the basolateral membrane, whereas apical NaCl was replaced by equimolar sodium gluconate (titrated to pH 7.4 with NaOH) to give a large Cl^- concentration gradient across the epithelium. All experiments were performed with intact monolayers. To fully activate ΔF508 -CFTR, forskolin (10 mM) and the PDE inhibitor, IBMX (100 mM), were applied followed by the addition of the CFTR potentiator, genistein (50 mM).

As observed in other cell types, incubation at low temperatures of FRT cells stably expressing ΔF508 -CFTR increases the functional density of CFTR in the plasma membrane. To determine the activity of correction compounds, the cells were incubated with 10 mM of the test compound for 24 hours at 37°C and were subsequently washed 3X prior to recording. The cAMP- and genistein-mediated I_{sc} in compound-treated cells was normalized to the 27°C and 37°C controls and expressed as percentage activity. Preincubation of the cells with the correction compound significantly increased the cAMP- and genistein-mediated I_{sc} compared to the 37°C controls.

Identification of Potentiator Compounds

Typical protocol utilized a basolateral to apical membrane Cl^- concentration gradient. To set up this gradient, normal ringers was used on the basolateral membrane and was permeabilized with nystatin (360 $\mu\text{g/ml}$), whereas apical NaCl was replaced by equimolar sodium gluconate (titrated to pH 7.4 with NaOH) to give a large Cl^- concentration gradient across the epithelium. All experiments were performed 30 min after nystatin

permeabilization. Forskolin (10 mM) and all test compounds were added to both sides of the cell culture inserts. The efficacy of the putative $\Delta F508$ -CFTR potentiators was compared to that of the known potentiator, genistein.

Solutions

Basolateral solution (in mM): NaCl (135), CaCl_2 (1.2), MgCl_2 (1.2), K_2HPO_4 (2.4), KHPO_4 (0.6), N-2-hydroxyethylpiperazine-N'-2-ethanesulfonic acid (HEPES) (10), and dextrose (10). The solution was titrated to pH 7.4 with NaOH.

Apical solution (in mM): Same as basolateral solution with NaCl replaced with Na Gluconate (135).

Cell Culture

Fisher rat epithelial (FRT) cells expressing $\Delta F508$ -CFTR ($\text{FRT}^{\Delta F508\text{-CFTR}}$) were used for Ussing chamber experiments for the putative $\Delta F508$ -CFTR modulators identified from our optical assays. The cells were cultured on Costar Snapwell cell culture inserts and cultured for five days at 37 °C and 5% CO_2 in Coon's modified Ham's F-12 medium supplemented with 5% fetal calf serum, 100 U/ml penicillin, and 100 $\mu\text{g/ml}$ streptomycin. Prior to use for characterizing the potentiator activity of compounds, the cells were incubated at 27 °C for 16 - 48 hrs to correct for the $\Delta F508$ -CFTR. To determine the activity of corrections compounds, the cells were incubated at 27 °C or 37 °C with and without the compounds for 24 hours.

2. *Whole-cell recordings*

The macroscopic $\Delta F508$ -CFTR current ($I_{\Delta F508}$) in temperature and test compound-corrected NIH3T3 cells stably expressing $\Delta F508$ -CFTR were monitored using the perforated-patch, whole-cell recording. Briefly, voltage-clamp recordings of $I_{\Delta F508}$ were performed at room temperature using an Axopatch 200B patch-clamp amplifier (Axon Instruments Inc., Foster City, CA). All recordings were acquired at a sampling frequency of 10 kHz and low-pass filtered at 1 kHz. Pipettes had a resistance of 5–6 M Ω when filled with the intracellular solution. Under these recording conditions, the calculated reversal potential for Cl^- (E_{Cl}) at room temperature was -28 mV. All recordings had a seal resistance > 20 G Ω and a series resistance < 15 M Ω . Pulse generation, data acquisition, and analysis were performed using a PC equipped with a Digidata 1320 A/D interface in conjunction with Clampex 8 (Axon Instruments Inc.). The bath contained < 250 ml of saline and was continuously perfused at a rate of 2 ml/min using a gravity-driven perfusion system.

Identification of Correction Compounds

To determine the activity of correction compounds for increasing the density of functional $\Delta F508$ -CFTR in the plasma membrane, we used the above-described perforated-patch-recording techniques to measure the current density following 24-hr treatment with the correction compounds. To fully activate $\Delta F508$ -CFTR, 10 mM forskolin and 20 mM genistein were added to the cells. Under our recording conditions, the current density following 24-hr incubation at 27°C was higher than that observed following 24-hr incubation at 37 °C. These results are consistent with the known effects of low-temperature incubation on

the density of $\Delta F508$ -CFTR in the plasma membrane. To determine the effects of correction compounds on CFTR current density, the cells were incubated with 10 μM of the test compound for 24 hours at 37°C and the current density was compared to the 27°C and 37°C controls (% activity). Prior to recording, the cells were washed 3X with extracellular recording medium to remove any remaining test compound. Preincubation with 10 μM of correction compounds significantly increased the cAMP- and genistein-dependent current compared to the 37°C controls.

Identification of Potentiator Compounds

The ability of $\Delta F508$ -CFTR potentiators to increase the macroscopic $\Delta F508$ -CFTR Cl^- current ($I_{\Delta F508}$) in NIH3T3 cells stably expressing $\Delta F508$ -CFTR was also investigated using perforated-patch-recording techniques. The potentiators identified from the optical assays evoked a dose-dependent increase in $I_{\Delta F508}$ with similar potency and efficacy observed in the optical assays. In all cells examined, the reversal potential before and during potentiator application was around -30 mV, which is the calculated E_{Cl} (-28 mV).

Solutions

Intracellular solution (in mM): Cs-aspartate (90), CsCl (50), $MgCl_2$ (1), HEPES (10), and 240 $\mu g/ml$ amphotericin-B (pH adjusted to 7.35 with CsOH).

Extracellular solution (in mM): N-methyl-D-glucamine (NMDG)-Cl (150), $MgCl_2$ (2), $CaCl_2$ (2), HEPES (10) (pH adjusted to 7.35 with HCl).

Cell Culture

NIH3T3 mouse fibroblasts stably expressing $\Delta F508$ -CFTR are used for whole-cell recordings. The cells are maintained at 37 °C in 5% CO₂ and 90 % humidity in Dulbecco's modified Eagle's medium supplemented with 2 mM glutamine, 10 % fetal bovine serum, 1 X NEAA, β -ME, 1 X pen/strep, and 25 mM HEPES in 175 cm² culture flasks. For whole-cell recordings, 2,500-5,000 cells were seeded on poly-L-lysine-coated glass coverslips and cultured for 24 - 48 hrs at 27 °C before use to test the activity of potentiators; and incubated with or without the correction compound at 37 °C for measuring the activity of correctors.

3. Single-channel recordings

The single-channel activities of temperature-corrected $\Delta F508$ -CFTR stably expressed in NIH3T3 cells and activities of potentiator compounds were observed using excised inside-out membrane patch. Briefly, voltage-clamp recordings of single-channel activity were performed at room temperature with an Axopatch 200B patch-clamp amplifier (Axon Instruments Inc.). All recordings were acquired at a sampling frequency of 10 kHz and low-pass filtered at 400 Hz. Patch pipettes were fabricated from Corning Kovar Sealing #7052 glass (World Precision Instruments, Inc., Sarasota, FL) and had a resistance of 5 - 8 M Ω when filled with the extracellular solution. The $\Delta F508$ -CFTR was activated after excision, by adding 1 mM Mg-ATP, and 75 nM of the cAMP-dependent protein kinase, catalytic subunit (PKA; Promega Corp. Madison, WI). After channel activity stabilized, the patch was perfused using a gravity-driven microperfusion system. The inflow was placed adjacent to the patch, resulting in complete solution exchange within 1-2 sec.

To maintain $\Delta F508$ -CFTR activity during the rapid perfusion, the nonspecific phosphatase inhibitor F^- (10 mM NaF) was added to the bath solution. Under these recording conditions, channel activity remained constant throughout the duration of the patch recording (up to 60 min). Currents produced by positive charge moving from the intra- to extracellular solutions (anions moving in the opposite direction) are shown as positive currents. The pipette potential (V_p) was maintained at 80 mV.

Channel activity was analyzed from membrane patches containing ≤ 2 active channels. The maximum number of simultaneous openings determined the number of active channels during the course of an experiment. To determine the single-channel current amplitude, the data recorded from 120 sec of $\Delta F508$ -CFTR activity was filtered "off-line" at 100 Hz and then used to construct all-point amplitude histograms that were fitted with multigaussian functions using Bio-Patch Analysis software (Bio-Logic Comp. France). The total microscopic current and open probability (P_o) were determined from 120 sec of channel activity. The P_o was determined using the Bio-Patch software or from the relationship $P_o = I/i(N)$, where I = mean current, i = single-channel current amplitude, and N = number of active channels in patch.

Solutions

Extracellular solution (in mM): NMDG (150), aspartic acid (150), $CaCl_2$ (5), $MgCl_2$ (2), and HEPES (10) (pH adjusted to 7.35 with Tris base).

Intracellular solution (in mM): NMDG-Cl (150), $MgCl_2$ (2), EGTA (5), TES (10), and

Tris base (14) (pH adjusted to 7.35 with HCl).

Cell Culture

NIH3T3 mouse fibroblasts stably expressing $\Delta F508$ -CFTR are used for excised-membrane patch-clamp recordings. The cells are maintained at 37 °C in 5% CO₂ and 90 % humidity in Dulbecco's modified Eagle's medium supplemented with 2 mM glutamine, 10% fetal bovine serum, 1 X NEAA, β -ME, 1 X pen/strep, and 25 mM HEPES in 175 cm² culture flasks. For single channel recordings, 2,500 - 5,000 cells were seeded on poly-L-lysine-coated glass coverslips and cultured for 24 - 48 hrs at 27 °C before use.

The compounds of the present invention were found to modulate CFTR activity when tested using the above methods. The activity of exemplary compounds of the present invention is recited below in Table 3.

EC50: "+++" means <10 uM; "++" means between 10uM to 25 uM; "+" means between 25 uM to 60uM.

% Efficacy: "+" means < 25%; "++" means between 25% to 100%; "+++" means > 100%.

Table 3

Cmpd #	EC50 uM	% Activity
1	++	++
2	+++	+++
3	++	++
4	+++	++
5	++	++
6	++	++
7	++	+++
8	+++	+++
9	+++	++
10	++	+++
11	++	+++
12	+++	+++
13	+++	++

Cmpd #	EC50 uM	% Activity
14	+++	++
15	+++	++
16	+++	+++
17	+++	++
18	++	++
19	++	++
20	++	++
21	++	++
22	++	++
23	+++	++
24	++	+++
25	+++	++
26	++	++

Cmpd #	EC50 uM	% Activity
27	+++	++
28	++	++
29	+++	++
30	+++	+++
31	+++	++
32	++	++
33	++	++
34	++	++
35	++	++
36	++	++
37	+++	+++
38	+++	++
39	+++	+++
40	+++	++
41	+++	++
42	+++	++
43	+++	++
44	++	+++
45	++	++
46	+++	++
47	++	++
48	+++	+++
49	++	++
50	+++	+++
51	+++	+++
52	+++	+++
53	+++	+++
54	++	++
55	++	++
56	+++	++
57	+++	++
58	++	++
59	++	++
60	++	++
61	++	++
62	+++	+++
63	++	+++
64	+++	+++
65	++	++
66	++	++
67	++	++
68	+++	++
69	++	+++
70	+++	+++
71	+++	++
72	+++	+++
73	++	++
74	++	++
75	++	++
76	++	+++
77	++	++
78	++	++
79	++	++
80	++	++
81	++	++
82	++	++

Cmpd #	EC50 uM	% Activity
83	++	+++
84	++	++
85	++	++
86	++	++
87	++	++
88	++	+++
89	+++	+++
90	+++	+++
91	+++	+++
92	+++	++
93	+++	+++
94	+++	++
95	+++	++
96	++	++
97	+++	+++
98	+++	++
99	+++	+++
100	++	++
101	+++	++
102	+++	+++
103	+++	+++
104	+++	++
105	+++	+++
106	+++	++
107	+++	+++
108	+++	++
109	+++	++
110	+++	+++
111	+++	+++
112	+++	++
113	+++	++
114	+++	++
115	+++	+++
116	+++	++
117	+++	+++
118	+++	++
119	+++	+++
120	+++	++
121	+++	+++
122	+++	+++
123	+++	+++
124	+++	++
125	+++	++
126	+	++
127	+++	++
128	+++	+++
129	+++	++
130	+++	+++
131	+++	+++
132	+++	++
133	+++	++
134	+++	++
135	+++	++
136	+++	++
137	+++	++
138	+++	++

Cmpd #	EC50 uM	% Activity
139	+++	++
140	+++	++
141	++	++
142	+++	++
143	+	++
144	+	++
145	++	++
146	+++	++
147	+++	++
148	+++	++
149	++	++
150	++	++
151	+++	++
152	++	++
153	+++	++
154	+++	++
155	+++	++
156	+++	++
157	+++	++
158	+++4	++
159	+++	++
160	++	+++
161	+	++
162	+	++
163	++	++
164	++	++
165	+++	+++
166	++	++
167	+++	++
168	+++	++
169	+++	++
170	+++	++
171	+++	++
172	+++	+++
173	+++	++
174	+++	++
175	+++	++
176	+++	++
177	++	++
178	+++	+++
179	++	++
180	++	++
181	+	++
182	+	++
183	+	++
184	++	++
185	++	++
186	++	++
187	++	++
188	+	++
189	+	++
190	+	++
191	+++	++
192	++	++
193	++	++
194	+	++



Cmpd #	EC50 uM	% Activity
195	+++	++
196	+++	++
197	+++	++
198	+++	++
199	+++	+++
200	+	++
201	+++	+++
202	+++	++
203	+++	++
204	+++	++
205	+++	++
206	+++	++
207	+++	++
208	+	++
209	+++	++
210	++	++
211	+++	++
212	+++	++
213	++	++
214	++	++
215	+++	++
216	++	++
217	+	++
218	+	++
219	+	++
220	+	++
221	++	++
222	+	++
223	+	++
224	+++	++
225	+++	++
226	++	++
227	++	++
228	++	++
229	++	++
230	+++	+++
231	++	+++
232	++	++
233	+	++
234	++	++
235	+++	++
236	+++	++
237	+++	+++
238	+++	+++
239	+++	++
240	+++	++
241	+++	++
242	+++	+++
243	+++	+++
244	+++	++
245	+++	++
246	+++	++
247	+++	+++
248	+++	++
249	+++	++
250	+++	++

Cmpd #	EC50 uM	% Activity
251	+++	++
252	++	++
253	+	++
254	+++	++
255	+++	++
256	+++	++
257	+	++
258	++	++
259	+++	++
260	+++	++
261	+++	++
262	+	++
263	+++	++
264	+	++
265	+++	+++
266	+++	++
267	+++	++
268	+++	++
269	+++	++
270	+++	++

Cmpd #	EC50 uM	% Activity
271	+++	++
272	+++	+++
273	+++	++
274	+	++
275	+	++
276	+	++
277	+	++
278	+++	++
279	+	++
280	+++	++
281	++	++
282	+++	++
283	+++	++
284	+++	++
285	+++	++
286	+++	++
287	+++	++
288	+++	++
289	+	++

Claims

1. A method of modulating ABC transporter activity, comprising the step of contacting said ABC transporter with a compound of formula I or formula I':



or a pharmaceutically acceptable salt thereof;
wherein:

Y' is O, S, or NR;

p is 0-2;



X is a bond, O, S, S(O), S(O)₂, CF₂, CH₂, -CHOR-, -C(O)-, -O-C(O)-, -C(O)-O-, -C(O)-NR-, -NR-C(O)-, -NR-C(O)-O-, -O-C(O)-NR-, -NR-C(O)-NR-, or NR;

R is H, R², or R⁶;

A is aliphatic, aryl, heteroaryl, heterocyclic, or cycloalkyl;

C is a phenyl or 5-8 membered cycloaliphatic ring;

Q is selected from:



each B is independently selected from 3-7 membered monocyclic or 8-14 membered bicyclic or tricyclic, saturated, unsaturated or aromatic ring containing 0-4

heteroatoms in each ring, wherein each said heteroatom is independently selected from N, NH, S, or O;

wherein each A, B, and C is independently and optionally substituted with up to 4 substituents independently selected from R¹, R², R³, R⁴, or R⁵;

R^L is -OR^A, -SR^A, or -N(R^{AB})₂;

each R^A is independently hydrogen, C1-C6 aliphatic, or a 3-7 membered carbocyclic or heterocyclic ring, saturated or unsaturated ring, having up to 3 heteroatoms selected from O, N, or S, wherein each R^A is optionally substituted with up to 3 substituents independently selected from R¹, R⁴ or R⁷,

each R^{AB} is independently hydrogen or C1-C6 aliphatic optionally substituted with up to 3 substituents independently selected from R¹, R⁴ or R⁷;

wherein up to two methylene units in R^A or R^{AB} are optionally replaced with -CO-, -CS-, -COCO-, -CONR-, -CO₂-, -OCO-, -NRCO₂-, -O-, -NRCONR-, -OCONR-, -NRCO-, -S-, -SO, -SO₂-, -NR-, -SO₂NR-, NRSO₂-, or -NRSO₂NR; or

two R^{AB}, taken together with the nitrogen atom, is a 3-7 membered heterocyclic or heteroaryl ring containing up to 4 heteroatoms selected from O, N, or S, wherein said ring is optionally substituted with up to 2 substituents selected from oxo or (C₁₋₄aliphatic)_p-Y;

R^M is C1-C6 aliphatic, optionally substituted with up to two substituents selected from R¹, R², R³, or R⁴;

each of X₁ and X₂ is independently selected from O, S, or NR;

R^N is C1-C6 aliphatic or phenyl, wherein R^N is optionally substituted with up to two substituents selected from R¹, R², R³, or R⁴;

R^P is C1-C6 aliphatic, optionally substituted with up to two substituents selected from R¹, R², R³, or R⁴;

R^2 is C1-C6 aliphatic or aryl, wherein R^2 is optionally substituted with up to two substituents selected from R^1 , R^2 , R^3 , or R^4 ;

R^1 is oxo, R^6 or ((C1-C4)aliphatic) $_n$ -Y;

n is 0 or 1;

Y is halo, CN, NO₂, CF₃, OCF₃, OH, SR⁶, S(O)R⁶, SO₂R⁶, NH₂, NHR⁶, N(R⁶)₂, NR⁶R⁸, N(R⁸)₂, COOH, COOR⁶ or OR⁶; or

two R^1 on adjacent ring atoms, taken together, form 1,2-methylenedioxy or 1,2-ethylenedioxy;

R^2 is aliphatic, wherein each R^2 optionally comprises up to 2 substituents independently selected from R^1 , R^4 , or R^5 ;

R^3 is a cycloaliphatic, aryl, heterocyclic, or heteroaryl ring optionally comprising up to 3 substituents, independently selected from R^1 , R^2 , R^4 or R^5 ;

R^4 is OR⁵, OR⁶, OC(O)R⁶, OC(O)R⁵, OC(O)OR⁶, OC(O)OR⁵, OC(O)N(R⁶)₂, OC(O)N(R⁵)₂, OC(O)N(R⁶R⁵), SR⁶, SR⁵, S(O)R⁶, S(O)R⁵, SO₂R⁶, SO₂R⁵, SO₂N(R⁶)₂, SO₂N(R⁵)₂, SO₂NR⁵R⁶, SO₃R⁶, SO₃R⁵, C(O)R⁵, C(O)OR⁵, C(O)R⁶, C(O)OR⁶, C(O)N(R⁶)₂, C(O)N(R⁵)₂, C(O)N(R⁵R⁶), C(O)N(OR⁶)R⁶, C(O)N(OR⁵)R⁶, C(O)N(OR⁶)R⁵, C(O)N(OR⁵)R⁵, C(NOR⁶)R⁶, C(NOR⁶)R⁵, C(NOR⁵)R⁶, C(NOR⁵)R⁵, N(R⁶)₂, N(R⁵)₂, N(R⁵R⁶), NR⁵C(O)R⁵, NR⁶C(O)R⁶, NR⁶C(O)R⁵, NR⁵C(O)R⁶, NR⁶C(O)OR⁶, NR⁵C(O)OR⁶, NR⁶C(O)OR⁵, NR⁵C(O)OR⁵, NR⁶C(O)N(R⁶)₂, NR⁶C(O)NR⁵R⁶, NR⁶C(O)N(R⁵)₂, NR⁵C(O)N(R⁶)₂, NR⁵C(O)NR⁵R⁶, NR⁵C(O)N(R⁵)₂, NR⁶SO₂R⁶, NR⁶SO₂R⁵, NR⁵SO₂R⁵, NR⁵SO₂R⁶, NR⁶SO₂N(R⁶)₂, NR⁶SO₂NR⁵R⁶, NR⁶SO₂N(R⁵)₂, NR⁵SO₂N(R⁶)₂,

$\text{NR}^5\text{SO}_2\text{NR}^5\text{R}^6$, $\text{NR}^5\text{SO}_2\text{N}(\text{R}^5)_2$, $\text{N}(\text{OR}^6)\text{R}^6$, $\text{N}(\text{OR}^6)\text{R}^5$, $\text{N}(\text{OR}^5)\text{R}^5$,
or $\text{N}(\text{OR}^5)\text{R}^6$;

R^5 is a cycloaliphatic, aryl, heterocyclic, or heteroaryl ring, optionally comprising up to 3 R^1 substituents;

R^6 is H or aliphatic, wherein R^6 optionally comprises a R^7 substituent;

R^7 is a cycloaliphatic, aryl, heterocyclic, or heteroaryl ring, and each R^7 optionally comprises up to 2 substituents independently chosen from H, $(\text{C}_1\text{-C}_6)$ -straight or branched alkyl, $(\text{C}_2\text{-C}_6)$ straight or branched alkenyl or alkynyl, 1,2-methylenedioxy, 1,2-ethylenedioxy, or $(\text{CH}_2)_n\text{-Z}$;

Z is selected from halo, CN, NO_2 , CF_3 , OCF_3 , OH, S-aliphatic, S(O)-aliphatic, SO_2 -aliphatic, NH_2 , NH-aliphatic, $\text{N}(\text{aliphatic})_2$, $\text{N}(\text{aliphatic})\text{R}^8$, NHR^8 , $\text{N}(\text{R}^8)_2$, COOH, $\text{C}(\text{O})\text{O}(-\text{aliphatic})$, or O-aliphatic; and

R^8 is an amino-capping group.

2. The method according to claim 1, wherein Y' is S or O.

3. The method according to claim 2, wherein p is 1 and X is attached to the carbon adjacent to Y' atom.

4. The method according to claim 2, wherein p is 1 and X is attached to the carbon adjacent to the ring nitrogen atom.

5. The method according to claim 2, wherein, p is 2.

6. The method according to claim 1, wherein X is a bond, O, S, CH₂, CF₂, CHOR, C(O)NR, C(O)O, NRC(O), or NR.

7. The method according to claim 6, wherein X is a bond, CH₂, CHOH, C(O), or C(O)O.


8. The method according to claim 1, wherein A is optionally substituted (C1-C10)aliphatic.


9. The method according to claim 8, wherein A is optionally substituted methyl, ethyl, propyl, butyl, pentyl, or hexyl.

10. The method according to claim 1, wherein A is optionally substituted C6-C10 aryl ring.

11. The method according to claim 10, wherein A is optionally substituted phenyl or naphthyl.

12. The method according to claim 1, wherein A, X, and the ring attached thereto, taken together, is selected from:





wherein:

R^{Ph} is independently R^1 , R^2 , or R^3 ; and

r is 0-3.


X_5 is CH_2 , $\text{C}(\text{O})$, or CHOR ;

X_6 is O or NR^2 ; and


R^{Ak} is C1-C6 aliphatic, optionally substituted with R^1 , R^2 , or R^3 .

13. The method according to claim 12, wherein each B is independently an optionally substituted phenyl or naphthyl.

14. The method according to claim 1, wherein said compound has formula I-a:



15. The method according to claim 14, wherein said B is selected from:



wherein X_3 is O, S, or NR.

16. The method according to claim 15, wherein B is substituted with up to two R^1 substituents.

17. The method according to claim 16, wherein said substituent is selected from C1-C4 alkyl, -O-C1-C4 alkyl, CN, halo, COOH, -C(O)NH₂, -C(O)O(C1-C4 alkyl), -C(O)NH(C1-C4 alkyl), -C(O)N(C1-C4 alkyl)₂, or phenyl optionally substituted with up to two substituents selected from C1-C4 alkyl, -O-C1-C4 alkyl, CN, halo, COOH, -C(O)NH₂, -C(O)O(C1-C4 alkyl), -C(O)NH(C1-C4 alkyl), -C(O)N(C1-C4 alkyl)₂.

18. The method according to claim 15, wherein said B is selected from rings i, iii, iv, v, vi, or vii.

19. The method according to claim 15, wherein said B is ring **x**.

20. The method according to claim 15, wherein said B is selected from ring **xi**, **xii**, **xiii**, or **xiv**.

21. The method according to claim 15, wherein X is a bond and A is optionally substituted phenyl and A is attached to the carbon atom adjacent to the nitrogen ring atom.

22. The method according to claim 21, wherein A is phenyl optionally substituted with up to two substituents selected from C1-C4 alkyl, C1-C4 alkoxy, cyano, halo, N-pyrrolidinyl, N-piperidinyl, or methylenedioxy.

23. The method according claim 22, wherein A is phenyl, 3-methoxyphenyl, 2-methoxyphenyl, 4-cyanophenyl, 4-chlorophenyl, 4-(N-pyrrolidinyl)phenyl, 4-tolyl, 3,4-methylenedioxyphenyl, 3-chlorophenyl, 2,4-dimethoxyphenyl, 2-chlorophenyl, 4-bromophenyl, 2,5-dimethylphenyl, or 2,4-dimethylphenyl.

24. The method according to claim 15, wherein A is selected from cyclopropyl, cyclopentyl, cyclohexyl, cycloheptyl, or adamantyl.


25. The method according to claim 15, wherein the compounds have one or more of the following features:

- a) R is hydrogen;
- b) Y' is S;
- c) A is phenyl optionally substituted with up to two substituents selected from C1-C4 alkyl, C1-C4

alkoxy, cyano, halo, N-pyrrolidinyl, N-piperidinyl, or methylenedioxy; and

d) B is phenyl optionally substituted with up to two substituents selected from C1-C4 alkyl, -O-C1-C4 alkyl, CN, halo, COOH, -C(O)NH₂, -C(O)O(C1-C4 alkyl), -C(O)NH(C1-C4 alkyl), -C(O)N(C1-C4 alkyl)₂, or phenyl optionally substituted with up to two substituents selected from C1-C4 alkyl, -O-C1-C4 alkyl, CN, halo, COOH, -C(O)NH₂, -C(O)O(C1-C4 alkyl), -C(O)NH(C1-C4 alkyl), -C(O)N(C1-C4 alkyl)₂.

26. The method according to claim 1, wherein said compounds have formula **I-b**:



27. The method according to claim 26, wherein p is 1, X is a bond, and A is an optionally substituted phenyl ring.

28. The method according to claim 26 wherein p is 2, each X is a bond, and each A is an optionally substituted phenyl.


29. The method according to claim 26 wherein X is CH₂, CHOH, or C(O), and A is an optionally substituted phenyl.

30. The method according to claim 26 wherein said C1-C6 aliphatic is C1-C4 straight or branched alkylidene.

31. The method according to claim 30, wherein said alkylidene is selected from $-\text{CH}_2-$, $-\text{CH}(\text{Me})-$, $-\text{C}(\text{Me})_2-$, $-\text{CH}(\text{Et})-$, $-\text{C}(\text{Et})_2-$, or $-\text{CH}_2-\text{CH}(\text{Me})-$.

32. The method according to claim 26, wherein B is selected from C3-C8 cycloalkyl, phenyl, piperidyl, or pyrrolidinyl, optionally substituted with up to two R^1 substituents.

33. The method according to claim 26, wherein said (C1-C6 aliphatic)-B, together, is selected from:



wherein:

Ak is C1-C6 straight or branched alkylidene;

X_4 is CH_2 , O or S;

Ar' is phenyl optionally substituted with up to two R^1 ; and

B is optionally substituted with up to two R^1 .


34. The method according to claim 33, wherein Ak is selected from CH_2 , $\text{CH}(\text{CH}_3)$, $\text{C}(\text{CH}_3)_2$, $\text{CH}(\text{Et})$, $\text{C}(\text{Et})_2$, $\text{CH}(\text{n-propyl})$, $\text{CH}(\text{i-Pr})$, $\text{CH}(\text{n-butyl})$, $\text{CH}(\text{but-2-yl})$, or $\text{CH}(\text{t-butyl})$.

35. The method according to claim 33, wherein Ar' is phenyl optionally substituted with halo, C1-C4 alkyl, or O-(C1-C4 alkyl).

36. The method according to claim 33, wherein X_4 is S or O.

37. The method according to claim 26 or 33, wherein R is hydrogen.

38. The method according to claim 1, wherein said compound has formula **I-f**:



wherein:

Y' is O or S;

X₁ is O, S, or NR;

R^M is C1-C6 aliphatic or phenyl, wherein R^M is optionally substituted with up to two substituents independently selected from R¹, R², or R³;

R^N is C1-C6 aliphatic or a 3-7 membered monocyclic, saturated, unsaturated or aromatic ring containing 0-4 heteroatoms in each ring, wherein each said heteroatom is independently selected from N, NH, S, or O;

wherein R^N is optionally substituted with up to 4 substituents independently selected from R¹, R², R³, R⁴, or R⁵.

39. The method according to claim 38, wherein p is 1.

40. The method according to claim 38, wherein X is CH₂, CHOH, or C(O).

41. The method according to claim 38, wherein X is a bond, and A is an optionally substituted phenyl.

42. The method according to claim 38, wherein p is 2, each X is a bond, and each A is an optionally substituted phenyl.

43. The method according to claim 38, wherein X_1 is NH or N(C1-C4 alkyl).

44. The method according to claim 38, wherein X_1 is O.

45. The method according to claim 38, wherein R^M is optionally substituted phenyl.

46. The method according to claim 38, wherein R^M is C1-C4 alkyl.


47. The method according to claim 38, wherein R^N is optionally substituted C3-C7 cycloaliphatic, phenyl, or benzyl.

48. The method according to claim 38, wherein R^N is C1-C6 aliphatic.

49. The method according to claim 38, R is hydrogen.

50. The method according to claim 38, wherein Y' is S.

51. The method according to claim 1, wherein said compound has formula **I-g**:



I-g

wherein:

Y' is O or S;

R^P is C1-C8 aliphatic optionally substituted with up to two substituents independently selected from R¹, R², or R³.

52. The method according to claim 51, wherein p is 1 and X is a bond.

53. The method according to claim 51, wherein p is 1, X is CH₂, CHOH, or C(O).

54. The method according to claim 51, wherein X is a bond, and A is an optionally substituted phenyl.

55. The method according to claim 51, wherein p is 2, each X is a bond, and each A is an optionally substituted phenyl.

56. The method according to claim 51, wherein R^P is C1-C4 alkyl, optionally substituted with up to two R¹.

57. The method according to claim 56, wherein R^P is selected from ethyl, n-propyl, i-propyl, n-butyl, but-2-yl, isoamyl, or t-butyl, optionally substituted with halo, CN, COOH, or CONH₂.


58. The method according to claim 51, wherein R is hydrogen.

59. The method according to claim 51, wherein said compounds have one or more of the following features:

a) Y' is S;

- b) R is hydrogen;
- c) p is 2 and each A is phenyl;
- d) R^p is isoamyl, t-butyl, ethyl, isopropyl, n-propyl, 1-carboxy-prop-3-yl, or 1-carboxy-2-methyl-prop-3-yl.

60. The method according to claim 1, wherein said compound has formula **I-h**:



wherein:

- Y' is O or S;
- X₂ is O, S, or NR;
- R^Q is C1-C6 aliphatic or phenyl, optionally substituted with up to two substituents independently selected from R¹, R², or R³.

61. The method according to claim 60, wherein p is 1 and X is a bond.

62. The method according to claim 60, wherein X is CH₂, CHOH, or C(O).

63. The method according to claim 60, wherein X is a bond, and A is an optionally substituted phenyl.

64. The method according to claim 60, wherein p is 2, each X is a bond, and each A is an optionally substituted phenyl.

65. The method according to claim 60, wherein X₂ is S.

66. The method according to claim 60, wherein X_2 is O.


67. The method according to claim 60, wherein R^0 is C1-C4 alkyl, optionally substituted with up to three R^1 .

68. The method according to claim 60, wherein R^0 is C1-C4 alkyl, optionally substituted with up to one R^1 .

69. The method according to claim 60, wherein R^0 is phenyl optionally substituted with C1-C4 alkyl, or R^1 .

70. The method according to claim 60, wherein R is hydrogen.

71. The method according to claim 1, wherein said compound has formula (IA):




wherein:

X is a bond, CH_2 , CHOR, C(O), NR, or O;

A is aliphatic, aryl, heteroaryl, heterocyclic, or cycloaliphatic;

Q is selected from:



each B is independently selected from 3-7 membered monocyclic or 8-14 membered bicyclic or tricyclic,

saturated, unsaturated or aromatic ring containing 0-4 heteroatoms in each ring, wherein each said heteroatom is independently selected from N, NH, S, or O;

R is H, R², or R⁶;

wherein each A and B is independently and optionally substituted with up to 4 substituents independently selected from R¹, R², R³, R⁴, or R⁵; and

R¹, R², R³, R⁴, or R⁵ are as defined in claim 1.

72. The method according to claim 71, wherein X is CH₂.

73. The method according to claim 71, X is a bond.

74. The method according to claim 71, wherein A is phenyl or a 5-6 membered heteroaryl, wherein A is optionally substituted with up to 3 substituents selected from R¹, R², R³, or R⁴.

75. The method according to claim 71, wherein A is optionally substituted phenyl.

76. The method according to claim 71, wherein Q is B.

77. The method according to claim 71, wherein Q is -(C1-C6)- aliphatic-B.


78. The method according to claim 71, wherein Q is CH(B)₂.

79. The method according to claim 71, wherein Q is $C(B)_3$.

80. The method according to claim 71, wherein B is phenyl.

81. The method according to claim 71, wherein:
 (i) X is a bond, CHOH, C(O), CH_2 , or O;
 (ii) A is optionally substituted phenyl; and
 (iii) Q is diphenylmethyl.

82. The method according to claim 1, wherein said compound has formula (IB):




wherein:

X is a bond, CH_2 , CHOR, C(O), NR, or O;

A is aliphatic, aryl, heteroaryl, heterocyclic, or cycloaliphatic;

Q is selected from:



each B is independently selected from 3-7 membered monocyclic or 8-14 membered bicyclic or tricyclic, saturated, unsaturated or aromatic ring containing 0-4 heteroatoms in each ring, wherein each said heteroatom is independently selected from N, NH, S, or O;

R is H, R^2 , or R^6 ;

wherein each A and B is independently and optionally substituted with up to 4 substituents independently selected from R¹, R², R³, R⁴, or R⁵.

83. The method according to claim 82, X is a bond or CH₂.

84. The method according to claim 82, wherein X is a bond.

85. The method according to claim 82, wherein A is phenyl or a 5-6 membered heteroaryl, wherein A is optionally substituted with up to 3 substituents selected from R¹, R², R³, or R⁴.

86. The method according to claim 82, wherein Q is B.

87. The method according to claim 82, wherein Q is -(C1-C6)- aliphatic-B.

88. The method according to claim 82, wherein Q is CH(B)₂.


89. The method according to claim 82, wherein Q is C(B)₃.

90. The method according to claim 82, wherein B is phenyl.

91. The method according to claim 82, wherein:
(i) X is a bond, CHOH, C(O), CH₂, or O;
(ii) A is optionally substituted phenyl; and

(iii) Q is diphenylmethyl.

92. The method according to claim 1, wherein said compound has formula (IC):




wherein:

each X is independently a bond, CH₂, CHOR, C(O), NR, or O;

A is aliphatic, aryl, heteroaryl, heterocyclic, or cycloaliphatic;

Q is selected from:



each B is independently selected from 3-7 membered monocyclic or 8-14 membered bicyclic or tricyclic, saturated, unsaturated or aromatic ring containing 0-4 heteroatoms in each ring, wherein each said heteroatom is independently selected from N, NH, S, or O;

R is H, R², or R⁶;

wherein each A and B is independently and optionally substituted with up to 4 substituents independently selected from R¹, R², R³, R⁴, or R⁵.

93. The method according to claim 92, X is a bond or CH₂.

94. The method according to claim 92, wherein X is CH_2 .

95. The method according to claim 92, wherein X is a bond.

96. The method according to claim 92, wherein A is phenyl or a 5-6 membered heteroaryl, wherein A is optionally substituted with up to 3 substituents selected from R^1 , R^2 , R^3 , or R^4 .

97. The method according to claim 92, wherein Q is B.

98. The method according to claim 92, wherein Q is - (C1-C6)- aliphatic-B.

99. The method according to claim 92, wherein Q is $\text{CH}(\text{B})_2$.

100. The method according to claim 92, wherein Q is $\text{C}(\text{B})_3$.

101. The method according to claim 92, wherein B is phenyl.

102. The method according to claim 92, wherein:


(i) each X is a bond or CH_2 ;

(ii) each A is optionally substituted phenyl, (C1-C6)aliphatic, or CF_3 ; and

(iii) Q is optionally substituted phenyl, (C1-C6)aliphatic, or diphenylmethyl.

103. The method according to claim 102, wherein X is CH₂.

104. A compound having formula (II):



or a pharmaceutically acceptable salt thereof;
wherein:

X₁ is a bond, O, S, CF₂, CHOR, C(O), C(O)O, CH₂, or NR;

R is H or R²

A₁ is (C₂-C₁₀) aliphatic, aryl, heteroaryl, heterocyclic, or cycloalkyl;

each B₁ is independently selected from 3-7 membered monocyclic, saturated, unsaturated or aromatic ring containing 0-4 heteroatoms selected from N, NH, S, or O;

wherein each A₁ is optionally substituted with up to 4 substituents independently selected from R¹, R², R³, R⁴, or R⁵;

R¹ is R⁶ or ((C₁-C₄)aliphatic)_n-Y;

n is 0 or 1;

Y is halo, CN, NO₂, CF₃, OCF₃, OH, SR⁶, S(O)R⁶, SO₂R⁶, NH₂, NHR⁶, N(R⁶)₂, NR⁶R⁸, COOH, COOR⁶ or OR⁶; or

two R¹ on adjacent ring atoms, taken together, form 1,2-methylenedioxy or 1,2-ethylenedioxy;

R² is aliphatic, wherein each R² optionally comprises up to 2 substituents independently selected from R¹, R⁴, or R⁵;

R^3 is a cycloaliphatic, aryl, heterocyclic, or heteroaryl ring optionally comprising up to 3 substituents, independently selected from R^1 , R^2 , R^4 or R^5 ;

R^4 is OR^5 , OR^6 , $OC(O)R^6$, $OC(O)R^5$, $OC(O)OR^6$, $OC(O)OR^5$, $OC(O)N(R^6)_2$, $OC(O)N(R^5)_2$, $OC(O)N(R^6R^5)$, SR^6 , SR^5 , $S(O)R^6$, $S(O)R^5$, SO_2R^6 , SO_2R^5 , $SO_2N(R^6)_2$, $SO_2N(R^5)_2$, $SO_2NR^5R^6$, SO_3R^6 , SO_3R^5 , $C(O)R^5$, $C(O)OR^5$, $C(O)R^6$, $C(O)OR^6$, $C(O)N(R^6)_2$, $C(O)N(R^5)_2$, $C(O)N(R^5R^6)$, $C(O)N(OR^6)R^6$, $C(O)N(OR^5)R^6$, $C(O)N(OR^6)R^5$, $C(O)N(OR^5)R^5$, $C(NOR^6)R^6$, $C(NOR^6)R^5$, $C(NOR^5)R^6$, $C(NOR^5)R^5$, $N(R^6)_2$, $N(R^5)_2$, $N(R^5R^6)$, $NR^5C(O)R^5$, $NR^6C(O)R^6$, $NR^6C(O)R^5$, $NR^5C(O)R^6$, $NR^6C(O)OR^6$, $NR^5C(O)OR^6$, $NR^6C(O)OR^5$, $NR^5C(O)OR^5$, $NR^6C(O)N(R^6)_2$, $NR^6C(O)NR^5R^6$, $NR^6C(O)N(R^5)_2$, $NR^5C(O)N(R^6)_2$, $NR^5C(O)NR^5R^6$, $NR^5C(O)N(R^5)_2$, $NR^6SO_2R^6$, $NR^6SO_2R^5$, $NR^5SO_2R^5$, $NR^5SO_2R^6$, $NR^6SO_2N(R^6)_2$, $NR^6SO_2NR^5R^6$, $NR^6SO_2N(R^5)_2$, $NR^5SO_2N(R^6)_2$, $NR^5SO_2NR^5R^6$, $NR^5SO_2N(R^5)_2$, $N(OR^6)R^6$, $N(OR^6)R^5$, $N(OR^5)R^5$, or $N(OR^5)R^6$;

R^5 is a cycloaliphatic, aryl, heterocyclic, or heteroaryl ring, optionally comprising up to 3 R^1 substituents;

R^6 is H or aliphatic, wherein R^6 optionally comprises a R^7 substituent;

R^7 is a cycloaliphatic, aryl, heterocyclic, or heteroaryl ring, and each R^7 optionally comprises up to 2 substituents independently chosen from H, (C_1-C_6) -straight or branched alkyl, (C_2-C_6) straight or branched alkenyl or alkynyl, 1,2-methylenedioxy, 1,2-ethylenedioxy, or $(CH_2)_n-Z$;

Z is selected from halo, CN, NO₂, CF₃, OCF₃, OH, S-aliphatic, S(O)-aliphatic, SO₂-aliphatic, NH₂, N-aliphatic, N(aliphatic)₂, N(aliphatic)R⁸, COOH, C(O)O(-aliphatic, or O-aliphatic; and

R⁸ is an amino protecting group;
provided that:

- (i) when both B₁ are simultaneously phenyl and X₁ is CH₂, then A is not 4-fluoro-phenyl, 4-phenyl-piperidyl, phenyl, 2,4-dichloro-phenyl, 4-methoxy-phenyl, 3,4-dichloro-phenyl, 2,5-dichloro-phenyl, 4-nitro-phenyl, 4-bromo-phenyl, 4-methyl-phenyl, 2-chloro-phenyl, 1-naphthyl, 3-trifluoromethyl-phenyl, 2,3-dichlorophenyl, N-morpholinyl, 4-chloro-phenyl, 3-chloro-phenyl, or 3-nitro-phenyl;
- (ii) when X₁ is a bond or CH₂, one B₁ is a substituted phenyl and the other B₁ is cycloaliphatic, then A₁ is not (C2-C8)aliphatic; and
- (iii) when X₁ is a bond, then A₁ is not an optionally substituted 6-membered heteroaryl ring with 1-3 nitrogen ring atoms.

105. The compound according to claim 104, wherein X₁ is CH₂, CHOR, C(O), C(O)O, CF₂, or O.

106. The compound according to claim 104, wherein X₁ is CH₂, CHOR, C(O), C(O)O.

107. The compound according to claim 104, wherein X₁ is CH₂.

108. The compound according to claim 104, wherein A₁ is selected from phenyl, triazinyl, pyrazinyl, pyrimidinyl, pyridazinyl, pyridyl, thiadiazolyl,

triazolyl, oxadiazolyl, isothiazolyl, pyrazolyl, imidazolyl, thiazolyl, oxazolyl, pyrrolyl, thienyl, furanyl, indoliziny, indolyl, isoindolyl, benzofuranyl, benzo[b]thienyl, 1H-indazolyl, benzimidazolyl, benzthiazolyl, purinyl, quinolinyl, isoquinolinyl, cinnolinyl, phthazinyl, quinazolinyl, quinoxalinyl, 1,8-naphthyridinyl, pteridinyl, carbazolyl, acridinyl, phenazinyl, phenothiazinyl, phenoxazinyl, indenyl, naphthyl, azuliny, or anthracenyl.

109. The compound according to claim 104, wherein each B₁ is independently selected from optionally substituted C6-C10 aryl.

110. The compound according to claim 104, wherein each B₁ is independently an optionally substituted phenyl or naphthyl.

111. The compound according to claim 104, wherein each B₁ is an unsubstituted phenyl.


112. The compound according to claim 104, wherein each B₁ is independently selected from optionally substituted C5-C12 heteroaryl.

113. The compound according to claim 112, wherein each B₁ is independently and optionally substituted C5-C7 heteroaryl.

114. The compound according to claim 113, wherein each B₁ is independently selected from optionally substituted pyrazolyl or imidazolyl.

115. The compound according to claim 104, wherein B₁ is selected from optionally substituted aziridine, oxirane, thiirane, pyrrolidyl, tetrahydrofuranyl, tetrahydrothienyl, dioxolanyl, pyrrolinyl, pyranyl, pyrazolinyl, pyrazolidinyl, piperidinyl, 1,4-dioxanyl, morpholinyl, 1,4-dithianyl, thiomorpholinyl, piperazinyl, 3H-indolyl, or indolyl.

116. A compound having formula (III):



or a pharmaceutically acceptable salt thereof;
wherein:

X₁ is a bond, O, S, CF₂, CHOR, C(O), C(O)O, CH₂, or NR;

R is H or R²

A₁ is (C2-C10) aliphatic, aryl, heteroaryl, heterocyclic, or cycloaliphatic;

each B₁ is independently selected from 3-7 membered monocyclic, saturated, unsaturated or aromatic ring containing 0-4 heteroatoms selected from N, NH, S, or O;

wherein each A₁ is optionally substituted with up to 4 substituents independently selected from R¹, R², R³, R⁴, or R⁵;

R¹ is R⁶ or ((C1-C4)aliphatic)_n-Y;

n is 0 or 1;

Y is halo, CN, NO₂, CF₃, OCF₃, OH, SR⁶, S(O)R⁶, SO₂R⁶, NH₂, NHR⁶, N(R⁶)₂, NR⁶R⁸, COOH, COOR⁶ or OR⁶; or

two R^1 on adjacent ring atoms, taken together, form 1,2-methylenedioxy or 1,2-ethylenedioxy;

R^2 is aliphatic, wherein each R^2 optionally comprises up to 2 substituents independently selected from R^1 , R^4 , or R^5 ;

R^3 is a cycloaliphatic, aryl, heterocyclic, or heteroaryl ring optionally comprising up to 3 substituents, independently selected from R^1 , R^2 , R^4 or R^5 ;

R^4 is OR^5 , OR^6 , $OC(O)R^6$, $OC(O)R^5$, $OC(O)OR^6$, $OC(O)OR^5$, $OC(O)N(R^6)_2$, $OC(O)N(R^5)_2$, $OC(O)N(R^6R^5)$, SR^6 , SR^5 , $S(O)R^6$, $S(O)R^5$, SO_2R^6 , SO_2R^5 , $SO_2N(R^6)_2$, $SO_2N(R^5)_2$, $SO_2NR^5R^6$, SO_3R^6 , SO_3R^5 , $C(O)R^5$, $C(O)OR^5$, $C(O)R^6$, $C(O)OR^6$, $C(O)N(R^6)_2$, $C(O)N(R^5)_2$, $C(O)N(R^5R^6)$, $C(O)N(OR^6)R^6$, $C(O)N(OR^5)R^6$, $C(O)N(OR^6)R^5$, $C(O)N(OR^5)R^5$, $C(NOR^6)R^6$, $C(NOR^6)R^5$, $C(NOR^5)R^6$, $C(NOR^5)R^5$, $N(R^6)_2$, $N(R^5)_2$, $N(R^5R^6)$, $NR^5C(O)R^5$, $NR^6C(O)R^6$, $NR^6C(O)R^5$, $NR^5C(O)R^6$, $NR^6C(O)OR^6$, $NR^5C(O)OR^6$, $NR^6C(O)OR^5$, $NR^5C(O)OR^5$, $NR^6C(O)N(R^6)_2$, $NR^6C(O)NR^5R^6$, $NR^6C(O)N(R^5)_2$, $NR^5C(O)N(R^6)_2$, $NR^5C(O)NR^5R^6$, $NR^5C(O)N(R^5)_2$, $NR^6SO_2R^6$, $NR^6SO_2R^5$, $NR^5SO_2R^5$, $NR^5SO_2R^6$, $NR^6SO_2N(R^6)_2$, $NR^6SO_2NR^5R^6$, $NR^6SO_2N(R^5)_2$, $NR^5SO_2N(R^6)_2$, $NR^5SO_2NR^5R^6$, $NR^5SO_2N(R^5)_2$, $N(OR^6)R^6$, $N(OR^6)R^5$, $N(OR^5)R^5$, or $N(OR^5)R^6$;

R^5 is a cycloaliphatic, aryl, heterocyclic, or heteroaryl ring, optionally comprising up to 3 R^1 substituents;


R^6 is H or aliphatic, wherein R^6 optionally comprises a R^7 substituent;

R^7 is a cycloaliphatic, aryl, heterocyclic, or heteroaryl ring, and each R^7 optionally comprises up to 2 substituents independently chosen from H, (C_1-C_6) -straight or branched alkyl, (C_2-C_6) straight or branched alkenyl or alkynyl, 1,2-methylenedioxy, 1,2-ethylenedioxy, or $(CH_2)_n-Z$;

Z is selected from halo, CN, NO_2 , CF_3 , OCF_3 , OH, S-aliphatic, S(O)-aliphatic, SO_2 -aliphatic, NH_2 , N-(aliphatic), $N(aliphatic)_2$, $N(aliphatic)R^8$, NHR^8 , $N(R^8)_2$, COOH, $C(O)O(-aliphatic)$, or O-aliphatic; and

R^8 is an amino protecting group; provided that:

(i) when X_1 is a bond, one B_1 is phenyl and the other B_1 is N-piperidyl, then A is not:



ii) when X_1 is a bond, then A_1 is not an optionally substituted 6-membered heteroaryl ring with 1-3 nitrogen ring atoms

117. The compound according to claim 116, wherein X_1 is a bond, CH_2 , CHOR, $C(O)$, $C(O)O$, CF_2 , or O.

118. The compound according to claim 116, wherein X_1 is bond, CH_2 , CHOR, $C(O)$, $C(O)O$.

119. The compound according to claim 116, wherein A_1 is optionally substituted (C_1-C_{10}) aliphatic.

120. The compound according to claim 119, wherein A₁ is optionally substituted methyl, ethyl, propyl, butyl, pentyl, or hexyl.

121. The compound according to claim 116, wherein A₁ is optionally substituted C6-C10 aryl ring.

122. The compound according to claim 121, wherein A₁ is optionally substituted phenyl or naphthyl.

123. The compound according to claim 116, wherein A₁ is optionally substituted C5-C12 heteroaryl ring.

124. The compound according to claim 123, wherein A is pyridinyl.

125. The compound according to claim 116, wherein each B₁ is independently selected from optionally substituted C6-C10 aryl.

126. The compound according to claim 125, wherein each B₁ is independently an optionally substituted phenyl or naphthyl.


127. The compound according to claim 126, wherein each B₁ is an unsubstituted phenyl.

128. The compound according to claim 116, wherein each B₁ is independently an optionally substituted 3-12 membered heterocyclic ring having up to 4 heteroatoms selected from O, S, or NR.

129. The compound according to claim 128, wherein each B₁ is independently selected from optionally

substituted aziridine, oxirane, thiirane, pyrrolidyl, tetrahydrofuranyl, tetrahydrothienyl, dioxolanyl, pyrrolinyl, pyranyl, pyrazolinyl, pyrazolidinyl, piperidinyl, 1,4-dioxanyl, morpholinyl, 1,4-dithianyl, thiomorpholinyl, piperazinyl, 3H-indolyl, or indolinyl.

130. A compound of formula IIA:



IIA;

or a pharmaceutically acceptable salt thereof;

wherein:

Y' is O or S;

B is a 3-8 membered, saturated, monocyclic, ring having 0-4 heteroatoms selected from O, S, or N; and

ring G and B are optionally substituted with up to four substituents independently selected from R¹, R², R³, R⁴, or R⁵;

R¹ is oxo, R⁶ or ((C1-C4)aliphatic)_n-Y;

n is 0 or 1;

Y is halo, CN, NO₂, CF₃, OCF₃, OH, SR⁶, S(O)R⁶, SO₂R⁶, NH₂, NHR⁶, N(R⁶)₂, NR⁶R⁸, N(R⁸)₂, COOH, COOR⁶ or OR⁶; or

two R¹ on adjacent ring atoms, taken together, form 1,2-methylenedioxy or 1,2-ethylenedioxy;

R² is aliphatic, wherein each R² optionally comprises up to 2 substituents independently selected from R¹, R⁴, or R⁵;

R³ is a cycloaliphatic, aryl, heterocyclic, or heteroaryl ring optionally comprising up to 3

substituents, independently selected from R^1 , R^2 , R^4 or R^5 ;

R^4 is OR^5 , OR^6 , $OC(O)R^6$, $OC(O)R^5$, $OC(O)OR^6$, $OC(O)OR^5$, $OC(O)N(R^6)_2$, $OC(O)N(R^5)_2$, $OC(O)N(R^6R^5)$, SR^6 , SR^5 , $S(O)R^6$, $S(O)R^5$, SO_2R^6 , SO_2R^5 , $SO_2N(R^6)_2$, $SO_2N(R^5)_2$, $SO_2NR^5R^6$, SO_3R^6 , SO_3R^5 , $C(O)R^5$, $C(O)OR^5$, $C(O)R^6$, $C(O)OR^6$, $C(O)N(R^6)_2$, $C(O)N(R^5)_2$, $C(O)N(R^5R^6)$, $C(O)N(OR^6)R^6$, $C(O)N(OR^5)R^6$, $C(O)N(OR^6)R^5$, $C(O)N(OR^5)R^5$, $C(NOR^6)R^6$, $C(NOR^6)R^5$, $C(NOR^5)R^6$, $C(NOR^5)R^5$, $N(R^6)_2$, $N(R^5)_2$, $N(R^5R^6)$, $NR^5C(O)R^5$, $NR^6C(O)R^6$, $NR^6C(O)R^5$, $NR^5C(O)R^6$, $NR^6C(O)OR^6$, $NR^5C(O)OR^6$, $NR^6C(O)OR^5$, $NR^5C(O)OR^5$, $NR^6C(O)N(R^6)_2$, $NR^6C(O)NR^5R^6$, $NR^6C(O)N(R^5)_2$, $NR^5C(O)N(R^6)_2$, $NR^5C(O)NR^5R^6$, $NR^5C(O)N(R^5)_2$, $NR^6SO_2R^6$, $NR^6SO_2R^5$, $NR^5SO_2R^5$, $NR^5SO_2R^6$, $NR^6SO_2N(R^6)_2$, $NR^6SO_2NR^5R^6$, $NR^6SO_2N(R^5)_2$, $NR^5SO_2N(R^6)_2$, $NR^5SO_2NR^5R^6$, $NR^5SO_2N(R^5)_2$, $N(OR^6)R^6$, $N(OR^6)R^5$, $N(OR^5)R^5$, or $N(OR^5)R^6$;

R^5 is a cycloaliphatic, aryl, heterocyclic, or heteroaryl ring, optionally comprising up to 3 R^1 substituents;

R^6 is H or aliphatic, wherein R^6 optionally comprises a R^7 substituent;

R^7 is a cycloaliphatic, aryl, heterocyclic, or heteroaryl ring, and each R^7 optionally comprises up to 2 substituents independently chosen from H, (C_1-C_6) -straight or branched alkyl, (C_2-C_6) straight or branched alkenyl or alkynyl, 1,2-methylenedioxy, 1,2-ethylenedioxy, or $(CH_2)_n-Z$;

Z is selected from halo, CN, NO_2 , CF_3 , OCF_3 , OH, S-aliphatic, S(O)-aliphatic, SO_2 -aliphatic, NH_2 , NH-

aliphatic, $N(\text{aliphatic})_2$, $N(\text{aliphatic})R^8$, NHR^8 , $N(R^8)_2$,
COOH, $C(O)O(-\text{aliphatic})$, or O-aliphatic; and

R^8 is an amino capping group;

provided that when Y' is S, and:

a) when B is cyclohexyl, tetrahydrofuran-2-yl, or cyclopropyl, and ring G has 1-3 halo substituents, then ring G has at least one additional substituent other than halo; and

b) when B is tetrahydrofuran-2-yl, then ring G is not phenyl, trifluoromethylphenyl, methoxyphenyl, or tolyl;

c) when B is cyclohexyl, then ring G is not phenyl or trifluoromethylphenyl.

131. The compound according to claim 130, wherein B is tetrahydrofuranyl, piperidyl, morpholinyl, or thiomorpholinyl.

132. The compound according to claim 130, wherein B is C3-C8 saturated, carbocyclic, monocyclic ring.

133. The compound according to claim 130, wherein B is selected from cyclopropyl, cyclopentyl, cyclohexyl, or cycloheptyl.


134. The compound according to claim 130, wherein ring G is phenyl optionally substituted with up to two R^1 .

135. The compound according to claim 134, wherein ring G is optionally substituted with up to two substituents selected from halo, cyano, C1-C4 alkyl, or O-(C1-C4 alkyl).

136. The compound according to claim 135, wherein said compound has one or more of the following features:

- a) Y' is S;
- b) ring G is halo-substituted phenyl;
- c) B is phenyl optionally substituted with halo, cyano, C1-C4 alkyl, or O-(C1-C4 alkyl).

137. A compound formula IIB:



IIB;

or a pharmaceutically acceptable salt thereof;

wherein:

Y' is S or O;

R is H, R², or R⁶;

R^B is C1-C6 aliphatic or a 3-7 membered monocyclic or 8-14 membered bicyclic or tricyclic, saturated, unsaturated or aromatic ring containing 0-4 heteroatoms in each ring, wherein each said heteroatom is independently selected from N, NH, S, or O;

wherein each of ring G, ring H, and R^B is independently and optionally substituted with up to 4 substituents selected from R¹, R², R³, R⁴, or R⁵;

R¹ is oxo, R⁶ or ((C1-C4)aliphatic)_n-Y;

n is 0 or 1;

Y is halo, CN, NO₂, CF₃, OCF₃, OH, SR⁶, S(O)R⁶, SO₂R⁶, NH₂, NHR⁶, N(R⁶)₂, NR⁶R⁸, N(R⁸)₂, COOH, COOR⁶ or OR⁶; or

two R¹ on adjacent ring atoms, taken together, form 1,2-methylenedioxy or 1,2-ethylenedioxy;

R^2 is aliphatic, wherein each R^2 optionally comprises up to 2 substituents independently selected from R^1 , R^4 , or R^5 ;

R^3 is a cycloaliphatic, aryl, heterocyclic, or heteroaryl ring optionally comprising up to 3 substituents, independently selected from R^1 , R^2 , R^4 or R^5 ;

R^4 is OR^5 , OR^6 , $OC(O)R^6$, $OC(O)R^5$, $OC(O)OR^6$, $OC(O)OR^5$, $OC(O)N(R^6)_2$, $OC(O)N(R^5)_2$, $OC(O)N(R^6R^5)$, SR^6 , SR^5 , $S(O)R^6$, $S(O)R^5$, SO_2R^6 , SO_2R^5 , $SO_2N(R^6)_2$, $SO_2N(R^5)_2$, $SO_2NR^5R^6$, SO_3R^6 , SO_3R^5 , $C(O)R^5$, $C(O)OR^5$, $C(O)R^6$, $C(O)OR^6$, $C(O)N(R^6)_2$, $C(O)N(R^5)_2$, $C(O)N(R^5R^6)$, $C(O)N(OR^6)R^6$, $C(O)N(OR^5)R^6$, $C(O)N(OR^6)R^5$, $C(O)N(OR^5)R^5$, $C(NOR^6)R^6$, $C(NOR^6)R^5$, $C(NOR^5)R^6$, $C(NOR^5)R^5$, $N(R^6)_2$, $N(R^5)_2$, $N(R^5R^6)$, $NR^5C(O)R^5$, $NR^6C(O)R^6$, $NR^6C(O)R^5$, $NR^5C(O)R^6$, $NR^6C(O)OR^6$, $NR^5C(O)OR^6$, $NR^6C(O)OR^5$, $NR^5C(O)OR^5$, $NR^6C(O)N(R^6)_2$, $NR^6C(O)NR^5R^6$, $NR^6C(O)N(R^5)_2$, $NR^5C(O)N(R^6)_2$, $NR^5C(O)NR^5R^6$, $NR^5C(O)N(R^5)_2$, $NR^6SO_2R^6$, $NR^6SO_2R^5$, $NR^5SO_2R^5$, $NR^5SO_2R^6$, $NR^6SO_2N(R^6)_2$, $NR^6SO_2NR^5R^6$, $NR^6SO_2N(R^5)_2$, $NR^5SO_2N(R^6)_2$, $NR^5SO_2NR^5R^6$, $NR^5SO_2N(R^5)_2$, $N(OR^6)R^6$, $N(OR^6)R^5$, $N(OR^5)R^5$, or $N(OR^5)R^6$;

R^5 is a cycloaliphatic, aryl, heterocyclic, or heteroaryl ring, optionally comprising up to 3 R^1 substituents;

R^6 is H or aliphatic, wherein R^6 optionally comprises a R^7 substituent;

R^7 is a cycloaliphatic, aryl, heterocyclic, or heteroaryl ring, and each R^7 optionally comprises up to 2 substituents independently chosen from H, $(C_1-C_6)-$

straight or branched alkyl, (C₂-C₆) straight or branched alkenyl or alkynyl, 1,2-methylenedioxy, 1,2-ethylenedioxy, or (CH₂)_n-Z;

Z is selected from halo, CN, NO₂, CF₃, OCF₃, OH, S-aliphatic, S(O)-aliphatic, SO₂-aliphatic, NH₂, NH-aliphatic, N(aliphatic)₂, N(aliphatic)R⁸, NHR⁸, N(R⁸)₂, COOH, C(O)O(-aliphatic, or O-aliphatic; and

R⁸ is an amino capping group;
provided that when Y is S, and:

a) when R^B is hydrogen, and ring G and ring H both have 1-3 halo substituents, then at least one of ring G and ring H has an additional substituent other than halo;

b) when R^B is hydrogen and ring H is unsubstituted phenyl, then ring G is not phenyl or phenyl substituted with methyl, CF₃, -OMe, NO₂, or 1-3 halo;

c) when R^B is hydrogen and ring H is phenyl with methyl, 1-2 methoxy or 1-2 halo substituents, then ring G is not phenyl substituted with CF₃ or 1-2 halo;

d) when R^B is methyl and ring H phenyl substituted with butyl, then ring G is not phenyl substituted with methyl, or 1-2 halo; and

e) when R^B and ring H are both unsubstituted phenyl, then ring G is not unsubstituted phenyl, or phenyl substituted with methyl, CF₃, OMe, NO₂, or 1-2 halo.

138. The compound according to claim 137, wherein ring G is phenyl optionally substituted with up to two R¹.

139. The compound according to claim 138, wherein R¹ is selected from C1-C4 alkyl, O-(C1-C4 alkyl), halo, or cyano.

140. The compound according to claim 137, wherein R^B is C1-6 aliphatic optionally substituted with up to 4 substituents selected from R¹, R², R³, R⁴, or R⁵.

141. The compound according to claim 140, wherein R^B is C1-C4 alkyl optionally substituted with up to 2 substituents selected from R¹.

142. The compound according to claim 141, wherein R^B is selected from methyl, ethyl, n-propyl, isopropyl, sec-butyl, n-butyl, or t-butyl.

143. The compound according to claim 137, wherein R^B is a 3-7 membered monocyclic saturated, unsaturated or aromatic ring containing 0-4 heteroatoms optionally substituted with up to 4 substituents selected from R¹, R², R³, R⁴, or R⁵.

144. The compound according to claim 143, wherein R^B is a 3-7 membered monocyclic saturated, carbocyclic ring optionally substituted with up to 2 substituents selected from R¹.

145. The compound according to claim 144, wherein R^B is selected from cyclopropyl, cyclopentyl, cyclohexyl, or cycloheptyl.


146. The compound according to claim 143, wherein R^B is a 3-7 membered monocyclic saturated, unsaturated or aromatic ring containing 1-3 heteroatoms optionally substituted with up to 4 substituents selected from R¹, R², R³, R⁴, or R⁵.

147. The compound according to claim 143, R^B is a 3-7 membered monocyclic saturated ring containing 1-3 heteroatoms optionally substituted with up to 2 substituents selected from R^1 .

148. The compound according to claim 147, wherein R^B is selected from piperidinyl, morpholinyl, or thiomorpholinyl.

149. The compound according to claim 137, wherein R^B is a 8-14 membered bicyclic or tricyclic, saturated, unsaturated or aromatic ring containing 0-4 heteroatoms in each ring, wherein each said heteroatom is independently selected from N, NH, S, or O, optionally substituted with up to 4 substituents selected from R^1 , R^2 , R^3 , R^4 , or R^5 .

150. A compound of formula IIC:



IIC;

wherein:

Y' is O or S;

X_7 is O, S, or NR' ;

R' is hydrogen, R^2 , or R^6 ;

R is hydrogen, R^2 , or R^6 ;

R^B is C1-6 aliphatic or a 3-7 membered monocyclic or 8-14 membered bicyclic or tricyclic, saturated, unsaturated or aromatic ring containing 0-4 heteroatoms in each ring, wherein each said heteroatom is independently selected from N, NH, S, or O;

R^C is C1-C6 aliphatic;

wherein each of ring G, R^B , and R^C is independently and optionally substituted with up to 4 substituents selected from R^1 , R^2 , R^3 , R^4 , or R^5 ;

R^1 is oxo, R^6 or ((C1-C4)aliphatic) $_n$ -Y;

n is 0 or 1;

Y is halo, CN, NO₂, CF₃, OCF₃, OH, SR⁶, S(O)R⁶, SO₂R⁶, NH₂, NHR⁶, N(R⁶)₂, NR⁶R⁸, N(R⁸)₂, COOH, COOR⁶ or OR⁶; or

two R^1 on adjacent ring atoms, taken together, form 1,2-methylenedioxy or 1,2-ethylenedioxy;

R^2 is aliphatic, wherein each R^2 optionally comprises up to 2 substituents independently selected from R^1 , R^4 , or R^5 ;

R^3 is a cycloaliphatic, aryl, heterocyclic, or heteroaryl ring optionally comprising up to 3 substituents, independently selected from R^1 , R^2 , R^4 or R^5 ;

R^4 is OR⁵, OR⁶, OC(O)R⁶, OC(O)R⁵, OC(O)OR⁶, OC(O)OR⁵, OC(O)N(R⁶)₂, OC(O)N(R⁵)₂, OC(O)N(R⁶R⁵), SR⁶, SR⁵, S(O)R⁶, S(O)R⁵, SO₂R⁶, SO₂R⁵, SO₂N(R⁶)₂, SO₂N(R⁵)₂, SO₂NR⁵R⁶, SO₃R⁶, SO₃R⁵, C(O)R⁵, C(O)OR⁵, C(O)R⁶, C(O)OR⁶, C(O)N(R⁶)₂, C(O)N(R⁵)₂, C(O)N(R⁵R⁶), C(O)N(OR⁶)R⁶, C(O)N(OR⁵)R⁶, C(O)N(OR⁶)R⁵, C(O)N(OR⁵)R⁵, C(NOR⁶)R⁶, C(NOR⁶)R⁵, C(NOR⁵)R⁶, C(NOR⁵)R⁵, N(R⁶)₂, N(R⁵)₂, N(R⁵R⁶), NR⁵C(O)R⁵, NR⁶C(O)R⁶, NR⁶C(O)R⁵, NR⁵C(O)R⁶, NR⁶C(O)OR⁶, NR⁵C(O)OR⁶, NR⁶C(O)OR⁵, NR⁵C(O)OR⁵, NR⁶C(O)N(R⁶)₂, NR⁶C(O)NR⁵R⁶, NR⁶C(O)N(R⁵)₂, NR⁵C(O)N(R⁶)₂, NR⁵C(O)NR⁵R⁶, NR⁵C(O)N(R⁵)₂, NR⁶SO₂R⁶, NR⁶SO₂R⁵, NR⁵SO₂R⁵, NR⁵SO₂R⁶, NR⁶SO₂N(R⁶)₂, NR⁶SO₂NR⁵R⁶, NR⁶SO₂N(R⁵)₂, NR⁵SO₂N(R⁶)₂,

$\text{NR}^5\text{SO}_2\text{NR}^5\text{R}^6$, $\text{NR}^5\text{SO}_2\text{N}(\text{R}^5)_2$, $\text{N}(\text{OR}^6)\text{R}^6$, $\text{N}(\text{OR}^6)\text{R}^5$, $\text{N}(\text{OR}^5)\text{R}^5$,
or $\text{N}(\text{OR}^5)\text{R}^6$;

R^5 is a cycloaliphatic, aryl, heterocyclic, or heteroaryl ring, optionally comprising up to 3 R^1 substituents;

R^6 is H or aliphatic, wherein R^6 optionally comprises a R^7 substituent;

R^7 is a cycloaliphatic, aryl, heterocyclic, or heteroaryl ring, and each R^7 optionally comprises up to 2 substituents independently chosen from H, $(\text{C}_1\text{-C}_6)$ -straight or branched alkyl, $(\text{C}_2\text{-C}_6)$ straight or branched alkenyl or alkynyl, 1,2-methylenedioxy, 1,2-ethylenedioxy, or $(\text{CH}_2)_n\text{-Z}$;

Z is selected from halo, CN, NO_2 , CF_3 , OCF_3 , OH, S-aliphatic, S(O)-aliphatic, SO_2 -aliphatic, NH_2 , NH-aliphatic, $\text{N}(\text{aliphatic})_2$, $\text{N}(\text{aliphatic})\text{R}^8$, NHR^8 , $\text{N}(\text{R}^8)_2$, COOH, $\text{C}(\text{O})\text{O}(-\text{aliphatic})$, or O-aliphatic; and

R^8 is an amino capping group.

151. The compound according to claim 150, wherein X_7 is O.

152. The compound according to claim 150, wherein X_7 is S.

153. The compound according to claim 150, wherein X_7 is NR' .

154. The compound according to claim 150, wherein R^c is $\text{C}_1\text{-C}_6$ alkyl, optionally substituted with up to two substituents selected from R^1 , R^2 , R^3 , R^4 , or R^5 .

155. The compound according to claim 154, wherein R^C is C1-C6 alkyl.

156. The compound according to claim 155, wherein R^C is selected from methyl, ethyl, isopropyl, n-propyl, n-butyl, sec-butyl, or t-butyl.

157. The compound according to claim 150, wherein R^B is phenyl optionally substituted with up to two R^1 substituents.


158. The compound according to claim 157, wherein R^B is phenyl.

159. The compound according to claim 150, wherein R^B is selected from methyl, ethyl, isopropyl, n-propyl, n-butyl, sec-butyl, or t-butyl.

160. The compound according to claim 150, wherein said compound has one or more of the following features:

- a) ring G is optionally substituted with one R^1 substituent;
- b) Y' is S and R is hydrogen;
- c) R^C is C1-C4 alkyl;
- d) X_7 is NH or NR' wherein R' is C1-C4 alkyl; and
- e) R^B is C1-C4 alkyl.

161. A compound of formula IID:



IID;

wherein:

Y' is O or S;

R is hydrogen or R²;

X₈ is O, S, or NR';

R' is hydrogen, R², or R⁶;

R^{BB} is C1-6 aliphatic or a 3-7 membered monocyclic or 8-14 membered bicyclic or tricyclic, saturated, unsaturated or aromatic ring containing 0-4 heteroatoms in each ring, wherein each said heteroatom is independently selected from N, S, or O;

wherein each of ring G and R^{BB} is independently and optionally substituted with up to 4 substituents selected from R¹, R², R³, R⁴, or R⁵;

R¹ is oxo, R⁶ or ((C1-C4)aliphatic)_n-Y;

n is 0 or 1;

Y is halo, CN, NO₂, CF₃, OCF₃, OH, SR⁶, S(O)R⁶, SO₂R⁶, NH₂, NHR⁶, N(R⁶)₂, NR⁶R⁸, N(R⁸)₂, COOH, COOR⁶ or OR⁶; or

two R¹ on adjacent ring atoms, taken together, form 1,2-methylenedioxy or 1,2-ethylenedioxy;

R² is aliphatic, wherein each R² optionally comprises up to 2 substituents independently selected from R¹, R⁴, or R⁵;

R³ is a cycloaliphatic, aryl, heterocyclic, or heteroaryl ring optionally comprising up to 3 substituents, independently selected from R¹, R², R⁴ or R⁵;

R⁴ is OR⁵, OR⁶, OC(O)R⁶, OC(O)R⁵, OC(O)OR⁶, OC(O)OR⁵, OC(O)N(R⁶)₂, OC(O)N(R⁵)₂, OC(O)N(R⁶R⁵), SR⁶, SR⁵, S(O)R⁶, S(O)R⁵, SO₂R⁶, SO₂R⁵, SO₂N(R⁶)₂, SO₂N(R⁵)₂, SO₂NR⁵R⁶, SO₃R⁶, SO₃R⁵, C(O)R⁵, C(O)OR⁵, C(O)R⁶, C(O)OR⁶,

$C(O)N(R^6)_2$, $C(O)N(R^5)_2$, $C(O)N(R^5R^6)$, $C(O)N(OR^6)R^6$,
 $C(O)N(OR^5)R^6$, $C(O)N(OR^6)R^5$, $C(O)N(OR^5)R^5$, $C(NOR^6)R^6$,
 $C(NOR^6)R^5$, $C(NOR^5)R^6$, $C(NOR^5)R^5$, $N(R^6)_2$, $N(R^5)_2$, $N(R^5R^6)$,
 $NR^5C(O)R^5$, $NR^6C(O)R^6$, $NR^6C(O)R^5$, $NR^5C(O)R^6$, $NR^6C(O)OR^6$,
 $NR^5C(O)OR^6$, $NR^6C(O)OR^5$, $NR^5C(O)OR^5$, $NR^6C(O)N(R^6)_2$,
 $NR^6C(O)NR^5R^6$, $NR^6C(O)N(R^5)_2$, $NR^5C(O)N(R^6)_2$, $NR^5C(O)NR^5R^6$,
 $NR^5C(O)N(R^5)_2$, $NR^6SO_2R^6$, $NR^6SO_2R^5$, $NR^5SO_2R^5$, $NR^5SO_2R^6$,
 $NR^6SO_2N(R^6)_2$, $NR^6SO_2NR^5R^6$, $NR^6SO_2N(R^5)_2$, $NR^5SO_2N(R^6)_2$,
 $NR^5SO_2NR^5R^6$, $NR^5SO_2N(R^5)_2$, $N(OR^6)R^6$, $N(OR^6)R^5$, $N(OR^5)R^5$,
 or $N(OR^5)R^6$;

R^5 is a cycloaliphatic, aryl, heterocyclic, or heteroaryl ring, optionally comprising up to 3 R^1 substituents;

R^6 is H or aliphatic, wherein R^6 optionally comprises a R^7 substituent;

R^7 is a cycloaliphatic, aryl, heterocyclic, or heteroaryl ring, and each R^7 optionally comprises up to 2 substituents independently chosen from H, (C_1-C_6) -straight or branched alkyl, (C_2-C_6) straight or branched alkenyl or alkynyl, 1,2-methylenedioxy, 1,2-ethylenedioxy, or $(CH_2)_n-Z$;

Z is selected from halo, CN, NO_2 , CF_3 , OCF_3 , OH, S-aliphatic, S(O)-aliphatic, SO_2 -aliphatic, NH_2 , NH-(aliphatic), $N(aliphatic)_2$, $N(aliphatic)R^8$, NHR^8 , $N(R^8)_2$, COOH, $C(O)O(-aliphatic)$, or O-aliphatic; and

R^8 is an amino capping group.

162. The compound according to claim 161, wherein X_8 is O.

163. The compound according to claim 161, wherein X_8 is S.

164. The compound according to claim 161, wherein X_8 is NR' .

165. The compound according to claim 161, wherein R^{BB} is phenyl optionally substituted with up to two R^1 substituents.

166. The compound according to claim 161, wherein R^{BB} is phenyl.

167. The compound according to claim 161, wherein R^{BB} is selected from methyl, ethyl, isopropyl, n-propyl, n-butyl, sec-butyl, or t-butyl.

168. The compound according to claim 161, wherein R^{BB} is an optionally substituted C3-C8 cycloalkyl.

169. The compound according to claim 168, wherein R^{BB} is selected from cyclopropyl, cyclopentyl, or cyclohexyl.


170. The compound according to claim 161, wherein R^{BB} is optionally substituted benzyl.

171. The compound according to claim 161, wherein said compound has one or more of the following features:

- a) Y' is S and R is hydrogen;
- b) each ring G is unsubstituted phenyl;
- d) X_8 is NR' , and R' is hydrogen or C1-C4 alkyl; and

e) R^{BB} is C1-C4 alkyl, benzyl, cyclopentyl, or cyclohexyl.

172. A compound of formula IV:



IV;

or a pharmaceutically acceptable salt thereof;

wherein:

Y' is O, S, or NR;


p is 0-2;

X is a bond, O, S, S(O), S(O)₂, CF₂, CH₂, -CHOR-, -C(O)-, -O-C(O)-, -C(O)-O, -C(O)-NR, -NR-C(O)-, -NR-C(O)-O-, -O-C(O)-NR-, -NR-C(O)-NR-, or NR;

R is H, R², or R⁶;

A is aliphatic, aryl, heteroaryl, heterocyclic, or cycloalkyl;

B is selected from:



wherein:

X_3 is O or S;

wherein A and B each is independently and optionally substituted with up to 4 substituents selected from R¹, R², R³, R⁴ or R⁵;

R¹ is oxo, R⁶ or ((C1-C4)aliphatic)_n-Y;

n is 0 or 1;

Y is halo, CN, NO₂, CF₃, OCF₃, OH, SR⁶, S(O)R⁶, SO₂R⁶, NH₂, NHR⁶, N(R⁶)₂, NR⁶R⁸, N(R⁸)₂, COOH, COOR⁶ or OR⁶; or

two R¹ on adjacent ring atoms, taken together, form 1,2-methylenedioxy or 1,2-ethylenedioxy;

R² is aliphatic, wherein each R² optionally comprises up to 2 substituents independently selected from R¹, R⁴, or R⁵;

R³ is a cycloaliphatic, aryl, heterocyclic, or heteroaryl ring optionally comprising up to 3 substituents, independently selected from R¹, R², R⁴ or R⁵;

R⁴ is OR⁵, OR⁶, OC(O)R⁶, OC(O)R⁵, OC(O)OR⁶, OC(O)OR⁵, OC(O)N(R⁶)₂, OC(O)N(R⁵)₂, OC(O)N(R⁶R⁵), SR⁶, SR⁵, S(O)R⁶, S(O)R⁵, SO₂R⁶, SO₂R⁵, SO₂N(R⁶)₂, SO₂N(R⁵)₂, SO₂NR⁵R⁶, SO₃R⁶, SO₃R⁵, C(O)R⁵, C(O)OR⁵, C(O)R⁶, C(O)OR⁶, C(O)N(R⁶)₂, C(O)N(R⁵)₂, C(O)N(R⁵R⁶), C(O)N(OR⁶)R⁶, C(O)N(OR⁵)R⁶, C(O)N(OR⁶)R⁵, C(O)N(OR⁵)R⁵, C(NOR⁶)R⁶, C(NOR⁶)R⁵, C(NOR⁵)R⁶, C(NOR⁵)R⁵, N(R⁶)₂, N(R⁵)₂, N(R⁵R⁶), NR⁵C(O)R⁵, NR⁶C(O)R⁶, NR⁶C(O)R⁵, NR⁵C(O)R⁶, NR⁶C(O)OR⁶, NR⁵C(O)OR⁶, NR⁶C(O)OR⁵, NR⁵C(O)OR⁵, NR⁶C(O)N(R⁶)₂, NR⁶C(O)NR⁵R⁶, NR⁶C(O)N(R⁵)₂, NR⁵C(O)N(R⁶)₂, NR⁵C(O)NR⁵R⁶, NR⁵C(O)N(R⁵)₂, NR⁶SO₂R⁶, NR⁶SO₂R⁵, NR⁵SO₂R⁵, NR⁵SO₂R⁶, NR⁶SO₂N(R⁶)₂, NR⁶SO₂NR⁵R⁶, NR⁶SO₂N(R⁵)₂, NR⁵SO₂N(R⁶)₂, NR⁵SO₂NR⁵R⁶, NR⁵SO₂N(R⁵)₂, N(OR⁶)R⁶, N(OR⁶)R⁵, N(OR⁵)R⁵, or N(OR⁵)R⁶;

R⁵ is a cycloaliphatic, aryl, heterocyclic, or heteroaryl ring, optionally comprising up to 3 R¹ substituents;

R^6 is H or aliphatic, wherein R^6 optionally comprises a R^7 substituent;

R^7 is a cycloaliphatic, aryl, heterocyclic, or heteroaryl ring, and each R^7 optionally comprises up to 2 substituents independently chosen from H, (C_1-C_6) -straight or branched alkyl, (C_2-C_6) straight or branched alkenyl or alkynyl, 1,2-methylenedioxy, 1,2-ethylenedioxy, or $(CH_2)_n-Z$;

Z is selected from halo, CN, NO_2 , CF_3 , OCF_3 , OH, S-aliphatic, S(O)-aliphatic, SO_2 -aliphatic, NH_2 , NH-aliphatic, $N(aliphatic)_2$, $N(aliphatic)R^8$, NHR^8 , $N(R^8)_2$, COOH, $C(O)O(-aliphatic)$, or O-aliphatic; and

R^8 is an amino-capping group.


173. The compound according to claim 172, wherein X_3 is O.

174. The compound according to claim 172, wherein R is hydrogen.

175. The compound according to claim 172, wherein Y' is S.

176. The compound according to claim 172, wherein Y' is O.

177. A compound of formula V:



or a pharmaceutically acceptable salt thereof;
wherein:

Y' is O or S;


p is 0-2;

X is a bond, O, S, S(O), S(O)₂, CF₂, CH₂, -CHOR-, -C(O)-, -O-C(O)-, -C(O)-O-, -C(O)-NR-, -NR-C(O)-, -NR-C(O)-O-, -O-C(O)-NR-, -NR-C(O)-NR-, or NR;


R is H, R², or R⁶;

A is aliphatic, aryl, heteroaryl, heterocyclic, or cycloalkyl;


B is selected from:



(i)



(ii)



(iii);

wherein:

Ak is C1-C6 alkylidene;

X₄ is CH₂, O or S;

Ar' is phenyl optionally substituted with up to two R¹;

B is optionally substituted with up to two R¹;

wherein A is independently and optionally substituted with up to 4 substituents selected from R¹, R², R³, R⁴ or R⁵;

R¹ is oxo, R⁶ or ((C1-C4)aliphatic)_n-Y;

n is 0 or 1;

Y is halo, CN, NO₂, CF₃, OCF₃, OH, SR⁶, S(O)R⁶, SO₂R⁶, NH₂, NHR⁶, N(R⁶)₂, NR⁶R⁸, N(R⁸)₂, COOH, COOR⁶ or OR⁶; or

two R¹ on adjacent ring atoms, taken together, form 1,2-methylenedioxy or 1,2-ethylenedioxy;

R² is aliphatic, wherein each R² optionally comprises up to 2 substituents independently selected from R¹, R⁴, or R⁵;

R^3 is a cycloaliphatic, aryl, heterocyclic, or heteroaryl ring optionally comprising up to 3 substituents, independently selected from R^1 , R^2 , R^4 or R^5 ;

R^4 is OR^5 , OR^6 , $OC(O)R^6$, $OC(O)R^5$, $OC(O)OR^6$, $OC(O)OR^5$, $OC(O)N(R^6)_2$, $OC(O)N(R^5)_2$, $OC(O)N(R^6R^5)$, SR^6 , SR^5 , $S(O)R^6$, $S(O)R^5$, SO_2R^6 , SO_2R^5 , $SO_2N(R^6)_2$, $SO_2N(R^5)_2$, $SO_2NR^5R^6$, SO_3R^6 , SO_3R^5 , $C(O)R^5$, $C(O)OR^5$, $C(O)R^6$, $C(O)OR^6$, $C(O)N(R^6)_2$, $C(O)N(R^5)_2$, $C(O)N(R^5R^6)$, $C(O)N(OR^6)R^6$, $C(O)N(OR^5)R^6$, $C(O)N(OR^6)R^5$, $C(O)N(OR^5)R^5$, $C(NOR^6)R^6$, $C(NOR^6)R^5$, $C(NOR^5)R^6$, $C(NOR^5)R^5$, $N(R^6)_2$, $N(R^5)_2$, $N(R^5R^6)$, $NR^5C(O)R^5$, $NR^6C(O)R^6$, $NR^6C(O)R^5$, $NR^5C(O)R^6$, $NR^6C(O)OR^6$, $NR^5C(O)OR^6$, $NR^6C(O)OR^5$, $NR^5C(O)OR^5$, $NR^6C(O)N(R^6)_2$, $NR^6C(O)NR^5R^6$, $NR^6C(O)N(R^5)_2$, $NR^5C(O)N(R^6)_2$, $NR^5C(O)NR^5R^6$, $NR^5C(O)N(R^5)_2$, $NR^6SO_2R^6$, $NR^6SO_2R^5$, $NR^5SO_2R^5$, $NR^5SO_2R^6$, $NR^6SO_2N(R^6)_2$, $NR^6SO_2NR^5R^6$, $NR^6SO_2N(R^5)_2$, $NR^5SO_2N(R^6)_2$, $NR^5SO_2NR^5R^6$, $NR^5SO_2N(R^5)_2$, $N(OR^6)R^6$, $N(OR^6)R^5$, $N(OR^5)R^5$, or $N(OR^5)R^6$;

R^5 is a cycloaliphatic, aryl, heterocyclic, or heteroaryl ring, optionally comprising up to 3 R^1 substituents;

R^6 is H or aliphatic, wherein R^6 optionally comprises a R^7 substituent;

R^7 is a cycloaliphatic, aryl, heterocyclic, or heteroaryl ring, and each R^7 optionally comprises up to 2 substituents independently chosen from H, (C_1-C_6) -straight or branched alkyl, (C_2-C_6) straight or branched alkenyl or alkynyl, 1,2-methylenedioxy, 1,2-ethylenedioxy, or $(CH_2)_n-Z$;

Z is selected from halo, CN, NO₂, CF₃, OCF₃, OH, S-aliphatic, S(O)-aliphatic, SO₂-aliphatic, NH₂, NH-aliphatic, N(aliphatic)₂, N(aliphatic)R⁸, NHR⁸, N(R⁸)₂, COOH, C(O)O(-aliphatic, or O-aliphatic; and

R⁸ is an amino capping group.

178. The compound according to claim 177, wherein Ak is selected from CH₂, CH(CH₃), C(CH₃)₂, CH(Et), C(Et)₂, CH(n-propyl), CH(i-Pr), CH(n-butyl), CH(but-2-yl), or CH(t-butyl).

179. The compound according to claim 177, wherein Ar' is phenyl optionally substituted with halo, C1-C4 alkyl, or O-(C1-C4 alkyl).

180. The compound according to claim 177, wherein X₄ is CH₂.

181. The compound according to claim 177, wherein X₄ is O or S.


182. The compound according to claim 177, wherein R is hydrogen.

183. The compound according to claim 177, wherein p is 2, X is a bond, and each A is optionally substituted phenyl.

184. The compound according to claim 177, wherein said compound has one or more of the following features:

- a) Y' is S;
- b) R is hydrogen;
- c) p is 2, X is a bond, and each A is phenyl;
- d) B is ring (iii) above, wherein Ak is CH(CH₃) and Ar' is phenyl optionally substituted with halo, C1-C4 alkyl, or O-(C1-C4 alkyl).

185. A compound of formula VI:



VI;

or a pharmaceutically acceptable salt thereof;

Y' is O or S;

R is hydrogen or R²;

B is phenyl, 3-7 membered, monocyclic, saturated, carbocyclic ring, or 3-10 membered saturated or unsaturated, monocyclic or bicyclic heterocyclic ring having up to 4 heteroatoms selected from O, S, or N, or 5-10 membered monocyclic or bicyclic heteroaryl ring having up to 4 heteroatoms selected from O, S, or N;

wherein each ring G₁, G₂, and B is independently substituted with up to 4 substituents selected from R¹, R², R³, R⁴, or R⁵;

R¹ is oxo, R⁶ or ((C1-C4)aliphatic)_n-Y;

n is 0 or 1;

Y is halo, CN, NO₂, CF₃, OCF₃, OH, SR⁶, S(O)R⁶, SO₂R⁶, NH₂, NHR⁶, N(R⁶)₂, NR⁶R⁸, N(R⁸)₂, COOH, COOR⁶ or OR⁶; or

two R¹ on adjacent ring atoms, taken together, form 1,2-methylenedioxy or 1,2-ethylenedioxy;

R^2 is aliphatic, wherein each R^2 optionally comprises up to 2 substituents independently selected from R^1 , R^4 , or R^5 ;

R^3 is a cycloaliphatic, aryl, heterocyclic, or heteroaryl ring optionally comprising up to 3 substituents, independently selected from R^1 , R^2 , R^4 or R^5 ;

R^4 is OR^5 , OR^6 , $OC(O)R^6$, $OC(O)R^5$, $OC(O)OR^6$, $OC(O)OR^5$, $OC(O)N(R^6)_2$, $OC(O)N(R^5)_2$, $OC(O)N(R^6R^5)$, SR^6 , SR^5 , $S(O)R^6$, $S(O)R^5$, SO_2R^6 , SO_2R^5 , $SO_2N(R^6)_2$, $SO_2N(R^5)_2$, $SO_2NR^5R^6$, SO_3R^6 , SO_3R^5 , $C(O)R^5$, $C(O)OR^5$, $C(O)R^6$, $C(O)OR^6$, $C(O)N(R^6)_2$, $C(O)N(R^5)_2$, $C(O)N(R^5R^6)$, $C(O)N(OR^6)R^6$, $C(O)N(OR^5)R^6$, $C(O)N(OR^6)R^5$, $C(O)N(OR^5)R^5$, $C(NOR^6)R^6$, $C(NOR^6)R^5$, $C(NOR^5)R^6$, $C(NOR^5)R^5$, $N(R^6)_2$, $N(R^5)_2$, $N(R^5R^6)$, $NR^5C(O)R^5$, $NR^6C(O)R^6$, $NR^6C(O)R^5$, $NR^5C(O)R^6$, $NR^6C(O)OR^6$, $NR^5C(O)OR^6$, $NR^6C(O)OR^5$, $NR^5C(O)OR^5$, $NR^6C(O)N(R^6)_2$, $NR^6C(O)NR^5R^6$, $NR^6C(O)N(R^5)_2$, $NR^5C(O)N(R^6)_2$, $NR^5C(O)NR^5R^6$, $NR^5C(O)N(R^5)_2$, $NR^6SO_2R^6$, $NR^6SO_2R^5$, $NR^5SO_2R^5$, $NR^5SO_2R^6$, $NR^6SO_2N(R^6)_2$, $NR^6SO_2NR^5R^6$, $NR^6SO_2N(R^5)_2$, $NR^5SO_2N(R^6)_2$, $NR^5SO_2NR^5R^6$, $NR^5SO_2N(R^5)_2$, $N(OR^6)R^6$, $N(OR^6)R^5$, $N(OR^5)R^5$, or $N(OR^5)R^6$;

R^5 is a cycloaliphatic, aryl, heterocyclic, or heteroaryl ring, optionally comprising up to 3 R^1 substituents;

R^6 is H or aliphatic, wherein R^6 optionally comprises a R^7 substituent;

R^7 is a cycloaliphatic, aryl, heterocyclic, or heteroaryl ring, and each R^7 optionally comprises up to 2 substituents independently chosen from H, $(C_1-C_6)-$

straight or branched alkyl, (C₂-C₆) straight or branched alkenyl or alkynyl, 1,2-methylenedioxy, 1,2-ethylenedioxy, or (CH₂)_n-Z;

Z is selected from halo, CN, NO₂, CF₃, OCF₃, OH, S-aliphatic, S(O)-aliphatic, SO₂-aliphatic, NH₂, NH-aliphatic, N(aliphatic)₂, N(aliphatic)R⁸, NHR⁸, N(R⁸)₂, COOH, C(O)O(-aliphatic, or O-aliphatic; and

R⁸ is an amino capping group;
provided that when R is hydrogen:

a) B is not quinolin-2-yl or 1,2-dihydro-2-oxo-quinolin-4-yl;

b) when G₁ and G₂ both are phenyl, and Y' is S, then B is not 1,4-benzodioxin-2-yl, cyclopropyl, cyclohexyl, thien-2-yl, 1H-thieno[2,3-c]pyrazol-1-phenyl-3-methyl-5-yl, 5-methyl-thien-3-yl, 2,5-dichloro-thien-3-yl, 2-phenyl-quinolin-4-yl, furan-2-yl, thien-5-(4,5-diphenyl-2-thiazolyl-carboxamide)-2-yl, benzo[b]thiophen-2-yl, pyridin-2-(4,5-diphenyl-2-thiazolyl-carboxamide)-6-yl, 5-nitro-thien-2-yl, 3-chloro-benzo[b]thiophen-2-yl, 4H-1-benzopyran-3-yl or 2H-1-benzopyran-3,4-dihydro-3-oxo-4-yl, 4H-1-benzopyran-3-yl or 2H-1-benzopyran-3,4-dihydro-3-oxo-4-yl;


c) when G₁ and G₂ both are phenyl, and Y' is O, then B is not 1,2-dihydro-2-oxo-quinolin-4-yl or 3,4-dihydro-3-phenyl-phthalazin-1-yl or thien-2-yl;

d) the following compounds are excluded:

Y'	G ₁	G ₂	B
S	Ph	4-Me-Ph	2-Cl-thien-5-yl or 2,5-dichloro-thien-3-yl
S	Ph or 3,4-dimethyl phenyl	3,4-dimethyl phenyl or Ph	pyridin-3-yl
S	3,4-dimethyl phenyl	Ph	pyridin-4-yl
S	4-Cl-Ph	Ph	thien-2-yl

S	3,4-dimethylphenyl	Ph	pyridin-4-yl
S	Ph	4-Me-phenyl	thien-2-yl or benzothiazol-2-yl
S	4-NO ₂ -Ph or 4-Me-Ph	Ph or 4-OMe-Ph or 2,4-dimethylphenyl	thien-2-yl or furan-2-yl
S	4-OMe-Ph or 2,4-dimethylphenyl	4-NO ₂ -Ph	furan-2-yl or thien-2-yl
S	4-OMe-Ph or Ph	4-OMe-Ph	furan-2-yl or thien-2-yl
S	Ph	4-Me-Ph	furan-2-yl
S	Ph	4-Me-Ph	5-nitro-thien-2-yl
S	4-Me-Ph	4-Me-Ph	2-chloro-pyridin-3-yl
S	Ph	4-Me-Ph	3-chloro-benzo[b]thiophen-2-yl
S	4-Me-Ph or Ph	4-NO ₂ -Ph	thien-2-yl or furan-2-yl
S	2,4-dimethylphenyl	4-NO ₂ -Ph	pyridin-3-yl or thien-2-yl
S	4-Cl-Ph	4-NO ₂ -Ph	thien-2-yl
S	Ph or 2-OMe-Ph or 3-OMe-Ph or 4-OMe-Ph	2,4-dimethoxyphenyl	1H-indol-2-yl.

e) when Y' is S, G₁ and G₂ are both phenyl, then B is



186. The compound according to claim 185, wherein G₁ and G₂ are both phenyl.

187. The compound according to claim 185, wherein each of G₁ and G₂ is independently and optionally substituted with up to two substituents selected from halo, or C1-C4 alkyl.

188. The compound according to claim 185, wherein B is phenyl optionally substituted with up to two substituents selected from halo, C1-C4 alkyl, O-(C1-C4 alkyl), COOH, COO(C1-C4 alkyl), or cyano.

189. The compound according to claim 185, wherein B is phenyl, phenyl-2-carboxylic acid methyl ester, 3,4-dichlorophenyl, 4-chlorophenyl, 4-methoxyphenyl, 4-methylphenyl, 2,6-difluorophenyl, 2-methylphenyl, 3-methylphenyl, phenyl-2-carboxylic acid, 2-chlorophenyl, 4-cyanophenyl, 2-methoxyphenyl, 3-chlorophenyl, or 3-methoxyphenyl.


190. The compound according to claim 185, wherein B is 3-7 membered, monocyclic, saturated, carbocyclic ring.

191. The compound according to claim 190, wherein B is selected from cyclopropyl, cyclopentyl, cyclohexyl, or cycloheptyl.

192. The compound according to claim 185, wherein B is a 3-10 membered saturated or unsaturated, monocyclic or bicyclic heterocyclic ring having up to 4 heteroatoms selected from O, S, or N.

193. The compound according to claim 192, wherein B is selected from tetrahydrofuranyl.

194. A compound of formula VII:



VII;

or a pharmaceutically acceptable salt thereof;

wherein:

Y' is O or S;

R is hydrogen or R²;

R^{Alk} is C1-C6 aliphatic, optionally substituted with up to 3 substituents independently selected from R¹, R², or R³;

B is phenyl, 3-7 membered, monocyclic, saturated, carbocyclic ring, or 3-10 membered saturated or unsaturated, monocyclic or bicyclic heterocyclic ring having up to 4 heteroatoms selected from O, S, or N, or 5-10 membered monocyclic or bicyclic heteroaryl ring having up to 4 heteroatoms selected from O, S, or N;

wherein each ring G₁, G₂, and B is independently substituted with up to 4 substituents selected from R¹, R², R³, R⁴, or R⁵;

R¹ is oxo, R⁶ or ((C1-C4)aliphatic)_n-Y;

n is 0 or 1;

Y is halo, CN, NO₂, CF₃, OCF₃, OH, SR⁶, S(O)R⁶, SO₂R⁶, NH₂, NHR⁶, N(R⁶)₂, NR⁶R⁸, N(R⁸)₂, COOH, COOR⁶ or OR⁶; or

two R¹ on adjacent ring atoms, taken together, form 1,2-methylenedioxy or 1,2-ethylenedioxy;

R² is aliphatic, wherein each R² optionally comprises up to 2 substituents independently selected from R¹, R⁴, or R⁵;

R³ is a cycloaliphatic, aryl, heterocyclic, or heteroaryl ring optionally comprising up to 3 substituents, independently selected from R¹, R², R⁴ or R⁵;

R^4 is OR^5 , OR^6 , $OC(O)R^6$, $OC(O)R^5$, $OC(O)OR^6$, $OC(O)OR^5$, $OC(O)N(R^6)_2$, $OC(O)N(R^5)_2$, $OC(O)N(R^6R^5)$, SR^6 , SR^5 , $S(O)R^6$, $S(O)R^5$, SO_2R^6 , SO_2R^5 , $SO_2N(R^6)_2$, $SO_2N(R^5)_2$, $SO_2NR^5R^6$, SO_3R^6 , SO_3R^5 , $C(O)R^5$, $C(O)OR^5$, $C(O)R^6$, $C(O)OR^6$, $C(O)N(R^6)_2$, $C(O)N(R^5)_2$, $C(O)N(R^5R^6)$, $C(O)N(OR^6)R^6$, $C(O)N(OR^5)R^6$, $C(O)N(OR^6)R^5$, $C(O)N(OR^5)R^5$, $C(NOR^6)R^6$, $C(NOR^6)R^5$, $C(NOR^5)R^6$, $C(NOR^5)R^5$, $N(R^6)_2$, $N(R^5)_2$, $N(R^5R^6)$, $NR^5C(O)R^5$, $NR^6C(O)R^6$, $NR^6C(O)R^5$, $NR^5C(O)R^6$, $NR^6C(O)OR^6$, $NR^5C(O)OR^6$, $NR^6C(O)OR^5$, $NR^5C(O)OR^5$, $NR^6C(O)N(R^6)_2$, $NR^6C(O)NR^5R^6$, $NR^6C(O)N(R^5)_2$, $NR^5C(O)N(R^6)_2$, $NR^5C(O)NR^5R^6$, $NR^5C(O)N(R^5)_2$, $NR^6SO_2R^6$, $NR^6SO_2R^5$, $NR^5SO_2R^5$, $NR^5SO_2R^6$, $NR^6SO_2N(R^6)_2$, $NR^6SO_2NR^5R^6$, $NR^6SO_2N(R^5)_2$, $NR^5SO_2N(R^6)_2$, $NR^5SO_2NR^5R^6$, $NR^5SO_2N(R^5)_2$, $N(OR^6)R^6$, $N(OR^6)R^5$, $N(OR^5)R^5$, or $N(OR^5)R^6$;

R^5 is a cycloaliphatic, aryl, heterocyclic, or heteroaryl ring, optionally comprising up to 3 R^1 substituents;

R^6 is H or aliphatic, wherein R^6 optionally comprises a R^7 substituent;

R^7 is a cycloaliphatic, aryl, heterocyclic, or heteroaryl ring, and each R^7 optionally comprises up to 2 substituents independently chosen from H, (C_1-C_6) -straight or branched alkyl, (C_2-C_6) straight or branched alkenyl or alkynyl, 1,2-methylenedioxy, 1,2-ethylenedioxy, or $(CH_2)_n-Z$;

Z is selected from halo, CN, NO_2 , CF_3 , OCF_3 , OH, S-aliphatic, $S(O)$ -aliphatic, SO_2 -aliphatic, NH_2 , NH-aliphatic, $N(\text{aliphatic})_2$, $N(\text{aliphatic})R^8$, NHR^8 , $N(R^8)_2$, COOH, $C(O)O(-\text{aliphatic})$, or O-aliphatic; and

R^8 is an amino-capping group;
provided that when each of G_1 , G_2 , and B is unsubstituted phenyl, and R is hydrogen, then B is not 3,4,5-trimethoxyphenyl.

195. The compound according to claim 194, wherein R^{Ak} is selected from methyl, ethyl, isopropyl, n-propyl, sec-butyl, n-butyl, or t-butyl.

196. The compound according to claim 194, wherein B is optionally substituted phenyl.

197. The compound according to claim 194, wherein B is 3-7 membered, monocyclic, saturated, carbocyclic ring.


198. The compound according to claim 197, wherein B is selected from cyclopropyl, cyclopentyl, cyclohexyl, or cycloheptyl.

199. The compound according to claim 194, wherein B is a 3-10 membered saturated or unsaturated, monocyclic or bicyclic heterocyclic ring having up to 4 heteroatoms selected from O, S, or N.

200. The compound according to claim 199, wherein B is selected from tetrahydrofuranyl, tetrahydrothiophenyl, pyrrolidinyl, or piperazinyl.

201. The compound according to claim 194, wherein B is a 5-10 membered monocyclic or bicyclic heteroaryl ring having up to 4 heteroatoms selected from O, S, or N.

202. A compound of formula I':



I';

or a pharmaceutically acceptable salt thereof;



wherein:

Y' is O, S, or NR;

R is H, R², or R⁶;

C is a phenyl or 5-8 membered cycloaliphatic ring;

Q is selected from:



each B is independently selected from 3-7 membered monocyclic or 8-14 membered bicyclic or tricyclic, saturated, unsaturated or aromatic ring containing 0-4 heteroatoms in each ring, wherein each said heteroatom is independently selected from N, NH, S, or O;

wherein each B and C is independently and optionally substituted with up to 4 substituents independently selected from R¹, R², R³, R⁴, or R⁵;

R^L is -OR^A, -SR^A, or -N(R^{AB})₂;

each R^A is independently hydrogen, C1-C6 aliphatic, or a 3-7 membered carbocyclic or heterocyclic ring, saturated or unsaturated ring, having up to 3 heteroatoms selected from O, N, or S, wherein each R^A is optionally

substituted with up to 3 substituents independently selected from R^1 , R^4 or R^7 ,

each R^{AB} is independently hydrogen or C1-C6 aliphatic optionally substituted with up to 3 substituents independently selected from R^1 , R^4 or R^7 ;

wherein up to two methylene units in R^A or R^{AB} are optionally replaced with -CO-, -CS-, -COCO-, -CONR-, -CO₂-, -OCO-, -NRCO₂-, -O-, -NRCONR-, -OCONR-, -NRCO-, -S-, -SO-, -SO₂-, -NR-, -SO₂NR-, NRSO₂-, or -NRSO₂NR; or

two R^{AB} , taken together with the nitrogen atom, is a 3-7 membered heterocyclic or heteroaryl ring containing up to 4 heteroatoms selected from O, N, or S, wherein said ring is optionally substituted with up to 2 substituents selected from oxo or (C₁₋₄aliphatic)_p-Y;

R^M is C1-C6 aliphatic, optionally substituted with up to two substituents selected from R^1 , R^2 , R^3 , or R^4 ;

each of X_1 and X_2 is independently selected from O, S, or NR;

R^N is C1-C6 aliphatic or phenyl, wherein R^N is optionally substituted with up to two substituents selected from R^1 , R^2 , R^3 , or R^4 ;

R^P is C1-C6 aliphatic, optionally substituted with up to two substituents selected from R^1 , R^2 , R^3 , or R^4 ;

R^Q is C1-C6 aliphatic or aryl, wherein R^Q is optionally substituted with up to two substituents selected from R^1 , R^2 , R^3 , or R^4 ;

R^1 is oxo, R^6 or ((C1-C4)aliphatic)_n-Y;

n is 0 or 1;

Y is halo, CN, NO₂, CF₃, OCF₃, OH, SR⁶, S(O)R⁶, SO₂R⁶, NH₂, NHR⁶, N(R⁶)₂, NR⁶R⁸, N(R⁸)₂, COOH, COOR⁶ or OR⁶; or

two R^1 on adjacent ring atoms, taken together, form 1,2-methylenedioxy or 1,2-ethylenedioxy;

R^2 is aliphatic, wherein each R^2 optionally comprises up to 2 substituents independently selected from R^1 , R^4 , or R^5 ;

R^3 is a cycloaliphatic, aryl, heterocyclic, or heteroaryl ring optionally comprising up to 3 substituents, independently selected from R^1 , R^2 , R^4 or R^5 ;

R^4 is OR^5 , OR^6 , $OC(O)R^6$, $OC(O)R^5$, $OC(O)OR^6$, $OC(O)OR^5$, $OC(O)N(R^6)_2$, $OC(O)N(R^5)_2$, $OC(O)N(R^6R^5)$, SR^6 , SR^5 , $S(O)R^6$, $S(O)R^5$, SO_2R^6 , SO_2R^5 , $SO_2N(R^6)_2$, $SO_2N(R^5)_2$, $SO_2NR^5R^6$, SO_3R^6 , SO_3R^5 , $C(O)R^5$, $C(O)OR^5$, $C(O)R^6$, $C(O)OR^6$, $C(O)N(R^6)_2$, $C(O)N(R^5)_2$, $C(O)N(R^5R^6)$, $C(O)N(OR^6)R^6$, $C(O)N(OR^5)R^6$, $C(O)N(OR^6)R^5$, $C(O)N(OR^5)R^5$, $C(NOR^6)R^6$, $C(NOR^6)R^5$, $C(NOR^5)R^6$, $C(NOR^5)R^5$, $N(R^6)_2$, $N(R^5)_2$, $N(R^5R^6)$, $NR^5C(O)R^5$, $NR^6C(O)R^6$, $NR^6C(O)R^5$, $NR^5C(O)R^6$, $NR^6C(O)OR^6$, $NR^5C(O)OR^6$, $NR^6C(O)OR^5$, $NR^5C(O)OR^5$, $NR^6C(O)N(R^6)_2$, $NR^6C(O)NR^5R^6$, $NR^6C(O)N(R^5)_2$, $NR^5C(O)N(R^6)_2$, $NR^5C(O)NR^5R^6$, $NR^5C(O)N(R^5)_2$, $NR^6SO_2R^6$, $NR^6SO_2R^5$, $NR^5SO_2R^5$, $NR^5SO_2R^6$, $NR^6SO_2N(R^6)_2$, $NR^6SO_2NR^5R^6$, $NR^6SO_2N(R^5)_2$, $NR^5SO_2N(R^6)_2$, $NR^5SO_2NR^5R^6$, $NR^5SO_2N(R^5)_2$, $N(OR^6)R^6$, $N(OR^6)R^5$, $N(OR^5)R^5$, or $N(OR^5)R^6$;

R^5 is a cycloaliphatic, aryl, heterocyclic, or heteroaryl ring, optionally comprising up to 3 R^1 substituents;

R^6 is H or aliphatic, wherein R^6 optionally comprises a R^7 substituent;

R^7 is a cycloaliphatic, aryl, heterocyclic, or heteroaryl ring, and each R^7 optionally comprises up to 2 substituents independently chosen from H, (C_1-C_6) -straight or branched alkyl, (C_2-C_6) straight or branched alkenyl or alkynyl, 1,2-methylenedioxy, 1,2-ethylenedioxy, or $(CH_2)_n-Z$;

Z is selected from halo, CN, NO_2 , CF_3 , OCF_3 , OH, S-aliphatic, S(O)-aliphatic, SO_2 -aliphatic, NH_2 , NH-aliphatic, $N(aliphatic)_2$, $N(aliphatic)R^8$, NHR^8 , $N(R^8)_2$, COOH, $C(O)O(-aliphatic)$, or O-aliphatic; and

R^8 is an amino-capping group.

203. The compound according to claim 202, wherein Y' is O.

204. The compound according to claim 202, wherein Y' is S.

205. The compound according to claim 202, wherein R is hydrogen.

206. The compound according to claim 202, wherein Q in formula I' is B (structure (a)).

207. The compound according to claim 202, wherein Q in formula I' is $-(C_1-C_6 \text{ aliphatic})-B$ (structure (b)).

208. The compound according to claim 207, wherein said C_1-C_6 aliphatic is C_1-C_4 straight or branched alkylidene.

209. The compound according to claim 208, wherein said alkylidene is selected from $-\text{CH}_2-$, $-\text{CH}(\text{Me})-$, $-\text{C}(\text{Me})_2-$, $-\text{CH}(\text{Et})-$, $-\text{C}(\text{Et})_2-$, or $-\text{CH}_2-\text{CH}(\text{Me})-$.

210. The compound according to claim 202, wherein B is selected from optionally substituted phenyl, or C3-C8 cycloaliphatic.

211. The compound according to claim 210, wherein B is phenyl or C3-C8 cycloalkyl optionally substituted with up to two substituents selected from R^1 , R^2 , or phenyl optionally substituted with up to two substituents selected from R^1 or R^2 .

212. The compound according to claim 202, wherein B is selected from optionally substituted C3-C8 cycloalkyl, phenyl, piperidyl, or pyrrolidinyl.

213. The compound according to claim 212, wherein B is phenyl, cyclopentyl, cyclohexyl, or piperidyl, optionally substituted with up to two R^1 or R^2 substituents.

214. The compound according to claim 202, wherein ring C is phenyl optionally substituted with up to two R^1 .

215. The compound according to claim 202, wherein ring C is cyclohexenyl optionally substituted with up to two R^1 .

216. The method according to claim 1, wherein said ABC-transporter or a fragment thereof is *in vivo*.

217. The method according to claim 1, wherein said ABC-transporter or a fragment thereof is *in vitro*.

218. The method according to claim 1, wherein said ABC-transporter is CFTR.

219. A method of treating an ABC transporter mediated disease in a mammal, comprising the step of administering to said mammal a composition comprising a compound according to any one of claims 1-104, or a compound according to any one of claims 105-215.

220. The method according to claim 219, wherein said disease is selected from cystic fibrosis, hereditary emphysema, hereditary hemochromatosis, coagulation-cibrinolysis deficiencies, such as protein C deficiency, Type 1 hereditary angioedema, lipid processing deficiencies, such as familial hypercholesterolemia, Type 1 chylomicronemia, abetalipoproteinemia, lysosomal storage diseases, such as I-cell disease/Pseudo-Hurler, secretory diarrhea or polycystic kidney disease, mucopolysaccharidoses, Sandhof/Tay-Sachs, Crigler-Najjar type II, polyendocrinopathy/Hyperinsulemia, Diabetes mellitus, Laron dwarfism, myeloperoxidase deficiency, primary hypoparathyroidism, melanoma, glycanosis CDG type 1, hereditary emphysema, congenital hyperthyroidism, osteogenesis imperfecta, hereditary hypofibrinogenemia, ACT deficiency, Diabetes insipidus (DI), neurophyseal DI, Neprogenic DI, Charcot-Marie Tooth syndrome, Perlizaeus-Merzbacher disease, neurodegenerative diseases such as Alzheimer's disease, Parkinson's disease, Amyotrophic lateral sclerosis, progressive supranuclear palsy, Pick's disease, several polyglutamine neurological disorders asuch as Huntington, Spinocerebular ataxia type I,

spinal and bulbar muscular atrophy, Dentatorubal pallidoluysian, and Myotonic dystrophy, as well as Spongiform encephalopathies, such as Hereditary Creutzfeldt-Jakob disease (due to Prion protein processing defect), Fabry disease, Strausler-Scheinker syndrome, COPD, dry eye disease, or Sjogren's disease.

221. The method according to claim 220, wherein said disease is cystic fibrosis.

222. A method of modulating activity of an anion channel *in vitro* or *in vivo*, comprising the step of contacting said channel with a compound according to any one of claims 1-104, or a compound according to any one of claims 105-215.

223. The method according to claim 222, wherein said anion channel is a chloride channel or a bicarbonate channel.

224. The method according to claim 223, wherein said anion channel is a chloride channel.

225. A method of treating an anion channel mediated disease in a mammal, comprising the step of administering to said mammal a composition comprising a compound according to any one of claims 1-104, or a compound according to any one of claims 105-220.

226. The method according to claim 225, wherein said disease is cystic fibrosis.

227. A pharmaceutical composition comprising a compound according to any one of claims 104-215, and a pharmaceutically acceptable carrier.

228. A pharmaceutical composition comprising:

- (i) a compound according to any one of claims 105-215;
- (ii) a pharmaceutically acceptable carrier; and
- (iii) an additional agent selected from a mucolytic agent, bronchodilator, an anti-biotic, an anti-infective agent, an anti-inflammatory agent, CFTR modulator, or a nutritional agent.

229. A kit for use in measuring the activity of a ABC transporter or a fragment thereof in a biological sample *in vitro* or *in vivo*, comprising:

- (i) a composition comprising a compound according to any one of claims 105-215; and
- (ii) instructions for:
 - a) contacting the composition with the biological sample;
 - b) measuring activity of said ABC transporter or a fragment thereof.

230. The kit according to claim 229, wherein said ABC transporter is CFTR.