




(86) Date de dépôt PCT/PCT Filing Date: 1991/05/16
(87) Date publication PCT/PCT Publication Date: 1992/11/26
(45) Date de délivrance/Issue Date: 2002/07/30
(85) Entrée phase nationale/National Entry: 1993/10/06
(86) N° demande PCT/PCT Application No.: US 1991/003421
(87) N° publication PCT/PCT Publication No.: 1992/020340

(51) Cl.Int.⁵/Int.Cl.⁵ A61K 31/52
(72) Inventeurs/Inventors:
Bolund, Lars, DK;
Jensen, Peter K. A., DK;
Bjerring, Peter, DK
(73) Propriétaire/Owner:
Senetek, plc, US
(74) Agent: MACRAE & CO.

(54) Titre : METHODE ET COMPOSITION POUR LE TRAITEMENT DES MALADIES HYPERPROLIFERATIVES DE LA
PEAU A L'AIDE DE CYTOKININES 6-AMINOPURINES
(54) Title: METHOD AND COMPOSITION FOR TREATING HYPERPROLIFERATIVE SKIN DISEASES USING 6-
AMINOPURINE CYTOKININS



(57) Abrégé/Abstract:

It has been discovered that 6-(substituted amino)purine cytokinins, such as kinetin, induce differentiation, and consequently reduce or eliminate the abnormally high rate of proliferation, of hyperproliferating cells associated with hyperproliferative skin diseases, such as psoriasis. Thus, the present invention provides methods and compositions for treating hyperproliferative skin diseases in mammals, particularly such diseases associated with hyperproliferating epidermal cells in humans, by administering to the hyperproliferating cells associated with such a disease, in the skin of a mammal suffering therefrom, a differentiation-inducing effective amount of a 6-(substituted amino)purine cytokinin.





PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : A61K 31/52	A1	(11) International Publication Number: WO 92/20340 (43) International Publication Date: 26 November 1992 (26.11.92) 2107896
(21) International Application Number: PCT/US91/03421 (22) International Filing Date: 16 May 1991 (16.05.91) (71) Applicant: SENETEK, PLC [GB/US]; Six The Pines Court, Suite D, St. Louis, MO 63141 (US). (72) Inventors: BOLUND, Lars ; Rylevej 22, DK-8240 Risskov (DK). JENSEN, Peter, K., A. ; Egevej 16, DK-8680 Ry (DK). BJERRING, Peter ; Permelillevej 15, DK-8240 Risskov (DK). (74) Agents: WATT, Phillip, H.; Fitch, Even, Tabin & Flannery, Room 900, 135 South LaSalle Street, Chicago, IL 60603 (US) et al.		(81) Designated States: AT (European patent), BE (European patent), CA, CH (European patent), DE (European patent), DK (European patent), ES (European patent), FR (European patent), GB (European patent), GR (European patent), IT (European patent), JP, LU (European patent), NL (European patent), SE (European patent). Published <i>With international search report.</i>
(54) Title: METHOD AND COMPOSITION FOR TREATING HYPERPROLIFERATIVE SKIN DISEASES USING 6-AMINOPURINE CYTOKININS (57) Abstract It has been discovered that 6-(substituted amino)purine cytokinins, such as kinetin, induce differentiation, and consequently reduce or eliminate the abnormally high rate of proliferation, of hyperproliferating cells associated with hyperproliferative skin diseases, such as psoriasis. Thus, the present invention provides methods and compositions for treating hyperproliferative skin diseases in mammals, particularly such diseases associated with hyperproliferating epidermal cells in humans, by administering to the hyperproliferating cells associated with such a disease, in the skin of a mammal suffering therefrom, a differentiation-inducing effective amount of a 6-(substituted amino)purine cytokinin.		

-1- 2107896

METHOD AND COMPOSITION FOR TREATING HYPERPROLIFERATIVE SKIN DISEASES USING 6-AMINOPURINE CYTOKININS

TECHNICAL FIELD

The present invention concerns methods and compositions for treating diseases associated with hyperproliferating epidermal cells, particularly disorders of skin in humans.

BACKGROUND OF THE INVENTION

Among diseases afflicting mammals, and particularly humans, are many which are due to hyperproliferation of epidermal cells. Among such diseases are skin diseases such as basal cell carcinoma, malignant melanoma, squamous cell carcinoma, actinic keratosis, Bowen's Disease, papilloma, seborrheic keratosis, toxic eczema, allergic eczema, atopic dermatitis, ichthyosis, and psoriasis.

The present invention provides methods and compositions for treating such hyperproliferative skin diseases.

The methods and compositions of the invention involve cytokinins. Cytokinins are plant hormones involved in promoting growth and cell division in explants of plant tissue in culture in standard media, which contain auxins (another class of plant hormones) as well as vitamins, mineral salts, and sugar.

Among cytokinins are those which are 6-(substituted amino)purines, including kinetin (6-(furfuryl)aminopurine), zeatin (6-(3-hydroxymethyl, 3-methylallyl)aminopurine, 6-(3,3-dimethylallyl)aminopurine, 6-(benzyl)aminopurine, 6-(phenyl)aminopurine, 6-(n-alkyl)aminopurine, wherein the alkyl group has 4, 5 or 6 carbon atoms, and 6-(cyclohexyl)methylaminopurine.

Cytokinins have been observed to have various effects on mammalian cells in culture. Orr and McSwain, Cancer 10, 617 - 624 (1957); Johnson et al., Exptl. Cell Res. 85, 47 - 56 (1974); Hecht et al., Proc. Natl. Acad. Sci. (USA) 71, 4670 - 4674 (1974); Mookerjee et al., J. Reticuloendoth. Soc. 25, 299 - 314 (1979);

SUBSTITUTE SHEET

2107896

-2-

Gordon and Seglen, Arch. Biochem. Biophys. 217, 282 - 294 (1982); Vesely et al., Endocrinology 116, 1887 - 1892 (1985).

5

SUMMARY OF THE INVENTION

It has been found unexpectedly that hyperproliferating epidermal cells associated with hyperproliferative skin diseases of mammals, when exposed
10 to 6-(substituted amino)purine cytokinins, are caused to differentiate and slow in their rate of proliferation.

Thus, the present invention entails methods and compositions for treating hyperproliferative skin diseases of mammals, including humans. According to the
15 methods of the invention, differentiation-inducing-effective amounts of a 6-(substituted amino)purine are administered to the hyperproliferating epidermal cells associated with such a disease in a mammal suffering therefrom.

20

DETAILED DESCRIPTION OF THE INVENTION

In one of its aspects, the present invention entails a method of treating an hyperproliferative skin disease in a mammal suffering therefrom which method
25 comprises administering to the hyperproliferating cells, associated with said disease, of the skin of said mammal a differentiation-inducing effective amount of a 6-(substituted amino)purine cytokinin.

In another of its aspects, the invention entails
30 a composition for treating an hyperproliferative skin disease of a mammal, which composition comprises a 6-(substituted amino)purine cytokinin and a physiologically acceptable vehicle for application of said cytokinin to the outer surface of the skin of a mammal
35 suffering from said disease such that said cytokinin is brought into contact with (i.e., administered to) the hyperproliferating cells associated with the disease.

SUBSTITUTE SHEET

2107896


-3-

The present invention is preferably employed to treat hyperproliferative skin diseases of humans.

Such diseases, in humans and other mammals, are due to hyperproliferating epidermal cells of the skin. Among such diseases, that can be treated in accordance with the present invention, are basal cell carcinoma, malignant melanoma, squamous cell carcinoma, actinic keratosis, Bowen's Disease, papilloma, seborrheic keratosis, toxic eczema, allergic eczema, atopic dermatitis, ichthyosis, and psoriasis.

In accordance with the method of the invention, a 6-(substituted amino)purine cytokinin is brought into contact with (i.e., administered to) the hyperproliferating cells associated with an hyperproliferating skin disease in an amount which is effective to induce such cells to differentiate.

Among the 6-(substituted amino)purine cytokinins which can be employed in accordance with the invention are those of formula I:



I,

wherein R_1 is selected from the group consisting of furfuryl, phenyl, benzyl, n-alkyl of 4, 5 or 6 carbon atoms, (cyclohexyl)methyl ($-\text{CH}_2(\text{C}_6\text{H}_{11})$),

(3-hydroxymethyl-3-methyl)allyl ($-\text{CH}_2\text{CH}=\text{C}\begin{matrix} \text{CH}_2\text{OH} \\ \text{CH}_3 \end{matrix}$), and 3,3-dimethylallyl ($-\text{CH}_2\text{CH}=\text{C}(\text{CH}_3)_2$). Most preferred among these is kinetin (R_1 being furfuryl).

For carrying out the method of the invention, the 6-(substituted amino)purine cytokinin is combined, at

SUBSTITUTE SHEET

2107896

-4-

a concentration of between about 0.01 % (w/v) and about 0.5 % (w/v), preferably about 0.1 % (w/v) with physiologically acceptable vehicle, such as a standard, commonly used skin-moisturizing vehicle. One example of the many such skin-moisturizing vehicles is a composition of 4 - 8 wt. % (e.g., 6.2 wt. %) glycerol monostearate 40 - 50, 12 - 16 wt. % (e.g., 14.3 wt. %) white petroleum, 0.5 - 2 wt. % (e.g., 1 wt. %) lanolin, 9 -13 wt. % (e.g., 11.4 wt. %) mineral oil, 0.5 - 2 wt. % (e.g., 1 wt. %) propylene glycol, 1 - 2 wt. % (e.g., 1.4 wt. %) triethanolamine (bp-80), 1 - 5 wt. % (e.g., 3.3 wt. %) stearin, 0.5 - 2 wt. % (e.g., 1 wt. %) parahydroxybenzoic acid, and the remainder (e.g., 60.4 wt. %) water. Alternatively, the 6-(substituted amino)purine cytokinin can be combined with a lipogel-type vehicle, such as that being sold by Rhone-Poulenc (Courbevoie, France) under the tradename CERIDAL™, at a concentration of between about 1 and about 5 mg/ml. The resulting composition according to the invention is applied topically to the surface of the area of the skin affected by the hyperproliferative disease (i.e., the area under, or as a part of, which the hyperproliferating epidermal cells associated with the disease are located), usually with light rubbing or message or the like as commonly employed in the application of moisturizing vehicles or lipogel-based compositions.

It is contemplated that the method of the invention will be carried out under the guidance of a dermatologist or other physician or, in the case of a non-human mammal, veterinarian.

The frequency of application most suitable for treating a particular disease in a particular mammal with a particular composition according to the invention will be readily determined by the skilled and will depend, as the skilled will understand, on numerous factors, such as

SUBSTITUTE SHEET

-5-

2107896

the species of mammal, the nature of the disease, the severity of the disease, the moisturizing vehicle, lipogel-based vehicle or other physiologically acceptable vehicle employed in the composition, and the concentration of the 6-(substituted amino)purine cytokinin in the composition. Generally, application of a composition according to the invention to an affected area of the skin between once and four times daily (preferably twice) for between about a week and about a year, more typically about a month, will be effective to induce hyperproliferating cells under, or in, the area to differentiate to a non-proliferating or normally proliferating state and thereby substantially cure the hyperproliferative disease of the affected area.

The invention will now be illustrated in the following Example.

EXAMPLE

Primary explant cultures of human epidermal cells from psoriatic skin of two persons suffering from psoriasis were established by taking skin samples, cutting the samples into small (approx. $2 \times 2 \text{ mm}^2$) pieces, and explanting the pieces into ordinary tissue culture flasks, where culture growth was carried out at 35°C with a standard keratinocyte culture medium (Dulbecco's Modified Eagle's Medium + 20 % fetal calf serum + 10 ng/ml epidermal growth factor + $0.4 \mu\text{g/ml}$ hydrocortisone).

Cultures were stripped by a standard procedure after growth for one month. That is, they were incubated for 72 hours in low Ca^{++} medium prior to selectively stripping off all the supra basal layers. After the stripping, the cultures were re-fed with medium with normal Ca^{++} concentration

In half of the cultures from each patient, kinetin was present in the culture medium at $200 \mu\text{M}$ concentration.

SUBSTITUTE SHEET

2107896

-6-

The morphology of cells in the primary, unstripped cultures was examined after one month plus ten days of growth. In the cultures fed with kinetin, in comparison with those not so fed, there was a large decrease in mitotic figures and some indication of increased differentiation (accumulation of numerous, filament-containing cells in discrete, elevated areas).

The morphology of cells in the stripped cultures was also examined, after seven days of growth after the stripping. In the cultures fed with kinetin, in comparison with those not so fed, there was much more heterogeneity in morphology. In the cultures fed with kinetin, in comparison with those not so fed, there was a reduction of mitotic figures, reduction of multilayering, no lateral growth, early in the observation period a vacuolization of the cytoplasm in many cells, late in the observation period areas with numerous cornified envelope-containing cells alternating with empty looking areas.

At seven days post-stripping, cultures were labeled using ^3H -labeled thymidine for 30 minutes. S-phase cells were sorted out onto glass slides, on the basis of EtBr-Mithramycin staining using a FACS-II cell sorter, and then subjected to autoradiography. The numbers of labeled and unlabeled S-phase cells, and the number of grains per nucleus in the labeled cells, were then measured microscopically. In cultures with kinetin, more than 70 % of S-phase cells were unlabeled (i.e., 0 - 3 grains per nucleus) and about 5 % of S-phase cells were highly labeled (>25 grains per nucleus). In cultures without kinetin, about 30 % of S-phase cells were unlabeled and about 40 % of S-phase cells were highly labeled. These results indicate that kinetin inhibited the regenerative response in psoriatic cells.

The cornified envelope formed by cultured keratinocytes is regarded as a marker of differentiation.

SUBSTITUTE SHEET

The fraction of cells in a culture that has acquired a cornified envelope can be measured by a standard method, wherein the cells in the culture are counted, then
5 treated with SDS and reducing agents, which destroys cells without the envelope but to which the envelope is resistant, and then recounted. With cultures of psoriatic cells from one of the persons suffering from psoriasis, in which cells with cornified envelope were
10 determined 7-days post-stripping, 37 % of the cells in cultures treated with kinetin had a cornified envelope while 18 % of the cells in cultures not treated with kinetin had a cornified envelope. With cultures of psoriatic cells from the other of the persons suffering
15 from psoriasis, in which cells with cornified envelope were determined 7-days post-stripping, 14 % of the cells in cultures treated with kinetin had a cornified envelope while 12 % of the cells in cultures not treated with kinetin had a cornified envelope. These results indicate
20 that kinetin stimulated differentiation of the psoriatic cells.

While the invention has been described with some specificity in the instant specification, persons of
25 skill in the art will recognize modifications and variations of what has been described that fall within the spirit of the invention. It is intended that such modifications and variations be encompassed by the following claims.

30

35

SUBSTITUTE SHEET

-8-


WHAT IS CLAIMED IS:

1. For use in topically treating a hyperproliferative skin disease in a human suffering therefrom, a differentiation-inducing-effective amount of a compound which is a 6-(substituted amino)purine cytokinin.

2. A compound as defined in claim 1, wherein the disease is selected from the group consisting of basal cell carcinoma, malignant melanoma, squamous cell carcinoma, actinic keratosis, Bowen's Disease, papilloma, seborrheic keratosis, toxic eczema, allergic eczema, atopic dermatitis, ichthyosis, and psoriasis.

3. A compound as defined in Claim 2, wherein the disease is psoriasis.

4. A compound as defined in Claim 1, wherein the 6-(substituted amino)purine cytokinin is of formula I:




I,

wherein R_1 is selected from the group consisting of furfuryl, phenyl, benzyl, n-alkyl of 4, 5 or 6 carbon atoms, (cyclohexyl)methyl, (3-hydroxymethyl-3-methyl)allyl, and 3,3-dimethylallyl.

-9-


5. A compound as defined in Claim 2, wherein the 6-(substituted amino)purine cytokinin is of formula I:



I,

wherein R₁ is selected from the group consisting of
furfuryl, phenyl, benzyl, n-alkyl of 4, 5 or 6 carbon
atoms, (cyclohexyl)methyl, (3-hydroxymethyl-3-methyl)allyl,
and 3,3-dimethylallyl.

6. A compound as defined in Claim 3, wherein the 6-(substituted amino)purine cytokinin is of formula I:



I,

wherein R₁ is selected from the group consisting of
furfuryl, phenyl, benzyl, n-alkyl of 4, 5 or 6 carbon
atoms, (cyclohexyl)methyl, (3-hydroxymethyl-3-methyl)allyl,
and 3,3-dimethylallyl.

7. A compound as defined in Claim 4 wherein the 6-(substituted amino)purine cytokinin is kinetin.

8. A compound as defined in Claim 5 wherein the 6-(substituted amino)purine cytokinin is kinetin.

-10-

9. A compound as defined in Claim 6 wherein the 6-(substituted amino)purine cytokinin is kinetin.

5 10. A commercial package comprising a container containing therein a compound according to Claim 1, 4 or 7 and written matter which states that the medicine can or should be used for treating a hyperproliferative skin disease.

10 11. A commercial package comprising a container containing therein a compound according to Claim 1, 4 or 7 and written matter which states that the medicine can or should be used for treating a hyperproliferative skin disease selected from the group consisting of basal cell carcinoma, malignant melanoma, squamous cell carcinoma, 15 actinic keratosis, Bowen's Disease, papilloma, seborrheic keratosis, toxic eczema, allergic eczema, atopic dermatitis, ichthyosis, and psoriasis.

HNR,

