



US00PP36427P2

(12) **United States Plant Patent**
Van Dijk

(10) **Patent No.:** **US PP36,427 P2**

(45) **Date of Patent:** **Feb. 4, 2025**

(54) **ANTHURIUM PLANT NAMED ‘AN2809193’**

(50) Latin Name: *Anthurium andraeanum* L.
Varietal Denomination: **AN2809193**

(71) Applicant: **ANTHURA B.V.**, Bleiswijk (NL)

(72) Inventor: **Jan Van Dijk**, Bleiswijk (NL)

(73) Assignee: **Anthura B.V.**, Bleiswijk (NL)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 342 days.

(21) Appl. No.: **17/803,690**

(22) Filed: **Oct. 11, 2022**

(51) **Int. Cl.**
A01H 5/02 (2018.01)
A01H 6/00 (2018.01)

(52) **U.S. Cl.**
USPC **Plt./365**
CPC **A01H 6/00** (2018.05)

(58) **Field of Classification Search**

USPC Plt./365

CPC A01H 5/02

See application file for complete search history.

Primary Examiner — Kent L Bell

(74) *Attorney, Agent, or Firm* — Jondle & Associates, P.C.

(57) **ABSTRACT**

A new *Anthurium* plant named ‘AN2809193’ particularly distinguished by having weakly blistered and medium glossy, red with white venation, cordate, durable spathes that retain the original color for a very long period of time with green, ovate-cordate, durable leaves, purplish-pink spadices with red tips, early and rich flowering continuously throughout the year, and a plant height of 32.0 cm to 37.0 cm is disclosed.

3 Drawing Sheets

1

Genus and species: *Anthurium andraeanum* L.
Variety denomination: ‘AN2809193’.

BACKGROUND OF THE NEW PLANT

The present invention comprises a new and distinct variety of *Anthurium*, botanically known as *Anthurium andraeanum* L., and hereinafter referred to by the variety name ‘AN2809193’. The new *Anthurium* plant is a product of a planned breeding program conducted by the inventor in Bleiswijk, the Netherlands. The objective of this breeding program was to create a new plant with a height of 32.0 cm to 37.0 cm having weakly blistered and medium glossy, red with white venation, cordate, and durable spathes.

The new variety originated from a cross-pollination made in July 2014 in Bleiswijk, the Netherlands. The female parent was a red *Anthurium* pot plant designated ‘28244-03’ (unpatented), and the male parent was an orange *Anthurium* pot plant designated ‘28516-01’ (unpatented).

A single plant was selected from the progeny of the stated cross in May 2016. Asexual reproduction of the new variety by tissue culture in 2017 in Bleiswijk, the Netherlands, has demonstrated that the new variety reproduces true to type with all of the characteristics, as herein described, firmly fixed and retained through successive generations.

Community Plant Variety Rights for this variety have been applied for in the European Union on Nov. 19, 2021 (Application no. 2021/2975), by Applicant who obtained the subject matter disclosed directly from the inventor. ‘AN2809193’ has not been made publicly available or sold anywhere in the world prior to the effective filing date of this application with the exception of sales or disclosures made one year or less before the effective filing date of this claimed invention by Applicant who obtained ‘AN2809193’ directly from the inventor.

2

SUMMARY OF THE INVENTION

The following are the most outstanding and distinguishing characteristics of this new variety when grown under normal horticultural practices in Bleiswijk, the Netherlands:

- 1) Weakly blistered and medium glossy, red with white venation, cordate, durable spathes;
- 2) Purplish-pink spadices with red tips;
- 3) Green, ovate-cordate leaves; and
- 4) Position of the spathes compared to the leaves is slightly above.

DESCRIPTION OF THE PHOTOGRAPHS

This new *Anthurium* plant is illustrated by the accompanying photographs which show the overall plant habit including blooms, buds, and foliage of the plant; the colors shown are as true as can be reasonably obtained by conventional photographic procedures. The photographs are of a 44-week-old plant grown in a greenhouse in Bleiswijk, the Netherlands, in August 2022. Colors in the photographs may differ from the color values cited in the detailed botanical description, which accurately describe the actual colors of the new variety.

FIG. 1 shows the overall plant habit, including blooms, buds, and foliage.

FIG. 2 shows a close-up of the mature spathe.

FIG. 3 shows a close-up of the upper leaf blade surface.

DESCRIPTION OF THE NEW VARIETY

The following detailed description sets forth the distinctive characteristics of ‘AN2809193’. The data which define these characteristics were collected from asexual reproductions carried out in Bleiswijk, the Netherlands. The plant history was taken on 44-week-old plants which were planted

from tissue culture in 14-centimeter (diameter) pots and grown in a glass greenhouse between 19° C. and 24° C. Observations were made in August 2022. Color readings were taken under 5000 lux natural light in the greenhouse. Color references are primarily to The R.H.S. Colour Chart of The Royal Horticultural Society of London (R.H.S.) (2015).

DETAILED BOTANICAL DESCRIPTION

Classification:

Family.—Araceae.

Botanical.—*Anthurium andraeanum* L.

Common name.—*Anthurium*.

Denomination.—‘AN2809193’.

Parentage:

Female parent.—*Anthurium* plant ‘28244-03’ (unpatented).

Male parent.—*Anthurium* plant ‘28516-01’ (unpatented).

Plant:

Propagation.—Tissue culture.

Root description.—Fleshy-creamy (RHS 161D) colored roots with small hairy lateral roots having yellow (RHS 6A) colored root tips.

Time to produce a finished flowering plant.—42 to 46 weeks after planting in a 14- cm (diameter) pot.

Growth habit.—Upright.

Height (measured from soil, including inflorescence).—32.0 cm to 37.0 cm.

Width (measured from leaf tips).—34.0 cm to 36.0 cm.

Leaves:

Immature leaves.—Length: 15.0 cm to 17.0 cm. Width: 9.0 cm to 10.0 cm. Color: Upper surface: RHS 146A.

Lower surface: RHS 146B. Texture (both upper and lower surfaces): Glossy, leathery, and thin.

Mature leaves.—Length (fully expanded): 17.0 cm to 19.0 cm. Width: 10.0 cm to 12.0 cm. Shape: Ovate cordate. Apex: Acuminate. Base: Cordate. Leaf blade angle with the petiole: Between 100 degrees and 120 degrees. Leaf margin: Entire. Color: Upper surface: RHS 147A. Lower surface: RHS 146B. Texture (both upper and lower surfaces): Glossy, leathery, and thick. Venation: Pinnate veining; the mid-vein and primary veins (the veins that radiate out from the junction of petiole and leaf) protrude at the underside of the leaf blade. Venation color: Upper surface: RHS 144A. Lower surface: RHS 144B.

Lobes.—Present. Arrangement: Leaf blade has two lobes extending past the petiole. The lobes are non-touching. Length of lobes of mature leaf blades: 1.5 cm to 2.0 cm. Width of lobes of mature leaf blades: 3.5 cm to 4.5 cm. Distance from petiole/leaf junction to highest point on lobes of mature leaf blades: 3.0 cm to 4.0 cm.

Petiole.—Cross-section: Round. Diameter: 0.4 cm to 0.5 cm. Length: 16.0 cm to 18.0 cm for a mature leaf size. Color: Mature leaf: RHS 144A. Immature leaf: RHS 144B. Cataphyll color surrounding the petiole: Outside: RHS 144A. Inside: RHS 145C.

Geniculum.—Length: 1.5 cm to 2.0 cm. Width: 0.4 cm to 0.5 cm. Color: RHS 144B.

Inflorescence:

Arrangement.—Single.

Flowering habit (length of flowering season).—Continuous.

Number of inflorescences per plant.—7 to 9.

Fragrance.—Absent.

Longevity of inflorescence on plant.—Over a year.

Spathes:

Buds.—The spathe is tightly rolled around the spadix and extrudes from the peduncle sheath. After the spathe is fully opened the peduncle elongates some extra centimeters.

Arrangement.—Spathe angle with the peduncle is between 90 degrees and 110 degrees; the spathe stands on a wiry peduncle about 1.0 cm to 4.0 cm above the foliage.

Shape.—Cordate.

Apex.—Acuminate.

Base.—Cordate.

Texture.—Weakly blistered and medium glossy.

Margin.—Weakly undulated.

Size.—Length: 10.0 cm to 11.0 cm. Width: 9.5 cm to 10.5 cm.

Lobes.—Present. Arrangement: The spathe has two lobes extending past the peduncle. The lobes are non-touching. Length: 1.0 cm to 1.5 cm. Width: 3.5 cm to 4.5 cm.

Color.—Just fully open: Upper surface: RHS 45B with white (RHS N155C) venation. Lower surface: A color in between RHS 51B and RHS 51C. This red color remains for a very long period, at least more than 30 weeks after opening. The spathe turns green after some weeks.

Peduncle:

Shape.—Erect.

Cross-section.—Round.

Length.—24.0 cm to 27.0 cm.

Diameter.—0.3 cm to 0.4 cm.

Color.—RHS 144B.

Flowering time:

General.—One small, rooted, untreated tissue culture plant of 8.0 cm tall will flower, depending on the season, after 42 to 44 weeks and 7 to 8 blossoms appear. More blossoms appear after some additional weeks so that a full flowering and commercial plant will have 8 to 9 red spathes. Smaller blossoms may occur on immature plants.

Spadix:

Size.—Length: 4.5 cm to 5.0 cm (depending on flower size). Width (at apex): 0.6 cm to 0.7 cm. Width (at base): 0.7 cm to 0.8 cm.

Shape.—Columnar.

Angle of spadix tip with peduncle.—150 degrees to 170 degrees.

Texture.—When the spathe is unfurling, the spadix is smooth. When the spadix matures, small stigmata protrude. The stigmata are evenly distributed around the spadix. The spadix matures from base to top, slowly giving the spadix a somewhat rough appearance.

Color.—Immature: RHS 45C. Mature: RHS 186D. Ages to: RHS N199A.

Flowers:

Quantity per spadix.—150 to 200.

Spadix flower arrangement.—Bisexual, rounded in cross-section.

Shape.—Rounded.

Size.—Length: 0.05 cm to 0.10 cm. Diameter (maximum): 0.10 cm.

Color.—RHS 69D.
 Reproductive organs:
Stamens.—Not visible.
Pollen amount.—Absent.
Pistil.—Quantity: 150 to 200. Length: Less than 0.01 5
 cm. Color: RHS 69D.
Style.—Not observed to date.
Stigma.—Shape: Ovoid. Diameter: Less than 0.01 cm.
 Color: RHS 69D.
Ovary.—Rarely visible.
Ovary color.—Not measured.
 Fruit and seed set: None observed to date.
 Disease and pest resistance: No specific resistance or sus-
 ceptibility observed to pathogens or pests common to
Anthurium under commercial conditions to date.

COMPARISON WITH PARENTAL AND
 SIMILAR COMMERCIAL VARIETIES

‘AN2809193’ differs from the female parent plant 20
 ‘28244-03’ (unpatented) in that ‘AN2809193’ has ovate-
 cordate leaves and longer spathes, whereas ‘28244-03’ has
 deltoid leaves and shorter spathes.
 ‘AN2809193’ differs from the male parent plant ‘28516-
 01’ (unpatented) in that ‘AN2809193’ has weakly blistered, 25

venation pattern spathes having a main color of red on the
 upper side of the spathes, whereas ‘28516-01’ has medium
 blistered, even pattern spathes having a main color of orange
 on the upper side of the spathes.

‘AN2809193’ differs from the similar commercial variety
 ‘ANTHFYTWAL’ (U.S. Plant Pat. No. 29,774) in that
 ‘AN2809193’ has poor shoot formation, ovate-cordate leaf
 blades, and the shape of the leaves in the cross section of the
 middle zone is weakly concave, whereas ‘ANTHFYTWAL’
 10 has rich shoot formation, deltoid leaf blades, and the shape
 of the leaves in the cross section of the middle zone is
 strongly concave.

‘AN2809193’ differs from the similar commercial variety
 ‘ANTHGNOBIC’ (U.S. Plant Pat. No. 34,430) in that
 15 ‘AN2809193’ has green cataphylls, purplish-pink spadices
 at the base, with medium spathes, whereas ‘ANTHGN-
 OBIC’ has red cataphylls, creamy-white spadices at the base,
 with small spathes.

I claim:

1. A new and distinct variety of *Anthurium* plant named
 ‘AN2809193’, substantially as illustrated and described
 herein.

* * * * *



FIG. 1




FIG. 2




FIG. 3