



(51) International Patent Classification:

A61K 39/12 (2006.01) C07K 14/18 (2006.01)
A61P 31/14 (2006.01) C07K 16/10 (2006.01)
A61P 37/04 (2006.01) G01N 33/53 (2006.01)
C07K 14/08 (2006.01)

(21) International Application Number:

PCT/US2016/015074

(22) International Filing Date:

27 January 2016 (27.01.2016)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/113,811 9 February 2015 (09.02.2015) US

(71) Applicant: **ACADEMIA SINICA**; 128 Sec. 2, Academia Road, Nankang, Taipei, Taiwan, Republic of China, 11529 (CN).

(71) Applicant (for CL only): **LIANG, Chi-Ming** [US/US]; 6627, River Road, Bethesda, Maryland 20817 (US).

(72) Inventors: **WU, Han-Chung**; 12F.-1, No.10, Lane 43, Sanmin Rd., Songshan District, Taipei City, Taiwan, Republic of China, 105 (CN). **TANG, Chung-Tao**; 1F., No.14, Ln. 127, Xiufeng St., Zhonghe Dist., New Taipei City, Taiwan, Republic of China, 235 (CN).

(74) Agent: **SAUNDERS, Hsiu-Ming**; Ipc Intellectual Property Connections, Inc., 3346 Ramona Street, Palo Alto, California 94306 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

- with international search report (Art. 21(3))
- with sequence listing part of description (Rule 5.2(a))

(54) Title: AN EPITOPE-SUBSTITUTED VACCINE FOR USE IN IMPROVING SAFETY AND IMMUNOGENICITY AGAINST DENGUE VIRUSES

(57) Abstract: Isolated mutant dengue virus E protein variants are disclosed. The variant comprises an amino acid sequence that is at least 80% identical to SEQ ID NO: 1 and has one or more amino acid residue substitutions at position corresponding to Asn8 (N8), Arg9 (R9), Val12 (V12) and/or Glu13 (E13). The variant may comprise an amino acid sequence that is at least 90% identical to the SEQ ID NO: 1 and lack an infection-enhancing antibody-binding motif comprising the amino acid sequence of SEQ ID NO: 28 at domain I. An isolated nucleic acid sequence encoding the variant, a plasmid expressing the variant, a plasmid expressing a virus-like particle comprising the variant, a DNA vaccine, and a method of detecting the presence of a dengue virus in a biological sample are also disclosed.



AN EPITOPE-SUBSTITUTED VACCINE FOR USE IN IMPROVING SAFETY AND IMMUNOGENICITY AGAINST DENGUE VIRUSES

FIELD OF THE INVENTION

The present invention relates generally to vaccines, and more specifically to vaccines for
5 dengue viruses.

BACKGROUND OF THE INVENTION

Dengue virus (DENV), a global disease, is divided into four serotypes (DENV1-4). Cross-reactive
and non-neutralizing antibodies against envelope protein (E) of DENV bind to the Fcγ receptors
(FcγR) of cells, and thereby exacerbate viral infection by heterologous serotypes via antibody-
10 dependent enhancement (ADE). Identification and modification of enhancing epitopes may mitigate
enhancement of DENV infection.

Therefore, identification of B-cell epitopes of DENV E protein, which induce cross-reactive and
non-neutralizing antibodies, may provide valuable information for vaccine development. A safe and
effective vaccine against DENV is not yet available. Thus, there is a need to identify and substitute
15 the epitopes recognized by poorly neutralizing and highly enhancing antibodies to improve dengue
vaccines.

SUMMARY OF THE INVENTION

In one aspect, the invention relates to an isolated mutant dengue virus E protein variant
comprising an amino acid sequence that is at least 80% identical to SEQ ID NO: 1 and having one or
20 more amino acid residue substitutions at position corresponding to Asn8 (N8), Arg9 (R9), Val12
(V12) and/or Glu13 (E13).

In another aspect, the invention relates to an isolated mutant dengue virus E protein variant,
wherein the nonmutated wild-type dengue virus E protein comprises the amino acid sequence of
SEQ ID NO: 1, the variant comprising an amino acid sequence that is at least 90% identical to the
25 SEQ ID NO: 1 and lacking an infection-enhancing antibody-binding motif comprising the amino
acid sequence of SEQ ID NO: 28 at domain I.

In one embodiment of the invention, the variant comprises one or more mutations selected from
the group consisting of: (a) a substitution of the 8th Asn residue; (b) a substitution of the 9th Arg
residue; (c) a substitution of the 12th Val residue; and (d) a substitution of the 13th Glu residue.

30 In one embodiment of the invention, the mutant dengue virus E protein variant has Asparagine at
position 8 being substituted. For example, the variant has a substitution of Asn for Ala, Glu, or Arg.

In another aspect, the invention relates to an isolated nucleic acid sequence encoding a mutant
dengue virus E protein variant as aforementioned.

In another aspect, the invention relates to a plasmid expressing a mutant dengue virus E protein variant as aforementioned.

In another aspect, the invention relates to a plasmid expressing a virus-like particle, the virus-like particle comprising a mutant dengue virus E protein variant as aforementioned.

5 In one embodiment of the invention, the aforementioned plasmid comprises (a) an isolated nucleic sequence comprising the nucleotide sequence of SEQ ID NO: 27; and (b) a promoter operably linked to the isolated nucleic sequence.

10 In another aspect, the invention relates to a DNA vaccine comprising: (a) a recombinant DNA comprising a nucleotide sequence encoding the mutant dengue virus E protein variant as aforementioned and a promoter operably linked to the nucleotide sequence; and (b) gold or tungsten, wherein the recombinant DNA and gold or tungsten forms a complex.

15 In another aspect, the invention relates to immunized human sera comprising antibodies, the antibodies exhibiting the following characteristics: (a) having specific binding affinity to the wild-type dengue virus E protein and the mutant dengue virus E protein variant as aforementioned; (b) possessing neutralizing and protective activities against dengue virus infection; (c) free of antibody dependent enhancement of dengue virus infection; (d) the antibodies being elicited by the mutant dengue virus E protein variant; and (e) having no binding activity to dengue virus infection-enhancing epitopes Asn8, Arg9, Val12 and/or Glu13 located within domain I of the wild-type dengue virus E protein.

20 In another aspect, the invention relates to use of immunized sera as aforementioned or use of a mutant dengue virus E protein variant as aforementioned in the manufacture of a medicament for eliciting neutralizing and protecting activity against one or more dengue virus serotypes in a subject in need thereof, and/or in the manufacture of a medicament for neutralizing a dengue virus, reducing, alleviating antibody dependent enhancement of dengue virus infection, and/or increasing survival rate in a subject in need thereof.

25 In one embodiment of the invention, the medicament is for alleviating the antibody dependent enhancement in a subject whose serum sample exhibits reactivity with an isolated peptide comprising the amino acid sequence of SEQ ID NO: 28.

30 Alternatively, the invention relates to immunized sera as aforementioned or a mutant dengue virus E protein variant as aforementioned for use in eliciting neutralizing and protecting activity against one or more dengue virus serotypes in a subject in need thereof, or for use in neutralizing a dengue virus, reducing, alleviating antibody dependent enhancement of dengue virus infection, and/or increasing survival rate in a subject in need thereof.

Alternatively, the invention relates to a method of eliciting neutralizing and protecting activity against one or more dengue virus serotypes in a subject in need thereof, or a method of neutralizing a dengue virus, reducing, alleviating antibody dependent enhancement of dengue virus infection, and/or increasing survival rate in a subject in need thereof, comprising administering the immunized
5 sera as aforementioned or a mutant dengue virus E protein variant as aforementioned to the subject in need thereof.

In one embodiment of the invention, immunized sera or a mutant dengue virus E protein variant as aforementioned is for use in alleviating the antibody dependent enhancement in a subject whose serum sample exhibits reactivity with an isolated peptide comprising the amino acid sequence of
10 SEQ ID NO: 28.

In one embodiment of the invention, the isolated peptide is less than 15 amino acid residues in length, and is at least 80%, 85%, 90%, 95%, or 100% identical to an amino acid sequence selected from the group consisting of SEQ ID NOs: 2-12, and 14-26.

Further in another aspect, the invention relates to a method of detecting the presence of a dengue
15 virus in a biological sample, comprising:

(a) contacting the biological sample with a composition comprising a mutant dengue virus E protein variant as aforementioned or an isolated peptide comprising the amino acid sequence of SEQ ID NO: 28;

(b) allowing dengue virus antibodies in the biological sample and the variant or the peptide to
20 form a complex; and

(c) detecting the presence of the dengue virus antibodies in the complex.

Yet in another aspect, the invention relates to a method of detecting the presence of dengue virus infection-enhancing antibodies in a biological sample, comprising:

(a) contacting the biological sample with a composition comprising an isolated peptide
25 comprising the amino acid sequence of SEQ ID NO: 28;

(b) allowing dengue virus infection-enhancing antibodies in the sample and the peptide to form a complex; and

(c) detecting the presence of the infection-enhancing antibodies in the complex.

The biological sample may be selected from the group consisting of: blood, serum, plasma,
30 saliva, cerebrospinal fluid, and urine.

These and other aspects will become apparent from the following description of the preferred embodiment taken in conjunction with the following drawings.

The accompanying drawings illustrate one or more embodiments of the invention and, together with the written description, serve to explain the principles of the invention. Wherever possible, the

same reference numbers are used throughout the drawings to refer to the same or like elements of an embodiment.

BRIEF DESCRIPTION OF THE DRAWINGS

FIGs. 1A-B show ADE of DENV infection mediated by cross-reactive mAbs. (A) Infection of K562 cells with DENV1 (MOI=1), DENV2 (MOI=1), DENV3 (MOI=5), or DENV4 (MOI=1) in the presence of serially-diluted mAbs was examined at 72 hours post-infection by staining with 4G2 and RPE-conjugated goat anti-mouse IgG, followed by flow cytometry. The percentages of infected K562 cells are shown. (B) The indicated mAbs were serially diluted and incubated with DENV2 (MOI=1 or 10) at 4°C for 1 hour. The mixtures were then added to THP-1 cells, and the infected cells were examined by flow cytometry at 72 hours post-infection. The infected cells were stained with hDB32-6 and RPE-conjugated goat anti-human IgG. The percentage of infected THP-1 cells is shown. NMIG was used as a control. Data shown in FIGs. A and B are the mean $\pm$ SEM from three independent experiments.

FIGs. 2A-E show characterization of *in vivo* ADE and competition of patient sera by DB21-6 and DB39-2 mAbs. (A) AG129 mice were i.v. infected on day 0 with 1×10^5 pfu DENV2 S221, and given i.p. injections of 5 μ g of DB21-6 (n=8) or isotype control (n=5) on days -1 and 1. Survival rates were recorded for 30 days. (** P <0.001, compared to isotype control). (B) AG129 mice were i.v. infected on day 0 with 1×10^5 pfu DENV2 S221, and given i.p. injections of 5 μ g of DB39-2 (n=7) or isotype control (n=6) on days -1 and 1. Survival rates were recorded for 30 days. (** P <0.01, compared to isotype control). Data shown in FIGs. 2A and 2B are from one representative experiment of two independent experiments. (C-D) The viremia was measured in DENV2 S221-infected AG129 mice following treatment with DB21-6 or DB39-2 mAbs using quantitative RT-PCR. The viral loads were significantly increased at day 4 and 5 post-infection. Data shown are the mean $\pm$ SD. (** P <0.01, *** P <0.001) (E) Competition between mAbs and patient sera for binding to DENV2. (* P <0.05, *** P <0.001, NS not significant).

FIGs. 3A-B show screening of a phage-displayed peptide library with DB21-6 and DB39-2 mAbs. (A) After 3 rounds of biopanning, phage titers were increased by 12,871-fold (DB21-6) and 5,000-fold (DB39-2), respectively. (B) The immunopositive phage clones selected by DB21-6 and DB39-2 were identified by ELISA. NMIG was used as a negative control.

FIGs. 4A-D show identification of the epitope residues of DB21-6 and DB39-2 mAbs. (A) BHK-21 cells were transfected with vectors expressing WT or mutant DENV2 E protein. After 2 days, the collected cells were reacted with mAbs, and then analyzed with flow cytometry. The relative indices were measured. Substitutions of N8, R9, V12, and E13 caused a significant loss of binding activity of DB21-6. Substitutions of N8, R9, and E13 caused a loss of binding activity of DB39-2. 4G2 was

used as a control. Data shown are from one representative experiment of two independent experiments. (B) The model is based on the DENV2 E protein model (PDB: 1OAN) with the positions shown from the bottom of the structure. The epitopes of cross-reactive mAbs are located at residues N8, R9, V12, and E13 in EDI. (C) Structure analysis determines the distance (°A) between epitope residues from the same (shaded) or adjacent monomer by PYMOL™ program. (D) The secreted VLPs were captured by polyclonal rabbit anti-DENV2 E protein sera. The WT and mutant VLPs were detected with DB32-6 and 4G2 by VLP-capture ELISA. Data shown are from one representative experiment of two independent experiments.

FIGs. 5A-E show *in vitro* and *in vivo* neutralization assays of DENV2 with immunized sera. (A) Mice were immunized with vector, WT, or N8R plasmids at three-week intervals. After the 3rd immunization, serum samples were collected and pooled for each immunized group. The immunized sera were evaluated by ELISA using plates containing C6/36 cells infected with DENV2 16681. The OD was measured at 490 nm. (***P*<0.01, NS not significant). (B) After the 3rd immunization, the pooled sera were serially diluted and incubated with DENV2 (MOI=1) for 1 hour at 4°C. The mixtures were then used to infect BHK-21 cells. After 3 days, the cells were stained with 4G2, and analyzed by flow cytometry. (n=3) (C) Table showing titers conferring 50% inhibition against DENV2 infection. (D and E) After the 3rd immunization, the pooled sera were serially diluted and incubated with 25-fold LD₅₀ of DENV2 for 0.5 hour at 4 °C . The mixtures were injected into ICR suckling mice by the intracerebral route (i.c.). The survival rates were recorded for 28 days. The number of animals tested for each immunized sera ranged from 4 to 16 per group. (****P*<0.001, ***P*<0.01, compared to vector-immunized sera). Data shown in FIGs. 5A, 5B, 5D and 5E are from one representative experiment of two independent experiments.

FIGs. 6A-F show *in vivo* enhancement of mortality in AG129 mice treated with immunized sera. (A) AG129 mice were i.v. infected with 1×10^5 pfu DENV2 S221 on day 0, and given i.p. injections of WT- (n=4), N8R- (n=4), or vector- (n=4) immunized sera (1:25 dilution) on days -1 and 1. The survival rates of mice treated with WT- (*P*=0.3834) or N8R- (*P*=0.1278) immunized sera did not significantly differ from that of mice treated with vector-immunized sera. (B) AG129 mice were i.v. infected with 1×10^5 pfu DENV2 (S221) on day 0, and given i.p. injections of WT- (n=5), N8R- (n=5), or vector- (n=7) immunized sera (1:100 dilution) on days -1 and 1. The survival rate of mice treated with WT- (****P*=0.0007) immunized sera was significantly lower than that of mice treated with vector-immunized sera. However, the survival rate of mice treated with N8R- (*P*=0.3538) immunized sera did not differ from that of mice treated with vector-immunized sera. (C) AG129 mice were i.v. infected with 1×10^5 pfu DENV2 (S221) on day 0, and given i.p. injections with WT- (n=7), N8R- (n=7), or vector- (n=6) immunized sera (1:400 dilution) on days -1 and 1. The survival

rate of mice treated with WT- ($P=0.8701$) or N8R- ($P=0.1587$) immunized sera did not significantly differ from that of mice treated with vector-immunized sera. (D) Schematic describing the competition assay of mAbs and immunized sera. The ELISA plates were coated with polyclonal rabbit anti-DENV hyper-immune sera at 4°C overnight. After blocking, the diluted DENV2 viral supernatants were added for 2 hours at RT. The HRP-conjugated enhancing mAbs (Innova Biosciences HRP Conjugation Kit) and immunized sera (1:40 dilution) were incubated for 2 hours at RT. After washing, the OPD was added, and the reaction was stopped with 3N HCl. The OD was measured at 490 nm. (E-F) Competition between mAbs and immunized sera for binding to DENV2. Normal mouse serum (NMS) was used as a control. The percentage of competition was calculated as follows: competition (%) = $[1 - (\text{OD of immunized serum-mAb mixture} / \text{OD of NMS-mAb mixture})] \times 100$. ($***P<0.001$)

FIGs. 7A-C show characterization of DB21-6 and DB39-2 against DENV and E protein of DENV2 by immunofluorescence assay (IFA). (A) BHK-21 cells were infected with DENV1 (MOI=1), DENV2 (MOI=1), DENV3 (MOI=10), or DENV4 (MOI=1). After 2 days, the infected cells were detected using DB21-6 and DB39-2. Uninfected cells (Mock) and NMIGG were used as negative controls. The results are shown at 400× magnification. (B and C) DENV2 E, comprising amino acids 1-400 of the E protein, was cloned into the pcDNA3.1 plasmid. DENV2 EDI-II (amino acids 1-295) and EDIII (amino acids 295-400) were also respectively inserted into the pcDNA3.1 plasmid. After transfection, IFA was used to reveal that DB21-6 and DB39-2 recognized BHK-21 cells expressing DENV2 E (B) or EDI-II (C) protein.

FIGs. 8A-B show evaluation of neutralizing activity of mAbs against DENV1-4. (A) The neutralizing activity of DB21-6, DB39-2, or 4G2 against DENV1-4 was examined with inhibition assays. DENV1 (MOI=5), DENV2 (MOI=1), DENV3 (MOI=5), or DENV4 (MOI=1) was incubated with mAbs at 4°C for 1 hour, and then used to infect BHK-21 cells. After 3 days, the cells were fixed and stained with 4G2. Titers are expressed as inhibition percentages. Data shown are from one representative experiment of two independent experiments. (B) Summary of 50% inhibition concentrations of DB21-6, DB39-2, and 4G2 against DENV1-4.

FIGs. 9A-B show *in vitro* measurement of mAb-mediated ADE of DENV2 S221 infection in K562 (A) and THP-1 (B) cells. DENV2 S221 was incubated with dilutions of mAbs for 1 hour at 4°C, and the resulting mixture was used to infect K562 or THP-1 cells. After 3 days, the cells were stained with 4G2, and analyzed by flow cytometry.

FIGs. 10A-B show competition assay using mAbs and patient sera. Competition for binding to DENV2 between mAbs and patient sera at a dilution of 1:50 (A) or 1:200 (B).

FIGs. 11A-D show evaluation of humoral immune responses against DENV2 by ELISA. (A) Mice were immunized with vector control, WT, or N8R plasmids at three-week intervals. The serum samples were collected after one, two, and three immunizations, and pooled. The sera were evaluated using plates containing C6/36 cells infected with DENV2 16681. (B) Collected sera
 5 diluted 1:200 and detected with anti-IgG1 or IgG2a antibodies. (C) IgG1/IgG2a ratios. (D) Immunized sera collected and examined after the 3rd immunization (***P*<0.001)

FIGs. 12A-B show neutralization of infections with different DENV2 strains. (A) BHK-21 cells were infected with DENV2 16681, PL046, Malaysia 07587, or NGC at an MOI of 1, 1, 1, or 10, respectively, at 37°C for 2 hours. After 3 days, the cells were fixed, stained with 4G2, and RPE-
 10 conjugated goat anti-mouse IgG. The percentages of infected cells were analyzed by flow cytometry. BHK-21 cells were used as a negative control (Mock). PE, phycoerythrin. FSC, forward scatter. (B) Serial dilutions of 3rd immunized pooled sera were incubated with DENV2 16681, PL046, Malaysia 07587, or NGC at an MOI of 1, 1, 1, or 10, respectively, at 4°C for 1 hour. The resulting mixtures were then used to infect BHK-21 cells. After 3 days, the cells were stained with 4G2, and analyzed
 15 by flow cytometry.

FIG. 13 shows a vector map of N8R plasmid. CMV: cytomegalovirus promoter. JEV SS: signal peptide sequence of Japanese encephalitis virus (JEV). JEV-E: Envelope (E) protein sequence of JEV. The nucleotide sequence of SEQ ID NO: 27 contains regions of JEV SS and prM (nucleotides 1-570), N8R (nucleotides 571-1755), and JEV-E (nucleotides 1756-2058).

20 DETAILED DESCRIPTION OF THE INVENTION

The present invention is more particularly described in the following examples that are intended as illustrative only since numerous modifications and variations therein will be apparent to those skilled in the art. Various embodiments of the invention are now described in detail. Referring to the drawings, like numbers indicate like components throughout the views. As used in the
 25 description herein and throughout the claims that follow, the meaning of “a”, “an”, and “the” includes plural reference unless the context clearly dictates otherwise. Also, as used in the description herein and throughout the claims that follow, the meaning of “in” includes “in” and “on” unless the context clearly dictates otherwise. Moreover, titles or subtitles may be used in the specification for the convenience of a reader, which shall have no influence on the scope of the
 30 present invention. Additionally, some terms used in this specification are more specifically defined below.

DEFINITIONS

The terms used in this specification generally have their ordinary meanings in the art, within the context of the invention, and in the specific context where each term is used. Certain terms that

are used to describe the invention are discussed below, or elsewhere in the specification, to provide additional guidance to the practitioner regarding the description of the invention. For convenience, certain terms may be highlighted, for example using italics and/or quotation marks. The use of highlighting has no influence on the scope and meaning of a term; the scope and meaning of a term is the same, in the same context, whether or not it is highlighted. It will be appreciated that same thing can be said in more than one way. Consequently, alternative language and synonyms may be used for any one or more of the terms discussed herein, nor is any special significance to be placed upon whether or not a term is elaborated or discussed herein. Synonyms for certain terms are provided. A recital of one or more synonyms does not exclude the use of other synonyms. The use of examples anywhere in this specification including examples of any terms discussed herein is illustrative only, and in no way limits the scope and meaning of the invention or of any exemplified term. Likewise, the invention is not limited to various embodiments given in this specification.

Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention pertains. In the case of conflict, the present document, including definitions will control.

As used herein, “around”, “about” or “approximately” shall generally mean within 20 percent, preferably within 10 percent, and more preferably within 5 percent of a given value or range. Numerical quantities given herein are approximate, meaning that the term “around”, “about” or “approximately” can be inferred if not expressly stated.

The term “treating” or “treatment” refers to administration of an effective amount of a therapeutic agent to a subject in need thereof, who has a disease, or a symptom or predisposition toward such a disease, with the purpose of cure, alleviate, relieve, remedy, ameliorate, or prevent the disease, the symptoms of it, or the predisposition towards it, or reduce incidence of symptoms. Such a subject can be identified by a health care professional based on results from any suitable diagnostic method.

“An effective amount” refers to the amount of an active compound that is required to confer a therapeutic effect on the treated subject. Effective doses will vary, as recognized by those skilled in the art, depending on route of administration, excipient usage, and the possibility of co-usage with other therapeutic treatment.

The “Guidance for Industry and Reviewers Estimating the Safe Starting Dose in Clinical Trials for Therapeutics in Adult Healthy Volunteers” published by the U.S. Department of Health and Human Services Food and Drug Administration discloses “a human equivalent dose” may be obtained by calculations from the following formula:

$$\text{HED} = \text{animal dose in mg/kg} \times (\text{animal weight in kg/human weight in kg})^{0.33}.$$

As used herein, when a number or a range is recited, ordinary skill in the art understand it intends to encompass an appropriate, reasonable range for the particular field related to the invention.

Detection reagent or labeling agent for labeling an antibody has been disclosed in US8,920,804, which is incorporated herein by reference in its entirety.

The amino acid 1-395 sequence of wild-type DENV2 (16681) E protein (The GenBank *accession number*: AAB58782) is shown in SEQ ID NO: 1.

Abbreviations: virus-like particles (VLPs); dengue fever (DF); dengue hemorrhagic fever (DHF); E protein domain I (EDI); antibody-dependent enhancement (ADE); glutamic acid (Glu; E); valine (Val; V); asparagine (Asn; N); arginine (Arg; R); pre-membrane (prM); envelope (E) proteins; multiplicity of infection (MOI).

We characterized the cross-reactive DB21-6 and DB39-2 monoclonal antibodies (mAbs) against domain I-II of DENV; these antibodies poorly neutralized and potently enhanced DENV infection both *in vitro* and *in vivo*. Two enhancing mAbs, DB21-6 and DB39-2, were observed to compete with sera antibodies from patients infected with dengue. The epitopes of these enhancing mAbs were identified. N8, R9, V12, and E13 are the reactive residues of DB21-6, while N8, R9, and E13 are the reactive residues of DB39-2. N8 substitution tended to maintain VLP secretion, and decrease the binding activity of DB21-6 and DB39-2. The immunized sera from N8 substitution (N8R) DNA vaccine exerted greater neutralizing and protecting activity than wild-type (WT) immunized sera, both *in vitro* and *in vivo*. Treatment with N8R-immunized sera reduced the enhancement of mortality in AG129 mice. These results support identification and substitution of enhancing epitope as a novel strategy for developing safe dengue vaccines.

EXAMPLES

Exemplary instruments, apparatus, methods and their related results according to the embodiments of the present invention are given below.

Materials and Methods

DENV, cell lines, and mAbs

Four dengue virus serotypes, DENV1 Hawaii, DENV2 16681, DENV3 H87, and DENV4 H241, were prepared as described in Wu et al. (2003) ("Identification of a dengue virus type 2 (DEN-2) serotype-specific B-cell epitope and detection of DEN-2-immunized animal serum samples using an epitope-based peptide antigen" *J Gen Virol* 84, 2771-2779). C6/36 cells were grown in medium having 50% Mitsumashi and Maramorsch insect medium (SIGMA-ALDRICH®) plus 50% Dulbecco's modified Eagle's medium (DMEM) containing 10% fetal bovine serum (FBS) and 100 U/ml penicillin, 100 µg/ml streptomycin, and 0.25 µg/ml amphotericin B. The C6/36 cells were infected with DENV at a multiplicity of infection (MOI) of 0.1-1, and incubated at 28°C for 7 to 9

days. The viruses were harvested from supernatant and titrated in a baby hamster kidney fibroblast cell line (BHK-21) by plaque assay. The aliquots were stored at -80°C. BHK-21 cells were grown in minimal essential medium (MEM) supplemented with 10% FBS, 100 U/ml penicillin, 100 µg/ml streptomycin, and 0.25 µg/ml amphotericin B. Human erythroleukaemic K562 and monocytic THP-1
 5 cells were grown in RPMI medium containing 10% FBS. The mouse mAbs, including DB21-6 and DB39-2, were previously generated by immunization of BALB/c mice with DENV2, and produced in hybridoma cells (Li et al., (2012). "Development of a Humanized Antibody with High Therapeutic Potential against Dengue Virus Type 2" PLoS Negl Trop Dis 6, e1636). DB21-6 and DB39-2 were isotypized as IgG1 and purified using PROTEIN G SEPHAROSE™ 4B gels (GE Healthcare).

10 *In vitro measurement of ADE with mAbs*

Serial dilutions of mAbs were incubated with DENV1 Hawaii (MOI=1), DENV2 16681 (MOI=1), DENV3 H87 (MOI=5), and DENV4 H241 (MOI=1) for 1 hour at 4°C. The mixtures were used to infect K562 cells for 2 hours at 37°C. After washing, the cells were incubated with 2% FBS in RPMI medium at 37°C for 3 days. The infected cells were collected and fixed with 3.7%
 15 formaldehyde for 10 minutes at 4°C. For staining, the cells were permeabilized with 2% FBS in PBS containing 0.1% saponin (Sigma), followed by staining with 4 µg/ml 4G2 for 0.5 hours at 4°C. The cells were washed and incubated with R-phycoerythrin (RPE)-conjugated goat anti-mouse IgG (Jackson ImmunoResearch Laboratories) for 1 hour at 4°C. The cells were washed, and the percentages of infected cells were determined by flow cytometry. For infection of THP-1 cells,
 20 DENV2 16681 (MOI=1 or 10) was incubated with diluted mAbs for 1 hour at 4°C, and then incubated with cells for 2 hours at 37°C. After 3 days, the cells were fixed, permeabilized, and stained with hDB32-6. After washing, the cells were incubated with an RPE-conjugated goat anti-human IgG, and were subsequently analyzed by flow cytometry.

Measurement of in vivo ADE with mAbs in AG129 mice

25 Type I and II interferon receptor-deficient mice (AG129; 5- to 6-weeks-old) were purchased from B&K Universal. The AG129 mice were given 5 µg mAbs in 200 µl PBS on days 1 and -1 (i.p.). The mouse IgG1 isotype antibody was used as a negative control. On day 0, mice were inoculated (i.v.) with 1×10^5 pfu of the mouse-adapted DENV2 S221 (obtained from S. Shrestha) in 100 µl PBS. The survival rates of AG129 mice were recorded for 30 days.

30 *Measurement of viremia by quantitative RT-PCR*

AG129 mice were infected with 1×10^5 pfu DENV2 S221 (i.v.) on day 0, and treated with 5 µg mAbs via i.p. on days -1 and 1. Viral RNA was extracted from pooled and infected mice sera using QIAAMP® viral RNA minikit (Qiagen). Quantitative RT-PCR was performed, using LIGHTCYCLER® 480 system (ROCHE™). The standard curve was generated with DENV2 S221

(at concentrations from 10^1 to 10^7 pfu/ml). Viremia measurements were expressed as pfu equivalents/ml, which was calculated based on the threshold cycle value (Ct) according to the standard curve for DENV2 S221.

Competitive ELISA of mAbs and patient serum samples

A total of 21 DENV2-infected patient serum samples were collected from 11 DF and 10 DHF patients during an outbreak between 2002 and 2003 in Taiwan. Diagnosis of DENV infection was based on IgM antibody-capture ELISA (MAC-ELISA), reverse-transcriptase PCR (RT-PCR), or virus isolation in cell cultures. These serum samples were collected between days 4 and 22 from the onset of symptoms; such sera contained anti-dengue antibodies. All of these patients were determined to have classical DF or DHF based on the criteria published by the World Health Organization (WHO) in 2009. Table 2 shows the characteristics of patient serum samples enrolled in this study. Competition ELISA was performed. Briefly, the plates were coated with polyclonal rabbit anti-DENV hyper-immune sera at 4°C overnight. After blocking, the diluted DENV2 viral supernatants (1×10^6 pfu) were added for 2 hours at room temperature (R.T.). The diluted mAbs and patient sera (1:100 dilution) were incubated for 2 hours at R.T. After washing, horseradish peroxidase (HRP)-conjugated anti-mouse IgG (Jackson ImmunoResearch Laboratories) was added for 1 hour at R.T. The peroxidase substrate *o*-phenylenediamine dihydrochloride (OPD, SIGMA-ALDRICH™) was then added, and the reaction stopped with 3N HCl. The optical density (OD) was measured at 490 nm. Normal human serum (NHS) was used as a control. The percentage of competition was calculated as follows: competition (%) = $[1 - (\text{OD of patient serum-mAb mixture} / \text{OD of NHS-mAb mixture})] \times 100$.

Phage display biopanning

Phage display biopanning was performed. Briefly, the plate was coated with 100 µg/ml mAbs at 4°C for 6 hours. After washing and blocking, 4×10^{10} pfu of phage-displayed peptide library (New England BioLabs, Inc.) were incubated for 50 mins at R.T. After washing, bound phage was eluted with 100 µl 0.2 M glycine/HCl (pH 2.2) and neutralized with 15 µl 1 M Tris/HCl (pH 9.1). The eluted phage was then amplified in ER2738 for subsequent rounds of selection. The phage was titrated onto LB medium plates containing IPTG and X-Gal. The second and third rounds of selection were identical to the first round except for the addition of 2×10^{11} pfu of amplified phage.

Identification of immunopositive phage clones by ELISA

Plates were coated with 50 µg/ml mAbs. After washing and blocking, the amplified phage was added, and incubated for 1 hour at R.T. After washing, diluted HRP-conjugated anti-M13 antibody (GE Healthcare) was added at RT for 1 hour. The plates were developed and subsequently terminated by 3N HCl. The optical density (OD) was measured at 490 nm.

Identification of epitopes using flow cytometry-based binding assay to cells expressing WT and mutant DENV2 prM/E proteins

The pCBD2-2J-2-9-1 plasmid expressing prM-E proteins of DENV2 has been previously characterized and described (Chang et al., (2003) "Enhancing biosynthesis and secretion of premembrane and envelope proteins by the chimeric plasmid of dengue virus type 2 and Japanese encephalitis virus" *Virology* 306, 170-180). Site-directed mutagenesis was performed to replace each of the selected amino acid residues. After mutagenesis, the plasmids were sequenced to ensure the absence of any further mutations at non-target sites. BHK-21 cells were transfected with constructs expressing the wild-type (WT) or mutant DENV2 E protein using POLYJET™ *in vitro* DNA transfection reagent (SignGen Laboratories). After 2 days, the cells were fixed, and permeabilized with 2% FBS in PBS containing 0.1% saponin (SIGMA™). For staining, cells were incubated with DB21-6, DB39-2, 4G2, and mixed mAbs (DB32-6, 3H5, and DB25-2) at a concentration of 1, 1, 1 and 1 µg/ml, respectively, at 4°C for 0.5 hours. After washing, the cells were incubated with RPE-conjugated goat anti-mouse IgG, and analyzed by flow cytometry. The relative index of a mAb to a mutant E protein was measured using the formula: [intensity of the mutant E/intensity of WT E (recognized by a mAb)]/[intensity of mutant E/intensity of WT E (recognized by mixed mAbs)].

Detection of secreted VLPs by capture ELISA

BHK-21 cells were transfected with vectors expressing WT or mutant E protein of DENV2, as described above. At 48 hours post-transfection, culture supernatants were collected. The plates were coated with polyclonal rabbit anti-DENV hyper-immune sera at 4°C overnight. After blocking, two-fold dilutions of supernatants containing WT or mutant VLPs were added for 2 hour at RT. The wells were then incubated with diluted DB32-6 and 4G2 at RT for 2 hour. After washing, a 1:2000 dilution of HRP-conjugated anti-mouse IgG was added for 1 hour at RT. Finally, the plates were developed, and the reaction was subsequently terminated with 3N HCl. The OD was measured at 490 nm.

Preparation of plasmids for immunization

Plasmids expressing WT E protein (SEQ ID NO: 1) of DENV2 or a mutant E protein in which the N8 residue was substituted with R (N8R) were used for immunization (FIG. 13). For coating, 25 mg of 1.0 µm gold powder was resuspended with 50 mM spermidine. Then, 50 µg of plasmid DNA was added, followed by the addition of 1M CaCl₂; the solution was mixed and precipitated for 10 mins at RT. After collection by centrifugation, the gold-DNA complex was washed with absolute ethanol and resuspended in 0.1 mg/ml of polyvinylpyrrolidone (PVP) (360 kDa) solution. The slurry was injected into a TefzelR tube (MCMASTER-CARR™, Chicago, IL), and then coated. After the ethanol had dried off, the tube was cut into 0.5-inch bullets and stored at -20°C. The gold in each

bullet contained 1 µg of DNA. Before use, the bullets were loaded into the Helios gene gun device (Bio-Rad, Hercules, CA) for delivery of plasmids.

Immunization of mice

The abdominal epidermis of 6 week-old female BALB/c mice was injected with a gene gun using a helium pressure setting of 400 lb/inch². Each mouse was immunized by administering 4 bullets containing 1 µg plasmid DNA. Mice were immunized at 0, 3, and 6 weeks. Serum samples were collected before immunization and 3 weeks after the third immunization (pre-, 1st, 2nd, 3rd immunized sera). The serum samples were pooled from five to six mice for each group and evaluated by ELISA, neutralization assay, and *in vivo* ADE assay.

Evaluation of immunized sera against DENV2 by ELISA

C6/36 cells infected with DENV2 16681 were used as antigens. C6/36 cells were seeded into each well (2×10^4 cells/well) of 96-well ELISA plates. After one day, 2×10^3 pfu of DENV2 16681 (MOI=0.1) was added to infect the cells at 37°C for 2 hours. The wells were washed with PBS, and then cultured in 2% FBS culture medium at 28°C for 5 days. The infected cells were fixed with 1:1 methanol/acetone at 4°C for 10 mins. The plates were blocked with 5% skimmed milk at 4°C for 24 hours. Diluted immunized sera were then added for incubation at RT for 2 hours. The plates were washed three times with phosphate-buffered saline containing 0.1% (w/v) TWEEN® 20 (PBST_{0.1}), and subsequently incubated with HRP-conjugated anti-mouse IgG. Finally, the plates were developed, and the reaction terminated with 3N HCl. The OD was measured at 490 nm.

In vitro and in vivo neutralization assays with immunized sera

DENV2 16681 (MOI=1) was incubated with the 3rd immunized sera for 1 hour at 4°C. The mixtures were used to infect BHK-21 cells for 2 hours at 37°C. After 3 days, the cells were fixed, permeabilized, and stained with 4 µg/ml 4G2. After washing, the cells were incubated with RPE-conjugated goat anti-mouse IgG, and analyzed by flow cytometry. Inhibition percentage (%) = $[1 - (\text{the percentage of infected cells incubated with immunized sera/without immunized sera})] \times 100$.

The ICR mice were purchased from the Laboratory Animal Center, National Taiwan University College of Medicine. Serially-diluted immunized sera were incubated with 1×10^4 pfu (25-fold lethal dose, 25-fold LD₅₀) of DENV2 16681 for 0.5 hours at 4°C. Two-day-old suckling mice were inoculated with 20 µl of the mixtures through intracranial (i.c.) injection. After challenge, the survival rates were recorded for 28 days.

Studies of in vivo ADE with immunized sera

AG129 mice were given (i.p.) dilutions of immunized sera on days -1 and 1, and inoculated (i.v.) with 1×10^5 pfu of DENV2 S221 on day 0. The survival rates were recorded for 30 days.

Statistical analysis

Survival rate was expressed using Kaplan-Meier survival curves, and statistical analyses were performed using GraphPad Prism 5. For competition assays of mAbs and patient sera, Student's *t* tests were used to identify significant differences and calculate *P* values (**P*<0.05, ****P*<0.001, NS not significant). For evaluation of immunized sera against DENV2 by ELISA, two-way ANOVA with Bonferroni *post-hoc* test was used to determine the significant differences and calculate *P* values (***P*<0.01, NS not significant). GRAPHPAD PRISM® 5 was used to analyze 50% inhibition titers against DENV2, based on inhibition percentages from pooled immunized sera.

Results

Characterization of cross-reactive DB21-6 and DB39-2 against DENV

FIG. 7A shows mAbs DB21-6 and DB39-2 could recognize cells infected with DENV1-4. These mAbs also recognized transfected BHK-21 cells expressing DENV2 E and EDI-II proteins, respectively (FIGs. 7B and 7C). Thus, the cross-reactive DB21-6 and DB39-2 recognized DENV1-4 and domain I-II on E protein.

To estimate the *in vitro* neutralizing activity of these mAbs, we infected BHK-21 cells with a mixture of individual mAbs and DENV1-4. FIGs. 8A-B show that 4G2, an anti-flavivirus antibody with neutralizing and enhancing activity at certain concentrations, exerted higher neutralization activity than DB21-6 and DB39-2 against DENV2. FIG. 8B, a Table showing a summary of 50% inhibition concentration of mAbs against DENV1-4, shows that DB21-6 and DB39-2 exhibited non-neutralizing activity against DENV1-4 (50% inhibition concentration >33 µg/ml).

Enhancing activities of DB21-6 and DB39-2

To investigate *in vitro* enhancement of DENV infection through ADE, we performed *in vitro* ADE assays, and detected the increases in the percentage of dengue-infected cells by flow cytometry. The FcγRIIA-bearing K562 cells, which do not express type 1 interferon (IFN), were used to measure the enhancement of infected cells through extrinsic ADE. The serially-diluted mAbs were incubated with DENV1-4, and then used to infect K562 cells. The infection percentage was measured by flow cytometry. FIG. 1A shows infection enhancement over a broad range of mAb concentrations. 4G2 caused enhancement of DENV1-4 infection in K562 cells at lower antibody concentrations. DB21-6 and DB39-2 enhanced DENV1-4 infection in K562 cells at high antibody concentrations.

To further confirm enhancement of infection, we examined the enhancement of DENV2 16681 infection by DB21-6 and DB39-2 in FcγRI- and FcγRIIA-bearing THP-1 cells. Infection in THP-1 cells was enhanced to a greater extent by DB21-6 and DB39-2 than by 4G2 (FIG. 1B).

DENV2 S221 was previously used to study enhancement of mortality via ADE in AG129 mice. To evaluate the *in vitro* enhancement of DENV2 S221 infection by mAbs, we performed ADE

assays using K562 cells and THP-1 cells. As for DENV1-4 infection, high concentrations of DB21-6 and DB39-2 enhanced DENV2 S221 infection in K562 cells (FIG. 9A). DB21-6 and DB39-2 enhanced DENV2 S221 infection in THP-1 cells at high concentrations of antibody (FIGs. 9A-B). These results suggest that DB21-6 and DB39-2 can enhance DENV2 S221 infection *in vitro*.

5 We confirmed the *in vivo* enhancing activities in AG129 mice. The AG129 mice treated with 5 μ g DB21-6 and infected with DENV2 S221 exhibited increased mortality as compared to control infected mice (FIG. 2A). AG129 mice treated with 5 μ g of DB39-2 also exhibited elevated mortality (FIG. 2B). To determine viremia in DENV2 S221-infected AG129 mice following treatment with DB21-6 or DB39-2, the viral RNA levels were measured by quantitative RT-PCR. The results
10 indicate that viral loads were significantly increased after DB21-6 or DB39-2 treatment of infected AG129 mice, as compared to isotype control Ab treatment (FIGs. 2C-D). The results indicate that DB21-6 and DB39-2 have non-neutralizing activities, and enhance mortality in AG129 mice.

Competition assay of mAbs and infected patient sera

We performed competition assay to determine whether sera antibodies from dengue patients
15 compete with mAbs for binding to DENV2. Table 2 shows the characteristics of patient serum samples. The results indicate that the sera antibodies from infected patients competed with DB21-6 and DB39-2. FIG. 2E shows that the competition percentages of DB21-6 and DB39-2 were significantly higher in serum samples from DHF patients than those from DF patients, while the competition percentage of neutralizing DB32-6 was similar for sera from either DF or DHF patients.
20 The same experiment was performed with more concentrated (1:50 dilution) or diluted serum (1:200 dilution) and obtained similar results (FIGs. 10A-B). The results suggest that serum samples from DHF patients contain higher levels of antibodies, which compete with DB21-6 and DB39-2 mAbs.

Identification of enhancing epitopes of DB21-6 and DB39-2

To identify the enhancing epitopes of DB21-6 and DB39-2, we used a phage-displayed peptide
25 library to screen for the reactive phage clones. After three biopanning rounds, the phage titers were increased to 12,871-fold (DB21-6) and 5,000-fold (DB39-2), respectively, compared to that of the first round (FIG. 3A). The individual phage clones from the third round of biopanning were randomly selected. As shown by ELISA, most selected phage clones exhibited significant reactivity to the mAbs, but not to normal mouse IgG (NMiGg). Of the 30 selected phage clones, 29 clones
30 reacted with DB21-6 (FIG. 3B). The immunopositive phage clones were amplified, and their phage DNA was isolated for DNA sequencing. Table 1 shows eleven phage clones with individual peptide sequences identified. Similarly, of the 47 selected phage clones, 46 reacted with DB39-2 (FIG. 3B). Thirteen of the 46 immunopositive phage clones that reacted with DB39-2 possess individual peptide sequences. Alignment of peptide sequences revealed the binding motif of DB21-6 and DB39-2 to be

N-R-x-x-V-E (SEQ ID NO: 28). Modeling of the peptide sequences with the pepitope server predicted that the epitope residues on the E protein are N8, R9, V12, and E13.

To further verify the epitopes of DB21-6 and DB39-2, we performed site-directed mutagenesis of the phage-displayed epitope using pCBD2-2J-2-9-1 as template. After confirmation of variants by sequencing, we transfected cells with mutant plasmids, and detected binding activity by flow cytometry. The binding percentages for each transfectant were normalized to those of anti-EDIII mAbs (DB32-6, 3H5, and DB25-2), and relative indices were calculated and as shown in FIG. 4A. 4G2, which binds to residues at the fusion-loop of EDII, was used as a control to verify the structural change of E proteins caused by mutations. Based on the relative indices, we found that mutations at N8, R9, V12, and E13 prevented binding by DB21-6. The same method was used to identify the epitope residues of DB39-2 as N8, R9, and E13. Structural modeling was applied to show that the recognition residues are located in domain I of E protein (FIG. 4B). The distance between these residues from the same monomers was analyzed using a structure modeling program, and was found to be less than 30°A (FIG. 4C). This distance can be spanned by a single IgG molecule. It suggests that the N8, R9, V12, and E13 residues constitute the epitope of DB21-6, and the N8, R9, and E13 residues constitute the epitope of DB39-2. Alignments revealed that the binding motif of DB21-6 and DB39-2 corresponds to the N8, R9, V12, and E13 residues, which are conserved in DENV1-4 (Tables 3 and 4). Finally, we used VLP-capture ELISA to demonstrate that the mutations at R9, V12, and E13 affect DENV2 VLP secretion (FIG. 4D). The effects of these mutations on the ability to secrete VLPs might be due to a change in the structure of E protein. However, the N8R substitution did not affect DENV2 VLP secretion (FIG. 4D). N8 substitution tends to maintain VLP secretion and reduces the binding activity of DB21-6 and DB39-2.

Examination of humoral immune responses in mice

The BALB/c mice were immunized with vector, WT, or N8R plasmids at 0, 3, and 6 weeks. After three rounds of immunization, the serum samples were collected and pooled within each group. The immunized sera were examined by ELISA. FIG. 11A shows a remarkable increase of antibody titer against DENV2 was observed after immunization. The 3rd WT- and N8R-immunized sera against DENV2 exhibited significantly higher absorbance values than those of vector-immunized sera (FIG. 5A). Analysis of immunized sera with anti-IgG1 and IgG2a antibodies revealed that the IgG1/IgG2a ratios increased between the second and third immunization (FIGs. 11B and 11C). The immunized mice maintained their anti-DENV2 responses after 15 weeks (FIG. 11D).

Evaluation of neutralizing activity of the immunized sera

The immunized sera were evaluated for neutralizing activity against DENV2. Both WT- and N8R-immunized sera exhibited high neutralizing activities, while vector-immunized sera did not

(FIG. 5B). DENV2 infection was more effectively neutralized by N8R-immunized sera than by WT-immunized sera (FIG. 5C). To further evaluate whether immunized sera could broadly neutralize the diverse DENV2 strains, BHK-21 cells were infected with mixtures of immunized sera and four different DENV2 strains: 16681, NGC, PL046, and Malaysia 07587. Remarkably, FIGs. 12A-B show that the WT- and N8R-immunized sera exhibited high neutralizing activities against various types of DENV2 strain.

The protective effect of immunized sera against DENV2 16681 *in vivo* was examined. The survival rates of mice treated with WT-immunized sera at dilutions of 1:100 and 1:200 were significantly higher than that of mice treated with vector-immunized sera at a dilution of 1:100 (FIG. 5D), while the survival rates of mice treated with N8R-immunized sera at dilutions of 1:100, 1:200, and 1:400 were significantly higher than that of mice treated with vector-immunized sera at a dilution of 1:100 (FIG. 5E). Treatment with WT-immunized sera afforded 50% protection at a dilution of 1:200, while N8R-immunized sera afforded 50% protection at a dilution of 1:400 (FIGs. 5D and 5E). Hence, N8R-immunized sera possessed higher neutralizing and protective activity than WT-immunized sera both *in vitro* and *in vivo*.

Reduction of in vivo enhancing activity of the immunized sera

To study the *in vivo* enhancement of mortality, we passively transferred different dilutions of WT-, N8R-, or vector-immunized sera into AG129 mice. Following infection with DENV2 S221, the survival rate of mice treated with WT- or N8R-immunized sera (1:25 dilution) was higher than that of mice treated with vector-immunized sera (FIG. 6A). FIG. 6B shows that mice treated with WT-immunized sera at a dilution of 1:100 exhibited higher mortality than mice treated with vector-immunized sera. The survival rate of mice treated with N8R-immunized sera at a dilution of 1:100 was higher than that of mice treated with vector-immunized sera. No enhancement of mortality was observed in mice treated with N8R-immunized sera. Treatment with WT- or N8R-immunized sera at a dilution of 1:400 did not have a neutralizing or enhancing effect on the survival rates of mice. These results indicate that the N8R substitution of E protein can reduce *in vivo* enhancement of mortality.

To further characterize these enhancing antibodies are produced in immunized sera, we performed competitive ELISA to inhibit the binding of HRP-conjugated DB21-6 or DB39-2 mAbs by immunized sera (FIG. 6D). The competition percentages of HRP-conjugated DB21-6 and DB39-2 were significantly higher in WT-immunized sera than those in N8R-immunized sera (FIGs. 6E-F). These results suggest that N8R substitution would redirect immunodominance by reducing the generation of enhancing antibodies.

We characterized the ability of DB21-6 and DB39-2 to increase the percentage of dengue virus-infected cells. These mAbs enhance mortality in AG129 mice. The sera antibodies from infected patients compete with these mAbs for binding. We mapped the epitopes of enhancing mAbs DB21-6 and DB39-2 on EDI protein. To investigate how to reduce the enhancing effects while maintaining neutralizing activity, we substituted the N8 residue of E protein, and immunized mice with WT or N8R plasmids with a gene gun delivery system. After three immunizations, N8R-immunized sera produced neutralizing activity against DENV2, and reduced enhancement of mortality as compared to WT-immunized sera. Thus, substitution of enhancing epitope residues can increase the immune response against viral infection while reducing the potential for ADE.

We demonstrated that cross-reactive DB21-6 and DB39-2 against EDI-II had poor neutralizing activities against DENV1-4 (FIGs. 8A-B). DB21-6 and DB39-2 have strong ADE activities *in vitro* (FIG. 1). 4G2 has partially neutralizing activity against DENV1-4 (FIGs. 8A-B) and enhances *in vitro* viral infections at low antibody concentrations (FIG. 1A). DB21-6 and DB39-2 enhanced DENV1-4 infection in K562 cells at high concentrations (FIG. 1A). Infection of DENV2 was enhanced to a greater extent by DB21-6 and DB39-2 than by 4G2 in THP-1 cells (FIG. 1B). DB21-6 and DB39-2 enhanced mortality in AG129 mice (FIGs. 2A and 2B) and increased the viral loads in infected mice sera (FIG. 2C-D). These results indicate that DB21-6 and DB39-2 have strong enhancing activity both *in vitro* and *in vivo*.

ADE is regarded as an important mechanism leading to the development of severe dengue disease, including DHF/DSS. Cross-reactive and non-neutralizing antibodies binding to viruses can enhance infection of FcγR-bearing cells by ADE, resulting in increased viral load and/or production of cytokines. High viral load is correlated with dengue disease severity and DHF. Thus, there is a need to be able to confirm the presence of enhancing antibodies in dengue patient sera. Our results indicate that the competition percentages of DB21-6 and DB39-2 were significantly higher in DHF patient sera than those in DF patient sera (FIG. 2E), suggesting that the higher levels of enhancing antibodies, DB21-6 and DB39-2, in serum samples of dengue patients are associated with severe dengue disease. We hypothesize that the DENV infected patients might suffer more severe symptoms, such as DHF, when the expression level of the enhancing antibodies is higher.

The phage clones selected using DB21-6 and DB39-2 mAbs displayed peptide sequences containing a consensus motif, N-R-x-x-V-E (Table 1). These displayed peptide sequences may be suitable for detecting enhancing antibodies in serum samples from dengue patients, and for providing information on the pathogenesis of dengue. By alignment of displayed peptide sequences and structural modeling, the candidate epitopes were predicted and verified using VLP mutants (FIGs. 3 and 4). The epitope residues of enhancing mAb DB21-6 are N8, R9, V12, and E13 in domain I of

DENV2 E protein (FIGs. 4A and 4B), and the epitope residues of enhancing mAb DB39-2 are N8, R9, and E13 in domain I of DENV2 E protein. The N8, R9, V12, and E13 residues were conserved in DENV1-4. Thus, cross-reactive DB21-6 and DB39-2 can bind to DENV1-4. A previous report indicated that G106 and L107 are the epitope residues of enhancing mAb 4G2. The data in FIG. 4A confirmed that W101, G106, L107, and F108 in the fusion loop are the epitope residues of 4G2. The epitope residues recognized by 4G2 are different from those recognized by DB21-6 and DB39-2. The enhancing epitopes of DB21-6 (N8, R9, V12, and E13) and DB39-2 (N8, R9, and E13) are novel and have not previously been reported.

The N8R substitution does not affect DENV2 VLP secretion (FIG. 4D). Both WT- and N8R-immunized sera exerted protective activities against DENV2. N8R-immunized sera had higher *in vitro* neutralizing activity and *in vivo* protective activity than the WT-immunized sera (FIGs. 5B-5E). These results suggest that immunization with the N8R DNA vaccine may increase neutralizing and protective immunity against DENV2.

We passively transferred diluted vector-, WT-, or N8R-immunized sera, and then challenged AG129 mice with DENV2 S221. WT- and N8R-immunized sera were protective at a 1:25 dilution, as compared to vector-immunized sera. The mortality of mice was enhanced by treatment with WT-immunized sera at a 1:100 dilution, as compared to treatment with vector-immunized sera. The mortality of mice treated with N8R-immunized sera at a 1:100 dilution was not enhanced (FIG. 6B). When the dilution was increased to 1:400, no enhanced mortality was observed (FIG. 6C). Our results indicate that substituting the enhancing epitope can reduce the ADE phenomenon and increase protective activity *in vivo*.

Substitution of enhancing epitope and preservation of neutralizing epitope in immunized mice provide protective immunity. Such an approach would redirect immunodominance (FIGs. 6E-F) and improve immunogenicity by satisfying the required neutralizing occupancy. In summary, we have identified novel enhancing epitopes and illustrated, through N8R substitution in DENV2 E protein, a useful way to enabling us to reduce the potential for ADE. This may be a viable approach for developing new dengue vaccines that can increase the anti-DENV immune response.

Table 1 shows alignment of phage-displayed peptide sequences selected by DB21-6 and DB39-2. Table 2 shows the DENV2-infected patient serum samples used. Table 3 shows comparison of the amino acid sequences of EDI-II proteins of DENV1, 2, 3, and 4^a. Table 4 shows the database, gene / protein and accession / ID numbers.

Table 1

DB21-6 phage clones	Peptide sequences ^a (SEQ ID NO:)
PC21-15	N G S N R D I V E V Q R (2)

PC21-10		N Q I Y N R D Y T E P T	(3)
PC21-9		Y N R D M L E T D Y V	N(4)
PC21-13		Q N T W N R D S I E E T	(5)
PC21-17	F P E V S V N R L V V E		(6)
PC21-20		H V N R L H V E G P V	P(7)
PC21-18	K M T L P M N R S H V E		(8)
PC21-2	S Y V T G G N R Y A V E		(9)
PC21-1	S S Y L S N R L F T E A		(10)
PC21-16	S A T T M S N R Y Y T E		(11)
PC21-5		Q P Y N R S Y I D F M V	(12)
DENV1-4 ^b		N ⁸ R D F V E ¹³	(13)
DB39-2	phage clones	Peptide sequences ^a (SEQ ID NO:)	
DB39-38		L S N R L H V E S L E L	(14)
DB39-40		N Q T N R H F V E I V H	(15)
DB39-11	S G L D R N R Q L V E R		(16)
DB39-39		N R T L V E L G Y A M L	(17)
DB39-3		V N R P W V E T T T Q G	(18)
DB39-28	I V P Y S N R T V T E T		(19)
DB39-31		N R V S N E P F W D I A	(20)
DB39-34		D Y L N R S T N E P A L	(21)
DB39-36	S M P L S G R A V V E G		(22)
DB39-47	H T S L H S G R N S V E		(23)
DB39-4	S S P G V I S R F L V E		(24)
DB39-43		D R Y L V E Y S S G R W	(25)
DB39-1	M P S G G R F L V E G A		(26)
DENV1-4 ^b		N ⁸ R D F V E ¹³	(13)

^a The phage-displayed consensus amino acids are indicated by boldface type.

^b The amino acid sequences 8 to 13 in E protein of DENV1-4 were retrieved from GenBank (accession number A1U47321, AAB58782, AAA99437, and AAX48017).

Table 2

Patient number	Severity ^a	Days post onset of symptoms	Primary /Secondary infection
1	DHF	17	Primary
2	DHF	22	Primary
3	DHF	5	Secondary
4	DHF	6	Secondary
5	DHF	7	Secondary
6	DHF	8	Secondary
7	DHF	4	Secondary
8	DHF	7	Secondary
9	DHF	8	Secondary
10	DHF	4	Secondary
11	DF	6	Primary
12	DF	10	Primary
13	DF	11	Primary
14	DF	7	Primary
15	DF	6	Secondary

16	DF	6	Secondary
17	DF	5	Secondary
18	DF	7	Secondary
19	DF	7	Secondary
20	DF	7	Secondary
21	DF	4	Secondary

^aDHF, dengue hemorrhagic fever; DF, dengue fever.

Table 3

Serotypes of DENV (strain)	Accession number (GenBank)	Amino acid of EDI-II protein			
		8 ^{b, c}	9 ^{b, c}	12 ^b	13 ^{b, c}
DENV2 (16681)	AAB58782	N	R	V	E
DENV2 (NGC)	AAA42941	N	R	V	E
DENV2 (PL046)	AHZ61501	N	R	V	E
DENV2 (PM33974)	ABO33322	N	R	V	E
DENV2 (IQT2913)	AAD32963	N	R	V	E
DENV2 (ThD2_0038_74)	ABA61185	N	R	V	E
DENV2 (ThD2_0168_79)	ABA61184	N	R	V	E
DENV2 (ThD2_0498_84)	ABA61183	N	R	V	E
DENV2 (ThD2_0263_95)	ABA61179	N	R	V	E
DENV2 (ThD2_0017_98)	ABA61178	N	R	V	E
DENV2 (ThD2_0284_90)	ABA61180	N	R	V	E
DENV2 (Jamaica/N.1409)	AAA42942	N	R	V	E
DENV2 (TSV01)	AAK67712	N	R	V	E
DENV2 (98900663 DHFBAD42415 DV-2)		N	R	V	E
DENV2 (Tonga/74)	AAV70829	N	R	V	E
DENV2 (I348600)	AAW31413	N	R	V	E
DENV1 (Hawaii)	AIU47321	N	R	V	E
DENV1 (16007)	AAF59977	N	R	V	E
DENV3 (ThD3_0183_85)	AAV88402	N	R	V	E
DENV3 (H87)	AAA99437	N	R	V	E
DENV4 (H241)	AAX48017	N	R	V	E
DENV4 (B5)	AAG30148	N	R	V	E
DENV4 (ThD4_0348_91)	AAU89377	N	R	V	E
DENV4 (ThD4_0087_77)	AAU89378	N	R	V	E
DENV4 (ThD4_0485_01)	AAU89379	N	R	V	E

DENV4 (ThD4_0734_00)	AAU89380	N	R	V	E
DENV4 (814669)	AAK01233	N	R	V	E
DENV4 (Indonesia 1976)	AAB70680	N	R	V	E
DENV4 (ThD4_0476_97)	AAU89375	N	R	V	E

^aEDI-II protein of strains of four dengue virus serotypes are aligned. Single letter amino acid abbreviations are shown.

^{b,c}The critical residues are shown in boldface type and identified by loss of binding for DB21-6^b and DB39-2^c, respectively.

Table 4

Gene / Protein	Database	Accession / ID number
DENV2 E protein	Protein Data Bank	1OAN
DENV2 (16681)	GenBank	AAB58782
DENV2 (NGC)	GenBank	AAA42941
DENV2 (PL046)	GenBank	AHZ61501
DENV2 (PM33974)	GenBank	ABO33322
DENV2 (IQT2913)	GenBank	AAD32963
DENV2 (ThD2_0038_74)	GenBank	ABA61185
DENV2 (ThD2_0168_79)	GenBank	ABA61184
DENV2 (ThD2_0498_84)	GenBank	ABA61183
DENV2 (ThD2_0263_95)	GenBank	ABA61179
DENV2 (ThD2_0017_98)	GenBank	ABA61178
DENV2 (ThD2_0284_90)	GenBank	ABA61180
DENV2 (Jamaica/N.1409)	GenBank	AAA42942
DENV2 (TSV01)	GenBank	AAK67712
DENV2 (98900663 DHF DV-2)	GenBank	BAD42415
DENV2 (Tonga/74)	GenBank	AAV70829
DENV2 (1348600)	GenBank	AAW31413
DENV1 (Hawaii)	GenBank	AIU47321
DENV1 (16007)	GenBank	AAF59977
DENV3 (ThD3_0183_85)	GenBank	AAV88402
DENV3 (H87)	GenBank	AAA99437
DENV4 (H241)	GenBank	AAX48017
DENV4 (B5)	GenBank	AAG30148
DENV4 (ThD4_0348_91)	GenBank	AAU89377
DENV4 (ThD4_0087_77)	GenBank	AAU89378
DENV4 (ThD4_0485_01)	GenBank	AAU89379
DENV4 (ThD4_0734_00)	GenBank	AAU89380
DENV4 (814669)	GenBank	AAK01233
DENV4 (Indonesia 1976)	GenBank	AAB70680
DENV4 (ThD4_0476_97)	GenBank	AAU89375

We found that the cross-reactive mAbs DB21-6 and DB39-2 exhibit poor neutralizing activity and high capacity for enhancing DENV infection. We identified the epitopes recognized by DB21-6 and DB39-2. To further improve the DNA vaccines against DENV2, in the illustrated example we

substituted the N8 residue of wild-type (WT) DENV2 E protein with arginine (N8R) in a plasmid for immunization. N8R-immunized sera produced higher neutralizing and protective activity than WT-immunized sera. Treatment of AG129 mice with N8R-immunized sera reduced ADE and mortality, as compared with mice treated with WT-immunized sera. In conclusion, we have identified a novel cross-reactive and infection-enhancing epitope in E protein, and demonstrated that substitution of this enhancing epitope is a promising strategy for development of a safe dengue vaccine.

All references cited and discussed in this specification are incorporated herein by reference in their entireties and to the same extent as if each reference was individually incorporated by reference.

CLAIMS

What is claimed is:

1. An isolated mutant dengue virus E protein variant, comprising an amino acid sequence that is at least 80% identical to SEQ ID NO: 1 and having one or more amino acid residue
5 substitutions at position corresponding to Asn8, Arg9, Val12, and/or Glu13.
2. An isolated mutant dengue virus E protein variant, wherein the nonmutated wild-type dengue virus E protein comprises the amino acid sequence of SEQ ID NO: 1, the variant comprising an amino acid sequence that is at least 90% identical to the SEQ ID NO: 1 and lacking an infection-enhancing antibody-binding motif comprising the amino acid sequence of SEQ ID
10 NO: 28 at domain I.
3. The variant of claim 2, comprising one or more mutations selected from the group consisting of:
 - (a) a substitution of the 8th Asn residue;
 - (b) a substitution of the 9th Arg residue;
 - 15 (c) a substitution of the 12th Val residue; and
 - (d) a substitution of the 13th Glu residue.
4. A mutant dengue virus E protein variant as claimed in any one of claims 1-3, wherein the variant has Asparagine at position 8 being substituted.
5. An isolated nucleic acid sequence encoding a mutant dengue virus E protein variant as
20 claimed in any one of claims 1-3.
6. A plasmid expressing a mutant dengue virus E protein variant as claimed in any one of claims 1-3.
7. A plasmid expressing a virus-like particle, the virus-like particle comprising a mutant dengue virus E protein variant as claimed in any one of claims 1-3.
- 25 8. The plasmid of claim 7, comprising:
 - (a) an isolated nucleic sequence comprising the nucleotide sequence of SEQ ID NO: 27; and
 - (b) a promoter operably linked to the isolated nucleic sequence.
9. A DNA vaccine comprising:
 - (a) a recombinant DNA comprising a nucleotide sequence encoding the mutant dengue virus
30 E protein variant of claim 4 and a promoter operably linked to the nucleotide sequence; and
 - (b) gold or tungsten, wherein the recombinant DNA and gold or tungsten forms a complex.
10. Immunized human sera comprising antibodies, the antibodies exhibiting the following characteristics:

- (a) having specific binding affinity to the wild-type dengue virus E protein and the mutant dengue virus E protein variant as claimed in any one of claims 1-3;
 - (b) possessing neutralizing and protective activities against dengue virus infection;
 - (c) free of antibody dependent enhancement of dengue virus infection;
 - 5 (d) the antibodies being elicited by the mutant dengue virus E protein variant; and
 - (e) having no binding activity to dengue virus infection-enhancing epitopes Asn8, Arg9, Val12 and/or Glu13 located within domain I of the wild-type dengue virus E protein.
11. Use of immunized sera as claimed in claim 10 or a mutant dengue virus E protein variant as claimed in any one of claims 1-3 in the manufacture of a medicament for eliciting
- 10 neutralizing and protecting activity against one or more dengue virus serotypes in a subject in need thereof, and/or in the manufacture of a medicament for neutralizing a dengue virus, reducing, alleviating antibody dependent enhancement of dengue virus infection, and/or increasing survival rate in a subject in need thereof.
12. The use of claim 11, wherein the medicament is for alleviating the antibody dependent
- 15 enhancement in a subject whose serum sample exhibits reactivity with an isolated peptide comprising the amino acid sequence of SEQ ID NO: 28.
13. The use of claim 12, wherein the isolated peptide is less than 15 amino acid residues in length, and is at least 80% identical to an amino acid sequence selected from the group consisting of SEQ ID NOs: 2-12, and 14-26.
- 20 14. A method of detecting the presence of a dengue virus in a biological sample, comprising:
- (a) contacting the biological sample with a composition comprising a mutant dengue virus E protein variant as claimed in any one of claims 1-3 or an isolated peptide comprising the amino acid sequence of SEQ ID NO: 28;
 - (b) allowing dengue virus antibodies in the biological sample and the variant or the peptide to
 - 25 form a complex; and
 - (c) detecting the presence of the dengue virus antibodies in the complex.
15. A method of detecting the presence of dengue virus infection-enhancing antibodies in a biological sample, comprising:
- (a) contacting the biological sample with a composition comprising an isolated peptide
 - 30 comprising the amino acid sequence of SEQ ID NO: 28;
 - (b) allowing dengue virus infection-enhancing antibodies in the sample and the peptide to form a complex; and
 - (c) detecting the presence of the infection-enhancing antibodies in the complex.

FIG. 1A




FIG. 1B




FIG. 2A




FIG. 2B




FIG. 2C




FIG. 2D




FIG. 2E




FIG. 3A





FIG. 3B



4/11

FIG. 4A




FIG. 4B




FIG. 4C

Distance °A	N8	R9	V12	E13
N8	70.68	7.19	12.31	16.51
R9	81.74	88.68	8.79	12.33
V12	81.4	88.37	92.66	5.67
E13	88.77	93.73	96.93	101.38

FIG. 4D




FIG. 5A




FIG. 5B




FIG. 5C

Groups	The 50% inhibition titers against DENV2 (mean $\pm$ SD)
Vector	<25
WT	859.4 $\pm$ 83.32
N8R	1425 $\pm$ 109.3

FIG. 5D




FIG. 5E




FIG. 6A

1:25 dilution




FIG. 6B

1:100 dilution




FIG. 6C

1:400 dilution




FIG. 6D




FIG. 6E

DB21-6




FIG. 6F

DB39-2




FIG. 7A




FIG. 7B





FIG. 7C



8/11

FIG. 8A




FIG. 8B

mAbs	The concentration of 50% inhibition percentage (µg/ml)			
	DENV1	DENV2	DENV3	DENV4
DB21-6	>33	>33	>33	>33
DB39-2	>33	>33	>33	>33
4G2	>33	7.1	>33	>33

FIG. 9A




FIG. 9B




FIG. 10A




FIG. 10B




FIG. 11A




FIG. 11B




FIG. 11C




FIG. 11D




FIG. 12A




FIG. 12B





FIG. 13



INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 16/15074

A. CLASSIFICATION OF SUBJECT MATTER

IPC(8) - A61K 39/12, A61P 31/14, A61P 37/04, C07K 14/08, C07K 14/18, C07K 16/10, G01N 33/53 (2016.01)

CPC - A61K 39/12, C07K 16/1081, A61K 2039/5258, A61K 2039/545, C07K 2317/21, C07K 2317/33

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC(8)- A61K 39/12, A61P 31/14, A61P 37/04, C07K 14/08, C07K 14/18, C07K 16/10, G01N 33/53 (2016.01)

CPC - A61K 39/12, C07K 16/1081, A61K 2039/5258, A61K 2039/545, C07K 2317/21, C07K 2317/33

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

CPC - C07K 2317/34, C12N 2770/24122, C07K 14/18, C12N 2770/24134, G01N 33/56983, G01N 2333/185, G01N 2469/20

USPC- 424/186.1, 530/350, 435/7.1

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

PubWEST(PGPB,USPT,USOC,EPAB,JPAB); PatBase, Google/Scholar: Dengue, DENV, E protein; STF2 protein; Viral envelope glycoprotein, B-cell epitopes of DENV E protein, antibody-dependent enhancement, ADE, AIDE, E protein domain I, EDI, DB21-6 and DB3972 mAbs, ACADEMIA SINICA...

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	de Alwis, et al. Dengue viruses are enhanced by distinct populations of serotype cross-reactive antibodies in human immune sera. PLoS Pathog. 2014, 10(10):e1004386; pg 7, Fig 5 and its legend; pg 10, col 1	15
A	Messer, et al. Dengue virus envelope protein domain I/II hinge determines long-lived serotype-specific dengue immunity. Proc Natl Acad Sci USA 2014, 111(5):1939-1944; pg 1940, Fig. 1(C) and its legend	15
A	US 2008/0248064 A1 (Chang et al.) 09 October 2008 (09.10.2008) para [0107], SEQ ID NO 14, amino acids 1-395	1-14
A	US 2011/0065686 A1 (Reyes, et al.) 17 March 2011 (17.03.2011) claims 1, 14; para [0073]	1, (4-14)/1
A	US 7,943,148 B1 (Sagripanti, et al.) 17 May 2011 (17.05.2011) col 2, ln 37-40, col 15, ln 20-37	1-14
A	WO 2014/016360 A1 (SANOFI PASTEUR) 30 January 2014 (30.01.2014) pg 47, ln 27 to pg 48, ln 10; pg 50, ln 18 to pg 51, ln 15	1, (4-14)/1
A	US 2012/0282287 A1 (Apt, et al.) 08 November 2012 (08.11.2012) para [0014]	2, 3, (4-14)/(2,3)

☒ Further documents are listed in the continuation of Box C.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

28 March 2016

Date of mailing of the international search report

19 MAY 2016

Name and mailing address of the ISA/US

Mail Stop PCT, Attn: ISA/US, Commissioner for Patents
P.O. Box 1450, Alexandria, Virginia 22313-1450

Facsimile No. 571-273-8300

Authorized officer:

Lee W. Young

PCT Helpdesk: 571-272-4300

PCT OSP: 571-272-7774

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 16/15074

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X,P ✓	Tang, et al. An Epitope-Substituted DNA Vaccine Improves Safety and Immunogenicity against Dengue Virus Type 2. PLoS Negl Trop Dis. 02 July 2015, 9(7):e0003903; in entirety	1-15

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 16/15074

Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying additional fees, this Authority did not invite payment of additional fees.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.
- ☐ The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.
- ☐ No protest accompanied the payment of additional search fees.