



US00PP11320P

United States Patent [19]

[11] Patent Number: Plant 11,320

Dümmen

[45] Date of Patent: Apr. 4, 2000

[54] NEW GUINEA IMPATIENS PLANT NAMED 'DUERIPINKEYE'

[75] Inventor: **Marga Dümmen**, Rheinberg, Germany

[73] Assignee: **Dümmen Jungpflanzenkulturen**, Rheinberg, Germany

[21] Appl. No.: **09/038,772**

[22] Filed: **Mar. 11, 1998**

[51] Int. Cl.<sup>7</sup> ..... **A01H 5/00**

[52] U.S. Cl. .... **Plt./318**

[58] Field of Search ..... Plt./317, 318

[56] **References Cited**  
U.S. PATENT DOCUMENTS

P.P. 10,255 2/1998 Neeman ..... Plt./318  
*Primary Examiner*—Howard J. Locker  
*Attorney, Agent, or Firm*—C. A. Whealy

[57] **ABSTRACT**

A new and distinct variety of New Guinea Impatiens plant named 'Dueripinkeye', characterized by its rounded flowers that are white with a bright pink eye and faint radiating stripe on the upper most petal; high degree of floriferousness; freely branching plant habit; and dark green leaves with reddish overtones that contrast well with the flower color.

**1 Drawing Sheet**

**1**

**BACKGROUND OF THE INVENTION**

The present invention relates to a new and distinct variety of New Guinea Impatiens plant, botanically known as *Impatiens hawkeri*, and hereinafter referred to by the name 'Dueripinkeye'.

The new variety is a product of a planned breeding program conducted by the inventor in Rheinberg, Germany. The objective of the breeding program is to develop New Guinea Impatiens that are freely branching; early flowering; and that have desirable flower and leaf colors.

The new cultivar originated from a cross made by the inventor of two proprietary seedling selections of *Impatiens hawkeri*. 'Dueripinkeye' was discovered and selected in 1995 by the inventor as a single flowering plant within the progeny of the stated cross in a controlled environment in Rheinberg, Germany.

Asexual reproduction of the new variety by terminal cuttings taken at Rheinberg, Germany, has shown that the unique features of this new variety are stable and reproduced true to type in successive generations of asexual reproduction.

**BRIEF SUMMARY OF THE INVENTION**

The following traits have been repeatedly observed and are determined to be the unique characteristics of 'Dueripinkeye'. These characteristics in combination distinguish 'Dueripinkeye' as a new and distinct variety:

1. Rounded flowers that are white with a bright pink eye and fainter radiating stripe on the upper petal.
2. High degree of floriferousness.
3. Freely branching plant habit.
4. Dark green leaves with reddish overtones that contrast well with the white and pink flower color.

The new variety can be compared to the commercial variety 'Samoa' (disclosed in U.S. Plant Pat. No. 8,422). However, in side-by-side comparisons conducted by the inventor in Rheinberg, Germany, plants of the new variety differed from plants of 'Samoa', in the following characteristics:

1. Plants of the new variety are taller than plants of 'Samoa'.
2. Plants of the new variety have shorter leaves than plants of 'Samoa'.

**2**

3. Leaves of plants of the new variety are dark green with reddish tones in color whereas leaves of plants of 'Samoa' are dark green.

4. Plants of the new variety are more floriferousness than plants of 'Samoa'.

5. Plants of the new variety have slightly smaller flowers than plants of 'Samoa'.

6. The pink eye covers a larger area on flowers of plants of the new variety than flowers of plants of 'Samoa'.

7. Peduncles of plants of the new variety are shorter and stronger than peduncles of plants of 'Samoa'. In addition, peduncles of plants of the new variety are dark red whereas peduncles of plants of 'Samoa' are green.

**BRIEF DESCRIPTION OF THE PHOTOGRAPH**

The accompanying colored photograph illustrates the overall appearance of the new variety, showing the colors as true as it is reasonably possible to obtain in colored reproductions of this type. The photograph comprises a top perspective view of a typical plant of 'Dueripinkeye' grown in a 12-cm container.

**DETAILED BOTANICAL DESCRIPTION**

Plants of 'Dueripinkeye' have not been observed under all possible environmental conditions. The phenotype may vary somewhat with variations in environment such as temperature, daylength and light intensity, without, however, any variance in genotype. The following observations and measurements describe plants grown in 12-cm containers during the spring in Rheinberg, Germany, under commercial practice in a glass-covered greenhouse with day and night temperatures about 18° C. and light levels about 45 thousand lux.

In the following description, color references are made to The Royal Horticultural Society Colour Chart except where general terms of ordinary dictionary significance are used.

Botanical classification: *Impatiens hawkeri* 'Dueripinkeye'.  
Parentage:

*Male, or pollen, parent*.—Proprietary seedling selection of *Impatiens hawkeri*.

*Female, or seed, parent*.—Proprietary seedling selection of *Impatiens hawkeri*.

## Propagation:

*Type cutting.*—Terminal cuttings.

*Time to initiate roots.*—Summer: About 10 days at 22° C. Winter: About 12 days at 22° C.

*Time to develop roots.*—Summer: About 21 days at 22° C. Winter: About 26 days at 22° C.

*Rooting habit.*—Freely branching.

## Plant Description:

*General appearance.*—Rounded; dense; very freely and basally branching; upright and spreading; moderate growth rate and vigor; suitable for 10 to 16-cm pots.

*Plant height.*—About 21 cm from soil level to top of plant plane.

*Lateral branches.*—Length: About 16 cm. Diameter: About 5.1 mm. Internode length: About 6.5 cm. Color: 136A.

*Foliage description.*—Quantity of leaves per lateral branch: About 14. Shape: Elliptic with acuminate tip and attenuate base. Length: About 7.8 cm. Width: About 3.4 cm. Texture: Smooth, leathery and glabrous. Margin: Serrulate. Color: Young foliage, upper surface: 131A. Young foliage, lower surface: 187B. Mature foliage, upper surface: Dark green, 131A, with reddish, close to 185A, tones. Mature foliage, lower surface: 187C. Veins, upper surface: 187B. Veins, lower surface: 187B. Petiole: Length: About 5 mm. Diameter: About 2.1 mm. Color: 185A.

## Flower description:

*Flowering habit.*—Freely flowering.

*Natural flowering season.*—Year-round. Plants flower about 8 weeks after planting.

*Flowers borne.*—Flower buds develop in apical leaf axils. Open flowers are displayed above the foliage.

*Quantity of inflorescences.*—Extremely floriferous; usually about 85 buds and flowers per plant.

*Flower shape.*—Rounded and slightly cupped in aspect.

*Flower diameter.*—About 4.8 cm.

*Flower depth (height).*—About 5.8 mm.

*Petals.*—Shape: Reniform/cordate with emarginate apex and acute base. Quantity, arrangement: 5 petals overlapping. Aspect: Flat. Length: About 2.6 cm. Width: About 2.7 cm. Texture: Smooth, satiny, and glabrous. Margin: Entire. Color: When opening, upper surface: White, 155D, background with bright pink, 64D/66D, eye with fainter radiating stripe, close to 66D, on the uppermost petal. When opening, lower surface: White, 155D, background with bright pink, 64D/66D, eye with fainter radiating stripe, close to 66D, on the uppermost petal. Fully opened, upper surface: White, 155D, background with bright pink, 64D/66D, eye with fainter radiating stripe, close to 66D, on the uppermost petal. Fully opened, lower surface: White, 155D, background with bright pink, 64D/66D, eye with fainter radiating stripe, close to 66D, on the uppermost petal.

*Peduncle.*—Angle: Erect. Length: About 3.8 cm. Color: 183B.

*Flower bud.*—Shape: Ovoid with spur. Length: About 1.5 cm. Diameter: About 9.2 mm. Color: 63C.

*Spur.*—Shape: Needle-shaped, curved at end. Length: About 4.8 cm. Color: 185B.

*Reproductive organs.*—Androecium: Stamen number: 5. Anther shape: Oval. Anther size: About 4 mm. Anther color: 64C. Pollen color: 27B. Amount of pollen: Moderate. Gynoecium: Pistil length: About 3.9 mm. Style color: 131A. Stigma color: 135A. Ovary color: 135B.

Disease resistance: No fungal, bacterial nor viral problems observed.

Seed development: Seed production is very rarely observed. It is claimed:

1. A new and distinct variety of New Guinea Impatiens plant named 'Dueripinkeye', as illustrated and described.

\* \* \* \* \*

**U.S. Patent**

**Apr. 4, 2000**

**Plant 11,320**

