



(51) International Patent Classification:

A61K 39/395 (2006.01) C07K 16/28 (2006.01)
G01N 33/15 (2006.01) C12N 5/10 (2006.01)
G01N 33/50 (2006.01) C12N 15/09 (2006.01)
C07K 14/705 (2006.01)

(21) International Application Number:

PCT/US2016/065788

(22) International Filing Date:

9 December 2016 (09.12.2016)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/267,409 15 December 2015 (15.12.2015) US

(71) Applicant: THE BOARD OF REGENTS OF THE UNIVERSITY OF TEXAS SYSTEM [US/US]; 201 West 7th Street, Austin, TX 78701 (US).

(72) Inventors: LIU, Qingyun; C/o Univ. Of Tx Health Science Center, At Houston, 7000 Fannin, Suite 1200, Houston, TX 77030 (US). GONG, Xing; C/o Univ. Of Tx Health Science Center, At Houston, 7000 Fannin, Suite 1200, Houston, TX 77030 (US). CARMON, Kendra; C/o Univ. Of Tx Health Science Center, At Houston, 7000 Fannin, Suite 1200, Houston, TX 77030 (US).

(74) Agent: BYRD, Marshall, P.; Parker Highlander PLLC, 1120 S. Capital Of Texas Highway, Building One, Suite 200, Austin, TX 78746 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH,

[Continued on next page]

(54) Title: LGR4 SPECIFIC MONOCLONAL ANTIBODIES AND METHODS OF THEIR USE




FIG. 1

(57) Abstract: Isolated or recombinant anti-LGR4 monoclonal antibodies are provided. In some cases, antibodies of the embodiments can be used for the detection, diagnosis and/or therapeutic treatment of human diseases, such as cancer.



GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

- *with international search report (Art. 21(3))*
- *before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))*
- *with sequence listing part of description (Rule 5.2(a))*

DESCRIPTION

LGR4 SPECIFIC MONOCLONAL ANTIBODIES AND METHODS OF THEIR USE

[0001] This application claims the benefit of United States Provisional Patent Application No. 62/267,409, filed December 15, 2015, the entirety of which is incorporated
5 herein by reference.

[0002] This invention was made with government support under Grant No. GM102485 awarded by the National Institutes of Health. The government has certain rights in the invention.

BACKGROUND OF THE INVENTION

1. Field of the Invention

[0003] The present invention relates generally to the field of cancer biology. More particularly, it concerns LGR4 targeting monoclonal antibodies for the treatment and detection of cancer.

2. Description of Related Art

15 [0004] The leucine-rich repeat-containing G-protein coupled receptor 4 (LGR4), also called G-protein coupled receptor 48 (GPR48), consists of a large extracellular domain (ECD) with 17 leucine-rich repeats and a seven transmembrane (7TM) domain typical of the rhodopsin family of G protein-coupled receptors. It has been demonstrated that LGR4, along with its closely related homologs LGR5 and LGR6, function as receptors of the R-spondin
20 group of stem cell factors to potentiate Wnt signaling (8-10). This function of RSPO-LGR4 signaling is mediated by the inhibition of two E3 ubiquitin ligases (RNF43 and ZNRF3) of Wnt receptors and the recruitment of IQGAP1 to coordinate LRP6 phosphorylation as well as actin
dynamics. As a receptor for R-spondins, LGR4 potentiates the canonical Wnt signaling pathway and is
25 involved in the formation of various organs. Upon binding to R-spondins (RSPO1, RSPO2, RSPO3 or RSPO4), LGR4 associates with phosphorylated LRP6 and frizzled receptors that are activated by extracellular Wnt receptors, triggering the canonical Wnt signaling pathway to increase expression of target genes. In contrast to classical G-protein coupled receptors, LGR4 does not activate heterotrimeric G-proteins to transduce the signal. LGR4 activation of
30 the Wnt signaling pathway is required for the development of various organs, including liver,

kidney, intestine, bone, reproductive tract and eye. LGR4 is also required during spermatogenesis to activate the Wnt signaling pathway in peritubular myoid cells and for the maintenance of intestinal stem cells and Paneth cell differentiation in postnatal intestinal crypts. LGR4 also acts as a regulator of bone formation and remodeling and is involved in kidney development. LGR4 is also involved in the development of the anterior segment of the eye; required during erythropoiesis; and also acts as a negative regulator of innate immunity by inhibiting TLR2/TLR4 associated pattern-recognition and proinflammatory cytokine production. LGR4 is also believed to play an important role in regulating the circadian rhythms of plasma lipids, partially through regulating the rhythmic expression of MTTP. The complex interactions of LGR4 receptors and ligands provide a great potential for significant therapeutic intervention.

SUMMARY OF THE INVENTION

[0005] Described herein are LGR4 monoclonal antibodies that potently block LGR4 signaling and inhibit cancer cell proliferation. Thus, in a first embodiment, there is provided an isolated or recombinant monoclonal antibody that specifically binds to a LGR4. In certain aspects, an antibody that competes for the binding of a LGR4 with the 8D2 or 3G6 monoclonal antibody is provided. In certain aspects, the antibody may comprise all or part of the heavy chain variable region and/or light chain variable region of the 8D2 or 3G6 monoclonal antibodies. In a further aspect, the antibody may comprise an amino acid sequence that corresponds to a first, second, and/or third complementarity determining region (CDR) from the light variable and/or heavy variable chain of the 8D2 or 3G6 monoclonal antibodies of the present embodiments.

[0006] In certain aspects, the isolated antibody comprises CDR sequences at least 80%, 90%, or 95% identical to the CDR regions of the 8D2 or 3G6 heavy and light chain amino acid sequences. In further aspects, an antibody comprises CDR regions identical to the 8D2 or 3G6 CDR regions, except for one or two amino acid substitutions, deletions, or insertions at one or more of the CDRs. For example, the antibody can comprise CDRs wherein the CDR sequences comprise 1 or 2 amino acid substitutions in the V_H CDR1, V_H CDR2, V_H CDR3, V_L CDR1, V_L CDR2 and/or V_L CDR3 relative to the CDRs of a 8D2 or 3G6 monoclonal antibody. Thus, in some specific aspects, an antibody of the embodiments comprises (a) a first V_H CDR at least 80% identical to V_H CDR1 of 8D2 (SEQ ID NO: 3) or 3G6 (SEQ ID NO: 13); (b) a second V_H CDR at least 80% identical to V_H CDR2 of 8D2

(SEQ ID NO: 4) or 3G6 (SEQ ID NO: 14); (c) a third V_H CDR at least 80% identical to V_H CDR3 of 8D2 (SEQ ID NO: 5) or 3G6 (SEQ ID NO: 15); (d) a first V_L CDR at least 80% identical to V_L CDR1 of 8D2 (SEQ ID NO: 8) or 3G6 (SEQ ID NO: 18); (e) a second V_L CDR at least 80% identical to V_L CDR2 of 8D2 (SEQ ID NO: 9) or 3G6 (SEQ ID NO: 19);
5 and (f) a third V_L CDR at least 80% identical to V_L CDR3 of 8D2 (SEQ ID NO: 10) or 3G6 (SEQ ID NO: 20). In certain aspects, such an antibody is a humanized or de-immunized antibody comprising the foregoing CDRs on a human IgGs (*e.g.*, IgG1, IgG2, IgG4, or a genetically modified IgG) backbone.

[0007] In further aspects, the isolated antibody comprises a first V_H, a second V_H, a
10 third V_H, a first V_L, a second V_L, and a third V_L CDR sequence at least 80% identical to the corresponding CDR sequence of monoclonal antibody 8D2, which are represented by SEQ ID NOs: 3, 4, 5, 8, 9, and 10, respectively. In one aspect, the isolated antibody comprises CDR sequences that are identical to the CDR sequences of monoclonal antibody 8D2.

[0008] In another aspect, the isolated antibody comprises a V_H domain at least about
15 80% identical to the the V_H domain of 8D2 (SEQ ID NO: 1) or the humanized V_H domain of 8D2 mAB; and a V_L domain at least about 80% identical to the V_L domain of 8D2 (SEQ ID NO: 6) or the humanized V_L domain of 8D2 mAB. For example, the antibody can comprise a V_H domain at least 95% identical to the V_H domain of the humanized 8D2 mAB and a V_L domain at least 95% identical to the V_L domain of the humanized 8D2 mAB. Thus, in some
20 aspects, an antibody comprises a V_H domain identical to the V_H domain of humanized 8D2 mAB and a V_L domain identical to the V_L domain of the humanized 8D2 mAB. In a specific example, the isolated antibody can comprise V_H and V_L domains identical to those of monoclonal antibody 8D2.

[0009] In further aspects, the isolated antibody comprises a first V_H, a second V_H, a
25 third V_H, a first V_L, a second V_L, and a third V_L CDR sequence at least 80% identical to the corresponding CDR sequence of monoclonal antibody 3G6, which are represented by SEQ ID NOs: 13, 14, 15, 18, 19, and 20, respectively. In one aspect, the isolated antibody comprises CDR sequences that are identical to the CDR sequences of monoclonal antibody 3G6.

[0010] In another aspect, the isolated antibody comprises a V_H domain at least about
30 80% identical to the V_H domain of 3G6 (SEQ ID NO: 11) or the humanized 3G6 mAB; and a

V_L domain at least about 80% identical to the V_L domain of 3G6 (SEQ ID NO: 16) or the humanized 3G6 mAB. Thus, in some aspects, the antibody comprises a V_H domain at least 95% identical to the V_H domain of the humanized 3G6 mAB and a V_L domain at least 95% identical to the V_L domain of the humanized 3G6 mAB. For instance the antibody can
5 comprise a V_H domain identical to the V_H domain of the humanized 3G6 mAB and a V_L domain identical to the V_L domain of the humanized 3G6 mAB. In a specific example, the isolated antibody can comprise V_H and V_L domains identical to those of monoclonal antibody 3G6.

[0011] In some aspects, an antibody of the embodiments may be an IgG (*e.g.*, IgG1,
10 IgG2, IgG3 or IgG4), IgM, IgA, genetically modified IgG isotype, or an antigen binding fragment thereof. The antibody may be a Fab', a F(ab')₂, a F(ab')₃, a monovalent scFv, a bivalent scFv, a bispecific or a single domain antibody. The antibody may be a human, humanized, or de-immunized antibody. In a further aspect, the isolated antibody is the 8D2 or 3G6 antibody.

15 [0012] In some aspects, the antibody may be conjugated to an imaging agent, a chemotherapeutic agent, a toxin, or a radionuclide.

[0013] In one embodiment, there is provided a recombinant polypeptide comprising an antibody V_H domain comprising CDRs 1-3 of the V_H domain of 8D2 (SEQ ID NOs: 3, 4, and 5) or CDRs 1-3 of the V_H domain of 3G6 (SEQ ID NOs: 13, 14, and 15). In another
20 embodiment, there is provided a recombinant polypeptide comprising an antibody V_L domain comprising CDRs 1-3 of the V_L domain of 8D2 (SEQ ID NOs: 8, 9, and 10) or 3G6 (SEQ ID NOs: 18, 19, and 20).

[0014] In some embodiments, there is provided an isolated polynucleotide molecule comprising nucleic acid sequence encoding an antibody or a polypeptide comprising an
25 antibody V_H or V_L domain disclosed herein.

[0015] In further embodiments, a host cell is provided that produces a monoclonal antibody or recombinant polypeptide of the embodiments. In some aspects, the host cell is a mammalian cell, a yeast cell, a bacterial cell, a ciliate cell, or an insect cell. In certain aspects, the host cell is a hybridoma cell.

[0016] In still further embodiments, there is provided a method of manufacturing an antibody of the present invention comprising expressing one or more polynucleotide molecule(s) encoding a V_L or V_H chain of an antibody disclosed herein in a cell and purifying the antibody from the cell.

5 [0017] In additional embodiments, there are pharmaceutical compositions comprising an antibody or antibody fragment as discussed herein. Such a composition further comprises a pharmaceutically acceptable carrier and may or may not contain additional active ingredients.

10 [0018] In embodiments of the present invention, there is provided a method for treating a subject having a cancer comprising administering an effective amount of an antibody disclosed herein. In certain aspects, the antibody is a monoclonal antibody of the embodiments herein, such as the 8D2 or 3G6 antibody or a recombinant polypeptide comprising antibody segment derived therefrom.

15 [0019] In certain aspects, the cancer may be a breast cancer, lung cancer, head & neck cancer, prostate cancer, esophageal cancer, tracheal cancer, brain cancer, liver cancer, bladder cancer, stomach cancer, pancreatic cancer, ovarian cancer, uterine cancer, cervical cancer, testicular cancer, colon cancer, rectal cancer or skin cancer.

20 [0020] In one aspect, the antibody may be administered systemically. In additional aspects, the antibody may be administered intravenously, intradermally, intratumorally, intramuscularly, intraperitoneally, subcutaneously, or locally. The method may further comprise administering at least a second anticancer therapy to the subject. Examples of the second anticancer therapy include, but are not limited to, surgical therapy, chemotherapy, radiation therapy, cryotherapy, hormonal therapy, immunotherapy, or cytokine therapy.

25 [0021] In further aspects, the method may further comprise administering a composition of the present invention more than one time to the subject, such as, for example, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20 or more times.

30 [0022] In another embodiment, there is provided a method for detecting a cancer in a subject comprising testing for the presence of elevated LGR4 relative to a control in a sample from the subject, wherein the testing comprises contacting the sample with an antibody disclosed herein. For example, the method may be an *in vitro* or *in vivo* method.

[0023] Certain embodiments are directed to an antibody or recombinant polypeptide composition comprising an isolated and/or recombinant antibody or polypeptide that specifically binds LGR4. In certain aspects the antibody or polypeptide has a sequence that is, is at least, or is at most 80, 85, 90, 95, 96, 97, 98, 99, or 100% identical (or any range derivable therein) to all or part of any monoclonal antibody provided herein. In still further aspects the isolated and/or recombinant antibody or polypeptide has, has at least, or has at most 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100 or more contiguous amino acids from any of the sequences provided herein or a combination of such sequences.

[0024] In still further aspects, an antibody or polypeptide of the embodiments comprises one or more amino acid segments of the any of the amino acid sequences disclosed herein. For example, the antibody or polypeptide can comprise 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more amino acid segments comprising about, at least or at most 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25 to 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199 or 200 amino acids in length, including all values and ranges there between, that are at least 80, 85, 90, 95, 96, 97, 98, 99, or 100% identical to any of the amino acid sequences disclosed herein. In certain aspects the amino segment(s) are selected from one of the amino acid sequences of a LGR4-binding antibody as provided in Table 1.

[0025] In still further aspects, an antibody or polypeptide of the embodiments comprises an amino acid segment of the any of the amino acid sequences disclosed herein, wherein the segment begins at amino acid position 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25 to 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37,

38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, or 200 in any sequence provided herein and ends at amino acid position 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25 to 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, or 200 in the same provided sequence. In certain aspects the amino segment(s), or portions thereof, are selected from one of the amino acid sequences of a LGR4-binding antibody as provided in Table 1.

[0026] In yet further aspects, an antibody or polypeptide of the embodiments comprises an amino acid segment that is at least 80, 85, 90, 95, 96, 97, 98, 99, or 100% identical (or any range derivable therein) to a V, VJ, VDJ, D, DJ, J or CDR domain of a LGR4-binding antibody (as provided in Table 1). For example, a polypeptide may comprise 1, 2 or 3 amino acid segment that are at least 80, 85, 90, 95, 96, 97, 98, 99, or 100% identical (or any range derivable therein) to CDRs 1, 2, and/or 3 a LGR4-binding antibody as provided in Table 1.

[0027] Embodiments discussed in the context of methods and/or compositions of the invention may be employed with respect to any other method or composition described herein. Thus, an embodiment pertaining to one method or composition may be applied to other methods and compositions of the invention as well.

[0028] As used herein the specification, “a” or “an” may mean one or more. As used herein in the claim(s), when used in conjunction with the word “comprising”, the words “a” or “an” may mean one or more than one.

5 [0029] The use of the term “or” in the claims is used to mean “and/or” unless explicitly indicated to refer to alternatives only or the alternatives are mutually exclusive, although the disclosure supports a definition that refers to only alternatives and “and/or.” As used herein “another” may mean at least a second or more.

10 [0030] Throughout this application, the term “about” is used to indicate that a value includes the inherent variation of error for the device, the method being employed to determine the value, or the variation that exists among the study subjects.

15 [0031] Other objects, features and advantages of the present invention will become apparent from the following detailed description. It should be understood, however, that the detailed description and the specific examples, while indicating preferred embodiments of the invention, are given by way of illustration only, since various changes and modifications within the spirit and scope of the invention will become apparent to those skilled in the art from this detailed description.

BRIEF DESCRIPTION OF THE DRAWINGS

20 [0032] The following drawings form part of the present specification and are included to further demonstrate certain aspects of the present invention. The invention may be better understood by reference to one or more of these drawings in combination with the detailed description of specific embodiments presented herein.

25 [0033] **FIG. 1.** A schematic diagram illustrating the working principles of ADC (antibody-drug conjugate). Upon binding to its target antigen, the mAb-antigen complex is internalized into endosomes which is then fused with lysosomes where the mAb is degraded and the drug is released.

30 [0034] **FIGS. 2A-2B.** LGR4 is highly expressed in primary tumors and cell lines of major epithelial cancers. 2A: RNA-seq data of LGR4 from TCGA are presented by a box and whiskers plot (top and lower bar are max and min values, respectively. The box represents the 95% confidence range of the mean for each tumor type). 2B: Microarray

expression data (normalized) of LGR4 in box and whiskers plot. Numbers in each box represent the number of cases/cell lines with expression data. CRC = colorectal adenocarcinoma; LUAD = lung adenocarcinoma; LUSQ = lung squamous cell carcinoma; BrCa = breast cancer; HCC = hepatocellular carcinoma; OvCA = ovarian cancer; RCC = kidney renal clear cell carcinoma; LuCA, lung cancer; KiCA, kidney cancer. All data were downloaded from the cBioportal website (cbioportal.org).

[0035] FIGS. 3A-3F. Identification and characterization of anti-LGR4 mAbs. 3A: Eight mAbs were purified to homogeneity. HC, heavy chain; LC, light chain. 3B: Seven purified rat mAbs gave dose-dependent binding to HEK293 cells stably expressing human LGR4. 3C: mAb 8D2 only detected recombinant LGR4 in WB. The lanes are: 1, HEK293, 2, HEK293-LGR4-mouse, 3, HEK293-LGR4-human, 4, breast cancer cell line BT474, 5, HeLa cells. 3D: ICC of 8D2 bound to LGR4 on the cell surface in HEK293 cells stably expressing LGR4. 3E: Binding of 8D2 to lung cancer cell line A549 cells led to its internalization (left panel), which was competed off by conditioned media containing LGR4-ECD (mid-panel) but not by control conditioned media (right panel). 3F: 8D2 (green) and RSPO1-Fc (red) were co-internalized in A549 cells.

[0036] FIG. 4. The LGR4-mAb complex is internalized into the lysosome. 8D2 was incubated with SKOV3 cells for 4 hours, fixed, permeabilized, co-stained with anti-rat(red) and anti-LAMP1 (green) antibodies, and viewed by confocal microscopy.

[0037] FIGS. 5A-5D. LGR4 mAb ADC showed cytotoxic activity in LGR4-high cancer cell lines. 5A: diagram of the 8D2-MC-VC-PAB-MMAE (8D2-MMAE) ADC. 5B-5D: Dose response curves of 8D2-MMAE and naked 8D2 on the viability of OVCAR3 (B), IGROV1 (C), and SKOV3(D) cells after 4 days of antibody treatment. Cell viability was determined using the CelltiterGlo™ assay. Error bars are S.E.M. (n = 3).

[0038] FIG. 6. LGR4 ADC inhibits the growth of xenograft tumors *in vivo*. OVCAR3 cells were implanted subcutaneously into nude mice. The animals were dosed with vehicle or the indicated substances every 10 days when tumors reached an average size of ~100 mm³.

[0039] FIG. 7. Binding of anti-LGR4 mAbs 8D2 and 3G6 to LGR4. Serial dilutions of the purified 8D2 and 3G6 were incubated with HEK293 cells overexpressing LGR4,

washed, fixed, and incubated with Alexa-488-labeled anti-rat IgG and then washed. Binding was based on amount of fluorescence retained by the cells.

[0040] **FIG. 8.** Comparison of cytostatic effect of 3G6 and 8D2 in the presence of MMAF-conjugated anti-rat IgG. Serial dilution of 8D2 or 3G6 mixed with MMAF-conjugated anti-rat IgG (ratio = 1:2) were incubated with the ovarian cancer cell line OVCAR3 cells or HEK293 cells overexpression LGR4 for 4 days. Cell viability was determined using the cell titer glo assays.

DESCRIPTION OF ILLUSTRATIVE EMBODIMENTS

I. Antibodies of the Embodiments

[0041] In certain embodiments, an antibody or a fragment thereof that binds to at least a portion of LGR4 protein and inhibits LGR4 signaling and cancer cell proliferation are contemplated. As used herein, the term “antibody” is intended to refer broadly to any immunologic binding agent, such as IgG, IgM, IgA, IgD, IgE, and genetically modified IgG as well as polypeptides comprising antibody CDR domains that retain antigen binding activity. The antibody may be selected from the group consisting of a chimeric antibody, an affinity matured antibody, a polyclonal antibody, a monoclonal antibody, a humanized antibody, a human antibody, or an antigen-binding antibody fragment or a natural or synthetic ligand. Preferably, the anti-LGR4 antibody is a monoclonal antibody or a humanized antibody.

[0042] Thus, by known means and as described herein, polyclonal or monoclonal antibodies, antibody fragments, and binding domains and CDRs (including engineered forms of any of the foregoing) may be created that are specific to LGR4 protein, one or more of its respective epitopes, or conjugates of any of the foregoing, whether such antigens or epitopes are isolated from natural sources or are synthetic derivatives or variants of the natural compounds.

[0043] Examples of antibody fragments suitable for the present embodiments include, without limitation: (i) the Fab fragment, consisting of V_L , V_H , C_L , and C_{H1} domains; (ii) the “Fd” fragment consisting of the V_H and C_{H1} domains; (iii) the “Fv” fragment consisting of the V_L and V_H domains of a single antibody; (iv) the “dAb” fragment, which consists of a V_H domain; (v) isolated CDR regions; (vi) $F(ab')_2$ fragments, a bivalent fragment comprising

two linked Fab fragments; (vii) single chain Fv molecules ("scFv"), wherein a V_H domain and a V_L domain are linked by a peptide linker that allows the two domains to associate to form a binding domain; (viii) bi-specific single chain Fv dimers (see U.S. Pat. No. 5,091,513); and (ix) diabodies, multivalent or multispecific fragments constructed by gene fusion (US Patent App. Pub. 20050214860). Fv, scFv, or diabody molecules may be stabilized by the incorporation of disulphide bridges linking the V_H and V_L domains. Minibodies comprising a scFv joined to a CH3 domain may also be made (Hu *et al.*, 1996).

[0044] Antibody-like binding peptidomimetics are also contemplated in embodiments. Liu *et al.* (2003) describe "antibody like binding peptidomimetics" (ABiPs), which are peptides that act as pared-down antibodies and have certain advantages of longer serum half-life as well as less cumbersome synthesis methods.

[0045] Animals may be inoculated with an antigen, such as a LGR4 extracellular domain (ECD) protein, in order to produce antibodies specific for LGR4 protein. The ECD may have 17 leucine-rich repeats and a seven transmembrane (7TM) domain typical of the rhodopsin family of G protein-coupled receptors. GenBank entries include: AF346711, AF346709, AF346710 (Genomic DNA), AAK31153.1 (Translation), BC033039, AAH33039.1 (mRNA and translation) and UniProtKB - Q9BXB1 (LGR4_Human). The LGR4 extracellular domain proteins may include amino acids 25-544, 597-620, 683-703, and 778-783 of UniProtKB Accession No. Q9BXB1. Frequently an antigen is bound or conjugated to another molecule to enhance the immune response. As used herein, a conjugate is any peptide, polypeptide, protein, or non-proteinaceous substance bound to an antigen that is used to elicit an immune response in an animal. Antibodies produced in an animal in response to antigen inoculation comprise a variety of non-identical molecules (polyclonal antibodies) made from a variety of individual antibody producing B lymphocytes. A polyclonal antibody is a mixed population of antibody species, each of which may recognize a different epitope on the same antigen. Given the correct conditions for polyclonal antibody production in an animal, most of the antibodies in the animal's serum will recognize the collective epitopes on the antigenic compound to which the animal has been immunized. This specificity is further enhanced by affinity purification to select only those antibodies that recognize the antigen or epitope of interest.

[0046] A monoclonal antibody is a single species of antibody wherein every antibody molecule recognizes the same epitope because all antibody producing cells are derived from a

single B-lymphocyte cell line. The methods for generating monoclonal antibodies (mAbs) generally begin along the same lines as those for preparing polyclonal antibodies. In some embodiments, rodents such as mice and rats are used in generating monoclonal antibodies. In some embodiments, rabbit, sheep, or frog cells are used in generating monoclonal antibodies.

5 The use of rats is well known and may provide certain advantages. Mice (e.g., BALB/c mice) are routinely used and generally give a high percentage of stable fusions.

[0047] Hybridoma technology involves the fusion of a single B lymphocyte from a mouse previously immunized with a LGR4 antigen with an immortal myeloma cell (usually mouse myeloma). This technology provides a method to propagate a single antibody-

10 producing cell for an indefinite number of generations, such that unlimited quantities of structurally identical antibodies having the same antigen or epitope specificity (monoclonal antibodies) may be produced.

[0048] Plasma B cells (CD45+CD5-CD19+) may be isolated from freshly prepared rabbit peripheral blood mononuclear cells of immunized rabbits and further selected for

15 LGR4 binding cells. After enrichment of antibody producing B cells, total RNA may be isolated and cDNA synthesized. DNA sequences of antibody variable regions from both heavy chains and light chains may be amplified, constructed into a phage display Fab expression vector, and transformed into E. coli. LGR4 specific binding Fab may be selected out through multiple rounds enrichment panning and sequenced. Selected LGR4 binding hits

20 may be expressed as full length IgG in rabbit and rabbit/human chimeric forms using a mammalian expression vector system in human embryonic kidney (HEK293) cells (Invitrogen) and purified using a protein G resin with a fast protein liquid chromatography (FPLC) separation unit.

[0049] In one embodiment, the antibody is a chimeric antibody, for example, an

25 antibody comprising antigen binding sequences from a non-human donor grafted to a heterologous non-human, human, or humanized sequence (e.g., framework and/or constant domain sequences). Methods have been developed to replace light and heavy chain constant domains of the monoclonal antibody with analogous domains of human origin, leaving the variable regions of the foreign antibody intact. Alternatively, "fully human" monoclonal

30 antibodies are produced in mice transgenic for human immunoglobulin genes. Methods have also been developed to convert variable domains of monoclonal antibodies to more human form by recombinantly constructing antibody variable domains having both rodent, for

example, mouse, and human amino acid sequences. In “humanized” monoclonal antibodies, only the hypervariable CDR is derived from mouse monoclonal antibodies, and the framework and constant regions are derived from human amino acid sequences (see U.S. Pat. Nos. 5,091,513 and 6,881,557). It is thought that replacing amino acid sequences in the antibody that are characteristic of rodents with amino acid sequences found in the corresponding position of human antibodies will reduce the likelihood of adverse immune reaction during therapeutic use. A hybridoma or other cell producing an antibody may also be subject to genetic mutation or other changes, which may or may not alter the binding specificity of antibodies produced by the hybridoma.

[0050] Methods for producing polyclonal antibodies in various animal species, as well as for producing monoclonal antibodies of various types, including humanized, chimeric, and fully human, are well known in the art and highly predictable. For example, the following U.S. patents and patent applications provide enabling descriptions of such methods: U.S. Patent Application Nos. 2004/0126828 and 2002/0172677; and U.S. Pat. Nos. 3,817,837; 3,850,752; 3,939,350; 3,996,345; 4,196,265; 4,275,149; 4,277,437; 4,366,241; 4,469,797; 4,472,509; 4,606,855; 4,703,003; 4,742,159; 4,767,720; 4,816,567; 4,867,973; 4,938,948; 4,946,778; 5,021,236; 5,164,296; 5,196,066; 5,223,409; 5,403,484; 5,420,253; 5,565,332; 5,571,698; 5,627,052; 5,656,434; 5,770,376; 5,789,208; 5,821,337; 5,844,091; 5,858,657; 5,861,155; 5,871,907; 5,969,108; 6,054,297; 6,165,464; 6,365,157; 6,406,867; 6,709,659; 6,709,873; 6,753,407; 6,814,965; 6,849,259; 6,861,572; 6,875,434; and 6,891,024. All patents, patent application publications, and other publications cited herein and therein are hereby incorporated by reference in the present application.

[0051] Antibodies may be produced from any animal source, including birds and mammals. Preferably, the antibodies are ovine, murine (e.g., mouse and rat), rabbit, goat, guinea pig, camel, horse, or chicken. In addition, newer technology permits the development of and screening for human antibodies from human combinatorial antibody libraries. For example, bacteriophage antibody expression technology allows specific antibodies to be produced in the absence of animal immunization, as described in U.S. Pat. No. 6,946,546, which is incorporated herein by reference. These techniques are further described in: Marks (1992); Stemmer (1994); Gram et al. (1992); Barbas et al. (1994); and Schier et al. (1996).

[0052] It is fully expected that antibodies to LGR4 will have the ability to neutralize or counteract the effects of LGR4 regardless of the animal species, monoclonal cell line, or

other source of the antibody. Certain animal species may be less preferable for generating therapeutic antibodies because they may be more likely to cause allergic response due to activation of the complement system through the “Fc” portion of the antibody. However, whole antibodies may be enzymatically digested into “Fc” (complement binding) fragment, and into antibody fragments having the binding domain or CDR. Removal of the Fc portion reduces the likelihood that the antigen antibody fragment will elicit an undesirable immunological response, and thus, antibodies without Fc may be preferential for prophylactic or therapeutic treatments. As described above, antibodies may also be constructed so as to be chimeric or partially or fully human, so as to reduce or eliminate the adverse immunological consequences resulting from administering to an animal an antibody that has been produced in, or has sequences from, other species.

[0053] Substitutional variants typically contain the exchange of one amino acid for another at one or more sites within the protein, and may be designed to modulate one or more properties of the polypeptide, with or without the loss of other functions or properties. Substitutions may be conservative, that is, one amino acid is replaced with one of similar shape and charge. Conservative substitutions are well known in the art and include, for example, the changes of: alanine to serine; arginine to lysine; asparagine to glutamine or histidine; aspartate to glutamate; cysteine to serine; glutamine to asparagine; glutamate to aspartate; glycine to proline; histidine to asparagine or glutamine; isoleucine to leucine or valine; leucine to valine or isoleucine; lysine to arginine; methionine to leucine or isoleucine; phenylalanine to tyrosine, leucine or methionine; serine to threonine; threonine to serine; tryptophan to tyrosine; tyrosine to tryptophan or phenylalanine; and valine to isoleucine or leucine. Alternatively, substitutions may be non-conservative such that a function or activity of the polypeptide is affected. Non-conservative changes typically involve substituting a residue with one that is chemically dissimilar, such as a polar or charged amino acid for a nonpolar or uncharged amino acid, and vice versa.

[0054] Proteins may be recombinant, or synthesized in vitro. Alternatively, a non-recombinant or recombinant protein may be isolated from bacteria. It is also contemplated that a bacteria containing such a variant may be implemented in compositions and methods. Consequently, a protein need not be isolated.

[0055] It is contemplated that in compositions there is between about 0.001 mg and about 10 mg of total polypeptide, peptide, and/or protein per ml. Thus, the concentration of

protein in a composition can be about, at least about or at most about 0.001, 0.010, 0.050, 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9, 1.0, 1.5, 2.0, 2.5, 3.0, 3.5, 4.0, 4.5, 5.0, 5.5, 6.0, 6.5, 7.0, 7.5, 8.0, 8.5, 9.0, 9.5, 10.0 mg/ml or more (or any range derivable therein). Of this, about, at least about, or at most about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, or 100% may be an antibody that binds LGR4.

[0056] An antibody or preferably an immunological portion of an antibody, can be chemically conjugated to, or expressed as, a fusion protein with other proteins. For purposes of this specification and the accompanying claims, all such fused proteins are included in the definition of antibodies or an immunological portion of an antibody.

[0057] Antibody-drug conjugates have emerged as a breakthrough approach to the development of cancer therapeutics. Cancer is one of the leading causes of deaths in the world. Antibody-drug conjugates (ADCs) comprise monoclonal antibodies (mAbs) that are covalently linked to cell-killing drugs (see, e.g., FIG. 1). Thus, embodiments provide antibodies and antibody-like molecules against LGR4, polypeptides and peptides that are linked to at least one agent to form an antibody conjugate or payload. In some aspects, this approach combines the high specificity of mAbs against their antigen targets with highly potent cytotoxic drugs, resulting in “armed” mAbs that deliver the payload (drug) to tumor cells with enriched levels of the antigen (Carter *et al.*, 2008; Teicher 2014; Leal *et al.*, 2014). Targeted delivery of the drug also minimizes its exposure in normal tissues, resulting in decreased toxicity and improved therapeutic index. Such a molecule or moiety may be, but is not limited to, at least one effector or reporter molecule. Effector molecules comprise molecules having a desired activity, e.g., cytotoxic activity. Non-limiting examples of effector molecules that have been attached to antibodies include toxins, therapeutic enzymes, antibiotics, radio-labeled nucleotides and the like. By contrast, a reporter molecule is defined as any moiety that may be detected using an assay. Non-limiting examples of reporter molecules that have been conjugated to antibodies include enzymes, radiolabels, haptens, fluorescent labels, phosphorescent molecules, chemiluminescent molecules, chromophores, luminescent molecules, photoaffinity molecules, colored particles or ligands, such as biotin.

[0058] In light of the noted oncogenic association of LGR4 and its broad expression at high levels in many major types of epithelial tumor cells and this added identification of spontaneous internalization, LGR4 presented itself as an excellent target of ADC-based drug development for the treatment of LGR4-high tumors. Indeed, the anti-LGR4 monoclonal antibodies disclosed herein demonstrate that an auristatin-conjugated LGR4 ADC had a strong cytotoxic effect *in vitro* on multiple LGR4-high cancer cell lines. As exemplified below, novel anti-LGR4 monoclonal antibodies with improved properties can be used to generate LGR4 ADC displaying strong cytotoxic effects on multiple LGR4-expressing human cancer cells, both *in vitro* and *in vivo*.

[0059] Several methods are known in the art for the attachment or conjugation of an antibody to its conjugate moiety. Some attachment methods involve the use of a metal chelate complex employing, for example, an organic chelating agent such as diethylenetriaminepentaacetic acid anhydride (DTPA); ethylenetriaminetetraacetic acid; N-chloro-p-toluenesulfonamide; and/or tetrachloro-3-6-diphenylglycouril-3 attached to the antibody. Monoclonal antibodies may also be reacted with an enzyme in the presence of a coupling agent such as glutaraldehyde or periodate. Conjugates with fluorescein markers are prepared in the presence of these coupling agents or by reaction with an isothiocyanate.

II. Treatment of Diseases

[0060] Certain aspects of the present embodiments can be used to prevent or treat a disease or disorder associated with LGR4 signaling. Signaling of LGR4 may be reduced by any suitable drugs to prevent cancer cell proliferation. Preferably, such substances would be an anti- LGR4 antibody.

[0061] “Treatment” and “treating” refer to administration or application of a therapeutic agent to a subject or performance of a procedure or modality on a subject for the purpose of obtaining a therapeutic benefit of a disease or health-related condition. For example, a treatment may include administration of a pharmaceutically effective amount of an antibody that inhibits the LGR4 signaling.

[0062] “Subject” and “patient” refer to either a human or non-human, such as primates, mammals, and vertebrates. In particular embodiments, the subject is a human.

[0063] The term “therapeutic benefit” or “therapeutically effective” as used throughout this application refers to anything that promotes or enhances the well-being of the subject with respect to the medical treatment of this condition. This includes, but is not limited to, a reduction in the frequency or severity of the signs or symptoms of a disease. For example, treatment of cancer may involve, for example, a reduction in the size of a tumor, a reduction in the invasiveness of a tumor, reduction in the growth rate of the cancer, or prevention of metastasis. Treatment of cancer may also refer to prolonging survival of a subject with cancer.

A. Pharmaceutical Preparations

[0064] Where clinical application of a therapeutic composition containing an inhibitory antibody is undertaken, it will generally be beneficial to prepare a pharmaceutical or therapeutic composition appropriate for the intended application. In certain embodiments, pharmaceutical compositions may comprise, for example, at least about 0.1% of an active compound. In other embodiments, an active compound may comprise between about 2% to about 75% of the weight of the unit, or between about 25% to about 60%, for example, and any range derivable therein.

[0065] The therapeutic compositions of the present embodiments are advantageously administered in the form of injectable compositions either as liquid solutions or suspensions; solid forms suitable for solution in, or suspension in, liquid prior to injection may also be prepared. These preparations also may be emulsified.

[0066] The phrases “pharmaceutical or pharmacologically acceptable” refers to molecular entities and compositions that do not produce an adverse, allergic, or other untoward reaction when administered to an animal, such as a human, as appropriate. The preparation of a pharmaceutical composition comprising an antibody or additional active ingredient will be known to those of skill in the art in light of the present disclosure. Moreover, for animal (e.g., human) administration, it will be understood that preparations should meet sterility, pyrogenicity, general safety, and purity standards as required by FDA Office of Biological Standards.

[0067] As used herein, “pharmaceutically acceptable carrier” includes any and all aqueous solvents (e.g., water, alcoholic/aqueous solutions, saline solutions, parenteral vehicles, such as sodium chloride, Ringer's dextrose, etc.), non-aqueous solvents (e.g.,

propylene glycol, polyethylene glycol, vegetable oil, and injectable organic esters, such as ethylolate), dispersion media, coatings, surfactants, antioxidants, preservatives (e.g., antibacterial or antifungal agents, anti-oxidants, chelating agents, and inert gases), isotonic agents, absorption delaying agents, salts, drugs, drug stabilizers, gels, binders, excipients, disintegration agents, lubricants, sweetening agents, flavoring agents, dyes, fluid and nutrient replenishers, such like materials and combinations thereof, as would be known to one of ordinary skill in the art. The pH and exact concentration of the various components in a pharmaceutical composition are adjusted according to well-known parameters.

[0068] The term “unit dose” or “dosage” refers to physically discrete units suitable for use in a subject, each unit containing a predetermined quantity of the therapeutic composition calculated to produce the desired responses discussed above in association with its administration, i.e., the appropriate route and treatment regimen. The quantity to be administered, both according to number of treatments and unit dose, depends on the effect desired. The actual dosage amount of a composition of the present embodiments administered to a patient or subject can be determined by physical and physiological factors, such as body weight, the age, health, and sex of the subject, the type of disease being treated, the extent of disease penetration, previous or concurrent therapeutic interventions, idiopathy of the patient, the route of administration, and the potency, stability, and toxicity of the particular therapeutic substance. For example, a dose may also comprise from about 1 $\mu\text{g/kg/body weight}$ to about 1000 mg/kg/body weight (this such range includes intervening doses) or more per administration, and any range derivable therein. In non-limiting examples of a derivable range from the numbers listed herein, a range of about 5 $\mu\text{g/kg/body weight}$ to about 100 mg/kg/body weight , about 5 $\mu\text{g/kg/body weight}$ to about 500 mg/kg/body weight , etc., can be administered. The practitioner responsible for administration will, in any event, determine the concentration of active ingredient(s) in a composition and appropriate dose(s) for the individual subject.

[0069] The active compounds can be formulated for parenteral administration, e.g., formulated for injection via the intravenous, intramuscular, sub-cutaneous, or even intraperitoneal routes. Typically, such compositions can be prepared as either liquid solutions or suspensions; solid forms suitable for use to prepare solutions or suspensions upon the addition of a liquid prior to injection can also be prepared; and, the preparations can also be emulsified.

[0070] The pharmaceutical forms suitable for injectable use include sterile aqueous solutions or dispersions; formulations including sesame oil, peanut oil, or aqueous propylene glycol; and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersions. In all cases the form must be sterile and must be fluid to the extent that it may be easily injected. It also should be stable under the conditions of manufacture and storage and must be preserved against the contaminating action of microorganisms, such as bacteria and fungi.

[0071] The proteinaceous compositions may be formulated into a neutral or salt form. Pharmaceutically acceptable salts, include the acid addition salts (formed with the free amino groups of the protein) and which are formed with inorganic acids such as, for example, hydrochloric or phosphoric acids, or such organic acids as acetic, oxalic, tartaric, mandelic, and the like. Salts formed with the free carboxyl groups can also be derived from inorganic bases such as, for example, sodium, potassium, ammonium, calcium, or ferric hydroxides, and such organic bases as isopropylamine, trimethylamine, histidine, procaine and the like.

[0072] A pharmaceutical composition can include a solvent or dispersion medium containing, for example, water, ethanol, polyol (for example, glycerol, propylene glycol, and liquid polyethylene glycol, and the like), suitable mixtures thereof, and vegetable oils. The proper fluidity can be maintained, for example, by the use of a coating, such as lecithin, by the maintenance of the required particle size in the case of dispersion, and by the use of surfactants. The prevention of the action of microorganisms can be brought about by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, sorbic acid, thimerosal, and the like. In many cases, it will be preferable to include isotonic agents, for example, sugars or sodium chloride. Prolonged absorption of the injectable compositions can be brought about by the use in the compositions of agents delaying absorption, for example, aluminum monostearate and gelatin.

B. Combination Treatments

[0073] In certain embodiments, the compositions and methods of the present embodiments involve an antibody or an antibody fragment against LGR4 to inhibit its activity in cancer cell proliferation, in combination with a second or additional therapy. Such therapy can be applied in the treatment of any disease that is associated with LGR4-mediated cell proliferation. For example, the disease may be cancer.

[0074] The methods and compositions, including combination therapies, enhance the therapeutic or protective effect, and/or increase the therapeutic effect of another anti-cancer or anti-hyperproliferative therapy. Therapeutic and prophylactic methods and compositions can be provided in a combined amount effective to achieve the desired effect, such as the killing of a cancer cell and/or the inhibition of cellular hyperproliferation. This process may involve contacting the cells with both an antibody or antibody fragment and a second therapy. A tissue, tumor, or cell can be contacted with one or more compositions or pharmacological formulation(s) comprising one or more of the agents (i.e., antibody or antibody fragment or an anti-cancer agent), or by contacting the tissue, tumor, and/or cell with two or more distinct compositions or formulations, wherein one composition provides 1) an antibody or antibody fragment, 2) an anti-cancer agent, or 3) both an antibody or antibody fragment and an anti-cancer agent. Also, it is contemplated that such a combination therapy can be used in conjunction with chemotherapy, radiotherapy, surgical therapy, or immunotherapy.

[0075] The terms “contacted” and “exposed,” when applied to a cell, are used herein to describe the process by which a therapeutic construct and a chemotherapeutic or radiotherapeutic agent are delivered to a target cell or are placed in direct juxtaposition with the target cell. To achieve cell killing, for example, both agents are delivered to a cell in a combined amount effective to kill the cell or prevent it from dividing.

[0076] An inhibitory antibody may be administered before, during, after, or in various combinations relative to an anti-cancer treatment. The administrations may be in intervals ranging from concurrently to minutes to days to weeks. In embodiments where the antibody or antibody fragment is provided to a patient separately from an anti-cancer agent, one would generally ensure that a significant period of time did not expire between the time of each delivery, such that the two compounds would still be able to exert an advantageously combined effect on the patient. In such instances, it is contemplated that one may provide a patient with the antibody therapy and the anti-cancer therapy within about 12 to 24 or 72 h of each other and, more particularly, within about 6-12 h of each other. In some situations it may be desirable to extend the time period for treatment significantly where several days (2, 3, 4, 5, 6, or 7) to several weeks (1, 2, 3, 4, 5, 6, 7, or 8) lapse between respective administrations.

[0077] In certain embodiments, a course of treatment will last 1-90 days or more (this such range includes intervening days). It is contemplated that one agent may be given on any

day of day 1 to day 90 (this such range includes intervening days) or any combination thereof, and another agent is given on any day of day 1 to day 90 (this such range includes intervening days) or any combination thereof. Within a single day (24-hour period), the patient may be given one or multiple administrations of the agent(s). Moreover, after a course of treatment, it is contemplated that there is a period of time at which no anti-cancer treatment is administered. This time period may last 1-7 days, and/or 1-5 weeks, and/or 1-12 months or more (this such range includes intervening days), depending on the condition of the patient, such as their prognosis, strength, health, etc. It is expected that the treatment cycles would be repeated as necessary.

- 10 **[0078]** Various combinations may be employed. For the example below an antibody therapy is “A” and an anti-cancer therapy is “B”:

A/B/A B/A/B B/B/A A/A/B A/B/B B/A/A A/B/B/B B/A/B/B
 B/B/B/A B/B/A/B A/A/B/B A/B/A/B A/B/B/A B/B/A/A
 B/A/B/A B/A/A/B A/A/A/B B/A/A/A A/B/A/A A/A/B/A

- 15 **[0079]** Administration of any compound or therapy of the present embodiments to a patient will follow general protocols for the administration of such compounds, taking into account the toxicity, if any, of the agents. Therefore, in some embodiments there is a step of monitoring toxicity that is attributable to combination therapy.

i. Chemotherapy

- 20 **[0080]** A wide variety of chemotherapeutic agents may be used in accordance with the present embodiments. The term “chemotherapy” refers to the use of drugs to treat cancer. A “chemotherapeutic agent” is used to connote a compound or composition that is administered in the treatment of cancer. These agents or drugs are categorized by their mode of activity within a cell, for example, whether and at what stage they affect the cell cycle.
- 25 Alternatively, an agent may be characterized based on its ability to directly cross-link DNA, to intercalate into DNA, or to induce chromosomal and mitotic aberrations by affecting nucleic acid synthesis.

- 30 **[0081]** Examples of chemotherapeutic agents include alkylating agents, such as thiotepa and cyclophosphamide; alkyl sulfonates, such as busulfan, improsulfan, and pipsulfan; aziridines, such as benzodopa, carboquone, meturedopa, and uredopa; ethylenimines and methylamelamines, including altretamine, triethylenemelamine,

triethylenephosphoramidate, triethylenethiophosphoramidate, and trimethylololmelamine;
 acetogenins (especially bullatacin and bullatacinone); a camptothecin (including the synthetic
 analogue topotecan); bryostatin; callystatin; CC-1065 (including its adozelesin, carzelesin
 and bizelesin synthetic analogues); cryptophycins (particularly cryptophycin 1 and
 5 cryptophycin 8); dolastatin; duocarmycin (including the synthetic analogues, KW-2189 and
 CB1-TM1); eleutherobin; pancratistatin; a sarcodictyin; spongistatin; nitrogen mustards, such
 as chlorambucil, chlornaphazine, cholophosphamide, estramustine, ifosfamide,
 mechlorethamine, mechlorethamine oxide hydrochloride, melphalan, novembichin,
 phenesterine, prednimustine, trofosfamide, and uracil mustard; nitrosureas, such as
 10 carmustine, chlorozotocin, fotemustine, lomustine, nimustine, and ranimustine; antibiotics,
 such as the enediyne antibiotics (e.g., calicheamicin, especially calicheamicin gammall and
 calicheamicin omegaII); dynemicin, including dynemicin A; bisphosphonates, such as
 clodronate; an esperamicin; as well as neocarzinostatin chromophore and related
 chromoprotein enediyne antibiotic chromophores, aclacinomysins, actinomycin,
 15 authrarnycin, azaserine, bleomycins, cactinomycin, carabycin, carminomycin, carzinophilin,
 chromomycinis, dactinomycin, daunorubicin, detorubicin, 6-diazo-5-oxo-L-norleucine,
 doxorubicin (including morpholino-doxorubicin, cyanomorpholino-doxorubicin, 2-pyrrolino-
 doxorubicin and deoxydoxorubicin), epirubicin, esorubicin, idarubicin, marcellomycin,
 mitomycins, such as mitomycin C, mycophenolic acid, nogalarnycin, olivomycins,
 20 peplomycin, potfiromycin, puromycin, quelamycin, rodorubicin, streptonigrin, streptozocin,
 tubercidin, ubenimex, zinostatin, and zorubicin; anti-metabolites, such as methotrexate and 5-
 fluorouracil (5-FU); folic acid analogues, such as denopterin, pteropterin, and trimetrexate;
 purine analogs, such as fludarabine, 6-mercaptopurine, thiamiprine, and thioguanine;
 pyrimidine analogs, such as ancitabine, azacitidine, 6-azauridine, carmofur, cytarabine,
 25 dideoxyuridine, doxifluridine, enocitabine, and floxuridine; androgens, such as calusterone,
 dromostanolone propionate, epitiostanol, mepitiothane, and testolactone; anti-adrenals, such
 as mitotane and trilostane; folic acid replenisher, such as frolinic acid; aceglatone;
 aldophosphamide glycoside; aminolevulinic acid; eniluracil; amsacrine; bestrabucil;
 bisantrene; edatraxate; defofamine; demecolcine; diaziqune; elformithine; elliptinium
 30 acetate; an epothilone; etoglucid; gallium nitrate; hydroxyurea; lentinan; lonidainine;
 maytansinoids, such as maytansine and ansamitocins; mitoguazone; mitoxantrone;
 mopidanmol; nitraerine; pentostatin; phenamet; pirarubicin; losoxantrone; podophyllinic
 acid; 2-ethylhydrazide; procarbazine; PSK polysaccharide complex; razoxane; rhizoxin;
 sizofiran; spirogermanium; tenuazonic acid; triaziquone; 2,2',2''-trichlorotriethylamine;

trichothecenes (especially T-2 toxin, verracurin A, roridin A and anguidine); urethan; vindesine; dacarbazine; mannomustine; mitobronitol; mitolactol; pipobroman; gacytosine; arabinoside ("Ara-C"); cyclophosphamide; taxoids, e.g., paclitaxel and docetaxel; gemcitabine; 6-thioguanine; mercaptopurine; platinum coordination complexes, such as cisplatin, oxaliplatin, and carboplatin; vinblastine; platinum; etoposide (VP-16); ifosfamide; mitoxantrone; vincristine; vinorelbine; novantrone; teniposide; edatrexate; daunomycin; aminopterin; xeloda; ibandronate; irinotecan (e.g., CPT-11); topoisomerase inhibitor RFS 2000; difluoromethylornithine (DMFO); retinoids, such as retinoic acid; capecitabine; carboplatin, procarbazine, plicomycin, gemcitabine, navelbine, farnesyl-protein transferase inhibitors, transplatinum, and pharmaceutically acceptable salts, acids, or derivatives of any of the above.

ii. Radiotherapy

[0082] Other factors that cause DNA damage and have been used extensively include what are commonly known as γ -rays, X-rays, and/or the directed delivery of radioisotopes to tumor cells. Other forms of DNA damaging factors are also contemplated, such as microwaves, proton beam irradiation (U.S. Patents 5,760,395 and 4,870,287), and UV-irradiation. It is most likely that all of these factors affect a broad range of damage on DNA, on the precursors of DNA, on the replication and repair of DNA, and on the assembly and maintenance of chromosomes. Dosage ranges for X-rays range from daily doses of 50 to 200 roentgens for prolonged periods of time (3 to 4 wk), to single doses of 2000 to 6000 roentgens. Dosage ranges for radioisotopes vary widely, and depend on the half-life of the isotope, the strength and type of radiation emitted, and the uptake by the neoplastic cells.

iii. Immunotherapy

[0083] The skilled artisan will understand that additional immunotherapies may be used in combination or in conjunction with methods of the embodiments. In the context of cancer treatment, immunotherapeutics, generally, rely on the use of immune effector cells and molecules to target and destroy cancer cells. Rituximab (RITUXAN®) is such an example. The immune effector may be, for example, an antibody specific for some marker on the surface of a tumor cell. The antibody alone may serve as an effector of therapy or it may recruit other cells to actually affect cell killing. The antibody also may be conjugated to a drug or toxin (chemotherapeutic, radionuclide, ricin A chain, cholera toxin, pertussis toxin, etc.) and serve as a targeting agent. Alternatively, the effector may be a lymphocyte carrying

a surface molecule that interacts, either directly or indirectly, with a tumor cell target. Various effector cells include cytotoxic T cells and NK cells

[0084] In one aspect of immunotherapy, the tumor cell must bear some marker that is amenable to targeting, i.e., is not present on the majority of other cells. Many tumor markers exist and any of these may be suitable for targeting in the context of the present embodiments. Common tumor markers include CD20, carcinoembryonic antigen, tyrosinase (p97), gp68, TAG-72, HMFG, Sialyl Lewis Antigen, MucA, MucB, PLAP, laminin receptor, erb B, and p155. An alternative aspect of immunotherapy is to combine anticancer effects with immune stimulatory effects. Immune stimulating molecules also exist including: cytokines, such as IL-2, IL-4, IL-12, GM-CSF, gamma-IFN, chemokines, such as MIP-1, MCP-1, IL-8, and growth factors, such as FLT3 ligand.

[0085] Examples of immunotherapies currently under investigation or in use are immune adjuvants, e.g., Mycobacterium bovis, Plasmodium falciparum, dinitrochlorobenzene, and aromatic compounds (U.S. Patents 5,801,005 and 5,739,169; Hui and Hashimoto, 1998; Christodoulides et al., 1998); cytokine therapy, e.g., interferons α , β , and γ , IL-1, GM-CSF, and TNF (Bukowski et al., 1998; Davidson et al., 1998; Hellstrand et al., 1998); gene therapy, e.g., TNF, IL-1, IL-2, and p53 (Qin et al., 1998; Austin-Ward and Villaseca, 1998; U.S. Patents 5,830,880 and 5,846,945); and monoclonal antibodies, e.g., anti-CD20, anti-ganglioside GM2, and anti-p185 (Hollander, 2012; Hanibuchi et al., 1998; U.S. Patent 5,824,311). It is contemplated that one or more anti-cancer therapies may be employed with the antibody therapies described herein.

iv. Surgery

[0086] Approximately 60% of persons with cancer will undergo surgery of some type, which includes preventative, diagnostic or staging, curative, and palliative surgery. Curative surgery includes resection in which all or part of cancerous tissue is physically removed, excised, and/or destroyed and may be used in conjunction with other therapies, such as the treatment of the present embodiments, chemotherapy, radiotherapy, hormonal therapy, gene therapy, immunotherapy, and/or alternative therapies. Tumor resection refers to physical removal of at least part of a tumor. In addition to tumor resection, treatment by surgery includes laser surgery, cryosurgery, electrosurgery, and microscopically-controlled surgery (Mohs' surgery).

[0087] Upon excision of part or all of cancerous cells, tissue, or tumor, a cavity may be formed in the body. Treatment may be accomplished by perfusion, direct injection, or local application of the area with an additional anti-cancer therapy. Such treatment may be repeated, for example, every 1, 2, 3, 4, 5, 6, or 7 days, or every 1, 2, 3, 4, and 5 weeks or
5 every 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, or 12 months. These treatments may be of varying dosages as well.

v. Other Agents

[0088] It is contemplated that other agents may be used in combination with certain aspects of the present embodiments to improve the therapeutic efficacy of treatment. These
10 additional agents include agents that affect the upregulation of cell surface receptors and GAP junctions, cytostatic and differentiation agents, inhibitors of cell adhesion, agents that increase the sensitivity of the hyperproliferative cells to apoptotic inducers, or other biological agents. Increases in intercellular signaling by elevating the number of GAP junctions would increase the anti-hyperproliferative effects on the neighboring
15 hyperproliferative cell population. In other embodiments, cytostatic or differentiation agents can be used in combination with certain aspects of the present embodiments to improve the anti-hyperproliferative efficacy of the treatments. Inhibitors of cell adhesion are contemplated to improve the efficacy of the present embodiments. Examples of cell adhesion inhibitors are focal adhesion kinase (FAKs) inhibitors and Lovastatin. It is further
20 contemplated that other agents that increase the sensitivity of a hyperproliferative cell to apoptosis, such as the antibody c225, could be used in combination with certain aspects of the present embodiments to improve the treatment efficacy.

III. Kits and Diagnostics

[0089] In various aspects of the embodiments, a kit is envisioned containing
25 therapeutic agents and/or other therapeutic and delivery agents. In some embodiments, the present embodiments contemplates a kit for preparing and/or administering a therapy of the embodiments. The kit may comprise one or more sealed vials containing any of the pharmaceutical compositions of the present embodiments. The kit may include, for example, at least one LGR4 antibody as well as reagents to prepare, formulate, and/or administer the
30 components of the embodiments or perform one or more steps of the inventive methods. In some embodiments, the kit may also comprise a suitable container, which is a container that will not react with components of the kit, such as an eppendorf tube, an assay plate, a syringe,

a bottle, or a tube. The container may be made from sterilizable materials such as plastic or glass.

[0090] The kit may further include an instruction sheet that outlines the procedural steps of the methods set forth herein, and will follow substantially the same procedures as described herein or are known to those of ordinary skill in the art. The instruction information may be in a computer readable media containing machine-readable instructions that, when executed using a computer, cause the display of a real or virtual procedure of delivering a pharmaceutically effective amount of a therapeutic agent.

IV. Examples

[0091] The following examples are included to demonstrate preferred embodiments of the invention. It should be appreciated by those of skill in the art that the techniques disclosed in the examples which follow represent techniques discovered by the inventor to function well in the practice of the invention, and thus can be considered to constitute preferred modes for its practice. However, those of skill in the art should, in light of the present disclosure, appreciate that many changes can be made in the specific embodiments which are disclosed and still obtain a like or similar result without departing from the spirit and scope of the invention.

Example 1 – LGR4 is highly upregulated in many major types of solid tumors

[0092] LGR4 is highly upregulated in colorectal cancer (CRC) and non-small cell lung cancer (NSCLC). Recurrent, gain-of-expression gene fusions of RSPO2 (to EIF3E) and RSPO3 (PTPRK) occur in a fraction of colorectal and other cancers as a driving mechanism. Recently, the inventors demonstrated that in lung adenocarcinoma (LUAD) gain-of-expression of RSPO3 was driven by NRF2 activation instead of gene fusion (Gong, *et al.*, 2014). This aberrant RSPO3-LGR4 signaling promotes tumor aggressiveness. Utilizing the RNA-sequence data provided by The Cancer Genome Atlas (TCGA, cancergenome.nih.gov) for a large number cases across multiple types of solid tumors, the expression of LGR4 across the major types of solid tumors was examined and it was confirmed that the receptor is highly expressed in CRC and NSCLC (**FIG. 2A**). LGR4 was also found to be expressed at similar or even higher levels in hepatocellular carcinoma (HCC), ovarian serous adenocarcinoma (OvCA), and kidney renal cell carcinoma (RCC) (**FIG. 2A**). Importantly, expression level of LGR4 is also high across cancer cell lines of the same tumor types (**FIG. 2B**) based on

microarray data of gene expression in the CCLE (Cancer Cell Line Encyclopedia) database (Barretina, *et al.* 2012), indicating that LGR4 is expressed in cancer cells of primary tumors instead of stromal cells.

Example 2 - Anti-LGR4 monoclonal antibodies

5 **[0093]** Anti-LGR4 monoclonal antibodies were generated under contract by the company Aldevron/Genovac (Aldevron, Fargo, ND, USA, aldevron.com) using DNA-based immunization of rats with a vector containing full-length human LGR4. After a series of steps from antibody purification to characterization, sixteen (16) clones were identified that produced LGR4-binding monoclonal antibodies. To further characterize the monoclonal
10 antibodies, eight clones with the best apparent binding affinity were scaled up and purified using protein G-based affinity chromatography (**FIG. 3A**). Analysis of their binding affinity using live HEK293 cells stably expressing human LGR4 revealed that the clone 8D2 was the best with $K_d = \sim 0.1 \mu\text{g/ml}$ (0.7 nM) (**FIG. 3B**). It also binds native mouse receptor with similar affinity (human and mouse LGR4 are 93% identical in the ECD at the amino acid
15 level). With denatured proteins (Western blots), 8D2 detected recombinant LGR4 of human and mouse (lower affinity) origin at the expected size without non-specific binding (**FIG. 3C**), and it bound to native LGR4 on the cell surface expressed on HEK293 cells (**FIG. 3D**). Next, the binding of 8D2 to the lung cancer cell line A549 cells which express high levels of LGR4 endogenously was determined. Incubation of 8D2 with live A549 cells at 37 degrees
20 C led to its rapid internalization (**FIG. 3E**, left panel). Importantly, the internalized signal was completely blocked by conditioned media containing LGR4-ECD but not by control conditioned media (**FIG. 3E**). Furthermore, internalized 8D2 was totally co-localized the LGR4 ligand RSPO1 when 8D2 and RSPO1-Fc were incubated together (**FIG. 3F**), indicating that 8D2 binds to LGR4 specifically without blocking its ligand binding site. In
25 addition, a specificity cross check analysis showed that none of the 8 monoclonal antibodies bound to LGR5 or LGR6.

Example 3 – Internalization of LGR4-mAb-8D2

[0094] LGR4-mAb-8D2 complex was co-internalized into lysosome. Current antibody–drug conjugate (ADC) approaches generally require internalization of the antigen-
30 ADC complex into the lysosome where the payload cytotoxin can be released by protease-mediated digestion. Therefore to investigate the potential of LGR4-mAb-8D2 complex for

use in ADC, the fate of the mAb 8D2 following incubating with ovarian cancer cell line SKOV3 cells which express high levels of LGR4 (LGR4-high cancer cells) and it was determined that the antibody was completely co-localized with the lysosome marker LAMP1 (FIG. 4), indicating that LGR4 is a suitable target for ADCs that are designed to rely on lysosomal proteases.

Example 4 - *In vitro* activity of anti LGR4-8D2-MMAE conjugate

[0095] LGR4-mAb-8D2 conjugated to auristatin (8D2-MMAE) inhibits the growth of cancer cell lines. To directly demonstrate that LGR4-targeted ADCs can inhibit the growth of LGR4-high cancer cells, the mAb 8D2 was conjugated with the toxin auristatin through a MC-VC-PAB (maleimidocaproyl-valine-citrulline-p-aminobenzyloxycarbonyl) linker in which the valine-citrulline peptide bond is subject to cleavage by proteases in the lysosome (as described in Doronina *et al.*, 2003, and diagramed in FIG. 5A). Auristatin is a highly potent inhibitor of microtubule assembly and is the payload used in the clinically approved drug ADCETRIS® (brentuximab vedotin).

[0096] A549 cells were used to confirm the binding affinity of mAb 8D2 was conjugated with auristatin (8D2-MMAE) to LGR4. A panel of cell lines with high levels of LGR4 that are represent the major types of epithelial cancers expressing LGR4 were used to validate that 8D2-MMAE was high toxic to a series of ovarian cancer cell lines (FIGS. 5B-5D).

[0097] The IC₅₀ for 8D2-MMAE on the cell lines were: OVCAR3, ~30 ng/ml; IGROV1, ~300 ng/ml; and SKOV3, 2.4 ug/ml (FIG. 5B-D). In contrast, 8D2 antibody that was not coupled to toxin, showed no effect on the growth of these cells (FIGS. 5B-5D). These results strongly suggest that anti-LGR4 ADCs have particular therapeutic potential for the treatment of ovarian cancer.

Example 5 - *In vivo* activity of 8D2-MMAE

[0098] 8D2-MMAE inhibits the growth of OVCAR3 xenograft tumors *in vivo*. A pilot study was carried out to test if 8D2-MMAE could inhibit the growth of OVCAR3 xenograft tumor *in vivo*. Tolerability of 8D2-MMAE was first tested in non-tumor bearing mice and it was found that a single dose of the ADC was well tolerated at 10 mg/kg. OVCAR3 cells were implanted subcutaneously into the dorsal flanks of athymic nude mice

and when tumors reached an average size of ~100 mm³, 8D2-MMAE, along with vehicle (PBS) and 8D2 control (naked, unconjugated 8D2), were administered by intravenous (IV) injection through the tail vein at 2 mg/kg, once every 10 days. Tumor sizes were measured once per week. As shown in **FIG. 6**, after treatment with 8D2-MMAE, tumor growth in 2 of the 6 animals was in complete remission.

Example 6 –additional anti-LGR4 mAbs

[0099] Another LGR4 monoclonal antibody (mAb) clone was identified. Additional mAbs in the library were analyzed and it was determined that the clone 3G6 displayed higher affinity ($K_d = 0.1$ ug/ml) than mAb 8D2 ($K_d = 0.5$ ug/ml) in binding to human LGR4 as is shown in **FIG. 7**.

[00100] It was determined that 3G6 was more potent than 8D2 in inhibiting cell survival when combined with a secondary ADC. Serial dilutions of the LGR4 monoclonal antibodies 8D2 or 3G6 were mixed with cells and MMAF-conjugated anti-rat IgG (ratio = 1:2). When tested, 3G6 was more potent in inhibiting the growth of cells with high expression of LGR4. As shown in **FIG. 8**, 3G6 displayed high potency in reducing the growth of the ovarian cancer cell line OVCAR3 cells and HEK293 cells which overexpress LGR4. This suggests that while both LGR4 monoclonal antibodies have activity, 3G6 may be more effective at targeting tumor cells *in vivo*, thus increasing patient survival.

V. Sequences

[00101] The V_H and V_L sequences of 3G6 and 8D2 clone were amplified by PCR, cloned, and sequenced. Their DNA and protein sequences are shown below in Table 1.

[00102] **Table 1.** Anti-LGR4 antibody sequences.

mAb	Heavy Chain Sequence			Light Chain Sequence		
	CDR1	CDR2	CDR3	CDR1	CDR2	CDR3
	Amino acid sequence			Amino acid sequence		
	Nucleic acid sequence			Nucleic acid sequence		
8D2	GYTFSTSY (SEQ ID NO: 3)	IYVGAGGP (SEQ ID NO: 4)	ARVRGYAM DA (SEQ ID NO:	QDISTS (SEQ ID NO: 8)	YAS (SEQ ID NO: 9)	QQSYSLPV T (SEQ ID NO:

			5)			10)
	QVQLQQSGAELVKPGASVNLSCKTSGY TFSTSYMSWFKQIPGQTIEWVGLIYVG AGGPTYNQKFKGKASLTVDTSSTAYM DLSSLTAEDSAVYFCARVRGYAMDAWG QGTSVTVSS (SEQ ID NO: 1)			DIVLTQSPATLSVTPGESVSLSCRASQ DISTSIHWYQQKSNESPRLLIKYASQS ISGIPSRFSGGGSGTDFTLTINRVESE DFSVYYCQQSYSLPVTFGSGTKLELKR (SEQ ID NO: 6)		
	CAGGTCCAGCTGCAGCAATCTGGGGCT GAGTTGGTAAAGCCTGGTGCTTCAGTG AATTTGTCCTGCAAGACTTCTGGTTAC ACCTTCAGCACTAGCTACATGAGTTGG TTCAAGCAGATTCCTGGACAGACTATT GAGTGGGTTGGACTGATTTATGTTGGA GCTGGTGGTCCTACCTATAATCAGAAAG TTCAAGGGCAAGGCCTCACTTACTGTC GACACATCTTCCAGCACAGCGTACATG GATCTCAGCAGCCTGACAGCTGAGGAC TCTGCAGTCTATTTTTGTGCAAGGGTT CGGGGCTATGCTATGGATGCCTGGGGT CAAGGAACCTCAGTCACCGTCTCCTCA (SEQ ID NO: 2)			GACATTGTGCTCACCCAGTCTCCAGCC ACCCTGTCTGTGACTCCAGGAGAGAGT GTGAGTCTCTCCTGCAGGGCCAGTCAG GATATTAGCACTAGCATTTCATTGGTAT CAGCAAAAATCAAATGAGTCTCCAAGG CTTCTCATCAAGTATGCTTCCCAGTCC ATCTCTGGAATCCCCCTCCAGGTTCACT GGAGGTGGATCAGGGACAGATTTCACT CTCACATCAACAGAGTCGAATCTGAA GATTTTTTCAGTTTATTACTGTCAACAG AGTTACAGCTTGCCCGTCACGTTCCGA TCTGGGACCAAGCTGGAAGTGAACGG (SEQ ID NO: 7)		
3G6	GYTFTSNF (SEQ ID NO: 13)	IYPGDGDT (SEQ ID NO: 14)	SRSNSGYN WFAY (SEQ ID NO: 15)	QNINKN (SEQ ID NO: 18)	YTN (SEQ ID NO: 19)	YQYRTGWT (SEQ ID NO: 20)
	QVQLQQSGAELVKPGSSVKISCKASGY TFTSNFMHWIKQQPGNGLEWIGWIYPG DGD TDYNQKFIGKATLTADKSSSTAYM QLSSLTSEDSAVYFCRSNSNGYNWFAY WGQGTTLVTVSS (SEQ ID NO: 11)			DIVLTQSPPVLSASVGDRTLSCKASQ NINKNLDWYQQKHGEAPKLLIYYTNNL QTGIPSRFSGSGSGTDYTLTIRSLQPE DVATYYCYQYRTGWTFGGGTKLELKR (SEQ ID NO: 16)		
	CAGGTTTCAGCTGCAGCAGTCTGGGGCT GAACTGGTGAAGCCTGGGTCCTCAGTG AAAATTTCTGCAAGGCTTCTGGCTAC ACCTTCACCAAGTAACCTTATGCACTGG ATAAAACAGCAGCCTGGAAATGGCCTT GAGTGGATTGGGTGGATTTATCCTGGA GATGGTGATACAGATTACAATCAAAAG TTCATTGGGAAGGCAACACTCACTGCA GACAAATCCTCCAGCACAGCCTATATG CAGCTCAGCAGCCTGACATCTGAGGAC TCTGCAGTCTATTTCTGTTCAAGATCG AATTCGGGATACAATTGGTTTGCTTAC TGGGGCCAAGGCACTCTGGTCACCGTC TCCTCA (SEQ ID NO: 12)			GACATTGTGCTCACCCAGTCTCCTCCA GTCCTGTCTGCATCTGTGGGAGACAGA GTCACCTCTCAGCTGCAAAGCAAGTCAG AATATTAATAAGAACTTAGACTGGTAT CAACAAAAGCATGGAGAAGCTCCAAAA CTCCTGATATATTATACAAACAATTTG CAAACGGGTATCCCATCAAGGTTCACT GGCAGTGGATCTGGTACAGATTACACA CTCACCATCAGAAGCCTGCAGCCTGAG GATGTTGCCACATATTACTGCTATCAG TATAGAACTGGGTGGACCTTCGGTGGA GGCACCAAGCTGGAGTTGAAACGG (SEQ ID NO: 17)		

* * *

[00103] All of the methods disclosed and claimed herein can be made and executed without undue experimentation in light of the present disclosure. While the compositions and methods of this invention have been described in terms of preferred embodiments, it will be apparent to those of skill in the art that variations may be applied to the methods and in the steps or in the sequence of steps of the method described herein without departing from the concept, spirit and scope of the invention. More specifically, it will be apparent that certain agents which are both chemically and physiologically related may be substituted for the agents described herein while the same or similar results would be achieved. All such similar substitutes and modifications apparent to those skilled in the art are deemed to be within the spirit, scope and concept of the invention as defined by the appended claims.

REFERENCES

The following references, to the extent that they provide exemplary procedural or other details supplementary to those set forth herein, are specifically incorporated herein by reference.

U.S. Patent Application Nos. 2004/0126828 and 2002/0172677

U.S. Patent Nos. 3,817,837; 3,850,752; 3,939,350; 3,996,345; 4,196,265; 4,275,149; 4,277,437; 4,366,241; 4,469,797; 4,472,509; 4,606,855; 4,703,003; 4,742,159; 4,767,720; 4,816,567; 4,867,973; 4,938,948; 4,946,778; 5,021,236; 5,164,296; 5,196,066; 5,223,409; 5,403,484; 5,420,253; 5,565,332; 5,571,698; 5,627,052; 5,656,434; 5,770,376; 5,789,208; 5,821,337; 5,844,091; 5,858,657; 5,861,155; 5,871,907; 5,969,108; 6,054,297; 6,165,464; 6,365,157; 6,406,867; 6,709,659; 6,709,873; 6,753,407; 6,814,965; 6,849,259; 6,861,572; 6,875,434; and 6,891,024.

Lefranc *et al.*, *Methods in molecular biology*, 907, 3, 2012.

Haidar *et al.*, *Proteins* 80, 896, Mar., 2012.

Retter *et al.*, *Nucleic Acids Res* 33, D671, Jan 1, 2005.

Singer *et al.*, *J Immunol* 150, 2844, 1993.

Yu *et al.*, *PLoS ONE* 5, e9072, 2010.

1. Carter, P. J., and Senter, P. D. (2008) Antibody-drug conjugates for cancer therapy, *Cancer J* 14, 154-169.
2. Teicher, B. A. (2014) Antibody drug conjugates, *Current opinion in oncology* 26, 476-483.
3. Leal, M., Sapra, P., Hurvitz, S. A., Senter, P., Wahl, A., Schutten, M., Shah, D. K., Haddish-Berhane, N., and Kabbarah, O. (2014) Antibody-drug conjugates: an emerging modality for the treatment of cancer, *Ann N Y Acad Sci* 1321, 41-54.
4. Senter, P. D. (2009) Potent antibody drug conjugates for cancer therapy, *Current opinion in chemical biology* 13, 235-244.
5. Teicher, B. A. (2009) Antibody-drug conjugate targets, *Current cancer drug targets* 9, 982-1004.
6. Bander, N. H. (2013) Antibody-drug conjugate target selection: critical factors, *Methods Mol Biol* 1045, 29-40.

7. Hsu, S. Y., Kudo, M., Chen, T., Nakabayashi, K., Bhalla, A., van der Spek, P. J., van Duin, M., and Hsueh, A. J. (2000) The three subfamilies of leucine-rich repeat-containing G protein-coupled receptors (LGR): identification of LGR6 and LGR7 and the signaling mechanism for LGR7, *Mol Endocrinol* 14, 1257-1271.
8. Carmon, K. S., Gong, X., Lin, Q., Thomas, A., and Liu, Q. (2011) R-spondins function as ligands of the orphan receptors LGR4 and LGR5 to regulate Wnt/beta-catenin signaling, *Proc Natl Acad Sci U S A* 108, 11452-11457.
9. de Lau, W., Barker, N., Low, T. Y., Koo, B. K., Li, V. S., Teunissen, H., Kujala, P., Haegebarth, A., Peters, P. J., van de Wetering, M., Stange, D. E., van Es, J. E., Guardavaccaro, D., Schasfoort, R. B., Mohri, Y., Nishimori, K., Mohammed, S., Heck, A. J., and Clevers, H. (2011) Lgr5 homologues associate with Wnt receptors and mediate R-spondin signalling, *Nature* 476, 293-297.
10. Glinka, A., Dolde, C., Kirsch, N., Huang, Y. L., Kazanskaya, O., Ingelfinger, D., Boutros, M., Cruciat, C. M., and Niehrs, C. (2011) LGR4 and LGR5 are R-spondin receptors mediating Wnt/beta-catenin and Wnt/PCP signalling, *EMBO Rep* 12, 1055-1061.
11. Hao, H. X., Xie, Y., Zhang, Y., Charlat, O., Oster, E., Avello, M., Lei, H., Mickanin, C., Liu, D., Ruffner, H., Mao, X., Ma, Q., Zamponi, R., Bouwmeester, T., Finan, P. M., Kirschner, M. W., Porter, J. A., Serluca, F. C., and Cong, F. (2012) ZNRF3 promotes Wnt receptor turnover in an R-spondin-sensitive manner, *Nature* 485, 195-200.
12. Carmon, K. S., Gong, X., Yi, J., Thomas, A., and Liu, Q. (2014) RSPO-LGR4 functions via IQGAP1 to potentiate Wnt signaling, *Proc Natl Acad Sci U S A* 111, E1221-1229.
13. de Lau, W., Peng, W. C., Gros, P., and Clevers, H. (2014) The R-spondin/Lgr5/Rnf43 module: regulator of Wnt signal strength, *Genes Dev* 28, 305-316.
14. Gao, Y., Kitagawa, K., Hiramatsu, Y., Kikuchi, H., Isobe, T., Shimada, M., Uchida, C., Hattori, T., Oda, T., Nakayama, K., Nakayama, K. I., Tanaka, T., Konno, H., and Kitagawa, M. (2006) Up-regulation of GPR48 induced by down-regulation of p27Kip1 enhances carcinoma cell invasiveness and metastasis, *Cancer Res* 66, 11623-11631.
15. Gugger, M., White, R., Song, S., Waser, B., Cescato, R., Riviere, P., and Reubi, J. C. (2008) GPR87 is an overexpressed G-protein coupled receptor in squamous cell carcinoma of the lung, *Dis Markers* 24, 41-50.

16. Seshagiri, S., Stawiski, E. W., Durinck, S., Modrusan, Z., Storm, E. E., Conboy, C. B., Chaudhuri, S., Guan, Y., Janakiraman, V., Jaiswal, B. S., Guillory, J., Ha, C., Dijkgraaf, G. J., Stinson, J., Gnad, F., Huntley, M. A., Degenhardt, J. D., Haverty, P. M., Bourgon, R., Wang, W., Koeppen, H., Gentleman, R., Starr, T. K., Zhang, Z., Largaespada, D. A., Wu, T. D., and de Sauvage, F. J. (2012) Recurrent R-spondin fusions in colon cancer, *Nature* 488, 660-664.
17. Shinmura, K., Kahyo, T., Kato, H., Igarashi, H., Matsuura, S., Nakamura, S., Kurachi, K., Nakamura, T., Ogawa, H., Funai, K., Tanahashi, M., Niwa, H., and Sugimura, H. (2014) RSPO fusion transcripts in colorectal cancer in Japanese population, *Mol Biol Rep.*
18. Gong, X., Yi, J., Carmon, K. S., Crumbley, C. A., Xiong, W., Thomas, A., Fan, X., Guo, S., An, Z., Chang, J. T., and Liu, Q. J. (2014) Aberrant RSPO3-LGR4 signaling in Keap1-deficient lung adenocarcinomas promotes tumor aggressiveness, *Oncogene*.
19. Yi, J., Xiong, W., Gong, X., Bellister, S., Ellis, L. M., and Liu, Q. (2013) Analysis of LGR4 Receptor Distribution in Human and Mouse Tissues, *PLoS One* 8, e78144.
20. Barretina, J., Caponigro, G., Stransky, N., Venkatesan, K., Margolin, A. A., Kim, S., Wilson, C. J., Lehar, J., Kryukov, G. V., Sonkin, D., Reddy, A., Liu, M., Murray, L., Berger, M. F., Monahan, J. E., Morais, P., Meltzer, J., Korejwa, A., Jane-Valbuena, J., Mapa, F. A., Thibault, J., Bric-Furlong, E., Raman, P., Shipway, A., Engels, I. H., Cheng, J., Yu, G. K., Yu, J., Aspesi, P., Jr., de Silva, M., Jagtap, K., Jones, M. D., Wang, L., Hatton, C., Palessandolo, E., Gupta, S., Mahan, S., Sougnez, C., Onofrio, R. C., Liefeld, T., MacConaill, L., Winckler, W., Reich, M., Li, N., Mesirov, J. P., Gabriel, S. B., Getz, G., Ardlie, K., Chan, V., Myer, V. E., Weber, B. L., Porter, J., Warmuth, M., Finan, P., Harris, J. L., Meyerson, M., Golub, T. R., Morrissey, M. P., Sellers, W. R., Schlegel, R., and Garraway, L. A. (2012) The Cancer Cell Line Encyclopedia enables predictive modelling of anticancer drug sensitivity, *Nature* 483, 603-607.
21. Austin, C. D., De Maziere, A. M., Pisacane, P. I., van Dijk, S. M., Eigenbrot, C., Sliwkowski, M. X., Klumperman, J., and Scheller, R. H. (2004) Endocytosis and sorting of ErbB2 and the site of action of cancer therapeutics trastuzumab and geldanamycin, *Mol Biol Cell* 15, 5268-5282.
22. Erickson, H. K., Park, P. U., Widdison, W. C., Kovtun, Y. V., Garrett, L. M., Hoffman, K., Lutz, R. J., Goldmacher, V. S., and Blattler, W. A. (2006) Antibody-

- maytansinoid conjugates are activated in targeted cancer cells by lysosomal degradation and linker-dependent intracellular processing, *Cancer Res* 66, 4426-4433.
23. Doronina, S. O., Toki, B. E., Torgov, M. Y., Mendelsohn, B. A., Cervený, C. G., Chace, D. F., DeBlanc, R. L., Gearing, R. P., Bovee, T. D., Siegall, C. B., Francisco, J. A., Wahl, A. F., Meyer, D. L., and Senter, P. D. (2003) Development of potent monoclonal antibody auristatin conjugates for cancer therapy, *Nat Biotechnol* 21, 778-784.

WHAT IS CLAIMED IS:

1. An isolated monoclonal antibody, wherein the antibody specifically binds to LGR4 and wherein the antibody comprises:

(i)

- (a) a first V_H CDR is identical to SEQ ID NO: 3;
- (b) a second V_H CDR is identical to SEQ ID NO: 4;
- (c) a third V_H CDR is identical to SEQ ID NO: 5;
- (d) a first V_L CDR is identical to SEQ ID NO: 8;
- (e) a second V_L CDR is identical to SEQ ID NO: 9; and
- (f) a third V_L CDR is identical to SEQ ID NO: 10; or

(ii)

- (a) a first V_H CDR is identical to SEQ ID NO: 13;
- (b) a second V_H CDR is identical to SEQ ID NO: 14;
- (c) a third V_H CDR is identical to SEQ ID NO: 15;
- (d) a first V_L CDR is identical to SEQ ID NO: 18;
- (e) a second V_L CDR is identical to SEQ ID NO: 19; and
- (f) a third V_L CDR is identical to SEQ ID NO: 20.

2. The antibody of claim 1, comprising:

- (a) a first V_H CDR is identical to SEQ ID NO: 3;
- (b) a second V_H CDR is identical to SEQ ID NO: 4;
- (c) a third V_H CDR is identical to SEQ ID NO: 5;
- (d) a first V_L CDR is identical to SEQ ID NO: 8;
- (e) a second V_L CDR is identical to SEQ ID NO: 9; and
- (f) a third V_L CDR is identical to SEQ ID NO: 10.

3. The antibody of claim 1, comprising:

- (a) a first V_H CDR is identical to SEQ ID NO: 13;
- (b) a second V_H CDR is identical to SEQ ID NO: 14;
- (c) a third V_H CDR is identical to SEQ ID NO: 15;

- (d) a first V_L CDR is identical to SEQ ID NO: 18;
 - (e) a second V_L CDR is identical to SEQ ID NO: 19; and
 - (f) a third V_L CDR is identical to SEQ ID NO: 20.
4. The antibody of claim 1, wherein the antibody comprises:
- (i) a V_H domain at least about 80% identical to the V_H domain of 8D2 (SEQ ID NO: 1) or the humanized V_H domain of 8D2 mAB; and a V_L domain at least about 80% identical to the V_L domain of 8D2 (SEQ ID NO: 6) or the humanized V_L domain of 8D2 mAB; or
 - (ii) a V_H domain at least about 80% identical to the V_H domain of 3G6 (SEQ ID NO: 11) or the humanized V_H domain of 3G6 mAB; and a V_L domain at least about 80% identical to the V_L domain of 3G6 (SEQ ID NO: 16) or the humanized V_L domain of 3G6 mAB.
5. The antibody of claim 4, wherein the antibody comprises a V_H domain identical to the V_H domain of 8D2 (SEQ ID NO: 1) and a V_L domain identical to the V_L domain of 8D2 (SEQ ID NO: 6).
6. The antibody of claim 4, wherein the antibody comprises a V_H domain at least 95% identical to the V_H domain of the humanized 8D2 mAB and a V_L domain at least 95% identical to the V_L domain of the humanized 8D2 mAB.
7. The antibody of claim 6, wherein the antibody comprises a V_H domain identical to the V_H domain of the humanized 8D2 mAB and a V_L domain identical to the V_L domain of the humanized 8D2 mAB.
8. The antibody of claim 4, wherein the antibody comprises a V_H domain identical to the V_H domain of 3G6 (SEQ ID NO: 11) and a V_L domain identical to the V_L domain of 3G6 (SEQ ID NO: 16).
9. The antibody of claim 4, wherein the antibody comprises a V_H domain at least 95% identical to the V_H domain of the humanized 3G6 mAB and a V_L domain at least 95% identical to the V_L domain of the humanized 3G6 mAB.
10. The antibody of claim 9, wherein the antibody comprises a V_H domain identical to the V_H domain of the humanized 3G6 mAB and a V_L domain identical to the V_L domain of the humanized 3G6 mAB.

11. The antibody of any one of claims 1-10, wherein the antibody is recombinant.
12. The antibody of claim 1, wherein the antibody is an IgG, IgM, IgA or an antigen binding fragment thereof.
13. The antibody of any one of claims 1-10, wherein the antibody is a Fab', a F(ab')₂, a F(ab')₃, a monovalent scFv, a bivalent scFv, or a single domain antibody.
14. The antibody of any one of claims 1-3, wherein the antibody is a human, humanized antibody or de-immunized antibody.
15. The antibody of any one of claims 1-10, wherein the antibody is conjugated to an imaging agent, a chemotherapeutic agent, a toxin or a radionuclide.
16. The antibody of claim 15, wherein the antibody is conjugated to a toxin.
17. The antibody of claim 16, wherein the toxin is auristatin.
18. The antibody of claim 16, wherein the toxin is monomethyl auristatin E (MMAE).
19. A composition comprising an antibody of any one of claims 1-10 in a pharmaceutically acceptable carrier.
20. An isolated polynucleotide molecule comprising a nucleic acid sequence encoding an antibody of any one of claims 1-10.
21. A recombinant polypeptide comprising an antibody V_H domain comprising CDRs 1-3 of the V_H domain of 8D2 (SEQ ID NOs: 3, 4, and 5) or CDRs 1-3 of the V_H domain of 3G6 (SEQ ID NOs: 13, 14, and 15).
22. A recombinant polypeptide comprising an antibody V_L domain comprising CDRs 1-3 of the V_L domain of 8D2 (SEQ ID NOs: 8, 9, and 10) or 3G6 (SEQ ID NOs: 18, 19, and 20).
23. An isolated polynucleotide molecule comprising a nucleic acid sequence encoding a polypeptide of claim 21 or 22.
24. A host cell comprising one or more polynucleotide molecule(s) encoding an antibody of any one of claims 1-10 or a recombinant polypeptide of claim 21 or 22.

25. The host cell of claim 24, wherein the host cell is a mammalian cell, a yeast cell, a bacterial cell, a ciliate cell or an insect cell.
26. A method of manufacturing an antibody comprising:
- (a) expressing one or more polynucleotide molecule(s) encoding a V_L and V_H chain of an antibody of any one of claims 1-10 in a cell; and
 - (b) purifying the antibody from the cell.
27. A method for treating a subject having a cancer comprising administering an effective amount of an antibody of any one of claims 1-10 to the subject.
28. The method of claim 27, wherein the cancer is a breast cancer, lung cancer, head & neck cancer, prostate cancer, esophageal cancer, tracheal cancer, skin cancer brain cancer, liver cancer, bladder cancer, stomach cancer, pancreatic cancer, ovarian cancer, uterine cancer, cervical cancer, testicular cancer, colon cancer, rectal cancer or skin cancer.
29. The method of claim 27, wherein the cancer is an epithelial cancer.
30. The method of claim 27, wherein the cancer is a colorectal adenocarcinoma, lung adenocarcinoma, lung squamous cell carcinoma, breast cancer, hepatocellular carcinoma, ovarian cancer, kidney renal clear cell carcinoma, lung cancer or kidney cancer.
31. The method of claim 27, wherein the antibody is in a pharmaceutically acceptable composition.
32. The method of claim 27, wherein the antibody is administered systemically.
33. The method of claim 27, wherein the antibody is administered intravenously, intradermally, intratumorally, intramuscularly, intraperitoneally, subcutaneously, or locally.
34. The method of claim 27, further comprising administering at least a second anticancer therapy to the subject.
35. The method of claim 34, wherein the second anticancer therapy is a surgical therapy, chemotherapy, radiation therapy, cryotherapy, hormonal therapy, immunotherapy or cytokine therapy.

36. A method for detecting a cancer in a subject comprising testing for the presence of elevated LGR4 relative to a control in a sample from the subject, wherein the testing comprises contacting the sample with an antibody of any one of claims 4-10 or 15.

37. The method of claim 36, further defined as an *in vitro* method.

1/8




FIG. 1

2/8




FIG. 2A




FIG. 2B

3/8




FIG. 3A




FIG. 3B




FIG. 3C




FIG. 3D

4/8




FIG. 3E




FIG. 3F




FIG. 4

5/8




FIG. 5A

6/8




FIG. 5B




FIG. 5C

7/8




FIG. 5D




FIG. 6




FIG. 7




FIG. 8

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US16/65788

A. CLASSIFICATION OF SUBJECT MATTER

IPC - A61K 39/395; G01N 33/15, 33/50; C07K 14/705, 16/28; C12N 5/10, 15/09 (2017. 01)

CPC - A61K 39/395; G01N 33/15, 33/50; C07K 14/705, 16/28; C12N 5/10, 15/09

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

See Search History document

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

See Search History document

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

See Search History document

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 2006/103116 A1 (BIOTHERAPIX MOLECULAR MEDICINES S.L.U.) 5 October 2006. SEQ ID NO: 4; paragraph [0032]; claim 5; figure 9	1, 2, 4-7, 11-35
A	WO 2015/032906 A2 (BIONOVION HOLDING B.V.) 12 March 2015. SEQ ID NO: 84; table 1; claim 5	1, 2, 4-7, 11-35
A	US 2013/0231465 A1 (ZEIDLER, R et al.) 5 September 2013. SEQ ID NO: 3; page 10	1, 2, 4-7, 11-35
A	WO 2013/067057 A1 (BIONOMICS, INC.) 10 May 2013; entire document	1, 2, 4-7, 11-35

☐ Further documents are listed in the continuation of Box C.☐ See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

12 April 2017 (12.04.2017)

Date of mailing of the international search report

28 APR 2017

Name and mailing address of the ISA/

Mail Stop PCT, Attn: ISA/US, Commissioner for Patents
P.O. Box 1450, Alexandria, Virginia 22313-1450

Facsimile No. 571-273-8300

Authorized officer

Shane Thomas

PCT Helpdesk: 571-272-4300

PCT OSP: 571-272-7774

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US16/65788

Box No. 1 Nucleotide and/or amino acid sequence(s) (Continuation of item 1.c of the first sheet)

1. With regard to any nucleotide and/or amino acid sequence disclosed in the international application, the international search was carried out on the basis of a sequence listing:

a. ☒ forming part of the international application as filed:

☒ in the form of an Annex C/ST.25 text file.

☐ on paper or in the form of an image file.

b. ☐ furnished together with the international application under PCT Rule 13ter.1(a) for the purposes of international search only in the form of an Annex C/ST.25 text file.

c. ☐ furnished subsequent to the international filing date for the purposes of international search only:

☐ in the form of an Annex C/ST.25 text file (Rule 13ter.1(a)).

☐ on paper or in the form of an image file (Rule 13ter.1(b) and Administrative Instructions, Section 713).

2. ☐ In addition, in the case that more than one version or copy of a sequence listing has been filed or furnished, the required statements that the information in the subsequent or additional copies is identical to that forming part of the application as filed or does not go beyond the application as filed, as appropriate, were furnished.

3. Additional comments:

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US16/65788

Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☒ Claims Nos.: 36 and 37
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

-Please See Supplemental Page-

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying additional fees, this Authority did not invite payment of additional fees.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:
Groups I+, Claims 1, 2, 4-7, 11-35 and SEQ ID NO: 1 (HVR); SEQ ID NO: 3 (HCDR1); SEQ ID NO: 4 (HCDR2); SEQ ID NO: 5 (HCDR3); SEQ ID NO: 6 (LVR); SEQ ID NO: 8 (LCDR1); SEQ ID NO: 9 (LCDR2); SEQ ID NO: 10 (LCDR3)

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.
- ☐ The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.
- ☐ No protest accompanied the payment of additional search fees.