

(54) Title
Small molecule inhibitors of Bcl-2-associated death promoter (BAD) phosphorylation

(51) International Patent Classification(s)
C07D 211/06 (2006.01) **A61P 35/00** (2006.01)
A61K 31/445 (2006.01) **C07D 241/04** (2006.01)
A61K 31/496 (2006.01)

(21) Application No: **2018256285** (22) Date of Filing: **2018.04.18**

(87) WIPO No: **WO18/194520**

(30) Priority Data

(31) Number (32) Date (33) Country
1706162.3 **2017.04.19** **GB**

(43) Publication Date: **2018.10.25**

(44) Accepted Journal Date: **2022.08.04**

(71) Applicant(s)
National University of Singapore;University of Mysore;Bangalore University

(72) Inventor(s)
Lobie, Peter Edward;Pandey, Vijay Kumar;Kanchugarakoppal Subbegowda, Rangappa;Salundi, Bassappa;Chakrabhavi Dhananjaya, Mohan;Rangappa, Shobith;Venkatachalaiah, Srinivasa

(74) Agent / Attorney
WRAYS PTY LTD, L7 863 Hay St, Perth, WA, 6000, AU

(56) Related Art
US 20130096133 A1
WO 0107050 A1
KARUBE K. et al., "Comprehensive gene expression profiles of NK cell neoplasms identify vorinostat as an effective drug candidate", Cancer Lett., vol. 333, no. 1, (2013-01-21), pages 47 - 55, (2018-06-13)
STN CAS RN 1088182-85-7 Registry File, Entered STN: 22 Dec 2008
XINWEI HE ET AL, "FeCl 3 -Mediated One-Pot Domino Reactions for the Synthesis of 9-Aryl/9-Arylethynyl-2,3,4,9-tetrahydro-1 H -xanthen-1-ones", JOURNAL OF ORGANIC CHEMISTRY, 2016, vol. 81, no. 5, pages 2062 - 2069
ALAN R. KATRITZKY ET AL, "Amino(hetero)arylmethylation of Phenols with N -[[alpha]-Amino(hetero)arylmethyl]benzotriazoles", JOURNAL OF ORGANIC CHEMISTRY, US, (1999-08-01), vol. 64, no. 16, pages 6071 - 6075
S. TABASSUM ET AL, "Sonochemistry - an innovative opportunity towards a one-pot three-component synthesis of novel pyridylpiperazine derivatives ...", NEW JOURNAL OF CHEMISTRY, 2017, vol. 41, no. 9, pages 3515 - 3523
HOSSEINZADEH R. et al., "A green protocol for the one-pot multicomponent Petasis boronic Mannich reaction using ball milling", J. Iran. Chem. Soc., vol. 14, no. 2, (2016-10-04), pages 347 - 355, (2018-06-13)
CHRISTOV K. et al., "Short-term Modulation of Cell Proliferation and Apoptosis and Preventive/Therapeutic Efficacy of Various Agents in a Mammary Cancer Model", Clin. Cancer Res., vol. 13, no. 18, 2007, pages 5488 - 5496,

LIN Ho. J.-C. et al., "Induction of apoptosis and cell -cycle arrest in human colon cancer cells by meclizine", *Food Chem. Toxicol.*, vol. 45, no. 6, (2006-12-09), pages 935 - 944, (2018-06-13)

RICHARD D. J. et al., "Hydroxyquinoline-derived compounds and analoguing of selective Mcl-1 inhibitors using a functional biomarker", *Bioorg Med Chem.*, vol. 21, no. 21, (2013-08-15), pages 6642 - 6649, (2018-06-15)

THANKRAN A. K. et al., "Synthesis and Pharmacological Evaluation of 1- Benz-Hydryl Piperazine Derivatives", *Int. J. Pharm. Sci. Res.*, vol. 3, no. 1, (2012-01-01), pages 213 - 217, (2018-06-13)

(19) World Intellectual Property
Organization

International Bureau

(43) International Publication Date
25 October 2018 (25.10.2018)(10) International Publication Number
WO 2018/194520 A9

(51) International Patent Classification:

C07D 211/06 (2006.01) A61K 31/496 (2006.01)
C07D 241/04 (2006.01) A61P 35/00 (2006.01)
A61K 31/445 (2006.01)

(21) International Application Number:

PCT/SG2018/050194

(22) International Filing Date:

18 April 2018 (18.04.2018)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

1706162.3 19 April 2017 (19.04.2017) GB

(71) Applicants: NATIONAL UNIVERSITY OF SINGAPORE [SG/SG]; 21 Lower Kent Ridge Road, Singapore 119077 (SG). UNIVERSITY OF MYSORE [IN/IN]; Manasagangotri, Mysore 570006 (IN). BANGALORE UNIVERSITY [IN/IN]; Central College Campus, Palace Road, Bangalore 560001 (IN).

(72) Inventors: LOBIE, Peter Edward; c/o National University of Singapore, Cancer Science Institute of Singapore, 21 Lower Kent Ridge Road, Singapore 119077 (SG). PANDEY, Vijay Kumar; c/o National University of Singapore, Cancer Science Institute of Singapore, 21 Lower Kent Ridge Road, Singapore 119077 (SG). KANCHUGARAKOPPAL SUBBEGOWDA, Rangappa; c/o University of Mysore, Department of Studies in Chemistry, Manasagangotri, Mysore 570006 (IN). SALUNDI, Bassappa; c/o Bangalore University, Department of Chemistry, Central College Campus, Palace Road, Bangalore 560001 (IN). CHAKRABHAVI DHANANJAYA, Mohan; c/o University of Mysore, Department of Studies in Chemistry, Manasagangotri, Mysore 570006 (IN). RANGAPPA, Shobith; c/o University of Mysore, Department of Studies in Chemistry, Manasagangotri, Mysore 570006 (IN). VENKATACHALAI AH, Srinivasa; c/o Bangalore University, Department of Chemistry, Central College Campus, Palace Road, Bangalore 560001 (IN).

(74) Agent: AMICA LAW LLC; 77 Robinson Road, #22-01 Robinson 77, Singapore 068896 (SG).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

— of inventorship (Rule 4.17(iv))

Published:

— with international search report (Art. 21(3))
— in black and white; the international application as filed contained color or greyscale and is available for download from PATENTSCOPE


(48) Date of publication of this corrected version:

17 October 2019 (17.10.2019)

(15) Information about Correction:

see Notice of 17 October 2019 (17.10.2019)

(54) Title: SMALL MOLECULE INHIBITORS OF BCL-2-ASSOCIATED DEATH PROMOTER (BAD) PHOSPHORYLATION



(57) Abstract: The invention relates to compounds of general formula (I); wherein R^1 , n , R^{2a} , R^{2b} , and R^3 are as defined herein. The compounds are inhibitors of Bcl-2-associated death promoter (BAD) phosphorylation and have anti-apoptotic activity and are useful in the treatment of cancer, particularly breast cancer, endometrial cancer, ovarian cancer, liver cancer, colon cancer, prostate cancer or pancreatic cancer.



SMALL MOLECULE INHIBITORS OF BCL-2-ASSOCIATED DEATH
PROMOTER (BAD) PHOSPHORYLATION

The present invention relates to novel compounds, and in particular to compounds which are inhibitors of Bcl-2-associated death promoter (BAD) in cancer cells, to the compounds
5 for use in the treatment or prevention of cancer and to pharmaceutical compositions comprising the compounds.

Apoptosis or programmed cell death is a regulatory mechanism pivotally involved in elimination of auto-reactive immune cells, aged cells and cells with irreparable DNA
10 damage [1]. Apoptosis is executed by two distinct mechanisms in vertebrates – the extrinsic and the intrinsic pathway [2]. The intrinsic pathway is activated under hypoxic conditions, after severe genomic damage, oxidative stress or after growth factor depletion [3]. BCL-2 family proteins possess a pivotal role in regulating the intrinsic apoptotic pathway [4]. More than 20 BCL-2 family members have been identified and broadly
15 classified into anti-apoptotic, pro-apoptotic, and BH3-only proteins. BCL-2, BCL-xL and BCL-w are anti-apoptotic; BAK and BAX are pro-apoptotic; and BAD, BIM, PUMA, NOXA are the predominant pro-apoptotic BH3 only proteins [5-7]. Hence, the members of the BCL-2 family serve both pro- and anti-apoptotic functions with an equilibrium maintained between them, and pro-apoptotic BH3 only proteins are normally repressed to ensure
20 homeostasis [8].


Suppression of apoptosis contributes to inappropriate cell survival and cancer development and/or progression [9-11]. In most malignancies, expression of anti-apoptotic proteins is increased leading to the development of resistance to apoptosis. Bcl-2
25 associated death promoter (BAD) plays a critical role in regulating apoptosis by interacting with Bcl-2, Bcl-xL and Bcl-w [12]. Human BAD is phosphorylated at Ser75 (equivalent murine residue is Ser112) by p44/42 MAP kinase and Ser99 (equivalent murine residue is Ser136) by AKT/p70S6K [13]. Both serine residues are also phosphorylated by Pim family kinases to prevent apoptosis [14]. Phosphorylation of BAD at either of these residues
30 results in loss of the ability of hBAD to heterodimerize with BCL-xL or BCL-2 [15]. Phosphorylated BAD protein is heterodimerized with 14-3-3 protein and sequestered in the cytoplasm [16]. Upon initiation of apoptosis, BAD undergoes dephosphorylation and heterodimerizes with BCL-2, BCL-xL or BCL-w which allows BAK and BAX to promote the release of cytochrome C to the cytoplasm with subsequent promotion of the intrinsic
35 apoptotic pathway [17]. BAD has also been reported to possess other protein and functional interactions within the cell. For example, BAD expression is regulated by p53 and BAD complexes with p53 to promote apoptosis [18]. Bad is transcriptionally up-

regulated by p53 and forms a Bad/p53 complex at the mitochondria to induce apoptosis [18]. Clinically, high BAD expression is associated with high Gleason scores in prostate cancer [15] and BAD phosphorylation has been reported to predict poor overall survival in ovarian cancer [19] and to be associated with resistance to cisplatin [19]. Furthermore, phosphorylated BAD is observed in more than 80% of the CD44 positive cancer stem cell (CSC) population in breast cancer and BAD phosphorylation has been reported to be essential for CSC survival [20].

Numerous small molecule modulators of programmed cell death have been developed, targeting BCL-2 family proteins for therapeutic use against various human malignancies [21-23]. To date there has been no reported small molecule modulators of BH3-only proteins such as BAD. Thus, there is a necessity to develop better and efficient compounds/therapies for managing cancer by inhibiting BAD phosphorylation without affecting the upstream events.

15

In the present invention there is provided a compound of general formula (I):



wherein:

20 each R^1 is independently halo, OH, cyano, nitro, $NR^{10}R^{11}$, $C(O)R^{10}$, $C(O)OR^{10}$, $C(O)NR^{10}R^{11}$, $-S(O)_qNR^{10}R^{11}$, C_{1-6} alkyl, C_{1-6} haloalkyl, $-O(C_{1-6}$ alkyl), $-O(C_{1-6}$ haloalkyl), aryl, heteroaryl, $-O$ -aryl or $-O$ -heteroaryl,

wherein

25 each R^{10} and R^{11} is independently selected from H or C_{1-6} alkyl optionally substituted with one or more substituents selected from OH, halo, cyano, NH_2 , aryl or heteroaryl;

alkyl and haloalkyl groups R^1 are optionally substituted with one or more substituents selected from OH, cyano, $-S(O)_pNR^4R^5$, $-C(O)NR^4R^5$, aryl, heteroaryl, $-O$ -aryl or $-O$ -heteroaryl $-O(C_{1-6}$ alkyl) optionally substituted with aryl or $-O(C_{1-6}$ haloalkyl);

30

aryl or heteroaryl groups R^1 are optionally substituted with one or more substituents selected from halo, OH, cyano, nitro, $-NR^4R^5$, $-S(O)_pNR^4R^5$, $-C(O)NR^4R^5$, $-C(O)R^4$,

-C(O)OR⁴ or -C₁₋₆ alkyl or -O(C₁₋₆ alkyl) , either of which is optionally substituted with one or more substituents selected from OH, halo, aryl, heteroaryl, -O(C₁₋₆ alkyl), O(C₁₋₆ haloalkyl), -O-aryl or -O-heteroaryl);

p is 1 or 2;

5 each R⁴ and R⁵ is independently selected from H or C₁₋₄ alkyl or R⁴ and R⁵ together with a nitrogen atom to which they are attached may form a 3- or 8-membered heterocyclic ring, optionally containing one or more additional heteroatoms selected from O, N and S;

10 n is 0, 1, 2, 3 4, or 5;

R^{2a} and R^{2b} are each independently C₁₋₆ alkyl optionally substituted with one or more substituents selected from halo, OH, aryl or heteroaryl; or

15 R^{2a} and R^{2b} together with the nitrogen atom to which they are attached form a 5- or 6-membered heterocyclic ring optionally containing one or more further heteroatoms selected from O, N or S and optionally substituted with one or more substituents R⁶;

each R⁶ is independently selected from aryl, heteroaryl, -O-aryl, -O-heteroaryl, carbocyclyl, heterocyclyl, -O-carbocyclyl, -O-heterocyclyl, R¹², OR¹², C(O)R¹², C(O)OR¹¹, C(O)NR¹¹R¹², CN, OH,

20 each R¹¹ and R¹² is independently H or C₁₋₄ alkyl, either or which may be substituted with one or more aryl or heteroaryl groups, wherein aryl and heteroaryl groups are substituted with one or more substituent selected from halo, OH, cyano, nitro, -NR⁴R⁵, -S(O)_pNR⁴R⁵, -C(O)NR⁴R⁵, -C(O)R⁴, -C(O)OR⁴ or -C₁₋₆ alkyl or -O(C₁₋₆ alkyl) , either of which is optionally substituted with one or more substituents selected from OH, halo, aryl, heteroaryl, -O(C₁₋₆ alkyl), O(C₁₋₆ haloalkyl), -O-aryl or -O-heteroaryl);

25 wherein R⁴ and R⁵ are as defined above; or R¹¹ and R¹² may combine with a nitrogen atom to which they are attached to form a 3 to 8-membered heterocyclic ring optionally containing one or more further heteroatoms selected from N, O and S and optionally substituted with C₁₋₄ alkyl, C₁₋₄ haloalkyl or halo;

30 R³ is aryl, heteroaryl, carbocyclyl or heterocyclyl any of which is optionally substituted with one or more substituents R⁷ selected from halo, -C₁₋₄ alkyl optionally substituted with aryl, -O(C₁₋₄ alkyl) optionally substituted with aryl, -C₁₋₄ haloalkyl, -O(C₁₋₄ haloalkyl) or -C(O)NR⁸R⁹;

each R⁸ and R⁹ is independently selected from H, C₁₋₄ alkyl or C₃₋₆ cycloalkyl or R⁸ and R⁹ together with the nitrogen atom to which they are attached may form a 5- or 6-membered heterocyclic ring, optionally containing one or more additional heteroatoms selected from O, N and S;

5

or a pharmaceutically acceptable salt, solvate or hydrate thereof or a deuterated or tritiated variant thereof, including all stereoisomers.

10

The compounds of general formula (I) inhibit of site-specific BAD phosphorylation without affecting its upstream kinase and are therefore of use in promoting apoptosis in multiple cancer-derived cells.

15

In the present specification, except where the context requires otherwise due to express language or necessary implication, the word “comprises”, or variations such as “comprises” or “comprising” is used in an inclusive sense i.e. to specify the presence of the stated features but not to preclude the presence or addition of further features in various embodiments of the invention.

20

In the present specification the term “C₁₋₆ alkyl” refers to a fully saturated straight or branched hydrocarbon chain having from 1 to 6 carbon atoms. Examples include methyl, ethyl, n-propyl, isopropyl, t-butyl, n-hexyl.

25

The term “C₁₋₄ alkyl” has a similar meaning but refers to alkyl groups with between 1 and 4 carbon atoms.

30

The term “C₁₋₆ haloalkyl” refers to a C₁₋₆ alkyl group as defined above in which one or more hydrogen atoms are replaced by halo atoms. Haloalkyl groups may have any number of halo substituents from 1 to perhalosubstituted. Examples include chloromethyl, trifluoromethyl, 1-bromoethyl, 1,1,2,2-tetrafluoroethyl etc.

35

The term “carbocyclyl” refers to a non-aromatic hydrocarbon ring having from 3 to 7 carbon atoms and optionally containing one or more carbon-carbon double bond. Examples include C₃₋₇ cycloalkyl groups such as cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and cycloheptyl and cycloalkenyl groups such as cyclohexenyl.

The term “heterocyclyl” refers to a non-aromatic ring having from 5 to 7 carbon atoms and at least one ring heteroatom selected from N, O and S. Examples include

piperazinyl, piperidinyl, morpholinyl, pyrrolidinyl and imidazoliny.

In the present specification, "halo" refers to fluoro, chloro, bromo or iodo.

- 5 The term "aryl" in the context of the present specification refer to a ring system with aromatic character having from 5 to 14 ring carbon atoms unless specified otherwise, and containing up to three rings. Where an aryl group contains more than one ring, not all rings must be fully aromatic in character. Examples of aromatic moieties are benzene, naphthalene, indane and indene.

10


The term "heteroaryl" in the context of the specification refer to a ring system with aromatic character having from 5 to 14 ring atoms (unless specified otherwise) at least one of which is a heteroatom selected from N, O and S, and containing up to three rings. Where a heteroaryl group contains more than one ring, not all rings must be fully aromatic in character. Examples of heteroaryl groups include pyridine, pyrimidine, indole, isoindole, benzofuran, benzimidazole, benzoxazole, benzisoxazole and indolene.

15

In the compounds of general formula (I), n is suitably 2 or 3, for example 2. At least one R¹ group may be OH. In this case, the OH group is suitably at the position adjacent to the -CH(R³)NR^{2a}R^{2b} group.

20

Suitably, the compound of general formula (I) is a compound of general formula general formula (IA):



25

(IA)

wherein R¹, R^{2a}, R^{2b} and R³ are as defined above and m is 0, 1, 2, 3 or 4.

Suitably, in the compound of general formulae (I) and (IA):

- each R¹ is independently halo, C₁₋₆ alkyl, C₁₋₆ haloalkyl, aryl or heteroaryl, wherein aryl or heteroaryl groups are optionally substituted with one or more substituents selected from halo, OH, cyano, nitro, -S(O)_pNR⁴R⁵, -C(O)NR⁴R⁵, -C₁₋₆ alkyl optionally substituted with aryl, -C₁₋₆ haloalkyl, -O(C₁₋₆ alkyl) optionally substituted with aryl or -O(C₁₋₆ haloalkyl);
- 30

p is 0, 1 or 2;

each R⁴ and R⁵ is independently selected from H or C₁₋₄ alkyl or R⁴ and R⁵ together with the nitrogen atom to which they are attached may form a 5- or 6-membered heterocyclic ring, optionally containing one or more additional heteroatoms selected from O, N and S;

R^{2a} and R^{2b} are each independently C₁₋₆ alkyl optionally substituted with one or more substituents selected from halo, OH, aryl or heteroaryl; or

R^{2a} and R^{2b} together with the nitrogen atom to which they are attached form a 5- or 6-membered heterocyclic ring optionally containing one or more further heteroatoms selected from O, N or S and optionally substituted with one or more substituents R⁶;

each R⁶ is independently selected from aryl, heteroaryl, -O-aryl, -O-heteroaryl or C₁₋₄ alkyl substituted with one or more aryl or heteroaryl groups, wherein aryl and heteroaryl groups are substituted with one or more substituent selected from halo, C₁₋₄ alkyl, C₁₋₄ haloalkyl, -O(C₁₋₄ alkyl) or -O(C₁₋₄ haloalkyl);

R³ is aryl, heteroaryl, carbocyclyl or heterocyclyl any of which is optionally substituted with one or more substituents R⁷ selected from halo, -C₁₋₄ alkyl optionally substituted with aryl, -O(C₁₋₄ alkyl) optionally substituted with aryl, -C₁₋₄ haloalkyl, -O(C₁₋₄ haloalkyl) or -C(O)NR⁸R⁹;

each R⁸ and R⁹ is independently selected from H, C₁₋₄ alkyl or C₃₋₆ cycloalkyl or R⁸ and R⁹ together with the nitrogen atom to which they are attached may form a 5- or 6-membered heterocyclic ring, optionally containing one or more additional heteroatoms selected from O, N and S;

or a pharmaceutically acceptable salt, solvate or hydrate thereof or a deuterated or tritiated variant thereof, including all stereoisomers.

In some suitable compounds of general formula (IA), m is 0 so that the group R¹ is not present.

In other suitable compounds m is 1 or 2, but more usually 1.

In such cases, R¹ is suitably halo, particularly chloro or fluoro; or alternatively R¹ is an aryl or heteroaryl group optionally substituted as described above. More usually, R¹ is chloro, fluoro or aryl optionally substituted as described above.


When R¹ is an aryl or heteroaryl group it is suitably at the 4-position of the phenyl ring (where the carbon substituted with the OH group is designated as position 1).

- 5 Suitable substituents for aryl or heteroaryl groups R¹ include halo, OH, cyano, nitro, -SO₂NH₂, -C(O)NR⁴R⁵, -C₁₋₄ alkyl optionally substituted with aryl, -C₁₋₄ haloalkyl, -O(C₁₋₄ alkyl) optionally substituted with aryl or -O(C₁₋₄ haloalkyl), where R⁴ and R⁵ together with the nitrogen atom to which they are attached form a piperidine or pyrrolidine ring.
- 10 More suitable substituents for aryl or heteroaryl groups R¹ include chloro, fluoro, methyl, ethyl, trifluoromethyl, benzyl, methoxy, ethoxy, benzyloxy, trifluoromethoxy and piperidine-1-carbonyl.

15 In some suitable compounds of the invention, R^{2a} and R^{2b} together with the nitrogen atom to which they are attached form a 5- or 6-membered heterocyclic ring optionally containing one or more further heteroatoms selected from O, N or S and optionally substituted with one or more substituents R⁶, wherein R⁶ is as defined above.

20 More suitably, R^{2a} and R^{2b} together with the nitrogen atom to which they are attached form a 6-membered heterocyclic ring optionally substituted with one or more substituents R⁶.

Still more suitably, the 6-membered ring formed by R^{2a} and R^{2b} is a piperidine or piperazine ring, most suitably a piperazine ring. When R^{2a} and R^{2b} form a piperazine ring, it may optionally be substituted by one or more substituents R⁶ as defined above but is suitably substituted by a single R⁶ substituent at the piperazine 4-position such that the compound of general formula (I) is a compound of general formula (IB):




wherein R^1 , n , R^3 and R^6 are as defined for general formulae (I) and (IA).

Particularly suitable substituents R^6 include aryl, heteroaryl, -O-aryl, -O-heteroaryl or methyl substituted with one or two aryl or heteroaryl groups; wherein aryl and heteroaryl groups are substituted with one or more substituents selected from halo, methyl, ethyl, trifluoromethyl, methoxy, ethoxy or trifluoromethoxy.

Still more suitable substituents R^6 include phenyl, heteroaryl, -O-phenyl, -O-heteroaryl, benzyl, -CH(phenyl)₂, -CH₂-heteroaryl and -CH(heteroaryl)₂, where the heteroaryl group is selected from pyridinyl, indolyl, isoindolyl, benzoxazolyl and benzisoxazolyl and wherein any of the above R^6 groups may be substituted as described above, but more suitably with one or more substituents selected from halo, methyl ethyl and trifluoromethyl.

In more suitable compounds of general formula (I) and (IA), R^3 is aryl or heteroaryl optionally substituted with one or more substituents R^7 as described above.

Most suitably, R^3 is phenyl optionally substituted with one or more substituents R^7 such that the compound of general formula (I) is a compound of general formula (IC):




wherein R^1 , n , R^{2a} , R^{2b} and R^7 are as defined for general formula (I) or (IA) and z is 0 to 5.

More suitably, z is 0, 1, 2 or 3, most suitably 0, 1 or 2.

R^7 is as defined above but in some suitable compounds R^7 not present (i.e. z is 0), or R^7 is halo, -C₁₋₄ alkyl, benzyl, -O(C₁₋₄ alkyl) benzyloxy, -C₁₋₄ haloalkyl, -O(C₁₋₄ haloalkyl) or -C(O)NR⁸R⁹, where R^8 and R^9 together with the nitrogen atom to which they are attached form a piperidinyl ring or wherein R^8 is H and R^9 is C₃₋₇ cycloalkyl.


More suitably, R^7 not present or each R^7 is independently halo, especially chloro or fluoro, methyl, ethyl, benzyl, methoxy, ethoxy, benzyloxy, -C(O)-piperidinyl or -C(O)NH-C₃₋₇ cycloalkyl.

Some particularly suitable compounds of the invention are compounds of general formula (ID):



- 5 wherein R^1 , n , R^6 , R^7 and z are as defined above for general formulae (I), (IA), (IB) and (IC).

In one embodiment of the present invention, there is provided a compound of general formula (ID):



10

wherein:

- each R^1 is independently halo, OH, cyano, nitro, $NR^{10}R^{11}$, $C(O)R^{10}$, $C(O)OR^{10}$, $C(O)NR^{10}R^{11}$, $-S(O)_qNR^{10}R^{11}$, C_{1-6} alkyl, C_{1-6} haloalkyl, $-O(C_{1-6}$ alkyl), $-O(C_{1-6}$ haloalkyl), aryl, heteroaryl, $-O$ -aryl or $-O$ -heteroaryl, wherein an aryl includes a ring system with aromatic character having from 5 to 14 ring carbon atoms unless specified otherwise and contains up to three rings, and wherein an heteroaryl includes a ring system with aromatic character having from 5 to 14 ring carbon atoms unless specified otherwise, at least one of which is a heteroatom selected from N, O, and S and contains up to three rings;
- 15
- 20

wherein

each R¹⁰ and R¹¹ is independently selected from H or C₁₋₆ alkyl optionally substituted with one or more substituents selected from OH, halo, cyano, NH₂, aryl or heteroaryl;

5 alkyl and haloalkyl groups R¹ are optionally substituted with one or more substituents selected from OH, cyano, -S(O)_pNR⁴R⁵, -C(O)NR⁴R⁵, aryl, heteroaryl, -O-aryl, -O-heteroaryl, -O(C₁₋₆ alkyl) optionally substituted with aryl or -O(C₁₋₆ haloalkyl);

10 aryl or heteroaryl groups R¹ are optionally substituted with one or more substituents selected from halo, OH, cyano, nitro, -NR⁴R⁵, -S(O)_pNR⁴R⁵, -C(O)NR⁴R⁵, -C(O)R⁴, -C(O)OR⁴ or -C₁₋₆ alkyl or -O(C₁₋₆ alkyl), any of which is optionally substituted with one or more substituents selected from OH, halo, aryl, heteroaryl, -O(C₁₋₆ alkyl), O(C₁₋₆ haloalkyl), -O-aryl or -O-heteroaryl);

p is 1 or 2;

15 each R⁴ and R⁵ is independently selected from H or C₁₋₄ alkyl or R⁴ and R⁵ together with a nitrogen atom to which they are attached may form a 3- or 8-membered heterocyclic ring, optionally containing one or more additional heteroatoms selected from O, N and S;

20 n is 0, 1, 2, 3 or 4;

R⁶ is selected from aryl, -O-aryl, -O-heteroaryl, carbocyclyl, heterocyclyl, -O-carbocyclyl, -O-heterocyclyl, R¹², OR¹², C(O)R¹², C(O)OR¹¹, C(O)NR¹¹R¹², CN, OH,

25 each R¹¹ and R¹² is independently H or C₁₋₄ alkyl, either or which may be substituted with one or more aryl or heteroaryl groups, wherein aryl and heteroaryl groups are substituted with one or more substituent selected from halo, OH, cyano, nitro, -NR⁴R⁵, -S(O)_pNR⁴R⁵, -C(O)NR⁴R⁵, -C(O)R⁴, -C(O)OR⁴, -C₁₋₆ alkyl or -O(C₁₋₆ alkyl), either of which is optionally substituted with one or more substituents selected from OH, halo, aryl, heteroaryl, -O(C₁₋₆ alkyl), -O(C₁₋₆ haloalkyl), -O-aryl, or -O-heteroaryl;

30

wherein R⁴ and R⁵ are as defined above; or R¹¹ and R¹² may combine with a nitrogen atom to which they are attached to form a 3 to 8-membered heterocyclic ring optionally containing one or more further heteroatoms selected from N, O and S and optionally substituted with C₁₋₄ alkyl, C₁₋₄ haloalkyl or halo;

R⁷ selected from halo, -C₁₋₄ alkyl optionally substituted with aryl, -O(C₁₋₄ alkyl) optionally substituted with aryl, -C₁₋₄ haloalkyl, -O(C₁₋₄ haloalkyl) or -C(O)NR⁸R⁹; wherein

each R⁸ and R⁹ is independently selected from H, C₁₋₄ alkyl or C₃₋₆ cycloalkyl or R⁸ and R⁹ together with the nitrogen atom to which they are attached may form a 5- or 6-membered heterocyclic ring, optionally containing one or more additional heteroatoms selected from O, N and S; and

z is 0 to 5;

or a pharmaceutically acceptable salt thereof; wherein the compound is not 2-[(4-methylphenyl)(4-phenyl-1-piperazinyl)methyl]phenol.

Example compounds of general formula (I) include:

2-((2-chlorophenyl)(4-(4-methoxyphenyl)piperazin-1-yl)methyl)phenol (Compound 1);

2-((4-chlorophenyl)(4-(4-methoxyphenyl)piperazin-1-yl)methyl)phenol (Compound 2);

2-((4-(benzyloxy)-3-fluorophenyl)(4-(4-methoxyphenyl)piperazin-1-yl)methyl)phenol (Compound 3);

(4-((2-hydroxyphenyl)(4-(4-Methoxyphenyl)piperazinyl)methyl)phenyl)(piperidin-1-yl)methanone (Compound 4);

3-((5-chloro-2-hydroxyphenyl)(4-(4-methoxyphenyl)piperazin-1-yl)methyl)-N-cyclopentylbenzamide (Compound 5);

2-((4-(benzyloxy)-3-fluorophenyl)(4-(4-methoxyphenyl)piperazin-1-yl)methyl)-4-chlorophenol (Compound 6);

2-((4-(benzyloxy)-3-fluorophenyl)(4-(6-fluorobenzo[d]isoxazol-3-yl)piperidin-1-yl)methyl)phenol (Compound 7);

2-((4-(2, 3-dichlorophenyl)piperazin-1-yl)(o-tolyl)methyl)phenol (Compound 8);

N-cyclopentyl-3-((4-(2,3-dichlorophenyl)piperazin-1-yl)(2-hydroxyphenyl)methyl)benzamide (Compound 9, NPB);

2-((4-(benzyloxy)-3-fluorophenyl)(4-(2,3-dichlorophenyl)piperazin-1-yl)methyl)phenol

2018256285 14 Apr 2022

(Compound 10);

2-((4-((4-chlorophenyl)(phenyl)methyl)piperazin-1-yl)(phenyl)methyl)phenol

(Compound 11);

2-((4-((4-chlorophenyl)(phenyl)methyl)piperazin-1-yl)(p-tolyl)methyl)phenol


5 (Compound

- 12);
2-((4-chlorophenyl)(4-((4-chlorophenyl)(phenyl)methyl)piperazin-1-yl)methyl)phenol
(Compound 13);
2-((4-((4-chlorophenyl)(phenyl)methyl)piperazin-1-yl)(4-ethylphenyl)methyl)phenol
5 (Compound 14);
4-((4-((4-chlorophenyl)(phenyl)methyl)piperazin-1-yl)(2-hydroxyphenyl)methyl)phenyl
(piperidin-1-yl)methanone (Compound 15);
N-cyclopentyl-3-((4-(2,3-dichlorophenyl)piperazin-1-yl)(4-hydroxy-[1,1'-biphenyl]-3-
yl)methyl)benzamide (Compound 16);
10 N-cyclopentyl-3-((4-(2,3-dichlorophenyl)piperazin-1-yl)(4-hydroxy-2'-methyl-[1,1'-
biphenyl]-3-yl)methyl)benzamide (Compound 17);
N-cyclopentyl-3-((4-(2,3-dichlorophenyl)piperazin-1-yl)(4-hydroxy-3'-methyl-[1,1'-
biphenyl]-3-yl)methyl)benzamide (Compound 18);
N-cyclopentyl-3-((4-(2,3-dichlorophenyl)piperazin-1-yl)(4-hydroxy-4'-methyl-[1,1'-
15 biphenyl]-3-yl)methyl)benzamide (Compound 19);
3-((2'-chloro-4-hydroxy-[1,1'-biphenyl]-3-yl)(4-(2,3-dichlorophenyl)piperazin-1-yl)methyl)-
N-cyclopentylbenzamide (Compound 20);
3-((3'-chloro-4-hydroxy-[1,1'-biphenyl]-3-yl)(4-(2,3-dichlorophenyl)piperazin-1-yl)methyl)-
N-cyclopentylbenzamide (Compound 21);
20 3-((4'-chloro-4-hydroxy-[1,1'-biphenyl]-3-yl)(4-(2,3-dichlorophenyl)piperazin-1-yl)methyl)-
N-cyclopentylbenzamide (Compound 22);
N-cyclopentyl-3-((4-(2,3-dichlorophenyl)piperazin-1-yl)(4'-ethyl-4-hydroxy-[1,1'-biphenyl]-
3-yl)methyl)benzamide (Compound 23);
N-cyclopentyl-3-((4-(2,3-dichlorophenyl)piperazin-1-yl)(4-hydroxy-4'-(piperidine-1-
25 carbonyl)-[1,1'-biphenyl]-3-yl)methyl)benzamide (Compound 24);
N-cyclopentyl-3-((4-(2,3-dichlorophenyl)piperazin-1-yl)(4-hydroxy-4'-methoxy-[1,1'-
biphenyl]-3-yl)methyl)benzamide (Compound 25);
N-cyclopentyl-3-((4-(2,3-dichlorophenyl)piperazin-1-yl)(2'-ethyl-4-hydroxy-[1,1'-biphenyl]-
3-yl)methyl)benzamide (Compound 26);
30 N-cyclopentyl-3-((4-(2,3-dichlorophenyl)piperazin-1-yl)(2'-fluoro-4-hydroxy-[1,1'-
biphenyl]-3-yl)methyl)benzamide (Compound 27);
N-cyclopentyl-3-((4-(2,3-dichlorophenyl)piperazin-1-yl)(3'-fluoro-4-hydroxy-[1,1'-
biphenyl]-3-yl)methyl)benzamide (Compound 28);
N-cyclopentyl-3-((4-(2,3-dichlorophenyl)piperazin-1-yl)(4'-fluoro-4-hydroxy-[1,1'-
35 biphenyl]-3-yl)methyl)benzamide (Compound 29);
N-cyclopentyl-3-((4-(2,3-dichlorophenyl)piperazin-1-yl)(4-hydroxy-3'-nitro-[1,1'-biphenyl]-
3-yl)methyl)benzamide (Compound 30);

- N-cyclopentyl-3-((4-(2,3-dichlorophenyl)piperazin-1-yl)(4-hydroxy-3'-sulfamoyl-[1,1'-biphenyl]-3-yl)methyl)benzamide (Compound 31);
- N-cyclopentyl-3-((4-(2,3-dichlorophenyl)piperazin-1-yl)(4-hydroxy-2'-(trifluoromethyl)-[1,1'-biphenyl]-3-yl)methyl)benzamide (Compound 32);
- 5 N-cyclopentyl-3-((4-(2,3-dichlorophenyl)piperazin-1-yl)(4-hydroxy-3'-(trifluoromethyl)-[1,1'-biphenyl]-3-yl)methyl)benzamide (Compound 33);
- N-cyclopentyl-3-((4-(2,3-dichlorophenyl)piperazin-1-yl)(4-hydroxy-4'-(trifluoromethyl)-[1,1'-biphenyl]-3-yl)methyl)benzamide (Compound 34);
- 3-((2'-cyano-4-hydroxy-[1,1'-biphenyl]-3-yl)(4-(2,3-dichlorophenyl)piperazin-1-yl)methyl)-
- 10 N-cyclopentylbenzamide (Compound 35);
- 3-((3'-cyano-4-hydroxy-[1,1'-biphenyl]-3-yl)(4-(2,3-dichlorophenyl)piperazin-1-yl)methyl)-
- N-cyclopentylbenzamide (Compound 36)
- 3-((4'-cyano-4-hydroxy-[1,1'-biphenyl]-3-yl)(4-(2,3-dichlorophenyl)piperazin-1-yl)methyl)-
- N-cyclopentylbenzamide (Compound 37);
- 15 3-((2'-chloro-4-hydroxy-4'-(trifluoromethyl)-[1,1'-biphenyl]-3-yl)(4-(2,3-dichlorophenyl)piperazin-1-yl)methyl)-N-cyclopentylbenzamide (Compound 38);
- N-cyclopentyl-3-((2',4'-dichloro-4-hydroxy-[1,1'-biphenyl]-3-yl)(4-(2,3-dichlorophenyl)piperazin-1-yl)methyl)benzamide (Compound 39);
- 3-((4'-chloro-2',4-dihydroxy-[1,1'-biphenyl]-3-yl)(4-(2,3-dichlorophenyl)piperazin-1-
- 20 yl)methyl)-N-cyclopentylbenzamide (Compound 40);
- 3-((4-(4-chlorophenyl)piperazin-1-yl)(2-hydroxyphenyl)methyl)-N-cyclopentylbenzamide (Compound 41, NCK1);
- 2-((4-chlorophenyl)(4-(2,3-dichlorophenyl)piperazin-1-yl)methyl)phenol (Compound 42, NCK2);
- 25 2-((4-(2,3-dichlorophenyl)piperazin-1-yl)(3-methoxyphenyl)methyl)phenol (Compound 43, NCK3);
- 1-(5-((4-(2,3-dichlorophenyl)piperazin-1-yl)(2-hydroxyphenyl)methyl)thiophen-2-yl)ethanone (Compound 44, NCK4);
- 2-((4-(2,3-dichlorophenyl)piperazin-1-yl)(naphthalen-1-yl)methyl)phenol (Compound 45,
- 30 NCK5);
- 5-((4-(2,3-dichlorophenyl)piperazin-1-yl)(2-hydroxyphenyl)methyl)furan-2-carbaldehyde (Compound 46, NCK6);
- 2-((4-(5,6-dichlorocyclohexa-1,5-dien-1-yl)piperazin-1-yl)(2-fluoro-3-methylpyridin-4-yl)methyl)phenol (Compound 47, NCK7);
- 35 2-((4-(2,3-dichlorophenyl)piperazin-1-yl)(4-(trifluoromethyl)phenyl)methyl)phenol (Compound 48, NCK8);
- 2-((6-chloro-5-methylpyridin-3-yl)(4-(2,3-dichlorophenyl)piperazin-1-yl)methyl)phenol

- (Compound 49, NCK9);
 2-((4-(2,3-dichlorophenyl)piperazin-1-yl)(pyridin-3-yl)methyl)phenol (Compound 50, NCK10);
 1-(5-((4-(4-chlorophenyl)piperazin-1-yl)(2-hydroxyphenyl)methyl)thiophen-2-yl)ethanone
 5 (Compound 51, NCK14);
 3-((4-(4-chlorophenyl)piperazin-1-yl)(4-(diethylamino)-2-hydroxyphenyl)methyl)-N-cyclopentylbenzamide(Compound 52, NCK16);
 N-cyclopentyl-3-((4-(2,3-dichlorophenyl)piperazin-1-yl)(4-(diethylamino)-2-hydroxyphenyl)methyl)benzamide (Compound 53, NCK18);
 10 N-cyclopentyl-3-((4-(2,3-dichlorophenyl)piperazin-1-yl)(2-hydroxy-4,6-dimethoxyphenyl)methyl)benzamide (Compound 54, NCK19);
 2-((4-chlorophenyl)(4-(4-chlorophenyl)piperazin-1-yl)methyl)phenol (Compound 55, NCK20);
 2-((4-(4-chlorophenyl)piperazin-1-yl)(6-methylpyridin-3-yl)methyl)phenol (Compound 56,
 15 NCK21);
 2-(o-tolyl(4-(p-tolyl)piperazin-1-yl)methyl)phenol (Compound 57, SG1);
 2-((4-(p-tolyl)piperazin-1-yl)(4-(trifluoromethyl)phenyl)methyl)phenol (Compound 58, SG2);
 N-cyclopentyl-4-((2-hydroxyphenyl)(4-(p-tolyl)piperazin-1-yl)methyl)benzamide
 20 (Compound 59, SG3);
 2-((4-chlorophenyl)(4-(p-tolyl)piperazin-1-yl)methyl)phenol (Compound 60, SG4);
 2-((3-methoxyphenyl)(4-(p-tolyl)piperazin-1-yl)methyl)phenol (Compound 61, SG5);
 5-((2-hydroxyphenyl)(4-(p-tolyl)piperazin-1-yl)methyl)furan-2-carbaldehyde (Compound 62, SG6); and
 25 2-((6-methylpyridin-3-yl)(4-(p-tolyl)piperazin-1-yl)methyl)phenol (Compound 63, SG7).


Compounds of general formula (I) may be prepared by reacting an aldehyde of general formula (II):



30


wherein R^1 and n are as defined for general formula (I);

with a compound of general formula (III):



wherein R^{2a} and R^{2b} are as defined for general formula (I);

- 5 and a boronic acid of general formula (IV):



wherein R³ is as defined for general formula (I).

- 10 The reaction is a Petasis reaction, which is a 3-component boronic mannich- type reaction which uses boronic acids of general formula (IV) as a potential nucleophilic species to react with the salicylaldehyde derivative of general formula (II) and the amine of general formula (III) to form a new carbon-carbon bond. The Petasis reaction proceeds via the formation of an iminium species, which reacts with the boronic acid to yield tertiary amines.

15

Suitable reaction solvents include organic solvents, for example cyclic ethers such as dioxane.

Suitably, the process comprises:

- 20 i. reacting the compound of general formula (II) with the compound of general formula (III) to form a mixture comprising intermediate iminium species;
- ii. reacting the product of step (i) with the compound of general formula (IV).

- 25 Suitably step (ii) is carried out at elevated temperature, for example 50 to 100°C, for example at the reflux temperature of the solvent. When the solvent is dioxane, this is about 85 to 95°C, typically about 90°C.

The product may be extracted into a second solvent such as ethyl acetate, dried and purified using standard methods such as described below in the examples.

30

Compounds of general formulae (II), (III) and (IV) are known and are commercially available or may be prepared using known methods.

Compounds of general formula (I) may also be prepared from other compounds of general formula (I). For example, a compound of general formula (I) in which R¹ is optionally substituted aryl or heteroaryl may be prepared from compounds of general formula (I) in which R¹ is halo by reaction with a compound of general formula (V):



wherein R^{1a} is aryl or heteroaryl optionally substituted as defined for general formula (I).

The reaction may be carried out using a Suzuki coupling reaction over a palladium catalyst.

10

Compounds of general formula (V) are known and are commercially available or may be prepared using known methods.

The methods for preparing compounds of general formula (I) represent a further aspect of the present invention.

15

The compounds of general formula (I) are capable of inhibiting/downregulating/suppressing Bcl-2 associated death promoter (BAD) protein, particularly inhibiting/downregulating/suppressing phosphorylation of Bcl-2 associated death promoter (BAD) protein in a cancer cell. The inhibition/downregulation/suppression of BAD occurs without affecting the upstream Ser/Thr kinase called Akt/protein kinase B (PKB). Therefore, the compounds of general formula (I) preferentially inhibit/downregulate/suppresses phosphorylation of BAD.

20

The inhibition/downregulation of BAD is carried out by the compound of general formula (I) occupying the hydrophobic groove within the protein-protein interface of Bcl-2/BAD. Additionally, the compound of general formula (I) occupies an additional hydrophobic side pocket within the interface formed by the side-chains of Leu-97, Trp-144, and Phe-198 of the human protein with a dichlorophenyl moiety.

25

The compounds of formula (I) are potent inhibitors/downregulators of BAD in cancer cells, particularly breast cancer, endometrial cancer, ovarian cancer, liver cancer, colon cancer, prostate cancer and pancreatic cancer cells and induce apoptosis due to the downregulation/inhibition of BAD phosphorylation. Said compounds enhance cytotoxicity

30

of cancer cells and carry out target specific downregulation/inhibition of BAD phosphorylation without affecting the upstream kinases.

The compounds of general formula (I) were screened for their anticancer activity against MCF7 cells and NPB (Compound 9) was found to be the most potent compound. NPB was subsequently evaluated against panel of normal immortalized mammary epithelial cells (MCF10A and MCF12A), normal hepatocytes (LO2), mammary carcinoma (BT549, MDA-MB-231, MCF7, T47D, BT474), endometrial cancer (Ishikawa, Ecc1, RL95-2, AN3), ovarian cancer (SK-OV-3, OVCAR-2, Caov-3, HEY C2, Ovca433), liver cancer (Hep3B, H2P, H2M), colon cancer (HCT116, DLD-1, Caco-2), prostate cancer (PC3, LNCaP, DU145) and pancreatic cancer (AsPC-1, BxPC-3) cells. NPB significantly enhanced apoptosis in all the tested cancer cell lines and NPB exhibited no substantial activity against normal hepatocytes and immortalized mammary epithelial cells. Using a Laplacian-modified Naïve Bayes classifier, NPB is subjected to in silico analysis which anticipates the human target of NPB as BAD protein.

NPB was found to induce apoptotic cell death in a range of carcinoma cells but did not have cytotoxic effect on normal (*immortalized*) epithelial cells.

The hallmark features of cells undergoing apoptosis are phosphatidylserine externalization, caspase activation and genomic DNA fragmentation. In an embodiment, NPB exhibits apoptosis against MCF7 cells using FITC-annexin-V and propidium iodide staining. FITC-annexin-V staining confirms the loss of membrane integrity and PI staining confirms the late apoptotic events and these results are correlated with caspase 3/7 activity.

Therefore in a further aspect of the invention, there is provided a compound of general formula (I) for use in medicine.

The compounds of general formula (I) are of particular use in the treatment of cancer, suitably, a cancer in which there is BAD phosphorylation..

In a further aspect, therefore, there is provided a compound of general formula (I) for use in the treatment of cancer.

Further, there is provided the use of a compound of general formula (I) in the preparation of an agent for the treatment of cancer.

5 There is also provided a method for the treatment of cancer, the method comprising administering to a patient in need of such treatment an effective amount of a compound of general formula (I).

10 The compounds are particularly useful for treating cancer in which there is BAD phosphorylation and, for example, the cancer may be breast cancer, endometrial cancer, ovarian cancer, liver cancer, colon cancer, prostate cancer or pancreatic cancer or other epithelial-derived cancers in which BAD is phosphorylated.

15 The compounds of general formula (I) are suitably administered in a pharmaceutical composition and therefore in a further aspect of the invention there is provided a pharmaceutical composition comprising a compound of general formula (I) and a pharmaceutically acceptable excipient.

20 The pharmaceutically acceptable excipient may be an adjuvant, diluent, carrier, granulating agent, binding agent, lubricating agent, disintegrating agent, sweetening agent, glidant, anti-adherent, anti-static agent, surfactant, anti-oxidant, gum, coating agent, coloring agent, flavouring agent, coating agent, plasticizer, preservative, suspending agent, emulsifying agent, plant cellulosic material, spheronization agent, other conventionally known pharmaceutically acceptable excipient or any combination of excipients thereof.

25

In another embodiment, the pharmaceutical composition of the present invention is formulated for intraperitoneal administration, hepatoportal administration, intravenous administration, intra articular administration, pancreatic duodenal artery administration or intramuscular administration, or any combination thereof.

30

The invention will now be described in greater detail with reference to the examples and to the drawings in which:

Figures 1(a) and 1(b) depict the ^1H -NMR and ^{13}C -NMR spectra of Compound 1.

35

Figure 2 depicts the ^{13}C -NMR spectra of compound 2.

Figures 3(a) and 3(b) depict the ^1H -NMR and ^{13}C -NMR spectra of Compound 3.

Figures 4(a) and 4(b) depict the ^1H -NMR and ^{13}C -NMR spectra of Compound 4.

5 Figures 5(a) and 5(b) depict the ^1H -NMR and ^{13}C -NMR spectra of Compound 5.

Figure 6 depicts the ^1H -NMR spectra of compound 6.

Figures 7(a) and 7(b) depict the ^1H -NMR and ^{13}C -NMR spectra of Compound 7.

10

Figure 8 depicts the ^{13}C -NMR spectra of Compound 8.

Figures 9(a) and 9(b) depict the ^1H -NMR and LCMS spectra of Compound 9.

15 Figure 10 depicts the ^1H -NMR spectra of Compound 10.

Figure 11: IC50 values of NPB in a range of carcinoma cell lines.

20 **Figure 12: NPB suppresses cell viability and promote apoptosis in carcinoma cell lines.**

Effect of NPB (5 μM) on carcinoma cell viability including, mammary, endometrial, ovarian, liver, colon, prostate, and pancreatic carcinoma cell lines. (A) Cell viability (B) caspase 3/7 activities and (C) cytotoxicity were evaluated using ApoTox-Glo™ Triplex Assay Kit as described in methodology. Statistical significance was assessed by an unpaired two-tailed Student's *t*-test using GraphPad Prism5. The column represents mean of triplicate determinations; bars, $\pm\text{SD}$. ** $P < 0.001$, * $P < 0.05$. **Note:** RFU, relative fluorescence unit; RLU, relative luminescence unit, #; non-transformed, immortalized epithelial cells; MB-231, MDA-MB-231.

30 **Figure 13: NPB stimulates apoptotic cell death in MCF7 cells.**

(A) Apoptotic cell death of MCF7 cells measured after treatment with 10 μM NPB using flow cytometry analysis. Annexin V-FITC staining is indicated on the X-axis and PI staining on the Y-axis. The lower left quadrant represents live cells, the lower right quadrant represents early apoptotic cells, the upper left quadrant represents necrotic cells, and the upper right quadrants display late apoptotic cells. Acquisition of Annexin V and PI data were represented as a percentage (%) in each quadrant. (B) Cell cycle analysis of MCF7 cells measured after treatment with 10 μM NPB using flow cytometry analysis. (C) Cell

35

viability of performed colonies generated by MCF7 cell after exposure to NPB or DMSO cultured 14 days in 3D Matrigel using AlamarBlue® viability assay. Microscopic visualization (*below*) of Calcein AM stained colonies generated by MCF7 cells after exposure to NPB or DMSO cultured in 3D Matrigel. (D) Cell viability in colonies generated by MCF7 cell after exposure to NPB or DMSO cultured in Soft agar using AlamarBlue® viability assay. (E) Crystal Violet staining of foci colonies generated by MCF7 cells after exposure to NPB or DMSO. All assays performed as described in methodology. Statistical significance was assessed by an unpaired two-tailed *Student's t-test* using GraphPad Prism5. The column represents mean of triplicate determinations; bars, $\pm SD$. ** $P < 0.001$, * $P < 0.05$.

Figure 14: Cheminformatics and surface plasmon resonance (SPR) analysis predicts an interaction of NPB compound to BAD protein.

(A) Sensorgrams obtained by SPR analysis of NPB with the BAD protein subunit. The BAD protein subunit was immobilized onto the surface of a CM5 sensor chip. A solution of NPB at variable concentrations (20-100 μ M) was injected to generate result binding responses (RU) recorded as a function of time (sec). The results were analyzed using BIA evaluation 3.1. (B) Western blot (WB) analysis was used to assess the level of Ser99 phosphorylation of BAD in MCF7 cells after treatment with NPB. (*Below*) Calculated IC_{50} of NPB from dose-response for BAD phosphorylation (Ser99), BAD and β -ACTIN as in shown above by use of ImageJ software from NIH, USA (<http://imagej.nih.gov/ij/>). (C) WB analysis was used to assess the level of a multiple proteins involved upstream of BAD in MCF7 cells after treatment with NPB. (D) WB analysis was used to assess the level of a multiple protein involved cell survival and cell proliferation in MCF7 cells after treatment with NPB. For WB analysis, soluble whole cell extracts were run on an SDS-PAGE and immunoblotted as described in methodology. β -ACTIN (ACTB) was used as an input control for cell lysate. The sizes of detected protein bands in kDa are shown on the *left side*.

Figure 15: NPB specifically inhibits phosphorylation of BAD (at Ser99) in carcinoma cell lines independent of AKT signalling

(A) WB analysis was used to assess the levels of phosphorylated human BAD (at Ser75 and Ser99) and BAD protein in the range of carcinoma cell lines, including mammary, ovarian, pancreatic, endometrial, hepatocellular, colon and prostate cancer after treatment with NPB (5 μ M). Total BAD was used as an input control for cell lysate. (B) WB analysis was used to assess the levels of pBAD (Ser99) and pAKT (Ser473), AKT and BAD in MCF7, Caov-3, Ishikawa, and AsPC-1 cells. 5 μ M each of AKT inhibitor (IV) and NPB was

used to treat cells. Depletion of *AKT* expression was achieved using transient-transfection of *short hairpin (sh)-RNA* (1&2) directed to *AKT* transcript as described in methodology. β -ACTIN was used as an input control for cell lysate. For WB analysis, soluble whole cell extracts were run on an SDS-PAGE and immunoblotted as described in materials methodology. The sizes of detected protein bands in kDa are shown on the *left side*. **Note:** #; non-transformed *immortalized*-cell line.

Figure 16: *siRNA-mediated* depletion of *BAD* expression prevents the effect of NPB in carcinoma cell lines.

10 **(A)** WB analysis was used to assess the levels of pBAD (Ser99) activity and BAD protein in MCF7, BT474, Caov-3, Ishikawa, AsPC-1, and DLD-1 cells after treatment with 5 μ M NPB. Depletion of *BAD* expression was achieved using transient-transfection of *small interfering (si)-RNA* directed to the *BAD* transcript. Soluble whole cell extracts were run on an SDS-PAGE and immunoblotted as described in materials and methods. β -ACTIN was used as input control. Effects of NPB (5 μ M) in MCF7, BT474, Caov-3, Ishikawa, AsPC-1, and DLD-1 cells. **(B)** Cell viability and **(C)** caspase 3/7 activities were evaluated using the ApoTox-Glo™ Triplex Assay Kit. All assays performed as described in methodology. Statistical significance was assessed by an unpaired two-tailed *Student's t-test* ($P < 0.05$ was considered as significant) using GraphPad Prism5. The column represents mean of triplicate determinations; bars, \pm SD. **** $P < 0.001$, * $P < 0.05$** . **Note** RFU, relative fluorescence unit; RLU, relative luminescence unit.

Figure 17: NPB inhibits phosphorylation of BAD Ser99 in mammary carcinoma and inhibits tumour growth

25 **(A)** Measurement of tumour volume in BALB/c-nu female mice as described in materials and methods. Animals ($n=5$ each group) were treated with vehicle, 5 mg/kg NPB or 20mg/kg NPB, and relative tumour burden was recorded. Animal weight was measured daily for the duration of the experiment. **(B)** Tumours were excised after the NPB treatment regime and weighed. Representative resected tumours are shown in the *right side*. **(C)** WB of tumour tissue to determine levels of p-BAD (Ser99) and BAD. Soluble whole cell extracts were run on an SDS-PAGE and immunoblotted as described in methodology. β -ACTIN was used as an input control. The sizes of detected protein bands in kDa are shown on the *left side*. **(D)** Histological analyses of phospho-BAD, BAD, Ki67 and TUNEL staining. Tumour tissue sections were immunolabeled with goat anti-pBAD (Ser 136) polyclonal antibody (Santa Cruz Biotechnology), mouse anti-BAD monoclonal (Santa Cruz Biotechnology) and anti-Ki67 antibody (Abcam, ab15580) and stained with hematoxylin. Apoptotic DNA fragmentation was detected using TUNEL Apoptosis Detection Kit (Gen

30

35

Script USA Inc.) as described in methodology. Statistical significance was assessed by an unpaired two-tailed *Student's t-test* ($P < 0.05$ was considered as significant) using GraphPad Prism5. The point represents mean of triplicate experiments; bars, $\pm SD$. $**P < 0.001$, $*P < 0.05$.

5

Figure 18:

(A) Western blot analysis was used to assess the level of BAD Ser99 phosphorylation of pBAD, BAD, pAKT, and AKT in MCF7 cells after an increasing period of treatment with NPB (10 μ M). Soluble whole cell extracts were run on an SDS-PAGE and immunoblotted as described in methodology. The sizes of detected protein bands in kDa are shown on the *left side*. (B) Kinases and phosphorylated substrates were detected using a Western Blot array (Proteome Profiler Human Phospho-Kinase Array Kit. MCF7 cells treated with NPB (10 μ M) or DMSO for 12h at 37 $^{\circ}$ C before preparation of cell lysate. Mean pixel density was analysed using ImageJ software and is represented *below*.

10

15 **Figure 19** depicts the $^1\text{H-NMR}$ spectra of Compound NCK5.

Figure 20 depicts the $^1\text{H-NMR}$ spectra of Compound NCK16.

20 **Figure 21** depicts the $^1\text{H-NMR}$ spectra of Compound NCK18.

Figures 22A, 22B, 22C, 22D and 22E: IC₅₀ values of NPB structure-based analogues in a carcinoma cell lines.

Note: NV, no value

25

Materials employed to arrive at the Examples of the present disclosure*Cell culture and Reagents –*

The human *immortalized* mammary epithelial cell lines, MCF10A, and MCF12A; and immortalized hepatocellular epithelial cell line, LO2 were obtained from the American Type Culture Collection (ATCC, Rockville, MD) and were cultured as per ATCC propagation instructions. MC cell lines, MCF7, T47D, BT474, BT549, and MDA-MB-231 (denoted as MB-231); endometrial carcinoma cell lines, Ishikawa, ECC1, RL95-2 and AN3; hepatocellular carcinoma cell lines, Hep3B, H2P, and H2M; colon carcinoma cell lines, HCT116, DLD-1, and Caco-2; and prostate carcinoma cell lines, PC3, LNCaP, DU145 were obtained from the American Type Culture Collection (ATCC, Rockville, MD). Ovarian carcinoma cell lines, SK-OV-3, OVCAR-2, Caov-3, HEY C2, and Ovca433 were obtained from Dr Ruby Huang's laboratory at The Cancer Science Institute of Singapore, National

30

35

University of Singapore (NUS). Pancreatic carcinoma cell lines were obtained from Prof. H. Phillip Koeffler's laboratory at The Cancer Science Institute of Singapore, National University of Singapore (NUS). All carcinoma cell lines were cultured as per ATCC propagation instructions. AKT inhibitor IV was purchased from Calbiochem (San Diego, CA, USA). BAD directed stealth (sh)-RNA-BAD (shRNA-BAD1, 5'-GCUCCGCACCAUGAGUGACGAGUUU-3' and shRNA-BAD2, 5'-AAACUCGUCACUCAUCCUCCGGAGC3') was purchased from Life Technologies (Singapore). AKT directed shRNA (shRNA-AKT1, 5'-CCGGCGCGTGACC ATGAACGAGTTTCTCGAGAACTCGTTCATGGTCACGCGTTTTTGTG-3' and shRNA2-AKT, 5'-CCGGGGACTACCTGCACTCGGAGAACTCGAGTTCTCCGAGTGCAGGTAG TCCTTTTTTGTG-3') was purchased from Life Technologies (Singapore), and cloned in to PLKO.1 vector (Sigma, Singapore). Cells were transiently-transfected with 20nM shRNA (AKT or BAD) or universal negative control (Invitrogen, Carlsbad, CA, USA) using FuGENE HD (Promega) for 24h and further assays performed. Alanine transaminase (ALT), aspartate transaminase (AST), lactate dehydrogenase (LDH), creatine kinase (CK), blood urea nitrogen (BUN) commercial kits were purchased from AGAPPE Diagnostics Ltd, Kerala, India.

Example 1

Synthesis and Characterization of Formula I compounds

General Synthesis of Compound of Formula I

























Piperazines (0.8mmol) and salicylaldehyde (0.8mmol) are taken in an RBF and stirred for about 10 minutes using Dioxane as solvent. After about 10 minutes, Aryl boronic acid (0.8mmol) is added to the mixture and refluxed with continuous stirring for about 8 hours using Dioxane as solvent on a hot plate maintained at about 90°C. After about 8 hours, ethyl acetate and water are added to the reaction mixture and the ethyl acetate layer is separated using separating funnel and dried over anhydrous sodium sulfate. Ethyl acetate is evaporated to obtain the product. The desired phenolic compound product is obtained by separation using column chromatography.

























30

The specific reagents used to obtain Compounds 1 to 15 and 41 to 63 are provided in Table 1 below.

























TABLE 1





















No	Salicylaldehyde	Piperazine	Aryl boronic acid	Product
1				
2				
3				
4				
5				
6				
7				

No	Salicylaldehyde	Piperazine	Aryl boronic acid	Product
8				
9				
10				
11				
12				
13				

No	Salicylaldehyde	Piperazine	Aryl boronic acid	Product
14				
15				
41				 Molecular Weight: 490.0363
42				 Molecular Weight: 447.7846
43				 Molecular Weight: 443.3656
44				 Molecular Weight: 461.4040

No	Salicylaldehyde	Piperazine	Aryl boronic acid	Product
45				
				Molecular Weight: 463.3983
46				
				Molecular Weight: 431.3118
47				
				Molecular Weight: 448.3606
48				
				Molecular Weight: 481.338
49				
				Molecular Weight: 462.7993
50				
				Molecular Weight: 414.3276

No	Salicylaldehyde	Piperazine	Aryl boronic acid	Product
51				 Molecular Weight: 426.9589
52				 Molecular Weight: 561.16
53				 Molecular Weight: 595.60
54				 Molecular Weight: 584.53
55				 Molecular Weight: 413.34
56				

No	Salicylaldehyde	Piperazine	Aryl boronic acid	Product
58				 Molecular Weight: 426.4740
57				 Molecular Weight: 372.5026
59				
61				 Molecular Weight: 388.5020
60				 Molecular Weight: 392.9211

No	Salicylaldehyde	Piperazine	Aryl boronic acid	Product
62				
63				

The compounds shown in Table 2 were obtained by reacting NPB bromide (see column 1 of Table 2) with a boronic acid using a palladium catalysed Suzuki coupling reaction as shown in Scheme 1.

5

Scheme 1


























































Table 2













	NPB-Br	Boronic acid	Product
16			
17			
18			










<p>19</p>			
<p>20</p>			
<p>21</p>			
<p>22</p>			

23			
24			
25			

<p>26</p>			
<p>27</p>			
<p>28</p>			
<p>29</p>			

30			
31			
32			
33			

34			
35			
36			
37			

38			
39			
40			

Characterization of Compound 1:

^1H NMR (CDCl_3 , 400 MHz) δ : 3.691(s, 3H), 5.303(s, 1H, C-H), 1.184-3.691(m, 8H- piperazine protons), 7.597-7.615(d, 1H, $J=7.2$ Hz), 7.341-7.323(d, 1H, $J=7.2$ Hz), 7.060-7.189(m, 4H-ArH), 6.555-6.815(m, 5H-ArH), 6.956-6.974, (d, 1H, $J=7.2$ Hz) 11.85(s, 1H-OH brd peak); ^{13}C NMR (400 MHz, CDCl_3) δ : 50.854, 55.521, 69.50, 114.45, 117.17, 118.44, 119.547, 122.05, 127.78, 128.83, 129.01, 129.19, 129.84, 129.93, 133.89, 145.11, 156.60; Melting point 120-124°C. (FIGURE 1)

10 Characterization of Compound 2:

^1H NMR (CDCl_3 , 400 MHz) δ : 2.539-3.065(m, 8H), 3.674(s, 3H), 4.359(s, 1H), 6.651(m, 1H), 6.740-6.804(m, 5H), 6.855-6.872(d, 1H, $J=6.8$ Hz), 7.086(m, 1H), 7.165(m, 2H),

7.280(m, 1H), 7.361(m, 1H); ^{13}C NMR (400 MHz, CDCl_3) δ : 50.74, 51.76, 55.52, 75.86, 114.49, 117.24, 118.43, 119.62, 124.41, 126.58, 128.28, 128.54, 128.92, 129.20, 130.28, 134.66, 141.63, 145.07, 154, 156.18; Melting point 85-89°C. **(FIGURE 2)**

5 Characterization of Compound 3:

^1H NMR (CDCl_3 , 400 MHz) δ : 1.194-3.079(m,8H), 3.689(s, 3H), 4.336(s, 1H), 5.041(s, 2H), 6.672-6.702(m, 1H), 6.781-6.819(m, 5H), 6.863-6.867(m, 2H), 7.026-7.082(m, 2H), 7.196(m, 1H), 7.279-7.348(m, 5H); ^{13}C NMR (400 MHz, CDCl_3) δ : 50.770, 55.52, 71.26, 75.36, 114.46, 115.46, 117.16, 118.42, 119.54, 124.82, 127.34, 128.14, 128.62, 128.76,
10 129.14, 145.04, 156.14; Melting point 72-76°C. **(FIGURE 3)**

Characterization of Compound 4:

^1H NMR (CDCl_3 , 400 MHz) δ : 1.183-2.469(m, 10H), 2.557-3.684(m, 8H), 3.684(s, 3H), 4.412(s, 1H), 6.631(m, 2H), 6.745-6.813(m, 4H), 7.059-7.095(m, 1H), 7.191-7.216(m, 1H),
15 7.261-7.269(m, 2H), 7.351-7.409(m, 2H); ^{13}C NMR (400 MHz, CDCl_3) δ : 24.51, 26.525, 29.68, 43.60, 48.73, 50.71, 51.80, 55.51, 76.04, 114.45, 117.134, 118.366, 119.198, 119.52, 124.75, 126.74, 127.50, 128.04, 128.50, 128.79, 129.29, 136.18, 141.02, 145, 154, 156.2, 169.79(C=O); Melting point 78-82°C. **(FIGURE 4)**

20 Characterization of Compound 5:

^1H NMR (CDCl_3 , 400 MHz) δ : 1.186-1.650(m, 8H), 1.978-3.633(m, 8H), 3.689(s, 3H), 4.298-4.345(m, 1H), 4.421(s, 1H), 5.979-5.992(s, 1H) 6.736-6.803(m, 5H), 6.860(m, 1H), 7.192(s, 1H), 7.737(s, 1H), 7.304-7.339(m, 1H), 7.540-7.557(m, 2H); ^{13}C NMR (400 MHz, CDCl_3) δ : 23.79, 33.18, 50.67, 51.83, 55.51, 67.06, 75.06, 114.46, 118.45, 118.58, 124.02,
25 126.11, 126.42, 128.72, 128.87, 129.39, 139.39, 144.89, 154.19, 154.95, 166.68; Melting point 102-106°C. **(FIGURE 5)**

Characterization of Compound 6:

^1H NMR (CDCl_3 , 400 MHz) δ : 1.179-3.620(m, 8H), 3.676(s, 3H), 4.261(s, 1H), 5.029(s, 2H),
30 6.716-6.791(m, 5H), 6.823(m, 1H), 6.862-6.882(m, 1H), 6.973-7.015(m, 2H), 7.110-7.139(m, 1H), 7.243-7.258(m, 2H), 7.280-7.317(m, 1H), 7.331-7.350(m, 2H); ^{13}C NMR (400 MHz, CDCl_3) δ : 50.70, 51.59, 55.52, 71.27, 74.93, 114.48, 115.52, 118.58, 123.97, 124.39, 126.3, 127.3, 128.2, 128.8, 132.1, 136.27, 144.96, 146.74, 154.19, 154.93; Melting point 60-64°C. **(FIGURE 6)**

Characterization of Compound 7:

¹H NMR (CDCl₃, 400 MHz) δ: 2.648-3.820 (m, 8H) 4.245(s, 1H), 5.009(s, 1H), 5.651(s, 2H), 7.276(s, 1H), 7.436-7.485(m, 3H), 7.606-7.682(m, 3H), 7.797(m, 2H), 7.876-7.954(m, 5H), 8.185(s, 1H); ¹³C NMR (400 MHz, CDCl₃)δ: 30.972, 50.76, 67.60, 71.79, 75.85, 98.16, 100.9, 104.9, 113.16, 115.97, 117.65, 119.96, 122.84, 125.4, 127.8, 128.6, 129.1, 129.6, 133.2, 136.8, 157.06, 160.87, 164.51, 165.86; Melting point 58-62°C. (FIGURE 7)

Characterization of Compound 8:

¹H NMR (CDCl₃, 400 MHz) δ: 2.479(s, 3H), 4.927(s, 1H), 2.260-3.063(m, 8H), 6.533-6.551(d, 1H, J=7.2Hz), 6.631-6.692(m, 2H), 6.739-6.758(d, 1H, J=7.6Hz), 6.789-6.809(d, 1H, J=8Hz), 6.864(m, 3H), 7.092-7.183(m, 1H), 7.281-7.297(m, 1H), 7.537-7.552(d, 1H, J=6Hz); ¹³C NMR (400 MHz, CDCl₃) δ: 20.92, 51.16, 51.42, 73.44, 116.07, 116.96, 117.14, 118.716, 119.27, 119.83, 124.965, 125.24, 126.445, 127.12, 127.545, 128.266, 128.729, 129.260, 130.869, 134.072, 138.171, 150.600, 156.443 Melting point 108-112°C. (FIGURE 8)

Characterization of Compound 9 (NPB):

¹H NMR (CDCl₃, 400 MHz) δ: 1.183-1.647(m, 8H), 2.019-3.067(m, 8H), 4.509(s, 1H), 4.312-4.327(m, 1H, NH), 5.965(s, 1H), 6.668(m, 1H), 6.801-6.896(m, 3H), 7.073-7.190(m, 3H), 7.305(m, 1H), 7.527-7.542(m, 2H), 7.770(s, 1H); ¹³C NMR (400MHz, CDCl₃)δ: 23.78, 33.18, 51.22, 51.78, 76.10, 117.14, 118.59, 119.67, 124.69, 124.98, 126.22, 127.53, 128.85, 129.29, 131.08, 134.08, 135.53, 140.28, 150.5, 156.1, 166.73; m/z (M+2, 526.2, 527.2) Melting point 174-178°C. (FIGURE 9)

Characterization of Compound 10:

¹H NMR (CDCl₃, 400 MHz) δ: 1.183-3.074(m, 8H), 4.361(s, 1H), 5.034(s, 2H), 6.658-6.694(t, 1H, J=7.2Hz), 6.768(m, 1H), 6.856-6.873(m, 3H), 7.023(m, 1H), 7.055-7.094(m, 3H), 7.164(m, 1H), 7.231(m, 1H), 7.266(m, 1H), 7.304(m, 1H), 7.322-7.355(m, 2H); ¹³C NMR (400 MHz, CDCl₃)δ: 51.264, 71.260, 75.434, 117.15, 118.6, 119.6, 124.8, 124.9, 127.3, 127.5, 128.1, 128.6, 128.8, 129.18, 150.5, 156.09; Melting point 75-80°C. (FIGURE 10)

Example 2Compound of formula I decreases the cell viability of a range of carcinoma cellsOncogenicity Assay

We initially investigated the effect of newly synthesized small molecule compounds against MCF7 cells (ER+ MC cells) using an AlamarBlue® cell viability assay. Among the series

of novel small molecule compounds, NPB was identified as an efficacious small molecule compound reducing viability of MCF7 cells compared to vehicle (DMSO) treated cells.

Among the compounds, Compound 9 N-cyclopentyl-3-((4-(2,3-dichlorophenyl)piperazin-1-yl)(2-hydroxyphenyl)methyl)benzamide (NPB) is identified as the most potent antiproliferative compound with the IC₅₀ of 6.5 μM. We next determined the inhibitory concentration 50% (IC₅₀) of NPB in wide-range of carcinoma cell lines including those derived from ER- mammary (BT549, MDA-MB-231), ER+ mammary (MCF7, T47D, and BT474), endometrial (Ishikawa, ECC-1, RL95-2, and AN3), ovarian (SK-OV-3, OVCAR-2, Caov-3, HEY C2, and OVCA433), hepatocellular (Hep3B, H2P, and H2M), colon (HCT116, DLD-1, and Caco-2), prostate (PC3, LNCaP, and DU145) and pancreatic (AsPC-1, BxPC-3) carcinoma. As normal cell controls, we also included *immortalized* mammary epithelial cells (MCF10A and MCF12A), and *immortalized* hepatocytes (LO2) in the panel of cell lines. The IC₅₀ values for NPB in the carcinoma cell lines are tabulated in Figure 11.

Example 3

NPB induces apoptotic cell death in range of carcinoma cell lines

In addition, NPB is evaluated against several cancer-derived cell lines to determine effect on whole cell viability, apoptosis, and cytotoxicity using ApoTox-Glo™ Triplex Assay Kit, Promega (Singapore) according to manufacturer's instructions (Figure 12). In brief, cells are seeded in black opaque 96-well plates (Corning®, Singapore). After an overnight incubation of cells, the medium is changed to the indicated NPB concentration. After about 48 hours of incubation, the viability/cytotoxicity reagent containing both GF-AFC substrate and bis-AAF-R110 substrate are added to the cells as suggested by the supplier. After about 45 minutes incubation at about 37 °C, fluorescence is recorded at 400 nm excitation/505 nm emission for viability and 485 nm excitation/520 nm emission for cytotoxicity using a Tecan microplate reader for fluorescence (Tecan, Singapore). Caspase-Glo 3/7 Reagent is further added to the cells and after about 25 minutes of incubation at room temperature, luminescence is recorded using Tecan microplate reader. Numbers of viable, cytotoxic, and apoptotic cells are measured in triplicates.

Annexin V and propidium iodide (Annexin V–PI) apoptosis assay

Phosphatidylserine exposure and cell death are assessed by FACS analysis using Annexin-V-FLUOS Staining Kit (Life Technologies, Singapore) and PI-stained cells. Briefly, 1×10⁵ MCF cells/well (190μL/well) are seeded in 6-well plates and incubated with different concentrations of NPB for about 24 hours and DMSO treated samples are used

as control. Cells are then washed with Annexin V binding buffer (10mM HEPES/NaOH, pH7.4, 140 mM NaCl, 2.5 mM CaCl₂), stained with Annexin V FITC for about 30 minutes at room temperature in the dark, then washed again and re-suspended in Annexin V binding buffer containing PI. Samples are analyzed immediately on a BD FACSAria Cell
5 Sorter (BD Biosciences, San Jose, CA).

Loss of membrane integrity and translocation of phosphatidylserine to outer leaflet of plasma membrane are the early events of apoptosis which can be detected using FITC conjugated annexin-V and propidium iodide staining. It is observed that NPB induces apoptosis in MCF7 cells using FITC-annexin V and propidium iodide. On treatment with
10 NPB, increase in both early (PI negative, FITC-Annexin V positive) and late apoptotic cells (PI positive, FITC-Annexin V positive) is observed in a dose dependent manner as shown in Figure 13A.

Example 4

Molecular interaction of NPB with recombinant BAD protein

To provide molecular interaction of the most promising candidate, NPB binding affinity to BAD, we performed surface plasmon resonance (SPR) measurements with immobilized BAD subunit using NPB as the analyte. We know that BAD could bind to BAD subunit *in vitro*. Hence, we analyzed the interaction using the BIAcore system. The recombinant BAD
20 was immobilized on CM5 sensor chip. To determine the association and dissociation curves, various concentrations of NPB were injected individually onto the surface of a sensor chip coated with BAD. The overlaid sensorgrams shown in Figure 14 were analyzed collectively. The direct binding of BAD to NPB was demonstrated (Figure 14A). The calculation of kinetic parameters for the interaction of NPB with BAD revealed the
25 association rate constant of $(1.4 \pm 0.4) \times 10^3 \text{ M}^{-1} \text{ S}^{-1}$ and dissociation rate constant of $(5.4 \pm 0.38) \times 10^3 \text{ S}^{-1}$ of binding affinity, which yielded dissociation equilibrium constants (K_d) of 37.12 μM. These kinetic parameters shows affinity support for the interaction of BAD with NPB structure.

Surface plasmon resonance analysis

Molecular interactions were analyzed based on surface plasmon resonance using a BIAcore-2000 system (BIAcore AB, Uppsala, Sweden). Human recombinant BAD protein (Catalog No.MBS143012, MyBiosource, USA) was immobilized on a sensor chip as described by the manufacturer protocol. To examine the interaction of BAD with NPB,
35 various concentrations of NPB (20 to 100 μM) in the running buffer (HBS-EP, pH 7.4, BIAcore AB) were injected onto the surface of the BAD-immobilized sensor chip with a flow rate of 15 μl/min as per the manufacturer's directions. NPB was allowed to interact

with BAD subunit for 2 min for association and dissociation, respectively, after which the sensor chip was regenerated by injecting 1 M NaCl for 2 min before the next injection. Using BIA evaluation software 4.1 (BIAcore AB), the kinetic parameters were such as association and dissociation rate constants (k_a and k_d), dissociation equilibrium constants (Kd) using a 1:1 binding model with mass transfer. Sensograms obtained were overlaid using BIA evaluation software.

Example 5

Effect of NPB on BAD phosphorylation is specific for Ser99 (human)

10 Phosphorylation of hBAD at residues Ser-75 (mouse BAD serine residue 112) and Ser-99 (mouse BAD serine residue 136) are crucial in regulating the activity of the BCL-2 family of anti-apoptotic proteins [15]. hBAD phosphorylation either at Ser-75 or Ser99 (or the corresponding residues in mouse bad) results in loss of the ability of hBAD to heterodimerize with BCL-xL or BCL-2 [15]. To further validate the predicted target, we first
15 analyzed the effect of NPB on phosphorylation of hBAD at Ser99 by western blot analysis. Treatment of MCF7 cells with NPB produced a dose dependent decrease in phosphorylation of hBAD Ser99 without a significant change in total hBAD protein (Figure 14B). The calculated EC_{50} for inhibition of BAD Ser99 phosphorylation by NPB was $0.41 \pm 0.21 \mu\text{M}$.

20 We next analyzed the effect of NPB on phosphorylation of hBAD at both Ser75 and Ser99 by western blot analysis in 25 carcinoma cell lines derived seven different types of cancer. It was observed that NPB largely inhibits the phosphorylation of BAD at the Ser99 site in all the tested carcinoma cell lines; however, NPB demonstrated no effect on the phosphorylation of hBAD at the Ser75 site in the same cells indicating that NPB specifically
25 inhibited phosphorylation at Ser99 of hBAD (Figure 15).

siRNA-mediated depletion of BAD expression revert effect of NPB in carcinoma cell lines.

To confirm the functional specificity of NPB directed to the BAD protein, we further
30 examined the effect of NPB exposure after *siRNA-mediated* depletion of *BAD* expression in 6 carcinoma cell lines (MCF7, BT474, Caov-3, Ishikawa, AsPC-1 and DLD-1). Transient-transfection of the different carcinoma cells with *siRNA* directed to the *BAD* transcript decreased BAD expression and also decreased levels of phosphor-Ser99 BAD compared to their control cells (transfected with *scrambled oligo*) as observed by western blot
35 analysis (Figure 16A). Cell viability nor apoptosis were significantly altered upon *siRNA* mediated depletion of BAD as previously reported [26]. As described above, NPB treatment of the control transfected carcinoma cell lines decreased BAD phosphorylation

(Ser99) compared to vehicle-treated cells. Concomitantly, exposure of the same carcinoma cell lines to NPB decreased cell viability and increased caspase 3/7 activity compared to vehicle-exposed cells. In contrast, NPB did not affect cell viability nor caspase3/7 activity in carcinoma cell lines with depleted expression of *BAD* (Figure 5 16B&C).

Example 6

NPB inhibits BAD phosphorylation independent of AKT signaling in carcinoma cell lines

The upstream AKT Ser/Thr kinase regulates the phosphorylation of hBAD at Ser99 [13]. We therefore determined whether NPB inhibits the phosphorylation of hBAD (Ser99) via modulation of the activity of AKT (as indicated by phosphorylation of Ser473) using western blot analysis. We observed no change in the levels of pAKT or levels of total AKT protein after exposure of four different carcinoma cells lines (MCF7, Caov-3, Ishikawa, and AsPC-1) to 10 μ M NPB. However, all NPB treated carcinoma cell lines (MCF7, Caov-3, Ishikawa, and AsPC-1) exhibited inhibition of BAD phosphorylation at the Ser99 site and with no change in the level of total BAD protein (Figure 15B). Additionally, we examined BAD phosphorylation after depletion of AKT using two independent *shRNA* targeting *AKT* expression or inhibition of AKT activity with AKT inhibitor IV as a positive control in the different carcinoma cell lines. We observed that depleted expression of AKT in the carcinoma cell lines was associated with a concomitant decrease in pAKT (Ser474) and pBAD (Ser99) levels compared to control cells; indicative that BAD Ser99 phosphorylation is AKT dependent in all tested cancer-derived cell lines and as previously published by others [13, 15, 27-29]. NPB therefore specifically inhibits BAD phosphorylation at Ser99 without affecting the activity of the upstream kinase (AKT) [29]. These results are concordant with the *in silico* target prediction and NPB binding to BAD observed by SPR. Hence, NPB specifically inhibits phosphorylation of BAD at Ser99 independent of the upstream (AKT) kinase.

30

Example 7:

5- to 6-week-old BALB/c-nu female mice were subcutaneously implanted with 17 β -estradiol pellets (Innovative Research of America) at 0.72mg/pellet with a 60-day release in the scruff of the neck after three days mice were injected subcutaneously with 100 μ l of cell suspension (1x10⁷ cells) in right flanks. Tumour growth was monitored by measuring the tumour size using callipers. About 12 days after implantation, mice were randomized and divided into three groups (each group, n=8), according to treatments administered

35

200µl of NPB (dissolved in 5% DMSO, 50% PEG400 and 45% water pH 5.0) by intraperitoneal injection every day for seven days. The first group of mice was treated with vehicle, the second with 5mg/kg dose of NPB, and the third with 20mg/kg dose of NPB. Animal weight and tumour volumes were measured daily. After completion tumours were excised, photographed, weighed, and fixed or stored in liquid nitrogen for later analysis. Histological analysis was performed as previously described (30-32).

We examined the *in vivo* efficacy of NPB in a xenograft (MCF7) of MC. Randomly grouped mice with preformed tumours (volume ~150cm³) were injected intraperitoneally with vehicle or NPB at 5mg/kg or 20mg/kg. A significant reduction in tumour volume was observed in NPB-treated mice as compared to their vehicle-treated counterparts (Figure 17A). During this period, animal weight was not significantly different between the groups (Figure 17A, below). However, the tumour weight of NPB-treated animals was reduced compared to vehicle-treated mice and in a dose-dependent manner (Figure 17B). We further analysed the effect of NPB on hBAD Ser99 phosphorylation levels in tumour tissue using WB analysis (Figure 17C). NPB treatment significantly inhibited phosphorylation of BAD (at Ser99) in a tumour compared to control specimens. No change was observed in total levels of BAD protein between NPB treated, and the control treated tumours.

Histological analyses of tumour specimens resected from the animals treated with NPB showed significantly reduced p-BAD (Ser99) compared to vehicle-treated tumours (Figure 17D), whereas BAD protein was not significantly different between the groups. Animals treated with NPB exhibited a significantly decreased percentage of Ki67 positive cells in tumours and a significantly increased TUNEL positivity compared to vehicle-treated (Figure 17D).

Example 8:

To elucidate the possibility that NPB decreased hBAD Ser99 phosphorylation by modulation of kinase activity, we assessed the effects of NPB on various kinases using Human Phospho-Kinase Antibody Array Kit from R&D Systems. No significant changes in kinase activity or phosphorylated substrates were observed in MCF7 cells exposed to NPB compared to DMSO exposed cells despite NPB inhibition of hBAD Ser99 phosphorylation in the same extract (Figure 18).

Example 9:

We also generated further analogues of NPB (Figures 19,20,21) according to the claimed chemical template and which may exhibit better pharmacokinetic profiles than NPB. Their structure and in vitro efficacy as determined by IC50 is shown in Figure 22.

5

References


1. Rakesh KS, Jagadish S, Vinayaka AC, Hemshekhar M, Paul M, Thushara RM, Sundaram MS, Swaroop TR, Mohan CD and Sadashiva MP. A New Ibuprofen Derivative
10 Inhibits Platelet Aggregation and ROS Mediated Platelet Apoptosis. PloS one. 2014; 9(9):e107182.
2. Krautwald S, Ziegler E, Rolver L, Linkermann A, Keyser KA, Steen P, Wollert KC, Korf-Klingebiel M and Kunzendorf U. Effective blockage of both the extrinsic and intrinsic pathways of apoptosis in mice by TAT-crmA. The Journal of biological chemistry. 2010;
15 285(26):19997-20005.
3. Riedl SJ and Shi Y. Molecular mechanisms of caspase regulation during apoptosis. Nat Rev Mol Cell Biol. 2004; 5(11):897-907.
4. Hardwick JM and Soane L. Multiple functions of BCL-2 family proteins. Cold Spring Harbor perspectives in biology. 2013; 5(2).
- 20 5. Czabotar PE and Lessene G. Bcl-2 family proteins as therapeutic targets. Current pharmaceutical design. 2010; 16(28):3132-3148.
6. Anilkumar U and Prehn JH. Anti-apoptotic BCL-2 family proteins in acute neural injury. Frontiers in cellular neuroscience. 2014; 8:281.
7. Alberts B, Johnson A, Lewis J, Raff M, Roberts K and Walter P. Molecular Biology
25 of the Cell. New York: Garland Science; 2008. Classic textbook now in its 5th Edition. 2010.
8. Hutt KJ. The role of BH3-only proteins in apoptosis within the ovary. Reproduction (Cambridge, England). 2015; 149(2):R81-r89.
9. Ambrosini G, Adida C and Altieri DC. A novel anti-apoptosis gene, survivin,
30 expressed in cancer and lymphoma. Nature medicine. 1997; 3(8):917-921.
10. Keerthy HK, Mohan CD, Siveen KS, Fuchs JE, Rangappa S, Sundaram MS, Li F, Girish KS, Sethi G and Bender A. Novel synthetic biscoumarins target tumor necrosis factor- α in hepatocellular carcinoma in vitro and in vivo. Journal of Biological Chemistry. 2014; 289(46):31879-31890.
- 35 11. Smith AJ, Karpova Y, D'Agostino R, Jr., Willingham M and Kulik G. Expression of the Bcl-2 protein BAD promotes prostate cancer growth. PloS one. 2009; 4(7):e6224.

12. Doerflinger M, Glab JA and Puthalakath H. BH3-only proteins: A 20-year stock-take. *The FEBS journal*. 2015.
13. Hayakawa J, Ohmichi M, Kurachi H, Kanda Y, Hisamoto K, Nishio Y, Adachi K, Tasaka K, Kanzaki T and Murata Y. Inhibition of BAD phosphorylation either at serine 112
5 via extracellular signal-regulated protein kinase cascade or at serine 136 via Akt cascade sensitizes human ovarian cancer cells to cisplatin. *Cancer research*. 2000; 60(21):5988-5994.
14. Macdonald A, Campbell DG, Toth R, McLauchlan H, Hastie CJ and Arthur JS. Pim kinases phosphorylate multiple sites on Bad and promote 14-3-3 binding and dissociation
10 from Bcl-XL. *BMC cell biology*. 2006; 7:1.
15. Fang X, Yu S, Eder A, Mao M, Bast RC, Jr., Boyd D and Mills GB. Regulation of BAD phosphorylation at serine 112 by the Ras-mitogen-activated protein kinase pathway. *Oncogene*. 1999; 18(48):6635-6640.
16. Masters SC, Yang H, Datta SR, Greenberg ME and Fu H. 14-3-3 inhibits Bad-
15 induced cell death through interaction with serine-136. *Molecular pharmacology*. 2001; 60(6):1325-1331.
17. Harada H, Becknell B, Wilm M, Mann M, Huang LJ-s, Taylor SS, Scott JD and Korsmeyer SJ. Phosphorylation and inactivation of BAD by mitochondria-anchored protein kinase A. *Molecular cell*. 1999; 3(4):413-422.
- 20 18. Jiang P, Du W, Heese K and Wu M. The Bad guy cooperates with good cop p53: Bad is transcriptionally up-regulated by p53 and forms a Bad/p53 complex at the mitochondria to induce apoptosis. *Molecular and cellular biology*. 2006; 26(23):9071-9082.
19. Marchion DC, Cottrill HM, Xiong Y, Chen N, Bicaku E, Fulp WJ, Bansal N, Chon HS, Stickles XB, Kamath SG, Hakam A, Li L, Su D, Moreno C, Judson PL, Berchuck A, et
25 al. BAD phosphorylation determines ovarian cancer chemosensitivity and patient survival. *Clinical cancer research : an official journal of the American Association for Cancer Research*. 2011; 17(19):6356-6366.
20. Sastry KSR, Al-Muftah MA, Li P, Al-Kowari MK, Wang E, Ismail Chouchane A, Kizhakayil D, Kulik G, Marincola FM, Haoudi A and Chouchane L. Targeting proapoptotic
30 protein BAD inhibits survival and self-renewal of cancer stem cells. *Cell death and differentiation*. 2014; 21(12):1936-1949.
21. Anderson MA, Huang D and Roberts A. Targeting BCL2 for the treatment of lymphoid malignancies. *Seminars in hematology*. 2014; 51(3):219-227.
22. Oltersdorf T, Elmore SW, Shoemaker AR, Armstrong RC, Augeri DJ, Belli BA,
35 Bruncko M, Deckwerth TL, Dinges J and Hajduk PJ. An inhibitor of Bcl-2 family proteins induces regression of solid tumours. *Nature*. 2005; 435(7042):677-681.

23. Keerthy HK, Garg M, Mohan CD, Madan V, Kanojia D, Shobith R, Nanjundaswamy S, Mason DJ, Bender A and Rangappa KS. Synthesis and characterization of novel 2-amino-chromene-nitriles that target Bcl-2 in acute myeloid leukemia cell lines. *PloS one*. 2014; 9(9):e107118.
- 5 24. Bruncko M, Oost TK, Belli BA, Ding H, Joseph MK, Kunzer A, Martineau D, McClellan WJ, Mitten M, Ng SC, Nimmer PM, Oltersdorf T, Park CM, Petros AM, Shoemaker AR, Song X, et al. Studies leading to potent, dual inhibitors of Bcl-2 and Bcl-xL. *Journal of medicinal chemistry*. 2007; 50(4):641-662.
- 10 25. Petros AM, Nettlesheim DG, Wang Y, Olejniczak ET, Meadows RP, Mack J, Swift K, Matayoshi ED, Zhang H, Thompson CB and Fesik SW. Rationale for Bcl-xL/Bad peptide complex formation from structure, mutagenesis, and biophysical studies. *Protein science : a publication of the Protein Society*. 2000; 9(12):2528-2534.
26. Boisvert-Adamo K and Aplin AE. Mutant B-RAF mediates resistance to anoikis via Bad and Bim. *Oncogene*. 2008; 27(23):3301-3312.
- 15 27. Seow HF, Yip WK, Loh HW, Ithnin H, Por P and Rohaizak M. Immunohistochemical detection of phospho-Akt, phospho-BAD, HER2 and oestrogen receptors alpha and beta in Malaysian breast cancer patients. *Pathology oncology research : POR*. 2010; 16(2):239-248.
28. Kanamori Y, Kigawa J, Itamochi H, Shimada M, Takahashi M, Kamazawa S, Sato S, Akeshima R and Terakawa N. Correlation between loss of PTEN expression and Akt phosphorylation in endometrial carcinoma. *Clinical cancer research : an official journal of the American Association for Cancer Research*. 2001; 7(4):892-895.
29. Datta SR, Brunet A and Greenberg ME. Cellular survival: a play in three Akts. *Genes & development*. 1999; 13(22):2905-2927.
- 25 30. Pandey V, Wu ZS, Zhang M, Li R, Zhang J, Zhu T, et al. Trefoil factor 3 promotes metastatic seeding and predicts poor survival outcome of patients with mammary carcinoma. *Breast Cancer Res*. 2014;16(5):429.
31. Wang XN, Wang SJ, Pandey V, Chen P, Li Q, Wu ZS, et al. Trefoil factor 3 as a novel biomarker to distinguish between adenocarcinoma and squamous cell carcinoma. *Medicine (Baltimore)*. 2015;94(20):e860.
- 30 32. You M, Chen Y, Chong Q, Wu M, Pandey V, Chen R, et al. Trefoil factor 3 mediation of oncogenicity and chemoresistance in hepatocellular carcinoma is AKT-BCL-2 dependent. *Oncotarget*. 2017.

CLAIMS

1. A compound of general formula (ID):



wherein:

each R^1 is independently halo, OH, cyano, nitro, $NR^{10}R^{11}$, $C(O)R^{10}$, $C(O)OR^{10}$, $C(O)NR^{10}R^{11}$, $-S(O)_qNR^{10}R^{11}$, C_{1-6} alkyl, C_{1-6} haloalkyl, $-O(C_{1-6}$ alkyl), $-O(C_{1-6}$ haloalkyl), aryl, heteroaryl, $-O$ -aryl or $-O$ -heteroaryl, wherein an aryl includes a ring system with aromatic character having from 5 to 14 ring carbon atoms unless specified otherwise and contains up to three rings, and wherein a heteroaryl includes a ring system with aromatic character having from 5 to 14 ring carbon atoms unless specified otherwise, at least one of which is a heteroatom selected from N, O, and S and contains up to three rings;

wherein

each R^{10} and R^{11} is independently selected from H or C_{1-6} alkyl optionally substituted with one or more substituents selected from OH, halo, cyano, NH_2 , aryl or heteroaryl;

alkyl and haloalkyl groups R^1 are optionally substituted with one or more substituents selected from OH, cyano, $-S(O)_pNR^4R^5$, $-C(O)NR^4R^5$, aryl, heteroaryl, $-O$ -aryl, $-O$ -heteroaryl, $-O(C_{1-6}$ alkyl) optionally substituted with aryl or $-O(C_{1-6}$ haloalkyl);

aryl or heteroaryl groups R^1 are optionally substituted with one or more substituents selected from halo, OH, cyano, nitro, $-NR^4R^5$, $-S(O)_pNR^4R^5$, $-C(O)NR^4R^5$, $-C(O)R^4$, $-C(O)OR^4$ or $-C_{1-6}$ alkyl or $-O(C_{1-6}$ alkyl), any of which is optionally substituted with one or more substituents selected from OH, halo, aryl, heteroaryl, $-O(C_{1-6}$ alkyl), $O(C_{1-6}$ haloalkyl), $-O$ -aryl or $-O$ -heteroaryl);

p is 1 or 2;

each R^4 and R^5 is independently selected from H or C_{1-4} alkyl or R^4 and R^5 together with a nitrogen atom to which they are attached may form a 3- or 8-

membered heterocyclic ring, optionally containing one or more additional heteroatoms selected from O, N and S;

n is 0, 1, 2, 3 or 4;

R⁶ is selected from aryl, -O-aryl, -O-heteroaryl, carbocyclyl, heterocyclyl, -O-carbocyclyl, -O-heterocyclyl, R¹², OR¹², C(O)R¹², C(O)OR¹¹, C(O)NR¹¹R¹², CN,OH,

each R¹¹ and R¹² is independently H or C₁₋₄ alkyl, either or which may be substituted with one or more aryl or heteroaryl groups, wherein aryl and heteroaryl groups are substituted with one or more substituent selected from halo, OH, cyano, nitro, -NR⁴R⁵, -S(O)_pNR⁴R⁵, -C(O)NR⁴R⁵, -C(O)R⁴, -C(O)OR⁴, -C₁₋₆ alkyl or -O(C₁₋₆ alkyl), either of which is optionally substituted with one or more substituents selected from OH, halo, aryl, heteroaryl, -O(C₁₋₆ alkyl), -O(C₁₋₆ haloalkyl), -O-aryl, or -O-heteroaryl;

wherein R⁴ and R⁵ are as defined above; or R¹¹ and R¹² may combine with a nitrogen atom to which they are attached to form a 3 to 8-membered heterocyclic ring optionally containing one or more further heteroatoms selected from N, O and S and optionally substituted with C₁₋₄ alkyl, C₁₋₄ haloalkyl or halo;

R⁷ selected from halo, -C₁₋₄ alkyl optionally substituted with aryl, -O(C₁₋₄ alkyl) optionally substituted with aryl, -C₁₋₄ haloalkyl, -O(C₁₋₄ haloalkyl) or -C(O)NR⁸R⁹; wherein

each R⁸ and R⁹ is independently selected from H, C₁₋₄ alkyl or C₃₋₆ cycloalkyl or R⁸ and R⁹ together with the nitrogen atom to which they are attached may form a 5- or 6-membered heterocyclic ring, optionally containing one or more additional heteroatoms selected from O, N and S; and

z is 0 to 5;

or a pharmaceutically acceptable salt thereof; wherein the compound is not 2-[(4-methylphenyl)(4-phenyl-1-piperazinyl)methyl]phenol.

2. A compound according to claim 1 wherein:

each R¹ is independently halo, C₁₋₆ alkyl, C₁₋₆ haloalkyl, aryl or heteroaryl, wherein aryl or heteroaryl groups are optionally substituted with one or more substituents

selected from halo, OH, cyano, nitro, $-S(O)_pNR^4R^5$, $-C(O)NR^4R^5$, $-C_{1-6}$ alkyl optionally substituted with aryl, $-C_{1-6}$ haloalkyl, $-O(C_{1-6}$ alkyl) optionally substituted with aryl or $-O(C_{1-6}$ haloalkyl);

p is 0, 1 or 2;

5 each R^4 and R^5 is independently selected from H or C_{1-4} alkyl or R^4 and R^5 together with the nitrogen atom to which they are attached may form a 5- or 6-membered heterocyclic ring, optionally containing one or more additional heteroatoms selected from O, N and S;

10 R^6 is independently selected from aryl, $-O$ -aryl, $-O$ -heteroaryl or C_{1-4} alkyl substituted with one or more aryl or heteroaryl groups, wherein aryl and heteroaryl groups are substituted with one or more substituent selected from halo, C_{1-4} alkyl, C_{1-4} haloalkyl, $-O(C_{1-4}$ alkyl) or $-O(C_{1-4}$ haloalkyl);

or a pharmaceutically acceptable salt.

15 3. A compound according to claim 1 or claim 2, wherein n is 2 or 3.

4. A compound according to claim 2 or claim 3 wherein R^1 is selected from
(i) halo or an aryl or heteroaryl group optionally substituted according to claim 2; or

20 (ii) an aryl or heteroaryl group optionally substituted with halo, OH, cyano, nitro, $-SO_2NH_2$, $-C(O)NR^4R^5$, $-C_{1-4}$ alkyl optionally substituted with aryl, $-C_{1-4}$ haloalkyl, $-O(C_{1-4}$ alkyl) optionally substituted with aryl or $-O(C_{1-4}$ haloalkyl), wherein R^4 and R^5 together with the nitrogen atom to which they are attached form a piperidine or pyrrolidine ring; or

25 (iii) an aryl or heteroaryl group optionally substituted with chloro, fluoro, methyl, ethyl, trifluoromethyl, benzyl, methoxy, ethoxy, benzyloxy, trifluoromethoxy and piperidine-1-carbonyl.

30 5. A compound according to any one of claims 1 to 4, wherein R^6 is phenyl, $-O$ -phenyl, $-O$ -heteroaryl, benzyl, $-CH(phenyl)_2$, $-CH_2$ -heteroaryl and $-CH(heteroaryl)_2$, where the heteroaryl group is selected from pyridinyl, indolyl, isoindolyl, benzoxazolyl and benzisoxazolyl and wherein the above R^6 groups may be substituted as defined in claim 1.

35 6. A compound according to any one of claims 1-5, wherein z is 0, 1 or 2, R^7 not present (i.e. z is 0), or R^7 is halo, $-C_{1-4}$ alkyl, benzyl, $-O(C_{1-4}$ alkyl) benzyloxy, $-C_{1-4}$

haloalkyl, $-O(C_{1-4} \text{ haloalkyl})$ or $-C(O)NR^8R^9$, where R^8 and R^9 together with the nitrogen atom to which they are attached form a piperidiny ring or wherein R^8 is H and R^9 is C_{3-7} cycloalkyl.

- 5 7. A compound according to claim 1 selected from:
 2-((2-chlorophenyl)(4-(4-methoxyphenyl)piperazin-1-yl)methyl)phenol (Compound 1);
 2-((4-chlorophenyl)(4-(4-methoxyphenyl)piperazin-1-yl)methyl)phenol (Compound 2);
 2-((4-(benzyloxy)-3-fluorophenyl)(4-(4-methoxyphenyl)piperazin-1-yl)methyl)phenol
 (Compound 3);
 10 (4-((2-hydroxyphenyl)(4-(4-Methoxyphenyl)piperazinyl)methyl)phenyl)(piperidin-1-
 yl)methanone (Compound 4);
 3-((5-chloro-2-hydroxyphenyl)(4-(4-methoxyphenyl)piperazin-1-yl)methyl)-N-
 cyclopentylbenzamide (Compound 5);
 2-((4-(benzyloxy)-3-fluorophenyl)(4-(4-methoxyphenyl)piperazin-1-yl)methyl)-4-
 15 chlorophenol (Compound 6);
 2-((4-(2, 3-dichlorophenyl)piperazin-1-yl)(o-tolyl)methyl)phenol (Compound 8);
 N-cyclopentyl-3-((4-(2,3-dichlorophenyl)piperazin-1-yl)(2-hydroxyphenyl)methyl)
 benzamide (Compound 9, NPB);
 2-((4-(benzyloxy)-3-fluorophenyl)(4-(2,3-dichlorophenyl)piperazin-1-yl)methyl)phenol
 20 (Compound 10);
 2-((4-((4-chlorophenyl)(phenyl)methyl)piperazin-1-yl)(phenyl)methyl)phenol
 (Compound 11);
 2-((4-((4-chlorophenyl)(phenyl)methyl)piperazin-1-yl)(p-tolyl)methyl)phenol
 (Compound 12);
 25 2-((4-chlorophenyl)(4-((4-chlorophenyl)(phenyl)methyl)piperazin-1-yl)methyl)phenol
 (Compound 13);
 2-((4-((4-chlorophenyl)(phenyl)methyl)piperazin-1-yl)(4-ethylphenyl)methyl)phenol
 (Compound 14);
 (4-((4-((4-chlorophenyl)(phenyl)methyl)piperazin-1-yl)(2-
 30 hydroxyphenyl)methyl)phenyl) (piperidin-1-yl)methanone (Compound 15);
 N-cyclopentyl-3-((4-(2,3-dichlorophenyl)piperazin-1-yl)(4-hydroxy-[1,1'-biphenyl]-3-
 yl)methyl)benzamide (Compound 16);
 N-cyclopentyl-3-((4-(2,3-dichlorophenyl)piperazin-1-yl)(4-hydroxy-2'-methyl-[1,1'-
 biphenyl]-3-yl)methyl)benzamide (Compound 17);
 35 N-cyclopentyl-3-((4-(2,3-dichlorophenyl)piperazin-1-yl)(4-hydroxy-3'-methyl-[1,1'-
 biphenyl]-3-yl)methyl)benzamide (Compound 18);
 N-cyclopentyl-3-((4-(2,3-dichlorophenyl)piperazin-1-yl)(4-hydroxy-4'-methyl-[1,1'-

- biphenyl]-3-yl)methyl)benzamide (Compound 19);
3-((2'-chloro-4-hydroxy-[1,1'-biphenyl]-3-yl)(4-(2,3-dichlorophenyl)piperazin-1-yl)methyl)-N-cyclopentylbenzamide (Compound 20);
3-((3'-chloro-4-hydroxy-[1,1'-biphenyl]-3-yl)(4-(2,3-dichlorophenyl)piperazin-1-yl)methyl)-N-cyclopentylbenzamide (Compound 21);
5 3-((4'-chloro-4-hydroxy-[1,1'-biphenyl]-3-yl)(4-(2,3-dichlorophenyl)piperazin-1-yl)methyl)-N-cyclopentylbenzamide (Compound 22);
N-cyclopentyl-3-((4-(2,3-dichlorophenyl)piperazin-1-yl)(4'-ethyl-4-hydroxy-[1,1'-biphenyl]-3-yl)methyl)benzamide (Compound 23);
10 N-cyclopentyl-3-((4-(2,3-dichlorophenyl)piperazin-1-yl)(4-hydroxy-4'-(piperidine-1-carbonyl)-[1,1'-biphenyl]-3-yl)methyl)benzamide (Compound 24);
N-cyclopentyl-3-((4-(2,3-dichlorophenyl)piperazin-1-yl)(4-hydroxy-4'-methoxy-[1,1'-biphenyl]-3-yl)methyl)benzamide (Compound 25);
N-cyclopentyl-3-((4-(2,3-dichlorophenyl)piperazin-1-yl)(2'-ethyl-4-hydroxy-[1,1'-biphenyl]-3-yl)methyl)benzamide (Compound 26);
15 N-cyclopentyl-3-((4-(2,3-dichlorophenyl)piperazin-1-yl)(2'-fluoro-4-hydroxy-[1,1'-biphenyl]-3-yl)methyl)benzamide (Compound 27);
N-cyclopentyl-3-((4-(2,3-dichlorophenyl)piperazin-1-yl)(3'-fluoro-4-hydroxy-[1,1'-biphenyl]-3-yl)methyl)benzamide (Compound 28);
20 N-cyclopentyl-3-((4-(2,3-dichlorophenyl)piperazin-1-yl)(4'-fluoro-4-hydroxy-[1,1'-biphenyl]-3-yl)methyl)benzamide (Compound 29);
N-cyclopentyl-3-((4-(2,3-dichlorophenyl)piperazin-1-yl)(4-hydroxy-3'-nitro-[1,1'-biphenyl]-3-yl)methyl)benzamide (Compound 30);
N-cyclopentyl-3-((4-(2,3-dichlorophenyl)piperazin-1-yl)(4-hydroxy-3'-sulfamoyl-[1,1'-biphenyl]-3-yl)methyl)benzamide (Compound 31);
25 N-cyclopentyl-3-((4-(2,3-dichlorophenyl)piperazin-1-yl)(4-hydroxy-2'-(trifluoromethyl)-[1,1'-biphenyl]-3-yl)methyl)benzamide (Compound 32);
N-cyclopentyl-3-((4-(2,3-dichlorophenyl)piperazin-1-yl)(4-hydroxy-3'-(trifluoromethyl)-[1,1'-biphenyl]-3-yl)methyl)benzamide (Compound 33);
30 N-cyclopentyl-3-((4-(2,3-dichlorophenyl)piperazin-1-yl)(4-hydroxy-4'-(trifluoromethyl)-[1,1'-biphenyl]-3-yl)methyl)benzamide (Compound 34);
3-((2'-cyano-4-hydroxy-[1,1'-biphenyl]-3-yl)(4-(2,3-dichlorophenyl)piperazin-1-yl)methyl)-N-cyclopentylbenzamide (Compound 35);
3-((3'-cyano-4-hydroxy-[1,1'-biphenyl]-3-yl)(4-(2,3-dichlorophenyl)piperazin-1-yl)methyl)-N-cyclopentylbenzamide (Compound 36);
35 3-((4'-cyano-4-hydroxy-[1,1'-biphenyl]-3-yl)(4-(2,3-dichlorophenyl)piperazin-1-yl)methyl)-N-cyclopentylbenzamide (Compound 37);

- 3-((2'-chloro-4-hydroxy-4'-(trifluoromethyl)-[1,1'-biphenyl]-3-yl)(4-(2,3-dichlorophenyl) piperazin-1-yl)methyl)-N-cyclopentylbenzamide (Compound 38);
 N-cyclopentyl-3-((2',4'-dichloro-4-hydroxy-[1,1'-biphenyl]-3-yl)(4-(2,3-dichlorophenyl) piperazin-1-yl)methyl)benzamide (Compound 39);
- 5 3-((4'-chloro-2',4'-dihydroxy-[1,1'-biphenyl]-3-yl)(4-(2,3-dichlorophenyl)piperazin-1-yl)methyl)-N-cyclopentylbenzamide (Compound 40);
 3-((4-(4-chlorophenyl)piperazin-1-yl)(2-hydroxyphenyl)methyl)-N-cyclopentylbenzamide (Compound 41, NCK1)
- 10 2-((4-chlorophenyl)(4-(2,3-dichlorophenyl)piperazin-1-yl)methyl)phenol (Compound 42, NCK2)
 2-((4-(2,3-dichlorophenyl)piperazin-1-yl)(3-methoxyphenyl)methyl)phenol (Compound 43, NCK3)
- 15 1-(5-((4-(2,3-dichlorophenyl)piperazin-1-yl)(2-hydroxyphenyl)methyl)thiophen-2-yl)ethanone (Compound 44, NCK4)
 2-((4-(2,3-dichlorophenyl)piperazin-1-yl)(naphthalen-1-yl)methyl)phenol (Compound 45, NCK5)
- 20 5-((4-(2,3-dichlorophenyl)piperazin-1-yl)(2-hydroxyphenyl)methyl)furan-2-carbaldehyde (Compound 46, NCK6)
 2-((4-(5,6-dichlorocyclohexa-1,5-dien-1-yl)piperazin-1-yl)(2-fluoro-3-methylpyridin-4-yl)methyl)phenol (Compound 47, NCK7)
- 2-((4-(2,3-dichlorophenyl)piperazin-1-yl)(4-(trifluoromethyl)phenyl)methyl)phenol (Compound 48, NCK8)
 2-((6-chloro-5-methylpyridin-3-yl)(4-(2,3-dichlorophenyl)piperazin-1-yl)methyl)phenol (Compound 49, NCK9)
- 25 2-((4-(2,3-dichlorophenyl)piperazin-1-yl)(pyridin-3-yl)methyl)phenol (Compound 50, NCK10)
 1-(5-((4-(4-chlorophenyl)piperazin-1-yl)(2-hydroxyphenyl)methyl)thiophen-2-yl)ethanone (Compound 51, NCK14)
- 30 3-((4-(4-chlorophenyl)piperazin-1-yl)(4-(diethylamino)-2-hydroxyphenyl)methyl)-N-cyclopentylbenzamide(Compound 52, NCK16)
 N-cyclopentyl-3-((4-(2,3-dichlorophenyl)piperazin-1-yl)(4-(diethylamino)-2-hydroxyphenyl)methyl)benzamide (Compound 53, NCK18)
 N-cyclopentyl-3-((4-(2,3-dichlorophenyl)piperazin-1-yl)(2-hydroxy-4,6-dimethoxyphenyl)methyl)benzamide (Compound 54, NCK19)
- 35 2-((4-chlorophenyl)(4-(4-chlorophenyl)piperazin-1-yl)methyl)phenol (Compound 55, NCK20)
 2-((4-(4-chlorophenyl)piperazin-1-yl)(6-methylpyridin-3-yl)methyl)phenol (Compound

56, NCK21)

2-(o-tolyl(4-(p-tolyl)piperazin-1-yl)methyl)phenol (Compound 57, SG1)

2-((4-(p-tolyl)piperazin-1-yl)(4-(trifluoromethyl)phenyl)methyl)phenol (Compound 58, SG2)

5 N-cyclopentyl-4-((2-hydroxyphenyl)(4-(p-tolyl)piperazin-1-yl)methyl)benzamide (Compound 59, SG3)

2-((4-chlorophenyl)(4-(p-tolyl)piperazin-1-yl)methyl)phenol (Compound 60, SG4)

2-((3-methoxyphenyl)(4-(p-tolyl)piperazin-1-yl)methyl)phenol (Compound 61, SG5)

5-((2-hydroxyphenyl)(4-(p-tolyl)piperazin-1-yl)methyl)furan-2-carbaldehyde


10 (Compound 62, SG6)

2-((6-methylpyridin-3-yl)(4-(p-tolyl)piperazin-1-yl)methyl)phenol (Compound 63, SG7);

or a pharmaceutically acceptable salt.


15 8. A process for the preparation of a compound according to any one of claims 1 to 7 which comprises either:

i. reacting an aldehyde of general formula (II):




20 wherein R¹ and n are as defined in claim 1;

with a compound of general formula (III):



25 wherein R^{2a} and R^{2b} together with the nitrogen atom to which they are attached form a 6-membered heterocyclic ring containing one further nitrogen heteroatom and substituted with one substituent R⁶, wherein R⁶ is as defined in claim 1;


and a boronic acid of general formula (IV):



wherein R^3 is aryl, which is optionally substituted with one or more substituents R^7 as defined in claim 1; or

5

ii. reacting a compound of general formula (ID) in which R^1 is halo with a compound of general formula (V):



10 wherein R^{1a} is aryl or heteroaryl optionally substituted as defined for R^1 in claim 1; via a Suzuki coupling reaction in the presence of a palladium catalyst to obtain a compound of general formula (ID) in which R^1 is R^{1a} .

15 9. A compound according to any one of claims 1 to 7 to treat cancer, wherein the cancer is associated with Bcl-2-associated death promoter (BAD).

20 10. The use of a compound according to any one of claims 1 to 7 in the preparation of a medicament for the treatment of cancer, wherein the cancer is associated with Bcl-2-associated death promoter (BAD).

20

11. A method for the treatment of cancer, the method comprising administering to a patient in need of such treatment an effective amount of a compound according to any one of claims 1 to 7, wherein the cancer is associated with Bcl-2-associated death promoter (BAD).

25

12. A compound of claim 9, the use according to claim 10, or the method according to claim 11, wherein the cancer is a cancer in which there is BAD phosphorylation.

30 13. The compound, the use or the method according to claim 12, wherein the cancer is selected from the group comprising breast cancer, endometrial cancer,

ovarian cancer, liver cancer, colon cancer, prostate cancer or pancreatic cancer or any other epithelial derived cancer in which BAD is phosphorylated.

5 14. A pharmaceutical composition comprising a compound according to any one of claims 1 to 7 and a pharmaceutically acceptable excipient.

10 15. A pharmaceutical composition according to claim 14 formulated for intraperitoneal administration, hepatoportal administration, intravenous administration, intra articular administration, pancreatic duodenal artery administration or intramuscular administration, or any combination thereof.




Figure 1A




Figure 1B




Figure 2




Figure 3A




Figure 3B




Figure 4A




Figure 4B

¹H-NMR





Figure 5A



C-13 NMR




Figure 5B




Figure 6




Figure 7A





Figure 7B



C-13 NMR




Figure 8




Figure 9A




Figure 9B




Figure 10

Figure 11IC₅₀ values of NPB in a range of carcinoma cell lines


Panel of cell lines		NPB (µM)	
Tissue	Cells	IC ₅₀ ^A	±SD
Mammary	#MCF10A	NV	NV
	#MCF12A	NV	NV
	MCF7	6.5	1.06
	T47D	7.24	1.91
	BT474	5.31	2.04
	BT549	4.88	1.31
	MDA-MB-231	6.94	1.86
Endometrial	Ishikawa	7.51	2.08
	ECC1	2.61	0.97
	RL95-2	6.38	1.85
	AN3	11.37	2.61
Ovarian	SK-OV-3	7.34	2.03
	OVCAR-2	4.21	1.74
	Caov-3	3.95	0.93
	HEY C2	6.82	1.94
	Ovca433	9.79	2.48
Hepatocellular	#LO2	NV	NV
	Hep3B	6.94	1.07
	H2P	4.18	0.83
	H2M	5.57	2.61
Colon	HCT116	7.29	2.02
	DLD-1	2.46	0.91
	Caco-2	3.08	0.76
Prostate	PC3	3.77	1.51
	LNCaP	8.02	3.3
	DU145	6.99	1.72
Pancreatic	AsPC-1	3.83	1.14
	BxPC-3	7.25	2.09

Note: NV, No value; #, Normal cells;


^AIC₅₀ values calculated using GraphPad Prism software (Version 5.0). Cell viability measured Using AlamarBlue® cell viability assay.

Figure 12

A. Cell viability



B. Caspase 3/7 activity (Apoptosis)



C. Cytotoxicity






Figure 13:


A. Flow cytometry





B. Cell cycle analysis



C. 3D matrigel growth



D. Soft agar colony formation



E. Foci formation






Figure 14:


A. Surface Plasmon Resonance (SPR)



B. Western blot



C. Western blot



D. Western blot





Figure 15:

A. Western blot



B. Western blot






Figure 16:

A. Western blot



B. Cell viability



C. Caspase 3/7 activity






Figure 17


A. Xenograft assay



B. Tumour weight



C. Western blot








Figure 17D

Figure 18:

A. Western blot



B. Kinase array






Figure 19




Figure 20




Figure 21

Figure 22

A






Analogues	Structure	MCF7	A2780
		IC ₅₀	IC ₅₀
NPB	 <p>MW: 523</p>	9.52	5.33
NCK1	 <p>Molecular Weight: 490.0363</p>	NV	26.63
NCK2	 <p>Molecular Weight: 447.7846</p>	NV	19.74
NCK3	 <p>Molecular Weight: 443.3656</p>	NV	50.82
NCK4	 <p>Molecular Weight: 461.4040</p>	NV	NV

Figure 22

B






NCK5	 Molecular Weight: 463.3983	5.907	27.39
NCK6	 Molecular Weight: 431.3118	3.114	6.68
NCK7	 Molecular Weight: 448.3606	47.28	38.74
NCK8	 Molecular Weight: 481.338	59.52	23.66
NCK9	 Molecular Weight: 462.7993	NV	16.06

Figure 22

C






NCK10	 <p>Molecular Weight: 414.3276</p>	24.83	35.86
NCK14	 <p>Molecular Weight: 426.9589</p>	NV	23.26
NCK16	 <p>Molecular Weight: 561.16</p>	23.35	9.53
NCK18	 <p>Molecular Weight: 595.60</p>	20.93	2.97
NCK19	 <p>Molecular Weight: 584.53</p>	NV	48.02


Figure 22

D

NCK20	<p>Molecular Weight: 413.34</p>	253.8	8.395
NCK21	<p>Molecular Weight: 377.77</p>	20.91	21.90
SG2	<p>Molecular Weight: 426.4740</p>	155.6	89.69
SG1	<p>Molecular Weight: 372.5026</p>	NV	NV
SG3	<p>Molecular Weight: 469.6176</p>	56.47	20.69

Figure 22

E

SG7		49.87	41.61
SG 4, 5, and 6	NOT TESTED		