

(19) **DANMARK**

(10) **DK/EP 2970406 T3**



(12)

Oversættelse af europæisk patentskrift

Patent- og
Varemærkestyrelsen

-
- (51) Int.Cl.: **A 61 K 38/00 (2006.01)** **C 07 K 14/31 (2006.01)** **C 07 K 14/435 (2006.01)**
C 07 K 14/735 (2006.01) **C 07 K 14/74 (2006.01)** **C 07 K 14/745 (2006.01)**
- (45) Oversættelsen bekendtgjort den: **2018-02-05**
- (80) Dato for Den Europæiske Patentmyndigheds bekendtgørelse om meddelelse af patentet: **2017-12-13**
- (86) Europæisk ansøgning nr.: **14710879.9**
- (86) Europæisk indleveringsdag: **2014-03-17**
- (87) Den europæiske ansøgnings publiceringsdag: **2016-01-20**
- (86) International ansøgning nr.: **EP2014055299**
- (87) Internationalt publikationsnr.: **WO2014140366**
- (30) Prioritet: **2013-03-15 US 201361787305 P** **2013-03-15 EP 13159500**
- (84) Designerede stater: **AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO RS SE SI SK SM TR**
- (73) Patenthaver: **Affibody AB, Gunnar Asplunds Allé 24, 171 63 Solna, Sverige**
- (72) Opfinder: **Ekblad, Caroline, Huldrevägen 23, 132 39 Saltsjö-Boo, Sverige**
LINDBORG, Malin, Hindvägen 5, 132 48 Saltsjö-Boo, Sverige
ABRAHMSÉN, Lars, Vereliusvägen 17, 168 53 Bromma, Sverige
LÖFBLOM, John, Högmora Ringväg 29, 141 37 Huddinge, Sverige
Gräslund, Torbjörn, KorpMossevägen 36, 126 38 Hägersten, Sverige
SEIJSING, Johan, Lustgårdsgatan 5, 112 18 Stockholm, Sverige
GUNNERIUSSON, Elin, Älgövägen 21, 133 36 Saltsjöbaden, Sverige
- (74) Fuldmægtig i Danmark: **Awapatent A/S, Strandgade 56, 1401 København K, Danmark**
- (54) Benævnelse: **Nye polypeptider**
- (56) Fremdragne publikationer:
WO-A1-2010/054699
WO-A1-2011/056124
A. JONSSON ET AL: "Engineering of a femtomolar affinity binding protein to human serum albumin", PROTEIN ENGINEERING DESIGN AND SELECTION, vol. 21, no. 8, 1 January 2008 (2008-01-01), pages 515-527, XP055047773, ISSN: 1741-0126, DOI: 10.1093/protein/gzn028
J. T. ANDERSEN ET AL: "Extending Half-life by Indirect Targeting of the Neonatal Fc Receptor (FcRn) Using a Minimal Albumin Binding Domain", JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 286, no. 7, 18 February 2011 (2011-02-18), pages 5234-5241, XP055019819, ISSN: 0021-9258, DOI: 10.1074/jbc.M110.164848

DESCRIPTION

Technical field of the invention

[0001] The present disclosure relates to a class of engineered polypeptides having a binding affinity for the neonatal Fc receptor (in the following referred to as FcRn). The present disclosure also relates to the use of such an FcRn binding polypeptide as an agent for modifying pharmacokinetic and pharmacodynamic properties of a biomolecule, e.g. a pharmaceutical, and as a therapeutic agent.

Background

[0002] The neonatal Fc receptor (FcRn) is a heterodimeric protein consisting of a transmembrane MHC class I-like heavy chain (FcRn α -chain) and the β 2-microglobulin light chain, the latter also forming a part of MHC class I molecules (Simister and Mostov (1989) Nature 337:184-7; Burmeister et al. (1994) Nature 372:379-83).

[0003] FcRn is predominantly located in endosomes and is able to bind to serum albumin and immunoglobulin G (IgG) at (pH $\leq$ 6.5 and release them at pH $\geq$ 7.0 (reviewed in Roopenian (2007) Nat Rev Immunol 7:715-25).

[0004] FcRn carries out several distinct tasks in mammals (reviewed in Roopenian, *supra*). FcRn is involved in recycling of endocytosed IgG and serum albumin, thus avoiding their degradation in the lysosome, giving them longer half-life and higher availability in the blood than other serum proteins. When IgG, serum albumin and other serum proteins are passively pinocytosed by cells in contact with blood, the pH becomes gradually lower in the formed endosomes, which permits the binding of IgG and serum albumin to FcRn. The receptor is then, together with its bound ligand, transported via recycling endosomes back to the plasma membrane. After returning to the plasma membrane, the pH increases to above 7, at which point the bound ligand is released.

[0005] FcRn is also recognized for its ability to transport IgG over barriers such as the placenta, the upper airway epithelium, the blood-brain barrier and the proximal small intestine.

[0006] In mammals, the properties of FcRn are used to transcytose IgG from a mother to a fetus via the placenta, and to transcytose IgG from a mother's milk to the blood stream of an infant in the proximal small intestine.

[0007] The expression pattern of FcRn differs between species. However, FcRn is widely expressed by cells in the blood brain barrier, upper airway epithelium, kidneys and vascular endothelia, and by antigen presenting cells as well as by other cells of hematopoietic origin in

most species (reviewed in Roopenian (2007), *supra*).

[0008] Antibodies and peptides with affinity towards FcRn (Liu et al. (2007) J Immunol 179:2999-3011, Mezo et al. (2008) Proc Natl Acad Sci U S A 105:2337-42) and β 2-microglobulin (Getman and Balthasar (2005) J Pharm Sci 94:718-29) have been developed with a view to inhibit the binding between endogenous IgG and FcRn. Another approach has been to mutate the Fc region of the IgG to get a higher affinity for FcRn (Petkova et al. (2006) Int Immunol 18:1759-69, Vaccaro et al. (2005) Nat Biotechnol 23:1283-8).

[0009] Fusion to the Fc domain or to albumin is a widely used strategy to increase the *in vivo* half-life of proteins. However, the large size of such fusion proteins adversely affects tissue penetration and reduces the specificity to the fusion partner (Valles et al. (2011) J Interferon Cytokine Res 32:178-184). On the other hand, mutations have been made in the Fc fragment of antibodies administered to non human primates to prolong half-life (Hinton et al. (2004) J Biol Chem 279:6213-6). However, this approach is only limited in use to therapeutic antibodies, and cannot be extrapolated to other therapeutic proteins unless the proteins in question are fused to Fc fragments, which also results in large size molecules. A number of chemical and recombinant methods have been devised to improve protein half-life, such as PEGylation and genetic fusions of the protein to the Fc domain of IgG or albumin (reviewed in Schellenberger et al. (2009) Nat Biotechnol 21:1186-1190). PEGylation of proteins has been reported to decrease their potency and contribute to their immunoreactivity.

[0010] Fc-fusion proteins have also been used for oral and pulmonary delivery mediated by the FcRn (Low et al., (2005) Human reproduction Jul;20(7):1805-13), however similar problems relating to tissue penetration and reduced specificity remain, due to the size of the fusion molecules.

[0011] WO2010/054699 discloses a HER2 binding Z variant molecule, denoted SEQ ID NO:493.

[0012] Hence, there is large need in the field for the continued provision of molecules with high affinity for FcRn. In particular, small binding molecules are needed that, when present as a fusion partner, do not adversely affect the properties of the molecules they are fused to and do not contribute to immunoreactivity.

Summary of the invention

[0013] It is an object of the present disclosure to provide new FcRn binding agents for use in modifying pharmacokinetic and/or pharmacodynamic properties of a biomolecule, e.g. a pharmaceutical.

[0014] It is also an object of the present disclosure to provide new FcRn binding agents for use as therapeutic agents in their own right, alone or as combination treatment.

[0015] It is an object of the present disclosure to provide a molecule allowing for efficient targeting of FcRn, while alleviating the above-mentioned and other drawbacks of current therapies.

[0016] These and other objects which are evident to the skilled person from the present disclosure are met by different aspects of the invention as claimed in the appended claims and as generally disclosed herein.

[0017] Thus, in the first aspect of the disclosure, there is provided a neonatal Fc receptor (FcRn) binding polypeptide, comprising an FcRn binding motif, *BM*, which motif consists of the amino acid sequence

$$\text{EX}_2 \text{X}_3 \text{X}_4 \text{AX}_6 \text{X}_7 \text{EIR WLPNLX}_{16} \text{X}_{17} \text{X}_{18} \text{QR X}_{21} \text{AFIX}_{25} \text{X}_{26} \text{LX}_{28} \text{X}_{29}$$

wherein, independently from each other,

X₂ is selected from A, D, E, F, H, I, K, L, N, Q, R, S, T, V, W and Y;

X₃ is selected from A, D, E, F, G, H, I, K, L, M, N, Q, R, S, T, V, W and Y;

X₄ is selected from A, D, E, F, G, H, I, K, L, N, Q, R, S, T, V, W and Y;

X₆ is selected from A, D, E, F, G, H, I, K, L, N, Q, R, S, T, V, W and Y;

X₇ is selected from A, F, H, I, K, L, N, Q, R, S, T, V, W and Y;

X₁₆ is selected from N and T;

X₁₇ is selected from F, W and Y;

X₁₈ is selected from A, D, E and N;

X₂₁ is selected from A, S, V and W;

X₂₅ is selected from A, D, E, F, G, H, I, K, L, N, Q, R, S, T, V, W and Y;

X₂₆ is selected from K and S;

X₂₈ is selected from A, D, E, F, H, I, K, L, N, Q, R, S, T, V, W and Y; and

X₂₉ is selected from D and R.

[0018] The above definition of a class of sequence related, FcRn binding polypeptides is based on a statistical analysis of a number of random polypeptide variants of a parent scaffold,

that were selected for their interaction with FcRn in several different selection experiments. The identified FcRn binding motif, or "*BM*", corresponds to the target binding region of the parent scaffold, which region constitutes two alpha helices within a three-helical bundle protein domain. In the parent scaffold, the varied amino acid residues of the two *BM* helices constitute a binding surface for interaction with the constant Fc part of antibodies. In the present disclosure, the random variation of binding surface residues and subsequent selection of variants have replaced the Fc interaction capacity with a capacity for interaction with FcRn.

[0019] In one embodiment of said FcRn binding polypeptide, the *BM* consists of the amino acid sequence

$$\text{EX}_2 \text{X}_3 \text{X}_4 \text{AX}_6 \text{X}_7 \text{EIR WLPNLTX}_{17} \text{X}_{18} \text{QR X}_{21} \text{AFIX}_{25} \text{KLX}_{28} \text{D}$$

wherein, independently from each other,

X_2 is selected from A, D, E, F, H, I, K, L, N, Q, R, S, T, V, W and Y;

X_3 is selected from A, D, E, F, G, H, I, K, L, M, N, Q, R, S, T, V, W and Y;

X_4 is selected from A, D, E, F, G, H, I, K, L, N, Q, R, S, T, V, W and Y;

X_6 is selected from A, D, E, F, G, H, I, K, L, N, Q, R, S, T, V, W and Y;

X_7 is selected from A, F, H, I, K, L, N, Q, R, S, T, V, W and Y;

X_{17} is selected from F, W and Y;

X_{18} is selected from A, D, E and N;

X_{21} is selected from A, S, V and W;

X_{25} is selected from A, D, E, F, G, H, I, K, L, N, Q, R, S, T, V, W and Y; and

X_{28} is selected from A, D, E, F, H, I, K, L, N, Q, R, S, T, V, W and Y.

[0020] In another embodiment of the first aspect of the disclosure, said neonatal Fc receptor (FcRn) binding polypeptide comprises an FcRn binding motif, *BM*, which motif consists of the amino acid sequence

$$\text{EX}_2 \text{X}_3 \text{X}_4 \text{AX}_6 \text{X}_7 \text{EIR WLPNLX}_{16} \text{X}_{17} \text{X}_{18} \text{QR X}_{21} \text{AFIX}_{25} \text{X}_{26} \text{LX}_{28} \text{X}_{29}$$

wherein, independently from each other,

X_2 is selected from A, D, E, F, H, I, K, L, N, Q, R, S, T, V, W and Y;

X₃ is selected from A, D, E, F, G, H, I, K, L, M, N, Q, R, S, T, V, W and Y;

X₄ is selected from A, D, E, F, G, H, I, K, L, N, Q, R, S, T, V, W and Y;

X₆ is selected from A, E, F, G, H, I, K, Q, R, S and V;

X₇ is selected from A, F, H, K, N, Q, R, S and V;

X₁₆ is selected from N and T;

X₁₇ is selected from F, W and Y;

X₁₈ is selected from A, D, E and N;

X₂₁ is selected from A, S, V and W;

X₂₅ is selected from D, E, G, H, I, K, L, N, Q, R, S, T, V, W and Y;

X₂₆ is selected from K and S;

X₂₈ is selected from A, D, E, F, H, I, K, L, N, Q, R, S, T, V, W and Y; and

X₂₉ is selected from D and R.

[0021] In another embodiment of the first aspect, there is provided an FcRn binding polypeptide, wherein, independently from each other,

X₂ is selected from A, D, E, F, H, I, K, L, N, Q, R, S, T, V, W and Y;

X₃ is selected from A, D, E, F, H, I, K, L, M, N, Q, R, S, T, V, W and Y;

X₄ is selected from A, D, E, F, H, I, K, L, N, Q, R, S, T, V, W and Y;

X₆ is selected from A, E, F, G, H, I, K, Q, R and S;

X₇ is selected from A, F, H, K, N, Q, R, S and V;

X₁₆ is selected from N and T;

X₁₇ is selected from F and Y;

X₁₈ is D;

X₂₁ is V;

X₂₅ is selected from D, E, H, I, K, L, N, Q, R, S, T, V, W and Y;

X₂₆ is selected from K and S;

X₂₈ is selected from A, D, E, F, H, I, K, L, N, Q, R, S, T, V and W and.

X₂₉ is selected from D and R.

[0022] In another embodiment of the first aspect, the *BM* consists of an amino acid sequence selected from

1. i)

EX₂ X₃ X₄ AX₆ HEIR WLPNLTX₁₇ X₁₈ QR X₂₁ AFIX₂₅ KLX₂₈ D

wherein, independently from each other,

X₂ is selected from A, D, E, F, H, I, K, L, N, Q, R, S, T, V, W and Y;

X₃ is selected from A, D, E, G, H, K, L, M, N, Q, R, S, T, V and Y;

X₄ is selected from A, D, E, F, G, I, K, L, N, Q, R, S, T, V and Y;

X₆ is selected from A, G, K, R, S and V;

X₁₇ is selected from F, W and Y;

X₁₈ is selected from A, D, E and N;

X₂₁ is selected from A, S, V and W;

X₂₅ is selected from D, G, H, K, L, N, R, V and W;

X₂₈ is selected from A, D, E, H, K, L, N, Q, R, S, T, W and Y; and

2. ii) an amino acid sequence which has at least 96 % identity to said sequence.

[0023] In yet another embodiment of said aspect, the *BM* in sequence i) consists of an amino acid sequence selected from

EX₂ X₃ X₄ AX₆ HEIR WLPNLTX₁₇ X₁₈ QR X₂₁ AFIX₂₅ KLX₂₈ D

wherein, independently from each other,

X₂ is selected from A, D, E, F, N, Q, R, S and W;

X₃ is selected from D, E, G, H, K, M, N, Q, S and T;

X₄ is selected from A, D, E, G, N, Q, R, S, T, V and Y;

X₆ is selected from A, G, S and V;

X₁₇ is selected from F, W and Y;

X₁₈ is selected from A, D, E and N;

X₂₁ is selected from A, S, V and W;

X₂₅ is selected from D, G, H, K, L, N, R and V; and

X₂₈ is selected from A, E, H, L, N, Q, R, S, T, W and Y.

[0024] As the skilled person will realize, the function of any polypeptide, including the FcRn binding capacity of the polypeptide of the present disclosure, is dependent on the tertiary structure of the polypeptide. It is therefore possible to make minor changes to the sequence of amino acids in a polypeptide without affecting the function thereof. Thus, the disclosure encompasses modified variants of the FcRn binding polypeptide, which are such that the FcRn binding characteristics are retained.

[0025] Therefore, as described above, also encompassed by the present disclosure is a FcRn binding polypeptide comprising an amino acid sequence with 96 % or greater identity to a polypeptide as defined in i).

[0026] In some embodiments, such changes may be made in all positions of the sequences of the FcRn binding polypeptide as disclosed herein. In other embodiments, such changes may be made only in the non-variable positions, also denoted as scaffold amino acid residues. In such cases, changes are not allowed in the variable positions, i.e. positions denoted with an "X" in sequence i). For example, it is possible that an amino acid residue belonging to a certain functional grouping of amino acid residues (e.g. hydrophobic, hydrophilic, polar etc) could be exchanged for another amino acid residue from the same functional group.

[0027] The term "% identity", as used throughout the specification, may for example be calculated as follows. The query sequence is aligned to the target sequence using the CLUSTAL W algorithm (Thompson et al. (1994) Nucleic Acids Research 22:4673-4680). A comparison is made over the window corresponding to the shortest of the aligned sequences. The shortest of the aligned sequences may in some instances be the target sequence. In other instances, the query sequence may constitute the shortest of the aligned sequences. The amino acid residues at each position are compared, and the percentage of positions in the query sequence that have identical correspondences in the target sequence is reported as % identity.

[0028] Below follows a list of embodiments which further specify amino acid residue X_n, wherein n is an integer which denotes the position of said residue within the polypeptide described herein. To clarify, in cases where the BM comprised in the polypeptide may consist of either a given amino acid sequence or an amino acid sequence with at least a given % identity to said given amino acid sequence, the X_n as used herein refers to an amino acid residue in said given amino acid sequence. For example, when applicable, X_n refers to an amino acid residue in sequence i) above.

- [0029]** In one embodiment, X_2 is selected from A, D, E, F, I, L, N, Q, R, S, T, V, W and Y.
- [0030]** In one embodiment, X_2 is selected from A, D, F, I, L, N, Q, R, S, T, V, W and Y.
- [0031]** In one embodiment, X_2 is selected from A, D, F, I, L, N, Q, R, S, V and W.
- [0032]** In one embodiment, X_2 is selected from A, I, L, N, Q, R, S, T, V, W and Y.
- [0033]** In one embodiment, X_2 is selected from A, I, L, N, Q, S, T, V and W.
- [0034]** In one embodiment, X_2 is selected from A, I, L, N, Q, V and W.
- [0035]** In one embodiment, X_2 is selected from A, I, L, Q, V and W.
- [0036]** In one embodiment, X_2 is selected from A, I, L and Q.
- [0037]** In one embodiment, X_2 is selected from I, L and Q.
- [0038]** In one embodiment, X_2 is selected from I and Q.
- [0039]** In one embodiment, X_2 is I.
- [0040]** In one embodiment, X_2 is Q.
- [0041]** In one embodiment, X_3 is selected from A, D, E, G, H, K, L, M, N, Q, R, S, T, V and Y.
- [0042]** In one embodiment, X_3 is selected from A, D, E, H, K, L, M, N, Q, R, S, T, V and Y.
- [0043]** In one embodiment, X_3 is selected from A, D, E, G, H, K, L, M, N, Q, R, S and T.
- [0044]** In one embodiment, X_3 is selected from A, D, E, G, H, K, M, N, Q, S and T.
- [0045]** In one embodiment, X_3 is selected from A, D, E, G, H, M, N, Q, S and T.
- [0046]** In one embodiment, X_3 is selected from A, D, E, K, N, Q, S and T.
- [0047]** In one embodiment, X_3 is selected from A, D, E, K, Q, and T.
- [0048]** In one embodiment, X_3 is selected from A, D, E, Q and T.
- [0049]** In one embodiment, X_3 is selected from D, E and T.

[0050] In one embodiment, X_3 is selected from D and E.

[0051] In one embodiment, X_3 is D.

[0052] In one embodiment, X_3 is E.

[0053] In one embodiment, X_4 is selected from A, D, E, F, G, I, K, L, N, Q, R, S, T, V and Y.

[0054] In one embodiment, X_4 is selected from A, D, E, G, N, Q, R, S, T and V.

[0055] In one embodiment, X_4 is selected from A, D, E, F, I, K, L, N, Q, R, S, T and V.

[0056] In one embodiment, X_4 is selected from A, D, E, I, K, N, Q, R, S and T.

[0057] In one embodiment, X_4 is selected from A, D, E, I, K, Q, S and T.

[0058] In one embodiment, X_4 is selected from A, D, I, K, Q and S.

[0059] In one embodiment, X_4 is selected from A, D, E, K and S.

[0060] In one embodiment, X_4 is selected from A, D, K and S.

[0061] In one embodiment, X_4 is selected from A, D, E and K.

[0062] In one embodiment, X_4 is selected from A, D and K.

[0063] In one embodiment, X_4 is selected from A and D.

[0064] In one embodiment, X_4 is selected from A and E.

[0065] In one embodiment, X_4 is A.

[0066] In one embodiment, X_4 is D.

[0067] In one embodiment, X_4 is E.

[0068] In one embodiment, X_6 is selected from A, G, K, Q, R, S and V.

[0069] In one embodiment, X_6 is selected from A, G, K, R, S and V.

[0070] In one embodiment, X_6 is selected from A, G, K, R and S.

[0071] In one embodiment, X_6 is selected from A, G, K, S and V.

[0072] In one embodiment, X_6 is selected from A, G, K and V.

[0073] In one embodiment, X_6 is selected from A, G, K and S.

[0074] In one embodiment, X_6 is selected from A, G and K.

[0075] In one embodiment, X_6 is selected from A, G and V.

[0076] In one embodiment, X_6 is selected from A and G.

[0077] In one embodiment, X_6 is A.

[0078] In one embodiment, X_6 is G.

[0079] In one embodiment, X_7 is selected from A and H.

[0080] In one embodiment, X_7 is H.

[0081] In one embodiment, X_{16} is T.

[0082] In one embodiment, X_{16} is N.

[0083] In one embodiment, X_{17} is selected from F and Y.

[0084] In one embodiment, X_{17} is F.

[0085] In one embodiment, X_{18} is selected from A, D and E.

[0086] In one embodiment, X_{18} is selected from A and D.

[0087] In one embodiment, X_{18} is D.

[0088] In one embodiment, X_{21} is selected from V and W.

[0089] In one embodiment, X_{21} is V.

[0090] In one embodiment, X_{25} is selected from D, E, G, H, K, L, N, Q, R, V and W.

[0091] In one embodiment, X_{25} is selected from D, G, H, K, L, N, R, V and W.

[0092] In one embodiment, X_{25} is selected from D, G, H, K, L, N, R and V.

[0093] In one embodiment, X_{25} is selected from H, L, R, V and W.

[0094] In one embodiment, X_{25} is selected from H, R, V and W.

[0095] In one embodiment, X_{25} is selected from H, R and V.

[0096] In one embodiment, X_{25} is selected from H, L and R.

[0097] In one embodiment, X_{25} is selected from H and R.

[0098] In one embodiment, X_{25} is selected from H and V.

[0099] In one embodiment, X_{25} is H.

[0100] In one embodiment, X_{26} is K.

[0101] In one embodiment, X_{26} is S.

[0102] In one embodiment, X_{28} is selected from A, D, E, H, K, L, N, Q, R, S, T, W and Y.

[0103] In one embodiment, X_{28} is selected from A, D, E, K, L, N, Q, R, S, T, W and Y.

[0104] In one embodiment, X_{28} is selected from A, D, E, L, R, S, T, W and Y.

[0105] In one embodiment, X_{28} is selected from A, D, K, L, N, Q, R, S, T and W.

[0106] In one embodiment, X_{28} is selected from A, D and R.

[0107] In one embodiment, X_{28} is selected from A and R.

[0108] In one embodiment, X_{28} is selected from D and R.

[0109] In one embodiment, X_{28} is A.

[0110] In one embodiment, X_{28} is R.

[0111] In one embodiment, X_{29} is D.

[0112] In one embodiment, X_{29} is R.

[0113] In one embodiment, X_6X_7 is selected from AH and GH.

[0114] In one embodiment, X_6X_7 is AH.

[0115] In one embodiment, X_6X_7 is GH.

[0116] In one embodiment, $X_{17}X_{18}$ is selected from FD and YD.

[0117] In one embodiment, $X_{17}X_{18}$ is FD.

[0118] In a more specific embodiment defining a sub-class of the FcRn binding polypeptide, the sequence fulfills at least three of the six conditions I-VI:

1. I. X_6 is selected from A, G, K and S, such as in particular A;
2. II. X_7 is H;
3. III. X_{17} is selected from F and Y, such as in particular F;
4. IV. X_{18} is D;
5. V. X_{21} is selected from V and W, such as in particular V;
6. VI. X_{25} is selected from H and R, such as in particular H.

[0119] In some examples of an FcRn binding polypeptide according to the first aspect, said sequence fulfills at least four of the six conditions I-VI. More specifically, the sequence may fulfill at least five of the six conditions I-VI, such as all of the six conditions I-VI.

[0120] As described in detail in the experimental section to follow, the selection of FcRn binding polypeptide variants has led to the identification of a number of individual FcRn binding motif (*BM*) sequences. These sequences constitute individual embodiments according to this aspect. The sequences of individual FcRn binding motifs are presented in Figure 1 and as SEQ ID NO:1-353. Hence, in one embodiment of the FcRn binding polypeptide according to this aspect, the sequence is selected from the group consisting of SEQ ID NO:1-353. In one embodiment, the sequence is selected from the group consisting of SEQ ID NO:1-15, SEQ ID NO:17-140 and SEQ ID NO:353. In one embodiment, the sequence is selected from the group consisting of SEQ ID NO:1-2 and SEQ ID NO:17-140. In one embodiment, the sequence is selected from the group consisting of SEQ ID NO:1-2, SEQ ID NO:17-92, SEQ ID NO:94-103, SEQ ID NO:105-125 and SEQ ID NO:127-140. In one embodiment, the sequence is selected from the group consisting of SEQ ID NO:1-8, SEQ ID NO:13, SEQ ID NO:19-20, SEQ ID NO:23, SEQ ID NO:28, SEQ ID NO:41, SEQ ID NO:44, SEQ ID NO:65, SEQ ID NO:70, SEQ ID NO:73,

SEQ ID NO:75-77 and SEQ ID NO:353. In another embodiment, the sequence is selected from the group consisting of SEQ ID NO:1, SEQ ID NO:23, SEQ ID NO:28, SEQ ID NO:41, SEQ ID NO:44, SEQ ID NO:65, SEQ ID NO:73 and SEQ ID NO:75-77. In yet another embodiment, the sequence is selected from SEQ ID NO:1, SEQ ID NO:23, SEQ ID NO:44, SEQ ID NO:65, SEQ ID NO:75 and SEQ ID NO:77. In one embodiment, the sequence is selected from SEQ ID NO:1, SEQ ID NO:23 and SEQ ID NO:75. In one embodiment, the sequence is SEQ ID NO:1.

[0121] In some embodiments of the present disclosure, the *BM* as defined above "forms part of" a three-helix bundle protein domain. This is understood to mean that the sequence of the *BM* is "inserted" into or "grafted" onto the sequence of the original three-helix bundle domain, such that the *BM* replaces a similar structural motif in the original domain. For example, without wishing to be bound by theory, the *BM* is thought to constitute two of the three helices of a three-helix bundle, and can therefore replace such a two-helix motif within any three-helix bundle. As the skilled person will realize, the replacement of two helices of the three-helix bundle domain by the two *BM* helices has to be performed so as not to affect the basic structure of the polypeptide. That is, the overall folding of the C α backbone of the polypeptide according to this embodiment of the invention is substantially the same as that of the three-helix bundle protein domain of which it forms a part, e.g. having the same elements of secondary structure in the same order etc. Thus, a *BM* according to the disclosure "forms part" of a three-helix bundle domain if the polypeptide according to this embodiment of the aspect has the same fold as the original domain, implying that the basic structural properties are shared, those properties e.g. resulting in similar CD spectra. The skilled person is aware of other parameters that are relevant.

[0122] In particular embodiments, the FcRn binding motif (*BM*) thus forms part of a three-helix bundle protein domain. For example, the *BM* may essentially constitute two alpha helices with an interconnecting loop, within said three-helix bundle protein domain. In particular embodiments, said three-helix bundle protein domain is selected from domains of bacterial receptor proteins. Non-limiting examples of such domains are the five different three-helical domains of Protein A from *Staphylococcus aureus*, such as domain B, and derivatives thereof. In some embodiments, the three-helical bundle protein domain is a variant of protein Z, which is derived from domain B of staphylococcal Protein A.

[0123] In embodiments where the FcRn binding polypeptide of the invention forms part of a three-helix bundle protein domain, the FcRn binding polypeptide may comprise an amino acid sequence selected from:

iii)

$$K-[BM]-DPSQS X_a X_b LLX_c EAKKL X_d X_e X_f Q;$$

wherein

[*BM*] is an FcRn binding motif as defined herein, provided that X₂₉ is D;

X_a is selected from A and S;

X_b is selected from N and E;

X_c is selected from A, S and C;

X_d is selected from E, N and S;

X_e is selected from D, E and S;

X_f is selected from A and S;

and

iv) an amino acid sequence which has at least 93 % identity to a sequence defined by iii).

[0124] In embodiments where the FcRn binding polypeptide of the invention forms part of a three-helix bundle protein domain, the FcRn binding polypeptide may comprise an amino acid sequence selected from:

v)

K-[BM]-QPEQS X_aX_bLLX_c EAKKL X_dX_eX_fQ;

wherein

[BM] is an FcRn binding motif as defined herein, provided that X₂₉ is R;

X_a is selected from A and S;

X_b is selected from N and E;

X_c is selected from A, S and C;

X_d is selected from E, N and S;

X_e is selected from D, E and S;

X_f is selected from A and S;

and

vi) an amino acid sequence which has at least 93 % identity to a sequence defined by v).

[0125] As discussed above, polypeptides comprising minor changes as compared to the above amino acid sequences which do not largely affect the tertiary structure and the function thereof are also within the scope of the present disclosure. Thus, in some embodiments, sequence iv) or sequence vi) has at least 95 %, for example at least 97 % identity to a sequence defined by iii) and v), respectively.

[0126] In one embodiment, X_a in sequence iii) or v) is A. In an alternative embodiment, X_a in sequence iii) or v) is S.

[0127] In one embodiment, X_b in sequence iii) or v) is N. In an alternative embodiment, X_b in sequence iii) or v) is E.

[0128] In one embodiment, X_c in sequence iii) or v) is A. In an alternative embodiment, X_c in sequence iii) or v) is S. In yet another alternative embodiment, X_c in sequence iii) or v) is C.

[0129] In one embodiment, X_d in sequence iii) or v) is E.

[0130] In one embodiment, X_d in sequence iii) or v) is N.

[0131] In one embodiment, X_d in sequence iii) or v) is S.

[0132] In one embodiment, X_e in sequence iii) or v) is D.

[0133] In one embodiment, X_e in sequence iii) or v) is E.

[0134] In one embodiment, X_e in sequence iii) or v) is S.

[0135] In one embodiment, X_dX_e in sequence iii) or v) is selected from EE, ES, SE and SS.

[0136] In one embodiment, X_dX_e in sequence iii) or v) is ES.

[0137] In one embodiment, X_dX_e in sequence iii) or v) is SE.

[0138] In one embodiment, X_f in sequence iii) or v) is A. In an alternative embodiment, X_f in sequence iii) or v) is S.

[0139] In one embodiment, in sequence iii) or v), X_a is A; X_b is N; X_c is A and X_f is A.

[0140] In one embodiment, in sequence iii) or v), X_a is A; X_b is N; X_c is C and X_f is A.

[0141] In one embodiment, in sequence iii) or v), X_a is S; X_b is E; X_c is S and X_f is S.

[0142] In one embodiment, in sequence iii) or v), X_a is S; X_b is E; X_c is C and X_f is S.

[0143] In one embodiment, in sequence iii) or v), X_a is A; X_b is N; X_c is A; X_dX_e is ND and X_f is A.

[0144] In one embodiment, in sequence iii) or v), X_a is A; X_b is N; X_c is C; X_dX_e is ND and X_f is A.

[0145] In one embodiment, in sequence iii) or v), X_a is S; X_b is E; X_c is S; X_dX_e is ND and X_f is S.

[0146] In one embodiment, in sequence iii) or v), X_a is S; X_b is E; X_c is C; X_dX_e is ND and X_f is S.

[0147] In one embodiment, in sequence iii) or v), X_a is A; X_b is N; X_c is A; X_dX_e is SE and X_f is A.

[0148] In one embodiment, in sequence iii) or v), X_a is A; X_b is N; X_c is C; X_dX_e is SE and X_f is A.

[0149] In one embodiment, in sequence iii) or v), X_a is S; X_b is E; X_c is S; X_dX_e is SE and X_f is S.

[0150] In one embodiment, in sequence iii) or v), X_a is S; X_b is E; X_c is C; X_dX_e is SE and X_f is S.

[0151] In one embodiment, in sequence iii) or v), X_a is A; X_b is N; X_c is A; X_dX_e is ES and X_f is A.

[0152] In one embodiment, in sequence iii) or v), X_a is A; X_b is N; X_c is C; X_dX_e is ES and X_f is A.

[0153] In one embodiment, in sequence iii) or v), X_a is S; X_b is E; X_c is S; X_dX_e is ES and X_f is S.

[0154] In one embodiment, in sequence iii) or v), X_a is S; X_b is E; X_c is C; X_dX_e is ES and X_f is S.

[0155] In yet a further embodiment, sequence iii) in the definition of FcRn binding polypeptides above is selected from the group consisting of SEQ ID NO:354-706. In one embodiment, sequence iii) is selected from the group consisting of SEQ ID NO:354-368, SEQ ID NO:370-493 and SEQ ID NO:706. In one embodiment, sequence iii) is selected from the

group consisting of SEQ ID NO:354-355 and SEQ ID NO:370-493. In one embodiment, sequence iii) is selected from the group consisting of SEQ ID NO:354-355, SEQ ID NO:370-445, SEQ ID NO:447-456, SEQ ID NO:458-478 and SEQ ID NO:480-493. In one embodiment, sequence iii) is selected from the group consisting of SEQ ID NO:354-361, SEQ ID NO:366, SEQ ID NO:372-373, SEQ ID NO:376, SEQ ID NO:381, SEQ ID NO:394, SEQ ID NO:397, SEQ ID NO:418, SEQ ID NO:423, SEQ ID NO:426, SEQ ID NO:428-430 and SEQ ID NO:706. In another embodiment, sequence iii) is selected from the group consisting of SEQ ID NO:354, SEQ ID NO:376, SEQ ID NO:381, SEQ ID NO:394, SEQ ID NO:397, SEQ ID NO:418, SEQ ID NO:426 and SEQ ID NO:428-430. In yet another embodiment, sequence iii) is selected from SEQ ID NO:354, SEQ ID NO:376, SEQ ID NO:397, SEQ ID NO:418, SEQ ID NO:428 and SEQ ID NO:430. In one embodiment, sequence iii) is selected from SEQ ID NO:354, SEQ ID NO:376 and SEQ ID NO:428. In one embodiment, sequence iii) is SEQ ID NO:354.

[0156] Also, in a further embodiment, there is provided an FcRn binding polypeptide as defined above, which comprises an amino acid sequence selected from:

vii)

YAK-[BM]-DPSQS SELLX_C EAKKL NDSQA P;

wherein [BM] is an FcRn binding motif as defined above and X_C is selected from A, S and C; and

viii) an amino acid sequence which has at least 94 % identity to a sequence defined by vii).

[0157] Alternatively, there is provided an FcRn binding polypeptide as defined above, which comprises an amino acid sequence selected from:

ix)

FNK-[BM]-DPSQS ANLLX_C EAKKL NDAQA P;

wherein [BM] is an FcRn binding motif as defined above and X_C is selected from A and C; and

x) an amino acid sequence which has at least 94 % identity to a sequence defined by ix).

[0158] As discussed above, polypeptides comprising minor changes as compared to the above amino acid sequences that do not largely affect the tertiary structure and the function thereof are also within the scope of the present disclosure. Thus, in some embodiments, the FcRn binding polypeptide as defined above may comprise a sequence which is at least 96 %, such as at least 98 % identical to a sequence defined by vii) or ix).

[0159] In some embodiments, the FcRn binding motif may form part of a polypeptide comprising an amino acid sequence selected from

ADNNFNK-[BM]-DPSQSANLLSEAKKLNESQAPK;

ADNKFNK-[BM]-DPSQSANLLAEAKKLNDQAQAPK;

ADNKFNK-[BM]-DPSVSKEILAEAKKLNDQAQAPK;

ADAQQNNFNK-[BM]-DPSQSTNVLGEAKKLNESQAPK;

AQHDE-[BM]-DPSQSANVLGEAQKLNDQAPK;

VDNKFNK-[BM]-DPSQSANLLAEAKKLNDQAQAPK;

AEAKYAK-[BM]-DPSESSELLSEAKKLNKSQAPK;

VDAKYAK-[BM]-DPSQSSELLAEAKKLNDQAQAPK;

VDAKYAK-[BM]-DPSQSSELLAEAKKLNDQAPK;

AEAKYAK-[BM]-DPSQSSELLSEAKKLNDQAPK;

AEAKYAK-[BM]-DPSQSSELLSEAKKLSQAPK

AEAKYAK-[BM]-DPSQSSELLSEAKKLESSQAPK

VDAKYAK-[BM]-DPSQSSELLSEAKKLNDQAPK;

VDAKYAK-[BM]-DPSQSSELLSEAKKLSQAPK;

VDAKYAK-[BM]-DPSQSSELLSEAKKLESSQAPK;

VDAKYAK-[BM]-DPSQSSELLAEAKKLNKAQAPK; and

AEAKYAK-[BM]-DPSQSSELLAEAKKLNKAQAPK;

wherein [BM] is an FcRn binding motif as defined above.

[0160] In one embodiment, the FcRn binding polypeptide comprises an amino acid sequence selected from:

xi)

AEAKYAK-[BM]-DPSQSSELLSEAKKLNDQAPK;

wherein [BM] is an FcRn binding motif as defined above; and

xii) an amino acid sequence which has at least 94 % identity to the sequence defined in xi).

[0161] In one embodiment, sequence xi) is selected from the group consisting of SEQ ID NO:1060-1062.

[0162] In one embodiment, the FcRn binding polypeptide comprises an amino acid sequence selected from:

xiii)

VDAKYAK-[BM]-DPSQSSELLSEAKKLNDSQAPK;

wherein [BM] is an FcRn binding motif as defined above; and

xiv) an amino acid sequence which has at least 94 % identity to the sequence defined in xiii).

[0163] Sequence xiii) in such a polypeptide may for example be selected from the group consisting of SEQ ID NO:707-1059. In one embodiment, sequence xiii) is selected from the group consisting of SEQ ID NO:707-721, SEQ ID NO:723-846 and SEQ ID NO:1059. In one embodiment, sequence xiii) is selected from the group consisting of SEQ ID NO:707-708 and SEQ ID NO:723-846. In one embodiment, sequence xiii) is selected from the group consisting of SEQ ID NO:707-708, SEQ ID NO:723-798, SEQ ID NO:800-809, SEQ ID NO:811-831 and SEQ ID NO:833-846. In one embodiment, sequence xiii) is selected from the group consisting of SEQ ID NO:707-714, SEQ ID NO:719, SEQ ID NO:725-726, SEQ ID NO:729, SEQ ID NO:734, SEQ ID NO:747, SEQ ID NO:750, SEQ ID NO:771, SEQ ID NO:776, SEQ ID NO:779, SEQ ID NO:781-783 and SEQ ID NO:1059. In another embodiment, sequence xiii) is selected from the group consisting of SEQ ID NO:707, SEQ ID NO:729, SEQ ID NO:734, SEQ ID NO:747, SEQ ID NO:750, SEQ ID NO:771, SEQ ID NO:779 and SEQ ID NO:781-783. In yet another embodiment, sequence xiii) is selected from SEQ ID NO:707, SEQ ID NO:729, SEQ ID NO:750, SEQ ID NO:771, SEQ ID NO:781 and SEQ ID NO:783. In one embodiment, sequence xiii) is selected from SEQ ID NO:707, SEQ ID NO:729 and SEQ ID NO:781. In one embodiment, sequence xiii) is SEQ ID NO:707.

[0164] Again, polypeptides comprising minor changes as compared to the above amino acid sequences which do not largely affect the tertiary structure and the function thereof are also within the scope of the present disclosure. Thus, in some embodiments, the FcRn binding polypeptide as defined above may comprise a sequence which is at least 96 %, such as at least 98 % identical to a sequence defined by xi) or xiii).

[0165] The terms "FcRn binding" and "binding affinity for FcRn" as used in this specification refer to a property of a polypeptide which may be tested for example by the use of surface plasmon resonance (SPR) technology or ELISA.

[0166] For example as described in the examples below, FcRn binding affinity may be tested

in an experiment in which FcRn, or a correctly folded fragment thereof, is immobilized on a sensor chip of the instrument, and the sample containing the polypeptide to be tested is passed over the chip. Alternatively, the polypeptide to be tested is immobilized on a sensor chip of the instrument, and a sample containing FcRn, or a correctly folded fragment thereof, is passed over the chip. The skilled person may then interpret the results obtained by such experiments to establish at least a qualitative measure of the binding affinity of the polypeptide for FcRn. If a quantitative measure is desired, for example to determine a K_D value for the interaction, surface plasmon resonance methods may also be used. Binding values may for example be defined in a Biacore (GE Healthcare) or ProteOn XPR 36 (Bio-Rad) instrument. FcRn is suitably immobilized on a sensor chip of the instrument, and samples of the polypeptide whose affinity is to be determined are prepared by serial dilution and injected in random order. K_D values may then be calculated from the results using for example the 1:1 Langmuir binding model of the BIAevaluation 4.1 software, or other suitable software, provided by the instrument manufacturer.

[0167] Alternatively, as described in the examples below, FcRn binding affinity may be tested in an experiment in which samples of the polypeptide are captured on antibody coated ELISA plates, and biotinylated FcRn is added followed by streptavidin conjugated HRP. TMB substrate is added and the absorbance at 450 nm is measured using a multi-well plate reader, such as Victor³ (Perkin Elmer). The skilled person may then interpret the results obtained by such experiments to establish at least a qualitative measure of the binding affinity of the polypeptide for FcRn. If a quantitative measure is desired, for example to determine the K_D value (the half maximal effective concentration) for the interaction, ELISA may also be used. The response of the polypeptides against a dilution series of biotinylated FcRn are measured using ELISA as described above. The skilled person may then interpret the results obtained by such experiments and K_D values may be calculated from the results using for example GraphPad Prism 5 and non-linear regression.

[0168] In one embodiment, there is provided an FcRn binding polypeptide, which is capable of binding to FcRn at pH 6.0 such that the K_D value of the interaction is at most 1×10^{-6} M, such as at most 1×10^{-7} M, such as at most 1×10^{-8} M, such as at most 1×10^{-9} M, such as at most 1×10^{-10} M. An FcRn binding polypeptide according to this embodiment would bind, or remain bound, to FcRn in acidic pH conditions, such as pH 6.0, for example in a lysosome. If such a polypeptide were to enter an increasingly acidic intracellular environment, it would be recycled to the plasma membrane through its interaction with FcRn, and thus avoid degradation.

[0169] In one embodiment, the K_D value of the interaction between FcRn binding polypeptide and FcRn at pH 7.4 is higher than the K_D value of said interaction at pH 6.0. Thus, the FcRn binding polypeptide would bind to FcRn with higher affinity at pH 6.0 than at pH 7.4. In one embodiment, the K_D value of said interaction at pH 7.4 is at least 2 times higher, such as at least 5 times higher, such as at least 10 times higher, such as at least 50 times higher, such as at least 100 times higher than the K_D value of said interaction at pH 6.0.

[0170] In one embodiment, the K_D value of the interaction between FcRn binding polypeptide and FcRn at pH 7.4 is at least 1×10^{-8} M, such as at least 1×10^{-7} M, such as at least 1×10^{-6} M, such as at least 1×10^{-5} M. In some embodiments, the only criterion for the interaction between FcRn binding polypeptide and FcRn at pH 7.4 is that any FcRn binding polypeptide which has bound to FcRn during more acidic conditions is released more rapidly from FcRn when the pH value increases.

[0171] In an alternative embodiment, there is provided an FcRn binding polypeptide, for which the K_D of said interaction at pH 7.4 is the same as or lower than the K_D of said interaction at pH 6.0. An FcRn binding polypeptide according to this embodiment would bind or remain bound to FcRn in acidic pH conditions (i.e. would have an off-rate at pH 6.0 which is sufficiently slow to avoid release), for example in the lysosome, as well as in neutral or slightly basic pH conditions, for example on the plasma membrane. In a more specific embodiment, the K_D value of said interaction at pH 7.4 is at least 2 times lower, such as at least 5 times lower, such as at least 10 times lower, such as at least 50 times lower, such as at least 100 times lower than the K_D value of said interaction at pH 6.0.

[0172] In another embodiment, there is provided an FcRn binding polypeptide, which is capable of binding to FcRn at pH 7.4 such that the K_D value of the interaction is at most 1×10^{-6} M, such as at most 1×10^{-7} M, such as at most 1×10^{-8} M, such as at most 1×10^{-9} M, such as at most 1×10^{-10} M. An FcRn binding polypeptide according to this embodiment would bind or remain bound for an extended time to FcRn in neutral or slightly basic pH conditions, such as pH 7.4, for example on the plasma membrane. The term "remain bound" should be understood to mean an interaction having a slow off-rate at given conditions.

[0173] In general, the skilled person knows that the K_D value of an interaction is defined as the ratio between the off-rate (k_{off}) and the on-rate (k_{on}). Thus, a high K_D value may be due to either a high k_{off} , a low k_{on} or both, and conversely, a low K_D value may be due to either a low k_{off} , a high k_{on} or both.

[0174] The skilled person will understand that various modifications and/or additions can be made to an FcRn binding polypeptide according to any aspect disclosed herein in order to tailor the polypeptide to a specific application without departing from the scope of the present disclosure.

[0175] For example, in one embodiment there is provided an FcRn binding polypeptide as described herein, which polypeptide has been extended by one or more amino acids at the C terminal and/or N terminal end. Such a polypeptide should be understood as a polypeptide having one or more additional amino acid residues at the very first and/or the very last position in the polypeptide chain. Thus, an FcRn binding polypeptide may comprise any suitable number of additional amino acid residues, for example at least one additional amino acid

residue. Each additional amino acid residue may individually or collectively be added in order to, for example, improve production, purification, stabilization *in vivo* or *in vitro*, coupling, or detection of the polypeptide. Such additional amino acid residues may comprise one or more amino acid residues added for the purpose of chemical coupling. One example of this is the addition of a cysteine residue. Such additional amino acid residues may also provide a "tag" for purification or detection of the polypeptide, such as a His₆ tag or a "myc" (c-myc) tag or a "FLAG" tag for interaction with antibodies specific to the tag or immobilized metal affinity chromatography (IMAC) in the case of the hexahistidine tag.

[0176] The further amino acids as discussed above may be coupled to the FcRn binding polypeptide by means of chemical conjugation (using known organic chemistry methods) or by any other means, such as expression of the FcRn binding polypeptide as a fusion protein or joined in any other fashion, either directly or via a linker, for example an amino acid linker.

[0177] The further amino acids as discussed above may for example comprise one or more polypeptide domain(s). A further polypeptide domain may provide the FcRn binding polypeptide with another function, such as for example another binding function, or an enzymatic function, or a toxic function or a fluorescent signaling function, or combinations thereof.

[0178] A further polypeptide domain may moreover provide another FcRn binding moiety with the same FcRn binding function. Thus, in a further embodiment, there is provided an FcRn binding polypeptide in a multimeric form. Said multimer is understood to comprise at least two FcRn binding polypeptides as disclosed herein as monomer units, the amino acid sequences of which may be the same or different. Multimeric forms of the polypeptides may comprise a suitable number of domains, each having an FcRn binding motif, and each forming a monomer within the multimer. These domains may have the same amino acid sequence, but alternatively, they may have different amino acid sequences. In other words, the FcRn binding polypeptide of the invention may form homo- or heteromultimers, for example homo- or heterodimers. In one embodiment, there is provided an FcRn binding polypeptide, wherein said monomeric units are covalently coupled together. In another embodiment, said FcRn binding polypeptide monomer units are expressed as a fusion protein. In one embodiment, there is provided an FcRn binding polypeptide in dimeric form.

[0179] Additionally, "heterogenic" fusion polypeptides or proteins, or conjugates, in which an FcRn binding polypeptide described herein, or multimer thereof, constitutes a first domain, or first moiety, and the second and further moieties have other functions than binding FcRn, are also contemplated and fall within the ambit of the present disclosure. The second and further moiety/moieties of the fusion polypeptide or conjugate in such a protein suitably have a desired biological activity.

[0180] Thus, in a second aspect of the present disclosure, there is provided a fusion protein or a conjugate, comprising a first moiety consisting of an FcRn binding polypeptide according to the first aspect, and a second moiety consisting of a polypeptide having a desired biological

activity. In another embodiment, said fusion protein or conjugate may additionally comprise further moieties, comprising desired biological activities that can be either the same or different from the biological activity of the second moiety.

[0181] Such heterogenic fusion polypeptides could also be used to create heteromultimeric complexes of higher order. One example is a heterodimeric complex of two fusion polypeptides, each comprising an FcRn binding polypeptide according to the present disclosure in fusion with another moiety. Such a complex could for example form a heterodimer *in vivo* or *in vitro* and be held together by non-covalent and/or covalent interactions. A specific example of such a complex is a Fab fragment, in which both the light chain and heavy chain are produced in fusion with one FcRn binding polypeptide each, and which may include an inter-domain disulphide bond. Many biologically relevant, heterodimeric complexes known to the skilled person may be constructed using FcRn binding fusion proteins as monomer units.

[0182] In one embodiment of said fusion protein or conjugate, the total size of the molecule is below the threshold for efficient renal clearance upon administration to a mammalian subject.

[0183] In another embodiment of said fusion protein or conjugate, the total size of the molecule is above the threshold for efficient renal clearance upon administration to a mammalian subject.

[0184] In one embodiment, there is provided a fusion protein or conjugate, wherein the *in vivo* half-life of said fusion protein or conjugate is longer than the *in vivo* half-life of the polypeptide having the desired biological activity *per se*.

[0185] Non-limiting examples of a desired biological activity comprise a therapeutic activity, a binding activity, and an enzymatic activity.

[0186] In one embodiment, said desired biological activity is a binding activity to a selected target.

[0187] One example of such a binding activity is a binding activity, which increases the *in vivo* half-life of a fusion protein or conjugate. This fusion protein or conjugate may comprise at least one further moiety. In one particular embodiment, said target is albumin, binding to which increases the *in vivo* half-life of said fusion protein or conjugate. In one embodiment, said albumin binding activity is provided by an albumin binding domain (ABD) of streptococcal protein G or a derivative thereof. For example, said fusion protein or conjugate, comprising at least one further moiety, may comprise [FcRn binding polypeptide moiety] - [albumin binding moiety] - [moiety with affinity for selected target]. It is to be understood that the three moieties in this example may be arranged in any order from the N- to the C-terminal of the polypeptide.

[0188] In one embodiment, when a complex between a target and the fusion protein or conjugate as described herein is formed (or maintained) at acidic pH, such as pH 6.0, the target is rescued from elimination by lysosomal degradation. Thus, target half-life is extended.

Half-life extension implies that the elimination rate of a target is lower when interacting with said fusion protein or conjugate than the elimination rate of the target molecule in the absence of said fusion protein or conjugate. Furthermore, it is desirable in this embodiment that the binding of target by the fusion protein or conjugate should not interfere substantially with the function of the target.

[0189] On the other hand, when a complex between the target and the fusion protein or conjugate as described herein is not maintained or not formed at acidic pH, the target is directed to the subcellular lysosomes where it is degraded.

[0190] In one embodiment, there is provided a fusion protein or conjugate, wherein the rate of elimination of a selected, undesirable target from the subject is increased. Increased elimination of an undesirable target implies increased elimination rate of the target from the body of the multicellular organism, as compared to a "normal" elimination rate of the target molecule *per se*, i.e. without previous interaction with the fusion protein or conjugate.

[0191] In another embodiment, binding of a selected undesirable target could inactivate the function of the target, thereby blocking its biological activity in situations where this is desirable. Such biological activity may for example be activation or blocking of receptors or an enzymatic or otherwise toxic or undesirable activity. Such undesirable target may be an endogenous hormone, enzyme, cytokine, chemokine or a target having some other biological activity. By using an inactivating target binding, the biological activity is blocked until the target is delivered for degradation and released at a low pH value, and the target binding fusion protein is recycled to circulation. This recycling of the target binding fusion protein (via its FcRn binding moiety) enables it to "catalyze" the removal of more than one molecule of the selected undesirable target.

[0192] Undesirable targets may for example be foreign proteins and compounds, or naturally expressed proteins that display elevated levels in plasma following a medical condition and where a therapeutic effect may be achieved by elimination of said protein. The undesired target is not necessarily evenly distributed in the plasma but may be concentrated in certain regions, for example around a tumor or at sites of inflammation.

[0193] Non-limiting examples of targets are targets selected from the group consisting of allergens, amyloids, antibodies, auto-antigens, blood clotting factors, hormones, tumor cells, drug molecules, cytokines, chemokines, proteases, hypersensitivity mediators, proinflammatory factors, toxins such as bacterial toxins and snake venoms; pollutants, metals and anti-oxidants.

[0194] Under certain conditions, such as in certain cancer diseases, it is desired to remove endogenous molecules, for example VEGF, PDGF, HGF and other growth stimulatory hormones. Such molecules could also be targeted by a binding function in said fusion protein or conjugate.

[0195] Under other conditions, such as in certain immunological diseases, it may be desirable to remove endogenous molecules transiently, such as selected interleukines or TNF. Such molecules could also be targeted by a binding function in said fusion protein or conjugate.

[0196] In one embodiment, the second moiety having a desired biological activity is a therapeutically active polypeptide. Non-limiting examples of therapeutically active polypeptides are biomolecules, such as molecules selected from the group consisting of enzymes, for example algalidase α and β , glucocerebrosidase, laronidase, arylsulphatase, aglucosidase- α , asparaginase, Factor VII, Factor VIII, Factor IX and Factor Xa; hormones and growth factors, for example growth hormone, transforming growth factor- β 2, erythropoietin, insulin, insulin-like growth factor-1, myostatin, bone-derived growth factor and glucagon-like peptide-1; chemokines, for example CCL17, CCL19, CCL20, CCL21, CCL22, CCL27, XCL1 and CXCL3; and cytokines, for example interleukin (IL)-2, IL-4, IL-7, IL-10, IL-12, IL-15, IL-18, IL-22, IL-27, interferon (IFN)- α , IFN- β , IFN- γ , tumor necrosis factor, granulocyte-colony stimulating factor (G-CSF), macrophage-CSF, and granulocyte/macrophage-CSF.

[0197] As the skilled person understands, the FcRn binding polypeptide according to the first aspect may be useful in a fusion protein or as a conjugate partner to any other moiety. Therefore, the above lists of therapeutically active polypeptides should not be construed as limiting in any way.

[0198] Other possibilities for the creation of fusion polypeptides or conjugates are also contemplated. Thus, an FcRn binding polypeptide according to the first aspect of the invention may be covalently coupled to a second or further moiety or moieties, which in addition to or instead of target binding exhibit other functions. One example is a fusion between one or more FcRn binding polypeptide(s) and an enzymatically active polypeptide serving as a reporter or effector moiety.

[0199] With regard to the description above of fusion proteins or conjugates incorporating an FcRn binding polypeptide according to the disclosure, it is to be noted that the designation of first, second and further moieties is made for clarity reasons to distinguish between FcRn binding polypeptide or polypeptides according to the disclosure on the one hand, and moieties exhibiting other functions on the other hand. These designations are not intended to refer to the actual order of the different domains in the polypeptide chain of the fusion protein or conjugate. Thus, for example, said first moiety may without restriction appear at the N-terminal end, in the middle, or at the C-terminal end of the fusion protein or conjugate.

[0200] In one embodiment, there is provided an FcRn binding polypeptide, fusion protein or conjugate, which binds to FcRn such that binding of IgG to FcRn is at least partially inhibited. This inhibition may be due to binding of the FcRn binding polypeptide, fusion protein or conjugate to the same, or an at least partially overlapping, region of FcRn as IgG. Alternatively, the FcRn binding polypeptide, fusion protein or conjugate may bind to a different region of FcRn than IgG but sterically hinder the binding of IgG to FcRn. Thus, the rate of elimination or clearance of IgG from the circulatory system would increase due to increased lysosomal

degradation of IgG, because the FcRn mediated recycling of IgG would be wholly or partially unavailable due to the occupation of FcRn binding sites by the FcRn binding polypeptide according to the present disclosure. In other words, administration of FcRn binding polypeptide, fusion protein or conjugate or composition according to the present disclosure will act to increase the catabolism of circulating IgG antibodies.

[0201] In one embodiment, the K_D value of the interaction between the FcRn binding polypeptide, fusion protein or conjugate and FcRn is lower than the K_D of the interaction between IgG and FcRn. This relationship may be true at both pH 6.0 and pH 7.4, or at pH 6.0 only.

[0202] The above aspects furthermore encompass polypeptides in which the FcRn binding polypeptide according to the first aspect, or the FcRn binding polypeptide as comprised in a fusion protein or conjugate according to the second aspect, further comprises a label, such as a label selected from the group consisting of fluorescent dyes and metals, chromophoric dyes, chemiluminescent compounds and bioluminescent proteins, enzymes, radionuclides and particles. Such labels may for example be used for detection of the polypeptide.

[0203] In other embodiments, the labeled FcRn binding polypeptide is present as a moiety in a fusion protein or conjugate also comprising a second moiety having a desired biological activity and/or comprising a binding function as described above. The label may in some instances be coupled only to the FcRn binding polypeptide, and in some instances both to the FcRn binding polypeptide and to the second moiety of the conjugate or fusion protein. Furthermore, it is also possible that the label may be coupled to a second moiety only and not to the FcRn binding moiety. Hence, in yet another embodiment there is provided an FcRn binding polypeptide comprising a second moiety, wherein said label is coupled to the second moiety only.

[0204] When reference is made to a labeled polypeptide, this should be understood as a reference to all aspects of polypeptides as described herein, including fusion proteins and conjugates comprising an FcRn binding polypeptide and a second and optionally further moieties. Thus, a labeled polypeptide may contain only the FcRn binding polypeptide and e.g. a therapeutic radionuclide, which may be chelated or covalently coupled to the FcRn binding polypeptide, or contain the FcRn binding polypeptide, a therapeutic radionuclide and a second moiety such as a small molecule having a desired biological activity, for example resulting in a therapeutic efficacy.

[0205] In embodiments where the FcRn binding polypeptide, fusion protein or conjugate is radiolabeled, such a radiolabeled polypeptide may comprise a radionuclide. A majority of radionuclides have a metallic nature, are used in the ionic form, and are typically incapable of forming stable covalent bonds with elements presented in proteins and peptides. For this reason, labeling of proteins and peptides with radioactive metals is performed with the use of chelators, i.e. multidentate ligands, which form non-covalent compounds, called chelates, with the metal ions. In an embodiment of the FcRn binding polypeptide, fusion protein or conjugate,

the incorporation of a radionuclide is enabled through the provision of a chelating environment, through which the radionuclide may be coordinated, chelated or complexed to the polypeptide.

[0206] One example of a chelator is the polyaminopolycarboxylate type of chelator. Two classes of such polyaminopolycarboxylate chelators can be distinguished: macrocyclic and acyclic chelators.

[0207] In one embodiment, the FcRn binding polypeptide, fusion protein or conjugate comprises a chelating environment provided by a polyaminopolycarboxylate chelator conjugated to the FcRn binding polypeptide via a thiol group of a cysteine residue or an epsilon amine group of a lysine residue.

[0208] The most commonly used macrocyclic chelators for radioisotopes of indium, gallium, yttrium, bismuth, radioactinides and radiolanthanides are different derivatives of DOTA (1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid). In one embodiment, a chelating environment of the FcRn binding polypeptide, fusion protein or conjugate is provided by DOTA or a derivative thereof. More specifically, in one embodiment, the chelating polypeptides encompassed by the present disclosure are obtained by reacting the DOTA derivative 1,4,7,10-tetraazacyclododecane-1,4,7-tris-acetic acid-10-maleimidoethylacetamide (maleimidomonoamide-DOTA) with said polypeptide.

[0209] Additionally, 1,4,7-triazacyclononane-1,4,7-triacetic acid (NOTA) and derivatives thereof may be used as chelators. Hence, in one embodiment, there is provided an FcRn binding polypeptide, fusion protein or conjugate, wherein the polyaminopolycarboxylate chelator is 1,4,7-triazacyclononane-1,4,7-triacetic acid or a derivative thereof.

[0210] The most commonly used acyclic polyaminopolycarboxylate chelators are different derivatives of DTPA (diethylenetriamine-pentaacetic acid). Hence, polypeptides having a chelating environment provided by diethylenetriaminepentaacetic acid or derivatives thereof are also encompassed by the present disclosure.

[0211] In a further embodiment, the FcRn binding polypeptide, produced recombinantly through expression of a polynucleotide or synthetically, is conjugated to one or more synthetic polymers, in order for example to increase its hydrodynamic radius. Polyethylene glycol (PEG) is commonly used for this purpose, but other polymers have also been used in the art. Such "PEGylation" may be used to increase the size of the FcRn binding polypeptide of any of the types described herein to a size above the threshold for effective renal excretion.

[0212] In one embodiment, a synthetic polymer is conjugated to one or more chemically synthesized, monomeric FcRn binding polypeptides. Other functionalities may also be conjugated to the same synthetic polymer. If the FcRn binding polypeptide and other components are chemically synthesized, none of the components will have to be made in a biological system if this is not desired.

[0213] In a preferred embodiment, one or more synthetically or biologically manufactured FcRn binding polypeptides are conjugated to a synthetic polymer, to achieve a size exceeding the size associated with efficient renal clearance and used for blocking binding of IgG to FcRn. A unique cysteine in each FcRn binding polypeptide may be used for site specific conjugation, for example a C-terminally located cysteine introduced for this purpose. With a branched synthetic polymer, more than two FcRn binding moieties may be conjugated to the same polymer, to enhance the avidity and therefore the blocking potency.

[0214] In a third aspect of the present disclosure, there is provided a polynucleotide encoding an FcRn binding polypeptide or a fusion protein as described herein. Also encompassed by this disclosure is a method of producing a polypeptide or fusion protein as described above comprising expressing a polynucleotide; an expression vector comprising the polynucleotide; and a host cell comprising the expression vector.

[0215] Also encompassed is a method of producing a polypeptide, comprising culturing said host cell under conditions permissive of expression of said polypeptide from its expression vector, and isolating the polypeptide.

[0216] The FcRn binding polypeptide of the present disclosure may alternatively be produced by non-biological peptide synthesis using amino acids and/or amino acid derivatives having protected reactive side-chains, the non-biological peptide synthesis comprising

- step-wise coupling of the amino acids and/or the amino acid derivatives to form a polypeptide according to the first aspect having protected reactive side-chains,
- removal of the protecting groups from the reactive side-chains of the polypeptide, and
- folding of the polypeptide in aqueous solution.

[0217] In a fourth aspect of the disclosure, there is provided a composition comprising an FcRn binding polypeptide, fusion protein or conjugate as described herein and at least one pharmaceutically acceptable excipient or carrier. In one embodiment thereof, said composition further comprises at least one additional active agent, such as at least two additional active agents, such as at least three additional active agents. Non-limiting examples of additional active agents that may prove useful in such a combination are immunosuppressing agents, anti-inflammatory agents, anti-microbial agents and enzymes.

[0218] In one embodiment of this aspect, said composition is adapted for administration by a route selected from the group consisting of oral administration, intranasal administration, pulmonar administration, vaginal administration, rectal administration, intravenous injection, intraperitoneal injection, intramuscular injection, subcutaneous injection and intradermal injection.

[0219] As used herein, the term "systemic administration" refers to a route of administration such the substance of interest enters into the circulatory system so that the entire body is

affected. The skilled person is aware that systemic administration can take place via enteral administration (absorption of the drug through the gastrointestinal tract) or parenteral administration (generally injection, infusion or implantation).

[0220] In one embodiment, said composition is adapted for administration systemically or locally. In certain embodiments, systemic administration of said compound may be used. In another embodiment, said composition is adapted for administration by a local route. For example, local administration may be topical in an ointment, paste, foam or cream. In another embodiment, said composition is adapted for administration across an endothelial or epithelial layer. Here, the composition may be transcytosed across said layer.

[0221] In one embodiment, the rate of uptake of a composition comprising a fusion protein or conjugate as described herein is higher than the rate of uptake of polypeptides corresponding to second or further moieties *per se*. In one embodiment, the rate of uptake is at least 2 times higher, such as at least 5 times higher, such as at least 10 times higher, such as at least 25 times higher than the rate of uptake of the at second or further moieties *per se*.

[0222] It should be understood from the above disclosure that the FcRn binding polypeptide fusion protein or conjugate or the composition as described herein may for example be useful as a therapeutic agent, and/or as a means for extending the *in vivo* half-life of a fusion partner, and/or as a means for increasing the rate of elimination of undesirable targets.

[0223] Hence, in a fifth aspect of the present disclosure, there is provided an FcRn binding polypeptide, fusion protein, conjugate or composition as disclosed herein for use as a medicament.

[0224] In a related, sixth, aspect of the present disclosure, there is provided a method of treatment of a subject in need thereof, comprising the step of administering a therapeutically active amount of an FcRn binding polypeptide, fusion protein, conjugate or composition as disclosed herein.

[0225] In one embodiment of any one of these two latter aspects, the medicament or method is intended for treatment in which the capacity of the FcRn binding polypeptide to at least partially block binding of IgG to FcRn is exploited, i.e. treatment in which increased catabolism of IgG antibodies is desired. In one embodiment, a condition in which such treatment may be indicated is an auto-immune condition. As non-limiting examples of indicated conditions, mention is made of myasthenia gravis, Guillain-Barré syndrome, autoimmune limbic encephalitis, pediatric autoimmune neuropsychiatric disorders associated with streptococcal infection (PANDAS), neuromyotonia (Isaac's syndrome), morvan syndrome, multiple sclerosis, pemphigus vulgaris, foliaceus, bullous pemphigoid, epidermolysis bullosa acquisita, pemphigoid gestationis, mucous membrane pemphigoid, lichen sclerosus, antiphospholipid syndrome, relapsing polychondritis, autoimmune anemia, idiopathic thrombocytic purpura, autoimmune Grave's disease, dilated cardiomyopathy, vasculitis, Goodpasture's syndrome, idiopathic membranous nephropathy, rheumatoid arthritis and systemic lupus erythematosus.

[0226] In another embodiment, there is provided an FcRn binding polypeptide, fusion protein, conjugate or composition as described herein for use in blocking or removal of an undesirable target from the circulation. In one embodiment, said undesirable target is selected from the group comprising allergens, amyloids, antibodies, auto-antigens, blood clotting factors, hormones, tumor cells, drug molecules, cytokines, chemokines, hypersensitivity mediators, pro-inflammatory factors, toxins such as bacterial toxins and snake venoms, pollutants, metals and anti-oxidants.

[0227] While the invention has been described with reference to various exemplary aspects and embodiments, it will be understood by those skilled in the art that various changes may be made and equivalents may be substituted for elements thereof without departing from the scope of the invention. In addition, many modifications may be made to adapt a particular situation or molecule to the teachings of the invention without departing from the essential scope thereof. Therefore, it is intended that the invention not be limited to any particular embodiment contemplated, but that the invention will include all embodiments falling within the scope of the appended claims.

Brief description of the figures

[0228]

Figure 1 is a listing of the amino acid sequences of examples of FcRn binding motifs comprised in FcRn binding polypeptides of the invention (SEQ ID NO:1-353), examples of 49-mer FcRn binding polypeptides according to the disclosure (SEQ ID NO:354-706), examples of 58-mer FcRn binding polypeptides according to the disclosure (SEQ ID NO:707-1062) as well as the amino acid sequences of the albumin binding polypeptide variant PP013 (SEQ ID NO:1063), *Taq* polymerase binding Z variant Z03638 (SEQ ID NO:1064), human α FcRn (SEQ ID NO:1065), murine α FcRn (SEQ ID NO:1070), human β 2-microglobulin (SEQ ID NO:1066), murine β 2-microglobulin (SEQ ID NO:1067), human α FcRn (SEQ ID NO:1068) when in human FcRn-eGFP and murine α FcRn (SEQ ID NO:1069) when in murine FcRn-eGFP.

Figures 2A-2E show the binding to human FcRn at pH 6.0 and dissociations at pH 6.0 and 7.4 for His₆-tagged Z variants and for IgG as described in Example 3. Overlays of sensorgrams obtained from a Biacore instrument representing injection at pH 6.0 followed by dissociation at pH 6.0 (solid line) and injection at pH 6.0 followed by dissociation at pH 7.4 (dashed line) are displayed for (A) Z07918 (SEQ ID NO:707), (B) Z07960 (SEQ ID NO:710), (C) Z10109 (SEQ ID NO:709), (D) Z10193 (SEQ ID NO:708) and (E) IgG.

Figure 3 shows dot plots from a flow cytometry analysis of binding of FcRn binding Z variant to human (upper panel) and mouse (lower panel) FcRn-eGFP HeLa cells, as described in Example 4. Due to heterogeneous expression of FcRn-eGFP by HeLa cells, cells were gated according to FcRn-eGFP expression level. Cells in gate H are considered to be FcRn-eGFP negative and cells in gate I are considered to be positive. Incubation with Alexa647 labeled Z

variants resulted in a population positive both for Alexa647 and eGFP, whereas incubation with buffer (buffer control) did not. The figure shows that the three variants Z07960 (SEQ ID NO:710), Z07930 (SEQ ID NO:712) and Z07918 (SEQ ID NO:707) bind to human FcRn and mouse FcRn. The y-axis shows Alexa647 intensity and the x-axis shows eGFP activity.

Figure 4 shows mean fluorescence intensity (MFI) values of Alexa647 labeled Z07960 (SEQ ID NO:710), Z07930 (SEQ ID NO:712) and Z07918 (SEQ ID NO:707), measured in the cell binding assay described in Example 4. Diagram (A) shows MFI from HeLa cells transduced with human FcRn-eGFP and diagram (B) shows MFI from HeLa cells transduced with mouse FcRn-eGFP.

Figure 5 shows dot plots from flow cytometry analysis of human or mouse IgG Alexa647 binding to human (upper panel) and mouse (lower panel) FcRn-eGFP HeLa cells, as described in Example 5. Due to heterogeneous expression of FcRn-eGFP by HeLa cells, cells were gated according to the abundance of FcRn-eGFP on the cell surface. Cells in gate M are considered to be FcRn-eGFP negative and cells in gate N are considered to be positive. Binding of 100 nM human or mouse IgG-Alexa647 to FcRn transduced HeLa cells are shown in the left panel (0 nM). The figure shows that IgG binding was blocked by His₆-tagged Z07918 (SEQ ID NO:707) in a dose dependent manner (1, 10, 100 and 1000 nM). The y-axis shows Alexa647 intensity and the x-axis shows eGFP activity.

Figure 6 shows mean fluorescence intensity (MFI) values resulting from FcRn binding of IgG Alexa647 in the presence of different concentrations of His₆-tagged Z07918 (SEQ ID NO:707) on (A) human FcRn-eGFP transduced HeLa cells and (B) mouse FcRn-eGFP transduced HeLa cells, as described in Example 5. The figure shows dose dependent blocking of the IgG-FcRn binding by the Z variant.

Figures 7A-7C show kinetics of binding of three Z variants to human FcRn at pH 6.0, as described in Example 6, using a Biacore instrument. Sensorgrams for a concentration series of (A) Z11948 (SEQ ID NO:1060), (B) Z11946 (SEQ ID NO:1061) and (C) Z11947 (SEQ ID NO:1062), respectively, in fusion with the albumin binding polypeptide PP013 (SEQ ID NO:1063) and the control Z variant molecule Z03638 (SEQ ID NO:1064; not specific for FcRn), are displayed. Curves from 640 nM (dashed line), 160 nM (dotted line) and 40 nM (solid grey line) were subjected to kinetic analysis using the Langmuir 1:1 binding model. Kinetic parameters and affinities were calculated from fitted curves (solid black lines) and are shown in Table 5.

Figure 8 shows the pharmacokinetic profiles for three FcRn binding Z variants fused to the albumin binding polypeptide PP013 obtained as described in Example 6. The Z variants Z11947 (SEQ ID NO: 1062, open squares), Z11946 (SEQ ID NO:1061, open triangles) and Z11948 (SEQ ID NO:1060, open diamonds) all displayed prolonged half-life compared to the negative control PP013-Z03638 (open circles).

Figure 9 shows the blocking of human IgG to human FcRn by His₆-Z07918 (SEQ ID NO:707; black circles), IVIg (grey squares) and SCIg (grey triangles), respectively, assayed as described in Example 10.

Figure 10 shows that blocking of the IgG-FcRn interactions with FcRn specific Z variants in mice results in reduced levels of IgG. As further described in Example 11, mice were treated with five daily injections of Vehicle (+), the ABD fused Z variant Z07918-PP013 (open square) and Z11948 (SEQ ID NO:1060; closed circle). The concentration of endogenous IgG was measured by ELISA. The concentration of IgG in individual mice at 24, 72, 120 and 168 h were related to the level at 0 h and the results are therefore presented as percentage of IgG at 0 h.

Examples

Summary

[0229] The following Examples disclose the development of novel Z variant molecules targeting the neonatal Fc receptor (FcRn). The Z variants were obtained using phage display technology. The genes encoding FcRn binding polypeptides described herein were sequenced, and the corresponding amino acid sequences are listed in Figure 1, and denoted by the identifiers SEQ ID NO:707-1059. Also, the deduced binding motifs of these selected binding variants are listed in Figure 1 with sequence identifiers SEQ ID NO:1-353.

Example 1

Production of human α FcRn and human β 2-microglobulin (B2M)

[0230] In this Example, the extracellular domain (ECD) of human α FcRn (SEQ ID NO:1065) in complex with human β 2-microglobulin (SEQ ID NO:1066) (complex denoted FcRn) and human β 2-microglobulin in non-complexed form (denoted B2M) were produced as soluble proteins. Human FcRn and B2M produced in this Example were used for phage selection, ELISA and Biacore assays in Examples 2 and 3.

Materials and methods

[0231] Construction of plasmids containing the genes for human α FcRn and human β 2-microglobulin to be used for co-expression: The genes encoding human α FcRn (Genbank BC008734.2) and human β 2-microglobulin (B2M) (Genbank BC032589.1) were obtained from OpenBiosystems. Using PCR overlap extension, a gene fragment encoding amino acids 24-290 of human α FcRn (α FcRn_{ECD}) (SEQ ID NO:1065) was amplified to a construct consisting of

attB1-site/Kozak sequence followed by a gene encoding: an Ig kappa chain leader sequence, hFcRn_{ECD}, a GS-linker and a flag tag, followed by an attB2 site. A similar construct was made containing a gene fragment encoding amino acids 21-119 of human B2M (SEQ ID NO:1066), except that a His₆ tag replaced the flag tag. The constructs were inserted into the plasmid pDONOR221 (Invitrogen, cat. no. 12536-017) by recombination using the Gateway system (Invitrogen, cat. no. 11789020, Gateway® BP Clonase® II Enzyme mix), according to the manufacturer's recommendations. After verification of correct sequences, the human α FcRn_{ECD} construct was inserted into 2K7_{bsd} (Suter et al. (2006) Stem Cells 24:615-623) using multi-site gateway cloning together with the promoter-containing plasmid pENTR-CMV (Tai et al. (2012) PLoS One 7(9):e46269), resulting in the vector 2K7_{bsd}-CMV-hFcRn_{ECD}. The human B2M gene construct was similarly inserted into 2K7_{neo} (Suter *et al.*, *supra*), giving the vector 2K7_{neo}-CMV-hB2M.

[0232] Cell culture, preparation of recombinant lentiviral vectors and gene insertions into SKOV-3 cell line: The HEK293T and SKOV-3 cell lines were obtained from ATCC. Cells were grown at 37 °C in a humidified incubator in the presence of 5 % CO₂. Complete medium for the HEK293T cell line was Dulbeccos modified eagle medium (DMEM) supplemented with 10 % fetal bovine serum (FBS), 1 % Antibiotic Antimycotic Solution (AA) and 1 % MEM Non-essential Amino Acid Solution (NEAA). Complete medium for the SKOV-3 cell line was McCoy's 5A medium supplemented with 10 % FBS and 1 % AA.

[0233] The plasmids 2K7_{bsd}-CMV-hFcRn_{ECD} and 2K7_{neo}-CMV-hB2M were separately co-transfected together with VSV-G envelope and gag/pol packaging plasmid into HEK293T cells using calcium chloride transfection (Zufferey et al. (1997) Nat Biotechnol 15(9):871-5; Jakobsson et al. (2006) J Neurosci Res 84:58-67). HEK293 culture supernatants containing formed lentiviral particles with human α FcRn_{ECD} and human B2M transgenes, respectively, were cleared from cell debris by centrifugation and filtration. The two types of lentiviral particles were used to sequentially transduce SKOV-3 cells. Successful double integrants containing both the human α FcRn_{ECD} and the B2M genes were selected for by the addition of blasticidin (Invitrogen) and G418 sulfate (Invitrogen) to culture medium while passaging the cells for two weeks. The resulting, stably transduced SKOV-3 cell line was denoted SKOV-3 hFcRn_{ECD}/hB2M.

[0234] Expression of recombinant human FcRn: SKOV-3 cells, co-expressing human α FcRn_{ECD} and B2M resulting in human FcRn, were expanded and 1.5×10^7 cells were seeded in a HYPERFlask (Corning) in 560 ml complete growth medium. After five days, when the cells had settled and multiplied, the medium was changed to complete growth medium without FBS. After five days, the culture was terminated and the supernatant was collected, passed through a 45 μ m filter and frozen at -80 °C.

[0235] Purification of recombinant human FcRn using human IgG chromatography: Protein purification was carried out in an ÄKTA Explorer system (GE Healthcare). Human IgG

(Pharmacia), 1 ml in 0.2 M NaHCO₃, 0.5 M NaCl pH 8.3 at a concentration of 10 mg/ml, was coupled to a 1 ml HiTrap NHS-activated HP column (GE Healthcare) according to the manufacturer's instruction. The supernatant containing recombinant human FcRn from SKOV-3 cells was thawed and the pH was adjusted to 5.8 with HCl. The supernatant was subsequently loaded in batches of 100 ml onto the column previously equilibrated with 20 mM Bis-Tris pH 5.8. The column was washed with 20 ml of 20 mM Bis-Tris pH 5.8 and eluted in fractions of 1 ml using 50 mM Tris, pH 8.1. Buffer exchange to PBS (phosphate buffered saline, 10 mM phosphate, 137 mM NaCl, 2.68 mM KCl, pH 7.4) was performed using dialysis.

[0236] SDS-PAGE and Western blot: The purity of the eluted fractions from the protein purification was analyzed by SDS-PAGE and staining with GelCode Blue Stain Reagent (Pierce) and SilverXpress® Silver Staining Kit (Invitrogen). Western blotting was carried out using an Amersham Hybond™-C Extra nitrocellulose membrane (GE Healthcare). The membrane was blocked with 5 % non-fat dry milk (Semper) in TBS+T (50 mM Trizma base, 150 mM NaCl, 0.05 % Tween-20, pH 8) for 1 hour, then probed with a mixture of rabbit anti-FCGRT polyclonal antibody (Atlas Antibodies) at a concentration of 0.15 µg/ml and rabbit anti-B2M polyclonal antibody (Atlas Antibodies) at a concentration of 0.23 µg/ml in TBS+T. The membrane was subsequently incubated with stabilized goat anti-rabbit antibody conjugated with horse radish peroxidase (Pierce) diluted 1:10,000 in TBS+T. After addition of TMB Substrate (Pierce), an image of the membrane was acquired on Amersham Hyperfilm ECL (GE Healthcare). The Hyperfilm was processed using GBX developer and GBX fixer (Sigma-Aldrich).

[0237] Production of a non-complexed form of human B2M: Human B2M was produced in *E. coli*. The expression and purification was performed essentially as described in Sandalova et al. (2005) Acta Chryst F61:1090-1093 and Michaëlsson et al. (2001) J Immunol 166:7327-7334. The purified protein, consisting of amino acids 21-119 of human B2M, in urea was subjected to arginine refolding as follows; 0.5 mg of B2M was rapidly added to 2 ml refolding buffer (20 ml 1 M Tris-HCl pH 8.0, 16.87 g L-Arginine (buffered with HCl), 0.8 ml 0.5 M EDTA, 61 mg GSSG, 307 mg GSH and milli-Q water to a final volume of 200 ml, pH 8.0, and supplemented with protease inhibitor (Roche, cat. no. 11 873 580 001)). The refolding procedure was performed at 4 °C during 4 hours. Refolded B2M protein was buffer exchanged to PBS using a PD-10 column (GE Healthcare).

Results

[0238] Construction of plasmids containing the genes for human αFcRn and human β2-microglobulin to be used for co-expression: Genes encoding the extracellular domain of the α-chain of human FcRn (αFcRn_{ECD}) and human B2M were inserted into the lentiviral transfer plasmids 2K7_{bsd} and 2K7_{neo}, respectively. In both cases, the inserted gene is under the control of a CMV promoter. The genes were extended so that the resulting proteins would have an Ig kappa chain leader sequence in the N-terminus to target the protein for export through the

endoplasmic reticulum to the culture medium (the signal sequence was cleaved upon secretion). In addition, α FcRn_{ECD} had a C-terminal spacer sequence followed by a FLAG-tag for potential detection. Human B2M had a C-terminal spacer sequence followed by a His₆ tag for potential detection. The spacer sequence was added to enhance accessibility of the tag. The lentiviral transfer plasmids also contained two different antibiotic resistance genes to allow selection of cells where both constructs had been inserted.

[0239] Expression and purification of recombinant human FcRn: The genes encoding α FcRn_{ECD} and B2M were inserted into the genome of SKOV-3 by lentiviruses, and the resulting FcRn protein was secreted into the culture medium. To capture only FcRn having retained pH-dependent IgG binding, affinity chromatography using immobilized IgG was used where the receptor was captured at pH 5.8 and eluted at pH 8.1. Captured protein was eluted in three fractions.

[0240] SDS-PAGE and Western blot: To investigate the presence of two peptide chains (α FcRn_{ECD} and B2M) of the produced FcRn protein, and to analyze the purity of the eluted material, an SDS-PAGE analysis was performed on the eluted fractions. For the gel stained with GelCode Blue Stain, two bands were detected with molecular weights of 12 and 36 kDa, respectively. This corresponds approximately to the theoretical molecular weights of the non-glycosylated peptide chains of 12 kDa for B2M and 31 kDa for α FcRn_{ECD}. The α FcRn_{ECD} part of the protein contains one glycosylation site and it was therefore expected that its molecular mass would be higher than 31 kDa. The gel was also silver stained to increase sensitivity and possibly detect impurities. A band of approximately 66 kDa was detected in the first eluted fraction, which could correspond to BSA (bovine serum albumin) originating from cell attachment. The total amount of protein recovered in fraction 2 and 3 corresponded to 1.4 mg/l culture medium. A western blot analysis on the pooled material was carried out, which showed essentially only the two major bands and in addition a very weak band below 12 kDa which might correspond to a degradation product.

Example 2

Selection and ELISA binding of FcRn binding Z variants

[0241] In this Example, human FcRn was used as target in phage display selections using a phage library of Z variants. Selected clones were DNA sequenced, produced in *E. coli* periplasmic fractions and assayed against FcRn in ELISA (enzyme-linked immunosorbent assay).

Materials and methods

[0242] Biotinylation of target protein FcRn and of B2M: Human FcRn and human B2M, produced as described in Example 1, were biotinylated using No-Weigh EZ-Link Sulfo-NHS-LC-Biotin (Pierce, cat. no. 21327) at a 31 x (FcRn) and 10 x (B2M) molar excess, respectively, according to the manufacturer's recommendations. The reactions were performed at room temperature (RT) for 30 min. Subsequent buffer exchange to PBS was performed using Slide-a-lyzer dialysis cassettes (FcRn; Pierce, cat. no. 66380, 10,000 MWCO and B2M; Pierce, cat. no. 66333, 3,500 MWCO), according to the manufacturer's instructions.

[0243] Phage display selection of FcRn binding Z variants: A library of random variants of protein Z displayed on bacteriophage, constructed in phagemid pAY02592 essentially as described in Grönwall et al. (2007) J Biotechnol, 128:162-183, was used to select FcRn binding Z variants. In this library, an albumin binding domain (ABD, GA3 of protein G from *Streptococcus* strain G148) is used as fusion partner to the Z variants. The library is denoted Zlib006Naive.II and has a size of 1.5×10^{10} library members (Z variants). *E. coli* RRIΔM15 cells (Rüther et al., (1982) Nucleic Acids Res 10:5765-5772) from a glycerol stock containing the phagemid library Zlib006Naive.II, were inoculated in 20 l of a defined proline free medium [dipotassium hydrogenphosphate 7 g/l, trisodium citrate dihydrate 1 g/l, uracil 0.02 g/l, YNB (Difco™ Yeast Nitrogen Base w/o amino acids, Becton Dickinson) 6.7 g/l, glucose monohydrate 5.5 g/l, L-alanine 0.3 g/l, L-arginine monohydrochloride 0.24 g/l, L-asparagine monohydrate 0.11 g/l, L-cysteine 0.1 g/l, L-glutamic acid 0.3 g/l, L-glutamine 0.1 g/l, glycine 0.2 g/l, L-histidine 0.05 g/l, L-isoleucine 0.1 g/l, L-leucine 0.1 g/l, L-lysine monohydrochloride 0.25 g/l, L-methionine 0.1 g/l, L-phenylalanine 0.2 g/l, L-serine 0.3 g/l, L-threonine 0.2 g/l, L-tryptophane 0.1 g/l, L-tyrosine 0.05 g/l, L-valine 0.1 g/l], supplemented with 100 µg/ml ampicillin. The cultivations were grown at 37 °C in a fermenter (Belach Bioteknik, BR20). When the cells reached an optical density at 600 nm (OD₆₀₀) of 0.75, approximately 2.6 l of the cultivation was infected using a 10 x molar excess of M13K07 helper phage (New England Biolabs, cat. no. N0315S). The cells were incubated for 30 minutes, whereupon the fermenter was filled up to 20 l with TSB-YE (Tryptic Soy Broth-Yeast Extract; 30 g/l TSB, 5 g/l yeast extract) supplemented with 100 µM isopropyl-β-D-1-thiogalactopyranoside (IPTG) for induction of expression and with 25 µg/ml kanamycin and 12.5 µg/ml carbenicillin and grown at 30 °C for 22 h. The cells in the cultivation were pelleted by centrifugation at 15,900 g. The phage particles were precipitated from the supernatant twice in PEG/NaCl (polyethylene glycol/sodium chloride), filtered and dissolved in PBS and glycerol as described in Grönwall et al., *supra*. Phage stocks were stored at -80 °C before use.

[0244] Selections against biotinylated human FcRn were performed in four cycles divided in two different tracks. Phage stock preparation and selection procedure were performed essentially as described for selection against another biotinylated target in WO2009/077175. The amplification of phage between the selection cycles was performed by infecting *E. coli* RRIΔM15 with phage, then performing cultivation in solution as follows. Eluted phage and 10 x excess of M13K07 helper phage compared to bacteria were allowed to simultaneously infect log phase bacteria at 37 °C for 30 min without rotation, followed by 30 min with slow rotation. Prior to infection, bacteria were grown to log phase in the defined proline free medium

described above. Infected bacteria were pelleted by centrifugation at 4,300 *g* for 10 min and resuspended in 200 ml TSB+YE medium supplemented with 0.1 mM IPTG, 25 µg/ml kanamycin and 100 µg/ml ampicillin and cultivated at 30 °C overnight for phage production.

[0245] The selection buffer consisted of 100 mM sodium phosphate and 150 mM sodium chloride adjusted to pH 5.5 with hydrogen chloride and supplemented with 0.1 % gelatin and 0.1 % Tween-20. At selection, human serum albumin (HSA, Albucult, Novozymes) was added to the selection buffer to a final concentration of 1.5 µM. In order to reduce the amount of background binders, pre-selection was performed by incubation of phage stock with Dynabeads® M-280 Streptavidin (SA-beads, Dynal, cat. no. 112.06) for 1 hour at RT. A second pre-selection was performed during 30 min at RT against human B2M immobilized in immunotubes (Nunc, cat. no. 444474). 5 µg/ml of human B2M in carbonate buffer (Sigma, cat. no. 068K8214) was immobilized in the tube at 7 °C for >1 h. After washing twice with tap water, the tubes were blocked with PBS + 0.5 % casein (Sigma, cat. no. C8654) for 30 min at RT before use. All tubes and beads used in the selection were pre-blocked with PBS + 0.1 % gelatin. Selection was performed in solution at RT, followed by capture of target-phage complexes on SA-beads where 1 mg beads per 2.9 µg biotinylated FcRn were used. In cycle 1 of the selections, 100 nM biotinylated FcRn was used and two washes of two min each were performed using selection buffer. An increased stringency, using a lowered target concentration and an increased number of washes, was applied in the subsequent cycles: 50 nM/5 washes, 25 nM/8 washes and 10 nM/12 washes were applied in cycle 2, 3 and 4, respectively. After the washes, bound phage was eluted from the two selection tracks using two different procedures; 1) 500 µl 0.1 M glycine-HCl, pH 2.2, followed by immediate neutralization with 50 µl 1 M Tris-HCl, pH 8.0, and 450 µl PBS, or; 2) 500 µl of 100 mM sodium phosphate and 150 mM sodium chloride, pH 8.0 and neutralization with 500 µl PBS.

[0246] Sequencing: PCR fragments were amplified from single colonies using a standard PCR program and the primers AFFI-21 (5'-tgcttcggctcgtatgtgtgtg (SEQ ID NO:1071)) and AFFI-22 (5'-cggaaccagagccaccaccgg (SEQ ID NO:1072)). Sequencing of amplified fragments was performed using the biotinylated oligonucleotide AFFI-72 (5'-biotin-cggaaccagagccaccaccgg (SEQ ID NO:1073)) and a BigDye® Terminator v3.1 Cycle Sequencing Kit (Applied Biosystems), used in accordance with the manufacturer's protocol. The sequencing reactions were purified by binding to magnetic streptavidin coated beads (Detach Streptavidin Beads, Nordiag, cat. no. 2012-01) using a Magnatrix 8000 (Magnetic Biosolution), and analyzed on ABI PRISM® 3130xl Genetic Analyzer (PE Applied Biosystems).

[0247] Production of Z variants for ELISA: Sequenced Z variants were produced by inoculating single colonies from the selections into 10 ml TSB-YE medium supplemented with 100 µg/ml ampicillin and 0.1 mM IPTG and incubating for 24 h at 37 °C. Cells were pelleted by centrifugation, resuspended in 2 ml PBST (PBS supplemented with 0.05 % Tween-20), frozen at -80 °C and thawed in a water bath, to release the periplasmic fraction of the cells. The freeze-thawing procedure was repeated seven times and cells were then pelleted by centrifugation. The supernatant of the periplasmic extract contained the Z variants as fusions to ABD, expressed as AQHDEALE-[Z#####]-VDYV-[ABD]-YVPG (Grönwall *et al.*, *supra*).

Z##### refers to individual, 58 amino acid residue Z variants.

[0248] ELISA K_D analysis of Z variants: The binding of Z variants to FcRn was analyzed in ELISA assays. Half-area 96-well ELISA plates were coated with 2 µg/ml of an anti-ABD goat antibody (produced in-house) diluted in coating buffer (50 mM sodium carbonate, pH 9.6) at 4 °C overnight. The antibody solution was poured off and the wells were blocked with 100 µl of PBSC (PBS supplemented with 0.5 % casein) for 1.5 h at RT. The blocking solution was discarded and 50 µl periplasmic solution, diluted 1:4, was added to the wells and incubated for 1.5 h at RT under slow shaking. The solutions were poured off and the wells were washed four times with either 0.05% PCT buffer, pH 6.0 (McIlvaines phosphate-citrate buffer, pH 6.0, supplemented with 0.05 % Tween-20) or 0.05% PCT buffer, pH 7.4 (McIlvaines phosphate-citrate buffer, pH 7.4, supplemented with 0.05 % Tween-20). The target protein, biotinylated human FcRn, was added to the wells in a 1:3 diluted concentration series from 2 µg/ml (45 nM) to 0.3 ng/ml (6.9 pM) diluted in PCC buffer, pH 6.0 or pH 7.4, (McIlvaines phosphate-citrate buffer, pH 6.0 or pH 7.4, supplemented with 0.5 % casein), respectively. The plates were incubated for 1.5 h at RT followed by washes as described above. Streptavidin conjugated HRP (Thermo Scientific, cat. no. N100) was diluted 1:30 000 in PCC buffer, pH 6.0 or pH 7.4, respectively, and added to the wells followed by 45 min incubation. After washing as described above, 50 µl ImmunoPure TMB substrate (Thermo Scientific, cat. no. 34021) was added to the wells and the plates were treated according to the manufacturer's recommendations. Absorbance was measured at 450 nm using a multi-well plate reader, Victor³ (Perkin Elmer). A Z variant binding an irrelevant protein was used as negative control and a blank was created by omitting the periplasmic step. A Z variant which bound to FcRn in a pre-experiment (Z07918, SEQ ID NO:707) was used as positive control. Measured values were analyzed using GraphPad Prism 5 (GraphPad Software, LaJolla, CA, USA) and non-linear regression in order to determine the affinities (K_D) of the interactions.

[0249] ELISA specificity analysis of Z variants: In another ELISA experiment, the specificities of the Z variants were tested by assaying them against 2 µg/ml biotinylated human proteins B2M, PSMA (in house produced) and IgG (polyclonal, Pharmacia, Sweden) and against PCC buffer pH 6.0 or pH 7.4, respectively. The assay was performed at pH 6.0 and at pH 7.4, respectively, as described above. The biotinylated proteins or buffer were added to the wells instead of FcRn in the target protein step.

Results

[0250] Phage display selection of FcRn binding Z variants: Individual clones were obtained after four cycles of phage display selections against biotinylated human FcRn.

[0251] Sequencing: Sequencing was performed on clones picked at random from selection round four. Each Z variant was given a unique identification number ##### and individual variants are referred to as Z#####. The amino acid sequences of the 58 amino acid residues

long Z variants are listed in Figure 1 as SEQ ID NO:707-722 and SEQ ID NO:1059.

[0252] The deduced FcRn binding motifs of these Z variants are listed in Figure 1 as SEQ ID NO:1-16 and SEQ ID NO:353. The amino acid sequences of the 49 amino acid residues long polypeptides predicted to constitute the complete three-helix bundle within each of these Z variants are listed in Figure 1 as SEQ ID NO:354-369 and SEQ ID NO:706.

[0253] ELISA assays with Z variants: Sixteen clones were produced as ABD fusion proteins in *E. coli*. The periplasmic fractions were used in an ELISA against a dilution series of human FcRn. The clones were: Z07909 (SEQ ID NO:719), Z07918 (SEQ ID NO:707), Z07930 (SEQ ID NO:712), Z07960 (SEQ ID NO:710), Z10109 (SEQ ID NO:709), Z10111 (SEQ ID NO:714), Z10127 (SEQ ID NO:718), Z10129 (SEQ ID NO:715), Z10140 (SEQ ID NO:711), Z10141 (SEQ ID NO:716), Z10145 (SEQ ID NO:721), Z10152 (SEQ ID NO:720), Z10156 (SEQ ID NO:717), Z10161 (SEQ ID NO:722), Z10183 (SEQ ID NO:713) and Z10193 (SEQ ID NO:708). K_D values were determined for all variants at pH 6.0 and for three variants at pH 7.4 (Table 1). For thirteen variants, data was not obtained for a K_D analysis at pH 7.4. None of the sixteen variants displayed non-specific binding when assayed against human B2M, IgG or PSMA.

Table 1. ELISA K_D analysis of Z-ABD variants in *E. coli* periplasmic fractions.

Z variant	SEQ ID NO:	K_D pH 6.0 (M)	K_D pH 7.4 (M)
Z07909	719	24.5×10^{-9}	n.d.
Z07918	707	2.0×10^{-9}	10.9×10^{-9}
Z07930	712	10.4×10^{-9}	n.d.
Z07960	710	6.0×10^{-9}	n.d.
Z10109	709	3.9×10^{-9}	23.9×10^{-9}
Z10111	714	11.4×10^{-9}	n.d.
Z10127	718	21.3×10^{-9}	n.d.
Z10129	715	17.6×10^{-9}	n.d.
Z10140	711	8.8×10^{-9}	n.d.
Z10141	716	21.2×10^{-9}	n.d.
Z10145	721	42.0×10^{-9}	n.d.
Z10152	720	24.6×10^{-9}	n.d.
Z10156	717	21.3×10^{-9}	n.d.
Z10161	722	163.0×10^{-9}	n.d.
Z10183	713	10.9×10^{-9}	n.d.
Z10193	708	2.3×10^{-9}	25.9×10^{-9}
n.d.= not determinable			

Example 3**Production and characterization of FcRn binding Z variants**

[0254] In this Example, seventeen Z variants were produced in *E. coli*, purified and assayed against human FcRn in Biacore. A subset of said variants was also assayed against mouse FcRn. Circular dichroism (CD) spectroscopy was performed for a subset of Z variants for investigation of their secondary structure.

Materials and methods

[0255] Subcloning of Z variants: The DNA of seventeen FcRn binding Z variants (SEQ ID NO:707-722 and SEQ ID NO:1059) was amplified from the library vector pAY02592. A subcloning strategy for construction of monomeric Z variant molecules with N-terminal His₆ tag was applied using standard molecular biology techniques (essentially as described in detail in WO2009/077175 for Z variants binding another target). The Z gene fragments were subcloned into the expression vector pAY01448 resulting in the encoded sequence MGSSHHHHHHLQ-[Z#####]-VD.

[0256] In addition, the FcRn binding variant Z07918 (SEQ ID NO:707), but starting with the amino acids AE instead of VD and denoted Z11948 (SEQ ID NO:1060), was cloned as homodimeric constructs with two different linkers between the Z variants and followed by a C-terminal His₆ tag. This was performed using conventional molecular biology methods including DNA amplification, restriction with suitable restriction enzymes and ligation of the DNA. The two linkers were obtained from Thermo Fisher Scientific. The Z gene fragments were subcloned into the expression vector (pET-26 origin, Novagen) resulting in the encoded sequence [Z#####]-GT-(G₄S)-PR-[Z#####]-LEHHHHHH and [Z#####]-GT-(G₄S)₃-[Z#####]-LEHHHHHH, respectively.

[0257] Cultivation and purification: *E. coli* BL21 (DE3) cells (Novagen) were transformed with plasmids containing the gene fragment of each respective FcRn binding Z variant and cultivated at 37 °C in 800 or 1000 ml of TSB-YE medium supplemented with 50 µg/ml kanamycin. At OD₆₀₀ = 2, IPTG was added to induce expression at a final concentration of 0.17 or 0.2 mM and the culture was incubated at 37 °C for another 5 h. The cells were harvested by centrifugation.

[0258] Approximately 2-5 g of each cell pellet was resuspended in 10-25 ml binding buffer (20 mM sodium phosphate, 0.5 M NaCl, 20 mM imidazole, pH 7.4) supplemented with

Benzonase® (Merck, cat. no. 1.01654.0001) to a concentration of 15 U/ml and Lysozyme (Sigma, cat. no. L-7651) to a concentration of 0.5 mg/ml. After cell disruption by three freeze-thawing cycles or sonication, cell debris was removed by centrifugation and each supernatant was applied on a 1 ml His GraviTrap IMAC column (GE Healthcare, cat. no. 11-0033-99). Contaminants were removed by washing with wash buffer (20 mM sodium phosphate, 0.5 M NaCl, 20 or 60 mM imidazole, pH 7.4), and the FcRn binding Z variants were subsequently eluted with elution buffer 1 (20 mM sodium phosphate, 0.5 M sodium chloride, 250 mM imidazole, pH 7.4) or elution buffer 2 (0.1 M acetic acid, 0.5 M sodium chloride, pH 4.5). Purified Z variants were buffer exchanged to PBS using PD-10 columns (GE Healthcare), according to the manufacturer's protocol. Protein concentrations were determined by measuring the absorbance at 280 nm, using a NanoDrop® ND-1000 spectrophotometer, and using the extinction coefficient of the respective protein. The purity of the FcRn binding Z variants was analyzed by SDS-PAGE stained with Coomassie Blue. The identity of each purified FcRn binding Z variant was confirmed using LC/MS analysis.

[0259] CD analysis: Purified His₆-tagged Z variants were diluted to 0.5 mg/ml in PBS. For each diluted Z variant, a CD spectrum at 250-195 nm or 250-190 nm was obtained at 20 °C. In addition, a variable temperature measurement (VTM) was performed to determine the melting temperature (T_m). In the VTM, the absorbance was measured at 221 nm while the temperature was raised from 20 to 90 °C, with a temperature slope of 5 °C/min. A new CD spectrum was obtained at 20 °C after the heating procedure in order to study the refolding ability of the Z variants. The CD measurements were performed on a Jasco J-810 spectropolarimeter (Jasco Scandinavia AB) using a cell with an optical path-length of 1 mm.

[0260] Biacore binding and kinetic analysis: The interaction of FcRn binding His₆-tagged Z variants with human FcRn was analyzed in a Biacore 2000 instrument (GE Healthcare). Human FcRn was immobilized in a flow cell on the carboxylated dextran layer of a CM5 chip surface (GE Healthcare). The immobilization was performed using amine coupling chemistry according to the manufacturer's protocol and using HBS-EP (GE Healthcare) as running buffer. One flow cell surface on the chip was activated and deactivated for use as blank during analyte injections. In the two binding experiments presented below, Mcllvaines phosphate-citrate buffer pH 6.0 supplemented with 0.005 % Tween-20 (0.005% PCT) was used as running buffer. In all experiments, a flow rate of 50 µl/min was used.

[0261] In one experiment, the dissociation at pH 6.0 was compared to the dissociation at pH 7.4. His₆-tagged Z variants and a human monoclonal IgG1 were diluted in running buffer to a final concentration of 250 nM or 2.5 nM, respectively, and injected over the FcRn chip for 1 minute using the co-inject procedure. The second injection of the co-inject procedure, representing the dissociation phase of the interactions, contained either running buffer (pH 6.0) or 0.005% PCT pH 7.4. The Z variants were allowed to dissociate for 1 minute, except for Z07918 and Z10193, which were allowed to dissociate for 4 minutes, before a surface equilibration during 5 minutes in running buffer. IgG was allowed to dissociate for 4 minutes before equilibration. Buffer injections were performed in a similar way; co-injection of buffer pH 6.0 followed by pH 6.0 or co-injection of buffer pH 6.0 followed by pH 7.4. The results were

analyzed in BiaEvaluation software 4.1 (GE Healthcare). Curves of the blank surface were subtracted from the curves of the ligand surface. In addition, curves of buffer injections were subtracted from the Z variant curves and from the IgG curves to adjust for the buffer effects.

[0262] In another experiment, approximate kinetic constants (k_{on} and k_{off}) and affinities (K_D) were determined for a subset of His₆-tagged Z variants. Three concentrations of the Z variants were injected for 1 minute followed by dissociation in running buffer for 1 minute. The surfaces were equilibrated with running buffer during 7.5 minutes before the start of next cycle. Injected concentrations were either 675 nM, 225 nM and 75 nM (Z10140, Z10156 and Z10183) or 225 nM, 75 nM and 25 nM (Z07918 and Z10193). Kinetic constants were calculated from the sensorgrams using the Langmuir 1:1 model of BiaEvaluation software 4.1 (GE Healthcare).

[0263] In a separate experiment, the affinity of the interactions of Z variants to hFcRn (SEQ ID NO:1065) and mFcRn (SEQ ID NO:1070), respectively, was measured at both pH 6.0 and pH 7.4 on a Biacore 3000 instrument (GE Healthcare). hFcRn and mFcRn were produced essentially as described in Example 1 but using mouse 3T3 cells instead of human SKOV-3 cells for production of mFcRn, and immobilized on separate flow cells on a CM5 chip in acetate buffer at pH 4.65. The immobilization level was approximately 1000 RU for both receptors. A reference flow cell was created by activation and deactivation. 0.005% PCT pH 6.0 or 7.4 was used as running buffer and for dilution of the analytes. All analyses were performed at 25 °C. The affinity constants for the His₆-tagged Z variants Z07918 (SEQ ID NO:707), Z07960 (SEQ ID NO:710) and Z10193 (SEQ ID NO:708) were determined by injecting a dilution series from 1024 nM to 0.5 nM (pH 6.0) or from 10240 nM to 5 nM (pH 7.4). The affinities were derived using GraphPad Prism 5 software, using a one site binding saturation model.

[0264] AlphaLISA blocking assay: The potential of Z variants to inhibit binding of IgG to FcRn was analyzed in an AlphaLISA assay with an EnSpire multiplate reader 2300 (Perkin Elmer). Human IgG (Roactemra) was immobilized on AlphaLISA acceptor beads (Perkin Elmer, cat. no. 6772002) according to the manufacturer's recommendations. Stepwise serial dilutions 1:3 of His-tagged Z variants to final concentrations of 250 nM to 38 pM were made in a 384-well plate (Perkin Elmer, cat. no. G6005350) and incubated for 45 min with 10 nM biotinylated human FcRn (Biorbyt, cat. no. orb84388; biotinylated essentially as described in Example 2) in AlphaLISA buffer (Perkin Elmer, cat. no. AL000F) adjusted to pH 6.0 using HCl. IgG-coated Acceptor beads were added to a final concentration of 10 µM and incubated for 45 min. Finally, streptavidin coated Donor beads (Perkin Elmer, cat. no. 6772002) were added to a final concentration of 40 µg/ml and incubated for 30 min. All incubations were performed at RT in the dark. The plate was analyzed in the EnSpire instrument and the IC₅₀ values were calculated using GraphPad Prism 5.

Results

[0265] Cultivation and purification: The seventeen FcRn binding Z variants (SEQ ID NO: 707-

722 and SEQ ID NO:1059), constructed with an N-terminal His₆ tag, were produced in *E. coli*. The amount of IMAC-purified protein from approximately 2-5 g bacterial pellets, determined spectrophotometrically by measuring the absorbance at 280 nm, ranged from approximately 10 mg to 20 mg for the different FcRn binding Z variants. SDS-PAGE analysis of each final protein preparation showed that these predominantly contained the FcRn binding Z variant. The correct identity and molecular weight of each FcRn binding Z variant was confirmed by HPLC-MS analysis.

[0266] CD analysis: The CD spectra determined for six Z variants showed that each had an α -helical structure at 20 °C. This result was also verified in the variable temperature measurements, wherein melting temperatures (T_m) were determined (Table 2). A reversible folding was seen for the six Z variants when overlaying spectra measured before and after heating to 90 °C.

Table 2. Melting temperatures for a selection of Z variants.

Z variant	SEQ ID NO:	T _m (°C)
Z07909	719	56
Z07918	707	49
Z07930	712	56
Z07960	710	58
Z10109	709	61
Z10193	708	59

[0267] Biacore binding and kinetic analyses: The binding of seventeen Z variants to human FcRn and the dissociation at different pH were tested in a Biacore instrument by sequentially injecting each of the Z variants at pH 6.0 and either buffer pH 6.0 or pH 7.4 over a chip surface containing FcRn. The ligand immobilization level of the surface was 1668 RU human FcRn. The seventeen Z variants showed binding to FcRn at pH 6.0, and for all variants, faster off-rates were seen at pH 7.4 compared to pH 6.0. The result for IgG was similar, displaying a faster off-rate at pH 7.4. The variants Z07918 and Z10193 showed the slowest dissociation curves. Sensorgrams for a subset of variants and IgG are displayed in Figure 2 A-E.

Table 3. Biacore kinetic constants and affinities for hFcRn binding at pH 6.0.

Z variant	SEQ ID NO:	k _{on} (M ⁻¹ s ⁻¹)	k _{off} (s ⁻¹)	K _D (M)
Z07918	707	1.4 x 10 ⁶	0.022	1.6 x 10 ⁻⁸
Z10140	711	1.4 x 10 ⁶	0.12	8.6 x 10 ⁻⁸
Z10156	717	7.6 x 10 ⁵	0.28	3.7 x 10 ⁻⁷
Z10183	713	1.0 x 10 ⁶	0.13	1.3 x 10 ⁻⁷
Z10193	708	1.5 x 10 ⁶	0.033	2.2 x 10 ⁻⁸

[0268] The kinetic constants of five Z variants interacting with FcRn at pH 6.0 were determined (see Table 3). The immobilization level of the surface was 2015 RU human FcRn. For each Z variant, kinetic constants were calculated using a curve set of three injected concentrations.

[0269] Affinity (K_D) constants were also determined for His₆-tagged Z variants Z07918 (SEQ ID NO:707), Z07960 (SEQ ID NO:710) and Z10193 (SEQ ID NO:708) interacting with human and mouse FcRn at pH 6.0 and pH 7.4 (Table 4). For all three variants, K_D values were lower at pH 6.0 compared to pH 7.4.

Table 4. Biacore affinities for hFcRn and mFcRn at pH 6.0 and pH 7.4.

Z variant	SEQ ID NO:	K_D (M) hFcRn		K_D (M) mFcRn	
		pH 6.0	pH 7.4	pH 6.0	pH 7.4
Z07918	707	1.2×10^{-8}	$>5 \times 10^{-7}$	9.0×10^{-8}	$>5 \times 10^{-7}$
Z07960	710	5.0×10^{-8}	$>1 \times 10^{-6}$	3.5×10^{-7}	$>5 \times 10^{-6}$
Z10193	708	1.4×10^{-8}	$>5 \times 10^{-7}$	9.5×10^{-8}	$>5 \times 10^{-7}$

Table 5: Calculated IC₅₀ values from AlphaLISA blocking assay.

Z variant	SEQ ID NO of Z variant	IC ₅₀ (M)
Z07909	719	4.6×10^{-8}
Z07918	707	2.1×10^{-9}
Z07930	712	4.2×10^{-8}
Z07960	710	4.2×10^{-8}
Z10109	709	5.7×10^{-8}
Z10111	714	4.6×10^{-8}
Z10140	711	5.6×10^{-8}
Z10183	713	3.9×10^{-8}
Z10193	708	1.2×10^{-8}
Z13993	1059	1.3×10^{-7}
Z11948-G ₄ S-Z11948	1060	3.8×10^{-10}
Z11948-(G ₄ S) ₃ -Z11948	1060	4.1×10^{-10}

[0270] AlphaLISA blocking assay: The ability of seventeen His₆-tagged monomeric Z variants (SEQ ID NO:707-722 and SEQ ID NO: 059) and two dimeric variant, Z11948-G₄S-Z11948 and Z11948-(G₄S)₃-Z11948 to inhibit IgG binding to FcRn was tested in an AlphaLISA blocking assay. Serial dilutions of the Z variants were incubated with biotinylated human FcRn and the

blocking ability of each respective variant was measured after addition of IgG coated Acceptor beads and subsequently streptavidin coated Donor beads. Inhibition could be measured as a decrease in AlphaLISA counts for positive Z variants. The calculated IC50 values for the ten monomeric variants and the two dimeric variants that were shown to block IgG binding to FcRn in this assay are shown in Table 5.

Example 4

Binding of FcRn binding Z variants to human or mouse FcRn/eGFP transfected HeLa cells

[0271] In this example, the binding ability of FcRn binding Z variants was investigated. The production of HeLa cells expressing human and murine FcRn-eGFP gene transgene and the use of these cells for flow cytometry analysis with Alexa647 labeled Z variants is described.

Materials and methods

[0272] Cloning of FcRn-eGFP and B2M viral vectors: The genes encoding murine FcRn (mFcRn, Genbank BC003786.1, OpenBiosystems) and murine B2M (mB2M, Genbank BC085164.1, OpenBiosystems) were amplified in a similar way as the genes for human FcRn and human B2M as described in Example 1. Human and murine FcRn and B2M genes were amplified as follows: for hFcRn, the sequence encoding amino acids 1-365 (SEQ ID NO:1068) was amplified; for hB2M, the sequence encoding amino acids 21-119 (SEQ ID NO:1066) was amplified; for mFcRn, the sequence encoding amino acids 1-369 (SEQ ID NO:1069) was amplified; and for mB2M, the sequence encoding amino acids 21-119 (SEQ ID NO:1067) was amplified. The vector pHR-cPPT-CMV-EGFP (Jakobsson et al. (2003) J Neurosci Res 73:876-85) and FcRn PCR amplicons (human and murine) were cut using the restriction enzymes *Bam*HI (human) or *Bcl*I (murine) and *Mlu*I (New England Biolabs, cat. nos. R0136M, R0160L and R0198L, respectively), and ligated using T4 DNA Ligase (New England Biolabs, cat. no. M0202M). The ligation mix was chemically transformed into *E. coli* RRIΔM15 and spread on ampicillin plates. Colonies were picked and screened with suitable primer pairs. The construct encoding the original signal peptide, human or murine FcRn and eGFP at the cytoplasmic tail were verified by sequencing and denoted pHR-cPPT-CMV-hFcRn-eGFP and pHR-cPPT-CMV-mFcRn-eGFP, respectively.

[0273] The human and murine B2M PCR amplicons were inserted into the plasmid pDONOR221 (Invitrogen, cat. no. 12536-017) by recombination using the Gateway system (Invitrogen, cat. no. 11789020, Gateway® BP Clonase® II Enzyme mix) according to the manufacturer's recommendations. After verification of correct sequences, human or murine B2M was inserted into p2k7_gtc (Suter *et al.*, *supra*) using a multi-site gateway cloning system

(Invitrogen, cat. no. 11791020, Gateway® LR Clonase® II Enzyme mix) together with the promoter containing plasmid pENTR-CMV (Tai *et al. supra*), resulting in the vectors 2k7_{neo}-CMV-hB2M and 2k7_{neo}-CMV-mB2M, respectively.

[0274] Lentiviral transduction of HeLa cells: The vector pairs 2k7_{neo}-CMV-hB2M and pHR-cPPT-CMV-hFcRn-eGFP or 2k7_{neo}-CMV-mB2M and pHR-cPPT-CMV-mFcRn-eGFP were co-transfected together with VSV-G envelope and gag/pol packaging plasmid into HEK293T cells using calcium chloride transfection (Zufferey *et al., supra*; Jakobsson *et al. (2006) supra*). HEK293T culture supernatants containing formed lentiviral particles with FcRn and B2M transgenes respectively were used to sequentially transduce HeLa Cervix adenocarcinoma cells (Cell Line Service) at low passage number. The resulting two stably transduced HeLa cell lines are in the following denoted hFcRn-eGFP (transduced with genes for human FcRn-eGFP and hB2M) and mFcRn-eGFP (transduced with genes for mouse FcRn-eGFP and mB2M).

[0275] Alexa647 labeling of FcRn binding Z variants: The three His₆-tagged Z variants Z07918, Z07930 and Z07960 were labeled with Alexa Fluor® 647 Carboxylic Acid Succinimidyl Ester (Invitrogen cat. no. A20106). Before labeling, buffer was exchanged to 0.2 M carbonate buffer, pH 8.3, using Vivaspin500 centrifugal filter units (10 kDa MWCO, Vivaproducts cat. no. 512-2838) spun at 10,000 g. The labeling was performed in the Vivaspin500 and 1 µl of Alexa647 Succinimidyl Ester dye (40 µg/µl in DMSO corresponding to 1.3 x molar excess) was added to 200 µg/25 µl Z variant. The mixes were incubated at RT in the dark for 40 minutes in a wiggling rota mixer. The reaction mixes were subsequently put on ice for 3.5 hours and free dye was removed by washing with 15 x 100 µl PBS in the Vivaspin500.

[0276] Immunofluorescence staining of human and mouse FcRn-eGFP transfected HeLa-cells with FcRn binding Z variants: hFcRn-eGFP and mFcRn-eGFP HeLa cells were harvested by trypsination and washed twice in PBS at pH 6.0 before counting. 100,000 cells were pipetted per well of a v-bottomed 96 well plate (Nunc, cat no 277143) and the cells in the plate were subsequently pelleted at 1,700 rpm for 4 min at 4 °C. The supernatants were removed and the cells were fixed with 50 µl of 2 % formaldehyde (Sigma Aldrich, cat. no. F8775) in PBS at pH 6.0 for 10 min at RT. Cells were thereafter washed with 2 x 100 µl PBS pH 6.0, saturated with casein (PBSC), and resuspended in PBSC plus 0.1 % saponin (AppliChem, cat no A4518.0100) containing 620 nM of Alexa647 labeled His₆-tagged Z variants; Z07960, Z07930 and Z07918. Transduced HeLa cells, incubated with buffer alone, were used as control. The cells were incubated for 1 h at 8 °C on a shaker in the dark, washed with 2 x 100 µl PBSC and resuspended in 180 µl of PBS pH 6.0 plus 1 % BSA (fraction V, Merck, cat. no. 1.12018.0100). 10,000 cells/well were analyzed in a Gallios Flow Cytometer (Beckman Coulter) and the data was analyzed using Kaluza software (Beckman Coulter).

Results

[0277] Flow cytometry analysis was utilized to determine whether the FcRn binding Z variants

could bind to human and/or mouse FcRn on human or mouse FcRn-eGFP transduced HeLa cells. The experiment was performed at pH 6.0 with Alexa647 labeled Z07960, Z07930 and Z07918 (SEQ ID NO:710, 712 and 707, respectively). Dot plot analysis (y-axis: Alexa647, x-axis: eGFP) showed that the transduced cell population could be divided into FcRn-eGFP negative and positive population (Figure 3, gate H and I, respectively) indicating heterogeneous expression of the FcRn-eGFP fusion protein by HeLa cells (Figure 3). Accordingly, the mean fluorescence intensity (MFI) values for Alexa647 in gate I were subtracted by background MFI values of Alexa647 in gate H. The calculated MFI values are presented in Figure 4. The results show that Z07960, Z07930 and Z07918 are capable of binding HeLa cells displaying human (Figure 4A) or murine (Figure 4B) FcRn-eGFP.

Example 5

Blocking of IgG binding to FcRn with the FcRn binding Z variant Z07918

[0278] In this example, the potential competition of FcRn binding Z variants with IgG for binding to FcRn was investigated in a cell based assay. Such binding will result in blocking of the IgG-FcRn interaction.

Materials and methods

[0279] Blocking of IgG-FcRn immunofluorescence staining: Human or murine FcRn-eGFP transduced HeLa cells were prepared as described in Example 4. Fixed cells were resuspended in 50 µl of a mix of either 100 nM Alexa647-conjugated human or mouse IgG (Jackson laboratories, cat. no. 009-600-003 and 015-600-003, respectively) and 1000, 100, 10, 1 or 0 (buffer control) nM His₆-tagged Z07918 diluted in PBS-casein, pH 6.0, plus 0.1 % saponin (AppliChem). The cells were incubated for 1 h at 37 °C on a shaker in the dark, washed with 2 x 100 µl PBS-casein pH 6.0 and re-suspended in 180 µl of PBS, pH 6.0, plus 1 % BSA. Data from 10,000 cells/well (except somewhat fewer cells for mouse 100 nM mIgG-Alexa647) were obtained using a Gallios Flow Cytometer (Beckman Coulter) and the data was analyzed using Kaluza software (Beckman Coulter).

Results

[0280] The experiment was performed to determine if the FcRn binding Z variant Z07918 (SEQ ID NO:707) blocks the IgG-FcRn interaction. Human or murine FcRn-eGFP transduced HeLa cells were incubated with human or mouse Alexa647-conjugated IgG. The binding was blocked with unlabeled Z07918 at different concentrations. Due to the heterogeneous expression of FcRn by the transduced HeLa cells (described in Example 4), the MFI values for

Alexa647 in gate N of each sample was subtracted by the corresponding MFI values in gate M (Figure 5). The percent IgG Alexa647 binding was calculated by dividing the different MFI values with the MFI for the blank control. The results showed that Z07918 effectively blocked hIgG binding to hFcRn (Figure 6A) in a dose dependent manner. Furthermore, Z07918 also blocked mIgG binding to mFcRn (Figure 6B) although less efficiently compared to hIgG-binding.

Example 6

Pharmacokinetic study of three FcRn binding Z variants

[0281] In this example, the ability of FcRn binding Z variants to prolong serum half-life of a non-specific Z variant was investigated by a pharmacokinetic study performed in mice.

Materials and methods

[0282] Subcloning of Z variants: A subset of Z variants (Z07918, Z07960 and Z10193) was submitted to a second subcloning. DNA from the subcloned His₆-tagged variants in Example 3 was used as template. First, PCR amplification using suitable primer pairs was performed to create genes encoding Z variants starting with the amino acids AE instead of VD. The mutated Z variants are listed in Figure 1 and were denoted Z11948 (SEQ ID NO:1060), Z11946 (SEQ ID NO:1061) and Z11947 (SEQ ID NO:1062), corresponding to mutated Z07918, Z07960 and Z10193, respectively. Genes encoding the new Z variants were restriction cleaved and ligated into a vector harboring the genes encoding albumin binding variant PP013 (SEQ ID NO:1063) and Z03638 (SEQ ID NO:1064) with spacer sequences resulting in a gene fusion encoding [Z#####]-GAP(G₄S)₄TS-[PP013]-GT(G₄S)₄PR-[Z03638] (also denoted "Z#####-PP013-Z03638" or "Z variant in fusion with PP013-Z03638"). The negative control molecule [Z03638]-GAP(G₄S)₄TS-[PP013] was subcloned in a similar way by ligating Z03638 into a vector containing a (G₄S)₄ linker and the sequence for PP013. The subsequent steps for vector transformation into *E. coli* were performed as in Example 3.

[0283] Cultivation and purification: Z variants in fusion with PP013-Z03638 were produced in *E. coli* as described in Example 3. Approximately 3 g of each cell pellet was re-suspended in 30 ml TST-buffer (25 mM Tris-HCl, 1 mM EDTA, 200 mM NaCl, 0.05 % Tween20, pH 8.0) supplemented with Benzonase® (Merck). After cell disruption by sonication and clarification by centrifugation, each supernatant was applied on a gravity flow column with 5 ml agarose immobilized with an anti-ABD ligand (produced in-house). After washing with TST-buffer and 5 mM NH₄Ac buffer, pH 5.5, the Z variants were eluted with 0.1 M HAc. Acetonitrile (ACN) was added to a final concentration of 10 % to the eluted fractions from the anti-ABD agarose affinity

chromatography purification step and the samples were loaded on a 3 ml Resource 15RPC column (GE Healthcare), previously equilibrated with RPC solvent A (0.1 % trifluoroacetic acid (TFA), 10 % ACN, 90 % water). After column wash with RPC solvent A, bound protein was eluted with a linear gradient 0-50 % RPC solvent B (0.1 % TFA, 80 % ACN, 20 % water) during 60 ml. Fractions containing pure Z variant were identified by SDS-PAGE analysis and pooled. After the RPC purification, the buffer of the pools was exchanged to PBS using a HiPrep 26/10 Desalting column (GE Healthcare). Finally, the Z variants were purified on 1 ml EndoTrap red columns (Hyglos, cat. no. 321063) to ensure low endotoxin content.

[0284] Protein concentrations, purities and the identity of each purified Z variant were analyzed as described in Example 3.

[0285] Biacore analysis: Expressed and purified Z variants fused to PP013-Z03638 were assayed against human FcRn at pH 6.0 essentially as described for the kinetic analysis in Example 3. The Z variants and the negative control Z03638-PP013 were injected at 40 nM, 160 nM and 640 nM during 1 minute followed by dissociation for 2.5 minutes and equilibration for 1 minute. Kinetic constants and affinities were determined for the Z variants using the BiaEvaluation software.

[0286] Pharmacokinetic study: Z11947, Z11946 and Z11948 fused to PP013-Z03638 were administered intravenously (i.v.) to male NMRI mice (Charles River, Germany) at a dose of 92 nmol/kg body weight. Sera from groups of three mice were obtained at 0.08, 6, 18, 78, 120, 168 and 240 hours. The concentration of respective Z variant was determined by ELISA.

[0287] ELISA: Half-area 96-well ELISA plates were coated at 4 °C overnight with 50 µl/well of an Z specific goat antibody (produced in-house) diluted to 4 µg/ml in coating buffer (50 mM sodium carbonate, pH 9.6). The antibody solution was poured off and the wells were blocked with 100 µl of PBSC for 1.5 h at RT. The sera were diluted in PBSC plus 1 % mouse serum (matrix) from 1:100 to 1:51,200 in a two-fold dilution series in a dilutions plate. A standard titration for respective Z variant and four quality controls (very low, low, medium and high control) diluted in matrix were included on each plate. 50 µl of the dilutions were transferred per well and the ELISA plates were incubated for 1.5 h at RT. The plates were washed four times with PBST. Bound Z variants were detected with 50 µl/well of rabbit anti-PP013 Ig (produced in-house) diluted to 4 µg/ml in PBSC. The plates were subsequently incubated for 1.5 h at RT followed by washes as described above. HRP conjugated donkey anti-rabbit HRP obtained from Jackson laboratories (cat. no. 711-035-152), diluted 1:20,000 in PBSC, was added and the plates were incubated for 1 hour. After washing as described above, 50 µl of ImmunoPure TMB substrate was added to the wells and the plates were developed according to the manufacturer's recommendations. After 15 minutes of development, the absorbance was measured at 450 nm using a multi-well plate reader (Victor³). The absorbance values were analyzed using GraphPad Prism 5 to determine the concentrations (cubic-spline curve fit) and area under curve (AUC). The concentrations were then plotted as their natural logarithms against time. The resulting curves followed a two compartment model and the terminal half-life was calculated as $\ln 2$ divided by the slope based on the last three time points.

Results

[0288] Cultivation and purification: The three FcRn binding Z variants Z11947, Z11946 and Z11948 (SEQ ID NO:1062, 1061 and 1060), constructed as Z#####-PP013-Z03638, and the negative control Z03638-PP013, were produced in *E. coli*. The amount of purified protein from approximately 3 g bacterial pellets, determined spectrophotometrically by measuring the absorbance at 280 nm, ranged from approximately 10 to 25 mg for the different FcRn binding Z variants. SDS-PAGE analysis of each final protein preparation showed that they predominantly contained respective FcRn binding Z variant. The correct molecular weight of each FcRn binding Z variant was confirmed by LC/MS analysis.

Table 6. Kinetic constants and affinities for FcRn at pH 6.0 of Z variants produced as fusions to PP013-Z03638.

Z variant	SEQ ID NO:	$k_{on} (M^{-1}s^{-1})$	$k_{off} (s^{-1})$	$K_D (M)$
Z11948	1060	7.73×10^5	0.047	6.2×10^{-8}
Z11946	1061	3.35×10^5	0.275	8.2×10^{-7}
Z11947	1062	6.54×10^5	0.064	9.8×10^{-8}

[0289] Biacore analysis: The binding to FcRn was analyzed for the three PP013-Z03638 fused Z variants. The immobilization level of the surface was 548 RU of human FcRn. The resulting rough kinetic constants and affinities for the target binding at pH 6.0 are displayed in Table 6. Fitted curves are displayed in Figure 7A-C. The negative control Z03638-PP013 was negative against FcRn.

[0290] Pharmacokinetic study: The pharmacokinetic profiles of the above-mentioned constructs of Z variants fused to PP013-Z03638 were compared to the negative control Z03638-PP013 in a mouse pharmacokinetic study. In previous work, e.g. as described in PCT application WO2009/016043, it is shown that ABD fusion proteins have a long half-life in serum, caused by ABD binding to serum albumin. In accordance with the previous results, terminal half-life of ABD-fused Z variant molecule (Z03638-PP013) was approximately 43 hours, which is comparable to half-life of mouse albumin (35 hours). The terminal half-lives of the constructs containing FcRn binding Z variant molecule in addition to ABD were two- to three-fold longer (Figure 8). The calculated terminal half-lives were 99 hours (Z11947), 69 hours (Z11946) and 58 hours (Z11948), suggesting that FcRn binding of the Z variants contributed to the prolonged half-life.

Example 7

Design and construction of a maturation library of FcRn binding Z variants

[0291] In this Example, a matured library was constructed. The library was used for selections of FcRn binding Z variants. Selections from matured libraries are usually expected to result in binders with increased affinity (Orlova et al., (2006) Cancer Res 66(8):4339-48). In this study, randomized single stranded linkers were generated using split-pool synthesis enabling incorporation of defined codons in desired positions in the synthesis.

Materials and methods

[0292] Library design: The library was based on the sixteen sequences of the human FcRn binding Z variants in Table 1 and further described in Examples 2-6. In the new library, 13 variable positions in the Z molecule scaffold were biased towards certain amino acid residues, according to a strategy mainly based on the binding motifs of the Z variants defined in SEQ ID NO:707-722. A DNA linker was generated using split-pool synthesis containing the 147 bp partially randomized helix 1 and 2 of the amino acid sequence: 5'- AAATAAT CTC GAG GTA GAT GCC AAA TAC GCC AAA GAA NNN NNN NNN GCG NNN NNN GAG ATC NNN NNN TTA CCT AAC TTA ACC NNN NNN CAA NNN NNN GCC TTC ATC NNN AAA TTA NNN GAT GAC CCAAGC CAG AGC TCA TTA TTT A-3' (SEQ ID NO:1074; randomized codons are illustrated as NNN) flanked by restriction sites *Xho*I and *Sac*I, was ordered from DNA 2.0 (Menlo Park, CA, USA). The theoretical distributions of amino acid residues in the new library, including eight variable amino acid positions (9, 10, 11, 13, 14, 24, 32 and 35) and five constant amino acid positions (17, 18, 25, 27 and 28) in the Z molecule scaffold are given in Table 8. The resulting theoretical library size is 5.3×10^8 variants.

Table 7: Design of library for maturation.

Amino acid position in the Z variant	Randomization (amino acid abbreviations)	No of amino acids	Proportion
9	A,D,E,F,H,I,K,L,N,Q,R,S,T,V,W,Y	16	1/16
10	A,D,E,F,H,I,K,L,M,N,Q,R,S,T,V,W,Y	17	1/17
11	A,D,E,F,H,I,K,L,N,Q,R,S,T,V,W,Y	16	1/16
13	A,D,E,F,G,H,I,K,L,N,Q,R,S,T,V,W,Y	17	1/17
14	A,F,H (25 %),I,K,L,N,Q,R,S,T,V,W,Y	14	3/52, 13/52 (H)
17	R	1	1
18	W	1	1
24	F,Y	2	1/2
25	D	1	1
27	R	1	1

Amino acid position in the Z variant	Randomization (amino acid abbreviations)	No of amino acids	Proportion
28	V	1	1
32	A,D,E,F,H,I,K,L,N,Q,R,S,T,V,W,Y	16	1/16
35	A,D,E,F,H,I,K,L,N,Q,R,S,T,V,W,Y	16	1/16

[0293] Library construction: The library was amplified using *AmpliTaq* Gold polymerase (Applied Biosystems, cat. no. 4311816) during 12 cycles of PCR and pooled products were purified with QIAquick PCR Purification Kit (QIAGEN, cat. no. 28106) according to the supplier's recommendations. The purified pool of randomized library fragments was digested with restriction enzymes *Xho*I and *Sac*I-HF (New England Biolabs, cat. no. R0146L, and cat. no. R3156M) and concentrated using a PCR Purification Kit. Subsequently, the product was subjected to preparative 2.5 % agarose (Nuisieve GTC agarose, Cambrex, Invitrogen) gel electrophoresis and purified using QIAGEN gel extraction Kit (QIAGEN, cat. no. 28706) according to the supplier's recommendations.

[0294] The phagemid vector pAY02592 (essentially as pAffi1 described in Grönwall *et al.*, *supra*) was restricted with the same enzymes, purified using phenol/chloroform extraction and ethanol precipitation. The restricted fragments and the restricted vector were ligated in a molar ratio of 5:1 with T4 DNA ligase (Fermentas, cat. no. EL0011) for 2 hours at RT, followed by overnight incubation at 4 °C. The ligated DNA was recovered by phenol/chloroform extraction and ethanol precipitation, followed by dissolution in 10 mM Tris-HCl, pH 8.5. Thus, the resulting library in vector pAY02592 encoded Z variants, each fused to an albumin binding domain (ABD) derived from streptococcal protein G.

[0295] The ligation reactions (approximately 160 ng DNA/transformation) were electroporated into electrocompetent *E. coli* ER2738 cells (50 µl, Lucigen, Middleton, WI, USA). Immediately after electroporation, approximately 1 ml of recovery medium (supplied with the ER2738 cells) was added. The transformed cells were incubated at 37 °C for 60 min. Samples were taken for titration and for determination of the number of transformants. The cells were thereafter pooled and cultivated overnight at 37 °C in 1 l of TSB-YE medium, supplemented with 2 % glucose, 10 µg/ml tetracycline and 100 µg/ml ampicillin. The cells were pelleted for 7 min at 4,000 g and resuspended in a PBS/glycerol solution (approximately 40 % glycerol). The cells were aliquoted and stored at -80 °C. Clones from the library of Z variants were sequenced in order to verify the content and to evaluate the outcome of the constructed library vis-à-vis the library design. Sequencing was performed as described in Example 1 and the amino acid distribution was verified.

[0296] Preparation of phage stock: Phage stock containing the phagemid library was prepared in a 20 l fermenter (Belach Bioteknik). Cells from a glycerol stock containing the phagemid library were inoculated in 10 l of TSB-YE (Tryptic Soy Broth-Yeast Extract; 30 g/l TSB, 5 g/l

yeast extract) supplemented with 1 g/l glucose, 100 mg/l ampicillin and 10 mg/l tetracycline. When the cells reached an optical density at 600 nm (OD₆₀₀) of 0.6, approximately 1.5 l of the cultivation was infected using a 5 x molar excess of M13K07 helper phage. The cells were incubated for 30 min, whereupon the fermenter was filled up to 10 l with complex fermentation medium [2.5 g/l (NH₄)₂SO₄; 5.0 g/l yeast extract; 30 g/l tryptone, 2 g/l K₂HPO₄; 3 g/l KH₂PO₄, 1.25 g/l; Na₃C₆H₅O₇ • 2 H₂O; Breox FMT30 antifoaming agent 0.1 ml/l]. The following components were added: 10 ml carbenicillin 25 mg/ml; 5 ml kanamycin 50 mg/ml; 1 ml 1 M isopropyl-β-D-1-thiogalactopyranoside (IPTG); 17.5 ml/l of 300 g/l MgSO₄, and 5 ml of a trace element solution [35 g/l FeCl₃ • 6 H₂O; 10.56 g/l ZnSO₄ • 7 H₂O; 2.64 g/l CuSO₄ 5 H₂O; 13.2 g/l MnSO₄ • H₂O; 13.84 g/l CaCl₂ 2 H₂O, dissolved in 1.2 M HCl]. A glucose limited fed-batch cultivation was started where a 600 g/l glucose solution was fed to the reactor (3.5 g/h in the start, 37.5 g/h after 20 h and until the end of the cultivation). pH was controlled at pH 7 through the automatic addition of 25 % NH₄OH, air was supplemented (5 l/min), and the stirrer was set at 500 rpm. After 24 h of fed-batch cultivation the OD₆₀₀ was 33.2. The cells in the cultivation were pelleted by centrifugation at 15,900 g. The phage particles were precipitated from the supernatant twice in PEG/NaCl, filtered and dissolved in PBS and glycerol as in Example 2. Phage stocks were stored at -80 °C until use in selection.

Results

[0297] Library construction: The new library was designed based on a set of 16 FcRn binding Z variants with verified binding properties (Example 2-6). The theoretical size of the designed library was 5.3×10^8 Z variants. The actual size of the library, determined by titration after transformation to *E. coli* ER2738 cells, was 4.5×10^9 transformants.

[0298] The library quality was tested by sequencing of 96 transformants and by comparing their actual sequences with the theoretical design. The contents of the actual library compared to the designed library were shown to be satisfying. A matured library of potential binders to FcRn was thus successfully constructed.

Example 8

Selection and screening of Z variants from a matured library

Materials and methods

[0299] Phage display selection of matured FcRn binding Z variants: The target proteins

human FcRn (Biorbyt, cat. no. orb84388) and murine FcRn (Biorbyt, cat. no. orb99076) were biotinylated essentially as described in Example 2 using biotin at 10x molar excess. Phage display selections, using the new library of Z variant molecules described in Example 7, were performed in four cycles against human FcRn or murine FcRn essentially as in Example 2 but with the following exceptions. Selection buffers were 0.1% PCTG buffer, pH 5.5 (Mcllvaines phosphate-citrate buffer, pH 5.5, supplemented with 0.1 % Tween-20 and 0.1% gelatin) or 0.1% PCTG buffer, pH 7.4, (Mcllvaines phosphate-citrate buffer, pH 7.4, supplemented with 0.1 % Tween-20 and 0.1% gelatin) respectively. Prior to selection, HSA was added to the selection buffers to a final concentration of 1.5 μ M. All tubes and beads used in the selection were pre-blocked with either of the two different selections buffers. A pre-selection step, by incubation of phage stock with SA-beads for 45 min, was performed in cycle 1. For capture of phage-target complexes, 1 mg beads per 1.1 μ g biotinylated human FcRn or 1.6 μ g biotinylated murine FcRn was used. Washes were performed with 0.1% PCT buffer pH 5.5 or pH 7.4 except for tracks 2-1-2-1 and 2-1-2-2 where 0.1% PCT supplemented with 25 nM IgG (Herceptin®) or 10 nM IgG, respectively, was used as outlined in Table 7.

[0300] The five tracks (1-5) in cycle 1 were divided in the second to fourth cycles, resulting in totally seven tracks (1-1 to 5-1) in cycle 2, eleven tracks (1-1-1 to 5-1-1) in cycle 3 and fourteen tracks (1-1-1-1 to 5-1-1-1) in cycle 4. The bound phage particles were eluted as described in Example 2.

[0301] An overview of the selection strategy, describing an increased stringency in subsequent cycles, using a lowered target concentration and an increased number of washes, is shown in Table 8.

Table 8. Overview of the maturation selection data.

Cycle	Selection track	Phage stock from library or selection track	Target species	Target conc. (nM)	Selection pH	Wash pH	Number of washes
1	1	Zlib006FcRn.I	human	100	7.4	7.4	2
1	2	Zlib006FcRn.I	human	100	7.4	5.5	2
1	3	Zlib006FcRn.I	human	25	5.5	5.5	4
1	4	Zlib006FcRn.I	murine	100	7.4	7.4	2
1	5	Zlib006FcRn.I	murine	100	5.5	5.5	2
2	1-1	1	human	50	7.4	7.4	4
2	2-1	2	human	50	7.4	5.5	4
2	2-2	2	human	25	5.5	7.4	6
2	3-1	3	human	5	5.5	7.4	4
2	3-2	3	human	5	5.5	5.5	8
2	4-1	4	murine	50	7.4	5.5	2
2	5-1	5	murine	100	5.5	5.5	2
3	1-1-1	1-1	human	10	7.4	7.4	8

Cycle	Selection track	Phage stock from library or selection track	Target species	Target conc. (nM)	Selection pH	Wash pH	Number of washes
3	1-1-2	1-1	human	5	5.5	7.4	8
3	2-1-1	2-1	human	10	7.4	5.5	8
3	2-1-2	2-1	human	5	7.4	5.5	12
3	2-2-1	2-2	human	10	7.4	5.5	12
3	2-2-2	2-2	human	5	7.4	5.5	15
3	3-1-1	3-1	human	1	5.5	7.4	8
3	3-2-1	3-2	human	0.5	5.5	5.5	12
3	3-2-2	3-2	human	0.25	5.5	5.5	16
3	4-1-1	4-1	murine	10	7.4	5.5	6
3	5-1-1	5-1	murine	5	5.5	5.5	8
4	1-1-1-1	1-1-1	human	1	7.4	7.4	12
4	1-1-1-2	1-1-1	human	0.25	7.4	7.4	15
4	1-1-2-1	1-1-2	human	0.5	7.4	5.5	15
4	1-1-2-2	1-1-2	human	0.1	5.5	7.4	15
4	2-1-1-1	2-1-1	human	1	7.4	5.5	15
4	2-1-1-2	2-1-1	human	0.5	7.4	5.5	15
4	2-1-2-1	2-1-2	human	0.25	7.4	5.5	20 (+IgG)
4	2-1-2-2	2-1-2	human	0.1	7.4	5.5	20 (+IgG)
4	2-2-1-1	2-2-1 and 2-2-2	human	0.5	5.5	7.4	15
4	2-2-2-1	2-2-1 and 2-2-2	human	0.5	7.4	5.5	20
4	3-1-1-1	3-1-1	human	1	5.5	7.4	12
4	3-2-1-1	3-2-1 and 3-2-2	human	0.5	5.5	5.5	16
4	4-1-1-1	4-1-1	murine	1	7.4	5.5	12
4	5-1-1-1	5-1-1	murine	0.5	5.5	5.5	15

[0302] Amplification of phage particles: Amplification of phage particles between selection cycle 1 and 2 was performed essentially as described in Example 2, with the following exceptions. *E. coli* ER2738 was used for phage amplification and M13K07 helper phage was used in 5 x excess. The amplification of phage particles between the selection cycles 2 and 4 was done by performing infection of bacteria in solution as follows. After infection of log phase *E. coli* ER2738 with phage particles, TSB supplemented with 2 % glucose, 10 µg/ml tetracycline and 100 µg/ml ampicillin was added, followed by incubation with rotation for 30 min at 37 °C. Thereafter, the bacteria were infected with M13K07 helper phage in 5 x excess. The infected

bacteria were pelleted by centrifugation, re-suspended in TSB-YE medium supplemented with 100 μ M IPTG, 25 μ g/ml kanamycin and 100 μ g/ml ampicillin, and grown overnight at 30 °C. The overnight cultures were pelleted in a centrifuge, and phage particles in the supernatant were precipitated twice with PEG/NaCl buffer. Finally, the phage particles were re-suspended in selection buffer before entering the next selection cycle.

[0303] In the final selection cycle, log phase bacteria were infected with eluate and diluted before spreading onto TBAB plates (30 g/l tryptose blood agar base, Oxoid cat. no. CMO233B) supplemented with 0.2 g/l ampicillin in order to form single colonies to be used in ELISA screening.

[0304] Sequencing of potential binders: Individual clones from the different selection tracks were picked for sequencing. All clones run in the ELISA screening were sequenced. Amplification of gene fragments and sequence analysis of gene fragments were performed essentially as described in Example 2.

[0305] ELISA screening of Z variants: Single colonies containing Z variants (expressed as Z variant ABD fusion proteins as described in Example 2) were randomly picked from the selected clones of the FcRn matured library and grown in 1 ml cultivations essentially as described in Example 2. Preparation of the periplasmic supernatants was performed as in Example 2 with eight freeze thawing cycles and the periplasmic fractions were used undiluted in the ELISA screening. ELISA screenings were performed at both pH 6.0 and pH 7.4 essentially as described in Example 2 using biotinylated human FcRn at a concentration of 2 nM in each well. The periplasmic fraction of the primary FcRn binder Z10193 (SEQ ID NO:708; assayed in above experiments) was used as a positive control. Periplasm containing the ABD moiety only was used as a negative control.

[0306] ELISA K_D analysis of FcRn binding Z variants: A selection of FcRn binders was subjected to an analysis of the response against a dilution series of biotinylated human FcRn using ELISA at both pH 6.0 and pH 7.4 as described above. Biotinylated human FcRn was added at a concentration of 30 nM and diluted stepwise 1:3 down to 14 pM. As a background control, all Z variants were also assayed with no target protein added. Periplasm samples containing the primary FcRn binder Z07918 (SEQ ID.NO:707) was included and analyzed as a positive control. Periplasm containing the ABD moiety only was used as a negative control. Data were analyzed using GraphPad Prism 5 and non-linear regression and K_D values (the half maximal effective concentration) were calculated.

Results

Phage display selection of matured FcRn binding Z variants:

[0307] Selection was performed in totally 14 parallel tracks containing four cycles each. The different selection tracks differed in target concentration, target type (human FcRn or murine FcRn), selection time, and wash conditions.

[0308] Sequencing of potential binders: Randomly picked clones were sequenced. Each individual Z variant was given an identification number, Z#####, as described in Example 2. In total, 445 new unique Z variant molecules were identified.

[0309] The amino acid sequences of the 58 amino acid residues long Z variants are listed in Figure 1 and in the sequence listing as SEQ ID NO:723-1058. The deduced FcRn binding motifs of these Z variants are listed in Figure 1 and in the sequence listing as SEQ ID NO:17-352. The amino acid sequences of the 49 amino acid residues long polypeptides predicted to constitute the complete three-helix bundle within each of these Z variants are listed in Figure 1 and in the sequence listing as SEQ ID NO:370-705.

[0310] ELISA screening of Z variants: Clones obtained after four selection cycles were produced in 96-well plates and screened for FcRn binding activity using ELISA. All randomly picked clones were analyzed. At pH 6.0, 333 of the 445 unique Z variants were found to give a response of 0.3 AU or higher (corresponding to at least 3x the negative control) against human FcRn at a concentration of 2 nM. At pH 7.4, 278 of the 445 unique Z variants were found to give a response of 0.3 AU or higher (corresponding to at least 3x the negative control) against human FcRn at a concentration of 2 nM. Clones with a positive signal against human FcRn were found in all tracks (including those with murine target) except 1-1-1-1. The negative controls had absorbances of 0.070-0.096 AU (pH 6.0) and 0.060-0.112 AU (pH 7.4), respectively. The average response of the blank controls was 0.070 AU (pH 6.0) and 0.062 (pH 7.4).

[0311] ELISA K_D analysis of FcRn binding Z variants: A subset of Z variants was selected based on the result in the ELISA experiment described above (highest ELISA value at pH 6.0 and/or pH 7.4) and subjected to a target titration in ELISA format. Periplasm samples were incubated with a serial dilution of biotinylated human FcRn. A periplasm sample with the primary binder Z07918 (SEQ ID NO:707) was also assayed as a positive control. Obtained values were analyzed and their respective K_D values were calculated (Table 9).

Table 9: Calculated K_D values from ELISA titration analysis of Z-ABD variants from the maturation.

Z variant	SEQ ID NO:	K_D pH 6.0 (M)	K_D pH 7.4 (M)	Z variant	SEQ ID NO:	K_D pH 6.0 (M)	K_D pH 7.4 (M)
Z13573	723	1.1×10^{-9}	3.8×10^{-9}	Z13684	785	1.0×10^{-9}	2.2×10^{-9}
Z13574	724	1.2×10^{-9}	5.0×10^{-9}	Z13688	786	1.3×10^{-9}	2.1×10^{-9}
Z13577	725	9.9×10^{-10}	1.4×10^{-9}	Z13691	787	1.8×10^{-9}	2.7×10^{-9}

Z variant	SEQ ID NO:	K _D pH 6.0 (M)	K _D pH 7.4 (M)	Z variant	SEQ ID NO:	K _D pH 6.0 (M)	K _D pH 7.4 (M)
Z13578	726	1.0×10^{-9}	2.5×10^{-9}	Z13692	788	1.3×10^{-9}	3.7×10^{-9}
Z13579	727	1.2×10^{-9}	5.3×10^{-9}	Z13694	789	9.8×10^{-9}	3.6×10^{-9}
Z13581	728	1.1×10^{-9}	3.3×10^{-9}	Z13695	790	1.8×10^{-9}	5.3×10^{-9}
Z13583	729	8.0×10^{-10}	1.5×10^{-9}	Z13697	791	1.2×10^{-9}	2.4×10^{-9}
Z13585	730	1.2×10^{-9}	1.7×10^{-9}	Z13706	792	2.0×10^{-9}	6.4×10^{-9}
Z13586	731	1.2×10^{-9}	2.3×10^{-9}	Z13708	793	1.9×10^{-9}	4.4×10^{-9}
Z13587	732	1.4×10^{-9}	6.9×10^{-9}	Z13710	794	1.6×10^{-9}	2.6×10^{-9}
Z13588	733	1.0×10^{-9}	2.3×10^{-9}	Z13711	795	2.1×10^{-9}	4.9×10^{-9}
Z13592	734	9.5×10^{-10}	1.8×10^{-9}	Z13714	796	2.1×10^{-9}	6.0×10^{-9}
Z13594	735	1.3×10^{-9}	6.3×10^{-9}	Z13716	797	1.8×10^{-9}	5.8×10^{-9}
Z13596	736	1.5×10^{-9}	3.6×10^{-9}	Z13719	798	2.6×10^{-9}	7.3×10^{-9}
Z13597	737	1.4×10^{-9}	6.0×10^{-9}	Z13720	799	2.5×10^{-9}	4.5×10^{-7}
Z13598	738	1.1×10^{-9}	1.7×10^{-9}	Z13721	800	1.9×10^{-9}	2.9×10^{-9}
Z13600	739	1.4×10^{-9}	4.0×10^{-9}	Z13725	801	1.8×10^{-9}	4.9×10^{-9}
Z13604	740	1.3×10^{-9}	4.1×10^{-9}	Z13727	802	2.1×10^{-9}	5.9×10^{-9}
Z13605	741	1.3×10^{-9}	3.8×10^{-9}	Z13728	803	2.6×10^{-9}	6.7×10^{-9}
Z13609	742	1.3×10^{-9}	2.7×10^{-9}	Z13732	804	2.1×10^{-9}	9.4×10^{-9}
Z13611	743	1.3×10^{-9}	2.5×10^{-9}	Z13735	805	1.6×10^{-9}	9.1×10^{-9}

Z variant	SEQ ID NO:	K _D pH 6.0 (M)	K _D pH 7.4 (M)	Z variant	SEQ ID NO:	K _D pH 6.0 (M)	K _D pH 7.4 (M)
Z13612	744	1.2×10^{-9}	8.6×10^{-9}	Z13736	806	1.7×10^{-9}	3.0×10^{-9}
Z13613	745	1.2×10^{-9}	4.3×10^{-9}	Z13740	807	2.0×10^{-9}	5.0×10^{-9}
Z13615	746	1.2×10^{-9}	3.1×10^{-9}	Z13742	808	2.4×10^{-9}	7.6×10^{-9}
Z13616	747	9.6×10^{-10}	1.7×10^{-9}	Z13747	809	1.3×10^{-9}	2.3×10^{-9}
Z13617	748	1.2×10^{-9}	1.9×10^{-9}	Z13749	810	2.8×10^{-9}	1.2×10^{-8}
Z13620	749	1.4×10^{-9}	3.3×10^{-9}	Z13750	811	2.7×10^{-9}	8.4×10^{-9}
Z13621	750	8.6×10^{-10}	1.4×10^{-9}	Z13751	812	2.0×10^{-9}	3.8×10^{-9}
Z13622	751	1.1×10^{-9}	2.1×10^{-9}	Z13752	813	2.0×10^{-9}	5.8×10^{-9}
Z13624	752	1.3×10^{-9}	3.4×10^{-9}	Z13758	814	1.9×10^{-9}	6.5×10^{-9}
Z13625	753	1.3×10^{-9}	2.8×10^{-9}	Z13759	815	2.1×10^{-9}	5.6×10^{-9}
Z13626	754	1.2×10^{-9}	2.7×10^{-9}	Z13760	816	2.1×10^{-9}	5.8×10^{-9}
Z13627	755	1.2×10^{-9}	2.9×10^{-9}	Z13761	817	1.9×10^{-9}	3.7×10^{-9}
Z13628	756	1.3×10^{-9}	5.5×10^{-9}	Z13771	818	1.5×10^{-9}	2.0×10^{-9}
Z13629	757	1.2×10^{-9}	8.5×10^{-9}	Z13773	819	2.5×10^{-9}	4.9×10^{-9}
Z13633	758	1.5×10^{-9}	6.2×10^{-9}	Z13776	820	2.2×10^{-9}	5.5×10^{-9}
Z13634	759	1.1×10^{-9}	2.3×10^{-9}	Z13777	821	2.4×10^{-9}	4.6×10^{-9}
Z13635	760	1.0×10^{-9}	1.7×10^{-9}	Z13780	822	2.1×10^{-9}	4.0×10^{-9}
Z13637	761	1.3×10^{-9}	4.8×10^{-9}	Z13782	823	2.2×10^{-9}	4.2×10^{-9}

Z variant	SEQ ID NO:	K _D pH 6.0 (M)	K _D pH 7.4 (M)	Z variant	SEQ ID NO:	K _D pH 6.0 (M)	K _D pH 7.4 (M)
Z13638	762	1.2×10^{-9}	2.9×10^{-9}	Z13783	824	1.4×10^{-9}	2.2×10^{-9}
Z13639	763	1.3×10^{-9}	3.0×10^{-9}	Z13786	825	2.3×10^{-9}	4.7×10^{-9}
Z13640	764	1.1×10^{-9}	1.9×10^{-9}	Z13792	826	2.0×10^{-9}	2.9×10^{-9}
Z13641	765	1.1×10^{-9}	1.8×10^{-9}	Z13796	827	2.3×10^{-9}	4.2×10^{-9}
Z13644	766	1.3×10^{-9}	2.8×10^{-9}	Z13799	828	1.9×10^{-9}	5.6×10^{-9}
Z13645	767	1.2×10^{-9}	2.5×10^{-9}	Z13806	829	1.6×10^{-9}	3.1×10^{-9}
Z13648	768	1.6×10^{-9}	3.3×10^{-9}	Z13808	830	2.4×10^{-9}	5.5×10^{-9}
Z13651	769	1.2×10^{-9}	2.7×10^{-9}	Z13811	831	2.0×10^{-9}	3.1×10^{-9}
Z13652	770	1.4×10^{-9}	2.9×10^{-9}	Z13812	832	2.3×10^{-9}	1.1×10^{-8}
Z13654	771	9.5×10^{-10}	2.9×10^{-9}	Z13823	833	2.9×10^{-9}	3.8×10^{-9}
Z13655	772	1.1×10^{-9}	2.4×10^{-9}	Z13824	834	1.9×10^{-9}	3.8×10^{-9}
Z13656	773	1.1×10^{-9}	3.7×10^{-9}	Z13838	835	2.6×10^{-9}	5.4×10^{-9}
Z13657	774	2.1×10^{-9}	3.9×10^{-9}	Z13840	836	2.2×10^{-9}	4.1×10^{-9}
Z13659	775	2.2×10^{-9}	3.1×10^{-9}	Z13842	837	2.2×10^{-9}	5.5×10^{-9}
Z13663	776	9.3×10^{-10}	1.5×10^{-9}	Z13845	838	2.6×10^{-9}	4.2×10^{-9}
Z13664	777	2.4×10^{-9}	4.2×10^{-9}	Z13846	839	2.3×10^{-9}	4.3×10^{-9}
Z13667	778	1.2×10^{-9}	2.3×10^{-9}	Z13848	840	2.1×10^{-9}	3.1×10^{-9}
Z13669	779	9.2×10^{-10}	1.7×10^{-9}	Z13849	841	2.1×10^{-9}	3.0×10^{-9}

Z variant	SEQ ID NO:	K _D pH 6.0 (M)	K _D pH 7.4 (M)	Z variant	SEQ ID NO:	K _D pH 6.0 (M)	K _D pH 7.4 (M)
Z13672	780	2.5×10^{-9}	5.6×10^{-9}	Z13860	842	2.3×10^{-9}	8.7×10^{-9}
Z13674	781	9.2×10^{-10}	1.3×10^{-9}	Z13865	843	2.5×10^{-9}	5.6×10^{-9}
Z13675	782	9.6×10^{-10}	2.2×10^{-9}	Z13866	844	2.0×10^{-9}	2.8×10^{-9}
Z13676	783	9.4×10^{-10}	3.1×10^{-9}	Z13875	845	2.0×10^{-9}	3.4×10^{-9}
Z13678	784	2.0×10^{-9}	3.3×10^{-9}	Z13879	846	2.1×10^{-9}	3.0×10^{-9}

Example 9**Production and characterization of Z variants from a matured library**

[0312] In this Example, twelve Z variants were produced in *E. coli*, purified and assayed for binding to FcRn as well as for inhibition of IgG binding to FcRn.

Materials and methods

[0313] Subcloning of Z variants into expression vectors: The DNA of twelve FcRn binding Z variants (Z13577 (SEQ ID NO:725), Z13578 (SEQ ID NO:726), Z13583 (SEQ ID NO:729), Z13592 (SEQ ID NO:734), Z13616 (SEQ ID NO:747), Z13621 (SEQ ID NO:750), Z13654 (SEQ ID NO:771), Z13663 (SEQ ID NO:776), Z13669 (SEQ ID NO:779), Z13674 (SEQ ID NO:781), Z13675 (SEQ ID NO:782) and Z13676 (SEQ ID NO:783)) were amplified from the library vector pAY02592. The subcloning was performed as described in Example 3. The Z gene fragments were subcloned into the expression vector pAY01448 resulting in the encoded sequence MGSSHHHHHHLQ-[Z#####]-VD.

[0314] Production of Z variants: Cultivation and purification of the His₆-tagged Z variants was performed essentially as described in Example 3.

[0315] Biacore binding and kinetic analyses: The interaction of FcRn binding His₆-tagged Z variants with human FcRn was analyzed in a Biacore 2000 instrument essentially as described in Example 3. Human FcRn purchased from Biorbyt (cat. no. orb84388) was used as target protein. The analytes were injected during 2 minutes at 30 µl/min. The dissociation phase was 4 minutes and the equilibration time between the analyte injections was 30 minutes.

[0316] In one experiment, the Z variants were injected at pH 6.0 followed by dissociation in buffers of pH 6.0 or pH 7.4, respectively, using the co-inject procedure. The concentration of the Z variants was 100 nM.

[0317] In another experiment, approximate kinetic constants (k_{on} and k_{off}) and affinities (K_D) were determined for a subset of Z variants. Injected concentrations were 540 nM, 180 nM, 60 nM, 20 nM and 6.7 nM.

[0318] AlphaLISA blocking assay: The potential of Z variants to inhibit binding of IgG to FcRn was analyzed in the AlphaLISA assay described in Example 3.

Results

[0319] Production of Z variants: The twelve FcRn binding Z variants constructed with an N-terminal His₆ tag were produced in *E. coli*. SDS-PAGE analysis of each final protein preparation showed that these predominantly contained the FcRn binding Z variant. The correct identity and molecular weight of each FcRn binding Z variant was confirmed by HPLC-MS analysis.

[0320] Biacore binding and kinetic analyses: The binding of the twelve Z variants to human FcRn and the dissociation at different pH were tested in a Biacore instrument by sequentially injecting each of the Z variants at pH 6.0 and either buffer pH 6.0 or pH 7.4 over a chip surface containing FcRn. The ligand immobilization level of the surface was 890 RU human FcRn. The twelve Z variants showed binding to FcRn at pH 6.0, and for all variants, faster off-rates were seen at pH 7.4 compared to pH 6.0.

[0321] The kinetic constants of the Z variants Z13577 (SEQ ID NO:725) and Z13621 (SEQ ID NO:750) interacting with FcRn at pH 6.0 were determined (see Table 10). Kinetic constants were calculated using curve sets of two or four injected concentrations of Z13577 and Z13621, respectively.

Table 10. Biacore kinetic constants and affinities for FcRn binding at pH 6.0.

Z variant	SEQ ID NO:	$k_{on}(M^{-1}s^{-1})$	$k_{off}(s^{-1})$	$K_D(M)$
Z13577	725	3.0×10^5	4.0×10^{-3}	13×10^{-9}
Z13621	750	6.4×10^5	3.7×10^{-3}	6×10^{-9}

[0322] AlphaLISA blocking analysis: The ability of twelve matured His₆-tagged monomeric Z variants to inhibit IgG binding to FcRn was tested in an AlphaLISA blocking assay. Serial dilutions of the Z variants were incubated with biotinylated human FcRn and the blocking ability of each respective variant was measured after addition of IgG coated Acceptor beads and subsequently streptavidin coated Donor beads. Inhibition could be measured as a decrease in

AlphaLISA counts for positive Z variants. All twelve tested Z variants were shown to block IgG binding to FcRn and the calculated IC50 values are shown in Table 11.

Table 11: Calculated IC50 values from AlphaLISA blocking assay.

Z variant	SEQ ID NO:	IC50 (M)
Z13577	725	1.2×10^{-8}
Z13578	726	1.2×10^{-8}
Z13583	729	2.7×10^{-9}
Z13592	734	6.4×10^{-9}
Z13616	747	7.4×10^{-9}
Z13621	750	3.2×10^{-9}
Z13654	771	3.5×10^{-9}
Z13663	776	1.1×10^{-8}
Z13669	779	5.2×10^{-9}
Z13674	781	2.5×10^{-9}
Z13675	782	8.2×10^{-9}
Z13676	783	3.9×10^{-9}

Example 10

Comparison of blocking capacity of IgG binding to FcRn

[0323] In this example, the IgG blocking capacity of the FcRn binding Z variant His₆-Z07918 (SEQ ID NO:707) was compared to Intravenous immunoglobulin (IVIg) and Subcutaneous immunoglobulin (SCIg) currently used in the treatment of some autoimmune disorders.

Materials and methods

[0324] Blocking of IgG-FcRn immunofluorescence staining: Human or murine FcRn-eGFP transduced HeLa cells were prepared as described in Example 4. Fixed cells were resuspended in 50 µl of a mix of 50 nM Alexa647-conjugated human IgG (Jackson laboratories, cat. no. 009-600-003) and His₆-tagged Z07918, IVIg (Octagam®, Octapharma) or SCIg (Gammanorm®, Octapharma), respectively, diluted at concentrations of 1000, 100, 10, 1, 0.1 or 0 (buffer control) nM in Mcllvanes, pH 6.0, plus 2.5 % FCS (Ultra low IgG, Life

technologies) and 0.1 % saponin (AppliChem). The cells were incubated for 1 h at 37 °C in the dark, washed with 2 x 100 µl Mcllvanes, pH 6.0, plus 2.5 % FCS (Ultra low IgG) pH 6.0 and re-suspended in 180 µl of Mcllvanes, pH 6.0, plus 1 % BSA. Data from 10,000 GFP/FcRn positive cells were obtained using a FACS Calibur (Beckman Coulter) and the data was analyzed using Flowing software 2.5.0 (Turku University).

Results

[0325] The experiment was performed to determine if the FcRn binding Z variant His₆-Z07918 (SEQ ID NO:707) blocks the IgG-FcRn interaction and compare the blocking effect to IVIg and SCIg. Human or murine FcRn-eGFP transduced HeLa cells were incubated with human Alexa647-conjugated IgG. The binding was blocked with unlabeled His₆-Z07918, IVIg or SCIg at different concentrations. The results showed that His₆-Z07918 effectively blocked hIgG binding to hFcRn to a similar extent as IVIg or SCIg (Figure 9).

Example 11

Increased IgG catabolism by FcRn binding Z variants in mice

[0326] The ability of the FcRn binding Z variant Z07918 to block IgG binding to FcRn *in vitro* was shown in Example 10. In this example, the blocking ability of the same Z variant was evaluated *in vivo*. Blocking of IgG-FcRn interactions *in vivo* will lead to increased IgG catabolism and concomitant reduced levels of IgG (Mezo 2008, *supra*).

Materials and methods

[0327] Animal study: The FcRn-binding Z variants Z11948 (SEQ ID NO:1060) and Z07918-PP013 (Z07918 (SEQ ID NO:707) identical to Z11948 but with the N-terminus starting with the amino acids VD instead of AE, in fusion with the ABD variant PP013 (SEQ ID NO:1063)) or vehicle (PBS buffer), were administered to male NMRI (Charles River), at a dose of 16.3 µmol/kg. The mice were treated with five intravenous injections given at 0, 24, 48, 72 and 96 h. Serum samples were taken at 0, 72, 120 and 168 h (termination of study) and stored at -20 °C. The concentration of mouse IgG in serum was quantified by ELISA.

[0328] Mouse IgG ELISA: The concentration of mouse IgG in mouse serum samples was analyzed by a mouse IgG ELISA kit (Mabtech 3825-1AD-6) and performed as described by the manufacturer. The concentration of mIgG was calculated from a standard curve provided and GraphPad prism5 using a nonlinear regression formula. The concentration of IgG in individual

mice at 24, 72, 120 and 168 h were related to the level at 0 h and the results are therefore presented as percentage of IgG (0 h).

Results

[0329] The results showed a reduction of mouse IgG concentration in mice treated with FcRn-specific Z variants. Both Z11948 and the ABD-fused variant Z07918-PP013 lowered the concentration of endogenous IgG in mice *in vivo*. Most pronounced effects were obtained with the ABD-fused variant and after 120 hours. Thus, the results indicates that the FcRn-specific Z variants blocked recycling of IgG resulting in increased IgG catabolism and subsequent lower levels of IgG in mice.

Example 12

In vitro transcytosis of FcRn binding Z variants

[0330] In this example, the FcRn binding Z variants are tested for their ability to be transported through epithelial or endothelial cells or recycled by FcRn *in vitro*. A drug containing a Z variant with the power of transcytosis will facilitate drug uptake after for example oral or pulmonary administration.

Materials and methods

[0331] Cells, for example T84, MDCK, HeLa, CaCo2, CaLu-1 and/or CaLu-3 cells, with or without endogenous or recombinant expression of FcRn, are grown in respective growth medium on a membrane in a transwell to form a monolayer. The integrity of monolayers can be evaluated by measuring the electrical resistance or adding a probe that is not able to penetrate or being actively transported over the cell monolayer. A defined monolayer of cells is pulsed from the apical or basolateral side with ligand such as FcRn binding Z variants, HSA or IgG in a buffer such as HBSS (Hanks' Balanced Salt Solution, SigmaAldrich, cat. no. H9269) or growth medium at a suitable pH and temperature, and chased with buffers such as HBSS or growth medium at a suitable pH and temperature on the opposite side.

[0332] In a variant of this assay, ligands can be chased with buffers such as HBSS or growth medium at suitable pH and temperature on the same side as administration to measure recycled ligand as well. This can be done in a transwell or in a cell culture dish. Cells are seeded into transwell or cell culture dishes and pulsed with ligands such as FcRn binding Z variants, HSA or IgG. Endocytosed ligands will bind to FcRn and return to the cell surface at the same or opposite side as they were loaded. After pulsing, free ligands are removed by

washing the cells with cold buffer. To chase ligands, warm buffer or medium is added to the cells and, after a period in the range from 10 minutes to several hours, the buffer or medium is removed and assayed for the presence of ligands.

[0333] In a variant of this assay, ligands such as FcRn binding Z variants, HSA or IgG can be used to block the binding to FcRn by ligands such as other FcRn binding Z variants, HSA or IgG by administering them at the same time or sequentially to the cells.

[0334] The amount of ligand can be quantified by methods such as ELISA, HPLC-MS, fluorescent dye or radio labeling.

[0335] The results of the experiment described above are expected to show that the FcRn-specific Z variants can be transcytosed and/or recycled *in vitro*.

ITEMIZED LISTING OF EMBODIMENTS

[0336]

1. FcRn binding polypeptide, comprising an FcRn binding motif *BM*, which motif consists of the amino acid sequence

$$\text{EX}_2 \text{X}_3 \text{X}_4 \text{AX}_6 \text{X}_7 \text{EIR WLPNLX}_{16} \text{X}_{17} \text{X}_{18} \text{QR X}_{21} \text{AFIX}_{25} \text{X}_{26} \text{LX}_{28} \text{X}_{29}$$

wherein, independently from each other,

X_2 is selected from A, D, E, F, H, I, K, L, N, Q, R, S, T, V, W and Y;

X_3 is selected from A, D, E, F, G, H, I, K, L, M, N, Q, R, S, T, V, W and Y;

X_4 is selected from A, D, E, F, G, H, I, K, L, N, Q, R, S, T, V, W and Y;

X_6 is selected from A, E, F, G, H, I, K, Q, R, S and V;

X_7 is selected from A, F, H, K, N, Q, R, S and V;

X_{16} is selected from N and T;

X_{17} is selected from F, W and Y;

X_{18} is selected from A, D, E and N;

X_{21} is selected from A, S, V and W;

X_{25} is selected from D, E, G, H, I, K, L, N, Q, R, S, T, V, W and Y;

X₂₆ is selected from K and S;

X₂₈ is selected from A, D, E, F, H, I, K, L, N, Q, R, S, T, V, W and Y; and

X₂₉ is selected from D and R.

2. 2. FcRn binding polypeptide according to item 1, wherein, independently from each other,

X₂ is selected from A, D, E, F, H, I, K, L, N, Q, R, S, T, V, W and Y;

X₃ is selected from A, D, E, F, H, I, K, L, M, N, Q, R, S, T, V, W and Y;

X₄ is selected from A, D, E, F, H, I, K, L, N, Q, R, S, T, V, W and Y;

X₆ is selected from A, E, F, G, H, I, K, Q, R and S;

X₇ is selected from A, F, H, K, N, Q, R, S and V;

X₁₆ is selected from N and T;

X₁₇ is selected from F and Y;

X₁₈ is D;

X₂₁ is V;

X₂₅ is selected from D, E, H, I, K, L, N, Q, R, S, T, V, W and Y;

X₂₆ is selected from K and S;

X₂₈ is selected from A, D, E, F, H, I, K, L, N, Q, R, S, T, V and W; and

X₂₉ is selected from D and R.

3. 3. FcRn binding polypeptide according to item 1, wherein the BM consists of an amino acid sequence selected from

1. i)

EX₂ X₃ X₄ AX₆ HEIR WLPNLTX₁₇ X₁₈ QR X₂₁ AFIX₂₅ KLX₂₈ D

wherein, independently from each other,

X₂ is selected from A, D, E, F, H, I, K, L, N, Q, R, S, T, V, W and Y;

X₃ is selected from A, D, E, G, H, K, L, M, N, Q, R, S, T, V and Y;

X₄ is selected from A, D, E, F, G, I, K, L, N, Q, R, S, T, V and Y;

X_6 is selected from A, G, K, R, S and V;

X_{17} is selected from F, W and Y;

X_{18} is selected from A, D, E and N;

X_{21} is selected from A, S, V and W;

X_{25} is selected from D, G, H, K, L, N, R, V and W;

X_{28} is selected from A, D, E, H, K, L, N, Q, R, S, T, W and Y;

and

2. ii) an amino acid sequence which has at least 96 % identity to a sequence defined by i).

4. 4. FcRn binding polypeptide according to any preceding item, wherein X_2 is selected from A, D, E, F, I, L, N, Q, R, S, T, V, W and Y.
5. 5. FcRn binding polypeptide according to item 4, wherein X_2 is selected from A, D, F, I, L, N, Q, R, S, T, V, W and Y.
6. 6. FcRn binding polypeptide according to item 5, wherein X_2 is selected from A, D, F, I, L, N, Q, R, S, V and W.
7. 7. FcRn binding polypeptide according to item 5, wherein X_2 is selected from A, I, L, N, Q, R, S, T, V, W and Y.
8. 8. FcRn binding polypeptide according to item 7, wherein X_2 is selected from A, I, L, N, Q, S, T, V and W.
9. 9. FcRn binding polypeptide according to item 6 or 8, wherein X_2 is selected from A, I, L, N, Q, V and W.
10. 10. FcRn binding polypeptide according to item 9, wherein X_2 is selected from A, I, L, Q, V and W.
11. 11. FcRn binding polypeptide according to item 10, wherein X_2 is selected from A, I, L and Q.
12. 12. FcRn binding polypeptide according to item 11, wherein X_2 is selected from I, L and Q.
13. 13. FcRn binding polypeptide according to item 12, wherein X_2 is selected from I and Q.
14. 14. FcRn binding polypeptide according to item 13, wherein X_2 is I.
15. 15. FcRn binding polypeptide according to item 13, wherein X_2 is Q.
16. 16. FcRn binding polypeptide according to any one of items 1 and 3-15, wherein X_3 is selected from A, D, E, G, H, K, L, M, N, Q, R, S, T, V and Y.
17. 17. FcRn binding polypeptide according to item 2 or 16, wherein X_3 is selected from A, D, E, H, K, L, M, N, Q, R, S, T, V and Y.
18. 18. FcRn binding polypeptide according to item 16, wherein X_3 is selected from A, D, E, G, H, K, L, M, N, Q, R, S and T.

19. 19. FcRn binding polypeptide according to item 18, wherein X_3 is selected from A, D, E, G, H, K, M, N, Q, S and T.
20. 20. FcRn binding polypeptide according to item 19, wherein X_3 is selected from A, D, E, G, H, M, N, Q, S and T.
21. 21. FcRn binding polypeptide according to item 19, wherein X_3 is selected from A, D, E, K, N, Q, S and T.
22. 22. FcRn binding polypeptide according to item 21, wherein X_3 is selected from A, D, E, K, Q and T.
23. 23. FcRn binding polypeptide according to item 22, wherein X_3 is selected from A, D, E, Q and T.
24. 24. FcRn binding polypeptide according to item 23, wherein X_3 is selected from D, E and T.
25. 25. FcRn binding polypeptide according to item 24, wherein X_3 is selected from D and E.
26. 26. FcRn binding polypeptide according to item 25, wherein X_3 is D.
27. 27. FcRn binding polypeptide according to item 25, wherein X_3 is E.
28. 28. FcRn binding polypeptide according to any one of items 1 and 3-27, wherein X_4 is selected from A, D, E, F, G, I, K, L, N, Q, R, S, T, V and Y.
29. 29. FcRn binding polypeptide according to item 28, wherein X_4 is selected from A, D, E, G, N, Q, R, S, T and V.
30. 30. FcRn binding polypeptide according to item 2 or 28, wherein X_4 is selected from A, D, E, F, I, K, L, N, Q, R, S, T and V.
31. 31. FcRn binding polypeptide according to item 30, wherein X_4 is selected from A, D, E, I, K, N, Q, R, S and T.
32. 32. FcRn binding polypeptide according to item 31, wherein X_4 is selected from A, D, E, I, K, Q, S and T.
33. 33. FcRn binding polypeptide according to item 32, wherein X_4 is selected from A, D, I, K, Q and S.
34. 34. FcRn binding polypeptide according to item 32, wherein X_4 is selected from A, D, E, K and S.
35. 35. FcRn binding polypeptide according to item 33 or 34, wherein X_4 is selected from A, D, K and S.
36. 36. FcRn binding polypeptide according to item 34, wherein X_4 is selected from A, D, E and K.
37. 37. FcRn binding polypeptide according to item 35 or 36, wherein X_4 is selected from A, D and K.
38. 38. FcRn binding polypeptide according to item 37, wherein X_4 is selected from A and D.
39. 39. FcRn binding polypeptide according to item 36, wherein X_4 is selected from A and E.
40. 40. FcRn binding polypeptide according to item 38 or 39, wherein X_4 is A.
41. 41. FcRn binding polypeptide according to item 38, wherein X_4 is D.
42. 42. FcRn binding polypeptide according to item 39, wherein X_4 is E.
43. 43. FcRn binding polypeptide according to any one of items 1 and 4-42, wherein X_6 is

- selected from A, G, K, Q, R, S and V.
44. 44. FcRn binding polypeptide according to item 3 or 43, wherein X_6 is selected from A, G, K, R, S and V.
 45. 45. FcRn binding polypeptide according to item 2 or 44, wherein X_6 is selected from A, G, K, R and S.
 46. 46. FcRn binding polypeptide according to item 44, wherein X_6 is selected from A, G, K, S and V.
 47. 47. FcRn binding polypeptide according to item 46, wherein X_6 is selected from A, G, K and V.
 48. 48. FcRn binding polypeptide according to item 45 or 46, wherein X_6 is selected from A, G, K and S.
 49. 49. FcRn binding polypeptide according to item 47 or 48, wherein X_6 is selected from A, G and K.
 50. 50. FcRn binding polypeptide according to item 47, wherein X_6 is selected from A, G and V.
 51. 51. FcRn binding polypeptide according to item 49 or 50, wherein X_6 is selected from A and G.
 52. 52. FcRn binding polypeptide according to item 51, wherein X_6 is A.
 53. 53. FcRn binding polypeptide according to item 51, wherein X_6 is G.
 54. 54. FcRn binding polypeptide according to any one of items 1, 2 and 4-53, wherein X_7 is selected from A and H.
 55. 55. FcRn binding polypeptide according to item 54, wherein X_7 is H.
 56. 56. FcRn binding polypeptide according to any one of items 1, 2 and 4-55, wherein X_{16} is T.
 57. 57. FcRn binding polypeptide according to any one of items 1, 2 and 4-55, wherein X_{16} is N.
 58. 58. FcRn binding polypeptide according to any one of items 1 and 3-57, wherein X_{17} is selected from F and Y.
 59. 59. FcRn binding polypeptide according to any preceding item, wherein X_{17} is F.
 60. 60. FcRn binding polypeptide according to any one of items 1 and 3-59, wherein X_{18} is selected from A, D and E.
 61. 61. FcRn binding polypeptide according to item 60, wherein X_{18} is selected from A and D.
 62. 62. FcRn binding polypeptide according to item 61, wherein X_{18} is D.
 63. 63. FcRn binding polypeptide according to any one of items 1 and 3-62, wherein X_{21} is selected from V and W.
 64. 64. FcRn binding polypeptide according to item 63, wherein X_{21} is V.
 65. 65. FcRn binding polypeptide according to any one of items 1 and 4-64, wherein X_{25} is selected from D, E, G, H, K, L, N, Q, R, V and W.
 66. 66. FcRn binding polypeptide according to item 65, wherein X_{25} is selected from D, G, H,

K, L, N, R, V and W.

67. 67. FcRn binding polypeptide according to any one of items 2, 3 and 66, wherein X_{25} is selected from H, L, R, V and W.
68. 68. FcRn binding polypeptide according to item 67, wherein X_{25} is selected from H, R, V and W.
69. 69. FcRn binding polypeptide according to item 68, wherein X_{25} is selected from H, R and V.
70. 70. FcRn binding polypeptide according to item 67, wherein X_{25} is selected from H, L and R.
71. 71. FcRn binding polypeptide according to item 69 or 70, wherein X_{25} is selected from H and R.
72. 72. FcRn binding polypeptide according to item 69, wherein X_{25} is selected from H and V.
73. 73. FcRn binding polypeptide according to item 71 or 72, wherein X_{25} is H.
74. 74. FcRn binding polypeptide according to any one of items 1, 2 and 4-73, wherein X_{26} is K.
75. 75. FcRn binding polypeptide according to any one of items 1, 2 and 4-73, wherein X_{26} is S.
76. 76. FcRn binding polypeptide according to any one of items 1 and 3-75, wherein X_{28} is selected from A, D, E, H, K, L, N, Q, R, S, T, W and Y.
77. 77. FcRn binding polypeptide according to item 76, wherein X_{28} is selected from A, D, E, K, L, N, Q, R, S, T, W and Y.
78. 78. FcRn binding polypeptide according to item 77, wherein X_{28} is selected from A, D, E, L, R, S, T, W and Y.
79. 79. FcRn binding polypeptide according to item 2 or 77, wherein X_{28} is selected from A, D, K, L, N, Q, R, S, T and W.
80. 80. FcRn binding polypeptide according to item 78 or 79, wherein X_{28} is selected from A, D and R.
81. 81. FcRn binding polypeptide according to item 80, wherein X_{28} is selected from A and R.
82. 82. FcRn binding polypeptide according to item 80, wherein X_{28} is selected from D and R.
83. 83. FcRn binding polypeptide according to item 81, wherein X_{28} is A.
84. 84. FcRn binding polypeptide according to item 81 or 82, wherein X_{28} is R.
85. 85. FcRn binding polypeptide according to item 82, wherein X_{28} is D.
86. 86. FcRn binding polypeptide according to any one of items 1, 2 and 4-85, wherein X_{29} is D.
87. 87. FcRn binding polypeptide according to any one of items 1, 2 and 4-85, wherein X_{29} is R.
88. 88. FcRn binding polypeptide according to any one of items 1, 2 and 4-87, wherein X_6X_7 is selected from AH and GH.

89. 89. FcRn binding polypeptide according to item 88, wherein X_6X_7 is AH.
90. 90. FcRn binding polypeptide according to item 88, wherein X_6X_7 is GH.
91. 91. FcRn binding polypeptide according to any preceding item, wherein $X_{17}X_{18}$ is selected from FD and YD.
92. 92. FcRn binding polypeptide according to item 91, wherein $X_{17}X_{18}$ is FD.
93. 93. FcRn binding polypeptide according to any preceding item, wherein the sequence fulfills at least three of the six conditions I-VI:
 1. I. X_6 is selected from A, G, K and S, such as in particular A;
 2. II. X_7 is H;
 3. III. X_{17} is selected from F and Y, such as in particular F;
 4. IV. X_{18} is D;
 5. V. X_{21} is selected from V and W, such as in particular V;
 6. VI. X_{25} is selected from H and R, such as in particular H.
94. 94. FcRn binding polypeptide according to item 93, wherein the sequence fulfills at least four of the six conditions I-VI.
95. 95. FcRn binding polypeptide according to item 94, wherein the sequence fulfills at least five of the six conditions I-VI.
96. 96. FcRn binding polypeptide according to item 95, wherein the sequence fulfills all of the six conditions I-VI.
97. 97. FcRn binding polypeptide according to any preceding item, wherein the sequence is selected from the group consisting of SEQ ID NO:1-353.
98. 98. FcRn binding polypeptide according to item 97, wherein the sequence is selected from the group consisting of SEQ ID NO:1-15, SEQ ID NO:17-140 and SEQ ID NO:353.
99. 99. FcRn binding polypeptide according to item 98, wherein the sequence is selected from the group consisting of SEQ ID NO:1-2 and SEQ ID NO:17-140.
100. 100. FcRn binding polypeptide according to item 99, wherein the sequence is selected from the group consisting of SEQ ID NO:1-2, SEQ ID NO:17-92, SEQ ID NO:94-103, SEQ ID NO:105-125 and SEQ ID NO:127-140.
101. 101. FcRn binding polypeptide according to item 98, wherein the sequence is selected from the group consisting of SEQ ID NO:1-8, SEQ ID NO:13, SEQ ID NO:19-20, SEQ ID NO:23, SEQ ID NO:28, SEQ ID NO:41, SEQ ID NO:44, SEQ ID NO:65, SEQ ID NO:70, SEQ ID NO:73, SEQ ID NO:75-77 and SEQ ID NO:353.
102. 102. FcRn binding polypeptide according to item 100 or 101, wherein the sequence is selected from the group consisting of SEQ ID NO:1, SEQ ID NO:23, SEQ ID NO:28, SEQ ID NO:41, SEQ ID NO:44, SEQ ID NO:65, SEQ ID NO:73 and SEQ ID NO:75-77.
103. 103. FcRn binding polypeptide according to item 102, wherein the sequence is selected from the group consisting of SEQ ID NO:1, SEQ ID NO:23, SEQ ID NO:44, SEQ ID NO:65, SEQ ID NO:75 and SEQ ID NO:77.
104. 104. FcRn binding polypeptide according to item 103, wherein the sequence is selected from the group consisting of SEQ ID NO:1, SEQ ID NO:23 and SEQ ID NO:75.
105. 105. FcRn binding polypeptide according to item 104, wherein the sequence is SEQ ID NO:1.
106. 106. FcRn binding polypeptide according to any preceding item, wherein said FcRn

binding motif forms part of a three-helix bundle protein domain.

107. 107. FcRn binding polypeptide according to item 106, wherein said FcRn binding motif essentially forms part of two helices with an interconnecting loop, within said three-helix bundle protein domain.
108. 108. FcRn binding polypeptide according to item 107, wherein said three-helix bundle protein domain is selected from bacterial receptor domains.
109. 109. FcRn binding polypeptide according to item 108, wherein said three-helix bundle protein domain is selected from domains of protein A from *Staphylococcus aureus* or derivatives thereof.
110. 110. FcRn binding polypeptide according to any preceding item, which comprises an amino acid sequence selected from:

iii)

$$K-[BM]-DPSQS X_a X_b LLX_c EAKKL X_d X_e X_f Q;$$

wherein

[BM] is an FcRn binding motif as defined herein, provided that X₂₉ is D;

X_a is selected from A and S;

X_b is selected from N and E;

X_c is selected from A, S and C;

X_d is selected from E, N and S;

X_e is selected from D, E and S;

X_f is selected from A and S;

and

iv) an amino acid sequence which has at least 93 % identity to a sequence defined by iii).

111. 111. FcRn binding polypeptide according to any one of items 1-109, which comprises an amino acid sequence selected from

v)

$$K-[BM]-QPEQS X_a X_b LLX_c EAKKL X_d X_e X_f Q;$$

wherein

[BM] is an FcRn binding motif as defined herein, provided that X₂₉ is R;

X_a is selected from A and S;

X_b is selected from N and E;

X_c is selected from A, S and C;

X_d is selected from E, N and S;

X_e is selected from D, E and S;

X_f is selected from A and S;

and

vi) an amino acid sequence which has at least 93 % identity to a sequence defined by v).

112. 112. FcRn binding polypeptide according to item 110 or 111, wherein X_a in sequence iii) or v) is A.
113. 113. FcRn binding polypeptide according to item 110 or 111, wherein X_a in sequence iii) or v) is S.
114. 114. FcRn binding polypeptide according to any one of items 110-113, wherein X_b in sequence iii) or v) is N.
115. 115. FcRn binding polypeptide according to any one of items 110-113, wherein X_b in sequence iii) or v) is E.
116. 116. FcRn binding polypeptide according to any one of items 110-115, wherein X_c in sequence iii) or v) is A.
117. 117. FcRn binding polypeptide according to any one of items 110-115, wherein X_c in sequence iii) or v) is S.
118. 118. FcRn binding polypeptide according to any one of items 110-115, wherein X_c in sequence iii) or v) is C.
119. 119. FcRn binding polypeptide according to any one of items 110-118, wherein X_d in sequence iii) or v) is E.
120. 120. FcRn binding polypeptide according to any one of items 110-118, wherein X_d in sequence iii) or v) is N.
121. 121. FcRn binding polypeptide according to any one of items 110-118, wherein X_d in sequence iii) or v) is S.
122. 122. FcRn binding polypeptide according to any one of items 110-121, wherein X_e in sequence iii) or v) is D.
123. 123. FcRn binding polypeptide according to any one of items 110-121, wherein X_e in sequence iii) or v) is E.
124. 124. FcRn binding polypeptide according to any one of items 110-121, wherein X_e in sequence iii) or v) is S.
125. 125. FcRn binding polypeptide according to any one of items 110-119, 121, 123 and

- 124, wherein X_dX_e in sequence iii) or v) is selected from EE, ES, SE and SS.
126. 126. FcRn binding polypeptide according to item 125, wherein X_dX_e in sequence iii) or v) is ES.
127. 127. FcRn binding polypeptide according to item 125, wherein X_dX_e in sequence iii) or v) is SE.
128. 128. FcRn binding polypeptide according to any one of items 110-127, wherein X_f in sequence iii) or v) is A.
129. 129. FcRn binding polypeptide according to any one of items 110-127, wherein X_f in sequence iii) or v) is S.
130. 130. FcRn binding polypeptide according to item 110 or 111, wherein in sequence iii) or v), X_a is A; X_b is N; X_c is A and X_f is A.
131. 131. FcRn binding polypeptide according to item 110 or 111, wherein in sequence iii) or v), X_a is A; X_b is N; X_c is C and X_f is A.
132. 132. FcRn binding polypeptide according to item 110 or 111, wherein in sequence iii) or v), X_a is S; X_b is E; X_c is S and X_f is S.
133. 133. FcRn binding polypeptide according to item 110 or 111, wherein in sequence iii) or v), X_a is S; X_b is E; X_c is C and X_f is S.
134. 134. FcRn binding polypeptide according to item 110 or 111, wherein in sequence iii) or v), X_a is A; X_b is N; X_c is A; X_dX_e is ND and X_f is A.
135. 135. FcRn binding polypeptide according to item 110 or 111, wherein in sequence iii) or v), X_a is A; X_b is N; X_c is C; X_dX_e is ND and X_f is A.
136. 136. FcRn binding polypeptide according to item 110 or 111, wherein in sequence iii) or v), X_a is S; X_b is E; X_c is S; X_dX_e is ND and X_f is S.
137. 137. FcRn binding polypeptide according to item 110 or 111, wherein in sequence iii) or v), X_a is S; X_b is E; X_c is C; X_dX_e is ND and X_f is S.
138. 138. FcRn binding polypeptide according to item 110 or 111, wherein in sequence iii) or v), X_a is A; X_b is N; X_c is A; X_dX_e is SE and X_f is A.
139. 139. FcRn binding polypeptide according to item 110 or 111, wherein in sequence iii) or v), X_a is A; X_b is N; X_c is C; X_dX_e is SE and X_f is A.
140. 140. FcRn binding polypeptide according to item 110 or 111, wherein in sequence iii) or v), X_a is S; X_b is E; X_c is S; X_dX_e is SE and X_f is S.
141. 141. FcRn binding polypeptide according to item 110 or 111, wherein in sequence iii) or v), X_a is S; X_b is E; X_c is C; X_dX_e is SE and X_f is S.
142. 142. FcRn binding polypeptide according to item 110 or 111, wherein in sequence iii) or v), X_a is A; X_b is N; X_c is A; X_dX_e is ES and X_f is A.
143. 143. FcRn binding polypeptide according to item 110 or 111, wherein in sequence iii) or v), X_a is A; X_b is N; X_c is C; X_dX_e is ES and X_f is A.
144. 144. FcRn binding polypeptide according to item 110 or 111, wherein in sequence iii) or v), X_a is S; X_b is E; X_c is S; X_dX_e is ES and X_f is S.
145. 145. FcRn binding polypeptide according to item 110 or 111, wherein in sequence iii) or v), X_a is S; X_b is E; X_c is C; X_dX_e is ES and X_f is S.

146. 146. FcRn binding polypeptide according to any one of items 110 and 112-145, wherein sequence iii) is selected from the group consisting of SEQ ID NO:354-706.
147. 147. FcRn binding polypeptide according to item 146, wherein sequence iii) is selected from the group consisting of SEQ ID NO:354-368, SEQ ID NO:370-493 and SEQ ID NO:706.
148. 148. FcRn binding polypeptide according to item 147, wherein sequence iii) is selected from the group consisting of SEQ ID NO:354-355 and SEQ ID NO:370-493.
149. 149. FcRn binding polypeptide according to item 148, wherein sequence iii) is selected from the group consisting of SEQ ID NO:354-355, SEQ ID NO:370-445, SEQ ID NO:447-456, SEQ ID NO:458-478 and SEQ ID NO:480-493.
150. 150. FcRn binding polypeptide according to item 147, wherein sequence iii) is selected from the group consisting of SEQ ID NO:354-361, SEQ ID NO:366, SEQ ID NO:372-373, SEQ ID NO:376, SEQ ID NO:381, SEQ ID NO:394, SEQ ID NO:397, SEQ ID NO:418, SEQ ID NO:423, SEQ ID NO:426, SEQ ID NO:428-430 and SEQ ID NO:706.
151. 151. FcRn binding polypeptide according to item 149 or 150, wherein sequence iii) is selected from the group consisting of SEQ ID NO:354, SEQ ID NO:376, SEQ ID NO:381, SEQ ID NO:394, SEQ ID NO:397, SEQ ID NO:418, SEQ ID NO:426 and SEQ ID NO:428-430.
152. 152. FcRn binding polypeptide according to item 151, wherein sequence iii) is selected from the group consisting of SEQ ID NO:354, SEQ ID NO:376, SEQ ID NO:397, SEQ ID NO:418, SEQ ID NO:428 and SEQ ID NO:430.
153. 153. FcRn binding polypeptide according to item 152, wherein sequence iii) is selected from the group consisting of SEQ ID NO:354, SEQ ID NO:376 and SEQ ID NO:428.
154. 154. FcRn binding polypeptide according to item 153, wherein sequence iii) is SEQ ID NO:354.
155. 155. FcRn binding polypeptide according to any one of items 1-109, which comprises an amino acid sequence selected from:

vii)

YAK-[BM]-DPSQS SELLX_C EAKKL NDSQA P;

wherein [BM] is an FcRn binding motif as defined in any one of items 1-105 and X_C is selected from A, S and C; and

viii) an amino acid sequence which has at least 94 % identity to a sequence defined by vii).

156. 156. FcRn binding polypeptide according to any one of items 1-109, which comprises an amino acid sequence selected from:

ix)

FNK-[BM]-DPSQS ANLLX_C EAKKL NDAQA P;

wherein [BM] is an FcRn binding motif as defined in any one of items 1-105 and X_C is selected from A and C; and

x) an amino acid sequence which has at least 94 % identity to a sequence defined by ix).

157. 157. FcRn binding polypeptide according to item 109, which comprises an amino acid sequence selected from:

ADNNFNK-[BM]-DPSQSANLLSEAKKLNESQAPK;

ADNKFNK-[BM]-DPSQSANLLAEAKKLNDQAAPK;

ADNKFNK-[BM]-DPSVSKEILAEAKKLNDQAAPK;

ADAQQNNFNK-[BM]-DPSQSTNVLGEAKKLNESQAPK;

AQHDE-[BM]-DPSQSANVLGEAQLNDSQAPK;

VDNKFNK-[BM]-DPSQSANLLAEAKKLNDQAAPK;

AEAKYAK-[BM]-DPSESSELLSEAKKLNKSQAPK;

VDAKYAK-[BM]-DPSQSSELLAEAKKLNDQAAPK;

VDAKYAK-[BM]-DPSQSSELLAEAKKLNDSQAPK;

AEAKYAK-[BM]-DPSQSSELLSEAKKLNDSQAPK;

AEAKYAK-[BM]-DPSQSSELLSEAKKLSQAPK

AEAKYAK-[BM]-DPSQSSELLSEAKKLESSQAPK

VDAKYAK-[BM]-DPSQSSELLSEAKKLNDSQAPK;

VDAKYAK-[BM]-DPSQSSELLSEAKKLSQAPK;

VDAKYAK-[BM]-DPSQSSELLSEAKKLESSQAPK;

VDAKYAK-[BM]-DPSQSSELLAEAKKLN KAQAPK; and

AEAKYAK-[BM]-DPSQSSELLAEAKKLNKAQAPK;

wherein [BM] is an FcRn binding motif as defined in any one of items 1-105.

158. 158. FcRn binding polypeptide according to any preceding item, which comprises an amino acid sequence selected from:

xi)

AEAKYAK-[BM]-DPSQSSELLSEAKKLNDSQAPK;

wherein [BM] is an FcRn binding motif as defined in any one of items 1-105; and

xii) an amino acid sequence which has at least 94 % identity to the sequence defined in xi).

159. 159. FcRn binding polypeptide according to item 158, in which sequence xi) is selected from the group consisting of SEQ ID NO:1060-1062.

160. 160. FcRn binding polypeptide according to any preceding item, which comprises an amino acid sequence selected from:

xiii)

VDAKYAK-[BM]-DPSQSSELLSEAKKLNDSQAPK;

wherein [BM] is an FcRn binding motif as defined in any one of items 1-105; and

xiv) an amino acid sequence which has at least 94 % identity to the sequence defined in xiii).

161. 161. FcRn binding polypeptide according to item 160, in which sequence xiii) is selected from the group consisting of SEQ ID NO: 707-1059.

162. 162. FcRn binding polypeptide according to item 161, in which sequence xiii) is selected from the group consisting of SEQ ID NO:707-721, SEQ ID NO:723-846 and SEQ ID NO:1059.

163. 163. FcRn binding polypeptide according to item 162, in which sequence xiii) is selected from the group consisting of SEQ ID NO:707-708 and SEQ ID NO:723-846.

164. 164. FcRn binding polypeptide according to item 163, in which sequence xiii) is selected from the group consisting of SEQ ID NO:707-708, SEQ ID NO:723-798, SEQ ID NO:800-809, SEQ ID NO:811-831 and SEQ ID NO:833-846.

165. 165. FcRn binding polypeptide according to item 162, in which sequence xiii) is selected from the group consisting of SEQ ID NO:707-714, SEQ ID NO:719, SEQ ID NO:725-726, SEQ ID NO:729, SEQ ID NO:734, SEQ ID NO:747, SEQ ID NO:750, SEQ ID NO:771, SEQ ID NO:776, SEQ ID NO:779, SEQ ID NO:781-783 and SEQ ID NO:1059.

166. 166. FcRn binding polypeptide according to item 163 or 165, in which sequence xiii) is selected from the group consisting of SEQ ID NO:707, SEQ ID NO:729, SEQ ID NO:734, SEQ ID NO:747, SEQ ID NO:750, SEQ ID NO:771, SEQ ID NO:779 and SEQ ID NO:781-783.

167. 167. FcRn binding polypeptide according to item 166, in which sequence xiii) is selected from the group consisting of SEQ ID NO:707, SEQ ID NO:729, SEQ ID NO:750, SEQ ID NO:771, SEQ ID NO:781 and SEQ ID NO:783.

168. 168. FcRn binding polypeptide according to item 167, in which sequence xiii) is selected from the group consisting of SEQ ID NO:707, SEQ ID NO:729 and SEQ ID NO:781.

169. 169. FcRn binding polypeptide according to item 168, in which sequence xiii) is SEQ ID NO:707.

170. 170. FcRn binding polypeptide according to any preceding item, which is capable of binding to FcRn at pH 6.0 such that the K_D value of the interaction is at most 1×10^{-6} M,

- such as at most 1×10^{-7} M, such as at most 1×10^{-8} M, such as at most 1×10^{-9} M, such as at most 1×10^{-10} M.
171. 171. FcRn binding polypeptide according to any preceding item, wherein the K_D value of the interaction between FcRn binding polypeptide and FcRn at pH 7.4 is higher than the K_D value of said interaction at pH 6.0, such as at least 2 times higher, such as at least 5 times higher, such as at least 10 times higher, such as at least 50 times higher, such as at least 100 times higher than the K_D value of said interaction at pH 6.0.
 172. 172. FcRn binding polypeptide according to any preceding item, wherein the K_D value of said interaction at pH 7.4 is at least 1×10^{-8} M, such as at least 1×10^{-7} M, such as at least 1×10^{-6} M, such as at least 1×10^{-5} M.
 173. 173. FcRn binding polypeptide according to any one of items 1-170, wherein the K_D value of said interaction at pH 7.4 is the same as or lower than the K_D value of said interaction at pH 6.0.
 174. 174. FcRn binding polypeptide according to any one of items 1-170, wherein the K_D value of said interaction at pH 7.4 is at most 1×10^{-6} M, such as at most 1×10^{-7} M, such as at most 1×10^{-8} M, such as at most 1×10^{-9} M, such as at most 1×10^{-10} M.
 175. 175. FcRn binding polypeptide according to any preceding item, which comprises at least one additional amino acid at the C-terminal and/or N-terminal end.
 176. 176. FcRn binding polypeptide according to item 175, wherein said at least one additional amino acid extension improves production, purification, stabilization *in vivo* or *in vitro*, coupling or detection of the polypeptide.
 177. 177. FcRn binding polypeptide according to any preceding item in multimeric form, comprising at least two FcRn binding polypeptide monomer units, whose amino acid sequences may be the same or different.
 178. 178. FcRn binding polypeptide according to item 177, wherein said FcRn binding polypeptide monomer units are covalently coupled together.
 179. 179. FcRn binding polypeptide according to item 177, wherein the FcRn binding polypeptide monomer units are expressed as a fusion protein.
 180. 180. FcRn binding polypeptide according to any one of items 177-179, in dimeric form.
 181. 181. Fusion protein or conjugate comprising
 - a first moiety consisting of an FcRn binding polypeptide according to any preceding item; and
 - a second moiety consisting of a polypeptide having a desired biological activity.
 182. 182. Fusion protein or conjugate according to item 181, wherein the *in vivo* half-life of said fusion protein or conjugate is longer than the *in vivo* half-life of the polypeptide having a desired biological activity *per se*.
 183. 183. Fusion protein or conjugate according to any one of items 181-182, wherein said desired biological activity is a therapeutic activity.
 184. 184. Fusion protein or conjugate according to any one of items 181-182, wherein said desired biological activity is a binding activity to a selected target.
 185. 185. Fusion protein or conjugate according to item 184, wherein said selected target is albumin.

186. 186. Fusion protein or conjugate according to item 185, wherein said albumin binding activity is provided by the albumin binding domain of streptococcal protein G, or a derivative thereof.
187. 187. Fusion protein or conjugate according to any one of items 185-189, wherein said albumin binding activity increases *in vivo* half-life of the fusion protein or conjugate.
188. 188. Fusion protein or conjugate according to any one of items 181-182, wherein said desired biological activity is an enzymatic activity.
189. 189. Fusion protein or conjugate according to any one of items 181-183, wherein the second moiety having a desired biological activity is a therapeutically active polypeptide.
190. 190. Fusion protein or conjugate according to any one of items 181-183 and 188-189, wherein the second moiety having a desired biological activity is selected from the group consisting of enzymes, hormones, growth factors, chemokines and cytokines.
191. 191. FcRn binding polypeptide, fusion protein or conjugate according to any preceding item, which inhibits binding of IgG to FcRn.
192. 192. FcRn binding polypeptide, fusion protein or conjugate according to item 191, wherein the K_D value of the interaction between said FcRn binding polypeptide, fusion protein or conjugate and FcRn is lower than the K_D value of the interaction between IgG and FcRn.
193. 193. FcRn binding polypeptide, fusion protein or conjugate according to any preceding item, further comprising a label.
194. 194. FcRn binding polypeptide, fusion protein or conjugate according to item 193, wherein said label is selected from the group consisting of fluorescent dyes and metals, chromophoric dyes, chemiluminescent compounds and bioluminescent proteins, enzymes, radionuclides and particles.
195. 195. FcRn binding polypeptide, fusion protein or conjugate according to any preceding item, comprising a chelating environment provided by a polyaminopolycarboxylate chelator conjugated to the FcRn binding polypeptide via a thiol group of a cysteine residue or an amine group of a lysine residue.
196. 196. FcRn binding polypeptide, fusion protein or conjugate according to item 195, wherein the polyaminopolycarboxylate chelator is 1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid or a derivative thereof.
197. 197. FcRn binding polypeptide, fusion protein or conjugate according to item 196, wherein the 1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid derivative is 1,4,7,10-tetraazacyclododecane-1,4,7-tris-acetic acid-10-maleimidoethylacetamide.
198. 198. FcRn binding polypeptide, fusion protein or conjugate according to item 195, wherein the polyaminopolycarboxylate chelator is 1,4,7-triazacyclononane-1,4,7-triacetic acid or a derivative thereof.
199. 199. FcRn binding polypeptide, fusion protein or conjugate according to item 195, wherein the polyaminopolycarboxylate chelator is diethylenetriaminepentaacetic acid or derivatives thereof.
200. 200. A polynucleotide encoding a polypeptide according to any one of items 1-192.
201. 201. Expression vector comprising a polynucleotide according to item 200.
202. 202. Host cell comprising an expression vector according to item 201.
203. 203. Method of producing a polypeptide according to any one of items 1-192, comprising

- culturing a host cell according to item 202 under conditions permissive of expression of said polypeptide from said expression vector, and
 - isolating said polypeptide.
- 204. 204. Composition comprising an FcRn binding polypeptide, fusion protein or conjugate according to any one of items 1-199 and at least one pharmaceutically acceptable excipient or carrier.
- 205. 205. Composition according to item 204, further comprising at least one additional active agent.
- 206. 206. Composition according to any one of items 204-205, which is adapted for administration by a route selected from the group consisting of oral administration, intranasal administration, pulmonar administration, vaginal administration, rectal administration, intravenous injection, intraperitoneal injection, intramuscular injection, subcutaneous injection and intradermal injection.
- 207. 207. FcRn binding polypeptide, fusion protein or conjugate according to any one of items 1-199 or composition according to any one of items 204-206 for use as a medicament.
- 208. 208. FcRn binding polypeptide, fusion protein, conjugate or composition for use according to item 207, wherein said medicament is intended for treatment of an auto-immune condition.
- 209. 209. FcRn binding polypeptide, fusion protein, conjugate or composition for use according to item 207 or 208, wherein said medicament is intended for treatment of a condition selected from the group consisting of myasthenia gravis, Guillain-Barré syndrome, autoimmune limbic encephalitis, pediatric autoimmune neuropsychiatric disorders associated with streptococcal infection (PANDAS), neuromyotonia (Isaac's syndrome), morvan syndrome, multiple sclerosis, pemphigus vulgaris, foliaceus, bullous pemphigoid, epidermolysis bullosa acquisita, pemphigoid gestationis, mucous membrane pemphigoid, lichen sclerosus, antiphospholipid syndrome, erlapsing polychondritis, autoimmune anemia, idiopathic trombocytic purpura, autoimmune Grave's disease, dilated cardiomyopathy, vasculitis, Goodpasture's syndrome, idiopathic membranous nephropathy, rheumatoid arthritis and systemic lupus erythematosus.
- 210. 210. Method of treatment of a subject in need thereof, comprising administering to the subject a therapeutically active amount of an FcRn binding polypeptide, fusion protein or conjugate according to any one of items 1-199 or composition according to any one of items 204-206.
- 211. 211. Method according to item 210, for treatment of an auto-immune condition.
- 212. 212. Method according to item 210 or 211, wherein said subject is suffering from a condition selected from the group consisting of myasthenia gravis, Guillain-Barré syndrome, autoimmune limbic encephalitis, pediatric autoimmune neuropsychiatric disorders associated with streptococcal infection (PANDAS), neuromyotonia (Isaac's syndrome), morvan syndrome, multiple sclerosis, pemphigus vulgaris, foliaceus, bullous pemphigoid, epidermolysis bullosa acquisita, pemphigoid gestationis, mucous membrane pemphigoid, lichen sclerosus, antiphospholipid syndrome, erlapsing polychondritis, autoimmune anemia, idiopathic trombocytic purpura, autoimmune Grave's disease, dilated cardiomyopathy, vasculitis, Goodpasture's syndrome, idiopathic membranous nephropathy, rheumatoid arthritis and systemic lupus erythematosus.

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- [WO2010054699A \[0011\]](#)
- [WO2009077175A \[0244\] \[0255\]](#)
- [WO2009016043A \[0290\]](#)

Non-patent literature cited in the description

- **SIMISTERMOSTOV** Nature, 1989, vol. 337, 184-7 [\[0002\]](#)
- **BURMEISTER et al.** Nature, 1994, vol. 372, 379-83 [\[0002\]](#)
- **ROOPENIAN** Nat Rev Immunol, 2007, vol. 7, 715-25 [\[0003\]](#)
- **LIU et al.** J Immunol, 2007, vol. 179, 2999-3011 [\[0008\]](#)
- **MEZO et al.** Proc Natl Acad Sci U S A, 2008, vol. 105, 2337-42 [\[0008\]](#)
- **GETMANBALTHASAR** J Pharm Sci, 2005, vol. 94, 718-29 [\[0008\]](#)
- **PETKOVA et al.** Int Immunol, 2006, vol. 18, 1759-69 [\[0008\]](#)
- **VACCARO et al.** Nat Biotechnol, 2005, vol. 23, 1283-8 [\[0008\]](#)
- **VALLES et al.** J Interferon Cytokine Res, 2011, vol. 32, 178-184 [\[0009\]](#)
- **HINTON et al.** J Biol Chem, 2004, vol. 279, 6213-6 [\[0009\]](#)
- **SCHELLENBERGER et al.** Nat Biotechnol, 2009, vol. 21, 1186-1190 [\[0009\]](#)
- **LOW et al.** Human reproduction, 2005, vol. 20, 71805-13 [\[0010\]](#)
- **THOMPSON et al.** Nucleic Acids Research, 1994, vol. 22, 4673-4680 [\[0027\]](#)
- **SUTER et al.** Stem Cells, 2006, vol. 24, 615-623 [\[0231\]](#)
- **TAI et al.** PLoS One, 2012, vol. 7, 9e46269- [\[0231\]](#)
- **ZUFFEREY et al.** Nat Biotechnol, 1997, vol. 15, 9871-5 [\[0233\]](#)
- **JAKOBSSON et al.** J Neurosci Res, 2006, vol. 84, 58-67 [\[0233\]](#)
- **SANDALOVA et al.** Acta Chryst, 2005, vol. F61, 1090-1093 [\[0237\]](#)
- **MICHAËLSSON et al.** J Immunol, 2001, vol. 166, 7327-7334 [\[0237\]](#)

- **GRÖNWALL et al.**J Biotechnol, 2007, vol. 128, 162-183 [\[0243\]](#)
- **RÜTHER et al.**Nucleic Acids Res, 1982, vol. 10, 5765-5772 [\[0243\]](#)
- **JAKOBSSON et al.**J Neurosci Res, 2003, vol. 73, 876-85 [\[0272\]](#)
- **ORLOVA et al.**Cancer Res, 2006, vol. 66, 84339-48 [\[0291\]](#)

P A T E N T K R A V

1. FcRn-bindende polypeptid, omfattende et FcRn-bindende motiv *BM*, hvilket motiv består af aminosyresekvensen

5 $EX_2 X_3 X_4 AX_6 X_7 EIR WLPNLX_{16}X_{17} X_{18} QR X_{21}AFIX_{25} X_{26}LX_{28} X_{29}$

hvor, uafhængigt af hinanden,

X_2 er udvalgt blandt A, D, E, F, H, I, K, L, N, Q, R, S, T, V, W og Y;

X_3 er udvalgt blandt A, D, E, F, G, H, I, K, L, M, N, Q, R, S, T, V, W og Y;

X_4 er udvalgt blandt A, D, E, F, G, H, I, K, L, N, Q, R, S, T, V, W og Y;

10 X_6 er udvalgt blandt A, E, F, G, H, I, K, Q, R, S og V;

X_7 er udvalgt blandt A, F, H, K, N, Q, R, S og V;

X_{16} er udvalgt blandt N og T;

X_{17} er udvalgt blandt F, W og Y;

X_{18} er udvalgt blandt A, D, E og N;

15 X_{21} er udvalgt blandt A, S, V og W;

X_{25} er udvalgt blandt D, E, G, H, I, K, L, N, Q, R, S, T, V, W og Y;

X_{26} er udvalgt blandt K og S;

X_{28} er udvalgt blandt A, D, E, F, H, I, K, L, N, Q, R, S, T, V, W og Y; og

X_{29} er udvalgt blandt D og R.

20 2. FcRn-bindende polypeptid ifølge krav 1, hvor BM består af en aminosyresekvens udvalgt blandt

i)

$EX_2 X_3 X_4 AX_6 HEIR WLPNLTX_{17} X_{18} QR X_{21}AFIX_{25} KLX_{28} D$

hvor, uafhængigt af hinanden,

25 X_2 er udvalgt blandt A, D, E, F, H, I, K, L, N, Q, R, S, T, V, W og Y;

X_3 er udvalgt blandt A, D, E, G, H, K, L, M, N, Q, R, S, T, V og Y;

X_4 er udvalgt blandt A, D, E, F, G, I, K, L, N, Q, R, S, T, V og Y;

X_6 er udvalgt blandt A, G, K, R, S og V;

X_{17} er udvalgt blandt F, W og Y;

30 X_{18} er udvalgt blandt A, D, E og N;

X_{21} er udvalgt blandt A, S, V og W;

X_{25} er udvalgt blandt D, G, H, K, L, N, R, V og W;

X_{28} er udvalgt blandt A, D, E, H, K, L, N, Q, R, S, T, W og Y;

og

35 ii) en aminosyresekvens, som har mindst 96 % similaritet med nævnte sekvens.

3. FcRn-bindende polypeptid ifølge et hvilket som helst af de foregående krav, hvor sekvensen i) opfylder mindst tre af de seks betingelser I-VI:

I. X_6 er udvalgt blandt A, G, K og S, såsom især A;

II. X_7 er H;

40 III. X_{17} er udvalgt blandt F og Y, såsom især F;

IV. X_{18} er D;

V. X_{21} er udvalgt blandt V og W, såsom især V;

VI. X_{25} er udvalgt blandt H og R, såsom især H.

4. FcRn-bindende polypeptid ifølge et hvilket som helst af de foregående krav, hvor sekvensen er udvalgt fra gruppen bestående af SEQ ID NO:1-353, såsom udvalgt fra
- 5 gruppen bestående af SEQ ID NO:1-15, SEQ ID NO:17-140 og SEQ ID NO:353, såsom udvalgt fra gruppen bestående af SEQ ID NO:1-2 og SEQ ID NO:17-140, såsom udvalgt fra gruppen bestående af SEQ ID NO:1-2, SEQ ID NO:17-92, SEQ ID NO:94-103, SEQ ID NO:105-125 og SEQ ID NO:127-140 eller udvalgt fra gruppen bestående af SEQ ID NO:1-8, SEQ ID NO:13, SEQ ID NO:19-20, SEQ ID NO:23, SEQ ID NO:28, SEQ ID NO:41, SEQ
- 10 ID NO:44, SEQ ID NO:65, SEQ ID NO:70, SEQ ID NO:73, SEQ ID NO:75-77 og SEQ ID NO:353, såsom udvalgt fra gruppen bestående af SEQ ID NO:1, SEQ ID NO:23, SEQ ID NO:28, SEQ ID NO:41, SEQ ID NO:44, SEQ ID NO:65, SEQ ID NO:73 og SEQ ID NO:75-77, såsom udvalgt fra gruppen bestående af SEQ ID NO:1, SEQ ID NO:23, SEQ ID NO:44, SEQ ID NO:65, SEQ ID NO:75 og SEQ ID NO:77, såsom udvalgt fra gruppen bestående af
- 15 SEQ ID NO:1, SEQ ID NO:23 og SEQ ID NO:75, såsom SEQ ID NO:1.

5. FcRn-bindende polypeptid ifølge et hvilket som helst af de foregående krav, som omfatter en aminosyresekvens udvalgt blandt:

iii)

$K-[BM]-DPSQS X_a X_b LLX_c EAKKL X_d X_e X_f Q;$

20 hvor

[BM] er et FcRn-bindingsmotiv som defineret heri, på den betingelse, at X_{29} er D;

X_a er udvalgt blandt A og S;

X_b er udvalgt blandt N og E;

X_c er udvalgt blandt A, S og C;

25 X_d er udvalgt blandt E, N og S;

X_e er udvalgt blandt D, E og S;

X_f er udvalgt blandt A og S;

og

iv) en aminosyresekvens, som har mindst 93 % similaritet med en sekvens defineret af

30 iii);

hvor nævnte sekvens iii) eventuelt er udvalgt fra gruppen bestående af SEQ ID NO:354-706, såsom udvalgt fra gruppen bestående af SEQ ID NO:354-368, SEQ ID NO:370-493 og SEQ ID NO:706, såsom udvalgt fra gruppen bestående af SEQ ID NO:354-355 og SEQ ID NO:370-493, såsom udvalgt fra gruppen bestående af SEQ ID NO:354-355, SEQ ID

35 NO:370-445, SEQ ID NO:447-456, SEQ ID NO:458-478 og SEQ ID NO:480-493 eller udvalgt fra gruppen bestående af SEQ ID NO:354-361, SEQ ID NO:366, SEQ ID NO:372-373, SEQ ID NO:376, SEQ ID NO:381, SEQ ID NO:394, SEQ ID NO:397, SEQ ID NO:418, SEQ ID NO:423, SEQ ID NO:426, SEQ ID NO:428-430 og SEQ ID NO:706, såsom udvalgt fra gruppen bestående af SEQ ID NO:354, SEQ ID NO:376, SEQ ID NO:381, SEQ ID

NO:394, SEQ ID NO:397, SEQ ID NO:418, SEQ ID NO:426 og SEQ ID NO:428-430, såsom udvalgt fra gruppen bestående af SEQ ID NO:354, SEQ ID NO:376, SEQ ID NO:397, SEQ ID NO:418, SEQ ID NO:428 og SEQ ID NO:430, såsom udvalgt fra gruppen bestående af SEQ ID NO:354, SEQ ID NO:376 og SEQ ID NO:428, såsom SEQ ID NO:354.

- 5 6. FcRn-bindende polypeptid ifølge et hvilket som helst af de foregående krav, som omfatter en aminosyresekvens udvalgt blandt:

xi)

AEAKYAK-[BM]-DPSQSSELLSEAKKLNDSQAPK;

hvor [BM] er et FcRn-bindingsmotiv som defineret i et hvilket som helst af kravene 1-4; og

- 10 xii) en aminosyresekvens, som har mindst 94 % similaritet med sekvensen defineret i xi);
hvor sekvensen xi) eventuelt er udvalgt fra gruppen bestående af SEQ ID NO:1060-1062.

7. FcRn-bindende polypeptid ifølge et hvilket som helst af de foregående krav, som omfatter en aminosyresekvens udvalgt blandt:

xiii)

- 15 VDAKYAK-[BM]-DPSQSSELLSEAKKLNDSQAPK;

hvor [BM] er et FcRn-bindende motiv som defineret i et hvilket som helst af kravene 1-4; og

- xiv) en aminosyresekvens, som har mindst 94 % similaritet med sekvensen defineret i
xiii);
20 hvor sekvensen xiii) eventuelt er udvalgt fra gruppen bestående af SEQ ID NO:707-1059, såsom udvalgt fra gruppen bestående af SEQ ID NO:707-721, SEQ ID NO:723-846 og SEQ ID NO:1059, såsom udvalgt fra gruppen bestående af SEQ ID NO:707-708 og SEQ ID NO:723-846, såsom udvalgt fra gruppen bestående af SEQ ID NO:707-708, SEQ ID NO:723-798, SEQ ID NO:800-809, SEQ ID NO:811-831 og SEQ ID NO:833-846 eller
25 udvalgt fra gruppen bestående af SEQ ID NO:707-714, SEQ ID NO:719, SEQ ID NO:725-726, SEQ ID NO:729, SEQ ID NO:734, SEQ ID NO:747, SEQ ID NO:750, SEQ ID NO:771, SEQ ID NO:776, SEQ ID NO:779, SEQ ID NO:781-783 og SEQ ID NO:1059, såsom udvalgt fra gruppen bestående af SEQ ID NO:707, SEQ ID NO:729, SEQ ID NO:734, SEQ ID NO:747, SEQ ID NO:750, SEQ ID NO:771, SEQ ID NO:779 og SEQ ID NO:781-783, såsom
30 udvalgt fra gruppen bestående af SEQ ID NO:707, SEQ ID NO:729, SEQ ID NO:750, SEQ ID NO:771, SEQ ID NO:781 og SEQ ID NO:783, såsom udvalgt fra gruppen bestående af SEQ ID NO:707, SEQ ID NO:729 og SEQ ID NO:781, såsom SEQ ID NO:707.

8. FcRn-bindende polypeptid ifølge et hvilket som helst af de foregående krav,

- som er stand til at binde sig til FcRn ved pH 6,0, således at K_D -værdien for interaktionen
35 er højst 1×10^{-6} M, såsom højst 1×10^{-7} M, såsom højst 1×10^{-8} M, såsom højst 1×10^{-9} M, såsom højst 1×10^{-10} M; og/eller
- hvor K_D -værdien for interaktionen mellem FcRn-bindende polypeptid og FcRn ved pH 7,4 er højere end K_D -værdien for nævnte interaktion ved pH 6,0, såsom mindst 2 gange højere, såsom mindst 5 gange højere, såsom mindst 10 gange højere, såsom mindst 50

gange højere, såsom mindst 100 gange højere end K_D -værdien for nævnte interaktion ved pH 6,0; og/eller

- hvor K_D -værdien for interaktionen mellem FcRn-bindende polypeptid og FcRn ved pH 7,4 er mindst 1×10^{-8} M, såsom mindst 1×10^{-7} M, såsom mindst 1×10^{-6} M, såsom mindst 1×10^{-5} M; og/eller

- hvor K_D -værdien for interaktionen mellem FcRn-bindende polypeptid og FcRn ved pH 7,4 er højst 1×10^{-6} M, såsom højst 1×10^{-7} M, såsom højst 1×10^{-8} M, såsom højst 1×10^{-9} M, såsom højst 1×10^{-10} M.

9. Fusionsprotein eller konjugat omfattende

10 - en første del bestående af et FcRn-bindende polypeptid ifølge et hvilket som helst af de foregående krav; og

- en anden del bestående af et polypeptid med en ønsket biologisk aktivitet.

10. FcRn-bindende polypeptid, fusionsprotein eller konjugat ifølge et hvilket som helst af de foregående krav, som hæmmer binding af IgG til FcRn.

15 11. Polynukleotid kodende for et polypeptid ifølge et hvilket som helst af kravene 1-10.

12. Sammensætning omfattende et FcRn-bindende polypeptid, fusionsprotein eller konjugat ifølge et hvilket som helst af kravene 1-10 og mindst en farmaceutisk acceptabel excipients eller bærestof.

20 13. FcRn-bindende polypeptid, fusionsprotein eller konjugat ifølge et hvilket som helst af kravene 1-10 eller sammensætning ifølge krav 12 til anvendelse som lægemiddel.

14. FcRn-bindende polypeptid, fusionsprotein, konjugat eller sammensætning til anvendelse ifølge krav 13, hvor lægemidlet er tiltænkt behandling af en autoimmun tilstand.

25 15. FcRn-bindende polypeptid, fusionsprotein, konjugat eller sammensætning til anvendelse ifølge krav 13 eller 14, hvor lægemidlet er tiltænkt behandling af en tilstand udvalgt fra gruppen bestående af Erb-Goldflam-syndrom, polyradikuloneurit, autoimmun limbisk encefalit, pædiatriske autoimmune neuropsykiatriske forstyrrelser forbundet med streptokokinfektion (PANDAS), neuromyotoni, Morvans syndrom, dissemineret sklerose, pemphigus vulgaris, foliaceus, bulløs pemfigoid, erhvervet bulløs epidermolysis, pemfigoid gestationis, slimhinde-pemfigoid, lichen sclerosus, antiphospholipid-syndrom, recidiverende polychondrit, autoimmun anæmi, idiopatisk trombocyt-purpura, autoimmun basedow-syge, dilateret kardiomyopati, vaskulitis, Goodpastures syndrom, idiopatisk membran-nefropati, leddegigt og systemisk lupus erythematosus.

DRAWINGS

Figure 1A

Designation	Sequence	SEQ ID NO:
BM07918	EQDAAAHEIRWLPNLTFDQQRVAFIHKLD	SEQ ID NO:1
BM10193	EWMDAAHEIRWLPNLTFDQQRVAFIHKLE	SEQ ID NO:2
BM10109	EANTAAAHEIRWLPNLTFDQQRVAFIRKLWD	SEQ ID NO:3
BM07960	EFESAAHEIRWLPNLTQDQQRVAFIHKLS	SEQ ID NO:4
BM10140	ERSAAAHAEIRWLPNLTFDQQRVAFILKLT	SEQ ID NO:5
BM07930	ESDSAVHEIRWLPNLTFDQQRVAFIHKLL	SEQ ID NO:6
BM10183	EADNAGHEIRWLPNLTWAQRVAFIHKLL	SEQ ID NO:7
BM10111	EDDTAAHEIRWLPNLTQEQRVAFIHKLY	SEQ ID NO:8
BM10129	EQHDAAEIRWLPNLTQDQQRVAFIRKLH	SEQ ID NO:9
BM10141	ENQGAHAEIRWLPNLTWDQQRVAFIRKLQD	SEQ ID NO:10
BM10156	ERTQASHEIRWLPNLTQDQQRVAFINKLR	SEQ ID NO:11
BM10127	ERKDAGHEIRWLPNLTQDQRSAFIKKLE	SEQ ID NO:12
BM07909	ERQEAHAEIRWLPNLTQDQQRVAFIVKLL	SEQ ID NO:13
BM10152	EEDVAAHEIRWLPNLTFNQRAAFIDKLL	SEQ ID NO:14
BM10145	ENQDAAHEIRWLPNLTQDQQRVAFIGKLN	SEQ ID NO:15
BM10161	ESGYAVHEIRWLPNLTQDQQRVAFINKLQ	SEQ ID NO:16
BM13573	ESKDAHAEIRWLPNLTQDQQRVAFIHKLR	SEQ ID NO:17
BM13574	EKKEAAHEIRWLPNLTQDQQRVAFIHKLR	SEQ ID NO:18
BM13577	EWHOAAHEIRWLPNLTQDQQRVAFIHKLR	SEQ ID NO:19
BM13578	EWTDAAHEIRWLPNLTQDQQRVAFIHKLR	SEQ ID NO:20
BM13579	EISAAHAEIRWLPNLTQDQQRVAFIHKLR	SEQ ID NO:21
BM13581	EQQAAHAEIRWLPNLTQDQQRVAFIHKLR	SEQ ID NO:22
BM13583	ELEKAGHEIRWLPNLTQDQQRVAFIHKLR	SEQ ID NO:23
BM13585	EYLDAAHEIRWLPNLTQDQQRVAFIHKLR	SEQ ID NO:24
BM13586	ELKDAGHEIRWLPNLTQDQQRVAFIHKLR	SEQ ID NO:25
BM13587	EHVDAHAEIRWLPNLTQDQQRVAFIHKLR	SEQ ID NO:26
BM13588	EYAAAAHAEIRWLPNLTQDQQRVAFIHKLR	SEQ ID NO:27
BM13592	EVDIAHAEIRWLPNLTQDQQRVAFIHKLR	SEQ ID NO:28

Figure 1B

Designation	Sequence	SEQ ID NO:
BM13594	EIDEAKHEIRWLPNLTFFDQRVAFIHKLDD	SEQ ID NO:29
BM13596	ELRQAGHEIRWLPNLTFFDQRVAFIHKLRD	SEQ ID NO:30
BM13597	ELQSAHEIRWLPNLTFFDQRVAFIHKIKD	SEQ ID NO:31
BM13598	ELEKAAHEIRWLPNLTFFDQRVAFIHKLRD	SEQ ID NO:32
BM13600	EAHEAAHEIRWLPNLTFFDQRVAFIHKLRD	SEQ ID NO:33
BM13604	ELQAASHEIRWLPNLTFFDQRVAFIHKLRD	SEQ ID NO:34
BM13605	EIESAKHEIRWLPNLTFFDQRVAFIHKLND	SEQ ID NO:35
BM13609	EWKVAHEIRWLPNLTFFDQRVAFIHKLRD	SEQ ID NO:36
BM13611	EWKAAAHEIRWLPNLTFFDQRVAFIHKLRD	SEQ ID NO:37
BM13612	EIDLAKHEIRWLPNLTFFDQRVAFIHKLDD	SEQ ID NO:38
BM13613	ELEAARHEIRWLPNLTFFDQRVAFIHKLND	SEQ ID NO:39
BM13615	EAATAGHEIRWLPNLTFFDQRVAFIHKLRD	SEQ ID NO:40
BM13616	EWQQAHEIRWLPNLTFFDQRVAFIHKLRD	SEQ ID NO:41
BM13617	EADQAGHEIRWLPNLTFFDQRVAFIRKLRD	SEQ ID NO:42
BM13620	EQSKAAHEIRWLPNLTFFDQRVAFIHKLRD	SEQ ID NO:43
BM13621	EADAAGHEIRWLPNLTFFDQRVAFIHKLRD	SEQ ID NO:44
BM13622	EFMDAGHEIRWLPNLTFFDQRVAFIHKLRD	SEQ ID NO:45
BM13624	ESKQAAHEIRWLPNLTFFDQRVAFIHKLRD	SEQ ID NO:46
BM13625	EVSDAAHEIRWLPNLTFFDQRVAFIHKLRD	SEQ ID NO:47
BM13626	EADSASHEIRWLPNLTFFDQRVAFIHKLRD	SEQ ID NO:48
BM13627	ELMEAAHEIRWLPNLTFFDQRVAFIHKLRD	SEQ ID NO:49
BM13628	ELNTAAHEIRWLPNLTFFDQRVAFIHKLKD	SEQ ID NO:50
BM13629	EVHEAGHEIRWLPNLTFFDQRVAFIHKLRD	SEQ ID NO:51
BM13633	ESTAAAHEIRWLPNLTFFDQRVAFIHKLRD	SEQ ID NO:52
BM13634	EWYNAHEIRWLPNLTFFDQRVAFIHKLRD	SEQ ID NO:53
BM13635	EWNDAAHEIRWLPNLTFFDQRVAFIHKLRD	SEQ ID NO:54
BM13637	EVEVAKHEIRWLPNLTFFDQRVAFIHKLDD	SEQ ID NO:55
BM13638	EENFAAHEIRWLPNLTFFDQRVAFIHKLRD	SEQ ID NO:56

Figure 1C

Designation	Sequence	SEQ ID NO:
BM13639	EHDSAAHEIRWLPNLTFFDQRFVAFIHKLRD	SEQ ID NO:57
BM13640	EWMDAAHEIRWLPNLTFFDQRFVAFIHKLRD	SEQ ID NO:58
BM13641	EFSAAGHEIRWLPNLTFFDQRFVAFIHKLRD	SEQ ID NO:59
BM13644	ELNSAAHEIRWLPNLTFFDQRFVAFIHKLRD	SEQ ID NO:60
BM13645	EVDTAAGHEIRWLPNLTFFDQRFVAFIHKLRD	SEQ ID NO:61
BM13648	ESQIAGHEIRWLPNLTFFDQRFVAFIHKLRD	SEQ ID NO:62
BM13651	EVSAASHEIRWLPNLTFFDQRFVAFIHKLRD	SEQ ID NO:63
BM13652	EDQDAAGHEIRWLPNLTFFDQRFVAFIHKLRD	SEQ ID NO:64
BM13654	ELEAAKHEIRWLPNLTFFDQRFVAFIHKLRD	SEQ ID NO:65
BM13655	ESKRAAGHEIRWLPNLTFFDQRFVAFIHKLRD	SEQ ID NO:66
BM13656	EYVKAGHEIRWLPNLTFFDQRFVAFIHKLRD	SEQ ID NO:67
BM13657	EFSRAGHEIRWLPNLTFFDQRFVAFIHKLRD	SEQ ID NO:68
BM13659	EWQFAAGHEIRWLPNLTFFDQRFVAFIHKLRD	SEQ ID NO:69
BM13663	EWQIAAGHEIRWLPNLTFFDQRFVAFIHKLRD	SEQ ID NO:70
BM13664	ELQEAAGHEIRWLPNLTFFDQRFVAFIHKLRD	SEQ ID NO:71
BM13667	EYRAAGHEIRWLPNLTFFDQRFVAFIHKLRD	SEQ ID NO:72
BM13669	ELASAGHEIRWLPNLTFFDQRFVAFIHKLRD	SEQ ID NO:73
BM13672	EVQSASHEIRWLPNLTFFDQRFVAFIHKLRD	SEQ ID NO:74
BM13674	EIEDAGHEIRWLPNLTFFDQRFVAFIHKLRD	SEQ ID NO:75
BM13675	ENQAAAGHEIRWLPNLTFFDQRFVAFIHKLRD	SEQ ID NO:76
BM13676	EATSAGHEIRWLPNLTFFDQRFVAFIVKLRD	SEQ ID NO:77
BM13678	EDEQAAGHEIRWLPNLTFFDQRFVAFIHKLRD	SEQ ID NO:78
BM13684	EQNQAGHEIRWLPNLTFFDQRFVAFIHKLRD	SEQ ID NO:79
BM13688	EYTSAAHEIRWLPNLTFFDQRFVAFIHKLRD	SEQ ID NO:80
BM13691	EWDAAGHEIRWLPNLTFFDQRFVAFIHKLRD	SEQ ID NO:81
BM13692	EEMQAGHEIRWLPNLTFFDQRFVAFIHKLRD	SEQ ID NO:82
BM13694	ELSDAAGHEIRWLPNLTFFDQRFVAFIHKLRD	SEQ ID NO:83
BM13695	EIDAAKHEIRWLPNLTFFDQRFVAFIHKLRD	SEQ ID NO:84

Figure 1D

Designation	Sequence	SEQ ID NO:
BM13697	EAERAGHEIRWLPNLTFFDQRVAFIRKLRD	SEQ ID NO:85
BM13706	EEDSAAHEIRWLPNLTFFDQRVAFIHKLRD	SEQ ID NO:86
BM13708	EQKAAGHEIRWLPNLTFFDQRVAFIHKLRD	SEQ ID NO:87
BM13710	EWQDAAHEIRWLPNLTFFDQRVAFIHKLRD	SEQ ID NO:88
BM13711	EAKAAAHEIRWLPNLTFFDQRVAFIHKLRD	SEQ ID NO:89
BM13714	ELSEAAHEIRWLPNLTFFDQRVAFIHKLRD	SEQ ID NO:90
BM13716	ETEAAKHEIRWLPNLTFFDQRVAFIHKLRD	SEQ ID NO:91
BM13719	EAKSAAHEIRWLPNLTFFDQRVAFIHKLRD	SEQ ID NO:92
BM13720	EQSAAAHEIRWLPNLTFFDQRVAFIHKLRD	SEQ ID NO:93
BM13721	EKERAAHEIRWLPNLTFFDQRVAFIHKLRD	SEQ ID NO:94
BM13725	EWDEAAHEIRWLPNLTFFDQRVAFIHKLRD	SEQ ID NO:95
BM13727	EKDDAAHEIRWLPNLTFFDQRVAFIHKLRD	SEQ ID NO:96
BM13728	EIENAAHEIRWLPNLTFFDQRVAFIHKLRD	SEQ ID NO:97
BM13732	ETKEAAHEIRWLPNLTFFDQRVAFIHKLRD	SEQ ID NO:98
BM13735	ELEAAKHEIRWLPNLTFFDQRVAFIHKLRD	SEQ ID NO:99
BM13736	EWAEAAHEIRWLPNLTFFDQRVAFIHKLRD	SEQ ID NO:100
BM13740	ESQEAGHEIRWLPNLTFFDQRVAFIHKLRD	SEQ ID NO:101
BM13742	ELSTAAHEIRWLPNLTFFDQRVAFIHKLRD	SEQ ID NO:102
BM13747	EIEEAGHEIRWLPNLTFFDQRVAFIHKLRD	SEQ ID NO:103
BM13749	ELQTASHEIRWLPNLTFFDQRVAFIHKLRD	SEQ ID NO:104
BM13750	EQDSAAHEIRWLPNLTFFDQRVAFIRKLRD	SEQ ID NO:105
BM13751	ESASAGHEIRWLPNLTFFDQRVAFIHKLRD	SEQ ID NO:106
BM13752	EVAKASHEIRWLPNLTFFDQRVAFIHKLRD	SEQ ID NO:107
BM13758	EVQEEAAHEIRWLPNLTFFDQRVAFIHKLRD	SEQ ID NO:108
BM13759	ESYEAAHEIRWLPNLTFFDQRVAFIHKLRD	SEQ ID NO:109
BM13760	ETAEEAGHEIRWLPNLTFFDQRVAFIHKLRD	SEQ ID NO:110
BM13761	ELEEEAAHEIRWLPNLTFFDQRVAFIHKLRD	SEQ ID NO:111
BM13771	EAAAAGHEIRWLPNLTFFDQRVAFIHKLRD	SEQ ID NO:112

Figure 1E

Designation	Sequence	SEQ ID NO:
BM13773	EYVDAGHEIRWLPNLTFFDQRVAFIRKLRD	SEQ ID NO:113
BM13776	EIQEAGHEIRWLPNLTFFDQRVAFIHKLRD	SEQ ID NO:114
BM13777	ESATAAHEIRWLPNLTFFDQRVAFIHKLRD	SEQ ID NO:115
BM13780	EWMSAAHEIRWLPNLTFFDQRVAFIHKLRD	SEQ ID NO:116
BM13782	EREQASHEIRWLPNLTFFDQRVAFIHKLRD	SEQ ID NO:117
BM13783	EIEQAGHEIRWLPNLTFFDQRVAFIHKLRD	SEQ ID NO:118
BM13786	EHNAAAHEIRWLPNLTFFDQRVAFIHKLRD	SEQ ID NO:119
BM13792	EIEVAAHEIRWLPNLTFFDQRVAFIHKLRD	SEQ ID NO:120
BM13796	ERAEAAHEIRWLPNLTFFDQRVAFIHKLRD	SEQ ID NO:121
BM13799	ESELAHEIRWLPNLTFFDQRVAFIHKLRD	SEQ ID NO:122
BM13806	EYRAAGHEIRWLPNLTFFDQRVAFIHKLRD	SEQ ID NO:123
BM13808	ETANAAHEIRWLPNLTFFDQRVAFIHKLRD	SEQ ID NO:124
BM13811	EWYEAAHEIRWLPNLTFFDQRVAFIHKLRD	SEQ ID NO:125
BM13812	EEQEAAHEIRWLPNLTFFDQRVAFIHKLRD	SEQ ID NO:126
BM13823	EHDDAAHEIRWLPNLTFFDQRVAFIHKLRD	SEQ ID NO:127
BM13824	EWYSAAHEIRWLPNLTFFDQRVAFIHKLRD	SEQ ID NO:128
BM13838	EISDAAHEIRWLPNLTFFDQRVAFIHKLRD	SEQ ID NO:129
BM13840	EYTTAAHEIRWLPNLTFFDQRVAFIHKLRD	SEQ ID NO:130
BM13842	EISQAAHEIRWLPNLTFFDQRVAFIHKLRD	SEQ ID NO:131
BM13845	ENDDAAHEIRWLPNLTFFDQRVAFIHKLRD	SEQ ID NO:132
BM13846	ESELAGHEIRWLPNLTFFDQRVAFIRKLRD	SEQ ID NO:133
BM13848	EQADAAHEIRWLPNLTFFDQRVAFIHKLRD	SEQ ID NO:134
BM13849	ETESAAHEIRWLPNLTFFDQRVAFIHKLRD	SEQ ID NO:135
BM13860	EISDAKHEIRWLPNLTFFDQRVAFIHKLRD	SEQ ID NO:136
BM13865	EHLNAAHEIRWLPNLTFFDQRVAFIHKLRD	SEQ ID NO:137
BM13866	EWLDAAHEIRWLPNLTFFDQRVAFIHKLRD	SEQ ID NO:138
BM13875	ENAAAAHEIRWLPNLTFFDQRVAFIHKLRD	SEQ ID NO:139
BM13879	EAELAGHEIRWLPNLTFFDQRVAFIRKLRD	SEQ ID NO:140

Figure 1F

Designation	Sequence	SEQ ID NO:
BM13480	EVTDAHEIRWLPNLTTFDQRFVAFIEKLKD	SEQ ID NO:141
BM13481	ELDSASHEIRWLPNLTTFDQRFVAFINKLLD	SEQ ID NO:142
BM13482	EINLAKHEIRWLPNLTTFDQRFVAFIEKIND	SEQ ID NO:143
BM13483	ESEVAKHEIRWLPNLTTFDQRFVAFIHKLSD	SEQ ID NO:144
BM13484	ESAEAGHEIRWLPNLTTFDQRFVAFIHKLQD	SEQ ID NO:145
BM13485	EYSNAAHEIRWLPNLTTFDQRFVAFIDKLSD	SEQ ID NO:146
BM13487	ETNNAGHEIRWLPNLTTFDQRFVAFIIKLRD	SEQ ID NO:147
BM13488	EVEFAAHEIRWLPNLTTFDQRFVAFIHKLND	SEQ ID NO:148
BM13489	EVELAGHEIRWLPNLTTFDQRFVAFIEKLHD	SEQ ID NO:149
BM13490	EVLKASHEIRWLPNLTTFDQRFVAFITKLQD	SEQ ID NO:150
BM13491	EIANAGHEIRWLPNLTTFDQRFVAFIRKLDD	SEQ ID NO:151
BM13493	EYMKAGHEIRWLPNLTTFDQRFVAFIVKLRD	SEQ ID NO:152
BM13495	EHANAQHEIRWLPNLTTFDQRFVAFIHKLQD	SEQ ID NO:153
BM13496	EVDIASHEIRWLPNLTTFDQRFVAFIWKLRD	SEQ ID NO:154
BM13497	EEVFAAHEIRWLPNLTTFDQRFVAFIVKLRD	SEQ ID NO:155
BM13499	EFNTAAHEIRWLPNLTTFDQRFVAFINKLDD	SEQ ID NO:156
BM13501	EVDVAGHEIRWLPNLTTFDQRFVAFIRKLND	SEQ ID NO:157
BM13502	EWSLAAHEIRWLPNLTTFDQRFVAFIVKLRD	SEQ ID NO:158
BM13503	ELDDAAHEIRWLPNLTTFDQRFVAFIQKLQD	SEQ ID NO:159
BM13505	ERHEAGHEIRWLPNLTTFDQRFVAFIRKLND	SEQ ID NO:160
BM13506	EISDAIHEIRWLPNLTTFDQRFVAFIHKLQD	SEQ ID NO:161
BM13507	EWETAGHEIRWLPNLTTFDQRFVAFIVKLSD	SEQ ID NO:162
BM13508	ERYWASHEIRWLPNLTTFDQRFVAFIHKLND	SEQ ID NO:163
BM13512	EIDWAGHEIRWLPNLTTFDQRFVAFIHKLLD	SEQ ID NO:164
BM13515	EQSKAGHEIRWLPNLTTFDQRFVAFIHKLKD	SEQ ID NO:165
BM13518	EIEAAQHEIRWLPNLTTFDQRFVAFINKLRD	SEQ ID NO:166
BM13519	EHEQAAHEIRWLPNLTTFDQRFVAFIRKLVD	SEQ ID NO:167
BM13520	EAEQAGHEIRWLPNLTTFDQRFVAFINKLQD	SEQ ID NO:168

Figure 1G

Designation	Sequence	SEQ ID NO:
BM13522	EVYAAHEIRWLPNLTFFDQRFVAFIHKLWD	SEQ ID NO:169
BM13525	EYSAAGHEIRWLPNLTFFDQRFVAFIEKLRD	SEQ ID NO:170
BM13561	ELATASHEIRWLPNLTFFDQRFVAFIHKLRD	SEQ ID NO:171
BM13572	EYRVAGHEIRWLPNLTFFDQRFVAFIQKLRD	SEQ ID NO:172
BM13575	EVVSAKHEIRWLPNLTFFDQRFVAFIHKLDD	SEQ ID NO:173
BM13576	ESAQAHEIRWLPNLTFFDQRFVAFIRKLRD	SEQ ID NO:174
BM13584	EYSAAGHEIRWLPNLTFFDQRFVAFIHKLRD	SEQ ID NO:175
BM13589	EQKEAAAEIRWLPNLTFFDQRFVAFIHKLRD	SEQ ID NO:176
BM13590	EAAIAGKEIRWLPNLTFFDQRFVAFIHKLRD	SEQ ID NO:177
BM13591	EISKAKHEIRWLPNLTFFDQRFVAFIHKLDD	SEQ ID NO:178
BM13593	ESVAAAHEIRWLPNLTFFDQRFVAFIWKLRD	SEQ ID NO:179
BM13595	EIQQAKHEIRWLPNLTFFDQRFVAFIHKLDD	SEQ ID NO:180
BM13599	EITSAKHEIRWLPNLTFFDQRFVAFIHKLDD	SEQ ID NO:181
BM13601	EQDVAHEIRWLPNLTFFDQRFVAFIRKLRD	SEQ ID NO:182
BM13603	ELERAAHEIRWLPNLTFFDQRFVAFINKLRD	SEQ ID NO:183
BM13607	ENQLAAHEIRWLPNLTFFDQRFVAFIHKLRD	SEQ ID NO:184
BM13608	EISQAKHEIRWLPNLTFFDQRFVAFIHKLDD	SEQ ID NO:185
BM13610	EIANASHEIRWLPNLTFFDQRFVAFIYKLRD	SEQ ID NO:186
BM13614	EWQAAAHEIRWLPNLTFFDQRFVAFIRKLRD	SEQ ID NO:187
BM13618	ERKDAGHEIRWLPNLTFFDQRFVAFIRKLRD	SEQ ID NO:188
BM13619	EITQAKHEIRWLPNLTFFDQRFVAFIHKLDD	SEQ ID NO:189
BM13623	EFIQAGHEIRWLPNLTFFDQRFVAFIHKLRD	SEQ ID NO:190
BM13630	EWNTASHEIRWLPNLTFFDQRFVAFIHKLRD	SEQ ID NO:191
BM13631	EKFVAAHEIRWLPNLTFFDQRFVAFIHKLRD	SEQ ID NO:192
BM13632	EADSAGAEIRWLPNLTFFDQRFVAFIHKLRD	SEQ ID NO:193
BM13636	ESSVAAAHEIRWLPNLTFFDQRFVAFIHKLRD	SEQ ID NO:194
BM13642	EVDLAGREIRWLPNLTFFDQRFVAFIHKLRD	SEQ ID NO:195
BM13643	EQERAAAEIRWLPNLTFFDQRFVAFIHKLRD	SEQ ID NO:196

Figure 1H

Designation	Sequence	SEQ ID NO:
BM13646	EIWQAKHEIRWLPNLTFFDQRVAFIHKLDD	SEQ ID NO:197
BM13647	ELNQAQHEIRWLPNLTFFDQRVAFIHKLED	SEQ ID NO:198
BM13649	ELQQASHEIRWLPNLTFFDQRVAFIHKLRD	SEQ ID NO:199
BM13650	EINQAQHEIRWLPNLTFFDQRVAFIHKLDD	SEQ ID NO:200
BM13653	ELVLACHHEIRWLPNLTFFDQRVAFIHKLRD	SEQ ID NO:201
BM13658	ELTSAAHEIRWLPNLTFFDQRVAFIHKLRD	SEQ ID NO:202
BM13661	EWNAAAHEIRWLPNLTFFDQRVAFIHKLRD	SEQ ID NO:203
BM13662	EILHAKHEIRWLPNLTFFDQRVAFIHKLDD	SEQ ID NO:204
BM13665	EVLTAQHEIRWLPNLTFFDQRVAFIHKLND	SEQ ID NO:205
BM13666	ENSKAAHEIRWLPNLTFFDQRVAFIHKLAD	SEQ ID NO:206
BM13668	EVMTAKHEIRWLPNLTFFDQRVAFIHKLDD	SEQ ID NO:207
BM13671	EARDAQHEIRWLPNLTFFDQRVAFIHKLSD	SEQ ID NO:208
BM13673	ERSKAAHEIRWLPNLTFFDQRVAFIHKLKD	SEQ ID NO:209
BM13677	EIYSAKHEIRWLPNLTFFDQRVAFIHKLDD	SEQ ID NO:210
BM13679	EVQSAKHEIRWLPNLTFFDQRVAFIHKLDD	SEQ ID NO:211
BM13680	ETLEAAHEIRWLPNLTFFDQRVAFIHKLKD	SEQ ID NO:212
BM13681	EQMRAAHEIRWLPNLTFFDQRVAFIRKLRD	SEQ ID NO:213
BM13685	ENKNAQHEIRWLPNLTFFDQRVAFIHKLRD	SEQ ID NO:214
BM13687	ETESAKHEIRWLPNLTFFDQRVAFIHKLTD	SEQ ID NO:215
BM13689	ETVQAQHEIRWLPNLTFFDQRVAFIHKLDD	SEQ ID NO:216
BM13693	EIASAKHEIRWLPNLTFFDQRVAFIHKLDD	SEQ ID NO:217
BM13698	EVMDAKHEIRWLPNLTFFDQRVAFIHKLDD	SEQ ID NO:218
BM13699	ETDAAKHEIRWLPNLTFFDQRVAFIHKLRD	SEQ ID NO:219
BM13701	ELQIAKHEIRWLPNLTFFDQRVAFIHKLDD	SEQ ID NO:220
BM13702	EWKDAQHEIRWLPNLTFFDQRVAFIHKLRD	SEQ ID NO:221
BM13703	ERDSAQHEIRWLPNLTFFDQRVAFIRKLRD	SEQ ID NO:222
BM13704	EIAAAKHEIRWLPNLTFFDQRVAFIHKLDD	SEQ ID NO:223
BM13707	ESVKAKHEIRWLPNLTFFDQRVAFIHKLDD	SEQ ID NO:224

Figure 11

Designation	Sequence	SEQ ID NO:
BM13709	ENERAKHEIRWLPNLTFFDQRVAFIHKLDD	SEQ ID NO:225
BM13712	EYKRAGHEIRWLPNLTFFDQRVAFIRKLRD	SEQ ID NO:226
BM13713	EVRAASHEIRWLPNLTFFDQRVAFIHKLRD	SEQ ID NO:227
BM13717	EDKRAAHEIRWLPNLTFFDQRVAFIRKLRD	SEQ ID NO:228
BM13718	ESEKAGHEIRWLPNLTFFDQRVAFIRKLRD	SEQ ID NO:229
BM13722	EINRAKHEIRWLPNLTFFDQRVAFIHKLDD	SEQ ID NO:230
BM13724	ETQQASHEIRWLPNLTFFDQRVAFIHKLRD	SEQ ID NO:231
BM13729	ENQSAHEIRWLPNLTFFDQRVAFIRKLRD	SEQ ID NO:232
BM13730	EAKQASHEIRWLPNLTFFDQRVAFIVKLRD	SEQ ID NO:233
BM13731	EAAQAAHEIRWLPNLTFFDQRVAFIHKLRD	SEQ ID NO:234
BM13733	EVQYASHEIRWLPNLTFFDQRVAFIHKLRD	SEQ ID NO:235
BM13734	ELRNAAHEIRWLPNLTFFDQRVAFIHKLRD	SEQ ID NO:236
BM13737	EQRAAAHEIRWLPNLTFFDQRVAFIHKLRD	SEQ ID NO:237
BM13739	EASEAAAEIRWLPNLTFFDQRVAFIHKLRD	SEQ ID NO:238
BM13741	ESVIAAHEIRWLPNLTFFDQRVAFIHKLRD	SEQ ID NO:239
BM13744	EILRAKHEIRWLPNLTFFDQRVAFIHKLDD	SEQ ID NO:240
BM13746	ESKTAASHEIRWLPNLTFFDQRVAFIQKLRD	SEQ ID NO:241
BM13748	ELAEASHEIRWLPNLTFFDQRVAFIHKLFD	SEQ ID NO:242
BM13753	EATTAKHEIRWLPNLTFFDQRVAFIHKLDD	SEQ ID NO:243
BM13754	EIENAAHEIRWLPNLTFFDQRVAFIHKLRD	SEQ ID NO:244
BM13755	EAKDAAHEIRWLPNLTFFDQRVAFIRKLRD	SEQ ID NO:245
BM13757	ERLEAAAEIRWLPNLTFFDQRVAFIHKLRD	SEQ ID NO:246
BM13762	EQMEAAAEIRWLPNLTFFDQRVAFIHKLRD	SEQ ID NO:247
BM13763	EVKTASHEIRWLPNLTFFDQRVAFIRKLRD	SEQ ID NO:248
BM13764	ESFEASHEIRWLPNLTFFDQRVAFIQKLRD	SEQ ID NO:249
BM13765	EIKSAKHEIRWLPNLTFFDQRVAFIHKLDD	SEQ ID NO:250
BM13766	EIKNAKHEIRWLPNLTFFDQRVAFIHKLDD	SEQ ID NO:251
BM13768	ELQEAKHEIRWLPNLTFFDQRVAFIHKLDD	SEQ ID NO:252

Figure 1J

Designation	Sequence	SEQ ID NO:
BM13769	ERQNAAEIRWLPNLTFFDQRVAFIRKLRD	SEQ ID NO:253
BM13770	EVLQAKHEIRWLPNLTFFDQRVAFIRKLRD	SEQ ID NO:254
BM13772	EANVASHEIRWLPNLTFFDQRVAFIRKLRD	SEQ ID NO:255
BM13774	ELDAAKHEIRWLPNLTFFDQRVAFIRKLRD	SEQ ID NO:256
BM13775	ETASAAHEIRWLPNLTFFDQRVAFIRKLRD	SEQ ID NO:257
BM13778	EWKQAASEIRWLPNLTFFDQRVAFIRKLRD	SEQ ID NO:258
BM13779	ETASASHEIRWLPNLTFFDQRVAFIRKLRD	SEQ ID NO:259
BM13781	ESIVAGHEIRWLPNLTFFDQRVAFIRKLRD	SEQ ID NO:260
BM13784	EIKQAKHEIRWLPNLTFFDQRVAFIRKLRD	SEQ ID NO:261
BM13785	EQATASHEIRWLPNLTFFDQRVAFIRKLRD	SEQ ID NO:262
BM13787	ELNAAKHEIRWLPNLTFFDQRVAFIRKLRD	SEQ ID NO:263
BM13788	EVKRAGHEIRWLPNLTFFDQRVAFIRKLRD	SEQ ID NO:264
BM13789	ESRNAAEIRWLPNLTFFDQRVAFIRKLRD	SEQ ID NO:265
BM13791	ESITASAEIRWLPNLTFFDQRVAFIRKLRD	SEQ ID NO:266
BM13793	EAATAAEIRWLPNLTFFDQRVAFIRKLRD	SEQ ID NO:267
BM13794	EYAAKHEIRWLPNLTFFDQRVAFIRKLRD	SEQ ID NO:268
BM13795	EISRAKHEIRWLPNLTFFDQRVAFIRKLRD	SEQ ID NO:269
BM13798	EYVTAHEIRWLPNLTFFDQRVAFIRKLRD	SEQ ID NO:270
BM13800	EHIDAGHEIRWLPNLTFFDQRVAFIRKLRD	SEQ ID NO:271
BM13801	EILQAKHEIRWLPNLTFFDQRVAFIRKLRD	SEQ ID NO:272
BM13802	ENSQAKHEIRWLPNLTFFDQRVAFIRKLRD	SEQ ID NO:273
BM13803	EYRVAKEIRWLPNLTFFDQRVAFIRKLRD	SEQ ID NO:274
BM13804	EIYNAGHEIRWLPNLTFFDQRVAFIRKLRD	SEQ ID NO:275
BM13805	ESNEAAAEIRWLPNLTFFDQRVAFIRKLRD	SEQ ID NO:276
BM13809	ESQLAAAEIRWLPNLTFFDQRVAFIRKLRD	SEQ ID NO:277
BM13810	ELKEAGHEIRWLPNLTFFDQRVAFIRKLRD	SEQ ID NO:278
BM13813	ETRVASVEIRWLPNLTFFDQRVAFIRKLRD	SEQ ID NO:279
BM13818	ELRTAAHEIRWLPNLTFFDQRVAFIRKLRD	SEQ ID NO:280

Figure 1K

Designation	Sequence	SEQ ID NO:
BM13820	EKTYAHFEIRWLPNLTFFDQRFVAFISKLWD	SEQ ID NO:281
BM13821	EEAQASHEIRWLPNLTFFDQRFVAFIHKLRD	SEQ ID NO:282
BM13822	EITSAASHEIRWLPNLTFFDQRFVAFIHKLRD	SEQ ID NO:283
BM13825	EVKTASHEIRWLPNLTFFDQRFVAFIHKLRD	SEQ ID NO:284
BM13826	ETKVAASHEIRWLPNLTFFDQRFVAFIHKLRD	SEQ ID NO:285
BM13829	EDLVAQHEIRWLPNLTFFDQRFVAFIRKLRD	SEQ ID NO:286
BM13830	ETQTAFNEIRWLPNLTFFDQRFVAFIHKLRD	SEQ ID NO:287
BM13831	EIKDAGHEIRWLPNLTFFDQRFVAFIRKLRD	SEQ ID NO:288
BM13836	EYKEAGHEIRWLPNLTFFDQRFVAFIVKLRD	SEQ ID NO:289
BM13839	EAALAAHEIRWLPNLTFFDQRFVAFIHKLRD	SEQ ID NO:290
BM13843	EQERAAHEIRWLPNLTFFDQRFVAFIRKLRD	SEQ ID NO:291
BM13844	EWFDAAHEIRWLPNLTFFDQRFVAFIRKLRD	SEQ ID NO:292
BM13847	EIIQAKHEIRWLPNLTFFDQRFVAFIHKLRD	SEQ ID NO:293
BM13850	ELTNAKHEIRWLPNLTFFDQRFVAFIHKLRD	SEQ ID NO:294
BM13851	EIQLAKHEIRWLPNLTFFDQRFVAFIHKLRD	SEQ ID NO:295
BM13854	EIHDAKHEIRWLPNLTFFDQRFVAFIHKLRD	SEQ ID NO:296
BM13869	EVKIASHEIRWLPNLTFFDQRFVAFIHKLRD	SEQ ID NO:297
BM13870	EQHSAASHEIRWLPNLTFFDQRFVAFIRKLRD	SEQ ID NO:298
BM13872	EVFAASAEIRWLPNLTFFDQRFVAFIHKLRD	SEQ ID NO:299
BM13874	ETDLAGHEIRWLPNLTFFDQRFVAFIRKLRD	SEQ ID NO:300
BM13877	EANFAGHEIRWLPNLTFFDQRFVAFIRKLRD	SEQ ID NO:301
BM13878	EFETAGHEIRWLPNLTFFDQRFVAFITKLRD	SEQ ID NO:302
BM13880	EVNLAKHEIRWLPNLTFFDQRFVAFIHKLRD	SEQ ID NO:303
BM13881	EADTAAHEIRWLPNLTFFDQRFVAFIYKLRD	SEQ ID NO:304
BM13882	EFVDAGHEIRWLPNLTFFDQRFVAFIQKLRD	SEQ ID NO:305
BM13883	EDHKAHEIRWLPNLTFFDQRFVAFIHKLRD	SEQ ID NO:306
BM13884	ETVDAGHEIRWLPNLTFFDQRFVAFIHKLRD	SEQ ID NO:307
BM13885	ESQRAGHEIRWLPNLTFFDQRFVAFITKLRD	SEQ ID NO:308

Figure 1L

Designation	Sequence	SEQ ID NO:
BM13886	EWSSAAHEIRWLPNLTTFDQRVAFIHKLQD	SEQ ID NO:309
BM13887	EVAVAGHEIRWLPNLTTFDQRVAFIVKLRD	SEQ ID NO:310
BM13888	ESAEAGHEIRWLPNLTTFDQRVAFIEKIRD	SEQ ID NO:311
BM13889	EAVAAGHEIRWLPNLTTFDQRVAFIHKLND	SEQ ID NO:312
BM13890	EFQIAGHEIRWLPNLTTFDQRVAFINKLRD	SEQ ID NO:313
BM13891	ELMVAKHEIRWLPNLTTFDQRVAFIHKLDD	SEQ ID NO:314
BM13892	EYDAAHEIRWLPNLTTFDQRVAFILKLRD	SEQ ID NO:315
BM13893	EVLEAKHEIRWLPNLTTFDQRVAFIHKLDD	SEQ ID NO:316
BM13894	ESIAASHEIRWLPNLTTFDQRVAFIHKLRD	SEQ ID NO:317
BM13895	EVAEAGHEIRWLPNLTTFDQRVAFISKLRD	SEQ ID NO:318
BM13896	EQAKAAHEIRWLPNLTTFDQRVAFIHKLND	SEQ ID NO:319
BM13897	ERDDAAHEIRWLPNLTTFDQRVAFIEKLRD	SEQ ID NO:320
BM13898	EAKDASHEIRWLPNLTTFDQRVAFIHKLRD	SEQ ID NO:321
BM13899	EASSAAHEIRWLPNLTTFDQRVAFIQKLRD	SEQ ID NO:322
BM13900	EWMEASHEIRWLPNLTTFDQRVAFIVKLRD	SEQ ID NO:323
BM13901	EQKNAHEIRWLPNLTTFDQRVAFIEKLRD	SEQ ID NO:324
BM13902	EIENAKHEIRWLPNLTTFDQRVAFIHKLDD	SEQ ID NO:325
BM13903	EVNRASHEIRWLPNLTTFDQRVAFIHKLDD	SEQ ID NO:326
BM13904	ERLLAGHEIRWLPNLTTFDQRVAFINKLRD	SEQ ID NO:327
BM13906	EVSIAGHEIRWLPNLTTFDQRVAFIRKLRD	SEQ ID NO:328
BM13907	EKEVAAHEIRWLPNLTTFDQRVAFIHKLDD	SEQ ID NO:329
BM13908	ESERASHEIRWLPNLTTFDQRVAFIHKLRD	SEQ ID NO:330
BM13909	EWNEAAHEIRWLPNLTTFDQRVAFIHKLRD	SEQ ID NO:331
BM13910	ENVDAHEIRWLPNLTTFDQRVAFIHKLRD	SEQ ID NO:332
BM13911	EADAASHEIRWLPNLTTFDQRVAFIRKLRD	SEQ ID NO:333
BM13912	ELESASHEIRWLPNLTTFDQRVAFIHKLDD	SEQ ID NO:334
BM13913	EEQLAAHEIRWLPNLTTFDQRVAFIRKLRD	SEQ ID NO:335
BM13914	EEFELAGHEIRWLPNLTTFDQRVAFIHKLDD	SEQ ID NO:336

Figure 1M

Designation	Sequence	SEQ ID NO:
BM13915	EAFVAQHEIRWLPNLTFDQRVAFIVKLRD	SEQ ID NO:337
BM13916	EALKASHEIRWLPNLTFDQRVAFINKLRD	SEQ ID NO:338
BM13917	ELERAGHEIRWLPNLTFDQRVAFIKKLTID	SEQ ID NO:339
BM13918	EVEWAKHEIRWLPNLTFDQRVAFIHKIQD	SEQ ID NO:340
BM13919	EKASAQHEIRWLPNLTFDQRVAFIHKLIDD	SEQ ID NO:341
BM13920	ETEIAKHEIRWLPNLTFDQRVAFIHKLIDD	SEQ ID NO:342
BM13921	EVNLAQHEIRWLPNLTFDQRVAFIHKLTID	SEQ ID NO:343
BM13922	EAEAAHEIRWLPNLTFDQRVAFIHKLRD	SEQ ID NO:344
BM13923	ETDRAKHEIRWLPNLTFDQRVAFIHKLND	SEQ ID NO:345
BM13925	EFAQAGHEIRWLPNLTFDQRVAFIHKLKD	SEQ ID NO:346
BM13926	ETDEASHEIRWLPNLTFDQRVAFIRKLRD	SEQ ID NO:347
BM13927	ENADAGHEIRWLPNLTFDQRVAFIQKLRD	SEQ ID NO:348
BM13928	ESTQAAHEIRWLPNLTFDQRVAFIHKIQD	SEQ ID NO:349
BM13929	EQALAAHEIRWLPNLTFDQRVAFIHKLND	SEQ ID NO:350
BM13930	EAAHAQHEIRWLPNLTFDQRVAFIHKLIDD	SEQ ID NO:351
BM13932	EVDNAGHEIRWLPNLTFDQRVAFIQKLRD	SEQ ID NO:352
BM13993	EAGRAAHEIRWLPNLTWDQRVAFIWKLRD	SEQ ID NO:353
PP07918	KEQDAAAHEIRWLPNLTFDQRVAFIHKLADDPQSSELLSEAKKLNDSDQ	SEQ ID NO:354
PP10193	KEWMRAAHEIRWLPNLTFDQRVAFIHKLEDDPSQSSELLSEAKKLNDSDQ	SEQ ID NO:355
PP10109	KEANTAHEIRWLPNLTFDQRVAFIRKLWDDPSQSSELLSEAKKLNDSDQ	SEQ ID NO:356
PP07960	KEFESAAHEIRWLPNLTFDQRVAFIHKLSDDPSQSSELLSEAKKLNDSDQ	SEQ ID NO:357
PP10140	KERSAAAHEIRWLPNLTFDQRVAFILKLTDDPSQSSELLSEAKKLNDSDQ	SEQ ID NO:358
PP07930	KESDSAVHEIRWLPNLTFDQRVAFIHKLLDDPSQSSELLSEAKKLNDSDQ	SEQ ID NO:359
PP10183	KEADNAGHEIRWLPNLTWAQRVAFIHKLLDDPSQSSELLSEAKKLNDSDQ	SEQ ID NO:360
PP10111	KEDDTAAHEIRWLPNLTYEQWAFIHKLYDDPSQSSELLSEAKKLNDSDQ	SEQ ID NO:361
PP10129	KEQHDAHEIRWLPNLTFDQRVAFIRKLHDDPSQSSELLSEAKKLNDSDQ	SEQ ID NO:362
PP10141	KENQGAHEIRWLPNLTWDQRVAFIRKLQDDPSQSSELLSEAKKLNDSDQ	SEQ ID NO:363
PP10156	KERTQASHEIRWLPNLTFDQRVAFINKLRDDPSQSSELLSEAKKLNDSDQ	SEQ ID NO:364

Figure 1N

Designation	Sequence	SEQ ID NO :
PP10127	KERKDAAGHEIRWLPNLTFFDQQRVAFIKKLEDDPSQSSELLSEAKKLNDSSQ	SEQ ID NO:365
PP07909	KERQEAAGHEIRWLPNLTFFDQQRVAFIVKLLDDPSQSSELLSEAKKLNDSSQ	SEQ ID NO:366
PP10152	KEEDVAAAGHEIRWLPNLTFFNQRAAFIDKLLDDPSQSSELLSEAKKLNDSSQ	SEQ ID NO:367
PP10145	KENQDAAAGHEIRWLPNLTFFDQQRVAFIGKLNDDPSQSSELLSEAKKLNDSSQ	SEQ ID NO:368
PP10161	KESGYAVAGHEIRWLPNLTFFDQQRVAFINKIQDDPSQSSELLSEAKKLNDSSQ	SEQ ID NO:369
PP13573	KESKDAAGHEIRWLPNLTFFDQQRVAFIHKLRDDPSQSSELLSEAKKLNDSSQ	SEQ ID NO:370
PP13574	KEKKEAAGHEIRWLPNLTFFDQQRVAFIHKLRDDPSQSSELLSEAKKLNDSSQ	SEQ ID NO:371
PP13577	KEWHQAAAGHEIRWLPNLTFFDQQRVAFIHKLRDDPSQSSELLSEAKKLNDSSQ	SEQ ID NO:372
PP13578	KEWTDAAAGHEIRWLPNLTFFDQQRVAFIHKLRDDPSQSSELLSEAKKLNDSSQ	SEQ ID NO:373
PP13579	KEISAASAGHEIRWLPNLTFFDQQRVAFIHKLRDDPSQSSELLSEAKKLNDSSQ	SEQ ID NO:374
PP13581	KEQQAAGHEIRWLPNLTFFDQQRVAFIHKLRDDPSQSSELLSEAKKLNDSSQ	SEQ ID NO:375
PP13583	KELEKAGHEIRWLPNLTFFDQQRVAFIHKLRDDPSQSSELLSEAKKLNDSSQ	SEQ ID NO:376
PP13585	KEYLDAAGHEIRWLPNLTFFDQQRVAFIHKLRDDPSQSSELLSEAKKLNDSSQ	SEQ ID NO:377
PP13586	KELKDAAGHEIRWLPNLTFFDQQRVAFIHKLRDDPSQSSELLSEAKKLNDSSQ	SEQ ID NO:378
PP13587	KEHVDAAGHEIRWLPNLTFFDQQRVAFIHKLRDDPSQSSELLSEAKKLNDSSQ	SEQ ID NO:379
PP13588	KEYAAAAAGHEIRWLPNLTFFDQQRVAFIHKLRDDPSQSSELLSEAKKLNDSSQ	SEQ ID NO:380
PP13592	KEVDIAAGHEIRWLPNLTFFDQQRVAFIHKLRDDPSQSSELLSEAKKLNDSSQ	SEQ ID NO:381
PP13594	KEIDEAKHEIRWLPNLTFFDQQRVAFIHKLRDDPSQSSELLSEAKKLNDSSQ	SEQ ID NO:382
PP13596	KELRQAGHEIRWLPNLTFFDQQRVAFIHKLRDDPSQSSELLSEAKKLNDSSQ	SEQ ID NO:383
PP13597	KELOSAAGHEIRWLPNLTFFDQQRVAFIHKLRDDPSQSSELLSEAKKLNDSSQ	SEQ ID NO:384
PP13598	KELEKAAAGHEIRWLPNLTFFDQQRVAFIHKLRDDPSQSSELLSEAKKLNDSSQ	SEQ ID NO:385
PP13600	KEAHEAAGHEIRWLPNLTFFDQQRVAFIHKLRDDPSQSSELLSEAKKLNDSSQ	SEQ ID NO:386
PP13604	KELOAASAGHEIRWLPNLTFFDQQRVAFIHKLRDDPSQSSELLSEAKKLNDSSQ	SEQ ID NO:387
PP13605	KEITESAKHEIRWLPNLTFFDQQRVAFIHKLRDDPSQSSELLSEAKKLNDSSQ	SEQ ID NO:388
PP13609	KEWKVAAAGHEIRWLPNLTFFDQQRVAFIHKLRDDPSQSSELLSEAKKLNDSSQ	SEQ ID NO:389
PP13611	KEWKAAAGHEIRWLPNLTFFDQQRVAFIHKLRDDPSQSSELLSEAKKLNDSSQ	SEQ ID NO:390
PP13612	KEIDLAKHEIRWLPNLTFFDQQRVAFIHKLRDDPSQSSELLSEAKKLNDSSQ	SEQ ID NO:391
PP13613	KELEAARAGHEIRWLPNLTFFDQQRVAFIHKLRDDPSQSSELLSEAKKLNDSSQ	SEQ ID NO:392

Figure 10

Designation	Sequence	SEQ ID NO:
PP13615	KEAATAGHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:393
PP13616	KEWQAAHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:394
PP13617	KEADQACHEIRWLPNLTFFDQVAFIRKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:395
PP13620	KEQSKAAHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:396
PP13621	KEADAACHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:397
PP13622	KEFMDACHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:398
PP13624	KESQAAHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:399
PP13625	KEVSDAAHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:400
PP13626	KEADSASHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:401
PP13627	KELMEAAHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:402
PP13628	KELNTAAHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:403
PP13629	KEVHEACHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:404
PP13633	KESTAAHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:405
PP13634	KEWYNAHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:406
PP13635	KEWNDAAHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:407
PP13637	KEVEVAKHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:408
PP13638	KEFNFAAHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:409
PP13639	KEHDSAAHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:410
PP13640	KEWMDAAHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:411
PP13641	KEFSAACHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:412
PP13644	KEINSAACHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:413
PP13645	KEVDTAAHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:414
PP13648	KESQIACHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:415
PP13651	KEVSAASHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:416
PP13652	KEDQDAAHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:417
PP13654	KELEAARHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:418
PP13655	KESKRAAHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:419
PP13656	KEYVKACHEIRWLPNLTFFDQVAFIRKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:420

Figure 1P

Designation	Sequence	SEQ ID NO:
PP13657	KEFSRACHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:421
PP13659	KEWQFAAHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:422
PP13663	KEWQIAAHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:423
PP13664	KELQEAHAHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:424
PP13667	KEYRAAAHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:425
PP13669	KELASACHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:426
PP13672	KEVQSACHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:427
PP13674	KEIEDACHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:428
PP13675	KENQAAAHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:429
PP13676	KEATSACHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:430
PP13678	KEDEQAAHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:431
PP13684	KEQNQAGHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:432
PP13688	KEYTSAACHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:433
PP13691	KEWDAAAHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:434
PP13692	KEEMQACHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:435
PP13694	KELSDAAHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:436
PP13695	KEIDAAKHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:437
PP13697	KEAERACHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:438
PP13706	KEEDSAAHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:439
PP13708	KEQKAACHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:440
PP13710	KEWDQAAHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:441
PP13711	KEAKAAAHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:442
PP13714	KELSEAAHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:443
PP13716	KETEAAKHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:444
PP13719	KEAKSAAHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:445
PP13720	KEQSAHAHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:446
PP13721	KEKERAHAHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:447
PP13725	KEWDEAAHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:448

Figure 1Q

Designation	Sequence	SEQ ID NO:
PP13727	KEEKDAAEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO: 449
PP13728	KEIENAAAEIRWLPNLTFFDQVAFIWKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO: 450
PP13732	KETKEAAAEIRWLPNLTFFDQVAFIHKLWDDPSQSSELLSEAKKLNDSQ	SEQ ID NO: 451
PP13735	KELEAAAEIRWLPNLTFFDQVAFIHKLNDDPSQSSELLSEAKKLNDSQ	SEQ ID NO: 452
PP13736	KEWAAAEAEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO: 453
PP13740	KESQEAGAEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO: 454
PP13742	KELSTAAAEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO: 455
PP13747	KEIEEAGAEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO: 456
PP13749	KELQTASAEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO: 457
PP13750	KEQDSAAAEIRWLPNLTFFDQVAFIRKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO: 458
PP13751	KESASAGAEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO: 459
PP13752	KEVAKASAEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO: 460
PP13758	KEVQEAHAEIRWLPNLTFFDQVAFIHKLWDDPSQSSELLSEAKKLNDSQ	SEQ ID NO: 461
PP13759	KESYEAHAEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO: 462
PP13760	KETAEGAEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO: 463
PP13761	KELEAAAEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO: 464
PP13771	KEAAAAGAEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO: 465
PP13773	KEYVDAGAEIRWLPNLTFFDQVAFIRKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO: 466
PP13776	KEIQEAGAEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO: 467
PP13777	KESATAHAEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO: 468
PP13780	KEWMSAAHAEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO: 469
PP13782	KEREQAGAEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO: 470
PP13783	KEIEQAGAEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO: 471
PP13786	KEHNAAGAEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO: 472
PP13792	KEIEVAAAEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO: 473
PP13796	KERAFAAEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO: 474
PP13799	KESELAHAEIRWLPNLTFFDQVAFIWKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO: 475
PP13806	KEYRAAGAEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO: 476

Figure 1R

Designation	Sequence	SEQ ID NO:
PP13808	KETANA AHEIRWLPNLTFFDQRVAFIHKLRDDPSQSSELLSEAKKLND SQ	SEQ ID NO:477
PP13811	KEWYEA AHEIRWLPNLTFFDQRVAFIHKLRDDPSQSSELLSEAKKLND SQ	SEQ ID NO:478
PP13812	KEEQEA AHEIRWLPNLTFFDQRVAFIHKLRDDPSQSSELLSEAKKLND SQ	SEQ ID NO:479
PP13823	KEHDAA AHEIRWLPNLTFFDQRVAFIHKLRDDPSQSSELLSEAKKLND SQ	SEQ ID NO:480
PP13824	KEWYSA AHEIRWLPNLTFFDQRVAFIHKLRDDPSQSSELLSEAKKLND SQ	SEQ ID NO:481
PP13838	KEISDA AHEIRWLPNLTFFDQRVAFIHKLRDDPSQSSELLSEAKKLND SQ	SEQ ID NO:482
PP13840	KEYTTA AHEIRWLPNLTFFDQRVAFIHKLRDDPSQSSELLSEAKKLND SQ	SEQ ID NO:483
PP13842	KEISQA AHEIRWLPNLTFFDQRVAFIHKLRDDPSQSSELLSEAKKLND SQ	SEQ ID NO:484
PP13845	KENDDA AHEIRWLPNLTFFDQRVAFIHKLRDDPSQSSELLSEAKKLND SQ	SEQ ID NO:485
PP13846	KESIEA GHEIRWLPNLTFFDQRVAFIHKLRDDPSQSSELLSEAKKLND SQ	SEQ ID NO:486
PP13848	KEQADA AHEIRWLPNLTFFDQRVAFIHKLRDDPSQSSELLSEAKKLND SQ	SEQ ID NO:487
PP13849	KETESA AHEIRWLPNLTFFDQRVAFIHKLRDDPSQSSELLSEAKKLND SQ	SEQ ID NO:488
PP13860	KEISDA KHEIRWLPNLTFFDQRVAFIHKLRDDPSQSSELLSEAKKLND SQ	SEQ ID NO:489
PP13865	KEHLNA AHEIRWLPNLTFFDQRVAFIHKLRDDPSQSSELLSEAKKLND SQ	SEQ ID NO:490
PP13866	KEWLDA AHEIRWLPNLTFFDQRVAFIHKLRDDPSQSSELLSEAKKLND SQ	SEQ ID NO:491
PP13875	KENAAA AHEIRWLPNLTFFDQRVAFIHKLRDDPSQSSELLSEAKKLND SQ	SEQ ID NO:492
PP13879	KEAELA GHEIRWLPNLTFFDQRVAFIHKLRDDPSQSSELLSEAKKLND SQ	SEQ ID NO:493
PP13480	KEVTDAGHEIRWLPNLTFFDQRVAFIEKLRDDPSQSSELLSEAKKLND SQ	SEQ ID NO:494
PP13481	KELDSA SHEIRWLPNLTFFDQRVAFINKLRDDPSQSSELLSEAKKLND SQ	SEQ ID NO:495
PP13482	KEINLA KHEIRWLPNLTFFDQRVAFIEKLRDDPSQSSELLSEAKKLND SQ	SEQ ID NO:496
PP13483	KESEVA KHEIRWLPNLTFFDQRVAFIHKLRDDPSQSSELLSEAKKLND SQ	SEQ ID NO:497
PP13484	KESAEAGHEIRWLPNLTFFDQRVAFIHKLRDDPSQSSELLSEAKKLND SQ	SEQ ID NO:498
PP13485	KEYSNA AHEIRWLPNLTFFDQRVAFIDKLRDDPSQSSELLSEAKKLND SQ	SEQ ID NO:499
PP13487	KETNNA GHEIRWLPNLTFFDQRVAFIHKLRDDPSQSSELLSEAKKLND SQ	SEQ ID NO:500
PP13488	KEVEFA AHEIRWLPNLTFFDQRVAFIHKLRDDPSQSSELLSEAKKLND SQ	SEQ ID NO:501
PP13489	KEVELAGHEIRWLPNLTFFDQRVAFIEKLRDDPSQSSELLSEAKKLND SQ	SEQ ID NO:502
PP13490	KEVLKA SHEIRWLPNLTFFDQRVAFITKLQDDPSQSSELLSEAKKLND SQ	SEQ ID NO:503
PP13491	KELANAGHEIRWLPNLTFFDQRVAFIRKLRDDPSQSSELLSEAKKLND SQ	SEQ ID NO:504

Figure 1S

Designation	Sequence	SEQ ID NO:
PP13493	KEYMKACHEIRWLPNLTFFDQRVAFIVKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:505
PP13495	KEHANAQHEIRWLPNLTFFDQRVAFIHKLQDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:506
PP13496	KEVDIASHEIRWLPNLTFFDQRVAFIWKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:507
PP13497	KEEVFAAHEIRWLPNLTFFDQRVAFIVKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:508
PP13499	KEFNTAAHEIRWLPNLTFFDQRVAFINKLDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:509
PP13501	KEVDVACHEIRWLPNLTFFDQRVAFIRKLNDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:510
PP13502	KEWSLAAHEIRWLPNLTFFDQRVAFIVKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:511
PP13503	KELDDAAHEIRWLPNLTFFDQRVAFIQKLQDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:512
PP13505	KERHEAGHEIRWLPNLTFFDQRVAFIRKLNDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:513
PP13506	KEISDAIHEIRWLPNLTFFDQRVAFIHKLQDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:514
PP13507	KEWETACHEIRWLPNLTFFDQRVAFIVKLSDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:515
PP13508	KERYWASHEIRWLPNLTFFDQRVAFIHKLNDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:516
PP13512	KEIDWACHEIRWLPNLTFFDQRVAFIHKLLDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:517
PP13515	KEQSKACHEIRWLPNLTFFDQRVAFIHKLKDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:518
PP13518	KEIEAAQHEIRWLPNLTFFDQRVAFINKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:519
PP13519	KEHEQAAHEIRWLPNLTFFDQRVAFIRKLVDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:520
PP13520	KEAEQACHEIRWLPNLTFFDQRVAFINKLQDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:521
PP13522	KEVDYAAHEIRWLPNLTFFDQRVAFIHKLWDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:522
PP13525	KEYSAAACHEIRWLPNLTFFDQRVAFIEKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:523
PP13561	KELATASHEIRWLPNLTFFDQRVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:524
PP13572	KEYRVACHEIRWLPNLTFFDQRVAFIQKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:525
PP13575	KEVVSAKHEIRWLPNLTFFDQRVAFIHKLDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:526
PP13576	KESAQAAHEIRWLPNLTFFDQRVAFIRKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:527
PP13584	KEYSAAACHEIRWLPNLTFFDQRVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:528
PP13589	KEQKEAAAHEIRWLPNLTFFDQRVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:529
PP13590	KEAAIACKEIRWLPNLTFFDQRVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:530
PP13591	KEISKAKHEIRWLPNLTFFDQRVAFIHKLDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:531
PP13593	KESVAAACHEIRWLPNLTFFDQRVAFIWKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:532

Figure 1T

Designation	Sequence	SEQ ID NO:
PP13595	KEIQAKHEIRWLPNLTFFQQRVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:533
PP13599	KEITSAKHEIRWLPNLTFFQQRVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:534
PP13601	KEQDVAACHEIRWLPNLTFFQQRVAFIRKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:535
PP13603	KELERAAHEIRWLPNLTFFQQRVAFINKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:536
PP13607	KENQLAAHEIRWLPNLTFFQQRVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:537
PP13608	KEISQAKHEIRWLPNLTFFQQRVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:538
PP13610	KEIANACHEIRWLPNLTFFQQRVAFIYKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:539
PP13614	KEWQAAACHEIRWLPNLTFFQQRVAFIRKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:540
PP13618	KERKDACHEIRWLPNLTFFQQRVAFIRKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:541
PP13619	KEITQAKHEIRWLPNLTFFQQRVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:542
PP13623	KEFIQACHEIRWLPNLTFFQQRVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:543
PP13630	KEWNTASHEIRWLPNLTFFQQRVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:544
PP13631	KEKFVAAHEIRWLPNLTFFQQRVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:545
PP13632	KEADSAGAEIRWLPNLTFFQQRVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:546
PP13636	KESSVAAAEIRWLPNLTFFQQRVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:547
PP13642	KEVDLACREIRWLPNLTFFQQRVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:548
PP13643	KEQERAAAEIRWLPNLTFFQQRVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:549
PP13646	KEIWQAKHEIRWLPNLTFFQQRVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:550
PP13647	KELNQAKHEIRWLPNLTFFQQRVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:551
PP13649	KELQQASHEIRWLPNLTFFQQRVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:552
PP13650	KEINQAKHEIRWLPNLTFFQQRVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:553
PP13653	KELVLACHEIRWLPNLTFFQQRVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:554
PP13658	KELTSAAHEIRWLPNLTFFQQRVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:555
PP13661	KEWNAAREIRWLPNLTFFQQRVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:556
PP13662	KEILHAKHEIRWLPNLTFFQQRVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:557
PP13665	KEVLTAKHEIRWLPNLTFFQQRVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:558
PP13666	KENSKAAHEIRWLPNLTFFQQRVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:559
PP13668	KEVMTAKHEIRWLPNLTFFQQRVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:560

Figure 1U

Designation	Sequence	SEQ ID NO:
PP13671	KEARDAAEIRWLPNLTFFDQVAFIHKLSDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:561
PP13673	KERSKAAHEIRWLPNLTFFDQVAFIHKLKDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:562
PP13677	KEIYSARHEIRWLPNLTFFDQVAFIHKLDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:563
PP13679	KEVQSAKHEIRWLPNLTFFDQVAFIHKLDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:564
PP13680	KETLEAAHEIRWLPNLTFFDQVAFIHKLKDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:565
PP13681	KEQMRAAHEIRWLPNLTFFDQVAFIRKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:566
PP13685	KENKNAAEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:567
PP13687	KETESAKHEIRWLPNLTFFDQVAFIHKLTDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:568
PP13689	KETVQAKHEIRWLPNLTFFDQVAFIHKLDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:569
PP13693	KEIASAKHEIRWLPNLTFFDQVAFIHKLDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:570
PP13698	KEVMDAKHEIRWLPNLTFFDQVAFIHKLDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:571
PP13699	KETDAAKHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:572
PP13701	KELQIAKHEIRWLPNLTFFDQVAFIHKLDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:573
PP13702	KEWKDAAQEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:574
PP13703	KERDSAHEIRWLPNLTFFDQVAFIRKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:575
PP13704	KEIAAAKHEIRWLPNLTFFDQVAFIHKLDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:576
PP13707	KESVKAKHEIRWLPNLTFFDQVAFIHKLDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:577
PP13709	KENERAKHEIRWLPNLTFFDQVAFIHKLDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:578
PP13712	KEYKRACHEIRWLPNLTFFDQVAFIRKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:579
PP13713	KEVRAASHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:580
PP13717	KEDKRAAHEIRWLPNLTFFDQVAFIRKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:581
PP13718	KESEKAKKEIRWLPNLTFFDQVAFIRKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:582
PP13722	KEINRAKHEIRWLPNLTFFDQVAFIHKLDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:583
PP13724	KETQQASHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:584
PP13729	KENQSAAEIRWLPNLTFFDQVAFIRKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:585
PP13730	KEAKQASHEIRWLPNLTFFDQVAFIVKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:586
PP13731	KEAAQAAHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:587
PP13733	KEVQYASHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:588

Figure 1V

Designation	Sequence	SEQ ID NO:
PP13734	KELRNAAHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:589
PP13737	KEQRAAAHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:590
PP13739	KEASEAAAAEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:591
PP13741	KESVIAAHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:592
PP13744	KEILRAKHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:593
PP13746	KESKTAHEIRWLPNLTFFDQVAFIQKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:594
PP13748	KELAEASHEIRWLPNLTFFDQVAFIHKLFDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:595
PP13753	KEATTAKHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:596
PP13754	KEIENAAHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:597
PP13755	KEAKDAAHEIRWLPNLTFFDQVAFIRKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:598
PP13757	KERLEAAAAEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:599
PP13762	KEQMEAAAAEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:600
PP13763	KEVKTASHEIRWLPNLTFFDQVAFIRKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:601
PP13764	KESFEASHEIRWLPNLTFFDQVAFIQKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:602
PP13765	KEIKSAKHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:603
PP13766	KEIKNAKHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:604
PP13768	KELQEAKEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:605
PP13769	KERQNAAEIRWLPNLTFFDQVAFIRKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:606
PP13770	KEVLQAKHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:607
PP13772	KEANVASHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:608
PP13774	KELDAAKHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:609
PP13775	KETASAAHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:610
PP13778	KEWKQAASEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:611
PP13779	KETASASHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:612
PP13781	KESIVACHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:613
PP13784	KEIKQAKHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:614
PP13785	KEQATASHEIRWLPNLTFFDQVAFIRKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:615
PP13787	KELNAAKHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:616

Figure 1X

Designation	Sequence	SEQ ID NO:
PP13788	KEVKRACHEIRWLPNLTFFDQVAFIRKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:617
PP13789	KESRNAACHEIRWLPNLTFFDQVAFIHKLQDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:618
PP13791	KESITASAEIRWLPNLTFFDQVAFIRKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:619
PP13793	KEAATAACHEIRWLPNLTFFDQVAFIHKLQDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:620
PP13794	KEVYAAKHEIRWLPNLTFFDQVAFIHKLDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:621
PP13795	KEISRACHEIRWLPNLTFFDQVAFIHKLDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:622
PP13798	KEYVTAACHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:623
PP13800	KEHIDACHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:624
PP13801	KEILQAKHEIRWLPNLTFFDQVAFIHKLNDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:625
PP13802	KENSQAKHEIRWLPNLTFFDQVAFIHKLNDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:626
PP13803	KEYRVAACHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:627
PP13804	KEIYNAGHEIRWLPNLTFFDQVAFIRKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:628
PP13805	KESNEAAAEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:629
PP13809	KESQLAAAEIRWLPNLTFFDQVAFIRKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:630
PP13810	KELKEACHEIRWLPNLTFFDQVAFIHKLQDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:631
PP13813	KETRVAACHEIRWLPNLTFFDQVAFIQKLNDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:632
PP13818	KELRTAAACHEIRWLPNLTFFDQVAFIHKLNDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:633
PP13820	KEKTYAHFEIRWLPNLTFFDQVAFISKLWDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:634
PP13821	KEEAQASHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:635
PP13822	KEITSAACHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:636
PP13825	KEVKTASHEIRWLPNLTFFDQVAFIHKLKKDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:637
PP13826	KETKVAACHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:638
PP13829	KEDLVAACHEIRWLPNLTFFDQVAFIRKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:639
PP13830	KETQTAFNEIRWLPNLTYPQRAAFILKLWDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:640
PP13831	KEIKDACHEIRWLPNLTFFDQVAFIRKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:641
PP13836	KEYKEACHEIRWLPNLTYPQRAAFIVKLRRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:642
PP13839	KEAALAAACHEIRWLPNLTFFDQVAFIHKLKKDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:643
PP13843	KEQERAAACHEIRWLPNLTFFDQVAFIRKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:644

Figure 1Y

Designation	Sequence	SEQ ID NO:
PP13844	KEWFDAAEIRWLPNLTFFDQVAFIRKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:645
PP13847	KEIIQAKHEIRWLPNLTFFDQVAFIHKLDDDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:646
PP13850	KELTNAKHEIRWLPNLTFFDQVAFIHKLDDDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:647
PP13851	KEIQLAKHEIRWLPNLTFFDQVAFIHKLDDDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:648
PP13854	KEIHDAKHEIRWLPNLTFFDQVAFIHKLQDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:649
PP13869	KEVKIASHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:650
PP13870	KEQSAHAHEIRWLPNLTFFDQVAFIRKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:651
PP13872	KEVFAASAEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:652
PP13874	KETDLAGEIRWLPNLTFFDQVAFIRKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:653
PP13877	KEANFACHEIRWLPNLTFFDQVAFIRKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:654
PP13878	KEFETACHEIRWLPNLTFFDQVAFITKLWDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:655
PP13880	KEVNLAKIHEIRWLPNLTFFDQVAFIHKLDDDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:656
PP13881	KEADTAHEIRWLPNLTFFDQVAFIYKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:657
PP13882	KEFVDACHEIRWLPNLTFFDQVAFIQKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:658
PP13883	KEDHKAHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:659
PP13884	KETVDACHEIRWLPNLTFFDQVAFIHKLADDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:660
PP13885	KESQRACHEIRWLPNLTFFDQVAFITKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:661
PP13886	KEWSSAAHEIRWLPNLTFFDQVAFIHKLQDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:662
PP13887	KEVAVACHEIRWLPNLTFFDQVAFIVKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:663
PP13888	KESAEACHEIRWLPNLTFFDQVAFIEKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:664
PP13889	KEAVAACHEIRWLPNLTFFDQVAFIHKLNDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:665
PP13890	KEFQIACHEIRWLPNLTFFDQVAFINKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:666
PP13891	KELMVAKHEIRWLPNLTFFDQVAFIHKLDDDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:667
PP13892	KEYDSAHEIRWLPNLTFFDQVAFILKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:668
PP13893	KEVLEAKHEIRWLPNLTFFDQVAFIHKLDDDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:669
PP13894	KESIAASHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:670
PP13895	KEVAEACHEIRWLPNLTFFDQVAFISKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:671
PP13896	KEQAKAAHEIRWLPNLTFFDQVAFIHKLNDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:672

Figure 1Z

Designation	Sequence	SEQ ID NO:
PP13897	KERDDAAHEIRWLPNLTFFDQVAFIEKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:673
PP13898	KEAKDASHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:674
PP13899	KEASSAAHEIRWLPNLTFFDQVAFIQKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:675
PP13900	KEWMEASHEIRWLPNLTFFDQVAFIVKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:676
PP13901	KEQKNAASHEIRWLPNLTFFDQVAFIEKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:677
PP13902	KEIENAKHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:678
PP13903	KEVNRASHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:679
PP13904	KERLLACHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:680
PP13906	KEVSIAGHEIRWLPNLTFFDQVAFIRKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:681
PP13907	KEKEVAAHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:682
PP13908	KESERASHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:683
PP13909	KEWNEAAHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:684
PP13910	KENVDAASHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:685
PP13911	KEADAASHEIRWLPNLTFFDQVAFIRKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:686
PP13912	KELESASHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:687
PP13913	KEEQLAASHEIRWLPNLTFFDQVAFIRKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:688
PP13914	KEFELACHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:689
PP13915	KEAFVAQHEIRWLPNLTFFDQVAFIVKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:690
PP13916	KEALKASHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:691
PP13917	KELERACHEIRWLPNLTFFDQVAFIKKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:692
PP13918	KEVEWAKHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:693
PP13919	KEKASACHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:694
PP13920	KETETAKHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:695
PP13921	KEVNLAASHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:696
PP13922	KEAEAAAHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:697
PP13923	KETDRAKHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:698
PP13925	KEFAQACHEIRWLPNLTFFDQVAFIHKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:699
PP13926	KETDEASHEIRWLPNLTFFDQVAFIRKLRDDPSQSSELLSEAKKLNDSQ	SEQ ID NO:700

Figure 1AA

Designation	Sequence	SEQ ID NO:
PP13927	KENADAGHEIRWLPNLTFDQRVAFIQKLRDDPSQSSELLSEAKKLNDSSQ	SEQ ID NO:701
PP13928	KESTQAAHEIRWLPNLTFDQRVAFIHKLQDDPSQSSELLSEAKKLNDSSQ	SEQ ID NO:702
PP13929	KEQALAAHEIRWLPNLTFDQRVAFIHKLNDPSQSSELLSEAKKLNDSSQ	SEQ ID NO:703
PP13930	KEAHAASHEIRWLPNLTFDQRVAFIHKLDDPSQSSELLSEAKKLNDSSQ	SEQ ID NO:704
PP13932	KEVDNAGHEIRWLPNLTFDQRVAFIQKLRDDPSQSSELLSEAKKLNDSSQ	SEQ ID NO:705
PP13993	KEAGRAAHEIRWLPNLTWQRVAFIWKLRDDPSQSSELLSEAKKLNDSSQ	SEQ ID NO:706
Z07918	VDKAYAKEQDAAAEIRWLPNLTFDQRVAFIHKLADDPQSSELLSEAKKLNDSSQAPK	SEQ ID NO:707
Z10193	VDKAYAKEWMRAAHEIRWLPNLTFDQRVAFIHKLEDDPSQSSELLSEAKKLNDSSQAPK	SEQ ID NO:708
Z10109	VDKAYAKEANTAHEIRWLPNLTFDQRVAFIRKLWDDPSQSSELLSEAKKLNDSSQAPK	SEQ ID NO:709
Z07960	VDKAYAKEFESAHEIRWLPNLTQDQRVAFIHKLDDPSQSSELLSEAKKLNDSSQAPK	SEQ ID NO:710
Z10140	VDKAYAKERSAAAHEIRWLPNLTFDQRVAFITKLTDPSQSSELLSEAKKLNDSSQAPK	SEQ ID NO:711
Z07930	VDKAYAKESDAAVHEIRWLPNLTFDQRVAFIHKLDDPSQSSELLSEAKKLNDSSQAPK	SEQ ID NO:712
Z10183	VDKAYAKEADNAGHEIRWLPNLTWQRVAFIHKLDDPSQSSELLSEAKKLNDSSQAPK	SEQ ID NO:713
Z10111	VDKAYAKEDDTAAHEIRWLPNLTQEQRWAFIHKLYDDPSQSSELLSEAKKLNDSSQAPK	SEQ ID NO:714
Z10129	VDKAYAKEQHDAAHEIRWLPNLTQDQRVAFIRKLHDDPSQSSELLSEAKKLNDSSQAPK	SEQ ID NO:715
Z10141	VDKAYAKENQGAHEIRWLPNLTWQDQRVAFIRKLQDDPSQSSELLSEAKKLNDSSQAPK	SEQ ID NO:716
Z10156	VDKAYAKERTQASHEIRWLPNLTQDQRVAFIRKLDDPSQSSELLSEAKKLNDSSQAPK	SEQ ID NO:717
Z10127	VDKAYAKERKDAGHEIRWLPNLTQDQSAFIRKLDDPSQSSELLSEAKKLNDSSQAPK	SEQ ID NO:718
Z07909	VDKAYAKERQEAHEIRWLPNLTQDQRVAFIRKLDDPSQSSELLSEAKKLNDSSQAPK	SEQ ID NO:719
Z10152	VDKAYAKEEDVAAHEIRWLPNLTQDQRAAFIRKLDDPSQSSELLSEAKKLNDSSQAPK	SEQ ID NO:720
Z10145	VDKAYAKENQDAAHEIRWLPNLTQDQRVAFIRKLDDPSQSSELLSEAKKLNDSSQAPK	SEQ ID NO:721
Z10161	VDKAYAKESGVAVHEIRWLPNLTQDQRVAFIRKLDDPSQSSELLSEAKKLNDSSQAPK	SEQ ID NO:722
Z13573	VDKAYAKESKDAHEIRWLPNLTQDQRVAFIRKLDDPSQSSELLSEAKKLNDSSQAPK	SEQ ID NO:723
Z13574	VDKAYAKEKKAHEIRWLPNLTQDQRVAFIRKLDDPSQSSELLSEAKKLNDSSQAPK	SEQ ID NO:724
Z13577	VDKAYAKEWHQAAHEIRWLPNLTQDQRVAFIRKLDDPSQSSELLSEAKKLNDSSQAPK	SEQ ID NO:725
Z13578	VDKAYAKEWTDAHEIRWLPNLTQDQRVAFIRKLDDPSQSSELLSEAKKLNDSSQAPK	SEQ ID NO:726
Z13579	VDKAYAKEISAAHEIRWLPNLTQDQRVAFIRKLDDPSQSSELLSEAKKLNDSSQAPK	SEQ ID NO:727
Z13581	VDKAYAKEQQAHEIRWLPNLTQDQRVAFIRKLDDPSQSSELLSEAKKLNDSSQAPK	SEQ ID NO:728

Figure 1BB

Designation	Sequence	SEQ ID NO:
Z13583	VDAKYAKELEKAGHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:729
Z13585	VDAKYAKELYDAAHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:730
Z13586	VDAKYAKELKDAGHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:731
Z13587	VDAKYAKEHVDAHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:732
Z13588	VDAKYAKYAAAAHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:733
Z13592	VDAKYAKEVDIAAHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:734
Z13594	VDAKYAKEIDEAKHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:735
Z13596	VDAKYAKELRQAGHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:736
Z13597	VDAKYAKELQSAHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:737
Z13598	VDAKYAKELEKAAHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:738
Z13600	VDAKYAKEAHEAAHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:739
Z13604	VDAKYAKELQAA SHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:740
Z13605	VDAKYAKEIE SAKHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:741
Z13609	VDAKYAKEWKVAAHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:742
Z13611	VDAKYAKEWKAAAHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:743
Z13612	VDAKYAKEIDLAKHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:744
Z13613	VDAKYAKELEAARHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:745
Z13615	VDAKYAKEAAATAGHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:746
Z13616	VDAKYAKEWQAAAHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:747
Z13617	VDAKYAKEADQAGHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:748
Z13620	VDAKYAKEQSKAAHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:749
Z13621	VDAKYAKEADAAGHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:750
Z13622	VDAKYAKEFMDAGHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:751
Z13624	VDAKYAKESKQAAHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:752
Z13625	VDAKYAKEVSDAAHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:753
Z13626	VDAKYAKEFADSASHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:754
Z13627	VDAKYAKELMEAAHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:755
Z13628	VDAKYAKELNTAAHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:756

Figure 1CC

Designation	Sequence	SEQ ID NO:
Z13629	VDAKYAKEVHEAGHEIRWLPNLTFDQRVAFTHKLDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:757
Z13633	VDAKYAKESTAAAEHEIRWLPNLTFDQRVAFTHKLDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:758
Z13634	VDAKYAKFWYNAAEHEIRWLPNLTFDQRVAFTHKLDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:759
Z13635	VDAKYAKENWDAAEHEIRWLPNLTFDQRVAFTHKLDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:760
Z13637	VDAKYAKEVEVAKHEIRWLPNLTFDQRVAFTHKLDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:761
Z13638	VDAKYAKEFNFAAEHEIRWLPNLTFDQRVAFTHKLDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:762
Z13639	VDAKYAKEHDSA AEHEIRWLPNLTFDQRVAFTHKLDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:763
Z13640	VDAKYAKEMWDA AEHEIRWLPNLTFDQRVAFTHKLDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:764
Z13641	VDAKYAKEFSAAGHEIRWLPNLTFDQRVAFTHKLDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:765
Z13644	VDAKYAKELN SAAAEHEIRWLPNLTFDQRVAFTHKLDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:766
Z13645	VDAKYAKEVD TAAAEHEIRWLPNLTFDQRVAFTHKLDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:767
Z13648	VDAKYAKESQIAGHEIRWLPNLTFDQRVAFTHKLDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:768
Z13651	VDAKYAKEVSAASHEIRWLPNLTFDQRVAFTHKLDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:769
Z13652	VDAKYAKEQDAAAEHEIRWLPNLTFDQRVAFTHKLDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:770
Z13654	VDAKYAKELEAAKHEIRWLPNLTFDQRVAFTHKLDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:771
Z13655	VDAKYAKESKRAAEHEIRWLPNLTFDQRVAFTHKLDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:772
Z13656	VDAKYAKERYVKAGHEIRWLPNLTFDQRVAFTHKLDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:773
Z13657	VDAKYAKEFFSRAGHEIRWLPNLTFDQRVAFTHKLDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:774
Z13659	VDAKYAKEWQFAAEHEIRWLPNLTFDQRVAFTHKLDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:775
Z13663	VDAKYAKEWQIAAEHEIRWLPNLTFDQRVAFTHKLDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:776
Z13664	VDAKYAKELQEA AEHEIRWLPNLTFDQRVAFTHKLDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:777
Z13667	VDAKYAKERYRAAAAEHEIRWLPNLTFDQRVAFTHKLDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:778
Z13669	VDAKYAKELASAGHEIRWLPNLTFDQRVAFTHKLDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:779
Z13672	VDAKYAKEVQSASHEIRWLPNLTFDQRVAFTHKLDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:780
Z13674	VDAKYAKEI EDAGHEIRWLPNLTFDQRVAFTHKLDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:781
Z13675	VDAKYAKENQAAA AEHEIRWLPNLTFDQRVAFTHKLDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:782
Z13676	VDAKYAKEAT SAGHEIRWLPNLTFDQRVAFTHKLDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:783
Z13678	VDAKYAKEDEQAAA AEHEIRWLPNLTFDQRVAFTHKLDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:784

Figure 1DD

Designation	Sequence	SEQ ID NO:
Z13684	VDAKYAKENQAGHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:785
Z13688	VDAKYAKETSAAGHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:786
Z13691	VDAKYAKEDWAAAHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:787
Z13692	VDAKYAKEMQAGHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:788
Z13694	VDAKYAKELSDAAHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:789
Z13695	VDAKYAKEIDAAKHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:790
Z13697	VDAKYAKEAERAGHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:791
Z13706	VDAKYAKEDSDAAHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:792
Z13708	VDAKYAKEQKAAAGHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:793
Z13710	VDAKYAKEDWQAAHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:794
Z13711	VDAKYAKAKAAAHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:795
Z13714	VDAKYAKELSEAAHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:796
Z13716	VDAKYAKETEAAKHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:797
Z13719	VDAKYAKAKSAAHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:798
Z13720	VDAKYAKESAAAHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:799
Z13721	VDAKYAKEKERAHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:800
Z13725	VDAKYAKEDWDEAAHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:801
Z13727	VDAKYAKEEKDAAHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:802
Z13728	VDAKYAKEIENAAHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:803
Z13732	VDAKYAKETKEAAHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:804
Z13735	VDAKYAKELEAAKHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:805
Z13736	VDAKYAKEWAAEAAHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:806
Z13740	VDAKYAKESQAGHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:807
Z13742	VDAKYAKELSTAAHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:808
Z13747	VDAKYAKEIEEAGHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:809
Z13749	VDAKYAKELIQTASHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:810
Z13750	VDAKYAKEDSDAAHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:811
Z13751	VDAKYAKESASAGHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:812

Figure 1EE

Designation	Sequence	SEQ ID NO:
Z13752	VDAKYAKEVAKASHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:813
Z13758	VDAKYAKEVQEAHAHEIRWLPNLTFDQRVAFTHKLWDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:814
Z13759	VDAKYAKESYEAHAHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:815
Z13760	VDAKYAKETAEGHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:816
Z13761	VDAKYAKELEEAHAHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:817
Z13771	VDAKYAKEAAAAGHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:818
Z13773	VDAKYAKYVDAGHEIRWLPNLTFDQRVAFIRKLDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:819
Z13776	VDAKYAKEIQEAGHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:820
Z13777	VDAKYAKESATAAHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:821
Z13780	VDAKYAKEMMSAAHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:822
Z13782	VDAKYAKEREQASHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:823
Z13783	VDAKYAKEIEQAGHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:824
Z13786	VDAKYAKEHNAAAHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:825
Z13792	VDAKYAKEIEVAAHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:826
Z13796	VDAKYAKERAEEAHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:827
Z13799	VDAKYAKESELAHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:828
Z13806	VDAKYAKERYAAGHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:829
Z13808	VDAKYAKETANAAHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:830
Z13811	VDAKYAKEWYEAHAHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:831
Z13812	VDAKYAKEEQEAHAHEIRWLPNLTFDQRVAFTHKLWDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:832
Z13823	VDAKYAKEHDDAAHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:833
Z13824	VDAKYAKEWYSAHAHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:834
Z13838	VDAKYAKETSDAAHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:835
Z13840	VDAKYAKETYTTAAHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:836
Z13842	VDAKYAKEISQAAHAHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:837
Z13845	VDAKYAKENDDDAAHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:838
Z13846	VDAKYAKESEIAGHEIRWLPNLTFDQRVAFIRKLDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:839
Z13848	VDAKYAKEQADAAHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:840

Figure 1FF

Designation	Sequence	SEQ ID NO :
Z13849	VDKAYAKETESAAHEIRWLPNLTFDQRFVFIHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:841
Z13860	VDKAYAKEISDAKHEIRWLPNLTFDQRFVFIHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:842
Z13865	VDKAYAKEHINA AHEIRWLPNLTFDQRFVFIHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:843
Z13866	VDKAYAKEWLDAAHEIRWLPNLTFDQRFVFIHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:844
Z13875	VDKAYAKENAAAAHEIRWLPNLTFDQRFVFIHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:845
Z13879	VDKAYAKEAELAGHEIRWLPNLTFDQRFVFIHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:846
Z13480	VDKAYAKEVTDAGHEIRWLPNLTFDQRFVFIHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:847
Z13481	VDKAYAKELDASHEIRWLPNLTFDQRFVFIHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:848
Z13482	VDKAYAKEINIAKHEIRWLPNLTFDQRFVFIHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:849
Z13483	VDKAYAKETEYAKHEIRWLPNLTFDQRFVFIHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:850
Z13484	VDKAYAKESAEAGHEIRWLPNLTFDQRFVFIHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:851
Z13485	VDKAYAKEYSNAAHEIRWLPNLTFDQRFVFIHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:852
Z13487	VDKAYAKETNAGHEIRWLPNLTFDQRFVFIHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:853
Z13488	VDKAYAKEVEFAAHEIRWLPNLTFDQRFVFIHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:854
Z13489	VDKAYAKEVELAGHEIRWLPNLTFDQRFVFIHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:855
Z13490	VDKAYAKEVLKASHEIRWLPNLTFDQRFVFIHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:856
Z13491	VDKAYAKEIANAGHEIRWLPNLTFDQRFVFIHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:857
Z13493	VDKAYAKEMYKAGHEIRWLPNLTFDQRFVFIHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:858
Z13495	VDKAYAKEHANAQHEIRWLPNLTFDQRFVFIHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:859
Z13496	VDKAYAKEVDIASHEIRWLPNLTFDQRFVFIHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:860
Z13497	VDKAYAKEEVFAAHEIRWLPNLTFDQRFVFIHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:861
Z13499	VDKAYAKEFNTAAHEIRWLPNLTFDQRFVFIHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:862
Z13501	VDKAYAKEVDVAGHEIRWLPNLTFDQRFVFIHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:863
Z13502	VDKAYAKEWSLAAHEIRWLPNLTFDQRFVFIHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:864
Z13503	VDKAYAKELDDAAHEIRWLPNLTFDQRFVFIHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:865
Z13505	VDKAYAKERHEAGHEIRWLPNLTFDQRFVFIHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:866
Z13506	VDKAYAKEISDAIHEIRWLPNLTFDQRFVFIHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:867
Z13507	VDKAYAKEWEIAGHEIRWLPNLTFDQRFVFIHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:868

Figure 1GG

Designation	Sequence	SEQ ID NO:
Z13508	VDKAKERYWASHEIRWLPNLTFDQRVAFTHKLNDDPSQSSELLSEAKKLNDSQAPK	SEQ ID NO:869
Z13512	VDKAKELDWAGHEIRWLPNLTFDQRVAFTHKLLDDPSQSSELLSEAKKLNDSQAPK	SEQ ID NO:870
Z13515	VDKAKRFQSKAGHEIRWLPNLTFDQRVAFTHKLDKDDPSQSSELLSEAKKLNDSQAPK	SEQ ID NO:871
Z13518	VDKAKELIEAAQHEIRWLPNLTFDQRVAFTHKLNDDPSQSSELLSEAKKLNDSQAPK	SEQ ID NO:872
Z13519	VDKAKREHEQAAGHEIRWLPNLTFDQRVAFTHKLVDDPSQSSELLSEAKKLNDSQAPK	SEQ ID NO:873
Z13520	VDKAKAEAEQAGHEIRWLPNLTFDQRVAFTHKLDKDDPSQSSELLSEAKKLNDSQAPK	SEQ ID NO:874
Z13522	VDKAKEDVYAAHEIRWLPNLTFDQRVAFTHKLNDDPSQSSELLSEAKKLNDSQAPK	SEQ ID NO:875
Z13525	VDKAKESYSAAGHEIRWLPNLTFDQRVAFTEKLRDDPSQSSELLSEAKKLNDSQAPK	SEQ ID NO:876
Z13561	VDKAKELATASHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLNDSQAPK	SEQ ID NO:877
Z13572	VDKAKERYRVAGHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLNDSQAPK	SEQ ID NO:878
Z13575	VDKAKKEVVSAGHEIRWLPNLTFDQRVAFTHKLLDDPSQSSELLSEAKKLNDSQAPK	SEQ ID NO:879
Z13576	VDKAKESQAQAAGHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLNDSQAPK	SEQ ID NO:880
Z13584	VDKAKESYSAAGHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLNDSQAPK	SEQ ID NO:881
Z13589	VDKAKKEQKEAAAEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLNDSQAPK	SEQ ID NO:882
Z13590	VDKAKKEAAIAGKEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLNDSQAPK	SEQ ID NO:883
Z13591	VDKAKKEISKAKHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLNDSQAPK	SEQ ID NO:884
Z13593	VDKAKKESVAAAHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLNDSQAPK	SEQ ID NO:885
Z13595	VDKAKKEIQAKHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLNDSQAPK	SEQ ID NO:886
Z13599	VDKAKAKEITSAKHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLNDSQAPK	SEQ ID NO:887
Z13601	VDKAKKEQDVAAGHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLNDSQAPK	SEQ ID NO:888
Z13603	VDKAKKELERAAGHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLNDSQAPK	SEQ ID NO:889
Z13607	VDKAKENQLAAAGHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLNDSQAPK	SEQ ID NO:890
Z13608	VDKAKKEISQAKHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLNDSQAPK	SEQ ID NO:891
Z13610	VDKAKKEIANASHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLNDSQAPK	SEQ ID NO:892
Z13614	VDKAKKEWQAAAAGHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLNDSQAPK	SEQ ID NO:893
Z13618	VDKAKKERKDGAGHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLNDSQAPK	SEQ ID NO:894
Z13619	VDKAKKEITQAKHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLNDSQAPK	SEQ ID NO:895
Z13623	VDKAKKEFIQAGHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLNDSQAPK	SEQ ID NO:896

Figure 1HH

Designation	Sequence	SEQ ID NO:
Z13630	VDKAYAKENWTASHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLNDQSAPK	SEQ ID NO:897
Z13631	VDKAYAKEKFVAAHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLNDQSAPK	SEQ ID NO:898
Z13632	VDKAYAREADSAGAEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLNDQSAPK	SEQ ID NO:899
Z13636	VDKAYAKESVAAAEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLNDQSAPK	SEQ ID NO:900
Z13642	VDKAYAKEVDLAGREIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLNDQSAPK	SEQ ID NO:901
Z13643	VDKAYAKEQERAAAEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLNDQSAPK	SEQ ID NO:902
Z13646	VDKAYAKEIWQAKHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLNDQSAPK	SEQ ID NO:903
Z13647	VDKAYAKELNQAKHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLNDQSAPK	SEQ ID NO:904
Z13649	VDKAYAKELQQASHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLNDQSAPK	SEQ ID NO:905
Z13650	VDKAYAKEINQAKHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLNDQSAPK	SEQ ID NO:906
Z13653	VDKAYAKELVLAGHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLNDQSAPK	SEQ ID NO:907
Z13658	VDKAYAKELTSAHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLNDQSAPK	SEQ ID NO:908
Z13661	VDKAYAKEWNAAREIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLNDQSAPK	SEQ ID NO:909
Z13662	VDKAYAKELIHAKHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLNDQSAPK	SEQ ID NO:910
Z13665	VDKAYAKEVLTAKEHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLNDQSAPK	SEQ ID NO:911
Z13666	VDKAYAKENSKAAHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLNDQSAPK	SEQ ID NO:912
Z13668	VDKAYAKEVMTAKHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLNDQSAPK	SEQ ID NO:913
Z13671	VDKAYAKEARDAAHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLNDQSAPK	SEQ ID NO:914
Z13673	VDKAYAKERSKAAHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLNDQSAPK	SEQ ID NO:915
Z13677	VDKAYAKEIYSKHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLNDQSAPK	SEQ ID NO:916
Z13679	VDKAYAKEVQSAKHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLNDQSAPK	SEQ ID NO:917
Z13680	VDKAYAKETLEAAHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLNDQSAPK	SEQ ID NO:918
Z13681	VDKAYAKEQMAAHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLNDQSAPK	SEQ ID NO:919
Z13685	VDKAYAKENKNAHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLNDQSAPK	SEQ ID NO:920
Z13687	VDKAYAKETESAKHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLNDQSAPK	SEQ ID NO:921
Z13689	VDKAYAKETVQAKHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLNDQSAPK	SEQ ID NO:922
Z13693	VDKAYAKEIASAKHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLNDQSAPK	SEQ ID NO:923
Z13698	VDKAYAKEVMDAKHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLNDQSAPK	SEQ ID NO:924

Figure 1II

Designation	Sequence	SEQ ID NO:
Z13699	VDKAYAKETDAAKHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLNDQSAPK	SEQ ID NO:925
Z13701	VDKAYAKELQIAKHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLNDQSAPK	SEQ ID NO:926
Z13702	VDKAYAKRWKDAQAQHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLNDQSAPK	SEQ ID NO:927
Z13703	VDKAYAKERDAAHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLNDQSAPK	SEQ ID NO:928
Z13704	VDKAYAKEIAAAKHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLNDQSAPK	SEQ ID NO:929
Z13707	VDKAYAKESVKAKHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLNDQSAPK	SEQ ID NO:930
Z13709	VDKAYAKENERAKHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLNDQSAPK	SEQ ID NO:931
Z13712	VDKAYAKYKRAKHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLNDQSAPK	SEQ ID NO:932
Z13713	VDKAYAKEVRAASHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLNDQSAPK	SEQ ID NO:933
Z13717	VDKAYAKEDKRAAHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLNDQSAPK	SEQ ID NO:934
Z13718	VDKAYAKESKAGKEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLNDQSAPK	SEQ ID NO:935
Z13722	VDKAYAKEINRAKHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLNDQSAPK	SEQ ID NO:936
Z13724	VDKAYAKETQQAASHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLNDQSAPK	SEQ ID NO:937
Z13729	VDKAYAKENQSAASHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLNDQSAPK	SEQ ID NO:938
Z13730	VDKAYAKEAKQASHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLNDQSAPK	SEQ ID NO:939
Z13731	VDKAYAKEAAQAAHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLNDQSAPK	SEQ ID NO:940
Z13733	VDKAYAKEVQYASHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLNDQSAPK	SEQ ID NO:941
Z13734	VDKAYAKELRNAASHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLNDQSAPK	SEQ ID NO:942
Z13737	VDKAYAKEQRAAAHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLNDQSAPK	SEQ ID NO:943
Z13739	VDKAYAKEASEAAAEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLNDQSAPK	SEQ ID NO:944
Z13741	VDKAYAKESVIAAHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLNDQSAPK	SEQ ID NO:945
Z13744	VDKAYAKEILRAKHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLNDQSAPK	SEQ ID NO:946
Z13746	VDKAYAKESKTAASHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLNDQSAPK	SEQ ID NO:947
Z13748	VDKAYAKELAEASHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLNDQSAPK	SEQ ID NO:948
Z13753	VDKAYAKEATTAKEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLNDQSAPK	SEQ ID NO:949
Z13754	VDKAYAKEIFENAAHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLNDQSAPK	SEQ ID NO:950
Z13755	VDKAYAKEAKDAASHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLNDQSAPK	SEQ ID NO:951
Z13757	VDKAYAKERLEAAAEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLNDQSAPK	SEQ ID NO:952

Figure 1JJ

Designation	Sequence	SEQ ID NO:
Z1 3762	VDKAYAKEQMEAAAEIRWLPNLTFDQRVAFTHKLDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO: 953
Z1 3763	VDKAYAKEVKTASHEIRWLPNLTFDQRVAFTHKLDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO: 954
Z1 3764	VDKAYAKESFEASHEIRWLPNLTFDQRVAFTHKLDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO: 955
Z1 3765	VDKAYAKEIKSAKHEIRWLPNLTFDQRVAFTHKLDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO: 956
Z1 3766	VDKAYAKEIKNAKHEIRWLPNLTFDQRVAFTHKLDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO: 957
Z1 3768	VDKAYAKEIQEAKHEIRWLPNLTFDQRVAFTHKLDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO: 958
Z1 3769	VDKAYAKERQNAAEIRWLPNLTFDQRVAFTHKLDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO: 959
Z1 3770	VDKAYAKEVLQAKHEIRWLPNLTFDQRVAFTHKLDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO: 960
Z1 3772	VDKAYAKEANVASHEIRWLPNLTFDQRVAFTHKLDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO: 961
Z1 3774	VDKAYAKEIDAAKHEIRWLPNLTFDQRVAFTHKLDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO: 962
Z1 3775	VDKAYAKETASAAAEIRWLPNLTFDQRVAFTHKLDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO: 963
Z1 3778	VDKAYAKEWKQAAAEIRWLPNLTFDQRVAFTHKLDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO: 964
Z1 3779	VDKAYAKETASASHEIRWLPNLTFDQRVAFTHKLDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO: 965
Z1 3781	VDKAYAKESIVAGHEIRWLPNLTFDQRVAFTHKLDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO: 966
Z1 3784	VDKAYAKEIKQAKHEIRWLPNLTFDQRVAFTHKLDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO: 967
Z1 3785	VDKAYAKEQATASHEIRWLPNLTFDQRVAFTHKLDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO: 968
Z1 3787	VDKAYAKEINAAKHEIRWLPNLTFDQRVAFTHKLDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO: 969
Z1 3788	VDKAYAKEVVKRAGHEIRWLPNLTFDQRVAFTHKLDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO: 970
Z1 3789	VDKAYAKESRNAAEIRWLPNLTFDQRVAFTHKLDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO: 971
Z1 3791	VDKAYAKESITASAEIRWLPNLTFDQRVAFTHKLDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO: 972
Z1 3793	VDKAYAKEAATAAEIRWLPNLTFDQRVAFTHKLDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO: 973
Z1 3794	VDKAYAKEVYAAKHEIRWLPNLTFDQRVAFTHKLDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO: 974
Z1 3795	VDKAYAKEISRAKHEIRWLPNLTFDQRVAFTHKLDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO: 975
Z1 3798	VDKAYKEYVTAAEIRWLPNLTFDQRVAFTHKLDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO: 976
Z1 3800	VDKAYAKEHIDAGHEIRWLPNLTFDQRVAFTHKLDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO: 977
Z1 3801	VDKAYAKEITLQAKHEIRWLPNLTFDQRVAFTHKLDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO: 978
Z1 3802	VDKAYAKENSQAKHEIRWLPNLTFDQRVAFTHKLDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO: 979
Z1 3803	VDKAYAYEYRVAKEIRWLPNLTFDQRVAFTHKLDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO: 980

Figure 1KK

Designation	Sequence	SEQ ID NO:
Z13804	VDAKYAKEIYNAGHEIRWLPNLTFDQRVAFIRKLRDDPSQSSSELLSEAKKLND SQAPK	SEQ ID NO:981
Z13805	VDAKYAKESNEAAAAEIRWLPNLTFDQRVAFIRKLRDDPSQSSSELLSEAKKLND SQAPK	SEQ ID NO:982
Z13809	VDAKYAKESQLAAAEIRWLPNLTFDQRVAFIRKLRDDPSQSSSELLSEAKKLND SQAPK	SEQ ID NO:983
Z13810	VDAKYAKELKEAGHEIRWLPNLTFDQRVAFIRKLRDDPSQSSSELLSEAKKLND SQAPK	SEQ ID NO:984
Z13813	VDAKYAKETRVASVEIRWLPNLTFDQRVAFIRKLRDDPSQSSSELLSEAKKLND SQAPK	SEQ ID NO:985
Z13818	VDAKYAKELRTAAHEIRWLPNLTFDQRVAFIRKLRDDPSQSSSELLSEAKKLND SQAPK	SEQ ID NO:986
Z13820	VDAKYAKEITYAHFEIRWLPNLTFDQRVAFIRKLRDDPSQSSSELLSEAKKLND SQAPK	SEQ ID NO:987
Z13821	VDAKYAKEEAQAASHEIRWLPNLTFDQRVAFIRKLRDDPSQSSSELLSEAKKLND SQAPK	SEQ ID NO:988
Z13822	VDAKYAKEITSAASHEIRWLPNLTFDQRVAFIRKLRDDPSQSSSELLSEAKKLND SQAPK	SEQ ID NO:989
Z13825	VDAKYAKEVKTASHEIRWLPNLTFDQRVAFIRKLRDDPSQSSSELLSEAKKLND SQAPK	SEQ ID NO:990
Z13826	VDAKYAKETKVAASHEIRWLPNLTFDQRVAFIRKLRDDPSQSSSELLSEAKKLND SQAPK	SEQ ID NO:991
Z13829	VDAKYAKEDLVAQHEIRWLPNLTFDQRVAFIRKLRDDPSQSSSELLSEAKKLND SQAPK	SEQ ID NO:992
Z13830	VDAKYAKETQTAFNEIRWLPNLTFDQRAAFIRKLRDDPSQSSSELLSEAKKLND SQAPK	SEQ ID NO:993
Z13831	VDAKYAKEIKDAGHEIRWLPNLTFDQRVAFIRKLRDDPSQSSSELLSEAKKLND SQAPK	SEQ ID NO:994
Z13836	VDAKYAKEYKEAGHEIRWLPNLTFDQRVAFIRKLRDDPSQSSSELLSEAKKLND SQAPK	SEQ ID NO:995
Z13839	VDAKYAKEAALAAHEIRWLPNLTFDQRVAFIRKLRDDPSQSSSELLSEAKKLND SQAPK	SEQ ID NO:996
Z13843	VDAKYAKEQERAAHEIRWLPNLTFDQRVAFIRKLRDDPSQSSSELLSEAKKLND SQAPK	SEQ ID NO:997
Z13844	VDAKYAKEDWFDAAHEIRWLPNLTFDQRVAFIRKLRDDPSQSSSELLSEAKKLND SQAPK	SEQ ID NO:998
Z13847	VDAKYAKEI IQAKHEIRWLPNLTFDQRVAFIRKLRDDPSQSSSELLSEAKKLND SQAPK	SEQ ID NO:999
Z13850	VDAKYAKELTNAKHEIRWLPNLTFDQRVAFIRKLRDDPSQSSSELLSEAKKLND SQAPK	SEQ ID NO:1000
Z13851	VDAKYAKEIQLAKHEIRWLPNLTFDQRVAFIRKLRDDPSQSSSELLSEAKKLND SQAPK	SEQ ID NO:1001
Z13854	VDAKYAKEIHDAAHEIRWLPNLTFDQRVAFIRKLRDDPSQSSSELLSEAKKLND SQAPK	SEQ ID NO:1002
Z13869	VDAKYAKEVKIASHEIRWLPNLTFDQRVAFIRKLRDDPSQSSSELLSEAKKLND SQAPK	SEQ ID NO:1003
Z13870	VDAKYAKEQHSAAHEIRWLPNLTFDQRVAFIRKLRDDPSQSSSELLSEAKKLND SQAPK	SEQ ID NO:1004
Z13872	VDAKYAKEVFFAASAEIRWLPNLTFDQRVAFIRKLRDDPSQSSSELLSEAKKLND SQAPK	SEQ ID NO:1005
Z13874	VDAKYAKETDLAGHEIRWLPNLTFDQRVAFIRKLRDDPSQSSSELLSEAKKLND SQAPK	SEQ ID NO:1006
Z13877	VDAKYAKEANFAGHEIRWLPNLTFDQRVAFIRKLRDDPSQSSSELLSEAKKLND SQAPK	SEQ ID NO:1007
Z13878	VDAKYAKEFETAGHEIRWLPNLTFDQRVAFIRKLRDDPSQSSSELLSEAKKLND SQAPK	SEQ ID NO:1008

Figure 1LL

Designation	Sequence	SEQ ID NO:
Z13880	VDAKYAKENVNLAKEIRWLPNLTFDQRVAFTHKLDLDDPSQSSELLSEAKKLNDQSAPK	SEQ ID NO:1009
Z13881	VDAKYAKEADTAAGHEIRWLPNLTFDQRVAFYKLRDDPSQSSELLSEAKKLNDQSAPK	SEQ ID NO:1010
Z13882	VDAKYAKRFVDAGHEIRWLPNLTFDQRVAFYKLRDDPSQSSELLSEAKKLNDQSAPK	SEQ ID NO:1011
Z13883	VDAKYAKEDHKAEGHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLNDQSAPK	SEQ ID NO:1012
Z13884	VDAKYAKETVDAGHEIRWLPNLTFDQRVAFTHKLDLDDPSQSSELLSEAKKLNDQSAPK	SEQ ID NO:1013
Z13885	VDAKYAKESQAGHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLNDQSAPK	SEQ ID NO:1014
Z13886	VDAKYAKESWSSAAHEIRWLPNLTFDQRVAFTHKLDLDDPSQSSELLSEAKKLNDQSAPK	SEQ ID NO:1015
Z13887	VDAKYAKEVAVAGHEIRWLPNLTFDQRVAFYKLRDDPSQSSELLSEAKKLNDQSAPK	SEQ ID NO:1016
Z13888	VDAKYAKESAEAGHEIRWLPNLTFDQRVAFTEKLRDDPSQSSELLSEAKKLNDQSAPK	SEQ ID NO:1017
Z13889	VDAKYAKEAVAAAGHEIRWLPNLTFDQRVAFTHKLDLDDPSQSSELLSEAKKLNDQSAPK	SEQ ID NO:1018
Z13890	VDAKYAKRFQIAGHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLNDQSAPK	SEQ ID NO:1019
Z13891	VDAKYAKELMVAKHEIRWLPNLTFDQRVAFTHKLDLDDPSQSSELLSEAKKLNDQSAPK	SEQ ID NO:1020
Z13892	VDAKYAKEDYDAAHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLNDQSAPK	SEQ ID NO:1021
Z13893	VDAKYAKEVLEAKHEIRWLPNLTFDQRVAFTHKLDLDDPSQSSELLSEAKKLNDQSAPK	SEQ ID NO:1022
Z13894	VDAKYAKESIAASHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLNDQSAPK	SEQ ID NO:1023
Z13895	VDAKYAKEVAEAGHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLNDQSAPK	SEQ ID NO:1024
Z13896	VDAKYAKEQAKAAHEIRWLPNLTFDQRVAFTHKLDLDDPSQSSELLSEAKKLNDQSAPK	SEQ ID NO:1025
Z13897	VDAKYAKERDDAAHEIRWLPNLTFDQRVAFTEKLRDDPSQSSELLSEAKKLNDQSAPK	SEQ ID NO:1026
Z13898	VDAKYAKEAKDASHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLNDQSAPK	SEQ ID NO:1027
Z13899	VDAKYAKEASSAAHEIRWLPNLTFDQRVAFYKLRDDPSQSSELLSEAKKLNDQSAPK	SEQ ID NO:1028
Z13900	VDAKYAKEWMEASHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLNDQSAPK	SEQ ID NO:1029
Z13901	VDAKYAKEQKNAAGHEIRWLPNLTFDQRVAFTEKLRDDPSQSSELLSEAKKLNDQSAPK	SEQ ID NO:1030
Z13902	VDAKYAKETENAKHEIRWLPNLTFDQRVAFTHKLDLDDPSQSSELLSEAKKLNDQSAPK	SEQ ID NO:1031
Z13903	VDAKYAKENVNRASHEIRWLPNLTFDQRVAFTHKLDLDDPSQSSELLSEAKKLNDQSAPK	SEQ ID NO:1032
Z13904	VDAKYAKERLLAGHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLNDQSAPK	SEQ ID NO:1033
Z13906	VDAKYAKEVSIAGHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLNDQSAPK	SEQ ID NO:1034
Z13907	VDAKYAKEKEVAAGHEIRWLPNLTFDQRVAFTHKLDLDDPSQSSELLSEAKKLNDQSAPK	SEQ ID NO:1035
Z13908	VDAKYAKESERASHEIRWLPNLTFDQRVAFTHKLRDDPSQSSELLSEAKKLNDQSAPK	SEQ ID NO:1036

Figure 1MM

Designation	Sequence	SEQ ID NO :
Z13909	VDKAYAKENNEAAHEIRWLPNLTFDQRFVFIHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:1037
Z13910	VDKAYAKENVDAAAHEIRWLPNLTFDQRFVFIHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:1038
Z13911	VDKAYAKEADAASHEIRWLPNLTFDQRFVFIHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:1039
Z13912	VDKAYAKELESASHEIRWLPNLTFDQRFVFIHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:1040
Z13913	VDKAYAKEEQIAAHEIRWLPNLTFDQRFVFIHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:1041
Z13914	VDKAYAKEFELAGHEIRWLPNLTFDQRFVFIHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:1042
Z13915	VDKAYAKEAFVAQHEIRWLPNLTFDQRFVFIHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:1043
Z13916	VDKAYAKEALKASHEIRWLPNLTFDQRFVFIHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:1044
Z13917	VDKAYAKELEERAGHEIRWLPNLTFDQRFVFIHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:1045
Z13918	VDKAYAKEVEWAKHEIRWLPNLTFDQRFVFIHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:1046
Z13919	VDKAYAKEKASQAHEIRWLPNLTFDQRFVFIHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:1047
Z13920	VDKAYAKEETEIAKHEIRWLPNLTFDQRFVFIHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:1048
Z13921	VDKAYAKEVNLAHEIRWLPNLTFDQRFVFIHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:1049
Z13922	VDKAYAKEEEAAHEIRWLPNLTFDQRFVFIHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:1050
Z13923	VDKAYAKETDRAKHEIRWLPNLTFDQRFVFIHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:1051
Z13925	VDKAYAKEFAQAQHEIRWLPNLTFDQRFVFIHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:1052
Z13926	VDKAYAKETDEASHEIRWLPNLTFDQRFVFIHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:1053
Z13927	VDKAYAKENADAGHEIRWLPNLTFDQRFVFIHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:1054
Z13928	VDKAYAKESTQAAHEIRWLPNLTFDQRFVFIHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:1055
Z13929	VDKAYAKEQALAAHEIRWLPNLTFDQRFVFIHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:1056
Z13930	VDKAYAKEAHAASHEIRWLPNLTFDQRFVFIHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:1057
Z13932	VDKAYAKEVDNAGHEIRWLPNLTFDQRFVFIHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:1058
Z13993	VDKAYAKEAGRAAHEIRWLPNLTWDQRFVFIHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:1059
Z11948	AEAKYAKEQDAAAHEIRWLPNLTFDQRFVFIHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:1060
Z11946	AEAKYAKEFESAAHEIRWLPNLTVDQRFVFIHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:1061
Z11947	AEAKYAKEWMRAAHEIRWLPNLTFDQRFVFIHKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:1062
PP013	LAEAKEAANAELDSYGVSDFYKRLIDKAKTVEGVKALKDAILAALP	SEQ ID NO:1063
Z03638	AEAKYAKELGWATWEIENLPNLTGVQVKAFIDKLRDDPSQSSELLSEAKKLND SQAPK	SEQ ID NO:1064

Figure 1NN

Designation	Sequence	SEQ ID NO:
Human α FcRn	AESHLSLLYHLTAVSSPAGTPAFWVGWLGPPQQYLSYNSLRGEALPCGAWWENQVSWYWEKE TTDLRIKEKLFLEAFKALGGKPYTLQGLGCELGPDNTSVPTAKFALNGEEFMFDLKQGTWG GDWPEALAIISQRWQQDKAANKELTFLLFSCPHRLREHLERGRGNLEWKEPPSMRLKARPSPG FSVLTCSAFSFPPELQLRFLRNLGLAAGTGQDGFNPNSDGSFHASSLTVKSGDEHHYCCIVQH AGLAQPLRVEL	SEQ ID NO:1065
	IQRTPKIQVYSRHPAENGKSNFNCYVSGFHPSPDIEVDLLKNGERIEKVEHSDLSFSKDWSEFYL LYYTEFTPTTEKDEYACRVNHNHVLISQPKIVKWDRDM	SEQ ID NO:1066
	IQKTPQIQVYSRHPPENGKPNILNCYVTFPHPPHIEIQMLKNGKKIPKVMSDMSFSKDWSEFYI LAHTEFTPTTETDITYACRVKHASMAEPKTVYWDKDM	SEQ ID NO:1067
	MGVPRPQFWALGILLFLPGSLGAESHLSLLYHLTAVSSPAGTPAFWVGWLGPPQQYLSYNSL RGEALPCGAWWENQVSWYWEKETDRLRIKEKLFLEAFKALGGKPYTLQGLGCELGPDNTSV PTAKFALNGEEFMFDLKQGTWGDWPEALAIISQRWQQDKAANKELTFLLFSCPHRLREHLER GRGNLEWKEPPSMRLKARPSPGFSVLTCSAFSFPPELQLRFLRNLGLAAGTGQDGFNPNSDGS FHASSLTVKSGDEHHYCCIVQHAGLAQPLRVELESPAKSSVLVVGIVIGVLLLTAAAVGGALL WRRMRSGLPAPWISLRGDDTGVLPTPGEAQDADLKDVNVIPTATA	SEQ ID NO:1068
mFcRn-eGFP	MGMLPWLALSILLVLPLQTWGSETRPPIMYHLTAVSNPSTGLPSFWATGWLGPPQQYLTYNLSLRQ EADPCGAWWENQVSWYWEKETDRLKSKEQLFLEALKTLKILNGQKRGTYTLQGLLGCELASD NSSVPTAVFALNGEEFMKNPRIGNWTGEWPETEIVANLWMKQPDAAARKESEFLLNSCPERLLG HLERGRNLEWKEPPSMRLKARPGNSGSSVLTCAAFSFPPELKFRLRNLGLASGSGNCSTGPN GDGSHAWSLLEVKRGDEHHYQCQVEHEGLAQPLTVDLDSARSSVPVVGIVLGLLLVVVAIAG GVLLWGRMRSGLPAPWLSLGGDSDGLLPGGNLPPAEPOGANAFPATIS	SEQ ID NO:1069
Murine α FcRn	SETRPPLMYHLTAVSNPSTGLPSFWATGWLGPPQQYLTYNLSLQEADPCGAWWENQVSWYWEKE TTDLKSKEQLFLEALKTLKILNGQKRGTYTLQGLLGCELASDSSVPTAVFALNGEEFMKNP RIGNWTGEWPETEIVANLWMKQPDAAARKESEFLLNSCPERLLGHLERGRNLEWKEPPSMRLKA RPGNSGSSVLTCAAFSFPPELKFRLRNLGLASGSGNCSTGPNGDGSHAWSLLEVKRGDEHHY QCQVEHEGLAQPLTVDL	SEQ ID NO:1070

Figure 2A




Figure 2B




Figure 2C




Figure 2D




Figure 2E




Figure 3




Figure 4




Figure 5




Figure 6




Figure 7A




Figure 7B




Figure 7C




Figure 8




Figure 9




Figure 10

