



(86) Date de dépôt PCT/PCT Filing Date: 2001/10/26
(87) Date publication PCT/PCT Publication Date: 2002/05/02
(45) Date de délivrance/Issue Date: 2009/10/06
(85) Entrée phase nationale/National Entry: 2003/04/25
(86) N° demande PCT/PCT Application No.: CA 2001/001508
(87) N° publication PCT/PCT Publication No.: 2002/034047
(30) Priorité/Priority: 2000/10/26 (CA2,324,677)

(51) Cl.Int./Int.Cl. *A01N 25/30* (2006.01),
A01P 13/00 (2006.01), *C05G 3/00* (2006.01),
C05G 3/06 (2006.01)

(72) Inventeurs/Inventors:
STEWART, JAMES F., CA;
REINARTZ, HEINRICH J., CA;
BROWN, WILLIAM G., CA

(73) Propriétaire/Owner:
ADJUVANTS PLUS INC., CA

(74) Agent: BORDEN LADNER GERVAIS LLP

(54) Titre : COMPOSITION D'ADJUVANT DE FORMULATION AGROCHIMIQUE ET UTILISATIONS DE CELLE-CI
(54) Title: AGROCHEMICAL FORMULATION AID COMPOSITION AND USES THEREOF

(57) **Abrégé/Abstract:**

There is provided an agrochemical formulation aid composition for preparing bioactive and sprayable agrochemicals, wherein various components for the composition were selected from mineral oil paraffinic distillate and/or aromatic hydrocarbon distillate; 2N-octanol; oleyl-cetyl alcohol; polyoxyethylene (2) oleylether ; polyoxyethylene (8) nonylphenoethin and/or ethoxylated tallow amine blend; lauryl sulphate; fatty alcohol alkoxylate; terpenes, diammonium phosphate; tetrasodium ethylene diamine tetracetate; cab-o-sil; fatty acid methyl ester; (C18) free fatty acid blend; N-butanol; and methyl alcohol. Also provided are methods of preparing the formulation aid composition on site by mixing various components and methods of preparing sprayable and bioactive agrochemical systems using the formulation aid and non-formulated or formulated agrochemicals. Also provided are uses of the formulation aid in preparing sprayable and bioactive agrochemical systems for controlling pests.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
2 May 2002 (02.05.2002)

PCT

(10) International Publication Number
WO 02/34047 A1

(51) International Patent Classification⁷: **A01N 25/30**,
25/02

(21) International Application Number: PCT/CA01/01508

(22) International Filing Date: 26 October 2001 (26.10.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
2,324,677 26 October 2000 (26.10.2000) CA

(71) Applicant (for all designated States except US): **AD-
JUVANTS PLUS INC.** [CA/CA]; 1338 Lincoln Road,
Kingsville, Ontario N9Y 2S5 (CA).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **STEWART, James,
F.** [CA/CA]; 10 Caryndale Drive, Kitchener, Ontario N2R
1C7 (CA). **REINARTZ, Heinrich, J.** [CA/CA]; 4096
Hwy 640, Harvey Station, New Brunswick E0H 1H0 (CA).
BROWN, William, G. [CA/CA]; 1338 Lincoln Road,
Kingsville, Ontario N9Y 2S5 (CA).

(74) Agents: **CONN, David, L.** et al.; Borden Ladner Ger-
vais LLP, 1000-60 Queen Street, Ottawa, Ontario K1P 5Y7
(CA).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG,
SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN,
YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,
CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD,
TG).


Published:

- with international search report
- before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: AGROCHEMICAL FORMULATION AID COMPOSITION AND USES THEREOF

(57) Abstract: There is provided an agrochemical formulation aid composition for preparing bioactive and sprayable agrochemicals, wherein various components for the composition were selected from mineral oil paraffinic distillate and/or aromatic hydrocarbon distillate; 2N-octanol; oleyl-cetyl alcohol; polyoxyethylene (2) oleylether ; polyoxyethylene (8) nonylphenoletin and/or ethoxy-
lated tallow amine blend; lauryl sulphate; fatty alcohol alkoxylate; terpenes, diammonium phosphate; tetrasodium ethylene diamine tetracetate; cab-o-sil; fatty acid methyl ester; (C18) free fatty acid blend; N-butanol; and methyl alcohol. Also provided are methods of preparing the formulation aid composition on site by mixing various components and methods of preparing sprayable and bioac-
tive agrochemical systems using the formulation aid and non-formulated or formulated agrochemicals. Also provided are uses of the
formulation aid in preparing sprayable and bioactive agrochemical systems for controlling pests.



WO 02/34047 A1

AGROCHEMICAL FORMULATION AID COMPOSITION AND USES THEREOF**FIELD OF THE INVENTION**

This invention relates to agrochemical formulation aid compositions, their uses,
5 and processes of preparing agrochemical systems of non-formulated and formulated
agrochemicals using the formulation aid compositions to obtain improved sprayability and
bioactivity.

BACKGROUND OF THE INVENTION

10 Agriculture chemicals (agrochemicals) such as pesticides including insecticides,
fungicides, rodenticides, and herbicides are materials that provide control of agricultural
pests including insects, pathogens, rodents, and weeds. In order for many agrochemicals
to be used as a means of controlling pests, these agrochemicals have to be incorporated
into an agriculturally acceptable carrier. Generally, the agrochemical is modified into a
15 soluble ester, amine salt, or dissolved in a solvent system and added to this are products to
create an agriculturally acceptable spray solution or suspension with water as a carrier.
The carrier systems may also include several other components.

Agricultural adjuvants are materials that modify the performance of an
agrochemical and may also improve the physical properties of an agricultural formulation.

20 For example, an activator adjuvant increases the biological efficacy of an agrochemical.
Also, fertilizers such as urea or diammonium phosphate are frequently used as adjuvants
or are used with other adjuvants to improve the efficacy of agrochemical formulations. A
compatibility agent prevents the chemical interaction of two or more agrochemical
components in a mixture. It could also improve the homogeneity of additional
25 components such as fertilizers with other agrochemicals in a mixture. A wetting agent or
spreading agent increases the surface area covered by a given volume of a spray mixture.

Herein, the term "non-formulated agrochemical" includes pesticides that are sold
as technical acids or as technical acid grade products. The terms "formulated
agrochemical" and "pre-formulated agrochemical" have been used interchangeably and
30 includes pesticides that are sold as ester, amine salts, or in a solvent.

Currently, most agrochemicals have to be pre-formulated e.g., as emulsifiable

concentrates, flowables, soluble powders, or soluble liquids to enable their application e.g., by spraying, on a crop and to make them bioactive for a targeted substrate e.g., a pest. However, the formulated products sold in the form of amines or esters have higher vapor pressure. Therefore, they have a tendency to move off the site of application to trespass
5 and damage adjacent crops and horticultural plantings. The use of solvents such as xylene, isobutanol, and dimethyl amine in formulated products presents the problem of toxicity, odor, and potential explosiveness to the user and neighbors.

However, all prior art adjuvants lack versatility or they often limit the form in which an agrochemical may be pre-formulated as a manufactured product. Further, a pre-
10 formulated manufactured product when shipped must be stored under appropriate environmental conditions to ensure that the agrochemical is not adversely affected in terms of its activity and to ensure that the formulation remains stable. For example, agrochemical formulations often include emulsions in which water is one of the phases. These agrochemical formulations must be stored under conditions that protect the
15 emulsion from freezing. Further, it should be noted that different levels of adjuvants are used with a given level of the pesticide for different crop applications, different stages of crop growth, different weather or climate conditions, and target species. Pre-formulating a given agrochemical to meet such diverse needs thus requires the construction of formulation facilities that use costly energy and create toxic wastes as byproducts such as
20 aromatic petrochemicals. Further, the pre-formulated agrochemical products often require the addition of an adjuvant or water conditioner to provide enhanced bioactivity and/or spray applicability.

The present invention provides agrochemical formulation-aid compositions, which overcome disadvantages exhibited by the prior art. The present invention provides
25 agrochemical formulation-aid compositions which permit on-site formulation of an agrochemical, a mixture of agrochemicals, or pre-formulated products prior to use. It thus becomes unnecessary to first formulate the active ingredient to render it bioactive and sprayable. It also significantly ameliorates any storage problems since the present invention makes it possible to formulate the agrochemical mixture as needed and just prior
30 to its use. In most applications, the only equipment required to use the agrochemical formulation aid of the present invention is a means to measure quantities reasonably

accurately, and an adequately powerful stirrer. Thus, both the equipment and energy requirements for formulation are significantly reduced, thereby reducing environmental and toxic byproducts in comparison with the known manufacturing processes used to produce both pre-formulated products, and the materials used in them in addition to the agrochemicals. The agrochemical formulation aid of the present invention is a free flowing material that disperses completely and rapidly in water. The agrochemical formulation aids of the present invention also improve the uptake and performance of pre-formulated agrochemicals, such improvement has not been provided hitherto by a range of other adjuvants. Technical acids of herbicides tend have lower vapor pressure when solubilized in a carrier. The formulation aid of the present invention is non-toxic and odorless unlike the solvents used in formulated products.

SUMMARY OF THE INVENTION

It is an object of the present invention to provide an agrochemical formulation aid composition for formulating an agrochemical, a mixture of agrochemicals, or pre-formulated products prior to use in a bioactive and sprayable form. The agrochemical formulation aid may be used on-site prior to use.

It is another object of the present invention to provide an agrochemical formulation aid composition for formulating non-formulated pesticides and pre-formulated pesticides.

It is yet another object of the present invention to provide a formulation aid that is a free flowing material that disperses completely and rapidly in water.

It is yet another object of the present invention to provide a formulation aid that improves the bioactivity of even pre-formulated agrochemicals

It is yet another object of the present invention to provide a formulation aid that is easy to store and requires simple equipment for measuring quantities reasonably accurately, a mixing-container and a stirrer.

It is yet another object of the present invention to provide formulation aid that has reduced energy requirements and reduced environmental and toxic byproducts.

It is yet another object of the present invention to provide a formulation aid composition that can be used for different crop applications, different stages of crop growth, different weather or climate conditions, and target species.

It is yet another object of the present invention to provide a formulation aid composition that is applicable at lower vapor pressure and has no odor problems.

It is yet another object of the present invention to provide a means to further improve the bioactivity of pesticides by adding a fertilizer to the formulation aid.

5 Accordingly, there is provided an agrochemical formulation aid composition for preparing bioactive and sprayable agrochemicals, wherein various components for the composition were selected from mineral oil paraffinic distillate and/or aromatic hydrocarbon distillate; 2N-octanol; oleyl-cetyl alcohol; polyoxyethylene (2) oleylether; polyoxyethylene (8) nonylphenolethin and/or ethoxylated tallow amine blend; lauryl
10 sulphate; fatty alcohol alkoxylate; terpenes, diammonium phosphate; tetrasodium ethylene diamine tetracetate; cab-o-sil; fatty acid methyl ester; (C18) free fatty acid blend; N-butanol; and methyl alcohol. Also provided are methods of preparing the formulation aid composition on site by mixing various components and methods of preparing sprayable and bioactive agrochemical system using the formulation aid and non-formulated and/or
15 formulated agrochemicals. Also provided are uses of the formulation aid in preparing sprayable and bioactive agrochemical systems for controlling pests.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

The present invention relates to agrochemical formulation aid compositions, their
20 uses, and methods of preparations. The invention also relates to methods of preparing agrochemical systems of non-formulated and pre-formulated agrochemicals using the formulation aid compositions to obtain improved tank-spray capability and bioactivity.

In an embodiment, the formulation aid composition comprises effective amounts of adjuvants, dispersants, emulsifiers, penetrants, surfactants, distillates, water conditioners,
25 and fertilizers for improved bioactivity and sprayability.

In a preferred embodiment the adjuvants, dispersants, emulsifiers, penetrants, surfactants, distillates, water conditioners, and fertilizers for improved bioactivity and spray applicability are selected from mineral oil paraffinic distillate; 2N-octanol; oleyl-cetyl alcohol; polyoxyethylene (2) oleylether; polyoxyethylene (8) nonylphenolethin; sodium lauryl sulphate; fatty alcohol alkoxylate; terpenes, preferably from a plant source;
30 diammonium phosphate; tetrasodium ethylene diamine tetracetate; and cab-o-sil. The

diammonium phosphate may be replaced with an alternative nitrogen-containing nutrient such as ammonia, ammonium nitrate, or ammonium sulphate.

More preferably, the formulation aid comprises from about 135 to about 165 parts by weight mineral oil paraffinic distillate; from about 23 to about 29 parts by weight 2N-octanol; from about 42 to about 52 parts by weight oleyl-cetyl alcohol; from about 46 to about 56 parts by weight polyoxyethylene (2) oleylether; from about 9 to about 11 parts by weight polyoxyethylene (8) nonylphenoethin; from about 1.2 to about 1.4 parts by weight sodium lauryl sulphate; from about 36 to about 43 parts by weight fatty alcohol alkoxylate; from about 7.5 to about 9.5 parts by weight terpenes, preferably from a plant source; from about 2.2 to about 2.8% w/v tetrasodium ethylene diamine tetracetate; and from about 8 to about 22 parts by weight of cab-o-sil for improved bioactivity and sprayability. From about 610 to about 670 parts by weight of a mixture containing from about 0.8 to about 1.2 parts by weight of diammonium phosphate in 2.7 to 3.3 parts by weight water may also be added. The diammonium phosphate may be replaced with an alternative nitrogen-containing nutrient such as ammonia, ammonium nitrate, or ammonium sulphate. The amount of polyoxyethylene (8) nonylphenoethin may be selected from about 3 to about 60 parts by weight. In an alternate embodiment, the formulation aid composition may further comprise of from about 5 to about 15 parts by weight of methyl alcohol.

Particularly preferably, the formulation aid comprises from about 143 to about 158 parts by weight mineral oil paraffinic distillate; from about 21.5 to about 27.5 parts by weight 2N-octanol; from about 44 to about 49 parts by weight oleyl-cetyl alcohol; from about 48 to about 53 parts by weight polyoxyethylene (2) oleylether; from about 9.5 to about 10.5 parts by weight polyoxyethylene (8) nonylphenoethin; from about 1.2 to about 1.4 parts by weight sodium lauryl sulphate; from about 38 to about 42 parts by weight fatty alcohol alkoxylate; from about 8 to about 9 parts by weight terpenes, preferably from a plant source; from about 2.35 to about 2.65 % w/v tetrasodium ethylene diamine tetracetate; from about 9 to about 21 parts by weight of cab-o-sil for improved bioactivity and sprayability. From about 636 to about 704 parts by weight of a mixture containing from about 0.9 to about 1.1 parts by weight of diammonium phosphate in 2.85 to 3.15 parts by weight water may be added further. The diammonium phosphate may be replaced with an alternative nitrogen-containing nutrient such as ammonia, ammonium nitrate, or

ammonium sulphate.

In a specially preferred embodiment, the formulation aid comprises about 150 parts by weight mineral oil paraffinic distillate; about 26 parts by weight 2N-octanol; about 47.2 parts by weight oleyl-cetyl alcohol; about 50.2 parts by weight polyoxyethylene (2) oleylether; about 10 parts by weight polyoxyethylene (8) nonylphenolethin; 1.0 parts by weight sodium lauryl sulphate; about 39.4 parts by weight fatty alcohol alkoxylate; about 8.6 parts by weight terpenes, preferably from a plant source; about 2.5% w/v tetrasodium ethylene diamine tetracetate; and from about 10 to about 20 parts by weight of cab-o-sil for improved bioactivity and sprayability. About 667 parts by weight of a mixture containing about 1 part by weight of diammonium phosphate in 3 parts by weight water may be added further. The diammonium phosphate may be replaced with an alternative nitrogen-containing nutrient such as ammonia, ammonium nitrate, or ammonium sulphate.

The above described formulation aids of the present invention were prepared by a three-part process comprising a step A followed by steps B and C, wherein step A comprises the following sub steps: placing from about 135 to about 165 parts by weight mineral oil paraffinic distillate in a suitable container with stirring means; adding from about 23 to about 29 parts by weight 2N-octanol and mixing; adding from about 42 to about 52 parts by weight oleyl-cetyl alcohol to the mixture and continuing mixing for up to about 8 minutes at a temperature of about 25°C until a homogenous mixture is obtained; adding from about 1.2 to about 1.4 parts by weight sodium lauryl sulphate and mixing it slowly until it dissolved; adding from about 46 to about 56 parts by weight polyoxyethylene (2) oleylether slowly and mixing it until dissolved; adding from about 9 to about 11 parts by weight polyoxyethylene (8) nonylphenolethin slowly and mixing it until dissolved; adding from about 36 to about 43 parts by weight fatty alcohol alkoxylate and mixing until it dissolved; adding from about 7.5 to about 9.5 parts by weight terpenes, preferably from a plant source, and mixing until the resulting mixture is homogenous. The range for polyoxyethylene (8) nonylphenolethin may be selected from about 3 to about 60 parts by weight for preparing the formulation aid.

The preparation of the formulation aid may additionally comprise of adding from about 5 to about 15 parts by weight of methyl alcohol to the mixture and mixing.

Step B comprises adding from about 0.8 to about 1.2 parts by weight of diammonium phosphate to 2.7 to 3.3 parts by weight water at a temperature of about 60°C and mixing until it dissolved; adding from about 2.2 to about 2.8% w/v tetrasodium
5 ethylene diamine tetracetate to the mixture; and adding from about 8 to about 22 parts by weight of cab-o-sil and mixing until it dispersed;

Step C comprises combining from about 610 to about 670 parts by weight of the product of step B to the product of step A and continuing mixing until a stable uniform homogenous blend is obtained.

10 In another embodiment the adjuvants, dispersants, emulsifiers, penetrants, surfactants, distillates, water conditioners, and fertilizers for improved bioactivity and sprayability may be selected from aromatic hydrocarbon distillate, 2N-octanol, oleyl-cetyl alcohol, polyoxyethylene (8) nonylphenoethin, polyoxyalkylated fatty alcohol, and ethoxylated tallow amine blend.

15 Preferably, the formulation aid of present invention comprises from about 10 to 40% by weight aromatic hydrocarbon distillate, from about 3 to 6% by weight of 2N-octanol, from about 5 to 10% by weight oleyl-cetyl alcohol, from about 0 to about 4% by weight of polyoxyethylene (8) nonylphenoethin, from about 25 to 35% by weight of polyoxyalkylated fatty alcohol, and from about 20% to 35% by weight of ethoxylated
20 tallow amine blend.

In a preferred embodiment, the formulation aid comprises about 21.5% by weight aromatic hydrocarbon distillate, about 3.1% by weight 2N-octanol, about 5.7% by weight oleyl-cetyl alcohol, about 3.1% by weight polyoxyethylene (8) nonylphenoethin, about 33.3% by weight polyoxyalkylated fatty alcohol, and from about 33.3% by weight
25 ethoxylated tallow amine blend.

In a more preferred embodiment, the formulation aid comprises about 23.6% by weight aromatic hydrocarbon distillate, about 3.5% by weight 2N-octanol, about 6.3% by weight oleyl-cetyl alcohol, about 33.3% by weight polyoxyalkylated fatty alcohol, and from about 33.3% by weight ethoxylated tallow amine blend.

5 In an especially preferred embodiment, the formulation aid comprises about 35.4% by weight aromatic hydrocarbon distillate, about 5.2% by weight 2N-octanol, about 9.4% by weight oleyl-cetyl alcohol, about 25.0% by weight polyoxyalkylated fatty alcohol, and about 25.0% weight ethoxylated tallow amine blend.

The above described formulation aids of the present invention can be prepared by a
10 one-part process comprising the following steps: placing from about 10 to 40% by weight aromatic hydrocarbon distillate in a suitable container containing stirring means; adding from about 3 to 6% by weight of 2N-octanol to the mixture; adding from about 5 to 10% by weight oleyl-cetyl alcohol to the mixture and continuing mixing for up to 8 minutes at a temperature of about 25° until a homogenous mixture is obtained; adding from 0 to about
15 4% by weight of polyoxyethylene (8) nonylphenoethin slowly and mixing until it dissolved; adding from about 25 to 35% by weight of polyoxyalkylated fatty alcohol and mixing until it dissolved; and adding from about 20% to 35% by weight of ethoxylated tallow amine blend and agitating until the resulting mixture is homogenous.

In yet another embodiment the adjuvants, dispersants, emulsifiers, penetrants,
20 surfactants, distillates, water conditioners, and fertilizers for improved bioactivity and spray applicability are selected from mineral oil paraffinic distillate, aromatic hydrocarbon, surfactant blend, 2N-octanol, oleyl-cetyl alcohol, polyoxyalkylated fatty alcohol, and ethoxylated tallow amine blend.

In a preferred embodiment, the formulation aid of present invention contained
25 about 6 to 8% by weight of mineral oil paraffinic distillate, from about 10 to 15% by weight aromatic hydrocarbon distillate, from about 5 to 7% by weight of surfactant blend, from about 1 to 3% by weight of 2N-octanol, from about 2 to 5% by weight of oleyl-cetyl alcohol, from about 25 to 35% by weight of polyoxyalkylated fatty alcohol, and from about 25-35% by weight of ethoxylated tallow amine blend.

30 In an especially preferred embodiment, the formulation aid of present invention comprises about 6.7% by weight of mineral oil paraffinic distillate; about 14.9% by weight

of aromatic hydrocarbon distillate; about 5.7% by weight of surfactant blend; about 2.2% by weight of 2N-octanol; about 3.9% by weight of oleyl-cetyl alcohol; about 33.3% by weight of polyoxyalkylated fatty alcohol; and about 33.3% by weight of ethoxylated tallow amine blend for improved bioactivity and sprayability.

5 In yet another embodiment the adjuvants, dispersants, emulsifiers, penetrants, surfactants, distillates, water conditioners, and fertilizers for improved bioactivity and sprayability may be selected from fatty acid methyl ester, preferably from a vegetable source; (C18) free fatty acid blend; oleyl-cetyl alcohol; N-butanol; polyoxyalkylated fatty alcohol; and ethoxylated tallow amine blend.

10 In a preferred embodiment, the formulation aid of present invention contained from about 20-25% by weight of a fatty acid methyl ester, preferably from a vegetable source; from about 0.1-3% by weight of (C18) free fatty acid blend; from about 0.5 to 3% by weight of 2N-octanol, from about 1-6 by weight of oleyl-cetyl alcohol; from about 0.1 to 1% by weight of N-butanol; from about 25 to 35% by weight of polyoxyalkylated fatty
15 alcohol; and from about 25 to 35% by weight of ethoxylated tallow amine surfactant blend for improved bioactivity and sprayability.

In a especially preferred embodiment, the formulation aid of present invention comprises about 24.4% by weight of vegetable fatty acid methyl ester, about 2% by weight of (C18) free fatty acid blend; about 2% by weight of 2N-octanol, about 4% by weight of
20 oleyl-cetyl alcohol; about 1% by weight of N-butanol; about 33.3% by weight of polyoxyalkylated fatty alcohol; and about 33.3% by weight of ethoxylated tallow amine surfactant blend for improved bioactivity and sprayability.

The above described formulation aids of the present invention can be prepared by a one-part process comprising the following steps: placing from about 20-25% by weight of
25 fatty acid methyl ester, preferably from a vegetable source in a suitable container with stirring apparatus and mixing it; adding from about 0.1 – 3% by weight of (C18) free fatty acid blend to the mixture; adding from about 2% by weight of 2N-octanol and mixing; adding from about 1-6 by weight of oleyl-cetyl alcohol to the mixture; adding from about 0.1 to 1% by weight of N-butanol to the mixture; adding from about 25 to 35% by weight
30 of polyoxyalkylated fatty alcohol to the mixture and mixing until it dissolved; and adding

from about 25 to 35% by weight of ethoxylated tallow amine surfactant blend and mixing it until the resulting mixture is homogenous.

In yet another embodiment, the formulation aid of present invention comprises about 27 parts by weight water; about 2 parts by weight free fatty acid blend; about 2 parts
5 by weight 2N-octanol; about 4 parts by weight oleyl-cetyl alcohol, about 1 part by weight N-butanol; about 33.3 parts by weight polyoxyalkylated fatty alcohol; and about 33.3 parts by weight ethoxylated tallow amine surfactant blend.

The agrochemical formulation aids described above can be stored under reasonable conditions, such as a temperature of below about 35°C. Although some of the formulation
10 aids contain water, freezing does not damage the formulation aid. If minor separation of phases does occur in the formulation aid, the agrochemical formulation aid can be readily re-dispersed to become homogenous by mixing. Some formulation aids do not contain water and are fully stable with prolonged shelf life.

The formulation aids described above have been used to prepare sprayable and
15 bioactive mixtures with a number of non-formulated i.e., technical acids or formulated herbicides alone or in combination. These sprayable and bioactive mixtures were sprayed on a number of plant species under variety of conditions in a variety of locations to show their effectiveness. Percentage control of weeds or percentage injury to specific crops was recorded as a measure of their effectiveness.

20 The invention will be illustrated with reference to the following examples.

Example 1

Preparation of sprayable and bioactive agrochemical and formulation aid mixture

Prior to any preparation all spray-mixing and application equipment were cleaned
25 thoroughly by observing cleaning directions provided on the agrochemical labeling and accompanying materials provided by the maker. Half of the required water was placed in a spray tank. A required amount of the formulation aid was added in the spray tank for the required rate of glyphosate application as indicated in Table 1. A required amount of herbicide e.g., glyphosate technical acid active ingredient or product, as indicated in Table
30 1, was added to the spray tank. Mixing was continued and remainder of the water was

added to the spray tank. Mixing was continued for another 5-10 minutes. The ready to use herbicide and formulation aid mixture was applied by spraying at the required rate.

Table 1 shows the amounts of glyphosate technical acid (N-phosphonomethylglycine, 95-99.9%) active ingredient or product and formulation aid
5 required to prepare sprayable and bioactive glyphosate and formulation aid mixture.

Example 2

Control of broad-spectrum weeds by using formulation aid with non-formulated glyphosate technical acid

10 Table 2 shows control of broad-spectrum weeds at commercial levels by using formulation aid of the present invention with a non-formulated herbicide, glyphosate technical acid at various rates. The results shown in Table 2 were obtained by applying glyphosate technical acid (95%) and the formulation aid mixture to 10 fields infected with various annual and perennial weeds (indicated in Table 2). The experiments were
15 conducted between the period of September 1998 to August 2000.

Hundred percent weed control was obtained 21 days after treatment (DAT) at a spray volume of 100 l/ha at different rates. No weed control was observed when formulation aid was replaced with water. Additional supporting efficacy was independently collected on weeds such as giant foxtail, yellow foxtail, velvet leaf, smart
20 weeds, ragweed species, venice mallow, entire leaf morning glory from research trials (small plot and grower trials) in various countries. The results demonstrate the utility of the formulation aid of the present invention in preparing a sprayable and bioactive mixture of technical acid of a herbicide for controlling weeds.

The formulation aid tested in this example contained about 21.5% by weight
25 aromatic hydrocarbon distillate, about 3.1% by weight 2N-octanol, about 5.7% by weight oleyl-cetyl alcohol, about 3.1% by weight polyoxyethylene (8) nonylphenolethin, about 33.3% by weight polyoxyalkylated fatty alcohol, and about 33.3% by weight ethoxylated tallow amine blend.

*Example 3***Control of broad-spectrum weeds by using formulation aid with non-formulated dicamba technical acid**

5 Table 3 shows control of broad-spectrum weeds at commercial levels by using formulation aid of the present invention with another non-formulated herbicide, Dicamba technical acid at various rates. The results shown in Table 3 were obtained by applying the mixture of dicamba technical acid (87%) and the formulation aid to two fields infected with various annual and perennial weeds (indicated in Table 3). The experiments were
10 conducted between the period of September 1998 to August 2000.

Hundred percent weed control was obtained 21 days after treatment (DAT) at a spray volume of 200 l/ha at different rates. No weed control was observed when formulation aid was replaced with water. Similar results were obtained independently on weeds such as giant foxtail, yellow foxtail, Velvetleaf, Smartweeds, ragweed species,
15 venice mallow, entire leaf morning glory from research trials (small plot and grower trials) in various countries. The results demonstrate the utility of the formulation aid of the present invention in preparing sprayable and bioactive of dicamba technical acid in controlling weeds.

The formulation aid used in this example contained about 150 parts by weight
20 mineral oil paraffinic distillate; about 26 parts by weight 2N-octanol; about 47.2 parts by weight oleyl-cetyl alcohol; about 50.8 parts by weight polyoxyethylene (2) oleylether; about 10 parts by weight polyoxyethylene (8) nonylphenoletin; 1.3 parts by weight sodium lauryl sulphate; about 39.4 parts by weight fatty alcohol alkoxylate; about 8.6 parts by weight plant terpenes; about 667 parts by weight of a mixture containing about 1 part
25 by weight of diammonium phosphate in 3 parts by weight water about 2.5% w/v tetrasodium ethylene diamine tetracetate (EDTA); and from about 10 to about 20 parts by weight of cab-o-sil for improved bioactivity and sprayability.

Table 4 shows the amounts of dicamba technical acid (3, 6-dichloro-2-methoxybenzoic acid, 88-95%) and formulation aid required to prepare a sprayable and
30 bioactive mixture.

Example 4

Effective weed control with sprayable and bioactive mixture of formulation aid and glyphosate technical acid

Table 5 shows the effectiveness of using a mixture of glyphosate technical acid
5 95% or glyphosate 97.3% wet cake isopropyl amine (IPA) technical acid and the
formulation aid in comparison to Roundup Transorb, a pre-formulated herbicide
preparation, 27 days after treatment. The plots (2 x 10 meters) were treated with a
precision plot spray system and applied with 100 l/ha water at 220 kPa through a four
nozzle (50 cm spacing) hand-held (2 meter) boom using TeeJet flat fan 8002 nozzles. The
10 application was done in the summer of 1999. The results from 4 replicates were not
significantly different among different treatments after 27 days. All combinations
performed equally well on the four weed species and provided excellent weed control at
the rates used. The results indicate that a herbicide mixture prepared with the formulation
aid of the present invention performed equally well as compared to a pre-formulated
15 herbicide.

The agrochemical formulation aid tested in the above example with glyphosate
technical acid contained about 150 parts by weight mineral oil paraffinic distillate; from
about 26 parts by weight 2N-octanol; from about 48 parts by weight oleyl-cetyl alcohol;
about 50 parts by weight polyoxyethylene (2) oleylether; about 5 parts by weight
20 polyoxyethylene (8) nonylphenolethin; about 1.3 parts by weight sodium lauryl sulphate;
about 40 parts by weight fatty alcohol alkoxylate; about 8 parts by weight plant terpenes;
about 667 parts by weight of diammonium phosphate in three parts by weight water. In an
alternate embodiment, diammonium phosphate was replaced with an alternative nitrogen-
containing nutrient such as ammonium sulphate or ammonium nitrate.

25

Example 5

Control of pre-plant vegetation using a sprayable and bioactive mixture of glyphosate technical acid and formulation aid

Table 6 shows pre-plant vegetation control when glyphosate technical acid 87%
30 was applied with one of the formulation aids of the present invention in comparison to
Roundup Ultra, a pre-formulated glyphosate herbicide. The application was done in

Summer of 2000. Table 6 shows percent control 14 days after treatment (DAT).

Application was made with a research plot sprayer with 42 lbs pressure and 20 gallons of water through DG 11002 nozzles to plots measuring 6.67 ft x 20 ft.

As shown in Table 6, formulation aid and formulation aid with ammonium sulphate controlled all plant species tested with slight variations. The formulation aid and glyphosate technical acid 95% proved slightly less effective on Velvetleaf but addition of ammonium sulphate improved Velvetleaf control in comparison to Roundup Ultra. The formulation aid and Glyphosate technical acid 95% with or without ammonium sulphate proved superior for control of Smartweed in comparison to Roundup Ultra.

The agrochemical formulation aid tested in the above example with glyphosate technical acid contained about 150 parts by weight mineral oil paraffinic distillate; from about 26 parts by weight 2N-octanol; from about 48 parts by weight oleyl-cetyl alcohol; about 50 parts by weight polyoxyethylene (2) oleylether; about 50 parts by weight polyoxyethylene (8) nonylphenoethin; about 1.3 parts by weight sodium lauryl sulphate; about 40 parts by weight fatty alcohol alkoxylate; about 8 parts by weight plant terpenes; about 667 parts by weight of diammonium phosphate in three parts by weight water. In an alternate embodiment, diammonium phosphate was replaced with an alternative nitrogen-containing nutrient such as ammonium sulphate or ammonium nitrate.

Example 6

Control of winter wheat using a sprayable and bioactive mixture of formulation aid with glyphosate technical acid and ammonium sulphate in comparison to Roundup Transorb.

Table 7 shows the use of formulation aid with glyphosate technical acid 95% with or without ammonium sulphate in comparison to Roundup Transorb, a pre-formulated herbicide, for controlling winter wheat. Percent control 7, 14, 28 & 56 days after treatment (DAT) of winter wheat was recorded. Plots (2x10 meter) were applied in the summer of 2000 using a precision plot sprayer at 200 l/ha. Formulation aid and glyphosate technical acid 95% alone was less effective in controlling winter wheat than Roundup Transorb or formulation aid with ammonium sulphate at 7, 14, 28 DAT.

However, formulation aid and glyphosate technical acid 95% alone were equally effective

in controlling winter wheat than Roundup Transorb or formulation aid with ammonium sulphate at 56 DAT.

The agrochemical formulation aid tested in this example with glyphosate technical acid contained about 150 parts by weight mineral oil paraffinic distillate; about 26 parts by weight 2N-octanol; about 47.2 parts by weight oleyl-cetyl alcohol; about 50.2 parts by weight polyoxyethylene (2) oleylether; about 50 parts by weight polyoxyethylene (8) nonylphenoethin; about 1 part by weight sodium lauryl sulphate; about 39.4 parts by weight fatty alcohol alkoxylate; about 8.6 parts by weight plant terpenes; about 667 parts by weight of diammonium phosphate in three parts by weight water. In an alternate embodiment, diammonium phosphate was replaced with an alternative nitrogen-containing nutrient such as ammonium sulphate or ammonium nitrate.

Example 7

Post emergent broadleaf weed control using formulation aid with dicamba technical acid in comparison to Banvel II

Table 8 shows post emergent broadleaf weed control using the formulation aid with dicamba technical acid 87% in comparison to Banvel II, a pre-formulated dicamba herbicide. The mixture was applied to 5-leaf stage corn in 2x10 meter plots with a precision plot sprayer at 200 l/ha in the summer of 2000. Percent weed control 28 days after treatment (DAT) was recorded. Formulation aid and dicamba technical acid 87% were more effective than Banvel II at both treatment rates. In another experiment, the formulation aid was used in combination with Prosulfuron, another pre-formulated herbicide and dicamba technical acid. Percentage control was as good as when Prosulfuron and Banvel II were used with a commercial adjuvant, Agral 90.

The agrochemical formulation aid tested in this example contained about 150 parts by weight mineral oil paraffinic distillate; about 26 parts by weight 2N-octanol; about 47.2 parts by weight oleyl-cetyl alcohol; about 50.2 parts by weight polyoxyethylene (2) oleylether; about 50 parts by weight polyoxyethylene (8) nonylphenoethin; about 1 part by weight sodium lauryl sulphate; about 39.4 parts by weight fatty alcohol alkoxylate; about 8.6 parts by weight plant terpenes; about 667 parts by weight of diammonium

phosphate in three parts by weight water. In an alternate embodiment, diammonium phosphate was replaced with an alternative nitrogen-containing nutrient such as ammonium sulphate or ammonium nitrate.

5

Example 8

Broadleaf weed control in Turf grass using the formulation aid with 2,4-D technical acid or dicamba technical acid alone and in combination in comparison to commercial formulations

10 This example demonstrates the use of formulation aid of the present invention in rendering 2,4-D technical acid, dicamba technical acid, and a combination thereof into an agriculturally and horticulturally acceptable form. This preparation is bioactive and spray-applicable at lower vapor pressure and has no odor problems.

15 Table 9 shows broadleaf weed control in Turf grass using the formulation aid with 2,4-D 96% technical acid or dicamba 98% technical acid alone and in combination in comparison to commercial formulations of 2,4-D and dicamba i.e., 2,4-D amine and Banvel 11. Treatments were applied using a precision plot sprayer at 200 l/ha to 2x8 meter plots during the fall of 2000. Percentage of injury on crop and percentage control of weeds was recorded 30 days after treatment

20 The agrochemical formulation aid tested in this example contained about 150 parts by weight mineral oil paraffinic distillate; about 26 parts by weight 2N-octanol; about 47 parts by weight oleyl-cetyl alcohol; about 51 parts by weight polyoxyethylene(2)oleylether; about 10 parts by weight polyoxyethylene(8) nonylphenoletin; about 1 part by weight sodium lauryl sulphate; about 39 parts by weight fatty alcohol alkoxylate; about 8 parts by weight plant terpenes; about 667 parts
25 by weight of diammonium phosphate in three parts by weight water; about 2.5% weight by volume of tetrasodium ethylene diamine tetracetate; and from about 10 to 20 parts by weight of cab-o-sil. In an alternate embodiment, diammonium phosphate was replaced with an alternative nitrogen-containing nutrient such as ammonium sulphate or ammonium nitrate.

30

*Example 9***Fall control of Alfalfa using the formulation aid with glyphosate technical acid and 2,4-D technical acid or glyphosate technical acid and dicamba technical acid in comparison to pre-formulated products**

5 This example demonstrates the use of formulation aid of the present invention in rendering the combination of glyphosate technical acid and 2,4-D technical acid or glyphosate technical acid and dicamba technical acid into an agriculturally or horticulturally acceptable form that is bioactive and spray applicable.

10 Table 10 shows fall control of Alfalfa using the formulation aid with glyphosate technical acid 95% and 2,4-D technical acid 96% or glyphosate technical acid 95% and dicamba technical acid 98% in comparison to pre-formulated products. Plots (2x8 meter) were sprayed with a precision plot sprayer at 200 l/ha in the fall, 2000 by Agriculture Canada, Harrow, Ontario, Canada. In comparison to use of pre-formulated products, the use of technical acids with the formulation aid of the present invention caused less injury
15 to the crop.

 The agrochemical formulation aid tested in this example contained about 150 parts by weight mineral oil paraffinic distillate; about 26 parts by weight 2N-octanol; about 47 parts by weight oleyl-cetyl alcohol; about 50 parts by weight polyoxyethylene (2) oleylether; about 10 parts by weight polyoxyethylene (8) nonylphenoletin; about 1 part
20 by weight sodium lauryl sulphate; about 39 parts by weight fatty alcohol alkoxylate; about 8 parts by weight plant terpenes; about 5 parts by weight methyl alcohol; about 667 parts by weight of diammonium phosphate in three parts by weight water; and about 2 parts weight by volume of tetrasodium ethylene diamine tetracetate; and from about 10 to 20 parts by weight of cab-o-sil. In an alternate embodiment, diammonium phosphate was
25 replaced with an alternative nitrogen-containing nutrient such as ammonium sulphate or ammonium nitrate.

*Example 10**Effectiveness of the formulation aid with glyphosate technical acid or pre-formulated agrochemical pesticides under controlled greenhouse conditions*

This example demonstrates the use of formulation aid of the present invention in rendering glyphosate technical acid or pre-formulated agrochemical pesticides into an agriculturally or horticulturally acceptable form that is bioactive and spray applicable.

Table 11, 12, 13 and 14 show effectiveness of the formulation aid with glyphosate technical acid or pre-formulated agrochemical pesticides in controlled greenhouse conditions during summer of 2001 in controlling several weeds such as Velvetleaf, Barnyard grass, and Lambsquarters. Herbicides, technical acid with the formulation aid or formulated products, were applied at 0.25 lb ae/h. The formulation aid was used at 0.3% v/v. AMS was used at 1% or 8.5 lb/100 gallon of water. The mixture was sprayed at 20 Gallons Per Acre at 30 Pressure Square Inch. The results indicate that the formulation aid works with formulated products also. The formulated products may contain different adjuvants. The results also indicate that the addition of AMS improved effectiveness of herbicides in controlling weeds.

The agrochemical formulation aid tested in this example contained about 6.7% by weight of mineral oil paraffinic distillate; about 14.9% by weight of aromatic hydrocarbon distillate; about 5.7% by weight of ATPLUS 300 F; about 2.2% by weight of 2N-octanol; about 3.9% by weight of oleyl-cetyl alcohol; about 33.3% by weight of polyoxyalkylated fatty alcohol; and about 33.3% by weight of ethoxylated tallow amine blend.

The commercial names and the suppliers of the components used in formulation aids are:

mineral oil paraffinic distillate: Sun Cropspray 11N, Sunoco Inc., Ten Penn Center, 1801 Market Street, Philadelphia, PA 19103;

aromatic hydrocarbon distillate: SOLVESSO 200, Imperial Oil, Products and Chemicals Division, 111 St. Clair Avenue West PO Box 4029 Stn A, Toronto, Ontario, Canada;

surfactant blend: ATPLUS 300 F, Uniqema, 1000 Uniqema Boulevard, New Castle, Delaware, USA;

2N-octanol: Jarchem Industries Inc., 414 Wilson Avenue, Newark, New Jersey,

USA;

oleyl-cetyl alcohol: HD Ocenol 80/85, Cognis Corporation, 5051 Estecreek Drive, Cincinnati, Ohio, USA;

polyoxyalkylated fatty alcohol: KLEARFAC AA270 surfactant, BASF

5 Corporation 3000 Continental Drive North, Mount Olive, New Jersey, USA; and

ethoxylated tallow amine blend: HENKEL 6821 A, Agnique GPU Booster 6821A, Cognis Corporation, 5051 Estecreek Drive, Cincinnati, Ohio, USA.

polyoxyethylene (2) oleylether: Brij 93, , ICI Americas Inc., Wilmington, Delaware

10 polyoxyethylene (8) nonylphenolethin::Renex 688 , ICI Americas Inc, Wilmington, Delaware, USA;

Fatty alcohol alkoxylate: Plurafac LF 700, BASF Canada Inc., Toronto, Ontario Canada;

plant terpenes: Orange Terpenes, Gerard-Roure Inc., Brampton, Ontario, Canada.

15 Other components are routinely available.

The above-described embodiments of the invention are intended to be examples of the present invention. Alterations, modifications and variations may be effected to the particular embodiments by those of skill in the art, without departing from the scope of the invention which is defined solely by the claims appended hereto.

20

Table 1. Amounts of glyphosate technical acid (N-phosphonomethylglycine, 95-99.9%) active ingredient or product and formulation aid required to prepare a sprayable and bioactive mixture.

Component A		Component B	
Glyphosate technical acid (95%)(Kg ae*)/hectare	Glyphosate technical grade (95%) (Kg product)/hectare	Formulation aid per 100 to 150 liters/ha water volume	Formulation aid per 50 liters/ha water volume

0.225	0.24	0.5	0.5
0.500	0.53	0.5	1.1
0.675	0.7	0.9	1.6
0.9	0.94	1.1	2.3
1.5	1.6	1.9	2.3
2.0	2.1	2.5	2.3

*Active ingredient

Table 2. Control of broad-spectrum weeds by using formulation aid of the present invention with non-formulated herbicide, glyphosate technical acid, at various application rates.

Glyphosate + formulation aid*	Canada Thistle	Lambs- quarters	Quack- grass**	Annual grasses ***	Dande -lions	Sow Thistle	Pig Weed	Catnip
0.94 + 0.9	100%	100%	100%	100%	100%	100%	100%	100%
0.7 + 0.7	100%	100%	100%	100%	100%	100%	100%	100%
0.53 + 0.5	100%	100%	100%	100%	100%	100%	100%	100%
0.24 + 0.5	80%	80%	80%	90%	80%	90%	100%	80%
0.94 + water	0%	0%	0%	0%	0%	0%	0%	0%
0.53 + water	0%	0%	0%	0%	0%	0%	0%	0%
0.24 + water	0%	0%	0%	0%	0%	0%	0%	0%

*Glyphosate technical acid (95%) (Kg or l)/ ha + formulation aid (l)/ ha

10 ** Includes other perennial grasses such as wirestem muhly

*** Annual grasses included foxtails, annual bluegrass, crabgrass, witchgrass

Table 3. Control of broad-spectrum weeds by using formulation aid of the present invention with non-formulated herbicide, Dicamba technical acid, at various application rates.

Dicamba + formulation aid*	Canada Thistle	Lambs- quarters	Quack Grass **	Various annual grasses ***	Dande -lions	Sow Thistle	Pig Weed	Catni p
0.140 + 1.0	N/A	100%	N/A	N/A	N/A	N/A	100%	N/A
0.280 + 1.0	N/A	100%	N/A	N/A	N/A	N/A	100%	N/A
0.140 + water	N/A	0%	N/A	N/A	N/A	N/A	0%	N/A
0.280 + water	N/A	0%	N/A	N/A	N/A	N/A	0%	N/A

* Dicamba 95% (Kg) + formulation aid (L)/ha

** includes other perennial grasses such as wirestem muhly

*** various annual grasses including foxtails, annual bluegrass, crabgrass, witch grass

5 N/A: Not applicable

Table 4. Amounts of Dicamba technical acid (3, 6-dichloro-2-methoxybenzoic acid, 88-95%) and formulation aid required to prepare a sprayable and bioactive mixture for spraying.

10

Component A	Component B
Dicamba technical acid (88-95%) (Kg)/ hectare	Formulation aid (l)/water volume
0.140	0.5% v/v
0.280	0.5% v/v

Table 5. Control of weeds using a mixture of glyphosate technical acid or glyphosate wet cake isopropyl amine technical acid and the formulation aid and Roundup Transorb.

Treatments	Rate	Redroot Pigweed	Lambs quarters	Green Foxtail	Smooth Crabgrass
		% Control			
Glyphosate 95% ta * + Formulation aid	225 gm ae**/ha 1.5% v/v	81.3%	63.8%	83.8%	67.5%
Glyphosate 97%wc***+ Formulation aid	225 gm ae/ha 1.5% v/v	83.8%	78.8%	83.8%	61.3%
Roundup Transorb	225 gm ae/ha	68.5%	61.3%	81.6%	73.5%

* technical acid

** active ingredient

*** wet cake isopropyl amine technical acid.

5

Table 6. Control of pre-plant vegetation using a sprayable and bioactive mixture of glyphosate technical acid and the formulation aid.

Treatments	Rates	Giant Foxtail	Yellow oxtail	Velvet leaf	Smart weed
Roundup Ultra	450 gm ae*/ha	100	98	85	30
Glyphosate ta** + Formulation aid	450 gm ae/ha 450 ml/ha	100	90	82	65
Glyphosate ta + Formulation aid + AMS***	450 gm ae/ha 450 ml/ha 1%v/v	100	95	93	85

10 * Active ingredient

** Technical acid

*** Ammonium sulphate

Table 7. Control of winter wheat using a mixture of formulation aid with glyphosate technical acid with and without ammonium sulphate in comparison to Roundup Transorb, a pre-formulated herbicide.

Treatments	Rates	% Control			
		7 DAT	14 DAT	28 DAT	56 DAT
Roundup Transorb	450 gm ae*/ha	15 c	23 c	53 b	100 a
Glyphosate ta** + Formulation aid	450 gm ae/ha 450 ml/ha	4 de	6 e	20 d	100 a
Glyphosate ta+ Formulation aid + AMS***	450 gm ae/ha 450 ml/ha 1%v/v	20 b	29 b	83 a	100 a

5

Means followed by the same letter do not significantly differ (P= .05, LSD)

*Active ingredient

**Technical acid

**Ammonium sulphate

10

Table 8. Post emergent broadleaf weed control using the formulation aid with dicamba technical acid 87% in comparison to Banvel II, a pre-formulated dicamba herbicide.

Treatment	Rate	Velvet	Lambs	Redroot
		Leaf	Quarters	Pigweed
		% Control		
Banvel II	70 g ae*/ha	23 d	18 e	18 c
Dicamba tech** + Formulation aid	70 g ae/ha 0.5% v/v	36 bc	63 c	43 b
Banvel II	140 g ae/ha	35 cd	35 d	58 b
Dicamba tech +	140 g ae/ha			

Formulation aid	0.5% v/v	49 b	79 b	58 b
Prosulfuron +	10 g ae/ha			
Banvel II +	140 g ae/ha	98 a	93 a	97 a
Agral 90	0.2% v/v			
Prosulfuron +	10 g ae/ha			
Dicamba tech +	140 g ae/ha	99 a	96 a	98 a
Formulation aid	0.5% v/v			

Means followed by the same letter do not significantly differ (P= .05, LSD).

*active ingredient

** dicamba technical acid

5

Table 9. Control of broadleaf weeds in Turfgrass using the formulation aid with 2,4-D technical acid or dicamba technical acid alone or in combination in comparison to commercial formulations of 2,4-D or dicamba.

Treatment	Rate	% Crop Injury	Dandelion	Chickweed	White clover
			% Control		
2,4-D amine	0.7 k ae*/ha	0.0 c	56.7 bc	56.7 a	53.3 b
2,4-D 96% ta** + Formulation aid	0.7 k ae/ha 0.5% v/v	0.0 c	50.0 bc	63.7 a	70.0 ab
Banvel II	0.6 k ae/ha	0.0 c	71.7 abc	100.0 a	100.0 a
Dicamba 98% ta + Formulation aid	0.6 k ae/ha 1.0% v/v	15 a	88.3 a	100.0 a	100.0 a
2,4-D amine	1.4 k ae/ha	6.7 b	63.3 abc	83.3 a	100.0 a
2,4-D 96% ta + Formulation aid	1.4 k ae/ha 1.0% v/v	0.0 c	70.0 abc	90.0 a	83.3 ab
2,4-D amine +	0.7 k ae/ha	0.0 c	70.0 abc	100.0 a	70.0 ab

Banvel II	0.07 k ae/ha				
2,4-D 96% ta + dicamba 98% ta + Formulation aid	0.7 k ae/ha 0.07 k ae/ha 0.5% v/v	0.0 c	41.7 c	73.3 a	93.3 a
2,4-D amine + Banvel II	1.4 k ae/ha 0.14 k ae/ha	0.0 c	78.3 ab	100.0 a	100.0 a
2,4-D 96% ta + dicamba 98% ta + formulation aid	1.4 k ae/ha 0.14 k ae/ha 1.0% v/v	0.0 c	68.3 abc	73.3 a	100.0 a

Means followed by the same letter do not significantly differ (P= .05, Duncan's New MRT).

* Active ingredient

5

** Technical acid

Table 10. Control of Alfalfa using the formulation aid with glyphosate technical acid and 2,4-D technical acid or glyphosate technical acid and dicamba technical acid in comparison to pre-formulated products.

10

Treatments	Rates	% injury	% control
Roundup Transorb	0.5 kg ae*/ha		
2,4-D amine	0.5 kg ae/ha	71.3 ab	100
Glyphosate 95% ta**+	0.5 kg ae/ha		
2,4-D 96% ta + formulation aid	0.5 kg ae/ha 0.5% v/v	47.5 cd	100
Roundup Transorb + 2,4-D amine	1.0 kg ae/ha 1.0 kg ae/ha	77.5 a	100
Glyphosate 95% ta + 2,4-D 96% ta + formulation aid	1.0 kg ae/ha 1.0 kg ae/ha 1.0% v/v	75.0 ab	100

Roundup Transorb + Banvel II	0.5 kg ae/ha 0.5 kg ae/ha	73.8 ab	100
Glyphosate 95% ta + Dicamba 98% ta + Formulation aid	0.5 kg ae/ha 0.5 kg ae/ha 0.5% v/v	47.5 cd	100
Roundup Transorb + Banvel II	1.0 kg ae/ha 1.0 kg ae/ha	75.0 ab	100
Glyphosate 95% ta + Dicamba 98% ta + Formulation aid	1.0 kg ae/ha 1.0 kg ae/ha 1.0% v/v	77.5 a	100

Means followed by the same letter do not significantly differ (P= .05, Duncan's New MRT).

* Active ingredient

5 ** Technical acid

Table 11. Control of Velvetleaf using the formulation aid with glyphosate technical acid or formulated agrochemical pesticides in controlled greenhouse conditions.

TRT NO.	Treatment	% Control		
		7 DAT	10 DAT	14 DAT
1	Control	0 G	0 E	0 E
2	Glyphos*	13 g	10 e	23 d
3	Glyphos + formulation aid	12 g	10 e	15 d e
4	Glyphos + formulation aid + AMS	40 e f	43 d	70 a b c
5	Glyphomax**	52 b c d e	60 c d	67 a b c
6	Glyphomax + formulation aid	43 d e f	77 a b c	81 a b c
7	Glyphomax + formulation aid + AMS	38 e f	78 a b c	82 a b c
8	RoundUp Ultra max***	10 g	13 e	7 d e

9	RoundUp Ultra max + AMS	60 a b c	70 a b c	78 a b c
	LSD (0.05)	16	19	23

Means followed by the same letter do not significantly differ (P= .05, Duncan's New MRT).

* glyphosate amine (IPA salt)

5 ** glyphosate amine (IPA salt)

*** glyphosate amine (IPA salt)

Table 12. Control of barnyard grass using the formulation aid with glyphosate technical acid or formulated agrochemical pesticides in controlled greenhouse conditions.

10

TRT NO.	Treatment	% Control		
		7 DAT	10 DAT	14 DAT
1	Control	0 h	0 k	0 h
2	Glyphos*	38 e f	40 h i j	30 g
3	Glyphomax**	55 b c d	36 i j	40 e f
4	Glyphos + formulation aid	53 c d	48 f g h	48 e
5	Glyphomax + formulation aid	28 f g	45 g h i	34 f g
6	Glyphos + formulation aid + AMS	63 a b c	64 b c d	70 a b
7	Glyphomax + formulation aid + AMS	55 b c d	66 a b c	73 a b
8	Glyphosate ta* + formulation aid + AMS	71 a	74 a	69 b c
9	RoundUp Ultra Max***	18 g	33 j	0 h
10	RoundUp Ultra Max + AMS	66 a b	68 a b c	69 b c
	LSD (0.05)	13	10	10

Means followed by the same letter do not significantly differ (P= .05, Duncan's New MRT).

* glyphosate amine (IPA salt)

** glyphosate amine (IPA salt)

5 *** glyphosate amine (IPA salt)

Table 13. Control of Velvetleaf using the formulation aid with glyphosate technical acid or formulated agrochemical pesticides in controlled greenhouse conditions.

TRT	Treatment	% Control		
NO.		7 DAT	10 DAT	14 DAT
1	Control	0 i	0 j	0 e
2	Glyphos*	13 g h	13 i	13 d e
3	Glyphos + formulation aid	45 f	49 h	53 c
4	Glyphos + formulation aid + AMS	65 a b c d e	68 c d e f	65 a b c
5	Glyphomax**	23 g	15 i	15 d
6	Glyphomax + formulation aid	63 b c d e	50 g h	53 c
7	Glyphomax + formulation aid + AMS	70 a b c	71 a b c d e	70 a b
8	Glyphosate ta + formulation aid + AMS	60 c d e	63 e f	63 b c
9	RoundUp Ultra Max***	10 h i	14 i	18 d
10	RoundUp Ultra Max + AMS	60 c d e	66 d e f	68 a b
	LSD (0.05)	11	11	13

10

Means followed by the same letter do not significantly differ (P= .05, Duncan's New MRT).

* glyphosate amine (IPA salt)

** glyphosate amine (IPA salt)

15 *** glyphosate amine (IPA salt)

Table 14. Control of Lambsquarters using the formulation aid with glyphosate technical acid or formulated agrochemical pesticides in controlled greenhouse conditions.

TRT NO.	Treatment	% Control		
		7 DAT	10 DAT	14 DAT
1	Control	1 g h	0 j	0 i
2	Glyphos*	9 f g h	1 j	0 i
3	Glyphos + formulation aid	9 f g h	10 h i j	0 i
4	Glyphos + formulation aid + AMS	23 b c d	30 b c d e	18 e f g h
5	Glyphomax**	11 e f g	15 g h i	5 h i
6	Glyphomax + formulation aid	18 c d e f	25 c d e f g	15 f g h
7	Glyphomax + formulation aid + AMS	16 c d e f	20 e f g h	33 a b c d
8	Glyphosate ta + formulation aid + AMS	40 a	40 a b	44 a
9	Round Up Ultra Max***	3 g h	0 j	0 i
10	Round Up Ultra Max + AMS	33 a b	33 a b c d	36 a b c
	LSD (0.05)	11	11	15

5 Means followed by the same letter do not significantly differ ($P = .05$, Duncan's New MRT).

* glyphosate amine (IPA salt)

** glyphosate amine (IPA salt)

*** glyphosate amine (IPA salt)

CLAIMS:

1. An agrochemical formulation aid composition for use with technical acid, non-formulated, partially formulated or pre-formulated actives for in-tank preparation of a bioactive and sprayable agrochemical or mixture of agrochemicals comprising:
 - (a) 2N-octanol;
 - (b) oleyl-cetyl alcohol; and
 together with at least one of:
 - (c) fatty alcohol alkoxylate;
 - (d) polyoxyethylene (2) oleylether;
 - (e) aromatic hydrocarbon distillate;
 - (f) methylated seed oil;
 - (g) polyoxyethylene (8) nonylphenol;
 - (h) sodium lauryl sulphate;
 - (i) a nitrogen-containing nutrient selected from the group consisting of diammonium phosphate, ammonia, ammonium nitrate, diammonium sulphate;
 - (j) tetrasodium ethylene diamine tetraacetate;
 - (k) colloidal silica;
 - (l) a terpene;
 - (m) polyethoxylated amine;
 - (n) polyoxyalkylated alcohol;
 - (o) C18 free fatty acid blend;
 - (p) N-butanol;
 - (q) methanol;
 - (r) water.

2. The agrochemical formulation aid composition according to claim 1, wherein said nitrogen-containing nutrient comprises diammonium phosphate.

3. The agrochemical formulation aid composition according to claim 1, wherein said nitrogen-containing nutrient is selected from the group consisting of ammonia, ammonium nitrate, and ammonium sulphate.
4. The agrochemical formulation aid composition according to claim 1, comprising from about 135 to about 165 parts by weight mineral oil paraffinic distillate; from about 23 to about 29 parts by weight 2N-octanol; from about 42 to about 52 parts by weight oleyl-cetyl alcohol; from about 46 to about 56 parts by weight polyoxyethylene (2) oleylether; from about 9 to about 11 parts by weight polyoxyethylene (8) nonylphenol; from about 1.2 to about 1.4 parts by weight sodium lauryl sulphate; from about 36 to about 43 parts by weight fatty alcohol alkoxylate; from about 7.5 to about 9.5 parts by weight terpenes; from about 2.2 to about 2.8% w/v tetrasodium ethylene diamine tetraacetate; from about 8 to about 22 parts by weight of colloidal silica; and from about 610 to about 670 parts by weight of a mixture containing from about 0.8 to about 1.2 parts by weight of diammonium phosphate in 2.7 to 3.3 parts by weight water.
5. The agrochemical formulation aid composition according to claim 1, comprising from about 143 to about 158 parts by weight mineral oil paraffinic distillate; from about 21.5 to about 27.5 parts by weight 2N-octanol; from about 44 to about 49 parts by weight oleyl-cetyl alcohol; from about 48 to about 53 parts by weight polyoxyethylene (2) oleylether; from about 9.5 to about 10.5 parts by weight polyoxyethylene (8) nonylphenol; from about 1.2 to about 1.4 parts by weight sodium lauryl sulphate; from about 38 to about 42 parts by weight fatty alcohol alkoxylate; from about 8 to about 9 parts by weight terpenes; from about 2.35 to about 2.65% w/v tetrasodium ethylene diamine tetraacetate; from about 9 to about 21 parts by weight of colloidal silica; and from about 636 to about 704 parts by weight of a mixture containing from about 0.9 to about 1.1 parts by weight of diammonium phosphate in 2.85 to 3.15 parts by weight water.
6. The agrochemical formulation aid composition according to claim 1, comprising about 150 parts by weight mineral oil paraffinic distillate; about 26 parts by weight 2N-octanol; about 47.2 parts by weight oleyl-cetyl alcohol; about 50.2 parts by

weight polyoxyethylene (2) oleylether; about 10 parts by weight polyoxyethylene (8) nonylphenol; about 1.0 parts by weight sodium lauryl sulphate; about 39.4 parts by weight fatty alcohol alkoxylate; about 8.6 parts by weight terpenes; about 2.5% w/v tetrasodium ethylene diamine tetraacetate; from about 10 to about 20 parts by weight of colloidal silica; and about 667 parts by weight of a mixture containing about 1 part by weight of diammonium phosphate in 3 parts by weight water.

7. The agrochemical formulation aid composition according to claim 1, comprising from about 10 to about 40% by weight aromatic hydrocarbon distillate, from about 3 to about 6% by weight of 2N-octanol, from about 5 to about 10% by weight oleyl-cetyl alcohol, from about 0 to about 4% by weight of polyoxyethylene (8) nonylphenol, from about 25 to about 35% by weight of polyoxyalkylated fatty alcohol, and from about 20% to about 35% by weight of ethoxylated tallow amine blend.
8. The agrochemical formulation aid composition according to claim 1, comprising about 21.5% by weight aromatic hydrocarbon distillate, about 3.1% by weight 2N-octanol, about 5.7% by weight oleyl-cetyl alcohol, about 3.1% by weight polyoxyethylene (8) nonylphenol, about 33.3% by weight polyoxyalkylated fatty alcohol, and about 33.3% by weight ethoxylated tallow amine blend.
9. The agrochemical formulation aid composition according to claim 1, comprising about 23.6% by weight aromatic hydrocarbon distillate, about 3.5% by weight 2N-octanol, about 6.3% by weight oleyl-cetyl alcohol, about 33.3% by weight polyoxyalkylated fatty alcohol, and about 33.3% by weight ethoxylated tallow amine blend.
10. The agrochemical formulation aid composition according to claim 1, comprising about 35.4% by weight aromatic hydrocarbon distillate, about 5.2% by weight 2N-octanol, about 9.4% by weight oleyl-cetyl alcohol, about 25.0% by weight polyoxyalkylated fatty alcohol, and about 25.0% weight ethoxylated tallow amine blend.

11. The agrochemical formulation aid composition according to claim 1, comprising about 6 to about 8% by weight of mineral oil paraffinic distillate, from about 10 to about 15% by weight aromatic hydrocarbon distillate, from about 5 to about 7% by weight of surfactant blend, from about 1 to about 3% by weight of 2N-octanol, from about 25 to about 35% by weight of oleyl-cetyl alcohol, from about 25 to about 35% by weight of polyoxyalkylated fatty alcohol, and from about 25 to about 35% by weight of ethoxylated tallow amine blend.
12. The agrochemical formulation aid composition according to claim 1, comprising about 6.7% by weight mineral oil paraffinic distillate, about 14.9% by weight aromatic hydrocarbon distillate, about 5.7% by weight surfactant blend, about 2.2% by weight 2N-octanol, about 3.9% by weight oleyl-cetyl alcohol, about 33.3% by weight polyoxyalkylated fatty alcohol, and about 33.3% by weight ethoxylated tallow amine blend.
13. The agrochemical formulation aid composition according to claim 1, comprising from about 20 to 25% by weight fatty acid methyl ester; from about 0.1 to 3% by weight C18 free fatty acid blend; from about 0.5-3.0% by weight 2N-octanol, from about 1-6% by weight oleyl-cetyl alcohol; from about 0.1 to 1% by weight N-butanol; from about 25 to 35% by weight polyoxyalkylated fatty alcohol; and from about 25 to 35% by weight ethoxylated tallow amine blend.
14. The agrochemical formulation aid composition according to claim 1, comprising about 24.4% by weight vegetable fatty acid methyl ester, about 2% by weight of C18 free fatty acid blend; about 0.5-3.0% by weight of 2N-octanol, about 4% by weight oleyl-cetyl alcohol; about 1% by weight N-butanol; about 33.3% by weight polyoxyalkylated fatty alcohol; and about 33.3% by weight ethoxylated tallow amine blend.
15. The agricultural formulation aid composition according to claim 1 in substantially aqueous form for control of weeds.

16. The agricultural formulation aid composition according to claim 1 additionally comprising a fertiliser.
17. A method of controlling weed pests which comprises applying the agrochemical formulation aid composition as defined in claim 1 in conjunction with a herbicide in a bioactive and sprayable form.