

(19) **DANMARK**

(10) **DK/EP 2035564 T3**



(12) **Oversættelse af
europæisk patentskrift**

Patent- og
Varemærkestyrelsen

-
- (51) Int.Cl.: **C 12 N 15/864 (2006.01)** **C 07 K 14/015 (2006.01)** **C 12 N 5/10 (2006.01)**
- (45) Oversættelsen bekendtgjort den: **2016-07-18**
- (80) Dato for Den Europæiske Patentmyndigheds bekendtgørelse om meddelelse af patentet: **2016-04-20**
- (86) Europæisk ansøgning nr.: **07747521.8**
- (86) Europæisk indleveringsdag: **2007-06-20**
- (87) Den europæiske ansøgnings publiceringsdag: **2009-03-18**
- (86) International ansøgning nr.: **NL2007050298**
- (87) Internationalt publikationsnr.: **WO2007148971**
- (30) Prioritet: **2006-06-21 US 815262 P** **2006-06-21 EP 06115804**
- (84) Designerede stater: **AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HU IE IS IT LI LT LU LV MC MT NL PL PT RO SE SI SK TR**
- (73) Patenthaver: **UniQure IP B.V., Meibergdreef 61, 1105 BA Amsterdam Zuidoost, Holland**
- (72) Opfinder: **HAAST, Saskia Jacoba Petronella, Graaf Floris 2F, 1276 XB Huizen, Holland**
BIESMANS, Dennis Johan, Reigersweide 132, 1383 KJ Weesp, Holland
HERMENS, Wilhelmus Theodorus Johannes, Maria Christiaan, Sprietzeil 83, 1319 EB Almere, Holland
BAKKER, Andrew Christian, Leguaanstraat 23, 1338 HB Almere, Holland
- (74) Fuldmægtig i Danmark: **Zacco Denmark A/S, Arne Jacobsens Allé 15, 2300 København S, Danmark**
- (54) Benævnelse: **VEKTORER MED MODIFICERET INITIATIONSKODON TIL TRANSLATION AF AAV-REP78 TIL FREMSTILLING AF AAV I INSEKTCELLER**
- (56) Fremdragne publikationer:
WO-A-99/61641
WO-A-2005/072364
US-A1- 2003 148 506
URABE MASASHI ET AL: "Insect cells as a factory to produce adeno-associated virus type 2 vectors" HUMAN GENE THERAPY, vol. 13, no. 16, 1 November 2002 (2002-11-01), pages 1935-1943, XP002394454 ISSN: 1043-0342 cited in the application
KOHLBRENNER ET AL: "Successful Production of Pseudotyped rAAV Vectors Using a Modified Baculovirus Expression System" MOLECULAR THERAPY, ACADEMIC PRESS, SAN DIEGO, CA,, US, vol. 12, no. 6, December 2005 (2005-12), pages 1217-1225, XP005176630 ISSN: 1525-0016 cited in the application
CHEJANOVSKY N CARTER B J: "MUTAGENESIS OF AUG CODON IN THE ADENO-ASSOCIATED VIRUS REP GENE: EFFECTS ON VIRAL DNA REPLICATION" VIROLOGY, vol. 173, no. 1, November 1989 (1989-11), pages 120-128, XP000561298 ISSN: 0042-6822

DESCRIPTION

Field of the invention

[0001] The present invention relates to the production of adeno-associated virus in insect cells and to adeno-associated virus with improvements in expression and stability of the viral rep proteins that increase the productivity of adeno-associated viral vectors in insect cells.

Background of the invention

[0002] Adeno-associated virus (AAV) may be considered as one of the most promising viral vectors for human gene therapy. AAV has the ability to efficiently infect dividing as well as non-dividing human cells, the AAV viral genome integrates into a single chromosomal site in the host cell's genome, and most importantly, even though AAV is present in many humans it has never been associated with any disease. In view of these advantages, recombinant adeno-associated virus (rAAV) is being evaluated in gene therapy clinical trials for hemophilia B, malignant melanoma, cystic fibrosis, and other diseases.

[0003] Host cells that sustain AAV replication *in vitro* are all derived from mammalian cell types. Therefore, rAAV for use in gene therapy has thus far mainly been produced on mammalian cell lines such as e.g. 293 cells, COS cells, HeLa cells, KB cells, and other mammalian cell lines (see e.g. US 6,156,303, US 5,387,484, US 5,741,683, US 5,691,176, US 5,688,676, US 20020081721, WO 00/47757, WO 00/24916, and WO 96/17947). rAAV vectors are typically produced in such mammalian cell culture systems by providing DNA plasmids that contain the therapeutic gene flanked by the origin of AAV replication (inverted terminal repeats or ITRs), genes for AAV replication proteins Rep78, Rep68, Rep52, and Rep40, and genes for virion or structural proteins VP1, VP2, and VP3. In addition, a plasmid containing early genes from adenovirus (E2A, E4ORF6, VARNA) is provided to enhance the expression of the AAV genes and improve vector yield (see e.g. Grimm et al., 1998, Hum. Gene Ther. 9: 2745-2760). However, in most of these mammalian cell culture systems, the number of AAV particles generated per cell is on the order of 10^4 particles (reviewed in Clark, 2002, Kidney Int. 61(Suppl. 1): 9-15). For a clinical study, more than 10^{15} particles of rAAV may be required. To produce this number of rAAV particles, transfection and culture with approximately 10^{11} cultured human 293 cells, the equivalent of 5,000 175-cm² flasks of cells, would be required., which means transfecting up to 10^{11} 293 cells. Therefore, large scale production of rAAV using mammalian cell culture systems to obtain material for clinical trials has already proven to be problematic, production at commercial scale may not even be feasible. Furthermore there is always the risk, that a vector for clinical use that is produced in a mammalian cell culture will be contaminated with undesirable, perhaps pathogenic, material present in the mammalian host cell.

[0004] To overcome these problems of mammalian productions systems, recently, an AAV production system has been developed using insect cells (Urabe et al., 2002, Hum. Gene Ther. 13: 1935-1943; US 20030148506 and US 20040197895). For production of AAV in insect cells some modifications were necessary in order to achieve the correct stoichiometry of the three AAV capsid proteins (VP1, VP2 and VP3), which relies on a combination of alternate usage of two splice acceptor sites and the suboptimal utilization of an ACG initiation codon for VP2 that is not accurately reproduced by insect cells. To mimic the correct stoichiometry of the capsid proteins in insect cells Urabe et al. (2002, *supra*) use a construct that is transcribed into a single polycistronic messenger that is able to express all three VP proteins without requiring splicing and wherein the most upstream initiator codon is replaced by the suboptimal initiator codon ACG. In co-pending application (PCT/NL2005/050018) the present inventors have further improved the infectivity of baculovirus-produced rAAV vectors based production by further optimisation of the stoichiometry of AAV capsid proteins in insect cells.

[0005] For expression of the AAV Rep proteins in the AAV insect cell expression system as initially developed by Urabe et al. (2002, *supra*), a recombinant baculovirus construct is used that harbours two independent Rep expression units (one for Rep78 and one for Rep52), each under the control of a separate insect cell promoter, the Δ IE1 and PolH promoters, respectively. In this system, the Δ IE1 promoter, a much weaker promoter than the PolH promoter, was chosen for driving Rep78 expression since it is known that in mammalian cells a less abundant expression of Rep78 as compared to Rep52 favours high vector yields (Li et al., 1997, J Virol. 71: 5236-43; Grimm et al., 1998, *supra*).

[0006] More recently however, Kohlbrenner et al. (2005, Mol. Ther. 12: 1217-25) reported that the baculovirus construct for expression of the two Rep protein, as used by Urabe et al., suffers from an inherent instability. By splitting the palindromic orientation of the two Rep genes in Urabe's original vector and designing two separate baculovirus vectors for expressing Rep52

and Rep78, Kohlbrenner et al. (2005, *supra*) increased the passaging stability of the vector. However, despite the consistent expression of Rep78 and Rep52 from the two independent baculovirus-Rep constructs in insect cells over at least 5 passages, rAAV vector yield is 5 to 10-fold lower as compared to the original baculovirus-Rep construct designed by Urabe et al. (2002, *supra*).

[0007] There is thus still a need to overcome the above serious limitations of large scale (commercial) production of AAV vectors in insect cells. Thus it is an object of the present invention to provide for means and methods that allow for stable and high yield (large scale) production of AAV vectors in insect cells.

Description of the invention

Definitions

[0008] As used herein, the term "operably linked" refers to a linkage of polynucleotide (or polypeptide) elements in a functional relationship. A nucleic acid is "operably linked" when it is placed into a functional relationship with another nucleic acid sequence. For instance, a transcription regulatory sequence is operably linked to a coding sequence if it affects the transcription of the coding sequence. Operably linked means that the DNA sequences being linked are typically contiguous and, where necessary to join two protein encoding regions, contiguous and in reading frame.

[0009] "Expression control sequence" refers to a nucleic acid sequence that regulates the expression of a nucleotide sequence to which it is operably linked. An expression control sequence is "operably linked" to a nucleotide sequence when the expression control sequence controls and regulates the transcription and/or the translation of the nucleotide sequence. Thus, an expression control sequence can include promoters, enhancers, internal ribosome entry sites (IRES), transcription terminators, a start codon in front of a protein-encoding gene, splicing signal for introns, and stop codons. The term "expression control sequence" is intended to include, at a minimum, a sequence whose presence are designed to influence expression, and can also include additional advantageous components. For example, leader sequences and fusion partner sequences are expression control sequences. The term can also include the design of the nucleic acid sequence such that undesirable, potential initiation codons in and out of frame, are removed from the sequence. It can also include the design of the nucleic acid sequence such that undesirable potential splice sites are removed. It includes sequences or polyadenylation sequences (pA) which direct the addition of a polyA tail, i.e., a string of adenine residues at the 3'-end of a mRNA, sequences referred to as polyA sequences. It also can be designed to enhance mRNA stability. Expression control sequences which affect the transcription and translation stability, e.g., promoters, as well as sequences which effect the translation, e.g., Kozak sequences, are known in insect cells. Expression control sequences can be of such nature as to modulate the nucleotide sequence to which it is operably linked such that lower expression levels or higher expression levels are achieved.

[0010] As used herein, the term "promoter" or "transcription regulatory sequence" refers to a nucleic acid fragment that functions to control the transcription of one or more coding sequences, and is located upstream with respect to the direction of transcription of the transcription initiation site of the coding sequence, and is structurally identified by the presence of a binding site for DNA-dependent RNA polymerase, transcription initiation sites and any other DNA sequences, including, but not limited to transcription factor binding sites, repressor and activator protein binding sites, and any other sequences of nucleotides known to one of skill in the art to act directly or indirectly to regulate the amount of transcription from the promoter. A "constitutive" promoter is a promoter that is active in most tissues under most physiological and developmental conditions. An "inducible" promoter is a promoter that is physiologically or developmentally regulated, e.g. by the application of a chemical inducer. A "tissue specific" promoter is only active in specific types of tissues or cells.

[0011] The terms "substantially identical", "substantial identity" or "essentially similar" or "essential similarity" means that two peptide or two nucleotide sequences, when optimally aligned, such as by the programs GAP or BESTFIT using default parameters, share at least a certain percentage of sequence identity as defined elsewhere herein. GAP uses the Needleman and Wunsch global alignment algorithm to align two sequences over their entire length, maximizing the number of matches and minimizes the number of gaps. Generally, the GAP default parameters are used, with a gap creation penalty = 50 (nucleotides) / 8 (proteins) and gap extension penalty = 3 (nucleotides) / 2 (proteins). For nucleotides the default scoring matrix used is nws gapdna and for proteins the default scoring matrix is Blosum62 (Henikoff & Henikoff, 1992, PNAS 89, 915-919). It is clear than when RNA sequences are said to be essentially similar or have a certain degree of sequence identity with DNA sequences, thymine (T) in the DNA sequence is considered equal to uracil (U) in the RNA sequence. Sequence alignments and scores for percentage sequence identity may be determined using computer programs, such as the GCG Wisconsin Package, Version 10.3,

available from Accelrys Inc., 9685 Scranton Road, San Diego, CA 92121-3752 USA or the open-source software Emboss for Windows (current version 2.7.1-07). Alternatively percent similarity or identity may be determined by searching against databases such as FASTA, BLAST, etc.

[0012] Nucleotide sequences encoding parvoviral Rep proteins of the invention may also be defined by their capability to hybridise with the nucleotide sequence of SEQ ID NO.10, respectively, under moderate, or preferably under stringent hybridisation conditions. Stringent hybridisation conditions are herein defined as conditions that allow a nucleic acid sequence of at least about 25, preferably about 50 nucleotides, 75 or 100 and most preferably of about 200 or more nucleotides, to hybridise at a temperature of about 65°C in a solution comprising about 1 M salt, preferably 6 x SSC or any other solution having a comparable ionic strength, and washing at 65°C in a solution comprising about 0.1 M salt, or less, preferably 0.2 x SSC or any other solution having a comparable ionic strength. Preferably, the hybridisation is performed overnight, i.e. at least for 10 hours and preferably washing is performed for at least one hour with at least two changes of the washing solution. These conditions will usually allow the specific hybridisation of sequences having about 90% or more sequence identity.

[0013] Moderate conditions are herein defined as conditions that allow a nucleic acid sequences of at least 50 nucleotides, preferably of about 200 or more nucleotides, to hybridise at a temperature of about 45°C in a solution comprising about 1 M salt, preferably 6 x SSC or any other solution having a comparable ionic strength, and washing at room temperature in a solution comprising about 1 M salt, preferably 6 x SSC or any other solution having a comparable ionic strength. Preferably, the hybridisation is performed overnight, i.e. at least for 10 hours, and preferably washing is performed for at least one hour with at least two changes of the washing solution. These conditions will usually allow the specific hybridisation of sequences having up to 50% sequence identity. The person skilled in the art will be able to modify these hybridisation conditions in order to specifically identify sequences varying in identity between 50% and 90%.

Detailed description of the invention

[0014] The present invention relates the use of animal parvoviruses, in particular dependoviruses such as infectious human or simian AAV, and the components thereof (e.g., an animal parvovirus genome) for use as vectors for introduction and/or expression of nucleic acids in mammalian cells. In particular, the invention relates to improvements in productivity of such parvoviral vectors when produced in insect cells.

[0015] Viruses of the Parvoviridae family are small DNA animal viruses. The family Parvoviridae may be divided between two subfamilies: the Parvovirinae, which infect vertebrates, and the Densovirinae, which infect insects. Members of the subfamily Parvovirinae are herein referred to as the parvoviruses and include the genus Dependovirus. As may be deduced from the name of their genus, members of the Dependovirus are unique in that they usually require coinfection with a helper virus such as adenovirus or herpes virus for productive infection in cell culture. The genus Dependovirus includes AAV, which normally infects humans (e.g., serotypes 1, 2, 3A, 3B, 4, 5, and 6) or primates (e.g., serotypes 1 and 4), and related viruses that infect other warm-blooded animals (e.g., bovine, canine, equine, and ovine adeno-associated viruses). Further information on parvoviruses and other members of the Parvoviridae is described in Kenneth I. Berns, "Parvoviridae: The Viruses and Their Replication," Chapter 69 in Fields Virology (3d Ed. 1996). For convenience the present invention is further exemplified and described herein by reference to AAV. It is however understood that the invention is not limited to AAV but may equally be applied to other parvoviruses.

[0016] The genomic organization of all known AAV serotypes is very similar. The genome of AAV is a linear, single-stranded DNA molecule that is less than about 5,000 nucleotides (nt) in length. Inverted terminal repeats (ITRs) flank the unique coding nucleotide sequences for the non-structural replication (Rep) proteins and the structural (VP) proteins. The VP proteins (VP1, -2 and -3) form the capsid. The terminal 145 nt are self-complementary and are organized so that an energetically stable intramolecular duplex forming a T-shaped hairpin may be formed. These hairpin structures function as an origin for viral DNA replication, serving as primers for the cellular DNA polymerase complex. Following wtAAV infection in mammalian cells the Rep genes (i.e. Rep78 and Rep52) are expressed from the P5 promoter and the P19 promoter, respectively and both Rep proteins have a function in the replication of the viral genome. A splicing event in the Rep ORF results in the expression of actually four Rep proteins (i.e. Rep78, Rep68, Rep52 and Rep40). However, it has been shown that the unspliced mRNA, encoding Rep78 and Rep52 proteins, in mammalian cells are sufficient for AAV vector production. Also in insect cells the Rep78 and Rep52 proteins suffice for AAV vector production.

[0017] A "recombinant parvoviral or AAV vector" (or "rAAV vector") herein refers to a vector comprising one or more polynucleotide sequences of interest, genes of interest or "transgenes" that are flanked by parvoviral or AAV inverted terminal

repeat sequences (ITRs). Such rAAV vectors can be replicated and packaged into infectious viral particles when present in an insect host cell that is expressing AAV rep and cap gene products (i.e. AAV Rep and Cap proteins). When an rAAV vector is incorporated into a larger nucleic acid construct (e.g. in a chromosome or in another vector such as a plasmid or baculovirus used for cloning or transfection), then the rAAV vector is typically referred to as a "pro-vector" which can be "rescued" by replication and encapsidation in the presence of AAV packaging functions and necessary helper functions.

[0018] In a first aspect the invention relates to a nucleotide sequence comprising an open reading frame comprising nucleotide sequences encoding animal parvoviruses Rep proteins, wherein the initiation codon for translation of the parvoviral Rep78 protein is a suboptimal initiation codon. The suboptimal initiation codon preferably is an initiation codon that effects partial exon skipping. Partial exon skipping is herein understood to mean that at least part of the ribosomes do not initiate translation at the suboptimal initiation codon of the Rep78 protein but at an initiation codon further downstream, whereby preferably the initiation codon further downstream is the initiation codon of the Rep52 protein. The suboptimal initiation codon preferably effects partial exon skipping upon expression of the nucleotide sequence in an insect cell. Preferably, the suboptimal initiation codon effects partial exon skipping in an insect cell so as to produce in the insect cell a molar ratio of Rep78 to Rep52 in the range of 1:10 to 10:1, 1:5 to 5:1, or 1:3 to 3:1, preferably at about 20 - 40 hours post infection, more preferably at about 30 - 40 hours post infection, using a baculovirus expression. The molar ratio of the Rep78 and Rep52 may be determined by means of Western blotting as described in Example 1.1.3, preferably using a monoclonal antibody that recognizes a common epitope of both Rep78 and Rep52, or using the antibody described in Example 1.1.3.

[0019] The term "suboptimal initiation codon" herein not only refers to the tri-nucleotide initiation codon itself but also to its context. Thus, a suboptimal initiation codon may consist of an "optimal" ATG codon in a suboptimal context, e.g. a non-Kozak context. However, more preferred are suboptimal initiation codons wherein the tri-nucleotide initiation codon itself is suboptimal, i.e. is not ATG. Suboptimal is herein understood to mean that the codon is less efficient in the initiation of translation in an otherwise identical context as compared to the normal ATG codon. Preferably, the efficiency of suboptimal codon is less than 90, 80, 60, 40 or 20% of the efficiency of the normal ATG codon in an otherwise identical context. Methods for comparing the relative efficiency of initiation of translation are known per se to the skilled person. Preferred suboptimal initiation codons may be selected from ACG, TTG, CTG, and GTG. More preferred is ACG.

[0020] A nucleotide sequence encoding animal parvoviruses Rep proteins, is herein understood as a nucleotide sequence encoding the non-structural Rep proteins that are required and sufficient for parvoviral vector production in insect cells such the Rep78 and Rep52 proteins. The animal parvovirus nucleotide sequence preferably is from a dependovirus, more preferably from a human or simian adeno-associated virus (AAV) and most preferably from an AAV which normally infects humans (e.g., serotypes 1, 2, 3A, 3B, 4, 5, and 6) or primates (e.g., serotypes 1 and 4). An example of a nucleotide sequence encoding animal parvoviruses Rep proteins is given in SEQ ID No. 10, which depicts a part of the AAV serotype-2 sequence genome encoding the Rep proteins. The Rep78 coding sequence comprises nucleotides 11 - 1876 and the Rep52 coding sequence comprises nucleotides 683 - 1876. It is understood that the exact molecular weights of the Rep78 and Rep52 proteins, as well as the exact positions of the translation initiation codons may differ between different parvoviruses. However, the skilled person will know how to identify the corresponding position in nucleotide sequence from other parvoviruses than AAV-2. A nucleotide sequence encoding animal parvoviruses Rep proteins may thus also be defined as a nucleotide sequence:

1. a) that encodes a polypeptide comprising an amino acid sequence that has at least 50, 60, 70, 80, 88, 89, 90, 95, 97, 98, or 99% sequence identity with the amino acid sequence of SEQ ID NO. 11;
2. b) that has at least 50, 60, 70, 80, 81, 82, 85, 90, 95, 97, 98, or 99% sequence identity with the nucleotide sequence of positions 11 - 1876 of SEQ ID NO. 10;
3. c) the complementary strand of which hybridises to a nucleic acid molecule sequence of (a) or (b);
4. d) nucleotide sequences the sequence of which differs from the sequence of a nucleic acid molecule of (c) due to the degeneracy of the genetic code.

Preferably, the nucleotide sequence encodes animal parvoviruses Rep proteins that are required and sufficient for parvoviral vector production in insect cells.

[0021] A further preferred nucleotide sequence of the invention comprises an expression control sequence that comprising a nine nucleotide sequence of SEQ. ID NO: 7 or a nucleotide sequence substantially homologous to SEQ. ID NO: 7, upstream of the initiation codon of the nucleotide sequence encoding the parvoviral Rep78 protein. A sequence with substantial identity to the nucleotide sequence of SEQ. ID NO: 7 and that will help increase expression of the parvoviral Rep78 protein is e.g. a sequence which has at least 60%, 70%, 80% or 90% identity to the nine nucleotide sequence of SEQ ID NO: 7.

[0022] Elimination of possible false translation initiation sites in the Rep protein coding sequences, other than the Rep78 and

Rep52 translation initiation sites, of other parvoviruses will be well understood by an artisan of skill in the art, as will be the elimination of putative splice sites that may be recognised in insect cells. The various modifications of the wild-type parvoviral sequences for proper expression in insect cells is achieved by application of well-known genetic engineering techniques such as described e.g. in Sambrook and Russell (2001) "Molecular Cloning: A Laboratory Manual (3rd edition), Cold Spring Harbor Laboratory, Cold Spring Harbor Laboratory Press, New York. Various further modifications of Rep protein coding regions are known to the skilled artisan which could increase yield of Rep protein. These modifications are within the scope of the present invention.

[0023] In a further aspect the invention relates to a nucleic acid construct comprising a nucleotide sequence encoding parvoviral Rep proteins as defined above. Preferably, in the construct, the nucleotide sequence encoding the parvoviral Rep proteins is operably linked to expression control sequences for expression in an insect cell. These expression control sequences will at least include a promoter that is active in insect cells. Techniques known to one skilled in the art for expressing foreign genes in insect host cells can be used to practice the invention. Methodology for molecular engineering and expression of polypeptides in insect cells is described, for example, in Summers and Smith. 1986. A Manual of Methods for Baculovirus Vectors and Insect Culture Procedures, Texas Agricultural Experimental Station Bull. No. 7555, College Station, Tex .; Luckow. 1991. In Prokop et al., Cloning and Expression of Heterologous Genes in Insect Cells with Baculovirus Vectors' Recombinant DNA Technology and Applications, 97-152; King, L. A. and R. D. Possee, 1992, The baculovirus expression system, Chapman and Hall, United Kingdom; O'Reilly, D. R., L. K. Miller, V. A. Luckow, 1992, Baculovirus Expression Vectors: A Laboratory Manual, New York ; W. H. Freeman and Richardson, C. D., 1995, Baculovirus Expression Protocols, Methods in Molecular Biology, volume 39; US 4,745,051; US2003148506; and WO 03/074714. A particularly suitable promoter for transcription of the nucleotide sequence of the invention encoding of the parvoviral Rep proteins is e.g. the polyhedron promoter. However, other promoters that are active in insect cells are known in the art, e.g. the p10, p35, IE-1 or Δ IE-1 promoters and further promoters described in the above references.

[0024] Preferably the nucleic acid construct for expression of the parvoviral Rep proteins in insect cells is an insect cell-compatible vector. An "insect cell-compatible vector" or "vector" is understood to a nucleic acid molecule capable of productive transformation or transfection of an insect or insect cell. Exemplary biological vectors include plasmids, linear nucleic acid molecules, and recombinant viruses. Any vector can be employed as long as it is insect cell-compatible. The vector may integrate into the insect cells genome but the presence of the vector in the insect cell need not be permanent and transient episomal vectors are also included. The vectors can be introduced by any means known, for example by chemical treatment of the cells, electroporation, or infection. In a preferred embodiment, the vector is a baculovirus, a viral vector, or a plasmid. In a more preferred embodiment, the vector is a baculovirus, i.e. the construct is a baculoviral vector. Baculoviral vectors and methods for their use are described in the above cited references on molecular engineering of insect cells.

[0025] In another aspect the invention relates to an insect cell that comprises no more than one type of nucleotide sequence comprising a single open reading frame encoding a parvoviral Rep protein. Preferably the single open reading frame encodes one or more of the parvoviral Rep proteins, more preferably the open reading frame encodes all of the parvoviral Rep proteins, most preferably the open reading frame encodes the full-length Rep 78 protein from which preferably at least both Rep 52 and Rep 78 proteins may be expressed in the insect cell. It is understood herein that the insect cell may comprise more than one copy of the single type of nucleotide sequence, e.g. in a multicopy episomal vector, but that these are multiple copies of essentially one and the same nucleic acid molecule, or at least nucleic acid molecules that encode one and the same Rep amino acid sequence, e.g. nucleic acid molecules that only differ between each other due to the degeneracy of the genetic code. The presence of only a single type of nucleic acid molecule encoding the parvoviral Rep proteins avoids recombination between homologous sequences as may be present in different types of vectors comprising Rep sequences, which may give rise to defective Rep expression constructs that affect (stability of) parvoviral production levels in insect cells. Preferably, in the insect cell, the nucleotide sequence comprising the single open reading frame encoding one or more parvoviral Rep proteins is part of a nucleic acid construct wherein the nucleotide sequence is operably linked to expression control sequences for expression in an insect cell. A further preferred insect cell comprises as a "first" nucleotide sequence a nucleotide sequence as defined above encoding parvoviral Rep proteins, preferably a coding sequence with a suboptimal initiation codon as defined above, or a nucleic acid construct as defined above or the insect cell comprises as a "first" nucleic acid construct a nucleic acid construct as defined above comprising such nucleotide sequences.

[0026] Any insect cell which allows for replication of a recombinant parvoviral (rAAV) vector and which can be maintained in culture can be used in accordance with the present invention. For example, the cell line used can be from *Spodoptera frugiperda*, *drosophila* cell lines, or mosquito cell lines, e.g., *Aedes albopictus* derived cell lines. Preferred insect cells or cell lines are cells from the insect species which are susceptible to baculovirus infection, including e.g. Se301, SeIZD2109, SeUCR1, Sf9, Sf900+, Sf21, BTI-TN-5B1-4, MG-1, Tn368, HzAm1, Ha2302, Hz2E5, High Five (Invitrogen, CA, USA) and *expresSF+*[®] (US 6,103,526; Protein Sciences Corp., CT, USA).

[0027] A preferred insect cell according to the invention, in addition to the above described "first" nucleotide sequence or a nucleic acid construct, further comprises:

1. a) a second nucleotide sequence comprising at least one parvoviral inverted terminal repeat (ITR) nucleotide sequence; and,
2. b) a third nucleotide sequence comprising parvoviral Cap protein coding sequences operably linked to expression control sequences for expression in an insect cell.

[0028] In the context of the invention "at least one parvoviral ITR nucleotide sequence" is understood to mean a palindromic sequence, comprising mostly complementary, symmetrically arranged sequences also referred to as "A," "B," and "C" regions. The ITR functions as an origin of replication, a site having a "cis" role in replication, i.e., being a recognition site for trans acting replication proteins such as e.g. Rep 78 (or Rep68) which recognize the palindrome and specific sequences internal to the palindrome. One exception to the symmetry of the ITR sequence is the "D" region of the ITR. It is unique (not having a complement within one ITR). Nicking of single-stranded DNA occurs at the junction between the A and D regions. It is the region where new DNA synthesis initiates. The D region normally sits to one side of the palindrome and provides directionality to the nucleic acid replication step. A parvovirus replicating in a mammalian cell typically has two ITR sequences. It is, however, possible to engineer an ITR so that binding sites are on both strands of the A regions and D regions are located symmetrically, one on each side of the palindrome. On a double-stranded circular DNA template (e.g., a plasmid), the Rep78- or Rep68-assisted nucleic acid replication then proceeds in both directions and a single ITR suffices for parvoviral replication of a circular vector. Thus, one ITR nucleotide sequence can be used in the context of the present invention. Preferably, however, two or another even number of regular ITRs are used. Most preferably, two ITR sequences are used. A preferred parvoviral ITR is an AAV ITR. For safety reasons it may be desirable to construct a recombinant parvoviral (rAAV) vector that is unable to further propagate after initial introduction into a cell. Such a safety mechanism for limiting undesirable vector propagation in a recipient may be provided by using rAAV with a chimeric ITR as described in US2003148506.

[0029] The number of nucleic acid constructs employed in the insect cell for the production of the recombinant parvoviral (rAAV) vector is not limiting in the invention. For example, one, two, three, four, five, or more separate constructs can be employed to produce rAAV in insect cells in accordance with the methods of the present invention. If five constructs are employed, one construct encodes AAV VP 1, another construct encodes AAV VP2, yet another construct encodes AAV VP3, still yet another construct encodes the Rep protein as defined above and a final construct comprises at least one AAV ITR. If fewer than five constructs are used, the constructs can comprise various combinations of the at least one AAV ITR and the VP1, VP2, VP3, and the Rep protein coding sequences. Preferably, two constructs or three constructs are used, with two constructs being more preferred as described above. If two constructs are used, preferably the insect cell comprises: (a) a first nucleic acid construct for expression of the Rep proteins as defined above, which construct further comprises the third nucleotide sequences as defined in (b) above (comprising parvoviral Cap protein coding sequences operably linked to at least one expression control sequence for expression in an insect cell; see also below); and (c) a second nucleic acid construct comprising the second nucleotide sequence as defined in (a) above (comprising at least one parvoviral/AAV ITR nucleotide sequence). If three constructs are used, preferably the same configuration as used for two constructs is used except that separate constructs are used for expression of the capsid proteins and for expression of the Rep proteins. The sequences on each construct can be in any order relative to each other. For example, if one construct comprises ITRs and an ORF comprising nucleotide sequences encoding VP capsid proteins, the VP ORF can be located on the construct such that, upon replication of the DNA between ITR sequences, the VP ORF is replicated or not replicated. For another example, the Rep coding sequences and/or the ORF comprising nucleotide sequences encoding VP capsid proteins can be in any order on a construct. It is understood that also the second, third and further nucleic acid construct(s) preferably are an insect cell-compatible vectors, preferably a baculoviral vectors as described above. Alternatively, in the insect cell of the invention, one or more of the first nucleotide sequence, second nucleotide sequence, third nucleotide sequence, and fourth nucleotide sequence and optional further nucleotide sequences may be stably integrated in the genome of the insect cell. One of ordinary skill in the art knows how to stably introduce a nucleotide sequence into the insect genome and how to identify a cell having such a nucleotide sequence in the genome. The incorporation into the genome may be aided by, for example, the use of a vector comprising nucleotide sequences highly homologous to regions of the insect genome. The use of specific sequences, such as transposons, is another way to introduce a nucleotide sequence into a genome.

[0030] In the invention, the third nucleotide sequence comprising parvoviral capsid (Cap) protein coding sequences is herein understood to comprises sequences encoding each of the three parvoviral capsid proteins, VP1, -2 and -3. The third nucleotide sequence comprising the capsid protein coding sequences may be present in various forms. E.g. separate coding sequences for each of the capsid proteins VP1, -2 and -3 may be used, whereby each coding sequence is operably linked to expression control

sequences for expression in an insect cell. More preferably, however, the third nucleotide sequence comprises a single open reading frame encoding all three of the animal parvoviral (AAV) VP1, VP2, and VP3 capsid proteins, wherein the initiation codon for translation of the VP1 capsid protein is a suboptimal initiation codon that is not ATG as e.g. described by Urabe et al. (2002, *supra*). A suboptimal initiation codon for the VP1 capsid protein may be as defined above for the Rep78 protein. More preferred suboptimal initiation codons for the VP1 capsid protein may be selected from ACG, TTG, CTG and GTG, of which CTG and GTG are most preferred. A preferred third nucleotide sequence for the expression of the capsid proteins further comprises an expression control sequence comprising a nine nucleotide sequence of SEQ. ID NO: 7 or a nucleotide sequence substantially homologous to SEQ. ID NO: 7, upstream of the initiation codon of the nucleotide sequence encoding the VP1 capsid protein. A sequence with substantial identity to the nucleotide sequence of SEQ. ID NO: 7 and that will help increase expression of VP1 is e.g. a sequence which has at least 60%, 70%, 80% or 90% identity to the nine nucleotide sequence of SEQ ID NO: 7. A further preferred third nucleotide sequence for expression of the capsid proteins further preferably comprises at least one modification of the nucleotide sequence encoding the VP1 capsid protein selected from among a C at nucleotide position 12, an A at nucleotide position 21, and a C at nucleotide position 24 (with reference to position 1 being the first nucleotide of the translation initiation codon; see SEQ ID NO.1). Elimination of possible false initiation codons for translation of VP1 of other serotypes will be well understood by an artisan of skill in the art, as will be the elimination of putative splice sites that may be recognised in insect cells. Various further modifications of VP coding regions are known to the skilled artisan which could either increase yield of VP and virion or have other desired effects, such as altered tropism or reduce antigenicity of the virion. These modifications are within the scope of the present invention. Preferably the nucleotide sequence of the invention encoding the parvoviral capsid proteins is operably linked to expression control sequences for expression in an insect cell, which will at least include a promoter that is active in insect cells. Such control sequences and further techniques and materials (e.g. vectors) for expressing parvoviral capsid proteins in insect host cells are already described above for the Rep proteins.

[0031] In a preferred embodiment of the invention, the second nucleotide sequence present in the insect cells of the invention, i.e. the sequence comprising at least one parvoviral (AAV) ITR, further comprises at least one nucleotide sequence encoding a gene product of interest, whereby preferably the at least one nucleotide sequence encoding a gene product of interest becomes incorporated into the genome of a recombinant parvoviral (rAAV) vector produced in the insect cell. Preferably, at least one nucleotide sequence encoding a gene product of interest is a sequence for expression in a mammalian cell. Preferably, the second nucleotide sequence comprises two parvoviral (AAV) ITR nucleotide sequences and wherein the at least one nucleotide sequence encoding a gene product of interest is located between the two parvoviral (AAV) ITR nucleotide sequences. Preferably, the nucleotide sequence encoding a gene product of interest (for expression in the mammalian cell) will be incorporated into the recombinant parvoviral (rAAV) vector produced in the insect cell if it is located between two regular ITRs, or is located on either side of an ITR engineered with two D regions.

[0032] The second nucleotide sequence defined herein above may thus comprise a nucleotide sequence encoding at least one "gene product of interest" for expression in a mammalian cell, located such that it will be incorporated into an recombinant parvoviral (rAAV) vector replicated in the insect cell. Any nucleotide sequence can be incorporated for later expression in a mammalian cell transfected with the recombinant parvoviral (rAAV) vector produced in accordance with the present invention. The nucleotide sequence may e.g. encode a protein it may express an RNAi agent, i.e. an RNA molecule that is capable of RNA interference such as e.g. a shRNA (short hairpinRNA) or an siRNA (short interfering RNA). "siRNA" means a small interfering RNA that is a short-length double-stranded RNA that are not toxic in mammalian cells (Elbashir et al., 2001, Nature 411: 494-98; Caplen et al., 2001, Proc. Natl. Acad. Sci. USA 98: 9742-47). In a preferred embodiment, the second nucleotide sequence may comprise two nucleotide sequences and each encodes one gene product of interest for expression in a mammalian cell. Each of the two nucleotide sequences encoding a product of interest is located such that it will be incorporated into a recombinant parvoviral (rAAV) vector replicated in the insect cell.

[0033] The product of interest for expression in a mammalian cell may be a therapeutic gene product. A therapeutic gene product can be a polypeptide, or an RNA molecule (siRNA), or other gene product that, when expressed in a target cell, provides a desired therapeutic effect such as e.g. ablation of an undesired activity, e.g. the ablation of an infected cell, or the complementation of a genetic defect, e.g. causing a deficiency in an enzymatic activity. Examples of therapeutic polypeptide gene products include CFTR, Factor IX, Lipoprotein lipase (LPL, preferably LPL S447X; see WO 01/00220), Apolipoprotein A1, Uridine Diphosphate Glucuronosyltransferase (UGT), Retinitis Pigmentosa GTPase Regulator Interacting Protein (RP-GRIP), and cytokines or interleukins like e.g. IL-10.

[0034] Alternatively, or in addition as a second gene product, second nucleotide sequence defined herein above may comprise a nucleotide sequence encoding a polypeptide that serve as marker proteins to assess cell transformation and expression. Suitable marker proteins for this purpose are e.g. the fluorescent protein GFP, and the selectable marker genes HSV thymidine kinase (for selection on HAT medium), bacterial hygromycin B phosphotransferase (for selection on hygromycin B), Tn5 aminoglycoside phosphotransferase (for selection on G418), and dihydrofolate reductase (DHFR) (for selection on methotrexate), CD20, the low

affinity nerve growth factor gene. Sources for obtaining these marker genes and methods for their use are provided in Sambrook and Russel (2001) "Molecular Cloning: A Laboratory Manual (3rd edition), Cold Spring Harbor Laboratory, Cold Spring Harbor Laboratory Press, New York. Furthermore, second nucleotide sequence defined herein above may comprise a nucleotide sequence encoding a polypeptide that may serve as a fail-safe mechanism that allows to cure a subject from cells transduced with the recombinant parvoviral (rAAV) vector of the invention, if deemed necessary. Such a nucleotide sequence, often referred to as a suicide gene, encodes a protein that is capable of converting a prodrug into a toxic substance that is capable of killing the transgenic cells in which the protein is expressed. Suitable examples of such suicide genes include e.g. the *E.coli* cytosine deaminase gene or one of the thymidine kinase genes from Herpes Simplex Virus, Cytomegalovirus and Varicella-Zoster virus, in which case ganciclovir may be used as prodrug to kill the transgenic cells in the subject (see e.g. Clair et al., 1987, Antimicrob. Agents Chemother. 31: 844-849).

[0035] In another embodiment one of the gene products of interest can be an AAV protein. In particular, a Rep protein, such as Rep78 or Rep68, or a functional fragment thereof. A nucleotide sequence encoding a Rep78 and/or a Rep68, if present on the genome of a recombinant parvoviral (rAAV) vector of the invention and expressed in a mammalian cell transduced with the vector, allows for integration of the recombinant parvoviral (rAAV) vector into the genome of the transduced mammalian cell. Expression of Rep78 and/or Rep68 in an rAAV-transduced or infected mammalian cell can provide an advantage for certain uses of the recombinant parvoviral (rAAV) vector, by allowing long term or permanent expression of any other gene product of interest introduced in the cell by the vector.

[0036] In the recombinant parvoviral (rAAV) vectors of the invention the at least one nucleotide sequence(s) encoding a gene product of interest for expression in a mammalian cell, preferably is/are operably linked to at least one mammalian cell-compatible expression control sequence, e.g., a promoter. Many such promoters are known in the art (see Sambrook and Russel, 2001, *supra*). Constitutive promoters that are broadly expressed in many cell-types, such as the CMV promoter may be used. However, more preferred will be promoters that are inducible, tissue-specific, cell-type-specific, or cell cycle-specific. For example, for liver-specific expression a promoter may be selected from an α 1-anti-trypsin promoter, a thyroid hormone-binding globulin promoter, an albumin promoter, LPS (thyroxine-binding globulin) promoter, HCR-ApoCII hybrid promoter, HCR-hAAT hybrid promoter and an apolipoprotein E promoter. Other examples include the E2F promoter for tumor-selective, and, in particular, neurological cell tumor-selective expression (Parr et al., 1997, Nat. Med. 3:1145-9) or the IL-2 promoter for use in mononuclear blood cells (Hagenbaugh et al., 1997, J Exp Med; 185: 2101-10).

[0037] AAV is able to infect a number of mammalian cells. See, e.g., Tratschin et al. (1985, Mol. Cell Biol. 5:3251-3260) and Grimm et al. (1999, Hum. Gene Ther. 10:2445-2450). However, AAV transduction of human synovial fibroblasts is significantly more efficient than in similar murine cells, Jennings et al., Arthritis Res, 3:1 (2001), and the cellular tropicity of AAV differs among serotypes. See, e.g., Davidson et al. (2000, Proc. Natl. Acad. Sci. USA, 97:3428-3432), who discuss differences among AAV2, AAV4, and AAV5 with respect to mammalian CNS cell tropism and transduction efficiency.

[0038] AAV sequences that may be used in the present invention for the production of recombinant AAV vectors in insect cells can be derived from the genome of any AAV serotype. Generally, the AAV serotypes have genomic sequences of significant homology at the amino acid and the nucleic acid levels, provide an identical set of genetic functions, produce virions which are essentially physically and functionally equivalent, and replicate and assemble by practically identical mechanisms. For the genomic sequence of the various AAV serotypes and an overview of the genomic similarities see e.g. GenBank Accession number U89790; GenBank Accession number J01901; GenBank Accession number AF043303; GenBank Accession number AF085716; Chlorini et al. (1997, J. Vir. 71: 6823-33); Srivastava et al. (1983, J. Vir. 45:555-64); Chlorini et al. (1999, J. Vir. 73:1309-1319); Rutledge et al. (1998, J. Vir. 72:309-319); and Wu et al. (2000, J. Vir. 74: 8635-47). AAV serotypes 1, 2, 3, 4 and 5 are preferred source of AAV nucleotide sequences for use in the context of the present invention. Preferably the AAV ITR sequences for use in the context of the present invention are derived from AAV1, AAV2, and/or AAV4. Likewise, the Rep (Rep78 and Rep52) coding sequences are preferably derived from AAV1, AAV2, and/or AAV4. The sequences coding for the VP1, VP2, and VP3 capsid proteins for use in the context of the present invention may however be taken from any of the known 42 serotypes, more preferably from AAV1, AAV2, AAV3, AAV4, AAV5, AAV6, AAV7, AAV8 or AAV9 or newly developed AAV-like particles obtained by e.g. capsid shuffling techniques and AAV capsid libraries.

[0039] AAV Rep and ITR sequences are particularly conserved among most serotypes. The Rep78 proteins of various AAV serotypes are e.g. more than 89% identical and the total nucleotide sequence identity at the genome level between AAV2, AAV3A, AAV3B, and AAV6 is around 82% (Bantel-Schaal et al., 1999, J. Virol., 73(2):939-947). Moreover, the Rep sequences and ITRs of many AAV serotypes are known to efficiently cross-complement (i.e., functionally substitute) corresponding sequences from other serotypes in production of AAV particles in mammalian cells. US2003148506 reports that AAV Rep and ITR sequences also efficiently cross-complement other AAV Rep and ITR sequences in insect cells.

[0040] The AAV VP proteins are known to determine the cellular tropicity of the AAV virion. The VP protein-encoding sequences are significantly less conserved than Rep proteins and genes among different AAV serotypes. The ability of Rep and ITR sequences to cross-complement corresponding sequences of other serotypes allows for the production of pseudotyped rAAV particles comprising the capsid proteins of a serotype (e.g., AAV3) and the Rep and/or ITR sequences of another AAV serotype (e.g., AAV2). Such pseudotyped rAAV particles are a part of the present invention.

[0041] Modified "AAV" sequences also can be used in the context of the present invention, e.g. for the production of rAAV vectors in insect cells. Such modified sequences e.g. include sequences having at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, or more nucleotide and/or amino acid sequence identity (e.g., a sequence having about 75-99% nucleotide sequence identity) to an AAV1, AAV2, AAV3, AAV4, AAV5, AAV6, AAV7, AAV8 or AAV9 ITR, Rep, or VP can be used in place of wild-type AAV ITR, Rep, or VP sequences.

[0042] Although similar to other AAV serotypes in many respects, AAV5 differs from other human and simian AAV serotypes more than other known human and simian serotypes. In view thereof, the production of AAV5 can differ from production of other serotypes in insect cells. Where methods of the invention are employed to produce AAV5, it is preferred that one or more constructs comprising, collectively in the case of more than one construct, a nucleotide sequence comprising an AAV5 ITR, a nucleotide sequence comprises an AAV5 Rep coding sequence (i.e. a nucleotide sequence comprises an AAV5 Rep78). Such ITR and Rep sequences can be modified as desired to obtain efficient production of AAV5 or pseudotyped AAV5 vectors in insect cells. E.g., the start codon of the Rep sequences can be modified, VP splice sites can be modified or eliminated, and/or the VP1 start codon and nearby nucleotides can be modified to improve the production of AAV5 vectors in the insect cell.

[0043] In another aspect the invention thus relates to a method for producing a recombinant parvoviral (rAAV) virion (comprising a recombinant parvoviral (rAAV) vector as defined above) in an insect cell. Preferably, the method comprises the steps of: (a) culturing an insect cell as defined in herein above under conditions such that recombinant parvoviral (rAAV) vector is produced; and, (b) recovery of the recombinant parvoviral (rAAV) vector. It is understood here that the recombinant parvoviral (rAAV) vector produced in the method preferably is an infectious parvoviral or AAV virion that comprise the recombinant parvoviral (rAAV) vector nucleic acids. Growing conditions for insect cells in culture, and production of heterologous products in insect cells in culture are well-known in the art and described e.g. in the above cited references on molecular engineering of insects cells.

[0044] Preferably the method further comprises the step of affinity-purification of the (virions comprising the) recombinant parvoviral (rAAV) vector using an anti-AAV antibody, preferably an immobilised antibody. The anti-AAV antibody preferably is a monoclonal antibody. A particularly suitable antibody is a single chain camelid antibody or a fragment thereof as e.g. obtainable from camels or llamas (see e.g. Muyldermans, 2001, Biotechnol. 74: 277-302). The antibody for affinity-purification of rAAV preferably is an antibody that specifically binds an epitope on a AAV capsid protein, whereby preferably the epitope is an epitope that is present on capsid protein of more than one AAV serotype. E.g. the antibody may be raised or selected on the basis of specific binding to AAV2 capsid but at the same time also it may also specifically bind to AAV1, AAV3 and AAV5 capsids.

[0045] In a further aspect the invention relates to a rAAV virion produced in the above described methods of the invention, using the nucleic acid constructs and cells as defined above. Preferably the rAAV virion comprises in its genome at least one nucleotide sequence encoding a gene product of interest, whereby the at least one nucleotide sequence is not a native AAV nucleotide sequence, and whereby in the stoichiometry of the AAV VP1, VP2, and VP3 capsid proteins the amount of VP1: (a) is at least 100, 105, 110, 120, 150, 200 or 400% of the amount of VP2; or (b) is at least 8, 10, 10.5, 11, 12, 15, 20 or 40% of the amount of VP3; or (c) is at least as defined in both (a) and (b). Preferably, the amount of VP1, VP2 and VP3 is determined using an antibody recognising an epitope that is common to each of VP1, VP2 and VP3. Various immunoassays are available in the art that will allow quantify the relative amounts of VP1, VP2 and/or VP3 (see e.g. Using Antibodies, E. Harlow and D. Lane, 1999, Cold Spring Harbor Laboratory Press, New York). A suitable antibody recognising an epitope that is common to each of the three capsid proteins is e.g. the mouse anti-Cap B1 antibody (as is commercially available from Progen, Germany). A preferred rAAV virion according to the invention is a virion comprising in its genome at least one nucleotide sequence encoding a gene product of interest, whereby the at least one nucleotide sequence is not a native AAV nucleotide sequence, and whereby the AAV virion comprises a VP1 capsid protein comprises a leucine or a valine at amino acid position 1. A more preferred AAV virion according to the invention has the ratio's of capsid proteins as defined above and comprises a VP1 capsid protein comprises a leucine or a valine at amino acid position 1.

[0046] In further aspects the invention pertains to the following numbered embodiments:

1. 1. A nucleotide sequence comprising an open reading frame comprising nucleotide sequences encoding parvoviral Rep proteins, wherein the initiation codon for translation of the parvoviral Rep78 protein is an initiation codon that effects partial

exon skipping upon expression in insect cells.

2. 2. A nucleotide sequence according to embodiment 1, wherein the initiation codon is selected from ACG, TTG, CTG, and GTG.
3. 3. A nucleotide sequence according to embodiments 1 or 2, wherein the nucleotide sequence comprises an expression control sequence comprising a nine nucleotide sequence of SEQ. ID NO: 7 or a nucleotide sequence substantially homologous to SEQ. ID NO: 7, upstream of the initiation codon of the nucleotide sequence encoding the AAV Rep78 protein.
4. 4. A nucleotide sequence according to any one of embodiments 1 - 3, wherein the parvoviral Rep proteins are adeno-associated virus (AAV) Rep proteins.
5. 5. A nucleic acid construct comprising a nucleotide sequence according to any one of embodiments 1 - 4, wherein the nucleotide sequence is operably linked to expression control sequences for expression in an insect cell.
6. 6. A nucleic acid construct according to embodiment 5, wherein the nucleotide sequence is operably linked to a polyhedron promoter.
7. 7. A nucleic acid construct according to embodiments 5 or 6, wherein the construct is an insect cell-compatible vector, preferably a baculoviral vector.
8. 8. An insect cell comprising no more than one type of nucleotide sequence comprising a single open reading frame encoding one or more parvoviral Rep proteins.
9. 9. An insect cell according to embodiment 8, wherein the single open reading frame encodes the full-length Rep 78 protein.
10. 10. An insect cell according to embodiment 8 or 9, wherein the nucleotide sequence comprising the single open reading frame encoding one or more parvoviral Rep proteins is part of a nucleic acid construct wherein the nucleotide sequence is operably linked to expression control sequences for expression in an insect cell.
11. 11. An insect cell according to any one of embodiments 8 - 10, wherein the insect cell comprises a first nucleotide sequence comprising a single open reading frame encoding one or more parvoviral Rep proteins as defined in any one of embodiment 1 - 4, or a first nucleic acid construct wherein the nucleotide sequence is operably linked to expression control sequences for expression in an insect cell as defined in any one of embodiments 5 - 7.
12. 12. An insect cell according to embodiment 11, wherein the insect cell further comprises:
 1. a) a second nucleotide sequence comprising at least one parvoviral inverted terminal repeat (ITR) nucleotide sequence; and,
 2. b) a third nucleotide sequence comprising parvoviral capsid protein coding sequences operably linked to expression control sequences for expression in an insect cell.
13. 13. A insect cell according to embodiment 12, wherein the insect cell comprises:
 1. a) a first nucleic acid construct according to any of embodiments 5 - 7, whereby the first nucleic acid construct further comprises a third nucleotide sequence as defined in (b) of embodiment 9; and,
 2. b) a second nucleic acid construct comprising the second nucleotide sequence as defined in (a) of embodiment 9.
14. 14. An insect cell according to embodiment 13, wherein the second nucleic acid construct is an insect cell-compatible vector, preferably a baculoviral vector.
15. 15. An insect cell according to any one of embodiments 12 - 14, wherein the second nucleotide sequence further comprises at least one nucleotide sequence encoding a gene product of interest and whereby the at least one nucleotide sequence encoding a gene product of interest becomes incorporated into the genome of an parvoviral vector produced in the insect cell.
16. 16. An insect cell according to embodiment 15, wherein the second nucleotide sequence comprises two parvoviral ITR nucleotide sequences and wherein the at least one nucleotide sequence encoding a gene product of interest is located between the two parvoviral ITR nucleotide sequences.
17. 17. An insect cell according to any one of embodiments 12 - 16, wherein the third nucleotide sequence comprising an open reading frame comprising nucleotide sequences encoding parvoviral VP1, VP2, and VP3 capsid proteins, wherein the initiation codon for translation of the parvoviral VP1 capsid protein is selected from ACG, TTG, CTG, and GTG.
18. 18. An insect cell according to embodiment 17, wherein the third nucleotide sequence comprises an expression control sequence comprising a nine nucleotide sequence of SEQ. ID NO: 7 or a nucleotide sequence substantially homologous to SEQ. ID NO: 7, upstream of the initiation codon of the nucleotide sequence encoding the parvoviral VP1 capsid protein.
19. 19. An insect cell according to embodiments 17 or 18, wherein the third nucleotide sequence further comprises at least one modification of the nucleotide sequence encoding the parvoviral VP1 capsid protein selected from among a C at nucleotide position 12, an A at nucleotide position 21, and a C at nucleotide position 24.
20. 20. An insect cell according to embodiment any one of embodiments 12 - 19, wherein at least one of the first nucleotide sequence, second nucleotide sequence, and third nucleotide sequence are stably integrated in the genome of the insect cell.
21. 21. An insect cell according to embodiment any one of embodiments 8 - 20, wherein the parvovirus is AAV.
22. 22. A method for producing an a recombinant parvoviral virion in an insect cell, the virion comprising a second nucleotide sequence as defined in any one of embodiments 12, 15 and 16, the method comprising the comprising the steps of:

1. a) culturing an insect cell as defined in any one of embodiments 12 - 21 under conditions such that recombinant parvoviral virion is produced; and,
2. b) recovery of the recombinant parvoviral virion.
23. 23. A method according to embodiment 22, further comprising the step of affinity-purification of the virion using an anti-parvoviral antibody, preferably an immobilised antibody.
24. 24. A method according to embodiment 23, wherein the anti-parvoviral antibody is a single chain cameloid antibody or a fragment thereof.
25. 25. A method according to any one of embodiments 22 - 24, wherein the recombinant parvoviral virion is a recombinant AAV virion.

[0047] In this document and in its claims, the verb "to comprise" and its conjugations is used in its non-limiting sense to mean that items following the word are included, but items not specifically mentioned are not excluded. In addition, reference to an element by the indefinite article "a" or "an" does not exclude the possibility that more than one of the element is present, unless the context clearly requires that there be one and only one of the elements. The indefinite article "a" or "an" thus usually means "at least one".

Description of the figures

[0048]

Figure 1:

1. A) Organisation of Rep expression in the wild type AAV genome. The Rep78 and Rep 52 genes are expressed from respectively the P5 and P19 promoter. Expression of Rep68 and Rep40 (which are the spliced variants of resp. Rep78 and Rep52) are not shown. Both expression units contain a ATG-initiation site.
2. B) The construct of the invention has the Rep ORF under the control of a single promoter (e.g. the polyhedron (PolH) promoter). This promoter drives the expression of both Rep78 and Rep52 because the Rep78 initiation codon ATG is converted to the alternate ACG initiation codon and partially skipped by the ribosome.
3. C) The original construct by Urabe et al. (2002, *supra*) drives Rep78 and Rep52 independently from two different promoters (resp. Δ IE1 and polH).

Figure 2: Western blot analysis of Rep proteins expressed from recombinant baculovirus that was passaged 5 times on insect cells. The original baculovirus designed by Urabe et al., 2002 (original REP/Bac-to-Bac) results in a slow decrease of Rep78/52 expression over 5 passages. The expression unit for Rep78 and 52 designed by Urabe et al., 2002 inserted in baculovirus backbone PSC (original REP / PSC) also results in a decrease of Rep78/52 expression following passaging on insect cells. However, the baculovirus with the REP expression unit containing the ACG initiation codon in the PSC backbone (REP-ACG / PSC) results in stable expression of Rep78/52 over at least 5 passages. Western blot analysis was performed as described in Example 1.1.3.

Figure 3: Results of Table 1 plotted in a graph.

Figure 4: Comparison of the stabilities of various rAAV constructs in insect cells. rAAV production in SF+ cells was performed as described above in Example 1. For all productions the ITR containing baculovirus and the capsid gene containing baculovirus were identical, the passage number was the same as the Rep gene containing baculoviruses. 4 different Rep gene containing baculoviruses were used: 1) The REP-ACG / PSC, 2) SLR: the original construct by Urabe et al. (2002, *supra*), 3) Rep52 + Rep78(B2B): Two separate Bac-to-Bac baculoviruses, one containing the Rep 78 gene and the other one containing the Rep 52 gene. 4) Rep52 + Rep78(PSC): Two separate protein sciences baculoviruses one containing the Rep 78 gene and the other one containing the Rep 52 gene.

Figure 5: Stability of the REP-ACG / PSC baculovirus constructs up to passage 8. rAAV productions in SF+ cells were performed as described in Example 1.

Figure 6: Comparison of the effect of passage effect on rep protein expression of the original construct from Urabe et al. (2002, *supra*) with a REP-ACG / PSC construct in accordance with the invention. The baculovirus passages and the western blot were done as described in Example 1. During a normal passage of the rep baculoviruses, samples were taken at 40 hours after addition of the baculoviruses to the SF cells and western blot was performed.

Examples**Example 1: Rep constructs****1.1. Materials &Methods****1.1.1 Baculovirus plasmid construction**

[0049] In order to express Rep78 and Rep52 from a sole bicistronic messenger RNA, the ATG initiation codon of Rep78 situated on the expression vector pFastBacDualSLR (Urabe et al., 2002, *supra*) was converted to ACG. The upstream primer used was:

BamHI

5'-cgcgatcctgtaagACGGCGGGGTTTTACGAGATTGTGATTAAGGTC-3' (SEQ ID NO.8)

PRIMER SEQUENCE forward

[0050] The 3'-primer that was used in the PCR reaction was flanking the REP78 gene and contains a XbaI site (TCTAGA):

XbaI

5'-AGGCTCTAGATTCGAAAGCGGCCCG-3'

(SEQ ID NO.9)

PRIMER SEQUENCE reverse

[0051] The sequence between the above-mentioned primer set was amplified by PCR (reaction volume 50 µl; 1x Pfx Amp. Buffer, 0.3mM dNTP's, 1mM MgSO₄, 150mM primer forw., 150mM primer rev., 2x enhancer solution, template 50ng (pFastBacDualSLR), 1 U Platinum Pfx (Invitrogen, Carlsbad, CA, USA) using the following protocol: 1 cycle of 95°C, 5 min; 35 cycles of 95°C, 15 sec; 55°C, 30 sec; 72°C, 2 min; 1 cycle of 72°C, 10 min; 4°C, for ever). The PCR product was cloned in PCR-blunt II-TOPO using the Zero Blunt TOPO PCR cloning kit (Invitrogen). The Rep78 was subcloned into pFastBacDual (Invitrogen) using the restriction sites SpeI and XbaI. The mutated Rep expression cassette was finally cloned (using restriction enzymes BstZ17I and AvrII) into the baculovirus expression construct (cut open with EcoRV and XbaI) pPSC10 (Protein Sciences Corporation, Meriden, CT, USA). The sequence analysis of the construct was verified by Baseclear, Leiden, the Netherlands.

1.1.2 Recombinant baculovirus production

[0052] Recombinant baculoviruses derived from the Autographa californica nuclear polyhydrosis virus (AcNPV) were produced using the GeneXpress BaculoKIT (Protein Sciences Corporation). Transfection was performed as follows: in a round bottom 14ml tube 200 µl GRACE medium was mixed with 6 µl cellfectine (Invitrogen), and in a eppendorf tube 200 µl GRACE medium was mixed with 50 µl viral DNA (protein sciences) and 2 µg transfer plasmid (REP). The contents from the eppendorf tube were added to the tube and mixed carefully. After an incubation period of 30 minutes at RT 1,300 µl GRACE was added to the transfection mix. Insect cells in a T25 flask were washed with GRACE medium and the transfection mixture was added dropwise to the cell layer. After an incubation of 6 hours at 28°C SF900II serum supplemented with 10% FBS was added carefully and the T25 flask

was put in a 28°C stove for 5 days after which the recombinant baculovirus was harvested.

1.1.3 Western blot analysis

[0053] Insect cells (SF+) were infected with baculovirus-REP. At 16, 40, and 64 hours post-infection cells a sample was taken and cells were lysed by adding 0.1V 10 x TRIS lysis buffer (1.5M NaCl, 0.5M TRIS, 0.01M MgCl, 1% TRITON X-100, pH8.5, filter sterilised) and incubated at 28°C for 30 minutes in a shaker (Innova 44, New Brunswick). Free DNA and RNA was degraded by incubation with benzonase at 37°C for 30 minutes. Cell lysate was centrifuged (1,900 x g; 15 min; 4°C). NuPAGE LDS sample buffer (4x, Invitrogen) was added to a sample of the supernatant and was loaded onto a 4-12% Bis-Tris gel (120V). Proteins were blotted onto a PVDF membrane (BioRad) for 30 minutes, 10V (Semidry blotting). Western immunochrometry was performed by blocking the membrane with Superblock-PBS blocking buffer (PIERCE) and subsequent incubation with mouse anti-Rep (303.9, Progen, Germany; dilution 1:50) and rabbit anti-mouse-HRP (DAKO, dilution 1:500). The Rep-proteins were visualized by chemoluminescent staining with lumi-light plus Western-blotting substrate (Roche).

1.2 Results

[0054] The performance of the newly designed Rep-construct of the invention (REP-ACG / PSC) was compared with the original Rep constructs in both 1) PSC baculovirus backbone and in 2) Bac-to-Bac baculovirus backbone (Urabe et al., 2002). All three constructs were serially passaged until passage 5. AAV1-LPL production experiments were performed using the passage 2, 3, 4 and 5 Rep-constructs in combination with an AAV-LPL and a AAV-Cap recombinant baculovirus of respectively passage 2, 3, 4 and 5 (AAV-LPL and AAV-Cap recombinant Baculovirus used here are described below in Example 2). AAV1-LPL production yields were determined by qPCR and are shown in Table 1. The original baculovirus designed by Urabe et al., 2002 (original REP/Bac-to-Bac) results in a fast decrease of AAV production over 5 passages. The expression unit for Rep designed by Urabe et al., 2002 inserted in baculovirus backbone PSC (original REP / PSC) also results in a decrease of AAV production following passaging on insect cells. However, the baculovirus with the REP expression unit containing the ACG initiation codon in the PSC backbone (REP-ACG / PSC) results in stable AAV production over at least 5 passages. Therefore, reproducible production yields of AAV-LPL over several passages (e.g. 2 to 5) were only obtained using baculoviruses containing the REP-ACG construct.

Table 1: Production of rAAV virions using the baculovirus constructs of several passages: Sf9 cells were infected with three recombinant baculoviruses encoding a LPL-vector unit of passage 2, 3, 4 or 5, a Rep-expression unit of passage 2, 3, 4 or 5 and a Cap-expression unit of passage 2, 3, 4 or 5. After three days cells were harvested and AAV yields (vector genomes per ml; vg/ml) were determined by qPCR.

passage	original REP / PSC vg/ml	REP-ACG / PSC vg/ml	original REP / Bac-to-Bac vg/ml
2	5.38E+09	3.04E+09	3.62E+10
3	9.57E+09	4.77E+09	7.28E+09
4	1.66E+09	7.81E+09	7.59E+08
5	7.35E+08	9.90E+09	2.03E+08

Table 2: Q-PCR performed on the various Bac-Rep constructs following passaging on insect cells (Passage 2-5).

	titer (gc's/ml)				
	ORF	Rep78	Rep52	Ratio ORF/Rep78	Ratio ORF/Rep52
original REP/Bac-to-Bac P2	1,4E+09	2,2E+08	2,4E+08	6,42	5,82
original REP/Bac-to-Bac P3	6,4E+08	5,6E+07	5,0E+07	11,43	12,93
original REP/Bac-to-Bac P4	2,1E+09	7,1E+07	6,5E+07	29,47	32,02
original REP/Bac-to-Bac P5	1,7E+09	3,2E+07	2,5E+07	53,68	69,67
REP-ACG / PSC (C4) P2	3,0E+09	2,7E+09	2,9E+09	1,11	1,04
REP-ACG / PSC (C4) P3	2,3E+09	2,0E+09	2,2E+09	1,11	1,05
REP-ACG / PSC (C4) P4	2,5E+09	2,2E+09	2,3E+09	1,13	1,08
REP-ACG / PSC (C4) P5	2,7E+09	2,1E+09	2,5E+09	1,26	1,07

	titer (gc's/ml)				
	ORF	Rep78	Rep52	Ratio ORF/Rep78	Ratio ORF/Rep52
REP-ACG / PSC (A3) P2	2,5E+09	2,2E+09	2,5E+09	1,18	1,00
REP-ACG / PSC (A3) P3	4,2E+09	3,9E+09	4,0E+09	1,08	1,04
REP-ACG / PSC (A3) P4	2,7E+09	2,4E+09	2,5E+09	1,10	1,05
REP-ACG / PSC (A3) P5	1,5E+09	1,5E+09	1,5E+09	1,03	0,98
original REP/Bac-to-Bac P2	1,0E+09	1,1E+09	1,1E+09	0,95	0,87
original REP/Bac-to-Bac P3	7,1E+08	6,7E+08	8,1E+08	1,07	0,88
original REP/Bac-to-Bac P4	1,3E+08	1,1E+08	1,3E+08	1,18	1,03
original REP/Bac-to-Bac P5	1,3E+08	5,3E+07	6,9E+07	2,34	1,82

[0055] Table 2 shows the results of a quantitative PCR (Q-PCR) assay that was designed for the Rep-expression unit in the recombinant baculoviruses and for a flanking baculovirus ORF (gene copies per ml; gc's/ml). The ratio between the Q-PCR values determines the presence of deletions in the Rep-baculovirus. A ratio of 1 theoretically means that all baculoviruses in the batch contain a recombinant Rep78 or 52-sequence. The original baculovirus designed by Urabe et al., 2002 (original REP/Bac-to-Bac) shows significant amounts of the recombinant baculovirus at passage 5 have deletions in the Rep sequences. The expression unit for Rep78 and 52 designed by Urabe et al., 2002 inserted in baculovirus backbone PSC (original REP / PSC) shows a very early and dramatic loss of recombinant baculovirus. However, the baculovirus with the REP expression unit containing the ACG initiation codon in the PSC backbone (REP-ACG / PSC) (clone C4 and A3) show stable recombinant baculoviruses over at least 5 passages.

Example 2: Cap constructs

2.1.1 Baculovirus plasmid construction

[0056] In order to express VP1,2,3 from a sole polycistronic messenger RNA, the baculovirus-AAV-Cap construct was designed as described by (Urabe et al., 2002, supra). Briefly, the ATG initiation codon of VP1 was mutated to ACG. A potential ATG initiation codon at position 11 has been changed to ACG. The splice acceptor site downstream of the VP1 initiation codon was destroyed (mutation at position 21 and 24). The mutated Cap expression cassette was cloned into a baculovirus expression construct; pFastBacDual (pFBDAAV1VPm11) with BamHI/StuI restriction sites. This plasmid (pFBDAAV1VPm11) was the starting material for introduction of alternate initiation codons for VP1. The forward primer used by Urabe et al. (2002, supra) in order to introduce the mentioned mutations was:

BamHI 1 11 21 24

5'-cgcggtatcctgttaagACGGCTGCCGACGGTTATCTACCCGATTGGCTC-3'

(SEQ ID NO. 1)

[0057] The following forward primers were used to make the expression constructs using pFBDAAV1VPm11 (Urabe et al., 2002, supra) as starting material:

5'-cgcggtatcctgttaagTTGGCTGCCGACGGTTATCTACCCGATTGGCTC-3'

(SEQ ID NO.2)

5'-cgcggtatcctgttaagATTGCTGCCGACGGTTATCTACCCGATTGGCTC-3'

(SEQ ID NO.3)

5'-cgcggtatcctgttaagGTGGCTGCCGACGGTTATCTACCCGATTGGCTC-3'

(SEQ ID NO. 4)

5'-cgcggtatcctgttaagCTGGCTGCCGACGGTTATCTACCCGATTGGCTC-3'

(SEQ ID NO. 5)

[0058] The backward-primer that was used in the PCR reactions with the above forward primers was directed to position ~ 230 bp downstream of the VP1 initiation codon and contains a unique Stu I site (AGGCCT).

5'-GTCGTAGGCCTTGTCGTGCTCGAGGGCCGC-3'

(SEQ ID NO. 6)

[0059] Fragments were amplified with the above-mentioned sets of forward and backward primer pairs by PCR. Following digestion of PCR products with BamHI and StuI the PCR products were subcloned into the BamHI / StuI sites of pFBDAAV1vpm11 resulting in the various to be tested baculovirus-AAV-Cap constructs. DNA constructs were verified by sequence analysis at Baseclear, Leiden, the Netherlands.

2.1.2 Recombinant baculovirus production

[0060] Recombinant baculoviruses derived from the Autographa californica nuclear polyhydrosis virus (AcNPV) were produced using the Bac-to-Bac baculovirus expression system (Invitrogen). rBac-Cap was amplified by infecting 2×10^6 Sf9 cells per ml at an moi of 0.1. Three days after infection the cells were spun down and the supernatant containing the virus recovered.

2.1.3 recombinant AAV production

[0061] rAAV batches were produced using three recombinant baculoviruses according to Urabe et al., 2002. However, for this study one baculovirus harboured an expression construct for the LPL^{S447X}-transgene. The therapeutically active agent expressed from the transgene is a naturally occurring variant of human lipoprotein lipase, a single chain polypeptide of 448 amino acids. The LPL^{S447X} variant has a deletion of two amino acids at the C-terminus of the protein. The second baculovirus harboured an expression construct for the AAV replication genes, Rep 78 and Rep 52. The third baculovirus harboured the AAV1 capsid sequence with either an ACG or a TTG, CTG, GTG initiation codon for VP1.

[0062] Mammalian-rAAV batches produced with the plasmid-transfection system were produced according to Grimm et al., 1998 (Novel tools for production and purification of recombinant adeno-associated virus vectors. Hum Gene Ther. 1998 Dec 10;9(18):2745-60).

2.1.3 Western blot analysis

[0063] Insect cells were infected with baculovirus-Cap. At three days post-infection cells were centrifuged (3,000 g; 15 min). The supernatant was filtered through a 0.22µm Millex filter. NuPAGE LDS sample buffer (Invitrogen) was added to a sample of the supernatant and was loaded onto a 4-12% Bis-Tris gel. The gel was run at 100V. Proteins were blotted onto a nitrocellulose membrane (BioRad) for 1 hr, 100V, 350mA. Western immunochemistry was performed by blocking the membrane with 1% marvel, dried skimmed milk and subsequently incubation with mouse anti-Cap (B1 from Progen, Germany; dilution 1:50) and rabbit anti-mouse-HRP (DAKO, dilution 1:100). VP1 ,2 and 3 were visualized by chemoluminescent staining with lumi-light plus Western-blotting substrate (Roche).

2.1.4 Biochemical measurements

[0064] Human LPL^{S447X} activity was assayed as previously described using a radioactive trioleoylglycerol emulsion substrate (Nilsson-Ehle and Scholtz, 1976). Human LPL^{S447X} immunoreactive mass was assayed using a sandwich ELISA with chicken IgY and mouse 5D2 anti-hLPL antibodies (Liu et al., 2000). Plasma triglyceride levels were measured by using commercial kits following manufacturer protocols (Boehringer Mannheim, #450032).

2.2 Results

2.2.1 Construction of recombinant baculovirus

[0065] In order to introduce different alternate initiation codons for VP1 expression in the baculovirus plasmid designed by Urabe et al. (2002, *supra*) a series of upstream primers were designed containing a BamHI restriction site and either a TTG, ATT, GTG or CTG codon in place of the ACG initiation codon of VP1. PCR using these primers in combination with a downstream primer containing a StuI site resulted in amplified fragments that were subcloned into the BamHI/StuI site of pFBDVPm11 (Bac-Cap). The resulting baculovirus plasmids were used for the preparation of recombinant baculoviruses using the Bac-to-Bac baculovirus expression system. The prepared recombinant baculoviruses were infected on insect cells in order to produce AAV capsids. At three days following infection viral protein expression of the different baculovirus batches were determined on Western blots. From the Western blots it became clear that the baculovirus construct containing the TTG initiation codon for VP1 expressed this protein to a higher level compared to the previously used ACG initiation codon. The ratio between VP1 and VP2 using the TTG codon was found to be 1:1 which is similar to what is reported for wild type AAV (not shown).

2.2.2 Infection of rAAV batches on cells in culture

[0066] In order to investigate the infectivity of the AAV capsids derived from recombinant baculoviruses with the TTG initiation codon rAAV was generated. Also a rAAV batch was generated by plasmid transfection on mammalian HEK293 cells. A vector genome titer of both rAAV batches was determined by qPCR. This titer was used to infect HEK 293 cells in a microtiter plate at an increasing moi. At two days following infection an quantitative assay (LPL^{S447X}-mass assay) for the transgene product (LPL^{S447X}) was performed on the medium of the infected cells. The assay showed that the amount of LPL^{S447X} produced by baculovirus-produced rAAV was similar to the LPL produced by the plasmid-produced rAAV (not shown).

2.2.3 Injection of rAAV batches in mice

[0067] The rAAV batches produced with the baculovirus-production system and with the conventional mammalian plasmid-production system were injected intramuscularly in mice to follow LPL^{S447X}-protein activity and triglyceride fasting in vivo. At 3 days, 7 days and at 2 weeks following injection blood samples were taken and evaluated. Between 3 and 7 days post virus administration blood-plasma sampled from both mice injected with mammalian-rAAV and one mouse injected with baculo-rAAV was turned from milky to completely clear. Blood plasma derived from one baculo-rAAV-injected mouse remained relatively milky however fat level was clearly reduced. Triglyceride levels were lowered respectively of all treated mice (not shown). On day 14 TG levels in both mammalian-AAV and baculovirus-(TTG)-AAV treated mice TG levels were reduced for 96%. Plasma samples taken at two weeks after virus administration showed that the LPL^{S447X}-activity of the mice treated with baculovirus-AAV and mammalian-AAV was similar (not shown).

Example 3: Stability of rAAV constructs with modified Rep 78 initiation codon in insect cells

3.1 Comparison of the stabilities of various rAAV constructs in insect cells

[0068] rAAV productions in SF+ cells were performed as described above in Example 1. For all productions the ITR containing baculovirus and the capsid gene containing baculovirus were identical, the passage number was the same as the Rep gene containing baculoviruses. 4 different Rep gene containing baculoviruses were used: 1) The REP-ACG / PSC, 2) SLR: the original construct by Urabe et al. (2002, *supra*), 3) Rep52 + Rep78(B2B): Two separate Bac-to-Bac baculoviruses, one containing the Rep 78 gene and the other one containing the Rep 52 gene. 4) Rep52 + Rep78(PSC): Two separate protein sciences baculoviruses one containing the Rep 78 gene and the other one containing the Rep 52 gene.

[0069] Results are shown in Figure 4. At fifth baculovirus passage rAAV production is already improved by more than a factor 10

using a REP-ACG / PSC in accordance with invention as compared to the original Rep construct and compared to the split Rep constructs.

3.2 Stability of the baculovirus constructs up to passage 8

[0070] rAAV productions in SF+ cells were performed as described in Example 1. For all productions the ITR containing baculovirus and the capsid gene containing baculovirus were identical, the passage number was the same as the REP-ACG / PSC baculovirus. Results are shown in Figure 5. The REP-ACG / PSC baculovirus is stable to at least passage 8. rAAV production titers of REP-ACG / PSC are stable up to at least 8th passage of the baculovirus.

3.3 Passage effect on rep protein expression

[0071] The effect of passage number on the expression of Rep protein for the original construct from Urabe et al. (2002, *supra*) was compared to a REP-ACG / PSC construct in accordance with the invention. The baculovirus passages and the western blot were done as described in Example 1. During a normal passage of the rep baculoviruses, samples were taken at 40 hours after addition of the baculoviruses to the SF cells and western blot was performed. Figure 6 clearly shows diminished Rep expression in higher passages compared to earlier passages for the original Urabe construct (SLR), while the Rep expression in the REP-ACG / PSC construct stays the same in the higher passages compared to the lower ones.

SEQUENCE LISTING

[0072]

<110> Amsterdam Molecular Therapeutics B.V.

<120> AAV vectors with improved Rep coding sequences for production in insect cells

<130> P6009362PCT

<140> P6009362PCT

<141> 2007-06-21

<150> EP06115804.4

<151> 2006-06-21

<150> US 60/815,262

<151> 2006-06-21

<160> 11

<170> PatentIn version 3.3

<210> 1

<211> 49

<212> DNA

<213> artificial synthetic sequence

<220>

<223> primer

<400> 1

cgcggaatcct gtaagacgg ctgccgacgg ttatctaccc gattggctc 49

<210> 2

<211> 49

<212> DNA

<213> artificial synthetic sequence

<220>
 <223> primer

<400> 2
 cgcggatcct gtaagttgg ctgccgacgg ttatctaccc gattggctc 49

<210> 3
 <211> 49
 <212> DNA
 <213> artificial synthetic sequence

<220>
 <223> primer

<400> 3
 cgcggatcct gtaagattg ctgccgacgg ttatctaccc gattggctc 49

<210> 4
 <211> 49
 <212> DNA
 <213> artificial synthetic sequence

<220>
 <223> primer

<400> 4
 cgcggatcct gtaaggtgg ctgccgacgg ttatctaccc gattggctc 49

<210> 5
 <211> 49
 <212> DNA
 <213> artificial synthetic sequence

<220>
 <223> primer

<400> 5
 cgcggatcct gtaagctgg ctgccgacgg ttatctaccc gattggctc 49

<210> 6
 <211> 30
 <212> DNA
 <213> artificial synthetic sequence

<220>
 <223> primer

<400> 6
 gtcgtaggcc ttgctgtgt cgagggccgc 30

<210> 7
 <211> 9
 <212> DNA
 <213> adeno-associated virus 2

<400> 7
 cctgttaag 9

<210> 8
 <211> 49
 <212> DNA
 <213> Artificial

<220>
 <223> primer

 <400> 8
 cgcggtatcct gttaagacgg cggggtttta cgagattgtg attaaggtc 49

 <210> 9
 <211> 25
 <212> DNA
 <213> Artificial

 <220>
 <223> primer

 <400> 9
 aggctctaga ttcgaaagcg gcccg 25

 <210> 10
 <211> 1876
 <212> DNA
 <213> adeno-associated virus 2

 <220>
 <221> CDS
 <222> (11)..(1876)
 <223> Rep 78

 <220>
 <221> misc_feature
 <222> (683)..(1876)
 <223> coding sequence Rep52

 <400> 10

cgagccgcc atg ccg ggg ttt tac gag att gtg att aag gtc ccc agc Met Pro Gly Phe Tyr Glu Ile Val Ile Lys Val Pro Ser 1 5 10	49
gac ctt gac gag cat ctg ccc ggc att tct gac agc ttt gtg aac tgg Asp Leu Asp Glu His Leu Pro Gly Ile Ser Asp Ser Phe Val Asn Trp 15 20 25	97
gtg gcc gag aag gaa tgg gag ttg ccg cca gat tct gac atg gat ctg Val Ala Glu Lys Glu Trp Glu Leu Pro Pro Asp Ser Asp Met Asp Leu 30 35 40 45	145
aat ctg att gag cag gca ccc ctg acc gtg gcc gag aag ctg cag cgc Asn Leu Ile Glu Gln Ala Pro Leu Thr Val Ala Glu Lys Leu Gln Arg 50 55 60	193
gac ttt ctg acg gaa tgg cgc cgt gtg agt aag gcc ccg gag gcc ctt Asp Phe Leu Thr Glu Trp Arg Arg Val Ser Lys Ala Pro Glu Ala Leu 65 70 75	241
ttc ttt gtg caa ttt gag aag gga gag agc tac ttc cac atg cac gtg Phe Phe Val Gln Phe Glu Lys Gly Glu Ser Tyr Phe His Met His Val 80 85 90	289
ctc gtg gaa acc acc ggg gtg aaa tcc atg gtt ttg gga cgt ttc ctg Leu Val Glu Thr Thr Gly Val Lys Ser Met Val Leu Gly Arg Phe Leu 95 100 105	337
agt cag att cgc gaa aaa ctg att cag aga att tac cgc ggg atc gag Ser Gln Ile Arg Glu Lys Leu Ile Gln Arg Ile Tyr Arg Gly Ile Glu 110 115 120 125	385
ccg act ttg cca aac tgg ttc gcg gtc aca aag acc aga aat ggc gcc Pro Thr Leu Pro Asn Trp Phe Ala Val Thr Lys Thr Arg Asn Gly Ala 130 135 140	433
gga ggc ggg aac aag gtg gtg gat gag tgc tac atc ccc aat tac ttg Gly Gly Gly Asn Lys Val Val Asp Glu Cys Tyr Ile Pro Asn Tyr Leu 145 150 155	481
ctc ccc aaa acc cag cct gag ctg cag tgg gcg tgg act aat atg gaa Leu Pro Lys Thr Gln Pro Glu Leu Gln Trp Ala Trp Thr Asn Met Glu 160 165 170	529
cag tat tta agc gcc tgt ttg aat ctg acg gag cgt aaa cgg ttg gtg Gln Tyr Leu Ser Ala Cys Leu Asn Leu Thr Glu Arg Lys Arg Leu Val 175 180 185	577
gcg cag cat ctg acg cac gtg tcg cag acg cag gag cag aac aaa gag Ala Gln His Leu Thr His Val Ser Gln Thr Gln Glu Gln Asn Lys Glu 190 195 200 205	625
aat cag aat ccc aat tct gat gcg ccg gtg atc aga tca aaa act tca Asn Gln Asn Pro Asn Ser Asp Ala Pro Val Ile Arg Ser Lys Thr Ser 210 215 220	673
gcc agg tac atg gag ctg gtc ggg tgg ctg gtg gac aag ggg att acc Ala Arg Tyr Met Glu Leu Val Gly Trp Leu Val Asp Lys Gly Ile Thr 225 230 235	721
tcg gag aag cag tgg atc cag gag gac cag gcc tca tac atc tcc ttc Ser Glu Lys Gln Trp Ile Gln Glu Asp Gln Ala Ser Tyr Ile Ser Phe 240 245 250	769

aat gcg gcc tcc aac tcg cgg tcc caa atc aag gct gcc ttg gac aat Asn Ala Ala Ser Asn Ser Arg Ser Gln Ile Lys Ala Ala Leu Asp Asn 255 260 265	817
gcg gga aag att atg agc ctg act aaa acc gcc ccc gac tac ctg gtg Ala Gly Lys Ile Met Ser Leu Thr Lys Thr Ala Pro Asp Tyr Leu Val 270 275 280 285	865
ggc cag cag ccc gtg gag gac att tcc agc aat cgg att tat aaa att Gly Gln Gln Pro Val Glu Asp Ile Ser Ser Asn Arg Ile Tyr Lys Ile 290 295 300	913
ttg gaa cta aac ggg tac gat ccc caa tat gcg gct tcc gtc ttt ctg Leu Glu Leu Asn Gly Tyr Asp Pro Gln Tyr Ala Ala Ser Val Phe Leu 305 310 315	961
gga tgg gcc acg aaa aag ttc ggc aag agg aac acc atc tgg ctg ttt Gly Trp Ala Thr Lys Lys Phe Gly Lys Arg Asn Thr Ile Trp Leu Phe 320 325 330	1009
ggg cct gca act acc ggg aag acc aac atc gcg gag gcc ata gcc cac Gly Pro Ala Thr Thr Gly Lys Thr Asn Ile Ala Glu Ala Ile Ala His 335 340 345	1057
act gtg ccc ttc tac ggg tgc gta aac tgg acc aat gag aac ttt ccc Thr Val Pro Phe Tyr Gly Cys Val Asn Trp Thr Asn Glu Asn Phe Pro 350 355 360 365	1105
ttc aac gac tgt gtc gac aag atg gtg atc tgg tgg gag gag ggg aag Phe Asn Asp Cys Val Asp Lys Met Val Ile Trp Trp Glu Glu Gly Lys 370 375 380	1153
atg acc gcc aag gtc gtg gag tgc gcc aaa gcc att ctc gga gga agc Met Thr Ala Lys Val Val Glu Ser Ala Lys Ala Ile Leu Gly Gly Ser 385 390 395	1201
aag gtg cgc gtg gac cag aaa tgc aag tcc tgc gcc cag ata gac ccg Lys Val Arg Val Asp Gln Lys Cys Lys Ser Ser Ala Glu Ile Asp Pro 400 405 410	1249
act ccc gtg atc gtc acc tcc aac acc aac atg tgc gcc gtg att gac Thr Pro Val Ile Val Thr Ser Asn Thr Asn Met Cys Ala Val Ile Asp 415 420 425	1297
ggg aac tca acg acc ttc gaa cac cag cag ccg ttg caa gac cgg atg Gly Asn Ser Thr Thr Phe Glu His Gln Gln Pro Leu Gln Asp Arg Met 430 435 440 445	1345
ttc aaa ttt gaa ctc acc cgc cgt ctg gat cat gac ttt ggg aag gtc Phe Lys Phe Glu Leu Thr Arg Arg Leu Asp His Asp Phe Gly Lys Val 450 455 460	1393
acc aag cag gaa gtc aaa gac ttt ttc cgg tgg gca aag gat cac gtg Thr Lys Gln Glu Val Lys Asp Phe Phe Arg Trp Ala Lys Asp His Val 465 470 475	1441
gtt gag gtg gag cat gaa ttc tac gtc aaa aag ggt gga gcc aag aaa Val Glu Val Glu His Glu Phe Tyr Val Lys Lys Gly Gly Ala Lys Lys 480 485 490	1489
aga ccc gcc ccc agt gac gca gat ata agt gag ccc aaa cgg gtg cgc Arg Pro Ala Pro Ser Asp Ala Asp Ile Ser Glu Pro Lys Arg Val Arg 495 500 505	1537
gag tca gtt gcg cag cca tgc acg tca gac gcg gaa gct tgc atc aac Glu Ser Val Ala Gln Ser Ser Thr Ser Asp Ala Glu Ala Ser Ile Asn 510 515 520 525	1585
tac gca gac agg tac caa aac aaa tgt tct cgt cac gtg ggc atg aat Tyr Ala Asp Arg Tyr Gln Asn Lys Cys Ser Arg His Val Gly Met Asn 530 535 540	1633
ctg atg ctg ttt ccc tgc aga caa tgc gag aga atg aat cag aat tca Leu Met Leu Phe Pro Cys Arg Gln Cys Glu Arg Met Asn Gln Asn Ser 545 550 555	1681
aat atc tgc ttc act cac gga cag aaa gac tgt tta gag tgc ttt ccc Asn Ile Cys Phe Thr His Gly Lys Gln Lys Asp Cys Leu Glu Cys Phe Pro 560 565 570	1729
gtg tca gaa tct caa ccc gtt tct gtc gtc aaa aag gcg tat cag aaa Val Ser Glu Ser Gln Pro Val Ser Val Val Lys Lys Ala Tyr Gln Lys 575 580 585	1777
ctg tgc tac att cat cat atc atg gga aag gtg cca gac gct tgc act Leu Cys Tyr Ile His His Ile Met Gly Lys Val Pro Asp Ala Cys Thr 590 595 600 605	1825
gcc tgc gat ctg gtc aat gtg gat ttg gat gac tgc atc ttt gaa caa Ala Cys Asp Leu Val Asn Val Asp Leu Asp Asp Cys Ile Phe Glu Gln 610 615 620	1873
taa	1876
<210> 11	
<211> 621	
<212> PRT	
<213> adeno-associated virus 2	
<400> 11	

Met Pro Gly Phe Tyr Glu Ile Val Ile Lys Val Pro Ser Asp Leu Asp
 1 5 10 15
 Glu His Leu Pro Gly Ile Ser Asp Ser Phe Val Asn Trp Val Ala Glu
 20 25 30
 Lys Glu Trp Glu Leu Pro Pro Asp Ser Asp Met Asp Leu Asn Leu Ile
 35 40 45
 Glu Gln Ala Pro Leu Thr Val Ala Glu Lys Leu Gln Arg Asp Phe Leu
 50 55 60
 Thr Glu Trp Arg Arg Val Ser Lys Ala Pro Glu Ala Leu Phe Phe Val
 65 70 75 80
 Gln Phe Glu Lys Gly Glu Ser Tyr Phe His Met His Val Leu Val Glu
 85 90 95
 Thr Thr Gly Val Lys Ser Met Val Leu Gly Arg Phe Leu Ser Gln Ile
 100 105 110
 Arg Glu Lys Leu Ile Gln Arg Ile Tyr Arg Gly Ile Glu Pro Thr Leu
 115 120 125
 Pro Asn Trp Phe Ala Val Thr Lys Thr Arg Asn Gly Ala Gly Gly Gly
 130 135 140
 Asn Lys Val Val Asp Glu Cys Tyr Ile Pro Asn Tyr Leu Leu Pro Lys
 145 150 155 160
 Thr Gln Pro Glu Leu Gln Trp Ala Trp Thr Asn Met Glu Gln Tyr Leu
 165 170 175
 Ser Ala Cys Leu Asn Leu Thr Glu Arg Lys Arg Leu Val Ala Gln His
 180 185 190
 Leu Thr His Val Ser Gln Thr Gln Glu Gln Asn Lys Glu Asn Gln Asn
 195 200 205
 Pro Asn Ser Asp Ala Pro Val Ile Arg Ser Lys Thr Ser Ala Arg Tyr
 210 215 220
 Met Glu Leu Val Gly Trp Leu Val Asp Lys Gly Ile Thr Ser Glu Lys
 225 230 235 240
 Gln Trp Ile Gln Glu Asp Gln Ala Ser Tyr Ile Ser Phe Asn Ala Ala
 245 250 255
 Ser Asn Ser Arg Ser Gln Ile Lys Ala Ala Leu Asp Asn Ala Gly Lys
 260 265 270
 Ile Met Ser Leu Thr Lys Thr Ala Pro Asp Tyr Leu Val Gly Gln Gln
 275 280 285
 Pro Val Glu Asp Ile Ser Ser Asn Arg Ile Tyr Lys Ile Leu Glu Leu
 290 295 300
 Asn Gly Tyr Asp Pro Gln Tyr Ala Ala Ser Val Phe Leu Gly Trp Ala
 305 310 315 320
 Thr Lys Lys Phe Gly Lys Arg Asn Thr Ile Trp Leu Phe Gly Pro Ala
 325 330 335
 Thr Thr Gly Lys Thr Asn Ile Ala Glu Ala Ile Ala His Thr Val Pro
 340 345 350
 Phe Tyr Gly Cys Val Asn Trp Thr Asn Glu Asn Phe Pro Phe Asn Asp
 355 360 365
 Cys Val Asp Lys Met Val Ile Trp Trp Glu Glu Gly Lys Met Thr Ala
 370 375 380
 Lys Val Val Glu Ser Ala Lys Ala Ile Leu Gly Gly Ser Lys Val Arg
 385 390 395 400
 Val Asp Gln Lys Cys Lys Ser Ser Ala Gln Ile Asp Pro Thr Pro Val
 405 410 415

Ile Val Thr Ser Asn Thr Asn Met Cys Ala Val Ile Asp Gly Asn Ser
 420 425 430
 Thr Thr Phe Glu His Gln Gln Pro Leu Gln Asp Arg Met Phe Lys Phe
 435 440 445
 Glu Leu Thr Arg Arg Leu Asp His Asp Phe Gly Lys Val Thr Lys Gln
 450 455 460
 Glu Val Lys Asp Phe Phe Arg Trp Ala Lys Asp His Val Val Glu Val
 465 470 475 480
 Glu His Glu Phe Tyr Val Lys Lys Gly Gly Ala Lys Lys Arg Pro Ala
 485 490 495
 Pro Ser Asp Ala Asp Ile Ser Glu Pro Lys Arg Val Arg Glu Ser Val
 500 505 510
 Ala Gln Pro Ser Thr Ser Asp Ala Glu Ala Ser Ile Asn Tyr Ala Asp
 515 520 525
 Arg Tyr Gln Asn Lys Cys Ser Arg His Val Gly Met Asn Leu Met Leu
 530 535 540
 Phe Pro Cys Arg Gln Cys Glu Arg Met Asn Gln Asn Ser Asn Ile Cys
 545 550 555 560
 Phe Thr His Gly Gln Lys Asp Cys Leu Glu Cys Phe Pro Val Ser Glu
 565 570 575
 Ser Gln Pro Val Ser Val Val Lys Lys Ala Tyr Gln Lys Leu Cys Tyr
 580 585 590
 Ile His His Ile Met Gly Lys Val Pro Asp Ala Cys Thr Ala Cys Asp
 595 600 605
 Leu Val Asn Val Asp Leu Asp Asp Cys Ile Phe Glu Gln
 610 615 620

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- [US6156303A \[0003\]](#)
- [US5387484A \[0003\]](#)
- [US5741683A \[0003\]](#)
- [US5691176A \[0003\]](#)
- [US568676A \[0003\]](#)
- [US20020081721A \[0003\]](#)
- [WO0047757A \[0003\]](#)
- [WO0024916A \[0003\]](#)
- [WO9617947A \[0003\]](#)
- [US20030148506A \[0004\]](#)
- [US20040197895A \[0004\]](#)
- [NL2005050018W \[0004\]](#)
- [US4745051A \[0023\]](#)
- [US2003148506A \[0023\] \[0028\] \[0039\]](#)
- [WO03074714A \[0023\]](#)
- [US6103526A \[0026\]](#)

- [WO0100220A \[0033\]](#)
- [EP06115804A \[0072\]](#)
- [US60815262B \[0072\]](#)

Non-patent literature cited in the description

- **GRIMM et al.** Hum. Gene Ther., 1998, vol. 9, 2745-2760 [\[0003\]](#)
- **CLARK** Kidney Int., 2002, vol. 61, 19-15 [\[0003\]](#)
- **URABE et al.** Hum. Gene Ther, 2002, vol. 13, 1935-1943 [\[0004\]](#)
- **LI et al.** J Virol., 1997, vol. 71, 5236-43 [\[0005\]](#)
- **KOHLBRENNER et al.** Mol. Ther, 2005, vol. 12, 1217-25 [\[0006\]](#)
- **HENIKOFF HENIKOFF** PNAS, 1992, vol. 89, 915-919 [\[0011\]](#)
- Parvoviridae: The Viruses and Their Replication **KENNETH I. BERN** Fields Virology 19960000 [\[0015\]](#)
- **SAMBROOK RUSSELL** Molecular Cloning: A Laboratory Manual Cold Spring Harbor Laboratory Press 20010000 [\[0022\]](#)
- **SUMMERSSMITH** A Manual of Methods for Baculovirus Vectors and Insect Culture Procedures 19860000 [\[0023\]](#)
- **LUCKOW et al.** Cloning and Expression of Heterologous Genes Insect Cells with Baculovirus Vectors' Recombinant DNA Technology and Applications, 1991, 97-152 [\[0023\]](#)
- **KING, L. A. R. D. POSSE** The baculovirus expression system Chapman and Hall 19920000 [\[0023\]](#)
- **O'REILLY, D. R. L. K. MILLER V. A. LUCKOW** Baculovirus Expression Vectors: A Laboratory Manual 19920000 [\[0023\]](#)
- **W. H. FREEMAN RICHARDSON, C. D.** Baculovirus Expression Protocols, Methods Molecular Biology, 1995, vol. 39, [\[0023\]](#)
- **ELBASHIR et al.** Nature, 2001, vol. 411, 494-98 [\[0032\]](#)
- **CAPLEN et al.** Proc. Natl. Acad. Sci. USA, 2001, vol. 98, 9742-47 [\[0032\]](#)
- **SAMBROOK RUSSEL** Molecular Cloning: A Laboratory Manual Cold Spring Harbor Laboratory Press 20010000 [\[0034\]](#)
- **CLAIR et al.** Antimicrob. Agents Chemother., 1987, vol. 31, 844-849 [\[0034\]](#)
- **PARR et al.** Nat. Med., 1997, vol. 3, 1145-9 [\[0036\]](#)
- **HAGENBAUGH et al.** J Exp Med, 1997, vol. 185, 2101-10 [\[0036\]](#)
- **TRATSCHIN et al.** Mol. Cell Biol., 1985, vol. 5, 3251-3260 [\[0037\]](#)
- **GRIMM et al.** Hum. Gene Ther., 1999, vol. 10, 2445-2450 [\[0037\]](#)
- **JENNINGS et al.** Arthritis Res, 2001, vol. 3, 1- [\[0037\]](#)
- **DAVIDSON et al.** Proc. Natl. Acad. Sci. USA, 2000, vol. 97, 3428-3432 [\[0037\]](#)
- **CHLORINI et al.** J. Vir., 1997, vol. 71, 6823-33 [\[0038\]](#)
- **SRIVASTAVA et al.** J. Vir., 1983, vol. 45, 555-64 [\[0038\]](#)
- **CHLORINI et al.** J. Vir, 1999, vol. 73, 1309-1319 [\[0038\]](#)
- **RUTLEDGE et al.** J. Vir., 1998, vol. 72, 309-319 [\[0038\]](#)
- **WU et al.** J. Vir., 2000, vol. 74, 8635-47 [\[0038\]](#)
- **BANTEL-SCHAAL et al.** J. Virol., 1999, vol. 73, 2939-947 [\[0039\]](#)
- **MUYLDERMANS** Biotechnol., 2001, vol. 74, 277-302 [\[0044\]](#)
- **E. HARLOW D. LANE** Using Antibodies Cold Spring Harbor Laboratory Press 19990000 [\[0045\]](#)
- **GRIMM et al.** Novel tools for production and purification of recombinant adeno-associated virus vectors Hum Gene Ther., 1998, vol. 9, 182745-60 [\[0052\]](#)

P a t e n t k r a v

- 5 **1.** Nukleotidsekvens omfattende en enkelt åben læseramme, der koder for parvovirus-Rep78- og -Rep52-proteiner, hvor initiationskodonet til translation af parvovirus-Rep78-proteinet er et suboptimalt initiationskodon sammenlignet med det normale ATG-kodon, hvor nukleotidsekvensen omfatter en ekspressionskontrolsekvens omfattende en sekvens på ni nukleotider af SEQ. ID NO: 7 opstrøms for initiationskodonet af nukleotidsekvensen, der koder for parvovirus-Rep78-protein.
- 10 **2.** Nukleotidsekvens ifølge krav 1, hvor det suboptimale initiationskodon er således, at mindst en del af ribosomerne ikke initierer translation ved det suboptimale kodon af Rep78-proteinet, men ved et initiationskodon længere nedstrøms.
- 15 **3.** Nukleotidsekvens ifølge krav 1 eller 2, hvor initiationskodonet er udvalgt blandt ACG, TTG, CTG og GTG.
- 20 **4.** Nukleotidsekvens ifølge et hvilket som helst af kravene 1 - 3, hvor parvovirus-Rep-proteinerne er adeno-associerede virus (AAV)-Rep-proteiner.
- 25 **5.** Nukleinsyrekonstrukt omfattende en nukleotidsekvens ifølge et hvilket som helst af kravene 1 - 4, hvor nukleotidsekvensen er funktionsmæssigt forbundet med ekspressionskontrolsekvenser til ekspression i en insektcelle, fortrinsvis med en polyeder-promoter, og hvor nukleinsyrekonstruktet fortrinsvis er en insektcellekompatibel vektor, mere fortrinsvis en baculovirusvektor.
- 30 **6.** Insektcelle omfattende en nukleotidsekvens omfattende en enkelt åben læseramme, der koder for parvovirus-Rep78- og Rep52-proteiner, hvor initiationskodonet til translation af parvovirus-Rep78-proteinet er et suboptimalt initiationskodon sammenlignet med det normale ATG-kodon.
- 35 **7.** Insektcelle ifølge krav 6, hvor nukleotidsekvensen koder for fuldlængde-Rep78-proteinet.

- 5 **8.** Insektcelle ifølge krav 6 eller 7, hvor det suboptimale initiationskodon er således, at mindst en del af ribosomerne ikke initierer translation ved det suboptimale kodon af Rep78-proteinet, men ved et initiationskodon længere nedstrøms.
- 10 **9.** Insektcelle ifølge et hvilket som helst af kravene 6 - 8, hvor initiationskodonet er udvalgt blandt ACG, TTG, CTG og GTG.
- 10 **10.** Insektcelle ifølge et hvilket som helst af kravene 6 til 9, hvor nukleotidsekvensen omfatter en ekspressionskontrolsekvens omfattende en sekvens på ni nukleotider af SEQ. ID NO: 7 opstrøms for initiationskodonet af nukleotidsekvensen, der koder for AAV-Rep78-proteinet.
- 15 **11.** Insektcelle ifølge et hvilket som helst af kravene 6 til 10, hvor parvovirus-Rep-proteiner er adeno-associerede virus (AAV)-Rep-proteiner.
- 20 **12.** Insektcelle ifølge et hvilket som helst af kravene 6 til 11, hvor nukleotidsekvensen ifølge et hvilket som helst af kravene 1 - 4 er del af et nukleinsyrekonstrukt og er funktionsmæssigt forbundet med ekspressionskontrolsekvenser til ekspresion i en insektcelle.
- 25 **13.** Insektcelle ifølge et hvilket som helst af kravene 6 til 12, hvor insektcellen yderligere omfatter:
a) en anden nukleotidsekvens omfattende mindst en parvovirus-inverted terminal repeat (ITR)-nukleotidsekvens; og
b) en tredje nukleotidsekvens omfattende parvoviruscapsidprotein-kodende sekvenser, der er funktionsmæssigt forbundet med ekspresionskontrolsekvenser til ekspresion i en insektcelle.
- 30 **14.** Insektcelle ifølge krav 13, hvor insektcellen omfatter:
a) et første nukleotidkonstrukt, hvor nukleotidsekvensen er en del af en nukleinsyrekonstrukt og er funktionsmæssigt forbundet med ekspresionskontrolsekvenser til ekspresion i en insektcelle;
b) er andet nukleinsyrekonstrukt omfattende mindst en parvovirus-inverted
35 terminal repeat (ITR)-nukleotidsekvens; og

c) et tredje nukleotidsyrekonstrukt omfattende parvoviruscapsidprotein-kodende sekvenser, der er funktionsmæssigt forbundet med ekspressionkontrolsekvenser til ekspression i en insektcelle.

5 hvor det første, andet og tredje nukleinsyrekonstrukt fortrinsvis er insektcellekompatible vektorer, fortrinsvis baculovirusvektorer.

10 **15.** Insektcelle ifølge krav 13 eller 14, hvor den anden nukleotidsekvens yderligere omfatter mindst en nukleotidsekvens, der koder for et genprodukt af interesse, og hvor den mindst ene nukleotidsekvens, der koder for et genprodukt af interesse, inkorporeres i genomet af en parvovirusvektor produceret i insektcellen, og fortrinsvis hvor den anden nukleotidsekvens omfatter to parvovirus-ITR-nukleotidsekvenser, og hvor den mindst ene nukleotidsekvens, der koder for et genprodukt af interesse, er lokaliseret mellem de to parvovirus-ITR-nukleotidsekvenser.

15 **16.** Insektcelle ifølge et hvilket som helst af kravene 13 - 15, hvor den tredje nukleotidsekvens omfatter en åben læseramme omfattende nukleotidsekvenser, der koder for parvovirus-VP1-, -VP2- og -VP3-capsidproteiner, hvor initiationskodonet til translation af parvovirus-VP1-capsidproteinet er udvalgt blandt ACG, TTG, CTG og GTG, og hvor den tredje nukleotidsekvens eventuelt omfatter en ekspressionskontrolsekvens omfattende en sekvens på ni nukleotider af SEQ. ID NO: 7 opstrøms for initiationskodonet af nukleotidsekvensen, der koder for parvovirus-VP1-capsidproteinet.

20 **17.** Insektcelle ifølge krav 16, hvor den tredje nukleotidsekvens yderligere omfatter mindst en modificering af nukleotidsekvensen, der koder for parvovirus-VP1-capsidproteinet udvalgt blandt et C i nukleotidposition 12, et A i nukleotidposition 21 og et C i nukleotidposition 24.

30 **18.** Insektcelle ifølge et hvilket som helst af kravene 13 - 17, hvor den mindst ene af den første nukleotidsekvens, anden nukleotidsekvens og tredje nukleotidsekvens er stabilt integreret i insektcellens genom.

35 **19.** Insektcelle et hvilket som helst af kravene 6 - 18, hvor parvoviruset er AAV.

20. Fremgangsmåde til fremstilling af en rekombinant parvovirusvirion, fortrinsvis en AAV-virion, i en insektcelle, hvilken virion omfatter en anden nukleotidsekvens omfattende mindst en parvovirus-inverted terminal repeat (ITR)-nukleotidsekvens, hvilken fremgangsmåde omfatter trinnene:

- 5 a) dyrkning af en insektcelle ifølge et af kravene 13 - 19 under sådanne betingelser, at der fremstilles en rekombinant parvovirusvirion; og
b) genvinding af den rekombinante parvovirusvirion;
og eventuelt yderligere omfattende trinnet:
c) affinitetsoprensning af virionen under anvendelse af et anti-parvovirus-
10 antistof, fortrinsvis et immobiliseret antistof, hvor anti-parvovirus-antistoffet fortrinsvis er et enkeltkæde-camelid-antistof eller et fragment deraf.

- 21.** Fremgangsmåde ifølge krav 20, hvor den anden nukleotidsekvens yderligere omfatter mindst en nukleotidsekvens, der koder for et genprodukt af
15 interesse, og hvor den mindst ene nukleotidsekvens, der koder for et genprodukt af interesse, inkorporeres i genomet af en parvovirusvektor produceret i insektcellen, og fortrinsvis hvor den anden nukleotidsekvens omfatter to parvovirus-ITR-nukleotidsekvenser, og hvor den mindst ene nukleotidsekvens, der koder for et genprodukt af interesse, er lokaliseret mellem de to
20 parvovirus-ITR-nukleotidsekvenser.

- 22.** Nukleotidsekvens ifølge et hvilket som helst af kravene 1 til 4, nukleinsyrekonstrukt ifølge krav 5 eller insektcelle ifølge et hvilket som helst af kravene 6 til 19, hvori et eller flere mulige falske translationsinitiationssteder i de Rep-proteinkodende sekvenser, som ikke er Rep78- og Rep52-
25 translationsinitiationsstederne, er elimineret.

- 23.** Fremgangsmåde ifølge krav 20 eller 21, hvori et eller flere mulige falske translationsinitiationssteder i de Rep-proteinkodende sekvenser, som ikke er
30 Rep78- og Rep52-translationsinitiationsstederne, elimineres.

DRAWINGS

Fig 1






Fig 2

16 hours post infection:



40 hours post infection:



64 hours post infection:




Fig 3




Fig 4





Fig 5

Fig 6

