

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau

(43) International Publication Date
10 June 2021 (10.06.2021)



(10) International Publication Number
WO 2021/113698 A1

(51) International Patent Classification:

C07D 207/16 (2006.01) C07D 413/04 (2006.01)
C07D 403/04 (2006.01) C07D 471/04 (2006.01)

TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW,
KM, ML, MR, NE, SN, TD, TG).

(21) International Application Number:

PCT/US2020/063391

Published:

- with international search report (Art. 21(3))
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))
- with sequence listing part of description (Rule 5.2(a))

(22) International Filing Date:

04 December 2020 (04.12.2020)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/943,629 04 December 2019 (04.12.2019) US

(71) Applicant: **OMEROS CORPORATION** [US/US]; 201
Elliott Avenue West, Seattle, Washington 98119 (US).

(72) Inventors: **CUTSHALL, Neil S.**; c/o 201 Elliott Avenue
West, Seattle, Washington 98119 (US). **GAGE, Jennifer
Lynn**; c/o 201 Elliott Avenue West, Seattle, Washington
98119 (US). **GOLDSTEIN, Sara Rebecca**; c/o 201 Elliott
Avenue West, Seattle, Washington 98119 (US). **KWON,
Do Yeon**; c/o 201 Elliott Avenue, Seattle, Washington
98119 (US). **LITTLE, Thomas L.**; c/o 201 Elliott Avenue
West, Seattle, Washington 98119 (US). **METZ, Markus**;
c/o 201 Elliott Avenue West, Seattle, Washington 98119
(US). **NOLLERT VON SPECHT, Peter Kurt**; c/o 201
Elliott Avenue West, Seattle, Washington 98119 (US).
TSOUNG, Jennifer; c/o 201 Elliott Avenue West, Seattle,
Washington 98119 (US).

(74) Agent: **LIGON, Toby J.** et al.; Seed Intellectual Proper-
ty Law Group LLP, Suite 5400, 701 Fifth Avenue, Seattle,
Washington 98104-7064 (US).

(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ,
CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO,
DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN,
HR, HU, ID, IL, IN, IR, IS, IT, JO, JP, KE, KG, KH, KN,
KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD,
ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO,
NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW,
SA, SC, SD, SE, SG, SK, SL, ST, SV, SY, TH, TJ, TM, TN,
TR, TT, TZ, UA, UG, US, UZ, VC, VN, WS, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ,
UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ,
TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK,
EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV,
MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM,

(54) Title: MASP-2 INHIBITORS AND METHODS OF USE

(57) Abstract: The present disclosure provides, inter alia, compounds with MASP-2 inhibitory activity, compositions of such com-
pounds, and methods of making and using such compounds.

WO 2021/113698 A1

MASP-2 INHIBITORS AND METHODS OF USE

STATEMENT REGARDING SEQUENCE LISTING

The Sequence Listing associated with this application is provided in text format in lieu of a paper copy, and is hereby incorporated by reference into the specification. The name of the text file containing the Sequence Listing is 700128_421WO_SEQUENCE_LISTING.txt. The text file is 6.0 KB, was created on December 3, 2020, and is being submitted electronically via EFS-Web.

TECHNICAL FIELD

The present disclosure is directed generally to compositions and methods that are useful in the field of medicine. More specifically, the disclosure provides synthetic inhibitors of mannan-binding lectin-associated serine protease-2 (MASP-2), including inhibitors that selectively inhibit MASP-2 over thrombin, compositions thereof, and methods for the manufacture and use thereof.

BACKGROUND


The complement system plays a role in the inflammatory response and becomes activated because of tissue damage or microbial infection. Complement activation must be tightly regulated to ensure selective targeting of invading microorganisms and avoid self-inflicted damage (Ricklin et al., Nat. Immunol. 11 :785-797, 2010). Currently, it is widely accepted that the complement system can be activated through three distinct pathways: the classical pathway, the lectin pathway, and the alternative pathway. The classical pathway is usually triggered by a complex composed of host antibodies bound to a foreign particle (*i.e.*, an antigen) and generally requires prior exposure to an antigen for the generation of a specific antibody response. Since activation of the classical pathway depends on a prior adaptive immune response by the host, the classical pathway is part of the acquired immune system. In contrast, both the lectin and alternative pathways are independent of adaptive immunity and are part of the innate immune system.

Mannan-binding lectin-associated serine protease-2 (MASP-2) has been shown to be required for the function of the lectin pathway, one of the principal complement activation pathways (Vorup-Jensen et al., J. Immunol 165:2093-2100, 2000; Ambrus et al., J Immunol. 170: 1374-1382, 2003; Schwaeble et al., PNAS 108:7523-7528, 2011). Importantly, inhibition of MASP-2 does not appear to interfere with the antibody-dependent classical complement activation pathway, which is a critical component of the acquired immune response to infection. As described in U.S. Patent No. 9,011,860 (assigned to Omeros Corporation), which is hereby incorporated by reference, discloses a fully human monoclonal antibody targeting human MASP-2 has been generated which binds to human MASP-2 with high affinity and blocks the lectin pathway complement activity and is therefore useful to treat various lectin complement pathway-associated diseases and disorders.

MASP-2-dependent complement activation has been implicated as contributing to the pathogenesis of numerous acute and chronic disease states. Therefore, a need exists for compounds that are suitable for administration/treatment of subject suffering from MASP-2 complement pathway-associated diseases and disorders, including diseases that are not suitably or efficiently treated with large molecule biologic inhibitors.


BRIEF SUMMARY

One embodiment provides a compound having the following Structure (I):



or a stereoisomer, tautomer, or pharmaceutically acceptable salt thereof, wherein R^1 , R^{2a} , R^{2b} , R^{2c} , R^{2d} , R^3 , R^4 , and n are as defined herein.


Another embodiment provides a compound having the following Structure (II):



(II)

5 or a stereoisomer, tautomer, or pharmaceutically acceptable salt thereof, wherein R^1 , R^{2a} , R^{2b} , R^{2c} , R^{2d} , R^3 , R^4 , and n are as defined herein.

Another embodiment provides a compound having the following Structure (III):



(III)

10 or a stereoisomer, tautomer, or pharmaceutically acceptable salt thereof, wherein R^{25} , R^{26a} , R^{26b} , R^{27} , R^{28} , and p are as defined herein.

Additional embodiments of the present disclosure provide a pharmaceutical composition comprising a compound of Structure (I), (II) or (III), or a stereoisomer, tautomer, or pharmaceutically acceptable salt thereof, and at least one pharmaceutically acceptable carrier or excipient.

The compounds of Structures (I), (II) and (III) are useful in the treatment of MASP-2-associated diseases and disorders, and in the manufacture of medicaments for treating MASP-2-associated diseases and disorders. Accordingly, other embodiments of the disclosure provide methods of treating a MASP-2-associated disease and disorder comprising administering to a patient a therapeutically effective amount of a compound

of Structure (I), (II) or (III), or a stereoisomer, tautomer, or pharmaceutically acceptable salt thereof.

These and other aspects, objects and embodiments will become more apparent when read with the detailed description and figures which follow.

5 DETAILED DESCRIPTION

I. Definitions

Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this disclosure belongs. Although methods and materials similar or equivalent to those described herein can be used in the practice or testing of the subject matter of the present disclosure, suitable methods and materials are described below. In addition, the materials, methods, and examples are illustrative only and not intended to be limiting.

In certain embodiments herein, reference is made to features and aspects of the disclosure, including method steps. All possible combinations of such features and aspects within the embodiments of the disclosure are included, at least to the extent that such combinations are non-contradictory. For example, if an embodiment presents aspects A, B, and C, it is understood that this also discloses embodiments including both aspects A and B, both aspects B and C, and both aspects A and C, as well as an embodiment with aspects A, B, and C.

The terms "a," "an," or "the" not only include aspects with one member, but also include aspects with more than one member. For instance, the singular forms "a," "an," and "the" include plural referents unless the context clearly dictates otherwise. Thus, for example, reference to "a cell" includes a plurality of such cells and reference to "the agent" includes reference to one or more agents known to those skilled in the art.

The terms "about" and "approximately" refer to an acceptable degree of error for the quantity measured given the nature or precision of the measurements. Typical, exemplary degrees of error are within ± 20 percent (%); preferably, within $\pm 10\%$; and more preferably, within $\pm 5\%$ of a given value or range of values. Any reference to "about X" specifically indicates at least the values X, 0.95X, 0.96X, 0.97X,

0.98X, 0.99X, 1.01X, 1.02X, 1.03X, 1.04X, and 1.05X. Thus, "about X" is intended to teach and provide written support for a claim limitation of, *e.g.*, "0.98X." Alternatively, in biological systems, the terms "about" and "approximately" may mean values that are within an order of magnitude, preferably within 5-fold, and more preferably within 2-fold of a given value. Numerical quantities given herein are approximate unless stated otherwise, meaning that the term "about" or "approximately" can be inferred when not expressly stated. When "about" is applied to the beginning of a numerical range, it applies to both ends of the range. Thus, "from about 5 to 20%" is equivalent to "from about 5% to about 20%." When "about" is applied to the first value of a set of values, it applies to all values in that set. Thus, "about 7, 9, or 11 mg/kg" is equivalent to "about 7, about 9, or about 11 mg/kg."

The term "MASP-2" refers to mannan-binding lectin-associated serine protease-2. Human MASP-2 protein with UniProt accession code O00187 (SEQ ID NO:1). The Serine Protease Domain ('B-chain' = Mannan-binding lectin serine protease 2 B chain, based on UniProtKB - O00187 (MASP-2_HUMAN)) includes residues 445 to 686 (or consists of residues 445 to 686).

The term "MASP-2-dependent complement activation" refers to MASP-2-dependent activation of the lectin pathway, which occurs under physiological conditions (*i.e.*, in the presence of Ca^{++}) leading to the formation of the lectin pathway C3 convertase C4b2a and upon accumulation of the C3 cleavage product C3b subsequently to the C5 convertase C4b2a(C3b)_n.

The term "MASP-2-dependent complement-associated disease or disorder" refers to a disease or disorder that is associated with MASP-2-dependent complement activation.

The term "MASP-2-associated disease or disorder" refers to a disease or disorder that is associated with activation or activity of MASP-2, including MASP-2-dependent complement-associated disease or disorders, and wherein inhibition of MASP-2 is or is expected to be therapeutically beneficial.

The term "lectin pathway" refers to complement activation that occurs via the specific binding of serum and non-serum carbohydrate-binding proteins including mannan-binding lectin (MBL), CL-11 and the ficolins (H-ficolin, M-ficolin, or L-ficolin).

The term "classical pathway" refers to complement activation that is
5 triggered by an antibody bound to a foreign particle and requires binding of the recognition molecule Clq.

Amino acid residues are abbreviated as follows: alanine (Ala; A), asparagine (Asn; N), aspartic acid (Asp; D), arginine (Arg; R), cysteine (Cys; C), glutamic acid (Glu; E), glutamine (Gln; Q), glycine (Gly; G), histidine (His; H),
10 isoleucine (Ile), leucine (Leu), lysine (Lys; K), methionine (Met; M), phenylalanine (Phe; F), proline (Pro; P), serine (Ser; S), threonine (Thr; T), tryptophan (Trp; W), tyrosine (Tyr; Y), and valine (Val; V).

In the broadest sense, the naturally occurring amino acids can be divided into groups based upon the chemical characteristic of the side chain of the respective
15 amino acids. By "hydrophobic" amino acid is meant either His, Leu, Met, Phe, Trp, Tyr, Val, Ala, Cys or Pro. By "hydrophilic" amino acid is meant either Gly, Asn, Gln, Ser, Thr, Asp, Glu, Lys, Arg or His. This grouping of amino acids can be further sub-classed as follows: by "uncharged hydrophilic" amino acid is meant either Ser, Thr, Asn or Gln. By "acidic" amino acid is meant either Glu or Asp. By "basic" amino acid is meant either
20 Lys, Arg or His.

The term "conservative amino acid substitution" is illustrated by a substitution among amino acids within each of the following groups:

- (1) glycine, alanine, valine, leucine, and isoleucine;
- (2) phenylalanine, tyrosine, and tryptophan;
- 25 (3) serine and threonine;
- (4) aspartate and glutamate;
- (5) glutamine and asparagine; and
- (6) lysine, arginine and histidine.

The term "a subject" includes all mammals, including without limitation, humans, non-human primates, dogs, cats, horses, sheep, goats, cows, rabbits, pigs, and rodents.

"Mammal" includes humans and both domestic animals such as laboratory
5 animals and household pets, (*e.g.*, cats, dogs, swine, cattle, sheep, goats, horses, rabbits), and non-domestic animals such as wildlife and the like.

The terms "small molecule" and "small organic molecule" refers to a small carbon-containing molecule that has a molecular weight of about 2500 daltons or lower. In some embodiments, a small molecule has a molecular weight of about 2000 daltons or
10 lower. In some embodiments, a small molecule has a molecular weight of about 1500 daltons or lower. In some embodiments, a small molecule has a molecular weight of about 1000 daltons or lower. In some embodiments, a small molecule has a molecular weight of about 750 daltons or lower. In some embodiments, a small molecule has a molecular weight of about 500 daltons or lower. In some embodiments, a small molecule has a
15 molecular weight of about 50 daltons or greater. In some embodiments, a small molecule has a molecular weight of about 75 daltons or greater. In some embodiments, a small molecule has a molecular weight of about 100 daltons or greater. In some embodiments, a small molecule has a molecular weight of about 150 daltons or greater. In some embodiments, a small molecule has a molecular weight of about 250 daltons or greater.
20 In some embodiments, small molecules may have a molecular weight in the range from about 50 daltons to about 500 daltons, from about 50 daltons to about 750 daltons, from about 50 daltons to about 1000 daltons, from about 50 daltons to about 1500 daltons, from about 50 daltons to about 2000 daltons, or from about 50 daltons to about 2500 daltons. When the term "compound" is used herein, the term is explicitly intended to
25 include small molecule compounds as defined herein (including any of the embodiments thereof).

As used herein, the terms "disease" and "condition" may be used interchangeably or may be different in that the particular malady or condition may not have a known causative agent (so that etiology has not yet been worked out) and it is
30 therefore not yet recognized as a disease but only as an undesirable condition or

syndrome, wherein a more or less specific set of symptoms have been identified by clinicians. In some embodiments, a disease is a pathological condition of an organ, a body part, or a system resulting from various causes, such as infection, genetic defect, or environmental stress that is characterized by an identifiable group of symptoms.

5 "Therapeutically effective amount," "effective amount," or "effective dose" refers to that amount of a compound of the disclosure that, when administered to a mammal (*e.g.*, a human), is sufficient to effect treatment, as defined below, of a disease or condition in the mammal, preferably a human. The amount of a compound of the disclosure which constitutes a "therapeutically effective amount" will vary depending on
10 the compound, the condition and its severity, the manner of administration, and the age of the mammal to be treated, but can be determined routinely by one of ordinary skill in the art having regard to his own knowledge and to this disclosure.

The term "subcutaneous administration" refers to administration of a formulation under all layers of the skin of a subject.

15 The term "histidine" specifically includes L-histidine unless otherwise specified.

The term "isotonic" refers to a formulation that has essentially the same osmotic pressure as human blood. Isotonic formulations will generally have an osmotic pressure from about 250 to about 350 mOsm/L. Isotonicity can be measured using a
20 vapor pressure or freezing point depression osmometer, for example.

The term "hypertonic" refers to a formulation with an osmotic pressure above that of human (*i.e.*, greater than 350 mOsm/L).

The term "hydrogen-bonding" is a partially electrostatic attraction between a hydrogen (H) which is bound to a more electronegative atom such as nitrogen
25 (N) or oxygen (O) and another adjacent atom bearing a lone pair of electrons. For example, when it is stated that the nitrogen acts as a "hydrogen bond donor" it means that a hydrogen (H) bound to a nitrogen (N) is donated by the nitrogen as it electrostatically attracted to or accepted by an adjacent atom bearing a lone pair of electrons such as an oxygen. Similarly, when it is stated that an oxygen acts as a "hydrogen bond acceptor,"
30 it means that a hydrogen (H) bound to a more electronegative atom such as nitrogen (N)

is electrostatically attracted to or "accepted by" an adjacent atom such as oxygen bearing a lone pair of electrons. Sometimes the hydrogen bonded atoms are called out without explicitly stating the origin and presence of an intermediate hydrogen atom. The term "hydrogen bonding" is used wherever LigPlot+ software predicts a hydrogen bonding
5 interaction using its algorithm and applied parameters of 3.35 Å for maximum distance between hydrogen bond donor and acceptor. Not all hydrogen bonds may actually be in place simultaneously; this is evident for atoms that are shown to form 4 putative hydrogen bonds, where however, at any given time only 3 hydrogen bonds are chemically possible. In general, although crystal structures such as the co-crystal structural information herein
10 does not directly show or detect hydrogen bonding, the software used to describe the co-crystal does predict such H-bonding exists. Therefore, throughout the disclosure when a H-bond is present and described, it may be said to be "predicted" by software to be present.

The term ionic bonding includes a type of chemical bond that involves the
15 electrostatic attraction between oppositely charged ions, and is the primary interaction occurring in ionic compounds.

The term "van der Waals" interaction includes weak, short-range electrostatic attractive forces between uncharged molecules, arising from the interaction of permanent or transient electric dipole moments. As determined by LigPlot+ software
20 employing models derived from the corresponding crystallographic MASP-2 compound co-structures, such interactions include all contacts that are computed using non-bonded contact parameters between hydrophobic to any contacts for interactions with a maximum contact distance of 3.90 Å.

The term " π - π interaction" or " π - π stacking" interaction includes
25 attractive, non-covalent interactions between aromatic rings that are oriented either roughly parallel or roughly perpendicular (such as in "edge-face" interactions) to each other, since they contain π -bonds.

Typically, the active site of serine proteases such as MASP-2 is shaped as a cleft where the polypeptide substrate or inhibitor binds. Schechter and Berger labeled
30 amino acid residues from the N to C terminus of the polypeptide substrate as follows:

Pi,..., P3, P2, P1, P1', P2', P3',..., Pj) and their respective binding sub-sites Si,..., S3, S2, S1, S1', S2', S3',..., Sj. The cleavage is catalyzed between P1 and P1' (Schechter, I. & Berger, A. On the size of the active site in proteases. I. Papain. Biochem. Biophys. Res. Commun. 27 (1967)).

5 The term "binding site" is an area on the protein wherein a small molecule can interact with, such as a region on the surface of MASP-2. The binding site or region may not or only partially overlap with the active site, but nevertheless render the MASP-2 molecule less active or inactive.

 The term "or" refers to an alternative and should in general be construed
10 non-exclusively. For example, a claim to "a composition comprising A or B" would typically present an aspect with a composition comprising both A and B. "Or" should, however, be construed to exclude those aspects presented that cannot be combined without contradiction (*e.g.*, a composition pH that is between 9 and 10 or between 7 and 8).

15 The group "A or B" is equivalent to the group "selected from the group consisting of A and B."

 The linking term "comprising" or "comprise" is not closed. For example, "a composition comprising A" must include at least the component A, but it may also include one or more other components (*e.g.*, B; B and C; B, C, and D; and the like). The
20 term "comprising" therefore should in general be construed as not excluding additional ingredients. For example, a claim to "a composition comprising A" would cover compositions that include A and B; A, B, and C; A, B, C, and D; A, B, C, D, and E; and the like.

 The term "hypertonic" refers to a formulation with an osmotic pressure
25 above that of human (*i.e.*, greater than 350 mOsm/KgH₂O).

 The term "agent" refers to a compound or mixture of compounds that, when added to a composition, tend to produce an effect on the composition's properties. For example, a composition comprising a thickening agent is likely to be more viscous than an otherwise identical comparative composition that lacks the thickening agent.

A "synthetic" compound means a compound that is not naturally occurring and that has been synthesized by humans. Reference to a compound herein may be understood to include reference to synthetic compounds, unless the context indicates otherwise.

5 The expressions, "ambient temperature" and "room temperature," as used herein, are understood in the art, and refer generally to a temperature, *e.g.*, a reaction temperature, that is about the temperature of the room in which the reaction is carried out, *e.g.*, a temperature from about 20 °C to about 30 °C.

 At various places in the present specification, certain features of the
10 compounds are disclosed in groups or in ranges. It is specifically intended that such a disclosure include each and every individual sub-combination of the members of such groups and ranges. For example, the terms "C₁₋₆ alkyl" and "C₁-C₆ alkyl" are specifically intended to individually disclose (without limitation) methyl, ethyl, C₃ alkyl, C₄ alkyl, C₅ alkyl, and C₆ alkyl.

15 The term "substituted" means that an atom or group of atoms formally replaces hydrogen as a "substituent" attached to another group. The term "substituted", unless otherwise indicated, refers to any level of substitution, *e.g.*, mono-, di-, tri-, tetra-, penta-, or higher substitution, where such substitution is permitted (*e.g.*, results in a stable compound). The substituents are independently selected, and substitution may be at any
20 chemically accessible position. It is to be understood that substitution at a given atom is limited by valency. The phrase "optionally substituted" means substituted or unsubstituted. The term "substituted" means that at least hydrogen atom is replaced with a substituent. A single divalent substituent, *e.g.*, oxo, can replace two hydrogen atoms.

 The terms "C_{n-m}" and "C_n-C_m" where n and m are integers indicates a
25 group that contains from n to m carbon atoms. Examples include C₁₋₄, C₁₋₆, and the like. The term is intended to expressly disclose every member in the range, *i.e.*, C_n, C_{n+1}, C_{n+2} ... C_{m-2}, C_{m-1}, C_m. For example, C₁₋₆ is intended to disclose C₁, C₂, C₃, C₄, C₅, and C₆. As used herein, "C_{n-m}" means the same as "C_n-C_m".

 The term "n-membered," where n is an integer (*e.g.*, 6-membered),
30 typically describes the number of ring-forming atoms in a moiety where the number of

ring-forming atoms is n . The term "n-m membered" wherein n and m are integers (*e.g.*, 6-10 membered) describes a range where the number of ring forming atoms is from n to m . For example, piperidinyl is an example of a 6-membered heterocyclyl ring, pyrazolyl is an example of a 5-membered heteroaryl ring, pyridyl is an example of a 6-membered heteroaryl ring and 1,2,3,4-tetrahydro-naphthalene is an example of a 10-membered cycloalkyl group.

"Alkyl" refers to a straight or branched hydrocarbon chain radical consisting solely of carbon and hydrogen atoms, containing no unsaturation, having from one to twelve carbon atoms, preferably one to eight carbon atoms, more preferably one to six carbon atoms, which is attached to the rest of the molecule by a single bond, *e.g.*, methyl, ethyl, *n*-propyl, 1-methylethyl (*iso*-propyl), *n*-butyl, *n*-pentyl, 1,1-dimethylethyl (*t*-butyl), 3-methylhexyl, 2-methylhexyl, and the like. In certain specific embodiments, an alkyl group may be optionally substituted by one of the following groups: alkyl, alkenyl, halo, haloalkenyl, cyano, nitro, aryl, cycloalkyl, heterocyclyl, heteroaryl, oxo, trimethylsilanyl, $-OR^{100}$, $-OC(O)R^{100}$, $-N(R^{100})_2$, $-C(O)R^{100}$, $-C(O)OR^{100}$, $-C(O)N(R^{100})_2$, $-N(R^{20})C(O)OR^{102}$, $-N(R^{100})C(O)R^{102}$, $-N(R^{102})S(O)_pR^{102}$ (where p is 1 to 2), $-S(O)_pOR^{102}$ (where p is 1 to 2), $-S(O)_tR^{102}$ (where t is 0 to 2), and $-S(O)_pN(R^{100})_2$ (where p is 1 to 2) where each R^{100} is independently hydrogen, alkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, aryl, arylalkyl, heterocyclyl, heterocyclylalkyl, heteroaryl or heteroarylalkyl; and each R^{102} is alkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, aryl, arylalkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, or heteroarylalkyl. In some embodiments, alkyl is a C_1 - C_{12} alkyl, a C_1 - C_8 alkyl, a C_1 - C_6 alkyl, or C_1 - C_4 alkyl.

"Alkenyl" refers to a straight or branched hydrocarbon chain radical group consisting solely of carbon and hydrogen atoms, containing at least one double bond, having from two to twelve carbon atoms, preferably two to eight carbon atoms and which is attached to the rest of the molecule by a single bond, *e.g.*, ethenyl, prop-1-enyl, but-1-enyl, pent-1-enyl, penta-1,4-dienyl, and the like. In certain embodiments, an alkyl or alkenyl group may be optionally substituted by one of the following groups: alkyl, alkenyl, halo, haloalkenyl, cyano, nitro, aryl, cycloalkyl, heterocyclyl, heteroaryl, oxo, trimethylsilanyl, $-OR^{100}$, $-OC(O)R^{100}$, $-N(R^{100})_2$, $-C(O)R^{100}$, $-C(O)OR^{100}$, $-C(O)N(R^{100})_2$,

-N(R²⁰)C(O)OR¹⁰², -N(R¹⁰⁰)C(O)R¹⁰², -N(R¹⁰²)S(O)_pR¹⁰² (where p is 1 to 2), -S(O)_pOR¹⁰² (where p is 1 to 2), -S(O)_tR¹⁰² (where t is 0 to 2), and -S(O)_pN(R¹⁰⁰)₂ (where p is 1 to 2) where each R¹⁰⁰ is independently hydrogen, alkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, aryl, arylalkyl, heterocyclyl, heterocyclalkyl, heteroaryl or
 5 heteroarylalkyl; and each R¹⁰² is alkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, aryl, arylalkyl, heterocyclyl, heterocyclalkyl, heteroaryl, or heteroarylalkyl.

"Alkynyl" refers to a straight-chain or branched hydrocarbon group corresponding to an alkyl group as defined above having one or more triple carbon-carbon bonds. An alkynyl group formally corresponds to an alkyne with one C-H bond
 10 replaced by the point of attachment of the alkyl group to the remainder of the compound. The term "C_{n-m} alkynyl" and "C_n-C_m alkynyl" refer to an alkynyl group having n to m carbons. Example alkynyl groups include, but are not limited to, ethynyl, propyn-1-yl, propyn-2-yl and the like. In some embodiments, the alkynyl moiety contains 2 to 6, 2 to 4, or 2 to 3 carbon atoms. Unless indicated otherwise, alkynyl groups are optionally
 15 substituted.

"Alkylene" or "alkylene chain" refers to a straight or branched divalent hydrocarbon chain linking the rest of the molecule to a radical group or linking two parts of the molecule, consisting solely of carbon and hydrogen, containing no unsaturation and having from one to twelve carbon atoms, *e.g.*, methylene, ethylene, propylene, n-
 20 butylene, and the like. The alkylene chain may optionally contain one or more heteroatoms wherein a carbon in the alkylene chain is replaced with a heteroatom selected from oxygen, nitrogen or sulfur. The alkylene chain is attached to the rest of the molecule through a single bond and to the radical group through a single bond or is attached to two parts of the molecule through a single bond at each point of attachment. In some
 25 embodiments, an alkyl group may be optionally substituted by one of the following groups: alkyl, alkenyl, halo, haloalkenyl, cyano, nitro, aryl, cycloalkyl, heterocyclyl, heteroaryl, oxo, trimethylsilanyl, -OR¹⁰⁰, -OC(O)R¹⁰⁰, -N(R¹⁰⁰)₂, -C(O)R¹⁰⁰, -C(O)OR¹⁰⁰, -C(O)N(R¹⁰⁰)₂, -N(R²⁰)C(O)OR¹⁰², -N(R¹⁰⁰)C(O)R¹⁰², -N(R¹⁰²)S(O)_pR¹⁰² (where p is 1 to 2), -S(O)_pOR¹⁰² (where p is 1 to 2), -S(O)_tR¹⁰² (where t is 0 to 2), and -S(O)_pN(R¹⁰⁰)₂
 30 (where p is 1 to 2) where each R¹⁰⁰ is independently hydrogen, alkyl, haloalkyl,

cycloalkyl, cycloalkylalkyl, aryl, arylalkyl, heterocyclyl, heterocyclalkyl, heteroaryl or heteroarylalkyl; and each R^{102} is alkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, aryl, arylalkyl, heterocyclyl, heterocyclalkyl, heteroaryl, or heteroarylalkyl.


The term "hydroxyalkyl" refers to an alkyl group as defined above in which one or more of the hydrogen atoms has been replaced by a hydroxy group (*i.e.*, -OH). The term " C_{n-m} hydroxyalkyl" refers to a C_{n-m} alkyl group having n to m carbon atoms and from at least one hydroxy group. In some embodiments, the hydroxyalkyl group comprises one hydroxy group. In certain aspects, the hydroxyalkyl group comprises two or more hydroxy groups (*e.g.*, a "dihydroxyalkyl"), each on the same or a different carbon atom(s). In certain aspects, the hydroxyalkyl group has 1, 2, 3, 4, 5, 6, or more hydroxy groups. Examples may include, but are not limited to, hydroxymethyl, 2-hydroxyethyl, and 1-hydroxyethyl. In some embodiments, hydroxyalkyl is a C_1 - C_{12} hydroxyalkyl, a C_1 - C_8 hydroxyalkyl, a C_1 - C_6 hydroxyalkyl, or C_1 - C_4 hydroxyalkyl.

"Aminylalkyl" refers to an alkyl group as defined above in which one or more hydrogen atoms have been replaced by an aminyl group (*i.e.*, $-NR^{100}R^{101}$ wherein R^{100} and R^{101} are each independently hydrogen, alkyl, alkenyl, or alkynyl as defined herein). In some embodiments, the aminylalkyl comprises one aminyl group. In some embodiments, the aminyl group is $-NH_2$.

"Carboxyalkyl" refers to an alkyl group as defined above in which one or more hydrogen atoms have been replaced by a carboxy group (*i.e.*, $-C(O)OH$). In some embodiments, the carboxyalkyl comprises one carboxy group. In some embodiments, carboxyalkyl is a C_1 - C_{12} carboxyalkyl, a C_1 - C_8 carboxyalkyl, a C_1 - C_6 carboxyalkyl, or C_1 - C_4 carboxyalkyl.

"Aryl" refers to a hydrocarbon ring system radical comprising hydrogen, 6 to 18 carbon atoms and at least one aromatic ring. For purposes of this disclosure, the aryl radical may be a monocyclic, bicyclic, tricyclic or tetracyclic ring system, which may include fused or bridged ring systems. Aryl radicals include, but are not limited to, aryl radicals derived from aceanthrylene, acenaphthylene, acephenanthrylene, anthracene, azulene, benzene, chrysene, fluoranthene, fluorene, *as*-indacene, *s*-indacene, indane, indene, naphthalene, phenalene, phenanthrene, pleiadene, pyrene, and

triphenylene. In some embodiments, an aryl group may be optionally substituted by one or more substituents independently selected from the group consisting of alkyl, alkenyl, halo, haloalkyl, haloalkenyl, cyano, nitro, aryl, arylalkyl, cycloalkyl, cycloalkylalkyl, heterocyclyl, heterocyclalkyl, heteroaryl, heteroarylalkyl, $-R^{101}-OR^{100}$, $-R^{101}OC(O)R^{100}$, $-R^{101}-N(R^{100})_2$, $-R^{101}-N(R^{100})-R^{103}-OR^{100}$, $-R^{101}-C(O)R^{100}$, $-R^{101}-C(O)OR^{100}$, $-R^{101}-C(O)N(R^{100})_2$, $-R^{101}-N(R^{100})C(O)OR^{102}$, $-R^{101}-N(R^{100})C(O)R^{102}$, $-R^{101}-N(R^{100})S(O)_pR^{102}$ (where p is 1 to 2), $-R^{101}-N=C(OR^{100})R^{100}$, $-R^{101}-S(O)_pOR^{102}$ (where p is 1 to 2), $-R^{101}-S(O)_tR^{102}$ (where t is 0 to 2), and $-R^{101}-S(O)_pN(R^{100})_2$ (where p is 1 to 2) where each R^{100} is independently hydrogen, alkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, aryl, arylalkyl, heterocyclyl, heterocyclalkyl, heteroaryl or heteroarylalkyl; each R^{101} is independently a direct bond or a straight or branched alkylene chain; each R^{102} is alkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, aryl, arylalkyl, heterocyclyl, heterocyclalkyl, heteroaryl or heteroarylalkyl, and each R^{103} is a direct bond or a straight or branched alkylene chain. In some embodiments an aryl group has the following structure:



"Arylalkyl" or "aralkyl" refers to a group of formula -alkylene-aryl wherein the alkylene group and aryl groups are as defined herein, respectively. In some embodiments, arylalkyl is C_{6-10} aryl- C_{1-3} alkyl. In some embodiments, arylalkyl is C_{6-10} aryl- C_{1-4} alkyl. In some embodiments, arylalkyl is C_{6-10} aryl- C_{1-3} alkyl. In some embodiments, arylalkyl is phenyl- C_{1-3} alkyl. Examples include, but are not limited to, benzyl, 1-phenylethyl, 4-methylbenzyl, and 1,1,-dimethyl-1-phenylmethyl. In some embodiments, arylalkyl is optionally substituted benzyl.

"Aryloxy" refers to a group with the formula -O-aryl wherein aryl is a group as defined above. In some embodiments, the aryloxy group is -O- C_{6-10} aryl. In some embodiments, the aryloxy is a substituted or unsubstituted phenyloxy (*i.e.*, -O- C_6 aryl).

"Arylalkoxy" refers to a group with the formula -alkoxy-aryl wherein alkoxy and aryl are groups as defined above, respectively. In some embodiments,

arylalkoxy is C₆₋₁₀ aryl-C₁₋₃ alkoxy. In some embodiments, arylalkoxy is C₆₋₁₀ aryl-C₁₋₄ alkoxy. In some embodiments, arylalkoxy is C₆₋₁₀ aryl-C₁₋₃ alkoxy. In some embodiments, arylalkoxy is phenyl-C₁₋₃ alkoxy (*e.g.*, methoxy).

"Cycloalkyl" refers to a stable non-aromatic monocyclic or polycyclic hydrocarbon radical consisting solely of carbon and hydrogen atoms, which may include fused or bridged ring systems, having from three to fifteen carbon atoms, preferably having from three to ten carbon atoms, and which is saturated or unsaturated and attached to the rest of the molecule by a single bond. Monocyclic radicals include, for example, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, and cyclooctyl. Polycyclic radicals include, for example, adamantyl, norbornyl, decaliny, and the like. In some embodiments, a cycloalkyl group may be optionally substituted by one or more substituents independently selected from the group consisting of alkyl, alkenyl, halo, haloalkyl, haloalkenyl, cyano, nitro, oxo, aryl, arylalkyl, cycloalkyl, cycloalkylalkyl, heterocyclyl, heterocyclalkyl, heteroaryl, heteroarylalkyl, -R¹⁰¹-OR¹⁰⁰, -R¹⁰¹-OC(O)-R¹⁰⁰, -R¹⁰¹-N(R¹⁰⁰)-R¹⁰³-OR¹⁰⁰, -R¹⁰¹-N(R¹⁰⁰)₂, -R¹⁰¹-C(O)R¹⁰⁰, -R¹⁰¹-C(O)OR¹⁰⁰, -R¹⁰¹-C(O)N(R¹⁰⁰)₂, -R¹⁰¹-N(R¹⁰⁰)C(O)OR¹⁰², -R¹⁰¹-N(R¹⁰⁰)C(O)R¹⁰², -R¹⁰¹-N(R¹⁰⁰)S(O)_pR¹⁰² (where p is 1 to 2), -R¹⁰¹-N=C(OR¹⁰⁰)R¹⁰⁰, -R¹⁰¹-S(O)_pOR¹⁰² (where p is 1 to 2), -R¹⁰¹-S(O)_tR¹⁰² (where t is 0 to 2), and -R¹⁰¹-S(O)_pN(R¹⁰⁰)₂ (where p is 1 to 2) where each R¹⁰⁰ is independently hydrogen, alkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, aryl, arylalkyl, heterocyclyl, heterocyclalkyl, heteroaryl or heteroarylalkyl; each R¹⁰¹ is independently a direct bond or a straight or branched alkylene chain; each R¹⁰² is alkyl, haloalkyl, cycloalkyl, cycloalkylalkyl, aryl, arylalkyl, heterocyclyl, heterocyclalkyl, heteroaryl or heteroarylalkyl, and each R¹⁰³ is a direct bond or a straight or branched alkylene chain.

"Cycloalkylalkyl" refers to a radical of the formula -R¹⁰⁰R¹⁰¹ where R¹⁰⁰ is an alkylene chain as defined above and R¹⁰¹ is a cycloalkyl radical as defined above. When specifically stated in the specification, the alkylene chain and/or the cycloalkyl radical may be optionally substituted as defined above for optionally substituted alkylene chain and optionally substituted cycloalkyl.

"Alkoxy" refers to a radical group having the following formula "-O-alkyl," wherein the alkyl group is as defined herein above. Example alkoxy groups

include methoxy, ethoxy, propoxy (*e.g.*, *n*-propoxy and isopropoxy), *t*-butoxy, and the like. In some embodiments, the alkyl group has 1 to 6, 1 to 4, or 1 to 3 carbon atoms. Unless indicated otherwise, alkoxy groups are optionally substituted.

"Alkoxyalkyl" refers to a radical having the following formula "-alkylene-O-alkyl," wherein the alkylene and alkyl groups are as defined herein above, respectively. In some embodiments, the alkoxyalkyl group comprises one -O-alkyl group. In some embodiments, the alkoxyalkyl group comprises two or more alkoxy groups. Examples may include, but are not limited to, methoxymethyl, ethoxymethyl, 3-ethoxyethyl, and 1-methoxyethyl. Unless indicated otherwise, alkoxyalkyl groups are optionally substituted.

"Oxo" refers to a =O group. For example, an oxo connected to a carbon atom forms a carbonyl group (*i.e.*, C=O). Alternatively, when an oxo group is attached to a heteroatom, for example, a sulfoxide, sulfone group, an *N*-oxide group is formed.

"Sulfido" refers to a =S group.

"Amino" refers to a -NH₂ group.

"Carbamyl" refers to a -C(O)NH₂ group.

"Carboxy" refers to a -C(O)OH group.

"Carbonyl" refers to a C(=O) group, which also may be written as C(O).

"Cyano" or "nitrile" refers to a -C≡N group, which also may be written as -CN.

"Nitro" refers to a -NO₂ group.

"Hydroxy" or "hydroxyl" refers to an -OH group.

"Halo" or "halogen" refers to bromo, chloro, fluoro or iodo.

"Haloalkyl" refers to an alkyl radical, as defined above, that is substituted by one or more halo radicals, as defined above, *e.g.*, trifluoromethyl, difluoromethyl, trichloromethyl, 2,2,2-trifluoroethyl, 1-fluoromethyl-2-fluoroethyl, 3-bromo-2-fluoropropyl, 1-bromomethyl-2-bromoethyl, and the like. The alkyl part of the haloalkyl radical may be optionally substituted as defined above for an alkyl group.

The term "haloalkoxy", employed alone or in combination with other terms, refers to a group of formula -O-haloalkyl, wherein the haloalkyl group is as defined

above. Example haloalkoxy groups include trifluoromethoxy, difluoromethoxy, pentafluoroethoxy, and the like.


"Heterocyclyl" refers to a stable 3- to 18-membered non-aromatic ring radical which consists of two to twelve carbon atoms and from one to six heteroatoms selected from the group consisting of nitrogen, oxygen, and sulfur. Unless stated otherwise specifically in the specification, the heterocyclyl radical may be a monocyclic, bicyclic, tricyclic, or tetracyclic ring system, which may include fused, bridged, and spiro ring systems; and the nitrogen, carbon, or sulfur atoms in the heterocyclyl radical may be optionally oxidized; the nitrogen atom may be optionally quaternized; and the heterocyclyl radical may be partially or fully saturated. Examples of such heterocyclyl radicals include, but are not limited to, azetidiny, 3-azabicyclo[3.1.0]hexan-3-yl, 1-azaspiro[3.3]heptan-1-yl, 5-azaspiro[2.3]hexan-5-yl, 2-oxa-6-azaspiro[3.3]heptan-6-yl, 1-oxa-6-azaspiro[3.4]octan-6-yl, 1-oxa-6-azaspiro[3.3]heptan-6-yl, 6-oxa-1-azaspiro[3.3]heptan-1-yl, 6-azaspiro[3.4]octan-6-yl, 7-oxa-2-azaspiro[3.5]nonan-2-yl, 2,6-diazaspiro[3.3]heptan-2-yl, dioxolanyl, dioxinyl, thienyl[1,3]dithianyl, decahydroisoquinolyl, imidazoliny, imidazolidiny, isothiazolidiny, isoxazolidiny, morpholiny, octahydroindolyl, octahydroisoindolyl, 2-oxopiperazinyl, 2-oxopiperidinyl, 2-oxopyrrolidinyl, oxazolidiny, piperidinyl, piperazinyl, 4-piperidonyl, pyrrolidinyl, pyrazolidiny, quinuclidiny, thiazolidiny, 1,2,4-thiadiazol-5(4H)-ylidene, tetrahydrofuryl, trioxanyl, trithianyl, triazinanyl, tetrahydropyranyl, thiomorpholiny, thiamorpholiny, 1-oxo-thiomorpholiny, and 1,1-dioxo-thiomorpholiny. In certain embodiments, a heterocyclyl group may be optionally substituted by one or more substituents selected from the group consisting of alkyl, alkenyl, halo, haloalkyl, haloalkenyl, cyano, oxo, thioxo, nitro, aryl, arylalkyl, cycloalkyl, cycloalkylalkyl, heterocyclyl, heterocyclylalkyl, heteroaryl, heteroarylalkyl, $-R^{101}-OR^{100}$, $-R^{101}-OC(O)-R^{100}$, $-R^{101}-N(R^{100})-R^{103}-OR^{100}$, $-R^{101}-N(R^{100})_2$, $-R^{101}-C(O)R^{100}$, $-R^{101}-C(O)OR^{100}$, $-R^{101}-C(O)N(R^{100})_2$, $-R^{101}-N(R^{100})C(O)OR^{102}$, $-R^{101}-N(R^{100})C(O)R^{102}$, $-R^{101}-N(R^{100})S(O)_pR^{102}$ (where p is 1 to 2), $-R^{101}-N=C(OR^{100})R^{102}$, $-R^{101}-S(O)_pOR^{102}$ (where p is 1 to 2), $-R^{101}-S(O)_tR^{102}$ (where t is 0 to 2), and $-R^{101}-S(O)_pN(R^{100})_2$ (where p is 1 to 2) where each R^{100} is independently hydrogen, alkyl, alkenyl, haloalkyl, cycloalkyl, cycloalkylalkyl, aryl,

arylalkyl, heterocyclyl, heterocyclalkyl, heteroaryl or heteroarylalkyl; each R^{101} is independently a direct bond or a straight or branched alkylene chain; each R^{102} is alkyl, alkenyl, haloalkyl, cycloalkyl, cycloalkylalkyl, aryl, arylalkyl, heterocyclyl, heterocyclalkyl, heteroaryl or heteroarylalkyl, and each R^{103} is a direct bond or a
 5 straight or branched alkylene chain.

"Heterocyclalkyl" refers to a radical of the formula $-R^{100}R^{101}$ where R^{100} is an alkylene chain as defined above and R^{101} is a heterocyclyl radical as defined above, and if the heterocyclyl is a nitrogen-containing heterocyclyl, the heterocyclyl may be attached to the alkyl radical at the nitrogen atom. In some embodiments, the alkylene
 10 chain of the heterocyclalkyl radical may be optionally substituted as defined above for an optionally substituted alkylene chain. In some embodiments, the heterocyclyl part of the heterocyclalkyl radical may be optionally substituted as defined above for an optionally substituted heterocyclyl group.

"Heteroaryl" refers to a 4- to 14-membered ring system radical comprising
 15 hydrogen atoms, one to thirteen carbon atoms, one to six heteroatoms selected from the group consisting of nitrogen, oxygen, and sulfur, and at least one aromatic ring. For purposes of this disclosure, the heteroaryl radical may be a monocyclic, bicyclic, tricyclic, or tetracyclic ring system, which may include fused or bridged ring systems; and the nitrogen, carbon, or sulfur atoms in the heteroaryl radical may be optionally
 20 oxidized; the nitrogen atom may be optionally quaternized. Examples include, but are not limited to, azepinyl, acridinyl, benzimidazolyl, benzthiazolyl, benzindolyl, benzodioxolyl, benzofuranyl, benzooxazolyl, benzothiazolyl, benzothiadiazolyl, benzo[b][1,4]dioxepinyl, 1,4-benzodioxanyl, benzonaphthofuranyl, benzoxazolyl, benzodioxolyl, benzodioxinyl, benzopyranyl, benzopyranonyl, benzofuranyl,
 25 benzofuranonyl, benzothienyl (benzothiophenyl), benzotriazolyl, benzo[4,6]imidazo[1,2-a]pyridinyl, benzoxazolinonyl, benzimidazolthionyl, carbazolyl, cinnolinyl, dibenzofuranyl, dibenzothiophenyl, furanyl, furanonyl, isothiazolyl, imidazolyl, indazolyl, indolyl, indazolyl, isoindolyl, indolinyl, isoindolinyl, isoquinolyl, indoliziny, isoxazolyl, naphthyridinyl, oxadiazolyl, 2-oxoazepinyl, oxazolyl, oxiranyl,
 30 1-oxidopyridinyl, 1-oxidopyrimidinyl, 1-oxidopyrazinyl, 1-oxidopyridazinyl, 1-phenyl-

1H-pyrrolyl, phenaziny, phenothiaziny, phenoxaziny, phthalaziny, pteridinyl, pteridinonyl, purinyl, pyrrolyl, pyrazolyl, pyridinyl, pyridinonyl, pyrazinyl, pyrimidinyl, pyrimidinonyl, pyridaziny, pyrrolyl, pyrido[2,3-d]pyrimidinonyl, quinazolinyl, quinazolinonyl, quinoxaliny, quinoxalinonyl, quinolinyl, isoquinolinyl, 5 tetrahydroquinolinyl, thiazolyl, thiadiazolyl, thieno[3,2-d]pyrimidin-4-onyl, thieno[2,3-d]pyrimidin-4-onyl, triazolyl, tetrazolyl, triazinyl, and thiophenyl (*i.e.* thienyl). In certain embodiments, a heteroaryl group may be optionally substituted by one or more substituents selected from the group consisting of alkyl, alkenyl, halo, haloalkyl, haloalkenyl, cyano, oxo, thioxo, nitro, thioxo, aryl, arylalkyl, cycloalkyl, cycloalkylalkyl, 10 heterocyclyl, heterocyclalkyl, heteroaryl, heteroarylalkyl, $-R^{101}-OR^{100}$, $-R^{101}-OC(O)-R^{100}$, $-R^{101}-N(R^{100})-R^{103}-OR^{100}$, $-R^{101}-N(R^{100})_2$, $-R^{101}-C(O)R^{100}$, $-R^{101}-C(O)OR^{100}$, $-R^{101}-C(O)N(R^{100})_2$, $-R^{101}-N(R^{100})C(O)OR^{102}$, $-R^{101}-N(R^{100})C(O)R^{100}C(O)R^{102}$, $-R^{101}-N(R^{100})S(O)_pR^{102}$ (where p is 1 to 2), $-R^{101}-N=C(OR^{100})R^{100}$, $-R^{101}-S(O)_pOR^{102}$ (where p is 1 to 2), $-R^{101}-S(O)_tR^{102}$ (where t is 0 to 2), and $-R^{101}-S(O)_pN(R^{100})_2$ (where p is 1 to 2) 15 where each R^{100} is independently hydrogen, alkyl, alkenyl, haloalkyl, cycloalkyl, cycloalkylalkyl, aryl, arylalkyl, heterocyclyl, heterocyclalkyl, heteroaryl or heteroarylalkyl; each R^{101} is independently a direct bond or a straight or branched alkylene chain; each R^{102} is alkyl, alkenyl, haloalkyl, cycloalkyl, cycloalkylalkyl, aryl, arylalkyl, heterocyclyl, heterocyclalkyl, heteroaryl, or heteroarylalkyl, and each R^{103} is 20 a direct bond or a straight or branched alkylene chain. Preferably, the optional substituents on an optionally substituted bicyclic heteroaryl group for R^1 herein are halo. Preferably, the optional substituents on an optionally substituted monocyclic heteroaryl group for R^1 herein are alkyl. The term "heteroaryl" includes, *e.g.*, the following structures:



"N-heteroaryl" refers to a heteroaryl radical as defined above containing at least one nitrogen. The point of attachment of the N-heteroaryl to the rest of the

molecule can be through a nitrogen atom or a carbon atom in the N-heteroaryl. When specifically stated in the specification, an N-heteroaryl radical may be optionally substituted as described above for an optionally substituted heteroaryl radical.

"Heteroarylalkyl" refers to a radical of the formula $\text{—R}^{100}\text{R}^{101}$ where R^{100} is an alkylene chain as defined above and R^{101} is a heteroaryl radical as defined above. When specifically stated in the specification, the heteroaryl part of the heteroarylalkyl radical may be optionally substituted as defined above for an optionally substituted heteroaryl group. In some specific embodiments, the alkylene chain part of the heteroarylalkyl radical may be optionally substituted as defined above for an optionally substituted alkylene chain.

The compounds and methods of the present disclosure are also meant to encompass all pharmaceutically acceptable compounds of Structures (I), (II) and (III) being isotopically-labelled by having one or more atoms replaced by an atom having a different atomic mass or mass number.

Examples of isotopes that can be incorporated into the disclosed compounds include isotopes of hydrogen, carbon, nitrogen, oxygen, phosphorous, fluorine, chlorine, and iodine, such as ^2H , ^3H , ^{11}C , ^{13}C , ^{14}C , ^{13}N , ^{15}N , ^{15}O , ^{17}O , ^{18}O , ^{31}P , ^{32}P , ^{35}S , ^{18}F , ^{36}Cl , ^{123}I , and ^{125}I , respectively. These radio-labelled compounds could be useful to help determine or measure the effectiveness of the compounds, by characterizing, for example, the site or mode of action or binding affinity. Certain isotopically-labelled compounds of Structures (I), (II) or (III), for example, those incorporating a radioactive isotope, are useful in drug and/or substrate tissue distribution studies. The radioactive isotopes tritium, *i.e.*, ^3H , and carbon-14, *i.e.*, ^{14}C , are particularly useful for this purpose in view of their ease of incorporation and ready means of detection.

Substitution with heavier isotopes such as deuterium, *i.e.*, ^2H , may afford certain therapeutic advantages resulting from greater metabolic stability, for example, increased *in vivo* half-life or reduced dosage requirements, and hence may be preferred in some circumstances. In one embodiment, the compounds of Structures (I), (II) or (III) are enriched with deuterium. Such deuterated compounds can be achieved by methods

known to one skilled in the art, such as exchanging protons with deuterium or by synthesizing the molecule with enriched starting materials.

Substitution with positron emitting isotopes, such as ^{11}C , ^{18}F , ^{15}O , and ^{13}N , can be useful in Positron Emission Topography (PET) studies for examining substrate
5 receptor occupancy. Isotopically-labeled compounds of Structures (I), (II) or (III) can generally be prepared by conventional techniques known to those skilled in the art or by processes analogous to those described in the Examples and Preparations as set out below using an appropriate isotopically-labeled reagent in place of the non-labeled reagent previously employed.

10 This disclosure is also meant to encompass the *in vivo* metabolic products of the disclosed compounds. Such products may result from, for example, the oxidation, reduction, hydrolysis, amidation, esterification, and the like of an administered compound, primarily due to enzymatic processes. Accordingly, this disclosure includes compounds produced by a process comprising contacting a compound of this disclosure
15 with a mammal for a period of time sufficient to yield a metabolic product thereof. Such products are typically identified by administering a radio-labelled compound in a detectable dose to an animal, such as rat, mouse, guinea pig, monkey, or to human, allowing sufficient time for metabolism to occur, and isolating its conversion products from the urine, blood, or other biological samples.

20 "Stable compound" and "stable structure" are meant to indicate a compound that is sufficiently robust to survive isolation to a useful degree of purity from a reaction mixture, and formulation into an efficacious therapeutic agent.

"Optional" or "optionally" means that the subsequently described event of circumstances may or may not occur, and that the description includes instances where
25 said event or circumstance occurs and instances in which it does not. For example, "optionally substituted aryl" means that the aryl radical may or may not be substituted and that the description includes both substituted aryl radicals and aryl radicals having no substitution ("unsubstituted"). When a functional group is described as "optionally substituted," and in turn, substituents on the functional group are also "optionally

substituted" and so on, for the purposes of this disclosure, such iterations are limited to five, preferably such iterations are limited to two.

"Pharmaceutically acceptable carrier, diluent or excipient" includes without limitation any adjuvant, carrier, excipient, glidant, sweetening agent, diluent, 5 preservative, dye/colorant, flavor enhancer, surfactant, wetting agent, dispersing agent, suspending agent, stabilizer, isotonic agent, solvent, or emulsifier which has been approved by the United States Food and Drug Administration as being acceptable for use in humans or domestic animals.

"Pharmaceutically acceptable salt" includes both acid and base addition 10 salts.

"Pharmaceutically acceptable acid addition salt" refers to those salts which retain the biological effectiveness and properties of the free bases, which are not biologically or otherwise undesirable, and which are formed with inorganic acids such as, but are not limited to, hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, 15 phosphoric acid and the like, and organic acids such as, but not limited to, acetic acid, 2,2-dichloroacetic acid, adipic acid, alginic acid, ascorbic acid, aspartic acid, benzenesulfonic acid, benzoic acid, 4-acetamidobenzoic acid, camphoric acid, camphor-10-sulfonic acid, capric acid, caproic acid, caprylic acid, carbonic acid, cinnamic acid, citric acid, cyclamic acid, dodecylsulfuric acid, ethane-1,2-disulfonic acid, 20 ethanesulfonic acid, 2-hydroxyethanesulfonic acid, formic acid, fumaric acid, galactaric acid, gentisic acid, glucoheptonic acid, gluconic acid, glucuronic acid, glutamic acid, glutaric acid, 2-oxo-glutaric acid, glycerophosphoric acid, glycolic acid, hippuric acid, isobutyric acid, lactic acid, lactobionic acid, lauric acid, maleic acid, malic acid, malonic acid, mandelic acid, methanesulfonic acid, mucic acid, naphthalene-1,5-disulfonic acid, 25 naphthalene-2-sulfonic acid, 1-hydroxy-2-naphthoic acid, nicotinic acid, oleic acid, orotic acid, oxalic acid, palmitic acid, pamoic acid, propionic acid, pyroglutamic acid, pyruvic acid, salicylic acid, 4-aminosalicylic acid, sebacic acid, stearic acid, succinic acid, tartaric acid, thiocyanic acid, p-toluenesulfonic acid, trifluoroacetic acid, undecylenic acid, and the like.

"Pharmaceutically acceptable base addition salt" refers to those salts which retain the biological effectiveness and properties of the free acids, which are not biologically or otherwise undesirable. These salts are prepared from addition of an inorganic base or an organic base to the free acid. Salts derived from inorganic bases include, but are not limited to, the sodium, potassium, lithium, ammonium, calcium, magnesium, iron, zinc, copper, manganese, aluminum salts, and the like. Preferred inorganic salts are the ammonium, sodium, potassium, calcium, and magnesium salts. Salts derived from organic bases include, but are not limited to, salts of primary, secondary, and tertiary amines, substituted amines including naturally occurring substituted amines, cyclic amines and basic ion exchange resins, such as ammonia, isopropylamine, trimethylamine, diethylamine, triethylamine, tripropylamine, diethanolamine, ethanolamine, deanol, 2-dimethylaminoethanol, 2-diethylaminoethanol, dicyclohexylamine, lysine, arginine, histidine, caffeine, procaine, hydrabamine, choline, betaine, benethamine, benzathine, ethylenediamine, glucosamine, methylglucamine, theobromine, triethanolamine, tromethamine, purines, piperazine, piperidine, N-ethylpiperidine, polyamine resins and the like. Particularly preferred organic bases are isopropylamine, diethylamine, ethanolamine, trimethylamine, dicyclohexylamine, choline, and caffeine.

Often crystallizations produce a solvate of the compound of the disclosure (*e.g.*, a compound of Structure (I), (II) or (III)). As used herein, the term "solvate" refers to an aggregate that comprises one or more molecules of a compound of the disclosure with one or more molecules of solvent. The solvent may be water, in which case the solvate may be a hydrate. Alternatively, the solvent may be an organic solvent. Thus, the compounds of the present disclosure may exist as a hydrate, including a monohydrate, dihydrate, hemihydrate, sesquihydrate, trihydrate, tetrahydrate, and the like, as well as the corresponding solvated forms. The compound of the disclosure may be true solvates, while in other cases, the compound of the disclosure may merely retain adventitious water or be a mixture of water plus some adventitious solvent.

A "pharmaceutical composition" refers to a formulation of a compound of the disclosure and a medium generally accepted in the art for the delivery of the

biologically active compound to mammals, *e.g.*, humans. Such a medium includes all pharmaceutically acceptable carriers, diluents, or excipients therefor.

"Treating" or "treatment" as used herein covers the treatment of the disease or condition of interest in a mammal, preferably a human, having the disease or
5 condition of interest, and includes:

(a) preventing the disease or condition from occurring in a mammal, in particular, when such mammal is predisposed to the condition but has not yet been diagnosed as having it;

(b) inhibiting the disease or condition, *i.e.*, arresting the disease or
10 condition's development;

(c) relieving (or ameliorating) the disease or condition, *i.e.*, causing regression of the disease or condition; or

(d) relieving (or ameliorating) the symptoms resulting from the disease or condition, *e.g.*, without addressing the underlying disease or condition.

15 As used herein, the terms "disease" and "condition" may be used interchangeably or may be different in that the particular malady or condition may not have a known causative agent (so that etiology has not yet been worked out) and it is therefore not yet recognized as a disease but only as an undesirable condition or syndrome, wherein a more or less specific set of symptoms have been identified by
20 clinicians.

The compounds of the disclosure, or their pharmaceutically acceptable salts may contain one or more stereocenter and may thus give rise to enantiomers, diastereomers, and other stereoisomeric forms that may be defined, in terms of absolute stereochemistry, as (*R*)- or (*S*)- or, as (*D*)- or (*L*)- for amino acids. The present disclosure
25 is meant to include all such possible isomers, as well as their racemic and optically pure forms. Optically active (+) and (−), (*R*)- and (*S*)-, or (*D*)- and (*L*)- isomers may be prepared using chiral synthons or chiral reagents, or resolved using conventional techniques, for example, chromatography and fractional crystallization. Conventional techniques for the preparation/isolation of individual enantiomers include chiral synthesis
30 from a suitable optically pure precursor or resolution of the racemate (or the racemate of

a salt or derivative) using, for example, chiral high-pressure liquid chromatography (HPLC). When the compounds described herein contain olefinic double bonds or other centres giving rise to geometric asymmetry, and unless specified otherwise, it is intended that the compounds include both E and Z geometric isomers. Likewise, all tautomeric
5 forms are also intended to be included.

A "stereoisomer" refers to a compound made up of the same atoms bonded by the same bonds but having different three-dimensional structures, which are not interchangeable. The present disclosure contemplates various stereoisomers and mixtures thereof and includes enantiomers, which refers to two stereoisomers whose molecules are
10 non-superimposable mirror images of one another. See, *e.g.*, Smith, M. B. and J. March, March's Advanced Organic Chemistry: Reactions, Mechanisms, and Structure, 6th edition (Wiley, 2007), for a detailed description of the structure and properties of enantiomers and stereoisomers.

A "tautomer" refers to a proton shift from one atom of a molecule to
15 another atom of the same molecule. The present disclosure includes tautomers of any said compounds.

The use of parentheses and brackets in substituent groups is used herein to conserve space. Accordingly, the use of parenthesis in a substituent group indicates that the group enclosed within the parentheses is attached directly to the atom preceding
20 the parenthesis. The use of brackets in a substituent group indicates that the group enclosed within the brackets is also attached directly to the atom preceding the parenthesis.

The chemical naming protocol and structure diagrams used herein are a modified form of the I.U.P.A.C. nomenclature system, using ChemBioDraw Ultra
25 Version 14.0 software program. For complex chemical names employed herein, a substituent group is named before the group to which it attaches. For example, cyclopropylethyl comprises an ethyl backbone with cyclopropyl substituent. In chemical structure diagrams, all bonds are identified, except for some carbon atoms, which are assumed to be bonded to sufficient hydrogen atoms to complete the valency.

At certain places, the definitions or embodiments may refer to specific rings (*e.g.*, an azetidine ring, a pyridine ring, *etc.*). Unless otherwise indicated, these rings can be attached to any ring member provided that the valency of the atom is not exceeded.

When any two groups or two instances of the same substituent group are
5 "independently selected" from a list of alternatives, the groups may be the same or different. For example, if R^a and R^b are independently selected from the group consisting of alkyl, fluoro, amino, and hydroxyalkyl, then a molecule with two R^a groups and two R^b groups could have all groups be alkyl group (*e.g.*, four different alkyl groups). Alternatively, the first R^a could be alkyl, the second R^a could be fluoro, the first R^b could
10 be hydroxyalkyl, and the second R^b could be amino (or any other substituents taken from the group). Alternatively, both R^a and the first R^b could be fluoro, while the second R^b could be alkyl (*i.e.*, some pairs of substituent groups may be the same, while other pairs may be different). Unless otherwise indicated, if two or more groups having the same definition are present, but the definition provides for alternatives, it should be understood
15 that each occurrence of the same group is independently selected from the possible alternatives. For example, if two or more R^a groups are present in a compound, and the definition of R^a provides that R^a can be A, B, or C, then it should be understood that each R^a group present in the compound is independently chosen from A, B, and C, so that the R^a groups present in the compound can be the same or different.

20 Compounds, and salts thereof, including pharmaceutically acceptable salts, can be found together with other substances such as water and solvents (*e.g.*, hydrates and solvates) or can be isolated. When in the solid state, the compounds described herein, and salts thereof may occur in various forms and may, *e.g.*, take the form of solvates, including hydrates. The compounds may be in any solid-state form,
25 such as a polymorph or solvate, so unless clearly indicated otherwise, reference to compounds and salts thereof should be understood as encompassing any solid-state form of the compound.

In some embodiments, the compounds described herein or salts thereof, are substantially isolated. "Substantially isolated" means the compound is at least
30 partially or substantially separated from the environment in which it was formed or


detected. Partial separation can include, *e.g.*, a composition enriched in the compounds of the disclosure. Substantial separation can include compositions containing at least about 50%, at least about 60%, at least about 70%, at least about 80%, at least about 90%, at least about 95%, at least about 97%, or at least about 99% by weight of the compounds of the disclosure, or salt thereof.

The following abbreviations may be used herein and, unless otherwise noted, have the meanings indicated below:

ACN or MeCN (acetonitrile); chrom. (chromatography); CH₂Cl₂ (dichloromethane); DIAD (Diisopropyl azodicarboxylate); L (liter(s)); *m*- (meta); Ms (methanesulfonyl); NCS (*N*-chlorosuccinimide); NIS (*N*-iodosuccinimide); Phth (Phthalimide); TPP (triphenylphosphine); and Tr (trityl). Other abbreviations may also be used and have the meanings that would be understood by the person having skill in the art.

II. Compounds

In certain aspects, the present disclosure provides a compound of Structure (I):



(I)

or a stereoisomer, tautomer, or pharmaceutically acceptable salt thereof, wherein:

X is a direct bond, -CR^{2e}R^{2f}-, or -CR^{2e}R^{2f}-CR^{2g}R^{2h}-;

Y is a direct bond or -CR²ⁱR^{2j}-;

R¹ is a substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, substituted or unsubstituted cycloalkyl, or a substituted or unsubstituted heterocyclyl;

R^{2a}, R^{2b}, R^{2c}, R^{2d}, R^{2e}, R^{2f}, R^{2g}, R^{2h}, R²ⁱ, and R^{2j} are each independently selected from the group consisting of hydrogen, halo, C(=O)OR⁵, OC(=O)R⁵,

hydroxyalkyl, alkoxy, alkoxyalkyl, haloalkoxy, cyano, aminylalkyl, carboxyalkyl, NR^5R^6 , $\text{C}(=\text{O})\text{NR}^5\text{R}^6$, $\text{N}(\text{R}^5)\text{C}(=\text{O})\text{R}^6$, $\text{NR}^5\text{C}(=\text{O})\text{NR}^6$, $\text{S}(\text{O})_t$, SR^5 , nitro, $\text{N}(\text{R}^5)\text{C}(\text{O})\text{OR}^6$, $\text{C}(=\text{NR}^5)\text{NR}^6\text{R}^7$, $\text{N}(\text{R}^5)\text{C}(=\text{NR}^6)\text{NR}^7\text{R}^8$, $\text{S}(\text{O})\text{R}^5$, $\text{S}(\text{O})\text{NR}^5\text{R}^6$, $\text{S}(\text{O})_2\text{R}^5$, $\text{N}(\text{R}^5)\text{S}(\text{O})_2\text{R}^6$, $\text{S}(\text{O})_2\text{NR}^5\text{R}^6$, aryl, heteroaryl, heterocyclyl, cycloalkyl, and oxo, provided that at least
 5 one of R^{2a} , R^{2b} , R^{2c} , R^{2d} , R^{2e} , R^{2f} , R^{2g} , R^{2h} , R^{2i} , and R^{2j} is not hydrogen;

R^3 is $\text{NR}^{3a}\text{R}^{3b}$;

R^{3a} and R^{3b} are each independently hydrogen, alkyl, hydroxyalkyl, haloalkyl, alkoxyalkyl, heterocyclyl, heteroaryl, heterocyclylalkyl, heteroarylalkyl, or cycloalkyl; or R^{3a} and R^{3b} , together with the nitrogen to which they are attached, form an
 10 optionally substituted 4-7 membered heteroaryl or an optionally substituted 4-7 membered heterocyclyl; or R^{3a} and R^4 together with the nitrogen and carbon to which they are attached, respectively, form an optionally substituted 4-7 membered heterocyclyl, and R^{3b} is hydrogen, alkyl, hydroxyalkyl, haloalkyl, alkoxyalkyl, heterocyclyl, heteroaryl, heterocyclylalkyl, heteroarylalkyl, cycloalkyl,
 15 $(\text{CH}_2)_n\text{C}(=\text{O})\text{OR}^6$, or $(\text{CH}_2)_n\text{P}(=\text{O})(\text{OR}^6)_2$;

R^4 is a substituted or unsubstituted aryl, a substituted or unsubstituted heteroaryl, a substituted or unsubstituted cycloalkyl, or a substituted or unsubstituted heterocyclyl when n is 2, 3, 4, 5, or 6; or R^4 is a substituted or unsubstituted monocyclic heteroaryl, or a substituted or unsubstituted heterocyclyl when n is 0 or 1; or R^4 and R^{3a}
 20 together with the carbon and nitrogen to which they are attached, respectively, form an optionally substituted 4-7 membered heterocyclyl;

R^5 , R^6 , R^7 , and R^8 are, at each occurrence, independently hydrogen, alkyl, hydroxyalkyl, haloalkyl, alkoxyalkyl, carboxyalkyl, heterocyclyl, heteroaryl, or cycloalkyl;

25 n is an integer from 0-6; and

t is an integer from 1-3.

In some embodiments, R^1 is a substituted or unsubstituted aryl. In certain embodiments, R^1 is a substituted or unsubstituted $\text{C}_6\text{-C}_{10}$ aryl. In some more specific embodiments, R^1 is a substituted or unsubstituted phenyl. In certain specific
 30 embodiments, R^1 is a substituted phenyl.

In some embodiments, R^1 is phenyl substituted with one or more of R^{1a} , R^{1b} , R^{1c} , R^{1d} , and R^{1e} wherein R^{1a} , R^{1b} , R^{1c} , R^{1d} , and R^{1e} are each independently selected from the group consisting of C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, halo, C_{1-6} haloalkyl, aminylalkyl, hydroxyalkyl, cyano, nitro, OR^9 , SR^9 , $C(O)R^9$, $C(O)NR^9R^{10}$, $C(O)OR^9$,
 5 $OC(O)R^9$, $OC(O)OR^9$, $OC(O)NR^9R^{10}$, NR^9R^{10} , $N(R^9)C(O)R^{10}$, $N(R^9)C(O)NR^{10}R^{11}$,
 $N(R^9)C(O)OR^{10}$, $C(=NR^9)NR^{10}R^{11}$, $C(=NOR^9)NR^{10}R^{11}$, $C(=NOC(O)R^9)NR^{10}R^{11}$,
 $C(=NR^9)N(R^{10})C(O)OR^{11}$, $N(R^9)C(=NR^{10})NR^{11}R^{12}$, $S(O)R^9$, $S(O)NR^9R^{10}$, $S(O)_2R^9$,
 $N(R^9)S(O)_2R^{10}$, $S(O)_2NR^9R^{10}$, oxo, substituted or unsubstituted C_{6-10} aryl, substituted or
 10 unsubstituted C_{6-10} arylalkyl, substituted or unsubstituted C_{6-10} aryloxy, substituted or
 unsubstituted C_{6-10} arylalkoxy, substituted or unsubstituted 5-10 membered heteroaryl,
 substituted or unsubstituted C_{3-10} cycloalkyl, and substituted or unsubstituted 4-10
 membered heterocyclyl, wherein R^9 , R^{10} , R^{11} , and R^{12} , are, at each occurrence,
 independently selected from the group consisting of hydrogen, C_{1-6} alkyl, C_{2-6} alkenyl,
 C_{2-6} alkynyl, hydroxyl, C_{1-6} alkoxy, aryl, arylalkyl, C_{1-6} haloalkyl, C_{1-6} haloalkoxy, C_{1-6}
 15 hydroxyalkyl, cycloalkyl, heterocyclyl, and heteroaryl.


In some more specific embodiments, R^{1a} , R^{1b} , R^{1c} , R^{1d} , or R^{1e} is optionally substituted with one or more substituents selected from the group consisting of halo, CN, OR^{13} , SR^{13} , $C(O)R^{13}$, $C(O)NR^{13}R^{14}$, $C(O)OR^{13}$, $OC(O)R^{13}$, $OC(O)NR^{13}R^{14}$, $NR^{13}R^{14}$,
 $NR^{13}C(O)R^{14}$, $NR^{13}C(O)NR^{14}R^{15}$, $NR^{13}C(O)OR^{14}$, $C(=NR^{13})NR^{14}R^{15}$,
 20 $NR^{13}C(=NR^{14})NR^{15}R^{16}$, $S(O)R^{13}$, $S(O)NR^{13}R^{14}$, $S(O)_2R^{13}$, $NR^{13}S(O)_2R^{14}$, $S(O)_2NR^{13}R^{14}$
 and oxo when R^{1a} , R^{1b} , R^{1c} , R^{1d} , or R^{1e} is a substituted C_{6-10} aryl, a substituted C_{6-10}
 arylalkyl, a substituted C_{6-10} aryloxy, a substituted C_{6-10} arylalkoxy, a substituted 5-10
 membered heteroaryl, a substituted C_{3-10} cycloalkyl, and a substituted 4-10 membered
 heterocyclyl, wherein R^{13} , R^{14} , R^{15} , and R^{16} are, at each occurrence, independently
 25 selected from the group consisting of hydrogen, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl,
 hydroxyl, C_{1-6} alkoxy, aryl, arylalkyl, C_{1-6} haloalkyl, C_{1-6} haloalkoxy, C_{1-6} hydroxyalkyl,
 cycloalkyl, heterocyclyl, and heteroaryl.


In certain embodiments, R^1 is phenyl substituted with at least one substituent selected from the group consisting of OR^9 , $C(=NR^9)NR^{10}R^{11}$,
 30 $C(=NOR^9)NR^{10}R^{11}$, $C(=NOC(O)R^9)NR^{10}R^{11}$, and $C(=NR^9)NR^{10}C(O)OR^{11}$. In certain

more specific embodiments, R^1 is substituted with at least one substituent selected from the group consisting of $C(=NR^9)NR^{10}R^{11}$, $C(=NOR^9)NR^{10}R^{11}$, $C(=NOC(O)R^9)NR^{10}R^{11}$, and $C(=NR^9)NR^{10}C(O)OR^{11}$. In some embodiments, R^1 is substituted with at least one $C(=NR^9)NR^{10}R^{11}$. In some specific embodiments, R^1 is substituted with at least one -


5 $C(=NH)NH_2$.

In certain more specific embodiments, R^1 has one of the following structures:






In still more specific embodiments, R^1 has one of the following structures:



In some embodiments, R^1 has one of the following structures:




In some of the foregoing embodiments, R^9 is C_{1-6} alkyl or C_{1-6} haloalkyl. In certain more specific embodiments, R^9 is methyl. In some embodiments, R^9 is trifluoromethyl. In more embodiments, R^1 is an unsubstituted phenyl.

In certain other embodiments, R^1 is a substituted or unsubstituted heteroaryl. In more specific embodiments, R^1 is a substituted or unsubstituted 5-10 membered heteroaryl. In some more specific embodiments, R^1 is a substituted or unsubstituted pyridinyl, pyrrolopyridinyl, or benzoimidazolyl. In some embodiments, R^1 is a substituted or unsubstituted pyridin-3-yl, 1H-pyrrolo[2,3-b]pyridine-5-yl, 1H-pyrrolo[3,2-c]pyridine-2-yl, or 1H-benzo[d]imidazol-6-yl. In more specific embodiments, R^1 is pyridin-3-yl, 1H-pyrrolo[2,3-b]pyridine-5-yl, 1H-pyrrolo[3,2-c]pyridine-2-yl, or 1H-benzo[d]imidazol-6-yl substituted with one or more of R^{1a} , R^{1b} , R^{1c} , R^{1d} , and R^{1e} wherein R^{1a} , R^{1b} , R^{1c} , R^{1d} , and R^{1e} are each independently selected from the group consisting of C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, halo, C_{1-6} haloalkyl, aminylalkyl, hydroxyalkyl, cyano, nitro, OR^9 , SR^9 , $C(O)R^9$, $C(O)NR^9R^{10}$, $C(O)OR^9$, $OC(O)R^9$, $OC(O)OR^9$, $OC(O)NR^9R^{10}$, NR^9R^{10} , $N(R^9)C(O)R^{10}$, $N(R^9)C(O)NR^{10}R^{11}$, $N(R^9)C(O)OR^{10}$, $C(=NR^9)NR^{10}R^{11}$, $C(=NOR^9)NR^{10}R^{11}$, $C(=NOC(O)R^9)NR^{10}R^{11}$, $C(=NR^9)N(R^{10})C(O)OR^{11}$, $N(R^9)C(=NR^{10})NR^{11}R^{12}$, $S(O)R^9$, $S(O)NR^9R^{10}$, $S(O)_2R^9$, $N(R^9)S(O)_2R^{10}$, $S(O)_2NR^9R^{10}$, oxo, substituted or unsubstituted C_{6-10} aryl, substituted or unsubstituted C_{6-10} arylalkyl, substituted or unsubstituted C_{6-10} aryloxy, substituted or unsubstituted C_{6-10} arylalkoxy, substituted or unsubstituted 5-10 membered heteroaryl, substituted or unsubstituted C_{3-10} cycloalkyl, and substituted or unsubstituted 4-10 membered heterocyclyl, wherein R^9 , R^{10} , R^{11} , and R^{12} , are, at each occurrence,


independently selected from the group consisting of hydrogen, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, hydroxyl, C₁₋₆ alkoxy, aryl, arylalkyl, C₁₋₆ haloalkyl, C₁₋₆ haloalkoxy, C₁₋₆ hydroxyalkyl, cycloalkyl, heterocyclyl, and heteroaryl.


In some embodiments, R^{1a}, R^{1b}, R^{1c}, R^{1d}, or R^{1e} is optionally substituted
 5 with one or more substituents selected from the group consisting of halo, CN, OR¹³, SR¹³, C(O)R¹³, C(O)NR¹³R¹⁴, C(O)OR¹³, OC(O)R¹³, OC(O)NR¹³R¹⁴, NR¹³R¹⁴, NR¹³C(O)R¹⁴, NR¹³C(O)NR¹⁴R¹⁵, NR¹³C(O)OR¹⁴, C(=NR¹³)NR¹⁴R¹⁵, NR¹³C(=NR¹⁴)NR¹⁵R¹⁶, S(O)R¹³, S(O)NR¹³R¹⁴, S(O)₂R¹³, NR¹³S(O)₂R¹⁴, S(O)₂NR¹³R¹⁴ and oxo when R^{1a}, R^{1b}, R^{1c}, R^{1d}, or R^{1e} is a substituted C₆₋₁₀ aryl, a substituted C₆₋₁₀ arylalkyl, a substituted C₆₋₁₀
 10 aryloxy, a substituted C₆₋₁₀ arylalkoxy, a substituted 5-10 membered heteroaryl, a substituted C₃₋₁₀ cycloalkyl, and a substituted 4-10 membered heterocyclyl, wherein R¹³, R¹⁴, R¹⁵, and R¹⁶ are, at each occurrence, independently selected from the group consisting of hydrogen, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, hydroxyl, C₁₋₆ alkoxy, aryl, arylalkyl, C₁₋₆ haloalkyl, C₁₋₆ haloalkoxy, C₁₋₆ hydroxyalkyl, cycloalkyl, heterocyclyl,
 15 and heteroaryl.


In certain specific embodiments, R¹ has one of the following structures:



5










- 5 In some specific embodiments, R^{1a} or R^{1b} is independently C_{1-6} alkyl, amino, $C(O)R^9$, $C(O)NR^9R^{10}$, $N(R^9)C(O)NR^{10}R^{11}$, $N(R^9)C(O)OR^{10}$, $OC(O)NR^9R^{10}$, or halo. In some embodiments, R^{1a} or R^{1b} is methyl or ethyl. In some specific embodiments, R^{1a} or R^{1b} is F, Cl, or Br. In certain embodiments, each R^{1a} or R^{1b} attached to nitrogen is C_{1-6} alkyl, $C(O)R^9$, $C(O)NR^9R^{10}$, $N(R^9)C(O)NR^{10}R^{11}$, $N(R^9)C(O)OR^{10}$, or
- 10 $OC(O)NR^9R^{10}$. In more specific embodiments, R^{1a} or R^{1b} is methyl or ethyl.


In some embodiments, R^1 has one of the following structures:







In certain embodiments, R^1 has one of the following structures:







In certain embodiments, R^1 has one of the following structures:



In some embodiments, R^1 is a substituted or unsubstituted cycloalkyl. In certain embodiments, R^1 is a substituted or unsubstituted C_3 - C_6 cycloalkyl. In some more specific embodiments, R^1 is a substituted C_3 - C_6 cycloalkyl.

In some embodiments, R^1 is a C_3 - C_6 cycloalkyl substituted with one or more of R^{1a} , R^{1b} , R^{1c} , R^{1d} , and R^{1e} wherein R^{1a} , R^{1b} , R^{1c} , R^{1d} , and R^{1e} are each independently selected from the group consisting of C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} haloalkyl, aminylalkyl, hydroxyalkyl, nitro, OR^9 , SR^9 , $C(O)R^9$, $C(O)NR^9R^{10}$,

$C(O)OR^9$, $OC(O)R^9$, $OC(O)OR^9$, $OC(O)NR^9R^{10}$, NR^9R^{10} , $N(R^9)C(O)R^{10}$,
 $N(R^9)C(O)NR^{10}R^{11}$, $N(R^9)C(O)OR^{10}$, $C(=NR^9)NR^{10}R^{11}$, $C(=NOR^9)NR^{10}R^{11}$,
 $C(=NOC(O)R^9)NR^{10}R^{11}$, $C(=NR^9)N(R^{10})C(O)OR^{11}$, $N(R^9)C(=NR^{10})NR^{11}R^{12}$, $S(O)R^9$,
 $S(O)NR^9R^{10}$, $S(O)_2R^9$, $N(R^9)S(O)_2R^{10}$, $S(O)_2NR^9R^{10}$, oxo, substituted or unsubstituted
5 C_{6-10} aryl, substituted or unsubstituted C_{6-10} arylalkyl, substituted or unsubstituted C_{6-10}
arylloxy, substituted or unsubstituted C_{6-10} arylalkoxy, substituted or unsubstituted 5-10
membered heteroaryl, substituted or unsubstituted C_{3-10} cycloalkyl, and substituted or
unsubstituted 4-10 membered heterocyclyl, wherein R^9 , R^{10} , R^{11} , and R^{12} , are, at each
occurrence, independently selected from the group consisting of hydrogen, C_{1-6} alkyl, C_{2-}
10 $_6$ alkenyl, C_{2-6} alkynyl, hydroxyl, C_{1-6} alkoxy, aryl, arylalkyl, C_{1-6} haloalkyl, C_{1-6}
haloalkoxy, C_{1-6} hydroxyalkyl, cycloalkyl, heterocyclyl, and heteroaryl.

In certain embodiments of the foregoing, R^{1a} , R^{1b} , R^{1c} , R^{1d} , or R^{1e} is
optionally substituted with one or more substituents selected from the group consisting
of OR^{13} , SR^{13} , $C(O)R^{13}$, $C(O)NR^{13}R^{14}$, $C(O)OR^{13}$, $OC(O)R^{13}$, $OC(O)NR^{13}R^{14}$, $NR^{13}R^{14}$,
15 $NR^{13}C(O)R^{14}$, $NR^{13}C(O)NR^{14}R^{15}$, $NR^{13}C(O)OR^{14}$, $C(=NR^{13})NR^{14}R^{15}$,
 $NR^{13}C(=NR^{14})NR^{15}R^{16}$, $S(O)R^{13}$, $S(O)NR^{13}R^{14}$, $S(O)_2R^{13}$, $NR^{13}S(O)_2R^{14}$, $S(O)_2NR^{13}R^{14}$
and oxo when R^{1a} , R^{1b} , R^{1c} , R^{1d} , or R^{1e} is a substituted C_{6-10} aryl, a substituted C_{6-10}
arylalkyl, a substituted C_{6-10} aryloxy, a substituted C_{6-10} arylalkoxy, a substituted 5-10
membered heteroaryl, a substituted C_{3-10} cycloalkyl, and a substituted 4-10 membered
20 heterocyclyl, wherein R^{13} , R^{14} , R^{15} , and R^{16} are, at each occurrence, independently
selected from the group consisting of hydrogen, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl,
hydroxyl, C_{1-6} alkoxy, aryl, arylalkyl, C_{1-6} haloalkyl, C_{1-6} haloalkoxy, C_{1-6} hydroxyalkyl,
cycloalkyl, heterocyclyl, and heteroaryl.

In some embodiments, R^1 is an unsubstituted C_3-C_6 cycloalkyl. In
25 certain embodiments, R^1 is a substituted or unsubstituted heterocyclyl. In certain specific
embodiments, R^1 is a substituted or unsubstituted 4-10 membered heterocyclyl. In some
embodiments, R^1 is a substituted 4-10 membered heterocyclyl.









In some embodiments, R^1 is a 4-10 membered heterocyclyl substituted
with one or more of R^{1a} , R^{1b} , R^{1c} , R^{1d} , and R^{1e} wherein R^{1a} , R^{1b} , R^{1c} , R^{1d} , and R^{1e} are each
30 independently selected from the group consisting of C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl,





halo, C₁₋₆ haloalkyl, aminylalkyl, hydroxyalkyl, cyano, nitro, OR⁹, SR⁹, C(O)R⁹, C(O)NR⁹R¹⁰, C(O)OR⁹, OC(O)R⁹, OC(O)OR⁹, OC(O)NR⁹R¹⁰, NR⁹R¹⁰, N(R⁹)C(O)R¹⁰, N(R⁹)C(O)NR¹⁰R¹¹, N(R⁹)C(O)OR¹⁰, C(=NR⁹)NR¹⁰R¹¹, C(=NOR⁹)NR¹⁰R¹¹, C(=NOC(O)R⁹)NR¹⁰R¹¹, C(=NR⁹)N(R¹⁰)C(O)OR¹¹, N(R⁹)C(=NR¹⁰)NR¹¹R¹², S(O)R⁹,
 5 S(O)NR⁹R¹⁰, S(O)₂R⁹, N(R⁹)S(O)₂R¹⁰, S(O)₂NR⁹R¹⁰, oxo, substituted or unsubstituted C₆₋₁₀ aryl, substituted or unsubstituted C₆₋₁₀ arylalkyl, substituted or unsubstituted C₆₋₁₀ aryloxy, substituted or unsubstituted C₆₋₁₀ arylalkoxy, substituted or unsubstituted 5-10 membered heteroaryl, substituted or unsubstituted C₃₋₁₀ cycloalkyl, and substituted or unsubstituted 4-10 membered heterocyclyl, wherein R⁹, R¹⁰, R¹¹, and R¹², are, at each
 10 occurrence, independently selected from the group consisting of hydrogen, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, hydroxyl, C₁₋₆ alkoxy, aryl, arylalkyl, C₁₋₆ haloalkyl, C₁₋₆ haloalkoxy, C₁₋₆ hydroxyalkyl, cycloalkyl, heterocyclyl, and heteroaryl.

In some of the foregoing embodiments, R^{1a}, R^{1b}, R^{1c}, R^{1d}, or R^{1e} is optionally substituted with one or more substituents selected from the group consisting
 15 of halo, CN, OR¹³, SR¹³, C(O)R¹³, C(O)NR¹³R¹⁴, C(O)OR¹³, OC(O)R¹³, OC(O)NR¹³R¹⁴, NR¹³R¹⁴, NR¹³C(O)R¹⁴, NR¹³C(O)NR¹⁴R¹⁵, NR¹³C(O)OR¹⁴, C(=NR¹³)NR¹⁴R¹⁵, NR¹³C(=NR¹⁴)NR¹⁵R¹⁶, S(O)R¹³, S(O)NR¹³R¹⁴, S(O)₂R¹³, NR¹³S(O)₂R¹⁴, S(O)₂NR¹³R¹⁴ and oxo when R^{1a}, R^{1b}, R^{1c}, R^{1d}, or R^{1e} is a substituted C₆₋₁₀ aryl, a substituted C₆₋₁₀ arylalkyl, a substituted C₆₋₁₀ aryloxy, a substituted C₆₋₁₀ arylalkoxy, a substituted 5-10
 20 membered heteroaryl, a substituted C₃₋₁₀ cycloalkyl, and a substituted 4-10 membered heterocyclyl, wherein R¹³, R¹⁴, R¹⁵, and R¹⁶ are, at each occurrence, independently selected from the group consisting of hydrogen, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, hydroxyl, C₁₋₆ alkoxy, aryl, arylalkyl, C₁₋₆ haloalkyl, C₁₋₆ haloalkoxy, C₁₋₆ hydroxyalkyl, cycloalkyl, heterocyclyl, and heteroaryl.

25 In some embodiments, R¹ is an unsubstituted 4-10 membered heterocyclyl.


In some embodiments, the compound has one of the following structures:






In some embodiments, at least one occurrence of R^{2a} , R^{2b} , R^{2c} , or R^{2d} is alkoxy, alkoxyalkyl, cyano, $C(=O)OR^5$, $OC(=O)R^5$, $C(=O)NR^5R^6$, $N(R^5)C(=O)R^6$, aryl, heteroaryl, heterocyclyl, or $C(=O)N(R^5)$ heterocyclyl, wherein R^5 and R^6 are, at each occurrence, independently hydrogen or C_1 - C_6 alkyl. In some embodiments, R^{2a} and R^{2b} are both halo (*e.g.*, fluoro).

In some more specific embodiments, X is CH_2 , Y is a direct bond, R^{2c} and R^{2d} are both hydrogen, and R^{2a} and R^{2b} are both fluoro. In more specific embodiments, R^{3b} is hydrogen and R^{3a} and R^4 together with the nitrogen and carbon to which they are attached, respectively, form an optionally substituted 4-7 membered heterocyclyl (*e.g.*, piperidinyl or pyrrolyl). In some more specific embodiments, the compound of Structure (I) has the following structure:



wherein:


R^{33} is a substituted or unsubstituted aryl or a substituted or unsubstituted heteroaryl. In some embodiments, R^{33} has the following structure:




In certain embodiments, at least one occurrence of R^{2a} , R^{2b} , R^{2c} , or R^{2d} is methoxy. In some embodiments, at least one of R^{2a} , R^{2b} , R^{2c} , and R^{2d} is methoxymethyl. In certain embodiments, at least one of R^{2a} , R^{2b} , R^{2c} , and R^{2d} is cyano. In some more specific embodiments, at least one of R^{2a} , R^{2b} , R^{2c} , and R^{2d} is $-C(=O)OH$. In certain more specific embodiments, at least one of R^{2a} , R^{2b} , R^{2c} , and R^{2d} is $-C(=O)OCH_3$ or $-OC(=O)CH_3$. In some embodiments, at least one of R^{2a} , R^{2b} , R^{2c} , and R^{2d} is $-C(=O)N(R')_2$ or $-NR'(C=O)alkyl$ wherein R' is, at each occurrence, independently a C_1 - C_6 alkyl. In more specific embodiments, each occurrence of R' is methyl. In some embodiments, at least one of R^{2a} , R^{2b} , R^{2c} , and R^{2d} is $-C(=O)NR''heterocyclyl$ wherein R'' is hydrogen or C_1 - C_6 alkyl (*e.g.*, methyl, ethyl, or isopropyl).

In some specific embodiments, at least one of R^{2a} , R^{2b} , R^{2c} , and R^{2d} is a substituted or unsubstituted phenyl. In some embodiments, at least one of R^{2a} , R^{2b} , R^{2c} , and R^{2d} is a substituted or unsubstituted heterocyclyl. In some embodiments, at least one of R^{2a} , R^{2b} , R^{2c} , and R^{2d} is a substituted or unsubstituted heteroaryl (*e.g.*, tetrazolyl, triazolyl, oxadiazolyl, imidazolidinyl, pyrazolyl, etc.). In some embodiments, R^{2a} is heteroaryl or heterocyclyl. In some more specific embodiments, R^{2c} is heteroaryl or heterocyclyl. In some embodiments, R^{2a} is heterocyclyl. In some more specific embodiments, R^{2c} is heterocyclyl. In certain specific embodiments, R^{2a} is heteroaryl. In certain more specific embodiments, R^{2c} is heteroaryl.

In some embodiments, at least one of R^{2a} , R^{2b} , R^{2c} , and R^{2d} has one of the following structures:




In more specific embodiments, at least one of R^{2a} , R^{2b} , R^{2c} , and R^{2d} has
 5 one of the following structures:



10


In some more specific embodiments, R^3 has one of the following
 structures:





In certain embodiments, R³ is -NH₂.

5 In some embodiments, n is 0. In certain embodiments, n is 1. In still other embodiments, n is 2. In some embodiments, n is 3, 4, 5, or 6.


In some embodiments, n is 2 and R⁴ is a substituted or unsubstituted aryl. In some more specific embodiments, R⁴ is a substituted aryl. In certain more specific embodiments, R⁴ is substituted with a substituent selected from the group consisting of alkyl, halo (*e.g.*, fluoro, chloro, bromo), haloalkyl, hydroxyl, amino, nitro, and combinations thereof. In some embodiments, R⁴ is an unsubstituted aryl. In certain embodiments, R⁴ has one of the following structures:










5 In some more specific embodiments, R^4 has one of the following structures:




In certain specific embodiments, n is 2 and R^4 is a substituted or unsubstituted heteroaryl. In some more specific embodiments, R^4 has one of the following structures:










In some embodiments, R⁴ has one of the following structures:








In some embodiments, R^4 has one of the following structures:




In certain related embodiments, n is 0 and R^4 has one of the following structures:




5

In some specific embodiments, R^4 has one of the following structures:



One embodiment provides a compound having the following Structure

(II):



10

(II)

or a stereoisomer, tautomer, or pharmaceutically acceptable salt thereof, wherein:

X is a direct bond, $-[C(R^{2e})R^{2f}]$ -, or $-[C(R^{2e})R^{2f}]-[C(R^{2g})R^{2h}]$;

Y is a direct bond or $-[C(R^{2i})R^{2j}]$;

R^1 is a substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, substituted or unsubstituted cycloalkyl, or a substituted or unsubstituted heterocyclyl;

R^{2a} , R^{2b} , R^{2c} , R^{2d} , R^{2e} , R^{2f} , R^{2g} , R^{2h} , R^{2i} , and R^{2j} are each independently
 5 selected from the group consisting of hydrogen, halo, OR^5 , $C(=O)OR^5$, $OC(=O)R^5$, hydroxyalkyl, alkoxy, alkoxyalkyl, haloalkoxy, cyano, aminylalkyl, carboxyalkyl, NR^5R^6 , $C(=O)NR^5R^6$, $N(R^5)C(=O)R^6$, $NR^5C(=O)NR^6$, $S(O)_t$, SR^5 , nitro, $N(R^5)C(O)OR^6$, $C(=NR^5)NR^6R^7$, $N(R^5)C(=NR^6)NR^7R^8$, $S(O)R^5$, $S(O)NR^5R^6$, $S(O)_2R^5$, $N(R^5)S(O)_2R^6$, $S(O)_2NR^5R^6$, aryl, heteroaryl, heterocyclyl, cycloalkyl, and oxo, provided that at least
 10 one of R^{2a} , R^{2b} , R^{2c} , R^{2d} , R^{2e} , R^{2f} , R^{2g} , R^{2h} , R^{2i} , and R^{2j} is not hydrogen;

R^3 is $NR^{3a}R^{3b}$;

R^{3a} and R^{3b} are each independently hydrogen, alkyl, hydroxyalkyl, haloalkyl, alkoxyalkyl, $-CH_2C(C=O)OH$, $-CH_2C(=O)Oalkyl$, heterocyclyl, heteroaryl, heterocyclylalkyl, heteroarylalkyl, or cycloalkyl; or R^{3a} and R^{3b} , together with the
 15 nitrogen to which they are attached, form an optionally substituted 4-7 membered heteroaryl or an optionally substituted 4-7 membered heterocyclyl;

R^4 is a substituted or unsubstituted aryl, a substituted or unsubstituted heteroaryl, a substituted or unsubstituted cycloalkyl, or a substituted or unsubstituted heterocyclyl when n is 2, 3, 4, 5, or 6; or

20 R^4 is a substituted or unsubstituted monocyclic heteroaryl, or a substituted or unsubstituted heterocyclyl when n is 0 or 1;


R^5 , R^6 , R^7 , and R^8 are, at each occurrence, independently hydrogen, alkyl, hydroxyalkyl, haloalkyl, alkoxyalkyl, carboxyalkyl, heterocyclyl, heteroaryl, or cycloalkyl;

25 n is an integer from 0-6; and

t is an integer from 1-3,


provided that:

a) when one of R^{2a} , R^{2b} , R^{2c} , R^{2d} , R^{2e} , R^{2f} , R^{2g} , R^{2h} , R^{2i} , and R^{2j} is OH, R^1 does not have the following structure:




b) when one of R^{2a} , R^{2b} , R^{2c} , R^{2d} , R^{2e} , R^{2f} , R^{2g} , and R^{2h} is -OH, n is an integer from 2-6; and

c) when one of R^{2a} , R^{2b} , R^{2c} , R^{2d} , R^{2e} , R^{2f} , R^{2g} , R^{2h} , R^{2i} , and R^{2j} is an unsubstituted phenyl, neither R^{3a} nor R^{3b} has the following structure:



In some embodiments, none of R^{2a} , R^{2b} , R^{2c} , R^{2d} , R^{2e} , R^{2f} , R^{2g} , R^{2h} , R^{2i} , or R^{2j} are alkyl. In some more embodiments, none of R^{2a} , R^{2b} , R^{2c} , R^{2d} , R^{2e} , R^{2f} , R^{2g} , R^{2h} , R^{2i} , or R^{2j} are alkenyl. In some embodiments, none of R^{2a} , R^{2b} , R^{2c} , R^{2d} , R^{2e} , R^{2f} , R^{2g} , R^{2h} , R^{2i} , or R^{2j} are arylalkyl. In certain embodiments, none of R^{2a} , R^{2b} , R^{2c} , R^{2d} , R^{2e} , R^{2f} , R^{2g} , R^{2h} , R^{2i} , or R^{2j} are arylalkoxy. In some embodiments, none of R^{2a} , R^{2b} , R^{2c} , R^{2d} , R^{2e} , R^{2f} , R^{2g} , R^{2h} , R^{2i} , or R^{2j} are cycloalkylalkoxy. In some embodiments, none of R^{2a} , R^{2b} , R^{2c} , R^{2d} , R^{2e} , R^{2f} , R^{2g} , R^{2h} , R^{2i} , or R^{2j} are halo (*e.g.*, fluoro). In some embodiments, none of R^{2a} , R^{2b} , R^{2c} , R^{2d} , R^{2e} , R^{2f} , R^{2g} , R^{2h} , R^{2i} , or R^{2j} are hydroxyl. In some embodiments, none of R^{2a} , R^{2b} , R^{2c} , R^{2d} , R^{2e} , R^{2f} , R^{2g} , R^{2h} , R^{2i} , or R^{2j} are alkoxy. In certain specific embodiments, none of R^{2a} , R^{2b} , R^{2c} , R^{2d} , R^{2e} , R^{2f} , R^{2g} , R^{2h} , R^{2i} , or R^{2j} have the following structure:





In still other embodiments, neither R^{3a} nor R^{3b} are carboxyalkyl. In some more embodiments, neither R^{3a} nor R^{3b} are arylalkylcarbonyl. In certain embodiments, neither R^{3a} nor R^{3b} are alkoxy carbonyl (*e.g.*, Boc). In some embodiments, neither R^{3a} nor R^{3b} are heteroarylsulfo or alkylsulfo (*e.g.*, $-S(O)_2CH_3$). In some embodiments, neither R^{3a} nor R^{3b} are arylalkoxy carbonyl (*e.g.*, Fmoc).

In some embodiments, R^1 is a substituted or unsubstituted aryl. In certain embodiments, R^1 is a substituted or unsubstituted C_6 - C_{10} aryl. In some embodiments, R^1 is a substituted or unsubstituted phenyl. In certain embodiments, R^1 is a substituted phenyl. In some more specific embodiments, R^1 is phenyl substituted with one or more
 5 of R^{1a} , R^{1b} , R^{1c} , R^{1d} , and R^{1e} wherein R^{1a} , R^{1b} , R^{1c} , R^{1d} , and R^{1e} are each independently selected from the group consisting of C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, halo, C_{1-6} haloalkyl, aminylalkyl, hydroxyalkyl, cyano, nitro, OR^9 , SR^9 , $C(O)R^9$, $C(O)NR^9R^{10}$, $C(O)OR^9$, $OC(O)R^9$, $OC(O)OR^9$, $OC(O)NR^9R^{10}$, NR^9R^{10} , $N(R^9)C(O)R^{10}$, $N(R^9)C(O)NR^{10}R^{11}$, $N(R^9)C(O)OR^{10}$, $C(=NR^9)NR^{10}R^{11}$, $C(=NOR^9)NR^{10}R^{11}$,
 10 $C(=NOC(O)R^9)NR^{10}R^{11}$, $C(=NR^9)N(R^{10})C(O)OR^{11}$, $N(R^9)C(=NR^{10})NR^{11}R^{12}$, $S(O)R^9$, $S(O)NR^9R^{10}$, $S(O)_2R^9$, $N(R^9)S(O)_2R^{10}$, $S(O)_2NR^9R^{10}$, oxo, substituted or unsubstituted C_{6-10} aryl, substituted or unsubstituted C_{6-10} arylalkyl, substituted or unsubstituted C_{6-10} aryloxy, substituted or unsubstituted C_{6-10} arylalkoxy, substituted or unsubstituted 5-10 membered heteroaryl, substituted or unsubstituted C_{3-10} cycloalkyl, and substituted or
 15 unsubstituted 4-10 membered heterocyclyl, wherein R^9 , R^{10} , R^{11} , and R^{12} , are, at each occurrence, independently selected from the group consisting of hydrogen, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, hydroxyl, C_{1-6} alkoxy, aryl, arylalkyl, C_{1-6} haloalkyl, C_{1-6} haloalkoxy, C_{1-6} hydroxyalkyl, cycloalkyl, heterocyclyl, and heteroaryl.


In some more specific embodiments, R^{1a} , R^{1b} , R^{1c} , R^{1d} , or R^{1e} is optionally
 20 substituted with one or more substituents selected from the group consisting of halo, CN, OR^{13} , SR^{13} , $C(O)R^{13}$, $C(O)NR^{13}R^{14}$, $C(O)OR^{13}$, $OC(O)R^{13}$, $OC(O)NR^{13}R^{14}$, $NR^{13}R^{14}$, $NR^{13}C(O)R^{14}$, $NR^{13}C(O)NR^{14}R^{15}$, $NR^{13}C(O)OR^{14}$, $C(=NR^{13})NR^{14}R^{15}$, $NR^{13}C(=NR^{14})NR^{15}R^{16}$, $S(O)R^{13}$, $S(O)NR^{13}R^{14}$, $S(O)_2R^{13}$, $NR^{13}S(O)_2R^{14}$, $S(O)_2NR^{13}R^{14}$ and oxo when R^{1a} , R^{1b} , R^{1c} , R^{1d} , or R^{1e} is a substituted C_{6-10} aryl, a substituted C_{6-10}
 25 arylalkyl, a substituted C_{6-10} aryloxy, a substituted C_{6-10} arylalkoxy, a substituted 5-10 membered heteroaryl, a substituted C_{3-10} cycloalkyl, and a substituted 4-10 membered heterocyclyl, wherein R^{13} , R^{14} , R^{15} , and R^{16} are, at each occurrence, independently selected from the group consisting of hydrogen, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, hydroxyl, C_{1-6} alkoxy, aryl, arylalkyl, C_{1-6} haloalkyl, C_{1-6} haloalkoxy, C_{1-6} hydroxyalkyl,
 30 cycloalkyl, heterocyclyl, and heteroaryl.

In certain more specific embodiments, R^1 is phenyl substituted with at least one substituent selected from the group consisting of OR^9 , $C(=NR^9)NR^{10}R^{11}$, $C(=NOR^9)NR^{10}R^{11}$, $C(=NOC(O)R^9)NR^{10}R^{11}$, and $C(=NR^9)NR^{10}C(O)OR^{11}$. In some embodiments, R^1 is substituted with at least one substituent selected from the group consisting of $C(=NR^9)NR^{10}R^{11}$, $C(=NOR^9)NR^{10}R^{11}$, $C(=NOC(O)R^9)NR^{10}R^{11}$, and $C(=NR^9)NR^{10}C(O)OR^{11}$. In some embodiments, R^1 is substituted with at least one $C(=NR^9)NR^{10}R^{11}$. In more specific embodiments, R^1 is substituted with at least one $C(=NH)NH_2$. In still more specific embodiments, R^1 has one of the following structures:






In some embodiments, R¹ has one of the following structures:



In certain specific embodiments, R¹ has one of the following structures:



In some of the foregoing embodiments, R^9 is C_{1-6} alkyl or C_{1-6} haloalkyl. In more specific embodiments, R^9 is methyl. In certain embodiments, R^9 is trifluoromethyl. In some embodiments, R^1 is an unsubstituted phenyl.


In some embodiments, R^1 is a substituted or unsubstituted heteroaryl. In certain embodiments, R^1 is a substituted or unsubstituted 5-10 membered heteroaryl. In some embodiments, R^1 is a substituted or unsubstituted pyridinyl, pyrrolopyridinyl, or benzoimidazolyl. In some more specific embodiments, R^1 is a substituted or unsubstituted pyridin-3-yl, 1H-pyrrolo[2,3-b]pyridine-5-yl, 1H-pyrrolo[3,2-c]pyridine-2-yl, or 1H-benzo[d]imidazol-6-yl.


In some embodiments, R^1 is pyridin-3-yl, 1H-pyrrolo[2,3-b]pyridine-5-yl, or 1H-benzo[d]imidazol-6-yl substituted with one or more of R^{1a} , R^{1b} , R^{1c} , R^{1d} , and R^{1e} wherein R^{1a} , R^{1b} , R^{1c} , R^{1d} , and R^{1e} are each independently selected from the group consisting of C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, halo, C_{1-6} haloalkyl, aminylalkyl, hydroxyalkyl, cyano, nitro, OR^9 , SR^9 , $C(O)R^9$, $C(O)NR^9R^{10}$, $C(O)OR^9$, $OC(O)R^9$, $OC(O)OR^9$, $OC(O)NR^9R^{10}$, NR^9R^{10} , $N(R^9)C(O)R^{10}$, $N(R^9)C(O)NR^{10}R^{11}$, $N(R^9)C(O)OR^{10}$, $C(=NR^9)NR^{10}R^{11}$, $C(=NOR^9)NR^{10}R^{11}$, $C(=NOC(O)R^9)NR^{10}R^{11}$, $C(=NR^9)N(R^{10})C(O)OR^{11}$, $N(R^9)C(=NR^{10})NR^{11}R^{12}$, $S(O)R^9$, $S(O)NR^9R^{10}$, $S(O)_2R^9$, $N(R^9)S(O)_2R^{10}$, $S(O)_2NR^9R^{10}$, oxo, substituted or unsubstituted C_{6-10} aryl, substituted or unsubstituted C_{6-10} arylalkyl, substituted or unsubstituted C_{6-10} aryloxy, substituted or unsubstituted C_{6-10} arylalkoxy, substituted or unsubstituted 5-10 membered heteroaryl, substituted or unsubstituted C_{3-10} cycloalkyl, and substituted or unsubstituted 4-10 membered heterocyclyl, wherein R^9 , R^{10} , R^{11} , and R^{12} , are, at each occurrence,


independently selected from the group consisting of hydrogen, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, hydroxyl, C₁₋₆ alkoxy, aryl, arylalkyl, C₁₋₆ haloalkyl, C₁₋₆ haloalkoxy, C₁₋₆ hydroxyalkyl, cycloalkyl, heterocyclyl, and heteroaryl.


In more specific embodiments, R^{1a}, R^{1b}, R^{1c}, R^{1d}, or R^{1e} is optionally substituted with one or more substituents selected from the group consisting of halo, CN, OR¹³, SR¹³, C(O)R¹³, C(O)NR¹³R¹⁴, C(O)OR¹³, OC(O)R¹³, OC(O)NR¹³R¹⁴, NR¹³R¹⁴, NR¹³C(O)R¹⁴, NR¹³C(O)NR¹⁴R¹⁵, NR¹³C(O)OR¹⁴, C(=NR¹³)NR¹⁴R¹⁵, NR¹³C(=NR¹⁴)NR¹⁵R¹⁶, S(O)R¹³, S(O)NR¹³R¹⁴, S(O)₂R¹³, NR¹³S(O)₂R¹⁴, S(O)₂NR¹³R¹⁴ and oxo when R^{1a}, R^{1b}, R^{1c}, R^{1d}, or R^{1e} is a substituted C₆₋₁₀ aryl, a substituted C₆₋₁₀ arylalkyl, a substituted C₆₋₁₀ aryloxy, a substituted C₆₋₁₀ arylalkoxy, a substituted 5-10 membered heteroaryl, a substituted C₃₋₁₀ cycloalkyl, and a substituted 4-10 membered heterocyclyl, wherein R¹³, R¹⁴, R¹⁵, and R¹⁶ are, at each occurrence, independently selected from the group consisting of hydrogen, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, hydroxyl, C₁₋₆ alkoxy, aryl, arylalkyl, C₁₋₆ haloalkyl, C₁₋₆ haloalkoxy, C₁₋₆ hydroxyalkyl, cycloalkyl, heterocyclyl, and heteroaryl.

In some specific embodiments, R¹ has one of the following structures:











In some specific embodiments, R^{1a} or R^{1b} is independently C₁₋₆ alkyl, amino, C(O)R⁹, C(O)NR⁹R¹⁰, N(R⁹)C(O)NR¹⁰R¹¹, N(R⁹)C(O)OR¹⁰, or OC(O)NR⁹R¹⁰, or halo. In more specific embodiments, R^{1a} or R^{1b} is methyl or ethyl. In other specific embodiments, R^{1a} or R^{1b} is F, Cl, or Br.


In some embodiments, each R^{1a} or R^{1b} attached to nitrogen is C₁₋₆ alkyl, C(O)R⁹, C(O)NR⁹R¹⁰, N(R⁹)C(O)NR¹⁰R¹¹, N(R⁹)C(O)OR¹⁰, or OC(O)NR⁹R¹⁰. In certain embodiments, R^{1a} or R^{1b} is methyl or ethyl.


In some embodiments, R^1 has one of the following structures:







In some specific embodiments, R¹ has one of the following structures:







In some embodiments, R^1 has one of the following structures:



In some embodiments, R^1 is a substituted or unsubstituted cycloalkyl. In certain specific embodiments, R^1 is a substituted or unsubstituted C_3 - C_6 cycloalkyl. In some embodiments, R^1 is a substituted C_3 - C_6 cycloalkyl.

In some more specific embodiments, R^1 is a C_3 - C_6 cycloalkyl substituted with one or more of R^{1a} , R^{1b} , R^{1c} , R^{1d} , and R^{1e} wherein R^{1a} , R^{1b} , R^{1c} , R^{1d} , and R^{1e} are each independently selected from the group consisting of C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} haloalkyl, aminylalkyl, hydroxyalkyl, nitro, OR^9 , SR^9 , $C(O)R^9$, $C(O)NR^9R^{10}$, $C(O)OR^9$, $OC(O)R^9$, $OC(O)OR^9$, $OC(O)NR^9R^{10}$, NR^9R^{10} , $N(R^9)C(O)R^{10}$, $N(R^9)C(O)NR^{10}R^{11}$, $N(R^9)C(O)OR^{10}$, $C(=NR^9)NR^{10}R^{11}$, $C(=NOR^9)NR^{10}R^{11}$, $C(=NOC(O)R^9)NR^{10}R^{11}$, $C(=NR^9)N(R^{10})C(O)OR^{11}$, $N(R^9)C(=NR^{10})NR^{11}R^{12}$, $S(O)R^9$,

S(O)NR⁹R¹⁰, S(O)₂R⁹, N(R⁹)S(O)₂R¹⁰, S(O)₂NR⁹R¹⁰, oxo, substituted or unsubstituted C₆₋₁₀ aryl, substituted or unsubstituted C₆₋₁₀ arylalkyl, substituted or unsubstituted C₆₋₁₀ aryloxy, substituted or unsubstituted C₆₋₁₀ arylalkoxy, substituted or unsubstituted 5-10 membered heteroaryl, substituted or unsubstituted C₃₋₁₀ cycloalkyl, and substituted or unsubstituted 4-10 membered heterocyclyl, wherein R⁹, R¹⁰, R¹¹, and R¹², are, at each occurrence, independently selected from the group consisting of hydrogen, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, hydroxyl, C₁₋₆ alkoxy, aryl, arylalkyl, C₁₋₆ haloalkyl, C₁₋₆ haloalkoxy, C₁₋₆ hydroxyalkyl, cycloalkyl, heterocyclyl, and heteroaryl.

In some more specific embodiments, R^{1a}, R^{1b}, R^{1c}, R^{1d}, or R^{1e} is optionally substituted with one or more substituents selected from the group consisting of OR¹³, SR¹³, C(O)R¹³, C(O)NR¹³R¹⁴, C(O)OR¹³, OC(O)R¹³, OC(O)NR¹³R¹⁴, NR¹³R¹⁴, NR¹³C(O)R¹⁴, NR¹³C(O)NR¹⁴R¹⁵, NR¹³C(O)OR¹⁴, C(=NR¹³)NR¹⁴R¹⁵, NR¹³C(=NR¹⁴)NR¹⁵R¹⁶, S(O)R¹³, S(O)NR¹³R¹⁴, S(O)₂R¹³, NR¹³S(O)₂R¹⁴, S(O)₂NR¹³R¹⁴ and oxo when R^{1a}, R^{1b}, R^{1c}, R^{1d}, or R^{1e} is a substituted C₆₋₁₀ aryl, a substituted C₆₋₁₀ arylalkyl, a substituted C₆₋₁₀ aryloxy, a substituted C₆₋₁₀ arylalkoxy, a substituted 5-10 membered heteroaryl, a substituted C₃₋₁₀ cycloalkyl, and a substituted 4-10 membered heterocyclyl, wherein R¹³, R¹⁴, R¹⁵, and R¹⁶ are, at each occurrence, independently selected from the group consisting of hydrogen, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, hydroxyl, C₁₋₆ alkoxy, aryl, arylalkyl, C₁₋₆ haloalkyl, C₁₋₆ haloalkoxy, C₁₋₆ hydroxyalkyl, cycloalkyl, heterocyclyl, and heteroaryl.

In some embodiments, R¹ is an unsubstituted C₃₋₆ cycloalkyl.

In some embodiments, R¹ is a substituted or unsubstituted heterocyclyl.

In certain embodiments, R¹ is a substituted or unsubstituted 4-10 membered heterocyclyl.

In some more specific embodiments, R¹ is a substituted 4-10 membered heterocyclyl.









In some embodiments, R¹ is a 4-10 membered heterocyclyl substituted with one or more of R^{1a}, R^{1b}, R^{1c}, R^{1d}, and R^{1e} wherein R^{1a}, R^{1b}, R^{1c}, R^{1d}, and R^{1e} are each independently selected from the group consisting of C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, halo, C₁₋₆ haloalkyl, aminylalkyl, hydroxyalkyl, cyano, nitro, OR⁹, SR⁹, C(O)R⁹, C(O)NR⁹R¹⁰, C(O)OR⁹, OC(O)R⁹, OC(O)OR⁹, OC(O)NR⁹R¹⁰, NR⁹R¹⁰, N(R⁹)C(O)R¹⁰, N(R⁹)C(O)NR¹⁰R¹¹, N(R⁹)C(O)OR¹⁰, C(=NR⁹)NR¹⁰R¹¹, C(=NOR⁹)NR¹⁰R¹¹,





$C(=NOC(O)R^9)NR^{10}R^{11}$, $C(=NR^9)N(R^{10})C(O)OR^{11}$, $N(R^9)C(=NR^{10})NR^{11}R^{12}$, $S(O)R^9$,
 $S(O)NR^9R^{10}$, $S(O)_2R^9$, $N(R^9)S(O)_2R^{10}$, $S(O)_2NR^9R^{10}$, oxo, substituted or unsubstituted
 C_{6-10} aryl, substituted or unsubstituted C_{6-10} arylalkyl, substituted or unsubstituted C_{6-10}
 aryloxy, substituted or unsubstituted C_{6-10} arylalkoxy, substituted or unsubstituted 5-10
 5 membered heteroaryl, substituted or unsubstituted C_{3-10} cycloalkyl, and substituted or
 unsubstituted 4-10 membered heterocyclyl, wherein R^9 , R^{10} , R^{11} , and R^{12} , are, at each
 occurrence, independently selected from the group consisting of hydrogen, C_{1-6} alkyl, C_{2-6}
 alkenyl, C_{2-6} alkynyl, hydroxyl, C_{1-6} alkoxy, aryl, arylalkyl, C_{1-6} haloalkyl, C_{1-6}
 haloalkoxy, C_{1-6} hydroxyalkyl, cycloalkyl, heterocyclyl, and heteroaryl.

10 In more specific embodiments, R^{1a} , R^{1b} , R^{1c} , R^{1d} , or R^{1e} is optionally
 substituted with one or more substituents selected from the group consisting of halo, CN,
 OR^{13} , SR^{13} , $C(O)R^{13}$, $C(O)NR^{13}R^{14}$, $C(O)OR^{13}$, $OC(O)R^{13}$, $OC(O)NR^{13}R^{14}$, $NR^{13}R^{14}$,
 $NR^{13}C(O)R^{14}$, $NR^{13}C(O)NR^{14}R^{15}$, $NR^{13}C(O)OR^{14}$, $C(=NR^{13})NR^{14}R^{15}$,
 $NR^{13}C(=NR^{14})NR^{15}R^{16}$, $S(O)R^{13}$, $S(O)NR^{13}R^{14}$, $S(O)_2R^{13}$, $NR^{13}S(O)_2R^{14}$, $S(O)_2NR^{13}R^{14}$
 15 and oxo when R^{1a} , R^{1b} , R^{1c} , R^{1d} , or R^{1e} is a substituted C_{6-10} aryl, a substituted C_{6-10}
 arylalkyl, a substituted C_{6-10} aryloxy, a substituted C_{6-10} arylalkoxy, a substituted 5-10
 membered heteroaryl, a substituted C_{3-10} cycloalkyl, and a substituted 4-10 membered
 heterocyclyl, wherein R^{13} , R^{14} , R^{15} , and R^{16} are, at each occurrence, independently
 selected from the group consisting of hydrogen, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl,
 20 hydroxyl, C_{1-6} alkoxy, aryl, arylalkyl, C_{1-6} haloalkyl, C_{1-6} haloalkoxy, C_{1-6} hydroxyalkyl,
 cycloalkyl, heterocyclyl, and heteroaryl.

In some embodiments, R^1 is an unsubstituted 4-10 membered heterocyclyl.

In certain embodiments, the compound has one of the following
 25 structures:







In some embodiments, at least one occurrence of R^{2a} , R^{2b} , R^{2c} , or R^{2d} is alkoxy, alkoxyalkyl, cyano, $C(=O)OR^5$, $OC(=O)R^5$, $C(=O)NR^5R^6$, $N(R^5)C(=O)R^6$, aryl, heteroaryl, heterocyclyl, or $C(=O)N(R^5)$ heterocyclyl, wherein R^5 and R^6 are, at each occurrence, independently hydrogen or C_1 - C_6 alkyl. In certain embodiments, at least one occurrence of R^{2a} , R^{2b} , R^{2c} , or R^{2d} is methoxy. In some embodiments, at least one occurrence of R^{2a} , R^{2b} , R^{2c} , or R^{2d} is methoxymethyl. In certain embodiments, at least one occurrence of R^{2a} , R^{2b} , R^{2c} , or R^{2d} is cyano. In some embodiments, at least one occurrence of R^{2a} , R^{2b} , R^{2c} , or R^{2d} is $-C(=O)OH$. In certain embodiments, at least one occurrence of R^{2a} , R^{2b} , R^{2c} , or R^{2d} is $-C(=O)OCH_3$ or $-OC(=O)CH_3$. In some more specific embodiments, at least one occurrence of R^{2a} , R^{2b} , R^{2c} , or R^{2d} is $-C(=O)N(R')_2$ or $-NR'(C=O)$ alkyl wherein R' is, at each occurrence, independently a C_1 - C_6 alkyl (*e.g.*, methyl, ethyl, or isopropyl). In some of the foregoing embodiments, each occurrence of R' is methyl. In certain embodiments, at least one occurrence of R^{2a} , R^{2b} , R^{2c} , or R^{2d} is $-C(=O)NR''$ heterocyclyl wherein R'' is hydrogen or C_1 - C_6 alkyl (*e.g.*, methyl, ethyl, or isopropyl).


In some embodiments, at least one occurrence of R^{2a}, R^{2b}, R^{2c}, or R^{2d} is a substituted or unsubstituted phenyl. In certain embodiments, at least one occurrence of R^{2a}, R^{2b}, R^{2c}, or R^{2d} is a substituted or unsubstituted heterocyclyl. In some embodiments, at least one occurrence of R^{2a}, R^{2b}, R^{2c}, or R^{2d} is a substituted or unsubstituted heteroaryl.

5 In some embodiments, at least one occurrence of R^{2a} , R^{2b} , R^{2c} , or R^{2d} has one of the following structures:




10 In some embodiments, at least one occurrence of R^{2a} , R^{2b} , R^{2c} , or R^{2d} has one of the following structures:






In some embodiments, $\mathbf{R}^3$ has one of the following structures:




In some embodiments, R^3 is $-NH_2$. In certain embodiments, n is 0. In some other embodiments, n is 1. In certain other embodiments, n is 2. In still other embodiments, n is 3, 4, 5, or 6.


10 In some more specific embodiments, n is 2 and R⁴ is a substituted or unsubstituted aryl. In certain more specific embodiments, R⁴ has one of the following structures:

5








In certain embodiments, R⁴ has one of the following structures:




In some embodiments, n is 2 and R^4 is a substituted or unsubstituted heteroaryl. In some more specific embodiments, R^4 has one of the following structures:










In some embodiments, R^4 has one of the following structures:








In certain specific embodiments, R^4 has one of the following structures:




In some embodiments, n is 0 and R^4 has one of the following structures:




5

In some specific embodiments, R^4 has one of the following structures:



Another embodiment provides a compound having the following Structure (III):



10

(III)

or a stereoisomer, tautomer, or pharmaceutically acceptable salt thereof, wherein:

X is a direct bond, $-CR^{2e}R^{2f}-$, or $-CR^{2e}R^{2f}-CR^{2g}R^{2h}-$;

Y is a direct bond or $-CR^{2i}R^{2j}-$;

Z is O or S;

R^{17} is a substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, substituted or unsubstituted cycloalkyl, or a substituted or unsubstituted heterocyclyl;

R^{18a} , R^{18b} , R^{18c} , R^{18d} , R^{18e} , R^{18f} , R^{18g} , R^{18h} , R^{18i} , and R^{18j} are each
 5 independently selected from the group consisting of hydrogen, halo, $-OR^{21}$, $C(=O)OR^{21}$, $OC(=O)R^{21}$, hydroxyalkyl, alkoxy, alkoxyalkyl, haloalkoxy, cyano, aminylalkyl, carboxyalkyl, $NR^{21}R^{22}$, $C(=O)NR^{21}R^{22}$, $N(R^{21})C(=O)R^{22}$, $NR^{21}C(=O)NR^{22}$, $S(O)_t$, SR^{21} , nitro, $N(R^{21})C(O)OR^{22}$, $C(=NR^{21})NR^{22}R^{23}$, $N(R^{21})C(=NR^{22})NR^{23}R^{24}$, $S(O)R^{21}$, $S(O)NR^{21}R^{22}$, $S(O)_2R^{21}$, $N(R^{21})S(O)_2R^{22}$, $S(O)_2NR^{21}R^{22}$, aryl, heteroaryl, heterocyclyl,
 10 cycloalkyl, and oxo, provided that at least one of R^{18a} , R^{18b} , R^{18c} , R^{18d} , R^{18e} , R^{18f} , R^{18g} , R^{18h} , R^{18i} , and R^{18j} is not hydrogen;

R^{19} is $NR^{19a}R^{19b}$.

R^{19a} and R^{19b} are each independently hydrogen, alkyl, hydroxyalkyl, haloalkyl, alkoxyalkyl, heterocyclyl, heteroaryl, heterocyclylalkyl, heteroarylalkyl,
 15 cycloalkyl, $(CH_2)_nC(=O)OR^5$, or $(CH_2)_nP(=O)(OR^5)_2$; or R^{19a} and R^{19b} , together with the nitrogen to which they are attached, form an optionally substituted 4-7 membered heteroaryl or an optionally substituted 4-7 membered heterocyclyl;


R^{20} is a substituted or unsubstituted aryl, a substituted or unsubstituted heteroaryl, a substituted or unsubstituted cycloalkyl, or a substituted or unsubstituted
 20 heterocyclyl;

R^{21} , R^{22} , R^{23} , and R^{24} are, at each occurrence, independently hydrogen, alkyl, hydroxyalkyl, haloalkyl, alkoxyalkyl, carboxyalkyl, heterocyclyl, heteroaryl, or cycloalkyl;


m is an integer from 0-6; and


25 t is an integer from 1-3.

In some embodiments, R^{17} has one of the following structures:





5


In some embodiments, R¹⁷ has one of the following structures:




In certain embodiments, R^{17} is an unsubstituted phenyl. In some more specific embodiments, R^{17} has one of the following structures:









In some more specific embodiments, R^{17} has one of the following structures:




In some embodiments, at least one of R^{18a} and R^{18b} is alkoxy, alkoxyalkyl, cyano, $C(=O)OR^{21}$, $OC(=O)R^{21}$, $C(=O)NR^{21}R^{22}$, $N(R^{21})C(=O)R^{22}$, aryl, heteroaryl, heterocyclyl, or $C(=O)N(R^{21})$ heterocyclyl, wherein R^{21} and R^{22} are, at each occurrence, independently hydrogen or C_1 - C_6 alkyl. In certain more specific embodiments, at least one of R^{18a} and R^{18b} has one of the following structures:





In some embodiments, R^{19} has one of the following structures:



In certain embodiments, R^{20} has one of the following structures:










5


In some embodiments, R^{20} has one of the following structures:










In some embodiments, $\mathbf{R}^{20}$ has one of the following structures:




5







Another embodiment provides a compound having the following Structure (III):



(III)

or a stereoisomer, tautomer, or pharmaceutically acceptable salt thereof, wherein:

X is a direct bond or -CR^{26c}R^{26d}-;

R²⁵ is a substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, substituted or unsubstituted cycloalkyl, or a substituted or unsubstituted heterocyclyl;

R^{26a} , R^{26b} , R^{26c} , and R^{26d} are each independently selected from the group consisting of hydrogen, halo, $-OR^{29}$, $C(=O)OR^{29}$, $OC(=O)R^{29}$, hydroxyalkyl, alkoxy, alkoxyalkyl, haloalkoxy, cyano, aminylalkyl, carboxyalkyl, $NR^{29}R^{30}$, $C(=O)NR^{29}R^{30}$, $N(R^{29})C(=O)R^{30}$, $NR^{29}C(=O)NR^{30}$, $S(O)_t$, SR^{29} , nitro, $N(R^{29})C(O)OR^{30}$, $C(=NR^{29})NR^{30}R^{31}$, $N(R^{29})C(=NR^{30})NR^{31}R^{32}$, $S(O)R^{29}$, $S(O)NR^{29}R^{30}$, $S(O)_2R^{30}$, $N(R^{29})S(O)_2R^{30}$, $S(O)_2NR^{29}R^{30}$, aryl, heteroaryl, heterocyclyl, cycloalkyl, and oxo, provided that at least one of R^{26a} , R^{26b} , R^{26c} , and R^{26d} is not hydrogen;

$$\mathbf{R}^{27} \text{ is } \mathbf{NR}^{27a}\mathbf{R}^{27b}.$$

20 R^{27a} and R^{27b} are each independently hydrogen, alkyl, hydroxyalkyl, haloalkyl, alkoxyalkyl, heterocyclyl, heteroaryl, heterocyclylalkyl, heteroarylalkyl, cycloalkyl, (CH₂)_nC(=O)OR²⁹, or (CH₂)_nP(=O)(OR²⁹)₂; or R^{27a} and R^{27b}, together with the nitrogen to which they are attached, form an optionally substituted 4-7 membered heteroaryl or an optionally substituted 4-7 membered heterocyclyl;


R^{28} is a substituted or unsubstituted aryl, a substituted or unsubstituted heteroaryl, a substituted or unsubstituted cycloalkyl, or a substituted or unsubstituted heterocyclyl;

R^{29} , R^{30} , R^{31} , and R^{32} are, at each occurrence, independently hydrogen, alkyl, hydroxyalkyl, haloalkyl, alkoxyalkyl, carboxyalkyl, heterocyclyl, heteroaryl, or cycloalkyl;

p is an integer from 0-6; and


t is an integer from 1-3.


In some embodiments, R^{25} has one of the following structures:




In some embodiments, R^{25} is an unsubstituted phenyl.


15 In some more specific embodiments, R^{25} has one of the following structures:









In some embodiments, R^{25} has one of the following structures:






In certain embodiments, at least one of R^{26a} and R^{26b} is alkoxy, alkoxyalkyl, cyano, $C(=O)OR^{29}$, $OC(=O)R^{29}$, $C(=O)NR^{29}R^{30}$, $N(R^{29})C(=O)R^{30}$, aryl, heteroaryl, heterocyclyl, or $C(=O)N(R^{29})$ heterocyclyl, wherein R^{29} and R^{30} are, at each occurrence, independently hydrogen or C_1 - C_6 alkyl. In more specific embodiments, at least one occurrence of R^{26a} or R^{26b} has one of the following structures:





10

In certain embodiments, R^{27} has one of the following structures:





In some embodiments, R^{28} has one of the following structures:







In some embodiments, $\mathbf{R}^{28}$ has one of the following structures:







In certain embodiments, R²⁸ has one of the following structures:









5 In some embodiments, the compounds of Structure (I), (II) or (III), and embodiments thereof, can be in the form of a salt such as a pharmaceutically acceptable salt.

The compounds of Structure (I), (II) or (III), and embodiments thereof, are useful as inhibitors of MASP-2 and for therapeutic use. The compounds of Structure (I), (II) or (III), and embodiments thereof, are useful in the treatment of MASP-2-associated diseases and disorders, and in the manufacture of medicaments for treating MASP-2-associated diseases and disorders. The present disclosure also provides methods of treating a MASP-2-associated disease and disorder comprising administering to a patient a therapeutically effective amount of a compound of Structure (I), (II) or (III), or an embodiment thereof, optionally in the form of a salt.

In some embodiments the compound Structure (I), (II) or (III) or an embodiment thereof is provided in the form of a pharmaceutical composition comprising the compound or a salt thereof, such as a pharmaceutically acceptable salt, and at least one pharmaceutically acceptable carrier or excipient.

20 In certain aspects, the compound is one or more selected from the compounds of Structure (I), (II) or (III) set forth in the Examples, including the compounds listed in Table 1, (e.g., compounds with selectivity for MASP-2 over thrombin). In certain aspects, one or more of the variables defining the compounds of Structure (I), (II) or (III) is selected from the corresponding substituents in the compounds of Structure (I), (II) or (III) in the Examples including the compounds listed in Table 1, preferably, those of the compounds with selectivity for MASP-2 over thrombin.

In certain aspects, the disclosure sets forth a stereochemically pure enantiomer or diastereomer (*e.g.*, an optically active compound with one or more stereocenter(s)). Unless specifically indicated otherwise, for any compound with one or more stereocenters is intended to include and to describe both the pure (+) and (-) enantiomers, any other diastereomers, mixtures that are enriched in an enantiomer or diastereomer (*e.g.*, 10%, 20%, 30%, 40%, 50%, 60%, 70% 75%, 80%, 85, 90%, or 95% enantiomeric or diastereomeric excess), and a racemic mixture of enantiomers or diastereomers.

Certain embodiments provide a pharmaceutically acceptable salt of the indicated chemical structure (*e.g.*, a hydrohalide, such as a hydrochloride or dihydrochloride). Examples of pharmaceutically acceptable salts are set forth in, *e.g.*, Burge, S. M. et al., J. Pharm. Sci 1977, 66, 1-19. They include chlorides, bromides, iodides, formates, acetates, propionates, oxalates, malonates, succinates, fumarates, maleates, tartrates, citrates, benzoates, phthalates, sulfonates, arylsulfonates, alkylsulfonates, salts of fatty acids, and the like. Salts can be prepared by a variety of methods known to the skilled artisan, including a precipitation with the conjugate acid or base (*e.g.*, treatment with gaseous HCl or an HCl solution).

In certain embodiments provide a prodrug. A prodrug is a compound that is converted to a biologically active form under physiological conditions, often by hydrolysis, oxidation, or reduction (*e.g.*, ester to acid form; carbamate to amino or hydroxy group; hydroxyamidine to amidine) Exemplary prodrugs are set forth in, *e.g.*, Tilley, J.W., "Prodrugs of Benzamide," *Prodrugs* 2007, 191-222; Peterlin-Masic et al. *Curr. Pharma. Design* 2006, 12, 73-91. Prodrugs for the amidine group include amidoximes, O-alkylamidoximes, acylamidines, carbamates, 1,2,4-oxadiazolin-4-ones, and the like.

In certain aspects, the compound is useful for selectively inhibiting MASP-2 over thrombin, the method comprising administering the compound as described herein. In certain aspects, the selectivity ratio of MASP-2:thrombin is at least 1.1:1, 1.25:1, 1.5:1, 1.75:1, 2:1, 3:1, 4:1, 5:1, 6:1, 7:1, 8:1, 9:1, 10:1, 11:1, 12:1, 13:1, 14:1, 15:1, 16:1, 17:1, 18:1, 19:1, 20:1, 21:1, 22:1, 23:1, 24:1, 25:1, or 30:1.

III. Synthesis

Compounds described herein, including salts thereof, can be prepared using known organic synthesis techniques and can be synthesized according to any of numerous possible synthetic routes, such as those illustrated in the Examples below.

5 The reactions for preparing compounds described herein can be carried out in suitable solvents which can be readily selected by one of skill in the art of organic synthesis. Suitable solvents can be substantially non-reactive with the starting materials (reactants), the intermediates or products at the temperatures at which the reactions are carried out, *e.g.*, temperatures which can range from the solvent's freezing temperature to
10 the solvent's boiling temperature. A given reaction can be carried out in one solvent or a mixture of more than one solvent. Depending on the particular reaction step, suitable solvents for a particular reaction step can be selected by the skilled artisan.

Preparation of compounds of the disclosure can involve the protection and deprotection of various chemical groups. The need for protection and deprotection, and
15 the selection of appropriate protecting groups, can be readily determined by one skilled in the art. The chemistry of protecting groups is described, *e.g.*, in Kocienski, *Protecting Groups*, (Thieme, 2007); Robertson, *Protecting Group Chemistry*, (Oxford University Press, 2000); Smith *et al.*, *March's Advanced Organic Chemistry: Reactions, Mechanisms, and Structure*, 6th Ed. (Wiley, 2007); Petursson *et al.*, "Protecting Groups
20 in Carbohydrate Chemistry," *J. Chem. Educ.*, 1997, 74(11), 1297; and Wuts *et al.*, *Protective Groups in Organic Synthesis*, 4th Ed., (Wiley, 2006).

Reactions can be monitored according to any suitable method known in the art. For example, product formation can be monitored by spectroscopic means, such as nuclear magnetic resonance spectroscopy (*e.g.*, ^1H or ^{13}C), infrared spectroscopy,
25 spectrophotometry (*e.g.*, UV-visible), mass spectrometry, or by chromatographic methods such as high-performance liquid chromatography (HPLC) or thin layer chromatography (TLC).

The particular synthetic methods used in the Examples provide general guidance in connection with preparing the compounds of the disclosure. One skilled in
30 the art would understand that the preparations can be modified or optimized using general

knowledge of organic chemistry to prepare various compounds within the scope of the present disclosure.

Starting materials, reagents and intermediates whose synthesis is not described herein are either commercially available, known in the literature, or may be
5 prepared by methods known to one skilled in the art.

It will be appreciated by one skilled in the art that the processes described are not the exclusive means by which compounds of the disclosure may be synthesized and that a broad repertoire of synthetic organic reactions is available to be potentially employed in synthesizing compounds of the disclosure. The person skilled in the art
10 knows how to select and implement appropriate synthetic routes. Suitable synthetic methods of starting materials, intermediates and products may be identified by reference to the literature, including reference sources such as: *Advances in Heterocyclic Chemistry*, Vols. 1-107 (Elsevier, 1963-2012); *Journal of Heterocyclic Chemistry*, Vols. 1-49 (Journal of Heterocyclic Chemistry, 1964-2012); Carreira, *et al.* (Ed.) *Science of*
15 *Synthesis*, Vols. 1-48 (2001-2010) and Knowledge Updates KU2010/1-4; 2011/1-4; 2012/1-2 (Thieme, 2001-2012); Katritzky, *et al.* (Ed.) *Comprehensive Organic Functional Group Transformations*, (Pergamon Press, 1996); Katritzky *et al.* (Ed.); *Comprehensive Organic Functional Group Transformations II* (Elsevier, 2nd Edition, 2004); Katritzky *et al.* (Ed.), *Comprehensive Heterocyclic Chemistry* (Pergamon Press,
20 1984); Katritzky *et al.*, *Comprehensive Heterocyclic Chemistry II* (Pergamon Press, 1996); Smith *et al.*, *March's Advanced Organic Chemistry: Reactions, Mechanisms, and Structure*, 6th Ed. (Wiley, 2007); Trost *et al.* (Ed.), *Comprehensive Organic Synthesis* (Pergamon Press, 1991).

IV. Methods of Treatment

25 In another aspect, the present disclosure provides a method of treating a patient suffering from, or at risk for developing a MASP-2-associated disease or disorder such as a MASP-2-dependent complement-associated disease or disorder comprising administering a small molecule inhibitor of MASP-2.

The compound can be any small molecule inhibitor of MASP-2. In some
30 embodiments, the compound can be a small molecule inhibitor of MASP-2 that binds to

the serine protease domain of MASP-2. In some embodiments, the compound can be a small molecule inhibitor such as a synthetic small molecule inhibitor of MASP-2. In some embodiments, the compound can be a small molecule inhibitor of MASP-2 that binds to the catalytic, substrate-binding region of MASP-2. In some embodiments, the compound
5 selectively inhibits MASP-2 relative to thrombin. For example, in some embodiments, the compound is a compound of Structure (I), (II) or (III) as described in any of the foregoing embodiments.

As described in U.S. Patent No. 7,919,094; U.S. Patent No. 8,840,893; U.S. Patent No. 8,652,477; U.S. Patent No. 8,951,522, U.S. Patent No. 9,011,860, U.S.
10 Patent No. 9,475,885, U.S. Patent No. 9,644,035, U.S. Patent Application Publication Nos. US 2013/0344073, US 2013/0266560, US 2015/0166675, US 2017/0137537, US 2017/0166660, US 2017/0189525, US 2017/0267781, US 2017/0283508, US 2017/0253667, US 2018/0105604, and PCT Publication Nos. WO 2018/045054, WO 2019/036460 and co-pending U.S. Patent Application Serial No. 62/688,611 (each of
15 which is assigned to Omeros Corporation, the assignee of the instant application, each of which is hereby incorporated by reference), MASP-2-dependent complement activation has been implicated as contributing to the pathogenesis of numerous acute and chronic disease states. For example, as described in U.S. Patent No. 8,951,522, the primary function of the complement system, a part of the innate immune system, is to protect the
20 host against infectious agents, however, inappropriate or over-activation of the complement system can lead to serious disease, such as thrombotic microangiopathies (TMAs, including aHUS, TTP and HUS) in which endothelial damage as well as fibrin and platelet-rich thrombi in the microvasculature lead to organ damage. The lectin pathway plays a dominant role in activating complement in settings of endothelial cell
25 stress or injury and preventing the activation of MASP-2 and the lectin pathway halts the sequence of enzymatic reactions that lead to the formation of the membrane attack complex, platelet activation and leukocyte recruitment. As described in U.S. Patent No. 8,652,477, in addition to initiation of the lectin pathway, MASP-2 can also activate the coagulation system and is capable of cleaving prothrombin to thrombin.

Accordingly, in some embodiments, the method comprises administering to a patient suffering from or at risk for developing a MASP-2-dependent complement-associated disease or disorder an amount of a compound of the disclosure in an amount sufficient to inhibit MASP-2 dependent complement activation in the mammalian subject
5 to thereby treat the disease or disorder. In some embodiments, the method can further comprise, prior to administering a compound of the disclosure to the patient, determining that the patient is afflicted with the lectin complement-associated disease or disorder.

In some embodiments, the MASP-2-dependent complement-associated disease or disorder is selected from the group consisting of a thrombotic microangiopathy
10 (TMA), a renal condition, an inflammatory reaction resulting from tissue or organ transplantation, an ischemia reperfusion injury, a complication associated with diabetes, a cardiovascular disease or disorder, an inflammatory gastrointestinal disorder, a pulmonary disorder, an ophthalmic disease or disorder, disseminated intravascular coagulation, graft-versus-host disease, veno-occlusive disease, diffuse alveolar
15 hemorrhage, or similar, as well as combinations thereof.

In some embodiments, the MASP-2-dependent complement-associated disease or disorder is a thrombotic microangiopathy (TMA) including thrombotic thrombocytopenic purpura (TTP), refractory TTP, Upshaw-Schulman Syndrome (USS), hemolytic uremic syndrome (HUS), atypical hemolytic syndrome (aHUS), non-Factor H-
20 dependent atypical hemolytic syndrome, aHUS secondary to an infection, plasma therapy-resistant aHUS, a TMA secondary to cancer, a TMA secondary to chemotherapy, a TMA secondary to transplantation, or a TMA associated with hematopoietic stem cell transplant.

In some embodiments, the method comprises administering to a patient
25 suffering from or at risk for developing graft-versus-host disease (GVHD), including acute GVHD, chronic GVHD or steroid-resistant GVHD an amount of a compound of the disclosure in an amount sufficient to inhibit MASP-2 dependent complement activation in the mammalian subject to thereby treat the disease or disorder. In some embodiments, the subject suffering from or at risk for developing GVHD has previously
30 undergone, is undergoing, or will undergo a hematopoietic stem cell transplant.

In some embodiments, the method comprises administering to a patient suffering from, or at risk for developing diffuse alveolar hemorrhage (DAH) an amount of a compound of the disclosure in an amount sufficient to inhibit MASP-2 dependent complement activation in the mammalian subject to thereby treat the disease or disorder.

5 In some embodiments, the subject suffering from, or at risk for developing DAH has previously undergone, is undergoing, or will undergo a hematopoietic stem cell transplant.

In some embodiments, the method comprises administering to a patient suffering from, or at risk for developing veno-occlusive disease (VOD) an amount of a
10 compound of the disclosure in an amount sufficient to inhibit MASP-2 dependent complement activation in the mammalian subject to thereby treat the disease or disorder. In some embodiments, the subject suffering from, or at risk for developing VOD has previously undergone, is undergoing, or will undergo a hematopoietic stem cell transplant.

15 In some embodiments, the method comprises administering to a patient suffering from, or at risk for developing idiopathic pneumonia syndrome (IPS) an amount of a compound of the disclosure in an amount sufficient to inhibit MASP-2 dependent complement activation in the mammalian subject to thereby treat the disease or disorder. In some embodiments, the subject suffering from, or at risk for developing IPS has
20 previously undergone, is undergoing, or will undergo a hematopoietic stem cell transplant.

In some embodiments, the method comprises administering to a patient suffering from, or at risk for developing capillary leak syndrome (CLS) an amount of a compound of the disclosure in an amount sufficient to inhibit MASP-2 dependent
25 complement activation in the mammalian subject to thereby treat the disease or disorder. In some embodiments, the subject suffering from, or at risk for developing CLS has previously undergone, is undergoing, or will undergo a hematopoietic stem cell transplant.

In some embodiments, the method comprises administering to a patient
30 suffering from, or at risk for developing engraftment syndrome (ES) an amount of a

compound of the disclosure in an amount sufficient to inhibit MASP-2 dependent complement activation in the mammalian subject to thereby treat the disease or disorder. In some embodiments, the subject suffering from, or at risk for developing ES has previously undergone, is undergoing, or will undergo a hematopoietic stem cell
5 transplant.

In some embodiments, the method comprises administering to a patient suffering from, or at risk for developing fluid overload (FO) an amount of a compound of the disclosure in an amount sufficient to inhibit MASP-2 dependent complement activation in the mammalian subject to thereby treat the disease or disorder. In some
10 embodiments, the subject suffering from, or at risk for developing FO has previously undergone, is undergoing, or will undergo a hematopoietic stem cell transplant.

In some embodiments, the method comprises administering to a patient suffering from any of the above-referenced diseases or conditions an amount of a compound as disclosed in PCT Application No. PCT/US19/34225, which is hereby
15 incorporated in its entirety.

In some embodiments, the MASP-2-dependent complement-associated disease or disorder is a renal condition including mesangioproliferative glomerulonephritis, membranous glomerulonephritis, membranoproliferative glomerulonephritis (mesangiocapillary glomerulonephritis), acute post infectious
20 glomerulonephritis (poststreptococcal glomerulonephritis), C3 glomerulopathy, cryoglobulinemic glomerulonephritis, pauci-immune necrotizing crescentic glomerulonephritis, lupus nephritis, Henoch-Schonlein purpura nephritis, IgA nephropathy, and the like, as well as combinations thereof.

In some embodiments, the MASP-2-dependent complement-associated
25 disease or disorder is renal fibrosis (*e.g.*, tubulointerstitial fibrosis) and/or proteinuria in a subject suffering from or at risk for developing chronic kidney disease, chronic renal failure, glomerular disease (*e.g.*, focal segmental glomerulosclerosis), an immune complex disorder (*e.g.*, IgA nephropathy, membranous nephropathy), lupus nephritis, nephrotic syndrome, diabetic nephropathy, tubulointerstitial damage and
30 glomerulonephritis (*e.g.*, C3 glomerulopathy), or a disease or condition associated with

proteinuria, including, but not limited to, nephrotic syndrome, pre-eclampsia, eclampsia, toxic lesions of kidneys, amyloidosis, collagen vascular diseases (*e.g.*, systemic lupus erythematosus), dehydration, glomerular diseases (*e.g.*, membranous glomerulonephritis, focal segmental glomerulonephritis, C3 glomerulopathy, minimal change disease, lipoid nephrosis), strenuous exercise, stress, benign orthostatis (postural) proteinuria, focal segmental glomerulosclerosis, IgA nephropathy (*i.e.*, Berger's disease), IgM nephropathy, membranoproliferative glomerulonephritis, membranous nephropathy, minimal change disease, sarcoidosis, Alport's syndrome, diabetes mellitus (diabetic nephropathy), drug-induced toxicity (*e.g.*, NSAIDs, nicotine, penicillamine, lithium carbonate, gold and other heavy metals, ACE inhibitors, antibiotics (*e.g.*, adriamycin), opiates (*e.g.*, heroin), or other nephrotoxins); Fabry's disease, infections (*e.g.*, HIV, syphilis, hepatitis A, B or C, poststreptococcal infection, urinary schistosomiasis); aminoaciduria, Fanconi syndrome, hypertensive nephrosclerosis, interstitial nephritis, sickle cell disease, hemoglobinuria, multiple myeloma, myoglobinuria, organ rejection (*e.g.*, kidney transplant rejection), ebola hemorrhagic fever, Nail patella syndrome, familial Mediterranean fever, HELLP syndrome, systemic lupus erythematosus, Wegener's granulomatosis, Rheumatoid arthritis, Glycogen storage disease type 1, Goodpasture's syndrome, Henoch-Schonlein purpura, urinary tract infection which has spread to the kidneys, Sjogren's syndrome and post-infections glomerulonephritis.

20 In some embodiments, the MASP-2-dependent complement-associated disease or disorder is an inflammatory reaction resulting from tissue or solid organ transplantation, including allotransplantation or xenotransplantation of whole organs (*e.g.*, kidney, heart, liver, pancreas, lung, cornea, and the like) or tissue grafts (*e.g.*, valves, tendons, bone marrow, and the like).

25 In some embodiments, the MASP-2-dependent complement-associated disorder is an ischemia reperfusion injury (I/R), including myocardial I/R, gastrointestinal I/R, renal I/R, and I/R following an aortic aneurism repair, I/R associated with cardiopulmonary bypass, cerebral I/R, stroke, organ transplant or reattachment of severed or traumatized limbs or digits; revascularization to transplants and/or replants,

and hemodynamic resuscitation following shock, surgical procedures, or similar, as well as combinations thereof.

In some embodiments, the MASP-2-dependent complement-associated disease or disorder is a complication associated with non-obese diabetes (Type-1 diabetes
5 or Insulin-dependent diabetes mellitus) and/or complications associated with Type-1 or Type-2 (adult onset) diabetes including diabetic angiopathy, diabetic neuropathy, diabetic retinopathy, diabetic macular edema, and the like, as well as combinations thereof.

In some embodiments, the MASP-2-dependent complement-associated
10 disease or disorder is a cardiovascular disease or disorder, including Henoch-Schonlein purpura nephritis, systemic lupus erythematosus-associated vasculitis, vasculitis associated with rheumatoid arthritis (also called malignant rheumatoid arthritis), immune complex vasculitis, and Takayasu's disease; dilated cardiomyopathy; diabetic angiopathy; Kawasaki's disease (arteritis); venous gas embolus (VGE); and inhibition of
15 restenosis following stent placement, rotational atherectomy, percutaneous transluminal coronary angioplasty (PTCA), and the like as well as combinations thereof.

In some embodiments, the MASP-2-dependent complement-associated disease or disorder is an inflammatory gastrointestinal disorder, including pancreatitis, diverticulitis and bowel disorders including Crohn's disease, ulcerative colitis, irritable
20 bowel syndrome, inflammatory bowel disease (IBD), or similar, as well as combinations thereof.

In some embodiments, the MASP-2-dependent complement-associated disease or disorder is a pulmonary disorder, including acute respiratory distress syndrome, transfusion-related acute lung injury, ischemia/reperfusion acute lung injury,
25 chronic obstructive pulmonary disease, asthma, Wegener's granulomatosis, antiglomerular basement membrane disease (Goodpasture's disease), meconium aspiration syndrome, aspiration pneumonia, bronchiolitis obliterans syndrome, idiopathic pulmonary fibrosis, acute lung injury secondary to burn, non-cardiogenic pulmonary edema, transfusion-related respiratory depression, emphysema, and the like, as well as
30 combinations thereof.

In some embodiments, the MASP-2-dependent complement-associated disease or disorder is an extracorporeal exposure-triggered inflammatory reaction and the method comprises treating a subject undergoing an extracorporeal circulation procedure. In some embodiments, the extracorporeal circulation procedure includes hemodialysis, 5 plasmapheresis, leukopheresis, extracorporeal membrane oxygenation (ECMO), heparin-induced extracorporeal membrane oxygenation LDL precipitation (HELP), cardiopulmonary bypass (CPB), and the like.

In some embodiments, the MASP-2-dependent complement-associated disease or disorder is selected from inflammatory or non-inflammatory arthritides and 10 other musculoskeletal disorders, *e.g.*, osteoarthritis, rheumatoid arthritis, juvenile rheumatoid arthritis, gout, neuropathic arthropathy, psoriatic arthritis, ankylosing spondylitis or other spondyloarthropathies and crystalline arthropathies, muscular dystrophy, systemic lupus erythematosus (SLE), or similar, as well as combinations thereof.

15 In some embodiments, the MASP-2-dependent complement-associated disease or disorder is a skin disorder; for example, psoriasis, autoimmune bullous dermatoses, eosinophilic spongiosis, bullous pemphigoid, epidermolysis bullosa acquisita, atopic dermatitis, herpes gestationis, and other skin disorders. In some 20 embodiments, the MASP-2-dependent complement-associated disease or disorder is a thermal burn, chemical burn, or combinations thereof, including capillary leakage caused thereby.

In some embodiments, the MASP-2-dependent complement-associated disease or disorder is a peripheral nervous system (PNS) and/or central nervous system (CNS) disorder or injury including multiple sclerosis (MS), myasthenia gravis (MG), 25 Huntington's disease (HD), amyotrophic lateral sclerosis (ALS), Guillain Barre syndrome, reperfusion following stroke, degenerative discs, cerebral trauma, Parkinson's disease (PD), Alzheimer's disease (AD), Miller-Fisher syndrome, cerebral trauma and/or hemorrhage, traumatic brain injury, demyelination, meningitis, or similar, as well as combinations thereof.

In some embodiments, the MASP-2-dependent complement-associated disease or disorder is sepsis or a condition resulting from sepsis including severe sepsis, septic shock, acute respiratory distress syndrome resulting from sepsis, hemolytic anemia, systemic inflammatory response syndrome, hemorrhagic shock, or the like, as well as combinations thereof.

In some embodiments, the MASP-2-dependent complement-associated disease or disorder is a urogenital disorder including painful bladder disease, sensory bladder disease, chronic abacterial cystitis and interstitial cystitis, male and female infertility, placental dysfunction and miscarriage, pre-eclampsia, or similar, as well as combinations thereof.

In some embodiments, the MASP-2-dependent complement-associated disease or disorder is an inflammatory reaction in a subject being treated with chemotherapeutics and/or radiation therapy, including for the treatment of cancerous conditions.

In some embodiments, the MASP-2-dependent complement-associated disease or disorder is an angiogenesis-dependent cancer, including a solid tumor(s), blood borne tumor(s), high-risk carcinoid tumors, tumor metastases, and the like, including combinations thereof.

In some embodiments, the MASP-2-dependent complement-associated disease or disorder is an angiogenesis-dependent benign tumor, including hemangiomas, acoustic neuromas, neurofibromas, trachomas, carcinoid tumors, pyogenic granulomas, or similar, as well as combinations thereof.

In some embodiments, the MASP-2-dependent complement-associated disease or disorder is an endocrine disorder including Hashimoto's thyroiditis, stress, anxiety, other potential hormonal disorders involving regulated release of prolactin, growth or insulin-like growth factor, adrenocorticotropin from the pituitary, or similar, as well as combinations thereof.

In some embodiments, the MASP-2-dependent complement-associated disease or disorder is an ophthalmic disease or disorder including age-related macular degeneration, glaucoma, endophthalmitis, and the like, as well as combinations thereof.

In some embodiments, the MASP-2-dependent complement-associated disease or disorder is an ocular angiogenic disease or condition including age-related macular degeneration, uveitis, ocular melanoma, corneal neovascularization, primary pterygium, HSV stromal keratitis, HSV-1-induced corneal lymphangiogenesis, proliferative diabetic retinopathy, diabetic macular edema, retinopathy of prematurity, retinal vein occlusion, corneal graft rejection, neovascular glaucoma, vitreous hemorrhage secondary to proliferative diabetic retinopathy, neuromyelitis optica, rubeosis, or similar, as well as combinations thereof.

In some embodiments, the MASP-2-dependent complement-associated disease or disorder is disseminated intravascular coagulation (DIC) or other complement mediated coagulation disorder, including DIC secondary to sepsis, severe trauma, including neurological trauma (*e.g.*, acute head injury; see Kumura et al, *Acta Neurochirurgica* 55:23-28 (1987), infection (*e.g.*, bacterial, viral, fungal, parasitic), cancer, obstetrical complications, liver disease, severe toxic reaction (*e.g.*, snake bite, insect bite, transfusion reaction), shock, heat stroke, transplant rejection, vascular aneurysm, hepatic failure, cancer treatment by chemotherapy or radiation therapy, burn, or accidental radiation exposure.

In some embodiments, the MASP-2-dependent complement-associated disease or disorder is selected from the group consisting of acute radiation syndrome, dense deposit disease, Degos Disease, Catastrophic Antiphospholipid Syndrome (CAPS), Behcet's disease, cryoglobulinemia, paroxysmal nocturnal hemoglobinuria (PNH), cold agglutinin disease, and combinations thereof.

In some embodiments, the MASP-2-dependent complement-associated disease or disorder is selected from the group consisting of aHUS, HSCT-TMA, IgAN, Lupus Nephritis (LN), and combinations thereof.

In some embodiments, the method comprises administering to a patient suffering from, or at risk for developing a disease, disorder or condition associated with fibrin-induced activation of the complement system and the associated activation of the coagulation and/or contact systems an amount of a compound according to any one of the foregoing embodiments (*e.g.*, a compound of Structure (I), (II) or (III)) in an amount

sufficient to inhibit MASP-2 dependent complement activation in the mammalian subject to thereby treat the disease or disorder. In some embodiments, the subject is suffering from, or at risk of developing, a disease, disorder or condition associated with complement-related inflammation, excessive coagulation or contact system activation
5 initiated by fibrin or activated platelets. In some embodiments, the subject is suffering from a disease or disorder selected from the group consisting of arterial thrombosis, venous thrombosis, deep vein thrombosis, post-surgical thrombosis, restenosis following coronary artery bypass graft and/or an interventional cardiovascular procedure (*e.g.*, angioplasty or stent placement), atherosclerosis, plaque rupture, plaque instability,
10 restenosis, hypotension, acute respiratory distress syndrome (ARDS), systemic inflammatory response syndrome (SIRS), disseminated intravascular coagulation (DIC), veno-occlusive disease (VOD), thrombotic microangiopathy, lupus nephritis, superficial thrombophlebitis, Factor V Leiden mutation, ischemic/reperfusion injury, human immunodeficiency virus (HIV) infection, undergoing hormone-replacement therapy
15 (HRT), Alzheimer's disease and/or suffering from a hypercoagulable state.

In some embodiments, the subject is suffering from, or at risk for developing an acquired hypercoagulable state due to at least one or more of the following: undergoing therapy with a drug selected from the group consisting of 5-FU, GM-CSF, cisplatin, heparin, COX-2 inhibitor, contrast media, corticosteroids and antipsychotics;
20 venous stasis (immobilization, surgery, *etc.*), antiphospholipid syndrome, cancer (promyelocytic leukemia, lung, breast, prostate, pancreas, stomach and colon tumors), tissue injury due to trauma or surgery, presence of a catheter in a central vein, acquired deficiency of a protein involved in clot formation (*e.g.*, protein C), paroxysmal nocturnal hemoglobinuria (PNH), elevated levels of homocysteine, heart failure, presence of a
25 mechanical valve, pulmonary hypertension with *in situ* thrombosis, atrial fibrillation, heparin-induced thrombocytopenia (HIT), heparin-induced thrombocytopenia and thrombosis (HITT), Kawasaki disease with *in situ* thrombus, Takayasu arteritis with *in situ* thrombus, thrombophilia of metastatic cancer, elevated Factor VIII levels, pregnancy, inflammatory bowel disease (IBD), or due to a genetic defect that causes or
30 increases the risk of developing, a hypercoagulable state, such as a genetic defect selected

from the group consisting of a Prothrombin 20210 gene mutation, an MTHFR mutation, a deficiency of protein C, a deficiency of protein S, a deficiency of protein A, a deficiency of protein Z, an antithrombin deficiency, and a genetic disorder producing thrombophilia.

In some embodiments, the subject is suffering from, or at risk for
5 developing, a disease or disorder that is amenable to treatment with a kallikrein inhibitor. In some embodiments, the subject is suffering from, or at risk for developing a disease or disorder amenable to treatment with a kallikrein inhibitor is selected from the group consisting of hereditary angioedema, diabetic macular edema and bleeding during cardiopulmonary bypass. In some embodiments, the subject is suffering from, or at risk
10 for developing, a disease or disorder that is amenable to treatment with a thrombin inhibitor, such as arterial thrombosis, venous thrombosis, pulmonary embolism, atrial fibrillation, heparin-induced thrombocytopenia, conversion from one anticoagulant to another, or off-label use for extracorporeal circuit patency of continuous renal replacement therapy (CRRT) in critically ill patients with HIT (maintenance).

15 In some embodiments, the subject has previously experienced, is currently suffering from, or is at risk for developing atrial fibrillation and the MASP-2 inhibitory compound (*e.g.*, a compound of Structure (I), (II) or (III)) is administered in an amount sufficient to reduce the risk of stroke in said subject. In some embodiments, the subject is suffering from, or at risk for developing, a disease or disorder that is amenable to
20 treatment with a factor XII inhibitor, such as deep vein thrombosis (both primary prophylaxis and extended therapy), pulmonary embolism, nonvalvular atrial fibrillation, prevention of recurrent ischemia after acute coronary syndrome in subjects with or without atrial fibrillation, end-stage renal disease, cerebral ischemia, angina, or to reduce or prevent clotting associated with medical devices (*e.g.*, valves, small caliber grafts, *etc.*)
25 and/or extracorporeal circuits.

In some embodiments, the subject has previously experienced, is currently suffering from, or is at risk for developing nonvalvular atrial fibrillation and the MASP-2 inhibitory compound (*e.g.*, a compound of Structure (I), (II) or (III)) is administered in an amount sufficient to reduce the risk of stroke and/or embolism in said subject. In some
30 embodiments, the subject has an acquired disease or disorder that increases the propensity

for thromboembolism, such as a disease or disorder selected from the group consisting of atherosclerosis, antiphospholipid antibodies, cancer (*e.g.*, promyelocytic leukemia, lung, breast, prostate, pancreatic, stomach and colon), hyperhomocysteinemia, infection, tissue injury, venous stasis (such as due to surgery, orthopedic or paralytic immobilization, heart failure, pregnancy, or obesity) and a subject taking oral contraceptives that contain estrogen.

In some embodiments, the subject is in need of anticoagulant therapy and the MASP-2 inhibitory compound (*e.g.*, a compound of Structure (I), (II) or (III)) is used as a replacement for standard anticoagulant therapy (*e.g.*, Warfarin). In some embodiments, the subject has a condition that normally prohibits standard anticoagulant therapy, such as CNS amyloid angiopathy. In some embodiments of the method, the MASP-2 inhibitory compound is administered as a bridging agent perioperatively in a subject otherwise on standard anticoagulation therapy. In some embodiments, the subject has sickle cell disease which is a vaso-occlusive disorder involving activation of platelets.

Atypical hemolytic uremic syndrome (aHUS) is part of a group of conditions termed "Thrombotic microangiopathies." In the atypical form of HUS (aHUS), the disease is associated with defective complement regulation and can be either sporadic or familial. Familial cases of aHUS are associated with mutations in genes coding for complement activation or complement regulatory proteins, including complement factor H, factor I, factor B, membrane cofactor CD46 as well as complement factor H-related protein 1 (CFHR1) and complement factor H-related protein 3 (CFHR3). (Zipfel, P.F., et al., PloS Genetics 3(3):e41 (2007)). The unifying feature of this diverse array of genetic mutations associated with aHUS is a predisposition to enhanced complement activation on cellular or tissue surfaces. A subject is at risk for developing aHUS upon the onset of at least one or more symptoms indicative of aHUS (*e.g.*, the presence of anemia, thrombocytopenia and/or renal insufficiency) and/or the presence of thrombotic microangiopathy in a biopsy obtained from the subject. The determination of whether a subject is at risk for developing aHUS comprises determining whether the subject has a genetic predisposition to developing aHUS, which may be carried out by assessing genetic information (*e.g.* from a database containing the genotype of the

subject), or performing at least one genetic screening test on the subject to determine the presence or absence of a genetic marker associated with aHUS (*i.e.*, determining the presence or absence of a genetic mutation associated with aHUS in the genes encoding complement factor H (CFH), factor I (CFI), factor B (CFB), membrane cofactor CD46, C3, complement factor H-related protein 1 (CFHR1), or THBD (encoding the anticoagulant protein thrombomodulin) or complement factor H-related protein 3 (CFHR3), or complement factor H-related protein 4 (CFHR4)) either via genome sequencing or gene-specific analysis (*e.g.*, PCR analysis), and/or determining whether the subject has a family history of aHUS. Methods of genetic screening for the presence or absence of a genetic mutation associated with aHUS are well established, for example, see Noris M et al. "Atypical Hemolytic-Uremic Syndrome," 2007 Nov 16 [Updated 2011 Mar 10]. In: Pagon RA, Bird TD, Dolan CR, et al., editors. GeneReviews™, Seattle (WA): University of Washington, Seattle.

Hematopoietic stem cell transplant-associated TMA (HSCT-TMA) is a life-threatening complication that is triggered by endothelial injury. The kidney is the most commonly affected organ, though HSCT-TMA can be a multi-system disease that also involves the lung, bowel, heart, and brain. The occurrence of even mild TMA is associated with long-term renal impairment. Development of post-allogeneic HSCT-associated TMA differs in frequency based on varying diagnostic criteria and conditioning and graft-versus-host disease prophylaxis regimens, with calcineurin inhibitors being the most frequent drugs implicated (Ho VT et al., Biol Blood Marrow Transplant, 11(8):571-5, 2005).

Immunoglobulin A nephropathy (IgAN) is an autoimmune kidney disease resulting in intrarenal inflammation and kidney injury. IgAN is the most common primary glomerular disease globally. With an annual incidence of approximately 2.5 per 100,000, it is estimated that 1 in 1400 persons in the U.S. will develop IgAN. As many as 40% of patients with IgAN will develop end-stage renal disease (ESRD). Patients typically present with microscopic hematuria with mild to moderate proteinuria and variable levels of renal insufficiency (Wyatt R.J., et al., NEnglJ Med 36S(25):2402-4, 2013). Clinical markers such as impaired kidney function, sustained hypertension, and heavy proteinuria

(over 1 g per day) are associated with poor prognosis (Goto M et al., Nephrol Dial Transplant 24(10):3068-74, 2009; Berthoux F. et al., J Am Soc Nephrol 22(4):752-61, 2011). Proteinuria is the strongest prognostic factor independent of other risk factors in multiple large observational studies and prospective trials (Coppo R. et al., J Nephrol 5 18(5):503-12, 2005; Reich H. N., et al., J Am Soc Nephrol 18(12):3177-83, 2007). It is estimated that 15-20% of patients reach ESRD within 10 years of disease onset if left untreated (D'Amico G., Am J Kidney Dis 36(2):227-37, 2000). The diagnostic hallmark of IgAN is the predominance of IgA deposits, alone or with IgG, IgM, or both, in the glomerular mesangium.

10 A main complication of systemic lupus erythematosus (SLE) is nephritis, also known as lupus nephritis, which is classified as a secondary form of glomerulonephritis. Up to 60% of adults with SLE have some form of kidney involvement later in the course of the disease (Koda-Kimble et al., Koda-Kimble and Young's Applied Therapeutics: the clinical use of drugs, 10th Ed, Lippincott Williams & 15 Wilkins: pages 792-9, 2012) with a prevalence of 20-70 per 100,000 people in the U.S. Lupus nephritis often presents in patients with other symptoms of active SLE, including fatigue, fever, rash, arthritis, serositis, or central nervous system disease (Pisetsky D.S. et al., Med Clin North Am 81(1): 113-28, 1997). Some patients have asymptomatic lupus nephritis; however, during regular follow-up, laboratory abnormalities such as elevated 20 serum creatinine levels, low albumin levels, or urinary protein or sediment suggest active lupus nephritis.

V. Compositions, Dosage, and Administration

 The compounds as described herein (*e.g.*, a compound of Structure (I), (II) or (III)) can be administered in a manner compatible with the dosage formulation, and in 25 such amount as will be effective or suitable for treatment. The quantity to be administered depends on a variety of factors including, *e.g.*, the age, body weight, physical activity, and diet of the individual, and the desired effect. In certain embodiments, the size of the dose may also be determined by the existence, nature, and extent of any adverse side effects that accompany the administration of the compound in a particular individual.

It will be understood, however, that the specific dose level and frequency of dosage for any particular patient may be varied by a physician and will depend upon a variety of factors including the activity of the specific compound employed, the metabolic stability and length of action of that compound, the age, body weight, hereditary
5 characteristics, general health, sex, diet, mode and time of administration, rate of excretion, drug combination, the severity of the particular condition, and the host undergoing therapy.

In certain embodiments, the dose may take the form of solid, semi-solid, or liquid forms, preferably in unit dosage forms suitable for simple administration of
10 precise dosages.

As used herein, the term "unit dosage form" refers to physically discrete units suitable as unitary dosages for humans and other mammals, each unit containing a predetermined quantity of an active agent calculated to produce the desired onset, tolerability, and/or efficacious effects, in association with a suitable pharmaceutical
15 excipient (*e.g.*, an ampoule). In addition, more concentrated dosage forms may be prepared, from which the more dilute unit dosage forms may then be produced.

The compounds described herein (*e.g.*, a compound of Structure (I), (II) or (III)) can be administered to a subject in need of treatment using methods known in the art, such as by oral administration or by injection. The injection can be, *e.g.*,
20 subcutaneous, intravenous, intraperitoneal, or intramuscular. As described herein, parenteral formulations can be prepared in dosage unit form for ease of administration and uniformity of dosage. As used herein the term "unit dosage form" refers to physically discrete units suited as unitary dosages for the subject to be treated; each unit containing a predetermined quantity of active compound calculated to produce the desired
25 therapeutic effect.

The pharmaceutical compositions of the present application comprise a therapeutically effective amount of a compound of the present disclosure ((*e.g.*, a compound of Structure (I), (II) or (III)) formulated together with one or more pharmaceutically acceptable carriers or excipient. As used herein, the term
30 "pharmaceutically acceptable carrier" means a non-toxic, inert solid, semi-solid or liquid

filler, diluent, encapsulating material or formulation auxiliary of any type. The pharmaceutical compositions of this application can be administered to humans and other animals orally, rectally, parenterally, intracisternally, intravaginally, intraperitoneally, topically (as by powders, ointments, or drops), buccally, or as an oral or nasal spray.

5 Liquid dosage forms for oral administration include pharmaceutically acceptable emulsions, microemulsions, solutions, suspensions, syrups and elixirs. In addition to the active compounds, the liquid dosage forms may contain inert diluents commonly used in the art such as, for example, water or other solvents, solubilizing agents and emulsifiers such as ethyl alcohol, isopropyl alcohol, ethyl carbonate, ethyl
10 acetate, benzyl alcohol, benzyl benzoate, propylene glycol, 1,3-butylene glycol, dimethylformamide, oils (in particular, cottonseed, groundnut, com, germ, olive, castor, and sesame oils), glycerol, tetrahydrofurfuryl alcohol, polyethylene glycols and fatty acid esters of sorbitan, and mixtures thereof. Besides inert diluents, the oral compositions can also include adjuvants such as wetting agents, emulsifying and suspending agents,
15 sweetening, flavoring, and perfuming agents.

 Injectable preparations include, for example, sterile injectable aqueous or oleaginous suspensions formulated according to the known art using suitable dispersing or wetting agents and suspending agents. The sterile injectable preparation may also be a sterile injectable solution, suspension or emulsion in a nontoxic parenterally acceptable
20 diluent or solvent. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution, U.S.P. and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose, any bland fixed oil can be employed including synthetic mono- or diglycerides. In addition, fatty acids such as oleic acid are used in the preparation of injectables.

25 In order to prolong the effect of a drug, it is often desirable to slow the absorption of the drug from subcutaneous or intramuscular injection. This may be accomplished using a liquid suspension of crystalline or amorphous material with poor water solubility. The rate of absorption of the drug then depends upon its rate of dissolution which, in turn, may depend upon crystal size and crystalline form.

Alternatively, delayed absorption of a parenterally administered drug form is accomplished by dissolving or suspending the drug in an oil vehicle.

Solid compositions of a similar type may also be employed as fillers in soft and hard filled gelatin capsules using such excipients as lactose or milk sugar as well
5 as high molecular weight polyethylene glycols and the like.

The active compounds can also be in micro-encapsulated form with one or more excipients as noted above. The solid dosage forms of tablets, dragees, capsules, pills, and granules can be prepared with coatings and shells such as enteric coatings, release controlling coatings and other coatings well known in the pharmaceutical
10 formulating art. In such solid dosage forms the active compound may be admixed with at least one inert diluent such as sucrose, lactose, or starch. Such dosage forms may also comprise, as is normal practice, additional substances other than inert diluents, *e.g.*, tableting lubricants and other tableting aids such as magnesium stearate and microcrystalline cellulose. In the case of capsules, tablets, and pills, the dosage forms
15 may also comprise buffering agents.

Dosage forms for topical or transdermal administration of a compound as disclosed in the foregoing embodiments (*e.g.*, a compound of Structure (I), (II) or (III)) include ointments, pastes, creams, lotions, gels, powders, solutions, sprays, inhalants, or patches. For example, the active component may be ad-mixed under sterile conditions
20 with a pharmaceutically acceptable carrier or excipient and any needed preservatives or buffers as may be required.

Transdermal patches have the added advantage of providing controlled delivery of a compound to the body. Such dosage forms can be made by dissolving or dispensing the compound in the proper medium. Absorption enhancers can also be used
25 to increase the flux of the compound across the skin. The rate can be controlled by either providing a rate controlling membrane or by dispersing the compound in a polymer matrix or gel.

According to the methods of treatment of the present disclosure, disorders are treated or prevented in a subject, such as a human or other animal, by administering
30 to the subject a therapeutically effective amount of a compound according to any one of

the foregoing embodiments, in such amounts and for such time as is necessary to achieve the desired result. As is well understood in the medical arts a therapeutically effective amount of a compound will be at a reasonable benefit/risk ratio applicable to any medical treatment.

5 In general, compounds (*e.g.*, compounds of Structure (I), (II) or (III)) will be administered in therapeutically effective amounts via any of the usual and acceptable modes known in the art, either singly or in combination with one or more other therapeutic agents. A therapeutically effective amount may vary widely depending on the severity of the disease, the age and relative health of the subject, the potency of the
10 compound used and other factors.

 In general, satisfactory results are indicated to be obtained systemically at daily dosages of from about 0.03 to 2.5 mg/kg per body weight. An indicated daily dosage in the larger mammal, *e.g.*, humans, is in the range from about 0.5 mg to about 250 mg, about 5 mg to about 150 mg, about 5 mg to about 100 mg, about 10 mg to about 75 mg,
15 about 10 mg to about 50 mg, such as 10, 20, 30, 40, or about 50 mg, conveniently administered, *e.g.*, in divided doses up to four times a day or in retard form. Suitable unit dosage forms for oral administration comprise from ca. 1 to 60 mg active ingredient.

 In certain embodiments, a therapeutic amount or dose of the compound (*e.g.*, compounds of Structure (I), (II) or (III)) may range from about 0.1 mg/kg to about
20 500 mg/kg, alternatively from about 1 mg/kg to about 50 mg/kg. In general, treatment regimens according to the present application comprise administration to a patient in need of such treatment from about 10 mg to about 1000 mg of the compound(s) per day in single or multiple doses. Therapeutic amounts or doses will also vary depending on route of administration, as well as the possibility of co-usage with other agents.

25 Upon improvement of a subject's condition, a maintenance dose of a compound, composition or combination of this application may be administered, if necessary. Subsequently, the dosage or frequency of administration, or both, may be reduced, as a function of the symptoms, to a level at which the improved condition is retained when the symptoms have been alleviated to the desired level, treatment should

cease. The subject may, however, require intermittent treatment on a long-term basis upon any recurrence of disease symptoms.

It will be understood, however, that the total daily usage of the compounds (*e.g.*, compounds of Structure (I), (II) or (III)) and compositions thereof will be decided
5 by an attending physician within the scope of sound medical judgment. The specific inhibitory dose for any particular patient will depend upon a variety of factors including the disorder being treated and the severity of the disorder; the activity of the specific compound employed; the specific composition employed; the age, body weight, general health, sex and diet of the patient; the time of administration, route of administration, and
10 rate of excretion of the specific compound employed; the duration of the treatment; drugs used in combination or coincidental with the specific compound employed; and like factors well known in the medical arts.

The application also provides for a pharmaceutical combination, *e.g.*, a kit, comprising:

- 15 a) a first agent which is a compound of the application as disclosed herein, in free form or in pharmaceutically acceptable salt form, and
- b) at least one co-agent. The kit can comprise instructions for its administration.

Methods for preparing such dosage forms are known to those skilled in
20 the art (see, *e.g.*, REMINGTON'S PHARMACEUTICAL SCIENCES, 18th ED., Mack Publishing Co., Easton, PA (1990)). The dosage forms typically include a conventional pharmaceutical carrier or excipient and may additionally include other medicinal agents, carriers, adjuvants, diluents, tissue permeation enhancers, solubilizers, and the like. Appropriate excipients can be tailored to the particular dosage form and route of
25 administration by methods well known in the art (see, *e.g.*, REMINGTON'S PHARMACEUTICAL SCIENCES, 18th ED., Mack Publishing Co., Easton, PA (1990)).

GENERAL METHODS

The following examples are provided by way of illustration only and not by way of limitation. Those of skill will readily recognize a variety of noncritical parameters which could be changed or modified to yield essentially similar results.

5 If not otherwise stated, chromatography refers to flash chromatography conducted on silica gel. Amine column refers to flash chromatography conducted on Redisep Rf Gold high performance amine column.

HPLC purification was performed on a Waters preparative reverse phase HPLC system with the combination of UV/MS detectors (254 nm and 280 nm) and
 10 XBridge Prep (19 × 50 mm) C₁₈ 10μM OBD column. Eluents were a mixture of water and acetonitrile (with 0.05 % trifluoroacetic acid). Flow rate was typically 50 mL/min with a linear gradient of water in acetonitrile from 5-95% in 8 minutes. The injection volume was from 0.2 to 1 mL with maximum 20 mg per load.


Abbreviations

15	μ	micro
	°C	degrees Celsius
	Ac	acetyl
	anhyd	anhydrous
	aq	aqueous
20	atm	atmosphere(s)
	Bn	benzyl
	Boc	<i>tert</i> -butoxycarbonyl
	Bu	butyl
	calcd	calculated
25	Cbz	benzyloxycarbonyl
	CPME	cyclopentyl methyl ether
	concd	concentrated
	conc	concentration
	DCC	<i>N,N'</i> -dicyclohexylcarbodiimide
30	DIEA	<i>N,N</i> -diisopropylethylamine
	DMAP	4-(<i>N,N</i> -dimethylamino)pyridine
	DMF	dimethylformamide
	DMSO	dimethylsulfoxide
	EDC	<i>N</i> -(3-dimethylaminopropyl)- <i>N'</i> -ethylcarbodiimide hydrochloride
35	equiv	equivalent
	ES	electrospray
	Et	ethyl
	Et ₂ O	diethyl ether
	g	gram(s)


	h	hour(s)
	HATU	<i>N</i> -[(Dimethylamino)-1 <i>H</i> -1,2,3-triazolo-[4,5- <i>b</i>]pyridin-1-ylmethylene]- <i>N</i> -methylmethanaminium hexafluorophosphate <i>N</i> -oxide
	HBTU	<i>N,N,N',N'</i> -Tetramethyl- <i>O</i> -(1 <i>H</i> -benzotriazol-1-yl)uronium
5	hexafluorophosphate, <i>O</i> -(Benzotriazol-1-yl)-	
	HPLC	high-performance liquid chromatography/high-performance liquid chromatography
	HOBt	1-hydroxybenzotriazole hydrate
	<i>i</i> PrOH	<i>iso</i> -propanol
10	L	liter(s)
	LiOH	lithium hydroxide
	m	milli
	M	molar
	MeCN	acetonitrile
15	min	minute(s)
	mL	milliliter
	mol	mole; molecular (as in mol wt)
	MS	mass spectrometry
	MW	molecular weight
20	NBS	<i>N</i> -bromosuccinimide
	NHS	<i>N</i> -hydroxysuccinimide
	NMM	4-methylmorpholine
	NMR	nuclear magnetic resonance
	o	ortho
25	obsd	observed
	p	para
	Ph	phenyl
	ppt	precipitate
	Pr	propyl
30	psi	pounds per square inch
	temp	temperature
	TFA	trifluoroacetic acid
	THF	tetrahydrofuran

EXAMPLE 1


PREPARATION OF METHYL (3*R*,5*S*)-1-((*R*)-2-AMINO-4-PHENYLBUTANOYL)-5-((4-CARBAMIMIDOYLBENZYL)CARBAMOYL)PYRROLIDINE-3-CARBOXYLATE
DITRIFLUOROACETATE (COMPOUND I-1)




Step 1: To 1-(*tert*-butyl) 2-methyl (2*S*,4*R*)-4-cyanopyrrolidine-1,2-dicarboxylate (90 mg, 0.38 mmol, 1 eq.) and *N*-hydroxysuccinamide (48 mg, 0.42 mmol, 1.1 eq.) in dichloromethane (3.8 mL, 0.1M), was added DCC (93 mg, 0.38 mmol, 1.1 eq.). After 1 h stirring, benzyl ((4-(aminomethyl)phenyl)(imino)methyl)carbamate hydrochloride (120 mg, 0.38 mmol, 1 eq.) suspended in saturated sodium bicarbonate (3.8 mL, 0.1 M) was added to the solution. The mixture was allowed to stir for another hour, after which, the layers were partitioned and the organic was run through a 0.45 micron filter and concentrated. The resulting residue was purified on a silica column using EtOAc/Heptane to yield *tert*-butyl (2*S*,4*R*)-2-((4-(*N*-((benzyloxy)carbonyl)carbamimidoyl)benzyl)carbamoyl)-4-cyanopyrrolidine-1-carboxylate (180 mg, 95%) as an off-white film.



Step 2: To a solution of *tert*-butyl (2*S*,4*R*)-2-((4-(*N*-((benzyloxy)carbonyl)carbamimidoyl)benzyl)carbamoyl)-4-cyanopyrrolidine-1-carboxylate (180 mg, 0.36 mmol, 1 eq.) in methanol (1.8 mL, 0.2M), was added HCl in dioxane (4M, 1.8 mL, 0.2M) and the reaction mixture was stirred until reaction was complete as monitored by LCMS. The reaction mixture was then concentrated *in vacuo* to afford methyl (3*R*,5*S*)-5-((4-(*N*-((benzyloxy)carbonyl)carbamimidoyl)benzyl)carbamoyl)pyrrolidine-3-carboxylate hydrochloride as a clear film which was carried forward without further purification.



Step 3: To a solution of methyl (3*R*,5*S*)-5-((4-(*N*-((benzyloxy)carbonyl)carbamimidoyl)benzyl)carbamoyl)pyrrolidine-3-carboxylate hydrochloride (0.36 mmol, 1 eq.) and *D*-homophenylalanine (121 mg, 0.43 mmol, 1.2 eq.), in anhydrous *N,N*-dimethylformamide (7.2 mL, 0.05M) was added DIEA (250 μ L, 1.2 mmol, 4 eq.) followed by hydroxybenzotriazole (60 mg, 0.43 mmol, 1.2 eq.) and 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide (83 mg, 0.43 mmol, 1.2 eq.). After stirring overnight, reaction mixture was diluted with ethyl acetate then extracted with saturated ammonium chloride, followed by saturated sodium bicarbonate, then water (2 times), then brine. The organic layer was then concentrated and purified on a silica column using EtOAc/Heptane to afford methyl (3*R*,5*S*)-5-((4-(*N*-((benzyloxy)carbonyl)carbamimidoyl)benzyl)carbamoyl)-1-((*R*)-2-((*tert*-butoxycarbonyl)amino)-4-phenylbutanoyl)pyrrolidine-3-carboxylate (34 mg, 13%).




Step 4: To a solution of methyl (3*R*,5*S*)-5-((4-(*N*-((benzyloxy)carbonyl)carbamimidoyl)benzyl)carbamoyl)-1-((*R*)-2-((*tert*-butoxycarbonyl)amino)-4-phenylbutanoyl)pyrrolidine-3-carboxylate (30 mg, 0.04 mmol, 1 eq.) in dichloromethane (100 μ L, 0.2M) was added trifluoroacetic acid (40 μ L, 0.45 mmol, 20 eq.). After stirring for 3 h reaction was complete as monitored by LCMS and the reaction mixture was concentrated to dryness and dissolved in anhydrous methanol (200 μ L, 0.1M) to which 10% palladium on carbon (5 mg, 15 wt %) was added and the reaction mixture was sparged with argon for 10 minutes, followed by hydrogen for 10 minutes. The reaction mixture was left under a balloon of hydrogen for approximately 30 mins until reaction completion which was monitored by LCMS. The

reaction mixture then sparged with argon for 10 minutes and filtered over celite. The filtrate was then concentrated *in vacuo* to afford methyl (3*R*,5*S*)-1-((*R*)-2-amino-4-phenylbutanoyl)-5-((4-carbamimidoylbenzyl)carbamoyl)pyrrolidine-3-carboxylate ditrifluoroacetate (27.1 mg, 98% over 2 steps) as a solid white powder following lyophilization.

EXAMPLE 2

PREPARATION OF (2*S*,4*S*)-1-((*R*)-2-AMINO-4-PHENYLBUTANOYL)-*N*-(4-CARBAMIMIDOYLBENZYL)-4-(METHOXYMETHYL)PYRROLIDINE-2-CARBOXAMIDE DIHYDROCHLORIDE (COMPOUND I-2)




10

(2*S*,4*S*)-1-((*R*)-2-amino-4-phenylbutanoyl)-*N*-(4-carbamimidoylbenzyl)-4-(methoxymethyl)pyrrolidine-2-carboxamide dihydrochloride was synthesized according to the procedures for Example 1 except that in Step 4, the Boc group was deprotected with HCl in isopropanol with anhydrous methanol as the solvent.

EXAMPLE 3

PREPARATION OF (2*S*,4*R*)-1-((*R*)-2-AMINO-4-PHENYLBUTANOYL)-*N*-(4-CARBAMIMIDOYLBENZYL)-4-PHENYLPYRROLIDINE-2-CARBOXAMIDE DIHYDROCHLORIDE (COMPOUND I-3)




5

(2*S*,4*R*)-1-((*R*)-2-amino-4-phenylbutanoyl)-*N*-(4-carbamimidoylbenzyl)-4-phenylpyrrolidine-2-carboxamide dihydrochloride was synthesized according to the procedures for Example 2.


EXAMPLE 4

10 PREPARATION OF ((*R*)-1-((2*S*,4*R*)-2-((4-CARBAMIMIDOYLBENZYL)CARBAMOYL)-4-PHENYLPYRROLIDIN-1-YL)-1-OXO-4-PHENYLBUTAN-2-YL)GLYCINE (COMPOUND I-4)



Step 1: To a solution of benzyl ((4-(((2*S*,4*R*)-1-((*R*)-2-amino-4-phenylbutanoyl)-4-phenylpyrrolidine-2-carboxamido)methyl)phenyl)(imino)methyl)carbamate hydrochloride (114 mg, 0.17 mmol, 1 eq.) in anhydrous acetonitrile (1.1 mL, 0.16M), DIEA (120 μ L, 1.5M) was added followed by benzyl bromoacetate (35 μ L, 0.22 mmol, 1.3 eq.). The reaction mixture was stirred for 16 h, at which point additional benzyl bromoacetate (35 μ L, 0.22 mmol, 1.3 eq.) was added and the reaction was stirred for a further 24 h. The reaction mixture was then diluted with ethyl acetate and washed four times with water, then one time with brine. The organic layer was concentrated and purified on an amine column using EtOAc/Heptane to afford benzyl ((*R*)-1-((2*S*,4*R*)-2-((4-*N*-


((benzyloxy)carbonyl)carbamimidoyl)benzyl)carbamoyl)-4-phenylpyrrolidin-1-yl)-1-oxo-4-phenylbutan-2-yl)glycinate (43 mg, 32%).



Step 2: To a solution of benzyl ((*R*)-1-((2*S*,4*R*)-2-((4-(*N*-
5 ((benzyloxy)carbonyl)carbamimidoyl)benzyl)carbamoyl)-4-phenylpyrrolidin-1-yl)-1-oxo-4-phenylbutan-2-yl)glycinate (43 mg, 0.056 mmol, 1 eq.) in anhydrous methanol (560 μ L, 0.1M) was added 10% palladium on carbon (20 mg, 50 wt %) and the reaction mixture was sparged with argon for 10 minutes, followed by hydrogen for 10 minutes. The reaction mixture was left under a balloon of hydrogen for approximately 2 hr until
10 reaction completion which was monitored by LCMS. The reaction mixture then sparged with argon for 10 minutes and filtered over Celite. The filtrate was then concentrated *in vacuo* to afford ((*R*)-1-((2*S*,4*R*)-2-((4-carbamimidoylbenzyl)carbamoyl)-4-phenylpyrrolidin-1-yl)-1-oxo-4-phenylbutan-2-yl)glycine (22.4 mg, 74%) as a solid white powder following lyophilization.

15 EXAMPLE 5


PREPARATION OF (2*S*,4*S*)-1-((*R*)-2-AMINO-4-PHENYLBUTANOYL)-*N*-(4-CARBAMIMIDOYLBENZYL)-4-PHENYLPYRROLIDINE-2-CARBOXAMIDE DIHYDROCHLORIDE
(COMPOUND I-11)




20 (2*S*,4*S*)-1-((*R*)-2-amino-4-phenylbutanoyl)-*N*-(4-carbamimidoylbenzyl)-4-phenylpyrrolidine-2-carboxamide dihydrochloride was synthesized according to the procedures for Example 2.

EXAMPLE 6


PREPARATION OF (2*S*)-1-((*R*)-2-AMINO-4-PHENYLBUTANOYL)-*N*-(4-CARBAMIMIDOYLBENZYL)-4-PHENYLPYRROLIDINE-2-CARBOXAMIDE DIHYDROCHLORIDE
(COMPOUNDS I-26 AND I-27)




Step 1: To di-*tert*-butyl (2*S*,4*R*)-4-hydroxypyrrolidine-1,2-dicarboxylate (500 mg, 1.7 mmol, 1 eq.) and triethylamine (840 μ L, 6 mmol, 3 eq.) in dichloromethane (8 mL, 0.25M), was added DMAP (24 mg, 0.2 mmol, 0.1 eq.) followed by tosyl chloride (460 mg, 2.4 mmol, 1.2 eq.). After 16 h stirring, the mixture was further diluted with
10 dichloromethane, washed twice with 1N HCl, followed by 1N NaOH 2 times then by brine. The organic layer was dried over sodium sulfate, concentrated under vacuum and the resulting residue purified on a silica column using EtOAc/Heptane to yield di-*tert*-butyl (2*S*,4*R*)-4-(tosyloxy)pyrrolidine-1,2-dicarboxylate (690 mg, 90%) as a white solid.




Step 2: A mixture consisting of di-*tert*-butyl (2*S*,4*R*)-4-(tosyloxy)pyrrolidine-1,2-dicarboxylate (100 mg, 0.23 mmol, 1 eq.), 1*H*-pyrazole (30 mg, 0.46 mmol, 2 eq.) and cesium carbonate (98 mg, 0.3 mmol, 1.5 eq.) in *N,N*-dimethylformamide (1 mL, 0.2M) was sealed and heated to 60 °C. After stirring for 16 h, the mixture was partitioned between ethyl acetate and saturated ammonium chloride.
20 The organic layer was further washed with water followed by brine then dried over sodium sulfate and concentrated. The resulting residue was purified on a silica column using EtOAc/Heptane to yield di-*tert*-butyl (2*S*)-4-(1*H*-pyrazol-1-yl)pyrrolidine-1,2-dicarboxylate (41.1 mg, 53%) as a white solid.




Step 3: To a solution of di-*tert*-butyl (2*S*)-4-(1*H*-pyrazol-1-yl)pyrrolidine-1,2-dicarboxylate (41 mg, 0.12 mmol, 1 eq.) in methanol (2.4 mL, 0.05M) was added hydrochloric acid in dioxane (4N solution, 300 μ L, 0.4 mmol, 20 eq.). After stirring for 16 h, the reaction was incomplete, so additional hydrochloric acid in dioxane (4N solution, 300 μ L, 0.4 mmol, 20 eq.) was added and the reaction mixture was heated to 40 $^{\circ}$ C. After 3d the reaction mixture was concentrated *in vacuo* and carried to the next step without further purification.



Step 4: To a solution of methyl (2*S*)-4-(1*H*-pyrazol-1-yl)pyrrolidine-2-carboxylate hydrochloride (0.12 mmol, 1 eq.) and *D*-homophenylalanine (40 mg, 0.14 mmol, 1.2 eq.), in anhydrous *N,N*-dimethylformamide (1.2 mL, 0.05M) was added DIEA (84 μ L, 0.48 mmol, 4 eq.) followed by hydroxybenzotriazole (20 mg, 0.14 mmol, 1.2 eq.) and 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide (28 mg, 0.14 mmol, 1.2 eq.). After stirring overnight, the reaction mixture was diluted with ethyl acetate. The resulting solution was washed with saturated ammonium chloride, followed by saturated sodium bicarbonate, water then brine. The organic layer was concentrated and the resulting residue purified on a silica column using EtOAc/Heptane to afford methyl (2*S*)-1-((*R*)-2-((*tert*-butoxycarbonyl)amino)-4-phenylbutanoyl)-4-(1*H*-pyrazol-1-yl)pyrrolidine-2-carboxylate (34 mg, 58%).





Step 5: Methyl (2*S*)-1-((*R*)-2-((*tert*-butoxycarbonyl)amino)-4-phenylbutanoyl)-4-(1*H*-pyrazol-1-yl)pyrrolidine-2-carboxylate was saponified using THF, H₂O, and LiOH and stirring at room temperature; the reaction mixture was conc'd to remove THF and to the aqueous solution was added 10% KHSO₄ solution. The resulting ppt collected via vacuum filtration and washed with hexanes. To the product of saponification, (2*S*)-1-((*R*)-2-((*tert*-butoxycarbonyl)amino)-4-phenylbutanoyl)-4-(1*H*-pyrazol-1-yl)pyrrolidine-2-carboxylic acid (0.07 mmol), benzyl ((4-(aminomethyl)phenyl)(imino)methyl)carbamate hydrochloride (29 mg, 0.09 mmol), was added DIEA (50 μ L, 0.28 mmol) and DMF (540 μ L). After cooling the reaction mixture to 0 $^{\circ}$ C, HBTU (35 mg, 0.12 mmol) was added in a single portion and the reaction was left to stir at RT until completion. The mixture was concentrated and purified by silica column using EtOAc/Heptane which produced two crude products, one which was dissolved in ethyl acetate and washed with 0.1N HCl to afford an isomer of benzyl ((4-(((2*S*)-1-((*R*)-2-amino-4-phenylbutanoyl)-4-(1*H*-pyrazol-1-yl)pyrrolidine-2-carboxamido)methyl)phenyl)(imino)methyl)carbamate (8.5 mg, 13%). The other crude material was purified on a second column using MeOH/dichloromethane to afford the another isomer of benzyl ((4-(((2*S*)-1-((*R*)-2-amino-4-phenylbutanoyl)-4-(1*H*-pyrazol-1-yl)pyrrolidine-2-carboxamido)methyl)phenyl)(imino)methyl)carbamate (9.8 mg, 15%).



Step 6: Each isomer of benzyl ((4-(((2*S*)-1-((*R*)-2-amino-4-phenylbutanoyl)-4-(1*H*-pyrazol-1-yl)pyrrolidine-2-carboxamido)methyl)phenyl)(imino)methyl)carbamate was deprotected using Step 4 of Example 1 except that the Boc group was deprotected with HCl in isopropanol with anhydrous methanol as the solvent to afford the title compounds.

EXAMPLE 7

PREPARATION OF (2*S*,4*S*)-*N*-(((1*H*-PYRROLO[3,2-*C*]PYRIDIN-2-*YL*)METHYL)-1-((*R*)-2-AMINO-4-PHENYLBUTANOYL)-4-(METHOXYMETHYL)PYRROLIDINE-2-CARBOXAMIDE DITRIFLUOROACETATE (COMPOUND I-24)




EXAMPLE 8

PREPARATION OF (2*S*,4*S*)-*N*-((1*H*-pyrrolo[3,2-*c*]pyridin-2-yl)methyl)-1-((*R*)-2-amino-4-phenylbutanoyl)-4-cyanopyrrolidine-2-carboxamide

DITRIFLUOROACETATE

(COMPOUND I-25)




(2*S*,4*S*)-*N*-((1*H*-pyrrolo[3,2-*c*]pyridin-2-yl)methyl)-1-((*R*)-2-amino-4-phenylbutanoyl)-4-cyanopyrrolidine-2-carboxamide ditrifluoroacetate was synthesized according to the procedures for Example 1 except the hydrogenation deprotection was not required and TFA/dichloromethane was used to deprotect the Boc group.

EXAMPLE 9

PREPARATION OF (2*S*,4*S*)-1-((*R*)-2-amino-4-phenylbutanoyl)-*N*-(4-carbamimidoylbenzyl)-4-cyanopyrrolidine-2-carboxamide

DITRIFLUOROACETATE (COMPOUND I-6)











(2*S*,4*S*)-1-((*R*)-2-amino-4-phenylbutanoyl)-*N*-(4-carbamimidoylbenzyl)-4-cyanopyrrolidine-2-carboxamide ditrifluoroacetate was synthesized according to the procedures for Example 1 except TFA/dichloromethane was used to deprotect the Boc group.





Table 1 lists compounds of the Examples described above, as well as additional compounds that may be prepared according to methods analogous to those





described for the compounds above and other methods known to a person having skill in the art. In some embodiments, the compound is a compound in Table 1.





Table 1. Exemplary compounds





Cmp No.	Structure	Salt	Exact Observed Mass (ES ⁺ ; M+H)	Exact Calc. Mass
I-1		2TFA	466.44	465.24
I-2		2HCl	452.46	451.26
I-3		2HCl	484.56	483.26
I-4		-	542.47	541.64


Cmp No.	Structure	Salt	Exact Observed Mass (ES ⁺ ; M+H)	Exact Calc. Mass
I-5		-	-	-
I-6		2TFA	433.0	432.23
I-7		-	-	-
I-8		-	-	-

Cmp No.	Structure	Salt	Exact Observed Mass (ES ⁺ ; M+H)	Exact Calc. Mass
I-9		-	-	-
I-10		-	-	-
I-11		2HCl	482.3 (M-H)	483.26
I-12		-	-	-

Cmp No.	Structure	Salt	Exact Observed Mass (ES ⁺ ; M+H)	Exact Calc. Mass
I-13		-	-	-
I-14		-	-	-
I-15		-	-	-
I-16		-	-	-

Cmp No.	Structure	Salt	Exact Observed Mass (ES ⁺ ; M+H)	Exact Calc. Mass
I-17		-	-	-
I-18		-	-	-
I-19		-	-	-
I-20		-	-	-

Cmp No.	Structure	Salt	Exact Observed Mass (ES ⁺ ; M+H)	Exact Calc. Mass
I-21		-	-	-
I-22		2HCl	474.5	473.25
I-23		2HCl	474.5	473.25
I-24		2TFA	450.4	449.24

Cmp No.	Structure	Salt	Exact Observ ed Mass (ES+; M+H)	Exact Calc. Mass
I-25		2TFA	431.4	430.21

EXAMPLE 10

ENZYMATIC ASSAY FOR MASP-2

The MASP-2 assay utilizes a fluorogenic substrate, based on the cleavage site for its natural substrate C2. The assay is run at room temperature in an assay buffer containing 20 mM HEPES, pH 7.4, 140 mM NaCl and 0.1% Tween 20. Assay parameters are adjusted such that the assay is linear with respect to time, enzyme, and substrate concentrations. Under these optimized assays conditions, IC₅₀ values are equivalent to Ki values, except in a few cases of "tight binding" inhibitors. Cases of "tight binding" or possible "slow binding" inhibitors are handled by the methods described in Copeland R.A. (2013) Evaluation of Enzyme Inhibitors in Drug Discovery. 2nd Ed., John Wiley and Sons, Inc., Chapters 5-7.

The MASP-2 assay protocol is carried out as follows. Test compounds are serially diluted in DMSO and then 100 nL of each dilution is transferred to the assay plate(s). 10 µL of Assay Buffer is added, followed by 15 µL of Enzyme MASP-2 (CCP1-CCP2-SP) in Assay Buffer. 15 µL of Substrate in Assay Buffer is then added and mixed to start the reactions. After 20 min at room temperature, 15 µL of a stop solution (0.1 M acetic acid) is added, mixed and the plates are read on a SpectraMax i3x Microplate Reader and exported as Excel files. Each assay plate included a "no inhibitor" (DMSO Only) control, a "no enzyme" control and a reference inhibitor control. % Activity values = 100*(average test comp. fluorescence – average "no enzyme" fluorescence) / (average

"DMSO only" fluorescence – average "no enzyme" fluorescence). IC₅₀ and Ki values are very reproducible, falling well within ± 2-fold.

The results of biological assays for the compounds listed in Table 1 are listed in Table 2, below.

5 Table 2. MASP-2 Inhibition for exemplary compounds

Compound	MASP-2 Ki (μM)	Compound	MASP-2 Ki (μM)	Compound	MASP-2 Ki (μM)
I-1	****	I-2	****	I-3	****
I-4	--	I-5	--	I-6	****
I-7	--	I-8	--	I-9	--
I-10	--	I-11	****	I-12	--
I-13	--	I-14	--	I-15	--
I-16	--	I-17	--	I-18	--
I-19	--	I-20	--	I-21	--
I-22	****	I-23	--	I-24	--
I-25	--	I-26	****	I-27	****
I-28	****	I-29	****	-	--

MASP-2 Inhibition Ki Values:

- 10
- *

**

--

K_i greater than 10 μM

K_i between 2.5 - 10 μM

K_i between 0.5 - 2.5 μM

K_i of less than 0.5 μM

not tested

15


It is understood that the examples and embodiments described herein are for illustrative purposes only and that various modifications or changes in light thereof will be suggested to persons skilled in the art and are to be included within the spirit and purview of this application and scope of the appended claims. Each reference, including without limitation all patent, patent applications, and publications, cited in the present application is incorporated herein by reference in its entirety for all purposes.

20

This application claims the benefit of priority to U.S. Provisional Application No. 62/943,629, filed December 4, 2019, which application is hereby incorporated by reference in its entirety.

CLAIMS

1. A compound having the following Structure (I):



(I)

or a stereoisomer, tautomer, or pharmaceutically acceptable salt thereof, wherein:

R¹ is a substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, substituted or unsubstituted cycloalkyl, or a substituted or unsubstituted heterocyclyl;

R^{2a} , R^{2b} , R^{2c} , R^{2d} , R^{2e} , R^{2f} , R^{2g} , R^{2h} , R^{2i} , and R^{2j} are each independently selected from the group consisting of hydrogen, halo, $C(=O)OR^5$, $OC(=O)R^5$, hydroxyalkyl, alkoxy, alkoxyalkyl, haloalkoxy, cyano, aminylalkyl, carboxyalkyl, NR^5R^6 , $C(=O)NR^5R^6$, $N(R^5)C(=O)R^6$, $NR^5C(=O)NR^6$, $S(O)_t$, SR^5 , nitro, $N(R^5)C(O)OR^6$, $C(=NR^5)NR^6R^7$, $N(R^5)C(=NR^6)NR^7R^8$, $S(O)R^5$, $S(O)NR^5R^6$, $S(O)_2R^5$, $N(R^5)S(O)_2R^6$, $S(O)_2NR^5R^6$, aryl, heteroaryl, heterocyclyl, cycloalkyl, and oxo, provided that at least one of R^{2a} , R^{2b} , R^{2c} , R^{2d} , R^{2e} , R^{2f} , R^{2g} , R^{2h} , R^{2i} , and R^{2j} is not hydrogen;

$$\mathbf{R}^3 \text{ is } \mathbf{NR}^{3a}\mathbf{R}^{3b}.$$

R^{3a} and R^{3b} are each independently hydrogen, alkyl, hydroxyalkyl, haloalkyl, alkoxyalkyl, heterocyclyl, heteroaryl, heterocyclylalkyl, heteroarylalkyl, cycloalkyl, $(CH_2)_nC(=O)OR^6$, or $(CH_2)_nP(=O)(OR^6)_2$; or R^{3a} and R^{3b} , together with the nitrogen to which they are attached, form an optionally substituted 4-7 membered heteroaryl or an optionally substituted 4-7 membered heterocyclyl; or R^{3a} and R^4 together with the nitrogen and carbon to which they are attached, respectively, form an optionally substituted 4-7 membered heterocyclyl, and R^{3b} is hydrogen, alkyl, hydroxyalkyl,

haloalkyl, alkoxyalkyl, heterocyclyl, heteroaryl, heterocyclylalkyl, heteroarylalkyl, cycloalkyl, $(\text{CH}_2)_n\text{C}(=\text{O})\text{OR}^6$, or $(\text{CH}_2)_n\text{P}(=\text{O})(\text{OR}^6)_2$;

R^4 is a substituted or unsubstituted aryl, a substituted or unsubstituted heteroaryl, a substituted or unsubstituted cycloalkyl, or a substituted or unsubstituted heterocyclyl when n is 2, 3, 4, 5, or 6; or R^4 is a substituted or unsubstituted monocyclic heteroaryl, or a substituted or unsubstituted heterocyclyl when n is 0 or 1; or R^4 and R^{3a} together with the carbon and nitrogen to which they are attached, respectively, form an optionally substituted 4-7 membered heterocyclyl;

R^5 , R^6 , R^7 , and R^8 are, at each occurrence, independently hydrogen, alkyl, hydroxyalkyl, haloalkyl, alkoxyalkyl, carboxyalkyl, heterocyclyl, heteroaryl, or cycloalkyl;

X is a direct bond, $-\text{CR}^{2e}\text{R}^{2f}-$, or $-\text{CR}^{2e}\text{R}^{2f}-\text{CR}^{2g}\text{R}^{2h}-$;

Y is a direct bond or $-\text{CR}^{2i}\text{R}^{2j}-$;

n is an integer from 0-6; and

t is an integer from 1-3.

2. The compound of claim 1, wherein R^1 is a substituted or unsubstituted aryl.

3. The compound of claim 1 or 2, wherein R^1 is a substituted or unsubstituted $\text{C}_6\text{-C}_{10}$ aryl.

4. The compound of any one of claims 1-3, wherein R^1 is a substituted or unsubstituted phenyl.

5. The compound of any one of claims 1-4, wherein R^1 is a substituted phenyl.

6. The compound of any one of claims 1-5, wherein R^1 is phenyl substituted with one or more of R^{1a} , R^{1b} , R^{1c} , R^{1d} , and R^{1e} wherein R^{1a} , R^{1b} , R^{1c} , R^{1d} , and

R^{1e} are each independently selected from the group consisting of C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, halo, C_{1-6} haloalkyl, aminylalkyl, hydroxyalkyl, cyano, nitro, OR^9 , SR^9 , $C(O)R^9$, $C(O)NR^9R^{10}$, $C(O)OR^9$, $OC(O)R^9$, $OC(O)OR^9$, $OC(O)NR^9R^{10}$, NR^9R^{10} , $N(R^9)C(O)R^{10}$, $N(R^9)C(O)NR^{10}R^{11}$, $N(R^9)C(O)OR^{10}$, $C(=NR^9)NR^{10}R^{11}$, $C(=NOR^9)NR^{10}R^{11}$, $C(=NOC(O)R^9)NR^{10}R^{11}$, $C(=NR^9)N(R^{10})C(O)OR^{11}$, $N(R^9)C(=NR^{10})NR^{11}R^{12}$, $S(O)R^9$, $S(O)NR^9R^{10}$, $S(O)_2R^9$, $N(R^9)S(O)_2R^{10}$, $S(O)_2NR^9R^{10}$, oxo, substituted or unsubstituted C_{6-10} aryl, substituted or unsubstituted C_{6-10} arylalkyl, substituted or unsubstituted C_{6-10} aryloxy, substituted or unsubstituted C_{6-10} arylalkoxy, substituted or unsubstituted 5-10 membered heteroaryl, substituted or unsubstituted C_{3-10} cycloalkyl, and substituted or unsubstituted 4-10 membered heterocyclyl,

wherein R^9 , R^{10} , R^{11} , and R^{12} , are, at each occurrence, independently selected from the group consisting of hydrogen, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, hydroxyl, C_{1-6} alkoxy, aryl, arylalkyl, C_{1-6} haloalkyl, C_{1-6} haloalkoxy, C_{1-6} hydroxyalkyl, cycloalkyl, heterocyclyl, and heteroaryl.

7. The compound of claim 6, wherein R^{1a} , R^{1b} , R^{1c} , R^{1d} , or R^{1e} is optionally substituted with one or more substituents selected from the group consisting of halo, CN, OR^{13} , SR^{13} , $C(O)R^{13}$, $C(O)NR^{13}R^{14}$, $C(O)OR^{13}$, $OC(O)R^{13}$, $OC(O)NR^{13}R^{14}$, $NR^{13}R^{14}$, $NR^{13}C(O)R^{14}$, $NR^{13}C(O)NR^{14}R^{15}$, $NR^{13}C(O)OR^{14}$, $C(=NR^{13})NR^{14}R^{15}$, $NR^{13}C(=NR^{14})NR^{15}R^{16}$, $S(O)R^{13}$, $S(O)NR^{13}R^{14}$, $S(O)_2R^{13}$, $NR^{13}S(O)_2R^{14}$, $S(O)_2NR^{13}R^{14}$ and oxo when R^{1a} , R^{1b} , R^{1c} , R^{1d} , or R^{1e} is a substituted C_{6-10} aryl, a substituted C_{6-10} arylalkyl, a substituted C_{6-10} aryloxy, a substituted C_{6-10} arylalkoxy, a substituted 5-10 membered heteroaryl, a substituted C_{3-10} cycloalkyl, and a substituted 4-10 membered heterocyclyl,

wherein R^{13} , R^{14} , R^{15} , and R^{16} are, at each occurrence, independently selected from the group consisting of hydrogen, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, hydroxyl, C_{1-6} alkoxy, aryl, arylalkyl, C_{1-6} haloalkyl, C_{1-6} haloalkoxy, C_{1-6} hydroxyalkyl, cycloalkyl, heterocyclyl, and heteroaryl.


8. The compound of any one of claims 1-7, wherein R^1 is phenyl substituted with at least one substituent selected from the group consisting of OR^9 , $C(=NR^9)NR^{10}R^{11}$, $C(=NOR^9)NR^{10}R^{11}$, $C(=NOC(O)R^9)NR^{10}R^{11}$, and $C(=NR^9)NR^{10}C(O)OR^{11}$.


9. The compound of any one of claims 1-8, wherein R^1 is substituted with at least one substituent selected from the group consisting of $C(=NR^9)NR^{10}R^{11}$, $C(=NOR^9)NR^{10}R^{11}$, $C(=NOC(O)R^9)NR^{10}R^{11}$, and $C(=NR^9)NR^{10}C(O)OR^{11}$.

10. The compound of claim 9, wherein R^1 is substituted with at least one $C(=NR^9)NR^{10}R^{11}$.


11. The compound of claim 9 or 10, wherein R^1 is substituted with at least one $-C(=NH)NH_2$.


12. The compound of any one of claims 7-11, wherein R^1 has one of the following structures:






13. The compound of any one of claims 1-12, wherein R¹ has one of the following structures:





14. The compound of claim 7, wherein R^1 has one of the following structures:



15. The compound of claim 14, wherein R^9 is C_{1-6} alkyl or C_{1-6} haloalkyl.

16. The compound of claim 15, wherein R^9 is methyl.

17. The compound of claim 15, wherein R^9 is trifluoromethyl.

18. The compound of any one of claims 1-4, wherein R^1 is an unsubstituted phenyl.

19. The compound of claim 1, wherein R^1 is a substituted or unsubstituted heteroaryl.

20. The compound of claim 1 or 19, wherein R^1 is a substituted or unsubstituted 5-10 membered heteroaryl.

21. The compound of claim 1 or 19-20, wherein R^1 is a substituted or unsubstituted pyridinyl, pyrrolopyridinyl, or benzoimidazolyl.

22. The compound of any one of claims 1 or 19-21, wherein R^1 is a substituted or unsubstituted pyridin-3-yl, 1H-pyrrolo[2,3-b]pyridine-5-yl, 1H-pyrrolo[3,2-c]pyridine-2-yl, or 1H-benzo[d]imidazol-6-yl.

23. The compound of any one of claims 1 or 19-22, wherein R^1 is pyridin-3-yl, 1H-pyrrolo[2,3-b]pyridine-5-yl, or 1H-benzo[d]imidazol-6-yl substituted with one or more of R^{1a} , R^{1b} , R^{1c} , R^{1d} , and R^{1e} wherein R^{1a} , R^{1b} , R^{1c} , R^{1d} , and R^{1e} are each independently selected from the group consisting of C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, halo, C_{1-6} haloalkyl, aminylalkyl, hydroxyalkyl, cyano, nitro, OR^9 , SR^9 , $C(O)R^9$, $C(O)NR^9R^{10}$, $C(O)OR^9$, $OC(O)R^9$, $OC(O)OR^9$, $OC(O)NR^9R^{10}$, NR^9R^{10} , $N(R^9)C(O)R^{10}$, $N(R^9)C(O)NR^{10}R^{11}$, $N(R^9)C(O)OR^{10}$, $C(=NR^9)NR^{10}R^{11}$, $C(=NOR^9)NR^{10}R^{11}$, $C(=NOC(O)R^9)NR^{10}R^{11}$, $C(=NR^9)N(R^{10})C(O)OR^{11}$, $N(R^9)C(=NR^{10})NR^{11}R^{12}$, $S(O)R^9$, $S(O)NR^9R^{10}$, $S(O)_2R^9$, $N(R^9)S(O)_2R^{10}$, $S(O)_2NR^9R^{10}$, oxo, substituted or unsubstituted C_{6-10} aryl, substituted or unsubstituted C_{6-10} arylalkyl, substituted or unsubstituted C_{6-10} aryloxy, substituted or unsubstituted C_{6-10} arylalkoxy, substituted or unsubstituted 5-10 membered heteroaryl, substituted or unsubstituted C_{3-10} cycloalkyl, and substituted or unsubstituted 4-10 membered heterocyclyl,


wherein R^9 , R^{10} , R^{11} , and R^{12} , are, at each occurrence, independently selected from the group consisting of hydrogen, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, hydroxyl, C_{1-6} alkoxy, aryl, arylalkyl, C_{1-6} haloalkyl, C_{1-6} haloalkoxy, C_{1-6} hydroxyalkyl, cycloalkyl, heterocyclyl, and heteroaryl.


24. The compound of claim 23, wherein R^{1a} , R^{1b} , R^{1c} , R^{1d} , or R^{1e} is optionally substituted with one or more substituents selected from the group consisting


of halo, CN, OR¹³, SR¹³, C(O)R¹³, C(O)NR¹³R¹⁴, C(O)OR¹³, OC(O)R¹³, OC(O)NR¹³R¹⁴, NR¹³R¹⁴, NR¹³C(O)R¹⁴, NR¹³C(O)NR¹⁴R¹⁵, NR¹³C(O)OR¹⁴, C(=NR¹³)NR¹⁴R¹⁵, NR¹³C(=NR¹⁴)NR¹⁵R¹⁶, S(O)R¹³, S(O)NR¹³R¹⁴, S(O)₂R¹³, NR¹³S(O)₂R¹⁴, S(O)₂NR¹³R¹⁴ and oxo when R^{1a}, R^{1b}, R^{1c}, R^{1d}, or R^{1e} is a substituted C₆₋₁₀ aryl, a substituted C₆₋₁₀ arylalkyl, a substituted C₆₋₁₀ aryloxy, a substituted C₆₋₁₀ arylalkoxy, a substituted 5-10 membered heteroaryl, a substituted C₃₋₁₀ cycloalkyl, and a substituted 4-10 membered heterocyclyl,


wherein R¹³, R¹⁴, R¹⁵, and R¹⁶ are, at each occurrence, independently selected from the group consisting of hydrogen, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, hydroxyl, C₁₋₆ alkoxy, aryl, arylalkyl, C₁₋₆ haloalkyl, C₁₋₆ haloalkoxy, C₁₋₆ hydroxyalkyl, cycloalkyl, heterocyclyl, and heteroaryl.

25. The compound of any one of claims 1 or 19-23, wherein R¹ has one of the following structures:









26. The compound of claim 25, wherein R^{1a} or R^{1b} is independently C_{1-6} alkyl, amino, $C(O)R^9$, $C(O)NR^9R^{10}$, $N(R^9)C(O)NR^{10}R^{11}$, $N(R^9)C(O)OR^{10}$, $OC(O)NR^9R^{10}$, or halo.


27. The compound of claim 26, wherein R^{1a} or R^{1b} is methyl or ethyl.

28. The compound of claim 26, wherein R^{1a} or R^{1b} is F, Cl, or Br.


29. The compound of claim 25, wherein each R^{1a} or R^{1b} attached to nitrogen is C_{1-6} alkyl, $C(O)R^9$, $C(O)NR^9R^{10}$, $N(R^9)C(O)NR^{10}R^{11}$, $N(R^9)C(O)OR^{10}$, or $OC(O)NR^9R^{10}$.


30. The compound of claim 29, wherein R^{1a} or R^{1b} is methyl or ethyl.

31. The compound of any one of claims 1 or 19-20, wherein R^1 has one of the following structures:



32. The compound of any one of claims 1 or 19-20, wherein R¹ has one of the following structures:





33. The compound of claim 1, wherein R¹ is a substituted or unsubstituted cycloalkyl.

34. The compound of claim 1 or 33, wherein R¹ is a substituted or unsubstituted C₃-C₆ cycloalkyl.

35. The compound of any one of claims 1 or 33-34, wherein R¹ is a substituted C₃-C₆ cycloalkyl.

36. The compound of claim 35, wherein R¹ is a C₃-C₆ cycloalkyl substituted with one or more of R^{1a}, R^{1b}, R^{1c}, R^{1d}, and R^{1e} wherein R^{1a}, R^{1b}, R^{1c}, R^{1d}, and R^{1e} are each independently selected from the group consisting of C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, C₁₋₆ haloalkyl, aminylalkyl, hydroxyalkyl, nitro, OR⁹, SR⁹, C(O)R⁹, C(O)NR⁹R¹⁰, C(O)OR⁹, OC(O)R⁹, OC(O)OR⁹, OC(O)NR⁹R¹⁰, NR⁹R¹⁰, N(R⁹)C(O)R¹⁰, N(R⁹)C(O)NR¹⁰R¹¹, N(R⁹)C(O)OR¹⁰, C(=NR⁹)NR¹⁰R¹¹, C(=NOR⁹)NR¹⁰R¹¹, C(=NOC(O)R⁹)NR¹⁰R¹¹, C(=NR⁹)N(R¹⁰)C(O)OR¹¹, N(R⁹)C(=NR¹⁰)NR¹¹R¹², S(O)R⁹, S(O)NR⁹R¹⁰, S(O)₂R⁹, N(R⁹)S(O)₂R¹⁰, S(O)₂NR⁹R¹⁰, oxo, substituted or unsubstituted C₆₋₁₀ aryl, substituted or unsubstituted C₆₋₁₀ arylalkyl, substituted or unsubstituted C₆₋₁₀ aryloxy, substituted or unsubstituted C₆₋₁₀ arylalkoxy, substituted or unsubstituted 5-10 membered heteroaryl, substituted or unsubstituted C₃₋₁₀ cycloalkyl, and substituted or unsubstituted 4-10 membered heterocyclyl,

wherein R⁹, R¹⁰, R¹¹, and R¹², are, at each occurrence, independently selected from the group consisting of hydrogen, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl,

hydroxyl, C₁₋₆ alkoxy, aryl, arylalkyl, C₁₋₆ haloalkyl, C₁₋₆ haloalkoxy, C₁₋₆ hydroxyalkyl, cycloalkyl, heterocyclyl, and heteroaryl.

37. The compound of claim 36, wherein R^{1a}, R^{1b}, R^{1c}, R^{1d}, or R^{1e} is optionally substituted with one or more substituents selected from the group consisting of halo, CN, OR¹³, SR¹³, C(O)R¹³, C(O)NR¹³R¹⁴, C(O)OR¹³, OC(O)R¹³, OC(O)NR¹³R¹⁴, NR¹³R¹⁴, NR¹³C(O)R¹⁴, NR¹³C(O)NR¹⁴R¹⁵, NR¹³C(O)OR¹⁴, C(=NR¹³)NR¹⁴R¹⁵, NR¹³C(=NR¹⁴)NR¹⁵R¹⁶, S(O)R¹³, S(O)NR¹³R¹⁴, S(O)₂R¹³, NR¹³S(O)₂R¹⁴, S(O)₂NR¹³R¹⁴ and oxo when R^{1a}, R^{1b}, R^{1c}, R^{1d}, or R^{1e} is a substituted C₆₋₁₀ aryl, a substituted C₆₋₁₀ arylalkyl, a substituted C₆₋₁₀ aryloxy, a substituted C₆₋₁₀ arylalkoxy, a substituted 5-10 membered heteroaryl, a substituted C₃₋₁₀ cycloalkyl, and a substituted 4-10 membered heterocyclyl,

wherein R¹³, R¹⁴, R¹⁵, and R¹⁶ are, at each occurrence, independently selected from the group consisting of hydrogen, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, hydroxyl, C₁₋₆ alkoxy, aryl, arylalkyl, C₁₋₆ haloalkyl, C₁₋₆ haloalkoxy, C₁₋₆ hydroxyalkyl, cycloalkyl, heterocyclyl, and heteroaryl.

38. The compound of claim 34, wherein R¹ is an unsubstituted a C₃-C₆ cycloalkyl.

39. The compound of claim 1, wherein R¹ is a substituted or unsubstituted heterocyclyl.

40. The compound of claim 1 or 39, wherein R¹ is a substituted or unsubstituted 4-10 membered heterocyclyl.

41. The compound of any one of claims 1 or 39-40, wherein R¹ is a substituted 4-10 membered heterocyclyl.

42. The compound of claim 41, wherein R^1 is a 4-10 membered heterocyclyl substituted with one or more of R^{1a} , R^{1b} , R^{1c} , R^{1d} , and R^{1e} wherein R^{1a} , R^{1b} , R^{1c} , R^{1d} , and R^{1e} are each independently selected from the group consisting of C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, halo, C_{1-6} haloalkyl, aminylalkyl, hydroxyalkyl, cyano, nitro, OR^9 , SR^9 , $C(O)R^9$, $C(O)NR^9R^{10}$, $C(O)OR^9$, $OC(O)R^9$, $OC(O)OR^9$, $OC(O)NR^9R^{10}$, NR^9R^{10} , $N(R^9)C(O)R^{10}$, $N(R^9)C(O)NR^{10}R^{11}$, $N(R^9)C(O)OR^{10}$, $C(=NR^9)NR^{10}R^{11}$, $C(=NOR^9)NR^{10}R^{11}$, $C(=NOC(O)R^9)NR^{10}R^{11}$, $C(=NR^9)N(R^{10})C(O)OR^{11}$, $N(R^9)C(=NR^{10})NR^{11}R^{12}$, $S(O)R^9$, $S(O)NR^9R^{10}$, $S(O)_2R^9$, $N(R^9)S(O)_2R^{10}$, $S(O)_2NR^9R^{10}$, oxo, substituted or unsubstituted C_{6-10} aryl, substituted or unsubstituted C_{6-10} arylalkyl, substituted or unsubstituted C_{6-10} aryloxy, substituted or unsubstituted C_{6-10} arylalkoxy, substituted or unsubstituted 5-10 membered heteroaryl, substituted or unsubstituted C_{3-10} cycloalkyl, and substituted or unsubstituted 4-10 membered heterocyclyl,







wherein R^9 , R^{10} , R^{11} , and R^{12} , are, at each occurrence, independently selected from the group consisting of hydrogen, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, hydroxyl, C_{1-6} alkoxy, aryl, arylalkyl, C_{1-6} haloalkyl, C_{1-6} haloalkoxy, C_{1-6} hydroxyalkyl, cycloalkyl, heterocyclyl, and heteroaryl.







43. The compound of claim 42, wherein R^{1a} , R^{1b} , R^{1c} , R^{1d} , or R^{1e} is optionally substituted with one or more substituents selected from the group consisting of OR^{13} , SR^{13} , $C(O)R^{13}$, $C(O)NR^{13}R^{14}$, $C(O)OR^{13}$, $OC(O)R^{13}$, $OC(O)NR^{13}R^{14}$, $NR^{13}R^{14}$, $NR^{13}C(O)R^{14}$, $NR^{13}C(O)NR^{14}R^{15}$, $NR^{13}C(O)OR^{14}$, $C(=NR^{13})NR^{14}R^{15}$, $NR^{13}C(=NR^{14})NR^{15}R^{16}$, $S(O)R^{13}$, $S(O)NR^{13}R^{14}$, $S(O)_2R^{13}$, $NR^{13}S(O)_2R^{14}$, $S(O)_2NR^{13}R^{14}$ and oxo when R^{1a} , R^{1b} , R^{1c} , R^{1d} , or R^{1e} is a substituted C_{6-10} aryl, a substituted C_{6-10} arylalkyl, a substituted C_{6-10} aryloxy, a substituted C_{6-10} arylalkoxy, a substituted 5-10 membered heteroaryl, a substituted C_{3-10} cycloalkyl, and a substituted 4-10 membered heterocyclyl,

wherein R^{13} , R^{14} , R^{15} , and R^{16} are, at each occurrence, independently selected from the group consisting of hydrogen, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, hydroxyl, C_{1-6} alkoxy, aryl, arylalkyl, C_{1-6} haloalkyl, C_{1-6} haloalkoxy, C_{1-6} hydroxyalkyl, cycloalkyl, heterocyclyl, and heteroaryl.

44. The compound of claim 39, wherein R^1 is an unsubstituted 4-10 membered heterocyclyl.

45. The compound of any one of claims 1-44, wherein the compound has one of the following structures:





46. The compound of any one of claims 1-45, wherein at least one of R^{2a}, R^{2b}, R^{2c}, and R^{2d} is alkoxy, alkoxyalkyl, cyano, halo, C(=O)OR⁵, OC(=O)R⁵, C(=O)NR⁵R⁶, N(R⁵)C(=O)R⁶, aryl, heteroaryl, heterocyclyl, or C(=O)N(R⁵)heterocyclyl, wherein R⁵ and R⁶ are, at each occurrence, independently hydrogen or C₁-C₆ alkyl.

47. The compound of any one of claims 1-46, wherein at least one of R^{2a}, R^{2b}, R^{2c}, and R^{2d} is methoxy.

48. The compound of any one of claims 1-46, wherein at least one of R^{2a} , R^{2b} , R^{2c} , and R^{2d} is methoxymethyl.

49. The compound of any one of claims 1-46, wherein at least one of R^{2a} , R^{2b} , R^{2c} , and R^{2d} is cyano.

50. The compound of any one of claims 1-46, wherein at least one of R^{2a} , R^{2b} , R^{2c} , and R^{2d} is $-C(=O)OH$.

51. The compound of any one of claims 1-46, wherein at least one of R^{2a} , R^{2b} , R^{2c} , and R^{2d} is $-C(=O)OCH_3$ or $-OC(=O)CH_3$.


52. The compound of any one of claims 1-46, wherein at least one of R^{2a} , R^{2b} , R^{2c} , and R^{2d} is $-C(=O)N(R')_2$ or $-NR'(C=O)alkyl$ wherein R' is, at each occurrence, independently a C_1 - C_6 alkyl.

53. The compound of claim 52, wherein each occurrence of R' is methyl.


54. The compound of any one of claims 1-46, wherein at least one of R^{2a} , R^{2b} , R^{2c} , and R^{2d} is $-C(=O)NR''heterocyclyl$ wherein R'' is hydrogen or C_1 - C_6 alkyl.

55. The compound of any one of claims 1-46, wherein at least one of R^{2a} , R^{2b} , R^{2c} , and R^{2d} is a substituted or unsubstituted phenyl.

56. The compound of any one of claims 1-46, wherein at least one of R^{2a} , R^{2b} , R^{2c} , and R^{2d} is a substituted or unsubstituted heterocyclyl.



60. The compound of any one of claims 1-59, wherein R³ has one of the following structures:



61. The compound of any one of claims 1-60, wherein R³ is -NH₂.

62. The compound of any one of claims 1-61, wherein n is 0.


63. The compound of any one of claims 1-61, wherein n is 1.


64. The compound of any one of claims 1-61, wherein n is 2.


65. The compound of any one of claims 1-61, wherein n is 3, 4, 5, or 6.

66. The compound of any one of claims 1-61, wherein n is 2 and R⁴ is a substituted or unsubstituted aryl.


67. The compound of claim 66, wherein R⁴ has one of the following structures:









68. The compound of claim 66, wherein R^4 has one of the following structures:




69. The compound of any one of claims 1-61, wherein n is 2 and R^4 is a substituted or unsubstituted heteroaryl.


70. The compound of claim 69, wherein R^4 has one of the following structures:







71. The compound of claim 69, wherein R^4 has one of the following structures:









72. The compound of claim 1, wherein R^4 has one of the following structures:




73. The compound of any one of claims 1-61, wherein n is 0 and R^4 has one of the following structures:






74. The compound of claim 73, wherein R⁴ has one of the following structures:



75. A compound having the following Structure (II):


$$(II)$$

or a stereoisomer, tautomer, or pharmaceutically acceptable salt thereof, wherein:

X is a direct bond, $-\text{C}(\text{R}^{2\text{e}})\text{R}^{2\text{f}}-$, or $-\text{C}(\text{R}^{2\text{e}})\text{R}^{2\text{f}}-\text{C}(\text{R}^{2\text{g}})\text{R}^{2\text{h}}-$;

Y is a direct bond or $-\text{[C(R}^{2i})\text{R}^{2j}\text{]}-$;

R¹ is a substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, substituted or unsubstituted cycloalkyl, or a substituted or unsubstituted heterocyclyl;

R^{2a} , R^{2b} , R^{2c} , R^{2d} , R^{2e} , R^{2f} , R^{2g} , R^{2h} , R^{2i} , and R^{2j} are each independently selected from the group consisting of hydrogen, halo, OR^5 , $C(=O)OR^5$, $OC(=O)R^5$,

hydroxyalkyl, alkoxy, alkoxyalkyl, haloalkoxy, cyano, aminylalkyl, carboxyalkyl, NR^5R^6 , $\text{C}(=\text{O})\text{NR}^5\text{R}^6$, $\text{N}(\text{R}^5)\text{C}(=\text{O})\text{R}^6$, $\text{NR}^5\text{C}(=\text{O})\text{NR}^6$, $\text{S}(\text{O})_t$, SR^5 , nitro, $\text{N}(\text{R}^5)\text{C}(\text{O})\text{OR}^6$, $\text{C}(=\text{NR}^5)\text{NR}^6\text{R}^7$, $\text{N}(\text{R}^5)\text{C}(=\text{NR}^6)\text{NR}^7\text{R}^8$, $\text{S}(\text{O})\text{R}^5$, $\text{S}(\text{O})\text{NR}^5\text{R}^6$, $\text{S}(\text{O})_2\text{R}^5$, $\text{N}(\text{R}^5)\text{S}(\text{O})_2\text{R}^6$, $\text{S}(\text{O})_2\text{NR}^5\text{R}^6$, aryl, heteroaryl, heterocyclyl, cycloalkyl, and oxo provided that at least one occurrence of R^{2a} , R^{2b} , R^{2c} , R^{2d} , R^{2e} , R^{2f} , R^{2g} , R^{2h} , R^{2i} , or R^{2j} is not hydrogen;

R^3 is $\text{NR}^{3a}\text{R}^{3b}$;

R^{3a} and R^{3b} are each independently hydrogen, alkyl, hydroxyalkyl, haloalkyl, alkoxyalkyl, $-\text{CH}_2\text{C}(\text{C}=\text{O})\text{OH}$, $-\text{CH}_2\text{C}(=\text{O})\text{Oalkyl}$, heterocyclyl, heteroaryl, heterocyclylalkyl, heteroarylalkyl, or cycloalkyl; or R^{3a} and R^{3b} , together with the nitrogen to which they are attached, form an optionally substituted 4-7 membered heteroaryl or an optionally substituted 4-7 membered heterocyclyl;

R^4 is a substituted or unsubstituted aryl, a substituted or unsubstituted heteroaryl, a substituted or unsubstituted cycloalkyl, or a substituted or unsubstituted heterocyclyl when n is 2, 3, 4, 5, or 6; or R^4 is a substituted or unsubstituted monocyclic heteroaryl, or a substituted or unsubstituted heterocyclyl when n is 0 or 1;


R^5 , R^6 , R^7 , and R^8 are, at each occurrence, independently hydrogen, alkyl, hydroxyalkyl, haloalkyl, alkoxyalkyl, carboxyalkyl, heterocyclyl, heteroaryl, or cycloalkyl;

n is an integer from 0-6; and

t is an integer from 1-3,


provided that:

a) when one of R^{2a} , R^{2b} , R^{2c} , R^{2d} , R^{2e} , R^{2f} , R^{2g} , R^{2h} , R^{2i} , and R^{2j} is OH, R^1 does not have the following structure:



b) when one of R^{2a} , R^{2b} , R^{2c} , R^{2d} , R^{2e} , R^{2f} , R^{2g} , and R^{2h} is $-\text{OH}$, n is an integer from 2-6; and

c) when one of R^{2a} , R^{2b} , R^{2c} , R^{2d} , R^{2e} , R^{2f} , R^{2g} , R^{2h} , R^{2i} , and R^{2j} is an unsubstituted phenyl, neither R^{3a} nor R^{3b} has the following structure:



76. The compound of claim 75, wherein R^1 is a substituted or unsubstituted aryl.

77. The compound of claim 75 or 76, wherein R^1 is a substituted or unsubstituted C_6 - C_{10} aryl.

78. The compound of any one of claims 75-77, wherein R^1 is a substituted or unsubstituted phenyl.

79. The compound of any one of claims 75-78, wherein R^1 is a substituted phenyl.

80. The compound of any one of claims 75-79, wherein R^1 is phenyl substituted with one or more of R^{1a} , R^{1b} , R^{1c} , R^{1d} , and R^{1e} wherein R^{1a} , R^{1b} , R^{1c} , R^{1d} , and R^{1e} are each independently selected from the group consisting of C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, halo, C_{1-6} haloalkyl, aminylalkyl, hydroxyalkyl, cyano, nitro, OR^9 , SR^9 , $C(O)R^9$, $C(O)NR^9R^{10}$, $C(O)OR^9$, $OC(O)R^9$, $OC(O)OR^9$, $OC(O)NR^9R^{10}$, NR^9R^{10} , $N(R^9)C(O)R^{10}$, $N(R^9)C(O)NR^{10}R^{11}$, $N(R^9)C(O)OR^{10}$, $C(=NR^9)NR^{10}R^{11}$, $C(=NOR^9)NR^{10}R^{11}$, $C(=NOC(O)R^9)NR^{10}R^{11}$, $C(=NR^9)N(R^{10})C(O)OR^{11}$, $N(R^9)C(=NR^{10})NR^{11}R^{12}$, $S(O)R^9$, $S(O)NR^9R^{10}$, $S(O)_2R^9$, $N(R^9)S(O)_2R^{10}$, $S(O)_2NR^9R^{10}$, oxo, substituted or unsubstituted C_{6-10} aryl, substituted or unsubstituted C_{6-10} arylalkyl, substituted or unsubstituted C_{6-10} aryloxy, substituted or unsubstituted C_{6-10} arylalkoxy, substituted or unsubstituted 5-10 membered heteroaryl, substituted or unsubstituted C_{3-10} cycloalkyl, and substituted or unsubstituted 4-10 membered heterocyclyl,

wherein R^9 , R^{10} , R^{11} , and R^{12} , are, at each occurrence, independently selected from the group consisting of hydrogen, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl,

hydroxyl, C₁₋₆ alkoxy, aryl, arylalkyl, C₁₋₆ haloalkyl, C₁₋₆ haloalkoxy, C₁₋₆ hydroxyalkyl, cycloalkyl, heterocyclyl, and heteroaryl.

81. The compound of claim 80, wherein R^{1a}, R^{1b}, R^{1c}, R^{1d}, or R^{1e} is optionally substituted with one or more substituents selected from the group consisting of halo, CN, OR¹³, SR¹³, C(O)R¹³, C(O)NR¹³R¹⁴, C(O)OR¹³, OC(O)R¹³, OC(O)NR¹³R¹⁴, NR¹³R¹⁴, NR¹³C(O)R¹⁴, NR¹³C(O)NR¹⁴R¹⁵, NR¹³C(O)OR¹⁴, C(=NR¹³)NR¹⁴R¹⁵, NR¹³C(=NR¹⁴)NR¹⁵R¹⁶, S(O)R¹³, S(O)NR¹³R¹⁴, S(O)₂R¹³, NR¹³S(O)₂R¹⁴, S(O)₂NR¹³R¹⁴ and oxo when R^{1a}, R^{1b}, R^{1c}, R^{1d}, or R^{1e} is a substituted C₆₋₁₀ aryl, a substituted C₆₋₁₀ arylalkyl, a substituted C₆₋₁₀ aryloxy, a substituted C₆₋₁₀ arylalkoxy, a substituted 5-10 membered heteroaryl, a substituted C₃₋₁₀ cycloalkyl, and a substituted 4-10 membered heterocyclyl,

wherein R¹³, R¹⁴, R¹⁵, and R¹⁶ are, at each occurrence, independently selected from the group consisting of hydrogen, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, hydroxyl, C₁₋₆ alkoxy, aryl, arylalkyl, C₁₋₆ haloalkyl, C₁₋₆ haloalkoxy, C₁₋₆ hydroxyalkyl, cycloalkyl, heterocyclyl, and heteroaryl.


82. The compound of any one of claims 75-81, wherein R¹ is phenyl substituted with at least one substituent selected from the group consisting of OR⁹, C(=NR⁹)NR¹⁰R¹¹, C(=NOR⁹)NR¹⁰R¹¹, C(=NOC(O)R⁹)NR¹⁰R¹¹, and C(=NR⁹)NR¹⁰C(O)OR¹¹.


83. The compound of any one of claims 75-82, wherein R¹ is substituted with at least one substituent selected from the group consisting of C(=NR⁹)NR¹⁰R¹¹, C(=NOR⁹)NR¹⁰R¹¹, C(=NOC(O)R⁹)NR¹⁰R¹¹, and C(=NR⁹)NR¹⁰C(O)OR¹¹.

84. The compound of claim 83, wherein R¹ is substituted with at least one C(=NR⁹)NR¹⁰R¹¹.


85. The compound of claim 83 or 84, wherein R^1 is substituted with at least one $-C(=NH)NH_2$.

86. The compound of any one of claims 81-85, wherein R^1 has one of the following structures:






87. The compound of any one of claims 75-86, wherein R^1 has one of the following structures:



88. The compound of claim 81, wherein R^1 has one of the following structures:



89. The compound of claim 88, wherein R^9 is C_{1-6} alkyl or C_{1-6} haloalkyl.
90. The compound of claim 89, wherein R^9 is methyl.
91. The compound of claim 89, wherein R^9 is trifluoromethyl.
92. The compound of any one of claims 75-78, wherein R^1 is an unsubstituted phenyl.
93. The compound of claim 75, wherein R^1 is a substituted or unsubstituted heteroaryl.
94. The compound of claim 75 or 93, wherein R^1 is a substituted or unsubstituted 5-10 membered heteroaryl.
95. The compound of claim 75 or 93-94, wherein R^1 is a substituted or unsubstituted pyridinyl, pyrrolopyridinyl, or benzoimidazolyl.

96. The compound of any one of claims 75 or 93-95, wherein R^1 is a substituted or unsubstituted pyridin-3-yl, 1H-pyrrolo[2,3-b]pyridine-5-yl, 1H-pyrrolo[3,2-c]pyridine-2-yl, or 1H-benzo[d]imidazol-6-yl.

97. The compound of any one of claims 75 or 93-96, wherein R^1 is pyridin-3-yl, 1H-pyrrolo[2,3-b]pyridine-5-yl, or 1H-benzo[d]imidazol-6-yl substituted with one or more of R^{1a} , R^{1b} , R^{1c} , R^{1d} , and R^{1e} wherein R^{1a} , R^{1b} , R^{1c} , R^{1d} , and R^{1e} are each independently selected from the group consisting of C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, halo, C_{1-6} haloalkyl, aminylalkyl, hydroxyalkyl, cyano, nitro, OR^9 , SR^9 , $C(O)R^9$, $C(O)NR^9R^{10}$, $C(O)OR^9$, $OC(O)R^9$, $OC(O)OR^9$, $OC(O)NR^9R^{10}$, NR^9R^{10} , $N(R^9)C(O)R^{10}$, $N(R^9)C(O)NR^{10}R^{11}$, $N(R^9)C(O)OR^{10}$, $C(=NR^9)NR^{10}R^{11}$, $C(=NOR^9)NR^{10}R^{11}$, $C(=NOC(O)R^9)NR^{10}R^{11}$, $C(=NR^9)N(R^{10})C(O)OR^{11}$, $N(R^9)C(=NR^{10})NR^{11}R^{12}$, $S(O)R^9$, $S(O)NR^9R^{10}$, $S(O)_2R^9$, $N(R^9)S(O)_2R^{10}$, $S(O)_2NR^9R^{10}$, oxo, substituted or unsubstituted C_{6-10} aryl, substituted or unsubstituted C_{6-10} arylalkyl, substituted or unsubstituted C_{6-10} aryloxy, substituted or unsubstituted C_{6-10} arylalkoxy, substituted or unsubstituted 5-10 membered heteroaryl, substituted or unsubstituted C_{3-10} cycloalkyl, and substituted or unsubstituted 4-10 membered heterocyclyl,


wherein R^9 , R^{10} , R^{11} , and R^{12} , are, at each occurrence, independently selected from the group consisting of hydrogen, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, hydroxyl, C_{1-6} alkoxy, aryl, arylalkyl, C_{1-6} haloalkyl, C_{1-6} haloalkoxy, C_{1-6} hydroxyalkyl, cycloalkyl, heterocyclyl, and heteroaryl.


98. The compound of claim 97, wherein R^{1a} , R^{1b} , R^{1c} , R^{1d} , or R^{1e} is optionally substituted with one or more substituents selected from the group consisting of halo, CN, OR^{13} , SR^{13} , $C(O)R^{13}$, $C(O)NR^{13}R^{14}$, $C(O)OR^{13}$, $OC(O)R^{13}$, $OC(O)NR^{13}R^{14}$, $NR^{13}R^{14}$, $NR^{13}C(O)R^{14}$, $NR^{13}C(O)NR^{14}R^{15}$, $NR^{13}C(O)OR^{14}$, $C(=NR^{13})NR^{14}R^{15}$, $NR^{13}C(=NR^{14})NR^{15}R^{16}$, $S(O)R^{13}$, $S(O)NR^{13}R^{14}$, $S(O)_2R^{13}$, $NR^{13}S(O)_2R^{14}$, $S(O)_2NR^{13}R^{14}$ and oxo when R^{1a} , R^{1b} , R^{1c} , R^{1d} , or R^{1e} is a substituted C_{6-10} aryl, a substituted C_{6-10} arylalkyl, a substituted C_{6-10} aryloxy, a substituted C_{6-10} arylalkoxy, a substituted 5-10


membered heteroaryl, a substituted C₃₋₁₀ cycloalkyl, and a substituted 4-10 membered heterocyclyl,


wherein R¹³, R¹⁴, R¹⁵, and R¹⁶ are, at each occurrence, independently selected from the group consisting of hydrogen, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, hydroxyl, C₁₋₆ alkoxy, aryl, arylalkyl, C₁₋₆ haloalkyl, C₁₋₆ haloalkoxy, C₁₋₆ hydroxyalkyl, cycloalkyl, heterocyclyl, and heteroaryl.

99. The compound of any one of claims 75 or 93-97, wherein R¹ has one of the following structures:









100. The compound of claim 99, wherein R^{1a} or R^{1b} is independently C_{1-6} alkyl, amino, $C(O)R^9$, $C(O)NR^9R^{10}$, $N(R^9)C(O)NR^{10}R^{11}$, $N(R^9)C(O)OR^{10}$, or $OC(O)NR^9R^{10}$, or halo.


101. The compound of claim 100, wherein R^{1a} or R^{1b} is methyl or ethyl.


102. The compound of claim 100, wherein R^{1a} or R^{1b} is F, Cl, or Br.

103. The compound of claim 99, wherein each R^{1a} or R^{1b} attached to nitrogen is C_{1-6} alkyl, $C(O)R^9$, $C(O)NR^9R^{10}$, $N(R^9)C(O)NR^{10}R^{11}$, $N(R^9)C(O)OR^{10}$, or $OC(O)NR^9R^{10}$.


104. The compound of claim 103, wherein R^{1a} or R^{1b} is methyl or ethyl.

105. The compound of any one of claims 75 or 93-94, wherein R^1 has one of the following structures:





106. The compound of any one of claims 75 or 93-94, wherein R^1 has one of the following structures:



107. The compound of claim 75, wherein R^1 is a substituted or unsubstituted cycloalkyl.

108. The compound of claim 75 or 107, wherein R^1 is a substituted or unsubstituted C_3 - C_6 cycloalkyl.

109. The compound of any one of claims 75 or 107-108, wherein R^1 is a substituted C_3 - C_6 cycloalkyl.

110. The compound of claim 109, wherein R^1 is a C_3 - C_6 cycloalkyl substituted with one or more of R^{1a} , R^{1b} , R^{1c} , R^{1d} , and R^{1e} wherein R^{1a} , R^{1b} , R^{1c} , R^{1d} , and R^{1e} are each independently selected from the group consisting of C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} haloalkyl, aminylalkyl, hydroxyalkyl, nitro, OR^9 , SR^9 , $C(O)R^9$, $C(O)NR^9R^{10}$, $C(O)OR^9$, $OC(O)R^9$, $OC(O)OR^9$, $OC(O)NR^9R^{10}$, NR^9R^{10} , $N(R^9)C(O)R^{10}$, $N(R^9)C(O)NR^{10}R^{11}$, $N(R^9)C(O)OR^{10}$, $C(=NR^9)NR^{10}R^{11}$, $C(=NOR^9)NR^{10}R^{11}$, $C(=NOC(O)R^9)NR^{10}R^{11}$, $C(=NR^9)N(R^{10})C(O)OR^{11}$, $N(R^9)C(=NR^{10})NR^{11}R^{12}$, $S(O)R^9$, $S(O)NR^9R^{10}$, $S(O)_2R^9$, $N(R^9)S(O)_2R^{10}$, $S(O)_2NR^9R^{10}$, oxo, substituted or unsubstituted C_{6-10} aryl, substituted or unsubstituted C_{6-10} arylalkyl, substituted or unsubstituted C_{6-10} aryloxy, substituted or unsubstituted C_{6-10} arylalkoxy, substituted or unsubstituted 5-10 membered heteroaryl, substituted or unsubstituted C_{3-10} cycloalkyl, and substituted or unsubstituted 4-10 membered heterocyclyl,

wherein R^9 , R^{10} , R^{11} , and R^{12} , are, at each occurrence, independently selected from the group consisting of hydrogen, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, hydroxyl, C_{1-6} alkoxy, aryl, arylalkyl, C_{1-6} haloalkyl, C_{1-6} haloalkoxy, C_{1-6} hydroxyalkyl, cycloalkyl, heterocyclyl, and heteroaryl.

111. The compound of claim 110, wherein R^{1a} , R^{1b} , R^{1c} , R^{1d} , or R^{1e} is optionally substituted with one or more substituents selected from the group consisting of halo, CN, OR^{13} , SR^{13} , $C(O)R^{13}$, $C(O)NR^{13}R^{14}$, $C(O)OR^{13}$, $OC(O)R^{13}$, $OC(O)NR^{13}R^{14}$, $NR^{13}R^{14}$, $NR^{13}C(O)R^{14}$, $NR^{13}C(O)NR^{14}R^{15}$, $NR^{13}C(O)OR^{14}$, $C(=NR^{13})NR^{14}R^{15}$, $NR^{13}C(=NR^{14})NR^{15}R^{16}$, $S(O)R^{13}$, $S(O)NR^{13}R^{14}$, $S(O)_2R^{13}$, $NR^{13}S(O)_2R^{14}$, $S(O)_2NR^{13}R^{14}$ and oxo when R^{1a} , R^{1b} , R^{1c} , R^{1d} , or R^{1e} is a substituted C_{6-10} aryl, a substituted C_{6-10} arylalkyl, a substituted C_{6-10} aryloxy, a substituted C_{6-10} arylalkoxy, a substituted 5-10

membered heteroaryl, a substituted C₃₋₁₀ cycloalkyl, and a substituted 4-10 membered heterocyclyl,

wherein R¹³, R¹⁴, R¹⁵, and R¹⁶ are, at each occurrence, independently selected from the group consisting of hydrogen, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, hydroxyl, C₁₋₆ alkoxy, aryl, arylalkyl, C₁₋₆ haloalkyl, C₁₋₆ haloalkoxy, C₁₋₆ hydroxyalkyl, cycloalkyl, heterocyclyl, and heteroaryl.

112. The compound of claim 108, wherein R¹ is an unsubstituted a C₃-C₆ cycloalkyl.

113. The compound of claim 75, wherein R¹ is a substituted or unsubstituted heterocyclyl.

114. The compound of claim 75 or 113, wherein R¹ is a substituted or unsubstituted 4-10 membered heterocyclyl.

115. The compound of any one of claims 75 or 73-84, wherein R¹ is a substituted 4-10 membered heterocyclyl.

116. The compound of claim 115, wherein R¹ is a 4-10 membered heterocyclyl substituted with one or more of R^{1a}, R^{1b}, R^{1c}, R^{1d}, and R^{1e} wherein R^{1a}, R^{1b}, R^{1c}, R^{1d}, and R^{1e} are each independently selected from the group consisting of C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, halo, C₁₋₆ haloalkyl, aminylalkyl, hydroxyalkyl, cyano, nitro, OR⁹, SR⁹, C(O)R⁹, C(O)NR⁹R¹⁰, C(O)OR⁹, OC(O)R⁹, OC(O)OR⁹, OC(O)NR⁹R¹⁰, NR⁹R¹⁰, N(R⁹)C(O)R¹⁰, N(R⁹)C(O)NR¹⁰R¹¹, N(R⁹)C(O)OR¹⁰, C(=NR⁹)NR¹⁰R¹¹, C(=NOR⁹)NR¹⁰R¹¹, C(=NOC(O)R⁹)NR¹⁰R¹¹, C(=NR⁹)N(R¹⁰)C(O)OR¹¹, N(R⁹)C(=NR¹⁰)NR¹¹R¹², S(O)R⁹, S(O)NR⁹R¹⁰, S(O)₂R⁹, N(R⁹)S(O)₂R¹⁰, S(O)₂NR⁹R¹⁰, oxo, substituted or unsubstituted C₆₋₁₀ aryl, substituted or unsubstituted C₆₋₁₀ arylalkyl, substituted or unsubstituted C₆₋₁₀ aryloxy, substituted or unsubstituted C₆₋₁₀ arylalkoxy,

substituted or unsubstituted 5-10 membered heteroaryl, substituted or unsubstituted C₃₋₁₀ cycloalkyl, and substituted or unsubstituted 4-10 membered heterocyclyl,









wherein R⁹, R¹⁰, R¹¹, and R¹², are, at each occurrence, independently selected from the group consisting of hydrogen, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, hydroxyl, C₁₋₆ alkoxy, aryl, arylalkyl, C₁₋₆ haloalkyl, C₁₋₆ haloalkoxy, C₁₋₆ hydroxyalkyl, cycloalkyl, heterocyclyl, and heteroaryl.




117. The compound of claim 116, wherein R^{1a}, R^{1b}, R^{1c}, R^{1d}, or R^{1e} is optionally substituted with one or more substituents selected from the group consisting of OR¹³, SR¹³, C(O)R¹³, C(O)NR¹³R¹⁴, C(O)OR¹³, OC(O)R¹³, OC(O)NR¹³R¹⁴, NR¹³R¹⁴, NR¹³C(O)R¹⁴, NR¹³C(O)NR¹⁴R¹⁵, NR¹³C(O)OR¹⁴, C(=NR¹³)NR¹⁴R¹⁵, NR¹³C(=NR¹⁴)NR¹⁵R¹⁶, S(O)R¹³, S(O)NR¹³R¹⁴, S(O)₂R¹³, NR¹³S(O)₂R¹⁴, S(O)₂NR¹³R¹⁴ and oxo when R^{1a}, R^{1b}, R^{1c}, R^{1d}, or R^{1e} is a substituted C₆₋₁₀ aryl, a substituted C₆₋₁₀ arylalkyl, a substituted C₆₋₁₀ aryloxy, a substituted C₆₋₁₀ arylalkoxy, a substituted 5-10 membered heteroaryl, a substituted C₃₋₁₀ cycloalkyl, and a substituted 4-10 membered heterocyclyl,

wherein R¹³, R¹⁴, R¹⁵, and R¹⁶ are, at each occurrence, independently selected from the group consisting of hydrogen, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, hydroxyl, C₁₋₆ alkoxy, aryl, arylalkyl, C₁₋₆ haloalkyl, C₁₋₆ haloalkoxy, C₁₋₆ hydroxyalkyl, cycloalkyl, heterocyclyl, and heteroaryl.


118. The compound of claim 113, wherein R¹ is an unsubstituted 4-10 membered heterocyclyl.

119. The compound of any one of claims 75-118, wherein the compound has one of the following structures:





or



120. The compound of any one of claims 75-119, wherein at least one of R^{2a} , R^{2b} , R^{2c} , and R^{2d} is alkoxy, alkoxyalkyl, cyano, $C(=O)OR^5$, $OC(=O)R^5$, $C(=O)NR^5R^6$, $N(R^5)C(=O)R^6$, aryl, heteroaryl, heterocyclyl, or $C(=O)N(R^5)$ heterocyclyl, wherein R^5 and R^6 are, at each occurrence, independently hydrogen or C_1 - C_6 alkyl.

121. The compound of any one of claims 75-120, wherein at least one of R^{2a} , R^{2b} , R^{2c} , and R^{2d} is methoxy.

122. The compound of any one of claims 75-120, wherein at least one of R^{2a} , R^{2b} , R^{2c} , and R^{2d} is methoxymethyl.

123. The compound of any one of claims 75-120, wherein at least one of R^{2a} , R^{2b} , R^{2c} , and R^{2d} is cyano.

124. The compound of any one of claims 75-120, wherein at least one of R^{2a} , R^{2b} , R^{2c} , and R^{2d} is $-C(=O)OH$.

125. The compound of any one of claims 75-120, wherein at least one of R^{2a} , R^{2b} , R^{2c} , and R^{2d} is $-C(=O)OCH_3$ or $-OC(=O)CH_3$.

126. The compound of any one of claims 75-120, at least one of R^{2a} , R^{2b} , R^{2c} , and R^{2d} is $-C(=O)N(R')_2$ or $-NR'(C=O)alkyl$ wherein R' is, at each occurrence, independently a C_1 - C_6 alkyl.

127. The compound of claim 126, wherein each occurrence of R' is methyl.


128. The compound of any one of claims 75-120, wherein at least one of R^{2a} , R^{2b} , R^{2c} , and R^{2d} is $-C(=O)NR''heterocyclyl$ wherein R'' is hydrogen or C_1 - C_6 alkyl.

129. The compound of any one of claims 75-120, wherein at least one of R^{2a} , R^{2b} , R^{2c} , and R^{2d} is a substituted or unsubstituted phenyl.


130. The compound of any one of claims 75-120, wherein at least one of R^{2a} , R^{2b} , R^{2c} , and R^{2d} is a substituted or unsubstituted heterocyclyl.


131. The compound of any one of claims 75-120, wherein at least one of R^{2a} , R^{2b} , R^{2c} , and R^{2d} is a substituted or unsubstituted heteroaryl.

132. The compound of any one of claims 75-131, wherein at least one of R^{2a} , R^{2b} , R^{2c} , and R^{2d} has one of the following structures:




133. The compound of any one of claims 75-132, wherein at least one of R^{2a} , R^{2b} , R^{2c} , and R^{2d} has one of the following structures:





134. The compound of any one of claims 75-133, wherein R³ has one of the following structures:





135. The compound of any one of claims 75-134, wherein R³ is -NH₂.

136. The compound of any one of claims 75-135, wherein n is 0.


137. The compound of any one of claims 75-135, wherein n is 1.

138. The compound of any one of claims 75-135, wherein n is 2.







142. The compound of claim 140, wherein R⁴ has one of the following structures:




143. The compound of any one of claims 75-135, wherein n is 2 and R⁴ is a substituted or unsubstituted heteroaryl.


144. The compound of claim 143, wherein R⁴ has one of the following structures:







145. The compound of claim 143, wherein R⁴ has one of the following structures:









146. The compound of claim 139, wherein R^4 has one of the following structures:



147. The compound of any one of claims 75-135, wherein n is 0 and R^4 has one of the following structures:




148. The compound of claim 147, wherein R⁴ has one of the following structures:



149. A compound having a structure in Table 1 or a stereoisomer, tautomer, or pharmaceutically acceptable salt thereof.

150. A compound having the following Structure (III):



(III)

or a stereoisomer, tautomer, or pharmaceutically acceptable salt thereof, wherein:

X is a direct bond, $-\text{CR}^{2e}\text{R}^{2f}-$, or $-\text{CR}^{2e}\text{R}^{2f}-\text{CR}^{2g}\text{R}^{2h}-$;

Y is a direct bond or $-\text{CR}^{2i}\text{R}^{2j}-$;

Z is O or S;

R^{17} is a substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, substituted or unsubstituted cycloalkyl, or a substituted or unsubstituted heterocyclyl;

R^{18a} , R^{18b} , R^{18c} , R^{18d} , R^{18e} , R^{18f} , R^{18g} , R^{18h} , R^{18i} , and R^{18j} are each independently selected from the group consisting of hydrogen, halo, $-\text{OR}^{21}$, $\text{C}(=\text{O})\text{OR}^{21}$, $\text{OC}(=\text{O})\text{R}^{21}$, hydroxyalkyl, alkoxy, alkoxyalkyl, haloalkoxy, cyano, aminylalkyl, carboxyalkyl, $\text{NR}^{21}\text{R}^{22}$, $\text{C}(=\text{O})\text{NR}^{21}\text{R}^{22}$, $\text{N}(\text{R}^{21})\text{C}(=\text{O})\text{R}^{22}$, $\text{NR}^{21}\text{C}(=\text{O})\text{NR}^{22}$, $\text{S}(\text{O})_t$, SR^{21} , nitro, $\text{N}(\text{R}^{21})\text{C}(\text{O})\text{OR}^{22}$, $\text{C}(=\text{NR}^{21})\text{NR}^{22}\text{R}^{23}$, $\text{N}(\text{R}^{21})\text{C}(=\text{NR}^{22})\text{NR}^{23}\text{R}^{24}$, $\text{S}(\text{O})\text{R}^{21}$, $\text{S}(\text{O})\text{NR}^{21}\text{R}^{22}$, $\text{S}(\text{O})_2\text{R}^{21}$, $\text{N}(\text{R}^{21})\text{S}(\text{O})_2\text{R}^{22}$, $\text{S}(\text{O})_2\text{NR}^{21}\text{R}^{22}$, aryl, heteroaryl, heterocyclyl, cycloalkyl, and oxo provided that at least one of R^{18a} , R^{18b} , R^{18c} , R^{18d} , R^{18e} , R^{18f} , R^{18g} , R^{18h} , R^{18i} , and R^{18j} is not hydrogen;

R^{19} is $\text{NR}^{19a}\text{R}^{19b}$;

R^{19a} and R^{19b} are each independently hydrogen, alkyl, hydroxyalkyl, haloalkyl, alkoxyalkyl, heterocyclyl, heteroaryl, heterocyclylalkyl, heteroarylalkyl, cycloalkyl, $(\text{CH}_2)_n\text{C}(=\text{O})\text{OR}^5$, or $(\text{CH}_2)_n\text{P}(=\text{O})(\text{OR}^5)_2$; or R^{19a} and R^{19b} , together with the nitrogen to which they are attached, form an optionally substituted 4-7 membered heteroaryl or an optionally substituted 4-7 membered heterocyclyl;


R^{20} is a substituted or unsubstituted aryl, a substituted or unsubstituted heteroaryl, a substituted or unsubstituted cycloalkyl, or a substituted or unsubstituted heterocyclyl;

R^{21} , R^{22} , R^{23} , and R^{24} are, at each occurrence, independently hydrogen, alkyl, hydroxyalkyl, haloalkyl, alkoxyalkyl, carboxyalkyl, heterocyclyl, heteroaryl, or cycloalkyl;

m is an integer from 0-6; and


t is an integer from 1-3.


151. The compound of claim 150, wherein R^{17} has one of the following structures:




152. The compound of claim 150, wherein R^{17} is an unsubstituted phenyl.


153. The compound of claim 150, wherein R^{17} has one of the following structures:








154. The compound of claim 150, wherein R¹⁷ has one of the following structures:




155. The compound of any one of claims 150-154, wherein at least one of R^{18a} or R^{18b} is alkoxy, alkoxyalkyl, cyano, C(=O)OR²¹, OC(=O)R²¹, C(=O)NR²¹R²²,

$N(R^{21})C(=O)R^{22}$, aryl, heteroaryl, heterocyclyl, or $C(=O)N(R^{21})$ heterocyclyl, wherein R^{21} and R^{22} are, at each occurrence, independently hydrogen or C_1 - C_6 alkyl.


156. The compound of any one of claims 150-154, wherein at least one of R^{18a} or R^{18b} has one of the following structures:




157. The compound of any one of claims 150-156, wherein R^{19} has one of the following structures:





158. The compound of any one of claims 150-157, wherein R^{20} has one of the following structures:







159. The compound of any one of claims 150-157, wherein R^{20} has one of the following structures:







160. The compound of any one of claims 150-157, wherein R^{20} has one of the following structures:







161. A compound having the following Structure (III):



or a stereoisomer, tautomer, or pharmaceutically acceptable salt thereof, wherein:

X is a direct bond or -CR^{26c}R^{26d}-;

R²⁵ is a substituted or unsubstituted aryl, substituted or unsubstituted heteroaryl, substituted or unsubstituted cycloalkyl, or a substituted or unsubstituted heterocyclyl;

R^{26a}, R^{26b}, R^{26c}, and R^{26d} are each independently selected from the group consisting of hydrogen, halo, -OR²⁹, C(=O)OR²⁹, OC(=O)R²⁹, hydroxyalkyl, alkoxy, alkoxyalkyl, haloalkoxy, cyano, aminylalkyl, carboxyalkyl, NR²⁹R³⁰, C(=O)NR²⁹R³⁰, N(R²⁹)C(=O)R³⁰, NR²⁹C(=O)NR³⁰, S(O)_t, SR²⁹, nitro, N(R²⁹)C(O)OR³⁰, C(=NR²⁹)NR³⁰R³¹, N(R²⁹)C(=NR³⁰)NR³¹R³², S(O)R²⁹, S(O)NR²⁹R³⁰, S(O)₂R³⁰,

$N(R^{29})S(O)_2R^{30}$, $S(O)_2NR^{29}R^{30}$, aryl, heteroaryl, heterocyclyl, cycloalkyl, and oxo, provided that at least one of R^{26a} , R^{26b} , R^{26c} , and R^{26d} is not hydrogen;

R^{27} is $NR^{27a}R^{27b}$,

R^{27a} and R^{27b} are each independently hydrogen, alkyl, hydroxyalkyl, haloalkyl, alkoxyalkyl, heterocyclyl, heteroaryl, heterocyclylalkyl, heteroarylalkyl, cycloalkyl, $(CH_2)_nC(=O)OR^{29}$, or $(CH_2)_nP(=O)(OR^{29})_2$; or R^{27a} and R^{27b} , together with the nitrogen to which they are attached, form an optionally substituted 4-7 membered heteroaryl or an optionally substituted 4-7 membered heterocyclyl;


R^{28} is a substituted or unsubstituted aryl, a substituted or unsubstituted heteroaryl, a substituted or unsubstituted cycloalkyl, or a substituted or unsubstituted heterocyclyl;


R^{29} , R^{30} , R^{31} , and R^{32} are, at each occurrence, independently hydrogen, alkyl, hydroxyalkyl, haloalkyl, alkoxyalkyl, carboxyalkyl, heterocyclyl, heteroaryl, or cycloalkyl;

p is an integer from 0-6; and

t is an integer from 1-3.


162. The compound of claim 161, wherein R^{25} has one of the following structures:







163. The compound of claim 161, wherein R^{25} is an unsubstituted phenyl.


164. The compound of claim 161, wherein R^{25} has one of the following structures:








165. The compound of claim 161, wherein R²⁵ has one of the following structures:




166. The compound of any one of claims 161-165, wherein at least one of R^{26a} and R^{26b} is alkoxy, alkoxyalkyl, cyano, $C(=O)OR^{29}$, $OC(=O)R^{29}$, $C(=O)NR^{29}R^{30}$, $N(R^{29})C(=O)R^{30}$, aryl, heteroaryl, heterocyclyl, or $C(=O)N(R^{29})$ heterocyclyl, wherein R^{29} and R^{30} are, at each occurrence, independently hydrogen or C_1 - C_6 alkyl.


167. The compound of any one of claims 161-165, wherein at least one of R^{26a} and R^{26b} has one of the following structures:





168. The compound of any one of claims 161-167, wherein R²⁷ has one of the following structures:




169. The compound of any one of claims 161-168, wherein R²⁸ has one of the following structures:










170. The compound of any one of claims 161-168, wherein R²⁸ has one of the following structures:






171. The compound of any one of claims 161-168, wherein R^{28} has one of the following structures:







172. A pharmaceutical composition comprising a compound or pharmaceutically acceptable salt thereof of any one of claims 1 to 171 and a pharmaceutically acceptable carrier or excipient.

173. A method for treating a MASP-2-associated disease or disorder in a subject in need thereof, the method comprising administering to the subject an effective amount of a compound of any one of claims 1-171 or a pharmaceutical composition according to claim 172.

174. The method of claim 173, wherein the compound is administered in an amount sufficient to inhibit MASP-2 dependent complement activation in the subject.

175. The method of claim 173, wherein the subject has been diagnosed as being in need of treatment for a lectin complement-associated disease or disorder.

176. The method of claim 173, wherein disease or disorder is a thrombotic microangiopathy (TMA), a renal condition, an inflammatory reaction resulting from tissue or organ transplantation, an ischemia reperfusion injury, a complication associated with diabetes, a cardiovascular disease or disorder, an inflammatory gastrointestinal disorder, a pulmonary disorder, an ophthalmic disease or disorder, disseminated intravascular coagulation, graft-versus-host disease, veno-occlusive disease, diffuse alveolar hemorrhage, idiopathic pneumonia syndrome, capillary leak syndrome, engraftment syndrome, fluid overload, or combinations thereof.

177. The method of claim 173, wherein the disease or disorder is a thrombotic microangiopathy (TMA) thrombotic thrombocytopenic purpura (TTP), refractory TTP, Upshaw-Schulman Syndrome (USS), hemolytic uremic syndrome (HUS), atypical hemolytic syndrome (aHUS), non-Factor H-dependent atypical hemolytic syndrome, aHUS secondary to an infection, plasma therapy-resistant aHUS, a TMA secondary to cancer, a TMA secondary to chemotherapy, a TMA secondary to transplantation, a TMA associated with hematopoietic stem cell transplant, or combinations thereof.

178. The method of claim 173, wherein the disease or disorder is graft-versus-host disease.

179. The method of claim 173, wherein the disease or disorder is diffuse alveolar hemorrhage (DAH).

180. The method of claim 173, wherein the disease or disorder is veno-occlusive disease (VOD).

181. The method of claim 173, wherein the disease or disorder is a renal condition.

182. The method of claim 173, wherein the renal condition is mesangioproliferative glomerulonephritis, membranous glomerulonephritis, membranoproliferative glomerulonephritis (mesangiocapillary glomerulonephritis), acute post infectious glomerulonephritis (poststreptococcal glomerulonephritis), C3 glomerulopathy, cryoglobulinemic glomerulonephritis, pauci-immune necrotizing crescentic glomerulonephritis, lupus nephritis, Henoch-Schonlein purpura nephritis, IgA nephropathy, or combinations thereof.

183. The method of claim 173, wherein the disease or disorder is renal fibrosis, proteinuria, or combinations thereof.

184. The method of claim 173, wherein the disease or disorder is an inflammatory reaction resulting from tissue or solid organ transplantation.

185. The method of claim 173, wherein the disease or disorder is an ischemia reperfusion injury (I/R).

186. The method of claim 173, wherein the disease or disorder is a complication associated with non-obese diabetes, complications associated with Type-1 diabetes, Type-2 (adult onset) diabetes, or combinations thereof.

187. The method of claim 173, wherein the disease or disorder is a cardiovascular disease or disorder.

188. The method of claim 173, wherein the disease or disorder is an inflammatory gastrointestinal disorder.

189. The method of claim 173, wherein the disease or disorder is a pulmonary disorder.

190. The method of claim 173, wherein the disease or disorder is an extracorporeal exposure-triggered inflammatory reaction.

191. The method of claim 190, wherein the method further comprises treating a subject undergoing an extracorporeal circulation procedure.

192. The method of claim 173, wherein the disease or disorder is an inflammatory arthritis, non-inflammatory arthritis, a musculoskeletal disorder, or combinations thereof.

193. The method of claim 173, wherein the disease or disorder is a skin disorder.

194. The method of claim 173, wherein the disease or disorder is peripheral nervous system (PNS) disorder or injury, a central nervous system (CNS) disorder or injury, or combinations thereof.

195. The method of claim 173, wherein the disease or disorder is sepsis or a condition resulting from sepsis.

196. The method of claim 173, wherein the disease or disorder is a urogenital disorder.

197. The method of claim 173, wherein the disease or disorder is an inflammatory reaction in a subject being treated with chemotherapeutics, radiation therapy, or combinations thereof.

198. The method of claim 173, wherein the disease or disorder is an angiogenesis-dependent cancer.

199. The method of claim 173, wherein the disease or disorder is an angiogenesis-dependent benign tumor.

200. The method of claim 173, wherein the disease or disorder is an endocrine disorder.

201. The method of claim 173, wherein the disease or disorder is an ophthalmic disease or disorder.

202. The method of claim 173, wherein the disease or disorder is an ocular angiogenic disease or condition.

203. The method of claim 173, wherein the disease or disorder is disseminated intravascular coagulation (DIC), a complement mediated coagulation disorder, or combinations thereof.

204. The method of claim 173, wherein the disease or disorder is acute radiation syndrome, dense deposit disease, Degos Disease, Catastrophic Antiphospholipid Syndrome (CAPS), Behcet's disease, cryoglobulinemia; paroxysmal nocturnal hemoglobinuria ("PNH"), cold agglutinin disease, or combinations thereof.

205. The method of claim 173, wherein the disease or disorder is atypical hemolytic uremic syndrome (aHUS).

206. The method of claim 173, wherein the disease or disorder is hematopoietic stem cell transplant-associated TMA.

207. The method of claim 173, wherein the disease or disorder is immunoglobulin A nephropathy (IgAN)

208. The method of claim 173, wherein the disease or disorder is lupus nephritis (LN).

INTERNATIONAL SEARCH REPORT

International application No
PCT/US2020/063391

A. CLASSIFICATION OF SUBJECT MATTER

INV. C07D207/16 C07D403/04 C07D413/04 C07D471/04
ADD.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EPO-Internal, CHEM ABS Data, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	W0 2019/211585 A1 (UNIV LEEDS INNOVATIONS LTD [GB]) 7 November 2019 (2019-11-07) page 83; compounds M05062,M05071 -----	1-92, 96-128, 130,131, 134-171, 173-208
X	W0 2019/186164 A1 (UNIV LEEDS INNOVATIONS LTD [GB]) 3 October 2019 (2019-10-03) page 133; compound M05125 ----- -/-	75, 93-95, 119,129, 132-135, 138,140, 141,172



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

9 April 2021

Date of mailing of the international search report

15/04/2021

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040,
Fax: (+31-70) 340-3016

Authorized officer

Bedel, Christian

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US2020/063391

Box No. I Nucleotide and/or amino acid sequence(s) (Continuation of item 1.c of the first sheet)

1. With regard to any nucleotide and/or amino acid sequence disclosed in the international application, the international search was carried out on the basis of a sequence listing:
 - a. ☒ forming part of the international application as filed:
 - ☒ in the form of an Annex C/ST.25 text file.
 - ☐ on paper or in the form of an image file.
 - b. ☐ furnished together with the international application under PCT Rule 13~~ter~~.1(a) for the purposes of international search only in the form of an Annex C/ST.25 text file.
 - c. ☐ furnished subsequent to the international filing date for the purposes of international search only:
 - ☐ in the form of an Annex C/ST.25 text file (Rule 13~~ter~~.1(a)).
 - ☐ on paper or in the form of an image file (Rule 13~~ter~~.1(b) and Administrative Instructions, Section 713).
2. ☐ In addition, in the case that more than one version or copy of a sequence listing has been filed or furnished, the required statements that the information in the subsequent or additional copies is identical to that forming part of the application as filed or does not go beyond the application as filed, as appropriate, were furnished.
3. Additional comments:

INTERNATIONAL SEARCH REPORT

International application No

PCT/US2020/063391

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 9 011 860 B2 (DUDLER THOMAS [US]; GOMBOTZ WAYNE R [US] ET AL.) 21 April 2015 (2015-04-21) cited in the application the whole document -----	1-208
A	WO 2006/101860 A1 (JANSSEN PHARMACEUTICA NV [BE]; SUBASINGHE NALIN L [US] ET AL.) 28 September 2006 (2006-09-28) the whole document -----	1-208
A	STAAS D D ET AL: "Discovery of potent, selective 4-fluoroproline-based thrombin inhibitors with improved metabolic stability", BIOORGANIC & MEDICINAL CHEMISTRY,, vol. 14, no. 20, 15 October 2006 (2006-10-15), pages 6900-6916, XP027992972, ISSN: 0968-0896 [retrieved on 2006-10-15] table 2; compounds 41-45a -----	1-208
A	LANGE UDO E W ET AL: "Orally active thrombin inhibitors. Part 2: Optimization of the P2-moiety", BIORGANIC & MEDICINAL CHEMISTRY LETTERS, vol. 16, no. 10, 3 February 2006 (2006-02-03), pages 2648-2653, XP028674507, ISSN: 0960-894X, DOI: 10.1016/J.BMCL.2006.01.046 page 2649 - page 2650; table 2 -----	1-208
A	WO 2008/085608 A1 (IRM LLC; TULLY DAVID C [US] ET AL.) 17 July 2008 (2008-07-17) claims 1,13; examples -----	1-208
A	DANIEL RICKLIN ET AL: "Complement: a key system for immune surveillance and homeostasis", NATURE IMMUNOLOGY, vol. 11, no. 9, 19 August 2010 (2010-08-19), pages 785-797, XP055306904, New York ISSN: 1529-2908, DOI: 10.1038/ni.1923 the whole document -----	1-208

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/US2020/063391

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 2019211585	A1	07-11-2019	
		AU 2019264253 A1	19-11-2020
		CA 3097811 A1	07-11-2019
		CN 112204023 A	08-01-2021
		EP 3788039 A1	10-03-2021
		KR 20210005629 A	14-01-2021
		US 2021087172 A1	25-03-2021
		WO 2019211585 A1	07-11-2019

WO 2019186164	A1	03-10-2019	
		AU 2019244400 A1	08-10-2020
		CA 3094220 A1	03-10-2019
		CN 112165938 A	01-01-2021
		EP 3773555 A1	17-02-2021
		KR 20200135846 A	03-12-2020
		WO 2019186164 A1	03-10-2019

US 9011860	B2	21-04-2015	
		AU 2012250627 A1	21-03-2013
		BR 112013028429 A2	21-07-2020
		CA 2832871 A1	08-11-2012
		CA 3025762 A1	08-11-2012
		CL 2013003140 A1	23-05-2014
		CN 103687620 A	26-03-2014
		CN 107011443 A	04-08-2017
		DK 2704743 T3	25-05-2020
		EP 2704743 A1	12-03-2014
		EP 3725811 A2	21-10-2020
		ES 2795667 T3	24-11-2020
		HR P20200887 T1	04-09-2020
		HU E049154 T2	28-09-2020
		IL 229208 A	25-03-2021
		JP 5891298 B2	22-03-2016
		JP 6352226 B2	04-07-2018
		JP 6669800 B2	18-03-2020
		JP 2014515922 A	07-07-2014
		JP 2015227380 A	17-12-2015
		JP 2018123148 A	09-08-2018
		JP 2020097630 A	25-06-2020
		KR 20140006096 A	15-01-2014
		KR 20160087917 A	22-07-2016
		KR 20180086298 A	30-07-2018
		LT 2704743 T	10-09-2020
		MX 347691 B	09-05-2017
		MX 361175 B	29-11-2018
		NZ 617487 A	31-03-2016
		NZ 715226 A	28-07-2017
		PL 2704743 T3	24-08-2020
		PT 2704743 T	17-06-2020
		RU 2013153527 A	10-06-2015
		RU 2017137872 A	30-04-2019
		SI 2704743 T1	31-08-2020
		US 2012282263 A1	08-11-2012
		US 2016002349 A1	07-01-2016
		US 2017088632 A1	30-03-2017
		US 2019119402 A1	25-04-2019
		US 2021054099 A1	25-02-2021
		WO 2012151481 A1	08-11-2012
		ZA 201308943 B	27-05-2015

WO 2006101860	A1	28-09-2006	
		AT 450522 T	15-12-2009

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/US2020/063391

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
		AU 2006227778 A1	28-09-2006
		CA 2601523 A1	28-09-2006
		EP 1863785 A1	12-12-2007
		JP 5081807 B2	28-11-2012
		JP 2008533156 A	21-08-2008
		KR 20080002832 A	04-01-2008
		US 2007066667 A1	22-03-2007
		US 2008255108 A1	16-10-2008
		WO 2006101860 A1	28-09-2006

WO 2008085608	A1	17-07-2008	
		AR 064841 A1	29-04-2009
		AT 493174 T	15-01-2011
		AU 2007342223 A1	17-07-2008
		BR PI0720799 A2	11-03-2014
		CA 2673803 A1	17-07-2008
		CL 2008000060 A1	18-07-2008
		CO 6190626 A2	19-08-2010
		CU 20090122 A7	11-07-2011
		CY 1111340 T1	05-08-2015
		DK 2104535 T3	04-04-2011
		EA 200900920 A1	26-02-2010
		EC SP099517 A	28-08-2009
		EP 2104535 A1	30-09-2009
		EP 2279777 A2	02-02-2011
		GT 200900196 A	15-03-2010
		HK 1129860 A1	11-12-2009
		HN 2009001282 A	16-04-2012
		HR P20110201 T1	31-08-2011
		JP 2010515729 A	13-05-2010
		KR 20090087511 A	17-08-2009
		MA 31037 B1	01-12-2009
		NZ 577939 A	31-03-2011
		PE 20081566 A1	14-12-2008
		PL 2104535 T3	31-05-2011
		PT 2104535 E	31-03-2011
		RS 51644 B	31-10-2011
		SV 2009003330 A	28-07-2011
		TN 2009000294 A1	31-12-2010
		TW 200845981 A	01-12-2008
		US 2008176901 A1	24-07-2008
		WO 2008085608 A1	17-07-2008
