



(86) Date de dépôt PCT/PCT Filing Date: 2011/08/02
(87) Date publication PCT/PCT Publication Date: 2012/02/09
(45) Date de délivrance/Issue Date: 2019/03/19
(85) Entrée phase nationale/National Entry: 2013/02/04
(86) N° demande PCT/PCT Application No.: EP 2011/063291
(87) N° publication PCT/PCT Publication No.: 2012/016984
(30) Priorité/Priority: 2010/08/02 (EP10171626.4)

(51) Cl.Int./Int.Cl. *C07K 1/16* (2006.01),
C07K 1/18 (2006.01), *C07K 1/22* (2006.01),
C07K 14/46 (2006.01), *C07K 14/47* (2006.01)
(72) Inventeurs/Inventors:
ANGERMANN, AXEL, DE;
SCHECKERMANN, CHRISTIAN, DE;
SCHMIDT, KARSTEN, DE
(73) Propriétaire/Owner:
RATIOPHARM GMBH, DE
(74) Agent: GILBERT'S LLP

(54) Titre : PROCEDE DE PRODUCTION ET DE PURIFICATION D'UNE SIALYLTRANSFERASE SOLUBLE ACTIVE
(54) Title: METHOD OF PRODUCING AND PURIFYING AN ACTIVE SOLUBLE SIALYLTRANSFERASE

(57) **Abrégé/Abstract:**

The present invention relates to a method for the production and purification of a sialyltransferase polypeptide, in particular a N-Acetylgalactosamine (Gal NAc)-a-2,6-sialyltransferase I (ST6GalNAcI) polypeptide. The method comprises the steps of producing the sialyltransferase polypeptide in a Chinese Hamster Ovary (CHO) cell and purifying the polypeptide with a combination of chromatography steps. The method results in high yield of sialyltransferase polypeptide which is highly pure and active. The obtained sialyltransferase, especially ST6GalNAcI, can be employed for the glycosylation of therapeutic proteins such as G-CSF.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
9 February 2012 (09.02.2012)

PCT

(10) International Publication Number
WO 2012/016984 A1

(51) International Patent Classification:

C07K 1/16 (2006.01) **C07K 14/46** (2006.01)
C07K 1/18 (2006.01) **C07K 14/47** (2006.01)
C07K 1/22 (2006.01)

(21) International Application Number:

PCT/EP2011/063291

(22) International Filing Date:

2 August 2011 (02.08.2011)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

10171626.4 2 August 2010 (02.08.2010) EP

(71) Applicant (for all designated States except US): **BIO-GENERIX AG** [DE/DE]; Janderstrasse 3, High-Tech-Park Mannheim, 68199 Mannheim (DE).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **ANGERMAN, Axel** [DE/DE]; An der Tränk 23, 63266 Dreieich (DE). **SCHECKERMANN, Christian** [DE/DE]; Sägegasse 1c, 79238 Ehrenkirchen (DE). **SCHMIDT, Karsten** [DE/DE]; Kösliner Str. 60, 76139 Karlsruhe (DE).(74) Agent: **LASAR, Andrea**; Maiwald Patentanwalts GmbH, Elisenhof, Elisenstraße 3, Munich 80335 (DE).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report (Art. 21(3))
- with sequence listing part of description (Rule 5.2(a))

(54) Title: METHOD OF PRODUCING AND PURIFYING AN ACTIVE SOLUBLE SIALYLTRANSFERASE

(57) Abstract: The present invention relates to a method for the production and purification of a sialyltransferase polypeptide, in particular a N-Acetylgalactosamine (Gal NAc)- α -2,6-sialyltransferase I (ST6GalNAcI) polypeptide. The method comprises the steps of producing the sialyltransferase polypeptide in a Chinese Hamster Ovary (CHO) cell and purifying the polypeptide with a combination of chromatography steps. The method results in high yield of sialyltransferase polypeptide which is highly pure and active. The obtained sialyltransferase, especially ST6GalNAcI, can be employed for the glycosylation of therapeutic proteins such as G-CSF.

WO 2012/016984 A1

5

Method of producing and purifying an active soluble sialyltransferase

FIELD OF THE INVENTION

10

The present invention relates to a method for the production and purification of a sialyltransferase polypeptide, in particular a *N*-acetylgalactosamine (GalNAc)-alpha-2,6-sialyltransferase I (ST6GalNAcI) polypeptide. The method comprises the steps of producing the sialyltransferase polypeptide in Chinese Hamster Ovary (CHO)

15

cells and purifying the polypeptide with a combination of chromatography steps. The method results in high yield of sialyltransferase polypeptide which is highly pure and enzymatically active. The obtained sialyltransferase, especially ST6GalNAcI, may be employed for the glycosylation of therapeutic proteins such as G-CSF.

20

BACKGROUND OF THE INVENTION

A great diversity of oligosaccharide structures and many types of glycopeptides are found in nature, and these are synthesized, in part, by a large number of glycosyltransferases. Glycosyltransferases catalyze the synthesis of glycolipids, glycopeptides and polysaccharides by transferring an activated mono- or oligosaccharide residue from a donor to an existing acceptor molecule to initiate or elongate a carbohydrate chain. A catalytic reaction is believed to involve the recognition of both the donor and acceptor by suitable domains of the glycosyltransferase, as well as the catalytic site of the enzyme.

30

More than 30% of all therapeutic proteins and many potential peptide therapeutics are glycosylated peptides. It is well known in the art that the attachment of the correct glycan structure can play a key role in the folding, biological activity,

- 2 -

biodistribution and pharmacological efficacy of therapeutic peptides. Furthermore, glycosylation is a critically important factor influencing the *in vivo* half life and immunogenicity of therapeutic peptides. Indeed, humans will typically tolerate only those biotherapeutics that have particular types of carbohydrate attachments and will often reject glycoproteins that include non-mammalian oligosaccharide attachments. For instance, poorly glycosylated peptides are recognized by the liver as being "old" and thus, are more quickly eliminated from the body than are properly glycosylated peptides. In contrast, hyperglycosylated peptides or incorrectly glycosylated peptides can be immunogenic.

10

The production of a recombinant glycopeptide, in contrast to a recombinant non-glycosylated peptide, requires that a recombinantly produced peptide is subjected to additional processing steps, either *in vivo* within the cell or *in vitro* after the peptide has been produced by the cell. The peptide can be treated enzymatically, using a glycosyltransferase to introduce one or more glycosyl groups onto the peptide by covalently attaching the glycosyl group or groups to the peptide.

15

The production of a glycopeptide by external *in vitro*-steps of peptide processing can be time consuming and costly. This is due, in part, to the burden and cost of producing recombinant glycosyltransferases for the *in vitro* glycosylation of peptides and glycopeptides to produce glycopeptide therapeutics. As the demand and usage of recombinant glycotherapeutics increases, new methods are required in order to prepare glycopeptides more efficiently.

20

Moreover, as more and more glycopeptides are discovered to be useful for the treatment of a variety of diseases, there is a need for methods that lower the cost of their production. Further, there is also a need in the art to develop methods of more efficiently producing recombinant glycopeptides for use in developing and improving glycopeptide therapeutics.

25

30

- 3 -

Glycosyltransferases and their use for the glycosylation of proteins are disclosed in WO 2003/031464 A2 .

Sialyltransferases constitute a family of glycosyltransferases that catalyze the
5 posttranslational transfer of sialic acid (*N*-acetylneuraminic acid) to acceptor
oligosaccharide substrates at terminal positions on glycoproteins and glycolipids
(Paulson et al., 1989, *J. Biol. Chem.* 264: 17615–17618). It is estimated that the
human genome encodes more than 20 different sialyltransferases required to
synthesize all known sialo-oligosaccharide structures present in mammalian cells,
10 but only 16 distinct human sialyltransferase cDNAs have been cloned (Tsuiji S et al.,
1996, *Glycobiology* 6: 5–7; Tsuiji S, 1996, *J. Biochem.* 120: 1–13; Weinstein J et al.,
1982, *J. Biol. Chem.* 257: 13835–13844). Originally, sialyltransferases were
biochemically purified and their cDNAs were cloned using N-terminal sequences.
Comparison of the obtained cDNA sequences revealed two highly conserved regions,
15 termed the *L*- and *S*-sialylmotifs, that participate in substrate binding. Subsequently,
several sialyltransferases were cloned by PCR using degenerate primers designed
within the sialylmotifs or by expression cloning (Nara K et al., 1994, *Proc. Natl.*
Acad. Sci. USA 91: 7952–7956; Nakayama J et al., 1996, *J. Biol. Chem.* 271: 3684–
3691 ; Nakayama J et al., 1995, *Proc. Natl. Acad. Sci. USA* 92: 7031–7035). Gene
20 cloning by differential display adds an entirely different approach to the
identification of novel sialyltransferases with putative functional significance in
disease-related processes.

Sialyltransferases differ in their substrate specificity and tissue distribution, and they
25 are classified into four families according to the carbohydrate linkages they
synthesize: the ST3Gal-, ST6Gal-, ST6GalNAc-, and ST8Sia-families. The members
of each family exhibit strong activity towards certain acceptor groups, although the
substrate specificities of these enzymes overlap; one linkage can be synthesized by
multiple enzymes.

30

- 4 -

One such particular sialyltransferase that has utility in the development and production of therapeutic glycopeptides is *N*-acetylgalactosamine- α 2,6-sialyltransferase (ST6GalNAcI) which catalyzes the transfer of sialic acid from a sialic acid donor to a sialic acid acceptor. Full length chicken ST6GalNAcI enzyme, 5 for example, is disclosed by Kurosawa et al. (1994, *J. Biol. Chem.*, 269:1402-1409).

In the past, there have been efforts to increase the availability of recombinant sialyltransferases for the *in vitro* production of glycopeptides.

10 EP 0 737 745 A1 and US 5,032,519 of the Institute of Physical & Chemical Research refer to the use of *E. coli* for producing a secreted version of a protein comprising a portion, i.e. active domain, that is derived from ST6GalNAcI and is responsible for its activity.

15 WO 2007/056524 A2 of Neose Technologies Inc. describes methods of producing a modified ST6GalNAcI polypeptide, the method comprising growing a recombinant prokaryotic host cell under conditions suitable for expression of the modified ST6GalNAcI polypeptide in prokaryotic host cells. These modified ST6GalNAcI polypeptides are chimeric polypeptides comprising a first portion from a Gal-
20 β 1,3GalNAc- α 2,3-sialyltransferase (ST3GalI) polypeptide and a second portion from a GalNAc- α -2,6-sialyltransferase I (ST6GalNAcI) polypeptide. Modified ST6GalNAcI polypeptides can further be truncated polypeptides lacking all or portion of the ST6GalNAcI signal domain, all or a portion of the ST6GalNAcI transmembrane domain, and/or all or a portion of the ST6GalNAcI stem domain in
25 eukaryotic or prokaryotic host cells.

US 2006/0234345 A1 of Neose Technologies Inc. discloses a method of producing a soluble eukaryotic glycosyltransferase in a prokaryotic microorganism that has an oxidizing environment, by a) expressing a nucleic acid that encodes the eukaryotic
30 glycosyltransferase in the prokaryotic microorganism; and then b) growing the

- 5 -

prokaryotic microorganism under conditions that allow expression of the soluble active eukaryotic glycosyltransferase within a cellular compartment of the prokaryotic microorganism.

- 5 Skretas et al. (2009, *Microbial Cell Factories*, 8:50) relate to a method for the expression of the human sialyltransferase ST6GalNAcI in engineered *E. coli* strains which possess certain types of oxidative cytoplasm or which co-express the molecular chaperones/co-chaperones trigger factor, DnaK/DnaJ, GroEL/GroES, and Skp, and can produce greatly enhanced amounts of soluble ST6GalNAcI.

10

However, the capacity of *E. coli* for protein folding and forming disulfide bonds is not sufficient although there are a number of tools developed to overcome these limitations. Furthermore, the high expression yield of recombinant proteins in *E.coli* expression systems can often lead to the accumulation of aggregated, insoluble proteins that form inclusion bodies which can be a significant hindrance in obtaining soluble, active proteins (Brondyk W.H., 2009, *Methods in Enzymology*, Vol. 463, Ch.11).

15

To overcome the problems associated with recombinant sialyltransferase production in *E. coli* cultures, insect cell culture systems have been developed.

20

US 2006/0246544 A1 and US 2008/0207487 A1 disclose a method of making a composition that includes a recombinant polypeptide, e.g. sialyltransferases, wherein the polypeptide is expressed in an insect cell (e.g., using a baculoviral expression system) and wherein the composition is essentially free of endoglycanase activity. The method includes subjecting a mixture including the polypeptide to mixed-mode chromatography including the steps of: (i) contacting the mixture and a mixed-mode chromatography medium; and (ii) eluting the polypeptide from the mixed-mode chromatography medium generating a flow-through fraction comprising the polypeptide.

25

30

- 6 -

However, the complexity of the baculovirus-insect cell expression system, the limited storage stability of the required viral seed stocks and the requirement of very high virus titers for an efficient infection can limit its use for large-scale
5 bioproduction. Furthermore, viral vectors such as baculovirus have been shown to be able to infect mammalian cells, particularly human cells (Boyce FM and Buchner NL, 1996, *Proc. Natl. Acad. Sci. USA* 93:2348-2352; Lundstrom et al., 2001, *Cytotechnology*, 35: 213-221). Thus, these vectors pose a threat concerning safety issues, especially when applied for large-scale recombinant protein production,
10 where large volumes of infected cells are handled.

An alternative to overcome the described limitations of the use of insect cell culture systems is the use of mammalian cell systems for the manufacture of recombinant sialyltransferases.

15 WO 2005/121332 A2 of Neose Technologies Inc. discloses methods of producing isolated truncated ST6GaINAcI polypeptide that lacks all or a portion of the ST6GaINAcI signal domain, all or a portion of the ST6GalNAcI transmembrane domain, and/or all or a portion of the ST6GaINAcI stem domain in prokaryotic and
20 insect host cells and generally mentions that the polypeptide may also be produced in mammalian cells.

US 5,032,519 of the University of California describes methods of transfecting a host cell, e.g. a CHO cell, with a vector carrying a gene which expresses a glycosyl-
25 transferase that has the membrane anchor and most of the stem region replaced with a cleavable secretion signal segment. The resulting soluble glycosyltransferase, when expressed in the cell, is secreted by the cell. The secreted soluble glycosyltransferase is then separated from the cell culture media for use in industrial applications or carbohydrate synthesis research. Further, US 5,032,519 discloses a method of
30 purifying a soluble glycosyltransferase by using an affinity chromatography.

- 7 -

However, none of the mentioned documents relating to the production of recombinant sialyltransferases in mammalian cells discloses a method for providing a recombinant sialyltransferase which is highly active, purified to a pharmaceutical
5 grade and amenable to large scale production.

Therefore, a need still exists for efficient methods of production of recombinant sialyltransferases having activity and purity that are suitable for “pharmaceutical-scale” processes and reactions, especially for the production of glycopeptide
10 therapeutics. Thus, the problem underlying the present invention is to provide such methods for producing recombinant sialyltransferases.

SUMMARY OF THE INVENTION

15

This problem is solved according to the present invention by providing, in one aspect, a method of producing a sialyltransferase polypeptide, which comprises the steps of:

- 20 a) expressing a sialyltransferase polypeptide in a CHO cell; collecting the culture medium containing the expressed sialyltransferase polypeptide; and
- b) purifying the sialyltransferase polypeptide from the culture medium by subjecting the cell culture medium to (i) at least one affinity chromatography and/or one mixed-mode chromatography step and (ii) at
25 least one anion exchange chromatography and/or one cation exchange chromatography step.

In one embodiment, the sialyltransferase polypeptide is a truncated sialyltransferase polypeptide lacking all or portion of the sialyltransferase signal domain, all or
30 portion of the sialyltransferase transmembrane domain, and/or all or portion of the

- 8 -

sialyltransferase stem domain. Preferably, the sialyltransferase polypeptide only comprises the sialyltransferase active domain.

In a preferred embodiment, the sialyltransferase is ST6GalNAcI. Typically, the
5 ST6GalNAcI is selected from the group consisting of: human, chimpanzee, orangutan, pig, cow, dog, rat, mouse and chicken ST6GalNAcI. In a preferred embodiment, the ST6GalNAcI is a chicken ST6GalNAcI. Most preferably, the ST6GalNAcI polypeptide comprises an amino acid sequence according to SEQ ID NO:4 or SEQ ID NO: 6. According to an embodiment of the present invention, an
10 expression cassette encoding an EPO signal sequence and a sialyltransferase polypeptide sequence is used for expressing the sialyltransferase polypeptide in CHO cells.

Preferably, step a) of the production method is performed by using a serum-free fed-
15 batch culture and/or by performing an incubation temperature shift from 37°C +/- 1°C to 32°C +/- 1°C after reaching a predefined cell density.

Further, the invention includes methods wherein step c) comprises two affinity chromatography processes or one affinity chromatography process and one mixed-
20 mode chromatography process, one anion exchange chromatography process and one cation exchange chromatography process. In a most preferred embodiment, step c) is performed in the following order:

- i. an anion exchange chromatography;
- ii. a first affinity chromatography;
- 25 iii. a second affinity chromatography or a mixed-mode chromatography;
- iv. a cation exchange chromatography.

Typically, the anion exchange chromatography is performed with anion exchange resins or membranes that contain Diethylaminoethyl-groups (DEAE), quaternary
30 aminoethyl-groups (QAE), quaternary ammonium-groups (Q), Dimethylaminoethyl-

- 9 -

groups (DMAE) and/or Trimethylaminoethyl-groups (TMAE) as functional groups. Preferably it is performed with a commercially available Q-Sepharose Fast Flow resin, using a NaCl / Tris-HCl buffer as eluent at a pH in the range between 7.0 and 8.0.

5

Generally, the affinity chromatography is performed with Ni-NTA resins, Talon resins, dye chromatography resins, antibody-affinity resins, lectin affinity resins and/or peptide-ligand-affinity resins. In a preferred embodiment, the (first) affinity chromatography is performed with a commercially available Blue Sepharose FF
10 resin, preferably using a L-arginine hydrochloride / potassium phosphate buffer at a pH in the range between 7.0 and 8.0 as eluent and the mixed-mode chromatography is performed with a commercially available hydroxyapatite resin, preferably using a NaCl / potassium phosphate buffer at a pH in the range between 7.0 and 8.0 as eluent.

15

Typically, the cation exchange chromatography is performed with a resin that contains sulfopropyl cation exchange material or a resin having similar characteristics. Preferably it is performed with a commercially available SP-Sepharose High Performance resin, more preferably using a NaCl / potassium
20 phosphate buffer at a pH in the range between 6.0 and 7.0 as eluent.

In another aspect, the present invention provides a sialyltransferase polypeptide produced by any one of the methods according to the invention, wherein the sialyltransferase polypeptide is at least 98% pure, preferably at least 99% pure, most
25 preferably more than 99% pure. In a preferred embodiment, the sialyltransferase polypeptide is ST6GalNAcI.

In a further aspect, the present invention includes the use of the ST6GalNAcI polypeptide produced by any one of the methods according to the invention for the
30 glycosylation of therapeutic proteins, such as for the glycosylation of human

- 10 -

cytokines, in particular for the glycosylation of human granulocyte-colony stimulating factor, G-CSF.

5 DETAILED DESCRIPTION OF THE INVENTION

The present invention provides a method for the production and purification of an improved sialyltransferase which is highly active and pure. This objective is achieved by expressing a sialyltransferase polypeptide in a CHO cell, collecting the
10 cell culture medium containing the expressed sialyltransferase polypeptide, and purifying the sialyltransferase polypeptide from the culture medium by subjecting the culture medium to (i) at least one affinity chromatography and/or one mixed-mode chromatography process and (ii) at least one anion exchange chromatography and/or one cation exchange chromatography process.

15

More specifically, the present invention relates to a method where the purification of the sialyltransferase polypeptide is performed in the following order:

- i. an anion exchange chromatography;
- ii. a first affinity chromatography;
- 20 iii. a second affinity chromatography or a mixed-mode chromatography;
- iv. a cation exchange chromatography.

Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which
25 this invention belongs. Although any methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention, the preferred methods and materials are described herein.

A. Sialyltransferase polypeptide and expression cassette

30

- 11 -

In one embodiment, the sialyltransferase polypeptide is a truncated sialyltransferase polypeptide lacking all or portion of the sialyltransferase signal domain, all or portion of the sialyltransferase transmembrane domain, and/or all or portion of the sialyltransferase stem domain. Preferably, the sialyltransferase polypeptide only
5 comprises the sialyltransferase active domain. The sialyltransferase polypeptide can further comprise a signal peptide.

In a preferred embodiment, an expression cassette encoding an EPO signal sequence and a sialyltransferase polypeptide sequence is used for expressing a sialyltransferase
10 polypeptide in CHO cells.

"Polypeptide", alternatively referred to as a "protein", refers to a polymer in which the monomers are amino acids and are joined together through amide bonds. Additionally, unnatural amino acids, for example, P-alanine, phenylglycine and
15 homoarginine are also included. Amino acids that are not gene-encoded may also be used in the present invention. Furthermore, amino acids that have been modified to include reactive groups, glycosylation sites, polymers, therapeutic moieties, biomolecules and the like may also be used in the invention. All of the amino acids used in the present invention may be either the D- or L-isomer. The L-isomer is
20 generally preferred. In addition, other peptidomimetics are also useful in the present invention. As used herein, "peptide" refers to both glycosylated and unglycosylated peptides. Also included are peptides that are incompletely glycosylated by a system that expresses the peptide. For a general review, see Spatola, A. F., in "Chemistry and biochemistry of amino acids, peptides and proteins", B. Weinstein, eds., Marcel
25 Dekker, New York, p. 267 (1983). The term polypeptide includes molecules that are commonly referred to as proteins or peptides.

The sialyltransferase polypeptide can be any sialyltransferase polypeptide known to those of ordinary skill in the art, comprising but not being limited to: α -
30 (2,3)sialyltransferase (ST3Gal3) (Kitagawa and Paulson, 1994, *J. Biol. Chem.* 269:

- 12 -

1394–1401), α -N-acetylgalactosamide α -2,6-sialyltransferase I (ST6GalNAcI) (Kurosawa et al., 1994, *J. Biol. Chem.*, 269: 1402-1409), and β 1,3GalNAc- α 2,3-sialyltransferase (ST3GalI) (Gillespie et al., 1992, *J. Biol. Chem.* 267(29): 21004-10).

5

A "truncated sialyltransferase polypeptide" refers to a sialyltransferase that has fewer amino acid residues than a naturally occurring sialyltransferase, but that retains enzymatic activity. Any number of amino acid residues can be deleted as long as the enzyme retains activity. In some embodiments, domains or portions of domains can be deleted. In a preferred embodiment of the present invention, the sialyltransferase

10 polypeptide only comprises the sialyltransferase active domain.

An "active domain" or "catalytic domain" refers to a protein domain, or a subsequence thereof, that catalyzes an enzymatic reaction performed by the enzyme.

15 For example, a catalytic domain of a sialyltransferase will include a subsequence of the sialyltransferase sufficient to transfer a sialic acid residue from a donor to an acceptor saccharide. A catalytic domain can include an entire enzyme, a subsequence thereof, or can include additional amino acid sequences that are not attached to the naturally occurring enzyme, or a subsequence thereof. An exemplary catalytic region

20 is, but is not limited to, the catalytic domain of chicken ST6GalNAcI, which comprises amino acid residues from 232 to 566 of the full-length sequence according to SEQ ID NO: 2. The catalytic domain of chicken ST6GalNAcI is depicted in SEQ ID No. 4. In a preferred embodiment, the sialyltransferase is ST6GalNAcI. Typically, the ST6GalNAcI is selected from the group consisting of: human,

25 chimpanzee, orangutan, pig, cow, dog, rat, mouse and chicken ST6GalNAcI (Kurosawa et al., 1994, *J. Biol. Chem.*, 269:1402-1409; Skretas et al., 2009, *Microbial cell factories*, 8:50; WO 2005/121332). In a most preferred embodiment, the ST6GalNAcI is a chicken ST6GalNAcI which is encoded by the nucleotide sequence of SEQ ID NO: 1. Preferably, the ST6GalNAcI polypeptide is encoded by

- 13 -

the nucleotide sequence according to SEQ ID NO: 3 or SEQ ID NO: 5 and has an amino acid sequence according to SEQ ID NO: 4 or SEQ ID NO: 6, respectively.

B. Expression system

5

According to the present invention, the sialyltransferase polypeptide is expressed in a CHO cell or an equivalent cell line known by one of ordinary skill in the art. The particular procedure used to introduce the respective genetic material into the host cell for expression of the soluble sialyltransferase is not particularly critical. As will
10 be understood by the skilled artisan, the choice of promoter, as well as methods and strategies for introducing genetic material into the host cell used for expressing a sialyltransferase polypeptide of the invention are well-known in the art.

These include the use of plasmid vectors, viral vectors and any of the other well
15 known methods for introducing cloned genomic DNA, cDNA, synthetic DNA or other foreign genetic material into a host cell. It is only necessary that the particular genetic engineering procedure utilized be capable of successfully introducing at least one gene into the host cell which is capable of expressing the full-length or genetically modified or truncated sialyltransferase.

20

A "vector" is a composition of matter which comprises an isolated nucleic acid and which can be used to deliver the isolated nucleic acid to the interior of a cell. Various kinds of vectors are known in the art including, but not limited to, linear nucleic acids, nucleic acids associated with ionic or amphiphilic compounds, plasmids, and
25 viruses. Thus, the term "vector" includes an autonomously replicating plasmid or a genetically modified virus. The term should also be construed to include non-plasmid and non-viral compounds which facilitate transfer of nucleic acid into cells, such as, for example, polylysine compounds, liposomes, and the like.

30 Suitable vectors include pSV40, pEF-1- α , pSV2, pT-Rex, pSecTag2, pBudCE4.1, or

- 14 -

pCDNA/His Max vector, along with, for example, a CMV promoter. In a preferred embodiment of the present invention, the pMOZ-G8 vector which is designed for high level expression of a heterologous gene is used for the expression of the sialyltransferase polypeptide. Suitable selection markers are, among others,
5 neomycin, puromycin, hygromycin and dihydrofolate reductase (DHFR).

C. CHO cell culture

According to an embodiment of the present invention, step a) of the production
10 method is performed by using a serum-free fed-batch culture and by using an incubation temperature shift from 37°C +/- 1°C to 32°C +/- 1°C after reaching a predefined cell density of $\geq 5 \times 10^5$ viable cells/mL, preferably $\geq 1,5 \times 10^6$ viable cells/mL. The present inventors have found that this temperature shift increases the viability of the cells so that they can be kept in culture for a longer period. This in
15 turn results in a higher product yield.

For example, without temperature shift the cells can only be cultured for five days and a product yield of about 24 mg/l is obtained. In contrast, if a temperature shift is applied, the cells can be cultured for nine days and a product yield of 68 mg/l is
20 obtained.

The sialyltransferase polypeptide is produced in serum-free fed-batch culture using a CHO derived cell line engineered to overexpress ST6GalNAc1. The inoculum for the fermentation process is grown from one vial of the master cell bank (MCB) using T-
25 flasks or spinner flasks followed by cultivation in a seed bioreactor. The inoculum is then transferred to the production bioreactor where further cell expansion occurs until the cell density has reached an appropriate level for production.

A temperature shift is initiated at a defined cell density and the glucose level is
30 maintained within a defined range by feeding of a glucose solution. At the end of the

- 15 -

cultivation the culture is harvested. Cells and debris in the culture are removed from the harvest by depth filtration. The harvest is stored at 2-8 °C until the start of the protein purification.

- 5 Figure 1 illustrates the different cell culture and harvest steps of a preferred embodiment of the present invention.

D. Purification of sialyltransferase

- 10 According to a further aspect of the present invention, the purification of the sialyltransferase polypeptide from the culture medium is performed by subjecting the culture medium to at least one affinity chromatography step and/or mixed mode chromatography step and at least one anion and/or cation exchange chromatography step.

15

In a preferred embodiment, the purification is achieved in the following order:

- 20 i. an anion exchange chromatography, which concentrates enzyme containing solution and provides a first reduction of contaminants such as DNA, host cell proteins (HCP) and fermentation medium compounds;
- ii. a first affinity chromatography, which is used as intermediate purification step to enrich the sialyltransferase enzyme and deplete host cell proteins;
- 25 iii. a second affinity chromatography or a mixed-mode chromatography, which is used to effectively reduce host cell protein level; and
- iv. a cation exchange chromatography, which is used to remove any residual host cell protein and other contaminants.

- 30 Figure 2 illustrates the different purification steps of a preferred embodiment of the present invention.

I. Anion exchange chromatography step

According to an embodiment of the present invention, the sialyltransferase
5 polypeptide purification process includes an anion exchange chromatography (AEC)
step.

AEC relies on charge-charge interactions between the proteins in the sample and the
charges immobilized on the resin. In anion exchange chromatography, the binding
10 ions of the proteins are negative, and the functional group immobilized on the resin is
positive. Commonly used anion exchange resins are Q-resin, a quaternary amine, and
DEAE resin (DiEthylAminoEthane). However, in general the anion exchange
chromatography step can be performed with all common commercially available
anion exchange resins or membranes. Anion exchange resins may be used in the
15 form of pre-packed columns. Alternatively, columns may be self-prepared. There are
no specific limitations as to the capacity and the geometry of the columns other than
the usual ones. The person skilled in the art knows that the amount of anion
exchange resin to be used depends on the overall protein content of the cell culture
fluid or any other fluid, e.g., the eluate of a preceding chromatography step, applied
20 to the column in the capture step.

Typical strong anion exchange resins that can be used for the purpose of the
invention comprise functional groups such as: quaternary aminoethyl (QAE)
moieties, quaternary ammonium (Q) moieties and trimethylammoniumethyl (TMAE)
groups.

25 Resins having quaternary aminoethyl (QAE) moieties include, e.g., Toyopearl™
QAE (available from Tosoh Bioscience, Germany), Selectacel™ QAE (a quaternary
aminoethyl derivative of cellulose, available from Polysciences Inc., Pennsylvania
USA) and others. Resins having quaternary ammonium (Q) moieties include, e.g., Q
Sepharose™ XL, Q Sepharose™ FF, Q Sepharose™ HP, Resource Q™ (available

from GE Healthcare, Germany), Macro Prep High Q™ (Bio-Rad, California, USA),
Toyopearl Super Q (available from Tosoh Bioscience, Germany) and UNOsphere
Q™ (available from Bio-Rad, California, USA). Resins having
trimethylammoniummethyl (TMAE) groups include, e.g., Fractogel™ EMD TMAE
5 (available from Merck, Germany).

The anion exchange chromatography is preferably a strong anion exchange
chromatography which is performed using a strong anion exchange resin having –
N⁺(CH₃)₃ functional groups, or a resin having similar characteristics. Preferred
examples of strong anion exchange resins which can be used for the purpose of the
10 invention are quaternary ammonium strong anion exchanger resins known in the art
as UNOsphere Q, Q Sepharose HP, Q Sepharose FF and other resins having
quaternary ammonium (Q) moieties. In a most preferred embodiment of the present
invention, the anion exchange chromatography is performed with a commercially
available Q-Sepharose Fast Flow resin.

15
The step of anion exchange chromatography is preferably carried out using
equilibration buffers having a mildly alkaline pH. Suitable buffers include, for
example, borate buffer, triethanolamine/iminodiacetic acid, Tris, ammonium acetate,
tricine, bicine, TES, HEPES, TAPS. The use of a Tris buffer is preferred, more
20 preferably the buffer contains 20 mM Tris at a pH of 7.6. The anion exchange resin
is washed one or more times with the equilibration buffer and the flow-through
fraction is discarded. Elution from the anion exchange resin is usually achieved by
increasing the conductivity of the mobile phase through the addition of salt,
preferably sodium chloride. Preferably, the anion exchange chromatography is
25 performed using a NaCl / Tris-HCl buffer as eluent at a pH in the range between 7.0
and 8.0. More preferably, the elution buffer contains 800 mM NaCl and 20 mM
Tris/HCl at a pH of 7.6.

II. First affinity chromatography step

According to an embodiment of the present invention, the sialyltransferase polypeptide purification process includes an affinity chromatography step which preferably is a dye affinity chromatography.

5

In a preferred embodiment the method of purifying the sialyltransferase polypeptide comprises a single affinity chromatography step, more preferably the single affinity chromatography step is a dye affinity chromatography step.

- 10 Dye affinity chromatography is based on the high affinity of immobilized dyes for binding sites on the proteins of the sample. The step of dye affinity chromatography is carried out using a resin having as an immobilised ligand a dye compound which is well known to a person skilled in the art, i.e., Cibacron Blue™ F3G-A. The term "immobilized" is well understood by a person skilled in the art and means that the
- 15 ligand is derivatised in the sense that it is chemically linked to the resin. A particularly preferred resin is Blue Sepharose FF (obtainable from Amersham Biosciences Inc.).

- It is understood that the method may be performed with alternative resins, having similar characteristics. Examples of alternative resins include: Toyopearl AF-blue-HC- 650M (Tosoh Bioscience), Toyopearl SuperButyl 550, Toyopearl Phenyl 650,
- 20 Blue Cellthru BigBead™ (Sterogene), SwellGel Blue™ (Pierce), Cibachrome blue™ 3GA-agarose 100 (Sigma), Affi-Gel Blue™ (BioRad), Econo-Pac blue™ cartridges (Bio-Rad), Blue sepharose HP (Amersham), Cibacron Blue 3GA (Sigma), Blue Sepharose 6FF (GE Healthcare), ProSep PB™ (Millipore), Methyl Sepharose and
- 25 CDP Sepharose (Calbiochem).

The step of dye affinity chromatography is preferably carried out using equilibration buffers having a mildly alkaline pH. Suitable buffers include, for example, MES, Bis-Tris, ADA, PIPES, ACES, BES, MOPS, TES, HEPES. Preferably, the

- 19 -

equilibration buffer comprises potassium phosphate buffer and NaCl. More preferably, the equilibration buffer comprises 25 mM potassium phosphate buffer, pH 7.5 and 50 mM NaCl. The affinity resin is washed one or more times with the equilibration buffer and the flow-through fraction is discarded. In a more preferred
5 embodiment, the dye affinity chromatography is performed using an L-arginine hydrochloride / potassium phosphate buffer at a pH in the range between 7.0 and 8.0 as eluent. Most preferably, the protein is eluted with a buffer containing 500 mM L-arginine hydrochloride and 25 mM potassium phosphate at a pH of 7.5.

10 III. Second affinity chromatography step or mixed-mode chromatography

According to an embodiment of the present invention, the sialyltransferase polypeptide purification process includes a second affinity chromatography step as described in previous chapter II or a mixed-mode chromatography, preferably a
15 hydroxyapatite chromatography.

Hydroxyapatite chromatography is a mixed mode chromatography that utilizes an insoluble hydroxylated calcium phosphate $\text{Ca}_{10}(\text{PO}_4)_6(\text{OH})_2$, which forms both the matrix and ligand. Functional groups consist of pairs of positively charged calcium
20 ions (C-sites) and clusters of negatively charged phosphate groups (P-sites). The interactions between hydroxyapatite and proteins are complex and multi-mode. In one method of interaction, positively charged amino groups on proteins associate with the negatively charged P-sites and negatively charged carboxyl groups on the protein interact by coordination complexation to C-sites (Shepard (2000) *J. of*
25 *Chromatography* 891:93-98).

Crystalline hydroxyapatite was the first type of hydroxyapatite used in chromatography. Ceramic hydroxyapatite chromatography is a further development in hydroxyapatite chromatography. Ceramic hydroxyapatite refers to a form of
30 hydroxyapatite in which nanocrystals are agglomerated into particles and fused at

- 20 -

high temperature to create stable ceramic microspheres suitable for chromatography applications. Commercial examples of ceramic hydroxyapatite include, but are not limited to, CHT Type I and CHT Type II.

- 5 Ceramic hydroxyapatite has high durability, good protein binding capacity, and can be used at higher flow rates and pressures than crystalline hydroxyapatite. (Voia et al. (1993) *BioTechniques* 14:650-655). Hydroxyapatite has been used in the chromatographic separation of proteins, nucleic acids, as well as antibodies. In hydroxyapatite chromatography, the column is usually equilibrated, and the sample
10 applied, in a low concentration of phosphate buffer and the adsorbed proteins are then eluted in a concentration gradient of phosphate buffer (Giovannini, (2000) *Biotechnology and Bioengineering* 73:522-529).

- Any hydroxyapatite resin may be used to carry out the mixed mode chromatography
15 step of the method according to the invention. In a preferred embodiment, it is carried out on a ceramic hydroxyapatite resin, such as a type I or type II hydroxyapatite resin. The hydroxyapatite resin may have particles of any size such as 20, 40 or 80 μm . In a highly preferred embodiment, the ceramic hydroxyapatite resin comprises particles having a size of 40 μm . A hydroxyapatite resin that is
20 particularly suitable is a column commercially available under the name CHT Ceramic hydroxyapatite Type I, 40 μm .

- The equilibration buffer preferably comprises a potassium phosphate buffer at a pH between 6.3 and 7.3, more preferably it comprises 5 mM potassium phosphate
25 buffer, pH 6.8. The hydroxyapatite resin is washed one or more times with the equilibration buffer and the flow-through fraction is discarded. In a more preferred embodiment of the present invention, the hydroxyapatite affinity chromatography is performed using a NaCl / potassium phosphate buffer at a pH in the range between 6.3 and 7.3 as eluent. Most preferably, the elution buffer contains 5 mM potassium
30 phosphate buffer and 1 M NaCl at a pH of 6.8.

IV. Cation exchange chromatography step

According to an embodiment of the present invention, the sialyltransferase
5 polypeptide purification process includes a cation exchange chromatography (CEC)
step.

CEC relies on charge-charge interactions between the proteins in the sample and the
charges immobilized on the resin. In cation exchange chromatography, the binding
10 ions of the proteins are positive and the immobilized functional group is negative.
Commonly used cation exchange resins are S-resin, sulfate derivatives, and CM
(carboxymethyl) resins, carboxylated derived ions.

However, in general the cation exchange chromatography step can be performed
15 with all common commercially available cation exchange resins or membranes.
Cation exchange resins may be used in the form of pre-poured columns or
membranes on which the functional group, e.g., sulfonic acid, is fixed. Alternatively
columns may be self-prepared. There are no specific limitations as to the capacity
and the geometry of the columns other than the usual ones. The person skilled in the
20 art knows that the amount of cation exchange resin to be used depends on the overall
protein content of the cell culture fluid or any other fluid, e.g., the eluate of a
preceding chromatography step.

Different types of cation exchange materials are available under different names and
25 from a multitude of suppliers such as Bio-Rex.RTM. (e.g., type 70), Chelex.RTM.
(e.g., type 100), Macro-Prep.RTM. (e.g., type CM, High S, 25 S), AG.RTM. (e.g.,
type 50W, MP) (all available from BioRad Laboratories); WCX 2 (available from
Ciphergen), Dowex.RTM. MAC-3 (available from Dow Chemical company),
Mustang™ C and Mustang™ S (available from Pall Corporation), Cellulose CM
30 (e.g., type 23, 52), hyper-D, partisphere (available from Whatman plc.).

Amberlite.RTM. IRC (e.g., type 76, 747, 748), Amberlite.RTM. GT 73,
Toyopearl.RTM. (e.g., type SP, CM, 650M) (all available from Tosoh Bioscience
GmbH), CM 1500 and CM 3000 (available from BioChrom Labs), SP-
Sephacrose.TM., CM-Sephacrose.TM. (available from GE Healthcare), Porous resins
5 (available from PerSeptive Biosystems). Asahipak™ ES (e.g., type 502C), CXpak™
P, IEC CM (e.g., type 825, 2825, 5025, LG), IEC SP (e.g., type 420N, 825), IEC QA
(e.g., type LG, 825) (available from Shoko America Inc.), 50W cation exchange
resin (available from Eichrom Technologies Inc.). Preferably the cation exchange
material is a strong cation exchange material such as Macro-Prep.RTM. High S or
10 25S, MacroCap™ SP, Toyopearl.RTM. SP 650M, Source S™, SP Sepharose, or
POLYCAT A™.

In a preferred embodiment of the present invention, the cation exchange step is
performed with a resin that contains sulfopropyl cation exchange material or a resin
15 having similar characteristics. In a most preferred embodiment of the present
invention, the cation exchange chromatography is performed with a commercially
available SP-Sepharose High Performance resin.

The step of cation exchange chromatography is preferably carried out using an
20 equilibration buffer having a mildly acidic pH. Suitable buffers include, for example,
maleic acid, malonic acid, citric acid, lactic acid, formic acid, butaneandioic acid,
acetic acid, phosphate, HEPES and BICINE. The use of a phosphate buffer is
preferred, more preferably the equilibration buffer contains 25 mM potassium
phosphate buffer at a pH of 6.0. The cation exchange resin is washed one or more
25 times with the equilibration buffer and the flow-through fraction is discarded.
Elution from the cation exchange resin is usually achieved by increasing the
conductivity of the mobile phase through the addition of salt, preferably sodium
chloride. Preferably, the cation exchange chromatography is performed using a NaCl
/ potassium phosphate buffer at a pH in the range between 6.0 and 7.0 as eluent.

More preferably, the elution buffer contains 250 mM NaCl and 25 mM potassium phosphate having a pH of 6.0.

V. Further purification steps

5

Furthermore, the method can comprise filtration steps. Filtration processes are well known by the person skilled in the art and all common methods can be used. The method can comprise one or more ultrafiltration processes at diverse steps of the method. The method can comprise nanofiltration, e.g., for final virus clearance.

10

Ultrafiltration is a form of membrane filtration in which hydrostatic pressure forces a liquid against a semipermeable membrane. Suspended solids and solutes of high molecular weight are retained, while water and low molecular weight solutes pass through the membrane. Ultrafiltration is a commonly used method of separation for purifying and concentrating macromolecular solutions, especially protein solutions. Ultrafiltration is similar to nanofiltration, however, differing in terms of the size of the molecules it retains. In the framework of the present invention, a molecular weight cut off of 10 kDa is preferred (10 kDa UF). UF membranes may also be used for diafiltration to remove salts and other microspecies from solution via repeated or continuous dilution and reconcentration.

20

Preferably, the process of purification comprises one or more ultrafiltration/diafiltration steps. These filtration steps can be performed before, between and/or after the chromatography steps. Preferably, one diafiltration step is performed between the chromatography steps, for example between the mixed-mode chromatography step and the cation exchange chromatography step, and one ultrafiltration/diafiltration step is performed after the chromatography steps. These filtration steps can be performed using commercially available filtration devices, e.g., available from GE Healthcare or Sartorius. The ultrafiltration is preferably performed using the Sartocon™ cassettes and Sartocon™ Slice cassettes supplied by Sartorius.

25

30

- 24 -

E. Produced sialyltransferase polypeptide

In another aspect, the present invention provides sialyltransferase polypeptides
5 produced by any one of the methods according to the invention.

In one embodiment, the sialyltransferase polypeptide is a truncated sialyltransferase polypeptide lacking all or portion of the sialyltransferase signal domain, all or portion of the sialyltransferase transmembrane domain, and/or all or portion of the
10 sialyltransferase stem domain. In a preferred embodiment of the present invention, the sialyltransferase polypeptide only comprises the sialyltransferase active domain, said polypeptide being soluble. In a most preferred embodiment, the sialyltransferase is ST6GalNAcI, preferably from chicken, most preferably the sialyltransferase is the protein according to SEQ ID NO:4 or SEQ ID NO: 6.

15

The sialyltransferase polypeptides produced according to the methods of the present invention have several advantages over sialyltransferase polypeptides from the prior art. The novel and improved production methods of the present invention provide sialyltransferase polypeptides which are highly active, purified to a pharmaceutical
20 grade and amenable to large scale production.

The term "pure" refers to a sialyltransferase polypeptide that is substantially or essentially free from components which normally accompany the material in the mixture used to prepare the polypeptide, e.g. DNA or host cell proteins. Typically,
25 the sialyltransferase polypeptide produced by the method of the present invention is at least 98% pure, preferably at least 99% pure. Purity is determined by any art-recognized method of analysis (e.g., band intensity on a silver stained gel, polyacrylamide gel electrophoresis, HPLC, RP-HPLC, ELISA, or a similar means).

- 25 -

The term “active” refers to the specific activity of a sialyltransferase polypeptide produced by the method of the present invention, that means the catalytic activity of the sialyltransferase, which is transferring a sialic acid moiety from a donor molecule to an acceptor molecule.

5

The catalytic activity of ST6GalNAcI refers to the transfer of a sialic acid moiety from CMP-sialic acid by an α 2,6 linkage onto an *N*-acetylgalactosamine (GalNAc) residue O-linked to the amino acid threonine/serine of a glycoprotein. The specific activity may be expressed in activity units. As used herein, one activity unit catalyzes the formation of 1 μ mol of product per minute at a given acceptor-substrate, temperature and pH value. In the present invention, the specific activity of ST6GalNAcI is in a range between 5 U/mg and 10 U/mg, preferably 6 U/mg to 9 U/mg, most preferably 7 U/mg to 8 U/mg.

15 The sialyltransferase activity may be determined by known methods. Such methods include fluorescence assay, RP-HPLC-based assays and radioactive approaches (Spiegel et al., 1992, *J Chromatogr.*, 573(1):23-7; Gross et al., 1990, *Anal Biochem.*, 186(1):127-34; Skretas et al., 2009, *Microbial cell factories*, 8:50).

20 F. Use of sialyltransferases

In a further aspect, the present invention includes the use of the ST6GalNAcI produced by the method according to the invention for the glycosylation of therapeutic proteins, in particular for the glyco-PEGylation of human G-CSF, erythropoietin, IFN, hGH, FSH, insulin or antibodies.

The glycosyltransferase ST6GalNAcI is an essential reagent for glycosylation of therapeutic proteins. Additionally, ST6GalNAcI is an important reagent for research and development of therapeutically important glycopeptides and oligosaccharide therapeutics.

30

- 26 -

The modified ST6GalNAcI sialyltransferase enzymes of the present invention are useful for *in vivo* and *in vitro* preparation of glycosylated peptides, as well as for the production of oligosaccharides containing the specific glycosyl residues that can be transferred by the modified glycosyltransferase enzymes of the present invention. Modified forms of ST6GalNAcI polypeptides can possess biological activities comparable to, and in some instances, in excess of their full-length polypeptide counterparts.

10 The term "glycosylation" as used herein, refers to the enzymatically mediated conjugation of a modified sugar species to an amino acid or glycosyl residue of a polypeptide, e.g., an erythropoietin peptide, by the sialyltransferase prepared by the method of the present invention. Subgenera of glycosylation are "glycoconjugation" and "glyco-PEGylation," in which the modifying group of the modified sugar is
15 poly(ethylene glycol), an alkyl derivative (e.g., m-PEG) or reactive derivative (e.g., H₂N-PEG, HOOC-PEG) thereof.

A "therapeutic protein" as used herein, refers to a protein, peptide, glycoprotein or glycopeptide that is administered to a subject to treat a disease or dysfunction or to
20 improve health of the subject. In a preferred embodiment the subject is a human. In a further preferred embodiment, the therapeutic protein is a human protein. In an additional embodiment, the therapeutic protein is glycosylated or otherwise modified by one or more glycosyltransferases produced in CHO cells.

25 G-CSF (Granulocyte-colony stimulating factor) is a hematopoietic growth factor that stimulates the proliferation and differentiation of hematopoietic precursor cells and the activation of mature neutrophils. G-CSF is capable of supporting neutrophil proliferation *in vitro* and *in vivo*. The human form of G-CSF was cloned by groups from Japan and the USA in 1986 (see e.g. Nagata et al., 1986, *Nature* 319: 415-418).

30 The natural human glycoprotein exists in two forms, one having 174 and the other

- 27 -

having 177 amino acids. The more abundant and more active 174 amino acid form has been used in the development of pharmaceutical products by recombinant DNA technology.

- 5 The following examples refer to the production and purification of a truncated chicken ST6GalNAcI polypeptide and are provided merely to further illustrate the method of the invention. The scope of the invention shall not be construed as merely consisting of the following examples.

10

EXAMPLES

1. Construction of the expression cassette

- 15 An expression cassette was generated for expression of an ST6GalNAcI polypeptide consisting of the chicken glycosyltransferase (α -N-acetyl-neuraminy-2,3- β -galactosyl-1,3-N-Acetylgalactosaminide α -2,6-sialyltransferase I) N-terminally truncated at amino acid K232 to which 8 amino acids of an intervening sequence were attached to the N-terminus. The entire amino acid sequence used for expression
20 is depicted in SEQ ID NO: 6.

To add a translation start signal and to increase the expression rate, the nucleic acid encoding the truncated ST6GalNAcI polypeptide was introduced into an expression cassette of genomic human erythropoietin (EPO) to replace the coding sequences of
25 the EPO.

Overall the ST6GalNAcI transcription unit was designed to contain the following elements:

- 28 -

- EPO signal sequence (consisting of the human erythropoietin exon 1 [coding for 4 amino acids], intron 1 and a portion of exon 2 [coding for 27 amino acids: MGVHECPAWLWLLLSLLSL PLGLPVLG]);
 - ST6GalNAcI and intervening sequence nucleotide sequence (SEQ ID NO: 5);
 - 3'-UTR of human erythropoietin 285 bp;
 - 45 bp linker with 5'-Not I site; the 3'-MunI site is an integral part of the SV40 sequence.
- 10 Based on the amino acid sequence as presented in SEQ ID NO: 6, the nucleotide sequence of the mature ST6GalNAcI was optimised for codon usage in CHO cells. However, the nucleotides derived from the genomic EPO sequence were not adapted and correspond to the natural human Erythropoietin sequence. The resulting expression cassette is abbreviated EcST6. The nucleotide sequence of EcST6 was
- 15 confirmed by sequencing (SEQ ID NO: 7). Sequence congruence was 100%.

2. Construction of the expression vector

The synthetic EcST6 fragment was cloned as a *NotI/MunI* fragment into the

20 expression vector pMOZ-G8.

This vector is based on a pSV2 backbone which contains the β -lactamase expression unit which confers ampicillin resistance and the pBR322 origin of replication as prokaryotic elements and the eukaryotic selection marker dihydrofolate reductase

25 (DHFR) driven by a SV40 early gene promoter and the corresponding SV40 3'-UTR. The original mouse DHFR selection marker was modified by partial fusion with a CHO DHFR gene to obtain a recombinant hybrid DHFR (DHFR_{ec}), which shows a 10-fold lower affinity to the selective agent methotrexate (MTX). Cells transfected with the DHFR_{ec} selection marker can be selected with higher amounts of MTX to

30 block endogenous DHFR which is present in the progenitor CHOSI CHO DHFR-

- 29 -

cell line first, and to enrich for genetically altered cells harbouring plasmid derived DHFR^{rec} only. Gene expression is under control of a strong mCMV promoter. Downstream of mCMV promoter is the multiple cloning site (MCS) and a rabbit β -globin intron as well as a 3'-UTR enabling the efficient expression of cDNA transcripts.

Insertion of the cassette into the vector pMOZ-G8 was verified by PCR.

E. coli cells were transformed with the expression plasmid pMO7-G8EcST6. Colony screening of different transformants was performed and plasmid DNA was prepared from a selected clone. To confirm the identity of pMO7-G8 EcST6, the entire nucleotide sequence was verified by DNA sequencing.

3. CHO cell line

For expression of the secreted form of the chicken ST6Gal-NAcI, a Chinese hamster ovary cell line was used. The parental cell line used for production of ST6GalNAcI is a derivative of a CHOSI 4, a Chinese hamster ovary cell line (CHO dhfr-) deficient in dihydrofolate reductase (dhfr) activity which has been adapted for growth in serum and protein free medium.

Summary of host cell line CHOSI 4:

- The CHOSI 4 host cell line was derived from CHO dhfr- (ATCC number CRL-9096; ACC126 from DSMZ, Braunschweig, Germany).
- The CHOSI 4 host cell line was adapted for growth in chemically defined, serum- and protein-free medium by long term cultivation in MAM-PF2 medium.
- The morphology of CHOSI 4 is round in shape. The culture grows as a single cell suspension.
- CHOSI 4 is adapted for growth in animal component free, chemical defined MAM-PF media, which do not contain serum, proteins, peptides or hydrolysates.

- 30 -

Nevertheless, the cell line can be cultivated in other commercially available serum-free media.

- CHOSI 4 is subcultured by a total split ratio of 1:50 per week. The cells are split first 1:20 into a new flask and after 4 days obtain 3/2 of the initial volume of fresh medium. CHOSI 4 shows a specific growth rate of more than one per day ($D > 1 \times d^{-1}$) and cell densities in stirred culture systems of 2×10^6 cells/mL in batch mode and 1×10^7 cells/mL in fed-batch mode.
- Addition of attachment factors or foetal calf serum to the cell culture medium leads to a change of the cells morphology, and CHOSI 4 cells revert to an epithelial layer.

4. Cell culture and harvest

Thawing of Cells and Inoculum Expansion in T-flasks and Spinner flasks

- 15 A single vial of the MCB was taken out of storage in the vapour phase of liquid nitrogen and warmed in a water bath at $37 \pm 1^\circ\text{C}$. The cells were collected by centrifugation into a T-flask containing fresh, pre-warmed culture medium. Further inoculum expansion was achieved by dilution and subcultivation steps in spinner flasks with increasing volumes of culture medium to a final volume which is
- 20 adequate for use as seed in the bioreactor.

Cell Expansion in the Seed Bioreactor

- After cell expansion in spinner flasks the inoculum was transferred into the seed bioreactor and the volume was expanded with fresh medium to a final working
- 25 volume of 10 L. Cells were cultured in the seed bioreactor for 3 days until the adequate cell number had been reached for inoculation of the production bioreactor.

- 31 -

Cultivation in the Production Bioreactor

Following growth in the seed bioreactor, the contents of the seed bioreactor was aseptically transferred into the production bioreactor and the transferred culture was expanded to the final working volume with fresh culture medium. After reaching a predefined cell density of $\geq 1.5 \times 10^6$ viable cells/mL, the incubation temperature was lowered from $37^\circ \pm 1^\circ\text{C}$ to $32^\circ \pm 1^\circ\text{C}$. The glucose level was maintained within a defined range by feeding of a glucose solution at a concentration between 4 g/L and 8 g/L. The fermentation process was terminated 14 – 17 days after start of the cultivation in the production bioreactor.

10

Harvesting and Storage

The cell suspension of the production bioreactor was cleared from cells and debris by depth filtration of the harvest in sterile disposable bags after termination of the fermentation. A disposable depth filter was used for the filtration of the harvest. The harvest was stored at $2-8^\circ\text{C}$ until further downstream processing.

15

5. Purification

➤ Purification - Example 1

20

The purification process of ST6GalNAcI started with a capture chromatography step using an anion exchange column. The subsequent purification steps included affinity chromatography, mixed-mode chromatography, ultra-/diafiltration, cation exchange chromatography and a final nano-filtration through a 15-nm Planova® membrane.

25

The chromatographic steps were performed using gradient chromatography systems and ran automatically. UV adsorption, conductivity, pH, flow rate and back pressure were recorded at every chromatographic step by a suitable software.

The column types and resins used are presented in Table 1:

- 32 -

Step	Column type	Resin type	Gel dimensions (D x H) [mm]
Anion Exchange Chromatography	Borosilicate glass column BPG	Q-Sepharose Fast Flow	200 x 205
Affinity Chromatography	Borosilicate glass column BPG	Blue Sepharose 6-Fast Flow	140 x 110
Mixed Mode Chromatography	Borosilicate glass column BPG	Hydroxyapatite Type I 40 μ m	100 x 250
Cation Exchange Chromatography	Borosilicate glass column BPG	SP-Sepharose High performance	100 x 150

1st Column: Anion Exchange Chromatography (Q-Sepharose FF)

Anion exchange chromatography with Q-Sepharose Fast Flow resin was used as the first chromatographic step to capture ST6GalNAcI in the harvest and for volume reduction.

ST6GalNAcI was eluted by applying a salt step of 800 mM NaCl in 20 mM Tris HCl, pH 7.6. The eluate was collected based on the UV absorption profile.

Chromatographic conditions are summarized in Table 2:

Parameter	Conditions
Resin	Q-Sepharose Fast Flow (Amersham Biociences / GE-Healthcare)
Column diameter	200 mm
Bed height	205 mm +/- 15 mm
Theoretical plates	> 2,200 N/m
Asymmetry factor	0.8 - 1.5
Equilibration buffer	20 mM Tris, pH 7.6
Elution Buffer	800 mM NaCl in 20 mM Tris HCl, pH 7.6
Pooling criteria	Start: OD280 > 0.04 AU Stop: 0.6 CV after start

2nd Column: Affinity Chromatography (Blue Sepharose 6 FF)

Blue Sepharose 6FF is an agarose resin covalently linked to the dye Cibacron Blue[®] and was used to preferentially bind ST6GalNAcI in the presence of contaminants contained in the Q-Sepharose eluate. ST6GalNAcI was eluted with 500 mM L-arginine hydrochloride in 25 mM potassium phosphate buffer, pH 7.5. The eluate was collected based on the UV absorption profile.

- 33 -

Chromatographic conditions are summarized in Table 3:

Parameter	Conditions
Resin	Blue Sepharose FF (Amersham Biosciences / GE-Healthcare)
Column diameter	140 mm
Bed height	110 mm +/- 20 mm
Theoretical plates	> 2,778 N/m
Asymmetry factor	0.8 - 1.5
Equilibration buffer	25 mM potassium phosphate buffer pH 7.5, 50 mM NaCl
Elution Buffer	500 mM L-arginine hydrochloride 25 mM potassium phosphate pH 7.5
Pooling criteria	Start: OD > 0.010 AU Stop: OD < 0.05 AU

5 *3rd Column: Mixed-mode Chromatography (Hydroxyapatite)*

Mixed-mode chromatography on a hydroxyapatite resin was used as an intermediate step to further reduce the levels of HCP and DNA. ST6GalNAcI was eluted with 1.0 M NaCl in 5 mM sodium phosphate buffer pH 6.8. The eluate was collected based on the UV absorption profile.

10

Chromatographic conditions are summarized in Table 4:

Parameter	Conditions
Resin	Hydroxyapatite CHT Type I 40 µm (BioRad)
Column diameter	100 mm
Bed height	250 mm +/- 20 mm
Theoretical plates	> 4,760 N/m
Asymmetry factor	0.8 - 2.3
Equilibration buffer	5 mM Potassium phosphate buffer pH 6.8
Elution Buffer	5 mM Potassium phosphate buffer pH 6.8, 1 M NaCl
Pooling criteria	Start: OD > 0.050 AU Stop: 1.34 CV

- 34 -

Ultrafiltration/Diafiltration

The hydroxyapatite eluate was diafiltered using a Sartocan slice module with a molecular weight cut-off (MWCO) of 30 kDa. The tangential flow filtration was conducted to desalt and concentrate the product and finally exchange the buffer to 25 mM potassium phosphate buffer to pH 6.0.

4th Column: Cation Exchange Chromatography (SP Sepharose)

Cation exchange chromatography was used as the final polishing step to remove any possibly remaining residual HCP. Under the specific conditions applied ST6GalNAcI was eluted with 250 mM NaCl in 25 mM potassium phosphate buffer pH 6.0.

Chromatographic conditions are summarised in Table 5:

Parameter	Conditions
Resin	SP-Sepharose High Performance
Column diameter	100 mm
Bed height	150 mm +/- 20 mm
Theoretical plates	> 7,353 N/m
Asymmetry factor	0.8 - 1.5
Equilibration buffer	25 mM potassium phosphate buffer pH 6.0
Elution Buffer	250 mM Sodium chloride, 25 mM Potassium phosphate pH 6.0
Pooling criteria	Start OD > 0.050 AU Stop: OD < 0.200 AU + 0.6 CV

Ultrafiltration/Diafiltration and Formulation

The SP-Sepharose eluate was diafiltrated in a Sartocan slice module with a MWCO of 30 kDa. The tangential flow filtration was conducted to desalt and concentrate the product and exchange the buffer to 50 mM BisTris, pH 6.5, 100 mM NaCl, 5% Sorbitol.

After completion of the diafiltration Tween 80 was added to a final concentration of 0.003%.

- 35 -

➤ Purification - Example 2

1st Column: Anion Exchange Chromatography (Q Sepharose FF)

- 5 This process step is used for capture of cST6 by anion exchange chromatography from clarified, sterile filtered bioreactor harvest. 1000 mL supernatant from bioreactor was diluted 1:3 in Buffer A (20 mM Tris pH 7.6) and loaded on an omni 25/150 Q Sepharose FF column. The cST6 was eluted by a linear NaCl gradient (0 - 1 M, 20 CV).

10

2nd Column: Affinity Chromatography (CDP affinity matrix)

- This affinity chromatography process step is an intermediate step which is important to increase the cST6 purity. The pool of fractions 9-11 from the Q Sepharose capture step was diluted 1:3 in Buffer A (10 mM MES pH 6.8, 25% glycerol) and loaded on
15 a XK 10/55 CDP Affinity column. To avoid overloading of the CDP column the load was applied in two identical CDP runs. The cST6 was eluted by a linear NaCl gradient (0 - 1 M, 10 CV).

3rd Column: Anion Exchange Chromatography (Mono Q Sepharose)

- 20 This anion exchange chromatography was used as the final polishing step. The pool of the cST6 fractions from the CDP intermediate step was diluted 1:10 in Buffer A (20 mM Tris pH 7.6) and loaded on a Mono Q HR 10/10 column. The cST6 was eluted by a linear NaCl gradient (0 - 0.6 M, 5 CV).

25 ➤ Purification - Example 3

1st Column: Anion Exchange Chromatography (Q Sepharose FF)

Anion exchange chromatography with Q-Sepharose Fast Flow resin was used as the first chromatographic step to capture ST6GalNAcI in the harvest and for volume

- 36 -

reduction. 300 mL clarified and sterile filtered supernatant from fed-batch cultivation bioreactor was purified.

Chromatographic conditions are summarized in Table 6:

Column	Omnifit 25, bed height = 15 cm, CV = 73.6 mL
Equilibration buffer A	20 mM Tris, pH 7.6
Elution buffer B	20 mM Tris, pH 7.6, 1 M NaCl
Dilution factor for load	Sample dilution 1:3 in equilibration buffer A
Gradient	Steps: 4 % (4CV), 20 % (4 CV) and 100 % (2 CV) elution buffer B

5

2nd Column: Intermediate Affinity Chromatography (Blue Sepharose FF)

This affinity chromatography process step is an intermediate step which is important to increase the cST6 purity. Eluate of the capture chromatography step (29 mL) was diluted and purified by a Blue Sepharose chromatography step.

10

Chromatographic conditions are summarized in Table 7:

Column	Omnifit 10, bed height = 7.0 cm, CV = 5.5 mL
Equilibration buffer A	25 mM potassium phosphate, pH 7.5, 50 mM NaCl
Elution buffer B	25 mM potassium phosphate, pH 7.5. 1 M arginine-HCl
Dilution factor for load	Sample dilution 1:3 in equilibration buffer A
Gradient	Steps: 10 % B, 75 % B, 100 % B, (each step 5CV)

3rd Column: Polishing Affinity Chromatography (Prosep PB)

ProSep-PB media is composed of a synthetic m-aminophenyl ligand immobilized on controlled pore size glass beads.

15

This affinity chromatography was used as the final polishing step. 15 mL intermediate eluate was purified in this chromatography step.

20

- 37 -

Chromatographic conditions are summarized in Table 8:

Column	Omnifit 10, bed height = 5.8 cm, CV = 4.55 mL
Equilibration buffer A12	25 mM potassiumphosphate, pH 6.5, 0.5 M NaCl
Equilibration buffer A11	50 mM BisTris, pH 6.5, 0.5 M NaCl
Elution buffer B	50 mM BisTris, pH 6.5, 0.5 M NaCl, 1.37 M Sorbitol
Dilution factor for sample load	Dilution 1:5 in equilibration buffer A12
Gradient	Step, 100 % B over 5 CV
Dilution buffer	50 mM BisTris, pH 6.5, 0.015 % Tween80
Formulation buffer	50 mM BisTris, pH 6.5, 0.1 M NaCl, 5 % Sorbitol, 0.003 % Tween80

5

6. ST6GalNAcI purity, specific activity and yield of production

➤ Example 1

10 The purity of the ST6GalNAcI purified according to example 1 was measured by Coomassie stained SDS-PAGE (see Figure 3), RP-HPLC, ELISA, and Threshold assay in three batches. The specific activity of the ST6GalNAcI was measured by RP-HPLC.

15 Purity and specific activities were as shown in Table 9:

Test	Method	Batch # 1	Batch # 2	Batch # 3
Identity	SDS-PAGE	Main band at approx. 66 kDa.	Main band at approx. 66 kDa.	Main band at approx. 66 kDa.
Purity	RP-HPLC	100.0 %	100.0 %	99.5 %
Residual CHO host-cell protein (Generic	ELISA	35 ng/mg (ppm)	35 ng/mg (ppm)	37 ng/mg (ppm)

- 38 -

assay)				
Residual host-cell DNA	Threshold assay	≤ 784 pg/mg	≤ 833 pg/mg	≤ 755 pg/mg
Specific activity	RP-HPLC	5,6 U/mg	7,8 U/mg	7,7 U/mg

The yield of production of the purified ST6GalNAcI was in a range from 65 mg/L to 75 mg/L.

- 5 A comparability study was made to compare the expression of ST6GalNAcI in CHO cells with the expression of ST6GalNAcI in insect cells. Insect cell-derived ST6GalNAcI showed a specific activity of 2.5 U/mg and the CHO-derived ST6GalNAcI showed a higher specific activity of 7.8 U/mg.

10 ➤ Example 2

The purified cST6 fractions according to example 2 were pooled and analyzed for purity via Coomassie-stained SDS-PAGE (see Figure 4). The cST6 sample pools coming from three different clones C1, C2 and C3 were loaded undiluted, diluted 1:5 and diluted 1:10.

Table 10 shows sample preparation and loading:

Well	<i>Sample</i>	Sample volume [μl]	Sample buffer [μl]	Loading volume [μl]	Conditions
1	C1 pool after 3 rd column (Mono Q)	10	10	20	boiling / reduced
2	C2 pool after 3 rd column (Mono Q)	10			
3	C3 pool after 3 rd column (Mono Q)	10			
4	-	-	10	10	
5	C1 pool after 3 rd column (Mono Q)	2+8 H2O	10	20	
6	C2 pool after 3 rd column (Mono Q)	2+8 H2O			
7	C3 pool after 3 rd column (Mono Q)	2+8 H2O			
8	-	-	10	10	
9	C1 pool after 3 rd column (Mono Q)	1+9 H2O	10	20	
10	C2 pool after 3 rd column (Mono Q)	1+9 H2O			
11	C3 pool after 3 rd column (Mono Q)	1+9 H2O			
12	Low molecular weight marker			10	

- 39 -

As shown in Fig. 4, the undiluted pools show minor contaminations below 20 kDa. The signal of the 1:10-diluted pools was stronger than the signal of the impurities in the undiluted samples. Thus the purity can be estimated to be > 90%.

5 ➤ Example 3

The purity of the ST6GalNAcI purified according to example 3 was measured by Coomassie-stained SDS-PAGE, RP-HPLC, ELISA, and Threshold assay in three batches. The specific activity of the ST6GalNAcI was measured by RP-HPLC.

10

FIGURE LEGENDS

Figure 1:

15 Sialyltransferase manufacturing process: cell culture and harvest

Figure 2:

Sialyltransferase manufacturing process: purification

20 Figure 3:

Protein gel (SDS-PAGE) that has been stained with Coomassie Blue after the different purification steps of example 1. The band at around 66 kDa corresponds to the purified ST6GalNAcI polypeptide according to SEQ ID NO: 6.

25 Figure 4:

Protein gel (SDS-PAGE) stained with Coomassie Blue of cST6 pools from three different clones C1, C2 and C3 after third column of the purification according to example 2. The band at around 66 kDa corresponds to the purified ST6GalNAcI polypeptide according to SEQ ID NO:6.

30

What is claimed is:

1. A method of producing a sialyltransferase polypeptide, which comprises the steps of:
 - (a) expressing a sialyltransferase polypeptide in CHO cells; collecting the cell culture medium containing the expressed sialyltransferase polypeptide; and
 - (b) purifying the sialyltransferase polypeptide from the cell culture medium by subjecting the cell culture medium to, in the following order,:
 - (i) an anion exchange chromatography;
 - (ii) a dye affinity chromatography;
 - (iii) a mixed-mode chromatography using a hydroxyapatite resin;
 - (iv) a cation exchange chromatography,thereby providing the sialyltransferase polypeptide.
2. The method according to claim 1, wherein the sialyltransferase is ST6GalNAcI.
3. The method according to claim 2, wherein the ST6GalNAcI is selected from the group consisting of: human, chimpanzee, orangutan, pig, cow, dog, rat, mouse and chicken ST6GalNAcI.
4. The method according to claim 3, wherein the ST6GalNAcI is a chicken ST6GalNAcI.
5. The method according to claim 4, wherein the ST6GalNAcI polypeptide comprises an amino acid sequence according to SEQ ID NO: 4 or SEQ ID NO: 6.
6. The method according to any one of claims 1 to 5, wherein the sialyltransferase polypeptide only comprises the sialyltransferase active domain.

7. The method according to any one of claims 1 to 6, wherein an expression cassette encoding an EPO signal sequence and a sialyltransferase polypeptide sequence is used for expressing the sialyltransferase polypeptide in CHO cells.
8. The method according to any one of claims 1 to 7, wherein step a) is performed by applying an incubation temperature shift from 37°C +/- 1°C to 32° C +/- 1°C after reaching a predefined cell density.
9. The method according to any one of claims 1 to 8, wherein the anion exchange chromatography is performed using a resin carrying a quaternary ammonium as a functional group.
10. The method according to claim 9, wherein the anion exchange chromatography is performed using a NaCl / Tris-HCl buffer as eluent at a pH in the range between 7.0 and 8.0.
11. The method according to claim 1, wherein the dye affinity chromatography is performed using a Blue Sepharose™ Fast Flow resin.
12. The method according to claim 11, wherein the Blue Sepharose™ affinity chromatography is performed using a L-arginine hydrochloride / potassium phosphate buffer as eluent at a pH in the range between 7.0 and 8.0.
13. The method according to claim 1, wherein the mixed mode chromatography using a hydroxyapatite resin is performed using a NaCl / potassium phosphate buffer as eluent at a pH in the range between 6.3 and 7.3.
14. The method according to any one of claims 1 to 13, wherein the cation exchange chromatography is performed with a resin that contains sulfopropyl cation exchange material.

15. The method according to claim 14, wherein the cation exchange chromatography is performed using SP-Sepharose™ High Performance resin.

16. The method according to claim 15, wherein the cation exchange chromatography is performed using a NaCl / potassium phosphate buffer as eluent at a pH in the range between 6.0 and 7.0.

17. The method according to any one of claims 1 to 16, wherein step (b) is performed in the following order:

- (i) an anion exchange chromatography using a Q-Sepharose™ Fast Flow resin;
- (ii) a dye affinity chromatography using a Blue Sepharose™ Fast Flow resin;
- (iii) a mixed-mode chromatography using a hydroxyapatite resin;
- (iv) a cation exchange chromatography using a SP-Sepharose™ High Performance resin.


Figure 1


Figure 2

Figure 3




Figure 4