



(86) Date de dépôt PCT/PCT Filing Date: 2002/04/01
(87) Date publication PCT/PCT Publication Date: 2002/10/10
(45) Date de délivrance/Issue Date: 2015/10/13
(85) Entrée phase nationale/National Entry: 2003/09/26
(86) N° demande PCT/PCT Application No.: US 2002/010196
(87) N° publication PCT/PCT Publication No.: 2002/079126
(30) Priorités/Priorities: 2001/03/30 (US60/280,590);
2001/03/30 (US60/280,462); 2001/04/16 (US60/284,213);
2001/04/20 (US60/285,597); 2001/12/14 (US60/340,588)

(51) Cl.Int./Int.Cl. *C07F 15/00* (2006.01),
B01J 31/12 (2006.01), *C07C 17/275* (2006.01),
C07C 29/40 (2006.01), *C07C 319/20* (2006.01),
C07C 45/69 (2006.01), *C07C 67/293* (2006.01),
C07C 67/343 (2006.01), *C07D 317/12* (2006.01),
C07D 317/20 (2006.01), *C07D 317/24* (2006.01),
C07F 7/18 (2006.01), ...

(72) Inventeurs/Inventors:
GRUBBS, ROBERT H., US;
CHATTERJEE, ARNAB K., US;
CHOI, TAE-LIM, US;
GOLDBERG, STEVEN D., US;
LOVE, JENNIFER A., US; ...

(54) Titre : REACTION DE METATHESE CROISEE D'OLEFINES FONCTIONNALISEES ET SUBSTITUEES AU MOYEN
DE COMPLEXES DE CARBENE METALLIQUE DE TRANSITION DU GROUPE 8 EN TANT QUE CATALYSEURS
DE METATHESE

(54) Title: CROSS-METATHESIS REACTION OF FUNCTIONALIZED AND SUBSTITUTED OLEFINS USING GROUP 8
TRANSITION METAL CARBENE COMPLEXES AS METATHESIS CATALYSTS

(57) **Abrégé/Abstract:**

The invention pertains to the use of Group 8 transition metal carbene complexes as catalysts for olefin cross-metathesis reactions. In particular, ruthenium and osmium alkylidene complexes substituted with an N-heterocyclic carbene ligand are used to catalyze cross-metathesis reactions to provide a variety of substituted and functionalized olefins, including phosphonate-substituted olefins, directly halogenated olefins, 1,1,2- trisubstituted olefins, and quaternary allylic olefins. The invention further provides a method for creating functional diversity using the aforementioned complexes to catalyze cross-metathesis reactions of a first olefinic reactant, which may or may not be substituted with a functional group, with each of a plurality of different olefinic reactants, which may or may not be substituted with functional groups, to give a plurality of structurally distinct olefinic products. The methodology of the invention is also useful in facilitating the stereoselective synthesis of 1,2-disubstituted olefins in the cis configuration.

(51) Cl.Int./Int.Cl. (suite/continued) *C07F 9/40* (2006.01), *C40B 50/00* (2006.01), *C07D 233/28* (2006.01)

(72) Inventeurs(suite)/Inventors(continued): MORGAN, JOHN P., US; SANDERS, DANIEL P., US; SCHOLL, MATTHIAS, US; TOSTE, F. DEAN, US; TRNKA, TINA M., US

(73) Propriétaires(suite)/Owners(continued): CALIFORNIA INSTITUTE OF TECHNOLOGY, US

(74) Agent: ADE & COMPANY INC.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
10 October 2002 (10.10.2002)

PCT

(10) International Publication Number
WO 02/079126 A1

(51) International Patent Classification⁷: **C07C 6/02**,
6/04, 6/06

(21) International Application Number: PCT/US02/10196

(22) International Filing Date: 1 April 2002 (01.04.2002)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

60/280,590	30 March 2001 (30.03.2001)	US
60/280,462	30 March 2001 (30.03.2001)	US
60/284,213	16 April 2001 (16.04.2001)	US
60/285,597	20 April 2001 (20.04.2001)	US
60/340,588	14 December 2001 (14.12.2001)	US

(71) Applicant (for all designated States except US): **CALIFORNIA INSTITUTE OF TECHNOLOGY** [US/US];
1200 East California Boulevard, Pasadena, CA 91125 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **GRUBBS, Robert, H.** [US/US]; 1700 Spruce Street, South Pasadena, CA 91030 (US). **CHATTERJEE, Arnab, K.** [US/US]; 71 N. Wilson Avenue #20, Pasadena, CA 91106 (US). **CHOI, Tae-Lim** [KR/US]; 434 S. Catalina Avenue, #102, Pasadena, CA 91106 (US). **GOLDBERG, Steven, D.** [US/US]; 86 West Glenarm Street, Pasadena, CA 91105 (US). **LOVE, Jennifer, A.** [US/US]; 222 S. Holliston Avenue, #204, Pasadena, CA 92215 (US). **MORGAN, John, P.** [US/US]; 2148 E. Crary Street, Pasadena, CA 91104 (US). **SANDERS, Daniel, P.** [US/US]; 442 S. Catalina Avenue, #302, Pasadena, CA 91106 (US). **SCHOLL,**

Matthias [DE/US]; 1770 Massachusetts Avenue, #599, Cambridge, MA 02139 (US). **TOSTE, F., Dean** [CA/US]; 696 Mira Monte Place, #2, Pasadena, CA 91101 (US). **TRNKA, Tina, M.** [US/US]; 278 South Oak Knoll Avenue, #2, Pasadena, CA 91101 (US).

(74) Agents: **REED, Dianne, E.** et al.; Reed & Associates, 800 Menlo Avenue, Suite 210, Menlo Park, CA 94025 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: CROSS-METATHESIS REACTION OF FUNCTIONALIZED AND SUBSTITUTED OLEFINS USING GROUP 8 TRANSITION METAL CARBENE COMPLEXES AS METATHESIS CATALYSTS

(57) Abstract: The invention pertains to the use of Group 8 transition metal carbene complexes as catalysts for olefin cross-metathesis reactions. In particular, ruthenium and osmium alkylidene complexes substituted with an N-heterocyclic carbene ligand are used to catalyze cross-metathesis reactions to provide a variety of substituted and functionalized olefins, including phosphonate-substituted olefins, directly halogenated olefins, 1,1,2-trisubstituted olefins, and quaternary allylic olefins. The invention further provides a method for creating functional diversity using the aforementioned complexes to catalyze cross-metathesis reactions of a first olefinic reactant, which may or may not be substituted with a functional group, with each of a plurality of different olefinic reactants, which may or may not be substituted with functional groups, to give a plurality of structurally distinct olefinic products. The methodology of the invention is also useful in facilitating the stereoselective synthesis of 1,2-disubstituted olefins in the *cis* configuration.



WO 02/079126 A1

CROSS-METATHESIS REACTION OF FUNCTIONALIZED AND SUBSTITUTED
OLEFINS USING GROUP 8 TRANSITION METAL CARBENE COMPLEXES
AS METATHESIS CATALYSTS

5

TECHNICAL FIELD

This invention relates generally to a method for carrying out an olefin metathesis reaction using a Group 8 transition metal complex as a catalyst, and more particularly relates to a method for carrying out cross-metathesis reactions using the aforementioned catalyst wherein at least one of the olefinic reactants is a functionalized olefin, a geminal disubstituted olefin, a trisubstituted
 10 olefin, and/or a quaternary allylic olefin. Methods are also provided for the catalysis of stereoselective olefin metathesis reactions, and for the creation of chemical diversity by carrying out a plurality of olefin metathesis reactions using a single olefinic substrate and different metathesis partners, to generate a plurality of structurally distinct products.

15


BACKGROUND OF THE INVENTION

To the synthetic organic or polymer chemist, simple methods for forming carbon-carbon bonds are extremely important and valuable tools. One method of C-C bond formation that has proved particularly useful is transition-metal catalyzed olefin metathesis. "Olefin metathesis," as is understood in the art, refers to the metal-catalyzed redistribution of carbon-carbon bonds. See
 20 Trnka and Grubbs (2001) *Acc. Chem. Res.* 34:18-29. Over two decades of intensive research effort has culminated in the discovery of well-defined ruthenium and osmium carbenes that are highly active olefin metathesis catalysts and stable in the presence of a variety of functional groups.

These ruthenium and osmium carbene complexes have been described in U.S.
 25 Patent Nos. 5,312,940, 5,342,909, 5,831,108, 5,969,170, 6,111,121, and 6,211,391 to Grubbs et al., assigned to the California Institute of Technology. The ruthenium and osmium carbene complexes disclosed in these patents all possess metal centers that are formally in the +2 oxidation state, have an electron count of 16, and are penta-coordinated. These catalysts are of the general formula (I)


30

(I)



where M is a Group 8 transition metal such as ruthenium or osmium, X and X' are anionic ligands, L and L' are neutral electron donors, and R and R' are specific substituents, e.g., one may be H and the other may be a substituted or unsubstituted hydrocarbyl group such as phenyl or C=C(CH₃)₂. Such complexes have been disclosed as useful in catalyzing a variety of olefin metathesis reactions, including ring opening metathesis polymerization ("ROMP"), ring closing metathesis ("RCM"), acyclic diene metathesis polymerization ("ADMET"), ring-opening metathesis ("ROM"), and cross-metathesis ("CM" or "XMET") reactions.





For the most part, such metathesis catalysts have been prepared with phosphine ligands, e.g., triphenylphosphine or dimethylphenylphosphine, exemplified by the well-defined, metathesis-active ruthenium alkylidene complexes (II) and (III)



wherein "Cy" is a cycloalkyl group such as cyclohexyl or cyclopentyl. See U.S. Patent No. 5,917,071 to Grubbs et al. and Trnka and Grubbs, cited *supra*. These compounds are highly reactive catalysts useful for catalyzing a variety of olefin metathesis reactions, and are tolerant of many different functional groups. However, as has been recognized by those in the field, the compounds display low thermal stability, decomposing at relatively low temperatures. Jafarpour and Nolan (2000) *Organometallics* 19(11):2055-2057.


Recently, however, significant interest has focused on the use of N-heterocyclic carbene ligands as superior alternatives to phosphines. See, e.g., Trnka and Grubbs, *supra*; Bourissou et al. (2000) *Chem. Rev.* 100:39-91; Scholl et al. (1999) *Tetrahedron Lett.* 40:2247-2250; Scholl et al. (1999) *Organic Lett.* 1(6):953-956; and Huang et al. (1999) *J. Am. Chem. Soc.* 121:2674-2678. N-heterocyclic carbene ligands offer many advantages, including readily tunable steric bulk, vastly increased electron donor character, and compatibility with a variety of metal species. In addition, replacement of one of the phosphine ligands in these complexes significantly improves thermal stability. The vast majority of research on these carbene ligands has focused on their generation and isolation, a feat finally accomplished by Arduengo and coworkers within the last ten years (see, e.g., Arduengo et al. (1999) *Acc. Chem. Res.* 32:913-921). Representative of these second generation catalysts are the four ruthenium complexes (IVA), (IVB), (VA) and (VB):

-3-




15 In the above structures, Cy is as defined previously, "IMes" represents 1,3-dimesityl-imidazol-2-ylidene

20 IMes:



and "IMesH₂" represents 1,3-dimesityl-4,5-dihydroimidazol-2-ylidene

25 IMesH₂:



30 Other complexes formed from N-heterocyclic carbene ligands are also known.

35 These transition metal carbene complexes, particularly those containing a ligand having the 4,5-dihydroimidazol-2-ylidene structure, such as in IMesH₂, have been found to address a number of previously unsolved problems in olefin metathesis reactions, particularly cross-metathesis reactions. These problems span a variety of reactions and starting materials. The following discussion focuses on representative problems in the art that have now been addressed by way of the present invention.

Use of Olefinic Phosphonates and Other Functionalized Olefins as Cross-Metathesis

Reactants:

Olefins that contain phosphonate functionality are used extensively in synthetic organic chemistry. For example, allylic phosphonates are employed in the preparation of dienes and polyenes by Horner-Emmons olefination, providing products with improved stereoselectivity as compared to the corresponding phosphonium salts; see Crombie et al. (1969) *J. Chem. Soc., Chem. Commun.* at 1024; and Whang et al. (1992) *J. Org. Chem.* 56 :7177. The reaction of organic halides with trialkyl phosphites (Michaelis-Arbuzov reaction) is used primarily for the synthesis of allylphosphonates; see Bhattacharya et al. (1981) *Chem. Rev.* 81 :415. However, elimination and/or loss of olefin stereochemical integrity are often competitive with product formation. Palladium catalyzed cross-coupling of hydrogen phosphonates to conjugated dienes and allenes has also been developed, but requires high reaction temperatures and provide low regioselectivity in highly substituted phosphonates products. See Hirao et al. (1980) *Tetrahedron Lett.* 21:3595; Hirao et al. (1982) *Bull. Chem. Soc. Jpn.* 55: 909; Imamoto et al. (1990) *J. Am. Chem. Soc.* 112 :5244; Zhao et al. (2000) *Organometallics* 19:4196.

Vinylphosphonates are important synthetic intermediates and have been investigated as biologically active compounds. Vinylphosphonates have been used as intermediates in stereoselective synthesis of trisubstituted olefins and in heterocycle synthesis; see Shen et al. (2000) *Synthesis*, p. 99; Tago et al. (2000) *Org. Lett.* 2:1975; Kouno et al. (1998) *J. Org. Chem.* 63:6239; and Kouno et al. (2000) *J. Org. Chem.* 65:4326. The synthesis of vinylphosphonates has also been widely examined and a variety of non-catalytic approaches have been described in the literature. Recent metal-catalyzed methods include palladium catalyzed cross-coupling (see, e.g., Holt et al. (1989), *Tetrahedron Lett.* 30:5393; Han et al. (1996), *J. Am. Chem. Soc.* 118:1571; Kazankova et al. (1999), *Tetrahedron Lett.* 40:569; Okauchi et al. (1999), *Tetrahedron Lett.* 40:5337; Zhong et al. (2000), *Synth. Commun.* 30:273; and Han et al. (2000), *J. Am. Chem. Soc.* 122:5407) and Heck coupling of aryldiazonium salts with vinyl phosphonates (Brunner et al. (2000) *Synlett.* at p. 201), but are limited by the requirement of highly reactive functional groups in the substrates. Therefore, a more mild, general and stereoselective method for the synthesis of vinyl and allylphosphonates using commercially available starting materials would be quite valuable, and would provide an additional degree of orthogonality to the previously reported syntheses. An ideal such method would also be applicable in other contexts as well, for example in the synthesis of olefins substituted with functional groups other than phosphonates. The invention, in one embodiment, is directed to this pressing need in the art, and provides a method that not only accomplishes the aforementioned goals, but is also useful in a more generalized process for creating functional group diversity in a population of olefinic products prepared using cross-metathesis.

Cross-Metathesis of α -Halogenated Olefins and Synthesis of Directly Halogenated Olefins:

Since the discovery of the olefin metathesis reaction in the 1950s, the metathesis of halogen-containing olefins has received very little attention. The metathesis of allyl bromide, allyl chloride, and related substrates with the heterogeneous $\text{Re}_2\text{O}_7/\text{Al}_2\text{O}_3/\text{Me}_4\text{Sn}$ catalyst system are among the few examples. Kawai et al. (1998) *J. Mol. Catal. A* 133:51; Bogolepova et al. (1992) *Petrol. Chem.* 32:461; Mol et al. (1979) *J. Chem. Soc. Chem. Commun.*, at pp. 330-331 Nakamura et al. (1977) *Chem. Lett.*, at p. 1127; Fridman et al. (1997) *Doklady Akad. Nauk S.S.S.R.* 234:1354. Most recently, the cross-metathesis of 3,3,4,4,5,5,6,6,6-nonafluoro-1-hexene with terminal olefin and the dimerization of vinyl gem-difluorocyclopropane derivatives have been achieved using catalyst (VB). Chatterjee et al. (2000) *J. Am. Chem. Soc.* 122:3783; International Patent Publication No. WO 02/00590 to Grubbs et al.; Itoh et al. (2000) *Org. Lett.* 2:1431. In these cases, the substrates are challenging because of the electron-withdrawing nature of the pendent halogens. A particularly challenging situation arises when the olefin is directly halogenated, because the metathesis reaction will then involve a monohalo $[\text{M}]=\text{CXR}$ or dihalo $[\text{M}]=\text{CX}_2$ carbene complex as the propagating species (where X = halide), rather than the more usual alkylidene $[\text{M}]=\text{CR}_2$ (where R = H, alkyl, aryl). To the best of applicants' knowledge, there has been only one report of metathesis involving directly halogenated olefins, namely the cross-metathesis of 1-chloro- and 1-bromoethylene with propylene using $\text{Re}_2\text{O}_7/\text{Al}_2\text{O}_3/\text{Me}_4\text{Sn}$ (Fridman et al. (1977) *Doklady Akad. Nauk S.S.S.R.* 234:1354).

Accordingly, there are very few methods available for the mild and selective synthesis of directly halogenated olefins, and in particular, directly fluorinated olefins. The present invention now provides a straightforward method for carrying out an olefin metathesis reaction using an α -halogenated olefin, which may be an α -fluorinated olefin, in order to provide a directly halogenated (e.g., fluorinated) olefinic product.

Catalyzed Cross-Metathesis of Highly Substituted Olefins, Including Geminal Disubstituted Olefins and Quaternary Allylic Olefins:

In prior applications of olefin metathesis, particularly olefin cross-metathesis, there has been no method available for generation of highly substituted olefins, such as trisubstituted olefins (wherein the substituents may be the same or different) and olefins that contain quaternary carbons at the allylic position. Trisubstituted and quaternary allylic olefinic substituents are, of course, present in a diverse array of organic molecules, including pharmaceuticals, natural products, and functionalized polymers, and the difficulty in generating such compounds has been a substantial limitation. The methodology of the present invention overcomes this limitation and now provides an efficient and versatile way to synthesize 1,1,2-trisubstituted olefins as well as 1,2-disubstituted olefins containing one quaternary allylic carbon atom.

Stereoselective Synthesis of 1,2-Disubstituted Olefins via Cross-metathesis: Another limitation in known olefin metathesis reactions is that there is no general method for controlling the stereoselectivity of the newly formed olefins. In many cases, the more thermodynamically stable *trans* olefin geometry was selectively formed, with minimal, if any, of the *cis* olefin produced. See Blackwell et al. (2000), "New approaches to olefin cross-metathesis," *J. Am. Chem. Soc.* 122(1):58-71; and Chatterjee et al. (2000), "Synthesis of functionalized olefins by cross and ring-closing metathesis," *J. Am. Chem. Soc.* 122(15):3783-3784. The present invention also addresses this need in the art by providing a stereoselective method for synthesizing a 1,2-disubstituted olefin in primarily the *cis* configuration.

SUMMARY OF THE INVENTION

The present invention is addressed to the aforementioned needs in the art, and provides a novel process for using certain Group 8 transition metal complexes to catalyze a variety of olefin metathesis reactions, primarily cross-metathesis reactions. The complexes used are metal carbenes comprised of a Group 8 transition metal, particularly ruthenium or osmium, which preferably, although not necessarily, contain an N-heterocyclic carbene ligand. Such complexes are highly active catalysts of olefin metathesis reactions, including the cross-metathesis reactions described in detail herein. In contrast to previous catalysts used in olefin cross-metathesis, the present complexes allow an olefinic reactant to be substituted with a functional group without compromising the efficiency or selectivity of a metathesis reaction involving that olefin. The present invention also allows the second reactant, i.e., the olefin metathesis partner, to be substituted with a functional group. The functional group may or may not be a ligand for the metal complex; the present method is not limited in this regard. The olefinic reactant may also be di-substituted at one of the olefinic carbon atoms, as is the case with 2-methyl-2-butene, for example, or may be a quaternary allylic olefin, i.e., an olefin directly substituted at one or both of the olefinic carbon atoms with the moiety $-\text{CH}_2-\text{CR}_3$ where R is other than hydrogen.

These cross-metathesis reactions are carried out with a catalyst having the structure of formula (VI)



in which:

M is a Group 8 transition metal, particularly Ru or Os;


X¹ and X² may be the same or different, and are anionic ligands or polymers;

R^1 is selected from the group consisting of hydrogen, hydrocarbyl, substituted hydrocarbyl, heteroatom-containing hydrocarbyl, substituted heteroatom-containing hydrocarbyl, and carboxyl;

R^2 is selected from the group consisting of hydrogen, hydrocarbyl, substituted hydrocarbyl, heteroatom-containing hydrocarbyl, and substituted heteroatom-containing hydrocarbyl;

L is a neutral electron donor ligand; and

L^1 is a neutral electron donor ligand having the structure of formula (VII)



In structure (VII):

X and Y are heteroatoms selected from N, O, S, and P;

p is zero when X is O or S, and is 1 when X is N or P;

q is zero when Y is O or S, and is 1 when Y is N or P;


Q^1 , Q^2 , Q^3 , and Q^4 are linkers, e.g., hydrocarbylene (including substituted hydrocarbylene, heteroatom-containing hydrocarbylene, and substituted heteroatom-containing hydrocarbylene, such as substituted and/or heteroatom-containing alkylene) or $-(CO)-$;

w , x , y and z are independently zero or 1; and


R^3 , R^{3A} , R^4 , and R^{4A} are independently selected from the group consisting of hydrogen, hydrocarbyl, substituted hydrocarbyl, heteroatom-containing hydrocarbyl, and substituted heteroatom-containing hydrocarbyl,

wherein any two or more of X^1 , X^2 , L , R^1 , R^2 , R^3 , R^{3A} , R^4 , and R^{4A} can be taken together to form a chelating multidentate ligand.


Accordingly, the complex of formula (V) may also be represented as (VIA)



In a preferred embodiment, L is an N-heterocyclic carbene having the structure of formula (VIIA)



wherein R³ and R⁴ are defined above, with preferably at least one of R³ and R⁴, and more preferably both R³ and R⁴, being alicyclic or aromatic of one to about five rings, and optionally containing one or more heteroatoms and/or substituents. Q is a linker, typically a hydrocarbylene linker, including substituted hydrocarbylene, heteroatom-containing hydrocarbylene, and substituted heteroatom-containing hydrocarbylene linkers, wherein two or more substituents on adjacent atoms within Q may also be linked to form an additional cyclic structure, which may be similarly substituted to provide a fused polycyclic structure of two to about five cyclic groups. Q is often, although again not necessarily, a two-atom linkage or a three-atom linkage. Accordingly, the metal carbene complex of formula (VIA) may also be represented as follows:



In one embodiment, then, a method is provided for synthesizing olefins substituted with a functional group by cross-metathesis using a Group 8 transition metal catalyst having the structure of formula (VI). At least one of the two olefinic reactants is substituted with one or more functional groups, which may or may not be in protected form (e.g., a hydroxyl group may be protected as an acyloxy or benzyloxy group). More specifically, at least one of the two olefinic reactants has the structure of formula (VIII)



wherein:

Fn is a functional group such as phosphonato, phosphoryl, phosphanyl, phosphino, sulfonato, C₁-C₂₀ alkylsulfanyl, C₅-C₂₀ arylsulfanyl, C₁-C₂₀ alkylsulfonyl, C₅-C₂₀ arylsulfonyl, C₁-

C₂₀ alkylsulfinyl, C₅-C₂₀ arylsulfinyl, sulfonamido, amino, amido, imino, nitro, nitroso, hydroxyl, C₁-C₂₀ alkoxy, C₅-C₂₀ aryloxy, C₂-C₂₀ alkoxy carbonyl, C₅-C₂₀ aryloxy carbonyl, carboxyl, carboxylato, mercapto, formyl, C₁-C₂₀ thioester, cyano, cyanato, carbamoyl, epoxy, styrenyl, silyl, silyloxy, silanyl, siloxazanyl, boronato, or boryl, or a metal-containing or metalloid-containing group (wherein the metal may be, for example, Sn or Ge);

n is zero or 1;


Z is a hydrocarbylene or a substituted and/or heteroatom-containing hydrocarbylene linking group such as an alkylene, substituted alkylene, heteroalkylene, substituted heteroalkene, arylene, substituted arylene, heteroarylene, or substituted heteroarylene linkage; and

R⁵, R⁶, and R⁷ are independently selected from the group consisting of hydrogen, -(Z)_n-Fn, hydrocarbyl, substituted hydrocarbyl, heteroatom-containing hydrocarbyl, and substituted heteroatom-containing hydrocarbyl, and, if substituted hydrocarbyl or substituted heteroatom-containing hydrocarbyl, wherein one or more of the substituents may be -(Z)_n-Fn.

In one preferred embodiment, Fn is a phosphonate and Z is CH₂, such that the reactant is an allylphosphonate (when n is 1) and a vinylphosphonate (when n is zero). The product of the cross-metathesis reaction is also an olefin substituted with a -(Z)_n-Fn group.

In another embodiment, a method is provided for synthesizing directly halogenated olefins by cross-metathesis using a catalyst having the structure of formula (VI). In this embodiment, at least one of the olefinic reactants has the structure of formula (IX)


(IX)




wherein X³ is halo, and R⁸, R⁹, and R¹⁰ are independently selected from the group consisting of hydrogen, halo, hydrocarbyl, substituted hydrocarbyl, heteroatom-containing hydrocarbyl, substituted heteroatom-containing hydrocarbyl, and -(Z)_n-Fn where n, Z and Fn are as defined above.

In a further embodiment, a method is provided for synthesizing substituted olefins, particularly trisubstituted and quaternary allylic olefins, wherein the method comprises using the complex of formula (VI) to catalyze a cross-metathesis reaction between a geminal disubstituted olefin or a quaternary allylic olefin, and a second olefin. If it is a geminal disubstituted olefin, the first olefin has the structure (X)

(X)



wherein R^{11} , R^{12} , R^{13} , and R^{14} are selected from the group consisting of hydrogen, hydrocarbyl, substituted hydrocarbyl, heteroatom-containing hydrocarbyl, substituted heteroatom-containing hydrocarbyl, and $-(Z)_n$ -Fn where n , Z and F_n are as defined above, with the proviso that R^{11} and R^{12} , or R^{13} and R^{14} , are other than hydrogen. If it is a quaternary allylic olefin, the first olefin has the structure (XI)



wherein R^{11} and R^{12} are as defined previously, and R^{15} , R^{16} , and R^{17} are any nonhydrogen substituents, e.g., alkyl, aryl, heteroalkyl, heteroaryl, $-(Z)_n$ -Fn (where n , Z , and F_n are as defined above with respect to formula (VIII)), or the like.

In the above-described embodiments, the second olefin has a molecular structure given by $R^{18}R^{19}C=CR^{20}R^{21}$ wherein R^{18} , R^{19} , R^{20} , and R^{21} may be hydrogen, hydrocarbyl, substituted hydrocarbyl, heteroatom-containing hydrocarbyl, substituted heteroatom-containing hydrocarbyl, $-(Z)_n$ -Fn, etc. As such, it will be appreciated that the second olefin may have a molecular structure encompassed by any one of the above generic formulae (VIII), (IX), (X), and (XI), or may be a simple structure such as ethylene *per se*.


The invention is additionally useful in providing a method for controlling the stereoselectivity of an olefin cross-metathesis reaction, and in providing a cross-metathesis product in which the thermodynamically less favored *cis* configuration predominates. The reaction is carried out using selected olefinic reactants, with one olefinic reactant substituted in a 1,2-*cis* configuration. The catalyst used has the structure of formula (VI), with R^3 and R^4 representing bulky ligands, e.g., bicyclic or polycyclic ligands that may or may not be aromatic.

In a still further embodiment of the invention, complexes of formula (VI) are used to catalyze a plurality of cross-metathesis reactions from a common olefinic reactant to generate chemical diversity, i.e., to provide a plurality of products having related structures but retaining a distinguishing feature, such that each synthesized compound is different from each other synthesized compound. Each olefinic reactant can be substituted with functional groups, yielding cross-metathesis products containing those groups, and thus providing the option of further derivatization. While prior olefin cross-metathesis reactions have been used to synthesize alkenes bearing a range of functional groups, these prior reactions have been limited to olefins that do not

-11-

contain any functional groups that could behave as ligands for the catalyst employed. By contrast, the present method can be used with olefinic starting materials in which functional groups are present that could act as ligands for the metal complex selected as a metathesis catalyst.

According to an aspect of the invention, there is provided a method for synthesizing a functionalized olefin via a cross-metathesis reaction, comprising contacting (a) a first olefinic reactant directly substituted with a functional group Fn selected from the group consisting of: phosphonato; phosphoryl; phosphanyl; phosphino; sulfonato; C₁-C₂₀ alkylsulfanyl; C₅-C₂₀ arylsulfanyl; C₁-C₂₀ alkylsulfinyl; C₅-C₂₀ arylsulfinyl; sulfonamido; amino; imino; nitro; nitroso; hydroxyl; unsubstituted C₁-C₂₀ alkoxy; substituted C₂-C₂₀ alkoxy; C₅-C₂₀ aryloxy; C₅-C₂₀ aryloxycarbonyl; mercapto; C₁-C₂₀ thioester; cyano; cyanato; carbamoyl; styrenyl; siloxazanyl; boronato; boryl; stannyl; and germyl, or indirectly substituted with a functional group Fn selected from the group consisting of: phosphonato; phosphoryl; phosphanyl; phosphino; sulfonato; C₁-C₂₀ alkylsulfanyl; C₅-C₂₀ arylsulfanyl; C₁-C₂₀ alkylsulfinyl; C₅-C₂₀ arylsulfinyl; sulfonamido; amino; imino; nitro; nitroso; hydroxyl; unsubstituted C₁-C₂₀ alkoxy; substituted C₄-C₂₀ alkoxy; C₅-C₂₀ aryloxy; C₅-C₂₀ aryloxycarbonyl; carboxylato; mercapto; C₁-C₂₀ thioester; cyano; cyanato; carbamoyl; styrenyl; siloxazanyl; boronato; boryl; stannyl; and germyl, with (b) a second olefinic reactant in the presence of (c) a catalyst composed of a Group 8 transition metal alkylidene complex under conditions and for a time period effective to allow cross-metathesis to occur, wherein the catalyst has the structure of formula (VIA)



in which:

M is a Group 8 transition metal;

X¹ and X² may be the same or different, and are anionic ligands or polymers;

R¹ is selected from the group consisting of: hydrogen; hydrocarbyl; substituted hydrocarbyl; heteroatom-containing hydrocarbyl; substituted heteroatom-containing hydrocarbyl; and carboxyl;

R² is selected from the group consisting of: hydrogen; hydrocarbyl; substituted hydrocarbyl; heteroatom-containing hydrocarbyl; and substituted heteroatom-containing hydrocarbyl;

L is a neutral electron donor ligand;

X and Y are heteroatoms selected from the group consisting of: N; O; S; and P;

-11a-

p is zero when X is O or S, and is 1 when X is N or P;

q is zero when Y is O or S, and is 1 when Y is N or P;

Q¹, Q², Q³, and Q⁴ are selected from the group consisting of: hydrocarbylene; substituted hydrocarbylene; heteroatom-containing hydrocarbylene; substituted heteroatom-containing hydrocarbylene; and -(CO)-;


w, x, y and z are independently zero or 1; and

R³, R^{3A}, R⁴, and R^{4A} are independently selected from the group consisting of hydrogen, hydrocarbyl, substituted hydrocarbyl, heteroatom-containing hydrocarbyl, and substituted heteroatom-containing hydrocarbyl,

wherein any two or more of X¹, X², L, R¹, R², R³, R^{3A}, R⁴, and R^{4A} can be taken together to form a chelating multidentate ligand,

wherein the first olefinic reactant has the structure of formula (VIII)

(VIII)



wherein:

n is zero or 1;

Z is a hydrocarbylene or a substituted or heteroatom-containing hydrocarbylene linking group; and

R⁵, R⁶ and R⁷ are independently selected from the group consisting of hydrogen, hydrocarbyl, substituted hydrocarbyl, heteroatom-containing hydrocarbyl, substituted heteroatom-containing hydrocarbyl, and -(Z)_n-Fn,


wherein 'substituted' indicates that in the "substituted" compound, at least one hydrogen atom bound to a carbon (or other) atom is replaced with one or more non-hydrogen substituents, and

wherein 'heteroatom' is selected from the group consisting of: N, O, S, P, and Si.

According to another aspect of the invention, there is provided a method for synthesizing a plurality of functionalized olefins from a common olefinic reactant via a cross-metathesis reaction, comprising:

(a) contacting an olefinic substrate with a first olefinic reactant in the presence of a catalyst composed of a Group 8 transition metal alkylidene complex under conditions and for a time period effective to allow cross-metathesis to occur, wherein the olefinic substrate is substituted with at least one -(Z)_n-Fn moiety in which n is zero or 1, Z is a

wherein the first olefinic reactant has the structure of formula (VIII)



-11c-

M is a Group 8 transition metal;

X^1 and X^2 may be the same or different, and are anionic ligands or polymers;

R^1 is selected from the group consisting of: hydrogen; hydrocarbyl; substituted hydrocarbyl;
5 heteroatom-containing hydrocarbyl; substituted heteroatom-containing hydrocarbyl; and carboxyl;

R^2 is selected from the group consisting of: hydrogen; hydrocarbyl; substituted hydrocarbyl;
heteroatom-containing hydrocarbyl; and substituted heteroatom-containing hydrocarbyl;

L is a neutral electron donor ligand;

X and Y are heteroatoms selected from the group consisting of: N; O; S; and P;

10 p is zero when X is O or S, and is 1 when X is N or P;

q is zero when Y is O or S, and is 1 when Y is N or P;

Q^1 , Q^2 , Q^3 , and Q^4 are selected from the group consisting of: hydrocarbylene; substituted
hydrocarbylene; heteroatom-containing hydrocarbylene; substituted heteroatom-containing
hydrocarbylene; and $-(CO)-$;

15 w, x, y and z are independently zero or 1; and


R^3 , R^{3A} , R^4 , and R^{4A} are independently selected from the group consisting of hydrogen,
hydrocarbyl, substituted hydrocarbyl, heteroatom-containing hydrocarbyl, and substituted
heteroatom-containing hydrocarbyl,

20 wherein any two or more of X^1 , X^2 , L, R^1 , R^2 , R^3 , R^{3A} , R^4 , and R^{4A} can be taken together to
form a chelating multidentate ligand,

The second olefinic reactant may also be substituted with at least one $-(Z)_n$ -Fn moiety.


According to another aspect of the invention, there is provided a method for synthesizing a
functionalized olefin of the formula

25



30

via a cross-metathesis reaction, comprising contacting (a) a first olefinic reactant of the formula



with (b) a second olefinic reactant of the formula


40



-11d-

in the presence of (c) a catalyst composed of a Group 8 transition metal alkylidene complex under conditions and for a time period effective to allow cross-metathesis to occur, wherein the catalyst has the structure of formula (VIA)

5



in which:

M is a Group 8 transition metal;

15 X^1 and X^2 may be the same or different, and are anionic ligands or polymers;

R^1 is selected from the group consisting of hydrogen, hydrocarbyl, substituted hydrocarbyl, heteroatom-containing hydrocarbyl, substituted heteroatom-containing hydrocarbyl, and carboxyl;

R^2 is selected from the group consisting of hydrogen, hydrocarbyl, substituted hydrocarbyl, heteroatom-containing hydrocarbyl, and substituted heteroatom-containing hydrocarbyl;

20 L is a neutral electron donor ligand;

X and Y are heteroatoms selected from the group consisting of N, O, S, and P;

p is zero when X is O or S, and is 1 when X is N or P;

q is zero when Y is O or S, and is 1 when Y is N or P;

25 Q^1 , Q^2 , Q^3 , and Q^4 are selected from hydrocarbylene, substituted hydrocarbylene, heteroatom-containing hydrocarbylene, substituted heteroatom-containing hydrocarbylene, and $-(CO)-$;

w, x, y and z are independently zero or 1; and

30 R^3 , R^{3A} , R^4 , and R^{4A} are independently selected from the group consisting of hydrogen, hydrocarbyl, substituted hydrocarbyl, heteroatom-containing hydrocarbyl, and substituted heteroatom-containing hydrocarbyl,

wherein any two or more of X^1 , X^2 , L, R^1 , R^2 , R^3 , R^{3A} , R^4 , and R^{4A} can be taken together to form a chelating multidentate ligand, and

wherein 'substituted' indicates that in the "substituted" compound, at least one hydrogen atom bound to a carbon (or other) atom is replaced with one or more non-hydrogen substituents,

35 and

-11e-

wherein 'heteroatom' is selected from the group consisting of: N, O, S, P, and Si.

DETAILED DESCRIPTION OF THE INVENTION

I. DEFINITIONS AND NOMENCLATURE:

It is to be understood that unless otherwise indicated this invention is not limited to specific reactants, reaction conditions, ligands, metal complexes, or the like, as such may vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only and is not intended to be limiting.

As used in the specification and the appended claims, the singular forms "a," "an" and "the" include plural referents unless the context clearly dictates otherwise. Thus, for example, reference to "a compound" encompasses a combination or mixture of different compounds as well as a single compound, reference to "a functional group" includes a single functional group as well as two or more functional groups that may or may not be the same, and the like.

In this specification and in the claims that follow, reference will be made to a number of terms, which shall be defined to have the following meanings:

As used herein, the phrase "having the formula" or "having the structure" is not intended to be limiting and is used in the same way that the term "comprising" is commonly used.

The term "alkyl" as used herein refers to a linear, branched or cyclic saturated hydrocarbon group typically although not necessarily containing 1 to about 20 carbon atoms, such as methyl, ethyl, *n*-propyl, isopropyl, *n*-butyl, isobutyl, *t*-butyl, octyl, decyl, and the like, as well as cycloalkyl groups such as cyclopentyl, cyclohexyl and the like. Generally, although again not necessarily, alkyl groups herein contain 1 to about 12 carbon atoms. The term "lower alkyl" intends an alkyl group of 1 to 6 carbon atoms, and the specific term "cycloalkyl" intends a cyclic alkyl group, typically having 4 to 8, preferably 5 to 7, carbon atoms. The term "substituted alkyl" refers to alkyl substituted with one or more substituent groups, and the terms "heteroatom-containing alkyl" and "heteroalkyl" refer to alkyl in which at least one carbon atom is replaced with a heteroatom. If not otherwise indicated, the terms "alkyl" and "lower alkyl" include linear, branched, cyclic, unsubstituted, substituted, and/or heteroatom-containing alkyl and lower alkyl, respectively.

The term "alkylene" as used herein refers to a difunctional linear, branched or cyclic alkyl group, where "alkyl" is as defined above.

The term "alkenyl" as used herein refers to a linear, branched or cyclic hydrocarbon group of 2 to 20 carbon atoms containing at least one double bond, such as ethenyl, *n*-propenyl,

isopropenyl, *n*-butenyl, isobutenyl, octenyl, decenyl, tetradecenyl, hexadecenyl, eicosenyl, tetracosenyl, and the like. Preferred alkenyl groups herein contain 2 to 12 carbon atoms. The term "lower alkenyl" intends an alkenyl group of 2 to 6 carbon atoms, and the specific term "cycloalkenyl" intends a cyclic alkenyl group, preferably having 5 to 8 carbon atoms. The term "substituted alkenyl" refers to alkenyl substituted with one or more substituent groups, and the terms "heteroatom-containing alkenyl" and "heteroalkenyl" refer to alkenyl in which at least one carbon atom is replaced with a heteroatom. If not otherwise indicated, the terms "alkenyl" and "lower alkenyl" include linear, branched, cyclic, unsubstituted, substituted, and/or heteroatom-containing alkenyl and lower alkenyl, respectively.

The term "alkenylene" as used herein refers to a difunctional linear, branched or cyclic alkenyl group, where "alkenyl" is as defined above.

The term "alkynyl" as used herein refers to a linear or branched hydrocarbon group of 2 to 20 carbon atoms containing at least one triple bond, such as ethynyl, *n*-propynyl, and the like. Preferred alkynyl groups herein contain 2 to 12 carbon atoms. The term "lower alkynyl" intends an alkynyl group of 2 to 6 carbon atoms. The term "substituted alkynyl" refers to alkynyl substituted with one or more substituent groups, and the terms "heteroatom-containing alkynyl" and "heteroalkynyl" refer to alkynyl in which at least one carbon atom is replaced with a heteroatom. If not otherwise indicated, the terms "alkynyl" and "lower alkynyl" include linear, branched, unsubstituted, substituted, and/or heteroatom-containing alkynyl and lower alkynyl, respectively.

The term "alkoxy" as used herein intends an alkyl group bound through a single, terminal ether linkage; that is, an "alkoxy" group may be represented as -O-alkyl where alkyl is as defined above. A "lower alkoxy" group intends an alkoxy group containing 1 to 6 carbon atoms. Analogously, "alkenyloxy" and "lower alkenyloxy" respectively refer to an alkenyl and lower alkenyl group bound through a single, terminal ether linkage, and "alkynyloxy" and "lower alkynyloxy" respectively refer to an alkynyl and lower alkynyl group bound through a single, terminal ether linkage.

The term "aryl" as used herein, and unless otherwise specified, refers to an aromatic substituent containing a single aromatic ring or multiple aromatic rings that are fused together, directly linked, or indirectly linked (such that the different aromatic rings are bound to a common group such as a methylene or ethylene moiety). Preferred aryl groups contain one aromatic ring or 2 to 4 fused or linked aromatic rings, e.g., phenyl, naphthyl, biphenyl, and the like. "Substituted aryl" refers to an aryl moiety substituted with one or more substituent groups, and the terms "heteroatom-containing aryl" and "heteroaryl" refer to aryl in which at least one carbon atom is replaced with a heteroatom. Unless otherwise indicated, the terms "aromatic," "aryl," and "arylene" include heteroaromatic, substituted aromatic, and substituted heteroaromatic species.

The term "aryloxy" as used herein refers to an aryl group bound through a single, terminal ether linkage. An "aryloxy" group may be represented as -O-aryl where aryl is as defined above.

The term "aralkyl" refers to an alkyl group with an aryl substituent, and the term "aralkylene" refers to an alkylene group with an aryl substituent; the term "alkaryl" refers to an aryl group that has an alkyl substituent, and the term "alkarylene" refers to an arylene group with an alkyl substituent.

The term "alicyclic" refers to an aliphatic cyclic moiety, which may or may not be bicyclic or polycyclic.

The terms "halo" and "halogen" are used in the conventional sense to refer to a chloro, bromo, fluoro or iodo substituent. The terms "haloalkyl," "haloalkenyl" or "haloalkynyl" (or "halogenated alkyl," "halogenated alkenyl," or "halogenated alkynyl") refers to an alkyl, alkenyl or alkynyl group, respectively, in which at least one of the hydrogen atoms in the group has been replaced with a halogen atom.

"Hydrocarbyl" refers to univalent hydrocarbyl radicals containing 1 to about 30 carbon atoms, preferably 1 to about 20 carbon atoms, most preferably 1 to about 12 carbon atoms, including linear, branched, cyclic, saturated and unsaturated species, such as alkyl groups, alkenyl groups, aryl groups, and the like. The term "lower hydrocarbyl" intends a hydrocarbyl group of 1 to 6 carbon atoms, and the term "hydrocarbylene" intends a divalent hydrocarbyl moiety containing 1 to about 30 carbon atoms, preferably 1 to about 20 carbon atoms, most preferably 1 to about 12 carbon atoms, including linear, branched, cyclic, saturated and unsaturated species. The term "lower hydrocarbylene" intends a hydrocarbylene group of 1 to 6 carbon atoms. "Substituted hydrocarbyl" refers to hydrocarbyl substituted with one or more substituent groups, and the terms "heteroatom-containing hydrocarbyl" and "heterohydrocarbyl" refer to hydrocarbyl in which at least one carbon atom is replaced with a heteroatom. Similarly, "substituted hydrocarbylene" refers to hydrocarbylene substituted with one or more substituent groups, and the terms "heteroatom-containing hydrocarbylene" and "heterohydrocarbylene" refer to hydrocarbylene in which at least one carbon atom is replaced with a heteroatom. Unless otherwise indicated, the term "hydrocarbyl" and "hydrocarbylene" are to be interpreted as including substituted and/or heteroatom-containing hydrocarbyl and hydrocarbylene moieties, respectively.

The term "heteroatom-containing" as in a "heteroatom-containing hydrocarbyl group" refers to a hydrocarbon molecule or a hydrocarbyl molecular fragment in which one or more carbon atoms is replaced with an atom other than carbon, e.g., nitrogen, oxygen, sulfur, phosphorus or silicon, typically nitrogen, oxygen or sulfur. Similarly, the term "heteroalkyl" refers to an alkyl substituent that is heteroatom-containing, the term "heterocyclic" refers to a cyclic substituent that is heteroatom-containing, the terms "heteroaryl" and "heteroaromatic"

respectively refer to "aryl" and "aromatic" substituents that are heteroatom-containing, and the like. It should be noted that a "heterocyclic" group or compound may or may not be aromatic, and further that "heterocycles" may be monocyclic, bicyclic, or polycyclic as described above with respect to the term "aryl."

5 By "substituted" as in "substituted hydrocarbyl," "substituted alkyl," "substituted aryl," and the like, as alluded to in some of the aforementioned definitions, is meant that in the hydrocarbyl, alkyl, aryl, or other moiety at least one hydrogen atom bound to a carbon (or other) atom is replaced with one or more non-hydrogen substituents. Examples of such substituents include, without limitation: functional groups such as halogen, phosphonato, phosphoryl,
10 phosphanyl, phosphino, sulfonato, C₁-C₂₀ alkylsulfanyl, C₅-C₂₀ arylsulfanyl, C₁-C₂₀ alkylsulfonyl, C₅-C₂₀ arylsulfonyl, C₁-C₂₀ alkylsulfinyl, C₅-C₂₀ arylsulfinyl, sulfonamido, amino, amido, imino, nitro, nitroso, hydroxyl, C₁-C₂₀ alkoxy, C₅-C₂₀ aryloxy, C₂-C₂₀ alkoxycarbonyl, C₅-C₂₀ aryloxcarbonyl, carboxyl, carboxylato, mercapto, formyl, C₁-C₂₀ thioester, cyano, cyanato, carbamoyl, epoxy, styrenyl, silyl, silyloxy, silanyl, siloxazanyl, boronato, or boryl, or a metal-
15 containing or metalloid-containing group (wherein the metal may be, for example, Sn or Ge); and the hydrocarbyl moieties C₁-C₂₀ alkyl, C₂-C₂₀ alkenyl, C₂-C₂₀ alkynyl, C₅-C₂₀ aryl, C₅-C₃₀ aralkyl, and C₅-C₃₀ alkaryl.

In addition, the aforementioned functional groups may, if a particular group permits, be further substituted with one or more additional functional groups or with one or more hydrocarbyl
20 moieties such as those specifically enumerated above. Analogously, the above-mentioned hydrocarbyl moieties may be further substituted with one or more functional groups or additional hydrocarbyl moieties such as those specifically enumerated.

When the term "substituted" appears prior to a list of possible substituted groups, it is intended that the term apply to every member of that group. That is, the phrase "substituted alkyl,
25 alkenyl and alkynyl" is to be interpreted as "substituted alkyl, substituted alkenyl and substituted alkynyl." Similarly, "optionally substituted alkyl, alkenyl and alkynyl" is to be interpreted as "optionally substituted alkyl, optionally substituted alkenyl and optionally substituted alkynyl."

The term "amino" is used herein to refer to the group -NZ¹Z², where each of Z¹ and Z² is independently selected from the group consisting of hydrogen and optionally substituted alkyl,
30 alkenyl, alkynyl, aryl, aralkyl, alkaryl and heterocyclic.

The term "stereoselective" refers to a chemical reaction that preferentially results in one stereoisomer relative to a second stereoisomer, i.e., gives rise to a product of which the ratio of a desired stereoisomer to a less desired stereoisomer is greater than 1:1.


"Optional" or "optionally" means that the subsequently described circumstance may or
35 may not occur, so that the description includes instances where the circumstance occurs and instances where it does not. For example, the phrase "optionally substituted" means that a non-

hydrogen substituent may or may not be present on a given atom, and, thus, the description includes structures wherein a non-hydrogen substituent is present and structures wherein a non-hydrogen substituent is not present.

In the molecular structures herein, the use of bold and dashed lines to denote particular conformation of groups follows the IUPAC convention. A bond indicated by a broken line indicates that the group in question is below the general plane of the molecule as drawn (the " α " configuration), and a bond indicated by a bold line indicates that the group at the position in question is above the general plane of the molecule as drawn (the " β " configuration).

II. THE CATALYST:

The cross-metathesis reactions of the invention are carried out catalytically, using a Group 8 transition metal complex that preferably contains two different ligands. These transition metal carbene complexes include a metal center in a +2 oxidation state, have an electron count of 16, and are penta-coordinated. More specifically, the preferred catalysts herein have the structure of formula (VIA)



wherein the various substituents are as follows:

M, which serves as the transition metal center in the +2 oxidation state, is a Group 8 transition metal, particularly ruthenium or osmium. In a preferred embodiment, M is ruthenium.

X¹ and X² are anionic ligands or polymers, and may be the same or different, or are linked together to form a cyclic group, typically although not necessarily a five- to eight-membered ring.

In preferred embodiments, X¹ and X² are each independently hydrogen, halide, or one of the

following groups: C₁-C₂₀ alkyl, C₅-C₂₀ aryl, C₁-C₂₀ alkoxy, C₅-C₂₀ aryloxy, C₃-C₂₀ alkylidicketonate, C₅-C₂₀ arylidicketonate, C₂-C₂₀ alkoxycarbonyl, C₅-C₂₀ aryloxy carbonyl, C₂-C₂₀ acyl, C₁-C₂₀ alkylsulfonato, C₅-C₂₀ arylsulfonato, C₁-C₂₀ alkylsulfanyl, C₅-C₂₀ arylsulfanyl, C₁-C₂₀ alkylsulfinyl, or C₅-C₂₀ arylsulfinyl. Optionally, X¹ and X² may be substituted with one or more moieties selected from the group consisting of C₁-C₁₀ alkyl, C₁-C₁₀ alkoxy, aryl, and halide, which may, in turn, with the exception of halide, be further substituted with one or more groups selected from halide, C₁-C₆ alkyl, C₁-C₆ alkoxy, and phenyl. In more preferred embodiments, X¹ and X² are

halide, benzoate, C₂-C₆ acyl, C₂-C₆ alkoxy carbonyl, C₁-C₆ alkyl, phenoxy, C₁-C₆ alkoxy, C₁-C₆ alkylsulfanyl, aryl, or C₁-C₆ alkylsulfonyl. In even more preferred embodiments, X¹ and X² are each halide, CF₃CO₂, CH₃CO₂, CFH₂CO₂, (CH₃)₃CO, (CF₃)₂(CH₃)CO, (CF₃)(CH₃)₂CO, PhO, MeO, EtO, tosylate, mesylate, or trifluoromethanesulfonate. In the most preferred embodiments, X¹ and X² are each chloride. The complex may also be attached to a solid support, such as to a polymeric substrate, and this attachment may be effected by means of X¹ and/or X², in which case X¹ and/or X² is a polymer.


R¹ is selected from the group consisting of hydrogen, hydrocarbyl (e.g., alkyl, alkenyl, alkynyl, aryl, aralkyl, alkaryl, etc.), substituted hydrocarbyl (e.g., substituted alkyl, alkenyl, alkynyl, aryl, aralkyl, alkaryl, etc.), heteroatom-containing hydrocarbyl (e.g., heteroatom-containing alkyl, alkenyl, alkynyl, aryl, aralkyl, alkaryl, etc.), and substituted heteroatom-containing hydrocarbyl (e.g., substituted heteroatom-containing alkyl, alkenyl, alkynyl, aryl, aralkyl, alkaryl, etc.), and carboxyl, and R² is selected from the group consisting of hydrogen, hydrocarbyl (e.g., alkyl, alkenyl, alkynyl, aryl, aralkyl, alkaryl, etc.), substituted hydrocarbyl (e.g., substituted alkyl, alkenyl, alkynyl, aryl, aralkyl, alkaryl, etc.), heteroatom-containing hydrocarbyl (e.g., heteroatom-containing alkyl, alkenyl, alkynyl, aryl, aralkyl, alkaryl, etc.), and substituted heteroatom-containing hydrocarbyl (e.g., substituted heteroatom-containing alkyl, alkenyl, alkynyl, aryl, aralkyl, alkaryl, etc.). R¹ and R² may also be linked to form a cyclic group, which may be aliphatic or aromatic, and may contain substituents and/or heteroatoms. Generally, such a cyclic group will contain 4 to 12, preferably 5 to 8, ring atoms.

In preferred catalysts, the R¹ substituent is hydrogen and the R² substituent is selected from the group consisting of C₁-C₂₀ alkyl, C₂-C₂₀ alkenyl, and aryl. More preferably, R² is phenyl, vinyl, methyl, isopropyl, or t-butyl, optionally substituted with one or more moieties selected from the group consisting of C₁-C₆ alkyl, C₁-C₆ alkoxy, phenyl, and a functional group Fn. Still more preferably, R² is phenyl or vinyl substituted with one or more moieties selected from the group consisting of methyl, ethyl, chloro, bromo, iodo fluoro, nitro, dimethylamino, methyl, methoxy, and phenyl. In the most preferred embodiments, the R² substituent is phenyl or -C=C(CH₃)₂.

L is a neutral electron donor ligand, and may or may not be linked to R². Examples of suitable L moieties include, without limitation, phosphine, sulfonated phosphine, phosphite, phosphinite, phosphonite, arsine, stibine, ether (including cyclic ethers), amine, amide, imine, sulfoxide, carboxyl, nitrosyl, pyridine, substituted pyridine (e.g., halogenated pyridine), imidazole, substituted imidazole (e.g., halogenated imidazole), pyrazine (e.g., substituted pyrazine), and thioether. In more preferred embodiments, L is a phosphine of the formula PR⁵R⁶R⁷, where R⁵, R⁶, and R⁷ are each independently aryl or C₁-C₁₀ alkyl, particularly primary

alkyl, secondary alkyl or cycloalkyl. In the most preferred embodiments, L is selected from the group consisting of -P(cyclohexyl)₃, -P(cyclopentyl)₃, -P(isopropyl)₃, -P(phenyl)₃, -P(phenyl)₂(R⁷) and -P(phenyl)(R⁷)₂, in which R⁷ is alkyl, typically lower alkyl. Also preferred are weaker ligands such as the nitrogen-containing heterocycles, which enhance catalytic activity presumably because of the requirement that the L ligand dissociate for initiation to occur.

Examples of complexes wherein L and R² are linked include the following:



X and Y are heteroatoms typically selected from N, O, S, and P. Since O and S are divalent, p is necessarily zero when X is O or S, and q is necessarily zero when Y is O or S. However, when X is N or P, then p is 1, and when Y is N or P, then q is 1. In a preferred embodiment, both X and Y are N.

Q¹, Q², Q³, and Q⁴ are linkers, e.g., hydrocarbylene (including substituted hydrocarbylene, heteroatom-containing hydrocarbylene, and substituted heteroatom-containing hydrocarbylene, such as substituted and/or heteroatom-containing alkylene) or -(CO)-, and w, x, y and z are independently zero or 1, meaning that each linker is optional. Preferably, w, x, y and z are all zero.


R³, R^{3A}, R⁴, and R^{4A} are independently selected from the group consisting of hydrogen, hydrocarbyl, substituted hydrocarbyl, heteroatom-containing hydrocarbyl, and substituted heteroatom-containing hydrocarbyl, wherein R^{3A} and R^{4A} may be linked to form a cyclic group.

It should be emphasized that any two or more (typically two, three or four) of X¹, X², L, R¹, R², R³, R^{3A}, R⁴, and R^{4A} can be taken together to form a chelating multidentate ligand, as disclosed, for example, in U.S. Patent No. 5,312,940 to Grubbs et al. Examples of bidentate

ligands include, but are not limited to, bisphosphines, dialkoxides, alkylidicketonates, and arylidicketonates. Specific examples include $-\text{P}(\text{Ph})_2\text{CH}_2\text{CH}_2\text{P}(\text{Ph})_2-$, $-\text{As}(\text{Ph})_2\text{CH}_2\text{CH}_2\text{As}(\text{Ph})_2-$, $-\text{P}(\text{Ph})_2\text{CH}_2\text{CH}_2\text{C}(\text{CF}_3)_2\text{O}-$, binaphtholate dianions, pinacolate dianions, $-\text{P}(\text{CH}_3)_2(\text{CH}_2)_2\text{P}(\text{CH}_3)_2-$ and $-\text{OC}(\text{CH}_3)_2(\text{CH}_3)_2\text{CO}-$. Preferred bidentate ligands are $-\text{P}(\text{Ph})_2\text{CH}_2\text{CH}_2\text{P}(\text{Ph})_2-$ and $-\text{P}(\text{CH}_3)_2(\text{CH}_2)_2\text{P}(\text{CH}_3)_2-$. Tridentate ligands include, but are not limited to, $(\text{CH}_3)_2\text{NCH}_2\text{CH}_2\text{P}(\text{Ph})\text{CH}_2\text{CH}_2\text{N}(\text{CH}_3)_2$. Other preferred tridentate ligands are those in which any three of X^1 , X^2 , L , R^1 , R^2 , R^3 , R^{3A} , R^4 , and R^{4A} (e.g., X , L , and any one of R^3 , R^{3A} , R^4 , and R^{4A}) are taken together to be cyclopentadienyl, indenyl or fluorenyl, each optionally substituted with C_2 - C_{20} alkenyl, C_2 - C_{20} alkynyl, C_1 - C_{20} alkyl, C_5 - C_{20} aryl, C_1 - C_{20} alkoxy, C_2 - C_{20} alkenyloxy, C_2 - C_{20} alkynyloxy, C_5 - C_{20} aryloxy, C_2 - C_{20} alkoxycarbonyl, C_1 - C_{20} alkylthio, C_1 - C_{20} alkylsulfonyl, or C_1 - C_{20} alkylsulfinyl, each of which may be further substituted with C_1 - C_6 alkyl, halogen, C_1 - C_6 alkoxy or with a phenyl group optionally substituted with halogen, C_1 - C_6 alkyl or C_1 - C_6 alkoxy. More preferably, in compounds of this type, X , L , and any one of R^3 , R^{3A} , R^4 , and R^{4A} are taken together to be cyclopentadienyl or indenyl, each optionally substituted with vinyl, C_1 - C_{10} alkyl, C_5 - C_{20} aryl, C_1 - C_{10} carboxylate, C_2 - C_{10} alkoxycarbonyl, C_1 - C_{10} alkoxy, C_5 - C_{20} aryloxy, each optionally substituted with C_1 - C_6 alkyl, halogen, C_1 - C_6 alkoxy or with a phenyl group optionally substituted with halogen, C_1 - C_6 alkyl or C_1 - C_6 alkoxy. Most preferably, X , L , and any one of R^3 , R^{3A} , R^4 , and R^{4A} may be taken together to be cyclopentadienyl, optionally substituted with vinyl, hydrogen, Me or Ph. Tetradentate ligands include, but are not limited to $\text{O}_2\text{C}(\text{CH}_2)_2\text{P}(\text{Ph})(\text{CH}_2)_2\text{P}(\text{Ph})(\text{CH}_2)_2\text{CO}_2$, phthalocyanines, and porphyrins.

In a preferred embodiment, the catalyst has the structure of formula (VIB)


(VIB)



wherein R^3 and R^4 are defined above, with preferably at least one of R^3 and R^4 , and more preferably both R^3 and R^4 , being alicyclic or aromatic of one to about five rings, and optionally containing one or more heteroatoms and/or substituents. Q is a linker, typically a hydrocarbylene linker, including substituted hydrocarbylene, heteroatom-containing hydrocarbylene, and substituted heteroatom-containing hydrocarbylene linkers, wherein two or more substituents on adjacent atoms within Q may also be linked to form an additional cyclic structure, which may be similarly substituted to provide a fused polycyclic structure of two to five cyclic groups. Q is

often, although again not necessarily, a two-atom linkage or a three-atom linkage, e.g., $-\text{CH}_2-\text{CH}_2-$, $-\text{CH}(\text{Ph})-\text{CH}(\text{Ph})-$ where Ph is phenyl; $=\text{CR}-\text{N}=$, giving rise to an unsubstituted (when $\text{R} = \text{H}$) or substituted ($\text{R} = \text{other than H}$) triazolyl group; and $-\text{CH}_2-\text{SiR}_2-\text{CH}_2$ (where R is H, alkyl, alkoxy, etc.).


In a more preferred embodiment, Q is a two-atom linkage having the structure $-\text{CR}^{22}\text{R}^{22\text{A}}-\text{CR}^{23}\text{R}^{23\text{A}}-$ or $-\text{CR}^{22}=\text{CR}^{23}-$, more preferably $-\text{CR}^{22}\text{R}^{22\text{A}}-\text{CR}^{23}\text{R}^{23\text{A}}-$, in which case the complex has the structure of formula (VIC)



wherein R^{22} , $\text{R}^{22\text{A}}$, R^{23} , and $\text{R}^{23\text{A}}$ are independently selected from the group consisting of hydrogen, hydrocarbyl, substituted hydrocarbyl, heteroatom-containing hydrocarbyl, substituted heteroatom-containing hydrocarbyl, and functional groups (i.e., Fn, as defined previously), e.g., C_1-C_{20} alkyl, C_2-C_{20} alkenyl, C_2-C_{20} alkynyl, aryl, C_1-C_{20} carboxylate, C_1-C_{20} alkoxy, C_2-C_{20} alkenyloxy, C_2-C_{20} alkynyloxy, aryloxy, C_2-C_{20} alkoxycarbonyl, C_1-C_{20} alkylthio, arylthio, C_1-C_{20} alkylsulfonyl, and C_1-C_{20} alkylsulfinyl, optionally substituted with one or more moieties selected from the group consisting of C_1-C_{10} alkyl, C_1-C_{10} alkoxy, aryl, hydroxyl, sulfhydryl, $-(\text{CO})-\text{H}$, halide, and functional groups (Fn, again, as defined previously).


Additionally, R^{22} , $\text{R}^{22\text{A}}$, R^{23} , and $\text{R}^{23\text{A}}$ may be linked to form a substituted or unsubstituted, saturated or unsaturated ring structure, e.g., a C_4-C_{12} alicyclic group or a C_5 or C_6 aryl group, which may itself be substituted, e.g., with linked or fused alicyclic or aromatic groups, or with other substituents.

Examples of N-heterocyclic carbene ligands incorporated into complex (VIC) thus include, but are not limited to, the following:




-20-

5




10



15

R^3 and R^4 are preferably aromatic, substituted aromatic, heteroaromatic, substituted heteroaromatic, alicyclic, or substituted alicyclic, composed of from one to about five cyclic groups. When R^3 and R^4 are aromatic, they are typically although not necessarily composed of one or two aromatic rings, which may or may not be substituted, e.g., R^3 and R^4 may be phenyl, substituted phenyl, biphenyl, substituted biphenyl, or the like. In one preferred embodiment, R^3 and R^4 are the same and have the structure (XII)

25 (XII)




30

in which R^{24} , R^{25} , and R^{26} are each independently hydrogen, C_1 - C_{20} alkyl, substituted C_1 - C_{20} alkyl, C_1 - C_{20} heteroalkyl, substituted C_1 - C_{20} heteroalkyl, C_5 - C_{20} aryl, substituted C_5 - C_{20} aryl, C_5 - C_{20} heteroaryl, C_5 - C_{30} aralkyl, C_5 - C_{30} alkaryl, or halogen.


35

In especially preferred embodiments, R^{24} , R^{25} , and R^{26} are each independently selected from the group consisting of hydrogen, methyl, ethyl, propyl, isopropyl, hydroxyl, halogen, phenyl, and lower alkyl-substituted phenyl (e.g., dimethylphenyl). In the most preferred embodiments, R^{24} , R^{25} , and R^{26} are the same and are each methyl.

When R^3 and R^4 are alicyclic, they are generally composed of a C_7 - C_{20} , preferably a C_7 - C_{12} , alicyclic structure, e.g., diisopinocampheyl. Complexes formed with such ligands are exemplified by the complex containing the diisopinocampheyl-substituted ligand shown in structural formula (XIV). In the most preferred embodiments, R^{24} , R^{25} , and R^{26} are the same and are each methyl. In another preferred embodiment, R^3 and R^4 are each biphenyl or substituted biphenyl. Catalysts formed with such ligands are exemplified by the complex containing the 2,4,2',6'-tetramethylbiphenyl- (i.e., 2,6-dimethyl-3-(2',6'-dimethylphenyl)phenyl) substituted ligand shown below as structural formula (XIII), preparation of which is described in detail in as illustrated in Example 8.



When R^3 and R^4 are alicyclic, they are generally composed of a C_7 - C_{20} , preferably a C_7 - C_{12} , alicyclic structure, e.g., diisopinocampheyl. Complexes formed with such ligands, exemplified by the complex containing the diisopinocampheyl-substituted ligand shown in structural formula (XIV), are novel compositions of matter and claimed as such herein.



Ligands containing bulky, electron-donating groups such as those illustrated in the complexes of formulae (XIII) and (XIV) provide for very highly active olefin metathesis catalysts. Such catalysts are thus suitable to catalyze reactions for which other, less active catalysts are ineffective, and are also useful in enhancing the stereoselectivity of a catalyzed cross-metathesis reaction.

In the above molecular structures, "Mes" represents mesityl (2,4,6-trimethylphenyl), "iPr" is isopropyl, "Ph" is phenyl, and "Cy" is cyclohexyl.

III. CROSS-METATHESIS OF FUNCTIONALIZED AND SUBSTITUTED OLEFINS:

The present invention, in one embodiment, provides a method for using olefin cross-metathesis to synthesize olefins substituted with functional groups. The reaction is carried out with a functional group-substituted olefinic reactant, and may in fact be carried out with two such functionalized olefins as cross-metathesis reactants. The reaction is catalyzed using a transition metal carbene complex as described in part (II) of this section, and involves reaction between a first olefinic reactant substituted with one or more functional groups, and a second olefinic reactant that may or may not be substituted. With respect to the first olefinic reactant, the functional groups may or may not be in protected form (e.g., a hydroxyl group may be protected as an acyloxy or benzyloxy group). More specifically, the first olefinic reactant has the structure of formula (VIII)



wherein:

Fn is a functional group such as phosphonato, phosphoryl, phosphanyl, phosphino, sulfonato, C₁-C₂₀ alkylsulfanyl, C₅-C₂₀ arylsulfanyl, C₁-C₂₀ alkylsulfonyl, C₅-C₂₀ arylsulfonyl, C₁-C₂₀ alkylsulfinyl, C₅-C₂₀ arylsulfinyl, sulfonamido, amino, amido, imino, nitro, nitroso, hydroxyl, C₁-C₂₀ alkoxy, C₅-C₂₀ aryloxy, C₂-C₂₀ alkoxycarbonyl, C₅-C₂₀ aryloxycarbonyl, carboxyl, carboxylato, mercapto, formyl, C₁-C₂₀ thioester, cyano, cyanato, carbamoyl, epoxy, styrenyl, silyl, silyloxy, silanyl, siloxazanyl, boronato, or boryl, or a metal-containing or metalloid-containing group (wherein the metal may be, for example, Sn or Ge);

n is zero or 1;

Z is a hydrocarbylene or a substituted and/or heteroatom-containing hydrocarbylene linking group such as an alkylene, substituted alkylene, heteroalkylene, substituted heteroalkene, arylene, substituted arylene, heteroarylene, or substituted heteroarylene linkage; and

R⁵, R⁶, and R⁷ are independently selected from the group consisting of hydrogen, -(Z)_n-Fn, hydrocarbyl, substituted hydrocarbyl, heteroatom-containing hydrocarbyl, and substituted heteroatom-containing hydrocarbyl, and, if substituted hydrocarbyl or substituted heteroatom-containing hydrocarbyl, one or more substituents may be -(Z)_n-Fn.

The functional group will generally not be directly bound to the olefinic carbon through a heteroatom containing one or more lone pairs of electrons, e.g., an oxygen, sulfur, nitrogen or phosphorus atom, or through an electron-rich metal or metalloid such as Ge, Sn, As, Sb, Se, Te, etc. With such functional groups, there will normally be an intervening linkage Z, i.e., n is 1.

5 The second olefinic reactant has a molecular structure given by $R^{18}R^{19}C=CR^{20}R^{21}$ wherein R^{18} , R^{19} , R^{20} , and R^{21} may be hydrogen, hydrocarbyl, substituted hydrocarbyl, heteroatom-containing hydrocarbyl, substituted heteroatom-containing hydrocarbyl, or $-(Z)_n-Fn$, wherein n, Z and Fn are as defined earlier.

10 In a preferred embodiment, with respect to the first reactant, R^5 and at least one of R^6 and R^7 is hydrogen, Fn is a phosphonate, and Z is lower alkylene, and in a most preferred embodiment, R^5 , R^6 and R^7 are hydrogen, and Z is methylene, such that the first olefinic reactant is a vinylphosphonate having the structure of formula (XII)



when n is zero, and an allylphosphonate having the structure of formula (XIII)



when n is 1. In formulae (XV) and (XVI), R^{27} and R^{28} are hydrocarbyl, preferably lower hydrocarbyl, and most preferably are lower alkyl such as methyl or ethyl.


25 With respect to the second reactant, it is preferred that R^{18} , R^{19} , and R^{20} are hydrogen, such that reactant has the structure $H_2C=C(H)R^{21}$.

The capability of the methods of the invention with respect to such reactants are illustrated by a series of experiments summarized in the following tables, using the ruthenium catalyst (V)



35 in which IMesH₂ is as defined previously, Cy is cyclohexyl, and Ph is phenyl.

Terminal olefins were reacted with commercially available diethyl vinylphosphonate as described in Example 4. As may be seen in Table 1, cross-metathesis with an olefinic ester resulted in a 95% yield of product, almost exclusively as the (*E*) isomer (Table 1, Entry 1). No dimerization of the vinylphosphonate was detected by ¹H-NMR, allowing for selective cross-metathesis. Alkyl halide (Entry 2) and unprotected aldehyde functionalities (Entry 3) were well tolerated with the ruthenium catalyst (V). Allyl benzene also gave the desired metathesis product, without olefin isomerization (Entry 4). The reaction also gave good yields with a variety of styrenes, which were converted to (*E*)-cinnamylphosphonates in high yield (Table 1, Entry 5).

Table 1

Entry	Cross Metathesis Partner	Product	Isolated Yield ^a
1			95%
2			82%
3			77%
4			90%
5			97% R = H 97% R = 4-OMe 93% R = 4-Br 77% R = 2,4-(CH ₃) ₂

^a >20:1 *E/Z* as determined by ¹H-NMR

Second, diethylallylphosphonate was investigated as a cross-metathesis partner. As indicated by the data in Table 2, allylphosphonates are viable cross-metathesis partners using the present method, providing enhanced cross-metathesis ratios relative to the predicted statistical mixture.

Table 2

10

$$\xrightarrow[\text{CH}_2\text{Cl}_2, 40^\circ\text{C}, 12\text{ hr}]{\text{V (5 mol\%), } \text{CH}_2=\text{CH-R (2.0 eq.)}}$$

15

20

25

Entry	Metathesis Partner	Product	Isolated Yield	E/Z ratio ^a
1			70%	>20:1
2			93%	>20:1
3			73%	>20:1
4			74%	5.4:1
5			85%	3.5:1
6			90%	2.5:1


^a Determined by ¹H-NMR

In addition, some sterically challenging styrenes proved to be excellent CM partners (Table 2, Entry 2 and 3), providing the *E*-isomer exclusively. A trisubstituted olefin is also formed


in excellent yield with modest stereoselectivity (Table 2, Entry 6). All of the reaction products were easily separated from their respective homodimers by column chromatography.

As noted above, the functional group Fn is not necessarily phosphonate. A significant advantage of the present methodology is that the olefinic reactants can be substituted with one or more of a host of functional groups, even if those functional groups are potential ligands for the catalyst.

For example, catalyst (V) has been used to effect cross-metathesis reactions using allylboronates as starting materials. Such reactions are quite useful in the stereoselective synthesis of homoallylic alcohols. Prior to the present invention, the accessibility of functionalized allyl boron reagents was quite limited, and such complexes are traditionally prepared by allylmetal addition to haloboranes or hydroboration of 1,3-dienes, methods that can be incompatible with complex substrates and/or many desired functional groups. The present invention, however, enables a one pot cross-metathesis/allylboration reaction that affords densely functionalized homoallylic alcohols, as illustrated using pinacol allyl boronate according to the following scheme:



The general procedures for carrying out such reactions are described in detail in Example 9. The effects of varying the catalyst and the relative stoichiometries of components of the cross-metathesis/allylation reaction were explored using (*Z*)-1,4-diacetoxy-2-butene and benzaldehyde (Table 3). As may be seen, *E* allylboronates afforded *anti* products with high diastereoselectivity, and the catalyst of the invention, complex (V), was more *E* selective than the bis-tricyclohexylphosphine ruthenium alkylidene complex (II), indicating that catalyst (V) provides the homoallylic product with greater *anti* selectivity. In fact, using the complex of the invention, catalyst loading could be reduced to 2 mol% (Table 3, entry 3) without effecting the yield or diastereoselectivity. The fact that the homoallylic alcohol was formed in 57% yield using a stoichiometric cross partner (Table 3, entry 4), greater than is expected statistically (50%), indicates that the asymmetrically terminated product was favored over symmetrical dimers.

Table 3

entry	eq of cross partner	eq of benzaldehyde	catalyst (mol%)	yield(%)	anti/syn
1	3	1.5	II (5)	32	1.8/1
2	3	1.5	V (5)	75	4.5/1
3	3	1.5	V (2)	75	4.5/1
4	0.5	1.5	V (5)	57	4.7/1
5	3	0.75	V (5)	75	4.5/1

A number of experiments, summarized in Table 4, were carried out in which the aforementioned reaction was used to generate homoallylic alcohols with protected hydroxymethyl, protected aldehyde, and halomethyl side chains from pinacol allyl boronate.

Table 4











entry	cross partner	mol% of V	yield (%)	anti/syn	product
1		5	44–67	4/1	
2		5 2	60 63	4.7/1 3.2/1	
3		5 2	68 69	>20/1 >20/1	
4		5 2	73 72	3.8/1 3.6/1	
5		5 2	78 79	4.9/1 4.6/1	

As may be seen in entries 1 and 2, silyl and benzyl allylic ethers were efficiently transformed to the corresponding homoallylic alcohols 2 and 3, respectively, indicating that the methodology enables facile tuning of the olefinic substrate to conform to preexisting protecting


group strategies. 2-Vinyl-1,3-dioxolane was effectively converted into alcohol 4 in 69% yield as a single diastereomer (entry 3), indicating that an increase in steric bulk at the allylic carbon atom favors the formation of trans olefins. The present method is also effective in achieving incorporation of a halomethyl group directly by allylboration, a reaction that has not been achieved previously. The bromomethyl (entry 4) and chloromethyl (entry 5) allylation products, 5 and 6 respectively, were synthesized in good yields from the corresponding 1,4-dihalo-2-butenes. The present method thus enables a one-step, one-pot synthesis of halogenated targets that would require several steps to prepare by traditional methods.

As another example, catalyst (V) has been used to prepare secondary allylic alcohols from other protected or unprotected secondary allylic reactants. Examples of such reactions are summarized in Table 5.







Table 5

Allylic Substit. Olefin	Cross Partner	Equiv.	Product	Isolated Yield(%)	E/Z ratio
		2.0 eq.		92	13:1
		2.0 eq.		88	>20:1
		2.0 eq.		38	18:1
		2.0 eq.		82	11:1
		0.5 eq.		53	6.7:1
		0.5 eq.		61	>20:1
		1.5 eq.		60	6:1


In addition, catalyst (V) has been used to dimerize the allylic sulfide 3-methylsulfanylpropene according to the following scheme:




As additional examples, complex (V) has been used to catalyze cross-metathesis reactions with other functionalized olefins, as described in Example 6 and as indicated in Table 6:

Table 6**Entry**

As may be seen above, the present method is applicable not only to dimerization of functionalized allylic olefins, but extends to catalytic reaction of such compounds as substrates for cross-metathesis, regardless of the oxidation state of a particular atom in the functional group (e.g., phosphorus-containing functional groups in the form of phosphines, protected phosphines, and phosphonates) or the nature of the functional group (e.g., the reaction proceeds with an allyl amine as well). This versatility is further evidenced by applicants' use of (V) as a catalyst for the preparation of oxazolyphenols as illustrated below:




In another aspect of the invention, the versatility of the present methodology is applied to create functional diversity, i.e., to create a plurality of different olefinic products from a single olefinic reactant. This is carried out by conducting a plurality of olefin metathesis reactions each employing a common first olefinic reactant but a different second olefinic reactant. In this way, a plurality of analogs is provided sharing some structural commonality but having a distinguishing feature. As each olefinic reactant may be substituted with functional groups, cross-metathesis products result that contain those groups, thus providing the option of further derivatization. This can be illustrated by reference to the following schemes:



In the above olefins, n and Z are as defined previously, and Fn^1 , Fn^2 , Fn^3 , and Fn^4 are as defined for Fn or may include other functional groups, e.g., carboxylate, alkoxy, etc. The olefinic reactants may be further substituted on the olefinic carbon atoms with additional $-(Z)_n\text{-Fn}$ groups, or with other moieties such as R^5 , R^6 , and R^7 , defined above with respect to the olefins of formula (VIII).

As a specific example, a family of related potential oxazolyphenol ligands was prepared by cross-metathesis of a single olefinic reactant with a plurality of different second olefinic reactants, again using complex (V) as catalyst, as illustrated below:



It will be appreciated that the capability of the invention in this regard enables the generation of diverse libraries of related but structurally distinct compounds, which may then be screened using any of various processes to ascertain utility, e.g., as potential ligands, reactants, biologically active agents, and the like. The method may be generally characterized as a process for generating a plurality of structurally diverse functionalized olefins from a common olefinic reactant via a cross-metathesis reaction, the method involving the following steps:

(a) contacting a functionalized olefinic substrate with a first olefinic reactant in the presence of a catalyst composed of a Group 8 transition metal alkylidene complex containing an N-heterocyclic carbene ligand, under conditions and for a time period effective to allow cross-metathesis to occur;

(b) in a separate reaction, contacting the first olefinic reactant with a second olefinic reactant having a molecular structure that is different from that of the first olefinic reactant, in the presence of the Group 8 transition metal alkylidene complex, under conditions and for a time period effective to allow cross-metathesis to occur; and


(c) optionally repeating step (b) with a plurality of olefinic reactants each having a different molecular structure.

In another embodiment, the present invention provides a straightforward method for carrying out an olefin cross-metathesis reaction using an α -halogenated olefin in order to provide a directly halogenated olefinic product. In this embodiment, the catalyst used may be the complex of formula (VIB), or it may be an alternative complex of formula (VI) wherein L^1 is a neutral electron donor other than an N-heterocyclic carbene. For example, the catalyst may be a bis(phosphine), in which case both L and L^1 of formula (VI) are phosphine ligands such as triphenylphosphine. At least one of the olefinic reactants has the structure of formula (IX)



wherein X^3 is halo, and R^8 , R^9 , and R^{10} are independently selected from the group consisting of hydrogen, halo, hydrocarbyl, substituted hydrocarbyl, heteroatom-containing hydrocarbyl, substituted heteroatom-containing hydrocarbyl, and $-(Z)_n$ -Fn where n , Z and Fn are as defined previously with respect to formula (VIII). The second olefinic reactant has the same structure, or the structure $R^{18}R^{19}C=CR^{20}R^{21}$ wherein R^{18} , R^{19} , R^{20} , and R^{21} are as defined previously.

The following schemes exemplify cross-metathesis reactions of this type:



The reaction is straightforward and provides a facile method for obtaining an α -halogenated olefin product.


For example, using the procedures described in Example 5, an olefin metathesis catalyst $(L)(L^1)X^1X^2Ru=CR^1R^2$ such as $(H_2IMes)(PCy_3)Cl_2Ru=CHPh$ reacts with 1,1-difluoroethylene to yield the corresponding methylidene $(H_2IMes)(PCy_3)Cl_2Ru=CH_2$ and difluorocarbene $(H_2IMes)(PCy_3)Cl_2Ru=CF_2$ complexes. At elevated temperatures, greater than 98% of the difluorocarbene complex forms, and it can be isolated in pure form by column chromatography.

Although this reaction is not catalytic, the $\text{H}_2\text{C}=\text{CF}_2$ double bond is cleaved in a metathesis fashion, and as such, it is the first example of metathesis involving a directly halide-substituted olefin. In addition, it should be emphasized that $(\text{H}_2\text{IMes})(\text{PCy}_3)\text{Cl}_2\text{Ru}=\text{CF}_2$ is active for subsequent metathesis reactions, such as the ring-closing metathesis of diethyl diallylmalonate and the ring-opening metathesis polymerization of norbornene derivatives. The activity of $(\text{H}_2\text{IMes})(\text{PCy}_3)\text{Cl}_2\text{Ru}=\text{CF}_2$ can be enhanced by the addition of HCl or CuCl , which aid in the dissociation of PCy_3 from the metal center. The bis(pyridine) derivative of the catalyst, $(\text{H}_2\text{IMes})(\text{py})_2\text{Cl}_2\text{Ru}=\text{CF}_2$, is somewhat more active for subsequent metathesis reactions than the PCy_3 complex, presumably because the pyridine ligands are less basic and thus more labile. Likewise, the bis(phosphine) olefin metathesis catalyst $(\text{PCy}_3)_2\text{Cl}_2\text{Ru}=\text{CHPh}$ reacts with 1,1-difluoroethylene to yield the corresponding methyldiene $(\text{PCy}_3)_2\text{Cl}_2\text{Ru}=\text{CH}_2$ and difluorocarbene $(\text{PCy}_3)_2\text{Cl}_2\text{Ru}=\text{CF}_2$ complexes.

In a further embodiment, a method is provided for synthesizing substituted olefins, particularly geminal disubstituted olefins, 1,1,2-trisubstituted olefins and quaternary allylic olefins, wherein the method comprises using the complex of formula (VI) to catalyze a cross-metathesis reaction between a geminal disubstituted olefin, a 1,1,2-trisubstituted olefin, or a quaternary allylic olefin, and a second olefin. If it is a geminal disubstituted olefin or a 1,1,2-trisubstituted olefin, the first olefin has the structure (X)



wherein R^{11} , R^{12} , R^{13} , and R^{14} are selected from the group consisting of hydrogen, halo, hydrocarbyl, substituted hydrocarbyl, heteroatom-containing hydrocarbyl, substituted heteroatom-containing hydrocarbyl, and $-(\text{Z})_n\text{-Fn}$ where n , Z and Fn are as defined above, with the provisos that R^{11} and R^{12} , or R^{13} and R^{14} are other than hydrogen for a geminal disubstituted olefin, and that R^{11} , R^{12} , and R^{13} are other than hydrogen for a 1,1,2-trisubstituted olefin. If it is a quaternary allylic olefin, the first olefin has the structure (XI)



wherein R^{11} and R^{12} are as defined previously, and R^{15} , R^{16} , and R^{17} are nonhydrogen substituents.

In the aforementioned cross-metathesis reaction, the second olefin has a molecular structure given by $R^{18}R^{19}C=CR^{20}R^{21}$ wherein R^{18} , R^{19} , R^{20} , and R^{21} may be hydrogen, hydrocarbyl, substituted hydrocarbyl, heteroatom-containing hydrocarbyl, or substituted heteroatom-containing hydrocarbyl.

Generally, the reaction is carried out with the two olefinic reactants in a mole ratio in the range of about 1:3 to 3:1, at a temperature in the range of about 20 °C to about 40 °C, for a time period in the range of about 4 to 16 hours. Typically, about 0.01 to 7.5 mole % catalyst is used. However, the reaction is also viable if there is a large excess of one reactant, such as is the case when one reactant serves as a solvent for the reaction mixture.





















Another example of such a reaction is the synthesis of 1,1-dimethyl olefins through a cross-metathesis reaction of α -olefins with isobutylene or 2-methyl-2-butene. The capability of the methods of the invention with respect to such a reaction is illustrated by a series of experiments summarized in Table 7, using 2-methyl-2-butene as the geminal disubstituted olefin and $(\text{IMesH}_2)(\text{PCy}_3)\text{Cl}_2\text{Ru}=\text{C}(\text{H})\text{Ph}$ (complex (V)) as the catalyst.

Table 7

Entry	Terminal α -Olefin	Product	Yield
5	1		97
	2		97
10	3		96
	4		91
15	5		91
20	6		99
	7		80
25	8		83

Additional experiments, summarized in Table 8, were carried out using catalyst (V) to generate trisubstituted olefins from symmetrical 1,1-disubstituted olefins as starting materials.































Table 8

Entry	1,1-Disubstituted Olefin	Temp(°C)	Metathesis Partner	Product	Isolated Yield
5	1				
		40			97
	2				
		40			88
10	3				
		40			48
	4				
		40			65
15	5				
		40			96
	6				
		40			42
20	7				
		40			83



Further experiments illustrating the versatility of the present methodology were carried out in order to generate 1,2-disubstituted olefins with quaternary allylic carbons using the catalyst (V), the results of which are summarized in Table 9.

Table 9

Entry	Quat. Allylic Olefin	Equiv.	CM Partner	Product	Yield
5 1		0.5 eq.			95
2		2.0 eq.			97
3		0.5 eq.			66
10 4		2.0 eq. 1.0 eq.			90 69
5		3 eq.			88
15 6		~50eq.			93
7		3 eq.			44
8		2 eq.			93
20 9		1 eq.			95
10		2 eq.			71

25 These reactions were stereoselective, resulting in virtually exclusive formation of the *trans* olefin isomer, as may be seen under the column heading "E/Z Ratio" in the figure. This stereoselectivity is an important feature of the method, insofar as prior to the present invention, there was no general method for controlling the stereoselectivity of newly formed olefins.

30 In a related embodiment of the invention, a stereoselective method for carrying out an olefin cross-metathesis reaction is provided, wherein the stereochemistry of the olefinic product may be either *cis* or *trans*, as desired. The catalyst used has the structure of formula (VIB), wherein the nitrogen atoms of the N-heterocyclic carbene ligand are substituted with bulky substituents, i.e., R³ and R⁴ are aromatic, substituted aromatic, heteroaromatic, substituted aromatic, alicyclic, or substituted alicyclic. For a stereoselective synthesis that will preferentially
 35 result in a *cis*-1,2-disubstituted olefin, bulky R³ and R⁴ substituent are preferred, e.g., bicyclic or polycyclic ligands that may or may not be aromatic. If R³ and R⁴ are aromatic, they are generally


composed of two to five aromatic rings that may be fused or linked (e.g., biphenyl or substituted biphenyl), and if R³ and R⁴ are alicyclic, they are generally composed of a C₇-C₂₀, preferably a C₇-C₁₂, alicyclic structure that may or may not be substituted. Representative such R³ and R⁴ groups thus include the alicyclic groups norbornyl, adamantyl, camphenyl, isobornyl, any of which may be substituted, e.g., with a lower alkyl group (as in diisopinocampheyl, as shown in the structure of formula (XIV)), and the bicyclic groups biphenyl and 2', 6' -dimethyl-3'-(2'',6''-dimethylphenyl)phenyl (as shown in the structure of formula (XIII)).

It is to be understood that while the invention has been described in conjunction with the preferred specific embodiments thereof, that the foregoing description as well as the examples that follow are intended to illustrate and not limit the scope of the invention. Other aspects, advantages and modifications within the scope of the invention will be apparent to those skilled in the art to which the invention pertains.

EXPERIMENTAL

IMesH₂Cl was prepared according to a modified version of the procedure described in Scholl et al. (1999) *Org. Lett.* 1:953-956 and Jafarpour et al. (2000) *Organometallics* 19:2055-2057. Unless otherwise specified, all other reagents were purchased from commercial suppliers and used without further purification. All other solvents were purified by passage through a solvent column (containing activated A-2 alumina; see Pangborn et al. (1996) *Organometallics* 15:1518-1520.). Analytical thin-layer chromatography (TLC) was performed using silica gel 60 F254 precoated plates (0.25 mm thickness) with a fluorescent indicator. Flash column chromatography was performed using silica gel 60 (230-400 mesh) from EM Science. ¹H, ¹³C, and ³¹P NMR spectra were obtained on a Varian 300 MHz Fourier Transform spectrometer (300 MHz ¹H, 75.4 MHz ¹³C, 121.4 MHz ³¹P). All chemical shift values are given in parts-per million (δ) and are referenced with respect to residual solvent (¹H and ¹³C) or phosphoric acid (³¹P).

Preparation of IMesH₂Cl: IMesH₂Cl, used as a starting material in Examples 1 through 3, was synthesized according to the following scheme:




To a solution of glyoxal (9 mL, 79 mmol, 40% wt in H₂O) in isopropanol (100 mL) and H₂O (200 mL) was added mesitylamine (25 mL, 2.2 eq.) at 0 °C. The reaction mixture was stirred while allowing to warm to room temperature. Immediately upon addition of amine, yellow precipitates were formed. After 24 hrs of stirring at ambient temperature, the precipitates were filtered and washed with H₂O (1 x 100 mL) and hexanes (3 x 100 mL). The yellow precipitates obtained were dried *in vacuo* to yield the diimine (20.6 g, 89%).

To a solution of diimine (8.0 g, 27.3 mmol) in THF (100 mL) was added NaBH₄ (4.24 g, 112.1 mmol) at 0 °C. Concentrated HCl (4.5 mL, 2 eq.) was added dropwise over 30 minutes. After the HCl addition, the reaction mixture was stirred at 0 °C for 20 min. Then, 3 M HCl (250 mL) was added carefully to the flask at 0 °C and the mixture was stirred for an additional 1 hr, allowing the temperature to rise to ambient temperature. The resulting white precipitates were filtered and washed with water (200 mL) and 5% acetone-ether (150 mL). The product (9.4 g, 93%) was obtained as a white solid and dried *in vacuo*. To a suspension of the HCl salt (8.5 g, 23 mmol) in HC(OEt)₃ (35 mL, 162 mmol) was added 2 drops of HCO₂H (adding about 1 mol%). The reaction mixture was then heated at 120 °C for 5 hr under Ar. Then, the reaction mixture was cooled to an ambient temperature and hexane (200 mL) was added. The mixture was stirred for 1 hr and the white precipitates were filtered, washed with hexane (~ 200 mL) and dried *in vacuo* to yield the IMesH₂HCl salt (7.6 g, 96%).

EXAMPLE 1

REPRESENTATIVE PROCEDURE FOR SYNTHESIS OF RUTHENIUM ALKYLIDENE CATALYSTS

Synthesis of RuCl₂(=CH-CH=C(CH₃)₂)(IMesH₂)(PCy₃) (complex (2), Scheme 1):





SCHEME 1

[Ru(COD)Cl₂]_n (300 mg, 1 mmol), IMesH₂Cl (1.47 g, 4 mmol), tricyclohexylphosphine (300 mg, 1 mmol), and KN(SiMe₃)₂ (540 mg, 2.5 mmol) were weighed directly into a 600 mL Schlenk tube. The flask was evacuated and filled with dry argon (2x). Degassed benzene (300 mL) was added and the flask was pressurized to 30 psi with H₂. The suspension was vigorously stirred for 12 hours at 90 °C, yielding a bright yellow solution and white precipitate (1). After cooling the reaction to 5 °C, propargyl chloride (0.3 mL, 4 mmol) was slowly added via syringe and the reaction mixture was allowed to warm to room temperature. The resulting brown benzene solution was washed with degassed 1M HCl (2x), degassed brine (2x), filtered through Celite and concentrated *in vacuo* to afford compound (2) as a brown solid in 90% yield (~ 95% purity). The brown solid displayed catalytic behavior identical with previously synthesized second-generation catalysts. Analytically pure (2) was obtained by column chromatography on silica gel (degassed 3:1 hexanes/Et₂O). ¹H NMR (CD₂Cl₂): δ 18.49 (d, J = 11.1 Hz, 1H), 7.26 (d, J = 10.9 Hz, 1H), 6.97 (s, 2H), 6.77 (s, 2H), 3.92 (m, 4H), 2.58 (s, 6H), 2.37 (s, 6H), 2.29 (s, 3H), 2.23 (s, 3H), 0.88-1.584 (m, 33H), 1.06 (s, 3H), 1.08 (s, 3H). ³¹P NMR (CD₂Cl₂): δ 28.9. The reaction was repeated several times with one or more reaction conditions modified so as to optimize the yield of the product. It was found that the yield could be increased to greater than 95% by reducing the reaction temperature from 90 °C to 80 °C.

Analogous ruthenium alkylidene complexes can be prepared using the aforementioned protocol and differently substituted phosphines, alkynes, etc., as indicated in the following two examples.

EXAMPLE 2

Synthesis of RuCl₂(=CH-CH=C(CH₃)₂)(IMesH₂)(PPh₃) (complex (4), Scheme 2):




The procedure of Example 2 was employed using [Ru(COD)Cl₂]_n (300 mg, 1 mmol), IMesH₂Cl (0.74 g, 2 mmol), triphenylphosphine (280 mg, 1 mmol), and KN(SiMe₃)₂ (380 mg, 1.9

mmol), giving 550 mg (68%) of complex (3). ^{31}P NMR (CD_2Cl_2): δ 24.0. ^1H NMR (CD_2Cl_2): δ 18.49 (d, $J = 11.1$ Hz, 1H).

EXAMPLE 3

Synthesis of $\text{RuCl}_2(=\text{CH}-\text{CH}-\text{Ph})(\text{IMesH}_2)(\text{PCy}_3)$ (complex (5), Scheme 3):



SCHEME 3

15

$\text{RuCl}_2(=\text{CHPh})(\text{PCy}_3)_2$ (phenylmethylene-bis(tricyclohexylphosphine) ruthenium dichloride, "catalyst (I)") (6.00 g, 7.29 mmol, 1.0 eq.), IMesH_2HCl salt prepared above (2 eq.), and potassium t-butoxide (2 eq.) were placed in a Schlenk flask. 60 mL of anhydrous degassed hexanes (Aldrich SureSeal bottle) were added. A vacuum was applied to further degas the reaction mixture, which was then heated to 60°C for 24 hours. The suspension changed color from purple to orange-brown over the reaction time. After approximately 24 hr, the mixture was cooled to room temperature, and an excess of 1:1 isopropanol:water (180 mL) was added. The mixture was stirred rapidly in air for 30 min., then filtered using a medium porosity frit, and washed with isopropanol-water (3 x 100 mL) and hexanes (3 x 100 mL). The solids were dried in *in vacuo*, and the yield was approximately 75%. ^1H NMR (CD_2Cl_2 , 400 MHz) δ 19.16 (s, 1H), 7.37-7.05 (m, 9H), 3.88 (s, 4H), 2.56-0.15 (m, 51H); ^{31}P NMR (CD_2Cl_2 , 161.9 MHz) δ 31.41; HRMS (FAB) $\text{C}_{45}\text{H}_{65}\text{Cl}_2\text{N}_2\text{PRu}$ [M^+] 848.3306, found 848.3286.

20

25

EXAMPLE 4

30

REPRESENTATIVE PROCEDURES FOR CROSS-METATHESIS REACTIONS USED TO SYNTHESIZE FUNCTIONALIZED OLEFINS

Preparation of Olefinic Phosphonates (Tables 1 and 2): Terminal olefin (0.75 mmol) and diethyl vinylphosphonate (Aldrich) or diethyl allylphosphonate (Acros Organics, 0.51 mmol) were added simultaneously *via* syringe to a stirring solution of (5) (21 mg, 0.026 mmol, 5.2 mol %) in CH_2Cl_2 (2.5 mL, 0.2M in phosphonate) under a nitrogen atmosphere. The flask was fitted


35

with a condenser and refluxed under nitrogen for 12 hours. The reaction mixture was then reduced in volume to 0.5 mL and purified directly on a silica gel column (2x10 cm), eluting with 1:1 hexane:ethyl acetate to provide cross products as viscous oils.

5

EXAMPLE 5

**REPRESENTATIVE PROCEDURES FOR SYNTHESIS OF
DIRECTLY HALOGENATED OLEFINS**

**SCHEME 4**

30

Synthesis and characterization of $[(\text{IMesH}_2)(\text{PCy}_3)(\text{Cl})_2\text{Ru}=\text{CF}_2]$: A solution of 0.32 g (0.37 mmol) $[(\text{IMesH}_2)(\text{PCy}_3)(\text{Cl})_2\text{Ru}=\text{CHPh}]$ (**5**) in dry, degassed benzene (15 mL) in a thick-walled glass ampule was put under ~1.5 atm of 1,1-difluoroethylene. The reaction was heated at 60°C for 12 hrs, during which time it changed from reddish to brown in color. The solution was then concentrated to 5 mL and purified by column chromatography in air (silica gel, 5:1 pentane/THF). The orange fraction was stripped of solvent and dried under vacuum: yield 0.26 g (86%). ^1H NMR (499.852 MHz, 25°C, CD_2Cl_2): δ 1.118 [br, 15H, PCy_3], 1.626 [br, 15H, PCy_3], 2.248 [s, 3H, p- CH_3 of Mes], 2.285 [s, 3H, p- CH_3 of Mes], 2.385 [m, 3H, PCy_3], 2.480 [s, 6H, o- CH_3 of Mes], 2.551 [s, 6H, o- CH_3 of Mes], 4.003 [s, 4H, $\text{NCH}_2\text{CH}_2\text{N}$], 6.921 [s, 4H, m-H of Mes]. $^{13}\text{C}\{^1\text{H}\}$ NMR (125.705 MHz, 30°C, C_6D_6): δ 19.44 [s, CH_3 of Mes], 20.65 [s, CH_3 of Mes], 21.49 [s, CH_3 of Mes], 21.50 [s, CH_3 of Mes], 26.92 [d, $J = 1.3$ Hz, PCy_3], 28.50 [d, $J = 10$ Hz, PCy_3], 30.14 [s, PCy_3], 33.34 [d, $J = 18$ Hz, PCy_3], 51.86 [d, $^4J_{\text{PC}} = 2.6$ Hz, $\text{NCH}_2\text{CH}_2\text{N}$], 52.61

40

[d, $^4J_{PC} = 3.5$ Hz, NCH₂CH₂N], 127.30 [s, Mes], 128.17 [s, Mes], 129.26 [s, Mes], 129.51 [s, Mes], 130.11 [s, Mes], 130.52 [s, Mes], 134.68 [d, $^4J_{PC} = 0.7$ Hz, *ipso*-C of Mes], 136.85 [s, *ipso*-C of Mes], 138.91 [s, Mes], 138.93 [s, Mes], 139.03 [s, Mes], 139.67 [s, Mes], 217.23 [d, $^2J_{CP} = 87$ Hz, NCN], 218.09 [td, $^2J_{CP} = 12$ Hz, $^1J_{CF} = 430$ Hz, Ru=CF₂]. ¹⁹F NMR (282.192 MHz, 25°C, CD₂Cl₂): δ 133.74 [d, $^3J_{FP} = 4.5$ Hz]. ³¹P{¹H} NMR (121.392 MHz, 25°C, CD₂Cl₂): δ 32.15 (t, $^3J_{PF} = 4.4$ Hz]. IR (KBr pellet): 1167 and 1172 (ν_{C-F}).

When the reaction is performed at room temperature, the product mixture contains approximately 40% methylidene and 60% difluorocarbene, as well as styrene (H₂C=CHPh) and β,β-difluorostyrene (F₂C=CHPh). The amount of difluorocarbene complex formed increased to greater than 98% when the reaction was carried out at 60°C instead. In a similar fashion, the bis(phosphine) olefin metathesis catalyst [(PCy₃)₂Cl₂Ru=CHPh] reacts with 1,1-difluoroethylene to yield the corresponding methylidene [(PCy₃)₂Cl₂Ru=CH₂] and difluorocarbene [(PCy₃)₂Cl₂Ru=CF₂] complexes.

15

EXAMPLE 6

**REPRESENTATIVE PROCEDURES FOR SYNTHESIS OF
SUBSTITUTED ALLYLIC OLEFINS**

20

Allyldiphenylphosphine oxide (53mg, 0.22 mmol) and catalyst (5) (14mg, .0165 mmol) were weighed directly into a dried 25mL round bottom flask with a Teflon stirbar. Dry methylene chloride (1.5mL, 0.3M) and *cis*-2-butene-1,4-diacetate (TCI) (70 μL, 0.44 mmol) were added *via* syringe under a nitrogen atmosphere. The flask was fitted with a condenser and refluxed under nitrogen for 12 hours. The reaction mixture was then reduced in volume to 0.5 mL and purified directly on a silica gel column (2x10 cm), eluting with 1:1 hexane:ethyl acetate to provide the cross product (62mg, 90% yield) as viscous oil/semi solid as confirmed by ¹H and ¹³C-NMR.

25

EXAMPLE 7

**REPRESENTATIVE PROCEDURES FOR SYNTHESIS OF
TRISUBSTITUTED AND QUATERNARY ALLYLIC OLEFINS**

30

General procedure for isobutylene CM: To an oven dried, 100 mL Fischer-Porter bottle with Teflon stir bar, ruthenium metathesis catalyst (15.0 mg, 0.018 mmol, 0.01-0.02 equiv.) was added. The bottle was capped with a rubber septum and flushed with dry nitrogen and cooled to -78 °C (or temperature sufficient to freeze substrate). Substrate (0.9-1.9 mmol) was injected into the bottle. Once the substrate was frozen, a pressure regulator was attached to the bottle. The bottle was evacuated and backfilled with dry nitrogen 3 times. Subsequently, isobutylene (5-10 mL, 50-100 equiv.) was condensed into the bottle. The bottle was backfilled to ~2 psi with nitrogen, sealed, and allowed to slowly warm to room temperature, at which time it was

35

transferred to an oil bath at 40 °C. After stirring for 12-18 hours, the bottle was removed from the oil bath and allowed to cool to room temperature. The isobutylene was slowly vented off at room temperature until the pressure apparatus could be safely disassembled. The remaining mixture was taken up in organic solvent for subsequent silica gel chromatography and/or spectrographic characterization.

Representative procedure for CM with 2-methyl-2-butene (Table 7, Entry 5): Pw Pentafluoroallylbenzene (225 μ L, 1.468 mmol) from Aldrich Chem. Co. and 2-methyl-2-butene (3.2 mL) from Aldrich Chem. Co. were added simultaneously via syringe to a stirring solution of catalyst (5) (12.5 mg, 0.015 mmol, 1.0 mol %) under a nitrogen atmosphere. The reaction mixture was allowed to stir at room temperature for 12 hours, and was then reduced in volume to 0.5 mL and purified directly on a silica gel column (2x10 cm), eluting with 20:1 hexane:ethyl acetate to provide the cross-metathesis product (316mg, 1.337 mmol, 91% yield) as a viscous oils.

Representative procedure for CM with 3,3-dimethyl-1-butene (Table 8, Entry 6): *Cis*-2-butene-1,4-diacetate (50 μ L, .3168 mmol) from TCI America and 3,3-dimethyl-1-butene (3.2mL, 0.15M) from Aldrich Chem. Co. were added simultaneously via syringe to a stirring solution of catalyst (5) (10 mg, 0.012 mmol, 3.7 mol %) under a nitrogen atmosphere. The flask was allowed to stir at room temperature for 12 hours. The reaction mixture was then reduced in volume to 0.5 mL and purified directly on a silica gel column (2x10 cm), eluting with 50:1 hexane:ethyl acetate to provide the cross-metathesis product (92 mg, .5891 mmol, 93% yield) as a viscous oils.

Representative 40 °C procedure with 3,3-dimethyl-1-hexene (Table 8, Entry 5): Allylbenzene (40 μ L, 0.30 mmol) from Aldrich Chem. Co. and 3,3-dimethyl-1-hexene (140 μ L, .90 mmol, 3 equiv.) from Aldrich Chem. Co were added simultaneously *via* syringe to a stirring solution of catalyst (5) (20 mg, 0.024 mmol, 7.8 mol %) in CH₂Cl₂ (2.0 mL, 0.15M in allylbenzene) under a nitrogen atmosphere. The flask was fitted with a condenser and refluxed under nitrogen for 12 hours at 40C. The reaction mixture was then reduced in volume to 0.5 mL and purified directly on a silica gel column (2x10 cm), eluting with 20:1 hexane:ethyl acetate to provide the cross product (54 mg, .27 mmol, 88% yield) as a viscous oils.

EXAMPLE 8

REPRESENTATIVE PROCEDURES FOR SYNTHESIS OF CIS-1,2-DISUBSTITUTED OLEFINS

(a) General considerations: All manipulations were performed using a combination of glovebox, high vacuum, and Schlenk techniques under a nitrogen atmosphere, unless otherwise specified. Solvents were dried and degassed by standard procedures. ¹H and ¹³C NMR spectra were measured on a Varian 300 or an Inova 500 spectrometer. Chemical shifts are reported in ppm relative to SiMe₄ (δ = 0) and were referenced internally with respect to the protio solvent

impurity ($\delta = 5.32$ for CDHCl_2) and the ^{13}C resonances ($\delta = 54.00$ for CD_2Cl_2). Coupling constants are in hertz. The silica gel used for the purification of organometallic complexes was obtained from TSI Scientific, Cambridge, MA (60 Å, pH 6.5-7.0).

(b) Preparation of representative catalysts useful for stereoselective synthesis of cis-1,2-disubstituted olefins: The ligand precursors 1,3-(+)-diisopinocampheyl-4,5-dihydroimidazolium tetrafluoroborate salt [IPCimid(H)][BF_4] and 1,3-bis[2',6'-dimethyl-3'-(2'',6''-dimethylphenyl)phenyl]-4,5-dihydroimidazolium chloride salt were prepared by analogy to the method of Kaloustian et al. (see Saba et al. (1991) *Tet. Lett.* 32:5031-34).

(b-i) In a nitrogen-filled glovebox, a large Schlenk flask was charged with 0.475 g [IPCimid(H)][BF_4] (1.120 mmol), 0.131 g potassium *tert*-butoxide (1.120 mmol), and 30 mL anhydrous, degassed benzene. This mixture was stirred at room temperature for 6 hrs. Then, a solution of 0.400 g [$(\text{PCy}_3)_2(\text{Cl})_2 \text{Ru}=\text{CHPh}$] (0.486 mmol) in 15 mL benzene was added, and the reaction was stirred for 30 min at room temperature, during which time the mixture changed from purple to brown. The reaction was concentrated to a third of its original volume under vacuum and transferred to a silica gel column (1.5×16"). The product was quickly eluted with 5:1 heptane:ether. The second, brown band was collected and stripped of solvent. The oily residue that remained was redissolved in a minimum amount of benzene and lyophilized to yield 0.080 g of the desired product as a brown powder (19%). ^1H NMR (299.817 MHz, 20°C, CD_2Cl_2): 20.583 and 20.577 [two s, two orientations of $\text{Ru}=\text{CH}_a$], 8.54 [br s], 7.60 [t, $J = 7.3$], 7.34 (t, $J = 7.8$), 5.16 (qt, $J = 5.1$), 3.46 -3.96 [m], 2.86 (t, $J = 12.4$), 2.34 -2.50 [m], 1.44 -2.20 [m], 1.43 (s), 1.41 (s), 0.82 -1.31 [m], 1.26 [s], 1.12 [s], 1.01 [s], 0.57 [d, $J = 6.9$], 0.25 [s]. ^1H NMR (299.817 MHz, -70°C, CD_2Cl_2): 20.32 [s, $\text{Ru}=\text{CH}_a$], 9.07 [d, $J = 7.8$], 7.87 [t, $J = 7.1$], 7.59 [t, $J = 7.4$], 7.35 [m], 4.92 [br], 3.30 -3.90 [m], 2.69 [m], 2.44 -0.78 [m], 1.33 [s], 1.16 [s], 1.02 [s], 0.90 [s], 0.88 [s], 0.86 [s], 0.80 [s], 0.78 [s], 0.43 [s], 0.11 [br d, $J = 5.7$]. $^{31}\text{P}\{^1\text{H}\}$ MR (121.39 MHz, 25°C, CD_2Cl_2): 21.72 [s]. $^{31}\text{P}\{^1\text{H}\}$ NMR (121.39 MHz, -65°C, CD_2Cl_2): 21.95 [s], 21.16 [s].

(b-ii) 2-*tert*-butoxy-1,3-bis[2',6'-dimethyl-3'-(2'',6''-dimethylphenyl)phenyl]-4,5-dihydroimidazol-2-ylidene was prepared by stirring a suspension of potassium *tert*-butoxide (9 mg, 0.080 mmol) and 1,3-bis[2',6'-dimethyl-3'-(2'',6''-dimethylphenyl)-phenyl]-4,5-dihydroimidazol-2-ylidene (50 mg, 0.079 mmol) in benzene (1 mL) for 1 h at room temperature. To this suspension was added phenylmethylene-bis(tricyclohexylphosphine) ruthenium dichloride (65 mg, 0.079 mmol) in benzene (1 mL). The solution, which immediately became pinkish purple, was stirred at 50 °C for 16 h. After this time, the solution was cooled and the solvent was evaporated to near dryness. The residue was passed through a plug of TSI silica gel, using 1:1 ether/pentane as the eluant. After concentrating, the solids were washed with pentane (5 x 1 mL). The solid material was dissolved in benzene (1 mL) and was frozen (dry ice/acetone). The solvent was removed by sublimation to give phenylmethylene 1,3-bis[2',6'-dimethyl-3'-(2'',6''-dimethyl-

phenyl)phenyl]-4,5-dihydroimidazol-2-ylidene (50 mg, 62%) as a pink solid. ¹H NMR (500 MHz, toluene-*d*₈): δ = 19.46 (s, 1H), 9.59 (br s, 1H), 7.35-6.18 (multiple peaks, 14H), 3.68-3.22 (multiple peaks, 4H), 2.98 (s, 3H), 2.61 (s, 3H), 2.46 (s, 3H), 2.35 (s, 3H), 2.19 (s, 3H), 1.98 (s, 3H), 1.92 (s, 3H), 2.09-1.10 (multiple peaks, 36H) ppm. ³¹P NMR (202 MHz, toluene-*d*₈): δ = 34.54 ppm (s).

The same procedures may be followed for preparation of a ruthenium catalyst containing other ligands.

Representative cross-metathesis reactions using the catalysts prepared in (b-i) and (b-ii):

(c-i) Upon isolation of the catalyst prepared with the 1,3-(+)-diisopinocampheyl-4,5-dihydroimidazole-2-ylidene ligand as described in (b), a representative cross-metathesis reaction can be conducted with 5 mol% of the catalyst in a reaction with a 2:1 ratio of *cis*-2-butene-1,4-diacetate and an α-terminal olefin at 40 °C in methylene chloride for 12 hours to generate the cross-metathesis allylic acetate product as a 1.3:1 mixture of *trans* and *cis* isomers in 85% overall yield. Upon isolation of the second catalyst prepared as described in (a), i.e., phenylmethylene 1,3-bis[2',6'-dimethyl-3'-(2'',6''-dimethylphenyl)phenyl]-4,5-dihydroimidazol-2-ylidene tricyclohexylphosphine ruthenium dichloride, a representative cross-metathesis reaction can be conducted with 5 mol% of the catalyst in a reaction with a 2:1 ratio of *cis*-2-butene-1,4-diacetate and an α-terminal olefin at 40 °C in methylene chloride for 6 hours to generate the cross-metathesis allylic acetate product as a 2.2:1 mixture of *trans* and *cis* isomers in 60% overall yield.

(c-ii) Allylbenzene (15 mg, 0.13 mmol), *cis*-1,4-diacetoxy-2-butene (45 mg, 0.26 mmol), and phenylmethylene 1,3-bis[2',6'-dimethyl-3'-(2'',6''-dimethylphenyl)phenyl]-4,5-dihydroimidazol-2-ylidene (4 mg, 0.004 mmol) were dissolved in CD₂Cl₂ (0.7 mL) and added to a screw-cap NMR tube. The tube was heated at 40 °C and the reaction progress was monitored periodically by NMR. After 12 h at 40 °C, NMR analysis indicated that the reaction had proceeded to 77% completion. The tube was cooled to room temperature and the solution was transferred to a 5 mL flask and the solvent was removed in vacuo. The residue was taken up in CH₂Cl₂ and passed through a small plug of silica. The solution was concentrated and the residue was taken up in CDCl₃ and was added to an NMR tube. NMR analysis indicated a 2.4:1 (*E*:*Z*) ratio of *E*/*Z*-1-acetoxy-4-phenyl-2-butene.

EXAMPLE 9

ALLYLBORONATES AS CROSS-METATHESIS SUBSTRATES


General procedures for carrying out cross-metathesis reactions with pinacol allyl boronate: A flame-dried round-bottomed flask was charged with pinacol allyl boronate (1 eq.) and the olefin cross partner CH₂=CHR (3.0 equiv.). a rubber septum was attached, dichloromethane was added (0.2 - 0.3 N in pinacol allyl boronate), and argon was bubbled

through the resultant solution for 10 min. Under a stream of argon, catalyst (5) (0.050 equiv.) was added to the degassed solution as a solid. A reflux condenser was attached immediately, and the entire system was flushed with argon for 2 min. The colored solution was then heated at reflux for 2 - 12 h, and the reaction was monitored by thin-layer chromatography. Upon consumption of the allylboronate reactant, the aldehyde R'CHO (1.5 eq.) was added to the reaction mixture through a syringe, and the resultant solution was stirred at 23 °C in vacuo, and the residue was purified by means of silica gel chromatography to yield the allylic alcohol product having the structure $\text{CH}_2=\text{CHR}-\text{CH}(\text{OH})-\text{R}'$.

-50-

CLAIMS

1. A method for synthesizing a functionalized olefin via a cross-metathesis reaction, comprising contacting (a) a first olefinic reactant directly substituted with a functional group Fn
 5 selected from the group consisting of: phosphonato; phosphoryl; phosphanyl; phosphino; sulfonato; C₁-C₂₀ alkylsulfanyl; C₅-C₂₀ arylsulfanyl; C₁-C₂₀ alkylsulfinyl; C₅-C₂₀ arylsulfinyl; sulfonamido; amino; imino; nitro; nitroso; hydroxyl; unsubstituted C₁-C₂₀ alkoxy; substituted C₂-C₂₀ alkoxy; C₅-C₂₀ aryloxy; C₅-C₂₀ aryloxycarbonyl; mercapto; C₁-C₂₀ thioester; cyano; cyanato; carbamoyl; styrenyl; siloxazanyl; boronato; boryl; stannyl; and germyl, or indirectly substituted with a functional group Fn
 10 selected from the group consisting of: phosphonato; phosphoryl; phosphanyl; phosphino; sulfonato; C₁-C₂₀ alkylsulfanyl; C₅-C₂₀ arylsulfanyl; C₁-C₂₀ alkylsulfinyl; C₅-C₂₀ arylsulfinyl; sulfonamido; amino; imino; nitro; nitroso; hydroxyl; unsubstituted C₁-C₂₀ alkoxy; substituted C₄-C₂₀ alkoxy; C₅-C₂₀ aryloxy; C₅-C₂₀ aryloxycarbonyl; carboxylato; mercapto; C₁-C₂₀ thioester; cyano; cyanato; carbamoyl; styrenyl; siloxazanyl; boronato; boryl; stannyl; and germyl, with (b) a second olefinic reactant in the
 15 presence of (c) a catalyst composed of a Group 8 transition metal alkylidene complex under conditions and for a time period effective to allow cross-metathesis to occur, wherein the catalyst has the structure of formula (VIA)



25 in which:

- M is a Group 8 transition metal;
 X¹ and X² may be the same or different, and are anionic ligands or polymers;
 R¹ is selected from the group consisting of: hydrogen; hydrocarbyl; substituted hydrocarbyl;
 heteroatom-containing hydrocarbyl; substituted heteroatom-containing hydrocarbyl; and carboxyl;
 30 R² is selected from the group consisting of: hydrogen; hydrocarbyl; substituted hydrocarbyl;
 heteroatom-containing hydrocarbyl; and substituted heteroatom-containing hydrocarbyl;
 L is a neutral electron donor ligand;
 X and Y are heteroatoms selected from the group consisting of: N; O; S; and P;
 p is zero when X is O or S, and is 1 when X is N or P;
 35 q is zero when Y is O or S, and is 1 when Y is N or P;

-51-

Q^1 , Q^2 , Q^3 , and Q^4 are selected from the group consisting of: hydrocarbylene; substituted hydrocarbylene; heteroatom-containing hydrocarbylene; substituted heteroatom-containing hydrocarbylene; and $-(CO)-$;

w, x, y and z are independently zero or 1; and

5 R^3 , R^{3A} , R^4 , and R^{4A} are independently selected from the group consisting of hydrogen, hydrocarbyl, substituted hydrocarbyl, heteroatom-containing hydrocarbyl, and substituted heteroatom-containing hydrocarbyl,

wherein any two or more of X^1 , X^2 , L, R^1 , R^2 , R^3 , R^{3A} , R^4 , and R^{4A} can be taken together to form a chelating multidentate ligand,

10 wherein the first olefinic reactant has the structure of formula (VIII)



wherein:

n is zero or 1;

Z is a hydrocarbylene or a substituted or heteroatom-containing hydrocarbylene linking group; and

20 R^5 , R^6 and R^7 are independently selected from the group consisting of hydrogen, hydrocarbyl, substituted hydrocarbyl, heteroatom-containing hydrocarbyl, substituted heteroatom-containing hydrocarbyl, and $-(Z)_n$ -Fn,

wherein 'substituted' indicates that in the "substituted" compound, at least one hydrogen atom bound to a carbon (or other) atom is replaced with one or more non-hydrogen substituents, and

25 wherein 'heteroatom' is selected from the group consisting of: N, O, S, P, and Si.

2. The method of claim 1, wherein the first olefinic reactant is directly substituted with the functional group.

3. The method of claim 1, wherein the first olefinic reactant is indirectly substituted with the functional group.

30 4. The method of claim 1, wherein the second olefinic reactant is also directly substituted with a functional group Fn or indirectly substituted with a functional group Fn.

5. The method of claim 1, wherein:

X^1 and X^2 are anionic ligands, and are optionally linked to form a cyclic group;


L is a neutral electron donor ligand that is optionally linked to R^2 , X^1 , or X^2 through a spacer

35 moiety; and

-52-

R^{3A} and R^{4A} are optionally linked to form a cyclic group.


6. The method of claim 5, wherein w , x , y and z are zero, X and Y are N, and R^{3A} and R^{4A} are linked to form $-Q-$, such that the catalyst has the structure of formula (VIB)



wherein Q is a hydrocarbylene, substituted hydrocarbylene, heteroatom-containing hydrocarbylene, or substituted heteroatom-containing hydrocarbylene linker, and further wherein two or more substituents on adjacent atoms within Q may be linked to form an additional cyclic group.

7. The method of claim 6, wherein Q has the structure $-CR^{22}R^{22A}-CR^{23}R^{23A}-$ or $-CR^{22}=CR^{23}-$, wherein R^{22} , R^{22A} , R^{23} , and R^{23A} are independently selected from the group consisting of hydrogen, hydrocarbyl, substituted hydrocarbyl, heteroatom-containing hydrocarbyl, substituted heteroatom-containing hydrocarbyl, and functional groups, or wherein any two of R^{22} , R^{22A} , R^{23} , and R^{23A} may be linked together to form a substituted or unsubstituted, saturated or unsaturated ring, each respective functional group being selected from the group consisting of C_1 - C_{20} alkyl, C_2 - C_{20} alkenyl, C_2 - C_{20} alkynyl, aryl, C_1 - C_{20} carboxylate, C_1 - C_{20} alkoxy, C_2 - C_{20} alkenyloxy, C_2 - C_{20} alkynyloxy, aryloxy, C_2 - C_{20} alkoxycarbonyl, C_1 - C_{20} alkylthio, arylthio, C_1 - C_{20} alkylsulfonyl, and C_1 - C_{20} alkylsulfinyl, optionally substituted with one or more moieties selected from the group consisting of C_1 - C_{10} alkyl, C_1 - C_{10} alkoxy, aryl, hydroxyl, sulfhydryl, $-(CO)-H$, and halide.

8. The method of claim 7, wherein Q has the structure $-CR^{22}R^{22A}-CR^{23}R^{23A}-$, such that the catalyst has the structure of formula (VIC)



9. The method of claim 8, wherein:

-53-

M is Ru;

X^1 and X^2 are independently selected from the group consisting of: hydrogen; halide; C_1 - C_{20} alkyl; C_5 - C_{20} aryl; C_1 - C_{20} alkoxy; C_5 - C_{20} aryloxy; C_3 - C_{20} alkylidicketonate; C_5 - C_{20} arylidicketonate; C_2 - C_{20} alkoxy-carbonyl; C_5 - C_{20} aryloxy-carbonyl; C_2 - C_{20} acyl; C_1 - C_{20} alkylsulfonate; C_5 - C_{20} arylsulfonate; 5 C_1 - C_{20} alkylsulfanyl; C_5 - C_{20} arylsulfanyl; C_1 - C_{20} alkylsulfinyl; and C_5 - C_{20} arylsulfinyl, any of which, with the exception of halide, are optionally further substituted with one or more groups selected from the group consisting of: halide; C_1 - C_6 alkyl; C_1 - C_6 alkoxy; and phenyl;

R^1 is hydrogen and R^2 is selected from the group consisting of phenyl, vinyl, methyl, isopropyl, and t-butyl, optionally substituted with one or more moieties selected from the group 10 consisting of C_1 - C_6 alkyl, C_1 - C_6 alkoxy, phenyl, and a functional group Fn, wherein Fn is phosphonato, phosphoryl, phosphanyl, phosphino, sulfonate, C_1 - C_{20} alkylsulfanyl, C_5 - C_{20} arylsulfanyl, C_1 - C_{20} alkylsulfonyl, C_5 - C_{20} arylsulfonyl, C_1 - C_{20} alkylsulfinyl, C_5 - C_{20} arylsulfinyl, sulfonamido, amino, amido, imino, nitro, nitroso, hydroxyl, C_1 - C_{20} alkoxy, C_5 - C_{20} aryloxy, C_2 - C_{20} alkoxy-carbonyl, C_5 - C_{20} aryloxy-carbonyl, carboxyl, carboxylate, mercapto, formyl, C_1 - C_{20} thioester, cyano, cyanato, 15 carbamoyl, epoxy, styrenyl, silyl, silyloxy, silanyl, siloxazanyl, boronate, boryl, halogen, stannyl, and germlyl;

L is a neutral electron donor ligand selected from the group consisting of: phosphine; sulfonated phosphine; phosphite; phosphinite; phosphonite; arsine; stibine; ether; amine; amide; imine; sulfoxide; carboxyl; nitrosyl; pyridine; substituted pyridine; imidazole; substituted imidazole; 20 pyrazine; and thioether;

R^3 and R^4 are aromatic, substituted aromatic, heteroaromatic, substituted heteroaromatic, alicyclic, substituted alicyclic, heteroatom-containing alicyclic, or substituted heteroatom-containing alicyclic, composed of from one to five rings; and

R^{22} and R^{23} are hydrogen, and R^{22A} and R^{23A} are hydrogen, C_1 - C_6 alkyl or phenyl, or are linked 25 to form a cyclic group.

10. The method of claim 9, wherein:

R^1 is hydrogen, and R^2 is phenyl, vinyl, methyl, isopropyl, or t-butyl, optionally substituted with one or more moieties selected from the group consisting of C_1 - C_6 alkyl, C_1 - C_6 alkoxy, phenyl, and a functional group Fn, wherein Fn is phosphonato, phosphoryl, phosphanyl, phosphino, sulfonate, C_1 - 30 C_{20} alkylsulfanyl, C_5 - C_{20} arylsulfanyl, C_1 - C_{20} alkylsulfonyl, C_5 - C_{20} arylsulfonyl, C_1 - C_{20} alkylsulfinyl, C_5 - C_{20} arylsulfinyl, sulfonamido, amino, amido, imino, nitro, nitroso, hydroxyl, C_1 - C_{20} alkoxy, C_5 - C_{20} aryloxy, C_2 - C_{20} alkoxy-carbonyl, C_5 - C_{20} aryloxy-carbonyl, carboxyl, carboxylate, mercapto, formyl, C_1 - C_{20} thioester, cyano, cyanato, carbamoyl, epoxy, styrenyl, silyl, silyloxy, silanyl, siloxazanyl, boronate, boryl, halogen, stannyl, or germlyl; and

-54-

L is a phosphine of the formula $PR^{27}R^{28}R^{29}$, where R^{27} , R^{28} , and R^{29} are each independently aryl or C_1 - C_{10} alkyl.

11. The method of claim 10, wherein:

X^1 and X^2 are independently selected from the group consisting of halide, CF_3CO_2 , CH_3CO_2 , CFH_2CO_2 , $(CH_3)_3CO$, $(CF_3)_2(CH_3)CO$, $(CF_3)(CH_3)_2CO$, PhO , MeO , EtO , tosylate, mesylate, and trifluoromethanesulfonate;

L is selected from the group consisting of $-P(cyclohexyl)_3$, $-P(cyclopentyl)_3$, $-P(isopropyl)_3$, $-P(phenyl)_3$, $-P(phenyl)_2(R^7)$ and $-P(phenyl)(R^7)_2$, in which R^7 is C_1 - C_6 alkyl; and

R^3 and R^4 are the same and are either aromatic or C_7 - C_{12} alicyclic, if aromatic, each having the structure of formula (XI)



in which R^{24} , R^{25} , and R^{26} are each independently hydrogen, C_1 - C_{10} alkyl, C_1 - C_{10} alkoxy, aryl, substituted aryl, halogen, or a functional group, said functional group being selected from the group consisting of C_1 - C_{20} alkyl, C_2 - C_{20} alkenyl, C_2 - C_{20} alkynyl, aryl, C_1 - C_{20} carboxylate, C_1 - C_{20} alkoxy, C_2 - C_{20} alkenyloxy, C_2 - C_{20} alkynyloxy, aryloxy, C_2 - C_{20} alkoxy carbonyl, C_1 - C_{20} alkylthio, arylthio, C_1 - C_{20} alkylsulfonyl, and C_1 - C_{20} alkylsulfinyl, optionally substituted with one or more moieties selected from the group consisting of C_1 - C_{10} alkyl, C_1 - C_{10} alkoxy, aryl, hydroxyl, sulfhydryl, $-(CO)-H$, and halide.

12. The method of claim 11, wherein:

X^1 and X^2 are halide;

R^2 is phenyl or 2,2-dimethylvinyl;

R^3 and R^4 are mesityl, isopinocampheyl, or 2,4,2',6'-tetramethylbiphenyl;

L is selected from the group consisting of $-P(cyclohexyl)_3$ and $-P(cyclopentyl)_3$; and

R^{22} and R^{23} are hydrogen.


13. The method of claim 1, wherein n is zero.

14. The method of claim 13, wherein Fn is a phosphonate.

-55-

15. The method of claim 14, wherein R^5 , R^6 and R^7 are hydrogen, such that the first olefinic reactant is a vinylphosphonate having the structure of formula (XII)

5 (XII)



wherein R^{27} and R^{28} are C_1 - C_6 alkyl.

16. The method of claim 1, wherein n is 1.

10 17. The method of claim 16, wherein F_n is a phosphonate, a hydroxyl group, or boronate.

18. The method of claim 17, wherein Z is methylene, and R^5 , R^6 and R^7 are hydrogen.

19. The method of claim 1, wherein the second olefinic reactant has the molecular structure $R^{18}R^{19}C=CR^{20}R^{21}$ wherein R^{18} , R^{19} , R^{20} , and R^{21} are independently selected from the group consisting of hydrogen, hydrocarbyl, substituted hydrocarbyl, heteroatom-containing hydrocarbyl, substituted
15 heteroatom-containing hydrocarbyl, and functional groups F_n .

20. A method for synthesizing a plurality of functionalized olefins from a common olefinic reactant via a cross-metathesis reaction, comprising:


(a) contacting an olefinic substrate with a first olefinic reactant in the presence of a catalyst composed of a Group 8 transition metal alkylidene complex under conditions and for a time period
20 effective to allow cross-metathesis to occur, wherein the olefinic substrate is substituted with at least one $-(Z)_n-F_n$ moiety in which n is zero or 1, Z is a hydrocarbylene or a substituted or heteroatom-containing hydrocarbylene linking group, and F_n is selected from the group consisting of phosphonate, phosphoryl, phosphanyl, phosphino, sulfonate, C_1 - C_{20} alkylsulfanyl, C_5 - C_{20} arylsulfanyl, C_1 - C_{20} alkylsulfinyl, C_5 - C_{20} arylsulfinyl, sulfonamido, amino, imino, nitro, nitroso, hydroxyl,
25 unsubstituted C_1 - C_{20} alkoxy, C_5 - C_{20} aryloxy, C_5 - C_{20} aryloxycarbonyl, mercapto, C_1 - C_{20} thioester, cyano, cyanato, carbamoyl, styrenyl, siloxazanyl, boronate, boryl, stannyl, and germlyl;

(b) in a separate reaction, contacting the first olefinic reactant with a second olefinic reactant having a molecular structure that is different from that of the first olefinic reactant, in the presence of the Group 8 transition metal alkylidene complex, under conditions and for a time period effective to
30 allow cross-metathesis to occur; and

(c) optionally repeating step (b) with a plurality of olefinic reactants each having a different molecular structure,

wherein the first olefinic reactant has the structure of formula (VIII)

-56-



n is zero or 1;


Z is a hydrocarbylene or a substituted or heteroatom-containing hydrocarbylene linking group; and

10 R^5 , R^6 and R^7 are independently selected from the group consisting of hydrogen, hydrocarbyl, substituted hydrocarbyl, heteroatom-containing hydrocarbyl, substituted heteroatom-containing hydrocarbyl, and $-(Z)_n-Fn$,

wherein 'substituted' indicates that in the "substituted" compound, at least one hydrogen atom bound to a carbon (or other) atom is replaced with one or more non-hydrogen substituents,

15 wherein "heteroatom" is selected from the group consisting of: N, O, S, P, and Si and

wherein the catalyst has the structure of formula (VIA)



M is a Group 8 transition metal;

25 X^1 and X^2 may be the same or different, and are anionic ligands or polymers;

R^1 is selected from the group consisting of: hydrogen; hydrocarbyl; substituted hydrocarbyl; heteroatom-containing hydrocarbyl; substituted heteroatom-containing hydrocarbyl; and carboxyl;

R^2 is selected from the group consisting of: hydrogen; hydrocarbyl; substituted hydrocarbyl; heteroatom-containing hydrocarbyl; and substituted heteroatom-containing hydrocarbyl;

30 L is a neutral electron donor ligand;

X and Y are heteroatoms selected from the group consisting of: N; O; S; and P;

p is zero when X is O or S, and is 1 when X is N or P;

q is zero when Y is O or S, and is 1 when Y is N or P;

-57-

Q^1 , Q^2 , Q^3 , and Q^4 are selected from the group consisting of: hydrocarbylene; substituted hydrocarbylene; heteroatom-containing hydrocarbylene; substituted heteroatom-containing hydrocarbylene; and $-(CO)-$;

w, x, y and z are independently zero or 1; and

5 R^3 , R^{3A} , R^4 , and R^{4A} are independently selected from the group consisting of hydrogen, hydrocarbyl, substituted hydrocarbyl, heteroatom-containing hydrocarbyl, and substituted heteroatom-containing hydrocarbyl,

wherein any two or more of X^1 , X^2 , L, R^1 , R^2 , R^3 , R^{3A} , R^4 , and R^{4A} can be taken together to form a chelating multidentate ligand,

10 21. The method of claim 20, wherein the second olefinic reactant is also substituted with at least one $-(Z)_n$ -Fn moiety.

22. The method of claim 1, wherein the second olefinic reactant has the molecular structure $R^{18}R^{19}C=CR^{20}R^{21}$ wherein R^{18} , R^{19} , R^{20} , and R^{21} are independently selected from the group consisting of hydrogen, hydrocarbyl and $-(Z)_n$ -Fn, wherein n is zero or 1, Z is a hydrocarbylene or a substituted or heteroatom-containing hydrocarbylene linking group, and Fn is a functional group selected from
15 phosphonate, hydroxyl, boronate, and carboxylate.

23. The method of claim 22, wherein R^{18} , R^{19} and R^{20} are hydrogen.


24. The method of claim 23, wherein the functional group Fn for R^{21} is a carboxylate group.

25. The method of claim 1, wherein the second olefinic reactant has the molecular structure
20 $R^{18}R^{19}C=CR^{20}R^{21}$ wherein R^{18} , R^{19} , R^{20} , and R^{21} are independently selected from the group consisting of hydrogen, hydrocarbyl and $-(Z)_n$ -Fn, wherein n is zero or 1, Z is a hydrocarbylene or a substituted and/or heteroatom-containing hydrocarbylene linking group, and Fn is a functional group selected from the group consisting of phosphonate, hydroxyl and boronate.

26. The method of claim 25, wherein R^{18} , R^{19} and R^{20} are hydrogen.


25 27. The method of claim 26, wherein the functional group Fn for R^{21} is a carboxylate group.

28. A method for synthesizing a functionalized olefin of the formula



30 via a cross-metathesis reaction, comprising contacting (a) a first olefinic reactant of the formula

-58-



5


with (b) a second olefinic reactant of the formula



- 10 in the presence of (c) a catalyst composed of a Group 8 transition metal alkylidene complex under conditions and for a time period effective to allow cross-metathesis to occur, wherein the catalyst has the structure of formula (VIA)

15

(VIA)



20

in which:

M is a Group 8 transition metal;

X¹ and X² may be the same or different, and are anionic ligands or polymers;

- 25 R¹ is selected from the group consisting of hydrogen, hydrocarbyl, substituted hydrocarbyl, heteroatom-containing hydrocarbyl, substituted heteroatom-containing hydrocarbyl, and carboxyl;

R² is selected from the group consisting of hydrogen, hydrocarbyl, substituted hydrocarbyl, heteroatom-containing hydrocarbyl, and substituted heteroatom-containing hydrocarbyl;

L is a neutral electron donor ligand;

X and Y are heteroatoms selected from the group consisting of N, O, S, and P;

- 30 p is zero when X is O or S, and is 1 when X is N or P;

q is zero when Y is O or S, and is 1 when Y is N or P;

Q¹, Q², Q³, and Q⁴ are selected from hydrocarbylene, substituted hydrocarbylene, heteroatom-containing hydrocarbylene, substituted heteroatom-containing hydrocarbylene, and -(CO)-;

w, x, y and z are independently zero or 1; and

- 35 R³, R^{3A}, R⁴, and R^{4A} are independently selected from the group consisting of hydrogen,

-59-

hydrocarbyl, substituted hydrocarbyl, heteroatom-containing hydrocarbyl, and substituted heteroatom-containing hydrocarbyl,

wherein any two or more of X^1 , X^2 , L, R^1 , R^2 , R^3 , R^{3A} , R^4 , and R^{4A} can be taken together to form a chelating multidentate ligand, and

5 wherein 'substituted' indicates that in the "substituted" compound, at least one hydrogen atom bound to a carbon (or other) atom is replaced with one or more non-hydrogen substituents, and

wherein 'heteroatom' is selected from the group consisting of: N, O, S, P, and Si.

29. The method of claim 28, wherein:


X^1 and X^2 are anionic ligands, and are optionally linked to form a cyclic group;

10 L is a neutral electron donor ligand that is optionally linked to R^2 , X^1 , or X^2 through a spacer moiety; and

R^{3A} and R^{4A} are optionally linked to form a cyclic group.

30. The method of claim 29, wherein w, x, y and z are zero, X and Y are N, and R^{3A} and R^{4A} are linked to form -Q-, such that the catalyst has the structure of formula (VIB)

15



wherein Q is a hydrocarbylene, substituted hydrocarbylene, heteroatom-containing hydrocarbylene, or substituted heteroatom-containing hydrocarbylene linker, and further wherein two or more substituents on adjacent atoms within Q may be linked to form an additional cyclic group.


25 31. The method of claim 30, wherein Q has the structure $-CR^{22}R^{22A}-CR^{23}R^{23A}-$ or $-CR^{22}=CR^{23}-$, wherein R^{22} , R^{22A} , R^{23} , and R^{23A} are independently selected from the group consisting of hydrogen, hydrocarbyl, substituted hydrocarbyl, heteroatom-containing hydrocarbyl, substituted heteroatom-containing hydrocarbyl, a functional group, said functional group selected from the group consisting of phosphonato, phosphoryl, phosphanyl, phosphino, sulfonato, C_1 - C_{20} alkylsulfanyl, C_5 - C_{20} arylsulfanyl, C_1 - C_{20} alkylsulfonyl, C_5 - C_{20} arylsulfonyl, C_1 - C_{20} alkylsulfinyl, C_5 - C_{20} arylsulfinyl, sulfonamido, amino, amido, imino, nitro, nitroso, hydroxyl, C_1 - C_{20} alkoxy, C_5 - C_{20} aryloxy, C_2 - C_{20} alkoxycarbonyl, C_5 - C_{20} aryloxycarbonyl, carboxyl, carboxylato, mercapto, formyl, C_1 - C_{20} thioester, cyano, cyanato, carbamoyl, epoxy, styrenyl, silyl, silyloxy, silanyl, siloxazanyl, boronato, and boryl; and a metal-containing or metalloid-containing group, or wherein any two of R^{22} , R^{22A} , R^{23} , and R^{23A}

30

35 may be linked together to form a substituted or unsubstituted, saturated or unsaturated ring.

-60-

32. The method of claim 31, wherein Q has the structure $-\text{CR}^{22}\text{R}^{22\text{A}}-\text{CR}^{23}\text{R}^{23\text{A}}-$, such that the catalyst has the structure of formula (VIC)



33. The method of claim 32, wherein:

M is Ru;

X^1 and X^2 are independently selected from the group consisting of hydrogen, halide, C_1 - C_{20} alkyl, C_5 - C_{20} aryl, C_1 - C_{20} alkoxy, C_5 - C_{20} aryloxy, C_3 - C_{20} alkyldiketonate, C_5 - C_{20} aryldiketonate, C_2 - C_{20} alkoxy carbonyl, C_5 - C_{20} aryloxy carbonyl, C_2 - C_{20} acyl, C_1 - C_{20} alkylsulfonate, C_5 - C_{20} arylsulfonate, C_1 - C_{20} alkylsulfanyl, C_5 - C_{20} arylsulfanyl, C_1 - C_{20} alkylsulfinyl, or C_5 - C_{20} arylsulfinyl, any of which, with the exception of halide, are optionally further substituted with one or more groups selected from the group consisting of halide, C_1 - C_6 alkyl, C_1 - C_6 alkoxy, and phenyl;

R^1 is hydrogen and R^2 is selected from the group consisting of hydrogen, C_1 - C_{20} alkyl, C_7 - C_{20} alkenyl, and aryl optionally substituted with one or more moieties selected from the group consisting of C_1 - C_6 alkyl, C_1 - C_6 alkoxy, phenyl, and a functional group Fn, wherein Fn is phosphonato, phosphoryl, phosphanyl, phosphino, sulfonate, C_1 - C_{20} alkylsulfanyl, C_5 - C_{20} arylsulfanyl, C_1 - C_{20} alkylsulfonyl, C_5 - C_{20} arylsulfonyl, C_1 - C_{20} alkylsulfinyl, C_5 - C_{20} arylsulfinyl, sulfonamido, amino, amido, imino, nitro, nitroso, hydroxyl, C_1 - C_{20} alkoxy, C_5 - C_{20} aryloxy, C_2 - C_{20} alkoxy carbonyl, C_5 - C_{20} aryloxy carbonyl, carboxyl, carboxylate, mercapto, formyl, C_1 - C_{20} thioester, cyano, cyanate, carbamoyl, epoxy, styrenyl, silyl, silyloxy, silanyl, siloxazanyl, boronate, boryl, halogen, stannyl, and germlyl;

L is a neutral electron donor ligand selected from the group consisting of phosphine, sulfonated phosphine, phosphite, phosphinite, phosphonite, arsine, stibine, ether, amine, amide, imine, sulfoxide, carboxyl, nitrosyl, pyridine, substituted pyridine, imidazole, substituted imidazole, pyrazine, and thioether;

R^3 and R^4 are aromatic, substituted aromatic, heteroaromatic, substituted heteroaromatic, alicyclic, substituted alicyclic, heteroatom-containing alicyclic, or substituted heteroatom-containing alicyclic, composed of from one to five rings; and

R^{22} and R^{23} are hydrogen, and $\text{R}^{22\text{A}}$ and $\text{R}^{23\text{A}}$ are hydrogen, C_1 - C_6 alkyl or phenyl, or are linked to form a cyclic group.

-61-

34. The method of claim 33, wherein:

R^1 is hydrogen, and R^2 is hydrogen, phenyl, vinyl, methyl, isopropyl, or t-butyl, optionally substituted with one or more moieties selected from the group consisting of C_1 - C_6 alkyl, C_1 - C_6 alkoxy, phenyl, and a functional group Fn, wherein Fn is phosphonato, phosphoryl, phosphanyl, phosphino, sulfonato, C_1 - C_{20} alkylsulfanyl, C_5 - C_{20} arylsulfanyl, C_1 - C_{20} alkylsulfonyl, C_5 - C_{20} arylsulfonyl, C_1 - C_{20} alkylsulfinyl, C_5 - C_{20} arylsulfinyl, sulfonamido, amino, amido, imino, nitro, nitroso, hydroxyl, C_1 - C_{20} alkoxy, C_5 - C_{20} aryloxy, C_2 - C_{20} alkoxycarbonyl, C_5 - C_{20} aryloxycarbonyl, carboxyl, carboxylato, mercapto, formyl, C_1 - C_{20} thioester, cyano, cyanato, carbamoyl, epoxy, styrenyl, silyl, silyloxy, silanyl, siloxazanyl, boronato, boryl, halogen, stannyl, and germlyl; and


L is a phosphine of the formula $PR^{27}R^{28}R^{29}$, where R^{27} , R^{28} , and R^{29} are each independently aryl or C_1 - C_{10} alkyl.

35. The method of claim 34, wherein:

X^1 and X^2 are independently selected from the group consisting of halide, CF_3CO_2 , CH_3CO_2 , CFH_2CO_2 , $(CH_3)_3CO$, $(CF_3)_2(CH_3)CO$, $(CF_3)(CH_3)_2CO$, PhO, MeO, EtO, tosylate, mesylate, and trifluoromethanesulfonate;

L is selected from the group consisting of -P(cyclohexyl)₃, -P(cyclopentyl)₃, -P(isopropyl)₃, -P(phenyl)₃, -P(phenyl)₂(R^7) and -P(phenyl)(R^7)₂, in which R^7 is C_1 - C_6 alkyl; and

R^3 and R^4 are the same and are either aromatic or C_7 - C_{12} alicyclic, if aromatic, each having the structure of formula (XI)



in which R^{24} , R^{25} , and R^{26} are each independently hydrogen, C_1 - C_{10} alkyl, C_1 - C_{10} alkoxy, aryl, substituted aryl, halogen, or a functional group, said functional group selected from the group consisting of phosphonato, phosphoryl, phosphanyl, phosphino, sulfonato, C_1 - C_{20} alkylsulfanyl, C_5 - C_{20} arylsulfanyl, C_1 - C_{20} alkylsulfonyl, C_5 - C_{20} arylsulfonyl, C_1 - C_{20} alkylsulfinyl, C_5 - C_{20} arylsulfinyl, sulfonamido, amino, amido, imino, nitro, nitroso, hydroxyl, C_1 - C_{20} alkoxy, C_5 - C_{20} aryloxy, C_2 - C_{20} alkoxycarbonyl, C_5 - C_{20} aryloxycarbonyl, carboxyl, carboxylato, mercapto, formyl, C_1 - C_{20} thioester, cyano, cyanato, carbamoyl, epoxy, styrenyl, silyl, silyloxy, silanyl, siloxazanyl, boronato, and boryl, or a metal-containing or metalloid-containing group.

36. The method of claim 35, wherein:

X^1 and X^2 are halide;

-62-

R^2 is hydrogen or 2,2-dimethylvinyl;

R^3 and R^4 are mesityl, isopinocampheyl, or 2,4,2',6'-tetramethylbiphenyl;

L is selected from the group consisting of $-P(\text{cyclohexyl})_3$ and $-P(\text{cyclopentyl})_3$; and

R^{22} and R^{23} are hydrogen.