

(12) 特許協力条約に基づいて公開された国際出願

(19) 世界知的所有権機関
国際事務局

(43) 国際公開日
2018年11月29日(29.11.2018)



(10) 国際公開番号

WO 2018/216795 A1

(51) 国際特許分類:

A61K 31/551 (2006.01) A61K 31/4709 (2006.01)
A61K 31/203 (2006.01) A61K 31/4725 (2006.01)
A61K 31/282 (2006.01) A61K 31/4745 (2006.01)
A61K 31/337 (2006.01) A61K 31/495 (2006.01)
A61K 31/40 (2006.01) A61K 31/498 (2006.01)
A61K 31/401 (2006.01) A61K 31/4995 (2006.01)
A61K 31/4025 (2006.01) A61K 31/502 (2006.01)
A61K 31/403 (2006.01) A61K 31/517 (2006.01)
A61K 31/404 (2006.01) A61K 31/519 (2006.01)
A61K 31/407 (2006.01) A61K 31/538 (2006.01)
A61K 31/416 (2006.01) A61K 31/5386 (2006.01)
A61K 31/4184 (2006.01) A61K 31/55 (2006.01)
A61K 31/4192 (2006.01) A61K 31/704 (2006.01)
A61K 31/423 (2006.01) A61K 31/7048 (2006.01)
A61K 31/428 (2006.01) A61K 31/706 (2006.01)
A61K 31/437 (2006.01) A61K 31/7068 (2006.01)
A61K 31/438 (2006.01) A61K 31/7084 (2006.01)
A61K 31/439 (2006.01) A61K 33/24 (2006.01)
A61K 31/4439 (2006.01) A61K 45/00 (2006.01)
A61K 31/445 (2006.01) A61P 35/00 (2006.01)
A61K 31/4468 (2006.01) A61P 43/00 (2006.01)
A61K 31/46 (2006.01)

(71) 出願人: 大鵬薬品工業株式会社
(TAIHO PHARMACEUTICAL CO., LTD.) [JP/
JP]; 〒1018444 東京都千代田区神田錦町
1-27 Tokyo (JP).

(72) 発明者: 長田 亜希子(OSADA, Akiko); 〒3002611
茨城県つくば市大久保3 大鵬薬品工
業株式会社内 Ibaraki (JP).

(74) 代理人: 特許業務法人三枝国際特許事務
所(SAEGUSA & PARTNERS); 〒5410045 大阪
府大阪市中央区道修町1-7-1 北
浜T N Kビル Osaka (JP).

(81) 指定国(表示のない限り、全ての種類の国内保
護が可能): AE, AG, AL, AM, AO, AT, AU, AZ,
BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH,
CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO,
DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT,
HN, HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH,
KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY,
MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ,
NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT,
QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL,
SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA,
UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) 指定国(表示のない限り、全ての種類の広域保
護が可能): ARIPO (BW, GH, GM, KE, LR, LS,
MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM,
ZW), ユーラシア (AM, AZ, BY, KG, KZ, RU, TJ,
TM), ヨーロッパ (AL, AT, BE, BG, CH, CY, CZ,
DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT,

(21) 国際出願番号: PCT/JP2018/020101

(22) 国際出願日: 2018年5月25日(25.05.2018)


(25) 国際出願の言語: 日本語

(26) 国際公開の言語: 日本語

(30) 優先権データ:
特願 2017-104841 2017年5月26日(26.05.2017) JP

(54) Title: ANTI-TUMOR EFFECT POTENTIATOR USING NOVEL BIPHENYL COMPOUND

(54) 発明の名称: 新規なビフェニル化合物を用いた抗腫瘍効果増強剤



(57) Abstract: An anti-cancer agent containing a biphenyl compound represented by general formula (I) [wherein ring A, ring B, R1 to R6, l, m and n are as defined in the description] and having a LSD1-inhibiting activity or a salt thereof and another anti-tumor agent which are administered in combination.

(57) 要約: 一般式 (I) [式中、環 A、環 B、R1 ~ R6、l、m、n は明細書中で定義した通りの意味を有する] で表される LSD1 阻害活性を有するビフェニル化合物又はその塩と、他の抗腫瘍剤を併用投与する抗腫瘍剤。

LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS,
SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM,
GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

添付公開書類：

- 一 国際調査報告（条約第21条(3)）

DESCRIPTION

Title of Invention: ANTI-TUMOR EFFECT POTENTIATOR USING NOVEL
BIPHENYL COMPOUND

5 Technical Field
 [0001]

 The present invention relates to an antitumor
formulation comprising a combination of a biphenyl compound or a
salt thereof and one or more other antitumor agents, and an
10 antitumor effect potentiator for the one or more other antitumor
agents.

Background Art
 [0002]

15 Histone methylation modification is one of the
epigenetic mechanisms, which regulate gene expressions. Histone
methylation modification regulates various processes including
cellular maintenance, growth, and differentiation.
 [0003]

20 LSD1 (KDM1A), one of the enzymes that regulate histone
methylation modification, is an FAD (flavin adenine
dinucleotide)-dependent histone demethylase, and mainly
demethylates the lysine residue at position 4 (K4) and the lysine
residue at position 9 (K9) on histone H3 (Non-patent Literature
25 (NPL) 1). With such functions, LSD1 is believed to positively or
negatively regulate various gene transcriptions, and regulate
stem cell self-renewal and cell differentiation in each normal
tissue.
 [0004]

30 In general, abnormalities in cell self-renewal capacity
or differentiation are believed to lead to cell cancerization.
Thus, aberrant control of LSD1, which plays a key role in these
processes, can possibly cause cell cancerization. In fact, in
terms of various solid and blood cancers, many reports have been
35 made regarding the correlation of overexpression of LSD1 and

their prognosis (NPL 2). Further, in cell lines from carcinomas or in non-clinical models, LSD1 inhibition has been reported to have resulted in induction of cellular differentiation, growth inhibition, and an *in vivo* antitumor effect (NPL 3 and NPL 4), which strongly suggests that LSD1 serves as one of the important target molecules in cancer therapy. These carcinomas in which LSD1 is involved, such as small-cell lung cancer (SCLC) and acute myelocytic leukemia (AML), have an extremely short lifetime, and existing therapeutic methods cannot achieve a satisfactory therapeutic effect.

[0005]

Accordingly, LSD1 inhibitory drugs are expected to provide effective therapeutic means based on novel mechanisms to treat intractable cancers, for which no therapeutic methods currently exist.

[0006]

Further, according to some reports, LSD1, which is involved in neuron programs and functions, can also possibly serve as a target in the treatment of diseases other than cancers, such as Alzheimer's disease, Huntington's disease, Rett syndrome, and other cranial nerve diseases (NPL 2); Herpesvirus infections, in which LSD1 function has been implicated (NPL 5); and sickle cell diseases (NPL 6).

[0007]

There are some reports regarding antitumor effects of combinations of LSD1 inhibitors with other antitumor agents. For example, an antitumor effect of a combination of LSD1 inhibitor SP2509 with histone deacetylase inhibitor panobinostat on acute myelocytic leukemia has been reported (NPL 7). In addition, an antitumor effect of a combination of LSD1 inhibitor INCB059872 with all-trans retinoic acid (ATRA) on acute myelocytic leukemia has been reported (NPL 8).

Citation List

Non-patent Literature

[0008]

NPL 1: Biochim. Biophys. Acta, 1829 (10), pp. 981-986 (2013)

NPL 2: Epigenomics, 7 (4), pp. 609-626 (2015)

NPL 3: Cancer Cell, 21 (4), pp. 473-487 (2012)

5 NPL 4: Cancer Cell, 28 (1), pp. 57-69 (2015)

NPL 5: Sci. Transl. Med., 6 (265), 265ra169 (2014)

NPL 6: Nat. Med., 19 (3), pp. 291-294 (2013)

NPL 7: Leukemia, 28(11), pp. 2155-2164 (2014)

NPL 8: Cancer Res., 74 (14 Suppl) Abstract, 4696 (2016)

10

Summary of Invention

Technical Problem

[0009]

15 An object of the present invention is to provide a method for potentiating an antitumor effect using a compound or a salt thereof that potently inhibits LSD1.

Solution to Problem

[0010]


20 The present inventors conducted extensive research to achieve the above object, and found that a compound represented by the following Formula (I) or a salt thereof has excellent antitumor effect potentiating action in combined use with one or more other compounds having an antitumor effect (other antitumor
25 agents). The present invention has been accomplished based on this finding.

[0011]

More specifically, the present invention provides the following:

30 Item 1. An antitumor formulation comprising a biphenyl compound represented by Formula (I) or a salt thereof and one or more other antitumor agents that are administered in combination:

[0012]



[0013]

wherein

ring A represents a monocyclic, bridged cyclic, or spirocyclic 4-
 5 to 14-membered nitrogen-containing saturated heterocyclic group
 having 1 to 3 nitrogen atoms, 0 to 1 sulfur atoms, and 0 to 2
 oxygen atoms as heteroatoms;

ring B represents monocyclic or bicyclic 5- to 14-membered
 unsaturated hydrocarbon or a monocyclic or bicyclic 5- to 14-
 10 membered unsaturated heterocyclic group that may be substituted
 with oxo, that has 0 to 4 nitrogen atoms, 0 to 2 sulfur atoms,
 and 0 to 3 oxygen atoms as heteroatoms, and that has at least one
 of nitrogen, sulfur, and oxygen;

R1 represents nitro or cyano;

15 R2 represents halogen;

R3 represents amino, mono- or di(C1-C6 alkyl)amino, (C3-C7
 cycloalkyl)amino, or C1-C6 alkyl;

R4 represents halogen, nitro, cyano, carboxy, substituted or
 unsubstituted C1-C8 alkyl, substituted or unsubstituted C2-C6
 20 alkenyl, substituted or unsubstituted C1-C6 alkoxy, substituted
 or unsubstituted C3-C7 cycloalkyl, mono- or di(C1-C6 alkyl)amino,
 or substituted or unsubstituted carbamoyl;

when at least one R4 represents substituted C1-C8 alkyl,
 substituted C2-C6 alkenyl, substituted C1-C6 alkoxy, substituted
 25 C3-C7 cycloalkyl, or substituted carbamoyl, the substituent is
 halogen, carboxy, C1-C6 alkoxy, hydroxy, C1-C6 alkyl that may be
 substituted with hydroxy, monocyclic 5- to 10-membered
 unsaturated hydrocarbon, carbamoyl that may be substituted with
 C1-C6 alkyl or monocyclic 5- to 10-membered unsaturated

hydrocarbon, (C2-C7 acyl)oxy, amino that may be substituted with C1-C6 alkyl or C2-C7 acyl, C3-C7 cycloalkyl that may be substituted with hydroxy, or (C1-C6 alkoxy) (C1-C6 alkyl); when two or more of the substituents are present, the substituents may

5 be identical or different;

R5 and R6 may be identical or different, and each represent hydrogen or C1-C6 alkyl; R5 and R6 may otherwise form oxo or thioxo together;

when R5 and R6 are identical or different, and each represent
10 hydrogen or C1-C6 alkyl, at least one of R5 and R6 is hydrogen;

l represents an integer of 0 to 2;

m represents an integer of 0 to 2;

n represents an integer of 0 to 5;

when l is 2, two R2s may be identical or different;

15 when m is 2, two R3s may be identical or different;

when n is 2 to 5, two to five R4s may be identical or different.


[0014]

Item 2. The antitumor formulation according to Item 1, wherein the biphenyl compound satisfies the following conditions

20 in Formula (I):

ring A represents pyrrolidinyl, piperidinyl, piperazinyl, azepanyl, diazepanyl,

[0015]



25 [0016]

, 2,7-diazaspiro[3.4]octanyl, 3,7-diazaspiro[3.4]octanyl, 2,7-diazaspiro[3.5]nonanyl, 2,8-diazaspiro[3.5]nonanyl, 3,7-diazaspiro[3.5]nonanyl, 3,8-diazaspiro[4.4]nonanyl, 3,8-diazaspiro[4.5]decanyl, or 9-oxa-diazaspiro[3.5]nonanyl;

30 ring B represents phenyl, naphthyl, pyridyl, pyrazolopyridyl, pyrazolopyrimidinyl, indolyl, indolinyl, 2-oxo-indolinyl,

indazolyl, benzoimidazolyl, benzoisoxazolyl, benzothiazolyl, benzotriazolyl, imidazopyridinyl, quinolinyl, isoquinolinyl, quinazolinyl, quinoxalinyl, phthalazinyl, 2-oxo-2,3-dihydrobenzo[d]oxazolyl, 1,3-dihydroisobenzofuranyl,

5 dihydrobenzooxazinyl, benzodioxolyl, dihydrobenzodioxynyl, or 2-oxo-2,3-dihydrobenzo[d]thiazolyl;

R1 represents nitro or cyano;

R2 represents fluorine, and is present at the ortho position relative to R1 on the phenyl;

10 R3 represents amino, methylamino, ethylamino, isopropylamino, dimethylamino, cyclobutylamino, or methyl (when two or more R3s are present, R3s may be identical or different);

R4 represents fluorine, chlorine, bromine, iodine, nitro, cyano, carboxy, methyl, ethyl, n-propyl, isopropyl, tert-butyl,

15 difluoromethyl, trifluoromethyl, fluoroethyl, aminoethyl, hydroxymethyl, hydroxyethyl, hydroxypropyl, hydroxydimethylethyl, hydroxymethylpropyl, hydroxymethylbutyl, hydroxyethylbutyl, carboxymethyl, carbamoylmethyl, methylcarbamoylmethyl, dimethylcarbamoylmethyl, acetylaminoethyl, methoxyethyl,

20 hydroxycyclopropylmethyl, hydroxycyclopropylethyl, hydroxycyclobutylmethyl, methylcarbonyloxyethyl, isobutenyl, methoxy, hydroxypropoxy, cyclopropyl, hydroxymethyl cyclopropyl, methoxymethyl cyclopropyl, hydroxycyclopropyl cyclopropyl, phenylcarbamoyl cyclopropyl, benzyloxy, dimethylamino, carbamoyl, methylcarbamoyl, or dimethylcarbamoyl (when two or more R4s are
25 present, R4s may be identical or different);

R5 and R6 may be identical or different, and each represent hydrogen or C1-C6 alkyl; R5 and R6 may otherwise form oxo or thioxo together;

30 when R5 and R6 are identical or different, and each represent hydrogen or C1-C6 alkyl, at least one of R5 and R6 is hydrogen; n represents an integer of 0 to 3; and

when n is 2 to 3, two to three R4s may be identical or different.
[0017]

35 Item 3. The antitumor formulation according to Item 1

or 2, wherein the biphenyl compound is selected from the following (1) to (7):

(1) 4-[5-[(3S)-3-aminopyrrolidine-1-carbonyl]-2-[2-fluoro-4-(2-hydroxy-2-methyl-propyl)phenyl]phenyl]-2-fluoro-benzonitrile;

5 (2) 4-[5-[(3-endo)-3-amino-8-azabicyclo[3.2.1]octane-8-carbonyl]-2-[2-fluoro-4-(2-hydroxy-2-methyl-propyl)phenyl]phenyl]-2-fluoro-benzonitrile;

(3) 5'-((1S,2S,4R)-rel-2-amino-7-azabicyclo[2.2.1]heptane-7-carbonyl)-2'',3-difluoro-4''-(2-hydroxy-2-methylpropyl)-
10 [1,1':2',1''-terphenyl]-4-carbonitrile-isomer-B;

(4) 5'-((1S,2S,4R)-rel-2-amino-7-azabicyclo[2.2.1]heptane-7-carbonyl)-2'-(6,7-difluoro-1-(2-hydroxy-2-methylpropyl)-1H-benzo[d][1,2,3]triazol-5-yl)-3-fluoro-[1,1'-biphenyl]-4-carbonitrile-isomer-B;

15 (5) 5'-((1S,2S,4R)-rel-2-amino-7-azabicyclo[2.2.1]heptane-7-carbonyl)-2'-(1-(2-ethyl-2-hydroxybutyl)-6,7-difluoro-1H-benzo[d][1,2,3]triazol-5-yl)-3-fluoro-[1,1'-biphenyl]-4-carbonitrile-isomer-X;

(6) 5'-((1S,2S,4R)-rel-2-amino-7-azabicyclo[2.2.1]heptane-7-carbonyl)-2'-(7-bromo-6-fluoro-1-(2-hydroxy-2-methylpropyl)-1H-benzo[d][1,2,3]triazol-5-yl)-3-fluoro-[1,1'-biphenyl]-4-carbonitrile-isomer-X;

(7) 5'-(((1S,2S,4R)-rel-2-amino-7-azabicyclo[2.2.1]heptan-7-yl)methyl)-2'-(6,7-difluoro-1-(2-hydroxy-2-methylpropyl)-1H-benzo[d][1,2,3]triazol-5-yl)-3-fluoro-[1,1'-biphenyl]-4-carbonitrile-isomer-X.
25

[0018]

Item 4. The antitumor formulation according to any one of Items 1 to 3, wherein the one or more other antitumor agents
30 are at least one member selected from antimetabolites, antitumor antibiotics, molecular target drugs, platinum-based drugs, and plant alkaloid-based drugs.

[0019]


Item 5. The antitumor formulation according to any one
35 of Items 1 to 3, wherein the one or more other antitumor agents

are at least one member selected from cytarabine, azacitidine, decitabine, guadecitabine, daunorubicin, tretinoin (ATRA), RG7388, carboplatin, cisplatin, paclitaxel, irinotecan (SN-38), and etoposide.

5 [0020]

Item 6. An antitumor effect potentiator for one or more other antitumor agents, the antitumor effect potentiator comprising as an active ingredient a biphenyl compound represented by Formula (I) or a salt thereof:

10 [0021]



[0022]

wherein

15 ring A represents a monocyclic, bridged cyclic, or spirocyclic 4- to 14-membered nitrogen-containing saturated heterocyclic group having 1 to 3 nitrogen atoms, 0 to 1 sulfur atoms, and 0 to 2 oxygen atoms as heteroatoms;

ring B represents monocyclic or bicyclic 5- to 14-membered
20 unsaturated hydrocarbon or a monocyclic or bicyclic 5- to 14-membered unsaturated heterocyclic group that may be substituted with oxo, that has 0 to 4 nitrogen atoms, 0 to 2 sulfur atoms, and 0 to 3 oxygen atoms as heteroatoms, and that has at least one of nitrogen, sulfur, and oxygen;

25 R1 represents nitro or cyano;

R2 represents halogen;


R3 represents amino, mono- or di(C1-C6 alkyl)amino, (C3-C7 cycloalkyl)amino, or C1-C6 alkyl;

R4 represents halogen, nitro, cyano, carboxy, substituted or

unsubstituted C1-C8 alkyl, substituted or unsubstituted C2-C6 alkenyl, substituted or unsubstituted C1-C6 alkoxy, substituted or unsubstituted C3-C7 cycloalkyl, mono- or di(C1-C6 alkyl)amino, or substituted or unsubstituted carbamoyl;

- 5 when at least one R4 represents substituted C1-C8 alkyl, substituted C2-C6 alkenyl, substituted C1-C6 alkoxy, substituted C3-C7 cycloalkyl, or substituted carbamoyl, the substituent is halogen, carboxy, C1-C6 alkoxy, hydroxy, C1-C6 alkyl that may be substituted with hydroxy, monocyclic 5- to 10-membered
- 10 unsaturated hydrocarbon, carbamoyl that may be substituted with C1-C6 alkyl or monocyclic 5- to 10-membered unsaturated hydrocarbon, (C2-C7 acyl)oxy, amino that may be substituted with C1-C6 alkyl or C2-C7 acyl, C3-C7 cycloalkyl that may be substituted with hydroxy, or (C1-C6 alkoxy)(C1-C6 alkyl); when
- 15 two or more of the substituents are present, the substituents may be identical or different;
- R5 and R6 may be identical or different, and each represent hydrogen or C1-C6 alkyl; R5 and R6 may otherwise form oxo or thioxo together;
- 20 when R5 and R6 are identical or different, and each represent hydrogen or C1-C6 alkyl, at least one of R5 and R6 is hydrogen;
- l represents an integer of 0 to 2;
- m represents an integer of 0 to 2;
- n represents an integer of 0 to 5;
- 25 when l is 2, two R2s may be identical or different;
- when m is 2, two R3s may be identical or different;
- when n is 2 to 5, two to five R4s may be identical or different.
- [0023]

- Item 7. An antitumor agent comprising a biphenyl
- 30 compound or a salt thereof for treating a cancer patient administered one or more other antitumor agents, wherein the biphenyl compound is represented by Formula (I):
- [0024]



[0025]

wherein

ring A represents a monocyclic, bridged cyclic, or spirocyclic 4-
 5 to 14-membered nitrogen-containing saturated heterocyclic group
 having 1 to 3 nitrogen atoms, 0 to 1 sulfur atoms, and 0 to 2
 oxygen atoms as heteroatoms;

ring B represents monocyclic or bicyclic 5- to 14-membered
 unsaturated hydrocarbon or a monocyclic or bicyclic 5- to 14-
 10 membered unsaturated heterocyclic group that may be substituted
 with oxo, that has 0 to 4 nitrogen atoms, 0 to 2 sulfur atoms,
 and 0 to 3 oxygen atoms as heteroatoms, and that has at least one
 of nitrogen, sulfur, and oxygen;

R1 represents nitro or cyano;

15 R2 represents halogen;

R3 represents amino, mono- or di(C1-C6 alkyl)amino, (C3-C7
 cycloalkyl)amino, or C1-C6 alkyl;

R4 represents halogen, nitro, cyano, carboxy, substituted or
 unsubstituted C1-C8 alkyl, substituted or unsubstituted C2-C6
 20 alkenyl, substituted or unsubstituted C1-C6 alkoxy, substituted
 or unsubstituted C3-C7 cycloalkyl, mono- or di(C1-C6 alkyl)amino,
 or substituted or unsubstituted carbamoyl;

when at least one R4 represents substituted C1-C8 alkyl,
 substituted C2-C6 alkenyl, substituted C1-C6 alkoxy, substituted
 25 C3-C7 cycloalkyl, or substituted carbamoyl, the substituent is
 halogen, carboxy, C1-C6 alkoxy, hydroxy, C1-C6 alkyl that may be
 substituted with hydroxy, monocyclic 5- to 10-membered
 unsaturated hydrocarbon, carbamoyl that may be substituted with
 C1-C6 alkyl or monocyclic 5- to 10-membered unsaturated

hydrocarbon, (C2-C7 acyl)oxy, amino that may be substituted with C1-C6 alkyl or C2-C7 acyl, C3-C7 cycloalkyl that may be substituted with hydroxy, or (C1-C6 alkoxy) (C1-C6 alkyl); when two or more of the substituents are present, the substituents may

5 be identical or different;

R5 and R6 may be identical or different, and each represent hydrogen or C1-C6 alkyl; R5 and R6 may otherwise form oxo or thioxo together;

when R5 and R6 are identical or different, and each represent
10 hydrogen or C1-C6 alkyl, at least one of R5 and R6 is hydrogen;

l represents an integer of 0 to 2;

m represents an integer of 0 to 2;

n represents an integer of 0 to 5;

when l is 2, two R2s may be identical or different;


15 when m is 2, two R3s may be identical or different;

when n is 2 to 5, two to five R4s may be identical or different.

[0026]

Item 8. An antitumor formulation comprising a combination of a biphenyl compound or a salt thereof and one or
20 more other antitumor agents, wherein the biphenyl compound is represented by Formula (I):

[0027]



[0028]

25 wherein

ring A represents a monocyclic, bridged cyclic, or spirocyclic 4- to 14-membered nitrogen-containing saturated heterocyclic group having 1 to 3 nitrogen atoms, 0 to 1 sulfur atoms, and 0 to 2 oxygen atoms as heteroatoms;

ring B represents monocyclic or bicyclic 5- to 14-membered unsaturated hydrocarbon or a monocyclic or bicyclic 5- to 14-membered unsaturated heterocyclic group that may be substituted with oxo, that has 0 to 4 nitrogen atoms, 0 to 2 sulfur atoms,
5 and 0 to 3 oxygen atoms as heteroatoms, and that has at least one of nitrogen, sulfur, and oxygen;

R1 represents nitro or cyano;

R2 represents halogen;

R3 represents amino, mono- or di(C1-C6 alkyl)amino, (C3-C7
10 cycloalkyl)amino, or C1-C6 alkyl;

R4 represents halogen, nitro, cyano, carboxy, substituted or unsubstituted C1-C8 alkyl, substituted or unsubstituted C2-C6 alkenyl, substituted or unsubstituted C1-C6 alkoxy, substituted or unsubstituted C3-C7 cycloalkyl, mono- or di(C1-C6 alkyl)amino,
15 or substituted or unsubstituted carbamoyl;

when at least one R4 represents substituted C1-C8 alkyl, substituted C2-C6 alkenyl, substituted C1-C6 alkoxy, substituted C3-C7 cycloalkyl, or substituted carbamoyl, the substituent is halogen, carboxy, C1-C6 alkoxy, hydroxy, C1-C6 alkyl that may be
20 substituted with hydroxy, monocyclic 5- to 10-membered unsaturated hydrocarbon, carbamoyl that may be substituted with C1-C6 alkyl or monocyclic 5- to 10-membered unsaturated hydrocarbon, (C2-C7 acyl)oxy, amino that may be substituted with C1-C6 alkyl or C2-C7 acyl, C3-C7 cycloalkyl that may be
25 substituted with hydroxy, or (C1-C6 alkoxy)(C1-C6 alkyl); when two or more of the substituents are present, the substituents may be identical or different;

R5 and R6 may be identical or different, and each represent hydrogen or C1-C6 alkyl; R5 and R6 may otherwise form oxo or
30 thioxo together;

when R5 and R6 are identical or different, and each represent hydrogen or C1-C6 alkyl, at least one of R5 and R6 is hydrogen;


l represents an integer of 0 to 2;

m represents an integer of 0 to 2;

35 n represents an integer of 0 to 5;

when l is 2, two R2s may be identical or different;
when m is 2, two R3s may be identical or different;
when n is 2 to 5, two to five R4s may be identical or different.
[0029]

- 5 Item 9. A pharmaceutical composition for the prevention and/or treatment of a tumor, the composition comprising a biphenyl compound represented by Formula (I) or a salt thereof and one or more other antitumor agents:




- 10 wherein
ring A represents a monocyclic, bridged cyclic, or spirocyclic 4- to 14-membered nitrogen-containing saturated heterocyclic group having 1 to 3 nitrogen atoms, 0 to 1 sulfur atoms, and 0 to 2 oxygen atoms as heteroatoms;
15 ring B represents monocyclic or bicyclic 5- to 14-membered unsaturated hydrocarbon or a monocyclic or bicyclic 5- to 14-membered unsaturated heterocyclic group that may be substituted with oxo, that has 0 to 4 nitrogen atoms, 0 to 2 sulfur atoms, and 0 to 3 oxygen atoms as heteroatoms, and that has at least one
20 of nitrogen, sulfur, and oxygen;
R1 represents nitro or cyano;
R2 represents halogen;
R3 represents amino, mono- or di(C1-C6 alkyl)amino, (C3-C7 cycloalkyl)amino, or C1-C6 alkyl;
25 R4 represents halogen, nitro, cyano, carboxy, substituted or unsubstituted C1-C8 alkyl, substituted or unsubstituted C2-C6 alkenyl, substituted or unsubstituted C1-C6 alkoxy, substituted or unsubstituted C3-C7 cycloalkyl, mono- or di(C1-C6 alkyl)amino, or substituted or unsubstituted carbamoyl;

when at least one R4 represents substituted C1-C8 alkyl, substituted C2-C6 alkenyl, substituted C1-C6 alkoxy, substituted C3-C7 cycloalkyl, or substituted carbamoyl, the substituent is halogen, carboxy, C1-C6 alkoxy, hydroxy, C1-C6 alkyl that may be substituted with hydroxy, monocyclic 5- to 10-membered unsaturated hydrocarbon, carbamoyl that may be substituted with C1-C6 alkyl or monocyclic 5- to 10-membered unsaturated hydrocarbon, (C2-C7 acyl)oxy, amino that may be substituted with C1-C6 alkyl or C2-C7 acyl, C3-C7 cycloalkyl that may be substituted with hydroxy, or (C1-C6 alkoxy) (C1-C6 alkyl); when two or more of the substituents are present, the substituents may be identical or different; R5 and R6 may be identical or different, and each represent hydrogen or C1-C6 alkyl; R5 and R6 may otherwise form oxo or thioxo together; when R5 and R6 are identical or different, and each represent hydrogen or C1-C6 alkyl, at least one of R5 and R6 is hydrogen; l represents an integer of 0 to 2; m represents an integer of 0 to 2; n represents an integer of 0 to 5; when l is 2, two R2s may be identical or different; when m is 2, two R3s may be identical or different; when n is 2 to 5, two to five R4s may be identical or different.

[0030]

Item 10. A biphenyl compound represented by Formula (I) or a salt thereof for potentiating an antitumor effect of one or more other antitumor agents:



wherein

ring A represents a monocyclic, bridged cyclic, or spirocyclic 4- to 14-membered nitrogen-containing saturated heterocyclic group having 1 to 3 nitrogen atoms, 0 to 1 sulfur atoms, and 0 to 2 oxygen atoms as heteroatoms;

5 ring B represents monocyclic or bicyclic 5- to 14-membered unsaturated hydrocarbon or a monocyclic or bicyclic 5- to 14-membered unsaturated heterocyclic group that may be substituted with oxo, that has 0 to 4 nitrogen atoms, 0 to 2 sulfur atoms, and 0 to 3 oxygen atoms as heteroatoms, and that has at least one
10 of nitrogen, sulfur, and oxygen;

R1 represents nitro or cyano;

R2 represents halogen;

R3 represents amino, mono- or di(C1-C6 alkyl)amino, (C3-C7 cycloalkyl)amino, or C1-C6 alkyl;

15 R4 represents halogen, nitro, cyano, carboxy, substituted or unsubstituted C1-C8 alkyl, substituted or unsubstituted C2-C6 alkenyl, substituted or unsubstituted C1-C6 alkoxy, substituted or unsubstituted C3-C7 cycloalkyl, mono- or di(C1-C6 alkyl)amino, or substituted or unsubstituted carbamoyl;

20 when at least one R4 represents substituted C1-C8 alkyl, substituted C2-C6 alkenyl, substituted C1-C6 alkoxy, substituted C3-C7 cycloalkyl, or substituted carbamoyl, the substituent is halogen, carboxy, C1-C6 alkoxy, hydroxy, C1-C6 alkyl that may be substituted with hydroxy, monocyclic 5- to 10-membered

25 unsaturated hydrocarbon, carbamoyl that may be substituted with C1-C6 alkyl or monocyclic 5- to 10-membered unsaturated hydrocarbon, (C2-C7 acyl)oxy, amino that may be substituted with C1-C6 alkyl or C2-C7 acyl, C3-C7 cycloalkyl that may be substituted with hydroxy, or (C1-C6 alkoxy)(C1-C6 alkyl); when

30 two or more of the substituents are present, the substituents may be identical or different;

R5 and R6 may be identical or different, and each represent hydrogen or C1-C6 alkyl; R5 and R6 may otherwise form oxo or thioxo together;

35 when R5 and R6 are identical or different, and each represent

hydrogen or C1-C6 alkyl, at least one of R5 and R6 is hydrogen;

l represents an integer of 0 to 2;

m represents an integer of 0 to 2;

n represents an integer of 0 to 5;

5 when l is 2, two R2s may be identical or different;


when m is 2, two R3s may be identical or different;

when n is 2 to 5, two to five R4s may be identical or different.

[0031]

Item 11. Use of a biphenyl compound represented by

10 Formula (I) or a salt thereof for potentiating an antitumor effect of one or more other antitumor agents:



wherein

ring A represents a monocyclic, bridged cyclic, or spirocyclic 4-
15 to 14-membered nitrogen-containing saturated heterocyclic group having 1 to 3 nitrogen atoms, 0 to 1 sulfur atoms, and 0 to 2 oxygen atoms as heteroatoms;

ring B represents monocyclic or bicyclic 5- to 14-membered
unsaturated hydrocarbon or a monocyclic or bicyclic 5- to 14-
20 membered unsaturated heterocyclic group that may be substituted with oxo, that has 0 to 4 nitrogen atoms, 0 to 2 sulfur atoms, and 0 to 3 oxygen atoms as heteroatoms, and that has at least one of nitrogen, sulfur, and oxygen;

R1 represents nitro or cyano;


25 R2 represents halogen;

R3 represents amino, mono- or di(C1-C6 alkyl)amino, (C3-C7 cycloalkyl)amino, or C1-C6 alkyl;

R4 represents halogen, nitro, cyano, carboxy, substituted or unsubstituted C1-C8 alkyl, substituted or unsubstituted C2-C6

alkenyl, substituted or unsubstituted C1-C6 alkoxy, substituted
or unsubstituted C3-C7 cycloalkyl, mono- or di(C1-C6 alkyl)amino,
or substituted or unsubstituted carbamoyl;
when at least one R4 represents substituted C1-C8 alkyl,
5 substituted C2-C6 alkenyl, substituted C1-C6 alkoxy, substituted
C3-C7 cycloalkyl, or substituted carbamoyl, the substituent is
halogen, carboxy, C1-C6 alkoxy, hydroxy, C1-C6 alkyl that may be
substituted with hydroxy, monocyclic 5- to 10-membered
unsaturated hydrocarbon, carbamoyl that may be substituted with
10 C1-C6 alkyl or monocyclic 5- to 10-membered unsaturated
hydrocarbon, (C2-C7 acyl)oxy, amino that may be substituted with
C1-C6 alkyl or C2-C7 acyl, C3-C7 cycloalkyl that may be
substituted with hydroxy, or (C1-C6 alkoxy)(C1-C6 alkyl); when
two or more of the substituents are present, the substituents may
15 be identical or different;
R5 and R6 may be identical or different, and each represent
hydrogen or C1-C6 alkyl; R5 and R6 may otherwise form oxo or
thioxo together;
when R5 and R6 are identical or different, and each represent
20 hydrogen or C1-C6 alkyl, at least one of R5 and R6 is hydrogen;
l represents an integer of 0 to 2;
m represents an integer of 0 to 2;
n represents an integer of 0 to 5;
when l is 2, two R2s may be identical or different;
25 when m is 2, two R3s may be identical or different;
when n is 2 to 5, two to five R4s may be identical or different.
[0032]

Item 12. Use of a biphenyl compound represented by
Formula (I) or a salt thereof for the production of an antitumor
30 effect potentiator for one or more other antitumor agents:



wherein

ring A represents a monocyclic, bridged cyclic, or spirocyclic 4- to 14-membered nitrogen-containing saturated heterocyclic group having 1 to 3 nitrogen atoms, 0 to 1 sulfur atoms, and 0 to 2 oxygen atoms as heteroatoms;

ring B represents monocyclic or bicyclic 5- to 14-membered unsaturated hydrocarbon or a monocyclic or bicyclic 5- to 14-membered unsaturated heterocyclic group that may be substituted with oxo, that has 0 to 4 nitrogen atoms, 0 to 2 sulfur atoms, and 0 to 3 oxygen atoms as heteroatoms, and that has at least one of nitrogen, sulfur, and oxygen;

R1 represents nitro or cyano;

R2 represents halogen;

R3 represents amino, mono- or di(C1-C6 alkyl)amino, (C3-C7 cycloalkyl)amino, or C1-C6 alkyl;

R4 represents halogen, nitro, cyano, carboxy, substituted or unsubstituted C1-C8 alkyl, substituted or unsubstituted C2-C6 alkenyl, substituted or unsubstituted C1-C6 alkoxy, substituted or unsubstituted C3-C7 cycloalkyl, mono- or di(C1-C6 alkyl)amino, or substituted or unsubstituted carbamoyl;

when at least one R4 represents substituted C1-C8 alkyl, substituted C2-C6 alkenyl, substituted C1-C6 alkoxy, substituted C3-C7 cycloalkyl, or substituted carbamoyl, the substituent is halogen, carboxy, C1-C6 alkoxy, hydroxy, C1-C6 alkyl that may be substituted with hydroxy, monocyclic 5- to 10-membered unsaturated hydrocarbon, carbamoyl that may be substituted with C1-C6 alkyl or monocyclic 5- to 10-membered unsaturated hydrocarbon, (C2-C7 acyl)oxy, amino that may be substituted with

C1-C6 alkyl or C2-C7 acyl, C3-C7 cycloalkyl that may be substituted with hydroxy, or (C1-C6 alkoxy) (C1-C6 alkyl); when two or more of the substituents are present, the substituents may be identical or different;

5 R5 and R6 may be identical or different, and each represent hydrogen or C1-C6 alkyl; R5 and R6 may otherwise form oxo or thioxo together;

when R5 and R6 are identical or different, and each represent hydrogen or C1-C6 alkyl, at least one of R5 and R6 is hydrogen;

10 l represents an integer of 0 to 2;

m represents an integer of 0 to 2;

n represents an integer of 0 to 5;


when l is 2, two R2s may be identical or different;

when m is 2, two R3s may be identical or different;

15 when n is 2 to 5, two to five R4s may be identical or different.

[0033]

Item 13. A method for preventing and/or treating a tumor, comprising administering a combination of a biphenyl compound represented by Formula (I) or a salt thereof and one or
20 more other antitumor agents in an amount effective for prevention and/or treatment to a patient:



wherein

ring A represents a monocyclic, bridged cyclic, or spirocyclic 4-
25 to 14-membered nitrogen-containing saturated heterocyclic group having 1 to 3 nitrogen atoms, 0 to 1 sulfur atoms, and 0 to 2 oxygen atoms as heteroatoms;

ring B represents monocyclic or bicyclic 5- to 14-membered unsaturated hydrocarbon or a monocyclic or bicyclic 5- to 14-

membered unsaturated heterocyclic group that may be substituted with oxo, that has 0 to 4 nitrogen atoms, 0 to 2 sulfur atoms, and 0 to 3 oxygen atoms as heteroatoms, and that has at least one of nitrogen, sulfur, and oxygen;

5 R1 represents nitro or cyano;

R2 represents halogen;

R3 represents amino, mono- or di(C1-C6 alkyl)amino, (C3-C7 cycloalkyl)amino, or C1-C6 alkyl;

10 R4 represents halogen, nitro, cyano, carboxy, substituted or unsubstituted C1-C8 alkyl, substituted or unsubstituted C2-C6 alkenyl, substituted or unsubstituted C1-C6 alkoxy, substituted or unsubstituted C3-C7 cycloalkyl, mono- or di(C1-C6 alkyl)amino, or substituted or unsubstituted carbamoyl;

when at least one R4 represents substituted C1-C8 alkyl,
15 substituted C2-C6 alkenyl, substituted C1-C6 alkoxy, substituted C3-C7 cycloalkyl, or substituted carbamoyl, the substituent is halogen, carboxy, C1-C6 alkoxy, hydroxy, C1-C6 alkyl that may be substituted with hydroxy, monocyclic 5- to 10-membered unsaturated hydrocarbon, carbamoyl that may be substituted with
20 C1-C6 alkyl or monocyclic 5- to 10-membered unsaturated hydrocarbon, (C2-C7 acyl)oxy, amino that may be substituted with C1-C6 alkyl or C2-C7 acyl, C3-C7 cycloalkyl that may be substituted with hydroxy, or (C1-C6 alkoxy)(C1-C6 alkyl); when two or more of the substituents are present, the substituents may
25 be identical or different;

R5 and R6 may be identical or different, and each represent hydrogen or C1-C6 alkyl; R5 and R6 may otherwise form oxo or thioxo together;

when R5 and R6 are identical or different, and each represent
30 hydrogen or C1-C6 alkyl, at least one of R5 and R6 is hydrogen;

l represents an integer of 0 to 2;

m represents an integer of 0 to 2;


n represents an integer of 0 to 5;

when l is 2, two R2s may be identical or different;

35 when m is 2, two R3s may be identical or different;

when n is 2 to 5, two to five R4s may be identical or different.
[0034]

Item 14. A method for preventing and/or treating a tumor, comprising administering a biphenyl compound represented by Formula (I) or a salt thereof in an amount effective for prevention and/or treatment to a cancer patient administered one or more other antitumor agents:



wherein

ring A represents a monocyclic, bridged cyclic, or spirocyclic 4- to 14-membered nitrogen-containing saturated heterocyclic group having 1 to 3 nitrogen atoms, 0 to 1 sulfur atoms, and 0 to 2 oxygen atoms as heteroatoms;

ring B represents monocyclic or bicyclic 5- to 14-membered unsaturated hydrocarbon or a monocyclic or bicyclic 5- to 14-membered unsaturated heterocyclic group that may be substituted with oxo, that has 0 to 4 nitrogen atoms, 0 to 2 sulfur atoms, and 0 to 3 oxygen atoms as heteroatoms, and that has at least one of nitrogen, sulfur, and oxygen;

R1 represents nitro or cyano;


R2 represents halogen;

R3 represents amino, mono- or di(C1-C6 alkyl)amino, (C3-C7 cycloalkyl)amino, or C1-C6 alkyl;

R4 represents halogen, nitro, cyano, carboxy, substituted or unsubstituted C1-C8 alkyl, substituted or unsubstituted C2-C6 alkenyl, substituted or unsubstituted C1-C6 alkoxy, substituted or unsubstituted C3-C7 cycloalkyl, mono- or di(C1-C6 alkyl)amino, or substituted or unsubstituted carbamoyl;

when at least one R4 represents substituted C1-C8 alkyl, substituted C2-C6 alkenyl, substituted C1-C6 alkoxy, substituted C3-C7 cycloalkyl, or substituted carbamoyl, the substituent is halogen, carboxy, C1-C6 alkoxy, hydroxy, C1-C6 alkyl that may be substituted with hydroxy, monocyclic 5- to 10-membered unsaturated hydrocarbon, carbamoyl that may be substituted with C1-C6 alkyl or monocyclic 5- to 10-membered unsaturated hydrocarbon, (C2-C7 acyl)oxy, amino that may be substituted with C1-C6 alkyl or C2-C7 acyl, C3-C7 cycloalkyl that may be substituted with hydroxy, or (C1-C6 alkoxy) (C1-C6 alkyl); when two or more of the substituents are present, the substituents may be identical or different; R5 and R6 may be identical or different, and each represent hydrogen or C1-C6 alkyl; R5 and R6 may otherwise form oxo or thioxo together; when R5 and R6 are identical or different, and each represent hydrogen or C1-C6 alkyl, at least one of R5 and R6 is hydrogen; l represents an integer of 0 to 2; m represents an integer of 0 to 2; n represents an integer of 0 to 5; when l is 2, two R2s may be identical or different; when m is 2, two R3s may be identical or different; when n is 2 to 5, two to five R4s may be identical or different. [0035]

Item 15. A method for potentiating an antitumor effect, comprising administering a biphenyl compound represented by Formula (I) or a salt thereof in an amount effective for prevention and/or treatment to a cancer patient administered one or more other antitumor agents:



wherein

ring A represents a monocyclic, bridged cyclic, or spirocyclic 4- to 14-membered nitrogen-containing saturated heterocyclic group having 1 to 3 nitrogen atoms, 0 to 1 sulfur atoms, and 0 to 2 oxygen atoms as heteroatoms;

ring B represents monocyclic or bicyclic 5- to 14-membered unsaturated hydrocarbon or a monocyclic or bicyclic 5- to 14-membered unsaturated heterocyclic group that may be substituted with oxo, that has 0 to 4 nitrogen atoms, 0 to 2 sulfur atoms, and 0 to 3 oxygen atoms as heteroatoms, and that has at least one of nitrogen, sulfur, and oxygen;

R1 represents nitro or cyano;

R2 represents halogen;

R3 represents amino, mono- or di(C1-C6 alkyl)amino, (C3-C7 cycloalkyl)amino, or C1-C6 alkyl;

R4 represents halogen, nitro, cyano, carboxy, substituted or unsubstituted C1-C8 alkyl, substituted or unsubstituted C2-C6 alkenyl, substituted or unsubstituted C1-C6 alkoxy, substituted or unsubstituted C3-C7 cycloalkyl, mono- or di(C1-C6 alkyl)amino, or substituted or unsubstituted carbamoyl;

when at least one R4 represents substituted C1-C8 alkyl, substituted C2-C6 alkenyl, substituted C1-C6 alkoxy, substituted C3-C7 cycloalkyl, or substituted carbamoyl, the substituent is halogen, carboxy, C1-C6 alkoxy, hydroxy, C1-C6 alkyl that may be substituted with hydroxy, monocyclic 5- to 10-membered unsaturated hydrocarbon, carbamoyl that may be substituted with C1-C6 alkyl or monocyclic 5- to 10-membered unsaturated hydrocarbon, (C2-C7 acyl)oxy, amino that may be substituted with

C1-C6 alkyl or C2-C7 acyl, C3-C7 cycloalkyl that may be substituted with hydroxy, or (C1-C6 alkoxy) (C1-C6 alkyl); when two or more of the substituents are present, the substituents may be identical or different;

5 R5 and R6 may be identical or different, and each represent hydrogen or C1-C6 alkyl; R5 and R6 may otherwise form oxo or thioxo together;

when R5 and R6 are identical or different, and each represent hydrogen or C1-C6 alkyl, at least one of R5 and R6 is hydrogen;

10 l represents an integer of 0 to 2;

m represents an integer of 0 to 2;

n represents an integer of 0 to 5;


when l is 2, two R2s may be identical or different;

when m is 2, two R3s may be identical or different;

15 when n is 2 to 5, two to five R4s may be identical or different.

[0036]

Item 16. A product as a combined preparation to be used simultaneously, sequentially, or at an interval upon prevention and/or treatment of a tumor, the product comprising a biphenyl
20 compound or a salt thereof and one or more other antitumor agents, wherein the biphenyl compound is represented by Formula (I):



wherein


25 ring A represents a monocyclic, bridged cyclic, or spirocyclic 4- to 14-membered nitrogen-containing saturated heterocyclic group having 1 to 3 nitrogen atoms, 0 to 1 sulfur atoms, and 0 to 2 oxygen atoms as heteroatoms;

ring B represents monocyclic or bicyclic 5- to 14-membered

unsaturated hydrocarbon or a monocyclic or bicyclic 5- to 14-membered unsaturated heterocyclic group that may be substituted with oxo, that has 0 to 4 nitrogen atoms, 0 to 2 sulfur atoms, and 0 to 3 oxygen atoms as heteroatoms, and that has at least one
5 of nitrogen, sulfur, and oxygen;
R1 represents nitro or cyano;
R2 represents halogen;
R3 represents amino, mono- or di(C1-C6 alkyl)amino, (C3-C7 cycloalkyl)amino, or C1-C6 alkyl;
10 R4 represents halogen, nitro, cyano, carboxy, substituted or unsubstituted C1-C8 alkyl, substituted or unsubstituted C2-C6 alkenyl, substituted or unsubstituted C1-C6 alkoxy, substituted or unsubstituted C3-C7 cycloalkyl, mono- or di(C1-C6 alkyl)amino, or substituted or unsubstituted carbamoyl;
15 when at least one R4 represents substituted C1-C8 alkyl, substituted C2-C6 alkenyl, substituted C1-C6 alkoxy, substituted C3-C7 cycloalkyl, or substituted carbamoyl, the substituent is halogen, carboxy, C1-C6 alkoxy, hydroxy, C1-C6 alkyl that may be substituted with hydroxy, monocyclic 5- to 10-membered
20 unsaturated hydrocarbon, carbamoyl that may be substituted with C1-C6 alkyl or monocyclic 5- to 10-membered unsaturated hydrocarbon, (C2-C7 acyl)oxy, amino that may be substituted with C1-C6 alkyl or C2-C7 acyl, C3-C7 cycloalkyl that may be substituted with hydroxy, or (C1-C6 alkoxy)(C1-C6 alkyl); when
25 two or more of the substituents are present, the substituents may be identical or different;
R5 and R6 may be identical or different, and each represent hydrogen or C1-C6 alkyl; R5 and R6 may otherwise form oxo or thioxo together;
30 when R5 and R6 are identical or different, and each represent hydrogen or C1-C6 alkyl, at least one of R5 and R6 is hydrogen;
l represents an integer of 0 to 2;
m represents an integer of 0 to 2;
n represents an integer of 0 to 5;
35 when l is 2, two R2s may be identical or different;

when m is 2, two R3s may be identical or different;
when n is 2 to 5, two to five R4s may be identical or different.
[0037]

Item 17. A combination of a biphenyl compound
5 represented by Formula (I) or a salt thereof and one or more
other antitumor agents:



wherein

ring A represents a monocyclic, bridged cyclic, or spirocyclic 4-
10 to 14-membered nitrogen-containing saturated heterocyclic group
having 1 to 3 nitrogen atoms, 0 to 1 sulfur atoms, and 0 to 2
oxygen atoms as heteroatoms;

ring B represents monocyclic or bicyclic 5- to 14-membered
unsaturated hydrocarbon or a monocyclic or bicyclic 5- to 14-
15 membered unsaturated heterocyclic group that may be substituted
with oxo, that has 0 to 4 nitrogen atoms, 0 to 2 sulfur atoms,
and 0 to 3 oxygen atoms as heteroatoms, and that has at least one
of nitrogen, sulfur, and oxygen;

R1 represents nitro or cyano;

20 R2 represents halogen;

R3 represents amino, mono- or di(C1-C6 alkyl)amino, (C3-C7
cycloalkyl)amino, or C1-C6 alkyl;

R4 represents halogen, nitro, cyano, carboxy, substituted or
unsubstituted C1-C8 alkyl, substituted or unsubstituted C2-C6
25 alkenyl, substituted or unsubstituted C1-C6 alkoxy, substituted
or unsubstituted C3-C7 cycloalkyl, mono- or di(C1-C6 alkyl)amino,
or substituted or unsubstituted carbamoyl;

when at least one R4 represents substituted C1-C8 alkyl,
substituted C2-C6 alkenyl, substituted C1-C6 alkoxy, substituted

C3-C7 cycloalkyl, or substituted carbamoyl, the substituent is halogen, carboxy, C1-C6 alkoxy, hydroxy, C1-C6 alkyl that may be substituted with hydroxy, monocyclic 5- to 10-membered unsaturated hydrocarbon, carbamoyl that may be substituted with
5 C1-C6 alkyl or monocyclic 5- to 10-membered unsaturated hydrocarbon, (C2-C7 acyl)oxy, amino that may be substituted with C1-C6 alkyl or C2-C7 acyl, C3-C7 cycloalkyl that may be substituted with hydroxy, or (C1-C6 alkoxy) (C1-C6 alkyl); when two or more of the substituents are present, the substituents may
10 be identical or different;
R5 and R6 may be identical or different, and each represent hydrogen or C1-C6 alkyl; R5 and R6 may otherwise form oxo or thioxo together;
when R5 and R6 are identical or different, and each represent
15 hydrogen or C1-C6 alkyl, at least one of R5 and R6 is hydrogen;
l represents an integer of 0 to 2;
m represents an integer of 0 to 2;
n represents an integer of 0 to 5;
when l is 2, two R2s may be identical or different;
20 when m is 2, two R3s may be identical or different;
when n is 2 to 5, two to five R4s may be identical or different.

Advantageous Effects of Invention

[0038]


25 The present invention enables cancer therapy that achieves an excellent antitumor effect while suppressing occurrence of side effects.

Description of Embodiments

30 [0039]

 In the present invention, the LSD1 inhibitor that provides excellent synergism with one or more other antitumor agents is a biphenyl compound represented by the following Formula (I) or a salt thereof.

35 [0040]



wherein ring A, ring B, R1 to R6, l, m, and n are as defined above.

[0041]

5 In the present specification, unless otherwise specified, examples of the "substituent" include halogen, hydroxy, cyano, nitro, alkyl, hydroxyalkyl, halogenoalkyl, cycloalkyl, hydroxycycloalkyl, cycloalkyl-alkyl, aralkyl, alkenyl, alkynyl, alkoxy, halogenoalkoxy, cycloalkoxy, cycloalkyl-alkoxy, unsaturated hydrocarbon ring-alkoxy, alkylthio, cycloalkyl-alkylthio, amino, mono- or di-alkylamino, cycloalkylamino, cycloalkyl-alkylamino, acyl, acyloxy, oxo, carboxy, alkoxycarbonyl, aralkyloxycarbonyl, carbamoyl that may be substituted with an unsaturated hydrocarbon ring, saturated or unsaturated heterocyclic group, unsaturated hydrocarbon ring (e.g., aromatic hydrocarbon), saturated heterocyclic oxy, and the like. The number of the substituents, when present, is typically one, two, or three.

[0042]

20 In the present specification, examples of the "halogen" include fluorine, chlorine, bromine, iodine, and the like, with fluorine, chlorine, bromine, or iodine being preferable, and fluorine or chlorine being more preferable.

[0043]

25 In the present specification, the "alkyl" may be straight or branched. Examples include C1-C6 alkyl, such as methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl, tert-butyl, n-pentyl, isopentyl, tert-pentyl, and n-hexyl.

[0044]

In the present specification, examples of the "hydroxyalkyl" include the above-listed alkyl groups that have at least one hydroxy group (e.g., one or two hydroxy groups). Specific examples include hydroxymethyl, 2-hydroxyethyl, 1-
5 hydroxyethyl, 3-hydroxypropyl, 2-hydroxypropyl, 1-methyl-2-hydroxyethyl, 4-hydroxybutyl, 2,2-dimethyl-2-hydroxyethyl, 5-hydroxypentyl, 3,3-dimethyl-3-hydroxypropyl, 6-hydroxyhexyl, dihydroxymethyl, 1,2-dihydroxyethyl, 2,3-dihydroxypropyl, 3,4-dihydroxybutyl, 4,5-dihydroxypentyl, 5,6-dihydroxyhexyl, and the
10 like, with hydroxyalkyl having one hydroxy group being preferable.

[0045]

In the present specification, the "halogenoalkyl" is straight or branched C1-C6 alkyl having 1 to 13 halogen atoms
15 (halogeno C1-C6 alkyl). Examples include halogeno C1-C6 alkyl, such as fluoromethyl, difluoromethyl, trifluoromethyl, trichloromethyl, fluoroethyl, 1,1,1-trifluoroethyl, monofluoro-n-propyl, perfluoro-n-propyl, and perfluoroisopropyl, with halogeno C1-C4 alkyl being preferable, and halogeno C1-C4 alkyl having 1
20 to 7 halogen atoms being more preferable.

[0046]

In the present specification, specific examples of the "cycloalkyl" include C3-C7 cycloalkyl, such as cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, and cycloheptyl.

25 [0047]

In the present specification, examples of the "hydroxycycloalkyl" include the above-listed C3-C7 cycloalkyl groups that have at least one hydroxy group (e.g., one or two hydroxy groups). Specific examples include 1-hydroxycyclopropyl,
30 2-hydroxycyclopropyl, 1-hydroxycyclobutyl, 3-hydroxycyclobutyl, 1-hydroxycyclopentyl, 3,4-dihydroxycyclopentyl, 1-hydroxycyclohexyl, 4-hydroxycyclohexyl, 1-hydroxycycloheptyl, and the like, with hydroxycycloalkyl having one hydroxy group being preferable.

35 [0048]

In the present specification, examples of the "cycloalkyl-alkyl" include C3-C7 cycloalkyl-substituted C1-C4 alkyl, such as cyclopropylmethyl, cyclobutylmethyl, cyclopentylmethyl, cyclohexylmethyl, and cycloheptylmethyl.

5 [0049]

In the present specification, examples of the "aralkyl" include C7-C13 aralkyl, such as benzyl, phenethyl, naphthylmethyl, and fluorenylmethyl.

[0050]

10 In the present specification, the "alkenyl" may be straight, branched, or cyclic, and refers to unsaturated hydrocarbon having at least one double bond (e.g., one or two double bonds). Examples include C2-C6 alkenyl, such as vinyl, allyl, 1-propenyl, 2-methyl-2-propenyl, isopropenyl, 1-butenyl, 15 2-butenyl, 3-butenyl, isobutenyl, 2-pentenyl, 3-pentenyl, 4-pentenyl, 2-methyl-2-butenyl, 3-methyl-2-butenyl, 5-hexenyl, 1-cyclopentenyl, 1-cyclohexenyl, and 3-methyl-3-butenyl.

[0051]

In the present specification, the "alkynyl" may be 20 straight, branched, or cyclic, and refers to unsaturated hydrocarbon having at least one triple bond (e.g., one or two triple bonds). Examples include C2-C6 alkynyl, such as ethynyl, 1-propynyl, 2-propynyl, 1-butyne, 2-butyne, 3-butyne, and 1-methyl-2-propynyl.

25 [0052]

In the present specification, the "alkoxy" may be straight or branched. Examples include C1-C6 alkoxy, such as methoxy, ethoxy, n-propoxy, isopropoxy, n-butoxy, isobutoxy, sec-butoxy, tert-butoxy, pentyloxy, isopentyloxy, and hexyloxy.

30 [0053]

In the present specification, the "halogenoalkoxy" refers to straight or branched C1-C6 alkoxy having 1 to 13 halogen atoms (halogeno C1-C6 alkoxy). Examples include halogeno C1-C6 alkoxy, such as fluoromethoxy, difluoromethoxy, 35 trifluoromethoxy, trichloromethoxy, fluoroethoxy, 1,1,1-

trifluoroethoxy, monofluoro-n-propoxy, perfluoro-n-propoxy, and perfluoro-isopropoxy, with halogeno C1-C4 alkoxy being preferable, and halogeno C1-C4 alkoxy having 1 to 7 halogen atoms being more preferable.

5 [0054]

In the present specification, examples of the "cycloalkoxy" include C3-C7 cycloalkoxy, such as cyclopropoxy, cyclobutoxy, cyclopentyloxy, cyclohexyloxy, and cycloheptyloxy.
[0055]

10 In the present specification, examples of the "cycloalkyl-alkoxy" include C3-C7 cycloalkyl-substituted C1-C4 alkoxy, such as cyclopropylmethoxy, cyclobutylmethoxy, cyclopentylmethoxy, cyclohexylmethoxy, and cycloheptylmethoxy.
[0056]

15 In the present specification, the "alkylthio" may be straight or branched. Examples include C1-C6 alkylthio, such as methylthio, ethylthio, n-propylthio, isopropylthio, n-butylthio, isobutylthio, tert-butylthio, n-pentylthio, isopentylthio, and hexylthio.

20 [0057]

In the present specification, examples of the "cycloalkyl-alkylthio" include C3-C7 cycloalkyl-substituted C1-C4 alkylthio, such as cyclopropylmethylthio, cyclobutylmethylthio, cyclopentylmethylthio, cyclohexylmethylthio, and
25 cycloheptylmethylthio.
[0058]

In the present specification, examples of the "monoalkylamino" include amino monosubstituted with straight or branched C1-C6 alkyl, such as methylamino, ethylamino, n-
30 propylamino, isopropylamino, n-butylamino, isobutylamino, tert-butylamino, n-pentylamino, isopentylamino, and hexylamino.
[0059]

In the present specification, examples of the "dialkylamino" include amino disubstituted with the same or
35 different straight or branched C1-C6 alkyl groups, such as

dimethylamino, diethylamino, di(n-propyl)amino, diisopropylamino, di(n-butyl)amino, diisobutylamino, di(tert-butyl)amino, di(n-pentyl)amino, diisopentylamino, dihexylamino, methylethylamino, and methylisopropylamino.

5 [0060]

In the present specification, examples of the "cycloalkylamino" include amino having one or two cycloalkyl groups mentioned above. Specific examples include N-cyclopropylamino, N,N-dicyclopropylamino, N-cyclobutylamino, N-cyclopentylamino, N-cyclohexylamino, N-cycloheptylamino, and the like.

[0061]

In the present specification, examples of the "cycloalkyl-alkylamino" include C3-C7 cycloalkyl-substituted C1-C4 alkylamino, such as cyclopropylmethylamino, cyclobutylmethylamino, cyclopentylmethylamino, cyclohexylmethylamino, and cycloheptylmethylamino.

[0062]

In the present specification, the "acyl" refers to alkylcarbonyl or arylcarbonyl.

[0063]

In the present specification, examples of the "alkylcarbonyl" include straight or branched (C1-C6 alkyl)carbonyl, such as methylcarbonyl, ethylcarbonyl, n-propylcarbonyl, isopropylcarbonyl, n-butylcarbonyl, isobutylcarbonyl, tert-butylcarbonyl, n-pentylcarbonyl, isopentylcarbonyl, and hexylcarbonyl.

[0064]

In the present specification, examples of the "arylcarbonyl" include (C6-C13 aryl)carbonyl, such as phenylcarbonyl, naphthylcarbonyl, fluorenylcarbonyl, anthrylcarbonyl, biphenylylcarbonyl, tetrahydronaphthylcarbonyl, chromanylcarbonyl, 2,3-dihydro-1,4-dioxanaphthalenylcarbonyl, indanylcarbonyl, and phenanthrylcarbonyl.

35 [0065]

In the present specification, the "acylamino" refers to alkylcarbonylamino or arylcarbonylamino.

[0066]

In the present specification, examples of the
5 "alkylcarbonylamino" include straight or branched (C1-C6
alkyl)carbonylamino, such as methylcarbonylamino,
ethylcarbonylamino, n-propylcarbonylamino,
isopropylcarbonylamino, n-butylcarbonylamino,
isobutylcarbonylamino, tert-butylcarbonylamino, n-
10 pentylcarbonylamino, isopentylcarbonylamino, and
hexylcarbonylamino.

[0067]

In the present specification, examples of the
"arylcarbonylamino" include (C6-C13 aryl)carbonylamino, such as
15 phenylcarbonylamino, naphthylcarbonylamino,
fluorenylcarbonylamino, anthrylcarbonylamino,
biphenylylcarbonylamino, tetrahydronaphthylcarbonylamino,
chromanylcarbonylamino, 2,3-dihydro-1,4-
dioxanaphthalenylcarbonylamino, indanylcarbonylamino, and
20 phenanthrylcarbonylamino.

[0068]

In the present specification, the "acyloxy" refers to alkylcarbonyloxy or arylcarbonyloxy.

[0069]

25 In the present specification, examples of the
"alkylcarbonyloxy" include straight or branched (C1-C6
alkyl)carbonyloxy, such as methylcarbonyloxy, ethylcarbonyloxy,
n-propylcarbonyloxy, isopropylcarbonyloxy, n-butylcarbonyloxy,
isobutylcarbonyloxy, tert-butylcarbonyloxy, n-pentylcarbonyloxy,
30 isopentylcarbonyloxy, and hexylcarbonyloxy.

[0070]

In the present specification, examples of the
"arylcarbonyloxy" include (C6-C13 aryl)carbonyloxy, such as
phenylcarbonyloxy, naphthylcarbonyloxy, fluorenylcarbonyloxy,
35 anthrylcarbonyloxy, biphenylylcarbonyloxy,

tetrahydronaphthylcarbonyloxy, chromanylcarbonyloxy, 2,3-dihydro-1,4-dioxanaphthalenylcarbonyloxy, indanylcarbonyloxy, and phenanthrylcarbonyloxy.

[0071]

5 In the present specification, the "alkoxycarbonyl" may be straight or branched. Examples include (C1-C6 alkoxy)carbonyl, such as methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, isopropoxycarbonyl, butoxycarbonyl, isobutoxycarbonyl, tert-butoxycarbonyl, pentyloxycarbonyl, isopentyloxycarbonyl, and
10 hexyloxycarbonyl.

[0072]

 In the present specification, examples of the "aralkyloxycarbonyl" include (C7-C13 aralkyl)oxycarbonyl, such as benzyloxycarbonyl, phenethyloxycarbonyl,
15 naphthylmethyloxycarbonyl, and fluorenylmethyloxycarbonyl.

[0073]

 In the present specification, the "saturated heterocyclic group" refers to a monocyclic or polycyclic saturated heterocyclic group having one or more (preferably 1 to
20 3) heteroatoms selected from nitrogen, oxygen, and sulfur. Specific examples include morpholinyl, pyrrolidinyl, piperidinyl, piperazinyl, azepanyl, diazepanyl, tetrahydrofuranyl, tetrahydropyranyl, tetrahydrothiophenyl, thiazolidinyl, oxazolidinyl, and the like.

25 [0074]

 In the present specification, the "unsaturated heterocyclic group" refers to a monocyclic or polycyclic, completely or partially unsaturated heterocyclic group having one or more (preferably 1 to 3) heteroatoms selected from nitrogen,
30 oxygen, and sulfur. Specific examples include imidazolyl, thienyl, furyl, pyrrolyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, thiadiazolyl, pyrazolyl, triazolyl, tetrazolyl, pyridyl, pyrazyl, pyrimidinyl, pyridazinyl, pyrazolopyridyl, pyrazolopyrimidinyl, indolyl, isoindolyl, indazolyl,
35 triazolopyridyl, benzoimidazolyl, benzoxazolyl, benzoisoxazolyl,

benzothiazolyl, benzothienyl, benzofuranyl, 1,3-dihydroisobenzofuranyl, purinyl, benzotriazolyl, imidazopyridinyl, quinolinyl, isoquinolinyl, quinazolinyl, quinoxalinyl, phthalazinyl, methylenedioxyphenyl, ethylenedioxyphenyl, dihydrobenzofuranyl, and the like.
[0075]

In the present specification, examples of the "unsaturated hydrocarbon" include a monocyclic or polycyclic C5-14 hydrocarbon ring having at least one unsaturated bond (e.g., 1 to 8 unsaturated bonds), and the like. The "unsaturated hydrocarbon" is preferably aromatic hydrocarbon, or monocyclic or bicyclic 5- to 14-membered unsaturated hydrocarbon.
[0076]

In the present specification, examples of the "aromatic hydrocarbon" include C6-C14 aromatic hydrocarbons, such as phenyl, naphthyl, anthracenyl, phenanthryl, fluorenyl, and tetrahydronaphthyl.
[0077]

In the present specification, examples of the "monocyclic or bicyclic 5- to 14-membered unsaturated hydrocarbon" include cyclopentadienyl, phenyl, naphthyl, tetrahydronaphthyl, azulenyl, heptalenyl, and the like.
[0078]

In the present specification, examples of the "monocyclic 5- to 10-membered unsaturated hydrocarbon" include cyclopentadienyl, phenyl, cyclooctatetraenyl, and the like.
[0079]

In the present specification, the "saturated heterocyclic oxy" refers to saturated heterocyclic oxy having a heteroatom selected from nitrogen, oxygen, and sulfur. Specific examples include morpholinyl, 1-pyrrolidinyl, piperidinyl, piperazinyl, 4-methyl-1-piperazinyl, tetrahydrofuran, tetrahydropyran, tetrahydrothiophene, thiazolidine, and oxazolidine, with saturated heterocyclic oxy having 1 to 3 heteroatoms

selected from nitrogen, oxygen, and sulfur being preferable.

[0080]

In the present specification, the term "CA-CB" used in the description of a group indicates that the group has A- to B-
5 number of carbon atoms. For example, "C1-C6 alkyl" refers to alkyl having 1 to 6 carbon atoms, and "C6-C14 aromatic hydrocarbon oxy" refers to oxy to which C6-C14 aromatic hydrocarbon is bonded. Further, the term "A- to B-membered" indicates that the number of atoms (number of ring members) that
10 constitute a ring is A to B. For example, "4- to 10-membered nitrogen-containing saturated heterocyclic group" refers to a nitrogen-containing saturated heterocyclic group containing 4 to 10 ring members.

[0081]

15 In the compound represented by Formula (I), ring A refers to a nitrogen-containing saturated heterocyclic group that may be crosslinked or spirocyclic. As shown in Formula (I) above, the nitrogen on ring A is bonded to carbonyl, carbothionyl or methylene.

20 [0082]

Examples of the monocyclic nitrogen-containing saturated heterocyclic group in the "monocyclic, bridged cyclic, or spirocyclic nitrogen-containing saturated heterocyclic group" represented by ring A include monocyclic nitrogen-containing
25 saturated heterocyclic groups, such as pyrrolidinyl, piperidinyl, piperazinyl, azepanyl, and diazepanyl. The monocyclic nitrogen-containing saturated heterocyclic group is preferably a monocyclic nitrogen-containing saturated heterocyclic group having 1 to 3 nitrogen atoms, 0 to 1 sulfur atoms, and 0 to 2
30 oxygen atoms as heteroatoms, more preferably a monocyclic nitrogen-containing saturated heterocyclic group having 1 to 2 nitrogen atoms as heteroatoms, more preferably a monocyclic 4- to 10-membered nitrogen-containing saturated heterocyclic group having 1 to 2 nitrogen atoms as heteroatoms, more preferably
35 pyrrolidinyl, piperidinyl, piperazinyl, azepanyl, or diazepanyl,

5 Examples of the bridged cyclic nitrogen-containing saturated heterocyclic group in the "monocyclic, bridged cyclic, or spirocyclic nitrogen-containing saturated heterocyclic group" represented by ring A include

Chemical structures of various indole derivatives, including indole, 2,3-dihydroindole, 2,3,4,5-tetrahydroindole, 2,3,4,5,6-pentahydroindole, 2,3,4,5,6,7-hexahydroindole, 2,3,4,5,6,7,8-heptahydroindole, 2,3,4,5,6,7,8,9-octahydroindole, and 2,3,4,5,6,7,8,9,10-nonahydroindole.


and the like. The bridged cyclic nitrogen-containing saturated heterocyclic group is preferably

, more preferably

Chemical structures of 10 bicyclic and tricyclic nitrogen-containing compounds, including indole, norbornene derivatives, and complex polycyclic systems.

, more preferably

[0090]



[0091]

, more preferably

[0092]




5

[0093]

, and more preferably

[0094]



10 [0095]

[0096]

Examples of the spirocyclic nitrogen-containing saturated heterocyclic group in the "monocyclic, bridged cyclic, or spirocyclic nitrogen-containing saturated heterocyclic group" represented by ring A include spirocyclic groups having 0 to 2 oxygen atoms in which any two of 4- to 7-membered nitrogen-containing saturated heterocyclic rings are bonded to each other. The spirocyclic nitrogen-containing saturated heterocyclic group is preferably a 7- to 12-membered spirocyclic group having 2 nitrogen atoms and 0 to 1 oxygen atoms in which any two of 4- to 7-membered nitrogen-containing saturated heterocyclic rings are bonded to each other, more preferably diazaspiroheptanyl, diazaspirooctanyl, diazaspirononanyl, diazaspirodecanyl, diazaspiroundecanyl, oxadiazaspiroheptanyl, oxadiazaspirooctanyl, oxadiazaspirononanyl, oxadiazaspirodecanyl, or oxadiazaspiroundecanyl, more preferably diazaspirooctanyl, diazaspirononanyl, diazaspirodecanyl, or oxadiazaspirononanyl,

more preferably 2,7-diazaspiro[3.4]octanyl, 3,7-diazaspiro[3.4]octanyl, 2,7-diazaspiro[3.5]nonanyl, 2,8-diazaspiro[3.5]nonanyl, 3,7-diazaspiro[3.5]nonanyl, 3,8-diazaspiro[4.4]nonanyl, 3,8-diazaspiro[3.5]decanyl, or 9-oxa-

5 diazaspiro[3.5]nonanyl, more preferably 2,7-diazaspiro[3.4]octanyl, 3,7-diazaspiro[3.4]octanyl, 2,7-diazaspiro[3.5]nonanyl, 2,8-diazaspiro[3.5]nonanyl, or 9-oxa-diazaspiro[3.5]nonanyl, more preferably 2,7-diazaspiro[3.4]octanyl, 3,7-diazaspiro[3.4]octanyl, or 2,8-

10 diazaspiro[3.5]nonanyl, and more preferably 2,8-diazaspiro[3.5]nonanyl.


[0097]

The "monocyclic, bridged cyclic, or spirocyclic nitrogen-containing saturated heterocyclic group" represented by

15 ring A is preferably a monocyclic, bridged cyclic, or spirocyclic 4- to 14-membered nitrogen-containing saturated heterocyclic group having 1 to 3 nitrogen atoms, 0 to 1 sulfur atoms, and 0 to 2 oxygen atoms as heteroatoms, more preferably a monocyclic 4- to 10-membered nitrogen-containing saturated heterocyclic group

20 having 1 to 2 nitrogen atoms as heteroatoms, a bridged cyclic nitrogen-containing saturated heterocyclic group, such as

[0098]




[0099]

or a spirocyclic group having 0 to 2 oxygen atoms in which any two of 4- to 7-membered nitrogen-containing saturated heterocyclic rings are bonded to each other, more preferably a monocyclic 4- to 10-membered nitrogen-containing saturated heterocyclic group having 1 to 2 nitrogen atoms as heteroatoms, a

30 bridged cyclic nitrogen-containing saturated heterocyclic group, such as


[0100]



[0101]

or a 7- to 12-membered spirocyclic group having 2 nitrogen atoms
 5 and 0 to 1 oxygen atoms in which any two of 4- to 7-membered
 nitrogen-containing saturated heterocyclic rings are bonded to
 each other, more preferably pyrrolidinyl, piperidinyl,
 piperazinyl, azepanyl, diazepanyl,

[0102]




10

[0103]

, 2,7-diazaspiro[3.4]octanyl, 3,7-diazaspiro[3.4]octanyl, 2,7-
 diazaspiro[3.5]nonanyl, 2,8-diazaspiro[3.5]nonanyl, 3,7-
 diazaspiro[3.5]nonanyl, 3,8-diazaspiro[4.4]nonanyl, 3,8-
 15 diazaspiro[4.5]decanyl, or 9-oxa-diazaspiro[3.5]nonanyl, more
 preferably pyrrolidinyl, piperidinyl, azepanyl, diazepanyl,


[0104]



[0105]

20 , 2,7-diazaspiro[3.4]octanyl, 3,7-diazaspiro[3.4]octanyl, 2,7-
 diazaspiro[3.5]nonanyl, 2,8-diazaspiro[3.5]nonanyl, or 9-oxa-
 diazaspiro[3.5]nonanyl, more preferably pyrrolidinyl,


[0106]



[0107]

, or 2,8-diazaspiro[3.5]nonanyl, more preferably pyrrolidinyl,

[0108]



5

[0109]

[0110]

In the compound represented by Formula (I), ring B represents monocyclic or bicyclic unsaturated hydrocarbon, or a monocyclic or bicyclic unsaturated heterocyclic group that may be substituted with oxo.

[0111]


The "monocyclic or bicyclic unsaturated hydrocarbon" represented by ring B is preferably monocyclic or bicyclic 5- to 14-membered unsaturated hydrocarbon, more preferably phenyl or naphthyl, and more preferably phenyl.

[0112]

The "monocyclic or bicyclic unsaturated heterocyclic group" represented by ring B refers to a monocyclic or bicyclic, completely or partially unsaturated heterocyclic group having a heteroatom selected from nitrogen, oxygen, and sulfur, preferably a 5- to 14- membered monocyclic or bicyclic completely unsaturated heterocyclic group or partially unsaturated heterocyclic group having 0 to 4 nitrogen atoms, 0 to 2 sulfur atoms, and 0 to 3 oxygen atoms as heteroatoms and having at least one of nitrogen, sulfur, and oxygen, more preferably imidazolyl, thienyl, furyl, pyrrolyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, thiadiazolyl, pyrazolyl, triazolyl, tetrazolyl, pyridyl, pyrazyl, pyrimidinyl, pyridazinyl, pyrazolopyridyl, pyrazolopyrimidinyl, indolyl, isoindolyl, indolinyl, indazolyl, triazolopyridyl, benzoimidazolyl, benzoxazolyl, benzoisoxazolyl,


benzothiazolyl, benzotriazolyl, benzothienyl, benzofuranyl,
purinyl, imidazopyridinyl, quinolinyl, isoquinolinyl,
quinazolinyl, quinoxalinyl, phthalazinyl, methylenedioxyphenyl,
ethylenedioxyphenyl, dihydrobenzofuranyl, 1,3-
5 dihydroisobenzofuranyl, dihydrobenzoxazolyl (e.g., 2,3-
dihydrobenzo[d]oxazolyl), dihydrobenzooxazinyl (e.g., 3,4-
dihydro-2H-benzo[b][1,4]oxazinyl), benzodioxolyl (e.g.,
benzo[d][1,3]dioxolyl), dihydrobenzodioxynyl (e.g., 2,3-
dihydrobenzo[b][1,4]dioxynyl), or dihydrobenzothiazolyl (e.g.,
10 2,3-dihydrobenzo[d]thiazolyl), more preferably pyridyl,
pyrazolopyridyl, pyrazolopyrimidinyl, indolyl, indolinyl,
indazolyl, benzoimidazolyl, benzoisoxazolyl, benzothiazolyl,
benzotriazolyl, imidazopyridinyl, quinolinyl, isoquinolinyl,
quinoxalinyl, quinazolinyl, phthalazinyl, dihydrobenzoxazolyl,
15 1,3-dihydroisobenzofuranyl, dihydrobenzooxazinyl, benzodioxolyl,
dihydrobenzodioxynyl, or dihydrobenzothiazolyl, more preferably
pyridyl, pyrazolopyridyl, indolyl, indolinyl, indazolyl,
benzoimidazolyl, benzoisoxazolyl, benzotriazolyl, quinolinyl,
dihydrobenzoxazolyl, 1,3-dihydroisobenzofuranyl,
20 dihydrobenzooxazinyl, or dihydrobenzothiazolyl, more preferably
indolyl, indazolyl, benzoisoxazolyl, or benzotriazolyl, and more
preferably benzotriazolyl. The monocyclic or bicyclic unsaturated
heterocyclic group represented by ring B may be substituted with
oxo. Examples of the monocyclic or bicyclic unsaturated
25 heterocyclic group that is substituted with oxo include 2-oxo-
indolinyl,

[0113]



[0114]


30 2-oxo-2,3-dihydrobenzo[d]oxazolyl,
[0115]



[0116]

2-oxo-2,3-dihydrobenzo[d]thiazolyl,

[0117]



5

[0118]

and the like. The monocyclic or bicyclic unsaturated heterocyclic group that is substituted with oxo is preferably 2-oxo-indolinyl, 2-oxo-2,3-dihydrobenzo[d]oxazolyl, or 2-oxo-2,3-dihydrobenzo[d]thiazolyl, and more preferably 2-oxo-2,3-dihydrobenzo[d]oxazolyl or 2-oxo-2,3-dihydrobenzo[d]thiazolyl.

10

[0119]

Ring B is preferably monocyclic or bicyclic 5- to 14-membered unsaturated hydrocarbon or a monocyclic or bicyclic 5- to 14-membered unsaturated heterocyclic group that may be substituted with oxo, that has 0 to 4 nitrogen atoms, 0 to 2 sulfur atoms, and 0 to 3 oxygen atoms as heteroatoms, and that has at least one of nitrogen, sulfur, and oxygen, more preferably phenyl, naphthyl, pyridyl, pyrazolopyridyl, pyrazolopyrimidinyl, indolyl, indolinyl, 2-oxo-indolinyl, indazolyl, benzoimidazolyl, benzoisoxazolyl, benzothiazolyl, benzotriazolyl, imidazopyridinyl, quinolinyl, isoquinolinyl, quinazolinyl, quinoxalinyl, phthalazinyl, 2-oxo-2,3-dihydrobenzo[d]oxazolyl, 1,3-dihydroisobenzofuranyl, dihydrobenzooxazinyl, benzodioxolyl, dihydrobenzodioxynyl, or 2-oxo-2,3-dihydrobenzo[d]thiazolyl, more preferably phenyl, naphthyl, pyridyl, pyrazolopyridyl, indolyl, indolinyl, indazolyl, benzoimidazolyl, benzoisoxazolyl, benzotriazolyl, quinolinyl, 2-oxo-2,3-dihydrobenzo[d]oxazolyl, 1,3-dihydroisobenzofuranyl, dihydrobenzooxazinyl, or 2-oxo-2,3-dihydrobenzo[d]thiazolyl, more preferably phenyl, indolyl, indazolyl, benzoisoxazolyl, or benzotriazolyl, and more

20

25

30

preferably benzotriazolyl.

[0120]

In the compound represented by Formula (I), R1 represents nitro or cyano, and preferably cyano.

5 [0121]

In the compound represented by Formula (I), R2 represents halogen, and preferably fluorine. When two or more R2s are present, R2s may be identical or different.

[0122]

10 In the compound represented by Formula (I), l is an integer of 0 to 2, and preferably an integer of 0 to 1.

[0123]

In the compound represented by Formula (I), R3 represents substituted or unsubstituted amino, C1-C6 alkyl, halogen, cyano, oxo, hydroxy, carbamoyl, sulfo, C1-C6 alkoxy, or amino (C1-C6 alkyl). When two or more R3s are present, R3s may be identical or different.

[0124]

20 The "C1-C6 alkyl" represented by R3 may be straight or branched. Examples include methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, tert-butyl, n-pentyl, isopentyl, hexyl, and the like, with C1-C4 alkyl being preferable, and methyl being more preferable.

[0125]

25 Examples of the "mono(C1-C6 alkyl)amino" represented by R3 include methylamino, ethylamino, n-propylamino, isopropylamino, n-butylamino, isobutylamino, tert-butylamino, n-pentylamino, isopentylamino, hexylamino, and the like, with mono(C1-C4 alkyl)amino being preferable, and methylamino, ethylamino, or isopropylamino being more preferable.

[0126]

35 Examples of the "di(C1-C6 alkyl)amino" represented by R3 include dimethylamino, diethylamino, di(n-propyl)amino, diisopropylamino, di(n-butyl)amino, diisobutylamino, di(tert-butyl)amino, di(n-pentyl)amino, diisopentylamino, dihexylamino,

methylethylamino, methylisopropylamino, and the like. The "di(C1-C6 alkyl)amino" is preferably dimethylamino, diethylamino, di(n-propyl)amino, diisopropylamino, di(n-butyl)amino, diisobutylamino, di(tert-butyl)amino, di(n-pentyl)amino, diisopentylamino, dihexylamino, methylethylamino, or methylisopropylamino, more preferably di(C1-C4 alkyl)amino, and more preferably dimethylamino.

[0127]

Examples of the "(C3-C7 cycloalkyl)amino" represented by R3 include (C3-C7 cycloalkyl)amino, such as cyclopropylamino, cyclobutylamino, cyclopentylamino, cyclohexylamino, and cycloheptylamino. The "(C3-C7 cycloalkyl)amino" is preferably cyclopropylamino, cyclobutylamino, cyclopentylamino, cyclohexylamino, or cycloheptylamino, and more preferably cyclobutylamino.

[0128]

R3 is preferably substituted or unsubstituted amino, C1-C6 alkyl, halogen, cyano, oxo, hydroxy, carbamoyl, sulfo, C1-C6 alkoxy, or amino(C1-C6 alkyl), more preferably amino that may be substituted with one to two C1-C6 alkyl or C3-C7 cycloalkyl groups, C1-C6 alkyl, halogen, cyano, oxo, hydroxy, carbamoyl, sulfo, C1-C6 alkoxy, or amino(C1-C6 alkyl), more preferably amino, mono- or di(C1-C6 alkyl)amino, (C3-C7 cycloalkyl)amino, or C1-C6 alkyl, more preferably amino, methylamino, ethylamino, isopropylamino, dimethylamino, cyclobutylamino, or methyl, more preferably amino or methyl, and more preferably amino.

[0129]

In the compound represented by Formula (I), m is an integer of 0 to 2, and preferably an integer of 0 to 1.

[0130]

In the compound represented by Formula (I), R4 represents halogen, hydroxy, nitro, cyano, amino, carboxy, (C2-C7 acyl)amino, (C2-C7 acyl)oxy, substituted or unsubstituted C1-C8 alkyl, substituted or unsubstituted C2-C6 alkenyl, substituted or unsubstituted C1-C6 alkoxy, substituted or unsubstituted C3-C7

cycloalkyl, mono- or di(C1-C6 alkyl)amino, substituted or unsubstituted carbamoyl, substituted or unsubstituted C2-C6 alkynyl, substituted or unsubstituted (C1-C6 alkyl)carbonyl, substituted or unsubstituted 4- to 14-membered nitrogen-
5 containing saturated heterocyclic group, or substituted or unsubstituted C6-C14 aromatic hydrocarbon. When two or more R4s are present, R4s may be identical or different.

[0131]

In the present invention, when at least one R4
10 represents substituted C1-C8 alkyl, substituted C2-C6 alkenyl, substituted C1-C6 alkoxy, substituted C3-C7 cycloalkyl, or substituted carbamoyl, examples of the substituents include halogen, carboxy, C1-C6 alkoxy, hydroxy, C1-C6 alkyl that may be substituted with hydroxy, monocyclic 5- to 10-membered
15 unsaturated hydrocarbon, carbamoyl that may be substituted with C1-C6 alkyl or monocyclic 5- to 10-membered unsaturated hydrocarbon, (C2-C7 acyl)oxy, amino that may be substituted with C1-C6 alkyl or C2-C7 acyl, C3-C7 cycloalkyl that may be substituted with hydroxy, (C1-C6 alkoxy) (C1-C6 alkyl), and the
20 like. When two or more of the substituents are present, the substituents may be identical or different.

[0132]

The "C1-C8 alkyl" in the "substituted or unsubstituted C1-C8 alkyl" represented by R4 is preferably methyl, ethyl, n-
25 propyl, isopropyl, n-butyl, isobutyl, sec-butyl, tert-butyl, pentyl, hexyl, heptyl, or octyl, more preferably C1-C6 alkyl, more preferably methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl, tert-butyl, pentyl, or hexyl, and more preferably methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl,
30 sec-butyl, or tert-butyl.

[0133]

The substituent in the "substituted or unsubstituted C1-C8 alkyl" represented by R4 may be, for example, the substituents mentioned above, preferably halogen, amino, hydroxy,
35 carboxy, carbamoyl, alkylcarbamoyl, acylamino, alkoxy,

hydroxycycloalkyl, or acyloxy, more preferably halogen, amino, hydroxy, carboxy, carbamoyl, (C1-C6 alkyl)carbamoyl, (C2-C7 acyl)amino, C1-C6 alkoxy, C3-C7 cycloalkyl, hydroxy(C3-C7 cycloalkyl), or (C2-C7 acyl)oxy, more preferably halogen, amino, hydroxy, carboxy, carbamoyl, (C1-C6 alkyl)carbamoyl, (C1-C6 alkyl)carbonylamino, C1-C6 alkoxy, C3-C7 cycloalkyl, hydroxy(C3-C7 cycloalkyl), or (C1-C6 alkyl)carbonyloxy, and more preferably fluorine, amino, hydroxy, carboxy, carbamoyl, methylcarbamoyl, dimethylcarbamoyl, acetylamino, methoxy, hydroxycyclopropyl, or methylcarbonyloxy.

[0134]

The "substituted or unsubstituted C1-C8 alkyl" represented by R4 is preferably unsubstituted C1-C8 alkyl, or C1-C8 alkyl that may be substituted with halogen, amino, hydroxy, carboxy, carbamoyl, (C1-C6 alkyl)carbamoyl, (C1-C6 alkyl)carbonylamino, C1-C6 alkoxy, C3-C7 cycloalkyl, hydroxy(C3-C7 cycloalkyl), or (C1-C6 alkyl)carbonyloxy, more preferably methyl, ethyl, n-propyl, isopropyl, tert-butyl, difluoromethyl, trifluoromethyl, fluoroethyl, aminoethyl, hydroxymethyl, hydroxyethyl, hydroxypropyl, hydroxydimethylethyl, hydroxymethylpropyl, hydroxymethylbutyl, hydroxyethylbutyl, carboxymethyl, carbamoylmethyl, methylcarbamoylmethyl, dimethylcarbamoylmethyl, acetylaminoethyl, methoxyethyl, hydroxycyclopropylmethyl, hydroxycyclopropylethyl, hydroxycyclobutylmethyl, or methylcarbonyloxyethyl, more preferably methyl, ethyl, n-propyl, tert-butyl, difluoromethyl, hydroxyethyl, hydroxymethylpropyl, hydroxymethylbutyl, hydroxyethylbutyl, carbamoylmethyl, methylcarbamoylmethyl, dimethylcarbamoylmethyl, methoxyethyl, hydroxycyclopropylmethyl, hydroxycyclobutylmethyl, or methylcarbonyloxyethyl, more preferably methyl, difluoromethyl, hydroxymethylpropyl, hydroxymethylbutyl, hydroxycyclobutylmethyl, methoxyethyl, or hydroxycyclobutylmethyl, and more preferably methyl, difluoromethyl, hydroxymethylpropyl, hydroxyethylbutyl, or hydroxycyclobutylmethyl.

[0135]

The "substituted or unsubstituted C2-C6 alkenyl" represented by R4 is preferably unsubstituted C2-C6 alkenyl, more preferably vinyl, allyl, 1-propenyl, 2-methyl-2-propenyl, isopropenyl, 1-, 2- or 3-butenyl, isobutenyl, 2-, 3-, or 4-pentenyl, 2-methyl-2-butenyl, 3-methyl-2-butenyl, 5-hexenyl, 1-cyclopentenyl, 1-cyclohexenyl, or 3-methyl-3-butenyl, and more preferably isobutenyl.

[0136]

Examples of the "C2-C6 alkynyl" in the "substituted or unsubstituted C2-C6 alkynyl" represented by R4 include ethynyl, 1- or 2-propynyl, 1-, 2- or 3-butyne, 1-methyl-2-propynyl, and the like. The "substituted or unsubstituted C2-C6 alkynyl" is preferably unsubstituted C2-C6 alkynyl.

[0137]

The "C1-C6 alkoxy" in the "substituted or unsubstituted C1-C6 alkoxy" represented by R4 is preferably methoxy, ethoxy, n-propoxy, isopropoxy, n-butoxy, isobutoxy, sec-butoxy, tert-butoxy, pentyloxy, or hexyloxy, and more preferably methoxy, ethoxy, n-propoxy, isopropoxy, n-butoxy, isobutoxy, sec-butoxy, and tert-butoxy.

[0138]

The substituent in the "substituted or unsubstituted C1-C6 alkoxy" represented by R4 may be, for example, those mentioned above, and is preferably hydroxy or 5- to 14-membered unsaturated hydrocarbon, more preferably hydroxy or monocyclic 5- to 10-membered unsaturated hydrocarbon, and more preferably hydroxy or phenyl.

[0139]

The "substituted or unsubstituted C1-C6 alkoxy" represented by R4 is preferably C1-C6 alkoxy that may be substituted with hydroxy or 5- to 14-membered unsaturated hydrocarbon, more preferably C1-C6 alkoxy that may be substituted with hydroxy or monocyclic 5- to 10-membered unsaturated hydrocarbon, more preferably C1-C6 alkoxy that may be substituted

with hydroxy or phenyl, and more preferably methoxy, hydroxypropoxy, or benzyloxy.

[0140]

The "substituted or unsubstituted C3-C7 cycloalkyl" represented by R4 is preferably C3-C7 cycloalkyl that may be substituted with hydroxyalkyl, alkoxyalkyl, hydroxycycloalkyl, or unsaturated hydrocarbon carbamoyl, more preferably C3-C7 cycloalkyl that may be substituted with hydroxy(C1-C4 alkyl), (C1-C4 alkoxy) (C1-C4 alkyl), hydroxy(C3-C7 cycloalkyl), or (C6-C14 aromatic hydrocarbon) carbamoyl, more preferably C3-C7 cycloalkyl that may be substituted with hydroxy(C1-C4 alkyl), (C1-C4 alkoxy) (C1-C4 alkyl), hydroxy(C3-C7 cycloalkyl), or phenylcarbamoyl, more preferably cyclopropyl, hydroxymethyl cyclopropyl, methoxymethyl cyclopropyl, hydroxycyclopropyl cyclopropyl, or phenylcarbamoyl cyclopropyl, more preferably cyclopropyl or hydroxymethyl cyclopropyl, and more preferably cyclopropyl.

[0141]

The "mono- or di(C1-C6 alkyl)amino" represented by R4 is preferably methylamino, ethylamino, n-propylamino, isopropylamino, n-butylamino, isobutylamino, tert-butylamino, n-pentylamino, isopentylamino, hexylamino, dimethylamino, diethylamino, di(n-propyl)amino, diisopropylamino, di(n-butyl)amino, diisobutylamino, di(tert-butyl)amino, di(n-pentyl)amino, diisopentylamino, dihexylamino, methylethylamino, or methylisopropylamino, more preferably methylamino, ethylamino, n-propylamino, isopropylamino, n-butylamino, isobutylamino, tert-butylamino, dimethylamino, diethylamino, di(n-propyl)amino, diisopropylamino, di(n-butyl)amino, diisobutylamino, di(tert-butyl)amino, methylethylamino, or methylisopropylamino, and more preferably dimethylamino.

[0142]

The "substituted or unsubstituted carbamoyl" represented by R4 is preferably carbamoyl that may be substituted with alkyl, more preferably carbamoyl that may be substituted

with C1-C6 alkyl, and more preferably carbamoyl, methylcarbamoyl, or dimethylcarbamoyl.

[0143]

Examples of the alkylcarbonyl in the "substituted or
5 unsubstituted (C1-C6 alkyl)carbonyl" represented by R4 include
straight or branched (C1-C6 alkyl)carbonyl, such as
methylcarbonyl, ethylcarbonyl, n-propylcarbonyl,
isopropylcarbonyl, n-butylcarbonyl, isobutylcarbonyl, tert-
butylcarbonyl, n-pentylcarbonyl, isopentylcarbonyl, and
10 hexylcarbonyl.

[0144]

Examples of the "nitrogen-containing saturated
heterocyclic group" in the "substituted or unsubstituted 4- to
14-membered nitrogen-containing saturated heterocyclic group"
15 represented by R4 include morpholinyl, azetidiny, pyrrolidinyl,
piperidinyl, piperazinyl, and the like.

[0145]

Examples of the "substituted or unsubstituted C6-C14
aromatic hydrocarbon" represented by R4 include C6-C14 aromatic
20 hydrocarbon that may be substituted with methyl, such as phenyl,
toluyl, xylyl, naphthyl, anthracenyl, phenanthryl, fluorenyl, and
tetrahydronaphthyl.

[0146]

R4 is preferably halogen, hydroxy, nitro, cyano, amino,
25 carboxy, (C2-C7 acyl)amino, (C2-C7 acyl)oxy, substituted or
unsubstituted C1-C8 alkyl, substituted or unsubstituted C2-C6
alkenyl, substituted or unsubstituted C1-C6 alkoxy, substituted
or unsubstituted C3-C7 cycloalkyl, mono- or di(C1-C6 alkyl)amino,
substituted or unsubstituted carbamoyl, substituted or
30 unsubstituted C2-C6 alkynyl, substituted or unsubstituted (C1-C6
alkyl)carbonyl, substituted or unsubstituted 4- to 14-membered
nitrogen-containing saturated heterocyclic group, or substituted
or unsubstituted C6-C14 aromatic hydrocarbon, more preferably
halogen, nitro, cyano, carboxy, substituted or unsubstituted C1-
35 C8 alkyl, substituted or unsubstituted C2-C6 alkenyl, substituted

or unsubstituted C1-C6 alkoxy, substituted or unsubstituted C3-C7 cycloalkyl, mono- or di(C1-C6 alkyl)amino, or substituted or unsubstituted carbamoyl, more preferably halogen, nitro, cyano, carboxy, C1-C6 alkyl that may be substituted with halogen, amino, hydroxy, carboxy, carbamoyl, (C1-C6 alkyl)carbamoyl, (C1-C6 alkyl)carbonylamino, C1-C6 alkoxy, C3-C7 cycloalkyl, hydroxy(C3-C7 cycloalkyl), or (C1-C6 alkyl)carbonyloxy, C2-C6 alkenyl, C1-C6 alkoxy that may be substituted with hydroxy or monocyclic 5- to 10-membered unsaturated hydrocarbon, C3-C7 cycloalkyl that may be substituted with hydroxy, hydroxy(C1-C4 alkyl), (C1-C4 alkoxy)(C1-C4 alkyl), hydroxy(C3-C7 cycloalkyl), or (C6-C14 aromatic hydrocarbon)-substituted carbamoyl, mono- or di(C1-C6 alkyl)amino, or carbamoyl that may be substituted with C1-C6 alkyl, more preferably fluorine, chlorine, bromine, iodine, nitro, cyano, carboxy, methyl, ethyl, n-propyl, isopropyl, tert-butyl, difluoromethyl, trifluoromethyl, fluoroethyl, aminoethyl, hydroxymethyl, hydroxyethyl, hydroxypropyl, hydroxydimethylethyl, hydroxymethylpropyl, hydroxymethylbutyl, hydroxyethylbutyl, carboxymethyl, carbamoylmethyl, methylcarbamoylmethyl, dimethylcarbamoylmethyl, acetylaminoethyl, methoxyethyl, hydroxycyclopropylmethyl, hydroxycyclopropylethyl, hydroxycyclobutylmethyl, methylcarbonyloxyethyl, isobutenyl, methoxy, hydroxypropoxy, cyclopropyl, hydroxymethyl cyclopropyl, methoxymethyl cyclopropyl, hydroxycyclopropyl cyclopropyl, phenylcarbamoyl cyclopropyl, benzyloxy, dimethylamino, carbamoyl, methylcarbamoyl, or dimethylcarbamoyl, more preferably fluorine, chlorine, bromine, nitro, cyano, carboxy, methyl, ethyl, n-propyl, tert-butyl, difluoromethyl, hydroxyethyl, hydroxymethylpropyl, hydroxymethylbutyl, hydroxyethylbutyl, carbamoylmethyl, methylcarbamoylmethyl, dimethylcarbamoylmethyl, methoxyethyl, hydroxycyclopropylmethyl, hydroxycyclobutylmethyl, methylcarbonyloxyethyl, methoxy, cyclopropyl, hydroxymethyl cyclopropyl, dimethylamino, carbamoyl, methylcarbamoyl, or dimethylcarbamoyl, more preferably fluorine, chlorine, bromine, cyano, methyl, difluoromethyl, hydroxymethylpropyl,

hydroxymethylbutyl, hydroxyethylbutyl, methoxyethyl,
hydroxycyclobutylmethyl, or cyclopropyl, and more preferably
fluorine, chlorine, bromine, cyano, methyl, difluoromethyl,
hydroxymethylpropyl, hydroxyethylbutyl, or
5 hydroxycyclobutylmethyl.

[0147]

In the compound represented by Formula (I), n is an
integer of 0 to 5, and preferably an integer of 0 to 3.

[0148]

10 In the compound represented by Formula (I), R5 and R6
may be identical or different, and each represent hydrogen or C1-
C6 alkyl; R5 and R6 may otherwise form oxo or thioxo together.
When R5 and R6 are identical or different, and each represent
hydrogen or C1-C6 alkyl, at least one of R5 and R6 is hydrogen.

15 [0149]

The "C1-C6 alkyl" represented by R5 and R6 may be
straight or branched. Examples include methyl, ethyl, n-propyl,
isopropyl, n-butyl, isobutyl, tert-butyl, n-pentyl, isopentyl,
hexyl, and the like, with C1-C4 alkyl being preferable, and
20 methyl being more preferable.

[0150]

R5 and R6 are preferably identical or different, and
each represent hydrogen or C1-C6 alkyl; R5 and R6 may otherwise
form oxo or thioxo together. When R5 and R6 are identical or
25 different, and each represent hydrogen or C1-C6 alkyl, at least
one of R5 and R6 is hydrogen; more preferably, they are hydrogen,
or may form oxo together.

[0151]

As the biphenyl compound represented by Formula (I),
30 preferred is a compound represented by Formula (I) or a salt
thereof, wherein
ring A represents a monocyclic, bridged cyclic, or spirocyclic 4-
to 14-membered nitrogen-containing saturated heterocyclic group
having 1 to 3 nitrogen atoms, 0 to 1 sulfur atoms, and 0 to 2
35 oxygen atoms as heteroatoms;

ring B represents monocyclic or bicyclic 5- to 14-membered unsaturated hydrocarbon or a monocyclic or bicyclic 5- to 14-membered unsaturated heterocyclic group that may be substituted with oxo, that has 0 to 4 nitrogen atoms, 0 to 2 sulfur atoms,
5 and 0 to 3 oxygen atoms as heteroatoms, and that has at least one of nitrogen, sulfur, and oxygen;

R1 represents nitro or cyano;

R2 represents halogen;

R3 represents amino, mono- or di(C1-C6 alkyl)amino, (C3-C7
10 cycloalkyl)amino, or C1-C6 alkyl;

R4 represents halogen, nitro, cyano, carboxy, substituted or unsubstituted C1-C8 alkyl, substituted or unsubstituted C2-C6 alkenyl, substituted or unsubstituted C1-C6 alkoxy, substituted or unsubstituted C3-C7 cycloalkyl, mono- or di(C1-C6 alkyl)amino,
15 or substituted or unsubstituted carbamoyl;

when at least one R4 represents substituted C1-C8 alkyl, substituted C2-C6 alkenyl, substituted C1-C6 alkoxy, substituted C3-C7 cycloalkyl, or substituted carbamoyl, the substituent is halogen, carboxy, C1-C6 alkoxy, hydroxy, C1-C6 alkyl that may be
20 substituted with hydroxy, monocyclic 5- to 10-membered unsaturated hydrocarbon, carbamoyl that may be substituted with C1-C6 alkyl or monocyclic 5- to 10-membered unsaturated hydrocarbon, (C2-C7 acyl)oxy, amino that may be substituted with C1-C6 alkyl or C2-C7 acyl, C3-C7 cycloalkyl that may be
25 substituted with hydroxy, or (C1-C6 alkoxy)(C1-C6 alkyl); when two or more of the substituents are present, the substituents may be identical or different;

R5 and R6 may be identical or different, and each represent hydrogen or C1-C6 alkyl; R5 and R6 may otherwise form oxo or
30 thioxo together;

when R5 and R6 are identical or different, and each represent hydrogen or C1-C6 alkyl, at least one of R5 and R6 is hydrogen;

l is an integer of 0 to 2;

m is an integer of 0 to 2;


35 n is an integer of 0 to 5;

when l is 2, two R₂s may be identical or different;
 when m is 2, two R₃s may be identical or different;
 when n is 2 to 5, two to five R₄s may be identical or different.
 [0152]

5 More preferred is a compound represented by Formula (I)
 or a salt thereof, wherein

ring A represents pyrrolidinyl, piperidinyl, piperazinyl,
 azepanyl, diazepanyl,

[0153]



[0154]

10 , 2,7-diazaspiro[3.4]octanyl, 3,7-diazaspiro[3.4]octanyl, 2,7-
 diazaspiro[3.5]nonanyl, 2,8-diazaspiro[3.5]nonanyl, 3,7-
 diazaspiro[3.5]nonanyl, 3,8-diazaspiro[4.4]nonanyl, 3,8-
 15 diazaspiro[4.5]decanyl, or 9-oxa-diazaspiro[3.5]nonanyl;
 ring B represents monocyclic or bicyclic 5- to 14-membered
 unsaturated hydrocarbon or a monocyclic or bicyclic 5- to 14-
 membered unsaturated heterocyclic group that may be substituted
 with oxo, that has 0 to 4 nitrogen atoms, 0 to 2 sulfur atoms,
 20 and 0 to 3 oxygen atoms as heteroatoms, and that has at least one
 of nitrogen, sulfur, and oxygen;

R₁ represents nitro or cyano;

R₂ represents halogen;


25 R₃ represents amino, methylamino, ethylamino, isopropylamino,
 dimethylamino, cyclobutylamino, or methyl;

R₄ represents halogen, nitro, cyano, carboxy, substituted or
 unsubstituted C₁-C₈ alkyl, substituted or unsubstituted C₂-C₆
 alkenyl, substituted or unsubstituted C₁-C₆ alkoxy, substituted
 or unsubstituted C₃-C₇ cycloalkyl, mono- or di(C₁-C₆ alkyl)amino,
 30 or substituted or unsubstituted carbamoyl;

when at least one R₄ is substituted C₁-C₈ alkyl, substituted C₂-

- C6 alkenyl, substituted C1-C6 alkoxy, substituted C3-C7 cycloalkyl, or substituted carbamoyl, the substituent is halogen, carboxy, C1-C6 alkoxy, hydroxy, C1-C6 alkyl that may be substituted with hydroxy, a monocyclic 5- to 10-membered
- 5 unsaturated hydrocarbon, carbamoyl that may be substituted with C1-C6 alkyl or monocyclic 5- to 10-membered unsaturated hydrocarbon, C2-C7 acyl, amino that may be substituted with C1-C6 alkyl or C2-C7 acyl, C3-C7 cycloalkyl that may be substituted with hydroxy, or (C1-C6 alkoxy) (C1-C6 alkyl);
- 10 when two or more of the substituents are present, the substituents may be identical or different;
- R5 and R6 may be identical or different, and each represent hydrogen or C1-C6 alkyl; R5 and R6 may otherwise form oxo or thioxo together;
- 15 when R5 and R6 are identical or different, and each represent hydrogen or C1-C6 alkyl, at least one of R5 and R6 is hydrogen;
- l is an integer of 0 to 2;
- m is an integer of 0 to 2;
- n is an integer of 0 to 5;
- 20 when l is 2, two R2s may be identical or different;
- when m is 2, two R3s may be identical or different;
- when n is 2 to 5, two to five R4s may be identical or different.
- [0155]

- More preferred is a compound represented by Formula (I)
- 25 or a salt thereof, wherein
- ring A represents pyrrolidinyl, piperidinyl, piperazinyl, azepanyl, and diazepanyl,
- [0156]



- 30 [0157]
- , 2,7-diazaspiro[3.4]octanyl, 3,7-diazaspiro[3.4]octanyl, 2,7-

diazaspiro[3.5]nonanyl, 2,8-diazaspiro[3.5]nonanyl, 3,7-diazaspiro[3.5]nonanyl, 3,8-diazaspiro[4.4]nonanyl, 3,8-diazaspiro[4.5]decanyl, or 9-oxa-diazaspiro[3.5]nonanyl; ring B represents monocyclic or bicyclic 5- to 14-membered

5 unsaturated hydrocarbon or a monocyclic or bicyclic 5- to 14-membered unsaturated heterocyclic group that may be substituted with oxo, that has 0 to 4 nitrogen atoms, 0 to 2 sulfur atoms, and 0 to 3 oxygen atoms as heteroatoms, and that has at least one of nitrogen, sulfur, and oxygen;

10 R1 represents nitro or cyano;

R2 represents halogen;

R3 represents amino, methylamino, ethylamino, isopropylamino, dimethylamino, cyclobutylamino, or methyl;

R4 represents halogen, nitro, cyano, carboxy, C1-C8 alkyl that
15 may be substituted with halogen, amino, hydroxy, carboxy, carbamoyl, (C1-C6 alkyl)carbamoyl, (C1-C6 alkyl)carbonylamino, C1-C6 alkoxy, (C1-C6 alkyl)carbonyl, C3-C7 cycloalkyl,

hydroxy(C3-C7 cycloalkyl), or (C1-C6 alkyl)carbonyloxy, C2-C6 alkenyl, C1-C6 alkoxy that may be substituted with hydroxy or
20 monocyclic 5- to 10-membered unsaturated hydrocarbon, C3-C7 cycloalkyl that may be substituted with hydroxy, hydroxy(C1-C4 alkyl), (C1-C4 alkoxy)(C1-C4 alkyl), hydroxy(C3-C7 cycloalkyl), or (C6-C14 aromatic hydrocarbon)-substituted carbamoyl, mono- or di(C1-C6 alkyl)amino, or carbamoyl that may be substituted with
25 C1-C6 alkyl;

R5 and R6 may be identical or different, and each represent hydrogen or C1-C6 alkyl; R5 and R6 may otherwise form oxo or thioxo together;

when R5 and R6 are identical or different, and each represent
30 hydrogen or C1-C6 alkyl, at least one of R5 and R6 is hydrogen;

l is an integer of 0 to 2;

m is an integer of 0 to 2;

n is an integer of 0 to 5;


when l is 2, two R2s may be identical or different;

35 when m is 2, two R3s may be identical or different;

when n is 2 to 5, two to five R4s may be identical or different.
[0158]

More preferred is a compound represented by Formula (I)
or a salt thereof, wherein

5 ring A represents pyrrolidinyl, piperidinyl, piperazinyl,
azepanyl, diazepanyl,
[0159]



[0160]

10 , 2,7-diazaspiro[3.4]octanyl, 3,7-diazaspiro[3.4]octanyl, 2,7-
diazaspiro[3.5]nonanyl, 2,8-diazaspiro[3.5]nonanyl, 3,7-
diazaspiro[3.5]nonanyl, 3,8-diazaspiro[4.4]nonanyl, 3,8-
diazaspiro[4.5]decanyl, or 9-oxa-diazaspiro[3.5]nonanyl;
ring B represents phenyl, naphthyl, pyridyl, pyrazolopyridyl,
15 pyrazolopyrimidinyl, indolyl, indolinyl, 2-oxo-indolinyl,
indazolyl, benzoimidazolyl, benzoisoxazolyl, benzothiazolyl,
benzotriazolyl, imidazopyridinyl, quinolinyl, isoquinolinyl,
quinoxalinyl, quinazolinyl, phthalazinyl, 2-oxo-2,3-
dihydrobenzo[d]oxazolyl, 1,3-dihydroisobenzofuranyl,
20 dihydrobenzooxazinyl, benzodioxolyl, dihydrobenzodioxynyl, or 2-
oxo-2,3-dihydrobenzo[d]thiazolyl;
R1 represents nitro or cyano;
R2 represents fluorine, and is present at the ortho position
relative to R1 on the phenyl;
25 R3 represents amino, methylamino, ethylamino, isopropylamino,
dimethylamino, cyclobutylamino, or methyl (when two or more R3s
are present, R3s may be identical or different);
R4 represents fluorine, chlorine, bromine, iodine, nitro, cyano,
carboxy, methyl, ethyl, n-propyl, isopropyl, tert-butyl,
30 difluoromethyl, trifluoromethyl, fluoroethyl, aminoethyl,
hydroxymethyl, hydroxyethyl, hydroxypropyl, hydroxydimethylethyl,

hydroxymethylpropyl, hydroxymethylbutyl, hydroxyethylbutyl,
 carboxymethyl, carbamoylmethyl, methylcarbamoylmethyl,
 dimethylcarbamoylmethyl, acetylaminoethyl, methoxyethyl,
 hydroxycyclopropylmethyl, hydroxycyclopropylethyl,
 5 hydroxycyclobutylmethyl, methylcarbonyloxyethyl, isobutenyl,
 methoxy, hydroxypropoxy, cyclopropyl, hydroxymethyl cyclopropyl,
 methoxymethyl cyclopropyl, hydroxycyclopropyl cyclopropyl,
 phenylcarbamoyl cyclopropyl, benzyloxy, dimethylamino, carbamoyl,
 methylcarbamoyl, or dimethylcarbamoyl;

10 R5 and R6 may be identical or different, and each represent
 hydrogen or C1-C6 alkyl; R5 and R6 may otherwise form oxo or
 thioxo together;

when R5 and R6 are identical or different, and each represent
 hydrogen or C1-C6 alkyl, at least one of R5 and R6 is hydrogen,

15 l is an integer of 0 to 2,

m is an integer of 0 to 2, and

n is an integer of 0 to 3,

when m is 2, two R3s may be identical or different;


when n is 2 to 3, two to three R4s may be identical or different.

20 [0161]

More preferred is a compound represented by Formula (I)
 or a salt thereof, wherein

ring A represents pyrrolidinyl, piperidinyl, azepanyl,
 diazepanyl,

25 [0162]



[0163]

, 2,7-diazaspiro[3.4]octanyl, 3,7-diazaspiro[3.4]octanyl, 2,7-
 diazaspiro[3.5]nonanyl, 2,8-diazaspiro[3.5]nonanyl, or 9-oxa-
 30 diazaspiro[3.5]nonanyl;

ring B represents phenyl, naphthyl, pyridyl, pyrazolopyridyl,

indolyl, indolinyl, indazolyl, benzoimidazolyl, benzoisoxazolyl, benzotriazolyl, quinolinyl, 2-oxo-2,3-dihydrobenzo[d]oxazolyl, 1,3-dihydroisobenzofuranyl, dihydrobenzooxazinyl, or 2-oxo-2,3-dihydrobenzo[d]thiazolyl;

5 R1 represents cyano;

R2 represents fluorine, and is present at the ortho position relative to R1 on the phenyl;

R3 represents amino, methylamino, ethylamino, isopropylamino, dimethylamino, cyclobutylamino, or methyl (when two or more R3s
10 are present, R3s may be identical or different);

R4 represents fluorine, chlorine, bromine, nitro, cyano, carboxy, methyl, ethyl, n-propyl, tert-butyl, difluoromethyl, hydroxyethyl, hydroxymethylpropyl, hydroxymethylbutyl, hydroxyethylbutyl, carbamoylmethyl, methylcarbamoylmethyl,
15 dimethylcarbamoylmethyl, methoxyethyl, hydroxycyclopropylmethyl, hydroxycyclobutylmethyl, methylcarbonyloxyethyl, methoxy, cyclopropyl, hydroxymethylcyclopropyl, dimethylamino, carbamoyl, methylcarbamoyl, or dimethylcarbamoyl;

R5 and R6 may be identical or different, and each represent
20 hydrogen or C1-C6 alkyl; R5 and R6 may otherwise form oxo or thioxo together;

when R5 and R6 are identical or different, and each represent hydrogen or C1-C6 alkyl, at least one of R5 and R6 is hydrogen;
l is an integer of 0 to 2;

25 m is an integer of 0 to 2;

n is an integer of 0 to 3;


when m is 2, two R3s may be identical or different;

when n is 2 to 3, two to three R4s may be identical or different.

[0164]

30 More preferred is a compound represented by Formula (I) or a salt thereof, wherein ring A represents pyrrolidinyl,

[0165]



[0166]

or 2,8-diazaspiro[3.5]nonanyl;

ring B represents phenyl, indolyl, indazolyl, benzoisoxazolyl, or
5 benzotriazolyl;

R1 represents cyano;

R2 represents fluorine, and is present at the ortho position
relative to R1 on the phenyl;

R3 represents amino or methyl (when two or more R3s are present,

10 R3s may be identical or different);

R4 represents fluorine, chlorine, bromine, cyano, methyl,
difluoromethyl, hydroxymethylpropyl, hydroxymethylbutyl,
hydroxyethylbutyl, methoxyethyl, hydroxycyclobutylmethyl, or
cyclopropyl;

15 R5 and R6 may represent hydrogen or form oxo together;

l is an integer of 0 to 2;

m is an integer of 0 to 2;

n is an integer of 0 to 3;


when m is 2, two R3s may be identical or different;

20 when n is 2 to 3, two to three R4s may be identical or different.

[0167]

More preferred is a compound represented by Formula (I)
or a salt thereof, wherein ring A represents pyrrolidinyl,

[0168]



25

[0169]

ring B represents phenyl, indolyl, indazolyl, benzoisoxazolyl, or
benzotriazolyl;

R1 represents cyano;

30 R2 represents fluorine, and is present at the ortho position
relative to R1 on the phenyl;

R3 represents amino (when two or more R3s are present, R3s may be identical or different);

R4 represents fluorine, chlorine, bromine, cyano, methyl, difluoromethyl, hydroxymethylpropyl, hydroxyethylbutyl, or

5 hydroxycyclobutylmethyl;

R5 and R6 may represent hydrogen or form oxo together;

l is an integer of 0 to 2;

m is an integer of 0 to 2;

n is an integer of 0 to 3;


10 when m is 2, two R3s may be identical or different;

when n is 2 to 3, two to three R4s may be identical or different.

[0170]

More preferred is a compound represented by Formula (I) or a salt thereof, wherein ring A represents pyrrolidinyl,

15 [0171]



[0172]

ring B represents phenyl or benzotriazolyl;

R1 represents cyano;

20 R2 represents fluorine, and is present at the ortho position relative to R1 on the phenyl;

R3 represents amino (when two or more R3s are present, R3s may be identical or different);

R4 represents fluorine, bromine, hydroxymethylpropyl, or

25 hydroxyethylbutyl;

R5 and R6 may represent hydrogen or form oxo together;

l is an integer of 0 to 2;

m is an integer of 0 to 2;

n is an integer of 0 to 3;

30 when m is 2, two R3s may be identical or different;

when n is 2 to 3, two to three R4s may be identical or different.

[0173]

In an embodiment of the present invention, preferred as

the biphenyl compound represented by Formula (I) is a compound represented by Formula (I) or a salt thereof, wherein

ring A represents a monocyclic or bridged cyclic 4- to 8-membered nitrogen-containing saturated heterocyclic group having 1 to 3

nitrogen atoms as heteroatoms;
ring B represents monocyclic 5- to 8-membered unsaturated hydrocarbon or a bicyclic 8- to 14-membered unsaturated heterocyclic group having 1 to 4 nitrogen atoms as heteroatoms;

R1 represents cyano;

R2 represents halogen;

R3 represents amino;

R4 represents halogen or C1-C8 alkyl that may be substituted with hydroxy; when two or more of the substituents are present, the substituents may be identical or different;

R5 and R6 may represent hydrogen or form oxo together;

l is an integer of 0 to 2;

m is an integer of 0 to 2;

n is an integer of 0 to 3;

when l is 2, two R2s may be identical or different;


when m is 2, two R3s may be identical or different;

when n is 2 to 3, two to three R4s may be identical or different.

[0174]

More preferred is a compound represented by Formula (I) or a salt thereof, wherein ring A represents pyrrolidinyl;

[0175]



[0176]

ring B represents phenyl or benzotriazolyl;

R1 represents cyano;

R2 represents halogen;

R3 represents amino;

R4 represents halogen or C1-C8 alkyl that may be substituted with

hydroxy; when two or more of the substituents are present, the substituents may be identical or different;

R5 and R6 may represent hydrogen or form oxo together;

l is an integer of 0 to 2;

5 m is an integer of 0 to 2;

n is an integer of 0 to 3;

when l is 2, two R2s may be identical or different;


when m is 2, two R3s may be identical or different;

when n is 2 to 3, two to three R4s may be identical or different.

10 [0177]

More preferred is a compound represented by Formula (I) or a salt thereof, wherein ring A represents pyrrolidinyl,

[0178]



15 [0179]

ring B represents phenyl or benzotriazolyl;

R1 represents cyano;

R2 represents fluorine;

R3 represents amino;

20 R4 represents fluorine, bromine, or hydroxy(C1-C8 alkyl); when two or more of the substituents are present, the substituents may be identical or different;

R5 and R6 may represent hydrogen or form oxo together;

l is an integer of 0 to 2;

25 m is an integer of 0 to 1;

n is an integer of 0 to 3;


when l is 2, two R2s may be identical or different;

when n is 2 to 3, two to three R4s may be identical or different.

[0180]

30 More preferred is a compound represented by Formula (I) or a salt thereof, wherein ring A represents pyrrolidinyl,

[0181]



[0182]

ring B represents phenyl or benzotriazolyl;

R1 represents cyano;

5 R2 represents fluorine;

R3 represents amino;

R4 represents fluorine, bromine, hydroxymethylpropyl, or hydroxyethylbutyl; when two or more of the substituents are present, the substituents may be identical or different,

10 R5 and R6 may represent hydrogen or form oxo together;

l is an integer of 0 to 2;

m is an integer of 0 to 1;

n is an integer of 0 to 3;

when l is 2, two R2s may be identical or different;

15 when n is 2 to 3, two to three R4s may be identical or different.

[0183]

Specific examples of the biphenyl compounds according to the present invention include, but are not limited to, Example compounds 1 to 295 produced in the Examples below.

20 [0184]

The following are examples of preferable biphenyl compounds:

4-[5-[(3S)-3-aminopyrrolidine-1-carbonyl]-2-[2-fluoro-4-(2-hydroxy-2-methyl-propyl)phenyl]phenyl]-2-fluoro-benzonitrile;

25 4-[5-[(3-endo)-3-amino-8-azabicyclo[3.2.1]octane-8-carbonyl]-2-[2-fluoro-4-(2-hydroxy-2-methyl-propyl)phenyl]phenyl]-2-fluoro-benzonitrile;

5'-((1S,2S,4R)-rel-2-amino-7-azabicyclo[2.2.1]heptane-7-carbonyl)-2'',3-difluoro-4''-(2-hydroxy-2-methylpropyl)-

30 [1,1':2',1''-terphenyl]-4-carbonitrile-isomer-B;

5'-((1S,2S,4R)-rel-2-amino-7-azabicyclo[2.2.1]heptane-7-carbonyl)-2'-(6,7-difluoro-1-(2-hydroxy-2-methylpropyl)-1H-benzo[d][1,2,3]triazol-5-yl)-3-fluoro-[1,1'-biphenyl]-4-

carbonitrile-isomer-B;

5'-((1S,2S,4R)-rel-2-amino-7-azabicyclo[2.2.1]heptane-7-carbonyl)-2'-(1-(2-ethyl-2-hydroxybutyl)-6,7-difluoro-1H-benzo[d][1,2,3]triazol-5-yl)-3-fluoro-[1,1'-biphenyl]-4-

5 carbonitrile-isomer-X;

5'-((1S,2S,4R)-rel-2-amino-7-azabicyclo[2.2.1]heptane-7-carbonyl)-2'-(7-bromo-6-fluoro-1-(2-hydroxy-2-methylpropyl)-1H-benzo[d][1,2,3]triazol-5-yl)-3-fluoro-[1,1'-biphenyl]-4-carbonitrile-isomer-X;

10 5'-((1S,2S,4R)-rel-2-amino-7-azabicyclo[2.2.1]heptan-7-yl)methyl)-2'-(6,7-difluoro-1-(2-hydroxy-2-methylpropyl)-1H-benzo[d][1,2,3]triazol-5-yl)-3-fluoro-[1,1'-biphenyl]-4-carbonitrile-isomer-X.

[0185]


15 Next, the methods for producing the compounds of the present invention are described.

[0186]

The biphenyl compound represented by Formula (I) may be produced, for example, through the production methods below or
20 the methods described in the Examples. However, the methods for producing the biphenyl compound represented by Formula (I) are not limited to these reaction examples.

In steps 1 to 9, in the formulas, L1, L2, and L3 each individually represent a leaving group, NH₂, or OH,
25 W represents hydroxy, C1-C6 alkoxy, or


[0187]



[0188]

, Q1 represents L1 or


30 [0189]



[0190]

, Q2 represents L2 or


[0191]



[0192]

and X represents oxygen or sulfur, and E1 represents hydrogen or substituted or unsubstituted C1-C6 alkyl, wherein when E1 is substituted or unsubstituted C1-C6 alkyl, E1, taken together with the BOO, may form a ring. Ring A, ring B, R1, R2, R3, R4, R5, R6, l, m, and n are as defined above. Further, R5a represents C1-C6 alkyl.

10 [0193]



[0194]

Step 1: Suzuki Reaction

This step represents a method for producing a compound represented by Formula (IV) through a Suzuki reaction using a compound represented by Formula (II).

[0195]

This step may be performed in accordance with a commonly known method (e.g., the method disclosed in Chemical Reviews, Vol. 95, p. 2457, 1995). Protection of a substituent, removal or conversion of the protecting group, and conversion of leaving groups L1, L2, and L3 can be suitably performed.

[0196]

Examples of the leaving groups represented by L1, L2, and L3 include halogen, such as chlorine, bromine, and iodine;

organic sulfonyloxy groups, such as trifluoromethylsulfonyloxy and p-tolylsulfonyloxy; and the like.

[0197]

The amount of the aromatic boronic acid or aromatic
5 boronic acid ester (III) used may be 0.5 to 10 moles, and preferably 0.8 to 3 moles, per mole of the compound represented by Formula (II).

[0198]

Examples of transition metal catalysts include
10 palladium catalysts, such as palladium acetate, tetrakis(triphenylphosphine)palladium, 1,1'-bis(diphenylphosphino)ferrocene-palladium(II) dichloride, bis(dibenzylideneacetone)dipalladium(0), and tris(dibenzylideneacetone)dipalladium(0); nickel catalysts, such
15 as nickel chloride; and the like.

[0199]

As necessary, a ligand may be added. Examples of ligands include triphenylphosphine, tricyclohexylphosphine, (diphenylphosphino)ferrocene, 2-dicyclohexylphosphino-2',4',6'-
20 triisopropylbiphenyl, Silica-SMAP, and the like. The amount of the transition metal catalyst used varies depending on the type of catalyst. The amount of the transition metal catalyst used is usually 0.0001 to 1 mole, and preferably 0.01 to 0.5 moles, per mole of the compound represented by Formula (II). The amount of
25 the ligand used is usually 0.0001 to 4 moles, and preferably 0.01 to 2 moles, per mole of the compound represented by Formula (II).

[0200]

Examples of bases include organic amines, such as triethylamine; alkali metal salts, such as sodium carbonate,
30 potassium carbonate, cesium carbonate, sodium phosphate, tripotassium phosphate, sodium hydroxide, and potassium hydroxide; alkali metal alkoxides, such as sodium methoxide, sodium ethoxide, sodium tert-butoxide, and potassium tert-butoxide; and the like. The amount of the base used is usually
35 0.1 to 10 moles, and preferably 1 to 5 moles, per mole of the

compound represented by Formula (II).

[0201]

The solvent is not limited as long as it does not adversely affect the reaction. Examples include toluene, acetonitrile, 1,2-dimethoxyethane, tetrahydrofuran, 1,4-dioxane, ethanol, N,N-dimethylformamide, water, mixed solvents thereof, and the like. The reaction time is 0.1 to 7 days, and preferably 0.5 to 24 hours. The reaction temperature is 0°C to the boiling temperature of the solvent, and preferably 20°C to 160°C.


10 [0202]

The thus-obtained compound represented by Formula (IV) can be subjected to the subsequent step after or without isolation or purification by known isolation and purification means, such as concentration, vacuum concentration, crystallization, solvent extraction, reprecipitation, and chromatography.

[0203]

It is also possible to first perform a reaction for converting L2 into

20 [0204]



[0205]

by reacting the compound represented by Formula (V) or (VI) with the compound represented by Formula (II), as in step 2 described below.

25

[0206]

Step 2: Suzuki Reaction


This step represents a method for producing a compound represented by Formula (I') through a Suzuki reaction using the compound represented by Formula (IV).

30

[0207]

This step can be performed as in step 1. When L2 (if the reaction for converting L2 into

[0208]




[0209]

is performed first, then L1) is a boronic acid or a boronic acid ester derivative, compound (VI) is used for the reaction.

5 [0210]

L3 in (VI) is the same as L1 and L2 in step 1, and the amount of (VI) used is usually 1 to 10 moles, and preferably 1 to 5 moles, per mole of the compound represented by Formula (IV).

[0211]



10

[0212]

Step 3: Boronic Acid Esterification Reaction

This step represents a method for producing a compound represented by Formula (IX), in which L2 has been converted into a boronic acid ester through a boronic acid esterification reaction, using a compound represented by Formula (IV) and diborane compound (VIII) in the presence of a transition metal catalyst and a base, optionally using a ligand.

15

[0213]

The amount of diborane compound (VIII) used is 1 to 10 moles, and preferably 1 to 5 moles, per mole of the compound represented by Formula (IV).

20

[0214]

The transition metal catalyst may be the same as in step 1.

25

[0215]

As a base, potassium acetate, sodium acetate, and the like may be used, in addition to those mentioned in step 1.

[0216]

The ligand may be the same as in step 1, with Silica-

30

SMAP being preferable.

[0217]

The solvent may be the same as in step 1.


[0218]

5 The reaction temperature is usually 0 to 200°C, and preferably 50 to 160°C. The reaction time is usually 5 minutes to 3 days, preferably 5 minutes to 10 hours.

[0219]

10 Before performing step 3, it is possible to first introduce


[0220]



[0221]

15 into a compound represented by Formula (II); afterward, a boronic acid esterification reaction with respect to L1 may be performed as in step 3.

[0222]



[0223]

20 Step 4: Amidation Reaction

This step represents a method for producing a compound represented by Formula (XI) through an amidation reaction using a carboxylic acid compound represented by Formula (VII), an amine compound represented by Formula (X), and a condensation agent.

25 [0224]

The amount of amine compound (X) used is 0.5 to 10 moles, and preferably 0.8 to 5 moles, per mole of the compound represented by Formula (VII).

[0225]

30 Examples of condensation agents include benzotriazol-1-yloxy-trisdimethylaminophosphonium salts, 4-(4,6-dimethoxy-1,3,5-

triazin-2-yl)-4-methylmorpholinium chloride, a combination of 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide and 1-hydroxybenzotriazole, O-(7-azabenzotriazo-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate, and the like. The amount
5 added is usually 1 to 100 moles, and preferably 1 to 5 moles, per mole of the compound represented by Formula (VII).

[0226]

A base is optionally added during the above reaction. Examples of bases include organic bases, such as triethylamine,
10 diisopropylethylamine, and pyridine; and inorganic bases, such as potassium carbonate. The amount added is usually 1 to 100 moles, preferably 1 to 10 moles, per mole of the compound represented by Formula (VII).

[0227]

15 The solvent is not particularly limited, and any solvent that does not adversely affect the reaction can be used. Examples of the solvent include toluene, chloroform, tetrahydrofuran, N,N-dimethylformamide, dimethylacetamide, N-methylpyrrolidin-2-one, mixed solvent thereof, and the like.

20 [0228]

The reaction temperature is usually -78 to 200°C, and preferably 0 to 50°C. The reaction time is usually 5 minutes to 3 days, preferably 5 minutes to 10 hours.

[0229]

25 The thus-obtained compound represented by Formula (XI) can be subjected to the subsequent step after or without isolation or purification by known isolation and purification means, such as concentration, vacuum concentration, crystallization, solvent extraction, reprecipitation, and
30 chromatography.

[0230]



[0231]

Step 5: Thionation Reaction

This step represents a method for producing a thioamide compound represented by Formula (XII) through a reaction that uses a compound represented by Formula (XI) and a thionation reagent.

[0232]

Examples of thionation reagents include Lawesson's reagent and the like. The amount of this reagent added may be 1 to 10 moles, and preferably 1 to 5 moles, per mole of the compound represented by Formula (XI).

[0233]

The solvent may be the same as in step 1.


[0234]

The reaction temperature is usually 0 to 200°C, and preferably 0 to 100°C. The reaction time is usually 5 minutes to 3 days, and preferably 5 minutes to 10 hours.

[0235]

The thus-obtained compound represented by Formula (XII) can be subjected to the subsequent step after or without isolation or purification by known isolation and purification means, such as concentration, vacuum concentration, crystallization, solvent extraction, reprecipitation, and chromatography.

[0236]



[0237]

Step 6: Reaction of Reduction to Primary Alcohol


This step represents a method for producing a primary alcohol compound (XIII) by a reaction using the carboxylic acid compound represented by Formula (VII) and a reducing agent. Examples of reducing agents include borane, lithium aluminum hydride, and the like. Further, sodium borohydride, lithium

borohydride and the like are also exemplified after the formation of the active ester in the system. Examples of active esterification agents include WSC HCl and the like used together with HATU and HOBT. The amount of the reducing agent to be added is usually 1 to 10 moles, and preferably 1 to 5 moles, per mole of the compound represented by Formula (VII). The amount of the active esterification agent to be added is usually 1 to 10 moles, and preferably 1 to 5 moles, per mole of the compound represented by Formula (VII). The solvent may be the same as in step 1. The reaction temperature is usually 0 to 200°C, and preferably 0 to 100°C. The reaction time is usually 5 minutes to 3 days, preferably 5 minutes to 10 hours.

[0238]

The thus-obtained compound represented by Formula (XIII) can be subjected to the subsequent step after or without isolation or purification by known isolation and purification means, such as concentration, vacuum concentration, crystallization, solvent extraction, reprecipitation, and chromatography.

[0239]



[0240]

Step 7: Alkylation Reaction

This step represents a method for producing a secondary alcohol compound (XV) through a reaction that uses an aldehyde compound represented by Formula (XIV) and an alkyl metal reagent. Examples of alkyl metal reagents include alkyl lithium, alkyl magnesium reagents, and the like. The amount of the alkyl metal reagent to be added is usually 1 to 10 moles, and preferably 1 to 5 moles, per mole of the compound represented by Formula (XIV). The solvent may be the same as in step 5. The reaction

5 The thus-obtained compound represented by Formula (XV)
can be subjected to the subsequent step after or without
isolation or purification by known isolation and purification
means, such as concentration, vacuum concentration,
crystallization, solvent extraction, reprecipitation, and
10 chromatography.




Step 8: Reaction of Oxidation to Aldehyde or Ketone

[0244]

Examples of oxidants include manganese dioxide, tetrapropylammonium perruthenate, sulfur trioxide pyridine complex, Dess-Martin periodinane, pyridinium chlorochromate, and the like. The amount of the oxidant to be added is usually 1 to 10 moles, and preferably 1 to 5 moles, per mole of the compound represented by Formula (XVI). The solvent may be the same as in step 5. The reaction temperature is usually -78 to 200°C, and preferably 0 to 100°C. The reaction time is usually 5 minutes to 3 days, preferably 5 minutes to 10 hours. The thus-obtained compound represented by Formula (XVII) can be subjected to the subsequent step after or without isolation or purification by known isolation and purification means, such as concentration, vacuum concentration, crystallization, solvent extraction,

reprecipitation, and chromatography.

[0245]



[0246]

5 Step 9: Reductive Amination Reaction

This step represents a method for producing a compound represented by Formula (XVIII) using an amine compound represented by Formula (X) and a reducing agent with the aldehyde or ketone compound represented by Formula (XVII).

10 [0247]

Examples of reducing agents include sodium triacetoxyborohydride, sodium cyanoborohydride, sodium borohydride, and the like. The amount of the reducing agent to be added is usually 1 to 10 moles, and preferably 1 to 5 moles, per

15 mole of the compound represented by Formula (XVII).

[0248]

The solvent is not particularly limited, and any solvent that does not interfere with the reaction can be used. Examples of solvents include toluene, chloroform,

20 tetrahydrofuran, dichloromethane, methanol, ethanol, and the like; as well as a mixed solvent thereof.

[0249]

The thus-obtained compound represented by Formula (XVIII) can be subjected to the subsequent step after or without isolation or purification by known isolation and purification means, such as concentration, vacuum concentration, crystallization, solvent extraction, reprecipitation, and chromatography.

[0250]

30 The conversion of substituents W and X, and leaving

groups L1, L2, and L3, may be suitably performed.

[0251]

In any of steps 1 to 9, protection of a substituent, and removal or conversion of the protecting group, can be
5 suitably performed. For example, for functional groups such as amino, imino, hydroxy, carboxy, carbonyl, and amide groups, as well as functional groups having an active proton, such as indole, protected reagents can be used, or a protecting group can be introduced into such a functional group according to a usual
10 method; afterward, the protecting group can be removed in an appropriate step in each production method.

[0252]

The protecting group of an amino group or protecting group of an imino group is not particularly limited, insofar as
15 it has a protecting function. Examples of such protecting groups include aralkyl groups, such as benzyl, p-methoxybenzyl, 3,4-dimethoxybenzyl, o-nitrobenzyl, p-nitrobenzyl, benzhydryl, trityl, and cumyl; lower alkanoyl groups, such as formyl, acetyl, propionyl, butyryl, pivaloyl, trifluoroacetyl, and
20 trichloroacetyl; benzoyl; arylalkanoyl groups, such as phenylacetyl and phenoxyacetyl; lower alkoxy carbonyl groups, such as methoxycarbonyl, ethoxycarbonyl, propyloxycarbonyl, and tert-butoxycarbonyl; aralkyloxycarbonyl groups, such as p-nitrobenzyloxycarbonyl and phenethyloxycarbonyl; lower alkylsilyl
25 groups, such as trimethylsilyl and tert-butyldimethylsilyl; tetrahydropyranyl; trimethylsilylethoxymethyl; lower alkylsulfonyl groups, such as methylsulfonyl, ethylsulfonyl, and tert-butylsulfonyl; lower alkylsulfinyl groups, such as tert-butylsulfinyl; arylsulfonyl groups, such as benzenesulfonyl and
30 toluenesulfonyl; and imido groups, such as phthalimido. In particular, trifluoroacetyl, acetyl, tert-butoxycarbonyl, benzyloxycarbonyl, trimethylsilylethoxymethyl, cumyl, and the like are preferable.

[0253]

35 The protecting group of a hydroxy group is not

particularly limited insofar as it has a protecting function.

Examples of such protecting groups include lower alkyl groups, such as methyl, ethyl, propyl, isopropyl, and tert-butyl; lower alkylsilyl groups, such as trimethylsilyl and tert-

5 butyldimethylsilyl; lower alkoxymethyl groups, such as methoxymethyl and 2-methoxyethoxymethyl; tetrahydropyranyl; trimethylsilylethoxymethyl; aralkyl groups, such as benzyl, p-methoxybenzyl, 2,3-dimethoxybenzyl, o-nitrobenzyl, p-nitrobenzyl, and trityl; and acyl groups, such as formyl, acetyl, and
10 trifluoroacetyl. In particular, methyl, methoxymethyl, tetrahydropyranyl, trimethylsilylethoxymethyl, tert-butyldimethylsilyl, and acetyl are preferable.

[0254]

The protecting group of a carboxy group is not
15 particularly limited insofar as it has a protecting function. Examples of such protecting groups include lower alkyl groups, such as methyl, ethyl, propyl, isopropyl, and tert-butyl; halo-lower-alkyl groups, such as 2,2,2-trichloroethyl; lower alkenyl groups, such as allyl; trimethylsilylethoxymethyl; and aralkyl
20 groups, such as benzyl, p-methoxybenzyl, p-nitrobenzyl, benzhydryl, and trityl. In particular, methyl, ethyl, tert-butyl, allyl, benzyl, p-methoxybenzyl, trimethylsilylethoxymethyl, and the like are preferable.

[0255]

25 The protecting group of a carbonyl group is not particularly limited insofar as it has a protecting function. Examples of such protecting groups include ethylene ketal, trimethylene ketal, dimethyl ketal, ethylene acetal, trimethylene acetal, dimethyl acetal, and like ketals and acetals.

30 [0256]

The protecting group of an amide group or the protecting group of a functional group having an active proton, such as indole, is not particularly limited, insofar as it has a protecting function. Examples of such protecting groups include
35 lower alkyl groups, such as methyl, ethyl, propyl, isopropyl, and

tert-butyl; lower alkylsilyl groups, such as trimethylsilyl and tert-butyldimethylsilyl; lower alkoxyethyl groups, such as methoxyethyl and 2-methoxyethoxyethyl; tetrahydropyranyl; trimethylsilylethoxyethyl; aralkyl groups, such as benzyl, p-methoxybenzyl, 2,3-dimethoxybenzyl, o-nitrobenzyl, p-nitrobenzyl, and trityl; and acyl groups, such as formyl, acetyl, and trifluoroacetyl. In particular, methyl, methoxyethyl, tetrahydropyranyl, trimethylsilylethoxyethyl, tert-butyldimethylsilyl, and acetyl are preferable.

10 [0257]

The method for removing such a protecting group may vary depending on the type of protecting group, stability of the target compound, etc. For example, the following methods can be used: solvolysis using an acid or a base according to the method disclosed in a publication (Protective Groups in Organic Synthesis, third edition, T.W. Green, John Wiley & Sons (1999)) or a similar method, i.e., a method comprising reacting with 0.01 moles or a large excess of an acid, preferably trifluoroacetic acid, formic acid, or hydrochloric acid, or an equimolar to large excessive molar amount of a base, preferably potassium hydroxide or calcium hydroxide; chemical reduction using a metal hydride complex etc.; or catalytic reduction using a palladium-carbon catalyst, Raney nickel catalyst, etc.

[0258]

25 The biphenyl compound represented by Formula (I) can be easily isolated and purified by common isolation and purification means. Examples of such means include solvent extraction, recrystallization, preparative reversed-phase high-performance liquid chromatography, column chromatography, preparative thin-layer chromatography, and the like.

30 [0259]

When the biphenyl compound represented by Formula (I) has isomers such as optical isomers, stereoisomers, rotational isomers, and tautomers, any of the isomers and mixtures thereof is included within the scope of the compound, unless otherwise

35

specified. For example, when the the biphenyl compound represented by Formula (I) has optical isomers, the optical isomer separated from a racemic mixture is also included within the scope of the compound, unless otherwise specified. Each of

5 such isomers can be obtained as a single compound by known synthesis and separation means (e.g., concentration, solvent extraction, column chromatography, and recrystallization).

[0260]

As stated above, unless otherwise specified, the

10 biphenyl compound represented by Formula (I) includes all of the enantiomers and mixtures thereof. The biphenyl compound represented by Formula (I) may be a mixture of R and S enantiomers. Such a mixture may be a mixture comprising 90% or more, 95% or more, or 99% or more of R enantiomer; a mixture

15 comprising 90% or more, 95% or more, or 99% or more of S enantiomer; or the like.

[0261]

Methods for chiral resolution include, for example: a diastereomer method of causing a chiral resolving agent to act on

20 the biphenyl compound represented by Formula (I) to form a salt, and resolving one of the enantiomers using a solubility difference etc. of the obtained salt; a preferential crystallization method of adding one of the enantiomers to a supersaturated solution of a racemate as a seed for

25 crystallization; and column chromatography such as HPLC using a chiral column. A chiral resolving agent that can be used in the diastereomer method can be appropriately selected from, for example, acid resolving agents such as tartaric acid, malic acid, lactic acid, mandelic acid, 10-camphorsulfonic acid, and

30 derivatives thereof; and basic resolving agents such as brucine, strychnine, quinine, and like alkaloid compounds, amino acid derivatives, cinchonidine, and α -methylbenzylamine. One of the enantiomers of the biphenyl compound represented by Formula (I) alone can be obtained not only by obtaining the biphenyl compound

35 represented by Formula (I) as a mixture of enantiomers and then

conducting chiral resolution as above, but also by obtaining one enantiomer of the biphenyl compound represented by Formula (I) through chiral resolution as above or by other methods, and using it as a synthetic raw material of the biphenyl compound

5 represented by Formula (I). Furthermore, methods for obtaining one of the enantiomers of the biphenyl compound represented by Formula (I) or its raw material compound include a method of preferentially obtaining one of the enantiomers by adjusting reaction conditions for a catalyst or the like in a reaction step
10 of generating asymmetric carbon.

[0262]

The biphenyl compound represented by Formula (I) or a salt thereof may be in the form of crystals. Single crystals and polymorphic crystal mixtures are included within the scope of the
15 compound or a salt thereof. Such crystals can be produced by crystallization according to a crystallization method known *per se* in the art. The biphenyl compound represented by Formula (I) or a salt thereof may be a solvate (e.g., a hydrate) or a non-solvate. Any of such forms are included within the scope of the
20 biphenyl compound represented by Formula (I) or a salt thereof. Compounds labeled with an isotope (e.g., ^3H , ^{14}C , ^{35}S , and ^{125}I) are also included within the scope of the biphenyl compound represented by Formula (I) or a salt thereof.

[0263]

25 The salts of the biphenyl compound represented by Formula (I) or of the intermediates thereof refer to common salts used in the field of organic chemistry. The salts are preferably pharmaceutically acceptable salts. Examples of such salts include base addition salts to a carboxy group when the compound has a
30 carboxy group, and acid addition salts to an amino or basic heterocyclic group when the compound has an amino or basic heterocyclic group.

[0264]

Examples of base addition salts include alkali metal
35 salts, such as sodium salts and potassium salts; alkaline earth

metal salts, such as calcium salts and magnesium salts; ammonium salts; and organic amine salts, such as trimethylamine salts, triethylamine salts, dicyclohexylamine salts, ethanolamine salts, diethanolamine salts, triethanolamine salts, procaine salts, and
5 N,N'-dibenzylethylenediamine salts.
[0265]

Examples of acid addition salts include inorganic acid salts, such as hydrochloride, sulfate, nitrate, phosphate, and perchlorate; organic acid salts, such as acetate, formate,
10 maleate, fumarate, tartrate, citrate, ascorbate, and trifluoroacetate; and sulfonates such as methanesulfonate, isethionate, benzenesulfonate, and p-toluenesulfonate.
[0266]

Due to their excellent LSD1 inhibitory activity, the
15 biphenyl compound represented by Formula (I) or a salt thereof is useful as a pharmaceutical preparation for preventing and treating, preferably treating, LSD1-related diseases.
[0267]

Using the biphenyl compound represented by Formula (I)
20 or a salt thereof in combination with one or more other antitumor agents potentiates the antitumor effect. The form of a combination of the biphenyl compound represented by Formula (I) or a salt thereof and one or more other antitumor agents may be a single preparation (i.e., a combination drug) or two or more
25 separate preparations to be administered in combination.
[0268]

In the present invention, the antitumor effect can be evaluated as, for example, reduced tumor volume, tumor growth stasis, or prolonged survival time.
30 [0269]

An embodiment provides an antitumor formulation comprising a combination of the biphenyl compound represented by Formula (I) or a salt thereof and one or more other antitumor agents. Another embodiment provides an antitumor effect
35 potentiators for an antitumor agent, the potentiators comprising

the biphenyl compound represented by Formula (I) or a salt thereof as an active ingredient.

[0270]

The other antitumor agents are not particularly
5 limited. Examples thereof include antimetabolites, antitumor antibiotics, molecular target drugs, platinum-based drugs, and plant alkaloid-based drugs.

[0271]

Examples of antimetabolites include 5-fluorouracil (5-
10 FU), 5-fluoro-2'-deoxyuridine (FdUrd), tegafur, combination drugs of tegafur and uracil (e.g., UFT), combination drugs of tegafur, gimeracil, and oteracil (e.g., TS-1), pemetrexed, trifluridine, combination drugs of trifluridine and *tipiracil hydrochloride* (e.g., Lonsurf), gemcitabine, capecitabine, nelarabine,
15 clofarabine, cytarabine, DNA methylation inhibitors (such as azacitidine, decitabine, and guadecitabine), and the like, with cytarabine or DNA methylation inhibitors, such as azacitidine, decitabine, and guadecitabine, being preferable, and cytarabine, azacitidine, decitabine, or guadecitabine being more preferable.

20 [0272]

Examples of antitumor antibiotics include daunorubicin, doxorubicin, amrubicin, idarubicin, epirubicin, and like anthracycline-based antitumor antibiotics, mitomycin C, bleomycin, and the like, with anthracycline-based antitumor
25 antibiotics being preferable, and daunorubicin being more preferable.

[0273]

Examples of molecular target drugs include all-trans retinoic acid or derivatives thereof, human MDM2 (mouse double
30 minute 2) (HDM2; human double minute 2) inhibitors, and HDAC inhibitors.

[0274]

The all-trans retinoic acid or a derivative thereof is preferably tretinoin (ATRA) or tamibarotene, and more preferably
35 tretinoin (ATRA).

[0275]

The human MDM2 (HDM2) inhibitor is preferably RG7388 (RO5503781), AMG-232, DS-3032b, RG7112 (RO5045337), SAR405838, or MK-8242, and more preferably RG7388.

5 [0276]

Examples of HDAC inhibitors include vorinostat, panobinostat, romidepsin, belinostat, and the like.

[0277]

The molecular target drug is preferably all-trans
10 retinoic acid or a derivative thereof, a human MDM2 (HDM2) inhibitor, or an HDAC inhibitor, and more preferably tretinoin (ATRA) or RG7388.

[0278]

Examples of platinum-based drugs include oxaliplatin,
15 carboplatin, cisplatin, nedaplatin, and the like, with carboplatin or cisplatin being preferable.

[0279]

Examples of plant alkaloid-based drugs include microtubule inhibitors, such as paclitaxel, docetaxel,
20 vinblastine, vincristine, vindesine, vinorelbine, and eribulin, and topoisomerase inhibitors, such as irinotecan (SN-38), nogitecan, and etoposide, with taxane microtubule inhibitors such as paclitaxel and docetaxel or topoisomerase inhibitors such as irinotecan (SN-38), nogitecan, and etoposide being preferable,
25 and paclitaxel, irinotecan (SN-38), or etoposide being more preferable.

[0280]

The one or more other antitumor agents are preferably an antimetabolite, an antitumor antibiotic, a molecular target
30 drug, a platinum-based drug, or a plant alkaloid-based drug, more preferably an antimetabolite, an antitumor antibiotic, all-trans retinoic acid or a derivative thereof, a human MDM2 (HDM2) inhibitor, an HDAC inhibitor, a platinum-based drug, or a plant alkaloid-based drug, more preferably an antimetabolite, an
35 antitumor antibiotic, all-trans retinoic acid or a derivative

thereof, a human MDM2 (HDM2) inhibitor, a platinum-based drug, or a plant alkaloid-based drug, more preferably an antimetabolite, an antitumor antibiotic, all-trans retinoic acid or a derivative thereof, a human MDM2 (HDM2) inhibitor, a platinum-based drug, a topoisomerase inhibitor, or a taxane microtubule inhibitor, more preferably cytarabine, a DNA methylation inhibitor, an anthracycline-based antitumor antibiotic, all-trans retinoic acid or a derivative thereof, a platinum-based drug, a topoisomerase inhibitor, or a taxane microtubule inhibitor, and more preferably cytarabine, azacitidine, decitabine, guadecitabine, daunorubicin, tretinoin (ATRA), RG7388, carboplatin, cisplatin, paclitaxel, irinotecan (SN-38), or etoposide.
[0281]

The tumor targeted in the present invention is not particularly limited as long as the antitumor effect is potentiated. The tumor is preferably a tumor in which the biphenyl compound represented by Formula (I) or a salt thereof exerts an antitumor effect, and more preferably LSD1-related malignant tumors.

[0282]

Examples of the "LSD1-related malignant tumors" include malignant tumors whose incidence can be reduced, and whose symptoms can be remitted, relieved, and/or completely cured by eliminating, suppressing, and/or inhibiting LSD1 function. The type of malignant tumor to be treated by the compound or a salt thereof of the present invention is not particularly limited. Examples of such malignant tumors include head and neck cancers, esophagus cancer, gastric cancer, colon cancer, rectum cancer, liver cancer, gallbladder cancer, cholangiocarcinoma, biliary tract cancer, pancreatic cancer, lung cancer, breast cancer, ovarian cancer, cervical cancer, endometrial cancer, renal cancer, bladder cancer, prostate cancer, testicular tumor, osteosarcoma, soft-tissue sarcoma, leukemia, myelodysplastic syndrome, chronic myeloproliferative disease, malignant lymphoma, multiple myeloma, skin cancer, brain tumor, mesothelioma, and the

like. Preferable examples include lung cancers (e.g., non-small cell lung cancer and small cell lung cancer), leukemia, and myelodysplastic syndromes.

[0283]

5 When the biphenyl compound or a salt thereof and one or more other antitumor agents are used as a pharmaceutical preparation, a pharmaceutical carrier can be added, if required, thereby forming one or more suitable dosage forms according to prevention and treatment purposes. Examples of the dosage form(s)
10 include oral preparations, injections, suppositories, ointments, patches, and the like. In the case of the biphenyl compound or a salt thereof, oral preparations are preferable. Such dosage forms can be formed by methods conventionally known to persons skilled in the art.

15 [0284]

 The administration schedule of the biphenyl compound represented by Formula (I) or a salt thereof and one or more other antitumor agents is suitably selected within the range in which an antitumor effect is potentiated. The active ingredients
20 may be administered simultaneously or separately at intervals. When they are administered separately, the order of administration is not limited.

[0285]

 The biphenyl compound represented by Formula (I) or a
25 salt thereof and one or more other antitumor agents may be formulated into multiple dosage forms, or may be formulated into a single dosage form (i.e., formulated as a combination drug), based on the administration form and administration schedule of each active ingredient. Further, preparations may be produced and
30 sold in a single package suitable for combined use, or preparations may be produced and sold in separate packages.

[0286]

 As the pharmaceutical carrier, various conventional organic or inorganic carrier materials used as preparation
35 materials may be blended as an excipient, binder, disintegrant,

lubricant, or coating agent in solid preparations; or as a solvent, solubilizing agent, suspending agent, isotonizing agent, pH adjuster, buffer, or soothing agent in liquid preparations.

Moreover, pharmaceutical preparation additives, such as

5 antiseptics, antioxidants, colorants, taste-masking or flavoring agents, and stabilizers, may also be used, if required.

[0287]

When oral solid preparations are prepared, an excipient, a binder, a disintegrant, a lubricant, a colorant, a
10 taste-masking or flavoring agent, etc., may be added; and the resulting mixture may be formulated into tablets, coated tablets, granules, powders, capsules, or the like by ordinary methods.

[0288]

Examples of excipients include lactose, sucrose, D-
15 mannitol, glucose, starch, calcium carbonate, kaolin, microcrystalline cellulose, and silicic acid anhydride. Examples of binders include water, ethanol, 1-propanol, 2-propanol, simple syrup, liquid glucose, liquid α -starch, liquid gelatin, D-mannitol, carboxymethyl cellulose, hydroxypropyl cellulose,
20 hydroxypropyl starch, methyl cellulose, ethyl cellulose, shellac, calcium phosphate, polyvinylpyrrolidone, and the like. Examples of disintegrators include dry starch, sodium alginate, powdered agar, sodium hydrogen carbonate, calcium carbonate, sodium lauryl sulfate, stearic acid monoglyceride, lactose, and the like.

25 Examples of lubricants include purified talc, stearic acid salt sodium, magnesium stearate, borax, polyethylene glycol, and the like. Examples of colorants include titanium oxide, iron oxide, and the like. Examples of taste-masking or flavoring agents include sucrose, bitter orange peel, citric acid, tartaric acid,
30 and the like.

[0289]

When liquid preparations for oral administration are prepared, a taste-masking agent, a buffer, a stabilizer, a flavoring agent, etc., may be added; and the resulting mixture
35 may be formulated into oral liquid preparations, syrups, elixirs,

or the like by ordinary methods.

[0290]

In this case, the same taste-masking or flavoring agent as those mentioned above may be used. Examples of the buffer
5 include sodium citrate and the like, and examples of the stabilizer include tragacanth, gum arabic, gelatin, and the like. As necessary, these preparations for oral administration may be coated according to methods known in the art with an enteric coating or other coating for the purpose of, for example,
10 persistence of effects. Examples of such coating agents include hydroxypropyl methylcellulose, ethyl cellulose, hydroxymethyl cellulose, hydroxypropyl cellulose, polyoxyethylene glycol, and Tween 80 (registered trademark).

[0291]

15 When injections are prepared, a pH adjuster, a buffer, a stabilizer, an isotonizing agent, a topical anesthetic, etc., may be added; and the resulting mixture may be formulated into subcutaneous, intramuscular, and intravenous injections by ordinary methods.

20 [0292]

Examples of usable pH adjusters and buffers include sodium citrate, sodium acetate, sodium phosphate, and the like. Examples of usable stabilizers include sodium pyrosulfite, EDTA, thioglycolic acid, and thiolactic acid. Examples of usable
25 topical anesthetics include procaine hydrochloride, lidocaine hydrochloride, and the like. Examples of usable isotonizing agents include sodium chloride, glucose, D-mannitol, glycerin, and the like.

[0293]

30 The preparations of the one or more other antitumor agents also include drug delivery system (DDS) preparations for them. For example, "paclitaxel" includes albumin-bound paclitaxel (e.g., Abraxane), paclitaxel micelles (e.g., NK105), and the like; and "cisplatin" includes cisplatin micelles (e.g., NC-6004)
35 and the like.

[0294]

The amount of the biphenyl compound to be incorporated in each of such dosage unit forms depends on the condition of the patient to whom the compound is administered, the dosage form, etc. In general, in the case of an oral agent, an injection, and a suppository, the amount of the biphenyl compound is preferably 0.05 to 1000 mg, 0.01 to 500 mg, and 1 to 1000 mg, respectively, per dosage unit form.

[0295]

The daily dose of the biphenyl compound in such a dosage form depends on the condition, body weight, age, gender, etc., of the patient, and cannot be generalized. For example, the daily dose of the biphenyl compound for an adult (body weight: 50 kg) may be usually 0.05 to 5000 mg, and preferably 0.1 to 1000 mg; and is preferably administered in one dose, or in two to three divided doses, per day.

Examples

[0296]

The present invention is described below in more detail with reference to Examples. However, the scope of the present invention is not limited to these Examples. The present invention is fully described below by way of Examples; however, it is understood that various changes and modifications by a skilled artisan are possible. Therefore, such changes and modifications are included in the present invention as long as they do not depart from the scope of the invention.

[0297]

The various reagents used in the Examples were obtained from commercial suppliers, unless otherwise specified. For silica gel column chromatography, a SNAP-Ultra (registered trademark) silica prepacked column produced by Biotage was used. Alternatively, for basic silica gel column chromatography, a KP-NH (registered trademark) prepacked column produced by Biotage was used. NMR spectra were measured by using an AL400 (400 MHz;

produced by JEOL), a Mercury 400 (400 MHz; produced by Agilent Technologies, Inc.), or a 500-MHz Bruker Avance III HD NMR Spectrometer (500 MHz; Bruker). When the deuterated solvent contained tetramethylsilane, tetramethylsilane was used as the internal reference. Otherwise, an NMR solvent was used as the internal reference. All of the δ values are shown in ppm. The microwave reaction was performed using an Initiator produced by Biotage.

[0298]

LCMS spectra were measured using an Acquity SQD (quadrupole) produced by Waters Corporation under the following conditions.
Column: Acquity UPLC (registered trademark) BEH C18, 2.1 x 50 mm, 1.7 μ m (produced by Waters Corporation)
MS detection: ESI positive
UV detection: 254 and 280 nm
Column flow rate: 0.5 mL/min
Mobile phase: Water/acetonitrile (0.1% formic acid)
Injection volume: 1 μ L

Gradient (Table 1)

Time (min)	Water	Acetonitrile
0	95	5
0.1	95	5
2.1	5	95
3.0	STOP	

[0299]

Preparative reversed-phase HPLC purification was performed under the following conditions using a preparative separation system available from Gilson, Inc.

Column: Xselect CSH Prep C18 5 μ m OBD (19 x 50 mm) + (19 x 100 mm), produced by Waters Corporation
UV detection: 254 nm
Column flow rate: 18 mL/min
Mobile phase: Water/acetonitrile (0.1% formic acid)
Injection volume: 0.1 to 0.5 mL

[0300]

The symbols stand for the following.

[0301]

s: Singlet

5 d: Doublet

t: Triplet

q: Quartet

dd: Double doublet

dt: Double triplet

10 td: Triple doublet

tt: Triple triplet

ddd: Double double doublet

ddt: Double double triplet

dtd: Double triple doublet

15 tdd: Triple double doublet

m: Multiplet

br: Broad

brs: Broad singlet

THF: Tetrahydrofuran

20 DMF: N,N-dimethylformamide

DME: 1,2-Dimethoxyethane

DMSO: Dimethylsulfoxide

HATU: O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium
hexafluorophosphate

25 TEA: Triethylamine

WSC HCl: 1-Ethyl-3-(3-dimethylaminopropyl)carbodiimide
hydrochloride

t-BuOH: Tertiary butanol

DMAP: N,N-dimethylaminopyridine

30 Pd(PPh₃)₄: Tetrakis(triphenylphosphine)palladium(0)

Pd(dba)₂: Bis(dibenzylideneacetone)palladium(0)

PCy₃: Tricyclohexylphosphine

TFA: Trifluoroacetic acid

Pd(OAc)₂: Palladium acetate

35 KOAc: Potassium acetate

$\text{PdCl}_2(\text{dppf})$: [1,1'-Bis(diphenylphosphino)ferrocene]palladium(II) dichloride

$\text{PdCl}_2(\text{dppf})\text{CH}_2\text{Cl}_2$: [1,1'-Bis(diphenylphosphino)ferrocene]palladium(II) dichloride

5 dichloromethane complex

DMEAD: Di-2-methoxyethyl azodicarboxylate

PPh_3 : Triphenylphosphine

DMA: Dimethylacetamide

MeMgBr : Methylmagnesium bromide

10 EtMgBr : Ethylmagnesium bromide

MTBE: Methyltertiary-butyl ether

DCM: Dichloromethane

Boc_2O : Di-tert-butyl dicarbonate

NBS: N-bromosuccinimide

15 X-phos: 2-dicyclohexylphosphino-2',4',6'-triisopropylbiphenyl

MeOH: Methanol

EtOH: Ethanol

IPE: Diisopropyl ether

TBAF: Tetrabutylammonium fluoride

20 $\text{Pd}_2(\text{dba})_3$: Tris(dibenzylideneacetone)dipalladium(0)

$\text{PdCl}_2(\text{PPh}_3)_2$: Bistriphenylphosphine palladium chloride

S-Phos: 2-Dicyclohexylphosphino-2,6-dimethoxybiphenyl

HOBt: 1-Hydroxybenzotriazole

Pd/C : Palladium on carbon

25 NMP: N-methyl-2-pyrrolidinone.

[0302]

Example 1: Synthesis of 4-[5-[(3S)-3-aminopyrrolidine-1-carbonyl]-2-(p-tolyl)phenyl]benzonitrile

Step 1

30 3-Bromo-4-chloro-benzoic acid (19 g) was dissolved in DMF (160 mL). At 25°C, DMAP (20 g) and WSC HCl (31 g) were added thereto, followed by the addition of t-BuOH (38 mL). The resulting mixture was stirred at room temperature overnight. Ethyl acetate was added thereto, and the resulting mixture was
35 washed sequentially with water and saturated brine. After the

organic layer was dried over anhydrous sodium sulfate, the solvent was distilled off. The residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate) to give tert-butyl 3-bromo-4-chloro-benzoate.

5 [0303]

Step 2

The tert-butyl 3-bromo-4-chloro-benzoate (1.3 g) obtained in step 1 above was dissolved in 1,4-dioxane (8.7 mL). At room temperature, (4-cyanophenyl)boronic acid (768 mg),
10 Pd(PPh₃)₄ (151 mg), and a 2 M Na₂CO₃ aqueous solution (5.4 mL) were added thereto, and the reaction mixture was stirred in a microwave reactor at 120°C for 30 minutes. The reaction mixture was then vacuum-concentrated, and the residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl
15 acetate) to give tert-butyl 4-chloro-3-(4-cyanophenyl)benzoate.

[0304]

Step 3

The tert-butyl 4-chloro-3-(4-cyanophenyl)benzoate (1.1 g) obtained in step 2 above was dissolved in 1,4-dioxane (17 mL).
20 At room temperature, p-tolylboronic acid (932 mg), Pd(dba)₂ (157 mg), tripotassium phosphate (1.5 g), and a solution of 1 M PCy₃ in THF (0.57 mL) were added thereto, and the reaction mixture was stirred in a microwave reactor at 160°C for 30 minutes. After the addition of chloroform, the insoluble matter was filtered off,
25 and the solvent was distilled off. The residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate), and the solvent was distilled off. The residue was dissolved in TFA (2 mL). The solvent was distilled off. Ethyl acetate was added thereto, and the mixture was washed
30 sequentially with water and saturated brine. After the organic layer was dried over anhydrous sodium sulfate, the solvent was distilled off to give 3-(4-cyanophenyl)-4-(p-tolyl)benzoic acid.

[0305]

Step 4

35 The 3-(4-cyanophenyl)-4-(p-tolyl)benzoic acid (10 mg)

obtained in step 3 above, tert-butyl N-[(3S)-pyrrolidin-3-yl]carbamate (6 mg), and HATU (24 mg) were dissolved in THF (0.5 mL). At room temperature, TEA (0.013 mL) was added thereto, followed by stirring at 50°C overnight. The reaction mixture was vacuum-concentrated, and the solvent was distilled off. The residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate) to give tert-butyl N-[(3S)-1-[3-(4-cyanophenyl)-4-(p-tolyl)benzoyl]pyrrolidin-3-yl]carbamate.

[0306]

10 Step 5

The tert-butyl N-[(3S)-1-[3-(4-cyanophenyl)-4-(p-tolyl)benzoyl]pyrrolidin-3-yl]carbamate (15 mg) obtained in step 4 above was dissolved in TFA (0.3 mL), and the progress of the reaction was confirmed by LCMS, followed by vacuum concentration. The residue was purified by reversed-phase HPLC (mobile phase: water/acetonitrile) to give the title compound.

[0307]

Example 2: Synthesis of 4-[5-[(3S)-3-aminopyrrolidine-1-carbothioyl]-2-(p-tolyl)phenyl]benzonitrile

20 The 4-[5-[(3S)-3-aminopyrrolidine-1-carbonyl]-2-(p-tolyl)phenyl]benzonitrile (6 mg) obtained in Example 1 (step 5) was dissolved in THF (0.8 mL). At room temperature, Lawesson's reagent (3.8 mg) was added thereto, followed by stirring at room temperature for 30 minutes. Chloroform was added thereto, and the mixture was partitioned with sodium bicarbonate water. After the organic layer was dried over anhydrous sodium sulfate, the solvent was distilled off. The residue was purified by reversed-phase HPLC (mobile phase: water/acetonitrile) to give the title compound.

30 [0308]

Example 3: Synthesis of 4-[5-(4-aminopiperidine-1-carbonyl)-2-(p-tolyl)phenyl]benzonitrile

Step 1

35 The 3-(4-cyanophenyl)-4-(p-tolyl)benzoic acid (20 mg) obtained in Example 1 (step 3) was dissolved in THF (1 mL). At

room temperature, tert-butyl N-(4-piperidyl)carbamate (13 mg), HATU (49 mg), and TEA (0.027 mL) were added thereto, followed by stirring at 50°C overnight. The reaction mixture was vacuum-concentrated, and the solvent was distilled off. The residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate) to give tert-butyl N-[1-[3-(4-cyanophenyl)-4-(p-tolyl)benzoyl]-4-piperidyl]carbamate.

Step 2

The tert-butyl N-[1-[3-(4-cyanophenyl)-4-(p-tolyl)benzoyl]-4-piperidyl]carbamate (30 mg) obtained in step 1 above was dissolved in TFA (0.3 mL), and the progress of the reaction was confirmed by LCMS, followed by vacuum concentration. The residue was purified by reversed-phase HPLC (mobile phase: water/acetonitrile) to give the title compound.

[0309]

Example 4: Synthesis of 4-[5-(2,8-diazaspiro[3.5]nonane-2-carbonyl)-2-(p-tolyl)phenyl]benzonitrile

The procedure of steps 1 to 5 in Example 1 was conducted using tert-butyl 2,8-diazaspiro[3.5]nonane-8-carboxylate hydrochloride instead of tert-butyl N-[(3S)-pyrrolidin-3-yl]carbamate to give the title compound.

[0310]

Example 5: Synthesis of 4-[5-(2,7-diazaspiro[3.4]octane-7-carbonyl)-2-(p-tolyl)phenyl]benzonitrile

The procedure of steps 1 to 5 in Example 1 was conducted using tert-butyl 2,7-diazaspiro[3.4]octane-2-carboxylate instead of tert-butyl N-[(3S)-pyrrolidin-3-yl]carbamate to give the title compound.

[0311]

Example 6: Synthesis of 4-[5-(3,8-diazaspiro[4.4]nonane-8-carbonyl)-2-(p-tolyl)phenyl]benzonitrile

The procedure of steps 1 to 5 in Example 1 was conducted using tert-butyl 3,8-diazaspiro[4.4]nonane-8-carboxylate instead of tert-butyl N-[(3S)-pyrrolidin-3-yl]carbamate to give the title compound.

[0312]

Example 7: Synthesis of 4-[5-[(3-exo)-3-amino-8-azabicyclo[3.2.1]octane-8-carbonyl]-2-(p-tolyl)phenyl]benzonitrile

5 Step 1

3-Bromo-4-chloro-benzoic acid (500 mg) was dissolved in DMA (5.3 mL). At room temperature, HATU (1 g), TEA (0.59 mL), and tert-butyl N-[(3-exo)-8-azabicyclo[3.2.1]octan-3-yl]carbamate (480 mg) were added thereto, followed by stirring at room
10 temperature for 1 hour. Ethyl acetate was added thereto, and the resulting mixture was washed sequentially with water and saturated brine. After the organic layer was dried over anhydrous sodium sulfate, the solvent was distilled off. The residue was purified by silica gel column chromatography (mobile phase:
15 hexane/ethyl acetate) to give tert-butyl N-[(3-exo)-8-(3-bromo-4-chloro-benzoyl)-8-azabicyclo[3.2.1]octan-3-yl]carbamate.

[0313]

Step 2

The tert-butyl N-[(3-exo)-8-(3-bromo-4-chloro-benzoyl)-
20 8-azabicyclo[3.2.1]octan-3-yl]carbamate (200 mg) obtained in step 1 above was dissolved in 1,4-dioxane (2.3 mL). At room temperature, (4-cyanophenyl)boronic acid (60 mg), Pd(PPh₃)₄ (16 mg), and a 2 M Na₂CO₃ aqueous solution (1.1 mL) were added thereto, and the reaction mixture was stirred in a microwave
25 reactor at 120°C for 30 minutes. The solvent was distilled off, and the residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate) to give tert-butyl N-[(3-exo)-8-[4-chloro-3-(4-cyanophenyl)benzoyl]-8-azabicyclo[3.2.1]octan-3-yl]carbamate.

30 [0314]

Step 3

The tert-butyl N-[(3-exo)-8-[4-chloro-3-(4-cyanophenyl)benzoyl]-8-azabicyclo[3.2.1]octan-3-yl]carbamate (15 mg) obtained in step 2 above was dissolved in 1,4-dioxane (0.322
35 mL). At room temperature, p-tolylboronic acid (5.3 mg), Pd(dba)₂

(0.93 mg), a solution of 1 M PCy₃ in THF (0.003 mL), and tripotassium phosphate (21 mg) were added thereto, and the reaction mixture was stirred in a microwave reactor at 160°C for 30 minutes. The solvent was distilled off, and the residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate) to give tert-butyl N-[(3-exo)-8-[3-(4-cyanophenyl)-4-(p-tolyl)benzoyl]-8-azabicyclo[3.2.1]octan-3-yl]carbamate.

[0315]

10 Step 4

The tert-butyl N-[(3-exo)-8-[3-(4-cyanophenyl)-4-(p-tolyl)benzoyl]-8-azabicyclo[3.2.1]octan-3-yl]carbamate (15 mg) obtained in step 3 above was dissolved in TFA (0.3 mL), and the progress of the reaction was confirmed by LCMS, followed by vacuum concentration. The residue was purified by reversed-phase HPLC (mobile phase: water/acetonitrile) to give the title compound.

[0316]

20 Example 8: Synthesis of 4-[5-[(3S)-3-amino-3-methyl-pyrrolidine-1-carbonyl]-2-(p-tolyl)phenyl]benzonitrile

The procedure of steps 1 to 5 in Example 1 was conducted using tert-butyl N-[(3S)-3-methylpyrrolidin-3-yl]carbamate instead of tert-butyl N-[(3S)-pyrrolidin-3-yl]carbamate to give the title compound.

25 [0317]

Example 9: Synthesis of 4-[5-[(3S)-3-aminopyrrolidine-1-carbonyl]-2-(2-chloro-4-methyl-phenyl)phenyl]benzonitrile

Step 1

3-Bromo-4-chloro-benzoic acid (10 g) was dissolved in DMA (85 mL). At room temperature, HATU (24 g), TEA (12 mL), and tert-butyl N-[(3S)-pyrrolidin-3-yl]carbamate (8.7 g) were added thereto, followed by stirring at room temperature for 1 hour. Ethyl acetate was added thereto, and the resulting mixture was washed sequentially with water and saturated brine. After the organic layer was dried over anhydrous sodium sulfate, the

solvent was distilled off. The residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate) to give tert-butyl N-[(3S)-1-(3-bromo-4-chloro-benzoyl)pyrrolidin-3-yl]carbamate.

5 [0318]

Step 2

The tert-butyl N-[(3S)-1-(3-bromo-4-chloro-benzoyl)pyrrolidin-3-yl]carbamate (2.2 g) obtained in step 1 above was dissolved in 1,4-dioxane (13.6 mL). At room
10 temperature, 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzonitrile (1.5 g), Pd(PPh₃)₄ (189 mg), and a 2 M Na₂CO₃ aqueous solution (6.8 mL) were added thereto, and the reaction mixture was stirred in a microwave reactor at 120°C for 30 minutes. Ethyl acetate was added thereto, and the resulting
15 mixture was washed sequentially with water and saturated brine. After the organic layer was dried over anhydrous sodium sulfate, the solvent was distilled off. The residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate) to give tert-butyl N-[(3S)-1-[4-chloro-3-(4-
20 cyanophenyl)benzoyl]pyrrolidin-3-yl]carbamate.

[0319]

Step 3

The tert-butyl N-[(3S)-1-[4-chloro-3-(4-cyanophenyl)benzoyl]pyrrolidin-3-yl]carbamate (500 mg) obtained
25 in step 2 above was dissolved in 1,4-dioxane (9.8 mL). At room temperature, Pd(OAc)₂ (26 mg), KOAc (346 mg), bis(pinacolato)diboron (596 mg), and Silica-SMAP (150 mg, produced by Wako Pure Chemical Industries, Ltd.) were added thereto, followed by stirring at 160°C overnight. The mixture was
30 passed through Celite, and the filtrate was vacuum-concentrated. The residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate) to give tert-butyl N-[(3S)-1-[3-(4-cyanophenyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoyl]pyrrolidin-3-yl]carbamate.

35 [0320]

Step 4

The tert-butyl N-[(3S)-1-[3-(4-cyanophenyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoyl]pyrrolidin-3-yl]carbamate (15 mg) obtained in step 3 above, 1-bromo-2-chloro-4-methyl-benzene (12 mg), and Pd(PPh₃)₄ (1.7 mg) were suspended in 1,4-dioxane (1.5 mL). At room temperature, a 2 M Na₂CO₃ aqueous solution (0.7 mL) was added thereto, followed by stirring at 120°C for 30 minutes. After the reaction mixture was filtrated, the solvent was distilled off to give tert-butyl N-[(3S)-1-[4-(2-chloro-4-methyl-phenyl)-3-(4-cyanophenyl)benzoyl]pyrrolidin-3-yl]carbamate.

[0321]

Step 5

The tert-butyl N-[(3S)-1-[4-(2-chloro-4-methyl-phenyl)-3-(4-cyanophenyl)benzoyl]pyrrolidin-3-yl]carbamate (15 mg) obtained in step 4 above was dissolved in TFA (0.3 mL), and the progress of the reaction was confirmed by LCMS, followed by vacuum concentration. The residue was purified by reversed-phase HPLC (mobile phase: water/acetonitrile) to give the title compound.

[0322]

Example 10: Synthesis of 4-[5-[(3S)-3-aminopyrrolidine-1-carbonyl]-2-(3-chloro-4-methyl-phenyl)phenyl]benzonitrile

The procedure of steps 1 to 5 in Example 9 was conducted using 4-bromo-2-chloro-1-methyl-benzene instead of 1-bromo-2-chloro-4-methyl-benzene to give the title compound.

[0323]

Example 11: Synthesis of 4-[5-[(3S)-3-aminopyrrolidine-1-carbonyl]-2-[3-fluoro-4-

(trifluoromethyl)phenyl]phenyl]benzonitrile

The procedure of steps 1 to 5 in Example 9 was conducted using 4-bromo-2-fluoro-1-(trifluoromethyl)benzene instead of 1-bromo-2-chloro-4-methyl-benzene to give the title compound.

[0324]

Example 12: Synthesis of 4-[5-[(3S)-3-aminopyrrolidine-1-carbonyl]-2-(4-methyl-2-nitro-phenyl)phenyl]benzonitrile

Step 1

The tert-butyl N-[(3S)-1-[4-chloro-3-(4-cyanophenyl)benzoyl]pyrrolidin-3-yl]carbamate (15 mg) obtained in Example 9 (step 2), 4-methyl-2-nitrophenylboronic acid pinacol ester (18 mg), Pd(dba)₂ (1.6 mg), a solution of 1 M PCy₃ in THF (0.003 mL), and tripotassium phosphate (15 mg) were added thereto, and the reaction mixture was stirred in a microwave reactor at 160°C for 30 minutes. The solvent was distilled off, and the residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate) to give tert-butyl N-[(3S)-1-[3-(4-cyanophenyl)-4-(4-methyl-2-nitro-phenyl)benzoyl]pyrrolidin-3-yl]carbamate.

[0325]

Step 2

The tert-butyl N-[(3S)-1-[3-(4-cyanophenyl)-4-(4-methyl-2-nitro-phenyl)benzoyl]pyrrolidin-3-yl]carbamate (10 mg) obtained in step 1 above was dissolved in TFA (0.3 mL), and the progress of the reaction was confirmed by LCMS, followed by vacuum concentration. The residue was purified by reversed-phase HPLC (mobile phase: water/acetonitrile) to give the title compound.

[0326]

Example 13: Synthesis of 4-[5-[(3S)-3-aminopyrrolidine-1-carbonyl]-2-[4-(difluoromethyl)phenyl]phenyl]benzonitrile

The procedure of steps 1 to 5 in Example 9 was conducted using 1-bromo-4-(difluoromethyl)benzene instead of 1-bromo-2-chloro-4-methyl-benzene to give the title compound.

[0327]

Example 14: Synthesis of 4-[5-[(3S)-3-aminopyrrolidine-1-carbonyl]-2-[4-(trifluoromethyl)phenyl]phenyl]benzonitrile

The procedure of steps 1 to 2 in Example 12 was conducted using [4-(trifluoromethyl)phenyl]boronic acid instead of 4-methyl-2-nitrophenylboronic acid pinacol ester to give the

title compound.

[0328]

Example 15: Synthesis of 4-[5-[(3S)-3-aminopyrrolidine-1-carbonyl]-2-(2-fluoro-4-methyl-phenyl)phenyl]benzonitrile

5 Step 1

The tert-butyl N-[(3S)-1-[4-chloro-3-(4-cyanophenyl)benzoyl]pyrrolidin-3-yl]carbamate (1.7 g) obtained in Example 9 (step 2) was dissolved in 1,4-dioxane (20 mL). At room temperature, (2-fluoro-4-methyl-phenyl)boronic acid (980 mg),
10 Pd(dba)₂ (110 mg), a solution of 1 M PCy₃ in THF (0.4 mL), and tripotassium phosphate (2.5 g) were added thereto, and the reaction mixture was stirred in a microwave reactor at 160°C for 45 minutes. The mixture was purified by NH-silica gel and washed with methanol/ethyl acetate, and the solvent was distilled off to
15 give tert-butyl N-[(3S)-1-[3-(4-cyanophenyl)-4-(2-fluoro-4-methyl-phenyl)benzoyl]pyrrolidin-3-yl]carbamate.

[0329]

Step 2

The tert-butyl N-[(3S)-1-[3-(4-cyanophenyl)-4-(2-fluoro-4-methyl-phenyl)benzoyl]pyrrolidin-3-yl]carbamate (1.7 g)
20 obtained in step 1 above was dissolved in TFA (44 mL), followed by stirring for 10 minutes. The solvent was distilled off, and the residue was purified by reversed-phase HPLC (mobile phase: water/acetonitrile) to give the title compound.

25 [0330]

Example 16: Synthesis of 4-[5-[(3S)-3-aminopyrrolidine-1-carbonyl]-2-(p-tolyl)phenyl]-2-fluoro-benzonitrile

Step 1

The tert-butyl N-[(3S)-1-(3-bromo-4-chloro-benzoyl)pyrrolidin-3-yl]carbamate (14 g) obtained in Example 9
30 (step 1) was dissolved in 1,4-dioxane (87 mL). At room temperature, (4-cyano-3-fluoro-phenyl)boronic acid (6.3 g), Pd(PPh₃)₄ (1.2 g), and a 2 M Na₂CO₃ aqueous solution (44 mL) were added thereto, followed by stirring at 90°C overnight. Ethyl
35 acetate was added thereto, and the resulting mixture was washed

sequentially with water and saturated brine. After the organic layer was dried over anhydrous sodium sulfate, the solvent was distilled off. The residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate) to give tert-butyl N-[(3S)-1-[4-chloro-3-(4-cyano-3-fluoro-phenyl)benzoyl]pyrrolidin-3-yl]carbamate.

[0331]

Step 2

The tert-butyl N-[(3S)-1-[4-chloro-3-(4-cyano-3-fluoro-phenyl)benzoyl]pyrrolidin-3-yl]carbamate (48 mg) obtained in step 1 above was dissolved in 1,4-dioxane (0.5 mL). At room temperature, p-tolylboronic acid (29 mg), Pd(dba)₂ (3.1 mg), a solution of 1 M PCy₃ in THF (0.005 mL), and tripotassium phosphate (68 mg) were added thereto, and the reaction mixture was stirred in a microwave reactor at 160°C for 45 minutes. The mixture was purified by NH-silica gel and washed with methanol/ethyl acetate, and the solvent was distilled off to give tert-butyl N-[(3S)-1-[3-(4-cyano-3-fluoro-phenyl)-4-(p-tolyl)benzoyl]pyrrolidin-3-yl]carbamate.

[0332]

Step 3

TFA (1.2 mL) was added to the tert-butyl N-[(3S)-1-[3-(4-cyano-3-fluoro-phenyl)-4-(p-tolyl)benzoyl]pyrrolidin-3-yl]carbamate (48 mg) obtained in step 2 above, followed by stirring for 10 minutes. The solvent was distilled off, and the residue was purified by reversed-phase HPLC (mobile phase: water/acetonitrile) to give the title compound.

[0333]

Example 17: Synthesis of 4-[5-[(3-endo)-3-amino-8-azabicyclo[3.2.1]octane-8-carbonyl]-2-(p-tolyl)phenyl]-2-fluorobenzonitrile

Step 1

3-Bromo-4-chloro-benzoic acid (700 mg) was dissolved in THF (15 mL). At room temperature, HATU (1.2 g), TEA (0.83 mL), and tert-butyl N-[(3-endo)-8-azabicyclo[3.2.1]octan-3-

yl]carbamate (700 mg) were added thereto, followed by stirring at 50°C for 1 hour. The solvent was distilled off, and the residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate) to give tert-butyl N-[(3-endo)-8-(3-bromo-4-chloro-benzoyl)-8-azabicyclo[3.2.1]octan-3-yl]carbamate.

[0334]

Step 2

The tert-butyl N-[(3-endo)-8-(3-bromo-4-chloro-benzoyl)-8-azabicyclo[3.2.1]octan-3-yl]carbamate (1.2 g) obtained in step 1 above was dissolved in 1,4-dioxane (6.7 mL). At room temperature, (4-cyano-3-fluoro-phenyl)boronic acid (461 mg), PdCl₂(dppf) (58 mg), and a 2 M Na₂CO₃ aqueous solution (3.3 mL) were added thereto, followed by stirring at 95°C overnight. Ethyl acetate was added thereto, and the resulting mixture was washed sequentially with water and saturated brine. After the organic layer was dried over anhydrous sodium sulfate, the solvent was distilled off. The residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate) to give tert-butyl N-[(3-endo)-8-[4-chloro-3-(4-cyano-3-fluoro-phenyl)benzoyl]-8-azabicyclo[3.2.1]octan-3-yl]carbamate.

[0335]

Step 3

The tert-butyl N-[(3-endo)-8-[4-chloro-3-(4-cyano-3-fluoro-phenyl)benzoyl]-8-azabicyclo[3.2.1]octan-3-yl]carbamate (17 mg) obtained in step 2 above was dissolved in 1,4-dioxane (0.5 mL). At room temperature, p-tolylboronic acid (9.6 mg), Pd(dba)₂ (1.6 mg), tripotassium phosphate (15 mg), and a solution of 1 M PCy₃ in THF (0.004 mL) were added thereto, and the mixture was stirred in a microwave reactor at 160°C for 30 minutes. The reaction mixture was filtered through NH-silica gel, and the solvent of the filtrate was distilled off to give tert-butyl N-[(3-endo)-8-[3-(4-cyano-3-fluoro-phenyl)-4-(p-tolyl)benzoyl]-8-azabicyclo[3.2.1]octan-3-yl]carbamate.

[0336]

Step 4

The tert-butyl N-[(3-endo)-8-[3-(4-cyano-3-fluoro-phenyl)-4-(p-tolyl)benzoyl]-8-azabicyclo[3.2.1]octan-3-yl]carbamate (15 mg) obtained in step 3 above was dissolved in TFA (0.3 mL), and the progress of the reaction was confirmed by LCMS, followed by vacuum concentration. The residue was purified by reversed-phase HPLC (mobile phase: water/acetonitrile) to give the title compound.

[0337]

Example 18: Synthesis of 4-[5-[(3-exo)-3-amino-8-azabicyclo[3.2.1]octane-8-carbonyl]-2-(1-methylindol-5-yl)phenyl]-2-fluoro-benzonitrile

Step 1

The tert-butyl N-[(3-exo)-8-(3-bromo-4-chloro-benzoyl)-8-azabicyclo[3.2.1]octan-3-yl]carbamate (300 mg) obtained in Example 7 (step 1) was dissolved in 1,4-dioxane (1.7 mL). At room temperature, (4-cyano-3-fluoro-phenyl)boronic acid (123 mg), PdCl₂(dppf) (17 mg), and a 2 M Na₂CO₃ aqueous solution (0.85 mL) were added thereto, and the reaction mixture was stirred in a microwave reactor at 120°C for 30 minutes. Ethyl acetate was added thereto, and the resulting mixture was washed sequentially with water and saturated brine. After the organic layer was dried over anhydrous sodium sulfate, the solvent was distilled off. The residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate) to give tert-butyl N-[(3-exo)-8-[4-chloro-3-(4-cyano-3-fluoro-phenyl)benzoyl]-8-azabicyclo[3.2.1]octan-3-yl]carbamate.

[0338]

Step 2

The tert-butyl N-[(3-exo)-8-[4-chloro-3-(4-cyano-3-fluoro-phenyl)benzoyl]-8-azabicyclo[3.2.1]octan-3-yl]carbamate (10 mg) obtained in step 1 above was dissolved in 1,4-dioxane (0.5 mL). At room temperature, (1-methylindol-5-yl)boronic acid (7.2 mg), Pd(dba)₂ (0.9 mg), tripotassium phosphate (8.8 mg), and a solution of 1 M PCy₃ in THF (0.002 mL) were added thereto, and the reaction mixture was stirred in a microwave reactor at 160°C

for 30 minutes. The reaction mixture was filtered through NH-silica gel, and the solvent of the filtrate was distilled off. The residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate) to give tert-butyl N-[(3-exo)-8-[3-(4-cyano-3-fluoro-phenyl)-4-(1-methylindol-5-yl)benzoyl]-8-azabicyclo[3.2.1]octan-3-yl]carbamate.

[0339]

Step 3

The tert-butyl N-[(3-exo)-8-[3-(4-cyano-3-fluoro-phenyl)-4-(1-methylindol-5-yl)benzoyl]-8-azabicyclo[3.2.1]octan-3-yl]carbamate (15 mg) obtained in step 2 above was dissolved in TFA (0.3 mL), and the progress of the reaction was confirmed with LCMS, followed by vacuum concentration. The residue was purified by reversed-phase HPLC (mobile phase: water/acetonitrile) to give the title compound.

[0340]

Example 19: Synthesis of 4-[5-[(3S)-3-aminopyrrolidine-1-carbonyl]-2-(p-tolyl)phenyl]-2,6-difluoro-benzonitrile

Step 1

4-Bromo-3-chloro-benzoic acid (2 g) was dissolved in DMA (17 mL). At room temperature, HATU (4.8 g), TEA (2.4 mL), and tert-butyl N-[(3S)-pyrrolidin-3-yl]carbamate (1.7 g) were added thereto, followed by stirring at room temperature for 1 hour. Ethyl acetate was added thereto, and the resulting mixture was washed sequentially with water and saturated brine. After the organic layer was dried over anhydrous sodium sulfate, the solvent was distilled off. The residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate) to give tert-butyl N-[(3S)-1-(4-bromo-3-chloro-benzoyl)pyrrolidin-3-yl]carbamate.

[0341]

Step 2

The tert-butyl N-[(3S)-1-(4-bromo-3-chloro-benzoyl)pyrrolidin-3-yl]carbamate obtained in step 1 above was dissolved in 1,4-dioxane (10.6 mL). At room temperature, Pd(PPh₃)₄

(147 mg), a 2 M Na₂CO₃ aqueous solution (5.3 mL), and p-tolylboronic acid (693 mg) were added thereto, and the reaction mixture was stirred in a microwave reactor at 120°C for 30 minutes. Ethyl acetate was added thereto, and the resulting mixture was washed sequentially with water and saturated brine. After the organic layer was dried over anhydrous sodium sulfate, the solvent was distilled off. The residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate) to give tert-butyl N-[(3S)-1-[3-chloro-4-(p-tolyl)benzoyl]pyrrolidin-3-yl]carbamate. [0342]

Step 3

The tert-butyl N-[(3S)-1-[3-chloro-4-(p-tolyl)benzoyl]pyrrolidin-3-yl]carbamate (666 mg) obtained in step 2 above was dissolved in 1,4-dioxane (16 mL). At room temperature, Pd(OAc)₂ (36 mg), KOAc (473 mg), bis(pinacolato)diboron (815 mg), and a solution of 1 M PCy₃ in THF (0.24 mL) were added thereto. After degassing and nitrogen substitution, the mixture was stirred at 80°C overnight. The reaction mixture was passed through Celite, and the solvent of the filtrate was distilled off. The residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate) to give tert-butyl N-[(3S)-1-[4-(p-tolyl)-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoyl]pyrrolidin-3-yl]carbamate. [0343]

Step 4

The tert-butyl N-[(3S)-1-[4-(p-tolyl)-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoyl]pyrrolidin-3-yl]carbamate (15 mg) obtained in step 3 above, 4-bromo-2,6-difluoro-benzonitrile (12.9 mg), and Pd(PPh₃)₄ (1.7 mg) were suspended in 1,4-dioxane (1.5 mL). At room temperature, a 2 M Na₂CO₃ aqueous solution (0.7 mL) was added thereto, and the reaction mixture was stirred in a microwave reactor at 120°C for 30 minutes. The reaction mixture was filtrated, and the solvent

was distilled off to give tert-butyl N-[(3S)-1-[3-(4-cyano-3,5-difluoro-phenyl)-4-(p-tolyl)benzoyl]pyrrolidin-3-yl]carbamate. [0344]

Step 5

5 The tert-butyl N-[(3S)-1-[3-(4-cyano-3,5-difluoro-phenyl)-4-(p-tolyl)benzoyl]pyrrolidin-3-yl]carbamate (15 mg) obtained in step 4 above was dissolved in TFA (0.3 mL), and the progress of the reaction was confirmed with LCMS, followed by vacuum concentration. The residue was purified by reversed-phase
10 HPLC (mobile phase: water/acetonitrile) to give the title compound.
[0345]

Example 20: Synthesis of 4-[5-[(3S)-3-aminopyrrolidine-1-carbonyl]-2-[2-fluoro-4-(2-methoxyethyl)phenyl]phenyl]-2-fluoro-
15 benzonitrile

Step 1

1-Bromo-2-fluoro-4-(2-methoxyethyl)benzene (4.5 g) was suspended in 1,4-dioxane (48 mL), followed by stirring. Then, bis(pinacolato)diboron (7.4 g), KOAc (3.8 g), and PdCl₂(dppf)
20 (0.71 g) were added thereto, followed by stirring at 90°C overnight. Ethyl acetate was added thereto, the mixture was passed through Celite, and the filtrate was washed sequentially with water and saturated brine. After the organic layer was dried over anhydrous sodium sulfate, the solvent was distilled off. The
25 residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate) to give 2-[2-fluoro-4-(2-methoxyethyl)phenyl]-4,4,5,5-tetramethyl-1,3,2-dioxaborolane.
[0346]

Step 2

30 The tert-butyl N-[(3S)-1-[4-chloro-3-(4-cyano-3-fluoro-phenyl)benzoyl]pyrrolidin-3-yl]carbamate (150 mg) obtained in Example 16 (step 1), the 2-[2-fluoro-4-(2-methoxyethyl)phenyl]-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (189 mg) obtained in step 1 above, Pd(dba)₂ (15 mg), tripotassium phosphate (144 mg), and a
35 solution of 1 M PCy₃ in THF (0.034 mL) were dissolved in 1,4-

dioxane (3.8 mL). The reaction mixture was stirred in a microwave reactor at 160°C for 45 minutes. The reaction mixture was filtered through NH-silica gel, and the solvent of the filtrate was distilled off. The residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate) to give tert-butyl N-[(3S)-1-[3-(4-cyanophenyl)-4-[2-fluoro-4-(2-methoxyethyl)phenyl]benzoyl]pyrrolidin-3-yl]carbamate.

[0347]

Step 3

The tert-butyl N-[(3S)-1-[3-(4-cyanophenyl)-4-[2-fluoro-4-(2-methoxyethyl)phenyl]benzoyl]pyrrolidin-3-yl]carbamate (150 mg) obtained in step 2 above was dissolved in TFA (10 mL), and the progress of the reaction was confirmed with LCMS, followed by vacuum concentration. The residue was purified by reversed-phase HPLC (mobile phase: water/acetonitrile) to give the title compound.

[0348]

Example 21: Synthesis of 4-[5-[(3-endo)-3-amino-8-azabicyclo[3.2.1]octane-8-carbonyl]-2-(2-fluoro-4-methyl-phenyl)phenyl]-2-fluoro-benzonitrile

The procedure of steps 1 to 4 in Example 17 was conducted using (2-fluoro-4-methyl-phenyl)boronic acid instead of p-tolylboronic acid to give the title compound.

[0349]

Example 22: Synthesis of 4-[5-[(3-endo)-3-amino-8-azabicyclo[3.2.1]octane-8-carbonyl]-2-[2-fluoro-4-(2-methoxyethyl)phenyl]phenyl]-2-fluoro-benzonitrile

Step 1

The tert-butyl 3-bromo-4-chloro-benzoate (1.00 g) obtained in Example 1 (step 1) was dissolved in 1,4-dioxane (8.6 mL). At room temperature, (4-cyano-3-fluoro-phenyl)boronic acid (509 mg), Pd(PPh₃)₄ (119 mg), and a 2 M Na₂CO₃ aqueous solution (4.3 mL) were added thereto, and the reaction mixture was stirred in a microwave reactor at 120°C for 30 minutes. The reaction mixture was filtrated, and the solvent was distilled off. The

residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate) to give tert-butyl 4-chloro-3-(4-cyano-3-fluoro-phenyl)benzoate.

[0350]

5 Step 2

The tert-butyl 4-chloro-3-(4-cyano-3-fluoro-phenyl)benzoate (1.00 g) obtained in step 1 above was dissolved in 1,4-dioxane (15 mL). At room temperature, the 2-[2-fluoro-4-(2-methoxyethyl)phenyl]-4,4,5,5-tetramethyl-1,3,2-dioxaborolane
10 (1.69 g) obtained in Example 20 (step 1), Pd(dba)₂ (138 mg), tripotassium phosphate (1.28 g), and a solution of 1 M PCy₃ in THF (0.30 mL) were added thereto, and the reaction mixture was stirred in a microwave reactor at 160°C for 30 minutes. After the addition of chloroform, the insoluble matter was filtered off,
15 and the solvent was distilled off. The residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate), and the solvent was distilled off. The residue was dissolved in TFA (2 mL), and the solvent was distilled off. Ethyl acetate was added thereto, and the mixture was washed
20 sequentially with water and saturated brine. After the organic layer was dried over anhydrous sodium sulfate, the solvent was distilled off to give 3-(4-cyano-3-fluoro-phenyl)-4-[2-fluoro-4-(2-methoxyethyl)phenyl]benzoic acid.

[0351]

25 Step 3

The 3-(4-cyano-3-fluoro-phenyl)-4-[2-fluoro-4-(2-methoxyethyl)phenyl]benzoic acid (10 mg) obtained in step 2 above, tert-butyl N-[(3-endo)-8-azabicyclo[3.2.1]octan-3-yl]carbamate (5.8 mg), and HATU (19 mg) were dissolved in THF
30 (0.5 mL). At room temperature, TEA (0.007 mL) was added thereto, followed by stirring at 50°C overnight. The reaction mixture was vacuum-concentrated, and the solvent was distilled off. The residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate) to give tert-butyl N-[(3-endo)-8-[3-(4-cyano-3-fluoro-phenyl)-4-[2-fluoro-4-(2-

methoxyethyl)phenyl]benzoyl]-8-azabicyclo[3.2.1]octan-3-yl]carbamate.

[0352]

Step 4

5 The tert-butyl N-[(3-endo)-8-[3-(4-cyano-3-fluoro-phenyl)-4-[2-fluoro-4-(2-methoxyethyl)phenyl]benzoyl]-8-azabicyclo[3.2.1]octan-3-yl]carbamate (10.9 mg) obtained in step 3 above was dissolved in TFA (0.3 mL), and the progress of the reaction was confirmed with LCMS, followed by vacuum
10 concentration. The residue was purified by reversed-phase HPLC (mobile phase: water/acetonitrile) to give the title compound.
[0353]

Example 23: Synthesis of 4-[5-[(3-exo)-3-amino-8-azabicyclo[3.2.1]octane-8-carbonyl]-2-(6-fluoro-1-methyl-indol-5-yl)phenyl]-2-fluoro-benzonitrile
15

Step 1

 The tert-butyl N-[(3-exo)-8-[4-chloro-3-(4-cyano-3-fluoro-phenyl)benzoyl]-8-azabicyclo[3.2.1]octan-3-yl]carbamate (270 mg) obtained in Example 18 (step 1) was dissolved in 1,4-
20 dioxane (2.8 mL). At room temperature, Pd(OAc)₂ (2.5 mg), KOAc (164 mg), bis(pinacolato)diboron (283 mg), and Silica-SMAP (4.6 mg) were added thereto, followed by stirring at 150°C overnight. The mixture was passed through Celite, and the filtrate was vacuum-concentrated. The residue was purified by silica gel
25 column chromatography (mobile phase: hexane/ethyl acetate) to give tert-butyl N-[(3-exo)-8-[3-(4-cyano-3-fluoro-phenyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoyl]-8-azabicyclo[3.2.1]octan-3-yl]carbamate.
[0354]

30 Step 2

 The tert-butyl N-[(3-exo)-8-[3-(4-cyano-3-fluoro-phenyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoyl]-8-azabicyclo[3.2.1]octan-3-yl]carbamate (10 mg) obtained in step 1 above, 5-bromo-6-fluoro-1-methyl-indole (4.8 mg), and
35 PdCl₂(dppf) (0.71 mg) were suspended in 1,4-dioxane (0.5 mL). At

room temperature, tripotassium phosphate (11 mg) was added thereto, followed by stirring at 125°C for 45 minutes. After the reaction mixture was filtrated, the solvent was distilled off to give tert-butyl N-[(3-exo)-8-[3-(4-cyano-3-fluoro-phenyl)-4-(6-fluoro-1-methyl-indol-5-yl)benzoyl]-8-azabicyclo[3.2.1]octan-3-yl]carbamate.

[0355]

Step 3

The tert-butyl N-[(3-exo)-8-[3-(4-cyano-3-fluoro-phenyl)-4-(6-fluoro-1-methyl-indol-5-yl)benzoyl]-8-azabicyclo[3.2.1]octan-3-yl]carbamate (8 mg) obtained in step 2 above was dissolved in TFA (0.3 mL), and the progress of the reaction was confirmed with LCMS, followed by vacuum concentration. The residue was purified by reversed-phase HPLC (mobile phase: water/acetonitrile) to give the title compound.

[0356]

Example 24: Synthesis of 4-[5-[(3-exo)-3-amino-8-azabicyclo[3.2.1]octane-8-carbonyl]-2-(6-fluoro-1-methyl-indazol-5-yl)phenyl]-2-fluoro-benzonitrile

Step 1

The tert-butyl N-[(3-exo)-8-[3-(4-cyano-3-fluoro-phenyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoyl]-8-azabicyclo[3.2.1]octan-3-yl]carbamate (10 mg) obtained in Example 23 (step 1) was dissolved in 1,4-dioxane (0.5 mL). At room temperature, 5-bromo-6-fluoro-1-methyl-indazole (4.8 mg), PdCl₂(dppf) (0.71 mg), and tripotassium phosphate (11 mg) were added thereto, and the mixture was stirred in a microwave reactor at 125°C for 45 minutes. The reaction mixture was filtrated, and the solvent was distilled off to give tert-butyl N-[(3-exo)-8-[3-(4-cyano-3-fluoro-phenyl)-4-(6-fluoro-1-methyl-indazol-5-yl)benzoyl]-8-azabicyclo[3.2.1]octan-3-yl]carbamate.

[0357]

Step 2

The tert-butyl N-[(3-exo)-8-[3-(4-cyano-3-fluoro-phenyl)-4-(6-fluoro-1-methyl-indazol-5-yl)benzoyl]-8-

azabicyclo[3.2.1]octan-3-yl]carbamate (15 mg) obtained in step 1 above was dissolved in TFA (0.3 mL), and the progress of the reaction was confirmed with LCMS, followed by vacuum concentration. The residue was purified by reversed-phase HPLC (mobile phase: water/acetonitrile) to give the title compound. [0358]

Example 25: Synthesis of 4-[5-[(3S)-3-amino-3-methyl-pyrrolidine-1-carbonyl]-2-[2-fluoro-4-(2-methoxyethyl)phenyl]phenyl]-2-fluoro-benzonitrile

Step 1

The 3-(4-cyano-3-fluoro-phenyl)-4-[2-fluoro-4-(2-methoxyethyl)phenyl]benzoic acid (10 mg) obtained in Example 22 (step 2) and tert-butyl N-[(3S)-3-methylpyrrolidin-3-yl]carbamate (5.1 mg) were dissolved in THF (0.5 mL). At room temperature, TEA (0.011 mL) and HATU (19 mg) were added thereto, followed by stirring at 50°C overnight. The reaction mixture was vacuum-concentrated, and the solvent was distilled off to give tert-butyl N-[(3S)-1-[3-(4-cyano-3-fluoro-phenyl)-4-[2-fluoro-4-(2-methoxyethyl)phenyl]benzoyl]-3-methyl-pyrrolidin-3-yl]carbamate. [0359]

Step 2

The tert-butyl N-[(3S)-1-[3-(4-cyano-3-fluoro-phenyl)-4-[2-fluoro-4-(2-methoxyethyl)phenyl]benzoyl]-3-methyl-pyrrolidin-3-yl]carbamate (15 mg) obtained in step 1 above was dissolved in TFA (0.3 mL), and the progress of the reaction was confirmed with LCMS, followed by vacuum concentration. The residue was purified by reversed-phase HPLC (mobile phase: water/acetonitrile) to give the title compound. [0360]

Example 26: Synthesis of 4-[5-[(3-exo)-3-amino-8-azabicyclo[3.2.1]octane-8-carbonyl]-2-[2-fluoro-4-(2-methoxyethyl)phenyl]phenyl]-2-fluoro-benzonitrile

Step 1

The 3-(4-cyano-3-fluoro-phenyl)-4-[2-fluoro-4-(2-methoxyethyl)phenyl]benzoic acid (10 mg) obtained in Example 22

(step 2), and tert-butyl N-[(3-exo)-8-azabicyclo[3.2.1]octan-3-yl]carbamate (5.8 mg) were dissolved in THF (0.5 mL). At room temperature, TEA (0.011 mL) and HATU (19 mg) were added thereto, followed by stirring at 50°C overnight. The reaction mixture was vacuum-concentrated, and the solvent was distilled off to give tert-butyl N-[(3-exo)-8-[3-(4-cyano-3-fluoro-phenyl)-4-[2-fluoro-4-(2-methoxyethyl)phenyl]benzoyl]-8-azabicyclo[3.2.1]octan-3-yl]carbamate.

[0361]

10 Step 2

The tert-butyl N-[(3-exo)-8-[3-(4-cyano-3-fluoro-phenyl)-4-[2-fluoro-4-(2-methoxyethyl)phenyl]benzoyl]-8-azabicyclo[3.2.1]octan-3-yl]carbamate (15 mg) obtained in step 1 above was dissolved in TFA (0.3 mL), and the progress of the reaction was confirmed with LCMS, followed by vacuum concentration. The residue was purified by reversed-phase HPLC (mobile phase: water/acetonitrile) to give the title compound.

[0362]

20 Example 27: Synthesis of 4-[5-(3,8-diazabicyclo[3.2.1]octane-8-carbonyl)-2-[2-fluoro-4-(2-methoxyethyl)phenyl]phenyl]-2-fluoro-benzonitrile

The procedure of steps 1 to 4 in Example 22 was conducted using tert-butyl 3,8-diazabicyclo[3.2.1]octane-3-carboxylic acid instead of tert-butyl N-[(3-endo)-8-azabicyclo[3.2.1]octan-3-yl]carbamate to give the title compound.

[0363]

Example 28: Synthesis of 4-[5-[(3-exo)-3-amino-8-azabicyclo[3.2.1]octane-8-carbonyl]-2-[2-fluoro-4-(2-hydroxy-2-methyl-propyl)phenyl]phenyl]-2-fluoro-benzonitrile

30 Step 1

Methyl 2-(4-bromo-3-fluoro-phenyl)acetate (500 mg) was dissolved in THF (2.2 mL). At -30°C, a solution of 3 M MeMgBr in ether (5.40 mL) was added thereto dropwise, followed by stirring at room temperature overnight. The reaction mixture was introduced into an aqueous ammonium chloride solution, ethyl

acetate was added thereto, and the mixture was washed sequentially with water and saturated brine. After the organic layer was dried over anhydrous sodium sulfate, the solvent was distilled off. The residue was purified by silica gel column chromatography (mobile phase: ethyl acetate/hexane = 10% → 50%) to give 1-(4-bromo-3-fluoro-phenyl)-2-methyl-propan-2-ol. [0364]

Step 2

The tert-butyl N-[(3-exo)-8-[3-(4-cyano-3-fluoro-phenyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoyl]-8-azabicyclo[3.2.1]octan-3-yl]carbamate (68 mg) obtained in Example 23 (step 1), the 1-(4-bromo-3-fluoro-phenyl)-2-methyl-propan-2-ol (107 mg) obtained in step 1 above, and Pd(PPh₃)₄ (6.42 mg) were suspended in 1,4-dioxane (0.93 mL). At room temperature, a 2 M Na₂CO₃ aqueous solution (0.46 mL) was added thereto, followed by stirring at 125°C for 45 minutes. After the reaction mixture was filtrated, the solvent was distilled off to give tert-butyl N-[(3-exo)-8-[3-(4-cyano-3-fluoro-phenyl)-4-[2-fluoro-4-(2-hydroxy-2-methyl-propyl)phenyl]benzoyl]-8-azabicyclo[3.2.1]octan-3-yl]carbamate. [0365]

Step 3

The tert-butyl N-[(3-exo)-8-[3-(4-cyano-3-fluoro-phenyl)-4-[2-fluoro-4-(2-hydroxy-2-methyl-propyl)phenyl]benzoyl]-8-azabicyclo[3.2.1]octan-3-yl]carbamate (90 mg) obtained in step 2 above was dissolved in TFA (0.3 mL), and the progress of the reaction was confirmed with LCMS, followed by vacuum concentration. The residue was purified by reversed-phase HPLC (mobile phase: water/acetonitrile) to give the title compound. [0366]

Example 29: Synthesis of 4-[5-[(3S)-3-aminopyrrolidine-1-carbonyl]-2-[4-(hydroxymethyl)phenyl]phenyl]benzonitrile

Step 1

[4-(4,4,5,5-Tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl]methanol (500 mg) and DMAP (26 mg) were dissolved in

THF (7.1 mL), followed by the addition of TEA (0.74 mL). At room temperature, acetylchloride (0.23 mL) was added thereto, followed by stirring for 1 hour. Ethyl acetate was added thereto, and the resulting mixture was washed sequentially with water and

5 saturated brine. After the organic layer was dried over anhydrous sodium sulfate, the solvent was distilled off. The residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate) to give [4-(4,4,5,5-tetramethyl-1,3,2-

10 [0367]

Step 2

The tert-butyl N-[(3S)-1-[4-chloro-3-(4-cyanophenyl)benzoyl]pyrrolidin-3-yl]carbamate (100 mg) obtained in Example 9 (step 2) and the [4-(4,4,5,5-tetramethyl-1,3,2-

15 dioxaborolan-2-yl)phenyl]methyl acetate (130 mg) obtained in step 1 above were dissolved in 1,4-dioxane (1.2 mL). At room temperature, Pd(dba)₂ (6.8 mg), tripotassium phosphate (100 mg), and a solution of 1 M PCy₃ in THF (0.02 mL) were added thereto, and the mixture was stirred in a microwave reactor at 160°C for 1

20 hour. The reaction mixture was filtered through NH-silica gel, and the solvent of the filtrate was distilled off. The residue was purified by silica gel column chromatography (mobile phase: ethyl acetate/hexane = 30% → 100%) to give [4-[4-[(3S)-3-(tert-butoxycarbonylamino)pyrrolidine-1-carbonyl]-2-(4-

25 cyanophenyl)phenyl]phenyl]methyl acetate.

[0368]

Step 3

The [4-[4-[(3S)-3-(tert-butoxycarbonylamino)pyrrolidine-1-carbonyl]-2-(4-

30 cyanophenyl)phenyl]phenyl]methyl acetate (100 mg) obtained in step 2 above was dissolved in MeOH (2 mL). At room temperature, K₂CO₃ (65 mg) was added thereto, followed by stirring at room temperature for 30 minutes. Chloroform was added thereto, the mixture was washed sequentially with a saturated aqueous ammonium

35 chloride solution and saturated brine, and dried over anhydrous

sodium sulfate, and the solvent was distilled off. The residue was purified by silica gel column chromatography (mobile phase: ethyl acetate/hexane = 40% → 100%) to give tert-butyl N-[(3S)-1-[3-(4-cyanophenyl)-4-[4-(hydroxymethyl)phenyl]benzoyl]pyrrolidin-3-yl]carbamate.

[0369]

Step 4

The tert-butyl N-[(3S)-1-[3-(4-cyanophenyl)-4-[4-(hydroxymethyl)phenyl]benzoyl]pyrrolidin-3-yl]carbamate (15 mg) obtained in step 3 above was dissolved in TFA (0.3 mL), and the progress of the reaction was confirmed with LCMS, followed by vacuum concentration. The residue was purified by reversed-phase HPLC (mobile phase: water/acetonitrile) to give the title compound.

[0370]

Example 30: Synthesis of 4-[5-[(3S)-3-aminopyrrolidine-1-carbonyl]-2-[4-(2-methoxyethyl)phenyl]phenyl]benzonitrile

Step 1

1-Bromo-4-(2-methoxyethyl)benzene (450 mg) was dissolved in 1,4-dioxane (5.2 mL). Then, bis(pinacolato)diboron (797 mg), KOAc (411 mg), and PdCl₂(dppf) (77 mg) were added thereto, followed by stirring at 90°C overnight. Ethyl acetate was added thereto, the mixture was passed through Celite, and the filtrate was washed sequentially with water and saturated brine. After the organic layer was dried over anhydrous sodium sulfate, the solvent was distilled off. The residue was purified by silica gel column chromatography (mobile phase: ethyl acetate/hexane = 2% → 20%) to give 2-[4-(2-methoxyethyl)phenyl]-4,4,5,5-tetramethyl-1,3,2-dioxaborolane.

[0371]

Step 2

The tert-butyl N-[(3S)-1-[4-chloro-3-(4-cyanophenyl)benzoyl]pyrrolidin-3-yl]carbamate (300 mg) obtained in Example 9 (step 2) and the 2-[4-(2-methoxyethyl)phenyl]-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (369 mg) obtained in step

1 above were dissolved in 1,4-dioxane (2 mL). At room temperature, Pd(dba)₂ (32 mg), tripotassium phosphate (300 mg), and a solution of 1 M PCy₃ in THF (0.07 mL) were added thereto, and the mixture was stirred in a microwave reactor at 160°C for

5 45 minutes. The reaction mixture was passed through Celite, and the solvent of the filtrate was distilled off. The residue was purified by reversed-phase HPLC (mobile phase: water/acetonitrile) to give tert-butyl N-[(3S)-1-[3-(4-cyanophenyl)-4-[4-(2-methoxyethyl)phenyl]benzoyl]pyrrolidin-3-yl]carbamate.

10 [0372]

Step 3

The tert-butyl N-[(3S)-1-[3-(4-cyanophenyl)-4-[4-(2-methoxyethyl)phenyl]benzoyl]pyrrolidin-3-yl]carbamate (15 mg) obtained in step 2 above was dissolved in TFA (0.3 mL), and the progress of the reaction was confirmed with LCMS, followed by vacuum concentration. The residue was purified by reversed-phase HPLC (mobile phase: water/acetonitrile) to give the title compound.

20 [0373]

Example 31: Synthesis of 4-[5-[(3S)-3-aminopyrrolidine-1-carbonyl]-2-[4-(2-hydroxyethyl)phenyl]phenyl]benzonitrile

The procedure of steps 1 to 4 in Example 29 was conducted using 2-[4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl]ethanol instead of [4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl]methanol to give the title compound.

25 [0374]

Example 32: Synthesis of 4-[5-[(3S)-3-aminopyrrolidine-1-carbonyl]-2-[4-(3-hydroxypropyl)phenyl]phenyl]benzonitrile

The procedure of steps 1 to 4 in Example 29 was conducted using 3-[4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl]propan-1-ol instead of [4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl]methanol to give the title compound.

30 [0375]

35 Example 33: Synthesis of 4-[5-[(3S)-3-aminopyrrolidine-1-

carbonyl]-2-[4-[1-(hydroxymethyl)cyclopropyl]phenyl]phenyl]benzonitrile

The procedure of steps 1 to 5 in Example 9 was conducted using 1-(4-bromophenyl)cyclopropylmethanol instead of 1-bromo-2-chloro-4-methyl-benzene to give the title compound.
[0376]

Example 34: Synthesis of 4-[5-[(3S)-3-aminopyrrolidine-1-carbonyl]-2-[4-(2-hydroxy-2-methylpropyl)phenyl]phenyl]benzonitrile

The procedure of steps 1 to 5 in Example 9 was conducted using 1-(4-bromophenyl)-2-methylpropan-2-ol instead of 1-bromo-2-chloro-4-methyl-benzene to give the title compound.
[0377]

Example 35: Synthesis of 4-[5-[(3S)-3-aminopyrrolidine-1-carbonyl]-2-[4-(2-hydroxypropoxy)phenyl]phenyl]benzonitrile
Step 1

The procedure of step 1 in Example 12 was conducted using (4-benzyloxyphenyl)boronic acid instead of 4-methyl-2-nitrophenylboronic acid pinacol ester to give tert-butyl N-[(3S)-1-[4-(4-benzyloxyphenyl)-3-(4-cyanophenyl)benzoyl]pyrrolidin-3-yl]carbamate.
[0378]

Step 2

The tert-butyl N-[(3S)-1-[4-(4-benzyloxyphenyl)-3-(4-cyanophenyl)benzoyl]pyrrolidin-3-yl]carbamate (800 mg) obtained in step 1 above and palladium hydroxide-carbon (160 mg) were suspended in EtOH (20 mL), and hydrogen substitution was carried out, followed by stirring at room temperature for 6 hours. The reaction mixture was passed through Celite, and the solvent was distilled off. The residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate) to give tert-butyl N-[(3S)-1-[3-(4-cyanophenyl)-4-(4-hydroxyphenyl)benzoyl]pyrrolidin-3-yl]carbamate.
[0379]

Step 3

The tert-butyl N-[(3S)-1-[3-(4-cyanophenyl)-4-(4-hydroxyphenyl)benzoyl]pyrrolidin-3-yl]carbamate (15 mg) obtained in step 2 above was dissolved in DMF (0.5 mL). At room temperature, K₂CO₃ (6.4 mg) and 2-methyloxirane (5.4 mg) were added thereto, followed by stirring at 120°C for 2 hours. Ethyl acetate was added thereto, the resulting mixture was washed sequentially with water and saturated brine, and dried over anhydrous sodium sulfate, and the solvent was distilled off. The residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate) to give tert-butyl N-[(3S)-1-[3-(4-cyanophenyl)-4-[4-(2-hydroxypropoxy)phenyl]benzoyl]pyrrolidin-3-yl]carbamate.

[0380]

Step 4

The tert-butyl N-[(3S)-1-[3-(4-cyanophenyl)-4-[4-(2-hydroxypropoxy)phenyl]benzoyl]pyrrolidin-3-yl]carbamate (15 mg) obtained in step 3 above was dissolved in TFA (0.3 mL), and the progress of the reaction was confirmed with LCMS, followed by vacuum concentration. The residue was purified by reversed-phase HPLC (mobile phase: water/acetonitrile) to give the title compound.

[0381]

Example 36: Synthesis of 4-[5-[(3S)-3-aminopyrrolidine-1-carbonyl]-2-(2-fluoro-4-methyl-phenyl)phenyl]-2-fluoro-benzonitrile

The procedure of steps 1 to 3 in Example 16 was conducted using (2-fluoro-4-methyl-phenyl)boronic acid instead of p-tolylboronic acid to give the title compound.

[0382]

Example 37: Synthesis of 4-[5-[(3S)-3-aminopyrrolidine-1-carbonyl]-2-[2-fluoro-4-(2-hydroxy-2-methyl-propyl)phenyl]phenyl]-2-fluoro-benzonitrile

Step 1

The tert-butyl N-[(3S)-1-[4-chloro-3-(4-cyano-3-fluoro-phenyl)benzoyl]pyrrolidin-3-yl]carbamate (4 g) obtained in

Example 16 (step 1) was dissolved in 1,4-dioxane (45 mL). At room temperature, Pd(OAc)₂ (0.40 g), KOAc (2.7 g), bis(pinacolato)diboron (4.6 g), and Silica-SMAP (0.72 g) were added thereto, followed by stirring at 150°C for 18 hours. The
5 reaction mixture was filtrated, and the solvent was distilled off. The residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate) to give tert-butyl N-[(3S)-1-[3-(4-cyano-3-fluoro-phenyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoyl]pyrrolidin-3-yl]carbamate.

10 [0383]

Step 2

The tert-butyl N-[(3S)-1-[3-(4-cyano-3-fluoro-phenyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoyl]pyrrolidin-3-yl]carbamate (30 mg) obtained in step 1
15 above and the 1-(4-bromo-3-fluoro-phenyl)-2-methyl-propan-2-ol (28 mg) obtained in Example 28 (step 1) were dissolved in 1,4-dioxane (0.8 mL). At room temperature, Pd(PPh₃)₄ (3.2 mg) and a 2 M Na₂CO₃ aqueous solution (0.4 mL) were added thereto, and the mixture was stirred in a microwave reactor at 120°C for 30
20 minutes. The reaction mixture was filtrated, and the solvent was distilled off. Ethyl acetate was added thereto, the resulting mixture was washed sequentially with water and saturated brine, and dried over anhydrous sodium sulfate, and the solvent was distilled off. The residue was purified by silica gel column
25 chromatography (mobile phase: chloroform/methanol) to give tert-butyl N-[(3S)-1-[3-(4-cyano-3-fluoro-phenyl)-4-[2-fluoro-4-(2-hydroxy-2-methyl-propyl)phenyl]benzoyl]pyrrolidin-3-yl]carbamate.
[0384]

Step 3

30 The tert-butyl N-[(3S)-1-[3-(4-cyano-3-fluoro-phenyl)-4-[2-fluoro-4-(2-hydroxy-2-methyl-propyl)phenyl]benzoyl]pyrrolidin-3-yl]carbamate (20 mg) obtained in step 2 above was dissolved in MeOH (1 mL). At room temperature, a 12 M HCl aqueous solution (1 mL) was added
35 thereto, followed by stirring at room temperature for 30 minutes.

The reaction mixture was neutralized by the addition of water (1 mL) and a 2 M aqueous sodium hydroxide solution (6 mL).

Chloroform was added thereto, the mixture was washed sequentially with water and saturated brine, and dried over anhydrous sodium sulfate. Thereafter, the solvent was distilled off to give the title compound.

[0385]

Example 38: Synthesis of 2-fluoro-4-[2-[2-fluoro-4-(2-methoxyethyl)phenyl]-5-(9-oxa-2,6-diazaspiro[3.5]nonane-2-carbonyl)phenyl]benzonitrile

The procedure of steps 1 to 4 in Example 22 was conducted using tert-butyl 9-oxa-2,6-diazaspiro[3.5]nonane-6-carboxylic acid instead of tert-butyl N-[(3-endo)-8-azabicyclo[3.2.1]octan-3-yl]carbamate to give the title compound.

[0386]

Example 39: Synthesis of 4-[5-(2,3,3a,4,6,6a-hexahydro-1H-pyrrolo[3,4-c]pyrrole-5-carbonyl)-2-[2-fluoro-4-(2-methoxyethyl)phenyl]phenyl]-2-fluoro-benzonitrile

The procedure of steps 1 to 4 in Example 22 was conducted using tert-butyl 2,3,3a,4,6,6a-hexahydro-1H-pyrrolo[3,4-c]pyrrole-5-carboxylic acid instead of tert-butyl N-[(3-endo)-8-azabicyclo[3.2.1]octan-3-yl]carbamate to give the title compound.

[0387]

Example 40: 4-[5-[(3S)-3-aminopyrrolidine-1-carbonyl]-2-[4-(2-hydroxy-2-methyl-propyl)phenyl]phenyl]-2-fluoro-benzonitrile

The procedure of steps 1 to 3 in Example 37 was conducted using 1-(4-bromophenyl)-2-methylpropan-2-ol instead of 1-(4-bromo-3-fluoro-phenyl)-2-methylpropan-2-ol to give the title compound.

[0388]

Example 41: Synthesis of 4-[5-[(3-endo)-3-amino-8-azabicyclo[3.2.1]octane-8-carbonyl]-2-[2-fluoro-4-(2-hydroxy-2-methyl-propyl)phenyl]phenyl]-2-fluoro-benzonitrile

Step 1

The tert-butyl 4-chloro-3-(4-cyano-3-fluoro-phenyl)benzoate (300 mg) obtained in Example 22 (step 1) was dissolved in 1,4-dioxane (5 mL). At room temperature, Pd(OAc)₂ (40 mg), KOAc (300 mg), bis(pinacolato)diboron (500 mg), and Silica-SMAP (50 mg) were added thereto, followed by stirring at 100°C for 26 hours. The reaction mixture was filtrated, and the solvent was distilled off. The residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate) to give tert-butyl 3-(4-cyano-3-fluoro-phenyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate.

[0389]

Step 2

The tert-butyl 3-(4-cyano-3-fluoro-phenyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (100 mg) obtained in step 1 above was dissolved in DCM (1.2 mL). At room temperature, TFA (1.00 mL) was added thereto, followed by stirring at room temperature for 30 minutes. The reaction mixture was vacuum-concentrated, and the solvent was distilled off. Chloroform was added thereto, the mixture was washed with water and dried over anhydrous sodium sulfate, and the solvent was distilled off to give 3-(4-cyano-3-fluoro-phenyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoic acid.

[0390]

Step 3

The 3-(4-cyano-3-fluoro-phenyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoic acid (500 mg) obtained in step 2 above and tert-butyl N-[(3-endo)-8-azabicyclo[3.2.1]octan-3-yl]carbamate (308 mg) were dissolved in THF (4.5 mL). At room temperature, TEA (0.57 mL) and HATU (1 g) were added thereto, followed by stirring at 50°C for 1 hour. The reaction mixture was vacuum-concentrated, and the residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate) to give tert-butyl N-[(3-endo)-8-[3-(4-cyano-3-fluoro-phenyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoyl]-8-azabicyclo[3.2.1]octan-3-yl]carbamate.

[0391]

Step 4

The tert-butyl N-[(3-endo)-8-[3-(4-cyano-3-fluoro-phenyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoyl]-8-azabicyclo[3.2.1]octan-3-yl]carbamate (30 mg) obtained in step 5 3 above and the 1-(4-bromo-3-fluoro-phenyl)-2-methyl-propan-2-ol (19 mg) obtained in Example 28 (step 1) were dissolved in 1,4-dioxane (0.5 mL). At room temperature, Pd(PPh₃)₄ (18 mg) and a 2 M Na₂CO₃ aqueous solution (0.3 mL) were added thereto, and the 10 mixture was stirred in a microwave reactor at 120°C for 30 minutes. The supernatant of the reaction mixture was collected and filtered through NH-silica gel, and the solvent was distilled off to give tert-butyl-N-[(3-endo)-8-[3-(4-cyano-3-fluoro-phenyl)-4-[2-fluoro-4-(2-hydroxy-2-methyl-propyl)phenyl]benzoyl]-15 8-azabicyclo[3.2.1]octan-3-yl]carbamate.

[0392]

Step 5

The tert-butyl-N-[(3-endo)-8-[3-(4-cyano-3-fluoro-phenyl)-4-[2-fluoro-4-(2-hydroxy-2-methyl-propyl)phenyl]benzoyl]-20 8-azabicyclo[3.2.1]octan-3-yl]carbamate (15 mg) obtained in step 4 above was dissolved in MeOH (0.5 mL). At room temperature, a 12 M HCl aqueous solution (0.5 mL) was added thereto, followed by stirring at room temperature for 30 minutes. Then, water and a 2 M aqueous sodium hydroxide solution (3 mL) were added thereto, 25 and the mixture was partitioned and extracted with chloroform. After the organic layer was dried over anhydrous sodium sulfate, the solvent was distilled off. The residue was purified by reversed-phase HPLC (mobile phase: water/acetonitrile) to give the title compound.

30 [0393]

Example 42: Synthesis of 4-[5-[(3S)-3-(methylamino)pyrrolidine-1-carbonyl]-2-(p-tolyl)phenyl]benzonitrile

The procedure of steps 1 to 5 in Example 1 was conducted using (S)-tert-butyl methyl(pyrrolidin-3-yl)carbamate 35 instead of tert-butyl N-[(3S)-pyrrolidin-3-yl]carbamate to give

the title compound.

[0394]

Example 43: Synthesis of 4-[5-[(3S)-3-aminopyrrolidine-1-carbonyl]-2-(4-benzyloxyphenyl)phenyl]benzonitrile

5 The procedure of steps 1 to 5 in Example 9 was conducted using 1-(benzyloxy)-4-bromobenzene instead of 1-bromo-2-chloro-4-methyl-benzene to give the title compound.

[0395]

Example 44: Synthesis of 1-[4-[4-[(3S)-3-aminopyrrolidine-1-carbonyl]-2-(4-cyanophenyl)phenyl]phenyl]-N-phenyl-cyclopropanecarboxamide

10 The procedure of steps 1 to 5 in Example 9 was conducted using 1-(4-bromophenyl)-N-phenylcyclopropanecarboxamide instead of 1-bromo-2-chloro-4-methyl-benzene to give the title
15 compound.

[0396]

Example 45: Synthesis of 2-[4-[4-[(3S)-3-aminopyrrolidine-1-carbonyl]-2-(4-cyanophenyl)phenyl]phenyl]ethyl acetate

Step 1

20 The procedure of step 1 in Example 29 was conducted using 2-(4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl)ethanol instead of [4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl]methanol to give 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenethyl acetate.

25 [0397]

Step 2

 The procedure of steps 1 to 2 in Example 12 was conducted using the 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenethyl acetate obtained in step 1 above instead of 4-methyl-2-nitrophenylboronic acid pinacol ester to give the title
30 compound.

[0398]

Example 46: Synthesis of 4-[2-[4-(2-hydroxyethyl)phenyl]-5-[(3S)-3-(methylamino)pyrrolidine-1-carbonyl]phenyl]benzonitrile

35 Step 1

The procedure of steps 1 to 5 in Example 1 was conducted using the 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenethyl acetate obtained in Example 45 (step 1) instead of p-tolylboronic acid, and using (S)-tert-butyl methyl(pyrrolidin-3-yl)carbamate instead of tert-butyl N-[(3S)-pyrrolidin-3-yl]carbamate to give 2-[4-[2-(4-cyanophenyl)-4-[(3S)-3-(methylamino)pyrrolidine-1-carbonyl]phenyl]phenyl]ethyl acetate. [0399]

Step 2

The procedure of step 3 in Example 29 was conducted using the 2-[4-[2-(4-cyanophenyl)-4-[(3S)-3-(methylamino)pyrrolidine-1-carbonyl]phenyl]phenyl]ethyl acetate obtained in step 1 above instead of [4-[4-[(3S)-3-(tert-butoxycarbonylamino)pyrrolidine-1-carbonyl]-2-(4-cyanophenyl)phenyl]phenyl]methyl acetate to give the title compound.

[0400]

Example 47: Synthesis of 4-[2-[4-(2-methoxyethyl)phenyl]-5-[(3S)-3-(methylamino)pyrrolidine-1-carbonyl]phenyl]benzonitrile

Step 1

The procedure of steps 1 to 5 in Example 1 was conducted using (4-(2-methoxyethyl)phenyl)boronic acid instead of p-tolylboronic acid, and using (S)-tert-butyl methyl(pyrrolidin-3-yl)carbamate instead of tert-butyl N-[(3S)-pyrrolidin-3-yl]carbamate to give the title compound.

[0401]

Example 48: Synthesis of 4-[5-[(3S)-3-(dimethylamino)pyrrolidine-1-carbonyl]-2-[4-[1-(hydroxymethyl)cyclopropyl]phenyl]phenyl]benzonitrile

The procedure of steps 1 to 4 in Example 1 was conducted using [4-[1-(hydroxymethyl)cyclopropyl]phenyl]boronic acid instead of p-tolylboronic acid, and using (S)-N,N-dimethylpyrrolidin-3-amine instead of tert-butyl N-[(3S)-pyrrolidin-3-yl]carbamate to give the title compound.

[0402]

Example 49: Synthesis of 4-[5-[(3S)-3-aminopyrrolidine-1-carbonyl]-2-(3-fluoro-4-methyl-phenyl)phenyl]benzonitrile

The procedure of steps 1 to 2 in Example 12 was conducted using (3-fluoro-4-methyl-phenyl)boronic acid instead of
5 4-methyl-2-nitrophenylboronic acid pinacol ester to give the title compound.

[0403]

Example 50: Synthesis of 4-[5-[(3S)-3-aminopyrrolidine-1-carbonyl]-2-(4-chlorophenyl)phenyl]benzonitrile

10 The procedure of steps 1 to 5 in Example 9 was conducted using 1-bromo-4-chloro-benzene instead of 1-bromo-2-chloro-4-methyl-benzene to give the title compound.

[0404]

Example 51: Synthesis of 4-[5-[(3S)-3-aminopyrrolidine-1-carbonyl]-2-(4-bromophenyl)phenyl]benzonitrile

15 The procedure of steps 1 to 5 in Example 9 was conducted using 1,4-dibromobenzene instead of 1-bromo-2-chloro-4-methyl-benzene to give the title compound.

[0405]

20 Example 52: Synthesis of 5'-((1S,4S)-2,5-diazabicyclo[2.2.1]heptane-2-carbonyl)-4'''-methyl-[1,1':2',1'''-terphenyl]-4-carbonitrile

The procedure of steps 1 to 5 in Example 1 was conducted using tert-butyl (1S,4S)-2,5-
25 diazabicyclo[2.2.1]heptane-2-carboxylate instead of tert-butyl N-[(3S)-pyrrolidin-3-yl]carbamate to give the title compound.

[0406]

Example 53: Synthesis of 4-[2-[4-(2-aminoethyl)phenyl]-5-[(3S)-3-aminopyrrolidine-1-carbonyl]phenyl]benzonitrile

30 Step 1

The tert-butyl N-[(3S)-1-[3-(4-cyanophenyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoyl]pyrrolidin-3-yl]carbamate (50 mg) obtained in Example 9 (step 3) was dissolved in 1,4-dioxane (0.48 mL). At room temperature, 2-(4-bromophenyl)ethanamine (29 mg), Pd(PPh₃)₄ (3.4 mg), and a 2 M
35

Na₂CO₃ aqueous solution (0.24 mL) were added thereto, and the mixture was stirred in a microwave reactor at 120°C for 30 minutes. The residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate) to give tert-butyl N-[(3S)-1-[4-[4-(2-aminoethyl)phenyl]-3-(4-

cyanophenyl)benzoyl]pyrrolidin-3-yl]carbamate.

[0407]

Step 2

The tert-butyl N-[(3S)-1-[4-[4-(2-aminoethyl)phenyl]-3-(4-cyanophenyl)benzoyl]pyrrolidin-3-yl]carbamate (15 mg) obtained in step 1 above was dissolved in TFA (0.3 mL), and the progress of the reaction was confirmed with LCMS, followed by vacuum concentration. The residue was purified by reversed-phase HPLC (mobile phase: water/acetonitrile) to give the title compound.

[0408]

Example 54: Synthesis of 4-[5-[(3S)-3-aminopyrrolidine-1-carbonyl]-2-(4-iodophenyl)phenyl]benzonitrile

The procedure of steps 1 to 5 in Example 9 was conducted using 1,4-diiodobenzene instead of 1-bromo-2-chloro-4-methyl-benzene to give the title compound.

[0409]

Example 55: Synthesis of N-[2-[4-[4-[(3S)-3-aminopyrrolidine-1-carbonyl]-2-(4-cyanophenyl)phenyl]phenyl]ethyl]acetamide

The tert-butyl N-[(3S)-1-[4-[4-(2-aminoethyl)phenyl]-3-(4-cyanophenyl)benzoyl]pyrrolidin-3-yl]carbamate (15 mg) obtained in Example 53 (step 1) was dissolved in THF. At room temperature, TEA (0.02 mL) and then acetylchloride (4.6 mg) were added thereto, followed by stirring at room temperature for 1 hour. TFA was added to the residue, and the progress of the reaction was confirmed with LCMS, followed by vacuum concentration. The residue was purified by reversed-phase HPLC (mobile phase: water/acetonitrile) to give the title compound.

[0410]

Example 56: Synthesis of 4-[5-[(3S)-3-aminopyrrolidine-1-carbonyl]-2-(4-propylphenyl)phenyl]benzonitrile

The procedure of steps 1 to 2 in Example 12 was conducted using (4-propylphenyl)boronic acid instead of 4-methyl-2-nitrophenylboronic acid pinacol ester to give the title compound.

5 [0411]

Example 57: Synthesis of 4-[5-[(3S)-3-aminopyrrolidine-1-carbonyl]-2-(2-naphthyl)phenyl]benzonitrile

The procedure of steps 1 to 2 in Example 12 was conducted using 2-naphthyl boronic acid instead of 4-methyl-2-nitrophenylboronic acid pinacol ester to give the title compound.
10 [0412]

Example 58: Synthesis of 4-[2-[4-[1-(hydroxymethyl)cyclopropyl]phenyl]-5-[(3S)-3-(methylamino)pyrrolidine-1-carbonyl]phenyl]benzonitrile

15 The procedure of steps 1 to 5 in Example 1 was conducted using [4-[1-(hydroxymethyl)cyclopropyl]phenyl]boronic acid instead of p-tolylboronic acid, and using (S)-tert-butyl methyl(pyrrolidin-3-yl)carbamate instead of tert-butyl N-[(3S)-pyrrolidin-3-yl]carbamate to give the title compound.

20 [0413]

Example 59: Synthesis of 4-[5-[(3S)-3-aminopyrrolidine-1-carbonyl]-2-[4-[1-(hydroxycyclopropyl)methyl]phenyl]phenyl]benzonitrile

The procedure of steps 1 to 5 in Example 9 was conducted using 1-[(4-bromophenyl)methyl]cyclopropanol instead of 1-bromo-2-chloro-4-methyl-benzene to give the title compound.
25 [0414]

Example 60: Synthesis of 4-[5-[(3S)-3-aminopyrrolidine-1-carbonyl]-2-[4-(2-methylprop-1-enyl)phenyl]phenyl]benzonitrile

30 The procedure of steps 1 to 5 in Example 9 was conducted using 1-bromo-4-(2-methylprop-1-enyl)benzene instead of 1-bromo-2-chloro-4-methyl-benzene to give the title compound.
[0415]

Example 61: Synthesis of 4-[5-[(3S)-3-aminopyrrolidine-1-carbonyl]-2-[4-(3-hydroxy-3-methyl-

35

butyl)phenyl]phenyl]benzonitrile

The procedure of steps 1 to 5 in Example 9 was conducted using 4-(4-bromophenyl)-2-methyl-butan-2-ol instead of 1-bromo-2-chloro-4-methyl-benzene to give the title compound.

5 [0416]

Example 62: Synthesis of 4-[5-[(3S)-3-aminopyrrolidine-1-carbonyl]-2-[4-[2-(1-hydroxycyclopropyl)ethyl]phenyl]phenyl]benzonitrile

10 The procedure of steps 1 to 5 in Example 9 was conducted using 1-[2-(4-bromophenyl)ethyl]cyclopropanol instead of 1-bromo-2-chloro-4-methyl-benzene to give the title compound. [0417]

Example 63: Synthesis of 4-[5-[(3S)-3-aminopyrrolidine-1-carbonyl]-2-[4-(2-hydroxyethyl)phenyl]phenyl]-2-fluoro-
15 benzonitrile

The procedure of steps 1 to 3 in Example 37 was conducted using 2-(4-bromophenyl)ethanol instead of 1-(4-bromo-3-fluoro-phenyl)-2-methyl-propan-2-ol to give the title compound. [0418]

20 Example 64: Synthesis of 4-[5-[(3S)-3-aminopyrrolidine-1-carbonyl]-2-[2-fluoro-4-(2-hydroxy-2-methyl-propyl)phenyl]phenyl]benzonitrile

The procedure of steps 1 to 5 in Example 9 was conducted using the 1-(4-bromo-3-fluoro-phenyl)-2-methyl-propan-2-ol obtained in Example 28 (step 1) instead of 1-bromo-2-chloro-4-methyl-benzene to give the title compound. [0419]

Example 65: Synthesis of 4-[5-[(3S)-3-aminopyrrolidine-1-carbonyl]-2-[4-(3-hydroxy-3-methyl-butyl)phenyl]phenyl]-2-fluoro-
30 benzonitrile

The procedure of steps 1 to 3 in Example 16 was conducted using 2-methyl-4-[4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl]butan-2-ol instead of p-tolylboronic acid to give the title compound.

35 [0420]

Example 66: Synthesis of 4-[5-[(3S)-3-aminopyrrolidine-1-carbonyl]-2-[4-[1-(methoxymethyl)cyclopropyl]phenyl]phenyl]benzonitrile

The procedure of steps 1 to 5 in Example 9 was conducted using 1-bromo-4-[1-(methoxymethyl)cyclopropyl]benzene instead of 1-bromo-2-chloro-4-methyl-benzene to give the title compound.

[0421]

Example 67: Synthesis of 4-[5-[(3S)-3-aminopyrrolidine-1-carbonyl]-2-[2-fluoro-4-[(1-hydroxycyclopropyl)methyl]phenyl]phenyl]benzonitrile

Step 1

Methyl-2-(4-bromo-3-fluoro-phenyl)acetate (500 mg) and titanium isopropoxide (0.84 mL) were dissolved in THF (5 mL). At 0°C, a solution of 3 M EtMgBr in diethyl ether (1.9 mL) was added thereto dropwise, followed by stirring at room temperature overnight. Ethyl acetate was added thereto, the mixture was washed sequentially with water and saturated brine, and dried over anhydrous sodium sulfate, and the solvent was distilled off. The residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate) to give 1-[(4-bromo-3-fluoro-phenyl)methyl]cyclopropanol.

[0422]

Step 2

The procedure of steps 1 to 5 in Example 9 was conducted using the 1-[(4-bromo-3-fluoro-phenyl)methyl]cyclopropanol obtained in step 1 above instead of 1-bromo-2-chloro-4-methyl-benzene to give the title compound.

[0423]

Example 68: Synthesis of 4-[5-[(3S)-3-aminopyrrolidine-1-carbonyl]-2-[4-[1-(1-hydroxycyclopropyl)cyclopropyl]phenyl]phenyl]benzonitrile

Step 1

The procedure of step 1 in Example 67 was conducted using methyl 1-(4-bromophenyl)cyclopropanecarboxylic acid instead

of methyl-2-(4-bromo-3-fluoro-phenyl)acetate to give 1-[1-(4-bromophenyl)cyclopropyl]cyclopropanol.

[0424]

Step 2

5 The procedure of steps 1 to 5 in Example 9 was conducted using the 1-[1-(4-bromophenyl)cyclopropyl]cyclopropanol obtained in step 1 above instead of 1-bromo-2-chloro-4-methylbenzene to give the title compound.

[0425]

10 Example 69: Synthesis of 4-[5-[(3S)-3-aminopyrrolidine-1-carbonyl]-2-[4-(2-methoxyethyl)phenyl]phenyl]-2-fluorobenzonitrile

 The procedure of steps 1 to 3 in Example 16 was conducted using the 2-[4-(2-methoxyethyl)phenyl]-4,4,5,5-tetramethyl-1,3,2-dioxaborolane obtained in Example 30 (step 1) instead of p-tolylboronic acid to give the title compound.

[0426]

20 Example 70: Synthesis of 4-[5-[(3S)-3-aminopyrrolidine-1-carbonyl]-2-[2-fluoro-4-(2-hydroxyethyl)phenyl]phenyl]benzonitrile

 The procedure of steps 1 to 5 in Example 9 was conducted using 2-(4-bromo-3-fluoro-phenyl)ethanol instead of 1-bromo-2-chloro-4-methylbenzene to give the title compound.

[0427]

25 Example 71: Synthesis of 4-[5-[(3S)-3-aminopyrrolidine-1-carbonyl]-2-[2-fluoro-4-(2-methoxyethyl)phenyl]phenyl]benzonitrile

 The procedure of steps 1 to 5 in Example 9 was conducted using 1-bromo-2-fluoro-4-(2-methoxyethyl)benzene instead of 1-bromo-2-chloro-4-methylbenzene to give the title compound.

[0428]

35 Example 72: Synthesis of 4-[5-[(3S)-3-aminopyrrolidine-1-carbonyl]-2-[4-(2-hydroxy-1,1-dimethylethyl)phenyl]phenyl]benzonitrile

The procedure of steps 1 to 5 in Example 9 was conducted using 2-(4-bromophenyl)-2-methyl-propan-1-ol instead of 1-bromo-2-chloro-4-methyl-benzene to give the title compound.
[0429]

5 Example 73: Synthesis of 4-[5-[(3S)-3-aminopyrrolidine-1-carbonyl]-2-[4-(2-fluoroethyl)phenyl]phenyl]benzonitrile

The procedure of steps 1 to 5 in Example 9 was conducted using 1-bromo-4-(2-fluoroethyl)benzene instead of 1-bromo-2-chloro-4-methyl-benzene to give the title compound.

10 [0430]

Example 74: Synthesis of 4-[5-(2,7-diazaspiro[3.4]octane-7-carbonyl)-2-[2-fluoro-4-(2-methoxyethyl)phenyl]phenyl]-2-fluorobenzonitrile

The procedure of steps 1 to 4 in Example 22 was conducted using tert-butyl 2,7-diazaspiro[3.4]octane-2-carboxylic acid instead of tert-butyl N-[(3-endo)-8-azabicyclo[3.2.1]octan-3-yl]carbamate to give the title compound.

[0431]

20 Example 75: Synthesis of 4-[5-(2,8-diazaspiro[3.5]nonane-2-carbonyl)-2-[2-fluoro-4-(2-methoxyethyl)phenyl]phenyl]-2-fluorobenzonitrile

The procedure of steps 1 to 4 in Example 22 was conducted using tert-butyl 2,8-diazaspiro[3.5]nonane-8-carboxylic acid instead of tert-butyl N-[(3-endo)-8-azabicyclo[3.2.1]octan-3-yl]carbamate to give the title compound.

[0432]

Example 76: Synthesis of 4-[5-(2,7-diazaspiro[3.4]octane-7-carbonyl)-2-[2-fluoro-4-(2-hydroxy-2-methyl-propyl)phenyl]phenyl]-2-fluorobenzonitrile

30 The procedure of steps 1 to 5 in Example 41 was conducted using tert-butyl 2,7-diazaspiro[3.4]octane-2-carboxylic acid instead of tert-butyl N-[(3-endo)-8-azabicyclo[3.2.1]octan-3-yl]carbamate to give the title compound.

[0433]

35 Example 77: Synthesis of 4-[5-(2,8-diazaspiro[3.5]nonane-2-

carbonyl)-2-[2-fluoro-4-(2-hydroxy-2-methyl-propyl)phenyl]phenyl]-2-fluoro-benzonitrile

The procedure of steps 1 to 5 in Example 41 was conducted using tert-butyl 2,8-diazaspiro[3.5]nonane-8-carboxylic acid instead of tert-butyl N-[(3-endo)-8-azabicyclo[3.2.1]octan-3-yl]carbamate to give the title compound.

[0434]

Example 78: Synthesis of 4-[5-(3,8-diazaspiro[4.5]decane-8-carbonyl)-2-[2-fluoro-4-(2-methoxyethyl)phenyl]phenyl]-2-fluoro-benzonitrile hydrochloride

The procedure of steps 1 to 4 in Example 22 was conducted using tert-butyl 3,8-diazaspiro[4.5]decane-3-carboxylic acid instead of tert-butyl N-[(3-endo)-8-azabicyclo[3.2.1]octan-3-yl]carbamate to give the title compound.

[0435]

Example 79: Synthesis of 4-[5-(2,8-diazaspiro[3.5]nonane-8-carbonyl)-2-[2-fluoro-4-(2-methoxyethyl)phenyl]phenyl]-2-fluoro-benzonitrile

The procedure of steps 1 to 4 in Example 22 was conducted using tert-butyl 2,8-diazaspiro[3.5]nonane-2-carboxylic acid instead of tert-butyl N-[(3-endo)-8-azabicyclo[3.2.1]octan-3-yl]carbamate to give the title compound.

[0436]

Example 80: Synthesis of 4-[5-(1,4-diazepane-1-carbonyl)-2-[2-fluoro-4-(2-methoxyethyl)phenyl]phenyl]-2-fluoro-benzonitrile hydrochloride

The procedure of steps 1 to 4 in Example 22 was conducted using tert-butyl-1,4-diazepane-1-carboxylic acid instead of tert-butyl N-[(3-endo)-8-azabicyclo[3.2.1]octan-3-yl]carbamate to give the title compound.

[0437]

Example 81: Synthesis of 4-[5-(3,7-diazaspiro[3.4]octane-7-carbonyl)-2-[2-fluoro-4-(2-methoxyethyl)phenyl]phenyl]-2-fluoro-benzonitrile

The procedure of steps 1 to 4 in Example 22 was

conducted using tert-butyl-3,7-diazaspiro[3.4]octane-3-carboxylic acid instead of tert-butyl N-[(3-endo)-8-azabicyclo[3.2.1]octan-3-yl]carbamate to give the title compound.

[0438]

5 Example 82: Synthesis of 4-[5-[(1S,4S)-2,5-diazabicyclo[2.2.1]heptane-2-carbonyl]-2-[2-fluoro-4-(2-methoxyethyl)phenyl]phenyl]-2-fluoro-benzonitrile

The procedure of steps 1 to 4 in Example 22 was conducted using tert-butyl (1S,4S)-2,5-

10 diazabicyclo[2.2.1]heptane-2-carboxylic acid instead of tert-butyl N-[(3-endo)-8-azabicyclo[3.2.1]octan-3-yl]carbamate to give the title compound.

[0439]

Example 83: Synthesis of 4-[5-(3,7-diazaspiro[3.5]nonane-7-carbonyl)-2-[2-fluoro-4-(2-methoxyethyl)phenyl]phenyl]-2-fluoro-benzonitrile

The procedure of steps 1 to 4 in Example 22 was conducted using tert-butyl 3,7-diazaspiro[3.5]nonane-3-carboxylic acid instead of tert-butyl N-[(3-endo)-8-azabicyclo[3.2.1]octan-3-yl]carbamate to give the title compound.

[0440]

Example 84: Synthesis of 4-[5-(2,7-diazaspiro[3.5]nonane-2-carbonyl)-2-[2-fluoro-4-(2-methoxyethyl)phenyl]phenyl]-2-fluoro-benzonitrile

25 The procedure of steps 1 to 4 in Example 22 was conducted using tert-butyl 2,7-diazaspiro[3.5]nonane-7-carboxylic acid hydrochloride instead of tert-butyl N-[(3-endo)-8-azabicyclo[3.2.1]octan-3-yl]carbamate to give the title compound.

[0441]

30 Example 85: Synthesis of 4-[5-[(1R,4R)-2,5-diazabicyclo[2.2.1]heptane-2-carbonyl]-2-[2-fluoro-4-(2-methoxyethyl)phenyl]phenyl]-2-fluoro-benzonitrile

The procedure of steps 1 to 4 in Example 22 was conducted using tert-butyl (1R,4R)-2,5-

35 diazabicyclo[2.2.1]heptane-2-carboxylic acid instead of tert-

butyl N-[(3-endo)-8-azabicyclo[3.2.1]octan-3-yl]carbamate to give the title compound.

[0442]

Example 86: Synthesis of 2-fluoro-4-[2-[2-fluoro-4-(2-methoxyethyl)phenyl]-5-[(1S,4S)-5-methyl-2,5-diazabicyclo[2.2.1]heptane-2-carbonyl]phenyl]benzonitrile

The procedure of steps 1 to 4 in Example 22 was conducted using (1S,4S)-2-methyl-2,5-diazabicyclo[2.2.1]heptane instead of tert-butyl N-[(3-endo)-8-azabicyclo[3.2.1]octan-3-yl]carbamate to give the title compound.

[0443]

Example 87: Synthesis of 2-fluoro-4-[2-[2-fluoro-4-(2-methoxyethyl)phenyl]-5-[(1R,4R)-5-methyl-2,5-diazabicyclo[2.2.1]heptane-2-carbonyl]phenyl]benzonitrile

The procedure of steps 1 to 4 in Example 22 was conducted using (1R,4R)-2-methyl-2,5-diazabicyclo[2.2.1]heptane instead of tert-butyl N-[(3-endo)-8-azabicyclo[3.2.1]octan-3-yl]carbamate to give the title compound.

[0444]

Example 88: Synthesis of 4-[5-(3,8-diazabicyclo[3.2.1]octane-3-carbonyl)-2-[2-fluoro-4-(2-methoxyethyl)phenyl]phenyl]-2-fluorobenzonitrile

The procedure of steps 1 to 4 in Example 22 was conducted using tert-butyl 3,8-diazabicyclo[3.2.1]octane-8-carboxylic acid instead of tert-butyl N-[(3-endo)-8-azabicyclo[3.2.1]octan-3-yl]carbamate to give the title compound.

[0445]

Example 89: Synthesis of 4-[5-[(3S)-3-aminopyrrolidine-1-carbonyl]-2-(1,3-benzothiazol-5-yl)phenyl]-2-fluorobenzonitrile

The procedure of steps 1 to 3 in Example 16 was conducted using 1,3-benzothiazol-5-yl boronic acid instead of p-tolylboronic acid to give the title compound.

[0446]

Example 90: Synthesis of 4-[5-[(3S)-3-aminopyrrolidine-1-carbonyl]-2-(1-methylpyrazolo[3,4-b]pyridin-5-yl)phenyl]-2-

fluoro-benzonitrile

The procedure of steps 1 to 3 in Example 16 was conducted using (1-methylpyrazolo[3,4-b]pyridin-5-yl)boronic acid instead of p-tolylboronic acid to give the title compound.

5 [0447]

Example 91: Synthesis of 4-[5-[(3S)-3-aminopyrrolidine-1-carbonyl]-2-(1-methylbenzimidazol-5-yl)phenyl]-2-fluoro-benzonitrile

10 The procedure of steps 1 to 3 in Example 37 was conducted using 5-bromo-1-methyl-benzimidazole instead of 1-(4-bromo-3-fluoro-phenyl)-2-methyl-propan-2-ol to give the title compound.

[0448]

15 Example 92: Synthesis of 4-[5-[(3S)-3-aminopyrrolidine-1-carbonyl]-2-(1-methylindazol-5-yl)phenyl]-2-fluoro-benzonitrile

The procedure of steps 1 to 3 in Example 16 was conducted using (1-methylindazol-5-yl)boronic acid instead of p-tolylboronic acid to give the title compound.

[0449]

20 Example 93: Synthesis of 4-[5-[(3S)-3-aminopyrrolidine-1-carbonyl]-2-(2-methylindazol-5-yl)phenyl]-2-fluoro-benzonitrile

25 The procedure of steps 1 to 3 in Example 16 was conducted using 2-methyl-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)indazole instead of p-tolylboronic acid to give the title compound.

[0450]

Example 94: Synthesis of 4-[5-[(3S)-3-aminopyrrolidine-1-carbothioyl]-2-[2-fluoro-4-(2-methoxyethyl)phenyl]phenyl]-2-fluoro-benzonitrile

30 The procedure of Example 2 was conducted using the 4-[5-[(3S)-3-aminopyrrolidine-1-carbonyl]-2-[2-fluoro-4-(2-methoxyethyl)phenyl]phenyl]-2-fluoro-benzonitrile obtained in Example 20 (step 3) instead of 4-[5-[(3S)-3-aminopyrrolidine-1-carbonyl]-2-(p-tolyl)phenyl]benzonitrile to give the title
35 compound.

[0451]

Example 95: Synthesis of 4-[5-[(3S)-3-aminopyrrolidine-1-carbonyl]-2-(6-fluoro-1-methyl-benzimidazol-5-yl)phenyl]-2-fluoro-benzonitrile

5 The procedure of steps 1 to 3 in Example 37 was conducted using 5-bromo-6-fluoro-1-methyl-benzimidazole instead of 1-(4-bromo-3-fluoro-phenyl)-2-methyl-propan-2-ol to give the title compound.

[0452]

10 Example 96: Synthesis of 4-[5-[(3S)-3-aminopyrrolidine-1-carbonyl]-2-(6-fluoro-1-methyl-benzotriazol-5-yl)phenyl]-2-fluoro-benzonitrile

Step 1

 The tert-butyl N-[(3S)-1-[3-(4-cyano-3-fluoro-phenyl)-
15 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoyl]pyrrolidin-3-yl]carbamate (15 mg) obtained in Example 37 (step 1) was dissolved in 1,4-dioxane (0.5 mL). At room temperature, 5-bromo-6-fluoro-1-methyl-benzotriazole (9.7 mg), PdCl₂(dppf) (1.0 mg), and tripotassium phosphate (18 mg) were
20 added thereto, and the mixture was stirred in a microwave reactor at 125°C for 30 minutes. Ethyl acetate was added thereto, and the mixture was put on NH-silica gel and washed with ethyl acetate/methanol. The solvent was distilled off to give tert-butyl N-[(3S)-1-[3-(4-cyano-3-fluoro-phenyl)-4-(6-fluoro-1-
25 methyl-benzotriazol-5-yl)benzoyl]pyrrolidin-3-yl]carbamate.

[0453]

Step 2

 The tert-butyl N-[(3S)-1-[3-(4-cyano-3-fluoro-phenyl)-4-(6-fluoro-1-methyl-benzotriazol-5-yl)benzoyl]pyrrolidin-3-
30 yl]carbamate (15 mg) obtained in step 1 above was dissolved in TFA (0.3 mL), and the progress of the reaction was confirmed with LCMS, followed by vacuum concentration. The residue was purified by reversed-phase HPLC (mobile phase: water/acetonitrile) to give the title compound.

35 [0454]

Example 97: Synthesis of 4-[5-[(3-exo)-3-amino-8-azabicyclo[3.2.1]octane-8-carbonyl]-2-(4-fluorophenyl)phenyl]benzonitrile

The procedure of steps 1 to 3 in Example 23 was
5 conducted using 1-bromo-4-fluorobenzene instead of 5-bromo-6-fluoro-1-methyl-indole to give the title compound.
[0455]

Example 98: Synthesis of 4-[5-[(3-exo)-3-amino-8-azabicyclo[3.2.1]octane-8-carbonyl]-2-(4-chlorophenyl)phenyl]benzonitrile

The procedure of steps 1 to 3 in Example 23 was
conducted using 1-bromo-4-chloro-benzene instead of 5-bromo-6-fluoro-1-methyl-indole to give the title compound.
[0456]

15 Example 99: Synthesis of [(3S)-3-aminopyrrolidin-1-yl]-[3-(4-nitrophenyl)-4-(p-tolyl)phenyl]methanone

The procedure of steps 1 to 5 in Example 19 was
conducted using 1-bromo-4-nitro-benzene instead of 4-bromo-2,6-difluoro-benzonitrile to give the title compound.

20 [0457]

Example 100: Synthesis of 4-[5-[(3S)-3-aminopyrrolidine-1-carbonyl]-2-[6-(dimethylamino)-3-pyridyl]phenyl]-2-fluoro-benzonitrile

The procedure of steps 1 to 3 in Example 37 was
25 conducted using 5-bromo-N,N-dimethylpyridin-2-amine instead of 1-(4-bromo-3-fluoro-phenyl)-2-methyl-propan-2-ol to give the title compound.
[0458]

Example 101: Synthesis of 4-[5-[(3S)-3-aminopyrrolidine-1-carbonyl]-2-(1-methylbenzotriazol-5-yl)phenyl]-2-fluoro-benzonitrile

The procedure of steps 1 to 3 in Example 37 was
conducted using 5-bromo-1-methyl-benzotriazole instead of 1-(4-bromo-3-fluoro-phenyl)-2-methyl-propan-2-ol to give the title
35 compound.

[0459]

Example 102: Synthesis of 4-[5-[(3S)-3-aminopyrrolidine-1-carbonyl]-2-(6,7-difluoro-1-methyl-benzimidazol-5-yl)phenyl]-2-fluoro-benzonitrile

5 The procedure of steps 1 to 3 in Example 37 was conducted using 5-bromo-6,7-difluoro-1-methyl-benzimidazole instead of 1-(4-bromo-3-fluoro-phenyl)-2-methyl-propan-2-ol to give the title compound.

[0460]

10 Example 103: Synthesis of 4-[5-[(3S)-3-aminopyrrolidine-1-carbonyl]-2-(1,2-dimethylbenzimidazol-5-yl)phenyl]-2-fluoro-benzonitrile

 The procedure of steps 1 to 3 in Example 37 was conducted using 5-bromo-1,2-dimethyl-benzimidazole instead of 1-
15 (4-bromo-3-fluoro-phenyl)-2-methyl-propan-2-ol to give the title compound.

[0461]

Example 104: Synthesis of 4-[5-[(3-exo)-3-amino-8-azabicyclo[3.2.1]octane-8-carbonyl]-2-(2-naphthyl)phenyl]-2-
20 fluoro-benzonitrile

 The procedure of steps 1 to 3 in Example 23 was conducted using 2-bromonaphthalene instead of 5-bromo-6-fluoro-1-methyl-indole to give the title compound.

[0462]

25 Example 105: Synthesis of 4-[5-[(3-exo)-3-amino-8-azabicyclo[3.2.1]octane-8-carbonyl]-2-(8-fluoro-7-quinolyl)phenyl]-2-fluoro-benzonitrile

 The procedure of steps 1 to 3 in Example 23 was conducted using 7-bromo-8-fluoroquinoline instead of 5-bromo-6-
30 fluoro-1-methyl-indole to give the title compound.

[0463]

Example 106: Synthesis of 4-[5-[(3-exo)-3-amino-8-azabicyclo[3.2.1]octane-8-carbonyl]-2-(4-methyl-2,3-dihydro-1,4-benzoxazin-7-yl)phenyl]-2-fluoro-benzonitrile

35 The procedure of steps 1 to 3 in Example 23 was

conducted using 7-bromo-4-methyl-3,4-dihydro-2H-benzo[b][1,4]oxazine instead of 5-bromo-6-fluoro-1-methyl-indole to give the title compound.

[0464]

5 Example 107: Synthesis of 4-[5-[(3-exo)-3-amino-8-azabicyclo[3.2.1]octane-8-carbonyl]-2-(7-quinonyl)phenyl]-2-fluoro-benzonitrile

The procedure of steps 1 to 3 in Example 23 was conducted using 7-bromoquinoline instead of 5-bromo-6-fluoro-1-methyl-indole to give the title compound.

[0465]

Example 108: Synthesis of 4-[5-[(3-exo)-3-amino-8-azabicyclo[3.2.1]octane-8-carbonyl]-2-(6-fluoro-1-methyl-benzimidazol-5-yl)phenyl]-2-fluoro-benzonitrile

15 The procedure of steps 1 to 3 in Example 23 was conducted using 5-bromo-6-fluoro-1-methyl-benzimidazole instead of 5-bromo-6-fluoro-1-methyl-indole to give the title compound.

[0466]

Example 109: Synthesis of 4-[5-[(3-exo)-3-amino-8-azabicyclo[3.2.1]octane-8-carbonyl]-2-(6-fluoro-1-methyl-benzotriazol-5-yl)phenyl]-2-fluoro-benzonitrile

The procedure of steps 1 to 3 in Example 23 was conducted using 5-bromo-6-fluoro-1-methyl-benzotriazole instead of 5-bromo-6-fluoro-1-methyl-indole to give the title compound.

25 [0467]

Example 110: Synthesis of 4-[5-[(3-exo)-3-amino-8-azabicyclo[3.2.1]octane-8-carbonyl]-2-(4-fluoro-1-methyl-indazol-5-yl)phenyl]-2-fluoro-benzonitrile

The procedure of steps 1 to 3 in Example 23 was conducted using 5-bromo-4-fluoro-1-methyl-indazole instead of 5-bromo-6-fluoro-1-methyl-indole to give the title compound.

[0468]

Example 111: Synthesis of 4-[5-[(3-exo)-3-amino-8-azabicyclo[3.2.1]octane-8-carbonyl]-2-(2-methylindazol-5-yl)phenyl]benzonitrile

35

The procedure of steps 1 to 3 in Example 23 was conducted using 5-bromo-2-methyl-2H-indazole instead of 5-bromo-6-fluoro-1-methyl-indole to give the title compound.

[0469]

5 Example 112: Synthesis of 2-fluoro-4-[2-[2-fluoro-4-(2-methoxyethyl)phenyl]-5-[(3-exo)-3-(isopropylamino)-8-azabicyclo[3.2.1]octane-8-carbonyl]phenyl]benzonitrile

Acetone (0.002 mL) was added at 25°C to a solution of the 4-[5-[(3-exo)-3-amino-8-azabicyclo[3.2.1]octane-8-carbonyl]-2-[2-fluoro-4-(2-methoxyethyl)phenyl]phenyl]-2-fluoro-
10 benzonitrile obtained in Example 26 (step 2) in dichloromethane (0.05 mL). Subsequently, NaBH(OAc)₃ (8.45 mg) was added thereto, followed by stirring at room temperature for 1 hour. MeOH was added thereto, and the solvent was distilled off. Then, the
15 residue was purified by reversed-phase HPLC (mobile phase: water/acetonitrile) to give the title compound.

[0470]

Example 113: Synthesis of 2-fluoro-4-[2-[2-fluoro-4-(2-hydroxy-2-methyl-propyl)phenyl]-5-[(3-exo)-3-(isopropylamino)-8-azabicyclo[3.2.1]octane-8-carbonyl]phenyl]benzonitrile

The procedure of Example 112 was conducted using the 4-[5-[(3-exo)-3-amino-8-azabicyclo[3.2.1]octane-8-carbonyl]-2-[2-fluoro-4-(2-hydroxy-2-methyl-propyl)phenyl]phenyl]-2-fluoro-
25 benzonitrile obtained in Example 28 (step 3) instead of 4-[5-[(3-exo)-3-amino-8-azabicyclo[3.2.1]octane-8-carbonyl]-2-[2-fluoro-4-(2-methoxyethyl)phenyl]phenyl]-2-fluoro-benzonitrile to give the title compound.

[0471]

Example 114: Synthesis of 4-[5-[(3S)-3-(ethylamino)pyrrolidine-1-carbonyl]-2-(6-fluoro-1-methyl-benzotriazol-5-yl)phenyl]-2-fluoro-benzonitrile

Step 1

The tert-butyl N-[(3S)-1-[3-(4-cyano-3-fluoro-phenyl)-4-(6-fluoro-1-methyl-benzotriazol-5-yl)benzoyl]pyrrolidin-3-yl]carbamate (10 mg) obtained in Example 96 (step 1) was
35

dissolved in THF (0.5 mL). At room temperature, sodium hydride (0.85 mg), and then iodoethane (5.58 mg) were added thereto, the mixture was stirred at 50°C overnight, and the solvent was distilled off to give (S)-tert-butyl (1-(4'-cyano-3'-fluoro-6-(6-fluoro-1-methyl-1H-benzo[d][1,2,3]triazol-5-yl)-[1,1'-biphenyl]-3-carbonyl)pyrrolidin-3-yl)(ethyl)carbamate. The thus obtained product was used in the next step without purification.

[0472]

Step 2

10 The procedure of step 2 in Example 26 was conducted using the (S)-tert-butyl (1-(4'-cyano-3'-fluoro-6-(6-fluoro-1-methyl-1H-benzo[d][1,2,3]triazol-5-yl)-[1,1'-biphenyl]-3-carbonyl)pyrrolidin-3-yl)(ethyl)carbamate obtained in step 1 above instead of [(3-exo)-8-[3-(4-cyano-3-fluoro-phenyl)-4-[2-fluoro-4-(2-methoxyethyl)phenyl]benzoyl]-8-

15 azabicyclo[3.2.1]octan-3-yl]carbamate to give the title compound.

[0473]

Example 115: Synthesis of 2-fluoro-4-[2-(6-fluoro-1-methyl-benzotriazol-5-yl)-5-[(3S)-3-(isopropylamino)pyrrolidine-1-carbonyl]phenyl]benzonitrile

20 The procedure of Example 112 was conducted using the 4-[5-[(3S)-3-aminopyrrolidine-1-carbonyl]-2-(6-fluoro-1-methyl-benzotriazol-5-yl)phenyl]-2-fluoro-benzonitrile obtained in Example 96 (step 2) instead of 4-[5-[(3-exo)-3-amino-8-azabicyclo[3.2.1]octane-8-carbonyl]-2-[2-fluoro-4-(2-methoxyethyl)phenyl]phenyl]-2-fluoro-benzonitrile to give the title compound.

[0474]

Example 116: Synthesis of 4-[5-[(3S)-3-(cyclobutylamino)pyrrolidine-1-carbonyl]-2-(6-fluoro-1-methyl-benzotriazol-5-yl)phenyl]-2-fluoro-benzonitrile

30 The procedure of Example 112 was conducted using the 4-[5-[(3S)-3-aminopyrrolidine-1-carbonyl]-2-(6-fluoro-1-methyl-benzotriazol-5-yl)phenyl]-2-fluoro-benzonitrile obtained in Example 96 (step 2) instead of 4-[5-[(3-exo)-3-amino-8-

35

azabicyclo[3.2.1]octane-8-carbonyl]-2-[2-fluoro-4-(2-methoxyethyl)phenyl]phenyl]-2-fluoro-benzonitrile, and using cyclobutanone instead of acetone to give the title compound.
[0475]

5 Example 117: Synthesis of 4-[5-[(3S)-3-aminopyrrolidine-1-carbonyl]-2-(1-methylindolin-5-yl)phenyl]-2-fluoro-benzonitrile

The procedure of steps 1 to 3 in Example 37 was conducted using 5-bromo-1-methyl-indoline instead of 1-(4-bromo-3-fluoro-phenyl)-2-methyl-propan-2-ol to give the title compound.
10 [0476]

Example 118: Synthesis of 4-[5-[(3S)-3-aminopyrrolidine-1-carbonyl]-2-(4-methyl-2,3-dihydro-1,4-benzoxazin-7-yl)phenyl]-2-fluoro-benzonitrile

The procedure of steps 1 to 3 in Example 37 was
15 conducted using 7-bromo-4-methyl-3,4-dihydro-2H-benzo[b][1,4]oxazine instead of 1-(4-bromo-3-fluoro-phenyl)-2-methyl-propan-2-ol to give the title compound.
[0477]

Example 119: Synthesis of 4-[5-[(3S)-3-aminopyrrolidine-1-carbonyl]-2-(3-methyl-2-oxo-1,3-benzooxazol-6-yl)phenyl]-2-fluoro-benzonitrile

The procedure of steps 1 to 3 in Example 37 was
conducted using 6-bromo-3-methyl-1,3-benzooxazol-2-one instead of
1-(4-bromo-3-fluoro-phenyl)-2-methyl-propan-2-ol to give the
25 title compound.
[0478]

Example 120: Synthesis of 4-[5-[(3S)-3-aminopyrrolidine-1-carbonyl]-2-(3-methyl-2-oxo-1,3-benzothiazol-6-yl)phenyl]-2-fluoro-benzonitrile

30 The procedure of steps 1 to 3 in Example 37 was conducted using 6-bromo-3-methyl-1,3-benzothiazol-2-one instead of 1-(4-bromo-3-fluoro-phenyl)-2-methyl-propan-2-ol to give the title compound.
[0479]

35 Example 121: Synthesis of 4-[5-[(3S)-3-aminopyrrolidine-1-

carbonyl]-2-(2,3-dihydro-1,4-benzodioxin-6-yl)phenyl]-2-fluoro-benzonitrile

The procedure of steps 1 to 3 in Example 16 was conducted using 2,3-dihydro-1,4-benzodioxan-6-yl boronic acid instead of p-tolylboronic acid to give the title compound.
[0480]

Example 122: Synthesis of 4-[5-[(3S)-3-aminopyrrolidine-1-carbonyl]-2-(1,3-benzodioxol-5-yl)phenyl]-2-fluoro-benzonitrile

The procedure of steps 1 to 3 in Example 16 was conducted using 1,3-benzodioxol-5-yl boronic acid instead of p-tolylboronic acid to give the title compound.
[0481]

Example 123: Synthesis of 4-[5-[(3-endo)-3-amino-8-azabicyclo[3.2.1]octane-8-carbonyl]-2-(6-fluoro-1-methyl-indol-5-yl)phenyl]-2-fluoro-benzonitrile

The procedure of steps 1 to 5 in Example 41 was conducted using 5-bromo-6-fluoro-1-methyl-indole instead of 1-(4-bromo-3-fluoro-phenyl)-2-methyl-propan-2-ol to give the title compound.
[0482]

Example 124: Synthesis of 4-[5-[(3-endo)-3-amino-8-azabicyclo[3.2.1]octane-8-carbonyl]-2-(6-fluoro-1-methyl-indazol-5-yl)phenyl]-2-fluoro-benzonitrile

The procedure of steps 1 to 5 in Example 41 was conducted using 5-bromo-6-fluoro-1-methyl-indazole instead of 1-(4-bromo-3-fluoro-phenyl)-2-methyl-propan-2-ol to give the title compound.
[0483]

Example 125: Synthesis of 4-[5-[(3-endo)-3-amino-8-azabicyclo[3.2.1]octane-8-carbonyl]-2-(6-fluoro-1-methyl-benzotriazol-5-yl)phenyl]-2-fluoro-benzonitrile

The procedure of steps 1 to 5 in Example 41 was conducted using 5-bromo-6-fluoro-1-methyl-benzotriazole instead of 1-(4-bromo-3-fluoro-phenyl)-2-methyl-propan-2-ol to give the title compound.

[0484]

Example 126: Synthesis of 4-[5-[(3-endo)-3-amino-8-azabicyclo[3.2.1]octane-8-carbonyl]-2-(6,7-difluoro-1-methyl-benzimidazol-5-yl)phenyl]-2-fluoro-benzonitrile

5 The procedure of steps 1 to 5 in Example 41 was conducted using 5-bromo-6,7-difluoro-1-methyl-benzimidazole instead of 1-(4-bromo-3-fluoro-phenyl)-2-methyl-propan-2-ol to give the title compound.

[0485]

10 Example 127: Synthesis of 4-[5-[(3-exo)-3-amino-9-azabicyclo[3.3.1]nonane-9-carbonyl]-2-[2-fluoro-4-(2-methoxyethyl)phenyl]phenyl]-2-fluoro-benzonitrile

 The procedure of steps 1 to 4 in Example 22 was conducted using tert-butyl N-[(3-exo)-9-azabicyclo[3.3.1]nonan-3-yl]carbamate instead of tert-butyl N-[(3-endo)-8-azabicyclo[3.2.1]octan-3-yl]carbamate to give the title compound.

[0486]

Example 128: Synthesis of 4-[5-[(3-endo)-3-amino-8-azabicyclo[3.2.1]octane-8-carbonyl]-2-(4-methyl-2,3-dihydro-1,4-benzoxazin-7-yl)phenyl]-2-fluoro-benzonitrile

20 The procedure of steps 1 to 5 in Example 41 was conducted using 7-bromo-4-methyl-3,4-dihydro-2H-benzo[b][1,4]oxazine instead of 1-(4-bromo-3-fluoro-phenyl)-2-methyl-propan-2-ol to give the title compound.

25 [0487]

Example 129: Synthesis of 4-[5-[(3-endo)-3-amino-9-azabicyclo[3.3.1]nonane-9-carbonyl]-2-[2-fluoro-4-(2-methoxyethyl)phenyl]phenyl]-2-fluoro-benzonitrile

 The procedure of steps 1 to 4 in Example 22 was conducted using tert-butyl N-[(3-endo)-9-azabicyclo[3.3.1]nonan-3-yl]carbamate instead of tert-butyl N-[(3-endo)-8-azabicyclo[3.2.1]octan-3-yl]carbamate to give the title compound.

[0488]

Example 130: Synthesis of 4-[5-[(3-endo)-3-amino-8-azabicyclo[3.2.1]octane-8-carbonyl]-2-(6-fluoro-1-methyl-

35

benzimidazol-5-yl)phenyl]-2-fluoro-benzonitrile

The procedure of steps 1 to 5 in Example 41 was conducted using 5-bromo-6-fluoro-1-methyl-benzimidazole instead of 1-(4-bromo-3-fluoro-phenyl)-2-methyl-propan-2-ol to give the title compound.

[0489]

Example 131: Synthesis of 4-[5-[(3-endo)-3-amino-8-azabicyclo[3.2.1]octane-8-carbonyl]-2-[6-(dimethylamino)-3-pyridyl]phenyl]-2-fluoro-benzonitrile

The procedure of steps 1 to 5 in Example 41 was conducted using 5-bromo-N,N-dimethylpyridin-2-amine instead of 1-(4-bromo-3-fluoro-phenyl)-2-methyl-propan-2-ol to give the title compound.

[0490]

Example 132: Synthesis of 4-[5-[(3-endo)-3-amino-8-azabicyclo[3.2.1]octane-8-carbonyl]-2-(1,3,3-trimethyl-2-oxo-indolin-5-yl)phenyl]-2-fluoro-benzonitrile

The procedure of steps 1 to 5 in Example 41 was conducted using 5-bromo-1,3,3-trimethyl-indolin-2-one instead of 1-(4-bromo-3-fluoro-phenyl)-2-methyl-propan-2-ol to give the title compound.

[0491]

Example 133: Synthesis of 4-[5-[(3-endo)-3-amino-8-azabicyclo[3.2.1]octane-8-carbonyl]-2-(3-methyl-2-oxo-1,3-benzothiazol-6-yl)phenyl]-2-fluoro-benzonitrile

The procedure of steps 1 to 5 in Example 41 was conducted using 6-bromo-3-methyl-1,3-benzothiazol-2-one instead of 1-(4-bromo-3-fluoro-phenyl)-2-methyl-propan-2-ol to give the title compound.

[0492]

Example 134: Synthesis of (S)-5'-(3-aminopyrrolidine-1-carbonyl)-3-fluoro-2'-(6-fluoro-1-methyl-1H-indol-5-yl)-[1,1'-biphenyl]-4-carbonitrile

The procedure of steps 1 to 3 in Example 37 was conducted using 5-bromo-6-fluoro-1-methyl-1H-indole instead of 1-

(4-bromo-3-fluoro-phenyl)-2-methyl-propan-2-ol to give the title compound.

[0493]

Example 135: Synthesis of (S)-5'-(3-aminopyrrolidine-1-carbonyl)-3-fluoro-2'-(6-fluoro-1-methyl-1H-indazol-5-yl)-[1,1'-biphenyl]-4-carbonitrile

The procedure of steps 1 to 3 in Example 37 was conducted using 5-bromo-6-fluoro-1-methyl-1H-indazole instead of 1-(4-bromo-3-fluoro-phenyl)-2-methyl-propan-2-ol to give the title compound.

[0494]

Example 136: Synthesis of 5'-((3-endo)-3-amino-8-azabicyclo[3.2.1]octane-8-carbonyl)-3-fluoro-2'-(6-fluoro-1-(2-hydroxy-2-methylpropyl)-1H-indol-5-yl)-[1,1'-biphenyl]-4-carbonitrile

Step 1

5-Bromo-6-fluoro-1H-indole (50 mg) was dissolved in DMF (0.78 mL). At room temperature, Cs₂CO₃ (151 mg) and 2,2-dimethyloxirane (42 µL) were added thereto, followed by stirring at 90°C for 16 hours. The reaction was quenched with a saturated NH₄Cl aqueous solution, ethyl acetate was added thereto, and the mixture was washed sequentially with water and saturated brine. After the organic layer was dried over anhydrous sodium sulfate, the solvent was distilled off. The residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate) to give 1-(5-bromo-6-fluoro-indol-1-yl)-2-methyl-propan-2-ol.

[0495]

Step 2

The procedure of steps 1 to 5 in Example 41 was conducted using the 1-(5-bromo-6-fluoro-indol-1-yl)-2-methyl-propan-2-ol obtained in step 1 above instead of 1-(4-bromo-3-fluoro-phenyl)-2-methyl-propan-2-ol to give the title compound.

[0496]

Example 137: Synthesis of 5'-((3-endo)-3-amino-8-azabicyclo[3.2.1]octane-8-carbonyl)-2'-(1,3-dihydroisobenzofuran-

5-yl)-3-fluoro-[1,1'-biphenyl]-4-carbonitrile

The procedure of steps 1 to 5 in Example 41 was conducted using 5-bromo-1,3-dihydroisobenzofuran instead of 1-(4-bromo-3-fluoro-phenyl)-2-methyl-propan-2-ol to give the title compound.

[0497]

Example 138: Synthesis of 5'-((3-endo)-3-amino-8-azabicyclo[3.2.1]octane-8-carbonyl)-3-fluoro-2'-(3-isopropyl-2-oxo-2,3-dihydrobenzo[d]thiazol-6-yl)-[1,1'-biphenyl]-4-carbonitrile

Step 1

6-Bromo-3H-1,3-benzothiazol-2-one (100 mg) was dissolved in DMF (0.87 mL). At room temperature, potassium carbonate (90 mg) was added thereto, followed by stirring at 0°C for 15 minutes. At room temperature, 2-bromopropane (0.082 mL) was added thereto, followed by stirring at 100°C for 3 hours. The reaction was quenched with a saturated NH₄Cl aqueous solution, ethyl acetate was added thereto, and the resulting mixture was washed sequentially with water and saturated brine. After the organic layer was dried over anhydrous sodium sulfate, the solvent was distilled off. The residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate) to give 6-bromo-3-isopropyl-1,3-benzothiazol-2-one.

[0498]

Step 2

The procedure of steps 1 to 5 in Example 41 was conducted using the 6-bromo-3-isopropyl-1,3-benzothiazol-2-one obtained in step 1 above instead of 1-(4-bromo-3-fluoro-phenyl)-2-methyl-propan-2-ol to give the title compound.

[0499]

Example 139: Synthesis of 5'-((3-endo)-3-amino-8-azabicyclo[3.2.1]octane-8-carbonyl)-2'-(1-(tert-butyl)-6-fluoro-1H-benzo[d][1,2,3]triazol-5-yl)-3-fluoro-[1,1'-biphenyl]-4-carbonitrile

The procedure of steps 1 to 5 in Example 41 was

conducted using 5-bromo-1-(tert-butyl)-6-fluoro-1H-benzo[d][1,2,3]triazole instead of 1-(4-bromo-3-fluoro-phenyl)-2-methyl-propan-2-ol to give the title compound.

[0500]

5 Example 140: Synthesis of (S)-5'-(3-aminopyrrolidine-1-carbonyl)-2'-(1,3-dihydroisobenzofuran-5-yl)-3-fluoro-[1,1'-biphenyl]-4-carbonitrile

The procedure of steps 1 to 3 in Example 37 was conducted using 5-bromo-1,3-dihydroisobenzofuran instead of 1-(4-bromo-3-fluoro-phenyl)-2-methyl-propan-2-ol to give the title compound.

[0501]

Example 141: Synthesis of (S)-5'-(3-aminopyrrolidine-1-carbonyl)-3-fluoro-2'-(5-fluoro-3-methyl-2-oxo-2,3-dihydrobenzo[d]thiazol-6-yl)-[1,1'-biphenyl]-4-carbonitrile

Step 1

5-Fluoro-3H-1,3-benzothiazol-2-one (200 mg) was suspended in MeCN (1 mL). At room temperature, NBS (231 mg) was added thereto, followed by stirring at room temperature for 1 hour. The solvent was vacuum-concentrated, and the residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate) to give 6-bromo-5-fluoro-3H-1,3-benzothiazol-2-one.

[0502]

Step 2

The 6-bromo-5-fluoro-3H-1,3-benzothiazol-2-one (100 mg) obtained in step 1 above was dissolved in DMF (1.3 mL). At room temperature, potassium carbonate (84 mg) was added thereto, followed by stirring at 0°C for 15 minutes. At room temperature, iodomethane (0.050 mL) was added thereto, followed by stirring at room temperature for 0.5 hours. The reaction was quenched with a saturated NH₄Cl aqueous solution, ethyl acetate was added thereto, and the resulting mixture was washed sequentially with water and saturated brine. After the organic layer was dried over anhydrous sodium sulfate, the solvent was distilled off. The residue was

purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate) to give 6-bromo-5-fluoro-3-methyl-1,3-benzothiazol-2-one.

[0503]

5 Step 3

The procedure of steps 1 to 3 in Example 37 was conducted using the 6-bromo-5-fluoro-3-methyl-1,3-benzothiazol-2-one obtained in step 2 above instead of 1-(4-bromo-3-fluorophenyl)-2-methylpropan-2-ol to give the title compound.

10 [0504]

Example 142: Synthesis of 5'-((3-endo)-3-amino-8-azabicyclo[3.2.1]octane-8-carbonyl)-3-fluoro-2'-(6-fluoro-1-(2-hydroxy-2-methylpropyl)-1H-indazol-5-yl)-[1,1'-biphenyl]-4-carbonitrile

15 Step 1

5-Bromo-6-fluoro-1H-indazole (94 mg) was dissolved in DMF (1.5 mL). At room temperature, cesium carbonate (285 mg) and 2,2-dimethyloxirane (0.078 mL) were added thereto, followed by stirring at 90°C for 16 hours. The reaction was quenched with a saturated NH₄Cl aqueous solution, ethyl acetate was added thereto, and the resulting mixture was washed sequentially with water and saturated brine. After the organic layer was dried over anhydrous sodium sulfate, the solvent was distilled off. The residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate) to give 1-(5-bromo-6-fluoro-indazol-1-yl)-2-methylpropan-2-ol.

25 [0505]

Step 2

The procedure of steps 1 to 5 in Example 41 was conducted using the 1-(5-bromo-6-fluoro-indazol-1-yl)-2-methylpropan-2-ol obtained in step 1 above instead of 1-(4-bromo-3-fluorophenyl)-2-methylpropan-2-ol to give the title compound.

[0506]

Example 143: Synthesis of 4-[5-[(1S,3R,4R)-rel-3-amino-7-azabicyclo[2.2.1]heptane-7-carbonyl]-2-[2-fluoro-4-(2-

35

methoxyethyl)phenyl]phenyl]-2-fluoro-benzonitrile

Step 1

tert-Butyl (1S,3R,4R)-rel-3-amino-7-azabicyclo[2.2.1]heptane-7-carboxylate (50 mg) was dissolved in THF (1.2 mL). At 0°C, TEA (0.066 mL) and 2-nitrobenzene sulfonyl chloride (57 mg) were added thereto, followed by stirring at room temperature for 1 hour. Ethyl acetate was added thereto, and the mixture was washed sequentially with water and saturated brine. After the organic layer was dried over anhydrous sodium sulfate, the solvent was distilled off. The residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate), and the solvent was distilled off. The residue was dissolved in a 4 N hydrochloric acid-ethyl acetate solution (2 mL), followed by stirring at room temperature for 30 minutes. The reaction mixture was vacuum-concentrated to give N-[(1S,3R,4R)-rel-7-azabicyclo[2.2.1]heptan-3-yl]-2-nitrobenzenesulfonamide hydrochloride.

[0507]

Step 2

The procedure of step 3 in Example 22 was conducted using the N-[(1S,3R,4R)-rel-7-azabicyclo[2.2.1]heptan-3-yl]-2-nitrobenzenesulfonamide hydrochloride obtained in step 1 above instead of tert-butyl N-[(3-endo)-8-azabicyclo[3.2.1]octan-3-yl]carbamate to give N-[(1S,3R,4R)-rel-7-[3-(4-cyano-3-fluorophenyl)-4-[2-fluoro-4-(2-methoxyethyl)phenyl]benzoyl]-7-azabicyclo[2.2.1]heptan-3-yl]-2-nitrobenzenesulfonamide.

[0508]

Step 3

The N-[(1S,3R,4R)-rel-7-[3-(4-cyano-3-fluorophenyl)-4-[2-fluoro-4-(2-methoxyethyl)phenyl]benzoyl]-7-azabicyclo[2.2.1]heptan-3-yl]-2-nitrobenzenesulfonamide (20 mg) obtained in step 2 above was dissolved in DMF (0.5 mL). At room temperature, K₂CO₃ (21 mg) and 4-mercaptobenzoic acid (12 mg) were added thereto, followed by stirring at 40°C for 12 hours. Ethyl acetate was added thereto, and the mixture was washed

sequentially with water and saturated brine. After the organic layer was dried over anhydrous sodium sulfate, the solvent was distilled off. The residue was purified by reversed-phase HPLC (mobile phase: water/acetonitrile) to give the title compound.

5 [0509]

Example 144: Synthesis of 5'-((3-endo)-3-amino-8-azabicyclo[3.2.1]octane-8-carbonyl)-2'',3,3''-trifluoro-4''-methyl-[1,1':2',1''-terphenyl]-4-carbonitrile

The procedure of steps 1 to 5 in Example 41 was
10 conducted using 1-bromo-2,3-difluoro-4-methyl-benzene instead of 1-(4-bromo-3-fluoro-phenyl)-2-methyl-propan-2-ol to give the title compound.

[0510]

Example 145: Synthesis of 5'-((3-endo)-3-amino-8-azabicyclo[3.2.1]octane-8-carbonyl)-2'-(6,7-difluoro-1-methyl-1H-benzo[d][1,2,3]triazol-5-yl)-3-fluoro-[1,1'-biphenyl]-4-carbonitrile

Step 1

The tert-butyl N-[(3-endo)-8-[3-(4-cyano-3-fluoro-phenyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoyl]-8-azabicyclo[3.2.1]octan-3-yl]carbamate (200 mg) obtained in step 3 in Example 41 and 5-bromo-6,7-difluoro-1-methyl-benzotriazole (129 mg) were dissolved in 1,4-dioxane (1.74 mL). At room temperature, Pd(dba)₂ (16.0 mg), X-phos (26.5 mg), and
25 tripotassium phosphate (221 mg) were added thereto, and the mixture was stirred in a microwave reactor at 125°C for 1 hour. The reaction mixture was filtrated, and the solvent was distilled off. The residue was purified by silica gel column chromatography (mobile phase: ethyl acetate/hexane) to give tert-butyl N-[(3-endo)-8-[3-(4-cyano-3-fluoro-phenyl)-4-(6,7-difluoro-1-methyl-benzotriazol-5-yl)benzoyl]-8-azabicyclo[3.2.1]octan-3-yl]carbamate.
30

[0511]

Step 2

35 The tert-butyl N-[(3-endo)-8-[3-(4-cyano-3-fluoro-

phenyl)-4-(6,7-difluoro-1-methyl-benzotriazol-5-yl)benzoyl]-8-azabicyclo[3.2.1]octan-3-yl]carbamate (210 mg) obtained in step 1 above was dissolved in MeOH (1.60 mL). At room temperature, a 4 N hydrochloric acid-ethyl acetate solution (2.40 mL) was added thereto, followed by stirring at room temperature for 1 hour. The reaction mixture was vacuum-concentrated, and the solvent was distilled off. The residue was purified by reversed-phase HPLC (mobile phase: water/acetonitrile) to give the title compound. [0512]

10 Example 146: Synthesis of (S)-5'-(3-aminopyrrolidine-1-carbonyl)-3-fluoro-2'-(6-fluoro-1-(2-hydroxy-2-methylpropyl)-1H-indol-5-yl)-[1,1'-biphenyl]-4-carbonitrile

Step 1

The tert-butyl N-[(3S)-1-[3-(4-cyano-3-fluoro-phenyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoyl]pyrrolidin-3-yl]carbamate (60 mg) obtained in step 1 of Example 37 and the 1-(5-bromo-6-fluoro-indol-1-yl)-2-methylpropan-2-ol (48.1 mg) obtained in step 1 of Example 136 were dissolved in 1,4-dioxane (0.50 mL). At room temperature, Pd(dba)₂ (3.22 mg), X-phos (5.34 mg), and tripotassium phosphate (71.4 mg) were added thereto, and the mixture was stirred in a microwave reactor at 125°C for 1 hour. The reaction mixture was filtrated, and the solvent was distilled off. The residue was purified by silica gel column chromatography (mobile phase: chloroform/methanol) to give tert-butyl N-[(3S)-1-[3-(4-cyano-3-fluoro-phenyl)-4-[6-fluoro-1-(2-hydroxy-2-methyl-propyl)indol-5-yl]benzoyl]pyrrolidin-3-yl]carbamate. [0513]

Step 2

30 The tert-butyl N-[(3S)-1-[3-(4-cyano-3-fluoro-phenyl)-4-[6-fluoro-1-(2-hydroxy-2-methyl-propyl)indol-5-yl]benzoyl]pyrrolidin-3-yl]carbamate (68.0 mg) obtained in step 1 above was dissolved in MeOH (1.0 mL). At room temperature, 12 N hydrochloric acid (1.0 mL) was added thereto, followed by stirring at room temperature for 1 hour. A 2 N aqueous sodium

hydroxide solution (6.00 mL) and chloroform were added thereto, the mixture was washed sequentially with water and saturated brine, and dried over anhydrous sodium sulfate, and the solvent was distilled off. The residue was purified by reversed-phase
5 HPLC (mobile phase: water/acetonitrile) to give the title compound.

[0514]

Example 147: Synthesis of (S)-5'-(3-aminopyrrolidine-1-carbonyl)-3-fluoro-2'-(6-fluoro-1-(2-hydroxy-2-methylpropyl)-1H-indazol-5-yl)-[1,1'-biphenyl]-4-carbonitrile
10

Step 1

The tert-butyl N-[(3S)-1-[3-(4-cyano-3-fluoro-phenyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoyl]pyrrolidin-3-yl]carbamate (70 mg) obtained in step 1
15 of Example 37 and the 1-(5-bromo-6-fluoro-indazol-1-yl)-2-methylpropan-2-ol (56.3 mg) obtained in step 1 of Example 142 were dissolved in 1,4-dioxane (0.50 mL). At room temperature, Pd(dba)₂ (3.76 mg), X-phos (6.23 mg), and tripotassium phosphate (83.3 mg) were added thereto, and the mixture was stirred in a microwave
20 reactor at 125°C for 1 hour. The reaction mixture was filtrated, and the solvent was distilled off. The residue was purified by silica gel column chromatography (mobile phase: chloroform/methanol) to give tert-butyl N-[(3S)-1-[3-(4-cyano-3-fluoro-phenyl)-4-[6-fluoro-1-(2-hydroxy-2-methyl-propyl)indazol-5-yl]benzoyl]pyrrolidin-3-yl]carbamate.
25

[0515]

Step 2

The tert-butyl N-[(3S)-1-[3-(4-cyano-3-fluoro-phenyl)-4-[6-fluoro-1-(2-hydroxy-2-methyl-propyl)indazol-5-yl]benzoyl]pyrrolidin-3-yl]carbamate (70.0 mg) obtained in step 1
30 above was dissolved in MeOH (1.0 mL). At room temperature, 12 N hydrochloric acid (1.0 mL) was added thereto, followed by stirring at room temperature for 1 hour. A 2 N aqueous sodium hydroxide solution (6.00 mL) and chloroform were added thereto,
35 the mixture was washed sequentially with water and saturated

brine, and dried over anhydrous sodium sulfate, and the solvent was distilled off. The residue was purified by reversed-phase HPLC (mobile phase: water/acetonitrile) to give the title compound.

5 [0516]

Example 148: Synthesis of 5'-((3-endo)-3-amino-8-azabicyclo[3.2.1]octane-8-carbonyl)-3-fluoro-2'-(quinoxalin-6-yl)-[1,1'-biphenyl]-4-carbonitrile

The procedure of steps 1 to 5 in Example 41 was
10 conducted using 6-bromoquinoxaline instead of 1-(4-bromo-3-fluoro-phenyl)-2-methyl-propan-2-ol to give the title compound.
[0517]

Example 149: Synthesis of 5'-((3-endo)-3-amino-8-azabicyclo[3.2.1]octane-8-carbonyl)-3-fluoro-2'-(isoquinolin-6-yl)-[1,1'-biphenyl]-4-carbonitrile

The procedure of steps 1 to 5 in Example 41 was
conducted using 6-bromoisquinoline instead of 1-(4-bromo-3-fluoro-phenyl)-2-methyl-propan-2-ol to give the title compound.
[0518]

20 Example 150: Synthesis of 5'-((3-endo)-3-amino-8-azabicyclo[3.2.1]octane-8-carbonyl)-3-fluoro-2'-(isoquinolin-7-yl)-[1,1'-biphenyl]-4-carbonitrile

The procedure of steps 1 to 5 in Example 41 was
conducted using 7-bromoisquinoline instead of 1-(4-bromo-3-fluoro-phenyl)-2-methyl-propan-2-ol to give the title compound.
25 [0519]

Example 151: Synthesis of 5'-((3-endo)-3-amino-8-azabicyclo[3.2.1]octane-8-carbonyl)-3-fluoro-2'-(quinolin-6-yl)-[1,1'-biphenyl]-4-carbonitrile

30 The procedure of steps 1 to 5 in Example 41 was
conducted using 6-bromoquinoline instead of 1-(4-bromo-3-fluoro-phenyl)-2-methyl-propan-2-ol to give the title compound.
[0520]

Example 152: Synthesis of 5'-((3-endo)-3-amino-8-azabicyclo[3.2.1]octane-8-carbonyl)-3-fluoro-2'-(quinazolin-7-

35

yl)-[1,1'-biphenyl]-4-carbonitrile

The procedure of steps 1 to 5 in Example 41 was conducted using 7-bromoquinazolin instead of 1-(4-bromo-3-fluorophenyl)-2-methyl-propan-2-ol to give the title compound.

5 [0521]

Example 153: Synthesis of 5'-((3-endo)-3-amino-8-azabicyclo[3.2.1]octane-8-carbonyl)-3-fluoro-2'-(quinazolin-6-yl)-[1,1'-biphenyl]-4-carbonitrile

10 The procedure of steps 1 to 5 in Example 41 was conducted using 6-bromoquinazolin instead of 1-(4-bromo-3-fluorophenyl)-2-methyl-propan-2-ol to give the title compound.

[0522]

Example 154: Synthesis of 5'-((3-endo)-3-amino-8-azabicyclo[3.2.1]octane-8-carbonyl)-3-fluoro-2'-(phthalazin-6-yl)-[1,1'-biphenyl]-4-carbonitrile

15

The procedure of steps 1 to 5 in Example 41 was conducted using 6-bromophthalazine instead of 1-(4-bromo-3-fluorophenyl)-2-methyl-propan-2-ol to give the title compound.

[0523]

20 Example 155: Synthesis of 5'-((1R,2R,4S)-rel-2-amino-7-azabicyclo[2.2.1]heptane-7-carbonyl)-2'',3-difluoro-4''-(2-methoxyethyl)-[1,1':2',1''-terphenyl]-4-carbonitrile-isomer-B
Step 1

tert-Butyl (1S,3R,4R)-rel-3-amino-7-azabicyclo[2.2.1]heptane-7-carboxylate (550 mg) was dissolved in THF (13.0 mL). At 0°C, TEA (0.720 mL) and 2,4-dinitrobenzene sulfonyl chloride (829 mg) were added thereto, followed by stirring at room temperature for 1 hour. Ethyl acetate was added thereto, the mixture was washed sequentially with water and saturated brine, and dried over anhydrous sodium sulfate, and the solvent was distilled off. The residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate) to give tert-butyl (1S,3R,4R)-rel-3-[(2,4-dinitrophenyl)sulfonylamino]-7-azabicyclo[2.2.1]heptane-7-carboxylate.

25

30

35

[0524]

Step 2

The tert-butyl (1S,3R,4R)-rel-3-[(2,4-dinitrophenyl)sulfonylamino]-7-azabicyclo[2.2.1]heptane-7-carboxylate (440 mg) obtained in step 1 above was subjected to chiral separation using SFC (device: Thar SFC prep 80 system, column: CHIRALPAK IE 20 x 250 mm, flow rate: 50 g/min, mobile phase: CO₂/MeOH = 90/10) to give (1S,3R,4R)-rel-3-[(2,4-dinitrophenyl)sulfonylamino]-7-azabicyclo[2.2.1]heptane-7-carboxylate-isomer-A (faster isomer) and (1S,3R,4R)-rel-3-[(2,4-dinitrophenyl)sulfonylamino]-7-azabicyclo[2.2.1]heptane-7-carboxylate-isomer-B (slower isomer).

[0525]

Each isomer was analyzed under the following HPLC conditions.

Column: CHIRALPAK IE 4.6 x 150 mm

Mobile phase: hexane (0.1% triethylamine)/ethanol = 85/15

Flow rate: 1.0 mL/min

Retention time of each isomer:

(1S,3R,4R)-rel-3-[(2,4-dinitrophenyl)sulfonylamino]-7-azabicyclo[2.2.1]heptane-7-carboxylate-isomer-A: 10.903 min (faster isomer)

(1S,3R,4R)-rel-3-[(2,4-dinitrophenyl)sulfonylamino]-7-azabicyclo[2.2.1]heptane-7-carboxylate-isomer-B: 14.028 min (slower isomer)

Step 3

The (1S,3R,4R)-rel-3-[(2,4-dinitrophenyl)sulfonylamino]-7-azabicyclo[2.2.1]heptane-7-carboxylate-isomer-B (200 mg) obtained in step 2 above was dissolved in ethyl acetate (1.00 mL). At room temperature, a 4 N hydrochloric acid-ethyl acetate solution (2.00 mL) was added thereto, followed by stirring at room temperature for 2 hours. The reaction mixture was vacuum-concentrated to give N-((1R,2R,4S)-rel-7-azabicyclo[2.2.1]heptan-2-yl)-2,4-dinitrobenzenesulfonamide-isomer-B hydrochloride.

[0526]

Step 4

The 3-(4-cyano-3-fluoro-phenyl)-4-[2-fluoro-4-(2-methoxyethyl)phenyl]benzoic acid (8 mg) obtained in step 2 of
5 Example 22 and the N-((1R,2R,4S)-rel-7-azabicyclo[2.2.1]heptan-2-yl)-2,4-dinitrobenzenesulfonamide-isomer-B hydrochloride (8.47 mg) obtained in step 3 above were dissolved in THF (0.30 mL). At room temperature, TEA (8.49 μ L) and HATU (15.5 mg) were added thereto, followed by stirring at 50°C for 1 hour. The solvent was
10 distilled off under reduced pressure. The residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate) to give N-[(1S,3R,4R)-rel-7-[3-(4-cyano-3-fluoro-phenyl)-4-[2-fluoro-4-(2-methoxyethyl)phenyl]benzoyl]-7-azabicyclo[2.2.1]heptan-3-yl]-2,4-dinitrobenzenesulfonamide-
15 isomer-B.

[0527]

Step 5

The N-[(1S,3R,4R)-rel-7-[3-(4-cyano-3-fluoro-phenyl)-4-[2-fluoro-4-(2-methoxyethyl)phenyl]benzoyl]-7-
20 azabicyclo[2.2.1]heptan-3-yl]-2,4-dinitrobenzenesulfonamide-isomer-B (14.5 mg) obtained in step 4 above was dissolved in DCM (1 mL). At 0°C, mercaptoacetic acid (2.83 μ L) and TEA (7.49 μ L) were added thereto, followed by stirring at room temperature for 2 hours. Chloroform was added thereto, the mixture was washed
25 with a 4 N aqueous sodium hydroxide solution, and dried over anhydrous sodium sulfate, and the solvent was distilled off. The residue was purified by reversed-phase HPLC (mobile phase: water/acetonitrile) to give the title compound.

[0528]

30 Example 156: Synthesis of 5'-((1R,2R,4S)-rel-2-amino-7-azabicyclo[2.2.1]heptane-7-carbonyl)-2'',3-difluoro-4'''-(2-methoxyethyl)-[1,1':2',1'''-terphenyl]-4-carbonitrile-isomer-A
Step 1

The (1S,3R,4R)-rel-3-[(2,4-
35 dinitrophenyl)sulfonylamino]-7-azabicyclo[2.2.1]heptane-7-

carboxylate-isomer-A (200 mg) obtained in step 2 of Example 155 was dissolved in ethyl acetate (1.00 mL). At room temperature, a 4 N hydrochloric acid-ethyl acetate solution (2.00 mL) was added thereto, followed by stirring at room temperature for 2 hours.

5 The reaction mixture was vacuum-concentrated to give N-((1R,2R,4S)-rel-7-azabicyclo[2.2.1]heptan-2-yl)-2,4-dinitrobenzenesulfonamide-isomer-A hydrochloride.

[0529]

Step 2

10 The 3-(4-cyano-3-fluoro-phenyl)-4-[2-fluoro-4-(2-methoxyethyl)phenyl]benzoic acid (8 mg) obtained in step 2 of Example 22 and the N-((1r,2r,4s)-rel-7-azabicyclo[2.2.1]heptan-2-yl)-2,4-dinitrobenzenesulfonamide-isomer-A hydrochloride (8.47 mg) obtained in step 1 above were dissolved in THF (0.30 mL). At
15 room temperature, TEA (8.49 μ L) and HATU (15.5 mg) were added thereto, followed by stirring at 50°C for 1 hour. The solvent was distilled off under reduced pressure. The residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate) to give N-[(1S,3R,4R)-rel-7-[3-(4-cyano-3-fluoro-
20 phenyl)-4-[2-fluoro-4-(2-methoxyethyl)phenyl]benzoyl]-7-azabicyclo[2.2.1]heptan-3-yl]-2,4-dinitrobenzenesulfonamide-isomer-A.

[0530]

Step 3

25 The N-[(1S,3R,4R)-rel-7-[3-(4-cyano-3-fluoro-phenyl)-4-[2-fluoro-4-(2-methoxyethyl)phenyl]benzoyl]-7-azabicyclo[2.2.1]heptan-3-yl]-2,4-dinitrobenzenesulfonamide-isomer-A (14.5 mg) obtained in step 2 above was dissolved in DCM (1 mL). At 0°C, mercaptoacetic acid (2.83 μ L) and TEA (7.49 μ L)
30 were added thereto, followed by stirring at room temperature for 2 hours. Chloroform was added thereto, the mixture was washed with a 4 N aqueous sodium hydroxide solution, and dried over anhydrous sodium sulfate, and the solvent was distilled off. The residue was purified by reversed-phase HPLC (mobile phase:
35 water/acetonitrile) to give the title compound.

[0531]

Example 157: Synthesis of 5'-((3-endo)-3-amino-8-azabicyclo[3.2.1]octane-8-carbonyl)-3-fluoro-2'-(3-methylimidazo[1,5-a]pyridin-7-yl)-[1,1'-biphenyl]-4-carbonitrile

5 The procedure of steps 1 to 5 in Example 41 was conducted using 7-bromo-3-methyl-imidazo[1,5-a]pyridine instead of 1-(4-bromo-3-fluoro-phenyl)-2-methyl-propan-2-ol to give the title compound.

[0532]

10 Example 158: Synthesis of 5'-((3-endo)-3-amino-8-azabicyclo[3.2.1]octane-8-carbonyl)-3-fluoro-2'-(3-methylpyrazolo[1,5-a]pyrimidin-6-yl)-[1,1'-biphenyl]-4-carbonitrile

15 The procedure of steps 1 to 5 in Example 41 was conducted using 6-bromo-3-methyl-pyrazolo[1,5-a]pyrimidine instead of 1-(4-bromo-3-fluoro-phenyl)-2-methyl-propan-2-ol to give the title compound.

[0533]

20 Example 159: Synthesis of 5'-((3-endo)-3-amino-8-azabicyclo[3.2.1]octane-8-carbonyl)-3-fluoro-2'-(6-fluoro-2-(2-hydroxy-2-methylpropyl)-2H-indazol-5-yl)-[1,1'-biphenyl]-4-carbonitrile

Step 1

25 5-Bromo-6-fluoro-1H-indazole (94 mg) was dissolved in DMF (1.5 mL). At room temperature, cesium carbonate (285 mg) and 2,2-dimethyloxirane (0.078 mL) were added thereto, followed by stirring at 90°C for 16 hours. The reaction was quenched with a saturated NH₄Cl aqueous solution, ethyl acetate was added thereto, and the resulting mixture was washed sequentially with water and saturated brine. After the organic layer was dried over anhydrous sodium sulfate, the solvent was distilled off. The residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate) to give 1-(5-bromo-6-fluoro-indazol-2-yl)-2-methyl-propan-2-ol.

35 [0534]

Step 2

The procedure of steps 1 to 5 in Example 41 was conducted using the 1-(5-bromo-6-fluoro-indazol-2-yl)-2-methyl-propan-2-ol obtained in step 1 above instead of 1-(4-bromo-3-fluoro-phenyl)-2-methyl-propan-2-ol to give the title compound.
[0535]

Example 160: Synthesis of 5'-((3-endo)-3-amino-8-azabicyclo[3.2.1]octane-8-carbonyl)-2'-(1-ethyl-6-fluoro-1H-benzo[d][1,2,3]triazol-5-yl)-3-fluoro-[1,1'-biphenyl]-4-carbonitrile

The procedure of steps 1 to 5 in Example 41 was conducted using 5-bromo-1-ethyl-6-fluoro-benzotriazole instead of 1-(4-bromo-3-fluoro-phenyl)-2-methyl-propan-2-ol to give the title compound.

[0536]

Example 161: Synthesis of 5'-((3-endo)-3-amino-8-azabicyclo[3.2.1]octane-8-carbonyl)-2'-(6,7-difluoro-1-(2-hydroxy-2-methylpropyl)-1H-benzo[d][1,2,3]triazol-5-yl)-3-fluoro-[1,1'-biphenyl]-4-carbonitrile

Step 1

1-(2,3-Difluoro-6-nitro-anilino)-2-methyl-propan-2-ol (6.20 g) was dissolved in DMF (84.0 mL). At room temperature, NBS (5.80 g) was added thereto, followed by stirring at 90°C for 1 hour. Ethyl acetate was added thereto, the mixture was washed sequentially with water and saturated brine, and dried over anhydrous sodium sulfate, and the solvent was distilled off. The residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate) to give 1-(4-bromo-2,3-difluoro-6-nitro-anilino)-2-methyl-propan-2-ol.

[0537]

Step 2

The 1-(4-bromo-2,3-difluoro-6-nitro-anilino)-2-methyl-propan-2-ol (5.67 g) obtained in step 1 above was dissolved in ethanol (87.2 mL). At room temperature, ammonium chloride (5.67 g), iron (5.67 g), and water (87.2 mL) were added thereto,

followed by stirring at 60°C overnight. The reaction mixture was passed through Celite and washed with ethyl acetate. The filtrate was vacuum-concentrated, ethyl acetate was added thereto, the mixture was washed sequentially with water and saturated brine, and dried over anhydrous sodium sulfate, and the solvent was distilled off. The residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate) to give 1-(6-amino-4-bromo-2,3-difluoro-anilino)-2-methyl-propan-2-ol.

[0538]

10 Step 3

The 1-(6-amino-4-bromo-2,3-difluoro-anilino)-2-methyl-propan-2-ol (4.36 g) obtained in step 2 above was dissolved in water (28.4 mL) and THF (28.4 mL). At 0°C, 12 N hydrochloric acid (28.4 mL) and sodium nitrite (1.80 g) were added thereto, followed by stirring at room temperature for 1 hour. Ethyl acetate was added thereto, the mixture was washed sequentially with water and saturated brine, and dried over anhydrous sodium sulfate, and the solvent was distilled off. The residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate) to give 1-(5-bromo-6,7-difluoro-benzotriazol-1-yl)-2-methyl-propan-2-ol.

[0539]

Step 4

The tert-butyl N-[(3-endo)-8-[3-(4-cyano-3-fluoro-phenyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoyl]-8-azabicyclo[3.2.1]octan-3-yl]carbamate (50 mg) obtained in step 3 of Example 41 and the 1-(5-bromo-6,7-difluoro-benzotriazol-1-yl)-2-methyl-propan-2-ol (39.9 mg) obtained in step 3 above were dissolved in 1,4-dioxane (0.50 mL). At room temperature, Pd(dba)₂ (2.50 mg), X-phos (4.14 mg), and tripotassium phosphate (55.3 mg) were added thereto, and the mixture was stirred in a microwave reactor at 125°C for 1 hour. The reaction mixture was filtrated, and the solvent was distilled off. The residue was purified by silica gel column chromatography (mobile phase: chloroform/methanol) to give tert-butyl N-[(3-endo)-8-[3-(4-

cyano-3-fluoro-phenyl)-4-[6,7-difluoro-1-(2-hydroxy-2-methylpropyl)benzotriazol-5-yl]benzoyl]-8-azabicyclo[3.2.1]octan-3-yl]carbamate.

[0540]

5 Step 5

The tert-butyl N-[(3-endo)-8-[3-(4-cyano-3-fluoro-phenyl)-4-[6,7-difluoro-1-(2-hydroxy-2-methylpropyl)benzotriazol-5-yl]benzoyl]-8-azabicyclo[3.2.1]octan-3-yl]carbamate (55.0 mg) obtained in step 4 above was dissolved in
10 MeOH (1.0 mL). At room temperature, 12 N hydrochloric acid (1.0 mL) was added thereto, followed by stirring at room temperature for 1 hour. A 2 N aqueous sodium hydroxide solution (6.00 mL) and chloroform were added thereto, the mixture was washed sequentially with water and saturated brine, and dried over
15 anhydrous sodium sulfate, and the solvent was distilled off. The residue was purified by reversed-phase HPLC (mobile phase: water/acetonitrile) to give the title compound.

[0541]

Example 162: Synthesis of (S)-5'-(3-aminopyrrolidine-1-carbonyl)-
20 2'-(6,7-difluoro-1-(2-hydroxy-2-methylpropyl)-1H-
benzo[d][1,2,3]triazol-5-yl)-3-fluoro-[1,1'-biphenyl]-4-
carbonitrile

Step 1

The tert-butyl N-[(3S)-1-[3-(4-cyano-3-fluoro-phenyl)-
25 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoyl]pyrrolidin-3-yl]carbamate (100 mg) obtained in step 1 of Example 37 and the 1-(5-bromo-6,7-difluoro-benzotriazol-1-yl)-2-methyl-propan-2-ol (85.8 mg) obtained in step 3 of Example 161 were dissolved in 1,4-dioxane (0.934 mL). At room temperature,
30 Pd(dba)₂ (5.37 mg), X-phos (8.90 mg), and tripotassium phosphate (119 mg) were added thereto, and the mixture was stirred in a microwave reactor at 125°C for 1 hour. The reaction mixture was filtrated, and the solvent was distilled off. The residue was purified by silica gel column chromatography (mobile phase:
35 chloroform/methanol) to give tert-butyl N-[(3S)-1-[3-(4-cyano-3-

fluoro-phenyl)-4-[6,7-difluoro-1-(2-hydroxy-2-methyl-propyl)benzotriazol-5-yl]benzoyl]pyrrolidin-3-yl]carbamate.

[0542]

Step 2

5 The tert-butyl N-[(3S)-1-[3-(4-cyano-3-fluoro-phenyl)-4-[6,7-difluoro-1-(2-hydroxy-2-methyl-propyl)benzotriazol-5-yl]benzoyl]pyrrolidin-3-yl]carbamate (99.8 mg) obtained in step 1 above was dissolved in MeOH (1.0 mL). At room temperature, 12 N hydrochloric acid (1.0 mL) was added thereto, followed by
10 stirring at room temperature for 1 hour. Then, a 2 N aqueous sodium hydroxide solution (6.00 mL) and chloroform were added thereto, the mixture was washed sequentially with water and saturated brine, and dried over anhydrous sodium sulfate, and the solvent was distilled off. The residue was purified by reversed-
15 phase HPLC (mobile phase: water/acetonitrile) to give the title compound.

[0543]

Example 163: Synthesis of 2'',3-difluoro-4''-(2-methoxyethyl)-5''-(piperazine-1-carbonyl)-[1,1':2',1''-terphenyl]-4-carbonitrile

20 The procedure of steps 1 to 4 in Example 22 was conducted using tert-butyl piperazine-1-carboxylate instead of tert-butyl N-[(3-endo)-8-azabicyclo[3.2.1]octan-3-yl]carbamate to give the title compound.

[0544]

25 Example 164: Synthesis of (R)-5''-(3-aminopiperidine-1-carbonyl)-2'',3-difluoro-4''-(2-methoxyethyl)-[1,1':2',1''-terphenyl]-4-carbonitrile

 The procedure of steps 1 to 4 in Example 22 was conducted using tert-butyl N-[(3R)-3-piperidyl]carbamate instead
30 of tert-butyl N-[(3-endo)-8-azabicyclo[3.2.1]octan-3-yl]carbamate to give the title compound.

[0545]

35 Example 165: Synthesis of 5''-(4-aminoazepan-1-carbonyl)-2'',3-difluoro-4''-(2-methoxyethyl)-[1,1':2',1''-terphenyl]-4-carbonitrile

The procedure of steps 1 to 4 in Example 22 was conducted using tert-butyl N-(azepan-4-yl)carbamate instead of tert-butyl N-[(3-endo)-8-azabicyclo[3.2.1]octan-3-yl]carbamate to give the title compound.

5 [0546]

Example 166: Synthesis of 5'-((1S,2S,4R)-rel-2-amino-7-azabicyclo[2.2.1]heptane-7-carbonyl)-2'',3-difluoro-4''-(2-hydroxy-2-methylpropyl)-[1,1':2',1''-terphenyl]-4-carbonitrile-isomer-B

10 Step 1

The tert-butyl 3-(4-cyano-3-fluoro-phenyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (500 mg) obtained in step 1 of Example 41 and the 1-(4-bromo-3-fluoro-phenyl)-2-methyl-propan-2-ol (379 mg) obtained in step 1 of Example 28 were
15 dissolved in 1,4-dioxane (5.9 mL). At room temperature, Pd(dba)₂ (68 mg), X-phos (113 mg), and tripotassium phosphate (752 mg) were added thereto, followed by stirring at 100°C overnight. The solvent was distilled off, and the residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate) to
20 give tert-butyl 3-(4-cyano-3-fluoro-phenyl)-4-[2-fluoro-4-(2-hydroxy-2-methyl-propyl)phenyl]benzoate.

[0547]

Step 2

The tert-butyl 3-(4-cyano-3-fluoro-phenyl)-4-[2-fluoro-
25 4-(2-hydroxy-2-methyl-propyl)phenyl]benzoate (300 mg) obtained in step 1 above was dissolved in THF (0.9 mL). At 0°C, 12 N hydrochloric acid (0.9 mL) was added thereto, followed by stirring at room temperature for 2 hours. Then, MTBE was added thereto, the mixture was washed with water and dried over
30 anhydrous sodium sulfate, and the solvent was distilled off to give 3-(4-cyano-3-fluoro-phenyl)-4-[2-fluoro-4-(2-hydroxy-2-methyl-propyl)phenyl]benzoic acid.

[0548]

Step 3

35 The 3-(4-cyano-3-fluoro-phenyl)-4-[2-fluoro-4-(2-

hydroxy-2-methyl-propyl)phenyl]benzoic acid (10 mg) obtained in step 2 above and the N-((1R,2R,4S)-rel-7-azabicyclo[2.2.1]heptan-2-yl)-2,4-dinitrobenzenesulfonamide-isomer-B hydrochloride (10.2 mg) obtained in step 3 of Example 155 were dissolved in THF (0.12 mL). At room temperature, TEA (0.014 mL) and HATU (18.7 mg) were added thereto, followed by stirring at 50°C for 1 hour. The reaction mixture was vacuum-concentrated, and the residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate) to give N-[(1S,3R,4R)-rel-7-[3-(4-cyano-3-fluoro-phenyl)-4-[2-fluoro-4-(2-hydroxy-2-methyl-propyl)phenyl]benzoyl]-7-azabicyclo[2.2.1]heptan-3-yl]-2,4-dinitrobenzenesulfonamide-isomer-B.

[0549]

Step 4

The N-[(1S,3R,4R)-rel-7-[3-(4-cyano-3-fluoro-phenyl)-4-[2-fluoro-4-(2-hydroxy-2-methyl-propyl)phenyl]benzoyl]-7-azabicyclo[2.2.1]heptan-3-yl]-2,4-dinitrobenzenesulfonamide-isomer-B (15 mg) obtained in step 3 above was dissolved in DCM (0.2 mL). At 0°C, mercaptoacetic acid (2 µL) and TEA (8.6 µL) were added thereto, followed by stirring at room temperature for 2 hours. Ethyl acetate was added thereto, and the mixture was washed sequentially with water and saturated brine. After the organic layer was dried over anhydrous sodium sulfate, the solvent was distilled off. The residue was purified by reversed-phase HPLC (mobile phase: water/acetonitrile) to give the title compound.

[0550]

Example 167: Synthesis of (S)-5'-(3-aminopyrrolidine-1-carbonyl)-2'-(6,7-difluoro-1-methyl-1H-benzo[d][1,2,3]triazol-5-yl)-3-fluoro-[1,1'-biphenyl]-4-carbonitrile

Step 1

The tert-butyl N-[(3S)-1-[3-(4-cyano-3-fluoro-phenyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoyl]pyrrolidin-3-yl]carbamate (60 mg) obtained in step 1 of Example 37 and 5-bromo-6,7-difluoro-1-methyl-benzotriazole

(41.7 mg) were dissolved in 1,4-dioxane (0.56 mL). At room temperature, Pd(dba)₂ (3.22 mg), X-phos (5.34 mg), and tripotassium phosphate (71.4 mg) were added thereto, and the mixture was stirred in a microwave reactor at 125°C for 1 hour.

5 The reaction mixture was filtrated, and the solvent was distilled off. The residue was purified by silica gel column chromatography (mobile phase: chloroform/methanol) to give tert-butyl N-[(3S)-1-[3-(4-cyano-3-fluoro-phenyl)-4-(6,7-difluoro-1-methyl-benzotriazol-5-yl)benzoyl]pyrrolidin-3-yl]carbamate.

10 [0551]

Step 2

The tert-butyl N-[(3S)-1-[3-(4-cyano-3-fluoro-phenyl)-4-(6,7-difluoro-1-methyl-benzotriazol-5-yl)benzoyl]pyrrolidin-3-yl]carbamate (20 mg) obtained in step 1 above was dissolved in 15 TFA (0.40 mL), followed by stirring at room temperature for 5 minutes. After the completion of the reaction was confirmed by LCMS, DMSO (1.60 mL) was added thereto, and purification was performed by reversed-phase HPLC (mobile phase: water/acetonitrile) to give the title compound.

20 [0552]

Example 168: Synthesis of 5'-((3-endo)-3-amino-8-azabicyclo[3.2.1]octane-8-carbonyl)-3-fluoro-2'-(6-fluoro-1-propyl-1H-benzo[d][1,2,3]triazol-5-yl)-[1,1'-biphenyl]-4-carbonitrile

25 The procedure of steps 1 to 5 in Example 41 was conducted using 5-bromo-6-fluoro-1-propyl-benzotriazole instead of 1-(4-bromo-3-fluoro-phenyl)-2-methyl-propan-2-ol to give the title compound.

[0553]

30 Example 169: Synthesis of (S)-5'-(3-aminopyrrolidine-1-carbonyl)-2'-(6,7-difluoro-1-(2-methoxyethyl)-1H-benzo[d][1,2,3]triazol-5-yl)-3-fluoro-[1,1'-biphenyl]-4-carbonitrile

Step 1

The procedure of steps 1 to 3 in Example 161 was 35 conducted using 2,3-difluoro-N-(2-methoxyethyl)-6-nitro-aniline

instead of 1-(2,3-difluoro-6-nitro-anilino)-2-methyl-propan-2-ol to give 5-bromo-6,7-difluoro-1-(2-methoxyethyl)benzotriazole.

[0554]

Step 2

5 The procedure of steps 1 to 3 in Example 37 was conducted using the 5-bromo-6,7-difluoro-1-(2-methoxyethyl)benzotriazole obtained in step 1 above instead of 1-(4-bromo-3-fluoro-phenyl)-2-methyl-propan-2-ol to give the title compound.

10 [0555]

Example 170: Synthesis of (S)-5'-(3-aminopyrrolidine-1-carbonyl)-3-fluoro-2'-(6-fluoro-1-(2-hydroxyethyl)-1H-indol-5-yl)-[1,1'-biphenyl]-4-carbonitrile

15 The procedure of steps 1 to 3 in Example 37 was conducted using 2-(5-bromo-6-fluoro-indol-1-yl)ethanol instead of 1-(4-bromo-3-fluoro-phenyl)-2-methyl-propan-2-ol to give the title compound.

[0556]

20 Example 171: Synthesis of 5'-((1R,2R,4S)-rel-2-amino-7-azabicyclo[2.2.1]heptane-7-carbonyl)-3-fluoro-2'-(6-fluoro-1-(2-hydroxy-2-methylpropyl)-1H-indol-5-yl)-[1,1'-biphenyl]-4-carbonitrile-isomer-B

Step 1

25 The tert-butyl 3-(4-cyano-3-fluoro-phenyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (2.3 g) obtained in step 1 of Example 41 and the 1-(5-bromo-6-fluoro-indol-1-yl)-2-methyl-propan-2-ol (2.02 g) obtained in step 1 of Example 136 were dissolved in 1,4-dioxane (18.1 mL). At room temperature, Pd(dba)₂ (250 mg), X-phos (414 mg), and tripotassium phosphate (3.46 g) were added thereto, followed by stirring at 100°C overnight. The solvent was distilled off, the residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate), and the solvent was distilled off. The residue was dissolved in THF (40.0 mL). At 0°C, 12 N hydrochloric acid (30.0 mL) was added thereto, followed by stirring at room

35

temperature for 2 hours. Then, MTBE was added thereto, the mixture was washed with water and dried over anhydrous sodium sulfate, and the solvent was distilled off to give 3-(4-cyano-3-fluoro-phenyl)-4-[6-fluoro-1-(2-hydroxy-2-methyl-propyl)indol-5-yl]benzoic acid.

[0557]

Step 2

The 3-(4-cyano-3-fluoro-phenyl)-4-[6-fluoro-1-(2-hydroxy-2-methyl-propyl)indol-5-yl]benzoic acid (8 mg) obtained in step 1 above and the N-((1R,2R,4S)-rel-7-azabicyclo[2.2.1]heptan-2-yl)-2,4-dinitrobenzenesulfonamide-isomer-B hydrochloride (7.47 mg) obtained in step 3 of Example 155 were dissolved in THF (0.30 mL). At room temperature, TEA (0.00748 mL) and HATU (13.6 mg) were added thereto, followed by stirring at 50°C for 1 hour. The reaction mixture was vacuum-concentrated, and the residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate) to give N-[(1S,3R,4R)-rel-7-[3-(4-cyano-3-fluoro-phenyl)-4-[6-fluoro-1-(2-hydroxy-2-methyl-propyl)indol-5-yl]benzoyl]-7-azabicyclo[2.2.1]heptan-3-yl]-2,4-dinitrobenzenesulfonamide-isomer-B.

[0558]

Step 3

The N-[(1S,3R,4R)-rel-7-[3-(4-cyano-3-fluoro-phenyl)-4-[6-fluoro-1-(2-hydroxy-2-methyl-propyl)indol-5-yl]benzoyl]-7-azabicyclo[2.2.1]heptan-3-yl]-2,4-dinitrobenzenesulfonamide-isomer-B (13.8 mg) obtained in step 2 above was dissolved in DCM (1.0 mL). At 0°C, mercaptoacetic acid (2.49 µL) and TEA (7.48 µL) were added thereto, followed by stirring at room temperature for 1 hour. Chloroform and 4 N sodium hydroxide were added thereto, and the mixture was washed sequentially with water and saturated brine. After the organic layer was dried over anhydrous sodium sulfate, the solvent was distilled off. The residue was purified by reversed-phase HPLC (mobile phase: water/acetonitrile) to give the title compound.

[0559]

Example 172: Synthesis of (S)-5'-(3-aminopyrrolidine-1-carbonyl)-2'-(6,7-difluoro-1-(2-hydroxy-2-methylpropyl)-1H-indol-5-yl)-3-fluoro-[1,1'-biphenyl]-4-carbonitrile

5 Step 1

5-Bromo-6,7-difluoro-1H-indole (300 mg) was dissolved in DMF (4.31 mL). At room temperature, Cs₂CO₃ (843 mg) and 2,2-dimethyloxirane (0.230 mL) were added thereto, followed by stirring at 80°C for 3 hours. The reaction mixture was filtrated,
10 and the solvent was distilled off. Ethyl acetate was added thereto, the mixture was washed sequentially with a saturated aqueous ammonium chloride solution, water, and saturated brine, and dried over anhydrous sodium sulfate, and the solvent was distilled off. The residue was purified by silica gel column
15 chromatography (mobile phase: hexane/ethyl acetate) to give 1-(5-bromo-6,7-difluoro-indol-1-yl)-2-methyl-propan-2-ol.

[0560]

Step 2

The tert-butyl 3-(4-cyano-3-fluoro-phenyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (210 mg) obtained in
20 step 1 of Example 41 and the 1-(5-bromo-6,7-difluoro-indol-1-yl)-2-methyl-propan-2-ol (196 mg) obtained in step 1 above were dissolved in 1,4-dioxane (1.65 mL). At room temperature, Pd(dba)₂ (22.8 mg), X-phos (37.8 mg), and tripotassium phosphate (316 mg)
25 were added thereto, followed by stirring at 100°C overnight. The solvent was distilled off, the residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate), and the solvent was distilled off. The residue was dissolved in THF (2.63 mL). At 0°C, 12 N hydrochloric acid (2.1 mL) was added
30 thereto, followed by stirring at room temperature for 2 hours. MTBE was added thereto, the mixture was washed with water and dried over anhydrous sodium sulfate, and the solvent was distilled off to give 3-(4-cyano-3-fluoro-phenyl)-4-[6,7-difluoro-1-(2-hydroxy-2-methyl-propyl)indol-5-yl]benzoic acid.

35 [0561]

Step 3

The 3-(4-cyano-3-fluoro-phenyl)-4-[6,7-difluoro-1-(2-hydroxy-2-methyl-propyl)indol-5-yl]benzoic acid (30 mg) obtained in step 2 above and tert-butyl N-[(3S)-pyrrolidin-3-yl]carbamate (13.2 mg) were dissolved in THF (0.323 mL). At room temperature, TEA (0.027 mL) and HATU (49.1 mg) were added thereto, followed by stirring at 50°C for 1 hour. The solvent was distilled off under reduced pressure. The residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate) to give tert-butyl N-[(3S)-1-[3-(4-cyano-3-fluoro-phenyl)-4-[6,7-difluoro-1-(2-hydroxy-2-methyl-propyl)indol-5-yl]benzoyl]pyrrolidin-3-yl]carbamate.
[0562]

Step 4

The tert-butyl N-[(3S)-1-[3-(4-cyano-3-fluoro-phenyl)-4-[6,7-difluoro-1-(2-hydroxy-2-methyl-propyl)indol-5-yl]benzoyl]pyrrolidin-3-yl]carbamate (40 mg) obtained in step 3 above was dissolved in MeOH (0.80 mL). At room temperature, a 4 N hydrochloric acid-1,4-dioxane solution (0.80 mL) was added thereto, followed by stirring at room temperature for 1 hour. Chloroform and a 2 N aqueous sodium hydroxide solution (1.6 mL) were added thereto, the mixture was washed with water and dried over anhydrous sodium sulfate, and the solvent was distilled off. The residue was purified by reversed-phase HPLC (mobile phase: water/acetonitrile) to give the title compound.
[0563]

Example 173: Synthesis of 5'-(7-amino-2-azabicyclo[2.2.1]heptane-2-carbonyl)-3-fluoro-2'-(6-fluoro-1-(2-hydroxy-2-methylpropyl)-1H-indol-5-yl)-[1,1'-biphenyl]-4-carbonitrile

The 3-(4-cyano-3-fluoro-phenyl)-4-[6-fluoro-1-(2-hydroxy-2-methyl-propyl)indol-5-yl]benzoic acid (8 mg) obtained in step 1 of Example 171 was dissolved in THF (0.3 mL). At room temperature, tert-butyl N-(3-azabicyclo[2.2.1]heptan-7-yl)carbamate (3.80 mg), TEA (0.0075 mL), and HATU (13.6 mg) were added thereto, followed by stirring at 50°C for 3 hours. After

the completion of the reaction was confirmed by LCMS, the reaction mixture was concentrated. TFA (0.20 mL) was added to the residue, followed by stirring at room temperature for 5 minutes. After the completion of the reaction was confirmed by LCMS, DMSO (0.8 mL) was added to the reaction mixture, and purification was performed by reversed-phase HPLC (mobile phase: water/acetonitrile) to give the title compound.

[0564]

Example 174: Synthesis of 5'-(7-amino-2-azabicyclo[2.2.1]heptane-2-carbonyl)-2'',3-difluoro-4''-(2-hydroxy-2-methylpropyl)-[1,1':2',1''-terphenyl]-4-carbonitrile

Step 1

The 3-(4-cyano-3-fluoro-phenyl)-4-[2-fluoro-4-(2-hydroxy-2-methyl-propyl)phenyl]benzoic acid (100 mg) obtained in step 2 of Example 166 was dissolved in THF (0.982 mL). At room temperature, tert-butyl N-(3-azabicyclo[2.2.1]heptan-7-yl)carbamate (52.1 mg), TEA (0.103 mL), and HATU (187 mg) were added thereto, followed by stirring at 50°C for 3 hours. The residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate) to give tert-butyl N-[3-[3-(4-cyano-3-fluoro-phenyl)-4-[2-fluoro-4-(2-hydroxy-2-methyl-propyl)phenyl]benzoyl]-3-azabicyclo[2.2.1]heptan-7-yl]carbamate.

[0565]

Step 2

The tert-butyl N-[3-[3-(4-cyano-3-fluoro-phenyl)-4-[2-fluoro-4-(2-hydroxy-2-methyl-propyl)phenyl]benzoyl]-3-azabicyclo[2.2.1]heptan-7-yl]carbamate (30 mg) obtained in step 1 above was dissolved in MeOH (0.5 mL). At room temperature, 12 N hydrochloric acid (0.5 mL) was added thereto. After the mixture was stirred at room temperature for 0.5 hour, water and a 2 N aqueous sodium hydroxide solution (3.0 mL) were added thereto. The mixture was extracted with chloroform, and the solvent was distilled off. The residue was purified by reversed-phase HPLC (mobile phase: water/acetonitrile) to give the title compound.

[0566]

Example 175: Synthesis of 5'-((1S,2S,4R)-rel-2-amino-7-azabicyclo[2.2.1]heptane-7-carbonyl)-2'-(6,7-difluoro-1-methyl-1H-benzo[d][1,2,3]triazol-5-yl)-3-fluoro-[1,1'-biphenyl]-4-carbonitrile-isomer-B

5 Step 1

The tert-butyl 3-(4-cyano-3-fluoro-phenyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (90 mg) obtained in step 1 of Example 41 and 5-bromo-6,7-difluoro-1-methyl-benzotriazole (68.6 mg) were dissolved in 1,4-dioxane (0.71 mL).
10 At room temperature, Pd(dba)₂ (9.8 mg), X-phos (16 mg), and tripotassium phosphate (135 mg) were added thereto, followed by stirring at 100°C overnight. The solvent was distilled off, the residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate), and the solvent was distilled off.
15 The residue was dissolved in TFA (1.0 mL), followed by stirring at room temperature for 2 hours. MTBE was added thereto, the mixture was washed with water and dried over anhydrous sodium sulfate, and the solvent was distilled off to give 3-(4-cyano-3-fluoro-phenyl)-4-(6,7-difluoro-1-methyl-benzotriazol-5-yl)benzoic
20 acid.
[0567]

Step 2

The 3-(4-cyano-3-fluoro-phenyl)-4-(6,7-difluoro-1-methyl-benzotriazol-5-yl)benzoic acid (30 mg) obtained in step 1
25 above and the N-((1R,2R,4S)-rel-7-azabicyclo[2.2.1]heptan-2-yl)-2,4-dinitrobenzenesulfonamide-isomer-B hydrochloride (30.6 mg) obtained in step 3 of Example 155 were dissolved in THF (0.367 mL). At room temperature, TEA (0.042 mL) and HATU (55.9 mg) were added thereto, followed by stirring at 50°C for 1 hour. The
30 reaction mixture was vacuum-concentrated, and the residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate) to give N-[(1R,3S,4S)-rel-7-[3-(4-cyano-3-fluoro-phenyl)-4-(6,7-difluoro-1-methyl-benzotriazol-5-yl)benzoyl]-7-azabicyclo[2.2.1]heptan-3-yl]-2,4-
35 dinitrobenzenesulfonamide-isomer-B.

[0568]

Step 3

The N-[(1R,3S,4S)-rel-7-[3-(4-cyano-3-fluoro-phenyl)-4-(6,7-difluoro-1-methyl-benzotriazol-5-yl)benzoyl]-7-azabicyclo[2.2.1]heptan-3-yl]-2,4-dinitrobenzenesulfonamide-isomer-B (51 mg) obtained in step 2 above was dissolved in DCM (0.70 mL). At 0°C, mercaptoacetic acid (5.8 µL) and TEA (29.1 µL) were added thereto, followed by stirring at room temperature for 2 hours. Ethyl acetate was added thereto, and the mixture was washed sequentially with water and saturated brine. After the organic layer was dried over anhydrous sodium sulfate, the solvent was distilled off. The residue was purified by reversed-phase HPLC (mobile phase: water/acetonitrile) to give the title compound.

15 [0569]

Example 176: Synthesis of 5'-((1S,2S,4R)-rel-2-amino-7-azabicyclo[2.2.1]heptane-7-carbonyl)-3-fluoro-2'-(6-fluoro-1-(2-hydroxy-2-methylpropyl)-1H-indazol-5-yl)-[1,1'-biphenyl]-4-carbonitrile-isomer-B

20 Step 1

The tert-butyl 3-(4-cyano-3-fluoro-phenyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (2.3 g) obtained in step 1 of Example 41 and the 1-(5-bromo-6-fluoro-1H-indazol-1-yl)-2-methylpropan-2-ol (2.03 g) obtained in step 1 of Example 142 were dissolved in 1,4-dioxane (18.7 mL). At room temperature, Pd(dba)₂ (250 mg), X-phos (414 mg), and tripotassium phosphate (3.46 g) were added thereto, followed by stirring at 100°C overnight. The solvent was distilled off, the residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate), and the solvent was distilled off. The residue was dissolved in THF (10.0 mL). At 0°C, 12 N hydrochloric acid (10.0 mL) was added thereto, followed by stirring at room temperature for 2 hours. MTBE was added thereto, the mixture was washed with water and dried over anhydrous sodium sulfate, and the solvent was distilled off to give 3-(4-cyano-3-fluoro-

phenyl)-4-[6-fluoro-1-(2-hydroxy-2-methyl-propyl)indazol-5-yl]benzoic acid.

[0570]

Step 2

5 The 3-(4-cyano-3-fluoro-phenyl)-4-[6-fluoro-1-(2-hydroxy-2-methyl-propyl)indazol-5-yl]benzoic acid (30 mg) obtained in step 1 above and the N-((1R,2R,4S)-rel-7-azabicyclo[2.2.1]heptan-2-yl)-2,4-dinitrobenzenesulfonamide-isomer-B hydrochloride (27.9 mg) obtained in step 3 of Example
10 155 were dissolved in THF (0.34 mL). At room temperature, TEA (0.038 mL) and HATU (51.0 mg) were added thereto, followed by stirring at 50°C for 1 hour. The reaction mixture was vacuum-concentrated, and the residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate) to give N-
15 [(1S,3R,4R)-rel-7-[3-(4-cyano-3-fluoro-phenyl)-4-[6-fluoro-1-(2-hydroxy-2-methyl-propyl)indazol-5-yl]benzoyl]-7-azabicyclo[2.2.1]heptan-3-yl]-2,4-dinitrobenzenesulfonamide-isomer-B.

[0571]

20 Step 3

 The N-[(1S,3R,4R)-rel-7-[3-(4-cyano-3-fluoro-phenyl)-4-[6-fluoro-1-(2-hydroxy-2-methyl-propyl)indazol-5-yl]benzoyl]-7-azabicyclo[2.2.1]heptan-3-yl]-2,4-dinitrobenzenesulfonamide-isomer-B (45 mg) obtained in step 2 above was dissolved in DCM
25 (0.58 mL). At 0°C, mercaptoacetic acid (4.9 µL) and TEA (24 µL) were added thereto, followed by stirring at room temperature for 2 hours. Ethyl acetate was added thereto, and the mixture was washed sequentially with water and saturated brine. After the organic layer was dried over anhydrous sodium sulfate, the
30 solvent was distilled off. The residue was purified by reversed-phase HPLC (mobile phase: water/acetonitrile) to give the title compound.

[0572]

Example 177: Synthesis of 5'-((1S,2S,4R)-rel-2-amino-7-azabicyclo[2.2.1]heptane-7-carbonyl)-2'-(6,7-difluoro-1-(2-
35

hydroxy-2-methylpropyl)-1H-benzo[d][1,2,3]triazol-5-yl)-3-fluoro-
[1,1'-biphenyl]-4-carbonitrile-isomer-B

Step 1

The tert-butyl 3-(4-cyano-3-fluoro-phenyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (3.2 g) obtained in
5 step 1 of Example 41 and the 1-(5-bromo-6,7-difluoro-benzotriazol-1-yl)-2-methyl-propan-2-ol (3.01 g) obtained in step 3 of Example 161 were dissolved in 1,4-dioxane (25.2 mL). At room temperature, Pd(dba)₂ (348 mg), X-phos (577 mg), and tripotassium
10 phosphate (4.81 g) were added thereto, followed by stirring at 100°C overnight. The solvent was distilled off, the residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate), and the solvent was distilled off. The residue was dissolved in THF (15.0 mL). At 0°C, 12 N hydrochloric
15 acid (15.0 mL) was added thereto, followed by stirring at room temperature for 2 hours. MTBE was added thereto, the mixture was washed with water and dried over anhydrous sodium sulfate, and the solvent was distilled off to give 3-(4-cyano-3-fluoro-phenyl)-4-[6,7-difluoro-1-(2-hydroxy-2-methyl-propyl)benzotriazol-5-yl]benzoic acid.
20 [0573]

Step 2

The 3-(4-cyano-3-fluoro-phenyl)-4-[6,7-difluoro-1-(2-hydroxy-2-methyl-propyl)benzotriazol-5-yl]benzoic acid (30 mg)
25 obtained in step 1 above and the N-((1S,2S,4R)-rel-7-azabicyclo[2.2.1]heptan-2-yl)-2,4-dinitrobenzenesulfonamide-isomer-B hydrochloride (26.8 mg) obtained in step 3 of Example 155 were dissolved in THF (0.33 mL). At room temperature, TEA (0.037 mL) and HATU (48.9 mg) were added thereto, followed by
30 stirring at 50°C for 1 hour. The reaction mixture was vacuum-concentrated, and the residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate) to give N-[(1S,2S,4R)-rel-7-[3-(4-cyano-3-fluoro-phenyl)-4-[6,7-difluoro-1-(2-hydroxy-2-methyl-propyl)benzotriazol-5-yl]benzoyl]-7-azabicyclo[2.2.1]heptan-3-yl]-2,4-dinitrobenzenesulfonamide-
35

isomer-B.

[0574]

Step 3

The N-[(1S,2S,4R)-rel-7-[3-(4-cyano-3-fluoro-phenyl)-4-
5 [6,7-difluoro-1-(2-hydroxy-2-methyl-propyl)benzotriazol-5-
yl]benzoyl]-7-azabicyclo[2.2.1]heptan-3-yl]-2,4-
dinitrobenzenesulfonamide-isomer-B (43 mg) obtained in step 2
above was dissolved in DCM (0.54 mL). At 0°C, mercaptoacetic acid
(4.5 µL) and TEA (22.7 µL) were added thereto, followed by
10 stirring at room temperature for 2 hours. Ethyl acetate was added
thereto, and the mixture was washed sequentially with water and
saturated brine. After the organic layer was dried over anhydrous
sodium sulfate, the solvent was distilled off. The residue was
purified by reversed-phase HPLC (mobile phase:
15 water/acetonitrile) to give the title compound.

[0575]

Example 178: Synthesis of 5'-((3-endo)-3-amino-8-
azabicyclo[3.2.1]octane-8-carbonyl)-2'-(3-bromo-6-fluoro-1-(2-
hydroxy-2-methylpropyl)-1H-indazol-5-yl)-3-fluoro-[1,1'-
20 biphenyl]-4-carbonitrile

Step 1

The procedure of steps 1 to 4 in Example 41 was
conducted using the 1-(5-bromo-6-fluoro-indazol-1-yl)-2-methyl-
propan-2-ol obtained in step 1 of Example 142 instead of 1-(4-
25 bromo-3-fluoro-phenyl)-2-methyl-propan-2-ol to give tert-butyl N-
((3-endo)-8-(4'-cyano-3'-fluoro-6-(6-fluoro-1-(2-hydroxy-2-
methylpropyl)-1H-indazol-5-yl)-[1,1'-biphenyl]-3-carbonyl)-8-
azabicyclo[3.2.1]octan-3-yl)carbamate.

[0576]

30 Step 2

The tert-butyl N-((3-endo)-8-(4'-cyano-3'-fluoro-6-(6-
fluoro-1-(2-hydroxy-2-methylpropyl)-1H-indazol-5-yl)-[1,1'-
biphenyl]-3-carbonyl)-8-azabicyclo[3.2.1]octan-3-yl)carbamate (10
mg) obtained in step 1 above was dissolved in DMF (0.076 mL). NBS
35 (3.5 mg) was added thereto, followed by stirring at 80°C

overnight. The reaction mixture was diluted with DMSO to 1 mL, and purification was performed by reversed-phase HPLC (mobile phase: water/acetonitrile) to give the title compound.

[0577]

5 Example 179: Synthesis of 5'-((3-endo)-3-amino-8-azabicyclo[3.2.1]octane-8-carbonyl)-3-fluoro-2'-(5-fluoro-3-methylbenzo[d]isoxazol-6-yl)-[1,1'-biphenyl]-4-carbonitrile

Step 1

3-Bromo-4-fluorophenol (5 g) was dissolved in
10 dichloromethane (114 mL). At 0°C, TEA (5.5 mL) was added thereto, and acetyl chloride (2.8 mL) was added thereto dropwise. The reaction mixture was stirred at 20°C for 30 minutes and diluted with dichloromethane (100 mL). The resulting product was washed with 0.5 N hydrochloric acid, a saturated aqueous sodium hydrogen
15 carbonate solution, and saturated brine, and the solvent was distilled off to give 3-bromo-4-fluorophenyl acetate.

[0578]

Step 2

A boron trifluoride-acetic acid complex (53 mL) was
20 added to the 3-bromo-4-fluorophenyl acetate (6.2 g) obtained in step 1 above, followed by stirring at 155°C for 14 hours. The reaction mixture was cooled to 0°C, and ice was added thereto. The precipitate was collected by filtration, washed with water at 0°C, and dried. The obtained solid was purified by silica gel
25 column chromatography (mobile phase: hexane/ethyl acetate) to give 1-(4-bromo-5-fluoro-2-hydroxyphenyl)ethanone.

[0579]

Step 3

MeOH (30 mL) was added to the 1-(4-bromo-5-fluoro-2-
30 hydroxyphenyl)ethanone (2.16 g) obtained in step 2 above, hydroxylamine hydrochloride (1.29 g), and sodium acetate (1.14 g), followed by stirring at 60°C for 1 hour. Ice water was added to the reaction mixture, and the precipitate was collected by filtration, washed with water, and dried. The obtained solid was
35 dissolved in THF (31 mL), and TEA (1.68 mL) and N,N'-

carbonyldiimidazole (1.65 g) were added thereto, followed by stirring at 70°C for 1 hour. The solvent was distilled off, and the residue was purified by silica gel column chromatography (mobile phase: chloroform/ethyl acetate) to give 6-bromo-5-fluoro-3-methylbenzo[d]isoxazol.

[0580]

Step 4

The procedure of steps 1 to 5 in Example 41 was conducted using the 6-bromo-5-fluoro-3-methylbenzo[d]isoxazol obtained in step 3 above instead of 1-(4-bromo-3-fluoro-phenyl)-2-methyl-propan-2-ol to give the title compound.

[0581]

Example 180: Synthesis of (S)-5'-(3-aminopyrrolidine-1-carbonyl)-3-fluoro-2'-(5-fluoro-3-methylbenzo[d]isoxazol-6-yl)-[1,1'-biphenyl]-4-carbonitrile

The procedure of steps 1 to 3 in Example 37 was conducted using the 6-bromo-5-fluoro-3-methylbenzo[d]isoxazol obtained in step 3 of Example 179 instead of 1-(4-bromo-3-fluoro-phenyl)-2-methyl-propan-2-ol to give the title compound.

[0582]

Example 181: Synthesis of 5'-((3-endo)-3-amino-8-azabicyclo[3.2.1]octane-8-carbonyl)-3-fluoro-2'-(6-fluoro-1-(2-hydroxy-2-methylpropyl)-3-methyl-1H-indazol-5-yl)-[1,1'-biphenyl]-4-carbonitrile

Step 1

The tert-butyl N-((3-endo)-8-(4'-cyano-3'-fluoro-6-(6-fluoro-1-(2-hydroxy-2-methylpropyl)-1H-indazol-5-yl)-[1,1'-biphenyl]-3-carbonyl)-8-azabicyclo[3.2.1]octan-3-yl)carbamate (69 mg) obtained in step 1 of Example 178 was dissolved in DMF (0.53 mL), and NBS (38 mg) was added thereto, followed by stirring at 80°C overnight. The mixture was cooled to room temperature, and Boc₂O (200 mg) and DMAP (1 mg) were added thereto, followed by stirring at room temperature for 2 hours. Ethyl acetate was added thereto, and the resulting mixture was washed sequentially with water and saturated brine. After the organic layer was dried over

anhydrous sodium sulfate, the solvent was distilled off. The solvent was distilled off, and the residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate) to give tert-butyl ((3-endo)-8-(4'-cyano-3'-fluoro-6-(6-fluoro-1-(2-hydroxy-2-methylpropyl)-1H-indazol-5-yl)-[1,1'-biphenyl]-3-carbonyl)-8-azabicyclo[3.2.1]octan-3-yl)carbamate.

[0583]

Step 2

The tert-butyl ((3-endo)-8-(4'-cyano-3'-fluoro-6-(6-fluoro-1-(2-hydroxy-2-methylpropyl)-1H-indazol-5-yl)-[1,1'-biphenyl]-3-carbonyl)-8-azabicyclo[3.2.1]octan-3-yl)carbamate (15 mg) obtained in step 1 above, trimethylboroxine (7.7 mg), PdCl₂(dppf)CH₂Cl₂ (1 mg), and cesium carbonate (20 mg) were suspended in 1,4-dioxane, followed by stirring at 125°C for 30 minutes under microwave irradiation. The solvent was distilled off, and trifluoroacetic acid (0.2 mL) was added to the residue, followed by stirring at room temperature for 10 minutes. The reaction mixture was diluted with DMSO to 1 mL, and purification was performed by reversed-phase HPLC (mobile phase: water/acetonitrile) to give the title compound.

[0584]

Example 182: Synthesis of 5'-((1R,2S,4S)-rel-2-amino-7-azabicyclo[2.2.1]heptane-7-carbonyl)-2'',3-difluoro-4''-(2-hydroxy-2-methylpropyl)-[1,1':2',1''-terphenyl]-4-carbonitrile

Step 1

tert-Butyl (1S,3S,4R)-rel-3-amino-7-azabicyclo[2.2.1]heptane-7-carboxylate (919 mg) was dissolved in THF (14.4 mL). At 0°C, TEA (1.81 mL) and 2,4-dinitrobenzenesulfonyl chloride (1.73 g) were added thereto, followed by stirring at room temperature overnight. Ethyl acetate was added thereto, the mixture was washed sequentially with water and saturated brine, and dried over anhydrous sodium sulfate, and the solvent was distilled off. The residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate) to give tert-butyl (1S,3S,4R)-rel-3-[(2,4-

dinitrophenyl)sulfonylamino]-7-azabicyclo[2.2.1]heptane-7-carboxylate.

[0585]

Step 2

5 The tert-butyl (1S,3S,4R)-rel-3-[(2,4-dinitrophenyl)sulfonylamino]-7-azabicyclo[2.2.1]heptane-7-carboxylate (100 mg) obtained in step 1 above was dissolved in ethyl acetate (1.00 mL). At room temperature, a 4 N hydrochloric acid-ethyl acetate solution (2.00 mL) was added thereto, followed
10 by stirring at room temperature for 1 hour. The reaction mixture was vacuum-concentrated to give N-[(1S,3S,4R)-rel-7-azabicyclo[2.2.1]heptan-3-yl]-2,4-dinitrobenzenesulfonamide hydrochloride.

[0586]

15 Step 3

 The 3-(4-cyano-3-fluoro-phenyl)-4-[2-fluoro-4-(2-hydroxy-2-methyl-propyl)phenyl]benzoic acid (30 mg) obtained in step 2 of Example 166 and the N-[(1S,3S,4R)-rel-7-azabicyclo[2.2.1]heptan-3-yl]-2,4-dinitrobenzenesulfonamide
20 hydrochloride (30.7 mg) obtained in step 2 above were dissolved in THF (0.40 mL). At room temperature, TEA (0.0420 mL) and HATU (56.0 mg) were added thereto, followed by stirring at 50°C for 1 hour. The reaction mixture was vacuum-concentrated, and the residue was purified by silica gel column chromatography (mobile
25 phase: hexane/ethyl acetate) to give N-[(1S,3S,4R)-rel-7-[3-(4-cyano-3-fluoro-phenyl)-4-[2-fluoro-4-(2-hydroxy-2-methyl-propyl)phenyl]benzoyl]-7-azabicyclo[2.2.1]heptan-3-yl]-2,4-dinitrobenzenesulfonamide.

[0587]

30 Step 4

 The N-[(1S,3S,4R)-rel-7-[3-(4-cyano-3-fluoro-phenyl)-4-[2-fluoro-4-(2-hydroxy-2-methyl-propyl)phenyl]benzoyl]-7-azabicyclo[2.2.1]heptan-3-yl]-2,4-dinitrobenzenesulfonamide (55 mg) obtained in step 3 above was dissolved in DCM (0.752 mL). At
35 0°C, mercaptoacetic acid (6.27 µL) and TEA (31.4 µL) were added

thereto, followed by stirring at room temperature for 2 hours. Chloroform was added thereto, the mixture was washed with a 4 N aqueous sodium hydroxide solution, and dried over anhydrous sodium sulfate, and the solvent was distilled off. The residue
5 was purified by reversed-phase HPLC (mobile phase: water/acetonitrile) to give the title compound.

[0588]

Example 183: Synthesis of 5'-((3-endo)-3-amino-8-azabicyclo[3.2.1]octane-8-carbonyl)-2'-(6,7-difluoro-1-(2-hydroxy-2-methylpropyl)-1H-indol-5-yl)-3-fluoro-[1,1'-biphenyl]-4-carbonitrile

Step 1

The 3-(4-cyano-3-fluoro-phenyl)-4-[6,7-difluoro-1-(2-hydroxy-2-methyl-propyl)indol-5-yl]benzoic acid (30 mg) obtained
15 in step 2 of Example 172 and tert-butyl N-[(3-endo)-8-azabicyclo[3.2.1]octan-3-yl]carbamate (16.1 mg) were dissolved in THF (0.323 mL). At room temperature, TEA (0.027 mL) and HATU (49.1 mg) were added thereto, followed by stirring at 50°C for 1 hour. The solvent was distilled off under reduced pressure. The
20 residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate) to give tert-butyl N-[(3-endo)-8-[3-(4-cyano-3-fluoro-phenyl)-4-[6,7-difluoro-1-(2-hydroxy-2-methyl-propyl)indol-5-yl]benzoyl]-8-azabicyclo[3.2.1]octan-3-yl]carbamate.

[0589]

Step 2

The tert-butyl N-[(3-endo)-8-[3-(4-cyano-3-fluoro-phenyl)-4-[6,7-difluoro-1-(2-hydroxy-2-methyl-propyl)indol-5-yl]benzoyl]-8-azabicyclo[3.2.1]octan-3-yl]carbamate (42 mg)
30 obtained in step 1 above was dissolved in MeOH (0.84 mL). At room temperature, a 4 N hydrochloric acid-1,4-dioxane solution (0.84 mL) was added thereto, followed by stirring at room temperature for 1 hour. Chloroform and a 2 N aqueous sodium hydroxide solution (1.68 mL) were added thereto, the mixture was washed
35 with water and dried over anhydrous sodium sulfate, and the

solvent was distilled off. The residue was purified by reversed-phase HPLC (mobile phase: water/acetonitrile) to give the title compound.

[0590]

5 Example 184: Synthesis of 5'-((1R,2S,4S)-rel-2-amino-7-azabicyclo[2.2.1]heptane-7-carbonyl)-3-fluoro-2'-(6-fluoro-1-(2-hydroxy-2-methylpropyl)-1H-indol-5-yl)-[1,1'-biphenyl]-4-carbonitrile

10 The procedure of steps 1 to 3 in Example 171 was conducted using the N-[(1R,2S,4S)-rel-7-azabicyclo[2.2.1]heptan-3-yl]-2,4-dinitrobenzenesulfonamide hydrochloride obtained in step 2 of Example 182 instead of N-((1R,2R,4S)-rel-7-azabicyclo[2.2.1]heptan-2-yl)-2,4-dinitrobenzenesulfonamide-isomer-B hydrochloride to give the title compound.

15 [0591]

Example 185: Synthesis of 5'-((1S,2S,4R)-rel-2-amino-7-azabicyclo[2.2.1]heptane-7-carbonyl)-2'-(6,7-difluoro-1-methyl-1H-benzo[d][1,2,3]triazol-5-yl)-3-fluoro-[1,1'-biphenyl]-4-carbonitrile

20 The procedure of steps 3 to 4 in Example 172 was conducted using the 3-(4-cyano-3-fluoro-phenyl)-4-(6,7-difluoro-1-methyl-benzotriazol-5-yl)benzoic acid obtained in step 1 of Example 175 instead of 3-(4-cyano-3-fluoro-phenyl)-4-[6,7-difluoro-1-(2-hydroxy-2-methyl-propyl)indol-5-yl]benzoic acid, and using tert-butyl ((1S,2S,4R)-rel-7-azabicyclo[2.2.1]heptan-2-yl)carbamate hydrochloride instead of tert-butyl N-[(3S)-pyrrolidin-3-yl]carbamate to give the title compound.

[0592]

30 Example 186: Synthesis of 5'-((3-endo)-3-amino-8-azabicyclo[3.2.1]octane-8-carbonyl)-3-fluoro-2'-(6-fluoro-1-(2-hydroxy-2-methylpropyl)-7-methoxy-1H-indazol-5-yl)-[1,1'-biphenyl]-4-carbonitrile

Step 1

35 5-Bromo-2,3,4-trifluoro-benzaldehyde (480 mg) was dissolved in 1,2-dimethoxyethane (4.8 mL). At room temperature,

hydrazine monohydrate (7.68 mL) was added thereto, followed by stirring at 80°C for 5 hours. Ethyl acetate was added thereto, the mixture was washed sequentially with water and saturated brine, and dried over anhydrous sodium sulfate, and the solvent
5 was distilled off. The residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate) to give 5-bromo-6,7-difluoro-1H-indazole.

[0593]

Step 2

10 The 5-bromo-6,7-difluoro-1H-indazole (97 mg) obtained in step 1 above was dissolved in DMF (1.38 mL). At room temperature, methanol (0.1 mL), cesium carbonate (271 mg), and 2,2-dimethyloxirane (0.074 mL) were added thereto, followed by stirring at 80°C for 1 hour. Ethyl acetate was added thereto, the
15 mixture was washed sequentially with water and saturated brine, and dried over anhydrous sodium sulfate, and the solvent was distilled off. The residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate) to give 1-(5-bromo-6-fluoro-7-methoxy-indazol-1-yl)-2-methyl-propan-2-ol.

20 [0594]

Step 3

 The tert-butyl 3-(4-cyano-3-fluoro-phenyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (45.0 mg) obtained in step 1 of Example 41 and the 1-(5-bromo-6-fluoro-7-methoxy-
25 indazol-1-yl)-2-methyl-propan-2-ol (43.8 mg) obtained in step 2 above were dissolved in 1,4-dioxane (0.50 mL). At room temperature, Pd(dba)₂ (4.89 mg), X-phos (8.11 mg), and tripotassium phosphate (67.7 mg) were added thereto, followed by stirring at 100°C overnight. The solvent was distilled off, the
30 residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate), and the solvent was distilled off. The residue was dissolved in THF (0.45 mL). At 0°C, 12 N hydrochloric acid (0.56 mL) was added thereto, followed by stirring at room temperature for 2 hours. MTBE was added thereto,
35 and the mixture was washed with water and dried over anhydrous

sodium sulfate. Thereafter, the solvent was distilled off to give 3-(4-cyano-3-fluoro-phenyl)-4-[6-fluoro-1-(2-hydroxy-2-methyl-propyl)-7-methoxy-indazol-5-yl]benzoic acid.

[0595]

5 Step 4

The procedure of steps 3 to 4 in Example 172 was conducted using the 3-(4-cyano-3-fluoro-phenyl)-4-[6-fluoro-1-(2-hydroxy-2-methyl-propyl)-7-methoxy-indazol-5-yl]benzoic acid obtained in step 3 above instead of 3-(4-cyano-3-fluoro-phenyl)-4-[6,7-difluoro-1-(2-hydroxy-2-methyl-propyl)indol-5-yl]benzoic acid, and using tert-butyl ((3-endo)-8-azabicyclo[3.2.1]octan-3-yl)carbamate instead of tert-butyl N-[(3S)-pyrrolidin-3-yl]carbamate to give the title compound.

[0596]

15 Example 187: Synthesis of (S)-5'-(3-aminopyrrolidine-1-carbonyl)-2'-(6,7-difluoro-1-(2-hydroxy-2-methylpropyl)-1H-indazol-5-yl)-3-fluoro-[1,1'-biphenyl]-4-carbonitrile

Step 1

The 5-bromo-6,7-difluoro-1H-indazole (101 mg) obtained in step 1 of Example 186 was dissolved in DMF (1.44 mL). At room temperature, cesium carbonate (283 mg) and 2,2-dimethyloxirane (0.077 mL) were added thereto, followed by stirring at 80°C overnight. Ethyl acetate was added thereto, and the mixture was washed sequentially with water and saturated brine, and dried over anhydrous sodium sulfate. Thereafter, the solvent was distilled off. The residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate) to give 1-(5-bromo-6,7-difluoro-indazol-1-yl)-2-methyl-propan-2-ol.

Step 2

30 The procedure of steps 2 to 4 in Example 172 was conducted using the 1-(5-bromo-6,7-difluoro-indazol-1-yl)-2-methyl-propan-2-ol obtained in step 1 above instead of 1-(5-bromo-6,7-difluoro-indol-1-yl)-2-methyl-propan-2-ol to give the title compound.

35 [0597]

Example 188: Synthesis of 5'-((3-endo)-3-amino-8-azabicyclo[3.2.1]octane-8-carbonyl)-2'-(6,7-difluoro-1-(2-hydroxy-2-methylpropyl)-1H-indazol-5-yl)-3-fluoro-[1,1'-biphenyl]-4-carbonitrile

5 The procedure of steps 2 to 4 in Example 172 was conducted using the 1-(5-bromo-6,7-difluoro-indazol-1-yl)-2-methyl-propan-2-ol obtained in step 1 of Example 187 instead of 1-(5-bromo-6,7-difluoro-indol-1-yl)-2-methyl-propan-2-ol, and using tert-butyl ((3-endo)-8-azabicyclo[3.2.1]octan-3-yl)carbamate instead of tert-butyl N-[(3S)-pyrrolidin-3-yl]carbamate to give the title compound.
10 [0598]

Example 189: Synthesis of 5'-((1S,2S,4R)-rel-2-amino-7-azabicyclo[2.2.1]heptane-7-carbonyl)-3-fluoro-2'-(6-fluoro-1-(2-hydroxy-2-methylpropyl)-1H-indazol-5-yl)-[1,1'-biphenyl]-4-carbonitrile

15 The procedure of steps 2 to 4 in Example 172 was conducted using the 1-(5-bromo-6-fluoro-indazol-1-yl)-2-methyl-propan-2-ol obtained in step 1 of Example 142 instead of 1-(5-bromo-6,7-difluoro-indol-1-yl)-2-methyl-propan-2-ol, and using tert-butyl ((1S,2S,4R)-rel-7-azabicyclo[2.2.1]heptan-2-yl)carbamate hydrochloride instead of tert-butyl N-[(3S)-pyrrolidin-3-yl]carbamate to give the title compound.
20 [0599]

25 Example 190: Synthesis of (S)-5'-(3-amino-3-methylpyrrolidine-1-carbonyl)-3-fluoro-2'-(6-fluoro-1-(2-hydroxy-2-methylpropyl)-1H-indol-5-yl)-[1,1'-biphenyl]-4-carbonitrile

 The procedure of steps 2 to 4 in Example 172 was conducted using the 1-(5-bromo-6-fluoro-indol-1-yl)-2-methyl-propan-2-ol obtained in step 1 of Example 136 instead of 1-(5-bromo-6,7-difluoro-indol-1-yl)-2-methyl-propan-2-ol, and using tert-butyl (S)-(3-methylpyrrolidin-3-yl)carbamate instead of tert-butyl N-[(3S)-pyrrolidin-3-yl]carbamate to give the title compound.
30 [0600]

Example 191: Synthesis of (S)-5'-(3-amino-3-methylpyrrolidine-1-carbonyl)-2'-(6,7-difluoro-1-(2-hydroxy-2-methylpropyl)-1H-benzo[d][1,2,3]triazol-5-yl)-3-fluoro-[1,1'-biphenyl]-4-carbonitrile

5 The procedure of steps 3 to 4 in Example 172 was conducted using the 3-(4-cyano-3-fluoro-phenyl)-4-[6,7-difluoro-1-(2-hydroxy-2-methyl-propyl)benzotriazol-5-yl]benzoic acid obtained in step 1 of Example 177 instead of 3-(4-cyano-3-fluoro-phenyl)-4-[6,7-difluoro-1-(2-hydroxy-2-methyl-propyl)indol-5-yl]benzoic acid, and using tert-butyl (S)-(3-methylpyrrolidin-3-yl)carbamate instead of tert-butyl N-[(3S)-pyrrolidin-3-yl]carbamate to give the title compound.
[0601]

Example 192: Synthesis of 5'-((1S,2S,4R)-rel-2-amino-7-azabicyclo[2.2.1]heptane-7-carbonyl)-2'-(6,7-difluoro-1-(2-hydroxy-2-methylpropyl)-1H-benzo[d][1,2,3]triazol-5-yl)-3-fluoro-[1,1'-biphenyl]-4-carbonitrile

15 The procedure of steps 3 to 4 in Example 172 was conducted using the 3-(4-cyano-3-fluoro-phenyl)-4-[6,7-difluoro-1-(2-hydroxy-2-methyl-propyl)benzotriazol-5-yl]benzoic acid obtained in step 1 of Example 177 instead of 3-(4-cyano-3-fluoro-phenyl)-4-[6,7-difluoro-1-(2-hydroxy-2-methyl-propyl)indol-5-yl]benzoic acid, and using tert-butyl ((1S,2S,4R)-rel-7-azabicyclo[2.2.1]heptan-2-yl)carbamate hydrochloride instead of
20 tert-butyl N-[(3S)-pyrrolidin-3-yl]carbamate to give the title compound.
[0602]

Example 193: Synthesis of 5'-((1S,2S,4R)-rel-2-amino-7-azabicyclo[2.2.1]heptane-7-carbonyl)-2'',3-difluoro-4''-(2-hydroxy-2-methylpropyl)-[1,1':2',1''-terphenyl]-4-carbonitrile

30 The procedure of steps 3 to 4 in Example 172 was conducted using the 3-(4-cyano-3-fluoro-phenyl)-4-[2-fluoro-4-(2-hydroxy-2-methyl-propyl)phenyl]benzoic acid obtained in step 2 of Example 166 instead of 3-(4-cyano-3-fluoro-phenyl)-4-[6,7-difluoro-1-(2-hydroxy-2-methyl-propyl)indol-5-yl]benzoic acid,
35

and using tert-butyl ((1S,2S,4R)-rel-7-azabicyclo[2.2.1]heptan-2-yl)carbamate hydrochloride instead of tert-butyl N-[(3S)-pyrrolidin-3-yl]carbamate to give the title compound.

[0603]

5 Example 194: Synthesis of 5'-((3-endo)-3-amino-8-azabicyclo[3.2.1]octane-8-carbonyl)-3-fluoro-2'-(6-fluoro-1-(3-hydroxy-3-methylbutyl)-1H-indazol-5-yl)-[1,1'-biphenyl]-4-carbonitrile

Step 1

10 5-Bromo-6-fluoro-1H-indazole (200 mg) was dissolved in DMF (3.1 mL). At room temperature, cesium carbonate (606 mg), and 3-hydroxy-3-methyl-butyl ester of 4-methylbenzene sulfonic acid (481 mg) were added thereto, followed by stirring at 90°C for 16 hours. Ethyl acetate was added thereto, and the mixture was
15 washed sequentially with water and saturated brine, and dried over anhydrous sodium sulfate. Thereafter, the solvent was distilled off. The residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate) to give 4-(5-bromo-6-fluoro-indazol-1-yl)-2-methyl-butan-2-ol.

20 [0604]

Step 2

 The procedure of steps 2 to 4 in Example 172 was conducted using the 4-(5-bromo-6-fluoro-indazol-1-yl)-2-methyl-butan-2-ol obtained in step 1 above instead of 1-(5-bromo-6,7-difluoro-indol-1-yl)-2-methyl-propan-2-ol, and using tert-butyl
25 ((3-endo)-8-azabicyclo[3.2.1]octan-3-yl)carbamate instead of tert-butyl N-[(3S)-pyrrolidin-3-yl]carbamate to give the title compound.

[0605]

30 Example 195: Synthesis of 5'-((1S,2S,4R)-rel-2-amino-7-azabicyclo[2.2.1]heptane-7-carbonyl)-2'-(6,7-difluoro-1-(2-hydroxy-2-methylpropyl)-1H-indazol-5-yl)-3-fluoro-[1,1'-biphenyl]-4-carbonitrile

 The procedure of steps 2 to 4 in Example 172 was
35 conducted using the 1-(5-bromo-6,7-difluoro-indazol-1-yl)-2-

methyl-propan-2-ol obtained in step 1 of Example 187 instead of 1-(5-bromo-6,7-difluoro-indol-1-yl)-2-methyl-propan-2-ol, and using tert-butyl ((1S,2S,4R)-rel-7-azabicyclo[2.2.1]heptan-2-yl)carbamate hydrochloride instead of tert-butyl N-[(3S)-pyrrolidin-3-yl]carbamate to give the title compound.

[0606]

Example 196: Synthesis of 5'-((1S,2S,4R)-rel-2-amino-7-azabicyclo[2.2.1]heptane-7-carbonyl)-2'-(6,7-difluoro-1-(2-hydroxy-2-methylpropyl)-1H-indol-5-yl)-3-fluoro-[1,1'-biphenyl]-4-carbonitrile

The procedure of steps 1 to 4 in Example 172 was conducted using tert-butyl ((1S,2S,4R)-rel-7-azabicyclo[2.2.1]heptan-2-yl)carbamate hydrochloride instead of tert-butyl N-[(3S)-pyrrolidin-3-yl]carbamate to give the title compound.

[0607]

Example 197: Synthesis of (S)-5'-(3-amino-3-methylpyrrolidine-1-carbonyl)-2'-(6,7-difluoro-1-(2-hydroxy-2-methylpropyl)-1H-indazol-5-yl)-3-fluoro-[1,1'-biphenyl]-4-carbonitrile

The procedure of steps 2 to 4 in Example 172 was conducted using the 1-(5-bromo-6,7-difluoro-indazol-1-yl)-2-methyl-propan-2-ol obtained in step 1 of Example 187 instead of 1-(5-bromo-6,7-difluoro-indol-1-yl)-2-methyl-propan-2-ol, and using tert-butyl (S)-(3-methylpyrrolidin-3-yl)carbamate instead of tert-butyl N-[(3S)-pyrrolidin-3-yl]carbamate to give the title compound.

[0608]

Example 198: Synthesis of (S)-5'-(3-amino-3-methylpyrrolidine-1-carbonyl)-2'-(6,7-difluoro-1-(2-hydroxy-2-methylpropyl)-1H-indol-5-yl)-3-fluoro-[1,1'-biphenyl]-4-carbonitrile

The procedure of steps 1 to 4 in Example 172 was conducted using tert-butyl (S)-(3-methylpyrrolidin-3-yl)carbamate instead of tert-butyl N-[(3S)-pyrrolidin-3-yl]carbamate to give the title compound.

[0609]

Example 199: Synthesis of 3-fluoro-2'-(6-fluoro-1-(2-hydroxy-2-methylpropyl)-1H-indazol-5-yl)-5'-(2,7-diazaspiro[3.4]octane-6-carbonyl)-[1,1'-biphenyl]-4-carbonitrile

The procedure of steps 2 to 4 in Example 172 was
5 conducted using the 1-(5-bromo-6-fluoro-indazol-1-yl)-2-methylpropan-2-ol obtained in step 1 of Example 142 instead of 1-(5-bromo-6,7-difluoro-indol-1-yl)-2-methylpropan-2-ol, and using tert-butyl 2,7-diazaspiro[3.4]octane-2-carboxylate instead of tert-butyl N-[(3S)-pyrrolidin-3-yl]carbamate to give the title
10 compound.

[0610]

Example 200: Synthesis of 2'-(6,7-difluoro-1-(2-hydroxy-2-methylpropyl)-1H-indol-5-yl)-3-fluoro-5'-(2,7-diazaspiro[3.4]octane-6-carbonyl)-[1,1'-biphenyl]-4-carbonitrile

15 The procedure of steps 1 to 4 in Example 172 was conducted using tert-butyl 2,7-diazaspiro[3.4]octane-2-carboxylate instead of tert-butyl N-[(3S)-pyrrolidin-3-yl]carbamate to give the title compound.

[0611]

20 Example 201: Synthesis of 2'-(6,7-difluoro-1-(2-hydroxy-2-methylpropyl)-1H-indol-5-yl)-3-fluoro-5'-(2,8-diazaspiro[3.5]nonane-2-carbonyl)-[1,1'-biphenyl]-4-carbonitrile

The procedure of steps 1 to 4 in Example 172 was conducted using tert-butyl 2,8-diazaspiro[3.5]nonane-6-
25 carboxylate instead of tert-butyl N-[(3S)-pyrrolidin-3-yl]carbamate to give the title compound.

[0612]

Example 202: Synthesis of 5'-((1S,2S,4R)-rel-2-amino-7-azabicyclo[2.2.1]heptane-7-carbonyl)-3-fluoro-2'-(5-fluoro-3-methylbenzo[d]isoxazol-6-yl)-[1,1'-biphenyl]-4-carbonitrile-isomer-X

Step 1

tert-Butyl ((1S,2S,4R)-rel-7-azabicyclo[2.2.1]heptan-2-yl)carbamate hydrochloride (36 mg) was dissolved in DCM (2.89
35 mL). At room temperature, TEA (40 µL) and benzyl chloroformate

(25 µL) were added thereto, followed by stirring at room temperature for 1 hour. The solvent was distilled off, and chloroform and water were added thereto. The mixture was extracted twice with chloroform and washed with water and saturated brine. The solvent was distilled off, and the residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate) to give benzyl (1S,2S,4R)-rel-2-((tert-butoxycarbonyl)amino)-7-azabicyclo[2.2.1]heptane-7-carboxylate. [0613]

The benzyl (1S,2S,4R)-rel-2-((tert-butoxycarbonyl)amino)-7-azabicyclo[2.2.1]heptane-7-carboxylate was obtained as a 10 mg/mL ethanol solution, and separation was performed under the following conditions. [0614]

The isomer having a shorter retention time was defined as "isomer-X," and the isomer having a longer retention time was defined as "isomer-Y."

Column: Daicel CHIRALPAK IC 2.0 x 25 cm

Mobile phase: hexane/2-propanol = 85/15

Flow rate: 12.5 mL/min

Retention time of each isomer:

benzyl (1S,2S,4R)-rel-2-((tert-butoxycarbonyl)amino)-7-azabicyclo[2.2.1]heptane-7-carboxylate-isomer-X: 16.93 minutes

benzyl (1S,2S,4R)-rel-2-((tert-butoxycarbonyl)amino)-7-azabicyclo[2.2.1]heptane-7-carboxylate-isomer-Y: 23.82 minutes.

[0615]

Chiral analysis conditions:

Column: CHIRALPAK IC 4.6 x 150 mm

Mobile phase: hexane/2-propanol = 85/15

Flow rate: 1.0 mL/min

Retention time of each isomer:

benzyl (1S,2S,4R)-rel-2-((tert-butoxycarbonyl)amino)-7-azabicyclo[2.2.1]heptane-7-carboxylate-isomer-X: 6.972 minutes

benzyl (1S,2S,4R)-rel-2-((tert-butoxycarbonyl)amino)-7-

azabicyclo[2.2.1]heptane-7-carboxylate-isomer-Y: 9.895 minutes.

[0616]

Step 2

The benzyl (1S,2S,4R)-rel-2-((tert-butoxycarbonyl)amino)-7-azabicyclo[2.2.1]heptane-7-carboxylate-isomer-X (93 g) obtained in step 1 above and 10% Pd/C (10 g) were suspended in methanol (1.0 L). The mixture was stirred at room temperature for 5 hours under a hydrogen atmosphere (50 psi). The reaction mixture was filtrated, and the filtrate was concentrated to give tert-butyl ((1S,2S,4R)-rel-7-azabicyclo[2.2.1]heptan-2-yl)carbamate-isomer-X.

Step 3

The procedure of steps 2 to 4 in Example 172 was conducted using the 6-bromo-5-fluoro-3-methylbenzo[d]isoxazol obtained in step 3 of Example 179 instead of 1-(5-bromo-6,7-difluoro-indol-1-yl)-2-methyl-propan-2-ol, and using the tert-butyl ((1S,2S,4R)-rel-7-azabicyclo[2.2.1]heptan-2-yl)carbamate-isomer-X obtained in step 2 above instead of tert-butyl N-[(3S)-pyrrolidin-3-yl]carbamate to give the title compound.

[0617]

Example 203: Synthesis of 2'-(6,7-difluoro-1-(2-hydroxy-2-methylpropyl)-1H-indol-5-yl)-3-fluoro-5'-(octahydropyrrolo[3,4-c]pyrrole-2-carbonyl)-[1,1'-biphenyl]-4-carbonitrile

The procedure of steps 1 to 4 in Example 172 was conducted using tert-butyl hexahydropyrrolo[3,4-c]pyrrole-2(1H)-carboxylate instead of tert-butyl N-[(3S)-pyrrolidin-3-yl]carbamate to give the title compound.

Example 204: Synthesis of (S)-5'-(3-aminopyrrolidine-1-carbonyl)-2'-(1-(2-ethyl-2-hydroxybutyl)-6-fluoro-1H-indazol-5-yl)-3-fluoro-[1,1'-biphenyl]-4-carbonitrile

Step 1

5-Bromo-6-fluoro-1H-indazole (300 mg) was dissolved in DMF (4.65 mL). At room temperature, cesium carbonate (90.9 mg) and 2,2-diethyloxirane (0.20 mL) were added thereto, followed by stirring at 90°C for 16 hours. Ethyl acetate was added thereto,

and the mixture was washed sequentially with water and saturated brine, and dried over anhydrous sodium sulfate. Thereafter, the solvent was distilled off. The residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate) to give 3-[(5-bromo-6-fluoro-indazol-1-yl)methyl]pentan-3-ol.

[0618]

Step 2

The procedure of steps 2 to 4 in Example 172 was conducted using the 3-[(5-bromo-6-fluoro-indazol-1-yl)methyl]pentan-3-ol obtained in step 1 above instead of 1-(5-bromo-6,7-difluoro-indol-1-yl)-2-methyl-propan-2-ol to give the title compound.

[0619]

Example 205: Synthesis of 5'-((3-endo)-3-amino-8-

azabicyclo[3.2.1]octane-8-carbonyl)-2'-(1-(2-ethyl-2-hydroxybutyl)-6-fluoro-1H-indazol-5-yl)-3-fluoro-[1,1'-biphenyl]-4-carbonitrile

The procedure of steps 2 to 4 in Example 172 was conducted using the 3-[(5-bromo-6-fluoro-indazol-1-yl)methyl]pentan-3-ol obtained in step 1 of Example 204 instead of 1-(5-bromo-6,7-difluoro-indol-1-yl)-2-methyl-propan-2-ol, and using tert-butyl ((3-endo)-8-azabicyclo[3.2.1]octan-3-yl)carbamate instead of tert-butyl N-[(3S)-pyrrolidin-3-yl]carbamate to give the title compound.

[0620]

Example 206: Synthesis of 5'-((1S,2S,4R)-rel-2-amino-7-

azabicyclo[2.2.1]heptane-7-carbonyl)-2'-(1-(2-ethyl-2-hydroxybutyl)-6-fluoro-1H-indazol-5-yl)-3-fluoro-[1,1'-biphenyl]-4-carbonitrile-isomer-X

The procedure of steps 2 to 4 in Example 172 was conducted using the 3-[(5-bromo-6-fluoro-indazol-1-yl)methyl]pentan-3-ol obtained in step 1 of Example 204 instead of 1-(5-bromo-6,7-difluoro-indol-1-yl)-2-methyl-propan-2-ol, and using the tert-butyl ((1S,2S,4R)-rel-7-azabicyclo[2.2.1]heptan-2-yl)carbamate-isomer-X obtained in step 2 of Example 202 instead

of tert-butyl N-[(3S)-pyrrolidin-3-yl]carbamate to give the title compound.

[0621]

Example 207: Synthesis of 2-(5-(5-((1S,2S,4R)-rel-2-amino-7-
5 azabicyclo[2.2.1]heptane-7-carbonyl)-4'-cyano-3'-fluoro-[1,1'-
biphenyl]-2-yl)-6-fluoro-1H-indol-1-yl)acetic acid-isomer-X

Step 1

5-Bromo-6-fluoro-1H-indole (500 mg) was dissolved in
DMF (7.79 mL). At room temperature, cesium carbonate (1.67 g) and
10 ethyl 2-chloro acetate (573 mg) were added thereto, followed by
stirring at 90°C for 16 hours. The reaction was terminated with a
saturated aqueous ammonium chloride solution. Ethyl acetate was
added thereto, and the mixture was washed sequentially with water
and saturated brine, and dried over anhydrous sodium sulfate.
15 Thereafter, the solvent was distilled off. The residue was
purified by silica gel column chromatography (mobile phase:
hexane/ethyl acetate) to give ethyl 2-(5-bromo-6-fluoro-indol-1-
yl)acetate.

[0622]

20 Step 2

The 3-(4-cyano-3-fluoro-phenyl)-4-(4,4,5,5-tetramethyl-
1,3,2-dioxaborolan-2-yl)benzoic acid (2 g) obtained in step 2 of
Example 41, and the tert-butyl ((1S,2S,4R)-rel-7-
azabicyclo[2.2.1]heptan-2-yl)carbamate-isomer-X (1.24 g) obtained
25 in step 2 of Example 202 were dissolved in THF (21.8 mL). At room
temperature, TEA (1.52 mL) and HATU (2.28 g) were added thereto,
followed by stirring at 50°C for 1 hour. The reaction mixture was
vacuum-concentrated, and the residue was purified by silica gel
column chromatography (mobile phase: hexane/ethyl acetate) to
30 give tert-butyl ((1S,2S,4R)-rel-7-(4'-cyano-3'-fluoro-6-(4,4,5,5-
tetramethyl-1,3,2-dioxaborolan-2-yl)-[1,1'-biphenyl]-3-carbonyl)-
7-azabicyclo[2.2.1]heptan-2-yl)carbamate-isomer-X.

[0623]

Step 3

35 The tert-butyl ((1S,2S,4R)-rel-7-(4'-cyano-3'-fluoro-6-

(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-[1,1'-biphenyl]-3-carbonyl)-7-azabicyclo[2.2.1]heptan-2-yl)carbamate-isomer-X (100 mg) obtained in step 2 above, and the ethyl 2-(5-bromo-6-fluoro-indol-1-yl)acetate (69.5 mg) obtained in step 1 above were
5 suspended in 1,4-dioxane (0.59 mL). At room temperature, Pd(dba)₂ (8.2 mg), X-phos (13.6 mg), and tripotassium phosphate (113 mg) were added thereto, followed by degassing and nitrogen substitution. Under a nitrogen atmosphere, stirring was performed at an external temperature of 100°C overnight. The solvent was
10 distilled off, and the residue was purified by silica gel column chromatography (mobile phase: chloroform/methanol). The residue was dissolved in MeOH (1.0 mL), and a 5 N aqueous sodium hydroxide solution (1.0 mL) was added thereto, followed by stirring for 1 hour. MTBE was added thereto, and the aqueous
15 layer was extracted. The aqueous layer was acidified with hydrochloric acid, MTBE was added thereto, and the resulting mixture was washed sequentially with water and saturated brine. After the organic layer was dried over anhydrous sodium sulfate, the solvent was distilled off to give 2-(5-(5-((1S,2S,4R)-rel-2-
20 ((tert-butoxycarbonyl)amino)-7-azabicyclo[2.2.1]heptane-7-carbonyl)-4'-cyano-3'-fluoro-[1,1'-biphenyl]-2-yl)-6-fluoro-1H-indole-1-yl)acetic acid-isomer-X.

[0624]

Step 4

25 Acetonitrile (1.0 mL) and a 4 N hydrochloric acid-1,4-dioxane solution (1.0 mL) were added to the 2-(5-(5-((1S,2S,4R)-rel-2-((tert-butoxycarbonyl)amino)-7-azabicyclo[2.2.1]heptane-7-carbonyl)-4'-cyano-3'-fluoro-[1,1'-biphenyl]-2-yl)-6-fluoro-1H-indole-1-yl)acetic acid-isomer-X (10 mg) obtained in step 3
30 above, followed by stirring for 30 minutes. The solvent was distilled off, and the residue was purified by reversed-phase HPLC (mobile phase: water/acetonitrile) to give the title compound.

[0625]

35 Example 208: Synthesis of 2-(4'-((1S,2S,4R)-rel-2-amino-7-

azabicyclo[2.2.1]heptane-7-carbonyl)-4''-cyano-2,3''-difluoro-[1,1':2',1''-terphenyl]-4-yl)acetic acid-isomer-X

The procedure of steps 3 to 4 in Example 207 was conducted using methyl 2-(4-bromo-3-fluorophenyl)acetate instead of ethyl 2-(5-bromo-6-fluoro-indol-1-yl)acetate to give the title compound.

[0626]

Example 209: Synthesis of 5'-((1S,2S,4R)-rel-2-amino-7-azabicyclo[2.2.1]heptane-7-carbonyl)-2'-(7-chloro-6-fluoro-1-(2-hydroxy-2-methylpropyl)-1H-benzo[d][1,2,3]triazol-5-yl)-3-fluoro-[1,1'-biphenyl]-4-carbonitrile-isomer-X

Step 1

2-Chloro-1,3-difluoro-4-nitro-benzene (1 g) was dissolved in THF (12.9 mL). TEA (1.08 mL) and 1-amino-2-methylpropan-2-ol (0.59 mL) were added thereto, followed by stirring at room temperature for 1 hour. Ethyl acetate was added thereto, and the mixture was washed sequentially with water and saturated brine. The organic layer was dried over anhydrous sodium sulfate, and the solvent was distilled off to give 1-(2-chloro-3-fluoro-6-nitro-anilino)-2-methyl-propan-2-ol.

[0627]

Step 2

The 1-(2-chloro-3-fluoro-6-nitro-anilino)-2-methylpropan-2-ol (1.3 g) obtained in step 1 above was dissolved in DMF (9.9 mL). At room temperature, NBS (1.1 g) was added thereto, followed by stirring at 90°C for 1 hour. Ethyl acetate was added thereto, and the mixture was washed sequentially with water and saturated brine, and dried over anhydrous sodium sulfate. Thereafter, the solvent was distilled off. The residue was crystallized from IPE:hexane = 1:1, and washed twice with hexane to give 1-(4-bromo-2-chloro-3-fluoro-6-nitro-anilino)-2-methyl-propan-2-ol.

[0628]

Step 3

The 1-(4-bromo-2-chloro-3-fluoro-6-nitro-anilino)-2-

methyl-propan-2-ol (1.6 g) obtained in step 2 above, NH_4Cl (1.6 g), and iron (0.8 g) were suspended in EtOH (7.81 mL) and water (7.81 mL), followed by stirring at 60°C overnight. MTBE was added thereto, and the mixture was passed through Celite. MTBE was added thereto, and the mixture was washed sequentially with water and saturated brine, and dried over anhydrous sodium sulfate. The solvent was then distilled off to give 1-(6-amino-4-bromo-2-chloro-3-fluoro-anilino)-2-methyl-propan-2-ol.

[0629]

10 Step 4

The 1-(6-amino-4-bromo-2-chloro-3-fluoro-anilino)-2-methyl-propan-2-ol (352 mg) obtained in step 3 above was dissolved in water (0.70 mL) and THF (1.76 mL). At 0°C, 12 N hydrochloric acid (1.06 mL) and sodium nitrite (an aqueous solution (0.3 mL) in which 101 mg of sodium nitrite was dissolved) were added thereto dropwise, followed by stirring at room temperature for 1 hour. MTBE was added thereto, and the mixture was washed sequentially with water and saturated brine, and dried over anhydrous sodium sulfate. The solvent was then distilled off. IPE:hexane = 1:1 (68 mL) was added to the residue, and the target compound was collected by filtration and washed with IPE:hexane = 1:1 to give 1-(5-bromo-7-chloro-6-fluoro-benzotriazol-1-yl)-2-methyl-propan-2-ol.

[0630]

25 Step 5

The tert-butyl ((1S,2S,4R)-rel-7-(4'-cyano-3'-fluoro-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-[1,1'-biphenyl]-3-carbonyl)-7-azabicyclo[2.2.1]heptan-2-yl)carbamate-isomer-X (50 mg) obtained in step 2 of Example 207, and the 1-(5-bromo-7-chloro-6-fluoro-benzotriazol-1-yl)-2-methyl-propan-2-ol (37.4 mg) obtained in step 4 above were suspended in 1,4-dioxane (0.3 mL). At room temperature, $\text{Pd}(\text{dba})_2$ (4.1 mg), X-phos (6.8 mg), and tripotassium phosphate (56.7 mg) were added thereto. After nitrogen substitution, the mixture was stirred at 100°C for 2 hours. Ethyl acetate was added thereto, and the mixture was put

on NH-silica gel, and washed with ethyl acetate: methanol =10:1. The solvent was distilled off, and acetonitrile (1.0 mL) and a 4 N hydrochloric acid-1,4-dioxane solution (1.0 mL) were added to the residue, followed by stirring for 10 minutes. The solvent was
5 distilled off, the residue was dissolved in DMSO, and purification was performed by reversed-phase HPLC (mobile phase: water/acetonitrile) to give the title compound.

[0631]

10 Example 210: Synthesis of (S)-5'-(3-aminopyrrolidine-1-carbonyl)-2'-(7-chloro-6-fluoro-1-(2-hydroxy-2-methylpropyl)-1H-benzo[d][1,2,3]triazol-5-yl)-3-fluoro-[1,1'-biphenyl]-4-carbonitrile

The procedure of steps 1 to 5 in Example 209 was conducted using the tert-butyl N-[(3S)-1-[3-(4-cyano-3-fluoro-phenyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoyl]pyrrolidin-3-yl]carbamate obtained in step 1 of
15 Example 37 instead of tert-butyl ((1S,2S,4R)-rel-7-(4'-cyano-3'-fluoro-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-[1,1'-biphenyl]-3-carbonyl)-7-azabicyclo[2.2.1]heptan-2-yl)carbamate-
20 isomer-X to give the title compound.

[0632]

Example 211: Synthesis of 5'-((3-endo)-3-amino-8-azabicyclo[3.2.1]octane-8-carbonyl)-2'-(7-chloro-6-fluoro-1-(2-hydroxy-2-methylpropyl)-1H-benzo[d][1,2,3]triazol-5-yl)-3-fluoro-
25 [1,1'-biphenyl]-4-carbonitrile

The procedure of steps 1 to 5 in Example 209 was conducted using the tert-butyl N-[(3-endo)-8-[3-(4-cyano-3-fluoro-phenyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoyl]-8-azabicyclo[3.2.1]octan-3-yl]carbamate obtained in
30 step 3 of Example 41 instead of tert-butyl ((1S,2S,4R)-rel-7-(4'-cyano-3'-fluoro-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-[1,1'-biphenyl]-3-carbonyl)-7-azabicyclo[2.2.1]heptan-2-yl)carbamate-isomer-X to give the title compound.

[0633]

35 Example 212: Synthesis of 5-(5-((1S,2S,4R)-rel-2-amino-7-

azabicyclo[2.2.1]heptane-7-carbonyl)-4'-cyano-3'-fluoro-[1,1'-biphenyl]-2-yl)-6-fluoro-1-(2-hydroxy-2-methylpropyl)-1H-indazole-3-carboxylic acid-isomer-X

Step 1

5 The 3-(4-cyano-3-fluoro-phenyl)-4-[6-fluoro-1-(2-hydroxy-2-methyl-propyl)indazol-5-yl]benzoic acid (250 mg) obtained in step 1 of Example 176 was dissolved in THF (2.24 mL). At room temperature, HATU (234 mg), the tert-butyl ((1S,2S,4R)-rel-7-azabicyclo[2.2.1]heptan-2-yl)carbamate-isomer-X (125 mg)
10 obtained in step 2 of Example 202, and TEA (0.156 mL) were added thereto, followed by stirring at 50°C for 1 hour. The solvent was distilled off under reduced pressure. The residue was purified by silica gel column chromatography (mobile phase: chloroform/methanol) to give tert-butyl ((1S,2S,4R)-rel-7-(4'-cyano-3'-fluoro-6-(6-fluoro-1-(2-hydroxy-2-methylpropyl)-1H-indazol-5-yl)-[1,1'-biphenyl]-3-carbonyl)-7-azabicyclo[2.2.1]heptan-2-yl)carbamate-isomer-X.

[0634]

Step 2

20 The tert-butyl ((1S,2S,4R)-rel-7-(4'-cyano-3'-fluoro-6-(6-fluoro-1-(2-hydroxy-2-methylpropyl)-1H-indazol-5-yl)-[1,1'-biphenyl]-3-carbonyl)-7-azabicyclo[2.2.1]heptan-2-yl)carbamate-isomer-X (289 mg) obtained in step 1 above was dissolved in DMF (4.50 mL). At room temperature, NBS (120 mg) was added thereto,
25 followed by stirring at room temperature for 1 hour. Ethyl acetate was added thereto, and the mixture was washed sequentially with water and saturated brine. After the organic layer was dried over anhydrous sodium sulfate, the solvent was distilled off. The residue was purified by silica gel column
30 chromatography (mobile phase: hexane/ethyl acetate) to give tert-butyl ((1S,2S,4R)-rel-7-(6-(3-bromo-6-fluoro-1-(2-hydroxy-2-methylpropyl)-1H-indazol-5-yl)-4'-cyano-3'-fluoro-[1,1'-biphenyl]-3-carbonyl)-7-azabicyclo[2.2.1]heptan-2-yl)carbamate-isomer-X.

35 [0635]

Step 3

The tert-butyl ((1S,2S,4R)-rel-7-(6-(3-bromo-6-fluoro-1-(2-hydroxy-2-methylpropyl)-1H-indazol-5-yl)-4'-cyano-3'-fluoro-[1,1'-biphenyl]-3-carbonyl)-7-azabicyclo[2.2.1]heptan-2-yl)carbamate-isomer-X (50 mg) obtained in step 2 above and PdCl₂(PPh₃)₂ (2.4 mg) were suspended in 1-methyl-2-pyrrolidinone (0.5 mL). At room temperature, N,N-diethylethanolamine (0.046 mL) was added thereto, and after CO substitution, the mixture was stirred at 125°C for 1 hour. t-BuOH (0.5 mL) and a 2 N aqueous sodium hydroxide solution (0.25 mL) were added to the reaction mixture, followed by stirring at room temperature for 1 hour. MTBE was added thereto, and the aqueous layer was separated. The aqueous layer was acidified with hydrochloric acid, and extraction was performed with MTBE. After the organic layer was dried over anhydrous sodium sulfate, the solvent was distilled off. Acetonitrile (0.5 mL) and a 4 N hydrochloric acid-1,4-dioxane solution (0.5 mL) were added to the residue, followed by stirring for 10 minutes. The reaction mixture was concentrated, and the residue was purified by reversed-phase HPLC (mobile phase: water/acetonitrile) to give the title compound. [0636]

Example 213: Synthesis of 5'-((1S,2S,4R)-rel-2-amino-7-azabicyclo[2.2.1]heptane-7-carbonyl)-2'-(7-chloro-6-fluoro-1-(2-hydroxy-2-methylpropyl)-1H-benzo[d][1,2,3]triazol-5-yl)-3-fluoro-[1,1'-biphenyl]-4-carbonitrile

Step 1

The 3-(4-cyano-3-fluoro-phenyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoic acid (500 mg) obtained in step 2 of Example 41 and tert-butyl ((1S,2S,4R)-rel-7-azabicyclo[2.2.1]heptan-2-yl)carbamate hydrochloride (303.5 mg) were dissolved in THF (5.45 mL). At room temperature, TEA (0.379 mL) and HATU (569.5 mg) were added thereto, followed by stirring at 50°C for 1 hour. The reaction mixture was vacuum-concentrated, and the residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate) to give tert-butyl

((1S,2S,4R)-rel-7-(4'-cyano-3'-fluoro-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-[1,1'-biphenyl]-3-carbonyl)-7-azabicyclo[2.2.1]heptan-2-yl) carbamate.

Step 2

5 The procedure of steps 1 to 5 in Example 209 was conducted using the tert-butyl ((1S,2S,4R)-rel-7-(4'-cyano-3'-fluoro-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-[1,1'-biphenyl]-3-carbonyl)-7-azabicyclo[2.2.1]heptan-2-yl) carbamate obtained in step 1 above instead of tert-butyl ((1S,2S,4R)-rel-7-(4'-cyano-3'-fluoro-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-[1,1'-biphenyl]-3-carbonyl)-7-azabicyclo[2.2.1]heptan-2-yl) carbamate-isomer-X to give the title compound.
10 [0637]

Example 214: Synthesis of 5'-((1S,2S,4R)-rel-2-amino-7-azabicyclo[2.2.1]heptane-7-carbonyl)-2'-(1-(2-ethyl-2-hydroxybutyl)-6,7-difluoro-1H-benzo[d][1,2,3]triazol-5-yl)-3-fluoro-[1,1'-biphenyl]-4-carbonitrile-isomer-X

 The procedure of steps 1 to 5 in Example 161 was conducted using 3-[(2,3-difluoro-6-nitro-anilino)methyl]pentan-3-ol instead of 1-(2,3-difluoro-6-nitro-anilino)-2-methyl-propan-2-ol, and using the tert-butyl ((1S,2S,4R)-rel-7-(4'-cyano-3'-fluoro-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-[1,1'-biphenyl]-3-carbonyl)-7-azabicyclo[2.2.1]heptan-2-yl) carbamate-isomer-X obtained in step 2 of Example 207 instead of tert-butyl
20 N-[(3-endo)-8-[3-(4-cyano-3-fluoro-phenyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoyl]-8-azabicyclo[3.2.1]octan-3-yl] carbamate to give the title compound.
25 [0638]

Example 215: Synthesis of (S)-5'-(3-aminopyrrolidine-1-carbonyl)-2'-(1-(2-ethyl-2-hydroxybutyl)-6,7-difluoro-1H-benzo[d][1,2,3]triazol-5-yl)-3-fluoro-[1,1'-biphenyl]-4-carbonitrile

 The procedure of steps 1 to 5 in Example 161 was conducted using 3-[(2,3-difluoro-6-nitro-anilino)methyl]pentan-3-ol instead of 1-(2,3-difluoro-6-nitro-anilino)-2-methyl-propan-2-
35 ol

ol, and using the tert-butyl N-[(3S)-1-[3-(4-cyano-3-fluoro-phenyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoyl]pyrrolidin-3-yl]carbamate obtained in step 1 of Example 37 instead of tert-butyl N-[(3-endo)-8-[3-(4-cyano-3-fluoro-phenyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoyl]-8-azabicyclo[3.2.1]octan-3-yl]carbamate to give the title compound.

[0639]

Example 216: Synthesis of 5'-((3-endo)-3-amino-8-azabicyclo[3.2.1]octane-8-carbonyl)-2'-(1-(2-ethyl-2-hydroxybutyl)-6,7-difluoro-1H-benzo[d][1,2,3]triazol-5-yl)-3-fluoro-[1,1'-biphenyl]-4-carbonitrile

The procedure of steps 1 to 5 in Example 161 was conducted using 3-[(2,3-difluoro-6-nitro-anilino)methyl]pentan-3-ol instead of 1-(2,3-difluoro-6-nitro-anilino)-2-methyl-propan-2-ol to give the title compound.

[0640]

Example 217: Synthesis of 5'-((1S,2S,4R)-rel-2-amino-7-azabicyclo[2.2.1]heptane-7-carbonyl)-2'-(1-(2-ethyl-2-hydroxybutyl)-6,7-difluoro-1H-benzo[d][1,2,3]triazol-5-yl)-3-fluoro-[1,1'-biphenyl]-4-carbonitrile

The procedure of steps 1 to 5 in Example 161 was conducted using 3-[(2,3-difluoro-6-nitro-anilino)methyl]pentan-3-ol instead of 1-(2,3-difluoro-6-nitro-anilino)-2-methyl-propan-2-ol, and using the tert-butyl ((1S,2S,4R)-rel-7-(4'-cyano-3'-fluoro-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-[1,1'-biphenyl]-3-carbonyl)-7-azabicyclo[2.2.1]heptan-2-yl)carbamate obtained in step 1 of Example 213 instead of tert-butyl N-[(3-endo)-8-[3-(4-cyano-3-fluoro-phenyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoyl]-8-azabicyclo[3.2.1]octan-3-yl]carbamate to give the title compound.

[0641]

Example 218: Synthesis of 5'-((1S,2S,4R)-rel-2-amino-7-azabicyclo[2.2.1]heptane-7-carbonyl)-2'-(7-chloro-1-(2-ethyl-2-hydroxybutyl)-6-fluoro-1H-benzo[d][1,2,3]triazol-5-yl)-3-fluoro-

[1,1'-biphenyl]-4-carbonitrile-isomer-X

The procedure of steps 1 to 5 in Example 209 was conducted using 3-(aminomethyl)pentan-3-ol instead of 1-amino-2-methyl-propan-2-ol to give the title compound.

5 [0642]

Example 219: Synthesis of (S)-5'-(3-aminopyrrolidine-1-carbonyl)-2'-(7-chloro-1-(2-ethyl-2-hydroxybutyl)-6-fluoro-1H-benzo[d][1,2,3]triazol-5-yl)-3-fluoro-[1,1'-biphenyl]-4-carbonitrile

10 The procedure of steps 1 to 5 in Example 209 was conducted using 3-(aminomethyl)pentan-3-ol instead of 1-amino-2-methyl-propan-2-ol, and using the tert-butyl N-[(3S)-1-[3-(4-cyano-3-fluoro-phenyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoyl]pyrrolidin-3-yl]carbamate obtained in step 1 of
15 Example 37 instead of tert-butyl ((1S,2S,4R)-rel-7-(4'-cyano-3'-fluoro-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-[1,1'-biphenyl]-3-carbonyl)-7-azabicyclo[2.2.1]heptan-2-yl)carbamate-isomer-X to give the title compound.

[0643]

20 Example 220: Synthesis of 5'-((3-endo)-3-amino-8-azabicyclo[3.2.1]octane-8-carbonyl)-2'-(7-chloro-1-(2-ethyl-2-hydroxybutyl)-6-fluoro-1H-benzo[d][1,2,3]triazol-5-yl)-3-fluoro-[1,1'-biphenyl]-4-carbonitrile

The procedure of steps 1 to 5 in Example 209 was
25 conducted using 3-(aminomethyl)pentan-3-ol instead of 1-amino-2-methyl-propan-2-ol, and using the tert-butyl N-[(3-endo)-8-[3-(4-cyano-3-fluoro-phenyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoyl]-8-azabicyclo[3.2.1]octan-3-yl]carbamate obtained in step 3 of Example 41 instead of tert-butyl ((1S,2S,4R)-rel-7-(4'-cyano-3'-fluoro-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-
30 [1,1'-biphenyl]-3-carbonyl)-7-azabicyclo[2.2.1]heptan-2-yl)carbamate-isomer-X to give the title compound.

[0644]

Example 221: Synthesis of 5'-((1S,2S,4R)-rel-2-amino-7-azabicyclo[2.2.1]heptane-7-carbonyl)-2'-(7-chloro-1-(2-ethyl-2-

35

hydroxybutyl)-6-fluoro-1H-benzo[d][1,2,3]triazol-5-yl)-3-fluoro-
[1,1'-biphenyl]-4-carbonitrile

The procedure of steps 1 to 5 in Example 209 was conducted using 3-(aminomethyl)pentan-3-ol instead of 1-amino-2-methyl-propan-2-ol, and using the tert-butyl ((1S,2S,4R)-rel-7-(4'-cyano-3'-fluoro-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-[1,1'-biphenyl]-3-carbonyl)-7-azabicyclo[2.2.1]heptan-2-yl)carbamate obtained in step 1 of Example 213 instead of tert-butyl ((1S,2S,4R)-rel-7-(4'-cyano-3'-fluoro-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-[1,1'-biphenyl]-3-carbonyl)-7-azabicyclo[2.2.1]heptan-2-yl)carbamate-isomer-X to give the title compound.

[0645]

Example 222: Synthesis of 5-(5-((1S,2S,4R)-rel-2-amino-7-azabicyclo[2.2.1]heptane-7-carbonyl)-4'-cyano-3'-fluoro-[1,1'-biphenyl]-2-yl)-6-fluoro-1-(2-hydroxy-2-methylpropyl)-1H-indole-3-carboxylic acid-isomer-X

Step 1

The 3-(4-cyano-3-fluoro-phenyl)-4-[6-fluoro-1-(2-hydroxy-2-methyl-propyl)indol-5-yl]benzoic acid (250 mg) obtained in step 1 of Example 171 was dissolved in THF (2.24 mL). At room temperature, HATU (234 mg), the tert-butyl ((1S,2S,4R)-rel-7-azabicyclo[2.2.1]heptan-2-yl)carbamate-isomer-X (125 mg) obtained in step 2 of Example 202, and TEA (0.156 mL) were added thereto, followed by stirring at 50°C for 1 hour. The solvent was distilled off under reduced pressure. The residue was purified by silica gel column chromatography (mobile phase: chloroform/methanol) to give tert-butyl ((1S,2S,4R)-rel-7-(4'-cyano-3'-fluoro-6-(6-fluoro-1-(2-hydroxy-2-methylpropyl)-1H-indol-5-yl)-[1,1'-biphenyl]-3-carbonyl)-7-azabicyclo[2.2.1]heptan-2-yl)carbamate-isomer-X.

[0646]

Step 2

The tert-butyl ((1S,2S,4R)-rel-7-(4'-cyano-3'-fluoro-6-(6-fluoro-1-(2-hydroxy-2-methylpropyl)-1H-indol-5-yl)-[1,1'-

biphenyl]-3-carbonyl)-7-azabicyclo[2.2.1]heptan-2-yl) carbamate-isomer-X (289 mg) obtained in step 1 above was dissolved in DMF (4.50 mL). At room temperature, N-iodosuccinimide (120 mg) was added thereto, followed by stirring at room temperature for 1
5 hour. Ethyl acetate was added thereto, and the mixture was washed sequentially with water and saturated brine. After the organic layer was dried over anhydrous sodium sulfate, the solvent was distilled off. The residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate) to give tert-
10 butyl ((1S,2S,4R)-rel-7-(4'-cyano-3'-fluoro-6-(6-fluoro-1-(2-hydroxy-2-methylpropyl)-3-iodo-1H-indol-5-yl)-[1,1'-biphenyl]-3-carbonyl)-7-azabicyclo[2.2.1]heptan-2-yl) carbamate-isomer-X.
[0647]

Step 3

15 The tert-butyl ((1S,2S,4R)-rel-7-(4'-cyano-3'-fluoro-6-(6-fluoro-1-(2-hydroxy-2-methylpropyl)-3-iodo-1H-indol-5-yl)-[1,1'-biphenyl]-3-carbonyl)-7-azabicyclo[2.2.1]heptan-2-yl) carbamate-isomer-X (20 mg) obtained in step 2 above and PdCl₂(PPh₃)₂ (0.92 mg) were suspended in 1-methyl-2-pyrrolidinone
20 (0.2 mL). At room temperature, N,N-diethylethanolamine (0.0173 mL) was added thereto, and after CO substitution, the mixture was stirred at 100°C for 1 hour. t-BuOH (0.2 mL) and a 2 N aqueous sodium hydroxide solution (0.2 mL) were added to the reaction mixture, followed by stirring at room temperature overnight. MTBE
25 was added thereto, and the aqueous layer was separated. The aqueous layer was acidified with hydrochloric acid, and extraction was performed with MTBE. After the organic layer was dried over anhydrous sodium sulfate, the solvent was distilled off. Acetonitrile (0.5 mL) and a 4 N hydrochloric acid-1,4-
30 dioxane solution (0.5 mL) were added to the residue, followed by stirring for 10 minutes. The reaction mixture was concentrated, and the residue was purified by reversed-phase HPLC (mobile phase: water/acetonitrile) to give the title compound.
[0648]

35 Example 223: Synthesis of 5-(5-((1S,2S,4R)-rel-2-amino-7-

azabicyclo[2.2.1]heptane-7-carbonyl)-4'-cyano-3'-fluoro-[1,1'-biphenyl]-2-yl)-6-fluoro-1-methyl-1H-indazole-3-carboxylic acid-isomer-X

Step 1

5 The tert-butyl ((1S,2S,4R)-rel-7-(4'-cyano-3'-fluoro-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-[1,1'-biphenyl]-3-carbonyl)-7-azabicyclo[2.2.1]heptan-2-yl)carbamate-isomer-X (350 mg) obtained in step 2 of Example 207, and 5-bromo-6-fluoro-1-methyl-indazole (186 mg) were suspended in 1,4-dioxane (2.08 mL).
10 At room temperature, Pd(dba)₂ (28.7 mg), X-phos (47.6 mg), and tripotassium phosphate (397 mg) were added thereto, followed by degassing and nitrogen substitution. Under a nitrogen atmosphere, stirring was performed at an external temperature of 100°C overnight. The solvent was distilled off, and the residue was
15 purified by silica gel column chromatography (mobile phase: chloroform/methanol) to give tert-butyl ((1S,2S,4R)-rel-7-(4'-cyano-3'-fluoro-6-(6-fluoro-1-methyl-1H-indazol-5-yl)-[1,1'-biphenyl]-3-carbonyl)-7-azabicyclo[2.2.1]heptan-2-yl)carbamate-isomer-X.

20 Step 2

 The procedure of steps 2 to 3 in Example 212 was conducted using the tert-butyl ((1S,2S,4R)-rel-7-(4'-cyano-3'-fluoro-6-(6-fluoro-1-methyl-1H-indazol-5-yl)-[1,1'-biphenyl]-3-carbonyl)-7-azabicyclo[2.2.1]heptan-2-yl)carbamate-isomer-X
25 obtained in step 1 above instead of tert-butyl ((1S,2S,4R)-rel-7-(4'-cyano-3'-fluoro-6-(6-fluoro-1-(2-hydroxy-2-methylpropyl)-1H-indazol-5-yl)-[1,1'-biphenyl]-3-carbonyl)-7-azabicyclo[2.2.1]heptan-2-yl)carbamate-isomer-X to give the title compound.

30 [0649]

Example 224: Synthesis of 5-(5-((1S,2S,4R)-rel-2-amino-7-azabicyclo[2.2.1]heptane-7-carbonyl)-4'-cyano-3'-fluoro-[1,1'-biphenyl]-2-yl)-6-fluoro-1-methyl-1H-indole-3-carboxylic acid-isomer-X

35 Step 1

The tert-butyl ((1S,2S,4R)-rel-7-(4'-cyano-3'-fluoro-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-[1,1'-biphenyl]-3-carbonyl)-7-azabicyclo[2.2.1]heptan-2-yl)carbamate-isomer-X (350 mg) obtained in step 2 of Example 207, and 5-bromo-6-fluoro-1-methyl-indole (185 mg) were suspended in 1,4-dioxane (2.08 mL). At room temperature, Pd(dba)₂ (28.7 mg), X-phos (47.6 mg), and tripotassium phosphate (397 mg) were added thereto, followed by degassing and nitrogen substitution. Under a nitrogen atmosphere, stirring was performed at an external temperature of 100°C overnight. The solvent was distilled off, and the residue was purified by silica gel column chromatography (mobile phase: chloroform/methanol) to give tert-butyl ((1S,2S,4R)-rel-7-(4'-cyano-3'-fluoro-6-(6-fluoro-1-methyl-1H-indol-5-yl)-[1,1'-biphenyl]-3-carbonyl)-7-azabicyclo[2.2.1]heptan-2-yl)carbamate-isomer-X.

[0650]

Step 2

The tert-butyl ((1S,2S,4R)-rel-7-(4'-cyano-3'-fluoro-6-(6-fluoro-1-methyl-1H-indol-5-yl)-[1,1'-biphenyl]-3-carbonyl)-7-azabicyclo[2.2.1]heptan-2-yl)carbamate-isomer-X (209 mg) obtained in step 1 above was dissolved in DMF (3.6 mL). At room temperature, N-iodosuccinimide (121 mg) was added thereto, followed by stirring at room temperature for 1 hour. Ethyl acetate was added thereto, and the resulting mixture was washed sequentially with water and saturated brine. After the organic layer was dried over anhydrous sodium sulfate, the solvent was distilled off. The residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate) to give tert-butyl ((1S,2S,4R)-rel-7-(4'-cyano-3'-fluoro-6-(6-fluoro-3-iodo-1-methyl-1H-indol-5-yl)-[1,1'-biphenyl]-3-carbonyl)-7-azabicyclo[2.2.1]heptan-2-yl)carbamate-isomer-X.

[0651]

Step 3

The tert-butyl ((1S,2S,4R)-rel-7-(4'-cyano-3'-fluoro-6-(6-fluoro-3-iodo-1-methyl-1H-indol-5-yl)-[1,1'-biphenyl]-3-

carbonyl)-7-azabicyclo[2.2.1]heptan-2-yl) carbamate-isomer-X (211 mg) obtained in step 2 above and PdCl₂(PPh₃)₂ (10.5 mg) were suspended in 1-methyl-2-pyrrolidinone (2.11 mL). At room temperature, N,N-diethylethanolamine (0.197 mL) was added thereto, and after CO substitution, the mixture was stirred at 100°C for 1 hour. t-BuOH (0.2 mL) and a 2 N aqueous sodium hydroxide solution (0.2 mL) were added to the reaction mixture, and the resulting mixture was stirred at room temperature overnight. MTBE was added thereto, and the aqueous layer was separated. The aqueous layer was acidified with hydrochloric acid, and extraction was performed with MTBE. The organic layer was dried over anhydrous sodium sulfate, and the solvent was distilled off to give 5-(5-((1S,2S,4R)-rel-2-((tert-butoxycarbonyl)amino)-7-azabicyclo[2.2.1]heptane-7-carbonyl)-4'-cyano-3'-fluoro-[1,1'-biphenyl]-2-yl)-6-fluoro-1-methyl-1H-indole-3-carboxylic acid-isomer-X.

Step 4

Acetonitrile (0.5 mL) and a 4 N hydrochloric acid-1,4-dioxane solution (0.5 mL) were added to the 5-(5-((1S,2S,4R)-rel-2-((tert-butoxycarbonyl)amino)-7-azabicyclo[2.2.1]heptane-7-carbonyl)-4'-cyano-3'-fluoro-[1,1'-biphenyl]-2-yl)-6-fluoro-1-methyl-1H-indole-3-carboxylic acid-isomer-X (10 mg) obtained in step 3 above, followed by stirring for 10 minutes. The reaction mixture was concentrated, and the residue was purified by reversed-phase HPLC (mobile phase: water/acetonitrile) to give the title compound.

[0652]

Example 225: Synthesis of 5'-((1S,2S,4R)-rel-2-amino-7-azabicyclo[2.2.1]heptane-7-carbonyl)-2'-(1-(2-ethyl-2-hydroxybutyl)-6-fluoro-1H-indazol-5-yl)-3-fluoro-[1,1'-biphenyl]-4-carbonitrile

The tert-butyl ((1S,2S,4R)-rel-7-(4'-cyano-3'-fluoro-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-[1,1'-biphenyl]-3-carbonyl)-7-azabicyclo[2.2.1]heptan-2-yl) carbamate (40 mg) obtained in step 1 of Example 213 and the 3-[(5-bromo-6-fluoro-

indazol-1-yl)methyl]pentan-3-ol (29.2 mg) obtained in step 1 of Example 204 were suspended in 1,4-dioxane (0.5 mL). At room temperature, Pd(dba)₂ (3.3 mg), X-phos (5.5 mg), and tripotassium phosphate (45.4 mg) were added thereto, followed by stirring at 100°C for 1 hour. The reaction mixture was filtrated, and the solvent was distilled off. The residue was dissolved in acetonitrile (0.5 mL). At room temperature, a 4 N hydrochloric acid-1,4-dioxane solution (0.5 mL) was added thereto, followed by stirring at room temperature for 5 minutes. The reaction mixture was concentrated, and the residue was purified by reversed-phase HPLC (mobile phase: water/acetonitrile) to give the title compound.

[0653]

Example 226: Synthesis of 5'-((1S,2S,4R)-rel-2-amino-7-azabicyclo[2.2.1]heptane-7-carbonyl)-3-fluoro-2'-(6-fluoro-1-(2-hydroxy-2-methylpropyl)-1,3-dihydroisobenzofuran-5-yl)-[1,1'-biphenyl]-4-carbonitrile-isomer-X

Step 1

Methyl 5-bromo-4-fluoro-2-iodo-benzoate (2 g) was dissolved in diethyl ether (55.7 mL). At 0°C, a solution of 2.0 M LiBH₄ in THF (6.13 mL) and MeOH (0.56 mL) were added thereto, followed by stirring at 0°C for 1 hour. MTBE was added thereto, and the mixture was washed sequentially with water and saturated brine, and dried over anhydrous sodium sulfate. Thereafter, the solvent was distilled off. The residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate) to give (5-bromo-4-fluoro-2-iodo-phenyl)methanol.

[0654]

Step 2

The (5-bromo-4-fluoro-2-iodo-phenyl)methanol (1.39 g) obtained in step 1 above and 3,4-dihydro-2H-pyran (0.419 mL) were dissolved in DCM (8.4 mL). At room temperature, pyridinium p-toluenesulfonic acid (106 mg) was added thereto, followed by stirring at room temperature overnight. Ethyl acetate was added thereto, and the mixture was washed sequentially with water and

saturated brine, and dried over anhydrous sodium sulfate. Thereafter, the solvent was distilled off. The residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate) to give 2-[(5-bromo-4-fluoro-2-iodo-phenyl)methoxy]tetrahydropyran.

[0655]

Step 3

The 2-[(5-bromo-4-fluoro-2-iodo-phenyl)methoxy]tetrahydropyran (1.5 g) obtained in step 2 above, PdCl₂(PPh₃)₂ (130 mg), and CuI (34 mg) were suspended in THF (18 mL). At room temperature, TEA (18 mL) and 2-methyl-3-butyn-2-ol (0.42 mL) were added thereto, followed by stirring at room temperature for 4 hours. The reaction mixture was filtrated, and the solvent was distilled off. The residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate) to give 4-[4-bromo-5-fluoro-2-(tetrahydropyran-2-yloxymethyl)phenyl]-2-methyl-3-butyn-2-ol.

[0656]

Step 4

The tert-butyl ((1S,2S,4R)-rel-7-(4'-cyano-3'-fluoro-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-[1,1'-biphenyl]-3-carbonyl)-7-azabicyclo[2.2.1]heptan-2-yl)carbamate-isomer-X (150 mg) obtained in step 2 of Example 207, and the 4-[4-bromo-5-fluoro-2-(tetrahydropyran-2-yloxymethyl)phenyl]-2-methyl-3-butyn-2-ol (129 mg) obtained in step 3 above were suspended in 1,4-dioxane (0.89 mL). At room temperature, Pd(dba)₂ (12.3 mg) and X-phos (20.4 mg) were added thereto, followed by stirring at 100°C for 1 hour. The reaction mixture was filtrated, and the solvent was distilled off. The residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate). The residue was dissolved in THF (0.92 mL) and water (0.46 mL). At room temperature, p-toluenesulfonic acid monohydrate (6.9 mg) was added thereto, followed by stirring at 70°C for 1 hour. Ethyl acetate was added thereto, and the mixture was washed sequentially with water and saturated brine, and dried over

anhydrous sodium sulfate. Then, the solvent was distilled off. The residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate) to give tert-butyl ((1S,2S,4R)-rel-7-(4''-cyano-2,3''-difluoro-4-(3-hydroxy-3-methyl-1-butyn-1-yl)-5-(hydroxymethyl)-[1,1':2',1''-terphenyl]-4'-carbonyl)-7-azabicyclo[2.2.1]heptan-2-yl) carbamate-isomer-X. Step 5

The tert-butyl ((1S,2S,4R)-rel-7-(4''-cyano-2,3''-difluoro-4-(3-hydroxy-3-methyl-1-butyn-1-yl)-5-(hydroxymethyl)-[1,1':2',1''-terphenyl]-4'-carbonyl)-7-azabicyclo[2.2.1]heptan-2-yl) carbamate-isomer-X (30 mg) obtained in step 4 above was dissolved in 1,4-dioxane (0.24 mL). At room temperature, a solution of 1.0 M TBAF in THF (0.14 mL) was added thereto, followed by stirring at 100°C for 1 hour. EtOH (0.12 mL) and 10% Pd/C (30 mg) were added to the reaction mixture, and after hydrogen substitution, the resulting mixture was stirred at 70°C for 30 minutes. The reaction mixture was filtrated, and the filtrate was concentrated. The residue was dissolved in THF. At room temperature, TEA (0.013 mL), DMAP (1.1 mg), and Boc₂O (20.4 mg) were added thereto, followed by stirring at 70°C for 1 hour. Ethyl acetate was added thereto, the mixture was washed 5 times with phosphoric acid at a concentration of about 0.5 mol/L, washed with saturated brine, and dried over anhydrous sodium sulfate. Thereafter, the solvent was distilled off. The residue was dissolved in acetonitrile (0.5 mL). At room temperature, a 4 N hydrochloric acid-1,4-dioxane solution (0.5 mL) was added thereto, followed by stirring at room temperature for 5 minutes. After the completion of the reaction was confirmed by LCMS, the solvent was distilled off. The residue was purified by reversed-phase HPLC (mobile phase: water/acetonitrile) to give the title compound.

[0657]

Example 227: Synthesis of 5'-(S)-3-aminopyrrolidine-1-carbonyl)-3-fluoro-2'-(6-fluoro-1-(2-hydroxy-2-methylpropyl)-1,3-dihydroisobenzofuran-5-yl)-[1,1'-biphenyl]-4-carbonitrile

Step 1

The tert-butyl 3-(4-cyano-3-fluoro-phenyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (400 mg) obtained in step 1 of Example 41, and the 4-[4-bromo-5-fluoro-2-(tetrahydropyran-2-yloxymethyl)phenyl]-2-methyl-3-butyn-2-ol (456 mg) obtained in step 3 of Example 226 were suspended in 1,4-dioxane (3.15 mL). At room temperature, Pd(dba)₂ (43.5 mg), X-phos (144 mg), and tripotassium phosphate (601 mg) were added thereto, followed by stirring at 100°C for 1 hour. The reaction mixture was filtrated, and the solvent was distilled off. The residue was purified by silica gel column chromatography (mobile phase: chloroform/methanol). The residue was dissolved in THF (1.62 mL). At room temperature, water (0.81 mL) and p-toluenesulfonic acid monohydrate (12.3 mg) were added thereto, followed by stirring at 70°C for 1 hour. Ethyl acetate was added thereto, and the mixture was washed sequentially with water and saturated brine, and dried over anhydrous sodium sulfate. Thereafter, the solvent was distilled off. The residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate) to give tert-butyl 3-(4-cyano-3-fluoro-phenyl)-4-[2-fluoro-5-(hydroxymethyl)-4-(3-hydroxy-3-methyl-1-butenyl)phenyl]benzoate.

[0658]

Step 2

The tert-butyl 3-(4-cyano-3-fluoro-phenyl)-4-[2-fluoro-5-(hydroxymethyl)-4-(3-hydroxy-3-methyl-1-butenyl)phenyl]benzoate (90 mg) obtained in step 1 above was dissolved in 1,4-dioxane (0.9 mL). At room temperature, a solution of 1.0 M TBAF in THF (0.54 mL) was added thereto, followed by stirring at 100°C for 2 hours. EtOH (0.30 mL) and 10% Pd/C (90 mg) were added to the reaction mixture, and after hydrogen substitution, stirring was performed at 70°C overnight. The reaction mixture was filtrated, and the solvent was distilled off. The residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate). The residue was dissolved in THF (1.0 mL). At room temperature, 12 N hydrochloric acid (0.5 mL) was added thereto,

followed by stirring at room temperature for 1.5 hours. MTBE was added thereto, and extraction was performed twice with a 2 N aqueous sodium hydroxide solution. The aqueous layer was acidified with 2 N hydrochloric acid, and extraction was performed twice with MTBE. The organic layer was sequentially washed with saturated brine, and dried over anhydrous sodium sulfate, and the solvent was distilled off to give 3-(4-cyano-3-fluoro-phenyl)-4-[6-fluoro-1-(2-hydroxy-2-methyl-propyl)-1,3-dihydroisobenzofuran-5-yl]benzoic acid.

[0659]

Step 3

The 3-(4-cyano-3-fluoro-phenyl)-4-[6-fluoro-1-(2-hydroxy-2-methyl-propyl)-1,3-dihydroisobenzofuran-5-yl]benzoic acid (10 mg) obtained in step 2 above was dissolved in THF (0.5 mL). At room temperature, HATU (9.31 mg), tert-butyl N-[(3S)-pyrrolidin-3-yl]carbamate (43.5 mg), and TEA (6.2 µL) were added thereto, followed by stirring at 50°C for 1 hour. The solvent was distilled off, and MeOH (0.5 mL) and a 4 N hydrochloric acid-1,4-dioxane solution (0.5 mL) were added thereto, followed by stirring at room temperature for 30 minutes. The solvent was distilled off, and the residue was purified by reversed-phase HPLC (mobile phase: water/acetonitrile) to give the title compound.

[0660]

Example 228: Synthesis of 5'-((1S,2S,4R)-rel-2-amino-7-azabicyclo[2.2.1]heptane-7-carbonyl)-3-fluoro-2'-(6-fluoro-1-(2-hydroxy-2-methylpropyl)-7-methyl-1H-benzo[d][1,2,3]triazol-5-yl)-[1,1'-biphenyl]-4-carbonitrile-isomer-X

The procedure of steps 1 to 5 in Example 209 was conducted using 1,3-difluoro-2-methyl-4-nitro-benzene instead of 2-chloro-1,3-difluoro-4-nitro-benzene to give the title compound.

[0661]

Example 229: Synthesis of (S)-5'-(3-aminopyrrolidine-1-carbonyl)-3-fluoro-2'-(5-fluoro-3-(2-hydroxy-2-methylpropyl)benzo[d]isoxazol-6-yl)-[1,1'-biphenyl]-4-

carbonitrile

Step 1

The 1-(4-bromo-5-fluoro-2-hydroxyphenyl)ethanone (150 mg) obtained in step 2 of Example 179 was dissolved in THF (3.2 mL). At -25°C, lithium diisopropylamide (1.0 M, a THF solution) (3.2 mL) was added thereto, followed by stirring at -25°C for 1 hour. The mixture was cooled to -40°C, and acetone (0.118 mL) was added thereto, followed by stirring at -40°C for 1 hour. After a phosphoric acid aqueous solution was added thereto, ethyl acetate was added thereto, and the mixture was washed sequentially with water and saturated brine. After the organic layer was dried over anhydrous sodium sulfate, the solvent was distilled off. The residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate) to give 1-(4-bromo-5-fluoro-2-hydroxy-phenyl)-3-hydroxy-3-methyl-butan-1-one.

[0662]

Step 2

The 1-(4-bromo-5-fluoro-2-hydroxy-phenyl)-3-hydroxy-3-methyl-butan-1-one (60 mg) obtained in step 1 above, hydroxylamine hydrochloride (28.6 mg), and sodium acetate (25.4 mg) were dissolved in methanol (0.69 mL), followed by stirring at 60°C overnight. MTBE was added thereto, and the mixture was washed sequentially with water and saturated brine. After the organic layer was dried over anhydrous sodium sulfate, the solvent was distilled off. The residue was dissolved in THF (0.69 mL), and N,N'-carbonyldiimidazole (36.8 mg) and TEA (0.037 mL) were added thereto, followed by stirring at 70°C for 1 hour. The solvent was distilled off, and the residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate) to give 1-(6-bromo-5-fluoro-1,2-benzooxazol-3-yl)-2-methyl-propan-2-ol.

[0663]

Step 3

The procedure of steps 1 to 3 in Example 37 was conducted using the 1-(6-bromo-5-fluoro-1,2-benzooxazol-3-yl)-2-

methyl-propan-2-ol obtained in step 2 above instead of 1-(4-bromo-3-fluoro-phenyl)-2-methyl-propan-2-ol to give the title compound.

[0664]

5 Example 230: Synthesis of 2-(5-(5-((1S,2S,4R)-rel-2-amino-7-azabicyclo[2.2.1]heptane-7-carbonyl)-4'-cyano-3'-fluoro-[1,1'-biphenyl]-2-yl)-6-fluoro-1H-indol-1-yl)acetamide

Step 1

The tert-butyl ((1S,2S,4R)-rel-7-(4'-cyano-3'-fluoro-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-[1,1'-biphenyl]-3-carbonyl)-7-azabicyclo[2.2.1]heptan-2-yl)carbamate (100 mg) obtained in step 1 of Example 213, and the ethyl 2-(5-bromo-6-fluoro-indol-1-yl)acetate (69.5 mg) obtained in step 1 of Example 207 were suspended in 1,4-dioxane (0.59 mL). At room temperature, Pd(dba)₂ (8.2 mg), X-phos (13.6 mg), and tripotassium phosphate (113 mg) were added thereto, followed by degassing and nitrogen substitution. Under a nitrogen atmosphere, stirring was performed at an external temperature of 100°C overnight. The solvent was distilled off, and the residue was purified by silica gel column chromatography (mobile phase: chloroform/methanol). The residue was dissolved in MeOH (1.0 mL), and a 5 N aqueous sodium hydroxide solution (1.0 mL) was added thereto, followed by stirring for 1 hour. MTBE was added thereto, and the aqueous layer was extracted. The aqueous layer was acidified with hydrochloric acid, MTBE was added thereto, and the mixture was washed sequentially with water and saturated brine. The organic layer was dried over anhydrous sodium sulfate, and the solvent was distilled off to give 2-(5-(5-((1S,2S,4R)-rel-2-((tert-butoxycarbonyl)amino)-7-azabicyclo[2.2.1]heptane-7-carbonyl)-4'-cyano-3'-fluoro-[1,1'-biphenyl]-2-yl)-6-fluoro-1H-indol-1-yl)acetic acid.

[0665]

Step 2

The 2-(5-(5-((1S,2S,4R)-rel-2-((tert-butoxycarbonyl)amino)-7-azabicyclo[2.2.1]heptane-7-carbonyl)-4'-

cyano-3'-fluoro-[1,1'-biphenyl]-2-yl)-6-fluoro-1H-indol-1-yl)acetic acid (10 mg) obtained in step 1 above was dissolved in THF (0.32 mL). Then, N,N'-carbonyldiimidazole (5.2 mg) was added thereto, and the mixture was stirred at room temperature for 20 minutes. Twenty-eight percent aqueous ammonia (0.06 mL) was added thereto, and the mixture was stirred at room temperature for 20 minutes. The solvent was distilled off, and acetonitrile (0.2 mL) and a 4 N hydrochloric acid-1,4-dioxane solution (0.2 mL) were added to the residue, followed by stirring for 30 minutes. The solvent was distilled off, and the residue was purified by reversed-phase HPLC (mobile phase: water/acetonitrile) to give the title compound.

[0666]

Example 231: Synthesis of 2-(5-(5-((1S,2S,4R)-rel-2-amino-7-azabicyclo[2.2.1]heptane-7-carbonyl)-4'-cyano-3'-fluoro-[1,1'-biphenyl]-2-yl)-6-fluoro-1H-indol-1-yl)-N-methylacetamide

The 2-(5-(5-((1S,2S,4R)-rel-2-((tert-butoxycarbonyl)amino)-7-azabicyclo[2.2.1]heptane-7-carbonyl)-4'-cyano-3'-fluoro-[1,1'-biphenyl]-2-yl)-6-fluoro-1H-indol-1-yl)acetic acid (10 mg) obtained in step 1 of Example 230 was dissolved in THF (0.064 mL). At room temperature, HATU (6.7 mg), methylamine hydrochloride (2.2 mg), and TEA (6.7 µL) were added thereto, followed by stirring at 50°C for 1 hour. The solvent was distilled off, and acetonitrile (1.0 mL) and a 4 N hydrochloric acid-1,4-dioxane solution (1.0 mL) were added to the residue, followed by stirring for 10 minutes. The solvent was distilled off, the residue was dissolved in DMSO, and purification was performed by reversed-phase HPLC (mobile phase: water/acetonitrile) to give the title compound.

[0667]

Example 232: Synthesis of 2-(5-(5-((1S,2S,4R)-rel-2-amino-7-azabicyclo[2.2.1]heptane-7-carbonyl)-4'-cyano-3'-fluoro-[1,1'-biphenyl]-2-yl)-6-fluoro-1H-indol-1-yl)-N,N-dimethylacetamide

The procedure of Example 231 was conducted using dimethylamine hydrochloride instead of methylamine hydrochloride

to give the title compound.

[0668]

Example 233: Synthesis of 2-(4'-((1S,2S,4R)-rel-2-amino-7-azabicyclo[2.2.1]heptane-7-carbonyl)-4'''-cyano-2,3'''-difluoro-
5 [1,1':2',1'''-terphenyl]-4-yl)acetamide

Step 1

2-(4-Bromo-3-fluoro-phenyl)acetic acid (600 mg) was dissolved in THF (10.3 mL). At room temperature, HATU (1.08 g), NH₄Cl (275.4 mg), and TEA (1.08 mL) were added thereto, followed
10 by stirring at 50°C for 1 hour. The solvent was distilled off, and the residue was purified by silica gel column chromatography (mobile phase: chloroform/methanol) to give 2-(4-bromo-3-fluoro-phenyl)acetamide.

Step 2

15 The procedure of Example 225 was conducted using the 2-(4-bromo-3-fluoro-phenyl)acetamide obtained in step 1 above instead of 3-[(5-bromo-6-fluoro-indazol-1-yl)methyl]pentan-3-ol to give the title compound.

[0669]

20 Example 234: Synthesis of 2-(4'-((1S,2S,4R)-rel-2-amino-7-azabicyclo[2.2.1]heptane-7-carbonyl)-4'''-cyano-2,3'''-difluoro-
[1,1':2',1'''-terphenyl]-4-yl)-N-methylacetamide

Step 1

2-(4-Bromo-3-fluoro-phenyl)acetic acid (600 mg) was dissolved in THF (10.3 mL). At room temperature, HATU (1.08 g),
25 methylamine (ca. 9.8 mol/L in MeOH) (0.525 mL), and TEA (1.08 mL) were added thereto, followed by stirring at 50°C for 1 hour. The solvent was distilled off, and the residue was purified by silica gel column chromatography (mobile phase: chloroform/methanol) to
30 give 2-(4-bromo-3-fluoro-phenyl)-N-methyl-acetamide.

Step 2

The procedure of Example 225 was conducted using the 2-(4-bromo-3-fluoro-phenyl)-N-methyl-acetamide obtained in step 1
above instead of 3-[(5-bromo-6-fluoro-indazol-1-yl)methyl]pentan-
35 3-ol to give the title compound.

[0670]

Example 235: Synthesis of 2-(4'-((1S,2S,4R)-rel-2-amino-7-azabicyclo[2.2.1]heptane-7-carbonyl)-4''-cyano-2,3''-difluoro-[1,1':2',1''-terphenyl]-4-yl)-N,N-dimethylacetamide

5 Step 1

2-(4-Bromo-3-fluoro-phenyl)acetic acid (600 mg) was dissolved in THF (10.3 mL). At room temperature, HATU (1.08 g), dimethylamine hydrochloride (419.9 mg), and TEA (1.08 mL) were added thereto, followed by stirring at 50°C for 1 hour. The
10 solvent was distilled off, and the residue was purified by silica gel column chromatography (mobile phase: chloroform/methanol) to give 2-(4-bromo-3-fluoro-phenyl)-N,N-dimethyl-acetamide.

Step 2

The procedure of Example 225 was conducted using the 2-
15 (4-bromo-3-fluoro-phenyl)-N,N-dimethyl-acetamide obtained in step 1 above instead of 3-[(5-bromo-6-fluoro-indazol-1-yl)methyl]pentan-3-ol to give the title compound.

[0671]

Example 236: Synthesis of 5-(5-((1S,2S,4R)-rel-2-amino-7-azabicyclo[2.2.1]heptane-7-carbonyl)-4'-cyano-3'-fluoro-[1,1'-biphenyl]-2-yl)-6-fluoro-1-methyl-1H-indole-3-carboxamide-isomer-X

The 5-(5-((1S,2S,4R)-rel-2-((tert-butoxycarbonyl)amino)-7-azabicyclo[2.2.1]heptane-7-carbonyl)-4'-
25 cyano-3'-fluoro-[1,1'-biphenyl]-2-yl)-6-fluoro-1-methyl-1H-indole-3-carboxylic acid-isomer-X (10 mg) obtained in step 3 of Example 224 was dissolved in THF (0.32 mL). Then, N,N'-carbonyldiimidazole (5.2 mg) was added thereto, and the mixture was stirred at room temperature for 20 minutes. Twenty-eight
30 percent aqueous ammonia (0.1 mL) was added thereto, and the mixture was stirred at room temperature for 20 minutes. The solvent was distilled off, and acetonitrile (0.2 mL) and a 4 N hydrochloric acid-1,4-dioxane solution (0.2 mL) were added to the residue, followed by stirring for 30 minutes. The solvent was
35 distilled off, and the residue was purified by reversed-phase

HPLC (mobile phase: water/acetonitrile) to give the title compound.

[0672]

Example 237: Synthesis of 5-(5-((1S,2S,4R)-rel-2-amino-7-azabicyclo[2.2.1]heptane-7-carbonyl)-4'-cyano-3'-fluoro-[1,1'-biphenyl]-2-yl)-6-fluoro-N,1-dimethyl-1H-indole-3-carboxamide-isomer-X

The 5-(5-((1S,2S,4R)-rel-2-((tert-butoxycarbonyl)amino)-7-azabicyclo[2.2.1]heptane-7-carbonyl)-4'-cyano-3'-fluoro-[1,1'-biphenyl]-2-yl)-6-fluoro-1-methyl-1H-indole-3-carboxylic acid-isomer-X (10 mg) obtained in step 3 of Example 224 was dissolved in THF (0.064 mL). At room temperature, HATU (6.7 mg), methylamine hydrochloride (2.2 mg), and TEA (6.7 µL) were added thereto, followed by stirring at 50°C for 1 hour. The solvent was distilled off, and acetonitrile (0.5 mL) and a 4 N hydrochloric acid-1,4-dioxane solution (0.5 mL) were added to the residue, followed by stirring for 10 minutes. The solvent was distilled off, the residue was dissolved in DMSO, and purification was performed by reversed-phase HPLC (mobile phase: water/acetonitrile) to give the title compound.

[0673]

Example 238: Synthesis of 5-(5-((1S,2S,4R)-rel-2-amino-7-azabicyclo[2.2.1]heptane-7-carbonyl)-4'-cyano-3'-fluoro-[1,1'-biphenyl]-2-yl)-6-fluoro-N,N,1-trimethyl-1H-indole-3-carboxamide-isomer-X

The procedure of Example 237 was conducted using dimethylamine hydrochloride instead of methylamine hydrochloride to give the title compound.

[0674]

Example 239: Synthesis of (S)-5'-(3-aminopyrrolidine-1-carbonyl)-3-fluoro-2'-(6-fluoro-1-(2-hydroxy-2-methylpropyl)-7-methyl-1H-benzo[d][1,2,3]triazol-5-yl)-[1,1'-biphenyl]-4-carbonitrile

The procedure of steps 1 to 5 of Example 209 was conducted using 1,3-difluoro-2-methyl-4-nitro-benzene instead of 2-chloro-1,3-difluoro-4-nitro-benzene, and using the tert-butyl

N-[(3S)-1-[3-(4-cyano-3-fluoro-phenyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoyl]pyrrolidin-3-yl]carbamate

obtained in step 1 of Example 37 instead of tert-butyl

((1S,2S,4R)-rel-7-(4'-cyano-3'-fluoro-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-[1,1'-biphenyl]-3-carbonyl)-7-

5 azabicyclo[2.2.1]heptan-2-yl)carbamate-isomer-X to give the title compound.

[0675]

Example 240: Synthesis of 5'-((3-endo)-3-amino-8-

10 azabicyclo[3.2.1]octane-8-carbonyl)-3-fluoro-2'-(6-fluoro-1-(2-hydroxy-2-methylpropyl)-7-methyl-1H-benzo[d][1,2,3]triazol-5-yl)-[1,1'-biphenyl]-4-carbonitrile

The procedure of steps 1 to 5 in Example 209 was conducted using 1,3-difluoro-2-methyl-4-nitro-benzene instead of

15 2-chloro-1,3-difluoro-4-nitro-benzene, and using the tert-butyl

N-[(3-endo)-8-[3-(4-cyano-3-fluoro-phenyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoyl]-8-

azabicyclo[3.2.1]octan-3-yl]carbamate obtained in step 3 of

Example 41 instead of tert-butyl ((1S,2S,4R)-rel-7-(4'-cyano-3'-

20 fluoro-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-[1,1'-biphenyl]-3-carbonyl)-7-azabicyclo[2.2.1]heptan-2-yl)carbamate-isomer-X to give the title compound.

[0676]

Example 241: Synthesis of 5'-((1S,2S,4R)-rel-2-amino-7-

25 azabicyclo[2.2.1]heptane-7-carbonyl)-3-fluoro-2'-(6-fluoro-1-(2-hydroxy-2-methylpropyl)-7-methyl-1H-benzo[d][1,2,3]triazol-5-yl)-[1,1'-biphenyl]-4-carbonitrile

The procedure of steps 1 to 5 in Example 209 was conducted using 1,3-difluoro-2-methyl-4-nitro-benzene instead of

30 2-chloro-1,3-difluoro-4-nitro-benzene, and using the tert-butyl

((1S,2S,4R)-rel-7-(4'-cyano-3'-fluoro-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-[1,1'-biphenyl]-3-carbonyl)-7-

azabicyclo[2.2.1]heptan-2-yl)carbamate obtained in step 1 of

Example 213 instead of tert-butyl ((1S,2S,4R)-rel-7-(4'-cyano-3'-

35 fluoro-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-[1,1'-

biphenyl]-3-carbonyl)-7-azabicyclo[2.2.1]heptan-2-yl) carbamate-isomer-X to give the title compound.

[0677]

Example 242: Synthesis of 5-(5-((1S,2S,4R)-rel-2-amino-7-

5 azabicyclo[2.2.1]heptane-7-carbonyl)-4'-cyano-3'-fluoro-[1,1'-
biphenyl]-2-yl)-6-fluoro-1-methyl-1H-indole-3-carboxylic acid

Step 1

The tert-butyl ((1S,2S,4R)-rel-7-(4'-cyano-3'-fluoro-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-[1,1'-biphenyl]-3-carbonyl)-7-azabicyclo[2.2.1]heptan-2-yl) carbamate (200 mg)
10 obtained in step 1 of Example 213 and 5-bromo-6-fluoro-1-methyl-indole (105.6 mg) were suspended in 1,4-dioxane (1.19 mL). At room temperature, Pd(dba)₂ (16.4 mg), X-phos (27.2 mg), and tripotassium phosphate (226.9 mg) were added thereto, followed by
15 degassing and nitrogen substitution. Under a nitrogen atmosphere, stirring was performed at an external temperature of 100°C overnight. The solvent was distilled off, and the residue was purified by silica gel column chromatography (mobile phase: chloroform/methanol) to give tert-butyl ((1S,2S,4R)-rel-7-(4'-
20 cyano-3'-fluoro-6-(6-fluoro-1-methyl-1H-indol-5-yl)-[1,1'-biphenyl]-3-carbonyl)-7-azabicyclo[2.2.1]heptan-2-yl) carbamate.

[0678]

Step 2

The tert-butyl ((1S,2S,4R)-rel-7-(4'-cyano-3'-fluoro-6-(6-fluoro-1-methyl-1H-indol-5-yl)-[1,1'-biphenyl]-3-carbonyl)-7-azabicyclo[2.2.1]heptan-2-yl) carbamate (209 mg) obtained in step
25 1 above was dissolved in DMF (3.6 mL). At room temperature, N-iodosuccinimide (121 mg) was added thereto, followed by stirring at room temperature for 1 hour. Ethyl acetate was added thereto,
30 and the resulting mixture was washed sequentially with water and saturated brine. After the organic layer was dried over anhydrous sodium sulfate, the solvent was distilled off. The residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate) to give tert-butyl ((1S,2S,4R)-rel-7-(4'-
35 cyano-3'-fluoro-6-(6-fluoro-3-iodo-1-methyl-1H-indol-5-yl)-[1,1'-

biphenyl]-3-carbonyl)-7-azabicyclo[2.2.1]heptan-2-yl) carbamate.
[0679]

Step 3

The tert-butyl ((1S,2S,4R)-rel-7-(4'-cyano-3'-fluoro-6-
5 (6-fluoro-3-iodo-1-methyl-1H-indol-5-yl)-[1,1'-biphenyl]-3-
carbonyl)-7-azabicyclo[2.2.1]heptan-2-yl) carbamate (211 mg)
obtained in step 2 above and PdCl₂(PPh₃)₂ (10.5 mg) were suspended
in 1-methyl-2-pyrrolidinone (2.11 mL). At room temperature, N,N-
diethylethanolamine (0.197 mL) was added thereto, and after CO
10 substitution, the mixture was stirred at 100°C for 1 hour. t-BuOH
(0.2 mL) and a 2 N aqueous sodium hydroxide solution (0.2 mL)
were added to the reaction mixture, and the resulting mixture was
stirred at room temperature overnight. MTBE was added thereto,
and the aqueous layer was separated. The aqueous layer was
15 acidified with hydrochloric acid, and extraction was performed
with MTBE. The organic layer was dried over anhydrous sodium
sulfate, and the solvent was distilled off to give 5-(5-
((1S,2S,4R)-rel-2-((tert-butoxycarbonyl)amino)-7-
azabicyclo[2.2.1]heptane-7-carbonyl)-4'-cyano-3'-fluoro-[1,1'-
20 biphenyl]-2-yl)-6-fluoro-1-methyl-1H-indole-3-carboxylic acid.
[0680]

Step 4

Acetonitrile (0.5 mL) and a 4 N hydrochloric acid-1,4-
dioxane solution (0.5 mL) were added to the 5-(5-((1S,2S,4R)-rel-
25 2-((tert-butoxycarbonyl)amino)-7-azabicyclo[2.2.1]heptane-7-
carbonyl)-4'-cyano-3'-fluoro-[1,1'-biphenyl]-2-yl)-6-fluoro-1-
methyl-1H-indole-3-carboxylic acid (10 mg) obtained in step 3
above, followed by stirring for 10 minutes. The reaction mixture
was concentrated, and the residue was purified by reversed-phase
30 HPLC (mobile phase: water/acetonitrile) to give the title
compound.

[0681]

Example 243: Synthesis of 5-(5-((1S,2S,4R)-rel-2-amino-7-
azabicyclo[2.2.1]heptane-7-carbonyl)-4'-cyano-3'-fluoro-[1,1'-
35 biphenyl]-2-yl)-6-fluoro-N,1-dimethyl-1H-indole-3-carboxamide

The procedure of Example 237 was conducted using the 5-(5-((1S,2S,4R)-rel-2-((tert-butoxycarbonyl)amino)-7-azabicyclo[2.2.1]heptane-7-carbonyl)-4'-cyano-3'-fluoro-[1,1'-biphenyl]-2-yl)-6-fluoro-1-methyl-1H-indole-3-carboxylic acid
5 obtained in step 3 of Example 242 instead of 5-(5-((1S,2S,4R)-rel-2-((tert-butoxycarbonyl)amino)-7-azabicyclo[2.2.1]heptane-7-carbonyl)-4'-cyano-3'-fluoro-[1,1'-biphenyl]-2-yl)-6-fluoro-1-methyl-1H-indole-3-carboxylic acid-isomer-X to give the title compound.

10 [0682]

Example 244: Synthesis of 5-(5-((1S,2S,4R)-rel-2-amino-7-azabicyclo[2.2.1]heptane-7-carbonyl)-4'-cyano-3'-fluoro-[1,1'-biphenyl]-2-yl)-6-fluoro-N,N,1-trimethyl-1H-indole-3-carboxamide

The procedure of Example 237 was conducted using the 5-(5-((1S,2S,4R)-rel-2-((tert-butoxycarbonyl)amino)-7-azabicyclo[2.2.1]heptane-7-carbonyl)-4'-cyano-3'-fluoro-[1,1'-biphenyl]-2-yl)-6-fluoro-1-methyl-1H-indole-3-carboxylic acid
15 obtained in step 3 of Example 242 instead of 5-(5-((1S,2S,4R)-rel-2-((tert-butoxycarbonyl)amino)-7-azabicyclo[2.2.1]heptane-7-carbonyl)-4'-cyano-3'-fluoro-[1,1'-biphenyl]-2-yl)-6-fluoro-1-methyl-1H-indole-3-carboxylic acid-isomer-X, and using dimethylamine hydrochloride instead of methylamine hydrochloride to give the title compound.

[0683]

25 Example 245: Synthesis of 5-(5-((1S,2S,4R)-rel-2-amino-7-azabicyclo[2.2.1]heptane-7-carbonyl)-4'-cyano-3'-fluoro-[1,1'-biphenyl]-2-yl)-6-fluoro-1-methyl-1H-indole-3-carboxamide

The procedure of Example 236 was conducted using the 5-(5-((1S,2S,4R)-rel-2-((tert-butoxycarbonyl)amino)-7-azabicyclo[2.2.1]heptane-7-carbonyl)-4'-cyano-3'-fluoro-[1,1'-biphenyl]-2-yl)-6-fluoro-1-methyl-1H-indole-3-carboxylic acid
30 obtained in step 3 of Example 242 instead of 5-(5-((1S,2S,4R)-rel-2-((tert-butoxycarbonyl)amino)-7-azabicyclo[2.2.1]heptane-7-carbonyl)-4'-cyano-3'-fluoro-[1,1'-biphenyl]-2-yl)-6-fluoro-1-methyl-1H-indole-3-carboxylic acid-isomer-X to give the title
35

compound.

[0684]

Example 246: Synthesis of 5'-((1S,2S,4R)-rel-2-amino-7-azabicyclo[2.2.1]heptane-7-carbonyl)-2'-(7-(difluoromethyl)-6-fluoro-1-(2-hydroxy-2-methylpropyl)-1H-benzo[d][1,2,3]triazol-5-yl)-3-fluoro-[1,1'-biphenyl]-4-carbonitrile-isomer-X

The procedure of steps 1 to 5 in Example 209 was conducted using 2-(difluoromethyl)-1,3-difluoro-4-nitro-benzene instead of 2-chloro-1,3-difluoro-4-nitro-benzene to give the title compound.

[0685]

Example 247: Synthesis of 5-(5-((1S,2S,4R)-rel-2-amino-7-azabicyclo[2.2.1]heptane-7-carbonyl)-4'-cyano-3'-fluoro-[1,1'-biphenyl]-2-yl)-6-fluoro-1-(2-hydroxy-2-methylpropyl)-1H-benzo[d][1,2,3]triazole-7-carbonitrile-isomer-X

The procedure of steps 1 to 5 in Example 209 was conducted using 2,6-difluoro-3-nitro-benzonitrile instead of 2-chloro-1,3-difluoro-4-nitro-benzene, and using THF instead of EtOH to give the title compound.

[0686]

Example 248: Synthesis of (S)-5'-(3-aminopyrrolidine-1-carbonyl)-2'-(7-(difluoromethyl)-6-fluoro-1-(2-hydroxy-2-methylpropyl)-1H-benzo[d][1,2,3]triazol-5-yl)-3-fluoro-[1,1'-biphenyl]-4-carbonitrile

The procedure of steps 1 to 5 in Example 209 was conducted using 2-(difluoromethyl)-1,3-difluoro-4-nitro-benzene instead of 2-chloro-1,3-difluoro-4-nitro-benzene, and using the tert-butyl N-[(3S)-1-[3-(4-cyano-3-fluoro-phenyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoyl]pyrrolidin-3-yl]carbamate obtained in step 1 of Example 37 instead of tert-butyl ((1S,2S,4R)-rel-7-(4'-cyano-3'-fluoro-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-[1,1'-biphenyl]-3-carbonyl)-7-azabicyclo[2.2.1]heptan-2-yl)carbamate-isomer-X to give the title compound.

[0687]

Example 249: Synthesis of (S)-5-(5-(3-aminopyrrolidine-1-carbonyl)-4'-cyano-3'-fluoro-[1,1'-biphenyl]-2-yl)-6-fluoro-1-(2-hydroxy-2-methylpropyl)-1H-benzo[d][1,2,3]triazole-7-carbonitrile

The procedure of steps 1 to 5 in Example 209 was
5 conducted using 2,6-difluoro-3-nitro-benzonitrile instead of 2-chloro-1,3-difluoro-4-nitro-benzene, using THF instead of EtOH, and using the tert-butyl N-[(3S)-1-[3-(4-cyano-3-fluoro-phenyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoyl]pyrrolidin-3-yl]carbamate obtained in step 1 of
10 Example 37 instead of tert-butyl ((1S,2S,4R)-rel-7-(4'-cyano-3'-fluoro-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-[1,1'-biphenyl]-3-carbonyl)-7-azabicyclo[2.2.1]heptan-2-yl)carbamate-isomer-X to give the title compound.

[0688]

15 Example 250: Synthesis of 5'-((3-endo)-3-amino-8-azabicyclo[3.2.1]octane-8-carbonyl)-2'-(7-(difluoromethyl)-6-fluoro-1-(2-hydroxy-2-methylpropyl)-1H-benzo[d][1,2,3]triazol-5-yl)-3-fluoro-[1,1'-biphenyl]-4-carbonitrile

The procedure of steps 1 to 5 in Example 209 was
20 conducted using 2-(difluoromethyl)-1,3-difluoro-4-nitro-benzene instead of 2-chloro-1,3-difluoro-4-nitro-benzene, and using the tert-butyl N-[(3-endo)-8-[3-(4-cyano-3-fluoro-phenyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoyl]-8-azabicyclo[3.2.1]octan-3-yl]carbamate obtained in step 3 of
25 Example 41 instead of tert-butyl ((1S,2S,4R)-rel-7-(4'-cyano-3'-fluoro-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-[1,1'-biphenyl]-3-carbonyl)-7-azabicyclo[2.2.1]heptan-2-yl)carbamate-isomer-X to give the title compound.

[0689]

30 Example 251: Synthesis of 5-(5-((3-endo)-3-amino-8-azabicyclo[3.2.1]octane-8-carbonyl)-4'-cyano-3'-fluoro-[1,1'-biphenyl]-2-yl)-6-fluoro-1-(2-hydroxy-2-methylpropyl)-1H-benzo[d][1,2,3]triazole-7-carbonitrile

The procedure of steps 1 to 5 in Example 209 was
35 conducted using 2,6-difluoro-3-nitro-benzonitrile instead of 2-

chloro-1,3-difluoro-4-nitro-benzene, using THF instead of EtOH, and using the tert-butyl N-[(3-endo)-8-[3-(4-cyano-3-fluorophenyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoyl]-8-azabicyclo[3.2.1]octan-3-yl]carbamate obtained in step 3 of
5 Example 41 instead of tert-butyl ((1S,2S,4R)-rel-7-(4'-cyano-3'-fluoro-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-[1,1'-biphenyl]-3-carbonyl)-7-azabicyclo[2.2.1]heptan-2-yl)carbamate-isomer-X to give the title compound.

[0690]

10 Example 252: Synthesis of 5'-((1S,2S,4R)-rel-2-amino-7-azabicyclo[2.2.1]heptane-7-carbonyl)-2'-(7-(difluoromethyl)-6-fluoro-1-(2-hydroxy-2-methylpropyl)-1H-benzo[d][1,2,3]triazol-5-yl)-3-fluoro-[1,1'-biphenyl]-4-carbonitrile

The procedure of steps 1 to 5 in Example 209 was
15 conducted using 2-(difluoromethyl)-1,3-difluoro-4-nitro-benzene instead of 2-chloro-1,3-difluoro-4-nitro-benzene, and using the tert-butyl ((1S,2S,4R)-rel-7-(4'-cyano-3'-fluoro-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-[1,1'-biphenyl]-3-carbonyl)-7-azabicyclo[2.2.1]heptan-2-yl)carbamate obtained in step 1 of
20 Example 213 instead of tert-butyl ((1S,2S,4R)-rel-7-(4'-cyano-3'-fluoro-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-[1,1'-biphenyl]-3-carbonyl)-7-azabicyclo[2.2.1]heptan-2-yl)carbamate-isomer-X to give the title compound.

[0691]

25 Example 253: Synthesis of 5-(5-((1S,2S,4R)-rel-2-amino-7-azabicyclo[2.2.1]heptane-7-carbonyl)-4'-cyano-3'-fluoro-[1,1'-biphenyl]-2-yl)-6-fluoro-1-(2-hydroxy-2-methylpropyl)-1H-benzo[d][1,2,3]triazole-7-carbonitrile

The procedure of steps 1 to 5 in Example 209 was
30 conducted using 2,6-difluoro-3-nitro-benzonitrile instead of 2-chloro-1,3-difluoro-4-nitro-benzene, using THF instead of EtOH, and using the tert-butyl ((1S,2S,4R)-rel-7-(4'-cyano-3'-fluoro-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-[1,1'-biphenyl]-3-carbonyl)-7-azabicyclo[2.2.1]heptan-2-yl)carbamate obtained in
35 step 1 of Example 213 instead of tert-butyl ((1S,2S,4R)-rel-7-

(4'-cyano-3'-fluoro-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-[1,1'-biphenyl]-3-carbonyl)-7-azabicyclo[2.2.1]heptan-2-yl)carbamate-isomer-X to give the title compound.

[0692]

5 Example 254: Synthesis of 5'-((1S,2S,4R)-rel-2-amino-7-azabicyclo[2.2.1]heptane-7-carbonyl)-2'-(6,7-difluoro-1-((1-hydroxycyclobutyl)methyl)-1H-benzo[d][1,2,3]triazol-5-yl)-3-fluoro-[1,1'-biphenyl]-4-carbonitrile-isomer-X

The procedure of steps 1 to 5 in Example 209 was
10 conducted using 1,2,3-trifluoro-4-nitro-benzene instead of 2-chloro-1,3-difluoro-4-nitro-benzene, and using 1-(aminomethyl)cyclobutanol instead of 1-amino-2-methyl-propan-2-ol to give the title compound.

[0693]

15 Example 255: Synthesis of (S)-5'-(3-aminopyrrolidine-1-carbonyl)-2'-(6,7-difluoro-1-((1-hydroxycyclobutyl)methyl)-1H-benzo[d][1,2,3]triazol-5-yl)-3-fluoro-[1,1'-biphenyl]-4-carbonitrile

The procedure of steps 1 to 5 in Example 209 was
20 conducted using 1,2,3-trifluoro-4-nitro-benzene instead of 2-chloro-1,3-difluoro-4-nitro-benzene, using 1-(aminomethyl)cyclobutanol instead of 1-amino-2-methyl-propan-2-ol, and using the tert-butyl N-[(3S)-1-[3-(4-cyano-3-fluorophenyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoyl]pyrrolidin-3-yl]carbamate obtained in step 1 of
25 Example 37 instead of tert-butyl ((1S,2S,4R)-rel-7-(4'-cyano-3'-fluoro-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-[1,1'-biphenyl]-3-carbonyl)-7-azabicyclo[2.2.1]heptan-2-yl)carbamate-isomer-X to give the title compound.

30 [0694]

Example 256: Synthesis of 5'-((3-endo)-3-amino-8-azabicyclo[3.2.1]octane-8-carbonyl)-2'-(6,7-difluoro-1-((1-hydroxycyclobutyl)methyl)-1H-benzo[d][1,2,3]triazol-5-yl)-3-fluoro-[1,1'-biphenyl]-4-carbonitrile

35 The procedure of steps 1 to 5 in Example 209 was

conducted using 1,2,3-trifluoro-4-nitro-benzene instead of 2-chloro-1,3-difluoro-4-nitro-benzene, using 1-(aminomethyl)cyclobutanol instead of 1-amino-2-methyl-propan-2-ol, and using the tert-butyl N-[(3-endo)-8-[3-(4-cyano-3-fluorophenyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoyl]-8-azabicyclo[3.2.1]octan-3-yl]carbamate obtained in step 3 of
5 Example 41 instead of tert-butyl ((1S,2S,4R)-rel-7-(4'-cyano-3'-fluoro-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-[1,1'-biphenyl]-3-carbonyl)-7-azabicyclo[2.2.1]heptan-2-yl)carbamate-
10 isomer-X to give the title compound.

[0695]

Example 257: Synthesis of 5'-((1S,2S,4R)-rel-2-amino-7-azabicyclo[2.2.1]heptane-7-carbonyl)-2'-(7-chloro-6-fluoro-1-((1-hydroxycyclobutyl)methyl)-1H-benzo[d][1,2,3]triazol-5-yl)-3-fluoro-[1,1'-biphenyl]-4-carbonitrile-isomer-X
15

The procedure of steps 1 to 5 in Example 209 was conducted using 1-(aminomethyl)cyclobutanol instead of 1-amino-2-methyl-propan-2-ol to give the title compound.

[0696]

Example 258: Synthesis of (S)-5'-(3-aminopyrrolidine-1-carbonyl)-2'-(7-chloro-6-fluoro-1-((1-hydroxycyclobutyl)methyl)-1H-benzo[d][1,2,3]triazol-5-yl)-3-fluoro-[1,1'-biphenyl]-4-carbonitrile
20

The procedure of steps 1 to 5 in Example 209 was
25 conducted using 1-(aminomethyl)cyclobutanol instead of 1-amino-2-methyl-propan-2-ol, and using the tert-butyl N-[(3S)-1-[3-(4-cyano-3-fluorophenyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoyl]pyrrolidin-3-yl]carbamate obtained in step 1 of
Example 37 instead of tert-butyl ((1S,2S,4R)-rel-7-(4'-cyano-3'-fluoro-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-[1,1'-biphenyl]-3-carbonyl)-7-azabicyclo[2.2.1]heptan-2-yl)carbamate-
30 isomer-X to give the title compound.

[0697]

Example 259: Synthesis of 5'-((3-endo)-3-amino-8-azabicyclo[3.2.1]octane-8-carbonyl)-2'-(7-chloro-6-fluoro-1-((1-
35

hydroxycyclobutyl)methyl)-1H-benzo[d][1,2,3]triazol-5-yl)-3-fluoro-[1,1'-biphenyl]-4-carbonitrile

The procedure of steps 1 to 5 in Example 209 was conducted using 1-(aminomethyl)cyclobutanol instead of 1-amino-2-methyl-propan-2-ol, and using the tert-butyl N-[(3-endo)-8-[3-(4-cyano-3-fluoro-phenyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoyl]-8-azabicyclo[3.2.1]octan-3-yl]carbamate obtained in step 3 of Example 41 instead of tert-butyl ((1S,2S,4R)-rel-7-(4'-cyano-3'-fluoro-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-[1,1'-biphenyl]-3-carbonyl)-7-azabicyclo[2.2.1]heptan-2-yl)carbamate-isomer-X to give the title compound.
[0698]

Example 260: Synthesis of 5'-((1S,2S,4R)-rel-2-amino-7-azabicyclo[2.2.1]heptane-7-carbonyl)-2'-(7-(difluoromethyl)-6-fluoro-1-((1-hydroxycyclobutyl)methyl)-1H-benzo[d][1,2,3]triazol-5-yl)-3-fluoro-[1,1'-biphenyl]-4-carbonitrile-isomer-X

The procedure of steps 1 to 5 in Example 209 was conducted using 2-(difluoromethyl)-1,3-difluoro-4-nitro-benzene instead of 2-chloro-1,3-difluoro-4-nitro-benzene, and using 1-(aminomethyl)cyclobutanol instead of 1-amino-2-methyl-propan-2-ol to give the title compound.
[0699]

Example 261: Synthesis of (S)-5'-(3-aminopyrrolidine-1-carbonyl)-2'-(7-(difluoromethyl)-6-fluoro-1-((1-hydroxycyclobutyl)methyl)-1H-benzo[d][1,2,3]triazol-5-yl)-3-fluoro-[1,1'-biphenyl]-4-carbonitrile

The procedure of steps 1 to 5 in Example 209 was conducted using 2-(difluoromethyl)-1,3-difluoro-4-nitro-benzene instead of 2-chloro-1,3-difluoro-4-nitro-benzene, using 1-(aminomethyl)cyclobutanol instead of 1-amino-2-methyl-propan-2-ol, and using the tert-butyl N-[(3S)-1-[3-(4-cyano-3-fluoro-phenyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoyl]pyrrolidin-3-yl]carbamate obtained in step 1 of Example 37 instead of tert-butyl ((1S,2S,4R)-rel-7-(4'-cyano-3'-fluoro-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-[1,1'-

biphenyl]-3-carbonyl)-7-azabicyclo[2.2.1]heptan-2-yl) carbamate-isomer-X to give the title compound.

[0700]

Example 262: Synthesis of 5'-((3-endo)-3-amino-8-

5 azabicyclo[3.2.1]octane-8-carbonyl)-2'-(7-(difluoromethyl)-6-fluoro-1-((1-hydroxycyclobutyl)methyl)-1H-benzo[d][1,2,3]triazol-5-yl)-3-fluoro-[1,1'-biphenyl]-4-carbonitrile

The procedure of steps 1 to 5 in Example 209 was conducted using 2-(difluoromethyl)-1,3-difluoro-4-nitro-benzene
10 instead of 2-chloro-1,3-difluoro-4-nitro-benzene, using 1-(aminomethyl)cyclobutanol instead of 1-amino-2-methyl-propan-2-ol, and using the tert-butyl N-[(3-endo)-8-[3-(4-cyano-3-fluorophenyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoyl]-8-azabicyclo[3.2.1]octan-3-yl]carbamate obtained in step 3 of
15 Example 41 instead of tert-butyl ((1S,2S,4R)-rel-7-(4'-cyano-3'-fluoro-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-[1,1'-biphenyl]-3-carbonyl)-7-azabicyclo[2.2.1]heptan-2-yl) carbamate-isomer-X to give the title compound.

[0701]

20 Example 263: Synthesis of 5'-((1S,2S,4R)-rel-2-amino-7-

azabicyclo[2.2.1]heptane-7-carbonyl)-2'-(7-(difluoromethyl)-6-fluoro-1-((1-hydroxycyclobutyl)methyl)-1H-benzo[d][1,2,3]triazol-5-yl)-3-fluoro-[1,1'-biphenyl]-4-carbonitrile

The procedure of steps 1 to 5 in Example 209 was
25 conducted using 2-(difluoromethyl)-1,3-difluoro-4-nitro-benzene instead of 2-chloro-1,3-difluoro-4-nitro-benzene, using 1-(aminomethyl)cyclobutanol instead of 1-amino-2-methyl-propan-2-ol, and using the tert-butyl ((1S,2S,4R)-rel-7-(4'-cyano-3'-fluoro-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-[1,1'-
30 biphenyl]-3-carbonyl)-7-azabicyclo[2.2.1]heptan-2-yl) carbamate obtained in step 1 of Example 213 instead of tert-butyl ((1S,2S,4R)-rel-7-(4'-cyano-3'-fluoro-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-[1,1'-biphenyl]-3-carbonyl)-7-azabicyclo[2.2.1]heptan-2-yl) carbamate-isomer-X to give the title
35 compound.

[0702]

Example 264: Synthesis of 5-(5-((1S,2S,4R)-rel-2-amino-7-azabicyclo[2.2.1]heptane-7-carbonyl)-4'-cyano-3'-fluoro-[1,1'-biphenyl]-2-yl)-6-fluoro-1-((1-hydroxycyclobutyl)methyl)-1H-benzo[d][1,2,3]triazole-7-carbonitrile-isomer-X

5 The procedure of steps 1 to 5 in Example 209 was conducted using 2,6-difluoro-3-nitro-benzonitrile instead of 2-chloro-1,3-difluoro-4-nitro-benzene, and using 1-(aminomethyl)cyclobutanol instead of 1-amino-2-methyl-propan-2-ol
10 to give the title compound.

[0703]

Example 265: Synthesis of (S)-5-(5-(3-aminopyrrolidine-1-carbonyl)-4'-cyano-3'-fluoro-[1,1'-biphenyl]-2-yl)-6-fluoro-1-((1-hydroxycyclobutyl)methyl)-1H-benzo[d][1,2,3]triazole-7-carbonitrile

15 The procedure of steps 1 to 5 in Example 209 was conducted using 2,6-difluoro-3-nitro-benzonitrile instead of 2-chloro-1,3-difluoro-4-nitro-benzene, using 1-(aminomethyl)cyclobutanol instead of 1-amino-2-methyl-propan-2-ol, and using the tert-butyl N-[(3S)-1-[3-(4-cyano-3-fluorophenyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoyl]pyrrolidin-3-yl]carbamate obtained in step 1 of
20 Example 37 instead of tert-butyl ((1S,2S,4R)-rel-7-(4'-cyano-3'-fluoro-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-[1,1'-biphenyl]-3-carbonyl)-7-azabicyclo[2.2.1]heptan-2-yl)carbamate-isomer-X to give the title compound.
25

[0704]

Example 266: Synthesis of 5-(5-((3-endo)-3-amino-8-azabicyclo[3.2.1]octane-8-carbonyl)-4'-cyano-3'-fluoro-[1,1'-biphenyl]-2-yl)-6-fluoro-1-((1-hydroxycyclobutyl)methyl)-1H-benzo[d][1,2,3]triazole-7-carbonitrile

30 The procedure of steps 1 to 5 in Example 209 was conducted using 2,6-difluoro-3-nitro-benzonitrile instead of 2-chloro-1,3-difluoro-4-nitro-benzene, using 1-(aminomethyl)cyclobutanol instead of 1-amino-2-methyl-propan-2-ol
35

ol, and using the tert-butyl N-[(3-endo)-8-[3-(4-cyano-3-fluoro-phenyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoyl]-8-azabicyclo[3.2.1]octan-3-yl]carbamate obtained in step 3 of Example 41 instead of tert-butyl ((1S,2S,4R)-rel-7-(4'-cyano-3'-fluoro-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-[1,1'-biphenyl]-3-carbonyl)-7-azabicyclo[2.2.1]heptan-2-yl)carbamate-isomer-X to give the title compound.

[0705]

Example 267: Synthesis of 5-(5-((1S,2S,4R)-rel-2-amino-7-azabicyclo[2.2.1]heptane-7-carbonyl)-4'-cyano-3'-fluoro-[1,1'-biphenyl]-2-yl)-6-fluoro-1-((1-hydroxycyclobutyl)methyl)-1H-benzo[d][1,2,3]triazole-7-carbonitrile

The procedure of steps 1 to 5 in Example 209 was conducted using 2,6-difluoro-3-nitro-benzonitrile instead of 2-chloro-1,3-difluoro-4-nitro-benzene, using 1-(aminomethyl)cyclobutanol instead of 1-amino-2-methyl-propan-2-ol, and using the tert-butyl ((1S,2S,4R)-rel-7-(4'-cyano-3'-fluoro-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-[1,1'-biphenyl]-3-carbonyl)-7-azabicyclo[2.2.1]heptan-2-yl)carbamate obtained in step 1 of Example 213 instead of tert-butyl ((1S,2S,4R)-rel-7-(4'-cyano-3'-fluoro-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-[1,1'-biphenyl]-3-carbonyl)-7-azabicyclo[2.2.1]heptan-2-yl)carbamate-isomer-X to give the title compound.

[0706]

Example 268: Synthesis of 5'-((1S,2S,4R)-rel-2-amino-7-azabicyclo[2.2.1]heptane-7-carbonyl)-2'-(6,7-difluoro-1-((1-hydroxycyclobutyl)methyl)-1H-benzo[d][1,2,3]triazol-5-yl)-3-fluoro-[1,1'-biphenyl]-4-carbonitrile

The procedure of steps 1 to 5 in Example 209 was conducted using 1,2,3-trifluoro-4-nitro-benzene instead of 2-chloro-1,3-difluoro-4-nitro-benzene, using 1-(aminomethyl)cyclobutanol instead of 1-amino-2-methyl-propan-2-ol, and using the tert-butyl ((1S,2S,4R)-rel-7-(4'-cyano-3'-fluoro-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-[1,1'-

biphenyl]-3-carbonyl)-7-azabicyclo[2.2.1]heptan-2-yl) carbamate obtained in step 1 of Example 213 instead of tert-butyl ((1S,2S,4R)-rel-7-(4'-cyano-3'-fluoro-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-[1,1'-biphenyl]-3-carbonyl)-7-azabicyclo[2.2.1]heptan-2-yl) carbamate-isomer-X to give the title compound.

[0707]

Example 269: Synthesis of 5'-((1S,2S,4R)-rel-2-amino-7-azabicyclo[2.2.1]heptane-7-carbonyl)-2'-(7-chloro-6-fluoro-1-((1-hydroxycyclobutyl)methyl)-1H-benzo[d][1,2,3]triazol-5-yl)-3-fluoro-[1,1'-biphenyl]-4-carbonitrile

The procedure of steps 1 to 5 in Example 209 was conducted using 1-(aminomethyl)cyclobutanol instead of 1-amino-2-methyl-propan-2-ol, and using the tert-butyl ((1S,2S,4R)-rel-7-(4'-cyano-3'-fluoro-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-[1,1'-biphenyl]-3-carbonyl)-7-azabicyclo[2.2.1]heptan-2-yl) carbamate obtained in step 1 of Example 213 instead of tert-butyl ((1S,2S,4R)-rel-7-(4'-cyano-3'-fluoro-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-[1,1'-biphenyl]-3-carbonyl)-7-azabicyclo[2.2.1]heptan-2-yl) carbamate-isomer-X to give the title compound.

[0708]

Example 270: Synthesis of 5'-((1S,2S,4R)-rel-2-amino-7-azabicyclo[2.2.1]heptane-7-carbonyl)-3-fluoro-2'-(6-fluoro-1-((1-hydroxycyclobutyl)methyl)-7-methyl-1H-benzo[d][1,2,3]triazol-5-yl)-[1,1'-biphenyl]-4-carbonitrile-isomer-X

The procedure of steps 1 to 5 in Example 209 was conducted using 1,3-difluoro-2-methyl-4-nitro-benzene instead of 2-chloro-1,3-difluoro-4-nitro-benzene, and using 1-(aminomethyl)cyclobutanol instead of 1-amino-2-methyl-propan-2-ol to give the title compound.

[0709]

Example 271: Synthesis of (S)-5'-(3-aminopyrrolidine-1-carbonyl)-3-fluoro-2'-(6-fluoro-1-((1-hydroxycyclobutyl)methyl)-7-methyl-1H-benzo[d][1,2,3]triazol-5-yl)-[1,1'-biphenyl]-4-carbonitrile

The procedure of steps 1 to 5 in Example 209 was conducted using 1,3-difluoro-2-methyl-4-nitro-benzene instead of 2-chloro-1,3-difluoro-4-nitro-benzene, using 1-(aminomethyl)cyclobutanol instead of 1-amino-2-methyl-propan-2-ol, and using the tert-butyl N-[(3S)-1-[3-(4-cyano-3-fluorophenyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoyl]pyrrolidin-3-yl]carbamate obtained in step 1 of Example 37 instead of tert-butyl ((1S,2S,4R)-rel-7-(4'-cyano-3'-fluoro-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-[1,1'-biphenyl]-3-carbonyl)-7-azabicyclo[2.2.1]heptan-2-yl)carbamate-isomer-X to give the title compound.

[0710]

Example 272: Synthesis of 5'-((3-endo)-3-amino-8-azabicyclo[3.2.1]octane-8-carbonyl)-3-fluoro-2'-(6-fluoro-1-((1-hydroxycyclobutyl)methyl)-7-methyl-1H-benzo[d][1,2,3]triazol-5-yl)-[1,1'-biphenyl]-4-carbonitrile

The procedure of steps 1 to 5 in Example 209 was conducted using 1,3-difluoro-2-methyl-4-nitro-benzene instead of 2-chloro-1,3-difluoro-4-nitro-benzene, using 1-(aminomethyl)cyclobutanol instead of 1-amino-2-methyl-propan-2-ol, and using the tert-butyl N-[(3-endo)-8-[3-(4-cyano-3-fluorophenyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoyl]-8-azabicyclo[3.2.1]octan-3-yl]carbamate obtained in step 3 of Example 41 instead of tert-butyl ((1S,2S,4R)-rel-7-(4'-cyano-3'-fluoro-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-[1,1'-biphenyl]-3-carbonyl)-7-azabicyclo[2.2.1]heptan-2-yl)carbamate-isomer-X to give the title compound.

[0711]

Example 273: Synthesis of 5'-((1S,2S,4R)-rel-2-amino-7-azabicyclo[2.2.1]heptane-7-carbonyl)-2'-(7-bromo-6-fluoro-1-(2-hydroxy-2-methylpropyl)-1H-benzo[d][1,2,3]triazol-5-yl)-3-fluoro-[1,1'-biphenyl]-4-carbonitrile-isomer-X

Step 1

2-Bromo-1,3-difluoro-4-nitro-benzene (3 g) was dissolved in THF (31.5 mL). TEA (2.6 mL) and 1-amino-2-methyl-

propan-2-ol (1.4 mL) were added thereto, followed by stirring at room temperature for 1 hour. Ethyl acetate was added thereto, and the resulting mixture was washed sequentially with water and saturated brine. After the organic layer was dried over anhydrous sodium sulfate, the solvent was distilled off. The residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate) to give 1-(2-bromo-3-fluoro-6-nitro-anilino)-2-methyl-propan-2-ol.

[0712]

10 Step 2

The 1-(2-bromo-3-fluoro-6-nitro-anilino)-2-methyl-propan-2-ol (1.03 g) obtained in step 1 above was dissolved in acetic acid (6.7 mL). At room temperature, N-iodosuccinimide (981 mg) was added thereto, followed by stirring at 50°C for 3 hours. MTBE and water were added thereto, and extraction was performed twice with MTBE. The combined organic layers were dried over anhydrous sodium sulfate, and the solvent was distilled off. The residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate) to give 1-(2-bromo-3-fluoro-4-iodo-6-nitro-anilino)-2-methyl-propan-2-ol.

[0713]

Step 3

The 1-(2-bromo-3-fluoro-4-iodo-6-nitro-anilino)-2-methyl-propan-2-ol (1.33 g) obtained in step 2 above and iron (1.33 g) were dissolved in THF (10.2 mL) and a 2 N hydrochloric acid (10.2 mL), followed by stirring at 60°C for 1 hour. MTBE was added thereto, and the mixture was passed through Celite. MTBE was added thereto, and the mixture was washed sequentially with water and saturated brine, and dried over anhydrous sodium sulfate. Thereafter, the solvent was distilled off. The residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate) to give 1-(6-amino-2-bromo-3-fluoro-4-iodo-anilino)-2-methyl-propan-2-ol.

[0714]

35 Step 4

The 1-(6-amino-2-bromo-3-fluoro-4-iodo-anilino)-2-methyl-propan-2-ol (940 mg) obtained in step 3 above was dissolved in water (1.88 mL) and THF (4.7 mL). At 0°C, 12 N hydrochloric acid (2.82 mL) and an aqueous sodium nitrite solution (an aqueous solution obtained by dissolving 209 mg of sodium nitrite in 0.63 mL of water) were added thereto dropwise, followed by stirring at room temperature for 1 hour. MTBE was added thereto, and the mixture was washed sequentially with water and saturated brine, and dried over anhydrous sodium sulfate. Thereafter, the solvent was distilled off. The residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate) to give 1-(7-bromo-6-fluoro-5-iodo-benzotriazol-1-yl)-2-methyl-propan-2-ol.

[0715]

Step 5

The tert-butyl ((1S,2S,4R)-rel-7-(4'-cyano-3'-fluoro-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-[1,1'-biphenyl]-3-carbonyl)-7-azabicyclo[2.2.1]heptan-2-yl)carbamate-isomer-X (50 mg) obtained in step 2 of Example 207 and the 1-(7-bromo-6-fluoro-5-iodo-benzotriazol-1-yl)-2-methyl-propan-2-ol (47.9 mg) obtained in step 4 above were dissolved in 1,4-dioxane (0.3 mL). At room temperature, Pd(dba)₂ (4.1 mg), X-phos (6.8 mg), and tripotassium phosphate (56.7 mg) were added thereto. After nitrogen substitution, the mixture was stirred at 90°C overnight. Ethyl acetate was added thereto, and the mixture was put on NH-silica gel, and washed with ethyl acetate:methanol = 10:1. The solvent was distilled off, and acetonitrile (1.0 mL) and a 4 N hydrochloric acid-1,4-dioxane solution (1.0 mL) were added to the residue, followed by stirring for 10 minutes. The solvent was distilled off, the residue was dissolved in DMSO, and purification was performed by reversed-phase HPLC (mobile phase: water/acetonitrile) to give the title compound.

[0716]

Example 274: Synthesis of (S)-5'-(3-aminopyrrolidine-1-carbonyl)-2'-(7-bromo-6-fluoro-1-(2-hydroxy-2-methylpropyl)-1H-

benzo[d][1,2,3]triazol-5-yl)-3-fluoro-[1,1'-biphenyl]-4-carbonitrile

The procedure of steps 1 to 5 in Example 273 was conducted using the tert-butyl N-[(3S)-1-[3-(4-cyano-3-fluoro-phenyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoyl]pyrrolidin-3-yl]carbamate obtained in step 1 of Example 37 instead of tert-butyl ((1S,2S,4R)-rel-7-(4'-cyano-3'-fluoro-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-[1,1'-biphenyl]-3-carbonyl)-7-azabicyclo[2.2.1]heptan-2-yl)carbamate-isomer-X to give the title compound.

[0717]

Example 275: Synthesis of 5'-((3-endo)-3-amino-8-azabicyclo[3.2.1]octane-8-carbonyl)-2'-(7-bromo-6-fluoro-1-(2-hydroxy-2-methylpropyl)-1H-benzo[d][1,2,3]triazol-5-yl)-3-fluoro-[1,1'-biphenyl]-4-carbonitrile

The procedure of steps 1 to 5 in Example 273 was conducted using the tert-butyl N-[(3-endo)-8-[3-(4-cyano-3-fluoro-phenyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoyl]-8-azabicyclo[3.2.1]octan-3-yl]carbamate obtained in step 3 of Example 41 instead of tert-butyl ((1S,2S,4R)-rel-7-(4'-cyano-3'-fluoro-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-[1,1'-biphenyl]-3-carbonyl)-7-azabicyclo[2.2.1]heptan-2-yl)carbamate-isomer-X to give the title compound.

[0718]

Example 276: Synthesis of 5'-((1S,2S,4R)-rel-2-amino-7-azabicyclo[2.2.1]heptane-7-carbonyl)-3-fluoro-2'-(6-fluoro-1-((1-hydroxycyclobutyl)methyl)-7-methyl-1H-benzo[d][1,2,3]triazol-5-yl)-[1,1'-biphenyl]-4-carbonitrile

The procedure of steps 1 to 5 in Example 209 was conducted using 1,3-difluoro-2-methyl-4-nitro-benzene instead of 2-chloro-1,3-difluoro-4-nitro-benzene, using 1-(aminomethyl)cyclobutanol instead of 1-amino-2-methyl-propan-2-ol, and using the tert-butyl ((1S,2S,4R)-rel-7-(4'-cyano-3'-fluoro-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-[1,1'-biphenyl]-3-carbonyl)-7-azabicyclo[2.2.1]heptan-2-yl)carbamate

obtained in step 1 of Example 213 instead of tert-butyl ((1S,2S,4R)-rel-7-(4'-cyano-3'-fluoro-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-[1,1'-biphenyl]-3-carbonyl)-7-azabicyclo[2.2.1]heptan-2-yl)carbamate-isomer-X to give the title compound.

[0719]

Example 277: Synthesis of 5'-((1S,2S,4R)-rel-2-amino-7-azabicyclo[2.2.1]heptane-7-carbonyl)-2'-(7-bromo-6-fluoro-1-(2-hydroxy-2-methylpropyl)-1H-benzo[d][1,2,3]triazol-5-yl)-3-fluoro-[1,1'-biphenyl]-4-carbonitrile

The procedure of steps 1 to 5 in Example 273 was conducted using the tert-butyl ((1S,2S,4R)-rel-7-(4'-cyano-3'-fluoro-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-[1,1'-biphenyl]-3-carbonyl)-7-azabicyclo[2.2.1]heptan-2-yl)carbamate obtained in step 1 of Example 213 instead of tert-butyl ((1S,2S,4R)-rel-7-(4'-cyano-3'-fluoro-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-[1,1'-biphenyl]-3-carbonyl)-7-azabicyclo[2.2.1]heptan-2-yl)carbamate-isomer-X to give the title compound.

[0720]

Example 278: Synthesis of 5'-((1R,2R,4S)-rel-2-amino-7-azabicyclo[2.2.1]heptane-7-carbonyl)-2'-(7-cyclopropyl-6-fluoro-1-(2-hydroxy-2-methylpropyl)-1H-benzo[d][1,2,3]triazol-5-yl)-3-fluoro-[1,1'-biphenyl]-4-carbonitrile

Step 1

The 1-(2-bromo-3-fluoro-6-nitro-anilino)-2-methylpropan-2-ol (975 mg) obtained in step 1 of Example 273 was dissolved in 1,4-dioxane (10.6 mL). At room temperature, dichlorobis(tricyclohexylphosphine)palladium(II) (234 mg), cyclopropyl boronic acid (464 mg), and tripotassium phosphate (2.02 g) were added thereto, followed by stirring at 10°C overnight. The solvent was distilled off, and the residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate) to give 1-((2-cyclopropyl-3-fluoro-6-nitrophenyl)amino)-2-methylpropan-2-ol.

[0721]

Step 2

The 1-((2-cyclopropyl-3-fluoro-6-nitrophenyl)amino)-2-methylpropan-2-ol (204 mg) obtained in step 1 above was dissolved in acetonitrile (1.5 mL). At room temperature, NBS (196 mg) was added thereto, followed by stirring at 50°C for 1 hour. The solvent was distilled off, and the residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate) to give 1-((4-bromo-2-cyclopropyl-3-fluoro-6-nitrophenyl)amino)-2-methylpropan-2-ol.

[0722]

Step 3

The 1-((4-bromo-2-cyclopropyl-3-fluoro-6-nitrophenyl)amino)-2-methylpropan-2-ol (250 mg) obtained in step 2 above and iron (250 mg) were dissolved in THF (2.4 mL) and a 2 N hydrochloric acid (2.4 mL), followed by stirring at 60°C for 1 hour. Ethyl acetate was added thereto, and the mixture was passed through Celite. Ethyl acetate was added thereto, and the mixture was washed sequentially with water and saturated brine, and dried over anhydrous sodium sulfate. Thereafter, the solvent was distilled off. The residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate) to give 1-((6-amino-4-bromo-2-cyclopropyl-3-fluorophenyl)amino)-2-methylpropan-2-ol.

[0723]

Step 4

The 1-((6-amino-4-bromo-2-cyclopropyl-3-fluorophenyl)amino)-2-methylpropan-2-ol (192 mg) obtained in step 3 above was dissolved in THF (2.0 mL) and a 2 N hydrochloric acid (2.0 mL). An aqueous sodium nitrite solution (an aqueous solution obtained by dissolving 54 mg of sodium nitrite in 0.16 mL of water) was added thereto dropwise, followed by stirring at room temperature for 1 hour. Ethyl acetate was added thereto, and the mixture was washed sequentially with water and saturated brine, and dried over anhydrous sodium sulfate. Thereafter, the solvent

was distilled off. The residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate) to give 1-(5-bromo-7-cyclopropyl-6-fluoro-1H-benzo[d][1,2,3]triazol-1-yl)-2-methylpropan-2-ol.

5 [0724]

Step 5

The tert-butyl ((1S,2S,4R)-rel-7-(4'-cyano-3'-fluoro-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-[1,1'-biphenyl]-3-carbonyl)-7-azabicyclo[2.2.1]heptan-2-yl)carbamate (15 mg)
10 obtained in step 1 of Example 213, and the 1-(5-bromo-7-cyclopropyl-6-fluoro-1H-benzo[d][1,2,3]triazol-1-yl)-2-methylpropan-2-ol (10.5 mg) obtained in step 4 above were dissolved in 1,4-dioxane (0.2 mL). Pd(dba)₂ (1.2 mg), X-phos (2.0 mg), and tripotassium phosphate (17 mg) were added thereto. After
15 nitrogen substitution, the mixture was stirred at 100°C overnight. Ethyl acetate was added thereto, and the mixture was put on NH-silica gel, and washed with ethyl acetate:methanol = 10:1. The solvent was distilled off, and acetonitrile (1.0 mL) and a 4 N hydrochloric acid-1,4-dioxane solution (1.0 mL) were
20 added to the residue, followed by stirring for 10 minutes. The solvent was distilled off, the residue was dissolved in DMSO, and purification was performed by reversed-phase HPLC (mobile phase: water/acetonitrile) to give the title compound.

[0725]

25 Example 279: Synthesis of 5'-((1R,2R,4S)-rel-2-amino-7-azabicyclo[2.2.1]heptane-7-carbonyl)-2'-(7-cyclopropyl-6-fluoro-1-(2-hydroxy-2-methylpropyl)-1H-benzo[d][1,2,3]triazol-5-yl)-3-fluoro-[1,1'-biphenyl]-4-carbonitrile-isomer-X

[0726]

30 The procedure of steps 1 to 5 in Example 278 was conducted using the tert-butyl ((1S,2S,4R)-rel-7-(4'-cyano-3'-fluoro-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-[1,1'-biphenyl]-3-carbonyl)-7-azabicyclo[2.2.1]heptan-2-yl)carbamate-isomer-X obtained in step 2 of Example 207 instead of tert-butyl
35 ((1S,2S,4R)-rel-7-(4'-cyano-3'-fluoro-6-(4,4,5,5-tetramethyl-

1,3,2-dioxaborolan-2-yl)-[1,1'-biphenyl]-3-carbonyl)-7-azabicyclo[2.2.1]heptan-2-yl)carbamate to give the title compound.

[0727]

5 Example 280: Synthesis of (S)-5'-(3-aminopyrrolidine-1-carbonyl)-2'-(7-cyclopropyl-6-fluoro-1-(2-hydroxy-2-methylpropyl)-1H-benzo[d][1,2,3]triazol-5-yl)-3-fluoro-[1,1'-biphenyl]-4-carbonitrile

The procedure of steps 1 to 5 in Example 278 was
10 conducted using the tert-butyl N-[(3S)-1-[3-(4-cyano-3-fluorophenyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoyl]pyrrolidin-3-yl]carbamate obtained in step 1 of Example 37 instead of tert-butyl ((1S,2S,4R)-rel-7-(4'-cyano-3'-fluoro-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-[1,1'-
15 biphenyl]-3-carbonyl)-7-azabicyclo[2.2.1]heptan-2-yl)carbamate to give the title compound.

[0728]

Example 281: Synthesis of 5'-((3-endo)-3-amino-8-azabicyclo[3.2.1]octane-8-carbonyl)-2'-(7-cyclopropyl-6-fluoro-1-(2-hydroxy-2-methylpropyl)-1H-benzo[d][1,2,3]triazol-5-yl)-3-fluoro-[1,1'-biphenyl]-4-carbonitrile

The procedure of steps 1 to 5 in Example 278 was
conducted using the tert-butyl N-[(3-endo)-8-[3-(4-cyano-3-fluorophenyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-
25 yl)benzoyl]-8-azabicyclo[3.2.1]octan-3-yl]carbamate obtained in step 3 of Example 41 instead of tert-butyl ((1S,2S,4R)-rel-7-(4'-cyano-3'-fluoro-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-[1,1'-biphenyl]-3-carbonyl)-7-azabicyclo[2.2.1]heptan-2-yl)carbamate to give the title compound.

30 [0729]

Example 282: Synthesis of (S)-5'-((3-aminopyrrolidin-1-yl)methyl)-4'''-methyl-[1,1':2',1'''-terphenyl]-4-carbonitrile
Step 1

3-Bromo-4-chloro-benzaldehyde (100 mg) and tert-butyl
35 N-[(3S)-pyrrolidin-3-yl]carbamate (93.4 mg) were dissolved in

MeOH (1.0 mL). At 0°C, acetic acid (0.1 mL) and borane-2-picoline complex (146 mg) were added thereto, followed by stirring at room temperature overnight. Ethyl acetate was added thereto, and the resulting mixture was washed sequentially with water and saturated brine, and dried over anhydrous sodium sulfate. The solvent was then distilled off. The residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate) to give tert-butyl N-[(3S)-1-[(3-bromo-4-chloro-phenyl)methyl]pyrrolidin-3-yl]carbamate.

[0730]

Step 2

The tert-butyl N-[(3S)-1-[(3-bromo-4-chloro-phenyl)methyl]pyrrolidin-3-yl]carbamate (60 mg) obtained in step 1 above and (4-cyanophenyl)boronic acid (24.9 mg) were dissolved in 1,4-dioxane (0.77 mL). At room temperature, PdCl₂(dppf)CH₂Cl₂ (3.4 mg) and tripotassium phosphate (97.9 mg) were added thereto, followed by stirring at 125°C for 45 minutes. The reaction mixture was filtrated, and the solvent was distilled off. The residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate) to give tert-butyl N-[(3S)-1-[[4-chloro-3-(4-cyanophenyl)phenyl]methyl]pyrrolidin-3-yl]carbamate.

[0731]

Step 3

The tert-butyl N-[(3S)-1-[[4-chloro-3-(4-cyanophenyl)phenyl]methyl]pyrrolidin-3-yl]carbamate (20 mg) obtained in step 2 above, p-tolylboronic acid (13.2 mg), Pd₂(dba)₃ (2.23 mg), tripotassium phosphate (20.6 mg), and a solution of 1 M PCy₃ in THF (0.1 mL) were suspended in 1,4-dioxane (0.5 mL), followed by stirring at 160°C for 45 minutes. The reaction mixture was filtered through NH-silica gel, and the solvent of the filtrate was distilled off. The residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate) to give tert-butyl N-[(3S)-1-[[3-(4-cyanophenyl)-4-(p-tolyl)phenyl]methyl]pyrrolidin-3-yl]carbamate.

[0732]

Step 4

The tert-butyl N-[(3S)-1-[[3-(4-cyanophenyl)-4-(p-tolyl)phenyl]methyl]pyrrolidin-3-yl]carbamate (10 mg) obtained in step 3 above was dissolved in TFA (0.3 mL), and the progress of the reaction was confirmed by LCMS, followed by vacuum concentration. The residue was purified by reversed-phase HPLC (mobile phase: water/acetonitrile) to give the title compound. [0733]

Example 283: Synthesis of 5'-(((1S,2S,4R)-rel-2-amino-7-azabicyclo[2.2.1]heptan-7-yl)methyl)-2'',3-difluoro-4''-methyl-[1,1':2',1''-terphenyl]-4-carbonitrile-isomer-X

Step 1

3-Bromo-4-chloro-benzaldehyde (1.1 g) was dissolved in 1,4-dioxane (13 mL). At room temperature, (4-cyano-3-fluorophenyl)boronic acid (870 mg), PdCl₂(dppf)CH₂Cl₂ (110 mg), and a 2 M aqueous sodium carbonate solution (6.3 mL) were added thereto, followed by stirring at 90°C for 5 hours. Ethyl acetate was added thereto, and the resulting mixture was washed sequentially with water and saturated brine. After the organic layer was dried over anhydrous sodium sulfate, the solvent was distilled off. The residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate) to give 4-(2-chloro-5-formylphenyl)-2-fluorobenzonitrile. [0734]

Step 2

The 4-(2-chloro-5-formylphenyl)-2-fluorobenzonitrile (505 mg) obtained in step 1 above was dissolved in 1,4-dioxane (19.45 mL). At room temperature, (2-fluoro-4-methylphenyl)boronic acid (599 mg), Pd₂dba₃ (89 mg), a solution of 1 M PCy₃ in THF (0.1 mL), and tripotassium phosphate (1.24 g) were added thereto, and the mixture was stirred in a microwave reactor at 160°C for 1 hour. The solvent was distilled off, and the residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate) to give 2-fluoro-4-[2-(2-fluoro-4-methylphenyl)-5-formylphenyl]benzonitrile.

[0735]

Step 3

The 2-fluoro-4-[2-(2-fluoro-4-methyl-phenyl)-5-formyl-phenyl]benzonitrile (10 mg) obtained in step 2 above was
5 dissolved in DCM (0.6 mL). At room temperature, the tert-butyl
((1S,2S,4R)-rel-7-azabicyclo[2.2.1]heptan-2-yl)carbamate-isomer-X
(9.55 mg) synthesized in step 2 of Example 202 and sodium
triacetoxyborohydride (25.4 mg) were added thereto, followed by
stirring at room temperature for 1 hour. The solvent was
10 distilled off, chloroform was added thereto, and the insoluble
matter was filtered off. The solvent was distilled off, and
acetonitrile (0.2 mL) and 4 N hydrochloric acid-1,4-dioxane
solution (0.2 mL) were added thereto, followed by stirring for 10
minutes. DMSO (0.6 mL) was added thereto, and purification was
15 performed by reversed-phase HPLC (mobile phase:
water/acetonitrile) to give the title compound.

[0736]

Example 284: Synthesis of (S)-5'-((3-amino-3-methylpyrrolidin-1-yl)methyl)-2'',3-difluoro-4'''-methyl-[1,1':2',1'''-terphenyl]-4-
20 carbonitrile

The procedure of step 3 in Example 283 was conducted
using tert-butyl (S)-(3-methylpyrrolidin-3-yl)carbamate instead
of tert-butyl ((1S,2S,4R)-rel-7-azabicyclo[2.2.1]heptan-2-yl)carbamate-isomer-X to give the title compound.

25 [0737]

Example 285: Synthesis of 5'-(((1S,2S,4R)-rel-2-amino-7-azabicyclo[2.2.1]heptan-7-yl)methyl)-2'-(6,7-difluoro-1-(2-hydroxy-2-methylpropyl)-1H-benzo[d][1,2,3]triazol-5-yl)-3-fluoro-
[1,1'-biphenyl]-4-carbonitrile-isomer-X dihydrochloride

30 Step 1

3-Bromo-4-chlorobenzoic acid (19 g) was dissolved in
DMF (160 mL). At 25°C, DMAP (20 g) and WSC HCl (31 g) were added
thereto, followed by the addition of t-BuOH (38 mL). The
resulting mixture was stirred at room temperature overnight.
35 Ethyl acetate was added thereto, and the resulting mixture was

washed sequentially with water and saturated brine. After the organic layer was dried over anhydrous sodium sulfate, the solvent was distilled off. The residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate) to give tert-butyl 3-bromo-4-chloro-benzoate.

[0738]

Step 2

The tert-butyl 3-bromo-4-chloro-benzoate (1.00 g) obtained in step 1 above was dissolved in 1,4-dioxane (8.6 mL).

At room temperature, (4-cyano-3-fluoro-phenyl)boronic acid (509 mg), Pd(PPh₃)₄ (119 mg), and a 2 M aqueous sodium carbonate solution (4.3 mL) were added thereto, and the reaction mixture was stirred in a microwave reactor at 120°C for 30 minutes. The reaction mixture was filtrated, and the solvent was distilled off. The residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate) to give tert-butyl 4-chloro-3-(4-cyano-3-fluoro-phenyl)benzoate.

[0739]

Step 3

The tert-butyl 4-chloro-3-(4-cyano-3-fluoro-phenyl)benzoate (300 mg) obtained in step 2 above was dissolved in 1,4-dioxane (5 mL). At room temperature, Pd(OAc)₂ (40 mg), KOAc (300 mg), bispinacolatodiboron (500 mg), and Silica-SMAP (50 mg) were added thereto, followed by stirring at 100°C for 26 hours.

The reaction mixture was filtrated, and the solvent was distilled off. The residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate) to give tert-butyl 3-(4-cyano-3-fluoro-phenyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate.

[0740]

Step 4

1,2,3-Trifluoro-4-nitro-benzene (500 mg), 1-amino-2-methyl-propan-2-ol (302 mg), and triethylamine (0.590 mL) were dissolved in THF (5.65 mL), followed by stirring at 45°C for 5

hours. Ethyl acetate was added thereto, and the resulting mixture

was washed sequentially with water and saturated brine. After the organic layer was dried over anhydrous sodium sulfate, the solvent was distilled off. The residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate) to give 1-(2,3-difluoro-6-nitro-anilino)-2-methyl-propan-2-ol.

[0741]

Step 5

The 1-(2,3-difluoro-6-nitro-anilino)-2-methyl-propan-2-ol (74.1 g) obtained in step 4 above was dissolved in DMF (602 mL). At room temperature, NBS (64.3 g) was added thereto, followed by stirring at 90°C for 1 hour. Ethyl acetate was added thereto, and the mixture was washed 3 times with water.

Subsequently, washing was performed with saturated brine, drying was performed over anhydrous sodium sulfate, and the solvent was then distilled off. The residue was crystallized from IPE:hexane = 1:1, and the crystals were washed twice with hexane. The obtained crystals were dried to give 1-(4-bromo-2,3-difluoro-6-nitro-anilino)-2-methyl-propan-2-ol.

[0742]

Step 6

The 1-(4-bromo-2,3-difluoro-6-nitro-anilino)-2-methyl-propan-2-ol (82.2 g) obtained in step 5 above, ammonium chloride (82.2 g), and iron powder (41.1 g) were suspended in EtOH (421 mL) and water (421 mL), followed by stirring at 60°C overnight.

MTBE was added thereto, and the mixture was passed through Celite. MTBE was added thereto, and the mixture was washed sequentially with water and saturated brine, and dried over anhydrous sodium sulfate. The solvent was then distilled off to give 1-(6-amino-4-bromo-2,3-difluoro-anilino)-2-methyl-propan-2-ol.

[0743]

Step 7

The 1-(6-amino-4-bromo-2,3-difluoro-anilino)-2-methyl-propan-2-ol (68.1 g) obtained in step 6 above was dissolved in water (136 mL) and THF (341 mL). At 0°C, 12 N hydrochloric acid

(204 mL) and sodium nitrite (an aqueous solution (60 mL) in which 20.7 g of sodium nitrite was dissolved) were added thereto dropwise over 3 minutes, followed by stirring at room temperature for 1 hour. MTBE was added thereto, and the mixture was washed
5 sequentially with water and saturated brine, and dried over anhydrous sodium sulfate. The solvent was then distilled off. IPE:hexane = 1:1 (68 mL) was added to the residue, and the generated solid was collected by filtration and washed with IPE:hexane = 1:1. The obtained solid was dried to give 1-(5-
10 bromo-6,7-difluoro-benzotriazol-1-yl)-2-methyl-propan-2-ol.
[0744]

Step 8

The tert-butyl 3-(4-cyano-3-fluoro-phenyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (3.2 g) obtained in
15 step 3 above and the 1-(5-bromo-6,7-difluoro-benzotriazol-1-yl)-2-methyl-propan-2-ol (3.01 g) obtained in step 7 above were dissolved in 1,4-dioxane (25.2 mL). At room temperature, Pd(dba)₂ (348 mg), X-Phos (577 mg), and tripotassium phosphate (4.81 g) were added thereto, followed by stirring at 100°C overnight. The
20 solvent was distilled off, the residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate), and the solvent was distilled off. The residue was dissolved in THF (15.0 mL). At 0°C, 12 N hydrochloric acid (15.0 mL) was added thereto, followed by stirring at room temperature for 2 hours.
25 MTBE was added thereto, the mixture was washed with water, and dried over anhydrous sodium sulfate, and the solvent was distilled off to give 3-(4-cyano-3-fluoro-phenyl)-4-[6,7-difluoro-1-(2-hydroxy-2-methyl-propyl)benzotriazol-5-yl]benzoic acid.

30 [0745]

Step 9

The 3-(4-cyano-3-fluoro-phenyl)-4-[6,7-difluoro-1-(2-hydroxy-2-methyl-propyl)benzotriazol-5-yl]benzoic acid (120 mg) obtained in step 8 above was dissolved in THF (1.29 mL). At room
35 temperature, WSC HCl (98.6 mg) and HOBt (78.8 mg) were added

thereto, followed by stirring at room temperature for 20 minutes. Sodium borohydride (19.5 mg) was added thereto, followed by stirring for 1 hour. The solvent was distilled off under reduced pressure. The residue was purified by silica gel column

5 chromatography (mobile phase: chloroform/methanol) to give 4-[2-[6,7-difluoro-1-(2-hydroxy-2-methyl-propyl)benzotriazol-5-yl]-5-(hydroxymethyl)phenyl]-2-fluoro-benzonitrile.

[0746]

Step 10

10 The 4-[2-[6,7-difluoro-1-(2-hydroxy-2-methyl-propyl)benzotriazol-5-yl]-5-(hydroxymethyl)phenyl]-2-fluoro-benzonitrile (100 mg) obtained in step 9 above was dissolved in DCM (2.21 mL). At room temperature, Dess-Martin periodinane (103 mg) was added thereto, followed by stirring at room temperature
15 overnight. The reaction mixture was vacuum-concentrated, and the residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate) to give 4-[2-[6,7-difluoro-1-(2-hydroxy-2-methyl-propyl)benzotriazol-5-yl]-5-formyl-phenyl]-2-fluoro-benzonitrile.

20 [0747]

Step 11

The 4-[2-[6,7-difluoro-1-(2-hydroxy-2-methyl-propyl)benzotriazol-5-yl]-5-formyl-phenyl]-2-fluoro-benzonitrile (50 mg) obtained in step 10 above was dissolved in DCM (1.11 mL).
25 At room temperature, the tert-butyl ((1S,2S,4R)-rel-7-azabicyclo[2.2.1]heptan-2-yl)carbamate-isomer-X (47.1 mg) synthesized in step 2 of Example 202 was added thereto, followed by stirring at room temperature for 30 minutes. At room temperature, sodium triacetoxyborohydride (94.1 mg) was added
30 thereto, followed by stirring at room temperature overnight. After the addition of chloroform, the insoluble matter was filtered off, and the solvent was distilled off. The residue was purified by silica gel column chromatography (NH-silica gel, mobile phase: hexane/ethyl acetate), and the solvent was
35 distilled off. The residue was dissolved in MeOH (1.0 mL), and a

4 N hydrochloric acid-1,4-dioxane solution (1.0 mL) was added thereto, followed by stirring for 30 minutes. The solvent was removed from the reaction mixture to give the title compound. [0748]

5 Example 286: Synthesis of 5'-(((1S,2S,4R)-rel-2-amino-7-azabicyclo[2.2.1]heptan-7-yl)methyl)-2'-(7-chloro-6-fluoro-1-((1-hydroxycyclobutyl)methyl)-1H-benzo[d][1,2,3]triazol-5-yl)-3-fluoro-[1,1'-biphenyl]-4-carbonitrile-isomer-X dihydrochloride

The procedure of steps 4 to 11 in Example 285 was
10 conducted using 2-chloro-1,3-difluoro-4-nitro-benzene instead of 1,2,3-trifluoro-4-nitro-benzene, and using 1-(aminomethyl)cyclobutanol instead of 1-amino-2-methyl-propan-2-ol to give the title compound. [0749]

15 Example 287: Synthesis of 5'-(((1S,2S,4R)-rel-2-amino-7-azabicyclo[2.2.1]heptan-7-yl)methyl)-3-fluoro-2'-(6-fluoro-1-(2-hydroxy-2-methylpropyl)-1H-indazol-5-yl)-[1,1'-biphenyl]-4-carbonitrile-isomer-X dihydrochloride

Step 1

20 5-Bromo-6-fluoro-1H-indazole (94 mg) was dissolved in DMF (1.5 ml). At room temperature, cesium carbonate (285 mg) and 2,2-dimethyloxirane (0.078 mL) were added thereto, followed by stirring at 90°C for 16 hours. The reaction was quenched with a saturated aqueous ammonium chloride solution, ethyl acetate was
25 added thereto, and the resulting mixture was washed sequentially with water and saturated brine. After the organic layer was dried over anhydrous sodium sulfate, the solvent was distilled off. The residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate) to give 1-(5-bromo-6-fluoro-indazol-
30 1-yl)-2-methyl-propan-2-ol. [0750]

Step 2

The procedure of steps 8 to 11 in Example 285 was
conducted using the 1-(5-bromo-6-fluoro-indazol-1-yl)-2-methyl-
35 propan-2-ol obtained in step 1 above instead of 1-(5-bromo-6,7-

difluoro-benzotriazol-1-yl)-2-methyl-propan-2-ol to give the title compound.

[0751]

Example 288: Synthesis of 5'-(((1S,2S,4R)-rel-2-amino-7-azabicyclo[2.2.1]heptan-7-yl)methyl)-2'-(7-chloro-1-(2-ethyl-2-hydroxybutyl)-6-fluoro-1H-benzo[d][1,2,3]triazol-5-yl)-3-fluoro-[1,1'-biphenyl]-4-carbonitrile-isomer-X dihydrochloride

The procedure of steps 4 to 11 in Example 285 was conducted using 2-chloro-1,3-difluoro-4-nitro-benzene instead of 1,2,3-trifluoro-4-nitro-benzene, and using 3-(aminomethyl)pentan-3-ol instead of 1-amino-2-methyl-propan-2-ol to give the title compound.

[0752]

Example 289: Synthesis of 5'-(((1S,2S,4R)-rel-2-amino-7-azabicyclo[2.2.1]heptan-7-yl)methyl)-3-fluoro-2'-(6-fluoro-1-(2-hydroxy-2-methylpropyl)-1H-indol-5-yl)-[1,1'-biphenyl]-4-carbonitrile-isomer-X dihydrochloride

Step 1

5-Bromo-6-fluoro-1H-indole (50 mg) was dissolved in DMF (0.78 mL). At room temperature, cesium carbonate (151 mg) and 2,2-dimethyloxirane (42 µL) were added thereto, followed by stirring at 90°C for 16 hours. The reaction was quenched with a saturated aqueous ammonium chloride solution, ethyl acetate was added thereto, and the resulting mixture was washed sequentially with water and saturated brine. After the organic layer was dried over anhydrous sodium sulfate, the solvent was distilled off. The residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate) to give 1-(5-bromo-6-fluoro-indol-1-yl)-2-methyl-propan-2-ol.

[0753]

Step 2

The procedure of steps 8 to 11 in Example 285 was conducted using the 1-(5-bromo-6-fluoro-indol-1-yl)-2-methyl-propan-2-ol obtained in step 1 above instead of 1-(5-bromo-6,7-difluoro-benzotriazol-1-yl)-2-methyl-propan-2-ol to give the

title compound.

[0754]

Example 290: Synthesis of 5'-(((1S,2S,4R)-rel-2-amino-7-azabicyclo[2.2.1]heptan-7-yl)methyl)-2'',3-difluoro-4'''-(2-hydroxy-2-methylpropyl)-[1,1':2',1'''-terphenyl]-4-carbonitrile-isomer-X dihydrochloride

Step 1

Methyl 2-(4-bromo-3-fluoro-phenyl)acetate (500 mg) was dissolved in THF (2.2 mL). At -30°C, a solution of 3 M MeMgBr in diethyl ether (5.40 mL) was added thereto dropwise, followed by stirring at room temperature overnight. The reaction mixture was introduced into an aqueous ammonium chloride solution, ethyl acetate was added thereto, and the mixture was washed sequentially with water and saturated brine. After the organic layer was dried over anhydrous sodium sulfate, the solvent was distilled off. The residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate) to give 1-(4-bromo-3-fluoro-phenyl)-2-methyl-propan-2-ol.

[0755]

Step 2

The procedure of steps 8 to 11 in Example 285 was conducted using the 1-(4-bromo-3-fluoro-phenyl)-2-methyl-propan-2-ol obtained in step 1 above instead of 1-(5-bromo-6,7-difluoro-benzotriazol-1-yl)-2-methyl-propan-2-ol to give the title compound.

[0756]

Example 291: Synthesis of (S)-5'-((3-aminopyrrolidin-1-yl)methyl)-2'',3-difluoro-4'''-(2-hydroxy-2-methylpropyl)-[1,1':2',1'''-terphenyl]-4-carbonitrile dihydrochloride

The procedure of steps 8 to 11 in Example 285 was conducted using the 1-(4-bromo-3-fluoro-phenyl)-2-methyl-propan-2-ol obtained in step 1 of Example 290 instead of 1-(5-bromo-6,7-difluoro-benzotriazol-1-yl)-2-methyl-propan-2-ol, and using tert-butyl N-[(3S)-pyrrolidin-3-yl]carbamate instead of tert-butyl ((1S,2S,4R)-rel-7-azabicyclo[2.2.1]heptan-2-yl)carbamate-isomer-X

to give the title compound.

[0757]

Example 292: Synthesis of 5'-(1-((1S,2S,4R)-rel-2-amino-7-azabicyclo[2.2.1]heptan-7-yl)ethyl)-2'',3-difluoro-4''-methyl-[1,1':2',1''-terphenyl]-4-carbonitrile-isomer-X dihydrochloride

Step 1

1-(3-Bromo-4-chloro-phenyl)ethanone (2.00 g) was dissolved in 1,4-dioxane (14.3 mL). At room temperature, (4-cyano-3-fluoro-phenyl)boronic acid (1.55 g), PdCl₂(dppf)CH₂Cl₂ (188 mg), and a 2 M aqueous sodium carbonate solution (10.7 mL) were added thereto, followed by stirring at 90°C overnight. Ethyl acetate was added thereto, and the resulting mixture was washed sequentially with water and saturated brine. After the organic layer was dried over anhydrous sodium sulfate, the solvent was distilled off. Diethyl ether was added thereto, and the precipitate was collected by filtration to give 4-(5-acetyl-2-chloro-phenyl)-2-fluoro-benzonitrile.

[0758]

Step 2

The 4-(5-acetyl-2-chloro-phenyl)-2-fluoro-benzonitrile (200 mg) obtained in step 1 above and the tert-butyl ((1S,2S,4R)-rel-7-azabicyclo[2.2.1]heptan-2-yl)carbamate-isomer-X (160 mg) synthesized in step 2 of Example 202 were suspended in titanium isopropoxide (1.0 mL), followed by stirring for 3 days. EtOH (5 mL) and sodium borohydride (138 mg) were added thereto, followed by stirring for 1 hour. The mixture was diluted with THF, Celite was added thereto, filtration was performed through a Celite bed, and the solvent was distilled off. The residue was purified by silica gel column chromatography (NH silica, mobile phase: hexane/ethyl acetate) to give tert-butyl N-[(1S,3R,4R)-7-[1-[4-chloro-3-(4-cyano-3-fluoro-phenyl)phenyl]ethyl]-7-azabicyclo[2.2.1]heptan-3-yl]carbamate.

[0759]

Step 3

The tert-butyl N-[(1S,3R,4R)-7-[1-[4-chloro-3-(4-cyano-

3-fluoro-phenyl]phenyl]ethyl]-7-azabicyclo[2.2.1]heptan-3-yl]carbamate (20 mg) obtained in step 2 above was dissolved in DMF (0.50 mL). At room temperature, (2-fluoro-4-methyl-phenyl)boronic acid (19.7 mg), S-Phos (1.75 mg), Pd(dba)₂ (1.22 mg), and tripotassium phosphate (27.1 mg) were added thereto, and the mixture was stirred in a microwave reactor at 160°C for 45 minutes. MTBE was added thereto, and the mixture was washed 3 times with water. Subsequently, washing was performed with saturated brine, drying was performed over anhydrous sodium sulfate, and the solvent was then distilled off. The residue was purified by reversed-phase HPLC (mobile phase: water/acetonitrile), and the target fractions were combined, neutralized with an aqueous sodium hydrogen carbonate solution, and extracted twice with chloroform. The solvent was distilled off, the residue was dissolved in MeOH (1.0 mL), and a 4 N hydrochloric acid-1,4-dioxane solution (1.0 mL) was added thereto, followed by stirring for 30 minutes. The solvent was removed from the reaction mixture to give the title compound. [0760]

Example 293: Synthesis of 5'-(((1S,2S,4R)-rel-2-amino-7-azabicyclo[2.2.1]heptan-7-yl)methyl)-2'-(1-(2-ethyl-2-hydroxybutyl)-6,7-difluoro-1H-benzo[d][1,2,3]triazol-5-yl)-3-fluoro-[1,1'-biphenyl]-4-carbonitrile-isomer-X dihydrochloride

The procedure of steps 4 to 11 in Example 285 was conducted using 3-(aminomethyl)pentan-3-ol instead of 1-amino-2-methyl-propan-2-ol to give the title compound. [0761]

Example 294: Synthesis of 5'-(((1S,2S,4R)-rel-2-amino-7-azabicyclo[2.2.1]heptan-7-yl)methyl)-2'-(7-bromo-6-fluoro-1-(2-hydroxy-2-methylpropyl)-1H-benzo[d][1,2,3]triazol-5-yl)-3-fluoro-[1,1'-biphenyl]-4-carbonitrile-isomer-X dihydrochloride

Step 1

2-Bromo-1,3-difluoro-4-nitro-benzene (3.00 g) was dissolved in THF (31.5 mL). TEA (2.63 mL) and 1-amino-2-methyl-propan-2-ol (1.40 mL) were added thereto, followed by stirring at

room temperature for 1 hour. Ethyl acetate was added thereto, and the resulting mixture was washed sequentially with water and saturated brine. After the organic layer was dried over anhydrous sodium sulfate, the solvent was distilled off. The residue was
5 purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate) to give 1-(2-bromo-3-fluoro-6-nitro-anilino)-2-methyl-propan-2-ol.

[0762]

Step 2

10 The 1-(2-bromo-3-fluoro-6-nitro-anilino)-2-methyl-propan-2-ol (1.03 g) obtained in step 1 above was dissolved in acetic acid (6.7 mL). At room temperature, N-iodosuccinimide (981 mg) was added thereto, followed by stirring at 50°C for 3 hours. MTBE and water were added thereto, and the mixture was extracted
15 twice with MTBE. After the combined organic layers were dried over anhydrous sodium sulfate, the solvent was distilled off. The residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate) to give 1-(2-bromo-3-fluoro-4-iodo-6-nitro-anilino)-2-methyl-propan-2-ol.

20 [0763]

Step 3

 The 1-(2-bromo-3-fluoro-4-iodo-6-nitro-anilino)-2-methyl-propan-2-ol (1.33 g) obtained in step 2 above and iron powder (1.33 g) were suspended in THF (10.2 mL) and a 2 N
25 hydrochloric acid (10.2 mL), followed by stirring at 60°C for 1 hour. MTBE was added thereto, and the mixture was passed through Celite. MTBE was added thereto, and the mixture was washed sequentially with water and saturated brine, and dried over anhydrous sodium sulfate. The solvent was then distilled off. The
30 residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate) to give 1-(6-amino-2-bromo-3-fluoro-4-iodo-anilino)-2-methyl-propan-2-ol.

[0764]

Step 4

35 The 1-(6-amino-2-bromo-3-fluoro-4-iodo-anilino)-2-

methyl-propan-2-ol (940 mg) obtained in step 3 above was dissolved in water (1.88 mL) and THF (4.7 mL). At 0°C, 12 N hydrochloric acid (2.82 mL) and an aqueous sodium nitrite solution (an aqueous solution obtained by dissolving 209 mg of sodium nitrite in 0.63 mL of water) were added thereto, followed by stirring at room temperature for 1 hour. MTBE was added thereto, and the mixture was washed sequentially with water and saturated brine, and dried over anhydrous sodium sulfate. The solvent was then distilled off. The residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate) to give 1-(7-bromo-6-fluoro-5-iodo-benzotriazol-1-yl)-2-methyl-propan-2-ol.

[0765]

Step 5

The tert-butyl 3-(4-cyano-3-fluoro-phenyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoate (3 g) obtained in step 3 of Example 285 and the 1-(7-bromo-6-fluoro-5-iodo-benzotriazol-1-yl)-2-methyl-propan-2-ol (3.228 g) obtained in step 5 above were dissolved in 1,4-dioxane (23.6 mL). At room temperature, PdCl₂(PPh₃)₂ (398 mg) and tripotassium phosphate (4.513 g) were added thereto. After nitrogen substitution and degassing, the mixture was stirred at 100°C overnight. The reaction mixture was passed through Celite, and the solvent was distilled off. The residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate), and the solvent was distilled off. THF (45.0 mL) and 12 N hydrochloric acid (45.0 mL) were added to the residue, followed by stirring at room temperature for 2 hours. MTBE and water were added thereto for partition. After the organic layer was washed with saturated brine and dried over anhydrous sodium sulfate, the solvent was distilled off. The residue was crystallized from diethyl ether/hexane to give 4-[7-bromo-6-fluoro-1-(2-hydroxy-2-methyl-propyl)benzotriazol-5-yl]-3-(4-cyano-3-fluoro-phenyl)benzoic acid.

[0766]

Step 6

The 4-[7-bromo-6-fluoro-1-(2-hydroxy-2-methyl-propyl)benzotriazol-5-yl]-3-(4-cyano-3-fluoro-phenyl)benzoic acid (88.0 mg) obtained in step 5 above was dissolved in THF (0.834 mL). At room temperature, WSC HCl (64.0 mg) and HOBt (51.1 mg) were added thereto, followed by stirring at room temperature for 20 minutes. Sodium borohydride (18.9 mg) was added thereto, followed by stirring for 1 hour. The solvent was distilled off under reduced pressure. The residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate) to give 4-[2-[7-bromo-6-fluoro-1-(2-hydroxy-2-methyl-propyl)benzotriazol-5-yl]-5-(hydroxymethyl)phenyl]-2-fluorobenzonitrile.

[0767]

15 Step 7

The 4-[2-[7-bromo-6-fluoro-1-(2-hydroxy-2-methyl-propyl)benzotriazol-5-yl]-5-(hydroxymethyl)phenyl]-2-fluorobenzonitrile (43.5 mg) obtained in step 6 above was dissolved in DCM (0.847 mL). At room temperature, Dess-Martin periodinane (39.5 mg) was added thereto, followed by stirring at room temperature overnight. The reaction mixture was vacuum-concentrated, and the residue was purified by silica gel column chromatography (mobile phase: hexane/ethyl acetate) to give 4-[2-[7-bromo-6-fluoro-1-(2-hydroxy-2-methyl-propyl)benzotriazol-5-yl]-5-formyl-phenyl]-2-fluorobenzonitrile.

[0768]

Step 8

The 4-[2-[7-bromo-6-fluoro-1-(2-hydroxy-2-methyl-propyl)benzotriazol-5-yl]-5-formyl-phenyl]-2-fluorobenzonitrile (37.2 mg) obtained in step 7 above was dissolved in DCM (1.00 mL). At room temperature, the tert-butyl ((1S,2S,4R)-rel-7-azabicyclo[2.2.1]heptan-2-yl)carbamate-isomer-X (30.9 mg) synthesized in step 2 of Example 202 was added thereto, followed by stirring at room temperature for 30 minutes. At room temperature, sodium triacetoxyborohydride (27.8 mg) was added

thereto, followed by stirring at room temperature overnight.
After the addition of chloroform, the insoluble matter was
filtered off, and the solvent was distilled off. The residue was
purified by silica gel column chromatography (NH-silica gel,
5 mobile phase: hexane/ethyl acetate), and the solvent was
distilled off. The residue was dissolved in MeOH (1.0 mL), and a
4 N hydrochloric acid-1,4-dioxane solution (1.0 mL) was added
thereto, followed by stirring for 30 minutes. The solvent was
removed from the reaction mixture to give the title compound.

10 [0769]


Example 295: Synthesis of 5'-(((1S,2S,4R)-rel-2-amino-7-
azabicyclo[2.2.1]heptan-7-yl)methyl)-2'-(7-bromo-1-(2-ethyl-2-
hydroxybutyl)-6-fluoro-1H-benzo[d][1,2,3]triazol-5-yl)-3-fluoro-
[1,1'-biphenyl]-4-carbonitrile-isomer-X dihydrochloride

15 The procedure of steps 1 to 8 in Example 294 was
conducted using 3-(aminomethyl)pentan-3-ol instead of 1-amino-2-
methyl-propan-2-ol to give the title compound.

[0770]

20 The following are lists of the compounds of Examples 1
to 295.


[0771]



[0772]

In the following tables, if the structure:


25 [0773]



[0774]


is shown as:

[0775]



Racemate
Relative
configuration

or




Racemate
Relative
configuration


[0776]

then the compound represents a mixture of compounds having the structure:

5 [0777]




or



[0778]

if the structure is shown as:

[0779]




Racemate
Relative
configuration

10


[0780]

then the compound represents a mixture of compounds having the structure:

[0781]



or




15

[0782]

and if the structure is shown as:


[0783]



[0784]

then the compound has one of the following structures:

[0785]



5

[0786]

[0787]

The term "salt form" indicates a salt form. The term "free" indicates a free form.

10 [0788]

Table 1

Ex. No.					salt form	MS m/z (M+1)	NMR
1					free	382.4	¹ H-NMR (DMSO-D ₆) δ: 7.74 (2H, d, J = 8.1 Hz), 7.66-7.61 (1H, m), 7.53 (1H, d, J = 8.4 Hz), 7.49 (1H, d, J = 9.1 Hz), 7.33 (2H, d, J = 8.1 Hz), 7.09 (2H, d, J = 8.1 Hz), 7.01 (2H, d, J = 7.7 Hz), 3.70-3.40 (4H, m), 3.24-3.13 (1H, m), 2.27 (3H, s), 2.03-1.90 (1H, m), 1.70-1.59 (1H, m).
2					free	398.2	¹ H-NMR (DMSO-D ₆) δ: 7.74 (2H, d, J = 8.2 Hz), 7.52-7.49 (1H, m), 7.45-7.41 (2H, m), 7.32 (2H, dd, J = 8.4, 2.3 Hz), 7.09 (2H, d, J = 8.2 Hz), 7.01 (2H, d, J = 8.2 Hz), 3.99-3.51 (4H, m), 2.27 (3H, s), 2.15-2.01 (1H, m), 1.83-1.70 (1H, m), 1.31-1.19 (1H, m).
3					free	396.4	¹ H-NMR (DMSO-D ₆) δ: 7.74 (2H, d, J = 8.1 Hz), 7.49 (2H, s), 7.38 (1H, s), 7.33 (2H, d, J = 8.1 Hz), 7.09 (2H, d, J = 8.1 Hz), 7.01 (2H, d, J = 8.1 Hz), 4.38-4.23 (1H, m), 3.73-3.61 (1H, m), 3.18-3.05 (1H, m), 3.01-2.89 (1H, m), 2.87-2.79 (1H, m), 2.27 (3H, s), 1.87-1.64 (4H, m).
4					free	422.5	¹ H-NMR (DMSO-D ₆) δ: 7.76 (3H, d, J = 8.2 Hz), 7.63 (1H, s), 7.53 (1H, d, J = 7.6 Hz), 7.32 (2H, d, J = 8.2 Hz), 7.10 (2H, d, J = 7.6 Hz), 7.00 (2H, d, J = 7.6 Hz), 4.24 (1H, d, J = 8.9 Hz), 4.12 (1H, d, J = 8.9 Hz), 3.95 (1H, d, J = 9.5 Hz), 3.78 (1H, d, J = 10.2 Hz), 3.51 (2H, s), 2.96 (2H, s), 2.27 (3H, s), 1.86-1.60 (2H, m), 1.71-1.60 (2H, m).
5					free	408.5	¹ H-NMR (DMSO-D ₆) δ: 7.76 (2H, d, J = 8.2 Hz), 7.67-7.61 (1H, m), 7.58-7.47 (2H, m), 7.35-7.29 (2H, m), 7.14-7.07 (2H, m), 7.00 (2H, d, J = 8.2 Hz), 4.12-4.03 (1H, m), 3.96-3.84 (2H, m), 3.76-3.71 (2H, m), 3.59-3.47 (3H, m), 2.28 (3H, s), 2.19-2.11 (2H, m).
6					free	422.2	¹ H-NMR (DMSO-D ₆) δ: 7.76 (2H, t, J = 4.3 Hz), 7.65 (1H, dd, J = 7.9, 1.8 Hz), 7.56 (1H, dd, J = 18.9, 1.5 Hz), 7.50 (1H, d, J = 7.9 Hz), 7.34 (2H, d, J = 8.2 Hz), 7.11 (2H, d, J = 7.6 Hz), 7.01 (2H, dd, J = 8.2, 2.4 Hz), 3.61-3.47 (4H, m), 2.95 (4H, t, J = 1.7 Hz), 2.28 (3H, s), 1.92-1.72 (4H, m).
7					free	422.3	¹ H-NMR (DMSO-D ₆) δ: 7.73 (2H, d, J = 8.4 Hz), 7.56 (1H, d, J = 7.7 Hz), 7.50 (1H, d, J = 7.7 Hz), 7.44 (1H, s), 7.32 (2H, d, J = 8.4 Hz), 7.08 (2H, d, J = 8.1 Hz), 6.99 (2H, d, J = 7.7 Hz), 4.65-4.57 (1H, m), 4.19-4.11 (1H, m), 3.47-3.35 (2H, m), 2.26 (2H, s), 2.04-1.44 (8H, m).
8					free	396.1	¹ H-NMR (DMSO-D ₆) δ: 7.76 (2H, d, J = 8.2 Hz), 7.69-7.61 (1H, m), 7.51 (2H, d, J = 7.9 Hz), 7.33-7.32 (2H, m), 7.10 (2H, d, J = 7.9 Hz), 7.01 (2H, d, J = 7.0 Hz), 3.81-3.40 (4H, m), 2.27 (3H, s), 2.03-1.90 (2H, m), 1.44-1.22 (3H, m).



















[0789]

Table 2

9					free	416.0 418.0	¹ H-NMR (DMSO-D ₆) δ: 7.74 (2H, d, J = 8.2 Hz), 7.67-7.59 (2H, m), 7.44 (1H, d, J = 7.9 Hz), 7.34-7.23 (3H, m), 7.16-7.10 (2H, m), 3.95-3.45 (5H, m), 2.29 (3H, s), 2.28-2.19 (1H, m), 2.06-1.96 (1H, m).
10					free	416.2 418.1	¹ H-NMR (DMSO-D ₆) δ: 7.72 (2H, d, J = 8.2 Hz), 7.62-7.57 (1H, m), 7.50 (2H, d, J = 8.2 Hz), 7.28 (2H, d, J = 7.6 Hz), 7.17 (2H, d, J = 9.5 Hz), 6.85 (1H, d, J = 7.0 Hz), 3.82-3.52 (5H, m), 2.22 (3H, s), 2.19-2.12 (1H, m), 1.95-1.87 (1H, m).
11					free	454.2	¹ H-NMR (DMSO-D ₆) δ: 7.74 (2H, d, J = 8.2 Hz), 7.66-7.52 (4H, m), 7.35-7.26 (3H, m), 7.01 (1H, d, J = 8.2 Hz), 3.86-3.47 (5H, m), 2.23-2.10 (1H, m), 1.98-1.87 (1H, m).
12					free	427.2	¹ H-NMR (DMSO-D ₆) δ: 7.80 (2H, d, J = 8.2 Hz), 7.76 (1H, s), 7.73-7.67 (1H, m), 7.66-7.58 (2H, m), 7.43-7.35 (3H, m), 7.31 (1H, d, J = 7.6 Hz), 4.13-4.07 (1H, m), 3.92-3.67 (2H, m), 3.64-3.49 (2H, m), 3.16 (3H, s), 2.28-2.15 (1H, m), 2.02-1.91 (1H, m).
13					free	418.2	¹ H-NMR (DMSO-D ₆) δ: 7.70 (2H, d, J = 8.2 Hz), 7.65-7.59 (1H, m), 7.52 (2H, d, J = 8.2 Hz), 7.43 (2H, d, J = 8.2 Hz), 7.27 (2H, d, J = 7.0 Hz), 7.20 (2H, d, J = 8.2 Hz), 3.87-3.48 (5H, m), 3.14-3.06 (1H, m), 2.22-2.10 (1H, m), 1.99-1.86 (1H, m).
14					free	436.2	¹ H-NMR (DMSO-D ₆) δ: 7.71 (2H, d, J = 8.2 Hz), 7.65-7.58 (3H, m), 7.54 (2H, d, J = 7.6 Hz), 7.33-7.24 (4H, m), 3.83-3.53 (5H, m), 2.23-2.11 (1H, m), 1.98-1.89 (1H, m).
15					free	400.2	¹ H-NMR (DMSO-D ₆) δ: 7.68 (2H, d, J = 8.2 Hz), 7.63-7.56 (1H, m), 7.56-7.49 (1H, m), 7.44 (1H, d, J = 8.2 Hz), 7.24 (2H, d, J = 7.6 Hz), 7.10 (1H, t, J = 7.9 Hz), 6.95 (1H, d, J = 7.6 Hz), 6.85 (1H, d, J = 10.8 Hz), 3.83-3.53 (5H, m), 2.22 (3H, s), 2.19-2.10 (1H, m), 1.98-1.88 (1H, m).
16					free	400.1	¹ H-NMR (DMSO-D ₆) δ: 7.75 (1H, t, J = 7.6 Hz), 7.60 (1H, d, J = 7.6 Hz), 7.47 (1H, d, J = 7.6 Hz), 7.33-7.24 (1H, m), 7.07-6.95 (6H, m), 4.02-3.52 (5H, m), 2.22 (3H, s), 2.18-2.11 (1H, m), 1.98-1.87 (1H, m).
17					free	440.2	¹ H-NMR (DMSO-D ₆) δ: 7.80 (1H, t, J = 7.5 Hz), 7.61-7.58 (1H, m), 7.52-7.49 (2H, m), 7.38 (1H, dd, J = 10.5, 1.4 Hz), 7.12 (2H, d, J = 7.9 Hz), 7.11-7.09 (1H, m), 7.04 (2H, d, J = 7.9 Hz), 4.63-4.57 (1H, m), 4.14-4.06 (1H, m), 2.28 (3H, s), 2.24-2.02 (2H, m), 2.00-1.89 (1H, m), 1.84-1.51 (3H, m), 1.34-1.14 (3H, m).
18					free	479.2	¹ H-NMR (DMSO-D ₆) δ: 7.76 (1H, t, J = 7.5 Hz), 7.62-7.56 (2H, m), 7.51-7.50 (1H, m), 7.44-7.42 (1H, m), 7.41-7.37 (1H, m), 7.35-7.31 (2H, m), 7.11 (1H, d, J = 8.2 Hz), 6.83 (1H, d, J = 8.2 Hz), 6.40 (1H, d, J = 3.1 Hz), 4.65-4.58 (1H, m), 4.22-4.10 (1H, m), 3.77 (3H, s), 3.74-3.72 (1H, m), 2.04-1.19 (8H, m).





















[0790]

Table 3

19					free	418.1	¹ H-NMR (DMSO-D6) δ: 7.71-7.68 (1H, m), 7.62 (1H, s), 7.55 (1H, d, J = 7.9 Hz), 7.24-7.13 (4H, m), 7.06 (2H, d, J = 7.9 Hz), 3.90-3.53 (4H, m), 3.49-3.41 (1H, m), 2.30 (3H, s), 2.25-2.16 (1H, m), 1.99-1.93 (1H, m).
20					free	462.1	¹ H-NMR (DMSO-D6) δ: 7.80 (1H, t, J = 7.5 Hz), 7.70-7.66 (1H, m), 7.62 (1H, dd, J = 6.6, 1.5 Hz), 7.51 (1H, d, J = 7.7 Hz), 7.33 (1H, d, J = 10.6 Hz), 7.24 (1H, t, J = 7.9 Hz), 7.13-7.08 (2H, m), 7.00 (1H, d, J = 10.3 Hz), 3.68-3.56 (2H, m), 3.55-3.40 (4H, m), 3.23 (3H, s), 3.22-3.11 (1H, m), 2.81 (2H, t, J = 6.6 Hz), 2.03-1.88 (1H, m), 1.69-1.57 (1H, m).
21					free	458.2	¹ H-NMR (DMSO-D6) δ: 7.81 (1H, t, J = 7.6 Hz), 7.61 (1H, dd, J = 7.8, 1.7 Hz), 7.54 (1H, d, J = 1.5 Hz), 7.50 (1H, d, J = 7.6 Hz), 7.37 (1H, d, J = 9.2 Hz), 7.21 (1H, t, J = 7.9 Hz), 7.11 (1H, dd, J = 7.9, 1.5 Hz), 7.05 (1H, d, J = 8.2 Hz), 6.95 (1H, d, J = 11.3 Hz), 4.63-4.57 (1H, m), 4.12-4.05 (1H, m), 2.31 (3H, s), 2.25-2.02 (3H, m), 2.00-1.69 (3H, m), 1.65-1.50 (2H, m), 1.47-1.34 (1H, m).
22					free	502.2	¹ H-NMR (DMSO-D6) δ: 7.81 (1H, t, J = 7.5 Hz), 7.62 (1H, dd, J = 7.8, 1.7 Hz), 7.56-7.52 (2H, m), 7.35 (1H, dd, J = 10.5, 1.4 Hz), 7.24 (1H, t, J = 7.9 Hz), 7.13-7.09 (2H, m), 7.02-6.99 (1H, m), 4.62 (1H, s), 4.10 (1H, s), 3.53 (2H, t, J = 6.7 Hz), 3.40-3.34 (1H, m), 3.23 (3H, s), 2.81 (2H, t, J = 6.7 Hz), 2.33-1.90 (6H, m), 1.70-1.55 (2H, m).
23					free	497.2	¹ H-NMR (DMSO-D6) δ: 7.76 (1H, t, J = 7.6 Hz), 7.62 (1H, dd, J = 7.8, 1.7 Hz), 7.57-7.50 (3H, m), 7.39-7.34 (2H, m), 7.22 (1H, d, J = 11.3 Hz), 7.13-7.09 (1H, m), 6.46-6.43 (1H, m), 4.63 (1H, s), 4.15 (1H, s), 3.74 (3H, s), 3.19-3.13 (1H, m), 2.05-1.78 (4H, m), 1.77-1.67 (2H, m), 1.65-1.56 (1H, m), 1.48-1.39 (1H, m).
24					free	498.2	¹ H-NMR (DMSO-D6) δ: 8.09 (1H, d, J = 0.9 Hz), 7.81-7.75 (2H, m), 7.65 (1H, dd, J = 7.8, 1.7 Hz), 7.60-7.55 (2H, m), 7.46 (1H, d, J = 10.7 Hz), 7.41 (1H, dd, J = 10.5, 1.4 Hz), 7.11 (1H, dd, J = 7.9, 1.5 Hz), 4.64 (1H, s), 4.14 (1H, s), 3.99 (3H, s), 3.36-3.30 (1H, m), 2.06-1.44 (8H, m).
25					free	476.1	¹ H-NMR (DMSO-D6) δ: 7.85-7.80 (1H, m), 7.74-7.59 (2H, m), 7.55 (1H, d, J = 7.9 Hz), 7.35-7.29 (1H, m), 7.26-7.21 (1H, m), 7.11 (2H, d, J = 7.9 Hz), 7.01 (1H, d, J = 11.0 Hz), 3.79-3.44 (6H, m), 3.23 (3H, s), 2.85-2.79 (2H, m), 2.10-1.98 (2H, m), 1.47-1.27 (3H, m).






















[0791]

Table 4

26					free	502.1	¹ H-NMR (DMSO-D ₆) δ: 7.82 (1H, t, J = 7.5 Hz), 7.53 (1H, d, J = 7.9 Hz), 7.57-7.52 (2H, m), 7.34 (1H, d, J = 9.8 Hz), 7.24 (1H, t, J = 7.9 Hz), 7.15 (2H, m), 7.01 (1H, d, J = 11.3 Hz), 4.63 (1H, s), 4.13 (1H, s), 3.53 (2H, t, J = 6.6 Hz), 3.42-3.36 (1H, m), 3.23 (3H, s), 2.81 (2H, t, J = 6.6 Hz), 2.09-1.45 (8H, m).
27					free	468.2	¹ H-NMR (DMSO-D ₆) δ: 7.83-7.78 (1H, m), 7.71-7.48 (3H, m), 7.39-7.31 (1H, m), 7.27-7.21 (1H, m), 7.15-7.08 (2H, m), 7.03-6.98 (1H, m), 4.77-4.47 (1H, m), 4.29-3.95 (1H, m), 3.66-3.38 (4H, m), 3.23 (3H, d, J = 0.9 Hz), 2.99-2.88 (1H, m), 2.81 (2H, t, J = 6.5 Hz), 2.72-2.55 (1H, m), 2.01-1.43 (4H, m).
28					free	516.3	¹ H-NMR (DMSO-D ₆) δ: 7.80 (1H, t, J = 7.5 Hz), 7.64-7.61 (1H, m), 7.57-7.52 (2H, m), 7.29-7.26 (1H, m), 7.21 (1H, t, J = 7.9 Hz), 7.15 (1H, dd, J = 8.2, 1.2 Hz), 7.07 (1H, dd, J = 7.9, 0.9 Hz), 6.95 (1H, d, J = 11.3 Hz), 4.61 (1H, s), 4.40 (1H, s), 4.09 (1H, s), 3.19-3.11 (1H, m), 2.65 (2H, s), 2.11-1.59 (6H, m), 1.58-1.48 (1H, m), 1.41-1.32 (1H, m), 1.04 (6H, s).
29					free	398.6	¹ H-NMR (DMSO-D ₆) δ: 7.74 (2H, d, J = 8.5 Hz), 7.67-7.63 (1H, m), 7.54 (1H, dd, J = 10.5, 1.7 Hz), 7.50 (1H, d, J = 7.9 Hz), 7.34 (2H, d, J = 8.2 Hz), 7.22 (2H, d, J = 8.2 Hz), 7.08 (2H, d, J = 8.2 Hz), 5.19 (1H, t, J = 5.6 Hz), 4.47 (2H, d, J = 5.8 Hz), 3.71-3.50 (2H, m), 3.27-3.13 (2H, m), 2.55-2.53 (1H, m), 2.04-1.90 (1H, m), 1.70-1.61 (1H, m).
30					free	426.2	¹ H-NMR (DMSO-D ₆) δ: 7.73 (2H, d, J = 8.5 Hz), 7.66-7.62 (1H, m), 7.55-7.49 (2H, m), 7.32 (2H, d, J = 7.3 Hz), 7.15 (2H, d, J = 8.2 Hz), 7.03 (2H, d, J = 8.2 Hz), 3.68-3.46 (6H, m), 3.22 (3H, s), 3.20-3.12 (1H, m), 2.77 (2H, t, J = 6.9 Hz), 2.02-1.87 (1H, m), 1.69-1.57 (1H, m).
31					free	412.1	¹ H-NMR (DMSO-D ₆) δ: 7.74 (2H, d, J = 8.5 Hz), 7.66-7.62 (1H, m), 7.55-7.49 (2H, m), 7.35-7.32 (2H, m), 7.13 (2H, d, J = 7.9 Hz), 7.03 (2H, d, J = 8.2 Hz), 4.63 (1H, t, J = 5.2 Hz), 3.62-3.56 (2H, m), 3.30-3.14 (4H, m), 2.73-2.66 (2H, m), 2.55-2.53 (1H, m), 2.03-1.90 (1H, m), 1.67-1.59 (1H, m).
32					free	426.2	¹ H-NMR (DMSO-D ₆) δ: 7.49-7.38 (2H, m), 7.39-7.30 (1H, m), 7.27-7.14 (2H, m), 7.10-6.98 (2H, m), 6.90-6.67 (4H, m), 4.26-4.08 (1H, m), 3.47-3.11 (5H, m), 2.93-2.75 (2H, m), 2.38-2.14 (2H, m), 1.78-1.44 (2H, m), 1.42-1.27 (2H, m).
33					free	438.2	¹ H-NMR (DMSO-D ₆) δ: 7.68 (2H, d, J = 8.2 Hz), 7.63-7.53 (1H, m), 7.46 (2H, t, J = 8.2 Hz), 7.26 (2H, d, J = 7.0 Hz), 7.12 (2H, d, J = 7.6 Hz), 6.95 (2H, d, J = 8.2 Hz), 4.65 (1H, s), 3.92-3.60 (5H, m), 3.09 (2H, s), 2.24-2.08 (1H, m), 2.01-1.86 (1H, m), 0.77 (2H, s), 0.65 (2H, s).






















[0792]

Table 5

34					free	440.2	¹ H-NMR (DMSO-D ₆) δ: 7.63 (2H, d, J = 7.0 Hz), 7.56 (1H, d, J = 5.7 Hz), 7.46 (2H, t, J = 10.2 Hz), 7.24 (2H, d, J = 7.6 Hz), 7.04 (2H, d, J = 7.6 Hz), 6.93 (2H, d, J = 7.0 Hz), 4.28 (1H, s), 3.64-3.48 (5H, m), 2.55 (2H, s), 1.98-1.87 (1H, m), 1.65-1.54 (1H, m), 0.96 (6H, s).
35					free	442.2	¹ H-NMR (DMSO-D ₆) δ: 7.78 (2H, d, J = 8.5 Hz), 7.67-7.61 (1H, m), 7.57-7.51 (2H, m), 7.36-7.30 (2H, m), 7.03 (2H, d, J = 8.5 Hz), 6.85 (2H, d, J = 8.5 Hz), 3.96-3.87 (1H, m), 3.85-3.68 (4H, m), 3.66-3.46 (2H, m), 3.27-3.21 (1H, m), 2.59-2.51 (1H, m), 2.29-2.17 (1H, m), 2.04-1.94 (1H, m), 1.13 (3H, d, J = 6.4 Hz).
36					free	418.1	¹ H-NMR (DMSO-D ₆) δ: 7.85-7.81 (1H, m), 7.71-7.67 (1H, m), 7.63 (1H, s), 7.54 (1H, d, J = 7.6 Hz), 7.38-7.32 (1H, m), 7.24-7.18 (1H, m), 7.10 (1H, d, J = 7.6 Hz), 7.06 (1H, d, J = 8.2 Hz), 6.96 (1H, d, J = 11.4 Hz), 4.12-4.07 (1H, m), 3.92-3.53 (4H, m), 2.31 (3H, s), 2.26-2.17 (1H, m), 2.02-1.91 (1H, m).
37					free	476.1	¹ H-NMR (DMSO-D ₆) δ: 7.79 (1H, t, J = 7.5 Hz), 7.70-7.67 (1H, m), 7.63 (1H, dd, J = 6.6, 1.5 Hz), 7.52 (1H, d, J = 8.1 Hz), 7.27 (1H, d, J = 10.3 Hz), 7.21 (1H, t, J = 7.9 Hz), 7.14 (1H, dd, J = 8.1, 1.5 Hz), 7.06 (1H, d, J = 7.7 Hz), 6.95 (1H, d, J = 11.4 Hz), 4.39 (1H, s), 3.68-3.55 (2H, m), 3.54-3.41 (2H, m), 3.21-3.12 (1H, m), 2.65 (2H, s), 2.03-1.88 (1H, m), 1.70-1.59 (1H, m), 1.04 (6H, s).
38					free	504.2	¹ H-NMR (CDCl ₃) δ: 7.76-7.69 (2H, m), 7.49-7.43 (2H, m), 7.09-6.96 (4H, m), 6.83 (1H, dd, J = 10.8, 0.9 Hz), 4.23 (2H, dd, J = 23.5, 9.2 Hz), 4.10 (2H, s), 3.75-3.64 (2H, m), 3.60 (2H, t, J = 6.6 Hz), 3.35 (3H, s), 3.00 (2H, s), 2.90-2.80 (4H, m).
39					free	488.1	¹ H-NMR (DMSO-D ₆) δ: 7.80 (1H, t, J = 7.6 Hz), 7.65 (1H, d, J = 7.0 Hz), 7.57 (1H, d, J = 10.8 Hz), 7.51 (1H, d, J = 7.6 Hz), 7.33 (1H, d, J = 10.2 Hz), 7.23 (1H, t, J = 7.9 Hz), 7.14-7.06 (2H, m), 7.00 (1H, d, J = 11.4 Hz), 3.86-3.64 (2H, m), 3.58-3.47 (4H, m), 3.23 (3H, s), 3.03-2.86 (2H, m), 2.81 (2H, t, J = 6.7 Hz), 2.77-2.55 (4H, m).
40					free	458.3	¹ H-NMR (DMSO-D ₆) δ: 7.82-7.77 (1H, m), 7.69-7.64 (1H, m), 7.60 (1H, d, J = 8.9 Hz), 7.54 (1H, d, J = 7.6 Hz), 7.27 (1H, d, J = 10.8 Hz), 7.15 (3H, d, J = 7.0 Hz), 7.03 (2H, d, J = 7.0 Hz), 3.74-3.53 (4H, m), 3.52-3.49 (1H, m), 3.22-3.13 (1H, m), 2.63 (2H, s), 2.07-1.97 (1H, m), 1.76-1.67 (1H, m), 1.03 (6H, s).

[0793]

Table 6

41					free	516.2	¹ H-NMR (DMSO-D ₆) δ: 7.80 (1H, t, J = 7.6 Hz), 7.61 (1H, dd, J = 7.8, 1.7 Hz), 7.53 (2H, dd, J = 10.9, 4.7 Hz), 7.29 (1H, dd, J = 10.7, 1.2 Hz), 7.22 (1H, t, J = 7.9 Hz), 7.15 (1H, dd, J = 7.9, 1.5 Hz), 7.07 (1H, dd, J = 7.9, 1.2 Hz), 6.95 (1H, d, J = 11.3 Hz), 4.57 (1H, br s), 4.40 (1H, s), 4.05 (1H, br s), 2.65 (2H, s), 2.35-2.21 (2H, m), 2.12-2.03 (1H, m), 1.99-1.81 (3H, m), 1.61-1.47 (3H, m), 1.06 (6H, d, J = 14.6 Hz).
42					free	396.3	¹ H-NMR (DMSO-D ₆) δ: 7.71 (2H, d, J = 7.7 Hz), 7.61 (1H, dd, J = 7.7, 1.5 Hz), 7.50 (1H, s), 7.46 (1H, dd, J = 6.1, 2.6 Hz), 7.30 (2H, d, J = 8.4 Hz), 7.07 (2H, d, J = 8.1 Hz), 6.98 (2H, d, J = 7.7 Hz), 3.63-3.44 (3H, m), 3.26-3.07 (2H, m), 2.27-2.16 (6H, m), 1.94-1.85 (1H, m), 1.75-1.68 (1H, m).
43					free	474.5	¹ H-NMR (DMSO-D ₆) δ: 7.75 (2H, d, J = 7.6 Hz), 7.67-7.61 (1H, m), 7.56-7.50 (2H, m), 7.45-7.30 (7H, m), 7.04 (2H, d, J = 7.6 Hz), 6.93 (2H, d, J = 7.6 Hz), 5.07 (2H, s), 4.17-4.07 (1H, m), 3.91-3.52 (4H, m), 2.26-2.15 (1H, m), 2.00-1.90 (1H, m).
44					free	527.2	¹ H-NMR (DMSO-D ₆) δ: 8.96 (1H, s), 7.74 (2H, dd, J = 6.7, 1.8 Hz), 7.67-7.64 (1H, m), 7.54 (4H, t, J = 12.7, 4.8 Hz), 7.37 (2H, dd, J = 8.2, 1.5 Hz), 7.32-7.28 (4H, m), 7.13 (2H, d, J = 8.2 Hz), 7.07-7.04 (1H, m), 3.68-3.43 (5H, m), 1.99 (1H, d, J = 6.1 Hz), 1.66 (1H, dt, J = 7.0, 2.2 Hz), 1.44 (2H, dd, J = 6.7, 4.3 Hz), 1.12 (2H, dd, J = 6.9, 4.4 Hz).
45					free	454.5	¹ H-NMR (DMSO-D ₆) δ: 7.73 (2H, d, J = 8.2 Hz), 7.67-7.62 (1H, m), 7.56-7.50 (2H, m), 7.32 (2H, d, J = 6.7 Hz), 7.16 (2H, d, J = 8.2 Hz), 7.05 (2H, d, J = 8.2 Hz), 4.19 (2H, t, J = 6.9 Hz), 3.72-3.43 (4H, m), 3.22-3.13 (1H, m), 2.86 (2H, t, J = 6.9 Hz), 2.06-1.88 (4H, m), 1.69-1.58 (1H, m).
46					free	426.5	¹ H-NMR (DMSO-D ₆) δ: 7.74 (2H, d, J = 8.2 Hz), 7.64 (1H, d, J = 8.2 Hz), 7.53 (1H, s), 7.49 (1H, d, J = 8.2 Hz), 7.33 (2H, d, J = 8.2 Hz), 7.13 (2H, d, J = 8.2 Hz), 7.03 (2H, d, J = 7.6 Hz), 3.71-3.45 (7H, m), 3.26-3.12 (1H, m), 2.69 (2H, t, J = 6.7 Hz), 2.34-2.19 (3H, m), 2.06-1.90 (1H, m), 1.83-1.72 (1H, m).
47					free	440.5	¹ H-NMR (DMSO-D ₆) δ: 7.73 (2H, d, J = 8.2 Hz), 7.64 (1H, d, J = 8.9 Hz), 7.56-7.47 (2H, m), 7.33 (2H, d, J = 6.2 Hz), 7.15 (2H, d, J = 7.6 Hz), 7.03 (2H, d, J = 7.6 Hz), 3.69-3.38 (6H, m), 3.26-3.08 (4H, m), 2.78 (2H, t, J = 6.7 Hz), 2.32-2.19 (3H, m), 2.04-1.87 (1H, m), 1.81-1.69 (1H, m).

[0794]

Table 7

48					free	466.2	¹ H-NMR (DMSO-D6) δ: 7.67 (2H, d, J = 7.6 Hz), 7.57 (1H, t, J = 8.6 Hz), 7.43 (2H, dd, J = 19.4, 4.8 Hz), 7.27 (2H, d, J = 8.2 Hz), 7.11 (2H, d, J = 8.2 Hz), 6.95 (2H, d, J = 5.7 Hz), 4.65-4.53 (1H, m), 3.68-3.38 (5H, m), 3.17-3.08 (1H, m), 2.71-2.58 (1H, m), 2.11 (3H, s), 2.04 (3H, s), 1.99-1.91 (1H, m), 1.74-1.58 (1H, m), 0.76 (2H, s), 0.65 (2H, s).
49					free	400.2	¹ H-NMR (DMSO-D6) δ: 7.71 (2H, d, J = 6.2 Hz), 7.62-7.56 (1H, m), 7.49 (2H, d, J = 8.2 Hz), 7.28 (2H, d, J = 7.0 Hz), 7.11 (1H, t, J = 7.9 Hz), 6.87 (1H, d, J = 10.8 Hz), 6.73 (1H, d, J = 7.6 Hz), 3.86-3.47 (5H, m), 2.21-2.09 (4H, m), 2.09-1.86 (1H, m).
50					free	402.1 404.1	¹ H-NMR (DMSO-D6) δ: 7.71 (2H, d, J = 8.2 Hz), 7.64-7.56 (1H, m), 7.54-7.46 (2H, m), 7.33-7.22 (4H, m), 7.07 (2H, d, J = 6.2 Hz), 4.04-3.50 (5H, m), 2.20-2.04 (1H, m), 1.96-1.82 (1H, m).
51					free	446.0 448.0	¹ H-NMR (DMSO-D6) δ: 7.71 (2H, d, J = 7.6 Hz), 7.65-7.57 (1H, m), 7.55-7.46 (2H, m), 7.43 (2H, d, J = 8.2 Hz), 7.27 (2H, d, J = 7.0 Hz), 7.01 (2H, d, J = 8.2 Hz), 3.78-3.50 (5H, m), 2.17-2.06 (1H, m), 1.92-1.82 (1H, m).
52					free	394.5	¹ H-NMR (DMSO-D6) δ: 7.75 (2H, d, J = 8.2 Hz), 7.67-7.59 (3H, m), 7.33 (2H, t, J = 8.6 Hz), 7.10 (2H, d, J = 7.6 Hz), 7.01 (2H, d, J = 7.6 Hz), 4.79-4.75 (1H, m), 4.51-4.47 (1H, m), 4.06 (1H, d, J = 18.4 Hz), 3.73-3.52 (2H, m), 3.15-3.02 (1H, m), 2.27 (3H, s), 2.04-1.91 (1H, m), 1.82-1.66 (1H, m).
53					free	411.1	¹ H-NMR (DMSO-D6) δ: 7.74 (2H, dd, J = 6.6, 2.0 Hz), 7.65 (1H, s), 7.54-7.50 (2H, m), 7.34-7.33 (2H, m), 7.14 (2H, d, J = 8.2 Hz), 7.06 (2H, d, J = 6.2 Hz), 3.59 (5H, ddd, J = 14.0, 4.3, 2.5 Hz), 2.85 (2H, s), 2.69-2.67 (2H, m), 1.70-1.50 (1H, m), 1.24 (1H, s).
54					free	494	¹ H-NMR (DMSO-D6) δ: 7.71 (2H, d, J = 8.2 Hz), 7.63-7.56 (3H, m), 7.53-7.45 (2H, m), 7.26 (2H, d, J = 7.6 Hz), 6.86 (2H, d, J = 8.2 Hz), 3.87-3.47 (5H, m), 2.23-2.08 (1H, m), 1.98-1.86 (1H, m).
55					free	453.2	¹ H-NMR (DMSO-D6) δ: 7.85-7.79 (1H, m), 7.68 (2H, d, J = 8.2 Hz), 7.61-7.55 (1H, m), 7.52-7.44 (2H, m), 7.27-7.22 (2H, m), 7.06 (2H, d, J = 7.6 Hz), 6.97 (2H, d, J = 7.6 Hz), 3.83-3.52 (5H, m), 3.17-3.12 (2H, m), 2.60 (2H, t, J = 7.3 Hz), 2.22-2.10 (1H, m), 1.98-1.86 (1H, m), 1.69 (3H, s).
56					free	410.2	¹ H-NMR (DMSO-D6) δ: 7.67 (2H, d, J = 8.2 Hz), 7.63-7.55 (1H, m), 7.54-7.45 (2H, m), 7.25 (2H, d, J = 8.2 Hz), 7.04 (2H, d, J = 8.2 Hz), 6.95 (2H, d, J = 7.6 Hz), 3.88-3.48 (5H, m), 2.48-2.44 (2H, m), 2.22-2.08 (1H, m), 1.97-1.86 (1H, m), 1.49 (2H, dd, J = 14.9, 7.3 Hz), 0.80 (3H, t, J = 7.3 Hz).









[0795]

Table 8

57					free	418.2	¹ H-NMR (DMSO-D ₆) δ: 7.83-7.76 (3H, m), 7.73-7.61 (5H, m), 7.59-7.53 (1H, m), 7.49-7.41 (2H, m), 7.29 (2H, d, J = 7.6 Hz), 7.03 (1H, d, J = 8.9 Hz), 4.03-3.55 (5H, m), 2.24-2.12 (1H, m), 1.98-1.88 (1H, m).
58					free	452.1	¹ H-NMR (DMSO-D ₆) δ: 7.74 (2H, d, J = 8.2 Hz), 7.63 (1H, d, J = 8.2 Hz), 7.53-7.51 (1H, m), 7.49 (1H, d, J = 7.9 Hz), 7.34 (2H, d, J = 8.5 Hz), 7.19 (2H, d, J = 8.2 Hz), 7.03 (2H, d, J = 8.2 Hz), 4.67 (1H, t, J = 5.5 Hz), 3.71-3.47 (6H, m), 3.27-3.16 (1H, m), 2.35-2.21 (3H, m), 2.03-1.94 (1H, m), 1.81-1.72 (1H, m), 0.85-0.82 (2H, m), 0.74-0.70 (2H, m).
59					free	438.1	¹ H-NMR (DMSO-D ₆) δ: 7.76-7.62 (3H, m), 7.58-7.50 (2H, m), 7.36-7.26 (2H, m), 7.14-7.02 (4H, m), 5.34-5.31 (1H, m), 3.89-3.58 (6H, m), 3.23-3.13 (1H, m), 2.05-1.95 (1H, m), 1.74-1.65 (1H, m), 0.94-0.84 (4H, m).
60					free	422.1	¹ H-NMR (DMSO-D ₆) δ: 7.67 (2H, d, J = 7.6 Hz), 7.62-7.55 (1H, m), 7.51-7.43 (2H, m), 7.26 (2H, d, J = 7.6 Hz), 7.03 (4H, dd, J = 28.6, 7.6 Hz), 6.15 (1H, s), 3.68-3.48 (5H, m), 2.01-1.89 (1H, m), 1.79 (3H, s), 1.74 (3H, s), 1.67-1.59 (1H, m).
61					free	454.3	¹ H-NMR (DMSO-D ₆) δ: 7.74 (2H, d, J = 7.0 Hz), 7.67-7.61 (1H, m), 7.57-7.49 (2H, m), 7.33 (2H, d, J = 7.6 Hz), 7.11 (2H, d, J = 7.6 Hz), 7.02 (2H, d, J = 7.0 Hz), 3.82-3.56 (4H, m), 3.53-3.47 (1H, m), 3.21-3.13 (1H, m), 2.62-2.55 (2H, m), 2.13-2.03 (1H, m), 1.86-1.75 (1H, m), 1.64-1.57 (2H, m), 1.12 (6H, s).
62					free	452.2	¹ H-NMR (DMSO-D ₆) δ: 7.74 (2H, d, J = 8.2 Hz), 7.68-7.62 (1H, m), 7.58-7.51 (2H, m), 7.32 (2H, d, J = 7.0 Hz), 7.12 (2H, d, J = 7.6 Hz), 7.02 (2H, d, J = 7.0 Hz), 4.15-4.08 (1H, m), 3.94-3.55 (4H, m), 3.53-3.48 (2H, m), 3.19-3.14 (1H, m), 2.77-2.69 (2H, m), 2.41-2.39 (2H, m), 2.31-2.17 (1H, m), 2.02-1.94 (1H, m), 0.91-0.89 (2H, m).
63					free	430.3	¹ H-NMR (DMSO-D ₆) δ: 8.30 (1H, s), 7.80 (1H, t, J = 7.0 Hz), 7.69-7.64 (1H, m), 7.58 (1H, d, J = 7.6 Hz), 7.51 (1H, d, J = 8.2 Hz), 7.35 (1H, d, J = 11.4 Hz), 7.16 (2H, d, J = 7.6 Hz), 7.11 (1H, d, J = 8.2 Hz), 7.05 (2H, d, J = 7.0 Hz), 3.69-3.52 (6H, m), 3.24-3.17 (1H, m), 2.71 (2H, t, J = 6.3 Hz), 2.04-1.93 (1H, m), 1.73-1.63 (1H, m).
64					free	458.2	¹ H-NMR (DMSO-D ₆) δ: 7.73-7.63 (3H, m), 7.59 (1H, d, J = 10.8 Hz), 7.51 (1H, d, J = 8.2 Hz), 7.31 (2H, d, J = 7.6 Hz), 7.20-7.14 (1H, m), 7.03 (1H, d, J = 7.6 Hz), 6.92 (1H, d, J = 11.4 Hz), 3.70-3.54 (4H, m), 3.52-3.49 (1H, m), 3.24-3.20 (1H, m), 2.64 (2H, s), 2.08-1.96 (1H, m), 1.76-1.66 (1H, m), 1.04 (6H, s).

[0796]

Table 9

65					free	472.2	¹ H-NMR (DMSO-D ₆) δ: 7.80 (1H, t, J = 7.3 Hz), 7.69-7.64 (1H, m), 7.59 (1H, d, J = 8.2 Hz), 7.52 (1H, d, J = 8.2 Hz), 7.33 (1H, d, J = 10.8 Hz), 7.15-7.09 (3H, m), 7.04 (2H, d, J = 7.6 Hz), 3.70-3.57 (4H, m), 3.54-3.49 (1H, m), 3.24-3.17 (1H, m), 2.63-2.57 (2H, m), 2.07-1.93 (1H, m), 1.77-1.64 (1H, m), 1.64-1.58 (2H, m), 1.12 (6H, s).
66					free	452.2	¹ H-NMR (DMSO-D ₆) δ: 7.74 (2H, d, J = 7.6 Hz), 7.67-7.61 (1H, m), 7.56-7.49 (2H, m), 7.33 (2H, d, J = 8.2 Hz), 7.17 (2H, d, J = 7.0 Hz), 7.03 (2H, d, J = 7.0 Hz), 3.82-3.52 (5H, m), 3.51 (2H, s), 3.21 (3H, s), 2.15-2.01 (1H, m), 1.89-1.75 (1H, m), 0.89-0.78 (4H, m).
67					free	456.2	¹ H-NMR (DMSO-D ₆) δ: 7.73 (2H, d, J = 6.3 Hz), 7.67-7.61 (1H, m), 7.57-7.48 (2H, m), 7.39-7.24 (4H, m), 7.02 (1H, d, J = 7.6 Hz), 3.73-3.52 (4H, m), 3.51 (1H, s), 3.19-3.10 (1H, m), 2.54 (2H, s), 1.72-1.58 (1H, m), 1.48-1.35 (1H, m), 0.83-0.42 (4H, m).
68					free	464.2	¹ H-NMR (DMSO-D ₆) δ: 7.74 (2H, d, J = 8.5 Hz), 7.67-7.61 (1H, m), 7.57-7.48 (2H, m), 7.39-7.33 (3H, m), 7.28 (2H, d, J = 8.2 Hz), 7.02 (1H, d, J = 8.2 Hz), 5.16 (1H, s), 3.78-3.54 (4H, m), 3.21-3.13 (1H, m), 2.04-1.95 (1H, m), 1.67-1.58 (1H, m), 0.82-0.70 (4H, m), 0.57-0.42 (4H, m).
69					free	444.1	¹ H-NMR (DMSO-D ₆) δ: 7.81 (1H, s), 7.72-7.47 (3H, m), 7.37-7.31 (1H, m), 7.18 (2H, d, J = 7.6 Hz), 7.13-7.01 (3H, m), 4.29-4.11 (1H, m), 3.88-3.56 (4H, m), 3.52 (2H, t, J = 6.7 Hz), 3.22 (3H, s), 2.79 (2H, t, J = 6.4 Hz), 2.23-2.03 (1H, m), 1.54-1.34 (1H, m).
70					free	430	¹ H-NMR (DMSO-D ₆) δ: 7.74 (2H, d, J = 8.2 Hz), 7.69-7.63 (1H, m), 7.58 (1H, d, J = 12.5 Hz), 7.49 (1H, d, J = 7.9 Hz), 7.32 (2H, d, J = 7.6 Hz), 7.19 (1H, t, J = 7.9 Hz), 7.06 (1H, d, J = 7.6 Hz), 6.96 (1H, d, J = 11.0 Hz), 4.70-4.64 (1H, m), 3.72-3.46 (6H, m), 3.26-3.18 (1H, m), 2.74-2.68 (2H, m), 2.08-1.98 (1H, m), 1.78-1.68 (1H, m).
71					free	444.1	¹ H-NMR (DMSO-D ₆) δ: 7.73 (2H, d, J = 6.7 Hz), 7.67-7.64 (1H, m), 7.57 (1H, dd, J = 10.8, 1.4 Hz), 7.50 (1H, d, J = 7.9 Hz), 7.31 (2H, d, J = 7.6 Hz), 7.20 (1H, t, J = 7.8 Hz), 7.07 (1H, dd, J = 7.9, 1.2 Hz), 6.97 (1H, d, J = 11.3 Hz), 3.68-3.45 (6H, m), 3.23 (3H, s), 3.21-3.15 (1H, m), 2.80 (2H, t, J = 6.6 Hz), 2.04-1.93 (1H, m), 1.72-1.62 (1H, m).
72					free	440.1	¹ H-NMR (DMSO-D ₆) δ: 7.74 (2H, d, J = 8.2 Hz), 7.64 (1H, t, J = 7.3 Hz), 7.55-7.49 (2H, m), 7.34 (2H, d, J = 7.0 Hz), 7.28 (2H, d, J = 8.2 Hz), 7.05 (2H, d, J = 8.2 Hz), 4.69-4.65 (1H, m), 3.70-3.37 (6H, m), 3.25-3.18 (1H, m), 2.08-1.95 (1H, m), 1.76-1.68 (1H, m), 1.19 (6H, s).

















[0797]

Table 10

73					free	414.1	¹ H-NMR (DMSO-D ₆) δ: 7.75 (2H, d, J = 7.9 Hz), 7.65 (1H, t, J = 7.6 Hz), 7.58-7.49 (2H, m), 7.33 (2H, d, J = 7.3 Hz), 7.19 (2H, d, J = 7.9 Hz), 7.07 (2H, d, J = 7.6 Hz), 4.63 (2H, dt, J = 47.2, 6.2 Hz), 3.72-3.47 (4H, m), 3.27-3.21 (1H, m), 2.95 (2H, dt, J = 25.2, 6.2 Hz), 2.10-1.99 (1H, m), 1.81-1.71 (1H, m).
74					free	488.1	¹ H-NMR (DMSO-D ₆) δ: 7.84-7.78 (1H, m), 7.71-7.62 (2H, m), 7.56-7.51 (1H, m), 7.36-7.30 (1H, m), 7.25-7.21 (1H, m), 7.13-7.09 (2H, m), 7.03-6.98 (1H, m), 4.09-3.49 (10H, m), 3.22 (3H, s), 2.84-2.78 (2H, m), 2.19-2.10 (2H, m).
75					free	502.1	¹ H-NMR (DMSO-D ₆) δ: 7.84-7.79 (2H, m), 7.73-7.70 (1H, m), 7.57-7.52 (1H, m), 7.32 (1H, d, J = 10.4 Hz), 7.25-7.20 (1H, m), 7.11 (2H, d, J = 7.3 Hz), 7.00 (1H, d, J = 11.0 Hz), 4.19 (1H, d, J = 8.2 Hz), 4.10 (1H, d, J = 8.2 Hz), 3.90 (1H, d, J = 9.8 Hz), 3.75 (1H, d, J = 10.1 Hz), 3.55-3.50 (2H, m), 3.25-3.07 (7H, m), 2.88-2.78 (2H, m), 1.86-1.73 (2H, m), 1.67-1.52 (2H, m).
76					free	502.1	¹ H-NMR (DMSO-D ₆) δ: 7.82-7.77 (1H, m), 7.72-7.62 (2H, m), 7.57-7.51 (1H, m), 7.29-7.25 (1H, m), 7.22-7.18 (1H, m), 7.16-7.11 (1H, m), 7.09-7.05 (1H, m), 6.98-6.92 (1H, m), 4.40 (1H, s), 3.98-3.44 (6H, m), 2.67-2.64 (2H, m), 2.18-2.05 (2H, m), 2.04-1.90 (2H, m), 1.04 (6H, s).
77					free	516.1	¹ H-NMR (DMSO-D ₆) δ: 7.84-7.71 (3H, m), 7.57-7.54 (1H, m), 7.29-7.24 (1H, m), 7.21-7.17 (1H, m), 7.15-7.11 (1H, m), 7.08-7.05 (1H, m), 6.97-6.93 (1H, m), 4.39 (1H, s), 4.14-4.02 (2H, m), 3.85-3.68 (2H, m), 2.95-2.84 (2H, m), 2.71-2.62 (4H, m), 1.78-1.68 (2H, m), 1.52-1.42 (2H, m), 1.04 (6H, s).
78					free	516.2	¹ H-NMR (CD ₃ OD) δ: 7.64-7.55 (4H, m), 7.22-7.08 (4H, m), 6.92 (1H, d, J = 11.0 Hz), 4.01-3.68 (1H, m), 3.77-3.71 (2H, m), 3.67 (3H, s), 3.63-3.58 (3H, m), 3.47-3.40 (2H, m), 3.23-3.15 (2H, m), 2.87 (2H, t, J = 6.6 Hz), 2.09-2.01 (2H, m), 1.83-1.65 (4H, m).
79					free	502.1	¹ H-NMR (DMSO-D ₆) δ: 7.83-7.00 (9H, m), 3.82 (12H, tt, J = 147.2, 54.5 Hz), 3.24 (3H, d, J = 5.5 Hz), 2.82 (2H, t, J = 6.4 Hz), 1.76 (1H, s), 1.48 (1H, s).
80					free	476.2	¹ H-NMR (CD ₃ OD) δ: 7.65-7.54 (4H, m), 7.13 (4H, ddd, J = 29.3, 15.0, 6.6 Hz), 6.90 (1H, d, J = 11.4 Hz), 4.02-3.97 (1H, m), 3.90-3.82 (1H, m), 3.74-3.67 (2H, m), 3.59 (2H, t, J = 6.4 Hz), 3.49-3.43 (2H, m), 3.39-3.35 (2H, m), 3.31 (3H, s), 2.85 (2H, t, J = 6.6 Hz), 2.24-2.05 (2H, m).
81					free	488.2	¹ H-NMR (DMSO-D ₆) δ: 7.83-7.79 (1H, m), 7.72-7.60 (2H, m), 7.55-7.51 (1H, m), 7.36-7.30 (1H, m), 7.27-7.21 (1H, m), 7.15-7.08 (2H, m), 7.03-6.98 (1H, m), 3.88-3.38 (10H, m), 3.23 (3H, s), 2.85-2.79 (2H, m), 2.38-2.02 (2H, m).



















[0798]

Table 11

82					free	474.1	¹ H-NMR (DMSO-D6) δ: 7.82 (1H, t, J = 7.3 Hz), 7.74-7.60 (2H, m), 7.56 (1H, d, J = 7.9 Hz), 7.39-7.30 (1H, m), 7.24 (1H, t, J = 7.6 Hz), 7.14-7.09 (2H, m), 7.01 (1H, d, J = 11.0 Hz), 4.57 (1H, s), 4.27 (1H, s), 3.79-3.72 (1H, m), 3.66-3.61 (1H, m), 3.57-3.51 (2H, m), 3.46-3.40 (1H, m), 3.22 (3H, s), 3.21-3.17 (1H, m), 2.84-2.79 (2H, m), 2.14-2.05 (1H, m), 1.90-1.73 (1H, m).
83					free	502.1	¹ H-NMR (DMSO-D6) δ: 7.81 (1H, t, J = 7.6 Hz), 7.59-7.51 (3H, m), 7.33 (1H, dd, J = 10.7, 1.2 Hz), 7.24 (1H, t, J = 7.9 Hz), 7.11 (2H, d, J = 8.2 Hz), 7.00 (1H, d, J = 10.4 Hz), 3.79-3.57 (6H, m), 3.53 (2H, t, J = 6.7 Hz), 3.23 (3H, s), 2.81 (2H, t, J = 6.7 Hz), 2.25-2.17 (2H, m), 2.01-1.81 (4H, m).
84					free	502.2	¹ H-NMR (DMSO-D6) δ: 7.84-7.79 (2H, m), 7.72 (1H, s), 7.54 (1H, d, J = 7.9 Hz), 7.33 (1H, d, J = 10.4 Hz), 7.22 (1H, t, J = 7.9 Hz), 7.14-7.09 (2H, m), 7.00 (1H, d, J = 11.3 Hz), 4.15 (2H, s), 3.83 (2H, s), 3.53 (2H, t, J = 6.6 Hz), 3.22 (3H, s), 2.97-2.85 (4H, m), 2.81 (2H, t, J = 6.6 Hz), 1.89-1.63 (4H, m).
85					free	474.2	¹ H-NMR (DMSO-D6) δ: 7.81 (1H, t, J = 7.6 Hz), 7.71-7.62 (2H, m), 7.59-7.51 (1H, m), 7.38-7.30 (1H, m), 7.27-7.22 (1H, m), 7.15-7.08 (2H, m), 7.03-6.98 (1H, m), 4.72 (1H, s), 4.39 (1H, s), 3.86-3.61 (2H, m), 3.56-3.49 (2H, m), 3.22 (3H, s), 3.10-2.93 (2H, m), 2.84-2.79 (2H, m), 1.91-1.60 (2H, m).
86					free	488.2	¹ H-NMR (DMSO-D6) δ: 7.81 (1H, t, J = 7.5 Hz), 7.74-7.53 (3H, m), 7.38-7.31 (1H, m), 7.24 (1H, t, J = 7.9 Hz), 7.14-7.09 (2H, m), 7.01 (1H, d, J = 11.0 Hz), 4.76 (1H, s), 4.44 (1H, s), 4.02-3.85 (1H, m), 3.72-3.57 (1H, m), 3.53 (2H, t, J = 6.7 Hz), 3.22 (3H, s), 3.12-2.97 (2H, m), 2.81 (2H, t, J = 6.7 Hz), 2.63 (3H, s), 2.18-1.91 (2H, m).
87					free	488.1	¹ H-NMR (DMSO-D6) δ: 7.81 (1H, t, J = 7.5 Hz), 7.75-7.51 (3H, m), 7.34 (1H, t, J = 10.5 Hz), 7.24 (1H, t, J = 7.9 Hz), 7.14-7.08 (2H, m), 7.01 (1H, d, J = 11.3 Hz), 4.72 (1H, s), 4.39 (1H, s), 3.89-3.59 (2H, m), 3.53 (2H, t, J = 6.6 Hz), 3.22 (3H, s), 3.08-2.90 (2H, m), 2.81 (2H, t, J = 6.7 Hz), 2.54 (3H, s), 2.16-1.84 (2H, m).
88					free	488.1	¹ H-NMR (DMSO-D6) δ: 7.80-7.75 (1H, m), 7.55-7.49 (2H, m), 7.47 (1H, d, J = 1.5 Hz), 7.28 (1H, dd, J = 10.5, 1.4 Hz), 7.22 (1H, t, J = 7.9 Hz), 7.10 (2H, dt, J = 10.5, 3.8 Hz), 6.97 (1H, d, J = 11.3 Hz), 4.31-4.21 (1H, m), 3.54-3.29 (5H, m), 3.21 (3H, s), 3.00-2.93 (1H, m), 2.79 (2H, t, J = 6.6 Hz), 1.76-1.48 (4H, m).
89					free	443.1	¹ H-NMR (DMSO-D6) δ: 9.34 (1H, s), 7.89 (1H, s), 7.73-7.58 (5H, m), 7.42-7.33 (1H, m), 7.11 (1H, d, J = 8.2 Hz), 7.07-7.01 (1H, m), 3.85-3.50 (5H, m), 2.20-2.07 (1H, m), 1.97-1.82 (1H, m).

[0799]

Table 12

90					free	441.2	¹ H-NMR (DMSO-D ₆) δ: 8.19 (1H, d, J = 1.9 Hz), 8.08 (1H, s), 8.04 (1H, d, J = 1.9 Hz), 7.72 (1H, t, J = 7.6 Hz), 7.68-7.65 (1H, m), 7.61 (2H, d, J = 8.2 Hz), 7.44-7.37 (1H, m), 7.07-6.99 (1H, m), 3.97 (3H, s), 3.66-3.42 (5H, m), 2.23-2.09 (1H, m), 1.99-1.86 (1H, m).
91					free	440.2	¹ H-NMR (DMSO-D ₆) δ: 8.23 (1H, s), 7.69 (1H, t, J = 7.6 Hz), 7.66-7.60 (1H, m), 7.54 (2H, d, J = 7.6 Hz), 7.43 (2H, d, J = 6.3 Hz), 7.35-7.28 (1H, m), 7.06-7.00 (1H, m), 6.98-6.91 (1H, m), 3.86-3.53 (8H, m), 2.22-2.10 (1H, m), 1.99-1.89 (1H, m).
92					free	440.2	¹ H-NMR (DMSO-D ₆) δ: 7.75-7.71 (1H, m), 7.70-7.66 (1H, m), 7.64 (1H, s), 7.62-7.58 (2H, m), 7.50 (1H, d, J = 8.9 Hz), 7.37-7.32 (2H, m), 7.08 (1H, d, J = 7.9 Hz), 7.04 (1H, d, J = 8.5 Hz), 4.41-4.12 (4H, m), 3.32-3.24 (1H, m), 3.16 (3H, s), 2.21-2.10 (1H, m), 1.82-1.59 (1H, m).
93					free	440.2	¹ H-NMR (DMSO-D ₆) δ: 8.20 (1H, s), 7.75 (1H, t, J = 7.5 Hz), 7.69-7.64 (1H, m), 7.62-7.58 (2H, m), 7.55 (1H, s), 7.44 (1H, d, J = 9.2 Hz), 7.40-7.33 (1H, m), 7.08 (1H, d, J = 8.4 Hz), 6.86 (1H, d, J = 9.2 Hz), 4.13 (3H, s), 3.91-3.40 (4H, m), 3.19-3.09 (1H, m), 2.27-2.17 (1H, m), 2.06-1.93 (1H, m).
94					free	478.2	¹ H-NMR (DMSO-D ₆) δ: 7.80 (1H, t, J = 7.5 Hz), 7.57-7.54 (1H, m), 7.51-7.49 (1H, m), 7.47 (1H, d, J = 7.9 Hz), 7.31 (1H, ddd, J = 10.6, 5.3, 1.4 Hz), 7.24 (1H, td, J = 7.9, 2.4 Hz), 7.10 (2H, d, J = 7.6 Hz), 7.00 (1H, d, J = 11.3 Hz), 4.00-3.56 (4H, m), 3.53 (2H, t, J = 6.7 Hz), 3.32-3.27 (1H, m), 3.23 (3H, s), 2.81 (2H, t, J = 6.6 Hz), 2.16-2.04 (1H, m), 1.85-1.74 (1H, m).
95					free	458.2	¹ H-NMR (DMSO-D ₆) δ: 8.34 (1H, s), 7.78 (1H, t, J = 7.5 Hz), 7.71 (1H, t, J = 6.4 Hz), 7.67-7.63 (2H, m), 7.60 (1H, d, J = 7.9 Hz), 7.47-7.35 (2H, m), 7.11 (1H, d, J = 7.9 Hz), 3.81 (3H, s), 3.78-3.60 (5H, m), 2.31-2.19 (1H, m), 2.07-1.96 (1H, m).
96					free	459.2	¹ H-NMR (DMSO-D ₆) δ: 7.95-8.15 (2H, m), 7.60-7.79 (4H, m), 7.37-7.46 (1H, m), 7.07-7.13 (1H, m), 4.24 (3H, s), 4.09-3.49 (5H, m), 1.95-2.20 (2H, m).
97					free	426.3	¹ H-NMR (DMSO-D ₆) δ: 8.30 (1H, d, J = 9.9 Hz), 7.75 (2H, d, J = 8.4 Hz), 7.57 (1H, d, J = 8.1 Hz), 7.52 (1H, d, J = 7.7 Hz), 7.46 (1H, s), 7.32 (2H, d, J = 8.1 Hz), 7.16-7.09 (3H, m), 4.54-4.52 (1H, m), 4.19-4.07 (1H, m), 3.45-3.37 (1H, m), 2.11-1.40 (8H, m).
98					free	442.2 444.2	¹ H-NMR (DMSO-D ₆) δ: 8.30 (1H, d, J = 8.4 Hz), 7.76 (2H, d, J = 8.4 Hz), 7.58 (1H, d, J = 7.7 Hz), 7.53 (1H, d, J = 8.1 Hz), 7.46 (1H, s), 7.35-7.32 (3H, m), 7.13 (2H, d, J = 8.4 Hz), 4.65-4.58 (1H, m), 4.16-4.08 (1H, m), 3.44-3.38 (1H, m), 2.04-1.47 (8H, m).















[0800]

Table 13

99					free	402.2	¹ H-NMR (DMSO-D ₆) δ: 8.36 (2H, d, J = 6.7 Hz), 7.90 (1H, t, J = 7.8 Hz), 7.81 (1H, d, J = 13.1 Hz), 7.75 (1H, dd, J = 7.9, 1.8 Hz), 7.64 (2H, d, J = 7.6 Hz), 7.33 (2H, d, J = 6.7 Hz), 7.26 (2H, d, J = 6.4 Hz), 3.95-3.72 (4H, m), 3.51-3.45 (1H, m), 2.50 (3H, s), 2.31-2.21 (1H, m), 2.02-1.93 (1H, m).
100					free	430.3	¹ H-NMR (DMSO-D ₆) δ: 7.94-7.86 (3H, m), 7.73-7.65 (1H, m), 7.62-7.54 (2H, m), 7.50-7.43 (1H, m), 7.29-7.22 (1H, m), 7.19-7.14 (1H, m), 3.93-3.57 (5H, m), 3.03 (6H, s), 2.30-2.18 (1H, m), 2.06-1.94 (1H, m).
101					free	441.2	¹ H-NMR (DMSO-D ₆) δ: 7.93-7.86 (2H, m), 7.76-7.62 (4H, m), 7.41 (1H, t, J = 9.5 Hz), 7.19 (1H, d, J = 8.8 Hz), 7.10-7.05 (1H, m), 4.28 (3H, s), 3.72-3.41 (5H, m), 2.27-2.18 (1H, m), 2.05-1.95 (1H, m).
102					free	476.2	¹ H-NMR (DMSO-D ₆) δ: 8.27 (1H, s), 7.80 (1H, t, J = 7.6 Hz), 7.73 (1H, t, J = 5.5 Hz), 7.67 (1H, s), 7.63 (1H, d, J = 7.9 Hz), 7.48-7.40 (2H, m), 7.16-7.10 (1H, m), 3.98 (3H, s), 3.92-3.44 (5H, m), 2.31-2.20 (1H, m), 2.07-1.96 (1H, m).
103					free	454.2	¹ H-NMR (DMSO-D ₆) δ: 7.79-7.73 (1H, m), 7.70-7.65 (1H, m), 7.63-7.55 (2H, m), 7.39 (2H, d, J = 8.2 Hz), 7.32 (1H, d, J = 1.2 Hz), 7.12-7.07 (1H, m), 6.96-6.91 (1H, m), 3.72 (3H, t, J = 6.7 Hz), 3.67-3.48 (5H, m), 2.54 (3H, s), 1.99 (1H, dd, J = 9.0, 3.8 Hz), 1.67-1.62 (1H, m).
104					free	476.2	¹ H-NMR (DMSO-D ₆) δ: 7.91-7.86 (3H, m), 7.81-7.65 (4H, m), 7.57 (1H, d, J = 1.5 Hz), 7.53 (2H, dd, J = 6.6, 3.4 Hz), 7.46 (1H, dd, J = 10.4, 1.2 Hz), 7.16-7.12 (2H, m), 4.65 (1H, s), 4.19 (1H, s), 3.51-3.43 (1H, m), 2.11-1.49 (8H, m).
105					free	495.2	¹ H-NMR (DMSO-D ₆) δ: 8.93 (1H, d, J = 2.1 Hz), 8.44 (1H, d, J = 8.9 Hz), 7.81 (1H, d, J = 8.5 Hz), 7.78-7.61 (5H, m), 7.50-7.43 (2H, m), 7.13 (1H, d, J = 6.7 Hz), 4.65 (1H, s), 4.18 (1H, s), 3.49-3.40 (1H, m), 2.09-1.17 (8H, m).
106					free	497.3	¹ H-NMR (DMSO-D ₆) δ: 7.85 (1H, t, J = 7.6 Hz), 7.59-7.38 (4H, m), 7.16 (1H, dd, J = 7.9, 0.9 Hz), 6.58 (1H, d, J = 8.2 Hz), 6.52 (1H, d, J = 1.8 Hz), 6.48 (1H, dd, J = 9.2, 1.8 Hz), 4.61 (1H, s), 4.22-4.13 (3H, m), 3.45-3.40 (1H, m), 3.26-2.22 (2H, m), 2.81 (3H, s), 2.06-1.15 (8H, m).
107					free	477.3	¹ H-NMR (DMSO-D ₆) δ: 8.92 (1H, dd, J = 4.3, 1.8 Hz), 8.39-8.35 (1H, m), 7.95-7.88 (2H, m), 7.80-7.73 (2H, m), 7.69 (1H, dd, J = 7.8, 1.7 Hz), 7.59-7.55 (2H, m), 7.50 (1H, dd, J = 10.4, 1.5 Hz), 7.30 (1H, dd, J = 8.4, 1.7 Hz), 7.15 (1H, dd, J = 8.2, 1.5 Hz), 4.68 (1H, s), 4.25 (1H, s), 3.69-3.57 (1H, m), 2.14-1.56 (8H, m).














[0801]

Table 14

108					free	498.2	¹ H-NMR (DMSO-D ₆) δ: 8.23 (1H, s), 7.77 (1H, t, J = 7.5 Hz), 7.65-7.62 (2H, m), 7.59 (1H, d, J = 7.9 Hz), 7.55 (1H, d, J = 1.5 Hz), 7.43-7.38 (2H, m), 7.12 (1H, dd, J = 7.4, 1.5 Hz), 4.65 (1H, s), 4.18 (1H, s), 3.80 (3H, s), 3.39-3.35 (1H, m), 2.11-1.66 (7H, m), 1.56-1.44 (1H, m).
109					free	499.2	¹ H-NMR (DMSO-D ₆) δ: 8.14 (1H, d, J = 6.4 Hz), 7.77 (1H, t, J = 7.5 Hz), 7.72-7.63 (3H, m), 7.58 (1H, d, J = 1.5 Hz), 7.46 (1H, d, J = 10.4 Hz), 7.14 (1H, dd, J = 8.1, 1.4 Hz), 4.65 (1H, s), 4.26 (3H, s), 4.16 (1H, s), 3.27-3.33 (1H, m), 2.12-1.46 (6H, m).
110					free	498.2	¹ H-NMR (DMSO-D ₆) δ: 8.11 (1H, s), 7.76 (1H, t, J = 7.5 Hz), 7.65 (1H, dd, J = 7.9, 1.8 Hz), 7.61 (1H, d, J = 7.9 Hz), 7.58 (1H, d, J = 1.5 Hz), 7.50 (1H, d, J = 8.5 Hz), 7.42 (1H, dd, J = 10.5, 1.4 Hz), 7.25 (1H, dd, J = 8.5, 7.0 Hz), 7.10 (1H, dd, J = 7.9, 1.5 Hz), 4.64 (1H, s), 4.16 (1H, s), 4.06 (3H, s), 3.42-3.35 (1H, m), 2.08-1.44 (6H, m).
111					free	462.2	¹ H-NMR (DMSO-D ₆) δ: 8.32 (1H, s), 7.74 (2H, d, J = 8.5 Hz), 7.63 (2H, t, J = 5.5 Hz), 7.56 (1H, s), 7.50 (1H, s), 7.45 (1H, d, J = 8.9 Hz), 7.37 (2H, d, J = 8.5 Hz), 6.88 (2H, dd, J = 8.9, 1.8 Hz), 4.72-4.65 (1H, m), 4.32-4.22 (1H, m), 4.15 (3H, s), 2.12-1.54 (9H, m).
112					free	544.2	¹ H-NMR (DMSO-D ₆) δ: 7.82 (1H, t, J = 7.5 Hz), 7.64 (1H, dd, J = 7.8, 1.7 Hz), 7.56-7.53 (2H, m), 7.35 (1H, d, J = 10.7 Hz), 7.25 (1H, t, J = 7.9 Hz), 7.15-7.10 (2H, m), 7.01 (1H, d, J = 11.9 Hz), 4.67-4.61 (1H, m), 4.17-4.10 (1H, m), 3.54 (2H, t, J = 6.7 Hz), 3.47-3.45 (1H, m), 3.23 (3H, s), 3.02-2.94 (1H, m), 2.82 (2H, t, J = 6.7 Hz), 2.04-1.31 (8H, m), 1.01 (6H, d, J = 6.1 Hz).
113					free	558.2	¹ H-NMR (DMSO-D ₆) δ: 7.81 (1H, t, J = 7.5 Hz), 7.64 (1H, dd, J = 7.8, 1.7 Hz), 7.58-7.54 (2H, m), 7.29 (1H, d, J = 9.2 Hz), 7.22 (1H, t, J = 7.8 Hz), 7.16 (1H, dd, J = 8.1, 1.4 Hz), 7.07 (1H, d, J = 7.9 Hz), 6.96 (1H, d, J = 11.3 Hz), 4.66-4.61 (1H, m), 4.42-4.28 (1H, m), 4.16-4.10 (1H, m), 3.48-3.43 (1H, m), 3.02-2.93 (1H, m), 2.68-2.65 (2H, m), 2.06-1.32 (8H, m), 1.04 (6H, s), 1.00 (6H, d, J = 5.8 Hz).
114					free	487.2	¹ H-NMR (DMSO-D ₆) δ: 7.82-7.63 (6H, m), 7.49-7.40 (1H, m), 7.13 (1H, d, J = 7.5 Hz), 4.27 (3H, s), 3.77-3.57 (5H, m), 2.50-2.49 (2H, m), 1.15 (2H, s), 1.08 (3H, t, J = 6.9 Hz).
115					free	501.3	¹ H-NMR (DMSO-D ₆) δ: 8.16-8.14 (1H, m), 7.79-7.72 (2H, m), 7.70 (1H, d, J = 9.2 Hz), 7.66 (1H, s), 7.62 (1H, d, J = 7.6 Hz), 7.47-7.43 (1H, m), 7.14-7.12 (1H, m), 4.28 (3H, s), 3.74-3.52 (6H, m), 2.13-2.03 (1H, m), 1.81-1.72 (1H, m), 1.07-0.96 (6H, m).






















[0802]

Table 15

115					free	513.2	¹ H-NMR (DMSO-D ₆) δ: 8.14-8.13 (1H, m), 7.78-7.61 (5H, m), 7.45 (1H, dd, J = 9.9, 4.7 Hz), 7.13 (1H, dd, J = 7.9, 1.5 Hz), 4.27 (3H, s), 3.68-3.48 (6H, m), 2.15-1.95 (4H, m), 1.79-1.58 (4H, m).
117					free	441.2	¹ H-NMR (DMSO-D ₆) δ: 7.84 (1H, t, J = 7.5 Hz), 7.65-7.69 (1H, m), 7.55 (1H, s), 7.50 (1H, d, J = 8.2 Hz), 7.39-7.34 (1H, m), 7.12 (1H, d, J = 7.3 Hz), 6.88 (1H, s), 6.69 (1H, d, J = 8.2 Hz), 6.38 (1H, d, J = 7.9 Hz), 3.93-3.54 (6H, m), 3.20-3.15 (1H, m), 2.82 (2H, t, J = 8.1 Hz), 2.69 (3H, s), 2.30-2.18 (1H, m), 2.06-1.92 (1H, m).
118					free	457.2	¹ H-NMR (DMSO-D ₆) δ: 7.86 (1H, t, J = 7.5 Hz), 7.64-7.60 (1H, m), 7.54 (1H, s), 7.50 (1H, d, J = 7.9 Hz), 7.42-7.35 (1H, m), 7.17-7.11 (1H, m), 6.58 (1H, d, J = 8.5 Hz), 6.52 (1H, d, J = 2.1 Hz), 6.47 (1H, d, J = 9.8 Hz), 4.22-4.17 (2H, m), 3.94-3.53 (6H, m), 3.30-3.21 (3H, m), 2.81 (3H, s), 2.29-2.15 (1H, m), 2.02-1.89 (1H, m).
119					free	457.2	¹ H-NMR (DMSO-D ₆) δ: 7.82 (1H, t, J = 7.5 Hz), 7.71-7.67 (1H, m), 7.62-7.57 (2H, m), 7.41 (1H, t, J = 10.7 Hz), 7.22 (1H, s), 7.18 (1H, d, J = 8.2 Hz), 7.13-7.08 (1H, m), 6.94 (1H, d, J = 9.5 Hz), 3.96-3.56 (4H, m), 3.30-3.26 (3H, m), 3.24-3.17 (1H, m), 2.31-2.18 (1H, m), 2.07-1.95 (1H, m).
120					free	473.2	¹ H-NMR (DMSO-D ₆) δ: 7.82 (1H, t, J = 7.5 Hz), 7.69 (1H, t, J = 6.1 Hz), 7.62-7.56 (3H, m), 7.44 (1H, t, J = 9.2 Hz), 7.22 (1H, d, J = 8.2 Hz), 7.11-7.07 (1H, m), 7.07-7.04 (1H, m), 3.84-3.52 (4H, m), 3.31-3.28 (3H, m), 3.28-3.23 (1H, m), 2.21-2.12 (1H, m), 1.94-1.85 (1H, m).
121					free	444.1	¹ H-NMR (DMSO-D ₆) δ: 7.85 (1H, t, J = 7.5 Hz), 7.67-7.63 (1H, m), 7.57 (1H, s), 7.53 (1H, d, J = 7.6 Hz), 7.42-7.35 (1H, m), 7.14-7.11 (1H, m), 6.76 (1H, d, J = 8.5 Hz), 6.72 (1H, d, J = 2.1 Hz), 6.50 (1H, d, J = 7.3 Hz), 4.25-4.20 (4H, m), 3.98-3.56 (4H, m), 3.30-3.25 (1H, m), 2.33-2.15 (1H, m), 2.07-1.95 (1H, m).
122					free	430.2	¹ H-NMR (DMSO-D ₆) δ: 7.83 (1H, t, J = 7.5 Hz), 7.68-7.62 (1H, m), 7.59-7.55 (1H, m), 7.53 (1H, d, J = 7.9 Hz), 7.40-7.34 (1H, m), 7.14-7.10 (1H, m), 6.84 (1H, d, J = 7.9 Hz), 6.74 (1H, d, J = 1.5 Hz), 6.56 (1H, d, J = 7.9 Hz), 6.01 (2H, s), 4.23-3.74 (4H, m), 3.36-3.28 (1H, m), 2.30-2.14 (1H, m), 2.11-1.95 (1H, m).
123					free	497.2	¹ H-NMR (DMSO-D ₆) δ: 7.74 (1H, t, J = 7.5 Hz), 7.60 (1H, dd, J = 7.9, 1.6 Hz), 7.55-7.49 (3H, m), 7.37-7.33 (2H, m), 7.21 (1H, d, J = 11.0 Hz), 7.10 (1H, dd, J = 8.2, 1.5 Hz), 6.45 (1H, d, J = 3.1 Hz), 4.59 (1H, s), 4.09 (1H, s), 3.75-3.69 (3H, m), 2.22-1.50 (9H, m).





















[0803]

Table 16

124					free	498.2	¹ H-NMR (DMSO-D ₆) δ: 8.09 (1H, d, J = 0.9 Hz), 7.79 (1H, d, J = 7.3 Hz), 7.75 (1H, t, J = 7.5 Hz), 7.63 (1H, dd, J = 7.9, 1.8 Hz), 7.56-7.54 (2H, m), 7.44 (1H, d, J = 10.7 Hz), 7.39 (1H, dd, J = 10.5, 1.4 Hz), 7.10 (1H, dd, J = 7.9, 1.5 Hz), 4.61 (1H, s), 4.10 (1H, s), 3.98 (3H, s), 2.61-2.57 (2H, m), 2.47-2.42 (1H, m), 2.27-1.52 (6H, m).
125					free	499.2	¹ H-NMR (DMSO-D ₆) δ: 8.12 (1H, d, J = 6.4 Hz), 7.74 (1H, t, J = 7.5 Hz), 7.69-7.65 (1H, m), 7.62 (1H, d, J = 7.9 Hz), 7.58 (1H, d, J = 1.5 Hz), 7.44 (2H, dd, J = 10.5, 1.4 Hz), 7.13 (1H, dd, J = 7.9, 1.5 Hz), 4.64 (1H, s), 4.25 (3H, s), 4.11 (1H, s), 2.34-2.10 (3H, m), 2.07-1.94 (3H, m), 1.93-1.52 (3H, m).
126					free	516.2	¹ H-NMR (DMSO-D ₆) δ: 8.25 (1H, s), 7.76 (1H, t, J = 7.5 Hz), 7.66-7.63 (1H, m), 7.60-7.56 (2H, m), 7.45-7.41 (2H, m), 7.12 (1H, dd, J = 7.9, 1.5 Hz), 4.61 (1H, s), 4.10 (1H, s), 3.96 (3H, s), 2.30-2.17 (2H, m), 2.14-2.04 (1H, m), 2.01-1.51 (6H, m).
127					free	516.2	¹ H-NMR (DMSO-D ₆) δ: 7.83 (1H, t, J = 7.5 Hz), 7.56 (2H, d, J = 0.9 Hz), 7.46 (1H, d, J = 0.9 Hz), 7.34 (1H, dd, J = 10.5, 1.4 Hz), 7.25 (1H, t, J = 7.9 Hz), 7.13-7.10 (2H, m), 7.04-6.99 (1H, m), 4.78-4.72 (1H, m), 3.97-3.92 (1H, m), 3.54 (2H, t, J = 8.5 Hz), 3.29-3.28 (1H, m), 3.23 (3H, s), 2.82 (2H, t, J = 6.7 Hz), 2.07-1.56 (10H, m).
128					free	497.2	¹ H-NMR (DMSO-D ₆) δ: 7.84 (1H, t, J = 7.5 Hz), 7.54 (1H, dd, J = 7.9, 1.5 Hz), 7.44 (3H, dt, J = 17.3, 7.9 Hz), 7.16 (1H, dd, J = 8.1, 1.4 Hz), 6.59 (1H, d, J = 8.2 Hz), 6.53 (1H, d, J = 2.1 Hz), 6.48 (1H, dd, J = 8.2, 2.1 Hz), 4.58 (1H, s), 4.20 (2H, t, J = 4.4 Hz), 4.10 (1H, s), 3.24 (2H, t, J = 4.4 Hz), 2.82 (3H, s), 2.40-1.90 (7H, m), 1.60-1.52 (2H, m).
129					free	516.2	¹ H-NMR (DMSO-D ₆) δ: 7.82 (1H, t, J = 7.5 Hz), 7.54 (2H, s), 7.48 (1H, s), 7.34 (1H, d, J = 10.7 Hz), 7.25 (1H, t, J = 7.9 Hz), 7.16-7.08 (2H, m), 7.01 (1H, d, J = 11.0 Hz), 4.91-4.84 (1H, m), 4.07-4.01 (1H, m), 3.54 (2H, t, J = 6.6 Hz), 3.24 (3H, s), 2.93-2.86 (1H, m), 2.82 (2H, t, J = 6.6 Hz), 2.31-1.96 (3H, m), 1.67-1.50 (4H, m), 1.43-1.24 (3H, m).
130					free	498.1	¹ H-NMR (DMSO-D ₆) δ: 8.28 (1H, s), 7.77-7.73 (1H, m), 7.64 (2H, dd, J = 9.6, 7.9 Hz), 7.58 (2H, d, J = 7.9 Hz), 7.42 (2H, d, J = 9.8 Hz), 7.11 (1H, d, J = 7.9 Hz), 4.69 (1H, s), 4.18 (1H, s), 3.80 (3H, s), 2.45-2.28 (3H, m), 2.13-2.02 (2H, m), 1.88-1.80 (2H, m), 1.73-1.59 (2H, m).












[0804]

Table 17

131					free	470.2	¹ H-NMR (DMSO-D ₆) δ: 7.91 (1H, d, J = 2.4 Hz), 7.87 (1H, t, J = 7.5 Hz), 7.79-7.74 (1H, m), 7.60 (1H, d, J = 7.9 Hz), 7.55-7.46 (2H, m), 7.22 (1H, d, J = 7.9 Hz), 7.17 (1H, d, J = 7.9 Hz), 6.60-6.56 (1H, m), 4.66 (1H, s), 4.17 (1H, s), 3.02 (6H, s), 2.40-2.25 (2H, m), 2.10-2.01 (2H, m), 1.64-1.78 (2H, m), 1.70-1.59 (2H, m).
132					free	523.2	¹ H-NMR (DMSO-D ₆) δ: 7.80 (1H, t, J = 7.5 Hz), 7.62-7.55 (2H, m), 7.51 (1H, d, J = 1.5 Hz), 7.36 (1H, d, J = 10.4 Hz), 7.22 (1H, dd, J = 7.9, 1.8 Hz), 7.09 (1H, dd, J = 8.1, 1.4 Hz), 7.02 (1H, d, J = 8.2 Hz), 6.90 (1H, d, J = 1.8 Hz), 4.58 (1H, s), 4.07 (1H, s), 3.13 (3H, s), 2.25-1.84 (7H, m), 1.63-1.48 (2H, m), 1.07 (6H, s).
133					free	513.1	¹ H-NMR (DMSO-D ₆) δ: 7.81 (1H, t, J = 7.6 Hz), 7.62-7.58 (2H, m), 7.55 (1H, d, J = 7.9 Hz), 7.51 (1H, d, J = 1.8 Hz), 7.46 (1H, dd, J = 10.5, 1.4 Hz), 7.22 (1H, d, J = 8.2 Hz), 7.10 (1H, dd, J = 8.2, 1.5 Hz), 7.06 (1H, dd, J = 8.2, 1.8 Hz), 4.58 (1H, s), 4.07 (1H, s), 3.27-3.24 (3H, m), 2.27-1.84 (7H, m), 1.63-1.48 (2H, m).
134					free	457.1	¹ H-NMR (DMSO-D ₆) δ: 7.79-7.73 (1H, m), 7.72-7.66 (1H, m), 7.65-7.59 (1H, m), 7.58-7.49 (2H, m), 7.39-7.32 (2H, m), 7.22 (1H, d, J = 11.0 Hz), 7.14-7.09 (1H, m), 6.45 (1H, d, J = 3.1 Hz), 3.74 (3H, s), 3.69-3.60 (5H, m), 2.04-1.94 (1H, m), 1.71-1.65 (1H, m).
135					free	458.1	¹ H-NMR (DMSO-D ₆) δ: 8.10 (1H, s), 7.81-7.75 (2H, m), 7.71 (1H, t, J = 7.0 Hz), 7.66 (1H, d, J = 5.2 Hz), 7.59 (1H, d, J = 7.9 Hz), 7.46 (1H, d, J = 10.4 Hz), 7.42-7.38 (1H, m), 7.10 (1H, d, J = 8.2 Hz), 4.00 (3H, s), 3.79-3.66 (5H, m), 2.16-2.09 (1H, m), 1.88-1.80 (1H, m).
136					free	555.1	¹ H-NMR (DMSO-D ₆) δ: 7.74 (1H, t, J = 7.6 Hz), 7.61 (1H, dd, J = 7.9, 1.8 Hz), 7.58-7.53 (2H, m), 7.48 (1H, d, J = 7.6 Hz), 7.33 (3H, dd, J = 21.4, 7.0 Hz), 7.13 (1H, dd, J = 7.9, 1.5 Hz), 6.45 (1H, d, J = 3.1 Hz), 4.64 (1H, s), 4.62-4.57 (1H, m), 4.13-4.08 (1H, m), 4.02 (2H, s), 2.25-2.14 (2H, m), 2.03-1.86 (3H, m), 1.80-1.68 (1H, m), 1.63-1.51 (2H, m), 1.46-1.33 (1H, m), 1.06 (6H, s).
137					free	468.1	¹ H-NMR (DMSO-D ₆) δ: 7.81 (1H, t, J = 7.5 Hz), 7.61 (1H, d, J = 9.8 Hz), 7.56-7.50 (2H, m), 7.42 (1H, d, J = 10.7 Hz), 7.22 (1H, d, J = 7.9 Hz), 7.18 (1H, s), 7.11 (1H, d, J = 8.2 Hz), 6.98 (1H, d, J = 7.9 Hz), 4.99 (2H, s), 4.95 (2H, s), 4.62-4.56 (1H, m), 4.12-4.07 (1H, m), 2.22-2.10 (2H, m), 2.05-1.86 (3H, m), 1.82-1.69 (1H, m), 1.63-1.50 (2H, m), 1.45-1.24 (1H, m).













[0805]

Table 18

138					free	541.1	¹ H-NMR (DMSO-D ₆) δ: 7.82 (1H, t, J = 7.6 Hz), 7.62 (1H, d, J = 7.9 Hz), 7.58-7.54 (2H, m), 7.52 (1H, s), 7.48 (1H, d, J = 10.7 Hz), 7.40 (1H, d, J = 8.5 Hz), 7.12 (1H, d, J = 7.9 Hz), 7.04 (1H, d, J = 8.5 Hz), 4.78-4.71 (1H, m), 4.65-4.58 (1H, m), 4.14-4.08 (1H, m), 2.28-2.19 (1H, m), 2.15-1.90 (3H, m), 1.87-1.70 (2H, m), 1.65-1.52 (2H, m), 1.47 (6H, d, J = 7.0 Hz).
139					free	541.2	¹ H-NMR (DMSO-D ₆) δ: 8.13 (1H, d, J = 7.0 Hz), 7.93 (1H, d, J = 10.4 Hz), 7.77 (1H, t, J = 7.6 Hz), 7.67-7.62 (2H, m), 7.59 (1H, s), 7.48 (1H, d, J = 10.7 Hz), 7.15 (1H, dd, J = 8.1, 1.4 Hz), 4.66-4.59 (1H, m), 4.14-4.08 (1H, m), 2.24-2.05 (4H, m), 2.01-1.86 (3H, m), 1.77 (9H, s), 1.66-1.53 (2H, m).
140					free	428.1	¹ H-NMR (DMSO-D ₆) δ: 7.82 (1H, t, J = 7.6 Hz), 7.69 (1H, t, J = 7.6 Hz), 7.61 (1H, d, J = 6.4 Hz), 7.55 (1H, d, J = 7.9 Hz), 7.45-7.38 (1H, m), 7.23 (1H, d, J = 7.9 Hz), 7.18 (1H, s), 7.10 (1H, d, J = 7.6 Hz), 6.98 (1H, d, J = 6.4 Hz), 4.99 (2H, s), 4.95 (2H, s), 3.82-3.67 (5H, m), 2.13-2.06 (1H, m), 1.85-1.78 (1H, m).
141					free	491	¹ H-NMR (DMSO-D ₆) δ: 7.82 (1H, t, J = 7.3 Hz), 7.75 (1H, d, J = 7.0 Hz), 7.72-7.69 (1H, m), 7.64 (1H, d, J = 4.7 Hz), 7.54 (1H, d, J = 7.9 Hz), 7.45 (1H, d, J = 10.4 Hz), 7.24 (1H, d, J = 10.7 Hz), 7.12 (1H, dd, J = 8.2, 1.5 Hz), 3.67-3.48 (5H, m), 3.37 (3H, s), 2.02-1.91 (1H, m), 1.69-1.61 (1H, m).
142					free	556.1	¹ H-NMR (DMSO-D ₆) δ: 8.10 (1H, d, J = 0.9 Hz), 7.79-7.74 (2H, m), 7.63 (1H, dd, J = 7.8, 1.7 Hz), 7.59-7.55 (2H, m), 7.45 (1H, d, J = 11.0 Hz), 7.39 (1H, dd, J = 10.7, 1.5 Hz), 7.13 (1H, dd, J = 8.2, 1.5 Hz), 4.62 (1H, s), 4.60-4.56 (1H, m), 4.26 (2H, s), 4.08-4.05 (1H, m), 2.35-2.21 (2H, m), 2.12-2.06 (1H, m), 1.99-1.83 (3H, m), 1.64-1.48 (3H, m), 1.11 (6H, s).
143		 Racemate Relative configuration			free	448.1	¹ H-NMR (DMSO-D ₆) δ: 7.82 (1H, t, J = 7.5 Hz), 7.73-7.58 (2H, m), 7.53 (1H, d, J = 7.9 Hz), 7.39-7.33 (1H, m), 7.25 (1H, t, J = 7.9 Hz), 7.16-7.10 (2H, m), 7.01 (1H, d, J = 11.3 Hz), 4.49-3.86 (3H, m), 3.54 (2H, t, J = 6.6 Hz), 3.23 (3H, s), 2.82 (2H, t, J = 6.6 Hz), 2.22-2.10 (2H, m), 1.89-1.70 (1H, m), 1.64-1.48 (2H, m), 0.91 (1H, dd, J = 12.2, 4.3 Hz).
144					free	476.0	¹ H-NMR (DMSO-D ₆) δ: 7.83 (1H, dd, J = 7.9, 7.0 Hz), 7.63 (1H, dd, J = 7.8, 1.7 Hz), 7.57-7.53 (2H, m), 7.43 (1H, dd, J = 10.5, 1.4 Hz), 7.13-7.10 (2H, m), 7.03-6.99 (1H, m), 4.59 (1H, s), 4.05 (1H, s), 2.27 (3H, d, J = 1.2 Hz), 2.23-2.09 (3H, m), 2.04-1.85 (3H, m), 1.63-1.49 (3H, m).

[0806]

Table 19

145					free	517.0	¹ H-NMR (DMSO-D ₆) δ: 7.97 (1H, d, J = 4.6 Hz), 7.78 (1H, dd, J = 7.9, 7.0 Hz), 7.57 (1H, dd, J = 7.9, 1.8 Hz), 7.53 (1H, d, J = 7.6 Hz), 7.59 (1H, d, J = 1.5 Hz), 7.50 (2H, dd, J = 10.5, 1.4 Hz), 7.16 (1H, dd, J = 7.9, 1.5 Hz), 4.61 (1H, s), 4.40 (2H, s), 4.06 (1H, s), 2.24-1.86 (6H, m), 1.65-1.51 (2H, m).
146					free	515.0	¹ H-NMR (DMSO-D ₆) δ: 7.74 (1H, t, J = 7.5 Hz), 7.68 (1H, t, J = 7.2 Hz), 7.62 (1H, d, J = 8.5 Hz), 7.55 (1H, d, J = 7.9 Hz), 7.47 (1H, d, J = 7.6 Hz), 7.35 (1H, d, J = 3.1 Hz), 7.30 (2H, d, J = 11.9 Hz), 7.11 (1H, t, J = 4.7 Hz), 6.44 (1H, d, J = 3.1 Hz), 4.62 (1H, s), 4.01 (2H, s), 3.78-3.58 (4H, m), 3.13-2.96 (1H, m), 2.03-1.93 (1H, m), 1.72-1.61 (1H, m), 1.05 (5H, s).
147					free	516.0	¹ H-NMR (DMSO-D ₆) δ: 8.10 (1H, d, J = 0.9 Hz), 7.77-7.74 (2H, m), 7.71-7.69 (1H, m), 7.64 (1H, dd, J = 7.8, 1.7 Hz), 7.57-7.57 (1H, m), 7.44 (1H, d, J = 11.0 Hz), 7.37 (1H, d, J = 10.7 Hz), 7.11 (1H, dd, J = 8.1, 1.7 Hz), 4.61 (1H, s), 4.25 (2H, s), 3.69-3.57 (2H, m), 3.56-3.43 (2H, m), 3.24-3.14 (1H, m), 2.02-1.90 (1H, m), 1.68-1.61 (1H, m), 1.10 (6H, s).
148					free	478.1	¹ H-NMR (DMSO-D ₆) δ: 8.97 (2H, s), 8.03-7.98 (2H, m), 7.80-7.74 (2H, m), 7.70 (1H, dd, J = 7.9, 1.5 Hz), 7.62 (1H, d, J = 1.5 Hz), 7.57-7.52 (2H, m), 7.15 (1H, dd, J = 8.1, 1.7 Hz), 4.69-4.62 (1H, m), 4.20-4.13 (1H, m), 2.33-1.91 (7H, m), 1.74-1.53 (2H, m).
149					free	477.0	¹ H-NMR (DMSO-D ₆) δ: 9.40-9.31 (1H, m), 8.60-8.53 (1H, m), 8.07-7.95 (2H, m), 7.78-7.68 (4H, m), 7.62 (1H, s), 7.51 (1H, d, J = 9.5 Hz), 7.29-7.30 (1H, m), 7.09 (1H, d, J = 7.6 Hz), 4.76-4.63 (1H, m), 4.24-4.15 (1H, m), 2.36-1.81 (7H, m), 1.78-1.58 (2H, m).
150					free	477.0	¹ H-NMR (DMSO-D ₆) δ: 9.37-9.31 (1H, m), 8.56-8.52 (1H, m), 8.13-8.08 (1H, m), 7.92-7.85 (2H, m), 7.76-7.70 (3H, m), 7.62 (1H, s), 7.50 (1H, d, J = 10.4 Hz), 7.46-7.40 (1H, m), 7.10 (1H, d, J = 7.9 Hz), 4.73-4.66 (1H, m), 4.24-4.17 (1H, m), 2.34-2.00 (5H, m), 1.92-1.57 (4H, m).
151					free	477.0	¹ H-NMR (DMSO-D ₆) δ: 8.93 (1H, d, J = 4.3 Hz), 8.36 (1H, d, J = 8.5 Hz), 7.94 (1H, s), 7.90 (1H, d, J = 8.5 Hz), 7.76-7.69 (3H, m), 7.61 (1H, d, J = 1.5 Hz), 7.60-7.55 (1H, m), 7.51 (1H, d, J = 10.4 Hz), 7.40 (1H, d, J = 9.2 Hz), 7.10 (1H, dd, J = 7.9, 1.5 Hz), 4.74-4.65 (1H, m), 4.25-4.16 (1H, m), 2.31-1.58 (9H, m).
152					free	478.0	¹ H-NMR (DMSO-D ₆) δ: 9.60 (1H, s), 8.06 (1H, d, J = 8.5 Hz), 7.93 (1H, s), 7.75-7.74 (1H, m), 7.73-7.69 (1H, m), 7.63 (1H, s), 7.58-7.52 (2H, m), 7.45-7.41 (1H, m), 7.15-7.12 (2H, m), 4.72-4.63 (1H, m), 4.22-4.15 (1H, m), 2.13-2.04 (2H, m), 1.88-1.60 (7H, m).







[0807]

Table 20

153					free	478.0	¹ H-NMR (DMSO-D ₆) δ: 9.32 (1H, s), 8.12 (1H, d, J = 2.7 Hz), 7.90 (1H, d, J = 8.5 Hz), 7.75-7.71 (3H, m), 7.69-7.62 (3H, m), 7.53 (1H, d, J = 10.1 Hz), 7.11-7.09 (1H, m), 4.73-4.65 (1H, m), 4.24-4.15 (1H, m), 2.14-2.01 (3H, m), 1.87-1.65 (6H, m).
154					free	478.0	¹ H-NMR (DMSO-D ₆) δ: 9.66 (2H, d, J = 3.4 Hz), 8.20 (1H, s), 8.12 (1H, s), 8.03 (1H, d, J = 8.5 Hz), 7.75-7.66 (4H, m), 7.62 (1H, s), 7.51 (1H, d, J = 9.2 Hz), 7.07 (1H, dd, J = 7.9, 1.5 Hz), 4.66-4.58 (1H, m), 4.15-4.07 (1H, m), 2.31-1.88 (7H, m), 1.67-1.50 (2H, m).
155		 isomer-B Relative configuration			free	488.0	¹ H-NMR (DMSO-D ₆) δ: 7.81 (1H, t, J = 7.5 Hz), 7.73-7.56 (2H, m), 7.52 (1H, d, J = 7.9 Hz), 7.35 (1H, d, J = 10.4 Hz), 7.24 (1H, t, J = 7.9 Hz), 7.14-7.09 (2H, m), 7.01 (1H, d, J = 11.3 Hz), 4.42-3.89 (3H, m), 3.53 (2H, t, J = 6.6 Hz), 3.23 (3H, s), 2.81 (2H, t, J = 6.6 Hz), 2.25-2.08 (2H, m), 1.79-1.68 (1H, m), 1.64-1.44 (2H, m), 0.91-0.87 (1H, m).
156		 isomer-A Relative configuration			free	486.0	¹ H-NMR (DMSO-D ₆) δ: 7.81 (1H, t, J = 7.5 Hz), 7.73-7.56 (2H, m), 7.52 (1H, d, J = 7.5 Hz), 7.35 (1H, d, J = 10.7 Hz), 7.24 (1H, t, J = 7.9 Hz), 7.15-7.09 (2H, m), 7.01 (1H, d, J = 11.0 Hz), 4.46-3.85 (3H, m), 3.53 (2H, t, J = 6.7 Hz), 3.23 (3H, s), 2.81 (2H, t, J = 6.6 Hz), 2.21-2.10 (2H, m), 1.79-1.68 (1H, m), 1.64-1.46 (2H, m), 0.93-0.87 (1H, m).
157					free	480.0	¹ H-NMR (DMSO-D ₆) δ: 7.87 (1H, t, J = 7.5 Hz), 7.78 (3H, s), 7.65 (2H, s), 7.59-7.51 (3H, m), 7.24 (1H, d, J = 7.0 Hz), 4.67 (1H, s), 4.16 (1H, s), 2.52 (3H, s), 2.34-2.22 (2H, m), 2.14-2.02 (1H, m), 1.89-1.80 (3H, m), 1.75-1.57 (3H, m).
158					free	481.1	¹ H-NMR (DMSO-D ₆) δ: 9.04 (1H, d, J = 2.1 Hz), 8.11 (1H, d, J = 0.6 Hz), 8.02 (1H, d, J = 2.1 Hz), 7.85 (1H, t, J = 7.5 Hz), 7.75 (1H, d, J = 7.9 Hz), 7.67 (1H, dd, J = 7.8, 1.7 Hz), 7.62 (1H, dd, J = 10.4, 1.2 Hz), 7.59 (1H, d, J = 1.5 Hz), 7.28 (1H, dd, J = 7.9, 1.5 Hz), 4.52 (1H, s), 4.08 (1H, s), 2.26 (3H, s), 2.15-1.91 (6H, m), 1.68-1.51 (3H, m).
159					free	556.0	¹ H-NMR (DMSO-D ₆) δ: 8.37 (1H, d, J = 0.9 Hz), 7.81-7.75 (2H, m), 7.62 (1H, dd, J = 7.9, 1.8 Hz), 7.58-7.54 (2H, m), 7.41 (1H, dd, J = 10.7, 1.5 Hz), 7.26 (1H, d, J = 11.6 Hz), 7.14 (1H, dd, J = 8.2, 1.5 Hz), 4.87 (1H, s), 4.59 (1H, s), 4.32 (2H, s), 4.08 (1H, s), 2.28-2.10 (3H, m), 2.02-1.83 (3H, m), 1.63-1.49 (3H, m), 1.11 (6H, s).
160					free	513.0	¹ H-NMR (DMSO-D ₆) δ: 8.14 (1H, d, J = 6.4 Hz), 7.77-7.74 (2H, m), 7.62 (3H, ddd, J = 29.6, 12.8, 3.2 Hz), 7.46 (1H, dd, J = 10.5, 1.4 Hz), 7.14 (1H, dd, J = 7.9, 1.5 Hz), 4.68 (2H, q, J = 7.3 Hz), 4.61 (1H, s), 4.09 (1H, s), 2.23-1.88 (6H, m), 1.65-1.52 (3H, m), 1.49 (3H, t, J = 7.2 Hz).








[0808]

Table 21

161					free	575.0	¹ H-NMR (DMSO-D ₆) δ: 7.97 (1H, d, J = 4.6 Hz), 7.76 (1H, t, J = 7.6 Hz), 7.67-7.64 (2H, m), 7.59 (1H, d, J = 0.9 Hz), 7.48 (1H, dd, J = 10.5, 1.4 Hz), 7.16 (1H, dd, J = 7.9, 1.5 Hz), 4.81 (1H, s), 4.61 (2H, s), 4.57 (1H, s), 4.05 (1H, s), 2.34-2.21 (2H, m), 2.12-2.05 (1H, m), 1.99-1.83 (3H, m), 1.62-1.46 (3H, m), 1.15 (6H, s).
162					free	535.0	¹ H-NMR (DMSO-D ₆) δ: 7.97 (1H, d, J = 5.2 Hz), 7.78-7.64 (4H, m), 7.46 (1H, d, J = 10.4 Hz), 7.15 (1H, dd, J = 8.2, 1.5 Hz), 4.81 (1H, s), 4.61 (2H, s), 3.69-3.57 (2H, m), 3.55-3.46 (2H, m), 3.23-3.15 (1H, m), 2.04-1.90 (1H, m), 1.58-1.56 (1H, m), 1.15 (6H, s).
163					free	462.0	¹ H-NMR (DMSO-D ₆) δ: 7.82 (1H, t, J = 7.5 Hz), 7.63-7.60 (1H, m), 7.57-7.53 (2H, m), 7.34 (1H, d, J = 10.4 Hz), 7.26-7.21 (1H, m), 7.13-7.09 (2H, m), 7.01 (1H, d, J = 11.3 Hz), 3.76-3.42 (6H, m), 3.23 (3H, s), 3.13-2.92 (2H, m), 2.81 (2H, t, J = 6.7 Hz).
164					free	475.1	¹ H-NMR (DMSO-D ₆) δ: 7.82 (1H, t, J = 7.5 Hz), 7.59 (1H, dd, J = 7.8, 1.7 Hz), 7.56-7.52 (2H, m), 7.32 (1H, dd, J = 10.5, 1.4 Hz), 7.24 (1H, t, J = 7.9 Hz), 7.11 (2H, dd, J = 7.9, 1.5 Hz), 7.01 (1H, d, J = 11.3 Hz), 4.34-3.71 (2H, m), 3.53 (2H, t, J = 6.6 Hz), 3.23 (3H, s), 3.12-2.94 (2H, m), 2.82 (2H, t, J = 6.7 Hz), 1.99-1.91 (1H, m), 1.81-1.64 (1H, m), 1.56-1.41 (2H, m), 0.88-0.80 (1H, m).
165					free	490.1	¹ H-NMR (DMSO-D ₆) δ: 7.81 (1H, t, J = 7.5 Hz), 7.57-7.48 (3H, m), 7.35-7.31 (1H, m), 7.24 (1H, t, J = 7.9 Hz), 7.11 (2H, dd, J = 8.1, 1.4 Hz), 7.01 (1H, d, J = 11.2 Hz), 3.92-3.62 (2H, m), 3.58-3.41 (4H, m), 3.23 (3H, s), 2.81 (2H, t, J = 6.7 Hz), 2.14-2.03 (1H, m), 1.91-1.49 (5H, m), 0.88-0.80 (1H, m).
166		 isomer-B Relative configuration			free	502.6	¹ H-NMR (DMSO-D ₆) δ: 7.80 (1H, t, J = 7.6 Hz), 7.68 (1H, s), 7.60 (1H, s), 7.53 (1H, d, J = 7.9 Hz), 7.29 (1H, t, J = 5.3 Hz), 7.21 (1H, t, J = 7.9 Hz), 7.15 (1H, dd, J = 8.2, 1.5 Hz), 7.07 (1H, d, J = 7.9 Hz), 6.95 (1H, d, J = 11.6 Hz), 4.46-3.86 (3H, m), 2.65 (2H, s), 2.25-2.10 (2H, m), 1.74-1.49 (4H, m), 1.04 (6H, s), 0.87 (1H, dd, J = 12.2, 4.6 Hz).
167					free	477.0	¹ H-NMR (DMSO-D ₆) δ: 7.99-7.96 (1H, m), 7.80-7.73 (2H, m), 7.69-7.67 (1H, m), 7.64 (1H, d, J = 7.9 Hz), 7.50-7.46 (1H, m), 7.18-7.14 (1H, m), 4.40 (3H, d, J = 6.7 Hz), 3.70-3.64 (2H, m), 3.58-3.48 (2H, m), 3.25-3.16 (1H, m), 2.08-2.04 (1H, m), 1.78-1.74 (1H, m).
168					free	527.0	¹ H-NMR (DMSO-D ₆) δ: 8.14 (1H, d, J = 6.7 Hz), 7.77-7.74 (2H, m), 7.68-7.59 (3H, m), 7.45 (1H, dd, J = 10.4, 1.5 Hz), 7.14 (1H, dd, J = 7.9, 1.5 Hz), 4.64-4.60 (3H, m), 4.14-4.07 (1H, m), 2.30-1.87 (9H, m), 1.66-1.55 (2H, m), 0.84 (3H, t, J = 7.3 Hz).













[0809]

Table 22

169					free	521.0	¹ H-NMR (DMSO-D ₆) δ: 7.99 (1H, d, J = 4.9 Hz), 7.79-7.73 (2H, m), 7.70-7.64 (2H, m), 7.48-7.45 (1H, m), 7.16 (1H, dd, J = 8.2, 1.5 Hz), 4.91 (2H, t, J = 5.2 Hz), 3.81 (2H, t, J = 5.0 Hz), 3.69-3.45 (4H, m), 3.18-3.16 (4H, m), 2.01-1.94 (1H, m), 1.68-1.63 (1H, m).
170					free	487.0	¹ H-NMR (DMSO-D ₆) δ: 7.75 (1H, dd, J = 7.8, 7.2 Hz), 7.69-7.66 (1H, m), 7.62 (1H, dd, J = 7.9, 1.5 Hz), 7.51 (2H, dd, J = 16.9, 7.6 Hz), 7.39-7.35 (2H, m), 7.25 (1H, d, J = 11.3 Hz), 7.11 (1H, dd, J = 7.9, 1.5 Hz), 6.44 (1H, dd, J = 3.1, 0.6 Hz), 4.86 (1H, t, J = 5.2 Hz), 4.15 (2H, t, J = 5.3 Hz), 3.69-3.59 (4H, m), 3.54-3.47 (2H, m), 3.22-3.18 (1H, m), 1.99-1.94 (1H, m), 1.73-1.66 (1H, m).
171		 isomer-B Relative configuration			free	541.0	¹ H-NMR (DMSO-D ₆) δ: 7.74 (1H, t, J = 7.5 Hz), 7.68 (1H, d, J = 7.9 Hz), 7.61-7.55 (2H, m), 7.47 (1H, d, J = 7.6 Hz), 7.36-7.29 (3H, m), 7.13 (1H, dd, J = 7.9, 1.5 Hz), 6.44 (1H, d, J = 3.4 Hz), 4.62 (1H, s), 4.48-3.87 (5H, m), 2.19-2.14 (2H, m), 1.78-1.73 (1H, m), 1.62-1.49 (2H, m), 1.05 (6H, s), 0.88 (1H, dd, J = 12.1, 4.7 Hz).
172					free	533.0	¹ H-NMR (DMSO-D ₆) δ: 7.76 (1H, t, J = 7.6 Hz), 7.71-7.68 (1H, m), 7.65-7.64 (1H, m), 7.59 (1H, d, J = 7.9 Hz), 7.38 (1H, d, J = 3.4 Hz), 7.35-7.30 (2H, m), 7.12 (1H, dd, J = 7.9, 1.5 Hz), 6.51 (1H, t, J = 2.4 Hz), 4.70 (1H, s), 4.13 (2H, s), 3.70-3.55 (4H, m), 3.24-3.18 (1H, m), 2.01-1.99 (1H, m), 1.71-1.67 (1H, m), 1.05 (6H, s).
173					free	541.0	¹ H-NMR (DMSO-D ₆) δ: 7.76-7.53 (5H, m), 7.50-7.44 (1H, m), 7.35-7.28 (2H, m), 7.15-7.08 (1H, m), 6.44 (1H, t, J = 2.7 Hz), 4.62 (1H, s), 4.30-3.84 (3H, m), 3.78-3.59 (1H, m), 3.20-3.15 (1H, m), 3.09-3.04 (1H, m), 2.25-1.50 (5H, m), 1.05 (6H, dd, J = 12.4, 4.7 Hz).
174					free	502.1	¹ H-NMR (DMSO-D ₆) δ: 7.93-7.77 (1H, m), 7.67-7.49 (3H, m), 7.28-7.12 (3H, m), 7.09-7.05 (1H, m), 6.98-6.93 (1H, m), 4.40 (1H, d, J = 3.1 Hz), 4.11-3.72 (1H, m), 3.61-3.48 (1H, m), 3.22-3.16 (1H, m), 3.10-3.03 (1H, m), 2.66 (2H, s), 2.21-1.55 (5H, m), 1.04 (6H, s).
175		 isomer-B Relative configuration			free	503.0	¹ H-NMR (DMSO-D ₆) δ: 7.98 (1H, d, J = 5.2 Hz), 7.78 (1H, t, J = 7.5 Hz), 7.75-7.70 (1H, m), 7.63 (2H, d, J = 7.6 Hz), 7.55-7.48 (1H, m), 7.17 (1H, dd, J = 8.2, 1.5 Hz), 4.48-3.88 (6H, m), 2.24-2.12 (2H, m), 1.75-1.51 (3H, m), 0.89 (1H, dd, J = 12.1, 4.7 Hz).
176		 isomer-B Relative configuration			free	542.0	¹ H-NMR (DMSO-D ₆) δ: 8.10 (1H, d, J = 0.9 Hz), 7.80-7.74 (2H, m), 7.73-7.67 (1H, m), 7.64-7.56 (2H, m), 7.45 (1H, d, J = 11.0 Hz), 7.40 (1H, dd, J = 10.5, 1.4 Hz), 7.13 (1H, dd, J = 7.9, 1.5 Hz), 4.61 (1H, s), 4.49-3.89 (5H, m), 2.26-2.13 (2H, m), 1.81-1.50 (4H, m), 1.19 (6H, s).

[0810]

Table 23

177		 isomer-B Relative configuration			free	551.0	¹ H-NMR (DMSO-D ₆) δ: 7.97 (1H, d, J = 5.2 Hz), 7.78-7.73 (2H, m), 7.66 (2H, d, J = 7.6 Hz), 7.49 (1H, dd, J = 10.4, 1.2 Hz), 7.17 (1H, dd, J = 8.2, 1.5 Hz), 4.81 (1H, s), 4.61 (2H, s), 4.49-3.87 (3H, m), 2.28-2.11 (2H, m), 1.80-1.51 (3H, m), 1.15 (6H, s), 0.89 (1H, dd, J = 11.7, 4.7 Hz).
178					free	635.9	¹ H-NMR (DMSO-D ₆) δ: 7.77 (1H, t, J = 7.5 Hz), 7.64-7.56 (4H, m), 7.51 (1H, d, J = 10.7 Hz), 7.44 (1H, d, J = 11.6 Hz), 7.14 (1H, d, J = 9.2 Hz), 4.64 (1H, s), 4.58 (1H, s), 4.25 (2H, s), 4.06 (1H, s), 2.34-1.85 (6H, m), 1.64-1.44 (2H, m), 1.16-1.15 (1H, m), 1.13-1.08 (6H, m).
179					free	499.0	¹ H-NMR (DMSO-D ₆) δ: 7.84 (1H, d, J = 5.2 Hz), 7.78 (1H, t, J = 7.5 Hz), 7.69-7.64 (2H, m), 7.63-7.58 (2H, m), 7.47 (1H, d, J = 10.4 Hz), 7.13 (1H, dd, J = 8.2, 1.5 Hz), 4.63-4.57 (1H, m), 4.11-4.02 (1H, m), 2.53 (3H, s), 2.24-1.90 (7H, m), 1.64-1.48 (2H, m).
180					free	459.0	¹ H-NMR (DMSO-D ₆) δ: 7.83 (1H, d, J = 5.5 Hz), 7.79 (1H, t, J = 7.6 Hz), 7.74 (1H, t, J = 6.7 Hz), 7.70-7.62 (3H, m), 7.45 (1H, t, J = 9.0 Hz), 7.12 (1H, d, J = 8.9 Hz), 3.82-3.49 (5H, m), 2.53 (3H, s), 2.20-2.12 (1H, m), 1.94-1.84 (1H, m).
181					free	570.0	¹ H-NMR (DMSO-D ₆) δ: 7.80-7.73 (2H, m), 7.66-7.52 (3H, m), 7.39 (1H, d, J = 10.7 Hz), 7.32 (1H, d, J = 11.3 Hz), 7.12 (1H, dd, J = 8.2, 1.5 Hz), 4.62-4.55 (2H, m), 4.18-4.11 (2H, m), 4.10-4.03 (1H, m), 2.47 (3H, s), 2.35-1.84 (7H, m), 1.64-1.49 (2H, m), 1.10 (6H, s).
182		 Racemate Relative configuration			free	502.0	¹ H-NMR (DMSO-D ₆) δ: 7.83-7.64 (3H, m), 7.51 (1H, d, J = 7.3 Hz), 7.30-7.25 (1H, m), 7.21 (1H, t, J = 7.9 Hz), 7.16 (1H, dd, J = 8.2, 1.5 Hz), 7.07 (1H, dd, J = 7.9, 1.2 Hz), 6.95 (1H, d, J = 11.6 Hz), 4.51-3.85 (3H, m), 3.08-2.95 (1H, m), 2.65 (2H, s), 1.82-1.68 (2H, m), 1.36-1.23 (4H, m), 1.04 (6H, s).
183					free	573.1	¹ H-NMR (DMSO-D ₆) δ: 7.76 (1H, t, J = 7.6 Hz), 7.62 (1H, dd, J = 7.9, 1.5 Hz), 7.59 (1H, d, J = 7.9 Hz), 7.56 (1H, d, J = 1.5 Hz), 7.38 (1H, d, J = 3.1 Hz), 7.34 (1H, dd, J = 10.5, 1.4 Hz), 7.31 (1H, d, J = 5.8 Hz), 7.13 (1H, dd, J = 7.9, 1.5 Hz), 6.51 (1H, t, J = 2.4 Hz), 4.70 (1H, s), 4.59 (1H, s), 4.13 (2H, s), 4.09 (1H, s), 2.27-2.11 (3H, m), 2.03-1.85 (4H, m), 1.64-1.48 (2H, m), 1.05 (6H, s).
184		 Racemate Relative configuration			free	541.1	¹ H-NMR (DMSO-D ₆) δ: 7.79-7.72 (3H, m), 7.60-7.53 (1H, m), 7.47 (1H, d, J = 7.3 Hz), 7.37-7.28 (3H, m), 7.14-7.12 (1H, m), 6.45 (1H, d, J = 2.4 Hz), 4.62 (1H, s), 4.53-3.89 (4H, m), 3.10-3.04 (1H, m), 2.45-2.39 (1H, m), 1.87-1.79 (1H, m), 1.72-1.69 (2H, m), 1.41-1.17 (2H, m), 1.05 (6H, s).









[0811]

Table 24

185		 Racemate Relative configuration			free	503	¹ H-NMR (DMSO-D ₆) δ: 7.98 (1H, d, J = 5.2 Hz), 7.78 (1H, t, J = 7.5 Hz), 7.75-7.70 (1H, m), 7.63 (2H, d, J = 7.6 Hz), 7.55-7.48 (1H, m), 7.17 (1H, dd, J = 8.2, 1.5 Hz), 4.48-3.88 (6H, m), 2.24-2.12 (2H, m), 1.75-1.51 (3H, m), 0.89 (1H, dd, J = 12.1, 4.7 Hz).
186					free	586.1	¹ H-NMR (DMSO-D ₆) δ: 8.12 (1H, s), 7.76 (1H, t, J = 7.6 Hz), 7.65-7.59 (2H, m), 7.57 (1H, d, J = 1.2 Hz), 7.50 (1H, d, J = 6.1 Hz), 7.36 (1H, dd, J = 10.5, 1.4 Hz), 7.12 (1H, dd, J = 8.1, 1.7 Hz), 4.62 (1H, s), 4.58 (1H, s), 4.41 (2H, s), 4.06 (1H, s), 3.72 (3H, d, J = 1.5 Hz), 2.30-2.18 (2H, m), 2.14-2.08 (1H, m), 2.00-1.84 (3H, m), 1.62-1.49 (3H, m), 1.09 (6H, s).
187					free	534.1	¹ H-NMR (DMSO-D ₆) δ: 8.19 (1H, d, J = 1.8 Hz), 7.78-7.75 (1H, m), 7.73-7.70 (1H, m), 7.65 (1H, dd, J = 8.5, 1.5 Hz), 7.62-7.58 (2H, m), 7.42-7.39 (1H, m), 7.12 (1H, dd, J = 7.9, 1.5 Hz), 4.68 (1H, s), 4.33 (2H, s), 3.72-3.59 (4H, m), 3.20-3.15 (1H, m), 2.01-1.94 (1H, m), 1.69-1.63 (1H, m), 1.10 (6H, s).
188					free	574.1	¹ H-NMR (DMSO-D ₆) δ: 8.19 (1H, d, J = 1.8 Hz), 7.76 (1H, dd, J = 8.1, 7.2 Hz), 7.64 (1H, dd, J = 7.9, 1.5 Hz), 7.61-7.57 (3H, m), 7.42 (1H, dd, J = 10.5, 1.4 Hz), 7.14-7.11 (1H, m), 4.68 (1H, s), 4.57 (1H, s), 4.33 (2H, s), 4.05 (1H, s), 2.33-2.20 (2H, m), 2.12-2.07 (1H, m), 1.97-1.83 (3H, m), 1.61-1.48 (3H, m), 1.10 (6H, s).
189		 Racemate Relative configuration			free	542.1	¹ H-NMR (DMSO-D ₆) δ: 8.10 (1H, d, J = 0.9 Hz), 7.80-7.74 (2H, m), 7.73-7.67 (1H, m), 7.64-7.56 (2H, m), 7.45 (1H, d, J = 11.0 Hz), 7.40 (1H, dd, J = 10.5, 1.4 Hz), 7.13 (1H, dd, J = 7.9, 1.5 Hz), 4.61 (1H, s), 4.49-3.89 (5H, m), 2.26-2.13 (2H, m), 1.81-1.50 (4H, m), 1.10 (6H, s).
190					free	529.2	¹ H-NMR (DMSO-D ₆) δ: 7.76-7.73 (1H, m), 7.70-7.60 (2H, m), 7.56-7.54 (1H, m), 7.47 (1H, d, J = 7.3 Hz), 7.35 (1H, d, J = 3.1 Hz), 7.32-7.28 (2H, m), 7.11 (1H, d, J = 8.2 Hz), 6.44 (1H, d, J = 3.1 Hz), 4.62 (1H, s), 4.01 (2H, s), 3.75-3.50 (4H, m), 1.77-1.73 (2H, m), 1.25-1.16 (3H, m), 1.05 (6H, s).
191					free	549.1	¹ H-NMR (DMSO-D ₆) δ: 7.98-7.95 (1H, m), 7.78-7.64 (4H, m), 7.48-7.44 (1H, m), 7.17-7.14 (1H, m), 4.82-4.81 (1H, m), 4.63-4.60 (2H, m), 3.72-3.50 (4H, m), 1.80-1.73 (2H, m), 1.26-1.15 (9H, m).
192		 Racemate Relative configuration			free	561.1	¹ H-NMR (DMSO-D ₆) δ: 7.97 (1H, d, J = 5.2 Hz), 7.78-7.73 (2H, m), 7.66 (2H, d, J = 7.6 Hz), 7.49 (1H, dd, J = 10.4, 1.2 Hz), 7.17 (1H, dd, J = 8.2, 1.5 Hz), 4.81 (1H, s), 4.61 (2H, s), 4.49-3.87 (3H, m), 2.28-2.11 (2H, m), 1.80-1.51 (3H, m), 1.15 (6H, s), 0.89 (1H, dd, J = 11.7, 4.7 Hz).







[0812]

Table 25

193		 Racemate Relative configuration			free	502.2	¹ H-NMR (DMSO- <i>d</i> ₆) δ: 7.80 (1H, t, J = 7.6 Hz), 7.68 (1H, s), 7.60 (1H, s), 7.53 (1H, d, J = 7.9 Hz), 7.29 (1H, t, J = 5.3 Hz), 7.21 (1H, t, J = 7.9 Hz), 7.15 (1H, dd, J = 8.2, 1.5 Hz), 7.07 (1H, d, J = 7.9 Hz), 6.95 (1H, d, J = 11.6 Hz), 4.46-3.86 (3H, m), 2.65 (2H, s), 2.25-2.10 (2H, m), 1.74-1.49 (4H, m), 1.04 (6H, s), 0.87 (1H, dd, J = 12.2, 4.6 Hz).
194					free	570.2	¹ H-NMR (DMSO- <i>d</i> ₆) δ: 8.09 (1H, d, J = 0.6 Hz), 7.79-7.74 (2H, m), 7.63 (1H, dd, J = 7.9, 1.5 Hz), 7.57-7.55 (2H, m), 7.43 (1H, d, J = 10.7 Hz), 7.39 (1H, dd, J = 10.5, 1.4 Hz), 7.12 (1H, dd, J = 8.2, 1.5 Hz), 4.59 (1H, s), 4.48 (1H, s), 4.42-4.38 (2H, m), 4.08 (1H, s), 2.23-2.12 (3H, m), 1.98-1.87 (5H, m), 1.62-1.51 (3H, m), 1.12 (6H, s).
195		 Racemate Relative configuration			free	560.1	¹ H-NMR (DMSO- <i>d</i> ₆) δ: 8.19 (1H, d, J = 1.8 Hz), 7.79-7.75 (1H, m), 7.73-7.70 (1H, m), 7.64-7.57 (3H, m), 7.43 (1H, dd, J = 10.5, 1.4 Hz), 7.14 (1H, dd, J = 8.2, 1.7 Hz), 4.69 (1H, s), 4.33-3.66 (5H, m), 2.20-2.14 (2H, m), 1.78-1.72 (1H, m), 1.65-1.56 (2H, m), 1.10 (6H, s), 0.93-0.89 (1H, m).
196		 Racemate Relative configuration			free	559.1	¹ H-NMR (DMSO- <i>d</i> ₆) δ: 7.76 (1H, t, J = 7.6 Hz), 7.62-7.58 (2H, m), 7.38-7.30 (4H, m), 7.14 (1H, dd, J = 8.2, 1.5 Hz), 6.52-6.50 (1H, m), 4.71 (1H, s), 4.44-3.93 (5H, m), 2.19-2.13 (2H, m), 1.78-1.72 (1H, m), 1.64-1.50 (2H, m), 1.04 (6H, s), 0.92-0.88 (1H, m).
197					free	548.2	¹ H-NMR (DMSO- <i>d</i> ₆) δ: 8.19 (1H, d, J = 1.8 Hz), 7.79-7.69 (3H, m), 7.64-7.58 (2H, m), 7.40 (1H, dd, J = 9.9, 4.1 Hz), 7.12 (1H, dd, J = 8.1, 1.7 Hz), 4.69 (1H, s), 4.33 (2H, s), 3.75-3.64 (4H, m), 1.80 (2H, t, J = 7.3 Hz), 1.30-1.18 (3H, m), 1.10 (6H, s).
198					free	547.2	¹ H-NMR (DMSO- <i>d</i> ₆) δ: 7.76 (1H, t, J = 7.5 Hz), 7.71-7.58 (3H, m), 7.37 (1H, d, J = 3.1 Hz), 7.31-7.29 (1H, m), 7.12 (1H, dd, J = 8.1, 1.4 Hz), 6.56 (1H, s), 6.52-6.50 (1H, m), 4.71 (1H, s), 4.13 (2H, s), 3.74-3.63 (4H, m), 1.81-1.78 (2H, m), 1.29-1.17 (3H, m), 1.04 (6H, s).
199					free	542.1	¹ H-NMR (DMSO- <i>d</i> ₆) δ: 8.10 (1H, d, J = 3.7 Hz), 7.77-7.74 (2H, m), 7.72-7.69 (1H, m), 7.65 (1H, d, J = 9.2 Hz), 7.58 (1H, dd, J = 12.5, 7.9 Hz), 7.45 (1H, dd, J = 10.7, 4.6 Hz), 7.40-7.36 (1H, m), 7.13-7.09 (1H, m), 4.61 (1H, s), 4.25 (2H, s), 3.82-3.52 (8H, m), 2.13-2.07 (2H, m), 1.10 (6H, s).
200					free	559.1	¹ H-NMR (DMSO- <i>d</i> ₆) δ: 7.78-7.74 (1H, m), 7.72-7.68 (1H, m), 7.66-7.64 (1H, m), 7.62-7.58 (1H, m), 7.39-7.29 (3H, m), 7.14-7.10 (1H, m), 6.51 (1H, s), 4.71 (1H, s), 4.13 (2H, s), 3.76-3.50 (6H, m), 2.11-2.07 (2H, m), 1.05 (6H, s).









[0813]

Table 26

201					free	573.2	¹ H-NMR (DMSO-D ₆) δ: 7.82 (1H, dd, J = 7.9, 1.8 Hz), 7.77 (1H, t, J = 7.5 Hz), 7.74 (1H, d, J = 1.8 Hz), 7.61 (1H, d, J = 7.9 Hz), 7.38 (1H, d, J = 3.1 Hz), 7.34-7.29 (2H, m), 7.12 (1H, dd, J = 7.9, 1.5 Hz), 6.52-6.50 (1H, m), 4.71 (1H, s), 4.13-4.04 (4H, m), 3.75 (2H, dd, J = 46.2, 9.9 Hz), 3.25-3.16 (2H, m), 2.86-2.81 (2H, m), 1.73-1.68 (2H, m), 1.47-1.41 (2H, m), 1.04 (6H, s).
202		 isomer-X Relative configuration			free	485.2	¹ H-NMR (DMSO-D ₆) δ: 7.83 (1H, d, J = 5.5 Hz), 7.78 (1H, t, J = 7.5 Hz), 7.74-7.72 (1H, m), 7.66-7.61 (3H, m), 7.47 (1H, dd, J = 10.5, 1.4 Hz), 7.14 (1H, dt, J = 7.9, 1.5 Hz), 4.46-3.90 (2H, m), 3.22-3.16 (1H, m), 2.52 (3H, s), 2.21-2.12 (2H, m), 1.78-1.72 (1H, m), 1.64-1.50 (2H, m), 0.93-0.90 (1H, m).
203					free	559.1	¹ H-NMR (DMSO-D ₆) δ: 7.76 (1H, t, J = 7.5 Hz), 7.67 (1H, dd, J = 7.9, 1.8 Hz), 7.62-7.58 (2H, m), 7.38 (1H, d, J = 3.1 Hz), 7.33 (1H, dt, J = 10.5, 1.4 Hz), 7.30 (1H, d, J = 5.8 Hz), 7.12 (1H, dd, J = 7.9, 1.5 Hz), 6.52-6.50 (1H, m), 4.70 (1H, s), 4.13 (2H, s), 3.85-3.79 (4H, m), 3.07-2.75 (6H, m), 1.05 (6H, s).
204					free	544.2	¹ H-NMR (DMSO-D ₆) δ: 8.10 (1H, d, J = 0.6 Hz), 7.77-7.74 (2H, m), 7.71-7.66 (1H, m), 7.65-7.63 (1H, m), 7.57 (1H, d, J = 8.2 Hz), 7.43 (1H, d, J = 11.0 Hz), 7.37 (1H, dd, J = 10.5, 2.3 Hz), 7.12 (1H, dd, J = 8.2, 1.5 Hz), 4.33 (1H, s), 4.24 (2H, s), 3.66-3.60 (4H, m), 3.21-3.18 (1H, m), 2.04-1.99 (1H, m), 1.74-1.67 (1H, m), 1.38-1.30 (4H, m), 0.85 (6H, t, J = 7.3 Hz).
205					free	584.2	¹ H-NMR (DMSO-D ₆) δ: 8.10 (1H, d, J = 0.6 Hz), 7.77-7.74 (2H, m), 7.63 (1H, dd, J = 7.9, 1.8 Hz), 7.58-7.55 (2H, m), 7.44 (1H, d, J = 11.0 Hz), 7.38 (1H, dd, J = 10.7, 1.2 Hz), 7.12 (1H, dt, J = 9.1, 1.4 Hz), 4.59 (1H, s), 4.34 (1H, s), 4.24 (2H, s), 4.08 (1H, s), 2.23-2.13 (3H, m), 2.02-1.88 (3H, m), 1.80-1.50 (3H, m), 1.39-1.31 (4H, m), 0.85 (6H, t, J = 7.3 Hz).
206		 isomer-X Relative configuration			free	570.2	¹ H-NMR (DMSO-D ₆) δ: 8.10 (1H, d, J = 0.9 Hz), 7.77-7.74 (2H, m), 7.72-7.69 (1H, m), 7.62-7.56 (2H, m), 7.44 (1H, d, J = 11.0 Hz), 7.39 (1H, dd, J = 10.5, 1.4 Hz), 7.13 (1H, dd, J = 8.2, 1.5 Hz), 4.45-3.64 (6H, m), 2.19-2.13 (2H, m), 1.79-1.73 (1H, m), 1.65-1.51 (2H, m), 1.37-1.31 (4H, m), 0.92 (1H, dd, J = 12.1, 4.1 Hz), 0.85 (6H, t, J = 7.3 Hz).
207		 isomer-X Relative configuration			free	527.2	¹ H-NMR (DMSO-D ₆) δ: 7.75 (1H, t, J = 7.5 Hz), 7.70 (1H, d, J = 7.9 Hz), 7.61 (1H, s), 7.54 (1H, d, J = 7.9 Hz), 7.48 (1H, d, J = 7.3 Hz), 7.38 (1H, d, J = 10.1 Hz), 7.33 (1H, d, J = 3.1 Hz), 7.15 (1H, d, J = 11.0 Hz), 7.12 (1H, d, J = 6.9 Hz), 6.43 (1H, d, J = 2.7 Hz), 4.79 (2H, s), 4.46-2.48 (3H, m), 2.24-2.05 (2H, m), 1.80-1.76 (1H, m), 1.69-1.64 (1H, m), 1.57-1.53 (1H, m), 1.06-1.01 (1H, m).













[0814]

Table 27

208		 isomer-X Relative configuration			free	488.2	¹ H-NMR (DMSO-D ₆) δ: 7.80 (1H, t, J = 7.5 Hz), 7.71-7.68 (1H, m), 7.61-7.58 (1H, m), 7.53 (1H, d, J = 7.9 Hz), 7.37 (1H, dd, J = 10.5, 1.4 Hz), 7.26 (1H, t, J = 7.9 Hz), 7.14-7.11 (2H, m), 7.03 (1H, d, J = 11.3 Hz), 4.43-3.57 (5H, m), 2.19-2.13 (2H, m), 1.78-1.72 (1H, m), 1.59-1.50 (2H, m), 0.90 (1H, dd, J = 11.9, 4.3 Hz).
209		 isomer-X Relative configuration			free	577.2 579.2	¹ H-NMR (DMSO-D ₆) δ: 8.10 (1H, d, J = 6.1 Hz), 7.74-7.69 (2H, m), 7.63-7.59 (2H, m), 7.44 (1H, d, J = 10.4 Hz), 7.11 (1H, dd, J = 7.9, 1.5 Hz), 4.78 (2H, s), 4.71 (1H, s), 4.41-3.56 (3H, m), 2.18-2.10 (2H, m), 1.75-1.68 (1H, m), 1.60-1.54 (1H, m), 1.51-1.46 (1H, m), 1.13 (6H, s), 0.86 (1H, dd, J = 12.1, 4.7 Hz).
210		 isomer-X Relative configuration			free	551.2 553.2	¹ H-NMR (DMSO-D ₆) δ: 8.10 (1H, d, J = 6.1 Hz), 7.73-7.68 (2H, m), 7.65-7.61 (2H, m), 7.43-7.40 (1H, m), 7.10 (1H, dd, J = 8.1, 1.4 Hz), 4.78 (2H, s), 4.71 (1H, s), 3.65-3.46 (4H, m), 3.24-3.23 (1H, m), 1.98-1.93 (1H, m), 1.63-1.59 (1H, m), 1.13 (6H, s).
211		 isomer-X Relative configuration			free	591.2 593.2	¹ H-NMR (DMSO-D ₆) δ: 8.10 (1H, d, J = 6.1 Hz), 7.71 (1H, t, J = 7.5 Hz), 7.62-7.60 (2H, m), 7.55 (1H, s), 7.42 (1H, dd, J = 10.5, 1.4 Hz), 7.10 (1H, dd, J = 7.9, 1.5 Hz), 4.77 (2H, s), 4.70 (1H, s), 4.53 (1H, s), 4.01 (1H, s), 2.22-2.04 (3H, m), 1.92-1.80 (3H, m), 1.56-1.45 (2H, m), 1.12 (6H, s).
212		 isomer-X Relative configuration			free	586.2	¹ H-NMR (DMSO-D ₆) δ: 8.10 (1H, d, J = 7.3 Hz), 7.76-7.70 (2H, m), 7.63-7.58 (2H, m), 7.48 (1H, d, J = 10.7 Hz), 7.41 (1H, d, J = 10.4 Hz), 7.12 (1H, dd, J = 7.9, 1.5 Hz), 4.67 (1H, s), 4.46-3.50 (5H, m), 2.21-2.12 (2H, m), 1.80-1.74 (1H, m), 1.66-1.60 (1H, m), 1.56-1.51 (1H, m), 1.11 (6H, s), 0.98-0.93 (1H, m).
213		 Racemate Relative configuration			free	577.3 579.2	¹ H-NMR (DMSO-D ₆) δ: 8.14 (1H, d, J = 6.1 Hz), 7.78-7.73 (2H, m), 7.68-7.63 (2H, m), 7.48 (1H, d, J = 9.2 Hz), 7.15 (1H, dd, J = 8.1, 1.1 Hz), 4.81 (2H, s), 4.75 (1H, s), 4.45-3.82 (3H, m), 2.22-2.12 (2H, m), 1.79-1.73 (1H, m), 1.64-1.59 (1H, m), 1.55-1.50 (1H, m), 1.17 (6H, s), 0.91 (1H, dd, J = 11.2, 4.4 Hz).
214		 isomer-X Relative configuration			free	589.3	¹ H-NMR (DMSO-D ₆) δ: 7.97 (1H, d, J = 5.2 Hz), 7.79-7.72 (2H, m), 7.66-7.63 (2H, m), 7.49 (1H, dd, J = 10.4, 1.2 Hz), 7.17 (1H, dd, J = 7.9, 1.5 Hz), 4.61 (2H, s), 4.45 (1H, s), 4.33-3.50 (3H, m), 2.21-2.15 (2H, m), 1.79-1.72 (1H, m), 1.64-1.50 (2H, m), 1.44-1.35 (4H, m), 0.91-0.87 (7H, m).
215		 isomer-X Relative configuration			free	563.3	¹ H-NMR (DMSO-D ₆) δ: 7.97 (1H, d, J = 4.9 Hz), 7.76-7.72 (2H, m), 7.68 (1H, d, J = 8.2 Hz), 7.64 (1H, d, J = 7.6 Hz), 7.47 (1H, d, J = 10.4 Hz), 7.15 (1H, dd, J = 7.9, 1.5 Hz), 4.61 (2H, s), 4.46 (1H, s), 3.67-3.43 (4H, m), 3.22-3.16 (1H, m), 2.04-1.90 (1H, m), 1.70-1.62 (1H, m), 1.44-1.35 (4H, m), 0.89 (5H, t, J = 7.5 Hz).






















[0815]

Table 28

216					free	603.3	¹ H-NMR (DMSO-D ₆) δ: 7.97 (1H, d, J = 5.2 Hz), 7.77 (1H, t, J = 7.5 Hz), 7.67-7.59 (3H, m), 7.48 (1H, d, J = 11.3 Hz), 7.16 (1H, d, J = 8.9 Hz), 4.61 (2H, s), 4.58 (1H, s), 4.46 (1H, s), 4.05 (1H, s), 2.33-2.20 (2H, m), 2.12-2.07 (1H, m), 1.96-1.84 (3H, m), 1.61-1.49 (3H, m), 1.42-1.37 (4H, m), 0.89 (6H, t, J = 7.3 Hz).
217		 Racemate Relative configuration			free	589.3	¹ H-NMR (DMSO-D ₆) δ: 7.97 (1H, d, J = 5.2 Hz), 7.79-7.72 (2H, m), 7.66-7.63 (2H, m), 7.49 (1H, dd, J = 10.4, 1.2 Hz), 7.17 (1H, dd, J = 7.9, 1.5 Hz), 4.61 (2H, s), 4.45 (1H, s), 4.33-3.50 (3H, m), 2.21-2.15 (2H, m), 1.79-1.72 (1H, m), 1.64-1.50 (2H, m), 1.44-1.35 (4H, m), 0.91-0.87 (7H, m).
218		 isomer-X Relative configuration			free	605.2 607.2	¹ H-NMR (DMSO-D ₆) δ: 8.14 (1H, d, J = 6.1 Hz), 7.79-7.72 (2H, m), 7.67-7.63 (2H, m), 7.50-7.48 (1H, m), 7.15 (1H, dd, J = 7.9, 1.5 Hz), 4.81 (2H, s), 4.45-3.92 (4H, m), 2.22-2.13 (2H, m), 1.79-1.73 (1H, m), 1.64-1.59 (1H, m), 1.55-1.37 (5H, m), 0.93-0.87 (7H, m).
219					free	579.2 581.3	¹ H-NMR (DMSO-D ₆) δ: 8.14 (1H, d, J = 5.5 Hz), 7.78-7.72 (2H, m), 7.68 (1H, d, J = 7.9 Hz), 7.65 (1H, d, J = 7.9 Hz), 7.47 (1H, dd, J = 10.4, 3.7 Hz), 7.14 (1H, d, J = 7.9 Hz), 4.62 (2H, s), 4.37 (1H, s), 3.69-3.49 (4H, m), 3.23-3.20 (1H, m), 2.06-1.97 (1H, m), 1.75-1.67 (1H, m), 1.52-1.35 (4H, m), 0.90 (6H, t, J = 7.3 Hz).
220					free	619.3 621.2	¹ H-NMR (DMSO-D ₆) δ: 8.14 (1H, d, J = 6.1 Hz), 7.78-7.74 (1H, m), 7.68-7.59 (3H, m), 7.50-7.47 (1H, m), 7.14 (1H, dd, J = 7.9, 1.5 Hz), 4.82 (2H, s), 4.60 (1H, s), 4.37 (1H, s), 4.07 (1H, s), 2.24-2.13 (3H, m), 2.02-1.88 (3H, m), 1.63-1.38 (7H, m), 0.90 (6H, t, J = 7.5 Hz).
221		 Racemate Relative configuration			free	605.2 607.2	¹ H-NMR (DMSO-D ₆) δ: 8.14 (1H, d, J = 6.1 Hz), 7.79-7.72 (2H, m), 7.57-7.63 (2H, m), 7.50-7.48 (1H, m), 7.15 (1H, dd, J = 7.9, 1.5 Hz), 4.81 (2H, s), 4.45-3.92 (4H, m), 2.22-2.13 (2H, m), 1.79-1.73 (1H, m), 1.64-1.59 (1H, m), 1.55-1.37 (5H, m), 0.93-0.87 (7H, m).
222		 isomer-X Relative configuration			free	585.3	¹ H-NMR (DMSO-D ₆) δ: 7.99 (1H, s), 7.94 (1H, d, J = 7.5 Hz), 7.76-7.68 (2H, m), 7.62-7.58 (2H, m), 7.46 (1H, d, J = 11.3 Hz), 7.34 (1H, t, J = 5.3 Hz), 7.11 (1H, dd, J = 7.9, 1.5 Hz), 4.73 (1H, s), 4.44-3.51 (5H, m), 2.21-2.13 (2H, m), 1.78-1.72 (1H, m), 1.60-1.50 (2H, m), 1.06 (6H, s), 0.89 (1H, dd, J = 12.1, 4.4 Hz).
223		 isomer-X Relative configuration			free	526.2	¹ H-NMR (DMSO-D ₆) δ: 8.16 (1H, d, J = 7.3 Hz), 7.77-7.70 (2H, m), 7.62-7.60 (1H, m), 7.57 (1H, d, J = 7.9 Hz), 7.43 (2H, d, J = 10.4 Hz), 7.09 (1H, dd, J = 7.9, 1.5 Hz), 4.44-3.50 (6H, m), 2.19-2.14 (2H, m), 1.79-1.74 (1H, m), 1.64-1.59 (1H, m), 1.55-1.50 (1H, m), 0.94-0.90 (1H, m).








[0816]

Table 29

224		 isomer-X Relative configuration			free	527.2	¹ H-NMR (DMSO-D ₆) δ: 8.07 (1H, s), 7.96 (1H, d, J = 7.3 Hz), 7.77-7.69 (2H, m), 7.62-7.55 (2H, m), 7.41 (1H, dd, J = 10.5, 1.4 Hz), 7.36 (1H, d, J = 10.7 Hz), 7.10 (1H, dd, J = 8.2, 1.5 Hz), 4.44-3.51 (6H, m), 2.20-2.13 (2H, m), 1.79-1.73 (1H, m), 1.60-1.50 (2H, m), 0.91 (1H, dd, J = 12.2, 4.6 Hz).
225		 Racemate Relative configuration			free	570.3	¹ H-NMR (DMSO-D ₆) δ: 8.10 (1H, d, J = 0.9 Hz), 7.77-7.74 (2H, m), 7.72-7.69 (1H, m), 7.62-7.56 (2H, m), 7.44 (1H, d, J = 11.0 Hz), 7.39 (1H, dd, J = 10.5, 1.4 Hz), 7.13 (1H, dd, J = 8.2, 1.5 Hz), 4.45-3.64 (6H, m), 2.19-2.13 (2H, m), 1.79-1.73 (1H, m), 1.65-1.51 (2H, m), 1.37-1.31 (4H, m), 0.92 (1H, dd, J = 12.1, 4.1 Hz), 0.85 (6H, t, J = 7.3 Hz).
226		 isomer-X Relative configuration			free	544.3	¹ H-NMR (DMSO-D ₆) δ: 7.83-7.80 (1H, m), 7.70-7.67 (1H, m), 7.61-7.58 (1H, m), 7.53 (1H, d, J = 7.9 Hz), 7.41-7.38 (1H, m), 7.28 (1H, d, J = 6.7 Hz), 7.13 (1H, d, J = 7.9 Hz), 7.02 (1H, d, J = 10.1 Hz), 5.30 (1H, d, J = 8.9 Hz), 4.99-4.87 (2H, m), 4.44-3.53 (4H, m), 2.21-2.11 (2H, m), 1.92-1.88 (1H, m), 1.77-1.49 (4H, m), 1.23-1.18 (6H, m), 0.88 (1H, dd, J = 12.1, 4.7 Hz).
227		 isomer-X Relative configuration			free	518.2	¹ H-NMR (DMSO-D ₆) δ: 7.83-7.80 (1H, m), 7.70-7.67 (1H, m), 7.64-7.62 (1H, m), 7.52 (1H, dd, J = 7.9, 2.1 Hz), 7.37 (1H, d, J = 9.5 Hz), 7.29 (1H, d, J = 6.4 Hz), 7.11 (1H, dd, J = 7.9, 1.5 Hz), 7.01 (1H, d, J = 9.8 Hz), 5.30 (1H, d, J = 7.3 Hz), 4.99-4.88 (2H, m), 4.35-4.33 (1H, m), 3.67-3.50 (4H, m), 3.17-3.12 (1H, m), 2.01-1.88 (2H, m), 1.71-1.62 (2H, m), 1.22-1.19 (6H, m).
228		 isomer-X Relative configuration			free	557.2	¹ H-NMR (DMSO-D ₆) δ: 7.87 (1H, d, J = 6.1 Hz), 7.79-7.68 (2H, m), 7.65-7.56 (2H, m), 7.46 (1H, d, J = 10.4 Hz), 7.15 (1H, dd, J = 8.2, 1.5 Hz), 4.78 (1H, s), 4.69 (2H, s), 4.49-3.87 (3H, m), 2.49 (3H, s), 2.24-2.11 (2H, m), 1.81-1.48 (3H, m), 1.20 (6H, s), 0.90 (1H, dd, J = 12.2, 4.3 Hz).
229		 isomer-X Relative configuration			free	517.2	¹ H-NMR (DMSO-D ₆) δ: 7.81-7.62 (6H, m), 7.44 (1H, d, J = 8.9 Hz), 7.16-7.13 (1H, m), 4.70 (1H, s), 2.73-3.51 (5H, m), 3.03 (2H, s), 2.03-1.93 (1H, m), 1.70-1.61 (1H, m), 1.18 (6H, s).
230		 Racemate Relative configuration			free	526.2	¹ H-NMR (DMSO-D ₆) δ: 8.34 (2H, s), 7.76 (1H, t, J = 7.5 Hz), 7.53 (3H, dd, J = 15.4, 7.5 Hz), 7.40 (1H, d, J = 10.4 Hz), 7.34 (1H, d, J = 3.1 Hz), 7.24 (1H, s), 7.14 (2H, dd, J = 12.8, 6.1 Hz), 6.47 (1H, d, J = 3.4 Hz), 4.76 (2H, s), 4.45-3.94 (3H, m), 2.26-2.14 (2H, m), 1.82-1.70 (1H, m), 1.66-1.46 (2H, m), 0.90 (1H, dd, J = 12.2, 4.6 Hz).






















[0817]

Table 30

231		 Racemate Relative configuration			free	540.2	¹ H-NMR (DMSO- <i>d</i> ₆) δ: 8.03 (1H, d, <i>J</i> = 4.6 Hz), 7.91 (1H, d, <i>J</i> = 7.0 Hz), 7.76 (1H, t, <i>J</i> = 7.5 Hz), 7.56-7.48 (2H, m), 7.47-7.38 (2H, m), 7.34 (1H, d, <i>J</i> = 3.1 Hz), 7.20-7.09 (2H, m), 6.47 (1H, d, <i>J</i> = 2.7 Hz), 4.76 (2H, s), 4.46-3.92 (3H, m), 2.61 (3H, d, <i>J</i> = 4.5 Hz), 2.26-2.13 (2H, m), 1.82-1.72 (1H, m), 1.68-1.46 (2H, m), 0.90 (1H, dd, <i>J</i> = 12.1, 4.4 Hz).
232		 Racemate Relative configuration			free	554.2	¹ H-NMR (DMSO- <i>d</i> ₆) δ: 7.93 (1H, t, <i>J</i> = 4.1 Hz), 7.77 (1H, t, <i>J</i> = 7.6 Hz), 7.53 (2H, dd, <i>J</i> = 19.1, 7.5 Hz), 7.47 (1H, d, <i>J</i> = 7.9 Hz), 7.39 (1H, d, <i>J</i> = 10.4 Hz), 7.27 (1H, d, <i>J</i> = 3.1 Hz), 7.15 (2H, dt, <i>J</i> = 13.2, 6.9 Hz), 6.46 (1H, d, <i>J</i> = 3.4 Hz), 5.11 (2H, s), 4.46-3.87 (3H, m), 3.08 (3H, d, <i>J</i> = 12.2 Hz), 2.85 (3H, s), 2.25-2.15 (2H, m), 1.82-1.71 (1H, m), 1.67-1.50 (2H, m), 0.95-0.87 (1H, m).
233		 Racemate Relative configuration			free	487.2	¹ H-NMR (DMSO- <i>d</i> ₆) δ: 7.83 (1H, dd, <i>J</i> = 23.7, 7.2 Hz), 7.72-7.56 (1H, m), 7.52 (1H, t, <i>J</i> = 7.5 Hz), 7.33 (3H, ddd, <i>J</i> = 33.5, 19.5, 8.0 Hz), 7.05 (3H, dd, <i>J</i> = 53.3, 15.2 Hz), 4.40-3.87 (3H, m), 3.01 (2H, s), 2.22-2.14 (2H, m), 1.83-1.57 (3H, m), 0.91-0.85 (1H, m).
234		 Racemate Relative configuration			free	501.2	¹ H-NMR (DMSO- <i>d</i> ₆) δ: 7.83-7.79 (2H, m), 7.56-7.45 (2H, m), 7.38 (1H, d, <i>J</i> = 10.7 Hz), 7.27 (1H, t, <i>J</i> = 7.9 Hz), 7.12 (2H, td, <i>J</i> = 9.1, 3.7 Hz), 7.00 (1H, d, <i>J</i> = 11.0 Hz), 4.45-4.07 (3H, m), 3.02 (2H, s), 2.58 (3H, d, <i>J</i> = 4.5 Hz), 2.21-2.15 (2H, m), 1.79-1.72 (1H, m), 1.64-1.51 (2H, m), 0.93-0.85 (1H, m).
235		 Racemate Relative configuration			free	515.2	¹ H-NMR (DMSO- <i>d</i> ₆) δ: 7.95-7.91 (1H, m), 7.82 (1H, t, <i>J</i> = 7.6 Hz), 7.54 (1H, d, <i>J</i> = 7.9 Hz), 7.49-7.44 (1H, m), 7.35 (1H, d, <i>J</i> = 10.7 Hz), 7.27 (1H, t, <i>J</i> = 7.9 Hz), 7.15 (1H, dd, <i>J</i> = 8.2, 1.5 Hz), 7.06 (1H, d, <i>J</i> = 7.6 Hz), 6.97 (1H, d, <i>J</i> = 11.3 Hz), 4.47-3.89 (3H, m), 3.72 (2H, s), 3.00 (3H, s), 2.84 (3H, s), 2.22-2.13 (2H, m), 1.79-1.72 (1H, m), 1.67-1.48 (2H, m), 0.89 (1H, s).
236		 isomer-X Relative configuration			free	526.2	¹ H-NMR (DMSO- <i>d</i> ₆) δ: 8.14 (1H, d, <i>J</i> = 7.6 Hz), 8.01 (1H, s), 7.76 (1H, t, <i>J</i> = 7.6 Hz), 7.72-7.59 (1H, m), 7.60 (1H, s), 7.56 (1H, d, <i>J</i> = 7.9 Hz), 7.40 (1H, d, <i>J</i> = 10.4 Hz), 7.30 (1H, d, <i>J</i> = 10.7 Hz), 7.11 (1H, dd, <i>J</i> = 7.9, 1.5 Hz), 4.46-3.93 (3H, m), 3.77 (3H, s), 2.25-2.14 (2H, m), 1.81-1.72 (1H, m), 1.67-1.49 (2H, m), 0.93-0.86 (1H, m).
237		 isomer-X Relative configuration			free	540.2	¹ H-NMR (DMSO- <i>d</i> ₆) δ: 8.13 (1H, d, <i>J</i> = 7.6 Hz), 7.94 (1H, s), 7.92-7.89 (1H, m), 7.78-7.55 (4H, m), 7.39 (1H, d, <i>J</i> = 9.2 Hz), 7.30 (1H, d, <i>J</i> = 10.7 Hz), 7.12 (1H, d, <i>J</i> = 7.9 Hz), 4.47-3.91 (3H, m), 3.77 (3H, s), 2.76 (3H, d, <i>J</i> = 4.5 Hz), 2.23-2.15 (2H, m), 1.80-1.74 (1H, m), 1.65-1.50 (2H, m), 0.90 (1H, dd, <i>J</i> = 12.1, 4.7 Hz).











[0818]

Table 31

238		 isomer-X Relative configuration			free	554.2	¹ H-NMR (DMSO-D ₆) δ: 7.83 (1H, s), 7.90-7.75 (2H, m), 7.73-7.68 (1H, m), 7.57 (2H, t, J = 18.3 Hz), 7.40 (1H, dd, J = 10.7, 1.5 Hz), 7.32 (1H, d, J = 11.0 Hz), 7.11 (1H, dd, J = 7.9, 1.5 Hz), 4.48-3.92 (3H, m), 3.80 (3H, d, J = 7.0 Hz), 3.06 (6H, s), 2.25-2.14 (2H, m), 1.80-1.70 (1H, m), 1.65-1.48 (2H, m), 0.90 (1H, t, J = 8.4 Hz).
239					free	531.1	¹ H-NMR (DMSO-D ₆) δ: 7.87 (1H, d, J = 6.7 Hz), 7.60-7.64 (3H, m), 7.58 (1H, d, J = 7.9 Hz), 7.44 (1H, d, J = 10.4 Hz), 7.13 (1H, dd, J = 8.2, 1.5 Hz), 4.78 (1H, s), 4.69 (2H, s), 3.75-3.58 (5H, m), 2.49 (3H, s), 2.04-1.93 (1H, m), 1.74-1.64 (1H, m), 1.20 (6H, s).
240					free	571.2	¹ H-NMR (DMSO-D ₆) δ: 7.87 (1H, d, J = 6.7 Hz), 7.75 (1H, t, J = 7.5 Hz), 7.63 (1H, d, J = 7.3 Hz), 7.60-7.55 (2H, m), 7.45 (1H, d, J = 10.4 Hz), 7.14 (1H, d, J = 7.9 Hz), 4.78 (1H, s), 4.69 (2H, s), 4.64-4.56 (1H, m), 4.11-4.04 (1H, m), 2.49 (3H, s), 2.34-2.20 (2H, m), 2.13-2.06 (1H, m), 1.99-1.85 (3H, m), 1.66-1.47 (3H, m), 1.20 (6H, s).
241		 Racemate Relative configuration			free	557.2	¹ H-NMR (DMSO-D ₆) δ: 7.87 (1H, d, J = 6.1 Hz), 7.79-7.68 (2H, m), 7.65-7.56 (2H, m), 7.46 (1H, d, J = 10.4 Hz), 7.15 (1H, dd, J = 8.2, 1.5 Hz), 4.78 (1H, s), 4.69 (2H, s), 4.49-3.87 (3H, m), 2.49 (3H, s), 2.24-2.11 (2H, m), 1.81-1.48 (3H, m), 1.20 (6H, s), 0.90 (1H, dd, J = 12.2, 4.3 Hz).
242		 Racemate Relative configuration			free	527.1	¹ H-NMR (DMSO-D ₆) δ: 8.07 (1H, s), 7.96 (1H, d, J = 7.3 Hz), 7.77-7.69 (2H, m), 7.62-7.55 (2H, m), 7.41 (1H, dd, J = 10.5, 1.4 Hz), 7.36 (1H, d, J = 10.7 Hz), 7.10 (1H, dd, J = 8.2, 1.5 Hz), 4.44-3.51 (6H, m), 2.20-2.13 (2H, m), 1.79-1.73 (1H, m), 1.60-1.50 (2H, m), 0.91 (1H, dd, J = 12.2, 4.6 Hz).
243		 Racemate Relative configuration			free	540.3	¹ H-NMR (DMSO-D ₆) δ: 8.13 (1H, d, J = 7.6 Hz), 7.94 (1H, s), 7.92-7.89 (1H, m), 7.78-7.55 (4H, m), 7.39 (1H, d, J = 9.2 Hz), 7.30 (1H, d, J = 10.7 Hz), 7.12 (1H, d, J = 7.9 Hz), 4.47-3.91 (3H, m), 3.77 (3H, s), 2.76 (3H, d, J = 4.6 Hz), 2.23-2.15 (2H, m), 1.80-1.74 (1H, m), 1.66-1.50 (2H, m), 0.90 (1H, dd, J = 12.1, 4.7 Hz).
244		 Racemate Relative configuration			free	554.2	¹ H-NMR (DMSO-D ₆) δ: 7.83 (1H, s), 7.90-7.75 (2H, m), 7.73-7.68 (1H, m), 7.57 (2H, t, J = 18.3 Hz), 7.40 (1H, dd, J = 10.7, 1.5 Hz), 7.32 (1H, d, J = 11.0 Hz), 7.11 (1H, dd, J = 7.9, 1.5 Hz), 4.48-3.92 (3H, m), 3.80 (3H, d, J = 7.0 Hz), 3.06 (6H, s), 2.25-2.14 (2H, m), 1.80-1.70 (1H, m), 1.65-1.48 (2H, m), 0.90 (1H, t, J = 8.4 Hz).












[0819]

Table 32

245		 Racemate Relative configuration			free	526.3	¹ H-NMR (DMSO-D ₆) δ: 8.14 (1H, d, J = 7.6 Hz), 8.01 (1H, s), 7.76 (1H, t, J = 7.6 Hz), 7.72-7.69 (1H, m), 7.60 (1H, s), 7.56 (1H, d, J = 7.9 Hz), 7.40 (1H, d, J = 10.4 Hz), 7.30 (1H, d, J = 10.7 Hz), 7.11 (1H, dd, J = 7.9, 1.5 Hz), 4.46-3.93 (3H, m), 3.77 (3H, s), 2.25-2.14 (2H, m), 1.81-1.72 (1H, m), 1.67-1.49 (2H, m), 0.93-0.86 (1H, m).
246		 isomer-X Relative configuration			free	593.3	¹ H-NMR (DMSO-D ₆) δ: 8.40 (1H, d, J = 6.1 Hz), 7.80-7.62 (5H, m), 7.51-7.42 (1H, m), 7.15 (1H, dd, J = 8.1, 1.4 Hz), 5.03 (1H, s), 4.75 (2H, s), 4.48-3.87 (3H, m), 2.27-2.12 (2H, m), 1.81-1.72 (1H, m), 1.68-1.50 (2H, m), 1.16 (6H, s), 0.91-0.86 (1H, m).
247		 isomer-X Relative configuration			free	568.3	¹ H-NMR (DMSO-D ₆) δ: 8.55 (1H, d, J = 6.7 Hz), 7.80-7.67 (4H, m), 7.53 (1H, d, J = 9.2 Hz), 7.19 (1H, dd, J = 8.2, 1.5 Hz), 4.83 (1H, s), 4.73 (2H, s), 4.52-3.87 (3H, m), 2.29-2.10 (2H, m), 1.83-1.72 (1H, m), 1.69-1.49 (2H, m), 1.22 (6H, s), 0.95-0.86 (1H, m).
248					free	567.3	¹ H-NMR (DMSO-D ₆) δ: 8.40 (1H, d, J = 5.5 Hz), 7.87-7.62 (5H, m), 7.43 (1H, d, J = 10.4 Hz), 7.14 (1H, dd, J = 8.1, 1.4 Hz), 5.03 (1H, s), 4.75 (2H, s), 3.79-3.60 (5H, m), 2.06-1.92 (1H, m), 1.75-1.62 (1H, m), 1.17 (6H, s).
249					free	542.3	¹ H-NMR (DMSO-D ₆) δ: 8.56 (1H, d, J = 6.4 Hz), 7.83-7.66 (4H, m), 7.51 (1H, d, J = 10.4 Hz), 7.22-7.14 (1H, m), 4.83 (1H, s), 4.73 (2H, s), 3.80-3.60 (5H, m), 2.06-1.92 (1H, m), 1.77-1.59 (1H, m), 1.22 (6H, s).
250					free	607.3	¹ H-NMR (DMSO-D ₆) δ: 8.41 (1H, d, J = 6.1 Hz), 7.89-7.59 (5H, m), 7.45 (1H, d, J = 10.4 Hz), 7.15 (1H, dd, J = 7.9, 1.5 Hz), 5.03 (1H, s), 4.75 (2H, s), 4.64-4.56 (1H, m), 4.11-4.02 (1H, m), 2.33-1.84 (6H, m), 1.66-1.46 (3H, m), 1.17 (6H, s).
251					free	582.3	¹ H-NMR (DMSO-D ₆) δ: 8.56 (1H, d, J = 6.4 Hz), 7.84-7.76 (1H, m), 7.68 (2H, s), 7.61 (1H, s), 7.52 (1H, d, J = 10.4 Hz), 7.19 (1H, dd, J = 8.1, 1.4 Hz), 4.83 (1H, s), 4.73 (2H, s), 4.63-4.55 (1H, m), 4.11-4.02 (1H, m), 2.33-1.82 (6H, m), 1.65-1.46 (3H, m), 1.22 (6H, s).
252		 Racemate Relative configuration			free	593.3	¹ H-NMR (DMSO-D ₆) δ: 8.40 (1H, d, J = 6.1 Hz), 7.80-7.62 (5H, m), 7.51-7.42 (1H, m), 7.15 (1H, dd, J = 8.1, 1.4 Hz), 5.03 (1H, s), 4.75 (2H, s), 4.48-3.87 (3H, m), 2.27-2.12 (2H, m), 1.81-1.72 (1H, m), 1.68-1.50 (2H, m), 1.16 (6H, s), 0.91-0.86 (1H, m).
253		 Racemate Relative configuration			free	568.3	¹ H-NMR (DMSO-D ₆) δ: 8.55 (1H, d, J = 6.7 Hz), 7.80-7.67 (4H, m), 7.53 (1H, d, J = 9.2 Hz), 7.19 (1H, dd, J = 8.2, 1.5 Hz), 4.83 (1H, s), 4.73 (2H, s), 4.52-3.87 (3H, m), 2.29-2.10 (2H, m), 1.83-1.72 (1H, m), 1.69-1.49 (2H, m), 1.22 (6H, s), 0.95-0.86 (1H, m).
254		 isomer-X Relative configuration			free	573.2	¹ H-NMR (DMSO-D ₆) δ: 7.96 (1H, d, J = 5.5 Hz), 7.83-7.61 (4H, m), 7.50 (1H, d, J = 10.4 Hz), 7.17 (1H, d, J = 7.9 Hz), 5.43 (1H, s), 4.78 (2H, s), 4.49-3.87 (3H, m), 2.21-1.96 (6H, m), 1.82-1.50 (5H, m), 0.99-0.79 (1H, m).























[0820]

Table 33

255					free	547.3	¹ H-NMR (DMSO-D ₆) δ: 7.95 (1H, d, J = 4.9 Hz), 7.81-7.72 (2H, m), 7.70-7.63 (2H, m), 7.48 (1H, d, J = 10.7 Hz), 7.16 (1H, d, J = 8.5 Hz), 5.43 (1H, s), 4.78 (2H, s), 3.75-3.59 (5H, m), 2.26-2.17 (2H, m), 2.04-1.91 (3H, m), 1.77-1.59 (3H, m).
256					free	587.2	¹ H-NMR (DMSO-D ₆) δ: 7.97 (1H, d, J = 5.2 Hz), 7.77 (1H, t, J = 7.5 Hz), 7.69-7.63 (2H, m), 7.59 (1H, s), 7.49 (1H, d, J = 10.7 Hz), 7.17 (1H, d, J = 8.5 Hz), 5.44 (1H, s), 4.78 (2H, s), 4.62-4.55 (1H, m), 4.07-4.01 (1H, m), 2.32-1.50 (15H, m).
257		 isomer-X Relative configuration			free	589.2 591.2	¹ H-NMR (DMSO-D ₆) δ: 8.14 (1H, d, J = 5.8 Hz), 7.79-7.63 (4H, m), 7.50 (1H, d, J = 9.8 Hz), 7.16 (1H, d, J = 8.5 Hz), 5.38 (1H, s), 4.97 (2H, s), 4.49-3.87 (3H, m), 2.25-1.97 (6H, m), 1.76-1.50 (5H, m), 0.95-0.83 (1H, m).
258					free	583.2 565.2	¹ H-NMR (DMSO-D ₆) δ: 8.14 (1H, d, J = 6.4 Hz), 7.78-7.73 (2H, m), 7.71-7.64 (2H, m), 7.47 (1H, d, J = 9.8 Hz), 7.14 (1H, d, J = 8.2 Hz), 5.38 (1H, s), 4.97 (2H, s), 3.68-3.59 (5H, m), 2.31-2.21 (2H, m), 2.10-1.93 (3H, m), 1.76-1.55 (3H, m).
259					free	603.2 605.2	¹ H-NMR (DMSO-D ₆) δ: 8.14 (1H, d, J = 5.8 Hz), 7.77 (1H, t, J = 7.6 Hz), 7.69-7.59 (3H, m), 7.49 (1H, d, J = 10.4 Hz), 7.15 (1H, d, J = 8.2 Hz), 5.38 (1H, s), 4.97 (2H, s), 4.63-4.57 (1H, m), 4.08-4.02 (1H, m), 2.33-1.47 (15H, m).
260		 isomer-X Relative configuration			free	605.2	¹ H-NMR (DMSO-D ₆) δ: 8.40 (1H, d, J = 5.8 Hz), 7.85-7.59 (5H, m), 7.47 (1H, d, J = 10.4 Hz), 7.15 (1H, d, J = 7.9 Hz), 5.77 (1H, s), 4.92 (2H, s), 4.50-3.69 (3H, m), 2.17-1.45 (11H, m), 0.90 (1H, dd, J = 12.1, 4.4 Hz).
261					free	579.3	¹ H-NMR (DMSO-D ₆) δ: 8.40 (1H, d, J = 5.2 Hz), 7.82-7.62 (5H, m), 7.44 (1H, d, J = 10.7 Hz), 7.14 (1H, d, J = 7.9 Hz), 5.77 (1H, s), 4.92 (2H, s), 3.63-3.39 (5H, m), 2.21-1.66 (8H, m).
262					free	619.3	¹ H-NMR (DMSO-D ₆) δ: 8.40 (1H, d, J = 6.1 Hz), 7.84-7.59 (5H, m), 7.46 (1H, d, J = 10.7 Hz), 7.15 (1H, dd, J = 8.1, 1.4 Hz), 5.78 (1H, s), 4.92 (2H, s), 4.65-4.54 (1H, m), 4.11-4.02 (1H, m), 2.29-1.47 (15H, m).
263		 Racemate Relative configuration			free	605.2	¹ H-NMR (DMSO-D ₆) δ: 8.40 (1H, d, J = 5.8 Hz), 7.85-7.59 (5H, m), 7.47 (1H, d, J = 10.4 Hz), 7.15 (1H, d, J = 7.9 Hz), 5.77 (1H, s), 4.92 (2H, s), 4.50-3.89 (3H, m), 2.17-1.45 (11H, m), 0.90 (1H, dd, J = 12.1, 4.4 Hz).
264		 isomer-X Relative configuration			free	580.2	¹ H-NMR (DMSO-D ₆) δ: 8.54 (1H, d, J = 6.7 Hz), 7.91-7.38 (5H, m), 7.19 (1H, dd, J = 7.9, 1.5 Hz), 5.56 (1H, s), 4.89 (2H, s), 4.47-3.88 (3H, m), 2.27-1.48 (11H, m), 0.96-0.85 (1H, m).
265					free	554.2	¹ H-NMR (DMSO-D ₆) δ: 8.55 (1H, d, J = 6.4 Hz), 7.80-7.68 (4H, m), 7.57-7.49 (1H, m), 7.18 (1H, dd, J = 8.1, 1.4 Hz), 5.55 (1H, s), 4.89 (2H, s), 3.67-3.46 (5H, m), 2.26-1.57 (8H, m).
















[0821]

Table 34

266					free	594.2	¹ H-NMR (DMSO-d ₆) δ: 8.55 (1H, d, J = 6.4 Hz), 7.78 (1H, t, J = 7.5 Hz), 7.70-7.60 (3H, m), 7.53 (1H, dd, J = 10.4, 1.5 Hz), 7.21-7.16 (1H, m), 5.55 (1H, s), 4.89 (2H, s), 4.51-4.55 (1H, m), 4.08-4.02 (1H, m), 2.31-1.50 (15H, m).
267		 Racemate Relative configuration			free	580.2	¹ H-NMR (DMSO-d ₆) δ: 8.54 (1H, d, J = 6.7 Hz), 7.91-7.38 (5H, m), 7.19 (1H, dd, J = 7.9, 1.5 Hz), 5.56 (1H, s), 4.89 (2H, s), 4.47-3.88 (3H, m), 2.27-1.48 (11H, m), 0.96-0.85 (1H, m).
268		 Racemate Relative configuration			free	573.2	¹ H-NMR (DMSO-d ₆) δ: 7.96 (1H, d, J = 5.5 Hz), 7.83-7.51 (4H, m), 7.50 (1H, d, J = 10.4 Hz), 7.17 (1H, d, J = 7.8 Hz), 5.42 (1H, s), 4.78 (2H, s), 4.49-3.87 (3H, m), 2.21-1.96 (6H, m), 1.82-1.50 (5H, m), 0.99-0.79 (1H, m).
269		 Racemate Relative configuration			free	589.2 591.4	¹ H-NMR (DMSO-d ₆) δ: 8.14 (1H, d, J = 5.8 Hz), 7.79-7.53 (4H, m), 7.50 (1H, d, J = 9.8 Hz), 7.16 (1H, d, J = 8.5 Hz), 5.38 (1H, s), 4.97 (2H, s), 4.49-3.87 (3H, m), 2.25-1.97 (6H, m), 1.76-1.50 (5H, m), 0.95-0.83 (1H, m).
270		 isomer-X Relative configuration			free	569.4	¹ H-NMR (DMSO-d ₆) δ: 7.86 (1H, d, J = 6.4 Hz), 7.79-7.57 (4H, m), 7.46 (1H, dd, J = 10.5, 1.4 Hz), 7.17-7.14 (1H, m), 5.52 (1H, s), 4.84 (2H, s), 4.45-3.89 (3H, m), 2.54 (3H, s), 2.30-1.96 (6H, m), 1.79-1.51 (5H, m), 0.89 (1H, dd, J = 11.9, 4.6 Hz).
271					free	543.2	¹ H-NMR (DMSO-d ₆) δ: 7.86 (1H, d, J = 6.4 Hz), 7.76-7.59 (2H, m), 7.68-7.63 (1H, m), 7.60-7.56 (1H, m), 7.44 (1H, d, J = 10.7 Hz), 7.13 (1H, dd, J = 8.2, 1.5 Hz), 5.52 (1H, s), 4.84 (2H, s), 3.68-3.49 (5H, m), 2.52 (3H, s), 2.30-2.24 (2H, m), 2.02-1.93 (3H, m), 1.75-1.60 (3H, m).
272					free	583.2	¹ H-NMR (DMSO-d ₆) δ: 7.87 (1H, d, J = 6.4 Hz), 7.75 (1H, t, J = 7.6 Hz), 7.63 (1H, dd, J = 7.8, 1.7 Hz), 7.60-7.53 (2H, m), 7.45 (1H, dd, J = 10.5, 1.4 Hz), 7.14 (1H, dd, J = 7.9, 1.5 Hz), 5.52 (1H, s), 4.84 (2H, s), 4.64-4.54 (1H, m), 4.11-4.01 (1H, m), 2.52 (3H, s), 2.28-1.50 (15H, m).
273		 isomer-X Relative configuration			free	621.1 623.1	¹ H-NMR (DMSO-d ₆) δ: 8.17 (1H, d, J = 6.1 Hz), 7.89-7.62 (4H, m), 7.47 (1H, d, J = 9.5 Hz), 7.15 (1H, d, J = 8.2 Hz), 4.88 (2H, s), 4.74 (1H, s), 4.49-3.86 (3H, m), 2.27-2.12 (2H, m), 1.79-1.50 (3H, m), 1.18 (6H, s), 0.89 (1H, dd, J = 12.4, 4.4 Hz).
274					free	595.1 597.1	¹ H-NMR (DMSO-d ₆) δ: 8.21-8.16 (1H, m), 7.78-7.65 (4H, m), 7.45 (1H, d, J = 10.7 Hz), 7.13 (1H, dd, J = 8.2, 1.5 Hz), 4.88 (2H, s), 4.74 (1H, s), 3.70-3.48 (5H, m), 2.03-1.94 (1H, m), 1.71-1.63 (1H, m), 1.18 (6H, s).
275					free	635.1 637.1	¹ H-NMR (DMSO-d ₆) δ: 8.18 (1H, d, J = 6.1 Hz), 7.76 (1H, t, J = 7.5 Hz), 7.68-7.59 (3H, m), 7.46 (1H, dd, J = 10.5, 1.4 Hz), 7.14 (1H, dd, J = 7.9, 1.5 Hz), 4.88 (2H, s), 4.74 (1H, s), 4.62-4.55 (1H, m), 4.10-4.02 (1H, m), 2.26-1.88 (5H, m), 1.65-1.47 (3H, m), 1.18 (6H, s).



















[0822]

Table 35

276		 Racemate Relative configuration			free	569.4	¹ H-NMR (DMSO-D ₆) δ: 7.86 (1H, d, J = 6.4 Hz), 7.79-7.57 (4H, m), 7.46 (1H, dd, J = 10.5, 1.4 Hz), 7.17-7.14 (1H, m), 5.52 (1H, s), 4.84 (2H, s), 4.45-3.89 (3H, m), 2.54 (3H, s), 2.30-1.96 (6H, m), 1.79-1.51 (5H, m), 0.89 (1H, dd, J = 11.9, 4.6 Hz).
277		 Racemate Relative configuration			free	521.1 523.1	¹ H-NMR (DMSO-D ₆) δ: 8.17 (1H, d, J = 6.1 Hz), 7.80-7.62 (4H, m), 7.47 (1H, d, J = 9.5 Hz), 7.15 (1H, d, J = 8.2 Hz), 4.88 (2H, s), 4.74 (1H, s), 4.49-3.86 (3H, m), 2.27-2.12 (2H, m), 1.79-1.50 (3H, m), 1.18 (6H, s), 0.89 (1H, dd, J = 12.4, 4.4 Hz).
278		 Racemate Relative configuration			free	583.4	¹ H-NMR (DMSO-D ₆) δ: 7.99 (1H, d, J = 5.8 Hz), 7.83-7.58 (4H, m), 7.38 (1H, dd, J = 10.5, 1.4 Hz), 7.09 (1H, dd, J = 8.2, 1.5 Hz), 5.10-4.83 (2H, m), 4.72 (1H, s), 4.53-3.88 (3H, m), 2.29-2.13 (2H, m), 1.80-1.50 (3H, m), 1.15 (6H, s), 1.02-0.96 (2H, m), 0.89 (1H, dd, J = 11.9, 4.6 Hz), 0.53-0.12 (2H, m).
279		 isomer-X Relative configuration			free	583.4	¹ H-NMR (DMSO-D ₆) δ: 7.99 (1H, d, J = 5.8 Hz), 7.83-7.58 (4H, m), 7.38 (1H, dd, J = 10.5, 1.4 Hz), 7.09 (1H, dd, J = 8.2, 1.5 Hz), 5.10-4.83 (2H, m), 4.72 (1H, s), 4.53-3.88 (3H, m), 2.29-2.13 (2H, m), 1.80-1.50 (3H, m), 1.15 (6H, s), 1.02-0.96 (2H, m), 0.89 (1H, dd, J = 11.9, 4.6 Hz), 0.53-0.12 (2H, m).
280					free	557.4	¹ H-NMR (DMSO-D ₆) δ: 7.99 (1H, d, J = 5.5 Hz), 7.76-7.64 (4H, m), 7.36 (1H, dd, J = 10.5, 2.9 Hz), 7.07 (1H, dd, J = 8.2, 1.5 Hz), 5.05-4.81 (2H, m), 4.72 (1H, s), 3.70-3.44 (5H, m), 2.26-2.17 (1H, m), 2.11-1.97 (1H, m), 1.78-1.67 (1H, m), 1.15 (6H, s), 1.03-0.95 (2H, m), 0.40-0.16 (2H, m).
281					free	597.5	¹ H-NMR (DMSO-D ₆) δ: 7.99 (1H, d, J = 6.1 Hz), 7.74 (1H, t, J = 7.5 Hz), 7.68-7.56 (3H, m), 7.37 (1H, d, J = 10.7 Hz), 7.08 (1H, d, J = 7.9 Hz), 5.04-4.81 (2H, m), 4.72 (1H, s), 4.65-4.57 (1H, m), 4.13-4.05 (1H, m), 2.24-1.51 (10H, m), 1.15 (6H, s), 1.02-0.94 (2H, m), 0.42-0.16 (2H, m).
282					free	368.2	¹ H-NMR (DMSO-D ₆) δ: 8.23-7.56 (5H, m), 7.37-7.28 (2H, m), 7.17-7.09 (2H, m), 7.02-6.89 (2H, m), 5.76 (2H, s), 4.65-3.85 (5H, m), 2.28 (3H, s), 1.99-1.64 (2H, m).
283		 isomer-X Relative configuration			free	430.1	¹ H-NMR (DMSO-D ₆) δ: 7.80 (1H, t, J = 7.3 Hz), 7.53 (1H, d, J = 7.9 Hz), 7.44 (1H, s), 7.36 (1H, d, J = 7.6 Hz), 7.29 (1H, d, J = 10.4 Hz), 7.18 (1H, t, J = 7.8 Hz), 7.09 (1H, d, J = 7.9 Hz), 7.03 (1H, d, J = 7.9 Hz), 6.93 (1H, d, J = 11.3 Hz), 3.62 (2H, s), 3.29-3.25 (1H, m), 3.15-3.10 (1H, m), 3.05-2.99 (1H, m), 2.30 (3H, s), 2.06-1.96 (2H, m), 1.89-1.76 (1H, m), 1.66-1.56 (1H, m), 1.39-1.29 (1H, m), 0.64 (1H, dd, J = 11.6, 4.0 Hz).










[0823]

Table 36

284				$1,2\text{-CH}_2\text{-}1,1$	free	418.2	¹ H-NMR (DMSO-D ₆) δ : 7.81 (1H, t, J = 7.5 Hz), 7.52 (1H, d, J = 7.6 Hz), 7.47 (1H, s), 7.39 (1H, d, J = 7.9 Hz), 7.31 (1H, d, J = 10.4 Hz), 7.18 (1H, t, J = 8.1 Hz), 7.09 (1H, d, J = 7.9 Hz), 7.03 (1H, d, J = 7.3 Hz), 6.93 (1H, d, J = 11.0 Hz), 3.70 (2H, dd, J = 37.4, 13.6 Hz), 2.89-2.78 (1H, m), 2.64-2.60 (2H, m), 2.38-2.34 (1H, m), 2.31 (3H, s), 1.86-1.73 (2H, m), 1.28 (3H, s).
285		isomer-X Relative configuration 		$1,2\text{-CH}_2\text{-}1,1$	2HCl	547.2	¹ H-NMR (DMSO-D ₆) δ : 7.94-7.72 (5H, m), 7.54-7.43 (1H, m), 7.22-7.09 (1H, m), 4.90-4.74 (1H, m), 4.63 (2H, s), 4.53-4.40 (1H, m), 4.29-3.98 (3H, m), 2.24-1.33 (5H, m), 1.16 (6H, s), 0.93-0.81 (1H, m).
286		isomer-X Relative configuration 		$1,2\text{-CH}_2\text{-}1,1$	2HCl	575.1 576.1	¹ H-NMR (DMSO-D ₆) δ : 8.09 (1H, d, J = 5.8 Hz), 7.91-7.51 (5H, m), 7.15 (1H, d, J = 8.2 Hz), 5.45-5.34 (1H, m), 4.98 (2H, s), 4.49-4.03 (5H, m), 2.34-2.21 (3H, m), 2.05-1.87 (3H, m), 1.79-1.49 (3H, m), 1.28-1.23 (1H, m), 1.04 (1H, d, J = 6.1 Hz), 0.92-0.81 (1H, m).
287		isomer-X Relative configuration 		$1,2\text{-CH}_2\text{-}1,1$	2HCl	528.2	¹ H-NMR (DMSO-D ₆) δ : 8.16-8.08 (1H, m), 8.01-7.31 (8H, m), 7.19-6.98 (1H, m), 4.77-4.39 (3H, m), 4.30-3.96 (6H, m), 2.13-1.39 (4H, m), 1.11 (6H, s).
288		isomer-X Relative configuration 		$1,2\text{-CH}_2\text{-}1,1$	2HCl	591.1 592.0	¹ H-NMR (DMSO-D ₆) δ : 8.10 (1H, s), 7.97-7.36 (5H, m), 7.18-7.13 (1H, m), 4.82 (2H, s), 4.52-3.95 (6H, m), 2.33-1.37 (9H, m), 0.90 (6H, t, J = 7.3 Hz).
289		isomer-X Relative configuration 		$1,2\text{-CH}_2\text{-}1,1$	2HCl	527.1	¹ H-NMR (DMSO-D ₆) δ : 7.91-7.75 (3H, m), 7.64-7.59 (1H, m), 7.53-7.31 (4H, m), 7.17-7.11 (1H, m), 6.48-6.43 (1H, m), 4.70-3.94 (9H, m), 2.18-1.50 (4H, m), 1.06 (6H, s).
290		isomer-X Relative configuration 		$1,2\text{-CH}_2\text{-}1,1$	2HCl	488.2	¹ H-NMR (DMSO-D ₆) δ : 7.97-7.80 (3H, m), 7.66-7.54 (1H, m), 7.35-6.93 (5H, m), 4.53-3.92 (7H, m), 2.32-1.48 (6H, m), 1.05 (6H, s).
291				$1,2\text{-CH}_2\text{-}1,1$	2HCl	462.2	¹ H-NMR (DMSO-D ₆) δ : 7.72 (4H, t, J = 62.6 Hz), 7.12 (5H, ddd, J = 96.0, 48.2, 18.5 Hz), 4.58-3.75 (6H, m), 2.66 (2H, s), 2.30-1.99 (2H, m), 1.29-1.16 (1H, m), 1.05 (6H, s).
292		isomer-X Relative configuration 		$1,2\text{-CH}_2\text{-}1,1$	2HCl	444.1	¹ H-NMR (DMSO-D ₆) δ : 7.97-6.93 (9H, m), 4.63-4.49 (1H, m), 4.23-3.86 (3H, m), 2.32 (3H, s), 2.21-1.97 (2H, m), 1.83-1.64 (3H, m), 1.31-1.24 (2H, m), 1.07-0.82 (2H, m).

[0824]

Table 37

293		 isomer-X Relative configuration		//-CH2-//	2HCl	575.1	1H-NMR (DMSO-D6) δ : 7.94-7.49 (6H, m), 7.18-7.11 (1H, m), 4.63 (2H, s), 4.52-4.40 (2H, m), 4.19-3.70 (4H, m), 2.32-1.66 (5H, m), 1.44-1.35 (4H, m), 0.90 (6H, t, J = 7.3 Hz).
294		 isomer-X Relative configuration		//-CH2-//	2HCl	607.0 609.0	1H-NMR (DMSO-D6) δ : 8.13 (1H, d, J = 5.8 Hz), 8.00-7.71 (4H, m), 7.50-7.44 (1H, m), 7.19-7.10 (1H, m), 4.88 (2H, s), 4.75 (1H, s), 4.51-3.98 (5H, m), 2.30-1.64 (5H, m), 1.18 (6H, s), 0.92-0.82 (1H, m).
295		 isomer-X Relative configuration		//-CH2-//	2HCl	635.1 637.0	1H-NMR (DMSO-D6) δ : 8.13 (1H, d, J = 5.8 Hz), 7.96-7.69 (4H, m), 7.52-7.46 (1H, m), 7.14 (1H, d, J = 7.9 Hz), 4.89 (2H, s), 4.48-4.01 (5H, m), 2.14-1.24 (11H, m), 0.91 (6H, t, J = 7.5 Hz).

[0825]

Test Example 1: Measurement of LSD1 Inhibitory Activity (in vitro)

The conditions for measuring inhibitory activity of compounds against LSD1 activity were determined with reference to a document available from the website of PerkinElmer (U-TRF #38) and a patent of GlaxoSmithKline (WO2012135113).

[0826]

To measure the inhibitory activity, first, the Example compound was serially diluted in dimethylsulfoxide (DMSO). Sequentially, the serially diluted solution of the Example compound in DMSO (final concentration of DMSO: 5%) and human LSD1 protein (Abcam, ab80379) were added to a reaction buffer (25 mM Tris-HCl (pH 7.5), 50 mM KCl, 2 mM CHAPS, 1 mM DTT, 0.02% BSA). The mixture was preincubated at 25°C for 30 minutes. Thereafter, a H3K4 (Me1)-biotin-labeled peptide (Anaspec #64355) (final concentration: 200 nM) was added thereto and reacted for 60 minutes. Tranylcypromine (final concentration: 3 mM) was then added thereto to terminate the reaction. Thereafter, a detection solution containing an Eu-labeled anti-H3K4 antibody (PerkinElmer, TRF0404) and Streptavidin Alexa Fluor 647 (Thermo Fisher Scientific, S21374) was added thereto, and the mixture was allowed to stand at room temperature for 1 hour. Finally, the intensity of fluorescence under the excitation light with a

wavelength of 337 nm was measured with a PHERAstar FS (BMG Labtech) at two wavelengths: 620 nm and 665 nm. The demethylation level was calculated from the ratio of the fluorescence intensity at the two wavelengths, and the compound concentration at which
5 demethylation was inhibited by 50% was defined as IC₅₀ (nM). The following tables show the results.
[0827]

Table 38

Example No.	LSD1 Inhibitory activity IC50 (nM)	Example No.	LSD1 Inhibitory activity IC50 (nM)	Example No.	LSD1 Inhibitory activity IC50 (nM)	Example No.	LSD1 Inhibitory activity IC50 (nM)	Example No.	LSD1 Inhibitory activity IC50 (nM)
1	7.02	42	11.6	85	6.27	127	17.0	174	13.8
3	10.6	45	7.71	86	6.58	128	0.62	175	0.33
4	4.93	46	19.7	87	5.12	130	0.29	176	0.30
5	1.59	47	9.15	89	11.2	131	1.81	177	0.23
6	10.7	48	17.9	90	14.0	132	10.8	178	1.24
7	0.78	49	5.51	91	4.85	133	0.91	179	0.69
8	5.92	51	11.0	92	2.78	134	0.21	180	0.88
9	4.45	52	8.31	93	20.3	135	0.30	181	2.70
10	3.74	53	10.9	94	5.10	136	0.30	182	17.8
11	17.4	54	4.58	95	0.75	137	3.18	183	0.67
12	5.83	55	20.0	96	0.49	138	2.83	184	0.25
13	14.9	56	11.3	97	10.1	139	1.68	185	0.14
14	11.0	57	3.42	98	2.18	140	5.98	186	0.43
15	2.41	58	13.5	100	3.13	141	0.25	187	0.25
16	1.97	59	6.78	101	3.70	142	0.38	188	0.61
17	3.51	60	13.8	102	0.63	143	0.25	189	0.33
18	0.40	61	15.5	103	9.65	144	0.51	190	0.16
19	2.67	62	6.34	104	0.44	145	0.38	191	0.22
20	1.31	63	3.48	105	0.51	146	0.42	192	0.40
21	0.87	64	6.69	106	0.34	147	0.35	193	2.35
22	1.18	65	2.80	107	1.05	148	19.8	194	0.34
23	0.18	66	6.99	108	0.21	150	2.99	195	0.20
24	0.29	67	10.4	109	0.28	151	5.02	196	0.14
25	1.83	68	17.3	110	0.47	155	0.51	197	0.11
26	0.45	69	11.1	111	9.45	156	12.8	198	0.18
27	11.4	70	3.68	112	1.80	158	6.35	199	1.00
28	0.57	71	3.04	113	1.56	159	17.7	200	0.91
29	19.8	72	7.57	114	1.42	160	0.92	201	0.40
30	4.31	73	9.23	115	2.81	161	0.74	202	0.09
31	16.0	74	1.53	116	3.03	162	1.09	203	0.24
32	17.7	75	1.54	117	1.97	164	12.8	204	0.45
33	11.0	76	2.09	118	1.57	165	11.2	205	1.29
34	15.1	77	1.54	119	1.98	166	0.63	206	0.30
35	10.0	78	9.65	120	0.90	167	1.79	207	6.89
36	1.04	79	17.9	121	9.46	168	1.05	208	16.8
37	2.40	80	12.8	122	8.61	169	0.90	209	0.06
38	16.0	81	3.30	123	0.12	170	0.72	210	0.16
39	1.91	82	3.04	124	0.36	171	0.35	211	0.14
40	5.96	83	9.29	125	0.29	172	0.55	213	0.15
41	1.57	84	9.33	126	0.41	173	1.84	214	0.10

[0828]

Table 39

Example No.	LSD1 Inhibitory activity IC50 (nM)	Example No.	LSD1 Inhibitory activity IC50 (nM)	Example No.	LSD1 Inhibitory activity IC50 (nM)	Example No.	LSD1 Inhibitory activity IC50 (nM)	Example No.	LSD1 Inhibitory activity IC50 (nM)
215	0.25	233	4.88	251	0.96	269	0.41	286	0.093
216	0.50	234	2.43	252	0.36	270	0.15	287	0.10
217	0.21	235	2.09	254	0.13	271	0.23	288	0.075
218	0.07	237	0.20	255	0.23	272	0.32	289	0.18
219	0.10	238	2.35	256	0.24	273	0.19	290	0.23
220	0.12	239	0.59	257	0.12	274	0.32	291	0.80
221	0.20	240	0.34	258	0.13	275	0.21	292	0.36
222	8.31	241	0.29	259	0.14	276	0.39	293	0.076
224	0.49	242	2.69	260	0.14	277	0.42	294	0.14
225	0.20	243	0.65	261	0.37	278	0.87	295	0.15
226	0.54	244	6.54	262	0.30	279	0.46		
227	11.7	245	0.39	263	0.52	280	2.13		
228	0.11	246	0.17	264	0.24	281	2.62		
229	0.43	247	0.23	265	0.43	282	3.51		
230	0.94	248	0.46	266	0.35	283	0.062		
231	0.65	249	0.88	267	0.49	284	0.74		
232	0.63	250	0.38	268	0.29	285	0.091		

[0829]

The results of the test clarified that the biphenyl compound represented by Formula (I) exhibits LSD1 inhibitory activity.

[0830]

Test Example 2: Cell-Growth Inhibition Test

Under the following conditions, an in vitro cell-growth inhibition test was performed with respect to HEL cells (human acute myelocytic leukemia cell lines), NCI-H1417 cells (human small-cell lung cancer cell lines), and NCI-H146 cells (human small-cell lung cancer cell lines).

[0831]

HEL cells (JCRB, Cat#: JCRB0062), NCI-H1417 cells (ATCC, Cat#: CRL-5869), or NCI-H146 cells (ATCC, Cat#: HTB-173) cultured in a 10% FBS-containing RPMI1640 medium (Thermo Fisher Scientific, Cat#: A10491-01) were seeded in a 96-well flat-bottom microplate (Thermo Fisher Scientific, Cat#: 165305) so that each well contained 1500 HEL cells (100 μ L), 5000 NCI-H1417 cells (100 μ L), or 1200 NCI-H146 cells (100 μ L). The Example compound was serially diluted in dimethylsulfoxide to a concentration that was 500 times higher than the final concentration. The serially

diluted Example compound or dimethylsulfoxide alone was added to a 10% FBS-containing RPMI1640 medium to a concentration that was 2 times higher than the final concentration, and the resulting product was added in an amount of 100 μ L to each well of the culture plate containing HEL cells, NCI-H1417 cells, or NCI-H146 cells, so that the final concentrations of the Example compound were 3000, 1000, 300, 100, 30, 10, 3, 1, 0.3, 0.1, 0.03, and 0.01 nM. The final concentration of dimethylsulfoxide was adjusted to 0.2%. The cells with the Example compound or with dimethylsulfoxide alone were cultured at 37°C in a 5% carbon-dioxide-containing incubator for 5 days (HEL cells) or 10 days (NCI-H1417 cells and NCI-H146 cells). After culture, the plate was allowed to stand at room temperature for 30 minutes, and 100 μ L of the supernatant was removed from each well to leave 100 μ L of the cell culture solution. To each well containing the remaining 100 μ L of the cell culture solution, the same amount of CellTiter-Glo 2.0 Assay (Promega, Cat#: G9242) was added. The microplate was shaken with a plate mixer for 1 minute and then allowed to stand in a dark place for 10 minutes. Thereafter, the luminescence intensity of viable cells in each well was measured using a microplate reader (PerkinElmer, EnSpire). The cell growth rate was determined in accordance with the following equation, and the concentration at which the cell growth rate was 50%, i.e.; the concentration of each test compound at which the cell growth was inhibited by 50% (IC₅₀ (nM)) was determined.

[0832]

Cell growth rate (%) = $T/C \times 100$

T: The luminescence intensity in a well to which the test compound was added (count per second)

C: The luminescence intensity in a well to which dimethylsulfoxide alone was added (count per second)

[0833]

The tables below show the results.

[0834]

Table 40

Cell-Growth Inhibition Test: HEL Cells

Ex.No.	IC50 (nM)	Ex.No.	IC50 (nM)	Ex.No.	IC50 (nM)	Ex.No.	IC50 (nM)
1	39.8	49	32.1	125	7.57	177	0.38
4	28.8	59	33.0	128	5.39	178	5.57
5	34.5	63	26.3	128	5.70	179	1.52
7	14.8	85	23.9	130	22.2	180	1.18
8	42.7	69	29.9	131	15.2	181	9.06
9	38.7	71	19.5	133	17.7	182	22.4
10	31.7	74	20.4	134	1.06	183	0.54
12	36.7	75	5.78	135	2.37	184	2.28
15	17.3	77	12.8	138	0.90	185	0.88
16	34.3	80	43.6	139	16.9	186	5.30
18	3.55	81	11.7	141	1.50	187	1.99
19	24.4	82	11.1	142	3.86	188	1.95
20	7.73	84	41.7	143	0.96	189	0.69
21	8.34	85	32.2	144	6.37	190	0.41
22	3.51	86	13.4	145	1.61	191	3.91
23	0.73	87	17.4	146	1.15	192	1.11
24	3.56	92	9.81	147	3.04	193	2.31
25	4.72	94	25.4	155	0.83	194	15.5
26	4.83	96	11.0	160	7.03	195	0.45
27	24.7	102	12.8	161	6.88	196	0.33
28	15.7	104	28.8	162	17.7	197	1.0
30	27.1	108	5.74	168	0.92	198	0.43
36	15.4	109	36.4	167	4.05	199	22.0
37	10.1	109	15.1	168	5.29	200	3.72
38	34.0	110	7.41	170	3.44	201	1.55
39	17.4	113	9.00	171	0.060	202	0.24
40	27.2	117	8.54	172	0.47	203	2.48
41	4.98	120	7.60	173	1.85	204	2.14
47	29.3	123	1.39	175	0.20	205	4.95
48	28.9	124	2.84	178	0.20	208	0.40

[0835]

Table 41

5 Cell-Growth Inhibition Test: HEL Cells

Ex.No.	IC50 (nM)	Ex.No.	IC50 (nM)	Ex.No.	IC50 (nM)	Ex.No.	IC50 (nM)
209	0.24	232	2.04	258	1.27	278	3.76
210	3.30	234	42.6	259	0.58	279	1.59
211	1.41	235	8.42	260	0.13	280	36.6
213	0.65	237	28.5	261	2.67	281	23.9
214	0.44	238	26.1	262	0.93	282	40.7
215	4.91	239	12.5	263	0.39	283	0.79
216	4.19	240	4.20	264	0.84	284	5.78
217	0.86	241	2.54	265	17.6	285	0.046
218	0.22	246	0.37	266	4.63	286	0.036
219	1.70	247	3.02	267	1.98	287	0.045
220	1.09	248	6.94	268	0.40	288	0.062
221	0.49	249	41.6	269	0.29	289	0.038
222	38.0	250	2.36	270	0.26	290	0.18
224	46.2	251	23.9	271	4.29		
225	0.69	252	0.77	272	1.52		
226	2.56	253	6.11	273	0.27		
227	23.6	254	0.20	274	3.15		
228	0.68	255	2.73	275	0.99		
229	2.55	256	1.31	276	0.62		
231	13.4	257	0.13	277	0.65		

[0836]

Table 42

Cell-Growth Inhibition Test: NCI-H1417 Cells

Ex.No.	IC50 (nM)	Ex.No.	IC50 (nM)	Ex.No.	IC50 (nM)	Ex.No.	IC50 (nM)
7	28.0	102	4.99	147	1.66	193	6.66
18	4.73	104	49.8	155	2.80	194	5.63
20	17.4	105	11.7	160	9.20	195	0.79
21	32.8	106	6.15	161	2.49	196	0.78
22	8.20	107	18.2	162	4.69	197	0.92
23	0.99	108	3.99	166	1.42	198	0.61
24	0.97	109	1.81	167	4.45	199	4.55
25	13.4	110	3.54	168	7.29	200	1.23
26	3.15	112	20.5	169	8.20	201	2.23
27	22.5	113	13.7	170	2.51	202	0.62
28	4.40	114	30.6	171	0.15	203	2.14
37	19.2	117	38.1	172	0.70	204	1.99
39	22.7	118	29.5	173	3.81	205	6.25
41	6.12	119	23.8	175	0.60	206	0.95
65	47.6	120	15.7	176	0.27	209	0.18
74	20.7	123	3.57	177	0.24	210	1.27
75	14.1	124	9.02	178	3.03	211	0.79
76	43.3	125	8.43	179	3.36	213	0.57
77	16.5	126	5.02	180	3.12	214	0.47
81	26.7	130	8.25	181	4.60	215	2.67
82	26.6	131	30.9	183	1.01	216	4.40
86	26.4	133	17.9	184	2.73	217	1.10
87	49.9	134	6.93	185	1.43	218	0.21
88	40.5	135	9.08	186	5.95	219	1.11
91	30.3	136	2.39	187	1.52	220	1.03
92	26.8	141	2.95	188	1.96	221	0.64
95	11.8	142	3.79	189	0.71	222	41.8
96	7.14	143	4.88	190	0.55	225	1.99
100	42.8	145	3.29	191	1.98	226	5.36
101	53.2	146	1.42	192	0.75	227	26.5

[0837]

5 Table 43

Cell-Growth Inhibition Test: NCI-H1417 Cells

Ex.No.	IC50 (nM)	Ex.No.	IC50 (nM)	Ex.No.	IC50 (nM)	Ex.No.	IC50 (nM)
228	0.30	247	1.4	262	0.70	277	0.53
229	1.30	248	6.06	263	0.38	278	2.68
230	16.4	249	16.1	264	0.30	279	0.90
231	7.11	250	1.2	265	4.40	280	9.71
232	1.55	251	2.7	266	1.85	281	8.50
233	36.5	252	0.5	267	0.86	283	2.10
234	14.8	253	1.47	268	0.30	284	13.0
235	6.99	254	0.18	269	0.27	285	0.090
236	11.0	255	1.52	270	0.18	286	0.083
237	6.63	256	0.94	271	1.42	287	0.10
238	10.8	257	0.12	272	0.81	288	0.22
239	3.8	258	0.65	273	0.19	289	0.094
240	2.55	259	0.46	274	1.25	290	1.06
241	2.89	260	0.13	275	0.59		
246	0.81	261	1.30	276	0.50		

[0838]

Table 44

Cell-Growth Inhibition Test: NCI-H146 Cells

Ex.No.	IC50 (nM)	Ex.No.	IC50 (nM)	Ex.No.	IC50 (nM)	Ex.No.	IC50 (nM)
18	8.13	193	5.33	233	41.4	267	3.14
22	15.0	194	11.3	234	19.1	268	0.94
23	9.33	195	1.26	235	8.12	269	0.53
24	4.04	196	0.82	236	28.5	270	0.37
37	10.1	197	1.62	237	5.14	271	3.08
41	9.76	198	1.10	238	17.7	272	3.17
123	1.39	199	5.45	239	5.39	273	0.63
146	3.15	200	2.31	240	4.81	274	2.23
161	1.09	201	3.25	241	2.89	275	1.78
166	2.37	202	0.73	246	1.32	276	0.89
171	0.19	203	4.73	247	3.84	277	1.81
172	0.52	205	15.1	248	10.3	278	6.18
175	0.16	206	1.09	249	24.7	279	3.36
176	0.27	209	0.22	250	9.30	280	16.8
177	0.11	210	4.26	251	13.2	281	32.3
178	1.22	211	1.40	252	1.48	283	11.1
179	1.05	213	0.87	253	7.59	284	21.9
180	0.94	214	2.58	254	0.36	285	0.49
181	2.93	215	6.99	255	4.41	286	0.17
182	13.2	216	5.02	256	2.54	287	0.20
183	0.25	217	1.32	257	0.39	288	1.29
184	2.51	218	0.33	258	1.83	289	0.97
185	1.32	219	1.45	259	2.08	290	2.12
186	3.21	220	1.81	260	0.48		
187	1.06	221	0.94	261	3.35		
188	1.58	225	5.47	262	3.1		
189	0.33	226	10.1	263	0.62		
190	1.17	228	0.51	264	0.96		
191	3.84	229	1.26	265	24.6		
192	0.83	231	16.7	266	17.5		

[0839]

5 The results of this test revealed that the biphenyl compound represented by Formula (I) exhibits in vitro cell growth inhibitory effects, and that this compound not only inhibits the activity of recombinant human LSD1 protein but also inhibits cancer cell growth, suggesting that this compound is useful as an antitumor agent.

[0840]

Test Example 3: Antitumor Effect Test Using NCI-H146 Cells (Human Small-Cell Lung Cancer Cell Lines)

15 NCI-H146 cells, 3.5×10^6 cells (100 μ L), were subcutaneously implanted into BALB/cAJcl-nu/nu mice, and mice with a tumor volume within a range of 100 to 300 mm³ were divided into groups so that the groups had a uniform average tumor volume. To 5 mice in each group, a vehicle (0.5% hydroxymethylpropylcellulose containing 0.1 N HCL) or each

20 Example compound was orally administered. The administration was

performed once a day for 21 consecutive days (Example compound 41) or 28 consecutive days (Example compounds 37, 161, 166, 175, 176, and 177). The major axis and the minor axis of each tumor were measured twice a week with an electric caliper to calculate the tumor volume (TV). According to the tumor volumes thus obtained, a relative tumor volume (RTV) and a relative tumor volume change (T/C (%)) were calculated. The TV, RTV, and T/C (%) were calculated using the following equations.

[0841]

10
$$\text{Tumor volume TV (mm}^3\text{)} = (\text{major axis, mm}) \times (\text{minor axis, mm}) \times (\text{minor axis, mm}) / 2$$

$$\text{Relative tumor volume RTV} = \text{TV} / (\text{TV on the grouping day})$$

15
$$\text{T/C (\%)} = (\text{average RTV of administration group}) / (\text{average RTV of vehicle administration group}) \times 100.$$

[0842]

The table below shows the results.

[0843]

20 Table 45

Example compound No.	Dose (mg/kg)	T/C (%)
37	50	22
41	25	19
161	40	14
166	20	19
175	2	41
176	20	27
177	10	22

[0844]

The final measurement day was the day following the final administration day. The biphenyl compound represented by Formula (I) showed an antitumor effect on the above models for efficacy evaluation, and the percentage of body weight reduction on the final measurement day was less than 20% of the body weight before administration (day 0).

[0845]

The results revealed that the biphenyl compound represented by Formula (I) or a salt thereof exhibits excellent LSD1 inhibitory activity, shows a cancer cell growth inhibitory effect, has low toxicity, and is orally administrable. Therefore, the biphenyl compound represented by Formula (I) or a salt thereof is useful as an agent for preventing and/or treating cancer.

[0846]

Test Example 4: Combination of Biphenyl Compounds and Other Antitumor agents Using Human AML Cell Line or Human Myelodysplastic Syndrome (MDS) Cell Line (in vitro)

The following tables show the source of reagents, the source of tumor cell lines, the media used, and the number of seeded cells.

[0847]

Table 46

Reagent	Source
Fetal bovine serum (FBS)	Biological Industries
RPMI-1640 medium	Wako Pure Chemical Industries, Ltd.
RPMI-1640 medium (ATCC-formulated medium: containing L-glutamine, HEPES, sodium pyruvate)	Wako Pure Chemical Industries, Ltd.
Human recombinant IL-3	Peptrotech
2-Mercaptoethanol (2-ME)	SIGMA
Cytarabine	SIGMA-ALDRICH
Decitabine	SIGMA-ALDRICH
Azacitidine	SIGMA-ALDRICH
ATRA	KareBay Biochem
Daunorubicin	CALBIOCHEM

[0848]

Table 47

Tumor cell line (origin)	Source of cell line	Medium	Number of seeded cells per well (cells)
Kasumi-1 (human AML cell line)	ATCC	RPMI-1640 medium containing 20% FBS (ATCC-formulated medium)	1000
MDS-L (human MDS cell line)	Dr. Kaoru Tohyama, Kawasaki Medical University	RPMI-1640 medium containing 10% FBS, 10 ng/mL IL-3, and 50 μ M 2-ME	100

[0849]

Cells were seeded at 20 μ L/well in a 384-well culture plate (Corning). The plate seeded with cells was cultured in an incubator set at 37°C and 5% CO₂. On the day of seeding, drug solutions containing a combination of one of the biphenyl compounds and one of other compounds having an antitumor effect mixed at various concentrations were each added to the cells. Specifically, using an RPMI-1640 medium containing 20% FBS (ATCC-formulated medium) or an RPMI-1640 medium containing 10% FBS, 10 ng/mL IL-3, and 50 μ M 2-ME, ten-step serial dilutions (including 0 nM) of the biphenyl compounds (Example compounds 37, 41, 166, 177, 214, 273, and 285) were prepared, and eight-step serial dilutions (including 0 nM) of other compounds having an antitumor effect were prepared. Solutions in which the same amount of serial dilution of each compound was dispensed were mixed at a ratio of 1:1, and a total of 80 mixed drug solutions were prepared with brute-force combinations. The table below shows the maximum concentration (expressed as the final concentration) and geometric ratio of each compound. Each of these mixed solutions was added to the plate at 5 μ L/well (N = 4), and further cultured at 37°C under 5% CO₂ conditions for 10 days. Ten days later, CellTiter-Glo 2.0 Assay (Promega, Cat#: G9242) was added at 25 μ L/well, and chemiluminescence was measured with a plate reader (EnSpire (registered trademark) Multimode Plate Reader, PerkinElmer Japan Co., Ltd.).

[0850]

Table 48

Other compound having antitumor effect			Biphenyl compound			Tumor cell line
Name	Maximum concentration (nM)	Geometric ratio	Name	Maximum concentration (nM)	Geometric ratio	
Cytarabine	200	2	Example compound 37	100	2	Kasumi-1, MDS-L
			Example compound 166	10	2	
			Example compound 177	1	2	
			Example compound 273	1	2	
Decitabine	200	2	Example compound 37	100	2	Kasumi-1, MDS-L
			Example compound 166	10	2	
			Example compound 177	1	2	
			Example compound 273	1	2	
			Example compound 41	100	2	Kasumi-1
			Example compound 214	1	2	
			Example compound 285	0.1	2	
Azacitidine	10000	2	Example compound 37	100	2	Kasumi-1, MDS-L
			Example compound 166	10	2	
			Example compound 177	1	2	
			Example compound 273	1	2	
Tretinoin (ATRA)	10000	5	Example compound 37	100	2	Kasumi-1
			Example compound 166	10	2	
			Example compound 177	1	2	
			Example compound 273	1	2	
Daunorubicin	10	2	Example compound 37	100	2	Kasumi-1
			Example compound 166	10	2	
			Example compound 177	1	2	
			Example compound 273	1	2	

[0851]

The effect potentiated by the combined use of drugs was evaluated according to the methods described in known documents (Trends Pharmacol. Sci., 4, 450-454, 1983; Pharmacol Rev., 58 (3), 621-81, 2006).

Specifically, normalized cell viability T/C (treatment/control ratio, %) was determined by comparing the cells treated with each single agent of the biphenyl compounds and other compounds having an antitumor effect with control cells. From the obtained T/C values, the half maximal inhibitory concentration (IC50) of each drug was determined by a curve fitting regression analysis tool using a regression analysis tool XLfit 5.3 (IDBS Ltd.).

The IC50 ratio of each single agent was calculated by the following equation.

IC50 ratio = [IC50 of biphenyl compound] : [IC50 of other compound having antitumor effect].

5 [0852]

Regarding the concentration ratio combination of two compounds actually used in the experiment, the half maximal inhibitory concentration (ED50) when both compounds were used in combination was determined using their T/C. The combination index (CI) and fraction affected (Fa) in ED50 were determined. For the calculation of CI, median effect analysis software CalcuSyn 2.0 (CalcuSyn, Inc.) calculating the median-effect method was used.

10 [0853]

The effect of combined use was determined as shown in the following table (Pharmacol Rev., 58 (3), pp. 621-81, 2006).

15 [0854]

Table 49

Range of CI (upper limit)	Explanation
0.1	Very strong synergism
0.3	Strong synergism
0.7	Synergism
0.85	Moderate synergism
0.9	Slight synergism
1.0	Almost additive
1.2	Slight antagonism
1.45	Moderate antagonism
3.3	Antagonism
10	Strong antagonism
>10	Very strong antagonism

[0855]

The following tables show the results.

20 [0856]

Table 50

Other antitumor agent : cytarabine, cell line : Kasumi-1				
Example compound 37 (nM)	Other antitumor agent (nM)	Fa	CI	Combination ratio (Example compound 37 : other antitumor agent)
1.563	50	0.971	0.872	1 : 32
1.563	100	0.994	0.875	1 : 64
3.125	25	0.886	0.852	1 : 8
3.125	50	0.971	0.874	1 : 16
6.25	6.25	0.732	0.795	1 : 1
12.5	6.25	0.794	0.890	1 : 0.5
12.5	12.5	0.855	0.799	1 : 1
12.5	25	0.936	0.720	1 : 2
12.5	50	0.984	0.689	1 : 4
12.5	100	0.995	0.832	1 : 8
25	12.5	0.886	0.892	1 : 0.5
25	25	0.960	0.608	1 : 1
25	50	0.982	0.747	1 : 2
50	25	0.957	0.761	1 : 0.5
50	50	0.984	0.731	1 : 1
50	100	0.994	0.891	1 : 2
100	50	0.985	0.775	1 : 0.5
100	100	0.995	0.893	1 : 1

[0857]

Table 51

Other antitumor agent : cytarabine, cell line : Kasumi-1				
Example compound 166 (nM)	Other antitumor agent (nM)	Fa	CI	Combination ratio (Example compound 166 : other antitumor agent)
0.1563	50	0.968	0.894	1 : 320
0.3125	50	0.968	0.896	1 : 160
0.3125	100	0.994	0.891	1 : 320
0.625	25	0.904	0.805	1 : 40
0.625	50	0.978	0.776	1 : 80
1.25	25	0.955	0.589	1 : 20
1.25	50	0.987	0.632	1 : 40
2.5	12.5	0.894	0.857	1 : 5
2.5	25	0.942	0.779	1 : 10
2.5	50	0.981	0.763	1 : 20
5	25	0.961	0.740	1 : 5
5	50	0.988	0.668	1 : 10
5	100	0.997	0.738	1 : 20
5	200	0.999	0.900	1 : 40
10	50	0.986	0.776	1 : 5
10	100	0.996	0.786	1 : 10

[0858]

Table 52

Other antitumor agent : cytarabine, cell line : Kasumi-1				
Example compound 177 (nM)	Other antitumor agent (nM)	Fa	CI	Combination ratio (Example compound 177 : other antitumor agent)
0.00781	200	0.999	0.823	1 : 25600
0.01563	200	0.999	0.823	1 : 12800
0.03125	100	0.996	0.794	1 : 3200
0.0625	25	0.951	0.699	1 : 400
0.0625	50	0.983	0.800	1 : 800
0.0625	100	0.996	0.795	1 : 1600
0.0625	200	0.999	0.824	1 : 3200
0.125	3.125	0.788	0.693	1 : 25
0.125	6.25	0.787	0.877	1 : 50
0.125	25	0.952	0.730	1 : 200
0.125	50	0.984	0.787	1 : 400
0.125	100	0.995	0.887	1 : 800
0.25	12.5	0.894	0.888	1 : 50
0.25	25	0.954	0.778	1 : 100
0.25	50	0.986	0.748	1 : 200
0.25	100	0.996	0.801	1 : 400
0.25	200	0.999	0.825	1 : 800
0.5	12.5	0.927	0.900	1 : 25
0.5	25	0.969	0.703	1 : 50
0.5	50	0.992	0.605	1 : 100
0.5	100	0.996	0.819	1 : 200
0.5	200	0.999	0.826	1 : 400
1	25	0.974	0.769	1 : 25
1	50	0.990	0.705	1 : 50
1	100	0.997	0.726	1 : 100

[0859]

Table 53

Other antitumor agent : cytarabine, cell line : Kasumi-1				
Example compound 273 (nM)	Other antitumor agent (nM)	Fa	CI	Combination ratio (Example compound 273 : other antitumor agent)
0.00391	100	0.993	0.794	1 : 25600
0.00781	200	0.999	0.685	1 : 25600
0.01563	50	0.968	0.782	1 : 3200
0.01563	100	0.993	0.810	1 : 6400
0.01563	200	0.999	0.685	1 : 12800
0.03125	25	0.859	0.827	1 : 800
0.03125	50	0.968	0.785	1 : 1600
0.03125	100	0.991	0.897	1 : 3200
0.0625	25	0.878	0.817	1 : 400
0.0625	50	0.960	0.887	1 : 800
0.0625	100	0.991	0.893	1 : 1600
0.125	25	0.925	0.692	1 : 200
0.125	50	0.979	0.679	1 : 400
0.125	100	0.993	0.808	1 : 800
0.25	100	0.993	0.822	1 : 400
0.5	25	0.954	0.774	1 : 50
0.5	50	0.984	0.684	1 : 100
0.5	100	0.995	0.761	1 : 200
1	50	0.984	0.813	1 : 50
1	100	0.995	0.808	1 : 100
1	200	0.999	0.819	1 : 200

5

[0860]

Table 54

Other antitumor agent : cytarabine, cell line : MDS-L				
Example compound 37 (nM)	Other antitumor agent (nM)	Fa	CI	Combination ratio (Example compound 37 : other antitumor agent)
0.391	3.125	0.201	0.816	1 : 8
0.391	6.25	0.467	0.696	1 : 16
0.781	3.125	0.264	0.730	1 : 4
0.781	6.25	0.455	0.734	1 : 8
1.563	3.125	0.357	0.611	1 : 2
1.563	6.25	0.416	0.863	1 : 4
3.125	3.125	0.409	0.631	1 : 1
12.5	200	0.999	0.834	1 : 16
25	200	0.999	0.834	1 : 8
50	200	0.999	0.834	1 : 4
100	200	0.999	0.872	1 : 2

[0861]

Table 55

Other antitumor agent : cytarabine, cell line : MDS-L				
Example compound 166 (nM)	Other antitumor agent (nM)	Fa	CI	Combination ratio (Example compound 166 : other antitumor agent)
0.0391	3.125	0.220	0.721	1 : 80
0.0391	6.25	0.452	0.712	1 : 160
0.0781	3.125	0.252	0.746	1 : 40
0.0781	6.25	0.443	0.748	1 : 80
0.1563	3.125	0.328	0.668	1 : 20
0.3125	3.125	0.429	0.569	1 : 10
5	200	0.999	0.858	1 : 40
10	200	0.999	0.896	1 : 20

5

[0862]

Table 56

Other antitumor agent : cytarabine, cell line : MDS-L				
Example compound 177 (nM)	Other antitumor agent (nM)	Fa	CI	Combination ratio (Example compound 177 : other antitumor agent)
0.00781	3.125	0.291	0.599	1 : 400
0.00781	6.25	0.481	0.669	1 : 800
0.01563	3.125	0.377	0.513	1 : 200
0.01563	6.25	0.448	0.755	1 : 400
0.03125	3.125	0.447	0.480	1 : 100
0.03125	6.25	0.430	0.867	1 : 200
0.0625	3.125	0.429	0.668	1 : 50
0.0625	6.25	0.517	0.763	1 : 100
0.125	200	0.998	0.864	1 : 1600
0.5	200	0.998	0.864	1 : 400
1	6.25	0.747	0.784	1 : 6.25

[0863]

10

Table 57

Other antitumor agent : cytarabine, cell line : MDS-L				
Example compound 273 (nM)	Other antitumor agent (nM)	Fa	CI	Combination ratio (Example compound 273 : other antitumor agent)
0.00391	3.125	0.245	0.772	1 : 800
0.00391	6.25	0.477	0.718	1 : 1600
0.00781	3.125	0.328	0.609	1 : 400
0.00781	6.25	0.474	0.741	1 : 800
0.01563	3.125	0.392	0.548	1 : 200
0.01563	6.25	0.454	0.823	1 : 400
0.03125	3.125	0.474	0.493	1 : 100
0.0625	3.125	0.510	0.556	1 : 50
0.0625	6.25	0.588	0.680	1 : 100
0.125	200	0.997	0.381	1 : 1600
0.25	200	0.998	0.288	1 : 800
0.5	200	0.998	0.289	1 : 400
1	200	0.999	0.191	1 : 200

[0864]

Table 58

Other antitumor agent : decitabine, cell line : Kasumi-1				
Example compound 37 (nM)	Other antitumor agent (nM)	Fa	CI	Combination ratio (Example compound 37 : other antitumor agent)
0.391	6.25	0.326	0.674	1 : 16
0.391	12.5	0.404	0.897	1 : 32
0.391	25	0.758	0.657	1 : 64
0.391	50	0.872	0.827	1 : 128
0.781	25	0.788	0.615	1 : 32
0.781	50	0.879	0.806	1 : 64
0.781	100	0.958	0.833	1 : 128
1.563	3.125	0.410	0.776	1 : 2
1.563	6.25	0.433	0.879	1 : 4
1.563	12.5	0.544	0.861	1 : 8
1.563	25	0.788	0.660	1 : 16
1.563	50	0.943	0.512	1 : 32
1.563	100	0.973	0.643	1 : 64
1.563	200	0.991	0.687	1 : 128
3.125	3.125	0.580	0.665	1 : 1
3.125	6.25	0.592	0.754	1 : 2
3.125	12.5	0.660	0.778	1 : 4
3.125	25	0.806	0.691	1 : 8
3.125	50	0.943	0.527	1 : 16
3.125	100	0.982	0.514	1 : 32
3.125	200	0.992	0.640	1 : 64
6.25	3.125	0.697	0.696	1 : 0.5
6.25	6.25	0.715	0.724	1 : 1
6.25	12.5	0.779	0.664	1 : 2
6.25	25	0.905	0.456	1 : 4
6.25	50	0.958	0.458	1 : 8
6.25	100	0.982	0.524	1 : 16
6.25	200	0.995	0.518	1 : 32
12.5	3.125	0.819	0.620	1 : 0.25
12.5	6.25	0.850	0.543	1 : 0.5
12.5	12.5	0.868	0.569	1 : 1
12.5	25	0.951	0.331	1 : 2
12.5	50	0.978	0.326	1 : 4
12.5	100	0.987	0.444	1 : 8
12.5	200	0.996	0.439	1 : 16
25	3.125	0.872	0.747	1 : 0.125
25	6.25	0.886	0.694	1 : 0.25
25	12.5	0.894	0.726	1 : 0.5
25	25	0.955	0.409	1 : 1
25	50	0.979	0.355	1 : 2
25	100	0.991	0.371	1 : 4
25	200	0.995	0.506	1 : 8
50	6.25	0.923	0.804	1 : 0.125
50	12.5	0.933	0.751	1 : 0.25
50	25	0.970	0.405	1 : 0.5
50	50	0.988	0.287	1 : 1
50	100	0.995	0.279	1 : 2
50	200	0.997	0.385	1 : 4
100	25	0.967	0.705	1 : 0.25
100	50	0.985	0.438	1 : 0.5
100	100	0.992	0.418	1 : 1
100	200	0.997	0.433	1 : 2

[0865]

Table 59

Other antitumor agent : decitabine, cell line : Kasumi-1				
Example compound 166 (nM)	Other antitumor agent (nM)	Fa	CI	Combination ratio (Example compound 166 : other antitumor agent)
0.0391	3.125	0.369	0.775	1 : 80
0.0391	6.25	0.511	0.712	1 : 160
0.0781	100	0.979	0.650	1 : 1280
0.0781	200	0.992	0.664	1 : 2560
0.156	50	0.953	0.585	1 : 320
0.156	100	0.985	0.522	1 : 640
0.156	200	0.993	0.606	1 : 1280
0.3125	25	0.901	0.562	1 : 80
0.3125	50	0.953	0.595	1 : 160
0.3125	100	0.991	0.356	1 : 320
0.3125	200	0.994	0.558	1 : 640
0.625	3.125	0.758	0.699	1 : 5
0.625	6.25	0.813	0.569	1 : 10
0.625	12.5	0.843	0.625	1 : 20
0.625	25	0.906	0.597	1 : 40
0.625	50	0.970	0.439	1 : 80
0.625	100	0.991	0.364	1 : 160
0.625	200	0.996	0.434	1 : 320
1.25	3.125	0.829	0.717	1 : 2.5
1.25	6.25	0.863	0.592	1 : 5
1.25	12.5	0.882	0.624	1 : 10
1.25	25	0.942	0.453	1 : 20
1.25	50	0.977	0.376	1 : 40
1.25	100	0.993	0.308	1 : 80
1.25	200	0.996	0.414	1 : 160
2.5	3.125	0.876	0.810	1 : 1.25
2.5	6.25	0.909	0.571	1 : 2.5
2.5	12.5	0.919	0.595	1 : 5
2.5	25	0.959	0.402	1 : 10
2.5	50	0.984	0.306	1 : 20
2.5	100	0.994	0.281	1 : 40
2.5	200	0.997	0.341	1 : 80
5	3.125	0.924	0.743	1 : 0.625
5	6.25	0.935	0.639	1 : 1.25
5	12.5	0.947	0.562	1 : 2.5
5	25	0.976	0.307	1 : 5
5	50	0.987	0.285	1 : 10
5	100	0.993	0.325	1 : 20
5	200	0.998	0.260	1 : 40
10	12.5	0.958	0.710	1 : 1.25
10	25	0.975	0.457	1 : 2.5
10	50	0.988	0.317	1 : 5
10	100	0.996	0.235	1 : 10
10	200	0.998	0.309	1 : 20

[0866]

Table 60

Other antitumor agent : decitabine, cell line : Kasumi-1				
Example compound 177 (nM)	Other antitumor agent (nM)	Fa	CI	Combination ratio (Example compound 177 : other antitumor agent)
0.00781	25	0.744	0.878	1 : 3200
0.00781	50	0.893	0.893	1 : 6400
0.01563	50	0.941	0.605	1 : 3200
0.01563	100	0.976	0.672	1 : 6400
0.01563	200	0.992	0.666	1 : 12800
0.03125	25	0.776	0.877	1 : 800
0.03125	50	0.941	0.618	1 : 1600
0.03125	100	0.983	0.538	1 : 3200
0.03125	200	0.996	0.447	1 : 6400
0.0625	25	0.844	0.734	1 : 400
0.0625	50	0.957	0.522	1 : 800
0.0625	100	0.983	0.549	1 : 1600
0.0625	200	0.995	0.501	1 : 3200
0.125	3.125	0.719	0.758	1 : 25
0.125	6.25	0.767	0.705	1 : 50
0.125	12.5	0.843	0.612	1 : 100
0.125	25	0.945	0.393	1 : 200
0.125	50	0.967	0.471	1 : 400
0.125	100	0.987	0.477	1 : 800
0.125	200	0.997	0.368	1 : 1600
0.25	3.125	0.844	0.716	1 : 12.5
0.25	6.25	0.882	0.595	1 : 25
0.25	12.5	0.899	0.617	1 : 50
0.25	25	0.956	0.418	1 : 100
0.25	50	0.981	0.359	1 : 200
0.25	100	0.99	0.425	1 : 400
0.25	200	0.996	0.450	1 : 800
0.5	12.5	0.932	0.706	1 : 25
0.5	25	0.972	0.412	1 : 50
0.5	50	0.989	0.301	1 : 100
0.5	100	0.992	0.409	1 : 200
0.5	200	0.996	0.468	1 : 400
1	25	0.974	0.603	1 : 25
1	50	0.988	0.420	1 : 50
1	100	0.992	0.477	1 : 100
1	200	0.995	0.571	1 : 200

[0867]

Table 61

Other antitumor agent : decitabine, cell line : Kasumi-1				
Example compound 273 (nM)	Other antitumor agent (nM)	Fa	CI	Combination ratio (Example compound 273 : other antitumor agent)
0.00391	3.125	0.223	0.781	1 : 800
0.00781	100	0.971	0.813	1 : 12800
0.00781	200	0.989	0.839	1 : 25600
0.01563	50	0.932	0.752	1 : 3200
0.01563	100	0.972	0.799	1 : 6400
0.01563	200	0.993	0.620	1 : 12800
0.03125	50	0.932	0.768	1 : 1600
0.03125	100	0.984	0.550	1 : 3200
0.03125	200	0.996	0.419	1 : 6400
0.0625	25	0.859	0.775	1 : 400
0.0625	50	0.952	0.620	1 : 800
0.0625	100	0.984	0.555	1 : 1600
0.0625	200	0.996	0.436	1 : 3200
0.125	3.125	0.717	0.817	1 : 25
0.125	6.25	0.781	0.710	1 : 50
0.125	12.5	0.817	0.780	1 : 100
0.125	25	0.941	0.443	1 : 200
0.125	50	0.961	0.571	1 : 400
0.125	100	0.989	0.440	1 : 800
0.125	200	0.996	0.432	1 : 1600
0.25	3.125	0.836	0.774	1 : 12.5
0.25	6.25	0.849	0.790	1 : 25
0.25	12.5	0.899	0.644	1 : 50
0.25	25	0.942	0.551	1 : 100
0.25	50	0.978	0.419	1 : 200
0.25	100	0.992	0.369	1 : 400
0.25	200	0.996	0.440	1 : 800
0.5	6.25	0.914	0.781	1 : 12.5
0.5	12.5	0.941	0.619	1 : 25
0.5	25	0.974	0.386	1 : 50
0.5	50	0.985	0.375	1 : 100
0.5	100	0.994	0.333	1 : 200
0.5	200	0.997	0.374	1 : 400
1	25	0.976	0.543	1 : 25
1	50	0.987	0.434	1 : 50
1	100	0.995	0.339	1 : 100
1	200	0.997	0.407	1 : 200

[0868]

Table 62

Other antitumor agent : decitabine, cell line : Kasumi-1				
Example compound 41 (nM)	Other antitumor agent (nM)	Fa	CI	Combination ratio (Example compound 41 : other antitumor agent)
0.391	50	0.757	0.717	1 : 128
0.781	25	0.617	0.709	1 : 32
0.781	50	0.764	0.713	1 : 64
0.781	200	0.955	0.540	1 : 256
1.563	25	0.617	0.808	1 : 16
1.563	50	0.866	0.409	1 : 32
1.563	200	0.972	0.356	1 : 128
3.125	3.125	0.547	0.654	1 : 1
3.125	6.25	0.636	0.501	1 : 2
3.125	12.5	0.587	0.799	1 : 4
3.125	25	0.744	0.564	1 : 8
3.125	50	0.866	0.447	1 : 16
3.125	100	0.944	0.350	1 : 32
3.125	200	0.978	0.291	1 : 64
6.25	3.125	0.692	0.590	1 : 0.5
6.25	6.25	0.783	0.380	1 : 1
6.25	12.5	0.797	0.417	1 : 2
6.25	25	0.865	0.340	1 : 4
6.25	50	0.929	0.268	1 : 8
6.25	100	0.944	0.375	1 : 16
6.25	200	0.989	0.158	1 : 32
12.5	3.125	0.826	0.467	1 : 0.25
12.5	6.25	0.853	0.395	1 : 0.5
12.5	12.5	0.885	0.325	1 : 1
12.5	25	0.914	0.289	1 : 2
12.5	50	0.957	0.198	1 : 4
12.5	100	0.945	0.416	1 : 8
12.5	200	0.989	0.168	1 : 16
25	6.25	0.888	0.514	1 : 0.25
25	12.5	0.913	0.402	1 : 0.5
25	25	0.920	0.421	1 : 1
25	50	0.967	0.201	1 : 2
25	100	0.967	0.303	1 : 4
25	200	0.995	0.089	1 : 8
50	12.5	0.919	0.678	1 : 0.25
50	25	0.958	0.333	1 : 0.5
50	50	0.969	0.278	1 : 1
50	100	0.970	0.368	1 : 2
50	200	0.995	0.099	1 : 4
100	25	0.957	0.621	1 : 0.25
100	50	0.971	0.433	1 : 0.5
100	100	0.972	0.509	1 : 1
100	200	0.992	0.202	1 : 2

[0869]

Table 63

Other antitumor agent : decitabine, cell line : Kasumi-1				
Example compound 214 (nM)	Other antitumor agent (nM)	Fa	CI	Combination ratio (Example compound 214 : other antitumor agent)
0.00391	100	0.867	0.693	1 : 25600
0.00781	100	0.865	0.704	1 : 12800
0.01563	100	0.873	0.671	1 : 6400
0.01563	200	0.932	0.744	1 : 12800
0.03125	100	0.904	0.521	1 : 3200
0.0625	100	0.904	0.532	1 : 1600
0.0625	200	0.957	0.503	1 : 3200
0.125	50	0.722	0.890	1 : 400
0.125	100	0.910	0.526	1 : 800
0.125	200	0.96	0.486	1 : 1600
0.25	100	0.884	0.727	1 : 400
0.5	3.125	0.745	0.561	1 : 6.25
0.5	6.25	0.692	0.743	1 : 12.5
0.5	12.5	0.791	0.560	1 : 25
0.5	50	0.884	0.549	1 : 100
0.5	100	0.966	0.284	1 : 200
0.5	200	0.985	0.243	1 : 400
1	12.5	0.904	0.477	1 : 12.5
1	25	0.856	0.780	1 : 25
1	50	0.930	0.506	1 : 50
1	100	0.964	0.396	1 : 100
1	200	0.986	0.265	1 : 200

[0870]

Table 64

Other antitumor agent : decitabine, cell line : Kasumi-1				
Example compound 285 (nM)	Other antitumor agent (nM)	Fa	CI	Combination ratio (Example compound 285 : other antitumor agent)
0.000781	50	0.569	0.814	1 : 64000
0.00156	25	0.222	0.894	1 : 16000
0.00156	50	0.614	0.747	1 : 32000
0.00313	50	0.617	0.754	1 : 16000
0.00625	50	0.617	0.774	1 : 8000
0.00625	100	0.848	0.810	1 : 16000
0.0125	25	0.418	0.688	1 : 2000
0.0125	50	0.722	0.638	1 : 4000
0.0125	100	0.848	0.829	1 : 8000
0.025	25	0.516	0.667	1 : 1000
0.025	50	0.795	0.566	1 : 2000
0.025	100	0.911	0.635	1 : 4000
0.025	200	0.962	0.777	1 : 8000
0.05	12.5	0.803	0.293	1 : 250
0.05	25	0.709	0.548	1 : 500
0.05	50	0.938	0.315	1 : 1000
0.05	100	0.920	0.643	1 : 2000
0.05	200	0.990	0.404	1 : 4000
0.1	3.125	0.676	0.596	1 : 31.3
0.1	6.25	0.792	0.432	1 : 62.5
0.1	12.5	0.724	0.621	1 : 125
0.1	25	0.884	0.404	1 : 250
0.1	50	0.967	0.269	1 : 500
0.1	100	0.977	0.362	1 : 1000
0.1	200	0.977	0.656	1 : 2000
0.2	6.25	0.881	0.520	1 : 31.3
0.2	12.5	0.870	0.601	1 : 62.5
0.2	25	0.919	0.498	1 : 125
0.2	50	0.972	0.331	1 : 250
0.2	100	0.985	0.342	1 : 500
0.2	200	0.993	0.381	1 : 1000

[0871]

Table 65

Other antitumor agent : decitabine, cell line : MDS-L				
Example compound 37 (nM)	Other antitumor agent (nM)	Fa	CI	Combination ratio (Example compound 37 : other antitumor agent)
0.391	6.25	0.434	0.708	1 : 16
0.781	6.25	0.446	0.711	1 : 8
1.563	100	0.984	0.852	1 : 64
3.125	100	0.983	0.879	1 : 32
6.25	6.25	0.498	0.867	1 : 1
6.25	12.5	0.685	0.817	1 : 2
12.5	6.25	0.604	0.733	1 : 0.5
12.5	12.5	0.754	0.695	1 : 1
12.5	25	0.847	0.891	1 : 2
12.5	50	0.951	0.829	1 : 4
12.5	100	0.984	0.842	1 : 8
25	6.25	0.729	0.546	1 : 0.25
25	12.5	0.823	0.564	1 : 0.5
25	25	0.877	0.784	1 : 1
25	50	0.971	0.608	1 : 2
25	100	0.990	0.638	1 : 4
50	6.25	0.779	0.560	1 : 0.125
50	12.5	0.835	0.607	1 : 0.25
50	25	0.932	0.536	1 : 0.5
50	50	0.977	0.531	1 : 1
50	100	0.990	0.657	1 : 2
50	200	0.995	0.849	1 : 4
100	6.25	0.827	0.574	1 : 0.0625
100	12.5	0.904	0.427	1 : 0.125
100	25	0.949	0.459	1 : 0.25
100	50	0.979	0.503	1 : 0.5
100	100	0.992	0.577	1 : 1
100	200	0.995	0.849	1 : 2

[0872]

Table 66

Other antitumor agent : decitabine, cell line : MDS-L				
Example compound 166 (nM)	Other antitumor agent (nM)	Fa	CI	Combination ratio (Example compound 166 : other antitumor agent)
0.0391	100	0.986	0.706	1 : 2560
0.0781	6.25	0.467	0.742	1 : 80
0.1563	6.25	0.448	0.835	1 : 40
0.3125	200	0.994	0.866	1 : 640
0.625	6.25	0.566	0.720	1 : 10
0.625	12.5	0.674	0.891	1 : 20
1.25	6.25	0.654	0.615	1 : 5
1.25	12.5	0.765	0.681	1 : 10
1.25	25	0.856	0.860	1 : 20
1.25	50	0.951	0.797	1 : 40
1.25	100	0.985	0.737	1 : 80
2.5	6.25	0.762	0.454	1 : 2.5
2.5	12.5	0.825	0.557	1 : 5
2.5	25	0.925	0.541	1 : 10
2.5	50	0.956	0.750	1 : 20
2.5	100	0.987	0.673	1 : 40
5	3.125	0.669	0.862	1 : 0.625
5	6.25	0.792	0.491	1 : 1.25
5	12.5	0.848	0.544	1 : 2.5
5	25	0.927	0.544	1 : 5
5	50	0.971	0.570	1 : 10
5	100	0.990	0.584	1 : 20
5	200	0.995	0.734	1 : 40
10	6.25	0.820	0.559	1 : 0.625
10	12.5	0.886	0.475	1 : 1.25
10	25	0.935	0.520	1 : 2.5
10	50	0.976	0.505	1 : 5
10	100	0.990	0.560	1 : 10
10	200	0.994	0.788	1 : 20

[0873]

Table 67

Other antitumor agent : decitabine, cell line : MDS-L				
Example compound 177 (nM)	Other antitumor agent (nM)	Fa	CI	Combination ratio (Example compound 177 : other antitumor agent)
0.00391	50	0.953	0.828	1 : 12800
0.00781	3.125	0.374	0.539	1 : 400
0.00781	6.25	0.409	0.870	1 : 800
0.01563	3.125	0.417	0.537	1 : 200
0.03125	3.125	0.499	0.475	1 : 100
0.0625	3.125	0.458	0.784	1 : 50
0.125	12.5	0.705	0.859	1 : 100
0.125	50	0.952	0.840	1 : 400
0.125	100	0.984	0.848	1 : 800
0.25	3.125	0.633	0.649	1 : 12.5
0.25	6.25	0.698	0.618	1 : 25
0.25	12.5	0.817	0.578	1 : 50
0.25	25	0.871	0.829	1 : 100
0.25	50	0.965	0.694	1 : 200
0.5	3.125	0.698	0.662	1 : 6.25
0.5	6.25	0.780	0.497	1 : 12.5
0.5	12.5	0.853	0.515	1 : 25
0.5	25	0.925	0.575	1 : 50
0.5	50	0.973	0.581	1 : 100
0.5	100	0.989	0.690	1 : 200
0.5	200	0.995	0.837	1 : 400
1	6.25	0.812	0.535	1 : 6.25
1	12.5	0.879	0.480	1 : 12.5
1	25	0.931	0.557	1 : 25
1	50	0.977	0.533	1 : 50
1	100	0.993	0.526	1 : 100
1	200	0.995	0.837	1 : 200

[0874]

Table 68

Other antitumor agent : decitabine, cell line : MDS-L				
Example compound 273 (nM)	Other antitumor agent (nM)	Fa	CI	Combination ratio (Example compound 273 : other antitumor agent)
0.00391	50	0.945	0.884	1 : 12800
0.00391	100	0.984	0.836	1 : 25600
0.00781	50	0.961	0.720	1 : 6400
0.00781	100	0.983	0.866	1 : 12800
0.01563	3.125	0.365	0.527	1 : 200
0.125	6.25	0.530	0.858	1 : 50
0.125	12.5	0.682	0.856	1 : 100
0.125	25	0.852	0.860	1 : 200
0.25	6.25	0.665	0.633	1 : 25
0.25	12.5	0.817	0.549	1 : 50
0.25	25	0.855	0.867	1 : 100
0.25	50	0.956	0.771	1 : 200
0.25	100	0.986	0.772	1 : 400
0.5	3.125	0.618	0.899	1 : 6.25
0.5	6.25	0.736	0.593	1 : 12.5
0.5	12.5	0.849	0.509	1 : 25
0.5	25	0.891	0.728	1 : 50
0.5	50	0.975	0.550	1 : 100
0.5	100	0.991	0.605	1 : 200
0.5	200	0.996	0.735	1 : 400
1	6.25	0.808	0.525	1 : 6.25
1	12.5	0.899	0.402	1 : 12.5
1	25	0.910	0.666	1 : 25
1	50	0.971	0.605	1 : 50
1	100	0.992	0.566	1 : 100

[0875]

Table 69

Other antitumor agent : azacitidine, cell line : Kasumi-1				
Example compound 37 (nM)	Other antitumor agent (nM)	Fa	CI	Combination ratio (Example compound 37 : other antitumor agent)
0.391	5000	0.548	0.844	1 : 12800
0.781	5000	0.646	0.729	1 : 6400
0.781	10000	0.783	0.863	1 : 12800
1.563	1250	0.575	0.621	1 : 800
1.563	2500	0.661	0.564	1 : 1600
1.563	5000	0.752	0.605	1 : 3200
1.563	10000	0.903	0.521	1 : 6400
3.125	1250	0.709	0.560	1 : 400
3.125	2500	0.661	0.852	1 : 800
3.125	5000	0.915	0.293	1 : 1600
3.125	10000	0.927	0.465	1 : 3200
6.25	1250	0.827	0.465	1 : 200
6.25	2500	0.786	0.723	1 : 400
6.25	5000	0.915	0.361	1 : 800
6.25	10000	0.955	0.367	1 : 1600
12.5	156.25	0.827	0.769	1 : 12.5
12.5	312.5	0.839	0.702	1 : 25
12.5	625	0.818	0.860	1 : 50
12.5	1250	0.913	0.342	1 : 100
12.5	2500	0.931	0.308	1 : 200
12.5	5000	0.948	0.312	1 : 400
12.5	10000	0.980	0.240	1 : 800
25	156.25	0.895	0.748	1 : 6.25
25	1250	0.917	0.588	1 : 50
25	2500	0.940	0.442	1 : 100
25	5000	0.955	0.394	1 : 200
25	10000	0.982	0.260	1 : 400
50	1250	0.952	0.560	1 : 25
50	2500	0.963	0.446	1 : 50
50	5000	0.976	0.328	1 : 100
50	10000	0.987	0.255	1 : 200
100	1250	0.966	0.700	1 : 12.5
100	2500	0.977	0.458	1 : 25
100	5000	0.976	0.539	1 : 50
100	10000	0.989	0.295	1 : 100

[0876]

Table 70

Other antitumor agent : azacitidine, cell line : Kasumi-1				
Example compound 166 (nM)	Other antitumor agent (nM)	Fa	CI	Combination ratio (Example compound 166 : other antitumor agent)
0.0391	156.25	0.370	0.325	1 : 4000
0.0391	312.5	0.296	0.629	1 : 8000
0.0781	1250	0.484	0.729	1 : 16000
0.0781	2500	0.591	0.717	1 : 32000
0.0781	5000	0.733	0.591	1 : 64000
0.0781	10000	0.877	0.351	1 : 128000
0.1563	2500	0.829	0.213	1 : 16000
0.1563	5000	0.844	0.302	1 : 32000
0.1563	10000	0.931	0.177	1 : 64000
0.3125	156.25	0.643	0.535	1 : 500
0.3125	312.5	0.675	0.478	1 : 1000
0.3125	625	0.667	0.545	1 : 2000
0.3125	1250	0.753	0.395	1 : 4000
0.3125	2500	0.829	0.297	1 : 8000
0.3125	5000	0.893	0.225	1 : 16000
0.3125	10000	0.946	0.150	1 : 32000
0.625	156.25	0.757	0.565	1 : 250
0.625	312.5	0.794	0.457	1 : 500
0.625	625	0.748	0.638	1 : 1000
0.625	1250	0.799	0.502	1 : 2000
0.625	2500	0.860	0.355	1 : 4000
0.625	5000	0.893	0.318	1 : 8000
0.625	10000	0.980	0.056	1 : 16000
1.25	156.25	0.836	0.644	1 : 125
1.25	312.5	0.861	0.524	1 : 250
1.25	625	0.845	0.617	1 : 500
1.25	1250	0.893	0.402	1 : 1000
1.25	2500	0.925	0.279	1 : 2000
1.25	5000	0.961	0.146	1 : 4000
1.25	10000	0.980	0.083	1 : 8000
2.5	156.25	0.906	0.628	1 : 62.5
2.5	312.5	0.918	0.535	1 : 125
2.5	625	0.889	0.786	1 : 250
2.5	1250	0.946	0.332	1 : 500
2.5	2500	0.954	0.285	1 : 1000
2.5	5000	0.963	0.240	1 : 2000
2.5	10000	0.979	0.141	1 : 4000
5	1250	0.952	0.570	1 : 250
5	2500	0.966	0.387	1 : 500
5	5000	0.979	0.229	1 : 1000
5	10000	0.988	0.131	1 : 2000
10	1250	0.963	0.833	1 : 125
10	2500	0.976	0.511	1 : 250
10	5000	0.986	0.284	1 : 500
10	10000	0.993	0.135	1 : 1000

[0877]

Table 71

Other antitumor agent : azacitidine, cell line : Kasumi-1				
Example compound 177 (nM)	Other antitumor agent (nM)	Fa	CI	Combination ratio (Example compound 177 : other antitumor agent)
0.00391	156.25	0.324	0.309	1 : 40000
0.00391	312.5	0.225	0.643	1 : 80000
0.00781	10000	0.834	0.225	1 : 1280000
0.01563	2500	0.545	0.632	1 : 160000
0.01563	5000	0.720	0.365	1 : 320000
0.01563	10000	0.852	0.222	1 : 640000
0.03125	5000	0.880	0.153	1 : 160000
0.03125	10000	0.868	0.241	1 : 320000
0.0625	2500	0.732	0.667	1 : 40000
0.0625	5000	0.880	0.244	1 : 80000
0.0625	10000	0.936	0.136	1 : 160000
0.125	1250	0.830	0.611	1 : 10000
0.125	2500	0.851	0.530	1 : 20000
0.125	5000	0.934	0.201	1 : 40000
0.125	10000	0.973	0.078	1 : 80000
0.25	1250	0.903	0.576	1 : 5000
0.25	2500	0.897	0.626	1 : 10000
0.25	5000	0.965	0.175	1 : 20000
0.25	10000	0.980	0.096	1 : 40000
0.5	1250	0.939	0.638	1 : 2500
0.5	2500	0.947	0.539	1 : 5000
0.5	5000	0.976	0.217	1 : 10000
0.5	10000	0.984	0.139	1 : 20000
1	5000	0.976	0.422	1 : 5000
1	10000	0.984	0.265	1 : 10000

[0878]

Table 72

Other antitumor agent : azacitidine, cell line : Kasumi-1				
Example compound 273 (nM)	Other antitumor agent (nM)	Fa	CI	Combination ratio (Example compound 273 : other antitumor agent)
0.00391	156.25	0.206	0.316	1 : 40000
0.00391	312.5	0.295	0.283	1 : 80000
0.00781	1250	0.384	0.588	1 : 160000
0.00781	2500	0.437	0.794	1 : 320000
0.00781	5000	0.635	0.558	1 : 640000
0.00781	10000	0.747	0.556	1 : 1280000
0.01563	1250	0.384	0.733	1 : 80000
0.01563	2500	0.644	0.356	1 : 160000
0.01563	5000	0.691	0.463	1 : 320000
0.01563	10000	0.849	0.279	1 : 640000
0.03125	156.25	0.476	0.459	1 : 5000
0.03125	312.5	0.425	0.595	1 : 10000
0.03125	625	0.404	0.743	1 : 20000
0.03125	1250	0.492	0.659	1 : 40000
0.03125	2500	0.644	0.475	1 : 80000
0.03125	5000	0.858	0.193	1 : 160000
0.03125	10000	0.914	0.160	1 : 320000
0.0625	156.25	0.563	0.656	1 : 2500
0.0625	312.5	0.640	0.514	1 : 5000
0.0625	625	0.561	0.730	1 : 10000
0.0625	1250	0.704	0.463	1 : 20000
0.0625	2500	0.724	0.498	1 : 40000
0.0625	5000	0.858	0.279	1 : 80000
0.0625	10000	0.934	0.159	1 : 160000
0.125	156.25	0.785	0.534	1 : 1250
0.125	312.5	0.771	0.579	1 : 2500
0.125	625	0.683	0.870	1 : 5000
0.125	1250	0.781	0.590	1 : 10000
0.125	2500	0.860	0.393	1 : 20000
0.125	5000	0.910	0.281	1 : 40000
0.125	10000	0.951	0.180	1 : 80000
0.25	156.25	0.874	0.618	1 : 625
0.25	312.5	0.869	0.645	1 : 1250
0.25	625	0.839	0.799	1 : 2500
0.25	1250	0.903	0.497	1 : 5000
0.25	2500	0.936	0.345	1 : 10000
0.25	5000	0.963	0.220	1 : 20000
0.25	10000	0.979	0.141	1 : 40000
0.5	1250	0.933	0.695	1 : 2500
0.5	2500	0.943	0.606	1 : 5000
0.5	5000	0.968	0.375	1 : 10000
0.5	10000	0.985	0.197	1 : 20000
1	5000	0.973	0.628	1 : 5000
1	10000	0.985	0.383	1 : 10000

[0879]

Table 73

Other antitumor agent : azacitidine, cell line : MDS-L				
Example compound 37 (nM)	Other antitumor agent (nM)	Fa	CI	Combination ratio (Example compound 37 : other antitumor agent)
0.391	312.5	0.376	0.547	1 : 800
0.391	625	0.451	0.720	1 : 1600
0.391	1250	0.706	0.403	1 : 3200
0.391	2500	0.724	0.722	1 : 6400
0.781	156.25	0.245	0.789	1 : 200
0.781	312.5	0.368	0.624	1 : 400
0.781	625	0.451	0.753	1 : 800
0.781	1250	0.663	0.515	1 : 1600
0.781	2500	0.706	0.804	1 : 3200
1.563	1250	0.663	0.532	1 : 800
12.5	156.25	0.509	0.840	1 : 12.5
25	312.5	0.615	0.870	1 : 12.5

[0880]

Table 74

Other antitumor agent : azacitidine, cell line : MDS-L				
Example compound 166 (nM)	Other antitumor agent (nM)	Fa	CI	Combination ratio (Example compound 166 : other antitumor agent)
0.0391	312.5	0.219	0.897	1 : 8000
0.0391	625	0.433	0.604	1 : 16000
0.0781	312.5	0.338	0.543	1 : 4000
0.0781	625	0.433	0.636	1 : 8000
0.1563	312.5	0.445	0.391	1 : 2000
0.1563	625	0.455	0.637	1 : 4000
0.3125	156.25	0.344	0.712	1 : 500
0.3125	312.5	0.367	0.793	1 : 1000
0.625	1250	0.571	0.884	1 : 2000
0.625	2500	0.719	0.844	1 : 4000
1.25	312.5	0.540	0.657	1 : 250
1.25	625	0.589	0.646	1 : 500
1.25	1250	0.674	0.637	1 : 1000
1.25	2500	0.788	0.623	1 : 2000
1.25	5000	0.853	0.793	1 : 4000
2.5	312.5	0.604	0.722	1 : 125
2.5	625	0.695	0.487	1 : 250
2.5	1250	0.793	0.380	1 : 500
2.5	2500	0.848	0.448	1 : 1000
2.5	5000	0.896	0.566	1 : 2000
2.5	10000	0.936	0.691	1 : 4000
5	312.5	0.711	0.568	1 : 62.5
5	625	0.770	0.419	1 : 125
5	1250	0.818	0.393	1 : 250
5	2500	0.867	0.422	1 : 500
5	5000	0.938	0.346	1 : 1000
5	10000	0.959	0.459	1 : 2000
10	625	0.790	0.564	1 : 62.5
10	1250	0.827	0.499	1 : 125
10	2500	0.878	0.445	1 : 250
10	5000	0.911	0.535	1 : 500
10	10000	0.962	0.438	1 : 1000

[0881]

Table 75

Other antitumor agent : azacitidine, cell line : MDS-L

Example compound 177 (nM)	Other antitumor agent (nM)	Fa	CI	Combination ratio (Example compound 177 : other antitumor agent)
0.00781	156.25	0.250	0.452	1 : 20000
0.00781	312.5	0.357	0.439	1 : 40000
0.01563	156.25	0.337	0.357	1 : 10000
0.01563	312.5	0.344	0.539	1 : 20000
0.01563	625	0.414	0.691	1 : 40000
0.03125	156.25	0.297	0.662	1 : 5000
0.03125	312.5	0.369	0.599	1 : 10000
0.0625	156.25	0.394	0.569	1 : 2500
0.125	312.5	0.471	0.722	1 : 2500
0.125	625	0.552	0.641	1 : 5000
0.125	1250	0.639	0.681	1 : 10000
0.125	2500	0.765	0.675	1 : 20000
0.125	5000	0.840	0.839	1 : 40000
0.25	312.5	0.584	0.602	1 : 1250
0.25	625	0.643	0.549	1 : 2500
0.25	1250	0.739	0.485	1 : 5000
0.25	2500	0.796	0.609	1 : 10000
0.25	5000	0.890	0.579	1 : 20000
0.25	10000	0.945	0.578	1 : 40000
0.5	156.25	0.608	0.814	1 : 313
0.5	312.5	0.721	0.405	1 : 625
0.5	625	0.777	0.332	1 : 1250
0.5	1250	0.821	0.346	1 : 2500
0.5	2500	0.844	0.484	1 : 5000
0.5	5000	0.917	0.451	1 : 10000
0.5	10000	0.955	0.482	1 : 20000
1	312.5	0.749	0.568	1 : 313
1	625	0.788	0.471	1 : 625
1	1250	0.822	0.458	1 : 1250
1	2500	0.897	0.339	1 : 2500
1	5000	0.923	0.438	1 : 5000
1	10000	0.958	0.462	1 : 10000

[0882]

Table 76

Other antitumor agent : azacitidine, cell line : MDS-L				
Example compound 273 (nM)	Other antitumor agent (nM)	Fa	CI	Combination ratio (Example compound 273 : other antitumor agent)
0.00781	156.25	0.215	0.537	1 : 20000
0.00781	312.5	0.381	0.383	1 : 40000
0.00781	625	0.374	0.733	1 : 80000
0.01563	2500	0.696	0.801	1 : 160000
0.01563	5000	0.818	0.852	1 : 320000
0.03125	156.25	0.296	0.627	1 : 5000
0.03125	312.5	0.391	0.518	1 : 10000
0.0625	156.25	0.442	0.423	1 : 2500
0.125	625	0.561	0.614	1 : 5000
0.125	1250	0.616	0.741	1 : 10000
0.125	2500	0.768	0.622	1 : 20000
0.125	5000	0.827	0.833	1 : 40000
0.125	10000	0.920	0.720	1 : 80000
0.25	312.5	0.641	0.453	1 : 1250
0.25	625	0.645	0.564	1 : 2500
0.25	1250	0.724	0.527	1 : 5000
0.25	2500	0.810	0.527	1 : 10000
0.25	5000	0.921	0.369	1 : 20000
0.5	156.25	0.611	0.875	1 : 313
0.5	312.5	0.663	0.680	1 : 625
0.5	625	0.745	0.463	1 : 1250
0.5	1250	0.809	0.395	1 : 2500
0.5	2500	0.853	0.439	1 : 5000
0.5	5000	0.929	0.348	1 : 10000
0.5	10000	0.965	0.327	1 : 20000
1	312.5	0.711	0.893	1 : 313
1	625	0.801	0.489	1 : 625
1	1250	0.824	0.497	1 : 1250
1	2500	0.905	0.309	1 : 2500
1	5000	0.930	0.367	1 : 5000
1	10000	0.967	0.315	1 : 10000

[0883]

Table 77

Other antitumor agent : ATRA, cell line : Kasumi-1				
Example compound 37 (nM)	Other antitumor agent (nM)	Fa	CI	Combination ratio (Example compound 37 : other antitumor agent)
0.391	2000	0.635	0.566	1 : 5120
3.13	3.2	0.564	0.877	1 : 1.02
3.13	16	0.579	0.824	1 : 5.11
6.25	80	0.786	0.530	1 : 12.8

5

[0884]

Table 78

Other antitumor agent : ATRA, cell line : Kasumi-1				
Example compound 166 (nM)	Other antitumor agent (nM)	Fa	CI	Combination ratio (Example compound 166 : other antitumor agent)
0.0391	2000	0.528	0.869	1 : 51200
0.313	0.64	0.529	0.852	1 : 2.05
0.625	0.64	0.713	0.730	1 : 1.02
0.625	3.2	0.732	0.660	1 : 5.12
0.625	16	0.701	0.779	1 : 25.6
0.625	80	0.695	0.807	1 : 128

[0885]

Table 79

Other antitumor agent : ATRA, cell line : Kasumi-1				
Example compound 177 (nM)	Other antitumor agent (nM)	Fa	CI	Combination ratio (Example compound 177 : other antitumor agent)
0.00391	16	0.166	0.737	1 : 4090
0.0313	3.2	0.440	0.824	1 : 102
0.0625	80	0.651	0.703	1 : 1280

[0886]

5 Table 80

Other antitumor agent : ATRA, cell line : Kasumi-1				
Example compound 273 (nM)	Other antitumor agent (nM)	Fa	CI	Combination ratio (Example compound 273 : other antitumor agent)
0.00391	3.2	0.265	0.304	1 : 818
0.00391	16	0.292	0.369	1 : 4090
0.00391	80	0.314	0.770	1 : 20500
0.00781	3.2	0.258	0.597	1 : 410

[0887]

Table 81

Other antitumor agent : daunorubicin, cell line : Kasumi-1				
Example compound 37 (nM)	Other antitumor agent (nM)	Fa	CI	Combination ratio (Example compound 37 : other antitumor agent)
0.781	10	0.992	0.283	1 : 12.8
1.563	2.5	0.703	0.654	1 : 1.6
1.563	5	0.773	0.893	1 : 3.2
1.563	10	0.987	0.360	1 : 6.4
3.125	2.5	0.703	0.869	1 : 0.8
3.125	5	0.894	0.611	1 : 1.6
3.125	10	0.994	0.256	1 : 3.2
6.25	5	0.894	0.733	1 : 0.8
6.25	10	0.996	0.214	1 : 1.6
12.5	1.25	0.853	0.851	1 : 0.1
12.5	2.5	0.878	0.834	1 : 0.2
12.5	5	0.942	0.616	1 : 0.4
12.5	10	0.997	0.191	1 : 0.8
25	2.5	0.929	0.832	1 : 0.1
25	5	0.966	0.572	1 : 0.2
25	10	0.998	0.165	1 : 0.4
50	5	0.971	0.751	1 : 0.1
50	10	0.999	0.124	1 : 0.2
100	10	0.998	0.250	1 : 0.1

10 [0888]

Table 82

Other antitumor agent : daunorubicin, cell line : Kasumi-1				
Example compound 166 (nM)	Other antitumor agent (nM)	Fa	CI	Combination ratio (Example compound 166 : other antitumor agent)
0.0781	10	0.986	0.823	1 : 128
0.1563	10	0.986	0.824	1 : 64
0.3125	5	0.894	0.771	1 : 16
0.3125	10	0.995	0.622	1 : 32
0.625	0.625	0.706	0.875	1 : 1
0.625	1.25	0.751	0.767	1 : 2
0.625	2.5	0.801	0.778	1 : 4
0.625	5	0.893	0.830	1 : 8
0.625	10	0.995	0.645	1 : 16
1.25	0.1563	0.835	0.517	1 : 0.125
1.25	0.3125	0.828	0.581	1 : 0.25
1.25	0.625	0.827	0.640	1 : 0.5
1.25	1.25	0.838	0.681	1 : 1
1.25	2.5	0.870	0.701	1 : 2
1.25	5	0.940	0.697	1 : 4
1.25	10	0.996	0.594	1 : 8
2.5	0.1563	0.866	0.705	1 : 0.0625
2.5	0.3125	0.886	0.562	1 : 0.125
2.5	0.625	0.879	0.664	1 : 0.25
2.5	1.25	0.905	0.554	1 : 0.5
2.5	2.5	0.925	0.582	1 : 1
2.5	5	0.960	0.638	1 : 2
2.5	10	0.996	0.596	1 : 4
5	0.1563	0.912	0.692	1 : 0.0313
5	0.3125	0.928	0.523	1 : 0.0625
5	0.625	0.916	0.705	1 : 0.125
5	1.25	0.931	0.609	1 : 0.25
5	2.5	0.943	0.634	1 : 0.5
5	5	0.968	0.648	1 : 1
5	10	0.995	0.636	1 : 2
10	0.313	0.934	0.880	1 : 0.0313
10	0.625	0.944	0.721	1 : 0.0625
10	1.25	0.942	0.834	1 : 0.125
10	2.5	0.947	0.886	1 : 0.25
10	5	0.969	0.756	1 : 0.5
10	10	0.997	0.558	1 : 1

[0889]

Table 83

Other antitumor agent : daunorubicin, cell line : Kasumi-1				
Example compound 177 (nM)	Other antitumor agent (nM)	Fa	CI	Combination ratio (Example compound 177 : other antitumor agent)
0.01563	10	0.950	0.859	1 : 640
0.03125	5	0.768	0.826	1 : 160
0.03125	10	0.967	0.779	1 : 320
0.0625	10	0.980	0.697	1 : 160
0.125	0.3125	0.730	0.795	1 : 2.5
0.125	0.625	0.729	0.842	1 : 5
0.125	1.25	0.756	0.849	1 : 10
0.25	10	0.987	0.708	1 : 40
0.5	10	0.992	0.687	1 : 20
1	10	0.993	0.814	1 : 10

[0890]

Table 84

Other antitumor agent : daunorubicin, cell line : Kasumi-1				
Example compound 273 (nM)	Other antitumor agent (nM)	Fa	CI	Combination ratio (Example compound 273 : other antitumor agent)
0.00781	10	0.974	0.790	1 : 1280
0.01563	10	0.983	0.702	1 : 640
0.03125	10	0.977	0.778	1 : 320
0.0625	10	0.983	0.724	1 : 160
0.125	10	0.990	0.635	1 : 80
0.25	5	0.937	0.873	1 : 20
0.25	10	0.995	0.533	1 : 40
0.5	10	0.997	0.481	1 : 20
1	10	0.994	0.713	1 : 10

[0891]

5 These results indicated that the antitumor effects of the biphenyl compounds were synergistically potentiated when used in combination with an antimetabolite (cytarabine, decitabine, or azacitidine), an all-trans retinoic acid (tretinoin (ATRA)), or an antitumor antibiotic (daunorubicin).

[0892]

10 Test Example 5: Combination of Biphenyl Compounds and Other Antitumor agents Using Human Small-Cell Lung Cancer Cell Line (in vitro)

15 The following tables show the source of reagents, the source of tumor cell lines, the media used, and the number of seeded cells.

[0893]

Table 85

Reagent	Source
Fetal bovine serum (FBS)	Biological Industries
RPMI-1640 medium	GIBCO
F-12K medium	GIBCO
Cisplatin	SIGMA-ALDRICH
Carboplatin	SIGMA-ALDRICH
Etoposide	SIGMA-ALDRICH
Paclitaxel	Wako Pure Chemical Industries, Ltd.
SN-38	Tokyo Chemical Industry Co., Ltd.

[0894]

Table 86

Tumor cell line (origin)	Source of cell line	Medium	Number of seeded cells per well (cells)
NCI-H1417 (human small-cell lung cancer)	ATCC	RPMI-1640 medium containing 10% FBS	5000
NCI-H510A (human small-cell lung cancer)	ATCC	F-12K medium containing 10% FBS	7500

[0895]

Cells were seeded at 100 μ L/well in a 96-well culture plate (NUNC). The plate seeded with cells was cultured in an incubator set at 37°C and 5% CO₂. On the day after seeding, drug solutions containing a combination of biphenyl compounds and other compounds having an antitumor effect mixed at various concentrations were each added to the cells. Specifically, using an RPMI-1640 medium containing 10% FBS or an F-12K medium containing 10% FBS, eight-step serial dilutions (including 0 nM) of the biphenyl compounds (Example compounds 37, 166, 177, and 273) were prepared, and eight-step serial dilutions (including 0 nM) of other compounds having an antitumor effect were also prepared. Drug solutions in which the same amount of serial dilution of each compound was dispensed were mixed at a ratio of 1:1, and a total of 64 mixed drug solutions were prepared with brute-force combinations. The table below shows the maximum concentration (expressed as the final concentration) and geometric ratio of each drug. Each of these mixed drug solutions was added to the plate at 100 μ L/well (N = 1), and further cultured at 37°C under 5% CO₂ conditions for 10 days. Ten days later, 100 μ L of the culture supernatant was removed from each well to leave 100 μ L of the cell culture solution. CellTiter-Glo 2.0 Assay (Promega, Cat#: G9242) was added at 100 μ L/well to the remaining 100 μ L of the cell culture solution. The chemiluminescence measurement was the same as in Test Example 4.

[0896]

Table 87

Other compound having antitumor effect			Biphenyl compound			Tumor cell line
Name	Maximum concentration (nM)	Geometric ratio	Name	Maximum concentration (nM)	Geometric ratio	
Cisplatin	500	2	Example compound 37	30	2	NCI-H1417, NCI-H510A
			Example compound 166	10	2	
			Example compound 177	1	2	
			Example compound 273	1	2	
Carboplatin	25000	2	Example compound 37	30	2	NCI-H510A
			Example compound 166	3	2	
			Example compound 177	1	2	
			Example compound 273	1	2	
SN-38	1	2	Example compound 37	30	2	NCI-H1417, NCI-H510A
			Example compound 166	10	2	
			Example compound 177	1	2	
			Example compound 273	1	2	
Etoposide	100	2	Example compound 37	30	2	NCI-H1417
			Example compound 166	10	2	
			Example compound 177	1	2	
			Example compound 273	1	2	
Paclitaxel	10	2	Example compound 37	30	2	NCI-H1417
	4	1.25	Example compound 166	3	2	
	4		Example compound 177	1	2	
	4		Example compound 273	1	2	

[0897]

The effect potentiated by the combined use of drugs was
5 evaluated according to the methods described in known documents
(Trends Pharmacol. Sci., 4, 450-454, 1983; Pharmacol Rev., 58
(3), 621-81, 2006).

The specific method and the method for determining the
effect of combined use were the same as in Test Example 4.

10 [0898]

The following tables show the results.

[0899]

Table 88

Other antitumor agent : cisplatin, cell line : NCI-H1417				
Example compound 37 (nM)	Other antitumor agent (nM)	Fa	CI	Combination ratio (Example compound 37 : other antitumor agent)
0.938	250	0.52	0.437	1 : 267
0.938	500	0.70	0.195	1 : 533
1.88	250	0.54	0.475	1 : 133
1.88	500	0.75	0.166	1 : 267
3.75	125	0.46	0.784	1 : 33.3
3.75	250	0.61	0.450	1 : 66.7
3.75	500	0.82	0.143	1 : 133
7.50	250	0.73	0.398	1 : 33.3
7.50	500	0.89	0.134	1 : 66.7
15	125	0.74	0.679	1 : 8.33
15	250	0.83	0.409	1 : 16.7
15	500	0.93	0.156	1 : 33.3
30	125	0.87	0.587	1 : 4.17
30	250	0.89	0.492	1 : 8.33
30	500	0.96	0.175	1 : 16.7

[0900]

Table 89

Other antitumor agent : cisplatin, cell line : NCI-H1417				
Example compound 166 (nM)	Other antitumor agent (nM)	Fa	CI	Combination ratio (Example compound 166 : other antitumor agent)
0.313	7.81	0.20	0.830	1 : 25
0.313	15.6	0.26	0.700	1 : 50
0.625	125	0.56	0.696	1 : 200
0.625	250	0.70	0.566	1 : 400
0.625	500	0.88	0.288	1 : 800
1.25	7.81	0.58	0.706	1 : 6.25
1.25	15.6	0.55	0.804	1 : 12.5
1.25	31.3	0.60	0.710	1 : 25
1.25	62.5	0.58	0.838	1 : 50
1.25	125	0.68	0.670	1 : 100
1.25	250	0.80	0.477	1 : 200
1.25	500	0.92	0.243	1 : 400
2.5	7.81	0.70	0.899	1 : 3.12
2.5	15.6	0.73	0.804	1 : 6.25
2.5	31.3	0.71	0.894	1 : 12.5
2.5	62.5	0.73	0.859	1 : 25
2.5	125	0.78	0.737	1 : 50
2.5	250	0.88	0.444	1 : 100
2.5	500	0.94	0.272	1 : 200
5	250	0.90	0.659	1 : 50
5	500	0.95	0.384	1 : 100
10	500	0.96	0.568	1 : 50

[0901]

Table 90

Other antitumor agent : cisplatin, cell line : NCI-H1417				
Example compound 177 (nM)	Other antitumor agent (nM)	Fa	CI	Combination ratio (Example compound 177 : other antitumor agent)
0.0156	31.3	0.19	0.751	1 : 2000
0.0625	500	0.78	0.478	1 : 8000
0.125	250	0.65	0.732	1 : 2000
0.125	500	0.89	0.259	1 : 4000
0.25	62.5	0.68	0.756	1 : 250
0.25	125	0.71	0.716	1 : 500
0.25	250	0.84	0.389	1 : 1000
0.25	500	0.95	0.146	1 : 2000
0.5	15.6	0.79	0.747	1 : 31.3
0.5	31.3	0.80	0.710	1 : 62.5
0.5	62.5	0.78	0.831	1 : 125
0.5	125	0.85	0.535	1 : 250
0.5	250	0.87	0.501	1 : 500
0.5	500	0.95	0.211	1 : 1000
1	125	0.90	0.615	1 : 125
1	250	0.92	0.506	1 : 250
1	500	0.96	0.266	1 : 500

[0902]

Table 91

Other antitumor agent : cisplatin, cell line : NCI-H1417				
Example compound 273 (nM)	Other antitumor agent (nM)	Fa	CI	Combination ratio (Example compound 273 : other antitumor agent)
0.0313	500	0.73	0.321	1 : 16000
0.0625	250	0.55	0.791	1 : 4000
0.0625	500	0.83	0.181	1 : 5000
0.125	250	0.78	0.314	1 : 2000
0.125	500	0.90	0.133	1 : 4000
0.25	7.81	0.70	0.698	1 : 31.3
0.25	15.6	0.68	0.768	1 : 62.5
0.25	31.3	0.73	0.621	1 : 125
0.25	62.5	0.70	0.733	1 : 250
0.25	125	0.78	0.511	1 : 500
0.25	250	0.86	0.315	1 : 1000
0.25	500	0.94	0.133	1 : 2000
0.5	31.3	0.80	0.851	1 : 62.5
0.5	62.5	0.82	0.760	1 : 125
0.5	125	0.82	0.773	1 : 250
0.5	250	0.89	0.460	1 : 500
0.5	500	0.95	0.209	1 : 1000
1	125	0.89	0.889	1 : 125
1	250	0.92	0.647	1 : 250
1	500	0.96	0.328	1 : 500

[0903]

Table 92

Other antitumor agent : cisplatin, cell line : NCI-H510A				
Example compound 37 (nM)	Other antitumor agent (nM)	Fa	CI	Combination ratio (Example compound 37 : other antitumor agent)
0.938	500	0.54	0.169	1 : 533
1.875	250	0.37	0.606	1 : 133
1.875	500	0.68	0.165	1 : 267
3.75	250	0.57	0.496	1 : 66.7
3.75	500	0.80	0.180	1 : 133
7.5	125	0.63	0.775	1 : 16.7
7.5	250	0.75	0.464	1 : 33.3
8	500	0.90	0.172	1 : 66.7
15	250	0.87	0.448	1 : 16.7
15	500	0.93	0.241	1 : 33.3
30	125	0.87	0.896	1 : 4.17
30	250	0.88	0.825	1 : 8.33
30	500	0.96	0.283	1 : 16.7

[0904]

Table 93

Other antitumor agent : cisplatin, cell line : NCI-H510A				
Example compound 166 (nM)	Other antitumor agent (nM)	Fa	CI	Combination ratio (Example compound 166 : other antitumor agent)
0.313	250	0.47	0.483	1 : 800
0.313	500	0.77	0.129	1 : 1600
0.625	7.81	0.42	0.667	1 : 12.5
0.625	15.6	0.36	0.842	1 : 25
0.625	31.3	0.57	0.408	1 : 80
0.625	62.5	0.49	0.563	1 : 100
0.625	125	0.55	0.481	1 : 200
0.625	250	0.69	0.295	1 : 400
0.625	500	0.82	0.162	1 : 800
1.25	7.81	0.61	0.691	1 : 6.25
1.25	15.6	0.62	0.669	1 : 12.5
1.25	31.3	0.66	0.581	1 : 25
1.25	62.5	0.70	0.502	1 : 50
1.25	125	0.74	0.430	1 : 100
1.25	250	0.85	0.241	1 : 200
1.25	500	0.93	0.118	1 : 400
2.5	7.81	0.73	0.871	1 : 3.12
2.5	15.6	0.73	0.872	1 : 6.24
2.5	31.3	0.77	0.730	1 : 12.5
2.5	62.5	0.79	0.664	1 : 25
2.5	125	0.85	0.472	1 : 50
2.5	250	0.90	0.320	1 : 100
2.5	500	0.95	0.171	1 : 200
5	62.5	0.86	0.876	1 : 12.5
5	125	0.87	0.816	1 : 25
5	250	0.91	0.578	1 : 50
5	500	0.96	0.280	1 : 100
10	500	0.96	0.558	1 : 50

[0905]

Table 94

Other antitumor agent : cisplatin, cell line : NCI-H510A				
Example compound 177 (nM)	Other antitumor agent (nM)	Fa	CI	Combination ratio (Example compound 177 : other antitumor agent)
0.0313	250	0.34	0.741	1 : 8000
0.0313	500	0.53	0.342	1 : 16000
0.0625	500	0.74	0.194	1 : 8000
0.125	125	0.58	0.630	1 : 1000
0.125	250	0.68	0.446	1 : 2000
0.125	500	0.89	0.144	1 : 4000
0.25	15.6	0.67	0.880	1 : 62.5
0.25	31.3	0.70	0.786	1 : 125
0.25	62.5	0.73	0.698	1 : 250
0.25	125	0.79	0.533	1 : 500
0.25	250	0.86	0.357	1 : 1000
0.25	500	0.94	0.165	1 : 2000
0.5	62.5	0.85	0.758	1 : 125
0.5	125	0.86	0.710	1 : 250
0.5	250	0.92	0.425	1 : 500
0.5	500	0.96	0.232	1 : 1000
1	250	0.94	0.658	1 : 250
1	500	0.97	0.363	1 : 500

[0906]

Table 95

Other antitumor agent : cisplatin, cell line : NCI-H510A				
Example compound 273 (nM)	Other antitumor agent (nM)	Fa	CI	Combination ratio (Example compound 273 : other antitumor agent)
0.0313	250	0.26	0.895	1 : 8000
0.0313	500	0.57	0.162	1 : 16000
0.0625	250	0.42	0.526	1 : 4000
0.0625	500	0.74	0.164	1 : 8000
0.125	31.3	0.51	0.705	1 : 250
0.125	62.5	0.56	0.603	1 : 500
0.125	125	0.57	0.587	1 : 1000
0.125	250	0.72	0.348	1 : 2000
0.125	500	0.89	0.142	1 : 4000
0.25	7.81	0.69	0.777	1 : 31.3
0.25	15.6	0.71	0.722	1 : 62.5
0.25	31.3	0.66	0.865	1 : 125
0.25	62.5	0.74	0.642	1 : 250
0.25	125	0.80	0.492	1 : 500
0.25	250	0.85	0.375	1 : 1000
0.25	500	0.94	0.170	1 : 2000
0.5	125	0.86	0.704	1 : 250
0.5	250	0.90	0.523	1 : 500
0.5	500	0.96	0.244	1 : 1000
1	500	0.97	0.386	1 : 500

5

[0907]

Table 96

Other antitumor agent : carboplatin, cell line : NCI-H510A				
Example compound 37 (nM)	Other antitumor agent (nM)	Fa	CI	Combination ratio (Example compound 37 : other antitumor agent)
0.938	1563	0.94	0.824	1 : 1670
1.88	1563	0.96	0.716	1 : 831
3.75	781	0.75	0.853	1 : 208
3.75	1563	0.97	0.653	1 : 417
7.5	391	0.64	0.869	1 : 52
7.5	781	0.88	0.648	1 : 104
7.5	1563	0.99	0.447	1 : 208
15	391	0.79	0.773	1 : 26
15	781	0.92	0.611	1 : 52
15	1563	0.99	0.458	1 : 104
30	781	0.95	0.581	1 : 26
30	1563	0.99	0.479	1 : 52

[0908]

Table 97

Other antitumor agent : carboplatin, cell line : NCI-H510A				
Example compound 166 (nM)	Other antitumor agent (nM)	Fa	CI	Combination ratio (Example compound 166 : other antitumor agent)
0.0938	391	0.42	0.880	1 : 4160
0.0938	781	0.71	0.849	1 : 8330
0.0938	1563	0.97	0.622	1 : 16700
0.188	391	0.60	0.747	1 : 2080
0.188	781	0.85	0.647	1 : 4160
0.188	1563	0.99	0.421	1 : 8310
0.375	391	0.76	0.634	1 : 1040
0.375	781	0.91	0.554	1 : 2080
0.375	1563	0.99	0.425	1 : 4170
0.75	391	0.84	0.644	1 : 521
0.75	781	0.93	0.562	1 : 1041
0.75	1563	0.99	0.436	1 : 2080
1.5	391	0.88	0.771	1 : 260
1.5	781	0.95	0.573	1 : 521
1.5	1563	0.99	0.456	1 : 1040
3	781	0.96	0.666	1 : 260

5

[0909]

Table 98

Other antitumor agent : carboplatin, cell line : NCI-H510A				
Example compound 177 (nM)	Other antitumor agent (nM)	Fa	CI	Combination ratio (Example compound 177 : other antitumor agent)
0.0313	3125	0.99	0.880	1 : 100000
0.0625	1563	0.95	0.780	1 : 25000
0.125	781	0.76	0.865	1 : 6250
0.125	1563	0.98	0.581	1 : 12500
0.25	391	0.76	0.743	1 : 1560
0.25	781	0.90	0.670	1 : 3120
0.25	1563	0.99	0.478	1 : 6250
0.5	391	0.83	0.897	1 : 781
0.5	781	0.94	0.664	1 : 1560
0.5	1563	0.99	0.514	1 : 3130
1	781	0.95	0.840	1 : 781
1	1563	0.99	0.587	1 : 1560

[0910]

Table 99

Other antitumor agent : carboplatin, cell line : NCI-H510A				
Example compound 273 (nM)	Other antitumor agent (nM)	Fa	CI	Combination ratio (Example compound 273 : other antitumor agent)
0.0625	1563	0.95	0.790	1 : 250000
0.125	781	0.79	0.868	1 : 6250
0.125	1563	0.98	0.558	1 : 12500
0.25	391	0.78	0.739	1 : 1560
0.25	781	0.90	0.648	1 : 3130
0.25	1563	0.99	0.436	1 : 6250
0.5	391	0.86	0.724	1 : 781
0.5	781	0.94	0.574	1 : 1560
0.5	1563	0.99	0.448	1 : 3130
1	781	0.95	0.650	1 : 781
1	1563	0.99	0.471	1 : 1560

[0911]

Table 100

Other antitumor agent : SN-38, cell line : NCI-H1417				
Example compound 37 (nM)	Other antitumor agent (nM)	Fa	CI	Combination ratio (Example compound 37 : other antitumor agent)
0.938	1	0.968	0.302	1 : 1.07
1.88	1	0.978	0.217	1 : 0.532
3.75	0.5	0.862	0.688	1 : 0.133
3.75	1	0.986	0.148	1 : 0.267
7.5	1	0.992	0.092	1 : 0.133
15	0.0313	0.785	0.805	1 : 0.00209
15	0.0625	0.806	0.743	1 : 0.00417
15	0.125	0.860	0.545	1 : 0.00833
15	0.25	0.907	0.425	1 : 0.0167
15	0.5	0.972	0.179	1 : 0.0333
30	0.0313	0.915	0.420	1 : 0.00104
30	0.0625	0.866	0.813	1 : 0.00208
30	0.125	0.907	0.548	1 : 0.00417
30	0.25	0.954	0.281	1 : 0.00833
30	0.5	0.980	0.160	1 : 0.0167
30	1	0.996	0.056	1 : 0.0333

5

[0912]

Table 101

Other antitumor agent : SN-38, cell line : NCI-H1417				
Example compound 166 (nM)	Other antitumor agent (nM)	Fa	CI	Combination ratio (Example compound 166 : other antitumor agent)
0.313	0.5	0.815	0.715	1 : 1.6
0.313	1	0.983	0.159	1 : 3.19
0.625	0.5	0.910	0.394	1 : 0.8
0.625	1	0.989	0.115	1 : 1.6
1.25	0.125	0.729	0.799	1 : 0.1
1.25	0.25	0.830	0.602	1 : 0.2
1.25	0.5	0.949	0.276	1 : 0.4
1.25	1	0.993	0.084	1 : 0.8
2.5	0.25	0.874	0.659	1 : 0.1
2.5	0.5	0.958	0.300	1 : 0.2
2.5	1	0.993	0.095	1 : 0.4
5	0.5	0.967	0.346	1 : 0.2
5	1	0.994	0.103	1 : 0.2
10	0.5	0.974	0.443	1 : 0.05
10	1	0.995	0.120	1 : 0.1

[0913]

Table 102

Other antitumor agent : SN-38, cell line : NCI-H1417				
Example compound 177 (nM)	Other antitumor agent (nM)	Fa	CI	Combination ratio (Example compound 177 : other antitumor agent)
0.0313	1	0.956	0.227	1 : 32
0.0625	0.5	0.778	0.826	1 : 8
0.0625	1	0.977	0.141	1 : 16
0.125	0.5	0.883	0.498	1 : 4
0.125	1	0.989	0.101	1 : 8
0.25	0.5	0.938	0.421	1 : 2
0.25	1	0.990	0.147	1 : 4
0.5	0.5	0.951	0.598	1 : 1
0.5	1	0.992	0.222	1 : 2
1	1	0.994	0.351	1 : 1

[0914]

5 Table 103

Other antitumor agent : SN-38, cell line : NCI-H1417				
Example compound 273 (nM)	Other antitumor agent (nM)	Fa	CI	Combination ratio (Example compound 273 : other antitumor agent)
0.0313	1	0.960	0.228	1 : 32
0.0625	0.5	0.833	0.546	1 : 8
0.0625	1	0.985	0.098	1 : 16
0.125	0.125	0.625	0.831	1 : 1
0.125	0.25	0.752	0.654	1 : 2
0.125	0.5	0.914	0.322	1 : 4
0.125	1	0.990	0.073	1 : 8
0.25	0.0313	0.764	0.617	1 : 0.125
0.25	0.0625	0.759	0.675	1 : 0.25
0.25	0.125	0.760	0.755	1 : 0.5
0.25	0.25	0.851	0.537	1 : 1
0.25	0.5	0.960	0.199	1 : 2
0.25	1	0.992	0.070	1 : 4
0.5	0.25	0.876	0.721	1 : 0.5
0.5	0.5	0.959	0.295	1 : 1
0.5	1	0.992	0.089	1 : 2
1	0.25	0.935	0.662	1 : 0.25
1	0.5	0.973	0.321	1 : 0.5
1	1	0.993	0.113	1 : 1

[0915]

Table 104

Other antitumor agent : SN-38, cell line : NCI-H510A				
Example compound 37 (nM)	Other antitumor agent (nM)	Fa	CI	Combination ratio (Example compound 37 : other antitumor agent)
0.938	1	0.516	0.337	1 : 1.07
1.88	1	0.615	0.419	1 : 0.532
3.75	1	0.775	0.379	1 : 0.267
7.5	1	0.874	0.371	1 : 0.133
15	0.5	0.860	0.840	1 : 0.0333
15	1	0.937	0.340	1 : 0.0667
30	0.5	0.924	0.835	1 : 0.0167
30	1	0.968	0.328	1 : 0.0333

10 [0916]

Table 105

Other antitumor agent : SN-38, cell line : NCI-H510A				
Example compound 166 (nM)	Other antitumor agent (nM)	Fa	CI	Combination ratio (Example compound 166 : other antitumor agent)
0.313	1	0.629	0.363	1 : 3.19
0.625	0.25	0.503	0.735	1 : 0.4
0.625	0.5	0.551	0.660	1 : 0.8
0.625	1	0.809	0.215	1 : 1.6
1.25	0.0156	0.651	0.702	1 : 0.0125
1.25	0.0313	0.625	0.792	1 : 0.025
1.25	0.0625	0.662	0.674	1 : 0.05
1.25	0.125	0.661	0.686	1 : 0.1
1.25	0.25	0.701	0.581	1 : 0.2
1.25	0.5	0.771	0.419	1 : 0.4
1.25	1	0.872	0.222	1 : 0.8
2.5	0.0156	0.760	0.804	1 : 0.00624
2.5	0.0313	0.758	0.815	1 : 0.0125
2.5	0.0625	0.786	0.692	1 : 0.025
2.5	0.125	0.781	0.719	1 : 0.05
2.5	0.25	0.785	0.711	1 : 0.1
2.5	0.5	0.827	0.551	1 : 0.2
2.5	1	0.934	0.187	1 : 0.4
5	0.25	0.863	0.792	1 : 0.05
5	0.5	0.865	0.789	1 : 0.1
5	1	0.954	0.237	1 : 0.2
10	1	0.959	0.406	1 : 0.1

[0917]

Table 106

Other antitumor agent : SN-38, cell line : NCI-H510A				
Example compound 177 (nM)	Other antitumor agent (nM)	Fa	CI	Combination ratio (Example compound 177 : other antitumor agent)
0.0313	1	0.367	0.205	1 : 32
0.0625	0.5	0.260	0.553	1 : 8
0.0625	1	0.514	0.271	1 : 16
0.125	0.0156	0.361	0.767	1 : 0.125
0.125	0.0313	0.321	0.852	1 : 0.25
0.125	0.0625	0.321	0.853	1 : 0.5
0.125	0.125	0.308	0.886	1 : 1
0.125	0.25	0.403	0.694	1 : 2
0.125	0.5	0.496	0.556	1 : 4
0.125	1	0.785	0.258	1 : 8
0.25	0.0156	0.677	0.714	1 : 0.0625
0.25	0.0313	0.616	0.834	1 : 0.125
0.25	0.0625	0.619	0.828	1 : 0.25
0.25	0.125	0.609	0.849	1 : 0.5
0.25	0.25	0.699	0.672	1 : 1
0.25	0.5	0.725	0.624	1 : 2
0.25	1	0.906	0.293	1 : 4
0.5	0.5	0.855	0.781	1 : 1
0.5	1	0.924	0.512	1 : 2
1	1	0.954	0.750	1 : 1

[0918]

Table 107

Other antitumor agent : SN-38, cell line : NCI-H510A				
Example compound 273 (nM)	Other antitumor agent (nM)	Fa	CI	Combination ratio (Example compound 273 : other antitumor agent)
0.0313	1	0.486	0.255	1 : 32
0.0625	0.5	0.326	0.800	1 : 8
0.0625	1	0.614	0.271	1 : 16
0.125	0.25	0.446	0.886	1 : 2
0.125	0.5	0.607	0.516	1 : 4
0.125	1	0.824	0.204	1 : 8
0.25	0.0625	0.707	0.698	1 : 0.25
0.25	0.125	0.682	0.770	1 : 0.5
0.25	0.25	0.706	0.702	1 : 1
0.25	0.5	0.791	0.483	1 : 2
0.25	1	0.903	0.229	1 : 4
0.5	0.5	0.858	0.654	1 : 1
0.5	1	0.942	0.289	1 : 2
1	1	0.947	0.535	1 : 1

[0919]

Table 108

Other antitumor agent : etoposide, cell line : NCI-H1417				
Example compound 37 (nM)	Other antitumor agent (nM)	Fa	CI	Combination ratio (Example compound 37 : other antitumor agent)
0.938	100	0.56	0.588	1 : 106
1.88	100	0.64	0.472	1 : 53.2
3.75	100	0.68	0.518	1 : 26.7
7.5	50	0.62	0.833	1 : 6.67
7.5	100	0.80	0.423	1 : 13.3
15	50	0.78	0.748	1 : 3.33
15	100	0.89	0.388	1 : 6.67
30	25	0.86	0.875	1 : 0.833
30	50	0.88	0.766	1 : 1.67
30	100	0.95	0.348	1 : 3.33

5

[0920]

Table 109

Other antitumor agent : etoposide, cell line : NCI-H1417				
Example compound 166 (nM)	Other antitumor agent (nM)	Fa	CI	Combination ratio (Example compound 166 : other antitumor agent)
0.156	25	0.35	0.792	1 : 160
0.156	50	0.41	0.729	1 : 320
0.313	50	0.50	0.674	1 : 160
0.313	100	0.65	0.353	1 : 320
0.625	50	0.57	0.819	1 : 80
0.625	100	0.74	0.344	1 : 160
1.25	50	0.79	0.442	1 : 40
2.5	25	0.79	0.862	1 : 10
2.5	50	0.84	0.589	1 : 20
2.5	100	0.92	0.238	1 : 40
5	25	0.88	0.791	1 : 5
5	50	0.88	0.793	1 : 10
5	100	0.94	0.330	1 : 20
10	50	0.93	0.794	1 : 5

[0921]

Table 110

Other antitumor agent : etoposide, cell line : NCI-H1417				
Example compound 177 (nM)	Other antitumor agent (nM)	Fa	CI	Combination ratio (Example compound 177 : other antitumor agent)
0.0625	100	0.62	0.817	1 : 1600
0.125	50	0.60	0.779	1 : 400
0.125	100	0.73	0.697	1 : 800
0.25	50	0.80	0.606	1 : 200
0.25	100	0.89	0.443	1 : 400
0.5	50	0.86	0.786	1 : 100
0.5	100	0.92	0.562	1 : 200

[0922]

Table 111

Other antitumor agent : etoposide, cell line : NCI-H1417				
Example compound 273 (nM)	Other antitumor agent (nM)	Fa	CI	Combination ratio (Example compound 273 : other antitumor agent)
0.0156	50	0.31	0.887	1 : 3200
0.0313	100	0.51	0.531	1 : 3200
0.0625	100	0.63	0.404	1 : 1600
0.125	25	0.48	0.797	1 : 200
0.125	50	0.64	0.527	1 : 400
0.125	100	0.80	0.308	1 : 800
0.25	1.56	0.71	0.728	1 : 6025
0.25	3.13	0.69	0.776	1 : 12.5
0.25	6.25	0.66	0.853	1 : 25
0.25	12.5	0.67	0.836	1 : 50
0.25	25	0.72	0.726	1 : 100
0.25	50	0.80	0.550	1 : 200
0.25	100	0.89	0.349	1 : 400
0.5	50	0.87	0.767	1 : 100
0.5	100	0.93	0.492	1 : 200
1	100	0.95	0.775	1 : 100

5

[0923]

Table 112

Other antitumor agent : paclitaxel, cell line : NCI-H1417				
Example compound 37 (nM)	Other antitumor agent (nM)	Fa	CI	Combination ratio (Example compound 37 : other antitumor agent)
0.469	2.5	0.556	0.455	1 : 5.33
0.469	5	0.910	0.615	1 : 10.7
0.938	2.5	0.548	0.512	1 : 2.67
0.938	5	0.909	0.626	1 : 5.33
1.88	2.5	0.589	0.584	1 : 1.33
1.88	5	0.920	0.632	1 : 2.66
3.75	2.5	0.586	0.777	1 : 0.667
3.75	5	0.931	0.648	1 : 1.333
7.5	2.5	0.769	0.749	1 : 0.333
7.5	5	0.942	0.685	1 : 0.667
15.0	5	0.958	0.723	1 : 0.333
15.0	10	0.987	0.989	1 : 0.667

[0924]

10

Table 113

Other antitumor agent : paclitaxel, cell line : NCI-H1417				
Example compound 166 (nM)	Other antitumor agent (nM)	Fa	CI	Combination ratio (Example compound 166 : other antitumor agent)
0.0469	1.05	0.829	0.365	1 : 22.4
0.0469	2.05	0.882	0.621	1 : 43.70
0.0938	1.05	0.851	0.368	1' : 11.20
0.0938	2.05	0.897	0.607	1 : 21.90
0.0938	4	0.905	1.158	1 : 42.60
0.188	1.05	0.896	0.344	1 : 5.59
0.188	2.05	0.918	0.582	1 ₁ : 10.9
0.188	4	0.921	1.112	1 : 21.3
0.375	1.05	0.574	1.423	1 : 2.80
0.375	2.05	0.933	0.571	1 : 5.47
0.375	4	0.946	1.007	1 : 10.7
0.75	1.64	0.731	1.459	1' : 2.19
0.75	2.05	0.872	0.924	1 : 2.73
1.5	1.05	0.785	1.519	1 : 0.70
1.5	2.05	0.963	0.547	1 : 1.37
1.5	2.56	0.941	0.831	1 : 1.71
1.5	3.2	0.963	0.819	1 : 2.13
1.5	4	0.963	1.021	1 : 2.67
3	1.31	0.805	2.445	1 : 0.44
3	1.64	0.845	1.967	1 : 0.55
3	2.05	0.920	1.146	1' : 0.68
3	4	0.967	1.036	1 : 1.33

[0925]

Table 114

Other antitumor agent : paclitaxel, cell line : NCI-H1417				
Example compound 177 (nM)	Other antitumor agent (nM)	Fa	CI	Combination ratio (Example compound 177 : other antitumor agent)
0.0157	1.05	0.878	0.347	1' : 66.9
0.0157	2.05	0.890	0.661	1 ₁ : 131.0
0.0313	1.05	0.865	0.358	1 : 33.5
0.0313	2.05	0.884	0.660	1 : 65.5
0.0313	4	0.919	1.168	1 : 128.0
0.0625	1.05	0.878	0.370	1 : 16.8
0.0625	2.05	0.893	0.670	1 : 32.8
0.0625	4	0.906	1.255	1 : 64.0
0.125	1.05	0.365	2.248	1 : 8.4
0.125	2.05	0.910	0.652	1 : 16.4
0.125	4	0.924	1.186	1 : 32.0
0.25	1.05	0.924	0.368	1 : 4.2
0.25	1.31	0.621	1.525	1 : 5.24
0.25	1.64	0.685	1.383	1 : 6.56
0.25	2.05	0.849	0.943	1' : 8.20
0.25	4	0.946	1.093	1 : 16.00
0.5	1.05	0.754	1.274	1 : 2.10
0.5	2.05	0.953	0.589	1 : 4.10
0.5	2.56	0.940	0.808	1 : 5.12
0.5	3.2	0.953	0.909	1 : 6.40
0.5	4	0.954	1.129	1 : 8.00
1	1.31	0.832	1.436	1 : 1.31
1	1.64	0.841	1.472	1 : 1.64
1	2.05	0.920	0.946	1 : 2.05
1	4	0.964	1.057	1 : 4.00

[0926]

Table 115

Other antitumor agent : paclitaxel, cell line : NCI-H1417				
Example compound 273 (nM)	Other antitumor agent (nM)	Fa	CI	Combination ratio (Example compound 273 : other antitumor agent)
0.0157	1.05	0.846	0.387	1 : 66.9
0.0157	2.05	0.880	0.694	1 : 130.0
0.0313	1.05	0.844	0.395	1 : 33.5
0.0313	2.05	0.859	0.726	1 : 65.5
0.0313	4	0.895	1.285	1 : 128.0
0.0625	1.05	0.864	0.41	1 : 16.8
0.0625	2.05	0.885	0.711	1 : 32.8
0.0625	4	0.909	1.266	1 : 64.0
0.125	1.05	0.365	2.212	1 : 8.40
0.125	2.05	0.915	0.667	1 : 16.4
0.125	4	0.936	1.144	1 : 32.0
0.25	1.05	0.934	0.382	1 : 4.20
0.25	1.31	0.727	1.170	1 : 5.24
0.25	1.64	0.733	1.307	1 : 6.56
0.25	2.05	0.860	0.985	1 : 8.20
0.25	4	0.949	1.101	1 : 16.00
0.5	1.05	0.729	1.559	1 : 2.10
0.5	2.05	0.948	0.677	1 : 4.10
0.5	2.56	0.934	0.918	1 : 5.12
0.5	3.2	0.948	1.003	1 : 6.40
0.5	4	0.959	1.130	1 : 8.00
1	1.31	0.794	2.169	1 : 1.31
1	1.64	0.804	2.185	1 : 1.64
1	4	0.961	1.182	1 : 4.00

[0927]

These results indicated that the antitumor effects of the biphenyl compounds were synergistically potentiated when used in combination with a platinum-based drug (cisplatin or carboplatin) or a plant alkaloid-based drug (SN-38 (active metabolite of irinotecan)), etoposide, or paclitaxel).

[0928]

10 Test Example 6: Combination of Biphenyl Compounds and Other Antitumor agents Using Human AML Cell Line (in vitro)

The following tables show the source of reagents, the source of tumor cell lines, the media used, and the number of seeded cells.

15 [0929]

Table 116

Reagent	Source
Fetal bovine serum (FBS)	Biological Industries
RPMI-1640 medium	GIBCO
Guadecitabine	AdooQ Bioscience

RG7388	ChemScene

[0930]

Table 117

Tumor cell line (origin)	Source of cell line	Medium	Number of seeded cells per well (cells)
MOLM-13 (human AML cell line)	DSMZ	RPMI-1640 medium containing 10% FBS	100

[0931]

Cells were seeded at 100 μ L/well in a 96-well culture plate (NUNC). The plate seeded with cells was cultured in an incubator set at 37°C and 5% CO₂. On the day of seeding, drug solutions containing a combination of biphenyl compounds and other compounds having an antitumor effect mixed at various concentrations were each added to the cells. Specifically, using an RPMI-1640 medium containing 10% FBS, eight-step serial dilutions (including 0 nM) of the biphenyl compounds (Example compounds 37, 166, 177, and 273) were prepared, and eight-step serial dilutions (including 0 nM) of other compounds having an antitumor effect were also prepared. The table below shows the maximum concentration (expressed as the final concentration) and geometric ratio of each drug.

[0932]

The procedures after production of the serial dilutions were the same as in Test Example 5.

[0933]

Table 118

Other compound having antitumor effect			Biphenyl compound			Tumor cell line
Name	Maximum concentration (nM)	Geometric ratio	Name	Maximum concentration (nM)	Geometric ratio	
RG7388	15	1.5	Example compound 37	50	2	MOLM-13
			Example compound 166	5	2	
			Example compound 177	1	2	
			Example compound 273	1	2	
Guadecitabine	10	2	Example compound 37	100	2	MOLM-13
			Example compound 166	10	2	
			Example compound 177	3	2	
			Example compound 273	3	2	

[0934]

The effect potentiated by the combined use of drugs was evaluated according to the methods described in known documents (Trends Pharmacol. Sci., 4, 450-454, 1983; Pharmacol Rev., 58 (3), 621-81, 2006).

The specific method and the method for determining the effect of combined use were the same as in Test Example 4.

10 [0935]

The following tables show the results.

[0936]

Table 119

Other antitumor agent : RG7388, cell line : MOLM-13				
Example compound 37 (nM)	Other antitumor agent (nM)	Fa	CI	Combination ratio (Example compound 37 : other antitumor agent)
3.13	2.63	0.373	0.683	1 : 0.842
6.25	5.27	0.801	0.465	1 : 0.843

15 [0937]

Table 120

Other antitumor agent : RG7388, cell line : MOLM-13				
Example compound 166 (nM)	Other antitumor agent (nM)	Fa	CI	Combination ratio (Example compound 166 : other antitumor agent)
0.156	1.32	0.244	0.605	1 : 8.46
0.313	2.63	0.572	0.553	1 : 8.4
0.625	5.26	0.927	0.320	1 : 8.42

[0938]

Table 121

Other antitumor agent : RG7388, cell line : MOLM-13				
Example compound 177 (nM)	Other antitumor agent (nM)	Fa	CI	Combination ratio (Example compound 177 : other antitumor agent)
0.125	2.63	0.797	0.209	1 : 21

20

[0939]

Table 122

Other antitumor agent : RG7388, cell line : MOLM-13				
Example compound 273 (nM)	Other antitumor agent (nM)	Fa	CI	Combination ratio (Example compound 273 : other antitumor agent)
0.0625	2.63	0.477	0.712	1 : 42.1

[0940]

5 Table 123

Other antitumor agent : guadecitabine, cell line : MOLM-13				
Example compound 37 (nM)	Other antitumor agent (nM)	Fa	CI	Combination ratio (Example compound 37 : other antitumor agent)
1.56	2.5	0.290	0.872	1 : 1.6
1.56	5	0.734	0.807	1 : 3.2
3.13	2.5	0.648	0.520	1 : 0.8
3.13	5	0.749	0.810	1 : 1.6
3.13	10	0.992	0.452	1 : 3.2
6.25	1.25	0.592	0.425	1 : 0.2
6.25	2.5	0.706	0.526	1 : 0.4
6.25	5	0.833	0.705	1 : 0.8
6.25	10	0.995	0.388	1 : 1.6
12.5	1.25	0.680	0.475	1 : 0.1
12.5	2.5	0.969	0.201	1 : 0.2
12.5	5	0.987	0.275	1 : 0.4
12.5	10	0.998	0.288	1 : 0.8
25	2.5	0.898	0.415	1 : 0.1
25	5	0.997	0.169	1 : 0.2
25	10	0.993	0.443	1 : 0.4
50	1.25	0.899	0.426	1 : 0.025
50	2.5	0.968	0.277	1 : 0.05
50	5	0.963	0.488	1 : 0.1
100	1.25	0.969	0.280	1 : 0.0125
100	2.5	0.997	0.105	1 : 0.025
100	5	0.999	0.123	1 : 0.05

[0941]

Table 124

Other antitumor agent : guadecitabine, cell line : MOLM-13				
Example compound 166 (nM)	Other antitumor agent (nM)	Fa	CI	Combination ratio (Example compound 166: other antitumor agent)
0.156	2.5	0.543	0.786	1 : 16
0.156	5	0.829	0.311	1 : 32
0.313	5	0.870	0.227	1 : 16
0.313	10	0.985	0.037	1 : 32
0.625	1.25	0.519	0.892	1 : 2
0.625	2.5	0.729	0.412	1 : 4
0.625	5	0.931	0.111	1 : 8
0.625	10	0.992	0.019	1 : 16
1.25	1.25	0.750	0.349	1 : 1
1.25	2.5	0.935	0.068	1 : 2
1.25	5	0.950	0.083	1 : 4
1.25	10	0.999	0.002	1 : 8
2.5	0.625	0.679	0.848	1 : 0.25
2.5	1.25	0.702	0.806	1 : 0.5
2.5	2.5	0.964	0.041	1 : 1
2.5	5	0.981	0.030	1 : 2
2.5	10	0.998	0.004	1 : 4
5	0.625	0.785	0.720	1 : 0.125
5	1.25	0.955	0.063	1 : 0.25
5	2.5	0.937	0.127	1 : 0.5
5	5	0.991	0.014	1 : 1
5	10	0.999	0.002	1 : 2
10	1.25	0.962	0.087	1 : 0.125
10	2.5	0.996	0.005	1 : 0.25
10	5	0.991	0.019	1 : 0.5

[0942]

Table 125

Other antitumor agent : guadecitabine, cell line : MOLM-13				
Example compound 177 (nM)	Other antitumor agent (nM)	Fa	CI	Combination ratio (Example compound 177: other antitumor agent)
0.0469	2.5	0.555	0.602	1 : 53.3
0.0469	5	0.735	0.518	1 : 107
0.0934	2.5	0.611	0.549	1 : 26.8
0.0934	5	0.836	0.308	1 : 53.5
0.0934	10	0.980	0.065	1 : 107
0.188	0.625	0.395	0.895	1 : 3.32
0.188	1.25	0.542	0.631	1 : 6.65
0.188	2.5	0.690	0.486	1 : 13.3
0.188	5	0.961	0.075	1 : 26.6
0.188	10	0.998	0.007	1 : 53.2
0.375	1.25	0.679	0.553	1 : 3.33
0.375	2.5	0.928	0.119	1 : 6.67
0.375	5	0.990	0.024	1 : 13.3
0.375	10	0.999	0.004	1 : 26.7
0.75	2.5	0.991	0.022	1 : 3.33
0.75	5	0.990	0.032	1 : 6.67
1.5	1.25	0.880	0.500	1 : 0.833
1.5	2.5	0.879	0.552	1 : 1.67
3	2.5	0.950	0.384	1 : 0.833

[0943]

Table 126

Other antitumor agent : guadecitabine, cell line : MOLM-13				
Example compound 273 (nM)	Other antitumor agent (nM)	Fa	CI	Combination ratio (Example compound 273:other antitumor agent)
0.0469	5	0.854	0.806	1 : 107
0.0934	5	0.910	0.492	1 : 53.3
0.0934	10	0.987	0.157	1 : 107
0.188	1.25	0.631	0.809	1 : 6.67
0.188	2.5	0.936	0.180	1 : 13.3
0.188	5	0.972	0.161	1 : 26.7
0.188	10	0.999	0.015	1 : 53.3
0.375	0.156	0.576	0.684	1 : 0.417
0.375	0.313	0.712	0.321	1 : 0.833
0.375	0.625	0.701	0.460	1 : 1.67
0.375	1.25	0.653	0.895	1 : 3.33
0.375	2.5	0.838	0.506	1 : 6.67
0.375	5	0.976	0.140	1 : 13.3
0.375	10	0.998	0.028	1 : 26.7
0.75	0.156	0.665	0.683	1 : 0.208
0.75	0.313	0.794	0.271	1 : 0.417
0.75	0.625	0.745	0.506	1 : 0.833
0.75	1.25	0.727	0.780	1 : 1.67
0.75	2.5	0.897	0.327	1 : 3.33
0.75	5	0.973	0.158	1 : 6.67
0.75	10	0.999	0.015	1 : 13.3
1.5	2.5	0.964	0.115	1 : 1.67
1.5	5	0.994	0.039	1 : 3.33
3	2.5	0.958	0.154	1 : 0.833
3	5	0.993	0.046	1 : 1.67
3	10	0.999	0.015	1 : 3.33

[0944]

These results indicated that the antitumor effects of the biphenyl compounds were synergistically potentiated when used in combination with a human MDM2 (HDM2) inhibitor (RG7388) or a DNA methylation inhibitor (guadecitabine).


[0945]

As described above, the biphenyl compounds represented by Formula (I) showed an effect of potentiating their antitumor effects when used in combination with various other antitumor agents.

CLAIMS

[Claim 1]

An antitumor formulation comprising a biphenyl compound represented by Formula (I) or a salt thereof and one or more
5 other antitumor agents that are used in combination:



wherein

ring A represents a monocyclic, bridged cyclic, or spirocyclic 4-
to 14-membered nitrogen-containing saturated heterocyclic group
10 having 1 to 3 nitrogen atoms, 0 to 1 sulfur atoms, and 0 to 2
oxygen atoms as heteroatoms;

ring B represents monocyclic or bicyclic 5- to 14-membered
unsaturated hydrocarbon or a monocyclic or bicyclic 5- to 14-
membered unsaturated heterocyclic group that may be substituted
15 with oxo, that has 0 to 4 nitrogen atoms, 0 to 2 sulfur atoms,
and 0 to 3 oxygen atoms as heteroatoms, and that has at least one
of nitrogen, sulfur, and oxygen;

R1 represents nitro or cyano;

R2 represents halogen;

20 R3 represents amino, mono- or di(C1-C6 alkyl)amino, (C3-C7
cycloalkyl)amino, or C1-C6 alkyl;

R4 represents halogen, nitro, cyano, carboxy, substituted or
unsubstituted C1-C8 alkyl, substituted or unsubstituted C2-C6
alkenyl, substituted or unsubstituted C1-C6 alkoxy, substituted
25 or unsubstituted C3-C7 cycloalkyl, mono- or di(C1-C6 alkyl)amino,
or substituted or unsubstituted carbamoyl;


when at least one R4 represents substituted C1-C8 alkyl,
substituted C2-C6 alkenyl, substituted C1-C6 alkoxy, substituted
C3-C7 cycloalkyl, or substituted carbamoyl, the substituent is

- halogen, carboxy, C1-C6 alkoxy, hydroxy, C1-C6 alkyl that may be substituted with hydroxy, monocyclic 5- to 10-membered unsaturated hydrocarbon, carbamoyl that may be substituted with C1-C6 alkyl or monocyclic 5- to 10-membered unsaturated
- 5 hydrocarbon, (C2-C7 acyl)oxy, amino that may be substituted with C1-C6 alkyl or C2-C7 acyl, C3-C7 cycloalkyl that may be substituted with hydroxy, or (C1-C6 alkoxy) (C1-C6 alkyl); when two or more of the substituents are present, the substituents may be identical or different;
- 10 R5 and R6 may be identical or different, and each represent hydrogen or C1-C6 alkyl; R5 and R6 may otherwise form oxo or thioxo together;
- when R5 and R6 are identical or different, and each represent hydrogen or C1-C6 alkyl, at least one of R5 and R6 is hydrogen;
- 15 l represents an integer of 0 to 2;
- m represents an integer of 0 to 2;
- n represents an integer of 0 to 5;
- when l is 2, two R2s may be identical or different;
- when m is 2, two R3s may be identical or different;
- 20 when n is 2 to 5, two to five R4s may be identical or different.

[Claim 2]

- The antitumor formulation according to claim 1, wherein the biphenyl compound satisfies the following conditions in
- 25 Formula (I):

ring A represents pyrrolidinyl, piperidinyl, piperazinyl, azepanyl, diazepanyl,



- , 2,7-diazaspiro[3.4]octanyl, 3,7-diazaspiro[3.4]octanyl, 2,7-diazaspiro[3.5]nonanyl, 2,8-diazaspiro[3.5]nonanyl, 3,7-diazaspiro[3.5]nonanyl, 3,8-diazaspiro[4.4]nonanyl, 3,8-
- 30

diazaspiro[4.5]decanyl, or 9-oxa-diazaspiro[3.5]nonanyl;
ring B represents phenyl, naphthyl, pyridyl, pyrazolopyridyl,
pyrazolopyrimidinyl, indolyl, indolinyl, 2-oxo-indolinyl,
indazolyl, benzoimidazolyl, benzoisoxazolyl, benzothiazolyl,
5 benzotriazolyl, imidazopyridinyl, quinolinyl, isoquinolinyl,
quinoxalinyl, quinazolinyl, phthalazinyl, 2-oxo-2,3-
dihydrobenzo[d]oxazolyl, 1,3-dihydroisobenzofuranyl,
dihydrobenzooxazinyl, benzodioxolyl, dihydrobenzodioxynyl, or 2-
oxo-2,3-dihydrobenzo[d]thiazolyl;

10 R1 represents nitro or cyano;
R2 represents fluorine, and is present at the ortho position
relative to R1 on the phenyl;
R3 represents amino, methylamino, ethylamino, isopropylamino,
dimethylamino, cyclobutylamino, or methyl (when two or more R3s
15 are present, R3s may be identical or different);
R4 represents fluorine, chlorine, bromine, iodine, nitro, cyano,
carboxy, methyl, ethyl, n-propyl, isopropyl, tert-butyl,
difluoromethyl, trifluoromethyl, fluoroethyl, aminoethyl,
hydroxymethyl, hydroxyethyl, hydroxypropyl, hydroxydimethylethyl,
20 hydroxymethylpropyl, hydroxymethylbutyl, hydroxyethylbutyl,
carboxymethyl, carbamoylmethyl, methylcarbamoylmethyl,
dimethylcarbamoylmethyl, acetylaminoethyl, methoxyethyl,
hydroxycyclopropylmethyl, hydroxycyclopropylethyl,
hydroxycyclobutylmethyl, methylcarbonyloxyethyl, isobutenyl,
25 methoxy, hydroxypropoxy, cyclopropyl, hydroxymethyl cyclopropyl,
methoxymethyl cyclopropyl, hydroxycyclopropyl cyclopropyl,
phenylcarbamoyl cyclopropyl, benzyloxy, dimethylamino, carbamoyl,
methylcarbamoyl, or dimethylcarbamoyl (when two or more R4s are
present, R4s may be identical or different);

30 R5 and R6 may be identical or different, and each represent
hydrogen or C1-C6 alkyl; R5 and R6 may otherwise form oxo or
thioxo together;
when R5 and R6 are identical or different, and each represent
hydrogen or C1-C6 alkyl, at least one of R5 and R6 is hydrogen;

35 n represents an integer of 0 to 3; and

when n is 2 to 3, two to three R4s may be identical or different.

[Claim 3]

The antitumor formulation according to claim 1 or 2,
5 wherein the biphenyl compound is selected from the following (1)
to (7):
(1) 4-[5-[(3S)-3-aminopyrrolidine-1-carbonyl]-2-[2-fluoro-4-(2-
hydroxy-2-methyl-propyl)phenyl]phenyl]-2-fluoro-benzonitrile;
(2) 4-[5-[(3-endo)-3-amino-8-azabicyclo[3.2.1]octane-8-carbonyl]-
10 2-[2-fluoro-4-(2-hydroxy-2-methyl-propyl)phenyl]phenyl]-2-fluoro-
benzonitrile;
(3) 5'-((1S,2S,4R)-rel-2-amino-7-azabicyclo[2.2.1]heptane-7-
carbonyl)-2'',3-difluoro-4''-(2-hydroxy-2-methylpropyl)-
[1,1':2',1''-terphenyl]-4-carbonitrile-isomer-B;
15 (4) 5'-((1S,2S,4R)-rel-2-amino-7-azabicyclo[2.2.1]heptane-7-
carbonyl)-2'-(6,7-difluoro-1-(2-hydroxy-2-methylpropyl)-1H-
benzo[d][1,2,3]triazol-5-yl)-3-fluoro-[1,1'-biphenyl]-4-
carbonitrile-isomer-B;
(5) 5'-((1S,2S,4R)-rel-2-amino-7-azabicyclo[2.2.1]heptane-7-
20 carbonyl)-2'-(1-(2-ethyl-2-hydroxybutyl)-6,7-difluoro-1H-
benzo[d][1,2,3]triazol-5-yl)-3-fluoro-[1,1'-biphenyl]-4-
carbonitrile-isomer-X;
(6) 5'-((1S,2S,4R)-rel-2-amino-7-azabicyclo[2.2.1]heptane-7-
carbonyl)-2'-(7-bromo-6-fluoro-1-(2-hydroxy-2-methylpropyl)-1H-
25 benzo[d][1,2,3]triazol-5-yl)-3-fluoro-[1,1'-biphenyl]-4-
carbonitrile-isomer-X;
(7) 5'-(((1S,2S,4R)-rel-2-amino-7-azabicyclo[2.2.1]heptan-7-
yl)methyl)-2'-(6,7-difluoro-1-(2-hydroxy-2-methylpropyl)-1H-
benzo[d][1,2,3]triazol-5-yl)-3-fluoro-[1,1'-biphenyl]-4-
30 carbonitrile-isomer-X.

[Claim 4]

The antitumor formulation according to any one of
claims 1 to 3, wherein the one or more other antitumor agents are
35 at least one member selected from antimetabolites, antitumor


antibiotics, molecular target drugs, platinum-based drugs, and plant alkaloid-based drugs.

[Claim 5]

5 The antitumor formulation according to any one of claims 1 to 3, wherein the one or more other antitumor agents are at least one member selected from cytarabine, azacitidine, decitabine, guadecitabine, daunorubicin, tretinoin (ATRA), RG7388, carboplatin, cisplatin, paclitaxel, irinotecan (SN-38),
10 and etoposide.

[Claim 6]

 An antitumor effect potentiator for one or more other antitumor agentst, the antitumor effect potentiator comprising as
15 an active ingredient a biphenyl compound represented by Formula (I) or a salt thereof:



wherein

ring A represents a monocyclic, bridged cyclic, or spirocyclic 4-
20 to 14-membered nitrogen-containing saturated heterocyclic group having 1 to 3 nitrogen atoms, 0 to 1 sulfur atoms, and 0 to 2 oxygen atoms as heteroatoms;

ring B represents monocyclic or bicyclic 5- to 14-membered unsaturated hydrocarbon or a monocyclic or bicyclic 5- to 14-
25 membered unsaturated heterocyclic group that may be substituted with oxo, that has 0 to 4 nitrogen atoms, 0 to 2 sulfur atoms, and 0 to 3 oxygen atoms as heteroatoms, and that has at least one of nitrogen, sulfur, and oxygen;

R1 represents nitro or cyano;

R2 represents halogen;


R3 represents amino, mono- or di(C1-C6 alkyl)amino, (C3-C7 cycloalkyl)amino, or C1-C6 alkyl;

R4 represents halogen, nitro, cyano, carboxy, substituted or
5 unsubstituted C1-C8 alkyl, substituted or unsubstituted C2-C6
alkenyl, substituted or unsubstituted C1-C6 alkoxy, substituted
or unsubstituted C3-C7 cycloalkyl, mono- or di(C1-C6 alkyl)amino,
or substituted or unsubstituted carbamoyl;
when at least one R4 represents substituted C1-C8 alkyl,
10 substituted C2-C6 alkenyl, substituted C1-C6 alkoxy, substituted
C3-C7 cycloalkyl, or substituted carbamoyl, the substituent is
halogen, carboxy, C1-C6 alkoxy, hydroxy, C1-C6 alkyl that may be
substituted with hydroxy, monocyclic 5- to 10-membered
unsaturated hydrocarbon, carbamoyl that may be substituted with
15 C1-C6 alkyl or monocyclic 5- to 10-membered unsaturated
hydrocarbon, (C2-C7 acyl)oxy, amino that may be substituted with
C1-C6 alkyl or C2-C7 acyl, C3-C7 cycloalkyl that may be
substituted with hydroxy, or (C1-C6 alkoxy)(C1-C6 alkyl); when
two or more of the substituents are present, the substituents may
20 be identical or different;
R5 and R6 may be identical or different, and each represent
hydrogen or C1-C6 alkyl; R5 and R6 may otherwise form oxo or
thioxo together;
when R5 and R6 are identical or different, and each represent
25 hydrogen or C1-C6 alkyl, at least one of R5 and R6 is hydrogen;
l represents an integer of 0 to 2;
m represents an integer of 0 to 2;
n represents an integer of 0 to 5;
when l is 2, two R2s may be identical or different;
30 when m is 2, two R3s may be identical or different;
when n is 2 to 5, two to five R4s may be identical or different.

[Claim 7]

An antitumor agent comprising a biphenyl compound or a
35 salt thereof for treating a cancer patient administered one or

more other antitumor agents, wherein the biphenyl compound is represented by Formula (I):



wherein

- 5 ring A represents a monocyclic, bridged cyclic, or spirocyclic 4- to 14-membered nitrogen-containing saturated heterocyclic group having 1 to 3 nitrogen atoms, 0 to 1 sulfur atoms, and 0 to 2 oxygen atoms as heteroatoms;
- ring B represents monocyclic or bicyclic 5- to 14-membered
- 10 unsaturated hydrocarbon or a monocyclic or bicyclic 5- to 14-membered unsaturated heterocyclic group that may be substituted with oxo, that has 0 to 4 nitrogen atoms, 0 to 2 sulfur atoms, and 0 to 3 oxygen atoms as heteroatoms, and that has at least one of nitrogen, sulfur, and oxygen;
- 15 R1 represents nitro or cyano;
- R2 represents halogen;
- R3 represents amino, mono- or di(C1-C6 alkyl)amino, (C3-C7 cycloalkyl)amino, or C1-C6 alkyl;
- R4 represents halogen, nitro, cyano, carboxy, substituted or
- 20 unsubstituted C1-C8 alkyl, substituted or unsubstituted C2-C6 alkenyl, substituted or unsubstituted C1-C6 alkoxy, substituted or unsubstituted C3-C7 cycloalkyl, mono- or di(C1-C6 alkyl)amino, or substituted or unsubstituted carbamoyl;
- when at least one R4 represents substituted C1-C8 alkyl,
- 25 substituted C2-C6 alkenyl, substituted C1-C6 alkoxy, substituted C3-C7 cycloalkyl, or substituted carbamoyl, the substituent is halogen, carboxy, C1-C6 alkoxy, hydroxy, C1-C6 alkyl that may be substituted with hydroxy, monocyclic 5- to 10-membered unsaturated hydrocarbon, carbamoyl that may be substituted with

C1-C6 alkyl or monocyclic 5- to 10-membered unsaturated hydrocarbon, (C2-C7 acyl)oxy, amino that may be substituted with C1-C6 alkyl or C2-C7 acyl, C3-C7 cycloalkyl that may be substituted with hydroxy, or (C1-C6 alkoxy) (C1-C6 alkyl); when
 5 two or more of the substituents are present, the substituents may be identical or different;

R5 and R6 may be identical or different, and each represent hydrogen or C1-C6 alkyl; R5 and R6 may otherwise form oxo or thioxo together;

10 when R5 and R6 are identical or different, and each represent hydrogen or C1-C6 alkyl, at least one of R5 and R6 is hydrogen;

l represents an integer of 0 to 2;

m represents an integer of 0 to 2;

n represents an integer of 0 to 5;


15 when l is 2, two R2s may be identical or different;

when m is 2, two R3s may be identical or different;

when n is 2 to 5, two to five R4s may be identical or different.

[Claim 8]

20 An antitumor formulation comprising a combination of a biphenyl compound or a salt thereof and one or more other antitumor agents, wherein the biphenyl compound is represented by Formula (I):



25 wherein

ring A represents a monocyclic, bridged cyclic, or spirocyclic 4- to 14-membered nitrogen-containing saturated heterocyclic group having 1 to 3 nitrogen atoms, 0 to 1 sulfur atoms, and 0 to 2 oxygen atoms as heteroatoms;

ring B represents monocyclic or bicyclic 5- to 14-membered unsaturated hydrocarbon or a monocyclic or bicyclic 5- to 14-membered unsaturated heterocyclic group that may be substituted with oxo, that has 0 to 4 nitrogen atoms, 0 to 2 sulfur atoms,
5 and 0 to 3 oxygen atoms as heteroatoms, and that has at least one of nitrogen, sulfur, and oxygen;

R1 represents nitro or cyano;

R2 represents halogen;

R3 represents amino, mono- or di(C1-C6 alkyl)amino, (C3-C7
10 cycloalkyl)amino, or C1-C6 alkyl;

R4 represents halogen, nitro, cyano, carboxy, substituted or unsubstituted C1-C8 alkyl, substituted or unsubstituted C2-C6 alkenyl, substituted or unsubstituted C1-C6 alkoxy, substituted or unsubstituted C3-C7 cycloalkyl, mono- or di(C1-C6 alkyl)amino,
15 or substituted or unsubstituted carbamoyl;

when at least one R4 represents substituted C1-C8 alkyl, substituted C2-C6 alkenyl, substituted C1-C6 alkoxy, substituted C3-C7 cycloalkyl, or substituted carbamoyl, the substituent is halogen, carboxy, C1-C6 alkoxy, hydroxy, C1-C6 alkyl that may be
20 substituted with hydroxy, monocyclic 5- to 10-membered unsaturated hydrocarbon, carbamoyl that may be substituted with C1-C6 alkyl or monocyclic 5- to 10-membered unsaturated hydrocarbon, (C2-C7 acyl)oxy, amino that may be substituted with C1-C6 alkyl or C2-C7 acyl, C3-C7 cycloalkyl that may be
25 substituted with hydroxy, or (C1-C6 alkoxy)(C1-C6 alkyl); when two or more of the substituents are present, the substituents may be identical or different;

R5 and R6 may be identical or different, and each represent hydrogen or C1-C6 alkyl; R5 and R6 may otherwise form oxo or
30 thioxo together;

when R5 and R6 are identical or different, and each represent hydrogen or C1-C6 alkyl, at least one of R5 and R6 is hydrogen;

l represents an integer of 0 to 2;


m represents an integer of 0 to 2;

35 n represents an integer of 0 to 5;

when l is 2, two R2s may be identical or different;
 when m is 2, two R3s may be identical or different;
 when n is 2 to 5, two to five R4s may be identical or different.

5 [Claim 9]

A pharmaceutical composition for the prevention and/or treatment of a tumor, the composition comprising a biphenyl compound represented by Formula (I) or a salt thereof and one or more other antitumor agents:



10

wherein

ring A represents a monocyclic, bridged cyclic, or spirocyclic 4- to 14-membered nitrogen-containing saturated heterocyclic group having 1 to 3 nitrogen atoms, 0 to 1 sulfur atoms, and 0 to 2

15 oxygen atoms as heteroatoms;

ring B represents monocyclic or bicyclic 5- to 14-membered unsaturated hydrocarbon or a monocyclic or bicyclic 5- to 14-membered unsaturated heterocyclic group that may be substituted with oxo, that has 0 to 4 nitrogen atoms, 0 to 2 sulfur atoms, and 0 to 3 oxygen atoms as heteroatoms, and that has at least one of nitrogen, sulfur, and oxygen;

20

R1 represents nitro or cyano;

R2 represents halogen;

R3 represents amino, mono- or di(C1-C6 alkyl)amino, (C3-C7 cycloalkyl)amino, or C1-C6 alkyl;

25


R4 represents halogen, nitro, cyano, carboxy, substituted or unsubstituted C1-C8 alkyl, substituted or unsubstituted C2-C6 alkenyl, substituted or unsubstituted C1-C6 alkoxy, substituted or unsubstituted C3-C7 cycloalkyl, mono- or di(C1-C6 alkyl)amino,

or substituted or unsubstituted carbamoyl;
when at least one R4 represents substituted C1-C8 alkyl,
substituted C2-C6 alkenyl, substituted C1-C6 alkoxy, substituted
C3-C7 cycloalkyl, or substituted carbamoyl, the substituent is
5 halogen, carboxy, C1-C6 alkoxy, hydroxy, C1-C6 alkyl that may be
substituted with hydroxy, monocyclic 5- to 10-membered
unsaturated hydrocarbon, carbamoyl that may be substituted with
C1-C6 alkyl or monocyclic 5- to 10-membered unsaturated
hydrocarbon, (C2-C7 acyl)oxy, amino that may be substituted with
10 C1-C6 alkyl or C2-C7 acyl, C3-C7 cycloalkyl that may be
substituted with hydroxy, or (C1-C6 alkoxy) (C1-C6 alkyl); when
two or more of the substituents are present, the substituents may
be identical or different;
R5 and R6 may be identical or different, and each represent
15 hydrogen or C1-C6 alkyl; R5 and R6 may otherwise form oxo or
thioxo together;
when R5 and R6 are identical or different, and each represent
hydrogen or C1-C6 alkyl, at least one of R5 and R6 is hydrogen;
l represents an integer of 0 to 2;
20 m represents an integer of 0 to 2;
n represents an integer of 0 to 5;
when l is 2, two R2s may be identical or different;
when m is 2, two R3s may be identical or different;
when n is 2 to 5, two to five R4s may be identical or different.

25

[Claim 10]

A biphenyl compound represented by Formula (I) or a
salt thereof for potentiating an antitumor effect of one or more
other antitumor agents:



wherein

ring A represents a monocyclic, bridged cyclic, or spirocyclic 4- to 14-membered nitrogen-containing saturated heterocyclic group having 1 to 3 nitrogen atoms, 0 to 1 sulfur atoms, and 0 to 2 oxygen atoms as heteroatoms;

ring B represents monocyclic or bicyclic 5- to 14-membered unsaturated hydrocarbon or a monocyclic or bicyclic 5- to 14-membered unsaturated heterocyclic group that may be substituted with oxo, that has 0 to 4 nitrogen atoms, 0 to 2 sulfur atoms, and 0 to 3 oxygen atoms as heteroatoms, and that has at least one of nitrogen, sulfur, and oxygen;

R1 represents nitro or cyano;

R2 represents halogen;

R3 represents amino, mono- or di(C1-C6 alkyl)amino, (C3-C7 cycloalkyl)amino, or C1-C6 alkyl;

R4 represents halogen, nitro, cyano, carboxy, substituted or unsubstituted C1-C8 alkyl, substituted or unsubstituted C2-C6 alkenyl, substituted or unsubstituted C1-C6 alkoxy, substituted or unsubstituted C3-C7 cycloalkyl, mono- or di(C1-C6 alkyl)amino, or substituted or unsubstituted carbamoyl;

when at least one R4 represents substituted C1-C8 alkyl, substituted C2-C6 alkenyl, substituted C1-C6 alkoxy, substituted C3-C7 cycloalkyl, or substituted carbamoyl, the substituent is halogen, carboxy, C1-C6 alkoxy, hydroxy, C1-C6 alkyl that may be substituted with hydroxy, monocyclic 5- to 10-membered unsaturated hydrocarbon, carbamoyl that may be substituted with C1-C6 alkyl or monocyclic 5- to 10-membered unsaturated hydrocarbon, (C2-C7 acyl)oxy, amino that may be substituted with

C1-C6 alkyl or C2-C7 acyl, C3-C7 cycloalkyl that may be substituted with hydroxy, or (C1-C6 alkoxy) (C1-C6 alkyl); when two or more of the substituents are present, the substituents may be identical or different;

5 R5 and R6 may be identical or different, and each represent hydrogen or C1-C6 alkyl; R5 and R6 may otherwise form oxo or thioxo together;

when R5 and R6 are identical or different, and each represent hydrogen or C1-C6 alkyl, at least one of R5 and R6 is hydrogen;

10 l represents an integer of 0 to 2;

m represents an integer of 0 to 2;

n represents an integer of 0 to 5;


when l is 2, two R2s may be identical or different;

when m is 2, two R3s may be identical or different;

15 when n is 2 to 5, two to five R4s may be identical or different.

[Claim 11]

Use of a biphenyl compound represented by Formula (I) or a salt thereof for potentiating an antitumor effect of one or
20 more other antitumor agents:



wherein

ring A represents a monocyclic, bridged cyclic, or spirocyclic 4- to 14-membered nitrogen-containing saturated heterocyclic group
25 having 1 to 3 nitrogen atoms, 0 to 1 sulfur atoms, and 0 to 2 oxygen atoms as heteroatoms;

ring B represents monocyclic or bicyclic 5- to 14-membered unsaturated hydrocarbon or a monocyclic or bicyclic 5- to 14-membered unsaturated heterocyclic group that may be substituted

with oxo, that has 0 to 4 nitrogen atoms, 0 to 2 sulfur atoms, and 0 to 3 oxygen atoms as heteroatoms, and that has at least one of nitrogen, sulfur, and oxygen;

R1 represents nitro or cyano;

5 R2 represents halogen;

R3 represents amino, mono- or di(C1-C6 alkyl)amino, (C3-C7 cycloalkyl)amino, or C1-C6 alkyl;

R4 represents halogen, nitro, cyano, carboxy, substituted or unsubstituted C1-C8 alkyl, substituted or unsubstituted C2-C6

10 alkenyl, substituted or unsubstituted C1-C6 alkoxy, substituted or unsubstituted C3-C7 cycloalkyl, mono- or di(C1-C6 alkyl)amino, or substituted or unsubstituted carbamoyl;

when at least one R4 represents substituted C1-C8 alkyl, substituted C2-C6 alkenyl, substituted C1-C6 alkoxy, substituted

15 C3-C7 cycloalkyl, or substituted carbamoyl, the substituent is halogen, carboxy, C1-C6 alkoxy, hydroxy, C1-C6 alkyl that may be substituted with hydroxy, monocyclic 5- to 10-membered unsaturated hydrocarbon, carbamoyl that may be substituted with C1-C6 alkyl or monocyclic 5- to 10-membered unsaturated

20 hydrocarbon, (C2-C7 acyl)oxy, amino that may be substituted with C1-C6 alkyl or C2-C7 acyl, C3-C7 cycloalkyl that may be substituted with hydroxy, or (C1-C6 alkoxy)(C1-C6 alkyl); when two or more of the substituents are present, the substituents may be identical or different;

25 R5 and R6 may be identical or different, and each represent hydrogen or C1-C6 alkyl; R5 and R6 may otherwise form oxo or thioxo together;

when R5 and R6 are identical or different, and each represent hydrogen or C1-C6 alkyl, at least one of R5 and R6 is hydrogen;

30 l represents an integer of 0 to 2;

m represents an integer of 0 to 2;

n represents an integer of 0 to 5;


when l is 2, two R2s may be identical or different;

when m is 2, two R3s may be identical or different;

35 when n is 2 to 5, two to five R4s may be identical or different.

[Claim 12]

Use of a biphenyl compound represented by Formula (I)
or a salt thereof for the production of an antitumor effect
5 potentiator for one or more other antitumor agents:



wherein

ring A represents a monocyclic, bridged cyclic, or spirocyclic 4-
to 14-membered nitrogen-containing saturated heterocyclic group
10 having 1 to 3 nitrogen atoms, 0 to 1 sulfur atoms, and 0 to 2
oxygen atoms as heteroatoms;

ring B represents monocyclic or bicyclic 5- to 14-membered
unsaturated hydrocarbon or a monocyclic or bicyclic 5- to 14-
membered unsaturated heterocyclic group that may be substituted
15 with oxo, that has 0 to 4 nitrogen atoms, 0 to 2 sulfur atoms,
and 0 to 3 oxygen atoms as heteroatoms, and that has at least one
of nitrogen, sulfur, and oxygen;

R1 represents nitro or cyano;

R2 represents halogen;

20 R3 represents amino, mono- or di(C1-C6 alkyl)amino, (C3-C7
cycloalkyl)amino, or C1-C6 alkyl;


R4 represents halogen, nitro, cyano, carboxy, substituted or
unsubstituted C1-C8 alkyl, substituted or unsubstituted C2-C6
alkenyl, substituted or unsubstituted C1-C6 alkoxy, substituted
25 or unsubstituted C3-C7 cycloalkyl, mono- or di(C1-C6 alkyl)amino,
or substituted or unsubstituted carbamoyl;

when at least one R4 represents substituted C1-C8 alkyl,
substituted C2-C6 alkenyl, substituted C1-C6 alkoxy, substituted
C3-C7 cycloalkyl, or substituted carbamoyl, the substituent is

- halogen, carboxy, C1-C6 alkoxy, hydroxy, C1-C6 alkyl that may be substituted with hydroxy, monocyclic 5- to 10-membered unsaturated hydrocarbon, carbamoyl that may be substituted with C1-C6 alkyl or monocyclic 5- to 10-membered unsaturated
- 5 hydrocarbon, (C2-C7 acyl)oxy, amino that may be substituted with C1-C6 alkyl or C2-C7 acyl, C3-C7 cycloalkyl that may be substituted with hydroxy, or (C1-C6 alkoxy) (C1-C6 alkyl); when two or more of the substituents are present, the substituents may be identical or different;
- 10 R5 and R6 may be identical or different, and each represent hydrogen or C1-C6 alkyl; R5 and R6 may otherwise form oxo or thioxo together;
- when R5 and R6 are identical or different, and each represent hydrogen or C1-C6 alkyl, at least one of R5 and R6 is hydrogen;
- 15 l represents an integer of 0 to 2;
- m represents an integer of 0 to 2;
- n represents an integer of 0 to 5;
- when l is 2, two R2s may be identical or different;
- when m is 2, two R3s may be identical or different;
- 20 when n is 2 to 5, two to five R4s may be identical or different.

[Claim 13]

- A method for preventing and/or treating a tumor, comprising administering a combination of a biphenyl compound
- 25 represented by Formula (I) or a salt thereof and one or more other antitumor agents in an amount effective for prevention and/or treatment to a patient:



wherein

ring A represents a monocyclic, bridged cyclic, or spirocyclic 4- to 14-membered nitrogen-containing saturated heterocyclic group having 1 to 3 nitrogen atoms, 0 to 1 sulfur atoms, and 0 to 2 oxygen atoms as heteroatoms;

5 ring B represents monocyclic or bicyclic 5- to 14-membered unsaturated hydrocarbon or a monocyclic or bicyclic 5- to 14-membered unsaturated heterocyclic group that may be substituted with oxo, that has 0 to 4 nitrogen atoms, 0 to 2 sulfur atoms, and 0 to 3 oxygen atoms as heteroatoms, and that has at least one
10 of nitrogen, sulfur, and oxygen;

R1 represents nitro or cyano;

R2 represents halogen;

R3 represents amino, mono- or di(C1-C6 alkyl)amino, (C3-C7 cycloalkyl)amino, or C1-C6 alkyl;

15 R4 represents halogen, nitro, cyano, carboxy, substituted or unsubstituted C1-C8 alkyl, substituted or unsubstituted C2-C6 alkenyl, substituted or unsubstituted C1-C6 alkoxy, substituted or unsubstituted C3-C7 cycloalkyl, mono- or di(C1-C6 alkyl)amino, or substituted or unsubstituted carbamoyl;

20 when at least one R4 represents substituted C1-C8 alkyl, substituted C2-C6 alkenyl, substituted C1-C6 alkoxy, substituted C3-C7 cycloalkyl, or substituted carbamoyl, the substituent is halogen, carboxy, C1-C6 alkoxy, hydroxy, C1-C6 alkyl that may be substituted with hydroxy, monocyclic 5- to 10-membered

25 unsaturated hydrocarbon, carbamoyl that may be substituted with C1-C6 alkyl or monocyclic 5- to 10-membered unsaturated hydrocarbon, (C2-C7 acyl)oxy, amino that may be substituted with C1-C6 alkyl or C2-C7 acyl, C3-C7 cycloalkyl that may be substituted with hydroxy, or (C1-C6 alkoxy)(C1-C6 alkyl); when

30 two or more of the substituents are present, the substituents may be identical or different;

R5 and R6 may be identical or different, and each represent hydrogen or C1-C6 alkyl; R5 and R6 may otherwise form oxo or thioxo together;

35 when R5 and R6 are identical or different, and each represent

hydrogen or C1-C6 alkyl, at least one of R5 and R6 is hydrogen;

l represents an integer of 0 to 2;

m represents an integer of 0 to 2;

n represents an integer of 0 to 5;


5 when l is 2, two R2s may be identical or different;

when m is 2, two R3s may be identical or different;

when n is 2 to 5, two to five R4s may be identical or different.

[Claim 14]

10 A method for preventing and/or treating a tumor,
comprising administering a biphenyl compound represented by
Formula (I) or a salt thereof in an amount effective for
prevention and/or treatment to a cancer patient administered one
or more other antitumor agents:



15 wherein

ring A represents a monocyclic, bridged cyclic, or spirocyclic 4-
to 14-membered nitrogen-containing saturated heterocyclic group
having 1 to 3 nitrogen atoms, 0 to 1 sulfur atoms, and 0 to 2
20 oxygen atoms as heteroatoms;

ring B represents monocyclic or bicyclic 5- to 14-membered
unsaturated hydrocarbon or a monocyclic or bicyclic 5- to 14-
membered unsaturated heterocyclic group that may be substituted
with oxo, that has 0 to 4 nitrogen atoms, 0 to 2 sulfur atoms,
25 and 0 to 3 oxygen atoms as heteroatoms, and that has at least one
of nitrogen, sulfur, and oxygen;

R1 represents nitro or cyano;

R2 represents halogen;

R3 represents amino, mono- or di(C1-C6 alkyl)amino, (C3-C7

cycloalkyl)amino, or C1-C6 alkyl;

R4 represents halogen, nitro, cyano, carboxy, substituted or unsubstituted C1-C8 alkyl, substituted or unsubstituted C2-C6 alkenyl, substituted or unsubstituted C1-C6 alkoxy, substituted or unsubstituted C3-C7 cycloalkyl, mono- or di(C1-C6 alkyl)amino, or substituted or unsubstituted carbamoyl;

when at least one R4 represents substituted C1-C8 alkyl, substituted C2-C6 alkenyl, substituted C1-C6 alkoxy, substituted C3-C7 cycloalkyl, or substituted carbamoyl, the substituent is

halogen, carboxy, C1-C6 alkoxy, hydroxy, C1-C6 alkyl that may be substituted with hydroxy, monocyclic 5- to 10-membered unsaturated hydrocarbon, carbamoyl that may be substituted with C1-C6 alkyl or monocyclic 5- to 10-membered unsaturated hydrocarbon, (C2-C7 acyl)oxy, amino that may be substituted with C1-C6 alkyl or C2-C7 acyl, C3-C7 cycloalkyl that may be substituted with hydroxy, or (C1-C6 alkoxy)(C1-C6 alkyl); when two or more of the substituents are present, the substituents may be identical or different;

R5 and R6 may be identical or different, and each represent hydrogen or C1-C6 alkyl; R5 and R6 may otherwise form oxo or thioxo together;

when R5 and R6 are identical or different, and each represent hydrogen or C1-C6 alkyl, at least one of R5 and R6 is hydrogen; l represents an integer of 0 to 2;

m represents an integer of 0 to 2;

n represents an integer of 0 to 5;

when l is 2, two R2s may be identical or different;


when m is 2, two R3s may be identical or different;

when n is 2 to 5, two to five R4s may be identical or different.

[Claim 15]

A method for potentiating an antitumor effect, comprising administering a biphenyl compound represented by Formula (I) or a salt thereof in an amount effective for prevention and/or treatment to a cancer patient administered one

or more other antitumor agents:



wherein

ring A represents a monocyclic, bridged cyclic, or spirocyclic 4-
 5 to 14-membered nitrogen-containing saturated heterocyclic group
 having 1 to 3 nitrogen atoms, 0 to 1 sulfur atoms, and 0 to 2
 oxygen atoms as heteroatoms;

ring B represents monocyclic or bicyclic 5- to 14-membered
 unsaturated hydrocarbon or a monocyclic or bicyclic 5- to 14-
 10 membered unsaturated heterocyclic group that may be substituted
 with oxo, that has 0 to 4 nitrogen atoms, 0 to 2 sulfur atoms,
 and 0 to 3 oxygen atoms as heteroatoms, and that has at least one
 of nitrogen, sulfur, and oxygen;

R1 represents nitro or cyano;

15 R2 represents halogen;

R3 represents amino, mono- or di(C1-C6 alkyl)amino, (C3-C7
 cycloalkyl)amino, or C1-C6 alkyl;


R4 represents halogen, nitro, cyano, carboxy, substituted or
 unsubstituted C1-C8 alkyl, substituted or unsubstituted C2-C6
 20 alkenyl, substituted or unsubstituted C1-C6 alkoxy, substituted
 or unsubstituted C3-C7 cycloalkyl, mono- or di(C1-C6 alkyl)amino,
 or substituted or unsubstituted carbamoyl;

when at least one R4 represents substituted C1-C8 alkyl,
 substituted C2-C6 alkenyl, substituted C1-C6 alkoxy, substituted
 25 C3-C7 cycloalkyl, or substituted carbamoyl, the substituent is
 halogen, carboxy, C1-C6 alkoxy, hydroxy, C1-C6 alkyl that may be
 substituted with hydroxy, monocyclic 5- to 10-membered
 unsaturated hydrocarbon, carbamoyl that may be substituted with
 C1-C6 alkyl or monocyclic 5- to 10-membered unsaturated

hydrocarbon, (C2-C7 acyl)oxy, amino that may be substituted with C1-C6 alkyl or C2-C7 acyl, C3-C7 cycloalkyl that may be substituted with hydroxy, or (C1-C6 alkoxy) (C1-C6 alkyl); when two or more of the substituents are present, the substituents may
 5 be identical or different;
 R5 and R6 may be identical or different, and each represent hydrogen or C1-C6 alkyl; R5 and R6 may otherwise form oxo or thioxo together;
 when R5 and R6 are identical or different, and each represent
 10 hydrogen or C1-C6 alkyl, at least one of R5 and R6 is hydrogen;
 l represents an integer of 0 to 2;
 m represents an integer of 0 to 2;
 n represents an integer of 0 to 5;
 when l is 2, two R2s may be identical or different;
 15 when m is 2, two R3s may be identical or different;
 when n is 2 to 5, two to five R4s may be identical or different.

[Claim 16]

A product as a combined preparation to be used
 20 simultaneously, sequentially, or at an interval upon prevention and/or treatment of a tumor, the product comprising a biphenyl compound or a salt thereof and one or more other antitumor agents, wherein the biphenyl compound is represented by Formula (I):



25

wherein

ring A represents a monocyclic, bridged cyclic, or spirocyclic 4- to 14-membered nitrogen-containing saturated heterocyclic group having 1 to 3 nitrogen atoms, 0 to 1 sulfur atoms, and 0 to 2

oxygen atoms as heteroatoms;

ring B represents monocyclic or bicyclic 5- to 14-membered

unsaturated hydrocarbon or a monocyclic or bicyclic 5- to 14-

membered unsaturated heterocyclic group that may be substituted

5 with oxo, that has 0 to 4 nitrogen atoms, 0 to 2 sulfur atoms,

and 0 to 3 oxygen atoms as heteroatoms, and that has at least one

of nitrogen, sulfur, and oxygen;

R1 represents nitro or cyano;

R2 represents halogen;

10 R3 represents amino, mono- or di(C1-C6 alkyl)amino, (C3-C7
cycloalkyl)amino, or C1-C6 alkyl;

R4 represents halogen, nitro, cyano, carboxy, substituted or
unsubstituted C1-C8 alkyl, substituted or unsubstituted C2-C6

alkenyl, substituted or unsubstituted C1-C6 alkoxy, substituted

15 or unsubstituted C3-C7 cycloalkyl, mono- or di(C1-C6 alkyl)amino,
or substituted or unsubstituted carbamoyl;

when at least one R4 represents substituted C1-C8 alkyl,

substituted C2-C6 alkenyl, substituted C1-C6 alkoxy, substituted

C3-C7 cycloalkyl, or substituted carbamoyl, the substituent is

20 halogen, carboxy, C1-C6 alkoxy, hydroxy, C1-C6 alkyl that may be
substituted with hydroxy, monocyclic 5- to 10-membered

unsaturated hydrocarbon, carbamoyl that may be substituted with

C1-C6 alkyl or monocyclic 5- to 10-membered unsaturated

hydrocarbon, (C2-C7 acyl)oxy, amino that may be substituted with

25 C1-C6 alkyl or C2-C7 acyl, C3-C7 cycloalkyl that may be

substituted with hydroxy, or (C1-C6 alkoxy)(C1-C6 alkyl); when

two or more of the substituents are present, the substituents may
be identical or different;

R5 and R6 may be identical or different, and each represent

30 hydrogen or C1-C6 alkyl; R5 and R6 may otherwise form oxo or
thioxo together;

when R5 and R6 are identical or different, and each represent

hydrogen or C1-C6 alkyl, at least one of R5 and R6 is hydrogen;

l represents an integer of 0 to 2;


35 m represents an integer of 0 to 2;

n represents an integer of 0 to 5;
 when l is 2, two R2s may be identical or different;
 when m is 2, two R3s may be identical or different;
 when n is 2 to 5, two to five R4s may be identical or different.

5

[Claim 17]

A combination of a biphenyl compound represented by Formula (I) or a salt thereof and one or more other antitumor agents:



10

wherein

ring A represents a monocyclic, bridged cyclic, or spirocyclic 4- to 14-membered nitrogen-containing saturated heterocyclic group having 1 to 3 nitrogen atoms, 0 to 1 sulfur atoms, and 0 to 2

15 oxygen atoms as heteroatoms;

ring B represents monocyclic or bicyclic 5- to 14-membered unsaturated hydrocarbon or a monocyclic or bicyclic 5- to 14-membered unsaturated heterocyclic group that may be substituted with oxo, that has 0 to 4 nitrogen atoms, 0 to 2 sulfur atoms,

20 and 0 to 3 oxygen atoms as heteroatoms, and that has at least one of nitrogen, sulfur, and oxygen;

R1 represents nitro or cyano;

R2 represents halogen;

R3 represents amino, mono- or di(C1-C6 alkyl)amino, (C3-C7 cycloalkyl)amino, or C1-C6 alkyl;

25

R4 represents halogen, nitro, cyano, carboxy, substituted or unsubstituted C1-C8 alkyl, substituted or unsubstituted C2-C6 alkenyl, substituted or unsubstituted C1-C6 alkoxy, substituted or unsubstituted C3-C7 cycloalkyl, mono- or di(C1-C6 alkyl)amino,

or substituted or unsubstituted carbamoyl;

when at least one R4 represents substituted C1-C8 alkyl,
substituted C2-C6 alkenyl, substituted C1-C6 alkoxy, substituted

C3-C7 cycloalkyl, or substituted carbamoyl, the substituent is

5 halogen, carboxy, C1-C6 alkoxy, hydroxy, C1-C6 alkyl that may be
substituted with hydroxy, monocyclic 5- to 10-membered

unsaturated hydrocarbon, carbamoyl that may be substituted with

C1-C6 alkyl or monocyclic 5- to 10-membered unsaturated

hydrocarbon, (C2-C7 acyl)oxy, amino that may be substituted with

10 C1-C6 alkyl or C2-C7 acyl, C3-C7 cycloalkyl that may be

substituted with hydroxy, or (C1-C6 alkoxy) (C1-C6 alkyl); when

two or more of the substituents are present, the substituents may
be identical or different;

R5 and R6 may be identical or different, and each represent

15 hydrogen or C1-C6 alkyl; R5 and R6 may otherwise form oxo or
thioxo together;

when R5 and R6 are identical or different, and each represent

hydrogen or C1-C6 alkyl, at least one of R5 and R6 is hydrogen;

l represents an integer of 0 to 2;

20 m represents an integer of 0 to 2;

n represents an integer of 0 to 5;

when l is 2, two R2s may be identical or different;

when m is 2, two R3s may be identical or different;

when n is 2 to 5, two to five R4s may be identical or different.

25