

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
29 October 2009 (29.10.2009)

(10) International Publication Number  
WO 2009/130453 A1

(51) International Patent Classification:  
*A61K 31/519* (2006.01)    *A61P 35/00* (2006.01)  
*C07D 475/00* (2006.01)

(74) Agent: WALLS, Alan, James; P.O. Box 223, Tadworth, Surrey KT20 5YF (GB).

(21) International Application Number:  
PCT/GB2009/001019

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(22) International Filing Date:  
23 April 2009 (23.04.2009)

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
0807451.0    24 April 2008 (24.04.2008)    GB

(71) Applicant (for all designated States except US): CHROMA THERAPEUTICS LTD. [GB/GB]; 93 Milton Park, Abingdon, Oxfordshire OX14 4RY (GB).


(72) Inventors; and

(75) Inventors/Applicants (for US only): MOFFAT, David, Festus, Charles [GB/GB]; Chroma Therapeutics Ltd., 93, Milton Park, Abingdon, Oxfordshire OX14 4RY (GB). PATEL, Sanjay, Ratilal [GB/GB]; Chroma Therapeutics Ltd., 93, Milton Park, Abingdon, Oxfordshire OX14 4RY (GB). BAKER, Kenneth, William, John [GB/GB]; Chroma Therapeutics Ltd., 93, Milton Park, Abingdon, Oxfordshire OX14 4RY (GB). NORTH, Carl, Leslie [GB/GB]; Chroma Therapeutics Ltd., 93, Milton Park, Abingdon, Oxfordshire OX14 4RY (GB).

Published:

— with international search report (Art. 21(3))

(54) Title: INHIBITORS OF PLK



WO 2009/130453 A1

(57) Abstract: Compounds of formula (I) are PLK inhibitors, useful for the treatment of cell proliferative diseases: wherein R<sub>1</sub> is hydrogen, or an optionally substituted (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>6</sub>)alkenyl, (C<sub>2</sub>-C<sub>6</sub>)alkynyl or (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl group; R<sub>2</sub> is hydrogen, or an optionally substituted (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>6</sub>)alkenyl, (C<sub>2</sub>-C<sub>6</sub>)alkynyl or (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl group; R<sub>3</sub> is hydrogen, -CN, hydroxyl, halogen, optionally substituted (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>6</sub>)alkenyl, (C<sub>2</sub>-C<sub>6</sub>)alkynyl or (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl, -NR<sub>5</sub>R<sub>6</sub> or C<sub>1</sub>-C<sub>4</sub> alkoxy, wherein R<sub>5</sub> and R<sub>6</sub> are independently hydrogen or optionally substituted (C<sub>1</sub>-C<sub>6</sub>)alkyl; ring A is an optionally substituted mono- or bi-cyclic carbocyclic or heterocyclic ring or a ring system having up to 12 ring atoms; T is a radical of formula R-L<sup>1</sup>-Y<sup>1</sup>- wherein L<sup>1</sup> and Y<sup>1</sup> are as defined in the claims and R is a carbon-linked, alpha disubstituted amino acid or amino acid ester residue.

### Inhibitors of PLK

This invention relates to a series of amino acid esters, to compositions containing them, to processes for their preparation and to their use in medicine as Polo-like kinase 'PLK' inhibitors. Polo-like kinases (PLKs) are key enzymes that control mitotic entry of proliferating cells and regulate many aspects of mitosis necessary for successful cytokinesis. Of the four known human PLKs, PLK1 is the best characterized and is overexpressed in many tumour types with aberrant elevation frequently constituting a prognostic indicator of poor disease outcome. The compounds may be of use in the treatment of cell proliferative diseases such as cancer. The present invention encompasses compounds that are dihydropteridinine derivatives.

### Background to invention

The PLKs, a family of Ser/Thr protein kinases named after their functional and sequence similarity with the archetypal *polo* kinase from *Drosophila melanogaster*, play a variety of roles in mitosis (*Nat. Rev. Mol. Cell Biol.*, **2001**, 2, 21-32.). In yeasts (*Saccharomyces cerevisiae* and *S. pombe*) single PLKs exist, whereas four distinct PLKs have been identified to date in mammals. Human PLK1 (*Cell Growth Differ.*, **1994**, 5, 249-257), PLK2 (serum-inducible kinase, SNK, *Mol. Cell. Biol.*, **1992**, 12, 4164-4169), PLK3 (proliferation-related kinase, PRK *J. Biol. Chem.* **1997**, 272, 28646-28651) and PLK4 (*Oncol. Rep.*, **1997**, 4, 505-510) are structurally homologous and contain two conserved domains, the N-terminal catalytic kinase domain, as well as a C-terminal region composed of the so-called polo boxes. Whereas PLK1, PLK2, and PLK3 are expressed in all tissues, PLK4 appears to possess unique physiological roles and the distribution of PLK4 mRNA in adults is restricted to certain tissues such as testes and thymus. PLK1 is the best characterized member of the PLK family and it appears to fulfil most of the known functions of the single PLKs present in invertebrates (*Nat. Rev. Mol. Cell Biol.*, **2004**, 5, 429-441). PLK1 protein levels fluctuate in a cell-cycle-dependent manner and its kinase activity peaks at the transition between the second gap phase and the mitosis phases (G2/M) of the eukaryotic cell division cycle. Upon exit from mitosis PLK1 levels drop as a result of ubiquitin-dependent proteolysis. PLK1 has been reported to be involved in the initiation of mitosis through activation of the cyclin-dependent kinase CDK1/cyclin B complex, *i.e.* the master switch for mitotic entry (mitosis-promoting factor, MPF *Nature*, **1990**, 344, 503-508).


This occurs when PLK1 phosphorylates, and thus activates, the dual specificity phosphatase CDC25C, which in turn relieves premitotic MYT1- and WEE1- mediated suppression of CDK1/cyclin B activity through dephosphorylation at the CDK1 pThr14 and pTyr15 sites (*Cell*, 1991, 67, 197-211). Upon entry into mitosis, phosphorylation of CDC25C by PLK1 and PLK3 leads to its translocation into the nucleus. Apart from controlling entry into mitosis through CDK1 activation, PLK1 has additional roles in regulating progression through mitosis. It is involved in bipolar spindle formation, including centrosome maturation and regulation of the microtubule organizing centre, in the subsequent steps of mitosis involving sister chromatid separation, and finally in cytokinesis (*Dev. Cell*, 2003, 5, 127-138).

### **Brief Summary of the Invention**

Compounds of the invention are related to compounds disclosed in WO2004076454. They are inhibitors of PLK1 and the isoforms thereof. The compounds are thus of use in medicine, for example in the treatment of a variety of proliferative disease states, including cancers. The compounds are characterised by the presence in the molecule of an  $\alpha,\alpha$ -disubstituted glycine acid motif or an  $\alpha,\alpha$ -disubstituted glycine ester motif which is hydrolysable by an intracellular carboxylesterase. Compounds of the invention having the lipophilic  $\alpha,\alpha$ -disubstituted glycine ester motif cross the cell membrane, and are hydrolysed to the acid by the intracellular carboxylesterases. The polar hydrolysis product accumulates in the cell since it does not readily cross the cell membrane. Hence the PLK1 activity of the compound is prolonged and enhanced within the cell.

### **Detailed Description of the Invention**

According to the invention there is provided a compound of formula (I), or a salt thereof:



wherein

**R<sub>1</sub>** is hydrogen, or an optionally substituted (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>6</sub>)alkenyl, (C<sub>2</sub>-C<sub>6</sub>)alkynyl or (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl group;

**R<sub>2</sub>** is hydrogen, or an optionally substituted (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>6</sub>)alkenyl, (C<sub>2</sub>-C<sub>6</sub>)alkynyl or (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl group;

**R<sub>3</sub>** is hydrogen, -CN, hydroxyl, halogen, optionally substituted (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>6</sub>)alkenyl, (C<sub>2</sub>-C<sub>6</sub>)alkynyl or (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl, -NR<sub>5</sub>R<sub>6</sub> or (C<sub>1</sub>-C<sub>4</sub>)alkoxy, wherein R<sub>5</sub> and R<sub>6</sub> are independently hydrogen or optionally substituted (C<sub>1</sub>-C<sub>6</sub>)alkyl;

ring **A** is an optionally substituted mono- or bi-cyclic carbocyclic or heterocyclic ring or a ring system having up to 12 ring atoms;

**T** is a radical of formula R-L<sup>1</sup>-Y<sup>1</sup>- wherein

**Y<sup>1</sup>** is a bond, -O-, -S-, -NR<sub>6</sub>-, -(C=O)-, -S(O<sub>2</sub>)-, -(C=O)NR<sub>6</sub>-, -NR<sub>6</sub>(C=O)-, -S(O<sub>2</sub>)NR<sub>6</sub>-, -NR<sub>6</sub>S(O<sub>2</sub>)-, or -NR<sub>6</sub>(C=O)NR<sub>9</sub>-, wherein R<sub>6</sub> and R<sub>9</sub> are independently hydrogen or optionally substituted (C<sub>1</sub>-C<sub>6</sub>)alkyl;


**L<sup>1</sup>** is a divalent radical of formula -(Alk<sup>1</sup>)<sub>m</sub>(Q)<sub>n</sub>(Alk<sup>2</sup>)<sub>p</sub>- wherein

**m**, **n** and **p** are independently 0 or 1,

**Q** is (i) an optionally substituted divalent mono- or bicyclic carbocyclic or heterocyclic radical having 5 - 13 ring members, or (ii), in the case where p is 0, a divalent radical of formula -Q<sup>1</sup>-X<sup>2</sup>- wherein X<sup>2</sup> is -O-, -S- or NR<sup>A</sup>- wherein R<sup>A</sup> is hydrogen or optionally substituted C<sub>1</sub>-C<sub>3</sub> alkyl, and Q<sup>1</sup> is an optionally substituted divalent mono- or bicyclic carbocyclic or heterocyclic radical having 5 - 13 ring members,

**Alk<sup>1</sup>** and **Alk<sup>2</sup>** independently represent optionally substituted divalent C<sub>3</sub>-C<sub>7</sub> cycloalkyl radicals, or optionally substituted straight or branched, C<sub>1</sub>-C<sub>6</sub> alkylene, C<sub>2</sub>-C<sub>6</sub> alkenylene, or C<sub>2</sub>-C<sub>6</sub> alkynylene radicals which may optionally contain or terminate in an ether (-O-), thioether (-S-) or amino (-NR<sup>A</sup>-) link wherein R<sup>A</sup> is hydrogen or optionally substituted C<sub>1</sub>-C<sub>3</sub> alkyl;

**R** is a radical of formula (X)



wherein

**R**<sub>7</sub> is a carboxylic acid group (-COOH), or an ester group which is hydrolysable by one or more intracellular carboxylesterase enzymes to a carboxylic acid group;

**R**<sup>1</sup><sub>7</sub> is the side chain of a natural or non-natural alpha-amino acid, in which any functional groups are protected, but **R**<sup>1</sup><sub>7</sub> is not hydrogen;

**R**<sub>8</sub> is hydrogen; or optionally substituted C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>3</sub>-C<sub>7</sub> cycloalkyl, aryl or heteroaryl or -(C=O)R<sub>6</sub>, -(C=O)OR<sub>6</sub>, or -(C=O)NR<sub>6</sub> wherein R<sub>6</sub> is hydrogen or optionally substituted (C<sub>1</sub>-C<sub>6</sub>)alkyl.

In the compounds of the invention, when R<sub>1</sub> is other than hydrogen, the carbon atom to which the R<sub>1</sub> substituent is attached is asymmetric. Preferably the stereo chemistry at that asymmetric center is R.

In another broad aspect the invention provides the use of a compound of formula (I) as defined above, or an N-oxide, salt, hydrate or solvate thereof in the preparation of a composition for inhibiting the activity of PLK1.

The compounds with which the invention is concerned may be used for the inhibition of PLK1 activity *ex vivo* or *in vivo*.

In one aspect of the invention, the compounds of the invention may be used in the preparation of a composition for treatment of cell proliferative diseases such as cancer.

In another aspect, the invention provides a method for the treatment of the foregoing disease types, which comprises administering to a subject suffering such disease an effective amount of a compound of formula (I) as defined above.

### ***Terminology***

As used herein, the term " $(C_a-C_b)alkyl$ " wherein a and b are integers, refers to a straight or branched chain alkyl radical having from a to b carbon atoms. Thus when a is 1 and b is 6, for example, the term includes methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl, t-butyl, n-pentyl and n-hexyl.

As used herein, the term "divalent  $(C_a-C_b)alkylene$  radical", wherein a and b are integers, refers to a saturated hydrocarbon chain having from a to b carbon atoms and two unsatisfied valences.

As used herein, the term " $(C_a-C_b)alkenyl$ " wherein a and b are integers, refers to a straight or branched chain alkenyl moiety with a to b carbon atoms; having at least one double bond of either E or Z stereochemistry where applicable. The term includes, for example, vinyl, allyl, 1- and 2-butenyl and 2-methyl-2-propenyl.

As used herein, the term "divalent  $(C_a-C_b)alkenylene$  radical" means a hydrocarbon chain having from a to b carbon atoms, at least one double bond, and two unsatisfied valences.

As used herein the term " $C_a-C_b$  alkynyl", wherein a and b are integers refers to straight chain or branched chain hydrocarbon groups having from two to six carbon atoms and having in addition one triple bond. This term would include, for example, ethynyl, 1-propynyl, 1- and 2-butynyl, 2-methyl-2-propynyl, 2-pentynyl, 3-pentynyl, 4-pentynyl, 2-hexynyl, 3-hexynyl, 4-hexynyl and 5-hexynyl.

As used herein, the term "divalent  $(C_a-C_b)alkynylene$  radical", wherein a and b are integers refers to a divalent hydrocarbon chain having from two to six carbon atoms, and at least one triple bond.

As used herein, the term "carbocyclic" refers to a mono-, bi- or tricyclic radical having up to 16 ring atoms, all of which are carbon, and includes aryl and cycloalkyl.


As used herein, the term "cycloalkyl" refers to a monocyclic saturated carbocyclic radical having from 3-8 carbon atoms and includes, for example, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl and cyclooctyl.

As used herein, the unqualified term "aryl" refers to a mono-, bi- or tri-cyclic carbocyclic aromatic radical, and includes radicals having two monocyclic carbocyclic aromatic rings which are directly linked by a covalent bond. Illustrative of such radicals are phenyl, biphenyl and naphthyl.

As used herein, the unqualified term "heteroaryl" refers to a mono-, bi- or tri-cyclic aromatic radical containing one or more heteroatoms selected from S, N and O, and includes radicals having two such monocyclic rings, or one such monocyclic ring and one monocyclic aryl ring, which are directly linked by a covalent bond. Illustrative of such radicals are thienyl, benzthienyl, furyl, benzfuryl, pyrrolyl, imidazolyl, benzimidazolyl, thiazolyl, benzthiazolyl, isothiazolyl, benzisothiazolyl, pyrazolyl, oxazolyl, benzoxazolyl, isoxazolyl, benzisoxazolyl, isothiazolyl, triazolyl, benztriazolyl, thiadiazolyl, oxadiazolyl, pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl, triazinyl, indolyl and indazolyl.

As used herein, the unqualified term "heterocyclyl" or "heterocyclic" includes "heteroaryl" as defined above, and in its non-aromatic meaning relates to a mono-, bi- or tri-cyclic non-aromatic radical containing one or more heteroatoms selected from S, N and O, and to groups consisting of a monocyclic non-aromatic radical containing one or more such heteroatoms which is covalently linked to another such radical or to a monocyclic carbocyclic radical. Illustrative of such radicals are pyrrolyl, furanyl, thienyl, piperidinyl, imidazolyl, oxazolyl, isoxazolyl, thiazolyl, thiadiazolyl, pyrazolyl, pyridinyl, pyrrolidinyl, pyrimidinyl, morpholinyl, piperazinyl, indolyl, morpholinyl, benzfuranyl, pyranyl, isoxazolyl, benzimidazolyl, methylenedioxyphenyl, ethylenedioxyphenyl, maleimido and succinimido groups.

A "divalent phenylene, pyridinylene, pyrimidinylene, or pyrazinylene radical" is a benzene, pyridine, pyrimidine or pyrazine ring, with two unsatisfied valencies, and includes 1,3-phenylene, 1,4-phenylene, and the following:



Unless otherwise specified in the context in which it occurs, the term "substituted", as applied to any moiety herein, means substituted with up to four compatible substituents, each of which independently may be, for example, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, hydroxy, hydroxy(C<sub>1</sub>-C<sub>6</sub>)alkyl, mercapto, mercapto(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkylthio, phenyl, halo (including fluoro, bromo and chloro), trifluoromethyl, trifluoromethoxy, nitro, nitrile (-CN), oxo, -COOH, -COOR<sup>A</sup>, -COR<sup>A</sup>, -SO<sub>2</sub>R<sup>A</sup>, -CONH<sub>2</sub>, -SO<sub>2</sub>NH<sub>2</sub>, -CONHR<sup>A</sup>, -SO<sub>2</sub>NHR<sup>A</sup>, -CONR<sup>A</sup>R<sup>B</sup>, -SO<sub>2</sub>NR<sup>A</sup>R<sup>B</sup>, -NH<sub>2</sub>, -NHR<sup>A</sup>, -NR<sup>A</sup>R<sup>B</sup>, -OCONH<sub>2</sub>, -OCONHR<sup>A</sup>, -OCONR<sup>A</sup>R<sup>B</sup>, -NHCOR<sup>A</sup>, -NHCOOR<sup>A</sup>, -NR<sup>B</sup>COOR<sup>A</sup>, -NHSO<sub>2</sub>OR<sup>A</sup>, -NR<sup>B</sup>SO<sub>2</sub>OH, -NR<sup>B</sup>SO<sub>2</sub>OR<sup>A</sup>, -NHCONH<sub>2</sub>, -NR<sup>A</sup>CONH<sub>2</sub>, -NHCONHR<sup>B</sup>, -NR<sup>A</sup>CONHR<sup>B</sup>, -NHCONR<sup>A</sup>R<sup>B</sup>, or -NR<sup>A</sup>CONR<sup>A</sup>R<sup>B</sup> wherein R<sup>A</sup> and R<sup>B</sup> are independently a (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>3</sub>-C<sub>6</sub>) cycloalkyl, phenyl or monocyclic heteroaryl having 5 or 6 ring atoms, or R<sup>A</sup> and R<sup>B</sup> when attached to the same nitrogen atom form a cyclic amino group (for example morpholino, piperidinyl, piperazinyl, or tetrahydropyrrolyl). An "optional substituent" may be one of the foregoing substituent groups.

The term "side chain of a natural or non-natural alpha-amino acid" refers to the group R<sup>Y</sup> in a natural or non-natural amino acid of formula NH<sub>2</sub>-CH(R<sup>Y</sup>)-COOH.

Examples of side chains of natural alpha amino acids include those of alanine, arginine, asparagine, aspartic acid, cysteine, cystine, glutamic acid, histidine, 5-hydroxylysine, 4-hydroxyproline, isoleucine, leucine, lysine, methionine, phenylalanine, proline, serine, threonine, tryptophan, tyrosine, valine,  $\alpha$ -amino adipic acid,  $\alpha$ -amino-n-butyric acid, 3,4-

dihydroxyphenylalanine, homoserine,  $\alpha$ -methylserine, ornithine, pipecolic acid, and thyroxine.

Natural alpha-amino acids which contain functional substituents, for example amino, carboxyl, hydroxy, mercapto, guanidyl, imidazolyl, or indolyl groups in their characteristic side chains include arginine, lysine, glutamic acid, aspartic acid, tryptophan, histidine, serine, threonine, tyrosine, and cysteine. When  $R^1$ , in the compounds of the invention is one of those side chains, the functional substituent may optionally be protected.

The term "protected" when used in relation to a functional substituent in a side chain of a natural alpha-amino acid means a derivative of such a substituent which is substantially non-functional. For example, carboxyl groups may be esterified (for example as a  $C_1-C_6$  alkyl ester), amino groups may be converted to amides (for example as a  $NHCOC_1-C_6$  alkyl amide) or carbamates (for example as an  $NHC(=O)OC_1-C_6$  alkyl or  $NHC(=O)OCH_2Ph$  carbamate), hydroxyl groups may be converted to ethers (for example an  $OC_1-C_6$  alkyl or a  $O(C_1-C_6$  alkyl)phenyl ether) or esters (for example a  $OC(=O)C_1-C_6$  alkyl ester) and thiol groups may be converted to thioethers (for example a tert-butyl or benzyl thioether) or thioesters (for example a  $SC(=O)C_1-C_6$  alkyl thioester).

As used herein the term "salt" includes base addition, acid addition and quaternary salts. Compounds of the invention which are acidic can form salts, including pharmaceutically acceptable salts, with bases such as alkali metal hydroxides, e.g. sodium and potassium hydroxides; alkaline earth metal hydroxides e.g. calcium, barium and magnesium hydroxides; with organic bases e.g. N-methyl-D-glucamine, choline tris(hydroxymethyl)amino-methane, L-arginine, L-lysine, N-ethyl piperidine, dibenzylamine and the like. Those compounds (I) which are basic can form salts, including pharmaceutically acceptable salts with inorganic acids, e.g. with hydrohalic acids such as hydrochloric or hydrobromic acids, sulphuric acid, nitric acid or phosphoric acid and the like, and with organic acids e.g. with acetic, tartaric, succinic, fumaric, maleic, malic, salicylic, citric, methanesulphonic, p-toluenesulphonic, benzoic, benzenesulphonic, glutamic, lactic, and mandelic acids and the like.

It is expected that compounds of the invention may be recovered in N-oxide, hydrate or solvate form, and such forms are expected to have the activity of the non-hydrated, non-

solvated or non-N-oxidised forms. The term 'solvate' is used herein to describe a molecular complex comprising the compound of the invention and a stoichiometric amount of one or more pharmaceutically acceptable solvent molecules, for example, ethanol. The term 'hydrate' is employed when said solvent is water.

Compounds of the invention which contain one or more actual or potential chiral centres, because of the presence of asymmetric carbon atoms, can exist as a number of diastereoisomers with R or S stereochemistry at each chiral centre. The invention includes all such diastereoisomers and mixtures thereof.

The term "ester" or "esterified carboxyl group" in connection with substituent R<sub>7</sub> above means a group R<sub>10</sub>O(C=O)- in which R<sub>10</sub> is the group characterising the ester, notionally derived from the alcohol R<sub>10</sub>OH.

*The substituents R<sub>1</sub>-R<sub>3</sub>*

R<sub>1</sub> is hydrogen, (C<sub>1</sub>-C<sub>6</sub>)alkyl, for example methyl, ethyl, n- or iso-propyl, (C<sub>2</sub>-C<sub>6</sub>)alkenyl, for example allyl, (C<sub>2</sub>-C<sub>6</sub>)alkynyl, for example -CH<sub>2</sub>C≡CH or (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl, for example cyclopropyl, cyclopentyl or cyclohexyl. In one subclass of compounds of the invention R<sub>1</sub> is ethyl.

R<sub>2</sub> is hydrogen, (C<sub>1</sub>-C<sub>6</sub>)alkyl, for example methyl, ethyl, n- or iso-propyl, (C<sub>2</sub>-C<sub>6</sub>)alkenyl, for example allyl, (C<sub>2</sub>-C<sub>6</sub>)alkynyl, for example -CH<sub>2</sub>C≡CH or (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl, for example cyclopropyl, cyclopentyl or cyclohexyl, or C<sub>6-14</sub> aryl for example phenyl or naphthyl. In one subclass of compounds of the invention R<sub>2</sub> is cyclopentyl.

R<sub>3</sub> is hydrogen, -CN, hydroxyl, halogen, (C<sub>1</sub>-C<sub>6</sub>)alkyl, for example methyl, ethyl, n- or iso-propyl, (C<sub>2</sub>-C<sub>6</sub>)alkenyl, for example allyl, (C<sub>2</sub>-C<sub>6</sub>)alkynyl, for example -CH<sub>2</sub>C≡CH or (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl, for example cyclopropyl, cyclopentyl or cyclohexyl, -NR<sub>5</sub>R<sub>6</sub> and C<sub>1</sub>-C<sub>4</sub> alkoxy, wherein R<sub>5</sub> and R<sub>6</sub> are independently hydrogen or optionally substituted (C<sub>1</sub>-C<sub>6</sub>)alkyl, for example methyl or ethyl. In one subclass of compounds of the invention R<sub>3</sub> is hydrogen.

*The ring A*

Ring A is a mono- or bi-cyclic carbocyclic or heterocyclic ring or a ring system having up to 12 ring atoms. Examples of such rings are piperidine, piperazine, pyridine, pyrimidine, pyrazoline, triazoline, furan, thiophene, pyrrole, thiazole, isothiazole, oxazole, isoxazole, and thiadiazole rings. Currently preferred rings A are phenyl, pyridinyl and pyrimidinyl.

Ring A may be substituted by any of the optional substituents referred to above, for example chloro, bromo or fluoro, trifluoromethyl, methoxy, and trifluoromethoxy.

*The substituent T*

This substituent contains the  $\alpha,\alpha$ -disubstituted glycine acid or  $\alpha,\alpha$ -disubstituted glycine ester moiety of formula (X) linked through a linker radical to ring A.

The ester compounds of the invention are converted by intracellular esterases to the carboxylic acid. Both the esters and carboxylic acids may have PLK inhibitory activity in their own right. The compounds of the invention therefore include not only the ester, but also the corresponding carboxylic acid hydrolysis products.

*The ester group R<sub>7</sub>*

The ester group R<sub>7</sub> present in substituent T must be one which in the compound of the invention is hydrolysable by one or more intracellular carboxylesterase enzymes to a carboxylic acid group. Intracellular carboxylesterase enzymes capable of hydrolysing the ester group of a compound of the invention to the corresponding acid include the three known human enzyme isotypes hCE-1, hCE-2 and hCE-3. Although these are considered to be the main enzymes other enzymes such as biphenylhydrolase (BPH) may also have a role in hydrolysing the conjugates. In general, if the carboxylesterase hydrolyses the free amino acid ester to the parent acid it will also hydrolyse the ester motif when covalently conjugated to the modulator. Hence, the broken cell assay described herein provides a straightforward, quick and simple first screen for esters which have the required hydrolysis profile. Ester motifs selected in that way may then be re-assayed in the same carboxylesterase assay when conjugated to the rest of the molecule via the chosen conjugation chemistry, to confirm that it is still a carboxylesterase substrate in that background.

Subject to the requirement that they be hydrolysable by intracellular carboxylesterase enzymes, examples of particular ester groups  $R_7$  include those of formula  $-(C=O)OR_{10}$  wherein  $R_{10}$  is  $R_{11}R_{12}R_{13}C-$  wherein

- (i)  $R_{11}$  is hydrogen, fluorine or optionally substituted  $(C_1-C_3)alkyl-(Z^1)_a-[(C_1-C_3)alkyl]_b-$  or  $(C_2-C_3)alkenyl-(Z^1)_a-[(C_1-C_3)alkyl]_b-$  wherein  $a$  and  $b$  are independently 0 or 1 and  $Z^1$  is  $-O-$ ,  $-S-$ , or  $-NR_{14}-$  wherein  $R_{14}$  is hydrogen or  $(C_1-C_3)alkyl$ ; and  $R_{12}$  and  $R_{13}$  are independently hydrogen or  $(C_1-C_3)alkyl-$ ;
- (ii)  $R_{11}$  is hydrogen or optionally substituted  $R_{15}R_{16}N-(C_1-C_3)alkyl-$  wherein  $R_{15}$  is hydrogen or  $(C_1-C_3)alkyl$  and  $R_{16}$  is hydrogen or  $(C_1-C_3)alkyl$ ; or  $R_{15}$  and  $R_{16}$  together with the nitrogen to which they are attached form an optionally substituted monocyclic heterocyclic ring of 5- or 6- ring atoms or bicyclic heterocyclic ring system of 8 to 10 ring atoms, and  $R_{12}$  and  $R_{13}$  are independently hydrogen or  $(C_1-C_3)alkyl$ ; or
- (iii)  $R_{11}$  and  $R_{12}$  taken together with the carbon to which they are attached form an optionally substituted monocyclic carbocyclic ring of from 3 to 7 ring atoms or bicyclic carbocyclic ring system of 8 to 10 ring atoms, and  $R_{13}$  is hydrogen.

In cases (i), (ii) and (iii) above, "alkyl" includes fluoroalkyl.

Within these classes,  $R_{10}$  may be, for example, methyl, trifluoromethyl, ethyl, n- or isopropyl, n-, sec- or tert-butyl, cyclohexyl, allyl, phenyl, benzyl, 2-, 3- or 4-pyridylmethyl, N-methylpiperidin-4-yl, tetrahydrofuran-3-yl, methoxyethyl, indanyl, norbonyl, dimethylaminoethyl, or morpholinoethyl. Currently preferred is where  $R_{10}$  is cyclopentyl.

#### The group $R_7^1$

The group  $R_7^1$  present in substituent T is the side chain of a natural or non-natural alpha-amino acid, in which any functional groups are protected, but  $R_7^1$  is not hydrogen.

For example,  $R_7^1$  may be phenyl, or heteroaryl such as pyridyl, or a group of formula  $-CR_aR_bR_c$  in which:

each of R<sub>a</sub>, R<sub>b</sub> and R<sub>c</sub> is independently hydrogen, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>6</sub>)alkenyl, (C<sub>2</sub>-C<sub>6</sub>)alkynyl, phenyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>3</sub>-C<sub>8</sub>)cycloalkyl; or

R<sub>c</sub> is hydrogen and R<sub>a</sub> and R<sub>b</sub> are independently phenyl or heteroaryl such as pyridyl; or

R<sub>c</sub> is hydrogen, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>6</sub>)alkenyl, (C<sub>2</sub>-C<sub>6</sub>)alkynyl, phenyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, or (C<sub>3</sub>-C<sub>8</sub>)cycloalkyl, and R<sub>a</sub> and R<sub>b</sub> together with the carbon atom to which they are attached form a 3 to 8 membered cycloalkyl or a 5- to 6-membered heterocyclic ring; or

R<sub>a</sub>, R<sub>b</sub> and R<sub>c</sub> together with the carbon atom to which they are attached form a tricyclic ring (for example adamantyl); or

R<sub>a</sub> and R<sub>b</sub> are each independently (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>6</sub>)alkenyl, (C<sub>2</sub>-C<sub>6</sub>)alkynyl, phenyl(C<sub>1</sub>-C<sub>6</sub>)alkyl, or a group as defined for R<sub>c</sub> below other than hydrogen, or R<sub>a</sub> and R<sub>b</sub> together with the carbon atom to which they are attached form a cycloalkyl or heterocyclic ring, and R<sub>c</sub> is hydrogen, -OH, -SH, halogen, -CN, -CO<sub>2</sub>H, (C<sub>1</sub>-C<sub>4</sub>)perfluoroalkyl, -CH<sub>2</sub>OH, -O(C<sub>1</sub>-C<sub>6</sub>)alkyl, -O(C<sub>2</sub>-C<sub>6</sub>)alkenyl, -S(C<sub>1</sub>-C<sub>6</sub>)alkyl, -SO(C<sub>1</sub>-C<sub>6</sub>)alkyl, -SO<sub>2</sub>(C<sub>1</sub>-C<sub>6</sub>)alkyl, -S(C<sub>2</sub>-C<sub>6</sub>)alkenyl, -SO(C<sub>2</sub>-C<sub>6</sub>)alkenyl, -SO<sub>2</sub>(C<sub>2</sub>-C<sub>6</sub>)alkenyl or a group -Q-W wherein Q represents a bond or -O-, -S-, -SO- or -SO<sub>2</sub>- and W represents a phenyl, phenylalkyl, (C<sub>3</sub>-C<sub>8</sub>)cycloalkyl, (C<sub>3</sub>-C<sub>8</sub>)cycloalkylalkyl, (C<sub>4</sub>-C<sub>8</sub>)cycloalkenyl, (C<sub>4</sub>-C<sub>8</sub>)cycloalkenylalkyl, heteroaryl or heteroarylalkyl group, which group W may optionally be substituted by one or more substituents independently selected from, hydroxyl, halogen, -CN, -CONH<sub>2</sub>, -CONH(C<sub>1</sub>-C<sub>6</sub>)alkyl, -CONH(C<sub>1</sub>-C<sub>6</sub>alkyl)<sub>2</sub>, -CHO, -CH<sub>2</sub>OH, (C<sub>1</sub>-C<sub>4</sub>)perfluoroalkyl, -O(C<sub>1</sub>-C<sub>6</sub>)alkyl, -S(C<sub>1</sub>-C<sub>6</sub>)alkyl, -SO(C<sub>1</sub>-C<sub>6</sub>)alkyl, -SO<sub>2</sub>(C<sub>1</sub>-C<sub>6</sub>)alkyl, -NO<sub>2</sub>, -NH<sub>2</sub>, -NH(C<sub>1</sub>-C<sub>6</sub>)alkyl, -N((C<sub>1</sub>-C<sub>6</sub>)alkyl)<sub>2</sub>, -NHCO(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>6</sub>)alkenyl, (C<sub>2</sub>-C<sub>6</sub>)alkynyl, (C<sub>3</sub>-C<sub>8</sub>)cycloalkyl, (C<sub>4</sub>-C<sub>8</sub>)cycloalkenyl, phenyl or benzyl.

In some cases, R<sup>17</sup> is H-Alk<sup>4</sup>-, phenyl, monocyclic heterocycl, C<sub>3</sub>-C<sub>7</sub> cycloalkyl, phenyl(Alk<sup>4</sup>)-, heterocycl(Alk<sup>4</sup>)-, or C<sub>3</sub>-C<sub>7</sub> cycloalkyl(Alk<sup>4</sup>)-, wherein the heterocycl part is monocyclic heterocycl having 3-7 ring atoms, and wherein -Alk<sup>4</sup>- is a straight or

branched, divalent ( $C_1$ - $C_6$ )alkylene, ( $C_2$ - $C_6$ )alkenylene, or ( $C_2$ - $C_6$ )alkynylene radical which may optionally be interrupted by, or terminate in, an ether (-O-), thioether (-S-) or amino (-NR<sup>A</sup>-) link wherein R<sup>A</sup> is hydrogen or optionally substituted ( $C_1$ - $C_3$ )alkyl, and wherein the Alk<sup>4</sup>-, or cyclic part is optionally substituted. For example, R<sup>17</sup> may be  $C_1$ - $C_6$  alkyl substituent, for example methyl, ethyl, n- or iso-propyl, or n-, sec- or tert-butyl.

In a particular case, R<sup>17</sup> is methyl.

#### The group R<sub>8</sub>

R<sub>8</sub> may be, for example, optionally substituted ( $C_1$ - $C_6$ )alkyl, ( $C_3$ - $C_6$ )cycloalkyl, aryl or heteroaryl, for example methyl, ethyl, n- or isopropyl, cyclopropyl, cyclopentyl, cyclohexyl, phenyl, or pyridyl. R<sub>8</sub> may also be, for example hydrogen or -(C=O)R<sub>16</sub>, wherein R<sub>16</sub> is optionally substituted ( $C_1$ - $C_6$ )alkyl such as methyl, ethyl, n- or isopropyl, or n-, iso- or sec-butyl, ( $C_3$ - $C_6$ )cycloalkyl such as cyclopropyl, cyclopentyl, cyclohexyl, phenyl, pyridyl, thiienyl, phenyl( $C_1$ - $C_6$ alkyl)-, thiienyl( $C_1$ - $C_6$ alkyl)- or pyridyl( $C_1$ - $C_6$ alkyl)- such as benzyl, 4-methoxyphenylmethylcarbonyl, thiienylmethyl or pyridylmethyl.

R<sub>8</sub> may also be, for example -(C=O)OR<sub>17</sub>, or -(C=O)NHR<sub>17</sub> wherein R<sub>17</sub> is hydrogen or optionally substituted ( $C_1$ - $C_6$ )alkyl such as methyl, ethyl, or n- or isopropyl.

Currently it is preferred that R<sub>8</sub> be hydrogen.

For compounds of the invention which are to be administered systemically, esters with a slow rate of esterase cleavage are preferred, since they are less susceptible to pre-systemic metabolism. Their ability to reach their target tissue intact is therefore increased, and the ester can be converted inside the cells of the target tissue into the acid product. However, for local administration, where the ester is either directly applied to the target tissue or directed there by, for example, inhalation, it will often be desirable that the ester has a rapid rate of esterase cleavage, to minimise systemic exposure and consequent unwanted side effects. If a carbon atom to which the group R is attached is unsubstituted, ie R is attached to a methylene (-CH<sub>2</sub>)- radical, then the esters tend to be cleaved more rapidly than if that carbon is substituted, or is part of a ring system such as a phenyl or cyclohexyl ring.

The radical -L<sup>1</sup>-Y<sup>1</sup>-

This radical (or bond) arises from the particular chemistry strategy chosen to link the amino acid ester motif R in substituent T to ring A of the inhibitor. Clearly the chemistry strategy for that coupling may vary widely, and thus many combinations of the variables Y<sup>1</sup> and L<sup>1</sup> are possible. However, when the inhibitor is bound to the enzyme at its active site, the amino acid ester motif generally extends in a direction away from the enzyme, and thus minimises or avoids interference with the binding mode of the inhibitor. Hence the precise combination of variable making up the linking chemistry between the amino acid ester motif and the rest of the molecule will often be irrelevant to the primary binding mode of the compound as a whole.

With the foregoing general observations in mind, taking the variables making up the radical -L<sup>1</sup>-Y<sup>1</sup>- in turn:

Y<sup>1</sup> may be, for example, -NR<sub>3</sub>-, -S-, -O-, -C(=O)NR<sub>3</sub>-, -NR<sub>3</sub>C(=O)-, or -C(=O)O-, wherein R<sub>3</sub> is hydrogen or optionally substituted C<sub>1</sub>-C<sub>6</sub> alkyl such as -CH<sub>2</sub>CH<sub>2</sub>OH;

In the radical L<sup>1</sup>, examples of Alk<sup>1</sup> and Alk<sup>2</sup> radicals, when present, include -CH<sub>2</sub>-, -CH<sub>2</sub>CH<sub>2</sub>-, -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-, -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-, -CH=CH-, -CH=CHCH<sub>2</sub>-, -CH<sub>2</sub>CH=CH-, CH<sub>2</sub>CH=CHCH<sub>2</sub>-, -C≡C-, -C≡CCH<sub>2</sub>-, -CH<sub>2</sub>C≡C-, and CH<sub>2</sub>C≡CCH<sub>2</sub>. Additional examples of Alk<sup>1</sup> and Alk<sup>2</sup> include -CH<sub>2</sub>W-, -CH<sub>2</sub>CH<sub>2</sub>W-, -CH<sub>2</sub>CH<sub>2</sub>WCH<sub>2</sub>-, -CH<sub>2</sub>CH<sub>2</sub>WCH(CH<sub>3</sub>)-, -CH<sub>2</sub>WCH<sub>2</sub>CH<sub>2</sub>-, -CH<sub>2</sub>WCH<sub>2</sub>CH<sub>2</sub>WCH<sub>2</sub>-, and -WCH<sub>2</sub>CH<sub>2</sub>- where W is -O-, -S-, -NH-, -N(CH<sub>3</sub>)-, or -CH<sub>2</sub>CH<sub>2</sub>N(CH<sub>2</sub>CH<sub>2</sub>OH)CH<sub>2</sub>-.


Alk<sup>1</sup> and Alk<sup>2</sup> when present may also be branched chain alkyl such as -CH(CH<sub>3</sub>)-, -C(CH<sub>3</sub>)<sub>2</sub>- or in either orientation -CH<sub>2</sub>CH(CH<sub>3</sub>)-, -CH<sub>2</sub>C(CH<sub>3</sub>)<sub>2</sub>-.

In L<sup>1</sup>, when n is 0, the radical is a hydrocarbon chain (optionally substituted and perhaps having an ether, thioether or amino linkage). Presently it is preferred that there be no optional substituents in L<sup>1</sup>. When both m and p are 0, L<sup>1</sup> is a divalent mono- or bicyclic carbocyclic or heterocyclic radical with 5 - 13 ring atoms (optionally substituted). When n is 1 and at least one of m and p is 1, L<sup>1</sup> is a

divalent radical including a hydrocarbon chain or chains and a mono- or bicyclic carbocyclic or heterocyclic radical with 5 - 13 ring atoms (optionally substituted). When present, Q may be, for example, a divalent phenyl, naphthyl, cyclopropyl, cyclopentyl, or cyclohexyl radical, or a mono-, or bi-cyclic heterocyclic radical having 5 to 13 ring members, such as piperidinyl, piperazinyl, indolyl, pyridyl, thienyl, or pyrrolyl radical, but 1,4-phenylene is presently preferred.

Specifically, in some embodiments of the invention, L<sup>1</sup>, m and p may be 0 with n being 1. In other embodiments, n and p may be 0 with m being 1. In further embodiments, m, n and p may be all 0. In still further embodiments m may be 0, n may be 1 with Q being a monocyclic heterocyclic radical, and p may be 0 or 1. Alk<sup>1</sup> and Alk<sup>2</sup>, when present, may be selected from -CH<sub>2</sub>-, -CH<sub>2</sub>CH<sub>2</sub>-, and -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>- and Q may be 1,4-phenylene.

Specific examples of the radical -L<sup>1</sup>-Y<sup>1</sup>-[CH<sub>2</sub>]<sub>z</sub>- include -(CH<sub>2</sub>)<sub>3</sub>NH-, -CH<sub>2</sub>C(=O)NH-, -CH<sub>2</sub>CH<sub>2</sub>C(=O)NH-, -CH<sub>2</sub>C(O)O-, -CH<sub>2</sub>S-, -CH<sub>2</sub>CH<sub>2</sub>C(O)O-, -(CH<sub>2</sub>)<sub>4</sub>NH-, -CH<sub>2</sub>CH<sub>2</sub>S-, -CH<sub>2</sub>O-, -CH<sub>2</sub>CH<sub>2</sub>O-,



In a specific preferred example, useful when ring A is a phenyl ring, of the radical -L<sup>1</sup>-Y<sup>1</sup>-, L<sup>1</sup> is C<sub>1</sub>-C<sub>3</sub> alkylene, eg -CH<sub>2</sub>-, -CH<sub>2</sub>CH<sub>2</sub>- or -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>-, and Y<sup>1</sup> is -NHC(=O)-.

As mentioned above, the compounds with which the invention is concerned are inhibitors of PLK1 kinase activity and are therefore of use for treatment of cell proliferative diseases such as cancer.

It will be understood that the specific dose level for any particular patient will depend upon a variety of factors including the activity of the specific compound employed, the

age, body weight, general health, sex, diet, time of administration, route of administration, rate of excretion, drug combination and the severity of the particular disease undergoing treatment. Optimum dose levels and frequency of dosing will be determined by clinical trial.

The compounds with which the invention is concerned may be prepared for administration by any route consistent with their pharmacokinetic properties. The orally administrable compositions may be in the form of tablets, capsules, powders, granules, lozenges, liquid or gel preparations, such as oral, topical, or sterile parenteral solutions or suspensions. Tablets and capsules for oral administration may be in unit dose presentation form, and may contain conventional excipients such as binding agents, for example syrup, acacia, gelatin, sorbitol, tragacanth, or polyvinyl-pyrrolidone; fillers for example lactose, sugar, maize-starch, calcium phosphate, sorbitol or glycine; tabletting lubricant, for example magnesium stearate, talc, polyethylene glycol or silica; disintegrants for example potato starch, or acceptable wetting agents such as sodium lauryl sulphate. The tablets may be coated according to methods well known in normal pharmaceutical practice. Oral liquid preparations may be in the form of, for example, aqueous or oily suspensions, solutions, emulsions, syrups or elixirs, or may be presented as a dry product for reconstitution with water or other suitable vehicle before use. Such liquid preparations may contain conventional additives such as suspending agents, for example sorbitol, syrup, methyl cellulose, glucose syrup, gelatin hydrogenated edible fats; emulsifying agents, for example lecithin, sorbitan monooleate, or acacia; non-aqueous vehicles (which may include edible oils), for example almond oil, fractionated coconut oil, oily esters such as glycerine, propylene glycol, or ethyl alcohol; preservatives, for example methyl or propyl p-hydroxybenzoate or sorbic acid, and if desired conventional flavouring or colouring agents.

For topical application to the skin, the drug may be made up into a cream, lotion or ointment. Cream or ointment formulations which may be used for the drug are conventional formulations well known in the art, for example as described in standard textbooks of pharmaceutics such as the British Pharmacopoeia.

For topical application by inhalation, the drug may be formulated for aerosol delivery for example, by pressure-driven jet atomizers or ultrasonic atomizers, or preferably by

propellant-driven metered aerosols or propellant-free administration of micronized powders, for example, inhalation capsules or other "dry powder" delivery systems. Excipients, such as, for example, propellants (e.g. Frigen in the case of metered aerosols), surface-active substances, emulsifiers, stabilizers, preservatives, flavourings, and fillers (e.g. lactose in the case of powder inhalers) may be present in such inhaled formulations. For the purposes of inhalation, a large number of apparae are available with which aerosols of optimum particle size can be generated and administered, using an inhalation technique which is appropriate for the patient. In addition to the use of adaptors (spacers, expanders) and pear-shaped containers (e.g. Nebulator®, Volumatic®), and automatic devices emitting a puffer spray (Autohaler®), for metered aerosols, in particular in the case of powder inhalers, a number of technical solutions are available (e.g. Diskhaler®, Rotadisk®, Turbohaler® or the inhalers for example as described in European Patent Application EP 0 505 321).

For topical application to the eye, the drug may be made up into a solution or suspension in a suitable sterile aqueous or non aqueous vehicle. Additives, for instance buffers such as sodium metabisulphite or disodium edetate; preservatives including bactericidal and fungicidal agents such as phenyl mercuric acetate or nitrate, benzalkonium chloride or chlorhexidine, and thickening agents such as hypromellose may also be included.

The active ingredient may also be administered parenterally in a sterile medium. Depending on the vehicle and concentration used, the drug can either be suspended or dissolved in the vehicle. Advantageously, adjuvants such as a local anaesthetic, preservative and buffering agents can be dissolved in the vehicle.

The compounds of the invention may be used in conjunction with a number of known pharmaceutically active substances. For example, the compounds of the invention may be used with cytotoxics, HDAC inhibitors, kinase inhibitors, aminopeptidase inhibitors, protease inhibitors, bcl-2 antagonists, inhibitors of mTor and monoclonal antibodies (for example those directed at growth factor receptors). Preferred cytotoxics include, for example, taxanes, platins, anti-metabolites such as 5-fluoracil, topoisomerase inhibitors and the like. The medicaments of the invention comprising amino acid derivatives of formula (I), tautomers thereof or pharmaceutically acceptable salts, N-oxides, hydrates

or solvates thereof therefore typically further comprise a cytotoxic, an HDAC inhibitor, a kinase inhibitor, an aminopeptidase inhibitor and/or a monoclonal antibody.

Further, the present invention provides a pharmaceutical composition comprising:

- (a) a compound (I), or a pharmaceutically acceptable salt, N-oxide, hydrate or solvate thereof;
- (b) a cytotoxic agent, an HDAC inhibitor, a kinase inhibitor, an aminopeptidase inhibitor, a protease inhibitor, a bcl-2 antagonist, an inhibitor of mTor and/or a monoclonal antibody; and
- (c) a pharmaceutically acceptable carrier or diluent.

Also provided is a product comprising:

- (a) a compound (I), or a pharmaceutically acceptable salt, N-oxide, hydrate or solvate thereof; and
- (b) a cytotoxic agent, an HDAC inhibitor, a kinase inhibitor, an aminopeptidase inhibitor, a protease inhibitor, a bcl-2 antagonist, an inhibitor of mTor and/or a monoclonal antibody,

for the separate, simultaneous or sequential use in the treatment of the human or animal body.

### **Synthesis**

There are multiple synthetic strategies for the synthesis of the compounds (I) with which the present invention is concerned, but all rely on known chemistry, known to the synthetic organic chemist. Thus, compounds according to formula (I) can be synthesised according to procedures described in the standard literature and are well-known to those skilled in the art. Typical literature sources are "*Advanced organic chemistry*", 4<sup>th</sup> Edition (Wiley), J March; "*Comprehensive Organic Transformation*", 2<sup>nd</sup> Edition (Wiley), R.C. Larock, "*Handbook of Heterocyclic Chemistry*", 2<sup>nd</sup> Edition (Pergamon), A.R. Katritzky; review articles such as found in "*Synthesis*", "*Acc. Chem. Res.*", "*Chem. Rev.*", or primary literature sources identified by standard literature searches online or from secondary sources such as "*Chemical Abstracts*" or "*Beilstein*".

The compounds of the invention may be prepared by a number of processes some of which are described specifically in the Examples below. In the reactions described below, it may be necessary to protect reactive functional groups, for example hydroxyl,

amino and carboxy groups, where these are desired in the final product, to avoid their unwanted participation in the reactions [see for example, "*Protecting Groups in Organic Synthesis*", 3<sup>rd</sup> Edition, (Wiley), T.W. Greene]. Conventional protecting groups may be used in conjunction with standard practice. In some instances deprotection may be the final step in the synthesis of a compound of general formula (I), and the processes according to the invention described herein after are understood to extend to such removal of protecting groups.

### Abbreviations

AcOH = acetic acid  
EDC = 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide  
EtOAc = ethyl acetate  
EtOH = ethanol  
Et<sub>2</sub>O = diethyl ether  
Et<sub>3</sub>N = triethylamine  
Boc or boc = *tert*-butoxycarbonyl  
Boc<sub>2</sub>O = Di-*tert*-butyldicarbonate  
Cbz = benzyloxycarbonyl  
DCE = dichloroethane  
DCM = dichloromethane  
DIPEA = diisopropylethylamine  
DMAP = dimethylamino pyridine  
DMF = dimethylformamide  
DMSO = dimethyl sulfoxide  
HCl = hydrochloric acid  
K<sub>2</sub>CO<sub>3</sub> = potassium carbonate  
LiOH = lithium hydroxide  
MeOH = methanol  
MgSO<sub>4</sub> = magnesium sulphate  
Na<sub>2</sub>CO<sub>3</sub> = sodium carbonate  
NaH = sodium hydride  
NaHCO<sub>3</sub> = sodium hydrogen carbonate  
NaI = sodium iodide  
NaOH = sodium hydroxide

NBu<sub>4</sub>Br = tetrabutylammonium bromide

Pd(dppf)Cl<sub>2</sub> = dichloro-(1,2-bis-(diphenylphosphino)ethane)-palladium(II)

Pd/C = palladium on carbon

STAB = sodium triacetoxyborohydride

TBTU = O-benzotriazol-1-yl-N,N,N',N'-tetramethyluronium tetrafluoroborate

TFA = trifluoroacetic acid

THF = tetrahydrofuran

aq = aqueous

g = gram(s)

LCMS = high performance liquid chromatography/mass spectrometry

mg = milligram(s)

min = minutes

mL = milliliter(s)

µL = microlitre(s)

mol = mole(s)

mmol = millimole(s)

NMR = nuclear magnetic resonance

RT or rt = room temperature

sat = saturated

Commercially available reagents and solvents (HPLC grade) were used without further purification. Solvents were removed using a Buchi rotary evaporator. Microwave irradiation was carried out using a Biotage Initiator™ Eight microwave synthetiser. Purification of compounds by flash chromatography column was performed using silica gel, particle size 40–63µm (230-400 mesh) obtained from Fluorochrom. Purification of compounds by preparative HPLC was performed on Gilson systems using reverse phase Axia™ prep Luna C18 columns (10µm, 100 x 21.2mm), gradient 0-100% B (A = water / 0.05% TFA, B = acetonitrile) over 10 min, flow = 25mL/min, UV detection at 254nm.

<sup>1</sup>H NMR spectra were recorded on a Bruker 300 MHz AV spectrometer in deuterated solvents. Chemical shifts ( $\delta$ ) are in parts per million. Thin-layer chromatography (TLC)

analysis was performed with Kieselgel 60 F<sub>254</sub> (Merck) plates and visualized using UV light.

Analytical HPLC/MS was performed on an Agilent HP1100 LC system using reverse phase Luna C18 columns (3μm, 50 x 4.6mm), gradient 5-95% B ( A = water / 0.1% Formic acid, B = acetonitrile / 0.1% Formic acid) over 13.0 min, flow = 1.25mL/min. UV spectra were recorded at 220 and 254nm using a G1315B DAD detector. Mass spectra were obtained over the range m/z 150 to 800 on a LC/MSD SL G1956B detector. Data were integrated and reported using ChemStation and ChemStation Data Browser softwares.

### Intermediates

The intermediates for the preparation of the examples described herein are shown below (Figure 1):






Figure 1

**Intermediate A:**(7R)-2-Chloro-8-cyclopentyl-7-ethyl-5-methyl-7,8-dihydropteridin-6(5H)-one

The title compound was prepared using methodology described in WO2004076454.

**Intermediate B:**


4-{[(7R)-8-Cyclopentyl-7-ethyl-5-methyl-6-oxo-5,6,7,8-tetrahydropteridin-2-yl]amino}-3-methoxybenzoic acid




The title compound was prepared using methodology described in WO2004076454.

**Intermediate C:**

4-{[(7R)-8-Cyclopentyl-7-ethyl-5-methyl-6-oxo-5,6,7,8-tetrahydropteridin-2-yl]amino}-3-methoxy-N-piperidin-4-ylbenzamide



The title compound was prepared from **Intermediate B** by the following methodology:



**Scheme 1**

**Stage 1** - *tert*-butyl 4-[(4-[(7*R*)-8-cyclopentyl-7-ethyl-5-methyl-6-oxo-5,6,7,8-tetrahydropteridin-2-yl]amino)-3-methoxybenzoyl]amino)piperidine-1-carboxylate

To a suspension of **Intermediate B** (500mg, 1.18mmol) in DCM (20mL) was added TBTU (415mg, 1.29mmol) and DIPEA (0.41mL, 2.35mmol). The reaction mixture was stirred at RT for 30 min and *tert*-butyl 4-aminopiperidine-1-carboxylate (282mg, 1.41mmol) was added. The reaction mixture was stirred at RT for 30 min, diluted with DCM (30mL), washed with water (2 x 30mL), dried ( $\text{MgSO}_4$ ), and concentrated under reduced pressure to leave a thick brown oil. Trituration with  $\text{Et}_2\text{O}$ /heptane (1:3) afforded the title product as a beige solid (528mg, 74%). ESMS m/z: 608 [M+H]<sup>+</sup>.


**Stage 2** - 4-[(7*R*)-8-Cyclopentyl-7-ethyl-5-methyl-6-oxo-5,6,7,8-tetrahydropteridin-2-yl]amino)-3-methoxy-N-piperidin-4-ylbenzamide

**Stage 1** product (528mg, 0.87mmol) was suspended in a 4N HCl in dioxane (10mL) and the reaction mixture was stirred at RT for 1 hour and then concentrated under reduced pressure. The residue was triturated with  $\text{Et}_2\text{O}$  and then partitioned between DCM (100mL) and sat  $\text{Na}_2\text{CO}_3$  (50mL). The organic layer was separated, washed with sat  $\text{Na}_2\text{CO}_3$  (50mL), dried ( $\text{MgSO}_4$ ) and concentrated under reduced pressure to afford the title intermediate as a thick yellow oil, which solidified on standing (407mg, 92%). ESMS m/z 508 [M+H]<sup>+</sup>. <sup>1</sup>H NMR (300 MHz,  $\text{CDCl}_3$ )  $\delta$ : 8.56 (1H, dd, J=8.4, 3.5 Hz), 7.57-7.76 (2H, m), 7.39-7.44 (1H, m), 4.53 (1H, br.s.), 4.08-4.34 (2H, m), 3.98 (3H, d, J=4.7Hz),


3.39-3.65 (2H, m), 3.29-3.38 (3H, m), 2.81-3.15 (2H, m), 1.41-2.44 (14H, m), 0.75-0.97 (3H, m).

**Intermediate D:**

Methyl N-(tert-butoxycarbonyl)-4-hydroxyisovalinate



The title compound was prepared by the following methodology:



**Scheme 2**

**Stage 1 - 2-amino-4-(benzyloxy)-2-methylbutanenitrile**

A 2-neck round-bottomed flask (100mL) was attached via an adaptor and rubber tubing to a funnel submerged in bleach. To this round-bottomed flask was added 98% sodium cyanide (1.81g, 36.9mmol) in water (3mL), followed by ammonium chloride (2.17g, 40.6mmol) in lukewarm water (5mL). Ammonium hydroxide (2.88mL, 73.8mmol) was then added, followed by 4-benzyloxy-2-butanone (6.58g, 36.9mmol) in ethanol (11mL).

The resulting mixture was stirred at RT for 15 min, then at 60°C for 2 hours. After cooling to RT the reaction mixture was concentrated under reduced pressure and partitioned between water (100mL) and DCM (100mL). The aqueous layer was extracted with DCM (3 x 75mL). The combined organic layers were dried ( $\text{MgSO}_4$ ), filtered and concentrated under reduced pressure. Purification on a 12g silica column using a CombiFlash® Companion® (Teledyne Isco Inc) (EtOAc in heptane) gave the title product as a colourless oil (4.70g, 62%). ESMS m/z: 205 [M+H]<sup>+</sup>; <sup>1</sup>H NMR (300 MHz,  $\text{CDCl}_3$ )  $\delta$ : 7.36 (2H, br. s.), 7.33-7.25 (3H, m), 4.56-4.49 (2H, m), 3.73 (2H, t,  $J$ =6.1 Hz), 2.67-2.27 (2H, m), 1.49 (3H, s).

#### Stage 2 - Methyl 4-(benzyloxy)isovalinate


A 3-neck round-bottomed flask (500mL) was attached via an adaptor and rubber tubing to a funnel submerged in saturated  $\text{NaHCO}_3$ . The round-bottomed flask was cooled to 0°C with an ice bath and anhydrous methanol (50mL) was added. The methanol was saturated with HCl (g) for 5 min. **Stage 1** product (2.0g, 9.8mmol) in methanol (7mL) was added to the reaction mixture and stirred at 65°C for 48 hours. The reaction mixture was cooled to RT, concentrated under reduced pressure and partitioned between water (50mL) and EtOAc (50mL). The aqueous layer was isolated, saturated  $\text{NaHCO}_3$  added and the product extracted into EtOAc (3 x 50mL). The combined organic layers were dried ( $\text{MgSO}_4$ ), filtered and concentrated under reduced pressure to afford the crude product as a dark yellow oil (2.32g, 100%). ESMS m/z: 238 [M+H]<sup>+</sup>; <sup>1</sup>H NMR (300 MHz,  $\text{CDCl}_3$ )  $\delta$ : 7.35-7.26 (5H, m), 5.39 (2H, br. s.), 4.57 (2H, d,  $J$ =5.8 Hz), 3.64 (3H, s), 3.61-3.52 (2H, m), 2.28-2.14 (2H, m), 1.53 (3H, s).

#### Stage 3 - Methyl 4-(benzyloxy)-N-(*tert*-butoxycarbonyl)isovalinate

To a solution of **stage 2** product (2.32g, 9.78mmol) in DCM (20mL) was added  $\text{Et}_3\text{N}$  (6.8mL, 48.9mmol). The reaction mixture was cooled to 0°C in an ice bath and to it was added  $\text{Boc}_2\text{O}$  (2.56g, 11.7mmol) in 2 portions. The reaction was stirred at RT for 24 hours and the product was extracted onto silica under reduced pressure. Purification on a 12g silica column using a CombiFlash® Companion® (Teledyne Isco Inc) (EtOAc in heptane) gave the title compound as a colourless oil (0.37g, 15%). ESMS m/z: 360 [M+Na]<sup>+</sup>; <sup>1</sup>H NMR (300 MHz,  $\text{CDCl}_3$ )  $\delta$ : 7.46-7.18 (5H, m), 6.03 (1H, br. s.), 4.45 (2H, m), 3.64 (3H, s), 3.55 (2H, t,  $J$ =5.6 Hz), 2.41-2.07 (2H, m), 1.58 (3H, s), 1.44 (9H, s).


**Stage 4 - Methyl N-(*tert*-butoxycarbonyl)-4-hydroxyisovalinate**

To a solution of **stage 3** product (0.370g, 1.10mmol) in EtOAc (15mL) was added 10% Pd/C (0.074g, 20% w.w.). The system was evacuated and put under a hydrogen atmosphere (using a 3-way tap apparatus and hydrogen-filled balloon), this was repeated twice and the mixture was allowed to stir for 24 h at RT under a hydrogen atmosphere. The system was evacuated of hydrogen and the palladium residues filtered over Celite. The Celite was washed thoroughly with ethyl acetate and the filtrate solvent removed under reduced pressure. The crude residue was purified by column chromatography (50% EtOAc in heptane) to afford the title compound as a colourless oil (0.251g, 93%). ESMS m/z: 270 [M+Na]<sup>+</sup>; <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ: 5.84 (1H, br. s.), 3.80-3.64 (5H, s), 2.30-2.07 (2H, m), 1.58 (3H, s), 1.43 (9H, s).


**Intermediate E:**Cyclopentyl N-(*tert*-butoxycarbonyl)-4-hydroxyisovalinate

The title compound was prepared using the same methodology described for **Intermediate D**, replacing the methanol in stage 2 with cyclopentanol. ESMS m/z: 324 [M+Na]<sup>+</sup>; <sup>1</sup>H NMR (300 MHz, CDCl<sub>3</sub>) δ: 5.99-5.75 (1H, m), 5.15 (1H, t, J=5.6 Hz), 3.67 (2H, t, J=5.9 Hz), 2.17-1.92 (2H, m), 1.89-1.51 (8H, m), 1.49 (3H, s), 1.38 (9H, s).

**Example 1:**Methyl 4-{4-[(4-[(7*R*)-8-cyclopentyl-7-ethyl-5-methyl-6-oxo-5,6,7,8-tetrahydropteridin-2-yl]amino)-3-methoxybenzoyl]amino}piperidin-1-yl}isovalinate



The title compound was prepared by the following methodology:



*Scheme 3*

#### Stage 1 - Methyl 4-oxoisovalinate

A solution of DCM (5mL) and oxalyl chloride (0.055mL, 0.62mmol) was stirred under a nitrogen atmosphere and cooled to an internal temperature of -70°C. DMSO (0.040mL, 0.56mmol) was added slowly, ensuring the temperature remained below -65°C. A solution of **Intermediate D** (0.068g, 0.26mmol) in DCM (2mL) was added slowly, followed by Et<sub>3</sub>N (0.310mL, 2.20mmol), keeping the temperature below -65°C at all times. The reaction was then allowed to warm to RT and the solvent was removed under reduced pressure to afford the crude product as a pale yellow solid. This crude mixture was taken on to the reductive amination step without further purification.

#### Stage 2 - Methyl *N*-(*tert*-butoxycarbonyl)-4-{4-[(4-[(7*R*)-8-cyclopentyl-7-ethyl-5-methyl-6-oxo-5,6,7,8-tetrahydropteridin-2-yl]amino)-3-methoxybenzoyl]amino}piperidin-1-yl} isovalinate


A solution of **Intermediate C** (0.131g, 0.26mmol) and **stage 1** product (0.064g, 0.26mmol) in DCE (3.5mL) was stirred at RT for 15 min. Sodium triacetoxyborohydride (0.165g, 0.78mmol) was added and the reaction mixture stirred at RT overnight.

Saturated NaHCO<sub>3</sub> (10mL) and DCM (20mL) were added and the aqueous layer extracted with DCM (3 x 15mL). The combined organic layers were dried (MgSO<sub>4</sub>), filtered and concentrated under reduced pressure to afford the crude product as a yellow oil (0.210g). ESMS m/z: 737 [M+H]<sup>+</sup>. <sup>1</sup>H NMR (300 MHz, MeOD), 7.90 (1H, d, J=8.4Hz), 7.64 (2H, s), 7.59 (1H, d J=9.8Hz), 4.52 (1H, dd, J=3.2, 6.4Hz), 4.33 (1H, t J=8.5Hz), 4.30-4.18 (1H, m), 4.00 (3H, m), 3.93 (3H, s), 3.76-3.69 (2H, m), 3.33 (3H, s), 3.27-3.15 (4H, m), 2.46 (2H, t, J=8.4Hz), 2.28 (2H, d J=12.8Hz), 2.08-1.90 (8H, m), 1.67-1.56 (7H, m), 1.38 (9H, s), 0.88 (3H, t, J=7.4Hz).

**Stage 3 - Methyl 4-{4-[(4-{[(7R)-8-cyclopentyl-7-ethyl-5-methyl-6-oxo-5,6,7,8-tetrahydropteridin-2-yl]amino}-3-methoxybenzoyl)amino]piperidin-1-yl}isovalinate**  
 To a solution of **stage 2** product (0.030g, 0.04mmol) in DCM (1mL) was added 4N HCl in dioxane (3mL). The reaction was stirred at RT for 2 hours and the solvent was removed under reduced pressure. The resulting residue was taken up in a 1:1 ratio of acetonitrile and water (1.3mL) and purified by preparative HPLC. The pure fractions were combined and dried on the freeze-drier to afford the title compound as a white solid. (0.023g, 12%). ESMS m/z: 637 [M+H]<sup>+</sup>; <sup>1</sup>H NMR (300 MHz, MeOD), 7.90 (1H, d, J=8.4Hz), 7.64 (2H, s), 7.59 (1H, d J=9.8Hz), 4.52 (1H, dd, J=3.2, 6.4Hz), 4.33 (1H, t J=8.5Hz), 4.30-4.18 (1H, m), 4.00 (3H, m), 3.93 (3H, s), 3.76-3.69 (2H, m), 3.33 (3H, s), 3.27-3.15 (4H, m), 2.46 (2H, t, J=8.4Hz), 2.28 (2H, d J=12.8Hz), 2.08-1.90 (8H, m), 1.67-1.56 (7H, m), 0.88 (3H, t, J=7.4Hz).

**Example 2:**


Cyclopentyl 4-{4-[(4-{[(7R)-8-cyclopentyl-7-ethyl-5-methyl-6-oxo-5,6,7,8-tetrahydropteridin-2-yl]amino}-3-methoxybenzoyl)amino]piperidin-1-yl}isovalinate




This compound was prepared from **Intermediate E** using the same methodology described for **Example 1**. ESMS m/z: 691 [M+H]<sup>+</sup>; <sup>1</sup>H NMR (300 MHz, MeOD), 7.89 (1H, d, J=8.3Hz), 7.63 (2H, s), 7.58 (1H, dd, J=8.4, 1.6Hz), 5.35 (1H, t, J=5.6Hz), 4.51 (1H, dd, J=6.4, 3.0Hz), 4.39-4.12 (2H, m), 3.99 (3H, s), 3.71 (2H, br.s.), 3.33 (3H, br.s.), 3.29-3.07 (4H, m), 2.43 (2H, t, J=8.5Hz), 2.27 (2H, d, J=14.3Hz), 2.16-1.66 (20H, m), 1.64 (3H, s), 0.87 (3H, t, J=7.4Hz).

**Example 3:**

4-{4-[4-{[(7R)-8-cyclopentyl-7-ethyl-5-methyl-6-oxo-5,6,7,8-tetrahydropteridin-2-yl]amino}-3-methoxybenzoyl]amino}piperidin-1-yl}isovaline



The title compound was prepared by the following methodology:



*Scheme 4*

To a solution of **Example 1 stage 2** product (0.150g, 0.20mmol) in THF (5mL) and water (5mL) was added LiOH (0.098g, 4.08mmol). The reaction mixture was stirred at 40°C overnight. THF was removed under reduced pressure and the resulting solution was acidified to pH 1-2 with 2N HCl. The product was extracted into *tert*-butanol (3 x 30mL) and concentrated to dryness under reduced pressure. The product was slurried in EtOAc and isolated by filtration to give 0.09g of solid. 0.03g was treated with 4N HCl in dioxane (1mL) for 1 hour. Solvent was removed under reduced pressure, product taken in 1:1

ratio of AcCN / H<sub>2</sub>O (1.3mL) and purified by preparative HPLC. The pure fractions were combined and dried on the freeze-drier to afford the title compound as a white solid. (0.010g, 7.9%). ESMS m/z: 623 [M+H]<sup>+</sup>; <sup>1</sup>H NMR (300 MHz, MeOD), 7.95 (1H, d, J=8.3Hz), 7.64 (2H, s), 7.58 (1H, dd J=1.8, 8.3Hz), 4.51 (1H, dd, J=4.0, 6.6Hz), 4.35 (1H, t J=9.1Hz), 4.27-4.18 (1H, m), 4.00 (3H, m), 3.72-3.663 (2H, m), 3.33 (3H, s), 3.27-3.15 (4H, m), 2.46-2.41 (2H, m), 2.30-2.24 (2H, m), 2.10-1.88 (8H, m), 1.70-1.63 (7H, m), 0.88 (3H, t, J=7.4Hz).

### **Measurement of Biological Activity**

#### ***PLK1 Enzyme Assay***

The ability of compounds to inhibit PLK-1 kinase activity was measured in an assay performed by Invitrogen (Paisley, UK). The Z'-LYTE™ biochemical assay employs a fluorescence-based, coupled-enzyme format and is based on the differential sensitivity of phosphorylated and non-phosphorylated peptides to proteolytic cleavage. The peptide substrate is labelled with two fluorophores—one at each end—that make up a FRET pair. In the primary reaction, the kinase transfers the gamma-phosphate of ATP to a single serine or threonine residue in a synthetic FRET-peptide. In the secondary reaction, a site-specific protease recognizes and cleaves non-phosphorylated FRET-peptides. Phosphorylation of FRET-peptides suppresses cleavage by the Development Reagent. Cleavage disrupts FRET between the donor (i.e., coumarin) and acceptor (i.e., fluorescein) fluorophores on the FRET-peptide, whereas uncleaved, phosphorylated FRET-peptides maintain FRET. A radiometric method, which calculates the ratio (the Emission Ratio) of donor emission to acceptor emission after excitation of the donor fluorophore at 400nm, is used to quantitate reaction progress.

The final 10  $\mu$ L Kinase Reaction consists of 2.8-25.3ng PLK1, 2 $\mu$ M Ser/Thr 16 Peptide substrate and ATP in 50mM HEPES pH 7.5, 0.01% BRIJ-35, 10mM MgCl<sub>2</sub>, 1mM EGTA. The assay is performed at an ATP concentration at, or close to, the Km. After the 60 minute Kinase Reaction incubation at RT, 5 $\mu$ L of a 1:8 dilution of Development Reagent is added. The assay plate is incubated for a further 60 minutes at RT and read on a fluorescence plate reader.

Duplicate data points are generated from a 1/3 log dilution series of a stock solution of test compound in DMSO. Nine dilutions steps are made from a top concentration of 10 $\mu$ M, and a 'no compound' blank is included. Data is collected and analysed using *XLfit* software from IDBS. The dose response curve is curve fitted to model number 205 (sigmoidal dose-response model). From the curve generated, the concentration giving 50% IC50 inhibition is determined and reported.

IC50 results were allocated to one of 3 ranges as follows:

Range A: IC50<100nM,

Range B: IC50 from 100nM to 500nM;

and Range C: IC50 >500nM.

NT = Not tested

The results obtained for compounds of the Examples herein are shown below (Table 1).

### ***Cell inhibition Assay***

Cells were seeded in 96W tissue culture plates (1 well = 30mm<sup>2</sup>) in 50 $\mu$ l of the appropriate culture medium (see below). Seeding densities were cell-line dependent: HCT-116 = 1000 cells per well, Hut-78 = 2250 cells per well, U937 cells = 2000 cells per well. 24hrs later 50 $\mu$ L of the compound prepared in the same medium was added as 4 fold dilutions to give final concentrations in the range 0.15nM-2500nM (n=6 for each concentration). The plates were then incubated at 37°C, 5% CO<sub>2</sub> for 72hrs. Cell proliferation was assessed using WST-1 (a metabolic indicator dye, Roche Cat no. 1 644 807) according to the manufacturers instructions. The results were calculated as percentage of vehicle response and IC50 values represent the concentration of compound that inhibited the vehicle response by 50%.

HCT-116 Culture Medium - Dulbeccos MEM ( Sigma D6546) plus 10% heat inactivated fetal calf serum (Hyclone SH30071 Thermo Fischer Scientific) containing 2mM Glutamine (Sigma cat no G-7513) and 50U/mL Penicillin and Streptomycin Sulphate (Sigma Cat no P-0781).

Hut-78 & U937 culture media: RPMI1640 (Sigma R0883) plus 10% heat inactivated fetal calf serum, as above and supplemented with 2mM Glutamine and 50U/mL Penicillin and Streptomycin Sulphate (details as above).

IC50 results were allocated to one of 3 ranges as follows:

Range A: IC50 <100nM,

Range B: IC50 from 100nM to 500nM;

and Range C: IC50 >500nM.

NT = Not tested

The results obtained for compounds of the Examples herein are shown below (Table 1).

Example Number	Inhibitor Activity vs . PLK1	Inhibitor Activity vs HCT 116 cell line	Inhibitor Activity vs Hut-78 cell line	Inhibitor Activity vs U937 cell line
1	A	B	A	A
2	A	A	A	A
3	A	NT	NT	NT

Table 1

#### Broken Cell Carboxylesterase Assay

Any given compound of the present invention wherein R<sub>7</sub> is an ester group, may be tested to determine whether it meets the requirement that it be hydrolysed by intracellular esterases, by testing in the following assay.

#### *Preparation of cell extract*

U937 or HCT 116 tumour cells (~ 10<sup>9</sup>) were washed in 4 volumes of Dulbeccos PBS (~ 1litre) and pelleted at 525g for 10min at 4°C. This was repeated twice and the final cell pellet was resuspended in 35mL of cold homogenising buffer (Trizma 10mM, NaCl 130mM, CaCl<sub>2</sub> 0.5mM pH 7.0 at 25°C). Homogenates were prepared by nitrogen cavitation (700psi for 50min at 4°C). The homogenate was kept on ice and supplemented with a cocktail of inhibitors at final concentrations of:

Leupeptin 1µM

Aprotinin 0.1µM

E64 8µM  
Pepstatin 1.5µM  
Bestatin 162µM  
Chymostatin 33µM

After clarification of the cell homogenate by centrifugation at 525g for 10min, the resulting supernatant was used as a source of esterase activity and was stored at -80°C until required.

#### ***Measurement of ester cleavage***

Hydrolysis of esters to the corresponding carboxylic acids can be measured using the cell extract, prepared as above. To this effect cell extract (~30µg / total assay volume of 0.5mL) was incubated at 37°C in a Tris- HCl 25mM, 125mM NaCl buffer, pH 7.5 at 25°C. At zero time the ester (substrate) was then added at a final concentration of 2.5 µM and the samples were incubated at 37°C for the appropriate time (usually 0 or 80min).

Reactions were stopped by the addition of 3 x volumes of acetonitrile. For zero time samples the acetonitrile was added prior to the ester compound. After centrifugation at 12000g for 5 min, samples were analysed for the ester and its corresponding carboxylic acid at RT by LCMS (Sciex API 3000, HP1100 binary pump, CTC PAL).

Chromatography was based on an AceCN (75x2.1mm) column and a mobile phase of 5-95% acetonitrile in water /0.1% formic acid.

The table below (Table 2) presents data showing that several amino acid ester motifs, conjugated to various intracellular enzyme inhibitors by several different linker chemistries are all hydrolysed by intracellular carboxyesterases to the corresponding acid.




















Structure of amino acid ester conjugate	R	Linker	Hydrolysis Rate Range U937Cells (pg/mL/min)	Preparation of amino ester conjugate
		$-\text{CH}_2\text{CH}_2\text{O}-$	100-1000	WO2006117552
		$-(\text{CH}_2)_3\text{O}-\text{C}_6\text{H}_4-\text{CH}_2\text{NHCH}_2-$	1000-50000	WO2006117548
		$-\text{CH}_2-\text{C}_6\text{H}_4-\text{CH}_2\text{NHCH}_2-$	>50000	WO2006117549
		$-\text{CH}_2\text{CH}_2\text{O}-$	>50000	WO2006117567
		$-\text{CH}_2\text{CH}_2\text{O}-$	1000-50000	WO2006117567
		$-\text{CH}_2-$	1000-50000	WO2006117567
		$-\text{CO}-$	>50000	WO2006117567
		$-\text{CH}_2-\text{C}_6\text{H}_4-\text{CH}_2\text{NHCH}_2-$	>50000	WO2006117549
		$-\text{CH}_2-\text{C}_6\text{H}_4-\text{CH}_2\text{NHCH}_2-$	>50000	WO2006117549

Table 2

**Claims:**

1. A compound of formula (I), or a salt thereof:



wherein

**R<sub>1</sub>** is hydrogen, or an optionally substituted (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>6</sub>)alkenyl, (C<sub>2</sub>-C<sub>6</sub>)alkynyl or (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl group;

**R<sub>2</sub>** is hydrogen, or an optionally substituted (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>6</sub>)alkenyl, (C<sub>2</sub>-C<sub>6</sub>)alkynyl or (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl group;

**R<sub>3</sub>** is hydrogen, -CN, hydroxyl, halogen, optionally substituted (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>2</sub>-C<sub>6</sub>)alkenyl, (C<sub>2</sub>-C<sub>6</sub>)alkynyl or (C<sub>3</sub>-C<sub>6</sub>)cycloalkyl, -NR<sub>5</sub>R<sub>6</sub> or C<sub>1</sub>-C<sub>4</sub> alkoxy, wherein R<sub>5</sub> and R<sub>6</sub> are independently hydrogen or optionally substituted (C<sub>1</sub>-C<sub>6</sub>)alkyl;

ring **A** is an optionally substituted mono- or bi-cyclic carbocyclic or heterocyclic ring or a ring system having up to 12 ring atoms;

**T** is a radical of formula R-L<sup>1</sup>-Y<sup>1</sup>- wherein

**Y<sup>1</sup>** is a bond, -O-, -S-, -NR<sub>6</sub>-, -(C=O)-, -S(O<sub>2</sub>)-, -(C=O)NR<sub>6</sub>-, -NR<sub>6</sub>(C=O)-, -S(O<sub>2</sub>)NR<sub>6</sub>-, -NR<sub>6</sub>S(O<sub>2</sub>)-, or -NR<sub>6</sub>(C=O)NR<sub>9</sub>-, wherein R<sub>6</sub> and R<sub>9</sub> are independently hydrogen or optionally substituted (C<sub>1</sub>-C<sub>6</sub>)alkyl;

**L<sup>1</sup>** is a divalent radical of formula -(Alk<sup>1</sup>)<sub>m</sub>(Q)<sub>n</sub>(Alk<sup>2</sup>)<sub>p</sub>- wherein


**m**, **n** and **p** are independently 0 or 1,

**Q** is (i) an optionally substituted divalent mono- or bicyclic carbocyclic or heterocyclic radical having 5 - 13 ring members, or (ii), in the case where p is 0, a

divalent radical of formula  $-Q^1-X^2-$  wherein  $X^2$  is  $-O-$ ,  $-S-$  or  $NR^A-$  wherein  $R^A$  is hydrogen or optionally substituted  $C_1-C_3$  alkyl, and  $Q^1$  is an optionally substituted divalent mono- or bicyclic carbocyclic or heterocyclic radical having 5 - 13 ring members,

**Alk<sup>1</sup>** and **Alk<sup>2</sup>** independently represent optionally substituted divalent ( $C_3-C_6$ )cycloalkyl radicals, or optionally substituted straight or branched, ( $C_1-C_6$ )alkylene, ( $C_2-C_6$ )alkenylene, or ( $C_2-C_6$ )alkynylene radicals which may optionally contain or terminate in an ether ( $-O-$ ), thioether ( $-S-$ ) or amino ( $-NR^A-$ ) link wherein  $R^A$  is hydrogen or optionally substituted ( $C_1-C_3$ )alkyl;

**R** is a radical of formula (X)



wherein

**R<sub>7</sub>** is a carboxylic acid group ( $-COOH$ ), or an ester group which is hydrolysable by one or more intracellular carboxylesterase enzymes to a carboxylic acid group;

**R<sup>1</sup><sub>7</sub>** is the side chain of a natural or non-natural alpha-amino acid, in which any functional groups are protected, but **R<sup>1</sup><sub>7</sub>** is not hydrogen;

**R<sub>8</sub>** is hydrogen; or optionally substituted  $C_1-C_6$  alkyl,  $C_3-C_7$  cycloalkyl, aryl or heteroaryl or  $-(C=O)R_6$ ,  $-(C=O)OR_6$ , or  $-(C=O)NR_6$  wherein  $R_6$  is hydrogen or optionally substituted ( $C_1-C_6$ )alkyl;

2. A compound as claimed in claim 1 wherein  $R_1$  is ethyl.
3. A compound as claimed in claim 1 or claim 2 wherein  $R_2$  is cyclopentyl.
4. A compound as claimed in any of the preceding claims wherein ring A is an optionally substituted phenyl ring.

5. A compound as claimed in any of the preceding claims wherein  $R_7$  is of formula  $-(C=O)OR_{10}$  wherein  $R_{10}$  is  $R_{11}R_{12}R_{13}C-$  wherein

- (i)  $R_{11}$  is hydrogen, fluorine or optionally substituted  $(C_1-C_3)alkyl-(Z^1)_a-[(C_1-C_3)alkyl]_b-$  or  $(C_2-C_3)alkenyl-(Z^1)_a-[(C_1-C_3)alkyl]_b-$  wherein a and b are independently 0 or 1 and  $Z^1$  is  $-O-$ ,  $-S-$ , or  $-NR_{14}-$  wherein  $R_{14}$  is hydrogen or  $(C_1-C_3)alkyl$ ; and  $R_{12}$  and  $R_{13}$  are independently hydrogen or  $(C_1-C_3)alkyl-$ ;
- (ii)  $R_{11}$  is hydrogen or optionally substituted  $R_{15}R_{16}N-(C_1-C_3)alkyl-$  wherein  $R_{15}$  is hydrogen or  $(C_1-C_3)alkyl$  and  $R_{16}$  is hydrogen or  $(C_1-C_3)alkyl$ ; or  $R_{15}$  and  $R_{16}$  together with the nitrogen to which they are attached form an optionally substituted monocyclic heterocyclic ring of 5- or 6- ring atoms or bicyclic heterocyclic ring system of 8 to 10 ring atoms, and  $R_{12}$  and  $R_{13}$  are independently hydrogen or  $(C_1-C_3)alkyl-$ ; or
- (iii)  $R_{11}$  and  $R_{12}$  taken together with the carbon to which they are attached form an optionally substituted monocyclic carbocyclic ring of from 3 to 7 ring atoms or bicyclic carbocyclic ring system of 8 to 10 ring atoms, and  $R_{13}$  is hydrogen.

and wherein in cases (i), (ii) and (iii) above, "alkyl" includes fluoroalkyl.

6. A compound as claimed in claim 5 wherein  $R_{10}$  is methyl, trifluoromethyl, ethyl, n- or iso-propyl, n-, sec- or tert-butyl, cyclohexyl, allyl, phenyl, benzyl, 2-, 3- or 4-pyridylmethyl, N-methylpiperidin-4-yl, tetrahydrofuran-3-yl, methoxyethyl, indanyl, norbonyl, dimethylaminoethyl, or morpholinoethyl.

7. A compound as claimed in claim 5 wherein  $R_{10}$  is cyclopentyl.

8. A compound as claimed in any of the preceding claims wherein  $R_7$  is phenyl, or heteroaryl such as pyridyl, or a group of formula  $-CR_aR_bR_c$  in which:

each of  $R_a$ ,  $R_b$  and  $R_c$  is independently hydrogen,  $(C_1-C_6)$ alkyl,  $(C_2-C_6)$ alkenyl,  $(C_2-C_6)$ alkynyl, phenyl( $C_1-C_6$ )alkyl,  $(C_3-C_8)$ cycloalkyl; or

$R_c$  is hydrogen and  $R_a$  and  $R_b$  are independently phenyl or heteroaryl such as pyridyl; or

$R_c$  is hydrogen,  $(C_1-C_6)$ alkyl,  $(C_2-C_6)$ alkenyl,  $(C_2-C_6)$ alkynyl, phenyl( $C_1-C_6$ )alkyl, or  $(C_3-C_8)$ cycloalkyl, and  $R_a$  and  $R_b$  together with the carbon atom to which they are attached form a 3 to 8 membered cycloalkyl or a 5- to 6-membered heterocyclic ring; or

$R_a$ ,  $R_b$  and  $R_c$  together with the carbon atom to which they are attached form a tricyclic ring (for example adamantyl); or

$R_a$  and  $R_b$  are each independently  $(C_1-C_6)$ alkyl,  $(C_2-C_6)$ alkenyl,  $(C_2-C_6)$ alkynyl, phenyl( $C_1-C_6$ )alkyl, or a group as defined for  $R_c$  below other than hydrogen, or  $R_a$  and  $R_b$  together with the carbon atom to which they are attached form a cycloalkyl or heterocyclic ring, and  $R_c$  is hydrogen, -OH, -SH, halogen, -CN, - $CO_2H$ ,  $(C_1-C_4)$ perfluoroalkyl, - $CH_2OH$ , - $O(C_1-C_6)$ alkyl, - $O(C_2-C_6)$ alkenyl, - $S(C_1-C_6)$ alkyl, - $SO(C_1-C_6)$ alkyl, - $SO_2(C_1-C_6)$ alkyl, - $S(C_2-C_6)$ alkenyl, - $SO(C_2-C_6)$ alkenyl, - $SO_2(C_2-C_6)$ alkenyl or a group -Q-W wherein Q represents a bond or -O-, -S-, -SO- or -SO<sub>2</sub>- and W represents a phenyl, phenylalkyl,  $(C_3-C_8)$ cycloalkyl,  $(C_3-C_8)$ cycloalkylalkyl,  $(C_4-C_8)$ cycloalkenyl,  $(C_4-C_8)$ cycloalkenylalkyl, heteroaryl or heteroarylalkyl group, which group W may optionally be substituted by one or more substituents independently selected from, hydroxyl, halogen, -CN, - $CONH_2$ , - $CONH(C_1-C_6)$ alkyl, - $CONH(C_1-C_6)$ alkyl<sub>2</sub>, -CHO, - $CH_2OH$ ,  $(C_1-C_4)$ perfluoroalkyl, - $O(C_1-C_6)$ alkyl, - $S(C_1-C_6)$ alkyl, - $SO(C_1-C_6)$ alkyl, - $SO_2(C_1-C_6)$ alkyl, - $NO_2$ , - $NH_2$ , - $NH(C_1-C_6)$ alkyl, - $N((C_1-C_6)$ alkyl)<sub>2</sub>, - $NHCO(C_1-C_6)$ alkyl,  $(C_1-C_6)$ alkyl,  $(C_2-C_6)$ alkenyl,  $(C_2-C_6)$ alkynyl,  $(C_3-C_8)$ cycloalkyl,  $(C_4-C_8)$ cycloalkenyl, phenyl or benzyl.

9. A compound as claimed in any of claims 1 to 7 wherein  $R^1_7$  is H-Alk<sup>4</sup>-, phenyl, monocyclic heterocycl,  $C_3-C_7$  cycloalkyl, phenyl(Alk<sup>4</sup>)-, heterocycl(Alk<sup>4</sup>)-, or  $C_3-C_7$  cycloalkyl(Alk<sup>4</sup>)-, wherein the heterocycl part is monocyclic heterocycl having 3-7 ring

atoms, and wherein -Alk<sup>4</sup>- is a straight or branched, divalent (C<sub>1</sub>-C<sub>6</sub>)alkylene, (C<sub>2</sub>-C<sub>6</sub>)alkenylene, or (C<sub>2</sub>-C<sub>6</sub>)alkynylene radical which may optionally be interrupted by, or terminate in, an ether (-O-), thioether (-S-) or amino (-NR<sup>A</sup>-) link wherein R<sup>A</sup> is hydrogen or optionally substituted (C<sub>1</sub>-C<sub>3</sub>)alkyl, and wherein the Alk<sup>4</sup>-, or cyclic part is optionally substituted.

10. A compound as claimed in any of claims 1 to 7 wherein R<sup>1</sup><sub>7</sub> is methyl, ethyl, n- or iso-propyl, or n-, sec- or tert-butyl.

11. A compound as claimed in any of claims 1 to 7 wherein R<sup>1</sup><sub>7</sub> is methyl.

12. A compound as claimed in any of the preceding claims wherein R<sub>8</sub> is hydrogen.

13. A compound as claimed in any of the preceding claims wherein L<sup>1</sup> is -CH<sub>2</sub>- , -CH<sub>2</sub>CH<sub>2</sub>- or -CH<sub>2</sub>CH<sub>2</sub>CH<sub>2</sub>- , and Y<sup>1</sup> is -NHC(=O)-.

14. A compound as claimed in claim 1 which is the subject of any of the Examples herein.

15. A pharmaceutical composition comprising a compound as claimed in any of the preceding claims, together with a pharmaceutically acceptable carrier.

16. The use of a compound as claimed in any of claims 1 to 14 in the preparation of a composition for inhibition of PLK1 activity in vitro or in vivo.

17. A method of treatment of conditions mediated by PLK1 activity, which comprises administering to a subject suffering such disease an effective amount of a compound of formula (I) as claimed in any of claims 1 to 14.

18. The use as claimed in claim 16 or a method as claimed in claim 17 for treatment of cell proliferative diseases

19. The use as claimed in claim 16 or a method as claimed in claim 17 for treatment of cancer.

# INTERNATIONAL SEARCH REPORT

International application No  
PCT/GB2009/001019

A. CLASSIFICATION OF SUBJECT MATTER  
INV. A61K31/519 C07D475/00 A61P35/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)  
A61K C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, BEILSTEIN Data, CHEM ABS Data, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 2006/046990 A1 (STADTMUELLER HEINZ [AT] ET AL) 2 March 2006 (2006-03-02) claims 1-6, 9, 10 -----	1-19
P, X	WO 2008/050096 A (CHROMA THERAPEUTICS LTD [GB]; MOFFAT DAVID FESTUS CHARLES [GB]; PATEL) 2 May 2008 (2008-05-02) the whole document -----	1-19

Further documents are listed in the continuation of Box C.

See parent family annex.

\* Special categories of cited documents :

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \*&\* document member of the same patent family

Date of the actual completion of the international search

2 July 2009

Date of mailing of the international search report

14/07/2009

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040,  
Fax: (+31-70) 340-3016

Authorized officer

Sotoca Usina, E

# INTERNATIONAL SEARCH REPORT

## Information on patent family members

International application No

PCT/GB2009/001019

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 2006046990	A1 02-03-2006	CA 2578560 A1 WO 2006021548 A1 JP 2008510771 T	02-03-2006 02-03-2006 10-04-2008
WO 2008050096	A 02-05-2008	NONE	