



Office de la Propriété
Intellectuelle
du Canada

Un organisme
d'Industrie Canada

Canadian
Intellectual Property
Office

An agency of
Industry Canada

CA 2954841 A1 2016/01/21

(21) **2 954 841**

**(12) DEMANDE DE BREVET CANADIEN
CANADIAN PATENT APPLICATION**

(13) A1

(86) **Date de dépôt PCT/PCT Filing Date:** 2015/07/16
(87) **Date publication PCT/PCT Publication Date:** 2016/01/21
(85) **Entrée phase nationale/National Entry:** 2017/01/10
(86) **N° demande PCT/PCT Application No.:** EP 2015/066263
(87) **N° publication PCT/PCT Publication No.:** 2016/008976
(30) **Priorité/Priority:** 2014/07/16 (EP14306153.9)

(51) **Cl.Int./Int.Cl. A61K 35/768** (2015.01),
A61P 35/00 (2006.01), **C07K 16/18** (2006.01)

(71) **Demandeur/Applicant:**
TRANSGENE SA, FR

(72) **Inventeurs/Inventors:**
SILVESTRE, NATHALIE, FR;
GEIST, MICHEL, FR;
RITTNER, KAROLA, FR;
MARCHAND, JEAN-BAPTISTE, FR;
THIOUDELLET, CHRISTINE, FR

(74) **Agent:** NORTON ROSE FULBRIGHT CANADA
LLP/S.E.N.C.R.L., S.R.L.

(54) **Titre : VIRUS ONCOLYTIQUES POUR L'EXPRESSION DE MODULATEURS DE POINTS DE CONTROLE IMMUNITAIRE**
(54) **Title: ONCOLYTIC VIRUS FOR EXPRESSION OF IMMUNE CHECKPOINT MODULATORS**

(57) Abrégé/Abstract:

The present invention provides an oncolytic virus comprising nucleotide sequence(s) encoding one or more immune checkpoint modulator(s). It also concerns a pharmaceutical composition comprising effective amount of said oncolytic virus and, eventually, a pharmaceutically acceptable vehicle and its use for treating proliferative diseases such as cancers.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(10) International Publication Number
WO 2016/008976 A1

(43) International Publication Date
21 January 2016 (21.01.2016)

(51) International Patent Classification:
A61K 35/768 (2015.01) C07K 16/18 (2006.01)

(21) International Application Number:
PCT/EP2015/066263

(22) International Filing Date:
16 July 2015 (16.07.2015)

(25) Filing Language:
English

(26) Publication Language:
English

(30) Priority Data:
14306153.9 16 July 2014 (16.07.2014) EP

(71) Applicant: **TRANSGENE SA [FR/FR]**; Parc d'Innovation Boulevard Gonthier d'Andernach, F-67400 Illkirch Grafenstaden (FR).

(72) Inventors: **SILVESTRE, Nathalie**; 6, rue des Iris, F-67120 Ergersheim (FR). **GEIST, Michel**; 18, rue Edouard Manet, F-67170 Brumath (FR). **RITTNER, Karola**; 51, rue des Pluviers, F-67100 Strasbourg (FR). **MARCHAND, Jean-Baptiste**; 1A rue des Hauts Pâturages, F-67210 Obernai (FR). **THIOUDELLET, Christine**; 4, boulevard de la Victoire, F-67000 Strasbourg (FR).

(74) Agent: **REGIMBEAU**; 20, rue de Chazelles, F-75847 Paris Cedex 17 (FR).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): **AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.**

AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): **ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).**

Declarations under Rule 4.17:

— of inventorship (Rule 4.17(iv))

Published:

— with international search report (Art. 21(3))

— with sequence listing part of description (Rule 5.2(a))



WO 2016/008976 A1

(54) Title: **ONCOLYTIC VIRUS FOR EXPRESSION OF IMMUNE CHECKPOINT MODULATORS**

(57) Abstract: The present invention provides an oncolytic virus comprising nucleotide sequence(s) encoding one or more immune checkpoint modulator(s). It also concerns a pharmaceutical composition comprising effective amount of said oncolytic virus and, eventually, a pharmaceutically acceptable vehicle and its use for treating proliferative diseases such as cancers.

Oncolytic virus for expression of immune checkpoint modulators

Field of the invention

5 The present invention generally relates to the field of oncolytic virotherapy and more specifically to compositions and methods to treat, prevent, or inhibit proliferative diseases, especially cancer. Embodiments include an oncolytic virus comprising nucleotide sequence(s) encoding one or more immune checkpoint modulator(s). Embodiments also include a pharmaceutical composition comprising such oncolytic virus and, eventually, a pharmaceutically acceptable vehicle and its use for treating proliferative diseases such as cancers.

Cancer is caused by both external factors (e.g. tobacco, infectious organisms, alimentary habits, chemicals, and radiation) and internal factors (e.g. inherited mutations, hormones, immune conditions, and mutations that occur from metabolism). Each year, cancer is diagnosed in more than 12 million subjects worldwide. In industrialized countries, approximately one person out five 15 will die of cancer. Although a vast number of chemotherapeutics exist, they are often ineffective, especially against malignant and metastatic tumors that establish at a very early stage of the disease. Moreover, antitumor immunity is often ineffective due to the fact that tumor cells have evolved mechanisms to escape host defense. One of the major mechanisms of immune suppression is a process known as “T-cell exhaustion”, which results from chronic exposure to antigens and is 20 characterized by the upregulation of inhibitory receptors. These inhibitory receptors serve as immune checkpoints in order to prevent uncontrolled immune reactions. Various immune checkpoints acting at different levels of T cell immunity have been described in the literature, including programmed cell death protein 1 (PD-1) and its ligands PD-L1 and PD-L2, CTLA-4 (cytotoxic T-lymphocyte associated protein-4), LAG3, B and T lymphocyte attenuator, T-cell immunoglobulin, 25 mucin domain-containing protein 3 (TIM-3), and V-domain immunoglobulin suppressor of T cell activation.

Whatever the mechanism of action, these immune checkpoints can inhibit the development of an efficient anti-tumor immune response. There is increasing interest in the possible therapeutic benefits of blocking such immune checkpoints as a means of inhibiting immune system tolerance to 30 tumors and thus rescue exhausted antitumor T cells (Leach et al., 1996, Science 271: 1734-6). A vast number of antagonistic antibodies have been developed during the last decade (e.g. anti Tim3, -PD-L1, -CTLA-4, -PD1, etc) and most importantly, some have been associated with objective clinical responses in cancer patients. Antibodies targeting CTLA-4 are already marketed (e.g. Ipilimumab, Yervoy, Bristol-Myers Squibb) for metastatic melanoma. BMS reported that from 1800 melanoma

patients treated with ipilimumab 22% are still alive 3 years later. Antibody therapies with anti PD-L1 (e.g. MPDL3280A, Roche), anti PD-1 (e.g. Nivolumab, BMS) are also ongoing.

Another therapeutic approach that is emerging in the field of cancer is oncolytic viruses 5 (Hermiston, 2006, Curr. Opin. Mol. Ther. 8: 322-30). Oncolytic viruses are capable of selective replication in dividing cells (e.g. cancer cell) while leaving non dividing cells (e.g. normal cells) unharmed. As the infected dividing cells are destroyed by lysis, they release new infectious virus particles to infect the surrounding dividing cells. Cancer cells are ideal hosts for many viruses because they have the antiviral interferon pathway inactivated or have mutated tumour suppressor 10 genes that enable viral replication to proceed unhindered (Chernajovsky et al., 2006, British Med. J. 332: 170-2). A number of viruses including adenovirus, reovirus, measles, herpes simplex, Newcastle disease virus and vaccinia have now been clinically tested as oncolytic agents.

Some viruses are naturally oncolytic (such as reovirus and the Seneca valley picornavirus) while others are engineered for tumor selectivity by modifying the viral genome. Such modifications 15 include functional deletions in essential viral genes, the use of tumor- or tissue-specific promoters to control the viral gene expression and tropism modification to redirect virus to the cancer cell surface.

The first oncolytic virus to be approved by a regulatory agency was a genetically modified adenovirus named H101 (Shanghai Sunway Biotech) that gained approval in 2005 from China's State 20 Food and Drug Administration (SFDA) for the treatment of head and neck cancer. Another oncolytic adenovirus, named ONYX-015 is in ongoing clinical trials for the treatment of various solid tumors (in phase III for the treatment of recurrent head and neck cancer) (Cohen et al., 2001, Curr. Opin. Investig. Drugs 2: 1770-5). As another example, oncolytic herpes simplex 1 (T-VEC) was genetically engineered to attenuate the virus virulence, increase selectivity for cancer cells and enhance 25 antitumor immune response (through GM-CSF expression). Clinical efficacy in unresectable melanoma has been demonstrated in Phase II and Phase III clinical trials (Senzer et al, 2009, J. Clin. Oncol. 27: 5763-71).

Vaccinia viruses (VV) possess many of the key attributes necessary for use in oncolytic virotherapy such as natural tropism for tumors, strong lytic ability, short life cycle with rapid cell-to- 30 cell spread, highly efficient gene expression and a large cloning capacity. In addition, they have been delivered to millions of individuals during the smallpox eradication campaign without major safety concerns. In this respect, a TK and VGF double deleted VV (Wyeth strain) expressing GM-CSF (named JX-963) showed significant cancer selectivity in tumor bearing mice (Thorne et al., 2007, J. Clin Invest. 117: 3350-8). On the same line, JX-594, a TK-deleted VV (Wyeth strain) armed with GM-

CSF, has shown promising clinical data, and a randomized Phase III trial in hepatocellular carcinoma is expected to start soon.

Combination therapies involving oncolytic virus and immune checkpoint inhibitors have been described in the literature. WO2014/022138 describes the combination of irradiated tumor 5 cells, an oncolytic adenovirus and an anti CTLA4 antibody for use for treating bladder or prostate cancer. WO2014/047350 envisages a recombinant oncolytic virus with a gene encoding an anti-PD-1 antibody inserted in the viral genome without providing any working example that would support utility of such an oncolytic virus.

10 **Technical problem**

One may expect that cancer will continue to be a serious global health threat for many years due to the high number of causal factors that may act together or separately to initiate or promote the development of a cancer. Moreover, malignant and especially metastatic tumors are often resistant to conventional therapies explaining the significant morbidity of some cancers.

15 Thus, there is an important need to develop more effective approaches, for improving prevention and treatment of such proliferative diseases, and especially metastatic cancers. The present invention provides a unique product combining oncolysis for killing dividing cells and immune checkpoint for breaking cancer-associated immune tolerance.

20 This technical problem is solved by the provision of the embodiments as defined in the claims.

Other and further aspects, features and advantages of the present invention will be apparent from the following description of the presently preferred embodiments of the invention. These embodiments are given for the purpose of disclosure.

25

Summary of the invention

The present invention concerns an oncolytic virus comprising inserted in its genome one or more nucleic acid molecule(s) encoding one or more immune checkpoint modulator(s).

30 The oncolytic virus is preferably selected from the group consisting of reovirus, New Castle Disease virus (NDV), vesicular stomatitis virus (VSV), measles virus, influenza virus, Sinbis virus, adenovirus, poxvirus and herpes virus (HSV) and the like. In one embodiment, the oncolytic virus is a vaccinia virus. In a preferred embodiment, the vaccinia virus is engineered to lack thymidine kinase activity (e.g. the genome of said VV has an inactivating mutation in J2R gene to produce a defective TK phenotype). Alternatively or in combination, the vaccinia virus is engineered to lack RR

activity (e.g. the genome of said VV has an inactivating mutation in I4L and/or F4L gene to produce a defective RR phenotype).

In one embodiment, the vaccinia virus further expresses at least one therapeutic gene, in particular a gene encoding a suicide gene product and/or an immunostimulatory protein.

5 In one embodiment, the encoded one or more immune checkpoint modulator(s) is an antagonist molecule that antagonizes the activity of PD-1, PD-L1 or CTLA4 with a specific preference for an anti PD-1 antibody and/or an anti CTLA4 antibody.

The present invention further provides a composition comprising said oncolytic virus, eventually with a pharmaceutical acceptable vehicle. In one embodiment, the composition is 10 formulated for intravenous or intratumoral administration.

The present invention also concerns the use of said oncolytic virus or composition thereof for treating a proliferative disease as well as a method of treatment relying on the administration of an effective amount of said oncolytic virus or composition thereof. In one embodiment, the proliferative disease treated by the method of the invention is cancer and especially melanoma, 15 renal cancer, prostate cancer, breast cancer, colorectal cancer, lung cancer and liver cancer. In one embodiment, the use or method comprises an additional step in which a pharmaceutically acceptable amount of a prodrug is administered to said mammal. The administration of said prodrug takes place preferably at least 3 days after the administration of said oncolytic virus or virus composition.

20

Detailed description

The present invention concerns an oncolytic virus comprising inserted in its genome one or more nucleic acid molecule(s) encoding one or more immune checkpoint modulator(s).

25

Definitions

As used throughout the entire application, the terms "a" and "an" are used in the sense that they mean "at least one", "at least a first", "one or more" or "a plurality" of the referenced components or steps, unless the context clearly dictates otherwise. For example, the term "a cell" 30 includes a plurality of cells, including mixtures thereof.

The term "one or more" refers to either one or a number above one (e.g. 2, 3, 4, 5, etc).

The term "and/or" wherever used herein includes the meaning of "and", "or" and "all or any other combination of the elements connected by said term".

The term "about" or "approximately" as used herein means within 20%, preferably within 10%, and more preferably within 5% of a given value or range.

As used herein, when used to define products, compositions and methods, the term "comprising" (and any form of comprising, such as "comprise" and "comprises"), "having" (and any 5 form of having, such as "have" and "has"), "including" (and any form of including, such as "includes" and "include") or "containing" (and any form of containing, such as "contains" and "contain") are open-ended and do not exclude additional, unrecited elements or method steps. Thus, a polypeptide "comprises" an amino acid sequence when the amino acid sequence might be part of the final amino acid sequence of the polypeptide. Such a polypeptide can have up to several 10 hundred additional amino acids residues. "Consisting essentially of" means excluding other components or steps of any essential significance. Thus, a composition consisting essentially of the recited components would not exclude trace contaminants and pharmaceutically acceptable carriers. A polypeptide "consists essentially of" an amino acid sequence when such an amino acid sequence is present with eventually only a few additional amino acid residues. "Consisting of" 15 means excluding more than trace elements of other components or steps. For example, a polypeptide "consists of" an amino acid sequence when the polypeptide does not contain any amino acids but the recited amino acid sequence.

The terms "polypeptide", "peptide" and "protein" refer to polymers of amino acid residues which comprise at least nine or more amino acids bonded via peptide bonds. The polymer can be 20 linear, branched or cyclic and may comprise naturally occurring and/or amino acid analogs and it may be interrupted by non-amino acids. As a general indication, if the amino acid polymer is more than 50 amino acid residues, it is preferably referred to as a polypeptide or a protein whereas if it is 50 amino acids long or less, it is referred to as a "peptide".

Within the context of the present invention, the terms "nucleic acid", "nucleic acid 25 molecule", "polynucleotide" and "nucleotide sequence" are used interchangeably and define a polymer of any length of either polydeoxyribonucleotides (DNA) (e.g. cDNA, genomic DNA, plasmids, vectors, viral genomes, isolated DNA, probes, primers and any mixture thereof) or polyribonucleotides (RNA) (e.g. mRNA, antisense RNA, SiRNA) or mixed polyribo-polydeoxyribonucleotides. They encompass single or double-stranded, linear or circular, natural or 30 synthetic, modified or unmodified polynucleotides. Moreover, a polynucleotide may comprise non-naturally occurring nucleotides and may be interrupted by non-nucleotide components.

The term "analog" or "variant" as used herein refers to a molecule (polypeptide or nucleic acid) exhibiting one or more modification(s) with respect to the native counterpart. Any modification(s) can be envisaged, including substitution, insertion and/or deletion of one or more

nucleotide/amino acid residue(s). Preferred are analogs that retain a degree of sequence identity of at least 80%, preferably at least 85%, more preferably at least 90%, and even more preferably at least 98% identity with the sequence of the native counterpart.

In a general manner, the term "identity" refers to an amino acid to amino acid or nucleotide 5 to nucleotide correspondence between two polypeptide or nucleic acid sequences. The percentage of identity between two sequences is a function of the number of identical positions shared by the sequences, taking into account the number of gaps which need to be introduced for optimal alignment and the length of each gap. Various computer programs and mathematical algorithms are available in the art to determine the percentage of identity between amino acid sequences, 10 such as for example the Blast program available at NCBI or ALIGN in Atlas of Protein Sequence and Structure (Dayhoffed, 1981, Suppl., 3: 482-9). Programs for determining identity between nucleotide sequences are also available in specialized data base (e.g. Genbank, the Wisconsin Sequence Analysis Package, BESTFIT, FASTA and GAP programs). For illustrative purposes, "at least 80% identity" means 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 15 94%, 95%, 96%, 97%, 98%, 99% or 100%.

As used herein, the term "isolated" refers to a protein, polypeptide, peptide, polynucleotide, vector, etc., that is removed from its natural environment (i.e. separated from at least one other component(s) with which it is naturally associated or found in nature). For example, a nucleotide sequence is isolated when it is separated of sequences normally associated with it in 20 nature (e.g. dissociated from a genome) but it can be associated with heterologous sequences.

The term "obtained from", "originating" or "originate" is used to identify the original source of a component (e.g. polypeptide, nucleic acid molecule) but is not meant to limit the method by which the component is made which can be, for example, by chemical synthesis or recombinant means.

25 As used herein, the term "host cell" should be understood broadly without any limitation concerning particular organization in tissue, organ, or isolated cells. Such cells may be of a unique type of cells or a group of different types of cells such as cultured cell lines, primary cells and dividing cells. In the context of the invention, the term "host cells" include prokaryotic cells, lower eukaryotic cells such as yeast, and other eukaryotic cells such as insect cells, plant and mammalian 30 (e.g. human or non-human) cells as well as cells capable of producing the oncolytic virus and/or the immune checkpoint modulator(s) for use in the invention. This term also includes cells which can be or has been the recipient of the vectors described herein as well as progeny of such cells.

As used herein, the term "oncolytic virus" refers to a virus capable of selectively replicating in dividing cells (e.g. a proliferative cell such as a cancer cell) with the aim of slowing the growth

and/or lysing said dividing cell, either in vitro or in vivo, while showing no or minimal replication in non-dividing cells. Typically, an oncolytic virus contains a viral genome packaged into a viral particle (or virion) and is infectious (i.e. capable of infecting and entering into a host cell or subject). As used herein, this term encompasses DNA or RNA vector (depending on the virus in question) as well as 5 viral particles generated thereof.

The term “treatment” (and any form of treatment such as “treating”, “treat”) as used herein encompasses prophylaxis (e.g. preventive measure in a subject at risk of having the pathological condition to be treated) and/or therapy (e.g. in a subject diagnosed as having the pathological condition), eventually in association with conventional therapeutic modalities. The 10 result of the treatment is to slow down, cure, ameliorate or control the progression of the targeted pathological condition. For example, a subject is successfully treated for a cancer if after administration of an oncolytic virus as described herein, the subject shows an observable improvement of its clinical status.

The term “administering” (or any form of administration such as “administered”) as used 15 herein refers to the delivery to a subject of a therapeutic agent such as the oncolytic virus described herein.

As used herein, the term “proliferative disease” encompasses any disease or condition resulting from uncontrolled cell growth and spread including cancers as well as diseases associated to an increased osteoclast activity (e.g. rheumatoid arthritis, osteoporosis, etc) and cardiovascular 20 diseases (restenosis that results from the proliferation of the smooth muscle cells of the blood vessel wall, etc). The term “cancer” may be used interchangeably with any of the terms “tumor”, “malignancy”, “neoplasm”, etc. These terms are meant to include any type of tissue, organ or cell, any stage of malignancy (e.g. from a prelesion to stage IV)

The term “subject” generally refers to an organism for whom any product and method of 25 the invention is needed or may be beneficial. Typically, the organism is a mammal, particularly a mammal selected from the group consisting of domestic animals, farm animals, sport animals, and primates. Preferably, the subject is a human who has been diagnosed as having or at risk of having a proliferative disease such as a cancer. The terms “subject” and “patients” may be used interchangeably when referring to a human organism and encompasses male and female. The 30 subject to be treated may be a newborn, an infant, a young adult or an adult.

The term “combination” or “association” as used herein refers to any arrangement possible of various components (e.g. an oncolytic virus and one or more substance effective in anticancer therapy). Such an arrangement includes mixture of said components as well as separate combinations for concomitant or sequential administrations. The present invention encompasses

combinations comprising equal molar concentrations of each component as well as combinations with very different concentrations. It is appreciated that optimal concentration of each component of the combination can be determined by the artisan skilled in the art.

The term "immune checkpoint modulator" refers to a molecule capable of modulating the 5 function of an immune checkpoint protein in a positive or negative way (in particular the interaction between an antigen presenting cell (APC) or a cancer cell and a T effector cell). The term "immune checkpoint" refers to a protein directly or indirectly involved in an immune pathway that under normal physiological conditions is crucial for preventing uncontrolled immune reactions and thus for the maintenance of self-tolerance and/or tissue protection. The one or more immune 10 checkpoint modulator(s) in use herein may independently act at any step of the T cell-mediated immunity including clonal selection of antigen-specific cells, T cell activation, proliferation, trafficking to sites of antigen and inflammation, execution of direct effector function and signaling through cytokines and membrane ligands. Each of these steps is regulated by counterbalancing stimulatory and inhibitory signals that fine tune the response. In the context of the present 15 invention, the term encompasses immune checkpoint modulator(s) capable of down-regulating at least partially the function of an inhibitory immune checkpoint (antagonist) and/or immune checkpoint modulator(s) capable of up-regulating at least partially the function of a stimulatory immune checkpoint (agonist).

20 Oncolytic virus

The oncolytic virus of the present invention can be obtained from any member of virus identified at present time provided that it is oncolytic by its propensity to selectively replicate and kill dividing cells as compared to non-dividing cells. It may be a native virus that is naturally oncolytic or may be engineered by modifying one or more viral genes so as to increase tumor 25 selectivity and/or preferential replication in dividing cells, such as those involved in DNA replication, nucleic acid metabolism, host tropism, surface attachment, virulence, lysis and spread (see for example Kirn et al., 2001, Nat. Med. 7: 781; Wong et al., 2010, Viruses 2: 78-106). One may also envisage placing one or more viral gene(s) under the control of event or tissue-specific regulatory elements (e.g. promoter).

30 Exemplary oncolytic viruses include without limitation reovirus, Seneca Valley virus (SVV), vesicular stomatitis virus (VSV), Newcastle disease virus (NDV), herpes simplex virus (HSV), morbillivirus virus, retrovirus, influenza virus, Sinbis virus, poxvirus, adenovirus, or the like.

In one embodiment, the oncolytic virus of the present invention is obtained from a reovirus. A representative example includes Reolysin (under development by Oncotherapy Biotech; NCT01166542).

In one embodiment, the oncolytic virus of the present invention is obtained from a Seneca 5 Valley virus. A representative example includes NTX-010 (Rudin et al., 2011, Clin. Cancer. Res. 17(4): 888-95).

In one embodiment, the oncolytic virus of the present invention is obtained from a vesicular stomatitis virus (VSV). Representative examples are described in the literature (e.g. Stojdl et al., 2000, Nat. Med. 6(7): 821-5; Stojdl et al., 2003, Cancer Cell 4(4): 263-75).

10 In one embodiment, the oncolytic virus of the present invention is obtained from a Newcastle disease virus. Representative examples include without limitation the 73-T PV701 and HDV-HUJ strains as well as those described in the literature (e.g. Phuangsab et al., 2001, Cancer Lett. 172(1): 27-36; Lorence et al., 2007, Curr. Cancer Drug Targets 7(2): 157-67; Freeman et al., 2006, Mol. Ther. 13(1): 221-8).

15 In one embodiment, the oncolytic virus of the present invention is obtained from a herpes virus. The Herpesviridae are a large family of DNA viruses that all share a common structure and are composed of relatively large double-stranded, linear DNA genomes encoding 100-200 genes encapsulated within an icosahedral capsid which is enveloped in a lipid bilayer membrane. Although the oncolytic herpes virus can be derived from different types of HSV, particularly preferred are 20 HSV1 and HSV2. The herpes virus may be genetically modified so as to restrict viral replication in tumors or reduce its cytotoxicity in non-dividing cells. For example, any viral gene involved in nucleic acid metabolism may be inactivated, such as thymidine kinase (Martuza et al., 1991, Science 252: 854-6), ribonucleotide reductase (RR) (Boviatsis et al., Gene Ther. 1: 323-31; Mineta et al., 1994, Cancer Res. 54: 3363-66), or uracil-N-glycosylase (Pyles et al., 1994, J. Virol. 68: 4963-72).

25 Another aspect involves viral mutants with defects in the function of genes encoding virulence factors such as the ICP34.5 gene (Chambers et al., 1995, Proc. Natl. Acad. Sci. USA 92: 1411-5). Representative examples of oncolytic herpes virus include NV1020 (e.g. Geevarghese et al., 2010, Hum. Gene Ther. 21(9): 1119-28) and T-VEC (Andtbacka et al., 2013, J. Clin. Oncol. 31, abstract number LBA9008).

30 In one embodiment, the oncolytic virus of the present invention is obtained from a morbillivirus which can be obtained from the paramyxoviridae family, with a specific preference for measles virus. Representative examples of oncolytic measles viruses include without limitation MV-Edm (McDonald et al., 2006; Breast Cancer Treat. 99(2): 177-84) and HMWMAA (Kaufmann et al., 2013, J. Invest. Dermatol. 133(4): 1034-42)

In one embodiment, the oncolytic virus of the present invention is obtained from an adenovirus. Methods are available in the art to engineer oncolytic adenoviruses. An advantageous strategy includes the replacement of viral promoters with tumor-selective promoters or modifications of the E1 adenoviral gene product(s) to inactivate its/their binding function with p53 or retinoblastoma (Rb) protein that are altered in tumor cells. In the natural context, the adenovirus E1B55kDa gene cooperates with another adenoviral product to inactivate p53 (p53 is frequently dysregulated in cancer cells), thus preventing apoptosis. Representative examples of oncolytic adenovirus include ONYX-015 (e.g. Khuri et al., 2000, Nat. Med 6(8): 879-85) and H101 also named Oncorine (Xia et al., 2004, Ai Zheng 23(12): 1666-70).

10 In one embodiment, the oncolytic virus of the present invention is a poxvirus. As used herein the term "poxvirus" refers to a virus belonging to the Poxviridae family, with a specific preference for a poxvirus belonging to the Chordopoxviridae subfamily and more preferably to the Orthopoxvirus genus. Sequences of the genome of various poxviruses, for example, the vaccinia virus, cowpox virus, Canarypox virus, Ectromelia virus, Myxoma virus genomes are available in the 15 art and specialized databases such as Genbank (accession number NC_006998, NC_003663, NC_005309, NC_004105, NC_001132 respectively).

Desirably, the oncolytic poxvirus is an oncolytic vaccinia virus. Vaccinia viruses are members of the poxvirus family characterized by a 200kb double-stranded DNA genome that encodes numerous viral enzymes and factors that enable the virus to replicate independently from the host 20 cell machinery. The majority of vaccinia virus particles is intracellular (IMV for intracellular mature virion) with a single lipid envelop and remains in the cytosol of infected cells until lysis. The other infectious form is a double enveloped particle (EEV for extracellular enveloped virion) that buds out from the infected cell without lysing it.

Although it can derive from any vaccinia virus strain, Elstree, Wyeth, Copenhagen and 25 Western Reserve strains are particularly preferred. The gene nomenclature used herein is that of Copenhagen vaccinia strain. It is also used herein for the homologous genes of other poxviridae unless otherwise indicated. However, gene nomenclature may be different according to the pox strain but correspondence between Copenhagen and other vaccinia strains are generally available in the literature.

30 Preferably, the oncolytic vaccinia virus of the present invention is modified by altering for one or more viral gene(s). Said modification(s) preferably lead(s) to the synthesis of a defective protein unable to ensure the activity of the protein produced under normal conditions by the unmodified gene (or lack of synthesis). Modifications encompass deletion, mutation and/or substitution of one or more nucleotide(s) (contiguous or not) within the viral gene or its regulatory

elements. Modification(s) can be made in a number of ways known to those skilled in the art using conventional recombinant techniques. Exemplary modifications are disclosed in the literature with a specific preference for those altering viral genes involved in DNA metabolism, host virulence, IFN pathway (see e.g. Guse et al., 2011, Expert Opinion Biol. Ther.11(5): 595-608) and the like.

5 More preferably, the oncolytic poxvirus of the present invention is modified by altering the thymidine kinase-encoding gene (locus J2R). The TK enzyme is involved in the synthesis of deoxyribonucleotides. TK is needed for viral replication in normal cells as these cells have generally low concentration of nucleotides whereas it is dispensable in dividing cells which contain high nucleotide concentration.

10 Alternatively or in combination, the oncolytic poxvirus of the present invention is modified by altering at least one gene or both genes encoding Ribonucleotide reductase (RR). In the natural context, this enzyme catalyzes the reduction of ribonucleotides to deoxyribonucleotides that represents a crucial step in DNA biosynthesis. The viral enzyme is similar in subunit structure to the mammalian enzyme, being composed of two heterologous subunits, designed R1 and R2 encoded 15 respectively by the I4L and F4L locus. Sequences for the I4L and F4L genes and their locations in the genome of various poxvirus are available in public databases, for example via accession number DQ437594, DQ437593, DQ377804, AH015635, AY313847, AY313848, NC_003391, NC_003389, NC_003310, M-35027, AY243312, DQ011157, DQ011156, DQ011155, DQ011154, DQ011153, Y16780, X71982, AF438165, U60315, AF410153, AF380138, U86916, L22579, NC_006998, 20 DQ121394 and NC_008291. In the context of the invention, either the I4L gene (encoding the R1 large subunit) or the F4L gene (encoding the R2 small subunit) or both may be inactivated.

Alternatively or in combination, other strategies may also be pursued to further increase the virus tumor-specificity. A representative example of suitable modification includes disruption of the VGF-encoding gene from the viral genome. VGF (for VV growth factor) is a secreted protein 25 which is expressed early after cell infection and its function seems important for virus spread in normal cells. Another example is the disruption of the A56R gene coding for hemagglutinin, eventually in combination with TK deletion (Zhang et al., 2007, Cancer Res. 67: 10038-46). Disruption of interferon modulating gene(s) may also be advantageous (e.g. the B8R or B18R gene) or the caspase-1 inhibitor B13R gene. Another suitable modification comprises the disruption of the 30 F2L gene which encodes the viral dUTPase involved in both maintaining the fidelity of DNA replication and providing the precursor for the production of TMP by thymidylate synthase (Broyles et al., 1993, Virol. 195: 863-5). Sequence of the vaccinia virus F2L gene is available in genbank via accession number M25392.

In a preferred embodiment, the oncolytic virus of this invention is a vaccinia virus defective for TK resulting from inactivating mutations in the J2R gene. In another preferred embodiment, the oncolytic virus of this invention is a vaccinia virus defective for both TK and RR activities resulting from inactivating mutations in both the J2R gene and the I4L and/or F4L gene(s) carried by the viral 5 genome (e.g. as described in WO2009/065546 and Foloppe et al., 2008, Gene Ther., 15: 1361-71). In another preferred embodiment, the oncolytic virus of this invention is a vaccinia virus defective for dUTPase resulting from inactivating mutations in the F2L gene (e.g. as described in WO2009/065547), eventually in combination with disruption of at least one of TK and RR activities or both (resulting in a virus with inactivating mutations in the F2L; F2L and J2R gene; F2L and I4L; or 10 F2L, J2R and I4L).

Therapeutic genes

In one embodiment, the oncolytic virus of this invention further expresses at least one therapeutic gene inserted in the viral genome. A "therapeutic gene" encodes a product capable of 15 providing a biological activity when administered appropriately to a subject, which is expected to cause a beneficial effect on the course or a symptom of the pathological condition to be treated by either potentiating anti-tumor efficacy or reinforcing the oncolytic nature of the virus. In the context of the invention, the therapeutic gene can be of mammal origin (e.g. human, murine, rabbit, etc) or not (e.g. of bacterial, yeast or viral origin).

20 A vast number of therapeutic genes may be envisaged in the context of the invention such as those encoding polypeptides that can compensate for defective or deficient proteins in the subject, or those that act through toxic effects to limit or remove harmful cells from the body or those that encode immunity conferring polypeptides. They may be native genes or genes obtained from the latter by mutation, deletion, substitution and/or addition of one or more nucleotides.

25 Advantageously, the oncolytic virus of the present invention carries a therapeutic gene selected from the group consisting of genes encoding suicide gene products and immunostimulatory proteins.

Suicide gene

30 The term "suicide gene" refers to a gene coding for a protein able to convert a precursor of a drug into a cytotoxic compound. Suicide genes comprise but are not limited to genes coding protein having a cytosine deaminase activity, a thymidine kinase activity, an uracil phosphoribosyl transferase activity, a purine nucleoside phosphorylase activity and a thymidylate kinase activity.

Examples of suicide genes and corresponding precursors of a drug comprising one nucleobase moiety are disclosed in the following table

Table 1

Suicide gene	prodrug
Thymidine Kinase	Ganciclovir; Ganciclovir elaidic acid ester; penciclovir; Acyclovir; Valacyclovir; (E)-5-(2-bromovinyl)-2'-deoxyuridine; zidovudine; 2'-Exo-methanocarbathymidine
Cytosine deaminase	5-Fluorocytosine
Purine nucleoside phosphorylase	6-Methylpurine deoxyriboside; Fludarabine
uracil phosphoribosyl transferase	5-Fluorocytosine; 5-Fluorouracil
thymidylate kinase.	Azidothymidine

5

Desirably, the suicide gene encodes a protein having at least a CDase activity. In the prokaryotes and lower eukaryotes (it is not present in mammals), CDase is involved in the pyrimidine metabolic pathway by which exogenous cytosine is transformed into uracil by means of a hydrolytic deamination. CDase also deaminates an analogue of cytosine, i.e. 5-fluorocytosine (5-FC), thereby forming 5-fluorouracil (5-FU), a compound which is highly cytotoxic when it is converted into 5-fluoro-UMP (5-FUMP). CDase encoding nucleic acid molecule can be obtained from any prokaryotes and lower eukaryotes such as *Saccharomyces cerevisiae* (FCY1 gene), *Candida Albicans* (FCA1 gene) and *Escherichia coli* (codA gene). The gene sequences and encoded CDase proteins have been published and are available in specialized data banks (SWISSPROT EMBL, 10 Genbank, Medline and the like). Functional analogues of these genes may also be used. Such analogues preferably have a nucleic acid sequence having a degree of identity of at least 70%, advantageously of at least 80%, preferably of at least 90%, and most preferably of at least 95% with 15 the nucleic acid sequence of the native gene.

Alternatively or in combination, the oncolytic virus of the invention carries in its viral 20 genome a suicide gene encoding a polypeptide having uracil phosphoribosyl transferase (UPRTase) activity. In prokaryotes and lower eukaryotes, uracil is transformed into UMP by the action of UPRTase. This enzyme converts 5-FU into 5-FUMP. By way of illustration, the nucleic acid sequences encoding the UPRTases from *E. coli* (Andersen et al., 1992, European J. Biochem. 204: 51-56), from *Lactococcus lactis* (Martinussen et al., 1994, J. Bacteriol. 176: 6457-63), from *Mycobacterium bovis*

(Kim et al., 1997, *Biochem. Mol. Biol. Internat.* 41: 1117-24) and from *Bacillus subtilis* (Martinussen et al., 1995, *J. Bacteriol.* 177: 271-4) may be used in the context of the invention. However, it is most particularly preferred to use a yeast UPRTase and in particular that encoded by the *S. cerevisiae* (FUR1 gene) whose sequence is disclosed in Kern et al. (1990, *Gene* 88: 149-57).

5 Functional UPRTase analogues may also be used such as the N-terminally truncated FUR1 mutant described in EP998568 (with a deletion of the 35 first residues up to the second Met residue present at position 36 in the native protein) which exhibits a higher UPRTase activity than that of the native enzyme.

Preferably, the suicide gene inserted in the viral genome of the oncolytic virus of the 10 present invention encodes a polypeptide having CDase and UPRTase activities. Such a polypeptide can be engineered by fusion of two enzymatic domains, one having the CDase activity and the second having the UPRTase activity. Exemplary polypeptides include without limitation fusion polypeptides codA::upp, FCY1::FUR1 and FCY1::FUR1[Delta] 105 (FCU1) and FCU1-8 described in WO96/16183, EP998568 and WO2005/07857. Of particular interest is the FCU1 suicide gene (or 15 FCY1::FUR1[Delta] 105 fusion) encoding a polypeptide comprising the amino acid sequence represented in the sequence identifier SEQ ID NO: 1 of WO2009/065546. The present invention encompasses analogs of such polypeptides providing they retain the CDase, and/or UPRTase activities. It is within the reach of the skilled person to isolate the CDase and/or UPRTase -encoding nucleic acid molecules from the published data, eventually engineer analogs thereof and test the 20 enzymatic activity in an acellular or cellular system according to conventional techniques (see e.g. EP998568).

Immunostimulatory therapeutic genes

As used herein, the term “immunostimulatory protein” refers to a protein which has the 25 ability to stimulate the immune system, in a specific or non-specific way. A vast number of proteins are known in the art for their ability to exert an immunostimulatory effect. Examples of suitable immunostimulatory proteins in the context of the invention include without limitation cytokines, with a specific preference for interleukins (e.g. IL-2, IL-6, IL-12, IL-15, IL-24), chemokines (e.g. CXCL10, CXCL9, CXCL11), interferons (e.g. IFNg, IFNAlpha), tumor necrosis factor (TNF), colony- 30 stimulating factors (e.g. GM-CSF, C-CSF, M-CSF...), APC (for Antigen Presenting Cell)-exposed proteins (e.g. B7.1, B7.2 and the like), growth factors (Transforming Growth Factor TGF, Fibroblast Growth Factor FGF, Vascular Endothelial Growth Factors VEGF, and the like), MHC antigens of class I or II, apoptosis inducers or inhibitors (e.g. Bax, Bcl2, BclX...), cytostatic agents (p21, p16, Rb...), immunotoxins, antigenic polypeptides (antigenic polypeptides, epitopes, and the like) and markers

(beta-galactosidase, luciferase....). Preferably, the immunostimulatory protein is an interleukin or a colony-stimulating factor, with a specific preference for GM-CSF.

Immune checkpoint modulator(s)

5 Immune checkpoints and modulators thereof as well as methods of using such compounds are described in the literature. In accordance with this invention, the one or more immune checkpoint modulator(s) may independently be a polypeptide comprising a domain capable of binding the targeted immune checkpoint and/or inhibiting the binding of a ligand to said targeted immune checkpoint so as to exert an antagonist function (i.e. being capable of antagonizing an 10 immune checkpoint-mediated inhibitory signal) or an agonist function (i.e. being capable of boosting an immune checkpoint-mediated stimulatory signal). Such one or more immune checkpoint modulator(s) can be independently selected from the group consisting of peptides (e.g. peptide ligands), soluble domains of natural receptors, RNAi, antisense molecules, antibodies and protein scaffolds.

15 In a preferred embodiment, the immune checkpoint modulator is an antibody. In the context of the invention, "antibody" ("Ab") is used in the broadest sense and encompasses naturally occurring and engineered by man as well as full length antibodies or functional fragments or analogs thereof that are capable of binding the target immune checkpoint or epitope (thus retaining the target-binding portion). The antibody encoded by the oncolytic virus of the invention 20 can be of any origin, e.g. human, humanized, animal (e.g. rodent or camelid antibody) or chimeric. It may be of any isotype (e.g. IgG1, IgG2, IgG3, IgG4, IgM isotype, etc). In addition, it may be glycosylated, partially glycosylated or non-glycosylated (e.g. by mutating one or more residue(s) within the site(s) of glycosylation). The term antibody also includes bispecific or multispecific antibodies so long as they exhibit the binding specificity described herein.

25 For illustrative purposes, full length antibodies are glycoproteins comprising at least two heavy (H) chains and two light (L) chains inter-connected by disulfide bonds. Each heavy chain is comprised of a heavy chain variable region (VH) and a heavy chain constant region which is made of three CH1, CH2 and CH3 domains (eventually with a hinge between CH1 and CH2). Each light chain is comprised of a light chain variable region (VL) and a light chain constant region which comprises 30 one CL domain. The VH and VL regions comprise hypervariable regions, named complementarity determining regions (CDR), interspersed with more conserved regions named framework regions (FR). Each VH and VL is composed of three CDRs and four FRs in the following order: FR1-CDR1-FR2-CDR2-FR3-CDR3-FR4. The CDR regions of the heavy and light chains are determinant for the binding specificity.

As used herein, a "humanized antibody" refers to a non-human (e.g. murine, camel, rat, etc) antibody whose protein sequence has been modified to increase its similarity to a human antibody (i.e. produced naturally in humans). The process of humanization is well known in the art (see e.g. Presta et al., 1997, *Cancer Res.* 57(20): 4593-9; US 5,225,539; US 5,530,101; US 6,180,370; 5 WO2012/110360). For example, the immune checkpoint antibody for use herein can be humanized by substituting one or more residue of the FR regions to look like human immunoglobulin sequence whereas the vast majority of the residues of the variable regions (especially the CDRs) are not modified and correspond to those of a non-human immunoglobulin. For general guidance, the number of these amino acid substitutions in the FR regions is typically no more than 20 in each 10 variable region VH or VL.

As used herein, a "chimeric antibody" refers to an antibody comprising one or more element(s) of one species and one or more element(s) of another species, for example, a non-human antibody comprising at least a portion of a constant region (Fc) of a human immunoglobulin.

Many forms of antibody can be expressed by the oncolytic virus of the invention. 15 Representative examples include without limitation Fab, Fab', F(ab')2, dAb, Fd, Fv, scFv, di-scFv and diabody, etc.. More specifically:

- (i) a Fab fragment represented by a monovalent fragment consisting of the VL, VH, CL and CH1 domains;
- (ii) a F(ab')2 fragment represented by a bivalent fragment comprising two Fab 20 fragments linked by at least one disulfide bridge at the hinge region;
- (iii) a Fd fragment consisting of the VH and CH1 domains;
- (iv) a Fv fragment consisting of the VL and VH domains of a single arm of an antibody;
- (v) a dAb fragment consisting of a single domain variable fragment (VH or VL domain);
- (vi) a single chain Fv (scFv) comprising the two domains of a Fv fragment, VL and VH, 25 that are fused together, eventually with a linker to make a single protein chain (see e.g. Bird et al., 1988, *Science* 242: 423-6; Huston et al., 1988, *Proc. Natl. Acad. Sci. USA* 85: 5879-83; US 4,946,778; US 5,258,498); and
- (vii) any other artificial antibody.

If needed, such fragments and analogs may be screened for functionality in the same 30 manner as with intact antibodies (e.g. by standard ELISA assay).

In a preferred embodiment, at least one of the one or more immune checkpoint modulator(s) encoded by the oncolytic virus of the present invention is a monoclonal antibody, with a specific preference for a human (in which both the framework regions are derived from human

germline immunoglobulin sequences) or a humanized antibody according to well-known humanization process.

Desirably, the one or more immune checkpoint modulator(s) encoded by the oncolytic virus of the present invention antagonize(s) at least partially (e.g. more than 50%) the activity of 5 inhibitory immune checkpoint(s), in particular those mediated by any of the following PD-1, PD-L1, PD-L2, LAG3, Tim3, BTLA and CTLA4, with a specific preference for a monoclonal antibody that specifically binds to any of such target proteins. The term "specifically binds to" refers to the capacity to a binding specificity and affinity for a particular target or epitope even in the presence of a heterogeneous population of other proteins and biologics. Thus, under designated assay 10 conditions, the antibody in use in the invention binds preferentially to its target and does not bind in a significant amount to other components present in a test sample or subject. Preferably, such an antibody shows high affinity binding to its target with an equilibrium dissociation constant equal or below 1×10^{-6} M (e.g. at least 0.5×10^{-6} , 1×10^{-7} , 1×10^{-8} , 1×10^{-9} , 1×10^{-10} , etc). Alternatively, the encoded 15 one or more immune checkpoint modulator(s) exert(s) an agonist function in the sense that it is capable of stimulating or reinforcing stimulatory signals, in particular those mediated by CD28 with a specific preference for any of ICOS, CD137 (4-1BB), OX40, CD27, CD40, GITR immune checkpoints. Standard assays to evaluate the binding ability of the antibodies toward immune checkpoints are known in the art, including for example, ELISAs, Western blots, RIAs and flow cytometry. The 20 binding kinetics (e.g., binding affinity) of the antibodies also can be assessed by standard assays known in the art, such as by Biacore analysis.

In a preferred embodiment, at least one of the one or more encoded checkpoint modulator(s) is a human or a humanized antibody capable of antagonizing an immune checkpoint involved in T cell-mediated response. A preferred example of immune checkpoint modulator is represented by a modulator capable of antagonizing at least partially the protein Programmed 25 Death 1 (PD-1), and especially an antibody that specifically binds to human PD-1. PD-1 is part of the immunoglobulin (Ig) gene superfamily and a member of the CD28 family. It is a 55 kDa type 1 transmembrane protein expressed on antigen-experienced cells (e.g. activated B cells, T cells, and myeloid cells) (Agata et al., 1996, Int. Immunol. 8: 765-72; Okazaki et al., 2002, Curr. Opin. Immunol. 14: 391779-82; Bennett et al., 2003, J. Immunol 170: 711-8). In normal context, it acts by limiting 30 the activity of T cells at the time of inflammatory response, thereby protecting normal tissues from destruction (Topalian, 2012, Curr. Opin. Immunol. 24: 207-12). Two ligands have been identified for PD-1, respectively PD-L1 (programmed death ligand 1) and PD-L2 (programmed death ligand 2) (Freeman et al., 2000, J. Exp. Med. 192: 1027-34; Carter et al., 2002, Eur. J. Immunol. 32: 634-43). PD-L1 was identified in 20-50% of human cancers (Dong et al., 2002, Nat. Med. 8: 787-9). The

interaction between PD-1 and PD-L1 resulted in a decrease in tumor infiltrating lymphocytes, a decrease in T-cell receptor mediated proliferation, and immune evasion by the cancerous cells (Dong et al., 2003, J. Mol. Med. 81: 281-7; Blank et al., 2005, Cancer Immunol. Immunother. 54: 307-314). The complete nucleotide and amino acid PD-1 sequences can be found under GenBank 5 Accession No U64863 and NP_005009.2. A number of anti PD1 antibodies are available in the art (see e.g. those described in WO2004/004771; WO2004/056875; WO2006/121168; WO2008/156712; WO2009/014708; WO2009/114335; WO2013/043569; and WO2014/047350). Preferably, the oncolytic virus of the present invention encodes and expresses an anti PD-1 antibody that is FDA approved or under advanced clinical development such as the ones marketed 10 or developed under the names of Nivolumab (also termed BMS-936558 under development by Bristol Myer Squibb), Pembrolizumab (also termed MK-3475 under development by Merck), and Pidilizumab (also termed CT-011 under development by CureTech). The corresponding nucleotide sequences can be cloned or isolated according to standard techniques based on the information disclosed in the available literature.

15 Another preferred example of immune checkpoint modulator suitable for expression by the oncolytic virus of the invention is represented by a modulator capable of antagonizing at least partially the PD-1 ligand termed PD-L1, and especially an antibody that recognizes human PD-L1. A number of anti PD-L1 antibodies are available in the art (see e.g. those described in EP1907000). Preferred anti PD-L1 antibodies are FDA approved or under advanced clinical development 20 (MPDL3280A under development by Genentech/Roche and BMS-936559 under development by Bristol Myer Squibb as well as anti-PD-L1 Fc fusions (e.g. AMP-224 developed by Medimmune and AstraZeneca).

One may also use an antagonist of the recently identified VISTA protein that was shown to negatively regulate T cell responses (Wang et al., 2011, J. Exp. Med. 208(3): 577-592). VISTA, also 25 designated PD-1H and PD-L3, resembles members of the PD-L1 family,. For example, anti-VISTA antagonists are described in US2013-0177557.

Still another preferred example of suitable immune checkpoint modulator is represented by a modulator capable of antagonizing at least partially the CTLA-4 protein, and especially an antibody that recognizes human CTLA-4. CTLA4 (for cytotoxic T-lymphocyte-associated antigen 4) also known 30 as CD152 was identified in 1987 (Brunet et al., 1987, Nature 328: 267-70) and is encoded by the CTLA4 gene (Dariavach et al., Eur. J. Immunol. 18: 1901-5). CTLA4 is a member of the immunoglobulin superfamily of receptors. It is expressed on the surface of helper T cells where it primarily regulates the amplitude of the early stages of T cell activation. Recent work has suggested that CTLA-4 may function *in vivo* by capturing and removing B7-1 and B7-2 from the membranes of

antigen-presenting cells, thus making these unavailable for triggering of CD28 (Qureshi et al., Science, 2011, 332: 600-3). The complete CTLA-4 nucleic acid sequence can be found under GenBank Accession No LI 5006. A number of anti CTLA-4 antibodies are available in the art (see e.g. those described in US 8,491,895). Preferred anti CTLA-4 antibodies in the context of this invention 5 are FDA approved or under advanced clinical development. One may cite more particularly ipilimumab marketed by Bristol Myer Squibb as Yervoy (see e.g. US 6,984,720; US 8,017,114), tremelimumab under development by Pfizer (see e.g. US 7,109,003 and US 8,143,379) and single chain anti-CTLA4 antibodies (see e.g. WO97/20574 and WO2007/123737).

The oncolytic virus of the present invention may also express an immune checkpoint 10 modulator for antagonizing the LAG3 receptor (see e.g. US 5,773,578).

Another example of suitable immune checkpoint modulator is represented by an OX40 agonist such as agonist ligand of OX40 (OX40L) (see e.g. US 5,457,035, US 7,622,444; WO03/082919) or an antibody directed to the OX40 receptor (see e.g. US 7,291,331 and WO03/106498).

15 Other examples of immune checkpoint modulators are represented by anti-KIR or anti-CD96 antibody targeting the inhibitory receptors harboured by CD8+ T cells and NK cells.

The present invention encompasses an oncolytic virus encoding more than one immune checkpoint modulator. A preferred example includes without limitation expression of an anti-CTLA-4 antibody and an anti-PD-1 antibody.

20

Expression of the one or more nucleic acid molecule(s) encoding the immune checkpoint modulator(s) and if any of the therapeutic genes inserted into the viral genome.

The immune checkpoint modulator-encoding nucleic acid molecule(s) and the therapeutic gene may be easily obtained by standard molecular biology techniques (e.g. PCR amplification, 25 cDNA cloning, chemical synthesis) using sequence data accessible in the art and the information provided herein. Methods for cloning antibodies, fragments and analogs thereof are known in the art (see e.g. Harlow and Lane, 1988, Antibodies – A laboratory manual; Cold Spring Harbor Laboratory, Cold Spring Harbor NY). For example, the nucleic acid molecule (e.g. cDNA) encoding the light and heavy chains of the antibody or their CDRs may be isolated from the producing 30 hybridoma (see e.g. Kohler and Milstein, 1975, Nature 256: 495-7; Cote et al., 1983, Proc. Natl. Acad. Sci. USA 80: 2026-30; Cole et al. in Monoclonal antibodies and Cancer Therapy; Alan Liss pp77-96), immunoglobulin gene libraries, or from any available source or the nucleotide sequence may be generated by chemical synthesis. Analogs and fragments may be generated using standard techniques of molecular biology.

The nucleic acid molecule(s) encoding the immune checkpoint modulator(s) and eventually the therapeutic gene(s) can independently be inserted at any location of the viral genome, with a specific preference for a non-essential locus. Insertion into the oncolytic virus can be performed by routine molecular biology, e.g. as described in Sambrook et al. (2001, Molecular Cloning-A 5 Laboratory Manual, Cold Spring Harbor Laboratory). Insertion into an adenoviral vector or a poxviral vector can be performed through homologous recombination as described respectively in Chartier et al. (1996, J. Virol. 70: 4805-10) and Paul et al. (2002, Cancer gene Ther. 9: 470-7). For example, TK, RR and F2L genes as well as intergenic regions are particularly appropriate for insertion in oncolytic vaccinia virus and E3 and E4 regions for insertion in oncolytic adenovirus virus.

10 In addition, the encoding nucleotide sequences can be optimized for providing high level expression in a particular host cell or subject. It has been indeed observed that, the codon usage patterns of organisms are highly non-random and the use of codons may be markedly different between different hosts. For example, the therapeutic gene may be from bacterial or lower 15 eukaryote origin (e.g. the suicide gene), and thus have an inappropriate codon usage pattern for efficient expression in higher eukaryotic cells (e.g. human). Typically, codon optimization is performed by replacing one or more "native" (e.g. bacterial or yeast) codon corresponding to a codon infrequently used in the host organism of interest by one or more codon encoding the same amino acid which is more frequently used. It is not necessary to replace all native codons corresponding to infrequently used codons since increased expression can be achieved even with 20 partial replacement.

Further to optimization of the codon usage, expression in the host cell or subject can further be improved through additional modifications of the nucleotide sequence(s). For example, various modifications may be envisaged so as to prevent clustering of rare, non-optimal codons being present in concentrated areas and/or to suppress or modify "negative" sequence elements 25 which are expected to negatively influence expression levels. Such negative sequence elements include without limitation the regions having very high (>80%) or very low (<30%) GC content; AT - rich or GC-rich sequence stretches; unstable direct or inverted repeat sequences; R A secondary structures; and/or internal cryptic regulatory elements such as internal TATA-boxes, chi-sites, ribosome entry sites, and/or splicing donor/acceptor sites.

30

In accordance with the present invention, each of the one or more nucleic acid molecule(s) encoding said immune checkpoint modulator(s) as well as the therapeutic gene(s) inserted in the genome of the oncolytic virus of the invention is operably linked to suitable regulatory elements for its expression in a host cell or subject. As used herein, the term "regulatory elements" or

"regulatory sequence" refers to any element that allows, contributes or modulates the expression of the encoding nucleic acid molecule(s) in a given host cell or subject, including replication, duplication, transcription, splicing, translation, stability and/or transport of the nucleic acid(s) or its derivative (i.e. m RNA). As used herein, "operably linked" means that the elements being linked are 5 arranged so that they function in concert for their intended purposes. For example, a promoter is operably linked to a nucleic acid molecule if the promoter effects transcription from the transcription initiation to the terminator of said nucleic acid molecule in a permissive host cell.

It will be appreciated by those skilled in the art that the choice of the regulatory sequences can depend on such factors as the nucleic acid molecule itself, the virus into which it is inserted, the 10 host cell or subject, the level of expression desired, etc. The promoter is of special importance. In the context of the invention, it can be constitutive directing expression of the nucleic acid molecule in many types of host cells or specific to certain host cells (e.g. liver-specific regulatory sequences) or regulated in response to specific events or exogenous factors (e.g. by temperature, nutrient additive, hormone, etc) or according to the phase of a viral cycle (e.g. late or early). One may also 15 use promoters that are repressed during the production step in response to specific events or exogenous factors, in order to optimize virus production and circumvent potential toxicity of the expressed polypeptide(s).

Promoters suitable for constitutive expression in mammalian cells include but are not limited to the cytomegalovirus (CMV) immediate early promoter (US 5,168,062), the RSV promoter, 20 the adenovirus major late promoter, the phosphoglycero kinase (PGK) promoter (Adra et al., 1987, Gene 60: 65-74), the thymidine kinase (TK) promoter of herpes simplex virus (HSV)-1 and the T7 polymerase promoter (WO98/10088). Vaccinia virus promoters are particularly adapted for expression in oncolytic poxviruses. Representative examples include without limitation the vaccinia 7.5K, H5R, 11K7.5 (Erbs et al., 2008, Cancer Gene Ther. 15(1): 18-28), TK, p28, p11, pB2R, pA35R 25 and K1L promoters, as well as synthetic promoters such as those described in Chakrabarti et al. (1997, Biotechniques 23: 1094-7; Hammond et al, 1997, J. Virol Methods 66: 135-8; and Kumar and Boyle, 1990, Virology 179: 151-8) as well as early/late chimeric promoters. Promoters suitable for oncolytic measles viruses include without limitation any promoter directing expression of measles transcription units (Bandler and Tangy, 2008, CIMID 31: 271).

30 In particular, when the encoded immune checkpoint modulator(s) comprise(s) an antibody and especially a mAb, with a specific preference for an anti-PD1 antibody, expression of the heavy chain or fragment thereof is placed under the control of a promoter that is stronger than the one used for expressing light chain or fragment thereof. In particular, the promoter for use for expressing the heavy component provides at least 10%, at least 15%, at least 20%, at least 25%, or

at least 30% more product as compared to expression of said light component. Appropriate promoters for expression can be tested *in vitro* (e.g. in a suitable cultured cell line) or *in vivo* (e.g. in a suitable animal model or in the subject). Examples of suitable promoters for expressing the heavy component of said immune checkpoint modulator comprise CMV, RSV and vaccinia virus pH5R and 5 p11K7.5 promoters. Examples of suitable promoters for expressing the light component of said immune checkpoint modulator comprise PGK, beta-actin and vaccinia virus p7.5K and pA35R promoters.

Those skilled in the art will appreciate that the regulatory elements controlling the expression of the nucleic acid molecule(s) inserted into the viral genome may further comprise 10 additional elements for proper initiation, regulation and/or termination of transcription (e.g. polyA transcription termination sequences), mRNA transport (e.g. nuclear localization signal sequences), processing (e.g. splicing signals), and stability (e.g. introns and non-coding 5' and 3' sequences), translation (e.g. an initiator Met, tripartite leader sequences, IRES ribosome binding sites, signal peptides, etc.).

15 When appropriate, it may be advantageous to include additional regulatory elements to facilitate expression, trafficking and biological activity of at least one of gene inserted into the viral genome of the oncolytic virus of the invention (i.e. the therapeutic gene(s) and/or the one or more immune checkpoint modulator(s)). For example, a signal peptide may be included for facilitating secretion from the infected cell. The signal peptide is typically inserted at the N-terminus of the 20 protein immediately after the Met initiator. The choice of signal peptides is wide and is accessible to persons skilled in the art. One may also envisage addition of a trans-membrane domain to facilitate anchorage of the encoded protein(s) in a suitable membrane (e.g. the plasmic membrane) of the infected cells. The trans-membrane domain is typically inserted at the C-terminus of the protein just before or at close proximity of the STOP codon. A vast variety of trans-membrane domains are 25 available in the art (see e.g. WO99/03885).

As an additional example, a peptide tag (typically a short peptide sequence able to be recognized by available antisera or compounds) may be also be added for following expression, trafficking or purification of the encoded gene product. A vast variety of tag peptides can be used in the context of the invention including, without limitation, PK tag, FLAG octapeptide, MYC tag, HIS 30 tag (usually a stretch of 4 to 10 histidine residues) and e-tag (US 6,686,152). The tag peptide(s) may be independently positioned at the N-terminus of the protein or alternatively at its C-terminus or alternatively internally or at any of these positions when several tags are employed. Tag peptides can be detected by immunodetection assays using anti-tag antibodies.

As another example, the glycosylation can be altered so as to increase biological activity of the encoded gene product (e.g. to increase). Such modifications can be accomplished, for example, by mutating one or more residues within the site(s) of glycosylation. Altered glycosylation patterns may increase the ADCC ability of antibodies and/or their affinity for their target.

5 Another approach that may be pursued in the context of the present invention is coupling of the gene product encoded by the oncolytic virus of the invention to an external agent such as a cytotoxic agent and/or a labelling agent. As used herein, the term "cytotoxic agent" refers to a compound that is directly toxic to cells, preventing their reproduction or growth such as toxins (e. g. an enzymatically active toxin of bacterial, fungal, plant or animal origin, or fragments thereof). As 10 used herein, "a labeling agent" refers to a detectable compound. The labeling agent may be detectable by itself (e. g., radioactive isotope labels or fluorescent labels) or, in the case of an enzymatic label, may catalyze chemical modification of a substrate compound which is detectable. The coupling may be performed by genetic fusion between the gene product (therapeutic gene(s) and/or immune checkpoint modulator(s)) and the external agent.

15

In a preferred embodiment, the oncolytic virus of the invention is a vaccinia virus (preferably from the Copenhagen strain) defective for both TK and RR activities (e.g. resulting from inactivating mutations in both the viral J2R and I4L genes) in the genome of which is inserted a nucleic acid molecule encoding an anti-PD1 antibody. Desirably, the heavy and light chain elements 20 (e.g. heavy and light chains for mAb expression, or variable fragments thereof for Fab and scFv expression) are respectively placed under the transcriptional control of the pH5R and p7.5K vaccinia promoters. Preferably, the anti-PD1-encoding nucleic acid molecule is inserted within TK locus of the viral genome. More preferably, said vaccinia virus is armed with a suicide gene with a special 25 preference for the FCU1 suicide gene described herein. Even more preferably, the suicide gene (e.g. FCU1) is under the transcriptional control of the p11K7.5 vaccinia promoter. Still more preferably, the FCU1 placed under the control of the vaccinia virus promoter is inserted within TK locus of the virus genome.

In an alternative and also preferred embodiment, the oncolytic virus of the invention is a vaccinia virus (preferably from the Wyeth strain) defective for TK activity (resulting from 30 inactivating mutations in the virus J2R gene) in the genome of which is inserted a nucleic acid molecule encoding an anti-PD1 antibody. More preferably, said vaccinia virus is armed with an immunostimulatory therapeutic gene with a special preference for the human GM-CSF gene described herein. Even more preferably, the therapeutic gene (e.g. GM-CSF) is under the

transcriptional control of a synthetic early-late promoter vaccinia promoter and is preferably inserted within TK locus.

Typically, the oncolytic virus of the present invention is produced into a suitable host cell line using conventional techniques including culturing the transfected or infected host cell under suitable conditions so as to allow the production of infectious viral particles and recovering the produced infectious viral particles from the culture of said cell and optionally purifying said recovered infectious viral particles. Suitable host cells for production of the oncolytic virus include without limitation human cell lines such as HeLa (ATCC), 293 cells (Graham et al., 1997, J. Gen. Virol. 10 36: 59-72), HER96, PER-C6 (Fallaux et al., 1998, Human Gene Ther. 9: 1909-17), avian cells such as those described in WO2005/042728, WO2006/108846, WO2008/129058, WO2010/130756, WO2012/001075, etc), hamster cell lines such as BHK-21 (ATCC CCL-10) as well as primary chicken embryo fibroblasts (CEF) prepared from chicken embryos obtained from fertilized eggs. The oncolytic virus can be at least partially isolated before being used according to the present invention. Various purification steps can be envisaged, including clarification, enzymatic treatment (e.g. benzonase, protease), chromatographic and filtration steps. Appropriate methods are described in the art (e.g. WO2007/147528; WO2008/138533, WO2009/100521, WO2010/130753, WO2013/022764).

20 Production of immune checkpoint modulator

In one embodiment, the oncolytic virus may also be utilized in the context of the invention for producing by recombinant means the one or more immune checkpoint modulator(s) that it encodes. It may advantageously comprise one or more additional element(s) enabling maintenance, propagation or expression of the nucleic acid molecule encoding the immune checkpoint modulator 25 in a host cell. Such additional elements comprise marker gene(s) in order to facilitate identification and isolation of the producer host cells (e.g. by complementation of a cell auxotrophy or by antibiotic resistance). Suitable marker genes include without limitation dihydrofolate reductase (dhfr) which confers resistance to methotrexate (Wigler et al., 1980, Proc. Natl. Acad. Sci. USA 77: 3567; O'Hare et al., 1981, Proc. Natl. Acad. Sci. USA 78: 1527); gpt which confers resistance to 30 mycophenolic acid (Mulligan and Berg, 1981, Proc. Natl. Acad. Sci. USA 78: 2072); neo which confers resistance to the aminoglycoside G-418 (Colberre-Garapin et al., 1981, J. Mol. Biol. 150: 1); zeo which confers resistance to zeomycin, kana which confers resistance to kanamycin; hygro, which confers resistance to hygromycin (Santerre et al., 1984, Gene 30: 147). Recombinant viruses lacking a functional TK (e.g. resulting from insertion of the nucleic acid molecule encoding the immune

checkpoint modulator into the TK locus) may be selected with media containing bromodeoxyuridine (BrdU). Indeed, TK- viruses are insensitive to the BrdU drug whereas the drug interferes with DNA synthesis in TK+ viruses. One may also rely on reporter luminescent or colorometric systems, e.g. based on GFP (green fluorescent protein), luciferase and beta-galactosidase.

5 The methods for recombinantly producing the immune checkpoint modulator are conventional in the art. Typically such methods comprise (a) introducing the oncolytic virus described herein into a suitable producer cell to produce a transfected or infected producer cell, (b) culturing in-vitro said transfected or infected producer cell under conditions suitable for its growth, (c) recovering the one or more immune checkpoint modulator(s) from the cell culture, and (d) 10 optionally, purifying the recovered immune checkpoint modulator(s). In the context of the invention, producer cells are preferably human or non-human eukaryotic cells. Preferred producer cells include without limitation BHK-21 (baby hamster kidney), CV-1 (African monkey kidney cell line), COS (e.g. COS-7) cells, Chinese hamster ovary (CHO) cells, mouse NIH/3T3 cells, mouse NSO myeloma cells, HeLa cells, Vero cells, HEK293 cells and PERC.6 cells and avian cells (e.g. chicken, 15 duck cells as described herein).

The producer cells can be cultured in conventional fermentation bioreactors, flasks, and petri plates. Culturing can be carried out at a temperature, pH and oxygen content appropriate for a given host cell. No attempts to describe in detail the various methods known for the production of proteins in eukaryotic cells will be made here. Production of the immune checkpoint modulator can 20 be intracellular or preferably secreted outside the producer cell (e.g. in the culture medium).

The immune checkpoint modulator can then be purified by well-known purification methods. The conditions and technology used to purify a particular protein will depend on factors such as the expression conditions, net charge, molecular weight, hydrophobicity, hydrophilicity and will be apparent to those having skill in the art. Moreover, the level of purification will depend on 25 the intended use. If necessary, especially when the immune checkpoint modulator is not secreted outside the producer cell or where it is not secreted completely, it can be recovered by standard lysis procedures, including freeze thaw, sonication, mechanical disruption, use of lysing agents and the like. If secreted, it can be recovered directly from the culture medium. Suitable purification techniques include without limitation ammonium sulfate precipitation, acid extraction, gel 30 electrophoresis, filtration and chromatographic methods (e.g. reverse phase, size exclusion, ion exchange, affinity, phosphocellulose, hydrophobic-interaction or hydroxylapatite chromatography, etc). Desirably, the immune checkpoint modulator recombinantly produced for the oncolytic virus of the invention is at least partially purified in the sense that it is substantially free of other antibodies having different antigenic specificities and/or other cellular material. Further, the

immune checkpoint modulator may be formulated according to the conditions conventionally used in the art (e.g. WO2009/073569).

Therapeutic use

5 The present invention also provides a composition comprising a therapeutically effective amount of the oncolytic virus of the invention, optionally with a pharmaceutically acceptable vehicle. Such a compositions may be administered once or several times and via the same or different routes

A “therapeutically effective amount” corresponds to the amount of oncolytic virus that is 10 sufficient for producing one or more beneficial results. Such a therapeutically effective amount may vary as a function of various parameters, in particular the mode of administration; the disease state; the age and weight of the subject; the ability of the subject to respond to the treatment; kind of concurrent treatment; the frequency of treatment; and/or the need for prevention or therapy.

When prophylactic use is concerned, the oncolytic virus or the composition of the invention is 15 administered at a dose sufficient to prevent or to delay the onset and/or establishment and/or relapse of a pathologic condition (e.g. a proliferative disease such as cancer), especially in a subject at risk. For “therapeutic” use, the oncolytic virus or the composition of the present invention is administered to a subject diagnosed as having a pathological condition (e.g. a proliferative disease such as cancer) with the goal of treating the disease, eventually in association with one or more 20 conventional therapeutic modalities. In particular, a therapeutically effective amount could be that amount necessary to cause an observable improvement of the clinical status over the baseline status or over the expected status if not treated, e.g. reduction in the tumor number; reduction in the tumor size, reduction in the number or extend of metastasis, increase in the length of remission, stabilization (i.e. not worsening) of the state of disease, delay or slowing of disease

25 progression or severity, amelioration or palliation of the disease state, prolonged survival, better response to the standard treatment, improvement of quality of life, reduced mortality, etc. A therapeutically effective amount could also be the amount necessary to cause the development of an effective non-specific (innate) and/or specific anti-tumor immune response. Typically, development of an immune response in particular T cell response can be evaluated in vitro, in 30 suitable animal models or using biological samples collected from the subject. For example, techniques routinely used in laboratories (e.g. flow cytometry, histology) may be used to perform tumor surveillance. One may also use various available antibodies so as to identify different immune cell populations involved in anti-tumor response that are present in the treated subjects, such as cytotoxic T cells, activated cytotoxic T cells, natural killer cells and activated natural killer

cells. An improvement of the clinical status can be easily assessed by any relevant clinical measurement typically used by physicians or other skilled healthcare staff.

The term “pharmaceutically acceptable vehicle” is intended to include any and all carriers, 5 solvents, diluents, excipients, adjuvants, dispersion media, coatings, antibacterial and antifungal agents, absorption agents and the like compatible with administration in mammals and in particular human subjects.

The oncolytic virus or the composition thereof can be placed in a solvent or diluent appropriate for human or animal use. The solvent or diluent is preferably isotonic, hypotonic or 10 weakly hypertonic and has a relatively low ionic strength. Representative examples include sterile water, physiological saline (e.g. sodium chloride), Ringer's solution, glucose, trehalose or saccharose solutions, Hank's solution, and other aqueous physiologically balanced salt solutions (see for example the most current edition of Remington : The Science and Practice of Pharmacy, A. Gennaro, Lippincott, Williams&Wilkins).

15 In one embodiment, the oncolytic virus composition is suitably buffered for human use. Suitable buffers include without limitation phosphate buffer (e.g. PBS), bicarbonate buffer and/or Tris buffer capable of maintaining a physiological or slightly basic pH (e.g. from approximately pH 7 to approximately pH 9).

The oncolytic virus compositions may also contain other pharmaceutically acceptable 20 excipients for providing desirable pharmaceutical or pharmacodynamic properties, including for example osmolarity, viscosity, clarity, colour, sterility, stability, rate of dissolution of the formulation, modifying or maintaining release or absorption into an the human or animal subject, promoting transport across the blood barrier or penetration in a particular organ.

The oncolytic virus compositions can also comprise one or more adjuvant(s) capable of 25 stimulating immunity (especially a T cell-mediated immunity) or facilitating infection of tumor cells upon administration, e.g. through toll-like receptors (TLR) such as TLR-7, TLR-8 and TLR-9, including without limitation alum, mineral oil emulsion such as, Freunds complete and incomplete (IFA), lipopolysaccharide or a derivative thereof (Ribi et al., 1986, Immunology and Immunopharmacology of Bacterial Endotoxins, Plenum Publ. Corp., NY, p407-419), saponins such as QS21 (Sumino et al., 30 1998, J.Virol. 72: 4931; WO98/56415), imidazo-quinoline compounds such as Imiquimod (Suader, 2000, J. Am Acad Dermatol. 43:S6), S-27609 (Smorlesi, 2005, Gene Ther. 12: 1324) and related compounds such as those described in WO2007/147529, cytosine phosphate guanosine oligodeoxynucleotides such as CpG (Chu et al., 1997, J. Exp. Med. 186: 1623; Tritel et al., 2003, J.

Immunol. 171: 2358) and cationic peptides such as IC-31 (Kritsch et al., 2005, J. Chromatogr Anal. Technol. Biomed. Life Sci. 822: 263-70).

In one embodiment, the oncolytic virus composition of the present invention may be formulated with the goal of improving its stability in particular under the conditions of manufacture 5 and long-term storage (i.e. for at least 6 months, with a preference for at least two years) at freezing (e.g. -70°C, -20°C), refrigerated (e.g. 4°C) or ambient temperatures. Various virus formulation are available in the art either in frozen, liquid form or lyophilized form (e.g. WO98/02522, WO01/66137, WO03/053463, WO2007/056847 and WO2008/114021, etc). Solid (e.g. dry powdered or lyophilized) compositions can be obtained by a process involving vacuum 10 drying and freeze-drying (see e.g. WO2014/053571). For illustrative purposes, buffered formulations including NaCl and sugar are particularly adapted to the preservation of viruses (e.g. Tris 10 mM pH 8 with saccharose 5 % (W/V), sodium glutamate 10 mM, and NaCl 50 mM or phosphate-buffered saline with glycerol (10%) and NaCl).

In certain embodiments, the oncolytic virus composition can be formulated to ensure 15 proper distribution or a delayed release *in vivo*. For example, it can be formulated in liposomes. Biodegradable, biocompatible polymers can be used, such as ethylene vinyl acetate, polyanhydrides, polyglycolic acid, collagen, polyorthoesters, and polylactic acid. Many methods for the preparation of such formulations are described by e.g. J. R. Robinson in "Sustained and Controlled Release Drug Delivery Systems", ed., Marcel Dekker, Inc., New York, 1978.

20 The appropriate dosage of oncolytic virus can be adapted as a function of various parameters and may be routinely determined by a practitioner in the light of the relevant circumstances. Suitable dosage for the oncolytic virus varies from approximately 10^5 to approximately 10^{13} vp (viral particles), iu (infectious unit) or pfu (plaque-forming units) depending on the virus and the quantitative technique used. As a general guidance, vaccinia virus doses from 25 approximately 10^5 to approximately 10^{13} pfu are suitable, preferably from approximately 10^6 pfu to approximately 10^{11} pfu, more preferably from approximately 10^7 pfu to approximately 5×10^9 pfu; doses of approximately 10^8 pfu to approximately 10^9 pfu being particularly preferred especially for human use. The quantity of virus present in a sample can be determined by routine titration techniques, e.g. by counting the number of plaques following infection of permissive cells using 30 permissive cells (e.g. BHK-21 or CEF), immunostaining (e.g. using anti-virus antibodies; Caroll et al., 1997, Virology 238: 198-211), by measuring the A260 absorbance (vp titers), or still by quantitative immunofluorescence (iu titers).

In a preferred embodiment, the administration(s) of said oncolytic virus composition permits to deliver at least 50ng/ml in a body sample obtained from the subject. In one

embodiment, said level of immune checkpoint is obtained after a single administration of approximately 10^7 pfu to approximately 5×10^9 pfu of the oncolytic virus described herein and can be further increased with subsequent administration(s). In another embodiment, the body sample is blood, serum, plasma, tumor biopsy, tumor homogenate, tumor fluid, etc and is collected at least 5 one hour to one month after said administration(s). More preferably, *in situ* production of the encoded immune checkpoint modulator reached at least 100ng (e.g., at least 200ng/ml, at least 400ng/ml, at least 500ng/ml, at least 700 ng/ml, at least 800ng/ml, at least 900ng/ml, at least 1000ng/ml, at least 1200ng/ml, at least 1500ng/ml, at least 1800ng/ml, at least 2000ng/ml, at least at least 2500ng/ml and at least 3000ng/ml) per ml of said biological sample collected from the 10 subject 2 to 8 days (e.g. 3 to 7 days or more preferably about 5 days) following the administration of said oncolytic virus. The levels of immune checkpoint inhibitor secreted *in situ* can be evaluated as described in the Examples.

Administration

15 The oncolytic virus or the composition of the present invention may be administered in a single dose (e.g. bolus injection) or multiple doses. If multiple administrations, they may be performed by the same or different routes and may take place at the same site or at alternative sites. It is also possible to proceed via sequential cycles of administrations that are repeated after a rest period. Intervals between each administration can be from several hours to one year (e.g. 24h, 20 48h, 72h, weekly, every two weeks, monthly or yearly). Intervals can also be irregular (e.g. following tumor progression). The doses can vary for each administration within the range described above.

Any of the conventional administration routes are applicable in the context of the invention including parenteral, topical or mucosal routes. Parenteral routes are intended for administration as an injection or infusion. Common parenteral injection types are intravenous (into a vein), intra- 25 arterial (into an artery), intradermal (into the dermis), subcutaneous (under the skin), intramuscular (into muscle) and intratumoral (into tumor or at its close proximity). Infusions typically are given by intravenous route. Mucosal administrations include without limitation oral/alimentary, intranasal, intratracheal, intrapulmonary, intravaginal or intra-rectal route. Topical administration can also be performed using transdermal means (e.g. patch and the like). Administrations may use conventional 30 syringes and needles (e.g. Quadrafuse injection needles) or any compound or device available in the art capable of facilitating or improving delivery of the active agent(s) in the subject. Preferred routes of administration for the oncolytic virus include intravenous and intratumoral routes.

In the context of the invention, the oncolytic virus may be administered once or several time (e.g. 2, 3, 4, 5, 6, 7 or 8 times etc) at a dose within the range of from 10^7 to 5×10^9 pfu. The time

interval between each administration can vary from approximately 1 day to approximately 8 weeks, advantageously from approximately 2 days to approximately 6 weeks, preferably from approximately 3 days to approximately 4 weeks and even more preferably from approximately 1 week to approximately 3 weeks (e.g. every two weeks for example). A preferred therapeutic 5 scheme involves from 2 to 5 (e.g. 3) intravenous or intratumoral administrations of 10^8 or 10^9 pfu of oncolytic vaccinia virus at approximately 1 or 2 weeks interval.

The present invention also relates to a method for treating a proliferative disease such as cancer comprising administering an oncolytic virus as described herein to a subject in need thereof.

The present invention also relates to a method for inhibiting tumor cell growth *in vivo* 10 comprising administering an oncolytic virus as described herein to a subject in need thereof.

The present invention also relates to a method for enhancing an immune response to tumor cells comprising administering an oncolytic virus as described herein to a subject in need thereof.

In one embodiment, the administration of the oncolytic virus for use in the present invention elicits, stimulates and/or re-orientates an immune response. In particular, the administration 15 induces a protective T or B cell response in the treated host, e.g. against said oncolytic virus or eventually against the product encoded by the therapeutic gene(s) inserted in the viral genome if any. The protective T response can be CD4+ or CD8+ or both CD4+ and CD8+ cell mediated. B cell response can be measured by ELISA and T cell response can be evaluated by conventional ELISpot, ICS assays from any sample (e.g. blood, organs, tumors, etc) collected from the immunized animal 20 or subject.

In one embodiment, the administration of the oncolytic virus permits to change tumor microenvironment with the goal of enhancing activity of effector cells in the tumor, especially effector T lymphocytes and/or promoting at least partial Treg depletion. Tumor infiltrating cells can be easily identified for examples by conventional immunostaining assays.

25 In one embodiment the method of the invention provides a higher therapeutic efficacy than the one obtained in the same conditions with a similar oncolytic virus (without immune checkpoint modulator) or the immune checkpoint modulator either individually or even in co-administration. In the context of the invention, the method of the invention provides at least 5%, at least 10%, at least 15%, at least 20%, or at least 25% more therapeutic efficacy than either the virus or the immune 30 check point modulator alone or in co-administration. A higher therapeutic efficacy could be evidenced as described above in connection with the term "therapeutically effective amount" with a specific preference for a longer survival.

Examples of proliferative diseases that may be treated using the oncolytic virus, composition or methods of the invention include bone cancer, liver cancer, pancreatic cancer,

stomach cancer, colon cancer, cancer of the esophagus, oropharyngeal cancer, lung cancer, cancer of the head or neck, skin cancer, melanoma, uterine cancer, cervix cancer, ovarian cancer, breast cancer, rectal cancer, cancer of the anal region, prostate cancer, lymphoma, cancer of the endocrine system, cancer of the thyroid gland, sarcoma of soft tissue, chronic or acute leukemias, 5 cancer of the bladder, renal cancer, neoplasm of the central nervous system (CNS), glioma, etc. The present invention is also useful for treatment of metastatic cancers, especially metastatic cancers that express PD-L1 (Iwai et al., 2005, *Int. Immunol.* 17: 133-44). Preferred cancers that may be treated using the oncolytic virus of the invention include cancers typically responsive to immunotherapy. Non-limiting examples of preferred cancers for treatment include melanoma (e.g. 10 metastatic malignant melanoma), renal cancer (e.g. clear cell carcinoma), prostate cancer (e.g. hormone refractory prostate adenocarcinoma), breast cancer, colorectal cancer, lung cancer (e.g. non-small cell lung cancer) and liver cancer (e.g. hepatocarcinoma).

According to an advantageous embodiment, especially when the oncolytic virus is armed with a suicide gene, the oncolytic virus therapy or methods according to the present invention may 15 comprise an additional step in which pharmaceutically acceptable quantities of a prodrug, advantageously an analog of cytosine, in particular 5-FC, are administered to the subject. By way of illustration, it is possible to use a dose of from 50 to 500 mg/kg/day, with a dose of 200 mg/kg/day or of 100 mg/kg/day being preferred. Within the context of the present invention, the prodrug is administered in accordance with standard practice (e.g. per os, systematically, etc). Preferably, the 20 administration taking place subsequent to the administration of the oncolytic virus, preferably at least 3 days, more preferably at least 4 days and even more preferably at least 7 days after the administration of the virus. The oral route is preferred. It is possible to administer a single dose of prodrug or doses which are repeated for a time which is sufficiently long to enable the toxic metabolite to be produced within the host organism or cell.

25

The oncolytic virus, composition or method according to the invention can be associated with one or more substances or therapy effective in anticancer therapy and the present invention also concerns a method which comprises the step of delivering to the subject an additional cancer therapy. In the context of the present invention, said additional cancer therapy comprises surgery, 30 radiation, chemotherapy, immunotherapy, hormone therapy or a combination thereof. In a preferred embodiment the method of the invention comprises the administration of one or more substances effective in anticancer therapy. Among pharmaceutical substances effective in anticancer therapy which may be used in association or in combination with the oncolytic virus, composition or method according to the invention, there may be mentioned more specifically:

- ✓ alkylating agents such as e.g. mitomycin C, cyclophosphamide, busulfan, ifosfamide, isosfamide, melphalan, hexamethylmelamine, thiotapec, chlorambucil, or dacarbazine;
- ✓ antimetabolites such as, e.g. gemcitabine, capecitabine, 5-fluorouracil, cytarabine, 2-fluorodeoxy cytidine, methotrexate, idarubicin, tomudex or trimetrexate;
- 5 ✓ topoisomerase II inhibitors such as, e.g. doxorubicin, epirubicin, etoposide, teniposide or mitoxantrone;
- ✓ topoisomerase I inhibitors such as, e.g. irinotecan (CPT-11), 7-ethyl-10-hydroxy-camptothecin (SN-38) or topotecan;
- ✓ antimitotic drugs such as, e.g., paclitaxel, docetaxel, vinblastine, vincristine or vinorelbine;
- 10 ✓ platinum derivatives such as, e.g., cisplatin, oxaliplatin, spiroplatinum or carboplatinum;
- ✓ inhibitors of tyrosine kinase receptors such as sunitinib (Pfizer) and sorafenib (Bayer);
- ✓ anti-neoplastic antibodies in particular antibodies that affect the regulation of cell surface receptors such as trastuzumab, cetuximab, panitumumab, zalutumumab, nimotuzumab, matuzumab, bevacizumab and ranibizumab;
- 15 ✓ EGFR (for Epidermal Growth Factor Receptor) inhibitors such as gefitinib, erlotinib and lapatinib; and
- ✓ immunomodulatory agents such as, e.g. alpha, beta or gamma interferon, interleukin (in particular IL-2, IL-6, IL-10 or IL-12) or tumor necrosis factor;

The oncolytic virus, composition or method according to the invention can also be used in
20 association with radiotherapy.

The present invention also provides kits including a different container (e.g., a sterile glass or plastic vial) for each virus dose to be administered. Optionally, the kit can include a device for performing the administration of the active agents. The kit can also include a package insert including information concerning the compositions or individual component and dosage forms in
25 the kit.

Brief Description of the Drawings

Figure 1 illustrates the quantity of purified anti-PD-1 mAb, Fab and scFv produced from infected CEF.

30 Figure 2 illustrates an analysis by electrophoresis of the purified recombinant mAb (Vacc.mAb) and scFv (Vacc.scFv) as well as commercial J43 (BioXCell) under reduced (R) and non reduced (NR) conditions.

Figure 3: *In vitro* replication efficacy of viruses in LoVo infected at a MOI of 0.0001, 0.001 and 0.01 with the indicated viruses at day 5 post infection. Values are represented in mean ± SD of three individual determinations.

Figure 4 illustrates the amino acid sequences of the heavy and light chains of the anti-PD1 5 antibodies expressed by the oncolytic vaccinia virus described herein.

Figure 5 illustrates immunoblot analysis of the recombinant mAb and Fab secreted in cell culture supernatants obtained from CEF infected with WRTG18618, WRTG18619, WRTG18620 and WRTG18621. Commercial J43 is used as reference; M represents MW markers and C negative control.

10 Figure 6 illustrates the concentration of recombinant J43 produced in the serum (Figure 6A) and in tumor homogenate (Figure 6B) of mice treated with a single administration of WRTG18618 (10^7 pfu) injected intratumorally (resulting from subcutaneous implantation of B16F10 tumors) or subcutaneously (mice without tumors) or with intratumoral administration of 1 μ g or 10 μ g of commercial J43 (Bioxcell). Concentration of recombinant J43 was measured by quantitative ELISA 1, 15 5 and 11 days after injection.

Examples

These examples illustrates oncolytic vaccinia virus engineered for expressing various forms of anti-immune checkpoint inhibitors. Preclinical evidence for the beneficial effects of expressing 20 immune checkpoint blockers from viral vectors was to be demonstrated in mouse tumor models. This implies the use of i) murine-specific anti-immune check point antibodies and ii) an oncolytic poxvirus capable of infecting murine cells with a higher efficacy.

The murine-specific hamster antibody J43 was chosen to target the immune checkpoint blocker murine PD-1 (mPD-1). This antibody was shown to block the interaction of mPD1 with PD-L1 25 (US patent 7,858,746). The antibody J43 and its isotype control (hamster IgG) are available from BioXCell. The anti mPD-1 antibodies and its isotype control were used to establish functional tests and quantitative ELISA *in vitro*. Further, J43 and various forms thereof were cloned in an oncolytic virus and tested *in vivo* in tumor animal models.

The oncolytic virus chosen for these studies is a vaccinia virus (VV) Western Reserve (WR) 30 strain defective for thymidine kinase (TK⁻) (locus J2R) and RR⁻ (locus I4L) rendering the virus non-replicative in healthy (non-dividing) cells. In contrast, the VV TK⁻RR⁻ is supposed to selectively and efficiently replicate in tumor cells.

Vectorization of anti mPD-1 molecules in oncolytic TK RR⁻ vaccinia virus.

The heavy chain of J43 showed high homology with the heavy chain of anti CD79b IgG. Thus, the sequence retained for cloning of the heavy chain was the variable chain of J43 and the constant 5 chain of anti CD79b. The light chain of J43 was cloned with signal sequence from the light chain of anti CD79b antibody. The heavy and light chains were put under the control of the viral promoters pH5R or p7.5K which have slightly different strengths. Further to “whole” antibody constructs, derivatives were generated, respectively His-tagged antigen binding fragments (Fab) constructs as well as a His-tagged single chain antibody (scFv) constructs. Two construct formats were also 10 generated for Fab depending on the light or heavy chain portions placed under each promoter. All five constructs were inserted in the backbone of TK, RR deleted WR VV. The constructs are summarized below:

WRTG18618 (or mAb1) corresponding to pH5R-HC – p7.5K-LC

WRTG18619 (or mAb2) corresponding to pH5R-LC – p7.5K-HC

15 WRTG18621 (or Fab1) corresponding to pH5R-(VH-CH1-6His) – p7.5K-LC

WRTG18620 (or Fab2) corresponding to pH5R-LC – p7.5K-(VH-CH1-6His)

WRTG18616 (scFv) corresponding to pH5R-VH-gs-VL-6His).

HC and LC represents the abbreviation of heavy and light chains, VH and VL of heavy and light variable domains, 6His of the HIS tag (6 histidines) and gs (poly Glycine-Serine linker)

20 Constructs 1 demonstrated the highest expression level of the recombinant mAb or Fab with the expected chain assembly profiles.

Virus stocks WRTG18616 (scFv), WRTG18618 (mAb1) and WRTG18621 (Fab1) were produced in BHK-21 cells and purified by tangential flow filtration (TFF).

25 *Construction of WRTG18618 (pH5R-HC – p7.5K-LC): mAb1*

Fragment containing HC followed by p7.5K promoter was generated by synthetic way (Geneart: Regensburg, Germany) and was cloned in vaccinia transfer plasmid pTG18496 restricted by *Pst*I and *Eco*RI to give pTG18614. Fragment containing LC was generated by synthetic way and was cloned in pTG18614 restricted by *Nsi*I and *Sa*II to give pTG18618.

30 The vaccinia transfer plasmid, pTG18496, is designed to permit insertion of the nucleotide sequence to be transferred by homologous recombination in TK gene of the VV genome. It contains the flanking sequences (BRGTK and BRDTK) surrounding the J2R gene and the pH5R promoter followed by multiple cloning sites.

The amino acid sequences of the anti-PD-1 HC and LC are given in SEQ ID NO: 1 and SEQ ID NO: 2, respectively and in Figure 4.

Generation of WRTG18618 was performed by homologous recombination in primary chicken embryos fibroblasts (CEF) infected with a RR-deleted WR and transfected by nucleofection 5 with pTG18618 (according to Amaxa Nucleofector technology). Viral selection was performed by plaque purification after growth in Thymidine kinase-deficient (TK⁻) 143B cells cultured in the presence of bromodeoxyuridine. This selection allows only TK⁻ rWR to remain viable. Absence of contamination by parental WR was verified by PCR.

10 *Construction of WRTG18619 (pH5R-LC – p7.5K-HC): mAb2*

Fragment containing LC followed by p7.5K promoter was generated by synthetic way and was cloned in vaccinia transfer plasmid pTG18496 restricted by *Pst*I and *Eco*RI to give pTG18615. Fragment containing HC was generated by synthetic way and was cloned in pTG18615 restricted by *Nsi*I and *Mlu*I to give pTG18619.

15 Generation of WRTG18619 virus was performed in CEF by homologous recombination as described above.

Construction of WRTG18621 (pH5R-(VH-CH1-6His) – p7.5K-LC): Fab1

Fragment containing VH-CH1-6His followed by p7.5K promoter was generated by synthetic 20 way and was cloned in vaccinia transfer plasmid pTG18496 restricted by *Pst*I and *Eco*RI to give pTG18617. Fragment containing LC was generated by synthetic way and was cloned in pTG18617 restricted by *Nsi*I and *Sal*I to give pTG18621.

Generation of WRTG18621 virus was performed in CEF by homologous recombination as described above.

25

Construction of WRTG18620 (pH5R-LC – p7.5K-(VH-CH1-6His): Fab2

Fragment containing LC followed by p7.5K promoter was generated by synthetic way and was cloned in vaccinia transfer plasmid pTG18496 restricted by *Pst*I and *Eco*RI to give pTG18615. Fragment containing VH-CH1-6His was generated by synthetic way and was cloned in pTG18615 30 restricted by *Nsi*I and *Mlu*I to give pTG18620.

Generation of WRTG18620 virus was performed in CEF by homologous recombination as described above.

Construction of WRTG18616 (pH5R-VH-gs-VL-6His): scFv

Fragment containing *VH-gs-VL-6His* was generated by synthetic way and was cloned in vaccinia transfer plasmid pTG18496 restricted by *Pst*I and *Eco*RI to give pTG18616.

Generation of WRTG18616 virus was performed in CEF by homologous recombination as described above.

5

In vitro characterization of the encoded anti-PD-1 antibodies

The recombinant oncolytic vectors described above were tested to assess the production of the encoded anti-mPD-1 molecules. To this, permissive primary cells or cell lines were infected, and supernatant harvested. Presence of anti-mPD-1 molecules can be determined by SDS-PAGE, 10 Western blot analysis, mass spectrometry analysis, ELISA and functional assays. Commercially available anti-mPD-1 antibodies were used as controls.

Supernatant from infected CEF (moi 0.2) were analyzed (after 48 and 72 h post infection) by Western Blot under non-reducing conditions using a polyclonal anti-hamster IgG antibody for detection. WRTG18618 (mAb1) and WRTG18621 (Fab1) could be detected by ELISA. Expression 15 profile of the product secreted by WRTG18618-infected cells has similar pattern as commercial J43 antibody and expected assembly of the product secreted by WRTG18621-infected cells was observed. ScFv expression was also detected in culture supernatant by Western Blot at the expected size.

20

mAb purification and In vitro assessment of anti PD-1 antibodies

CEF were cultured in F175 flasks and infected with 2.7×10^8 pfu of anti-PD-1-expressing viruses. After 48-72h, culture supernatants were collected and subjected to filtration on $0.2 \mu\text{m}$ filters. mAb1 was purified by Hitrap protein A (GE Healthcare) followed by Superdex 200 gel filtration (GE Healthcare) whereas Fab1 and scFv were purified by HIS-Trap (GE Healthcare) 25 followed by Superdex 75 gel filtration (GE Healthcare). scFv eluted from gel filtration in two peaks, the first one corresponding to dimers and the second one to monomers. Significant amounts of mAbs, Fab and scFv were produced as shown in Figure 1. The yield is between 0,5 and 2 $\mu\text{g}/\text{ml}$ supernatant according to the antibody format.

Recombinant purified mAb had an electrophoresis profile similar to the commercial mAbs 30 with the correct assembly of the two light and two heavy chains under non reducing conditions and the detection of the individual light and heavy chains under reducing conditions as illustrated in Figure 2. Moreover, purified scFv migrated on SDS-PAGE at the expected size (see Figure 2).

The recombinant products can be detected by ELISA and are functional in a competition ELISA (i.e. blocking of binding of mouse PD-L1 biot at $0.2 \mu\text{g}/\text{mL}$ to mouse PD-1 coated at $1 \mu\text{g}/\text{mL}$),

mAb being more efficient than Fab in this competition experiment. Moreover, the recombinant mAb1 is more efficient than its commercial J43 counterpart as well as VV-produced scFv dimeric fraction is more efficient than the monomeric one with a difference of approximately one log in EC50.

5 The ability of VV-encoded mAb1, Fab1 and scFv to interact with cell-surface PD-1 was studied in flow cytometry-based binding assays using the PD-1-positive T lymphoma cell lines EL4 and RMA. 10^5 EL4 or RMA cells were incubated for 45 min on ice with 100 μ l of either mAb1 (5 μ g/ml), commercially available anti-mPD-1 antibody J43 (5 μ g/ml, BioXCell) or negative control hamster IgG (5 μ g/ml, BioXCell) and washed. Binding to PD-1 was detected by incubating cells for 45 10 min on ice with 100 μ l of 10 μ g/ml FITC-conjugated monoclonal mouse anti-Armenian + Syrian antibody cocktail (BD Pharmingen). After washing, fluorescence intensity was measured on a NaviosTM flow cytometer (Beckman Coulter). Data were analysed using Kaluza 1.2 software (Beckman Coulter). WRTG18618-encoded mAb1 bound efficiently to EL4 and RMA cells.

To measure the binding of Fab1 and scFv, 5×10^5 EL4 or RMA cells were indirectly stained 15 with either 5 μ g/ml Fab1 or 2.5 μ g/ml monomeric scFv (which are both His-tagged) followed by PE-conjugated monoclonal mouse anti-His tag antibody (diluted 1/10, Miltenyi Biotec) and analysed as described above. Fab1 and monomeric scFv produced from WRTG18621 and WRTG18616 were shown to bind efficiently to the surface of EL4 and RMA cells. Specific binding could be demonstrated by co-incubating EL4 cells with Fab1 or monomeric scFv and full-length J43 or 20 negative control hamster IgG from BioXCell, followed by staining with PE-conjugated anti-His tag antibody. EL4 staining was reduced in the presence of full-length J43 but not negative control hamster IgG. Altogether, the results demonstrated that VV-encoded mAb1, Fab1 and scFv were able to recognize endogenously expressed PD-1 in two murine T lymphocyte cell lines.

25 A flow cytometry-based competition assay was established to compare blocking activity of anti-mPD-1 molecules. The assay is based on the murine T lymphoma cell line EL4 which is highly PD-1 positive. The binding of mPD-L1-hFc (murine PD-Ligand 1 comprising human IgG Fc fragment) to the cell-surface PD1 can be detected by flow cytometry using a PE-labeled anti-hFc monoclonal antibody. Binding of mPD-L1-hFc can be competed for by anti-PD-1 antibodies leading to reduction 30 of cell-bound PE signal.

The inhibitory activity of recombinant mAb1, Fab1 and scFv on PD-L1 binding to EL4 cells was assessed and compared to commercially available J43 anti-PD-1 antibody (BioXCell). More specifically, 10^5 EL4 cells were co-incubated with 2 μ g/ml of mPD-L1 hFc (R&D Systems) and increasing concentrations of the various anti-PD-1 antibodies in 100 μ l for 45 min on ice. After

washing, cells were incubated with 100 μ l of 5 μ g/ml PE-conjugated mouse anti-human IgG Fc (BioLegend) for 45 min on ice. Cells were washed and the mean fluorescence intensity was measured as described above. In these conditions, the control anti mPD-1 clone J43 showed an IC₅₀ of 1.6 μ g/ml.

5 As expected, the control hamster IgG did not show any blocking activity on PD-L1 binding to EL4 cells. The recombinant mAb1, Fab1, monomeric and dimeric scFv produced respectively from WRTG18618, WRTG18621 and WRTG18616 were able to block ligand binding to cell-surface receptor as J43 from BioXCell. Importantly, WRTG18618-encoded mAb1 exhibited enhanced blocking activity as compared to J43 from BioXCell and Fab1 and scFv versions in this assay.

10

A quantitative ELISA was established to quantify antibody concentrations produced in mouse serum or by recombinant WR-infected cells using coated rabbit-derived anti hamster antibodies to capture J43, which could be detected in return with a goat-derived anti hamster antibody. The assay is sensitive to mouse serum and standard curves have to be generated in the 15 presence of 50% mouse serum..

***In vitro* cytotoxicity assay (oncolytic activity)**

Human LoVo colon cancer cell line was transduced in suspension at a MOI of 0.001 and 0.0001 by TK- RR deleted WR vaccinia virus without transgene (WRTG18011) and by TK-RR-deleted 20 WR vaccinia virus expressing the different anti mPD-1 molecules (WRTG18616, WRTG18618 and WRTG8621). A total of 3x10⁵ cells/well were plated in 6-well culture dishes in 2 ml of medium supplemented with 10 % FCS. Cells were then cultured at 37°C and cell survival was determined 5 days later by trypan blue exclusion. As shown in Figure 3, the oncolytic activity of the different viruses resulted in 75-95% and 99% reduction in viable cell number at a MOI of 0.0001 and 0.001, 25 respectively. TK-RR-deleted WR virus without transgene and TK-RR deleted WR virus expressing the different anti mPD-1 molecules showed no difference in cytotoxicity. Thus the expression of the different anti mPD-1 molecules did not affect the oncolytic activity of the vaccinia virus.

Optimal design for antibody expression

30 The recombinant oncolytic vectors were tested by Immunoblotting to assess expression levels and assembly of the secreted antibodies. Immunoblot analyses were performed on cell culture supernatants collected 24h after CEF infection with mAb and Fab constructs described above. Supernatant were centrifuged 5 min at 16000 g and 25 μ l were prepared in Laemmli buffer without reducing agent and loaded on PrecastGel 4-15 % polyacrylamide gel (Biorad). Monoclonal

commercial J43 (BioXcell) was used as reference molecule. One μ g of each molecule was loaded on gel. Gel electrophoresis was performed in non-reducing conditions to preserve the assembly of light and heavy chains and allow an optimal detection (*i.e.* the polyclonal antibody used for detection did not recognize reduced IgG and Fab chains). Proteins were then transferred onto a PVDF 5 membrane using the Trans-blot Turbo system (Transblot Turbo Transfer pack Biorad) with the preprogrammed protocol (High MW: 10 min; 2.5 A constant; up to 25V). Membranes were saturated overnight at 4°C in blocking solution (8mM NaPO₄, 2mM KPO₄, 154mM NaCl pH 7.2 (PBS) supplemented with 0.05 % Tween20, 5 % Nonfat dry milk Biorad). Horseradish peroxidase (HRP) conjugated goat anti Armenian Hamster IgG (Jackson Immunoresearch) at 80 ng/mL in dilution 10 buffer (PBS, 0.05 % Tween20, 0.5 % Nonfat dry milk) was used for the antibody immunodetection. Development was performed with Amersham ECL Prime Western Blotting detection reagents and Molecular Imager ChemiDOCTM XRS was used to capture chemiluminescence.

As illustrated in Figure 5, both WRTG18618 and WRTG18619 secreted roughly the same quantities of an antibody product with an apparent size corresponding to two heavy and two light 15 chains linked together comparable to that of commercial J43 mAb. However, WRTG18619 also produced a non-expected product migrating between 43 and 55 kDa markers (see the extra band indicated by an arrow in the fourth lane “18619”). Similarly, an extra band was also present in the WRTG18620-infected culture supernatant (same position and same intensity in the fifth lane “18620”) whereas WRTG18621 produced high amount of correctly assembled Fab without any 20 detectable misassembled product (see arrow head in the sixth lane “18621”).

Analyses of cell culture supernatants generated after infection of mammal cell lines BHK21 and A549 generated the same profile (data not shown).

When looking at the construct design, it appears that in both WRTG18619 and WRTG18620 constructs, the antibody light chain was placed under the transcriptional control of pH5R promoter 25 (a strong promoter). One hypothesis is that in this configuration, antibody light chains could be overexpressed with respect to heavy chains and thus can assemble in homodimers. Therefore, this extra band, migrating between 43 and 55 kDa, could correspond to overproduced light chains that have assembled into a homodimer of a theoretical mass of 47 kDa.

Therefore, it is preferable to express heavy chain under the control of a stronger promoter 30 than the one used to express light chains in order to reduce the risk of generating aberrant assembly (*e.g.* homodimers of light chains)

All together these results confirmed the ability of the recombinant oncolytic viruses described herein to secrete mAb and Fab at detectable level. WRTG18618 and WRTG18621 were

selected for the rest of experiments due to their capacity to produce antibody products that are closer to expectation than those of WRTG18619 and WRTG18620.

In vivo expression of vectorised anti-PD1 antibody

5 Expression of the vectorized anti-PD-1 J43mAb was evaluated *in vivo* in mice with and without subcutaneously-implanted B16F10 tumors. WRTG18618 (mAb1 construct) was injected either intra-tumorally or subcutaneously and compared to commercial J43 (intra-tumoral injection).

In vivo experiments and sample collection

10 More specifically, 3×10^5 B16F10 cells (murine melanoma) were injected subcutaneously (S.C.) into the right flanks of six weeks old female C57BL/6 mice. At day 0 (D0), when tumor volumes reached $100-200 \text{ mm}^3$, 100 microliters of either WRTG18618 (10^7 pfu) or commercial J43 (1 μg or 10 μg , Bioxcell) were injected intra-tumorally (I.T.).

WRTG18618 was also injected S.C in mice without tumor.

15 Blood and tumors were collected at D1, D5 and D11. For each time point, 3 mice were anesthetized with 200 μL pentobarbital and the blood was collected by intra-cardiac puncture. The blood was stored at 4°C during 8 hours and the serum was recovered after two centrifugations and kept at -20°C until analysis. After blood sampling, mice were sacrificed by cervical elongation and the tumor recovered. Tumors were weighted, cut into small pieces and transferred in GentleMACS 20 C-type tubes (Miltenyi) containing 3 ml PBS. Tumors were mechanically dissociated applying program “m-imptumor01” (GentleMacs, Miltenyi). After centrifugation at 300g for 7 min, supernatants were recovered and kept at -20°C until analysis.

J43 concentration was measured by quantitative ELISA both in serum and in tumor homogenates at different time points after virus or commercial J43 mAb injections.

25

Quantitative ELISA

J43 concentrations was evaluated both in serum and tumor homogenates. More specifically, ninety six wells plates (Nunc immune plate Maxisorp) were incubated overnight at 4 °C with 100 μL /well of 0.8 $\mu\text{g}/\text{mL}$ of goat anti-hamster IgG (Southern Biotech) in coating solution 30 (0.05M Na carbonate pH 9.6, Sigma). Plates were washed three times with wash buffer (300 μL /well of PBS, 0.05 % Tween20) and incubated 1h at RT with 200 μL /well of blocking solution (PBS, 0.05% Tween 20, 5 % Non Fat Dry Milk). Plates were washed three times with washing buffer. A standard range of J43 (BioXcell) was prepared in 100 % mouse serum (Sigma) from 1000 to 0.488 ng/mL by 2 fold serial dilutions. Each standard was then further diluted 2 fold in blocking solution (final J43

concentration from 500 to 0.244 ng/mL) to have a final concentration of serum of 50%. One hundred μ L/well of each standard were added in duplicate on the plate. Samples of serum were diluted at least two fold in blocking buffer, and if necessary further diluted in 1Vol/1Vol mix of blocking buffer/serum and 100 μ L were added to the plates. The plates were incubated 2h at 37 °C.

5 After 3 washes with wash buffer, 100 μ L/well of HRP conjugated goat anti Armenian Hamster IgG (Jackson Immunoresearch) at 80 ng/mL were added and plates were incubated 1h at 37 °C. After 3 washes, 100 μ L/well of 3,3',5,5'-tétraméthylbenzidine (TMB , Sigma) were added and the plates incubated at RT for 30 min. The reaction was stopped with 100 μ L/well of 2M H₂SO₄ and the absorbance was measured at 450 nm with a plate reader (TECAN Infinite M200 PRO). The OD values 10 obtained were transferred to the software GraphPadPrism and samples concentrations were back-calculated using the standard curve fitted with 5 parameters.

As shown in Figure 6A, when injected subcutaneously in a naive mice (no tumors implanted), production of recombinant J43 from WRTG18618 was very weak (9ng/ml at day 1 and 4.5ng/ml at day 5). In contrast, high quantities of recombinant J43 were secreted in serum of mice 15 bearing B16F10 tumors following intratumoral WRTG18618 injection, respectively 9 and 1900 times higher at D1 and D5 (83 ng/ml and 9500 ng/ml) than the concentration of recombinant J43 measured after subcutaneous virus injection. This result suggests that the oncolytic virus is multiplying preferentially in tumors rather than in subcutaneous healthy tissues. In comparison, seric concentrations found after intratumoral injection of 10 μ g commercial J43 reached around 20 1000ng/ml (1306 ng/ml at D1 and 1049ng/ml at D5) whereas the seric concentrations were much lower with 1 μ g of commercial J43 (73ng/ml at D1 and 35ng/ml at D5).

In tumor homogenates, the production of recombinant J43 followed the same trend as in serum with a peak of concentration at D5 (see Figure 6B). The pharmacokinetics of J43 after a single IT injection of commercial mAb or WRTG18618 are very different. Following IT injection of 10 μ g of 25 commercial J43, the maximum antibody concentration was measured in serum and in tumor at D1 (1306ng/ml and 173ng/ml respectively) as expected, whereas IT injection of WRTG18618 resulted in maximum seric and tumor concentrations at D5 (9500ng/ml and 3800ng/ml, respectively). It should be highlighted that tumor concentrations of recombinant J43 produced after virus IT injection were much higher than those measured after IT injection of 10 μ g commercial J43 at all time points 30 (410ng/ml versus 173ng/ml at D1; 3800ng/ml versus 51ng/ml at D5 and 700ng/ml versus under detection limit at D11). Importantly, antibody production was still detected in tumors at D11 after virus treatment, which is not the case after commercial J43 injection.

These data suggest that vectorization of J43 in oncolytic viruses allow a higher and longer accumulation of antibody in tumors compared to IT injection of commercial antibody.

Antitumor activity in subcutaneous tumor model

Antitumor activity of the various anti-PD-1-expressing vaccinia viruses described above may be tested in conventional preclinical models after implantation of tumors followed by injection of 5 the constructs. For example, murine cancer cells are injected subcutaneously into the flanks of immunocompetent mice. When tumors reached a volume of 50-70 mm³, the mice are randomized in a blinded manner and treated with the indicated vaccinia virus at a dose of 1 x10⁷ PFU. The vectors or control vehicle (buffer used to resuspend the virus) are directly injected into the tumor at days 10, 12 and 14 after tumor implantation. Tumor size is measured twice weekly using callipers. In 10 particular, vectorised scFv and mAb1 antibody are at least as efficient as the co-administration of a WR vector with commercial J43 to delay tumor growth and increase survival rate.

Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, numerous equivalents to the specific method and reagents described herein, 15 including alternatives, variants, additions, deletions, modifications and substitutions. Such equivalents are considered to be within the scope of this invention and are covered by the following claims.

Claims

1. An oncolytic virus comprising inserted in its genome a nucleic acid molecule encoding one or more immune checkpoint modulator(s).
- 5 2. The oncolytic virus of claim 1, wherein said oncolytic viruses is selected from the group consisting of reovirus, Seneca Valley virus (SVV), vesicular stomatitis virus (VSV), Newcastle disease virus (NDV), herpes simplex virus (HSV), morbillivirus virus, retrovirus, influenza virus, Sinbis virus, poxvirus, and adenovirus.
- 10 3. The oncolytic virus of claim 2, wherein said poxvirus is a vaccinia virus and in particular a vaccinia virus selected from the group of Elstree, Wyeth, Copenhagen and Western Reserve strains.
- 15 4. The oncolytic virus of claim 3, wherein said oncolytic virus is a vaccinia virus defective for thymidine kinase (TK) resulting from inactivating mutations in the J2R viral gene.
5. The oncolytic virus of claim 3 or 4, wherein said virus is a vaccinia virus defective for Ribonucleotide reductase (RR) activity resulting from inactivating mutations in the viral I4L and/or F4L gene(s).
- 20 6. The oncolytic virus of anyone of claims 1 to 5, wherein said oncolytic virus further comprises at least one therapeutic gene inserted in the viral genome.
7. The oncolytic virus of claim 6, wherein said therapeutic gene is selected from the group consisting of genes encoding suicide gene products and genes encoding immunostimulatory proteins.
- 25 8. The oncolytic virus of claim 7, wherein said suicide gene is selected from the group consisting of genes coding protein having a cytosine deaminase (CDase) activity, a thymidine kinase activity, an uracil phosphoribosyl transferase (UPRTase) activity, a purine nucleoside phosphorylase activity and a thymidylate kinase activity.
- 30 9. The oncolytic virus of claim 8, wherein said suicide gene product has CDase and UPRTase activities, and preferably is selected from the group consisting of codA::upp, FCY1::FUR1 and FCY1::FUR1[Delta] 105 (FCU1) and FCU1-8 polypeptides.

10. The oncolytic virus of claim 8 or 9, wherein said oncolytic virus is a vaccinia virus defective for both TK and RR activities and comprising inserted into its genome the therapeutic FCU1 suicide gene.
- 5 11. The oncolytic virus of claim 7, wherein said immunostimulatory protein is an interleukin or a colony-stimulating factor, with a specific preference for GM-CSF.
12. The oncolytic virus of claim 11, wherein said oncolytic virus is a vaccinia virus defective for TK activity and comprising inserted into its genome the therapeutic human GM-CSF.
- 10 13. The oncolytic virus of anyone of claims 1 to 12, wherein said one or more immune checkpoint modulator(s) is selected from the group consisting of peptide ligands, soluble domains of natural receptors, RNAi, antisense molecules, antibodies and protein scaffolds.
- 15 14. The oncolytic virus of claim 13, wherein said one or more immune checkpoint modulator(s) antagonizes at least partially the activity of inhibitory immune checkpoint(s), in particular those mediated by any of the following PD-1, PD-L1, PD-L2, LAG3, Tim3, BTLA and CTLA4.
15. The oncolytic virus of claim 14, wherein said one or more immune checkpoint modulator(s) comprises an antibody that specifically binds to human PD-1 and in particular an anti-PD-1 antibody selected from the group consisting of Nivolumab, Pembrolizumab and pidilizumab.
- 20 16. The oncolytic virus of claim 14, wherein said one or more immune checkpoint modulator(s) comprises an antibody that specifically binds to human PD-L1 and in particular an anti PD-L1 antibody selected from the group consisting of MPDL3280A and BMS-936559.
- 25 17. The oncolytic virus of claim 14, wherein said one or more immune checkpoint modulator(s) comprises an antibody that specifically binds to human CTLA-4 and in particular an anti CTLA4 antibody selected from the group consisting of ipilimumab, tremelimumab and single chain anti-CTLA4 antibodies.
- 30 18. A pharmaceutical composition comprising an effective amount of the oncolytic virus of anyone of claims 1 to 17 and a pharmaceutical acceptable vehicle.
- 35 19. The pharmaceutical composition of claim 18 comprising from approximately 10^7 pfu to approximately 5×10^9 pfu of said oncolytic virus.

20. The pharmaceutical composition of claim 18 or 19 which is formulated for parenteral administration.

21. Use of the oncolytic virus of anyone of claims 1 to 17 or the pharmaceutical composition of
5 anyone of claims 18 to 20, for treating a proliferative disease.

22. The use according to claim 21, wherein said proliferative disease is a cancer, and preferably
a cancer selected from the group consisting of melanoma, renal cancer, prostate cancer,
breast cancer, colorectal cancer, lung cancer and liver cancer.

10

23. The use according to claim 21 or 22, wherein said oncolytic virus is administered by
intravenous or intratumoral route.

15

24. The use according to claim 23, which comprises from 2 to 5 intravenous or intratumoral
administrations of 10^8 or 10^9 pfu of oncolytic vaccinia virus at approximately 1 or 2 weeks
interval.

25. The use according to anyone of claims 21 to 24, which further comprises administration of a
prodrug and/or a substance effective in anticancer therapy.

20



Figure 1

Figure 2

R : reduced condition

NR : Non reduced condition


Figure 3

Figure 4**Heavy chain J43 sequence:**

MGLGLQWVFFVALLKGVHCEVRLLESGGLVKPEGSLKLSCVASGFTFSD
YFMSWVRQAPGKGLEWVAHIYTKSYNYATYYSGSVKGRFTISRDDSRSMV
YLQMNNLRTEDTATYYCTRDGSGYPSLDFWGQGTQVTVSSATTAPSVYP
LAPACDSTTSTTDVTLGCLVKGYFPEPVTVSWNSGALTSGVHTFPSVLH
SGLYSLSSSVTPSSTWPQPICTNVAHPASSTKVDKKIEPRTDTCPN
PPDPCPTCPTPDLLGGPSVIFPPKPKDVLMISSLTPKITCVVVDVSEEEP
DVQFNWYVNNVEDKTAQTETRQRQYNSTYRVVSVLPIKHQDWMSGKVFKC
KVNNNALPSPIEKTKPRGQVRVPQIYTFPPPIEQTVKKDVSVTCLVTG
FLPQDIHVEWESNGQPQPEQNYKNTQPVLSDGSYFLYSKLNVPKSRWDQ
GDSFTCSVIHEALHNHHMTKTISRSLGN

Light chain J43 sequence:

MAWTPGIFMVLSYLTGSFSYELTQPPSASVNGETVKITCSGDQLPKYFA
DWFHQRSQDTILQVIYDDNKRPSGIPERISGSSSGTTATLTIRDVRAEDE
GDYYCFSGYVDSDSKLYVFGSGTQLTVLGGPKSSPKVTVFPPSPEELRTN
KATLVCLVNDFYPGSATVTWKANGATINDGVTTKPSKQGQNYMTSSYLS
LTADQWKSHNRVSCQVTHEGETVEKSLSPAEL

Figure 5





Figure 6A

Figure 6B