



Office de la Propriété
Intellectuelle
du Canada

Un organisme
d'Industrie Canada

Canadian
Intellectual Property
Office

An agency of
Industry Canada

CA 2143648 C 2001/10/16

(11)(21) **2 143 648**

(12) **BREVET CANADIEN
CANADIAN PATENT**

(13) **C**

(86) Date de dépôt PCT/PCT Filing Date: 1994/06/29
(87) Date publication PCT/PCT Publication Date: 1995/01/26
(45) Date de délivrance/Issue Date: 2001/10/16
(85) Entrée phase nationale/National Entry: 1995/02/28
(86) N° demande PCT/PCT Application No.: US 94/07355
(87) N° publication PCT/PCT Publication No.: WO 95/02321
(30) Priorités/Priorities: 1993/07/14 (08/091,701) US;
1993/11/30 (08/161,211) US

(51) Cl.Int.⁶/Int.Cl.⁶ A01K 43/08, A23L 1/32, A01K 45/00,
B65D 85/32
(72) Inventeur/Inventor:
Brunnquell, John R., US
(73) Propriétaire/Owner:
Wisconsin Alumni Research Foundation, US
(74) Agent: GOUDREAU GAGE DUBUC

(54) Titre : OEUFs A FAIBLE TENEUR EN CHOLESTEROL
(54) Title: LOW CHOLESTEROL EGGS

(57) **Abrégé/Abstract:**

Reduced cholesterol, low fat eggs are obtained. At least a twenty-five percent reduction in cholesterol is achieved over normal levels for the grade size. The reduction can be achieved by weight monitoring of the egg, size monitoring of the yolk through candling, and selecting eggs from high producing, young, and high clutch chickens.



PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ :

A01K 43/06, 43/00, G01N 33/08, A23L
1/32

A1

(11) International Publication Number:

WO 95/02321

(43) International Publication Date:

26 January 1995 (26.01.95)

(21) International Application Number:

PCT/US94/07355

(22) International Filing Date:

29 June 1994 (29.06.94)

(30) Priority Data:

08/091,701

14 July 1993 (14.07.93)

US

08/161,211

30 November 1993 (30.11.93)

US

(71) Applicant: WISCONSIN ALUMNI RESEARCH FOUNDATION [US/US]; 614 Walnut Street, P.O. Box 7365, Madison, WI 53707-7365 (US).

(72) Inventor: BRUNNQUELL, John, R.; 3267 Highway W, Port Washington, WI 53074 (US).

(74) Agents: SCHWARTZ, Carl, R. et al.; Quarles & Brady, 411 East Wisconsin Avenue, Milwaukee, WI 53202-4497 (US).

(81) Designated States: CA, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).

Published

With international search report.

(54) Title: LOW CHOLESTEROL EGGS

(57) Abstract

Reduced cholesterol, low fat eggs are obtained. At least a twenty-five percent reduction in cholesterol is achieved over normal levels for the grade size. The reduction can be achieved by weight monitoring of the egg, size monitoring of the yolk through candling, and selecting eggs from high producing, young, and high clutch chickens.

LOW CHOLESTEROL EGGS

Background Of The Invention

Technical Field

5 The present invention relates to techniques for producing and selecting low cholesterol chicken eggs. It appears to be especially well suited for use in connection with "large" size eggs.

Background Art

10 Serum cholesterol has been implicated in connection with atherosclerosis, a form of heart disease. Efforts have therefore been made to lower serum cholesterol. One approach is to alter the diet by reducing cholesterol intake.

A source of dietary cholesterol is chicken eggs. For example, a "large" chicken egg contains on the
15 average 213 milligrams of total cholesterol. Completely eliminating egg product intake would adversely affect

-2-

many standard food recipes. While egg substitutes are lower in cholesterol, they are usually poor imitations of real eggs from a texture and taste standpoint. They also often differ in other characteristics that are important to consumers. In this regard, egg substitutes often either contain no egg yolks, or they contain egg yolks that have been treated in ways that are detrimental.

Another concern is the embarrassment factor for those who use egg substitutes. When purchasing or using such specialty dietary products there is a risk that private medical conditions will become public knowledge.

Efforts have therefore been made to try to reduce the cholesterol levels in the eggs themselves. One approach is by controlling the diet of chickens by using low fat, high fiber diets, and by reducing the stress levels for chickens (e.g. using reduced and/or red lights; providing a clean environment; reducing the number of chickens per cage). At best, this has lead to "large" eggs which average (in a randomly selected test sample) above 165 mg per 50 g of egg white and yolk.

U.S. regulatory authorities have promulgated rules that will prohibit advertisement of reduced cholesterol levels in foods unless there is at least a twenty-five percent reduction from "normal" levels. Thus, "large" eggs having (on average) 160 mg or less total cholesterol per 50 g of egg white and yolk are highly desired.

"Total cholesterol" can be determined using procedures based on M. Adams et al., 69 J. Assoc. Off. Anal. Chem. 844-846 (1986). See also Official Methods Of Analysis (1990), 15th Edition, Method 976.26 AOAC Arlington, Virginia.

There have been attempts to reduce the cholesterol content of eggs using drugs. In fact, sterols and triparanol have achieved greater than 25% reductions in chicken eggs. However, these drugs are not on the GRAS

2143648

22 05 95

-3-

(generally recognized as safe) listing. Thus, their use would cause adverse labeling and sales restrictions, and there may well be safety problems associated with their use.

5 Candling is a known technique of shining a light through an egg in order to spot cracks or blood spots by viewing the shadow. It has been reported that candling techniques are now being used with computer imaging systems so as to automate defect spotting. See Poultry
10 Times, p. 6, (March 1, 1993). However, the art has not applied candling techniques to assist in selecting low cholesterol eggs.

The need therefore still exists to achieve a 25% or more reduction in total cholesterol in chicken eggs using
15 techniques that do not require the use of sterols, triparanol, or other drugs that are not on the GRAS list.

Disclosure Of The Invention

In one aspect, the invention provides a container containing at least six (preferably a dozen) chicken
20 eggs. The eggs in the container average at least 1.5 ounce in weight per egg, and the average total cholesterol content of the eggs is less than 125 milligrams per 37.5 gram of egg white and yolk. The
cholesterol level is achieved without having to supply to
25 the chicken or chickens that laid the eggs exogenously supplied steroid, sterol, or triparanol so that the eggs do not contain exogenously supplied steroid, sterol, or triparanol.

In another aspect, the invention provides a method
30 for obtaining a collection of at least six whole chicken eggs, characterized in that the collection has an average total cholesterol content of 125 mg or less per 37.5 g of egg white and yolk, the eggs in the collection not contain exogenously supplied steroid, sterol, or
35 triparanol, and the eggs in the collection average at minimum 1.5 ounces in weight per egg. The method involves creating a supply of eggs by feeding a

2143648

22 00 00

-4-

population of chickens a diet free of steroid, sterol, or triparanol. One then selects from the supply only those eggs having a weight not more than .083 ounces above the minimum average weight and either (a) candles the eggs selected from the supply of eggs to select out and thus not include eggs having a yolk shadow above a selected size (so as to include eggs of a lower average cholesterol concentration in the white and yolk than the average of the supply); or (b) selects from the supply for the collection only those eggs that have been produced by a chicken or chickens that were fed yeast prior to laying the eggs and that were less than 38 weeks old when the chicken or chickens laid those eggs, and that were from a flock with at least 80% productivity when the selected eggs were laid.

Depending upon whether the eggs are medium, large, extra large, or jumbo, the minimum ounce averages per egg can be respectively 1.75, 2, 2.25, or 2.5, and the cholesterol maximum averages less than 145 mg per 43.75 g of egg white and yolk, 165 mg per 50 g of egg white and yolk, 185 mg per 56.25 g of egg white and yolk, or 205 mg per 62.5 g of egg white and yolk, respectively.

In an even more preferred form, only those eggs are selected where the layer or layers had their diet supplemented with Yea-Sacc yeast prior to the laying the eggs, and the eggs are from a Single Comb White Leghorn.

Best Modes For Carrying Out The Invention

Breed: The techniques described herein are believed to be applicable to a wide variety of chickens (Gallus Domesticus), and in particular to high production, commercially prevalent species thereof such as the White Leghorn (preferably the Single Comb White Leghorn) and the New Hampshire. Crosses between these species are also useable. The invention should also have applicability to low cholesterol mutant forms such as

that reported in A. Quershi, et al. 34 Nutritional Reports International 457-464 (1986).

Egg Size

In the United States, eggs are graded by weight based on the number of ounces (shell included) per dozen. A "small" egg is 18 to 20.99 ounces per dozen. A "medium" egg is 21 to 23.99 ounces per dozen. A "large" egg is 24 to 26.99 ounces per dozen. An "extra large" egg is 27 to 29.99 ounces per dozen. A "jumbo" egg is 30 or above ounces per dozen. The experiments described below were performed with large eggs. It should be appreciated that the same techniques can be applied to other sizes of eggs.

The large egg is preferred as persons who are on a diet are reluctant to select "extra large" or "jumbo" eggs. Also, medium and small eggs are too small for many consumers to feel that the product has value.

Eggs are first sorted by weight to divide them into the various sizes of small, medium, large, extra large and jumbo. In accordance with the present invention, there is then a further selection in which only those eggs in the lower third of the weight size range (preferably those in the lower sixth of the range) are selected (e.g. for large, below 2.083 ounces per egg).

25 Yolk Size

In addition to the above type of selection, one can candle (illuminate) the egg in order to determine the area of the shadow that the egg yolk throws. Conventional candling equipment currently used for spotting cracks or fertilized eggs can be used to project the shadow in the usual fashion. However, in accordance with the invention, the candling can project the shadow on a wall that has a bull's eye target type pattern.

When the yolk shadow exceeds a selected area, the egg can be selected out.

For example, this shadow can be compared to a selected norm (e.g. one that only selects on average the smallest 1/3 or 1/6th yolk shadow area for the grade). In this regard, almost all of the cholesterol in an egg is in its yolk. Therefore, while selection by overall weight is helpful, a further desirable step involves the selection by yolk size after the initial weight

10 selection.

In the alternative, the shadow can be projected onto computer imaging equipment, and a computer can then calculate even more accurately the overall yolk shadow area.

15 Young Layers

Eggs collected shortly after a chicken reaches maturity for egg laying typically have the lowest concentration of cholesterol per gram of yolk. The cholesterol concentration in the yolk typically increases over time until the first molt. Thereafter, it drops somewhat (but generally not all the way back to the beginning levels). It then increases again.

20

Thus, the total cholesterol concentration in the yolk zig-zags upwardly with chicken age, with the lowest cholesterol level being in the youngest chickens. To assist in achieving the results of the present invention it is preferred to select eggs from chickens that are between initial maturity and 38 weeks old. If extremely low levels need to be achieved, the selection could be skewed to the early part of the range (e.g. 80% or more 18 weeks old).

25

30

Productivity

The egg in a chicken is believed to be the excess cholesterol deposit site for the chicken. Typically, the more eggs that the chicken lays, the lower the concentration of cholesterol found in the average yolk. As such, it is preferred to select eggs from only flocks averaging over 80% production per day (80% of the chickens lay an egg that day).

Clutch Size

A clutch is the period of consecutive days in which a hen lays an egg. If a hen lays an egg on eight consecutive days, it would have a clutch size of eight. It has been learned that those eggs which appear later in a clutch typically have the lowest concentration of cholesterol. Thus, it is preferred to select eggs from clutch sizes of five or more, with eggs from days 5 or later being selected.

Low Stress

It has been learned that high levels of stress can increase the level of cholesterol in the egg. Under extremely low stress conditions, the cholesterol concentration in the chicken's egg is typically slightly reduced. Techniques that have proved to be of assistance in connection with lowering the level of cholesterol are housing the chickens in indirect light with high fresh air circulation; keeping the chickens caged and not crowded (not on the floor); keeping the cage areas very clean; and periodically providing soothing and relatively quiet music (e.g. classical music) for the chickens.

2143648

-8-

Diet

When one provides a low fat, high fiber diet to the chicken, cholesterol levels will decrease. A preferred feed contains about twenty to twenty-five pounds per day, per 100 birds, of the following feed: 50-75% corn, 10-30% soy bean meal, grit, standard vitamin, and calcium mix, 2-10% alfalfa, 20-100 gms/ton of niacin, 2 pounds/ton of Yea-Sacc 1026 or 1036 yeast (available from All-Tech).

Example

Applicants housed 5,600 Single Comb White Leghorns in a barn, and used the following selection process for the unfertilized eggs. They selected eggs only from chickens of 30 weeks or younger, and only eggs between 2 ounces and 2.083 ounces. A flock having over 80% productivity at the time of selection was used.

The standard environment for the chickens was 65-75°F, with 14-17 hours of light per day. (At week 18 of a chicken's life the light was 14 hours per day, with the light gradually being increased to 17 hours by the 28th week). This "pseudo-spring" further increases production. The birds were caged with an average of at least 54 square inches per bird.

The feed for the birds was 20 gms/ton niacin; 80 pounds/ton dehydrated alfalfa meal; 2 pounds/ton Yea-Sacc yeast; about 398 pounds/ton soy bean meal; 120 pounds/ton calcium; 1390 pounds/ton corn; and 10 pounds/ton grit and conventional nutritional requirements for the chicken ("N.R.C": standard vitamins and minerals). Eggs selected using the above procedures were then assayed for total egg cholesterol. The average (over twelve selected large eggs) was 165 milligrams cholesterol per 51.22 g of egg white and yolk (which equals roughly 161 mg on a 50 g basis).

-9-

In another separate test, a sample of eggs averaged 159.7 mg cholesterol per 50 g egg white and yolk.

Industrial Applicability

It should be appreciated that the above experiments were primarily intended to achieve the minimum government labelling requirements. Even further reductions in cholesterol content may be desirable. In this regard, the candling selection step can be added so as to further reduce the average cholesterol content.

For various health and regulatory reasons it is also desirable to reduce total fat and saturated fat levels in foods.

Surprisingly, the methods of the present invention also provide eggs with reduced levels of total and saturated fat. For purposes of the claims, total fat is determined by the ether extraction method. See e.g. Official Methods Of Analysis, 15th Edition, 925.32 AOAC, (1990).

Typical levels of total fat for prior art chicken eggs average as follows: small -3.8 gm/egg; medium -4.2 gm/egg; large -4.6 gm/egg; extra large -5.8 gm/egg; jumbo -6.2 gm/egg. In one test of the present invention, a sample of large eggs achieved an average total fat content of 3.8 grams fat per egg (compared with the normal -4.6 gm/egg), with the saturated fat portion reduced by more than 10%.

It should be appreciated that only the preferred embodiments have been described above. The following claims should therefore be looked to in order to judge the full scope of the invention.

-10-

Claims

I claim:

1. A container containing at least six whole chicken eggs, characterized in that eggs in the container average at least 1.5 ounce in weight per egg, wherein the average total cholesterol content of eggs in the container is less than 125 milligrams per 37.5 g of egg white and yolk, wherein the cholesterol content was achieved without having to supply to a chicken or chickens that laid the eggs exogenously supplied steroid, sterol, or triparanol, and wherein the eggs do not contain exogenously supplied steroid, sterol, or triparanol.

2. The container of claim 1, wherein the eggs in the container average at least 1.75 ounces in weight per egg and wherein the average total cholesterol content of the eggs in the container is less than 145 milligrams per 43.75 g of egg white and yolk.

3. The container of claim 1, wherein the eggs in the container average at least 2 ounces in weight per egg and wherein the average total cholesterol content of the eggs in the container is less than 165 milligrams per 50 g of egg white and yolk.

4. The container of claim 3, wherein the average total cholesterol content of the eggs in the container is less than 162 milligrams per 50 g of egg white and yolk.

22 05 98

-11-

5. The container of claim 4, wherein the average total cholesterol content of the eggs in the container is less than 160 milligrams per 50 g of egg white and yolk, and the eggs are Single Comb White Leghorn eggs.

6. The container of claim 1, wherein the eggs in the container average at least 2.25 ounces in weight per egg and wherein the average total cholesterol content of the eggs in the container is less than 185 milligrams per
5 56.25 g of egg white and yolk.

7. The container of claim 1, wherein the eggs in the container average at least 2.5 ounces in weight per egg, and wherein the average total cholesterol content of the eggs in the container is less than 205 milligrams per
5 62.5 g of egg white and yolk.

8. The container of claim 1, wherein the egg white and yolk in the eggs in the container average less than 3.5 gm total fat per egg.

9. The container of claim 2, wherein the egg white and yolk in the eggs in the container average less than 3.85 g total fat per egg.

-12-

10. The container of claim 3, wherein the egg white and yolk in the eggs in the container average less than 3.85 g total fat per egg.

11. The container of claim 6, wherein the egg white and yolk in the eggs in the container average less than 5.0 g total fat per egg.

12. The container of claim 7, wherein the egg white and yolk in the eggs in the container average less than 5.5 g total fat per egg.

-13-

13. A method for obtaining a collection of at least six whole chicken eggs, characterized in that the collection has an average total cholesterol content of 125 mg or less per 37.5 g of egg white and yolk, the eggs in the collection do not contain exogenously supplied steroid, sterol, or triparanol, and the eggs in the collection average at minimum 1.5 ounces in weight per egg, the method comprising:

creating a supply of eggs by feeding a population of chickens a diet free of steroid, sterol, or triparanol;

selecting from the supply only those eggs having a weight not more than .083 ounces above the minimum average weight; and either (a) candling the eggs selected from the supply to select out and thus not include eggs having a yolk shadow above a selected size so as to include eggs of a lower average cholesterol concentration in the egg white and yolk than the average of the supply; or (b) selecting from the supply for the collection only those eggs that have been produced by a chicken or chickens that were fed yeast prior to laying the egg and that were less than 38 weeks old when the chicken or chickens laid those eggs, and that were from a flock with at least 80% productivity when the selected eggs were laid.

14. The method of claim 13, wherein (a) applies and the average total cholesterol content of eggs in the collection is 145 mg or less per 43.75 g of egg white and yolk, the eggs in the collection weigh an average of at least 1.75 ounces per egg; and therefore none of the eggs in the collection exceeds 1.833 ounces.

-14-

15. The method of claim 13, wherein (a) applies and the average total cholesterol content of eggs in the collection is 165 mg or less per 50 g of egg white and yolk, the eggs in the collection weigh an average of at least 2 ounces per egg; and therefore none of the eggs in the collection exceeds 2.083 ounces.

16. The method of claim 13, wherein (a) applies and the average total cholesterol content of eggs in the collection is 185 mg or less per 56.25 g of egg white and yolk, the eggs in the collection weigh an average of at least 2.25 ounces per egg; and therefore none of the eggs in the collection exceeds 2.333 ounces.

17. The method of claim 13, wherein (a) applies and the average total cholesterol content of eggs in the collection is 205 mg or less per 62.6 g of egg white and yolk, the eggs in the collection weigh an average of at least 2.5 ounces per egg; and therefore none of the eggs in the collection exceeds 2.583 ounces.

18. The method of claim 13, wherein (b) applies and the chicken eggs in the collection are Single Comb White Leghorn chicken eggs.

-15-

19. The method of claim 13, wherein (b) applies and the collection contains eggs only from clutch day five or later.

QB2\157548.01

ATTACHED SHEET