



(43) International Publication Date
24 October 2013 (24.10.2013)

(10) International Publication Number
WO 2013/159108 A2

(51) International Patent Classification:
A61K 48/00 (2006.01)

(21) International Application Number:
PCT/US2013/037638

(22) International Filing Date:
22 April 2013 (22.04.2013)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
61/636,513 20 April 2012 (20.04.2012) US

(71) Applicant: **ISIS PHARMACEUTICALS, INC.** [US/US];
2855 Gazelle Court, Carlsbad, CA 92010 (US).

(72) Inventors: **FREIER, Susan, M.**; 2855 Gazelle Court,
Carlsbad, CA 92010 (US). **SWAYZE, Eric, E.**; 2855
Gazelle Court, Carlsbad, CA 92010 (US).

(74) Agents: **SCARR, Rebecca** et al.; Casimir Jones, 2275 De-
ming Way, Suite 310, Middleton, WI 53562 (US).

(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY,

BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM,
DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT,
HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP,
KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD,
ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI,
NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU,
RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ,
TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA,
ZM, ZW.

(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ,
UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ,
TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK,
EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV,
MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM,
TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW,
ML, MR, NE, SN, TD, TG).

Published:

- without international search report and to be republished
upon receipt of that report (Rule 48.2(g))
- with sequence listing part of description (Rule 5.2(a))

(54) Title: OLIGOMERIC COMPOUNDS COMPRISING BICYCLIC NUCLEOTIDES AND USES THEREOF

(57) Abstract: The present invention provides oligomeric compounds. Certain such oligomeric compounds are useful for hybridizing to a complementary nucleic acid, including but not limited to, nucleic acids in a cell. In certain embodiments, hybridization results in modulation of the amount activity or expression of the target nucleic acid in a cell.



WO 2013/159108 A2

OLIGOMERIC COMPOUNDS COMPRISING BICYCLIC NUCLEOTIDES AND USES THEREOF**SEQUENCE LISTING**

The present application is being filed along with a Sequence Listing in electronic format. The Sequence Listing is provided as a file entitled CORE0104USLSEQ.txt, created April 20, 2012, which is 4 Kb in size. The information in the electronic format of the sequence listing is incorporated herein by reference in its entirety.

BACKGROUND OF THE INVENTION

Antisense compounds have been used to modulate target nucleic acids. Antisense compounds comprising a variety of chemical modifications and motifs have been reported. In certain instances, such compounds are useful as research tools, diagnostic reagents, and as therapeutic agents. In certain instances antisense compounds have been shown to modulate protein expression by binding to a target messenger RNA (mRNA) encoding the protein. In certain instances, such binding of an antisense compound to its target mRNA results in cleavage of the mRNA. Antisense compounds that modulate processing of a pre-mRNA have also been reported. Such antisense compounds alter splicing, interfere with polyadenylation or prevent formation of the 5'-cap of a pre-mRNA.

Certain antisense compounds have been described previously. See for example United States Patent No. 7,399,845 and published International Patent Application No. WO 2008/049085, which are hereby incorporated by reference herein in their entirety.

SUMMARY OF THE INVENTION

In certain embodiments, the present invention provides compounds comprising oligonucleotides. In certain embodiments, such oligonucleotides comprise a gapmer region. In certain embodiments, such oligonucleotides consist of a gapmer region.

The present disclosure provides the following non-limiting numbered embodiments:

Embodiment 1: A compound comprising:

a modified oligonucleotide consisting of 10 to 20 linked nucleosides, wherein the modified oligonucleotide comprises:

a 5'-wing consisting of 2 to 5 linked nucleosides;

a 3'-wing consisting of 2 to 5 linked nucleosides; and

a gap between the 5'-wing and the 3'-wing consisting of 6 to 14 linked 2'-deoxynucleosides; and

wherein at least one of the 5'-wing and the 3'-wing comprises at least one bicyclic nucleoside; at least one of the 5'-wing and the 3'-wing comprises at least one 2'-substituted nucleoside; and
wherein the nucleobase sequence of the modified oligonucleotide is complementary to the nucleobase sequence of a target nucleic acid.

Embodiment 2: The compound of embodiment 1, wherein one of the 5'-wing or the 3'-wing comprises at least one 2'-deoxynucleoside.

Embodiment 3: The compound of embodiments 1-2, wherein each of the 5'-wing and the 3'-wing comprises at least one 2'-deoxynucleoside.

Embodiment 4: The compound of embodiments 1-3, wherein the 3'-wing comprises at least one 2'-deoxynucleoside.

Embodiment 5: The compound of embodiments 1-4, wherein the 5'-wing comprises at least one 2'-deoxynucleoside.

Embodiment 6: The compound of any of embodiments 1-5, wherein the 5'-wing comprises at least one bicyclic nucleoside.

Embodiment 7: The compound of any of embodiments 1-6, wherein the 3'-wing comprises at least one bicyclic nucleoside.

Embodiment 8: The compound of any of embodiments 1-7, wherein the 5'-wing comprises at least one 2'-substituted nucleoside.

Embodiment 9: The compound of any of embodiments 1-8, wherein the 3'-wing comprises at least one 2'-substituted nucleoside.

Embodiment 10: A compound comprising:
a modified oligonucleotide consisting of 10 to 20 linked nucleosides, wherein the modified oligonucleotide comprises:
a 5'-wing consisting of 2 to 5 linked nucleosides;
a 3'-wing of 2 to 5 linked nucleosides; and

a gap between the 5' wing and the 3' wing consisting of 6 to 14 linked 2'-deoxynucleosides; and
 wherein at least one of the 5'-wing and the 3'-wing comprises at least one constrained ethyl nucleoside; and
 at least one of the 5'-wing and the 3'-wing comprises at least one 2'-substituted nucleoside; and
 wherein the nucleobase sequence of the modified oligonucleotide is complementary to the nucleobase
 sequence of a target nucleic acid.

Embodiment 11: The compound of embodiments 1-10, wherein and at least one of the 5'-wing and the
 3'-wing comprises at least one 2'-deoxynucleoside.

Embodiment 12: The compound of embodiments 1-11, wherein at least one of the 5'-wing and the 3'-
 wing comprises both at least one constrained ethyl nucleoside and at least one 2'-substituted nucleoside.


Embodiment 13: The compound of embodiments 1-12, wherein the 5'-wing comprises at least one
 constrained ethyl nucleoside.

Embodiment 14: The compound of any of embodiments 10-13, wherein the 3'-wing comprises at least
 one constrained ethyl nucleoside.

Embodiment 15: The compound of any of embodiments 10-14, wherein the 5'-wing comprises at least
 one 2'-substituted nucleoside.

Embodiment 16: The compound of any of embodiments 10-15, wherein the 3'-wing comprises at least
 one 2'-substituted nucleoside.

Embodiment 17: The compound of any of embodiments 1-17, wherein the modified oligonucleotide
 has a sugar motif described by Formula I as follows:



wherein:

each A is independently a 2'-substituted nucleoside;

each B is independently a bicyclic nucleoside;

each J is independently either a 2'-substituted nucleoside or a 2'-deoxynucleoside;

each D is a 2'-deoxynucleoside;


m is 0-4; n is 0-2; p is 0-2; r is 0-2; t is 0-2; v is 0-2; w is 0-4; x is 0-2; y is 0-2; z is 0-4; and g is 6-
 14;

provided that:

at least one of m, n, and r is other than 0;
 at least one of w and y is other than 0;
 the sum of m, n, p, r, and t is from 2 to 5; and
 the sum of v, w, x, y, and z is from 2 to 5.

Embodiment 18: A compound comprising:

a modified oligonucleotide consisting of 10 to 20 linked nucleosides, wherein the modified oligonucleotide has a sugar motif described by Formula I as follows:



wherein:

each A is independently a 2'-substituted nucleoside;
 each B is independently a bicyclic nucleoside;
 each J is independently either a 2'-substituted nucleoside or a 2'-deoxynucleoside;
 each D is a 2'-deoxynucleoside;
 m is 0-4; n is 0-2; p is 0-2; r is 0-2; t is 0-2; v is 0-2; w is 0-4; x is 0-2; y is 0-2; z is 0-4; and g is 6-14;

provided that:

at least one of m, n, and r is other than 0;
 at least one of w and y is other than 0;
 the sum of m, n, p, r, and t is from 2 to 5; and
 the sum of v, w, x, y, and z is from 2 to 5.

Embodiment 19: The compound of embodiment 17 or 18, wherein at least one bicyclic nucleoside is a constrained ethyl nucleoside.

Embodiment 20: The compound of embodiment 17 or 18, wherein each bicyclic nucleoside is a constrained ethyl nucleoside.

Embodiment 21: The compound of any of embodiments 17-19, wherein at least one bicyclic nucleoside is an LNA nucleoside.

Embodiment 22: The compound of embodiment 17 or 18, wherein each bicyclic nucleoside is an LNA nucleoside.

Embodiment 23: The compound of any of embodiments 1-22, wherein the 2'-substituent of the at least one 2'-substituted nucleoside is selected from among: OCH_3 , F, OCH_2F , OCHF_2 , OCF_3 , OCH_2CH_3 , $\text{O}(\text{CH}_2)_2\text{F}$, OCH_2CHF_2 , OCH_2CF_3 , $\text{OCH}_2\text{-CH=CH}_2$, $\text{O}(\text{CH}_2)_2\text{-OCH}_3$, $\text{O}(\text{CH}_2)_2\text{-SCH}_3$, $\text{O}(\text{CH}_2)_2\text{-OCF}_3$, $\text{O}(\text{CH}_2)_3\text{-N}(\text{R}_4)(\text{R}_5)$, $\text{O}(\text{CH}_2)_2\text{-ON}(\text{R}_4)(\text{R}_5)$, $\text{O}(\text{CH}_2)_2\text{-O}(\text{CH}_2)_2\text{-N}(\text{R}_4)(\text{R}_5)$, $\text{OCH}_2\text{C(=O)-N}(\text{R}_4)(\text{R}_5)$, $\text{OCH}_2\text{C(=O)-N}(\text{R}_6)\text{-(CH}_2)_2\text{-N}(\text{R}_4)(\text{R}_5)$ and $\text{O}(\text{CH}_2)_2\text{-N}(\text{R}_6)\text{-C(=NR}_7)[\text{N}(\text{R}_4)(\text{R}_5)]$ wherein R_4 , R_5 , R_6 and R_7 are each, independently, H or $\text{C}_1\text{-C}_6$ alkyl.

Embodiment 24: The compound of embodiment 23, wherein the 2'-substituent of the at least one 2'-substituted nucleoside of is selected from among: OCH_3 , F, and $\text{O}(\text{CH}_2)_2\text{-OCH}_3$.

Embodiment 25: The compound of embodiment 24, wherein the 2'-substituent of the at least one 2'-substituted nucleoside is $\text{O}(\text{CH}_2)_2\text{-OCH}_3$.

Embodiment 26: The compound of any of embodiments 1-22, wherein the 2'-substituent of each 2'-substituted nucleoside is selected from among: OCH_3 , F, OCH_2F , OCHF_2 , OCF_3 , OCH_2CH_3 , $\text{O}(\text{CH}_2)_2\text{F}$, OCH_2CHF_2 , OCH_2CF_3 , $\text{OCH}_2\text{-CH=CH}_2$, $\text{O}(\text{CH}_2)_2\text{-OCH}_3$, $\text{O}(\text{CH}_2)_2\text{-SCH}_3$, $\text{O}(\text{CH}_2)_2\text{-OCF}_3$, $\text{O}(\text{CH}_2)_3\text{-N}(\text{R}_4)(\text{R}_5)$, $\text{O}(\text{CH}_2)_2\text{-ON}(\text{R}_4)(\text{R}_5)$, $\text{O}(\text{CH}_2)_2\text{-O}(\text{CH}_2)_2\text{-N}(\text{R}_4)(\text{R}_5)$, $\text{OCH}_2\text{C(=O)-N}(\text{R}_4)(\text{R}_5)$, $\text{OCH}_2\text{C(=O)-N}(\text{R}_6)\text{-(CH}_2)_2\text{-N}(\text{R}_4)(\text{R}_5)$ and $\text{O}(\text{CH}_2)_2\text{-N}(\text{R}_6)\text{-C(=NR}_7)[\text{N}(\text{R}_4)(\text{R}_5)]$ wherein R_4 , R_5 , R_6 and R_7 are each, independently, H or $\text{C}_1\text{-C}_6$ alkyl.

Embodiment 27: The compound of embodiment 26, wherein the 2'-substituent of each 2'-substituted nucleoside of is selected from among: OCH_3 , F, and $\text{O}(\text{CH}_2)_2\text{-OCH}_3$.

Embodiment 28: The compound of embodiment 27, wherein the 2'-substituent of each 2'-substituted nucleoside is $\text{O}(\text{CH}_2)_2\text{-OCH}_3$.

Embodiment 29: The compound of any of embodiments 1-28, wherein the 5'-wing does not comprise a bicyclic nucleotide.

Embodiment 30: The compound of any of embodiments 1-29, wherein the 3'-wing does not comprise a bicyclic nucleotide.

Embodiment 31: The compound of any of embodiments 1-30, wherein the target nucleic acid is not a Huntingtin gene transcript.

Embodiment 32: The compound of any of embodiments 1-31, wherein the modified oligonucleotide has a base sequence other than:

GTGCTACCCAACCTTTCTG (SEQ ID NO: 1); CACAGTGCTACCCAACCTT (SEQ ID NO: 2);
CAGTGCTACCCAACC (SEQ ID NO: 3); ATATCACAGTGCTACCCAA (SEQ ID NO: 4);
GATGCTGACTTGGGCCATT (SEQ ID NO: 5); GGGATGCTGACTTGG (SEQ ID NO: 6);
TGCCAAGGGATGCTGACTT (SEQ ID NO: 7); AATTGTCATCACCAGAAAA (SEQ ID NO: 8);
TAAATTGTCATCACC (SEQ ID NO: 9); ACAGTAGATGAGGGAGCAG (SEQ ID NO: 10);
ACACAGTAGATGAGG (SEQ ID NO: 11); AAGTGCACACAGTAGATGA (SEQ ID NO: 12);
AGCTGCAACCTGGCAACAA (SEQ ID NO: 13); GCAGCTGCAACCTGG (SEQ ID NO: 14); or
GCAAGAGCAGCTGCAACCT (SEQ ID NO: 15).

Embodiment 33: The compound of any of embodiments 1-31, wherein the oligonucleotide has a sugar motif other than:

E-K-K-(D)₉-K-K-E;
E-E-E-E-K-(D)₉-E-E-E-E-E;
E-K-K-K-(D)₉-K-K-K-E;
K-E-E-K-(D)₉-K-E-E-K;
K-D-D-K-(D)₉-K-D-D-K;
K-E-K-E-K-(D)₉-K-E-K-E-K;
K-D-K-D-K-(D)₉-K-D-K-D-K;
E-K-E-K-(D)₉-K-E-K-E;
E-E-E-E-E-K-(D)₈-E-E-E-E-E; or
E-K-E-K-E-(D)₉-E-K-E-K-E; wherein

K is a constrained ethyl nucleoside, E is a 2'-MOE substituted nucleoside, and D is a 2'-deoxynucleoside.

Embodiment 34: The compound of any of embodiments 1-30, wherein the 5'-wing consists of 2 linked nucleosides.

Embodiment 35: The compound of any of embodiments 1-30, wherein the 5'-wing consists of 3 linked nucleosides.

Embodiment 36: The compound of any of embodiments 1-30, wherein the 5'-wing consists of 4 linked nucleosides.

- Embodiment 37: The compound of any of embodiments 1-30, wherein the 5'-wing consists of 5 linked nucleosides.
- Embodiment 38: The compound of any of embodiments 1-34, wherein the 3'-wing consists of 2 linked nucleosides.
- Embodiment 39: The compound of any of embodiments 1-34, wherein the 3'-wing consists of 3 linked nucleosides.
- Embodiment 40: The compound of any of embodiments 1-34, wherein the 3'-wing consists of 4 linked nucleosides.
- Embodiment 41: The compound of any of embodiments 1-34, wherein the 3'-wing consists of 5 linked nucleosides.
- Embodiment 42: The compound of any of embodiments 1-38, wherein the gap consists of 6 linked 2'-deoxynucleosides.
- Embodiment 43: The compound of any of embodiments 1-38, wherein the gap consists of 7 linked 2'-deoxynucleosides.
- Embodiment 44: The compound of any of embodiments 1-38, wherein the gap consists of 8 linked 2'-deoxynucleosides.
- Embodiment 45: The compound of any of embodiments 1-38, wherein the gap consists of 9 linked 2'-deoxynucleosides.
- Embodiment 46: The compound of any of embodiments 1-38, wherein the gap consists of 10 linked 2'-deoxynucleosides.
- Embodiment 47: The compound of any of embodiments 1-38, wherein the gap consists of 11 linked 2'-deoxynucleosides.
- Embodiment 48: The compound of any of embodiments 1-38, wherein the gap consists of 12 linked 2'-deoxynucleosides.

- Embodiment 49: The compound of any of embodiments 1-38, wherein the gap consists of 13 linked 2'-deoxynucleosides.
- Embodiment 50: The compound of any of embodiments 1-38, wherein the gap consists of 14 linked 2'-deoxynucleosides.
- Embodiment 51: The compound of any of embodiments 1-50, wherein the oligonucleotide consists of 10 linked nucleosides.
- Embodiment 52: The compound of any of embodiments 1-50, wherein the oligonucleotide consists of 11 linked nucleosides.
- Embodiment 53: The compound of any of embodiments 1-50, wherein the oligonucleotide consists of 12 linked nucleosides.
- Embodiment 54: The compound of any of embodiments 1-50, wherein the oligonucleotide consists of 13 linked nucleosides.
- Embodiment 55: The compound of any of embodiments 1-50, wherein the oligonucleotide consists of 14 linked nucleosides.
- Embodiment 56: The compound of any of embodiments 1-50, wherein the oligonucleotide consists of 15 linked nucleosides.
- Embodiment 57: The compound of any of embodiments 1-50, wherein the oligonucleotide consists of 16 linked nucleosides.
- Embodiment 58: The compound of any of embodiments 1-50, wherein the oligonucleotide consists of 17 linked nucleosides.
- Embodiment 59: The compound of any of embodiments 1-50, wherein the oligonucleotide consists of 18 linked nucleosides.
- Embodiment 60: The compound of any of embodiments 1-50, wherein the oligonucleotide consists of 19 linked nucleosides.

- Embodiment 61: The compound of any of embodiments 1-50, wherein the oligonucleotide consists of 20 linked nucleosides.
- Embodiment 62: The compound of any of embodiments 1-50, wherein the oligonucleotide consists of 21 linked nucleosides.
- Embodiment 63: The compound of any of embodiments 1-50, wherein the oligonucleotide consists of 22 linked nucleosides.
- Embodiment 64: The compound of any of embodiments 1-30, wherein the gapmer motif is selected from among: 2-10-2, 2-10-3, 2-10-4, 2-10-5, 3-10-2, 3-10-3, 3-10-4, 3-10-5, 4-10-2, 4-10-3, 4-10-4, 4-10-5, 5-10-2, 5-10-3, 5-10-4, 5-10-5, 2-9-2, 2-9-3, 2-9-4, 2-9-5, 3-9-2, 3-9-3, 3-9-4, 3-9-5, 4-9-2, 4-9-3, 4-9-4, 4-9-5, 5-9-2, 5-9-3, 5-9-4, 5-9-5, 2-8-2, 2-8-3, 2-8-4, 2-8-5, 3-8-2, 3-8-3, 3-8-4, 3-8-5, 4-8-2, 4-8-3, 4-8-4, 4-8-5, 5-8-2, 5-8-3, 5-8-4, and 5-8-5.
- Embodiment 65: A compound comprising a modified oligonucleotide having a sugar motif selected from among sugar motifs 1-278 as shown in Table 4.
- Embodiment 66: The compound of any of embodiments 1-65, wherein the 5'-wing has a motif selected from among the 5'-wing motifs as shown in Tables 1-3.
- Embodiment 67: The compound of any of embodiments 1-66, wherein the 3'-wing has a motif selected from among the 3'-wing motifs as shown in Tables 4-6.
- Embodiment 68: The compound of any of embodiments 66-67, wherein each A, each B, and each C are independently selected from among: HNA and F-HNA.
- Embodiment 69: The compound of any of embodiments 1-68, wherein the 5'-wing comprises at least one F-HNA.
- Embodiment 70: The compound of any of embodiments 1-69, wherein the 3'-wing comprises at least one F-HNA.
- Embodiment 71: The compound of any of embodiments 1-68, wherein the 5'-wing comprises at least one modified nucleobase.

- Embodiment 72: The compound of any of embodiments 1-69, wherein the 3'-wing comprises at least one modified nucleobase.
- Embodiment 73: The compound of embodiment 72, wherein the modified nucleobase is 2-thio-thymidine.
- Embodiment 74: The compound of any of embodiments 1-73, wherein the 5'-wing has a motif selected from among the 5'-wing motifs as shown in Tables 1-3 and the 3'-wing has a motif selected from among the 3'-wing motifs as shown in Tables 4-6.
- Embodiment 75: The compound of any of embodiments 1-74, wherein the 5'-wing has an ABABA motif, wherein each A is a modified nucleoside and each B comprises a 2'-deoxynucleoside.
- Embodiment 76: The compound of embodiment 75, wherein the modified nucleoside is a bicyclic nucleoside.
- Embodiment 77: The compound of embodiment 76, wherein the bicyclic nucleoside is cEt.
- Embodiment 78: The compound of embodiment 76, wherein the bicyclic nucleoside is LNA.
- Embodiment 79: The compound of any of embodiments 75-78 wherein the 3'-wing has a motif selected from among: AA, AB, AC, BA, BB, BC, CA, CB, and CC.
- Embodiment 80: The compound of embodiment 79, wherein the 3'-wing has an AA motif.
- Embodiment 81: The compound of embodiment 80, wherein A is a 2'-substituted nucleoside.
- Embodiment 82: The compound of embodiment 80, wherein the 2'-substituted nucleoside is selected from among: OCH₃, F, OCH₂F, OCHF₂, OCF₃, OCH₂CH₃, O(CH₂)₂F, OCH₂CHF₂, OCH₂CF₃, OCH₂-CH=CH₂, O(CH₂)₂-OCH₃, O(CH₂)₂-SCH₃, O(CH₂)₂-OCF₃, O(CH₂)₃-N(R₄)(R₅), O(CH₂)₂-ON(R₄)(R₅), O(CH₂)₂-O(CH₂)₂-N(R₄)(R₅), OCH₂C(=O)-N(R₄)(R₅), OCH₂C(=O)-N(R₆)-(CH₂)₂-N(R₄)(R₅) and O(CH₂)₂-N(R₆)-C(=NR₇)[N(R₄)(R₅)] wherein R₄, R₅, R₆ and R₇ are each, independently, H or C₁-C₆ alkyl.
- Embodiment 83: The compound of embodiment 82, wherein the 2'-substituent of each 2'-substituted nucleoside of is selected from among: OCH₃, F, and O(CH₂)₂-OCH₃.

- Embodiment 84: The compound of embodiment 83, wherein the 2'-substituent of each 2'-substituted nucleoside is $\text{O}(\text{CH}_2)_2\text{-OCH}_3$.
- Embodiment 85: The compound of any of embodiments 76-84 wherein the gap between the 5'-wing and the 3'-wing consists of 6 to 11 linked 2'-deoxynucleosides.
- Embodiment 86: The compound of any of embodiments 76-84 wherein the gap between the 5'-wing and the 3'-wing consists of 7 to 10 linked 2'-deoxynucleosides.
- Embodiment 87: The compound of any of embodiments 76-84 wherein the gap between the 5'-wing and the 3'-wing consists of 10 linked 2'-deoxynucleosides.
- Embodiment 88: The compound of any of embodiments 75-87 having the sugar motif: K-D-K-D-K-(D)₆-E-E.
- Embodiment 89: The compound of any of embodiments 75-87 having the sugar motif: K-D-K-D-K-(D)₇-E-E.
- Embodiment 90: The compound of any of embodiments 75-87 having the sugar motif: K-D-K-D-K-(D)₈-E-E.
- Embodiment 91: The compound of any of embodiments 75-87 having the sugar motif: K-D-K-D-K-(D)₉-E-E.
- Embodiment 92: The compound of any of embodiments 75-87 having the sugar motif: K-D-K-D-K-(D)₁₀-E-E.
- Embodiment 93: The compound of any of embodiments 75-87 having the sugar motif: K-D-K-D-K-(D)₁₁-E-E.
- Embodiment 94: The compound of any of embodiments 75-87 having the sugar motif: K-D-K-D-K-(D)₁₂-E-E.
- Embodiment 95: The compound of any of embodiments 75-87 having the sugar motif: K-D-K-D-K-(D)₁₃-E-E.

- Embodiment 96: The compound of any of embodiments 75-87 having the sugar motif: K-D-K-D-K-(D)₁₄-E-E.
- Embodiment 97: The compound of any of embodiments 75-87 having the sugar motif: K-D-K-D-K-(D)₁₅-E-E.
- Embodiment 98: The compound of any of embodiments 1-97, wherein the 5'-wing has a BDBDB motif, wherein each B is a bicyclic nucleoside and each D comprises a 2'-deoxynucleoside.
- Embodiment 99: The compound of any of embodiments 1-97, wherein the 5'-wing has a BDBDB-(D)₆₋₁₅-AA motif, wherein each B is a bicyclic nucleoside and each D comprises a 2'-deoxynucleoside.
- Embodiment 100: The compound of any of embodiments 98-99, wherein B is selected from among: BNA, LNA, α -L-LNA, ENA and 2'-thio LNA.
- Embodiment 101: The compound of embodiment 100, wherein B comprises BNA.
- Embodiment 102: The compound of embodiment 100, wherein B comprises LNA.
- Embodiment 103: The compound of embodiment 100, wherein B comprises α -L-LNA.
- Embodiment 104: The compound of embodiment 100, wherein B comprises ENA.
- Embodiment 105: The compound of embodiment 100, wherein B comprises 2'-thio LNA.
- Embodiment 106: The compound of any of embodiments 100 to 105, wherein A comprises a 2'-substituted nucleoside.
- Embodiment 107: The compound of claim 106, wherein the 2' substituted nucleoside comprises MOE.
- Embodiment 108: The compound of any of embodiments 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: A-B-B-(D)₈-B-B-A, wherein each A is an independently selected 2'-substituted nucleoside, each B is an independently selected bicyclic nucleoside, and each D is a 2'-deoxynucleoside

- Embodiment 109: The compound of any of embodiments 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: A-B-B-(D)₉-B-B-A, wherein each A is an independently selected 2'-substituted nucleoside, each B is an independently selected bicyclic nucleoside, and each D is a 2'-deoxynucleoside
- Embodiment 110: The compound of any of embodiments 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: A-B-B-(D)₁₀-B-B-A, wherein each A is an independently selected 2'-substituted nucleoside, each B is an independently selected bicyclic nucleoside, and each D is a 2'-deoxynucleoside
- Embodiment 111: The compound of any of embodiments 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: A-A-B-(D)₈-B-B-A, wherein each A is an independently selected 2'-substituted nucleoside, each B is an independently selected bicyclic nucleoside, and each D is a 2'-deoxynucleoside
- Embodiment 112: The compound of any of embodiments 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: A-A-B-(D)₉-B-B-A, wherein each A is an independently selected 2'-substituted nucleoside, each B is an independently selected bicyclic nucleoside, and each D is a 2'-deoxynucleoside
- Embodiment 113: The compound of any of embodiments 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: A-A-B-(D)₁₀-B-B-A, wherein each A is an independently selected 2'-substituted nucleoside, each B is an independently selected bicyclic nucleoside, and each D is a 2'-deoxynucleoside
- Embodiment 114: The compound of any of embodiments 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: A-D-B-(D)₈-B-B-A, wherein each A is an independently selected 2'-substituted nucleoside, each B is an independently selected bicyclic nucleoside, and each D is a 2'-deoxynucleoside
- Embodiment 115: The compound of any of embodiments 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: A-D-B-(D)₉-B-B-A, wherein each A is an independently selected 2'-substituted nucleoside, each B is an independently selected bicyclic nucleoside, and each D is a 2'-deoxynucleoside

- Embodiment 116: The compound of any of embodiments 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: A-D-B-(D)₁₀-B-B-A, wherein each A is an independently selected 2'-substituted nucleoside, each B is an independently selected bicyclic nucleoside, and each D is a 2'-deoxynucleoside
- Embodiment 117: The compound of any of embodiments 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: B-D-A-(D)₈-B-B-A, wherein each A is an independently selected 2'-substituted nucleoside, each B is an independently selected bicyclic nucleoside, and each D is a 2'-deoxynucleoside
- Embodiment 118: The compound of any of embodiments 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: B-D-A-(D)₉-B-B-A, wherein each A is an independently selected 2'-substituted nucleoside, each B is an independently selected bicyclic nucleoside, and each D is a 2'-deoxynucleoside
- Embodiment 119: The compound of any of embodiments 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: B-D-A-(D)₁₀-B-B-A, wherein each A is an independently selected 2'-substituted nucleoside, each B is an independently selected bicyclic nucleoside, and each D is a 2'-deoxynucleoside
- Embodiment 120: The compound of any of embodiments 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: B-A-A-(D)₈-B-B-A, wherein each A is an independently selected 2'-substituted nucleoside, each B is an independently selected bicyclic nucleoside, and each D is a 2'-deoxynucleoside
- Embodiment 121: The compound of any of embodiments 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: B-A-A-(D)₉-B-B-A, wherein each A is an independently selected 2'-substituted nucleoside, each B is an independently selected bicyclic nucleoside, and each D is a 2'-deoxynucleoside
- Embodiment 122: The compound of any of embodiments 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: B-A-A-(D)₁₀-B-B-A, wherein each A is an independently selected 2'-substituted nucleoside, each B is an independently selected bicyclic nucleoside, and each D is a 2'-deoxynucleoside

- Embodiment 123: The compound of any of embodiments 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: B-B-B-(D)₈-B-B-A, wherein each A is an independently selected 2'-substituted nucleoside, each B is an independently selected bicyclic nucleoside, and each D is a 2'-deoxynucleoside
- Embodiment 124: The compound of any of embodiments 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: B-B-B-(D)₉-B-B-A, wherein each A is an independently selected 2'-substituted nucleoside, each B is an independently selected bicyclic nucleoside, and each D is a 2'-deoxynucleoside
- Embodiment 125: The compound of any of embodiments 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: B-B-B-(D)₁₀-B-B-A, wherein each A is an independently selected 2'-substituted nucleoside, each B is an independently selected bicyclic nucleoside, and each D is a 2'-deoxynucleoside
- Embodiment 126: The compound of any of embodiments 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: A-A-A-(D)₈-B-B-A, wherein each A is an independently selected 2'-substituted nucleoside, each B is an independently selected bicyclic nucleoside, and each D is a 2'-deoxynucleoside
- Embodiment 127: The compound of any of embodiments 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: A-A-A-(D)₉-B-B-A, wherein each A is an independently selected 2'-substituted nucleoside, each B is an independently selected bicyclic nucleoside, and each D is a 2'-deoxynucleoside
- Embodiment 128: The compound of any of embodiments 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: A-A-A-(D)₁₀-B-B-A, wherein each A is an independently selected 2'-substituted nucleoside, each B is an independently selected bicyclic nucleoside, and each D is a 2'-deoxynucleoside
- Embodiment 129: The compound of any of embodiments 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: B-B-B-(D)₈-B-B-B, wherein each B is an independently selected bicyclic nucleoside, and each D is a 2'-deoxynucleoside

Embodiment 130: The compound of any of embodiments 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: B-B-B-(D)₉-B-B-B, wherein each B is an independently selected bicyclic nucleoside, and each D is a 2'-deoxynucleoside

Embodiment 131: The compound of any of embodiments 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: B-B-B-(D)₁₀-B-B-B, wherein each B is an independently selected bicyclic nucleoside, and each D is a 2'-deoxynucleoside

Embodiment 132: The compound of any of embodiments 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: A-A-A-(D)₈-B-B-B, wherein each A is an independently selected 2'-substituted nucleoside, each B is an independently selected bicyclic nucleoside, and each D is a 2'-deoxynucleoside

Embodiment 133: The compound of any of embodiments 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: A-A-A-(D)₉-B-B-B, wherein each A is an independently selected 2'-substituted nucleoside, each B is an independently selected bicyclic nucleoside, and each D is a 2'-deoxynucleoside

Embodiment 134: The compound of any of embodiments 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: A-A-A-(D)₁₀-B-B-B, wherein each A is an independently selected 2'-substituted nucleoside, each B is an independently selected bicyclic nucleoside, and each D is a 2'-deoxynucleoside

Embodiment 135: The compound of any of embodiments 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: A-D-D-B-(D)₈-B-B-A, wherein each A is an independently selected 2'-substituted nucleoside, each B is an independently selected bicyclic nucleoside, and each D is a 2'-deoxynucleoside

Embodiment 136: The compound of any of embodiments 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: A-D-D-B-(D)₉-B-B-A, wherein each A is an independently selected 2'-substituted nucleoside, each B is an independently selected bicyclic nucleoside, and each D is a 2'-deoxynucleoside

Embodiment 137: The compound of any of embodiments 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: A-D-D-B-(D)₁₀-B-B-A, wherein each A is an

independently selected 2'-substituted nucleoside, each B is an independently selected bicyclic nucleoside, and each D is a 2'-deoxynucleoside

Embodiment 138: The compound of any of embodiments 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: B-D-D-A-(D)₈-B-B-A, wherein each A is an independently selected 2'-substituted nucleoside, each B is an independently selected bicyclic nucleoside, and each D is a 2'-deoxynucleoside

Embodiment 139: The compound of any of embodiments 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: B-D-D-A-(D)₉-B-B-A, wherein each A is an independently selected 2'-substituted nucleoside, each B is an independently selected bicyclic nucleoside, and each D is a 2'-deoxynucleoside

Embodiment 140: The compound of any of embodiments 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: B-D-D-A-(D)₁₀-B-B-A, wherein each A is an independently selected 2'-substituted nucleoside, each B is an independently selected bicyclic nucleoside, and each D is a 2'-deoxynucleoside

Embodiment 141: The compound of any of embodiments 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: A-A-A-B-(D)₈-B-B-A, wherein each A is an independently selected 2'-substituted nucleoside, each B is an independently selected bicyclic nucleoside, and each D is a 2'-deoxynucleoside

Embodiment 142: The compound of any of embodiments 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: A-A-A-B-(D)₉-B-B-A, wherein each A is an independently selected 2'-substituted nucleoside, each B is an independently selected bicyclic nucleoside, and each D is a 2'-deoxynucleoside

Embodiment 143: The compound of any of embodiments 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: A-A-A-B-(D)₁₀-B-B-A, wherein each A is an independently selected 2'-substituted nucleoside, each B is an independently selected bicyclic nucleoside, and each D is a 2'-deoxynucleoside

Embodiment 144: The compound of any of embodiments 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: B-A-A-A-(D)₈-B-B-A, wherein each A is an

independently selected 2'-substituted nucleoside, each B is an independently selected bicyclic nucleoside, and each D is a 2'-deoxynucleoside

Embodiment 145: The compound of any of embodiments 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: B-A-A-A-(D)₉-B-B-A, wherein each A is an independently selected 2'-substituted nucleoside, each B is an independently selected bicyclic nucleoside, and each D is a 2'-deoxynucleoside

Embodiment 146: The compound of any of embodiments 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: B-A-A-A-(D)₁₀-B-B-A, wherein each A is an independently selected 2'-substituted nucleoside, each B is an independently selected bicyclic nucleoside, and each D is a 2'-deoxynucleoside

Embodiment 147: The compound of any of embodiments 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: A-A-A-A-(D)₈-B-B-A, wherein each A is an independently selected 2'-substituted nucleoside, each B is an independently selected bicyclic nucleoside, and each D is a 2'-deoxynucleoside

Embodiment 148: The compound of any of embodiments 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: A-A-A-A-(D)₉-B-B-A, wherein each A is an independently selected 2'-substituted nucleoside, each B is an independently selected bicyclic nucleoside, and each D is a 2'-deoxynucleoside

Embodiment 149: The compound of any of embodiments 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: A-A-A-A-(D)₁₀-B-B-A, wherein each A is an independently selected 2'-substituted nucleoside, each B is an independently selected bicyclic nucleoside, and each D is a 2'-deoxynucleoside

Embodiment 150: The compound of any of embodiments 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: B-A-A-A-(D)₈-B-B-B, wherein each A is an independently selected 2'-substituted nucleoside, each B is an independently selected bicyclic nucleoside, and each D is a 2'-deoxynucleoside

Embodiment 151: The compound of any of embodiments 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: B-A-A-A-(D)₉-B-B-B, wherein each A is an

independently selected 2'-substituted nucleoside, each B is an independently selected bicyclic nucleoside, and each D is a 2'-deoxynucleoside

Embodiment 152: The compound of any of embodiments 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: B-A-A-A-(D)₁₀-B-B-B, wherein each A is an independently selected 2'-substituted nucleoside, each B is an independently selected bicyclic nucleoside, and each D is a 2'-deoxynucleoside

Embodiment 153: The compound of any of embodiments 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: B-D-D-B-(D)₈-B-B-A, wherein each A is an independently selected 2'-substituted nucleoside, each B is an independently selected bicyclic nucleoside, and each D is a 2'-deoxynucleoside

Embodiment 154: The compound of any of embodiments 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: B-D-D-B-(D)₉-B-B-A, wherein each A is an independently selected 2'-substituted nucleoside, each B is an independently selected bicyclic nucleoside, and each D is a 2'-deoxynucleoside

Embodiment 155: The compound of any of embodiments 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: B-D-D-B-(D)₁₀-B-B-A, wherein each A is an independently selected 2'-substituted nucleoside, each B is an independently selected bicyclic nucleoside, and each D is a 2'-deoxynucleoside

Embodiment 156: The compound of any of embodiments 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: A-A-A-A-(D)₈-B-B-B, wherein each A is an independently selected 2'-substituted nucleoside, each B is an independently selected bicyclic nucleoside, and each D is a 2'-deoxynucleoside

Embodiment 157: The compound of any of embodiments 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: A-A-A-A-(D)₉-B-B-B, wherein each A is an independently selected 2'-substituted nucleoside, each B is an independently selected bicyclic nucleoside, and each D is a 2'-deoxynucleoside

Embodiment 158: The compound of any of embodiments 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: A-A-A-A-(D)₁₀-B-B-B, wherein each A is an

independently selected 2'-substituted nucleoside, each B is an independently selected bicyclic nucleoside, and each D is a 2'-deoxynucleoside

Embodiment 159: The compound of any of embodiments 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: B-B-B-B-(D)₈-B-B-B, wherein each B is an independently selected bicyclic nucleoside, and each D is a 2'-deoxynucleoside

Embodiment 160: The compound of any of embodiments 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: B-B-B-B-(D)₉-B-B-B, each B is an independently selected bicyclic nucleoside, and each D is a 2'-deoxynucleoside

Embodiment 161: The compound of any of embodiments 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: B-B-B-B-(D)₁₀-B-B-B, wherein each B is an independently selected bicyclic nucleoside, and each D is a 2'-deoxynucleoside

Embodiment 162: The compound of any of embodiments 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: A-A-A-A-A-(D)₈-B-B-A, wherein each A is an independently selected 2'-substituted nucleoside, each B is an independently selected bicyclic nucleoside, and each D is a 2'-deoxynucleoside

Embodiment 163: The compound of any of embodiments 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: A-A-A-A-A-(D)₉-B-B-A, wherein each A is an independently selected 2'-substituted nucleoside, each B is an independently selected bicyclic nucleoside, and each D is a 2'-deoxynucleoside

Embodiment 164: The compound of any of embodiments 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: A-A-A-A-A-(D)₁₀-B-B-A, wherein each A is an independently selected 2'-substituted nucleoside, each B is an independently selected bicyclic nucleoside, and each D is a 2'-deoxynucleoside

Embodiment 165: The compound of any of embodiments 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: A-A-A-A-A-(D)₈-B-B-B, wherein each A is an independently selected 2'-substituted nucleoside, each B is an independently selected bicyclic nucleoside, and each D is a 2'-deoxynucleoside

- Embodiment 166: The compound of any of embodiments 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: A-A-A-A-A-(D)₉-B-B-B, wherein each A is an independently selected 2'-substituted nucleoside, each B is an independently selected bicyclic nucleoside, and each D is a 2'-deoxynucleoside
- Embodiment 167: The compound of any of embodiments 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: A-A-A-A-A-(D)₁₀-B-B-B, wherein each A is an independently selected 2'-substituted nucleoside, each B is an independently selected bicyclic nucleoside, and each D is a 2'-deoxynucleoside
- Embodiment 168: The compound of any of embodiments 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: A-D-A-D-B-(D)₈-B-B-A, wherein each A is an independently selected 2'-substituted nucleoside, each B is an independently selected bicyclic nucleoside, and each D is a 2'-deoxynucleoside
- Embodiment 169: The compound of any of embodiments 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: A-D-A-D-B-(D)₉-B-B-A, wherein each A is an independently selected 2'-substituted nucleoside, each B is an independently selected bicyclic nucleoside, and each D is a 2'-deoxynucleoside
- Embodiment 170: The compound of any of embodiments 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: A-D-A-D-B-(D)₁₀-B-B-A, wherein each A is an independently selected 2'-substituted nucleoside, each B is an independently selected bicyclic nucleoside, and each D is a 2'-deoxynucleoside
- Embodiment 171: The compound of any of embodiments 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: A-D-B-D-A-(D)₈-B-B-A, wherein each A is an independently selected 2'-substituted nucleoside, each B is an independently selected bicyclic nucleoside, and each D is a 2'-deoxynucleoside
- Embodiment 172: The compound of any of embodiments 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: A-D-B-D-A-(D)₉-B-B-A, wherein each A is an independently selected 2'-substituted nucleoside, each B is an independently selected bicyclic nucleoside, and each D is a 2'-deoxynucleoside

- Embodiment 173: The compound of any of embodiments 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: A-D-B-D-A-(D)₁₀-B-B-A, wherein each A is an independently selected 2'-substituted nucleoside, each B is an independently selected bicyclic nucleoside, and each D is a 2'-deoxynucleoside
- Embodiment 174: The compound of any of embodiments 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: B-D-A-D-A-(D)₈-B-B-A, wherein each A is an independently selected 2'-substituted nucleoside, each B is an independently selected bicyclic nucleoside, and each D is a 2'-deoxynucleoside
- Embodiment 175: The compound of any of embodiments 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: B-D-A-D-A-(D)₉-B-B-A, wherein each A is an independently selected 2'-substituted nucleoside, each B is an independently selected bicyclic nucleoside, and each D is a 2'-deoxynucleoside
- Embodiment 176: The compound of any of embodiments 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: B-D-A-D-A-(D)₁₀-B-B-A, wherein each A is an independently selected 2'-substituted nucleoside, each B is an independently selected bicyclic nucleoside, and each D is a 2'-deoxynucleoside
- Embodiment 177: The compound of any of embodiments 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: A-A-A-A-B-(D)₈-B-B-A, wherein each A is an independently selected 2'-substituted nucleoside, each B is an independently selected bicyclic nucleoside, and each D is a 2'-deoxynucleoside
- Embodiment 178: The compound of any of embodiments 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: A-A-A-A-B-(D)₉-B-B-A, wherein each A is an independently selected 2'-substituted nucleoside, each B is an independently selected bicyclic nucleoside, and each D is a 2'-deoxynucleoside
- Embodiment 179: The compound of any of embodiments 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: A-A-A-A-B-(D)₁₀-B-B-A, wherein each A is an independently selected 2'-substituted nucleoside, each B is an independently selected bicyclic nucleoside, and each D is a 2'-deoxynucleoside

- Embodiment 180: The compound of any of embodiments 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: A-A-B-A-A-(D)₈-B-B-A, wherein each A is an independently selected 2'-substituted nucleoside, each B is an independently selected bicyclic nucleoside, and each D is a 2'-deoxynucleoside
- Embodiment 181: The compound of any of embodiments 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: A-A-B-A-A-(D)₉-B-B-A, wherein each A is an independently selected 2'-substituted nucleoside, each B is an independently selected bicyclic nucleoside, and each D is a 2'-deoxynucleoside
- Embodiment 182: The compound of any of embodiments 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: A-A-B-A-A-(D)₁₀-B-B-A, wherein each A is an independently selected 2'-substituted nucleoside, each B is an independently selected bicyclic nucleoside, and each D is a 2'-deoxynucleoside
- Embodiment 183: The compound of any of embodiments 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: B-A-A-A-A-(D)₈-B-B-A, wherein each A is an independently selected 2'-substituted nucleoside, each B is an independently selected bicyclic nucleoside, and each D is a 2'-deoxynucleoside
- Embodiment 184: The compound of any of embodiments 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: B-A-A-A-A-(D)₉-B-B-A, wherein each A is an independently selected 2'-substituted nucleoside, each B is an independently selected bicyclic nucleoside, and each D is a 2'-deoxynucleoside
- Embodiment 185: The compound of any of embodiments 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: B-A-A-A-A-(D)₁₀-B-B-A, wherein each A is an independently selected 2'-substituted nucleoside, each B is an independently selected bicyclic nucleoside, and each D is a 2'-deoxynucleoside
- Embodiment 186: The compound of any of embodiments 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: e-e-k-k-(D)₉-e-k-e-e, wherein each k comprises a bicyclic nucleoside, each e comprises a 2'-modified nucleoside, and each D comprises a 2'-deoxynucleoside.

Embodiment 187: The compound of any of embodiments 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: k-d-k-d-k-(D)₁₀-e-e-e-e, wherein each k comprises a bicyclic nucleoside, each e comprises a 2'-modified nucleoside, and each D comprises a 2'-deoxynucleoside.

Embodiment 188: The compound of any of embodiments 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: k-e-k-(D)₁₀-k-e-k, wherein each k comprises a bicyclic nucleoside, each e comprises a 2'-modified nucleoside, and each D comprises a 2'-deoxynucleoside.

Embodiment 189: The compound of any of embodiments 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: k-d-k-d-k-(D)₈-e-e-e-e, wherein each k comprises a bicyclic nucleoside, each e comprises a 2'-modified nucleoside, and each D comprises a 2'-deoxynucleoside.

Embodiment 190: The compound of any of embodiments 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: k-d-k-d-k-(D)₈-e-e-e-e-e, wherein each k comprises a bicyclic nucleoside, each e comprises a 2'-modified nucleoside, and each D comprises a 2'-deoxynucleoside.

Embodiment 191: The compound of any of embodiments 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: k-e-k-(D)₈-e-e-e-e-e, wherein each k comprises a bicyclic nucleoside, each e comprises a 2'-modified nucleoside, and each D comprises a 2'-deoxynucleoside.

Embodiment 192: The compound of any of embodiments 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: e-k-(D)₁₀-k-e-k-e, wherein each k comprises a bicyclic nucleoside, each e comprises a 2'-modified nucleoside, and each D comprises a 2'-deoxynucleoside.

Embodiment 193: The compound of any of embodiments 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: e-e-e-(D)₁₀-k-k-k, wherein each k comprises a bicyclic nucleoside, each e comprises a 2'-modified nucleoside, and each D comprises a 2'-deoxynucleoside.

Embodiment 194: The compound of any of embodiments 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: e-e-e-k-k-(D)₈-e-e-e-e, wherein each k comprises a bicyclic nucleoside, each e comprises a 2'-modified nucleoside, and each D comprises a 2'-deoxynucleoside.

- Embodiment 195: The compound of any of embodiments 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: e-e-e-k-k-(D)₇-k-k-e-e-e, wherein each k comprises a bicyclic nucleoside, each e comprises a 2'-modified nucleoside, and each D comprises a 2'-deoxynucleoside.
- Embodiment 196: The compound of any of embodiments 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: e-e-e-k-(D)₉-k-e-e-e, wherein each k comprises a bicyclic nucleoside, each e comprises a 2'-modified nucleoside, and each D comprises a 2'-deoxynucleoside.
- Embodiment 197: The compound of any of embodiments 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: e-e-e-k-k-(D)₇-k-k-e-e-e, wherein each k comprises a bicyclic nucleoside, each e comprises a 2'-modified nucleoside, and each D comprises a 2'-deoxynucleoside.
- Embodiment 198: The compound of any of embodiments 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: e-e-e-k-k-(D)₇-k-k-e-e-e, wherein each k comprises a bicyclic nucleoside, each e comprises a 2'-modified nucleoside, and each D comprises a 2'-deoxynucleoside.
- Embodiment 199: The compound of any of embodiments 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: e-e-e-e-k-k-(D)₇-e-e-e-e, wherein each k comprises a bicyclic nucleoside, each e comprises a 2'-modified nucleoside, and each D comprises a 2'-deoxynucleoside.
- Embodiment 200: The compound of any of embodiments 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: e-k-e-k-(D)₉-e-e-e-e, wherein each k comprises a bicyclic nucleoside, each e comprises a 2'-modified nucleoside, and each D comprises a 2'-deoxynucleoside.
- Embodiment 201: The compound of any of embodiments 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: e-k-e-k-d-k-(D)₇-e-e-e-e, wherein each k comprises a bicyclic nucleoside, each e comprises a 2'-modified nucleoside, and each D comprises a 2'-deoxynucleoside.

- Embodiment 202: The compound of any of embodiments 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: e-e-k-k-(D)₇-k-k-e-e-e, wherein each k comprises a bicyclic nucleoside, each e comprises a 2'-modified nucleoside, and each D comprises a 2'-deoxynucleoside.
- Embodiment 203: The compound of any of embodiments 186 to 202, wherein each k comprises a cEt nucleoside.
- Embodiment 204: The compound of any of embodiments 186 to 202, wherein each k comprises an LNA nucleoside.
- Embodiment 205: The compound of any of embodiments 186 to 203, wherein each e comprises a 2'-MOE modified nucleoside.
- Embodiment 206: The compound of any of embodiments 186 to 203, wherein each e comprises a 2'-OMe modified nucleoside.
- Embodiment 207: The compound of any of embodiments 186 to 202, wherein each k comprises a cEt nucleoside and each e comprises a 2'-MOE modified nucleoside.
- Embodiment 208: The compound of any of embodiments 89-202, wherein at least one bicyclic nucleoside is a constrained ethyl nucleoside.
- Embodiment 209: The compound of any of embodiments 89-202, wherein each bicyclic nucleoside is a constrained ethyl nucleoside.
- Embodiment 210: The compound of any of embodiments, 89-202, wherein at least one bicyclic nucleoside is selected from among: BNA, LNA, α -L-LNA, ENA and 2'-thio LNA.
- Embodiment 211: The compound of any of embodiments, 89-202, wherein at least one bicyclic nucleoside is an LNA nucleoside.
- Embodiment 212: The compound of any of embodiments 89-202, wherein each bicyclic nucleoside is an LNA nucleoside.

Embodiment 213: The compound of any of embodiments 89-202, wherein the 2'-substituent of the at least one 2'-substituted nucleoside is selected from among: OCH_3 , F, OCH_2F , OCHF_2 , OCF_3 , OCH_2CH_3 , $\text{O}(\text{CH}_2)_2\text{F}$, OCH_2CHF_2 , OCH_2CF_3 , $\text{OCH}_2\text{-CH=CH}_2$, $\text{O}(\text{CH}_2)_2\text{-OCH}_3$, $\text{O}(\text{CH}_2)_2\text{-SCH}_3$, $\text{O}(\text{CH}_2)_2\text{-OCF}_3$, $\text{O}(\text{CH}_2)_3\text{-N}(\text{R}_4)(\text{R}_5)$, $\text{O}(\text{CH}_2)_2\text{-ON}(\text{R}_4)(\text{R}_5)$, $\text{O}(\text{CH}_2)_2\text{-O}(\text{CH}_2)_2\text{-N}(\text{R}_4)(\text{R}_5)$, $\text{OCH}_2\text{C(=O)-N}(\text{R}_4)(\text{R}_5)$, $\text{OCH}_2\text{C(=O)-N}(\text{R}_6)\text{-(CH}_2)_2\text{-N}(\text{R}_4)(\text{R}_5)$ and $\text{O}(\text{CH}_2)_2\text{-N}(\text{R}_6)\text{-C(=NR}_7)[\text{N}(\text{R}_4)(\text{R}_5)]$ wherein R_4 , R_5 , R_6 and R_7 are each, independently, H or $\text{C}_1\text{-C}_6$ alkyl.

Embodiment 214: The compound of embodiment 213, wherein the 2'-substituent of the at least one 2'-substituted nucleoside of is selected from among: OCH_3 , F, and $\text{O}(\text{CH}_2)_2\text{-OCH}_3$.

Embodiment 215: The compound of embodiment 214, wherein the 2'-substituent of the at least one 2'-substituted nucleoside is $\text{O}(\text{CH}_2)_2\text{-OCH}_3$.

Embodiment 216: The compound of any of embodiments 89-202, wherein the 2'-substituent of each 2'-substituted nucleoside is selected from among: OCH_3 , F, OCH_2F , OCHF_2 , OCF_3 , OCH_2CH_3 , $\text{O}(\text{CH}_2)_2\text{F}$, OCH_2CHF_2 , OCH_2CF_3 , $\text{OCH}_2\text{-CH=CH}_2$, $\text{O}(\text{CH}_2)_2\text{-OCH}_3$, $\text{O}(\text{CH}_2)_2\text{-SCH}_3$, $\text{O}(\text{CH}_2)_2\text{-OCF}_3$, $\text{O}(\text{CH}_2)_3\text{-N}(\text{R}_4)(\text{R}_5)$, $\text{O}(\text{CH}_2)_2\text{-ON}(\text{R}_4)(\text{R}_5)$, $\text{O}(\text{CH}_2)_2\text{-O}(\text{CH}_2)_2\text{-N}(\text{R}_4)(\text{R}_5)$, $\text{OCH}_2\text{C(=O)-N}(\text{R}_4)(\text{R}_5)$, $\text{OCH}_2\text{C(=O)-N}(\text{R}_6)\text{-(CH}_2)_2\text{-N}(\text{R}_4)(\text{R}_5)$ and $\text{O}(\text{CH}_2)_2\text{-N}(\text{R}_6)\text{-C(=NR}_7)[\text{N}(\text{R}_4)(\text{R}_5)]$ wherein R_4 , R_5 , R_6 and R_7 are each, independently, H or $\text{C}_1\text{-C}_6$ alkyl.

Embodiment 217: The compound of embodiment 216, wherein the 2'-substituent of each 2'-substituted nucleoside of is selected from among: OCH_3 , F, and $\text{O}(\text{CH}_2)_2\text{-OCH}_3$.

Embodiment 218: The compound of embodiment 217, wherein the 2'-substituent of each 2'-substituted nucleoside is $\text{O}(\text{CH}_2)_2\text{-OCH}_3$.

Embodiment 219: The compound of any of embodiments 1-218, wherein the oligonucleotide comprises at least one modified internucleoside linkage.

Embodiment 220: The compound of embodiment 219, wherein each internucleoside linkage is a modified internucleoside linkage.

Embodiment 221: The compound of embodiment 219 or 220, wherein the modified internucleoside linkage is a phosphorothioate linkage.

- Embodiment 222: The compound of embodiment 219 or 220, wherein the modified internucleoside linkage is a methylphosphonate.
- Embodiment 223: The compound of any of embodiments 1-222 comprising a conjugate.
- Embodiment 224: The compound of any of embodiments 1-223 comprising at least one 5-methyl cytosine nucleobase.
- Embodiment 225: The compound of any of embodiments 1-224 comprising at least one modified nucleobase.
- Embodiment 226: The compound of any of embodiments 1-225, wherein the compound is an antisense compound.
- Embodiment 227: The compound of embodiment 226, wherein the compound is capable of inhibiting expression of the target nucleic acid in a cell.
- Embodiment 228: The compound of embodiment 227, wherein the compound is capable of inhibiting expression of the target nucleic acid in a cell by at least 50%.
- Embodiment 229: The compound of embodiment 227, wherein the compound is capable of inhibiting expression of the target nucleic acid in a cell by at least 80%.
- Embodiment 230: The compound of any of embodiments 227-229, wherein the cell is in an animal.
- Embodiment 231: The compound of embodiment 230, wherein the animal is a human.
- Embodiment 232: The compound of any of embodiments 1 to 231, wherein bicyclic nucleoside is selected from among: BNA, LNA, α -L-LNA, ENA and 2'-thio LNA.
- Embodiment 233: A compound of any of embodiments 1-232, comprising not more than 6 bicyclic nucleosides.
- Embodiment 234: A compound of any of embodiments 1-232, comprising not more than 5 bicyclic nucleosides.

- Embodiment 235: A compound of any of embodiments 1-232, comprising not more than 4 bicyclic nucleosides.
- Embodiment 236: A compound of any of embodiments 1-232, comprising not more than 3 bicyclic nucleosides.
- Embodiment 237: A compound of any of embodiments 1-232, comprising not more than 2 bicyclic nucleosides.
- Embodiment 238: A compound of any of embodiments 1-232, comprising not more than 1 bicyclic nucleoside.
- Embodiment 239: The compound of any of embodiments 233-238, wherein the bicyclic nucleoside comprises cEt.
- Embodiment 240: The compound of any of embodiments 233-238, wherein the bicyclic nucleoside comprises LNA.
- Embodiment 241: A pharmaceutical composition comprising the compound according to any of embodiments 1-240 and a pharmaceutically acceptable diluent.
- Embodiment 242: A method of modulating expression of a target nucleic acid in a cell comprising contacting the cell with a compound according to any of embodiments 1-240.
- Embodiment 243: A method of modulating expression of a target nucleic acid in an animal comprising administering to the animal the pharmaceutical composition according to embodiment 242.
- Embodiment 244: A method of manufacturing a compound according to any of embodiments 1-241 comprising forming chemical bonds.
- Embodiment 245: The method of embodiment 244, wherein said chemical bonds are internucleoside linkages.
- Embodiment 246: The method embodiment 244 or 245, wherein the method is performed under conditions suitable for the preparation of products for administration to humans.

Embodiment 247: A method of manufacturing the pharmaceutical composition according to embodiment 246 comprising combining the compound according to any of embodiments 1-241 and the pharmaceutically acceptable diluent.

Embodiment 248: The method embodiment 247, wherein the method is performed under conditions suitable for the preparation of products for administration to humans.

Embodiment 249: A compound comprising a modified oligonucleotide having a sugar motif selected from among sugar motifs 279-615 as shown in Table 4.

Embodiment 250: A compound comprising:
a modified oligonucleotide consisting of 10 to 20 linked nucleosides, wherein the modified oligonucleotide comprises:

a 5'-wing consisting of 2 to 5 linked nucleosides;

a 3'-wing consisting of 2 to 5 linked nucleosides; and

a gap between the 5'-wing and the 3'-wing consisting of 6 to 14 linked 2'-deoxynucleosides; and

wherein the 5'-wing has a sugar modification motif selected from among the motifs in Table 1.

Embodiment 251: A compound comprising:
a modified oligonucleotide consisting of 10 to 20 linked nucleosides, wherein the modified oligonucleotide comprises:

a 5'-wing consisting of 2 to 5 linked nucleosides;

a 3'-wing consisting of 2 to 5 linked nucleosides; and

a gap between the 5'-wing and the 3'-wing consisting of 6 to 14 linked 2'-deoxynucleosides; and

wherein the 3'-wing has a sugar modification motif selected from among the motifs in Table 2.


Embodiment 252: A compound comprising:
a modified oligonucleotide consisting of 10 to 20 linked nucleosides, wherein the modified oligonucleotide comprises:

a 5'-wing consisting of 2 to 5 linked nucleosides;

a 3'-wing consisting of 2 to 5 linked nucleosides; and

a gap between the 5'-wing and the 3'-wing consisting of 6 to 14 linked 2'-deoxynucleosides; and wherein the 5'-wing has a sugar modification motif selected from among the motifs in Table 1 and the 3'-wing has a sugar modification motif selected from among the motifs in Table 2.

Embodiment 253: A compound of any of embodiments 1-16, wherein the modified oligonucleotide has a sugar motif described by Formula II as follows:



wherein:

each A is independently a 2'-substituted nucleoside;

each B is independently a bicyclic nucleoside;

each J is independently either a 2'-substituted nucleoside or a 2'-deoxynucleoside;

each D is a 2'-deoxynucleoside;

m is 0-4; n is 0-2; p is 0-2; r is 0-2; t is 0-2; v is 0-2; w is 0-4; x is 0-2; y is 0-2; z is 0-4; g is 6-14;

provided that:

at least one of m, n, and r is other than 0;


at least one of w and y is other than 0;

the sum of m, n, p, r, and t is from 1 to 5; and

the sum of v, w, x, y, and z is from 1 to 5.

Embodiment 254: A compound comprising:

a modified oligonucleotide consisting of 10 to 20 linked nucleosides, wherein the modified oligonucleotide has a sugar motif described by Formula II as follows:



wherein:

each A is independently a 2'-substituted nucleoside;

each B is independently a bicyclic nucleoside;

each J is independently either a 2'-substituted nucleoside or a 2'-deoxynucleoside;

each D is a 2'-deoxynucleoside;

m is 0-4; n is 0-2; p is 0-2; r is 0-2; t is 0-2; v is 0-2; w is 0-4; x is 0-2; y is 0-2; z is 0-4; g is 6-14;

provided that:

at least one of m, n, and r is other than 0;

at least one of w and y is other than 0;

the sum of m, n, p, r, and t is from 1 to 5; and

the sum of v, w, x, y, and z is from 1 to 5.

Embodiment 255: The compound of embodiment 253 or 254, wherein at least one bicyclic nucleoside is a constrained ethyl nucleoside.

Embodiment 256: The compound of embodiment 255, wherein each bicyclic nucleoside is a constrained ethyl nucleoside.

Embodiment 257: The compound of any of embodiments 253 -254, wherein at least one bicyclic nucleoside is an LNA nucleoside.

Embodiment 258: The compound of embodiments 250-254, wherein each bicyclic nucleoside is an LNA nucleoside.

Embodiment 259: A method of treating a disease or condition.

Embodiment 260: Use of a compound of any of embodiments 1 to 259 for the preparation of a medicament for the treatment of a disease or condition.

Embodiment 261: The use of embodiment 260, wherein the disease or condition is associated with a virus.

In certain embodiments, including but not limited to any of the above numbered embodiments, compounds including oligonucleotides described herein are capable of modulating expression of a target mRNA. In certain embodiments, the target mRNA is associated with a disease or disorder, or encodes a protein that is associated with a disease or disorder. In certain embodiments, the compounds or oligonucleotides provided herein modulate the expression of function of such mRNA to alleviate one or more symptom of the disease or disorder.

In certain embodiments, compounds including oligonucleotides describe herein are useful in vitro. In certain embodiments such compounds are used in diagnostics and/or for target validation experiments.

DETAILED DESCRIPTION OF THE INVENTION

Unless specific definitions are provided, the nomenclature used in connection with, and the procedures and techniques of, analytical chemistry, synthetic organic chemistry, and medicinal and pharmaceutical chemistry described herein are those well known and commonly used in the art. Standard techniques may be used for chemical synthesis, and chemical analysis. Certain such techniques and procedures may be found for example in "Carbohydrate Modifications in Antisense Research" Edited by Sangvi and Cook, American Chemical Society, Washington D.C., 1994; "Remington's Pharmaceutical Sciences," Mack Publishing Co., Easton, Pa., 21st edition, 2005; and "Antisense Drug Technology, Principles,

Strategies, and Applications” Edited by Stanley T. Crooke, CRC Press, Boca Raton, Florida; and Sambrook et al., “Molecular Cloning, A laboratory Manual,” 2nd Edition, Cold Spring Harbor Laboratory Press, 1989, which are hereby incorporated by reference for any purpose. Where permitted, all patents, applications, published applications and other publications and other data referred to throughout in the disclosure are incorporated by reference herein in their entirety.

Unless otherwise indicated, the following terms have the following meanings:

As used herein, “nucleoside” means a compound comprising a nucleobase moiety and a sugar moiety. Nucleosides include, but are not limited to, naturally occurring nucleosides (as found in DNA and RNA) and modified nucleosides. Nucleosides may be linked to a phosphate moiety.

As used herein, “chemical modification” means a chemical difference in a compound when compared to a naturally occurring counterpart. In reference to an oligonucleotide, chemical modification does not include differences only in nucleobase sequence. Chemical modifications of oligonucleotides include nucleoside modifications (including sugar moiety modifications and nucleobase modifications) and internucleoside linkage modifications.

As used herein, “furanosyl” means a structure comprising a 5-membered ring comprising four carbon atoms and one oxygen atom.

As used herein, “naturally occurring sugar moiety” means a ribofuranosyl as found in naturally occurring RNA or a deoxyribofuranosyl as found in naturally occurring DNA.

As used herein, “sugar moiety” means a naturally occurring sugar moiety or a modified sugar moiety of a nucleoside.

As used herein, “modified sugar moiety” means a substituted sugar moiety or a sugar surrogate.

As used herein, “substituted sugar moiety” means a furanosyl that is not a naturally occurring sugar moiety. Substituted sugar moieties include, but are not limited to furanosyls comprising substituents at the 2'-position, the 3'-position, the 5'-position and/or the 4'-position. Certain substituted sugar moieties are bicyclic sugar moieties.

As used herein, “2'-substituted sugar moiety” means a furanosyl comprising a substituent at the 2'-position other than H or OH. Unless otherwise indicated, a 2'-substituted sugar moiety is not a bicyclic sugar moiety (i.e., the 2'-substituent of a 2'-substituted sugar moiety does not form a bridge to another atom of the furanosyl ring).

As used herein, “MOE” means $\text{-OCH}_2\text{CH}_2\text{OCH}_3$.

As used herein the term "sugar surrogate" means a structure that does not comprise a furanosyl and that is capable of replacing the naturally occurring sugar moiety of a nucleoside, such that the resulting nucleoside is capable of (1) incorporation into an oligonucleotide and (2) hybridization to a complementary nucleoside. Such structures include rings comprising a different number of atoms than furanosyl (e.g., 4, 6, or 7-membered rings); replacement of the oxygen of a furanosyl with a non-oxygen atom (e.g., carbon, sulfur, or nitrogen); or both a change in the number of atoms and a replacement of the oxygen. Such structures may

also comprise substitutions corresponding to those described for substituted sugar moieties (e.g., 6-membered carbocyclic bicyclic sugar surrogates optionally comprising additional substituents). Sugar surrogates also include more complex sugar replacements (e.g., the non-ring systems of peptide nucleic acid). Sugar surrogates include without limitation morpholinos, cyclohexenyls and cyclohexitols.

As used herein, “bicyclic sugar moiety” means a modified sugar moiety comprising a 4 to 7 membered ring (including but not limited to a furanosyl) comprising a bridge connecting two atoms of the 4 to 7 membered ring to form a second ring, resulting in a bicyclic structure. In certain embodiments, the 4 to 7 membered ring is a sugar ring. In certain embodiments the 4 to 7 membered ring is a furanosyl. In certain such embodiments, the bridge connects the 2'-carbon and the 4'-carbon of the furanosyl.

As used herein, “nucleotide” means a nucleoside further comprising a phosphate linking group. As used herein, “linked nucleosides” may or may not be linked by phosphate linkages and thus includes, but is not limited to “linked nucleotides.” As used herein, “linked nucleosides” are nucleosides that are connected in a continuous sequence (i.e. no additional nucleosides are present between those that are linked).

As used herein, “nucleobase” means a group of atoms that can be linked to a sugar moiety to create a nucleoside that is capable of incorporation into an oligonucleotide, and wherein the group of atoms is capable of bonding with a complementary naturally occurring nucleobase of another oligonucleotide or nucleic acid. Nucleobases may be naturally occurring or may be modified.

As used herein, “heterocyclic base” or “heterocyclic nucleobase” means a nucleobase comprising a heterocyclic structure.

As used herein the terms, “unmodified nucleobase” or “naturally occurring nucleobase” means the naturally occurring heterocyclic nucleobases of RNA or DNA: the purine bases adenine (A) and guanine (G), and the pyrimidine bases thymine (T), cytosine (C) (including 5-methyl C), and uracil (U).

As used herein, “modified nucleobase” means any nucleobase that is not a naturally occurring nucleobase.

As used herein, “modified nucleoside” means a nucleoside comprising at least one chemical modification compared to naturally occurring RNA or DNA nucleosides. Modified nucleosides comprise a modified sugar moiety and/or a modified nucleobase.

As used herein, “bicyclic nucleoside” or “BNA” means a nucleoside comprising a bicyclic sugar moiety.

As used herein, “constrained ethyl nucleoside” or “cEt” means a nucleoside comprising a bicyclic sugar moiety comprising a 4'-CH(CH₃)-O-2' bridge.

As used herein, “locked nucleic acid nucleoside” or “LNA” means a nucleoside comprising a bicyclic sugar moiety comprising a 4'-CH₂-O-2' bridge.

As used herein, “2'-substituted nucleoside” means a nucleoside comprising a substituent at the 2'-position other than H or OH. Unless otherwise indicated, a 2'-substituted nucleoside is not a bicyclic nucleoside.

As used herein, "2'-deoxynucleoside" means a nucleoside comprising 2'-H furanosyl sugar moiety, as found in naturally occurring deoxyribonucleosides (DNA). In certain embodiments, a 2'-deoxynucleoside may comprise a modified nucleobase or may comprise an RNA nucleobase (e.g., uracil).

As used herein, "oligonucleotide" means a compound comprising a plurality of linked nucleosides. In certain embodiments, an oligonucleotide comprises one or more unmodified ribonucleosides (RNA) and/or unmodified deoxyribonucleosides (DNA) and/or one or more modified nucleosides.

As used herein "oligonucleoside" means an oligonucleotide in which none of the internucleoside linkages contains a phosphorus atom. As used herein, oligonucleotides include oligonucleosides.

As used herein, "modified oligonucleotide" means an oligonucleotide comprising at least one modified nucleoside and/or at least one modified internucleoside linkage.

As used herein "internucleoside linkage" means a covalent linkage between adjacent nucleosides in an oligonucleotide.

As used herein "naturally occurring internucleoside linkage" means a 3' to 5' phosphodiester linkage.

As used herein, "modified internucleoside linkage" means any internucleoside linkage other than a naturally occurring internucleoside linkage.

As used herein, "oligomeric compound" means a polymeric structure comprising two or more sub-structures. In certain embodiments, an oligomeric compound comprises an oligonucleotide. In certain embodiments, an oligomeric compound comprises one or more conjugate groups and/or terminal groups. In certain embodiments, an oligomeric compound consists of an oligonucleotide.

As used herein, "terminal group" means one or more atom attached to either, or both, the 3' end or the 5' end of an oligonucleotide. In certain embodiments a terminal group is a conjugate group. In certain embodiments, a terminal group comprises one or more terminal group nucleosides.

As used herein, "conjugate" means an atom or group of atoms bound to an oligonucleotide or oligomeric compound. In general, conjugate groups modify one or more properties of the compound to which they are attached, including, but not limited to pharmacodynamic, pharmacokinetic, binding, absorption, cellular distribution, cellular uptake, charge and/or clearance properties.

As used herein, "conjugate linking group" means any atom or group of atoms used to attach a conjugate to an oligonucleotide or oligomeric compound.

As used herein, "antisense compound" means a compound comprising or consisting of an oligonucleotide at least a portion of which is complementary to a target nucleic acid to which it is capable of hybridizing, resulting in at least one antisense activity.

As used herein, "antisense activity" means any detectable and/or measurable change attributable to the hybridization of an antisense compound to its target nucleic acid.

As used herein, "detecting" or "measuring" means that a test or assay for detecting or measuring is performed. Such detection and/or measuring may result in a value of zero. Thus, if a test for detection or measuring results in a finding of no activity (activity of zero), the step of detecting or measuring the activity

has nevertheless been performed.

As used herein, “detectable and/or measureable activity” means a measurable activity that is not zero.

As used herein, “essentially unchanged” means little or no change in a particular parameter, particularly relative to another parameter which changes much more. In certain embodiments, a parameter is essentially unchanged when it changes less than 5%. In certain embodiments, a parameter is essentially unchanged if it changes less than two-fold while another parameter changes at least ten-fold. For example, in certain embodiments, an antisense activity is a change in the amount of a target nucleic acid. In certain such embodiments, the amount of a non-target nucleic acid is essentially unchanged if it changes much less than the target nucleic acid does, but the change need not be zero.

As used herein, “expression” means the process by which a gene ultimately results in a protein. Expression includes, but is not limited to, transcription, post-transcriptional modification (e.g., splicing, polyadenylation, addition of 5’-cap), and translation.

As used herein, “target nucleic acid” means a nucleic acid molecule to which an antisense compound hybridizes.

As used herein, “single nucleotide polymorphism” or “SNP” means a single nucleotide variation between the genomes of individuals of the same species. In some cases, a SNP may be a single nucleotide deletion or insertion.

As used herein, “mRNA” means an RNA molecule that encodes a protein.

As used herein, “pre-mRNA” means an RNA transcript that has not been fully processed into mRNA. Pre-mRNA includes one or more intron.

As used herein, “object RNA” means an RNA molecule other than a target RNA, the amount, activity, splicing, and/or function of which is modulated, either directly or indirectly, by a target nucleic acid. In certain embodiments, a target nucleic acid modulates splicing of an object RNA. In certain such embodiments, an antisense compound modulates the amount or activity of the target nucleic acid, resulting in a change in the splicing of an object RNA and ultimately resulting in a change in the activity or function of the object RNA.

As used herein, “microRNA” means a naturally occurring, small, non-coding RNA that represses gene expression of at least one mRNA. In certain embodiments, a microRNA represses gene expression by binding to a target site within a 3’ untranslated region of an mRNA. In certain embodiments, a microRNA has a nucleobase sequence as set forth in miRBase, a database of published microRNA sequences found at <http://microrna.sanger.ac.uk/sequences/>. In certain embodiments, a microRNA has a nucleobase sequence as set forth in miRBase version 12.0 released September 2008, which is herein incorporated by reference in its entirety.

As used herein, “microRNA mimic” means an oligomeric compound having a sequence that is at least partially identical to that of a microRNA. In certain embodiments, a microRNA mimic comprises the microRNA seed region of a microRNA. In certain embodiments, a microRNA mimic modulates translation

of more than one target nucleic acids. In certain embodiments, a microRNA mimic is double-stranded.

As used herein, “targeting” or “targeted to” means the association of an antisense compound to a particular target nucleic acid molecule or a particular region of a target nucleic acid molecule. An antisense compound targets a target nucleic acid if it is sufficiently complementary to the target nucleic acid to allow hybridization under physiological conditions.

As used herein, “nucleobase complementarity” or “complementarity” when in reference to nucleobases means a nucleobase that is capable of base pairing with another nucleobase. For example, in DNA, adenine (A) is complementary to thymine (T). For example, in RNA, adenine (A) is complementary to uracil (U). In certain embodiments, complementary nucleobase means a nucleobase of an antisense compound that is capable of base pairing with a nucleobase of its target nucleic acid. For example, if a nucleobase at a certain position of an antisense compound is capable of hydrogen bonding with a nucleobase at a certain position of a target nucleic acid, then the position of hydrogen bonding between the oligonucleotide and the target nucleic acid is considered to be complementary at that nucleobase pair. Nucleobases comprising certain modifications may maintain the ability to pair with a counterpart nucleobase and thus, are still capable of nucleobase complementarity.

As used herein, “non-complementary” in reference to nucleobases means a pair of nucleobases that do not form hydrogen bonds with one another.

As used herein, “complementary” in reference to oligomeric compounds (e.g., linked nucleosides, oligonucleotides, or nucleic acids) means the capacity of such oligomeric compounds or regions thereof to hybridize to another oligomeric compound or region thereof through nucleobase complementarity under stringent conditions. Complementary oligomeric compounds need not have nucleobase complementarity at each nucleoside. Rather, some mismatches are tolerated. In certain embodiments, complementary oligomeric compounds or regions are complementary at 70% of the nucleobases (70% complementary). In certain embodiments, complementary oligomeric compounds or regions are 80% complementary. In certain embodiments, complementary oligomeric compounds or regions are 90% complementary. In certain embodiments, complementary oligomeric compounds or regions are 95% complementary. In certain embodiments, complementary oligomeric compounds or regions are 100% complementary.

As used herein, “hybridization” means the pairing of complementary oligomeric compounds (e.g., an antisense compound and its target nucleic acid). While not limited to a particular mechanism, the most common mechanism of pairing involves hydrogen bonding, which may be Watson-Crick, Hoogsteen or reversed Hoogsteen hydrogen bonding, between complementary nucleobases.

As used herein, “specifically hybridizes” means the ability of an oligomeric compound to hybridize to one nucleic acid site with greater affinity than it hybridizes to another nucleic acid site. In certain embodiments, an antisense oligonucleotide specifically hybridizes to more than one target site.

As used herein, “fully complementary” in reference to an oligonucleotide or portion thereof means that each nucleobase of the oligonucleotide or portion thereof is capable of pairing with a nucleobase of a

complementary nucleic acid or contiguous portion thereof. Thus, a fully complementary region comprises no mismatches or unhybridized nucleobases in either strand.

As used herein, “percent complementarity” means the percentage of nucleobases of an oligomeric compound that are complementary to an equal-length portion of a target nucleic acid. Percent complementarity is calculated by dividing the number of nucleobases of the oligomeric compound that are complementary to nucleobases at corresponding positions in the target nucleic acid by the total length of the oligomeric compound.

As used herein, “percent identity” means the number of nucleobases in a first nucleic acid that are the same type (independent of chemical modification) as nucleobases at corresponding positions in a second nucleic acid, divided by the total number of nucleobases in the first nucleic acid.

As used herein, “modulation” means a change of amount or quality of a molecule, function, or activity when compared to the amount or quality of a molecule, function, or activity prior to modulation. For example, modulation includes the change, either an increase (stimulation or induction) or a decrease (inhibition or reduction) in gene expression. As a further example, modulation of expression can include a change in splice site selection of pre-mRNA processing, resulting in a change in the absolute or relative amount of a particular splice-variant compared to the amount in the absence of modulation.

As used herein, “motif” means a pattern of chemical modifications in an oligomeric compound or a region thereof. Motifs may be defined by modifications at certain nucleosides and/or at certain linking groups of an oligomeric compound.

As used herein, “nucleoside motif” means a pattern of nucleoside modifications in an oligomeric compound or a region thereof. The linkages of such an oligomeric compound may be modified or unmodified. Unless otherwise indicated, motifs herein describing only nucleosides are intended to be nucleoside motifs. Thus, in such instances, the linkages are not limited.

As used herein, “sugar motif” means a pattern of sugar modifications in an oligomeric compound or a region thereof.

As used herein, “linkage motif” means a pattern of linkage modifications in an oligomeric compound or region thereof. The nucleosides of such an oligomeric compound may be modified or unmodified. Unless otherwise indicated, motifs herein describing only linkages are intended to be linkage motifs. Thus, in such instances, the nucleosides are not limited.

As used herein, “nucleobase modification motif” means a pattern of modifications to nucleobases along an oligonucleotide. Unless otherwise indicated, a nucleobase modification motif is independent of the nucleobase sequence.

As used herein, “sequence motif” means a pattern of nucleobases arranged along an oligonucleotide or portion thereof. Unless otherwise indicated, a sequence motif is independent of chemical modifications and thus may have any combination of chemical modifications, including no chemical modifications.

As used herein, “type of modification” in reference to a nucleoside or a nucleoside of a “type” means

the chemical modification of a nucleoside and includes modified and unmodified nucleosides. Accordingly, unless otherwise indicated, a “nucleoside having a modification of a first type” may be an unmodified nucleoside.

As used herein, “differently modified” mean chemical modifications or chemical substituents that are different from one another, including absence of modifications. Thus, for example, a MOE nucleoside and an unmodified DNA nucleoside are “differently modified,” even though the DNA nucleoside is unmodified. Likewise, DNA and RNA are “differently modified,” even though both are naturally-occurring unmodified nucleosides. Nucleosides that are the same but for comprising different nucleobases are not differently modified. For example, a nucleoside comprising a 2'-OMe modified sugar and an unmodified adenine nucleobase and a nucleoside comprising a 2'-OMe modified sugar and an unmodified thymine nucleobase are not differently modified.

As used herein, “the same type of modifications” refers to modifications that are the same as one another, including absence of modifications. Thus, for example, two unmodified DNA nucleoside have “the same type of modification,” even though the DNA nucleoside is unmodified. Such nucleosides having the same type modification may comprise different nucleobases.

As used herein, “separate regions” means portions of an oligonucleotide wherein the chemical modifications or the motif of chemical modifications of any neighboring portions include at least one difference to allow the separate regions to be distinguished from one another.

As used herein, “pharmaceutically acceptable carrier or diluent” means any substance suitable for use in administering to an animal. In certain embodiments, a pharmaceutically acceptable carrier or diluent is sterile saline. In certain embodiments, such sterile saline is pharmaceutical grade saline.

As used herein, “substituent” and “substituent group,” means an atom or group that replaces the atom or group of a named parent compound. For example a substituent of a modified nucleoside is any atom or group that differs from the atom or group found in a naturally occurring nucleoside (e.g., a modified 2'-substituent is any atom or group at the 2'-position of a nucleoside other than H or OH). Substituent groups can be protected or unprotected. In certain embodiments, compounds of the present invention have substituents at one or at more than one position of the parent compound. Substituents may also be further substituted with other substituent groups and may be attached directly or via a linking group such as an alkyl or hydrocarbyl group to a parent compound.

Likewise, as used herein, “substituent” in reference to a chemical functional group means an atom or group of atoms differs from the atom or a group of atoms normally present in the named functional group. In certain embodiments, a substituent replaces a hydrogen atom of the functional group (e.g., in certain embodiments, the substituent of a substituted methyl group is an atom or group other than hydrogen which replaces one of the hydrogen atoms of an unsubstituted methyl group). Unless otherwise indicated, groups amenable for use as substituents include without limitation, halogen, hydroxyl, alkyl, alkenyl, alkynyl, acyl (-C(O)R_{aa}), carboxyl (-C(O)O-R_{aa}), aliphatic groups, alicyclic groups, alkoxy, substituted oxy (-O-R_{aa}), aryl,

aralkyl, heterocyclic radical, heteroaryl, heteroarylalkyl, amino ($-\text{N}(\text{R}_{\text{bb}})(\text{R}_{\text{cc}})$), imino ($=\text{NR}_{\text{bb}}$), amido ($-\text{C}(\text{O})\text{N}(\text{R}_{\text{bb}})(\text{R}_{\text{cc}})$ or $-\text{N}(\text{R}_{\text{bb}})\text{C}(\text{O})\text{R}_{\text{aa}}$), azido ($-\text{N}_3$), nitro ($-\text{NO}_2$), cyano ($-\text{CN}$), carbamido ($-\text{OC}(\text{O})\text{N}(\text{R}_{\text{bb}})(\text{R}_{\text{cc}})$ or $-\text{N}(\text{R}_{\text{bb}})\text{C}(\text{O})\text{OR}_{\text{aa}}$), ureido ($-\text{N}(\text{R}_{\text{bb}})\text{C}(\text{O})\text{N}(\text{R}_{\text{bb}})(\text{R}_{\text{cc}})$), thioureido ($-\text{N}(\text{R}_{\text{bb}})\text{C}(\text{S})\text{N}(\text{R}_{\text{bb}})(\text{R}_{\text{cc}})$), guanidiny ($-\text{N}(\text{R}_{\text{bb}})\text{C}(=\text{NR}_{\text{bb}})\text{N}(\text{R}_{\text{bb}})(\text{R}_{\text{cc}})$), amidiny ($-\text{C}(=\text{NR}_{\text{bb}})\text{N}(\text{R}_{\text{bb}})(\text{R}_{\text{cc}})$ or $-\text{N}(\text{R}_{\text{bb}})\text{C}(=\text{NR}_{\text{bb}})(\text{R}_{\text{aa}})$), thiol ($-\text{SR}_{\text{bb}}$), sulfinyl ($-\text{S}(\text{O})\text{R}_{\text{bb}}$), sulfonyl ($-\text{S}(\text{O})_2\text{R}_{\text{bb}}$) and sulfonamidyl ($-\text{S}(\text{O})_2\text{N}(\text{R}_{\text{bb}})(\text{R}_{\text{cc}})$ or $-\text{N}(\text{R}_{\text{bb}})\text{S}(\text{O})_2\text{R}_{\text{bb}}$). Wherein each R_{aa} , R_{bb} and R_{cc} is, independently, H, an optionally linked chemical functional group or a further substituent group with a preferred list including without limitation, alkyl, alkenyl, alkynyl, aliphatic, alkoxy, acyl, aryl, aralkyl, heteroaryl, alicyclic, heterocyclic and heteroarylalkyl. Selected substituents within the compounds described herein are present to a recursive degree.

As used herein, "alkyl," as used herein, means a saturated straight or branched hydrocarbon radical containing up to twenty four carbon atoms. Examples of alkyl groups include without limitation, methyl, ethyl, propyl, butyl, isopropyl, n-hexyl, octyl, decyl, dodecyl and the like. Alkyl groups typically include from 1 to about 24 carbon atoms, more typically from 1 to about 12 carbon atoms ($\text{C}_1\text{-C}_{12}$ alkyl) with from 1 to about 6 carbon atoms being more preferred.

As used herein, "alkenyl," means a straight or branched hydrocarbon chain radical containing up to twenty four carbon atoms and having at least one carbon-carbon double bond. Examples of alkenyl groups include without limitation, ethenyl, propenyl, butenyl, 1-methyl-2-buten-1-yl, dienes such as 1,3-butadiene and the like. Alkenyl groups typically include from 2 to about 24 carbon atoms, more typically from 2 to about 12 carbon atoms with from 2 to about 6 carbon atoms being more preferred. Alkenyl groups as used herein may optionally include one or more further substituent groups.

As used herein, "alkynyl," means a straight or branched hydrocarbon radical containing up to twenty four carbon atoms and having at least one carbon-carbon triple bond. Examples of alkynyl groups include, without limitation, ethynyl, 1-propynyl, 1-butylnyl, and the like. Alkynyl groups typically include from 2 to about 24 carbon atoms, more typically from 2 to about 12 carbon atoms with from 2 to about 6 carbon atoms being more preferred. Alkynyl groups as used herein may optionally include one or more further substituent groups.

As used herein, "acyl," means a radical formed by removal of a hydroxyl group from an organic acid and has the general Formula $-\text{C}(\text{O})-\text{X}$ where X is typically aliphatic, alicyclic or aromatic. Examples include aliphatic carbonyls, aromatic carbonyls, aliphatic sulfonyls, aromatic sulfinyls, aliphatic sulfinyls, aromatic phosphates, aliphatic phosphates and the like. Acyl groups as used herein may optionally include further substituent groups.

As used herein, "alicyclic" means a cyclic ring system wherein the ring is aliphatic. The ring system can comprise one or more rings wherein at least one ring is aliphatic. Preferred alicyclics include rings having from about 5 to about 9 carbon atoms in the ring. Alicyclic as used herein may optionally include further substituent groups.

As used herein, "aliphatic" means a straight or branched hydrocarbon radical containing up to twenty four carbon atoms wherein the saturation between any two carbon atoms is a single, double or triple bond. An aliphatic group preferably contains from 1 to about 24 carbon atoms, more typically from 1 to about 12 carbon atoms with from 1 to about 6 carbon atoms being more preferred. The straight or branched chain of an aliphatic group may be interrupted with one or more heteroatoms that include nitrogen, oxygen, sulfur and phosphorus. Such aliphatic groups interrupted by heteroatoms include without limitation, polyalkoxys, such as polyalkylene glycols, polyamines, and polyimines. Aliphatic groups as used herein may optionally include further substituent groups.

As used herein, "alkoxy" means a radical formed between an alkyl group and an oxygen atom wherein the oxygen atom is used to attach the alkoxy group to a parent molecule. Examples of alkoxy groups include without limitation, methoxy, ethoxy, propoxy, isopropoxy, *n*-butoxy, *sec*-butoxy, *tert*-butoxy, *n*-pentoxy, neopentoxy, *n*-hexoxy and the like. Alkoxy groups as used herein may optionally include further substituent groups.

As used herein, "aminoalkyl" means an amino substituted C₁-C₁₂ alkyl radical. The alkyl portion of the radical forms a covalent bond with a parent molecule. The amino group can be located at any position and the aminoalkyl group can be substituted with a further substituent group at the alkyl and/or amino portions.

As used herein, "aralkyl" and "arylalkyl" mean an aromatic group that is covalently linked to a C₁-C₁₂ alkyl radical. The alkyl radical portion of the resulting aralkyl (or arylalkyl) group forms a covalent bond with a parent molecule. Examples include without limitation, benzyl, phenethyl and the like. Aralkyl groups as used herein may optionally include further substituent groups attached to the alkyl, the aryl or both groups that form the radical group.

As used herein, "aryl" and "aromatic" mean a mono- or polycyclic carbocyclic ring system radicals having one or more aromatic rings. Examples of aryl groups include without limitation, phenyl, naphthyl, tetrahydronaphthyl, indanyl, idenyl and the like. Preferred aryl ring systems have from about 5 to about 20 carbon atoms in one or more rings. Aryl groups as used herein may optionally include further substituent groups.

As used herein, "halo" and "halogen," mean an atom selected from fluorine, chlorine, bromine and iodine.

As used herein, "heteroaryl," and "heteroaromatic," mean a radical comprising a mono- or polycyclic aromatic ring, ring system or fused ring system wherein at least one of the rings is aromatic and includes one or more heteroatoms. Heteroaryl is also meant to include fused ring systems including systems where one or more of the fused rings contain no heteroatoms. Heteroaryl groups typically include one ring atom selected from sulfur, nitrogen or oxygen. Examples of heteroaryl groups include without limitation, pyridinyl, pyrazinyl, pyrimidinyl, pyrrolyl, pyrazolyl, imidazolyl, thiazolyl, oxazolyl, isooxazolyl, thiadiazolyl, oxadiazolyl, thiophenyl, furanyl, quinolinyl, isoquinolinyl, benzimidazolyl, benzooxazolyl,

quinoxalinyll and the like. Heteroaryl radicals can be attached to a parent molecule directly or through a linking moiety such as an aliphatic group or hetero atom. Heteroaryl groups as used herein may optionally include further substituent groups.

Oligomeric Compounds

In certain embodiments, the present invention provides oligomeric compounds. In certain embodiments, such oligomeric compounds comprise oligonucleotides optionally comprising one or more conjugate and/or terminal groups. In certain embodiments, an oligomeric compound consists of an oligonucleotide. In certain embodiments, oligonucleotides comprise one or more chemical modifications. Such chemical modifications include modifications one or more nucleoside (including modifications to the sugar moiety and/or the nucleobase) and/or modifications to one or more internucleoside linkage.

Certain Sugar Moieties

In certain embodiments, oligomeric compounds of the invention comprise one or more modified nucleosides comprising a modified sugar moiety. Such oligomeric compounds comprising one or more sugar-modified nucleosides may have desirable properties, such as enhanced nuclease stability or increased binding affinity with a target nucleic acid relative to oligomeric compounds comprising only nucleosides comprising naturally occurring sugar moieties. In certain embodiments, modified sugar moieties are substituted sugar moieties. In certain embodiments, modified sugar moieties are sugar surrogates. Such sugar surrogates may comprise one or more substitutions corresponding to those of substituted sugar moieties.

In certain embodiments, modified sugar moieties are substituted sugar moieties comprising one or more non-bridging sugar substituent, including but not limited to substituents at the 2' and/or 5' positions. Examples of sugar substituents suitable for the 2'-position, include, but are not limited to: 2'-F, 2'-OCH₃ ("OMe" or "O-methyl"), and 2'-O(CH₂)₂OCH₃ ("MOE"). In certain embodiments, sugar substituents at the 2' position is selected from allyl, amino, azido, thio, O-allyl, O-C₁-C₁₀ alkyl, O-C₁-C₁₀ substituted alkyl; OCF₃, O(CH₂)₂SCH₃, O(CH₂)₂-O-N(R_m)(R_n), and O-CH₂-C(=O)-N(R_m)(R_n), where each R_m and R_n is, independently, H or substituted or unsubstituted C₁-C₁₀ alkyl. Examples of sugar substituents at the 5'-position, include, but are not limited to: 5'-methyl (R or S); 5'-vinyl, and 5'-methoxy. In certain embodiments, substituted sugars comprise more than one non-bridging sugar substituent, for example, 2'-F-5'-methyl sugar moieties (*see, e.g.*, PCT International Application WO 2008/101157, for additional 5', 2'-bis substituted sugar moieties and nucleosides).

Nucleosides comprising 2'-substituted sugar moieties are referred to as 2'-substituted nucleosides. In certain embodiments, a 2'-substituted nucleoside comprises a 2'-substituent group selected from halo, allyl, amino, azido, SH, CN, OCN, CF₃, OCF₃, O, S, or N(R_m)-alkyl; O, S, or N(R_m)-alkenyl; O, S or N(R_m)-alkynyl; O-alkylenyl-O-alkyl, alkynyl, alkaryl, aralkyl, O-alkaryl, O-aralkyl, O(CH₂)₂SCH₃, O-(CH₂)₂-O-N(R_m)(R_n) or O-CH₂-C(=O)-N(R_m)(R_n), where each R_m and R_n is, independently, H, an amino protecting

group or substituted or unsubstituted C₁-C₁₀ alkyl. These 2'-substituent groups can be further substituted with one or more substituent groups independently selected from hydroxyl, amino, alkoxy, carboxy, benzyl, phenyl, nitro (NO₂), thiol, thioalkoxy (S-alkyl), halogen, alkyl, aryl, alkenyl and alkynyl.

In certain embodiments, a 2'-substituted nucleoside comprises a 2'-substituent group selected from F, NH₂, N₃, OCF₃, O-CH₃, O(CH₂)₃NH₂, CH₂-CH=CH₂, O-CH₂-CH=CH₂, OCH₂CH₂OCH₃, O(CH₂)₂SCH₃, O-(CH₂)₂-O-N(R_m)(R_n), O(CH₂)₂O(CH₂)₂N(CH₃)₂, and N-substituted acetamide (O-CH₂-C(=O)-N(R_m)(R_n)) where each R_m and R_n is, independently, H, an amino protecting group or substituted or unsubstituted C₁-C₁₀ alkyl.

In certain embodiments, a 2'-substituted nucleoside comprises a sugar moiety comprising a 2'-substituent group selected from F, OCF₃, O-CH₃, OCH₂CH₂OCH₃, O(CH₂)₂SCH₃, O-(CH₂)₂-O-N(CH₃)₂, -O(CH₂)₂O(CH₂)₂N(CH₃)₂, and O-CH₂-C(=O)-N(H)CH₃.

In certain embodiments, a 2'-substituted nucleoside comprises a sugar moiety comprising a 2'-substituent group selected from F, O-CH₃, and OCH₂CH₂OCH₃.

Certain modified sugar moieties comprise a bridging sugar substituent that forms a second ring resulting in a bicyclic sugar moiety. In certain such embodiments, the bicyclic sugar moiety comprises a bridge between the 4' and the 2' furanose ring atoms. Examples of such 4' to 2' sugar substituents, include, but are not limited to: -[C(R_a)(R_b)]_n-, -[C(R_a)(R_b)]_n-O-, -C(R_aR_b)-N(R)-O- or, -C(R_aR_b)-O-N(R)-; 4'-CH₂-2', 4'-(CH₂)₂-2', 4'-(CH₂)₃-2', 4'-(CH₂)-O-2' (LNA); 4'-(CH₂)-S-2'; 4'-(CH₂)₂-O-2' (ENA); 4'-CH(CH₃)-O-2' (cEt) and 4'-CH(CH₂OCH₃)-O-2', and analogs thereof (*see, e.g.*, U.S. Patent 7,399,845, issued on July 15, 2008); 4'-C(CH₃)(CH₃)-O-2' and analogs thereof, (*see, e.g.*, WO2009/006478, published January 8, 2009); 4'-CH₂-N(OCH₃)-2' and analogs thereof (*see, e.g.*, WO2008/150729, published December 11, 2008); 4'-CH₂-O-N(CH₃)-2' (*see, e.g.*, US2004/0171570, published September 2, 2004); 4'-CH₂-O-N(R)-2', and 4'-CH₂-N(R)-O-2', wherein each R is, independently, H, a protecting group, or C₁-C₁₂ alkyl; 4'-CH₂-N(R)-O-2', wherein R is H, C₁-C₁₂ alkyl, or a protecting group (*see*, U.S. Patent 7,427,672, issued on September 23, 2008); 4'-CH₂-C(H)(CH₃)-2' (*see, e.g.*, Chattopadhyaya, *et al.*, *J. Org. Chem.*, 2009, 74, 118-134); and 4'-CH₂-C(=CH₂)-2' and analogs thereof (*see*, published PCT International Application WO 2008/154401, published on December 8, 2008).

In certain embodiments, such 4' to 2' bridges independently comprise from 1 to 4 linked groups independently selected from -[C(R_a)(R_b)]_n-, -C(R_a)=C(R_b)-, -C(R_a)=N-, -C(=NR_a)-, -C(=O)-, -C(=S)-, -O-, -Si(R_a)₂-, -S(=O)_x-, and -N(R_a)-;

wherein:

x is 0, 1, or 2;


n is 1, 2, 3, or 4;

each R_a and R_b is, independently, H, a protecting group, hydroxyl, C₁-C₁₂ alkyl, substituted C₁-C₁₂ alkyl, C₂-C₁₂ alkenyl, substituted C₂-C₁₂ alkenyl, C₂-C₁₂ alkynyl, substituted C₂-C₁₂ alkynyl, C₅-C₂₀ aryl, substituted C₅-C₂₀ aryl, heterocycle radical, substituted heterocycle radical, heteroaryl, substituted heteroaryl,

C₅-C₇ alicyclic radical, substituted C₅-C₇ alicyclic radical, halogen, OJ₁, NJ₁J₂, SJ₁, N₃, COOJ₁, acyl (C(=O)-H), substituted acyl, CN, sulfonyl (S(=O)₂-J₁), or sulfoxyl (S(=O)-J₁); and

each J₁ and J₂ is, independently, H, C₁-C₁₂ alkyl, substituted C₁-C₁₂ alkyl, C₂-C₁₂ alkenyl, substituted C₂-C₁₂ alkenyl, C₂-C₁₂ alkynyl, substituted C₂-C₁₂ alkynyl, C₅-C₂₀ aryl, substituted C₅-C₂₀ aryl, acyl (C(=O)-H), substituted acyl, a heterocycle radical, a substituted heterocycle radical, C₁-C₁₂ aminoalkyl, substituted C₁-C₁₂ aminoalkyl, or a protecting group.

Nucleosides comprising bicyclic sugar moieties are referred to as bicyclic nucleosides or BNAs. Bicyclic nucleosides include, but are not limited to, (A) α -L-Methyleneoxy (4'-CH₂-O-2') BNA, (B) β -D-Methyleneoxy (4'-CH₂-O-2') BNA (also referred to as locked nucleic acid or LNA), (C) Ethyleneoxy (4'-(CH₂)₂-O-2') BNA, (D) Aminooxy (4'-CH₂-O-N(R)-2') BNA, (E) Oxyamino (4'-CH₂-N(R)-O-2') BNA, (F) Methyl(methyleneoxy) (4'-CH(CH₃)-O-2') BNA (also referred to as constrained ethyl or cEt), (G) methylene-thio (4'-CH₂-S-2') BNA, (H) methylene-amino (4'-CH₂-N(R)-2') BNA, (I) methyl carbocyclic (4'-CH₂-CH(CH₃)-2') BNA, and (J) propylene carbocyclic (4'-(CH₂)₃-2') BNA as depicted below.



wherein Bx is a nucleobase moiety and R is, independently, H, a protecting group, or C₁-C₁₂ alkyl.

Additional bicyclic sugar moieties are known in the art, for example: Singh et al., *Chem. Commun.*, 1998, 4, 455-456; Koshkin et al., *Tetrahedron*, 1998, 54, 3607-3630; Wahlestedt et al., *Proc. Natl. Acad. Sci.*


U. S. A., 2000, 97, 5633-5638; Kumar et al., *Bioorg. Med. Chem. Lett.*, 1998, 8, 2219-2222; Singh et al., *J. Org. Chem.*, 1998, 63, 10035-10039; Srivastava et al., *J. Am. Chem. Soc.*, 129(26) 8362-8379 (Jul. 4, 2007); Elayadi et al., *Curr. Opinion Invens. Drugs*, 2001, 2, 558-561; Braasch et al., *Chem. Biol.*, 2001, 8, 1-7; Orum et al., *Curr. Opinion Mol. Ther.*, 2001, 3, 239-243; U.S. Patent Nos. 7,053,207, 6,268,490, 6,770,748, 6,794,499, 7,034,133, 6,525,191, 6,670,461, and 7,399,845; WO 2004/106356, WO 1994/14226, WO 2005/021570, and WO 2007/134181; U.S. Patent Publication Nos. US2004/0171570, US2007/0287831, and US2008/0039618; U.S. Patent Serial Nos. 12/129,154, 60/989,574, 61/026,995, 61/026,998, 61/056,564, 61/086,231, 61/097,787, and 61/099,844; and PCT International Applications Nos. PCT/US2008/064591, PCT/US2008/066154, and PCT/US2008/068922.

In certain embodiments, bicyclic sugar moieties and nucleosides incorporating such bicyclic sugar moieties are further defined by isomeric configuration. For example, a nucleoside comprising a 4'-2' methylene-oxy bridge, may be in the α -L configuration or in the β -D configuration. Previously, α -L-methyleneoxy (4'-CH₂-O-2') bicyclic nucleosides have been incorporated into antisense oligonucleotides that showed antisense activity (Frieden et al., *Nucleic Acids Research*, 2003, 31, 6365-6372).

In certain embodiments, substituted sugar moieties comprise one or more non-bridging sugar substituent and one or more bridging sugar substituent (e.g., 5'-substituted and 4'-2' bridged sugars). (see, PCT International Application WO 2007/134181, published on 11/22/07, wherein LNA is substituted with, for example, a 5'-methyl or a 5'-vinyl group).

In certain embodiments, modified sugar moieties are sugar surrogates. In certain such embodiments, the oxygen atom of the naturally occurring sugar is substituted, e.g., with a sulfur, carbon or nitrogen atom. In certain such embodiments, such modified sugar moiety also comprises bridging and/or non-bridging substituents as described above. For example, certain sugar surrogates comprise a 4'-sulfur atom and a substitution at the 2'-position (see, e.g., published U.S. Patent Application US2005/0130923, published on June 16, 2005) and/or the 5' position. By way of additional example, carbocyclic bicyclic nucleosides having a 4'-2' bridge have been described (see, e.g., Freier et al., *Nucleic Acids Research*, 1997, 25(22), 4429-4443 and Albaek et al., *J. Org. Chem.*, 2006, 71, 7731-7740).

In certain embodiments, sugar surrogates comprise rings having other than 5-atoms. For example, in certain embodiments, a sugar surrogate comprises a six-membered tetrahydropyran. Such tetrahydropyrans may be further modified or substituted. Nucleosides comprising such modified tetrahydropyrans include, but are not limited to, hexitol nucleic acid (HNA), anitol nucleic acid (ANA), manitol nucleic acid (MNA) (see Leumann, C.J. *Bioorg. & Med. Chem.* (2002) 10:841-854), fluoro HNA (F-HNA), and those compounds having Formula VII:



VII

wherein independently for each of said at least one tetrahydropyran nucleoside analog of Formula VII:

Bx is a nucleobase moiety;

T₃ and T₄ are each, independently, an internucleoside linking group linking the tetrahydropyran nucleoside analog to the antisense compound or one of T₃ and T₄ is an internucleoside linking group linking the tetrahydropyran nucleoside analog to the antisense compound and the other of T₃ and T₄ is H, a hydroxyl protecting group, a linked conjugate group, or a 5' or 3'-terminal group;

q₁, q₂, q₃, q₄, q₅, q₆ and q₇ are each, independently, H, C₁-C₆ alkyl, substituted C₁-C₆ alkyl, C₂-C₆ alkenyl, substituted C₂-C₆ alkenyl, C₂-C₆ alkynyl, or substituted C₂-C₆ alkynyl; and

one of R₁ and R₂ is hydrogen and the other is selected from halogen, substituted or unsubstituted alkoxy, NJ₁J₂, SJ₁, N₃, OC(=X)J₁, OC(=X)NJ₁J₂, NJ₃C(=X)NJ₁J₂, and CN, wherein X is O, S or NJ₁, and each J₁, J₂, and J₃ is, independently, H or C₁-C₆ alkyl.

In certain embodiments, the modified THP nucleosides of Formula VII are provided wherein q₁, q₂, q₃, q₄, q₅, q₆ and q₇ are each H. In certain embodiments, at least one of q₁, q₂, q₃, q₄, q₅, q₆ and q₇ is other than H. In certain embodiments, at least one of q₁, q₂, q₃, q₄, q₅, q₆ and q₇ is methyl. In certain embodiments, THP nucleosides of Formula VII are provided wherein one of R₁ and R₂ is F. In certain embodiments, R₁ is fluoro and R₂ is H, R₁ is methoxy and R₂ is H, and R₁ is methoxyethoxy and R₂ is H.

Many other bicyclo and tricyclo sugar surrogate ring systems are also known in the art that can be used to modify nucleosides for incorporation into antisense compounds (*see, e.g.*, review article: Leumann, J. C, *Bioorganic & Medicinal Chemistry*, **2002**, 10, 841-854).

Combinations of modifications are also provided without limitation, such as 2'-F-5'-methyl substituted nucleosides (see PCT International Application WO 2008/101157 Published on 8/21/08 for other disclosed 5', 2'-bis substituted nucleosides) and replacement of the ribosyl ring oxygen atom with S and further substitution at the 2'-position (see published U.S. Patent Application US2005-0130923, published on June 16, 2005) or alternatively 5'-substitution of a bicyclic nucleic acid (see PCT International Application WO 2007/134181, published on 11/22/07 wherein a 4'-CH₂-O-2' bicyclic nucleoside is further substituted at the 5' position with a 5'-methyl or a 5'-vinyl group). The synthesis and preparation of carbocyclic bicyclic nucleosides along with their oligomerization and biochemical studies have also been described (*see, e.g.*, Srivastava *et al.*, *J. Am. Chem. Soc.* 2007, 129(26), 8362-8379).

Certain Nucleobases

In certain embodiments, nucleosides of the present invention comprise one or more unmodified nucleobases. In certain embodiments, nucleosides of the present invention comprise one or more modified nucleobases.

In certain embodiments, modified nucleobases are selected from: universal bases, hydrophobic bases, promiscuous bases, size-expanded bases, and fluorinated bases as defined herein. 5-substituted pyrimidines, 6-azapyrimidines and N-2, N-6 and O-6 substituted purines, including 2-aminopropyladenine, 5-propynyluracil; 5-propynylcytosine; 5-hydroxymethyl cytosine, xanthine, hypoxanthine, 2-aminoadenine, 6-methyl and other alkyl derivatives of adenine and guanine, 2-propyl and other alkyl derivatives of adenine and guanine, 2-thiouracil, 2-thiothymine and 2-thiocytosine, 5-halouracil and cytosine, 5-propynyl (-C≡C-CH₃) uracil and cytosine and other alkynyl derivatives of pyrimidine bases, 6-azo uracil, cytosine and thymine, 5-uracil (pseudouracil), 4-thiouracil, 8-halo, 8-amino, 8-thiol, 8-thioalkyl, 8-hydroxyl and other 8-substituted adenines and guanines, 5-halo particularly 5-bromo, 5-trifluoromethyl and other 5-substituted uracils and cytosines, 7-methylguanine and 7-methyladenine, 2-F-adenine, 2-amino-adenine, 8-azaguanine and 8-azaadenine, 7-deazaguanine and 7-deazaadenine, 3-deazaguanine and 3-deazaadenine, universal bases, hydrophobic bases, promiscuous bases, size-expanded bases, and fluorinated bases as defined herein. Further modified nucleobases include tricyclic pyrimidines such as phenoxazine cytidine([5,4-b][1,4]benzoxazin-2(3H)-one), phenothiazine cytidine (1H-pyrimido[5,4-b][1,4]benzothiazin-2(3H)-one), G-clamps such as a substituted phenoxazine cytidine (e.g. 9-(2-aminoethoxy)-H-pyrimido[5,4-b][1,4]benzoxazin-2(3H)-one), carbazole cytidine (2H-pyrimido[4,5-b]indol-2-one), pyridindole cytidine (H-pyrido[3',2':4,5]pyrrolo[2,3-d]pyrimidin-2-one). Modified nucleobases may also include those in which the purine or pyrimidine base is replaced with other heterocycles, for example 7-deaza-adenine, 7-deazaguanosine, 2-aminopyridine and 2-pyridone. Further nucleobases include those disclosed in United States Patent No. 3,687,808, those disclosed in *The Concise Encyclopedia Of Polymer Science And Engineering*, Kroschwitz, J.I., Ed., John Wiley & Sons, 1990, 858-859; those disclosed by Englisch *et al.*, *Angewandte Chemie*, International Edition, 1991, 30, 613; and those disclosed by Sanghvi, Y.S., Chapter 15, *Antisense Research and Applications*, Crooke, S.T. and Lebleu, B., Eds., CRC Press, 1993, 273-288.

Representative United States patents that teach the preparation of certain of the above noted modified nucleobases as well as other modified nucleobases include without limitation, U.S. 3,687,808; 4,845,205; 5,130,302; 5,134,066; 5,175,273; 5,367,066; 5,432,272; 5,457,187; 5,459,255; 5,484,908; 5,502,177; 5,525,711; 5,552,540; 5,587,469; 5,594,121; 5,596,091; 5,614,617; 5,645,985; 5,681,941; 5,750,692; 5,763,588; 5,830,653 and 6,005,096, certain of which are commonly owned with the instant application, and each of which is herein incorporated by reference in its entirety.

Certain Internucleoside Linkages

In certain embodiments, the present invention provides oligomeric compounds comprising linked nucleosides. In such embodiments, nucleosides may be linked together using any internucleoside linkage. The two main classes of internucleoside linking groups are defined by the presence or absence of a phosphorus atom. Representative phosphorus containing internucleoside linkages include, but are not limited to, phosphodiester (P=O), phosphotriesters, methylphosphonates, phosphoramidate, and phosphorothioates (P=S). Representative non-phosphorus containing internucleoside linking groups include, but are not limited to, methylenemethylimino (-CH₂-N(CH₃)-O-CH₂-), thiodiester (-O-C(O)-S-), thionocarbamate (-O-C(O)(NH)-S-); siloxane (-O-Si(H)₂-O-); and N,N'-dimethylhydrazine (-CH₂-N(CH₃)-N(CH₃)-). Modified linkages, compared to natural phosphodiester linkages, can be used to alter, typically increase, nuclease resistance of the oligomeric compound. In certain embodiments, internucleoside linkages having a chiral atom can be prepared as a racemic mixture, or as separate enantiomers. Representative chiral linkages include, but are not limited to, alkylphosphonates and phosphorothioates. Methods of preparation of phosphorous-containing and non-phosphorous-containing internucleoside linkages are well known to those skilled in the art.

The oligonucleotides described herein contain one or more asymmetric centers and thus give rise to enantiomers, diastereomers, and other stereoisomeric configurations that may be defined, in terms of absolute stereochemistry, as (R) or (S), α or β such as for sugar anomers, or as (D) or (L) such as for amino acids etc. Included in the antisense compounds provided herein are all such possible isomers, as well as their racemic and optically pure forms.

Neutral internucleoside linkages include without limitation, phosphotriesters, methylphosphonates, MMI (3'-CH₂-N(CH₃)-O-5'), amide-3 (3'-CH₂-C(=O)-N(H)-5'), amide-4 (3'-CH₂-N(H)-C(=O)-5'), formacetal (3'-O-CH₂-O-5'), and thioformacetal (3'-S-CH₂-O-5'). Further neutral internucleoside linkages include nonionic linkages comprising siloxane (dialkylsiloxane), carboxylate ester, carboxamide, sulfide, sulfonate ester and amides (See for example: *Carbohydrate Modifications in Antisense Research*; Y.S. Sanghvi and P.D. Cook, Eds., ACS Symposium Series 580; Chapters 3 and 4, 40-65). Further neutral internucleoside linkages include nonionic linkages comprising mixed N, O, S and CH₂ component parts.

Certain Motifs

In certain embodiments, the present invention provides oligomeric compounds comprising oligonucleotides. In certain embodiments, such oligonucleotides comprise one or more chemical modification. In certain embodiments, chemically modified oligonucleotides comprise one or more modified sugars. In certain embodiments, chemically modified oligonucleotides comprise one or more modified nucleobases. In certain embodiments, chemically modified oligonucleotides comprise one or more modified internucleoside linkages. In certain embodiments, the chemical modifications (sugar modifications, nucleobase modifications, and/or linkage modifications) define a pattern or motif. In certain embodiments, the patterns of chemical modifications of sugar moieties, internucleoside linkages, and nucleobases are each

independent of one another. Thus, an oligonucleotide may be described by its sugar modification motif, internucleoside linkage motif and/or nucleobase modification motif (as used herein, nucleobase modification motif describes the chemical modifications to the nucleobases independent of the sequence of nucleobases).

Certain sugar motifs

In certain embodiments, oligonucleotides comprise one or more type of modified sugar moieties and/or naturally occurring sugar moieties arranged along an oligonucleotide or region thereof in a defined pattern or sugar modification motif. Such motifs may include any of the sugar modifications discussed herein and/or other known sugar modifications.

In certain embodiments, the oligonucleotides comprise or consist of a region having a gapmer sugar modification motif, which comprises two external regions or “wings” and an internal region or “gap.” The three regions of a gapmer motif (the 5'-wing, the gap, and the 3'-wing) form a contiguous sequence of nucleosides wherein at least some of the sugar moieties of the nucleosides of each of the wings differ from at least some of the sugar moieties of the nucleosides of the gap. Specifically, at least the sugar moieties of the nucleosides of each wing that are closest to the gap (the 3'-most nucleoside of the 5'-wing and the 5'-most nucleoside of the 3'-wing) differ from the sugar moiety of the neighboring gap nucleosides, thus defining the boundary between the wings and the gap. In certain embodiments, the sugar moieties within the gap are the same as one another. In certain embodiments, the gap includes one or more nucleoside having a sugar moiety that differs from the sugar moiety of one or more other nucleosides of the gap. In certain embodiments, the sugar modification motifs of the two wings are the same as one another (symmetric gapmer). In certain embodiments, the sugar modification motifs of the 5'-wing differs from the sugar modification motif of the 3'-wing (asymmetric gapmer).

Certain 5'-wings

In certain embodiments, the 5'- wing of a gapmer consists of 1 to 5 linked nucleosides. In certain embodiments, the 5'- wing of a gapmer consists of 2 to 5 linked nucleosides. In certain embodiments, the 5'- wing of a gapmer consists of 3 to 5 linked nucleosides. In certain embodiments, the 5'- wing of a gapmer consists of 4 or 5 linked nucleosides. In certain embodiments, the 5'- wing of a gapmer consists of 1 to 4 linked nucleosides. In certain embodiments, the 5'- wing of a gapmer consists of 1 to 3 linked nucleosides. In certain embodiments, the 5'- wing of a gapmer consists of 1 or 2 linked nucleosides. In certain embodiments, the 5'- wing of a gapmer consists of 2 to 4 linked nucleosides. In certain embodiments, the 5'- wing of a gapmer consists of 2 or 3 linked nucleosides. In certain embodiments, the 5'- wing of a gapmer consists of 3 or 4 linked nucleosides. In certain embodiments, the 5'- wing of a gapmer consists of 1 nucleoside. In certain embodiments, the 5'- wing of a gapmer consists of 2 linked nucleosides. In certain embodiments, the 5'- wing of a gapmer consists of 3 linked nucleosides. In certain embodiments, the 5'- wing of a gapmer consists of 4 linked nucleosides. In certain embodiments, the 5'- wing of a gapmer consists of 5 linked nucleosides.

In certain embodiments, the 5'-wing of a gapmer comprises at least one bicyclic nucleoside. In certain embodiments, the 5'-wing of a gapmer comprises at least two bicyclic nucleosides. In certain embodiments, the 5'-wing of a gapmer comprises at least three bicyclic nucleosides. In certain embodiments, the 5'-wing of a gapmer comprises at least four bicyclic nucleosides. In certain embodiments, the 5'-wing of a gapmer comprises at least one constrained ethyl nucleoside. In certain embodiments, the 5'-wing of a gapmer comprises at least one LNA nucleoside. In certain embodiments, each nucleoside of the 5'-wing of a gapmer is a bicyclic nucleoside. In certain embodiments, each nucleoside of the 5'-wing of a gapmer is a constrained ethyl nucleoside. In certain embodiments, each nucleoside of the 5'-wing of a gapmer is a LNA nucleoside.

In certain embodiments, the 5'-wing of a gapmer comprises at least one non-bicyclic modified nucleoside. In certain embodiments, the 5'-wing of a gapmer comprises at least one 2'-substituted nucleoside. In certain embodiments, the 5'-wing of a gapmer comprises at least one 2'-MOE nucleoside. In certain embodiments, the 5'-wing of a gapmer comprises at least one 2'-OMe nucleoside. In certain embodiments, each nucleoside of the 5'-wing of a gapmer is a non-bicyclic modified nucleoside. In certain embodiments, each nucleoside of the 5'-wing of a gapmer is a 2'-substituted nucleoside. In certain embodiments, each nucleoside of the 5'-wing of a gapmer is a 2'-MOE nucleoside. In certain embodiments, each nucleoside of the 5'-wing of a gapmer is a 2'-OMe nucleoside.

In certain embodiments, the 5'-wing of a gapmer comprises at least one 2'-deoxynucleoside. In certain embodiments, each nucleoside of the 5'-wing of a gapmer is a 2'-deoxynucleoside. In certain embodiments, the 5'-wing of a gapmer comprises at least one ribonucleoside. In certain embodiments, each nucleoside of the 5'-wing of a gapmer is a ribonucleoside.

In certain embodiments, the 5'-wing of a gapmer comprises at least one bicyclic nucleoside and at least one non-bicyclic modified nucleoside. In certain embodiments, the 5'-wing of a gapmer comprises at least one bicyclic nucleoside and at least one 2'-substituted nucleoside. In certain embodiments, the 5'-wing of a gapmer comprises at least one bicyclic nucleoside and at least one 2'-MOE nucleoside. In certain embodiments, the 5'-wing of a gapmer comprises at least one bicyclic nucleoside and at least one 2'-OMe nucleoside. In certain embodiments, the 5'-wing of a gapmer comprises at least one bicyclic nucleoside and at least one 2'-deoxynucleoside.

In certain embodiments, the 5'-wing of a gapmer comprises at least one constrained ethyl nucleoside and at least one non-bicyclic modified nucleoside. In certain embodiments, the 5'-wing of a gapmer comprises at least one constrained ethyl nucleoside and at least one 2'-substituted nucleoside. In certain embodiments, the 5'-wing of a gapmer comprises at least one constrained ethyl nucleoside and at least one 2'-MOE nucleoside. In certain embodiments, the 5'-wing of a gapmer comprises at least one constrained ethyl nucleoside and at least one 2'-OMe nucleoside. In certain embodiments, the 5'-wing of a gapmer comprises at least one constrained ethyl nucleoside and at least one 2'-deoxynucleoside.

In certain embodiments, the 5'-wing of a gapmer comprises at least one LNA nucleoside and at least one non-bicyclic modified nucleoside. In certain embodiments, the 5'-wing of a gapmer comprises at least one LNA nucleoside and at least one 2'-substituted nucleoside. In certain embodiments, the 5'-wing of a gapmer comprises at least one LNA nucleoside and at least one 2'-MOE nucleoside. In certain embodiments, the 5'-wing of a gapmer comprises at least one LNA nucleoside and at least one 2'-OMe nucleoside. In certain embodiments, the 5'-wing of a gapmer comprises at least one LNA nucleoside and at least one 2'-deoxynucleoside.

In certain embodiments, the 5'-wing of a gapmer comprises at least one bicyclic nucleoside, at least one non-bicyclic modified nucleoside, and at least one 2'-deoxynucleoside. In certain embodiments, the 5'-wing of a gapmer comprises at least one constrained ethyl nucleoside, at least one non-bicyclic modified nucleoside, and at least one 2'-deoxynucleoside. In certain embodiments, the 5'-wing of a gapmer comprises at least one LNA nucleoside, at least one non-bicyclic modified nucleoside, and at least one 2'-deoxynucleoside.

In certain embodiments, the 5'-wing of a gapmer comprises at least one bicyclic nucleoside, at least one 2'-substituted nucleoside, and at least one 2'-deoxynucleoside. In certain embodiments, the 5'-wing of a gapmer comprises at least one constrained ethyl nucleoside, at least one 2'-substituted nucleoside, and at least one 2'-deoxynucleoside. In certain embodiments, the 5'-wing of a gapmer comprises at least one LNA nucleoside, at least one 2'-substituted nucleoside, and at least one 2'-deoxynucleoside.

In certain embodiments, the 5'-wing of a gapmer comprises at least one bicyclic nucleoside, at least one 2'-MOE nucleoside, and at least one 2'-deoxynucleoside. In certain embodiments, the 5'-wing of a gapmer comprises at least one constrained ethyl nucleoside, at least one 2'-MOE nucleoside, and at least one 2'-deoxynucleoside. In certain embodiments, the 5'-wing of a gapmer comprises at least one LNA nucleoside, at least one 2'-MOE nucleoside, and at least one 2'-deoxynucleoside.

In certain embodiments, the 5'-wing of a gapmer comprises at least one bicyclic nucleoside, at least one 2'-OMe nucleoside, and at least one 2'-deoxynucleoside. In certain embodiments, the 5'-wing of a gapmer comprises at least one constrained ethyl nucleoside, at least one 2'-OMe nucleoside, and at least one 2'-deoxynucleoside. In certain embodiments, the 5'-wing of a gapmer comprises at least one LNA nucleoside, at least one 2'-OMe nucleoside, and at least one 2'-deoxynucleoside.

In certain embodiments, the 5'- wing of a gapmer has a sugar motif selected from among those listed in the following non-limiting table:

Table 1
Certain 5'-Wing Sugar Motifs

5'-wing sugar motif #	motif	5'-wing sugar motif #	motif	5'-wing sugar motif #	motif
1a	A-B-B	1d	A-L-L	1g	A-K-K

2a	A-A-B	2d	A-A-L	2g	A-A-K
3a	A-D-B	3d	A-D-L	3g	A-D-K
4a	B-D-A	4d	L-D-A	4g	K-D-A
5a	B-A-A	5d	L-A-A	5g	K-A-A
6a	B-B-B	6d	L-L-L	6g	K-K-K
7a	A-A-A	7d	A-A-A	7g	A-A-A
8a	A-D-D-B	8d	A-D-D-L	8g	A-D-D-K
9a	B-D-D-A	9d	L-D-D-A	9g	K-D-D-A
10a	A-A-A-B	10d	A-A-A-L	10g	A-A-A-K
11a	B-A-A-A	11d	L-A-A-A	11g	K-A-A-A
12a	A-A-A-A	12d	A-A-A-A	12g	A-A-A-A
13a	B-D-D-B	13d	L-D-D-L	13g	K-D-D-K
14a	A-A-A-A	14d	A-A-A-A	14g	A-A-A-A
15a	B-B-B-B	15d	L-L-L-L	15g	K-K-K-K
16a	A-A-A-A-A	16d	A-A-A-A-A	16g	A-A-A-A-A
17a	A-D-A-D-B	17d	A-D-A-D-L	17g	A-D-A-D-K
18a	A-D-B-D-A	18d	A-D-L-D-A	18g	A-D-K-D-A
19a	B-D-A-D-A	19d	L-D-A-D-A	19g	K-D-A-D-A
20a	A-A-A-A-B	20d	A-A-A-A-L	20g	A-A-A-A-K
21a	A-A-B-A-A	21d	A-A-L-A-A	21g	A-A-K-A-A
22a	B-A-A-A-A	22d	L-A-A-A-A	22g	K-A-A-A-A
1b	E-B-B	1e	E-L-L	1h	E-K-K
2b	E-E-B	2e	E-E-L	2h	E-E-K
3b	E-D-B	3e	E-D-L	3h	E-D-K
4b	B-D-E	4e	L-D-E	4h	K-D-E
5b	B-E-E	5e	L-E-E	5h	K-E-E
6b	B-B-B	6e	L-L-L	6h	K-K-K
7b	E-E-E	7e	E-E-E	7h	E-E-E
8b	E-D-D-B	8e	E-D-D-L	8h	E-D-D-K
9b	B-D-D-E	9e	L-D-D-E	9h	K-D-D-E
10b	E-E-E-B	10e	E-E-E-L	10h	E-E-E-K
11b	B-E-E-E	11e	L-E-E-E	11h	K-E-E-E
12b	E-E-E-E	12e	E-E-E-E	12h	E-E-E-E
13b	B-D-D-B	13e	L-D-D-L	13h	K-D-D-K
14b	E-E-E-E	14e	E-E-E-E	14h	E-E-E-E
15b	B-B-B-B	15e	L-L-L-L	15h	K-K-K-K
16b	E-E-E-E-E	16e	E-E-E-E-E	16h	E-E-E-E-E
17b	E-D-E-D-B	17e	E-D-E-D-L	17h	E-D-E-D-K
18b	E-D-B-D-E	18e	E-D-L-D-E	18h	E-D-K-D-E
19b	B-D-E-D-E	19e	L-D-E-D-E	19h	K-D-E-D-E
20b	E-E-E-E-B	20e	E-E-E-E-L	20h	E-E-E-E-K
21b	E-E-B-E-E	21e	E-E-L-E-E	21h	E-E-K-E-E
22b	B-E-E-E-E	22e	L-E-E-E-E	22h	K-E-E-E-E
1c	M-B-B	1f	M-L-L	1i	M-K-K

2c	M-M-B	2f	M-M-L	2i	M-M-K
3c	M-D-B	3f	M-D-L	3i	M-D-K
4c	B-D-M	4f	L-D-M	4i	K-D-M
5c	B-M-M	5f	L-M-M	5i	K-M-M
6c	B-B-B	6f	L-L-L	6i	K-K-K
7c	M-M-M	7f	M-M-M	7i	M-M-M
8c	M-D-D-B	8f	M-D-D-L	8i	M-D-D-K
9c	B-D-D-M	9f	L-D-D-M	9i	K-D-D-M
10c	M-M-M-B	10f	M-M-M-L	10i	M-M-M-K
11c	B-M-M-M	11f	L-M-M-M	11i	K-M-M-M
12c	M-M-M-M	12f	M-M-M-M	12i	M-M-M-M
13c	B-D-D-B	13f	L-D-D-L	13i	K-D-D-K
14c	M-M-M-M	14f	M-M-M-M	14i	M-M-M-M
15c	B-B-B-B	15f	L-L-L-L	15i	K-K-K-K
16c	M-M-M-M-M	16f	M-M-M-M-M	16i	M-M-M-M-M
17c	M-D-M-D-B	17f	M-D-M-D-L	17i	M-D-M-D-K
18c	M-D-B-D-M	18f	M-D-L-D-M	18i	M-D-K-D-M
19c	B-D-M-D-M	19f	L-D-M-D-M	19i	K-D-M-D-M
20c	M-M-M-M-B	20f	M-M-M-M-L	20i	M-M-M-M-K
21c	M-M-B-M-M	21f	M-M-L-M-M	21i	M-M-K-M-M
22c	B-M-M-M-M	22f	L-M-M-M-M	22i	K-M-M-M-M
1j	A-L-K	1k	A-K-L	1l	E-L-K
2j	M-E-K	2k	M-E-L	2l	E-M-K
3j	L-D-K	3k	K-D-L	3l	B-D-K
4j	K-D-A	4k	L-D-K	4l	K-B-L
5j	B-M-E	5k	L-M-E	5l	K-M-E
6j	K-L-L	6k	L-K-L	6l	L-K-K
7j	E-M-E	7k	M-E-M	7l	M-E-E
8j	E-D-D-M	8k	K-D-D-L	8l	L-D-D-K
9j	M-D-D-E	9k	L-D-K-E	9l	K-D-L-E
10j	E-M-E-B	10k	E-M-E-L	10l	E-M-E-K
11j	B-E-E-M	11k	L-E-E-M	11l	K-E-E-M
12j	E-E-E-M	12k	M-E-E-E	12l	E-M-E-E
13j	K-L-D-K	13k	L-K-D-L	13l	K-D-L-K
14j	E-M-E-M	14k	M-EM-E	14l	E-E-M-E
15j	K-L-L-K	15k	L-K-L-K	15l	K-L-K-K
16j	E-E-M-E-E	16k	M-E-E-E-M	16l	E-E-M-M-E
17j	E-D-M-D-K	17k	E-D-M-D-L	17l	M-D-E-D-K
18j	E-D-K-D-M	18k	E-D-L-D-M	18l	M-D-K-D-E
19j	B-D-A-D-A	19k	L-D-A-D-A	19l	K-D-A-D-A
20j	E-M-E-E-L	20k	E-M-M-E-L	20l	M-E-E-E-K
21j	E-E-K-M-M	21k	E-E-L-M-M	21l	E-M-K-E-E
22j	B-E-M-E-A	22k	L-E-A-M-A	22l	K-E-A-A-A
23j	K-D-K-D-K	23k	E-K-E-K-D-K		

In the above table, “A” represents a nucleoside comprising a 2'-substituted sugar moiety; “B” represents a bicyclic nucleoside; “D” represents a 2'-deoxynucleoside; “K” represents a constrained ethyl nucleoside; “L” represents an LNA nucleoside; “E” represents a 2'-MOE nucleoside; and “M” represents a 2'-OMe nucleoside.

In certain embodiments, an oligonucleotide comprises any 5'-wing motif provided herein. In certain such embodiments, the oligonucleotide is a 5'-hemimer (does not comprise a 3'-wing). In certain embodiments, such an oligonucleotide is a gapmer. In certain such embodiments, the 3'-wing of the gapmer may comprise any sugar modification motif.

In certain embodiments, the 5'-wing of a gapmer has a sugar motif selected from among those listed in the following non-limiting tables:

Table 2
Certain 5'-Wing Sugar Motifs

Certain 5'-Wing Sugar Motifs				
AAAAA	ABCBB	BABCC	BCBBA	CBACC
AAAAB	ABCBC	BACAA	BCBBB	CBBA A
AAAAC	ABCCA	BACAB	BCBBC	CBBAB
AAABA	ABCCB	BACAC	BCBCA	CBBAC
AAABB	ABCCC	BACBA	BCBCB	CBBBA
AAABC	ACAAA	BACBB	BCBCC	CBBBB
AAACA	ACAAB	BACBC	BCCAA	CBBBC
AAACB	ACAAC	BACCA	BCCAB	CBBCA
AAACC	ACABA	BACCB	BCCAC	CBBCB
AABAA	ACABB	BACCC	BCCBA	CBGCC
AABAB	ACABC	BBAAA	BCCBB	CBCAA
AABAC	ACACA	BBAAB	BCCBC	CBCAB
AABBA	ACACB	BBAAC	BCCCA	CBCAC
AABBB	ACACC	BBABA	BCCCB	CBCBA
AABBC	ACBAA	BBABB	BCCCC	CBCBB
AABCA	ACBAB	BBABC	CAAAA	CBCBC
AABCB	ACBAC	BBACA	CAAAB	CBCCA
AABCC	ACBBA	BBACB	CAAAC	CBCCB
AACAA	ACBBB	BBACC	CAABA	CBCCC
AACAB	ACBBC	BBBAA	CAABB	CCAAA
AACAC	ACBCA	BBBAB	CAABC	CCAAB
AACBA	ACBCB	BBBAC	CAACA	CCAAC
AACBB	ACBCC	BBBBA	CAACB	CCABA
AACBC	ACCAA	BBBBB	CAACC	CCABB
AACCA	ACCAB	BBBBC	CABAA	CCABC
AACCB	ACCAC	BBBCA	CABAB	CCACA

AACCC	ACCBA	BBBCB	CABAC	CCACB
ABAAA	ACCBB	BBBCC	CABBA	CCACC
ABAAB	ACCBC	BBCAA	CABBB	CCBAA
ABAAC	ACCCA	BBCAB	CABBC	CCBAB
ABABA	ACCCB	BBCAC	CABCA	CCBAC
ABABB	ACCCC	BBCBA	CABCB	CCBBA
ABABC	BAAAA	BBCBB	CABCC	CCBBB
ABACA	BAAAB	BBCBC	CACAA	CCBBC
ABACB	BAAAC	BBCCA	CACAB	CCBCA
ABACC	BAABA	BBCCB	CACAC	CCBCB
ABBAA	BAABB	BBCCC	CACBA	CCBCC
ABBAB	BAABC	BCAAA	CACBB	CCCAA
ABBAC	BAACA	BCAAB	CACBC	CCCAB
ABBBA	BAACB	BCAAC	CACCA	CCCAC
ABBBB	BAACC	BCABA	CACCB	CCCBA
ABBBC	BABAA	BCABB	CACCC	CCCB
ABBCA	BABAB	BCABC	CBAAA	CCBC
ABBCB	BABAC	BCACA	CBAAB	CCCCA
ABGCC	BABBA	BCACB	CBAAC	CCCCB
ABCAA	BABBB	BCACC	CBABA	CCCCC
ABCAB	BABBC	BCBAA	CBABB	
ABCAC	BABCA	BCBAB	CBABC	
ABCBA	BABCB	BCBAC	CBACA	

Table 3
Certain 5'-Wing Sugar Motifs

Certain 5'-Wing Sugar Motifs				
AAAAA	BABC	CBAB	ABBB	BAA
AAAAB	BACA	CBAC	BAAA	BAB
AAABA	BACB	CBBA	BAAB	BBA
AAABB	BACC	CBBB	BABA	BBB
AABAA	BBAA	CBBC	BABB	AA
AABAB	BBAB	CBCA	BBAA	AB
AABBA	BBAC	CBCB	BBAB	AC
AABBB	BBBA	CBCC	BBBA	BA
ABAAA	BBBB	CCAA	BBBB	BB
ABAAB	BBBC	CCAB	AAA	BC
ABABA	BBCA	CCAC	AAB	CA
ABABB	BBBC	CCBA	AAC	CB
ABBAA	BBCC	CCBB	ABA	CC
ABBAB	BCAA	CCBC	ABB	AA
ABBBA	BCAB	CCCA	ABC	AB

ABBBB	BCAC	CCCB	ACA	BA
BAAAA	ABCB	BCBA	ACB	
BAAAB	ABCC	BCBB	ACC	
BAABA	ACAA	BCBC	BAA	
BAABB	ACAB	BCCA	BAB	
BABAA	ACAC	BCCB	BAC	
BABAB	ACBA	BCCC	BBA	
BABBA	ACBB	CAAA	BBB	
BABBB	ACBC	CAAB	BBC	
BBAAA	ACCA	CAAC	BCA	
BBAAB	ACCB	CABA	BCB	
BBABA	ACCC	CABB	BCC	
BBABB	BAAA	CABC	CAA	
BBBAA	BAAB	CACA	CAB	
BBBAB	BAAC	CACB	CAC	
BBBBA	BABA	CACC	CBA	
BBBBB	BABB	CBAA	CBB	
AAAA	AACC	CCCC	CBC	
AAAB	ABAA	AAAA	CCA	
AAAC	ABAB	AAAB	CCB	
AABA	ABAC	AABA	CCC	
AABB	ABBA	AABB	AAA	
AABC	ABBB	ABAA	AAB	
AACA	ABBC	ABAB	ABA	
AACB	ABCA	ABBA	ABB	

In certain embodiments, each A, each B, and each C located at the 3'-most 5'-wing nucleoside is a modified nucleoside. For example, in certain embodiments the 5'-wing motif is selected from among ABB, BBB, and CBB, wherein the underlined nucleoside represents the 3'-most 5'-wing nucleoside and wherein the underlined nucleoside is a modified nucleoside.

In certain embodiments, each A comprises an unmodified 2'-deoxyfuranose sugar moiety. In certain embodiments, each A comprises a modified sugar moiety. In certain embodiments, each A comprises a 2'-substituted sugar moiety. In certain embodiments, each A comprises a 2'-substituted sugar moiety selected from among F, ara-F, OCH₃ and O(CH₂)₂-OCH₃. In certain embodiments, each A comprises a bicyclic sugar moiety. In certain embodiments, each A comprises a bicyclic sugar moiety selected from among cEt, cMOE, LNA, α -L-LNA, ENA and 2'-thio LNA. In certain embodiments, each A comprises a modified nucleobase. In certain embodiments, each A comprises a modified nucleobase selected from among 2-thio-thymidine nucleoside and 5-propyne uridine nucleoside. In certain embodiments, each A comprises an HNA. In certain embodiments, each A comprises an F-HNA.

In certain embodiments, each B comprises an unmodified 2'-deoxyfuranose sugar moiety. In certain embodiments, each B comprises a modified sugar moiety. In certain embodiments, each B comprises a 2'-substituted sugar moiety. In certain embodiments, each B comprises a 2'-substituted sugar moiety selected from among F, (ara)-F, OCH₃ and O(CH₂)₂-OCH₃. In certain embodiments, each B comprises a bicyclic sugar moiety. In certain embodiments, each B comprises a bicyclic sugar moiety selected from among cEt, cMOE, LNA, α -L-LNA, ENA and 2'-thio LNA. In certain embodiments, each B comprises a modified nucleobase. In certain embodiments, each B comprises a modified nucleobase selected from among 2-thio-thymidine nucleoside and 5-propyne uridine nucleoside. In certain embodiments, each B comprises an HNA. In certain embodiments, each B comprises an F-HNA.

In certain embodiments, each C comprises an unmodified 2'-deoxyfuranose sugar moiety. In certain embodiments, each C comprises a modified sugar moiety. In certain embodiments, each C comprises a 2'-substituted sugar moiety. In certain embodiments, each C comprises a 2'-substituted sugar moiety selected from among F, (ara)-F, OCH₃ and O(CH₂)₂-OCH₃. In certain embodiments, each C comprises a 5'-substituted sugar moiety. In certain embodiments, each C comprises a 5'-substituted sugar moiety selected from among 5'-Me, and 5'-(R)-Me. In certain embodiments, each C comprises a bicyclic sugar moiety. In certain embodiments, each C comprises a bicyclic sugar moiety selected from among cEt, cMOE, LNA, α -L-LNA, ENA and 2'-thio LNA. In certain embodiments, each C comprises a modified nucleobase. In certain embodiments, each C comprises a modified nucleobase selected from among 2-thio-thymidine and 5-propyne uridine. In certain embodiments, each C comprises a 2-thio-thymidine nucleoside. In certain embodiments, each C comprises an HNA. In certain embodiments, each C comprises an F-HNA.

In certain embodiments, at least one of A or B comprises an unmodified 2'-deoxyfuranose sugar moiety, and the other comprises a 2'-substituted sugar moiety. In certain embodiments, at least one of A or B comprises an unmodified 2'-deoxyfuranose sugar moiety, and the other comprises a bicyclic sugar moiety.

In certain embodiments, at least one of A or B comprises a bicyclic sugar moiety, and the other comprises a 2'-substituted sugar moiety. In certain embodiments, one of A or B is an LNA nucleoside and the other of A or B comprises a 2'-substituted sugar moiety. In certain embodiments, one of A or B is a cEt nucleoside and the other of A or B comprises a 2'-substituted sugar moiety. In certain embodiments, one of A or B is an α -L-LNA nucleoside and the other of A or B comprises a 2'-substituted sugar moiety. In certain embodiments, one of A or B is an LNA nucleoside and the other of A or B comprises a 2'-MOE sugar moiety. In certain embodiments, one of A or B is a cEt nucleoside and the other of A or B comprises a 2'-MOE sugar moiety. In certain embodiments, one of A or B is an α -L-LNA nucleoside and the other of A or B comprises a 2'-MOE sugar moiety. In certain embodiments, one of A or B is an LNA nucleoside and the other of A or B comprises a 2'-F sugar moiety. In certain embodiments, one of A or B is a cEt nucleoside and the other of A or B comprises a 2'-F sugar moiety. In certain embodiments, one of A or B is an α -L-LNA nucleoside and the other of A or B comprises a 2'-F sugar moiety. In certain embodiments, one of A or B is an LNA nucleoside and the other of A or B comprises a 2'-(ara)-F sugar moiety. In certain

embodiments, one of A or B is a cEt nucleoside and the other of A or B comprises a 2'-(ara)-F sugar moiety. In certain embodiments, one of A or B is an α -L-LNA nucleoside and the other of A or B comprises a 2'-(ara)-F sugar moiety.

In certain embodiments, at least one of A or B comprises an unmodified 2'-deoxyfuranose sugar moiety, and the other comprises a 2'-substituted sugar moiety. In certain embodiments, one of A or B is an unmodified 2'-deoxyfuranose sugar moiety and the other of A or B comprises a 2'-substituted sugar moiety. In certain embodiments, one of A or B is an unmodified 2'-deoxyfuranose sugar moiety and the other of A or B comprises a 2'-MOE sugar moiety. In certain embodiments, one of A or B is an unmodified 2'-deoxyfuranose sugar moiety and the other of A or B comprises a 2'-F sugar moiety. In certain embodiments, one of A or B is an unmodified 2'-deoxyfuranose sugar moiety and the other of A or B comprises a 2'-(ara)-F sugar moiety. In certain embodiments, at least one of A or B comprises a bicyclic sugar moiety, and the other comprises an unmodified 2'-deoxyfuranose sugar moiety. In certain embodiments, one of A or B is an LNA nucleoside and the other of A or B comprises an unmodified 2'-deoxyfuranose sugar moiety. In certain embodiments, one of A or B is a cEt nucleoside and the other of A or B comprises an unmodified 2'-deoxyfuranose sugar moiety. In certain embodiments, one of A or B is an α -L-LNA nucleoside and the other of A or B comprises an unmodified 2'-deoxyfuranose sugar moiety. In certain embodiments, one of A or B is an LNA nucleoside and the other of A or B comprises an unmodified 2'-deoxyfuranose sugar moiety. In certain embodiments, one of A or B is a cEt nucleoside and the other of A or B comprises an unmodified 2'-deoxyfuranose sugar moiety. In certain embodiments, one of A or B is an α -L-LNA nucleoside and the other of A or B comprises an unmodified 2'-deoxyfuranose sugar moiety.

In certain embodiments, A comprises a bicyclic sugar moiety, and B comprises a 2'-substituted sugar moiety. In certain embodiments, A is an LNA nucleoside and B comprises a 2'-substituted sugar moiety. In certain embodiments, A is a cEt nucleoside and B comprises a 2'-substituted sugar moiety. In certain embodiments, A is an α -L-LNA nucleoside and B comprises a 2'-substituted sugar moiety.

In certain embodiments, A comprises a bicyclic sugar moiety, and B comprises a 2'-MOE sugar moiety. In certain embodiments, A is an LNA nucleoside and B comprises a 2'-MOE sugar moiety. In certain embodiments, A is a cEt nucleoside and B comprises a 2'-MOE sugar moiety. In certain embodiments, A is an α -L-LNA nucleoside and B comprises a 2'-MOE sugar moiety.

In certain embodiments, A comprises a bicyclic sugar moiety, and B comprises a 2'-F sugar moiety. In certain embodiments, A is an LNA nucleoside and B comprises a 2'-F sugar moiety. In certain embodiments, A is a cEt nucleoside and B comprises a 2'-F sugar moiety. In certain embodiments, A is an α -L-LNA nucleoside and B comprises a 2'-F sugar moiety.

In certain embodiments, A comprises a bicyclic sugar moiety, and B comprises a 2'-(ara)-F sugar moiety. In certain embodiments, A is an LNA nucleoside and B comprises a 2'-(ara)-F sugar moiety. In certain embodiments, A is a cEt nucleoside and B comprises a 2'-(ara)-F sugar moiety. In certain embodiments, A is an α -L-LNA nucleoside and B comprises a 2'-(ara)-F sugar moiety.

In certain embodiments, B comprises a bicyclic sugar moiety, and A comprises a 2'-MOE sugar moiety. In certain embodiments, B is an LNA nucleoside and A comprises a 2'-MOE sugar moiety. In certain embodiments, B is a cEt nucleoside and A comprises a 2'-MOE sugar moiety. In certain embodiments, B is an α -L-LNA nucleoside and A comprises a 2'-MOE sugar moiety.

In certain embodiments, B comprises a bicyclic sugar moiety, A comprises a 2'-MOE sugar moiety, and C comprises an unmodified 2'-deoxyfuranose sugar moiety. In certain embodiments, B is an LNA nucleoside, A comprises a 2'-MOE sugar moiety, and C comprises an unmodified 2'-deoxyfuranose sugar moiety. In certain embodiments, B is a cEt nucleoside, A comprises a 2'-MOE sugar moiety, and C comprises an unmodified 2'-deoxyfuranose sugar moiety. In certain embodiments, B is an α -L-LNA nucleoside and A comprises a 2'-MOE sugar moiety.

In certain embodiments, B comprises a bicyclic sugar moiety, and A comprises a 2'-F sugar moiety. In certain embodiments, B is an LNA nucleoside and A comprises a 2'-F sugar moiety. In certain embodiments, B is a cEt nucleoside and A comprises a 2'-F sugar moiety. In certain embodiments, B is an α -L-LNA nucleoside and A comprises a 2'-F sugar moiety.

In certain embodiments, B comprises a bicyclic sugar moiety, and A comprises a 2'-(ara)-F sugar moiety. In certain embodiments, B is an LNA nucleoside and A comprises a 2'-(ara)-F sugar moiety. In certain embodiments, B is a cEt nucleoside and A comprises a 2'-(ara)-F sugar moiety. In certain embodiments, B is an α -L-LNA nucleoside and A comprises a 2'-(ara)-F sugar moiety.

In certain embodiments, at least one of A or B comprises a bicyclic sugar moiety, another of A or B comprises a 2'-substituted sugar moiety and C comprises a modified nucleobase. In certain embodiments, one of A or B is an LNA nucleoside, another of A or B comprises a 2'-substituted sugar moiety, and C comprises a modified nucleobase. In certain embodiments, one of A or B is a cEt nucleoside, another of A or B comprises a 2'-substituted sugar moiety, and C comprises a modified nucleobase. In certain embodiments, one of A or B is an α -L-LNA nucleoside, another of A or B comprises a 2'-substituted sugar moiety, and C comprises a modified nucleobase.

In certain embodiments, one of A or B comprises a bicyclic sugar moiety, another of A or B comprises a 2'-MOE sugar moiety, and C comprises a modified nucleobase. In certain embodiments, one of A or B is an LNA nucleoside, another of A or B comprises a 2'-MOE sugar moiety, and C comprises a modified nucleobase. In certain embodiments, one of A or B is a cEt nucleoside, another of A or B comprises a 2'-MOE sugar moiety, and C comprises a modified nucleobase. In certain embodiments, one of A or B is an α -L-LNA nucleoside, another of A or B comprises a 2'-MOE sugar moiety, and C comprises a modified nucleobase.

In certain embodiments, one of A or B comprises a bicyclic sugar moiety, another of A or B comprises a 2'-F sugar moiety, and C comprises a modified nucleobase. In certain embodiments, one of A or B is an LNA nucleoside, another of A or B comprises a 2'-F sugar moiety, and C comprises a modified nucleobase. In certain embodiments, one of A or B is a cEt nucleoside, another of A or B comprises a 2'-F

sugar moiety, and C comprises a modified nucleobase. In certain embodiments, one of A or B is an α -L-LNA nucleoside, another of A or B comprises a 2'-F sugar moiety, and C comprises a modified nucleobase.

In certain embodiments, one of A or B comprises a bicyclic sugar moiety, another of A or B comprises a 2'-(ara)-F sugar moiety, and C comprises a modified nucleobase. In certain embodiments, one of A or B is an LNA nucleoside, another of A or B comprises a 2'-(ara)-F sugar moiety, and C comprises a modified nucleobase. In certain embodiments, one of A or B is a cEt nucleoside, another of A or B comprises a 2'-(ara)-F sugar moiety, and C comprises a modified nucleobase. In certain embodiments, one of A or B is an α -L-LNA nucleoside, another of A or B comprises a 2'-(ara)-F sugar moiety, and C comprises a modified nucleobase.

In certain embodiments, one of A or B comprises a bicyclic sugar moiety, another of A or B comprises a 2'-substituted sugar moiety, and C comprises a 2-thio-thymidine nucleobase. In certain embodiments, one of A or B is an LNA nucleoside, another of A or B comprises a 2'-substituted sugar moiety, and C comprises a 2-thio-thymidine nucleobase. In certain embodiments, one of A or B is a cEt nucleoside, another of A or B comprises a 2'-substituted sugar moiety, and C comprises a 2-thio-thymidine nucleobase. In certain embodiments, one of A or B is an α -L-LNA nucleoside, another of A or B comprises a 2'-substituted sugar moiety, and C comprises a 2-thio-thymidine nucleobase.

In certain embodiments, one of A or B comprises a bicyclic sugar moiety, another of A or B comprises a 2'-MOE sugar moiety, and C comprises a 2-thio-thymidine nucleobase. In certain embodiments, one of A or B is an LNA nucleoside, another of A or B comprises a 2'-MOE sugar moiety, and C comprises a 2-thio-thymidine nucleobase. In certain embodiments, one of A or B is a cEt nucleoside, another of A or B comprises a 2'-MOE sugar moiety, and C comprises a 2-thio-thymidine nucleobase. In certain embodiments, one of A or B is an α -L-LNA nucleoside, another of A or B comprises a 2'-MOE sugar moiety, and C comprises a 2-thio-thymidine nucleobase.

In certain embodiments, one of A or B comprises a bicyclic sugar moiety, another of A or B comprises a 2'-F sugar moiety, and C comprises a 2-thio-thymidine nucleobase. In certain embodiments, one of A or B is an LNA nucleoside, another of A or B comprises a 2'-F sugar moiety, and C comprises a 2-thio-thymidine nucleobase. In certain embodiments, one of A or B is a cEt nucleoside, another of A or B comprises a 2'-F sugar moiety, and C comprises a 2-thio-thymidine nucleobase. In certain embodiments, one of A or B is an α -L-LNA nucleoside, another of A or B comprises a 2'-F sugar moiety, and C comprises a 2-thio-thymidine nucleobase.

In certain embodiments, one of A or B comprises a bicyclic sugar moiety, another of A or B comprises a 2'-(ara)-F sugar moiety, and C comprises a 2-thio-thymidine nucleobase. In certain embodiments, one of A or B is an LNA nucleoside, another of A or B comprises a 2'-(ara)-F sugar moiety, and C comprises a 2-thio-thymidine nucleobase. In certain embodiments, one of A or B is a cEt nucleoside, another of A or B comprises a 2'-(ara)-F sugar moiety, and C comprises a 2-thio-thymidine nucleobase. In

certain embodiments, one of A or B is an α -L-LNA nucleoside, another of A or B comprises a 2'-(ara)-F sugar moiety, and C comprises 2-thio-thymidine nucleobase.

In certain embodiments, one of A or B comprises a bicyclic sugar moiety, another of A or B comprises a 2'-MOE sugar moiety, and C comprises a 5-propyne uridine nucleobase. In certain embodiments, one of A or B is an LNA nucleoside, another of A or B comprises a 2'-MOE sugar moiety, and C comprises a 5-propyne uridine nucleobase. In certain embodiments, one of A or B is a cEt nucleoside, another of A or B comprises a 2'-MOE sugar moiety, and C comprises a 5-propyne uridine nucleobase. In certain embodiments, one of A or B is an α -L-LNA nucleoside, another of A or B comprises a 2'-MOE sugar moiety, and C comprises a 5-propyne uridine nucleobase.

In certain embodiments, one of A or B comprises a bicyclic sugar moiety, another of A or B comprises a 2'-MOE sugar moiety, and C comprises an unmodified 2'-deoxyfuranose sugar moiety. In certain embodiments, one of A or B is an LNA nucleoside, another of A or B comprises a 2'-MOE sugar moiety, and C comprises a 5-propyne uridine nucleobase. In certain embodiments, one of A or B is a cEt nucleoside, another of A or B comprises a 2'-MOE sugar moiety, and C comprises a 5-propyne uridine nucleobase. In certain embodiments, one of A or B is an α -L-LNA nucleoside, another of A or B comprises a 2'-MOE sugar moiety, and C comprises a 5-propyne uridine nucleobase.

In certain embodiments, one of A or B comprises a bicyclic sugar moiety, another of A or B comprises a 2'-F sugar moiety, and C comprises a 5-propyne uridine nucleobase. In certain embodiments, one of A or B is an LNA nucleoside, another of A or B comprises a 2'-F sugar moiety, and C comprises a 5-propyne uridine nucleobase. In certain embodiments, one of A or B is a cEt nucleoside, another of A or B comprises a 2'-F sugar moiety, and C comprises a 5-propyne uridine nucleobase. In certain embodiments, one of A or B is an α -L-LNA nucleoside, another of A or B comprises a 2'-F sugar moiety, and C comprises a 5-propyne uridine nucleobase.

In certain embodiments, one of A or B comprises a bicyclic sugar moiety, another of A or B comprises a 2'-(ara)-F sugar moiety, and C comprises a 5-propyne uridine nucleobase. In certain embodiments, one of A or B is an LNA nucleoside, another of A or B comprises a 2'-(ara)-F sugar moiety, and C comprises a 5-propyne uridine nucleobase. In certain embodiments, one of A or B is a cEt nucleoside, another of A or B comprises a 2'-(ara)-F sugar moiety, and C comprises a 5-propyne uridine nucleobase. In certain embodiments, one of A or B is an α -L-LNA nucleoside, another of A or B comprises a 2'-(ara)-F sugar moiety, and C comprises a 5-propyne uridine nucleobase.

In certain embodiments, one of A or B comprises a bicyclic sugar moiety, another of A or B comprises a 2'-MOE sugar moiety, and C comprises a sugar surrogate. In certain embodiments, one of A or B is an LNA nucleoside, another of A or B comprises a 2'-MOE sugar moiety, and C comprises a sugar surrogate. In certain embodiments, one of A or B is a cEt nucleoside, another of A or B comprises a 2'-MOE sugar moiety, and C comprises a sugar surrogate. In certain embodiments, one of A or B is an α -L-LNA nucleoside, another of A or B comprises a 2'-MOE sugar moiety, and C comprises a sugar surrogate.

In certain embodiments, one of A or B comprises a bicyclic sugar moiety, another of A or B comprises a 2'-MOE sugar moiety, and C comprises a F-HNA sugar surrogate. In certain embodiments, one of A or B is an LNA nucleoside, another of A or B comprises a 2'-MOE sugar moiety, and C comprises a F-HNA sugar surrogate. In certain embodiments, one of A or B is a cEt nucleoside, another of A or B

comprises a 2'-MOE sugar moiety, and C comprises a F-HNA sugar surrogate. In certain embodiments, one of A or B is an α -L-LNA nucleoside, another of A or B comprises a 2'-MOE sugar moiety, and C comprises a F-HNA sugar surrogate.

In certain embodiments, one of A or B comprises a bicyclic sugar moiety, another of A or B comprises a 2'-F sugar moiety, and C comprises a F-HNA sugar surrogate. In certain embodiments, one of A or B is an LNA nucleoside, another of A or B comprises a 2'-F sugar moiety, and C comprises a F-HNA sugar surrogate. In certain embodiments, one of A or B is a cEt nucleoside, another of A or B comprises a 2'-F sugar moiety, and C comprises a F-HNA sugar surrogate. In certain embodiments, one of A or B is an α -L-LNA nucleoside, another of A or B comprises a 2'-F sugar moiety, and C comprises a F-HNA sugar surrogate.

In certain embodiments, one of A or B comprises a bicyclic sugar moiety, another of A or B comprises a 2'-(ara)-F sugar moiety, and C comprises a F-HNA sugar surrogate. In certain embodiments, one of A or B is an LNA nucleoside, another of A or B comprises a 2'-(ara)-F sugar moiety, and C comprises a F-HNA sugar surrogate. In certain embodiments, one of A or B is a cEt nucleoside, another of A or B comprises a 2'-(ara)-F sugar moiety, and C comprises a F-HNA sugar surrogate. In certain embodiments, one of A or B is an α -L-LNA nucleoside, another of A or B comprises a 2'-(ara)-F sugar moiety, and C comprises a F-HNA sugar surrogate.

In certain embodiments, one of A or B comprises a bicyclic sugar moiety, another of A or B comprises a 2'-MOE sugar moiety, and C comprises a 5'-Me DNA sugar moiety. In certain embodiments, one of A or B is an LNA nucleoside, another of A or B comprises a 2'-MOE sugar moiety, and C comprises a 5'-Me DNA sugar moiety. In certain embodiments, one of A or B is a cEt nucleoside, another of A or B comprises a 2'-MOE sugar moiety, and C comprises a 5'-Me DNA sugar moiety. In certain embodiments, one of A or B is an α -L-LNA nucleoside, another of A or B comprises a 2'-MOE sugar moiety, and C comprises a 5'-Me DNA sugar moiety.

In certain embodiments, one of A or B comprises a bicyclic sugar moiety, another of A or B comprises a 2'-F sugar moiety, and C comprises a 5'-Me DNA sugar moiety. In certain embodiments, one of A or B is an LNA nucleoside, another of A or B comprises a 2'-F sugar moiety, and C comprises a 5'-Me DNA sugar moiety. In certain embodiments, one of A or B is a cEt nucleoside, another of A or B comprises a 2'-F sugar moiety, and C comprises a 5'-Me DNA sugar moiety. In certain embodiments, one of A or B is an α -L-LNA nucleoside, another of A or B comprises a 2'-F sugar moiety, and C comprises a 5'-Me DNA sugar moiety.

In certain embodiments, one of A or B comprises a bicyclic sugar moiety, another of A or B comprises a 2'-(ara)-F sugar moiety, and C comprises a 5'-Me DNA sugar moiety. In certain embodiments, one of A or B is an LNA nucleoside, another of A or B comprises a 2'-(ara)-F sugar moiety, and C comprises a 5'-Me DNA sugar moiety. In certain embodiments, one of A or B is a cEt nucleoside, another of A or B comprises a 2'-(ara)-F sugar moiety, and C comprises a 5'-Me DNA sugar moiety. In certain embodiments,

one of A or B is an α -L-LNA nucleoside, another of A or B comprises a 2'-(ara)-F sugar moiety, and C comprises a 5'-Me DNA sugar moiety.

In certain embodiments, one of A or B comprises a bicyclic sugar moiety, another of A or B comprises a 2'-MOE sugar moiety, and C comprises a 5'-(R)-Me DNA sugar moiety. In certain embodiments, one of A or B is an LNA nucleoside, another of A or B comprises a 2'-MOE sugar moiety, and C comprises a 5'-(R)-Me DNA sugar moiety. In certain embodiments, one of A or B is a cEt nucleoside, another of A or B comprises a 2'-MOE sugar moiety, and C comprises a 5'-(R)-Me DNA sugar moiety. In certain embodiments, one of A or B is an α -L-LNA nucleoside, another of A or B comprises a 2'-MOE sugar moiety, and C comprises a 5'-(R)-Me DNA sugar moiety.

In certain embodiments, one of A or B comprises a bicyclic sugar moiety, another of A or B comprises a 2'-F sugar moiety, and C comprises a 5'-(R)-Me DNA sugar moiety. In certain embodiments, one of A or B is an LNA nucleoside, another of A or B comprises a 2'-F sugar moiety, and C comprises a 5'-(R)-Me DNA sugar moiety. In certain embodiments, one of A or B is a cEt nucleoside, another of A or B comprises a 2'-F sugar moiety, and C comprises a 5'-(R)-Me DNA sugar moiety. In certain embodiments, one of A or B is an α -L-LNA nucleoside, another of A or B comprises a 2'-F sugar moiety, and C comprises a 5'-(R)-Me DNA sugar moiety.

In certain embodiments, one of A or B comprises a bicyclic sugar moiety, another of A or B comprises a 2'-(ara)-F sugar moiety, and C comprises a 5'-(R)-Me DNA sugar moiety. In certain embodiments, one of A or B is an LNA nucleoside, another of A or B comprises a 2'-(ara)-F sugar moiety, and C comprises a 5'-(R)-Me DNA sugar moiety. In certain embodiments, one of A or B is a cEt nucleoside, another of A or B comprises a 2'-(ara)-F sugar moiety, and C comprises a 5'-(R)-Me DNA sugar moiety. In certain embodiments, one of A or B is an α -L-LNA nucleoside, another of A or B comprises a 2'-(ara)-F sugar moiety, and C comprises a 5'-(R)-Me DNA sugar moiety.

In certain embodiments, at least two of A, B or C comprises a 2'-substituted sugar moiety, and the other comprises a bicyclic sugar moiety. In certain embodiments, at least two of A, B or C comprises a bicyclic sugar moiety, and the other comprises a 2'-substituted sugar moiety.

In certain embodiments, at least two of A, B or C comprises a 2'-substituted sugar moiety, and the other comprises an unmodified 2'-deoxyfuranose sugar moiety.. In certain embodiments, at least two of A, B or C comprises a bicyclic sugar moiety, and the other comprises an unmodified 2'-deoxyfuranose sugar moiety.

Certain 3'-wings

In certain embodiments, the 3'- wing of a gapmer consists of 1 to 5 linked nucleosides. In certain embodiments, the 3'- wing of a gapmer consists of 2 to 5 linked nucleosides. In certain embodiments, the 3'- wing of a gapmer consists of 3 to 5 linked nucleosides. In certain embodiments, the 3'- wing of a gapmer consists of 4 or 5 linked nucleosides. In certain embodiments, the 3'- wing of a gapmer consists of 1 to 4 linked nucleosides. In certain embodiments, the 3'- wing of a gapmer consists of 1 to 3 linked nucleosides.

In certain embodiments, the 3'-wing of a gapmer consists of 1 or 2 linked nucleosides. In certain embodiments, the 3'-wing of a gapmer consists of 2 to 4 linked nucleosides. In certain embodiments, the 3'-wing of a gapmer consists of 2 or 3 linked nucleosides. In certain embodiments, the 3'-wing of a gapmer consists of 3 or 4 linked nucleosides. In certain embodiments, the 3'-wing of a gapmer consists of 1 nucleoside. In certain embodiments, the 3'-wing of a gapmer consists of 2 linked nucleosides. In certain embodiments, the 3'-wing of a gapmer consists of 3 linked nucleosides. In certain embodiments, the 3'-wing of a gapmer consists of 4 linked nucleosides. In certain embodiments, the 3'-wing of a gapmer consists of 5 linked nucleosides.

In certain embodiments, the 3'-wing of a gapmer comprises at least one bicyclic nucleoside. In certain embodiments, the 3'-wing of a gapmer comprises at least one constrained ethyl nucleoside. In certain embodiments, the 3'-wing of a gapmer comprises at least one LNA nucleoside. In certain embodiments, each nucleoside of the 3'-wing of a gapmer is a bicyclic nucleoside. In certain embodiments, each nucleoside of the 3'-wing of a gapmer is a constrained ethyl nucleoside. In certain embodiments, each nucleoside of the 3'-wing of a gapmer is a LNA nucleoside.

In certain embodiments, the 3'-wing of a gapmer comprises at least one non-bicyclic modified nucleoside. In certain embodiments, the 3'-wing of a gapmer comprises at least two non-bicyclic modified nucleosides. In certain embodiments, the 3'-wing of a gapmer comprises at least three non-bicyclic modified nucleosides. In certain embodiments, the 3'-wing of a gapmer comprises at least four non-bicyclic modified nucleosides. In certain embodiments, the 3'-wing of a gapmer comprises at least one 2'-substituted nucleoside. In certain embodiments, the 3'-wing of a gapmer comprises at least one 2'-MOE nucleoside. In certain embodiments, the 3'-wing of a gapmer comprises at least one 2'-OMe nucleoside. In certain embodiments, each nucleoside of the 3'-wing of a gapmer is a non-bicyclic modified nucleoside. In certain embodiments, each nucleoside of the 3'-wing of a gapmer is a 2'-substituted nucleoside. In certain embodiments, each nucleoside of the 3'-wing of a gapmer is a 2'-MOE nucleoside. In certain embodiments, each nucleoside of the 3'-wing of a gapmer is a 2'-OMe nucleoside.

In certain embodiments, the 3'-wing of a gapmer comprises at least one 2'-deoxynucleoside. In certain embodiments, each nucleoside of the 3'-wing of a gapmer is a 2'-deoxynucleoside. In certain embodiments, the 3'-wing of a gapmer comprises at least one ribonucleoside. In certain embodiments, each nucleoside of the 3'-wing of a gapmer is a ribonucleoside.

In certain embodiments, the 3'-wing of a gapmer comprises at least one bicyclic nucleoside and at least one non-bicyclic modified nucleoside. In certain embodiments, the 3'-wing of a gapmer comprises at least one bicyclic nucleoside and at least one 2'-substituted nucleoside. In certain embodiments, the 3'-wing of a gapmer comprises at least one bicyclic nucleoside and at least one 2'-MOE nucleoside. In certain embodiments, the 3'-wing of a gapmer comprises at least one bicyclic nucleoside and at least one 2'-OMe nucleoside. In certain embodiments, the 3'-wing of a gapmer comprises at least one bicyclic nucleoside and at least one 2'-deoxynucleoside.

In certain embodiments, the 3'-wing of a gapmer comprises at least one constrained ethyl nucleoside and at least one non-bicyclic modified nucleoside. In certain embodiments, the 3'-wing of a gapmer comprises at least one constrained ethyl nucleoside and at least one 2'-substituted nucleoside. In certain embodiments, the 3'-wing of a gapmer comprises at least one constrained ethyl nucleoside and at least one 2'-MOE nucleoside. In certain embodiments, the 3'-wing of a gapmer comprises at least one constrained ethyl nucleoside and at least one 2'-OMe nucleoside. In certain embodiments, the 3'-wing of a gapmer comprises at least one constrained ethyl nucleoside and at least one 2'-deoxynucleoside.

In certain embodiments, the 3'-wing of a gapmer comprises at least one LNA nucleoside and at least one non-bicyclic modified nucleoside. In certain embodiments, the 3'-wing of a gapmer comprises at least one LNA nucleoside and at least one 2'-substituted nucleoside. In certain embodiments, the 3'-wing of a gapmer comprises at least one LNA nucleoside and at least one 2'-MOE nucleoside. In certain embodiments, the 3'-wing of a gapmer comprises at least one LNA nucleoside and at least one 2'-OMe nucleoside. In certain embodiments, the 3'-wing of a gapmer comprises at least one LNA nucleoside and at least one 2'-deoxynucleoside.

In certain embodiments, the 3'-wing of a gapmer comprises at least one bicyclic nucleoside, at least one non-bicyclic modified nucleoside, and at least one 2'-deoxynucleoside. In certain embodiments, the 3'-wing of a gapmer comprises at least one constrained ethyl nucleoside, at least one non-bicyclic modified nucleoside, and at least one 2'-deoxynucleoside. In certain embodiments, the 3'-wing of a gapmer comprises at least one LNA nucleoside, at least one non-bicyclic modified nucleoside, and at least one 2'-deoxynucleoside.

In certain embodiments, the 3'-wing of a gapmer comprises at least one bicyclic nucleoside, at least one 2'-substituted nucleoside, and at least one 2'-deoxynucleoside. In certain embodiments, the 3'-wing of a gapmer comprises at least one constrained ethyl nucleoside, at least one 2'-substituted nucleoside, and at least one 2'-deoxynucleoside. In certain embodiments, the 3'-wing of a gapmer comprises at least one LNA nucleoside, at least one 2'-substituted nucleoside, and at least one 2'-deoxynucleoside.

In certain embodiments, the 3'-wing of a gapmer comprises at least one bicyclic nucleoside, at least one 2'-MOE nucleoside, and at least one 2'-deoxynucleoside. In certain embodiments, the 3'-wing of a gapmer comprises at least one constrained ethyl nucleoside, at least one 2'-MOE nucleoside, and at least one 2'-deoxynucleoside. In certain embodiments, the 3'-wing of a gapmer comprises at least one LNA nucleoside, at least one 2'-MOE nucleoside, and at least one 2'-deoxynucleoside.

In certain embodiments, the 3'-wing of a gapmer comprises at least one bicyclic nucleoside, at least one 2'-OMe nucleoside, and at least one 2'-deoxynucleoside. In certain embodiments, the 3'-wing of a gapmer comprises at least one constrained ethyl nucleoside, at least one 2'-OMe nucleoside, and at least one 2'-deoxynucleoside. In certain embodiments, the 3'-wing of a gapmer comprises at least one LNA nucleoside, at least one 2'-OMe nucleoside, and at least one 2'-deoxynucleoside.

In certain embodiments, the 3'-wing of a gapmer has a sugar motif selected from among those listed in the following non-limiting table:

Table 4
Certain 3'-Wing Sugar Motifs

3'-wing sugar motif #	motif	3'-wing sugar motif #	motif	3'-wing sugar motif #	motif
1a	B-B-A	1d	L-L-A	1g	K-K-A
2a	B-B-B	2d	L-L-L	2g	K-K-K
3a	A-A-B	3d	A-A-L	3g	A-A-K
4a	B-A-B	4d	L-A-L	4g	K-A-K
5a	B-A-B-A	5d	L-A-L-A	5g	K-A-K-A
6a	B-B-B-A	6d	L-L-L-A	6g	K-K-K-A
7a	B-D-B-A	7d	L-D-L-A	7g	K-D-K-A
8a	B-B-B-B	8d	L-L-L-L	8g	K-K-K-K
9a	B-D-D-B	9d	L-D-D-L	9g	K-D-D-K
10a	A-B-B-A	10d	A-L-L-A	10g	A-K-K-A
1b	B-B-E	1e	L-L-E	1h	K-K-E
2b	B-B-B	2e	L-L-L	2h	K-K-K
3b	E-E-B	3e	E-E-L	3h	E-E-K
4b	B-E-B	4e	L-E-L	4h	K-E-K
5b	B-E-B-E	5e	L-E-L-E	5h	K-E-K-E
6b	B-B-B-E	6e	L-L-L-E	6h	K-K-K-E
7b	B-D-B-E	7e	L-D-L-E	7h	K-D-K-E
8b	B-B-B-B	8e	L-L-L-L	8h	K-K-K-K
9b	B-D-D-B	9e	L-D-D-L	9h	K-D-D-K
10b	E-B-B-E	10e	E-L-L-E	10h	E-K-K-E
1c	B-B-M	1f	L-L-M	1i	K-K-M
2c	B-B-B	2f	L-L-L	2i	K-K-K
3c	M-M-B	3f	M-M-L	3i	M-M-K
4c	B-M-B	4f	L-M-L	4i	K-M-K
5c	B-M-B-M	5f	L-M-L-M	5i	K-M-K-M
6c	B-B-B-M	6f	L-L-L-M	6i	K-K-K-M
7c	B-D-B-M	7f	L-D-L-M	7i	K-D-K-M
8c	B-B-B-B	8f	L-L-L-L	8i	K-K-K-K
9c	B-D-D-B	9f	L-D-D-L	9i	K-D-D-K
10c	M-B-B-M	10f	M-L-L-M	10i	M-K-K-M
1j	K-K-A	1k	L-K-A	1l	K-L-E
2j	K-L-L	2k	K-K-L	2l	K-L-K
3j	E-M-B	3k	E-M-L	3l	E-K-K
4j	K-A-L	4k	L-A-K	4l	L-E-K
5j	K-A-L-A	5k	L-A-K-A	5l	K-E-L-E
6j	K-L-K-A	6k	K-K-L-A	6l	K-L-K-A

7j	L-D-K-A	7k	K-D-L-A	7l	K-D-L-E
8j	B-K-L-B	8k	K-L-L-L	8l	K-K-L-K
9j	K-D-D-B	9k	K-D-D-L	9l	L-D-D-K
10j	A-K-B-A	10k	A-K-L-A	10l	A-B-K-A
1m	E-E				

In the above table, “A” represents a nucleoside comprising a 2’-substituted sugar moiety; “B” represents a bicyclic nucleoside; “D” represents a 2’-deoxynucleoside; “K” represents a constrained ethyl nucleoside; “L” represents an LNA nucleoside; “E” represents a 2’-MOE nucleoside; and “M” represents a 2’-OMe nucleoside.

In certain embodiments, an oligonucleotide comprises any 3’-wing motif provided herein. In certain such embodiments, the oligonucleotide is a 3’-hemimer (does not comprise a 5’-wing). In certain embodiments, such an oligonucleotide is a gapmer. In certain such embodiments, the 5’-wing of the gapmer may comprise any sugar modification motif.

In certain embodiments, the 5’-wing of a gapmer has a sugar motif selected from among those listed in the following non-limiting tables:

Table 5
Certain 3’-Wing Sugar Motifs

Certain 3’-Wing Sugar Motifs				
AAAAA	ABCBB	BABCC	BCBBA	CBACC
AAAAB	ABCBC	BACAA	BCBBB	CBBA A
AAAAC	ABCCA	BACAB	BCBBC	CBBAB
AAABA	ABCCB	BACAC	BCBCA	CBBAC
AAABB	ABCCC	BACBA	BCBCB	CBBBA
AAABC	ACAAA	BACBB	BCBCC	CBBBB
AAACA	ACAAB	BACBC	BCCAA	CBBBC
AAACB	ACAAC	BACCA	BCCAB	CBBCA
AAACC	ACABA	BACCB	BCCAC	CBBCB
AABAA	ACABB	BACCC	BCCBA	CBBC C
AABAB	ACABC	BBAAA	BCCBB	CBCAA
AABAC	ACACA	BBAAB	BCCBC	CBCAB
AABBA	ACACB	BBAAC	BCCCA	CBCAC
AABBB	ACACC	BBABA	BCCCB	CBCBA
AABBC	ACBAA	BBABB	BCCCC	CBCBB
AABCA	ACBAB	BBABC	CAAAA	CBCBC
AABCB	ACBAC	BBACA	CAAAB	CBCCA
AABCC	ACBBA	BBACB	CAAAC	CBCCB
AACAA	ACBBB	BBACC	CAABA	CBCCC
AACAB	ACBBC	BBBAA	CAABB	CCAAA
AACAC	ACBCA	BBBAB	CAABC	CCAAB

AACBA	ACBCB	BBBAC	CAACA	CCAAC
AACBB	ACBCC	BBBBA	CAACB	CCABA
AACBC	ACCAA	BBBBB	CAACC	CCABB
AACCA	ACCAB	BBBBC	CABAA	CCABC
AACCB	ACCAC	BBBCA	CABAB	CCACA
AACCC	ACCBA	BBBCB	CABAC	CCACB
ABAAA	ACCB	BBBCC	CABBA	CCACC
ABAAB	ACCBC	BBCAA	CABBB	CCBAA
ABAAC	ACCCA	BBCAB	CABBC	CCBAB
ABABA	ACCCB	BBCAC	CABCA	CCBAC
ABABB	ACCCC	BBCBA	CABCB	CCBBA
ABABC	BAAAA	BBCBB	CABCC	CCBBB
ABACA	BAAAB	BBCBC	CACAA	CCBBC
ABACB	BAAAC	BBCCA	CACAB	CCBCA
ABACC	BAABA	BBCCB	CACAC	CCBCB
ABBAA	BAABB	BBCCC	CACBA	CCBCC
ABBAB	BAABC	BCAAA	CACBB	CCCAA
ABBAC	BAACA	BCAAB	CACBC	CC CAB
ABBBA	BAACB	BCAAC	CACCA	CCCAC
ABBBB	BAACC	BCABA	CACCB	CCCBA
ABBBC	BABAA	BCABB	CACCC	CCCB
ABBCA	BABAB	BCABC	CBAAA	CCCBC
ABBCB	BABAC	BCACA	CBAAB	CCCCA
ABBCC	BABBA	BCACB	CBAAC	CCCCB
ABCAA	BABBB	BCACC	CBABA	CCCCC
ABCAB	BABBC	BCBAA	CBABB	
ABCAC	BABCA	BCBAB	CBABC	
ABCBA	BABCB	BCBAC	CBACA	

Table 6
Certain 3'-Wing Sugar Motifs

Certain 3'-Wing Sugar Motifs				
AAAAA	BABC	CBAB	ABBB	BAA
AAAAB	BACA	CBAC	BAAA	BAB
AAABA	BACB	CBBA	BAAB	BBA
AAABB	BACC	CBBB	BABA	BBB
AABAA	BBAA	CBBC	BABB	AA
AABAB	BBAB	CBCA	BBAA	AB
AABBA	BBAC	CBCB	BBAB	AC
AABBB	BBBA	CBCC	BBBA	BA
ABAAA	BBBB	CCAA	BBBB	BB
ABAAB	BBBC	CCAB	AAA	BC

ABABA	BBCA	CCAC	AAB	CA
ABABB	BBCB	CCBA	AAC	CB
ABBAA	BBCC	CCBB	ABA	CC
ABBAB	BCAA	CCBC	ABB	AA
ABBBA	BCAB	CCCA	ABC	AB
ABBBB	BCAC	CCCB	ACA	BA
BAAAA	ABCB	BCBA	ACB	
BAAAB	ABCC	BCBB	ACC	
BAABA	ACAA	BCBC	BAA	
BAABB	ACAB	BCCA	BAB	
BABAA	ACAC	BCCB	BAC	
BABAB	ACBA	BCCC	BBA	
BABBA	ACBB	CAAA	BBB	
BABBB	ACBC	CAAB	BBC	
BBAAA	ACCA	CAAC	BCA	
BBAAB	ACCB	CABA	BCB	
BBABA	ACCC	CABB	BCC	
BBABB	BAAA	CABC	CAA	
BBBAA	BAAB	CACA	CAB	
BBBAB	BAAC	CACB	CAC	
BBBBA	BABA	CACC	CBA	
BBBBB	BABB	CBAA	CBB	
AAAA	AACC	CCCC	CBC	
AAAB	ABAA	AAAA	CCA	
AAAC	ABAB	AAAB	CCB	
AABA	ABAC	AABA	CCC	
AABB	ABBA	AABB	AAA	
AABC	ABBB	ABAA	AAB	
AACA	ABBC	ABAB	ABA	
AACB	ABCA	ABBA	ABB	

In certain embodiments, each A, each B, and each C located at the 5'-most 3'-wing region nucleoside is a modified nucleoside. For example, in certain embodiments the 3'-wing motif is selected from among ABB, BBB, and CBB, wherein the underlined nucleoside represents the the 5'-most 3'-wing region nucleoside and wherein the underlined nucleoside is a modified nucleoside.

In certain embodiments, each A comprises an unmodified 2'-deoxyfuranose sugar moiety. In certain embodiments, each A comprises a modified sugar moiety. In certain embodiments, each A comprises a 2'-substituted sugar moiety. In certain embodiments, each A comprises a 2'-substituted sugar moiety selected from among F, ara-F, OCH₃ and O(CH₂)₂-OCH₃. In certain embodiments, each A comprises a bicyclic sugar moiety. In certain embodiments, each A comprises a bicyclic sugar moiety selected from among cEt, cMOE,

LNA, α -L-LNA, ENA and 2'-thio LNA. In certain embodiments, each A comprises a modified nucleobase. In certain embodiments, each A comprises a modified nucleobase selected from among 2-thio-thymidine nucleoside and 5-propyne uridine nucleoside.

In certain embodiments, each B comprises an unmodified 2'-deoxyfuranose sugar moiety. In certain embodiments, each B comprises a modified sugar moiety. In certain embodiments, each B comprises a 2'-substituted sugar moiety. In certain embodiments, each B comprises a 2'-substituted sugar moiety selected from among F, (ara)-F, OCH₃ and O(CH₂)₂-OCH₃. In certain embodiments, each B comprises a bicyclic sugar moiety. In certain embodiments, each B comprises a bicyclic sugar moiety selected from among cEt, cMOE, LNA, α -L-LNA, ENA and 2'-thio LNA. In certain embodiments, each B comprises a modified nucleobase. In certain embodiments, each B comprises a modified nucleobase selected from among 2-thio-thymidine nucleoside and 5-propyne uridine nucleoside.

In certain embodiments, each C comprises an unmodified 2'-deoxyfuranose sugar moiety. In certain embodiments, each C comprises a modified sugar moiety. In certain embodiments, each C comprises a 2'-substituted sugar moiety. In certain embodiments, each C comprises a 2'-substituted sugar moiety selected from among F, (ara)-F, OCH₃ and O(CH₂)₂-OCH₃. In certain embodiments, each C comprises a 5'-substituted sugar moiety. In certain embodiments, each C comprises a 5'-substituted sugar moiety selected from among 5'-Me, and 5'-(R)-Me. In certain embodiments, each C comprises a bicyclic sugar moiety. In certain embodiments, each C comprises a bicyclic sugar moiety selected from among cEt, cMOE, LNA, α -L-LNA, ENA and 2'-thio LNA. In certain embodiments, each C comprises a modified nucleobase. In certain embodiments, each C comprises a modified nucleobase selected from among 2-thio-thymidine and 5-propyne uridine. In certain embodiments, each C comprises a 2-thio-thymidine nucleoside.

In certain embodiments, at least one of A or B comprises an unmodified 2'-deoxyfuranose sugar moiety, and the other comprises a 2'-substituted sugar moiety. In certain embodiments, at least one of A or B comprises an unmodified 2'-deoxyfuranose sugar moiety, and the other comprises a bicyclic sugar moiety.

In certain embodiments, at least one of A or B comprises a bicyclic sugar moiety, and the other comprises a 2'-substituted sugar moiety. In certain embodiments, one of A or B is an LNA nucleoside and the other of A or B comprises a 2'-substituted sugar moiety. In certain embodiments, one of A or B is a cEt nucleoside and the other of A or B comprises a 2'-substituted sugar moiety. In certain embodiments, one of A or B is an α -L-LNA nucleoside and the other of A or B comprises a 2'-substituted sugar moiety. In certain embodiments, one of A or B is an LNA nucleoside and the other of A or B comprises a 2'-MOE sugar moiety. In certain embodiments, one of A or B is a cEt nucleoside and the other of A or B comprises a 2'-MOE sugar moiety. In certain embodiments, one of A or B is an α -L-LNA nucleoside and the other of A or B comprises a 2'-MOE sugar moiety. In certain embodiments, one of A or B is an LNA nucleoside and the other of A or B comprises a 2'-F sugar moiety. In certain embodiments, one of A or B is a cEt nucleoside and the other of A or B comprises a 2'-F sugar moiety. In certain embodiments, one of A or B is an α -L-LNA nucleoside and the other of A or B comprises a 2'-F sugar moiety. In certain embodiments, one of A or

B is an LNA nucleoside and the other of A or B comprises a 2'-(ara)-F sugar moiety. In certain embodiments, one of A or B is a cEt nucleoside and the other of A or B comprises a 2'-(ara)-F sugar moiety. In certain embodiments, one of A or B is an α -L-LNA nucleoside and the other of A or B comprises a 2'-(ara)-F sugar moiety.

In certain embodiments, at least one of A or B comprises an unmodified 2'-deoxyfuranose sugar moiety, and the other comprises a 2'-substituted sugar moiety. In certain embodiments, one of A or B is an unmodified 2'-deoxyfuranose sugar moiety and the other of A or B comprises a 2'-substituted sugar moiety. In certain embodiments, one of A or B is an unmodified 2'-deoxyfuranose sugar moiety and the other of A or B comprises a 2'-MOE sugar moiety. In certain embodiments, one of A or B is an unmodified 2'-deoxyfuranose sugar moiety and the other of A or B comprises a 2'-F sugar moiety. In certain embodiments, one of A or B is an unmodified 2'-deoxyfuranose sugar moiety and the other of A or B comprises a 2'-(ara)-F sugar moiety. In certain embodiments, at least one of A or B comprises a bicyclic sugar moiety, and the other comprises an unmodified 2'-deoxyfuranose sugar moiety. In certain embodiments, one of A or B is an LNA nucleoside and the other of A or B comprises an unmodified 2'-deoxyfuranose sugar moiety. In certain embodiments, one of A or B is a cEt nucleoside and the other of A or B comprises an unmodified 2'-deoxyfuranose sugar moiety. In certain embodiments, one of A or B is an α -L-LNA nucleoside and the other of A or B comprises an unmodified 2'-deoxyfuranose sugar moiety. In certain embodiments, one of A or B is an LNA nucleoside and the other of A or B comprises an unmodified 2'-deoxyfuranose sugar moiety. In certain embodiments, one of A or B is a cEt nucleoside and the other of A or B comprises an unmodified 2'-deoxyfuranose sugar moiety. In certain embodiments, one of A or B is an α -L-LNA nucleoside and the other of A or B comprises an unmodified 2'-deoxyfuranose sugar moiety.

In certain embodiments, A comprises a bicyclic sugar moiety, and B comprises a 2'-substituted sugar moiety. In certain embodiments, A is an LNA nucleoside and B comprises a 2'-substituted sugar moiety. In certain embodiments, A is a cEt nucleoside and B comprises a 2'-substituted sugar moiety. In certain embodiments, A is an α -L-LNA nucleoside and B comprises a 2'-substituted sugar moiety.

In certain embodiments, A comprises a bicyclic sugar moiety, and B comprises a 2'-MOE sugar moiety. In certain embodiments, A is an LNA nucleoside and B comprises a 2'-MOE sugar moiety. In certain embodiments, A is a cEt nucleoside and B comprises a 2'-MOE sugar moiety. In certain embodiments, A is an α -L-LNA nucleoside and B comprises a 2'-MOE sugar moiety.

In certain embodiments, A comprises a bicyclic sugar moiety, and B comprises a 2'-F sugar moiety. In certain embodiments, A is an LNA nucleoside and B comprises a 2'-F sugar moiety. In certain embodiments, A is a cEt nucleoside and B comprises a 2'-F sugar moiety. In certain embodiments, A is an α -L-LNA nucleoside and B comprises a 2'-F sugar moiety.

In certain embodiments, A comprises a bicyclic sugar moiety, and B comprises a 2'-(ara)-F sugar moiety. In certain embodiments, A is an LNA nucleoside and B comprises a 2'-(ara)-F sugar moiety. In

certain embodiments, A is a cEt nucleoside and B comprises a 2'-(ara)-F sugar moiety. In certain embodiments, A is an α -L-LNA nucleoside and B comprises a 2'-(ara)-F sugar moiety.

In certain embodiments, B comprises a bicyclic sugar moiety, and A comprises a 2'-MOE sugar moiety. In certain embodiments, B is an LNA nucleoside and A comprises a 2'-MOE sugar moiety. In certain embodiments, B is a cEt nucleoside and A comprises a 2'-MOE sugar moiety. In certain embodiments, B is an α -L-LNA nucleoside and A comprises a 2'-MOE sugar moiety.

In certain embodiments, B comprises a bicyclic sugar moiety, A comprises a 2'-MOE sugar moiety, and C comprises an unmodified 2'-deoxyfuranose sugar moiety. In certain embodiments, B is an LNA nucleoside, A comprises a 2'-MOE sugar moiety, and C comprises an unmodified 2'-deoxyfuranose sugar moiety. In certain embodiments, B is a cEt nucleoside, A comprises a 2'-MOE sugar moiety, and C comprises an unmodified 2'-deoxyfuranose sugar moiety. In certain embodiments, B is an α -L-LNA nucleoside and A comprises a 2'-MOE sugar moiety.

In certain embodiments, B comprises a bicyclic sugar moiety, and A comprises a 2'-F sugar moiety. In certain embodiments, B is an LNA nucleoside and A comprises a 2'-F sugar moiety. In certain embodiments, B is a cEt nucleoside and A comprises a 2'-F sugar moiety. In certain embodiments, B is an α -L-LNA nucleoside and A comprises a 2'-F sugar moiety.

In certain embodiments, B comprises a bicyclic sugar moiety, and A comprises a 2'-(ara)-F sugar moiety. In certain embodiments, B is an LNA nucleoside and A comprises a 2'-(ara)-F sugar moiety. In certain embodiments, B is a cEt nucleoside and A comprises a 2'-(ara)-F sugar moiety. In certain embodiments, B is an α -L-LNA nucleoside and A comprises a 2'-(ara)-F sugar moiety.

In certain embodiments, at least one of A or B comprises a bicyclic sugar moiety, another of A or B comprises a 2'-substituted sugar moiety and C comprises a modified nucleobase. In certain embodiments, one of A or B is an LNA nucleoside, another of A or B comprises a 2'-substituted sugar moiety, and C comprises a modified nucleobase. In certain embodiments, one of A or B is a cEt nucleoside, another of A or B comprises a 2'-substituted sugar moiety, and C comprises a modified nucleobase. In certain embodiments, one of A or B is an α -L-LNA nucleoside, another of A or B comprises a 2'-substituted sugar moiety, and C comprises a modified nucleobase.

In certain embodiments, one of A or B comprises a bicyclic sugar moiety, another of A or B comprises a 2'-MOE sugar moiety, and C comprises a modified nucleobase. In certain embodiments, one of A or B is an LNA nucleoside, another of A or B comprises a 2'-MOE sugar moiety, and C comprises a modified nucleobase. In certain embodiments, one of A or B is a cEt nucleoside, another of A or B comprises a 2'-MOE sugar moiety, and C comprises a modified nucleobase. In certain embodiments, one of A or B is an α -L-LNA nucleoside, another of A or B comprises a 2'-MOE sugar moiety, and C comprises a modified nucleobase.

In certain embodiments, one of A or B comprises a bicyclic sugar moiety, another of A or B comprises a 2'-F sugar moiety, and C comprises a modified nucleobase. In certain embodiments, one of A or

B is an LNA nucleoside, another of A or B comprises a 2'-F sugar moiety, and C comprises a modified nucleobase. In certain embodiments, one of A or B is a cEt nucleoside, another of A or B comprises a 2'-F sugar moiety, and C comprises a modified nucleobase. In certain embodiments, one of A or B is an α -L-LNA nucleoside, another of A or B comprises a 2'-F sugar moiety, and C comprises a modified nucleobase.

In certain embodiments, one of A or B comprises a bicyclic sugar moiety, another of A or B comprises a 2'-(ara)-F sugar moiety, and C comprises a modified nucleobase. In certain embodiments, one of A or B is an LNA nucleoside, another of A or B comprises a 2'-(ara)-F sugar moiety, and C comprises a modified nucleobase. In certain embodiments, one of A or B is a cEt nucleoside, another of A or B comprises a 2'-(ara)-F sugar moiety, and C comprises a modified nucleobase. In certain embodiments, one of A or B is an α -L-LNA nucleoside, another of A or B comprises a 2'-(ara)-F sugar moiety, and C comprises a modified nucleobase.

In certain embodiments, one of A or B comprises a bicyclic sugar moiety, another of A or B comprises a 2'-substituted sugar moiety, and C comprises a 2-thio-thymidine nucleobase. In certain embodiments, one of A or B is an LNA nucleoside, another of A or B comprises a 2'-substituted sugar moiety, and C comprises a 2-thio-thymidine nucleobase. In certain embodiments, one of A or B is a cEt nucleoside, another of A or B comprises a 2'-substituted sugar moiety, and C comprises a 2-thio-thymidine nucleobase. In certain embodiments, one of A or B is an α -L-LNA nucleoside, another of A or B comprises a 2'-substituted sugar moiety, and C comprises a 2-thio-thymidine nucleobase.

In certain embodiments, one of A or B comprises a bicyclic sugar moiety, another of A or B comprises a 2'-MOE sugar moiety, and C comprises a 2-thio-thymidine nucleobase. In certain embodiments, one of A or B is an LNA nucleoside, another of A or B comprises a 2'-MOE sugar moiety, and C comprises a 2-thio-thymidine nucleobase. In certain embodiments, one of A or B is a cEt nucleoside, another of A or B comprises a 2'-MOE sugar moiety, and C comprises a 2-thio-thymidine nucleobase. In certain embodiments, one of A or B is an α -L-LNA nucleoside, another of A or B comprises a 2'-MOE sugar moiety, and C comprises a 2-thio-thymidine nucleobase.

In certain embodiments, one of A or B comprises a bicyclic sugar moiety, another of A or B comprises a 2'-F sugar moiety, and C comprises a 2-thio-thymidine nucleobase. In certain embodiments, one of A or B is an LNA nucleoside, another of A or B comprises a 2'-F sugar moiety, and C comprises a 2-thio-thymidine nucleobase. In certain embodiments, one of A or B is a cEt nucleoside, another of A or B comprises a 2'-F sugar moiety, and C comprises a 2-thio-thymidine nucleobase. In certain embodiments, one of A or B is an α -L-LNA nucleoside, another of A or B comprises a 2'-F sugar moiety, and C comprises a 2-thio-thymidine nucleobase.

In certain embodiments, one of A or B comprises a bicyclic sugar moiety, another of A or B comprises a 2'-(ara)-F sugar moiety, and C comprises a 2-thio-thymidine nucleobase. In certain embodiments, one of A or B is an LNA nucleoside, another of A or B comprises a 2'-(ara)-F sugar moiety, and C comprises a 2-thio-thymidine nucleobase. In certain embodiments, one of A or B is a cEt nucleoside,

another of A or B comprises a 2'-(ara)-F sugar moiety, and C comprises a 2-thio-thymidine nucleobase. In certain embodiments, one of A or B is an α -L-LNA nucleoside, another of A or B comprises a 2'-(ara)-F sugar moiety, and C comprises 2-thio-thymidine nucleobase.

In certain embodiments, one of A or B comprises a bicyclic sugar moiety, another of A or B comprises a 2'-MOE sugar moiety, and C comprises a 5-propyne uridine nucleobase. In certain embodiments, one of A or B is an LNA nucleoside, another of A or B comprises a 2'-MOE sugar moiety, and C comprises a 5-propyne uridine nucleobase. In certain embodiments, one of A or B is a cEt nucleoside, another of A or B comprises a 2'-MOE sugar moiety, and C comprises a 5-propyne uridine nucleobase. In certain embodiments, one of A or B is an α -L-LNA nucleoside, another of A or B comprises a 2'-MOE sugar moiety, and C comprises a 5-propyne uridine nucleobase.

In certain embodiments, one of A or B comprises a bicyclic sugar moiety, another of A or B comprises a 2'-MOE sugar moiety, and C comprises an unmodified 2'-deoxyfuranose sugar moiety. In certain embodiments, one of A or B is an LNA nucleoside, another of A or B comprises a 2'-MOE sugar moiety, and C comprises a 5-propyne uridine nucleobase. In certain embodiments, one of A or B is a cEt nucleoside, another of A or B comprises a 2'-MOE sugar moiety, and C comprises a 5-propyne uridine nucleobase. In certain embodiments, one of A or B is an α -L-LNA nucleoside, another of A or B comprises a 2'-MOE sugar moiety, and C comprises a 5-propyne uridine nucleobase.

In certain embodiments, one of A or B comprises a bicyclic sugar moiety, another of A or B comprises a 2'-F sugar moiety, and C comprises a 5-propyne uridine nucleobase. In certain embodiments, one of A or B is an LNA nucleoside, another of A or B comprises a 2'-F sugar moiety, and C comprises a 5-propyne uridine nucleobase. In certain embodiments, one of A or B is a cEt nucleoside, another of A or B comprises a 2'-F sugar moiety, and C comprises a 5-propyne uridine nucleobase. In certain embodiments, one of A or B is an α -L-LNA nucleoside, another of A or B comprises a 2'-F sugar moiety, and C comprises a 5-propyne uridine nucleobase.

In certain embodiments, one of A or B comprises a bicyclic sugar moiety, another of A or B comprises a 2'-(ara)-F sugar moiety, and C comprises a 5-propyne uridine nucleobase. In certain embodiments, one of A or B is an LNA nucleoside, another of A or B comprises a 2'-(ara)-F sugar moiety, and C comprises a 5-propyne uridine nucleobase. In certain embodiments, one of A or B is a cEt nucleoside, another of A or B comprises a 2'-(ara)-F sugar moiety, and C comprises a 5-propyne uridine nucleobase. In certain embodiments, one of A or B is an α -L-LNA nucleoside, another of A or B comprises a 2'-(ara)-F sugar moiety, and C comprises a 5-propyne uridine nucleobase.

In certain embodiments, one of A or B comprises a bicyclic sugar moiety, another of A or B comprises a 2'-MOE sugar moiety, and C comprises a sugar surrogate. In certain embodiments, one of A or B is an LNA nucleoside, another of A or B comprises a 2'-MOE sugar moiety, and C comprises a sugar surrogate. In certain embodiments, one of A or B is a cEt nucleoside, another of A or B comprises a 2'-MOE

sugar moiety, and C comprises a sugar surrogate. In certain embodiments, one of A or B is an α -L-LNA nucleoside, another of A or B comprises a 2'-MOE sugar moiety, and C comprises a sugar surrogate.

In certain embodiments, one of A or B comprises a bicyclic sugar moiety, another of A or B comprises a 2'-F sugar moiety, and C comprises a sugar surrogate. In certain embodiments, one of A or B is an LNA nucleoside, another of A or B comprises a 2'-F sugar moiety, and C comprises a sugar surrogate. In certain embodiments, one of A or B is a cEt nucleoside, another of A or B comprises a 2'-F sugar moiety, and C comprises a sugar surrogate. In certain embodiments, one of A or B is an α -L-LNA nucleoside, another of A or B comprises a 2'-F sugar moiety, and C comprises a sugar surrogate.

In certain embodiments, one of A or B comprises a bicyclic sugar moiety, another of A or B comprises a 2'-(ara)-F sugar moiety, and C comprises a sugar surrogate. In certain embodiments, one of A or B is an LNA nucleoside, another of A or B comprises a 2'-(ara)-F sugar moiety, and C comprises a sugar surrogate. In certain embodiments, one of A or B is a cEt nucleoside, another of A or B comprises a 2'-(ara)-F sugar moiety, and C comprises a sugar surrogate. In certain embodiments, one of A or B is an α -L-LNA nucleoside, another of A or B comprises a 2'-(ara)-F sugar moiety, and C comprises a sugar surrogate.

In certain embodiments, one of A or B comprises a bicyclic sugar moiety, another of A or B comprises a 2'-MOE sugar moiety, and C comprises a HNA sugar surrogate. In certain embodiments, one of A or B is an LNA nucleoside, another of A or B comprises a 2'-MOE sugar moiety, and C comprises a HNA sugar surrogate. In certain embodiments, one of A or B is a cEt nucleoside, another of A or B comprises a 2'-MOE sugar moiety, and C comprises a HNA sugar surrogate. In certain embodiments, one of A or B is an α -L-LNA nucleoside, another of A or B comprises a 2'-MOE sugar moiety, and C comprises a HNA sugar surrogate.

In certain embodiments, one of A or B comprises a bicyclic sugar moiety, another of A or B comprises a 2'-F sugar moiety, and C comprises a HNA sugar surrogate. In certain embodiments, one of A or B is an LNA nucleoside, another of A or B comprises a 2'-F sugar moiety, and C comprises a HNA sugar surrogate. In certain embodiments, one of A or B is a cEt nucleoside, another of A or B comprises a 2'-F sugar moiety, and C comprises a HNA sugar surrogate. In certain embodiments, one of A or B is an α -L-LNA nucleoside, another of A or B comprises a 2'-F sugar moiety, and C comprises a HNA sugar surrogate.

In certain embodiments, one of A or B comprises a bicyclic sugar moiety, another of A or B comprises a 2'-(ara)-F sugar moiety, and C comprises a HNA sugar surrogate. In certain embodiments, one of A or B is an LNA nucleoside, another of A or B comprises a 2'-(ara)-F sugar moiety, and C comprises a HNA sugar surrogate. In certain embodiments, one of A or B is a cEt nucleoside, another of A or B comprises a 2'-(ara)-F sugar moiety, and C comprises a HNA sugar surrogate. In certain embodiments, one of A or B is an α -L-LNA nucleoside, another of A or B comprises a 2'-(ara)-F sugar moiety, and C comprises a HNA sugar surrogate.

In certain embodiments, one of A or B comprises a bicyclic sugar moiety, another of A or B comprises a 2'-MOE sugar moiety, and C comprises a F-HNA sugar surrogate. In certain embodiments, one

of A or B is an LNA nucleoside, another of A or B comprises a 2'-MOE sugar moiety, and C comprises a F-HNA sugar surrogate. In certain embodiments, one of A or B is a cEt nucleoside, another of A or B comprises a 2'-MOE sugar moiety, and C comprises a F-HNA sugar surrogate. In certain embodiments, one of A or B is an α -L-LNA nucleoside, another of A or B comprises a 2'-MOE sugar moiety, and C comprises a F-HNA sugar surrogate.

In certain embodiments, one of A or B comprises a bicyclic sugar moiety, another of A or B comprises a 2'-F sugar moiety, and C comprises a F-HNA sugar surrogate. In certain embodiments, one of A or B is an LNA nucleoside, another of A or B comprises a 2'-F sugar moiety, and C comprises a F-HNA sugar surrogate. In certain embodiments, one of A or B is a cEt nucleoside, another of A or B comprises a 2'-F sugar moiety, and C comprises a F-HNA sugar surrogate. In certain embodiments, one of A or B is an α -L-LNA nucleoside, another of A or B comprises a 2'-F sugar moiety, and C comprises a F-HNA sugar surrogate.

In certain embodiments, one of A or B comprises a bicyclic sugar moiety, another of A or B comprises a 2'-(ara)-F sugar moiety, and C comprises a F-HNA sugar surrogate. In certain embodiments, one of A or B is an LNA nucleoside, another of A or B comprises a 2'-(ara)-F sugar moiety, and C comprises a F-HNA sugar surrogate. In certain embodiments, one of A or B is a cEt nucleoside, another of A or B comprises a 2'-(ara)-F sugar moiety, and C comprises a F-HNA sugar surrogate. In certain embodiments, one of A or B is an α -L-LNA nucleoside, another of A or B comprises a 2'-(ara)-F sugar moiety, and C comprises a F-HNA sugar surrogate.

In certain embodiments, one of A or B comprises a bicyclic sugar moiety, another of A or B comprises a 2'-MOE sugar moiety, and C comprises a 5'-Me DNA sugar moiety. In certain embodiments, one of A or B is an LNA nucleoside, another of A or B comprises a 2'-MOE sugar moiety, and C comprises a 5'-Me DNA sugar moiety. In certain embodiments, one of A or B is a cEt nucleoside, another of A or B comprises a 2'-MOE sugar moiety, and C comprises a 5'-Me DNA sugar moiety. In certain embodiments, one of A or B is an α -L-LNA nucleoside, another of A or B comprises a 2'-MOE sugar moiety, and C comprises a 5'-Me DNA sugar moiety.

In certain embodiments, one of A or B comprises a bicyclic sugar moiety, another of A or B comprises a 2'-F sugar moiety, and C comprises a 5'-Me DNA sugar moiety. In certain embodiments, one of A or B is an LNA nucleoside, another of A or B comprises a 2'-F sugar moiety, and C comprises a 5'-Me DNA sugar moiety. In certain embodiments, one of A or B is a cEt nucleoside, another of A or B comprises a 2'-F sugar moiety, and C comprises a 5'-Me DNA sugar moiety. In certain embodiments, one of A or B is an α -L-LNA nucleoside, another of A or B comprises a 2'-F sugar moiety, and C comprises a 5'-Me DNA sugar moiety.

In certain embodiments, one of A or B comprises a bicyclic sugar moiety, another of A or B comprises a 2'-(ara)-F sugar moiety, and C comprises a 5'-Me DNA sugar moiety. In certain embodiments, one of A or B is an LNA nucleoside, another of A or B comprises a 2'-(ara)-F sugar moiety, and C comprises

a 5'-Me DNA sugar moiety. In certain embodiments, one of A or B is a cEt nucleoside, another of A or B comprises a 2'-(ara)-F sugar moiety, and C comprises a 5'-Me DNA sugar moiety. In certain embodiments, one of A or B is an α -L-LNA nucleoside, another of A or B comprises a 2'-(ara)-F sugar moiety, and C comprises a 5'-Me DNA sugar moiety.

In certain embodiments, one of A or B comprises a bicyclic sugar moiety, another of A or B comprises a 2'-MOE sugar moiety, and C comprises a 5'-(R)-Me DNA sugar moiety. In certain embodiments, one of A or B is an LNA nucleoside, another of A or B comprises a 2'-MOE sugar moiety, and C comprises a 5'-(R)-Me DNA sugar moiety. In certain embodiments, one of A or B is a cEt nucleoside, another of A or B comprises a 2'-MOE sugar moiety, and C comprises a 5'-(R)-Me DNA sugar moiety. In certain embodiments, one of A or B is an α -L-LNA nucleoside, another of A or B comprises a 2'-MOE sugar moiety, and C comprises a 5'-(R)-Me DNA sugar moiety.

In certain embodiments, one of A or B comprises a bicyclic sugar moiety, another of A or B comprises a 2'-F sugar moiety, and C comprises a 5'-(R)-Me DNA sugar moiety. In certain embodiments, one of A or B is an LNA nucleoside, another of A or B comprises a 2'-F sugar moiety, and C comprises a 5'-(R)-Me DNA sugar moiety. In certain embodiments, one of A or B is a cEt nucleoside, another of A or B comprises a 2'-F sugar moiety, and C comprises a 5'-(R)-Me DNA sugar moiety. In certain embodiments, one of A or B is an α -L-LNA nucleoside, another of A or B comprises a 2'-F sugar moiety, and C comprises a 5'-(R)-Me DNA sugar moiety.

In certain embodiments, one of A or B comprises a bicyclic sugar moiety, another of A or B comprises a 2'-(ara)-F sugar moiety, and C comprises a 5'-(R)-Me DNA sugar moiety. In certain embodiments, one of A or B is an LNA nucleoside, another of A or B comprises a 2'-(ara)-F sugar moiety, and C comprises a 5'-(R)-Me DNA sugar moiety. In certain embodiments, one of A or B is a cEt nucleoside, another of A or B comprises a 2'-(ara)-F sugar moiety, and C comprises a 5'-(R)-Me DNA sugar moiety. In certain embodiments, one of A or B is an α -L-LNA nucleoside, another of A or B comprises a 2'-(ara)-F sugar moiety, and C comprises a 5'-(R)-Me DNA sugar moiety.

In certain embodiments, at least two of A, B or C comprises a 2'-substituted sugar moiety, and the other comprises a bicyclic sugar moiety. In certain embodiments, at least two of A, B or C comprises a bicyclic sugar moiety, and the other comprises a 2'-substituted sugar moiety.

In certain embodiments, at least two of A, B or C comprises a 2'-substituted sugar moiety, and the other comprises an unmodified 2'-deoxyfuranose sugar moiety.. In certain embodiments, at least two of A, B or C comprises a bicyclic sugar moiety, and the other comprises an unmodified 2'-deoxyfuranose sugar moiety.

Certain gaps

In certain embodiments, the gap of a gapmer consists of 6 to 20 linked nucleosides. In certain embodiments, the gap of a gapmer consists of 6 to 15 linked nucleosides. In certain embodiments, the gap of a gapmer consists of 6 to 12 linked nucleosides. In certain embodiments, the gap of a gapmer consists of 6 to

10 linked nucleosides. In certain embodiments, the gap of a gapmer consists of 6 to 9 linked nucleosides. In certain embodiments, the gap of a gapmer consists of 6 to 8 linked nucleosides. In certain embodiments, the gap of a gapmer consists of 6 or 7 linked nucleosides. In certain embodiments, the gap of a gapmer consists of 7 to 10 linked nucleosides. In certain embodiments, the gap of a gapmer consists of 7 to 9 linked nucleosides. In certain embodiments, the gap of a gapmer consists of 7 or 8 linked nucleosides. In certain embodiments, the gap of a gapmer consists of 8 to 10 linked nucleosides. In certain embodiments, the gap of a gapmer consists of 8 or 9 linked nucleosides. In certain embodiments, the gap of a gapmer consists of 6 linked nucleosides. In certain embodiments, the gap of a gapmer consists of 7 linked nucleosides. In certain embodiments, the gap of a gapmer consists of 8 linked nucleosides. In certain embodiments, the gap of a gapmer consists of 9 linked nucleosides. In certain embodiments, the gap of a gapmer consists of 10 linked nucleosides. In certain embodiments, the gap of a gapmer consists of 11 linked nucleosides. In certain embodiments, the gap of a gapmer consists of 12 linked nucleosides.

In certain embodiments, each nucleotide of the gap of a gapmer is a 2'-deoxynucleoside. In certain embodiments, the gap comprises one or more modified nucleosides. In certain embodiments, each nucleotide of the gap of a gapmer is a 2'-deoxynucleoside or is a modified nucleoside that is "DNA-like." In such embodiments, "DNA-like" means that the nucleoside has similar characteristics to DNA, such that a duplex comprising the gapmer and an RNA molecule is capable of activating RNase H. For example, under certain conditions, 2'-fluoro (arabino) nucleosides (also referred to as FANA) have been shown to support RNase H activation, and thus is DNA-like. In certain embodiments, one or more nucleosides of the gap of a gapmer is not a 2'-deoxynucleoside and is not DNA-like. In certain such embodiments, the gapmer nonetheless supports RNase H activation (e.g., by virtue of the number or placement of the non-DNA nucleosides).

Certain gapmer motifs

In certain embodiments, a gapmer comprises a 5'-wing, a gap, and a 3' wing, wherein the 5'-wing, gap, and 3' wing are independently selected from among those discussed above. For example, in certain embodiments, a gapmer has a 5'-wing selected from any of the 5'-wing motifs in Tables 1, 2, and 3 above and a 3'-wing selected from any of the 3'-wing motifs in Tables, 4, 5, and 6. For example, in certain embodiments, a gapmer has a 5'-wing, a gap, and a 3'-wing having features selected from among those listed in the following non-limiting table:

Table 7
Certain Gapmer Sugar Motifs

Gapmer motif #	5-wing	Gap	3'-wing
1	At least one non-bicyclic modified nucleoside	All 2'-deoxynucleosides	At least one bicyclic nucleoside
2	At least one non-bicyclic modified nucleoside	All 2'-deoxynucleosides	At least one LNA nucleoside

3	At least one non-bicyclic modified nucleoside	All 2'-deoxynucleosides	At least one cEt nucleoside
4	At least one 2'-substituted nucleoside	All 2'-deoxynucleosides	At least one bicyclic nucleoside
5	At least one 2'-substituted nucleoside	All 2'-deoxynucleosides	At least one LNA nucleoside
6	At least one 2'-substituted nucleoside	All 2'-deoxynucleosides	At least one cEt nucleoside
7	At least one 2'-MOE nucleoside	All 2'-deoxynucleosides	At least one bicyclic nucleoside
8	At least one 2'-MOE nucleoside	All 2'-deoxynucleosides	At least one LNA nucleoside
9	At least one 2'-MOE nucleoside	All 2'-deoxynucleosides	At least one cEt nucleoside
10	At least one 2'-OMe nucleoside	All 2'-deoxynucleosides	At least one bicyclic nucleoside
11	At least one 2'-OMe nucleoside	All 2'-deoxynucleosides	At least one LNA nucleoside
12	At least one 2'-OMe nucleoside	All 2'-deoxynucleosides	At least one cEt nucleoside
13	At least one 2'-deoxynucleoside	All 2'-deoxynucleosides	At least one bicyclic nucleoside
14	At least one 2'-deoxynucleoside	All 2'-deoxynucleosides	At least one LNA nucleoside
15	At least one 2'-deoxynucleoside	All 2'-deoxynucleosides	At least one cEt nucleoside
16	At least one bicyclic nucleoside	All 2'-deoxynucleosides	At least one non-bicyclic modified nucleoside
17	At least one LNA nucleoside	All 2'-deoxynucleosides	At least one non-bicyclic modified nucleoside
18	At least one cEt nucleoside	All 2'-deoxynucleosides	At least one non-bicyclic modified nucleoside
19	At least one bicyclic nucleoside	All 2'-deoxynucleosides	At least one 2'-substituted nucleoside
20	At least one LNA nucleoside	All 2'-deoxynucleosides	At least one 2'-substituted nucleoside
21	At least one cEt nucleoside	All 2'-deoxynucleosides	At least one 2'-substituted nucleoside
22	At least one bicyclic nucleoside	All 2'-deoxynucleosides	At least one 2'-MOE nucleoside
23	At least one LNA nucleoside	All 2'-deoxynucleosides	At least one 2'-MOE nucleoside
24	At least one cEt nucleoside	All 2'-deoxynucleosides	At least one 2'-MOE nucleoside
25	At least one bicyclic nucleoside	All 2'-deoxynucleosides	At least one 2'-OMe nucleoside
26	At least one LNA nucleoside	All 2'-deoxynucleosides	At least one 2'-OMe nucleoside
27	At least one cEt nucleoside	All 2'-deoxynucleosides	At least one 2'-OMe nucleoside

28	At least one bicyclic nucleoside	All 2'-deoxynucleosides	At least one 2'-deoxynucleoside
29	At least one LNA nucleoside	All 2'-deoxynucleosides	At least one 2'-deoxynucleoside
30	At least one cEt nucleoside	All 2'-deoxynucleosides	At least one 2'-deoxynucleoside
31	At least one bicyclic nucleoside and at least one 2'-substituted nucleoside	All 2'-deoxynucleosides	At least one bicyclic nucleoside and at least one 2'-substituted nucleoside
32	At least one bicyclic nucleoside and at least one 2'-substituted nucleoside	All 2'-deoxynucleosides	At least two bicyclic nucleosides
33	At least one cEt nucleoside and at least one 2'-substituted nucleoside	All 2'-deoxynucleosides	At least one bicyclic nucleoside and at least one 2'-substituted nucleoside
34	At least one cEt nucleoside and at least one 2'-substituted nucleoside	All 2'-deoxynucleosides	At least two bicyclic nucleosides
35	At least one LNA nucleoside and at least one 2'-substituted nucleoside	All 2'-deoxynucleosides	At least one bicyclic nucleoside and at least one 2'-substituted nucleoside
36	At least one LNA nucleoside and at least one 2'-substituted nucleoside	All 2'-deoxynucleosides	At least two bicyclic nucleosides
37	At least one bicyclic nucleoside and at least one 2'-substituted nucleoside	All 2'-deoxynucleosides	At least one LNA nucleoside and at least one 2'-substituted nucleoside
38	At least one bicyclic nucleoside and at least one 2'-substituted nucleoside	All 2'-deoxynucleosides	At least two LNA nucleosides
39	At least one cEt nucleoside and at least one 2'-substituted nucleoside	All 2'-deoxynucleosides	At least one LNA nucleoside and at least one 2'-substituted nucleoside
40	At least one cEt nucleoside and at least one 2'-substituted nucleoside	All 2'-deoxynucleosides	At least two LNA nucleosides
41	At least one LNA nucleoside and at least one 2'-substituted nucleoside	All 2'-deoxynucleosides	At least one LNA nucleoside and at least one 2'-substituted nucleoside
42	At least one LNA nucleoside and at least one 2'-substituted nucleoside	All 2'-deoxynucleosides	At least two LNA nucleosides
43	At least one bicyclic nucleoside and at least one 2'-deoxynucleoside	All 2'-deoxynucleosides	At least one bicyclic nucleoside and at least one 2'-substituted nucleoside

44	At least one bicyclic nucleoside and at least one 2'-deoxynucleoside	All 2'-deoxynucleosides	At least two bicyclic nucleosides
45	At least one cEt nucleoside and at least one 2'-deoxynucleoside	All 2'-deoxynucleosides	At least one bicyclic nucleoside and at least one 2'-substituted nucleoside
46	At least one cEt nucleoside and at least one 2'-deoxynucleoside	All 2'-deoxynucleosides	At least two bicyclic nucleosides
47	At least one LNA nucleoside and at least one 2'-deoxynucleoside	All 2'-deoxynucleosides	At least one bicyclic nucleoside and at least one 2'-substituted nucleoside
48	At least one LNA nucleoside and at least one 2'-deoxynucleoside	All 2'-deoxynucleosides	At least two bicyclic nucleosides
49	At least one bicyclic nucleoside and at least one 2'-deoxynucleoside	All 2'-deoxynucleosides	At least one LNA nucleoside and at least one 2'-substituted nucleoside
50	At least one bicyclic nucleoside and at least one 2'-deoxynucleoside	All 2'-deoxynucleosides	At least two LNA nucleosides
51	At least one cEt nucleoside and at least one 2'-deoxynucleoside	All 2'-deoxynucleosides	At least one LNA nucleoside and at least one 2'-substituted nucleoside
52	At least one cEt nucleoside and at least one 2'-deoxynucleoside	All 2'-deoxynucleosides	At least two LNA nucleosides
53	At least one LNA nucleoside and at least one 2'-deoxynucleoside	All 2'-deoxynucleosides	At least one LNA nucleoside and at least one 2'-substituted nucleoside
54	At least one LNA nucleoside and at least one 2'-deoxynucleoside	All 2'-deoxynucleosides	At least two LNA nucleosides
55	At least two 2'-substituted nucleosides	All 2'-deoxynucleosides	At least one bicyclic nucleoside and at least one 2'-substituted nucleoside
56	At least two 2'-substituted nucleosides	All 2'-deoxynucleosides	At least two bicyclic nucleosides
57	At least two 2'-substituted nucleosides	All 2'-deoxynucleosides	At least one LNA nucleoside and at least one 2'-substituted nucleoside
58	At least two 2'-substituted nucleosides	All 2'-deoxynucleosides	At least two LNA nucleosides

In certain embodiments, a gapmer comprises a 5'-wing, a gap, and a 3' wing, wherein the 5'-wing, gap, and 3' wing are independently selected from among those discussed above. For example, in certain embodiments, a gapmer has a 5'-wing, a gap, and a 3'-wing wherein the 5'-wing and the 3'-wing have features selected from among those listed in the tables above. In certain embodiments, any 5'-wing may be

paired with any 3'-wing. In certain embodiments the 5'-wing may comprise ABBBB and the 3'-wing may comprise BBA. In certain embodiments the 5'-wing may comprise ACACA and the 3'-wing may comprise BB. For example, in certain embodiments, a gapmer has a 5'-wing, a gap, and a 3'-wing having features selected from among those listed in the following non-limiting table, wherein each motif is represented as (5'-wing)-(gap)-(3'-wing), wherein each number represents the number of linked nucleosides in each portion of the motif, for example, a 5-10-5 motif would have a 5'-wing comprising 5 nucleosides, a gap comprising 10 nucleosides, and a 3'-wing comprising 5 nucleosides:

Table 8
Certain Gapmer Sugar Motifs

Certain Gapmer Sugar Motifs			
2-10-2	3-10-2	4-10-2	5-10-2
2-10-3	3-10-3	4-10-3	5-10-3
2-10-4	3-10-4	4-10-4	5-10-4
2-10-5	3-10-5	4-10-5	5-10-5
2-9-2	3-9-2	4-9-2	5-9-2
2-9-3	3-9-3	4-9-3	5-9-3
2-9-4	3-9-4	4-9-4	5-9-4
2-9-5	3-9-5	4-9-5	5-9-5
2-11-2	3-11-2	4-11-2	5-11-2
2-11-3	3-11-3	4-11-3	5-11-3
2-11-4	3-11-4	4-11-4	5-11-4
2-11-5	3-11-5	4-11-5	5-11-5
2-8-2	3-8-2	4-8-2	5-8-2
2-8-3	3-8-3	4-8-3	5-8-3
2-8-4	3-8-4	4-8-4	5-8-4
2-8-5	3-8-5	4-8-5	5-8-5

In certain embodiments, gapmers have a motif described by Formula I as follows:



wherein:

each A is independently a 2'-substituted nucleoside;

each B is independently a bicyclic nucleoside;

each J is independently either a 2'-substituted nucleoside or a 2'-deoxynucleoside;

each D is a 2'-deoxynucleoside;

m is 0-4; n is 0-2; p is 0-2; r is 0-2; t is 0-2; v is 0-2; w is 0-4; x is 0-2; y is 0-2; z is 0-4; g is 6; and h is 14;

provided that:

at least one of m, n, and r is other than 0;

at least one of w and y is other than 0;


the sum of m, n, p, r, and t is from 2 to 5; and

the sum of v, w, x, y, and z is from 2 to 5.

In certain embodiments, one or more 2'-substituted nucleoside is a 2'-MOE nucleoside. In certain embodiments, one or more 2'-substituted nucleoside is a 2'-OMe nucleoside. In certain In certain embodiments, one or more bicyclic nucleoside is a cEt nucleoside. In certain embodiments, one or more bicyclic nucleoside is an LNA nucleoside.

In certain embodiments, a gapmer of Formula I has a motif selected from among gapmer motifs 1-58.

In certain embodiments, gapmers have a motif described by Formula II as follows:



wherein:

each A is independently a 2'-substituted nucleoside;

each B is independently a bicyclic nucleoside;

each J is independently either a 2'-substituted nucleoside or a 2'-deoxynucleoside;

each D is a 2'-deoxynucleoside;

m is 0-4; n is 0-2; p is 0-2; r is 0-2; t is 0-2; v is 0-2; w is 0-4; x is 0-2; y is 0-2; z is 0-4; g is 6-14;

provided that:

at least one of m, n, and r is other than 0;

at least one of w and y is other than 0;

the sum of m, n, p, r, and t is from 1 to 5; and

the sum of v, w, x, y, and z is from 1 to 5.

In certain embodiments, one or more 2'-substituted nucleoside is a 2'-MOE nucleoside. In certain embodiments, one or more 2'-substituted nucleoside is a 2'-OMe nucleoside. In certain embodiments, one or more bicyclic nucleoside is a cEt nucleoside. In certain embodiments, one or more bicyclic nucleoside is an LNA nucleoside.

In certain embodiments, each 2'-substituted nucleoside is a 2'-MOE nucleoside. In certain embodiments, each 2'-substituted nucleoside is a 2'-OMe nucleoside. In certain embodiments, each bicyclic nucleoside is a cEt nucleoside. In certain embodiments, each bicyclic nucleoside is an LNA nucleoside.

In certain embodiments, each A is the same 2'-substituted nucleoside. In certain embodiments, each B is the same bicyclic nucleoside. In certain embodiments each A is the same 2'-modified nucleoside and each B is the same bicyclic nucleoside. In certain embodiments, each J is a 2'-modified nucleoside. In certain embodiments each J is the same 2'-modified nucleoside. In certain embodiments, each J and each A is the same 2'-modified nucleoside.

In certain embodiments, a gapmer of Formula II has a motif selected from among gapmer motifs 1-58.

In certain embodiments, a gapmer comprises a 5'-wing, a gap, and a 3' wing, independently selected from among those proved in the above tables, for example as provided in the following table:

Table 9
Certain Gapmer Sugar Motifs

Gapmer motif #	5-wing sugar motif (from table 1)	Gap	3'-wing sugar motif (from table 2)
59	1(a-i)	All 2'-deoxynucleosides	1(a-i)
60	2(a-i)	All 2'-deoxynucleosides	1(a-i)
61	3(a-i)	All 2'-deoxynucleosides	1(a-i)
62	4(a-i)	All 2'-deoxynucleosides	1(a-i)
63	5(a-i)	All 2'-deoxynucleosides	1(a-i)
64	6(a-i)	All 2'-deoxynucleosides	1(a-i)
65	7(a-i)	All 2'-deoxynucleosides	1(a-i)
66	8(a-i)	All 2'-deoxynucleosides	1(a-i)
67	9(a-i)	All 2'-deoxynucleosides	1(a-i)
68	10(a-i)	All 2'-deoxynucleosides	1(a-i)
69	11(a-i)	All 2'-deoxynucleosides	1(a-i)
70	12(a-i)	All 2'-deoxynucleosides	1(a-i)
71	13(a-i)	All 2'-deoxynucleosides	1(a-i)
72	14(a-i)	All 2'-deoxynucleosides	1(a-i)
73	15(a-i)	All 2'-deoxynucleosides	1(a-i)
74	16(a-i)	All 2'-deoxynucleosides	1(a-i)
75	17(a-i)	All 2'-deoxynucleosides	1(a-i)
76	18(a-i)	All 2'-deoxynucleosides	1(a-i)
77	19(a-i)	All 2'-deoxynucleosides	1(a-i)
78	20(a-i)	All 2'-deoxynucleosides	1(a-i)
79	21(a-i)	All 2'-deoxynucleosides	1(a-i)
80	22(a-i)	All 2'-deoxynucleosides	1(a-i)
81	1(a-i)	All 2'-deoxynucleosides	2(a-i)
82	2(a-i)	All 2'-deoxynucleosides	2(a-i)
83	3(a-i)	All 2'-deoxynucleosides	2(a-i)
84	4(a-i)	All 2'-deoxynucleosides	2(a-i)
85	5(a-i)	All 2'-deoxynucleosides	2(a-i)
86	6(a-i)	All 2'-deoxynucleosides	2(a-i)
87	7(a-i)	All 2'-deoxynucleosides	2(a-i)
88	8(a-i)	All 2'-deoxynucleosides	2(a-i)
89	9(a-i)	All 2'-deoxynucleosides	2(a-i)
90	10(a-i)	All 2'-deoxynucleosides	2(a-i)
91	11(a-i)	All 2'-deoxynucleosides	2(a-i)
92	12(a-i)	All 2'-deoxynucleosides	2(a-i)
93	13(a-i)	All 2'-deoxynucleosides	2(a-i)
94	14(a-i)	All 2'-deoxynucleosides	2(a-i)
94	15(a-i)	All 2'-deoxynucleosides	2(a-i)
96	16(a-i)	All 2'-deoxynucleosides	2(a-i)
97	17(a-i)	All 2'-deoxynucleosides	2(a-i)

98	18(a-i)	All 2'-deoxynucleosides	2(a-i)
99	19(a-i)	All 2'-deoxynucleosides	2(a-i)
100	20(a-i)	All 2'-deoxynucleosides	2(a-i)
101	21(a-i)	All 2'-deoxynucleosides	2(a-i)
102	22(a-i)	All 2'-deoxynucleosides	2(a-i)
103	1(a-i)	All 2'-deoxynucleosides	3(a-i)
104	2(a-i)	All 2'-deoxynucleosides	3(a-i)
105	3(a-i)	All 2'-deoxynucleosides	3(a-i)
106	4(a-i)	All 2'-deoxynucleosides	3(a-i)
107	5(a-i)	All 2'-deoxynucleosides	3(a-i)
108	6(a-i)	All 2'-deoxynucleosides	3(a-i)
109	7(a-i)	All 2'-deoxynucleosides	3(a-i)
110	8(a-i)	All 2'-deoxynucleosides	3(a-i)
111	9(a-i)	All 2'-deoxynucleosides	3(a-i)
112	10(a-i)	All 2'-deoxynucleosides	3(a-i)
113	11(a-i)	All 2'-deoxynucleosides	3(a-i)
114	12(a-i)	All 2'-deoxynucleosides	3(a-i)
115	13(a-i)	All 2'-deoxynucleosides	3(a-i)
116	14(a-i)	All 2'-deoxynucleosides	3(a-i)
117	15(a-i)	All 2'-deoxynucleosides	3(a-i)
118	16(a-i)	All 2'-deoxynucleosides	3(a-i)
119	17(a-i)	All 2'-deoxynucleosides	3(a-i)
120	18(a-i)	All 2'-deoxynucleosides	3(a-i)
121	19(a-i)	All 2'-deoxynucleosides	3(a-i)
122	20(a-i)	All 2'-deoxynucleosides	3(a-i)
123	21(a-i)	All 2'-deoxynucleosides	3(a-i)
124	22(a-i)	All 2'-deoxynucleosides	3(a-i)
125	1(a-i)	All 2'-deoxynucleosides	4(a-i)
126	2(a-i)	All 2'-deoxynucleosides	4(a-i)
127	3(a-i)	All 2'-deoxynucleosides	4(a-i)
128	4(a-i)	All 2'-deoxynucleosides	4(a-i)
129	5(a-i)	All 2'-deoxynucleosides	4(a-i)
130	6(a-i)	All 2'-deoxynucleosides	4(a-i)
131	7(a-i)	All 2'-deoxynucleosides	4(a-i)
132	8(a-i)	All 2'-deoxynucleosides	4(a-i)
133	9(a-i)	All 2'-deoxynucleosides	4(a-i)
134	10(a-i)	All 2'-deoxynucleosides	4(a-i)
135	11(a-i)	All 2'-deoxynucleosides	4(a-i)
136	12(a-i)	All 2'-deoxynucleosides	4(a-i)
137	13(a-i)	All 2'-deoxynucleosides	4(a-i)
138	14(a-i)	All 2'-deoxynucleosides	4(a-i)
139	15(a-i)	All 2'-deoxynucleosides	4(a-i)
140	16(a-i)	All 2'-deoxynucleosides	4(a-i)
141	17(a-i)	All 2'-deoxynucleosides	4(a-i)
142	18(a-i)	All 2'-deoxynucleosides	4(a-i)

143	19(a-i)	All 2'-deoxynucleosides	4(a-i)
144	20(a-i)	All 2'-deoxynucleosides	4(a-i)
145	21(a-i)	All 2'-deoxynucleosides	4(a-i)
146	22(a-i)	All 2'-deoxynucleosides	4(a-i)
147	1(a-i)	All 2'-deoxynucleosides	5(a-i)
148	2(a-i)	All 2'-deoxynucleosides	5(a-i)
149	3(a-i)	All 2'-deoxynucleosides	5(a-i)
150	4(a-i)	All 2'-deoxynucleosides	5(a-i)
151	5(a-i)	All 2'-deoxynucleosides	5(a-i)
152	6(a-i)	All 2'-deoxynucleosides	5(a-i)
153	7(a-i)	All 2'-deoxynucleosides	5(a-i)
154	8(a-i)	All 2'-deoxynucleosides	5(a-i)
155	9(a-i)	All 2'-deoxynucleosides	5(a-i)
156	10(a-i)	All 2'-deoxynucleosides	5(a-i)
157	11(a-i)	All 2'-deoxynucleosides	5(a-i)
158	12(a-i)	All 2'-deoxynucleosides	5(a-i)
159	13(a-i)	All 2'-deoxynucleosides	5(a-i)
160	14(a-i)	All 2'-deoxynucleosides	5(a-i)
161	15(a-i)	All 2'-deoxynucleosides	5(a-i)
162	16(a-i)	All 2'-deoxynucleosides	5(a-i)
163	17(a-i)	All 2'-deoxynucleosides	5(a-i)
164	18(a-i)	All 2'-deoxynucleosides	5(a-i)
165	19(a-i)	All 2'-deoxynucleosides	5(a-i)
166	20(a-i)	All 2'-deoxynucleosides	5(a-i)
167	21(a-i)	All 2'-deoxynucleosides	5(a-i)
168	22(a-i)	All 2'-deoxynucleosides	5(a-i)
169	1(a-i)	All 2'-deoxynucleosides	6(a-i)
170	2(a-i)	All 2'-deoxynucleosides	6(a-i)
171	3(a-i)	All 2'-deoxynucleosides	6(a-i)
172	4(a-i)	All 2'-deoxynucleosides	6(a-i)
173	5(a-i)	All 2'-deoxynucleosides	6(a-i)
174	6(a-i)	All 2'-deoxynucleosides	6(a-i)
175	7(a-i)	All 2'-deoxynucleosides	6(a-i)
176	8(a-i)	All 2'-deoxynucleosides	6(a-i)
177	9(a-i)	All 2'-deoxynucleosides	6(a-i)
178	10(a-i)	All 2'-deoxynucleosides	6(a-i)
179	11(a-i)	All 2'-deoxynucleosides	6(a-i)
180	12(a-i)	All 2'-deoxynucleosides	6(a-i)
181	13(a-i)	All 2'-deoxynucleosides	6(a-i)
182	14(a-i)	All 2'-deoxynucleosides	6(a-i)
183	15(a-i)	All 2'-deoxynucleosides	6(a-i)
184	16(a-i)	All 2'-deoxynucleosides	6(a-i)
184	17(a-i)	All 2'-deoxynucleosides	6(a-i)
186	18(a-i)	All 2'-deoxynucleosides	6(a-i)
187	19(a-i)	All 2'-deoxynucleosides	6(a-i)

188	20(a-i)	All 2'-deoxynucleosides	6(a-i)
189	21(a-i)	All 2'-deoxynucleosides	6(a-i)
190	22(a-i)	All 2'-deoxynucleosides	6(a-i)
191	1(a-i)	All 2'-deoxynucleosides	7(a-i)
192	2(a-i)	All 2'-deoxynucleosides	7(a-i)
193	3(a-i)	All 2'-deoxynucleosides	7(a-i)
194	4(a-i)	All 2'-deoxynucleosides	7(a-i)
195	5(a-i)	All 2'-deoxynucleosides	7(a-i)
196	6(a-i)	All 2'-deoxynucleosides	7(a-i)
197	7(a-i)	All 2'-deoxynucleosides	7(a-i)
198	8(a-i)	All 2'-deoxynucleosides	7(a-i)
199	9(a-i)	All 2'-deoxynucleosides	7(a-i)
200	10(a-i)	All 2'-deoxynucleosides	7(a-i)
201	11(a-i)	All 2'-deoxynucleosides	7(a-i)
202	12(a-i)	All 2'-deoxynucleosides	7(a-i)
203	13(a-i)	All 2'-deoxynucleosides	7(a-i)
204	14(a-i)	All 2'-deoxynucleosides	7(a-i)
205	15(a-i)	All 2'-deoxynucleosides	7(a-i)
206	16(a-i)	All 2'-deoxynucleosides	7(a-i)
207	17(a-i)	All 2'-deoxynucleosides	7(a-i)
208	18(a-i)	All 2'-deoxynucleosides	7(a-i)
209	19(a-i)	All 2'-deoxynucleosides	7(a-i)
210	20(a-i)	All 2'-deoxynucleosides	7(a-i)
211	21(a-i)	All 2'-deoxynucleosides	7(a-i)
212	22(a-i)	All 2'-deoxynucleosides	7(a-i)
213	1(a-i)	All 2'-deoxynucleosides	8(a-i)
214	2(a-i)	All 2'-deoxynucleosides	8(a-i)
215	3(a-i)	All 2'-deoxynucleosides	8(a-i)
216	4(a-i)	All 2'-deoxynucleosides	8(a-i)
217	5(a-i)	All 2'-deoxynucleosides	8(a-i)
218	6(a-i)	All 2'-deoxynucleosides	8(a-i)
219	7(a-i)	All 2'-deoxynucleosides	8(a-i)
220	8(a-i)	All 2'-deoxynucleosides	8(a-i)
221	9(a-i)	All 2'-deoxynucleosides	8(a-i)
222	10(a-i)	All 2'-deoxynucleosides	8(a-i)
223	11(a-i)	All 2'-deoxynucleosides	8(a-i)
224	12(a-i)	All 2'-deoxynucleosides	8(a-i)
225	13(a-i)	All 2'-deoxynucleosides	8(a-i)
226	14(a-i)	All 2'-deoxynucleosides	8(a-i)
227	15(a-i)	All 2'-deoxynucleosides	8(a-i)
228	16(a-i)	All 2'-deoxynucleosides	8(a-i)
229	17(a-i)	All 2'-deoxynucleosides	8(a-i)
230	18(a-i)	All 2'-deoxynucleosides	8(a-i)
231	19(a-i)	All 2'-deoxynucleosides	8(a-i)
232	20(a-i)	All 2'-deoxynucleosides	8(a-i)

233	21(a-i)	All 2'-deoxynucleosides	8(a-i)
234	22(a-i)	All 2'-deoxynucleosides	8(a-i)
235	1(a-i)	All 2'-deoxynucleosides	9(a-i)
236	2(a-i)	All 2'-deoxynucleosides	9(a-i)
237	3(a-i)	All 2'-deoxynucleosides	9(a-i)
238	4(a-i)	All 2'-deoxynucleosides	9(a-i)
239	5(a-i)	All 2'-deoxynucleosides	9(a-i)
240	6(a-i)	All 2'-deoxynucleosides	9(a-i)
241	7(a-i)	All 2'-deoxynucleosides	9(a-i)
242	8(a-i)	All 2'-deoxynucleosides	9(a-i)
243	9(a-i)	All 2'-deoxynucleosides	9(a-i)
244	10(a-i)	All 2'-deoxynucleosides	9(a-i)
245	11(a-i)	All 2'-deoxynucleosides	9(a-i)
246	12(a-i)	All 2'-deoxynucleosides	9(a-i)
247	13(a-i)	All 2'-deoxynucleosides	9(a-i)
248	14(a-i)	All 2'-deoxynucleosides	9(a-i)
249	15(a-i)	All 2'-deoxynucleosides	9(a-i)
250	16(a-i)	All 2'-deoxynucleosides	9(a-i)
251	17(a-i)	All 2'-deoxynucleosides	9(a-i)
252	18(a-i)	All 2'-deoxynucleosides	9(a-i)
253	19(a-i)	All 2'-deoxynucleosides	9(a-i)
254	20(a-i)	All 2'-deoxynucleosides	9(a-i)
255	21(a-i)	All 2'-deoxynucleosides	9(a-i)
256	22(a-i)	All 2'-deoxynucleosides	9(a-i)
257	1(a-i)	All 2'-deoxynucleosides	10(a-i)
258	2(a-i)	All 2'-deoxynucleosides	10(a-i)
259	3(a-i)	All 2'-deoxynucleosides	10(a-i)
260	4(a-i)	All 2'-deoxynucleosides	10(a-i)
261	5(a-i)	All 2'-deoxynucleosides	10(a-i)
262	6(a-i)	All 2'-deoxynucleosides	10(a-i)
263	7(a-i)	All 2'-deoxynucleosides	10(a-i)
264	8(a-i)	All 2'-deoxynucleosides	10(a-i)
265	9(a-i)	All 2'-deoxynucleosides	10(a-i)
266	10(a-i)	All 2'-deoxynucleosides	10(a-i)
267	11(a-i)	All 2'-deoxynucleosides	10(a-i)
268	12(a-i)	All 2'-deoxynucleosides	10(a-i)
269	13(a-i)	All 2'-deoxynucleosides	10(a-i)
270	14(a-i)	All 2'-deoxynucleosides	10(a-i)
271	15(a-i)	All 2'-deoxynucleosides	10(a-i)
272	16(a-i)	All 2'-deoxynucleosides	10(a-i)
273	17(a-i)	All 2'-deoxynucleosides	10(a-i)
274	18(a-i)	All 2'-deoxynucleosides	10(a-i)
275	19(a-i)	All 2'-deoxynucleosides	10(a-i)
276	20(a-i)	All 2'-deoxynucleosides	10(a-i)
277	21(a-i)	All 2'-deoxynucleosides	10(a-i)

278	22(a-i)	All 2'-deoxynucleosides	10(a-i)
279	1(a)-22(a)	All 2'-deoxynucleosides	1(a)-10(a)
280	1(b)-22(b)	All 2'-deoxynucleosides	1(a)-10(a)
281	1(c)-22(c)	All 2'-deoxynucleosides	1(a)-10(a)
282	1(d)-22(d)	All 2'-deoxynucleosides	1(a)-10(a)
283	1(e)-22(e)	All 2'-deoxynucleosides	1(a)-10(a)
284	1(f)-22(f)	All 2'-deoxynucleosides	1(a)-10(a)
285	1(g)-22(g)	All 2'-deoxynucleosides	1(a)-10(a)
286	1(h)-22(h)	All 2'-deoxynucleosides	1(a)-10(a)
287	1(i)-22(i)	All 2'-deoxynucleosides	1(a)-10(a)
288	1(a)-22(a)	All 2'-deoxynucleosides	1(b)-10(b)
289	1(b)-22(b)	All 2'-deoxynucleosides	1(b)-10(b)
290	1(c)-22(c)	All 2'-deoxynucleosides	1(b)-10(b)
291	1(d)-22(d)	All 2'-deoxynucleosides	1(b)-10(b)
292	1(e)-22(e)	All 2'-deoxynucleosides	1(b)-10(b)
293	1(f)-22(f)	All 2'-deoxynucleosides	1(b)-10(b)
294	1(g)-22(g)	All 2'-deoxynucleosides	1(b)-10(b)
295	1(h)-22(h)	All 2'-deoxynucleosides	1(b)-10(b)
296	1(i)-22(i)	All 2'-deoxynucleosides	1(b)-10(b)
297	1(a)-22(a)	All 2'-deoxynucleosides	1(c)-10(c)
298	1(b)-22(b)	All 2'-deoxynucleosides	1(c)-10(c)
299	1(c)-22(c)	All 2'-deoxynucleosides	1(c)-10(c)
300	1(d)-22(d)	All 2'-deoxynucleosides	1(c)-10(c)
301	1(e)-22(e)	All 2'-deoxynucleosides	1(c)-10(c)
302	1(f)-22(f)	All 2'-deoxynucleosides	1(c)-10(c)
303	1(g)-22(g)	All 2'-deoxynucleosides	1(c)-10(c)
304	1(h)-22(h)	All 2'-deoxynucleosides	1(c)-10(c)
305	1(i)-22(i)	All 2'-deoxynucleosides	1(c)-10(c)
306	1(a)-22(a)	All 2'-deoxynucleosides	1(d)-10(d)
307	1(b)-22(b)	All 2'-deoxynucleosides	1(d)-10(d)
308	1(c)-22(c)	All 2'-deoxynucleosides	1(d)-10(d)
309	1(d)-22(d)	All 2'-deoxynucleosides	1(d)-10(d)
310	1(e)-22(e)	All 2'-deoxynucleosides	1(d)-10(d)
311	1(f)-22(f)	All 2'-deoxynucleosides	1(d)-10(d)
312	1(g)-22(g)	All 2'-deoxynucleosides	1(d)-10(d)
313	1(h)-22(h)	All 2'-deoxynucleosides	1(d)-10(d)
314	1(i)-22(i)	All 2'-deoxynucleosides	1(d)-10(d)
315	1(a)-22(a)	All 2'-deoxynucleosides	1(e)-10(e)
316	1(b)-22(b)	All 2'-deoxynucleosides	1(e)-10(e)
317	1(c)-22(c)	All 2'-deoxynucleosides	1(e)-10(e)
318	1(d)-22(d)	All 2'-deoxynucleosides	1(e)-10(e)
319	1(e)-22(e)	All 2'-deoxynucleosides	1(e)-10(e)
320	1(f)-22(f)	All 2'-deoxynucleosides	1(e)-10(e)
321	1(g)-22(g)	All 2'-deoxynucleosides	1(e)-10(e)
322	1(h)-22(h)	All 2'-deoxynucleosides	1(e)-10(e)

323	1(i)-22(i)	All 2'-deoxynucleosides	1(e)-10(e)
324	1(a)-22(a)	All 2'-deoxynucleosides	1(f)-10(f)
325	1(b)-22(b)	All 2'-deoxynucleosides	1(f)-10(f)
326	1(c)-22(c)	All 2'-deoxynucleosides	1(f)-10(f)
327	1(d)-22(d)	All 2'-deoxynucleosides	1(f)-10(f)
328	1(e)-22(e)	All 2'-deoxynucleosides	1(f)-10(f)
329	1(f)-22(f)	All 2'-deoxynucleosides	1(f)-10(f)
330	1(g)-22(g)	All 2'-deoxynucleosides	1(f)-10(f)
331	1(h)-22(h)	All 2'-deoxynucleosides	1(f)-10(f)
332	1(i)-22(i)	All 2'-deoxynucleosides	1(f)-10(f)
333	1(a)-22(a)	All 2'-deoxynucleosides	1(g)-10(g)
334	1(b)-22(b)	All 2'-deoxynucleosides	1(g)-10(g)
335	1(c)-22(c)	All 2'-deoxynucleosides	1(g)-10(g)
336	1(d)-22(d)	All 2'-deoxynucleosides	1(g)-10(g)
337	1(e)-22(e)	All 2'-deoxynucleosides	1(g)-10(g)
338	1(f)-22(f)	All 2'-deoxynucleosides	1(g)-10(g)
339	1(g)-22(g)	All 2'-deoxynucleosides	1(g)-10(g)
340	1(h)-22(h)	All 2'-deoxynucleosides	1(g)-10(g)
341	1(i)-22(i)	All 2'-deoxynucleosides	1(g)-10(g)
342	1(a)-22(a)	All 2'-deoxynucleosides	1(h)-10(h)
343	1(b)-22(b)	All 2'-deoxynucleosides	1(h)-10(h)
344	1(c)-22(c)	All 2'-deoxynucleosides	1(h)-10(h)
345	1(d)-22(d)	All 2'-deoxynucleosides	1(h)-10(h)
346	1(e)-22(e)	All 2'-deoxynucleosides	1(h)-10(h)
347	1(f)-22(f)	All 2'-deoxynucleosides	1(h)-10(h)
348	1(g)-22(g)	All 2'-deoxynucleosides	1(h)-10(h)
349	1(h)-22(h)	All 2'-deoxynucleosides	1(h)-10(h)
350	1(i)-22(i)	All 2'-deoxynucleosides	1(h)-10(h)
351	1(a)-22(a)	All 2'-deoxynucleosides	1(i)-10(i)
352	1(b)-22(b)	All 2'-deoxynucleosides	1(i)-10(i)
353	1(c)-22(c)	All 2'-deoxynucleosides	1(i)-10(i)
354	1(d)-22(d)	All 2'-deoxynucleosides	1(i)-10(i)
355	1(e)-22(e)	All 2'-deoxynucleosides	1(i)-10(i)
356	1(f)-22(f)	All 2'-deoxynucleosides	1(i)-10(i)
357	1(g)-22(g)	All 2'-deoxynucleosides	1(i)-10(i)
358	1(h)-22(h)	All 2'-deoxynucleosides	1(i)-10(i)
359	1(i)-22(i)	All 2'-deoxynucleosides	1(i)-10(i)
360	1(a-l)	All 2'-deoxynucleosides	1(a-l)
361	2(a-l)	All 2'-deoxynucleosides	1(a-l)
362	3(a-l)	All 2'-deoxynucleosides	1(a-l)
363	4(a-l)	All 2'-deoxynucleosides	1(a-l)
364	5(a-l)	All 2'-deoxynucleosides	1(a-l)
365	6(a-l)	All 2'-deoxynucleosides	1(a-l)
366	7(a-l)	All 2'-deoxynucleosides	1(a-l)
367	8(a-l)	All 2'-deoxynucleosides	1(a-l)

368	9(a-l)	All 2'-deoxynucleosides	1(a-l)
369	10(a-l)	All 2'-deoxynucleosides	1(a-l)
370	11(a-l)	All 2'-deoxynucleosides	1(a-l)
371	12(a-l)	All 2'-deoxynucleosides	1(a-l)
372	13(a-l)	All 2'-deoxynucleosides	1(a-l)
373	14(a-l)	All 2'-deoxynucleosides	1(a-l)
374	15(a-l)	All 2'-deoxynucleosides	1(a-l)
375	16(a-l)	All 2'-deoxynucleosides	1(a-l)
376	17(a-l)	All 2'-deoxynucleosides	1(a-l)
377	18(a-l)	All 2'-deoxynucleosides	1(a-l)
378	19(a-l)	All 2'-deoxynucleosides	1(a-l)
379	20(a-l)	All 2'-deoxynucleosides	1(a-l)
380	21(a-l)	All 2'-deoxynucleosides	1(a-l)
381	22(a-l)	All 2'-deoxynucleosides	1(a-l)
382	1(a-l)	All 2'-deoxynucleosides	2(a-l)
383	2(a-l)	All 2'-deoxynucleosides	2(a-l)
384	3(a-l)	All 2'-deoxynucleosides	2(a-l)
385	4(a-l)	All 2'-deoxynucleosides	2(a-l)
386	5(a-l)	All 2'-deoxynucleosides	2(a-l)
387	6(a-l)	All 2'-deoxynucleosides	2(a-l)
388	7(a-l)	All 2'-deoxynucleosides	2(a-l)
389	8(a-l)	All 2'-deoxynucleosides	2(a-l)
390	9(a-l)	All 2'-deoxynucleosides	2(a-l)
391	10(a-l)	All 2'-deoxynucleosides	2(a-l)
392	11(a-l)	All 2'-deoxynucleosides	2(a-l)
393	12(a-l)	All 2'-deoxynucleosides	2(a-l)
394	13(a-l)	All 2'-deoxynucleosides	2(a-l)
395	14(a-l)	All 2'-deoxynucleosides	2(a-l)
396	15(a-l)	All 2'-deoxynucleosides	2(a-l)
397	16(a-l)	All 2'-deoxynucleosides	2(a-l)
398	17(a-l)	All 2'-deoxynucleosides	2(a-l)
399	18(a-l)	All 2'-deoxynucleosides	2(a-l)
400	19(a-l)	All 2'-deoxynucleosides	2(a-l)
401	20(a-l)	All 2'-deoxynucleosides	2(a-l)
402	21(a-l)	All 2'-deoxynucleosides	2(a-l)
403	22(a-l)	All 2'-deoxynucleosides	2(a-l)
404	1(a-l)	All 2'-deoxynucleosides	3(a-l)
405	2(a-l)	All 2'-deoxynucleosides	3(a-l)
406	3(a-l)	All 2'-deoxynucleosides	3(a-l)
407	4(a-l)	All 2'-deoxynucleosides	3(a-l)
408	5(a-l)	All 2'-deoxynucleosides	3(a-l)
409	6(a-l)	All 2'-deoxynucleosides	3(a-l)
410	7(a-l)	All 2'-deoxynucleosides	3(a-l)
411	8(a-l)	All 2'-deoxynucleosides	3(a-l)
412	9(a-l)	All 2'-deoxynucleosides	3(a-l)

413	10(a-l)	All 2'-deoxynucleosides	3(a-l)
414	11(a-l)	All 2'-deoxynucleosides	3(a-l)
415	12(a-l)	All 2'-deoxynucleosides	3(a-l)
416	13(a-l)	All 2'-deoxynucleosides	3(a-l)
417	14(a-l)	All 2'-deoxynucleosides	3(a-l)
418	15(a-l)	All 2'-deoxynucleosides	3(a-l)
419	16(a-l)	All 2'-deoxynucleosides	3(a-l)
420	17(a-l)	All 2'-deoxynucleosides	3(a-l)
421	18(a-l)	All 2'-deoxynucleosides	3(a-l)
422	19(a-l)	All 2'-deoxynucleosides	3(a-l)
423	20(a-l)	All 2'-deoxynucleosides	3(a-l)
424	21(a-l)	All 2'-deoxynucleosides	3(a-l)
425	22(a-l)	All 2'-deoxynucleosides	3(a-l)
426	1(a-l)	All 2'-deoxynucleosides	4(a-l)
427	2(a-l)	All 2'-deoxynucleosides	4(a-l)
428	3(a-l)	All 2'-deoxynucleosides	4(a-l)
429	4(a-l)	All 2'-deoxynucleosides	4(a-l)
430	5(a-l)	All 2'-deoxynucleosides	4(a-l)
431	6(a-l)	All 2'-deoxynucleosides	4(a-l)
432	7(a-l)	All 2'-deoxynucleosides	4(a-l)
433	8(a-l)	All 2'-deoxynucleosides	4(a-l)
434	9(a-l)	All 2'-deoxynucleosides	4(a-l)
435	10(a-l)	All 2'-deoxynucleosides	4(a-l)
436	11(a-l)	All 2'-deoxynucleosides	4(a-l)
437	12(a-l)	All 2'-deoxynucleosides	4(a-l)
438	13(a-l)	All 2'-deoxynucleosides	4(a-l)
439	14(a-l)	All 2'-deoxynucleosides	4(a-l)
440	15(a-l)	All 2'-deoxynucleosides	4(a-l)
441	16(a-l)	All 2'-deoxynucleosides	4(a-l)
442	17(a-l)	All 2'-deoxynucleosides	4(a-l)
443	18(a-l)	All 2'-deoxynucleosides	4(a-l)
444	19(a-l)	All 2'-deoxynucleosides	4(a-l)
445	20(a-l)	All 2'-deoxynucleosides	4(a-l)
446	21(a-l)	All 2'-deoxynucleosides	4(a-l)
447	22(a-l)	All 2'-deoxynucleosides	4(a-l)
448	1(a-l)	All 2'-deoxynucleosides	5(a-l)
449	2(a-l)	All 2'-deoxynucleosides	5(a-l)
450	3(a-l)	All 2'-deoxynucleosides	5(a-l)
451	4(a-l)	All 2'-deoxynucleosides	5(a-l)
452	5(a-l)	All 2'-deoxynucleosides	5(a-l)
453	6(a-l)	All 2'-deoxynucleosides	5(a-l)
454	7(a-l)	All 2'-deoxynucleosides	5(a-l)
455	8(a-l)	All 2'-deoxynucleosides	5(a-l)
456	9(a-l)	All 2'-deoxynucleosides	5(a-l)
457	10(a-l)	All 2'-deoxynucleosides	5(a-l)

458	11(a-l)	All 2'-deoxynucleosides	5(a-l)
459	12(a-l)	All 2'-deoxynucleosides	5(a-l)
460	13(a-l)	All 2'-deoxynucleosides	5(a-l)
461	14(a-l)	All 2'-deoxynucleosides	5(a-l)
462	15(a-l)	All 2'-deoxynucleosides	5(a-l)
463	16(a-l)	All 2'-deoxynucleosides	5(a-l)
464	17(a-l)	All 2'-deoxynucleosides	5(a-l)
465	18(a-l)	All 2'-deoxynucleosides	5(a-l)
466	19(a-l)	All 2'-deoxynucleosides	5(a-l)
467	20(a-l)	All 2'-deoxynucleosides	5(a-l)
468	21(a-l)	All 2'-deoxynucleosides	5(a-l)
469	22(a-l)	All 2'-deoxynucleosides	5(a-l)
470	1(a-l)	All 2'-deoxynucleosides	6(a-l)
471	2(a-l)	All 2'-deoxynucleosides	6(a-l)
472	3(a-l)	All 2'-deoxynucleosides	6(a-l)
473	4(a-l)	All 2'-deoxynucleosides	6(a-l)
474	5(a-l)	All 2'-deoxynucleosides	6(a-l)
475	6(a-l)	All 2'-deoxynucleosides	6(a-l)
476	7(a-l)	All 2'-deoxynucleosides	6(a-l)
477	8(a-l)	All 2'-deoxynucleosides	6(a-l)
478	9(a-l)	All 2'-deoxynucleosides	6(a-l)
479	10(a-l)	All 2'-deoxynucleosides	6(a-l)
480	11(a-l)	All 2'-deoxynucleosides	6(a-l)
481	12(a-l)	All 2'-deoxynucleosides	6(a-l)
482	13(a-l)	All 2'-deoxynucleosides	6(a-l)
483	14(a-l)	All 2'-deoxynucleosides	6(a-l)
484	15(a-l)	All 2'-deoxynucleosides	6(a-l)
485	16(a-l)	All 2'-deoxynucleosides	6(a-l)
486	17(a-l)	All 2'-deoxynucleosides	6(a-l)
487	18(a-l)	All 2'-deoxynucleosides	6(a-l)
488	19(a-l)	All 2'-deoxynucleosides	6(a-l)
489	20(a-l)	All 2'-deoxynucleosides	6(a-l)
490	21(a-l)	All 2'-deoxynucleosides	6(a-l)
491	22(a-l)	All 2'-deoxynucleosides	6(a-l)
492	1(a-l)	All 2'-deoxynucleosides	7(a-l)
493	2(a-l)	All 2'-deoxynucleosides	7(a-l)
494	3(a-l)	All 2'-deoxynucleosides	7(a-l)
495	4(a-l)	All 2'-deoxynucleosides	7(a-l)
496	5(a-l)	All 2'-deoxynucleosides	7(a-l)
497	6(a-l)	All 2'-deoxynucleosides	7(a-l)
498	7(a-l)	All 2'-deoxynucleosides	7(a-l)
499	8(a-l)	All 2'-deoxynucleosides	7(a-l)
500	9(a-l)	All 2'-deoxynucleosides	7(a-l)
501	10(a-l)	All 2'-deoxynucleosides	7(a-l)
502	11(a-l)	All 2'-deoxynucleosides	7(a-l)

503	12(a-l)	All 2'-deoxynucleosides	7(a-l)
504	13(a-l)	All 2'-deoxynucleosides	7(a-l)
505	14(a-l)	All 2'-deoxynucleosides	7(a-l)
506	15(a-l)	All 2'-deoxynucleosides	7(a-l)
507	16(a-l)	All 2'-deoxynucleosides	7(a-l)
508	17(a-l)	All 2'-deoxynucleosides	7(a-l)
509	18(a-l)	All 2'-deoxynucleosides	7(a-l)
510	19(a-l)	All 2'-deoxynucleosides	7(a-l)
511	20(a-l)	All 2'-deoxynucleosides	7(a-l)
512	21(a-l)	All 2'-deoxynucleosides	7(a-l)
513	22(a-l)	All 2'-deoxynucleosides	7(a-l)
514	1(a-l)	All 2'-deoxynucleosides	8(a-l)
515	2(a-l)	All 2'-deoxynucleosides	8(a-l)
516	3(a-l)	All 2'-deoxynucleosides	8(a-l)
517	4(a-l)	All 2'-deoxynucleosides	8(a-l)
518	5(a-l)	All 2'-deoxynucleosides	8(a-l)
519	6(a-l)	All 2'-deoxynucleosides	8(a-l)
520	7(a-l)	All 2'-deoxynucleosides	8(a-l)
521	8(a-l)	All 2'-deoxynucleosides	8(a-l)
522	9(a-l)	All 2'-deoxynucleosides	8(a-l)
523	10(a-l)	All 2'-deoxynucleosides	8(a-l)
524	11(a-l)	All 2'-deoxynucleosides	8(a-l)
525	12(a-l)	All 2'-deoxynucleosides	8(a-l)
526	13(a-l)	All 2'-deoxynucleosides	8(a-l)
527	14(a-l)	All 2'-deoxynucleosides	8(a-l)
528	15(a-l)	All 2'-deoxynucleosides	8(a-l)
529	16(a-l)	All 2'-deoxynucleosides	8(a-l)
530	17(a-l)	All 2'-deoxynucleosides	8(a-l)
531	18(a-l)	All 2'-deoxynucleosides	8(a-l)
532	19(a-l)	All 2'-deoxynucleosides	8(a-l)
533	20(a-l)	All 2'-deoxynucleosides	8(a-l)
534	21(a-l)	All 2'-deoxynucleosides	8(a-l)
535	22(a-l)	All 2'-deoxynucleosides	8(a-l)
536	1(a-l)	All 2'-deoxynucleosides	9(a-l)
537	2(a-l)	All 2'-deoxynucleosides	9(a-l)
538	3(a-l)	All 2'-deoxynucleosides	9(a-l)
539	4(a-l)	All 2'-deoxynucleosides	9(a-l)
540	5(a-l)	All 2'-deoxynucleosides	9(a-l)
541	6(a-l)	All 2'-deoxynucleosides	9(a-l)
542	7(a-l)	All 2'-deoxynucleosides	9(a-l)
543	8(a-l)	All 2'-deoxynucleosides	9(a-l)
544	9(a-l)	All 2'-deoxynucleosides	9(a-l)
545	10(a-l)	All 2'-deoxynucleosides	9(a-l)
546	11(a-l)	All 2'-deoxynucleosides	9(a-l)
547	12(a-l)	All 2'-deoxynucleosides	9(a-l)

548	13(a-l)	All 2'-deoxynucleosides	9(a-l)
549	14(a-l)	All 2'-deoxynucleosides	9(a-l)
550	15(a-l)	All 2'-deoxynucleosides	9(a-l)
551	16(a-l)	All 2'-deoxynucleosides	9(a-l)
552	17(a-l)	All 2'-deoxynucleosides	9(a-l)
553	18(a-l)	All 2'-deoxynucleosides	9(a-l)
554	19(a-l)	All 2'-deoxynucleosides	9(a-l)
555	20(a-l)	All 2'-deoxynucleosides	9(a-l)
556	21(a-l)	All 2'-deoxynucleosides	9(a-l)
557	22(a-l)	All 2'-deoxynucleosides	9(a-l)
558	1(a-l)	All 2'-deoxynucleosides	10(a-l)
559	2(a-l)	All 2'-deoxynucleosides	10(a-l)
560	3(a-l)	All 2'-deoxynucleosides	10(a-l)
561	4(a-l)	All 2'-deoxynucleosides	10(a-l)
562	5(a-l)	All 2'-deoxynucleosides	10(a-l)
563	6(a-l)	All 2'-deoxynucleosides	10(a-l)
564	7(a-l)	All 2'-deoxynucleosides	10(a-l)
565	8(a-l)	All 2'-deoxynucleosides	10(a-l)
566	9(a-l)	All 2'-deoxynucleosides	10(a-l)
567	10(a-l)	All 2'-deoxynucleosides	10(a-l)
568	11(a-l)	All 2'-deoxynucleosides	10(a-l)
569	12(a-l)	All 2'-deoxynucleosides	10(a-l)
570	13(a-l)	All 2'-deoxynucleosides	10(a-l)
571	14(a-l)	All 2'-deoxynucleosides	10(a-l)
572	15(a-l)	All 2'-deoxynucleosides	10(a-l)
573	16(a-l)	All 2'-deoxynucleosides	10(a-l)
574	17(a-l)	All 2'-deoxynucleosides	10(a-l)
575	18(a-l)	All 2'-deoxynucleosides	10(a-l)
576	19(a-l)	All 2'-deoxynucleosides	10(a-l)
577	20(a-l)	All 2'-deoxynucleosides	10(a-l)
578	21(a-l)	All 2'-deoxynucleosides	10(a-l)
579	22(a-l)	All 2'-deoxynucleosides	10(a-l)
580	1(j)-22(j)	All 2'-deoxynucleosides	1(a)-10(a)
581	1(k)-22(k)	All 2'-deoxynucleosides	1(a)-10(a)
582	1(l)-22(l)	All 2'-deoxynucleosides	1(a)-10(a)
583	1(j)-22(j)	All 2'-deoxynucleosides	1(b)-10(b)
584	1(k)-22(k)	All 2'-deoxynucleosides	1(b)-10(b)
585	1(l)-22(l)	All 2'-deoxynucleosides	1(b)-10(b)
586	1(j)-22(j)	All 2'-deoxynucleosides	1(c)-10(c)
587	1(k)-22(k)	All 2'-deoxynucleosides	1(c)-10(c)
588	1(l)-22(l)	All 2'-deoxynucleosides	1(c)-10(c)
589	1(j)-22(j)	All 2'-deoxynucleosides	1(d)-10(d)
590	1(k)-22(k)	All 2'-deoxynucleosides	1(d)-10(d)
591	1(l)-22(l)	All 2'-deoxynucleosides	1(d)-10(d)
592	1(j)-22(j)	All 2'-deoxynucleosides	1(e)-10(e)

593	1(k)-22(k)	All 2'-deoxynucleosides	1(e)-10(e)
594	1(l)-22(l)	All 2'-deoxynucleosides	1(e)-10(e)
595	1(j)-22(j)	All 2'-deoxynucleosides	1(f)-10(f)
596	1(k)-22(k)	All 2'-deoxynucleosides	1(f)-10(f)
597	1(l)-22(l)	All 2'-deoxynucleosides	1(f)-10(f)
598	1(j)-22(j)	All 2'-deoxynucleosides	1(g)-10(g)
599	1(k)-22(k)	All 2'-deoxynucleosides	1(g)-10(g)
600	1(l)-22(l)	All 2'-deoxynucleosides	1(g)-10(g)
601	1(j)-22(j)	All 2'-deoxynucleosides	1(h)-10(h)
602	1(k)-22(k)	All 2'-deoxynucleosides	1(h)-10(h)
603	1(l)-22(l)	All 2'-deoxynucleosides	1(h)-10(h)
604	1(j)-22(j)	All 2'-deoxynucleosides	1(i)-10(i)
605	1(k)-22(k)	All 2'-deoxynucleosides	1(i)-10(i)
606	1(l)-22(l)	All 2'-deoxynucleosides	1(i)-10(i)
607	1(j)-22(j)	All 2'-deoxynucleosides	1(j)-10(j)
608	1(k)-22(k)	All 2'-deoxynucleosides	1(j)-10(j)
609	1(l)-22(l)	All 2'-deoxynucleosides	1(j)-10(j)
610	1(j)-22(j)	All 2'-deoxynucleosides	1(k)-10(k)
611	1(k)-22(k)	All 2'-deoxynucleosides	1(k)-10(k)
612	1(l)-22(l)	All 2'-deoxynucleosides	1(k)-10(k)
612	1(j)-22(j)	All 2'-deoxynucleosides	1(l)-10(l)
614	1(k)-22(k)	All 2'-deoxynucleosides	1(l)-10(l)
615	1(l)-22(l)	All 2'-deoxynucleosides	1(l)-10(l)
616	1k	All 2'-deoxynucleosides	1m

In certain embodiments, a gapmer comprises a 5'-wing selected from among the 5'-wings provided herein and any 3'-wing. In certain embodiments, a gapmer comprises a 5'-wing selected from among 1(a-i) to 22(a-i). In certain embodiments, a gapmer comprises a 5'-wing selected from among 1(a-l) to 22(a-l). In certain embodiments, a gapmer comprises a 3'-wing selected from among the 3'-wings provided herein and any 5'-wing. In certain embodiments, a gapmer comprises a 3'-wing selected from among 1(a-i) to 10(a-i). In certain embodiments, a gapmer comprises a 3'-wing selected from among 1(a-l) to 10(a-l).

In certain embodiments, a gapmer has a sugar motif other than: E-K-K-(D)₉-K-K-E; E-E-E-E-K-(D)₉-E-E-E-E-E; E-K-K-K-(D)₉-K-K-K-E; K-E-E-K-(D)₉-K-E-E-K; K-D-D-K-(D)₉-K-D-D-K; K-E-K-E-K-(D)₉-K-E-K-E-K; K-D-K-D-K-(D)₉-K-D-K-D-K; E-K-E-K-(D)₉-K-E-K-E; E-E-E-E-E-K-(D)₈-E-E-E-E-E; or E-K-E-K-E-(D)₉-E-K-E-K-E. In certain embodiments, a gapmer not having one of the above motifs has a sugar motif of Formula I. In certain embodiments, a gapmer not having one of the above motifs has a sugar motif selected from motifs 1-58. In certain embodiments, a gapmer not having one of the above motifs has a sugar motif of Formula I and selected from sugar motifs 1-58. In certain embodiments, a gapmer not having one of the above motifs has a sugar motif of Formula II. In certain embodiments, a gapmer not having one of the above motifs has a sugar motif selected from motifs 1-615. In certain embodiments, a gapmer not having one of the above motifs has a sugar motif of Formula II and selected from sugar motifs 1-615.

In certain embodiments a gapmer comprises a A-(D)₄-A-(D)₄-A-(D)₄-AA motif. In certain embodiments a gapmer comprises a B-(D)₄-A-(D)₄-A-(D)₄-AA motif. In certain embodiments a gapmer comprises a A-(D)₄-B-(D)₄-A-(D)₄-AA motif. In certain embodiments a gapmer comprises a A-(D)₄-A-(D)₄-B-(D)₄-AA motif. In certain embodiments a gapmer comprises a A-(D)₄-A-(D)₄-A-(D)₄-BA motif. In certain embodiments a gapmer comprises a A-(D)₄-A-(D)₄-A-(D)₄-BB motif. In certain embodiments a gapmer comprises a K-(D)₄-K-(D)₄-K-(D)₄-K-E motif.

Certain Internucleoside Linkage Motifs

In certain embodiments, oligonucleotides comprise modified internucleoside linkages arranged along the oligonucleotide or region thereof in a defined pattern or modified internucleoside linkage motif. In certain embodiments, internucleoside linkages are arranged in a gapped motif, as described above for sugar modification motif. In such embodiments, the internucleoside linkages in each of two wing regions are different from the internucleoside linkages in the gap region. In certain embodiments the internucleoside linkages in the wings are phosphodiester and the internucleoside linkages in the gap are phosphorothioate. The sugar modification motif is independently selected, so such oligonucleotides having a gapped internucleoside linkage motif may or may not have a gapped sugar modification motif and if it does have a gapped sugar motif, the wing and gap lengths may or may not be the same.

In certain embodiments, oligonucleotides comprise a region having an alternating internucleoside linkage motif. In certain embodiments, oligonucleotides of the present invention comprise a region of uniformly modified internucleoside linkages. In certain such embodiments, the oligonucleotide comprises a region that is uniformly linked by phosphorothioate internucleoside linkages. In certain embodiments, the oligonucleotide is uniformly linked by phosphorothioate. In certain embodiments, each internucleoside linkage of the oligonucleotide is selected from phosphodiester and phosphorothioate. In certain embodiments, each internucleoside linkage of the oligonucleotide is selected from phosphodiester and phosphorothioate and at least one internucleoside linkage is phosphorothioate.

In certain embodiments, the oligonucleotide comprises at least 6 phosphorothioate internucleoside linkages. In certain embodiments, the oligonucleotide comprises at least 8 phosphorothioate internucleoside linkages. In certain embodiments, the oligonucleotide comprises at least 10 phosphorothioate internucleoside linkages. In certain embodiments, the oligonucleotide comprises at least one block of at least 6 consecutive phosphorothioate internucleoside linkages. In certain embodiments, the oligonucleotide comprises at least one block of at least 8 consecutive phosphorothioate internucleoside linkages. In certain embodiments, the oligonucleotide comprises at least one block of at least 10 consecutive phosphorothioate internucleoside linkages. In certain embodiments, the oligonucleotide comprises at least one block of at least 12 consecutive phosphorothioate internucleoside linkages. In certain such embodiments, at least one such block is located at the 3' end of the oligonucleotide. In certain such embodiments, at least one such block is located within 3 nucleosides of the 3' end of the oligonucleotide.

Certain Nucleobase Modification Motifs

In certain embodiments, oligonucleotides comprise chemical modifications to nucleobases arranged along the oligonucleotide or region thereof in a defined pattern or nucleobases modification motif. In certain such embodiments, nucleobase modifications are arranged in a gapped motif. In certain embodiments, nucleobase modifications are arranged in an alternating motif. In certain embodiments, each nucleobase is modified. In certain embodiments, none of the nucleobases is chemically modified.

In certain embodiments, oligonucleotides comprise a block of modified nucleobases. In certain such embodiments, the block is at the 3'-end of the oligonucleotide. In certain embodiments the block is within 3 nucleotides of the 3'-end of the oligonucleotide. In certain such embodiments, the block is at the 5'-end of the oligonucleotide. In certain embodiments the block is within 3 nucleotides of the 5'-end of the oligonucleotide.

In certain embodiments, nucleobase modifications are a function of the natural base at a particular position of an oligonucleotide. For example, in certain embodiments each purine or each pyrimidine in an oligonucleotide is modified. In certain embodiments, each adenine is modified. In certain embodiments, each guanine is modified. In certain embodiments, each thymine is modified. In certain embodiments, each cytosine is modified. In certain embodiments, each uracil is modified.

In certain embodiments, some, all, or none of the cytosine moieties in an oligonucleotide are 5-methyl cytosine moieties. Herein, 5-methyl cytosine is not a "modified nucleobase." Accordingly, unless otherwise indicated, unmodified nucleobases include both cytosine residues having a 5-methyl and those lacking a 5 methyl. In certain embodiments, the methylation state of all or some cytosine nucleobases is specified.

Certain Overall Lengths

In certain embodiments, the present invention provides oligomeric compounds including oligonucleotides of any of a variety of ranges of lengths. In certain embodiments, the invention provides oligomeric compounds or oligonucleotides consisting of X to Y linked nucleosides, where X represents the fewest number of nucleosides in the range and Y represents the largest number of nucleosides in the range. In certain such embodiments, X and Y are each independently selected from 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, and 50; provided that $X \leq Y$. For example, in certain embodiments, the invention provides oligomeric compounds which comprise oligonucleotides consisting of 8 to 9, 8 to 10, 8 to 11, 8 to 12, 8 to 13, 8 to 14, 8 to 15, 8 to 16, 8 to 17, 8 to 18, 8 to 19, 8 to 20, 8 to 21, 8 to 22, 8 to 23, 8 to 24, 8 to 25, 8 to 26, 8 to 27, 8 to 28, 8 to 29, 8 to 30, 9 to 10, 9 to 11, 9 to 12, 9 to 13, 9 to 14, 9 to 15, 9 to 16, 9 to 17, 9 to 18, 9 to 19, 9 to 20, 9 to 21, 9 to 22, 9 to 23, 9 to 24, 9 to 25, 9 to 26, 9 to 27, 9 to 28, 9 to 29, 9 to 30, 10 to 11, 10 to 12, 10 to 13, 10 to 14, 10 to 15, 10 to 16, 10 to 17, 10 to 18, 10 to 19, 10 to 20, 10 to 21, 10 to 22, 10 to 23,

10 to 24, 10 to 25, 10 to 26, 10 to 27, 10 to 28, 10 to 29, 10 to 30, 11 to 12, 11 to 13, 11 to 14, 11 to 15, 11 to 16, 11 to 17, 11 to 18, 11 to 19, 11 to 20, 11 to 21, 11 to 22, 11 to 23, 11 to 24, 11 to 25, 11 to 26, 11 to 27, 11 to 28, 11 to 29, 11 to 30, 12 to 13, 12 to 14, 12 to 15, 12 to 16, 12 to 17, 12 to 18, 12 to 19, 12 to 20, 12 to 21, 12 to 22, 12 to 23, 12 to 24, 12 to 25, 12 to 26, 12 to 27, 12 to 28, 12 to 29, 12 to 30, 13 to 14, 13 to 15, 13 to 16, 13 to 17, 13 to 18, 13 to 19, 13 to 20, 13 to 21, 13 to 22, 13 to 23, 13 to 24, 13 to 25, 13 to 26, 13 to 27, 13 to 28, 13 to 29, 13 to 30, 14 to 15, 14 to 16, 14 to 17, 14 to 18, 14 to 19, 14 to 20, 14 to 21, 14 to 22, 14 to 23, 14 to 24, 14 to 25, 14 to 26, 14 to 27, 14 to 28, 14 to 29, 14 to 30, 15 to 16, 15 to 17, 15 to 18, 15 to 19, 15 to 20, 15 to 21, 15 to 22, 15 to 23, 15 to 24, 15 to 25, 15 to 26, 15 to 27, 15 to 28, 15 to 29, 15 to 30, 16 to 17, 16 to 18, 16 to 19, 16 to 20, 16 to 21, 16 to 22, 16 to 23, 16 to 24, 16 to 25, 16 to 26, 16 to 27, 16 to 28, 16 to 29, 16 to 30, 17 to 18, 17 to 19, 17 to 20, 17 to 21, 17 to 22, 17 to 23, 17 to 24, 17 to 25, 17 to 26, 17 to 27, 17 to 28, 17 to 29, 17 to 30, 18 to 19, 18 to 20, 18 to 21, 18 to 22, 18 to 23, 18 to 24, 18 to 25, 18 to 26, 18 to 27, 18 to 28, 18 to 29, 18 to 30, 19 to 20, 19 to 21, 19 to 22, 19 to 23, 19 to 24, 19 to 25, 19 to 26, 19 to 27, 19 to 28, 19 to 29, 19 to 30, 20 to 21, 20 to 22, 20 to 23, 20 to 24, 20 to 25, 20 to 26, 20 to 27, 20 to 28, 20 to 29, 20 to 30, 21 to 22, 21 to 23, 21 to 24, 21 to 25, 21 to 26, 21 to 27, 21 to 28, 21 to 29, 21 to 30, 22 to 23, 22 to 24, 22 to 25, 22 to 26, 22 to 27, 22 to 28, 22 to 29, 22 to 30, 23 to 24, 23 to 25, 23 to 26, 23 to 27, 23 to 28, 23 to 29, 23 to 30, 24 to 25, 24 to 26, 24 to 27, 24 to 28, 24 to 29, 24 to 30, 25 to 26, 25 to 27, 25 to 28, 25 to 29, 25 to 30, 26 to 27, 26 to 28, 26 to 29, 26 to 30, 27 to 28, 27 to 29, 27 to 30, 28 to 29, 28 to 30, or 29 to 30 linked nucleosides. In embodiments where the number of nucleosides of an oligomeric compound or oligonucleotide is limited, whether to a range or to a specific number, the oligomeric compound or oligonucleotide may, nonetheless further comprise additional other substituents. For example, an oligonucleotide comprising 8-30 nucleosides excludes oligonucleotides having 31 nucleosides, but, unless otherwise indicated, such an oligonucleotide may further comprise, for example one or more conjugates, terminal groups, or other substituents. In certain embodiments, a gapmer oligonucleotide has any of the above lengths.

In certain embodiments, any of the gapmer motifs provided above, including but not limited to gapmer motifs 1-278 provided in Tables 3 and 4, may have any of the above lengths. One of skill in the art will appreciate that certain lengths may not be possible for certain motifs. For example: a gapmer having a 5'-wing region consisting of four nucleotides, a gap consisting of at least six nucleotides, and a 3'-wing region consisting of three nucleotides cannot have an overall length less than 13 nucleotides. Thus, one would understand that the lower length limit is 13 and that the limit of 10 in "10-20" has no effect in that embodiment.

Further, where an oligonucleotide is described by an overall length range and by regions having specified lengths, and where the sum of specified lengths of the regions is less than the upper limit of the overall length range, the oligonucleotide may have additional nucleosides, beyond those of the specified regions, provided that the total number of nucleosides does not exceed the upper limit of the overall length range. For example, an oligonucleotide consisting of 20-25 linked nucleosides comprising a 5'-wing

consisting of 5 linked nucleosides; a 3'-wing consisting of 5 linked nucleosides and a central gap consisting of 10 linked nucleosides ($5+5+10=20$) may have up to 5 nucleosides that are not part of the 5'-wing, the 3'-wing, or the gap (before reaching the overall length limitation of 25). Such additional nucleosides may be 5' of the 5'-wing and/or 3' of the 3' wing.

Certain Oligonucleotides

In certain embodiments, oligonucleotides of the present invention are characterized by their sugar motif, internucleoside linkage motif, nucleobase modification motif and overall length. In certain embodiments, such parameters are each independent of one another. Thus, each internucleoside linkage of an oligonucleotide having a gapmer sugar motif may be modified or unmodified and may or may not follow the gapmer modification pattern of the sugar modifications. Thus, the internucleoside linkages within the wing regions of a sugar-gapmer may be the same or different from one another and may be the same or different from the internucleoside linkages of the gap region. Likewise, such sugar-gapmer oligonucleotides may comprise one or more modified nucleobase independent of the gapmer pattern of the sugar modifications. One of skill in the art will appreciate that such motifs may be combined to create a variety of oligonucleotides, such as those provided in the non-limiting Table 5 below.

Table 10
Certain Oligonucleotides

Overall Length	Sugar motif	Internucleoside Linkage Motif	Nucleobase Mod. Motif
12	Gapmer motif selected from 1-278	uniform PS	uniform unmodified
14	Gapmer motif selected from 1-278	2-14-2 gapmer: PO in wings and PS in gap	uniform unmodified
14	Gapmer motif selected from 1-278	uniform PS	uniform unmodified; all C's are 5-meC
16	Gapmer of Formula I	uniform PS	uniform unmodified; no Cs are 5-meC)
16	Gapmer of Formula I	uniform PS	uniform unmodified; at least one nucleobase is a 5-meC
16	Gapmer of Formula I and having motif selected from 1-58	uniform PS	uniform unmodified
17	Gapmer of Formula I and having motif selected from 1-58	uniform PO	uniform unmodified
17	Gapmer motif selected from 1-278	uniform PS	uniform unmodified
17	Gapmer of Formula I	uniform PS	uniform unmodified

18	Gapmer of Formula I and having motif selected from 1-58	uniform PS	uniform unmodified
18	Gapmer motif selected from 1-278	uniform PS	uniform unmodified
20	Gapmer of Formula I	uniform PS	uniform unmodified
12	Gapmer motif selected from 1-359	uniform PS	uniform unmodified
14	Gapmer motif selected from 1-359	2-14-2 gapmer: PO in wings and PS in gap	uniform unmodified
14	Gapmer motif selected from 1-359	uniform PS	uniform unmodified; all C's are 5-meC
16	Gapmer of Formula II	uniform PS	uniform unmodified; no Cs are 5-meC)
16	Gapmer of Formula II	uniform PS	uniform unmodified; at least one nucleobase is a 5-meC
16	Gapmer of Formula II and having motif selected from 1-359	uniform PS	uniform unmodified
17	Gapmer of Formula II and having motif selected from 1-359	uniform PO	uniform unmodified
17	Gapmer motif selected from 1-359	uniform PS	uniform unmodified
17	Gapmer of Formula II	uniform PS	uniform unmodified
18	Gapmer of Formula I and having motif selected from 1-359	uniform PS	uniform unmodified
18	Gapmer motif selected from 1-359	uniform PS	uniform unmodified
20	Gapmer of Formula II	uniform PS	uniform unmodified
12	Gapmer motif selected from 1-615	uniform PS	uniform unmodified
14	Gapmer motif selected from 1-615	2-14-2 gapmer: PO in wings and PS in gap	uniform unmodified
14	Gapmer motif selected from 1-615	uniform PS	uniform unmodified; all C's are 5-meC
16	Gapmer of Formula I and having motif selected from 1-615	uniform PS	uniform unmodified
17	Gapmer of Formula I and having motif selected from 1-615	uniform PO	uniform unmodified
17	Gapmer motif selected from 1-615	uniform PS	uniform unmodified
18	Gapmer of Formula I and having motif selected from 1-615	uniform PS	uniform unmodified
18	Gapmer motif selected from 1-615	uniform PS	uniform unmodified

The above table is intended only to illustrate and not to limit the various combinations of the parameters of oligonucleotides of the present invention. Herein if a description of an oligonucleotide or oligomeric compound is silent with respect to one or more parameter, such parameter is not limited. Thus, an oligomeric compound described only as having a gapmer sugar motif without further description may have any length, internucleoside linkage motif, and nucleobase modification motif. Unless otherwise indicated, all chemical modifications are independent of nucleobase sequence.

Certain Conjugate Groups

In certain embodiments, oligomeric compounds are modified by attachment of one or more conjugate groups. In general, conjugate groups modify one or more properties of the attached oligomeric compound including but not limited to pharmacodynamics, pharmacokinetics, stability, binding, absorption, cellular distribution, cellular uptake, charge and clearance. Conjugate groups are routinely used in the chemical arts and are linked directly or via an optional conjugate linking moiety or conjugate linking group to a parent compound such as an oligomeric compound, such as an oligonucleotide. Conjugate groups includes without limitation, intercalators, reporter molecules, polyamines, polyamides, polyethylene glycols, thioethers, polyethers, cholesterol, thiocholesterol, cholic acid moieties, folate, lipids, phospholipids, biotin, phenazine, phenanthridine, anthraquinone, adamantane, acridine, fluoresceins, rhodamines, coumarins and dyes. Certain conjugate groups have been described previously, for example: cholesterol moiety (Letsinger et al., Proc. Natl. Acad. Sci. USA, 1989, 86, 6553-6556), cholic acid (Manoharan et al., Bioorg. Med. Chem. Lett., 1994, 4, 1053-1060), a thioether, e.g., hexyl-S-tritylthiol (Manoharan et al., Ann. N.Y. Acad. Sci., 1992, 660, 306-309; Manoharan et al., Bioorg. Med. Chem. Lett., 1993, 3, 2765-2770), a thiocholesterol (Oberhauser et al., Nucl. Acids Res., 1992, 20, 533-538), an aliphatic chain, e.g., do-decan-diol or undecyl residues (Saison-Behmoaras et al., EMBO J., 1991, 10, 1111-1118; Kabanov et al., FEBS Lett., 1990, 259, 327-330; Svinarchuk et al., Biochimie, 1993, 75, 49-54), a phospholipid, e.g., di-hexadecyl-rac-glycerol or triethyl-ammonium 1,2-di-O-hexadecyl-rac-glycero-3-H-phosphonate (Manoharan et al., Tetrahedron Lett., 1995, 36, 3651-3654; Shea et al., Nucl. Acids Res., 1990, 18, 3777-3783), a polyamine or a polyethylene glycol chain (Manoharan et al., Nucleosides & Nucleotides, 1995, 14, 969-973), or adamantane acetic acid (Manoharan et al., Tetrahedron Lett., 1995, 36, 3651-3654), a palmityl moiety (Mishra et al., Biochim. Biophys. Acta, 1995, 1264, 229-237), or an octadecylamine or hexylamino-carbonyl-oxycholesterol moiety (Crooke et al., J. Pharmacol. Exp. Ther., 1996, 277, 923-937).

In certain embodiments, a conjugate group comprises an active drug substance, for example, aspirin, warfarin, phenylbutazone, ibuprofen, suprofen, fen-bufen, ketoprofen, (S)-(+)-pranoprofen, carprofen, dansylsarcosine, 2,3,5-triiodobenzoic acid, flufenamic acid, folinic acid, a benzothiadiazide, chlorothiazide, a diazepam, indo-methicin, a barbiturate, a cephalosporin, a sulfa drug, an antidiabetic, an antibacterial or an antibiotic.

In certain embodiments, conjugate groups are directly attached to oligonucleotides in oligomeric

compounds. In certain embodiments, conjugate groups are attached to oligonucleotides by a conjugate linking group. In certain such embodiments, conjugate linking groups, including, but not limited to, bifunctional linking moieties such as those known in the art are amenable to the compounds provided herein. Conjugate linking groups are useful for attachment of conjugate groups, such as chemical stabilizing groups, functional groups, reporter groups and other groups to selective sites in a parent compound such as for example an oligomeric compound. In general a bifunctional linking moiety comprises a hydrocarbyl moiety having two functional groups. One of the functional groups is selected to bind to a parent molecule or compound of interest and the other is selected to bind essentially any selected group such as chemical functional group or a conjugate group. In some embodiments, the conjugate linker comprises a chain structure or an oligomer of repeating units such as ethylene glycol or amino acid units. Examples of functional groups that are routinely used in a bifunctional linking moiety include, but are not limited to, electrophiles for reacting with nucleophilic groups and nucleophiles for reacting with electrophilic groups. In some embodiments, bifunctional linking moieties include amino, hydroxyl, carboxylic acid, thiol, unsaturations (e.g., double or triple bonds), and the like.

Some nonlimiting examples of conjugate linking moieties include pyrrolidine, 8-amino-3,6-dioxaoctanoic acid (ADO), succinimidyl 4-(N-maleimidomethyl) cyclohexane-1-carboxylate (SMCC) and 6-aminohexanoic acid (AHX or AHA). Other linking groups include, but are not limited to, substituted C₁-C₁₀ alkyl, substituted or unsubstituted C₂-C₁₀ alkenyl or substituted or unsubstituted C₂-C₁₀ alkynyl, wherein a nonlimiting list of preferred substituent groups includes hydroxyl, amino, alkoxy, carboxy, benzyl, phenyl, nitro, thiol, thioalkoxy, halogen, alkyl, aryl, alkenyl and alkynyl.

Conjugate groups may be attached to either or both ends of an oligonucleotide (terminal conjugate groups) and/or at any internal position.

In certain embodiments, conjugate groups are at the 3'-end of an oligonucleotide of an oligomeric compound. In certain embodiments, conjugate groups are near the 3'-end. In certain embodiments, conjugates are attached at the 3' end of an oligomeric compound, but before one or more terminal group nucleosides. In certain embodiments, conjugate groups are placed within a terminal group. In certain embodiments, the present invention provides oligomeric compounds. In certain embodiments, oligomeric compounds comprise an oligonucleotide. In certain embodiments, an oligomeric compound comprises an oligonucleotide and one or more conjugate and/or terminal groups. Such conjugate and/or terminal groups may be added to oligonucleotides having any of the chemical motifs discussed above. Thus, for example, an oligomeric compound comprising an oligonucleotide having region of alternating nucleosides may comprise a terminal group.

Antisense Compounds

In certain embodiments, oligomeric compounds of the present invention are antisense compounds. Such antisense compounds are capable of hybridizing to a target nucleic acid, resulting in at least one

antisense activity. In certain embodiments, antisense compounds specifically hybridize to one or more target nucleic acid. In certain embodiments, a specifically hybridizing antisense compound has a nucleobase sequence comprising a region having sufficient complementarity to a target nucleic acid to allow hybridization and result in antisense activity and insufficient complementarity to any non-target so as to avoid non-specific hybridization to any non-target nucleic acid sequences under conditions in which specific hybridization is desired (e.g., under physiological conditions for *in vivo* or therapeutic uses, and under conditions in which assays are performed in the case of *in vitro* assays).

In certain embodiments, the present invention provides antisense compounds comprising oligonucleotides that are fully complementary to the target nucleic acid over the entire length of the oligonucleotide. In certain embodiments, oligonucleotides are 99% complementary to the target nucleic acid. In certain embodiments, oligonucleotides are 95% complementary to the target nucleic acid. In certain embodiments, such oligonucleotides are 90% complementary to the target nucleic acid.

In certain embodiments, such oligonucleotides are 85% complementary to the target nucleic acid. In certain embodiments, such oligonucleotides are 80% complementary to the target nucleic acid. In certain embodiments, an antisense compound comprises a region that is fully complementary to a target nucleic acid and is at least 80% complementary to the target nucleic acid over the entire length of the oligonucleotide. In certain such embodiments, the region of full complementarity is from 6 to 14 nucleobases in length.

Certain Antisense Activities and Mechanisms

In certain antisense activities, hybridization of an antisense compound results in recruitment of a protein that cleaves of the target nucleic acid. For example, certain antisense compounds result in RNase H mediated cleavage of target nucleic acid. RNase H is a cellular endonuclease that cleaves the RNA strand of an RNA:DNA duplex. The “DNA” in such an RNA:DNA duplex, need not be unmodified DNA. In certain embodiments, the invention provides antisense compounds that are sufficiently “DNA-like” to elicit RNase H activity. Such DNA-like antisense compounds include, but are not limited to gapmers having unmodified deoxyfuronose sugar moieties in the nucleosides of the gap and modified sugar moieties in the nucleosides of the wings.

Antisense activities may be observed directly or indirectly. In certain embodiments, observation or detection of an antisense activity involves observation or detection of a change in an amount of a target nucleic acid or protein encoded by such target nucleic acid; a change in the ratio of splice variants of a nucleic acid or protein; and/or a phenotypic change in a cell or animal.

In certain embodiments, compounds comprising oligonucleotides having a gapmer motif described herein have desirable properties compared to non-gapmer oligonucleotides or to gapmers having other motifs. In certain circumstances, it is desirable to identify motifs resulting in a favorable combination of potent antisense activity and relatively low toxicity. In certain embodiments, compounds of the present invention have a favorable therapeutic index (measure of potency divided by measure of toxicity).

Certain Target Nucleic Acids

In certain embodiments, antisense compounds comprise or consist of an oligonucleotide comprising a region that is complementary to a target nucleic acid. In certain embodiments, the target nucleic acid is an endogenous RNA molecule. In certain embodiments, the target nucleic acid is a non-coding RNA. In certain such embodiments, the target non-coding RNA is selected from: a long-non-coding RNA, a short non-coding RNA, an intronic RNA molecule, a snoRNA, a scaRNA, a microRNA (including pre-microRNA and mature microRNA), a ribosomal RNA, and promoter directed RNA. In certain embodiments, the target nucleic acid encodes a protein. In certain such embodiments, the target nucleic acid is selected from: an mRNA and a pre-mRNA, including intronic, exonic and untranslated regions. In certain embodiments, oligomeric compounds are at least partially complementary to more than one target nucleic acid. For example, antisense compounds of the present invention may mimic microRNAs, which typically bind to multiple targets.

In certain embodiments, the target nucleic acid is a nucleic acid other than a mature mRNA. In certain embodiments, the target nucleic acid is a nucleic acid other than a mature mRNA or a microRNA. In certain embodiments, the target nucleic acid is a non-coding RNA other than a microRNA. In certain embodiments, the target nucleic acid is a non-coding RNA other than a microRNA or an intronic region of a pre-mRNA. In certain embodiments, the target nucleic acid is a long non-coding RNA. In certain embodiments, the target RNA is an mRNA. In certain embodiments, the target nucleic acid is a pre-mRNA. In certain such embodiments, the target region is entirely within an intron. In certain embodiments, the target region spans an intron/exon junction. In certain embodiments, the target region is at least 50% within an intron. In certain embodiments, the target nucleic acid is selected from among non-coding RNA, including exonic regions of pre-mRNA. In certain embodiments, the target nucleic acid is a ribosomal RNA (rRNA). In certain embodiments, the target nucleic acid is a non-coding RNA associated with splicing of other pre-mRNAs. In certain embodiments, the target nucleic acid is a nuclear-retained non-coding RNA.

In certain embodiments, antisense compounds described herein are complementary to a target nucleic acid comprising a single-nucleotide polymorphism. In certain such embodiments, the antisense compound is capable of modulating expression of one allele of the single-nucleotide polymorphism-containing-target nucleic acid to a greater or lesser extent than it modulates another allele. In certain embodiments an antisense compound hybridizes to a single-nucleotide polymorphism-containing-target nucleic acid at the single-nucleotide polymorphism site. In certain embodiments an antisense compound hybridizes to a single-nucleotide polymorphism-containing-target nucleic acid near the single-nucleotide polymorphism site. In certain embodiments, the target nucleic acid is a Huntingtin gene transcript. In certain embodiments, the target nucleic acid is a single-nucleotide polymorphism-containing-target nucleic acid other than a Huntingtin gene transcript. In certain embodiments, the target nucleic acid is any nucleic acid other than a Huntingtin gene transcript.

Certain Pharmaceutical Compositions

In certain embodiments, the present invention provides pharmaceutical compositions comprising one or more antisense compound. In certain embodiments, such pharmaceutical composition comprises a suitable pharmaceutically acceptable diluent or carrier. In certain embodiments, a pharmaceutical composition comprises a sterile saline solution and one or more antisense compound. In certain embodiments, such pharmaceutical composition consists of a sterile saline solution and one or more antisense compound. In certain embodiments, the sterile saline is pharmaceutical grade saline. In certain embodiments, a pharmaceutical composition comprises one or more antisense compound and sterile water. In certain embodiments, a pharmaceutical composition consists of one or more antisense compound and sterile water. In certain embodiments, the sterile saline is pharmaceutical grade water. In certain embodiments, a pharmaceutical composition comprises one or more antisense compound and phosphate-buffered saline (PBS). In certain embodiments, a pharmaceutical composition consists of one or more antisense compound and sterile phosphate-buffered saline (PBS). In certain embodiments, the sterile saline is pharmaceutical grade PBS.

In certain embodiments, antisense compounds may be admixed with pharmaceutically acceptable active and/or inert substances for the preparation of pharmaceutical compositions or formulations. Compositions and methods for the formulation of pharmaceutical compositions depend on a number of criteria, including, but not limited to, route of administration, extent of disease, or dose to be administered.

Pharmaceutical compositions comprising antisense compounds encompass any pharmaceutically acceptable salts, esters, or salts of such esters. In certain embodiments, pharmaceutical compositions comprising antisense compounds comprise one or more oligonucleotide which, upon administration to an animal, including a human, is capable of providing (directly or indirectly) the biologically active metabolite or residue thereof. Accordingly, for example, the disclosure is also drawn to pharmaceutically acceptable salts of antisense compounds, prodrugs, pharmaceutically acceptable salts of such prodrugs, and other bioequivalents. Suitable pharmaceutically acceptable salts include, but are not limited to, sodium and potassium salts.

A prodrug can include the incorporation of additional nucleosides at one or both ends of an oligomeric compound which are cleaved by endogenous nucleases within the body, to form the active antisense oligomeric compound.

Lipid moieties have been used in nucleic acid therapies in a variety of methods. In certain such methods, the nucleic acid is introduced into preformed liposomes or lipoplexes made of mixtures of cationic lipids and neutral lipids. In certain methods, DNA complexes with mono- or poly-cationic lipids are formed without the presence of a neutral lipid. In certain embodiments, a lipid moiety is selected to increase distribution of a pharmaceutical agent to a particular cell or tissue. In certain embodiments, a lipid moiety is selected to increase distribution of a pharmaceutical agent to fat tissue. In certain embodiments, a lipid moiety is selected to increase distribution of a pharmaceutical agent to muscle tissue.

In certain embodiments, pharmaceutical compositions provided herein comprise one or more modified oligonucleotides and one or more excipients. In certain such embodiments, excipients are selected from water, salt solutions, alcohol, polyethylene glycols, gelatin, lactose, amylase, magnesium stearate, talc, silicic acid, viscous paraffin, hydroxymethylcellulose and polyvinylpyrrolidone.

In certain embodiments, a pharmaceutical composition provided herein comprises a delivery system. Examples of delivery systems include, but are not limited to, liposomes and emulsions. Certain delivery systems are useful for preparing certain pharmaceutical compositions including those comprising hydrophobic compounds. In certain embodiments, certain organic solvents such as dimethylsulfoxide are used.

In certain embodiments, a pharmaceutical composition provided herein comprises one or more tissue-specific delivery molecules designed to deliver the one or more pharmaceutical agents of the present invention to specific tissues or cell types. For example, in certain embodiments, pharmaceutical compositions include liposomes coated with a tissue-specific antibody.

In certain embodiments, a pharmaceutical composition provided herein comprises a co-solvent system. Certain of such co-solvent systems comprise, for example, benzyl alcohol, a nonpolar surfactant, a water-miscible organic polymer, and an aqueous phase. In certain embodiments, such co-solvent systems are used for hydrophobic compounds. A non-limiting example of such a co-solvent system is the VPD co-solvent system, which is a solution of absolute ethanol comprising 3% w/v benzyl alcohol, 8% w/v of the nonpolar surfactant Polysorbate 80™ and 65% w/v polyethylene glycol 300. The proportions of such co-solvent systems may be varied considerably without significantly altering their solubility and toxicity characteristics. Furthermore, the identity of co-solvent components may be varied: for example, other surfactants may be used instead of Polysorbate 80™; the fraction size of polyethylene glycol may be varied; other biocompatible polymers may replace polyethylene glycol, e.g., polyvinyl pyrrolidone; and other sugars or polysaccharides may substitute for dextrose.

In certain embodiments, a pharmaceutical composition provided herein is prepared for oral administration. In certain embodiments, pharmaceutical compositions are prepared for buccal administration.

In certain embodiments, a pharmaceutical composition is prepared for administration by injection (e.g., intravenous, subcutaneous, intramuscular, etc.). In certain of such embodiments, a pharmaceutical composition comprises a carrier and is formulated in aqueous solution, such as water or physiologically compatible buffers such as Hanks's solution, Ringer's solution, or physiological saline buffer. In certain embodiments, other ingredients are included (e.g., ingredients that aid in solubility or serve as preservatives). In certain embodiments, injectable suspensions are prepared using appropriate liquid carriers, suspending agents and the like. Certain pharmaceutical compositions for injection are presented in unit dosage form, e.g., in ampoules or in multi-dose containers. Certain pharmaceutical compositions for injection are suspensions, solutions or emulsions in oily or aqueous vehicles, and may contain formulatory agents such as suspending, stabilizing and/or dispersing agents. Certain solvents suitable for use in pharmaceutical compositions for

injection include, but are not limited to, lipophilic solvents and fatty oils, such as sesame oil, synthetic fatty acid esters, such as ethyl oleate or triglycerides, and liposomes. Aqueous injection suspensions may contain substances that increase the viscosity of the suspension, such as sodium carboxymethyl cellulose, sorbitol, or dextran. Optionally, such suspensions may also contain suitable stabilizers or agents that increase the solubility of the pharmaceutical agents to allow for the preparation of highly concentrated solutions.

In certain embodiments, a pharmaceutical composition is prepared for transmucosal administration. In certain of such embodiments penetrants appropriate to the barrier to be permeated are used in the formulation. Such penetrants are generally known in the art.

In certain embodiments, a pharmaceutical composition provided herein comprises an oligonucleotide in a therapeutically effective amount. In certain embodiments, the therapeutically effective amount is sufficient to prevent, alleviate or ameliorate symptoms of a disease or to prolong the survival of the subject being treated. Determination of a therapeutically effective amount is well within the capability of those skilled in the art.

In certain embodiments, one or more modified oligonucleotide provided herein is formulated as a prodrug. In certain embodiments, upon *in vivo* administration, a prodrug is chemically converted to the biologically, pharmaceutically or therapeutically more active form of an oligonucleotide. In certain embodiments, prodrugs are useful because they are easier to administer than the corresponding active form. For example, in certain instances, a prodrug may be more bioavailable (e.g., through oral administration) than is the corresponding active form. In certain instances, a prodrug may have improved solubility compared to the corresponding active form. In certain embodiments, prodrugs are less water soluble than the corresponding active form. In certain instances, such prodrugs possess superior transmittal across cell membranes, where water solubility is detrimental to mobility. In certain embodiments, a prodrug is an ester. In certain such embodiments, the ester is metabolically hydrolyzed to carboxylic acid upon administration. In certain instances the carboxylic acid containing compound is the corresponding active form. In certain embodiments, a prodrug comprises a short peptide (polyaminoacid) bound to an acid group. In certain of such embodiments, the peptide is cleaved upon administration to form the corresponding active form.

In certain embodiments, the present invention provides compositions and methods for reducing the amount or activity of a target nucleic acid in a cell. In certain embodiments, the cell is in an animal. In certain embodiments, the animal is a mammal. In certain embodiments, the animal is a rodent. In certain embodiments, the animal is a primate. In certain embodiments, the animal is a non-human primate. In certain embodiments, the animal is a human.

In certain embodiments, the present invention provides methods of administering a pharmaceutical composition comprising an oligomeric compound of the present invention to an animal. Suitable administration routes include, but are not limited to, oral, rectal, transmucosal, intestinal, enteral, topical, suppository, through inhalation, intrathecal, intracerebroventricular, intraperitoneal, intranasal, intraocular, intratumoral, and parenteral (e.g., intravenous, intramuscular, intramedullary, and subcutaneous). In certain

embodiments, pharmaceutical intrathecal are administered to achieve local rather than systemic exposures.

Nonlimiting disclosure and incorporation by reference

While certain compounds, compositions and methods described herein have been described with specificity in accordance with certain embodiments, the following examples serve only to illustrate the compounds described herein and are not intended to limit the same. Each of the references, GenBank accession numbers, and the like recited in the present application is incorporated herein by reference in its entirety.

Although the sequence listing accompanying this filing identifies each sequence as either “RNA” or “DNA” as required, in reality, those sequences may be modified with any combination of chemical modifications. One of skill in the art will readily appreciate that such designation as “RNA” or “DNA” to describe modified oligonucleotides is, in certain instances, arbitrary. For example, an oligonucleotide comprising a nucleoside comprising a 2'-OH sugar moiety and a thymine base could be described as a DNA having a modified sugar (2'-OH for the natural 2'-H of DNA) or as an RNA having a modified base (thymine (methylated uracil) for natural uracil of RNA).

Accordingly, nucleic acid sequences provided herein, including, but not limited to those in the sequence listing, are intended to encompass nucleic acids containing any combination of natural or modified RNA and/or DNA, including, but not limited to such nucleic acids having modified nucleobases. By way of further example and without limitation, an oligomeric compound having the nucleobase sequence “ATCGATCG” encompasses any oligomeric compounds having such nucleobase sequence, whether modified or unmodified, including, but not limited to, such compounds comprising RNA bases, such as those having sequence “AUCGAUCG” and those having some DNA bases and some RNA bases such as “AUCGATCG” and oligomeric compounds having other modified or naturally occurring bases, such as “AT^{me}CGAUCG,” wherein ^{me}C indicates a cytosine base comprising a methyl group at the 5-position.

EXAMPLES

The following examples illustrate certain embodiments of the present invention and are not limiting. Moreover, where specific embodiments are provided, the inventors have contemplated generic application of those specific embodiments. For example, disclosure of an oligonucleotide having a particular motif provides reasonable support for additional oligonucleotides having the same or similar motif. And, for example, where a particular high-affinity modification appears at a particular position, other high-affinity modifications at the same position are considered suitable, unless otherwise indicated.

Where nucleobase sequences are not provided, to allow assessment of the relative effects of nucleobase sequence and chemical modification, throughout the examples, oligomeric compounds are assigned a “Sequence Code.” Oligomeric compounds having the same Sequence Code have the same

nucleobase sequence. Oligomeric compounds having different Sequence Codes have different nucleobase sequences.

Example 1: Dose-dependent inhibition of chimeric antisense oligonucleotides targeting PTEN

A series of modified oligonucleotides were designed based on the parent gapmer, ISIS 482050, wherein the central gap region contains ten 2'-deoxynucleosides. These modified oligonucleotides were designed by having the central gap region shortened to nine, eight or seven 2'-deoxynucleosides and by introducing 2'-O-methoxyethyl (MOE) modifications at one or both wing regions. The newly designed oligonucleotides were evaluated for their effect in reducing PTEN mRNA levels *in vitro*.

The gapmers and their motifs are described in Table 60. The internucleoside linkages throughout each gapmer are phosphorothioate linkages (P=S). Nucleosides followed by a subscript "d" indicate 2'-deoxynucleosides. Nucleosides followed by a subscript "e" indicate 2'-O-methoxyethyl (MOE) nucleosides. Nucleosides followed by a subscript "k" indicate constrained ethyl (cEt) nucleosides. "N" indicates modified or naturally occurring nucleobases (A, T, C, G, U, or 5-methyl C).

The newly designed gapmers were tested *in vitro*. Mouse primary hepatocytes were plated at a density of 20,000 cells per well and transfected using electroporation with 0.6 μ M, 3.0 μ M and 15 μ M concentrations of antisense oligonucleotides. After a treatment period of approximately 24 hours, RNA was isolated from the cells and PTEN mRNA levels were measured by quantitative real-time PCR. Mouse PTEN primer probe set RTS186 was used to measure mRNA levels. PTEN mRNA levels were adjusted according to total RNA content, as measured by RIBOGREEN[®]. The results in Table 12 are presented as PTEN mRNA expression relative to untreated control cells (% UTC).

The parent gapmer, ISIS 482050 was included in the study as a bench mark oligonucleotide against which the activity of the newly designed gapmers targeting PTEN could be compared.

Table 11
Chimeric antisense oligonucleotides targeting PTEN

ISIS NO.	Sequence (5' to 3')	Motif	Gap chemistry	Wing chemistry		SEQ ID NO
				5'	3'	
482050	A _k T _k ^m C _k A _d T _d G _d G _d ^m C _d T _d G _d ^m C _d A _d G _d ^m C _k T _k T _k	3-10-3	Full deoxy	kkk	kkk	23
508033	A _k T _k ^m C _k A _d T _d G _d G _d ^m C _d T _d G _d ^m C _d A _d G _d ^m C _e T _e T _e	3-10-3	Full deoxy	kkk	eee	23
573351	A _e T _k ^m C _k A _d T _d G _d G _d ^m C _d T _d G _d ^m C _d A _d G _d ^m C _k T _k T _e	3-10-3	Full deoxy	ekk	kke	23
573352	A _e T _e ^m C _k A _k T _d G _d G _d ^m C _d T _d G _d ^m C _d A _d G _d ^m C _k T _k T _e	4-9-3	Full deoxy	eekk	kke	23
573353	A _e T _e ^m C _e A _k T _k G _d G _d ^m C _d T _d G _d ^m C _d A _d G _d ^m C _k T _k T _e	5-8-3	Full deoxy	eeekk	kke	23
573355	A _e T _k ^m C _k A _d T _d G _d G _d ^m C _d T _d G _d ^m C _d A _d G _d ^m C _k T _k T _e	3-9-4	Full deoxy	ekk	kkee	23

	${}^dA{}_dG{}_k{}^mC{}_kT_eT_e$					
573356	$A_eT_k{}^mC{}_kA_dT_dG_dG_d{}^mC_dT_dG_d{}^mC$ ${}^dA{}_dG{}_k{}^mC{}_kT_eT_e$	3-8-5	Full deoxy	ekk	kkeee	23
573357	$A_kT_k{}^mC{}_kA_dT_dG_dG_d{}^mC_dT_dG_d{}^mC$ ${}_kA{}_kG_e{}^mC{}_eT_eT_e$	3-7-6	Full deoxy	ekk	kkeeee	23
573358	$A_eT_e{}^mC{}_kA_kT_dG_dG_d{}^mC_dT_dG_d{}^mC$ ${}^dA{}_dG{}_k{}^mC{}_kT_eT_e$	4-8-4	Full deoxy	eekk	kkee	23
573359	$A_eT_e{}^mC{}_eA_kT_kG_dG_d{}^mC_dT_dG_d{}^mC$ ${}^dA{}_dG{}_k{}^mC{}_kT_eT_e$	5-7-4	Full deoxy	eeekk	kkee	23
573360	$A_eT_e{}^mC{}_kA_kT_dG_dG_d{}^mC_dT_dG_d{}^mC$ ${}^dA{}_dG{}_k{}^mC{}_kT_eT_e$	4-7-5	Full deoxy	eekk	kkeee	23

e = 2'-MOE, k = cEt, d = 2'-deoxynucleoside

Table 12

Dose-response effect of chimeric antisense oligonucleotides targeting PTEN

ISIS NO.	% UTC			Motif	Gap chemistry	Wing chemistry	
	0.6 μ M	3.0 μ M	15 μ M			5'	3'
482050	45.4	23.8	8.4	3-10-3	Full deoxy	kkk	kkk
508033	52.2	28.8	7.6	3-10-3	Full deoxy	kkk	eee
573351	66.0	24.0	12.4	3-10-3	Full deoxy	ekk	kke
573352	69.0	38.1	12.5	4-9-3	Full deoxy	eekk	kke
573353	59.8	36.5	13.8	5-8-3	Full deoxy	eeekk	kke
573355	52.1	37.4	11.4	3-9-4	Full deoxy	ekk	kkee
573356	52.9	46.4	15.4	3-8-5	Full deoxy	ekk	kkeee
573357	82.4	81.8	52.5	3-7-6	Full deoxy	ekk	kkeeee
573358	67.4	46.7	14.5	4-8-4	Full deoxy	eekk	kkee
573359	70.5	49.8	31.6	5-7-4	Full deoxy	eeekk	kkee
573360	62.2	50.8	17.6	4-7-5	Full deoxy	eekk	kkeee
Saline = 100							

e = 2'-MOE, k = cEt, d = 2'-deoxynucleoside

Example 2: Dose-dependent inhibition of chimeric antisense oligonucleotides targeting PTEN

Additional chimeric oligonucleotides were designed based on the parent gapmer, ISIS 482050, wherein the central gap region contains ten 2'-deoxynucleosides. These modified oligonucleotides were designed by having the central gap region shortened to eight 2'-deoxynucleosides and by introducing one or more 2'-O-methoxyethyl (MOE) modification(s) at the 3' wing region. The modified oligonucleotides designed by microwalk were evaluated for their effects in reducing PTEN mRNA levels *in vitro*.

The gapmers and their motifs are described in Table 13. The internucleoside linkages throughout each gapmer are phosphorothioate linkages (P=S). Nucleosides followed by a subscript "d" indicate 2'-deoxynucleoside. Nucleosides followed by a subscript "e" indicate 2'-O-methoxyethyl (MOE) nucleosides. Nucleosides followed by a subscript "k" indicate constrained ethyl (cEt) nucleosides. ^mC indicates a 5-methyl nucleoside.

The newly designed gapmers were tested *in vitro*. Mouse primary hepatocytes were plated at a density of 20,000 cells per well and transfected using electroporation with 0.6 μ M, 3.0 μ M and 15 μ M concentrations of antisense oligonucleotides. After a treatment period of approximately 24 hours, RNA was isolated from the cells and PTEN mRNA levels were measured by quantitative real-time PCR. Mouse PTEN primer probe set RTS186 was used to measure mRNA levels. PTEN mRNA levels were adjusted according to total RNA content, as measured by RIBOGREEN[®]. The results in Table 14 are presented as PTEN mRNA expression relative to untreated control cells (% UTC).

The parent gapmer, ISIS 482050 was included in the study as a bench mark oligonucleotide against which the activity of the newly designed gapmers targeting PTEN could be compared.

Table 13
Chimeric antisense oligonucleotides designed by microwalk targeting PTEN

ISIS NO.	Sequence (5' to 3')	Motif	Gap chemistry	Wing chemistry		SEQ ID NO.
				5'	3'	
482050	A _k T _k ^m C _k A _d T _d G _d G _d ^m C _d T _d G _d ^m C _d A _d G _d ^m C _k T _k T _k	3-10-3	Full deoxy	kkk	kkk	24
573797	T _k G _k G _k ^m C _d T _d G _d ^m C _d A _d G _d ^m C _d T _d T _k ^m C _e ^m C _e G _e A _e	3-8-5	Full deoxy	kkk	keeee	25
573798	A _k T _k G _k G _d ^m C _d T _d G _d ^m C _d A _d G _d ^m C _d T _d T _e ^m C _e ^m C _e G _e	3-8-5	Full deoxy	kkk	keeee	26
573799	^m C _k A _k T _k G _d G _d ^m C _d T _d G _d ^m C _d A _d G _d ^m C _k T _e ^m C _e ^m C _e	3-8-5	Full deoxy	kkk	keeee	27
573800	T _k ^m C _k A _k T _d G _d G _d ^m C _d T _d G _d ^m C _d A _d G _k ^m C _e T _e ^m C _e	3-8-5	Full deoxy	kkk	keeee	28
573801	A _k T _k ^m C _k A _d T _d G _d G _d ^m C _d T _d G _d ^m C _d A _d G _e ^m C _e T _e T _e	3-8-5	Full deoxy	kkk	keeee	24
573802	^m C _k A _k T _k ^m C _d A _d T _d G _d G _d ^m C _d T _d G _d ^m C _k A _e G _e ^m C _e T _e	3-8-5	Full deoxy	kkk	keeee	29
573803	^m C _k ^m C _k A _k T _d ^m C _d A _d T _d G _d G _d ^m C _d T _d G _k ^m C _e A _e G _e ^m C _e	3-8-5	Full deoxy	kkk	keeee	30
573804	T _k ^m C _k ^m C _k A _d T _d ^m C _d A _d T _d G _d G _d ^m C _d T _k G _e ^m C _e A _e G _e	3-8-5	Full deoxy	kkk	keeee	31
573805	T _k T _k ^m C _k ^m C _d A _d T _d ^m C _d A _d T _d G _d G _d ^m C _k T _e G _e ^m C _e A _e	3-8-5	Full deoxy	kkk	keeee	32

e = 2'-MOE, k = cEt, d = 2'-deoxynucleoside

Table 14

Dose-dependent inhibition of chimeric antisense oligonucleotides designed by microwalk targeting PTEN

ISIS NO.	% UTC			Motif	Gap chemistry	Wing chemistry	
	0.6 μ M	3.0 μ M	15 μ M			5'	3'
482050	45.4	23.8	8.4	3-10-3	Full deoxy	kkk	kkk
573797	56.8	55.4	13.1	3-8-5	Full deoxy	kkk	keeee
573798	50.9	33.5	9.6	3-8-5	Full deoxy	kkk	keeee
573799	62.6	27.7	10.3	3-8-5	Full deoxy	kkk	keeee

573800	68.6	38.9	12.0	3-8-5	Full deoxy	kkk	keeee
573801	54.6	46.3	11.8	3-8-5	Full deoxy	kkk	keeee
573802	60.7	40.4	13.0	3-8-5	Full deoxy	kkk	keeee
573803	47.0	29.8	8.5	3-8-5	Full deoxy	kkk	keeee
573804	62.5	34.1	11.3	3-8-5	Full deoxy	kkk	keeee
573805	70.3	31.6	15.2	3-8-5	Full deoxy	kkk	keeee
Saline = 100							

e = 2'-MOE, k = cEt, d= 2'-deoxynucleoside

Example 3: Antisense inhibition of Target-Z mRNA in HepG2 cells

Antisense oligonucleotides were designed targeting a Target-Z nucleic acid and were tested for their effects on Target-Z mRNA in vitro. The antisense oligonucleotides were tested in a series of experiments that had similar culture conditions. The results for each experiment are presented in separate tables shown below. ISIS 146786, 509934, ISIS 509959, and ISIS 510100, were also included in these studies for comparison. Cultured HepG2 cells at a density of 28,000 cells per well were transfected using LipofectAMINE2000® with 70 nM antisense oligonucleotide. After a treatment period of approximately 24 hours, RNA was isolated from the cells and Target-Z mRNA levels were measured by quantitative real-time PCR. Viral primer probe set RTS3370 (forward sequence CTTGGTCATGGGCCATCAG, designated herein as SEQ ID NO: 33; reverse sequence CGGCTAGGAGTTCGCAGTA, designated herein as SEQ ID NO: 34; probe sequence TGC GTGGAACCTTTTCGGCTCC, designated herein as SEQ ID NO: 35) was used to measure mRNA levels. Levels were also measured using primer probe set RTS3371 (forward sequence CCAAACCTTCGGACGAAA, designated herein as SEQ ID NO: 36; reverse sequence TGAGGCCCACTCCCATAGG, designated herein as SEQ ID NO: 37; probe sequence CCCATCATCCTGGGCTTTCGAAAAT, designated herein as SEQ ID NO: 38). Target-Z mRNA levels were adjusted according to total RNA content, as measured by RIBOGREEN®. Results are presented as percent inhibition of Target-Z, relative to untreated control cells.

The newly designed chimeric antisense oligonucleotides and their motifs are described in Tables 15-20. The gapmers are 16 nucleotides in length, wherein the central gap region comprises ten 2'-deoxynucleosides. Nucleosides followed by 'k' indicate constrained ethyl (cEt) nucleosides. Nucleosides followed by "e" indicate 2'-O-methoxyethyl (2'-MOE) nucleosides. The internucleoside linkages throughout each gapmer are phosphorothioate (P=S) linkages. All cytosine residues throughout each oligonucleotide are 5-methylcytosines.

Each gapmer listed in Tables 15-20 is targeted to the viral genomic sequence, designated herein as Target-Z. The activity of the newly designed oligonucleotides was compared with ISIS 146786, ISIS 509934, ISIS 509959, and ISIS 510100.

Table 15

Inhibition of viral Target-Z mRNA levels by chimeric antisense oligonucleotides measured with RTS3370

ISIS No	Motif	% inhibition
509934	eeee-d(10)- eeee	30
552787	ekk-d(10)-kke	57
552788	ekk-d(10)-kke	60
552789	ekk-d(10)-kke	67
552790	ekk-d(10)-kke	67
552791	ekk-d(10)-kke	65
552792	ekk-d(10)-kke	44
552793	ekkd(10)kke	0
552794	ekk-d(10)-kke	54
552795	ekk-d(10)-kke	55
552796	ekk-d(10)-kke	62
552797	ekk-d(10)-kke	59
552798	ekk-d(10)-kke	59
552799	ekk-d(10)-kke	58
552800	ekk-d(10)-kke	62
552801	ekk-d(10)-kke	65
552802	ekk-d(10)-kke	53
552803	ekk-d(10)-kke	67
552804	ekk-d(10)-kke	75
552805	ekk-d(10)-kke	72
552806	ekk-d(10)-kke	64
552807	ekk-d(10)-kke	68
552808	ekk-d(10)-kke	65
552809	ekk-d(10)-kke	60
552810	ekk-d(10)-kke	59
552811	ekk-d(10)-kke	64
552812	ekk-d(10)-kke	69
552813	ekk-d(10)-kke	64
552814	ekk-d(10)-kke	62
552815	ekk-d(10)-kke	61
552816	ekk-d(10)-kke	63
552817	ekk-d(10)-kke	42
552818	ekk-d(10)-kke	44
552819	ekk-d(10)-kke	56
552820	ekk-d(10)-kke	59
552821	ekk-d(10)-kke	76
552822	ekk-d(10)-kke	77
552823	ekk-d(10)-kke	73
552824	ekk-d(10)-kke	73

552825	ekk-d(10)-kke	51
552826	ekk-d(10)-kke	55
552827	ekk-d(10)-kke	67
552828	ekk-d(10)-kke	78
552829	ekk-d(10)-kke	72
552830	ekk-d(10)-kke	71
552831	ekk-d(10)-kke	69
552832	ekk-d(10)-kke	67
552833	ekk-d(10)-kke	65
552834	ekk-d(10)-kke	78
552835	ekk-d(10)-kke	70
552836	ekk-d(10)-kke	64
552837	ekk-d(10)-kke	65
552838	ekk-d(10)-kke	64
552839	ekk-d(10)-kke	60
552840	ekk-d(10)-kke	35
552841	ekk-d(10)-kke	62
552842	ekk-d(10)-kke	67
552843	ekk-d(10)-kke	77
552844	ekk-d(10)-kke	81
552845	ekk-d(10)-kke	63
552846	ekk-d(10)-kke	79
552847	ekk-d(10)-kke	47
552848	ekk-d(10)-kke	69
552849	ekk-d(10)-kke	59
552850	ekk-d(10)-kke	83
552851	ekk-d(10)-kke	90
552852	ekk-d(10)-kke	89
552853	ekk-d(10)-kke	83
552854	ekk-d(10)-kke	80
552855	ekk-d(10)-kke	75
552856	ekk-d(10)-kke	69
552857	ekk-d(10)-kke	68
552858	ekk-d(10)-kke	79
552859	ekk-d(10)-kke	79
552860	ekk-d(10)-kke	71
552861	ekk-d(10)-kke	68
552862	ekk-d(10)-kke	65
552863	ekk-d(10)-kke	70
552864	ekk-d(10)-kke	71

e = 2'-MOE, k = cEt, d = 2'-deoxynucleoside

Table 16

Inhibition of viral Target-Z mRNA levels by chimeric antisense oligonucleotides measured with RTS3371

ISIS No	Motif	% inhibition
552787	ekk-d(10)-kke	53
552788	ekk-d(10)-kke	45
552789	ekk-d(10)-kke	75
552790	ekk-d(10)-kke	68
552791	ekk-d(10)-kke	51
552792	ekk-d(10)-kke	38
552793	ekk-d(10)-kke	0
552794	ekk-d(10)-kke	44
552795	ekk-d(10)-kke	56
552796	ekk-d(10)-kke	45
552797	ekk-d(10)-kke	46
552798	ekk-d(10)-kke	53
552799	ekk-d(10)-kke	48
552800	ekk-d(10)-kke	54
552801	ekk-d(10)-kke	63
552802	ekk-d(10)-kke	49
552803	ekk-d(10)-kke	71
552804	ekk-d(10)-kke	64
552805	ekk-d(10)-kke	70
552806	ekk-d(10)-kke	67
552807	ekk-d(10)-kke	61
552808	ekk-d(10)-kke	83
552809	ekk-d(10)-kke	59
552810	ekk-d(10)-kke	56
552811	ekk-d(10)-kke	62
552812	ekk-d(10)-kke	66
552813	ekk-d(10)-kke	63
552814	ekk-d(10)-kke	65
552815	ekk-d(10)-kke	63
552816	ekk-d(10)-kke	88
552817	ekk-d(10)-kke	94
552818	ekk-d(10)-kke	82
552819	ekk-d(10)-kke	80
552820	ekk-d(10)-kke	84
552821	ekk-d(10)-kke	71
552822	ekk-d(10)-kke	85
552823	ekk-d(10)-kke	71
552824	ekk-d(10)-kke	81
552825	ekk-d(10)-kke	51
552826	ekk-d(10)-kke	64
552827	ekk-d(10)-kke	61

552828	ekk-d(10)-kke	76
552829	ekk-d(10)-kke	61
552830	ekk-d(10)-kke	59
552831	ekk-d(10)-kke	58
552832	ekk-d(10)-kke	64
552833	ekk-d(10)-kke	75
552834	ekk-d(10)-kke	84
552835	ekk-d(10)-kke	57
552836	ekk-d(10)-kke	51
552837	ekk-d(10)-kke	53
552838	ekk-d(10)-kke	48
552839	ekk-d(10)-kke	50
552840	ekk-d(10)-kke	54
552841	ekk-d(10)-kke	61
552842	ekk-d(10)-kke	71
552843	ekk-d(10)-kke	75
552844	ekk-d(10)-kke	78
552845	ekk-d(10)-kke	52
552846	ekk-d(10)-kke	76
552847	ekk-d(10)-kke	61
552848	ekk-d(10)-kke	72
552849	ekk-d(10)-kke	87
552850	ekk-d(10)-kke	76
552851	ekk-d(10)-kke	76
552852	ekk-d(10)-kke	79
552853	ekk-d(10)-kke	82
552854	ekk-d(10)-kke	85
552855	ekk-d(10)-kke	78
552856	ekk-d(10)-kke	77
552857	ekk-d(10)-kke	75
552858	ekk-d(10)-kke	75
552859	ekk-d(10)-kke	79
552860	ekk-d(10)-kke	71
552861	ekk-d(10)-kke	74
552862	ekk-d(10)-kke	66
552863	ekk-d(10)-kke	70
552864	ekk-d(10)-kke	73

e = 2'-MOE, k = cEt, d = 2'-deoxynucleoside

Table 17

Inhibition of viral Target-Z mRNA levels by chimeric antisense oligonucleotides measured with RTS3371

ISIS No	Motif	% inhibition
146786	eeee-d(10)-eeee	60
552889	ek-d(10)-keke	59
552890	ek-d(10)-keke	56
552891	ek-d(10)-keke	67
552892	ek-d(10)-keke	65
552893	ek-d(10)-keke	68
552894	ek-d(10)-keke	71
552895	ek-d(10)-keke	51
552896	ek-d(10)-keke	51
552897	ek-d(10)-keke	43
552898	ek-d(10)-keke	43
552899	ek-d(10)-keke	55
552900	ek-d(10)-keke	34
552901	ek-d(10)-keke	42
552902	ek-d(10)-keke	60
552903	ek-d(10)-keke	76
552904	ek-d(10)-keke	74
552905	ek-d(10)-keke	66
552907	ek-d(10)-keke	69
552908	ek-d(10)-keke	63
552909	ek-d(10)-keke	70
552910	ek-d(10)-keke	72
552911	ek-d(10)-keke	72
552912	ek-d(10)-keke	67
552913	ek-d(10)-keke	74
552914	ek-d(10)-keke	75
552915	ek-d(10)-keke	58
552916	ek-d(10)-keke	74
552917	ek-d(10)-keke	76
552918	ek-d(10)-keke	75
552919	ek-d(10)-keke	55
552920	ek-d(10)-keke	49
552921	ek-d(10)-keke	45
552922	ek-d(10)-keke	83
552923	ek-d(10)-keke	83
552924	ek-d(10)-keke	0
552925	ek-d(10)-keke	85
552926	ek-d(10)-keke	50
552927	ek-d(10)-keke	76

552928	ek-d(10)-keke	78
552929	ek-d(10)-keke	75
552930	ek-d(10)-keke	78
552931	ek-d(10)-keke	74
552932	ek-d(10)-keke	86
552933	ek-d(10)-keke	82
552934	ek-d(10)-keke	74
552935	ek-d(10)-keke	76
552936	ek-d(10)-keke	81
552937	ek-d(10)-keke	80
552938	ek-d(10)-keke	78
552939	ek-d(10)-keke	75
552940	ek-d(10)-keke	63
552941	ekk-d(10)-kke	78
552942	ek-d(10)-keke	80
552865	ekk-d(10)-kke	67
552866	ekk-d(10)-kke	68
552868	ekk-d(10)-kke	55
552869	ekk-d(10)-kke	48
552870	ekk-d(10)-kke	55
552871	ekk-d(10)-kke	57
552872	ekk-d(10)-kke	70
552873	ekk-d(10)-kke	49
552874	ekk-d(10)-kke	42
552875	ekk-d(10)-kke	41
552876	ekk-d(10)-kke	50
552877	ek-d(10)-keke	39
552878	ekk-d(10)-kke	31
552879	ekk-d(10)-kke	5
552880	ekk-d(10)-kke	5
552881	ekk-d(10)-kke	10
552882	ekk-d(10)-kke	11
552883	ekk-d(10)-kke	27
552884	ekk-d(10)-kke	36
552885	ekk-d(10)-kke	12
552886	ekk-d(10)-kke	32
552888	ekk-d(10)-kke	1

e = 2'-MOE, k = cEt, d = 2'-deoxynucleoside

Table 18

Inhibition of viral Target-Z mRNA levels by chimeric antisense oligonucleotides measured with RTS3371

ISIS No	Motif	% inhibition
146786	eeee-d(10)-eeee	59
552955	eee-d(10)-kkk	60
552956	eee-d(10)-kkk	60
552957	eee-d(10)-kkk	64
552958	eee-d(10)-kkk	56
552959	eee-d(10)-kkk	59
552960	eee-d(10)-kkk	42
552961	eee-d(10)-kkk	41
552962	eee-d(10)-kkk	35
552963	eee-d(10)-kkk	19
552964	eee-d(10)-kkk	34
552965	eee-d(10)-kkk	42
552966	eee-d(10)-kkk	60
552967	eee-d(10)-kkk	38
552968	eee-d(10)-kkk	35
552969	eee-d(10)-kkk	67
552970	eee-d(10)-kkk	56
552971	eee-d(10)-kkk	69
552972	eee-d(10)-kkk	75
552973	eee-d(10)-kkk	59
552974	eee-d(10)-kkk	71
552975	eee-d(10)-kkk	56
552976	eee-d(10)-kkk	50
552977	eee-d(10)-kkk	56
552978	eee-d(10)-kkk	43
552979	eee-d(10)-kkk	71
552980	eee-d(10)-kkk	80
552981	eee-d(10)-kkk	64
552982	ek-d(10)-keke	61
552983	eee-d(10)-kkk	77
552984	eee-d(10)-kkk	65
552985	eee-d(10)-kkk	41
552986	eee-d(10)-kkk	30
552987	eee-d(10)-kkk	41
552988	eee-d(10)-kkk	74
552989	eee-d(10)-kkk	85
552990	eee-d(10)-kkk	72
552991	eee-d(10)-kkk	73
552992	eee-d(10)-kkk	60

552993	eee-d(10)-kkk	52
552994	eee-d(10)-kkk	58
552995	eee-d(10)-kkk	70
552996	eee-d(10)-kkk	74
552997	eee-d(10)-kkk	59
552998	eee-d(10)-kkk	82
552999	eee-d(10)-kkk	70
553000	eee-d(10)-kkk	67
553001	eee-d(10)-kkk	67
553002	eee-d(10)-kkk	74
553003	eee-d(10)-kkk	72
553004	eee-d(10)-kkk	73
553005	eee-d(10)-kkk	67
553006	eee-d(10)-kkk	69
553007	eee-d(10)-kkk	60
553008	eee-d(10)-kkk	71
552943	ek-d(10)-keke	77
553009	eee-d(10)-kkk	78
552944	ek-d(10)-keke	74
553010	eee-d(10)-kkk	78
552945	ek-d(10)-keke	76
553011	eee-d(10)-kkk	72
552946	ek-d(10)-keke	71
553012	eee-d(10)-kkk	74
552947	ek-d(10)-keke	54
553013	eee-d(10)-kkk	39
552948	ek-d(10)-keke	50
553014	eee-d(10)-kkk	37
552949	ek-d(10)-keke	8
553015	eee-d(10)-kkk	45
552950	ek-d(10)-keke	44
553016	eee-d(10)-kkk	47
552951	ek-d(10)-keke	60
553017	eee-d(10)-kkk	47
552952	ek-d(10)-keke	35
553018	eee-d(10)-kkk	30
552953	ek-d(10)-keke	37
553019	eee-d(10)-kkk	37
552954	ek-d(10)-keke	40
553020	eee-d(10)-kkk	24

e = 2'-MOE, k = cEt, d = 2'-deoxynucleoside

Table 19

Inhibition of viral Target-Z mRNA levels by chimeric antisense oligonucleotides measured with RTS3370

ISIS No	Motif	% inhibition
552889	ek-d(10)-keke	42
552890	ek-d(10)-keke	56
552891	ek-d(10)-keke	55
552892	ek-d(10)-keke	53
552893	ek-d(10)-keke	56
552894	ek-d(10)-keke	53
552895	ek-d(10)-keke	38
552896	ek-d(10)-keke	43
552897	ek-d(10)-keke	40
552898	ek-d(10)-keke	50
552899	ek-d(10)-keke	37
552900	ek-d(10)-keke	43
552901	ek-d(10)-keke	56
552902	ek-d(10)-keke	43
552903	ek-d(10)-keke	78
552904	ek-d(10)-keke	75
552905	ek-d(10)-keke	52
552907	ek-d(10)-keke	75
552908	ek-d(10)-keke	57
552909	ek-d(10)-keke	66
552910	ek-d(10)-keke	60
552911	ek-d(10)-keke	65
552912	ek-d(10)-keke	37
552913	ek-d(10)-keke	76
552914	ek-d(10)-keke	79
552915	ek-d(10)-keke	71
552916	ek-d(10)-keke	82
552917	ek-d(10)-keke	78
552918	ek-d(10)-keke	64
552919	ek-d(10)-keke	38
552920	ek-d(10)-keke	43
552921	ek-d(10)-keke	49
552922	ek-d(10)-keke	90
552923	ek-d(10)-keke	92
552924	ek-d(10)-keke	30
552925	ek-d(10)-keke	81
552926	ek-d(10)-keke	39
552927	ek-d(10)-keke	53
552928	ek-d(10)-keke	48

552929	ek-d(10)-keke	68
552930	ek-d(10)-keke	87
552931	ek-d(10)-keke	87
552932	ek-d(10)-keke	88
552933	ek-d(10)-keke	75
552934	ek-d(10)-keke	76
552935	ek-d(10)-keke	71
552936	ek-d(10)-keke	80
552937	ek-d(10)-keke	81
552938	ek-d(10)-keke	85
552939	ek-d(10)-keke	82
552940	ek-d(10)-keke	76
552941	ekk-d(10)-kke	72
552942	ek-d(10)-keke	85
552865	ekk-d(10)-kke	70
552866	ekk-d(10)-kke	65
552868	ekk-d(10)-kke	36
552869	ekk-d(10)-kke	23
552870	ekk-d(10)-kke	49
552871	ekk-d(10)-kke	46
552872	ekk-d(10)-kke	73
552873	ekk-d(10)-kke	41
552874	ekk-d(10)-kke	18
552875	ekk-d(10)-kke	0
552876	ekk-d(10)-kke	49
552877	ek-d(10)-keke	37
552878	ekk-d(10)-kke	28
552879	ekk-d(10)-kke	0
552880	ekk-d(10)-kke	12
552881	ekk-d(10)-kke	0
552882	ekk-d(10)-kke	0
552883	ekk-d(10)-kke	12
552884	ekk-d(10)-kke	39
552885	ekk-d(10)-kke	37
552886	ekk-d(10)-kke	15
552888	ekk-d(10)-kke	0

e = 2'-MOE, k = cEt, d = 2'-deoxynucleoside

Table 20

Inhibition of viral Target-Z mRNA levels by chimeric antisense oligonucleotides measured with RTS3370

ISIS No	Motif	% inhibition
552955	eee-d(10)-kkk	67
552956	eee-d(10)-kkk	60

552957	eee-d(10)-kkk	73
552958	eee-d(10)-kkk	63
552959	eee-d(10)-kkk	58
552960	eee-d(10)-kkk	67
552961	eee-d(10)-kkk	78
552962	eee-d(10)-kkk	29
552963	eee-d(10)-kkk	25
552964	eee-d(10)-kkk	33
552965	eee-d(10)-kkk	55
552966	eee-d(10)-kkk	71
552967	eee-d(10)-kkk	23
552968	eee-d(10)-kkk	41
552969	eee-d(10)-kkk	76
552970	eee-d(10)-kkk	44
552971	eee-d(10)-kkk	77
552972	eee-d(10)-kkk	74
552973	eee-d(10)-kkk	61
552974	eee-d(10)-kkk	73
552975	eee-d(10)-kkk	66
552976	eee-d(10)-kkk	70
552977	eee-d(10)-kkk	65
552978	eee-d(10)-kkk	40
552979	eee-d(10)-kkk	79
552980	eee-d(10)-kkk	81
552981	eee-d(10)-kkk	74
552982	ek-d(10)-keke	52
552983	eee-d(10)-kkk	78
552984	eee-d(10)-kkk	71
552985	eee-d(10)-kkk	38
552986	eee-d(10)-kkk	48
552987	eee-d(10)-kkk	54
552988	eee-d(10)-kkk	85
552989	eee-d(10)-kkk	84
552990	eee-d(10)-kkk	79
552991	eee-d(10)-kkk	53
552992	eee-d(10)-kkk	68
552993	eee-d(10)-kkk	67
552994	eee-d(10)-kkk	69
552995	eee-d(10)-kkk	62
552996	eee-d(10)-kkk	82
552997	eee-d(10)-kkk	58
552998	eee-d(10)-kkk	86
552999	eee-d(10)-kkk	63

553000	eee-d(10)-kkk	67
553001	eee-d(10)-kkk	70
553002	eee-d(10)-kkk	84
553003	eee-d(10)-kkk	83
553004	eee-d(10)-kkk	68
553005	eee-d(10)-kkk	57
553006	eee-d(10)-kkk	74
553007	eee-d(10)-kkk	62
553008	eee-d(10)-kkk	50
552943	ek-d(10)-keke	86
553009	eee-d(10)-kkk	79
552944	ek-d(10)-keke	83
553010	eee-d(10)-kkk	74
552945	ek-d(10)-keke	79
553011	eee-d(10)-kkk	60
552946	ek-d(10)-keke	68
553012	eee-d(10)-kkk	78
552947	ek-d(10)-keke	51
553013	eee-d(10)-kkk	45
552948	ek-d(10)-keke	56
553014	eee-d(10)-kkk	53
552949	ek-d(10)-keke	1
553015	eee-d(10)-kkk	55
552950	ek-d(10)-keke	52
553016	eee-d(10)-kkk	65
552951	ek-d(10)-keke	59
553017	eee-d(10)-kkk	36
552952	ek-d(10)-keke	34
553018	eee-d(10)-kkk	20
552953	ek-d(10)-keke	55
553019	eee-d(10)-kkk	34
552954	ek-d(10)-keke	51
553020	eee-d(10)-kkk	28

e = 2'-MOE, k = cEt, d = 2'-deoxynucleoside

Example 4: Dose-dependent antisense inhibition of Target-Z mRNA in HepG2 cells

Antisense oligonucleotides from the study described in Example 46 exhibiting *in vitro* inhibition of Target-Z mRNA were selected and tested at various doses in HepG2 cells. Cells were plated at a density of 28,000 cells per well and transfected using LipofectAMINE2000® with 9.26 nM, 27.78 nM, 83.33 nM, and 250.00 nM concentrations of antisense oligonucleotide, as specified in Table 21. After a treatment period of approximately 16 hours, RNA was isolated from the cells and Target-Z mRNA levels were measured by

quantitative real-time PCR. Target-Z primer probe set RTS3371 was used to measure mRNA levels. Target-Z mRNA levels were adjusted according to total RNA content, as measured by RIBOGREEN[®]. Results are presented as percent inhibition of Target-Z, relative to untreated control cells.

As illustrated in Table 21, Target-Z mRNA levels were reduced in a dose-dependent manner in antisense oligonucleotide treated cells.

Table 21
Dose-dependent antisense inhibition of human Target-Z in HepG2 cells

ISIS No	Motif	9.2593 nM	27.7778 nM	83.3333 nM	250.0 nM
146786	eeee-d(10)-eeee	10	43	74	89
552808	ekk-d(10)-kke	13	14	55	70
552816	ekk-d(10)-kke	38	73	87	92
552818	ekk-d(10)-kke	29	63	87	85
552820	ekk-d(10)-kke	58	83	90	90
552821	ekk-d(10)-kke	33	49	71	88
552822	ekk-d(10)-kke	24	55	74	88
552824	ekk-d(10)-kke	8	24	65	87
552834	ekk-d(10)-kke	11	28	68	89
552849	ekk-d(10)-kke	12	25	73	84
552851	ekk-d(10)-kke	13	42	74	89
552852	ekk-d(10)-kke	4	35	70	87
552853	ekk-d(10)-kke	19	52	86	93
552854	ekk-d(10)-kke	28	57	80	89
552916	ek-d(10)-keke	5	26	64	82
552922	ek-d(10)-keke	25	44	77	89
552923	ek-d(10)-keke	22	49	82	91
552925	ek-d(10)-keke	33	56	80	92
552930	ek-d(10)-keke	12	49	79	89
552931	ek-d(10)-keke	12	40	62	82
552932	ek-d(10)-keke	24	62	84	91
552933	ek-d(10)-keke	20	40	75	89
552936	ek-d(10)-keke	18	36	75	88
552937	ek-d(10)-keke	22	51	82	88
552938	ek-d(10)-keke	12	36	67	80
552939	ek-d(10)-keke	17	40	65	79
552942	ek-d(10)-keke	21	48	74	88
552943	ek-d(10)-keke	5	39	70	85
552944	ek-d(10)-keke	14	33	70	77
552980	eee-d(10)-kkk	15	40	69	86
552988	eee-d(10)-kkk	4	36	58	84
552989	eee-d(10)-kkk	0	50	74	81
552996	eee-d(10)-kkk	0	25	53	72

552998	eee-d(10)-kkk	17	49	79	90
553002	eee-d(10)-kkk	0	32	68	86
553003	eee-d(10)-kkk	15	42	67	88

e = 2'-MOE, k = cEt, d = 2'-deoxynucleoside

Example 5: Efficacy of antisense oligonucleotides targeting Target-Z in transgenic mice

Mice harboring a Target-Z gene fragment (Guidotti, L. G. et al., *J. Virol.* **1995**, 69, 6158-6169) were used. The mice were treated with ISIS antisense oligonucleotides selected from studies described above as illustrated in Table 22 and evaluated for their efficacy in this model.

Treatment

Groups of 10 mice each were injected subcutaneously twice a week for the first with 50 mg/kg and, subsequently, twice a week for the next 3 weeks with 25 mg/kg of ISIS 146786 or ISIS 510100. Control groups of 10 mice each were treated in a similar manner with ISIS 141923 (5-10-5 MOE gapmer with no known murine target) or ISIS 459024 (3-10-4 MOE gapmer with no known murine target). Mice were euthanized 48 hours after the last dose, and organs and serum were harvested for further analysis.

Table 22

Antisense oligonucleotides targeting Target-Z in transgenic mice

ISIS NO.	Sequence (5' to 3')	Motif	SEQ ID NO.
146786	G _{es} T _{es} G _{es} A _{es} A _{es} G _{ds} C _{ds} G _{ds} A _{ds} A _{ds} G _{ds} T _{ds} G _{ds} C _{ds} A _{ds} C _{es} A _{es} C _{es} G _{es} G _{es}	e5-d(10)-e5	39
510100	G _{es} G _{es} ^m C _{es} A _{ds} T _{ds} A _{ds} G _{ds} ^m C _{ds} A _{ds} G _{ds} ^m C _{ds} A _{ds} G _{ds} G _{es} A _{es} T _{es} G _e	eee-d(10)-eeee	40
141923	^m C _{es} ^m C _{es} T _{es} T _{es} ^m C _{es} ^m C _{ds} ^m C _{ds} T _{ds} G _{ds} A _{ds} A _{ds} G _{ds} G _{ds} T _{ds} T _{ds} ^m C _{es} ^m C _{es} T _{es} ^m C _{es} ^m C _e	e5-d(10)-e5	41
459024	^m C _{es} G _{es} G _{es} T _{ds} ^m C _{ds} ^m C _{ds} T _{ds} T _{ds} G _{ds} G _{ds} A _{ds} G _{ds} G _{ds} A _{es} T _{es} G _{es} ^m C _e	eee-d(10)-eeee	42

e = 2'-MOE (e.g. e5 = eeeee), d = 2'-deoxynucleoside

DNA and RNA Analysis

RNA was extracted from liver tissue for real-time PCR analysis of Target-Z DNA, using primer probe sets RTS3370, RTS3371, or RTS3372 (forward sequence ATCCTATCAACACTTCCGGAAACT, designated SEQ ID NO: 43; reverse sequence CGACGCGGCGATTGAG, designated SEQ ID NO: 44; probe sequence AAGAACTCCCTCGCCTCGCAGACG, designated SEQ ID NO: 45). The DNA levels were normalized to picogreen. Target-Z RNA samples were also assayed with primer probe sets RTS3370 and RTS3371 after RT-PCR analysis. The mRNA levels were normalized to RIBOGREEN®. The data is presented in Table 23. Serum DNA samples were analyzed after the study period. The data is presented in

Table 24, expressed relative to the levels measured in the control group. As shown in Tables 23 and 24, the antisense oligonucleotides achieved reduction of Target-Z DNA and RNA over the PBS control. Treatment with either control oligonucleotide did not cause any changes in RNA or DNA levels, as expected.

Table 23

Percent inhibition of Target-Z RNA and DNA in the liver of transgenic mice

ISIS No	Motif	% inhibition DNA (RTS3370)	% inhibition DNA (RTS3371)	% inhibition DNA (RTS3372)	% inhibition RNA (RTS3370)	% inhibition RNA (RTS3371)	% inhibition RNA (RTS3372)
146786	e5-d(10)-e5	97	97	95	86	85	89
510100	eee-d(10)-eeee	95	94	94	56	64	77
141923	e5-d(10)-e5	2	0	13	0	7	31
459024	eee-d(10)-eeee	19	0	8	0	0	0

e = 2'-MOE (e.g. e5 = eeeee), d = 2'-deoxynucleoside

Table 24

Percent inhibition of Target-Z DNA in the serum of transgenic mice

ISIS No	Motif	% inhibition (RTS3370)	% inhibition (RTS3371)
146786	e5-d(10)-e5	98	98
510100	eee-d(10)-eeee	99	98
141923	e5-d(10)-e5	0	0
459024	eee-d(10)-eeee	0	0

e = 2'-MOE (e.g. e5 = eeeee), d = 2'-deoxynucleoside

Example 6: Efficacy of antisense oligonucleotides targeting Target-Z in transgenic mice

Transgenic mice were treated with ISIS antisense oligonucleotides selected from studies described above and evaluated for their efficacy in this model.

Treatment

A group of 6 mice was injected subcutaneously twice a week for 4 weeks with 25 mg/kg of ISIS 146786. Groups of 6 mice each were injected subcutaneously twice a week for 4 weeks with 10 mg/kg of ISIS 552803, ISIS 552903, ISIS 552817, ISIS 552822, and ISIS 552907. One group of 10 mice was injected subcutaneously twice a week for 4 weeks with PBS. Mice were euthanized 48 hours after the last dose, and organs and plasma were harvested for further analysis.

DNA and RNA Analysis

RNA was extracted from liver tissue for real-time PCR analysis of Target-Z DNA, using primer probe set RTS3371. The DNA levels were normalized to picogreen. Target-Z RNA samples were also assayed with primer probe set RTS3371 after RT-PCR analysis. The mRNA levels were normalized to

RIBOGREEN®. The data is presented in Table 25. Serum DNA samples were analyzed after the study period. The data is presented in Table 26, expressed relative to the levels measured in the control group. As shown in Tables 25 and 26, the antisense oligonucleotides achieved reduction of Target-Z DNA and RNA over the PBS control.

Table 25

Percent inhibition of Target-Z RNA and DNA in transgenic mice

ISIS No	Motif	Dose (mg/kg/wk)	% inhibition of RNA	% inhibition of DNA
146786	e5-d(10)-e5	50	81	91
552803	ekk-d(10)-kke	20	71	95
552817	ekk-d(10)-kke	20	86	51
552822	ekk-d(10)-kke	20	90	89
552903	ek-d(10)-keke	20	56	82
552907	ek-d(10)-keke	20	41	45

e = 2'-MOE (e.g. e5 = eeeee), d = 2'-deoxynucleoside

Table 26

Serum levels of Target-Z DNA in transgenic mice, relative to control levels

ISIS No	Motif	Dose (mg/kg/wk)	Post-dose DNA levels
146786	e5-d(10)-e5	50	0.1
552803	ekk-d(10)-kke	20	0.2
552817	ekk-d(10)-kke	20	1.3
552822	ekk-d(10)-kke	20	0.0
552903	ek-d(10)-keke	20	2.9
552907	ek-d(10)-keke	20	1.0

e = 2'-MOE (e.g. e5 = eeeee), d = 2'-deoxynucleoside

Liver function

To evaluate the effect of ISIS oligonucleotides on hepatic function, plasma concentrations of ALT were measured using an automated clinical chemistry analyzer (Hitachi Olympus AU400e, Melville, NY) (Nyblom, H. et al., Alcohol & Alcoholism 39: 336-339, 2004; Tietz NW (Ed): Clinical Guide to Laboratory Tests, 3rd ed. W. B. Saunders, Philadelphia, PA, 1995). The results are presented in Table 27 expressed in IU/L. All the ISIS oligonucleotides were considered tolerable in the mice, as demonstrated by their liver transaminase profile.

Table 27
ALT levels (IU/L) of transgenic mice

	Motif	Dose (mg/kg/wk)	ALT
PBS	--	-	77
ISIS 146786	e5-d(10)-e5	50	21
ISIS 552803	ekk-d(10)-kke	20	74
ISIS 552817	ekk-d(10)-kke	20	38
ISIS 552822	ekk-d(10)-kke	20	47
ISIS 552903	ek-d(10)-keke	20	57
ISIS 552907	ek-d(10)-keke	20	28

e = 2'-MOE (e.g. e5 = eeeee), d = 2'-deoxynucleoside

Example 7: Efficacy of antisense oligonucleotides targeting Target-Z in transgenic mice

Transgenic mice were treated with ISIS antisense oligonucleotides selected from studies described above and evaluated for their efficacy in this model.

Treatment

A group of 6 mice was injected subcutaneously twice a week for 4 weeks with 25 mg/kg of ISIS 146786. Groups of 6 mice each were injected subcutaneously twice a week for 4 weeks with 10 mg/kg of ISIS 552853, ISIS 552854, ISIS 552932, and ISIS 552938. One group of 10 mice was injected subcutaneously twice a week for 4 weeks with PBS. Mice were euthanized 48 hours after the last dose, and organs and plasma were harvested for further analysis.

DNA and RNA Analysis

RNA was extracted from liver tissue for real-time PCR analysis of Target-Z DNA, using primer probe set RTS3371. The DNA levels were normalized to picogreen. Target-Z RNA samples were also assayed with primer probe set RTS3371 after RT-PCR analysis. The mRNA levels were normalized to RIBOGREEN®. As shown in Table 28, the antisense oligonucleotides achieved reduction of Target-Z DNA and RNA over the PBS control. Results are presented as percent inhibition of Target-Z mRNA or DNA, relative to control.

Table 28
Percent inhibition of Target-Z DNA and RNA in transgenic mice

	Motif	Dose (mg/kg/wk)	% inhibition (DNA)	% inhibition (RNA)
PBS	--	-		
ISIS 146786	e5-d(10)-e5	50	90	60
ISIS 552853	ekk-d(10)-kke	20	94	60
ISIS 552854	ekk-d(10)-kke	20	61	23

ISIS 552932	ekk-d(10)-kke	20	75	70
ISIS 552938	ek-d(10)-keke	20	67	56

= 2'-MOE (e.g. e5 = eeeee), d = 2'-deoxynucleoside

Example 8: Efficacy of antisense oligonucleotides targeting Target-Z in transgenic mice

Transgenic mice were treated with ISIS antisense oligonucleotides selected from studies described above and evaluated for their efficacy in this model.

Treatment

A group of 6 mice was injected subcutaneously twice a week for 4 weeks with 25 mg/kg of ISIS 146786. Groups of 6 mice each were injected subcutaneously twice a week for 4 weeks with 10 mg/kg of ISIS 552922, ISIS 552923, ISIS 552942, ISIS 552872, ISIS 552925, ISIS 552937, and ISIS 552939. One group of 10 mice was injected subcutaneously twice a week for 4 weeks with PBS. Mice were euthanized 48 hours after the last dose, and organs and plasma were harvested for further analysis.

DNA and RNA Analysis

RNA was extracted from liver tissue for real-time PCR analysis of Target-Z DNA, using primer probe set RTS3371. The DNA levels were normalized to picogreen. Target-Z RNA samples were also assayed with primer probe set RTS3371 after RT-PCR analysis. The mRNA levels were normalized to RIBOGREEN®. As shown in Table 29, the antisense oligonucleotides achieved reduction of Target-Z DNA and RNA over the PBS control. Results are presented as percent inhibition of Target-Z mRNA or DNA, relative to control.

Table 29

Percent inhibition of Target-Z DNA and RNA in transgenic mice

ISIS No	Motif	Dose (mg/kg/wk)	% inhibition (DNA)	% inhibition (RNA)
146786	e5-d(10)-e5	50	52	57
552922	ek-d(10)-keke	20	61	50
552923	ek-d(10)-keke	20	89	76
552942	ek-d(10)-keke	20	58	52
552872	ekk-d(10)-kke	20	77	46
552925	ek-d(10)-keke	20	89	65
552937	ek-d(10)-keke	20	59	35
552939	ek-d(10)-keke	20	57	19

= 2'-MOE (e.g. e5 = eeeee), d = 2'-deoxynucleoside

Example 9: Antisense inhibition of Target-Z mRNA in HepG2 cells

Antisense oligonucleotides were designed targeting a Target-Z nucleic acid and were tested for their effects on Target-Z mRNA in vitro. The antisense oligonucleotides were tested in a series of experiments that had similar culture conditions. The results for each experiment are presented in separate tables. ISIS 146786, 509934, ISIS 509959, and ISIS 510100, from the studies described above, were also included. Cultured HepG2 cells at a density of 28,000 cells per well were transfected using LipofectAMINE2000® with 70 nM antisense oligonucleotide. After a treatment period of approximately 24 hours, RNA was isolated from the cells and Target-Z mRNA levels were measured by quantitative real-time PCR. Primer probe set RTS3370 (forward sequence CTTGGTCATGGGCCATCAG, designated herein as SEQ ID NO: 33; reverse sequence CGGCTAGGAGTTCCGCAGTA, designated herein as SEQ ID NO: 34; probe sequence TGCGTGGAACCTTTTCGGCTCC, designated herein as SEQ ID NO: 35) was used to measure mRNA levels. Levels were also measured using primer probe set RTS3371 (forward sequence CCAAACCTTCGGACGGAAA, designated herein as SEQ ID NO: 36; reverse sequence TGAGGCCCACTCCCATAGG, designated herein as SEQ ID NO: 37; probe sequence CCCATCATCCTGGGCTTTCGGAAAAT, designated herein as SEQ ID NO: 38). Target-Z mRNA levels were adjusted according to total RNA content, as measured by RIBOGREEN®. Results are presented as percent inhibition of Target-Z, relative to untreated control cells.

The newly designed chimeric antisense oligonucleotides and their motifs are described in Tables 30-47. The modified oligonucleotides are 16, 17 or 20 nucleotides in length, wherein the central gap segment comprises of nine or ten 2'-deoxynucleosides and is flanked by wing segments on the 5' direction and the 3' direction comprising 2'-O-methoxyethyl (2'-MOE) modifications. The internucleoside linkages throughout each gapmer are phosphorothioate (P=S) linkages. All cytosine residues throughout each oligonucleotide are 5-methylcytosines.

Each gapmer listed in the Tables is targeted to the viral genomic sequence, designated herein as Target-Z. The activity of the newly designed oligonucleotides was compared with ISIS 146786, 509934, ISIS 509959, and ISIS 510100, the information of which have been placed at the top of each table.

Table 30

Inhibition of viral Target-Z mRNA levels by chimeric antisense oligonucleotides measured with RTS3370

ISIS No	Motif	Wing chemistry	% inhibition
146786	5-10-5	2'-MOE	50
510100	3-10-4	2'-MOE	62
552276	5-9-3	2'-MOE	42
552277	5-9-3	2'-MOE	46
552278	5-9-3	2'-MOE	31

552279	5-9-3	2'-MOE	41
552280	5-9-3	2'-MOE	5
552281	5-9-3	2'-MOE	11
552282	5-9-3	2'-MOE	20
552283	5-9-3	2'-MOE	28
552230	4-9-4	2'-MOE	57
552284	5-9-3	2'-MOE	0
552231	4-9-4	2'-MOE	29
552285	5-9-3	2'-MOE	61
552232	4-9-4	2'-MOE	35
552286	5-9-3	2'-MOE	47
552233	4-9-4	2'-MOE	38
552287	5-9-3	2'-MOE	45
552234	4-9-4	2'-MOE	0
552288	5-9-3	2'-MOE	50
552235	4-9-4	2'-MOE	0
552289	5-9-3	2'-MOE	46
552236	4-9-4	2'-MOE	45
552290	5-9-3	2'-MOE	41
552237	4-9-4	2'-MOE	44
552291	5-9-3	2'-MOE	26
552239	4-9-4	2'-MOE	62
552293	5-9-3	2'-MOE	67
552240	4-9-4	2'-MOE	61
552294	5-9-3	2'-MOE	71
552241	4-9-4	2'-MOE	55
552295	5-9-3	2'-MOE	58
552242	4-9-4	2'-MOE	60
552296	5-9-3	2'-MOE	59
552243	4-9-4	2'-MOE	57
552297	5-9-3	2'-MOE	55
552244	4-9-4	2'-MOE	33
552298	5-9-3	2'-MOE	48
552245	4-9-4	2'-MOE	48
552299	5-9-3	2'-MOE	34
552246	4-9-4	2'-MOE	81
552300	5-9-3	2'-MOE	56
552247	4-9-4	2'-MOE	87
552301	5-9-3	2'-MOE	86
552248	4-9-4	2'-MOE	72
552302	5-9-3	2'-MOE	77
552249	4-9-4	2'-MOE	56
552303	5-9-3	2'-MOE	65

552250	4-9-4	2'-MOE	52
552304	5-9-3	2'-MOE	57
552251	4-9-4	2'-MOE	43
552305	5-9-3	2'-MOE	56
552252	4-9-4	2'-MOE	62
552306	5-9-3	2'-MOE	75
552253	4-9-4	2'-MOE	82
552307	5-9-3	2'-MOE	90
552254	4-9-4	2'-MOE	74
552255	4-9-4	2'-MOE	78
552256	4-9-4	2'-MOE	65
552257	4-9-4	2'-MOE	62
552258	4-9-4	2'-MOE	72
552259	4-9-4	2'-MOE	63
552260	4-9-4	2'-MOE	58
552261	4-9-4	2'-MOE	63
552262	4-9-4	2'-MOE	50
552263	4-9-4	2'-MOE	60
552264	4-9-4	2'-MOE	52
552265	4-9-4	2'-MOE	68
552266	4-9-4	2'-MOE	62
552267	4-9-4	2'-MOE	58
552268	4-9-4	2'-MOE	62
552269	4-9-4	2'-MOE	52
552270	4-9-4	2'-MOE	54
552271	4-9-4	2'-MOE	58
552272	4-9-4	2'-MOE	40
552273	4-9-4	2'-MOE	34
552274	4-9-4	2'-MOE	34
552275	4-9-4	2'-MOE	39

Table 31

Inhibition of viral Target-Z mRNA levels by chimeric antisense oligonucleotides measured with RTS3370

ISIS No	Motif	Wing chemistry	% inhibition
146786	5-10-5	2'-MOE	49
509959	3-10-3	2'-MOE	43
510100	3-10-4	2'-MOE	54
552384	2-9-5	2'-MOE	29
552440	3-9-4	2'-MOE	58
552385	2-9-5	2'-MOE	57
552441	3-9-4	2'-MOE	42
552386	2-9-5	2'-MOE	53

552442	3-9-4	2'-MOE	53
552387	2-9-5	2'-MOE	48
552443	3-9-4	2'-MOE	59
552388	2-9-5	2'-MOE	40
552444	3-9-4	2'-MOE	51
552389	2-9-5	2'-MOE	39
552445	3-9-4	2'-MOE	60
552390	2-9-5	2'-MOE	52
552446	3-9-4	2'-MOE	54
552391	2-9-5	2'-MOE	57
552447	3-9-4	2'-MOE	54
552392	2-9-5	2'-MOE	0
552448	3-9-4	2'-MOE	58
552393	2-9-5	2'-MOE	59
552449	3-9-4	2'-MOE	60
552394	2-9-5	2'-MOE	53
552450	3-9-4	2'-MOE	53
552395	2-9-5	2'-MOE	57
552451	3-9-4	2'-MOE	39
552396	2-9-5	2'-MOE	62
552452	3-9-4	2'-MOE	57
552238	4-9-4	2'-MOE	38
552292	5-9-3	2'-MOE	48
552346	6-9-2	2'-MOE	0
552397	2-9-5	2'-MOE	63
552453	3-9-4	2'-MOE	56
552398	2-9-5	2'-MOE	61
552454	3-9-4	2'-MOE	48
552399	2-9-5	2'-MOE	52
552400	2-9-5	2'-MOE	57
552401	2-9-5	2'-MOE	52
552402	2-9-5	2'-MOE	54
552403	2-9-5	2'-MOE	74
552404	2-9-5	2'-MOE	43
552405	2-9-5	2'-MOE	15
552406	2-9-5	2'-MOE	37
552407	2-9-5	2'-MOE	37
552408	2-9-5	2'-MOE	76
552409	2-9-5	2'-MOE	76
552410	2-9-5	2'-MOE	63
552411	2-9-5	2'-MOE	70
552412	2-9-5	2'-MOE	62
552413	2-9-5	2'-MOE	56

552414	2-9-5	2'-MOE	63
552415	2-9-5	2'-MOE	52
552416	2-9-5	2'-MOE	67
552417	2-9-5	2'-MOE	50
552418	2-9-5	2'-MOE	79
552419	2-9-5	2'-MOE	70
552420	2-9-5	2'-MOE	71
552421	2-9-5	2'-MOE	69
552422	2-9-5	2'-MOE	68
552423	2-9-5	2'-MOE	65
552424	2-9-5	2'-MOE	70
552425	2-9-5	2'-MOE	51
552426	2-9-5	2'-MOE	40
552427	2-9-5	2'-MOE	35
552428	2-9-5	2'-MOE	58
552429	2-9-5	2'-MOE	46
552430	2-9-5	2'-MOE	53
552431	2-9-5	2'-MOE	51
552432	2-9-5	2'-MOE	57
552433	2-9-5	2'-MOE	54
552434	2-9-5	2'-MOE	44
552435	2-9-5	2'-MOE	46
552436	2-9-5	2'-MOE	36
552437	2-9-5	2'-MOE	27
552438	2-9-5	2'-MOE	27
552439	2-9-5	2'-MOE	13

Table 32

Inhibition of viral Target-Z mRNA levels by chimeric antisense oligonucleotides measured with RTS3370

ISIS No	Motif	Wing chemistry	% inhibition
146786	5-10-5	2'-MOE	35
509959	3-10-3	2'-MOE	52
552496	4-9-3	2'-MOE	47
552497	4-9-3	2'-MOE	57
552498	4-9-3	2'-MOE	45
552499	4-9-3	2'-MOE	52
552500	4-9-3	2'-MOE	46
552501	4-9-3	2'-MOE	44
552502	4-9-3	2'-MOE	57
552503	4-9-3	2'-MOE	52
552504	4-9-3	2'-MOE	45
552505	4-9-3	2'-MOE	56

552506	4-9-3	2'-MOE	54
552507	4-9-3	2'-MOE	34
552508	4-9-3	2'-MOE	34
552509	4-9-3	2'-MOE	48
552510	4-9-3	2'-MOE	50
552455	3-9-4	2'-MOE	66
552511	4-9-3	2'-MOE	66
552456	3-9-4	2'-MOE	64
552512	4-9-3	2'-MOE	62
552457	3-9-4	2'-MOE	14
552513	4-9-3	2'-MOE	56
552458	3-9-4	2'-MOE	59
552514	4-9-3	2'-MOE	52
552459	3-9-4	2'-MOE	69
552515	4-9-3	2'-MOE	57
552460	3-9-4	2'-MOE	0
552516	4-9-3	2'-MOE	54
552461	3-9-4	2'-MOE	20
552517	4-9-3	2'-MOE	52
552462	3-9-4	2'-MOE	46
552518	4-9-3	2'-MOE	34
552463	3-9-4	2'-MOE	48
552519	4-9-3	2'-MOE	44
552464	3-9-4	2'-MOE	81
552520	4-9-3	2'-MOE	69
552465	3-9-4	2'-MOE	84
552521	4-9-3	2'-MOE	80
552466	3-9-4	2'-MOE	75
552522	4-9-3	2'-MOE	76
552467	3-9-4	2'-MOE	65
552523	4-9-3	2'-MOE	71
552468	3-9-4	2'-MOE	53
552524	4-9-3	2'-MOE	43
552469	3-9-4	2'-MOE	51
552525	4-9-3	2'-MOE	57
552470	3-9-4	2'-MOE	46
552526	4-9-3	2'-MOE	60
552471	3-9-4	2'-MOE	54
552527	4-9-3	2'-MOE	72
552472	3-9-4	2'-MOE	78
552528	4-9-3	2'-MOE	78
552473	3-9-4	2'-MOE	67
552529	4-9-3	2'-MOE	77

552474	3-9-4	2'-MOE	79
552530	4-9-3	2'-MOE	78
552475	3-9-4	2'-MOE	74
552531	4-9-3	2'-MOE	68
552476	3-9-4	2'-MOE	52
552477	3-9-4	2'-MOE	76
552478	3-9-4	2'-MOE	70
552479	3-9-4	2'-MOE	67
552480	3-9-4	2'-MOE	68
552481	3-9-4	2'-MOE	57
552482	3-9-4	2'-MOE	51
552483	3-9-4	2'-MOE	48
552484	3-9-4	2'-MOE	58
552485	3-9-4	2'-MOE	51
552486	3-9-4	2'-MOE	55
552487	3-9-4	2'-MOE	62
552488	3-9-4	2'-MOE	51
552489	3-9-4	2'-MOE	49
552490	3-9-4	2'-MOE	51
552491	3-9-4	2'-MOE	51
552492	3-9-4	2'-MOE	38
552493	3-9-4	2'-MOE	52
552494	3-9-4	2'-MOE	17
552495	3-9-4	2'-MOE	49

Table 33

Inhibition of viral Target-Z mRNA levels by chimeric antisense oligonucleotides measured with RTS3370

ISIS No	Motif	Wing chemistry	% inhibition
146786	5-10-5	2'-MOE	47
509959	3-10-3	2'-MOE	38
552552	5-9-2	2'-MOE	33
552553	5-9-2	2'-MOE	46
552554	5-9-2	2'-MOE	54
552555	5-9-2	2'-MOE	50
552556	5-9-2	2'-MOE	46
552557	5-9-2	2'-MOE	57
552558	5-9-2	2'-MOE	55
552559	5-9-2	2'-MOE	66
552560	5-9-2	2'-MOE	44
552561	5-9-2	2'-MOE	48
552562	5-9-2	2'-MOE	52
552563	5-9-2	2'-MOE	45

552564	5-9-2	2'-MOE	41
552565	5-9-2	2'-MOE	54
552566	5-9-2	2'-MOE	56
552567	5-9-2	2'-MOE	71
552568	5-9-2	2'-MOE	64
552569	5-9-2	2'-MOE	59
552570	5-9-2	2'-MOE	60
552571	5-9-2	2'-MOE	55
552572	5-9-2	2'-MOE	60
552573	5-9-2	2'-MOE	24
552574	5-9-2	2'-MOE	34
552575	5-9-2	2'-MOE	36
552576	5-9-2	2'-MOE	67
552577	5-9-2	2'-MOE	64
552578	5-9-2	2'-MOE	75
552579	5-9-2	2'-MOE	75
552580	5-9-2	2'-MOE	59
552581	5-9-2	2'-MOE	54
552582	5-9-2	2'-MOE	61
552583	5-9-2	2'-MOE	69
552584	5-9-2	2'-MOE	74
552585	5-9-2	2'-MOE	62
552586	5-9-2	2'-MOE	79
552587	5-9-2	2'-MOE	71
552532	4-9-3	2'-MOE	48
552588	5-9-2	2'-MOE	70
552533	4-9-3	2'-MOE	43
552589	5-9-2	2'-MOE	59
552534	4-9-3	2'-MOE	62
552590	5-9-2	2'-MOE	70
552535	4-9-3	2'-MOE	55
552591	5-9-2	2'-MOE	51
552536	4-9-3	2'-MOE	3
552592	5-9-2	2'-MOE	50
552537	4-9-3	2'-MOE	14
552593	5-9-2	2'-MOE	46
552538	4-9-3	2'-MOE	52
552594	5-9-2	2'-MOE	55
552539	4-9-3	2'-MOE	47
552595	5-9-2	2'-MOE	60
552540	4-9-3	2'-MOE	60
552596	5-9-2	2'-MOE	63
552541	4-9-3	2'-MOE	60

552597	5-9-2	2'-MOE	61
552542	4-9-3	2'-MOE	64
552598	5-9-2	2'-MOE	57
552543	4-9-3	2'-MOE	46
552600	5-9-2	2'-MOE	59
552544	4-9-3	2'-MOE	53
552602	5-9-2	2'-MOE	6
552545	4-9-3	2'-MOE	33
552604	5-9-2	2'-MOE	47
552546	4-9-3	2'-MOE	42
552606	5-9-2	2'-MOE	53
552547	4-9-3	2'-MOE	51
552608	5-9-2	2'-MOE	53
552548	4-9-3	2'-MOE	52
552610	5-9-2	2'-MOE	47
552549	4-9-3	2'-MOE	38
552612	5-9-2	2'-MOE	39
552550	4-9-3	2'-MOE	19
552614	5-9-2	2'-MOE	24
552551	4-9-3	2'-MOE	24
552616	5-9-2	2'-MOE	15

Table 34

Inhibition of viral Target-Z mRNA levels by chimeric antisense oligonucleotides measured with RTS3370

ISIS No	Motif	Wing chemistry	% inhibition
146786	5-10-5	2'-MOE	51
509934	5-10-5	2'-MOE	76
552007	6-10-4	2'-MOE	61
552039	7-10-3	2'-MOE	84
552008	6-10-4	2'-MOE	48
552040	7-10-3	2'-MOE	48
552009	6-10-4	2'-MOE	77
552041	7-10-3	2'-MOE	73
552010	6-10-4	2'-MOE	63
552042	7-10-3	2'-MOE	66
552011	6-10-4	2'-MOE	52
552043	7-10-3	2'-MOE	54
552012	6-10-4	2'-MOE	73
552044	7-10-3	2'-MOE	86
552013	6-10-4	2'-MOE	73
552045	7-10-3	2'-MOE	65
552014	6-10-4	2'-MOE	76

552046	7-10-3	2'-MOE	93
552015	6-10-4	2'-MOE	70
552047	7-10-3	2'-MOE	77
552016	6-10-4	2'-MOE	61
552048	7-10-3	2'-MOE	66
552017	6-10-4	2'-MOE	73
552049	7-10-3	2'-MOE	73
552018	6-10-4	2'-MOE	98
552050	7-10-3	2'-MOE	98
552019	6-10-4	2'-MOE	98
552051	7-10-3	2'-MOE	99
551986	4-10-6	2'-MOE	92
552020	6-10-4	2'-MOE	97
552052	7-10-3	2'-MOE	98
551987	4-10-6	2'-MOE	95
552021	6-10-4	2'-MOE	97
552053	7-10-3	2'-MOE	98
551988	4-10-6	2'-MOE	50
552005	5-10-5	2'-MOE	99
552022	6-10-4	2'-MOE	99
552054	7-10-3	2'-MOE	99
551989	4-10-6	2'-MOE	96
552023	6-10-4	2'-MOE	99
552055	7-10-3	2'-MOE	98
551990	4-10-6	2'-MOE	86
552024	6-10-4	2'-MOE	89
552056	7-10-3	2'-MOE	88
551991	4-10-6	2'-MOE	0
552025	6-10-4	2'-MOE	90
552057	7-10-3	2'-MOE	92
551992	4-10-6	2'-MOE	72
552026	6-10-4	2'-MOE	88
552058	7-10-3	2'-MOE	86
551993	4-10-6	2'-MOE	82
552027	6-10-4	2'-MOE	87
552059	7-10-3	2'-MOE	88
551994	4-10-6	2'-MOE	85
552028	6-10-4	2'-MOE	83
552060	7-10-3	2'-MOE	82
551995	4-10-6	2'-MOE	84
552029	6-10-4	2'-MOE	88
552061	7-10-3	2'-MOE	85
551996	4-10-6	2'-MOE	87

552030	6-10-4	2'-MOE	88
552062	7-10-3	2'-MOE	85
551997	4-10-6	2'-MOE	83
552031	6-10-4	2'-MOE	82
551998	4-10-6	2'-MOE	85
552032	6-10-4	2'-MOE	87
551999	4-10-6	2'-MOE	82
552033	6-10-4	2'-MOE	87
552000	4-10-6	2'-MOE	83
552006	5-10-5	2'-MOE	88
552034	6-10-4	2'-MOE	89
552001	4-10-6	2'-MOE	65
552035	6-10-4	2'-MOE	60
552002	4-10-6	2'-MOE	63
552036	6-10-4	2'-MOE	65
552003	4-10-6	2'-MOE	65
552037	6-10-4	2'-MOE	58
552004	4-10-6	2'-MOE	58
552038	6-10-4	2'-MOE	70

Table 35

Inhibition of viral Target-Z mRNA levels by chimeric antisense oligonucleotides measured with RTS3370

ISIS No	Motif	Wing chemistry	% inhibition
146786	5-10-5	2'-MOE	64
510100	3-10-4	2'-MOE	62
552168	3-9-5	2'-MOE	79
552222	4-9-4	2'-MOE	79
552169	3-9-5	2'-MOE	67
552223	4-9-4	2'-MOE	40
552170	3-9-5	2'-MOE	69
552224	4-9-4	2'-MOE	64
552171	3-9-5	2'-MOE	65
552225	4-9-4	2'-MOE	69
552172	3-9-5	2'-MOE	33
552226	4-9-4	2'-MOE	48
552173	3-9-5	2'-MOE	41
552227	4-9-4	2'-MOE	32
552174	3-9-5	2'-MOE	31
552228	4-9-4	2'-MOE	42
552175	3-9-5	2'-MOE	59
552176	3-9-5	2'-MOE	68
552177	3-9-5	2'-MOE	55

552178	3-9-5	2'-MOE	66
552179	3-9-5	2'-MOE	70
552180	3-9-5	2'-MOE	66
552181	3-9-5	2'-MOE	51
552182	3-9-5	2'-MOE	69
552183	3-9-5	2'-MOE	69
552184	3-9-5	2'-MOE	43
552185	3-9-5	2'-MOE	66
552186	3-9-5	2'-MOE	54
552187	3-9-5	2'-MOE	74
552188	3-9-5	2'-MOE	78
552189	3-9-5	2'-MOE	57
552190	3-9-5	2'-MOE	39
552191	3-9-5	2'-MOE	60
552192	3-9-5	2'-MOE	85
552193	3-9-5	2'-MOE	86
552194	3-9-5	2'-MOE	68
552195	3-9-5	2'-MOE	73
552196	3-9-5	2'-MOE	60
552197	3-9-5	2'-MOE	60
552198	3-9-5	2'-MOE	61
552199	3-9-5	2'-MOE	89
552200	3-9-5	2'-MOE	85
552201	3-9-5	2'-MOE	81
552202	3-9-5	2'-MOE	76
552203	3-9-5	2'-MOE	74
552204	3-9-5	2'-MOE	71
552151	2-9-6	2'-MOE	77
552205	3-9-5	2'-MOE	78
552152	2-9-6	2'-MOE	72
552206	3-9-5	2'-MOE	77
552153	2-9-6	2'-MOE	67
552207	3-9-5	2'-MOE	81
552154	2-9-6	2'-MOE	56
552208	3-9-5	2'-MOE	70
552155	2-9-6	2'-MOE	61
552209	3-9-5	2'-MOE	63
552156	2-9-6	2'-MOE	20
552210	3-9-5	2'-MOE	75
552157	2-9-6	2'-MOE	39
552211	3-9-5	2'-MOE	75
552158	2-9-6	2'-MOE	70
552212	3-9-5	2'-MOE	67

552159	2-9-6	2'-MOE	74
552213	3-9-5	2'-MOE	70
552160	2-9-6	2'-MOE	78
552214	3-9-5	2'-MOE	79
552161	2-9-6	2'-MOE	56
552215	3-9-5	2'-MOE	61
552162	2-9-6	2'-MOE	64
552216	3-9-5	2'-MOE	62
552163	2-9-6	2'-MOE	71
552217	3-9-5	2'-MOE	58
552164	2-9-6	2'-MOE	52
552218	3-9-5	2'-MOE	56
552165	2-9-6	2'-MOE	53
552219	3-9-5	2'-MOE	33
552166	2-9-6	2'-MOE	41
552220	3-9-5	2'-MOE	53
552167	2-9-6	2'-MOE	54
552221	3-9-5	2'-MOE	31

Table 36

Inhibition of viral Target-Z mRNA levels by chimeric antisense oligonucleotides measured with RTS3370

ISIS No	Motif	Wing chemistry	% inhibition
146786	5-10-5	2'-MOE	73
509934	5-10-5	2'-MOE	76
510100	3-10-4	2'-MOE	73
552071	8-10-2	2'-MOE	79
552114	2-9-6	2'-MOE	66
552115	2-9-6	2'-MOE	70
552116	2-9-6	2'-MOE	68
552117	2-9-6	2'-MOE	70
552072	8-10-2	2'-MOE	50
552118	2-9-6	2'-MOE	66
552119	2-9-6	2'-MOE	62
552120	2-9-6	2'-MOE	35
552121	2-9-6	2'-MOE	39
552073	8-10-2	2'-MOE	80
552122	2-9-6	2'-MOE	55
552074	8-10-2	2'-MOE	73
552123	2-9-6	2'-MOE	75
552075	8-10-2	2'-MOE	78
552124	2-9-6	2'-MOE	64
552076	8-10-2	2'-MOE	70

552125	2-9-6	2'-MOE	73
552077	8-10-2	2'-MOE	83
552126	2-9-6	2'-MOE	64
552078	8-10-2	2'-MOE	80
552127	2-9-6	2'-MOE	72
552079	8-10-2	2'-MOE	86
552128	2-9-6	2'-MOE	76
552080	8-10-2	2'-MOE	83
552129	2-9-6	2'-MOE	72
552131	2-9-6	2'-MOE	61
552132	2-9-6	2'-MOE	73
552133	2-9-6	2'-MOE	75
552081	8-10-2	2'-MOE	76
552134	2-9-6	2'-MOE	58
552135	2-9-6	2'-MOE	67
552136	2-9-6	2'-MOE	65
552137	2-9-6	2'-MOE	55
552082	8-10-2	2'-MOE	98
552138	2-9-6	2'-MOE	82
552083	8-10-2	2'-MOE	99
552139	2-9-6	2'-MOE	86
552084	8-10-2	2'-MOE	99
552140	2-9-6	2'-MOE	74
552085	8-10-2	2'-MOE	100
552141	2-9-6	2'-MOE	67
552086	8-10-2	2'-MOE	100
552142	2-9-6	2'-MOE	45
552087	8-10-2	2'-MOE	100
552143	2-9-6	2'-MOE	68
552144	2-9-6	2'-MOE	78
552145	2-9-6	2'-MOE	88
552146	2-9-6	2'-MOE	81
552088	8-10-2	2'-MOE	95
552147	2-9-6	2'-MOE	88
552089	8-10-2	2'-MOE	93
552148	2-9-6	2'-MOE	79
552090	8-10-2	2'-MOE	87
552149	2-9-6	2'-MOE	81
552091	8-10-2	2'-MOE	88
552092	8-10-2	2'-MOE	90
552093	8-10-2	2'-MOE	91
552094	8-10-2	2'-MOE	88
552063	7-10-3	2'-MOE	81

552095	8-10-2	2'-MOE	89
552064	7-10-3	2'-MOE	85
552096	8-10-2	2'-MOE	92
552065	7-10-3	2'-MOE	86
552097	8-10-2	2'-MOE	93
552066	7-10-3	2'-MOE	33
552098	8-10-2	2'-MOE	88
552067	7-10-3	2'-MOE	50
552099	8-10-2	2'-MOE	70
552068	7-10-3	2'-MOE	73
552100	8-10-2	2'-MOE	70
552069	7-10-3	2'-MOE	73
552101	8-10-2	2'-MOE	76
552070	7-10-3	2'-MOE	71
552102	8-10-2	2'-MOE	64

Table 37

Inhibition of viral Target-Z mRNA levels by chimeric antisense oligonucleotides measured with RTS3370

ISIS No	Motif	Wing chemistry	% inhibition
146786	5-10-5	2'-MOE	84
510100	3-10-4	2'-MOE	76
552330	6-9-2	2'-MOE	54
552331	6-9-2	2'-MOE	66
552332	6-9-2	2'-MOE	70
552333	6-9-2	2'-MOE	55
552334	6-9-2	2'-MOE	42
552335	6-9-2	2'-MOE	39
552336	6-9-2	2'-MOE	27
552337	6-9-2	2'-MOE	74
552338	6-9-2	2'-MOE	68
552339	6-9-2	2'-MOE	71
552340	6-9-2	2'-MOE	61
552341	6-9-2	2'-MOE	58
552342	6-9-2	2'-MOE	55
552343	6-9-2	2'-MOE	63
552344	6-9-2	2'-MOE	51
552345	6-9-2	2'-MOE	65
552346	6-9-2	2'-MOE	0
552347	6-9-2	2'-MOE	84
552348	6-9-2	2'-MOE	87
552349	6-9-2	2'-MOE	74
552350	6-9-2	2'-MOE	59

552351	6-9-2	2'-MOE	60
552352	6-9-2	2'-MOE	53
552353	6-9-2	2'-MOE	0
552354	6-9-2	2'-MOE	83
552355	6-9-2	2'-MOE	90
552356	6-9-2	2'-MOE	0
552357	6-9-2	2'-MOE	45
552358	6-9-2	2'-MOE	74
552359	6-9-2	2'-MOE	72
552360	6-9-2	2'-MOE	87
552361	6-9-2	2'-MOE	96
552308	5-9-3	2'-MOE	81
552362	6-9-2	2'-MOE	92
552309	5-9-3	2'-MOE	77
552363	6-9-2	2'-MOE	92
552310	5-9-3	2'-MOE	80
552364	6-9-2	2'-MOE	87
552311	5-9-3	2'-MOE	13
552365	6-9-2	2'-MOE	84
552150	2-9-6	2'-MOE	73
552312	5-9-3	2'-MOE	77
552366	6-9-2	2'-MOE	87
552313	5-9-3	2'-MOE	64
552367	6-9-2	2'-MOE	85
552314	5-9-3	2'-MOE	73
552368	6-9-2	2'-MOE	77
552315	5-9-3	2'-MOE	75
552369	6-9-2	2'-MOE	75
552316	5-9-3	2'-MOE	64
552370	6-9-2	2'-MOE	63
552317	5-9-3	2'-MOE	99
552371	6-9-2	2'-MOE	81
552318	5-9-3	2'-MOE	76
552372	6-9-2	2'-MOE	65
552319	5-9-3	2'-MOE	55
552373	6-9-2	2'-MOE	74
552320	5-9-3	2'-MOE	68
552374	6-9-2	2'-MOE	78
552321	5-9-3	2'-MOE	74
552375	6-9-2	2'-MOE	81
552322	5-9-3	2'-MOE	73
552376	6-9-2	2'-MOE	78
552323	5-9-3	2'-MOE	75

552377	6-9-2	2'-MOE	70
552324	5-9-3	2'-MOE	0
552378	6-9-2	2'-MOE	72
552325	5-9-3	2'-MOE	70
552379	6-9-2	2'-MOE	74
552326	5-9-3	2'-MOE	63
552380	6-9-2	2'-MOE	53
552327	5-9-3	2'-MOE	30
552381	6-9-2	2'-MOE	26
552328	5-9-3	2'-MOE	25
552382	6-9-2	2'-MOE	13
552329	5-9-3	2'-MOE	33
552383	6-9-2	2'-MOE	5

Table 38

Inhibition of viral Target-Z mRNA levels by chimeric antisense oligonucleotides measured with RTS3370

ISIS No	Motif	Wing chemistry	% inhibition
509934	5-10-5	2'-MOE	30
551909	2-10-8	2'-MOE	62
551941	3-10-7	2'-MOE	74
551973	4-10-6	2'-MOE	64
551910	2-10-8	2'-MOE	52
551942	3-10-7	2'-MOE	54
551974	4-10-6	2'-MOE	51
551911	2-10-8	2'-MOE	58
551943	3-10-7	2'-MOE	64
551975	4-10-6	2'-MOE	57
551912	2-10-8	2'-MOE	59
551944	3-10-7	2'-MOE	66
551976	4-10-6	2'-MOE	57
551913	2-10-8	2'-MOE	58
551945	3-10-7	2'-MOE	56
551977	4-10-6	2'-MOE	56
551914	2-10-8	2'-MOE	0
551946	3-10-7	2'-MOE	48
551978	4-10-6	2'-MOE	53
551915	2-10-8	2'-MOE	44
551947	3-10-7	2'-MOE	53
551979	4-10-6	2'-MOE	64
551916	2-10-8	2'-MOE	57
551948	3-10-7	2'-MOE	68
551980	4-10-6	2'-MOE	56

551917	2-10-8	2'-MOE	58
551949	3-10-7	2'-MOE	64
551981	4-10-6	2'-MOE	63
551918	2-10-8	2'-MOE	59
551950	3-10-7	2'-MOE	71
551982	4-10-6	2'-MOE	63
551919	2-10-8	2'-MOE	76
551951	3-10-7	2'-MOE	71
551983	4-10-6	2'-MOE	73
551920	2-10-8	2'-MOE	68
551952	3-10-7	2'-MOE	76
551984	4-10-6	2'-MOE	81
551921	2-10-8	2'-MOE	83
551953	3-10-7	2'-MOE	82
551985	4-10-6	2'-MOE	76
551922	2-10-8	2'-MOE	73
551954	3-10-7	2'-MOE	68
551923	2-10-8	2'-MOE	59
551955	3-10-7	2'-MOE	71
551924	2-10-8	2'-MOE	80
551956	3-10-7	2'-MOE	80
551925	2-10-8	2'-MOE	82
551957	3-10-7	2'-MOE	88
551926	2-10-8	2'-MOE	71
551958	3-10-7	2'-MOE	74
551927	2-10-8	2'-MOE	68
551959	3-10-7	2'-MOE	69
551928	2-10-8	2'-MOE	69
551960	3-10-7	2'-MOE	62
551929	2-10-8	2'-MOE	54
551961	3-10-7	2'-MOE	20
551930	2-10-8	2'-MOE	53
551962	3-10-7	2'-MOE	60
551931	2-10-8	2'-MOE	47
551963	3-10-7	2'-MOE	63
551932	2-10-8	2'-MOE	68
551964	3-10-7	2'-MOE	56
551933	2-10-8	2'-MOE	72
551965	3-10-7	2'-MOE	67
551934	2-10-8	2'-MOE	64
551966	3-10-7	2'-MOE	73
551935	2-10-8	2'-MOE	68
551967	3-10-7	2'-MOE	60

551936	2-10-8	2'-MOE	67
551968	3-10-7	2'-MOE	63
551937	2-10-8	2'-MOE	47
551969	3-10-7	2'-MOE	36
551938	2-10-8	2'-MOE	41
551970	3-10-7	2'-MOE	43
551939	2-10-8	2'-MOE	53
551971	3-10-7	2'-MOE	55
551940	2-10-8	2'-MOE	50
551972	3-10-7	2'-MOE	58

Table 39

Inhibition of viral Target-Z mRNA levels by chimeric antisense oligonucleotides measured with RTS3371

ISIS No	Motif	Wing chemistry	% inhibition
509934	5-10-5	2'-MOE	21
551909	2-10-8	2'-MOE	52
551941	3-10-7	2'-MOE	62
551973	4-10-6	2'-MOE	58
551910	2-10-8	2'-MOE	48
551942	3-10-7	2'-MOE	36
551974	4-10-6	2'-MOE	45
551911	2-10-8	2'-MOE	61
551943	3-10-7	2'-MOE	56
551975	4-10-6	2'-MOE	60
551912	2-10-8	2'-MOE	53
551944	3-10-7	2'-MOE	48
551976	4-10-6	2'-MOE	48
551913	2-10-8	2'-MOE	53
551945	3-10-7	2'-MOE	54
551977	4-10-6	2'-MOE	48
551914	2-10-8	2'-MOE	0
551946	3-10-7	2'-MOE	56
551978	4-10-6	2'-MOE	36
551915	2-10-8	2'-MOE	47
551947	3-10-7	2'-MOE	45
551979	4-10-6	2'-MOE	54
551916	2-10-8	2'-MOE	44
551948	3-10-7	2'-MOE	59
551980	4-10-6	2'-MOE	49
551917	2-10-8	2'-MOE	48
551949	3-10-7	2'-MOE	60
551981	4-10-6	2'-MOE	57

551918	2-10-8	2'-MOE	53
551950	3-10-7	2'-MOE	57
551982	4-10-6	2'-MOE	57
551919	2-10-8	2'-MOE	65
551951	3-10-7	2'-MOE	57
551983	4-10-6	2'-MOE	53
551920	2-10-8	2'-MOE	57
551952	3-10-7	2'-MOE	67
551984	4-10-6	2'-MOE	62
551921	2-10-8	2'-MOE	60
551953	3-10-7	2'-MOE	57
551985	4-10-6	2'-MOE	58
551922	2-10-8	2'-MOE	63
551954	3-10-7	2'-MOE	61
551923	2-10-8	2'-MOE	50
551955	3-10-7	2'-MOE	44
551924	2-10-8	2'-MOE	52
551956	3-10-7	2'-MOE	46
551925	2-10-8	2'-MOE	54
551957	3-10-7	2'-MOE	51
551926	2-10-8	2'-MOE	70
551958	3-10-7	2'-MOE	72
551927	2-10-8	2'-MOE	60
551959	3-10-7	2'-MOE	61
551928	2-10-8	2'-MOE	57
551960	3-10-7	2'-MOE	58
551929	2-10-8	2'-MOE	49
551961	3-10-7	2'-MOE	26
551930	2-10-8	2'-MOE	54
551962	3-10-7	2'-MOE	57
551931	2-10-8	2'-MOE	46
551963	3-10-7	2'-MOE	56
551932	2-10-8	2'-MOE	57
551964	3-10-7	2'-MOE	53
551933	2-10-8	2'-MOE	65
551965	3-10-7	2'-MOE	54
551934	2-10-8	2'-MOE	58
551966	3-10-7	2'-MOE	69
551935	2-10-8	2'-MOE	63
551967	3-10-7	2'-MOE	53
551936	2-10-8	2'-MOE	67
551968	3-10-7	2'-MOE	60
551937	2-10-8	2'-MOE	51

551969	3-10-7	2'-MOE	42
551938	2-10-8	2'-MOE	40
551970	3-10-7	2'-MOE	38
551939	2-10-8	2'-MOE	32
551971	3-10-7	2'-MOE	46
551940	2-10-8	2'-MOE	39
551972	3-10-7	2'-MOE	51

Table 40

Inhibition of viral Target-Z mRNA levels by chimeric antisense oligonucleotides measured with RTS3371

ISIS No	Motif	Wing chemistry	% inhibition
146786	5-10-5	2'-MOE	40
510100	3-10-4	2'-MOE	60
552276	5-9-3	2'-MOE	44
552277	5-9-3	2'-MOE	39
552278	5-9-3	2'-MOE	37
552279	5-9-3	2'-MOE	50
552280	5-9-3	2'-MOE	2
552281	5-9-3	2'-MOE	0
552282	5-9-3	2'-MOE	13
552229	4-9-4	2'-MOE	17
552283	5-9-3	2'-MOE	27
552230	4-9-4	2'-MOE	53
552284	5-9-3	2'-MOE	0
552231	4-9-4	2'-MOE	31
552285	5-9-3	2'-MOE	56
552232	4-9-4	2'-MOE	35
552286	5-9-3	2'-MOE	43
552233	4-9-4	2'-MOE	40
552287	5-9-3	2'-MOE	44
552234	4-9-4	2'-MOE	0
552288	5-9-3	2'-MOE	44
552235	4-9-4	2'-MOE	13
552289	5-9-3	2'-MOE	21
552236	4-9-4	2'-MOE	40
552290	5-9-3	2'-MOE	34
552237	4-9-4	2'-MOE	37
552291	5-9-3	2'-MOE	34
552239	4-9-4	2'-MOE	58
552293	5-9-3	2'-MOE	61
552240	4-9-4	2'-MOE	54
552294	5-9-3	2'-MOE	62

552241	4-9-4	2'-MOE	47
552295	5-9-3	2'-MOE	63
552242	4-9-4	2'-MOE	61
552296	5-9-3	2'-MOE	61
552243	4-9-4	2'-MOE	55
552297	5-9-3	2'-MOE	52
552244	4-9-4	2'-MOE	45
552298	5-9-3	2'-MOE	27
552245	4-9-4	2'-MOE	41
552299	5-9-3	2'-MOE	32
552246	4-9-4	2'-MOE	67
552300	5-9-3	2'-MOE	57
552247	4-9-4	2'-MOE	74
552301	5-9-3	2'-MOE	76
552248	4-9-4	2'-MOE	65
552302	5-9-3	2'-MOE	68
552249	4-9-4	2'-MOE	38
552303	5-9-3	2'-MOE	59
552250	4-9-4	2'-MOE	43
552304	5-9-3	2'-MOE	30
552251	4-9-4	2'-MOE	52
552305	5-9-3	2'-MOE	49
552252	4-9-4	2'-MOE	51
552306	5-9-3	2'-MOE	56
552253	4-9-4	2'-MOE	47
552307	5-9-3	2'-MOE	49
552254	4-9-4	2'-MOE	50
552255	4-9-4	2'-MOE	64
552256	4-9-4	2'-MOE	57
552257	4-9-4	2'-MOE	51
552258	4-9-4	2'-MOE	62
552259	4-9-4	2'-MOE	59
552260	4-9-4	2'-MOE	56
552261	4-9-4	2'-MOE	54
552262	4-9-4	2'-MOE	47
552263	4-9-4	2'-MOE	45
552264	4-9-4	2'-MOE	52
552265	4-9-4	2'-MOE	58
552266	4-9-4	2'-MOE	54
552267	4-9-4	2'-MOE	43
552268	4-9-4	2'-MOE	57
552269	4-9-4	2'-MOE	34
552270	4-9-4	2'-MOE	37

552271	4-9-4	2'-MOE	42
552272	4-9-4	2'-MOE	36
552273	4-9-4	2'-MOE	25
552274	4-9-4	2'-MOE	11
552275	4-9-4	2'-MOE	38

Table 41

Inhibition of viral Target-Z mRNA levels by chimeric antisense oligonucleotides measured with RTS3371

ISIS No	Motif	Wing chemistry	% inhibition
146786	5-10-5	2'-MOE	38
509959	3-10-3	2'-MOE	49
510100	3-10-4	2'-MOE	55
552384	2-9-5	2'-MOE	41
552440	3-9-4	2'-MOE	57
552385	2-9-5	2'-MOE	53
552441	3-9-4	2'-MOE	38
552386	2-9-5	2'-MOE	42
552442	3-9-4	2'-MOE	72
552387	2-9-5	2'-MOE	43
552443	3-9-4	2'-MOE	56
552388	2-9-5	2'-MOE	18
552444	3-9-4	2'-MOE	39
552389	2-9-5	2'-MOE	24
552445	3-9-4	2'-MOE	53
552390	2-9-5	2'-MOE	40
552446	3-9-4	2'-MOE	57
552391	2-9-5	2'-MOE	51
552447	3-9-4	2'-MOE	53
552392	2-9-5	2'-MOE	0
552448	3-9-4	2'-MOE	57
552393	2-9-5	2'-MOE	52
552449	3-9-4	2'-MOE	49
552394	2-9-5	2'-MOE	32
552450	3-9-4	2'-MOE	44
552395	2-9-5	2'-MOE	33
552451	3-9-4	2'-MOE	38
552396	2-9-5	2'-MOE	46
552452	3-9-4	2'-MOE	30
552130	2-9-6	2'-MOE	46
552184	3-9-5	2'-MOE	34
552238	4-9-4	2'-MOE	41
552292	5-9-3	2'-MOE	45

552346	6-9-2	2'-MOE	0
552397	2-9-5	2'-MOE	37
552453	3-9-4	2'-MOE	45
552398	2-9-5	2'-MOE	42
552454	3-9-4	2'-MOE	39
552399	2-9-5	2'-MOE	34
552400	2-9-5	2'-MOE	47
552401	2-9-5	2'-MOE	53
552402	2-9-5	2'-MOE	47
552403	2-9-5	2'-MOE	70
552404	2-9-5	2'-MOE	44
552405	2-9-5	2'-MOE	0
552406	2-9-5	2'-MOE	25
552407	2-9-5	2'-MOE	23
552408	2-9-5	2'-MOE	73
552409	2-9-5	2'-MOE	71
552410	2-9-5	2'-MOE	52
552411	2-9-5	2'-MOE	62
552412	2-9-5	2'-MOE	50
552413	2-9-5	2'-MOE	55
552414	2-9-5	2'-MOE	64
552415	2-9-5	2'-MOE	45
552416	2-9-5	2'-MOE	45
552417	2-9-5	2'-MOE	37
552418	2-9-5	2'-MOE	73
552419	2-9-5	2'-MOE	68
552420	2-9-5	2'-MOE	64
552421	2-9-5	2'-MOE	54
552422	2-9-5	2'-MOE	60
552423	2-9-5	2'-MOE	62
552424	2-9-5	2'-MOE	60
552425	2-9-5	2'-MOE	46
552426	2-9-5	2'-MOE	48
552427	2-9-5	2'-MOE	36
552428	2-9-5	2'-MOE	57
552429	2-9-5	2'-MOE	36
552430	2-9-5	2'-MOE	42
552431	2-9-5	2'-MOE	60
552432	2-9-5	2'-MOE	44
552433	2-9-5	2'-MOE	55
552434	2-9-5	2'-MOE	46
552435	2-9-5	2'-MOE	47
552436	2-9-5	2'-MOE	25

552437	2-9-5	2'-MOE	19
552438	2-9-5	2'-MOE	25
552439	2-9-5	2'-MOE	22

Table 42

Inhibition of viral Target-Z mRNA levels by chimeric antisense oligonucleotides measured with RTS3371

ISIS No	Motif	Wing chemistry	% inhibition
509959	3-10-3	2'-MOE	49
552496	4-9-3	2'-MOE	35
552497	4-9-3	2'-MOE	60
552498	4-9-3	2'-MOE	20
552499	4-9-3	2'-MOE	45
552500	4-9-3	2'-MOE	53
552501	4-9-3	2'-MOE	56
552502	4-9-3	2'-MOE	50
552503	4-9-3	2'-MOE	36
552504	4-9-3	2'-MOE	50
552505	4-9-3	2'-MOE	53
552506	4-9-3	2'-MOE	49
552507	4-9-3	2'-MOE	35
552508	4-9-3	2'-MOE	62
552509	4-9-3	2'-MOE	65
552510	4-9-3	2'-MOE	54
552455	3-9-4	2'-MOE	60
552511	4-9-3	2'-MOE	65
552456	3-9-4	2'-MOE	69
552512	4-9-3	2'-MOE	63
552457	3-9-4	2'-MOE	4
552513	4-9-3	2'-MOE	50
552458	3-9-4	2'-MOE	59
552514	4-9-3	2'-MOE	53
552459	3-9-4	2'-MOE	69
552515	4-9-3	2'-MOE	68
552460	3-9-4	2'-MOE	3
552516	4-9-3	2'-MOE	65
552461	3-9-4	2'-MOE	37
552517	4-9-3	2'-MOE	54
552462	3-9-4	2'-MOE	42
552518	4-9-3	2'-MOE	23
552463	3-9-4	2'-MOE	28
552519	4-9-3	2'-MOE	32
552464	3-9-4	2'-MOE	72

552520	4-9-3	2'-MOE	61
552465	3-9-4	2'-MOE	68
552521	4-9-3	2'-MOE	68
552466	3-9-4	2'-MOE	76
552522	4-9-3	2'-MOE	71
552467	3-9-4	2'-MOE	72
552523	4-9-3	2'-MOE	73
552468	3-9-4	2'-MOE	50
552524	4-9-3	2'-MOE	49
552469	3-9-4	2'-MOE	65
552525	4-9-3	2'-MOE	45
552470	3-9-4	2'-MOE	58
552526	4-9-3	2'-MOE	39
552471	3-9-4	2'-MOE	30
552527	4-9-3	2'-MOE	39
552472	3-9-4	2'-MOE	43
552528	4-9-3	2'-MOE	43
552473	3-9-4	2'-MOE	25
552529	4-9-3	2'-MOE	50
552474	3-9-4	2'-MOE	70
552530	4-9-3	2'-MOE	73
552475	3-9-4	2'-MOE	64
552531	4-9-3	2'-MOE	62
552476	3-9-4	2'-MOE	50
552477	3-9-4	2'-MOE	66
552478	3-9-4	2'-MOE	68
552479	3-9-4	2'-MOE	60
552480	3-9-4	2'-MOE	58
552481	3-9-4	2'-MOE	54
552482	3-9-4	2'-MOE	44
552483	3-9-4	2'-MOE	17
552484	3-9-4	2'-MOE	64
552485	3-9-4	2'-MOE	56
552486	3-9-4	2'-MOE	26
552487	3-9-4	2'-MOE	42
552488	3-9-4	2'-MOE	35
552489	3-9-4	2'-MOE	46
552490	3-9-4	2'-MOE	41
552491	3-9-4	2'-MOE	38
552492	3-9-4	2'-MOE	47
552493	3-9-4	2'-MOE	49
552494	3-9-4	2'-MOE	22
552495	3-9-4	2'-MOE	0

Table 43

Inhibition of viral Target-Z mRNA levels by chimeric antisense oligonucleotides measured with RTS3371

ISIS No	Motif	Wing chemistry	% inhibition
146786	5-10-5	2'-MOE	56
509959	3-10-3	2'-MOE	54
552552	5-9-2	2'-MOE	32
552553	5-9-2	2'-MOE	53
552554	5-9-2	2'-MOE	48
552555	5-9-2	2'-MOE	39
552556	5-9-2	2'-MOE	39
552557	5-9-2	2'-MOE	54
552558	5-9-2	2'-MOE	41
552559	5-9-2	2'-MOE	56
552560	5-9-2	2'-MOE	39
552561	5-9-2	2'-MOE	51
552562	5-9-2	2'-MOE	56
552563	5-9-2	2'-MOE	31
552564	5-9-2	2'-MOE	31
552565	5-9-2	2'-MOE	53
552566	5-9-2	2'-MOE	46
552567	5-9-2	2'-MOE	63
552568	5-9-2	2'-MOE	66
552569	5-9-2	2'-MOE	60
552570	5-9-2	2'-MOE	60
552571	5-9-2	2'-MOE	44
552572	5-9-2	2'-MOE	52
552573	5-9-2	2'-MOE	20
552574	5-9-2	2'-MOE	36
552575	5-9-2	2'-MOE	19
552576	5-9-2	2'-MOE	61
552577	5-9-2	2'-MOE	57
552578	5-9-2	2'-MOE	71
552579	5-9-2	2'-MOE	59
552580	5-9-2	2'-MOE	58
552581	5-9-2	2'-MOE	51
552582	5-9-2	2'-MOE	40
552583	5-9-2	2'-MOE	35
552584	5-9-2	2'-MOE	50
552585	5-9-2	2'-MOE	48
552586	5-9-2	2'-MOE	74
552587	5-9-2	2'-MOE	68
552532	4-9-3	2'-MOE	59

552588	5-9-2	2'-MOE	67
552533	4-9-3	2'-MOE	52
552589	5-9-2	2'-MOE	47
552534	4-9-3	2'-MOE	71
552590	5-9-2	2'-MOE	58
552535	4-9-3	2'-MOE	59
552591	5-9-2	2'-MOE	46
552536	4-9-3	2'-MOE	19
552592	5-9-2	2'-MOE	44
552537	4-9-3	2'-MOE	26
552593	5-9-2	2'-MOE	39
552538	4-9-3	2'-MOE	54
552594	5-9-2	2'-MOE	52
552539	4-9-3	2'-MOE	50
552595	5-9-2	2'-MOE	57
552540	4-9-3	2'-MOE	60
552596	5-9-2	2'-MOE	58
552541	4-9-3	2'-MOE	68
552597	5-9-2	2'-MOE	52
552542	4-9-3	2'-MOE	63
552598	5-9-2	2'-MOE	51
552543	4-9-3	2'-MOE	44
552600	5-9-2	2'-MOE	51
552544	4-9-3	2'-MOE	45
552602	5-9-2	2'-MOE	13
552545	4-9-3	2'-MOE	42
552604	5-9-2	2'-MOE	42
552546	4-9-3	2'-MOE	46
552606	5-9-2	2'-MOE	42
552547	4-9-3	2'-MOE	38
552608	5-9-2	2'-MOE	37
552548	4-9-3	2'-MOE	49
552610	5-9-2	2'-MOE	41
552549	4-9-3	2'-MOE	34
552612	5-9-2	2'-MOE	23
552550	4-9-3	2'-MOE	13
552614	5-9-2	2'-MOE	11
552551	4-9-3	2'-MOE	8
552616	5-9-2	2'-MOE	6

Table 44

Inhibition of viral Target-Z mRNA levels by chimeric antisense oligonucleotides measured with RTS3371

ISIS No	Motif	Wing chemistry	% inhibition
146786	5-10-5	2'-MOE	47
509934	5-10-5	2'-MOE	67
552007	6-10-4	2'-MOE	53
552039	7-10-3	2'-MOE	74
552008	6-10-4	2'-MOE	47
552040	7-10-3	2'-MOE	57
552009	6-10-4	2'-MOE	70
552041	7-10-3	2'-MOE	65
552010	6-10-4	2'-MOE	51
552042	7-10-3	2'-MOE	59
552011	6-10-4	2'-MOE	47
552043	7-10-3	2'-MOE	36
552012	6-10-4	2'-MOE	62
552044	7-10-3	2'-MOE	82
552013	6-10-4	2'-MOE	72
552045	7-10-3	2'-MOE	62
552014	6-10-4	2'-MOE	73
552046	7-10-3	2'-MOE	74
552015	6-10-4	2'-MOE	66
552047	7-10-3	2'-MOE	60
552016	6-10-4	2'-MOE	67
552048	7-10-3	2'-MOE	60
552017	6-10-4	2'-MOE	72
552049	7-10-3	2'-MOE	68
552018	6-10-4	2'-MOE	89
552050	7-10-3	2'-MOE	86
552019	6-10-4	2'-MOE	87
552051	7-10-3	2'-MOE	86
551986	4-10-6	2'-MOE	64
552020	6-10-4	2'-MOE	86
552052	7-10-3	2'-MOE	87
551987	4-10-6	2'-MOE	76
552021	6-10-4	2'-MOE	84
552053	7-10-3	2'-MOE	75
551988	4-10-6	2'-MOE	5
552005	5-10-5	2'-MOE	72
552022	6-10-4	2'-MOE	80
552054	7-10-3	2'-MOE	83
551989	4-10-6	2'-MOE	64

552023	6-10-4	2'-MOE	78
552055	7-10-3	2'-MOE	57
551990	4-10-6	2'-MOE	83
552024	6-10-4	2'-MOE	89
552056	7-10-3	2'-MOE	82
551991	4-10-6	2'-MOE	0
552025	6-10-4	2'-MOE	89
552057	7-10-3	2'-MOE	89
551992	4-10-6	2'-MOE	67
552026	6-10-4	2'-MOE	84
552058	7-10-3	2'-MOE	82
551993	4-10-6	2'-MOE	78
552027	6-10-4	2'-MOE	85
552059	7-10-3	2'-MOE	85
551994	4-10-6	2'-MOE	82
552028	6-10-4	2'-MOE	82
552060	7-10-3	2'-MOE	74
551995	4-10-6	2'-MOE	81
552029	6-10-4	2'-MOE	81
552061	7-10-3	2'-MOE	81
551996	4-10-6	2'-MOE	79
552030	6-10-4	2'-MOE	86
552062	7-10-3	2'-MOE	85
551997	4-10-6	2'-MOE	80
552031	6-10-4	2'-MOE	86
551998	4-10-6	2'-MOE	74
552032	6-10-4	2'-MOE	78
551999	4-10-6	2'-MOE	79
552033	6-10-4	2'-MOE	80
552000	4-10-6	2'-MOE	84
552006	5-10-5	2'-MOE	86
552034	6-10-4	2'-MOE	81
552001	4-10-6	2'-MOE	66
552035	6-10-4	2'-MOE	55
552002	4-10-6	2'-MOE	54
552036	6-10-4	2'-MOE	58
552003	4-10-6	2'-MOE	50
552037	6-10-4	2'-MOE	43
552004	4-10-6	2'-MOE	56
552038	6-10-4	2'-MOE	66

Table 45

Inhibition of viral Target-Z mRNA levels by chimeric antisense oligonucleotides measured with RTS3371

ISIS No	Motif	Wing chemistry	% inhibition
146786	5-10-5	2'-MOE	61
510100	3-10-4	2'-MOE	66
552168	3-9-5	2'-MOE	64
552222	4-9-4	2'-MOE	76
552169	3-9-5	2'-MOE	65
552223	4-9-4	2'-MOE	41
552170	3-9-5	2'-MOE	58
552224	4-9-4	2'-MOE	58
552171	3-9-5	2'-MOE	51
552225	4-9-4	2'-MOE	49
552172	3-9-5	2'-MOE	23
552226	4-9-4	2'-MOE	36
552173	3-9-5	2'-MOE	44
552227	4-9-4	2'-MOE	20
552174	3-9-5	2'-MOE	28
552228	4-9-4	2'-MOE	29
552175	3-9-5	2'-MOE	56
552176	3-9-5	2'-MOE	66
552177	3-9-5	2'-MOE	53
552178	3-9-5	2'-MOE	57
552179	3-9-5	2'-MOE	56
552180	3-9-5	2'-MOE	51
552181	3-9-5	2'-MOE	51
552182	3-9-5	2'-MOE	63
552183	3-9-5	2'-MOE	60
552185	3-9-5	2'-MOE	67
552186	3-9-5	2'-MOE	37
552187	3-9-5	2'-MOE	68
552188	3-9-5	2'-MOE	71
552189	3-9-5	2'-MOE	51
552190	3-9-5	2'-MOE	47
552191	3-9-5	2'-MOE	50
552192	3-9-5	2'-MOE	80
552193	3-9-5	2'-MOE	73
552194	3-9-5	2'-MOE	58
552195	3-9-5	2'-MOE	60
552196	3-9-5	2'-MOE	54
552197	3-9-5	2'-MOE	64
552198	3-9-5	2'-MOE	62

552199	3-9-5	2'-MOE	57
552200	3-9-5	2'-MOE	52
552201	3-9-5	2'-MOE	73
552202	3-9-5	2'-MOE	60
552203	3-9-5	2'-MOE	60
552204	3-9-5	2'-MOE	63
552151	2-9-6	2'-MOE	71
552205	3-9-5	2'-MOE	64
552152	2-9-6	2'-MOE	69
552206	3-9-5	2'-MOE	71
552153	2-9-6	2'-MOE	63
552207	3-9-5	2'-MOE	71
552154	2-9-6	2'-MOE	56
552208	3-9-5	2'-MOE	52
552155	2-9-6	2'-MOE	61
552209	3-9-5	2'-MOE	50
552156	2-9-6	2'-MOE	40
552210	3-9-5	2'-MOE	66
552157	2-9-6	2'-MOE	45
552211	3-9-5	2'-MOE	63
552158	2-9-6	2'-MOE	66
552212	3-9-5	2'-MOE	62
552159	2-9-6	2'-MOE	68
552213	3-9-5	2'-MOE	64
552160	2-9-6	2'-MOE	78
552214	3-9-5	2'-MOE	72
552161	2-9-6	2'-MOE	57
552215	3-9-5	2'-MOE	54
552162	2-9-6	2'-MOE	54
552216	3-9-5	2'-MOE	49
552163	2-9-6	2'-MOE	65
552217	3-9-5	2'-MOE	50
552164	2-9-6	2'-MOE	48
552218	3-9-5	2'-MOE	39
552165	2-9-6	2'-MOE	46
552219	3-9-5	2'-MOE	41
552166	2-9-6	2'-MOE	42
552220	3-9-5	2'-MOE	32
552167	2-9-6	2'-MOE	47
552221	3-9-5	2'-MOE	33

Table 46

Inhibition of viral Target-Z mRNA levels by chimeric antisense oligonucleotides measured with RTS3371

ISIS No	Motif	Wing chemistry	% inhibition
146786	5-10-5	2'-MOE	
509934	5-10-5	2'-MOE	56
510100	3-10-4	2'-MOE	69
552071	8-10-2	2'-MOE	73
552114	2-9-6	2'-MOE	64
552115	2-9-6	2'-MOE	61
552116	2-9-6	2'-MOE	53
552117	2-9-6	2'-MOE	69
552072	8-10-2	2'-MOE	39
552118	2-9-6	2'-MOE	49
552119	2-9-6	2'-MOE	49
552120	2-9-6	2'-MOE	21
552121	2-9-6	2'-MOE	27
552073	8-10-2	2'-MOE	73
552122	2-9-6	2'-MOE	48
552074	8-10-2	2'-MOE	69
552123	2-9-6	2'-MOE	68
552075	8-10-2	2'-MOE	78
552124	2-9-6	2'-MOE	47
552076	8-10-2	2'-MOE	63
552125	2-9-6	2'-MOE	72
552077	8-10-2	2'-MOE	62
552126	2-9-6	2'-MOE	64
552078	8-10-2	2'-MOE	59
552127	2-9-6	2'-MOE	65
552079	8-10-2	2'-MOE	80
552128	2-9-6	2'-MOE	78
552080	8-10-2	2'-MOE	74
552129	2-9-6	2'-MOE	68
552130	2-9-6	2'-MOE	46
552131	2-9-6	2'-MOE	61
552132	2-9-6	2'-MOE	66
552133	2-9-6	2'-MOE	78
552081	8-10-2	2'-MOE	69
552134	2-9-6	2'-MOE	68
552135	2-9-6	2'-MOE	59
552136	2-9-6	2'-MOE	39
552137	2-9-6	2'-MOE	36
552082	8-10-2	2'-MOE	86

552138	2-9-6	2'-MOE	80
552083	8-10-2	2'-MOE	85
552139	2-9-6	2'-MOE	80
552084	8-10-2	2'-MOE	86
552140	2-9-6	2'-MOE	70
552085	8-10-2	2'-MOE	83
552141	2-9-6	2'-MOE	72
552086	8-10-2	2'-MOE	83
552142	2-9-6	2'-MOE	58
552087	8-10-2	2'-MOE	77
552143	2-9-6	2'-MOE	70
552144	2-9-6	2'-MOE	66
552145	2-9-6	2'-MOE	78
552146	2-9-6	2'-MOE	63
552088	8-10-2	2'-MOE	90
552147	2-9-6	2'-MOE	80
552089	8-10-2	2'-MOE	87
552148	2-9-6	2'-MOE	74
552090	8-10-2	2'-MOE	85
552149	2-9-6	2'-MOE	79
552091	8-10-2	2'-MOE	84
552092	8-10-2	2'-MOE	86
552093	8-10-2	2'-MOE	82
552094	8-10-2	2'-MOE	84
552063	7-10-3	2'-MOE	79
552095	8-10-2	2'-MOE	85
552064	7-10-3	2'-MOE	83
552096	8-10-2	2'-MOE	88
552065	7-10-3	2'-MOE	86
552097	8-10-2	2'-MOE	90
552066	7-10-3	2'-MOE	35
552098	8-10-2	2'-MOE	86
552067	7-10-3	2'-MOE	53
552099	8-10-2	2'-MOE	66
552068	7-10-3	2'-MOE	70
552100	8-10-2	2'-MOE	67
552069	7-10-3	2'-MOE	68
552101	8-10-2	2'-MOE	65
552070	7-10-3	2'-MOE	64
552102	8-10-2	2'-MOE	54

Table 47

Inhibition of viral Target-Z mRNA levels by chimeric antisense oligonucleotides measured with RTS3371

ISIS No	Motif	Wing chemistry	% inhibition
146786	5-10-5	2'-MOE	63
510100	3-10-4	2'-MOE	59
552330	6-9-2	2'-MOE	50
552331	6-9-2	2'-MOE	46
552332	6-9-2	2'-MOE	50
552333	6-9-2	2'-MOE	48
552334	6-9-2	2'-MOE	42
552335	6-9-2	2'-MOE	30
552336	6-9-2	2'-MOE	23
552337	6-9-2	2'-MOE	42
552338	6-9-2	2'-MOE	40
552339	6-9-2	2'-MOE	50
552340	6-9-2	2'-MOE	45
552341	6-9-2	2'-MOE	44
552342	6-9-2	2'-MOE	51
552343	6-9-2	2'-MOE	44
552344	6-9-2	2'-MOE	24
552345	6-9-2	2'-MOE	41
552346	6-9-2	2'-MOE	0
552347	6-9-2	2'-MOE	75
552348	6-9-2	2'-MOE	72
552349	6-9-2	2'-MOE	65
552350	6-9-2	2'-MOE	42
552351	6-9-2	2'-MOE	45
552352	6-9-2	2'-MOE	43
552353	6-9-2	2'-MOE	20
552354	6-9-2	2'-MOE	70
552355	6-9-2	2'-MOE	66
552356	6-9-2	2'-MOE	62
552357	6-9-2	2'-MOE	53
552358	6-9-2	2'-MOE	57
552359	6-9-2	2'-MOE	46
552360	6-9-2	2'-MOE	45
552361	6-9-2	2'-MOE	44
552308	5-9-3	2'-MOE	38
552362	6-9-2	2'-MOE	51
552309	5-9-3	2'-MOE	76
552363	6-9-2	2'-MOE	73
552310	5-9-3	2'-MOE	58

552364	6-9-2	2'-MOE	66
552311	5-9-3	2'-MOE	38
552365	6-9-2	2'-MOE	64
552150	2-9-6	2'-MOE	68
552312	5-9-3	2'-MOE	75
552366	6-9-2	2'-MOE	55
552313	5-9-3	2'-MOE	66
552367	6-9-2	2'-MOE	67
552314	5-9-3	2'-MOE	56
552368	6-9-2	2'-MOE	41
552315	5-9-3	2'-MOE	46
552369	6-9-2	2'-MOE	52
552316	5-9-3	2'-MOE	55
552370	6-9-2	2'-MOE	35
552317	5-9-3	2'-MOE	53
552371	6-9-2	2'-MOE	58
552318	5-9-3	2'-MOE	59
552372	6-9-2	2'-MOE	68
552319	5-9-3	2'-MOE	56
552373	6-9-2	2'-MOE	63
552320	5-9-3	2'-MOE	62
552374	6-9-2	2'-MOE	70
552321	5-9-3	2'-MOE	63
552375	6-9-2	2'-MOE	64
552322	5-9-3	2'-MOE	52
552376	6-9-2	2'-MOE	58
552323	5-9-3	2'-MOE	45
552377	6-9-2	2'-MOE	42
552324	5-9-3	2'-MOE	49
552378	6-9-2	2'-MOE	37
552325	5-9-3	2'-MOE	48
552379	6-9-2	2'-MOE	57
552326	5-9-3	2'-MOE	50
552380	6-9-2	2'-MOE	48
552327	5-9-3	2'-MOE	13
552381	6-9-2	2'-MOE	22
552328	5-9-3	2'-MOE	9
552382	6-9-2	2'-MOE	20
552329	5-9-3	2'-MOE	18
552383	6-9-2	2'-MOE	18

Example 10: Dose-dependent antisense inhibition of Target-Z mRNA in HepG2 cells

Antisense oligonucleotides from the study described in Example 52 exhibiting *in vitro* inhibition of Target-Z mRNA were selected and tested at various doses in HepG2 cells. Cells were plated at a density of 28,000 cells per well and transfected using LipofectAMINE2000® with 9.26 nM, 27.78 nM, 83.33 nM, and 250.00 nM concentrations of antisense oligonucleotide, as specified in Table 48. After a treatment period of approximately 16 hours, RNA was isolated from the cells and Target-Z mRNA levels were measured by quantitative real-time PCR. Target-Z primer probe set RTS3371 was used to measure mRNA levels. Target-Z mRNA levels were adjusted according to total RNA content, as measured by RIBOGREEN®. Results are presented as percent inhibition of Target-Z, relative to untreated control cells.

As illustrated in Table 48, Target-Z mRNA levels were reduced in a dose-dependent manner in antisense oligonucleotide treated cells. 'n/a' indicates that the data for that dosage is not available.

Table 48
Dose-dependent antisense inhibition of human Target-Z in HepG2 cells

ISIS No	9.2593 nM	27.7778 nM	83.3333 nM	250.0 nM
146786	10	43	74	89
509934	12	31	52	79
509959	4	24	49	67
510100	11	28	60	77
510124	3	11	13	41
551926	1	26	51	76
551958	15	17	56	82
551987	4	40	65	81
551990	7	55	78	91
551993	15	30	70	80
551994	0	30	39	58
551995	6	41	73	85
551996	13	47	71	85
551997	16	38	68	89
551998	4	36	69	85
551999	10	31	67	86
552000	0	17	61	78
552006	6	37	74	89
552009	1	5	39	60
552013	0	28	3	72
552014	0	26	32	77
552018	6	27	63	81
552019	15	34	65	90
552020	2	35	65	91
552021	4	11	53	82

552022	6	35	57	79
552023	11	33	59	81
552024	15	43	69	91
552025	17	35	69	87
552026	14	26	66	86
552027	3	46	62	88
552028	9	43	58	78
552029	8	40	72	89
552030	18	48	77	92
552031	0	38	66	89
552032	42	48	80	88
552033	2	40	64	84
552034	6	40	70	81
552039	2	33	56	83
552044	19	30	63	84
552046	4	21	47	77
552050	15	44	70	92
552051	8	33	69	90
552052	17	38	71	91
552053	0	40	59	86
552054	7	15	58	75
552056	19	62	86	92
552057	11	33	69	86
552058	30	55	79	90
552059	11	25	69	90
552060	9	32	61	86
552061	6	40	69	88
552062	22	48	75	89
552064	23	49	69	90
552065	10	8	69	86
552069	11	4	28	60
552073	9	31	62	78
552075	21	18	33	65
552077	0	17	40	72
552079	1	12	44	70
552080	3	12	34	69
552082	13	29	66	87
552083	24	54	69	88
552084	10	25	48	82
552085	28	35	64	85
552086	0	24	65	84
552088	33	53	77	93
552089	0	41	69	92

552090	17	35	70	87
552091	13	31	69	89
552092	6	23	66	89
552093	0	17	61	89
552094	12	38	65	88
552095	20	42	73	88
552096	n/a	39	66	91
552097	24	43	67	88
552098	0	24	56	85
552101	3	13	28	61
552147	11	27	58	80
552160	20	25	69	89
552163	0	21	22	53
552176	16	11	40	66
552192	7	38	78	89
552222	0	24	65	79
552247	0	38	69	86
552255	5	27	69	81
552301	5	38	65	86
552309	8	26	62	85
552312	0	4	32	62
552347	2	15	38	75
552348	12	40	42	65
552354	10	35	44	76
552361	2	25	55	74
552363	20	36	54	76
552374	7	4	38	76
552379	0	12	24	46
552403	8	27	54	76
552408	2	25	44	77
552409	6	31	56	80
552418	0	30	72	84
552420	9	34	53	81
552442	4	23	46	56
552466	0	23	56	79
552474	11	34	66	87
552477	11	22	44	64
552530	25	37	73	87
552559	9	13	29	51

Example 11: Efficacy of antisense oligonucleotides targeting Target-Z in transgenic mice

Target-Z transgenic mice were treated with ISIS antisense oligonucleotides selected from studies

described above and evaluated for their efficacy in this model.

Treatment

Groups of 12 mice each were injected subcutaneously twice a week for 4 weeks with 50 mg/kg of ISIS 510106, ISIS 510116, ISIS 505347, or ISIS 509934. A control group of 12 mice was injected subcutaneously twice a week for 4 weeks with PBS. Mice were euthanized 48 hours after the last dose, and livers were harvested for further analysis.

DNA and RNA Analysis

RNA was extracted from liver tissue for real-time PCR analysis of Target-Z DNA, using primer probe sets RTS3370, RTS3371, and RTS3372. The DNA levels were normalized to picogreen. Target-Z RNA samples were also assayed with primer probe sets RTS3370 and RTS3371 after RT-PCR analysis. The mRNA levels were normalized to RIBOGREEN®. The data is presented in Table 49, expressed as percent inhibition compared to the control group. As shown in Table 49, most of the antisense oligonucleotides achieved reduction of Target-Z DNA and RNA over the PBS control. Results are presented as percent inhibition of Target-Z mRNA or DNA, relative to control.

Table 49
Percent inhibition of Target-Z RNA and DNA in the liver of transgenic mice

ISIS No	% inhibition DNA (RTS3370)	% inhibition DNA (RTS3371)	% inhibition DNA (RTS3372)	% inhibition RNA (RTS3370)	% inhibition RNA (RTS3371)	% inhibition RNA (RTS3372)
510106	0	0	51	0	0	12
510116	68	79	68	49	54	66
505347	72	79	75	54	28	30
509934	93	95	94	72	75	92

Example 12: Efficacy of antisense oligonucleotides targeting Target-Z in transgenic mice

Target-Z transgenic mice were treated with ISIS antisense oligonucleotides selected from studies described above and evaluated for their efficacy in this model.

Treatment

Groups of 6 mice each were injected subcutaneously twice a week for 4 weeks with 50 mg/kg of ISIS 146779, ISIS 505358, ISIS 146786, ISIS 509974, ISIS 509958, or ISIS 509959. A control group of 10 mice was injected subcutaneously twice a week for 4 weeks with PBS. Mice were euthanized 48 hours after the last dose, and livers were harvested for further analysis.

DNA and RNA Analysis

RNA was extracted from liver tissue for real-time PCR analysis of Target-Z DNA, using primer probe sets RTS3370. The DNA levels were normalized to picogreen. Target-Z RNA samples were also assayed with primer probe sets RTS3370 after RT-PCR analysis. The mRNA levels were normalized to RIBOGREEN®. The data is presented in Table 50, expressed as percent inhibition compared to the control group. As shown in Table 50, most of the antisense oligonucleotides achieved reduction of Target-Z DNA and RNA over the PBS control. Results are presented as percent inhibition of Target-Z mRNA or DNA, relative to control.

Table 50
Percent inhibition of Target-Z RNA and DNA in the liver of transgenic mice

ISIS No	% inhibition DNA	% inhibition RNA
146779	39	5
505358	84	77
146786	83	73
509974	56	28
509958	82	29
509959	54	30

Example 13: Efficacy of antisense oligonucleotides targeting Target-Z in transgenic mice

Transgenic mice were treated with ISIS antisense oligonucleotides selected from studies described above and evaluated for their efficacy in this model.

Treatment

Groups of 6 mice each were injected subcutaneously twice a week for 4 weeks with 25 mg/kg of ISIS 146786, ISIS 552176, and ISIS 552073. One group of 10 mice was injected subcutaneously twice a week for 4 weeks with PBS. Mice were euthanized 48 hours after the last dose, and organs and plasma were harvested for further analysis.

DNA and RNA Analysis

RNA was extracted from liver tissue for real-time PCR analysis of Target-Z DNA, using primer probe set RTS3371. The DNA levels were normalized to picogreen. Target-Z RNA samples were also assayed with primer probe set RTS3371 after RT-PCR analysis. The mRNA levels were normalized to RIBOGREEN®. The data is presented in Table 51. Serum DNA samples were analyzed after the study period. The data is presented in Table 52, expressed relative to the levels measured in the control group. As shown in Tables 51 and 52, the antisense oligonucleotides achieved reduction of Target-Z DNA and RNA over the PBS control. Results are presented as percent inhibition of Target-Z mRNA or DNA, relative to

control.

Table 51

Percent inhibition of Target-Z RNA and DNA in transgenic mice

ISIS No	Dose (mg/kg/wk)	% inhibition of RNA	% inhibition of DNA
146786	50	81	91
552073	50	39	22
552176	50	55	56

Table 52

Serum levels of Target-Z DNA in transgenic mice, relative to control levels

ISIS No	Dose (mg/kg/wk)	Post-dose DNA levels
146786	50	0.1
552073	50	2.9
552176	50	2.1

Liver function

To evaluate the effect of ISIS oligonucleotides on hepatic function, plasma concentrations of ALT were measured using an automated clinical chemistry analyzer (Hitachi Olympus AU400e, Melville, NY) (Nyblom, H. et al., Alcohol & Alcoholism 39: 336-339, 2004; Tietz NW (Ed): Clinical Guide to Laboratory Tests, 3rd ed. W. B. Saunders, Philadelphia, PA, 1995). The results are presented in Table 53 expressed in IU/L. Both the ISIS oligonucleotides were considered tolerable in the mice, as demonstrated by their liver transaminase profile.

Table 53

ALT levels (IU/L) of transgenic mice

	Dose (mg/kg/wk)	ALT
PBS	-	77
ISIS 146786	50	21
ISIS 552073	50	19
ISIS 552176	50	27

Example 14: Efficacy of antisense oligonucleotides targeting Target-Z in transgenic mice

Transgenic mice were treated with ISIS antisense oligonucleotides selected from studies described above and evaluated for their efficacy in this model.

Treatment

Groups of 6 mice each were injected subcutaneously twice a week for 4 weeks with 25 mg/kg of ISIS

146786, ISIS 552056, ISIS 552088, and ISIS 552309. One group of 10 mice was injected subcutaneously twice a week for 4 weeks with PBS. Mice were euthanized 48 hours after the last dose, and organs and plasma were harvested for further analysis.

DNA and RNA Analysis

RNA was extracted from liver tissue for real-time PCR analysis of Target-Z DNA, using primer probe set RTS3371. The DNA levels were normalized to picogreen. Target-Z RNA samples were also assayed with primer probe set RTS3371 after RT-PCR analysis. The mRNA levels were normalized to RIBOGREEN®. As shown in Table 54, the antisense oligonucleotides achieved reduction of Target-Z DNA and RNA over the PBS control. Results are presented as percent inhibition of Target-Z mRNA or DNA, relative to control.

Table 54
Percent inhibition of Target-Z DNA and RNA in transgenic mice

	Dose (mg/kg/wk)	% inhibition (RNA)	% inhibition (DNA)
ISIS 146786	50	60	90
ISIS 552056	50	25	58
ISIS 552088	50	8	0
ISIS 552309	50	35	84

Example 15: Antisense inhibition of Target-Z viral mRNA in HepG2 cells by deoxy, MOE and (S)-cEt gapmers

Additional antisense oligonucleotides were designed targeting a Target-Z viral nucleic acid and were tested for their effects on Target-Z mRNA in vitro. Cultured HepG2 cells at a density of 28,000 cells per well were transfected using LipofectAMINE2000® with 100 nM antisense oligonucleotide. After a treatment period of approximately 24 hours, RNA was isolated from the cells and Target-Z mRNA levels were measured by quantitative real-time PCR. Viral primer probe sets RTS3370 and RTS3371 and were used to separately measure mRNA levels. Target-Z mRNA levels were adjusted according to total RNA content, as measured by RIBOGREEN®. Results are presented as percent inhibition of Target-Z, relative to untreated control cells.

The newly designed chimeric antisense oligonucleotides in Table below were designed as MOE gapmers or deoxy, MOE and (S)-cEt gapmers. The 5-10-5 MOE gapmers are 20 nucleosides in length, wherein the central gap segment comprises of ten 2'-deoxynucleosides and is flanked on both sides (in the 5' and 3' directions) by wings comprising five nucleosides each. The deoxy, MOE and (S)-cEt gapmers are 16 nucleosides in length wherein the nucleoside have either a MOE sugar modification, an (S)-cEt sugar modification, or a deoxy modification. The 'Chemistry' column describes the sugar modifications of each

oligonucleotide. 'k' indicates an (S)-cEt sugar modification; the number indicates the number of deoxynucleosides; otherwise, 'd' indicates a deoxynucleoside; and 'e' indicates a MOE modification. The internucleoside linkages throughout each gapmer are phosphorothioate (P=S) linkages. All cytosine residues throughout each oligonucleotide are 5-methylcytosines.

Each gapmer listed in Table 55 is targeted to the viral Target-Z genomic sequence.

Table 55

Inhibition of viral Target-Z mRNA levels by chimeric antisense oligonucleotides measured with RTS3370 or RTS3371

ISIS No	Motif	% inhibition (RTS3370)	% inhibition (RTS3371)
5808	Uniform deoxy	57	64
524518	eeee-10-eeee	62	72
146781	eeee-10-eeee	72	93
582665	eeee-10-eeee	57	59
582666	eeee-10-eeee	49	92
566831	kdkdk-9-ee	96	73
577123	eekk-9-ekce	84	96
577124	kdkdk-8-eeee	92	96
577126	kkk-8-eeee	87	90
566830	kdkdk-9-ee	93	95
577130	eek-10-kke	87	94
577131	kdkdk-9-ee	83	93
566828	kdkdk-9-ee	97	90
146786	eeee-10-eeee	93	71
566829	kdkdk-9-ee	98	84
577120	kdkdk-10-eeee	94	93
577127	kkk-8-eeee	95	70
577134	kek-8-eeee	94	89
577135	kek-10-kek	96	94
552859	eek-10-kke	92	91
577121	kdkdk-10-eeee	91	74
577128	kkk-8-eeee	92	85
577132	kdkdk-9-ee	97	81
577136	kek-10-kek	95	95
566832	kdkdk-9-ee	95	78
552870	eek-10-kke	71	93
577122	kdkdk-10-eeee	70	96
577125	kdkdk-8-eeee	70	94
577129	kkk-8-eeee	76	51
577133	kdkdk-9-ee	80	52

9591	Uniform deoxy	30	14
------	---------------	----	----

Example 16: Antisense inhibition of Target-Z viral mRNA in HepG2 cells by deoxy, MOE and (S)-cEt gapmers

Additional antisense oligonucleotides were designed targeting a Target-Z viral nucleic acid and were tested for their effects on Target-Z mRNA in vitro. ISIS 577121, ISIS 577122, ISIS 577123, ISIS 577132, ISIS 577133, and ISIS 577134, disclosed in the study described above, were also included in the assay. Cultured HepG2 cells at a density of 28,000 cells per well were transfected using Cytofectin with 9.375 nM, 18.75 nM, 37.50 nM, 75.00 nM, 150.00 nM, or 300.00 nM antisense oligonucleotide. After a treatment period of approximately 24 hours, RNA was isolated from the cells and Target-Z mRNA levels were measured by quantitative real-time PCR. Viral primer probe set RTS3371 was used to measure mRNA levels. Target-Z mRNA levels were adjusted according to total RNA content, as measured by RIBOGREEN®. Results are presented as percent inhibition of Target-Z, relative to untreated control cells.

The newly designed chimeric antisense oligonucleotides in Tables below were designed as deoxy, MOE and (S)-cEt gapmers. The deoxy, MOE and (S)-cEt gapmers are 16, 17, or 18 nucleosides in length wherein the nucleosides have either a MOE sugar modification, an (S)-cEt sugar modification, or a deoxy modification. The 'Chemistry' column describes the sugar modifications of each oligonucleotide. 'k' indicates an (S)-cEt sugar modification; the number indicates the number of deoxynucleosides; otherwise, 'd' indicates a deoxynucleoside; and 'e' indicates a MOE modification. The internucleoside linkages throughout each gapmer are phosphorothioate (P=S) linkages. All cytosine residues throughout each oligonucleotide are 5-methylcytosines.

Each gapmer listed in Table 56 is targeted to the viral genomic sequence.

Table 56
Chimeric antisense oligonucleotides targeting viral Target-Z genomic sequence

ISIS No	Motif
585163	eeekk-8-eeee
585164	eeekk-7-kkeee
585165	eeek-9-keee
585170	eeekk-7-kkeee
585171	eeek-9-keee
585172	eeekk-7-eeee
585173	ek-9-eeee
585174	ekdk-7-eeee
585166	eeekk-7-kkeee
585167	eeek-9-keee
577119	kdkdk-8-eeee

585168	eeekk-7-kkeee
585169	eeek-9-keee

Table 57

Dose dependent inhibition of Target-Z mRNA levels by chimeric antisense oligonucleotides

ISIS No	9.375 nM	18.75 nM	37.5 nM	75.0 nM	150.0 nM	300.0 nM
146786	37	37	58	70	81	93
505358	30	26	28	57	74	85
510100	42	30	43	61	77	91
552859	21	30	39	61	79	91
577119	42	43	46	66	74	75
577121	10	15	42	64	82	89
577122	21	30	53	66	78	84
577123	27	29	45	56	78	84
577132	14	21	42	61	80	92
577133	12	14	32	47	62	77
577134	37	39	59	72	86	90
585174	31	28	48	61	80	90

Table 58

Dose dependent inhibition of Target-Z mRNA levels by chimeric antisense oligonucleotides

ISIS No	9.375 nM	18.75 nM	37.5 nM	75.0 nM	150.0 nM	300.0 nM
146786	25	34	57	71	85	92
509932	9	28	59	62	70	74
585163	17	32	52	68	77	81
585164	23	4	29	31	36	56
585165	6	31	42	58	66	82
585166	19	27	35	48	50	63
585167	22	25	50	69	76	88
585168	4	30	44	52	67	76
585169	32	32	42	62	76	80
585170	23	19	39	49	66	75
585171	28	27	42	59	81	88
585172	26	29	30	64	80	91
585173	29	30	41	71	86	88

We claim:

1. A compound comprising:

a modified oligonucleotide consisting of 10 to 20 linked nucleosides, wherein the modified oligonucleotide comprises:

a 5'-wing consisting of 2 to 5 linked nucleosides;

a 3'-wing consisting of 2 to 5 linked nucleosides; and

a gap between the 5'-wing and the 3'-wing consisting of 6 to 14 linked 2'-deoxynucleosides; and

wherein at least one of the 5'-wing and the 3'-wing comprises at least one bicyclic nucleoside; at least one of the 5'-wing and the 3'-wing comprises at least one 2'-substituted nucleoside; and

wherein the nucleobase sequence of the modified oligonucleotide is complementary to the nucleobase sequence of a target nucleic acid.

2. The compound of claim 1, wherein one of the 5'-wing or the 3'-wing comprises at least one 2'-deoxynucleoside.

3. The compound of any of claims 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: e-e-e-k-k-(D)₈-e-e-e-e, wherein each k comprises a bicyclic nucleoside, each e comprises a 2'-modified nucleoside, and each D comprises a 2'-deoxynucleoside.

4. The compound of any of claims 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: e-e-e-k-(D)₉-k-e-e-e, wherein each k comprises a bicyclic nucleoside, each e comprises a 2'-modified nucleoside, and each D comprises a 2'-deoxynucleoside.

5. The compound of any of claims 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: e-k-e-k-(D)₉-e-e-e-e, wherein each k comprises a bicyclic nucleoside, each e comprises a 2'-modified nucleoside, and each D comprises a 2'-deoxynucleoside.

6.

7. The compound of any of claims 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: k-d-k-d-k-(D)₁₀-e-e-e-e-e, wherein each k comprises a bicyclic nucleoside, each e comprises a 2'-modified nucleoside, and each D comprises a 2'-deoxynucleoside.

8. The compound of any of claims 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: k-d-k-d-k-(D)₈-e-e-e-e, wherein each k comprises a bicyclic nucleoside, each e comprises a 2'-modified nucleoside, and each D comprises a 2'-deoxynucleoside.
9. The compound of any of claims 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: e-e-e-k-k-(D)₇-k-k-e-e-e, wherein each k comprises a bicyclic nucleoside, each e comprises a 2'-modified nucleoside, and each D comprises a 2'-deoxynucleoside.
10. The compound of any of claims 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: k-d-k-d-k-(D)₉-e-e, wherein each k comprises a bicyclic nucleoside, each e comprises a 2'-modified nucleoside, and each D comprises a 2'-deoxynucleoside.
11. The compound of any of claims 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: k-e-k-(D)₁₀-k-e-k, wherein each k comprises a bicyclic nucleoside, each e comprises a 2'-modified nucleoside, and each D comprises a 2'-deoxynucleoside.
12. The compound of any of claims 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: k-d-k-d-k-(D)₈-e-e-e-e, wherein each k comprises a bicyclic nucleoside, each e comprises a 2'-modified nucleoside, and each D comprises a 2'-deoxynucleoside.
13. The compound of any of claims 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: k-k-k-(D)₈-e-e-e-e-e, wherein each k comprises a bicyclic nucleoside, each e comprises a 2'-modified nucleoside, and each D comprises a 2'-deoxynucleoside.
14. The compound of any of claims 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: k-e-k-(D)₈-e-e-e-e-e, wherein each k comprises a bicyclic nucleoside, each e comprises a 2'-modified nucleoside, and each D comprises a 2'-deoxynucleoside.
15. The compound of any of claims 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: e-k-(D)₁₀-k-e-k-e, wherein each k comprises a bicyclic nucleoside, each e comprises a 2'-modified nucleoside, and each D comprises a 2'-deoxynucleoside.
16. The compound of any of claims 1-2, wherein the compound comprises a modified oligonucleotide having the sugar motif: e-e-e-(D)₁₀-k-k-k, wherein each k comprises a bicyclic nucleoside, each e comprises a 2'-modified nucleoside, and each D comprises a 2'-deoxynucleoside.

17. A pharmaceutical composition comprising the compound according to any of claims 1-16 and a pharmaceutically acceptable diluent.
18. A method of modulating expression of a target nucleic acid in a cell comprising contacting the cell with a compound according to any of claims 1-16.
19. A method of modulating expression of a target nucleic acid in an animal comprising administering to the animal the pharmaceutical composition according to claim 17.
20. Use of a compound of any of claims 1 to 16 or the pharmaceutical composition of claim 17 for the preparation of a medicament for the treatment of a disease or condition.