

(12) PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. AU 200110756 B2
(10) Patent No. 781502

(54) Title
Improved water soluble fertilizer compositions

(51)⁷ International Patent Classification(s)
C05F 011/00

(21) Application No: **200110756**

(22) Application Date: **2000.10.06**

(87) WIPO No: **WO01/25168**

(30) Priority Data

(31) Number	(32) Date	(33) Country
09/414214	1999.10.07	US

(43) Publication Date : **2001.05.10**

(43) Publication Journal Date : **2001.07.05**

(44) Accepted Journal Date : **2005.05.26**

(71) Applicant(s)
OMS Investments, Inc.

(72) Inventor(s)
**Michael Gustaaf Eltink; Philip Van Roij; Edze Jan Tijsma; Johannes
Gijsbertus Antonius Terlingen; Hendrikus Gijsbertus Adrianus Van Kaathoven**

(74) Agent/Attorney
**PHILLIPS ORMONDE and FITZPATRICK, 367 Collins Street, MELBOURNE VIC
3000**

(56) Related Art
**US 5372626
US 5772723
US 5830255**

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
12 April 2001 (12.04.2001)

PCT

(10) International Publication Number
WO 01/25168 A1

- (51) International Patent Classification⁷: C05F 11/00 (74) Agents: RADEN, James, B. et al.; Jones, Day, Reavis & Pogue, 77 West Wacker Drive, Chicago, IL 60601-1692 (US).
- (21) International Application Number: PCT/US00/27846
- (22) International Filing Date: 6 October 2000 (06.10.2000) (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data: 09/414,214 7 October 1999 (07.10.1999) US (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- (71) Applicant: OMS INVESTMENTS, INC. [US/US]; Suite 1300, 1105 N. Market Street, Wilmington, DE 19899-8985 (US).
- (72) Inventors: ELTINK, Michael, Gustaaf; De Tichel 44, NL-6417 RC Heerlen (NL). VAN ROIJ, Philip; Burge-meester Kessensingel A-6, NL-6227 VE Maastricht (NL). TIJSMA, Edze, Jan; Heerderweg 173, NL-6224 LD Maastricht (NL). ~~TERLINGEN, Gijsbertus, Antonius~~; Aan de Put 14, NL-6373 VT Landgraaf (NL). VAN KAATHOVEN, Hendrikus, Gijsbertus, Adrianus; Beijerstraat 28, NL-6118 CS Nieuwstadt (NL). *Johannes Gijsbertus Antonius Terlingen*

Published:

--- With international search report.

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



(54) Title: IMPROVED WATER SOLUBLE FERTILIZER COMPOSITIONS

$$61 \frac{n}{M_{w, acid}}$$

(1)

(57) Abstract: The present invention relates to a solid water soluble fertilizer composition containing one or more fertilizer materials and a phosphate free, organic acid which is solid at ambient temperatures. The fertilizer materials include primary macro nutrients, secondary macro nutrients, micro nutrients and mixtures thereof. The organic acid has a water solubility of at least 10 g/l (at 25 °C) and an acidifying effect in the range of 0.5 to 1.3 g HCO₃⁻/g acid.

The acidifying effect is defined as the amount of HCO₃⁻ that can be transformed into H₂CO₃ per gram of acid and is calculated in accordance with formula (1), wherein M_{w, acid} is the molecular weight of said acid and n represents the number of dissociation constants (i.e., pK_a values) of the acid below the value of 6.35. The acid should be present in the composition in an amount sufficient to reduce the HCO₃⁻ level in water treated with the composition by between about 60 - 400 parts per million when applied to the water at a dosage of 1 gram per liter.

WO 01/25168 A1

IMPROVED WATER SOLUBLE FERTILIZER COMPOSITIONSBACKGROUND OF THE INVENTION5 1. Field of the Invention

 The present invention relates to improved water soluble fertilizer compositions. More particularly, it relates to solid fertilizer compositions having sufficiently high acidifying effect (i.e. alkalinity
10 reducing effect) and water solubility to provide stable, precipitate free stock and feed solutions, independent of the phosphorus content thereof.

15 2. Description of Related Art

 In greenhouses, nurseries and other intensive horticulture environments, best plant growth results are achieved when macro and micro nutrients are carefully delivered to the growing plants. Many plant growers choose to utilize compound high analysis water soluble
20 fertilizers for accomplishing this result. Typically, such high analysis fertilizers are marketed as solids which may be dissolved by the user to prepare concentrated stock solutions which are subsequently diluted into irrigation water employing proportioners or
25 injection devices, thereby forming so called "feed solutions".

 Generally, it is important to formulate these high analysis fertilizer compositions so that they dissolve quickly and completely with no precipitation.
30 Furthermore, these fertilizer compositions must provide good long-term stability in stock solutions. For example, precipitates in the stock solutions can cause clogging of the proportioners and irrigation lines. These functional requirements for water soluble
35 fertilizer compositions of this variety have presented

-2-

ongoing problems for producers and developers of such products and these problems have not been fully solved by previously available fertilizer compositions.

For example, when previously conventional
5 fertilizer compositions have been employed for preparing fertilizer irrigation solutions from hard water and/or water having high alkalinity, problems have been encountered with respect to precipitation. Irrigation water having a relatively high alkalinity normally
10 contains a high content of (hydrogen) carbonates and the presence of such materials very often results in precipitation of secondary macro nutrients and micro nutrients (trace elements) from the fertilizer solution. Specifically, calcium is an important secondary macro
15 nutrient which is required in many plant fertilizer compositions. However, a relatively high pH will cause calcium and non-chelated micro nutrients to precipitate in the form of (hydrogen) carbonates, phosphates, sulfates and/or hydroxides.

20 Heretofore, various solutions have been proposed to decrease the alkalinity and consequently to reduce the precipitation of nutrients. For example, in in US Patents 5,830,255 and 5,174,806, phosphorus containing acids were disclosed for use as acidifiers in the
25 fertilizer compositions.

US Patent 5,830,255 describes a liquid fertilizer composition including phosphorous acid (H_3PO_3) as the primary macro nutrient. The composition may further include other nutrients and, additionally, contains
30 citric acid. It is taught that this composition improves the uptake by plants of phosphites (PO_3^{3-}) which are the salts of phosphorous acid. Phosphites are said to be taken up by the foliage of some plants more readily than phosphates and are, therefore, preferred for these

-3-

plants.

The problem underlying the invention described in US Patent 5,830,255 is the precipitation of phosphites, which are used as primary macro nutrients, which
5 precipitation occurs in conventional fertilizer compositions comprising phosphorous acid and other nutrients. In an attempt to solve this problem, citric acid is included in the fertilizer composition which, when diluted with water having a pH of about 6.5-8.5,
10 results in a pH of 5.0-7.0. The effect of using citric acid is, therefore, lowering of the pH in order to prevent the precipitation of the phosphites.

US Patent 5,174,806 describes a method for preparing a fertilizer composition including phosphoric acid (H_3PO_4) as the primary macro nutrient, other
15 nutrients and additionally citric acid and urea. This fertilizer composition is said to be neutral and to prevent the evolution of heat of neutralization between the phosphoric acid and other nutrients such as
20 potassium hydroxide due to the presence of citric acid and urea in the formulation.

However, it should be noted that the use of phosphorus containing acids, such as phosphorous acid and phosphoric acid as proposed in the above-cited prior
25 art, has several disadvantages.

One disadvantage is that phosphorus containing acids are liquids. Therefore, a grower wishing to fertilize with, for example, both calcium and a phosphorus containing acid will need to inject these two
30 elements separately. This makes other non-phosphorus containing acids such as nitric acid, sulphuric acid, formic acid, acetic acid and the like, unuseable in the formulation. Another disadvantage is that phosphorus containing acids are hazardous in handling and

-4-

application. Thirdly, liquid acids cannot be blended properly with solid fertilizer nutrients in powder form to produce a fertilizer composition. Therefore, manufacturing of such compositions, especially with
5 phosphorus containing acids, is troublesome. Additionally, chelates (e.g., being used as micro nutrients) are unstable in combination with liquid acids.

In view of the foregoing disadvantages, other
10 solutions to the problem of decreasing the alkalinity and, consequently, reducing the precipitation of nutrients were sought and one proposal was to use certain phosphates in the formulation.

For example, International Publication WO 92/013813
15 describes the application of urea phosphate in water soluble solid fertilizer compositions. This publication discloses that the use of mono-ammonium phosphate (MAP) in fertilizer mixtures with calcium nitrate, magnesium nitrate, ferrous sulphate, manganese sulphate, copper
20 sulphate, zinc sulphate and the like results in solutions with precipitates. However, the publication teaches that when urea phosphate (UP) is employed as the principal phosphorus source in a fertilizer mixture with the same secondary and micro nutrients, the use of UP
25 will permit calcium nitrate, magnesium nitrate and/or metal sulfates to be present in clear, concentrated stock solutions. This is a result that MAP, as the principal phosphorus source, is incapable of providing.

Also, in view of the acidifying effect of UP on
30 irrigation water, alkalinity can be reduced considerably which results in a significant increase in solubility for calcium and non-chelated micro nutrients, especially in case of hard water and/or water of high alkalinity, and the reduction in the concentration of

hydrogencarbonate avoids precipitation in irrigation water of high alkalinity. Thus, depending on the alkalinity, water hardness and composition of the fertilizer, this publication teaches that a certain amount of UP in the fertilizer composition is needed.

A disadvantage to the solubility problem solution provided by International Publication WO 92/013813 is that more conventional phosphorus sources, such as mono-ammonium phosphate, mono-potassium phosphate and di-ammonium phosphate, cannot be used, for example, with calcium salts to produce a precipitate free solution. To the contrary, urea phosphate must be used as the principal phosphorus source instead of these more common materials.

Another disadvantage is that for some plants or in some stages of growth, a high phosphorus level in the irrigation water is not desirable. Problems occur when a certain amount of UP is needed for hydrogencarbonate neutralization, i.e. acidifying effect and, simultaneously, a relatively high phosphorus level is undesirable from a plant growth perspective.

Additionally, other solutions have been proposed heretofore for reducing the precipitation of nutrients in fertilizer compositions.

For example, the application of low-molecular-weight organic acids like oxalic acid, citric acid and the like to the soil to increase the absorption of phosphates, used as primary macro nutrients, by plants is known (Biology and Fertility Soils, Vol. 18, No. 4, 1994, pp. 311-319). The organic acids increase the availability of phosphorus in soils mainly through both decreased adsorption of phosphates to the soil and increased solubility of the phosphorus compounds. These organic acids are applied separately from the

AMENDED SHEET

phosphates. Accordingly, the fertilizer composition used does not comprise any organic acids.




Another widely postulated proposal for solving the solubility problem is the use of chelated micro nutrients (trace elements) in fertilizer compositions in order to keep the micro nutrients in solution in both stock solution and irrigation water, containing phosphate salts as well. The use of chelated micro nutrients is necessary since when non-chelated micro nutrients (such as simple nitrates or sulfates) are used with conventional phosphorous sources, the micro nutrients tend to precipitate. A disadvantage of such use is that chelated micro nutrients increase the cost of the fertilizer compositions considerably.

Thirdly, the application of several types of acids which have a complex-forming (i.e. chelating) effect are well known in fertilizer compositions to stabilize metals (such as micro nutrients) by avoiding precipitation of the metals. Examples of such acids include ethylene diamine tetraacetic acid (EDTA), diethylene triamine pentaacetic acid (DTPA) and the like. A disadvantage is that these chelating acids are not soluble enough (less than 10 g/l at 25°C) to be used in solid water soluble fertilizers. Moreover, these acids have a low acidifying effect.

The discussion of the background to the invention herein is included to explain the context of the invention. This is not to be taken as an admission that any of the material referred to was published, known or part of the common general knowledge in Australia as at the priority date of any of the claims.

SUMMARY OF THE INVENTION

In view of the above discussed disadvantages of prior art fertilizer compositions and, furthermore, in view of the demands of fertilizers in general, it is a primary object of the present invention to provide nutrients for plants in a fertilizer composition that renders these nutrients readily available to the plants through irrigation.



It is a further object of the present invention to provide fertilizer compositions in which the nutrients present do not precipitate, even when hard water and/or water of high alkalinity is used as the irrigation water.

The foregoing and other objects of the invention are achieved by providing a solid fertilizer composition having sufficiently high acidifying, i.e. alkalinity reducing, effect and water solubility to provide a stable, precipitate free stock and feed solution, independent of the phosphorus content of the composition.

More specifically, fertilizer compositions in accordance with the present invention are water soluble, solid compositions containing phosphorus free, organic acids which are solid at ambient temperature, and one or more fertilizer materials. Suitable fertilizer materials for inclusion in the fertilizer compositions include primary macro nutrients, secondary macro nutrients, micro nutrients and mixtures thereof. The primary macro nutrients include phosphorus, nitrogen and potassium containing macro nutrients. Preferably, phosphorus containing macro nutrients are present in the form of phosphates.

Solid, phosphorus free, organic acids suitable for use in the present invention must have a water solubility of at least 10 grams/liter (at 25°C) and an acidifying effect in the range of 0.5 to 1.3 g HCO_3^- /gram of acid. The acidifying effect of the acid is defined as being the amount of HCO_3^- that can be transformed into H_2CO_3 per gram of the acid and is calculated in accordance with the following formula:

$$61 \frac{n}{M_{\text{acid}}}$$

wherein $M_{w,acid}$ is the molecular weight of said acid and n represents the number of dissociation constants (i.e., pKa values) of the acid below 6.35.

5 Furthermore, in accordance with the present invention, the water soluble, solid fertilizer compositions having the above described solid, phosphorus free, organic acids incorporated therein must provide a sufficient acidifying effect when dissolved in
10 water such as a hard and/or high alkalinity irrigation water stream to prevent precipitation of the nutrients therein. In this regard, it has been found that the compositions of the present invention, preferably, should be dissolved in water in an amount sufficient to
15 reduce the HCO_3^- level in the water treated with the composition by between about 60 and about 400 parts per million when the composition is applied to the water at a dosage of 1 gram per liter as measured by the formula wherein the acidifying effect of the fertilizer
20 compositions is calculated by determining the sum of the acidifying effects of the particular acids present in the fertilizer composition on a percentage weight/weight basis:

$$Ac_{product} = \sum_{i=1}^m 61 \frac{n}{M_{w,acid i}} f_{acid i} \cdot 1000 [ppm HCO_3^-]$$

25 wherein:

$Ac_{product}$ is the overall acidifying effect of a fertilizer product in parts per million (ppm) HCO_3^- in grams/mole at a dosage of 1 gram of water soluble fertilizer per liter of water;

30 61 is the molecular weight of bicarbonate or HCO_3^-

-9-

n represents the number of dissociation constants (pKa) of the acid in the composition which are equal to or less than pKa-value of 6.35 (i.e., the pKa value of carbonic acid);

5 $M_{w, \text{acid } i}$ is the molecular weight of the acid and is expressed in grams/mole;

$f_{\text{acid } i}$ is the (dimensionless) weight fraction of the acid in the fertilizer composition;

m is the number of acids in the fertilizer
10 composition having an acidifying effect; and
1000 is a conversion factor for converting grams into milligrams or ppm.

Exemplary of the advantages resulting from use of
15 the fertilizer compositions of the present invention are the following:

1. The ability to prepare and apply a complete nutrient solution with only one stock solution, made from a fertilizer composition of the present invention, and
20 one proportioner.

2. The ability to use non-chelated secondary macro nutrients (such as simple calcium salts) and non-chelated micro nutrient trace elements without reduction in solubility in the stock solution which is not
25 possible if conventional dry phosphorus sources without said solid acids are used.

3. Solid acidic fertilizers are significantly less hazardous to the end user than liquid fertilizer compositions based on phosphorus containing acids.

30

4. The ability to prepare precipitate free feed solutions from stock solutions, made from the fertilizer composition of the present invention and hard water and/or water of high alkalinity. Both the reduction of

-10-

the amount of hydrogencarbonate (alkalinity) in the irrigation water and the decrease in pH cause good solubility.

5 5. The ability to formulate acidifying fertilizer compositions without influencing the phosphorus level directly. This is not possible using urea phosphate (UP) according to prior art techniques such as those described in International Publication WO. The hydrogencarbonate reducing (i.e. acidifying) effect of
10 the solid acids employed in the compositions of the present invention is much higher than that of UP and, as a result, a smaller amount of the composition is required.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

The present invention provides a water soluble, solid fertilizer composition comprising a phosphorus free organic acid, the acid being solid at ambient temperature, and one or more fertilizer materials selected from the group consisting of primary macro nutrients, secondary macro nutrients and micro nutrients. The fertilizer compositions are dry solid materials. They are particulate, flowing solids having a free water content of less than about 10% by weight of the total composition and contain a solid acid having an acidifying effect between about 0.5 and about 1.3 grams HCO_3^- per gram acid and a solubility of at least about 10 grams per liter water under standard conditions (25°C) when the fertilizer composition is dissolved in water to provide an aqueous solution. The acidifying effect is preferably between about 0.5 to about 1.3 grams HCO_3^- per gram of acid, more preferably between about 0.9 to about 1.3 grams HCO_3^- per gram of acid with the acidifying effect being defined as the amount of HCO_3^- that can be transformed into H_2CO_3 per gram of the acid, which amount is calculated employing the following formula:

25

$$61 \frac{n}{M_{w, acid}}$$

wherein $M_{w, acid}$ is the molecular weight of the acid and n represents the number of dissociation constants (i.e., pKa values) of the acid below 6.35 and any phosphorus containing macro nutrients in the composition are in the form of phosphates, preferably mono-ammonium phosphate, di-ammonium phosphate, mono-potassium phosphate and tri-

-12-

potassium phosphate.

The acid component of the composition is preferably present in an amount of about 2 to about 90%, and more preferably about 5 to about 40%, by weight of the composition.

A low acidifying effect, i.e. less than about 0.5 gram HCO_3^- per gram acid, is disadvantageous since the lower the acidifying effect the greater the amount of the acid that has to be incorporated in the fertilizer composition.

The solid acid is preferably selected from the group consisting of malonic acid (dicarboxylic acid), DL-malic acid ((+)-2-hydroxysuccinic acid), maleic acid (cis-butenedioic acid), succinic acid (butane dioic acid), itaconic acid (methylenesuccinic acid), glutaric acid (1,5-pentanedioic acid), glycolic acid (hydroxyacetic acid), tricarballic acid (1,2,3-propanetricarboxylic acid), adipic acid (hexanedioic acid), pimelic acid (heptanedioic acid), citric acid (2-hydroxy-1,2,3-propanetricarboxylic acid), maleic anhydride (2,5-furandione) and succinic anhydride.

Oxalic acid and DL-tartaric acid (DL-dihydrosuccinic acid) are not included as suitable solid acids.

The amount of solid acid to be added depends on the hardness and/or alkalinity of the water to be applied, the amount of calcium, phosphates and non-chelated trace elements needed in the product and the value for the acidifying effect of the solid acid.

Two types of water hardness which have been recognized heretofore are general hardness (GH) and carbonate hardness (KH). A third term commonly used in this field is total hardness which is a combination of GH and KH. Since it is important to know both the GH and KH, the use of total hardness can be misleading and

-13-

should be avoided. General Hardness is primarily the measure of calcium (Ca^{2+}) and magnesium (Mg^{2+}) ions in the water. GH is commonly expressed in parts per million (ppm) of calcium carbonate (CaCO_3), degrees hardness (dH) or, more properly, the molar concentration of CaCO_3 . One German degree hardness (dH) is 10 mg of calcium oxide (CaO) per liter. In the U.S., hardness is usually measured in ppm of CaCO_3 . A German dH is 17.8 ppm CaCO_3 . A molar concentration of 1 milli-equivalent per liter (mEq/l) = 2.8 dH = 50 ppm. Water hardness follows the guidelines:

	0 - 4 dH,	0 - 70 ppm	: very soft
	4 - 8 dH,	70 - 140 ppm	: soft
15	8 - 12 dH,	140 - 210 ppm	: medium hard
	12 - 18 dH,	210 - 320 ppm	: fairly hard
	18 - 30 dH,	320 - 530 ppm	: hard

Carbonate hardness (KH) is the measure of hydrogencarbonate (HCO_3^-) and carbonate (CO_3^{2-}) ions in water. Alkalinity is the measure of the total acid binding capacity (all the anions which can bind with free H^+) but is comprised mostly of carbonate hardness in fresh water systems. Thus, the terms carbonate hardness, acid binding, acid buffering capacity and alkalinity are used interchangeably. KH is generally referred to in degrees hardness and is expressed in CaCO_3 equivalents just like GH.

As employed herein, the term "hard water" refers to water containing relatively high levels of Ca^{2+} and/or Mg^{2+} (high GH) whereas the term "high alkalinity water" refers to water having a high level of hydrogencarbonate (high KH).

As previously discussed, the intrinsic acidifying

-14-

effect of the organic acid used in the fertilizer composition, has been defined as the quantity of hydrogencarbonate (in grams) that can be transformed into carbonic acid per gram of acid, and can be
5 determined by the following formula:

$$61 \frac{n}{M_{w, acid}}$$

10 wherein $M_{w, acid}$ is the molecular weight of the acid and n represents the number of dissociation constants of the acid below a pKa-value of 6.35. In other words, n corresponds to the number of dissociation/protonation steps having a value lower than 6.35, which is the first
15 logarithmic dissociation/protonation constant of carbonic acid (e.g., see Handbook of Chemistry and Physics, David R. Lide, 76th Edition, 1995-1996). In general it can be stated that n represents the number of protons available per molecule of acid able to transform
20 hydrogencarbonate, HCO_3^- , into carbonic acid, H_2CO_3 . Carbonic acid dissociates into water and carbon dioxide under normal conditions. The number 61 is the molecular weight of hydrogencarbonate (in grams/mole).

The determination of the acidifying effect of the
25 acids can be exemplified as follows for two representative acids:

Malonic acid:

$$\begin{aligned} M_{w, malonic} &= 104 \text{ g/mol} \\ n &= 2 \text{ (pK}_1 = 2.83, \text{ pK}_2 = 5.69) \end{aligned}$$

30 Accordingly, citric acid has an acidifying effect of 1.2 grams HCO_3^- /gram acid as calculated employing the foregoing formula for the acidifying effect of an acid.

-15-

Citric acid:

 $M_{w,citric} = 192 \text{ g/mol}$ $n = 2 \text{ (} pK_1 = 3.14, pK_2 = 4.77, pK_3 = 6.39 \text{)}$

Accordingly, malonic acid has an acidifying effect
 5 of 0.6 grams HCO_3^- /gram acid as calculated employing
 the foregoing formula for the acidifying effect of
 an acid. In this regard, it should be noted that
 the third protonation/dissociation constant of
 citric acid is not taken into account in
 10 calculating the acidifying effect of citric acid
 since the pK_3 value is greater than 6.35.

In accordance with the present invention, the total
 acidifying effect of a fertilizer composition containing
 15 the organic acids which is dissolved in water to provide
 an aqueous solution can be determined by multiplying the
 acidifying effect of each acid present in the
 composition by the weight fraction of the acid incorpo-
 rated in the fertilizer composition. Specifically, the
 20 fertilizer compositions of this invention are formulated
 so that the HCO_3^- level in water will be reduced by
 between about 60 and about 400 parts per million when
 the composition is dissolved in the water at a dosage of
 1 gram per liter of water as defined by the formula:

25

$$Ac_{product} = \sum_{i=1}^m 61 \frac{n}{M_{w,acid i}} f_{acid i} \cdot 1000 [\text{ppm HCO}_3^-]$$

wherein:

$Ac_{product}$ is the overall acidifying effect of a
 30 fertilizer product in parts per million (ppm) HCO_3^-

-16-

at a dosage of 1 gram of water soluble fertilizer per liter of water;

61 is the molecular weight of bicarbonate or HCO_3^- in grams/mole;

5 n represents the number of dissociation constants (pKa) of the acid in the composition which are equal to or less than pKa-value of 6.35 (i.e., the pKa value of carbonic acid);

10 $M_{w, \text{acid } i}$ is the molecular weight of the acid and is expressed in grams/mole;

$f_{\text{acid } i}$ is the (dimensionless) weight fraction of the acid in the fertilizer composition;

m is the number of acids in the fertilizer composition having an acidifying effect; and
15 1000 is a conversion factor for converting grams into milligrams or ppm.

Thus, for example, assuming that the carbonate hardness of water to be treated is 250 ppm HCO_3^- (which is considered to be fairly hard water) and
20 assuming that it would be desirable to end up with a residual hardness of only 70 ppm HCO_3^- in the feed solution after application of a solid water soluble fertilizer in a concentration of 1 gram per liter of water, a fertilizer composition having an
25 acidifying effect of 180 ppm HCO_3^- would be required. A composition in accordance with the present invention which enables this result is as follows:

-17-

Composition 1.	
Fertilizer 10-14-27	w/w %
NPK and acid containing compounds	
Mono ammonium phosphate	23
Potassium nitrate	58,6
Maleic acid	17,1
Trace elements containing compounds	
Boric acid	0,090
Cobalt sulfate	0,003
Copper sulfate	0,042
Iron-EDTA	0,900
Manganese sulfate	0,155
Sodium molybdate	0,025
Zinc sulfate	0,085
total	100,00

It should be noted that the only acid in the above formulation for the 10-14-27 fertilizer composition which contributes to the acidifying effect is maleic acid and maleic acid can be characterized as follows in terms of the variables in the foregoing formula for determining the acidifying effect of a composition of this invention:

10

Maleic Acid:

$$\begin{aligned}
 M_{w, \text{ maleic acid}} &= 116 \text{ g/mol} \\
 pK_1 &= 1.83 \\
 pK_2 &= 6.07 \\
 n &= 2 \\
 f_{\text{ maleic acid}} &= 0.17
 \end{aligned}$$

20

Accordingly, the acidifying effect of the 10-

-18-

14-27 fertilizer composition ($Ac_{product}$) can be calculated as being 179 ppm HCO_3^- as maleic acid is the only acid in the composition having an acidifying effect.

5 It should be further noted that the NPK-ratios of the above 10-14-27 fertilizer composition correspond to the needs of a particular plant in a particular stage of growth. To the contrary, the same NPK-ratios and acidifying effect cannot be
10 achieved employing urea phosphate (UP) as the acidifier in view of the significantly lower acidifying effect of UP relative to maleic acid and the negative influence UP exerts on both phosphorous and nitrogen content.

15 With regard to the primary macro nutrients for use in the compositions of this invention, they are selected from the group consisting of phosphorus, nitrogen and potassium containing macro nutrients.

20 Preferred nitrogen containing macro nutrients include nitrates, ammonium salts and urea derivatives. Preferred potassium containing macro nutrients include potassium salts.

25 The primary macro nutrients are preferably present in the composition in an amount of about 1 to about 99% by weight of the composition.

 The preferred secondary macro nutrients for use in the compositions of the present invention include non-chelated elements consisting of calcium, magnesium and sulfate salts.

30 The secondary macro nutrients are preferably present in the composition in an amount of about 0.1 to about 99% by weight of the composition.

 Preferred micro nutrients for use in the compositions include non-chelated elements consis-

-19-

ting of iron, molybdenum, manganese, copper, zinc and cobalt sulphates and nitrates, boric acid and molybdates.

5 The micro nutrients are preferably present in the compositions in an amount of about 0.1 to about 50% by weight of the composition.

According to one embodiment of this invention, the fertilizer composition contains about 2 to about 90% and, preferably, about 5 to about 40% of the acid, and about 10 to about 99% and, preferably, about 60 to about 95% of other fertilizer materials by weight of the composition.

According to another embodiment, the fertilizer composition of the present invention contain about 2 to about 90% and, preferably, about 5 to about 40% of the acid; about 1 to about 99% and, preferably, about 5 to about 95% mono-ammonium phosphate or mono-potassium phosphate; about 0.1 to about 99% and, preferably, about 5 to about 95% calcium nitrate, and about 0 to about 75% of other fertilizer materials by weight of the composition.

According to yet another embodiment of this invention, the fertilizer composition contains about 2 to about 90% and, preferably, about 5 to about 40% of the acid; about 1 to about 99% and, preferably, about 5 to about 94% mono-ammonium phosphate or mono-potassium phosphate; about 1 to about 99% and, preferably, about 5 to about 94% calcium nitrate; about 0.1 to about 50% and, preferably, about 1 to about 40% trace metal sulfates and nitrates, and about 0 to about 75% of other fertilizer materials (including magnesium nitrate) by weight of the composition.

In a still further embodiment of this

invention, the fertilizer composition includes about 2 to about 90% and, preferably, about 5 to about 40% of the acid; about 1 to about 99% and, preferably, about 5 to about 94% mono-ammonium phosphate or mono-potassium phosphate; about 0.1 to about 50% and, preferably, about 1 to about 40% trace metal sulfates and nitrates; and about 0 to about 75% of other fertilizer materials by weight of the composition.

10 In yet another embodiment of this invention, the fertilizer composition does not include any phosphorus containing macro nutrients therein.

15 Examples of phosphorus, nitrogen and potassium containing macro nutrients to be included in the fertilizer composition of the present invention are ammonium nitrate, urea, ammonium sulphate, sodium nitrate, mono-ammonium phosphate, di-ammonium phosphate, mono-potassium phosphate, tri-potassium phosphate, potassium nitrate, potassium sulphate, potassium chloride, magnesium nitrate, calcium nitrate and the like.

25 The range of fertilizer compositions falling within the scope of the present invention are those having a phosphorus content (as wt.% P_2O_5) of from about 0 to about 60%, nitrogen content (as wt.% N) of from about 0 to about 45% and potassium content (as wt.% K_2O) of from about 0 to about 60%.

30 In addition to phosphorus, nitrogen and potassium containing primary macro nutrients, elements such as calcium and magnesium (i.e., secondary macro nutrients) and iron, manganese, copper, boron, zinc, molybdenum and cobalt (i.e.,

-21-

micro nutrients) can be included in the fertilizer composition of the present invention as other fertilizer materials.

5 These elements may in practice for example be included in the concentration ranges listed in the following Table 1:.

IPEA/US 18 SEP 2001

.22.

TABLE 1

Nutrient	Concentration range in final dry product (wt.%)
Ca	0-15
Mg	0-5.0
Fe	0-1.0
Mn	0-1.0
Cu	0-0.5
B	0-1.0
Zn	0-1.0
Mo	0-0.2
Co	0-0.1

Importantly, the elements from the secondary macro nutrients and micro nutrients do not have to be included in chelated form, such as EDTA or DTPA chelates or the like, but can rather be added as simple metal salts, especially nitrates or sulphates. Boron may be included as boric acid. Molybdenum may be provided as an alkali metal or ammonium molybdate. Magnesium can be present as magnesium nitrate.

The non-chelated secondary macro nutrients and micro nutrients include: calcium nitrate, magnesium nitrate, magnesium sulphate, ammonium sulphate, potassium sulphate, ferrous sulphate, ferrous nitrate, manganese sulphate, manganese nitrate, copper sulphate, copper nitrate, boric acid, zinc sulphate, zinc nitrate, sodium molybdate, ammonium molybdate and the like.

Of course, although not necessary, chelated metal salts such as Calcium-EDTA, Magnesium-EDTA,

-23-

iron-EDTA, iron-DTPA and copper-EDTA may be used as secondary macro nutrients and micro nutrients.

In addition, the fertilizer composition of the present invention can contain additional materials
5 such as cofactors, if desired.

The above nutrients are mixed as solids. The resulting products are dry solids as defined above. They should be stored in a water resistant packaging to minimize caking or lumping. Also other
10 soluble inerts (dyes, anti-caking agents, etc) may be added to these fertilizer compositions.

The solid fertilizer compositions of the present invention are made up into stock solutions and finally into feed solutions by dissolving in
15 water. This should be carried out in clean equipment usually with some agitation. Commonly, the concentration of the fertilizer composition in the stock solution is from about 5% to about 40% by weight of the solution, more preferably from about
20 10 to 20% by weight of the solution. This stock solution material is diluted by a factor of from about 10 to 200 for application to the plants which gives the final feed concentrations. Preferably, the concentration of the fertilizer composition in
25 the feed solution is from about 0.05% to about 1% by weight of the solution, more preferably from about 0.1 to about 0.15% by weight of the solution.

The use of solid acids having a relatively high acidifying effect and water solubility in a
30 dry blended mixture of nutrients which may include mono-ammonium phosphate, calcium, and optionally magnesium and/or trace metals in non-chelated forms such as nitrates and/or sulfates, offers several advantages.

-24-

For one, the acid establishes a low pH condition when the blended mixture is added to water to make a concentrated stock solution. For a stock solution, a pH in the range of 1 to 4 may be achieved. This low solution pH maintains solubility and clarity of the concentrated stock solution. The acid mentioned before, through the effect it has on solution pH, prevents the formation of calcium phosphate, magnesium phosphate and micro nutrient trace metal phosphate, which are not soluble under regular circumstances. Similarly, the low pH helps to prevent calcium from precipitating in the presence of sulphate and/or phosphate ions which may be present. Therefore, when a solid acid as defined before is used in combination with mono-ammonium phosphate as a primary phosphorus source, it will be possible to include phosphorus and a metal such as calcium, magnesium or a micro nutrient trace element like iron in one compound fertilizer composition, without the use of chelates or the disadvantage of a precipitate forming. This allows the end user to prepare and apply a complete nutrient solution using one stock solution and utilizing one injector. It enables also the use of non-chelated trace nutrients into phosphorus-containing nutrient solutions without precipitation.

However, precipitation can also occur in the feed solution in case a regular water soluble fertilizer is used in combination with hard water and/or water of high alkalinity. This is caused by the relatively high pH, large amount of hydrogencarbonates and calcium ions. Micro nutrient trace elements also tend to precipitate under these

circumstances if they are not applied in a chelated state. Also calcium present in water will precipitate and form gypsum with sulphates coming from the fertilizer. Solid acids, as mentioned
5 before, result in a reduction of both the amount of hydrogencarbonates and pH which result in a better solubility. This avoids precipitation of calcium phosphates, sulphates and/or carbonates and non-chelated micro nutrient trace elements in
10 phosphate, carbonate, sulphate or hydroxide form.

The usage of solid acids in a compound fertilizer product also allows the fertilizer solution to have an increased acidifying effect on the growing medium if needed.

15 Oxalic acid having an acidifying effect of 1.35 g HCO_3^- /g acid and tartaric acid, which are solid acids at ambient temperature, are not considered to be appropriate acids as they precipitate with both calcium and micro nutrient trace
20 elements in non-chelated form.

Urea phosphate has an acidifying effect of only 0.35 g HCO_3^- /g. In case of water of high alkalinity a fertilizer composition containing a large amount of urea phosphate would be needed.
25 This directly results in a high phosphorus level which is not desirable for certain plants or stages of growth. The solid acids used in the fertilizer composition of the present invention do not influence phosphorous levels.

30 The fertilizer composition of the present invention will be further described and advantages thereof will be made apparent with reference to the following Examples which are provided to illustrate the practice of the invention and not to limit its

-26-

scope of the invention as defined by the appended claims. All percentages are by weight unless otherwise indicated.

5

EXAMPLE 1

Several sample stock solutions in demineralized water were prepared using various combinations of nutrients without the presence of an organic acid. The nutrients included materials
10 which are commonly used in the manufacture of water soluble fertilizers. The most widely used source of phosphorus is mono-ammonium phosphate (MAP) which was compared with urea phosphate (UP). The compositions of the various stock solutions are
15 included in the following Table 2.

-27-

TABLE 2

Sample No.	MAP (g/l)	UP (g/l)	Ca(NO ₃) ₂ (g/l)	FeSO ₄ (g/l)	Mg(NO ₃) ₂ (g/l)	MnSO ₄ (g/l)	CuSO ₄ (g/l)	ZnSO ₄ (g/l)
1	1		100					
2	10		10					
3	100		100					
4	10			10				
5	100			100				
6	150		150		150			
7	50		95	1		0.5	0.25	0.5
8		1	100					
9		10	10					
10		100	100					
11		10		10				
12		100		100				
13		150	150		150			
14		50	95	1		0.5	0.25	0.5

-28-

The stability of the water solubility of the nutrients dissolved in the various stock solutions (Samples 1-14) was assessed by observing whether precipitation occurred within a period of 30 days.

5 As a result of this observation, It appeared that when MAP is used as the sole phosphorus source in mixtures with calcium nitrate, magnesium nitrate, ferrous sulphate, manganese sulphate, copper sulphate and/or zinc sulphate (e.g, as in tabulated Samples
10 1-7), precipitation occurs and that when UP is used instead of MAP (see tabulated Samples 8-14) precipitation does not occur. These results confirm the findings disclosed in International Publication WO 92/013813 which is discussed above.

15

EXAMPLE 2

Several stock solutions were prepared and the stability thereof was assessed in the same manner as described in Examples 1, with the exception that MAP
20 was used as the sole phosphorus source and that an organic acid, such as citric acid, maleic acid or malonic acid was added to the composition. The compositions of the various stock solutions are included in the following Table 3.

TABLE 3

Sample No.	MAP (g/l)	citric acid (g/l)	maleic acid (g/l)	malonic acid (g/l)	Ca(NO ₃) ₂ (g/l)	FeSO ₄ (g/l)	Mg(NO ₃) ₂ (g/l)	MnSO ₄ (g/l)	CuSO ₄ (g/l)	ZnSO ₄ (g/l)
1	1	0.61			100					
2	10	6.1			10					
3	100	61			100					
4	10	6.1				10				
5	100	61				100				
6	150	91.5			150		150			
7	50	30			95	1		0.5	0.25	0.5
8	1		0.37		100					
9	10		3.7		10					
10	100		37		100					
11	10		3.7			10				
12	100		37			100				
13	150		55		150		150			
14	50		18		95	1		0.5	0.25	0.5
15	1			0.33	100					
16	10			3.3	10					
17	100			33	100					
18	10			3.3		10				
19	100			33		100				
20	150			49	150		150			
21	50			16	95	1		0.5	0.25	0.5

-30-

It was observed that within a period of 30 days following preparation of the stock solutions (Samples 1-21), precipitation did not appear in any of these stock solution in accordance with the present invention.

Accordingly, the data developed in Examples 2 is believed to demonstrate that the use of a solid organic acid in accordance with this invention will enable MAP, calcium nitrate, magnesium nitrate and/or metal sulfates to be present in stable, clear, concentrated stock solutions without precipitation. In this regard, it should be noted that the results observed in Example 1 demonstrate that MAP, as a phosphorus source alone, is incapable of providing such stable, clear, concentrated stock solutions. Furthermore, it should be understood that although UP is capable of providing a clear solution, the use of UP has some clear disadvantages as mentioned in the above discussion of International Publication WO 92/013813.

Therefore, for example, it can be concluded that by adding an organic acid in accordance with this invention to a stock solution including MAP as the sole phosphorus source, precipitation of the nutrients is prevented and, likewise, the water solubility thereof is stabilized.

EXAMPLE 3

The solubility of several water soluble fertilizer compositions having a relatively high phosphate level in hard water having high alkalinity was tested in both a stock and feed solution at room temperature (25°C). Aspects such as precipitation, turbidity and pH were taken into account. The

-31-

turbidity was measured by a turbidity meter (Orbeco-Hellige) and expressed in Number of Turbidity Units (NTU). The higher the turbidity the higher the amount of insolubles.

- 5 The components of the basic composition prepared for the various sample fertilizer compositions to be tested in this Example are listed in the following Table 4:

TABLE 4

Basic Composition Fertilizer 10-50-10	Weight percent (wt.%)
Primary macro nutrients	
Urea	4.18
Ammonium sulphate	4.64
Mono-ammonium phosphate	61.25
Mono-potassium phosphate	24.36
Potassium sulphate	4.18
Micro nutrients	
Boric acid	0.096
Cobalt sulphate	0.004
Copper sulphate	0.045
Iron-EDTA	0.956
Manganese sulphate	0.166
Sodium molybdate	0.024
Zinc sulphate	0.094
Total	100.00

25 Normally hard water contains sufficient calcium and magnesium with respect to the needs of the plant, therefore no calcium or magnesium is needed in this product. Additionally, it should be noted that the fertilizer composition contains a rather high amount of phosphates and contains sulfates and non-chelated trace elements.

30 With the addition of calcium chloride and sodium hydrogencarbonate in the water supply, water of an alkalinity of 350 ppm HCO_3^- and a hardness of 350 ppm CaCO_3 was prepared. The room temperature was 25°C. The obtained pH was 7.8.

-33-

Various organic acids were incorporated into the basic composition (see Table 4) in accordance with the present invention to form Samples 4-21 listed in Table 5 hereinafter. As controls, Sample 1 was prepared containing no acid additive and Samples 2-3 were prepared containing urea phosphate. The weight fractions (wt.%) of the acids added, based on the total amount of basic composition employed and the acid added are listed in Table 5 as follows:

15 4 02 10758

TABLE 5

Sample Nos.	Acid added	Weight fraction	Precipitation		Turbidity (NTU)		pH	
			Stock	Feed	Stock	Feed	Stock	Feed
1	none	-	Yes	Yes	390	20	4.3	6.2
2	urea phosphate	0.718	No	No	0	0	2.4	3.1
3	urea phosphate	0.359	No	No	0	0	2.7	5.4
4	oxalic acid	0.207	Yes	Yes	710	450	2.6	3.3
5	malonic acid	0.239	No	No	0	0	2.9	4.7
6	malonic acid	0.120	No	No	0	0	3.2	5.6
7	succinic acid	0.272	No	No	73	0	3.5	4.8
8	succinic anhydride	0.230	No	No	75	0	3.5	4.8
9	maleic acid	0.267	No	No	0	0	2.7	4.9
10	maleic acid	0.134	No	No	0	0	3.0	5.6
11	maleic anhydride	0.226	No	No	0	0	2.7	4.9
12	DL-malic acid	0.308	No	No	0	0	3.2	4.3
13	L(+)-tartaric acid	0.346	Yes	No	660	0	2.7	3.7
14	itaconic acid	0.298	No	No	0	0	3.4	4.9
15	glutaric acid	0.304	No	No	0	0	3.6	4.8
16	adipic acid	0.337	No	No	0	0	3.8	5.1
17	pimelic acid	0.368	No	No	0	0	3.9	5.3
18	glycolic acid	0.350	No	No	0	0	3.2	4.0
19	tricarballic acid	0.269	Yes	No	20	0	3.4	5.0
20	citric acid	0.444	No	No	0	0	3.0	4.0
21	citric acid	0.222	No	No	0	0	3.2	5.1

In preparing the Samples 1-15 and 18-21 for testing in this Example 3, the solid acids were mixed with the basic composition before dissolving the mixture in 1 liter of the high alkalinity, hard water. Then, a stock solution was prepared containing 100 grams of the basic composition having the acid dissolved therein per liter of water. Successively, this stock solution was diluted 100 times to obtain a final feed solution of 1 gram of basic composition having the acid dissolved therein per liter of water.

In the case of the samples prepared with adipic acid (Sample 16) and pimelic acid (Sample 17, a stock solution of only 10 grams/liter was prepared as the solubility of these acids had to be taken into account. However, for all acids present in the test Samples 2-21 listed in Table 5, an amount of acid was added corresponding to an acidifying effect of 280 ppm HCO_3^- to maintain a residual amount of HCO_3^- of about 70 ppm. In some cases (Samples 3, 6, 10 and 21) half of this amount was chosen to test in order to judge whether smaller amounts of acid can also result in clear solutions.

The stability of the water solubility of the nutrients dissolved in the various stock and feed solutions was assessed by observing whether precipitation occurred and by measuring the turbidity and pH of the solution. The results obtained are listed in Table 5 under the heading "Precipitation".

In reviewing the tabulated results, it should be noted that as a result of the relatively high level of phosphates in the water, the presence of sulfates and non-chelated trace elements in the basic

composition in combination with the hard water (high Ca level) of high alkalinity (high HCO_3^- level), precipitation normally occurs. When no acid is added to the composition (Sample 1) precipitation indeed occurs, both in the stock and feed solution.

However, in view of the addition of a specified amounts of the acids in the samples, precipitation was avoided, both in the stock and feed solutions.

In order to reduce the pH as much as possible and to avoid a pH drop, it is advisable to apply a fertilizer containing an amount of acid which will leave a residual amount of hydrogencarbonate in the feed solution of 70 ppm. Nevertheless, depending on both the composition of the fertilizer and the hardness and/or alkalinity of the water used, smaller or even higher amounts can be used advantageously.

For satisfactory application of the fertilizer compositions of this invention it is critical that the feed solution is absolutely precipitate free. This will guarantee that all nutrients will be available to the plant and that precipitates will not block the irrigation system. In this respect, compositions according to Sample 1 (no acid added) and Sample 4 (oxalic acid) of this Example will result in significant problems in practice and are, therefore, not desirable. Of course, stock solutions also need to be optically clear and precipitate free. Also, the higher the concentration of a stock solution, the better.

As can be seen from the results tabulated in Table 5, the solid acids dissolved in the formulations are able to avoid precipitation in both feed and stock solution, except for oxalic acid (Sample 4) and L(+)tartaric acid (Sample 13). It is

-37-

known that oxalate precipitates with calcium. From the results, it is expected that tartarate also precipitates with calcium. So oxalic and tartaric acid are not considered to be appropriate for
5 reducing the pH of high alkalinity hard water.

EXAMPLE 4

The solubility of several water soluble fertilizer compositions having a relatively high
10 phosphate level in soft water of low alkalinity was tested in both a stock and feed solution at room temperature (25°C).

Because soft water generally does not contain sufficient calcium and magnesium with respect to the
15 needs of a plant, calcium and magnesium compounds (CaO and MgO, respectively) were incorporated into a basic composition prepared with the components listed in the following Table 6. It should be noted that this basic composition contains a rather high amount
20 of phosphates and, also, contains sulfates and non-chelated trace elements.

TABLE 6

Basic Composition Fertilizer 9-40-9 + 2CaO + 2 MgO	Weight percent (wt.%)
Primary macro nutrients	
Potassium nitrate	1.87
Ammonium sulphate	3.75
Mono-ammonium phosphate	49.46
Mono-potassium phosphate	19.67
Potassium sulphate	3.37
Secondary macro nutrients	
Magnesium nitrate	13.11
Calcium nitrate	7.65
Micro nutrients	
Boric acid	0.078
Cobalt sulphate	0.003
Copper sulphate	0.037
Iron-EDTA	0.77
Manganese sulphate	0.135
Sodium molybdate	0.02
Zinc sulphate	0.076
Total	100.00

-39-

Organic acids were added to the basic composition of Samples 3 and 4 in this Example 4 as tabulated in the following Table 7. As control samples, Sample 1 contained no acid and Sample 2 contained urea phosphate. The acids added and the weight fractions (wt.%) of these acids based on the total amount of the basic composition are listed in Table 7.

TABLE 7

Sample Nos.	Acid added	Weight fraction (wt.%)	Precipitation	Turbidity (NTU)	pH
1	none	-	Yes	168	3.9
2	urea phosphate	2.5	Yes	94	3.3
3	malonic acid	1.5	Yes	45	3.5
4	malonic acid	2	No	0	3.4

In preparation, the acids were mixed with the basic composition before dissolving the mixture in 1 liter of soft water of low alkalinity. A stock solution of 100 grams of this basic composition having the acid therein per liter of water was prepared.

The stability of the water solubility of the nutrients dissolved in the various stock solutions was assessed by observing whether precipitation occurred and by measuring the turbidity and pH of the solution. The results obtained are set forth in Table 7.

From the data in Table 7, it can be seen that a fertilizer composition which contains non-chelated trace elements, calcium, sulfates and a relatively high level of phosphates will result in precipitation
5 in a stock solution of soft water of low alkalinity (see Sample 1). This can be avoided by the addition of an appropriate amount of an acid (see Samples 3 and 4). It can also be concluded from the results in Table 7 that urea phosphate (see Sample 2) is not as
10 effective as for the samples containing malonic acid (Samples 3 and 4).

Although the invention has been described in its preferred forms with a certain degree of particularity, it is to be understood that the
15 present disclosure has been made by way of example only. Numerous changes in the details of the compositions and in the operational steps of the methods and in the materials utilized therein will be apparent without departing from the spirit and scope
20 of the invention, as defined in the appended claims.

Throughout the description and claims of the specification the word "comprise" and variations of the word, such as "comprising" and "comprises", is not intended to exclude other additives, components, integers or steps.

CLAIMS

We claim:

1. A water soluble, solid fertilizer
5 composition for producing a stable, essentially
precipitate free fertilizer solution comprising:

at least one phosphorus free organic acid which
is solid at ambient temperature and has a water
solubility of at least 10 grams/liter (at 25°C), the
10 acid having an acidifying effect in the range of 0.5
to 1.3 g HCO₃⁻/grams acid, the acidifying effect being
defined as the amount of HCO₃⁻ that can be transformed
into H₂CO₃ per gram of said acid, which amount is
calculated employing the following formula:

15

$$61 \frac{n}{M_{w, acid}}$$

wherein M_{w, acid} is the molecular weight of said acid and
n represents the number of dissociation constants of
the acid below a pKa value of 6.35,

20

the acid being present in an amount sufficient
to reduce the HCO₃⁻ level in water by between about
60 - 400 parts per million when the composition is
applied to water at a dosage of 1 gram per liter of
water enabling formation of a stable, essentially
25 precipitate free aqueous fertilizer solution, the
amount of acid being measured by the formula:

$$Ac_{product} = \sum_{i=1}^m 61 \frac{n}{M_{w, acid i}} f_{acid i} 1000 [ppm HCO_3^-]$$

wherein Ac_{product} corresponds with the overall
acidifying effect of a fertilizer product in parts

per million HCO_3^- at a dosage of 1 gram of water soluble fertilizer per liter of water; 61 is the molecular weight of bicarbonate or HCO_3^- in grams/mole; n represents the number of dissociation constants of the acid below a pKa value of 6.35; $M_{\text{acid } i}$ is the molecular weight of the acid and is expressed in grams/mole; $f_{\text{acid } i}$ is the (dimensionless) weight fraction of the acid in the fertilizer composition; m is the number of acids in a product and 1000 is the conversion factor for converting grams into milligrams or parts per million; and

at least one fertilizer material selected from the group consisting of phosphorus, nitrogen and potassium containing primary macro nutrients; non-chelated secondary macro nutrients; non-chelated micro nutrients; and mixtures thereof; wherein the phosphorus containing primary macro nutrients are in the form of phosphates.

2. The fertilizer composition of claim 1 wherein the acidifying effect of the acid is about 0.9 - 1.3 g HCO_3^- /gram of acid.

3. The fertilizer composition of claim 1 wherein the acid is selected from the group consisting of malonic acid, malic acid, maleic acid, succinic acid, itaconic acid, glutaric acid, glycolic acid, tricarballic acid, adipic acid, pimelic acid, citric acid, maleic anhydride, succinic anhydride and mixtures thereof.

4. The fertilizer composition of claim 1 wherein the acid is present in the composition in an amount of about 2 - 90% by weight of the composition.

5. The fertilizer composition of claim 4 wherein the said acid is present in an amount of about 5 - 40% by weight of the composition.

6. The fertilizer composition of claim 1 wherein the phosphates are selected from the group consisting of mono-ammonium phosphate, di-ammonium phosphate, mono-potassium phosphate and tri-potassium phosphate.

7. The fertilizer composition of claim 1 wherein the nitrogen containing macro nutrients are selected from the group consisting of nitrates, ammonium salts and urea derivatives.

8. The fertilizer composition of claim 1 wherein the potassium containing macro nutrients are selected from the group consisting of potassium salts.

9. The fertilizer composition of claim 1 wherein the primary macro nutrients are present in the composition in an amount of about 1 - 99% by weight of the composition.

10. The fertilizer composition of claim 1 wherein the non-chelated secondary macro nutrients are selected from the group consisting of calcium salts, magnesium salts, sulfate salts and mixtures thereof.

11. The fertilizer composition of claim 1 wherein the secondary non-chelated macro nutrients are present in an amount of 0.1 to 99% by weight of

AMENDED SHEET

18 SEP 2001

-44-

the composition.

12. The fertilizer composition of claim 1 wherein the non-chelated micro nutrients are selected from the group consisting of iron sulfates and nitrates, manganese sulfates and nitrates, copper sulfates and nitrates, zinc sulfates and nitrates, cobalt sulfates and nitrates, boric acid, molybdates and mixtures thereof.

10

13. The fertilizer composition of claim 1 wherein the non-chelated micro nutrients are present in the composition in an amount of about 0.1 - 50% by weight of the composition.

15

14. The fertilizer composition of claim 1 comprising the organic acid in an amount of about 1 - 90% by weight of the composition and the fertilizer material in an amount of about 10 - 99% by weight of the composition.

20

15. The fertilizer composition of claim 1 comprising:

the organic acid in an amount of about 2 - 90% by weight of the composition;

25

mono-ammonium phosphate or mono-potassium phosphate in an amount of about 1 - 99% by weight of the composition;

calcium nitrate in an amount of about 0.1 to 99% by weight of the composition; and

30

other fertilizer materials in an amount of about 0 to 75% by weight of the composition.

16. The fertilizer composition of claim 1 comprising:

the organic acid in an amount of about 2 - 90% by weight of the

5 composition;

mono-ammonium phosphate or mono-potassium phosphate in an amount of about .1 - 99% by weight of the composition;

10 calcium nitrate in an amount of about 1 to 99% by weight of the composition;

trace metal sulfates and nitrates in an amount of about 0.1 to 50% by weight of the composition; and

other fertilizer materials in an amount of about 0 to 75% by weight of the composition.

15

17. The fertilizer composition of claim 16 wherein the other fertilizer materials include magnesium nitrate.

20 18. The fertilizer composition of claim 1 comprising:

the organic acid in an amount of about 2 - 90% by weight of the composition;

25 mono-ammonium phosphate or mono-potassium phosphate in an amount of about 1 - 99% by weight of the composition;

trace metal sulfates and nitrates in an amount of about 0.1 to 50% by weight of the composition; and

30 other fertilizer materials in an amount of about 0 to 75% by weight of the composition.

19. The fertilizer composition of claim 18 wherein the other fertilizer materials include

AMENDED SHEET

magnesium nitrate.

20. The aqueous fertilizer solution of claim 29 wherein the acid is selected from the group consisting of malonic acid, malic acid, maleic acid, succinic acid, itaconic acid, glutaric acid, glycolic acid, tricarballic acid, adipic acid, pimelic acid, citric acid, maleic anhydride, succinic anhydride and mixtures thereof.

10

21. The aqueous fertilizer solution of claim 29 wherein the phosphates are selected from the group consisting of mono-ammonium phosphate, di-ammonium phosphate, mono-potassium phosphate and tri-potassium phosphate.

15

22. The aqueous fertilizer solution of claim 29 wherein the nitrogen containing macro nutrients are selected from the group consisting of nitrates, ammonium salts and urea derivatives.

20

23. The aqueous fertilizer solution of claim 29 wherein the potassium containing macro nutrients are selected from the group consisting of potassium salts.

25

24. The aqueous fertilizer solution of claim 29 wherein the primary macro nutrients are present in the composition in an amount of about 1 - 99% by weight of the composition.

30

25. The aqueous fertilizer solution of claim 29 wherein the non-chelated secondary macro nutrients are selected from the group consisting of calcium

salts, magnesium salts, sulfate salts and mixtures thereof.

26. The aqueous fertilizer solution of claim 29
5 wherein the secondary macro nutrients are present in an amount of 0.1 to 99% by weight of the composition.

27. The aqueous fertilizer solution of claim 29
10 wherein the non-chelated micro nutrients are selected from the group consisting of iron sulfates and nitrates, manganese sulfates and nitrates, copper sulfates and nitrates, zinc sulfates and nitrates, cobalt sulfates and nitrates, boric acid, molybdates and mixtures thereof.

15 28. The aqueous fertilizer solution of claim 29 wherein the micro nutrients are present in the composition in an amount of about 0.1 - 50% by weight of the composition.

20 29. A stable, essentially precipitate free aqueous fertilizer solution formed by dissolving a water soluble, solid fertilizer composition in water, the fertilizer composition comprising:

25 at least one phosphorus free organic acid which is solid at ambient temperature and has a water solubility of at least 10 g/l (at 25°C) and an acidifying effect in the range of 0.5 to 1.3 g HCO_3^-
30 /grams acid, the acidifying effect of the acid being defined as the amount of HCO_3^- that can be transformed into H_2CO_3 per gram of said acid, which amount is calculated employing the following formula:

$$61 \frac{n}{M_{w, acid}}$$

wherein $M_{w, acid}$ is the molecular weight of said acid and
 5 n represents the number of dissociation constants of
 the acid below a pKa value of 6.35;

the acid being present in an amount sufficient
 to reduce the HCO_3^- level in the water by between
 about 60 - 400 parts per million when the composition
 10 is applied to the water at a dosage of 1 gram per
 liter of water enabling formation of a stable,
 essentially precipitate free aqueous fertilizer
 solution, the amount of acid being measured by the
 formula:

$$Ac_{product} = \sum_{i=1}^m 61 \frac{n}{M_{w, acid i}} f_{acid i} \cdot 1000 [ppm HCO_3^-]$$

15

wherein $Ac_{product}$ corresponds with the overall
 acidifying effect of a fertilizer product in parts
 per million HCO_3^- at a dosage of 1 gram of water
 soluble fertilizer per liter of water; 61 is the
 20 molecular weight of bicarbonate or HCO_3^- in
 grams/mole; n represents the number of dissociation
 constants of the acid below a pKa value of 6.35; $M_{w, acid i}$
 is the molecular weight of the acid and is
 expressed in grams/mole; $f_{acid i}$ is the (dimensionless)
 25 weight fraction of the acid in the fertilizer
 composition; m is the number of acids in a product
 and 1000 is the conversion factor for converting
 grams into milligrams or parts per million; and

at least one fertilizer material selected from

AMENDED SHEET

the group consisting of phosphorus, nitrogen and potassium containing primary macro nutrients; non-chelated secondary macro nutrients; non-chelated micro nutrients and mixtures thereof; wherein the phosphorus containing primary macro nutrients are in the form of phosphates.

30. The aqueous fertilizer solution of claim 29 wherein the fertilizer composition comprises about 0.05% - 1% by weight of the solution.

31. The aqueous fertilizer solution of claim 29 wherein the fertilizer composition comprises about 0.05% - 1% by weight of the solution.

32. A fertilizer composition according to claim 1 substantially as hereinbefore described, with reference to any of the examples.

33. An aqueous fertilizer solution according to claim 29 substantially as hereinbefore described, with reference to any of the examples.

DATED: 1 April, 2005

PHILLIPS ORMONDE & FITZPATRICK

Attorneys for: OMS Investments, Inc.

David B Fitzpatrick