



(11) (21) (C) **2,181,985**
(86) 1995/01/04
(87) 1995/08/03
(45) 2000/12/05

(72) Desai, Subhash, US

(72) Mancini, Alan Mark, US

(72) Schumann, Steven Charles, US

(73) G.D. SEARLE & CO., US

(51) Int.Cl.⁶ A61K 31/415, A61K 9/20

(30) 1994/01/27 (08/187,568) US

(54) **COMPOSITIONS DE METRONIDAZOLE A LIBERATION
MODIFIEE ET LEURS PROCEDES DE FABRICATION ET
D'UTILISATION**

(54) **MODIFIED-RELEASE METRONIDAZOLE COMPOSITIONS
AND METHODS FOR MAKING AND USING SAME**

(57) L'invention se rapporte à des compositions pharmaceutiques présentant un profil à libération modifiée pour une dose quotidienne de métronidazole, à leurs procédés de fabrication, et à des procédés de traitement d'une infection microbienne au moyen d'une dose quotidienne des compositions pharmaceutiques de cette invention. Les compositions administrées une fois par jour présentent une bioéquivalence substantielle par rapport au métronidazole à libération immédiate administré trois fois par jour. Les compositions de l'invention comprennent: (a) une première part de métronidazole d'environ 59 % en poids à environ 79 % en poids; (b) environ 1,5 % en poids à environ 3 % en poids d'un copolymère d'ester d'acide poly(meth)acrylique, insoluble, aqueux qui est perméable à l'eau, expansible à l'eau et indépendant du niveau de pH; (c) environ 0,1 % en poids à environ 2 % en poids d'un agent anti-adhésif; (d) 0 à environ 23 % en poids d'un premier diluant pharmaceutique soluble, aqueux; (e) 0 à environ 23 % en poids d'un second diluant soluble, aqueux qui est approprié pour constituer un comprimé pharmaceutique lorsqu'il est comprimé avec les granules de (a), le second diluant soluble, aqueux étant identique ou différent du premier diluant soluble, aqueux; (f) 0 à environ 20 % en poids d'une

(57) Pharmaceutical compositions having a modified release profile for once daily dosing of metronidazole, methods for making the pharmaceutical compositions, and methods for treating a microbial infection with once daily dosing of the pharmaceutical compositions of the invention are provided. The compositions given once daily exhibit substantial bioequivalence to immediate release metronidazole given three times per day. The compositions of the invention comprise: (a) a first portion of metronidazole which is about 59 wt.% to about 79 wt.% metronidazole; (b) about 1.5 wt.% to about 3.0 wt.% of an aqueous insoluble poly(meth)acrylic acid ester copolymer which is aqueous permeable, aqueous expandable and pH-independent; (c) about 0.1 wt.% to about 2.0 wt.% detackifier; (d) 0 to about 23 wt.% of a first aqueous soluble pharmaceutical diluent; (e) 0 to about 23 wt.% of a second aqueous soluble diluent which is suitable for forming a pharmaceutical tablet when compressed with the granules of (a), the second aqueous soluble diluent being the same as or different from the first aqueous soluble diluent; (f) 0 to about 20 wt.% of a second portion of metronidazole; (g) 0 to about 0.2 wt.% glidant; and (h) 0 to about 2 wt.% lubricant; wherein the composition comprises metronidazole containing granules





(11) (21) (C) **2,181,985**
(86) 1995/01/04
(87) 1995/08/03
(45) 2000/12/05

seconde part de métronidazole; (g) 0 à environ 0,2 % en poids d'un agent de glissement; et (h) 0 à environ 2 % en poids d'un lubrifiant. La composition comprend des granules renfermant le métronidazole comprenant (a), (b), (c) et (d), la somme des pourcentages en poids du métronidazole obtenue par (a) et (f) est comprise entre environ 72 % en poids et environ 79 % en poids, et la somme des pourcentages en poids du diluant soluble, aqueux obtenue par (d) et (e) est comprise entre environ 16 % en poids et environ 23 % en poids.

comprising (a), (b), (c) and (d), wherein the sum of the weight percentages of metronidazole provided by (a) and (f) is between about 72 wt.% and about 79 wt.%, and wherein the sum of the weight percentages of the aqueous soluble diluent provided by (d) and (e) is between about 16 wt.% and about 23 wt.%.



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ :
A61K 31/415, 9/20

A1

(11) International Publication Number: WO 95/20383

(43) International Publication Date: 3 August 1995 (03.08.95)

(21) International Application Number: PCT/US95/00005

(22) International Filing Date: 4 January 1995 (04.01.95)

(30) Priority Data:
08/187,568 27 January 1994 (27.01.94) US

(60) Parent Application or Grant
(63) Related by Continuation
US 08/187,568 (CON)
Filed on 27 January 1994 (27.01.94)

(71) Applicant (for all designated States except US): G.D. SEARLE & CO. [US/US]; Corporate Patent Dept., P.O. Box 5110, Chicago, IL 60680-5110 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): DESAI, Subhash [US/US]; 157 Partridge Court, Grayslake, IL 60030 (US). MANCINI, Alan, Mark [US/US]; 23 Westwood Drive, Indian Head Park, IL 60525 (US). SCHUMANN, Steven, Charles [US/US]; 10 N. 819 South Airlite Street, Elgin, IL 60123 (US).

(74) Agents: FEDER, Scott, B. et al.; G.D. Searle & Co., Corporate Patent Dept., P.O. Box 5110, Chicago, IL 60680-5110 (US).

(81) Designated States: AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, JP, KE, KG, KP, KR, KZ, LK, LR, LT, LU, LV, MD, MG, MN, MW, MX, NL, NO, NZ, PL, PT, RO, RU, SD, SE, SI, SK, TJ, TT, UA, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD, SZ).

Published


With international search report.

Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.

(54) Title: MODIFIED-RELEASE METRONIDAZOLE COMPOSITIONS AND METHODS FOR MAKING AND USING SAME

(57) Abstract

Pharmaceutical compositions having a modified release profile for once daily dosing of metronidazole, methods for making the pharmaceutical compositions, and methods for treating a microbial infection with once daily dosing of the pharmaceutical compositions of the invention are provided. The compositions given once daily exhibit substantial bioequivalence to immediate release metronidazole given three times per day. The compositions of the invention comprise: (a) a first portion of metronidazole which is about 59 wt.% to about 79 wt.% metronidazole; (b) about 1.5 wt.% to about 3.0 wt.% of an aqueous insoluble poly(meth)acrylic acid ester copolymer which is aqueous permeable, aqueous expandable and pH-independent; (c) about 0.1 wt.% to about 2.0 wt.% detackifier; (d) 0 to about 23 wt.% of a first aqueous soluble pharmaceutical diluent; (e) 0 to about 23 wt.% of a second aqueous soluble diluent which is suitable for forming a pharmaceutical tablet when compressed with the granules of (a), the second aqueous soluble diluent being the same as or different from the first aqueous soluble diluent; (f) 0 to about 20 wt.% of a second portion of metronidazole; (g) 0 to about 0.2 wt.% glidant; and (h) 0 to about 2 wt.% lubricant; wherein the composition comprises metronidazole containing granules comprising (a), (b), (c) and (d), wherein the sum of the weight percentages of metronidazole provided by (a) and (f) is between about 72 wt.% and about 79 wt.%, and wherein the sum of the weight percentages of the aqueous soluble diluent provided by (d) and (e) is between about 16 wt.% and about 23 wt.%.



**MODIFIED-RELEASE METRONIDAZOLE COMPOSITIONS
AND METHODS FOR MAKING AND USING SAME**

FIELD OF THE INVENTION

This invention relates generally to metronidazole containing pharmaceutical compositions and particularly to modified release metronidazole compositions which, when taken once daily, release a therapeutically effective amount of metronidazole over a 24 hour period.

10

BACKGROUND OF THE INVENTION

The use of metronidazole, 1-(2-hydroxyethyl)-2-methyl-5-nitroimidazole, has long been known for the treatment of trichomoniasis and more recently for the treatment of bacterial vaginosis. There are currently at least two effective ways to treat trichomoniasis or bacterial vaginosis with the administration of a metronidazole composition. In the first method, a single, large dose (~2 grams) of metronidazole is given as a bolus to the patient. The single treatment is clinically effective. A major drawback to the administration of a single large dose of metronidazole is occurrence of significant and undesirable side effects such as nausea.

25

A second and more generally accepted treatment entails orally administering 250 mgs of metronidazole three times a day for a period of 7 days. The lower dosages of metronidazole over a period of a week significantly reduce the occurrence and severity of side effects. However, patient compliance is a problem since patients can inadvertently forget to take one or more doses during the course of treatment, causing the plasma metronidazole levels to drop to below an

30

acceptable therapeutic level for a period of several hours or more.

The use of metronidazole is also known for the treatment of various other conditions, including
5 amebiasis (acute amebic dysentery), and *Helicobacter pylori* infections associated with duodenal ulcer disease, see, e.g., D. Graham, et al., *Annals of Internal Medicine*, 115, 266-269 (1991).

In order to reduce the number of daily doses of
10 metronidazole needed to treat a microbial infection, while maintaining the benefits of making bioavailable effective amounts of the drug over an extended time period, it would be desirable to be able to deliver a therapeutically effective amount of metronidazole in a
15 once daily dose.

Effectively maintaining acceptable bioavailability of metronidazole for up to 24 hours with a single dose and without increased side effects relative to the conventional multiple dose regimen cannot be
20 accomplished simply by increasing the amount of active drug in a single dose. Metronidazole is aqueous soluble and is rapidly absorbed by the bloodstream. Metronidazole is also rapidly cleared from the bloodstream. Thus, merely administering increased
25 amounts of immediate-release metronidazole results in a rapid peak, followed by a rapid decline in metronidazole levels. Such a profile is undesirable because of side effects caused by high peak levels of metronidazole. Also, the rapid clearing of the drug
30 does not permit plasma metronidazole to remain at acceptable levels for 24 hours.

On the other hand, adding amounts of excipients in ratios typical of conventional modified release formulations which are presently available would result
35 in a tablet which is too large for oral administration. For example, 750 mg of metronidazole represents at least about a 2-fold greater amount of active

ingredient than is presently available in other
pharmaceutical compositions which are available in
modified release form. Moreover, metronidazole itself
is not readily compressible, which presents a
5 significant problem with respect to forming modified
release tablets.

Thus, there is a need to be able to provide a
modified release metronidazole composition which is
capable of delivering acceptable bioavailability for up
10 to 24 hours. The composition should be readily
compressible such that the entire dose may be provided
in a single tablet suitable for oral administration.
In order to keep the size of a single tablet in the
range of about 1000-1100 mg, while providing about 750
15 mg of metronidazole, a minor amount of excipient (less
than about 30% by weight) must be capable of imparting
both compressibility properties (for tableting) and
modified release properties (for bioavailability).
Heretofore, there has not been a modified release oral
20 tablet dosage of metronidazole which is suitable for
once daily dosing, even though metronidazole has been
an accepted therapeutic treatment for trichomoniasis
for over 25 years.

25

SUMMARY OF THE INVENTION

The present invention provides pharmaceutical
compositions for once daily dosing of metronidazole
having substantial bioequivalence to immediate release
30 metronidazole given three times per day. The
compositions of the present invention comprise greater
than 70% metronidazole and less than 30% excipient,
yet, surprisingly, are capable of being tableted and
of providing modified release of metronidazole for up
35 to 24 hours. The compositions of the invention
comprise metronidazole containing granules and

pharmaceutically acceptable excipients. The compositions comprise:

(a) a first portion of metronidazole which is about 59 wt% to about 79 wt% metronidazole;

5 (b) about 1.5 wt% to about 3.0 wt% of an aqueous insoluble poly(meth)acrylic acid ester copolymer which is aqueous permeable, aqueous expandable and pH-independent;

(c) about 0.1 wt% to about 2.0 wt% detackifier;

10 (d) 0 to about 23 wt% of a first aqueous soluble pharmaceutical diluent;

(e) 0 to about 23 wt% of a second aqueous soluble diluent which is suitable for forming a pharmaceutical tablet when compressed with the granules of (a), the second aqueous soluble diluent being the same as or different from the first aqueous soluble diluent;

(f) 0 to about 20 wt% of a second portion of metronidazole;

(g) 0 to about 0.2 wt% glidant; and

20 (h) 0 to about 2 wt% lubricant;

wherein the composition comprises metronidazole containing granules comprising (a), (b), (c) and (d), wherein the sum of the weight percentages of metronidazole provided by (a) and (f) is between about 72 wt% and about 79 wt%, and wherein the sum of the weight percentages of the aqueous soluble diluent provided by (d) and (e) is between about 16 wt% and about 23 wt%.

A presently preferred modified release metronidazole composition comprises 74.05 wt% metronidazole, 8.22 wt% hydrous lactose, 2.47 wt% Eudragit® NE 30 D (solids), 1.23 wt% talc, 0.03 wt% simethicone emulsion, 12.90 wt% anhydrous lactose (direct tablet grade), 0.10 wt% colloidal silicon dioxide, and 0.99 wt% magnesium stearate.

35 The present invention also includes a method for making a composition of the invention, which method

comprises producing metronidazole containing granules
in a fluid bed granulation step employing aqueous
insoluble poly(meth)acrylic acid ester copolymer as the
retard agent. The use of poly(meth)acrylic acid ester
5 copolymer as a retard agent unexpectedly yields
granules which provide both modified release
characteristics useful for once daily oral
administration of metronidazole and compressibility
characteristics which allow the compositions of the
10 invention to be compressed into tablet form.

The present invention also entails a method for
treating an infection caused by a microorganism which
is susceptible to metronidazole, which method comprises
administering to a patient having the infection an
15 effective amount of a composition of the invention once
daily for a number of days sufficient to eliminate or
resolve the infection.

BRIEF DESCRIPTION OF THE DRAWINGS

20

Figure 1 is a flow diagram representing a
preferred method for making modified release
metronidazole composition of the invention;

Figure 2 is a graphical depiction of the
25 bioavailability of a commercially available immediate
release metronidazole composition, a modified release
metronidazole composition comprising 2.47 wt%
poly(meth)acrylic acid ester (formulation C) showing
very good bioavailability, and a modified release
30 metronidazole composition comprising 1.95 wt%
poly(meth)acrylic acid ester (formulation D) showing
very good bioavailability;

Figure 3 is a graphical depiction of the
bioavailability (for comparison to Figure 2) of a
35 commercially available immediate release metronidazole
composition, a metronidazole composition comprising
5.05 wt% ethylcellulose as a release agent (formulation

E) showing very poor modified release characteristics, and 6.38 wt% ethylcellulose as a release agent (formulation F), also showing very poor modified release characteristics;

5 Figure 4 is a graphical depiction of the bioavailability (for comparison to Figure 2) of a commercially available immediate release metronidazole composition, a modified release composition comprising 8.99 wt% ethylcellulose dispersion as the release
10 agent, showing poor bioavailability (formulation A) and a modified release metronidazole composition comprising 4.6 wt% poly(meth)acrylic acid ester as the release agent also showing poor bioavailability (formulation B); and

15 Figure 5 is a graphical depiction of *in vitro* dissolution rates for modified release metronidazole compositions comprising poly(meth)acrylic acid ester.

20

DETAILED DESCRIPTION OF THE INVENTION

In one of its aspects the invention provides
5 modified release, metronidazole-containing compositions
which permit a slow release of metronidazole into the
bloodstream over a 24 hour period. The pharmaceutical
compositions of the present invention comprise:

(a) a first portion of metronidazole which is
10 about 59 wt% to about 79 wt% metronidazole;

(b) about 1.5 wt% to about 3.0 wt% of an aqueous
insoluble poly(meth)acrylic acid ester copolymer which
is aqueous permeable, aqueous expandable and pH-
independent;

15 (c) about 0.1 wt% to about 2.0 wt% detackifier;

(d) 0 to about 23 wt% of a first aqueous soluble
pharmaceutical diluent;

(e) 0 to about 23 wt% of a second aqueous soluble
diluent which is suitable for forming a pharmaceutical
20 tablet when compressed with the granules of (a), the
second aqueous soluble diluent being the same as or
different from the first aqueous soluble diluent;

(f) 0 to about 20 wt% of a second portion of
metronidazole;

25 (g) 0 to about 0.2 wt% glidant; and

(h) 0 to about 2 wt% lubricant;

wherein the composition comprises metronidazole
containing granules comprising (a), (b), (c) and (d),
wherein the sum of the weight percentages of
30 metronidazole provided by (a) and (f) is between about
72 wt% and about 79 wt%, and wherein the sum of the
weight percentages of the aqueous soluble diluent
provided by (d) and (e) is between about 16 wt% and
about 23 wt%.

35 The compositions of present invention release
greater than about 90%-98% of their total metronidazole
content at a controlled rate over a 24-hour period.

The C_{max} and AUC attained after a single dose are comparable to the C_{max} and AUC values which are achieved by thrice daily oral administration of 250 mg immediate release Flagyl® (G.D. Searle & Co., Skokie, IL 60077).

5 As used herein the term " C_{max} " means the maximum plasma metronidazole concentration achieved after oral administration of metronidazole. The terms "AUC" or "area under the curve" mean the total amount of metronidazole absorbed by the bloodstream in a
10 predetermined time, generally 24 hours. AUC is a measure of bioavailability which is calculated by integrating plasma metronidazole levels with respect to time.

The term "granules" or "metronidazole-containing
15 granules" as used herein refers to particles produced in a granulation process, which particles comprise metronidazole, poly(meth)acrylic acid ester copolymer, a detackifying agent and preferably an aqueous soluble diluent.

20 As used herein, all percentages for ingredients are weight percentages, based on the total weight of the pharmaceutical composition, unless otherwise stated. In this respect, it should be noted that the pharmaceutical compositions of the invention are
25 defined as comprising metronidazole containing granules. The granules are blended with certain excipients to produce the pharmaceutical compositions. The major constituent of the granules is metronidazole (i.e., the "first portion of metronidazole").

30 Accordingly, it will be appreciated that while a first portion of metronidazole is present in an amount up to 79% by weight, with respect to the total weight of the pharmaceutical composition, the granules may comprise up to about 98 wt% of metronidazole (i.e., "first
35 portion"), depending upon the amount of other constituents in the granules. In particularly preferred embodiments of the invention, the granules

comprise about 85 wt%-90 wt% metronidazole with respect to the total weight of the granules.

The compositions of the present invention may comprise, optionally, a second portion of metronidazole which is present in an amount of up to 20% by weight. This optional second portion of metronidazole is not utilized in forming the granules, but is mixed in with the granules after they are made. The addition of a second portion of metronidazole is referred to in the Examples herein as being added "neat." The second portion of metronidazole, if added, typically augments the release profile of the compositions of the invention such as by increasing the C_{max} attained.

The synthesis of metronidazole is well known, for example as shown in U.S. Patent No. 2,944,061. Metronidazole is also commercially available from Farchemia, S.p.A. (Treviglio, Italy) as well as from a variety of other commercial sources. It is presently preferred to mill the metronidazole (first portion and second portion, if any) to a particle size of about 40 mesh or smaller.

Metronidazole containing granules further comprise a poly(meth)acrylic acid ester copolymer as a modified-release agent or retard agent. Poly(meth)acrylic acid ester copolymers which are contemplated for use in the compositions of the invention are pH-independent, aqueous insoluble, aqueous permeable and aqueous expandable, such as those copolymers which are commercially available under the name Eudragit® NE30D from Röhm Pharma (Weiterstadt, Germany). By "pH-independent" it is meant that the copolymers are substantially insoluble in gastric juices, intestinal juices and water. By use of the term "aqueous permeable" is meant that aqueous solutions can pass through pores in the structure of the copolymer. By the term "aqueous expandable" it is meant that the copolymer composition is capable of swelling in an

aqueous solution. Eudragit® NE30D copolymers have an average molecular weight of about 800,000 daltons. The compositions of the present invention comprise from about 1.5% by weight to about 3.0% by weight, more preferably from about 1.9% to about 2.5% by weight of a poly(meth)acrylic acid ester copolymer.

Metronidazole containing granules desirably, but not necessarily, further comprise a first aqueous soluble diluent. Among the pharmaceutically acceptable, first aqueous soluble diluents which may be used are lactose, sucrose, dextrose, mannitol, sorbitol, and the like. The first aqueous soluble diluent may comprise up to 23% by weight of the total weight of the pharmaceutical compositions of the invention. It is presently preferred that the first aqueous soluble diluent is present in an amount of from about 5% to about 11% by weight, more preferably from about 6% to about 9% by weight, based on the total weight of the pharmaceutical composition. A particularly preferred first aqueous soluble diluent is hydrous lactose.

It should be appreciated that, inasmuch as the amount of first aqueous soluble diluent employed is incorporated in the granulation process, the amount of first aqueous soluble diluent used, if any, reduces proportionately the amount of metronidazole which may be granulated, and thus the number of tablets which may be produced in a single batch.

Since metronidazole containing granules comprising poly(meth)acrylic acid ester copolymer are characteristically tacky, the granules should also comprise an effective amount of a detackifying agent. Suitable detackifying agents, which may be present in effective amounts of from about 0.1% to about 2% by weight, include talc, magnesium stearate, calcium sulfate, glyceryl monostearate, and the like. Finely

particulated talc (e.g., 500 mesh) is a preferred detackifying agent.

5 A presently preferred method for preparing metronidazole containing granules employs fluidized bed granulation. In conventional fluidized bed granulation methods, dry particles of the active ingredient and diluent are suspended in a rising air column and substantially continuously sprayed with a liquid which binds the particles so as to gradually build-up
10 granules as they are suspended in the air column. One advantage of fluidized bed granulation is that it produces dry granules. It may be desirable, although not necessary, to include an effective amount of an antifoam agent (e.g., 0.01 wt% - 0.05 wt% simethicone
15 emulsion) in the liquid suspension which is sprayed during the fluid bed granulation process.

After the granulation process is completed, an effective amount of glidant (e.g., about 0.01% - 0.2% by weight or more) may be added to the produced
20 granules, if desired, to improve flow characteristics of the mixture, thereby facilitating transfer of the mixture during the manufacturing steps. Several pharmaceutically acceptable glidants are well known to those of ordinary skill in the art. A presently
25 preferred glidant is colloidal silicon dioxide.

In a preferred embodiment, compositions of the present invention comprise a second aqueous soluble diluent, which may be the same as or different from the first aqueous soluble diluent. The second aqueous
30 soluble diluent is an excipient which is blended with the produced granules and which may impart good compressibility, hardness and friability qualities to the compositions of the invention. Examples of second aqueous soluble diluents, include lactose, sucrose,
35 dextrose, mannitol, sorbitol and the like. These pharmaceutical diluents are well known and commercially available in forms which provide desirable

compressibility traits (e.g., direct tablet grade (DTG) anhydrous lactose).

The compositions of the invention may comprise up to about 23% by weight, more preferably from about 8% to about 18% by weight, and most preferably from about 12% to about 17% by weight of the second aqueous soluble diluent. A presently preferred second aqueous soluble diluent is anhydrous lactose (direct tablet grade). While the first aqueous soluble diluent and the second aqueous soluble diluent may each comprise up to 23% by weight of the compositions of the invention, importantly, the sum of the weight percentages of the first and the second aqueous soluble diluents should be from about 16% to about 23% based on the total weight of the compositions.

The compositions of the present invention may be conventionally compressed into tablets after the metronidazole containing granules are mixed in with the above-described excipients. It is presently preferred that each tablet comprise between about 600 mg and about 1000 mg of metronidazole, more preferably about 750 mg of metronidazole, so as to provide a once daily dosage of metronidazole suitable for administration for a period of several days. It will be appreciated, however, that the compositions may be tableted into smaller weight units if it were desired to administer more than one tablet at a time.

To facilitate mechanical release of the tablets from the tablet press after compression, an effective amount of a pharmaceutically acceptable lubricant, as is well known in the art, may be added to the compositions. It is presently preferred to use magnesium stearate in an amount of from about 0.1% to about 2.0% by weight.

35

In another of its aspects, the present invention entails a method for making a modified release metronidazole tablet which is capable of delivering, in a single oral dosage, a therapeutically effective amount of metronidazole over a 24 hour period. The method of the invention comprises the steps of:

(a) contacting in a fluid bed granulator, under conditions suitable for producing granules, (1) a dry mixture comprising from about 59 to about 79 parts by weight of a first portion of metronidazole and optionally up to 23 parts by weight of a first aqueous soluble diluent with (2) an aqueous suspension comprising from about 1.5 to about 3.0 parts by weight of an aqueous insoluble, pH-independent, aqueous expandable polymethacrylic acid ester copolymer, and an effective amount of a detackifying agent;

(b) combining the granules produced in step (a) with an effective amount of a pharmaceutically acceptable glidant;

(c) if necessary, particle-sizing the mixture of (b) to provide a mixture with a substantially uniform particle size suitable for compressing into tablet form;

(d) optionally blending the mixture with up to about 20 parts by weight of a second portion of metronidazole;

(e) blending the mixture with up to 23 parts by weight of a second aqueous soluble diluent and an effective amount of a pharmaceutically acceptable lubricant; and

(f) compressing a predetermined amount of the blended mixture of step (e) to produce a tablet.

By use of the phrase "particle-sizing" with respect to a mixture comprising metronidazole granules is meant processing a blended mixture which may have a wide distribution of particle sizes to yield a mixture having a substantially uniform particle size. Milling

and sieving procedures, used alone or in combination, are well known in the art for achieving a mixture having a uniform particle size within a predetermined size range (e.g., 20 mesh).

5 With reference to FIG. 1, an especially preferred method 10 for making the compositions of the invention, utilizing fluid bed granulation, is described. Metronidazole 12 and hydrous lactose 14 are added as dry ingredients into the fluid bed granulator. The dry starting ingredients may have a particle size within a broad range so long as the particles may be suspended by the air flow during granulation. An aqueous suspension 16 comprising an aqueous dispersion of Eudragit® NE30D, talc and optionally simethicone 15 emulsion (antifoam) is produced with gentle stirring to provide the aqueous suspension for spraying during granulation. Fluid bed granulation 18 is carried out as is well known in the art. The granules which are produced are mixed with colloidal silicon dioxide 20, as a glidant, to improve the flow quality and sieved 22 through a screen having a suitable mesh size (e.g., 12 - 20 mesh), blended 24 in a conventional e.g., twin-shell blender with anhydrous lactose 26 and magnesium stearate (lubricant) 28, conventionally compressed 30 into tablets, and coated 32 with a rapidly dissolving aqueous soluble polymeric film material such as hydroxypropyl methylcellulose polymer, as is well known in the art. The preferred embodiment of the method is exemplified in Example 1, below.

30 In another of its aspects the present invention entails a method for treating a patient infected with a microorganism which is susceptible to metronidazole, which method comprises administering to a patient having the infection a composition of the invention 35 once daily for a number of days sufficient to eliminate or resolve an infection caused by the microorganisms.

Metronidazole is known to be effective for killing such microorganisms as *Trichomonas vaginalis*, *Entamoeba histolytica*, *Helicobacter* species, such as *Helicobacter pylori*, anaerobic gram-negative Bacilli, including
5 *Bacteroides fragilis*, *Bacteroides distasonis*,
Bacteroides ovatus, *Bacteroides thetaiotaomicron*,
Bacteroides vulgatus, *Fusobacterium* species, anaerobic gram-positive bacilli including *Clostridium* species and susceptible strains of *Eubacterium*, anaerobic gram-
10 positive cocci including *Peptococcus* species and *Peptostreptococcus* species, and the like.

Among the diseases or conditions which may be treated with once daily administration of the compositions of the present invention are
15 trichomoniasis, bacterial vaginosis and duodenal ulcer disease which is associated with the presence of *Helicobacter pylori*. With respect to *H. pylori*, it is well known to administer metronidazole in combination with other active ingredients such as bismuth and
20 amoxicillin or bismuth and tetracycline. Chiba et al., *Am. J. Gastroenterology*, 87, 1716-1727 (1992); E. Herschel et al., *New England J. Med.*, 328, 308-312 (1993).

A therapeutically effective regimen for treating
25 an infection caused by a microorganism which is susceptible to metronidazole entails once daily oral administration of a composition of the invention which has between about 600 mg metronidazole and about 1000 mg metronidazole, preferably about 750 mg metronidazole
30 for a period of between about 5 days and about 15 days or more. The clinical resolution of the infection or disease is readily determined by a clinician of ordinary skill in the art, such as by microbiological testing or disappearance of clinically characteristic
35 symptoms. The dosage of metronidazole given and/or the length of treatment may be increased or decreased based on the type of infection, the degree of susceptibility

of the microorganism to metronidazole, the age and general health of the patient, and like factors of which a clinician of ordinary skill in the art is aware and utilizes in the management of a patient.

5 The following non-limiting Examples are given by way of illustration.

EXAMPLE 1

10 PREPARATION OF MODIFIED RELEASE METRONIZADOLE COMPOSITION

This example demonstrates a method for preparing a once-daily metronidazole tablet of the invention, employing fluid bed granulation.

15 185.13 kg metronidazole (Farchemia, S.P.A., Treviglio, Italy) was milled using a Fitz-Mill fitted with a 40 mesh screen at medium speed, impact forward. The milled metronidazole and 20.55 kg of hydrous lactose (Sheffield, Corp., Norwich, NY) were combined
20 in a Vector Freund, Model # FLF-200 fluid bed granulator (Marion, IA), fitted with a single spray arm having 3 nozzles for spraying the coating suspension.

The coating suspension was prepared by combining the following in a mixing tank equipped with a suitable
25 propeller mixer and fluid recycling capability: (i) 20.58 kg of Eudragit® NE30D (30% (w/v) aqueous dispersion) (Röhm Pharma, Weiterstadt, Germany), (ii) 0.08 kg of Simethicone Emulsion (Dow Corning, Midland, MI), and (iii) 100 ml of water. After the Eudragit®
30 containing suspension had stirred for several minutes, 3.07 kg of Alphafill 500 Talc (Luzenac America, Englewood, CO) was slowly added with mixing over a period of 30 minutes to ensure wetting of the talc particles, importantly, without imparting substantial
35 shear to the Eudragit® polymer. The mixture was mixed and recirculated for an additional 60 minutes to provide a uniform suspension. Alternatively, the above-mentioned mixing tank may be fitted with an Arde

Dispershear apparatus (Arde Barinco, Norwood, NJ), which, in combination with fluid recirculation, should likewise suspend the talc without imparting substantial shear.

5 The granulator, loaded with the metronidazole and lactose, was run until the inlet air temperature stabilized at 40° C (about 5 minutes). Once the inlet air temperature stabilized at 40° C, spraying of the coating suspension was begun with the spray gun
10 operated at an atomizing pressure of 50-55 psig and a spray rate of about 600 grams per minute. The coating suspension was maintained with continuous stirring throughout the process. Inlet air temperature for the fluidization air was maintained at approximately 40° C
15 throughout the granulation process.

 When the entire coating suspension had been sprayed, the pump was stopped and fluidization was continued to reduce the moisture content of the granulation to less than 1%. Then the granulator was
20 stopped, 0.25 kg colloidal silicon dioxide (Aerosil 200, Degussa Corporation, Ridgefield Park, NJ) was added and mixed in by hand. The granulator was restarted and fluidization was continued for an additional 30 seconds to effect mixing.

25 The granulation was removed from the granulator and sieved by sifting it through a Sprout-Waldron sifter fitted with a 14 mesh screen. The sieved granulation was transferred into a Patterson-Kelly V-blender (30 cubic foot capacity), combined with 32.25
30 kg anhydrous lactose (direct tablet grade) (Sheffield Corp., Norwich, NY), and the mixture was blended for 10 minutes. Then 2.50 kg of magnesium stearate (Mallinckrodt Chemical, St. Louis, MO) was added to the mixture and blended for 5 minutes to provide a
35 metronidazole containing composition suitable for tableting.

The metronidazole-containing formulation was transferred to the hopper of a Stokes 454 (Stokes-Merrill, Bristol, PA) 35-station rotary tablet press which was fitted with 0.750" X 0.375" oval shape tablet tooling. The formulation was compressed into tablets at a theoretical weight of 1,013 mg to a thickness of 0.255" and a hardness of about 11 kilopounds.

The compressed tablets were then coated with Opadry. Opadry coating suspension was prepared by mixing 9.60 kg of Opadry Blue YS-1-4256 with 54.4 kg purified water for 30 minutes. The compressed tablets were loaded into a Vector Hi-Coater (Vector Corp., Marion, IA) and sprayed with the Opadry suspension at a rate of about 500 g/minute to produce the coated, metronidazole tablets.

EXAMPLE 2

BIOAVAILABILITY OF MODIFIED RELEASE METRONIDAZOLE COMPOSITIONS

This example demonstrates the ability of defined formulations comprising 750 mg of metronidazole to provide, with a single daily dose, bioavailability of metronidazole over a 24-hour period, which is comparable to the bioavailability provided by thrice daily doses of immediate release metronidazole (Flagyl®, 250 mg, G. D. Searle & Co., Skokie, IL) as determined by the area under the curve (AUC). This example describes two related *in vivo* studies which measured the plasma metronidazole levels achieved after oral administration of various modified-release formulation. The modified release formulations were made essentially as described in Example 1.

In one *in vivo* study, nine (9) healthy female volunteers were divided into 3 groups of 3. Three compositions were tested: (i) a modified release composition comprising ethylcellulose (Formulation A), (ii) a modified release composition comprising

Eudragit® NE30D (Formulation B), and (iii) Flagyl®, an immediate release metronidazole composition. Plasma metronidazole levels were determined over a 36-hour period, after a single oral administration of the
5 respective compositions. Each individual was administered each of the three formulations. The study was completed over three consecutive weekends to provide a one-week washout period between administrations.

10 In the other related *in vivo* study, the protocol was essentially as the above-described study, except as follows. Twelve (12) healthy female volunteers were divided into four groups of three; two Eudragit®-
15 containing formulations (C and D) and two ethylcellulose formulations (E and F) were tested; all individuals received Flagyl (250 mg) and either the two Eudragit®-containing formulations or the two Surelease®-containing formulations but there was no crossover between the Eudragit®-containing and
20 ethylcellulose-containing formulations. Because of the linear bioavailability kinetics exhibited with Flagyl, the plasma metronidazole levels attained with a 250 mg dose were statistically scaled to a 750 mg dose.

In the *in vivo* studies, following administration
25 of a formulation, individual plasma metronidazole levels were measured at predetermined times utilizing a validated assay method employing a Supelco-LC-18 high performance liquid chromatography column with UV spectrophotometry at 320 nm being used to detect the
30 concentration of metronidazole. A commercially available software program, PCNONLIN (ClinTrials, Inc., Lexington, KT 40504), was used to evaluate the plasma concentration values derived at each timepoint and determine the area under the curve (AUC) and maximum
35 concentration (C_{max}) afforded by each of the formulations.

The formulations which were administered are as follows:

<u>FORMULATION A</u>		
	mg/tablet	% w/w
5		
	750.0	80.95
	83.3	8.99
10	83.3	8.99
	0.7	0.08
	<u>9.2</u>	0.99
	926.5	
15		
<u>FORMULATION B</u>		
	mg/tablet	% w/w
	750.0	82.79
20	83.3	9.20
	41.7	4.60
	20.8	2.30
25	0.3	0.03
	0.8	0.09
	<u>9.0</u>	0.99
	905.9	
30		
<u>FORMULATION C</u>		
	mg/tablet	% w/w
	750.0	74.05
35	83.3	8.22
	25.0	2.47
	12.5	1.23
40	0.3	0.03
	130.7	12.90
	1.0	0.10
	<u>10.0</u>	0.99
	1012.8	

FORMULATION D

	mg/tablet	% w/w
5		
Metronidazole USP (20% of Metronidazole added neat)	750.0	72.99
Lactose, Hydrrous NF	66.6	6.48
Eudragit NE30D Solids (Acrylic Resin, 30% Solids)	20.0	1.95
Talc USP (500 mesh)	10.0	0.97
10		
Simethicone Emulsion USP (Antifoam C Emulsion)	0.3	0.03
Lactose, Anhydrous (DTG) NF	169.4	16.49
Colloidal Silicon Dioxide NF	1.0	0.10
Magnesium Stearate NF	<u>10.2</u>	0.99
15	1027.5	

FORMULATION E

	mg/tablet	% w/w
20		
Metronidazole USP (20% of Metronidazole added neat)	750.0	71.10
Microcrystalline Cellulose, PH101 NF Surelease Solids	66.6	6.31
25		
(Ethylcellulose Dispersion, 25% Solids)	53.3	5.05
Simethicone Emulsion USP (Antifoam C Emulsion)	0.5	0.05
Microcrystalline Cellulose, PH102 NF	174.0	16.50
Magnesium Stearate NF	<u>10.4</u>	0.99
30	1054.8	

FORMULATION F

	mg/tablet	% w/w
35		
Metronidazole USP	750.0	71.70
Microcrystalline Cellulose, PH101 NF Surelease Solids	83.3	7.96
(Ethylcellulose Dispersion, 25% Solids)	66.7	6.38
40		
Simethicone Emulsion USP (Antifoam C Emulsion)	0.6	0.06
Microcrystalline Cellulose, PH102 NF	135.0	12.91
Magnesium Stearate NF	<u>10.4</u>	0.99
45	1046.0	

The results of the studies are depicted graphically in Figures 2-4.

Formulations C and D, which comprise 2.47% and 1.95% Eudragit® NE30, respectively, provided bioavailability of metronidazole over 24 hours which was substantially equivalent to that of the immediate release Flagyl®, as determined by the PCNONLIN program. The ratio of $AUC_{(\text{formulation C})} : AUC_{(\text{Flagyl}^{\circ})}$ was 0.961, and the ratio of $AUC_{(\text{formulation D})} : AUC_{(\text{Flagyl}^{\circ})}$ was 1.01. Also these compositions exhibited desirable modified release characteristics; plasma metronidazole levels were maintained between about 2.5 mcg/ml and about 13.5 mcg/ml over the entire 24-hour period (Figure 2).

As Figure 3 demonstrates, the ethylcellulose-containing compositions, Formulations E and F, exhibited poor release characteristics. While the bioavailability of the ethylcellulose-containing compositions was similar to Flagyl® --the ratio of $AUC_{(\text{formulation E})} : AUC_{(\text{Flagyl}^{\circ})}$ was 1.07, and the ratio of $AUC_{(\text{formulation F})} : AUC_{(\text{Flagyl}^{\circ})}$ was 1.02-- the composition of Formulations E and F exhibited substantially no modified release characteristics.

Figure 4 demonstrates that Formulation A (comprising ~9% Surelease®) and Formulation B (comprising 4.6% Eudragit® NE30D) were able to slow the release of metronidazole, but only at the cost of greatly reducing the bioavailability of metronidazole. The ratio of $AUC_{(\text{formulation A})} : AUC_{(\text{Flagyl}^{\circ})}$ was 0.357, and the ratio of $AUC_{(\text{formulation B})} : AUC_{(\text{Flagyl}^{\circ})}$ was 0.397. Thus, these compositions were able to provide only about 35%-40% bioavailability of metronidazole, compared to Flagyl®.

In view of the above it will be seen that the compositions of the invention importantly provide bioavailability of metronidazole which is comparable to Flagyl®, while at the same time exhibiting a controlled rate of release so that plasma metronidazole levels are

Flagyl[®], while at the same time exhibiting a controlled rate of release so that plasma metronidazole levels are not caused to spike, but rather are maintained at levels of between about 2.5 mcg/ml and 13.5 mcg/ml over a 24 hour period, which provides sustained therapeutic effectiveness.

EXAMPLE 3

10 IN VITRO DISSOLUTION OF MODIFIED RELEASE METRONIDAZOLE COMPOSITIONS

This example demonstrates the rate of *in vitro* dissolution of modified release metronidazole tablets comprising poly(meth)acrylic acid ester. The dissolution rates for the compositions of the invention correlate to the *in vivo* bioavailability of metronidazole formulations comprising poly(meth)acrylic acid ester.

20 The dissolution tests were performed using the standard USP XXII Dissolution Apparatus 2 (paddles), run at 50 RPM, in Deionized Water at 37°C for 12 hours. Five modified release formulations containing 750 mg metronidazole, and an immediate release formulation
25 (Flagyl[®], 3 x 250 mg.) as a control were tested. Three of the modified release formulations were Formulation B, Formulation C and Formulation D which are disclosed in Example 2, above. The other two modified release formulation are as follows:

30

FORMULATION G

		mg/tablet	% w/w
5	Metronidazole USP (10% of metronidazole added neat)	750.0	78.48
	Lactose, Hydrous NF	75.0	7.85
	Eudragit NE30D Solids (Acrylic Resin, 30% Solids)	22.5	2.35
10	Talc USP (500 mesh)	11.3	1.18
	Simethicone Emulsion USP (Antifoam C Emulsion)	0.3	0.03
	Lactose, Anhydrous (DTG) NF	86.0	9.00
	Colloidal Silicon Dioxide NF	1.0	0.10
15	Magnesium Stearate NF	9.5	0.99
		955.6	

FORMULATION H

		mg/tablet	% w/w
20	Metronidazole USP	750.0	77.42
	Lactose, Hydrous NF	83.3	8.60
	Eudragit NE30D Solids (Acrylic Resin, 30% Solids)	25.0	2.58
25	Talc USP (500 mesh)	12.5	1.29
	Simethicone Emulsion USP (Antifoam C Emulsion)	0.3	0.03
	Lactose, Anhydrous (DTG) NF	87.1	8.99
30	Colloidal Silicon Dioxide NF	1.0	0.10
	Magnesium Stearate NF	9.6	0.99
		968.7	

35 The dissolution data are depicted graphically in Figure
5. As a comparison Figure 5 to Figures 2-4
demonstrates, desirable in vitro release rates for
modified release metronidazole compositions of the
invention, as compared to immediate release
40 metronidazole (Flagyl®), correlated well to in vivo
bioavailability.

While the present invention has been described in
some detail by way of illustration and examples,

modifications which may be apparent to those having ordinary skill in the art are intended to be within the scope of the following claims.

We Claim:

1. A pharmaceutical composition which is capable of being compressed into a modified release, once daily dosage form of metronidazole, the composition comprising:

(a) a first portion of metronidazole which is about 59 wt% to about 79 wt% metronidazole;

(b) about 1.5 wt% to about 3.0 wt% of an aqueous insoluble poly(meth)acrylic acid ester copolymer which is aqueous permeable, aqueous expandable and pH-independent;

(c) about 0.1 wt% to about 2.0 wt% detackifier;

(d) 0 to about 23 wt% of a first aqueous soluble pharmaceutical diluent;

(e) 0 to about 23 wt% of a second aqueous soluble diluent which is suitable for forming a pharmaceutical tablet, the second aqueous soluble diluent being the same as or different from the first aqueous soluble diluent;

(f) 0 to about 20 wt% of a second portion of metronidazole;

(g) 0 to about 0.2 wt% glidant; and

(h) 0 to about 2 wt% lubricant;

wherein the composition comprises metronidazole containing granules comprising (a), (b), (c) and (d), wherein the sum of the weight percentages of metronidazole provided by (a) and (f) is between about 72 wt% and about 79 wt%, and wherein the sum of the weight percentages of the aqueous soluble diluent provided by (d) and (e) is between about 16 wt% and about 23 wt%.

2. A pharmaceutical composition according to Claim 1, wherein the first aqueous soluble diluent is present in an amount of from about 5% and about 11% by weight and the second aqueous soluble diluent is present in an amount of from about 8% and about 18% by weight.

3. A pharmaceutical composition according to Claim 2, wherein the first aqueous soluble diluent and the second aqueous soluble diluent are lactose.

4. A pharmaceutical composition according to Claim 1, which comprises at least about 0.01% of a pharmaceutically acceptable glidant and at least about 0.1% of a pharmaceutically acceptable lubricant.

5. A pharmaceutical composition according to Claim 1 comprising (i) from about 1.9% to about 2.5% of the aqueous insoluble poly(meth)acrylic acid ester copolymer; (ii) from about 6% to about 9% of the first aqueous soluble diluent; and (iii) from about 12% to about 17% of the second aqueous soluble diluent; and wherein the sum of the percentage of the first portion of metronidazole and the second portion of metronidazole is from about 72.5% to about 74.5%.

6. A pharmaceutical composition according to Claim 1 wherein the detackifying agent is talc.

7. A pharmaceutical composition according to Claim 1 which comprises about 74.05% metronidazole, about 8.22% hydrous lactose, about 2.47% of the aqueous insoluble poly(meth)acrylic acid ester copolymer, about 1.23% talc, about 0.03% simethicone

emulsion, about 12.90% anhydrous lactose, about 0.10% colloidal silicon dioxide, and about 0.99% magnesium stearate.

8. A pharmaceutical composition according to Claim 1 which comprises about 72.99% metronidazole, about 6.48% hydrous lactose, about 1.95% of the aqueous insoluble poly(meth)acrylic acid ester copolymer, about 0.97% talc, about 0.03% simethicone emulsion, about 16.49% anhydrous lactose, about 0.10% colloidal silicon dioxide, and about 0.99% magnesium stearate.

9. A method for making a modified release metronidazole composition, the method comprising the steps of:

(a) contacting in a fluid bed granulator, under conditions suitable for producing granules, (1) a dry mixture comprising from about 59 to about 79 parts by weight of a first portion of metronidazole and optionally up to 23 parts by weight of a first aqueous soluble diluent with (2) an aqueous suspension comprising from about 1.5 to about 3.0 parts by weight of an aqueous insoluble, pH-independent, aqueous expandable polymethacrylic acid ester copolymer, and an effective amount of a detackifying agent;

(b) combining the granules produced in step (a) with an effective amount of a pharmaceutically acceptable glidant;

(c) if necessary, particle-sizing the mixture of (b) to provide a mixture with a substantially uniform particle size suitable for compressing into tablet form;

(d) optionally blending the mixture with up to about 20 parts by weight of a second portion of metronidazole;

(e) blending the mixture with up to 23 parts by weight of a second aqueous soluble diluent and an effective amount of a pharmaceutically acceptable lubricant; and

(f) compressing a predetermined amount of the blended mixture of step (e) to produce a tablet;

wherein the percentage of metronidazole provided by (a)(1) and (d) is from about 72% to about 79% by weight, based on the total weight of the tablet, and wherein the percentage of aqueous soluble diluent provided by (a)(1) and (e) is from about 16% and about 23% by weight, based on the total weight of the tablet.

10. A method according to Claim 9, wherein the first aqueous soluble diluent is present in an amount of from about 5% and about 11% by weight and the second aqueous soluble diluent is present in an amount of from about 8% and about 18% by weight.

11. A method according to Claim 10, wherein the first aqueous soluble diluent is present in an amount of from about 6% and about 9% by weight and the second aqueous soluble diluent is present in an amount of from about 12% and about 17% by weight.

12. A method according to Claim 10, wherein the first aqueous soluble diluent and the second aqueous soluble diluent are lactose.

13. A method according to Claim 9 wherein the detackifying agent is talc.

14. A method according to Claim 9 wherein the glidant is silicon dioxide and is present in amount of about 0.1% by weight.

15. A method according to Claim 9 wherein the lubricant is magnesium stearate and is present in amount of about 1% by weight.

16. A method according to Claim 9 wherein the aqueous suspension of (a)(ii) further comprises an amount of an antifoam agent effective to reduce foaming during step (a).

17. A method according to Claim 9 which further comprises (g) coating the tablet with an aqueous soluble polymeric coating material.

~~18~~. A modified release pharmaceutical tablet for once daily dosing of metronidazole, the tablet comprising a therapeutically effective amount of a pharmaceutical composition comprising:

(a) a first portion of metronidazole which is about 59 wt% to about 79 wt% metronidazole;

(b) about 1.5 wt% to about 3.0 wt% of an aqueous insoluble poly(meth)acrylic acid ester copolymer which is aqueous permeable, aqueous expandable and pH-independent;

(c) about 0.1 wt% to about 2.0 wt% detackifier;

(d) 0 to about 23 wt% of a first aqueous soluble pharmaceutical diluent;

(e) 0 to about 23 wt% of a second aqueous soluble diluent which is suitable for forming a pharmaceutical tablet, the second aqueous soluble diluent being the same as or different from the first aqueous soluble diluent;

(f) 0 to about 20 wt% of a second portion of metronidazole;

(g) 0 to about 0.2 wt% glidant; and

(h) 0 to about 2 wt% lubricant;

wherein the composition comprises metronidazole containing granules comprising (a), (b), (c) and (d), wherein the sum of the weight percentages of metronidazole provided by (a) and (f) is between about 72 wt% and about 79 wt%, and wherein the sum of the weight percentages of the aqueous soluble diluent provided by (d) and (e) is between about 16 wt% and about 23 wt%.

19. A modified release pharmaceutical tablet according to Claim 18, wherein the first aqueous soluble diluent is present in an amount of from about 5% and about 11% by weight and the second aqueous soluble diluent is present in an amount of from about 8% and about 18% by weight.

20. A modified release pharmaceutical tablet according to Claim 19, wherein the aqueous insoluble poly(meth)acrylic acid ester copolymer is Eudragit NE30.

21. A modified release pharmaceutical tablet according to Claim 20, wherein the first aqueous soluble diluent and the second aqueous soluble diluent are lactose.

22. A modified release pharmaceutical tablet according to Claim 21 which comprises at least about 0.01% of a pharmaceutically acceptable glidant and at least about 0.1% of a pharmaceutically acceptable lubricant.

23. A modified release pharmaceutical tablet according to Claim 18 which comprises about 750 milligrams of metronidazole.

24. A modified release pharmaceutical tablet according to Claim 18 comprising (i) from about 1.9% to about 2.5% of the aqueous insoluble poly(meth)acrylic acid ester copolymer; (ii) from about 6% to about 9% of the first aqueous soluble diluent;

and (iii) from about 12% to about 17% of the second aqueous soluble diluent; and wherein the sum of the percentage of the first portion of metronidazole and the second portion of metronidazole is from about 72.5% to about 74.5%.

25. A modified release pharmaceutical tablet according to Claim 24, wherein the aqueous insoluble poly(meth)acrylic acid ester copolymer is Eudragit NE30.

26. A modified release pharmaceutical tablet according to Claim 25 which comprises about 750 milligrams of metronidazole.

27. A modified release pharmaceutical tablet according to Claim 18 which comprises about 74.05% metronidazole, about 8.22% hydrous lactose, about 2.47% of the aqueous insoluble poly(meth)acrylic acid ester copolymer, about 1.23% talc, about 0.03% simethicone emulsion, about 12.90% anhydrous lactose, about 0.10% colloidal silicon dioxide, and about 0.99% magnesium stearate.

28. A modified release pharmaceutical tablet according to Claim 18 which comprises about 72.99% metronidazole, about 6.48% hydrous lactose, about 1.95% of the aqueous insoluble poly(meth)acrylic acid ester copolymer, about 0.97% talc, about 0.03% simethicone emulsion, about 16.49% anhydrous lactose, about 0.10% colloidal silicon dioxide, and about 0.99% magnesium stearate.

29. A modified release pharmaceutical tablet according to Claim 27, wherein the aqueous insoluble poly(meth)acrylic acid ester copolymer is Eudragit NE30.

30. A modified release pharmaceutical tablet according to Claim 28, wherein the aqueous insoluble poly(meth)acrylic acid ester copolymer is Eudragit NE30.

2181985

31. A modified release pharmaceutical tablet according to Claim 29 which comprises about 750 milligrams of metronidazole.

32. A modified release pharmaceutical tablet according to Claim 30 which comprises about 750 milligrams of metronidazole.

33. A modified release pharmaceutical tablet according to Claim 29, wherein the tablet is coated with an aqueous soluble polymeric coating material.

34. A modified release pharmaceutical tablet according to Claim 30, wherein the tablet is coated with an aqueous soluble polymeric coating material.

35. A use of the pharmaceutical tablet of claim 18 for treating a patient infected with a microorganism which is susceptible to metronidazole.

36. The use according to Claim 35 wherein the microorganism causes a disease which is selected from the group consisting of trichomoniasis, amoebiasis, and bacterial vaginosis.

37. The use according to Claim 35 wherein the microorganism is a species of *Helicobacter*.

38. The use according to Claim 37, wherein the microorganism is *Helicobacter pylori*.

39. The use according to Claim 35, wherein the pharmaceutical tablet comprises about 74.05% metronidazole, about 8.22% hydrous lactose, about 2.47% of the aqueous insoluble poly(meth)acrylic acid ester copolymer, about 1.23% talc, about 0.03% simethicone emulsion, about 12.90% anhydrous lactose, about 0.10% colloidal silicon dioxide, and about 0.99% magnesium stearate.

40. The use according to Claim 35, wherein the pharmaceutical tablet comprises about 72.99% metronidazole, about 6.48% hydrous lactose, about 1.95% of the aqueous insoluble poly(meth)acrylic acid ester copolymer, about 0.97% talc, about 0.03% simethicone emulsion, about 16.49% anhydrous lactose, about 0.10% colloidal silicon dioxide, and about 0.99% magnesium stearate.

41. The use according to Claim 39, wherein the aqueous insoluble poly(meth)acrylic acid ester copolymer is Eudragit NE30.

42. The use according to Claim 40, wherein the aqueous insoluble poly(meth)acrylic acid ester copolymer is Eudragit NE30.

43. The use according to Claim 41, wherein the pharmaceutical tablet comprises about 750 milligrams of metronidazole.

44. The use according to Claim 42, wherein the pharmaceutical tablet comprises about 750 milligrams of metronidazole.




Figure 1




FIG. 2




FIG. 3




FIG. 4




FIG. 5