

**(12) STANDARD PATENT
(19) AUSTRALIAN PATENT OFFICE**

(11) Application No. AU 2006316838 B2

(54) Title
Production of glycoproteins with reduced O-glycosylation

(51) International Patent Classification(s)
C12N 1/15 (2006.01) **C12N 1/16** (2006.01)
A61K 38/00 (2006.01) **C12N 1/19** (2006.01)

(21) Application No: **2006316838** (22) Date of Filing: **2006.11.10**

(87) WIPO No: **WO07/061631**

(30) Priority Data

(31) Number
60/737,108 (32) Date
2005.11.15 (33) Country
US

(43) Publication Date: **2007.05.31**
(44) Accepted Journal Date: **2012.04.12**

(71) Applicant(s)
GlycoFi, Inc

(72) Inventor(s)
Cook, James W.;Bobrowicz, Piotr;Kett, Warren

(74) Agent / Attorney
Spruson & Ferguson, Level 35 St Martins Tower 31 Market Street, Sydney, NSW, 2000

(56) Related Art
ENEYSKAYA, ELENA V. et al. Biochemical and Biophysical Research Communications. 1998. Vol. 245, pages 43-49
US 5,714,377 A
ORCHARD, MICHAEL G. et al. Bioorganic and Medicinal Chemistry Letters. 2004.
Vol. 14, pages 3975-3978
US 2002/0068325 A1

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
31 May 2007 (31.05.2007)

PCT

(10) International Publication Number
WO 2007/061631 A2

(51) International Patent Classification:
CI2P 21/06 (2006.01) *C07K 14/705* (2006.01)

(21) International Application Number:
PCT/US2006/043535

(22) International Filing Date:
10 November 2006 (10.11.2006)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/737,108 22 November 2005 (22.11.2005) US

(71) Applicant (for all designated States except US): **GLYCOFI, INC** [US/US]; Suite 200, 21 Lafayette Street, Lebanon, NH 03766 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **BOBROWICZ, Piotr** [PL/US]; Suite 200, 21 Lafayette Street, Lebanon, NH 03766 (US). **COOK, James, W.** [US/US]; Suite 200, 21 Lafayette Street, Lebanon, NH 03766 (US). **KETT, Warren** [AU/US]; Suite 200, 21 Lafayette Street, Lebanon, NH 03766 (US).

(74) Agent: **REILLY, John, David**; Merck & Co., Inc., 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.


(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: PRODUCTION OF GLYCOPROTEINS WITH REDUCED O-GLYCOSYLATION



A2

(57) Abstract: A method is described for producing protein compositions having reduced amounts of O-linked glycosylation. The method includes producing the protein in cells cultured in the presence of an inhibitor of Pmt-mediated O-linked glycosylation and/or in the presence of one or more α -1,2-mannosidases.

TITLE OF THE INVENTION

PRODUCTION OF GLYCOPROTEINS WITH REDUCED O-GLYCOSYLATION

BACKGROUND OF THE INVENTION

5 (1) Field of the Invention
The present invention relates to compositions and methods for producing proteins having specific glycosylation patterns. In particular, the present invention relates to compositions and methods for producing proteins having reduced O-linked glycosylation.

10 (2) Description of Related Art
Glycoproteins mediate many essential functions in humans and other mammals, including catalysis, signaling, cell-cell communication, and molecular recognition and association. Glycoproteins make up the majority of non-cytosolic proteins in eukaryotic organisms (Lis and Sharon, 1993, Eur. J. Biochem. 218:1-27). Many glycoproteins have been exploited for therapeutic purposes, and 15 during the last two decades, recombinant versions of naturally-occurring glycoproteins have been a major part of the biotechnology industry. Examples of recombinant glycosylated proteins used as therapeutics include erythropoietin (EPO), therapeutic monoclonal antibodies (mAbs), tissue plasminogen activator (tPA), interferon- β (IFN- β), granulocyte-macrophage colony stimulating factor (GM-CSF), and human chorionic gonadotrophin (hCH) (Cumming *et al.*, 1991, Glycobiology 1:115-130). Variations in 20 glycosylation patterns of recombinantly produced glycoproteins have recently been the topic of much attention in the scientific community as recombinant proteins produced as potential prophylactics and therapeutics approach the clinic.

In general, the glycosylation structures of glycoprotein oligosaccharides will vary depending upon the host species of the cells used to produce them. Therapeutic proteins produced in 25 non-human host cells are likely to contain non-human glycosylation which may elicit an immunogenic response in humans—e.g. hypermannosylation in yeast (Ballou, 1990, Methods Enzymol. 185:440-470); α (1,3)-fucose and β (1,2)-xylose in plants, (Cabanes-Macheteau *et al.*, 1999, Glycobiology, 9: 365-372); N-glycolylneuraminic acid in Chinese hamster ovary cells (Noguchi *et al.*, 1995, J. Biochem. 117: 5-62); and, Gal α -1,3Gal glycosylation in mice (Borrebaeck, *et al.*, 1993, Immun. Today, 14: 477-479). 30 Carbohydrate chains bound to proteins in animal cells include N-glycoside bond type carbohydrate chains (also called N-glycans; or N-linked glycosylation) bound to an asparagine (Asn) residue in the protein and O-glycoside bond type carbohydrate chains (also called O-glycans; or O-linked glycosylation) bound to a serine (Ser) or threonine (Thr) residue in the protein.

Because the oligosaccharide structures of glycoproteins produced by non-human 35 mammalian cells tend to be more closely related to those of human glycoproteins, most commercial glycoproteins are produced in mammalian cells. However, mammalian cells have several important disadvantages as host cells for protein production. Besides being costly, processes for producing

proteins in mammalian cells produce heterogeneous populations of glycoforms, have low volumetric titers, and require both ongoing viral containment and significant time to generate stable cell lines.

It is well recognized that the particular glycoforms on a protein can profoundly affect the properties of the protein, including its pharmacokinetic, pharmacodynamic, receptor-interaction, and tissue-specific targeting properties (Graddis *et al.*, 2002. *Curr Pharm Biotechnol.* 3: 285-297). For example, it has been shown that different glycosylation patterns of IgGs are associated with different biological properties (Jefferis and Lund, 1997, *Antibody Eng. Chem. Immunol.*, 65: 111-128; Wright and Morrison, 1997, *Trends Biotechnol.*, 15: 26-32). It has further been shown that galactosylation of a glycoprotein can vary with cell culture conditions, which may render some glycoprotein compositions immunogenic depending on the specific galactose pattern on the glycoprotein (Patel *et al.*, 1992. *Biochem J.* 285: 839-845). However, because it is not known which specific glycoform(s) contribute(s) to a desired biological function, the ability to enrich for specific glycoforms on glycoproteins is highly desirable. Because different glycoforms are associated with different biological properties, the ability to enrich for glycoproteins having a specific glycoform can be used to elucidate the relationship between a specific glycoform and a specific biological function of the glycoprotein. Also, the ability to enrich for glycoproteins having a specific glycoform enables the production of therapeutic glycoproteins having particular specificities. Thus, production of glycoprotein compositions that are enriched for particular glycoforms is highly desirable.

While the pathway for N-linked glycosylation has been the subject of much analysis, the process and function of O-linked glycosylation is not as well understood. However, it is known that in contrast to N-linked glycosylation, O-glycosylation is a posttranslational event, which occurs in the *cis*-Golgi (Varki, 1993, *Glycobiol.*, 3: 97-130). While a consensus acceptor sequence for O-linked glycosylation like that for N-linked glycosylation does not appear to exist, a comparison of amino acid sequences around a large number of O-linked glycosylation sites of several glycoproteins show an increased frequency of proline residues at positions -1 and +3 relative to the glycosylated residues and a marked increase of serine, threonine, and alanine residues (Wilson *et al.*, 1991, *Biochem. J.*, 275: 529-534). Stretches of serine and threonine residues in glycoproteins, may also be potential sites for O-glycosylation.

One gene family that has a role in O-linked glycosylation are the genes encoding the Dol-P-Man:Protein (Ser/Thr) Mannosyl Transferase (Pmt). These highly conserved genes have been identified in both higher eukaryotes such as humans, rodents, insects, and the like and lower eukaryotes such as fungi and the like. Yeast such as *Saccharomyces cerevisiae* and *Pichia pastoris* encode up to seven *PMT* genes encoding Pmt homologues (reviewed in Willer *et al.* *Curr. Opin. Struct. Biol.* 2003 Oct;13(5): 621-30.). In yeast, O-linked glycosylation starts by the addition of the initial mannose from dolichol-phosphate mannose to a serine or threonine residue of a nascent glycoprotein in the endoplasmic reticulum by one of the seven O-mannosyl transferases genes. While there appear to be seven *PMT* genes encoding Pmt homologues in yeast, O-mannosylation of secreted fungal and heterologous proteins

in yeast is primarily dependent on the genes encoding Pmt1 and Pmt2, which appear to function as a heterodimer. *PMT1* and *PMT2* and their protein products, Pmt1 and Pmt2, respectively, appear to be highly conserved among species.

5 Tanner *et al.* in U. S. Patent No. 5,714,377 describes the *PMT1* and *PMT2* genes of *Saccharomyces cerevisiae* and a method for making recombinant proteins having reduced O-linked glycosylation that uses fungal cells in which one or more of *PMT* genes have been genetically modified so that recombinant proteins are produced, which have reduced O-linked glycosylation.

10 Ng *et al.* in U.S. Published Patent Application No. 20020068325 discloses inhibition of O-glycosylation through the use of antisense or cosuppression or through the engineering of yeast host strains that have loss of function mutations in genes associated with O-linked glycosylation, in particular, one or more of the *PMT* genes.

15 UDP-N-acetyl-alpha-D-galactosamine:polypeptide N-acetyl galactosaminyl-transferases (GalNAc-transferases) are involved in mucin type O-linked glycosylation found in higher eukaryotes. These enzymes initiate O-glycosylation of specific serine and threonine amino acids in proteins by adding N-acetylgalactosamine to the hydroxy group of these amino acids to which mannose residues can then be added in a step-wise manner. Clausen *et al.* in U. S. Patent No. 5,871,990 and U.S. Published Patent Application No. 20050026266 discloses a family of nucleic acids encoding UDP-N-acetyl-alpha-D-galactosamine:polypeptide N-acetyl galactosaminyl-transferases (GalNAc-transferases). Clausen in U.S. Published Patent Application No. 20030186850 discloses the use of GalNAc-beta-benzyl to 20 selectively inhibit lectins of polypeptide GalNAc-transferases and not serve as substrates for other glycosyltransferases involved in O-glycan biosyntheses, thus inhibiting O-glycosylation.

25 Inhibitors of O-linked glycosylation have been described. For example, Orchard *et al.* in U.S. Patent No. 7,105,554 describes benzylidene thiazolidinediones and their use as antimycotic agents, e.g., antifungal agents. These benzylidene thiazolidinediones are reported to inhibit the Pmt1 enzyme, preventing the formation of the O-linked mannoproteins and compromising the integrity of the fungal cell wall. The end result is cell swelling and ultimately death through rupture.

30 Konrad *et al.* in U.S. Published Patent Application No. 20020128235 disclose a method for treating or preventing diabetes mellitus by pharmacologically inhibiting O-linked protein glycosylation in a tissue or cell. The method relies on treating a diabetic individual with (Z)-1-[N-(3-Ammoniopropyl)-N-(n-propyl)amino] diazen-ium-1,2-diolate or a derivative thereof, which binds O-linked N-acetylglucosamine transferase and thereby inhibits O-linked glycosylation.

35 Kojima *et al.* in U. S. Patent No. 5,268,364 disclose therapeutic compositions for inhibition of O-glycosylation using compounds such as benzyl- α -N-acetylgalactosamine, which inhibits extension of O-glycosylation leading to accumulation of O- α -GalNAc, to block expression of SLex or SLea by leukocytes or tumor cells and thereby inhibit adhesion of these cells to endothelial cells and platelets.

Boime et al. U.S. Patent No. 6,103,501 disclose variants of hormones in which O-linked glycosylation was altered by modifying the amino acid sequence at the site of glycosylation.

The present inventors have found that particular chemical compounds that are 5 inhibitors of Pmt proteins, which are generally lethal to fungi, can be used in a way which is not lethal to the host cells for production of recombinant proteins with reduced O-linked glycosylation. This enables O-linked glycosylation of proteins produced from fungi and yeast cells to be controlled. Other classes of chemical compounds, which the inventors believe to be non-lethal inhibitors of the PMT enzymes, are also useful in the 10 production of improved glycoproteins with reduced O-linked glycosylation. The present inventors have further found that addition to the host cell or cell culture of certain classes of enzymes, namely, α -1,2-mannosidases, alone or in combination with a chemical inhibitor of Pmt proteins effects a further reduction of O-glycosylation.

BRIEF SUMMARY OF THE INVENTION

15 In a first aspect of the invention there is provided a method of producing a protein having reduced O-linked glycosylation comprising:

- (a) providing a nucleic acid encoding a glycoprotein and a second nucleic acid encoding an α 1,2-mannosidase;
- (b) introducing the nucleic acid and the second nucleic acid into a host cell and 20 growing the host cell containing the nucleic acid and the second nucleic acid to produce a culture of the host cell;
- (c) contacting the culture of the host cell with one or more inhibitors of Pmt-mediated O-linked glycosylation; and
- (d) isolating the glycoprotein produced by the host cell in the presence of the one 25 or more inhibitors of Pmt-mediated O-linked glycosylation to produce the protein having reduced O-linked glycosylation.

In a second aspect of the invention there is provided a glycoprotein composition comprising a glycoprotein having a human-like or humanized N-glycosylation pattern and reduced O-linked glycosylation.

30 The present invention provides methods for producing proteins and glycoproteins having specific glycosylation patterns. In particular, the present invention provides a method for making a recombinant protein compositions in a host cell in which the O-linked glycosylation of the recombinant protein is reduced by contacting the host cells with one or more inhibitors of Pmt-mediated O-linked glycosylation of proteins in the

host cell or contacting the host cells or the recombinant protein with one or more α -1,2-mannosidases, or both. The amount of O-linked glycosylation of the recombinant protein or glycoprotein is reduced compared to the amount of O-linked glycosylation of the recombinant protein or glycoprotein produced by the host cell in the absence of the inhibitor.

Pmt-mediated O-linked glycosylation refers to O-linked glycosylation wherein transfer of mannose residues to the serine or threonine residues of a protein is mediated by a protein-O-D-mannosyltransferase (Pmt) or homologue encoded by a *PMT* gene or its homologue. The inhibitors of Pmt-mediated O-linked glycosylation include inhibitors that inhibit any one of the homologues of the PMT genes. In a currently preferred aspect, the inhibitor inhibits at least Pmt1 and/or Pmt2 activity of fungi and yeast, or the corresponding homologue in other organisms, including but not limited to, mammals, plants, and insects.

Currently, it is preferable that the amount of O-linked glycosylation has been reduced through the use of a chemical inhibitor, for example, a chemical inhibitor encompassed by the class of chemicals called benzylidene thiazolidinediones. In particular embodiments, the chemical inhibitor is selected from the group consisting of 5-[[3,4-bis(phenylmethoxy)phenyl]methylene]-4-oxo-2-thioxo-3-thiazolidineacetic acid; 5-[[3-(1-Phenylethoxy)-4-(2-phenylethoxy)]phenyl]methylene]-4-oxo-2-thioxo-3-thiazolidineacetic acid; 3-Hydroxy-4-(2-phenylethoxy)benzaldehyde; 3-(1-Phenylethoxy)-4-(2-phenylethoxy)-benzaldehyde; 5-[[3-(1-Phenyl-2-hydroxy)ethoxy]-4-(2-phenylethoxy)]phenyl]methylene]-4-oxo-2-thioxo-3-thiazolidineacetic acid.

In further aspects, provided is a method for producing recombinant protein compositions having reduced O-linked glycosylation, which use one or more inhibitors of the Pmt proteins involved in O-linked glycosylation and/or one or more α -1,2-mannosidase enzymes to produce the protein having reduced O-linked glycosylation. Currently preferred α -1,2-mannosidases may be isolated from 5 eukaryotic cells, including mammalian and yeast cells. In currently preferred embodiments, the α -1,2-mannosidase is that produced by *Trichoderma reesei*, *Saccharomyces* sp., or *Aspergillus* sp. In other currently preferred embodiments, the α -1,2-mannosidase may be produced from a chimeric construct comprising a nucleic acid sequence encoding the catalytic domain of an α -1,2-mannosidase operatively linked to a nucleic acid sequence encoding a cellular targeting signal peptide not normally associated 10 with the catalytic domain. In other embodiments, the α -1,2-mannosidase may be separately produced and added to the cell culture, or may be produced by co-expressing the α -1,2-mannosidase with the recombinant glycoprotein.

In particular aspects of the method, the recombinant protein composition comprises a 15 glycoprotein having N-linked glycosylation wherein the recombinant glycoprotein includes at least one predominant N-glycoform and has reduced O-linked glycosylation. Therefore, further provided are glycoprotein compositions comprising a predominant species of N-glycan structure and having reduced O-linked glycosylation compared to compositions of the glycoprotein which have been produced in host cells have not been incubated in the presence of an inhibitor of Pmt-mediated O-linked glycosylation or an α -1,2-mannosidase capable of trimming more than one mannose residue from a glycans structure. In 20 particular aspects, the glycoprotein composition comprises a glycoprotein having a predominant N-glycan structure selected from the group consisting of Man₅GlcNAc₂, Man₃GlcNAc₂, GlcNAcMan₅GlcNAc₂, GlcNAcMan₃GlcNAc₂, GlcNAc₂Man₃GlcNAc₂, GalGlcNAcMan₅GlcNAc₂, Gal(GlcNAc)₂Man₅GlcNAc₂, (GalGlcNAc)₂Man₅GlcNAc₂, NANAGalGlcNAcMan₃GlcNAc₂, NANA₂Gal₂GlcNAcMan₃GlcNAc₂, and GalGlcNAcMan₃GlcNAc₂ glycoforms. An important aspect 25 of the method is that it provides for a glycoprotein composition comprising reduced O-linked glycosylation and predominantly a specific N-linked glycoform in which the recombinant glycoprotein may exhibit increased biological activity and/or decreased undesired immunogenicity relative to compositions of the same glycoprotein produced from mammalian cell culture, such as CHO cells. An additional advantage of producing the glycoprotein composition comprising reduced O-linked 30 glycosylation and a predominant N-linked glycoform is that it avoids production of undesired or inactive glycoforms and heterogeneous mixtures, which may induce undesired effects and/or dilute the more effective glycoform. Thus, therapeutic pharmaceutical composition of glycoprotein molecules comprising, for example, predominantly Man₅GlcNAc₂, Man₃GlcNAc₂, GlcNAcMan₅GlcNAc₂, GlcNAcMan₃GlcNAc₂, GlcNAc₂Man₃GlcNAc₂, GalGlcNAcMan₅GlcNAc₂, Gal(GlcNAc)₂ 35 Man₅GlcNAc₂, (GalGlcNAc)₂Man₅GlcNAc₂, NANAGalGlcNAcMan₃GlcNAc₂, NANA₂Gal₂GlcNAcMan₃GlcNAc₂, and GalGlcNAcMan₃GlcNAc₂ glycoforms and having reduced O-linked glycosylation may well be effective at lower doses, thus having higher efficacy/potency.

Therefore, provided is a method of producing a protein having reduced O-linked glycosylation comprising providing a nucleic acid encoding a protein; introducing the nucleic acid into a host cell to provide a culture of the host cell; contacting the culture with one or more inhibitors of Pmt-mediated O-linked glycosylation; and isolating the glycoprotein produced by the host cell in the presence 5 of the inhibitor to produce the protein having reduced O-linked glycosylation.

In particular aspects of the method, the culture is grown for a time sufficient to provide a multiplicity of the host cells having the nucleic acid before contacting the culture with the one or more inhibitors of Pmt-mediated O-linked glycosylation or the culture is grown in the presence of the one or more inhibitors of Pmt-mediated O-linked glycosylation at the time the culture is established.

10 In a further aspect of the method, the nucleic acid encoding the protein is operably linked to an inducible promoter. Then the culture is grown for a time sufficient to provide a multiplicity of the host cells having the nucleic acid before contacting the culture with the one or more inhibitors of Pmt-mediated O-linked glycosylation and an inducer of the promoter to induce expression of the protein and isolating the protein produced by the host cell in the presence of the one or more inhibitors and the 15 inducer to produce the protein having reduced O-linked glycosylation or the culture is contacted with an inducer of the promoter to induce expression of the protein for a time before contacting the culture with the one or more inhibitors of Pmt-mediated O-linked glycosylation and isolating the protein produced by the host cell in the presence of the inhibitor and the inducer to produce the protein having reduced O-linked glycosylation.

20 Further provided is a method of producing a protein having reduced O-linked glycosylation comprising providing a nucleic acid encoding a protein; introducing the nucleic acid into a host cell to provide a culture of the host cell; contacting the culture with one or more α -1,2-mannosidase enzymes; and isolating the protein produced by the host cell in the presence of the one or more α -1,2-mannosidase enzymes to produce the glycoprotein having reduced O-linked glycosylation.

25 In particular aspects of the method, the culture is grown for a time sufficient to provide a multiplicity of the host cells having the nucleic acid before contacting the culture with the one or more α -1,2-mannosidase enzymes cosylation. In other aspects, the culture is grown in the presence of the one or more α -1,2-mannosidase enzymes.

30 In further aspects of the method, a second nucleic acid encoding the one or more α -1,2-mannosidase enzymes is provided and introducing the second nucleic acid into the host cell. In particular aspects, a second nucleic acid encoding the one or more α -1,2-mannosidase enzymes operably linked to an inducible promoter is provided and introducing the second nucleic acid into the host cell and the culture is grown for a time sufficient to provide a multiplicity of the host cells before inducing expression of the protein and the one or more α -1,2-mannosidase enzymes to produce the protein having reduced O-linked glycosylation or expression of the protein is induced for a time before inducing expression of the 35 one or more α -1,2-mannosidase enzymes to produce the protein having reduced O-linked glycosylation or

expression of the one or more α -1,2-mannosidase enzymes is induced for a time before inducing expression of the protein to produce the protein having reduced O-linked glycosylation.

Further provided is a method of producing a protein having reduced O-linked glycosylation comprising providing a nucleic acid encoding a protein operably linked to an inducible promoter; introducing the nucleic acid into a host cell and growing the host cell containing the nucleic acid to produce a culture of the host cell; contacting the culture with one or more inhibitors of Pmt-mediated O-linked glycosylation and one or more one or more α -1,2-mannosidase enzymes; and isolating the glycoprotein produced by the host cell in the presence of the one or more inhibitors and the one or more one or more α -1,2-mannosidase enzymes to produce the protein having reduced O-linked glycosylation.

In particular aspects of the method, the culture is grown for a time sufficient to provide a multiplicity of the host cells having the nucleic acid before contacting the culture with the one or more inhibitors of Pmt-mediated O-linked glycosylation or the culture is grown in the presence of the one or more inhibitors of Pmt-mediated O-linked glycosylation at the time the culture is established.

In a further aspect of the method, the nucleic acid encoding the protein is operably linked to an inducible promoter. Then the culture is grown for a time sufficient to provide a multiplicity of the host cells having the nucleic acid before contacting the culture with the one or more inhibitors of Pmt-mediated O-linked glycosylation and an inducer of the promoter to induce expression of the protein and isolating the protein produced by the host cell in the presence of the one or more inhibitors and the inducer to produce the protein having reduced O-linked glycosylation or the culture is contacted with an inducer of the promoter to induce expression of the protein for a time before contacting the culture with the one or more inhibitors of Pmt-mediated O-linked glycosylation and isolating the protein produced by the host cell in the presence of the inhibitor and the inducer to produce the protein having reduced O-linked glycosylation.

In particular aspects of the method, the culture is grown for a time sufficient to provide a multiplicity of the host cells having the nucleic acid before contacting the culture with the one or more α -1,2-mannosidase enzymes. In other aspects, the culture is grown in the presence of the one or more α -1,2-mannosidase enzymes.

In further aspects of the method, a second nucleic acid encoding the one or more α -1,2-mannosidase enzymes is provided and introducing the second nucleic acid into the host cell. In particular aspects, a second nucleic acid encoding the one or more α -1,2-mannosidase enzymes operably linked to an inducible promoter is provided and introducing the second nucleic acid into the host cell and the culture is grown for a time sufficient to provide a multiplicity of the host cells before inducing expression of the protein and the one or more α -1,2-mannosidase enzymes to produce the protein having reduced O-linked glycosylation or expression of the protein is induced for a time before inducing expression of the one or more α -1,2-mannosidase enzymes to produce the protein having reduced O-linked glycosylation or

expression of the one or more α -1,2-mannosidase enzymes is induced for a time before inducing expression of the protein to produce the protein having reduced O-linked glycosylation.

In further aspects of the above methods that use one or more inhibitors of a Pmt protein, currently, it is preferred that the one or more inhibitors is selected from the class of molecules

5 comprising benzylidene thiazolidinediones. Currently, it is preferable that the one or more inhibitors be selected from the group consisting of 5-[[3,4-bis(phenylmethoxy) phenyl]methylene]-4-oxo-2-thioxo-3-thiazolidineacetic Acid; 5-[[3-(1-Phenylethoxy)-4-(2-phenylethoxy)]phenyl]methylene]-4-oxo-2-thioxo-3-thiazolidineacetic Acid; and 5-[[3-(1-Phenyl-2-hydroxy)ethoxy)-4-(2-phenylethoxy)]phenyl]methylene]-4-oxo-2-thioxo-3-thiazolidineacetic Acid.

10 In particular aspects of the above methods that use an α -1,2-mannosidase, it is currently preferable that the α -1,2-mannosidase is selected from the group consisting of *Trichoderma reesei*, *Saccharomyces* sp., and *Aspergillus* sp. Currently, it is preferable that the α -1,2-mannosidase is from *Trichoderma reesei*. Alternatively, the host cell can include in addition to the first nucleic acid encoding the protein or glycoprotein, a second nucleic acid, which encodes the α -1,2-mannosidase, operably linked 15 to an inducible promoter. Expression of the α -1,2-mannosidase and the protein or glycoprotein can be induced simultaneously or expression of the protein or glycoprotein induced before expression of the α -1,2-mannosidase or vice versa.

20 While the method can be performed using any host cell that produced proteins having O-linked glycosylation, in currently preferred aspects, the host cell is a lower eukaryotic cell, preferably a fungal cell or a yeast cell. Currently, it is preferred that the host cell be selected from the group consisting of cells from *K. lactis*, *Pichia pastoris*, *Pichia methanolica*, and *Hansenula*. In further embodiments for producing recombinant glycoproteins in particular, the host cell is a yeast or filamentous fungal cell that has been genetically modified to produce glycoproteins with predominantly a particular N-glycan structure. In particularly preferred aspects, the host cells are genetically modified so 25 that they express recombinant glycoproteins in which the glycosylation pattern is human-like or humanized. In particular, the host cells can be modified so that they express recombinant glycoproteins having predominantly a particular desired N-glycan structure. A lower eukaryotic host cell when used herein in connection with glycosylation profiles, refers to any eukaryotic cell which ordinarily produces high mannose containing N-linked glycans, and thus, includes most typical lower eukaryotic cells, 30 including uni- and multi-cellular fungal and algal cells.

All publications, patents, patent applications, and other references mentioned herein are hereby incorporated by reference in their entireties.

BRIEF DESCRIPTION OF THE DRAWINGS

35 Figure 1 illustrates the effect of Pmt inhibitors on O-glycosylation of secreted recombinant reporter proteins in *Pichia pastoris*. The chemical inhibitors of Pmt reduced O-glycosylation to a level similar to that observed in a strain lacking PMT1. Western blotting using an anti-

polyhistidine antibody was used to detect His-tagged human Kringle 1-3 domain (K1-3) of human plasminogen in the growth media of wild-type (lanes 1-3) and pmt1 (lanes 4-5) strains. The slower migrating bands (seen as a higher molecular weight smear for K1-3 in lane 1) indicate O-glycosylated protein. Pmt1, PMT inhibitor 1.

5 Figure 2 shows a Western blot that demonstrates the effect of *T. reesei* α -mannosidase and the chemical inhibitor Pmt1 on the O-glycosylation of immunoglobulin light and heavy chain polypeptides. Both *T. reesei* α -mannosidase and chemical Pmt inhibitors reduced the level of O-glycosylation.

10 DETAILED DESCRIPTION OF THE INVENTION

The present invention provides a method for expressing a recombinant protein (includes polypeptides and glycoproteins), which is susceptible to O-linked glycosylation in a particular host cell, having a reduced amount of O-linked glycosylation (including no O-linked glycosylation) in that cell type. The method involves inducing expression of a protein of interest in a host cell in which the protein 15 is susceptible to O-linked glycosylation in the host cell in the presence of a chemical inhibitor of the activity of one or more of the Dol-P-Man:Protein (Ser/Thr) Mannosyl Transferase (Pmt) proteins involved in the transfer of mannose to a serine or threonine residue of the protein in the cell or one or more α 1,2-mannosidases, or both, at the time expression of the protein is induced. The protein that is expressed in the presence of the inhibitor or the one or more α 1,2-mannosidases has a reduced amount of 20 O-linked glycosylation compared to the amount of O-linked glycosylation that would have been present on the protein if it had been produced in the absence of the inhibitor or the one or more α 1,2-mannosidases, or both. The method is particularly useful because it provides a means for producing therapeutically relevant proteins where it is desired that the protein have a reduced amount of O-glycosylation in host cells such as lower eukaryotes, for example yeast, and bacteria, which would 25 normally produce proteins with O-linked glycans, having a reduced number of O-linked glycans. However, while the method is especially suitable for expressing proteins with reduced O-linked glycosylation in lower eukaryotic organisms, the method can also be practiced in higher eukaryotic organisms and bacteria.

The method is an improvement over prior art methods for producing proteins having 30 reduced O-linked glycosylation in host cells in which the proteins are susceptible to O-linked glycosylation. For example, Tanner *et al.* in U. S. Patent No. 5,714,377 describes a method for making recombinant proteins having reduced O-linked glycosylation using fungal cells such as yeast cells in which one or more of *PMT* genes encoding the Pmt protein have been genetically modified so that recombinant proteins are produced, which have reduced O-linked glycosylation. While deletion of either 35 the *PMT1* or *PMT2* genes in a fungal host cell enables production of a recombinant protein having reduced O-linked glycosylation in the fungal host cell, expression of the *PMT1* and *PMT2* genes are important for host cells growth and either deletion alone also adversely affects the ability of the fungal

host cell to grow thus making it difficult to produce a sufficient quantity of host cells or recombinant protein with a reduced amount of O-linked glycosylation. Deletion of both genes appears to be lethal to the fungal host cell. Therefore, genetic elimination of the *PMT1* and *PMT2* genes in a host cell would appear to be an undesirable means for producing recombinant proteins having reduced O-linked glycosylation.

5 In contrast, the *PMT* genes in the host cells used in the method of the present invention have not been modified or deleted, which enables the host cell to O-glycosylate those proteins that are important for cell growth until which time the activity of the Pmt proteins is inhibited. In general, this enables the host cells to be grown to higher levels than the levels that could be obtained if the *PMT* genes 10 had been deleted. In addition, in particular embodiments, expression of the recombinant protein in the host cell is controlled by an inducible promoter and the Pmt activity in the host cell is not inhibited or one or more α 1,2-mannosidases added, or both, until expression of the recombinant protein is induced. This enables large quantities of host cells containing a nucleic acid encoding a recombinant protein to be produced in culture before inducing expression of the recombinant protein and adding the Pmt inhibitor 15 and/or one or more α 1,2-mannosidases. This can enable production of larger amounts recombinant protein having reduced O-linked glycosylation to be produced in the culture in a shorter period of time than would occur for host cells which have had one or more *PMT* genes deleted and grow poorly in culture.

20 This improvement over the prior art also facilitates the production of glycoproteins having reduced O-linked glycosylation in host cells that have been genetically modified to produce glycoproteins having predominantly a particular N-linked glycan structure but which also O-glycosylate the glycoprotein. Methods for producing a wide variety of glycoproteins having predominantly particular N-linked glycoforms have been disclosed in U.S. Patent No. 7,029,872 and U.S. Published Application Nos. 20050170452, 20050260729, 20040230042, 20050208617, 20050208617, 20040171826, 25 20060160179, 20060040353, and 20060211085. Any one of the host cells described in the aforementioned patent and patent applications can be used to produce a glycoprotein having predominantly a particular N-linked glycan structure and having reduced O-linked glycosylation using the method disclosed herein. It has been found that some host cells that have been genetically modified to produce glycoproteins having predominantly a particular N-linked glycan structure can grow less well 30 in culture under particular conditions than host cells that have not been modified. For example, particular fungal and yeast cells in which genes involved in hypermannosylation have been deleted and other genes needed to produce particular mammalian or human like N-linked glycan structures have been added, can grow less well than fungal or yeast cells that do not the genetic modifications. In some of these genetically modified fungal or yeast cells, further introducing deletions of the *PMT1* or *PMT2* 35 genes either is lethal to the cells or adversely affects the ability of the cells to grow to sufficient quantities in culture. The method herein avoids the potential deleterious effects of deleting the *PMT1* and *PMT2* genes by allowing the cells to grow to sufficient quantities in culture before inducing

expression of the recombinant glycoprotein and adding an inhibitor of the activity of the Pmt proteins, or one or more α 1,2-mannosidases, or both, to produce the recombinant glycoprotein having predominantly particular N-linked glycan structures and reduced O-linked glycosylation.

Therefore, an important aspect of the method is that it provides for a glycoprotein 5 composition comprising reduced O-linked glycosylation and a predominantly a specific N-linked glycoform in which the recombinant glycoprotein may exhibit increased biological activity and/or decreased undesired immunogenicity relative to compositions of the same glycoprotein produced from mammalian cell culture, such as CHO cells. An additional advantage of producing the glycoprotein composition comprising reduced O-linked glycosylation and a predominant N-linked glycoform is that it 10 avoids production of undesired or inactive glycoforms and heterogeneous mixtures, which may induce undesired effects and/or dilute the more effective glycoform. Thus, therapeutic pharmaceutical composition of glycoprotein molecules comprising, for example, predominantly $\text{Man}_5\text{GlcNAc}_2$, $\text{Man}_3\text{GlcNAc}_2$, $\text{GlcNAcMan}_5\text{GlcNAc}_2$, $\text{GlcNAcMan}_3\text{GlcNAc}_2$, $\text{GlcNAc}_2\text{Man}_3\text{GlcNAc}_2$, $\text{GalGlcNAcMan}_5\text{GlcNAc}_2$, $\text{Gal}(\text{GlcNAc})_2\text{Man}_5\text{GlcNAc}_2$, $(\text{GalGlcNAc})_2\text{Man}_5\text{GlcNAc}_2$, 15 $\text{NANAGalGlcNAcMan}_3\text{GlcNAc}_2$, $\text{NANA}_2\text{Gal}_2\text{GlcNAcMan}_3\text{GlcNAc}_2$, and $\text{GalGlcNAcMan}_3\text{GlcNAc}_2$ glycoforms and having reduced O-linked glycosylation may well be effective at lower doses, thus having higher efficacy/potency.

In general, the method for producing proteins having reduced O-linked glycosylation comprises transforming a host cell with a nucleic acid encoding a recombinant or heterologous protein in 20 which it is desirable to produce the protein having reduced O-linked glycosylation. The nucleic acid encoding the recombinant protein is operably linked to regulatory sequences that allow expression of the recombinant protein. Such regulatory sequences include an inducible promoter and optionally an enhancer upstream, or 5', to the nucleic acid encoding the fusion protein and a transcription termination site 3' or down stream from the nucleic acid encoding the recombinant protein. The nucleic acid also 25 typically encodes a 5' UTR region having a ribosome binding site and a 3' untranslated region. The nucleic acid is often a component of a vector replicable in cells in which the recombinant protein is expressed. The vector can also contain a marker to allow recognition of transformed cells. However, some cell types, particularly yeast, can be successfully transformed with a nucleic acid lacking 30 extraneous vector sequences.

Nucleic acids encoding desired recombinant proteins can be obtained from several 35 sources. cDNA sequences can be amplified from cell lines known to express the protein using primers to conserved regions (see, for example, Marks *et al.*, *J. Mol. Biol.* 581-596 (1991)). Nucleic acids can also be synthesized de novo based on sequences in the scientific literature. Nucleic acids can also be synthesized by extension of overlapping oligonucleotides spanning a desired sequence (see, e.g., Caldas *et al.*, *Protein Engineering*, 13, 353-360 (2000)).

In one aspect, the nucleic acid encoding the protein is operably linked to an inducible promoter, which allows expression of the protein to be induced when desired. In another aspect, the

nucleic acid encoding the protein is operably linked to a constitutive promoter. To facilitate isolation of the expressed protein, it is currently preferable that the protein include a signal sequence that directs the protein to be excreted into the cell culture medium where it can then be isolated. In the first aspect, the transformed host cells are cultured for a time sufficient to produce a desired multiplicity of host cells

5 sufficient to produce the desired amount of protein before adding one or more inhibitors of Pmt-mediated O-linked glycosylation to the culture medium. The inducer and inhibitor can be added to the culture simultaneously or the inducer is added to the culture before adding the one or more Pmt inhibitors or the one or more Pmt inhibitors is added to the culture before adding the inducer. The induced protein is produced having reduced O-linked glycosylation and can be recovered from the culture medium or for

10 proteins not having a signal sequence, from the host cell by lysis. In the second aspect, wherein the nucleic acid encoding the protein is operably linked to a constitutive promoter, the one or more inhibitors of Pmt-mediated O-linked glycosylation is added to the culture medium at the same time the culture is established and the protein, which is produced having reduced O-linked glycosylation, can be recovered from the culture medium or for proteins not having a signal sequence, from the host cell by lysis. An

15 example illustrating the method using an inducible promoter is shown in Example 2 and an example illustrating the method using a constitutive promoter is shown in Example 3.

Inhibitors useful for producing proteins with reduced O-linked glycosylation are chemicals or compositions that inhibit the activity one or more of the Pmt proteins. When the host cell is a lower eukaryote such as fungi or yeast, it is desirable that the inhibitor inhibit at least the activity of Pmt1 or Pmt2, or both. In higher eukaryotes, it is desirable that the inhibitor inhibit activity of the homologue in the higher eukaryote that corresponds to the Pmt1 or Pmt2. Chemical inhibitors that can be used include the benzylidene thiazolidinediones identified in U.S. Patent No. 7,105,554, which includes 5-[[3,4-bis(phenylmethoxy) phenyl]methylene]-4-oxo-2-thioxo-3-thiazolidineacetic acid; 5-[[3-(1-phenylethoxy)-4-(2-phenylethoxy)]phenyl]methylene]-4-oxo-2-thioxo-3-thiazolidineacetic acid; 3-25 hydroxy-4-(2-phenylethoxy)benzaldehyde; 3-(1-phenylethoxy)-4-(2-phenylethoxy)-benzaldehyde; and, 5-[[3-(1-phenyl-2-hydroxy)ethoxy)-4-(2-phenylethoxy)]phenyl]methylene]-4-oxo-2-thioxo-3-thiazolidineacetic acid. Other compounds that might be useful are the structurally similar compounds disclosed in Voss *et al.* in WO 94/29287, which discloses methods of making arylidene-4-oxo-2-thioxo-3-thiazolidine carboxylic acids and which are disclosed to be useful in the prophylaxis and treatment of 30 late effects of diabetes as well as the prophylaxis and treatment of atherosclerosis and arteriosclerosis and in Esswein *et al.* in U.S. Patent No. 6,673,816, which discloses methods of making derivatives of rhodaninecarboxylic acids and their use for treatment of metabolic bone disorders.

In the examples, chemical inhibitors selected from the group consisting of 5-[[3,4-bis(phenylmethoxy) phenyl]methylene]-4-oxo-2-thioxo-3-thiazolidineacetic acid; 5-[[3-(1-phenylethoxy)-4-(2-phenylethoxy)]phenyl]methylene]-4-oxo-2-thioxo-3-thiazolidineacetic acid; and, 5-[[3-(1-phenyl-2-hydroxy)ethoxy)-4-(2-phenylethoxy)]phenyl]methylene]-4-oxo-2-thioxo-3-thiazolidineacetic acid are shown to be effective in producing recombinant proteins having reduced O-

linked glycosylation in *Pichia pastoris* strains that had intact, functional *PMT1* and *PMT2* genes. Table 1 of Example 2 shows that any one of the above three Pmt chemical inhibitors added to a culture of recombinant *Pichia pastoris* having intact, functional *PMT1* and *PMT2* genes and transformed with a nucleic acid encoding a recombinant, secretable Kringle 1-3 protein operably linked to an inducible promoter at the time expression of the recombinant protein was induced, produced a recombinant protein having a level of reduced O-linked glycosylation that was comparable to the level of O-linked glycosylation seen for *Pichia pastoris* cells containing a deletion of either the *PMT1* or *PMT2* gene. The above Pmti inhibitors have been used in amounts from about 0.03 μ M to 20 μ M to produce proteins having reduced O-linked glycosylation compared to the amount of O-linked glycosylation on the protein when grown in similar host cell cultures in the absence of the Pmti inhibitors. The results shown in Example 3 further shows that the host cell cultures can be grown in the presence of Pmti inhibitor at an amount that is sufficient to inhibit O-linked glycosylation without killing the host cells.

The method can include adding to the culture medium containing the one or more Pmt inhibitors one or more α -1,2-mannosidase enzymes to produce the recombinant protein having reduced O-linked glycosylation. The α -1,2-mannosidases are a conserved family of eukaryotic enzymes for maturation of N-glycans, which are capable of trimming $\text{Man}_9\text{GlcNAc}_2$ to $\text{Man}_8\text{GlcNAc}_2$ in yeast. (Vallee *et al.*, 2000, EMBO J., 19: 581-588). The α -1,2-mannosidases are also known as class I α -mannosidases and have been identified in mammalian, lower eukaryotic species, and insect cells (Kawar *et al.*, 2000, Glycobiology 10: 347-355). Mammalian cells are known to have several class I α -mannosidases, some of which are capable of trimming multiple mannose residues (Moremen *et al.*, 1994, Glycobiology 4: 113-125), while yeast appear to have fewer, more specialized α -1,2-mannosidases. For example, *Saccharomyces* has been disclosed to have a single α -1,2-mannosidase encoded by *MNN1*, which removes one specific mannose residue (for example, $\text{Man}_9\text{GlcNAc}_2$ to $\text{Man}_8\text{GlcNAc}_2$) (Herscovics, 1999, Biochim Biophys Acta., 1473: 96-107). Thus, the endogenous α -1,2-mannosidase present in many lower eukaryotes such as fungi and yeast and which cannot remove multiple mannose residues from Glycan structures, is not capable of enabling production of proteins having reduced O-linked glycosylation. Therefore, the method herein requires introduction into the culture medium containing the host cells an α -1,2-mannosidase capable of trimming multiple mannose residues from an O-linked Glycan or introduction into the host cell a nucleic acid encoding an α -1,2-mannosidase capable of trimming multiple mannose residues from an O-linked Glycan. The α -1,2-mannosidase herein includes the intact, native α -1,2-mannosidase; an α -1,2-mannosidase modified to enhance its α -1,2-mannosidase activity; an α -1,2-mannosidase modified to decrease its α -1,2-mannosidase activity; and, a recombinant α -1,2-mannosidase comprising at least the catalytic domain having the α -1,2-mannosidase activity (for example, a fusion protein comprising the catalytic domain having the α -1,2-mannosidase activity fused to heterologous proteins, polypeptides, or peptides).

In particular embodiments, the α -1,2-mannosidase, which is capable of trimming multiple mannose residues from an O-linked glycans and is added to the cell culture, is produced by

5 *Trichoderma* sp., *Saccharomyces* sp., or *Aspergillus* sp. Currently, preferred α -1,2-mannosidases are obtained from *Trichoderma reesei*, *Aspergillus niger*, or *Aspergillus oryzae*. *T. reesei* is also known as *Hypocrea jecorina*. In Example 3, a transformed yeast comprising an expression cassette, which expresses a recombinant α -1,2-mannosidase comprising the *Trichoderma reesei* α -1,2-mannosidase catalytic domain fused to the *Saccharomyces cerevisiae* α MAT pre signal sequence, was used to produce recombinant proteins having reduced O-linked glycosylation. Another example of a recombinant α -1,2-mannosidase that could be used in the method herein to produce proteins having reduced O-linked glycosylation is the recombinant *Trichoderma reesei* α -1,2-mannosidase disclosed in Maras *et al.*, 2000, J. Biotechnol. 77:255-263 wherein the *Trichoderma reesei* α -1,2-mannosidase catalytic domain was 10 fused to a *Saccharomyces cerevisiae* α -MAT prepro-signal peptide.

15 The α -1,2-mannosidase can also be produced from a chimeric nucleic acid comprising a nucleic acid sequence encoding at least the catalytic domain of an α -1,2-mannosidase, which is capable of trimming multiple mannose residues from an O-linked glycans, operatively linked to a nucleic acid sequence encoding a cellular targeting signal peptide not normally associated with the catalytic domain. 20 The chimeric nucleic acid can be operably linked to a constitutive or inducible promoter. The chimeric nucleic acid is transformed into a host cell to produce the α -1,2-mannosidase, which is then isolated and then added to the cell culture medium containing cells transformed with the nucleic acid encoding the heterologous protein at the time expression of the protein is induced. Alternatively, the host cell is transformed with the chimeric nucleic acid encoding the α -1,2-mannosidase and the nucleic acid 25 encoding the recombinant protein and co-expressing the α -1,2-mannosidase and the recombinant protein at the same time. In particular embodiments, both the chimeric nucleic acid encoding the α -1,2-mannosidase and the nucleic acid encoding the recombinant protein are both operably linked to an inducible promoter. In other embodiments, one or both of the promoters are constitutive. Example 3 30 illustrates the method wherein nucleic acids encoding both the α -mannosidase and the recombinant protein are operably linked to a constitutive promoter, introduced into a host cell, and a culture of the host cells is then incubated in the presence of one or more Pmt inhibitors to produce the recombinant protein having reduced O-linked glycosylation. Example 3 further shows that there appears that the Pmti 35 inhibitor and the α -1,2-mannosidase appear to synergistically reduce the amount of O-linked glycosylation compared to the amount of O-linked glycosylation in the presence of either alone.

30 In particular aspects, reduced O-linked glycosylation can be effected by adding only the one or more α -1,2-mannosidases and not one or more Pmt inhibitors to the culture medium. In one aspect, the nucleic acid encoding the recombinant protein is operably linked to an inducible promoter, which allows expression of the recombinant protein to be induced when desired. In another aspect, the nucleic acid encoding the protein is operably linked to a constitutive promoter. To facilitate isolation of 35 the expressed recombinant protein, it is currently preferable that the protein include a signal sequence that directs the recombinant protein to be excreted into the cell culture medium where it can then be isolated.

In the first aspect, the transformed host cells are cultured for a time sufficient to produce a desired multiplicity of host cells sufficient to produce the desired amount of the recombinant protein before adding the one or more α -1,2-mannosidases to the culture medium. The inducer and the one or more α -1,2-mannosidases can be added to the culture simultaneously or the inducer is added to the culture before adding the one or more α -1,2-mannosidases or the one or more α -1,2-mannosidases is added to the culture before adding the inducer. The induced recombinant protein is produced having reduced O-linked glycosylation and can be recovered from the culture medium or for proteins not having a signal sequence, from the host cell by lysis.

In the second aspect, wherein the nucleic acid encoding the recombinant protein is operably linked to a constitutive promoter, the one or more α -1,2-mannosidases is added to the culture medium at the same time the culture is established and the recombinant protein, which is produced having reduced O-linked glycosylation, can be recovered from the culture medium or for recombinant proteins not having a signal sequence, from the host cell by lysis.

In a further still aspect for producing proteins having reduced O-linked glycosylation without using an inhibitor of Pmt-mediated O-linked glycosylation, the host cell is transformed with a chimeric nucleic acid encoding the α -1,2-mannosidase and a nucleic acid encoding the recombinant protein and co-expressing the α -1,2-mannosidase and the recombinant protein to produce the recombinant protein having reduced O-linked glycosylation. In particular embodiments, both the chimeric nucleic acid encoding the α -1,2-mannosidase and the nucleic acid encoding the recombinant protein as both operably linked to an inducible promoter. In other embodiments, one or both of the promoters are constitutive. In the case of an inducible promoter, the host cells are grown to produce a desired multiplicity of host cells before inducing expression of the α -1,2-mannosidase and/or recombinant protein. Example 3 illustrates the method wherein nucleic acids encoding both the α -1,2-mannosidases and the recombinant protein are operably linked to a constitutive promoter are introduced into a host cell and a culture of the host cells is then incubated for a time to produce the recombinant protein, which has reduced O-linked glycosylation compared to the recombinant protein produced in cells in the absence of the α -1,2-mannosidase.

II. Host Cells

While host cells for the method herein includes both higher eukaryote cells and lower eukaryote cells, lower eukaryote cells, for example filamentous fungi or yeast cells, are currently preferred for expression of proteins because they can be economically cultured, give high yields of protein, and when appropriately modified are capable of producing proteins having suitable glycosylation patterns. Lower eukaryotes include yeast, fungi, collar-flagellates, microsporidia, alveolates (e.g., dinoflagellates), stramenopiles (e.g., brown algae, protozoa), rhodophyta (e.g., red algae), plants (e.g., green algae, plant cells, moss) and other protists. Yeast and fungi include, but are not limited to: *Pichia* sp. (for example, *Pichia pastoris*, *Pichia finlandica*, *Pichia trehalophila*, *Pichia koclamae*, *Pichia*

membranaefaciens, *Pichia minuta* (*Ogataea minuta*, *Pichia lindneri*), *Pichia opuntiae*, *Pichia thermotolerans*, *Pichia salictaria*, *Pichia guercuum*, *Pichia piperi*, *Pichia stiptis*, *Pichia methanolica*), *Saccharomyces* sp. (for example *Saccharomyces cerevisiae*), *Hansenula polymorpha*, *Kluyveromyces* sp. (for example, *Kluyveromyces lactis*), *Candida albicans*, *Aspergillus* sp (for example, *Aspergillus nidulans*, *Aspergillus niger*, *Aspergillus oryzae*), *Trichoderma reesei*, *Chrysosporium lucknowense*, *Fusarium* sp. (for example, *Fusarium gramineum*, *Fusarium venenatum*), *Physcomitrella patens* and *Neurospora crassa*. Yeast, in particular, are currently preferred because yeast offers established genetics

allowing for rapid transformations, tested protein localization strategies, and facile gene knock-out techniques. Suitable vectors have expression control sequences, such as promoters, including 3-phosphoglycerate kinase or other glycolytic enzymes, and an origin of replication, termination sequences, and the like as desired.

Various yeasts, such as *K. lactis*, *Pichia pastoris*, *Pichia methanolica*, and *Hansenula polymorpha* are currently preferred for cell culture because they are able to grow to high cell densities and secrete large quantities of recombinant protein. Likewise, filamentous fungi, such as *Aspergillus niger*, *Fusarium* sp, *Neurospora crass*, and others can be used to produce recombinant proteins at an industrial scale.

Lower eukaryotes, in particular filamentous fungi and yeast, can be genetically modified so that they express proteins or glycoproteins in which the glycosylation pattern is human-like or humanized. This can be achieved by eliminating selected endogenous glycosylation enzymes and/or supplying exogenous enzymes as described by Gerngross *et al.* in U.S. Patent No. US7029872, and U. S. Published Patent Application Nos. 20040018590, 20050170452, 20050260729, 20040230042, 20050208617, 20040171826, 20050208617, 20060160179, 20060040353, and 20060211085. Thus, a host cell can additionally or alternatively be engineered to express one or more enzymes or enzyme activities, which enable the production of particular N-glycan structures at a high yield. Such an enzyme can be targeted to a host subcellular organelle in which the enzyme will have optimal activity, for example, by means of signal peptide not normally associated with the enzyme. Host cells can also be modified to express a sugar nucleotide transporter and/or a nucleotide diphosphatase enzyme. The transporter and diphosphatase improve the efficiency of engineered glycosylation steps, by providing the appropriate substrates for the glycosylation enzymes in the appropriate compartments, reducing competitive product inhibition, and promoting the removal of nucleoside diphosphates. See, for example, Gerngross *et al.* in U.S. Published Patent Application No.20040018590 and Hamilton, 2003, Science 301: 1244-46 and the aforementioned U.S. patent and patent applications.

By way of example, a host cell (for example, yeast or fungal) can be selected or engineered to be depleted in 1,6-mannosyl transferase activities, which would otherwise add mannose residues onto the N-glycan of a glycoprotein, and to further include a nucleic acid for ectopic expression of an α -1,2 mannosidase activity, which enables production of recombinant glycoproteins having greater than 30 mole percent $\text{Man}_5\text{GlcNAc}_2$ N-glycans. When a glycoprotein is produced in the host cells

according to the method described herein, the host cells will produce a glycoprotein having predominantly a $\text{Man}_5\text{GlcNAc}_2$ N-glycan structure and reduced O-glycosylation compared to the glycoprotein produced in the cell otherwise. In a further aspect, the host cell is engineered to further include a nucleic acid for ectopic expression of GlcNAc transferase I activity, which enables production

5 of glycoproteins having predominantly $\text{GlcNAcMan}_5\text{GlcNAc}_2$ N-glycans. When a glycoprotein is produced in the host cells according to the method described herein, the host cells will produce a glycoprotein having predominantly a $\text{GlcNAcMan}_5\text{GlcNAc}_2$ N-glycan structure and reduced O-glycosylation compared to the glycoprotein produced in the cell otherwise. In a further still aspect, the host cell is engineered to further include a nucleic acid for ectopic expression of mannosidase II activity, which enables production of glycoproteins having predominantly $\text{GlcNAcMan}_3\text{GlcNAc}_2$ N-glycans.

10 When a glycoprotein is produced in the host cells according to the method described herein, the host cells will produce a glycoprotein having predominantly a $\text{GlcNAcMan}_3\text{GlcNAc}_2$ N-glycan structure and reduced O-glycosylation compared to the glycoprotein produced in the cell otherwise. In a further still aspect, the host cell is engineered to further include a nucleic acid for ectopic expression of GlcNAc 15 transferase II activity, which enables production of glycoproteins having predominantly $\text{GlcNAc}_2\text{Man}_3\text{GlcNAc}_2$ N-glycans. When a glycoprotein is produced in the host cells according to the method described herein, the host cells will produce a glycoprotein having predominantly a $\text{GlcNAc}_2\text{Man}_3\text{GlcNAc}_2$ N-glycan structure and reduced O-glycosylation compared to the glycoprotein produced in the cell otherwise. In further still aspects, the above host cells can be further engineered to 20 produce particular hybrid or complex N-glycan or human-like N-glycan structures by further including one or more higher eukaryote genes involved in N-linked glycosylation, in any combination, that encode for example, sialytransferase activities, class II and III mannosidase activities, GlcNAc transferase II, III, IV, V, VI, IX activity, and galactose transferase activity. It is currently preferable that the cells further include one or more of nucleic acids encoding UDP-specific diphosphatase activity, GDP-specific 25 diphosphatase activity, and UDP-GlcNAc transporter activity.

Plants and plant cell cultures may be used for expression of proteins and glycoproteins with reduced O-linked glycosylation as taught herein (See, for example, Lerrick & Fry, 1991, *Hum. Antibodies Hybridomas* 2: 172-89); Benvenuto *et al.*, 1991, *Plant Mol. Biol.* 17: 865-74); Durin *et al.*, 1990, *Plant Mol. Biol.* 15: 281-93); Hiatt *et al.*, 1989, *Nature* 342: 76-8). Preferable plant hosts include, 30 for example, *Arabidopsis*, *Nicotiana tabacum*, *Nicotiana rustica*, and *Solanum tuberosum*.

Insect cell culture can also be used to produce proteins and glycoproteins with reduced O-linked glycosylation, as taught herein for example, baculovirus-based expression systems (See, for example, Pultz *et al.*, 1990, *Bio/Technology* 8: 651-654).

35 Although not currently as economical to culture as lower eukaryotes and prokaryotes, mammalian tissue cell culture can also be used to express and produce proteins and glycoproteins with reduced O-linked glycosylation as taught herein (See Winnacker, *From Genes to Clones* (VCH Publishers, NY, 1987). Suitable hosts include CHO cell lines, various COS cell lines, HeLa cells,

preferably myeloma cell lines or the like or transformed B-cells or hybridomas. Expression vectors for these cells can include expression control sequences, such as an origin of replication, a promoter, an enhancer (Queen *et al.*, 1986I, *mmunol. Rev.* 89:49-68), and necessary processing information sites, such as ribosome binding sites, RNA splice sites, polyadenylation sites, and transcriptional terminator sequences. Expression control sequences are promoters derived from immunoglobulin genes, SV40, Adenovirus, bovine Papilloma Virus, cytomegalovirus and the like. Generally, a selectable marker, such as a neoR expression cassette, is included in the expression vector.

The nucleic acid encoding the protein to be expressed can be transferred into the host cell by conventional methods, which vary depending on the type of cellular host. For example, calcium phosphate treatment, protoplast fusion, natural breeding, lipofection, biolistics, viral-based transduction, or electroporation can be used for cellular hosts. Tungsten particle ballistic transgenesis is preferred for plant cells and tissues. (See, generally, Maniatis *et al.*, *Molecular Cloning: A Laboratory Manual* (Cold Spring Harbor Press, 1982))

Once expressed, the proteins or glycoproteins having reduced O-linked glycosylation can be purified according to standard procedures of the art, including ammonium sulfate precipitation, affinity columns, column chromatography, gel electrophoresis and the like (See, generally, Scopes, R., *Protein Purification* (Springer-Verlag, N.Y., 1982)). Substantially pure glycoproteins of at least about 90 to 95% homogeneity are preferred, and 98 to 99% or more homogeneity most preferred, for pharmaceutical uses. Once purified, partially or to homogeneity as desired, the proteins can then be used therapeutically (including extracorporeally) or in developing and performing assay procedures, immunofluorescent stainings, and the like. (See, generally, *Immunological Methods*, Vols. I and II (Lefkovits and Pernis, eds., Academic Press, NY, 1979 and 1981).

Therefore, further provided are glycoprotein compositions comprising a predominant species of N-glycan structure and having reduced O-linked glycosylation compared to compositions of the glycoprotein which have been produced in host cells have not been incubated in the presence of an inhibitor of Pmt-mediated O-linked glycosylation or an α -1,2-mannosidase capable of trimming more than one mannose residue from a glycans structure or both. In particular aspects, the glycoprotein composition comprises a glycoprotein having a predominant N-glycan structure selected from the group consisting of Man₅GlcNAc₂, Man₃GlcNAc₂, GlcNAcMan₅GlcNAc₂, GlcNAcMan₃GlcNAc₂, GlcNAc₂Man₃GlcNAc₂, GalGlcNAcMan₅GlcNAc₂, Gal(GlcNAc)₂Man₅GlcNAc₂, (GalGlcNAc)₂Man₅GlcNAc₂, NANAGalGlcNAcMan₃GlcNAc₂, NANA₂Gal₂GlcNAcMan₃GlcNAc₂, and GalGlcNAcMan₃GlcNAc₂ glycoforms.

III. Pharmaceutical Compositions

Proteins and glycoproteins having reduced O-linked glycosylation can be incorporated into pharmaceutical compositions comprising the glycoprotein as an active therapeutic agent and a variety of other pharmaceutically acceptable components (See, Remington's *Pharmaceutical Science*

(15th ed., Mack Publishing Company, Easton, Pennsylvania, 1980). The preferred form depends on the intended mode of administration and therapeutic application. The compositions can also include, depending on the formulation desired, pharmaceutically-acceptable, non-toxic carriers or diluents, which are defined as vehicles commonly used to formulate pharmaceutical compositions for animal or human administration. The diluent is selected so as not to affect the biological activity of the combination. Examples of such diluents are distilled water, physiological phosphate-buffered saline, Ringer's solutions, dextrose solution, and Hank's solution. In addition, the pharmaceutical composition or formulation can also include other carriers, adjuvants, or nontoxic, nontherapeutic, nonimmunogenic stabilizers, and the like.

10 Pharmaceutical compositions for parenteral administration are sterile, substantially isotonic, pyrogen-free and prepared in accordance with GMP of the FDA or similar body. Glycoproteins can be administered as injectable dosages of a solution or suspension of the substance in a physiologically acceptable diluent with a pharmaceutical carrier that can be a sterile liquid such as water oils, saline, glycerol, or ethanol. Additionally, auxiliary substances, such as wetting or emulsifying 15 agents, surfactants, pH buffering substances and the like can be present in compositions. Other components of pharmaceutical compositions are those of petroleum, animal, vegetable, or synthetic origin, for example, peanut oil, soybean oil, and mineral oil. In general, glycols such as propylene glycol or polyethylene glycol are preferred liquid carriers, particularly for injectable solutions. Glycoproteins can be administered in the form of a depot injection or implant preparation which can be formulated in 20 such a manner as to permit a sustained release of the active ingredient. Typically, compositions are prepared as injectables, either as liquid solutions or suspensions; solid forms suitable for solution in, or suspension in, liquid vehicles prior to injection can also be prepared. The preparation also can be emulsified or encapsulated in liposomes or micro particles such as polylactide, polyglycolide, or copolymer for enhanced adjuvant effect, as discussed above (See Langer, *Science* 249, 1527 (1990) and 25 Hanes, *Advanced Drug Delivery Reviews* 28, 97-119 (1997).

Unless otherwise defined herein, scientific and technical terms and phrases used in connection with the present invention shall have the meanings that are commonly understood by those of ordinary skill in the art. Further, unless otherwise required by context, singular terms shall include the plural and plural terms shall include the singular. Generally, nomenclatures used in connection with, and 30 techniques of biochemistry, enzymology, molecular and cellular biology, microbiology, genetics and protein and nucleic acid chemistry and hybridization described herein are those well known and commonly used in the art. The methods and techniques of the present invention are generally performed according to conventional methods well known in the art and as described in various general and more specific references that are cited and discussed throughout the present specification unless otherwise indicated. See, for example, Sambrook et al. *Molecular Cloning: A Laboratory Manual*, 2d ed., Cold 35 Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1989); Ausubel et al., *Current Protocols in Molecular Biology*, Greene Publishing Associates (1992, and Supplements to 2002); Harlow and Lane,


Antibodies: A Laboratory Manual, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1990); Taylor and Drickamer, Introduction to Glycobiology, Oxford Univ. Press (2003); Worthington Enzyme Manual, Worthington Biochemical Corp., Freehold, NJ; Handbook of Biochemistry: Section A Proteins, Vol I, CRC Press (1976); Handbook of Biochemistry: Section A Proteins, Vol II, CRC Press (1976); Essentials of Glycobiology, Cold Spring Harbor Laboratory Press (1999).

5 The following examples are intended to promote a further understanding of the present invention.

EXAMPLE 1


10 This example provides method for preparing various Pmt inhibitors. Unless otherwise stipulated all materials were obtained from Sigma-Aldrich Chemical Co. (St. Louis, MO) and used as received. The ^1H NMR spectra of all intermediates and final products were in accord with published data.

15 Preparation of Pmti-1, (5-[[3,4-bis(phenylmethoxy) phenyl]methylene]-4-oxo-2-thioxo-3-thiazolidineacetic Acid), is as follows.



20 The procedure was adapted from Orchard *et al.* in U.S. Patent No. 7,105,554. A solution of rhodanine-3-acetic acid (1 g, 5.20 mmol, 1 eq.), 3,4-dibenzylxybenzaldehyde (2.04 g, 6.25 mmol, 1.2 eq.), and sodium acetate (1.3 g, 15.6 mmol, 3 eq.) in acetic acid (30 mL) is heated to reflux, and stirred overnight. As the reaction mixture is cooled to room temperature, the product is precipitated and filtered and washed with acetic acid, then petroleum ether. The residue is dissolved in hot DMSO, filtered, and precipitated by addition of water. Upon cooling, the precipitate is filtered and recrystallized from ethyl acetate and petroleum ether to give a product which is suspended in water and freeze-dried overnight in vacuo to give the final product as a fluffy yellow powder.

Preparation of Pmti-2, 2 (5-[[3-(1-Phenylethoxy)-4-(2-phenylethoxy)]phenyl]methylene]-4-oxo-2-thioxo-3-thiazolidineacetic Acid), is as follows.




5

This product is synthesized according to the directions of Orchard *et al.* in U.S. Patent No. 7,105,554. A solution of rhodanine-3-acetic acid (375 mg, 1.96 mmol, 1 eq.), 3-(1-phenylethoxy)-4-(2-phenylethoxy)benzaldehyde (680 mg, 1.96 mmol, 1 eq.) and ammonium acetate (453 mg, 3 eq.) is heated to 70°C for ten minutes, then cooled to room temperature and diluted with ethyl acetate (100 mL). The 10 organic solution is washed with 1M HCl (2 x 200 mL) and brine (200 mL) then dried over sodium sulfate and evaporated. The product is purified by liquid chromatography using a 10 x 2.5cm glass column packed with 35–75 µm C18 (Alltech Associates, Deerfield, IL). Gradient elution is employed. Buffer A is 0.1% acetic acid and buffer B is 80% acetonitrile. The gradient is comprised of 20% B for three minutes, increasing to 75% B over 40 minutes. The flow rate is 8 mL/min. Detection is at 280 nm. The 15 appropriate fractions are pooled, concentrated, and freeze-dried *in vacuo* to give the product as a fluffy yellow powder.

Preparation of Pmti-3, (5-[[3-(1-Phenyl-2-hydroxy)ethoxy)-4-(2-phenylethoxy)]phenyl]methylene]-4-oxo-2-thioxo-3-thiazolidineacetic Acid), (Orchard *et al.* in U.S. Patent No. 7,105,554) is synthesized in three steps as follows.

20



Step 1: Production of (+)-(S)-2-Acetoxy-1-bromo-1-phenylethane. Cold HBr-acetic acid

5 (12.4 g, 52.2 mmol) is added dropwise to (−)-(R)-1-phenylethane-1,2-diol (2.4 g, 17.4 mmol) during about five minutes and the mixture stirred at room temperature for 40 minutes. Water (25 mL) is added and the solution is neutralized with sodium carbonate and extracted with ether (3 x 30 mL). The combined extracts are dried and evaporated to give (+)-(S)-2-acetoxy-1-bromo-1-phenylethane (3.93 g, 93%), $d_{25}^{1.415}$ g/mL, $[x]_D^{24} + 93.5^\circ$ (c 5.63 in CCl_4) 2.72 (5H, s), 4.98 (aH, dd, 6.7 and 7.0 Hz) and 5.56 (2H, d).

10 This product is not distilled. The isomeric homogeneity is established by comparison of the nmr spectrum (absence of $PhCH^*OAc$ resonance) with that of 1,2-diacetoxy-1-phenylethane. (Note that racemic reagents are substituted for the optical isomers listed).

Step 2: Production of 3-[(1-Phenyl-2-hydroxy)ethoxy]-4-(2-phenylethoxy)-benzaldehyde.

15 (2-Acetoxy-1-bromoethyl)benzene (3.32 g, 13.67 mmol, 1.2 eq) (the product of Step 1), is added to a stirred solution of 3-hydroxy-4-(2-phenylethoxy)-benzaldehyde (2.76 g, 11.39 mmol, 1 eq.) and cesium carbonate (2.97 g, 9.11 mmol, 0.8 eq.) in N,N -dimethylformamide (15 mL). The solution is stirred for 19 hours at room temperature, then 21 hours at 80°C. The reaction is worked up by partitioning between ethyl acetate and water (brine is added to help break up the emulsion that formed). The organic layer is washed twice more with water, brine, and then dried over sodium sulfate and evaporated to give a dark oil. The residue is purified by chromatography on silica gel and elution with diethyl ether gives an orange oil. This oil is dissolved in methanol (100 mL) and to the solution is added an aqueous solution of sodium hydroxide (7 mL, 1M). After 30 minutes, the mixture is evaporated (to remove the methanol) and the residue partitioned between dichloromethane and water. The organic layer is dried over sodium sulfate and evaporated. The residue is purified by chromatographed on silica gel and elution with petroleum ether:diethyl ether (1:2) gives the product as a cream coloured powder.

Step 3: Production of Pmti-3. A solution of rhodanine-3-acetic acid (158 mg, 0.828 mmol, 1 eq.), 3-(1-phenyl-2-hydroxy)ethoxy)-4-(2-phenylethoxy)benzaldehyde (300mg, 0.828 mmol, 1 eq.) (the product of Step 2), and ammonium acetate (191 mg, 3 eq) in toluene (10 mL) is heated to reflux for 3.5 hours, cooled to room temperature, and diluted with ethyl acetate (50 mL). The organic solution 5 is washed with 1M HCl (2 x 200 mL) and brine (200 mL) then dried over sodium sulfate and evaporated. After work-up, the residue is purified by chromatography on silica gel. Elution with ethyl acetate gives a yellow gum, which is recrystallized from diethyl ether and petroleum ether to give the product as a yellow powder.

10

EXAMPLE 2

This example shows that *Pichia pastoris* transformed with an expression vector encoding the Kringle 1-3 marker glycoprotein and treated with Pmt inhibitors produced a glycoprotein having reduced O-glycosylation.

15 Plasmid DNA encoding a His-tagged reporter glycoprotein consisting of human plasminogen domains K1, K2, and K3 (Kringle 1-3 protein) under the control of the *Pichia pastoris* alcohol oxidase 1 (AOX1) promoter was transformed into wild-type *Pichia pastoris* to produce strain yJC53. The Kringle reporter protein consisting of domains K1, K2, K3, and K4 has been discussed in Duman *et al.* Biotechnol. Appl. Biochem. (1998), v.28, p.39-45 and only domain K3 in Choi *et al.*, 2003, Proc. Natl. Acad. Sci. U. S. A. 100(9): 5022-5027. The amino acid sequence of the Kringle 1-3 protein 20 used in the Example is

SECKTGNGKNYRGTMSTKNGITCQKWSSTS PHRPRFSPATHPSEGLEENYCRNPNDPQGPWCYTTDPE
KRYDYCDILECEEECMHCSGENYDGKISKTMMSGLECQAWDSQS PHAHGYIPS KFPNKNLKKNYCRNPDRE
LRPWCFTTDPNKRWECDIPRCTTPPPSSGPTYQCLKGTGENYRGNVAVTVSGHTCQHWSAQTPHTHSRT
PENFPCKNLDENYCRNPDGKRAPWCHTTNSQVRWEYCKIPSCDSSPVSTEQLAPTAPPELTPVVQDGHHH

25 HHHHHHHH (SEQ ID NO:1). The Kringle 1-3 protein contains at least two potential mammalian O-glycosylation sites that conform to the purported consensus sequence P at -1 and +3: the serine residue, which is O-glycosylated, is capitalized in the amino acid sequence “pppSsgp” and the threonine residue, which is O-glycosylated, is capitalized in the amino acid sequence “lapTapp”. The O-glycosylation sites are underlined in the above amino acid sequence. The potential mammalian O-glycosylation sites are 30 located between the K1 and K2 domains and the K2 and K3 domains. However, as shown in Table 1, in yeast the protein has about 20 O-linked glycosylation sites. Thus, O-linked glycosylation can be a significant disadvantage to producing proteins in yeast without inhibiting O-linked glycosylation as shown by the methods herein. An N-glycosylation site resides in the K3 domain, which had been removed by replacing the asparagine at position 208 of SEQ ID NO:1 with a serine. Therefore, the only 35 glycans on the Kringle 1-3 protein would be the result of O-glycosylation.

Plasmid containing DNA encoding the Kringle 1-3 protein was prepared using forward primer K1-3/UP 5'-CGGAA TTCTC AGAGT GCAAG ACTGG GAATA GAA-3' (SEQ ID NO:2) and

reverse primer K1-3/LP1 (Reverse primer, 3Gly+2His, paired with K 1-3/UP) 5'-ATGAT GATGA CCACC ACCGT CCTGG ACCAC AGGGG TTAG-3' (SEQ ID NO:3) to produce a PCR product, which was then PCR amplified using reverse primer K1-3/LP2 (Reverse primer, 3Gly+ 9His+stop codon, paired with K 1-3/UP) 5'-TTAAT GATGA TGATG ATGAT GATGA TGATG ACCAC CACC-3' (SEQ ID NO:4). PCR conditions were as follows: after 1 cycle of 95°C for 2 minutes as a denaturation step, the PCR reaction was subjected to 30 cycles of 95°C for 30 seconds, 60°C for 30 seconds, and 72°C for 1 minute, and then 1 cycle of 72°C for 10 minutes. After the PCR reaction and column purification of the PCR products, nucleotide A overhangs of the PCR products were generated using ExTaq (1 cycle of 72°C for 15 minutes). The resulting PCR products were used for the second PCR reaction as a PCR template where the primers, K1-3/UP and K1-3/LP2, were used to amplify wild-type Kringle 1-3 +3Gly+9His, which was cloned into a pCR2.1 plasmid vector (Invitrogen) to produce pBK105. The following PCR primers were then used to generate an Asn to Ser mutation at position 208 in the Kringle 1-3 protein to produce amino acid sequence NRTP from amino acid sequence SRTP: forward primer K3f (Asn to Ser) 5'-ACCCCTCACACACATTCTAGGACACCAGAAAATTC-3' (SEQ ID NO:5) and reverse primer K3r 5'-CTGTGCACTCCAGTGCTGACAGGTGTG-3' (SEQ ID NO:6). The Asn to Ser mutation was then generated in pBK105 by the Inverse PCR. PCR conditions were as follows; after 1 cycle of 95°C for 2 minutes as a denaturation step, the PCR reaction was subjected to 35 cycles of 95°C for 30 seconds, 60°C for 30 seconds, and 72°C for 5 minutes, and then 1 cycle of 72°C for 10 minutes. The resulting PCR products were ligated to produce plasmid pBK118, which was sequenced to confirm 20 the mutation.

Plasmid pBK118 was digested with *Eco*RI and the DNA fragments were gel purified and cloned into the *Eco*RI sites of pPICZaA (Invitrogen, La Jolla, CA) to produce pBK119 (*Pichia* expression plasmid). Plasmid pPICZaA contains an α -factor secretion signal that allows the efficient secretion of most proteins from *Pichia pastoris*; 5'-AOX, a 942 bp fragment containing the AOX 1 promoter that allows methanol-inducible and high-level expression in *Pichia pastoris*; and, the ZEOCIN 25 resistance gene for positive selection in *E. coli* and *Pichia pastoris*. Plasmid pBK119 was linearized with *Pme*I before transforming into *Pichia pastoris* strains. Plasmid pBK was transformed into *Pichia pastoris* strain yJC53, a wild-type strain, and various *PMT* knockout strains.

PMT knockout yeast strains were created in *Pichia pastoris* following the procedure 30 outlined for *Saccharomyces cerevisiae* in Gentzsch and Tanner, EMBO J. 1996 Nov 1;15(21): 25752-5759). The *Pichia pastoris* *PMT* genes were identified in the nucleotide sequence of the *Pichia pastoris* genome obtained from Integrated Genomics, Chicago, IL by homology searching using the nucleotide 35 sequences for the *Saccharomyces cerevisiae* *PMT* genes. Deletion of *Pichia pastoris* *PMT* (Pp*PMT*) genes was as follows. The Pp*PMT* deletion alleles were generated by the PCR overlap method (See for example, Davidson *et al.*, 2004, Glycobiology 14:399-407; Ho *et al.*, 1989, Gene 77:51-9; Horton *et al.*, 1989, Gene 77:61-8). In the first PCR reaction, DNA comprising the nucleotide sequences for 5' and 3' flanking regions of the *PMT* genes and the *NAT* or *HYG* resistance markers (Goldstein and McCusker,

1999, Yeast 14:1541-1553; Goldstein *et al.*, 1999, Yeast 15:507-110) were PCR amplified. The primers sequences for the regions flanking the *PMT* genes were designed using the *Pichia pastoris* genome nucleotide sequence obtained from Integrated Genomics, Chicago, IL as a guide. *Pichia pastoris* genomic DNA was used as a template for the Pp*PMT* flanking regions PCR amplification, while *NAT* and *HYG* fragments were PCR amplified using plasmids as described in (Goldstein and McCusker, 1999, *ibid.*; Goldstein *et al.*, 1999, *ibid.*) as templates. Then, in a second PCR reaction, all three first round PCR products were used as templates to generate an overlap product that contained all three fragments as a single linear allele. The final PCR product was then directly employed for transformation. Transformants were selected on YPD medium containing 200 µg/mL of hygromycin or 100 µg/mL of nourseothricin. In each case the proper integration of the mutant allele was confirmed by PCR. The 5 *PMT* knockout strains created were yJC51 (*pmt3Δ,pmt5Δ,pmt6Δ*), yJC55 (*pmt1Δ*), yJC66 (*pmt2Δ*), and yJC65 (*pmt4Δ*). The *PMT* knockout strains were each transformed with plasmid pBK119 encoding the 10 above Kringle 1-3 protein.

Kringle 1-3 protein expression for the transformed yeast strains was carried out at in 15 shake flasks at 24°C with buffered glycerol-complex medium (BMGY) consisting of 1% yeast extract, 2% peptone, 100 mM potassium phosphate buffer pH 6.0, 1.34% yeast nitrogen base, 4 x 10-5 % biotin, and 1% glycerol. The induction medium for protein expression was buffered methanol-complex medium (BMMY) consisting of 1% methanol instead of glycerol in BMGY. Pmt inhibitor Pmti-1, Pmti-2, or Pmti-3 in methanol was added to the growth medium to a final concentration of 0.2 µM, 2 µM, or 20 µM 20 at the time the induction medium was added. Cells were harvested and centrifuged at 2,000 rpm for five minutes. The Pmt inhibitors Pmti-1, Pmti-2, and Pmti-3 are essentially interchangeable, with small variations in ease of use. For example, in the cell culture conditions described, the solubility of Pmti-3 is greater than that of Pmti-1 and Pmti-2 and, therefore, the most desirable of the three.

Seven µL of the supernatant from the yJC53 cultures treated with Pmti-1 or yJC55 was 25 separated by polyacrylamide gel electrophoresis (SDS-PAGE) according to Laemmli, U. K. (1970) Nature 227, 680-685 and then electroblotted onto nitrocellulose membranes (Schleicher & Schuell (now Whatman, Inc., Florham Park, NJ). Kringle 1-3 protein was detected on the Western blots using an anti-His antibody (H-15) from Santa Cruz Biotechnology Inc. (Santa Cruz, CA) and developed using the ImmunoPure Metal Enhanced DAB Substrate Kit (Pierce Biotechnology, Rockford, IL). As shown in 30 Lane 1 of the Western blot shown in Figure 1, Kringle 1-3 protein from untreated *Pichia pastoris* runs as a smear due to the presence of O-glycosylation. However, in contrast, Kringle 1-3 protein from yJC53 (*Pichia pastoris* treated with 2 or 20 µM Pmti-1, lanes 2 and 3, respectively) exhibits a distinct band, due to lack of O-glycosylation, similar to that of Kringle 1-3 protein expressed from yJC55 (a *pmt1Δ* knockout mutant of *Pichia pastoris*) (lanes 4 and 5). Figure 1 further shows that Pmti-1 reduced O-glycosylation to a level similar to that observed in a strain lacking Pmt1.

To measure O-glycosylation reduction by the Pmt inhibitors, the Kringle 1-3 protein was purified from the growth medium using nickel chelation chromatography (Choi *et al.*, 2003, Proc. Natl.

Acad. Sci. U. S. A. 100(9): 5022-5027) and the O-glycans released from and separated from Kringle 1-3 protein by alkaline elimination (beta -elimination) (Harvey, 1999 Mass Spectrometry Reviews 18, 349-451). This process also reduces the newly formed reducing terminus of the released O-glycan (either oligomannose or mannose) to mannitol. The mannitol group thus serves as a unique indicator of each O-glycan. 0.5 nmole or more of Kringle 1-3 protein, contained within a volume of 100 μ L PBS buffer, was required for beta elimination. The sample was treated with 25 μ L alkaline borohydride reagent and incubated at 50°C for 16 hours. About 20 μ L arabitol internal standard was added, followed by 10 μ L glacial acetic acid. The sample was then centrifuged through a Millipore filter containing both SEPABEADS and AG 50W-X8 resin and washed with water. The samples, including wash, were transferred to plastic autosampler vials and evaporated to dryness in a centrifugal evaporator. 150 μ L 1% AcOH/MeOH was added to the samples and the samples evaporated to dryness in a centrifugal evaporator. This last step was repeated five more times. 200 μ L of water was added and 100 μ L of the sample was analyzed by high pH anion-exchange chromatography coupled with pulsed electrochemical detection-Dionex HPLC (HPAEC-PAD). Average O-glycan occupancy was determined based upon the amount of mannitol recovered. The results are summarized in Table 1, which shows that any one of the *Pmt* chemical inhibitors reduced O-linked glycosylation of secreted Kringle 1-3 protein in the *Pichia pastoris* strains that contained intact *PMT1* and *PMT2* genes to a level that was comparable to the level of O-linked glycosylation seen for cells containing deletions of either the *PMT1* or *PMT2* gene. Table 1 also shows that while the protein has two potential mammalian O-linked glycosylation sites, in yeast the protein has about 20 O-linked glycosylation sites.

Table 1

Strain	Relevant Genotype	Treatment	O-Glycan Occupancy ¹
yJC53	Wild-Type	0	20
		2 μ M Pmti-1	9
yJC51	<i>pmtΔ3,Δ5,Δ6</i>	0	17
		2 μ M Pmti-1	6
		20 μ M Pmti-1	4
		0.2 μ M Pmti-2	3
		2 μ M Pmti-2	2
		0.2 μ M Pmti-3	6
		2 μ M Pmti-3	4
yJC55	<i>pmtΔ1</i>	0	3
		2 μ M Pmti-1	2
		20 μ M Pmti-1	2
yJC66	<i>pmtΔ2</i>	0	4
		2 μ M Pmti-1	4
		20 μ M Pmti-1	4
yJC65	<i>pmtΔ4</i>	0	18
		2 μ M Pmti-1	7
		20 μ M Pmti-1	4

¹average number of O-linked mannose chains per protein.

EXAMPLE 3

In this example, yeast cells transformed with DNA encoding the *T. reesei* α -mannosidase results in production of proteins with reduced O-glycosylation and that the O-glycosylation was further reduced when the cells were also incubated in the presence of a Pmt inhibitor.

The H + L chains of an anti-Her2 monoclonal antibody were expressed in *Pichia pastoris* strains GS115 (WT) and GS115 that was genetically engineered to co-expressed *T. reesei* α -mannosidase (+Trman). GS115 is available from Invitrogen (Carlsbad, CA) and, with the exception of a *HIS4* mutation to enable his4 selection, has an essentially wild type phenotype. The H + L chains were expressed as two separate genes from plasmid pJC284, which was derived from Invitrogen plasmid pAO815.

The H + L genes were generated using anti-Her2 antibody sequences obtained from GenBank. The GenBank accession number for the L chain is 1N8Z_A and the GenBank accession number for the H chain variable region plus CH1 domain is 1N8Z_B. The GenBank accession number

for the H chain Fc region is BC092518. Both the H and L chain DNA sequences were codon optimized according to *Pichia pastoris* codon usage to enhance translation in *Pichia pastoris*. Optimization of codons for use in *Pichia* sp. is well known in the art and has been described in, for example, Outchkourov *et al.*, 2002, *Protein Expr. Purif.* 24:18-24; Sharp and Li, 1987, *Nucleic Acids Res.* 15:1281-95; Woo JH, Liu *et al.*, 2002, *Protein Expression and Purification* 25:270-282, and, Nakamura, *et al.*, 2000, *Nucleic Acids Res.* 28:292. Constant regions of the H chain (human IgG1) and L chain (human Kappa) were synthesized by GeneArt Inc., Regensburg, Germany. Variable regions were made in-house using oligonucleotides purchased from IDT Inc. (Coralville, IA) in an overlapping PCR method. Full length H and L chains were assembled by overlapping PCR, and resulting H and L chains were cloned 5 into pCR2.1 TOPO vector (Invitrogen, La Jolla, CA) to generate pDX344 and pDX349, respectively. H + L chains from pDX344 and pDX349 were combined with GAPDH promoters and AOX1 terminator sequences in vector pDX580 (backbone derived from Invitrogen vector pGAPZA). Finally, the H + L chain expression cassettes were subcloned from pDX580 into vector pJC284. The nucleotide sequence 10 of the codon-optimized DNA encoding the light chain is shown in SEQ ID NO:7 and the nucleotide sequence of the codon-optimized DNA encoding the heavy chain is shown in SEQ ID NO:8. Plasmid pJC284 has GAPDH promoters for expressing the H + L genes and an intact his4 gene for selection of 15 transformants in strain GS115 and GS115(+Trman). Yeast strains GS115 and GS115(+Trman) were transformed with pJC285 and transformants with the plasmid integrated into the genome at the his4 locus were isolated to produce strains that produced the anti-Her2 antibody.

20 Construction of strain GS115(+Trman) was as follows. The *Trichoderma reesei* α -1,2-mannosidase was expressed from an expression cassette in plasmid pJC285. Plasmid pJC285 was derived from Invitrogen vector pGAPZA, which has the Zeocin resistant gene as the selectable marker, and contains an expression cassette comprising DNA encoding the *T. reesei* α -1,2-mannosidase catalytic domain (SEQ ID NO:9) with the first 84 base pairs encoding its signal sequence replaced with a DNA 25 encoding the *Saccharomyces cerevisiae* α MAT pre signal sequence (SEQ ID NO:10), which encodes just the ER-targeting amino acids, operably linked to DNA comprising the *Pichia pastoris* GAPH promoter (SEQ ID NO:11) at the 5' end and DNA comprising the *Pichia pastoris* AOX1 transcription termination sequence (SEQ ID NO:12) at the 3' end. The nucleotide sequence of the complete expression cassette is set forth in SEQ ID NO:13. Yeast strain GS115 was transformed with pJC285 and transformants with 30 the plasmid integrated into the genome at the GAPDH locus were isolated to produce strain GS115(+Trman).

Duplicate cultures of the strains were cultured in 200 mL of buffered dextrose-complex medium (BMDY) consisting of 1% yeast extract, 2% peptone, 100mM potassium phosphate pH 6.0, 1.34% Yeast Nitrogen Base, .00004% biotin, 2% dextrose, and with or without Pmti-2 at 0.3 or 0.03 μ M. 35 Following 72 hours of growth, culture supernatants were collected and centrifuged to remove yeast cells. Antibody in the remaining supernatant fraction (about 200 mL) was purified over a Protein A column and subjected to O-glycan analysis as described in Example 2. In addition to the mannitol assay, the average

length of O-linked mannose chains was determined by chromatographic analysis without hydrolysis. The results are summarized in Table 2.

There are 14 yeast O-glycan sites on the antibody. When the antibody was produced in the wild-type GS115 strain, all 14 O-glycan sites have glycan structures with 8% of the sites having just one mannose, 39% having a two mannose chain, 43% having a three mannose chain, and 9% having a four mannose chain (See Table 2). However, when the antibody was produced in wild-type cells treated with the chemical inhibitor Pmti-2, only two of the 14 O-glycan sites were occupied and for 76% of the two sites, the mannose chain had only one mannose residue. Neither of the two sites had a mannose chain with three or four mannose residues. It should be noted that the analysis did not determine which 5 two of the 14 sites were occupied. Either any combination of two O-glycan sites per antibody molecule were occupied or particular O-glycan sites are preferentially occupied. In the latter case, to provide 10 antibodies (or other proteins) completely devoid of O-glycans, the amino acid sequences comprising the preferred O-glycan sites can be modified to amino acid sequences that eliminates O-linked glycosylation at the sites.

15 Table 2 further shows that when the antibody was produced in cells that included DNA encoding the *Trichoderma reesei* α -1,2-mannosidase (strain GS115(+Trman)), only four of the 14 O-glycan sites were occupied and for 95% of the four sites, the mannose chain had only one mannose residue. None of the four sites had a mannose chain with three or four mannose residues. If particular 20 sites are preferentially O-glycosylated, to provide antibodies (or other proteins) completely devoid of O-glycans, the amino acid sequences comprising the preferred O-glycan sites can be modified to amino acid sequences that eliminates O-glycosylation at the sites.

Finally, Table 2 shows that when the antibody was produced in cells that included DNA encoding the *Trichoderma reesei* α -1,2-mannosidase and in the presence of Pmti-2, only one of the 14 O-glycan sites was occupied and for 91% of the sites, the mannose chain had only one mannose residue. 25 No mannose chain had three or four mannose residues. If only one or only a few sites are preferentially O-glycosylated, to provide antibodies (or other proteins) completely devoid of O-glycans, the amino acid sequences comprising the preferred O-glycan sites can be modified to amino acid sequences that eliminate O-glycosylation at the sites

Table 2 further shows that 0.3 uM of Pmti inhibitor is sufficient to reduced the 30 occupancy by about 86% and the chain length for 76% of the molecules to one mannose while allowing the culture to grow. Including the *Trichoderma reesei* α -1,2-mannosidase allowed the amount of Pmti inhibitor to be reduced by 10 fold and the occupancy reduced to 93% and the chain length for 87% of the molecules to one mannose. These results show that using an amount of Pmti inhibitor that does not kill the cells is sufficient to produce glycoproteins having reduced O-linked glycosylation. These results 35 further show that the Pmti inhibitor and the α -mannosidase appear to synergistically reduce the amount of O-linked glycosylation.

Table 2

Strain	Occupancy	Man1	Man2	Man3	Man4
GS115	14	8	39	43	9
GS115 + 0.3 μ M Pmti-2	2	76	24	0	0
GS115(+Trman)	4	95	5	0	0
GS115(+Trman) + 0.3 μ M Pmti-2	1	91	9	0	0
GS115(+Trman) + 0.03 μ M Pmti-2	1	87	13	0	0

Seven μ L of the supernatant for each of the above were reduced and subjected to SDS-PAGE and Western blotting using an HRP-conjugated anti-human IgG (H&L) to detect H and L chains.

5 The results are shown in Figure 2. The hyper O-glycosylated H chain is the slowest migrating band visible in the first pair of lanes in Figure 2. Figure 2 shows that there is a decrease in the amount of O-glycosylated heavy chain when the antibody was coexpressed with *Trichoderma reesei* α -1,2-mannosidase or the cells expressing the antibody was incubated in the presence of the Pmti-2 inhibitor, or when the antibody was coexpressed with *Trichoderma reesei* α -1,2-mannosidase and the cells expressing both

10 proteins was incubated in the presence of the Pmti-2 inhibitor.

15 While the present invention is described herein with reference to illustrated embodiments, it should be understood that the invention is not limited hereto. Those having ordinary skill in the art and access to the teachings herein will recognize additional modifications and embodiments within the scope thereof. Therefore, the present invention is limited only by the claims attached herein.

The claims defining the invention are as follows:

1. A method of producing a protein having reduced O-linked glycosylation comprising:

5 (a) providing a nucleic acid encoding a glycoprotein and a second nucleic acid encoding an α 1,2-mannosidase;

(b) introducing the nucleic acid and the second nucleic acid into a host cell and growing the host cell containing the nucleic acid and the second nucleic acid to produce a culture of the host cell;

10 (c) contacting the culture of the host cell with one or more inhibitors of Pmt-mediated O-linked glycosylation; and

(d) isolating the glycoprotein produced by the host cell in the presence of the one or more inhibitors of Pmt-mediated O-linked glycosylation to produce the protein having reduced O-linked glycosylation.

15 2. The method of Claim 1 wherein the α 1,2-mannosidase comprises the catalytic domain of an α 1,2-mannosidase operatively linked to a nucleic acid sequence encoding a cellular targeting signal peptide not normally associated with the catalytic domain.

20 3. The method of Claim 1 wherein the culture is grown for a time sufficient to provide a multiplicity of the host cells having the nucleic acid and second nucleic acid before contacting the culture with the one or more inhibitors of Pmt-mediated O-linked glycosylation.

4. The method of Claim 1 wherein the culture is grown in the presence of the one or more inhibitors of Pmt-mediated O-linked glycosylation.

25 5. The method of Claim 1 wherein the culture is grown for a time sufficient to provide a multiplicity of the host cells having the nucleic acid and the second nucleic acid before contacting the culture with the one or more inhibitors of Pmt-mediated O-linked glycosylation and an inducer of the promoters to induce expression of the protein and the α 1,2-mannosidase and isolating the glycoprotein produced by the host cell in the presence of the one or more inhibitors and the inducer to produce the protein having reduced O-linked glycosylation.

30 6. The method of claim 1 wherein the nucleic encoding the glycoprotein and the second nucleic acid encoding the α 1,2-mannosidase are operably linked to an inducible promoter.

35 7. The method of Claim 6 wherein the culture is contacted with an inducer of the promoters to induce expression of the protein and the α 1,2-mannosidase for a time before contacting the culture with the one or more inhibitors of Pmt-mediated O-linked

glycosylation and isolating the glycoprotein produced by the host cell in the presence of the one or more inhibitors and the inducer to produce the protein having reduced O-linked glycosylation.

8. The method of Claim 1 wherein the culture is grown for a time sufficient to 5 provide a multiplicity of the host cells before inducing expression of the protein and the one or more α 1,2-mannosidase enzymes to produce the glycoprotein having reduced O-linked glycosylation.

9. The method of Claim 1 wherein expression of the protein is induced for a time before inducing expression of the one or more α 1,2-mannosidase enzymes to 10 produce the glycoprotein having reduced O-linked glycosylation.

10. The method of Claim 1 wherein expression of the one or more α 1,2-mannosidase enzymes capable of inhibiting O-linked glycosylation is induced for a time before inducing expression of the glycoprotein to produce the glycoprotein having reduced O-linked glycosylation.

11. The method of Claim 1 wherein the one or more inhibitors are selected from 15 the group consisting of: 5-[[3,4-bis(phenylmethoxy) phenyl]methylene]-4-oxo-2-thioxo-3-thiazolidineacetic Acid; 5-[[3-(1-Phenylethoxy)-4-(2-phenylethoxy)]phenyl]methylene]-4-oxo-2-thioxo-3-thiazolidineacetic Acid; and 5-[[3-(1-Phenyl-2-hydroxy)ethoxy)-4-(2-phenylethoxy)]phenyl]methylene]-4-oxo-2-thioxo-3-thiazolidineacetic Acid. 20

12. The method of Claim 1 wherein the α 1,2-mannosidase is from *Trichoderma reesei*, *Saccharomyces* sp., or *Aspergillus* sp.

13. The method of Claim 1, wherein the host cell is a fungal cell or yeast cell.

14. The method of Claim 1, wherein the host cell is selected from the group 25 consisting of *K. lactis*, *Pichia pastoris*, *Pichia methanolica*, and *Hansenula*.

15. The method of Claim 1, wherein the host cell is a yeast or filamentous fungal cell that has been genetically modified to produce glycoproteins with a predominant N-glycan glycoform.

16. The method of Claim 1 wherein the host cells have been genetically modified 30 to produce glycoproteins in which the N-glycosylation pattern is human-like or humanized.

17. A glycoprotein produced by the method of any one of Claims 1-16.

18. A glycoprotein composition comprising the glycoprotein of claim 17 wherein 35 the glycoprotein has a human-like or humanized N-glycosylation pattern and reduced O-linked glycosylation.

19. The glycoprotein composition of Claim 18 wherein the human-like or humanized N-glycosylation pattern comprises predominantly an N-glycan structure selected from the group consisting of Man5GlcNAc2, Man3GlcNAc2, GlcNAcMan5GlcNAc2, GlcNAcMan3GlcNAc2, GlcNAc2Man3GlcNAc2, 5 GalGlcNAcMan5GlcNAc2, Gal(GlcNAc)2 Man5GlcNAc2, (GalGlcNAc)2Man5GlcNAc2, NANAGalGlcNAcMan3GlcNAc2, NANA2Gal2GlcNAcMan3GlcNAc2, and GalGlcNAcMan3GlcNAc2.

20. A method according to Claim 1 and substantially as hereinbefore described with reference to any one of the Examples or Figures.

10

Dated 16 March, 2012

GlycoFi, Inc

Patent Attorneys for the Applicant/Nominated Person
SPRUSON & FERGUSON

1/2

1 2 3 4 5




Fig. 1

2/2




FIG. 2

SEQUENCE LISTING

<110> GlycoFi, Inc.
Bobrowiza, Piotr
Cook, James W.
Kett, Warren

<120> PRODUCTION OF PROTEINS WITH REDUCED
O-GLYCOSYLATION

<130> PCT GF0005Y

<150> 60/737,108
<151> 2005-11-15

<160> 13

<170> FastSEQ for Windows Version 4.0

<210> 1
<211> 288
<212> PRT
<213> Artificial Sequence

<220>
<223> Kringle 1-3 protein

<400> 1
Ser Glu Cys Lys Thr Gly Asn Gly Lys Asn Tyr Arg Gly Thr Met Ser
1 5 10 15
Lys Thr Lys Asn Gly Ile Thr Cys Gln Lys Trp Ser Ser Thr Ser Pro
20 25 30
His Arg Pro Arg Phe Ser Pro Ala Thr His Pro Ser Glu Gly Leu Glu
35 40 45
Glu Asn Tyr Cys Arg Asn Pro Asp Asn Asp Pro Gln Gly Pro Trp Cys

50 55 60
Tyr Thr Thr Asp Pro Glu Lys Arg Tyr Asp Tyr Cys Asp Ile Leu Glu
65 70 75 80
Cys Glu Glu Glu Cys Met His Cys Ser Gly Glu Asn Tyr Asp Gly Lys
85 90 95
Ile Ser Lys Thr Met Ser Gly Leu Glu Cys Gln Ala Trp Asp Ser Gln
100 105 110
Ser Pro His Ala His Gly Tyr Ile Pro Ser Lys Phe Pro Asn Lys Asn
115 120 125
Leu Lys Lys Asn Tyr Cys Arg Asn Pro Asp Arg Glu Leu Arg Pro Trp
130 135 140
Cys Phe Thr Thr Asp Pro Asn Lys Arg Trp Glu Leu Cys Asp Ile Pro
145 150 155 160
Arg Cys Thr Thr Pro Pro Ser Ser Gly Pro Thr Tyr Gln Cys Leu
165 170 175
Lys Gly Thr Gly Glu Asn Tyr Arg Gly Asn Val Ala Val Thr Val Ser
180 185 190
Gly His Thr Cys Gln His Trp Ser Ala Gln Thr Pro His Thr His Ser
195 200 205
Arg Thr Pro Glu Asn Phe Pro Cys Lys Asn Leu Asp Glu Asn Tyr Cys
210 215 220
Arg Asn Pro Asp Gly Lys Arg Ala Pro Trp Cys His Thr Thr Asn Ser
225 230 235 240
Gln Val Arg Trp Glu Tyr Cys Lys Ile Pro Ser Cys Asp Ser Ser Pro
245 250 255
Val Ser Thr Glu Gln Leu Ala Pro Thr Ala Pro Pro Glu Leu Thr Pro
260 265 270
Val Val Gln Asp Gly Gly His His His His His His His His His
275 280 285

<210> 2
<211> 33
<212> DNA
<213> Artificial Sequence

<220>

<223> forward primer K1-3/UP

<400> 2

cggaattctc agagtgc aag actggaaata gaa

33

<210> 3

<211> 39

<212> DNA

<213> Artificial Sequence

<220>

<223> reverse primer K1-3/LP1

<400> 3

atgatgatga ccaccaccgt cctggaccac aggggttag

39

<210> 4

<211> 39

<212> DNA

<213> Artificial Sequence

<220>

<223> reverse primer K1-3/LP2

<400> 4

ttaatgatga ttagatgatgat gatgatgatg accaccacc

39

<210> 5

<211> 36

<212> DNA

<213> Artificial Sequence

<220>

<223> forward primer K3f (Asn to Ser)

<400> 5

acccttcaca cacattctag gacaccagaa aacttc

36

<210> 6
<211> 27
<212> DNA
<213> Artificial Sequence

<220>
<223> reverse primer K3r

<400> 6
ctgtgcactc cagtgcgtac aggtgtg

27

<210> 7
<211> 645
<212> DNA
<213> Artificial Sequence

<220>
<223> Codon-optimized sequence encoding light chain

<400> 7
gacattcaga tgacacagtc tccatcttct ttgtccgctt ccgtcggtga tagagttact 60
atcacctgta gagcttccca agacgtcaac accgctgtcg cctggtagcca acagaagccca 120
ggtaaggctc caaaaactttt gatctactct gctctttct tggactccgg tgttccatcc 180
agatttctg gttctagatc cggtaccgac ttcacccgttga ccatctcttc cttgcaacca 240
gaagacttcg ctacctacta ctgtcaacaa cactacacta ctccctccaaac tttcggtcaa 300
ggaactaagg ttgagattaa gagaactgtt gctgtccat ccgtttcat tttcccacca 360
tccgacgaac aattgaagtc tggtagatc tccgttggtt gtttggtagaa caacttctac 420
ccaagagagg ctaagggtca gtggaaagggtt gacaacgctt tgcaatccgg taactcccaa 480
gaatccgtta ctgagcagga ttctaaaggat tccacttact cttgtccctc cactttgact 540
ttgtccaaagg ctgattacga gaagcacaag gtttacgcat gcgagggtac acatcagggt 600
ttgtcctccc cagttactaa gtccttcaac agaggagagt gttaa

645

<210> 8
<211> 1353
<212> DNA
<213> Artificial Sequence

<220>

<223> Codon-optimized sequence encoding heavy chain

<400> 8

gaggtccaat tgggtgaatc tgggtggaggt ttgggtccaaac caggtggatc tctgagactt 60
tcttggctg cctctgggtt caacattaag gatacttaca tccactgggt tagacaggct 120
ccaggttaagg gtttggagtg gtttgctaga atctacccaa ccaacggta caccagatac 180
gctgattccg ttaagggttag attcaccatt tccgctgaca cttccaagaa cactgcttac 240
ttgcaaataa actctttgag agctgaggac actgccgtct actactgttc cagatgggt 300
ggtgacgggtt tctacgcccattt ggactactgg ggtcaaggta cttgggttac tgtctttcc 360
gcttctacta agggaccatc cggttttcca ttggctccat cctctaagtc tacttccgg 420
ggtaactgctg ctttggatg tttggtaag gactacttcc cagagcctgt tactgtttct 480
tggaactccg gtgctttgac ttctgggttt cacacttcc cagctgtttt gcaatcttcc 540
ggtttgtact ctttgcctc cgttgttact gttccatcct cttccctggg tactcagact 600
tacatctgtttaacca caagccatcc aacactaagg ttgacaagaa gggtgagcca 660
aagtccctgtg acaagacaca tacttgccttcc ccatgtccag ctccagaatt gttgggtgg 720
ccatccgttt tcttggttccc accaaagcca aaggacactt tgatgatctc cagaactcca 780
gaggttacat gtgttggatg tgacgtttctt caccggacc cagaggtaaa gttcaactgg 840
tacgttgacg gtgttgaagt tcacaacgct aagactaaggc caagagagga gcagtacaac 900
tccacttaca gagttgttcc cgttttgact gtttgcacc aggattgggtt gaacggaaag 960
gagtacaagt gtaagggttc caacaagggtt ttgccagctc caatcgaaaa gactatctcc 1020
aaggctaagg gtcaaccaag agagccacag gtttacactt tgccaccatc cagagatgag 1080
ttgactaaga accaggttcc ttgacttgtt ttggtaaag gattctaccc atccgacatt 1140
gctgttgagt gggaaatctaa cggtaacca gagaacaact acaagactac tccaccagtt 1200
ttggattctg acgggttcctt cttcttgcac tccaaatgttca ctgttgacaa gtccagatgg 1260
caacagggtt acgttttctc ctgttccgtt atgcattgagg ctttgcacaa ccactacact 1320
caaaagtccct tgcgtttgtc cccaggtaag taa 1353

<210> 9

<211> 500

<212> DNA

<213> Artificial Sequence

<220>

<223> Pichia pastoris GAPDH promoter

<400> 9

agatctttt ttagaaatg tctgggtgc ctgcgtccat caggtggcca tctctgaaat 60
atctggctcc gttgcaactc cgaacgacct gctggcaacg taaaattctc cggggtaaaa 120
cttaaatgtg gagtaatgga accagaaacg tcttccct tctctctcct tccaccgccc 180
gttaccgtcc ctaggaaatt ttactctgct ggagagcttc ttctacggcc cccttgcagc 240
aatgctcttc ccagcattac gttgcggta aaacggaggt cgtgtacccg acctagcagc 300
ccagggatgg aaaagtcccg gccgtcgctg gcaataatag cgggcggacg catgtcatga 360
gattattgga aaccaccaga atcgaatata aaaggcgaac acctttccca attttggttt 420
ctcctgaccc aaagacttta aatttaattt attttgcctt atttcaatca attgaacaac 480
tatttcgaaa cgaggaaatt 500

<210> 10

<211> 57

<212> DNA

<213> Artificial Sequence

<220>

<223> *Saccharomyces cerevisiae* alpha-MAT pre signal
sequence

<400> 10

atgagatttc cttcaatttt tactgctgtt ttattcgcag catcctccgc attagct 57

<210> 11

<211> 1494

<212> DNA

<213> Artificial Sequence

<220>

<223> *Trichoderma reesei* alph-1,2-mannosidase catalytic
domain

<400> 11

cgcgcggat ctcccaaccc tacgagggcg gcagcagtca aggccgcatt ccagacgtcg 60
tggAACGCTT accaccattt tgccTTccc catgacgacc tccacccggc cagcaacagc 120
tttgcgtatg agagaaacgg ctggggctcg tcggcaatcg atggcttgga cacggctatc 180
ctcatggggg atgcccacat tgtgaacacg atccttcagt atgtaccgca gatcaacttc 240

accacgactg cggttgccaa ccaaggcatc tccgtgttcg agaccaacat tcggcaccc 300
ggtggcctgc ttctgccta tgacctgttg cgaggtcctt tcagtcctt ggccacaaac 360
cagaccctgg taaacagcct tctgaggcag gctcaaacac tggccaacgg cctcaagggtt 420
gcgttccacca ctcccagcgg tgcgtccggac cctaccgtct tcttcaaccc tactgtccgg 480
agaagtggtg catctagcaa caacgtcgct gaaattggaa gcctggtgct cgagtggaca 540
cggttggcg acctgacggg aaacccgcag tatgcccage ttgcgcagaa gggcgagtcg 600
tatctcctga atccaaaggg aagcccgag gcatggcctg gcctgattgg aacgtttgc 660
agcacgagca acgttacctt tcaggatagc agcggcagct ggtccggcct catggacagc 720
ttctacgagt acctgatcaa gatgtacctg tacgaccggg ttgcgttgc acactacaag 780
gatcgctggg tccttgctgc cgactcgacc attgcgcate tcgcctctca cccgtcgacg 840
cgcaaggact tgacctttt gtcttcgtac aacggacagt ctacgtcgcc aaactcagga 900
catttggcca gtttgccgg tggcaacttc atcttggag gcattctcct gaacgagcaa 960
aagtacattg actttggaat caagcttgcc agctcgact ttgccacgta caaccagacg 1020
gcttctggaa tcggccccga aggcttcgcg tgggtggaca gcgtgacggg cgccggcggc 1080
tcgcccctt cgtcccgatc cgggttctac tcgtcggcag gattctgggt gacggcacccg 1140
tattacatcc tgccggcggg gacgctggag agcttgcact acgcataacgg cgtcacgggc 1200
gactccaagt ggcaggaccc ggcgtggaa gcgttcagtg ccattgagga cgcacgtccgc 1260
gccggcagcg cgtactcgtc catcaacgcgt gtcgacgcagg ccaacggcgg ggggtgcctct 1320
gacgatatgg agagcttctg gtttgccag ggcgtcaagt atgcgtaccc gatctttgcg 1380
gaggagtcgg atgtgcaggt gcaggccaaac ggcgggaaca aatttgcctt taacacggag 1440
gcgcacccct ttagcatccg ttcatcatca cgacggggcg gccaccttgc ttaa 1494

<210> 12

<211> 486

<212> DNA

<213> Artificial Sequence

<220>

<223> Pichia pastoris AOX1 transcription termination sequence

<400> 12

```
aagggcgaat tcaattcacg tggcccgagcc ggccgtctcg gatcggtacc tcgagcccg 60
gcggccgcca gcttgggccc gaacaaaaac tcatctcaga agaggatctg aatagcgccg 120
tcgaccatca tcatcatcat cattgagtt tagccttaga catgactgtt cctcagttca 180
agttgggcac ttacgagaag accggtcttg ctagattcta atcaagagga tgtcagaatg 240
ccatttccct qaaqatqca qqcttcattt ttqatacttt ttatqtatc acctatataq 300
```

tataggattt tttttgtcat tttgtttctt ctcgtacgag cttgctcctg atcagcctat 360
ctcgcagctg atgaatatct tgtggtaggg gtttgggaaa atcattcagag tttgatgttt 420
ttcttggtat ttcccactcc tcttcagagt acagaagatt aagtgagacc ttcgtttgt 480
cggatc 486

<210> 13
<211> 2550
<212> DNA
<213> Artificial Sequence

<220>
<223> Trichoderma reesei alpha-1,2-mannosidase expression
cassette

<221> promoter
<222> (1) ... (507)
<223> GAPDH promoter sequence

<221> sig_peptide
<222> (508) ... (564)
<223> encodes *Saccharomyces cerevisiae* alpha-MAT pre
signal sequence

<221> CDS
<222> (565) ... (2058)
<223> encodes Trichoderma reesei alpha-1,2-mannosidase
catalytic domain

<221> terminator
<222> (2065) ... (2550)
<223> *Pichia pastoris* AOX1 transcription termination
sequence

<400> 13
agatctttt tgttagaaatg tcttgggtgc ctcgtccat caggttagcca tctctgaaat 60
atctggctcc gttgcaactc cgaacgacct gctggcaacg taaaattctc cggggtaaaa 120
ctttaatgtg gagtaatgga accagaaacg tctctccct tctctctcct tccaccgccc 180

gttaccgtcc ctaggaaatt ttactctgct ggagagcttc ttctacggcc cccttgcagc 240
 aatgctcttc ccagcattac gttgcgggta aaacggaggt cgtgtacccg accttagcagc 300
 ccagggatgg aaaagtcccg gccgtcgctg gcaataatag cgggcggacg catgtcatga 360
 gattattgga aaccaccaga atcgaatata aaaggcgaac acctttccca attttggttt 420
 ctccctgaccc aaagacttta aatttaattt atttgtccct atttcaatca attgaacaac 480
 tatttcgaaa cgagggattt cgaaacg atg aga ttt cct tca att ttt act gct 534

Met Arg Phe Pro Ser Ile Phe Thr Ala

-15

gtt tta ttc gca gca tcc tcc gca tta gct cgc gcc gga tct ccc aac 582
 Val Leu Phe Ala Ala Ser Ser Ala Leu Ala Arg Ala Gly Ser Pro Asn
 -10 -5 1 5

cct acg agg gcg gca gca gtc aag gcc gca ttc cag acg tcg tgg aac 630
 Pro Thr Arg Ala Ala Ala Val Lys Ala Ala Phe Gln Thr Ser Trp Asn
 10 15 20

gct tac cac cat ttt gcc ttt ccc cat gac gac ctc cac ccg gtc agc 678
 Ala Tyr His His Phe Ala Phe Pro His Asp Asp Leu His Pro Val Ser
 25 30 35

aac agc ttt gat gat gag aga aac ggc tgg ggc tcg tcg gca atc gat 726
 Asn Ser Phe Asp Asp Glu Arg Asn Gly Trp Gly Ser Ser Ala Ile Asp
 40 45 50

ggc ttg gac acg gct atc ctc atg ggg gat gcc gac att gtg aac acg 774
 Gly Leu Asp Thr Ala Ile Leu Met Gly Asp Ala Asp Ile Val Asn Thr
 55 60 65 70

atc ctt cag tat gta ccg cag atc aac ttc acc acg act gcg gtt gcc 822
 Ile Leu Gln Tyr Val Pro Gln Ile Asn Phe Thr Thr Ala Val Ala
 75 80 85

aac caa ggc atc tcc gtg ttc gag acc aac att cgg tac ctc ggt ggc 870
 Asn Gln Gly Ile Ser Val Phe Glu Thr Asn Ile Arg Tyr Leu Gly Gly
 90 95 100

ctg ctt tct gcc tat gac ctg ttg cga ggt cct ttc agc tcc ttg gcg 918
 Leu Leu Ser Ala Tyr Asp Leu Leu Arg Gly Pro Phe Ser Ser Leu Ala

105 110 115

aca aac cag acc ctg gta aac agc ctt ctg agg cag gct caa aca ctg 966
 Thr Asn Gln Thr Leu Val Asn Ser Leu Leu Arg Gln Ala Gln Thr Leu
 120 125 130

gcc aac ggc ctc aag gtt gcg ttc acc act ccc agc ggt gtc ccg gac 1014
 Ala Asn Gly Leu Lys Val Ala Phe Thr Thr Pro Ser Gly Val Pro Asp
 135 140 145 150

cct acc gtc ttc ttc aac cct act gtc cgg aga agt ggt gca tct agc 1062
 Pro Thr Val Phe Phe Asn Pro Thr Val Arg Arg Ser Gly Ala Ser Ser
 155 160 165

aac aac gtc gct gaa att gga agc ctg gtg ctc gag tgg aca cgg ttg 1110
 Asn Asn Val Ala Glu Ile Gly Ser Leu Val Leu Glu Trp Thr Arg Leu
 170 175 180

agc gac ctg acg gga aac ccg cag tat gcc cag ctt gcg cag aag ggc 1158
 Ser Asp Leu Thr Gly Asn Pro Gln Tyr Ala Gln Leu Ala Gln Lys Gly
 185 190 195

gag tcg tat ctc ctg aat cca aag gga agc ccg gag gca tgg cct ggc 1206
 Glu Ser Tyr Leu Leu Asn Pro Lys Gly Ser Pro Glu Ala Trp Pro Gly
 200 205 210

ctg att gga acg ttt gtc agc acg agc aac ggt acc ttt cag gat agc 1254
 Leu Ile Gly Thr Phe Val Ser Thr Ser Asn Gly Thr Phe Gln Asp Ser
 215 220 225 230

agc ggc agc tgg tcc ggc ctc atg gac agc ttc tac gag tac ctg atc 1302
 Ser Gly Ser Trp Ser Gly Leu Met Asp Ser Phe Tyr Glu Tyr Leu Ile
 235 240 245

aag atg tac ctg tac gac ccg gtt gcg ttt gca cac tac aag gat cgc 1350

Lys Met Tyr Leu Tyr Asp Pro Val Ala Phe Ala His Tyr Lys Asp Arg
 250 255 260

tgg gtc ctt gct gcc gac tcg acc att gcg cat ctc gcc tct cac ccg 1398
 Trp Val Leu Ala Ala Asp Ser Thr Ile Ala His Leu Ala Ser His Pro
 265 270 275

tcg acg cgc aag gac ttg acc ttt ttg tct tcg tac aac gga cag tct 1446
 Ser Thr Arg Lys Asp Leu Thr Phe Leu Ser Ser Tyr Asn Gly Gln Ser
 280 285 290

acg tcg cca aac tca gga cat ttg gcc agt ttt gcc ggt ggc aac ttc 1494
 Thr Ser Pro Asn Ser Gly His Leu Ala Ser Phe Ala Gly Gly Asn Phe
 295 300 305 310

atc ttg gga ggc att ctc ctg aac gag caa aag tac att gac ttt gga 1542
 Ile Leu Gly Gly Ile Leu Leu Asn Glu Gln Lys Tyr Ile Asp Phe Gly
 315 320 325

atc aag ctt gcc agc tcg tac ttt gcc acg tac aac cag acg gct tct 1590
 Ile Lys Leu Ala Ser Ser Tyr Phe Ala Thr Tyr Asn Gln Thr Ala Ser
 330 335 340

gga atc ggc ccc gaa ggc ttc gcg tgg gtg gac agc gtg acg ggc gcc 1638
 Gly Ile Gly Pro Glu Gly Phe Ala Trp Val Asp Ser Val Thr Gly Ala
 345 350 355

ggc ggc tcg ccc tcg tcc cag tcc ggg ttc tac tcg tcg gca gga 1686
 Gly Gly Ser Pro Pro Ser Ser Gln Ser Gly Phe Tyr Ser Ser Ala Gly
 360 365 370

ttc tgg gtg acg gca ccg tat tac atc ctg cgg ccg gag acg ctg gag 1734
 Phe Trp Val Thr Ala Pro Tyr Tyr Ile Leu Arg Pro Glu Thr Leu Glu
 375 380 385 390

agc ttg tac tac gca tac cgc gtc acg ggc gac tcc aag tgg cag gac 1782
 Ser Leu Tyr Tyr Ala Tyr Arg Val Thr Gly Asp Ser Lys Trp Gln Asp

395	400	405	
ctg gcg tgg gaa gcg ttc agt gcc att gag gac gca tgc cgc gcc ggc 1830			
Leu Ala Trp Glu Ala Phe Ser Ala Ile Glu Asp Ala Cys Arg Ala Gly			
410	415	420	
agc gcg tac tcg tcc atc aac gac gtg acg cag gcc aac ggc ggg ggt 1878			
Ser Ala Tyr Ser Ser Ile Asn Asp Val Thr Gln Ala Asn Gly Gly Gly			
425	430	435	
gcc tct gac gat atg gag agc ttc tgg ttt gcc gag gcg ctc aag tat 1926			
Ala Ser Asp Asp Met Glu Ser Phe Trp Phe Ala Glu Ala Leu Lys Tyr			
440	445	450	
gcg tac ctg atc ttt gcg gag gag tcg gat gtg cag gtg cag gcc aac 1974			
Ala Tyr Leu Ile Phe Ala Glu Glu Ser Asp Val Gln Val Gln Ala Asn			
455	460	465	470
ggc ggg aac aaa ttt gtc ttt aac acg gag gcg cac ccc ttt agc atc 2022			
Gly Gly Asn Lys Phe Val Phe Asn Thr Glu Ala His Pro Phe Ser Ile			
475	480	485	
cgt tca tca tca cga cgg ggc cac ctt gct taa ttaaggagg 2068			
Arg Ser Ser Arg Arg Gly Gly His Leu Ala *			
490	495		
gcgaattcaa ttcacgtggc ccagccggcc gtctcgatc ggtacctcg a cccggcg 2128			
ccgcccagttt gggcccgaaac aaaaactcat ctcagaagag gatctgaata ggcgcgtcg 2188			
ccatcatcat catcatcatt gagtttttagc ctttagacatg actgttcctc agttcaagtt 2248			
gggcacttac gagaagaccg gtcttgctag attctaatca agaggatgtc agaatgccat 2308			
ttgcctgaga gatgcaggct tcattttga tactttta tttgtAACCT atatagtata 2368			
ggatTTTTT tgtcattttg tttcttctcg tacgagcttgc tcctgatca gcctatctcg 2428			
cagctgatga atatctgtg gtaggggttt gggaaatca ttgcagtttgc atgttttct 2488			
tggtatTTCC cactcctctt cagagtacag aagattaagt gagaccttcg tttgtgcgg 2548			
tc 2550			