



(51) International Patent Classification:

A61K 31/4458 (2006.01) A61B 5/046 (2006.01)
A61B 5/0464 (2006.01) A61M 15/00 (2006.01)

(21) International Application Number:

PCT/US2017/016018

(22) International Filing Date:

1 February 2017 (01.02.2017)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/289,473 1 February 2016 (01.02.2016) US

(71) Applicant: INCARDA THERAPEUTICS, INC.
[US/US]; 150 North Hill Drive #24, Brisbane, California
94005 (US).(72) Inventors: NARASIMHAN, Rangachari; 20661 Oak
Creek Lane, Saratoga, California 95070 (US). BE-
LARDINELLI, Luiz; 205 Manzanita Avenue, Palo Alto,
California 94306 (US). SCHULER, Carlos A.; 21866
Wilson Court, Cupertino, California 95014 (US).(74) Agents: MATA, David et al.; Wilson Sonsini Goodrich &
Rosati, 650 Page Mill Road, Palo Alto, California 94304
(US).(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY,
BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM,
DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT,
HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KH, KN,
KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA,
MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG,
NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS,
RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY,
TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN,
ZA, ZM, ZW.(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ,
TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU,
TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE,
DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU,
LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK,

[Continued on next page]


(54) Title: COMBINING ELECTRONIC MONITORING WITH INHALED PHARMACOLOGICAL THERAPY TO MANAGE
CARDIAC ARRHYTHMIAS INCLUDING ATRIAL FIBRILLATION

FIG. 1

(57) Abstract: Disclosed herein are methods of treating cardiac arrhythmia with electronic monitoring in a timely manner. Also disclosed herein are systems for electronic monitoring of cardiac arrhythmia.



SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG). **Published:**

— *with international search report (Art. 21(3))*

Declarations under Rule 4.17:

— *as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))*

**COMBINING ELECTRONIC MONITORING WITH INHALED
PHARMACOLOGICAL THERAPY TO MANAGE CARDIAC ARRHYTHMIAS
INCLUDING ATRIAL FIBRILLATION**

CROSS-REFERENCE

[0001] This application claims priority to US provisional application number 62/289,473, filed on February 1, 2016, which is herein incorporated by reference in its entirety.

BACKGROUND

[0002] The disclosure relates to the use of inhaled CV therapies in combination with electronic monitoring devices.

[0003] Cardiac arrhythmia (also dysrhythmia) is a term for any of a large and heterogeneous group of conditions in which there can be abnormal electrical activity in the heart. The heart beat may be too fast or too slow, and may be regular or irregular.

[0004] Cardiac Arrhythmia is a field with a high level of unmet clinical need. Many drugs used today have been on the market since the early 1980s and 1990s and are mostly inadequate due to either lack of efficacy or a side-effect profile that can be primarily cardiac related, that necessitates extensive monitoring of the subject.

[0005] Atrial Fibrillation is a type of Cardiac arrhythmia that occurs as both symptomatic and asymptomatic. When symptomatic the subject goes to the ER in the hospital for treatment while when asymptomatic that subject does not realize that they are in an arrhythmic state. Atrial fibrillation normally occurs asymptotically before it becomes symptomatic in short durations. These short episodes spontaneously convert into normal sinus rhythm. But the cumulative effect of these short episodes add to the overall AF burden and eventual progression of the disease.

[0006] Paroxysmal atrial fibrillation (PAF) is a subset of the overall Atrial Fibrillation (AF) population and is estimated to be a third of the overall AF population. AF affects about 34 million people worldwide and 11.3 million of these people worldwide are diagnosed as subjects with PAF.

[0007] Arrhythmias frequently result in emergency room (ER) visits, where intravenous drugs are administered, and sometimes necessitating extended stay in hospital and in some cases also leading to unplanned invasive procedures.

[0008] Ablation can be expensive and can be about 50% efficacious. Despite the high expense, ablation may not completely correct the arrhythmia. Often, multiple ablation procedures are required to achieve a satisfactory result in a subject's lifetime. Ablation also results in heart related medical complications associated with morbidity and mortality.

[0009] Electronic Monitoring systems are used in subjects to monitor if they are experiencing such arrhythmias and when an arrhythmia is sustained they are recorded and the treating physician is able to assess the course of therapy.

[0010] What is needed for slowing the progression of AF from its starting paroxysmal state to permanent AF.

[0011] None of the current oral or IV approved drug products are combined with an electronic monitoring system with the goal to terminate the arrhythmia at initiation thereby reducing the progression of the disease, the overall AF burden leading to significant reduction in the economies of hospital costs.

[0012] Exemplary monitoring devices include the AliveCor Mobile ECG, the Reveal LINQ Insertable Cardiac Monitor, the iRhythm ZIO XT Patch, the HeartCheck pen, the AfibAlert, smartphone or portable music player applications designed to detect arrhythmias, the Microlife AFIB Technology device, the WatchBP device, Holter monitors, Smart watches, wearable technology such as those that can vibrate as a means to detect an arrhythmia, or any other similar devices.

SUMMARY

[0013] Disclosed herein are methods of treating cardiac arrhythmia in a subject, comprising: (a) identifying cardiac arrhythmia in the subject with the aid of an electronic monitoring device; (b) aerosolizing a pharmaceutical composition in less than about 9 minutes using an inhaler, wherein the pharmaceutical composition can comprise of a therapeutically effective amount of at least one antiarrhythmic or a pharmaceutically

acceptable salt thereof; and (c) administering the aerosolized pharmaceutical composition to the subject; thereby treating the cardiac arrhythmia. In some embodiments, the identifying of (a) includes an establishing of a duration of the cardiac arrhythmia. In some embodiments, the duration of the cardiac arrhythmia can be at least about 2 hours. In some embodiments, the duration of the cardiac arrhythmia can be at least about 1 hour. In some embodiments, the duration of the cardiac arrhythmia can be at least about 0.5 hours. In some embodiments, the aerosolizing of the pharmaceutical composition occurs in less than 6 minutes. In some embodiments, the aerosolizing of the pharmaceutical composition occurs in less than 3 minutes. In some embodiments, the therapeutically effective amount can be an amount sufficient to convert the arrhythmia to normal sinus rhythm. In some embodiments, the therapeutically effective amount can be at least about 60 mg of the at least one antiarrhythmic or a pharmaceutically acceptable salt thereof. In some embodiments, the therapeutically effective amount can be at least about 50 mg of the at least one antiarrhythmic or a pharmaceutically acceptable salt thereof. In some embodiments, the therapeutically effective amount can be at least about 40 mg of the at least one antiarrhythmic or a pharmaceutically acceptable salt thereof. In some embodiments, the therapeutically effective amount can be at least about 30 mg of the at least one antiarrhythmic or a pharmaceutically acceptable salt thereof. In some embodiments, the therapeutically effective amount converts the arrhythmia to normal sinus rhythm from about 0 secs to about 2 hours after the administration. In some embodiments, the therapeutically effective amount converts the arrhythmia to normal sinus rhythm in about 60 mins after administration. In some embodiments, the treating occurs within about 120 minutes from an onset of the cardiac arrhythmia. In some embodiments, the treating occurs within about 90 minutes from an onset of the cardiac arrhythmia. In some embodiments, the treating occurs within about 60 minutes from an onset of the cardiac arrhythmia. In some embodiments, the administering can comprise inhalation. In some embodiments, prior to (a), an electronic monitoring chip can be inserted into the subject. In some embodiments, an electronic monitoring chip can be worn by the subject. In some embodiments, the electronic monitoring chip can be in communication with the electronic monitoring device. In

some embodiments, the communication can be a wireless communication. In some embodiments, prior to (a), an electronic monitoring chip can be inserted into the electronic monitoring device. In some embodiments, the identifying of (a) can comprise a communication of a result to the subject. In some embodiments, the communication can comprise an instruction to administer the pharmaceutical composition. In some embodiments, the instruction to administer can comprise a dosage of the aerosolized pharmaceutical composition to administer. In some embodiments, the administering can be performed by the subject. In some embodiments, the administering can be performed by a healthcare professional. In some embodiments, the healthcare professional can override a recommendation made by the electronic monitoring device. In some embodiments, the electronic monitoring device waits for a confirmation from the healthcare provider prior to allowing the administering of the aerosolized pharmaceutical composition to the subject. In some embodiments, the communication can comprise making a treatment recommendation. In some embodiments, the treatment recommendation can comprise restricting the administration of the pharmaceutical composition when the electronic monitoring device does not identify an arrhythmia. In some embodiments, the identifying of (a) can comprise a communication of a result to a healthcare professional. In some embodiments, the administering can be performed by the subject. In some embodiments, the administering can be performed by a healthcare professional. In some embodiments, the healthcare professional can be a paramedic. In some embodiments, the healthcare professional can be a physician. In some embodiments, the healthcare professional can be a monitoring professional. In some embodiments, the healthcare professional can override a recommendation made by the electronic monitoring device. In some embodiments, the electronic monitoring device waits for a confirmation from the healthcare provider prior to allowing the administering of the aerosolized pharmaceutical composition to the subject. In some embodiments, the communication can comprise making a treatment recommendation. In some embodiments, the treatment recommendation can comprise restricting the administration of the pharmaceutical composition when the electronic monitoring device does not identify an arrhythmia. In some embodiments, the at least one antiarrhythmic or salt

thereof can be a class I antiarrhythmic. In some embodiments, the class I antiarrhythmic can be a class Ia, Ib, or Ic antiarrhythmic. In some embodiments, the class I antiarrhythmic can be flecainide or a salt thereof. In some embodiments, the at least one antiarrhythmic or salt thereof can be a class II antiarrhythmic. In some embodiments, the at least one antiarrhythmic or salt thereof can be a class III antiarrhythmic. In some embodiments, the at least one antiarrhythmic or salt thereof can be a class IV antiarrhythmic. In some embodiments, the at least one antiarrhythmic or salt thereof can be a class V antiarrhythmic. In some embodiments, the at least one antiarrhythmic or salt thereof can comprise a sodium channel blocker. In some embodiments, the at least one antiarrhythmic or salt thereof can comprise a potassium channel blocker. In some embodiments, the at least one antiarrhythmic or salt thereof can comprise a calcium channel blocker. In some embodiments, the at least one antiarrhythmic or salt thereof can be rapidly absorbed through a subject's lungs. In some embodiments, the at least one antiarrhythmic or salt thereof can be delivered to a subject's heart after being absorbed through the subject's lungs. In some embodiments, the inhaler can be in communication with the electronic monitoring device. In some embodiments, the communication can be a wired communication. In some embodiments, the communication can be a wireless communication. In some embodiments, the administering of the aerosolized pharmacological composition by the inhaler can be communicated to the electronic monitoring device. In some embodiments, the electronic monitoring device verifies the administering of the aerosolized pharmaceutical composition. In some embodiments, the electronic monitoring device continues monitoring the progress of the arrhythmia after the administration. In some embodiments, a recommendation to stop the administering can be communicated by the electronic monitoring device when the arrhythmia can be converted to a normal sinus rhythm. In some embodiments, a recommendation to seek medical attention can be communicated by the electronic monitoring device when the arrhythmia can be not responding to the administration of the pharmaceutical composition as determined by a continued incidence of arrhythmia at least about 2 hours after the administering. In some embodiments, the electronic monitoring device can comprise a digital display. In some embodiments, the electronic monitoring device can

comprise auditory means capable of communicating to the subject. In some embodiments, the electronic monitoring device can be a smartphone. In some embodiments, the inhaler can comprise a digital display. In some embodiments, the inhaler can comprise auditory means capable of communicating to the subject.

[0014] Also disclosed herein are systems comprising: (a) an electronic monitoring chip; wherein the electronic monitoring chip monitors incidence of arrhythmia in a subject; (b) an electronic monitoring device; wherein the electronic monitoring device can be in wireless communication with the electronic monitoring chip; and (c) an inhaler; wherein the electronic monitoring device can be in communication with the inhaler. In some embodiments, the electronic monitoring device can be in communication with the inhaler through a wired connection. In some embodiments, the electronic monitoring device can be in communication with the inhaler through a wireless connection. In some embodiments, the electronic monitoring device can comprise a digital display. In some embodiments, the electronic monitoring device can comprise auditory means capable of communicating to the subject. In some embodiments, the electronic monitoring device can be a smartphone. In some embodiments, the electronic monitoring chip can be inserted into the subject. In some embodiments, the electronic monitoring chip can be worn by the subject. In some embodiments, the inhaler can comprise a digital display. In some embodiments, the inhaler can comprise auditory means capable of communicating to the subject. In some embodiments, the electronic monitoring device can comprise storage means. In some embodiments, subject data can be stored using the storage means. In some embodiments, the storage means can be a hard drive. In some embodiments, the storage means can be a cloud based storage.

INCORPORATION BY REFERENCE

[0015] All publications, patents, and patent applications mentioned in this specification are herein incorporated by reference to the same extent as if each individual publication, patent, or patent application was specifically and individually indicated to be incorporated by reference.

BRIEF DESCRIPTION OF THE DRAWINGS

[0016] The novel features are set forth with particularity in the appended claims. A better understanding of features and advantages will be obtained by reference to the following detailed description that sets forth illustrative embodiments, in which exemplary principles are utilized, and the accompanying drawings of which:

[0017] Fig. 1 depicts plots of the arterial and venous concentration of flecainide after intravenous or intra-tracheal administration.

[0018] Fig. 2 depicts an exemplary physiologically based pharmacokinetic (PBPK) model of the intravenous or intra-tracheal administration of flecainide.

[0019] Fig. 3 depicts a plot of the effect on left atrial conduction time after administration via intravenous injection or inhalation.

[0020] Fig. 4 depicts a plot of the effect of administration of flecainide on a subject's mean blood pressure after administration via intravenous injection or inhalation.

[0021] Fig. 5 depicts a plot of the effect of administration of flecainide on a subject's left ventricular pressure after administration via inhalation.

[0022] Fig. 6 depicts a plot of the effect of QTcf prolongation after administration of flecainide on a subject's left ventricular pressure after administration via inhalation.

[0023] Fig. 7 depicts an exemplary plot of sinus rhythm using a smart phone as a novel electronic monitoring device as compared to a Prucka System II.

[0024] Fig. 8 depicts an exemplary plot of Atrial Fibrillation using a smart phone as a novel electronic monitoring device as compared to a Prucka System II.

[0025] Fig. 9 depicts an exemplary plot of a treatment of Atrial Fibrillation with flecainide using a smart phone as a novel electronic monitoring device as compared to a Prucka System II.

[0026] Fig. 10 depicts an exemplary smart phone application allowing the smartphone to be used as an electronic monitoring device

[0027] Fig. 11 depicts a smart phone in use as an electronic monitoring device to monitor treatment of Atrial Fibrillation with flecainide.

DETAILED DESCRIPTION

[0028] The details of one or more inventive embodiments are set forth in the accompanying drawings, the claims, and the description herein. Other features, objects, and advantages of the inventive embodiments disclosed and contemplated herein can be combined with any other embodiment unless explicitly excluded.

[0029] Accordingly, disclosed herein are compositions and methods for treating a heart arrhythmia comprising combining the treatment with an electronic monitoring system. Other features and advantages will be set forth in the description that follows, and in part will be apparent from the description or may be learned by practice of exemplary embodiments. The disclosure will be realized and attained by the compositions and methods particularly pointed out in the written description and claims thereof.

[0030] In some embodiments, the method can be directed to effective monitoring of subjects with AF through electronic means. In some cases, an electronic monitor can have an ability to record with a subject interface to an alert center. In some cases, an electronic monitor can have an ability to transmit with a subject interface to an alert center. In some cases, an electronic monitor can have an ability to record and transmit with a subject interface to an alert center. In some cases, an electronic monitor can have an ability to record without a subject interface to an alert center. In some cases, an electronic monitor can have an ability to transmit without a subject interface to an alert center. In some cases, an electronic monitor can have an ability to record and transmit without a subject interface to an alert center.

[0031] In some embodiments, the method can comprise physicians and/or qualified personnel reading the alert signals from subjects. In some cases, the method can comprise providing a subject with timely advice to self-administer CV medications to terminate an arrhythmia.

[0032] In some embodiments, the method can be directed to effectively shorten the time of arrhythmia, irrespective of whether it is symptomatic or asymptomatic, by a system where the subject monitoring compliments with the therapeutic administration of an

antiarrhythmic. In some cases, the arrhythmia can be symptomatic. In some cases, the arrhythmia can be asymptomatic

[0033] The present invention is directed to an electronic monitoring system that is either inserted or is subject enabled combined with a inhalation therapy that is self-administered either by advise from a monitoring center authorized medical personnel or by self-diagnosis of the arrhythmia.

[0034] The present invention is directed to administering by inhalation an effective amount of at least one antiarrhythmic agent to a subject in need thereof, wherein a cardiac score from a monitor implementing an arrhythmia detection algorithm is detected early in its onset and the subject's ability to self-administer the inhaled medication to enable a transition from an arrhythmic state to normal sinus rhythm in the subject within 0 seconds to 2 hours after completion of administration.

[0035] Unless otherwise indicated, some embodiments herein contemplate numerical ranges. When a numerical range is provided, unless otherwise indicated, the range includes the range endpoints. Unless otherwise indicated, numerical ranges include all values and sub ranges therein as if explicitly written out.

[0036] The singular forms "a", "an", and "the" can be used herein to include plural references unless the context clearly dictates otherwise. Accordingly, unless the contrary is indicated, the numerical parameters set forth in this application can be approximations that may vary depending upon the desired properties sought to be obtained by the present invention.

[0037] As used herein, the term "salt" is intended to include, but not be limited to, pharmaceutically acceptable salts. And the term "pharmaceutically acceptable salt" is intended to mean those salts that retain one or more of the biological activities and properties of the free acids and bases and that are not biologically or otherwise undesirable. Illustrative examples of pharmaceutically acceptable salts include, but are not limited to, sulfates, pyrosulfates, bisulfates, sulfites, bisulfites, phosphates, monohydrogenphosphates, dihydrogenphosphates, metaphosphates, pyrophosphates, chlorides, bromides, iodides, acetates, propionates, decanoates, caprylates, acrylates, formates, isobutyrate, caproates, heptanoates, propiolates, oxalates, malonates,

succinates, suberates, sebacates, fumarates, maleates, butyne-1,4-dioates, hexyne-1,6-dioates, benzoates, chlorobenzoates, methylbenzoates, dinitrobenzoates, hydroxybenzoates, methoxybenzoates, phthalates, sulfonates, xylenesulfonates, phenylacetates, phenylpropionates, phenylbutyrates, citrates, lactates, γ -hydroxybutyrates, glycolates, tartrates, methanesulfonates, propanesulfonates, naphthalene-1-sulfonates, naphthalene-2-sulfonates, and mandelates. The term "pharmaceutically acceptable" is intended to include, but not be limited to, mean that a component is not biologically or otherwise undesirable, e.g., the component may be incorporated into a pharmaceutical composition of the invention and administered to a subject as described herein without causing any significant undesirable biological effects or interacting in a deleterious manner with any of the other components of the composition in which it is contained. When the term "pharmaceutically acceptable" is used to refer to an excipient, it is generally implied that the component has met the required standards of toxicological and manufacturing testing or that it is included on the Inactive Ingredient Guide prepared by the U.S. Food and Drug Administration.

[0038] Unless otherwise indicated, "aerosolizing a pharmaceutical composition" means at least about 90% by weight of the pharmaceutical composition is aerosolized. For example, "aerosolizing a pharmaceutical composition in less than 3 minutes" means at least 90% by weight of the pharmaceutical composition is aerosolized in less than 3 minutes.

[0039] Unless otherwise indicated, open terms for example "contain," "containing," "include," "including," and the like mean comprising.

[0040] Reference herein to "one embodiment," "one version," or "one aspect" shall include one or more such embodiments, versions or aspects, unless otherwise clear from the context.

[0041] As used herein, "tachycardia" means an arrhythmia in which the heart beat is too fast.

[0042] As used herein, the phrase "heart rhythm arrhythmia" means an arrhythmia in which the heart beat is irregular.

[0043] As used herein, the “amount of the at least one antiarrhythmic agent in blood in the coronary sinus of the heart” may be measured by extracting a sample from the coronary sinus of the heart by using a cannula. The amount of antiarrhythmic agent in the sample may then be determined by known means, such as bioanalytical techniques that employ analytical equipment such as LC-MS/MS. Thus, the amount of antiarrhythmic agent in the blood in the heart may be determined for any particular time.

[0044] As used herein, the terms "treating" and "treatment" refer to reduction in severity and/or frequency of symptoms, elimination of symptoms and/or underlying cause, reduction in likelihood of the occurrence of symptoms and/or underlying cause, and/or remediation of damage. Thus, "treating" a subject with an active agent as provided herein includes prevention of a particular condition, disease, or disorder in a susceptible individual as well as treatment of a clinically symptomatic individual.

[0045] As used herein, “nominal amount” refers to the amount contained within the unit dose receptacle(s) that can be administered.

[0046] As used herein, "effective amount" refers to an amount covering both therapeutically effective amounts and prophylactically effective amounts.

[0047] As used herein, a "therapeutically effective amount" of an active agent refers to an amount that is effective to achieve a desired therapeutic result. A therapeutically effective amount of a given active agent will typically vary with respect to factors such as the type and severity of the disorder or disease being treated and the age, gender, and weight of the subject.

[0048] Unless otherwise specified, the term "therapeutically effective amount" includes a "prophylactically effective amount," i.e., an amount of active agent that is effective to prevent the onset or recurrence of particular condition, disease, or disorder in a susceptible individual.

Pharmaceutical composition

[0049] In some embodiments, a composition can comprise an antiarrhythmic agent. Examples of antiarrhythmic agents include, but are not limited to, class Ia (sodium channel blockers, intermediate association/dissociation), class Ib (sodium channel

blockers, fast association/dissociation), class Ic (sodium channel blocker, slow association/dissociation), class II (beta blockers), class III (potassium channel blockers), class IV (calcium channel blockers), and class V (unknown mechanisms) antiarrhythmics.

[0050] Class Ia antiarrhythmics include, but are not limited to, quinidine, procainamide, and disopyramide. Class Ib antiarrhythmics include, but are not limited to, lidocaine, tocamide, phenytoin, moricizine, and mexiletine. Class Ic antiarrhythmics include, but are not limited to, flecainide, ajmaline, propafenone, and moricizine. Class II antiarrhythmics include, but are not limited to, propranolol, acebutolol, sotalol, esmolol, timolol, metoprolol, and atenolol. Class III antiarrhythmics include, but are not limited to, amiodarone, sotalol, bretylium, ibutilide, E-4031 (methanesulfonamide), vemakalant, and dofetilide. Class IV antiarrhythmics include, but are not limited to, bepridil, nitrendipine, amlodipine, isradipine, nifedipine, nicardipine, verapamil, and diltiazem. Class V antiarrhythmics include, but are not limited to, digoxin and adenosine.

[0051] The pharmaceutical composition can also include derivatives of the above antiarrhythmic pharmaceutical agents such as solvates, salts (e.g., acetate), solvated salts, esters, amides, hydrazides, N-alkyls, and/or N-amino acyls. In some cases, the pharmaceutical composition is a liquid pharmaceutical composition. Examples of ester derivatives include, but are not limited to, methyl esters, choline esters, and dimethylaminopropyl esters. Examples of amide derivatives include, but are not limited to, primary, secondary, and tertiary amides. Examples of hydrazide derivatives include, but are not limited to, N-methylpiperazine hydrazides. Examples of N-alkyl derivatives include, but are not limited to, N',N',N'-trimethyl and N',N'-dimethylaminopropyl succinimidyl derivatives of antiarrhythmic pharmaceutical agent methyl esters. Examples of N-aminoacyl derivatives include, but are not limited to, N-omithyl-, N-diaminopropionyl-, N-lysyl-, N-hexamethyllysyl-, and N-piperidine-propionyl- or N',N'-methyl-1-piperazine-propionyl-antiarrhythmic pharmaceutical agent methyl esters.

[0052] The pharmaceutical composition can comprise one or more antiarrhythmic pharmaceutical agents and, optionally, one or more pharmaceutically acceptable excipients. The pharmaceutically acceptable excipient can comprise lipids, metal ions,

surfactants, amino acids, carbohydrates, buffers, salts, polymers, and the like, and combinations thereof. In some cases, the pharmaceutically acceptable excipient is water. In some cases, the pharmaceutically acceptable excipient is not water. Examples of lipids include, but are not limited to, phospholipids, glycolipids, ganglioside GMI, sphingomyelin, phosphatidic acid, cardiolipin; lipids bearing polymer chains such as polyethylene glycol, chitin, hyaluronic acid, or polyvinylpyrrolidone; lipids bearing sulfonated mono-, di-, and polysaccharides; fatty acids such as palmitic acid, stearic acid, and oleic acid; cholesterol, cholesterol esters, and cholesterol hemisuccinate. The pharmaceutically acceptable excipient can comprise one or more osmolality adjuster, such as sodium chloride. For instance, sodium chloride may be added to solutions to adjust the osmolality of the solution. In one or more embodiments, an aqueous composition consists essentially of the antiarrhythmic pharmaceutical agent, the osmolality adjuster, and water. The pharmaceutically acceptable can also comprise a buffer or a pH adjusting agent, typically a salt prepared from an organic acid or base. For example, the pharmaceutical compositions typically have a pH ranging from 3.5 to 8.0, such as from 4.0 to 7.5, 4.5 to 7.0, or 5.0 to 6.5. Representative buffers comprise organic acid salts of citric acid, lactic acid, ascorbic acid, gluconic acid, carbonic acid, tartaric acid, succinic acid, acetic acid, or phthalic acid, tris, trimethamine hydrochloride, or phosphate buffers. Thus, the buffers include citrates, phosphates, phthalates, and lactates. It may be desirable to add other pharmaceutically acceptable excipients to the pharmaceutical composition to improve particle rigidity, production yield, emitted dose and deposition, shelf-life, and subject acceptance. Such optional pharmaceutically acceptable excipients include, but are not limited to: coloring agents, taste masking agents, buffers, hygroscopic agents, antioxidants, and chemical stabilizers.

[0053] As noted above, the pharmaceutical composition may include one or more surfactants. For instance, one or more surfactants may be in the liquid phase with one or more being associated with (e.g., the pharmaceutical compositions may incorporate, adsorb, absorb, be coated with, or be formed by the surfactant) solid particles or particles of the composition. By "associated with" it is meant that the pharmaceutical compositions may incorporate, adsorb, absorb, be coated with, or be formed by the

surfactant. Surfactants include, but are not limited to, fluorinated and nonfluorinated compounds, such as saturated and unsaturated lipids, nonionic detergents, nonionic block copolymers, ionic surfactants, and combinations thereof. Examples of ionic surfactants include, but are not limited to, sodium sulfosuccinate, and fatty acid soaps. Examples of buffers include, but are not limited to, tris or citrate. Examples of acids include, but are not limited to, carboxylic acids. Examples of carbohydrates include, but are not limited to, monosaccharides, disaccharides, and polysaccharides. For example, monosaccharides such as dextrose (e.g. anhydrous and monohydrate), galactose, mannitol, D-mannose, sorbitol, sorbose and the like; disaccharides such as lactose, maltose, sucrose, trehalose, and the like; trisaccharides such as raffinose and the like; and other carbohydrates such as starches (e.g., hydroxyethylstarch), cyclodextrins, and maltodextrins.

[0054] The pharmaceutical composition can comprise one or more antiarrhythmic pharmaceutical agents and, can include one or more additional active agents, for example, agents that may be delivered through the lungs. Additional active agents may comprise, for example, hypnotics and sedatives, psychic energizers, tranquilizers, respiratory drugs, anticonvulsants, muscle relaxants, anti-Parkinson agents (e.g. dopamine antagonists), analgesics, anti-inflammatories, antianxiety drugs (e.g. anxiolytics), appetite suppressants, antimigraine agents, muscle contractants, additional anti-infectives (e.g. antivirals, antifungals, vaccines) antiarthritics, antimalarials, antiemetics, anepileptics, cytokines, growth factors, anti-cancer agents, antithrombotic agents, antihypertensives, cardiovascular drugs, antiarrhythmics, antioxidants, anti-asthma agents, hormonal agents including contraceptives, sympathomimetics, diuretics, lipid regulating agents, antiandrogenic agents, antiparasitics, anticoagulants, neoplastics, antineoplastics, hypoglycemics, nutritional agents and supplements, growth supplements, antienteritis agents, vaccines, antibodies, diagnostic agents, and contrasting agents. The additional active agent, when administered by inhalation (e.g., as an aerosol), may act locally or systemically.

[0055] The amount of antiarrhythmic pharmaceutical agent in the pharmaceutical composition can vary. The amount of antiarrhythmic pharmaceutical agent(s) can be

about 0.1% to 100% by weight of the total amount of the pharmaceutical composition. In some cases, the amount of antiarrhythmic pharmaceutical agent(s) is at least about 0.1% by weight of the total amount of the pharmaceutical composition, for example, at least about 0.1%, 0.2%, 0.3%, 0.4%, 0.5%, 1%, 2%, 3%, 4%, 5%, 6%, 7%, 8%, 9%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, or 100% by weight of the total amount of the pharmaceutical composition. In some cases, the amount of antiarrhythmic pharmaceutical agent(s) is about 0.1%-100% by weight of the total amount of the pharmaceutical composition, for example, about 0.1%-1%, 0.1%-5%, 0.1-10%, 0.1%-20%, 0.5%-1%, 0.5%-5%, 0.5%-10%, 0.5%-20%, 1%-5%, 1%-10%, 1%-20%, 5%-10%, 5%-20%, 10%-20%, 10%-30%, 20%-30%, 20%-40%, 30%-40%, 30%-50%, 40%-50%, 40%-60%, 50%-60%, 50%-70%, 60%-70%, 60%-80%, 70%-80%, 70%-90%, 80%-90%, 80%-95%, 90%-95%, 90%-99%, 90%-100%, 95%-99%, or 99%-100% by weight of the total amount of the pharmaceutical composition.

Method of treatment

[0056] Described herein are methods of treating a condition in a subject pharmaceutical comprising delivering a pharmaceutical composition disclosed herein. The condition can be cardiac arrhythmia including, but not limited to, tachycardia, supraventricular, tachycardia (SVT), paroxysmal supraventricular tachycardia (PSVT), atrial fibrillation (AF), paroxysmal atrial fibrillation (PAF), permanent atrial fibrillation, persistent atrial fibrillation, atrial flutter, paroxysmal atrial flutter, and lone atrial fibrillation.

[0057] One aspect of embodiments disclosed herein is the surprising and unexpected rate at which the antiarrhythmic agents pass through the heart. While a skilled artisan might expect the rate to be too fast, modeling and empirical evidence indicates that the drug will not pass through the heart too fast. Thus, a therapeutic effect is achieved despite fast pass-through and despite only one pass-through at therapeutic levels

[0058] Also described are methods to treat acute episodes of and/or chronic arrhythmias. In certain embodiments, the treating can comprise acute treatment after detection of cardiac arrhythmia.

[0059] Flecainide can be effective via IV at doses of 75-150 mg administered in a hospital or other suitable medical facility. The equilibration between arterial concentration and venous concentration can occur rapidly. On the other hand, inhalation can target the drug to the pulmonary vein that originates in the lung capillaries, and the left atrium where PAF originates. Inhalation can be started much earlier at onset rather than the subject waiting to go to the hospital for an IV. Alternatively, Flecainide can be also administered orally as a bolus of 300 mg (max per day) dose at one time to terminate PAF. Via this route, it can take an average of 100 minutes to terminate the PAF. Oral flecainide is 30-50% effective. Subjects who take this maximum dose are restricted until 24 hours to take any further antiarrhythmics. Conversely, an inhalation dose can be several orders of magnitude less than the oral dose, and thus permit repeat doses of flecainide.

[0060] The method of treatment via inhalation can result in a pulsatile pharmacokinetic profile and transient pharmacodynamic effect mimicking the effect of an IV. The method can deliver high drug concentrations that are safe and effective to the heart, while the distribution to the rest of the body results in the drug being diluted to sub-therapeutic levels. This method is the shortest route of delivery to the heart next to intracardial injection. This provides the convenience of self-administration like the “pill-in-the-pocket” approach, but the effectiveness and fast onset of action of an IV. Although the delivery of medications through the lung for systemic effect is not new, it was widely thought it wouldn’t be effective to the heart, because of the fast passage of the drug through it, not having time to act on the target tissue. The PK/PD modeling and empirical evidence originating this invention shows that the drug exposure is sufficient for therapeutic effect at a much lower dose compared to other routes of administration. This method ensures drug concentrations in overall plasma are much lower than what is achieved by oral/IV hence minimizing drug-drug interactions and side effects.

[0061] The subject can be a mammal in need thereof, preferably such mammal is a human subject. Examples of subjects include, but are not limited to, pediatric subjects, adult subjects, and geriatric subjects. In some embodiments, the pharmaceutical composition is intended only as a treatment for rapid resolution of symptoms and is not

taken as a preventative, i.e., when the subject is well, there is no need for drug-this makes the therapy more effective and safe due to sporadic or intermittent dosing, and focused on reducing disabling symptoms.

[0062] Unit doses of the pharmaceutical compositions may be contained in a container. Examples of containers include, but are not limited to, syringes, capsules, blow fill seal, blisters, vials, ampoules, bottles, or container closure systems made of metal, polymer (e.g., plastic, elastomer), glass, or the like. For instance, the vial may be a colorless Type I borosilicate glass ISO 6R 10 mL vial with a chlorobutyl rubber siliconized stopper, and rip-off type aluminum cap with colored plastic cover.

[0063] The unit dose can comprise at least about 0.1 mL of the pharmaceutical composition, for example, at least about 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9, 1, 1.5, 2, 2.5, 3, 3.5, 4, 4.5, 5, 5.5, 6, 6.5, 7, 7.5, 8, 8.5, 9, 9.5, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, or 100 mL of the pharmaceutical composition. The unit dose can comprise about 0.1-100 mL of the pharmaceutical composition, for example, about 0.1-0.2, 0.1-0.5, 0.1-1.0, 0.2-0.5, 0.2-1.0, 0.5-1.0, 0.5-1.5, 0.5-2.0, 0.5-3.0, 0.5-4.0, 0.5-5.0, 1.0-2.0, 1.0-3.0, 1.0-4.0, 1.0-5.0, 2.0-3.0, 2.0-4.0, 2.0-5.0, 3.0-5.0, 5-10, 10-20, 20-30, 30-40, 40-50, 50-60, 60-70, 70-80, 80-90, or 90-100 mL of the pharmaceutical composition. In some cases, the unit dose of the pharmaceutical composition ranges from about 2 ml to about 15 ml, such as from about 3 ml to about 10 ml, about 4 ml to about 8 ml, or about 5 ml to about 6 ml.

[0064] The container may be inserted into an aerosolization device (e.g., inhaler). The container may be of a suitable shape, size, and material to contain the pharmaceutical composition and to provide the pharmaceutical composition in a usable condition.

[0065] The compositions of the present invention may be made by any of the various methods and techniques known and available to those skilled in the art. For instance, a solution of antiarrhythmic pharmaceutical agent may be made using the following procedure. Typically, manufacturing equipment is sterilized before use. A portion of the final volume, e.g., 70%, of solvent, e.g., water for injection, may be added into a suitable container. Antiarrhythmic pharmaceutical agent may then be added. The antiarrhythmic pharmaceutical agent may be mixed until dissolved. Additional solvent

may be added to make up the final batch volume. The batch may be filtered, e.g., through a 0.2 μm filter into a sterilized receiving vessel. Filling components may be sterilized before use in filling the batch into vials, e.g., 10 ml vials. As an example, the above-noted sterilizing may include the following. A 5 liter type 1 glass bottle and lid may be placed in an autoclave bag and sterilized at elevated temperature, e.g., 121 $^{\circ}\text{C}$, for 15 minutes, using an autoclave. Similarly, vials may be placed into suitable racks, inserted into an autoclave bag, and sterilized at elevated temperature, e.g., 121 $^{\circ}\text{C}$, for 15 minutes, using an autoclave. Also similarly, stoppers may be placed in an autoclave bag and sterilized at elevated temperature, e.g., 121 $^{\circ}\text{C}$, for 15 minutes, using an autoclave. Before sterilization, sterilizing filters may be attached to tubing, e.g., a 2 mm length of 7 mm x 13 mm silicone tubing. A filling line may be prepared by placed in an autoclave bag and sterilized at elevated temperature, e.g., 121 $^{\circ}\text{C}$, for 15 minutes, using an autoclave. Sterilization may also be accomplished using electronic beams or gamma rays. The filling may also be conducted under laminar flow protection. The filling line may be unwrapped and placed into the receiving bottle. The sterilized vials and stoppers may be unwrapped under laminar flow protection. Each vial may be filled, e.g., to a target fill of 5 g, and stoppered. A flip off collar may be applied to each vial. The sealed vials may be inspected for vial leakage, correct overseals, and cracks.

[0066] As another example, an antiarrhythmic may be prepared by lyophilizing the antiarrhythmic to form a powder for storage. The powder is then reconstituted prior to use. This technique may be used when the antiarrhythmic is unstable in solution. The solvent for the solution to be lyophilized may comprise water. The solution may be excipient-free. For instance, the solution may be cryoprotectant-free. In one or more embodiments, a suitable amount (e.g., 30 mg per mL of final solution) of drug substance may be dissolved, e.g., in about the 75% of the theoretical total amount of water for injection under nitrogen bubbling. The dissolution time may be recorded and appearance may be evaluated. Then, the dilution to the final volume with water for injection (WFI) may be carried out. Final volume may be checked. Density, pH, endotoxin, bioburden, and content by UV may be measured both before and after sterile filtration.

[0067] The solution may be filtered before lyophilizing. For instance, a double 0.2 μm filtration may be performed before filling. The filters may be tested for integrity and bubble point before and after the filtration. Pre-washed and autoclaved vials may be aseptically filled using an automatic filling line to a target of 5 ml per vial and then partially stoppered. In process check for fill volumes may be done by checking the fill weight every 15 minutes. The lyophilizing is generally conducted within about 72 hours, such as within about 8 hours, or within about 4 hours, of the dissolving. In one or more embodiments, the lyophilizing can comprise freezing the solution to form a frozen solution. The frozen solution is typically held at an initial temperature ranging from about $-40\text{ }^{\circ}\text{C}$ to about $-50\text{ }^{\circ}\text{C}$, such as about $-45\text{ }^{\circ}\text{C}$. During the initial temperature period, the pressure around the frozen solution is typically atmospheric pressure. The initial temperature period typically ranges from about 1 hour to about 4 hours, such as about 1.5 hours to about 3 hours, or about 2 hours. The lyophilizing may further comprise raising a temperature of the frozen solution to a first predetermined temperature, which may range from about $10\text{ }^{\circ}\text{C}$ to about $20\text{ }^{\circ}\text{C}$, such as about $15\text{ }^{\circ}\text{C}$. The time for the heat ramp from the initial temperature to the first predetermined temperature generally ranges from about 6 hours to about 10 hours, such as about 7 hours to about 9 hours. During the first predetermined temperature period, the pressure around the solution typically ranges from about 100 μbar to about 250 μbar , such as about 150 μbar to about 225 μbar . The solution may be held at the first predetermined temperature for a period ranging from about 20 hours to about 30 hours, such as from about 24 hours. The lyophilizing may still further comprise raising a temperature of the solution to a second predetermined temperature, which may range from about $25\text{ }^{\circ}\text{C}$ to about $35\text{ }^{\circ}\text{C}$, such as about $30\text{ }^{\circ}\text{C}$. During the second predetermined temperature period, the pressure around the frozen solution typically ranges from about 100 μbar to about 250 μbar , such as about 150 μbar to about 225 μbar . The solution may be held at the second predetermined temperature for a period ranging from about 10 hours to about 20 hours.

[0068] The lyophilization cycle may comprise a freezing ramp, e.g., from $20\text{ }^{\circ}\text{C}$ to $-45\text{ }^{\circ}\text{C}$ in 65 minutes, followed by a freeze soak, e.g., at $-45\text{ }^{\circ}\text{C}$ for 2 hours. Primary drying

may be accomplished with a heating ramp, e.g., from -45 °C to 15 °C in 8 hours, followed by a temperature hold, e.g., at 15 °C for 24 hours at a pressure of 200 μ bar. Secondary drying may be accomplished with a heating ramp, e.g., from 15 °C to 30 °C in 15 minutes, followed by a temperature hold at 30 °C for 15 hours at a pressure of 200 μ bar. At the end of the lyophilization cycle, the vacuum may be broken with sterile nitrogen, and the vials may be automatically stoppered. The water content of the lyophilized powder is typically less than about 7 wt %, such as less than about 5 wt %, less than about 4 wt %, less than about 3 wt %, less than about 2 wt %, or less than about 1 wt %. The powder is capable of being reconstituted with water at 25 °C and 1.0 atmosphere and with manual agitation, in less than about 60 seconds, such as less than about 30 seconds, less than about 15 seconds, or less than about 10 seconds. The powder typically has a large specific surface area that facilitates reconstitution. The specific surface area typically ranges from about 5 m²/g to 20 m²/g, such as about 8 m²/g to 15 m²/g, or about 10 m²/g to 12 m²/g. Upon reconstitution with water, the antiarrhythmic pharmaceutical agent solution typically has a pH that ranges from about 2.5 to about 7, such as about 3 to about 6.

[0069] The compositions of one or more embodiments of the present invention may be administered by inhalation. Moreover, the doses of composition that are inhaled can be much less than those administered by other routes and required to obtain similar effects. It may be due to the efficient targeting of the inhaled composition to the heart.

[0070] The dosage necessary and the frequency of dosing of the antiarrhythmic pharmaceutical agent depend on the composition and concentration of the antiarrhythmic pharmaceutical agent within the composition. In some cases, the dose is less than its normal intravenous dose. The pulmonary dose can be similar to intracardial doses.

[0071] Inhalation avoids the initial dilution of drug in the body as compared to intravenous or oral dosing. Inhalation also avoids first-pass metabolism, such as hepatic metabolism. For instance, calcium channel blockers, such as diltiazem, undergo significant hepatic metabolism when taken orally. Inhalation allows rapid delivery of the parent diltiazem compound to the heart as a bolus. Surprisingly, administration by

inhalation of diltiazem via the inhalation route according to the present invention converted atrial fibrillation to normal sinus rhythm and reduced heart rate. Thus, administration by inhalation of diltiazem is useful for treating both atrial fibrillation and supraventricular tachycardia (SVT). In contrast, administration by IV of diltiazem is typically only used for converting SVT to normal sinus rhythm and in atrial fibrillation to reduce heart rate (not for converting to normal sinus rhythm). Inhalation also avoids red blood cell metabolism. For instance, the reduced dilution and short route associated with inhalation reduces red blood cell metabolism of esmolol. Inhalation may also avoid reduced blood pressure and fainting. For instance, IV administration of beta blockers, such as esmolol, and/or Class I antiarrhythmics such as flecainide, may reduce mean arterial blood pressure (MAP). Inhalation can allow rapid delivery of esmolol and/or flecainide without reducing MAP. As a result, inhalation of beta blockers and/or Class I antiarrhythmics may result in an MAP of 10 mm Hg to 20 mm Hg greater than the MAP resulting from IV administration of the same beta blocker or Class I antiarrhythmic. With inhaled cardiotherapy the drug is directed to the heart from the lungs as a bolus. So, the heart sees a high concentration. The drug is rapidly diluted as it passes through the heart, but the exposure time is sufficient for the desired pharmacological action. Once the drug passes through the heart, the concentration of the drug in the overall blood is below the therapeutic concentration and is considered ineffective and thus safer. The therapeutic window is the range of dosage of a drug or of its concentration in a bodily system that provides safe effective therapy. Anything below the minimum amount is sub-therapeutic and hence ineffective in that concentration. In view of the dilution, unwanted side effects are minimized.

[0072] The dose may be administered during a single inhalation or may be administered during several inhalations. The fluctuations of antiarrhythmic pharmaceutical agent concentration can be reduced by administering the pharmaceutical composition more often or may be increased by administering the pharmaceutical composition less often. Therefore, the pharmaceutical composition of one or more embodiments of the present invention may be administered from about four times daily to about once a month, such as about once daily to about once every two weeks, about once every two days to about

once a week, and about once per week. The pharmaceutical composition can also be administered to the subject on an as-needed basis.

[0073] In some cases, the pharmaceutical composition in accordance with one or more embodiments of the invention may be administered from about 1 to about 4 times daily, such as from about 2 to about 3 times daily. In some cases, the antiarrhythmic may be administered daily. In some cases, the administration can be at least about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, or 62 times a month.

[0074] In some cases, an antiarrhythmic pharmaceutical agent or salt thereof can be administered at a dose of from about 1 mg to about 480 mg, from about 5 mg to about 480 mg, from about 10 mg to about 480 mg, from about 15 mg to about 480 mg, from about 20 mg to about 480 mg, from about 25 mg to about 480 mg, from about 30 mg to about 480 mg, from about 35 mg to about 480 mg, from about 40 mg to about 480 mg, from about 45 mg to about 480 mg, from about 50 mg to about 480 mg, from about 55 mg to about 480 mg, from about 60 mg to about 480 mg, from about 65 mg to about 480 mg, from about 70 mg to about 480 mg, from about 75 mg to about 480 mg, from about 80 mg to about 480 mg, from about 85 mg to about 480 mg, from about 90 mg to about 480 mg, from about 95 mg to about 480 mg, from about 100 mg to about 480 mg, from about 105 mg to about 480 mg, from about 110 mg to about 480 mg, from about 115 mg to about 480 mg, from about 120 mg to about 480 mg, from about 125 mg to about 480 mg, from about 130 mg to about 480 mg, from about 135 mg to about 480 mg, from about 140 mg to about 480 mg, from about 145 mg to about 480 mg, from about 150 mg to about 480 mg, from about 155 mg to about 480 mg, from about 160 mg to about 480 mg, from about 165 mg to about 480 mg, from about 170 mg to about 480 mg, from about 175 mg to about 480 mg, from about 180 mg to about 480 mg, from about 185 mg to about 480 mg, from about 190 mg to about 480 mg, from about 195 mg to about 480 mg, from about 200 mg to about 480 mg, from about 205 mg to about 480 mg, from about 210 mg to about 480 mg, from about 215 mg to about 480 mg, from about 220 mg to about 480 mg, from about 225 mg to about 480 mg, from about 230 mg to about 480

mg, from about 235 mg to about 480 mg, from about 240 mg to about 480 mg, from about 245 mg to about 480 mg, from about 250 mg to about 480 mg, from about 255 mg to about 480 mg, from about 260 mg to about 480 mg, from about 265 mg to about 480 mg, from about 270 mg to about 480 mg, from about 275 mg to about 480 mg, from about 280 mg to about 480 mg, from about 285 mg to about 480 mg, from about 290 mg to about 480 mg, from about 295 mg to about 480 mg, or from about 300 mg to about 480 mg.

[0075] In some cases, an antiarrhythmic pharmaceutical agent or salt thereof described herein can be administered at a dose of about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 184, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 242, 243, 244, 245, 246, 247, 248, 249, 250, 251, 252, 253, 254, 255, 256, 257, 258, 259, 260, 261, 262, 263, 264, 265, 266, 267, 268, 269, 270, 271, 272, 273, 274, 275, 276, 277, 278, 279, 280, 281, 282, 283, 284, 285, 286, 287, 288, 289, 290, 291, 292, 293, 294, 295, 296, 297, 298, 299, 300, 301, 302, 303, 304, 305, 306, 307, 308, 309, 310, 311, 312, 313, 314, 315, 316, 317, 318, 319, 320, 321, 322, 323, 324, 325, 326, 327, 328, 329, 330, 331, 332, 333, 334, 335, 336, 337, 338, 339, 340, 341, 342, 343, 344, 345, 346, 347, 348, 349, 350, 351, 352, 353, 354, 355, 356, 357, 358, 359, 360, 361, 362, 363, 364, 365, 366, 367, 368, 369, 370, 371, 372, 373, 374, 375, 376, 377, 378, 379, 380, 381, 382, 383, 384, 385, 386, 387, 388, 389, 390, 391, 392, 393, 394, 395, 396,

397, 398, 399, 400, 401, 402, 403, 404, 405, 406, 407, 408, 409, 410, 411, 412, 413, 414, 415, 416, 417, 418, 419, 420, 421, 422, 423, 424, 425, 426, 427, 428, 429, 430, 431, 432, 433, 434, 435, 436, 437, 438, 439, 440, 441, 442, 443, 444, 445, 446, 447, 448, 449, 450, 451, 452, 453, 454, 455, 456, 457, 458, 459, 460, 461, 462, 463, 464, 465, 466, 467, 468, 469, 470, 471, 472, 473, 474, 475, 476, 477, 478, 479, or 480 mg.

[0076] In some exemplary embodiments, a daily dosage of an antiarrhythmic pharmaceutical agent (e.g. flecainide) can range from about 0.1 mg to about 600 mg, such as about 0.5 mg to about 500 mg, about 1 mg to about 400 mg, about 2 mg to about 300 mg, and about 3 mg to about 200 mg.

[0077] For treating a subject suffering from arrhythmia, the amount per dose of antiarrhythmic pharmaceutical agent (e.g. flecainide) administered may be an amount that is effective to treat the arrhythmia. The amount of antiarrhythmic pharmaceutical agent (e.g. flecainide) for the treatment of arrhythmia can be at least about 0.001 mg/kg, such as at least about 0.001 mg/kg, 0.002 mg/kg, 0.003 mg/kg, 0.004 mg/kg, 0.005 mg/kg, 0.01 mg/kg, 0.02 mg/kg, 0.04 mg/kg, 0.06 mg/kg, 0.08 mg/kg, 0.1 mg/kg, 0.2 mg/kg, 0.3 mg/kg, 0.4 mg/kg, 0.5 mg/kg, 0.6 mg/kg, 0.7 mg/kg, 0.8 mg/kg, 0.9 mg/kg, 1 mg/kg, 1.5 mg/kg, 2 mg/kg, 2.5 mg/kg, 3 mg/kg, 3.5 mg/kg, 4 mg/kg, 4.5 mg/kg, 5 mg/kg, or 6 mg/kg. The amount of antiarrhythmic pharmaceutical agent (e.g. flecainide) for the treatment of arrhythmia can range from about 0.001 mg/kg to 6 mg/kg, such as from about 0.001 mg/kg to about 0.01 mg/kg, from about 0.01 mg/kg to about 0.05 mg/kg, from about 0.05 mg/kg to about 0.1 mg/kg, from about 0.1 mg/kg to about 0.2 mg/kg, from about 0.1 mg/kg to about 0.5 mg/kg, from about 0.1 mg/kg to about 1 mg/kg, from about 0.1 mg/kg to about 2 mg/kg, from about 0.1 mg/kg to about 3 mg/kg, from about 0.3 mg/kg to about 1 mg/kg, from about 0.3 mg/kg to about 2 mg/kg, from about 0.3 mg/kg to about 3 mg/kg, from about 0.5 mg/kg to about 1 mg/kg, from about 0.5 mg/kg to about 2 mg/kg, from about 0.5 mg/kg to about 3 mg/kg, from about 0.5 mg/kg to about 6 mg/kg, from about 0.7 mg/kg to about 1 mg/kg, from about 0.7 mg/kg to about 2 mg/kg, from about 0.7 mg/kg to about 4 mg/kg, from about 0.7 mg/kg to about 6 mg/kg, from about 1 mg/kg to about 2 mg/kg, from about 1 mg/kg to about 4 mg/kg, from about 1 mg/kg to about 6 mg/kg, from about 2 mg/kg to about 3 mg/kg,

from about 2 mg/kg to about 4 mg/kg, from about 2 mg/kg to about 6 mg/kg, or from about 3 mg/kg to about 6 mg/kg.

[0078] The amount of antiarrhythmic pharmaceutical agent (e.g. flecainide) for the treatment of arrhythmia can be at least about 0.1 mg, such as at least about 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9, 1, 1.5, 2, 2.5, 3, 3.5, 4, 4.5, 5, 5.5, 6, 6.5, 7, 7.5, 8, 8.5, 9, 9.5, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 30, 35, 40, 45, 50, 60, 70, 80, 90, 100, 110, 120, 130, 140, or 150 mg. The amount of antiarrhythmic pharmaceutical agent (e.g. flecainide) for the treatment of arrhythmia can range about 0.01-150 mg, such as about 0.1-150, 0.1-130, 0.1-110, 0.1-90, 0.1-70, 0.1-50, 0.1-30, 0.1-10, 0.1-5, 0.1-1.0, 0.1-0.5, 1-150, 1-130, 1-110, 1-90, 1-70, 1-50, 1-30, 1-10, 1-5, 5-150, 5-130, 5-110, 5-90, 5-70, 5-50, 5-30, 5-10, 10-150, 10-130, 10-110, 10-90, 10-70, 10-50, 10-30, 30-150, 30-130, 30-110, 30-90, 30-70, 30-50, 50-150, 50-130, 50-110, 50-90, 50-70, 70-150, 70-130, 70-110, 70-90, 90-150, 90-130, 90-110, 110-150, 110-130, or 130-150 mg. For example, the amount of antiarrhythmic pharmaceutical agent (e.g. flecainide) for the treatment of arrhythmia can range about from 0.1 to about 5 mg.

[0079] In some instances, an antiarrhythmic can be administered as a unit dose. In some cases, a unit dose can be a dose of from about 0.1 mg to about 100 mg or greater of an antiarrhythmic pharmaceutical agent to the lung in a single inhalation. The above described phospholipid hollow and/or porous dry powder particles allow for doses of about 5 mg or greater, often greater than about 10 mg, sometimes greater than about 15 mg, sometimes greater than about 20 mg, sometimes greater than about 25 mg, sometimes greater than about 30 mg, sometimes greater than about 35 mg, sometimes greater than about 40 mg, sometimes greater than about 45 mg, sometimes greater than about 50 mg, sometimes greater than about 55 mg, sometimes greater than about 60 mg, sometimes greater than about 65 mg, sometimes greater than about 70 mg, sometimes greater than about 75 mg, sometimes greater than about 80 mg, sometimes greater than about 85 mg, sometimes greater than about 90 mg, sometimes greater than about 95 mg, or sometimes greater than about 100 mg, to be delivered in a single inhalation and in an advantageous manner. Alternatively, a dosage may be delivered over two or more inhalations, such as at least 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20,

25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95 or 100 inhalations. A dosage may also be delivered over 1 to 100 inhalations, such as 1-3, 1-4, 1-5, 1-6, 1-10, 1-20, 1-50, 1-80, 1-100, 2-5, 2-6, 2-10, 2-20, 2-50, 2-100, 5-10, 5-20, 5-50, 5-100, 10-20, 10-50, 10-100, 20-50, 20-100, or 50-100 inhalations. For example, a 10 mg dosage may be delivered by providing two unit doses of 5 mg each, and the two unit doses may be separately inhaled. In certain embodiments, the overall dose of the antiarrhythmic pharmaceutical agent ranges from 0.1 mg to 200 mg, such as 0.5 mg to 150 mg, or 1 mg to 100 mg. In some instances the antiarrhythmic agent can be administered as-needed titrating the dosage to effect.

[0080] The concentration of antiarrhythmic pharmaceutical agent (e.g. flecainide) for the treatment of arrhythmia can be at least about 0.1 mg/mL, such as at least about 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9, 1, 1.5, 2, 2.5, 3, 3.5, 4, 4.5, 5, 5.5, 6, 6.5, 7, 7.5, 8, 8.5, 9, 9.5, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 110, 120, 130, 140, or 150 mg/mL. For example, the concentration of antiarrhythmic pharmaceutical agent (e.g. flecainide) for the treatment of arrhythmia can be at least about 30 mg/mL in an acetate buffer. The concentration of antiarrhythmic pharmaceutical agent (e.g. flecainide) for the treatment of arrhythmia can range about 0.1-150 mg/mL, such as about 0.1-130, 0.1-110, 0.1-90, 0.1-70, 0.1-50, 0.1-30, 0.1-10, 0.1-5, 0.1-1.0, 0.1-0.5, 1-150, 1-130, 1-110, 1-90, 1-70, 1-50, 1-30, 1-10, 1-5, 5-150, 5-130, 5-110, 5-90, 5-70, 5-50, 5-30, 5-10, 10-150, 10-130, 10-110, 10-90, 10-70, 10-50, 10-30, 30-150, 30-130, 30-110, 30-90, 30-70, 30-50, 50-150, 50-130, 50-110, 50-90, 50-70, 70-150, 70-130, 70-110, 70-90, 90-150, 90-130, 90-110, 110-150, 110-130, or 130-150 mg/mL. For example, the concentration of antiarrhythmic pharmaceutical agent (e.g. flecainide) for the treatment of arrhythmia can range about 30-50 mg/mL in an acetate buffer.

[0081] The present invention can be directed to a method of self-diagnosing and treating cardiac arrhythmia. The method can comprise self-diagnosing cardiac arrhythmia by detecting at least one of shortness of breath, heart palpitations, and above normal heart rate. The method also can comprise self-administering by inhalation an effective amount of at least one antiarrhythmic pharmaceutical agent within two hours, such as

within one hour, 30 minutes, or within 15 minutes, of the self-diagnosing. In some cases, the subject can self-titrate. For example, the subject can self-administer, e.g., by using an inhaler, until disabling symptoms disappear. In some cases, the self-administering continues until the subject no longer feels heart palpitations.

[0082] The time for onset of action is also typically short. For instance, the subject may have normal sinus rhythm within 20 minutes of initiating the administering, such as within 15 minutes, within 10 minutes, or within 5 minutes of initiating the administering. The rapid onset of action is advantageous because the longer a subject has had arrhythmia, the longer it typically takes to convert the subject to normal sinus rhythm. In some embodiments, the method of the present invention allows the subject to avoid other therapies, such as ablation and/or electrical cardioversion. In other embodiments, the method of the present invention is used in combination with other therapies, such as before or after electrical cardioversion and/or ablation therapy.

Device

[0083] The pharmaceutical composition may be delivered by an inhaler as described in WO 99116420, by a metered dose inhaler as described in WO 99116422, by a liquid dose instillation apparatus as described in WO 99116421. Inhalers impart energy into a liquid pharmaceutical composition to aerosolize the liquid, and to allow delivery to the pulmonary system, e.g., the lungs, of a subject. An inhaler can comprise a liquid delivery system, such as a container having a reservoir that contains a liquid pharmaceutical composition. The liquid pharmaceutical composition generally can comprise an active agent that is either in solution or suspended within a liquid medium.

[0084] In one type of inhaler, generally referred to as a jet inhaler, compressed gas is forced through an orifice in the container. The compressed gas forces liquid to be withdrawn through a nozzle, and the withdrawn liquid mixes with the flowing gas to form aerosol droplets. A cloud of droplets is then administered to the subject's respiratory tract. The compressed gas can be compressed air or compressed oxygen.

[0085] In another type of inhaler, generally referred to as a vibrating mesh inhaler, energy, such as mechanical energy, vibrates a mesh. This vibration of the mesh

aerosolizes the liquid pharmaceutical composition to create an aerosol cloud that is administered to the subject's lungs. The vibrating mesh inhaler can comprise a metal vibrating mesh. In some cases, iron constitutes at least about 30% weight, such as 30%, 40%, 50%, 60%, 70%, 80%, or 90% of the metal vibrating mesh. An inhaler of the vibrating mesh type, such as one that that forms droplets without the use of compressed gas, such as the Aeroneb® Pro can provide unexpected improvement in dosing efficiency and consistency. By generating fine droplets by using a vibrating perforated or unperforated membrane, rather than by introducing compressed air, the aerosolized pharmaceutical formulation can be introduced without substantially affecting the flow characteristics. In addition, the generated droplets when using an inhaler of this type are introduced at a low velocity, thereby decreasing the likelihood of the droplets being driven to an undesired region.

[0086] In still another type of inhaler, generally referred to as an ultrasonic wave inhaler, ultrasonic waves or ultrasonic agitations are generated to directly vibrate and aerosolize the pharmaceutical formulation.

[0087] Alternatively or additionally, the pharmaceutical composition may be in a liquid form and may be aerosolized using an inhaler as described in WO 2004/071368, which is herein incorporated by reference in its entirety, as well as U.S. Published application Nos. 2004/0011358 and 2004/0035413, which are both herein incorporated by reference in their entireties. Other examples of inhalers include, but are not limited to, the Aeroneb Go or Aeroneb®Pro nebulizers, available from Aerogen Ltd. of Galway, Ireland; the PARI eFlow and other PARI nebulizers available from PARI Respiratory Equipment, Inc. of Midlothian, Va.; the Lumiscope® Nebulizer 6600 or 6610 available from Lumiscope Company, Inc. of East Brunswick, N.J.; and the Omron NE-U22 available from Omron Healthcare, Inc. of Kyoto, Japan.

[0088] The inhaler has the ability to rapidly deliver the aerosol at a rate that assures availability of a bolus dose in the heart. The inhaler can aerosolize the pharmaceutical composition in short amount of time. In some cases, the inhaler can aerosolize the pharmaceutical composition (e.g. at least 30 mg of flecainide) in less than about 20 minutes, such as less than about 10 seconds, 20 seconds, 30 seconds, 1 minute, 2

minutes, 3 minutes, 4 minutes, 5 minutes, 6 minutes, 7 minutes, 8 minutes, 9 minutes, 10 minutes, 11 minutes, 12 minutes, 13 minutes, 14 minutes, 15 minutes, 16 minutes, 17 minutes, 18 minutes, 19 minutes, or 20 minutes. For example, the inhaler can aerosolize at least 30 mg of flecainide in less than 3 minutes. In some cases, the inhaler can aerosolize the pharmaceutical composition in about 10 seconds to 20 minutes, such as about 10-20 seconds, 10-30 seconds, 10 seconds to 1 minute, 10 seconds to 2 minute, 10 seconds to 3 minute, 20-30 seconds, 20 seconds to 1 minute, 20 seconds to 2 minute, 20 seconds to 3 minute, 30 seconds to 1 minute, 30 seconds to 2 minutes, 30 seconds to 3 minutes, 1-2 minutes, 1-5 minutes, 2-20 minutes, 2-10 minutes, 2-5 minutes, 5-20 minutes, 5-10 minutes, or 10-20 minutes. For example, the inhaler can aerosolize at least 30 mg of flecainide in about 30 seconds to 3 minutes.

[0089] One or more embodiments of the present invention may be used in conjunction with liquid dose instillation or LDI techniques as disclosed in, for example, WO 99116421, which is incorporated herein by reference in its entirety. Liquid dose instillation involves the direct administration of a formulation to the lung. With respect to LDI the formulations are preferably used in conjunction with partial liquid ventilation or total liquid ventilation. Moreover, one or more embodiments of the present invention may further comprise introducing a therapeutically beneficial amount of a physiologically acceptable gas (such as nitric oxide or oxygen) into the pharmaceutical micro-dispersion prior to, during or following administration.

[0090] The pharmaceutical composition of one or more embodiments of the present invention typically has improved emitted dose efficiency. Accordingly, high doses of the pharmaceutical composition may be delivered using a variety of aerosolization devices and techniques. The emitted dose (ED) of the particles of the present invention may be greater than about 30%, such as greater than about 40%, greater than about 50%, greater than about 60%, or greater than about 70%.

[0091] The present invention may involve a follow-up inhalation if no cardioversion occurs after an initial inhalation. Typically, if no cardioversion occurs within 30 minutes of the initial inhalation, the follow-up dosage is higher or the same as the initial dosage. The dosing may be guided by how the subject feels.

Electronic systems

[0092] In some cases, an electronic system can be used to monitor an occurrence or incidence of atrial fibrillation, and administer treatment to a subject in need thereof. Exemplary embodiments of a system can include an electronic monitoring chip, an electronic monitoring device, and an inhaler. Each component of a system can be in communication with any or all components of the system. The terms “communication,” “communicated,” “in communication with,” “transmission,” and “receiving” can refer to electronic exchange or communication between devices. In some instances, the exchange can occur through a wired connection between the devices such as through the use of a USB cord or fiber optic connection. In some instances, the exchange can occur through a wireless connection between the devices. In some cases, a wireless signal can be sent through a mobile data network such as a 4G LTE or 3G data signal. In some cases, a wireless signal can be sent through a Bluetooth connection. In some cases, a wireless signal can be sent through a Wi-Fi connection. In some cases, a wireless signal can be sent through an infra-red data association link.

[0093] An electronic monitoring chip can be inserted or implanted into a subject, and be used to detect a signal that correlates to an ECG in the subject. Alternatively, or in addition to insertion, an electronic monitoring chip can be worn by a subject. In some cases, an electronic monitoring device can be a component of a wearable electronic device such as a watch, ankle bracelet, and the like. An electronic monitoring chip can be in communication with an electronic monitoring device, an inhaler, or both. In some exemplary embodiments, a communication can be a wireless communication.

[0094] An electronic monitoring device may include but not limited to hand held heart monitors, hand held ECG monitors, Bluetooth App-Enabled Monitors, and the like. In some cases, a smartphone can be used as an electronic monitoring device. Such a device can include an application that can allow a smartphone to communicate with a that have the ability to transmit the cardiac readings to a monitoring center and/or chips and loop recorders inserted into a subject for the purpose of diagnosis and pharmacological intervention.

[0095] In some cases, an electronic monitoring device can include means to instruct a subject, or other personnel. In some cases, an audible source, a tactile source, and/or a visual source can be included. In some cases, an audible source can be a device producing sound. In some embodiments an audible source can be one or more speakers. A tactile source can comprise a tactile sensor. In some embodiments, a tactile sensor can detect a stimuli resulting from mechanical stimulation, or temperature. A visual source can be a light, multiple lights, gauge, display or the like. For example, in an exemplary embodiment the visual source can be a screen on a handheld device.

[0096] In some cases, an inhaler as described herein can be employed as part of a system. In some instances, the inhaler can be in communication with an electronic monitoring system. An electronic monitoring system can be used to direct administration of a pharmaceutical composition by an inhaler in communication therewith. In some cases, an electronic device can allow administration of the pharmaceutical composition by the inhaler. For example, an electronic device may allow administration when an electronic monitoring chip detects an incidence of atrial fibrillation. In some cases, an electronic device can prevent administration of the pharmaceutical composition by the inhaler. For example, an electronic device may prevent administration when an electronic monitoring chip does not detect an incidence of atrial fibrillation; when a healthcare professional overrides an administration; or when administration is deemed unsafe.

[0097] In some cases, an inhaler can contain visual, audible, or tactile sources as described above.

[0098] Furthermore, a system described herein can perform self-monitoring checks to detect and alert of potential fault conditions.

[0099] In some instances, dosing may be guided by a portable ECG. For instance, the dosing may be guided by a heart monitoring device (e.g. Holter monitor). In another version, the pharmaceutical composition is administered prophylactically to a subject who is likely to develop an arrhythmia. For example, a subject who has a history of arrhythmias can be prophylactically treated with a pharmaceutical composition comprising antiarrhythmic pharmaceutical agent to reduce the likelihood of developing

an arrhythmia. The pharmaceutical composition may be administered to a subject in any regimen which is effective to prevent an arrhythmia. Illustrative prophylactic regimes include administering an antiarrhythmic pharmaceutical agent as described herein 1 to 21 times per week. In some cases, an overall cumulative arrhythmic burden is controlled by monitoring.

[0100] In some cases, a result can be communicated to a healthcare profession. A healthcare professional can include a physician, a paramedic, a nurse, and the like. A system or device herein can be employed by a healthcare professional in a number of settings, including, but not limited to, a hospital, hospice, clinic, office, ambulance, nursing home, and the like. In some cases, a result can be communicated to a non-healthcare professional. In some cases, a result can be communicated to a monitoring center. In some instances, a monitoring center can contact the subject between 0 secs to 60 mins for the purpose of communicating the diagnosis and ascertaining the subject's ability to self-administer the dose of the inhaled antiarrhythmic.

[0101] Employment of a system described herein can produce an overall management of arrhythmia. This can result in fewer hospital visits, ER visits and extended hospitalizations.

[0102] In some cases, a system or component thereof can be capable of communicating with doctors, health providers, health provider organizations or their computerized systems to notify that the arrhythmia has been detected, the doses and nature of the pharmacological agent delivered, and the response of the arrhythmia to the pharmacological dose.

[0103] A method can comprise administering to free breathing subjects by way of an aerosol generator device and/or system for administration of aerosolized medicaments such as those disclosed in U.S. Published Application Nos. 20050235987, 20050211253, 20050211245, 20040035413, and 20040011358, the disclosures of which are incorporated herein by reference in their entireties.

[0104] Also disclosed herein are kits for treating cardiac arrhythmias. For instance, the kit may include an aerosolization device and/or a container, e.g., unit dose receptacle, containing aerosolizable antiarrhythmic pharmaceutical agent (e.g., flecainide), for

example, as a liquid or dry powder. The kit may further comprise a package, such as a bag, that contains the aerosolization device and/or the container. The kit may further comprise instructions for the aerosolization device and/or the container.

EXAMPLES

Example 1: Evaluation of rate of aerosolization in different buffer and solution systems

[0105] The methods disclosed herein relate to the nebulization of flecainide acetate solution for treating AF, PAF and PSVT, e.g. when they manifest as acute episodes. It can be essential that the inhaler has the ability to rapidly deliver the aerosol at a rate that assures availability of a bolus dose in the heart. When a pharmaceutical composition of flecainide around 30-40 mg/mL in an acetate buffer is used, one would require 1 to 2.0 ml of the drug to be dosed within 1-3 minutes to be effective.

[0106] The time required for aerosolizing solutions with different buffer and excipient systems were tested and measured using an Omron MicroAir device (Table 1). The aerosolization time for aerosolizing 0.3 mL of each solution was recorded. The device used was a vibrating mesh that is made of a Titanium-nickel alloy.

Table 1. Evaluation of rate of aerosolization in different buffer and solution systems using the Omron MicroAir

Sample	Aerosolization time (s) (0.3 mL)	Rate (mL / min)
0.9% sodium chloride (control)	27	0.67
30 mM acetate	34	0.53
22.5 mM acetate	27	0.67
15 mM acetate	30	0.60
7.5 mM acetate	31	0.58
40 mg/mL solution	281	0.06
30 mg/mL solution	260	0.07
20 mg/mL solution	213	0.08
10 mg/mL solution	131	0.14
40 mg/mL solution	281	0.06

30 mg/mL solution (22.5 mM acetate)	224	0.08
20 mg/mL solution (15 mM acetate)	182	0.10
10 mg/mL solution (7.5 mM acetate)	134	0.13

Example 2: Assessment of different inhalers with flecainide acetate (FA) solution

[0107] Different inhalers were then tested to assess compatibility of material to the different materials of construction and different types of inhalers (Table 2).

Table 2: Assessment of different inhalers with flecainide acetate (FA) solution at 30 mg/mL.

Device	Device Type	Material of construction	Solution	Average (mL/min)	St dev (mL/min)
Voyager	Vibrating Mesh	Not known	30 mg/mL FA	0.16	0.01
			Saline	0.29	0.01
AeroNeb Solo		Nickel-Palladium	30 mg/mL FA	0.06	0.01
			Saline	0.55	0.01
Pari eRapid		Stainless steel	30 mg/mL FA	0.61	0.06
			Saline	0.84	0.09
MiniBreeze	Ultrasonic	No mesh	30 mg/mL FA	0.13	0.04
			Saline	0.29	0.02
Lumiscope			30 mg/mL FA	0.85	0.03
			Saline	1.08	0.07
CompXP	Compressed air		30 mg/mL FA	0.14	0.02
			Saline	0.14	0.01
Trek S (PARI)			30 mg/mL FA	0.34	0.01
			Saline	0.26	0.01

[0108] The above results showed that a 30-40 mg dose of flecainide can be effectively delivered using the Pari e-rapid or Lumiscoppe inhalers in less than 2 minutes. The Trek S from PARI was able to deliver the same dose in about 3 minutes.

Example 3: Assessment of flecainide acetate (FA) efficacy using an electronic monitoring system

[0109] An exemplary smart phone app was installed on a smartphone in order to monitor the efficacy of FA at ceasing incidence of atrial fibrillation. The smartphone app allowed for communication between the smart phone and an electronic monitoring chip used to produce an ECG.

[0110] Figure 7 depicts an exemplary ECG produced from a smartphone as compared to a Prucka System in an animal in sinus rhythm. The traces show excellent correlation between the two devices.

[0111] Figure 8 depicts an exemplary ECG produced from a smartphone as compared to a Prucka System in an animal displaying atrial fibrillation. Again, the traces show excellent correlation between the two devices.

[0112] The smart phone device was then used to monitor the treatment efficacy of an intra-tracheal administration of 0.75 mg/kg flecainide in treating atrial fibrillation.

Figure 9 depicts an exemplary ECG produced from a smartphone as compared to a Prucka System in an animal displaying atrial fibrillation. Again, the traces show excellent correlation between the two devices. Each device shows a return to sinus rhythm upon administration. Figure 10 depicts the actual employment of the smart phone to monitor the return to sinus rhythm in the animal. Figure 12 depicts a comparison of the animal before and after administration of flecainide using the smartphone, which again showed a conversion back to sinus rhythm in the animal.

[0113] While preferred embodiments of the present invention have been shown and described herein, it will be obvious to those skilled in the art that such embodiments are provided by way of example only. Numerous variations, changes, and substitutions will now occur to those skilled in the art without departing from the invention. It should be understood that various alternatives to the embodiments of the invention described herein may be employed in practicing the invention. It is intended that the following claims define the scope of the invention and that methods and structures within the scope of these claims and their equivalents be covered thereby.

WHAT IS CLAIMED IS:

1. A method of treating cardiac arrhythmia in a subject, comprising:
 - (a) identifying cardiac arrhythmia in the subject with the aid of an electronic monitoring device;
 - (b) aerosolizing a pharmaceutical composition in less than about 9 minutes using an inhaler, wherein the pharmaceutical composition comprises of a therapeutically effective amount of at least one antiarrhythmic or a pharmaceutically acceptable salt thereof; and
 - (c) administering the aerosolized pharmaceutical composition to the subject; thereby treating the cardiac arrhythmia.
2. The method of claim 1, wherein the identifying of (a) includes an establishing of a duration of the cardiac arrhythmia.
3. The method of claim 2, wherein the duration of the cardiac arrhythmia is at least about 2 hours.
4. The method of claim 2, wherein the duration of the cardiac arrhythmia is at least about 1 hour.
5. The method of claim 2, wherein the duration of the cardiac arrhythmia is at least about 0.5 hours.
6. The method of claim 1, wherein the aerosolizing of the pharmaceutical composition occurs in less than 6 minutes.
7. The method of claim 1, wherein the aerosolizing of the pharmaceutical composition occurs in less than 3 minutes.
8. The method of claim 1, wherein the therapeutically effective amount is an amount sufficient to convert the arrhythmia to normal sinus rhythm.
9. The method of claim 8, wherein the therapeutically effective amount is at least about 60 mg of the at least one antiarrhythmic or a pharmaceutically acceptable salt thereof.
10. The method of claim 8, wherein the therapeutically effective amount is at least about 50 mg of the at least one antiarrhythmic or a pharmaceutically acceptable salt thereof.

11. The method of claim 8, wherein the therapeutically effective amount is at least about 40 mg of the at least one antiarrhythmic or a pharmaceutically acceptable salt thereof.
12. The method of claim 8, wherein the therapeutically effective amount is at least about 30 mg of the at least one antiarrhythmic or a pharmaceutically acceptable salt thereof.
13. The method of claim 8, wherein the therapeutically effective amount converts the arrhythmia to normal sinus rhythm from about 0 secs to about 2 hours after the administration.
14. The method of claim 13, wherein the therapeutically effective amount converts the arrhythmia to normal sinus rhythm in about 60 mins after administration.
15. The method of claim 1, wherein the treating occurs within about 120 minutes from an onset of the cardiac arrhythmia.
16. The method of claim 1, wherein the treating occurs within about 90 minutes from an onset of the cardiac arrhythmia.
17. The method of claim 1, wherein the treating occurs within about 60 minutes from an onset of the cardiac arrhythmia.
18. The method of claim 1, wherein the administering comprises inhalation.
19. The method of claim 1, wherein prior to (a), an electronic monitoring chip is inserted into the subject.
20. The method of claim 1, wherein an electronic monitoring chip is worn by the subject.
21. The method of claim 19, wherein the electronic monitoring chip is in communication with the electronic monitoring device.
22. The method of claim 21, wherein the communication is a wireless communication.
23. The method of claim 1, wherein prior to (a), an electronic monitoring chip is inserted into the electronic monitoring device.
24. The method of claim 1, wherein the identifying of (a) comprises a communication of a result to the subject.

25. The method of claim 24, wherein the communication comprises an instruction to administer the pharmaceutical composition.
26. The method of claim 25, wherein the instruction to administer comprises a dosage of the aerosolized pharmaceutical composition to administer.
27. The method of claim 25, wherein the administering is performed by the subject.
28. The method of claim 25, wherein the administering is performed by a healthcare professional.
29. The method of claim 28, wherein the healthcare professional can override a recommendation made by the electronic monitoring device.
30. The method of claim 28, wherein the electronic monitoring device waits for a confirmation from the healthcare provider prior to allowing the administering of the aerosolized pharmaceutical composition to the subject.
31. The method of claim 24, wherein the communication comprises making a treatment recommendation.
32. The method of claim 31, wherein the treatment recommendation comprises restricting the administration of the pharmaceutical composition when the electronic monitoring device does not identify an arrhythmia.
33. The method of claim 1, wherein the identifying of (a) comprises a communication of a result to a healthcare professional.
34. The method of claim 33, wherein the administering is performed by the subject.
35. The method of claim 33, wherein the administering is performed by a healthcare professional.
36. The method of claim 35, wherein the healthcare professional is a paramedic.
37. The method of claim 35, wherein the healthcare professional is a physician.
38. The method of claim 35, wherein the healthcare professional is a monitoring professional.
39. The method of claim 33, wherein the healthcare professional can override a recommendation made by the electronic monitoring device.

40. The method of claim 33, wherein the electronic monitoring device waits for a confirmation from the healthcare provider prior to allowing the administering of the aerosolized pharmaceutical composition to the subject.
41. The method of claim 33, wherein the communication comprises making a treatment recommendation.
42. The method of claim 41, wherein the treatment recommendation comprises restricting the administration of the pharmaceutical composition when the electronic monitoring device does not identify an arrhythmia.
43. The method of claim 1, wherein the at least one antiarrhythmic or salt thereof is a class I antiarrhythmic.
44. The method of claim 43, wherein the class I antiarrhythmic is a class Ia, Ib, or Ic antiarrhythmic.
45. The method of claim 43, wherein the class I antiarrhythmic is flecainide or a salt thereof.
46. The method of claim 1, wherein the at least one antiarrhythmic or salt thereof is a class II antiarrhythmic.
47. The method of claim 1, wherein the at least one antiarrhythmic or salt thereof is a class III antiarrhythmic.
48. The method of claim 1, wherein the at least one antiarrhythmic or salt thereof is a class IV antiarrhythmic.
49. The method of claim 1, wherein the at least one antiarrhythmic or salt thereof is a class V antiarrhythmic.
50. The method of claim 1, wherein the at least one antiarrhythmic or salt thereof comprises a sodium channel blocker.
51. The method of claim 1, wherein the at least one antiarrhythmic or salt thereof comprises a potassium channel blocker.
52. The method of claim 1, wherein the at least one antiarrhythmic or salt thereof comprises a calcium channel blocker.
53. The method of claim 1, wherein the at least one antiarrhythmic or salt thereof is rapidly absorbed through a subject's lungs.

- 54. The method of claim 53, wherein the at least one antiarrhythmic or salt thereof is delivered to a subject's heart after being absorbed through the subject's lungs.
- 55. The method of claim 1, wherein the inhaler is in communication with the electronic monitoring device.
- 56. The method of claim 55, wherein the communication is a wired communication.
- 57. The method of claim 55, wherein the communication is a wireless communication.
- 58. The method of claim 55, wherein the administering of the aerosolized pharmacological composition by the inhaler is communicated to the electronic monitoring device.
- 59. The method of claim 58, wherein the electronic monitoring device verifies the administering of the aerosolized pharmaceutical composition.
- 60. The method of claim 59, wherein the electronic monitoring device continues monitoring the progress of the arrhythmia after the administration.
- 61. The method of claim 60, wherein a recommendation to stop the administering is communicated by the electronic monitoring device when the arrhythmia is converted to a normal sinus rhythm.
- 62. The method of claim 60, wherein a recommendation to seek medical attention is communicated by the electronic monitoring device when the arrhythmia is not responding to the administration of the pharmaceutical composition as determined by a continued incidence of arrhythmia at least about 2 hours after the administering.
- 63. The method of claim 1, wherein the electronic monitoring device comprises a digital display.
- 64. The method of claim 1, wherein the electronic monitoring device comprises auditory means capable of communicating to the subject.
- 65. The method of claim 1, wherein the electronic monitoring device is a smartphone.
- 66. The method of claim 1, wherein the inhaler comprises a digital display.

67. The method of claim 1, wherein the inhaler comprises auditory means capable of communicating to the subject.
68. A system comprising:
- (a) an electronic monitoring chip; wherein the electronic monitoring chip monitors incidence of arrhythmia in a subject;
 - (b) an electronic monitoring device; wherein the electronic monitoring device is in wireless communication with the electronic monitoring chip; and
 - (c) an inhaler; wherein the electronic monitoring device is in communication with inhaler.
69. The system of claim 68, wherein the electronic monitoring device is in communication with the inhaler through a wired connection.
70. The system of claim 68, wherein the electronic monitoring device is in communication with the inhaler through a wireless connection.
71. The system of claim 68, wherein the electronic monitoring device comprises a digital display.
72. The system of claim 68, wherein the electronic monitoring device comprises auditory means capable of communicating to the subject.
73. The system of claim 68, wherein the electronic monitoring device is a smartphone.
74. The system of claim 68, wherein the electronic monitoring chip is inserted into the subject.
75. The system of claim 68, wherein the electronic monitoring chip is worn by the subject.
76. The system of claim 68, wherein the inhaler comprises a digital display.
77. The system of claim 68, wherein the inhaler comprises auditory means capable of communicating to the subject.
78. The system of claim 68, wherein the electronic monitoring device comprises storage means.
79. The system of claim 78, subject data is stored using the storage means.

- 80. The system of claim 78, wherein the storage means is a hard drive.
- 81. The system of claim 78, wherein the storage means is a cloud based storage.

1 / 11




FIG. 1




FIG. 2




FIG. 3




FIG. 4




FIG. 5




FIG. 6




FIG. 7




FIG. 8




FIG. 9

10 / 11




FIG. 10




FIG. 11

INTERNATIONAL SEARCH REPORTInternational application No.
PCT/US2017/016018**Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)**

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 1-67
because they relate to subject matter not required to be searched by this Authority, namely:
Claims 1-67 pertain to a method for treatment of the human body by therapy, and thus relate to a subject matter which this International Searching Authority is not required, under PCT Article 17(2)(a)(i) and PCT Rule 39.1(iv), to search.
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fees, this Authority did not invite payment of any additional fees.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.
- ☐ The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US2017/016018**A. CLASSIFICATION OF SUBJECT MATTER****A61K 31/4458(2006.01)i, A61B 5/0464(2006.01)i, A61B 5/046(2006.01)i, A61M 15/00(2006.01)i**

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

A61K 31/4458; A61P 9/00; A61P 9/06; A61B 5/08; A61B 5/02; A61M 16/00; A61N 1/365; A61B 5/0464; A61M 15/00

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Korean utility models and applications for utility models

Japanese utility models and applications for utility models

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

eKOMPASS(KIPO internal) & Keywords: monitoring chip, device, inhaler, arrhythmia

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 90-03144 A1 (HRUSHESKY, W. J.) 05 April 1990 See abstract; pages 4-8; and figures 1-3B.	68-81
A	US 8641631 B2 (SIERRA, G. et al.) 04 February 2014 See the whole document.	68-81
A	WO 2005-079897 A1 (RESMED LIMITED) 01 September 2005 See the whole document.	68-81
A	US 8974828 B2 (SCHULER, C. A. et al.) 10 March 2015 See the whole document.	68-81
A	US 7302295 B2 (STAHMANN, J. E. et al.) 27 November 2007 See the whole document.	68-81



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

17 April 2017 (17.04.2017)

Date of mailing of the international search report

18 April 2017 (18.04.2017)

Name and mailing address of the ISA/KR

International Application Division

Korean Intellectual Property Office

189 Cheongsa-ro, Seo-gu, Daejeon, 35208, Republic of Korea



Facsimile No. +82-42-481-8578

Authorized officer

CHO, Ki Yun

Telephone No. +82-42-481-5655



INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No.

PCT/US2017/016018

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 90-03144 A1	05/04/1990	US 4930518 A WO 90-03144 A1	05/06/1990 05/04/1990
US 8641631 B2	04/02/2014	EP 1740095 A1 EP 1740095 A4 EP 1740095 B1 JP 2007-532156 A JP 5090155 B2 US 2007-0282212 A1 US 2014-0180154 A1 WO 2005-096931 A1	10/01/2007 05/08/2009 23/01/2013 15/11/2007 05/12/2012 06/12/2007 26/06/2014 20/10/2005
WO 2005-079897 A1	01/09/2005	EP 1718356 A1 EP 1718356 A4 EP 1718356 B1 EP 2388036 A2 EP 2388036 A3 JP 2007-525267 A JP 4699443 B2 US 2008-0045813 A1 US 2015-0182713 A1 US 8794236 B2	08/11/2006 06/05/2009 21/09/2016 23/11/2011 13/03/2013 06/09/2007 08/06/2011 21/02/2008 02/07/2015 05/08/2014
US 8974828 B2	10/03/2015	AU 2010-226613 A1 AU 2010-226613 B2 EP 2413902 A1 EP 2413902 A4 JP 2012-520890 A JP 2015-193657 A US 2012-0003318 A1 US 2015-0313842 A1 WO 2010-107964 A1	29/09/2011 25/07/2013 08/02/2012 19/02/2014 10/09/2012 05/11/2015 05/01/2012 05/11/2015 23/09/2010
US 7302295 B2	27/11/2007	AU 2003-303632 A1 EP 1587422 A2 EP 1587422 B1 EP 1613395 A2 EP 1615692 A2 EP 1617894 A2 EP 1620165 A1 EP 1656181 A1 EP 1656181 B1 EP 1670547 A2 EP 1670547 B1 EP 1732642 A1 EP 1938862 A2 EP 1938862 A3 EP 1938862 B1 EP 2008581 A2	29/07/2004 26/10/2005 26/03/2014 11/01/2006 18/01/2006 25/01/2006 01/02/2006 17/05/2006 04/06/2008 21/06/2006 12/11/2008 20/12/2006 02/07/2008 25/08/2010 19/10/2011 31/12/2008

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No.

PCT/US2017/016018

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
		EP 2008581 A3	25/08/2010
		EP 2008581 B1	17/08/2011
		JP 2006-515191 A	25/05/2006
		JP 2006-522650 A	05/10/2006
		JP 2006-522659 A	05/10/2006
		JP 2006-522661 A	05/10/2006
		JP 2006-524106 A	26/10/2006
		JP 2007-502670 A	15/02/2007
		JP 2007-532178 A	15/11/2007
		JP 4521396 B2	11/08/2010
		JP 4750032 B2	17/08/2011
		JP 4794168 B2	19/10/2011
		US 2004-0127807 A1	01/07/2004
		US 2004-0204728 A1	14/10/2004
		US 2004-0215239 A1	28/10/2004
		US 2004-0215240 A1	28/10/2004
		US 2004-0215258 A1	28/10/2004
		US 2004-0220629 A1	04/11/2004
		US 2004-0220633 A1	04/11/2004
		US 2004-0230129 A1	18/11/2004
		US 2004-0230229 A1	18/11/2004
		US 2004-0230230 A1	18/11/2004
		US 2004-0230273 A1	18/11/2004
		US 2004-0230274 A1	18/11/2004
		US 2004-0230279 A1	18/11/2004
		US 2004-0230280 A1	18/11/2004
		US 2004-0230281 A1	18/11/2004
		US 2004-0230282 A1	18/11/2004
		US 2005-0004615 A1	06/01/2005
		US 2005-0038350 A1	17/02/2005
		US 2005-0039745 A1	24/02/2005
		US 2005-0042589 A1	24/02/2005
		US 2005-0043644 A1	24/02/2005
		US 2005-0043652 A1	24/02/2005
		US 2005-0043772 A1	24/02/2005
		US 2005-0061315 A1	24/03/2005
		US 2005-0061319 A1	24/03/2005
		US 2005-0061320 A1	24/03/2005
		US 2005-0061323 A1	24/03/2005
		US 2005-0065447 A1	24/03/2005
		US 2005-0065448 A1	24/03/2005
		US 2005-0065560 A1	24/03/2005
		US 2005-0065566 A1	24/03/2005
		US 2005-0065567 A1	24/03/2005
		US 2005-0065572 A1	24/03/2005
		US 2005-0074741 A1	07/04/2005
		US 2005-0076905 A1	14/04/2005
		US 2005-0076908 A1	14/04/2005
		US 2005-0076909 A1	14/04/2005
		US 2005-0080348 A1	14/04/2005

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No.

PCT/US2017/016018

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
		US 2005-0080461 A1	14/04/2005
		US 2005-0080463 A1	14/04/2005
		US 2005-0081847 A1	21/04/2005
		US 2005-0085738 A1	21/04/2005
		US 2005-0107838 A1	19/05/2005
		US 2005-0109338 A1	26/05/2005
		US 2005-0109339 A1	26/05/2005
		US 2005-0113710 A1	26/05/2005
		US 2005-0115561 A1	02/06/2005
		US 2005-0119708 A1	02/06/2005
		US 2005-0142070 A1	30/06/2005
		US 2005-0145246 A1	07/07/2005
		US 2006-0025827 A1	02/02/2006
		US 2007-0293896 A1	20/12/2007
		US 2008-0071317 A1	20/03/2008
		US 2008-0091242 A1	17/04/2008
		US 2008-0139948 A1	12/06/2008
		US 2008-0221468 A1	11/09/2008
		US 2008-0312548 A1	18/12/2008
		US 2009-0177702 A1	09/07/2009
		US 2009-0270750 A1	29/10/2009
		US 2009-0308395 A1	17/12/2009
		US 2010-0010378 A1	14/01/2010
		US 2010-0106211 A1	29/04/2010
		US 2010-0174154 A1	08/07/2010
		US 2010-0174207 A1	08/07/2010
		US 2010-0174335 A1	08/07/2010
		US 2010-0179613 A1	15/07/2010
		US 2011-0061647 A1	17/03/2011
		US 2011-0098587 A1	28/04/2011
		US 2011-0137197 A1	09/06/2011
		US 2011-0201955 A1	18/08/2011
		US 6949075 B2	27/09/2005
		US 7117035 B2	03/10/2006
		US 7218966 B2	15/05/2007
		US 7302294 B2	27/11/2007
		US 7336996 B2	26/02/2008
		US 7349742 B2	25/03/2008
		US 7364547 B2	29/04/2008
		US 7396333 B2	08/07/2008
		US 7468040 B2	23/12/2008
		US 7469697 B2	30/12/2008
		US 7493175 B2	17/02/2009
		US 7499758 B2	03/03/2009
		US 7510531 B2	31/03/2009
		US 7529592 B2	05/05/2009
		US 7532934 B2	12/05/2009
		US 7555335 B2	30/06/2009
		US 7566318 B2	28/07/2009
		US 7570997 B2	04/08/2009

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No.

PCT/US2017/016018

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
		US 7572225 B2	11/08/2009
		US 7575553 B2	18/08/2009
		US 7578794 B2	25/08/2009
		US 7591265 B2	22/09/2009
		US 7610094 B2	27/10/2009
		US 7616988 B2	10/11/2009
		US 7662101 B2	16/02/2010
		US 7664546 B2	16/02/2010
		US 7668591 B2	23/02/2010
		US 7678061 B2	16/03/2010
		US 7680537 B2	16/03/2010
		US 7702399 B2	20/04/2010
		US 7715916 B2	11/05/2010
		US 7720541 B2	18/05/2010
		US 7757690 B2	20/07/2010
		US 7787946 B2	31/08/2010
		US 7865233 B2	04/01/2011
		US 7887493 B2	15/02/2011
		US 7931600 B2	26/04/2011
		US 7938782 B2	10/05/2011
		US 7967756 B2	28/06/2011
		US 7970470 B2	28/06/2011
		US 7979122 B2	12/07/2011
		US 7981042 B2	19/07/2011
		US 7993279 B2	09/08/2011
		US 7996071 B2	09/08/2011
		US 8002553 B2	23/08/2011
		US 8024039 B2	20/09/2011
		US 8104470 B2	31/01/2012
		US 8192376 B2	05/06/2012
		US 8221327 B2	17/07/2012
		US 8251061 B2	28/08/2012
		US 8317719 B2	27/11/2012
		US 8321022 B2	27/11/2012
		US 8323204 B2	04/12/2012
		US 8380296 B2	19/02/2013
		US 8606356 B2	10/12/2013
		US 8655446 B2	18/02/2014
		US 8657756 B2	25/02/2014
		WO 2004-060166 A2	22/07/2004
		WO 2004-060166 A3	13/01/2005
		WO 2004-091715 A1	28/10/2004
		WO 2004-091717 A2	28/10/2004
		WO 2004-091717 A3	23/12/2004
		WO 2004-091719 A2	28/10/2004
		WO 2004-091719 A3	29/12/2004
		WO 2004-091720 A2	28/10/2004
		WO 2004-091720 A3	03/02/2005
		WO 2005-018737 A1	03/03/2005
		WO 2005-028029 A2	31/03/2005