



HU000025156T2

(19) **HU****MAGYARORSZÁG**
Szellemi Tulajdon Nemzeti Hivatala(11) Lajstromszám: **E 025 156**(13) **T2****EURÓPAI SZABADALOM**
SZÖVEGÉNEK FORDÍTÁSA(21) Magyar ügyszám: **E 10 712218**(22) A bejelentés napja: **2010. 03. 24.**(51) Int. Cl.: **A01N 43/50** (2006.01)**C07K 143/25** (2006.01)**C12N 15/82** (2006.01)

(96) Az európai bejelentés bejelentési száma:

EP 20100712218

(97) Az európai bejelentés közzétételi adatai:

EP 2419441 A1 **2010. 10. 21.**

(86) A nemzetközi (PCT) bejelentési szám:

PCT/US 10/028381

(97) Az európai szabadalom megadásának meghirdetési adatai:

EP 2419441 B1 **2015. 01. 21.**

(87) A nemzetközi közzétételi szám:

WO 10120452

(30) Elsőbbségi adatok:

170189 P **2009. 04. 17.** **US**

(73) Jogosult(ak):

Dow AgroSciences LLC, Indianapolis IN
46268-1054 (US)

(72) Feltaláló(k):

LIRA, Justin, Fishers IN 46038 (US)
BUTLER, Holly, Indianapolis IN 46260 (US)
SMITH, Doug, Noblesville IN 46062 (US)
NARVA, Kenneth, Zionsville IN 46077 (US)
MEADE, Thomas, Zionsville IN 46077 (US)

(74) Képviselő:

dr. Kompagne Hajnalka, DANUBIA Szabadalmi
és Jogi Iroda Kft., Budapest

(54)

Inszekticid DIG-3 CRY-toxinok

Az európai szabadalom ellen, megadásának az Európai Szabadalmi Közlönyben való meghirdetésétől számított kilenc hónapon belül, felszólalást lehet benyújtani az Európai Szabadalmi Hivatalnál. (Európai Szabadalmi Egyezmény 99. cikk(1))

A fordítást a szabadalmas az 1995. évi XXXIII. törvény 84/H. §-a szerint nyújtotta be. A fordítás tartalmi helyességét a Szellemi Tulajdon Nemzeti Hivatala nem vizsgálta.



(11) **EP 2 419 441 B1**

(12) **EUROPEAN PATENT SPECIFICATION**

(45) Date of publication and mention
of the grant of the patent:
21.01.2015 Bulletin 2015/04

(51) Int Cl.:
A01N 43/50 ^(2006.01) **C12N 15/82** ^(2006.01)
C07K 14/325 ^(2006.01)

(21) Application number: **10712218.6**

(86) International application number:
PCT/US2010/028381

(22) Date of filing: **24.03.2010**

(87) International publication number:
WO 2010/120452 (21.10.2010 Gazette 2010/42)

(54) **DIG-3 INSECTICIDAL CRY TOXINS**

INSEKTIZIDE DIG-3-CRY-TOXINE

TOXINES CRY INSECTICIDES COMPRENANT DIG-3

(84) Designated Contracting States:
**AT BE BG CH CY CZ DE DK EE ES FI FR GB GR
HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL
PT RO SE SI SK SM TR**
Designated Extension States:
RS

• **MEADE, Thomas**
Zionsville
IN 46077 (US)

(30) Priority: **17.04.2009 US 170189 P**

(74) Representative: **Zwicker, Jörk et al**
ZSP Patentanwälte
Partnerschaftsgesellschaft
Radlkoferstrasse 2
81373 München (DE)

(43) Date of publication of application:
22.02.2012 Bulletin 2012/08

(56) References cited:
WO-A1-2004/020636 WO-A1-2008/145406
WO-A2-01/19859 WO-A2-99/24581
US-A1- 2009 023 653

(73) Proprietor: **Dow AgroSciences LLC**
Indianapolis IN 46268-1054 (US)

(72) Inventors:
• **LIRA, Justin**
Fishers
IN 46038 (US)
• **BUTLER, Holly**
Indianapolis
IN 46260 (US)
• **SMITH, Doug**
Noblesville
IN 46062 (US)
• **NARVA, Kenneth**
Zionsville
IN 46077 (US)

• **KUO WHITE-SHANG ET AL: "Cloning of two new
cry genes from Bacillus thuringiensis subsp.
wuhanensis strain" CURRENT MICROBIOLOGY,**
vol. 40, no. 4, April 2000 (2000-04), pages 227-232,
XP002589373 ISSN: 0343-8651
• **DATABASE UniProt [Online] 1 October 2002**
(2002-10-01), "SubName: Full=CryIBII;"
XP002590686 retrieved from EBI accession no.
UNIPROT:Q8KNY2 Database accession no.
Q8KNY2

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

DescriptionCross Reference to Related Applications

[0001] This application claims benefit to U.S. Provisional Patent Application No. 61/170,189, filed April 17, 2009.

Field of the Invention

[0002] This invention concerns new insecticidal Cry toxins and their use to control insects.

Background of the Invention

[0003] *Bacillus thuringiensis* (*B.t.*) is a soil-borne bacterium that produces pesticidal crystal proteins known as delta endotoxins or Cry proteins. Cry proteins are oral intoxicants that function by acting on midgut cells of susceptible insects. An extensive list of delta endotoxins is maintained and regularly updated at http://www.lifesci.sussex.ac.uk/home/Neil_Crickmore/Bt/intro.html.

[0004] European corn borer (ECB), *Ostrinia nubilalis* (Hübner), is the most damaging insect pest of corn throughout the United States and Canada, and causes an estimated \$1 billion revenue loss each year due to crop yield loss and expenditures for insect management (Witkowski *et al.*, 2002). Transgenic corn expressing genes encoding Cry proteins, most notably CryIAb, Cry1Ac, or CryIF, provide commercial levels of efficacy against ECB.

[0005] Despite the success of ECB-resistant transgenic corn, the possibility of the development of resistant insect populations threatens the long-term durability of Cry proteins in ECB control and creates the need to discover and develop new Cry proteins to control ECB and other pests. Insect resistance to *B.t.* Cry proteins can develop through several mechanisms (Heckel *et al.*, 2007, Pigott and Ellar, 2007). Multiple receptor protein classes for Cry proteins have been identified within insects, and multiple examples exist within each receptor class. Resistance to a particular Cry protein may develop, for example, by means of a mutation within the toxin-binding portion of a cadherin domain of a receptor protein. A further means of resistance may be mediated through a protoxin-processing protease. Thus, resistance to Cry toxins in species of Lepidoptera has a complex genetic basis, with at least four distinct, major resistance genes. Lepidopteran insects resistant to Cry proteins have developed in the field within the species *Plutella xylostella* (Tabashnik, 1994), *Trichoplusia ni* (Janmaat and Myers 2003, 2005), and *Helicoverpa zea* (Tabashnik *et al.*, 2008). Development of new high potency Cry proteins would provide additional tools for management of ECB and other insect pests. Cry proteins with different modes of action produced in combination in transgenic corn would prevent the development ECB insect resistance and protect the long term utility of *B.t.* technology for insect pest control.

Brief Summary of the Invention

[0006] The present application relates to insecticidal Cry toxins, including the toxin designated herein as DIG-3 as well as variants of DIG-3, nucleic acids encoding these toxins, methods of controlling pests using the toxins, methods of producing the toxins in transgenic host cells, and transgenic plants that produce the toxins. The predicted amino acid sequence of the wild type DIG-3 toxin is given in SEQ ID NO:2.

[0007] As described in Example 1, a nucleic acid encoding the DIG-3 protein was isolated from a *B.t.* strain internally designated by Dow AgroSciences LLC as PS46L. The nucleic acid sequence for the full length coding region was determined, and the full length protein sequence was deduced from the nucleic acid sequence. The DIG-3 toxin has some similarity to Cry1BII (Genbank Accession No. AAM93496) and other *B. thuringiensis* Cry1B-type proteins (http://www.lifesci.sussex.ac.uk/home/Neil_Crickmore/Bt/intro.html).

[0008] Insecticidally active variants of the DIG-3 toxin are also described herein, and are referred to collectively as DIG-3 toxins.

[0009] DIG-3 toxins may also be used in combination with RNAi methodologies for control of other insect pests. For example, DIG-3 can be used in transgenic plants in combination with a dsRNA for suppression of an essential gene in corn rootworm or an essential gene in an insect pest. Such target genes include, for example, vacuolar ATPase, ARF-1, Act42A, CHD3, EF-1 α , and TFIIB. An example of a suitable target gene is vacuolar ATPase, as disclosed in WO2007/035650.

[0010] It is reported herein that DIG-3 toxins are active against populations of European corn borer and diamond back moth that are resistant to Cry1F and Cry1A toxins. Accordingly, DIG-3 toxins are ideal candidates for use to control of Lepidopteran pests. The toxins can be used alone or in combination with other Cry toxins, such as Cry1F, CryIAb, and CryIAc, to control development of resistant insect populations.

[0011] Insecticidally active fragments of SEQ ID NO:2, and nucleotides encoding such fragments, are aspects of the invention.

[0012] In one embodiment the invention provides an isolated DIG-3 toxin polypeptide comprising a core toxin segment selected from the group consisting of

(a) a polypeptide comprising the amino acid sequence of residues 113 to 643 of SEQ ID NO:2;

(b) a polypeptide comprising an amino acid sequence having at least 90% sequence identity to the amino acid sequence of residues 113 to 643 of SEQ ID NO:2;

(c) a polypeptide comprising an amino acid sequence of residues 113 to 643 of SEQ ID NO:2 with up to 20 amino acid substitutions, deletions, or modifications that do not adversely affect expression or activity of the toxin encoded by SEQ ID NO:2.

[0013] In one embodiment the invention provides an isolated DIG-3 toxin polypeptide comprising a core toxin segment selected from the group consisting of

(a) a polypeptide comprising the amino acid sequence of residues 73 to 643 of SEQ ID NO:2;

(b) a polypeptide comprising an amino acid sequence having at least 90% sequence identity to the amino acid sequence of residues 73 to 643 of SEQ ID NO:2;

(c) a polypeptide comprising an amino acid sequence of residues 73 to 643 of SEQ ID NO:2 with up to 20 amino acid substitutions, deletions, or modifications that do not adversely affect expression or activity of the toxin encoded by SEQ ID NO:2.

[0014] In another embodiment the invention provides an isolated DIG-3 toxin polypeptide comprising a DIG-3 core toxin segment selected from the group consisting of

(a) a polypeptide comprising the amino acid sequence of residues 1 to 643 of SEQ ID NO:2;

(b) a polypeptide comprising an amino acid sequence having at least 90% sequence identity to the amino acid sequence of residues 1 to 643 of SEQ ID NO:2;

(c) a polypeptide comprising an amino acid sequence of residues 1 to 643 of SEQ ID NO:2 with up to 20 amino acid substitutions, deletions, or modifications that do not adversely affect expression or activity of the toxin encoded by SEQ ID NO:2.

[0015] By "isolated" applicants mean that the polypeptide or DNA molecules have been removed from their native environment and have been placed in a different environment by the hand of man.

[0016] In another embodiment the invention provides a plant comprising a DIG-3 toxin.

[0017] In another embodiment the invention provides a method for controlling a pest population comprising contacting said population with a pesticidally effective amount of a DIG-3 toxin.

[0018] In another embodiment the invention provides an isolated nucleic acid that encodes a DIG-3 toxin.

[0019] In another embodiment the invention provides a DNA construct comprising a nucleotide sequence that encodes a DIG-3 toxin operably linked to a promoter that is not derived from *Bacillus thuringiensis* and is capable of driving expression in a plant. The invention also provides a transgenic plant that comprises the DNA construct stably incorporated into its genome and a method for protecting a plant from a pest comprising introducing the construct into said plant.

Brief Description of the Sequences

[0020] SEQ ID NO:1 DNA sequence encoding full-length DIG-3 toxin; 3771 nt.

[0021] SEQ ID NO:2 Full-length DIG-3 protein sequence; 1256 aa.

[0022] SEQ ID NO:3 Plant-optimized full length DIG-3 DNA sequence; 3771 nt.

[0023] SEQ ID NO:4 Cry1Ab protoxin segment; 545 aa.

[0024] SEQ ID NO:5 Chimeric toxin: DIG-3 Core toxin segment/Cry1Ab protoxin segment; 1188 aa.

[0025] SEQ ID NO:6 Dicot-optimized DNA sequence encoding the Cry1Ab protoxin segment; 1635 nt

[0026] SEQ ID NO:7 Maize-optimized DNA sequence encoding the Cry1Ab protoxin segment; 1635 nt

Detailed Description of the Invention

[0027] DIG-3 Toxins, and insecticidally active variants. In addition to the full length DIG-3 toxin of SEQ ID NO:2, the invention encompasses insecticidally active variants. By the term "variant", applicants intend to include fragments, certain deletion and insertion mutants, and certain fusion proteins. DIG-3 is a classic three-domain Cry toxin. As a preface to describing variants of the DIG-3 toxin that are included in the invention, it will be useful to briefly review the architecture of three-domain Cry toxins in general and of the DIG-3 protein toxin in particular.

[0028] A majority of *Bacillus thuringiensis* delta-endotoxin crystal protein molecules are composed of two functional segments. The protease-resistant core toxin is the first segment and corresponds to about the first half of the protein molecule. The fully ~130 kDa protoxin molecule is rapidly processed to the resistant core segment by proteases in the insect gut. The segment that is deleted by this processing will be referred to herein as the "protoxin segment". The protoxin segment is believed to participate in toxin crystal formation (Arvidson *et al.*, 1989). The protoxin segment may thus convey a partial insect specificity for the toxin by limiting the accessibility of the core to the insect by reducing the protease processing of the toxin molecule (Haider *et al.*, 1986) or by reducing toxin solubility (Aronson *et al.*, 1991). *B.t.* toxins, even within a certain class, vary to some extent in length and in the precise location of the transition from the core toxin segment to protoxin segment. The transition from core toxin segment to protoxin segment will typically occur at between about 50% to about 60% of the full length toxin. SEQ ID NO:2 discloses the 1256 amino acid sequence of the full-length DIG-3 polypeptide, of which the N-terminal 643 amino acids comprise the DIG-3 core toxin segment. The 5'-terminal 1929 nucleotides of SEQ ID NO:1 comprise the coding region for the core toxin segment.

[0029] Three dimensional crystal structures have been determined for Cry1Aa1, Cry2Aa1, Cry3Aa1, Cry3Bb1, Cry4Aa, Cry4Ba and Cry8Ea1. These structures for the core toxins are remarkably similar and are comprised of three distinct domains with the features described below (reviewed in de Maagd *et al.*, 2003).

[0030] Domain I is a bundle of seven alpha helices where α -helix 5 is surrounded by six amphipathic helices. This domain has been implicated in pore formation and shares homology with other pore forming proteins including hemolysins and colicins. Domain I of the DIG-3 protein comprises amino acid residues 56 to 278 of SEQ ID NO:2.

[0031] Domain II is formed by three anti-parallel beta sheets packed together in a beta prism. The loops of this domain play important roles in binding insect midgut receptors. In Cry1A proteins, surface exposed loops at the apices of Domain II beta sheets are involved in binding to Lepidopteran cadherin receptors. Cry3Aa Domain II loops bind a membrane-associated metalloprotease of *Leptinotarsa decemlineata* (Say) (Colorado potato beetle) in a similar fashion (Ochoa-Campuzano *et al.*, 2007). Domain II shares homology with certain carbohydrate-binding proteins including vitelline and jacaline. Domain II of the DIG-3 protein comprises amino acid residues 283 to 493 of SEQ ID NO:2.

[0032] Domain III is a beta sandwich of two anti-parallel beta sheets. Structurally this domain is related to carbohydrate-binding domains of proteins such as glucanases, galactose oxidase, sialidase and others. Domain III binds certain classes of receptor proteins and perhaps participates in insertion of an oligomeric toxin pre-pore that interacts with a second class of receptors, examples of which are aminopeptidase and alkaline phosphatase in the case of Cry1A proteins (Pigott and Ellar, 2007). Analogous Cry Domain III receptors have yet to be identified in Coleoptera. Conserved *B.t.* sequence blocks 2 and 3 map near the N-terminus and C-terminus of Domain 2, respectively. Hence, these conserved sequence blocks 2 and 3 are approximate boundary regions between the three functional domains. These regions of conserved DNA and protein homology have been exploited for engineering recombinant *B.t.* toxins (US Patent No. 6090931, WO 91/01087, WO 95/06730, WO 1998022595). Domain III of the DIG-3 protein comprises amino acid residues 503 to 641 of SEQ ID NO:2.

[0033] It has been reported that α -helix 1 of Domain I is removed following receptor binding. Aronson *et al.* (1999) demonstrated that Cry1Ac bound to BBMV was protected from proteinase K cleavage beginning at residue 59, just after α -helix 1; similar results were cited for Cry1Ab. Gomez *et al.* (2002) found that Cry1Ab oligomers formed upon BBMV receptor binding lacked the α -helix 1 portion of Domain I. Also, Soberon *et al.* (2007) have shown that N-terminal deletion mutants of Cry1Ab and Cry1Ac which lack approximately 60 amino acids encompassing α -helix 1 on the three dimensional Cry structure are capable of assembling monomers of molecular weight about 60 kDa into pre-pores in the absence of cadherin binding. These N-terminal deletion mutants were reported to be active on Cry-resistant insect larvae. Furthermore, Diaz-Mendoza *et al.* (2007) described Cry1Ab fragments of 43 kDa and 46 kDa that retained activity on Mediterranean corn borer (*Sesamia nonagrioides*). These fragments were demonstrated to include amino acid residues 116 to 423; however the precise amino acid sequences were not elucidated and the mechanism of activity of these proteolytic fragments is unknown. The results of Gomez *et al.* (2002), Soberon *et al.* (2007), and Diaz-Mendoza *et al.* (2007) contrast with those of Hofte *et al.* (1986), who reported that deletion of 36 amino acids from the N-terminus of Cry1Ab resulted in loss of insecticidal activity.

[0034] We have deduced the beginnings and ends of α -helix 1, α -helix 2A, α -helix 2B, and α -helix 3, and the location of the spacer regions between them in Domain I of the DIG-3 toxin by comparing the DIG-3 protein sequence with the protein sequence for Cry8Ea1, for which the structure is known. These locations are described in Table 1.

Table 1

Amino acid coordinates of projected α -helices of DIG-3 protein.							
	α -helix 1	spacer	α -helix 2A	spacer	α -helix 2B	spacer	α -helix 3
Residues of SEQ NO:2	53-70	71-76	77-91	92-99	100-108	109-113	114-138

[0035] Amino terminal deletion variants of DIG-3. In one of its aspects the invention provides DIG-3 variants in which all or part of α -helix 1, α -helix 2A, and α -helix 2B are deleted to improve insecticidal activity and avoid development of resistance by insects. These modifications are made to provide DIG-3 variants with improved attributes, such as improved target pest spectrum, potency, and insect resistance management. In some embodiments of the invention, the subject modifications may affect the efficiency of protoxin activation and pore formation, leading to insect intoxication. More specifically, to provide DIG-3 variants with improved attributes, step-wise deletions are described that remove part of the nucleic acid sequence encoding the N-terminus of the DIG-3 protein. The deletions remove all of α -helix 1 and all or part of α -helix 2 in Domain I, while maintaining the structural integrity of α -helices 3 through 7. The subject invention therefore relates in part to improvements to Cry protein efficacy made by engineering the α -helical components of Domain 1 for more efficient pore formation. More specifically, the subject invention relates in part to improved DIG-3 proteins designed to have N-terminal deletions in regions with putative secondary structure homology to α -helix 1 and α -helix 2 in Domain I of Cry1 proteins.

[0036] Deletions to improve the insecticidal properties of the DIG-3 toxins may initiate before the predicted α -helix 2A start, and may terminate after the α -helix 2B end, but preferably do not extend into α -helix 3.

[0037] In designing coding sequences for the N-terminal deletion variants, an ATG start codon, encoding methionine, is inserted at the 5' end of the nucleotide sequence designed to encode the deletion variant. For sequences designed for use in transgenic plants, it may be of benefit to adhere to the "N-end rule" of Varshavsky (1997). It is taught that some amino acids may contribute to protein instability and degradation in eukaryotic cells when displayed as the N-terminal residue of a protein. For example, data collected from observations in yeast and mammalian cells indicate that the N-terminal destabilizing amino acids are F, L, W, Y, R, K, H, I, N, Q, D, E and possibly P. While the specifics of protein degradation mechanisms may differ somewhat between organisms, the conservation of identity of N-terminal destabilizing amino acids seen above suggests that similar mechanisms may function in plant cells. For instance, Worley *et al.* (1998) found that in plants, the N-end rule includes basic and aromatic residues. It is a possibility that proteolytic cleavage by plant proteases near the start of α -helix 3 of subject *B.t.* insecticidal proteins may expose a destabilizing N-terminal amino acid. Such processing may target the cleaved proteins for rapid decay and limit the accumulation of the *B.t.* insecticidal proteins to levels insufficient for effective insect control. Accordingly, for N-terminal deletion variants that begin with one of the destabilizing amino acids, applicants prefer to add a codon that specifies a G (glycine) amino acid between the translational initiation methionine and the destabilizing amino acid.

[0038] Example 2 gives specific examples of amino-terminal deletion variants of DIG-3 in accordance with the invention. Additional useful fragments that retain toxicity can be identified by trypsin or chymotrypsin digestion of the full length solubilized crystal protein. Further examples of toxic DIG-3 protein fragments may be encoded by fragments of the DIG-3 coding region. Insect active DIG-3 variants will mostly have a short N-terminal truncation and a long C-terminal truncation. The N-terminal end of the smallest toxic fragment is conveniently determined by N-terminal amino acid sequence determination of trypsin- or chymotrypsin-treated soluble crystal protein by techniques routinely available in the art.

[0039] Chimeric Toxins. Chimeric proteins utilizing the core toxin segment of one Cry toxin fused to the protoxin segment of another Cry toxin have previously been reported. DIG-3 variants include toxins comprising an N-terminal core toxin segment of a DIG-3 toxin (which may be full length or have the N-terminal deletions described above) fused to a heterologous protoxin segment at some point past the end of the core toxin segment. The transition to the heterologous protoxin segment can occur at approximately the native core toxin/protoxin junction or, in the alternative, a portion of the native protoxin (extending past the core toxin segment) can be retained with the transition to the heterologous protoxin occurring downstream. As an example, a chimeric toxin of the subject invention has the full core toxin segment of DIG-3 (amino acids 1-643) and a heterologous protoxin segment (amino acids 643 to the C-terminus). In a preferred embodiment, the heterologous protoxin segment is derived from a Cry1Ab delta-endotoxin, as illustrated in SEQ ID NO:5.

[0040] SEQ ID NO:4 discloses the 545 amino acid sequence of a Cry1Ab protoxin segment useful in DIG-3 variants of the invention. Attention is drawn to the last about 100 to 150 amino acids of this protoxin segment, which it is most critical to include in the chimeric toxin of the subject invention.

[0041] Protease sensitivity variants. Insect gut proteases typically function in aiding the insect in obtaining needed amino acids from dietary protein. The best understood insect digestive proteases are serine proteases, which appear to be the most common type (Englemann and Geraerts (1980), particularly in Lepidopteran species. Coleopteran insects

have guts that are more neutral to acidic than are Lepidopteran guts. The majority of Coleopteran larvae and adults, for example Colorado potato beetle, have slightly acidic midguts, and cysteine proteases provide the major proteolytic activity (Wolfson and Murdock, 1990). More precisely, Thie and Houseman (1990) identified and characterized the cysteine proteases, cathepsin B-like and cathepsin H-like, and the aspartyl protease, cathepsin D-like, in Colorado potato beetle. Gillikin *et al.* (1992) characterized the proteolytic activity in the guts of western corn rootworm larvae and found primarily cysteine proteases. US Patent No. 7230167 disclosed that a protease activity attributed to cathepsin G exists in western corn rootworm. The diversity and different activity levels of the insect gut proteases may influence an insect's sensitivity to a particular *B.t.* toxin.

[0042] In another embodiment of the invention, protease cleavage sites may be engineered at desired locations to affect protein processing within the midgut of susceptible larvae of certain insect pests. These protease cleavage sites may be introduced by methods such as chemical gene synthesis or splice overlap PCR (Horton *et al.*, 1989). Serine protease recognition sequences, for example, can optionally be inserted at specific sites in the Cry protein structure to effect protein processing at desired deletion points within the midgut of susceptible larvae. Serine proteases that can be exploited in such fashion include Lepidopteran midgut serine proteases such as trypsin or trypsin-like enzymes, chymotrypsin, elastase, *etc.* (Christeller *et al.*, 1992). Further, deletion sites identified empirically by sequencing Cry protein digestion products generated with unfractionated larval midgut protease preparations or by binding to brush border membrane vesicles can be engineered to effect protein activation. Modified Cry proteins generated either by gene deletion or by introduction of protease cleavage sites have improved activity on Lepidopteran pests including *Ostrinia nubilalis*, *Diatraea grandiosella*, *Helicoverpa zea*, *Agrotis ipsilon*, *Spodoptera frugiperda*, *Spodoptera exigua*, *Diatraea saccharalis*, *Loxagrotis albicosta*, and other target pests.

[0043] Coleopteran serine proteases such as trypsin, chymotrypsin and cathepsin G-like protease, Coleopteran cysteine proteases such as cathepsins (B-like, L-like, O-like, and K-like proteases) (Koiwa *et al.*, (2000) and Bown *et al.*, (2004)), Coleopteran metalloproteases such as ADAM10 [Ochoa-Campuzano *et al.*, (2007)], and Coleopteran aspartic acid proteases such as cathepsins D-like and E-like, pepsin, plasmepsin, and chymosin may further be exploited by engineering appropriate recognition sequences at desired processing sites to affect Cry protein processing within the midgut of susceptible larvae of certain insect pests.

[0044] A preferred location for the introduction of such protease cleavage sites may be within the "spacer" region between α -helix 2B and α -helix 3, for example within amino acids 109 to 113 of the full length DIG-3 protein (SEQ ID NO:2 and Table 1). Modified Cry proteins generated either by gene deletion or by introduction of protease cleavage sites have improved activity on insect pests including but not limited to western corn rootworm, southern corn root worm, northern corn rootworm, and the like.

[0045] Various technologies exist to enable determination of the sequence of the amino acids which comprise the N-terminal or C-terminal residues of polypeptides. For example, automated Edman degradation methodology can be used in sequential fashion to determine the N-terminal amino acid sequence of up to 30 amino acid residues with 98% accuracy per residue. Further, determination of the sequence of the amino acids comprising the carboxy end of polypeptides is also possible [Bailey *et al.*, (1992); US Patent No. 6046053]. Thus, in some embodiments, *B.t.* Cry proteins which have been activated by means of proteolytic processing, for example, by proteases prepared from the gut of an insect, may be characterized and the N-terminal or C-terminal amino acids of the activated toxin fragment identified. DIG-3 variants produced by introduction or elimination of protease processing sites at appropriate positions in the coding sequence to allow, or eliminate, proteolytic cleavage of a larger variant protein by insect, plant or microorganism proteases are within the scope of the invention. The end result of such manipulation is understood to be the generation of toxin fragment molecules having the same or better activity as the intact (full length) toxin protein.

[0046] Domains of the DIG-3 toxin. The separate domains of the DIG-3 toxin, (and variants that are 90%, 95%, or 97% identical to such domains) are expected to be useful in forming combinations with domains from other Cry toxins to provide new toxins with increased spectrum of pest toxicity, improved potency, or increased protein stability. Domain I of the DIG-3 protein consists of amino acid residues 56 to 278 of SEQ ID NO:2. Domain II of the DIG-3 protein consists of amino acid residues 283 to 493 of SEQ ID NO:2. Domain III of the DIG-3 protein consists of amino acid residues 503 to 641 of SEQ ID NO:2. Domain swapping or shuffling is a mechanism for generating altered delta-endotoxin proteins. Domains II and III may be swapped between delta-endotoxin proteins, resulting in hybrid or chimeric toxins with improved pesticidal activity or target spectrum. Domain II is involved in receptor binding, and the DIG-3 Domain II is very divergent from other CryIB toxins. Domain III binds certain classes of receptor proteins and perhaps participates in insertion of an oligomeric toxin pre-pore. Some Domain III substitutions in other toxins have been shown to produce superior toxicity against *Spodoptera exigua* (de Maagd *et al.*, 1996), and guidance exists on the design of the Cry toxin domain swaps (Knight *et al.*, 2004).

[0047] Methods for generating recombinant proteins and testing them for pesticidal activity are well known in the art (see, for example, Naimov *et al.*, (2001), de Maagd *et al.*, (1996), Ge *et al.*, (1991), Schnepf *et al.*, (1990), Rang *et al.*, (1999)). Domain I from Cry1A and Cry3A proteins has been studied for the ability to insert and form pores in membranes. α -helix 4 and α -helix 5 of Domain I play key roles in membrane insertion and pore formation [Walters *et al.*, (1993), Gazit

et al., (1998); Nunez-Valdez *et al.*, (2001)], with the other alpha helices proposed to contact the membrane surface like the ribs of an umbrella (Bravo *et al.*, (2007); Gazit *et al.*, (1998)).

[0048] DIG-3 variants created by making a limited number of amino acid deletions, substitutions, or additions. Amino acid deletions, substitutions, and additions to the amino acid sequence of SEQ ID NO:2 can readily be made in a sequential manner and the effects of such variations on insecticidal activity can be tested by bioassay. Provided the number of changes is limited in number, such testing does not involve unreasonable experimentation. The invention includes insecticidally active variants of the core toxin segment (amino acids 1-643 of SEQ ID NO:2, or amino acids 73-643 of SEQ ID NO:2) in which up to 10, up to 15, or up to 20 independent amino acid additions, deletions, or substitutions have been made.

[0049] The invention includes DIG-3 variants having a core toxin segment that is 90%, 95% or 97% identical to amino acids 1-643 of SEQ ID NO:2 or amino acids 73-643 of SEQ ID NO:2.

[0050] Variants may be made by making random mutations or the variants may be designed. In the case of designed mutants, there is a high probability of generating variants with similar activity to the native toxin when amino acid identity is maintained in critical regions of the toxin which account for biological activity or are involved in the determination of three-dimensional configuration which ultimately is responsible for the biological activity. A high probability of retaining activity will also occur if substitutions are conservative. Amino acids may be placed in the following classes: non-polar, uncharged polar, basic, and acidic. Conservative substitutions whereby an amino acid of one class is replaced with another amino acid of the same type are least likely to materially alter the biological activity of the variant. Table 2 provides a listing of examples of amino acids belonging to each class.

Table 2

Class of Amino Acid	Examples of Amino Acids
Nonpolar Side Chains	Ala (A), Val (V), Leu (L), Ile (I), Pro (P), Met (M), Phe (F), Trp (W)
Uncharged Polar Side Chains	Gly (G), Ser (S), Thr (T), Cys (C), Tyr (Y), Asn (N), Gln (Q)
Acidic Side Chains	Asp (D), Glu (E)
Basic Side Chains	Lys (K), Arg (R), His (H)
Beta-branched Side Chains	Thr, Val, Ile
Aromatic Side Chains	Tyr, Phe, Trp, His

[0051] In some instances, non-conservative substitutions can also be made. The critical factor is that these substitutions must not significantly detract from the biological activity of the toxin. Variants include polypeptides that differ in amino acid sequence due to mutagenesis. Variant proteins encompassed by the present invention are biologically active, that is they continue to possess the desired biological activity of the native protein, namely, retaining pesticidal activity.

[0052] Variant proteins can also be designed that differ at the sequence level but that retain the same or similar overall essential three-dimensional structure, surface charge distribution, and the like. See e.g. U.S. Patent No. 7058515; Larson *et al.* (2002); Stemmer (1994a, 1994b, 1995); and Crameri *et al.* (1996a, 1996b, 1997).

[0053] Nucleic Acids. Isolated nucleic acids encoding DIG-3 toxins are one aspect of the present invention. This includes nucleic acids encoding SEQ ID NO:2 and SEQ ID NO:5, and complements thereof, as well as other nucleic acids that encode insecticidal variants of SEQ ID NO:2. Because of the redundancy of the genetic code, a variety of different DNA sequences can encode the amino acid sequences disclosed herein. It is well within the skill of a person trained in the art to create these alternative DNA sequences encoding the same, or essentially the same, toxins.

[0054] Gene synthesis. DNA sequences encoding the improved Cry proteins described herein can be made by a variety of methods well-known in the art. For example, synthetic gene segments and synthetic genes can be made by phosphite tri-ester and phosphoramidite chemistry (Caruthers *et al.*, 1987), and commercial vendors are available to perform DNA synthesis on demand. Sequences encoding full-length DIG-3 proteins can be assembled in a variety of ways including, for example, by ligation of restriction fragments or polymerase chain reaction assembly of overlapping oligonucleotides (Stewart and Burgin, 2005). Further, sequences encoding terminal deletions can be made by PCR amplification using site-specific terminal oligonucleotides.

[0055] Nucleic acids encoding DIG-3 toxins can be made for example, by synthetic construction by methods currently practiced by any of several commercial suppliers. (See for example, US Patent No. 7482119 B2). These nucleic acids, or portions or variants thereof, may also be constructed synthetically, for example, by use of a gene synthesizer and the design methods of, for example, US Patent No. 5380831. Alternatively, variations of synthetic or naturally occurring genes may be readily constructed using standard molecular biological techniques for making point mutations. Fragments of these genes can also be made using commercially available exonucleases or endonucleases according to standard

procedures. For example, enzymes such as *Bal*/31 or site-directed mutagenesis can be used to systematically cut off nucleotides from the ends of these genes. Also, gene fragments which encode active toxin fragments may be obtained using a variety of restriction enzymes.

[0056] Given the amino acid sequence for a DIG-3 toxin, a coding sequence can be designed by reverse translating the coding sequence using codons preferred by the intended host, and then refining the sequence using alternative codons to remove sequences that might cause problems and provide periodic stop codons to eliminate long open coding sequences in the non-coding reading frames,

[0057] Quantifying Sequence Identity. To determine the percent identity of two amino acid sequences or of two nucleic acid sequences, the sequences are aligned for optimal comparison purposes. The percent identity between the two sequences is a function of the number of identical positions shared by the sequences (*i.e.* percent identity = number of identical positions/total number of positions (*e.g.* overlapping positions) x100). In one embodiment, the two sequences are the same length. The percent identity between two sequences can be determined using techniques similar to those described below, with or without allowing gaps. In calculating percent identity, typically exact matches are counted.

[0058] The determination of percent identity between two sequences can be accomplished using a mathematical algorithm. A nonlimiting example of such an algorithm is that of Karlin and Altschul (1990), modified as in Karlin and Altschul (1993), and incorporated into the BLASTN and BLASTX programs. BLAST searches may be conveniently used to identify sequences homologous (similar) to a query sequence in nucleic or protein databases. BLASTN searches can be performed, (score = 100, word length = 12) to identify nucleotide sequences having homology to claimed nucleic acid molecules of the invention. BLASTX searches can be performed (score = 50, word length = 3) to identify amino acid sequences having homology to claimed insecticidal protein molecules of the invention.

[0059] Gapped BLAST (Altschul *et al.*, 1997) can be utilized to obtain gapped alignments for comparison purposes. Alternatively, PSI-Blast can be used to perform an iterated search that detects distant relationships between molecules (Altschul *et al.*, 1997). When utilizing BLAST, Gapped BLAST, and PSI-Blast programs, the default parameters of the respective programs can be used. See www.ncbi.nlm.nih.gov.

[0060] A non-limiting example of a mathematical algorithm utilized for the comparison of sequences is the ClustalW algorithm (Thompson *et al.*, 1994). ClustalW compares sequences and aligns the entirety of the amino acid or DNA sequence, and thus can provide data about the sequence conservation of the entire amino acid sequence or nucleotide sequence. The ClustalW algorithm is used in several commercially available DNA/amino acid analysis software packages, such as the ALIGNX module of the Vector NTI Program Suite (Invitrogen, Inc., Carlsbad, CA). When aligning amino acid sequences with ALIGNX, one may conveniently use the default settings with a Gap open penalty of 10, a Gap extend penalty of 0.1 and the blosum63mt2 comparison matrix to assess the percent amino acid similarity (consensus) or identity between the two sequences. When aligning DNA sequences with ALIGNX, one may conveniently use the default settings with a Gap open penalty of 15, a Gap extend penalty of 6.6 and the swgapdnamt comparison matrix to assess the percent identity between the two sequences.

[0061] Another non-limiting example of a mathematical algorithm utilized for the comparison of sequences is that of Myers and Miller (1988). Such an algorithm is incorporated into the wSTRETCHER program, which is part of the wEMB-BOSS sequence alignment software package (available at <http://emboss.sourceforge.net/>). wSTRETCHER calculates an optimal global alignment of two sequences using a modification of the classic dynamic programming algorithm which uses linear space. The substitution matrix, gap insertion penalty and gap extension penalties used to calculate the alignment may be specified. When utilizing the wSTRETCHER program for comparing nucleotide sequences, a Gap open penalty of 16 and a Gap extend penalty of 4 can be used with the scoring matrix file EDNAFULL. When used for comparing amino acid sequences, a Gap open penalty of 12 and a Gap extend penalty of 2 can be used with the EBLOSUM62 scoring matrix file.

[0062] A further non-limiting example of a mathematical algorithm utilized for the comparison of sequences is that of Needleman and Wunsch (1970), which is incorporated in the sequence alignment software packages GAP Version 10 and wNEEDLE (<http://emboss.sourceforge.net/>). GAP Version 10 may be used to determine sequence identity or similarity using the following parameters: for a nucleotide sequence, % identity and % similarity are found using GAP Weight of 50 and Length Weight of 3, and the nwsgapdna. cmp scoring matrix. For amino acid sequence comparison, % identity or % similarity are determined using GAP weight of 8 and length weight of 2, and the BLOSUM62 scoring program.

[0063] wNEEDLE reads two input sequences, finds the optimum alignment (including gaps) along their entire length, and writes their optimal global sequence alignment to file. The algorithm explores all possible alignments and chooses the best, using a scoring matrix that contains values for every possible residue or nucleotide match. wNEEDLE finds the alignment with the maximum possible score, where the score of an alignment is equal to the sum of the matches taken from the scoring matrix, minus penalties arising from opening and extending gaps in the aligned sequences. The substitution matrix and gap opening and extension penalties are user-specified. When amino acid sequences are compared, a default Gap open penalty of 10, a Gap extend penalty of 0.5, and the EBLOSUM62 comparison matrix are used. When DNA sequences are compared using wNEEDLE, a Gap open penalty of 10, a Gap extend penalty of 0.5, and the EDNAFULL comparison matrix are used.

[0064] Equivalent programs may also be used. By "equivalent program" is intended any sequence comparison program that, for any two sequences in question, generates an alignment having identical nucleotide or amino acid residue matches and an identical percent sequence identity when compared to the corresponding alignment generated by ALIGNX, wNEEDLE, or wSTRETCHER. The % identity is the percentage of identical matches between the two sequences over the reported aligned region (including any gaps in the length) and the % similarity is the percentage of matches between the two sequences over the reported aligned region (including any gaps in the length).

[0065] Alignment may also be performed manually by inspection.

[0066] Recombinant hosts. The toxin-encoding genes of the subject invention can be introduced into a wide variety of microbial or plant hosts. Expression of the toxin gene results, directly or indirectly, in the intracellular production and maintenance of the pesticidal protein. With suitable microbial hosts, e.g. *Pseudomonas*, the microbes can be applied to the environment of the pest, where they will proliferate and be ingested. The result is a control of the pest. Alternatively, the microbe hosting the toxin gene can be treated under conditions that prolong the activity of the toxin and stabilize the cell. The treated cell, which retains the toxic activity, then can be applied to the environment of the target pest.

[0067] Where the *B.t.* toxin gene is introduced via a suitable vector into a microbial host, and said host is applied to the environment in a living state, it is essential that certain host microbes be used. Microorganism hosts are selected which are known to occupy the "phytosphere" (phylloplane, phyllosphere, rhizosphere, and/or rhizoplane) of one or more crops of interest. These microorganisms are selected so as to be capable of successfully competing in the particular environment (crop and other insect habitats) with the wild-type indigenous microorganisms, provide for stable maintenance and expression of the gene expressing the polypeptide pesticide, and, desirably, provide for improved protection of the pesticide from environmental degradation and inactivation.

[0068] A large number of microorganisms are known to inhabit the phylloplane (the surface of the plant leaves) and/or the rhizosphere (the soil surrounding plant roots) of a wide variety of important crops. These microorganisms include bacteria, algae, and fungi. Of particular interest are microorganisms, such as bacteria, e.g. genera *Pseudomonas*, *Erwinia*, *Serratia*, *Klebsiella*, *Xanthomonas*, *Streptomyces*, *Rhizobium*, *Sinorhizobium*, *Rhodopseudomonas*, *Methylophilus*, *Agrobacterium*, *Acetobacter*, *Lactobacillus*, *Arthrobacter*, *Azotobacter*, *Leuconostoc*, and *Alcaligenes*; fungi, particularly yeast, e.g. genera *Saccharomyces*, *Cryptococcus*, *Kluyveromyces*, *Sporobolomyces*, *Rhodotorula*, and *Aureobasidium*. Of particular interest are such phytosphere bacterial species as *Pseudomonas syringae*, *Pseudomonas fluorescens*, *Serratia marcescens*, *Acetobacter xylinum*, *Agrobacterium tumefaciens*, *Agrobacterium radiobacter*, *Rhodopseudomonas spheroides*, *Xanthomonas campestris*, *Sinorhizobium meliloti* (formerly *Rhizobium meliloti*), *Alcaligenes eutrophus*, and *Azotobacter vinelandii* and phytosphere yeast species such as *Rhodotorula rubra*, *R. glutinis*, *R. marina*, *R. aurantiaca*, *Cryptococcus albidus*, *C. diffluens*, *C. laurentii*, *Saccharomyces rosei*, *S. pretoriensis*, *S. cerevisiae*, *Sporobolomyces roseus*, *S. odoratus*, *Kluyveromyces veronae*, and *Aureobasidium pollulans*. Of particular interest are the pigmented microorganisms.

Methods of controlling insect pests

[0069] When an insect comes into contact with an effective amount of toxin delivered via transgenic plant expression, formulated protein compositions(s), sprayable protein composition(s), a bait matrix or other delivery system, the results are typically death of the insect, or the insects do not feed upon the source which makes the toxins available to the insects.

[0070] The subject protein toxins can be "applied" or provided to contact the target insects in a variety of ways. For example, transgenic plants (wherein the protein is produced by and present in the plant) can be used and are well-known in the art. Expression of the toxin genes can also be achieved selectively in specific tissues of the plants, such as the roots, leaves, etc. This can be accomplished via the use of tissue-specific promoters, for example. Spray-on applications are another example and are also known in the art. The subject proteins can be appropriately formulated for the desired end use, and then sprayed (or otherwise applied) onto the plant and/or around the plant / to the vicinity of the plant to be protected - before an infestation is discovered, after target insects are discovered, both before and after, and the like. Bait granules, for example, can also be used and are known in the art.

Transgenic plants

[0071] The subject proteins can be used to protect practically any type of plant from damage by a Lepidopteran insect. Nonlimiting examples of such plants include maize, sunflower, soybean, cotton, canola, rice, sorghum, wheat, barley, vegetables, ornamentals, peppers (including hot peppers), sugar beets, fruit, and turf, to name but a few. Methods for transforming plants are well known in the art, and illustrative transformation methods are described in the Examples.

[0072] A preferred embodiment of the subject invention is the transformation of plants with genes encoding the subject insecticidal protein or its variants. The transformed plants are resistant to attack by an insect target pest by virtue of the presence of controlling amounts of the subject insecticidal protein or its variants in the cells of the transformed plant. By incorporating genetic material that encodes the insecticidal properties of the *B.t.* insecticidal toxins into the genome of

a plant eaten by a particular insect pest, the adult or larvae would die after consuming the food plant. Numerous members of the monocotyledonous and dicotyledonous classifications have been transformed. Transgenic agronomic crops as well as fruits and vegetables are of commercial interest. Such crops include, but are not limited to, maize, rice, soybeans, canola, sunflower, alfalfa, sorghum, wheat, cotton, peanuts, tomatoes, potatoes, and the like. Several techniques exist for introducing foreign genetic material into plant cells, and for obtaining plants that stably maintain and express the introduced gene. Such techniques include acceleration of genetic material coated onto microparticles directly into cells (US Patent No. 4945050 and US Patent No. 5141131). Plants may be transformed using *Agrobacterium* technology, see US Patent No. 5177010, US Patent No. 5104310, European Patent Application No. 0131624B1, European Patent Application No. 120516, European Patent Application No. 159418B1, European Patent Application No. 176112, US Patent No. 5149645, US Patent No. 5469976, US Patent No. 5464763, US Patent No. 4940838, US Patent No. 4693976, European Patent Application No. 116718, European Patent Application No. 290799, European Patent Application No. 320500, European Patent Application No. 604662, European Patent Application No. 627752, European Patent Application No. 0267159, European Patent Application No. 0292435, US Patent No. 5231019, US Patent No. 5463174, US Patent No. 4762785, US Patent No. 5004863, and US Patent No. 5159135. Other transformation technology includes WHISKERS™ technology, see US Patent No. 5302523 and US Patent No. 5464765. Electroporation technology has also been used to transform plants, see WO 87/06614, US Patent No. 5472869, US Patent No. 5384253, WO 9209696, and WO 9321335. In addition to numerous technologies for transforming plants, the type of tissue which is contacted with the foreign genes may vary as well. Such tissue would include but would not be limited to embryogenic tissue, callus tissue types I and II, hypocotyl, meristem, and the like. Almost all plant tissues may be transformed during dedifferentiation using appropriate techniques within the skill of an artisan.

[0073] Genes encoding DIG-3 toxins can be inserted into plant cells using a variety of techniques which are well known in the art as disclosed above. For example, a large number of cloning vectors comprising a marker that permits selection of the transformed microbial cells and a replication system functional in *Escherichia coli* are available for preparation and modification of foreign genes for insertion into higher plants. Such manipulations may include, for example, the insertion of mutations, truncations, additions, or substitutions as desired for the intended use. The vectors comprise, for example, pBR322, pUC series, M13mp series, pACYC184, etc. Accordingly, the sequence encoding the Cry protein or variants can be inserted into the vector at a suitable restriction site. The resulting plasmid is used for transformation of cells of *E. coli*, the cells of which are cultivated in a suitable nutrient medium, then harvested and lysed so that workable quantities of the plasmid are recovered. Sequence analysis, restriction fragment analysis, electrophoresis, and other biochemical-molecular biological methods are generally carried out as methods of analysis. After each manipulation, the DNA sequence used can be cleaved and joined to the next DNA sequence. Each manipulated DNA sequence can be cloned in the same or other plasmids.

[0074] The use of T-DNA-containing vectors for the transformation of plant cells has been intensively researched and sufficiently described in EP 120516; Lee and Gelvin (2008), Fraley *et al.* (1986), and An *et al.* (1985), and is well established in the field.

[0075] Once the inserted DNA has been integrated into the plant genome, it is relatively stable throughout subsequent generations. The vector used to transform the plant cell normally contains a selectable marker gene encoding a protein that confers on the transformed plant cells resistance to a herbicide or an antibiotic, such as bialaphos, kanamycin, G418, bleomycin, or hygromycin, *inter alia*. The individually employed selectable marker gene should accordingly permit the selection of transformed cells while the growth of cells that do not contain the inserted DNA is suppressed by the selective compound.

[0076] A large number of techniques are available for inserting DNA into a host plant cell. Those techniques include transformation with T-DNA delivered by *Agrobacterium tumefaciens* or *Agrobacterium rhizogenes* as the transformation agent. Additionally, fusion of plant protoplasts with liposomes containing the DNA to be delivered, direct injection of the DNA, biolistics transformation (microparticle bombardment), or electroporation, as well as other possible methods, may be employed.

[0077] In a preferred embodiment of the subject invention, plants will be transformed with genes wherein the codon usage of the protein coding region has been optimized for plants. See, for example, US Patent No. 5380831. Also, advantageously, plants encoding a truncated toxin will be used. The truncated toxin typically will encode about 55% to about 80% of the full length toxin. Methods for creating synthetic *B.t.* genes for use in plants are known in the art (Stewart, 2007).

[0078] Regardless of transformation technique, the gene is preferably incorporated into a gene transfer vector adapted to express the *B.t.* insecticidal toxin genes and variants in the plant cell by including in the vector a plant promoter. In addition to plant promoters, promoters from a variety of sources can be used efficiently in plant cells to express foreign genes. For example, one may use promoters of bacterial origin, such as the octopine synthase promoter, the nopaline synthase promoter, and the mannopine synthase promoter. Promoters of plant virus origin may be used, for example, the 35S and 19S promoters of Cauliflower Mosaic Virus, a promoter from Cassava Vein Mosaic Virus, and the like. Plant promoters include, but are not limited to, ribulose-1,6-bisphosphate (RUBP) carboxylase small subunit (ssu), beta-

conglycinin promoter, phaseolin promoter, ADH (alcohol dehydrogenase) promoter, heat-shock promoters, ADF (actin depolymerization factor) promoter, ubiquitin promoter, actin promoter, and tissue specific promoters. Promoters may also contain certain enhancer sequence elements that may improve the transcription efficiency. Typical enhancers include but are not limited to ADH1-intron 1 and ADH1-intron 6. Constitutive promoters may be used. Constitutive promoters direct continuous gene expression in nearly all cells types and at nearly all times (e.g., actin, ubiquitin, CaMV 35S). Tissue specific promoters are responsible for gene expression in specific cell or tissue types, such as the leaves or seeds (e.g., zein, oleosin, napin, ACP (Acyl Carrier Protein) promoters), and these promoters may also be used. Promoters may also be used that are active during a certain stage of the plants' development as well as active in specific plant tissues and organs. Examples of such promoters include but are not limited to promoters that are root specific, pollen-specific, embryo specific, corn silk specific, cotton fiber specific, seed endosperm specific, phloem specific, and the like.

[0079] Under certain circumstances it may be desirable to use an inducible promoter. An inducible promoter is responsible for expression of genes in response to a specific signal, such as: physical stimulus (e.g. heat shock genes); light (e.g. RUBP carboxylase); hormone (e.g. glucocorticoid); antibiotic (e.g. tetracycline); metabolites; and stress (e.g. drought). Other desirable transcription and translation elements that function in plants may be used, such as 5' untranslated leader sequences, RNA transcription termination sequences and poly-adenylate addition signal sequences. Numerous plant-specific gene transfer vectors are known to the art.

[0080] Transgenic crops containing insect resistance (IR) traits are prevalent in corn and cotton plants throughout North America, and usage of these traits is expanding globally. Commercial transgenic crops combining IR and herbicide tolerance (HT) traits have been developed by multiple seed companies. These include combinations of IR traits conferred by *B.t.* insecticidal proteins and HT traits such as tolerance to Acetolactate Synthase (ALS) inhibitors such as sulfonylureas, imidazolinones, triazolopyrimidine, sulfonanilides, and the like, Glutamine Synthetase (GS) inhibitors such as bialaphos, glufosinate, and the like, 4-HydroxyPhenylPyruvate Dioxygenase (HPPD) inhibitors such as mesotrione, isoxaflutole, and the like, 5-EnolPyruvylShikimate-3-Phosphate Synthase (EPSPS) inhibitors such as glyphosate and the like, and Acetyl-Coenzyme A Carboxylase (ACCase) inhibitors such as haloxyfop, quizalofop, diclofop, and the like. Other examples are known in which transgenically provided proteins provide plant tolerance to herbicide chemical classes such as phenoxy acids herbicides and pyridyloxyacetates auxin herbicides (see WO 2007/053482 A2), or phenoxy acids herbicides and aryloxyphenoxypropionates herbicides (see WO 2005107437 A2, A3). The ability to control multiple pest problems through IR traits is a valuable commercial product concept, and the convenience of this product concept is enhanced if insect control traits and weed control traits are combined in the same plant. Further, improved value may be obtained via single plant combinations of IR traits conferred by a *B.t.* insecticidal protein such as that of the subject invention, with one or more additional HT traits such as those mentioned above, plus one or more additional input traits (e.g. other insect resistance conferred by *B.t.*-derived or other insecticidal proteins, insect resistance conferred by mechanisms such as RNAi and the like, nematode resistance, disease resistance, stress tolerance, improved nitrogen utilization, and the like), or output traits (e.g. high oils content, healthy oil composition, nutritional improvement, and the like). Such combinations may be obtained either through conventional breeding (breeding stack) or jointly as a novel transformation event involving the simultaneous introduction of multiple genes (molecular stack). Benefits include the ability to manage insect pests and improved weed control in a crop plant that provides secondary benefits to the producer and/or the consumer. Thus, the subject invention can be used in combination with other traits to provide a complete agronomic package of improved crop quality with the ability to flexibly and cost effectively control any number of agronomic issues.

Target Pests

[0081] The DIG-3 toxins of the invention are particularly suitable for use in control of Lepidopteran insects. Lepidopterans are an important group of agricultural, horticultural, and household pests which cause a very large amount of damage each year. This insect order encompasses foliar- and root-feeding larvae and adults. Lepidopteran insect pests include, but are not limited to: *Achoria grisella*, *Acleris gloverana*, *Acleris variana*, *Adoxophyes orana*, *Agrotis ipsilon* (black cutworm), *Alabama argillacea*, *Alsophila pometaria*, *Amyelois transitella*, *Anagasta kuehniella*, *Anarsia lineatella*, *Anisota sanatoria*, *Antheraea pernyi*, *Anticarsia gemmatilis*, *Archips* sp., *Argyrotaenia* sp., *Athetis mindara*, *Bombyx mori*, *Bucculatrix thurberiella*, *Cadra cautella*, *Choristoneura* sp., *Cochylis hopes*, *Colias eurytheme*, *Corcyra cephalonica*, *Cydia latiferreanus*, *Cydia pomonella*, *Datana integerrima*, *Dendrolimus sibericus*, *Desmia fernalis*, *Diaphania hyalinata*, *Diaphania nitidalis*, *Diatraea grandiosella* (southwestern corn borer), *Diatraea saccharalis* (sugarcane borer), *Ennomos subsignaria*, *Eoreuma loftini*, *Esphestia elutella*, *Erannis tilaria*, *Estigmene acrea*, *Eulia salubricola*, *Eupocoellia ambigua*, *Eupoecilia ambigua*, *Euproctis chrysorrhoea*, *Euxoa messoria*, *Galleria mellonella*, *Grapholita molesta*, *Harrisina americana*, *Helicoverpa subflexa*, *Helicoverpa zea* (corn earworm), *Heliothis virescens* (tobacco budworm), *Hemileuca oliviae*, *Homoeosoma electellum* (sunflower head moth), *Hyphantia cunea*, *Keiferia lycopersicella*, *Lambdina fiscellaria fiscellaria*, *Lambdina fiscellaria lugubrosa*, *Leucoma salicis*, *Lobesia botrana*, *Loxagrotis albicosta* (western

bean cutworm), *Loxostege sticticalis*, *Lymantria dispar*, *Macalla thyrsalis*, *Malacosoma* sp., *Mamestra brassicae*, *Mamestra configurata* (bertha armyworm), *Manduca quinquemaculata*, *Manduca sexta* (tobacco hornworm), *Maruca testulalis*, *Melanchra picta*, *Operophtera brumata*, *Orgyia* sp., *Ostrinia nubilalis* (European corn borer), *Paleacrita vernata*, *Papiapema nebris* (common stalk borer), *Papilio cresphontes*, *Pectinophora gossypiella*, *Phryganidia californica*, *Phyl-
lonorycter blancardella*, *Pieris napi*, *Pieris rapae*, *Plathypena scabra*, *Platyhota flouendana*, *Platynota stultana*, *Platyptilia
carduidactyla*, *Plodia interpunctella*, *Plutella xylostella* (diamondback moth), *Pontia protodice*, *Pseudaletia unipuncta*
(armyworm), *Pseudoplasia includens*, *Rachiplusia nu* (Argentine looper), *Sabulodes aegrotata*, *Schizura concinna*, *Si-
totroga cerealella*, *Spilonta ocellana*, *Spodoptera frugiperda* (fall armyworm), *Spodoptera exigua* (beet armyworm),
Thaurnstopoea pityocampa, *Ensola bisselliella*, *Trichoplusia hi*, *Udea rubigalis*, *Xylomyges curialis*, and *Yponomeuta
padella*.

[0082] Use of DIG-3 toxins to control Coleopteran pests of crop plants is also contemplated. In some embodiments, Cry proteins may be economically deployed for control of insect pests that include but are not limited to, for example, rootworms such as *Diabrotica undecimpunctata howardi* (southern corn rootworm), *Diabrotica longicornis barberi* (north-
ern corn rootworm), and *Diabrotica virgifera* (western corn rootworm), and grubs such as the larvae of *Cyclocephala
borealis* (northern masked chafer), *Cyclocephala immaculate* (southern masked chafer), and *Popillia japonica* (Japanese
beetle).

[0083] Use of the DIG-3 toxins to control parasitic nematodes including, but not limited to, root knot nematode (*Mel-
oidogyne icognita*) and soybean cyst nematode (*Heterodera glycines*) is also contemplated.

Antibody detection of DIG-3 toxins

[0084] Anti-toxin antibodies. Antibodies to the toxins disclosed herein, or to equivalent toxins, or fragments of these
toxins, can readily be prepared using standard procedures in this art. Such antibodies are useful to detect the presence
of the DIG-3 toxins.

[0085] Once the *B.t.* insecticidal toxin has been isolated, antibodies specific for the toxin may be raised by conventional
methods that are well known in the art. Repeated injections into a host of choice over a period of weeks or months will
elicit an immune response and result in significant anti-*B.t.* toxin serum titers. Preferred hosts are mammalian species
and more highly preferred species are rabbits, goats, sheep and mice. Blood drawn from such immunized animals may
be processed by established methods to obtain antiserum (polyclonal antibodies) reactive with the *B.t.* insecticidal toxin.
The antiserum may then be affinity purified by adsorption to the toxin according to techniques known in the art. Affinity
purified antiserum may be further purified by isolating the immunoglobulin fraction within the antiserum using procedures
known in the art. The resulting material will be a heterogeneous population of immunoglobulins reactive with the *B.t.*
insecticidal toxin.

[0086] Anti-*B.t.* toxin antibodies may also be generated by preparing a semi-synthetic immunogen consisting of a
synthetic peptide fragment of the *B.t.* insecticidal toxin conjugated to an immunogenic carrier. Numerous schemes and
instruments useful for making peptide fragments are well known in the art. Many suitable immunogenic carriers such as
bovine serum albumin or keyhole limpet hemocyanin are also well known in the art, as are techniques for coupling the
immunogen and carrier proteins. Once the semi-synthetic immunogen has been constructed, the procedure for making
antibodies specific for the *B.t. insecticidal* toxin fragment is identical to those used for making antibodies reactive with
natural *B.t.* toxin.

[0087] Anti-*B.t.* toxin monoclonal antibodies (MAbs) are readily prepared using purified *B.t.* insecticidal toxin. Methods
for producing MAbs have been practiced for over 20 years and are well known to those of ordinary skill in the art.
Repeated intraperitoneal or subcutaneous injections of purified *B.t.* insecticidal toxin in adjuvant will elicit an immune
response in most animals. Hyperimmunized B-lymphocytes are removed from the animal and fused with a suitable
fusion partner cell line capable of being cultured indefinitely. Preferred animals whose B-lymphocytes may be hyperim-
munized and used in the production of MAbs are mammals. More preferred animals are rats and mice and most preferred
is the BALB/c mouse strain.

[0088] Numerous mammalian cell lines are suitable fusion partners for the production of hybridomas. Many such lines
are available from the American Type Culture Collection (ATCC, Manassas, VA) and commercial suppliers. Preferred
fusion partner cell lines are derived from mouse myelomas and the HL-1® Friendly myeloma-653 cell line (Ventrex,
Portland, ME) is most preferred. Once fused, the resulting hybridomas are cultured in a selective growth medium for
one to two weeks. Two well known selection systems are available for eliminating unfused myeloma cells, or fusions
between myeloma cells, from the mixed hybridoma culture. The choice of selection system depends on the strain of
mouse immunized and myeloma fusion partner used. The AAT selection system, described by Taggart and Samloff,
(1983), may be used; however, the HAT (hypoxanthine, aminopterin, thymidine) selection system, described by Littlefield,
(1964), is preferred because of its compatibility with the preferred mouse strain and fusion partner mentioned above.
Spent growth medium is screened for immunospecific MAb secretion. Enzyme linked immunosorbent assay (ELISA)
procedures are best suited for this purpose; though, radioimmunoassays adapted for large volume screening are also

acceptable. Multiple screens designed to consecutively pare down the considerable number of irrelevant or less desired cultures may be performed. Cultures that secrete MABs reactive with the *B.t.* insecticidal toxin may be screened for cross-reactivity with known *B.t.* insecticidal toxins. MABs that preferentially bind to the preferred *B.t.* insecticidal toxin may be isotyped using commercially available assays. Preferred MABs are of the IgG class, and more highly preferred MABs are of the IgG₁ and IgG_{2a} subisotypes.

[0089] Hybridoma cultures that secrete the preferred MABs may be sub-cloned several times to establish monoclonality and stability. Well known methods for sub-cloning eukaryotic, non-adherent cell cultures include limiting dilution, soft agarose and fluorescence activated cell sorting techniques. After each subcloning, the resultant cultures preferably are re-assayed for antibody secretion and isotype to ensure that a stable preferred MAB-secreting culture has been established.

[0090] The anti-*B.t.* toxin antibodies are useful in various methods of detecting the claimed *B.t.* insecticidal toxin of the instant invention, and variants or fragments thereof. It is well known that antibodies labeled with a reporting group can be used to identify the presence of antigens in a variety of milieus. Antibodies labeled with radioisotopes have been used for decades in radioimmunoassays to identify, with great precision and sensitivity, the presence of antigens in a variety of biological fluids. More recently, enzyme labeled antibodies have been used as a substitute for radiolabeled antibodies in the ELISA assay. Further, antibodies immunoreactive to the *B.t.* insecticidal toxin of the present invention can be bound to an immobilizing substance such as a polystyrene well or particle and used in immunoassays to determine whether the *B.t.* toxin is present in a test sample.

Detection using nucleic acid probes

[0091] A further method for identifying the toxins and genes of the subject invention is through the use of oligonucleotide probes. These probes are detectable nucleotide sequences. These sequences may be rendered detectable by virtue of an appropriate radioactive label or may be made inherently fluorescent as described in, for example, US Patent No. 6268132. As is well known in the art, if the probe molecule and nucleic acid sample hybridize by forming strong base-pairing bonds between the two molecules, it can be reasonably assumed that the probe and sample have substantial sequence homology. Preferably, hybridization is conducted under stringent conditions by techniques well-known in the art, as described, for example, in Keller and Manak (1993). Detection of the probe provides a means for determining in a known manner whether hybridization has occurred. Such a probe analysis provides a rapid method for identifying toxin-encoding genes of the subject invention. The nucleotide segments which are used as probes according to the invention can be synthesized using a DNA synthesizer and standard procedures. These nucleotide sequences can also be used as PCR primers to amplify genes of the subject invention.

Nucleic acid hybridization

[0092] As is well known to those skilled in molecular biology, similarity of two nucleic acids can be characterized by their tendency to hybridize. As used herein the terms "stringent conditions" or "stringent hybridization conditions" are intended to refer to conditions under which a probe will hybridize (anneal) to its target sequence to a detectably greater degree than to other sequences (e.g. at least 2-fold over background). Stringent conditions are sequence-dependent and will be different in different circumstances. By controlling the stringency of the hybridization and/or washing conditions, target sequences that are 100% complementary to the probe can be identified (homologous probing). Alternatively, stringency conditions can be adjusted to allow some mismatching in sequences so that lower degrees of similarity are detected (heterologous probing). Generally, a probe is less than about 1000 nucleotides in length, preferably less than 500 nucleotides in length.

[0093] Typically, stringent conditions will be those in which the salt concentration is less than about 1.5 M Na ion, typically about 0.01 to 1.0 M Na ion concentration (or other salts) at pH 7.0 to pH 8.3 and the temperature is at least about 30°C for short probes (e.g. 10 to 50 nucleotides) and at least about 60°C for long probes (e.g. greater than 50 nucleotides). Stringent conditions may also be achieved with the addition of destabilizing agents such as formamide. Exemplary low stringency conditions include hybridization with a buffer solution of 30% to 35% formamide, 1 M NaCl, 1% SDS (sodium dodecyl sulfate) at 37°C and a wash in 1X to 2X SSC (20X SSC = 3.0 M NaCl/0.3 M trisodium citrate) at 50°C to 55°C. Exemplary moderate stringency conditions include hybridization in 40% to 45% formamide, 1.0 M NaCl, 1% SDS at 37°C and a wash in 0.5X to 1X SSC at 55°C to 60°C. Exemplary high stringency conditions include hybridization in 50% formamide, 1 M NaCl, 1% SDS at 37°C and a wash in 0.1X SSC at 60°C to 65°C. Optionally, wash buffers may comprise about 0.1% to about 1% SDS. Duration of hybridization is generally less than about 24 hours, usually about 4 to about 12 hours.

[0094] Specificity is typically the function of post-hybridization washes, the critical factors being the ionic strength and temperature of the final wash solution. For DNA/DNA hybrids, the thermal melting point (T_m) is the temperature (under defined ionic strength and pH) at which 50% of a complementary target sequence hybridizes to a perfectly matched

probe. T_m is reduced by about 1°C for each 1% of mismatching; thus, T_m , hybridization conditions, and/or wash conditions can be adjusted to facilitate annealing of sequences of the desired identity. For example, if sequences with >90% identity are sought, the T_m can be decreased 10°C. Generally, stringent conditions are selected to be about 5°C lower than the T_m for the specific sequence and its complement at a defined ionic strength and pH. However, highly stringent conditions can utilize a hybridization and/or wash at 1°C, 2°C, 3°C, or 4°C lower than the T_m ; moderately stringent conditions can utilize a hybridization and/or wash at 6°C, 7°C, 8°C, 9°C, or 10°C lower than the T_m , and low stringency conditions can utilize a hybridization and/or wash at 11°C, 12°C, 13°C, 14°C, 15°C, or 20°C lower than the T_m .

[0095] T_m (in °C) may be experimentally determined or may be approximated by calculation. For DNA-DNA hybrids, the T_m can be approximated from the equation of Meinkoth and Wahl (1984):

$$T_m(^{\circ}\text{C}) = 81.5^{\circ}\text{C} + 16.6(\log M) + 0.41(\% \text{GC}) - 0.61(\% \text{ formamide}) - 500/L;$$

where M is the molarity of monovalent cations, %GC is the percentage of guanosine and cytosine nucleotides in the DNA, % formamide is the percentage of formamide in the hybridization solution, and L is the length of the hybrid in base pairs.

[0096] Alternatively, the T_m is described by the following formula (Beltz *et al.*, 1983).

$$T_m(^{\circ}\text{C}) = 81.5^{\circ}\text{C} + 16.6(\log[\text{Na}^+]) + 0.41(\% \text{GC}) - 0.61(\% \text{ formamide}) - 600/L$$

where $[\text{Na}^+]$ is the molarity of sodium ions, %GC is the percentage of guanosine and cytosine nucleotides in the DNA, % formamide is the percentage of formamide in the hybridization solution, and L is the length of the hybrid in base pairs.

[0097] Using the equations, hybridization and wash compositions, and desired T_m , those of ordinary skill will understand that variations in the stringency of hybridization and/or wash solutions are inherently described. If the desired degree of mismatching results in a T_m of less than 45°C (aqueous solution) or 32°C (formamide solution), it is preferred to increase the SSC concentration so that a higher temperature can be used. An extensive guide to the hybridization of nucleic acids is found in Tijssen (1993) and Ausubel *et al.* (1995). Also see Sambrook *et al.* (1989).

[0098] Hybridization of immobilized DNA on Southern blots with radioactively labeled gene-specific probes may be performed by standard methods Sambrook *et al.*, *supra.*). Radioactive isotopes used for labeling polynucleotide probes may include ^{32}P , ^{33}P , ^{14}C , or ^3H . Incorporation of radioactive isotopes into polynucleotide probe molecules may be done by any of several methods well known to those skilled in the field of molecular biology. (See, e.g. Sambrook *et al.*, *supra.*) In general, hybridization and subsequent washes may be carried out under stringent conditions that allow for detection of target sequences with homology to the claimed toxin encoding genes. For double-stranded DNA gene probes, hybridization may be carried out overnight at 20°-25°C below the T_m of the DNA hybrid in 6X SSPE, 5X Denhardt's Solution, 0.1% SDS, 0.1 mg/mL denatured DNA [20X SSPE is 3M NaCl, 0.2 M NaH_2PO_4 , and 0.02M EDTA (ethylenediamine tetra-acetic acid sodium salt); 100X Denhardt's Solution is 20 gm/L Polyvinylpyrrolidone, 20 gm/L Ficoll type 400 and 20 gm/L Bovine Serum Albumin (fraction V)].

[0099] Washes may typically be carried out as follows:

Twice at room temperature for 15 minutes in 1X SSPE, 0.1% SDS (low stringency wash).

Once at $T_m - 20^{\circ}\text{C}$ for 15 minutes in 0.2X SSPE, 0.1% SDS (moderate stringency wash).

[0100] For oligonucleotide probes, hybridization may be carried out overnight at 10°-20°C below the T_m of the hybrid in 6X SSPE, 5X Denhardt's solution, 0.1% SDS, 0.1 mg/mL denatured DNA. T_m for oligonucleotide probes may be determined by the following formula (Suggs *et al.*, 1981).

$$T_m(^{\circ}\text{C}) = 2(\text{number of T/A base pairs}) + 4(\text{number of G/C base pairs})$$

[0101] Washes may typically be carried out as follows:

Twice at room temperature for 15 minutes 1X SSPE, 0.1% SDS (low stringency wash).

Once at the hybridization temperature for 15 minutes in 1X SSPE, 0.1% SDS (moderate stringency wash).

[0102] Probe molecules for hybridization and hybrid molecules formed between probe and target molecules may be rendered detectable by means other than radioactive labeling. Such alternate methods are intended to be within the scope of this invention.

[0103] Unless specifically indicated or implied, the terms "a", "an", and "the" signify "at least one" as used herein. By the use of the term "genetic material" herein, it is meant to include all genes, nucleic acid, DNA and RNA.

[0104] For designations of nucleotide residues of polynucleotides, DNA, RNA, oligonucleotides, and primers, and for designations of amino acid residues of proteins, standard IUPAC abbreviations are employed throughout this document. Nucleic acid sequences are presented in the standard 5' to 3' direction, and protein sequences are presented in the standard amino (N) terminal to carboxy (C) terminal direction.

[0105] Following are examples that illustrate procedures for practicing the invention. It should be understood that the examples and embodiments described herein are for illustrative purposes only and that various modifications or changes in light thereof will be suggested to persons skilled in the art and are to be included within the purview of this application and the scope of the appended claims. These examples should not be construed as limiting. All percentages are by weight and all solvent mixture proportions are by volume unless otherwise noted. All temperatures are in degrees Celsius.

EXAMPLE 1

Isolation of a gene encoding DIG-3 toxin

[0106] Nucleic acid encoding the insecticidal Cry protein designated herein as DIG-3 was isolated from genomic DNA of *B.t.* strain PS46L by PCR using a degenerate forward primer that hybridized to bases 1286 to 1311 of SEQ ID NO:1, and a mismatched reverse primer that hybridized to the complement of bases 2480 to 2499 of SEQ ID NO: 1. This pair of primers was used to amplify a fragment of 1214 bp, corresponding to nucleotides 1286 to 2499 of SEQ ID NO: 1. This sequence was used as the anchor point to begin genome walking using methods adapted from the GenomeWalker™ Universal Kit (Clontech, Palo Alto, CA). The nucleic acid sequence of a fragment spanning the DIG-3 coding region was determined. SEQ ID NO:1 is the 3771 bp nucleotide sequence encoding the full length DIG-3 protein. SEQ ID NO:2 is the amino acid sequence of the full length DIG-3 protein deduced from SEQ ID NO: 1. It is noted that in *Bacillus* species, protein coding regions such as that of SEQ ID NO:1 may initiate with the TTG codon, which translationally represents the amino acid methionine.

EXAMPLE 2

Deletion of Domain I α -helices from DIG-3

[0107] To improve the insecticidal properties of the DIG-3 toxin, serial, step-wise deletions are made, each of which removes part of the N-terminus of the DIG-3 protein. The deletions remove part or all of α -helix 1 and part or all of α -helix 2 in Domain I, while maintaining the structural integrity of α -helix 3 through α -helix 7.

[0108] Deletions were designed as follows. This example utilizes the full length chimeric DNA sequence encoding the full-length DIG-3 protein e.g. SEQ ID NO:1 and SEQ ID NO:2, respectively) to illustrate the design principles with 67 specific variants. It utilizes the chimeric sequence of SEQ ID NO:5 (DNA encoding DIG-3 core toxin segment fused to Cry1Ab protoxin segment) to provide an additional 67 specific variants. One skilled in the art will realize that other DNA sequences encoding all or an N-terminal portion of the DIG-3 protein may be similarly manipulated to achieve the desired result. To devise the first deleted variant coding sequence, all of the bases that encode α -helix 1 up to the codon for the proline residue near the beginning of α -helix 2A (*i.e.* P73 for the full length DIG-3 protein of SEQ ID NO:2), are removed. Thus, elimination of bases 1 to 216 of SEQ ID NO:1 removes the coding sequence for amino acids 1 to 72 of SEQ ID NO:2. Reintroduction of a translation initiating ATG (methionine) codon at the beginning (*i.e.* in front of the codon corresponding to amino acid 73 of the full length protein) provides for the deleted variant coding sequence comprising an open reading frame of 3555 bases which encodes a deleted variant DIG-3 protein comprising 1185 amino acids (*i.e.* methionine plus amino acids 73 to 1256 of the full-length DIG-3 protein). Serial, stepwise deletions that remove additional codons for a single amino acid corresponding to residues 73 to 112 of the full-length DIG-3 protein of SEQ ID NO:2 provide variants lacking part or all of α -helix 2A and α -helix 2B. Thus a second designed deleted variant coding sequence requires elimination of bases 1 to 219 of SEQ ID NO:1, thereby removing the coding sequence for amino acids 1-73. Restoration of a functional open reading frame is again accomplished by reintroduction of a translation initiation methionine codon at the beginning of the remaining coding sequence, thus providing for a second deleted variant coding sequence having an open reading frame of 3552 bases encoding a deleted variant DIG-3 protein comprising 1184 amino acids (*i.e.* methionine plus amino acids 74 to 1256 of the full-length DIG-3 protein). The last designed deleted variant coding sequence requires removal of bases 1 to 336 of SEQ ID NO:1, thus eliminating the coding sequence for amino acids 1 to 112, and, after reintroduction of a translation initiation methionine codon, providing a deletion variant coding

EP 2 419 441 B1

sequence having an open reading frame of 3435 bases which encodes a deletion variant DIG-3 protein of 1145 amino acids (*i.e.* methionine plus amino acids 113 to 1256 of the full-length DIG-3 protein). As exemplified, after elimination of the deletion sequence, an initiator methionine codon is added to the beginning of the remaining coding sequence to restore a functional open reading frame. Also as described, an additional glycine codon is to be added between the methionine codon and the codon for the instability-determining amino acid in the instance that removal of the deleted sequence leaves exposed at the N-terminus of the remaining portion of the full-length protein one of the instability-determining amino acids as provided above.

[0109] Table 3 describes specific variants designed in accordance with the strategy described above.

Table 3

Deletion variant protein sequences of the full-length DIG-3 protein of SEQ ID NO:2 and the fusion protein sequence of SEQ ID NO:5.					
DIG-3Deletion Variant	Residues added at NH2 terminus	Residues of SEQ ID NO:2	DIG-3Deletion Variant	Residues added at NH2 terminus	Residues of SEQ ID NO:5
1	M	73-1256	68	M	73-1188
2	MG	73-1256	69	MG	73-1188
3	M	74-1256	70	M	74-1188
4	MG	74-1256	71	MG	74-1188
5	M	75-1256	72	M	75-1188
6	M	76-1256	73	M	76-1188
7	M	77-1256	74	M	77-1188
8	M	78-1256	75	M	78-1188
9	MG	78-1256	76	MG	78-1188
10	M	79-1256	77	M	79-1188
11	M	80-1256	78	M	80-1188
12	M	81-1256	79	M	81-1188
13	MG	81-1256	80	MG	81-1188
14	M	82-1256	81	M	82-1188
15	MG	82-1256	82	MG	82-1188
16	M	83-1256	83	M	83-1188
17	M	84-1256	84	M	84-1188
18	MG	84-1256	85	MG	84-1188
19	M	85-1256	86	M	85-1188
20	MG	85-1256	87	MG	85-1188
21	M	86-1256	88	M	86-1188
22	M	87-1256	89	M	87-1188
23	M	88-1256	90	M	88-1188
24	M	89-1256	91	M	89-1188
25	MG	89-1256	92	MG	89-1188
26	M	90-1256	93	M	90-1188
27	MG	90-1256	94	MG	90-1188
28	M	91-1256	95	M	91-1188
29	MG	91-1256	96	MG	91-1188

EP 2 419 441 B1

(continued)

Deletion variant protein sequences of the full-length DIG-3 protein of SEQ ID NO:2 and the fusion protein sequence of SEQ ID NO:5.					
DIG-3Deletion Variant	Residues added at NH2 terminus	Residues of SEQ ID NO:2	DIG-3Deletion Variant	Residues added at NH2 terminus	Residues of SEQ ID NO:5
30	M	92-1256	97	M	92-1188
31	M	93-1256	98	M	93-1188
32	M	94-1256	99	M	94-1188
33	M	95-1256	100	M	95-1188
34	MG	95-1256	101	MG	95-1188
35	M	96-1256	102	M	96-1188
36	MG	96-1256	103	MG	96-1188
37	M	97-1256	104	M	97-1188
38	MG	97-1256	105	MG	97-1188
39	M	98-1256	106	M	98-1188
40	MG	98-1256	107	MG	98-1188
41	M	99-1256	108	M	99-1188
42	MG	99-1256	109	MG	99-1188
43	M	100-1256	110	M	100-1188
44	MG	100-1256	111	MG	100-1188
45	M	101-1256	112	M	101-1188
46	MG	101-1256	113	MG	101-1188
47	M	102-1256	114	M	102-1188
48	MG	102-1256	115	MG	102-1188
49	M	103-1256	116	M	103-1188
50	MG	103-1256	117	MG	103-1188
51	M	104-1256	118	M	104-1188
52	M	105-1256	119	M	105-1188
53	MG	105-1256	120	MG	105-1188
54	M	106-1256	121	M	106-1188
55	MG	106-1256	122	MG	106-1188
56	M	107-1256	123	M	107-1188
57	MG	107-1256	124	MG	107-1188
58	M	108-1256	125	M	108-1188
59	MG	108-1256	126	MG	108-1188
60	M	109-1256	127	M	109-1188
61	MG	109-1256	128	MG	109-1188
62	M	110-1256	129	M	110-1188
63	MG	110-1256	130	MG	110-1188
64	M	111-1256	131	M	111-1188

(continued)

Deletion variant protein sequences of the full-length DIG-3 protein of SEQ ID NO:2 and the fusion protein sequence of SEQ ID NO:5.					
DIG-3 Deletion Variant	Residues added at NH2 terminus	Residues of SEQ ID NO:2	DIG-3 Deletion Variant	Residues added at NH2 terminus	Residues of SEQ ID NO:5
65	MG	111-1256	132	MG	111-1188
66	M	112-1356	133	M	112-1356
67	M	113-1256	134	M	113-1188

[0110] Nucleic acids encoding the toxins described in Table 3 are designed in accordance with the general principles for synthetic genes intended for expression in plants, as discussed above.

EXAMPLE 3

Design of a plant-optimized version of the coding sequence for the DIG-3 *B.t.* insecticidal protein

[0111] A DNA sequence having a plant codon bias was designed and synthesized to produce the DIG-3 protein in transgenic monocot and dicot plants. A codon usage table for maize (*Zea mays* L.) was calculated from 706 protein coding sequences (CDs) obtained from sequences deposited in GenBank. Codon usage tables for tobacco (*Nicotiana tabacum*, 1268 CDs), canola (*Brassica napus*, 530 CDs), cotton (*Gossypium hirsutum*, 197 CDs), and soybean (*Glycine max*; ca. 1000 CDs) were downloaded from data at the website <http://www.kazusa.or.jp/codon/>. A biased codon set that comprises highly used codons common to both maize and dicot datasets, in appropriate weighted average relative amounts, was calculated after omitting any redundant codon used less than about 10% of total codon uses for that amino acid in either plant type. To derive a plant optimized sequence encoding the DIG-3 protein, codon substitutions to the experimentally determined DIG-3 DNA sequence were made such that the resulting DNA sequence had the overall codon composition of the plant-optimized codon bias table. Further refinements of the sequence were made to eliminate undesirable restriction enzyme recognition sites, potential plant intron splice sites, long runs of A/T or C/G residues, and other motifs that might interfere with RNA stability, transcription, or translation of the coding region in plant cells. Other changes were made to introduce desired restriction enzyme recognition sites, and to eliminate long internal Open Reading Frames (frames other than +1). These changes were all made within the constraints of retaining the plant-biased codon composition. Synthesis of the designed sequence was performed by a commercial vendor (DNA2.0, Menlo Park, CA).

[0112] Additional guidance regarding the production of synthetic genes can be found in, for example, WO 97/13402 and US Patent No. 5380831.

[0113] A plant-optimized DNA sequence encoding the full length DIG-3 toxin is given in SEQ ID NO:3. A dicot-optimized DNA sequence encoding the Cry1Ab protoxin segment is disclosed as SEQ ID NO:6. A maize- optimized DNA sequence encoding the Cry1Ab protoxin segment is disclosed as SEQ ID NO:7.

EXAMPLE 4

Construction of expression plasmids encoding DIG-3 insecticidal toxin and expression in bacterial hosts

[0114] Standard cloning methods were used in the construction of *Pseudomonas fluorescens* (*Pj*) expression plasmids engineered to produce full-length DIG-3 proteins encoded by plant-optimized coding regions. Restriction endonucleases were obtained from New England BioLabs (NEB; Ipswich, MA) and T4 DNA Ligase (Invitrogen) was used for DNA ligation. Plasmid preparations were performed using the NucleoBond® Xtra Kit (Macherey-Nagel Inc, Bethlehem, PA) or the Plasmid Midi Kit® (Qiagen), following the instructions of the suppliers. DNA fragments were purified using the Millipore Ultrafree®-DA cartridge (Billerica, MA) after agarose Tris-acetate gel electrophoresis.

[0115] The basic cloning strategy entailed subcloning the DIG-3 toxin coding sequence (CDS) into pDOW1169 at the *SpeI* and *XhoI* restriction sites, whereby it is placed under the expression control of the *Ptac* promoter and the *rrnBT12* terminator from plasmid pKK223-3 (PL Pharmacia, Milwaukee, WI). pDOW1169 is a medium copy plasmid with the RSF1010 origin of replication, a *pyrF* gene, and a ribosome binding site preceding the restriction enzyme recognition sites into which DNA fragments containing protein coding regions may be introduced, (US Application 20080193974). The expression plasmid, designated pDAB4171, was transformed by electroporation into DC454 (a near wild-type *P. fluorescens* strain having mutations $\Delta pyrF$ and *lsc::lacI^Q*), or its derivatives, recovered in SOC-Soy hydrolysate medium, and plated on selective medium (M9 glucose agar lacking uracil, Sambrook *et al.*, *supra*). Details of the microbiological

manipulations are available in Squires et al., (2004), US Patent Application 20060008877, US Patent Application 20080193974, and US Patent Application 20080058262. Colonies were first screened by PCR and positive clones were then analyzed by restriction digestion of miniprep plasmid DNA. Plasmid DNA of selected clones containing inserts was sequenced, either by using Big Dye® Terminator version 3.1 as recommended by the supplier (Applied Biosystems/Invitrogen), or by contract with a commercial sequencing vendor (MWG Biotech, Huntsville, AL). Sequence data were assembled and analyzed using the Sequencher™ software (Gene Codes Corp., Ann Arbor, MI).

[0116] Growth and Expression Analysis in Shake Flasks. Production of DIG-3 toxin for characterization and insect bioassay was accomplished by shake-flask-grown *P. fluorescens* strains harboring expression constructs (e.g. clone DP2826). Seed cultures grown in M9 medium supplemented with 1% glucose and trace elements were used to inoculate 50 mL of defined minimal medium with 5% glycerol (Teknova Catalog No. 3D7426, Hollister, CA). Expression of the DIG-3 toxin gene via the *Ptac* promoter was induced by addition of isopropyl-β-D-1-thiogalactopyranoside (IPTG) after an initial incubation of 24 hours at 30° with shaking. Cultures were sampled at the time of induction and at various times post-induction. Cell density was measured by optical density at 600 nm (OD₆₀₀). Other culture media suitable for growth of *Pseudomonas fluorescens* may also be utilized, for example, as described in Huang et al. (2007) and US Patent Application 20060008877.

[0117] Cell Fractionation and SDS-PAGE Analysis of Shake Flask Samples. At each sampling time, the cell density of samples was adjusted to OD₆₀₀ = 20 and 1 mL aliquots were centrifuged at 14000 x g for five minutes. The cell pellets were frozen at -80°. Soluble and insoluble fractions from frozen shake flask cell pellet samples were generated using EasyLyse™ Bacterial Protein Extraction Solution (EPICENTRE® Biotechnologies, Madison, WI). Each cell pellet was resuspended in 1 mL EasyLyse™ solution and further diluted 1:4 in lysis buffer and incubated with shaking at room temperature for 30 minutes. The lysate was centrifuged at 14,000 rpm for 20 minutes at 4° and the supernatant was recovered as the soluble fraction. The pellet (insoluble fraction) was then resuspended in an equal volume of phosphate buffered saline (PBS; 11.9 mM Na₂HPO₄, 137 mM NaCl, 2.7 mM KCl, pH7.4).

[0118] Samples were mixed 1:1 with 2X Laemmli sample buffer containing β-mercaptoethanol (Sambrook *et al.*, *supra.*) and boiled for 5 minutes prior to loading onto Criterion XT® Bis-Tris 12% gels (Bio-Rad Inc., Hercules, CA.) Electrophoresis was performed in the recommended XT MOPS buffer. Gels were stained with Bio-Safe Coomassie Stain according to the manufacturer's (Bio-Rad) protocol and imaged using the Alpha Innotech Imaging system (San Leandro, CA).

[0119] Inclusion body preparation. Cry protein inclusion body (IB) preparations were performed on cells from *P. fluorescens* fermentations that produced insoluble *B.t.* insecticidal protein, as demonstrated by SDS-PAGE and MALDI-MS (Matrix Assisted Laser Desorption/Ionization Mass Spectrometry). *P. fluorescens* fermentation pellets were thawed in a 37° water bath. The cells were resuspended to 25% w/v in lysis buffer (50 mM Tris, pH 7.5, 200 mM NaCl, 20 mM EDTA disodium salt (Ethylenediaminetetraacetic acid), 1% Triton X-100, and 5 mM Dithiothreitol (DTT); 5 mL/L of bacterial protease inhibitor cocktail (P8465 Sigma-Aldrich, St. Louis, MO) were added just prior to use). The cells were suspended using a hand-held homogenizer at lowest setting (Tissue Tearor, BioSpec Products, Inc., Bartlesville, OK). Lysozyme (25 mg of Sigma L7651, from chicken egg white) was added to the cell suspension by mixing with a metal spatula, and the suspension was incubated at room temperature for one hour. The suspension was cooled on ice for 15 minutes, then sonicated using a Branson Sonifier 250 (two 1- minute sessions, at 50% duty cycle, 30% output). Cell lysis was checked by microscopy. An additional 25 mg of lysozyme were added if necessary, and the incubation and sonication were repeated. When cell lysis was confirmed via microscopy, the lysate was centrifuged at 11,500 x g for 25 minutes (4°) to form the IB pellet, and the supernatant was discarded. The IB pellet was resuspended with 100 mL lysis buffer, homogenized with the hand-held mixer and centrifuged as above. The IB pellet was repeatedly washed by resuspension (in 50 mL lysis buffer), homogenization, sonication, and centrifugation until the supernatant became colorless and the IB pellet became firm and off-white in color. For the final wash, the IB pellet was resuspended in sterile-filtered (0.22 μm) distilled water containing 2 mM EDTA, and centrifuged. The final pellet was resuspended in sterile-filtered distilled water containing 2 mM EDTA, and stored in 1 mL aliquots at -80°.

[0120] SDS-PAGE analysis and quantitation of protein in IB preparations was done by thawing a 1 mL aliquot of IB pellet and diluting 1:20 with sterile-filtered distilled water. The diluted sample was then boiled with 4X reducing sample buffer [250 mM Tris, pH6.8, 40% glycerol (v/v), 0.4% Bromophenol Blue (w/v), 8% SDS (w/v) and 8% β-Mercapto-ethanol (v/v)] and loaded onto a Novex® 4-20% Tris-Glycine, 12+2 well gel (Invitrogen) run with 1X Tris/Glycine/SDS buffer (BioRad). The gel was run for 60 min at 200 volts then stained with Coomassie Blue (50% G-250/50% R-250 in 45% methanol, 10% acetic acid), and destained with 7% acetic acid, 5% methanol in distilled water. Quantification of target bands was done by comparing densitometric values for the bands against Bovine Serum Albumin (BSA) samples run on the same gel to generate a standard curve.

[0121] Solubilization of Inclusion Bodies. Six mL of inclusion body suspension from Pf clone DP2826 (containing 32 mg/mL of DIG-3 protein) were centrifuged on the highest setting of an Eppendorf model 5415C microfuge (approximately 14,000 x g) to pellet the inclusions. The storage buffer supernatant was removed and replaced with 25 mL of 100 mM sodium carbonate buffer, pH11, in a 50 mL conical tube. Inclusions were resuspended using a pipette and vortexed to mix thoroughly. The tube was placed on a gently rocking platform at 4° overnight to extract the target protein. The extract

was centrifuged at 30,000 x g for 30 min at 4°, and the resulting supernatant was concentrated 5-fold using an Amicon Ultra-15 regenerated cellulose centrifugal filter device (30,000 Molecular Weight Cutoff; Millipore). The sample buffer was then changed to 10 mM CAPS [3-(cyclohexamino)1-propanesulfonic acid] pH 10, using disposable PD-10 columns (GE Healthcare, Piscataway, NJ).

[0122] Gel electrophoresis. The concentrated extract was prepared for electrophoresis by diluting 1:50 in NuPAGE® LDS sample buffer (Invitrogen) containing 5 mM dithiothreitol as a reducing agent and heated at 95° for 4 minutes. The sample was loaded in duplicate lanes of a 4-12% NuPAGE® gel alongside five BSA standards ranging from 0.2 to 2 µg/lane (for standard curve generation). Voltage was applied at 200V using MOPS SDS running buffer (Invitrogen) until the tracking dye reached the bottom of the gel. The gel was stained with 0.2% Coomassie Blue G-250 in 45% methanol, 10% acetic acid, and destained, first briefly with 45% methanol, 10% acetic acid, and then at length with 7% acetic acid, 5% methanol until the background cleared. Following destaining, the gel was scanned with a Biorad Fluor-S Multimager. The instrument's Quantity One v.4.5.2 Software was used to obtain background-subtracted volumes of the stained protein bands and to generate the BSA standard curve that was used to calculate the concentration of DIG-3 protein in the stock solution.

EXAMPLE 5

Insecticidal activity of modified DIG-3 protein produced in *Pseudomonas fluorescens*

[0123] DIG-3 *B.t.* insecticidal toxin was demonstrated to be active on Lepidopteran species including the European corn borer (ECB; *Ostrinia nubilalis* (Hübner)), cry1F-resistant ECB (rECB), diamondback moth (DBM; *Plutella xylostella* (Linnaeus)), cry1A-resistant DBM (rDBM), corn earworm (CEW; *Helicoverpa zea* (Boddie)), black cutworm (BCW; *Agrotis ipsilon* (Hufnagel)), tobacco budworm (TBW; *Heliothis virescens* (Fabricius)), and cabbage looper (CL; *Trichoplusia ni* (Hübner)). DIG-3 protein was also tested for activity on fall armyworm (FAW, *Spodoptera frugiperda*), Cry1F-resistant FAW (rFAW) and western corn rootworm (WCR, *Diabrotica virgifera virgifera* LeConte).

[0124] Sample preparation and bioassays. Inclusion body preparations in 10 mM CAPS pH10 were diluted appropriately in 10 mM CAPS pH 10, and all bioassays contained a control treatment consisting of this buffer, which served as a background check for mortality or growth inhibition.

[0125] Protein concentrations in bioassay buffer were estimated by gel electrophoresis using BSA to create a standard curve for gel densitometry, which was measured using a BioRad imaging system (Fluor-S Multimager with Quantity One software version 4.5.2). Proteins in the gel matrix were stained with Coomassie Blue-based stain and destained before reading.

[0126] Purified proteins were tested for insecticidal activity in bioassays conducted with neonate Lepidopteran larvae on artificial insect diet. Larvae of BCW, CEW, CL, DBM, rDBM, ECB, FAW and TBW were hatched from eggs obtained from a colony maintained by a commercial insectary (Benzon Research Inc., Carlisle, PA). WCR eggs were obtained from Crop Characteristics, Inc. (Farmington, MN). Larvae of rECB and rFAW were hatched from eggs harvested from proprietary colonies (Dow AgroSciences LLC, Indianapolis, IN).

[0127] The bioassays were conducted in 128-well plastic trays specifically designed for insect bioassays (C-D International, Pitman, NJ). Each well contained 1.0 mL of Multi-species Lepidoptera diet (Southland Products, Lake Village, AR). A 40 µL aliquot of protein sample was delivered by pipette onto the 1.5 cm² diet surface of each well (26.7 µL/cm²). Diet concentrations were calculated as the amount (ng) of DIG-3 protein per square centimeter (cm²) of surface area in the well. The treated trays were held in a fume hood until the liquid on the diet surface had evaporated or was absorbed into the diet.

[0128] Within a few hours of eclosion, individual larvae were picked up with a moistened camel hair brush and deposited on the treated diet, one larva per well. The infested wells were then sealed with adhesive sheets of clear plastic, vented to allow gas exchange (C-D International, Pitman, NJ). Bioassay trays were held under controlled environmental conditions (28°C, ~40% Relative Humidity, 16:8 [Light:Dark]) for 5 days, after which the total number of insects exposed to each protein sample, the number of dead insects, and the weight of surviving insects were recorded. Percent mortality and percent growth inhibition were calculated for each treatment. Growth inhibition (GI) was calculated as follows:

$$GI = [1 - (TWIT/TNIT)/(TWIBC/TNIBC)]$$

where TWIT is the Total Weight of Insects in the Treatment,

TNIT is the Total Number of Insects in the Treatment

TWIBC is the Total Weight of Insects in the Background Check (Buffer control), and

TNIBC is the Total Number of Insects in the Background Check (Buffer control).

[0129] The GI_{50} was determined to be the concentration of DIG-3 protein in the diet at which the GI value was 50%. The LC_{50} (50% Lethal Concentration) was recorded as the concentration of DIG-3 protein in the diet at which 50% of test insects were killed. Statistical analysis (One-way ANOVA) was done using JMP software (SAS, Cary, NC)

[0130] Table 6 presents the results of bioassay tests of DIG-3 protein on European corn borer and cry1F-resistant European corn borer (rECB). An unexpected and surprising finding is that the rECB test insects were as susceptible to the action of DIG-3 protein as were the wild type ECB insects.

Table 6

LC ₅₀ and GI ₅₀ values calculated for ECB and rECB, with Confidence Intervals (CI) of 95%				
Insect	LC ₅₀ (ng/cm ²)	95% CI	GI ₅₀ (ng/cm ²)	95% CI
ECB	591.9	308.1-1315.3	122.6	45.6-328.4
rECB	953.6	534.1-1953.6	270.9	53.0-1382.2

[0131] Table 7 presents the results of bioassays on a broad spectrum of Lepidopteran and a Coleopteran pest (WCR). The DIG-3 protein has unexpected and surprising activity on diamondback moth as well as rDBM. Further the DIG-3 Cry protein is effective in controlling the growth of several other Lepidopteran insects.

Table 7

Insecticidal and growth inhibitory effects of DIG-3 protein ingested by test insects		
Test Insect	Response at 9000 ng/cm ²	Statistical Analysis*
DBM	100% mortality	
rDBM	100% mortality	
CL	75% mortality, significant GI	(GI) P < 0.001, df = 1, α = 0.05
CEW	Significant GI	(GI) P = 0.02, df = 1, α = 0.05
TBW	Visible GI, some mortality	Not available
BCW	Visible GI	Not available
FAW	No activity observed	
rFAW	No activity observed	
WCR	No activity observed	
*GI = Growth Inhibition. P-value = test statistic. df = Degrees of Freedom, α (alpha) level 0.05 = the level of test significance.		

EXAMPLE 6

Agrobacterium transformation

[0132] Standard cloning methods are used in the construction of binary plant transformation and expression plasmids. Restriction endonucleases and T4 DNA Ligase are obtained from NEB. Plasmid preparations are performed using the NucleoSpin® Plasmid Preparation kit or the NucleoBond® AX Xtra Midi kit (both from Macherey-Nagel), following the instructions of the manufacturers. DNA fragments are purified using the QIAquick® PCR Purification Kit or the QIAEX II® Gel Extraction Kit (both from Qiagen) after gel isolation.

[0133] DNA fragments comprising nucleotide sequences that encode the DIG-3 protein in native or modified forms, or fragments thereof, may be synthesized by a commercial vendor (e.g. DNA2.0, Menlo Park, CA) and supplied as cloned fragments in standard plasmid vectors, or may be obtained by standard molecular biology manipulation of other constructs containing appropriate nucleotide sequences. Unique restriction sites internal to a DIG-3 coding region may be identified and DNA fragments comprising the sequences between the restriction sites of the DIG-3 coding region may be synthesized, each such fragment encoding a specific deletion, insertion or other DIG-3 variation. The DNA fragments encoding the modified DIG-3 fragments may be joined to other DIG-3 coding region fragments or other Cry coding region fragments at appropriate restriction sites to obtain a coding region encoding the desired full-length DIG-3 protein, deleted or variant DIG-3 protein, or fused protein. For example, one may identify an appropriate restriction

recognition site at the start of a first DIG-3 coding region, and a second restriction site internal to the DIG-3 coding region. Cleavage of this first DIG-3 coding region at these restriction sites would generate a DNA fragment comprising part of the first DIG-3 coding region. A second DNA fragment flanked by analogously-situated compatible restriction sites specific for another DIG-3 coding region or other Cry coding region may be used in combination with the first DNA restriction fragment to construct a variant or fused clone.

[0134] In a non-limiting example, a basic cloning strategy may be to subclone full length or modified DIG-3 coding sequences (CDS) into a plant expression plasmid at *NcoI* and *SacI* restriction sites. The resulting plant expression cassettes containing the appropriate DIG-3 coding region under the control of plant expression elements, (e.g., plant expressible promoters, 3' terminal transcription termination and polyadenylate addition determinants, and the like) are subcloned into a binary vector plasmid, utilizing, for example, Gateway® technology or standard restriction enzyme fragment cloning procedures. LR Clonase™ (Invitrogen) for example, may be used to recombine the full length and modified gene plant expression cassettes into a binary plant transformation plasmid if the Gateway® technology is utilized. It is convenient to employ a binary plant transformation vector that harbors a bacterial gene that confers resistance to the antibiotic spectinomycin when the plasmid is present in *E. coli* and *Agrobacterium* cells. It is also convenient to employ a binary vector plasmid that contains a plant-expressible selectable marker gene that is functional in the desired host plants. Examples of plant-expressible selectable marker genes include but are not limited those that encode the aminoglycoside phosphotransferase gene (*aphII*) of transposon Tn5, which confers resistance to the antibiotics kanamycin, neomycin and G418, as well as those genes which code for resistance or tolerance to glyphosate; hygromycin; methotrexate; phosphinothricin (bialaphos), imidazolinones, sulfonylureas and triazopyrimidine herbicides, such as chlorosulfuron, bromoxynil, dalapon and the like.

[0135] Electro-competent cells of *Agrobacterium tumefaciens* strain Z707S (a streptomycin-resistant derivative of Z707; Hepburn *et al.*, 1985) are prepared and transformed using electroporation (Weigel and Glazebrook, 2002). After electroporation, 1 mL of YEP broth (gm/L: yeast extract, 10; peptone, 10; NaCl, 5) are added to the cuvette and the cell-YEP suspension is transferred to a 15 mL culture tube for incubation at 28° in a water bath with constant agitation for 4 hours. The cells are plated on YEP plus agar (25 gm/L) with spectinomycin (200 µg/mL) and streptomycin (250 µg/mL) and the plates are incubated for 2-4 days at 28°. Well separated single colonies are selected and streaked onto fresh YEP + agar plates with spectinomycin and streptomycin as before, and incubated at 28° for 1-3 days.

[0136] The presence of the DIG-3 gene insert in the binary plant transformation vector is performed by PCR analysis using vector-specific primers with template plasmid DNA prepared from selected *Agrobacterium* colonies. The cell pellet from a 4 mL aliquot of a 15 mL overnight culture grown in YEP with spectinomycin and streptomycin as before is extracted using Qiagen Spin® Mini Preps, performed per manufacturer's instructions. Plasmid DNA from the binary vector used in the *Agrobacterium* electroporation transformation is included as a control. The PCR reaction is completed using Taq DNA polymerase from Invitrogen per manufacture's instructions at 0.5X concentrations. PCR reactions are carried out in a MJ Research Peltier Thermal Cycler programmed with the following conditions: Step 1) 94° for 3 minutes; Step 2) 94° for 45 seconds; Step 3) 55° for 30 seconds; Step 4) 72° for 1 minute per kb of expected product length; Step 5) 29 times to Step 2; Step 6) 72° for 10 minutes. The reaction is maintained at 4° after cycling. The amplification products are analyzed by agarose gel electrophoresis (e.g. 0.7 % to 1% agarose, w/v) and visualized by ethidium bromide staining. A colony is selected whose PCR product is identical to the plasmid control.

[0137] Alternatively, the plasmid structure of the binary plant transformation vector containing the DIG-3 gene insert is performed by restriction digest fingerprint mapping of plasmid DNA prepared from candidate *Agrobacterium* isolates by standard molecular biology methods well known to those skilled in the art of *Agrobacterium* manipulation.

[0138] Those skilled in the art of obtaining transformed plants *via Agrobacterium*-mediated transformation methods will understand that other *Agrobacterium* strains besides Z707S may be used to advantage, and the choice of strain may depend upon the identity of the host plant species to be transformed.

EXAMPLE 7

Production of DIG-3 *B.t.* insecticidal proteins and variants in dicot plants

[0139] *Arabidopsis* Transformation. *Arabidopsis thaliana* Col-01 is transformed using the floral dip method (Weigel and Glazebrook, 2002). The selected *Agrobacterium* colony is used to inoculate 1 mL to 15 mL cultures of YEP broth containing appropriate antibiotics for selection. The culture is incubated overnight at 28° with constant agitation at 220 rpm. Each culture is used to inoculate two 500 mL cultures of YEP broth containing appropriate antibiotics for selection and the new cultures are incubated overnight at 28° with constant agitation. The cells are pelleted at approximately 8700 x g for 10 minutes at room temperature, and the resulting supernatant is discarded. The cell pellet is gently resuspended in 500 mL of infiltration media containing: 1/2x Murashige and Skoog salts (Sigma-Aldrich)/Gamborg's B5 vitamins (Gold BioTechnology, St. Louis, MO), 10% (w/v) sucrose, 0.044 µM benzylaminopurine (10 µL/liter of 1 mg/mL stock in DMSO) and 300 µL/liter Silwet L-77. Plants approximately 1 month old are dipped into the media for 15 seconds, with care taken

to assure submergence of the newest inflorescence. The plants are then laid on their sides and covered (transparent or opaque) for 24 hours, washed with water, and placed upright. The plants are grown at 22°, with a 16-hour light/8-hour dark photoperiod. Approximately 4 weeks after dipping, the seeds are harvested.

[0140] Arabidopsis Growth and Selection. Freshly harvested T1 seed is allowed to dry for at least 7 days at room temperature in the presence of desiccant. Seed is suspended in a 0.1% agar/water (Sigma-Aldrich) solution and then stratified at 4° for 2 days. To prepare for planting, Sunshine Mix LP5 (Sun Gro Horticulture Inc., Bellevue, WA) in 10.5 inch x 21 inch germination trays (T.O. Plastics Inc., Clearwater, MN) is covered with fine vermiculite, sub-irrigated with Hoagland's solution (Hoagland and Arnon, 1950) until wet, then allowed to drain for 24 hours. Stratified seed is sown onto the vermiculite and covered with humidity domes (KORD Products, Bramalea, Ontario, Canada) for 7 days. Seeds are germinated and plants are grown in a Conviron (Models CMP4030 or CMP3244; Controlled Environments Limited, Winnipeg, Manitoba, Canada) under long day conditions (16 hours light/8 hours dark) at a light intensity of 120-150 $\mu\text{mol}/\text{m}^2\text{sec}$ under constant temperature (22°) and humidity (40-50%). Plants are initially watered with Hoagland's solution and subsequently with deionized water to keep the soil moist but not wet.

[0141] The domes are removed 5-6 days post sowing and plants are sprayed with a chemical selection agent to kill plants germinated from nontransformed seeds. For example, if the plant expressible selectable marker gene provided by the binary plant transformation vector is a *pat* or *bar* gene (Wehrmann *et al.*, 1996), transformed plants may be selected by spraying with a 1000X solution of Finale (5.78% glufosinate ammonium, Farnam Companies Inc., Phoenix, AZ.). Two subsequent sprays are performed at 5-7 day intervals. Survivors (plants actively growing) are identified 7-10 days after the final spraying and transplanted into pots prepared with Sunshine Mix LP5. Transplanted plants are covered with a humidity dome for 3-4 days and placed in a Conviron under the above-mentioned growth conditions.

[0142] Those skilled in the art of dicot plant transformation will understand that other methods of selection of transformed plants are available when other plant expressible selectable marker genes (*e.g.* herbicide tolerance genes) are used.

[0143] Insect Bioassays of transgenic Arabidopsis. Transgenic *Arabidopsis* lines expressing modified Cry proteins are demonstrated to be active against sensitive insect species in artificial diet overlay assays. Protein extracted from transgenic and non-transgenic *Arabidopsis* lines is quantified by appropriate methods and sample volumes are adjusted to normalize protein concentration. Bioassays are conducted on artificial diet as described above. Non-transgenic *Arabidopsis* and/or buffer and water are included in assays as background check treatments.

EXAMPLE 8

Agrobacterium transformation for generation of superbinary vectors

[0144] The *Agrobacterium* superbinary system is conveniently used for transformation of monocot plant hosts. Methodologies for constructing and validating superbinary vectors are well disclosed and incorporated herein by reference (Operating Manual for Plasmid pSB1, Version 3.1, available from Japan Tobacco, Inc., Tokyo, Japan). Standard molecular biological and microbiological methods are used to generate superbinary plasmids. Verification/validation of the structure of the superbinary plasmid is done using methodologies as described above for binary vectors, and may be modified as suggested in the Operating Manual for Plasmid pSB1.

EXAMPLE 9

Production of DIG-3 *B.t.* insecticidal proteins and variants in monocot plants

[0145] Agrobacterium-Mediated Transformation of Maize. Seeds from a High II F₁ cross (Armstrong *et al.*, 1991) are planted into 5-gallon-pots containing a mixture of 95% Metro-Mix 360 soilless growing medium (Sun Gro Horticulture, Bellevue, WA) and 5% clay/loam soil. The plants are grown in a greenhouse using a combination of high pressure sodium and metal halide lamps with a 16:8 hour Light:Dark photoperiod. For obtaining immature F₂ embryos for transformation, controlled sib-pollinations are performed. Immature embryos are isolated at 8-10 days post-pollination when embryos are approximately 1.0 to 2.0 mm in size.

[0146] Infection and co-cultivation. Maize ears are surface sterilized by scrubbing with liquid soap, immersing in 70% ethanol for 2 minutes, and then immersing in 20% commercial bleach (0.1% sodium hypochlorite) for 30 minutes before being rinsed with sterile water. A suspension of *Agrobacterium* cells containing a superbinary vector is prepared by transferring 1-2 loops of bacteria grown on YEP solid medium containing 100 mg/L spectinomycin, 10 mg/L tetracycline, and 250 mg/L streptomycin at 28° for 2-3 days into 5 mL of liquid infection medium (LS Basal Medium (Linsmaier and Skoog, 1965), N6 vitamins (Chu *et al.*, 1975), 1.5 mg/L 2,4-Dichlorophenoxyacetic acid (2,4-D), 68.5 gm/L sucrose, 36.0 gm/L glucose, 6 mM L-proline, pH 5.2) containing 100 μM acetosyringone. The solution was vortexed until a uniform suspension was achieved, and the concentration is adjusted to a final density of about 200 Klett units, using a Klett-Summerson colorimeter with a purple filter, or an optical density of approximately 0.4 at 550 nm. Immature embryos are

isolated directly into a micro centrifuge tube containing 2 mL of the infection medium. The medium is removed and replaced with 1 mL of the *Agrobacterium* solution with a density of 200 Klett units, and the *Agrobacterium* and embryo solution is incubated for 5 minutes at room temperature and then transferred to co-cultivation medium (LS Basal Medium, N6 vitamins, 1.5 mg/L 2,4-D, 30.0 gm/L sucrose, 6 mM L-proline, 0.85 mg/L AgNO₃, 100 µM acetosyringone, 3.0 gm/L Gellan gum (*PhytoTechnology Laboratories.*, Lenexa, KS), pH 5.8) for 5 days at 25° under dark conditions.

[0147] After co-cultivation, the embryos are transferred to selective medium after which transformed isolates are obtained over the course of approximately 8 weeks. For selection of maize tissues transformed with a superbinary plasmid containing a plant expressible *pat* or *bar* selectable marker gene, an LS based medium (LS Basal medium, N6 vitamins, 1.5 mg/L 2,4-D, 0.5 gm/L MES (2-(N-morpholino)ethanesulfonic acid monohydrate; *PhytoTechnologies Labr.*), 30.0 gm/L sucrose, 6 mM L-proline, 1.0 mg/L AgNO₃, 250 mg/L cefotaxime, 2.5 gm/L Gellan gum, pH 5.7) is used with Bialaphos (Gold BioTechnology). The embryos are transferred to selection media containing 3 mg/L Bialaphos until embryogenic isolates were obtained. Recovered isolates are bulked up by transferring to fresh selection medium at 2-week intervals for regeneration and further analysis.

[0148] Those skilled in the art of maize transformation will understand that other methods of selection of transformed plants are available when other plant expressible selectable marker genes (e.g. herbicide tolerance genes) are used.

[0149] Regeneration and seed production. For regeneration, the cultures are transferred to "28" induction medium (MS salts and vitamins, 30 gm/L sucrose, 5 mg/L Benzylaminopurine, 0.25 mg/L 2, 4-D, 3 mg/L Bialaphos, 250 mg/L cefotaxime, 2.5 gm/L Gellan gum, pH 5.7) for 1 week under low-light conditions (14 µEm⁻²s⁻¹) then 1 week under high-light conditions (approximately 89 µEm⁻²s⁻¹). Tissues are subsequently transferred to "36" regeneration medium (same as induction medium except lacking plant growth regulators). When plantlets grow to 3-5 cm in length, they were transferred to glass culture tubes containing SHGA medium (Schenk and Hildebrandt salts and vitamins (1972); *PhytoTechnologies Labr.*), 1.0 gm/L myo-inositol, 10 gm/L sucrose and 2.0 gm/L Gellan gum, pH 5.8) to allow for further growth and development of the shoot and roots. Plants are transplanted to the same soil mixture as described earlier herein and grown to flowering in the greenhouse. Controlled pollinations for seed production are conducted.

EXAMPLE 10

Bioassay of transgenic maize

[0150] Bioactivity of the DIG-3 protein and variants produced in plant cells is demonstrated by conventional bioassay methods (see, for example Huang *et al.*, 2006). One is able to demonstrate efficacy, for example, by feeding various plant tissues or tissue pieces derived from a plant producing a DIG-3 toxin to target insects in a controlled feeding environment. Alternatively, protein extracts may be prepared from various plant tissues derived from a plant producing the DIG-3 toxin and incorporate the extracted proteins in an artificial diet bioassay as previously described herein. It is to be understood that the results of such feeding assays are to be compared to similarly conducted bioassays that employ appropriate control tissues from host plants that do not produce the DIG-3 protein or variants, or to other control samples.

References

[0151]

An, G., Watson, B. D., Stachel, S., Gordon, M. P., Nester, E. W. (1985) New cloning vehicles for transformation of higher plants. EMBO J. 4:277-284.

Altschul, S. F., Gish, W., Miller, W., Myers, E. W., Lipman, D. J. (1990) Basic local alignment search tool. J. Mol. Biol. 215:403-410.

Altschul, S. F., Madden, T. L., Schäffer, A. A., Zhang, J., Zhang, Z., Miller, W., Lipman, D. J. (1997) Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. Nucl. Acids Res. 25:3389-3402.

Armstrong, C. L., Green, C. E., Phillips, R. L. (1991) Development and availability of germplasm with high Typell culture formation response. Maize Genet. Coop. Newslett. 65:92-93.

Aronson, A.I., Han, E.-S., McGaughey, W., Johnson, D. (1991) The solubility of inclusion proteins from *Bacillus thuringiensis* is dependent upon protoxin composition and is a factor in toxicity to insects. Appl. Environ. Microbiol. 57:981-986.

Aronson, A. I., Geng, C., Wu, L. (1999) Aggregation of *Bacillus thuringiensis* Cry1A toxins upon binding to target insect larval midgut vesicles. *Appl. Environ. Microbiol.* 65:2503-2507.

Arvidson, H., Dunn, P. E., Strand, S., Aronson, A. I. (1989) Specificity of *Bacillus thuringiensis* for lepidopteran larvae: factors involved in vivo and in the structure of a purified toxin. *Molec. Microbiol.* 3:1533-1543.

Ausubel et al., eds. (1995) *Current Protocols in Molecular Biology*, Chapter 2 (Greene Publishing and Wiley-Interscience, New York).

Bailey, J. M., Shenov, N. R., Ronk, M., and Shively, J. E., (1992) Automated carboxy-terminal sequence analysis of peptides. *Protein Sci.* 1:68-80.

Beltz, G. A., Jacobs, K. A., Eickbush, T. H., Cherbas, P. T., Kafatos, F. C. (1983) Isolation of multigene families and determination of homologies by filter hybridization methods. In Wu, R., Grossman, L., Moldave, K. (eds.) *Methods of Enzymology*, Vol. 100 Academic Press, New York pp.266-285.

Bown, D. P., Wilkinson, H. S., Jongsma, M. A., Gatehouse, J. A. (2004) Characterisation of cysteine proteinases responsible for digestive proteolysis in guts of larval western corn rootworm (*Diabrotica virgifera*) by expression in the yeast *Pichia pastoris*. *Insect Biochem. Molec. Biol.* 34,:305-320.

Bravo, A., Gill, S. S., Soberon, M. (2007) Mode of action of *Bacillus thuringiensis* Cry and Cyt toxins and their potential for insect control. *Toxicon* 49:423-435.

Caruthers, M. H., Kierzek, R., Tang, J. Y. (1987) Synthesis of oligonucleotides using the phosphoramidite method. *Bioactive Molecules (Biophosphates Their Analogues)* 3:3-21.

Christeller, J. T., Laing, W. A., Markwick, N. P., Burgess, E. P. J. (1992) Midgut protease activities in 12 phytophagous lepidopteran larvae: dietary and protease inhibitor interactions. *Insect Biochem. Molec. Biol.* 22:735-746.

Chu, C. C., Wand, C. C., Sun, C. S., Hsu, C., Yin, K. C., Chu, C. Y., Bi, F. Y. (1975) Establishment of an efficient medium for anther culture of rice through comparative experiments on the nitrogen sources. *Scientia Sinica* 18:659-668.

Cramer, A., Cwirla, S., Stemmer, W. P. C. (1996a) Construction and evolution of antibody-phage libraries by DNA shuffling. *Nat. Med.* 2:100-103.

Cramer, A., Dawes, G., Rodriguez, E., Silver, S., Stemmer, W.P.C. (1997) Molecular evolution of an arsenate detoxification pathway by DNA shuffling. *Nat. Biotech.* 15:436-438.

Cramer, A., Whitehorn, E.A., Tate, E., Stemmer, W. P. C. (1996b) Improved green fluorescent protein by molecular evolution using DNA shuffling. *Nat. Biotech.* 14:315-319.

de Maagd, R. A., Kwa, M. S., van der Klei, H., Yamamoto, T., Schipper, B., Vlak, J. M., Stiekema, W. J., Bosch, D. (1996) Domain III substitution in *Bacillus thuringiensis* delta-endotoxin CryIA(b) results in superior toxicity for *Spo-doptera exigua* and altered membrane protein recognition. *Appl. Environ. Microbiol.* 62:1537-1543.

de Maagd, R. A., Bravo, A., Berry, C., Crickmore, N., Schnepf, E. (2003) Structure, diversity, and evolution of protein toxins from spore-forming entomopathogenic bacteria. *Annu. Rev. Genet.* 37:409-433.

Diaz-Mendoza, M., Farinos, G. P., Castanera, P., Hernandez-Crespo, P., Ortego, F. (2007) Proteolytic processing of native Cry1Ab toxin by midgut extracts and purified trypsins from the Mediterranean corn borer *Sesamia non-grioide*. *J. Insect Physiol.* 53:428-435.

Englemann, F., Geraerts, W. P. M., (1980) The proteases and the protease inhibitor in the midgut of *Leucophaea maderae*. *J. Insect Physiol.* 26:703-710.

Fraley, R. T., Rogers, S. G., Horsch, R. B. (1986) Genetic transformation in higher plants. *Crit. Rev. Plant Sci.* 4:1-46.

Gazit, E., La Rocca, P., Sansom, M. S. P., Shai, Y. (1998) The structure and organization within the membrane of the helices composing the pore-forming domain of *Bacillus thuringiensis* delta-endotoxin are consistent with an "umbrella-like" structure of the pore. *Proc. Nat. Acad. Sci. USA* 95:12289-12294.

5 Ge, A., Rivers, D., Milne, R., Dean, D. H. (1991) Functional domains of *Bacillus thuringiensis* insecticidal crystal proteins. Refinement of *Heliothis virescens* and *Trichoplusia ni* specificity domains on CryIA(c). *J. Biol. Chem.* 266:17954-17958.

10 Gillikin, J. W., Bevilacqua, S., Graham, J. S. (1992) Partial characterization of digestive tract proteinases from western corn rootworm larvae, *Diabrotica virgifera*. *Arch. Insect Biochem. Physiol.* 19:285-298.

Gomez, I., Sanchez, J., Miranda, R., Bravo, A., Soberon, M. (2002) Cadherin-like receptor binding facilitates proteolytic cleavage of helix alpha-1 in domain I and oligomer pre-pore formation of *Bacillus thuringiensis* Cry1Ab toxin. *FEBS Lett.* 513:242-246.

15 Haider, M. Z., Knowles, B. H., Ellar, D. J. (1986) Specificity of *Bacillus thuringiensis* var. *colmeri* insecticidal δ -endotoxin is determined by differential proteolytic processing of the protoxin by larval gut proteases. *Eur. J. Biochem.* 156:531-540.

20 Heckel, D. G., Gahan, L. J., Baxter, S. W., Zhao, J.-Z., Shelton, A. M., Gould, F., Tabashnik, B. E. (2007) The diversity of Bt resistance genes in species of *Lepidoptera*. *J. Invert. Pathol.* 95:192-197.

25 Hepburn, A. G., White, J., Pearson, L., Maunders, M. J., Clarke, L. E., Prescott, A. G. Blundy, K. S. (1985) The use of pNJ5000 as an intermediate vector for the genetic manipulation of *Agrobacterium* Ti-plasmids. *J. Gen. Microbiol.* 131:2961-2969.

Hoagland, D. R., Arnon, D. I. (1950) The water-culture method of growing plants without soil. *Calif. Agr. Expt. Sta. Circ.* 347.

30 Hofte, H., de Greve, H., Seurinck, J., Jansens, S., Mahillon, J., Ampe, C., Vandekerckhove, J., Vanderbruggen, H., van Montagu, M., Zabeau, M., Vaeck, M. (1986) "Structural and functional analysis of a cloned delta endotoxin of *Bacillus thuringiensis* berliner 1715." *Eur. J. Biochem.* 161:273-280.

35 Horton, R.M., Hunt, H.D., Ho, S.N., Pullen, J.K., Pease, L.R. (1989) Engineering hybrid genes without the use of restriction enzymes: gene splicing by overlap extension. *Gene* 77:61-68.

Huang, F., Rogers, L. B., Rhett, G. H. (2006) Comparative susceptibility of European corn borer, southwestern corn borer, and sugarcane borer (*Lepidoptera*: Crambidae) to Cry1Ab protein in a commercial *Bacillus thuringiensis* corn hybrid. *J. Econ. Entomol.* 99:194-202.

40 Huang, K-X., Badger, M., Haney, K., Evans, S. L. (2007) Large scale production of *Bacillus thuringiensis* PS149B1 insecticidal proteins Cry34Ab1 and Cry35Ab1 from *Pseudomonas fluorescens*. *Prot. Express. Purific.* 53:325-330.

45 Janmaat, A. F., Myers, A. H. (2003) Rapid evolution and the cost of resistance to *Bacillus thuringiensis* in greenhouse populations of cabbage loopers, *Trichoplusia ni*. *Proc. Royal Soc. London. Ser. B, Biolog. Sci.* 270:2263-2270.

Janmaat, A. F., Myers, A. H. (2005) The cost of resistance to *Bacillus thuringiensis* varies with the host plant of *Trichoplusia ni*. *Proc. Royal Soc. London. Ser. B, Biolog. Sci.* 272:1031-1038.

50 Karlin, S., Altschul, S. F. (1990) Methods for assessing the statistical significance of molecular sequence features by using general scoring schemes. *Proc. Natl. Acad. Sci. USA* 87:2264-2268.

Karlin, S., Altschul, S. F. (1993) Applications and statistics for multiple high-scoring segments in molecular sequences. *Proc. Natl. Acad. Sci. USA* 90:5873-5877.

55 Keller, G.H., Manak, M. M. (1993) *DNA Probes, Background, Applications, Procedures*. Stockton Press, New York, NY.

Knight, J. S., Broadwell, A. H., Grant, W. N., Shoemaker, C. B. (2004) A Strategy for Shuffling Numerous *Bacillus thuringiensis* Crystal Protein Domains. *J. Econ. Entomol.* 97:1805-1813.

Koiwa, H., Shade, R. E., Zhu-Salzman, K., D'Urzo, M. P., Murdock, L. L., Bressan, R. A., Hasegawa, P. M. (2000) A plant defensive cystatin (soyacystatin) targets cathepsin L-like digestive cysteine proteinases (DvCALs) in the larval midgut of western corn rootworm *Diabrotica virgifera virgifera*. *FEBS Letters* 471:67-70.

Larson, S. M., England, J. L., Desjarlais, J. R., Pande, V. S. (2002) Thoroughly sampling sequence space: Large-scale protein design of structural ensembles. *Protein Sci.* 11:2804-2813.

Lee, L.-Y., Gelvin, S. B. (2008) T-DNA binary vectors and systems. *Plant Physiol.* 146: 325-332.

Linsmaier, E.M., Skoog, F. (1965) Organic growth factor requirements of tobacco tissue. *Physiologia Plantarum* 18:100-127.

Littlefield, J. W. (1964) Selection of hybrids from matings of fibroblasts in vitro and their presumed recombinants. *Science* 145:709-710.

Meinkoth, J., Wahl, G. (1984) Hybridization of nucleic acids immobilized on solid supports. *Anal. Biochem.* 138:267-284.

Myers, E., Miller, W. (1988) Optimal alignments in linear space. *CABIOS* 4:11-17.

Naimov, S., Weemen-Hendriks, M., Dukiandjiev, S., de Maagd, R.A. (2001) *Bacillus thuringiensis* delta-endotoxin Cry1 hybrid proteins with increased activity against the Colorado Potato Beetle. *Appl. Environ. Microbiol.* 11:5328-5330.

Needleman, S. B., Wunsch, C. D. (1970) A general method applicable to the search for similarities in the amino acid sequence of two proteins. *J. Mol. Biol.* 48:443-453.

Nunez-Valdez, M.-E., Sanchez, J., Lina, L., Guereca, L., Bravo, A. (2001) Structural and functional studies of alpha-helix 5 region from *Bacillus thuringiensis* Cry1Ab delta-endotoxin. *Biochim. Biophys. Acta, Prot. Struc. Molec. Enzymol.* 1546:122-131.

Ochoa-Campuzano, C., Real, M. D., Martinez-Ramirez, A. C., Bravo, A., Rausell, C. (2007) An ADAM metalloprotease is a Cry3Aa *Bacillus thuringiensis* toxin receptor. *Biochem. Biophys. Res. Commun.* 362:437-442.

Pigott, C. R., Ellar, D. J. (2007) Role of receptors in *Bacillus thuringiensis* crystal toxin activity. *Microbiol. Molec. Biol. Rev.* 71:255-281.

Rang, C., Vachon, V., de Maagd, R. A., Villalon, M., Schwartz, J.-L., Bosch, D., Frutos, R., Laprade R. (1999) Interaction between functional domains of *Bacillus thuringiensis* insecticidal crystal proteins. *Appl. Environ. Microbiol.* 65:2918-2925.

Sambrook, J., Fritsch, E. F., Maniatis, T. (1989) *Molecular Cloning: A Laboratory Manual* (2nd ed., Cold Spring Harbor Laboratory Press, Plainview, N.Y.)

Schenk, R. U., Hildebrandt, A. C. (1972) Medium and techniques for induction and growth of monocotyledonous and dicotyledonous plant cell cultures. *Can. J. Bot.* 50:199-204

Schnepf, H. E., Tomczak, K., Ortega, J. P., Whiteley, H. R. (1990) Specificity-determining regions of a Lepidopteran-specific insecticidal protein produced by *Bacillus thuringiensis*. *J. Biol. Chem.* 265:20923-20930.

Soberon, M., Pardo-Lopez, L., Lopez, I., Gomez, I., Tabashnik, B. E., Bravo, A. (2007) Engineering modified Bt toxins to counter insect resistance. *Science* 318:1640-1642.

Squires, C. H., Retallack, D. M., Chew, L. C., Ramseier, T. M., Schneider, J. C., Talbot, H. W. (2004) Heterologous protein production in *P. fluorescens*. *Bioprocess Intern.* 2:54-59.

Stemmer, W. P. C. (1994a) DNA shuffling by random fragmentation and reassembly: in vitro recombination for molecular evolution. *Proc. Natl. Acad. Sci. USA* 91:10747-10751

Stemmer, W. P. C. (1994b) Rapid evolution of a protein in vitro by DNA shuffling. *Nature* 370: 389-391.

Stemmer, W. P. C. (1995) Searching sequence space. *Bio/Technology* 13:549-553.

Stewart, L. (2007) Gene synthesis for protein production. *Encyclopedia of Life Sciences*. John Wiley and Sons, Ltd.

Stewart, L., Burgin, A. B., (2005) Whole gene synthesis: a gene-o-matic future. *Frontiers in Drug Design and Discovery* 1:297-341.

Suggs, S.V., Miyake, T., Kawashime, E. H., Johnson, M. J., Itakura, K., R.B. Wallace, R. B. (1981) ICN-UCLA Symposium. *Dev. Biol. Using Purified Genes*, D. D. Brown [ed.], Academic Press, New York, 23:683-69

Tabashnik, B. E., Finson, N., Groeters, F. R., Moar, W. J., Johnson, M. W., Luo, K., Adang, M. J. (1994) Reversal of resistance to *Bacillus thuringiensis* in *Plutella xylostella*. *Proc. Nat. Acad. Sci. USA* 91:4120-4124.

Tabashnik, B. E., Gassmann, A. J., Crowder, D. W., Carriere, T. (2008) Insect resistance to Bt crops: evidence versus theory. *Nat. Biotech.* 26:199-202.

Taggart, R. T., Samloff, I. M. (1983) Stable antibody-producing murine hybridomas. *Science* 219:1228-1230.

Thie, N. M. R., Houseman J. G. (1990) Identification of cathepsin B, D and H in the larval midgut of Colorado potato beetle, *Leptinotarsa decemlineata* say (Coleoptera: Chrysomelidae) *Insect Biochem.* 20:313-318.

Thompson, J. D., Higgins, D. G., Gibson, T. J. (1994) CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucl. Acids Res.* 22:4673-4680.

Tijssen, P. (1993) *Laboratory Techniques in Biochemistry and Molecular Biology Hybridization with Nucleic Acid Probes*, Part I, Chapter 2. P. C. van der Vliet [ed.], (Elsevier, N.Y.)

Varshavsky, A. (1997) The N-end rule pathway of protein degradation. *Genes to Cells* 2:13-28.

Walters, F. S., Slatin, S. L., Kulesza, C. A., English, L. H. (1993) Ion channel activity of N-terminal fragments from CryIA(c) delta-endotoxin. *Biochem. Biophys. Res. Commun.* 196:921-926.

Walters, F. S., Stacy, C. M., Lee, M. K., Palekar, N., Chen, J. S. (2008) An engineered chymotrypsin/cathepsin G site in domain I renders *Bacillus thuringiensis* Cry3A active against western corn rootworm larvae. *Appl. Environ. Microbiol.* 74:367-374.

Wehrmann, A., Van Vliet, A., Opsomer, C., Botterman, J., Schulz, A. (1996) The similarities of bar and pat gene products make them equally applicable for plant engineers. *Nat. Biotechnol.* 14:1274-1278.

Weigel, D., Glazebrook, J. [eds.] (2002) *Arabidopsis: A Laboratory Manual*. Cold Spring Harbor Press, Cold Spring Harbor, NY, 354 pages.

Witkowski, J. F., Wedberg, J. L., Steffey, K. L., Sloderbeck, P. E., Siegfried, B. D., Rice, M. E., Pilcher, C. D., Onstad, D. W., Mason, C. E., Lewis, L. C., Landis, D. A., Keaster, A.J., Huang, F., Higgins, R. A., Haas, M. J., Gray, M. E., Giles, K. L., Foster, J. E., Davis, P. M., Calvin, D. D., Buschman, L. L., Bolin, P. C., Barry, B. D., Andow, D. A., Alstad, D. N. (2002) Bt corn and European Corn Borer (Ostlie, K. R., Hutchison, W. D., Hellmich, R. L. (eds)). University of Minnesota Extension Service. Publ. WW-07055.

Wolfson, J. L., Murdock, L. L. (1990) Diversity in digestive proteinase activity among insects. *J. Chem. Ecol.* 16:1089-1102.

Worley, C. K., Ling, R., Callis, J. (1998) Engineering in vivo instability of firefly luciferase and *Escherichia coli* β -

EP 2 419 441 B1

glucuronidase in higher plants using recognition elements from the ubiquitin pathway. Plant Molec. Biol. 37:337-347.

SEQUENCE LISTING

5 **[0152]**

 <110> Lira, Justin Butler, Holly Smith, Doug Narva, Ken Meade, Thomas

 <120> Bacillus thuringiensis insecticidal proteins

10 <130> 67321

 <160> 7

15 <170> PatentIn version 3.5

 <210> 1

 <211> 3771

 <212> DNA

20 <213> Bacillus thuringiensis

 <220>

 <221> misc_feature

 <222> (1)..(3)

25 <223> translation start codon specifying Met

 <400> 1

30

35

40

45

50

55

EP 2 419 441 B1

	ttgacttcaa ataggaaaaa tgagaatgaa attataaatg ctttatcgat tccaacggta	60
	togaatcctt ccacgcaa at gaatctatca ccagatgctc gtattgaaga tagcttgtgt	120
5	gtagccgagg tgaacaatat tgatccattt gttagcgcat caacagtcca aacgggtata	180
	aacatagctg gtagaatatt gggcgtatta ggtgtgccgt ttgctggaca actagctagt	240
	ttttatagtt ttcttggttg tgaattatgg cctagtggca gagatccatg ggaaattttc	300
10	ctggaacatg tagaacaact tataagacaa caagtaacag aaaatactag gaatacggct	360
	attgctcgat tagaaggtct aggaagaggc tatagatctt accagcaggc tcttgaaact	420
	tggttagata accgaaatga tgcaagatca agaagcatta ttcttgagcg ctatgttgct	480
15	ttagaacttg acattactac tgctataccg cttttcagaa tacgaaatca agaggttcca	540
	ttattaatgg tatatgctca agctgcaaat ttacacctat tattattgag ggacgcatcc	600
	cttttttgta gtgaatgggg gacggcatct tccgatgtta accaatatta ccaagaacaa	660
20	atcagatata cagaggaata ttctaaccat tgcgtacaat ggtataatac ggggctaaat	720
	aacttaagag ggacaaatgc tgaaagttgg gtacggtata atcaattccg cagagaccta	780
	acattagggg tattagatct agtggcccta ttccaagtt atgacactcg cacttatcca	840
25	atcaatacga gtgctcagtt aacaagagaa gtttatacag acgcaattgg gaccgtacat	900
	ccgagtcaag cttttgcaag tacgacttgg tttaataata atgcaccatc gttttctgcc	960
	atagaagctg ccgttatcag gcctccgcat ctacttgatt ttccagaaca acttacaatt	1020
30	tacagcacat taagtogatg gagtaacact cagtttatga atatatgggc aggtcataga	1080
	cttgaatccc gcccaatagc agggctatta aatacctcta cacaaggatc taccaatact	1140
	tctattaatc ctgtaacatt acagtttacg tctcgagaca tttataggac tgaatcattg	1200
35	gcagggctaa atatatttat aactcaacct gttaatgggg ttcttgggt tagattta	1260
40		
45		
50		
55		

EP 2 419 441 B1

	tggagaaatc cctgaattc tcttagaggt agccttctct atacgatagg gtatactgga	1320
	gttgggacgc aattacaaga ttcagaaact gaattacccc cagaaacaac agaacgacca	1380
5	aattatgaat catatagtca tagattatct catataggac tcatttcac c atctcatgtg	1440
	agagcattgg tatattcttg gacgcaccgt agtgcagatc gtacgaatac gattggacca	1500
	aatagaatta ctcaaattcc tgcagtgaag ggaagatttc tttttaatgg ctctgtaatt	1560
10	tcaggaccag gatttactgg tggagacgta gttagattga ataggaataa tggtaaatatt	1620
	caaaatagag ggtatattga agttccaatt caattcacgt cgacatctac cagatatcga	1680
	gttcgagtac gttatgcttc tgtaacctcg attgagctca atgttaattg gggcaattca	1740
15	tcaattttta cgaacacatt accagcaaca gctgcatcat tagataatct acaatcaggg	1800
	gattttgggt atgttgaaat caacaatgct tttacatccg caacaggtaa tatagtaggt	1860
	gttagaaatt ttagtgcaaa tgcagaggta ataatagaca gatttgaatt tatcccagtt	1920
20	actgcaacct tcgaggcaaa atatgattta gaaagagcac aaaaggcggg gaatgctctg	1980
	tttacttcta caaatccaag aagattgaag acagatgtga cagattatca tattgaccaa	2040
	gtgtccaatc tgggtggatg tttatcagat gaattttgct tggatgagaa gcgagaatta	2100
25	tttgagaaag tgaaatatgc gaagcgactc agtgatgaaa gaaacttact ccaagatcca	2160
	aacttcacat tcatcaatgg gcaaccaagt tttgcatcca tcgatggaca atcaaaattc	2220
	acctctatta atgagctatc taatcatgga tgggtggggc gtgcgaatgt taccattcag	2280
30	gaagggaatg acgtatttta agagaattac gtcacactac cgggtacttt taatgagtgt	2340
	tatccaaatt atttatatca aaaaatagga gagtcagaat taaaggctta tacgcgctat	2400
	caattaagag ggtatattga agatagtc aa gatctagaga tttatttaat tcgttacaa	2460
35	gcaaagcatg aaacattaaa tgttccaggt accgagtccc tatggccgct ttcagttgaa	2520
	agcccaatcg gaagggtcgg agaacc aaat cgatgcgcac cacat tttgg atggaatcct	2580
	gatctagatt gttcctgcag agatagagaa aaatgtgcgc atcattccca tcatttcact	2640
40	ttggatattg atgttgatg cacagacttg caagaggatc taggcgtgtg ggttgatttc	2700
	aagattaaga cgcaggaagg ttatgcaaga ttaggaaatc tggaaatttat cgaagagaaa	2760
	ccattaattg gagaagcact gtctcgtgtg aagagagcgg aaaaaaatg gagagacaaa	2820
45	aggga aaac tacaagtga aacaaaacga gtatatatag acgcaaaaga agctgtggat	2880
	gctttattcg tagattctca atatgataga ttacaagcag atacaaacat cggatgatt	2940
	catgcggcag atagacttgt tcatcggatc cacgaggett atcttcaga actaccttc	3000
	attccaggaa taaatgtggt gatttttgaa gaattagaaa accgtatttc tactgcattt	3060
50	tccttatatg atgcgagaaa tgtcattaaa aatggcgatt tcaataatgg attgacatgc	3120
	tggaaactga aagggcatgt agaggtagag cagctgaaca atcatcggtc ggtccttgtc	3180
	atcccggaaat gggaagcaga agtttcacaa aaggtgcgcg tctgtccagg tcgtggctat	3240
55	attcttcgtg tcacagcgta caaagaggga tatggggaag gctgcgtaac tattcatgaa	3300

EP 2 419 441 B1

gtcgataata atacagacca attgaagttt agcaactgtg agaaaggaca agtatatcca 3360
 ggtaatacga tagcatgtaa tgattataat aagaatcatg gtgcgaatgc atgtagttct 3420
 5 cgtaatcgtg gatatgacga attctatgga aacaccccag ctgattattc tgcaaataca 3480
 aaagaatacg ggggtgcgta cacttccac aatcatgcat atggcgaatc ttatgaaagt 3540
 aattcgtcca taccagctga ttatgcgccc gtttatgaag aagaagcgta tacacatgga 3600
 10 cgaagaggta attcttgtga atataacaga gggatatacac cattaccagc tggttatgtg 3660
 acagcagagt tagaataactt cccagaaacg gatacagtat ggggtgagat tggagaaacg 3720
 gaaggaacat ttatcgtgga caatgtggaa ttactcctta tggaggaata g 3771

15 <210> 2
 <211> 1256
 <212> PRT
 <213> *Bacillus thuringiensis*

20 <400> 2

	Met	Thr	Ser	Asn	Arg	Lys	Asn	Glu	Asn	Glu	Ile	Ile	Asn	Ala	Leu	Ser	
	1				5					10					15		
25	Ile	Pro	Thr	Val	Ser	Asn	Pro	Ser	Thr	Gln	Met	Asn	Leu	Ser	Pro	Asp	
				20					25					30			
	Ala	Arg	Ile	Glu	Asp	Ser	Leu	Cys	Val	Ala	Glu	Val	Asn	Asn	Ile	Asp	
			35					40					45				
30	Pro	Phe	Val	Ser	Ala	Ser	Thr	Val	Gln	Thr	Gly	Ile	Asn	Ile	Ala	Gly	
		50					55					60					
	Arg	Ile	Leu	Gly	Val	Leu	Gly	Val	Pro	Phe	Ala	Gly	Gln	Leu	Ala	Ser	
	65					70					75					80	
35	Phe	Tyr	Ser	Phe	Leu	Val	Gly	Glu	Leu	Trp	Pro	Ser	Gly	Arg	Asp	Pro	
					85					90					95		
	Trp	Glu	Ile	Phe	Leu	Glu	His	Val	Glu	Gln	Leu	Ile	Arg	Gln	Gln	Val	
				100					105					110			
40	Thr	Glu	Asn	Thr	Arg	Asn	Thr	Ala	Ile	Ala	Arg	Leu	Glu	Gly	Leu	Gly	
			115					120					125				
	Arg	Gly	Tyr	Arg	Ser	Tyr	Gln	Gln	Ala	Leu	Glu	Thr	Trp	Leu	Asp	Asn	
		130					135					140					
45	Arg	Asn	Asp	Ala	Arg	Ser	Arg	Ser	Ile	Ile	Leu	Glu	Arg	Tyr	Val	Ala	
	145					150					155					160	
	Leu	Glu	Leu	Asp	Ile	Thr	Thr	Ala	Ile	Pro	Leu	Phe	Arg	Ile	Arg	Asn	
					165					170					175		
50	Gln	Glu	Val	Pro	Leu	Leu	Met	Val	Tyr	Ala	Gln	Ala	Ala	Asn	Leu	His	
				180					185					190			
	Leu	Leu	Leu	Leu	Arg	Asp	Ala	Ser	Leu	Phe	Gly	Ser	Glu	Trp	Gly	Thr	
			195					200					205				
55	Ala	Ser	Ser	Asp	Val	Asn	Gln	Tyr	Tyr	Gln	Glu	Gln	Ile	Arg	Tyr	Thr	
		210					215					220					
	Glu	Glu	Tyr	Ser	Asn	His	Cys	Val	Gln	Trp	Tyr	Asn	Thr	Gly	Leu	Asn	
	225					230					235					240	

EP 2 419 441 B1

	Asn	Leu	Arg	Gly	Thr	Asn	Ala	Glu	Ser	Trp	Val	Arg	Tyr	Asn	Gln	Phe	
					245					250					255		
5	Arg	Arg	Asp	Leu	Thr	Leu	Gly	Val	Leu	Asp	Leu	Val	Ala	Leu	Phe	Pro	
				260					265					270			
	Ser	Tyr	Asp	Thr	Arg	Thr	Tyr	Pro	Ile	Asn	Thr	Ser	Ala	Gln	Leu	Thr	
			275					280					285				
10	Arg	Glu	Val	Tyr	Thr	Asp	Ala	Ile	Gly	Thr	Val	His	Pro	Ser	Gln	Ala	
		290					295					300					
	Phe	Ala	Ser	Thr	Thr	Trp	Phe	Asn	Asn	Asn	Ala	Pro	Ser	Phe	Ser	Ala	
	305					310					315					320	
15	Ile	Glu	Ala	Ala	Val	Ile	Arg	Pro	Pro	His	Leu	Leu	Asp	Phe	Pro	Glu	
					325					330					335		
	Gln	Leu	Thr	Ile	Tyr	Ser	Thr	Leu	Ser	Arg	Trp	Ser	Asn	Thr	Gln	Phe	
				340					345					350			
20	Met	Asn	Ile	Trp	Ala	Gly	His	Arg	Leu	Glu	Ser	Arg	Pro	Ile	Ala	Gly	
		355						360					365				
	Ser	Leu	Asn	Thr	Ser	Thr	Gln	Gly	Ser	Thr	Asn	Thr	Ser	Ile	Asn	Pro	
		370					375					380					
25	Val	Thr	Leu	Gln	Phe	Thr	Ser	Arg	Asp	Ile	Tyr	Arg	Thr	Glu	Ser	Leu	
	385					390					395					400	
	Ala	Gly	Leu	Asn	Ile	Phe	Ile	Thr	Gln	Pro	Val	Asn	Gly	Val	Pro	Trp	
				405						410					415		
30	Val	Arg	Phe	Asn	Trp	Arg	Asn	Pro	Leu	Asn	Ser	Leu	Arg	Gly	Ser	Leu	
				420					425					430			
	Leu	Tyr	Thr	Ile	Gly	Tyr	Thr	Gly	Val	Gly	Thr	Gln	Leu	Gln	Asp	Ser	
			435					440					445				
35	Glu	Thr	Glu	Leu	Pro	Pro	Glu	Thr	Thr	Glu	Arg	Pro	Asn	Tyr	Glu	Ser	
		450					455					460					
	Tyr	Ser	His	Arg	Leu	Ser	His	Ile	Gly	Leu	Ile	Ser	Ser	Ser	His	Val	
	465					470					475					480	
40	Arg	Ala	Leu	Val	Tyr	Ser	Trp	Thr	His	Arg	Ser	Ala	Asp	Arg	Thr	Asn	
				485						490					495		
	Thr	Ile	Gly	Pro	Asn	Arg	Ile	Thr	Gln	Ile	Pro	Ala	Val	Lys	Gly	Arg	
				500					505					510			
45	Phe	Leu	Phe	Asn	Gly	Ser	Val	Ile	Ser	Gly	Pro	Gly	Phe	Thr	Gly	Gly	
		515						520					525				
	Asp	Val	Val	Arg	Leu	Asn	Arg	Asn	Asn	Gly	Asn	Ile	Gln	Asn	Arg	Gly	
		530					535					540					
50	Tyr	Ile	Glu	Val	Pro	Ile	Gln	Phe	Thr	Ser	Thr	Ser	Thr	Arg	Tyr	Arg	
	545					550					555					560	
	Val	Arg	Val	Arg	Tyr	Ala	Ser	Val	Thr	Ser	Ile	Glu	Leu	Asn	Val	Asn	
				565						570				575			
55	Trp	Gly	Asn	Ser	Ser	Ile	Phe	Thr	Asn	Thr	Leu	Pro	Ala	Thr	Ala	Ala	
				580					585					590			
	Ser	Leu	Asp	Asn	Leu	Gln	Ser	Gly	Asp	Phe	Gly	Tyr	Val	Glu	Ile	Asn	
			595					600					605				

EP 2 419 441 B1

	Asn	Ala	Phe	Thr	Ser	Ala	Thr	Gly	Asn	Ile	Val	Gly	Val	Arg	Asn	Phe
	610						615					620				
5	Ser	Ala	Asn	Ala	Glu	Val	Ile	Ile	Asp	Arg	Phe	Glu	Phe	Ile	Pro	Val
	625					630					635					640
	Thr	Ala	Thr	Phe	Glu	Ala	Lys	Tyr	Asp	Leu	Glu	Arg	Ala	Gln	Lys	Ala
					645					650					655	
10	Val	Asn	Ala	Leu	Phe	Thr	Ser	Thr	Asn	Pro	Arg	Arg	Leu	Lys	Thr	Asp
				660					665					670		
	Val	Thr	Asp	Tyr	His	Ile	Asp	Gln	Val	Ser	Asn	Leu	Val	Val	Cys	Leu
			675					680					685			
15	Ser	Asp	Glu	Phe	Cys	Leu	Asp	Glu	Lys	Arg	Glu	Leu	Phe	Glu	Lys	Val
		690					695					700				
	Lys	Tyr	Ala	Lys	Arg	Leu	Ser	Asp	Glu	Arg	Asn	Leu	Leu	Gln	Asp	Pro
	705					710					715					720
20	Asn	Phe	Thr	Phe	Ile	Asn	Gly	Gln	Pro	Ser	Phe	Ala	Ser	Ile	Asp	Gly
					725					730					735	
	Gln	Ser	Asn	Phe	Thr	Ser	Ile	Asn	Glu	Leu	Ser	Asn	His	Gly	Trp	Trp
				740					745					750		
25	Gly	Ser	Ala	Asn	Val	Thr	Ile	Gln	Glu	Gly	Asn	Asp	Val	Phe	Lys	Glu
			755					760					765			
	Asn	Tyr	Val	Thr	Leu	Pro	Gly	Thr	Phe	Asn	Glu	Cys	Tyr	Pro	Asn	Tyr
		770					775					780				
30	Leu	Tyr	Gln	Lys	Ile	Gly	Glu	Ser	Glu	Leu	Lys	Ala	Tyr	Thr	Arg	Tyr
	785					790					795					800
	Gln	Leu	Arg	Gly	Tyr	Ile	Glu	Asp	Ser	Gln	Asp	Leu	Glu	Ile	Tyr	Leu
					805					810					815	
35	Ile	Arg	Tyr	Asn	Ala	Lys	His	Glu	Thr	Leu	Asn	Val	Pro	Gly	Thr	Glu
				820					825					830		
	Ser	Leu	Trp	Pro	Leu	Ser	Val	Glu	Ser	Pro	Ile	Gly	Arg	Cys	Gly	Glu
			835					840					845			
40	Pro	Asn	Arg	Cys	Ala	Pro	His	Phe	Gly	Trp	Asn	Pro	Asp	Leu	Asp	Cys
		850					855					860				
	Ser	Cys	Arg	Asp	Arg	Glu	Lys	Cys	Ala	His	His	Ser	His	His	Phe	Thr
	865					870					875					880
45	Leu	Asp	Ile	Asp	Val	Gly	Cys	Thr	Asp	Leu	Gln	Glu	Asp	Leu	Gly	Val
					885					890					895	
	Trp	Val	Val	Phe	Lys	Ile	Lys	Thr	Gln	Glu	Gly	Tyr	Ala	Arg	Leu	Gly
				900					905					910		
50	Asn	Leu	Glu	Phe	Ile	Glu	Glu	Lys	Pro	Leu	Ile	Gly	Glu	Ala	Leu	Ser
			915					920					925			
	Arg	Val	Lys	Arg	Ala	Glu	Lys	Lys	Trp	Arg	Asp	Lys	Arg	Glu	Lys	Leu
		930					935					940				
55	Gln	Val	Glu	Thr	Lys	Arg	Val	Tyr	Ile	Asp	Ala	Lys	Glu	Ala	Val	Asp
	945					950					955					960
	Ala	Leu	Phe	Val	Asp	Ser	Gln	Tyr	Asp	Arg	Leu	Gln	Ala	Asp	Thr	Asn
					965					970					975	

EP 2 419 441 B1

	Ile Gly Met	Ile His Ala Ala Asp Arg Leu Val His Arg	Ile His Glu
	980	985	990
5	Ala Tyr Leu Pro Glu Leu Pro Phe	Ile Pro Gly Ile Asn	Val Val Ile
	995	1000	1005
	Phe Glu Glu Leu Glu Asn Arg	Ile Ser Thr Ala Phe	Ser Leu Tyr
	1010	1015	1020
10	Asp Ala Arg Asn Val Ile Lys	Asn Gly Asp Phe Asn	Asn Gly Leu
	1025	1030	1035
	Thr Cys Trp Asn Val Lys Gly	His Val Glu Val Gln	Gln Leu Asn
	1040	1045	1050
15	Asn His Arg Ser Val Leu Val	Ile Pro Glu Trp Glu	Ala Glu Val
	1055	1060	1065
	Ser Gln Lys Val Arg Val Cys	Pro Gly Arg Gly Tyr	Ile Leu Arg
	1070	1075	1080
20	Val Thr Ala Tyr Lys Glu Gly	Tyr Gly Glu Gly Cys	Val Thr Ile
	1085	1090	1095
	His Glu Val Asp Asn Asn Thr	Asp Gln Leu Lys Phe	Ser Asn Cys
	1100	1105	1110
25	Glu Lys Gly Gln Val Tyr Pro	Gly Asn Thr Ile Ala	Cys Asn Asp
	1115	1120	1125
	Tyr Asn Lys Asn His Gly Ala	Asn Ala Cys Ser Ser	Arg Asn Arg
	1130	1135	1140
30	Gly Tyr Asp Glu Phe Tyr Gly	Asn Thr Pro Ala Asp	Tyr Ser Ala
	1145	1150	1155
	Asn Gln Lys Glu Tyr Gly Gly	Ala Tyr Thr Ser His	Asn His Ala
	1160	1165	1170
35	Tyr Gly Glu Ser Tyr Glu Ser	Asn Ser Ser Ile Pro	Ala Asp Tyr
	1175	1180	1185
	Ala Pro Val Tyr Glu Glu Glu	Ala Tyr Thr His Gly	Arg Arg Gly
	1190	1195	1200
40	Asn Ser Cys Glu Tyr Asn Arg	Gly Tyr Thr Pro Leu	Pro Ala Gly
	1205	1210	1215
	Tyr Val Thr Ala Glu Leu Glu	Tyr Phe Pro Glu Thr	Asp Thr Val
	1220	1225	1230
45	Trp Val Glu Ile Gly Glu Thr	Glu Gly Thr Phe Ile	Val Asp Asn
	1235	1240	1245
	Val Glu Leu Leu Leu Met Glu	Glu	
	1250	1255	

<210> 3

50 <211> 3771

<212> DNA

<213> Artificial Sequence

<220>

55 <223> Synthetic molecule

<400> 3

EP 2 419 441 B1

	atgacctcca accgtaagaa cgaaaacgaa atcatcaatg cctcagcat acccaccgtt	60
5	agcaatccct cactcagat gaacctcagc ccagatgcc gaattgagga ctccctctgc	120
	gtggcagagg tgaacaacat tgaccctttt gtctcagcct cactgttca aactggcatc	180
	aacattgctg ggaggattct tgggtgcctc ggtgtcccat tcgctggcca gctcgttct	240
10	ttctactcct ttctggttgg tgagctgtgg ccttcggac gtgacccttg ggaaatcttc	300
	cttgagcacg ttgaacagct gattaggcag caagttactg aaaacacacg taacacagcc	360
	attgccagac tggagggctc cggcagagga tatcgcagct atcagcaagc actggaaacg	420
15	tggctggata acagaaatga tgcaaggtea cgctccatca ttctogaacg ctacgtggca	480
	ctggaacttg acataacaac agccatacct ctgttcagaa tcagaaacca agaagtcctc	540
	ctgttgatgg tctatgctca agcagccaac ttgacttgc ttcttttgag ggatgcctct	600
20	ctgttcggga gcgaatgggg aaccgcttcc agcgacgtga atcagtacta tcaagagcag	660
	ataagataca cggaagagta ctcaaatac tgcgttcagt ggtacaatac tggcttgaac	720
	aatctgcgtg gcaccaatgc cgaatcttgg gtgcgttaca atcagttcag aagagacctc	780
25	acacttgggg tgctcgacct cgttgctctc tttccaagct atgacacaag gacttatccc	840
	atcaatactt cagcacagct gacgaggga gtgtacacag acgcaatagg cacagttcac	900
	cctcccaag cctttgcctc cagcacatgg ttcaacaaca atgcacctc attctcagcc	960
30	atagaagcag ctgtgattcg tccaccccat ttgtggact ttccagagca gttgacgatc	1020
	tatagcacc tttcaagggt gtcaaacacg caattcatga acatttgggc tggatcataga	1080
	cttgagtcaa ggccaatcgc tggttctctt aacacatcaa cccaaggctc caccaacacc	1140
35	tccatcaacc cagtcacct ccagttcacc agcagagaca tctatcgac agaatccctt	1200
	gctggactca acatcttcat tacacagcca gtcaatggag tcccgtaggt gaggttcaac	1260
	tggaggaatc cttgaactc acttagggga agccttcttt acaccatagg ttacacgggt	1320
40	gtgggaacgc agttgcaaga ttacagagact gaactgcctc ccgagaccac cgaacgtccc	1380
	aactatgaat catactccca ccgtctgtcc cacattgggc ttatcagctc cagccacgtg	1440
	cgtgccctcg tctactcttg gacgcacgc tccgtgata ggaccaatac cataggtccg	1500
45	aatagaatca ccagatccc agcagtgaag ggcagattct tgttcaatgg ctctgtcatt	1560
	tctggctctg gtttactgg tggtagctc gttcgttga acagaaacaa tgggaacatt	1620
	caaaaccgtg gttacatcga ggtgcccatc cagtttactt ctacatcaac acgttacaga	1680
	gttcgcgtca gatacgctc tgtgacttct attgaattga acgtgaactg gggaaacagc	1740
50	tctatcttca ctaacacact tccagccacc gcagcttcac tggacaatct tcagtctggg	1800
	gactttgggt atgtggagat caacaatgct ttacgtctg cactgggaa cattgttgggt	1860
	gtgagaaact tctctgcaa tgccgaggtg atcatagata gatattgagtt cattccagtt	1920
55	actgccacct ttgaggcaaa gtacgatctt gagagagcac agaaggctgt caacgctctg	1980
	ttcacaagca ccaaccacag acgtctgaaa actgatgtga cagactatca catcgatcaa	2040
	gttagcaact tgggtggttg cctctcagat gagttctgct tggatgagaa gagggaaactc	2100

EP 2 419 441 B1

5	tttgagaagg	ttaagtatgc	aaaacgcctt	tcagacgaaa	gaaacttgct	gcaagacccc	2160
	aactttacgt	tcataaatgg	acaacctagc	ttcgcttcca	ttgatgggtca	aagcaacttt	2220
	acttctatca	atgagttgtc	taatcacggc	tggtggggaa	gcgcaaatgt	cacaattcaa	2280
	gaaggcaacg	atgtgttcaa	agagaactac	gtgacgctgc	ctggaacatt	caatgagtgt	2340
	tatccgaact	acttgatatca	gaagattggg	gaatcagagc	tgaaggctta	cactcgctat	2400
10	cagctgcgtg	gttacatcga	ggactcccaa	gaccttgaaa	tctatctcat	ccgctacaac	2460
	gctaaacacg	agacattgaa	tgttcctgga	acagagtccc	tttggcctct	gtcagttgag	2520
	tctcctatag	ggaggtgtgg	cgaacctaac	agatgtgctc	ctcactttgg	gtggaatccc	2580
15	gatttggtatt	gctcttgtag	ggatcgcgag	aagtgcgctc	accattcaca	ccactttacg	2640
	ctggacatag	atgttggtctg	cacggacctt	caagaggatc	tcggtgtgtg	ggtcgtgttc	2700
	aagatcaaaa	ctcaagaggg	ctacgcaaga	ctcggaatac	ttgagttcat	tgaggagaag	2760
20	cctctcattg	gcgaggctct	ctctaggggtg	aagagagccg	agaagaagtg	gcgtgacaag	2820
	agggagaaac	ttcaagttga	gacgaagagg	gtgtacattg	atgctaaaga	ggctgttgat	2880
	gcactgtttg	tggattcaca	gtatgatagg	ctccaagctg	acacaaacat	tggaatgatt	2940
25	catgcagcag	atcgctcgt	gcacgcgcatc	catgaggctt	atcttcggga	actcccgttc	3000
	atccctggga	tcaatgttgt	catctttgag	gagcttgaga	accgcatatc	cacagccttt	3060
	tcctctacg	atgctagaaa	tgttatcaag	aatggcgatt	tcaacaatgg	cttgacgtgt	3120
30	tggaatgtca	aagggcattg	tgagggtcaa	cagttgaaca	accataggtc	agtcctcgtc	3180
	attccggagt	gggaggcaga	ggtagccaa	aagggtgagag	tttgccttgg	acgtggctac	3240
	attctgagag	tcactgccta	caaggagggc	tacggagaag	ggtgcgtcac	catccatgaa	3300
35	gttgataaca	atactgacca	gttgaagttt	tccaactcgc	agaaagggca	agtgatatcct	3360
	gggaatacca	ttgcatgtaa	cgattacaac	aagaatcacg	gtgctaacgc	ttgctcatct	3420
	cgcaataggg	gatatgatga	gttctatggc	aatactccag	ctgactactc	tgcaaaccag	3480
40	aaagagtatg	gaggagctta	caccagccac	aacctgcct	atgggggaatc	ttacgaatcc	3540
	aacagcagca	tcccagcaga	ctatgctccg	gtctacgagg	aggaagccta	cacacatgga	3600
	aggaggggaa	actcatgtga	gtacaataga	ggctacactc	cgttgccagc	tggtctacgtc	3660
45	actgccgaat	tggagtactt	tccggaaacc	gacactgtct	gggttgagat	aggcgaaact	3720
	gagggcacct	tcatacgttga	caacgtggaa	cttctgctta	tggaaqaata	g	3771

<210> 4

<211> 545

<212> PRT

<213> *Bacillus thuringiensis*

<400> 4

55 Leu Glu Ala Glu Ser Asp Leu Glu Arg Ala Gln Lys Ala Val Asn Ala
 1 5 10 15

 Leu Phe Thr Ser Ser Asn Gln Ile Gly Leu Lys Thr Asp Val Thr Asp
 20 25 30

EP 2 419 441 B1

	Tyr	His	Ile	Asp	Arg	Val	Ser	Asn	Leu	Val	Glu	Cys	Leu	Ser	Asp	Glu	
			35					40					45				
5	Phe	Cys	Leu	Asp	Glu	Lys	Lys	Glu	Leu	Ser	Glu	Lys	Val	Lys	His	Ala	
		50					55					60					
	Lys	Arg	Leu	Ser	Asp	Glu	Arg	Asn	Leu	Leu	Gln	Asp	Pro	Asn	Phe	Arg	
	65					70					75					80	
10	Gly	Ile	Asn	Arg	Gln	Leu	Asp	Arg	Gly	Trp	Arg	Gly	Ser	Thr	Asp	Ile	
					85					90					95		
	Thr	Ile	Gln	Gly	Gly	Asp	Asp	Val	Phe	Lys	Glu	Asn	Tyr	Val	Thr	Leu	
				100					105						110		
15	Leu	Gly	Thr	Phe	Asp	Glu	Cys	Tyr	Pro	Thr	Tyr	Leu	Tyr	Gln	Lys	Ile	
			115					120					125				
	Asp	Glu	Ser	Lys	Leu	Lys	Ala	Tyr	Thr	Arg	Tyr	Gln	Leu	Arg	Gly	Tyr	
		130					135					140					
20	Ile	Glu	Asp	Ser	Gln	Asp	Leu	Glu	Ile	Tyr	Leu	Ile	Arg	Tyr	Asn	Ala	
	145					150					155					160	
	Lys	His	Glu	Thr	Val	Asn	Val	Pro	Gly	Thr	Gly	Ser	Leu	Trp	Pro	Leu	
					165					170					175		
25	Ser	Ala	Pro	Ser	Pro	Ile	Gly	Lys	Cys	Ala	His	His	Ser	His	His	Phe	
				180					185					190			
	Ser	Leu	Asp	Ile	Asp	Val	Gly	Cys	Thr	Asp	Leu	Asn	Glu	Asp	Leu	Gly	
			195					200					205				
30	Val	Trp	Val	Ile	Phe	Lys	Ile	Lys	Thr	Gln	Asp	Gly	His	Ala	Arg	Leu	
		210					215					220					
	Gly	Asn	Leu	Glu	Phe	Leu	Glu	Glu	Lys	Pro	Leu	Val	Gly	Glu	Ala	Leu	
	225					230					235					240	
35	Ala	Arg	Val	Lys	Arg	Ala	Glu	Lys	Lys	Trp	Arg	Asp	Lys	Arg	Glu	Lys	
				245						250					255		
	Leu	Glu	Trp	Glu	Thr	Asn	Ile	Val	Tyr	Lys	Glu	Ala	Lys	Glu	Ser	Val	
				260					265					270			
40	Asp	Ala	Leu	Phe	Val	Asn	Ser	Gln	Tyr	Asp	Arg	Leu	Gln	Ala	Asp	Thr	
			275					280					285				
	Asn	Ile	Ala	Met	Ile	His	Ala	Ala	Asp	Lys	Arg	Val	His	Ser	Ile	Arg	
		290					295					300					
45	Glu	Ala	Tyr	Leu	Pro	Glu	Leu	Ser	Val	Ile	Pro	Gly	Val	Asn	Ala	Ala	
	305					310					315					320	
	Ile	Phe	Glu	Glu	Leu	Glu	Gly	Arg	Ile	Phe	Thr	Ala	Phe	Ser	Leu	Tyr	
					325					330					335		
50	Asp	Ala	Arg	Asn	Val	Ile	Lys	Asn	Gly	Asp	Phe	Asn	Asn	Gly	Leu	Ser	
				340					345					350			
	Cys	Trp	Asn	Val	Lys	Gly	His	Val	Asp	Val	Glu	Glu	Gln	Asn	Asn	His	
			355					360					365				
55	Arg	Ser	Val	Leu	Val	Val	Pro	Glu	Trp	Glu	Ala	Glu	Val	Ser	Gln	Glu	
			370				375					380					
	Val	Arg	Val	Cys	Pro	Gly	Arg	Gly	Tyr	Ile	Leu	Arg	Val	Thr	Ala	Tyr	
	385					390					395					400	

EP 2 419 441 B1

	Lys	Glu	Gly	Tyr	Gly	Glu	Gly	Cys	Val	Thr	Ile	His	Glu	Ile	Glu	Asn
					405					410					415	
5	Asn	Thr	Asp	Glu	Leu	Lys	Phe	Ser	Asn	Cys	Val	Glu	Glu	Glu	Val	Tyr
				420					425					430		
	Pro	Asn	Asn	Thr	Val	Thr	Cys	Asn	Asp	Tyr	Thr	Ala	Thr	Gln	Glu	Glu
			435					440					445			
10	Tyr	Glu	Gly	Thr	Tyr	Thr	Ser	Arg	Asn	Arg	Gly	Tyr	Asp	Gly	Ala	Tyr
		450					455					460				
	Glu	Ser	Asn	Ser	Ser	Val	Pro	Ala	Asp	Tyr	Ala	Ser	Ala	Tyr	Glu	Glu
	465					470					475					480
15	Lys	Ala	Tyr	Thr	Asp	Gly	Arg	Arg	Asp	Asn	Pro	Cys	Glu	Ser	Asn	Arg
					485					490					495	
	Gly	Tyr	Gly	Asp	Tyr	Thr	Pro	Leu	Pro	Ala	Gly	Tyr	Val	Thr	Lys	Glu
				500					505					510		
20	Leu	Glu	Tyr	Phe	Pro	Glu	Thr	Asp	Lys	Val	Trp	Ile	Glu	Ile	Gly	Glu
			515					520					525			
	Thr	Glu	Gly	Thr	Phe	Ile	Val	Asp	Ser	Val	Glu	Leu	Leu	Leu	Met	Glu
		530					535					540				
25	Glu															
	545															
30	<210> 5															
	<211> 1188															
	<212> PRT															
	<213> Artificial Sequence															
35	<220>															
	<223> Chimeric protein; DIG3 core toxin with Cry1Ab protoxin															
	<400> 5															
40	Met	Thr	Ser	Asn	Arg	Lys	Asn	Glu	Asn	Glu	Ile	Ile	Asn	Ala	Leu	Ser
	1				5					10					15	
	Ile	Pro	Thr	Val	Ser	Asn	Pro	Ser	Thr	Gln	Met	Asn	Leu	Ser	Pro	Asp
				20					25					30		
	Ala	Arg	Ile	Glu	Asp	Ser	Leu	Cys	Val	Ala	Glu	Val	Asn	Asn	Ile	Asp
			35					40					45			
45	Pro	Phe	Val	Ser	Ala	Ser	Thr	Val	Gln	Thr	Gly	Ile	Asn	Ile	Ala	Gly
		50					55					60				
	Arg	Ile	Leu	Gly	Val	Leu	Gly	Val	Pro	Phe	Ala	Gly	Gln	Leu	Ala	Ser
	65					70				75						80
50	Phe	Tyr	Ser	Phe	Leu	Val	Gly	Glu	Leu	Trp	Pro	Ser	Gly	Arg	Asp	Pro
					85					90					95	
	Trp	Glu	Ile	Phe	Leu	Glu	His	Val	Glu	Gln	Leu	Ile	Arg	Gln	Gln	Val
				100					105					110		
55	Thr	Glu	Asn	Thr	Arg	Asn	Thr	Ala	Ile	Ala	Arg	Leu	Glu	Gly	Leu	Gly
			115					120					125			
	Arg	Gly	Tyr	Arg	Ser	Tyr	Gln	Gln	Ala	Leu	Glu	Thr	Trp	Leu	Asp	Asn
		130					135					140				

EP 2 419 441 B1

	Arg	Asn	Asp	Ala	Arg	Ser	Arg	Ser	Ile	Ile	Leu	Glu	Arg	Tyr	Val	Ala	
	145					150					155					160	
5	Leu	Glu	Leu	Asp	Ile	Thr	Thr	Ala	Ile	Pro	Leu	Phe	Arg	Ile	Arg	Asn	
					165					170					175		
	Gln	Glu	Val	Pro	Leu	Leu	Met	Val	Tyr	Ala	Gln	Ala	Ala	Asn	Leu	His	
				180					185					190			
10	Leu	Leu	Leu	Leu	Arg	Asp	Ala	Ser	Leu	Phe	Gly	Ser	Glu	Trp	Gly	Thr	
			195					200					205				
	Ala	Ser	Ser	Asp	Val	Asn	Gln	Tyr	Tyr	Gln	Glu	Gln	Ile	Arg	Tyr	Thr	
		210					215					220					
15	Glu	Glu	Tyr	Ser	Asn	His	Cys	Val	Gln	Trp	Tyr	Asn	Thr	Gly	Leu	Asn	
	225					230					235					240	
	Asn	Leu	Arg	Gly	Thr	Asn	Ala	Glu	Ser	Trp	Val	Arg	Tyr	Asn	Gln	Phe	
					245					250					255		
20	Arg	Arg	Asp	Leu	Thr	Leu	Gly	Val	Leu	Asp	Leu	Val	Ala	Leu	Phe	Pro	
				260					265					270			
	Ser	Tyr	Asp	Thr	Arg	Thr	Tyr	Pro	Ile	Asn	Thr	Ser	Ala	Gln	Leu	Thr	
			275					280					285				
25	Arg	Glu	Val	Tyr	Thr	Asp	Ala	Ile	Gly	Thr	Val	His	Pro	Ser	Gln	Ala	
		290					295					300					
	Phe	Ala	Ser	Thr	Thr	Trp	Phe	Asn	Asn	Asn	Ala	Pro	Ser	Phe	Ser	Ala	
	305					310					315					320	
30	Ile	Glu	Ala	Ala	Val	Ile	Arg	Pro	Pro	His	Leu	Leu	Asp	Phe	Pro	Glu	
					325					330					335		
	Gln	Leu	Thr	Ile	Tyr	Ser	Thr	Leu	Ser	Arg	Trp	Ser	Asn	Thr	Gln	Phe	
				340					345					350			
35	Met	Asn	Ile	Trp	Ala	Gly	His	Arg	Leu	Glu	Ser	Arg	Pro	Ile	Ala	Gly	
		355						360					365				
	Ser	Leu	Asn	Thr	Ser	Thr	Gln	Gly	Ser	Thr	Asn	Thr	Ser	Ile	Asn	Pro	
		370					375					380					
40	Val	Thr	Leu	Gln	Phe	Thr	Ser	Arg	Asp	Ile	Tyr	Arg	Thr	Glu	Ser	Leu	
	385					390					395					400	
	Ala	Gly	Leu	Asn	Ile	Phe	Ile	Thr	Gln	Pro	Val	Asn	Gly	Val	Pro	Trp	
				405						410					415		
45	Val	Arg	Phe	Asn	Trp	Arg	Asn	Pro	Leu	Asn	Ser	Leu	Arg	Gly	Ser	Leu	
				420					425					430			
	Leu	Tyr	Thr	Ile	Gly	Tyr	Thr	Gly	Val	Gly	Thr	Gln	Leu	Gln	Asp	Ser	
			435					440					445				
50	Glu	Thr	Glu	Leu	Pro	Pro	Glu	Thr	Thr	Glu	Arg	Pro	Asn	Tyr	Glu	Ser	
		450					455					460					
	Tyr	Ser	His	Arg	Leu	Ser	His	Ile	Gly	Leu	Ile	Ser	Ser	Ser	His	Val	
	465					470					475					480	
55	Arg	Ala	Leu	Val	Tyr	Ser	Trp	Thr	His	Arg	Ser	Ala	Asp	Arg	Thr	Asn	
					485					490					495		
	Thr	Ile	Gly	Pro	Asn	Arg	Ile	Thr	Gln	Ile	Pro	Ala	Val	Lys	Gly	Arg	
				500					505					510			

EP 2 419 441 B1

	Phe	Leu	Phe	Asn	Gly	Ser	Val	Ile	Ser	Gly	Pro	Gly	Phe	Thr	Gly	Gly	
		515						520					525				
5	Asp	Val	Val	Arg	Leu	Asn	Arg	Asn	Asn	Gly	Asn	Ile	Gln	Asn	Arg	Gly	
		530					535					540					
	Tyr	Ile	Glu	Val	Pro	Ile	Gln	Phe	Thr	Ser	Thr	Ser	Thr	Arg	Tyr	Arg	
	545					550					555					560	
10	Val	Arg	Val	Arg	Tyr	Ala	Ser	Val	Thr	Ser	Ile	Glu	Leu	Asn	Val	Asn	
					565					570					575		
	Trp	Gly	Asn	Ser	Ser	Ile	Phe	Thr	Asn	Thr	Leu	Pro	Ala	Thr	Ala	Ala	
				580					585					590			
15	Ser	Leu	Asp	Asn	Leu	Gln	Ser	Gly	Asp	Phe	Gly	Tyr	Val	Glu	Ile	Asn	
			595					600					605				
	Asn	Ala	Phe	Thr	Ser	Ala	Thr	Gly	Asn	Ile	Val	Gly	Val	Arg	Asn	Phe	
		610					615					620					
20	Ser	Ala	Asn	Ala	Glu	Val	Ile	Ile	Asp	Arg	Phe	Glu	Phe	Ile	Pro	Val	
	625					630					635					640	
	Thr	Ala	Thr	Leu	Glu	Ala	Glu	Ser	Asp	Leu	Glu	Arg	Ala	Gln	Lys	Ala	
					645					650					655		
25	Val	Asn	Ala	Leu	Phe	Thr	Ser	Ser	Asn	Gln	Ile	Gly	Leu	Lys	Thr	Asp	
				660					665					670			
	Val	Thr	Asp	Tyr	His	Ile	Asp	Arg	Val	Ser	Asn	Leu	Val	Glu	Cys	Leu	
			675					680					685				
30	Ser	Asp	Glu	Phe	Cys	Leu	Asp	Glu	Lys	Lys	Glu	Leu	Ser	Glu	Lys	Val	
		690					695					700					
	Lys	His	Ala	Lys	Arg	Leu	Ser	Asp	Glu	Arg	Asn	Leu	Leu	Gln	Asp	Pro	
	705					710					715					720	
35	Asn	Phe	Arg	Gly	Ile	Asn	Arg	Gln	Leu	Asp	Arg	Gly	Trp	Arg	Gly	Ser	
				725						730					735		
	Thr	Asp	Ile	Thr	Ile	Gln	Gly	Gly	Asp	Asp	Val	Phe	Lys	Glu	Asn	Tyr	
				740					745					750			
40	Val	Thr	Leu	Leu	Gly	Thr	Phe	Asp	Glu	Cys	Tyr	Pro	Thr	Tyr	Leu	Tyr	
			755					760					765				
	Gln	Lys	Ile	Asp	Glu	Ser	Lys	Leu	Lys	Ala	Tyr	Thr	Arg	Tyr	Gln	Leu	
		770					775					780					
45	Arg	Gly	Tyr	Ile	Glu	Asp	Ser	Gln	Asp	Leu	Glu	Ile	Tyr	Leu	Ile	Arg	
	785					790					795					800	
	Tyr	Asn	Ala	Lys	His	Glu	Thr	Val	Asn	Val	Pro	Gly	Thr	Gly	Ser	Leu	
					805					810					815		
50	Trp	Pro	Leu	Ser	Ala	Pro	Ser	Pro	Ile	Gly	Lys	Cys	Ala	His	His	Ser	
				820					825					830			
	His	His	Phe	Ser	Leu	Asp	Ile	Asp	Val	Gly	Cys	Thr	Asp	Leu	Asn	Glu	
			835					840					845				
	Asp	Leu	Gly	Val	Trp	Val	Ile	Phe	Lys	Ile	Lys	Thr	Gln	Asp	Gly	His	
		850					855					860					
55	Ala	Arg	Leu	Gly	Asn	Leu	Glu	Phe	Leu	Glu	Glu	Lys	Pro	Leu	Val	Gly	
						870					875					880	

EP 2 419 441 B1

	Glu	Ala	Leu	Ala	Arg	Val	Lys	Arg	Ala	Glu	Lys	Lys	Trp	Arg	Asp	Lys	
					885					890					895		
5	Arg	Glu	Lys	Leu	Glu	Trp	Glu	Thr	Asn	Ile	Val	Tyr	Lys	Glu	Ala	Lys	
				900					905					910			
	Glu	Ser	Val	Asp	Ala	Leu	Phe	Val	Asn	Ser	Gln	Tyr	Asp	Arg	Leu	Gln	
			915					920					925				
10	Ala	Asp	Thr	Asn	Ile	Ala	Met	Ile	His	Ala	Ala	Asp	Lys	Arg	Val	His	
		930					935					940					
	Ser	Ile	Arg	Glu	Ala	Tyr	Leu	Pro	Glu	Leu	Ser	Val	Ile	Pro	Gly	Val	
	945					950					955					960	
15	Asn	Ala	Ala	Ile	Phe	Glu	Glu	Leu	Glu	Gly	Arg	Ile	Phe	Thr	Ala	Phe	
				965						970					975		
	Ser	Leu	Tyr	Asp	Ala	Arg	Asn	Val	Ile	Lys	Asn	Gly	Asp	Phe	Asn	Asn	
				980				985						990			
20	Gly	Leu	Ser	Cys	Trp	Asn	Val	Lys	Gly	His	Val	Asp	Val	Glu	Glu	Gln	
		995						1000					1005				
	Asn	Asn	His	Arg	Ser	Val	Leu	Val	Val	Pro	Glu	Trp	Glu	Ala	Glu		
		1010					1015					1020					
25	Val	Ser	Gln	Glu	Val	Arg	Val	Cys	Pro	Gly	Arg	Gly	Tyr	Ile	Leu		
		1025					1030					1035					
	Arg	Val	Thr	Ala	Tyr	Lys	Glu	Gly	Tyr	Gly	Glu	Gly	Cys	Val	Thr		
		1040					1045					1050					
30	Ile	His	Glu	Ile	Glu	Asn	Asn	Thr	Asp	Glu	Leu	Lys	Phe	Ser	Asn		
		1055					1060					1065					
	Cys	Val	Glu	Glu	Glu	Val	Tyr	Pro	Asn	Asn	Thr	Val	Thr	Cys	Asn		
		1070					1075					1080					
35	Asp	Tyr	Thr	Ala	Thr	Gln	Glu	Glu	Tyr	Glu	Gly	Thr	Tyr	Thr	Ser		
		1085					1090					1095					
	Arg	Asn	Arg	Gly	Tyr	Asp	Gly	Ala	Tyr	Glu	Ser	Asn	Ser	Ser	Val		
		1100					1105					1110					
40	Pro	Ala	Asp	Tyr	Ala	Ser	Ala	Tyr	Glu	Glu	Lys	Ala	Tyr	Thr	Asp		
		1115					1120					1125					
	Gly	Arg	Arg	Asp	Asn	Pro	Cys	Glu	Ser	Asn	Arg	Gly	Tyr	Gly	Asp		
		1130					1135					1140					
45	Tyr	Thr	Pro	Leu	Pro	Ala	Gly	Tyr	Val	Thr	Lys	Glu	Leu	Glu	Tyr		
		1145					1150					1155					
	Phe	Pro	Glu	Thr	Asp	Lys	Val	Trp	Ile	Glu	Ile	Gly	Glu	Thr	Glu		
		1160					1165					1170					
50	Gly	Thr	Phe	Ile	Val	Asp	Ser	Val	Glu	Leu	Leu	Leu	Met	Glu	Glu		
		1175					1180					1185					

<210> 6

<211> 1635

<212> DNA

<213> Artificial Sequence

<220>

<223> Synthetic molecule

EP 2 419 441 B1

<400> 6

	ctcgaggctg aatctgatct cgaaagggca cagaaagctg taaacgcatt gtttacaagt	60
5	tctaatacaaa tcggactcaa aaccgatgtt acggactatc acatagatag ggtttcta	120
	cttgtggaat gtctttcaga tgagttttgt ttagatgaga agaaagaact ttcagaaaag	180
	gtcaagcacg ccaaaagact gtccgatgaa aggaatctcc ttcaagaccc aaactttcgt	240
10	ggaatcaata ggcagctcga cagaggttgg agaggagca cagatatac cattcaagga	300
	ggagatgacg ttttcaaaga gaactatgtc accttgtag gcaccttga tgagtgtat	360
	ccaacttatac tgtatcagaa gattgatgaa tccaagctga aggttacac aagatatacag	420
15	ctcagaggat acatcgagga ctcccaagat ttggagatat acttgattcg ttacaatgca	480
	aaacatgaga ccgtgaatgt tcctggtagt ggaagtctct ggccactgtc tgctccgtca	540
	cctattggga aatgtgccca tcaactccac catttctcat tggacataga cgttggtgc	600
20	acagatttga atgaagattt ggggtgttgg gtcatcttca agatcaaac tcaagacgga	660
	cacgctcgtt taggaaactt agagtttctt gaagagaagc ccttggttgg ggaggcactt	720
	gccagagtaa agagagctga aaagaagtgg agagataaga gggagaaact tgagtgggag	780
25	actaacattg tgtacaagga agccaaagaa agcgtggatg ctcttttcgt gaactctcag	840
	tatgataggt tacaagcaga caccaacata gcaatgatac atgcagctga caaaagagtc	900
	cattctattc gtgaggctta cttgccagaa cttagtgtga ttcccggtgt caacgctgcc	960
30	attttcgagg aattggaagg aagaatcttt acggctttca gcctctatga cgtaggaat	1020
	gttatcaaga atgggtgattt caacaatggc ctctcatgtt ggaatgtgaa aggtcatgtt	1080
	gatgtagagg agcaaaacaa tcaccgtagc gtgctggttg tcccagaatg ggaagccgaa	1140
35	gtaagccaag aagttagagt ttgccctgga agaggctaca ttctgcgtgt caccgcttac	1200
	aaagaaggat atggcgaagg gtgcgtgact attcatgaga ttgagaacaa tactgacgaa	1260
	cttaagtttt caaactgcgt cgaggaggaa gtgtatccta acaacacagt gacttgtaat	1320
40	gactatacag caacgcaaga ggaatacagag gggacataca ccagtcgtaa tcgtggttat	1380
	gatggtgctt atgaaagcaa ttcacccgtt ccagctgact atgccagtgc ctacgaagag	1440
	aaggcttaca cggatggcag aagagataac ccatgtgagt ccaacagagg ttatggtgat	1500
45	tacactcttc ttccagctgg ttacgtgact aaagagttag agtactttcc ggagactgat	1560
	aaggtttgga ttgaaatcgg agagacagaa gggacattca tagtagattc agttgagctt	1620
	cttctcatgg aagaa	1635

<210> 7

50 <211> 1635

<212> DNA

<213> Artificial Sequence

<220>

55 <223> Synthetic molecule

<400> 7

EP 2 419 441 B1

	ctcgaggctg aatcggtatct tgaaggggca cagaaggcag tcaacgctct cttcaccagc	60
5	tcaaatacaga ttggccttaa gaccgatgtt actgactatc atatcgacag agttttctaac	120
	cttgctcgagt gctctctccga cgagttctgt ctcgacgaaa agaaggaact ctccgagaaa	180
	gtgaagcacg cgaaacgcct ctcggtatgaa cggaacttgc tgcaagatcc gaacttcaga	240
10	ggcatcaatc gccagttgga tagaggctgg aggggatcaa ccgacataac cattcaaggt	300
	ggggatgatg tgttcaagga aaactacgtg acattgctgg gcaccttcga cgagtgtctat	360
	cccacgtatc tctatcagaa gattgacgag tccaagctca aagcctacac acgctatcag	420
15	ctcagaggct acattgagga ctctcaagac ctcgaaatct acttgatcag atacaacgcc	480
	aagcacgaga cgggtgaacgt ccctgggact gggtcactgt ggccactgtc ggcacctcgc	540
	ccaatcgga agtgcgctca ccacagccac cacttctccc ttgacataga tgttgggtgt	600
20	acggacttga atgaggatct ggggtgtgtg gtgatcttta agatcaagac ccaagatggt	660
	catgcgaggc ttggcaacct tgagttcctt gaagagaagc ctttggtcgg agaggcactg	720
	gctcgctga agagggctga gaagaaatgg agggacaaga gggagaaact ggagtgggag	780
25	accaacatag tgtacaagga ggccaaggag tcagtggacg cactgtttgt caattcccag	840
	tatgataggc tccaagcgga cacgaacatc gccatgatcc atgcagcgga caagagggtt	900
	cactccataa gggaggccta tcttccggag ctgtcagtga ttctctgggt caacgcagcc	960
	atctttgagg aattggaagg gaggatcttc accgctttct ctctgtacga cgctcggaac	1020
30	gtcatcaaga atggtgattt caacaatgga ctcagctgct ggaacgtgaa agggcatgtc	1080
	gatgttgaag aacagaacaa tcaccgcagc gtgctgggtg ttccggagtg ggaagccgag	1140
	gtctcacaag aagtcagagt gtgccctggg aggggttaca tcttgccgggt cacagcctac	1200
35	aaggaagggt atggcggaagg ctgtgtcacg atccatgaga tcgaaaacaa cacagacgag	1260
	ctgaagtttt ccaactgtgt tgaggaggag gtctatccta acaatactgt tacgtgcaac	1320
	gactacacag ccactcaaga ggagtacgag ggcacttaca cctctcgcaa cagaggctac	1380
40	gacggtgctt acgagtcaaa cagctccgtg ccagcggact acgcctcggc ttacgaagag	1440
	aaggcgtaca ccgacggtcg gagggataac ccgtgcgaga gcaatagagg ctatggcgac	1500
	tacactcctc tcccagctgg ctacgtgacc aaggagttgg agtactttcc ggagacagac	1560
45	aaagtctgga ttgagattgg agagacagaa ggcacgttca tcgtggactc tgttgaactc	1620
	ttgctgatgg aggag	1635

50 Claims

1. An isolated polypeptide comprising a core toxin segment selected from the group consisting of

- 55 (a) a polypeptide comprising the amino acid sequence of residues 113 to 643 of SEQ ID NO: 2;
 (b) a polypeptide comprising an amino acid sequence having at least 90% sequence identity to the amino acid sequence of residues 113 to 643 of SEQ ID NO:2;
 (c) a polypeptide comprising an amino acid sequence of residues 113 to 643 of SEQ ID NO: 2 with up to 20 amino acid substitutions, deletions, or modifications that do not adversely affect expression or activity of the

toxin encoded by SEQ ID NO: 2; or an insecticidally active fragment thereof.

2. The isolated polypeptide of claim 1 comprising a core toxin segment selected from the group consisting of
 - (a) a polypeptide comprising the amino acid sequence of residues 73 to 643 or 1 to 643 of SEQ ID NO: 2;
 - (b) a polypeptide comprising an amino acid sequence having at least 90% sequence identity to the amino acid sequence of residues 73 to 643 or 1 to 643 of SEQ ID NO:2;
 - (c) a polypeptide comprising an amino acid sequence of residues 73 to 643 or 1 to 643 of SEQ ID NO: 2 with up to 20 amino acid substitutions, deletions, or modifications that do not adversely affect expression or activity of the toxin encoded by SEQ ID NO: 2;or an insecticidally active fragment thereof.
3. The polypeptide of claim 1 wherein said polypeptide comprises an amino acid sequence of SEQ ID NO: 2 or SEQ ID NO:5.
4. A transgenic plant comprising the polypeptide of claim 1.
5. A method for controlling a pest population comprising contacting said population with a pesticidally effective amount of the polypeptide of claim 1.
6. An isolated nucleic acid that encodes a polypeptide of claim 1.
7. The isolated nucleic acid of claim 6 wherein said nucleic acid comprises SEQ ID NO: 1 or SEQ ID NO:3.
8. A DNA construct comprising the nucleotide sequence of claim 7 operably linked to a promoter that is not derived from *Bacillus thuringiensis* and is capable of driving expression in a plant.
9. A transgenic plant that comprises the DNA construct of claim 8 stably incorporated into its genome.
10. The transgenic plant of claim 9 further comprising a polynucleotide that encodes a Cry1F protein.
11. The transgenic plant of claim 10 wherein said plant is corn or a vegetable plant.
12. A maize plant cell comprising the DNA construct of claim 8 and optionally a polynucleotide that encodes a Cry1F protein.
13. A vegetable plant cell comprising the DNA construct of claim 8 and optionally a polynucleotide that encodes a Cry1A protein.
14. A method for protecting a plant from a pest comprising introducing into said plant the construct of claim 8.
15. The polypeptide of claim 1 having activity against European corn borer.
16. The method of claim 5, wherein said pest population is a population of European corn borer that is resistant to Cry1F, or said pest population is a population of diamond back moth that is resistant to a Cry1A.
17. A method of controlling a lepidopteran, said method comprising providing a polypeptide of claim 1 to said lepidopteran.
18. The method of claim 17, wherein said lepidopteran is a European corn borer, and said method further comprises providing a Cry1F to said lepidopteran.
19. The method of claim 17, wherein said lepidopteran is a diamond back moth, and said method further comprises providing a Cry1Ab and/or a Cry1Ac to said lepidopteran.
20. The method of claim 17, wherein said lepidopteran is a European corn borer that is resistant to Cry1F.
21. The method of claim 17, wherein said lepidopteran is a diamond back moth that is resistant to a Cry1A.

22. A transgenic seed of a plant of any of claims 4, and 9-11.

Patentansprüche

1. Isoliertes Polypeptid, umfassend ein Kerntoxinsegment, das ausgewählt ist aus der Gruppe bestehend aus:

- (a) einem Polypeptid, das die Aminosäuresequenz der Reste 113 bis 643 der SEQ ID Nr. 2 umfasst;
- (b) einem Polypeptid, das eine Aminosäuresequenz mit mindestens 90%-iger Sequenzidentität mit der Aminosäuresequenz der Reste 113 bis 643 der SEQ ID Nr. 2 umfasst;
- (c) einem Polypeptid, das eine Aminosäuresequenz der Reste 113 bis 643 der SEQ ID Nr. 2 umfasst, mit bis zu 20 Aminosäuresubstitutionen, Deletionen oder Modifikationen, welche die Expression oder Aktivität des durch SEQ ID Nr. 2 codierten Toxins nicht negative beeinflussen; oder einem insektizid aktiven Fragment davon.

2. Isoliertes Polypeptid gemäß Anspruch 1, umfassend ein Kerntoxinsegment, das ausgewählt ist aus der Gruppe bestehend aus:

- (a) einem Polypeptid, das die Aminosäuresequenz der Reste 73 bis 643 oder 1 bis 643 der SEQ ID Nr. 2 umfasst;
- (b) einem Polypeptid, das eine Aminosäuresequenz mit mindestens 90%-iger Sequenzidentität mit der Aminosäuresequenz der Reste 73 bis 643 oder 1 bis 643 der SEQ ID Nr. 2 umfasst;
- (c) einem Polypeptid, das eine Aminosäuresequenz der Reste 73 bis 643 oder 1 bis 643 der SEQ ID Nr. 2 umfasst, mit bis zu 20 Aminosäuresubstitutionen, Deletionen oder Modifikationen, welche die Expression oder Aktivität des durch SEQ ID Nr. 2 codierten Toxins nicht negative beeinflussen; oder einem insektizid aktiven Fragment davon.

3. Polypeptid gemäß Anspruch 1, wobei das Polypeptid eine Aminosäuresequenz der SEQ ID Nr. 2 oder der SEQ ID Nr. 5 umfasst.

4. Transgene Pflanze, die das Polypeptid gemäß Anspruch 1 umfasst.

5. Verfahren zur Kontrolle einer Schädlingspopulation, umfassend das Inkontaktbringen der Population mit einer pestizid wirkenden Menge des Polypeptids gemäß Anspruch 1.

6. Isolierte Nukleinsäure, die für ein Polypeptid gemäß Anspruch 1 codiert.

7. Isolierte Nukleinsäure gemäß Anspruch 6, wobei die Nukleinsäure SEQ ID Nr. 1 oder SEQ ID Nr. 3 umfasst.

8. DNA-Konstrukt, umfassend die Nukleotidsequenz nach Anspruch 7, funktionsfähig verknüpft mit einem Promotor, der nicht von *Bacillus thuringiensis* abstammt und die Expression in einer Pflanze steuern kann.

9. Transgene Pflanze, die das DNA-Konstrukt gemäß Anspruch 8 umfasst, stabil in deren Genom inkorporiert.

10. Transgene Pflanze gemäß Anspruch 9, ferner umfassend ein Polynukleotid, das für ein Cry1 F-Protein codiert.

11. Transgene Pflanze gemäß Anspruch 10, wobei die Pflanze Mais oder eine Gemüsepflanze ist.

12. Maispflanzenzelle, umfassend das DNA-Konstrukt gemäß Anspruch 8 und optional ein Polynukleotid, das für ein Cry1 F-Protein codiert.

13. Gemüsepflanzenzelle, umfassend das DNA-Konstrukt gemäß Anspruch 8 und optional ein Polynukleotid, das für ein Cry1 F-Protein codiert.

14. Verfahren zum Schützen einer Pflanze vor einem Schädling, umfassend das Einführen des Konstrukts gemäß Anspruch 8 in die Pflanze.

15. Polypeptid gemäß Anspruch 1 mit Aktivität gegen Maiszünsler.

16. Verfahren nach Anspruch 5, wobei die Schädlingspopulation eine gegen Cry1 F resistente Maiszünslerpopulation

ist, oder wobei die Schädlingspopulation eine gegen Cry1A resistente Kohlmottenpopulation ist.

17. Verfahren zur Kontrolle von Schmetterlingen, wobei das Verfahren das Bereitstellen eines Polypeptids gemäß Anspruch 1 an den Schmetterling umfasst.
18. Verfahren gemäß Anspruch 17, wobei der Schmetterling ein Maiszünsler ist, und wobei das Verfahren ferner das Bereitstellen eines Cry1 F an den Schmetterling umfasst.
19. Verfahren gemäß Anspruch 17, wobei der Schmetterling eine Kohlmotte ist, und wobei das Verfahren ferner das Bereitstellen eines Cry1 Ab und/oder eines Cry1 Ac an den Schmetterling umfasst.
20. Verfahren gemäß Anspruch 17, wobei der Schmetterling ein gegen Cry1 F resistenter Maiszünsler ist.
21. Verfahren gemäß Anspruch 17, wobei der Schmetterling eine gegen Cry1 A resistente Kohlmotte ist.
22. Transgener Samen einer Pflanze gemäß einem der Ansprüche 4 und 9 bis 11.

Revendications

1. Polypeptide isolé comprenant un segment de toxine de base choisi dans le groupe constitué par
 - (a) un polypeptide comprenant la séquence d'acides aminés des résidus 113 à 643 de la Séquence N° 2 ;
 - (b) un polypeptide comprenant une séquence d'acides aminés ayant une identité de séquence d'au moins 90 % avec la séquence d'acides aminés des résidus 113 à 643 de la Séquence N° 2 ;
 - (c) un polypeptide comprenant une séquence d'acides aminés des résidus 113 à 643 de la Séquence N° 2, comportant jusqu'à 20 remplacements, délétions ou modifications d'acides aminés qui n'ont pas d'incidence défavorable sur l'expression ou l'activité de la toxine codée par la Séquence N° 2 ;ou fragment à activité insecticide d'un tel polypeptide.
2. Polypeptide isolé selon la revendication 1, comprenant un segment de toxine de base choisi dans le groupe constitué par
 - (a) un polypeptide comprenant la séquence d'acides aminés des résidus 73 à 643 ou 1 à 643 de la Séquence N° 2 ;
 - (b) un polypeptide comprenant une séquence d'acides aminés ayant une identité de séquence d'au moins 90 % avec la séquence d'acides aminés des résidus 73 à 643 ou 1 à 643 de la Séquence N° 2 ;
 - (c) un polypeptide comprenant une séquence d'acides aminés des résidus 73 à 643 ou 1 à 643 de la Séquence N° 2, comportant jusqu'à 20 remplacements, délétions ou modifications d'acides aminés, qui n'ont pas d'incidence défavorable sur l'expression ou l'activité de la toxine codée par la Séquence N° 2 ;ou fragment à activité insecticide d'un tel polypeptide.
3. Polypeptide selon la revendication 1, ledit polypeptide comprenant une séquence d'acides aminés de la Séquence N° 2 ou de la Séquence N° 5.
4. Plante transgénique comprenant le polypeptide de la revendication 1.
5. Procédé pour combattre une population de ravageurs, comprenant la mise en contact de ladite population avec une quantité efficace en tant que pesticide du polypeptide selon la revendication 1.
6. Acide nucléique isolé qui code un polypeptide selon la revendication 1.
7. Acide nucléique isolé selon la revendication 6, ledit acide nucléique comprenant la Séquence N° 1 ou la Séquence N° 3.
8. Produit de construction d'ADN comprenant la séquence de nucléotides de la revendication 7 liée de façon opérationnelle à un promoteur qui n'est pas issu de *Bacillus thuringiensis* et est capable de diriger l'expression chez une plante.

9. Plante transgénique qui comprend le produit de construction d'ADN selon la revendication 8, incorporé de façon stable dans son génome.
10. Plante transgénique selon la revendication 9, comprenant en outre un polynucléotide qui code une protéine Cry1F.
11. Plante transgénique selon la revendication 10, ladite plante étant le maïs ou une plante maraîchère.
12. Cellule de plant de maïs comprenant le produit de construction d'ADN selon la revendication 8 et en option un polynucléotide qui code une protéine Cry1F.
13. Cellule de plante maraîchère comprenant le produit de construction d'ADN selon la revendication 8 et en option un polynucléotide qui code une protéine Cry1A.
14. Procédé pour la protection d'une plante contre un ravageur, comprenant l'introduction du produit de construction selon la revendication 8 dans ladite plante.
15. Polypeptide selon la revendication 1, ayant une activité contre la pyrale du maïs.
16. Procédé selon la revendication 5, dans lequel ladite population de ravageurs est une population de pyrales du maïs qui est résistante à Cry1F, ou ladite population de ravageurs est une population de teignes des crucifères qui est résistante à une protéine Cry1A.
17. Procédé de lutte contre un lépidoptère, ledit procédé comprenant l'apport d'un polypeptide selon la revendication 1 audit lépidoptère.
18. Procédé selon la revendication 17, dans lequel ledit lépidoptère est une pyrale du maïs, et ledit procédé comprend en outre l'apport d'une protéine Cry1F audit lépidoptère.
19. Procédé selon la revendication 17, dans lequel ledit lépidoptère est une teigne des crucifères, et ledit procédé comprend en outre l'apport d'une protéine Cry1Ab et/ou d'une protéine Cry1Ac audit lépidoptère.
20. Procédé selon la revendication 17, dans lequel ledit lépidoptère est une pyrale du maïs qui est résistante à Cry1F.
21. Procédé selon la revendication 17, dans lequel ledit lépidoptère est une teigne des crucifères qui est résistante à une protéine Cry1A.
22. Semence transgénique d'une plante selon l'une quelconques des revendications 4, et 9-11.

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- US 61170189 A [0001]
- US 6090931 A [0032]
- WO 9101087 A [0032]
- WO 9506730 A [0032]
- WO 1998022595 A [0032]
- US 7230167 B [0041]
- US 6046053 A, Bailey [0045]
- US 7058515 B [0052]
- US 7482119 B2 [0055]
- US 5380831 A [0055] [0077]
- US 4945050 A [0072]
- US 5141131 A [0072]
- US 5177010 A [0072]
- US 5104310 A [0072]
- EP 0131624 B1 [0072]
- EP 120516 A [0072] [0074]
- EP 159418 B1 [0072]
- EP 176112 A [0072]
- US 5149645 A [0072]
- US 5469976 A [0072]
- US 5464763 A [0072]
- US 4940838 A [0072]
- US 4693976 A [0072]
- EP 116718 A [0072]
- EP 290799 A [0072]
- EP 320500 A [0072]
- EP 604662 A [0072]
- EP 627752 A [0072]
- EP 0267159 A [0072]
- EP 0292435 A [0072]
- US 5231019 A [0072]
- US 5463174 A [0072]
- US 4762785 A [0072]
- US 5004863 A [0072]
- US 5159135 A [0072]
- US 5302523 A [0072]
- US 5464765 A [0072]
- WO 8706614 A [0072]
- US 5472869 A [0072]
- US 5384253 A [0072]
- WO 9209696 A [0072]
- WO 9321335 A [0072]
- WO 2007053482 A2 [0080]
- WO 2005107437 A2 [0080]
- US 6268132 B [0091]
- US 20080193974 A [0115]
- US 20060008877 A, Squires [0115] [0116]
- US 20080058262 A [0115]

Non-patent literature cited in the description

- AN, G. ; WATSON, B. D. ; STACHEL, S. ; GORDON, M. P. ; NESTER, E. W. New cloning vehicles for transformation of higher plants. *EMBO J.*, 1985, vol. 4, 277-284 [0151]
- ALTSCHUL, S. F. ; GISH, W. ; MILLER, W. ; MYERS, E. W. ; LIPMAN, D. J. Basic local alignment search tool. *J. Mol. Biol.*, 1990, vol. 215, 403-410 [0151]
- ALTSCHUL, S. F. ; MADDEN, T. L. ; SCHÄFFER, A. A. ; ZHANG, J. ; ZHANG, Z. ; MILLER, W. ; LIPMAN, D. J. Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucl. Acids Res.*, 1997, vol. 25, 3389-3402 [0151]
- ARMSTRONG, C. L. ; GREEN, C. E. ; PHILLIPS, R. L. Development and availability of germplasm with high Typell culture formation response. *Maize Genet. Coop. Newslett.*, 1991, vol. 65, 92-93 [0151]
- ARONSON, A.I. ; HAN, E.-S. ; MCGAUGHEY, W. ; JOHNSON, D. The solubility of inclusion proteins from *Bacillus thuringiensis* is dependent upon protoxin composition and is a factor in toxicity to insects. *Appl. Environ. Microbiol.*, 1991, vol. 57, 981-986 [0151]
- ARONSON, A. I. ; GENG, C. ; WU, L. Aggregation of *Bacillus thuringiensis* Cry1A toxins upon binding to target insect larval midgut vesicles. *Appl. Environ. Microbiol.*, 1999, vol. 65, 2503-2507 [0151]
- ARVIDSON, H. ; DUNN, P. E. ; STRAND, S. ; ARONSON, A. I. Specificity of *Bacillus thuringiensis* for lepidopteran larvae: factors involved in vivo and in the structure of a purified toxin. *Molec. Microbiol.*, 1989, vol. 3, 1533-1543 [0151]
- Current Protocols in Molecular Biology. Greene Publishing and Wiley-Interscience, 1995 [0151]
- BAILEY, J. M. ; SHENOV, N. R. ; RONK, M. ; SHIVELY, J. E. Automated carboxy-terminal sequence analysis of peptides. *Protein Sci.*, 1992, vol. 1, 68-80 [0151]

- Isolation of multigene families and determination of homologies by filter hybridization methods. **BELTZ, G.A. ; JACOBS, K. A. ; EICKBUSH, T. H. ; CHERBAS, P. T. ; KAFATOS, F. C.** Methods of Enzymology. Academic Press, 1983, vol. 100, 266-285 [0151]
- **BOWN, D. P. ; WILKINSON, H. S. ; JONGSMA, M. A. ; GATEHOUSE, J. A.** Characterisation of cysteine proteinases responsible for digestive proteolysis in guts of larval western corn rootworm (*Diabrotica virgifera*) by expression in the yeast *Pichia pastoris*. *Insect Biochem. Molec. Biol.*, 2004, vol. 34, 305-320 [0151]
- **BRAVO, A. ; GILL, S. S. ; SOBERON, M.** Mode of action of *Bacillus thuringiensis* Cry and Cyt toxins and their potential for insect control. *Toxicon*, 2007, vol. 49, 423-435 [0151]
- **CARUTHERS, M. H. ; KIERZEK, R. ; TANG, J. Y.** Synthesis of oligonucleotides using the phosphoramidite method. *Bioactive Molecules (Biophosphates Their Analogues)*, 1987, vol. 3, 3-21 [0151]
- **CHRISTELLER, J. T. ; LAING, W. A. ; MARKWICK, N. P. ; BURGESS, E. P. J.** Midgut protease activities in 12 phytophagous lepidopteran larvae: dietary and protease inhibitor interactions. *Insect Biochem. Molec. Biol.*, 1992, vol. 22, 735-746 [0151]
- **CHU, C. C. ; WAND, C. C. ; SUN, C. S. ; HSU, C. ; YIN, K. C. ; CHU, C. Y. ; BI, F. Y.** Establishment of an efficient medium for anther culture of rice through comparative experiments on the nitrogen sources. *Scientia Sinica*, 1975, vol. 18, 659-668 [0151]
- **CRAMERI, A. ; CWIRLA, S. ; STEMMER, W. P. C.** Construction and evolution of antibody-phage libraries by DNA shuffling. *Nat. Med.*, 1996, vol. 2, 100-103 [0151]
- **CRAMERI, A. ; DAWES, G. ; RODRIGUEZ, E. ; SILVER, S. ; STEMMER, W.P.C.** Molecular evolution of an arsenate detoxification pathway by DNA shuffling. *Nat. Biotech.*, 1997, vol. 15, 436-438 [0151]
- **CRAMERI, A. ; WHITEHOM, E.A. ; TATE, E. ; STEMMER, W. P. C.** Improved green fluorescent protein by molecular evolution using DNA shuffling. *Nat. Biotech.*, 1996, vol. 14, 315-319 [0151]
- **DE MAAGD, R. A. ; KWA, M. S. ; VAN DER KLEI, H. ; YAMAMOTO, T. ; SCHIPPER, B. ; VLAK, J. M. ; STIEKEMA, W. J. ; BOSCH, D.** Domain III substitution in *Bacillus thuringiensis* delta-endotoxin CryIA(b) results in superior toxicity for *Spodoptera exigua* and altered membrane protein recognition. *Appl. Environ. Microbiol.*, 1996, vol. 62, 1537-1543 [0151]
- **DE MAAGD, R. A. ; BRAVO, A. ; BERRY, C. ; CRICKMORE, N. ; SCHNEPPF, E.** Structure, diversity, and evolution of protein toxins from spore-forming entomopathogenic bacteria. *Annu. Rev. Genet.*, 2003, vol. 37, 409-433 [0151]
- **DIAZ-MENDOZA, M. ; FARINOS, G. P. ; CAS-TANERA, P. ; HERNANDEZ-CRESPO, P. ; ORTEGO, F.** Proteolytic processing of native Cry1Ab toxin by midgut extracts and purified trypsins from the Mediterranean corn borer *Sesamia nonagrioides*. *J. Insect Physiol.*, 2007, vol. 53, 428-435 [0151]
- **ENGLEMANN, F. ; GERAERTS, W. P. M.** The proteases and the protease inhibitor in the midgut of *Leucophaea maderae*. *J. Insect Physiol.*, 1980, vol. 26, 703-710 [0151]
- **FRALEY, R. T. ; ROGERS, S. G. ; HORSCH, R. B.** Genetic transformation in higher plants. *Crit. Rev. Plant Sci.*, 1986, vol. 4, 1-46 [0151]
- **GAZIT, E. ; LA ROCCA, P. ; SANSOM, M. S. P. ; SHAI, Y.** The structure and organization within the membrane of the helices composing the pore-forming domain of *Bacillus thuringiensis* delta-endotoxin are consistent with an "umbrella-like" structure of the pore. *Proc. Nat. Acad. Sci. USA*, 1998, vol. 95, 12289-12294 [0151]
- **GE, A. ; RIVERS, D. ; MILNE, R. ; DEAN, D. H.** Functional domains of *Bacillus thuringiensis* insecticidal crystal proteins. Refinement of *Heliothis virescens* and *Trichoplusia ni* specificity domains on CryIA(c). *J. Biol. Chem.*, 1991, vol. 266, 17954-17958 [0151]
- **GILLIKIN, J. W. ; BEVILACQUA, S. ; GRAHAM, J. S.** Partial characterization of digestive tract proteinases from western corn rootworm larvae, *Diabrotica virgifera*. *Arch. Insect Biochem. Physiol.*, 1992, vol. 19, 285-298 [0151]
- **GOMEZ, I. ; SANCHEZ, J. ; MIRANDA, R. ; BRAVO, A. ; SOBERON, M.** Cadherin-like receptor binding facilitates proteolytic cleavage of helix alpha-1 in domain I and oligomer pre-pore formation of *Bacillus thuringiensis* Cry1Ab toxin. *FEBS Lett.*, 2002, vol. 513, 242-246 [0151]
- **HAIDER, M. Z. ; KNOWLES, B. H. ; ELLAR, D. J.** Specificity of *Bacillus thuringiensis* var. *colmeri* insecticidal δ -endotoxin is determined by differential proteolytic processing of the protoxin by larval gut proteases. *Eur. J. Biochem.*, 1986, vol. 156, 531-540 [0151]
- **HECKEL, D. G. ; GAHAN, L. J. ; BAXTER, S. W. ; ZHAO, J.-Z. ; SHELTON, A. M. ; GOULD, F. ; TABASHNIK, B. E.** The diversity of Bt resistance genes in species of *Lepidoptera*. *J. Invert. Pathol.*, 2007, vol. 95, 192-197 [0151]
- **HEPBURN, A. G. ; WHITE, J. ; PEARSON, L. ; MAUNDERS, M. J. ; CLARKE, L. E. ; PRESCOTT, A. G. ; BLUNDY, K. S.** The use of pNJ5000 as an intermediate vector for the genetic manipulation of *Agrobacterium Ti*-plasmids. *J. Gen. Microbiol.*, 1985, vol. 131, 2961-2969 [0151]
- **HOAGLAND, D. R. ; ARNON, D. I.** The water-culture method of growing plants without soil. *Calif. Agr. Expt. Sta. Circ.*, 1950, 347 [0151]

- **HOFTE, H. ; DE GREVE, H. ; SEURINCK, J. ; JANSSENS, S. ; MAHILLON, J. ; AMPE, C. ; VANDEKERCKHOVE, J. ; VANDERBRUGGEN, H. ; VAN MONTAGU, M. ; ZABEAU, M.** Structural and functional analysis of a cloned delta endotoxin of *Bacillus thuringiensis* berliner 1715. *Eur. J. Biochem.*, 1986, vol. 161, 273-280 [0151]
- **HORTON, R.M. ; HUNT, H.D. ; HO, S.N. ; PULLEN, J.K. ; PEASE, L.R.** Engineering hybrid genes without the use of restriction enzymes: gene splicing by overlap extension. *Gene*, 1989, vol. 77, 61-68 [0151]
- **HUANG, F. ; ROGERS, L. B. ; RHETT, G. H.** Comparative susceptibility of European corn borer, southwestern corn borer, and sugarcane borer (Lepidoptera: Crambidae) to Cry1Ab protein in a commercial *Bacillus thuringiensis* corn hybrid. *J. Econ. Entomol.*, 2006, vol. 99, 194-202 [0151]
- **HUANG, K.-X. ; BADGER, M. ; HANEY, K. ; EVANS, S. L.** Large scale production of *Bacillus thuringiensis* PS149B1 insecticidal proteins Cry34Ab1 and Cry35Ab1 from *Pseudomonas fluorescens*. *Prot. Express. Purific.*, 2007, vol. 53, 325-330 [0151]
- **JANMAAT, A. F. ; MYERS, A. H.** Rapid evolution and the cost of resistance to *Bacillus thuringiensis* in greenhouse populations of cabbage loopers, *Trichoplusia ni*. *Proc. Royal Soc. London. Ser. B, Biolog. Sci.*, 2003, vol. 270, 2263-2270 [0151]
- **JANMAAT, A. F. ; MYERS, A. H.** The cost of resistance to *Bacillus thuringiensis* varies with the host plant of *Trichoplusia ni*. *Proc. Royal Soc. London. Ser. B, Biolog. Sci.*, 2005, vol. 272, 1031-1038 [0151]
- **KARLIN, S. ; ALTSCHUL, S. F.** Methods for assessing the statistical significance of molecular sequence features by using general scoring schemes. *Proc. Natl. Acad. Sci. USA*, 1990, vol. 87, 2264-2268 [0151]
- **KARLIN, S. ; ALTSCHUL, S. F.** Applications and statistics for multiple high-scoring segments in molecular sequences. *Proc. Natl. Acad. Sci. USA*, 1993, vol. 90, 5873-5877 [0151]
- **KELLER, G.H. ; MANAK, M. M.** DNA Probes, Background, Applications, Procedures. Stockton Press, 1993 [0151]
- **KNIGHT, J. S. ; BROADWELL, A. H. ; GRANT, W. N. ; SHOEMAKER, C. B.** A Strategy for Shuffling Numerous *Bacillus thuringiensis* Crystal Protein Domains. *J. Econ. Entomol.*, 2004, vol. 97, 1805-1813 [0151]
- **KOIWA, H. ; SHADE, R. E. ; ZHU-SALZMAN, K. ; D'URZO, M. P. ; MURDOCK, L. L. ; BRESSAN, R. A. ; HASEGAWA, P. M.** A plant defensive cystatin (soyacystatin) targets cathepsin L-like digestive cysteine proteinases (DvCALs) in the larval midgut of western corn rootworm *Diabrotica virgifera virgifera*. *FEBS Letters*, 2000, vol. 471, 67-70 [0151]
- **LARSON, S. M. ; ENGLAND, J. L. ; DESJARLAIS, J. R. ; PANDE, V. S.** Thoroughly sampling sequence space: Large-scale protein design of structural ensembles. *Protein Sci.*, 2002, vol. 11, 2804-2813 [0151]
- **LEE, L.-Y. ; GELVIN, S. B.** T-DNA binary vectors and systems. *Plant Physiol.*, 2008, vol. 146, 325-332 [0151]
- **LINSMAIER, E.M. ; SKOOG, F.** Organic growth factor requirements of tobacco tissue. *Physiologia Plantarum*, 1965, vol. 18, 100-127 [0151]
- **LITTLEFIELD, J. W.** Selection of hybrids from matings of fibroblasts in vitro and their presumed recombinants. *Science*, 1964, vol. 145, 709-710 [0151]
- **MEINKOTH, J. ; WAHL, G.** Hybridization of nucleic acids immobilized on solid supports. *Anal. Biochem.*, 1984, vol. 138, 267-284 [0151]
- **MYERS, E. ; MILLER, W.** Optimal alignments in linear space. *CABIOS*, 1988, vol. 4, 11-17 [0151]
- **NAIMOV, S. ; WEEMEN-HENDRIKS, M. ; DUKI-ANDJIEV, S. ; DE MAAGD, R.A.** *Bacillus thuringiensis* delta-endotoxin Cry1 hybrid proteins with increased activity against the Colorado Potato Beetle. *Appl. Environ. Microbiol.*, 2001, vol. 11, 5328-5330 [0151]
- **NEEDLEMAN, S. B. ; WUNSCH, C. D.** A general method applicable to the search for similarities in the amino acid sequence of two proteins. *J. Mol. Biol.*, 1970, vol. 48, 443-453 [0151]
- **NUNEZ-VALDEZ, M.-E. ; SANCHEZ, J. ; LINA, L. ; GUERECAL, L. ; BRAVO, A.** Structural and functional studies of alpha-helix 5 region from *Bacillus thuringiensis* Cry1Ab delta-endotoxin. *Biochim. Biophys. Acta, Prot. Struc. Molec. Enzymol.*, 2001, vol. 1546, 122-131 [0151]
- **OCHOA-CAMPUZANO, C. ; REAL, M. D. ; MARTINEZ-RAMIREZ, A. C. ; BRAVO, A. ; RAUSELL, C.** An ADAM metalloprotease is a Cry3Aa *Bacillus thuringiensis* toxin receptor. *Biochem. Biophys. Res. Commun.*, 2007, vol. 362, 437-442 [0151]
- **PIGOTT, C. R. ; ELLAR, D. J.** Role of receptors in *Bacillus thuringiensis* crystal toxin activity. *Microbiol. Molec. Biol. Rev.*, 2007, vol. 71, 255-281 [0151]
- **RANG, C. ; VACHON, V. ; DE MAAGD, R. A. ; VILLALON, M. ; SCHWARTZ, J.-L. ; BOSCH, D. ; FRUTOS, R. ; LAPRADE R.** Interaction between functional domains of *Bacillus thuringiensis* insecticidal crystal proteins. *Appl. Environ. Microbiol.*, 1999, vol. 65, 2918-2925 [0151]
- **SAMBROOK, J. ; FRITSCH, E. F. ; MANIATIS, T.** Molecular Cloning: A Laboratory Manual. Cold Spring Harbor Laboratory Press, 1989 [0151]
- **SCHENK, R. U. ; HILDEBRANDT, A. C.** Medium and techniques for induction and growth of monocotyledonous and dicotyledonous plant cell cultures. *Can. J. Bot.*, 1972, vol. 50, 199-204 [0151]

- **SCHNEPF, H. E. ; TOMCZAK, K. ; ORTEGA, J. P. ; WHITELEY, H. R.** Specificity-determining regions of a Lepidopteran-specific insecticidal protein produced by *Bacillus thuringiensis*. *J. Biol. Chem.*, 1990, vol. 265, 20923-20930 [0151]
- **SOBERON, M. ; PARDO-LOPEZ, L. ; LOPEZ, I. ; GOMEZ, I. ; TABASHNIK, B. E. ; BRAVO, A.** Engineering modified Bt toxins to counter insect resistance. *Science*, 2007, vol. 318, 1640-1642 [0151]
- **SQUIRES, C. H. ; RETALLACK, D. M. ; CHEW, L. C. ; RAMSEIER, T. M. ; SCHNEIDER, J. C. ; TALBOT, H. W.** Heterologous protein production in *P. fluorescens*. *Bioprocess Intern.*, 2004, vol. 2, 54-59 [0151]
- **STEMMER, W. P. C.** DNA shuffling by random fragmentation and reassembly: in vitro recombination for molecular evolution. *Proc. Natl. Acad. Sci. USA*, 1994, vol. 91, 10747-10751 [0151]
- **STEMMER, W. P. C.** Rapid evolution of a protein in vitro by DNA shuffling. *Nature*, 1994, vol. 370, 389-391 [0151]
- **STEMMER, W. P. C.** Searching sequence space. *BioTechnology*, 1995, vol. 13, 549-553 [0151]
- Gene synthesis for protein production. **STEWART, L.** Encyclopedia of Life Sciences. John Wiley and Sons, Ltd, 2007 [0151]
- **STEWART, L. ; BURGIN, A. B.** Whole gene synthesis: a gene-o-matic future. *Frontiers in Drug Design and Discovery*, 2005, vol. 1, 297-341 [0151]
- ICN-UCLA Symposium. **SUGGS, S.V. ; MIYAKE, T. ; KAWASHIME, E. H. ; JOHNSON, M. J. ; ITAKURA, K. ; R.B. WALLACE.** Dev. Biol. Using Purified Genes. Academic Press, 1981, vol. 23, 683-69 [0151]
- **TABASHNIK, B. E. ; FINSON, N. ; GROETERS, F. R. ; MOAR, W. J. ; JOHNSON, M. W. ; LUO, K. ; ADANG, M. J.** Reversal of resistance to *Bacillus thuringiensis* in *Plutella xylostella*. *Proc. Nat. Acad. Sci. USA*, 1994, vol. 91, 4120-4124 [0151]
- **TABASHNIK, B. E. ; GASSMANN, A. J. ; CROWDER, D. W. ; CARRIERE, T.** Insect resistance to Bt crops: evidence versus theory. *Nat. Biotech.*, 2008, vol. 26, 199-202 [0151]
- **TAGGART, R. T. ; SAMLOFF, I. M.** Stable antibody-producing murine hybridomas. *Science*, 1983, vol. 219, 1228-1230 [0151]
- **THIE, N. M. R. ; HOUSEMAN J. G.** Identification of cathepsin B, D and H in the larval midgut of Colorado potato beetle, *Leptinotarsa decemlineata* say (Coleoptera: Chrysomelidae). *Insect Biochem.*, 1990, vol. 20, 313-318 [0151]
- **THOMPSON, J. D. ; HIGGINS, D. G. ; GIBSON, T. J.** CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucl. Acids Res.*, 1994, vol. 22, 4673-4680 [0151]
- **TIJJSSEN, P.** Laboratory Techniques in Biochemistry and Molecular Biology Hybridization with Nucleic Acid Probes. Elsevier, 1993 [0151]
- **VARSHAVSKY, A.** The N-end rule pathway of protein degradation. *Genes to Cells*, 1997, vol. 2, 13-28 [0151]
- **WALTERS, F. S. ; SLATIN, S. L. ; KULESZA, C. A. ; ENGLISH, L. H.** Ion channel activity of N-terminal fragments from CryIA(c) delta-endotoxin. *Biochem. Biophys. Res. Commun.*, 1993, vol. 196, 921-926 [0151]
- **WALTERS, F. S. ; STACY, C. M. ; LEE, M. K. ; PALEKAR, N. ; CHEN, J. S.** An engineered chymotrypsin/cathepsin G site in domain I renders *Bacillus thuringiensis* Cry3A active against western corn rootworm larvae. *Appl. Environ. Microbiol.*, 2008, vol. 74, 367-374 [0151]
- **WEHRMANN, A. ; VAN VLIET, A. ; OPSOMER, C. ; BOTTERMAN, J. ; SCHULZ, A.** The similarities of bar and pat gene products make them equally applicable for plant engineers. *Nat. Biotechnol.*, 1996, vol. 14, 1274-1278 [0151]
- *Arabidopsis: A Laboratory Manual.* Cold Spring Harbor Press, 2002, 354 [0151]
- **WITKOWSKI, J. F. ; WEDBERG, J. L. ; STEFFEY, K. L. ; SLODERBECK, P. E. ; SIEGFRIED, B. D. ; RICE, M. E. ; PILCHER, C. D. ; ONSTAD, D. W. ; MASON, C. E. ; LEWIS, L. C.** Bt corn and European Corn Borer. University of Minnesota Extension Service, 2002 [0151]
- **WOLFSON, J. L. ; MURDOCK, L. L.** Diversity in digestive proteinase activity among insects. *J. Chem. Ecol.*, 1990, vol. 16, 1089-1102 [0151]
- **WORLEY, C. K. ; LING, R. ; CALLIS, J.** Engineering in vivo instability of firefly luciferase and *Escherichia coli* β -glucuronidase in higher plants using recognition elements from the ubiquitin pathway. *Plant Molec. Biol.*, 1998, vol. 37, 337-347 [0151]

INSZEKTICID DIG-3 CRY-TOXINOK

Szabadalmi igénypontok

1. Izolált polipeptid, amely a következők alkotta csoportból választott központi toxinszegmenst tartalmaz:

(a) polipeptid, amely SEQ ID NO: 2 szekvencia 113-643. aminosavai alkotta aminosav-szekvenciát tartalmaz,

(b) polipeptid, amely SEQ ID NO: 2 szekvencia 113-643. aminosavai alkotta aminosav-szekvenciával legalább 90% szekvenciaazonosságú aminosav-szekvenciát tartalmaz,

(c) polipeptid, amely SEQ ID NO: 2 szekvencia 113-643. aminosavai alkotta aminosav-szekvenciát tartalmaz, amelyben legfeljebb 20, a SEQ ID NO: 2 szekvencia által kódolt toxin expresszióját vagy aktivitását kedvezőtlenül nem befolyásoló aminosav-szubsztitúció, -delécio vagy -módosítás van; vagy inszekticid aktivitású fragmense.

2. Az 1. igénypont szerinti izolált polipeptid, amely a következők alkotta csoportból választott központi toxinszegmenst tartalmaz:

(a) polipeptid, amely SEQ ID NO: 2 szekvencia 73-643. vagy 1-643. aminosavai alkotta aminosav-szekvenciát tartalmaz,

(b) polipeptid, amely SEQ ID NO: 2 szekvencia 73-643. vagy 1-643. aminosavai alkotta aminosav-szekvenciával legalább 90% szekvenciaazonosságú aminosav-szekvenciát tartalmaz,

(c) polipeptid, amely SEQ ID NO: 2 szekvencia 73-643. vagy 1-643 aminosavai alkotta aminosav-szekvenciát tartalmaz, amelyben legfeljebb 20, a SEQ ID NO: 2 szekvencia által kódolt toxin expresszióját vagy aktivitását kedvezőtlenül nem befolyásoló aminosav-szubsztitúció, -delécio vagy -módosítás van, vagy inszekticid aktivitású fragmense.

3. Az 1. igénypont szerinti polipeptid, ahol a polipeptid SEQ ID NO:2 vagy SEQ ID NO:5 szerinti aminosav-szekvenciát tartalmaz.

4. Transzgenikus növény, amely 1. igénypont szerinti polipeptidet tartalmaz.

5. Eljárás kártevő-populáció irtására, amely magában foglalja a populáció érintkeztetését az 1. igénypont szerinti polipeptid peszticidként hatékony mennyiségével.

6. Izolált nukleinsav, amely 1. igénypont szerinti polipeptidet kódol.

7. A 6. igénypont szerinti izolált nukleinsav, amely SEQ ID NO: 1 vagy SEQ ID NO:3 szerinti szekvenciát tartalmaz.

8. DNS-konstrukció, amely 7. igénypont szerinti nukleotid-szekvenciát tartalmaz nem *Bacillus thuringiensis*-ből származó, növényben expresszió irányítására képes promóterhez kapcsolva működőképesen.

9. Transzgenikus növény, amely 8. igénypont szerinti DNS-konstrukciót tartalmaz genomjába stabilan beépítve.

10. A 9. igénypont szerinti transzgenikus növény, amely CryIF-fehérjét kódoló polinukleotidot is tartalmaz.

11. A 10. igénypont szerinti növény, amely kukorica vagy zöldségnövény.

12. Kukoricánövény sejtje, amely a 8. igénypont szerinti DNS-konstrukciót tartalmazza, és adott esetben CryIF-fehérjét kódoló polinukleotidot tartalmaz.

13. Zöldségnövény sejtje, amely a 8. igénypont szerinti DNS-konstrukciót tartalmazza, és adott esetben CryIA-fehérjét kódoló polinukleotidot tartalmaz.

14. Eljárás növények kártevővel szembeni védelmére, amely magában foglalja a 8. igénypont szerinti konstrukció bevitelét a növénybe.

15. Az 1. igénypont szerinti polipeptid, amely aktív kukoricamoly ellen.

16. Az 5. igénypont szerinti eljárás, ahol a kártevő-populáció CryIF-fel szemben rezisztens kukoricamoly-populáció vagy CryIA-val szemben rezisztens káposztamoly-populáció.

17. Eljárás lepkék irtására, amely magában foglalja 1. igénypont szerinti polipeptid biztosítását a lepkének.

18. A 17. igénypont szerinti eljárás, ahol a lepke kukoricamoly, és az eljárás magában foglalja CryIF biztosítását is a lepkének.

19. A 17. igénypont szerinti eljárás, ahol a lepke káposztamoly, és az eljárás magában foglalja CryIAb és/vagy CryIAc biztosítását is a lepkének.

20. A 17. igénypont szerinti eljárás, ahol a lepke CryIF-fel szemben rezisztens kukoricamoly.

21. A 17. igénypont szerinti eljárás, ahol a lepke CryIA-val szemben rezisztens káposztamoly.

22. A 4. és 9-11. igénypont bármelyike szerinti növény transzgenikus magja.