



(51) International Patent Classification:  
G01N 33/50 (2006.01)

(21) International Application Number:  
PCT/US2013/064116

(22) International Filing Date:  
9 October 2013 (09.10.2013)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
61/711,500 9 October 2012 (09.10.2012) US

(71) Applicant: THE PROCTER & GAMBLE COMPANY  
[US/US]; One Procter & Gamble Plaza, Cincinnati, Ohio 45202 (US).

(72) Inventors: OBLONG, John, Erich; One Procter & Gamble Plaza, Cincinnati, Ohio 45202 (US). ROVITO,

Holly, Ann; One Procter & Gamble Plaza, Cincinnati, Ohio 45202 (US).

(74) Agent: GUFFEY, Timothy B.; c/o The Procter & Gamble Company, Global Patent Services, 299 East 6th Street, Sycamore Building, 4th Floor, Cincinnati, Ohio 45202 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

[Continued on next page]

(54) Title: METHOD OF IDENTIFYING OR EVALUATING BENEFICIAL ACTIVES AND COMPOSITIONS CONTAINING THE SAME




FIG. 1A




FIG. 1B




FIG. 1C

(57) Abstract: A method of identifying or evaluating skin-care actives that improve the metabolism of a skin cell. The method includes contacting cells with a stressor and a test agent and determining a response of the cells to the stressor and test agent, based on the change in metabolic indicators associated with glycolysis and/or oxidative phosphorylation. The test agent may be identified as a skin-care active when the oxidative phosphorylation and/or glycolysis response corresponds to an improvement in cellular metabolism.

WO 2014/059008 A1



**(84) Designated States** (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK,

SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

**Published:**

— *with international search report (Art. 21(3))*

METHOD OF IDENTIFYING OR EVALUATING BENEFICIAL ACTIVES AND  
COMPOSITIONS CONTAINING THE SAME

FIELD OF THE INVENTION

The present invention is directed, generally, to a method of identifying or evaluating  
5 beneficial active ingredients for use in personal care compositions. More specifically, the present  
invention is directed to a method of identifying or evaluating actives that combat the effects of  
oxidative stress on skin cells.

BACKGROUND OF THE INVENTION

A fundamental basis for life is the need and ability of an organism to generate energy. In  
10 humans, food is taken in and converted into chemical compounds such as adenosine triphosphate  
("ATP") and nicotinamide adenine dinucleotide ("NAD"), which store the energy used by the  
cells of the body to perform the biological processes that sustain life. The metabolic pathways of  
the cells that convert the useful components of food (e.g., carbohydrates, fats and proteins) into  
usable energy are complex and may be affected by a variety of factors in ways that are not  
15 completely understood. Mammalian skin cells are no exception. Skin cells are known to include  
a variety of different kinds of cells that functions together in a dynamic, complex relationship to  
maintain the health of the skin. For example, keratinocytes proliferate and differentiate to  
provide continuous skin turnover. Melanocytes are known to provide melanin synthesis for skin  
pigmentation. And fibroblasts are known for synthesizing the extracellular matrix and collagen,  
20 which helps maintain the skin's thickness and elasticity. Similarly, other cells found in or around  
the skin or other bodily organs, such as myocytes, stem cells, sebocytes, neurocytes, and  
adipocytes all require energy derived from complex metabolic pathways, which can be  
undesirably impacted by a variety of different factors.

There is a growing awareness of the impact of various stressors on cellular bioenergetics  
25 and the impacts on cell aging, as well as other diseases (e.g., cancer, neurodegenerative diseases,  
diabetes, and cardiovascular disorders). One theory underlying some of these effects of altered  
metabolism in disease states is the Free Radical Theory of aging. Namely, that exposure of  
mammalian cells to reactive oxygen species ("ROS") causes damage to cellular structures and  
organelles such as the mitochondria. ROS are highly reactive molecules that contain oxygen  
30 (e.g., oxygen ions and peroxides). ROS are formed within cells as a natural byproduct of the  
normal metabolism of oxygen and play a role in cell signaling and homeostasis. However, when

a cell is exposed to a stressor such as heat or UV radiation, ROS levels can increase, and in some instances, dramatically.

As the damage caused by ROS accumulates over time, it causes more and more oxidative stress at the cellular level that ultimately may lead to tissue damage and/or organ dysfunction.

5 One effect of oxidative stress on cells is a diminished capacity of cellular bioenergetics, which can lead to reduced levels of ATP and/or NAD. This may be particularly problematic for human skin because the oxidative stress on human skin cells may manifest as visible signs of aging. Further, environmental stressors such as ultraviolet radiation (“UV”) and pollutants (e.g., cigarette smoke, car exhaust, ozone) can lead to heightened levels of ROS production. Over  
10 time, this may result in noticeable changes in the skin’s structure and morphology (e.g., “photodamage”) and, to a more extreme degree, skin carcinomas.

Human skin cells defend against ROS by using redox regulators such as glutathione and NAD as well as various enzymes that can neutralize ROS. However, these defenses can be overwhelmed by the elevated spike from stressor-induced ROS, leading to not just acute but also  
15 chronic alterations in energy homeostasis and metabolism efficiencies, causing overall cellular dysfunction. To complicate matters, the variety of cells types associated with human skin and the complexity of the metabolic pathways of these cells makes it difficult to identify suitable compounds to help combat the anti-aging effect associated with exposure to a particular oxidative stressor or combination of stressors. Certain oxidative stressors affect different types  
20 of cells and/or metabolic pathways differently. This makes it difficult to select a suitable skin care active or combination of actives (whose affect may also vary depending on type of cell or metabolic pathway) that combat the undesirable affects of a particular stressor or stressors. Since at least some consumers desire a skin care composition that combats the undesirable aging effects associated with oxidative stress, there is a need for a method of identifying suitable skin care  
25 compounds that can be used to tailor a skin care composition to help reduce, prevent and/or reverse the undesirable oxidative stress associated with various stressors, especially common environmental stressors.

As an initial step in finding suitable skin-care actives that combat metabolic effects of oxidative stress, a method capable of detecting the changes in cellular metabolism caused by  
30 stressors and actives must be identified. Various methods are known for evaluating the energy making processes of cells. For example, Clark-type electrode probes are known for measuring oxygen consumption. The Clark electrode provides kinetic information (i.e., rates of response) but introduces artifact (i.e., some undesirable and/or extraneous factor that influences the results

of a test) by its continuous consumption of oxygen, presenting a decreasing oxygen pressure to the cells or isolated mitochondria in the measurement chamber. Although oxygen consumption may provide an indication of mitochondrial function, it only measures one component of cellular bioenergetics and does not provide an assessment of other metabolic pathways that contribute to  
5 bioenergetic equilibrium, namely glycolysis.

Another conventional method for assessing cellular energy production is by measuring the amount of ATP in a cell. Luminescent ATP assay kits are commercially available for quantifying total energy metabolism. ATP assays are known to be relatively sensitive but they may not be an ideal metric of mitochondrial function as cells strive to maintain a particular ATP  
10 budget and will adjust metabolism accordingly. Thus, alterations in ATP levels are usually only detectable during pathophysiological changes. In addition, ATP assays are destructive (i.e., the cells are destroyed in order to measure the amount of ATP) and they lack kinetic information. Further, artifact has been reported from residual ATP present in dying or dead cells. And like the Clark-electrode assay, ATP cannot determine the relative contribution of different metabolic  
15 pathways to total ATP yield.

Still another convention method for assessing cell energetics is with commercially available MTT/XTT or alamarBlue™ kits. While these kits may provide a relatively simple way to assess cell health, they are not as sensitive as ATP assays. In addition, they have been reported to introduce error through cell toxicity, the very parameter they are supposed to be  
20 measuring. Further, both assays are destructive and do not provide kinetic information.

Accordingly, it would be desirable to provide a method of identifying and/or or evaluating actives that reduce, prevent and/or reverse the undesirable oxidative stress on a particular type of skin cell that is associated with a particular stressor. It would also be desirable to provide a skin care composition that includes skin care actives identified by the foregoing  
25 method. It would further be desirable to provide a method of treating skin damaged by the oxidative stress effects on a particular type of skin cell from a particular stressor.

#### BRIEF DESCRIPTION OF THE DRAWINGS

FIGS. 1A, 1B and 1C are illustrations of the oxygen consumption rate of keratinocytes.

FIG. 2 is an illustration of the oxygen consumption rate of keratinocytes.

30 FIGS. 3A, 3B and 3C are illustrations of the extracellular acidification rate of keratinocytes.

FIG 4 is an illustration of the extracellular acidification rate of keratinocytes.

FIGS. 5A, 5B, 5C, 5D and 5E are illustrations of the oxygen consumption rate of fibroblasts.

FIG. 6 is an illustration of the oxygen consumption rate of fibroblasts.

FIGS. 7A, 7B, 7C, 7D and 7E are illustrations of the extracellular acidification rate of fibroblasts.

FIG. 8 is an illustration of the extracellular acidification rate of fibroblasts.

FIG. 9 is an illustration of the extracellular acidification rate of fibroblasts.

5 FIG. 10 is an illustration of the extracellular acidification rate of fibroblasts.

FIG. 11 is an illustration of the oxygen consumption rate of fibroblasts.

#### SUMMARY OF THE INVENTION

In order to provide a solution to the aforementioned problems, disclosed herein is a method of identifying or evaluating skin-care actives that improve the metabolism of skin cells such and  
10 keratinocytes and fibroblasts. The method is directed to exposing skin cells to a stressor and a test agent, and determining a response to the test agent based on the change in metabolic indicators associated with glycolysis and oxidative phosphorylation. The test agent may be identified as a skin-care active when the oxidative phosphorylation and/or glycolysis response corresponds to an improvement in cellular metabolism.

15 Also disclosed herein is a method of making a personal care composition that provides a skin health benefit and is suitable for topical application to skin, the method comprising: identifying an active according to the method disclosed herein; and incorporating a safe and effective amount of the active into a pharmaceutically acceptable carrier.

Further disclosed herein is a method of improving skin health, comprising: identifying a  
20 target area of skin in need of a skin-care benefit; and applying a cosmetically effective amount of a personal care composition made according to the method disclosed herein to the target area.

Still further disclosed is a method of identifying a skin benefit agent that improves the metabolism of fibroblasts, comprising: providing a plurality of fibroblasts; exposing the fibroblasts to a stressor; non-lethally detecting a metabolic indicator associated with each of a  
25 glycolysis metabolic pathway and an oxidative phosphorylation metabolic pathway to provide a response of each metabolic pathway to the stressor; exposing the plurality of fibroblasts to a test agent; non-lethally detecting the metabolic indicators associated with each of the glycolysis and oxidative phosphorylation metabolic pathways to provide a response of each metabolic pathway to the test agent; and identifying the test agent as a skin benefit agent when at least one of the  
30 responses corresponds to an improvement in fibroblast metabolism relative to the same metabolic pathway.

#### DETAILED DESCRIPTION OF THE INVENTION

All percentages are by weight of the personal-care composition, unless otherwise specified. All ratios are weight ratios, unless specifically stated otherwise. All numeric ranges are inclusive of narrower ranges; delineated upper and lower range limits are interchangeable to create further ranges not explicitly delineated. The number of significant digits conveys neither  
5 limitation on the indicated amounts nor on the accuracy of the measurements. All measurements are understood to be made at about 25 °C and at ambient conditions, where “ambient conditions” means conditions under about one atmosphere of pressure and at about 50% relative humidity.

#### Definitions.

“Cosmetic” means providing a desired visual effect on an area of the human body. The  
10 visual cosmetic effect may be temporary, semi-permanent, or permanent. Some non-limiting examples of “cosmetic products” include products that leave color on the face, such as foundation, mascara, concealers, eye liners, brow colors, eye shadows, blushers, lip sticks, lip balms, face powders, solid emulsion compact, and the like.

“Dermatologically acceptable” means that the compositions or components thereof so  
15 described are suitable for use in contact with human skin without undue toxicity, incompatibility, instability, allergic response, and the like.

“Different types of cells” means cells that differ from one another in their intended biological function. Examples of different types of cells, with respect to one another, include, without limitation, fibroblasts, keratinocytes, melanocytes, myocytes, sebocytes, and adipocytes.

“Disposed” means an element is positioned in a particular place relative to another  
20 element.

“Non-lethal,” means that a test procedure is not intended to kill or destroy the cells being tested or observed. For example, non-lethally detecting a metabolic indicator means that at least  
25 75% of the cells are viable after the detection (e.g., 80%, 85%, 90%, 95% or even up to 99% or more of the cells remain viable). Ideally, 100% of the cells are viable after a non-lethal test, but it is to be appreciated that the death or destruction of some cells may be unavoidable and/or unrelated to the test.

“Oxidative Stressor” means an environmental element that causes the formation of undesirable reactive oxygen species in a cell. Some non-limiting examples of oxidative stressors  
30 include ultraviolet radiation, cigarette smoke, ozone, engine exhaust, diesel exhaust, smog, surfactants, and radiation from a computer monitor or television.

“Personal-care composition” means a composition suitable for topical application on mammalian skin. The personal care compositions of the present invention may be used in skin-care, cosmetic, and hair-care products; non-limiting uses of which include antiperspirants, deodorants, lotions (e.g. hand lotion and body lotion), skin-care products (e.g., face and neck  
5 lotions, serums, sprays), sunless tanners, cosmetics (e.g., foundation, concealer, blush, lipstick, lip gloss), depilatories, shampoos, conditioning shampoos, hair conditioners, hair dyes, body washes, moisturizing body washes, shower gels, skin cleansers, cleansing milks, hair and body washes, in-shower body moisturizers, pet shampoos, shaving preparations, after-shaves, razor moisturizing/lubricating strips, razor shave-gel bars, bar soaps, cleansing products, feminine-care  
10 products, oral-care products, and baby-care products. The methods of using any of the aforementioned compositions are also included within the meaning of personal-care composition.

“Regulate a skin condition” means maintaining skin appearance and/or feel with little to no degradation in appearance and/or feel.

“Safe and effective amount” means an amount of a compound or composition sufficient  
15 to significantly induce a positive benefit, preferably a positive skin or feel benefit, including independently or in combinations the benefits disclosed herein, but low enough to avoid serious side effects, i.e., to provide a reasonable benefit to risk ratio, within the scope of sound judgment of the skilled artisan.

“Skin” means the outermost protective covering of mammals that is composed of cells  
20 such as keratinocytes, fibroblasts and melanocytes. Skin includes an outer epidermal layer and an underlying dermal layer. Skin may also include hair and nails as well as other types of cells commonly associated with skin, such as, for example, myocytes, Merkel cells, Langerhans cells, macrophages, stem cells, sebocytes, nerve cells and adipocytes.

“Skin-care” means regulating and/or improving a skin condition. Some nonlimiting  
25 examples include improving skin appearance and/or feel by providing a smoother, more even appearance and/or feel; increasing the thickness of one or more layers of the skin; improving the elasticity or resiliency of the skin; improving the firmness of the skin; and reducing the oily, shiny, and/or dull appearance of skin, improving the hydration status or moisturization of the skin, improving the appearance of fine lines and/or wrinkles, improving skin exfoliation or  
30 desquamation, plumping the skin, improving skin barrier properties, improve skin tone, reducing the appearance of redness or skin blotches, and/or improving the brightness, radiancy, or translucency of skin. Some nonlimiting examples of “skin-care products” include skin creams, moisturizers, lotions, and body washes.



“Skin-care composition” means a composition that regulates and/or improves skin condition.

“Skin-care active” means a compound or combination of compounds that, when applied to skin, provide an acute and/or chronic benefit to skin or a type of cell commonly found therein.

5 Skin-care actives may regulate and/or improve skin or its associated cells (e.g., improve skin elasticity; improve skin hydration; improve skin condition; and improve cell metabolism).

“Topical application” means to apply or spread the compositions of the present invention onto the surface of the keratinous tissue.

As humans age, damage from external and internal stressors on the cells of the body  
10 accumulates (i.e., oxidative stress), which may lead to decreased efficiency and function of tissue and organs. It is not uncommon for oxidative stress to manifest as a reduction in the ability of the cells to produce energy. Skin-care actives that reduce, stop or even reverse the effects of oxidative damage to skin are known. But identifying new and/or better skin-care actives is difficult due to the complexity of the metabolic pathways of a cell and/or the number of different  
15 types of cells found in skin. Surprisingly, it has been found that certain oxidative stressors impact the bioenergetics of certain types of skin cells in previously unappreciated ways. This new learning may be used to identify skin-care actives that combat the specific metabolic effects of certain stressors on particular types of skin cells, which in turn may lead to a more holistic approach to skin care. In particular, a skin-care active, combination of actives and/or a skin-care  
20 regimen tailored to combat the effects of particular oxidative stressors on particular skin cell types may lead to an improvement in the overall health of a person’s skin. Accordingly, the novel method herein provides a convenient and accurate way to identify skin-care actives that combat the undesirable metabolic effects associated with a particular stressor on a particular type of skin cell.

25 While some examples herein may be directed to skin cells such as keratinocytes and fibroblasts in conjunction with UV radiation, it is to be appreciated that the method may be adapted to great advantage for use with any type of cell in conjunction with any stressor, as desired.

Two key metabolic pathways for mammalian cells to produce energy are the oxidative  
30 phosphorylation (“oxphos”) pathway and the glycolysis pathway. Both pathways are necessary to maintain a healthy energy balance within most mammalian cells. Oxidative phosphorylation involves the transfer of electrons from electron donors to electron acceptors such as oxygen in

redox reactions, which results in the release of energy that is used to form ATP. In mammals, the redox reactions are carried out by a series of protein complexes within the mitochondria membrane, and the linked sets of proteins are called electron transport chains. The energy released by electrons flowing through this electron transport chain is used to transport protons  
5 across the mitochondrial membrane, in a process called chemiosmosis, which generates potential energy in the form of a pH gradient and an electrical potential across this membrane. An enzyme commonly known as ATP synthase allows the potential energy to be used to generate ATP. Because the oxidative phosphorylation pathway uses oxygen to generate ATP, the rate at which a cell consumes oxygen may be used as a metabolic indicator of the cell. That is, the oxygen  
10 consumption rate of the cell may be directly correlated to energy production by the cell via oxidative phosphorylation. Additionally or alternatively, the carbon dioxide production rate may also be used as a metabolic indicator, since carbon dioxide is a by-product of cellular metabolism. A higher oxygen consumption rate or carbon dioxide production rate may indicate an increase in energy production from the oxidative phosphorylation pathway, and thus an  
15 improvement in the metabolism and/or health of a cell. Conversely, a lower oxygen consumption rate or carbon dioxide production rate may indicate a decrease in oxphos metabolism. As a result, it would be desirable to identify and/or evaluate compounds that reduce, prevent and/or reverse the decrease in oxphos metabolism caused by oxidative stress.

Glycolysis is the metabolic pathway that converts glucose into pyruvate. The free energy  
20 released in this process is used to form ATP and NADH (reduced NAD). Glycolysis is a definite sequence of ten reactions involving ten intermediate compounds (one of the steps involves two intermediates) that typically occurs in the cytosol of the cell. The intermediates provide entry points to glycolysis. For example, most monosaccharides such as fructose, glucose, and galactose, can be converted to one of these intermediates. The intermediates may also be directly  
25 useful. For example, the intermediate dihydroxyacetone phosphate (DHAP) is a source of the glycerol that combines with fatty acids to form fat. A by-product of glycolysis is lactic acid, which can form a lactate anion plus a proton in solution. Thus, lactic acid, lactate or proton concentration can be used as a metabolic indicator of glycolysis. That is, a change in extracellular pH or extracellular acidification rate may be directly correlated to energy production  
30 by the cell via the glycolysis pathway. A higher extracellular acidification rate may indicate an increase in energy production via the glycolysis pathway, and thus an improvement in the metabolism and/or health of a cell. Conversely, a lower extracellular acidification rate may indicate a decrease in glycolysis metabolism. As a result, it would be desirable to identify and/or

evaluate compounds that reduce, prevent and/or reverse the decrease in glycolysis metabolism caused by oxidative stress.

#### Keratinocytes

Keratinocytes are generally recognized as the predominant cell type in the epidermis, typically constituting about 95% of the cells found there. Keratinocytes are formed by differentiation from epidermal stem cells residing in the lower part of the stratum basale layer of the epidermis. Keratinocytes divide and differentiate as they move upward through the layers of the epidermis (e.g., stratum spinosum and stratum granulosum) to eventually become corneocytes in the stratum corneum. During the differentiation process, keratinocytes produce more and more keratin (“cornification”) and eventually permanently withdraw from the cell cycle to form the corneocytes that make up the hard outer layer of the stratum corneum. Corneocytes are eventually shed off through desquamation as new cells come in. When oxidative stress reduces the metabolism of keratinocytes, the rate at which the keratinocytes divide and differentiate may be reduced or even halted. This, in turn, may reduce the rate at which lost corneocytes are replaced in the stratum corneum and ultimately lead to an undesirable decrease in the barrier properties of the skin. Thus, it may be desirable to identify skin-care actives that reduce, prevent and/or reverse the undesirable metabolic effects of oxidative stress from certain stressors (e.g., UV-A, UV-B, cigarette/tobacco smoke, smog, ozone, engine exhaust, volatile organic compounds) on keratinocytes.

It is well known that exposing skin to UV radiation can cause oxidative stress to skin cells. However, before now, it was not fully appreciated how particular types of skin cells react metabolically to different wavelengths of UV radiation, or how the metabolic reaction of a particular type of skin cell to UV radiation may change with time after exposure to the UV radiation. It has now been found that the oxphos and glycolysis metabolic pathways of keratinocytes respond differently to UV-B radiation (i.e., electromagnetic radiation in the wavelength range of 315 – 280 nm). In particular, it has been found that exposure of keratinocytes to UV-B radiation causes a decrease in metabolic activity in the oxphos pathway, but appears to have no significant effect on glycolysis within the first 2 hours after exposure. Consequently, it may be desirable to identify skin-care actives that directly (i.e., without the presence of a stressor or ROS) and/or indirectly (e.g., only when the cell is exposed a stressor or ROS) reduce, prevent and/or reverse the undesirable metabolic effects of UV-B radiation on keratinocytes.

It has also been found that keratinocytes exhibit a previously unappreciated metabolic response to UV-B radiation at a particular dose and/or time after exposure. This discovery provides unique insight into screening for skin care actives to combat the effects of oxidative stress on keratinocytes from UV-B radiation. In particular, it is now known that between 5 milliJoules per square centimeter (“mJ/cm<sup>2</sup>”) and 50 mJ/cm<sup>2</sup> (e.g., from 7.5 – 40 mJ/cm<sup>2</sup>, 10 – 30 mJ/cm<sup>2</sup>, or even 15 – 30 mJ/cm<sup>2</sup>) of UV-B radiation provides sufficient energy to induce a measurable metabolic response in the oxphos and/or glycolysis pathway of keratinocytes, but does not kill the keratinocytes. Further, in some instances, it can be important to detect the desired metabolic indicator at least 1 hour after exposure of the keratinocytes to a stressor, but typically no more than 72 hours after exposure (e.g., from 2 to 24 hours; 3 – 23 hours; 4 – 22 hours; 5 – 21 hours; 6 – 20 hours; 7 – 19 hours; 8 – 18 hours; 9 – 17 hours; 10 – 16 hours; 11 – 15 hours; or even 12 – 14 hours). If the metabolic indicator is detected too soon, the cells may not have sufficient time to fully respond to the stressor. On the other hand, if too much time passes after exposure, a response may be missed (e.g., if the metabolism of the cell returns to the basal value). Further, it is now known that, in some instances, the kinetic data observed at particular times can provide important insights into the responses of keratinocytes to oxidative stressors and/or ROS, which may not be apparent when using a conventional static detection method (e.g., ATP assay).

In some instances, the method herein comprises exposing a plurality of keratinocytes to a stressor such as UV-B radiation and/or a ROS such as hydrogen peroxide and then contacting the keratinocytes with a test agent. The metabolic responses of the keratinocyte to the stressor and the test agent are obtained by detecting a metabolic indicator corresponding to each of the oxphos and glycolysis metabolic pathways. The metabolic responses are detected in real time in a controlled environment, and the oxphos and glycolysis indicators are obtained from the same cells at the same time. In some embodiments, it may be desirable to provide a basal value for each of the oxphos and glycolysis metabolic pathways of the keratinocyte, and compare the metabolic responses to the basal value and/or each other to determine the response of the metabolic pathway to the stressor and/or test agent. “Basal value” means the value of the metabolic indicator at a normal resting state prior to exposure of the cell to a stressor or test agent. The basal value of a cell may be provided by measuring the metabolic indicators and/or by consulting the scientific literature and/or other suitable sources. However, measuring the basal value may be preferred since it is well known that the basal metabolic values of cells may vary due to a variety of environmental and intrinsic factors.

FIGS. 1A, 1B and 1C illustrate kinetic data obtained by detecting the oxygen consumption rate (“OCR”) of keratinocytes after exposure to UV-B radiation (312 nanometers). The keratinocytes illustrated in the figures are primary keratinocytes obtained from Gibco Life Technologies, Grand Island, NY, USA. The cells were prepared and tested according to the method described in the Test Methods below. The upper plot 30A, 30B and 30C in each of FIGS. 1A, 1B and 1C illustrates the basal OCR value of the keratinocytes (i.e., OCR with no UV-B exposure). The lower plots 35A, 35B and 35C of each of FIGS. 1A, 1B and 1C illustrate the response OCR values of the keratinocytes after exposure to UV-B radiation doses of 7.5, 15 and 30 mJ/cm<sup>2</sup>, respectively. It is to be appreciated that providing the proper dose (i.e., 7.5, 15 or 30 mJ/cm<sup>2</sup>) can depend on a variety of well-known factors (e.g., dose level desired, age of light bulb, type of light bulb, whether the instrument is warm), and it is well within the ability of one of ordinary skill to provide the desired dose of UV radiation. The control plots 37A, 37B and 37C illustrate the control OCR values measured on a well that did not contain any keratinocytes, but included the same medium as the other wells. The control value enables correction of the basal value and response value for any background effect that may be present. The first data point in each plot is taken approximately one hour after exposure to UV-B radiation. As illustrated in FIGS. 1A, 1B and 1C, the response OCR values of the keratinocytes after exposure to UV-B were generally lower than the corresponding basal OCR values. It is believed, without being limited by theory, that the data illustrated in FIGS. 1A, 1B and 1C indicate that UV-B radiation decreases the oxphos metabolism of keratinocytes. Consequently, it would be desirable to identify skin-care actives that combat the undesirable effects of a stressor such as UV-B radiation on the oxphos metabolic pathway of keratinocytes and/or act directly to improve the oxphos metabolic pathway (i.e., cause an improvement even without the presence of a stressor or ROS), and incorporate a safe and effective amount of such actives into a personal care composition.

FIG. 2 illustrates the keratinocyte OCR response 24 hours after exposure to UV-B radiation (312 nm). The 24-hour values illustrated in FIG. 2 were obtained by exposing the keratinocytes to UV-B radiation and then incubating the plate at 37° C and 5% CO<sub>2</sub> for 24 hours. Twenty-four hours after UV exposure, the cells were analyzed according to the method described below. The 24-hour basal value 40 is shown at the far left side of the chart 49. Immediately to the right of the basal value 40 is the 24-hour response OCR value 41 for a 7.5 mJ/cm<sup>2</sup> dose of UV-B radiation, followed by the 24-hour response OCR value 42 for a 15 mJ/cm<sup>2</sup> dose. And at the far right of the chart 49 is the 24-hour response OCR value 43 for a 30 mJ/cm<sup>2</sup> dose. As illustrated in FIG. 2, the keratinocytes continued to show a reduction in OCR 24 hours after

exposure to UV-B doses of 7.5, 15 and 30 mJ/cm<sup>2</sup>. Thus, it may be important to measure the keratinocyte oxphos response up to 24-hours or more (e.g., 48 or 72 hours) after exposure to UV-B, particularly when screening for skin-care actives that block and/or mitigate the negative effects of UV-B on keratinocyte oxphos metabolism.

5           FIGS. 3A, 3B and 3C illustrate kinetic data obtained by detecting the extra cellular acidification rate (“ECAR”) of keratinocytes after exposure to UV-B radiation (312 nanometers). The keratinocytes are primary keratinocytes obtained in the same way as described above with regard to FIGS. 1A, 1B and 1C, and the test was performed according to the method described in the Test Methods section below. The upper plots 65A, 65B and 65C in each of FIGS. 3A, 3B  
10           and 3C illustrate the response ECAR values of the keratinocytes after exposure to UV-B radiation doses of 7.5, 15 and 30 mJ/cm<sup>2</sup>, respectively. The lower plot 60A, 60B and 60C illustrates the basal ECAR values of the keratinocytes (i.e., no UV-B exposure). The control plots 67A, 67B and 67C illustrate the control ECAR values measured on a well that did not contain any keratinocytes, but included the same medium as the other wells. As illustrated in FIGS. 3A,  
15           3B and 3C, the control values 67A, 67B and 67C did not show any measurable change over time, whereas the response ECAR values of the keratinocytes were generally higher than the corresponding basal ECAR values. A statistical analysis of the ECAR response data was conducted using a 1-way analysis of variance (“ANOVA”) with a Dunnett’s correction, and the statistical analysis showed that there was no statistically significant change in the ECAR  
20           response values 65A, 65B and 65C relative to the corresponding basal values 60A, 60B and 60C.

FIG. 4 illustrates the keratinocyte ECAR response at 24 hours after exposure to UV-B radiation (312 nm). The 24-hour values illustrated in FIG. 4 were obtained by exposing the keratinocytes to UV-B radiation and then incubating the plate at 37° C and 5% CO<sub>2</sub> for 24 hours. Twenty-four hours after UV exposure, the cells were analyzed according to the method described  
25           below. The 24-hour basal ECAR value 70 is shown at the far left side of the chart 79. Immediately to the right of the basal value 70 is the 24-hour response ECAR value 71 for a 7.5 mJ/cm<sup>2</sup> dose of UV-B radiation, followed by the 24-hour response ECAR value 72 for a 15 mJ/cm<sup>2</sup> dose. And at the far right of the chart 79 is the 24-hour response ECAR value 73 for a 30 mJ/cm<sup>2</sup> dose. A statistical analysis of the 24-hour ECAR response data was conducted using a 1-  
30           way ANOVA with a Dunnett’s correction. The analysis indicated that there was a statistically significant change in the ECAR response values relative to the basal value. Thus, the amount of time that passes after exposure of keratinocytes to a stressor such as UV-B radiation may be an important factor to consider when screening for skin-care actives that block and/or

mitigate the negative effects of UV-B on keratinocyte glycolysis metabolism. In particular, it may be desirable to wait more than 2 hours after exposure to a stressor such as UV-B (e.g., 4, 6, 8, 10, 12, 14, 16, 18, 20, 24, 48, or 72 hours) to detect a metabolic indicator of glycolysis in keratinocytes.

5           Fibroblasts

In some instances, it may be desirable to identify skin-care actives that reduce, prevent and/or reverse the undesirable effects of oxidative stress from certain stressors on fibroblasts. Fibroblasts are found in the dermal layer of the skin and the hypodermal (i.e., sub-cutaneous) layer, and are generally recognized as the cells that synthesize the extracellular matrix (“ECM”) and collagen to provide the structural framework for the tissues of mammals. The ECM and collagen help cushion the body from stress and strain by providing tensile strength and elasticity to the skin. Fibroblasts also play an important role in wound healing. When oxidative stress reduces the metabolism of fibroblasts, the body’s ability to synthesize collagen and the ECM may be reduced resulting in saggy, thinner looking skin. And the ability of the body to heal wounds may be impeded.

As mentioned previously, it has not been fully appreciated how particular types of skin cells react metabolically to different wavelengths of UV radiation, or that the metabolic reaction of a particular type of skin cell to UV radiation may change with time after exposure to the UV radiation. This adds uncertainty and difficulty to a process of identifying suitable skin-care actives. Surprisingly, it has been found that the oxphos and glycolysis metabolic pathways of fibroblasts respond differently to UV-A radiation. While most of the UV-B radiation that contacts the skin is absorbed or reflected by the keratinocytes, longer wavelength UV-A radiation can penetrate through the epidermis and damage the underlying fibroblasts and other cells commonly found in the dermis and hypodermis. In particular, it has been found that exposure of fibroblasts to UV-A radiation that exceeds a threshold energy level causes a decrease in metabolic activity in the oxphos pathway. Exposure of fibroblasts to UV-A radiation also may result in a decrease in metabolic activity in the glycolysis pathway. In addition, both metabolic pathways may exhibit changes in their responses at different energy levels and times after exposure. This discovery provides unique insight important for identifying skin-care actives that combat the effects of oxidative stressors such as UV-A radiation on fibroblasts. For example, it is now known that between 1 and 50 Joules per square centimeter (“J/cm<sup>2</sup>”) of UV-A radiation may provide sufficient energy to induce a measurable metabolic response in the oxphos and/or glycolysis pathway, but does not generally kill the fibroblasts. Suitable ranges of UV-A radiation

include between 5 and 40, 10 and 30, or even about 20 J/cm<sup>2</sup>. Further, it is also now known that in some instances it can be important to detect the metabolic indicator at least 1 hour after exposure of the fibroblasts to a stressor, but typically not more than 24 hours after exposure (e.g., from 2 to 24 hours; 3 – 23 hours; 4 – 22 hours; 5 – 21 hours; 6 – 20 hours; 7 – 19 hours; 8 – 18 hours; 9 – 17 hours; 10 – 16 hours; 11 – 15 hours; or 12 – 14 hours). If the metabolic indicator is detected too soon, the cells may not have sufficient time to fully respond to the stressor. On the other hand, if too much time elapses, an important transient response to the stressor may not be detected. Further, it is now known that, in some instances, the kinetic data observed at particular times can provide important insights into the responses of fibroblasts to oxidative stressors and/or ROS, which may not be apparent when using a conventional static detection method (e.g., ATP assay).

In some embodiments, the method herein includes exposing a plurality of fibroblasts to a stressor such as UV-A radiation and/or an ROS such as hydrogen peroxide and then contacting the fibroblasts with a test agent. The metabolic responses of the fibroblasts to the stressor and/or the test agent may be obtained by detecting a metabolic indicator corresponding to each of the oxphos and glycolysis metabolic pathways. The metabolic responses are detected in real time in a controlled environment, and the oxphos and glycolysis indicators are obtained from the same cells at the same time. In some embodiments, it may be desirable to provide a basal value for at least one of the oxphos and glycolysis metabolic pathways of the fibroblasts, and compare the metabolic responses to the basal value and/or each other to determine the response of the metabolic pathway to the stressor and/or test agent.

FIGS. 5A, 5B, 5C, 5D and 5E illustrate the kinetic data obtained by detecting the oxygen consumption rate of fibroblasts after exposure to UV-A radiation (365 nm). The fibroblasts are dermal fibroblasts obtained from ATCC, Bethesda, MD (BJ cell line). The cells were prepared and tested according to the method described in more detail below. The upper plots 90A, 90B, 90C, 90D and 90E in each of FIGS. 5A, 5B, 5C, 5D and 5E illustrate the basal OCR values of the fibroblasts (i.e., the OCR with no UV-A exposure). The lower plots 95A, 95B, 95C, 95D and 95E of each of FIGS. 5A, 5B, 5C, 5D and 5E illustrate the response OCR values of the fibroblasts after exposure to UV-A radiation at doses of 1, 5, 10, 20 and 30 J/cm<sup>2</sup>, respectively. The control plots 97A, 97B, 97C, 97D and 97E illustrate the OCR measured on a well that did not contain any fibroblasts, but included the same medium as the other wells. The control value enables correction of the basal value and response value for any background effect that may be present. As illustrated in FIGS. 5A, 5B, 5C, 5D and 5E, the response OCR values of



the fibroblasts after exposure to UV-A were generally lower than the corresponding basal OCR values. Thus, it is believed, without being limited by theory, that the data illustrated in FIGS. 5A, 5B, 5C, 5D and 5E indicate that UV-A radiation at doses of greater than  $1\text{J}/\text{cm}^2$  may decrease the oxphos metabolism of fibroblasts. Consequently, it would be desirable to identify skin-care actives that combat the undesirable effects of a stressor such as UV-A radiation on the oxphos metabolic pathway of fibroblasts and/or act directly to improve the oxphos metabolic pathway (i.e., cause an improvement even without the presence of a stressor or ROS), and incorporate a safe and effective amount of such actives into a personal care composition.

FIG. 6 illustrates fibroblast OCR responses at 24 hours after exposure to UV-A radiation (365 nm). The 24-hour values illustrated in FIG. 6 were obtained by exposing the fibroblasts to UV-B radiation and then incubating the plate at  $37^\circ\text{C}$  and  $5\%\text{CO}_2$  for 24 hours. Twenty-four hours after exposure to UV-A, the cells were analyzed according to the method described below. The 24-hour basal value 80 is shown at the far left side of the chart 89. Immediately to the right of the basal value 80 is the 24-hour response OCR value 81 for a  $1\text{J}/\text{cm}^2$  dose of UV-A radiation, followed by the 24-hour response OCR value 82 for a  $5\text{J}/\text{cm}^2$  dose. And at the far right of the chart 89 is the 24-hour response OCR value 83 for a  $10\text{J}/\text{cm}^2$  dose. As illustrated in FIG. 6, the fibroblasts appear to show a reduction in OCR after 24 hours only at a dose of  $10\text{J}/\text{cm}^2$ . Thus, it may be important to measure the fibroblast oxphos response up to 24-hours or more (e.g., 48 or 72 hours) after exposure to UV-A radiation when screening for skin-care actives to combat the undesirable effects of UV-A at a dose of, for example, greater than  $5\text{J}/\text{cm}^2$  or  $10\text{J}/\text{cm}^2$ . However, when screening for skin-care actives that combat the undesirable metabolic effects of UV-A radiation on fibroblasts at doses of less than  $10\text{J}/\text{cm}^2$  or  $5\text{J}/\text{cm}^2$ , time and resource consuming steps in the screening process may be avoided by not testing at the lower doses.

FIGS. 7A, 7B, 7C, 7D and 7E illustrate the kinetic data obtained by detecting the extra cellular acidification rate of fibroblasts. The fibroblasts are obtained in the same way as described above with regard to FIGS. 5A, 5B, 5C, 5D and 5E, and the test was performed according to the method described in the Test Methods below. In FIGS. 7A, 7B, 7C, 7D and 7E, the basal ECAR values are represented by plots 120A, 120B, 120C, 120D and 120E; the response ECAR values are represented by plots 125A, 125B, 125C, 125D and 125E; and the control ECAR values are represented by plots 127A, 127B, 127C, 127D and 127E. The basal values 120A, 120B, 120C, 120D and 120E and control values 127A, 127B, 127C, 127D and 127E are obtained in the same way as described above. The response values 125A, 125B, 125C, 125D and 125E correspond to the response of the fibroblasts after exposure to 1, 5, 10, 20 and 30

J/cm<sup>2</sup> of UV-A radiation (365 nm), respectively. As can be seen in FIG. 7A, the response ECAR value 125A of the fibroblasts after exposure to 1 J/cm<sup>2</sup> of UV-A was generally about the same as the basal value 120A, which suggests that the glycolysis metabolic pathway of the fibroblast is generally unaffected at this dose. But as illustrated in FIGS. 7B and 7C, the response ECAR values 125B and 125C of the fibroblasts after exposure to 5 and 10 J/cm<sup>2</sup> of UV-A, respectively, were higher than the corresponding basal values 120B and 120C. Surprisingly, when the fibroblasts were exposed to 20 J/cm<sup>2</sup> of UV-A, as illustrated in FIG. 7D, the response ECAR value 125D started out higher than the basal value 120D, but appears to decrease until eventually (i.e., approximately 85 to 102 minutes) crossing over the basal value 120D line to end up lower than the basal value 120D. The response ECAR value 125E of the fibroblasts after exposure to 30 J/cm<sup>2</sup>, as illustrated in FIG. 7E, is substantially lower than the basal value, which suggests that the glycolysis metabolic pathway of the fibroblasts has completely shut down. Thus, it is believed, without being limited by theory, that the data illustrated in FIGS. 7A, 7B, 7C, 7D and 7E indicate that both the time after exposure and dose are important factors to consider when screening skin-care actives that combat the effects of UV-A radiation on fibroblast glycolysis. In particular, doses above and below 20 J/cm<sup>2</sup> may have very different effects on the glycolysis metabolism of fibroblasts, and appreciating these differences can help identify beneficial combinations of skin-care actives that combat the undesirable effects of UV-A without inhibiting the desirable effects.

FIG. 8 illustrates fibroblast ECAR responses at 24 hours after exposure to UV-A radiation (365 nm). The 24-hour values illustrated in FIG. 8 were obtained by exposing the fibroblasts to UV-B radiation and then incubating the plate at 37° C and 5% CO<sub>2</sub> for 24 hours. Twenty-four hours after exposure to UV-A, the cells were analyzed according to the method described in the Test Methods below. The 24-hour basal ECAR value 90 is shown at the far left side of the chart 99. Immediately to the right of the basal value 90 is the 24-hour response ECAR value 91 for a 1 J/cm<sup>2</sup> dose of UV-A radiation, followed by the 24-hour response ECAR value 92 for a 5 J/cm<sup>2</sup> dose. And at the far right of the chart 99 is the 24-hour response ECAR value 93 for a 10 J/cm<sup>2</sup> dose. As illustrated in FIG. 8, the fibroblasts appear to show a reduction in ECAR after 24 hours only at a dose of 10 J/cm<sup>2</sup>. Thus, the time after exposure, especially up to 24 hours or more, may be an important factor to consider when screening for skin-care actives that combat effects of UV-A radiation on fibroblast glycolysis, depending on the dose of UV-A.

Personal Care Compositions and Method of Using the Same

Because of the skin health and/or appearance benefit provided by a healthy skin cells, it may be desirable to incorporate one or more skin-care actives identified according to the method herein into a cosmetic composition. That is, it may be desirable to identify a skin care active according to the method herein, and include the skin-care active(s) as an ingredient in the cosmetic composition. Such cosmetic compositions may include a dermatologically acceptable carrier and a skin-care active such as niacinamide that reduces, prevents and/or reverses the undesirable metabolic effects of oxidative stress on keratinocytes, fibroblasts and/or other types of cells commonly found in skin (e.g., melanocytes, myocytes, stem cells, sebocytes, neurocytes, and adipocytes). The cosmetic compositions herein may include one or more optional ingredients of the kind commonly included in the particular cosmetic compositing being provided. For example, the cosmetic composition may include additional skin-care actives known for regulating and/or improving the condition of mammalian skin. Nonlimiting examples of such optional ingredients include emollients, humectants, vitamins; peptides; and sugar amines. Other optional ingredients include sunscreen actives (or sunscreen agents) and/or ultraviolet light absorbers. In certain embodiments, the cosmetic composition may include a colorant, a surfactant, a film-forming composition, and/or a rheology modifier. Suitable cosmetic compositions herein may be in any one of a variety of forms known in the art, including, for example, an emulsion, lotion, milk, liquid, solid, cream, gel, mouse, ointment, paste, serum, stick, spray, tonic, aerosol, foam, pencil, and the like. The cosmetic compositions may also be incorporated into shave prep products, including, for example, gels, foams, lotions, and creams, and include both aerosol and non-aerosol versions. Other cosmetic compositions include antiperspirant, deodorant, and personal cleaning compositions such as soap and shampoo. Nonlimiting examples of cosmetic compositions and optional ingredients suitable for use therein are described in U.S. Publication No. 2010/0239510 filed by Ha, et al., on January 21, 2010.

Compositions incorporating skin-care actives identified by the novel methods described herein may be generally prepared according to conventional methods known in the art of making compositions and topical compositions. Such methods typically involve mixing of ingredients in or more steps to a relatively uniform state, with or without heating, cooling, application of vacuum, and the like. For example, emulsions may be prepared by first mixing the aqueous phase materials separately from the fatty phase materials and then combining the two phases as appropriate to yield the desired continuous phase. In certain embodiments, the compositions may be prepared to provide suitable stability (physical stability, chemical stability, photostability, etc.) and/or delivery of active materials. The composition may be provided in a package sized to store a sufficient amount of the composition for a treatment period. The size, shape, and design of the

package may vary widely. Some package examples are described in USPNs D570,707; D391,162; D516,436; D535,191; D542,660; D547,193; D547,661; D558,591; D563,221; and U.S. Publication Nos. 2009/0017080; 2007/0205226; and 2007/0040306.

5 The personal care compositions disclosed herein may be applied to the skin at an amount and frequency to improve skin elasticity, improve hydration, regulate or improve skin condition, maintain or improve the signs of skin aging, or maintain or improve insult-affected keratinous tissue. For example, it may be desirable to identify a target area of skin in need of a skin-care benefit applying a cosmetically safe and effective amount of the personal care composition to the target area. In some instances, the skin-care composition may be used as a specialized treatment  
10 for an entire face, with concentrated usage in areas with expression lines, wrinkles, undesirable tone or spots. Further, the personal-care compositions herein may be used to produce a chronic and/or acute skin or cosmetic benefit by topically applying to the skin of a mammal in need of such treatment a safe and effective amount of the skin-care composition.

#### Test Method

15 This method enables the non-lethal, simultaneous detection of metabolic indicators associated with the oxidative phosphorylation and glycolysis metabolic pathways in a controlled environment. The method also enables the collection of kinetic data. When assessing the metabolic response to a stressor or test agent, it is important to assess both metabolic pathways simultaneously to understand how the two metabolic pathways interact with the stressor or test  
20 agent and/or to one another. Additionally, it is important to monitor the metabolic pathways in real time (i.e., repeating periodic measurements) to observe trends and/or transient responses that may be missed when using methods that provide only static data. Thus, destructive tests are not suitable for use herein since they typically only allow for a single measurement.

It is also important to detect the metabolic indicator in a controlled environment to reduce  
25 the likelihood of artifact introduction. A suitable controlled environment should minimize or prevent any undesirable influence by external environmental conditions (e.g., temperature, pressure, humidity, light, and contact by undesirable gaseous, liquid and/or solid contaminants). For example, if the metabolic indicator being detected is oxygen consumption rate and the test sample is open to the environment, the measured oxygen concentration may not accurately reflect  
30 the amount of oxygen consumed by the cells, since at least some of the consumed oxygen may be replaced by environmental oxygen. Additionally, the test method itself should not introduce artifact into the measurement. For example, it is known that the Clark Electrode consumes oxygen, which is the very thing it is supposed to detect. Thus, the oxygen concentration

measured by a Clark Electrode device may not accurately reflect the amount of oxygen consumed by the cells in the test.

It is to be appreciated that environmental changes such as a change in the temperature of the medium may result in unwanted measurement errors. In particular, the capacity of the media to hold dissolved gasses changes with temperature, and therefore may result in an apparent change in dissolved gas concentration as the media seeks equilibrium with the surrounding environment. Further, the measurement properties of at least some types of sensors may change with temperature. Accordingly, it may be particularly desirable to control the environmental conditions such as the temperature of the medium in the test vessel and/or surrounding environment or apply a correction factor to the measurement. Similarly, any evaporation from the media due to other uncontrolled environmental conditions such as humidity or exposure to air currents may artificially impact the measurements made from various sensors including those of dissolved gases, ions, and temperature. Thus, providing a controlled environment to minimize or eliminate these factors can be important.

In some instances, the device used to detect the metabolic parameters may include a stage adapted to receive the test vessels (e.g., multiwell microplate) holding the cells. The device may also include a plunger configured to receive a barrier for isolating the environment within the test vessel from the external environment. The barrier may be configured to mechanically cooperate with a portion of the test vessel to seal the opening in the test vessel, for example, by mating with a seat or step in the test vessel wall. Additionally or alternatively, the plunger and barrier may be configured to reduce and/or expand the volume of the test vessel and the volume of the medium within the test vessel including at least a portion of the cells (e.g., 5 – 50%). For example, the barrier may be inserted into and/or retracted out of the test vessel by relative movement of the stage and the plunger. In some embodiments, the method may include perfusing additional media through the vessel and/or replenishing the medium. Reducing the volume of the medium enables the temporary creation of a highly concentrated volume of cells within a larger volume of cell media to improve the ability of the sensor(s) to detect sensitive changes in metabolic indicators in the medium that result from biological activity of the cells. By temporarily, rather than permanently, reducing the media volume (and therefore concentrating the cell/media mixture), cells are exposed to a non-normal environment for only a brief period of time, and therefore may not be adversely affected (e.g., killed) by the measurement process.

The instrument should also include a sensor capable of analyzing the desired metabolic indicator(s). In some embodiments, the sensor may be disposed on the barrier and/or plunger.

The sensor should be in sensing communication with the medium and configured to detect the desired metabolic indicator. The sensor may be configured to sense the presence and/or the concentration of the metabolic indicator; sense a rate of change of a concentration of the metabolic indicator; and/or sense a first concentration of a first metabolic indicator, sense a  
5 second concentration of a second metabolic indicator, and determine a relationship between the first concentration and the second concentration. It may be desirable to configure the sensor to detect the metabolic parameter without disturbing the cells. The instrument may also include a computer programmed to automate one or more aspects of the tests, including data collection and recording, cycling through one or more test steps and transferring a test agent or stressor to the  
10 extracellular environment. In some embodiments, the sensor may be coupled to the computer.

Suitable nonlimiting examples of devices that provide a controlled environment; non-lethal and simultaneous detection of metabolic indicators; and kinetic data are disclosed in U.S. Pat. Nos. 7,276,351; 7,638,321; and 7,851,201, to Teich et al.; and U.S. Pat. No. 8,202,702 to Neilson, et al. A particularly suitable device is the XF Extracellular Flux Analyzer available  
15 from Seahorse Bioscience, Massachusetts.

#### Cell Culture and Sample Preparation

Cells to be tested according to the method herein may be obtained by any suitable means known in the art. For example, the cells may be isolated from a natural environment (e.g., the  
20 skin of a person) or purchased from a suitable commercial source. The cells may be a primary cell line (i.e., a cell line that is isolated from a biological tissue source and propagated under normal tissue culture conditions) or an immortalized cell line (i.e., a cell line that has been modified via chemical or genetic modification such that its proliferation and doubling index are significantly greater than that available from primary cell lines). In some embodiments, frozen  
25 primary or immortalized cell lines may be obtained from a suitable commercial source and diluted with an appropriate growth medium into tissue culture flasks (e.g., collagen coated T-75 flask available from BD Biosciences) for incubation (e.g., at 37° C). Once the cells are ready for testing, they may be placed in a suitable test vessel (e.g., multi-well vessels, single-well vessels, one or more tubes, and conventional test plates such as a 12-well plate, 96-well plate or the like).  
30 A test medium may be provided in the test vessel to form an extracellular environment. The test medium should keep the cells alive and healthy for at least the duration of the test (e.g., at least 4 hours, 8 hours, 12 hours, 24 hours, 48 hours or 72 hours). The number of cells in each well should be sufficient to obtain a suitable measurement (e.g.,  $4 \times 10^4$  cells/well for keratinocytes and  $1 \times 10^5$  cells/well for fibroblasts). It is to be appreciated that during testing the cells may be

suspended in the test medium, attached to a suitable substrate disposed in the test vessel and/or attached to the test vessel.

By way of example, the keratinocytes discussed above with regard to FIGS. 1A, 1B, 1C, 2, 3A, 3B, 3C and 4 are frozen, human primary keratinocytes obtained from Gibco Life Sciences. The keratinocytes were grown to 70-80% confluence in EpiLife<sup>®</sup> brand keratinocyte medium (available from Invitrogen, Grand Island, NY) supplemented with human keratinocyte growth supplement and gentamicin/amphotericin B x500 solution, both available from Invitrogen. For each test, two separate donors of keratinocytes were cultured and equal cell numbers of each donor were combined in the test vessel. In this example,  $2 \times 10^4$  keratinocytes from each donor were placed in a Gelatin-coated, 24-well plate, for a total of approximately  $4 \times 10^4$  keratinocytes in each well. The number of cells present may be determined by any suitable means known in the art (e.g., using a Coulter counter). The keratinocytes were placed in the Gelatin-coated plates 24 hours prior to testing, and 100  $\mu$ L of the keratinocyte medium described above was added to each well. A test medium was made by modifying a commercially available EpiLife<sup>®</sup> medium that is devoid of buffers, calcium, glucose, pyruvate, and glutamine. The EpiLife<sup>®</sup> medium was modified by adding 10ng/ml insulin; 10ng/ml hydrocortisone; 60 $\mu$ M calcium chloride; 10mM of glucose; 1mM of pyruvate; and 2mM of glutamine. The test medium was warmed to 37° C and the pH adjusted to 7.4 with sodium hydroxide. The keratinocyte medium in the wells was replaced by the test medium 1 hour prior to taking measurements.

With regard to the fibroblasts discussed above in FIGS. 5A, 5B, 5C, 5D, 5E, 6, 7A, 7B, 7C, 7D, 7E and 8, frozen human dermal fibroblasts were obtained from the BJ cell line commercially available from ATCC, Bethesda, MD. The fibroblasts were grown to 70-80% confluence in a culture medium of Eagle's Minimum Essential Medium ("EMEM") supplemented with 10% fetal bovine serum ("FBS") and gentamicin/amphotericin B x500 solution (EMEM and FBS are available from ATCC, and gentamicin/amphotericin B x500 solution is available from Invitrogen). The fibroblasts were plated at  $1 \times 10^5$  cells per well in Gelatin-coated plates 24 hours prior to testing. Each well included 100  $\mu$ L of the fibroblast medium described above. A test medium was made from Dulbecco's Modified Eagle Medium (available from Seahorse Bioscience), 25mM glucose and 1mM pyruvate. The test medium was warmed to 37° C and the pH adjusted to 7.4. The cells were grown and plated at 37° C and 5% CO<sub>2</sub> air. The fibroblast medium in the wells was replaced by the test medium 1 hour prior to taking measurements.

The gelatin-coated plates may be prepared by coating each well in a multi-well plate (e.g., 24-well or 96-well plates available from Seahorse Biosciences, Massachusetts) with 0.2% gelatin (protein solution available from Sigma) diluted with phosphate buffered saline (“PBS”). The gelatin is diluted 1:10 to 0.2% in sterile PBS at 37° C, and then 50 uL of the diluted gelatin solution is added to each well and incubated at room temperature for 30 minutes. Excess liquid is removed by aspiration without touching the wells and the surface is allowed to dry for at least 2 hours.

#### Exposing Cells to a Stressor or Test Agent

The cells may be exposed to a stressor or test agent by any suitable means known in the art. In some embodiments, the cells may be exposed to the stressor prior to placing the cells in the instrument that will be used to detect the metabolic parameter. For example, the cells may be exposed to UV radiation using a BIO-SUN brand solar simulator (available from Vilber Lourmat, France) prior to placing the cells in an XF Extracellular Flux Analyzer. In this example, the test medium may be removed from plate wells and replaced with 100 uL of PBS to reduce any undesirable effects the test medium may have on the UV radiation (e.g., absorbance or reflectance). In embodiments where a metabolic indicator will not be detected within a suitable time after exposure (e.g., when obtaining a 24-hour measurement), the PBS may be replaced with the appropriate medium (e.g., the keratinocyte or fibroblast medium described above) immediately after exposure to the stressor but prior to placing the cells in an incubator. In order to obtain data from non-irradiated cells, the portion of the plate that will contain the non-irradiated cells may be covered with a UV impervious material such as aluminum foil. In some instances, a stressor and/or a test agent may be introduced into the test vessel before and/or after the cells are placed in the test device. For example, the test device may include one or more injector ports that enable the introduction of a substance directly into the test vessel before, during and/or after testing.

#### Measurement of Oxidative Phosphorylation and Glycolysis Metabolic Indicators

Oxygen consumption rate and extracellular acidification rate may be detected using an XF Extracellular Flux Analyzer or equivalent. The device should be capable of non-lethally and simultaneously detecting metabolic indicators of the oxphos and glycolysis pathways in a controlled environment, as well as providing kinetic data. The cells may be provided in a multi-well plate suitable for use with the instrument (e.g., 24-well plate or 96-well plate) and washed prior to testing. The cells may be washed by any suitable means known in the art (e.g., using a



Seahorse Biosciences XF prep station). When washing the cells, it may be desirable to remove the medium from the wells and wash the cells three times with a suitable amount of test medium (e.g., 180  $\mu$ L in a 96-well plate or 600  $\mu$ L in a 24-well plate). After washing the cells, a suitable amount of test medium is placed in each well, and the cells are equilibrated at 37° C in a CO<sub>2</sub>-free incubator for 1 - 1.5 hours prior to placing the plate in the instrument for testing. Following the equilibration period, load the plate containing the cells into the instrument and equilibrate according to manufacturer's instructions. The entire test is conducted at 37° C. In some embodiments, it may be desirable to set the instrument to provide a three minute mix cycle, a two minute wait cycle, and a 3 minute measurement cycle for keratinocytes and a two minute mix cycle, two minute wait cycle and 3 minute measurement cycle for fibroblasts. The cycles should be repeated for at least 88 minutes. It is to be appreciated that the cycles and times may be modified according to cell type and experiment design, as desired.

#### Example 1

The following example illustrates how the method herein can be used to identify skin care actives. Hydrogen peroxide is a well known ROS, and is used in this example to illustrate the undesirable metabolic effects associated with exposure of a cell to an oxidative stressor. The hydrogen peroxide was prepared at 10X working concentration in the fibroblast test medium described above. The test agent is niacinamide (also known as vitamin B<sub>3</sub>), which is commercially available from Sigma. The test agent was prepared at 10X the final working concentration in the fibroblast test medium described above to provide a test agent solution. The stressor solution and the test agent solution were both warmed to 37° C and pH adjusted to 7.4. The cells are frozen, human, dermal fibroblasts obtained from ATCC (BJ cell line). The cells were cultured and prepared according to the test method described above. The fibroblasts were plated in 2 gelatin-coated, 24-well, V7 plates. Each well contained approximately  $1 \times 10^5$  cells and 600  $\mu$ L of fibroblast test medium (i.e., DMEM, 25 mM glucose and 1 mM pyruvate) at 37° C and pH 7.4.

An XF Extracellular Flux Analyzer was used to detect metabolic indicators corresponding to the glycolysis and oxphos metabolic pathways (i.e., extracellular acidification rate and oxygen consumption rate). The plates were loaded into the analyzer and equilibrated according to the manufacturer's instructions. The test agent solution and stressor solution were each placed in an automated injection port of the XF cartridge plate. The analyzer was programmed to sequentially run a two minute mix cycle, a two minute wait cycle, and a 3 minute measurement cycle continuously for at least 88 minutes. Data points were collected and recorded by the analyzer.

The analyzer was allowed to complete three cycles prior to the addition of the stressor solution or test agent solution to provide a basal value for each metabolic indicator. After the third cycle was completed (i.e., about 21 minutes after detection began), the test solution and/or stressor solution was added to the wells by the analyzer from the appropriate injection port. The stressor solution was added to the well in sufficient amount to provide 1.5 mM hydrogen peroxide in the well. The test solution was added in sufficient amount to provide 0.1 mM, 0.25 mM, 0.5 mM or 1.0 mM niacinamide in the well. The first plate tested had six wells with only 1.5 mM hydrogen peroxide added (well nos. 4, 5, 6, 11, and 12); six wells with only 0.25 mM niacinamide added (well nos. 2, 3, 7, 8, and 9); six wells with a combination of 0.1 mM niacindamide and 1.5 mM hydrogen peroxide added simultaneously (well nos. 13, 14, 19, 20, and 12); and 6 wells with a combination of 0.25 mM niacinamide and 1.5 mM hydrogen peroxide added simultaneously (well nos. 16, 17, 18, 22, and 23. The second plate had six wells with only 1.5 mM hydrogen peroxide added (well nos. 4, 5, 6, 11, and 12); six wells with only 1.0 mM niacinamide added (well nos. 2, 3, 7, 8, and 9); six wells with a combination of 0.5 mM niacindamide and 1.5 mM hydrogen peroxide added simultaneously (well nos. 13, 14, 19, 20, 21); and 6 wells with a combination of 1.0 mM niacinamide and 1.5 mM hydrogen peroxide added simultaneously (well nos. 16, 17, 18, 22, 23). Well numbers 1, 10, 15, and 24 were blank wells. The results are illustrated in Table 1, Table 2 and Table 3 below and FIGS. 9, 10 and 11.

Table 1 – Glycolysis (Plate No. 1)

Well No.	ECAR (mpH/min)										
	1	2	3	4	5	6	7	8	9	10	11
1	2	1	1	-11	2	5	5	5	4	4	3
2	54	52	50	42	54	58	58	59	57	57	56
3	47	45	46	34	48	51	52	52	51	50	49
4	41	38	39	34	21	23	20	10	5	4	2
5	53	50	50	51	35	37	34	26	25	22	15
6	45	39	41	39	24	28	25	18	16	13	9
7	49	47	49	45	56	59	60	61	60	59	59
8	47	44	42	35	46	49	49	49	48	48	48
9	46	41	41	32	43	45	46	45	45	44	43
10	-3	-1	-1	6	-1	-3	-2	-3	-3	-2	-3
11	57	54	54	54	35	38	36	29	29	25	17
12	41	40	39	37	29	30	19	8	9	12	6
13	56	49	50	51	33	39	45	44	39	37	34
14	51	48	47	47	31	34	36	36	29	26	19
15	2	1	2	5	3	2	-2	-1	0	0	1
16	43	44	43	35	23	28	32	33	32	30	27

17	36	33	34	36	19	20	26	29	28	26	26
18	41	35	33	30	29	32	33	30	28	26	26
19	55	51	51	51	32	38	42	43	39	36	34
20	50	49	50	38	22	26	32	30	26	20	16
21	46	42	41	40	23	28	31	31	27	24	20
22	47	42	42	31	22	26	32	35	33	29	27
23	48	48	48	48	30	34	41	43	42	40	42
24	-1	-2	-1	1	-4	-4	-2	-1	-1	-2	-1

Table 1 illustrates the effects of hydrogen peroxide and niacinamide, alone and in combination, on the acidification rate of the extracellular environment of fibroblasts from the first plate of fibroblasts tested. The data from Table 1 was analyzed via XF Software Version 1.8 to calculate the averages for each treatment group at each time point and compared to the change from baseline as a percentage that is represented graphically in FIG. 9. As discussed above, the first three measurements were made prior to the addition of the hydrogen peroxide or niacinamide to provide a basal value. FIG. 9 shows a plot for each condition in Table 1, namely 1.5 mM H<sub>2</sub>O<sub>2</sub> alone 101; 0.25 mM niacinamide alone 100; 0.25 mM niacinamide and 1.5 mM H<sub>2</sub>O<sub>2</sub> in combination 103; and 0.1 mM niacinamide and 1.5 mM hydrogen peroxide in combination 102. As illustrated in Table 1 and FIG. 9, hydrogen peroxide by itself appears to causes a substantial decrease in extracellular acidification rate. Thus, a reasonable conclusion may be drawn that hydrogen peroxide causes a decrease in the glycolysis metabolism in human fibroblasts. Niacinamide, when added by itself at 0.25 mM, appears to cause an initial decrease in acidification rate before returning to the basal level or slightly above within about 20 minutes after its addition to the well. From this, a reasonable conclusion can be drawn that 0.25 mM niacinamide does not significantly affect glycolysis metabolism. However, when added in combination with hydrogen peroxide, niacinamide appears to lessen the detrimental effect of the hydrogen peroxide on glycolysis metabolism. That is, both the 0.1 mM and 0.25 mM niacinamide concentrations appear to slow the decrease in extracellular acidification rate and prevent the acidification rate from falling as far below the basal level when hydrogen peroxide is added alone.

Table 2 – Glycolysis (Plate No. 2)

Well No.	ECAR (mpH/min)										
	1	2	3	4	5	6	7	8	9	10	11
1	1	0	1	-9	3	3	3	3	2	2	1
2	48	45	45	41	49	50	49	49	48	48	48

3	38	37	35	25	39	39	38	38	37	36	35
4	45	41	41	36	21	22	15	5	2	1	0
5	52	50	50	52	33	34	30	25	22	20	15
6	38	35	35	36	21	19	15	10	7	6	3
7	36	34	34	28	38	38	38	37	37	37	36
8	51	48	45	44	52	52	52	52	52	52	51
9	39	40	39	34	44	44	43	42	42	41	40
10	1	1	0	6	1	1	2	1	1	2	2
11	52	51	50	52	36	35	33	29	29	28	24
12	55	55	54	48	42	43	24	8	11	16	13
13	43	39	40	36	23	24	28	32	34	35	35
14	49	45	47	55	40	40	45	48	50	50	50
15	-1	-1	-1	3	1	1	0	0	0	-1	0
16	40	40	38	31	18	22	24	26	30	32	34
17	45	43	45	46	27	30	35	39	43	46	48
18	81	81	81	56	57	64	70	73	77	77	78
19	44	45	41	42	24	27	31	36	37	42	42
20	52	49	50	55	32	33	40	44	48	49	50
21	36	36	37	25	17	17	19	22	24	25	24
22	41	40	40	33	22	25	27	29	32	34	35
23	66	66	66	59	42	47	54	60	64	67	68
24	-2	0	-1	-1	-5	-5	-5	-4	-3	-3	-3

Table 2 illustrates the effects of hydrogen peroxide and niacinamide, alone and in combination, on the acidification rate of the extracellular environment of fibroblasts from the second plate of fibroblasts tested. The data from Table 2 was analyzed via XF Software Version 5 1.8 to calculate the averages for each treatment group at each time point and compared to the change from baseline as a percentage that is represented graphically in FIG. 10. FIG. 10 shows a plot for each condition in Table 2, namely 1.5 mM H<sub>2</sub>O<sub>2</sub> alone 201; 1.0 mM niacinamide alone 200; 0.5 mM niacinamide and 1.5 mM H<sub>2</sub>O<sub>2</sub> in combination 202; and 1.0 mM niacinamide and 1.5 mM hydrogen peroxide in combination 203. As illustrated in Table 2 and FIG. 10, 1.5mM 10 hydrogen peroxide appears to cause the same metabolic response illustrated above in Table 1. Similarly, niacinamide, by itself at 1.0 mM, appears to cause substantially the same lack of metabolic response illustrated above in Table 1 for 0.25 mM niacinamide. However, when added in combination with hydrogen peroxide, the 0.5 mM and 1.0 mM niacinamide appears to further lessen the decrease in glycolysis when compared to the 0.1 mM and 0.25mM concentrations. 15 The 0.5 mM and 1.0 mM niacinamide concentrations even appear to prevent the acidification rate from decreasing below the basal value. Importantly, niacinamide by itself shows substantially no effect on fibroblast glycolysis, but when added with hydrogen peroxide the metabolic effect of the niacinamide appears to improve with increasing concentration. Thus, a reasonable

conclusion can be drawn that niacinamide combats the undesirable metabolic effects of the hydrogen peroxide on glycolysis, rather than directly improving glycolysis itself.

Table 3 – Oxidative Phosphorylation

Well No.	OCR (pMoles/min)										
	1	2	3	4	5	6	7	8	9	10	11
1	2	-1	1	-5	1	4	-6	-1	-1	6	-3
2	322	310	299	290	285	286	283	283	283	285	276
3	228	219	211	210	197	204	196	196	198	199	192
4	228	218	211	-	-6	28	56	69	53	34	-13
5	276	265	255	11	72	107	139	164	178	164	118
6	251	233	224	-	-	-	-	-41	1	1	-16
7	328	320	313	312	302	303	304	305	304	297	295
8	297	282	280	262	256	250	257	260	262	257	253
9	177	119	144	120	129	113	119	125	121	121	117
10	-6	-3	0	-1	-1	-6	0	-1	-1	-1	3
11	287	279	277	47	99	85	119	126	125	100	64
12	202	195	196	-	-77	-63	-	46	62	49	36
13	302	293	293	111	182	182	183	194	211	217	217
14	277	273	270	24	111	123	125	132	166	177	179
15	6	-1	-2	5	-4	-1	4	-1	1	2	0
16	247	245	240	33	111	109	124	126	159	169	170
17	285	271	275	35	118	120	123	126	144	157	169
18	206	195	190	0	18	19	21	27	48	61	61
19	332	327	317	90	203	202	182	202	212	219	232
20	309	307	294	87	192	176	177	183	208	217	228
21	316	309	301	106	161	137	128	135	145	152	163
22	238	236	230	-	-4	5	17	46	70	86	104
23	316	316	307	35	127	129	133	148	169	179	205
24	-1	6	1	1	5	3	1	3	2	-7	0

5 Table 3 illustrates the effects of hydrogen peroxide and niacinamide, alone and in combination, on the oxygen consumption rate of the extracellular environment of fibroblasts from the second plate of fibroblasts tested. The data from Table 3 was analyzed via XF Software Version 1.8 to calculate the averages for each treatment group at each time point and compared to change from baseline as a percentage that is represented graphically in FIG. 11. FIG. 11 shows a plot for each condition in Table 3, namely 1.5 mM H<sub>2</sub>O<sub>2</sub> alone 301; 1.0 mM niacinamide alone 300; 0.5 mM niacinamide and 1.5 mM H<sub>2</sub>O<sub>2</sub> in combination 302; and 0.1 mM niacinamide and 1.5 mM hydrogen peroxide in combination 303. As illustrated in Table 3 and FIG. 11, 1.5 mM hydrogen peroxide by itself initially causes a relatively sharp decrease in OCR followed by a less rapid increase. About 20 minutes or so after adding the hydrogen peroxide, the OCR begins

10

to approach a relatively constant rate, which is much lower than the basal rate. Thus, a reasonable conclusion may be drawn that hydrogen peroxide causes a decrease in the oxphos metabolism of human fibroblasts. Niacinamide, when added by itself at 1.0 mM, appears to have no significant affect on the oxphos metabolism of human fibroblasts. When niacinamide is added at a concentration of 0.25  $\mu$ M with 1.5 mM hydrogen peroxide, the niacinamide appears to have no significant impact on OCR. When niacinamide is added at a concentration of 1.0 mM in combination 1.5 mM hydrogen peroxide, it appears to lessen the initial decrease in OCR somewhat, but does not appear to have a significant impact beyond that. Thus, a reasonable conclusion can be drawn that 0.5 mM and 1.0 mM concentrations of niacinamide do not significantly combat the effects of hydrogen peroxide on the oxphos metabolism of human fibroblasts. While higher concentrations of niacinamide may further reduce the detrimental effects of hydrogen peroxide on oxphos metabolism, such concentrations may not be suitable for use in topical skin care compositions due to, for example, increases in production cost or undesirable side effects.

The dimensions and values disclosed herein are not to be understood as being strictly limited to the exact numerical values recited. Instead, unless otherwise specified, each such dimension is intended to mean both the recited value and a functionally equivalent range surrounding that value. For example, a dimension disclosed as "40 mm" is intended to mean "about 40 mm." Additionally, properties described herein may include one or more ranges of values. It is to be understood that these ranges include every value within the range, even though the individual values in the range may not be expressly disclosed.

All documents cited in the Detailed Description of the Invention are, in relevant part, incorporated herein by reference; the citation of any document is not to be construed as an admission that it is prior art with respect to the present invention. In particular, U.S. Provisional Application Ser. Nos. 61/711,500 and 61/711,521 are incorporated herein by reference in their entirety. To the extent that any meaning or definition of a term in this document conflicts with any meaning or definition of the same term in a document incorporated by reference, the meaning or definition assigned to that term in this document shall govern.

While particular embodiments of the present invention have been illustrated and described, it would be obvious to those skilled in the art that various other changes and modifications can be made without departing from the spirit and scope of the invention. It is therefore intended to cover in the appended claims all such changes and modifications that are within the scope of this invention.

## CLAIMS

What is claimed is:

1. A method of identifying a test agent as a skin-care active that improves the metabolism of keratinocytes, comprising:
  - a. providing a plurality of keratinocytes;
  - b. exposing the keratinocytes to a stressor;
  - c. non-lethally detecting a metabolic indicator associated with each of glycolysis and oxidative phosphorylation to provide a response of each to the stressor;
  - d. exposing the plurality of keratinocytes to a test agent;
  - e. non-lethally detecting a metabolic indicator associated with each of glycolysis and oxidative phosphorylation to provide a response of each to the test agent; and
  - f. identifying the test agent as a skin-care active when at least one of the responses of (e) indicates an improvement in keratinocyte metabolism relative to the corresponding response of (c).
2. The method of claim 1, wherein the stressor is selected from the group consisting of ultraviolet radiation, cigarette smoke, ozone, engine exhaust, diesel exhaust, smog, surfactants, and radiation from a computer monitor or television.
3. The method of claim 2, wherein the stressor is ultraviolet radiation having a wavelength of from about 315 nm to about 280 nm.
4. The method of any preceding claim, further comprising providing a basal value for each of the metabolic indicators associated with glycolysis and oxidative phosphorylation, and comparing the responses of (c) to the corresponding basal values to determine if one metabolic pathway exhibits a greater reduction in metabolism than the other.
5. The method of claim 4, wherein the test agent is identified as a skin-care active when one of the responses in (e) corresponds to an improvement in the metabolic pathway that exhibits a greater reduction in metabolism.
6. The method of claim 4, wherein the basal value is provided by measuring the metabolic indicators in keratinocytes that have not been exposed to the stressor.
7. The method of any preceding claim, wherein the metabolic indicators are detected in the same test vessel.

8. The method of any preceding claim, wherein the metabolic indicators are detected at substantially the same time.
9. The method of any preceding claim, wherein the detection of the metabolic indicators is done in a controlled environment.
10. The method of any preceding claim, wherein the metabolic indicator is detected at least 1 hour after exposure to the stressor.
11. The method of any preceding claim, wherein the metabolic indicator associated with the oxidative phosphorylation pathway is detected by measuring at least one of oxygen consumption rate and carbon dioxide generation rate.
12. The method of any preceding claim, wherein the metabolic indicator associated with the glycolysis pathway is detected by measuring at least one of extracellular acidification rate, extracellular lactic acid rate and lactate concentration rate.
13. The method of any preceding claim, further comprising repeating steps (a) to (f) for a plurality of non-keratinocyte cells selected from the group consisting of fibroblasts, melanocytes, myocytes, stem cells, sebocytes, neurocytes, and adipocytes, and identifying the test agent as a skin-care active when at least one of the responses of repeated step (e) indicates an improvement in the metabolism of the non-keratinocyte cells relative to the corresponding response of repeated step (c).
14. A method of making a personal care composition that provides a skin health benefit and is suitable for topical application to skin, the method comprising:
  - a. identifying an active according to the method of any preceding claim; and
  - b. incorporating a safe and effective amount of the active into a pharmaceutically acceptable carrier.
15. A method of improving skin health, comprising:
  - a. identifying a target area of skin in need of a skin-care benefit; and
  - b. applying a cosmetically effective amount of a personal care composition made according to the method of claim 14 to the target area.






FIG. 1A




FIG. 1B




FIG. 1C




FIG. 2




FIG. 3A




FIG. 3B




FIG. 3C




FIG. 4




FIG. 5A




FIG. 5B




FIG. 5C






FIG. 5D






FIG. 7A




FIG. 7B






FIG. 7C




FIG. 7D




FIG. 7E




FIG. 8




FIG. 9




FIG. 10




FIG. 11

INTERNATIONAL SEARCH REPORT

International application No  
PCT/US2013/064116

A. CLASSIFICATION OF SUBJECT MATTER  
INV. G01N33/50  
ADD.  
According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED  
Minimum documentation searched (classification system followed by classification symbols)  
G01N  
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)  
EPO-Internal, BIOSIS, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	SVOBODOVA ET AL: "Attenuation of UVA-induced damage to human keratinocytes by silymarin", JOURNAL OF DERMATOLOGICAL SCIENCE, ELSEVIER SCIENCE PUBLISHERS, SHANNON, IE, vol. 46, no. 1, 13 March 2007 (2007-03-13) , pages 21-30, XP005919541, ISSN: 0923-1811, DOI: 10.1016/J.JDERMSCI.2006.12.009 abstract; figures 3, 4 page 24, left-hand column page 28, right-hand column ----- -/--	1-15

Further documents are listed in the continuation of Box C.

See patent family annex.

\* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier application or patent but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- "&" document member of the same patent family

Date of the actual completion of the international search  27 November 2013	Date of mailing of the international search report  05/12/2013
Name and mailing address of the ISA/ European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Fax: (+31-70) 340-3016	Authorized officer  Weijland, Albert

## INTERNATIONAL SEARCH REPORT

International application No  
PCT/US2013/064116

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	"A formulation containing L-carnitine and other nutrients improves skin energy reserves and oxygen consumption in skin cells and improves the appearance of aging skin", JOURNAL OF THE AMERICAN ACADEMY OF DERMATOLOGY, C.V. MOSBY, ST. LOUIS, MO, US, vol. 60, no. 3, 1 March 2009 (2009-03-01), page AB29, XP025963988, ISSN: 0190-9622 [retrieved on 2009-02-24] abstract	1-15
A	----- US 2003/012405 A1 (HATTA KOICHI [JP] ET AL) 16 January 2003 (2003-01-16) abstract	1-15
A	----- US 2009/258841 A1 (MURPHY MICHAEL PATRICK [GB] ET AL) 15 October 2009 (2009-10-15) abstract -----	1-15

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/US2013/064116

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 2003012405	A1	16-01-2003	JP 2003016216 A
			US 2003012405 A1
-----			
US 2009258841	A1	15-10-2009	AU 2009251711 A1
			CA 2719658 A1
			CN 102046178 A
			EP 2271349 A1
			HK 1157225 A1
			JP 2011516483 A
			KR 20110015531 A
			NZ 588806 A
			RU 2010143806 A
			SG 189714 A1
			US 2009258841 A1
			WO 2009145982 A1
-----			