

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization

International Bureau

(43) International Publication Date
31 March 2022 (31.03.2022)



(10) International Publication Number
WO 2022/064490 A1

(51) International Patent Classification:

C07C 229/22 (2006.01)

(21) International Application Number:

PCT/IL2021/051150

(22) International Filing Date:

22 September 2021 (22.09.2021)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

277528 22 September 2020 (22.09.2020) IL

(71) Applicant: **AG PLENUS LTD** [IL/IL]; 13 Gad Feinsein St, 7612002 Rehovot (IL).

(72) Inventors: **INBAL, Boaz**; 218 Simtat Ha-Itstrubal St., P.O.B. 218, 9978800 Kfar Shmu'el (IL). **WEXSEL-BLATT, Ezequiel**; Hameyasdim 33, 9085500 Shoeva (IL). **FREUD, Yehoshua**; Hahatzotzra 2, 9839041 Maale Adumim (IL). **ATSMON-RAZ, Yoav**; Weizman 13/10, 7628026 Rehovot (IL). **LOTHAR, Willms**; Lindenstrasse 17, 56204 Hillscheid (DE).

(74) Agent: **COHEN, Mark S.** et al.; Pearl Cohen Zedek Latzer Baratz, P.O. Box 7198, 6107121 Tel Aviv (IL).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, IT, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, WS, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

- with international search report (Art. 21(3))
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))

(54) Title: HERBICIDAL COMPOUNDS AND METHODS OF USE THEREOF

(57) Abstract: The present invention relates to novel herbicidally active compounds, agrochemical composition thereof, methods of preparation thereof, and uses thereof for controlling the growth of undesirable plants (e.g., weeds), for example in crop fields.



WO 2022/064490 A1

HERBICIDAL COMPOUNDS AND METHODS OF USE THEREOF**FIELD OF THE INVENTION**

[001] The present invention relates to novel herbicidally active compounds, agrochemical
5 composition thereof, methods of preparation thereof, and uses thereof for controlling the growth of
undesirable plants (e.g., weeds), for example in crop fields.

BACKGROUND OF THE INVENTION

[002] Weeds often interfere with efficient utilization of land and water resources and typically
10 compete with desired plants for water, nutrients, light, carbon dioxide, and space. Many weeds are also
aesthetically displeasing, especially when the weeds appear within a stand of a desired plant, such as St.
Augustine grass or Kentucky bluegrass in a homeowner's lawn. Weeds may also obstruct visibility, become
fire hazards around buildings, and reduce the efficiency of irrigation systems. When weeds appear in
15 watercourses, such as rivers and lakes, the weeds may contribute to poor water quality, making the water
unsuitable for culinary and industrial uses. Furthermore, some weeds act in a poisonous fashion against
other plants, animals, and humans by secreting toxic substances known as allelopathic compounds or by
spreading agents that may cause allergies and/or disease. Finally, weeds provide shelter for insects and
rodents that spread disease or are otherwise harmful to desired plants, animals, or humans.

[003] Weeds cause agricultural losses to crops that consistently exceed losses caused by other classes
20 of agricultural pests, year after year. Besides reducing the quality of the crop, weed infestation may reduce
achievable crop yield by up to 100% of the theoretically achievable yield. A number of approaches,
including mechanical, agricultural, biological, and chemical techniques, have evolved in an attempt to
control weed infestation.

[004] Mechanical means, such as hand pulling, hoeing or cultivation, deep plowing, clipping, mowing,
25 burning and/or mulching, may be employed in an attempt to eradicate or control weeds. Also, cover crops
may be planted to keep the ground covered when not growing more valuable crops and thus weed infestation
that would ordinarily be expected to occur in bare ground areas is typically minimized. Crop rotation and
planting of "smother" crops that are adapted to grow more vigorously than weeds have also been attempted
as means of controlling weed infestations. Besides these mechanical and agricultural techniques, biological
30 methods of weed control, such as introduction of predator populations that feed on the weeds and thereby
reduce weed population, have also been attempted.

[005] Mechanical, agricultural, and biological methods of weed control, while sometimes helping to
reduce the extent of weed infestations, are not fully satisfactory. First, mechanical and agricultural
techniques are quite labor intensive and require use of limited physical and capital resources. Furthermore,
35 environmental factors beyond the control of the farmer or homeowner, such as excessive rainfall, may
diminish the effectiveness of these mechanical and agricultural techniques. Likewise, biological techniques,

such as introduction of predator populations, are not entirely satisfactory, since the predators may not be selective for only the weed population.

[006] Chemically active herbicides represent another potential weed control technique. These chemical herbicides may be broken down into pre-emergent herbicides and post-emergent herbicides. Pre-emergent herbicides typically interfere with germination of weed seeds, whereas post-emergent herbicides kill the weeds after the weed seeds have germinated and weed growth has begun.

[007] Pre-emergent herbicides may be effective when present at the required dosage at the time weed seed germination is ready to occur. However, this timing issue points out a major problem with respect to pre-emergent herbicides. Specifically, if the pre-emergent herbicide is not applied, or degrades, prior to weed seed germination, the weed seeds are free to germinate and begin growing into mature weeds. Additionally, pre-emergent herbicides are typically weed specific and are not equally effective against all types of weeds. The timing problem present with pre-emergent herbicides may be avoided by employing post-emergent herbicides and by applying the post-emergent herbicide only after the weed seeds have germinated and the weeds are actively growing. However, many presently available post-emergent herbicides are non-selective herbicides and therefore will kill desirable plants in addition to weeds.

[008] Many pre- and post-emergent herbicides also suffer from another problem. Specifically, many pre-emergent herbicides and post-emergent herbicides are either moderately or highly toxic to humans and animals and may thereby have damaging effects far beyond the intended weed control effect. Toxic herbicides may cause injury either immediately or over the long term to persons applying the herbicides and to persons present when the herbicides are applied. Also, residual concentrations of toxic herbicides that remain in the soil or water after application of the herbicide may pose a significant threat to human beings and to animals, including land-based animals and amphibians and fish, upon contact with the treated area or runoff from the treated area. Furthermore, public alarm about the use of toxic chemicals as herbicides and their potential widespread and long-term effects on environmental quality dictate against the continued use of these toxic herbicides.


[009] There is a need for an herbicidal solution that avoids the critical timing issues of pre-emergent herbicide applications. Furthermore, there is a need for an herbicidal solution that avoids the toxic effects of presently available pre-emergent and post-emergent herbicides on human beings, animals and the environment generally. Furthermore, there is a need for an economically efficient post-emergent weed control technique that selectively controls weeds without destroying or hindering the growth of desired plants. In addition, there is a need for a composition that reduces the amount of herbicides necessary to obtain sufficient weed control while minimizing the harm to crop plants.

[0010] As more weeds become resistant to herbicides, alternative compositions with high weed control are desired. Further, as no-till farming continues to increase in popularity, there is a greater need for effective herbicides. A composition with effective weed control and lower dosage rate will lead to increased crop plant yields, and decreased environmental, human, and mammalian health concerns.

SUMMARY OF THE INVENTION

[0011] In various embodiments, this invention is directed to a compound represented by the structure of formula **I**, **I(a)**-**I(ga)**, **X** and **X(a)**-**X(d)** as defined herein below or its agrochemically acceptable salt, stereoisomer, tautomer, hydrate, *N*-oxide, reverse amide analog, isotopic variant (e.g., deuterated analog), or any combination thereof.

[0012] In various embodiments, this invention is directed to a compound represented by the structure of formula **I(g)**:

**I(g)**

10 wherein

R₁, **R**_{1'}, **R**₂, **R**_{2'} and **R**₄₀ are each independently H, C₁-C₅ linear or branched, unsubstituted alkyl, methyl, ethyl, propyl, iso-propyl, t-Bu, iso-butyl, pentyl, benzyl, C(O)-R₁₀, or C(O)-CH₃;


R₃ is OH, or NH₂;

R₄ is NH₂, or OH;

15 wherein if **R**₃ is OH then **R**₄ is NH₂ and if **R**₃ is NH₂ then **R**₄ is OH;

wherein if **R**₃ is OH and **R**₄ is NH₂, then **n** + **m** cannot be equal to 3;

or **R**₃ and **R**₄ are joined together to form ring **A**, represented by the following structure:



20 **R**₅ is H, C₁-C₅ linear or branched, substituted or unsubstituted alkyl (e.g., methyl, CH₂SH, ethyl, butyl, CH₂-CCH, iso-propyl, CH₂-C(O)-OCH₃), C₂-C₅ linear or branched, substituted or unsubstituted alkenyl, C₂-C₅ linear or branched, substituted or unsubstituted alkynyl (e.g., CCH, CH₂-CCH), C₁-C₅ linear or branched haloalkyl (e.g., CF₃, CF₂CH₃, CH₂CF₃, CF₂CH₂CH₃, CH₂CH₂CF₃, CF₂CH(CH₃)₂, CF(CH₃)-CH(CH₃)₂), R₈-aryl (e.g., CH₂-Ph), C(=CH₂)-R₁₀ (e.g., C(=CH₂)-C(O)-OCH₃, C(=CH₂)-CN), substituted or unsubstituted alkyl sulfone (e.g., SO₂-CH₂-cyclopentyl), substituted or
 25 unsubstituted aryl (e.g., phenyl), substituted or unsubstituted heteroaryl (e.g., pyridine (2, 3, and 4-pyridine));

R₁₀ is H, CN, C₁-C₅ linear or branched alkyl, C(O)R, or S(O)₂R;

R is C₁-C₅ linear or branched alkyl, C₁-C₅ linear or branched alkoxy, phenyl, aryl or heteroaryl;

30 **m** is 1 or 2;

n is 0, 1, 2 or 3;

X₁ is S, O, or CH₂;


X₂ is S, O, or CH₂;

or its agrochemically acceptable salt, stereoisomer, tautomer, hydrate, *N*-oxide, reverse amide analog, isotopic variant (e.g., deuterated analog), or any combination thereof.

[0013] In some embodiments, ring **A** has two chiral centers. In some embodiments, the compound is not (*6R,7S*)-6-amino-7-hydroxyoctanoic acid or 5-((*4R,5S*)-5-methyl-2-oxooxazolidin-4-yl)pentanoic acid. In some embodiments, the compound is a substantially pure single stereoisomer. In some embodiments, the compound is a mixture of stereoisomers. In some embodiments, the compound is the substantially pure *SR* stereoisomer. In some embodiments, the compound is the substantially pure *RS* stereoisomer. In some embodiments, the substantially pure stereoisomer has a purity higher than 90%. In some embodiments, the substantially pure stereoisomer has a purity higher than 95%. In some embodiments, the substantially pure stereoisomer has a purity higher than 98%. In some embodiments, the compound is compound **101, 102, 104, 105, 113, 114, 115, 116, 117, 118, 119, 120, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 137, 138, 139, 140, 141, 142** or an agrochemically acceptable salt, stereoisomer, tautomer, hydrate, *N*-oxide, reverse amide analog, or isotopic variant thereof; each represents a separate embodiment according to this invention. In some embodiments, the compound is compound any one of the compounds listed in **Table 1** herein below or an agrochemically acceptable salt, stereoisomer, tautomer, hydrate, *N*-oxide, reverse amide analog, or isotopic variant thereof; each represents a separate embodiment according to this invention.

[0014] In various embodiments, this invention is directed to an herbicidal compound represented by the structure of formula **I, I(a)-I(ga), X** and **X(a)-X(d)** as defined herein below, or an agrochemically acceptable salt, stereoisomer, tautomer, hydrate, *N*-oxide, reverse amide analog, or isotopic variant (e.g., deuterated analog) thereof; each represents a separate embodiment according to this invention.


[0015] In various embodiments, this invention is directed to an herbicidal compound represented by the structure of formula **I(ga)**:



I(ga)

wherein

C_A and **C_B** are both chiral carbon centers, or **C_A** and **C_B** together with **R₃** and **R₄** are joined to form ring **A**, represented by the following structure:



R₁, R₁', **R₂, R₂'** and **R₄₀** are each independently H, C₁-C₅ linear or branched, unsubstituted alkyl, methyl, ethyl, propyl, iso-propyl, t-Bu, iso-butyl, pentyl, benzyl, C(O)-R₁₀, or C(O)-CH₃;

R₃ is OH, SH, NH₂, NHNH₂, NHR, N(R)₂, NHC(O)OBz, -NHC(O)-R₁₀, NHC(O)CH₃, C₁-C₅ linear or branched, substituted or unsubstituted alkyl, methyl, ethyl, propyl, iso-propyl, t-Bu, iso-butyl, pentyl, substituted or unsubstituted C₃-C₈ cycloalkyl, C₁-C₅ linear or branched or C₃-C₈ cyclic haloalkyl, C₁-C₅ linear or branched or C₃-C₈ cyclic alkoxy, substituted or unsubstituted C₃-C₈ heterocyclic ring, or substituted or unsubstituted aryl;

R₄ is NH₂, NHNH₂, N(R)₂, -NHC(O)-R₁₀, NHC(O)H, NHC(O)CH₃, C₁-C₅ linear or branched, substituted or unsubstituted alkyl, C₁-C₅ linear or branched or C₃-C₈ cyclic haloalkyl, C₁-C₅ linear or branched or C₃-C₈ cyclic alkoxy, substituted or unsubstituted C₃-C₈ cycloalkyl, substituted or unsubstituted C₃-C₈ heterocyclic ring, or substituted or unsubstituted aryl;

or **R₃** and **R₄** are joined together to form ring **A** as described above;

wherein **R₃** and **R₄** cannot both be NH₂, and

wherein if **R₃** is OH and **R₄** is NH₂, then **n + m** cannot be equal to 3;

R₅ is H, C₁-C₅ linear or branched, substituted or unsubstituted alkyl (e.g., methyl, CH₂SH, ethyl, butyl, CH₂-CCH, iso-propyl, CH₂-C(O)-OCH₃), C₂-C₅ linear or branched, substituted or unsubstituted alkenyl, C₂-C₅ linear or branched, substituted or unsubstituted alkynyl (e.g., CCH, CH₂-CCH), C₁-C₅ linear or branched haloalkyl (e.g., CF₃, CF₂CH₃, CH₂CF₃, CF₂CH₂CH₃, CH₂CH₂CF₃, CF₂CH(CH₃)₂, CF(CH₃)-CH(CH₃)₂), R₈-aryl (e.g., CH₂-Ph), C(=CH₂)-R₁₀ (e.g., C(=CH₂)-C(O)-OCH₃, C(=CH₂)-CN), substituted or unsubstituted alkyl sulfone (e.g., SO₂-CH₂-cyclopentyl), substituted or unsubstituted aryl (e.g., phenyl), substituted or unsubstituted heteroaryl (e.g., pyridine (2, 3, and 4-pyridine));

R₈ is [CH₂]_p

wherein **p** is between 1 and 10;

R₁₀ and **R₁₁** are each independently H, CN, C₁-C₅ linear or branched alkyl, C(O)R, or S(O)₂R;

or **R₁₀** and **R₁₁** are joined to form a substituted or unsubstituted C₃-C₈ heterocyclic ring;

R is C₁-C₅ linear or branched alkyl, C₁-C₅ linear or branched alkoxy, phenyl, aryl or heteroaryl, or two gem R substituents are joined together to form a 5 or 6 membered heterocyclic ring;

m is 1 or 2;

n is 0, 1, 2 or 3;


X₁ is S, O, or CH₂;

X₂ is S, O, or CH₂;

or its agrochemically acceptable salt, stereoisomer, tautomer, hydrate, *N*-oxide, reverse amide analog, isotopic variant (e.g., deuterated analog), or any combination thereof

[0016] In some embodiments, this invention is directed to an herbicidal compound represented by the structure of any one of compounds **101**, **102**, **104-120**, **123-134**, **137-178** or an agrochemically acceptable salt, stereoisomer, tautomer, hydrate, *N*-oxide, reverse amide analog, or isotopic variant thereof; each represents a separate embodiment according to this invention. In some embodiments, the compound is compound any one of the compounds listed in Table **2** herein below or an agrochemically acceptable salt, stereoisomer, tautomer, hydrate, *N*-oxide, reverse amide analog, or isotopic variant thereof; each represents a separate embodiment according to this invention.

[0017] In some embodiments, this invention is directed to an herbicidal compound represented by the structure of formula **X(a)**:



wherein

R₅ is H, C₁-C₅ linear or branched, substituted or unsubstituted alkyl (e.g., methyl, CH₂SH, ethyl, butyl, CH₂-CCH, iso-propyl, CH₂-C(O)-OCH₃), C₂-C₅ linear or branched, substituted or unsubstituted alkenyl, C₂-C₅ linear or branched, substituted or unsubstituted alkynyl (e.g., CCH, CH₂-CCH), C₁-C₅ linear or branched haloalkyl (e.g., CF₃, CF₂CH₃, CH₂CF₃, CF₂CH₂CH₃, CH₂CH₂CF₃, CF₂CH(CH₃)₂, CF(CH₃)-CH(CH₃)₂), R₈-aryl (e.g., CH₂-Ph), C(=CH₂)-R₁₀ (e.g., C(=CH₂)-C(O)-OCH₃, C(=CH₂)-CN), substituted or unsubstituted aryl (e.g., phenyl), substituted or unsubstituted heteroaryl (e.g., pyridine (2, 3, and 4-pyridine));

R₈ is [CH₂]_p

wherein **p** is between 1 and 10;

R₉ is [CH]_q, [C]_q

wherein **q** is between 2 and 10;

R₁₀ and **R₁₁** are each independently H, CN, C₁-C₅ linear or branched alkyl (e.g., methyl, ethyl), C(O)R (e.g., C(O)(OCH₃)), or S(O)₂R; or **R₁₀** and **R₁₁** are joined to form a substituted or unsubstituted C₃-C₈ heterocyclic ring (e.g., piperazine, piperidine),

wherein substitutions include: F, Cl, Br, I, OH, SH, C₁-C₅ linear or branched alkyl (e.g. methyl, ethyl), C₂-C₅ linear or branched alkenyl, C₂-C₅ linear or branched alkynyl (e.g. CCH), C₃-C₈ cycloalkyl, linear, branched or cyclic alkoxy, COOH, COO(R), NH₂, N(R)₂, CF₃, aryl, phenyl, heteroaryl (e.g., imidazole), C₃-C₈ cycloalkyl (e.g., cyclohexyl), CN, NO₂ or any combination thereof;

R is H, C₁-C₅ linear or branched alkyl (e.g., methyl, ethyl), C₁-C₅ linear or branched alkoxy (e.g., methoxy), phenyl, aryl or heteroaryl, or two gem R substituents are joined together to form a 5 or 6 membered heterocyclic ring;

X₁ is S, O, CH₂, CH(R) or C(R)₂;


n and **o** are each independently an integer number between 0 and 2;

m is an integer number between 1 and 3;

or its agrochemically acceptable salt, stereoisomer, tautomer, hydrate, *N*-oxide, reverse amide analog, isotopic variant (e.g., deuterated analog), or any combination thereof

[0018] In various embodiments, this invention is directed to a compound represented by the structure of formula **X(a)** as defined herein above, wherein **n** is 2; **o** is 0; and if X_1 is CH_2 , then R_5 cannot be H.

[0019] In some embodiments, this invention is directed to an herbicidal compound represented by the structure of formula **X(b)**:



X(b)

10 wherein

R_5 is H, C_1 - C_5 linear or branched, substituted or unsubstituted alkyl (e.g., methyl, CH_2SH , ethyl, butyl, CH_2-CCH , iso-propyl, $CH_2-C(O)-OCH_3$), C_2 - C_5 linear or branched, substituted or unsubstituted alkenyl, C_2 - C_5 linear or branched, substituted or unsubstituted alkynyl (e.g., CCH , CH_2-CCH), C_1 - C_5 linear or branched haloalkyl (e.g., CF_3 , CF_2CH_3 , CH_2CF_3 , $CF_2CH_2CH_3$, $CH_2CH_2CF_3$, $CF_2CH(CH_3)_2$, $CF(CH_3)-CH(CH_3)_2$), R_8 -aryl (e.g., CH_2-Ph), $C(=CH_2)-R_{10}$ (e.g., $C(=CH_2)-C(O)-OCH_3$, $C(=CH_2)-CN$), substituted or unsubstituted aryl (e.g., phenyl), substituted or unsubstituted heteroaryl (e.g., pyridine (2, 3, and 4-pyridine));

R_8 is $[CH_2]_p$

wherein **p** is between 1 and 10;

20 R_{10} is H, CN, C_1 - C_5 linear or branched alkyl (e.g., methyl, ethyl), $C(O)R$ (e.g., $C(O)(OCH_3)$), or $S(O)_2R$;

R is H, C_1 - C_5 linear or branched alkyl (e.g., methyl, ethyl), C_1 - C_5 linear or branched alkoxy (e.g., methoxy), phenyl, aryl or heteroaryl, or two gem **R** substituents are joined together to form a 5 or 6 membered heterocyclic ring;

25 X_1 is S, O, CH_2 , $CH(R)$ or $C(R)_2$;


X_2 is S, O, CH_2 , $CH(R)$ or $C(R)_2$;

n is an integer number between 0 and 2;

m is an integer number between 1 and 3;

or its agrochemically acceptable salt, stereoisomer, tautomer, hydrate, *N*-oxide, reverse amide analog, isotopic variant (e.g., deuterated analog), or any combination thereof

[0020] In various embodiments, this invention is directed to an herbicidal compound represented by the structure of formula **X(c)**:



X(c)

wherein

- 5 **R₅** is H, C₁-C₅ linear or branched, substituted or unsubstituted alkyl (e.g., methyl, CH₂SH, ethyl, butyl, CH₂-CCH, iso-propyl, CH₂-C(O)-OCH₃), C₂-C₅ linear or branched, substituted or unsubstituted alkenyl, C₂-C₅ linear or branched, substituted or unsubstituted alkynyl (e.g., CCH, CH₂-CCH), C₁-C₅ linear or branched haloalkyl (e.g., CF₃, CF₂CH₃, CH₂CF₃, CF₂CH₂CH₃, CH₂CH₂CF₃, CF₂CH(CH₃)₂, CF(CH₃)-CH(CH₃)₂), R₈-aryl (e.g., CH₂-Ph), C(=CH₂)-R₁₀ (e.g., C(=CH₂)-C(O)-OCH₃,
10 C(=CH₂)-CN), substituted or unsubstituted aryl (e.g., phenyl), substituted or unsubstituted heteroaryl (e.g., pyridine (2, 3, and 4-pyridine));

R₈ is [CH₂]_p

wherein **p** is between 1 and 10;

R₉ is [CH]_q, [C]_q

- 15 wherein **q** is between 2 and 10;

R₁₀ and **R₁₁** are each independently H, CN, C₁-C₅ linear or branched alkyl (e.g., methyl, ethyl), C(O)R (e.g., C(O)(OCH₃)), or S(O)₂R; or **R₁₀** and **R₁₁** are joined to form a substituted or unsubstituted C₃-C₈ heterocyclic ring (e.g., piperazine, piperidine),

- 20 wherein substitutions include: F, Cl, Br, I, OH, SH, C₁-C₅ linear or branched alkyl (e.g. methyl, ethyl), C₂-C₅ linear or branched alkenyl, C₂-C₅ linear or branched alkynyl (e.g. CCH), C₃-C₈ cycloalkyl, linear, branched or cyclic alkoxy, COOH, COO(R), NH₂, N(R)₂, CF₃, aryl, phenyl, heteroaryl (e.g., imidazole), C₃-C₈ cycloalkyl (e.g., cyclohexyl), CN, NO₂ or any combination thereof;

- 25 **R** is H, C₁-C₅ linear or branched alkyl (e.g., methyl, ethyl), C₁-C₅ linear or branched alkoxy (e.g., methoxy), phenyl, aryl or heteroaryl, or two gem R substituents are joined together to form a 5 or 6 membered heterocyclic ring;

X₁ is S, O, CH₂, CH(R) or C(R)₂;

X₂ is S, O, CH₂, CH(R) or C(R)₂;

n is an integer number between 0 and 2;


- 30 **m** is an integer number between 1 and 3;

or its agrochemically acceptable salt, stereoisomer, tautomer, hydrate, *N*-oxide, reverse amide analog, isotopic variant (e.g., deuterated analog), or any combination thereof.

[0021] In various embodiments, this invention is directed to a compound represented by the structure of formula X(c) as described hereinabove, wherein n is an integer number between 1 and 2; or its

agrochemically acceptable salt, stereoisomer, tautomer, hydrate, *N*-oxide, reverse amide analog, isotopic variant (e.g., deuterated analog), or any combination thereof.

[0022] In various embodiments, this invention is directed to an herbicidal compound represented by the structure of formula **X(d)**:



wherein

R₅ is H, C₁-C₅ linear or branched, substituted or unsubstituted alkyl (e.g., methyl, CH₂SH, ethyl, butyl, CH₂-CCH, iso-propyl, CH₂-C(O)-OCH₃), C₂-C₅ linear or branched, substituted or unsubstituted alkenyl, C₂-C₅ linear or branched, substituted or unsubstituted alkynyl (e.g., CCH, CH₂-CCH), C₁-C₅ linear or branched haloalkyl (e.g., CF₃, CF₂CH₃, CH₂CF₃, CF₂CH₂CH₃, CH₂CH₂CF₃, CF₂CH(CH₃)₂, CF(CH₃)-CH(CH₃)₂), R₈-aryl (e.g., CH₂-Ph), C(=CH₂)-R₁₀ (e.g., C(=CH₂)-C(O)-OCH₃, C(=CH₂)-CN), substituted or unsubstituted aryl (e.g., phenyl), substituted or unsubstituted heteroaryl (e.g., pyridine (2, 3, and 4-pyridine);

15 **R₈** is [CH₂]_p

wherein **p** is between 1 and 10;

R₉ is [CH]_q, [C]_q

wherein **q** is between 2 and 10;

R₁₀ and **R₁₁** are each independently H, CN, C₁-C₅ linear or branched alkyl (e.g., methyl, ethyl), C(O)R (e.g., C(O)(OCH₃)), or S(O)₂R; or **R₁₀** and **R₁₁** are joined to form a substituted or unsubstituted C₃-C₈ heterocyclic ring (e.g., piperazine, piperidine),

wherein substitutions include: F, Cl, Br, I, OH, SH, C₁-C₅ linear or branched alkyl (e.g. methyl, ethyl), C₂-C₅ linear or branched alkenyl, C₂-C₅ linear or branched alkynyl (e.g. CCH), C₃-C₈ cycloalkyl, linear, branched or cyclic alkoxy, COOH, COO(R), NH₂, N(R)₂, CF₃, aryl, phenyl, heteroaryl (e.g., imidazole), C₃-C₈ cycloalkyl (e.g., cyclohexyl), CN, NO₂ or any combination thereof;

R is H, C₁-C₅ linear or branched alkyl (e.g., methyl, ethyl), C₁-C₅ linear or branched alkoxy (e.g., methoxy), phenyl, aryl or heteroaryl, or two gem R substituents are joined together to form a 5 or 6 membered heterocyclic ring;

30 **X₂** is S, O, CH₂, CH(R) or C(R)₂;

n is an integer number between 0 and 2;

m is an integer number between 1 and 3;

or its agrochemically acceptable salt, stereoisomer, tautomer, hydrate, *N*-oxide, reverse amide analog, isotopic variant (e.g., deuterated analog), or any combination thereof.

35 [0023] In various embodiments, this invention is directed to a compound represented by the structure of formula **X(d)** as described hereinabove, wherein **m** is 1 or 3; or an agrochemically acceptable salt,

stereoisomer, tautomer, hydrate, *N*-oxide, reverse amide analog, isotopic variant (e.g., deuterated analog), or any combination thereof.

[0024] In various embodiments, this invention is directed to an agrochemical composition comprising an herbicidally effective amount of a compound according to this invention.


5 [0025] In various embodiments, this invention is directed to a method of controlling the growth of undesired plants, comprising applying a compound according to this invention or an agrochemical composition according to this invention, to crop fields.

[0026] In various embodiments, this invention is directed to a compound according to this invention, or an agrochemical composition according to this invention, for use in controlling the growth of
 10 undesired plants. In some embodiments, the plant is a eudicot (dicot) or a monocotyledon (monocot). In some embodiments, the plant is a weed. In some embodiments, the weed comprises: *Abutilon theophrasti*, *Amaranthus palmeri*, *Ambrosia artemisiifolia*, *Alopecurus myosuroides*, *Avena sterilis*, *Chenopodium album*, *Conyza Canadensis*, *Digitaria sanguinalis*, *Echinochloa colona*, *Euphorbia heterophylla*, *Lolium perenne*, *Lolium rigidum*, *Matricaria chamomilla*, *Phalaris paradoxa*, *Poa annua*, *Portulaca oleracea*, *Setaria viridis*, *Solanum nigrum* or any combination thereof. In some
 15 embodiments, the dicot plant is *Arabidopsis thaliana*, and/or the monocot plant is *Dactyloctenium aegyptium* or *Eragrostis teff*. In some embodiments, the compound is for use in pre-plant treatments, pre-emergence treatments, post-emergence treatments, or any combination thereof; each represents a separate embodiment according to this invention.

20

DETAILED DESCRIPTION OF THE INVENTION

[0027] In various embodiments, this invention is directed to a compound represented by the structure of formula (I):



25

wherein

A and **B** rings are absent, or are each independently a substituted or unsubstituted single or fused aromatic or heteroaromatic ring system (e.g., **B**: aryl, pyridine), or a substituted or unsubstituted single or fused C₃-C₁₀ cycloalkyl (e.g., cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl), or a substituted or
 30 unsubstituted single or fused C₃-C₁₀ heterocyclic ring (e.g., **A**: 5-methyloxazolidin-2-one, 1, 2, or 3-pyrrolidine, tetrahydropyridine, 5,6-dihydro-4H-1,3-thiazine, 4,5-dihydro-1H-imidazole, pyridine, tetrahydropyrimidine; piperidine, imidazole);

R₁, R₁', **R₂, R₂'** and **R₄₀** are each independently H, F, Cl, Br, I, OH, SH, R₈-OH (e.g., CH₂-OH), R₈-SH, -R₈-O-R₁₀, (e.g., -CH₂-O-CH₃), R₈-(C₃-C₈ cycloalkyl) (e.g., cyclohexyl), R₈-(C₃-C₈ heterocyclic ring) (e.g., CH₂-imidazole, CH₂-indazole), CF₃, CD₃, OCD₃, CN, NO₂, -CH₂CN, -R₈CN, NH₂, NHR, N(R)₂, R₈-N(R₁₀)(R₁₁) (e.g., CH₂-NH₂, CH₂-N(CH₃)₂), R₉-R₈-N(R₁₀)(R₁₁) (e.g., C≡C-CH₂-NH₂),
 5 B(OH)₂, -OC(O)CF₃, -OCH₂Ph, NHC(O)-R₁₀ (e.g., NHC(O)CH₃), NHC(O)-N(R₁₀)(R₁₁) (e.g., NHC(O)N(CH₃)₂), COOH, -C(O)Ph, C(O)O-R₁₀ (e.g. C(O)O-CH₃, C(O)O-CH(CH₃)₂, C(O)O-CH₂CH₃), R₈-C(O)-R₁₀ (e.g., CH₂C(O)CH₃), C(O)H, C(O)-R₁₀ (e.g., C(O)-CH₃, C(O)-CH₂CH₃, C(O)-CH₂CH₂CH₃), C₁-C₅ linear or branched C(O)-haloalkyl (e.g., C(O)-CF₃), -C(O)NH₂, C(O)NHR, C(O)N(R₁₀)(R₁₁) (e.g., C(O)N(CH₃)₂), SO₂R, SO₂N(R₁₀)(R₁₁) (e.g., SO₂N(CH₃)₂, SO₂NHC(O)CH₃), C₁-
 10 C₅ linear or branched, substituted or unsubstituted alkyl (e.g., methyl, 2, 3, or 4-CH₂-C₆H₄-Cl, ethyl, propyl, iso-propyl, t-Bu, iso-butyl, pentyl, benzyl), C₁-C₅ linear or branched, substituted or unsubstituted alkenyl (e.g., CH=C(Ph)₂), C₁-C₅ linear or branched or C₃-C₈ cyclic haloalkyl (e.g., CF₃, CF₂CH₃, CH₂CF₃, CF₂CH₂CH₃, CH₂CH₂CF₃, CF₂CH(CH₃)₂, CF(CH₃)-CH(CH₃)₂), C₁-C₅ linear or branched or C₃-
 15 C₈ cyclic alkoxy (e.g. methoxy, ethoxy, propoxy, isopropoxy, O-CH₂-cyclopropyl, O-cyclobutyl, O-cyclopentyl, O-cyclohexyl, 1-butoxy, 2-butoxy, O-tBu), optionally wherein at least one methylene group (CH₂) in the alkoxy is replaced with an oxygen atom (e.g., O-1-oxacyclobutyl, O-2-oxacyclobutyl), C₁-C₅ linear or branched thioalkoxy, C₁-C₅ linear or branched haloalkoxy (e.g., OCF₃, OCHF₂), C₁-C₅ linear or branched alkoxyalkyl, substituted or unsubstituted C₃-C₈ cycloalkyl (e.g., cyclopropyl, cyclopentyl), substituted or unsubstituted C₃-C₈ heterocyclic ring (e.g., 3-methyl-4H-1,2,4-
 20 triazole, 5-methyl-1,2,4-oxadiazole, thiophene, oxazole, oxadiazole, imidazole, furane, triazole, tetrazole, pyridine (2, 3, or 4-pyridine), 3-methyl-2-pyridine, pyrimidine, pyrazine, oxacyclobutane (1 or 2-oxacyclobutane), protonated or deprotonated pyridine oxide), substituted or unsubstituted aryl (e.g., phenyl), substituted or unsubstituted benzyl (e.g., benzyl, 4-Cl-benzyl, 4-OH-benzyl);

or **R₂** and **R₁** are joined to form a C₃-C₈ substituted or unsubstituted, carbocyclic or heterocyclic
 25 ring;

R₃ is H, F, Cl, Br, I, OH, SH, =O, R₈-OH (e.g., CH₂-OH), R₈-SH, -R₈-O-R₁₀, (e.g., CH₂-O-CH₃) CF₃, CD₃, OCD₃, CN, NO₂, -CH₂CN, -R₈CN, NH₂, NHNH₂, NHR, N(R)₂, R₈-N(R₁₀)(R₁₁) (e.g., CH₂-NH₂, CH₂-N(CH₃)₂), R₉-R₈-N(R₁₀)(R₁₁), B(OH)₂, -OC(O)CF₃, -OCH₂Ph, NHC(O)OBz, -NHC(O)-R₁₀ (e.g., NHC(O)CH₃), NHC(O)-N(R₁₀)(R₁₁) (e.g., NHC(O)N(CH₃)₂), COOH, -C(O)Ph, C(O)O-R₁₀ (e.g. C(O)O-CH₃, C(O)O-CH₂CH₃), R₈-C(O)-R₁₀ (e.g., CH₂C(O)CH₃), C(O)H, C(O)-R₁₀ (e.g., C(O)-CH₃, C(O)-CH₂CH₃, C(O)-CH₂CH₂CH₃), C₁-C₅ linear or branched C(O)-haloalkyl (e.g., C(O)-CF₃), -
 30 C(O)NH₂, C(O)NHR, C(O)N(R₁₀)(R₁₁) (e.g., C(O)N(CH₃)₂), SO₂R, SO₂N(R₁₀)(R₁₁) (e.g., SO₂N(CH₃)₂), C₁-C₅ linear or branched, substituted or unsubstituted alkyl (e.g., methyl, C(OH)(CH₃)(Ph), ethyl, propyl, iso-propyl, t-Bu, iso-butyl, pentyl), C₁-C₅ linear or branched or C₃-C₈ cyclic haloalkyl (e.g., CF₃, CF₂CH₃, CF₂-cyclobutyl, CH₂CF₃, CF₂CH₂CH₃, CH₂CH₂CF₃, CF₂CH(CH₃)₂, CF(CH₃)-CH(CH₃)₂), C₁-C₅
 35 linear or branched or C₃-C₈ cyclic alkoxy (e.g. methoxy, ethoxy, propoxy, isopropoxy, O-CH₂-cyclopropyl), C₁-C₅ linear or branched thioalkoxy, C₁-C₅ linear or branched haloalkoxy, C₁-C₅ linear or branched alkoxyalkyl, substituted or unsubstituted C₃-C₈ cycloalkyl (e.g., cyclopropyl, cyclopentyl), substituted or unsubstituted C₃-C₈ heterocyclic ring (e.g., 3-methyl-4H-1,2,4-triazole, 5-methyl-1,2,4-

oxadiazole, thiophene, oxazole, isoxazole, imidazole, furane, triazole, pyridine (2, 3, or 4-pyridine), pyrimidine, pyrazine, oxacyclobutane (1 or 2-oxacyclobutane)), substituted or unsubstituted aryl (e.g., phenyl);

5 or **R**₃ and **R**₂ are joined to form a C₃-C₈ substituted or unsubstituted, carbocyclic or heterocyclic ring (e.g., cyclopropyl);

R₄ is H, F, Cl, Br, I, OH, SH, =O, =NH-OH, R₈-OH (e.g., CH₂-OH), R₈-SH, -R₈-O-R₁₀, (e.g., CH₂-O-CH₃) CF₃, CD₃, OCD₃, CN, NO₂, -CH₂CN, -R₈CN, NH₂, NHNH₂, NHR, N(R)₂, R₈-N(R₁₀)(R₁₁) (e.g., CH₂-NH₂, CH₂-N(CH₃)₂) R₉-R₈-N(R₁₀)(R₁₁), B(OH)₂, -OC(O)CF₃, -OCH₂Ph, NHC(O)OBz, -NHC(O)-R₁₀ (e.g., NHC(O)H, NHC(O)CH₃), NHC(O)-N(R₁₀)(R₁₁) (e.g., NHC(O)N(CH₃)₂), COOH, -C(O)Ph, C(O)O-R₁₀ (e.g., C(O)O-CH₃, C(O)O-CH₂CH₃), R₈-C(O)-R₁₀ (e.g., CH₂C(O)CH₃), C(O)H, C(O)-R₁₀ (e.g., C(O)-CH₃, C(O)-CH₂CH₃, C(O)-CH₂CH₂CH₃), C₁-C₅ linear or branched C(O)-haloalkyl (e.g., C(O)-CF₃), -C(O)NH₂, C(O)NHR, C(O)N(R₁₀)(R₁₁) (e.g., C(O)N(CH₃)₂), SO₂R, SO₂N(R₁₀)(R₁₁) (e.g., SO₂N(CH₃)₂), C₁-C₅ linear or branched, substituted or unsubstituted alkyl (e.g., methyl, C(OH)(CH₃)(Ph), ethyl, propyl, iso-propyl, t-Bu, iso-butyl, pentyl), C₁-C₅ linear or branched or C₃-C₈ cyclic haloalkyl (e.g., CF₃, CF₂CH₃, CF₂-cyclobutyl, CH₂CF₃, CF₂CH₂CH₃, CH₂CH₂CF₃, CF₂CH(CH₃)₂, CF(CH₃)-CH(CH₃)₂), C₁-C₅ linear or branched or C₃-C₈ cyclic alkoxy (e.g. methoxy, ethoxy, propoxy, isopropoxy, O-CH₂-cyclopropyl), C₁-C₅ linear or branched thioalkoxy, C₁-C₅ linear or branched haloalkoxy, C₁-C₅ linear or branched alkoxyalkyl, substituted or unsubstituted C₃-C₈ cycloalkyl (e.g., cyclopropyl, cyclopentyl), substituted or unsubstituted C₃-C₈ heterocyclic ring (e.g., 3-methyl-4H-1,2,4-triazole, 5-methyl-1,2,4-oxadiazole, thiophene, oxazole, isoxazole, imidazole, furane, triazole, pyridine (2, 3, or 4-pyridine), pyrimidine, pyrazine, oxacyclobutane (1 or 2-oxacyclobutane), indole), substituted or unsubstituted aryl (e.g., phenyl);

or **R**₃ and **R**₄ are joined together to form ring A as defined above (e.g., cyclopropyl, 5-methyloxazolidin-2-one);

25 **R**₅ is H, C₁-C₅ linear or branched, substituted or unsubstituted alkyl (e.g., methyl, CH₂SH, ethyl, butyl, CH₂-CCH, iso-propyl, CH₂-C(O)-OCH₃), C₂-C₅ linear or branched, substituted or unsubstituted alkenyl, C₂-C₅ linear or branched, substituted or unsubstituted alkynyl (e.g., CCH, CH₂-CCH), C₁-C₅ linear or branched haloalkyl (e.g., CF₃, CF₂CH₃, CH₂CF₃, CF₂CH₂CH₃, CH₂CH₂CF₃, CF₂CH(CH₃)₂, CF(CH₃)-CH(CH₃)₂), R₈-aryl (e.g., CH₂-Ph), C(=CH₂)-R₁₀ (e.g., C(=CH₂)-C(O)-OCH₃, C(=CH₂)-CN), substituted or unsubstituted alkyl sulfone (e.g., SO₂-CH₂-cyclopentyl), substituted or unsubstituted aryl (e.g., phenyl), substituted or unsubstituted heteroaryl (e.g., pyridine (2, 3, and 4-pyridine));

R₈ is [CH₂]_p

wherein **p** is between 1 and 10;

35 **R**₉ is [CH]_q, [C]_q

wherein **q** is between 2 and 10;

R₁₀ and **R**₁₁ are each independently H, CN, C₁-C₅ linear or branched alkyl (e.g., methyl, ethyl), C(O)R (e.g., C(O)(OCH₃)), or S(O)₂R;

or **R**₁₀ and **R**₁₁ are joined to form a substituted or unsubstituted C₃-C₈ heterocyclic ring (e.g., piperazine, piperidine),

R is H, C₁-C₅ linear or branched alkyl (e.g., methyl, ethyl), C₁-C₅ linear or branched alkoxy (e.g., methoxy), phenyl, aryl or heteroaryl, or two gem R substituents are joined together to form a 5 or 6 membered heterocyclic ring;

m is an integer number between 1 and 5 (e.g., 1 or 2);

n is an integer number between 0 and 5 (e.g., 0, 1, 2 or 3);

X₁ is S, O, N-OH, CH₂, C(R)₂ or N-OMe;

X₂ is S, O, N-OH, CH₂, C(R)₂ or N-OMe;

or **X**₂ together with the carbon next to **X**₁ are joined to form ring B as defined above;

X₃ is O, NH or N-R₅₀;

R₅₀ is H or C₁-C₅ linear or branched, substituted or unsubstituted alkyl;

or its agrochemically acceptable salt, stereoisomer, tautomer, hydrate, *N*-oxide, reverse amide analog, isotopic variant (e.g., deuterated analog), or any combination thereof.

[0028] In some embodiments, **R**₃ and **R**₄ are joined together to form ring A. In some embodiments, ring B is absent. In some embodiments, **R**₃ is OH, or NH₂. In some embodiments, **R**₄ is NH₂, or OH. In some embodiments, **R**₃ and **R**₄ cannot both be NH₂. In some embodiments, if **R**₃ is OH then **R**₄ is NH₂. In some embodiments, if **R**₃ is NH₂ then **R**₄ is OH. In some embodiments, if **R**₃ is OH then **R**₄ is NH₂ and if **R**₃ is NH₂ then **R**₄ is OH. In some embodiments, if **R**₃ is OH and **R**₄ is NH₂, then **n** + **m** cannot be equal to 3. In some embodiments, the compound is compound **101**, **102**, **104**, **105**, **106**, **113**, **114**, **115**, **116**, **117**, **118**, **119**, or **120**; each represents a separate embodiment according to this invention. In some embodiments, ring A has two chiral centers. In some embodiments, the compound is not (6*R*,7*S*)-6-amino-7-hydroxyoctanoic acid. In some embodiments, the compound is not 5-((4*R*,5*S*)-5-methyl-2-oxooxazolidin-4-yl)pentanoic acid. In some embodiments, the compound is a substantially pure single stereoisomer. In some embodiments, **R**₁ and **R**₁' are both H. In some embodiments, **R**₂ is CH₃ or CH₂CH₃. In some embodiments, **R**₂' is H or CH₃. In some embodiments, **R**₄₀ is CH₃ or H. In some embodiments, **X**₁ is CH₂. In some embodiments, **X**₂ is CH₂. In some embodiments, **X**₃ is O, NH or N-CH₃. In some embodiments, **R**₅ is H, or C₁-C₅ linear or branched, substituted or unsubstituted alkyl, or substituted or unsubstituted alkyl sulfone. In some embodiments, **R**₅ is a substituted or unsubstituted alkyl. In some embodiments, **R**₅ is H. In some embodiments, **R**₅ is H, ethyl, butyl, CH₂-CCH, CH₂-C(O)-OCH₃ or SO₂-CH₂-cyclopentyl; each is a separate embodiment according to this invention. In some embodiments, **n** is 0, 1, 2 or 3. In some embodiments, **n** is 1. In some embodiments, **m** is 1 or 2. In some embodiments, **m** is 1.

[0029] In some embodiments, **n** is 0. In some embodiments, A is a substituted aryl (e.g., 2-amino-phenyl). In some embodiments, A is a substituted or unsubstituted cycloalkyl (e.g., cyclopentyl, cyclohexyl). In some embodiment, the substitution is at least one selected from: F, Cl, Br, I, OH, SH, C₁-C₅ linear or branched alkyl, C₃-C₈ cycloalkyl, linear, branched or cyclic alkoxy, COOH, COO(R),

NH₂, N(R)₂, CF₃, aryl, phenyl, heteroaryl, C₃-C₈ cycloalkyl, CN and NO₂; each represents a separate embodiment according to this invention.

[0030] In some embodiments, **A** is a substituted cycloalkyl. In some embodiments, **A** is substituted cyclopropyl. In some embodiments, **A** is substituted cyclobutyl. In some embodiments, **A** is substituted cyclopentyl. In some embodiments, **A** is substituted cyclohexyl. In some embodiments, **A** is substituted at least with an amine. In some embodiments, **A** is substituted at least with NH₂. In some embodiments, **A** is substituted with NH₂. In some embodiments, **A** is a cycloalkyl, substituted at least with an amine. In some embodiments, **A** is a cycloalkyl, substituted with NH₂. In some embodiments, **A** is a 5 or 6 membered nitrogen containing heterocyclic ring. In some embodiments, **A** is 1, 2, or 3-piperidine, oxazolidin-2-one, tetrahydropyrimidine, pyridine, dihydro-thiazine, dihydro-imidazole, tetrahydropyridine, or pyrrolidine, which may be substituted or unsubstituted; each is a separate embodiment according to this invention. In some embodiments, **A** is substituted with an amine. In some embodiments, **A** is substituted at least with NH₂.

[0031] In some embodiments, **B** is absent. In some embodiment **B** is pyridine.


[0032] In some embodiments, **R**₁, **R**₁' , **R**₂, **R**₂' and **R**₄₀ are each independently H or C₁-C₅ linear or branched, substituted or unsubstituted alkyl (e.g., methyl, ethyl).

[0033] In some embodiments, substitutions include: F, Cl, Br, I, OH, SH, C₁-C₅ linear or branched alkyl (e.g. methyl, ethyl), C₂-C₅ linear or branched alkenyl, C₂-C₅ linear or branched alkynyl (e.g. CCH), C₃-C₈ cycloalkyl, linear, branched or cyclic alkoxy, COOH, COO(R), NH₂, N(R)₂, CF₃, aryl, phenyl, heteroaryl (e.g., imidazole), C₃-C₈ cycloalkyl (e.g., cyclohexyl), CN, NO₂ or any combination thereof.

[0034] In some embodiments, the compound is any one of the compounds listed in **Table 1**. In various embodiments, the compound is an herbicidal compound. In various embodiments, the compound is for use in controlling the growth of undesired plants. In some embodiments, the compound is any one of the compounds listed in **Table 2**.

25

[0035] In various embodiments, this invention is directed to a compound represented by the structure of formula **I(a)**:




I(a)

30

wherein

C_A and **C**_B are both chiral carbon centers, or **C**_A and **C**_B together with **R**₃ and **R**₄ are joined to form ring **A**, represented by the following structure:



R₁, R₁', **R₂, R₂'** and **R₄₀** are each independently H, F, Cl, Br, I, OH, SH, R₈-OH (e.g., CH₂-OH), R₈-SH, -R₈-O-R₁₀, (e.g., -CH₂-O-CH₃), R₈-(C₃-C₈ cycloalkyl) (e.g., cyclohexyl), R₈-(C₃-C₈ heterocyclic ring) (e.g., CH₂-imidazole, CH₂-indazole), CF₃, CD₃, OCD₃, CN, NO₂, -CH₂CN, -R₈CN, NH₂, NHR, N(R)₂, R₈-N(R₁₀)(R₁₁) (e.g., CH₂-NH₂, CH₂-N(CH₃)₂), R₉-R₈-N(R₁₀)(R₁₁) (e.g., C≡C-CH₂-NH₂), B(OH)₂, -OC(O)CF₃, -OCH₂Ph, NHC(O)-R₁₀ (e.g., NHC(O)CH₃), NHC(O)-N(R₁₀)(R₁₁) (e.g., NHC(O)N(CH₃)₂), COOH, -C(O)Ph, C(O)O-R₁₀ (e.g., C(O)O-CH₃, C(O)O-CH(CH₃)₂, C(O)O-CH₂CH₃), R₈-C(O)-R₁₀ (e.g., CH₂C(O)CH₃), C(O)H, C(O)-R₁₀ (e.g., C(O)-CH₃, C(O)-CH₂CH₃, C(O)-CH₂CH₂CH₃), C₁-C₅ linear or branched C(O)-haloalkyl (e.g., C(O)-CF₃), -C(O)NH₂, C(O)NHR, C(O)N(R₁₀)(R₁₁) (e.g., C(O)N(CH₃)₂), SO₂R, SO₂N(R₁₀)(R₁₁) (e.g., SO₂N(CH₃)₂, SO₂NHC(O)CH₃), C₁-C₅ linear or branched, substituted or unsubstituted alkyl (e.g., methyl, 2, 3, or 4-CH₂-C₆H₄-Cl, ethyl, propyl, iso-propyl, t-Bu, iso-butyl, pentyl, benzyl), C₁-C₅ linear or branched, substituted or unsubstituted alkenyl (e.g., CH=C(Ph)₂), C₁-C₅ linear or branched or C₃-C₈ cyclic haloalkyl (e.g., CF₃, CF₂CH₃, CH₂CF₃, CF₂CH₂CH₃, CH₂CH₂CF₃, CF₂CH(CH₃)₂, CF(CH₃)-CH(CH₃)₂), C₁-C₅ linear or branched or C₃-C₈ cyclic alkoxy (e.g. methoxy, ethoxy, propoxy, isopropoxy, O-CH₂-cyclopropyl, O-cyclobutyl, O-cyclopentyl, O-cyclohexyl, 1-butoxy, 2-butoxy, O-tBu), optionally wherein at least one methylene group (CH₂) in the alkoxy is replaced with an oxygen atom (e.g., O-1-oxacyclobutyl, O-2-oxacyclobutyl), C₁-C₅ linear or branched thioalkoxy, C₁-C₅ linear or branched haloalkoxy (e.g., OCF₃, OCHF₂), C₁-C₅ linear or branched alkoxyalkyl, substituted or unsubstituted C₃-C₈ cycloalkyl (e.g., cyclopropyl, cyclopentyl), substituted or unsubstituted C₃-C₈ heterocyclic ring (e.g., 3-methyl-4H-1,2,4-triazole, 5-methyl-1,2,4-oxadiazole, thiophene, oxazole, oxadiazole, imidazole, furane, triazole, tetrazole, pyridine (2, 3, or 4-pyridine), 3-methyl-2-pyridine, pyrimidine, pyrazine, oxacyclobutane (1 or 2-oxacyclobutane), protonated or deprotonated pyridine oxide), substituted or unsubstituted aryl (e.g., phenyl), substituted or unsubstituted benzyl (e.g., benzyl, 4-Cl-benzyl, 4-OH-benzyl);

or **R₂** and **R₁** are joined to form a C₃-C₈ substituted or unsubstituted, carbocyclic or heterocyclic ring;

R₃ is H, F, Cl, Br, I, OH, SH, =O, R₈-OH (e.g., CH₂-OH), R₈-SH, -R₈-O-R₁₀, (e.g., CH₂-O-CH₃) CF₃, CD₃, OCD₃, CN, NO₂, -CH₂CN, -R₈CN, NH₂, NHNH₂, NHR, N(R)₂, R₈-N(R₁₀)(R₁₁) (e.g., CH₂-NH₂, CH₂-N(CH₃)₂), R₉-R₈-N(R₁₀)(R₁₁), B(OH)₂, -OC(O)CF₃, -OCH₂Ph, NHC(O)OBz, -NHC(O)-R₁₀ (e.g., NHC(O)CH₃), NHC(O)-N(R₁₀)(R₁₁) (e.g., NHC(O)N(CH₃)₂), COOH, -C(O)Ph, C(O)O-R₁₀ (e.g., C(O)O-CH₃, C(O)O-CH₂CH₃), R₈-C(O)-R₁₀ (e.g., CH₂C(O)CH₃), C(O)H, C(O)-R₁₀ (e.g., C(O)-CH₃, C(O)-CH₂CH₃, C(O)-CH₂CH₂CH₃), C₁-C₅ linear or branched C(O)-haloalkyl (e.g., C(O)-CF₃), -C(O)NH₂, C(O)NHR, C(O)N(R₁₀)(R₁₁) (e.g., C(O)N(CH₃)₂), SO₂R, SO₂N(R₁₀)(R₁₁) (e.g., SO₂N(CH₃)₂), C₁-C₅ linear or branched, substituted or unsubstituted alkyl (e.g., methyl, C(OH)(CH₃)(Ph), ethyl,

propyl, iso-propyl, t-Bu, iso-butyl, pentyl), C₁-C₅ linear or branched or C₃-C₈ cyclic haloalkyl (e.g., CF₃, CF₂CH₃, CF₂-cyclobutyl, CH₂CF₃, CF₂CH₂CH₃, CH₂CH₂CF₃, CF₂CH(CH₃)₂, CF(CH₃)-CH(CH₃)₂), C₁-C₅ linear or branched or C₃-C₈ cyclic alkoxy (e.g. methoxy, ethoxy, propoxy, isopropoxy, O-CH₂-cyclopropyl), C₁-C₅ linear or branched thioalkoxy, C₁-C₅ linear or branched haloalkoxy, C₁-C₅ linear or branched alkoxyalkyl, substituted or unsubstituted C₃-C₈ cycloalkyl (e.g., cyclopropyl, cyclopentyl), substituted or unsubstituted C₃-C₈ heterocyclic ring (e.g., 3-methyl-4H-1,2,4-triazole, 5-methyl-1,2,4-oxadiazole, thiophene, oxazole, isoxazole, imidazole, furane, triazole, pyridine (2, 3, or 4-pyridine), pyrimidine, pyrazine, oxacyclobutane (1 or 2-oxacyclobutane)), substituted or unsubstituted aryl (e.g., phenyl);

or **R₃** and **R₂** are joined to form a C₃-C₈ substituted or unsubstituted, carbocyclic or heterocyclic ring (e.g., cyclopropyl);

R₄ is H, F, Cl, Br, I, OH, SH, =O, =NH-OH, R₈-OH (e.g., CH₂-OH), R₈-SH, -R₈-O-R₁₀, (e.g., CH₂-O-CH₃) CF₃, CD₃, OCD₃, CN, NO₂, -CH₂CN, -R₈CN, NH₂, NHNH₂, NHR, N(R)₂, R₈-N(R₁₀)(R₁₁) (e.g., CH₂-NH₂, CH₂-N(CH₃)₂) R₉-R₈-N(R₁₀)(R₁₁), B(OH)₂, -OC(O)CF₃, -OCH₂Ph, NHC(O)OBz, -NHC(O)-R₁₀ (e.g., NHC(O)H, NHC(O)CH₃), NHC(O)-N(R₁₀)(R₁₁) (e.g., NHC(O)N(CH₃)₂), COOH, -C(O)Ph, C(O)O-R₁₀ (e.g. C(O)O-CH₃, C(O)O-CH₂CH₃), R₈-C(O)-R₁₀ (e.g., CH₂C(O)CH₃), C(O)H, C(O)-R₁₀ (e.g., C(O)-CH₃, C(O)-CH₂CH₃, C(O)-CH₂CH₂CH₃), C₁-C₅ linear or branched C(O)-haloalkyl (e.g., C(O)-CF₃), -C(O)NH₂, C(O)NHR, C(O)N(R₁₀)(R₁₁) (e.g., C(O)N(CH₃)₂), SO₂R, SO₂N(R₁₀)(R₁₁) (e.g., SO₂N(CH₃)₂), C₁-C₅ linear or branched, substituted or unsubstituted alkyl (e.g., methyl, C(OH)(CH₃)(Ph), ethyl, propyl, iso-propyl, t-Bu, iso-butyl, pentyl), C₁-C₅ linear or branched or C₃-C₈ cyclic haloalkyl (e.g., CF₃, CF₂CH₃, CF₂-cyclobutyl, CH₂CF₃, CF₂CH₂CH₃, CH₂CH₂CF₃, CF₂CH(CH₃)₂, CF(CH₃)-CH(CH₃)₂), C₁-C₅ linear or branched or C₃-C₈ cyclic alkoxy (e.g. methoxy, ethoxy, propoxy, isopropoxy, O-CH₂-cyclopropyl), C₁-C₅ linear or branched thioalkoxy, C₁-C₅ linear or branched haloalkoxy, C₁-C₅ linear or branched alkoxyalkyl, substituted or unsubstituted C₃-C₈ cycloalkyl (e.g., cyclopropyl, cyclopentyl), substituted or unsubstituted C₃-C₈ heterocyclic ring (e.g., 3-methyl-4H-1,2,4-triazole, 5-methyl-1,2,4-oxadiazole, thiophene, oxazole, isoxazole, imidazole, furane, triazole, pyridine (2, 3, or 4-pyridine), pyrimidine, pyrazine, oxacyclobutane (1 or 2-oxacyclobutane), indole), substituted or unsubstituted aryl (e.g., phenyl);

or **R₃** and **R₄** are joined together to form ring A as defined above (e.g., cyclopropyl, 5-methyloxazolidin-2-one);

R₅ is H, C₁-C₅ linear or branched, substituted or unsubstituted alkyl (e.g., methyl, CH₂SH, ethyl, butyl, CH₂-CCH, iso-propyl, CH₂-C(O)-OCH₃), C₂-C₅ linear or branched, substituted or unsubstituted alkenyl, C₂-C₅ linear or branched, substituted or unsubstituted alkynyl (e.g., CCH, CH₂-CCH), C₁-C₅ linear or branched haloalkyl (e.g., CF₃, CF₂CH₃, CH₂CF₃, CF₂CH₂CH₃, CH₂CH₂CF₃, CF₂CH(CH₃)₂, CF(CH₃)-CH(CH₃)₂), R₈-aryl (e.g., CH₂-Ph), C(=CH₂)-R₁₀ (e.g., C(=CH₂)-C(O)-OCH₃, C(=CH₂)-CN), substituted or unsubstituted alkyl sulfone (e.g., SO₂-CH₂-cyclopentyl), substituted or unsubstituted aryl (e.g., phenyl), substituted or unsubstituted heteroaryl (e.g., pyridine (2, 3, and 4-pyridine));

R₈ is [CH₂]_p

wherein **p** is between 1 and 10;

R₉ is [CH]_q, [C]_q

wherein **q** is between 2 and 10;

R₁₀ and **R₁₁** are each independently H, CN, C₁-C₅ linear or branched alkyl (e.g., methyl, ethyl),
5 C(O)R (e.g., C(O)(OCH₃)), or S(O)₂R;

or **R₁₀** and **R₁₁** are joined to form a substituted or unsubstituted C₃-C₈ heterocyclic ring (e.g., piperazine, piperidine),

R is H, C₁-C₅ linear or branched alkyl (e.g., methyl, ethyl), C₁-C₅ linear or branched alkoxy (e.g., methoxy), phenyl, aryl or heteroaryl, or two gem R substituents are joined together to form a 5 or
10 6 membered heterocyclic ring;


m is an integer number between 1 and 5 (e.g., 1 or 2);

n is an integer number between 0 and 5 (e.g., 0, 1, 2 or 3);

X₁ is S, O, N-OH, CH₂, C(R)₂ or N-OMe;

X₂ is S, O, N-OH, CH₂, C(R)₂ or N-OMe;

15 or **X₂** together with the carbon next to **X₁** are joined to form ring B, represented by the following structure: (in such case, **X₁** is **X₇**):



wherein

20 **X₄**, **X₅**, **X₆** and **X₇** are each independently C or N, wherein if any of **X₄**, **X₅**, **X₆** and **X₇** is N, then the respective substitution **R₉₀**, **R₆₀**, **R₇₀** or **R₈₀** is absent;

R₆₀, **R₈₀** and **R₉₀** are each independently selected from: H, F, Cl, Br, I, OH, SH, R₈-OH, R₈-SH, -R₈-O-R₁₀, CF₃, CN, NO₂, NH₂, NHR, N(R)₂, R₈-N(R₁₀)(R₁₁), -OC(O)CF₃, -OCH₂Ph, NHC(O)OBz, -NHC(O)-R₁₀, COOH, -C(O)Ph, C(O)O-R₁₀, C(O)H, C(O)-R₁₀, C₁-C₅
25 linear or branched C(O)-haloalkyl, -C(O)NH₂, C(O)NHR, C(O)N(R₁₀)(R₁₁), SO₂R, SO₂N(R₁₀)(R₁₁), C₁-C₅ linear or branched, substituted or unsubstituted alkyl, C₁-C₅ linear or branched or C₃-C₈ cyclic haloalkyl, C₁-C₅ linear or branched or C₃-C₈ cyclic alkoxy, C₁-C₅ linear or branched thioalkoxy, C₁-C₅ linear or branched haloalkoxy, C₁-C₅ linear or branched alkoxyalkyl, substituted or unsubstituted C₃-C₈ cycloalkyl, substituted or unsubstituted C₃-C₈
30 heterocyclic ring, substituted or unsubstituted aryl;

R₇₀ is selected from: H, F, Cl, Br, I, OH, SH, R₈-OH, R₈-SH, -R₈-O-R₁₀, CF₃, CN, NO₂, NH₂, NHR, N(R)₂, R₈-N(R₁₀)(R₁₁), -OC(O)CF₃, -OCH₂Ph, NHC(O)OBz, -NHC(O)-R₁₀, COOH, -C(O)Ph, C(O)O-R₁₀, C(O)H, C(O)-R₁₀, C₁-C₅ linear or branched C(O)-haloalkyl, -

C(O)NH₂, C(O)NHR, C(O)N(R₁₀)(R₁₁), SO₂R, SO₂N(R₁₀)(R₁₁), C₁-C₅ linear or branched or C₃-C₈ cyclic haloalkyl, C₁-C₅ linear or branched or C₃-C₈ cyclic alkoxy, C₁-C₅ linear or branched thioalkoxy, C₁-C₅ linear or branched haloalkoxy, C₁-C₅ linear or branched alkoxyalkyl, substituted or unsubstituted C₃-C₈ cycloalkyl, substituted or unsubstituted C₃-C₈ heterocyclic ring, substituted or unsubstituted aryl,

X₃ is O, NH or N-R₅₀;

R₅₀ is H or C₁-C₅ linear or branched, substituted or unsubstituted alkyl;

or its agrochemically acceptable salt, stereoisomer, tautomer, hydrate, *N*-oxide, reverse amide analog, isotopic variant (e.g., deuterated analog), or any combination thereof.

10


[0036] In some embodiments, **R₃** and **R₄** are joined together to form ring **A**. In some embodiments, ring **B** is absent. In some embodiment **B** is pyridine. In some embodiments, **R₃** is OH, or NH₂. In some embodiments, **R₄** is NH₂, or OH. In some embodiments, **R₃** and **R₄** cannot both be NH₂. In some embodiments, if **R₃** is OH then **R₄** is NH₂. In some embodiments, if **R₃** is NH₂ then **R₄** is OH. In some
 15
 20
 25
 30
 35
 40
 45
 50
 55
 60
 65
 70
 75
 80
 85
 90
 95
 100
 105
 110
 115
 120
 125
 130
 135
 140
 145
 150
 155
 160
 165
 170
 175
 180
 185
 190
 195
 200
 205
 210
 215
 220
 225
 230
 235
 240
 245
 250
 255
 260
 265
 270
 275
 280
 285
 290
 295
 300
 305
 310
 315
 320
 325
 330
 335
 340
 345
 350
 355
 360
 365
 370
 375
 380
 385
 390
 395
 400
 405
 410
 415
 420
 425
 430
 435
 440
 445
 450
 455
 460
 465
 470
 475
 480
 485
 490
 495
 500
 505
 510
 515
 520
 525
 530
 535
 540
 545
 550
 555
 560
 565
 570
 575
 580
 585
 590
 595
 600
 605
 610
 615
 620
 625
 630
 635
 640
 645
 650
 655
 660
 665
 670
 675
 680
 685
 690
 695
 700
 705
 710
 715
 720
 725
 730
 735
 740
 745
 750
 755
 760
 765
 770
 775
 780
 785
 790
 795
 800
 805
 810
 815
 820
 825
 830
 835
 840
 845
 850
 855
 860
 865
 870
 875
 880
 885
 890
 895
 900
 905
 910
 915
 920
 925
 930
 935
 940
 945
 950
 955
 960
 965
 970
 975
 980
 985
 990
 995

[0037] In some embodiments, substitutions include: F, Cl, Br, I, OH, SH, C₁-C₅ linear or branched alkyl (e.g. methyl, ethyl), C₂-C₅ linear or branched alkenyl, C₂-C₅ linear or branched alkynyl (e.g. CCH), C₃-C₈ cycloalkyl, linear, branched or cyclic alkoxy, COOH, COO(R), NH₂, N(R)₂, CF₃, aryl, phenyl, heteroaryl (e.g., imidazole), C₃-C₈ cycloalkyl (e.g., cyclohexyl), CN, NO₂ or any combination thereof.

[0038] In some embodiments, the compound is any one of the compounds listed in **Table 1**. In various embodiments, the compound is an herbicidal compound. In various embodiments, the compound is for


use in controlling the growth of undesired plants. In some embodiments, the compound is any one of the compounds listed in **Table 2**.

[0039] In various embodiments, this invention is directed to a compound represented by the structure of formula **I(b)**:



wherein

C_A and C_B are both chiral carbon centers, or C_A and C_B together with R_3 and R_4 are joined to form ring **A**, represented by the following structure:



R_1 , R_1' , R_2 , R_2' and R_{40} are each independently H, F, Cl, Br, I, OH, C_1 - C_5 linear or branched, substituted or unsubstituted alkyl (e.g., methyl, ethyl, propyl, iso-propyl, t-Bu, iso-butyl, pentyl, benzyl), or $C(O)$ - R_{10} (e.g., $C(O)$ - CH_3);

R_3 is OH, F, SH, R_8 -OH (e.g., CH_2 -OH), NH_2 , $NHNH_2$, NHR , $N(R)_2$, $NHC(O)OBz$, $-NHC(O)$ - R_{10} (e.g., $NHC(O)CH_3$), C_1 - C_5 linear or branched, substituted or unsubstituted alkyl (e.g., methyl, ethyl, propyl, iso-propyl, t-Bu, iso-butyl, pentyl), substituted or unsubstituted C_3 - C_8 cycloalkyl, C_1 - C_5 linear or branched or C_3 - C_8 cyclic haloalkyl, C_1 - C_5 linear or branched or C_3 - C_8 cyclic alkoxy, substituted or unsubstituted C_3 - C_8 heterocyclic ring, or substituted or unsubstituted aryl;

or R_3 and R_2 are joined to form a C_3 - C_8 substituted or unsubstituted, carbocyclic or heterocyclic ring (e.g., cyclopropyl);

R_4 is NH_2 , OH, $NHNH_2$, NHR , $N(R)_2$, $-NHC(O)$ - R_{10} , $NHC(O)H$, $NHC(O)CH_3$, C_1 - C_5 linear or branched, substituted or unsubstituted alkyl, C_1 - C_5 linear or branched or C_3 - C_8 cyclic haloalkyl, C_1 - C_5 linear or branched or C_3 - C_8 cyclic alkoxy, substituted or unsubstituted C_3 - C_8 cycloalkyl, substituted or unsubstituted C_3 - C_8 heterocyclic ring, or substituted or unsubstituted aryl;

or R_3 and R_4 are joined together to form ring **A** as described above (e.g., cyclopropyl, 5-methyloxazolidin-2-one [1,3]dioxole, furan-2(3H)-one, benzene, cyclopentane, imidazole);

R_5 is H, C_1 - C_5 linear or branched, substituted or unsubstituted alkyl (e.g., methyl, CH_2SH , ethyl, butyl, CH_2 -CCH, iso-propyl, CH_2 - $C(O)$ - OCH_3), C_2 - C_5 linear or branched, substituted or unsubstituted

alkenyl, C₂-C₅ linear or branched, substituted or unsubstituted alkynyl (e.g., CCH, CH₂-CCH), C₁-C₅ linear or branched haloalkyl (e.g., CF₃, CF₂CH₃, CH₂CF₃, CF₂CH₂CH₃, CH₂CH₂CF₃, CF₂CH(CH₃)₂, CF(CH₃)-CH(CH₃)₂), R₈-aryl (e.g., CH₂-Ph), C(=CH₂)-R₁₀ (e.g., C(=CH₂)-C(O)-OCH₃, C(=CH₂)-CN), substituted or unsubstituted alkyl sulfone (e.g., SO₂-CH₂-cyclopentyl), substituted or unsubstituted aryl (e.g., phenyl), substituted or unsubstituted heteroaryl (e.g., pyridine (2, 3, and 4-pyridine));

R₈ is [CH₂]_p

wherein **p** is between 1 and 10;

R₁₀ and **R₁₁** are each independently H, CN, C₁-C₅ linear or branched alkyl (e.g., methyl, ethyl), C(O)R (e.g., C(O)(OCH₃)), or S(O)₂R;

or **R₁₀** and **R₁₁** are joined to form a substituted or unsubstituted C₃-C₈ heterocyclic ring (e.g., piperazine, piperidine),

R is H, C₁-C₅ linear or branched alkyl (e.g., methyl, ethyl), C₁-C₅ linear or branched alkoxy (e.g., methoxy), phenyl, aryl or heteroaryl, or two gem R substituents are joined together to form a 5 or 6 membered heterocyclic ring;


m is an integer number between 1 and 5 (e.g., 1 or 2);

n is an integer number between 0 and 5 (e.g., 0, 1, 2 or 3);

X₁ is S, O, or CH₂;

X₂ is S, O, or CH₂;

or **X₂** together with the carbon next to **X₁** are joined to form ring B, represented by the following structure (in such case, **X₁** is **X₇**):



B

wherein

X₄, **X₅**, **X₆** and **X₇** are each independently C or N, wherein if any of **X₄**, **X₅**, **X₆** and **X₇** is N, then the respective substitution **R₉₀**, **R₆₀**, **R₇₀** or **R₈₀** is absent;

R₆₀, **R₈₀** and **R₉₀** are each independently selected from: H, F, Cl, Br, I, OH, SH, R₈-OH, R₈-SH, -R₈-O-R₁₀, CF₃, CN, NO₂, NH₂, NHR, N(R)₂, R₈-N(R₁₀)(R₁₁), -OC(O)CF₃, -OCH₂Ph, NHC(O)OBz, -NHC(O)-R₁₀, COOH, -C(O)Ph, C(O)O-R₁₀, C(O)H, C(O)-R₁₀, C₁-C₅ linear or branched C(O)-haloalkyl, -C(O)NH₂, C(O)NHR, C(O)N(R₁₀)(R₁₁), SO₂R, SO₂N(R₁₀)(R₁₁), C₁-C₅ linear or branched, substituted or unsubstituted alkyl, C₁-C₅ linear or branched or C₃-C₈ cyclic haloalkyl, C₁-C₅ linear or branched or C₃-C₈ cyclic alkoxy, C₁-C₅ linear or branched thioalkoxy, C₁-C₅ linear or branched haloalkoxy, C₁-C₅ linear or branched

alkoxyalkyl, substituted or unsubstituted C₃-C₈ cycloalkyl, substituted or unsubstituted C₃-C₈ heterocyclic ring, substituted or unsubstituted aryl;

R₇₀ is selected from: H, F, Cl, Br, I, OH, SH, R₈-OH, R₈-SH, -R₈-O-R₁₀, CF₃, CN, NO₂, NH₂, NHR, N(R)₂, R₈-N(R₁₀)(R₁₁), -OC(O)CF₃, -OCH₂Ph, NHC(O)OBz, -NHC(O)-R₁₀,
 5 COOH, -C(O)Ph, C(O)O-R₁₀, C(O)H, C(O)-R₁₀, C₁-C₅ linear or branched C(O)-haloalkyl, -C(O)NH₂, C(O)NHR, C(O)N(R₁₀)(R₁₁), SO₂R, SO₂N(R₁₀)(R₁₁), C₁-C₅ linear or branched or C₃-C₈ cyclic haloalkyl, C₁-C₅ linear or branched or C₃-C₈ cyclic alkoxy, C₁-C₅ linear or branched thioalkoxy, C₁-C₅ linear or branched haloalkoxy, C₁-C₅ linear or branched alkoxyalkyl, substituted or unsubstituted C₃-C₈ cycloalkyl, substituted or unsubstituted C₃-C₈ heterocyclic
 10 ring, substituted or unsubstituted aryl;

X₃ is O, NH or N-R₅₀;

R₅₀ is H or C₁-C₅ linear or branched, substituted or unsubstituted alkyl;

or its agrochemically acceptable salt, stereoisomer, tautomer, hydrate, *N*-oxide, reverse amide analog, isotopic variant (e.g., deuterated analog), or any combination thereof.

15


[0040] In some embodiments, **R**₃ and **R**₄ are joined together to form ring **A**. In some embodiments, ring **B** is absent. In some embodiment **B** is pyridine. In some embodiments, **R**₃ is OH, or NH₂. In some embodiments, **R**₄ is NH₂, or OH. In some embodiments, **R**₃ and **R**₄ cannot both be NH₂. In some embodiments, if **R**₃ is OH then **R**₄ is NH₂. In some embodiments, if **R**₃ is NH₂ then **R**₄ is OH. In some
 20 embodiments, if **R**₃ is OH then **R**₄ is NH₂ and if **R**₃ is NH₂ then **R**₄ is OH. In some embodiments, if **R**₃ is OH and **R**₄ is NH₂, then **n** + **m** cannot be equal to 3. In some embodiments, the compound is compound **101**, **102**, **104**, **105**, **106**, **113**, **114**, **115**, **116**, **117**, **118**, **119**, or **120**; each represents a separate embodiment according to this invention. In some embodiments, ring **A** has two chiral centers. In some
 25 embodiments, the compound is not (6*R*,7*S*)-6-amino-7-hydroxyoctanoic acid. In some embodiments, the compound is not 5-((4*R*,5*S*)-5-methyl-2-oxooxazolidin-4-yl)pentanoic acid. In some embodiments, the compound is a substantially pure single stereoisomer. In some embodiments, **R**₁, **R**₁', **R**₂, **R**₂' and **R**₄₀ are each independently H or C₁-C₅ linear or branched, substituted or unsubstituted alkyl (e.g., methyl, ethyl). In some embodiments, **R**₁ and **R**₁' are both H. In some
 30 embodiments, **R**₂ is CH₃ or CH₂CH₃. In some embodiments, **R**₂' is H or CH₃. In some embodiments, **R**₄₀ is CH₃ or H. In some embodiments, **X**₁ is CH₂. In some embodiments, **X**₂ is CH₂. In some embodiments, **X**₃ is O, NH or N-CH₃. In some embodiments, **R**₅ is H, or C₁-C₅ linear or branched, substituted or unsubstituted alkyl, or substituted or unsubstituted alkyl sulfone. In some embodiments, **R**₅ is a substituted or unsubstituted alkyl. In some embodiments, **R**₅ is H. In some embodiments, **R**₅ is H, ethyl, butyl, CH₂-CCH, CH₂-C(O)-OCH₃ or SO₂-CH₂-cyclopentyl; each is a separate embodiment
 35 according to this invention. In some embodiments, **n** is 0, 1, 2 or 3. In some embodiments, **n** is 1. In some embodiments, **m** is 1 or 2. In some embodiments, **m** is 1.

[0041] In some embodiments, substitutions include but not limited to: F, Cl, Br, I, OH, SH, C₁-C₅ linear or branched alkyl (e.g. methyl, ethyl), C₂-C₅ linear or branched alkenyl, C₂-C₅ linear or branched alkynyl

(e.g. CCH), C₃-C₈ cycloalkyl, linear, branched or cyclic alkoxy, COOH, COO(R), NH₂, N(R)₂, CF₃, aryl, phenyl, heteroaryl (e.g., imidazole), C₃-C₈ cycloalkyl (e.g., cyclohexyl), CN, NO₂ or any combination thereof.

[0042] In some embodiments, the compound is any one of the compounds listed in **Table 1**. In various
5
embodiments, the compound is an herbicidal compound. In various embodiments, the compound is for
use in controlling the growth of undesired plants. In some embodiments, the compound is any one of
the compounds listed in **Table 2**.

[0043] In various embodiments, this invention is directed to a compound represented by the structure
10
of formula **I(c)**:



I(c)

wherein

C_A and **C_B** are both chiral carbon centers;

15
R₁, **R₁'**, **R₂**, **R₂'** and **R₄₀** are each independently H, F, Cl, Br, I, OH, C₁-C₅ linear or branched, substituted or unsubstituted alkyl (e.g., methyl, ethyl, propyl, iso-propyl, t-Bu, iso-butyl, pentyl, benzyl), or C(O)-R₁₀ (e.g., C(O)-CH₃);

R₃ is OH, F, R₈-OH (e.g., CH₂-OH), SH, NH₂, NHNH₂, NHR, N(R)₂, NHC(O)OBz, -NHC(O)-R₁₀ (e.g., NHC(O)CH₃), C₁-C₅ linear or branched, substituted or unsubstituted alkyl (e.g., methyl, ethyl,
20
propyl, iso-propyl, t-Bu, iso-butyl, pentyl), substituted or unsubstituted C₃-C₈ cycloalkyl, C₁-C₅ linear or branched or C₃-C₈ cyclic haloalkyl, C₁-C₅ linear or branched or C₃-C₈ cyclic alkoxy, substituted or unsubstituted C₃-C₈ heterocyclic ring, or substituted or unsubstituted aryl;

or **R₃** and **R₂** are joined to form a C₃-C₈ substituted or unsubstituted, carbocyclic or heterocyclic ring (e.g., cyclopropyl);

25
R₄ is OH, NH₂, NHNH₂, NHR, N(R)₂, -NHC(O)-R₁₀, NHC(O)H, NHC(O)CH₃, C₁-C₅ linear or branched, substituted or unsubstituted alkyl (e.g., methyl, C(OH)(CH₃)(Ph), ethyl, propyl, iso-propyl, t-Bu, iso-butyl, pentyl), C₁-C₅ linear or branched or C₃-C₈ cyclic haloalkyl, C₁-C₅ linear or branched or C₃-C₈ cyclic alkoxy, substituted or unsubstituted C₃-C₈ cycloalkyl, substituted or unsubstituted C₃-C₈ heterocyclic ring, or substituted or unsubstituted aryl;

30
R₅ is H, C₁-C₅ linear or branched, substituted or unsubstituted alkyl (e.g., methyl, CH₂SH, ethyl, butyl, CH₂-CCH, iso-propyl, CH₂-C(O)-OCH₃), C₂-C₅ linear or branched, substituted or unsubstituted alkenyl, C₂-C₅ linear or branched, substituted or unsubstituted alkynyl (e.g., CCH, CH₂-CCH), C₁-C₅ linear or branched haloalkyl (e.g., CF₃, CF₂CH₃, CH₂CF₃, CF₂CH₂CH₃, CH₂CH₂CF₃, CF₂CH(CH₃)₂, CF(CH₃)-CH(CH₃)₂), R₈-aryl (e.g., CH₂-Ph), C(=CH₂)-R₁₀ (e.g., C(=CH₂)-C(O)-OCH₃,
35
C(=CH₂)-CN), substituted or unsubstituted alkyl sulfone (e.g., SO₂-CH₂-cyclopentyl), substituted or

unsubstituted aryl (e.g., phenyl), substituted or unsubstituted heteroaryl (e.g., pyridine (2, 3, and 4-pyridine));

R₈ is [CH₂]_p

wherein **p** is between 1 and 10;

5 **R₁₀** and **R₁₁** are each independently H, CN, C₁-C₅ linear or branched alkyl (e.g., methyl, ethyl), C(O)R (e.g., C(O)(OCH₃)), or S(O)₂R;

or **R₁₀** and **R₁₁** are joined to form a substituted or unsubstituted C₃-C₈ heterocyclic ring (e.g., piperazine, piperidine),

10 **R** is H, C₁-C₅ linear or branched alkyl, C₁-C₅ linear or branched alkoxy, phenyl, aryl or heteroaryl, or two gem R substituents are joined together to form a 5 or 6 membered heterocyclic ring;

m is an integer number between 1 and 5 (e.g., 1 or 2);

n is an integer number between 0 and 5 (e.g., 0, 1, 2 or 3);

X₁ is S, O, N-OH, CH₂, C(R)₂ or N-OMe;

X₂ is S, O, N-OH, CH₂, C(R)₂ or N-OMe;

15 **X₃** is O, NH or N-R₅₀;

R₅₀ is H or C₁-C₅ linear or branched, substituted or unsubstituted alkyl;

or its agrochemically acceptable salt, stereoisomer, tautomer, hydrate, *N*-oxide, reverse amide analog, isotopic variant (e.g., deuterated analog), or any combination thereof.

20 [0044] In some embodiments, **R₃** is OH, or NH₂. In some embodiments, **R₄** is NH₂, or OH. In some embodiments, **R₃** and **R₄** cannot both be NH₂. In some embodiments, if **R₃** is OH then **R₄** is NH₂. In some embodiments, if **R₃** is NH₂ then **R₄** is OH. In some embodiments, if **R₃** is OH then **R₄** is NH₂ and if **R₃** is NH₂ then **R₄** is OH. In some embodiments, if **R₃** is OH and **R₄** is NH₂, then **n + m** cannot be equal to 3. In some embodiments, the compound is compound **101, 102, 104, 105, 106, 113, 114, 115,**

25 **116, 117, 118, 119, or 120**; each represents a separate embodiment according to this invention. In some embodiments, the compound is not (6*R*,7*S*)-6-amino-7-hydroxyoctanoic acid. In some embodiments, the compound is a substantially pure single stereoisomer. In some embodiments, **R₁, R₁'**, **R₂, R₂'** and **R₄₀** are each independently H or C₁-C₅ linear or branched, substituted or unsubstituted alkyl (e.g., methyl, ethyl). In some embodiments, **R₁** and **R₁'** are both H. In some embodiments, **R₂** is CH₃ or


30 CH₂CH₃. In some embodiments, **R₂'** is H or CH₃. In some embodiments, **R₄₀** is CH₃ or H. In some embodiments, **X₁** is CH₂. In some embodiments, **X₂** is CH₂. In some embodiments, **X₃** is O, NH or N-CH₃. In some embodiments, **R₅** is H, or C₁-C₅ linear or branched, substituted or unsubstituted alkyl, or substituted or unsubstituted alkyl sulfone. In some embodiments, **R₅** is a substituted or unsubstituted alkyl. In some embodiments, **R₅** is H. In some embodiments, **R₅** is H, ethyl, butyl, CH₂-CCH, CH₂-

35 C(O)-OCH₃ or SO₂-CH₂-cyclopentyl; each is a separate embodiment according to this invention. In some embodiments, **n** is 0, 1, 2 or 3. In some embodiments, **n** is 1. In some embodiments, **m** is 1 or 2. In some embodiments, **m** is 1.

[0045] In some embodiments, substitutions include but not limited to: F, Cl, Br, I, OH, SH, C₁-C₅ linear or branched alkyl (e.g. methyl, ethyl), C₂-C₅ linear or branched alkenyl, C₂-C₅ linear or branched alkynyl (e.g. CCH), C₃-C₈ cycloalkyl, linear, branched or cyclic alkoxy, COOH, COO(R), NH₂, N(R)₂, CF₃, aryl, phenyl, heteroaryl (e.g., imidazole), C₃-C₈ cycloalkyl (e.g., cyclohexyl), CN, NO₂ or any combination thereof.

[0046] In various embodiments, the compound is an herbicidal compound. In various embodiments, the compound is for use in controlling the growth of undesired plants.

[0047] In various embodiments, this invention is directed to a compound represented by the structure of formula **I(d)**:



I(d)

wherein

R₁, R₁', R₂, R₂' are each independently H, C₁-C₅ linear or branched, substituted or unsubstituted alkyl (e.g., methyl, ethyl, propyl, iso-propyl, t-Bu, iso-butyl, pentyl, benzyl), or C(O)-R₁₀ (e.g., C(O)-CH₃);

R₅ is H, C₁-C₅ linear or branched, substituted or unsubstituted alkyl (e.g., methyl, CH₂SH, ethyl, butyl, CH₂-CCH, iso-propyl, CH₂-C(O)-OCH₃), C₂-C₅ linear or branched, substituted or unsubstituted alkenyl, C₂-C₅ linear or branched, substituted or unsubstituted alkynyl (e.g., CCH, CH₂-CCH), C₁-C₅ linear or branched haloalkyl (e.g., CF₃, CF₂CH₃, CH₂CF₃, CF₂CH₂CH₃, CH₂CH₂CF₃, CF₂CH(CH₃)₂, CF(CH₃)-CH(CH₃)₂), R₈-aryl (e.g., CH₂-Ph), C(=CH₂)-R₁₀ (e.g., C(=CH₂)-C(O)-OCH₃, C(=CH₂)-CN), substituted or unsubstituted alkyl sulfone (e.g., SO₂-CH₂-cyclopentyl), substituted or unsubstituted aryl (e.g., phenyl), substituted or unsubstituted heteroaryl (e.g., pyridine (2, 3, and 4-pyridine));

R₁₀ is H, CN, C₁-C₅ linear or branched alkyl, C(O)R, or S(O)₂R;

R is H, C₁-C₅ linear or branched alkyl, C₁-C₅ linear or branched alkoxy, phenyl, aryl or heteroaryl, or two gem R substituents are joined together to form a 5 or 6 membered heterocyclic ring;


m is an integer number between 1 and 5 (e.g., 1 or 2);

n is an integer number between 0 and 5 (e.g., 0, 1, 2 or 3);

X₁ is S, O, N-OH, CH₂, C(R)₂ or N-OMe;

X₂ is S, O, N-OH, CH₂, C(R)₂ or N-OMe;

or **X₂** together with the carbon next to **X₁** are joined to form ring B, represented by the following structure (in such case, **X₁** is **X₇**):



wherein

5 **X₄**, **X₅**, **X₆** and **X₇** are each independently C or N, wherein if any of **X₄**, **X₅**, **X₆** and **X₇** is N, then the respective substitution **R₉₀**, **R₆₀**, **R₇₀** or **R₈₀** is absent;

R₆₀, **R₇₀**, **R₈₀** and **R₉₀** are each independently selected from: H, F, Cl, Br, I, OH, SH, R₈-OH, R₈-SH, -R₈-O-R₁₀, CF₃, CN, NO₂, NH₂, NHR, N(R)₂, R₈-N(R₁₀)(R₁₁), -OC(O)CF₃, -OCH₂Ph, NHC(O)OBz, -NHC(O)-R₁₀, COOH, -C(O)Ph, C(O)O-R₁₀, C(O)H, C(O)-R₁₀, C₁-C₅ linear or branched C(O)-haloalkyl, -C(O)NH₂, C(O)NHR, C(O)N(R₁₀)(R₁₁), SO₂R, SO₂N(R₁₀)(R₁₁), C₁-C₅ linear or branched, substituted or unsubstituted alkyl, C₁-C₅ linear or branched or C₃-C₈ cyclic haloalkyl, C₁-C₅ linear or branched or C₃-C₈ cyclic alkoxy, C₁-C₅ linear or branched thioalkoxy, C₁-C₅ linear or branched haloalkoxy, C₁-C₅ linear or branched alkoxyalkyl, substituted or unsubstituted C₃-C₈ cycloalkyl, substituted or unsubstituted C₃-C₈ heterocyclic ring, substituted or unsubstituted aryl;

15 **X₃** is O, NH or N-R₅₀;

R₅₀ is H or C₁-C₅ linear or branched, substituted or unsubstituted alkyl;


 or its agrochemically acceptable salt, stereoisomer, tautomer, hydrate, *N*-oxide, reverse amide analog, isotopic variant (e.g., deuterated analog), or any combination thereof.

20 [0048] In some embodiments, the compound is not 5-((*4R,5S*)-5-methyl-2-oxooxazolidin-4-yl)pentanoic acid. In some embodiments, the compound is a substantially pure single stereoisomer. In some embodiments, **R₁**, **R₁'**, **R₂**, **R₂'** and **R₄₀** are each independently H or C₁-C₅ linear or branched, substituted or unsubstituted alkyl (e.g., methyl, ethyl). In some embodiments, **R₁** and **R₁'** are both H. In some embodiments, **R₂** is CH₃ or CH₂CH₃. In some embodiments, **R₂'** is H or CH₃. In some embodiments, **R₄₀** is CH₃ or H. In some embodiments, **X₁** is CH₂. In some embodiments, **X₂** is CH₂. In some embodiments, **X₃** is O, NH or N-CH₃. In some embodiments, **R₅** is H, or C₁-C₅ linear or branched, substituted or unsubstituted alkyl, or substituted or unsubstituted alkyl sulfone. In some embodiments, **R₅** is a substituted or unsubstituted alkyl. In some embodiments, **R₅** is H. In some embodiments, **R₅** is H, ethyl, butyl, CH₂-CCH, CH₂-C(O)-OCH₃ or SO₂-CH₂-cyclopentyl; each is a separate embodiment according to this invention. In some embodiments, n is 0, 1, 2 or 3. In some embodiments, n is 1. In some embodiments, m is 1 or 2. In some embodiments, m is 1. In some embodiments, the compound is compound **105**, **106**; each represents a separate embodiment according to this invention.

[0049] In some embodiments, substitutions include but not limited to: F, Cl, Br, I, OH, SH, C₁-C₅ linear or branched alkyl (e.g. methyl, ethyl), C₂-C₅ linear or branched alkenyl, C₂-C₅ linear or branched alkynyl (e.g. CCH), C₃-C₈ cycloalkyl, linear, branched or cyclic alkoxy, COOH, COO(R), NH₂, N(R)₂, CF₃, aryl, phenyl, heteroaryl (e.g., imidazole), C₃-C₈ cycloalkyl (e.g., cyclohexyl), CN, NO₂ or any combination thereof.

[0050] In various embodiments, the compound is an herbicidal compound. In various embodiments, the compound is for use in controlling the growth of undesired plants.

[0051] In various embodiments, this invention is directed to a compound represented by the structure of formula I(e):



I(e)

wherein

R₁, R₁', **R₂**, **R₂'** are each independently H, F, Cl, Br, I, OH, C₁-C₅ linear or branched, substituted or unsubstituted alkyl (e.g., methyl, ethyl, propyl, iso-propyl, t-Bu, iso-butyl, pentyl, benzyl), or C(O)-R₁₀ (e.g., C(O)-CH₃);

R₃ is OH, F, SH, R₈-OH (e.g., CH₂-OH), NH₂, NHNH₂, NHR, N(R)₂, NHC(O)OBz, -NHC(O)-R₁₀ (e.g., NHC(O)CH₃), C₁-C₅ linear or branched, substituted or unsubstituted alkyl (e.g., methyl, ethyl, propyl, iso-propyl, t-Bu, iso-butyl, pentyl), substituted or unsubstituted C₃-C₈ cycloalkyl, C₁-C₅ linear or branched or C₃-C₈ cyclic haloalkyl, C₁-C₅ linear or branched or C₃-C₈ cyclic alkoxy, substituted or unsubstituted C₃-C₈ heterocyclic ring, or substituted or unsubstituted aryl;

R₄ is NH₂, OH, NHNH₂, NHR, N(R)₂, -NHC(O)-R₁₀, NHC(O)H, NHC(O)CH₃, C₁-C₅ linear or branched, substituted or unsubstituted alkyl, C₁-C₅ linear or branched or C₃-C₈ cyclic haloalkyl, C₁-C₅ linear or branched or C₃-C₈ cyclic alkoxy, substituted or unsubstituted C₃-C₈ cycloalkyl, substituted or unsubstituted C₃-C₈ heterocyclic ring, or substituted or unsubstituted arylf;

or **R₃** and **R₄** are joined together to form a 5 or 6 membered substituted or unsubstituted, aliphatic or aromatic, carbocyclic or heterocyclic ring (e.g., cyclopropyl, 5-methyloxazolidin-2-one, [1,3]dioxole, furan-2(3H)-one, benzene, cyclopentane, imidazole);

R₅ is H, C₁-C₅ linear or branched, substituted or unsubstituted alkyl (e.g., methyl, CH₂SH, ethyl, butyl, CH₂-CCH, iso-propyl, CH₂-C(O)-OCH₃), C₂-C₅ linear or branched, substituted or unsubstituted alkenyl, C₂-C₅ linear or branched, substituted or unsubstituted alkynyl (e.g., CCH, CH₂-CCH), C₁-C₅ linear or branched haloalkyl (e.g., CF₃, CF₂CH₃, CH₂CF₃, CF₂CH₂CH₃, CH₂CH₂CF₃, CF₂CH(CH₃)₂, CF(CH₃)-CH(CH₃)₂), R₈-aryl (e.g., CH₂-Ph), C(=CH₂)-R₁₀ (e.g., C(=CH₂)-C(O)-OCH₃,

C(=CH₂)-CN), substituted or unsubstituted alkyl sulfone (e.g., SO₂-CH₂-cyclopentyl), substituted or unsubstituted aryl (e.g., phenyl), substituted or unsubstituted heteroaryl (e.g., pyridine (2, 3, and 4-pyridine));

R₈ is [CH₂]_p

5 wherein **p** is between 1 and 10;

R₁₀ and **R₁₁** are each independently H, CN, C₁-C₅ linear or branched alkyl (e.g., methyl, ethyl), C(O)R (e.g., C(O)(OCH₃)), or S(O)₂R;

or **R₁₀** and **R₁₁** are joined to form a substituted or unsubstituted C₃-C₈ heterocyclic ring (e.g., piperazine, piperidine),

10 **R** is H, C₁-C₅ linear or branched alkyl, C₁-C₅ linear or branched alkoxy, phenyl, aryl or heteroaryl, or two gem R substituents are joined together to form a 5 or 6 membered heterocyclic ring;

n is 0, 1, 2 or 3;

R₆₀, **R₇₀**, **R₈₀** and **R₉₀** are each independently selected from: H, F, Cl, Br, I, OH, SH, R₈-OH, R₈-SH, -R₈-O-R₁₀, CF₃, CN, NO₂, NH₂, NHR, N(R)₂, R₈-N(R₁₀)(R₁₁), -OC(O)CF₃, -OCH₂Ph, 15 NHC(O)OBz, -NHC(O)-R₁₀, COOH, -C(O)Ph, C(O)O-R₁₀, C(O)H, C(O)-R₁₀, C₁-C₅ linear or branched C(O)-haloalkyl, -C(O)NH₂, C(O)NHR, C(O)N(R₁₀)(R₁₁), SO₂R, SO₂N(R₁₀)(R₁₁), C₁-C₅ linear or branched, substituted or unsubstituted alkyl, C₁-C₅ linear or branched or C₃-C₈ cyclic haloalkyl, C₁-C₅ linear or branched or C₃-C₈ cyclic alkoxy, C₁-C₅ linear or branched thioalkoxy, substituted or unsubstituted C₃-C₈ cycloalkyl, substituted or unsubstituted C₃-C₈ heterocyclic ring, substituted or 20 unsubstituted aryl;

X₃ is O, NH or N-R₅₀;

R₅₀ is H or C₁-C₅ linear or branched, substituted or unsubstituted alkyl;

X₄, **X₅**, **X₆** and **X₇** are each independently C or N, wherein if any of **X₄**, **X₅**, **X₆** and **X₇** is N, then the respective substitution **R₉₀**, **R₆₀**, **R₇₀** or **R₈₀** is absent;


25 or its agrochemically acceptable salt, stereoisomer, tautomer, hydrate, *N*-oxide, reverse amide analog, isotopic variant (e.g., deuterated analog), or any combination thereof.

[0052] In some embodiments, if **R₄** is NH₂, then **R₃** cannot be OH. In some embodiments, if **R₃** is OH and **R₄** is NH₂, **n** cannot be 1. In some embodiments, if **R₃** is OH and **R₄** is NH₂, **X₇**-**R₉₀** cannot be CH.

30 [0053] In some embodiments, substitutions include but not limited to: F, Cl, Br, I, OH, SH, C₁-C₅ linear or branched alkyl (e.g. methyl, ethyl), C₂-C₅ linear or branched alkenyl, C₂-C₅ linear or branched alkynyl (e.g. CCH), C₃-C₈ cycloalkyl, linear, branched or cyclic alkoxy, COOH, COO(R), NH₂, N(R)₂, CF₃, aryl, phenyl, heteroaryl (e.g., imidazole), C₃-C₈ cycloalkyl (e.g., cyclohexyl), CN, NO₂ or any combination thereof.

35 [0054] In various embodiments, the compound is an herbicidal compound. In various embodiments, the compound is for use in controlling the growth of undesired plants. In some embodiments, the compound is compound **108**, **109**, **141**, **142**, **172**, **173**, **174**, **175**, **176**, **177** or **178**; each represents a separate embodiment according to this invention.

[0055] In various embodiments, this invention is directed to a compound represented by the structure of formula **I(f)**:



I(f)

5 **R₁, R₁'**, **R₂, R₂'** are each independently H, F, Cl, Br, I, OH, C₁-C₅ linear or branched, substituted or unsubstituted alkyl (e.g., methyl, ethyl, propyl, iso-propyl, t-Bu, iso-butyl, pentyl, benzyl), or C(O)-R₁₀ (e.g., C(O)-CH₃);

R₃ is SH, NHNH₂, or NHC(O)OBz;

R₄ is NH₂, OH, NHNH₂, =NH-OH, =O, =N-NH₂, NHR, N(R)₂, -NHC(O)-R₁₀, NHC(O)H,
10 NHC(O)CH₃, C₁-C₅ linear or branched, substituted or unsubstituted alkyl, C₁-C₅ linear or branched or C₃-C₈ cyclic haloalkyl, C₁-C₅ linear or branched or C₃-C₈ cyclic alkoxy, substituted or unsubstituted C₃-C₈ cycloalkyl, substituted or unsubstituted C₃-C₈ heterocyclic ring, or substituted or unsubstituted aryl;

R₅ is H, C₁-C₅ linear or branched, substituted or unsubstituted alkyl (e.g., methyl, CH₂SH, ethyl, butyl, CH₂-CCH, iso-propyl, CH₂-C(O)-OCH₃), C₂-C₅ linear or branched, substituted or unsubstituted
15 alkenyl, C₂-C₅ linear or branched, substituted or unsubstituted alkynyl (e.g., CCH, CH₂-CCH), C₁-C₅ linear or branched haloalkyl (e.g., CF₃, CF₂CH₃, CH₂CF₃, CF₂CH₂CH₃, CH₂CH₂CF₃, CF₂CH(CH₃)₂, CF(CH₃)-CH(CH₃)₂), R₈-aryl (e.g., CH₂-Ph), C(=CH₂)-R₁₀ (e.g., C(=CH₂)-C(O)-OCH₃, C(=CH₂)-CN), substituted or unsubstituted alkyl sulfone (e.g., SO₂-CH₂-cyclopentyl), substituted or
20 unsubstituted aryl (e.g., phenyl), substituted or unsubstituted heteroaryl (e.g., pyridine (2, 3, and 4-pyridine));

R₈ is [CH₂]_p

 wherein **p** is between 1 and 10;

R₁₀ and **R₁₁** are each independently H, CN, C₁-C₅ linear or branched alkyl, C(O)R, or S(O)₂R;
 or **R₁₀** and **R₁₁** are joined to form a substituted or unsubstituted C₃-C₈ heterocyclic ring,

25 **R** is H, C₁-C₅ linear or branched alkyl, C₁-C₅ linear or branched alkoxy, phenyl, aryl or heteroaryl, or two gem R substituents are joined together to form a 5 or 6 membered heterocyclic ring;


m is an integer number between 1 and 5 (e.g., 1 or 2);

n is an integer number between 0 and 5 (e.g., 0, 1, 2 or 3);

X₁ is S, O, N-OH, CH₂, C(R)₂ or N-OMe;

30 **X₂** is S, O, N-OH, CH₂, C(R)₂ or N-OMe;

 or **X₂** together with the carbon next to **X₁** are joined to form ring B, represented by the following structure (in such case, **X₁** is **X₇**):



wherein

X₄, **X₅**, **X₆** and **X₇** are each independently C or N, wherein if any of **X₄**, **X₅**, **X₆** and **X₇** is N, then the respective substitution **R₉₀**, **R₆₀**, **R₇₀** or **R₈₀** is absent;

R₆₀, **R₇₀**, **R₈₀** and **R₉₀** are each independently selected from: H, F, Cl, Br, I, OH, SH, R₈-OH, R₈-SH, -R₈-O-R₁₀, CF₃, CN, NO₂, NH₂, NHR, N(R)₂, R₈-N(R₁₀)(R₁₁), -OC(O)CF₃, -OCH₂Ph, NHC(O)OBz, -NHC(O)-R₁₀, COOH, -C(O)Ph, C(O)O-R₁₀, C(O)H, C(O)-R₁₀, C₁-C₅ linear or branched C(O)-haloalkyl, -C(O)NH₂, C(O)NHR, C(O)N(R₁₀)(R₁₁), SO₂R, SO₂N(R₁₀)(R₁₁), C₁-C₅ linear or branched, substituted or unsubstituted alkyl, C₁-C₅ linear or branched or C₃-C₈ cyclic haloalkyl, C₁-C₅ linear or branched or C₃-C₈ cyclic alkoxy, C₁-C₅ linear or branched thioalkoxy, C₁-C₅ linear or branched haloalkoxy, C₁-C₅ linear or branched alkoxyalkyl, substituted or unsubstituted C₃-C₈ cycloalkyl, substituted or unsubstituted C₃-C₈ heterocyclic ring, substituted or unsubstituted aryl;

X₃ is O, NH or N-R₅₀;


R₅₀ is H or C₁-C₅ linear or branched, substituted or unsubstituted alkyl;

or its agrochemically acceptable salt, stereoisomer, tautomer, hydrate, *N*-oxide, reverse amide analog, isotopic variant (e.g., deuterated analog), or any combination thereof.

[0056] In some embodiments, substitutions include but not limited to: F, Cl, Br, I, OH, SH, C₁-C₅ linear or branched alkyl (e.g. methyl, ethyl), C₂-C₅ linear or branched alkenyl, C₂-C₅ linear or branched alkynyl (e.g. CCH), C₃-C₈ cycloalkyl, linear, branched or cyclic alkoxy, COOH, COO(R), NH₂, N(R)₂, CF₃, aryl, phenyl, heteroaryl (e.g., imidazole), C₃-C₈ cycloalkyl (e.g., cyclohexyl), CN, NO₂ or any combination thereof.

[0057] In various embodiments, the compound is an herbicidal compound. In various embodiments, the compound is for use in controlling the growth of undesired plants.

[0058] In various embodiments, this invention is directed to a compound represented by the structure of formula **I(g)**:



I(g)


wherein

R₁, **R₁'**, **R₂**, **R₂'** and **R₄₀** are each independently H, F, Cl, Br, I, OH, C₁-C₅ linear or branched, substituted or unsubstituted alkyl (e.g., methyl, ethyl, propyl, iso-propyl, t-Bu, iso-butyl, pentyl, benzyl), or C(O)-R₁₀ (e.g., C(O)-CH₃);

R₃ is F, OH, SH, R₈-OH (e.g., CH₂-OH), NH₂, NHNH₂, NHR, N(R)₂, NHC(O)OBz, -NHC(O)-R₁₀ (e.g., NHC(O)CH₃), C₁-C₅ linear or branched, substituted or unsubstituted alkyl (e.g., methyl, ethyl, propyl, iso-propyl, t-Bu, iso-butyl, pentyl), substituted or unsubstituted C₃-C₈ cycloalkyl, C₁-C₅ linear or branched or C₃-C₈ cyclic haloalkyl, C₁-C₅ linear or branched or C₃-C₈ cyclic alkoxy, substituted or unsubstituted C₃-C₈ heterocyclic ring, or substituted or unsubstituted aryl;

R₄ is OH, NH₂, NHNH₂, NHR, N(R)₂, -NHC(O)-R₁₀, NHC(O)H, NHC(O)CH₃, C₁-C₅ linear or branched, substituted or unsubstituted alkyl, C₁-C₅ linear or branched or C₃-C₈ cyclic haloalkyl, C₁-C₅ linear or branched or C₃-C₈ cyclic alkoxy, substituted or unsubstituted C₃-C₈ cycloalkyl, substituted or unsubstituted C₃-C₈ heterocyclic ring, or substituted or unsubstituted aryl;

or **R₃** and **R₄** are joined to form ring **A**, represented by the following structure:



(e.g., 5-methyloxazolidin-2-one);

R₅ is H, C₁-C₅ linear or branched, substituted or unsubstituted alkyl (e.g., methyl, CH₂SH, ethyl, butyl, CH₂-CCH, iso-propyl, CH₂-C(O)-OCH₃), C₂-C₅ linear or branched, substituted or unsubstituted alkenyl, C₂-C₅ linear or branched, substituted or unsubstituted alkynyl (e.g., CCH, CH₂-CCH), C₁-C₅ linear or branched haloalkyl (e.g., CF₃, CF₂CH₃, CH₂CF₃, CF₂CH₂CH₃, CH₂CH₂CF₃, CF₂CH(CH₃)₂, CF(CH₃)-CH(CH₃)₂), R₈-aryl (e.g., CH₂-Ph), C(=CH₂)-R₁₀ (e.g., C(=CH₂)-C(O)-OCH₃, C(=CH₂)-CN), substituted or unsubstituted alkyl sulfone (e.g., SO₂-CH₂-cyclopentyl), substituted or unsubstituted aryl (e.g., phenyl), substituted or unsubstituted heteroaryl (e.g., pyridine (2, 3, and 4-pyridine));

R₈ is [CH₂]_p

wherein **p** is between 1 and 10;

R₁₀ and **R₁₁** are each independently H, CN, C₁-C₅ linear or branched alkyl (e.g., methyl, ethyl), C(O)R (e.g., C(O)(OCH₃)), or S(O)₂R;

or **R**₁₀ and **R**₁₁ are joined to form a substituted or unsubstituted C₃-C₈ heterocyclic ring (e.g., piperazine, piperidine),

R is H, C₁-C₅ linear or branched alkyl, C₁-C₅ linear or branched alkoxy, phenyl, aryl or heteroaryl, or two gem R substituents are joined together to form a 5 or 6 membered heterocyclic ring;

5 **m** is an integer number between 1 and 5 (e.g., 1 or 2);

n is an integer number between 0 and 5 (e.g., 0, 1, 2 or 3);

X₁ is S, O, N-OH, CH₂, C(R)₂ or N-OMe;

X₂ is S, O, N-OH, CH₂, C(R)₂ or N-OMe;

X₃ is O, NH or N-R₅₀;

10 **R**₅₀ is H or C₁-C₅ linear or branched, substituted or unsubstituted alkyl;


or its agrochemically acceptable salt, stereoisomer, tautomer, hydrate, *N*-oxide, reverse amide analog, isotopic variant (e.g., deuterated analog), or any combination thereof.

[0059] In various embodiments, **R**₃ and **R**₄ of compound of formula **I**(g) cannot both be NH₂. In various
 15 embodiments, if **R**₃ is OH and **R**₄ is NH₂, then **n** + **m** cannot be equal to 3. In various embodiments, **R**₃
 is OH, or NH₂. In various embodiments, **R**₄ is NH₂, or OH. In some embodiments, if **R**₃ is OH then **R**₄
 is NH₂. In some embodiments, if **R**₃ is NH₂ then **R**₄ is OH. In various embodiments, if **R**₃ is OH then **R**₄
 is NH₂ and if **R**₃ is NH₂ then **R**₄ is OH. In some embodiments, the compound is compound **101**, **102**,
 20 **104**, **105**, **106**, **113**, **114**, **115**, **116**, **117**, **118**, **119**, or **120**; each represents a separate embodiment
 according to this invention. In some embodiments, ring **A** has two chiral centers. In some embodiments,
 the compound is not (6*R*,7*S*)-6-amino-7-hydroxyoctanoic acid. In some embodiments, the
 compound is not 5-((4*R*,5*S*)-5-methyl-2-oxooxazolidin-4-yl)pentanoic acid. In some
 embodiments, the compound is a substantially pure single stereoisomer. In some embodiments, **R**₁, **R**₁'
 25 **R**₂, **R**₂' and **R**₄₀ are each independently H or C₁-C₅ linear or branched, substituted or unsubstituted alkyl
 (e.g., methyl, ethyl). In some embodiments, **R**₁ and **R**₁' are both H. In some embodiments, **R**₂ is CH₃ or
 CH₂CH₃. In some embodiments, **R**₂' is H or CH₃. In some embodiments, **R**₄₀ is CH₃ or H. In some
 embodiments, **X**₁ is CH₂. In some embodiments, **X**₂ is CH₂. In some embodiments, **X**₃ is O, NH or N-
 CH₃. In some embodiments, **R**₅ is H, or C₁-C₅ linear or branched, substituted or unsubstituted alkyl, or
 substituted or unsubstituted alkyl sulfone. In some embodiments, **R**₅ is a substituted or unsubstituted
 30 alkyl. In some embodiments, **R**₅ is H. In some embodiments, **R**₅ is H, ethyl, butyl, CH₂-CCH, CH₂-
 C(O)-OCH₃ or SO₂-CH₂-cyclopentyl; each is a separate embodiment according to this invention. In
 some embodiments, **n** is 0, 1, 2 or 3. In some embodiments, **n** is 1. In some embodiments, **m** is 1 or 2.
 In some embodiments, **m** is 1.

[0060] In some embodiments, substitutions include but not limited to: F, Cl, Br, I, OH, SH, C₁-C₅ linear
 35 or branched alkyl (e.g. methyl, ethyl), C₂-C₅ linear or branched alkenyl, C₂-C₅ linear or branched alkynyl
 (e.g. CCH), C₃-C₈ cycloalkyl, linear, branched or cyclic alkoxy, COOH, COO(R), NH₂, N(R)₂, CF₃,
 aryl, phenyl, heteroaryl (e.g., imidazole), C₃-C₈ cycloalkyl (e.g., cyclohexyl), CN, NO₂ or any
 combination thereof.

[0061] In various embodiments, the compound is an herbicidal compound. In various embodiments, the compound is for use in controlling the growth of undesired plants.


[0062] In various embodiments, this invention is directed to a compound represented by the structure of formula **I(ga)**:



I(ga)

wherein

C_A and C_B are both chiral carbon centers, or C_A and C_B together with R_3 and R_4 are joined to form ring **A**, represented by the following structure:



(e.g., 5-methyloxazolidin-2-one);

R_1 , R_1' , R_2 , R_2' and R_{40} are each independently H, F, Cl, Br, I, OH, C_1 - C_5 linear or branched, substituted or unsubstituted alkyl (e.g., methyl, ethyl, propyl, iso-propyl, t-Bu, iso-butyl, pentyl, benzyl), or $C(O)$ - R_{10} (e.g., $C(O)$ - CH_3);

R_3 is F, OH, SH, R_8 -OH (e.g., CH_2 -OH), NH_2 , $NHNH_2$, NHR , $N(R)_2$, $NHC(O)OBz$, $-NHC(O)$ - R_{10} (e.g., $NHC(O)CH_3$), C_1 - C_5 linear or branched, substituted or unsubstituted alkyl (e.g., methyl, ethyl, propyl, iso-propyl, t-Bu, iso-butyl, pentyl), substituted or unsubstituted C_3 - C_8 cycloalkyl, C_1 - C_5 linear or branched or C_3 - C_8 cyclic haloalkyl, C_1 - C_5 linear or branched or C_3 - C_8 cyclic alkoxy, substituted or unsubstituted C_3 - C_8 heterocyclic ring, or substituted or unsubstituted aryl;

R_4 is OH, NH_2 , $NHNH_2$, NHR , $N(R)_2$, $-NHC(O)$ - R_{10} , $NHC(O)H$, $NHC(O)CH_3$, C_1 - C_5 linear or branched, substituted or unsubstituted alkyl, C_1 - C_5 linear or branched or C_3 - C_8 cyclic haloalkyl, C_1 - C_5 linear or branched or C_3 - C_8 cyclic alkoxy, substituted or unsubstituted C_3 - C_8 cycloalkyl, substituted or unsubstituted C_3 - C_8 heterocyclic ring, or substituted or unsubstituted aryl;

R_5 is H, C_1 - C_5 linear or branched, substituted or unsubstituted alkyl (e.g., methyl, CH_2SH , ethyl, butyl, CH_2 -CCH, iso-propyl, CH_2 - $C(O)$ - OCH_3), C_2 - C_5 linear or branched, substituted or unsubstituted alkenyl, C_2 - C_5 linear or branched, substituted or unsubstituted alkynyl (e.g., CCH, CH_2 -CCH), C_1 - C_5 linear or branched haloalkyl (e.g., CF_3 , CF_2CH_3 , CH_2CF_3 , $CF_2CH_2CH_3$, $CH_2CH_2CF_3$, $CF_2CH(CH_3)_2$, $CF(CH_3)-CH(CH_3)_2$), R_8 -aryl (e.g., CH_2 -Ph), $C(=CH_2)$ - R_{10} (e.g., $C(=CH_2)$ - $C(O)$ - OCH_3 , $C(=CH_2)$ -CN), substituted or unsubstituted alkyl sulfone (e.g., SO_2 - CH_2 -cyclopentyl), substituted or

unsubstituted aryl (e.g., phenyl), substituted or unsubstituted heteroaryl (e.g., pyridine (2, 3, and 4-pyridine));

R₈ is [CH₂]_p

wherein **p** is between 1 and 10;

5 **R₁₀** and **R₁₁** are each independently H, CN, C₁-C₅ linear or branched alkyl (e.g., methyl, ethyl), C(O)R (e.g., C(O)(OCH₃)), or S(O)₂R;

or **R₁₀** and **R₁₁** are joined to form a substituted or unsubstituted C₃-C₈ heterocyclic ring (e.g., piperazine, piperidine),

10 **R** is H, C₁-C₅ linear or branched alkyl, C₁-C₅ linear or branched alkoxy, phenyl, aryl or heteroaryl, or two gem R substituents are joined together to form a 5 or 6 membered heterocyclic ring;

m is an integer number between 1 and 5 (e.g., 1 or 2);

n is an integer number between 0 and 5 (e.g., 0, 1, 2 or 3);

X₁ is S, O, N-OH, CH₂, C(R)₂ or N-OMe;

X₂ is S, O, N-OH, CH₂, C(R)₂ or N-OMe;

15 **X₃** is O, NH or N-R₅₀;

R₅₀ is H or C₁-C₅ linear or branched, substituted or unsubstituted alkyl;

or its agrochemically acceptable salt, stereoisomer, tautomer, hydrate, *N*-oxide, reverse amide analog, isotopic variant (e.g., deuterated analog), or any combination thereof.

20 [0063] In various embodiments, **R₃** and **R₄** of compound of formula **I(ga)** cannot both be NH₂. In various embodiments, if **R₃** is OH and **R₄** is NH₂, then **n** + **m** cannot be equal to 3. In various embodiments, **R₃** is OH, or NH₂. In various embodiments, **R₄** is NH₂, or OH. In various embodiments, if **R₃** is OH then **R₄** is NH₂ and if **R₃** is NH₂ then **R₄** is OH. In some embodiments, if **R₃** is OH then **R₄** is NH₂. In some embodiments, if **R₃** is NH₂ then **R₄** is OH. In some embodiments, the compound is
 25 compound **101**, **102**, **104**, **105**, **106**, **113**, **114**, **115**, **116**, **117**, **118**, **119**, or **120**; each represents a separate embodiment according to this invention. In some embodiments, ring **A** has two chiral centers. In some embodiments, the compound is not (6*R*,7*S*)-6-amino-7-hydroxyoctanoic acid. In some embodiments, the compound is not 5-((4*R*,5*S*)-5-methyl-2-oxooxazolidin-4-yl)pentanoic acid. In some embodiments, the compound is a substantially pure single stereoisomer. In some embodiments,
 30 **R₁**, **R₁'**, **R₂**, **R₂'** and **R₄₀** are each independently H or C₁-C₅ linear or branched, substituted or unsubstituted alkyl (e.g., methyl, ethyl). In some embodiments, **R₁** and **R₁'** are both H. In some embodiments, **R₂** is CH₃ or CH₂CH₃. In some embodiments, **R₂'** is H or CH₃. In some embodiments, **R₄₀** is CH₃ or H. In some embodiments, **X₁** is CH₂. In some embodiments, **X₂** is CH₂. In some embodiments, **X₃** is O, NH or N-CH₃. In some embodiments, **R₅** is H, or C₁-C₅ linear or branched,
 35 substituted or unsubstituted alkyl, or substituted or unsubstituted alkyl sulfone. In some embodiments, **R₅** is a substituted or unsubstituted alkyl. In some embodiments, **R₅** is H. In some embodiments, **R₅** is H, ethyl, butyl, CH₂-CCH, CH₂-C(O)-OCH₃ or SO₂-CH₂-cyclopentyl; each is a separate embodiment


according to this invention. In some embodiments, n is 0, 1, 2 or 3. In some embodiments, n is 1. In some embodiments, m is 1 or 2. In some embodiments, m is 1.

[0064] In some embodiments, substitutions include but not limited to: F, Cl, Br, I, OH, SH, C₁-C₅ linear or branched alkyl (e.g. methyl, ethyl), C₂-C₅ linear or branched alkenyl, C₂-C₅ linear or branched alkynyl (e.g. CCH), C₃-C₈ cycloalkyl, linear, branched or cyclic alkoxy, COOH, COO(R), NH₂, N(R)₂, CF₃, aryl, phenyl, heteroaryl (e.g., imidazole), C₃-C₈ cycloalkyl (e.g., cyclohexyl), CN, NO₂ or any combination thereof.

[0065] In various embodiments, the compound is an herbicidal compound. In various embodiments, the compound is for use in controlling the growth of undesired plants.

10

[0066] In various embodiments, this invention is directed to a compound represented by the structure of formula **I(h)**:



15 wherein

R₁, **R₁'**, **R₂**, **R₂'** are each independently H, F, Cl, Br, I, OH, SH, R₈-OH (e.g., CH₂-OH), R₈-SH, -R₈-O-R₁₀, (e.g., -CH₂-O-CH₃), R₈-(C₃-C₈ cycloalkyl) (e.g., cyclohexyl), R₈-(C₃-C₈ heterocyclic ring) (e.g., CH₂-imidazole, CH₂-indazole), CF₃, CD₃, OCD₃, CN, NO₂, -CH₂CN, -R₈CN, NH₂, NHR, N(R)₂, R₈-N(R₁₀)(R₁₁) (e.g., CH₂-NH₂, CH₂-N(CH₃)₂), R₉-R₈-N(R₁₀)(R₁₁) (e.g., C≡C-CH₂-NH₂), B(OH)₂, -OC(O)CF₃, -OCH₂Ph, NHC(O)-R₁₀ (e.g., NHC(O)CH₃), NHC(O)-N(R₁₀)(R₁₁) (e.g., NHC(O)N(CH₃)₂), COOH, -C(O)Ph, C(O)O-R₁₀ (e.g. C(O)O-CH₃, C(O)O-CH(CH₃)₂, C(O)O-CH₂CH₃), R₈-C(O)-R₁₀ (e.g., CH₂C(O)CH₃), C(O)H, C(O)-R₁₀ (e.g., C(O)-CH₃, C(O)-CH₂CH₃, C(O)-CH₂CH₂CH₃), C₁-C₅ linear or branched C(O)-haloalkyl (e.g., C(O)-CF₃), -C(O)NH₂, C(O)NHR, C(O)N(R₁₀)(R₁₁) (e.g., C(O)N(CH₃)₂), SO₂R, SO₂N(R₁₀)(R₁₁) (e.g., SO₂N(CH₃)₂, SO₂NHC(O)CH₃), C₁-C₅ linear or branched, substituted or unsubstituted alkyl (e.g., methyl, 2, 3, or 4-CH₂-C₆H₄-Cl, ethyl, propyl, iso-propyl, t-Bu, iso-butyl, pentyl, benzyl), C₁-C₅ linear or branched, substituted or unsubstituted alkenyl (e.g., CH=C(Ph)₂), C₁-C₅ linear or branched or C₃-C₈ cyclic haloalkyl (e.g., CF₃, CF₂CH₃, CH₂CF₃, CF₂CH₂CH₃, CH₂CH₂CF₃, CF₂CH(CH₃)₂, CF(CH₃)-CH(CH₃)₂), C₁-C₅ linear or branched or C₃-C₈ cyclic alkoxy (e.g. methoxy, ethoxy, propoxy, isopropoxy, O-CH₂-cyclopropyl, O-cyclobutyl, O-cyclopentyl, O-cyclohexyl, 1-butoxy, 2-butoxy, O-tBu), optionally wherein at least one methylene group (CH₂) in the alkoxy is replaced with an oxygen atom (e.g., O-1-oxacyclobutyl, O-2-oxacyclobutyl), C₁-C₅ linear or branched thioalkoxy, C₁-C₅ linear or branched haloalkoxy (e.g., OCF₃, OCHF₂), C₁-C₅ linear or branched alkoxyalkyl, substituted or unsubstituted C₃-C₈ cycloalkyl (e.g., cyclopropyl, cyclopentyl), substituted or unsubstituted C₃-C₈ heterocyclic ring (e.g., 3-methyl-4H-1,2,4-triazole, 5-methyl-1,2,4-

30

oxadiazole, thiophene, oxazole, oxadiazole, imidazole, furane, triazole, tetrazole, pyridine (2, 3, or 4-pyridine), 3-methyl-2-pyridine, pyrimidine, pyrazine, oxacyclobutane (1 or 2-oxacyclobutane), indole, protonated or deprotonated pyridine oxide), substituted or unsubstituted aryl (e.g., phenyl), substituted or unsubstituted benzyl (e.g., benzyl, 4-Cl-benzyl, 4-OH-benzyl);

5 or **R₂** and **R₁** are joined to form a C₃-C₈ substituted or unsubstituted, carbocyclic or heterocyclic ring;

R₃ is H, F, Cl, Br, I, OH, SH, =O, R₈-OH (e.g., CH₂-OH), R₈-SH, -R₈-O-R₁₀, (e.g., CH₂-O-CH₃) CF₃, CD₃, OCD₃, CN, NO₂, -CH₂CN, -R₈CN, NH₂, NHNH₂, NHR, N(R)₂, R₈-N(R₁₀)(R₁₁) (e.g., CH₂-NH₂, CH₂-N(CH₃)₂) R₉-R₈-N(R₁₀)(R₁₁), B(OH)₂, -OC(O)CF₃, -OCH₂Ph, NHC(O)OBz, -NHC(O)-R₁₀ 10 (e.g., NHC(O)CH₃), NHC(O)-N(R₁₀)(R₁₁) (e.g., NHC(O)N(CH₃)₂), COOH, -C(O)Ph, C(O)O-R₁₀ (e.g., C(O)O-CH₃, C(O)O-CH₂CH₃), R₈-C(O)-R₁₀ (e.g., CH₂C(O)CH₃), C(O)H, C(O)-R₁₀ (e.g., C(O)-CH₃, C(O)-CH₂CH₃, C(O)-CH₂CH₂CH₃), C₁-C₅ linear or branched C(O)-haloalkyl (e.g., C(O)-CF₃), -C(O)NH₂, C(O)NHR, C(O)N(R₁₀)(R₁₁) (e.g., C(O)N(CH₃)₂), SO₂R, SO₂N(R₁₀)(R₁₁) (e.g., SO₂N(CH₃)₂), C₁-C₅ linear or branched, substituted or unsubstituted alkyl (e.g., methyl, C(OH)(CH₃)(Ph), ethyl, 15 propyl, iso-propyl, t-Bu, iso-butyl, pentyl), C₁-C₅ linear or branched or C₃-C₈ cyclic haloalkyl (e.g., CF₃, CF₂CH₃, CF₂-cyclobutyl, CH₂CF₃, CF₂CH₂CH₃, CH₂CH₂CF₃, CF₂CH(CH₃)₂, CF(CH₃)-CH(CH₃)₂), C₁-C₅ linear or branched or C₃-C₈ cyclic alkoxy (e.g. methoxy, ethoxy, propoxy, isopropoxy, O-CH₂-cyclopropyl), C₁-C₅ linear or branched thioalkoxy, C₁-C₅ linear or branched haloalkoxy, C₁-C₅ linear or branched alkoxyalkyl, substituted or unsubstituted C₃-C₈ cycloalkyl (e.g., cyclopropyl, cyclopentyl), 20 substituted or unsubstituted C₃-C₈ heterocyclic ring (e.g., 3-methyl-4H-1,2,4-triazole, 5-methyl-1,2,4-oxadiazole, thiophene, oxazole, isoxazole, imidazole, furane, triazole, pyridine (2, 3, or 4-pyridine), pyrimidine, pyrazine, oxacyclobutane (1 or 2-oxacyclobutane), indole), substituted or unsubstituted aryl (e.g., phenyl);

R₄ is NHNH₂;

25 **R₅** is H, C₁-C₅ linear or branched, substituted or unsubstituted alkyl (e.g., methyl, CH₂SH, ethyl, butyl, CH₂-CCH, iso-propyl, CH₂-C(O)-OCH₃), C₂-C₅ linear or branched, substituted or unsubstituted alkenyl, C₂-C₅ linear or branched, substituted or unsubstituted alkynyl (e.g., CCH, CH₂-CCH), C₁-C₅ linear or branched haloalkyl (e.g., CF₃, CF₂CH₃, CH₂CF₃, CF₂CH₂CH₃, CH₂CH₂CF₃, CF₂CH(CH₃)₂, CF(CH₃)-CH(CH₃)₂), R₈-aryl (e.g., CH₂-Ph), C(=CH₂)-R₁₀ (e.g., C(=CH₂)-C(O)-OCH₃, C(=CH₂)-CN), substituted or unsubstituted alkyl sulfone (e.g., SO₂-CH₂-cyclopentyl), substituted or 30 unsubstituted aryl (e.g., phenyl), substituted or unsubstituted heteroaryl (e.g., pyridine (2, 3, and 4-pyridine);

R₈ is [CH₂]_p

wherein **p** is between 1 and 10;

35 **R₉** is [CH]_q, [C]_q

wherein **q** is between 2 and 10;

R₁₀ and **R₁₁** are each independently H, CN, C₁-C₅ linear or branched alkyl (e.g., methyl, ethyl), C(O)R (e.g., C(O)(OCH₃)), or S(O)₂R;

or **R**₁₀ and **R**₁₁ are joined to form a substituted or unsubstituted C₃-C₈ heterocyclic ring (e.g., piperazine, piperidine),

R is H, C₁-C₅ linear or branched alkyl (e.g., methyl, ethyl), C₁-C₅ linear or branched alkoxy (e.g., methoxy), phenyl, aryl or heteroaryl, or two gem R substituents are joined together to form a 5 or 6 membered heterocyclic ring;


m is an integer number between 1 and 5 (e.g., 1 or 2);

n is an integer number between 0 and 5 (e.g., 0, 1, 2 or 3);

X₁ is S, O, N-OH, CH₂, C(R)₂ or N-OMe;

X₂ is S, O, N-OH, CH₂, C(R)₂ or N-OMe;

or **X**₂ together with the carbon next to **X**₁ are joined to form ring B, represented by the following structure (in such case, **X**₁ is **X**₇):



wherein

X₄, **X**₅, **X**₆ and **X**₇ are each independently C or N, wherein if any of **X**₄, **X**₅, **X**₆ and **X**₇ is N, then the respective substitution **R**₉₀, **R**₆₀, **R**₇₀ or **R**₈₀ is absent;

R₆₀, **R**₇₀, **R**₈₀ and **R**₉₀ are each independently selected from: H, F, Cl, Br, I, OH, SH, R₈-OH, R₈-SH, -R₈-O-R₁₀, CF₃, CN, NO₂, NH₂, NHR, N(R)₂, R₈-N(R₁₀)(R₁₁), -OC(O)CF₃, -OCH₂Ph, NHC(O)OBz, -NHC(O)-R₁₀, COOH, -C(O)Ph, C(O)O-R₁₀, C(O)H, C(O)-R₁₀, C₁-C₅ linear or branched C(O)-haloalkyl, -C(O)NH₂, C(O)NHR, C(O)N(R₁₀)(R₁₁), SO₂R, SO₂N(R₁₀)(R₁₁), C₁-C₅ linear or branched, substituted or unsubstituted alkyl, C₁-C₅ linear or branched or C₃-C₈ cyclic haloalkyl, C₁-C₅ linear or branched or C₃-C₈ cyclic alkoxy, C₁-C₅ linear or branched thioalkoxy, C₁-C₅ linear or branched haloalkoxy, C₁-C₅ linear or branched alkoxyalkyl, substituted or unsubstituted C₃-C₈ cycloalkyl, substituted or unsubstituted C₃-C₈ heterocyclic ring, substituted or unsubstituted aryl;

X₃ is O, NH or N-R₅₀;


R₅₀ is H or C₁-C₅ linear or branched, substituted or unsubstituted alkyl;

or its agrochemically acceptable salt, stereoisomer, tautomer, hydrate, N-oxide, reverse amide analog, isotopic variant (e.g., deuterated analog), or any combination thereof.

[0067] In some embodiments, substitutions include but not limited to: F, Cl, Br, I, OH, SH, C₁-C₅ linear or branched alkyl (e.g. methyl, ethyl), C₂-C₅ linear or branched alkenyl, C₂-C₅ linear or branched alkynyl (e.g. CCH), C₃-C₈ cycloalkyl, linear, branched or cyclic alkoxy, COOH, COO(R), NH₂, N(R)₂, CF₃, aryl, phenyl, heteroaryl (e.g., imidazole), C₃-C₈ cycloalkyl (e.g., cyclohexyl), CN, NO₂ or any combination thereof.

[0068] In various embodiments, the compound is an herbicidal compound. In various embodiments, the compound is for use in controlling the growth of undesired plants.

[0069] In various embodiments, this invention is directed to a compound represented by the structure of formula (X):



(X)

wherein

A is a C₃-C₇ cycloalkyl or absent (e.g., cyclohexyl, cyclopropyl, cyclobutyl);

B is a 5-7 membered nitrogen-containing heterocyclic ring or absent (e.g., pyrrolidine, piperidine);

C is a C₅-C₇ a substituted or unsubstituted cycloalkyl, aromatic ring or absent (e.g., phenyl, cyclopentyl, cyclohexyl);

D is a C₅-C₇ cycloalkyl or absent (e.g., cyclopentyl);

E is substituted or unsubstituted 5-7 membered nitrogen-containing heterocyclic ring or absent (e.g., pyrrolidine, piperidine, oxazolidin-2-one);

wherein at least one of ring **A**, **B**, **C**, **D** or **E** is not absent;

R₅ is H, C₁-C₅ linear or branched, substituted or unsubstituted alkyl (e.g., methyl, CH₂SH, ethyl, butyl, CH₂-CCH, iso-propyl, CH₂-C(O)-OCH₃), C₂-C₅ linear or branched, substituted or unsubstituted alkenyl, C₂-C₅ linear or branched, substituted or unsubstituted alkynyl (e.g., CCH, CH₂-CCH), C₁-C₅ linear or branched haloalkyl (e.g., CF₃, CF₂CH₃, CH₂CF₃, CF₂CH₂CH₃, CH₂CH₂CF₃, CF₂CH(CH₃)₂, CF(CH₃)-CH(CH₃)₂), R₈-aryl (e.g., CH₂-Ph), C(=CH₂)-R₁₀ (e.g., C(=CH₂)-C(O)-OCH₃, C(=CH₂)-CN), substituted or unsubstituted aryl (e.g., phenyl), substituted or unsubstituted heteroaryl (e.g., pyridine (2, 3, and 4-pyridine));

R₈ is [CH₂]_p

wherein **p** is between 1 and 10;

R₉ is [CH]_q, [C]_q

wherein **q** is between 2 and 10;

R₁₀ and **R₁₁** are each independently H, CN, C₁-C₅ linear or branched alkyl (e.g., methyl, ethyl), C(O)R (e.g., C(O)(OCH₃)), or S(O)₂R; or **R₁₀** and **R₁₁** are joined to form a substituted or unsubstituted C₃-C₈ heterocyclic ring (e.g., piperazine, piperidine),

R is H, C₁-C₅ linear or branched alkyl (e.g., methyl, ethyl), C₁-C₅ linear or branched alkoxy (e.g., methoxy), phenyl, aryl or heteroaryl, or two gem R substituents are joined together to form a 5 or 6 membered heterocyclic ring;

n is an integer number between 0 and 2;

X₁ is S, O, CH₂, CH(R) or C(R)₂;

X₂ is S, O, CH₂, CH(R) or C(R)₂;


or its agrochemically acceptable salt, stereoisomer, tautomer, hydrate, *N*-oxide, reverse amide analog, isotopic variant (e.g., deuterated analog), or any combination thereof.

[0070] In some embodiments, at least one of rings **A** – **E** of compound of formula (**X**) is not absent. In some embodiments, only one of rings **A** – **E** is not absent. In some embodiments, **A** is absent. In some 15 embodiments, **A** is cyclohexyl. In some embodiments, **A** is cyclopropyl. In some embodiments, **A** is cyclobutyl. In some embodiments, **B** is absent. In some embodiments, **B** is pyrrolidine. In some embodiments, **B** is piperidine. In some embodiments, **C** is absent. In some embodiments, **C** is a C₅-C₇ substituted or unsubstituted cycloalkyl. In some embodiments, **C** is cyclopentyl. In some embodiments, **C** is cyclohexyl. In some embodiments, **C** is an aromatic ring. In some embodiments, **C** is a phenyl. In 20 some embodiments, **D** is absent. In some embodiments, **D** is cyclopentyl. In some embodiments, **E** is absent. In some embodiments, **E** is pyrrolidine. In some embodiments, **E** is piperidine. In some embodiments, **E** is oxazolidin-2-one. In some embodiments, **E** is substituted oxazolidin-2-one. In some embodiments, **X₁** is S. In some embodiments, **X₁** is O. In some embodiments, **X₁** is CH₂. In some 25 embodiments, **X₂** is S. In some embodiments, **X₂** is O. In some embodiments, **X₂** is CH₂. In some embodiments, **n** is 0. In some embodiments, **n** is 1. In some embodiments, **n** is 2. In some embodiments, **R₅** is H. In some embodiments, **R₅** is C₁-C₅ linear or branched, substituted or unsubstituted alkyl. In some embodiments, **R₅** is ethyl. In some embodiments, **R₅** is butyl. In some embodiments, **R₅** is CH₂-C≡CH.

[0071] In some embodiments, substitutions include but not limited to: F, Cl, Br, I, OH, SH, C₁-C₅ linear or branched alkyl (e.g. methyl, ethyl), C₂-C₅ linear or branched alkenyl, C₂-C₅ linear or branched alkynyl (e.g. CCH), C₃-C₈ cycloalkyl, linear, branched or cyclic alkoxy, COOH, COO(R), NH₂, N(R)₂, CF₃, aryl, phenyl, heteroaryl (e.g., imidazole), C₃-C₈ cycloalkyl (e.g., cyclohexyl), CN, NO₂ or any 30 combination thereof.

[0072] In various embodiments, the compound is an herbicidal compound. In various embodiments, 35 the compound is for use in controlling the growth of undesired plants. In various embodiments, the compound is compound **105**, **106**, **123**, **124**, **125**, **126**, **129**, **130**, **131**, **132**, **133**, **134**, **138**, **139**, **143**, **145**, **146**, **147**, **148**, **150**, **152**, **153**, **154**, **155**, **165**, **166**, **167**, **168** or **171**; each represents a separate embodiment according to this invention.

[0073] In various embodiments, this invention is directed to a compound represented by the structure of formula **X(a)**:



5 wherein

R₅ is H, C₁-C₅ linear or branched, substituted or unsubstituted alkyl (e.g., methyl, CH₂SH, ethyl, butyl, CH₂-CCH, iso-propyl, CH₂-C(O)-OCH₃), C₂-C₅ linear or branched, substituted or unsubstituted alkenyl, C₂-C₅ linear or branched, substituted or unsubstituted alkynyl (e.g., CCH, CH₂-CCH), C₁-C₅ linear or branched haloalkyl (e.g., CF₃, CF₂CH₃, CH₂CF₃, CF₂CH₂CH₃, CH₂CH₂CF₃, CF₂CH(CH₃)₂, CF(CH₃)-CH(CH₃)₂), R₈-aryl (e.g., CH₂-Ph), C(=CH₂)-R₁₀ (e.g., C(=CH₂)-C(O)-OCH₃, C(=CH₂)-CN), substituted or unsubstituted aryl (e.g., phenyl), substituted or unsubstituted heteroaryl (e.g., pyridine (2, 3, and 4-pyridine));

R₈ is [CH₂]_p

wherein **p** is between 1 and 10;

15 **R₉** is [CH]_q, [C]_q

wherein **q** is between 2 and 10;

R₁₀ and **R₁₁** are each independently H, CN, C₁-C₅ linear or branched alkyl (e.g., methyl, ethyl), C(O)R (e.g., C(O)(OCH₃)), or S(O)₂R;

or **R₁₀** and **R₁₁** are joined to form a substituted or unsubstituted C₃-C₈ heterocyclic ring (e.g., piperazine, piperidine),

R is H, C₁-C₅ linear or branched alkyl (e.g., methyl, ethyl), C₁-C₅ linear or branched alkoxy (e.g., methoxy), phenyl, aryl or heteroaryl, or two gem R substituents are joined together to form a 5 or 6 membered heterocyclic ring;

X₁ is S, O, CH₂, CH(R) or C(R)₂;

25 **n** and **o** are each independently an integer number between 0 and 2;

m is an integer number between 1 and 3;

or its agrochemically acceptable salt, stereoisomer, tautomer, hydrate, *N*-oxide, reverse amide analog, isotopic variant (e.g., deuterated analog), or any combination thereof.








30 [0074] In some embodiments, **m** compound of formula **X(a)** is 1. In some embodiments, **m** is 2. In some embodiments, **m** is 1 or 2. In some embodiments, **o** is 0. In some embodiments, **o** is 1. In some embodiments, **o** is 0 or 1. In some embodiments, **n** is 0. In some embodiments, **n** is 1. In some embodiments, **n** is 2. In some embodiments, **n** is 0 or 1. In some embodiments, **n** is 1 or 2. In some embodiments, **X₁** is CH₂. In some embodiments, **X₁** is S. In some embodiments, **X₁** is O. In some

35 embodiments, **R₅** is H. In some embodiments, **R₅** is C₁-C₅ linear or branched, substituted or unsubstituted alkyl. In some embodiments, **R₅** is ethyl. In some embodiments, **R₅** is butyl. In some

embodiments, R_5 is substituted alkyl. In some embodiments, R_5 is $CH_2-C\equiv CH$. In some embodiments, R_5 is not H. In some embodiments, X_1 is CH_2 , m is 2, n is 1, and R_5 is substituted or unsubstituted alkyl. In some embodiments, X_1 is CH_2 , m is 2, n is 1, o is 1, and R_5 is substituted or unsubstituted alkyl. In some embodiments, X_1 is O or S, m is 1 or 2, n is 1 or 2, o is 0 or 1, and R_5 is H. In some embodiments, if R_5 is H then X_1 is not CH_2 . In some embodiments, if X_1 is O and R_5 is H then n is not 1. In some embodiments, if X_1 is O then n is not 1.


[0075] In some embodiments, substitutions include but not limited to: F, Cl, Br, I, OH, SH, C_1 - C_5 linear or branched alkyl (e.g. methyl, ethyl), C_2 - C_5 linear or branched alkenyl, C_2 - C_5 linear or branched alkynyl (e.g. CCH), C_3 - C_8 cycloalkyl, linear, branched or cyclic alkoxy, COOH, COO(R), NH_2 , $N(R)_2$, CF_3 , aryl, phenyl, heteroaryl (e.g., imidazole), C_3 - C_8 cycloalkyl (e.g., cyclohexyl), CN, NO_2 or any combination thereof.

[0076] In various embodiments, the compound of formula **X(a)**, is represented by any one of the following structures:

Compound	Structure
143	
144	
145	
124	
125	
126	
139	

[0077] In various embodiments, the compound is an herbicidal compound. In various embodiments, the compound is for use in controlling the growth of undesired plants.

[0078] In various embodiments, this invention is directed to a compound represented by the structure of formula **X(b)**:



X(b)

wherein

R₅ is H, C₁-C₅ linear or branched, substituted or unsubstituted alkyl (e.g., methyl, CH₂SH, ethyl, butyl, CH₂-CCH, iso-propyl, CH₂-C(O)-OCH₃), C₂-C₅ linear or branched, substituted or unsubstituted alkenyl, C₂-C₅ linear or branched, substituted or unsubstituted alkynyl (e.g., CCH, CH₂-CCH), C₁-C₅ linear or branched haloalkyl (e.g., CF₃, CF₂CH₃, CH₂CF₃, CF₂CH₂CH₃, CH₂CH₂CF₃, CF₂CH(CH₃)₂, CF(CH₃)-CH(CH₃)₂), R₈-aryl (e.g., CH₂-Ph), C(=CH₂)-R₁₀ (e.g., C(=CH₂)-C(O)-OCH₃, C(=CH₂)-CN), substituted or unsubstituted aryl (e.g., phenyl), substituted or unsubstituted heteroaryl (e.g., pyridine (2, 3, and 4-pyridine));

R₈ is [CH₂]_p

10 wherein **p** is between 1 and 10;

R₉ is [CH]_q, [C]_q

 wherein **q** is between 2 and 10;

R₁₀ and **R₁₁** are each independently H, CN, C₁-C₅ linear or branched alkyl (e.g., methyl, ethyl), C(O)R (e.g., C(O)(OCH₃)), or S(O)₂R;

15 or **R₁₀** and **R₁₁** are joined to form a substituted or unsubstituted C₃-C₈ heterocyclic ring (e.g., piperazine, piperidine),

R is H, C₁-C₅ linear or branched alkyl (e.g., methyl, ethyl), C₁-C₅ linear or branched alkoxy (e.g., methoxy), phenyl, aryl or heteroaryl, or two gem R substituents are joined together to form a 5 or 6 membered heterocyclic ring;

20 **X₁** is S, O, CH₂, CH(R) or C(R)₂;

X₂ is S, O, CH₂, CH(R) or C(R)₂;

n is an integer number between 0 and 2;

m is an integer number between 1 and 3;

25 or its agrochemically acceptable salt, stereoisomer, tautomer, hydrate, *N*-oxide, reverse amide analog, isotopic variant (e.g., deuterated analog), or any combination thereof.

[0079] In some embodiments, m of compound of formula **X(b)** is 1. In some embodiments, m is 2. In some embodiments, m is 1 or 2. In some embodiments, n is 0. In some embodiments, n is 1. In some embodiments, n is 0 or 1. In some embodiments, **X₁** is CH₂. In some embodiments, **X₂** is CH₂. In some 30 embodiments, **X₁** and **X₂** are CH₂. In some embodiments, **R₅** is H.


[0080] In some embodiments, substitutions include but not limited to: F, Cl, Br, I, OH, SH, C₁-C₅ linear or branched alkyl (e.g. methyl, ethyl), C₂-C₅ linear or branched alkenyl, C₂-C₅ linear or branched alkynyl (e.g. CCH), C₃-C₈ cycloalkyl, linear, branched or cyclic alkoxy, COOH, COO(R), NH₂, N(R)₂, CF₃, aryl, phenyl, heteroaryl (e.g., imidazole), C₃-C₈ cycloalkyl (e.g., cyclohexyl), CN, NO₂ or any 35 combination thereof.

[0081] In various embodiments, the compound of formula **X(b)** represented by any one of the following structures:

Compound	Structure
148	
155	
166	

[0082] In various embodiments, the compound is an herbicidal compound. In various embodiments, the compound is for use in controlling the growth of undesired plants.

[0083] In various embodiments, this invention is directed to a compound represented by the structure of formula **X(c)**:



X(c)

wherein

R₅ is H, C₁-C₅ linear or branched, substituted or unsubstituted alkyl (e.g., methyl, CH₂SH, ethyl, butyl, CH₂-CCH, iso-propyl, CH₂-C(O)-OCH₃), C₂-C₅ linear or branched, substituted or unsubstituted alkenyl, C₂-C₅ linear or branched, substituted or unsubstituted alkynyl (e.g., CCH, CH₂-CCH), C₁-C₅ linear or branched haloalkyl (e.g., CF₃, CF₂CH₃, CH₂CF₃, CF₂CH₂CH₃, CH₂CH₂CF₃, CF₂CH(CH₃)₂, CF(CH₃)-CH(CH₃)₂), R₈-aryl (e.g., CH₂-Ph), C(=CH₂)-R₁₀ (e.g., C(=CH₂)-C(O)-OCH₃, C(=CH₂)-CN), substituted or unsubstituted aryl (e.g., phenyl), substituted or unsubstituted heteroaryl (e.g., pyridine (2, 3, and 4-pyridine);

R₈ is [CH₂]_p

wherein **p** is between 1 and 10;

R₉ is [CH]_q, [C]_q

wherein **q** is between 2 and 10;

R₁₀ and **R₁₁** are each independently H, CN, C₁-C₅ linear or branched alkyl (e.g., methyl, ethyl), C(O)R (e.g., C(O)(OCH₃)), or S(O)₂R;

or **R₁₀** and **R₁₁** are joined to form a substituted or unsubstituted C₃-C₈ heterocyclic ring (e.g., piperazine, piperidine),

R is H, C₁-C₅ linear or branched alkyl (e.g., methyl, ethyl), C₁-C₅ linear or branched alkoxy (e.g., methoxy), phenyl, aryl or heteroaryl, or two gem R substituents are joined together to form a 5 or 6 membered heterocyclic ring;

X₁ is S, O, CH₂, CH(R) or C(R)₂;

X_2 is S, O, CH_2 , $CH(R)$ or $C(R)_2$;

n is an integer number between 0 and 2;








m is an integer number between 1 and 3;


5 or its agrochemically acceptable salt, stereoisomer, tautomer, hydrate, *N*-oxide, reverse amide analog, isotopic variant (e.g., deuterated analog), or any combination thereof.

[0084] In some embodiments, m of compound of formula $X(c)$ is 1. In some embodiments, m is 2. In some embodiments, m is 1 or 2. In some embodiments, n is 0. In some embodiments, n is 1. In some embodiments, n is 0 or 1. In some embodiments, X_1 is CH_2 . In some embodiments, X_2 is CH_2 . In some
10 embodiments, X_2 is S. In some embodiments, X_2 is O. In some embodiments, X_1 and X_2 are CH_2 . In some embodiments, R_5 is H. In some embodiments, R_5 is substituted or unsubstituted alkyl. In some embodiments, R_5 is methyl. In some embodiments, X_1 and X_2 are CH_2 , n is 1, and m is 1 or 2. In some embodiments, n is 1, R_5 is H and m is 1 or 2.

[0085] In some embodiments, substitutions include but not limited to: F, Cl, Br, I, OH, SH, C_1 - C_5 linear or branched alkyl (e.g. methyl, ethyl), C_2 - C_5 linear or branched alkenyl, C_2 - C_5 linear or branched alkynyl (e.g. CCH), C_3 - C_8 cycloalkyl, linear, branched or cyclic alkoxy, COOH, COO(R), NH_2 , $N(R)_2$, CF_3 , aryl, phenyl, heteroaryl (e.g., imidazole), C_3 - C_8 cycloalkyl (e.g., cyclohexyl), CN, NO_2 or any
15 combination thereof.


20 [0086] In various embodiments, the compound of formula $X(c)$ is represented by any one of the following structures:

Compound	Structure
179	
146	
147	
152	
153	
131	
132	



[0087] In various embodiments, the compound is an herbicidal compound. In various embodiments, the compound is for use in controlling the growth of undesired plants.

[0088] In various embodiments, this invention is directed to a compound represented by the structure of formula **X(d)**:



wherein

R₅ is H, C₁-C₅ linear or branched, substituted or unsubstituted alkyl (e.g., methyl, CH₂SH, ethyl, butyl, CH₂-CCH, iso-propyl, CH₂-C(O)-OCH₃), C₂-C₅ linear or branched, substituted or unsubstituted alkenyl, C₂-C₅ linear or branched, substituted or unsubstituted alkynyl (e.g., CCH, CH₂-CCH), C₁-C₅ linear or branched haloalkyl (e.g., CF₃, CF₂CH₃, CH₂CF₃, CF₂CH₂CH₃, CH₂CH₂CF₃, CF₂CH(CH₃)₂, CF(CH₃)-CH(CH₃)₂), R₈-aryl (e.g., CH₂-Ph), C(=CH₂)-R₁₀ (e.g., C(=CH₂)-C(O)-OCH₃, C(=CH₂)-CN), substituted or unsubstituted aryl (e.g., phenyl), substituted or unsubstituted heteroaryl (e.g., pyridine (2, 3, and 4-pyridine);

R₈ is [CH₂]_p

wherein **p** is between 1 and 10;

R₉ is [CH]_q, [C]_q

wherein **q** is between 2 and 10;

R₁₀ and **R₁₁** are each independently H, CN, C₁-C₅ linear or branched alkyl (e.g., methyl, ethyl), C(O)R (e.g., C(O)(OCH₃)), or S(O)₂R;

or **R₁₀** and **R₁₁** are joined to form a substituted or unsubstituted C₃-C₈ heterocyclic ring (e.g., piperazine, piperidine),

R is H, C₁-C₅ linear or branched alkyl (e.g., methyl, ethyl), C₁-C₅ linear or branched alkoxy (e.g., methoxy), phenyl, aryl or heteroaryl, or two gem R substituents are joined together to form a 5 or 6 membered heterocyclic ring;

X₂ is S, O, CH₂, CH(R) or C(R)₂;

n is an integer number between 0 and 2;

m is an integer number between 1 and 3;

or its agrochemically acceptable salt, stereoisomer, tautomer, hydrate, *N*-oxide, reverse amide analog, isotopic variant (e.g., deuterated analog), or any combination thereof.

[0089] In some embodiments, **m** of compound of formula **X(d)** is 1. In some embodiments, **m** is 2. In some embodiments, **m** is 1 or 2. In some embodiments, **n** is 1. In some embodiments, **n** is 2. In some

embodiments, n is 1 or 2. In some embodiments, X_2 is CH_2 . In some embodiments, R_5 is H. In some embodiments, R_5 is H, n is 2 and m is 1. In some embodiments, R_5 is H, X_2 is CH_2 , n is 2 and m is 1.

[0090] In some embodiments, substitutions include but not limited to: F, Cl, Br, I, OH, SH, C_1 - C_5 linear or branched alkyl (e.g. methyl, ethyl), C_2 - C_5 linear or branched alkenyl, C_2 - C_5 linear or branched alkynyl (e.g. CCH), C_3 - C_8 cycloalkyl, linear, branched or cyclic alkoxy, COOH, COO(R), NH_2 , $N(R)_2$, CF_3 , aryl, phenyl, heteroaryl (e.g., imidazole), C_3 - C_8 cycloalkyl (e.g., cyclohexyl), CN, NO_2 or any combination thereof.

[0091] In various embodiments, the compound of formula **X(d)** is represented by any one of the following structures:



















Compound	Structure
123	
181	
154	









[0092] In various embodiments, the compound is an herbicidal compound. In various embodiments, the compound is for use in controlling the growth of undesired plants.

[0093] In various embodiments, this invention is directed to a compound represented by any one of the following structures:

Table 1:

Compound	Structure
101	
102	
104	
105	
106	






113	
114	
115	
123	
116	
117	
118	
119	
124	
125	
126	
120	
127	
128	
129	
130	
131	
132	





















133	
134	
137	
138	
139	
140	
141	
142	





















[0094] In various embodiments, the compound is an herbicidal compound. In various embodiments, the compound is for use in controlling the growth of undesired plants.























- 5 [0095] In various embodiments, this invention is directed to an herbicidal compound and/or to a use of a compound represented by any one of the following structures, or an agrochemical composition thereof, in controlling the growth of undesired plants:

Table 2:

Compound	Structure
101	
102	
104	
105	
106	

107	
108	
109	
110	
111	
112	
113	
114	
115	
116	
117	
118	
119	
120	
123	
124	
125	
126	
127	
128	

129	
130	
131	
132	
133	
134	
137	
138	
139	
140	
141	
142	
143	
144	
145	
146	
147	
148	
149	
150	

151	
152	
153	
154	
155	
156	
157	
158	
159	
160	
161	
162	
163	
164	
165	
166	
167	
168	
169	
170	
171	
172	

173	
174	
175	
176	
177	
178	

[0096] In various embodiments, ring **A** of compound of formula **I**, **I(a)**, **I(b)**, **I(g)** and **I(ga)** has one chiral center (i.e., R_2 and R_2' are identical). In various embodiments, C_B of ring **A** of formula **I(a)** or **I(b)** is achiral (i.e., R_2 and R_2' are identical). In various embodiments, C_B of ring **A** is chiral. In various

5 embodiments, ring **A** has two chiral centers (i.e., R_2 and R_2' are different)). In various embodiments, the compound is a single stereoisomer. In various embodiments, the compound is a single enantiomer.

[0097] In various embodiments, the compound comprises a substantially pure stereoisomer. By substantially pure, it is intended that a stereoisomer is at least about 90% pure, more preferably at least about 95% pure, even more preferably at least about 98% pure, most preferably at least about 99% pure.

10 In various embodiments, the compound comprises a single stereoisomer in a purity of >80%; >85%; >90%; >91%; >92%; >93%; >94%; >95%; >96%; >97%; >98%; >99%; >99.5% enantiomeric excess (*ee*); each represents a separate embodiment according to this invention. In various embodiments, the compound comprises a single stereoisomer in a purity >80%; >85%; >90%; >91%; >92%; >93%; >94%; >95%; >96%; >97%; >98%; >99%; >99.5% enantiomeric ratio (*er*); each represents a separate

15 embodiment according to this invention. In various embodiments, the compound comprises a single stereoisomer in a purity higher than 80%; 85%; 90%; 91%; 92%; 93%; 94%; 95%; 96%; 97%; 98%; 99%; 99.5%; each represents a separate embodiment according to this invention.

[0098] In various embodiments, the compound is a substantially pure single enantiomer. In various embodiments, the compound comprises a mixture of stereoisomers. In various embodiments, the

20 compound comprises a mixture of enantiomers. In various embodiments, the compound is a racemate.

[0099] In various embodiments, the compound has two chiral centers. In various embodiments, the compound comprises a mixture of stereoisomers. In various embodiments, the compound comprises a mixture of 2, 3, or 4 stereoisomers; each represents a separate embodiment according to this invention.

In various embodiments, the compound is a single stereoisomer. In various embodiments, the compound

25 is a substantially pure single stereoisomer. In various embodiments, the substantially pure stereoisomer is compound **104**, **105**, **106**, **114**, **115**, **116**, **117**, **118**, or **119** as described herein below; each represents

a separate embodiment according to this invention. In various embodiments, the substantially pure stereoisomer has at least 80%, 85%, 90%, 95%, 97%, 98%, 99% purity; each represents a separate embodiment according to this invention. In various embodiments, the compound is the substantially pure **RR** stereoisomer. In various embodiments, the compound is the substantially pure **SS** stereoisomer.

5 In various embodiments, the compound is the substantially pure **RS** stereoisomer. In various embodiments, the compound is 5-((4*R*,5*S*)-5-methyl-2-oxooxazolidin-4-yl)pentanoic acid. In various embodiments, the compound is Compound **106**. In various embodiments, the compound is the substantially pure **SR** stereoisomer. In various embodiments, the compound is (6*S*,7*R*)-6-amino-7-hydroxyoctanoic acid. In various embodiments, the compound is Compound **104**. In various
10 embodiments, the compound is 5-((4*S*,5*R*)-5-methyl-2-oxooxazolidin-4-yl)pentanoic acid. In various embodiments, the compound is Compound **105**. In various embodiments, the compound is not (6*R*,7*S*)-6-amino-7-hydroxyoctanoic acid. In various embodiments, the compound is not 5-((4*R*,5*S*)-5-methyl-2-oxooxazolidin-4-yl)pentanoic acid.

[00100] In some embodiments, **A** of formula **I**, **I(a)**, **I(b)**, **I(g)** or **I(ga)** is absent. In some embodiments,
15 **A** of formula **I**, is a substituted or unsubstituted single or fused aromatic or heteroaromatic ring system, or a substituted or unsubstituted single or fused C₃-C₁₀ cycloalkyl, or a single or fused C₃-C₁₀ heterocyclic ring. In some embodiments, **A** of formula **I**, is a substituted or unsubstituted single or fused aromatic or heteroaromatic ring system, or a substituted or unsubstituted single or fused C₄-C₁₀ cycloalkyl, or a single or fused C₄-C₁₀ heterocyclic ring. In some embodiments, **A** is a single aromatic
20 ring system. In some embodiments, **A** is a substituted aryl. In some embodiments, the aryl is substituted with NH₂. In some embodiments, **A** is a single heteroaromatic ring. In some embodiments, **A** is a single C₃-C₁₀ cycloalkyl. In some embodiments, **A** is a substituted single C₃-C₁₀ cycloalkyl. In some embodiments, **A** is a single C₄-C₁₀ cycloalkyl. In some embodiments, **A** is a substituted single C₄-C₁₀ cycloalkyl. In some embodiments, **A** is cyclopropyl. In some embodiments, **A** is cyclobutyl. In some
25 embodiments, **A** is cyclopentyl. In some embodiments, **A** is cyclohexyl. In some embodiments, **A** is further substituted with at least one substitution selected from: F, Cl, Br, I, OH, SH, C₁-C₅ linear or branched alkyl (e.g. methyl, ethyl), C₂-C₅ linear or branched alkenyl, C₂-C₅ linear or branched alkynyl (e.g. CCH), C₃-C₈ cycloalkyl, linear, branched or cyclic alkoxy, COOH, COO(R), NH₂, N(R)₂, CF₃, aryl, phenyl, heteroaryl (e.g., imidazole), C₃-C₈ cycloalkyl (e.g., cyclohexyl), CN and NO₂; each
30 represents a separate embodiment according to this invention. In some embodiments, **A** is further substituted with NH₂. In some embodiments, **A** is a cycloalkyl, substituted with NH₂. In some embodiments, the cycloalkyl is cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl; each represents a separate embodiment according to this invention. In some embodiments, **A** is a single C₃-C₁₀ heterocyclic ring. In some embodiments, **A** is a single C₄-C₁₀ heterocyclic ring. In some embodiments,
35 **A** is a substituted oxazolidin-2-one ring. In some embodiments, **A** is an alkyl substituted oxazolidin-2-one ring. In some embodiments, **A** is 5-methyloxazolidin-2-one. In some embodiments, **A** is 5-methyloxazolidin-2-one. In some embodiments, **A** is a nitrogen-containing heterocyclic ring. In some embodiments, **A** is 1, 2, or 3-pyrrolidine; each represents a separate embodiment according to this

invention. In some embodiments, **A** is tetrahydropyridine. In some embodiments, **A** is 5,6-dihydro-4H-1,3-thiazine. In some embodiments, **A** is 4,5-dihydro-1H-imidazole. In some embodiments, **A** is 2, 3 or 4-pyridine; each represents a separate embodiment according to this invention. In some embodiments, the pyridine is further substituted with at least NH₂. In some embodiments, **A** is tetrahydropyrimidine.

5 In some embodiments, **A** is 1, 2 or 3-piperidine; each represents a separate embodiment according to this invention. In some embodiments, **A** is imidazole. In some embodiments, the nitrogen containing heterocyclic ring is further substituted with at least NH₂. In some embodiments, **A** is a fused aromatic ring system. In some embodiments, **A** is a fused heteroaromatic ring system. In some embodiments, **A** is a fused C₃-C₁₀ cycloalkyl. In some embodiments, **A** is a fused C₃-C₁₀ heterocyclic ring system. In


10 some embodiments, **A** is a phenyl. In other embodiments, **A** is pyridinyl. In other embodiments, **A** is 2-pyridinyl. In other embodiments, **A** is 3-pyridinyl. In other embodiments, **A** is 4-pyridinyl. In other embodiments, **A** is naphthyl. In other embodiments, **A** is benzothiazolyl. In other embodiments, **A** is benzimidazolyl. In other embodiments, **A** is quinolinyl. In other embodiments, **A** is isoquinolinyl. In other embodiments, **A** is indolyl. In other embodiments, **A** is tetrahydronaphthyl. In other embodiments,

15 **A** is indenyl. In other embodiments, **A** is benzofuran-2(3H)-one. In other embodiments, **A** is benzo[d][1,3]dioxole. In other embodiments, **A** is naphthalene. In other embodiments, **A** is tetrahydrothiophene1,1-dioxide. In other embodiments, **A** is thiazole. In other embodiments, **A** is benzimidazole. In other embodiments, **A** is piperidine. In other embodiments, **A** is 1-methylpiperidine. In other embodiments, **A** is imidazole. In other embodiments, **A** is 1-methylimidazole. In other

20 embodiments, **A** is thiophene. In other embodiments, **A** is isoquinoline. In other embodiments, **A** is indole. In other embodiments, **A** is 1,3-dihydroisobenzofuran. In other embodiments, **A** is benzofuran. In other embodiments, **A** is single or fused C₃-C₁₀ cycloalkyl ring. In other embodiments, **A** is cyclohexyl. In some embodiments, **A** may be further substituted, with at least one substituent selected from: F, Cl, Br, I, OH, SH, C₁-C₅ linear or branched alkyl (e.g. methyl, ethyl), C₂-C₅ linear or branched

25 alkenyl, C₂-C₅ linear or branched alkynyl (e.g. CCH), C₃-C₈ cycloalkyl, linear, branched or cyclic alkoxy, COOH, COO(R), NH₂, N(R)₂, CF₃, aryl, phenyl, heteroaryl (e.g., imidazole), C₃-C₈ cycloalkyl (e.g., cyclohexyl), CN and NO₂; each represents a separate embodiment according to this invention. In some embodiments, **A** may be further substituted, with NH₂.

[00101] In some embodiments, ring **A** of formula **I**, **I(a)**, **I(b)**, **I(g)**, and/or **I(ga)**, is represented by the



30 following structure: $\text{wherein } R_2 \text{ and } R_2' \text{ are as defined below.}$

[00102] It is understood that the wavy line in ring **A** above, represents the connection point of **A** to the rest of the molecule (i.e., to $-(C(R_1)(R_1'))_n$) of formula **I**, **I(a)**, **I(b)**, **I(g)** or **I(ga)**.


[00103] In some embodiments, ring **B** of formula **I**, **I(a)**, **I(b)**, **I(f)** and/or **I(h)** is absent. In other embodiments, ring **B** is pyridine.

[00104] In some embodiments, ring **B** of formula **I**, is a single or fused aromatic or heteroaromatic ring system, or a single or fused C₃-C₁₀ cycloalkyl, or a single or fused C₃-C₁₀ heterocyclic ring. In some embodiments, ring **B** is a single aromatic ring system (i.e., arene). In some embodiments, ring **B** is a single heteroaromatic ring (e.g., pyridine). In some embodiments, ring **B** is a single C₃-C₁₀ cycloalkyl.

5 In some embodiments, ring **B** is a single C₃-C₁₀ heterocyclic ring. In some embodiments, ring **B** is a fused aromatic ring system. In some embodiments, ring **B** is a fused heteroaromatic ring system. In some embodiments, ring **B** is a fused C₃-C₁₀ cycloalkyl. In some embodiments, ring **B** is a fused C₃-C₁₀ heterocyclic ring system. In some embodiments, ring **B** is an arene. In other embodiments, ring **B** is pyridine ring. In other embodiments, ring **B** is pyrazine. In other embodiments, ring **B** is pyridazine. In

10 other embodiments, ring **B** is pyrimidine. In other embodiments, ring **B** is a triazine. In other embodiments, ring **B** is a tetrazine. In some embodiments, ring **B** may be further substituted, with at least one substituent selected from: F, Cl, Br, I, OH, SH, R₈-OH, R₈-SH, -R₈-O-R₁₀, CF₃, CN, NO₂, NH₂, NHR, N(R)₂, R₈-N(R₁₀)(R₁₁), -OC(O)CF₃, -OCH₂Ph, NHC(O)OBz, -NHC(O)-R₁₀, COOH, -C(O)Ph, C(O)O-R₁₀, C(O)H, C(O)-R₁₀, C₁-C₅ linear or branched C(O)-haloalkyl, -C(O)NH₂, C(O)NHR, C(O)N(R₁₀)(R₁₁), SO₂R, SO₂N(R₁₀)(R₁₁), C₁-C₅ linear or branched or C₃-C₈ cyclic haloalkyl, C₁-C₅ linear or branched or C₃-C₈ cyclic alkoxy, C₁-C₅ linear or branched thioalkoxy, C₁-C₅ linear or branched haloalkoxy, C₁-C₅ linear or branched alkoxyalkyl, substituted or unsubstituted C₃-C₈ cycloalkyl, substituted or unsubstituted C₃-C₈ heterocyclic ring, substituted or unsubstituted aryl, C(O)-CH₃, C₁-C₅ linear or branched, substituted or unsubstituted alkyl, methyl, ethyl, propyl, iso-propyl, t-Bu, iso-butyl, pentyl, benzyl, or any combination thereof; each represents a separate embodiment according to this invention.

[00105] In some embodiments, ring **B** of formula **I**, **I(a)**, **I(b)**, **I(f)** and/or **I(h)** is represented by the following structure:



25 In such case, **X₁** of formula **I**, **I(a)**, **I(b)**, **I(f)** and **I(h)** is represented by **X₇** in ring **B**, wherein **X₄**, **X₅**, **X₆** and **X₇**, **R₉₀**, **R₆₀**, **R₇₀** and **R₈₀** are as defined below.

[00106] It is understood that the wavy lines in ring **B** above, represent the connection points of ring **B** to the rest of the molecule (i.e., to -(C(R₁)(R₁'))_n from the left and to the carbonyl carbon atom from the right)

30 [00107] In some embodiments, **R₁** of compound of formula **I** and/or **I(a)**-**I(h)** is each independently H. In some embodiments, R₁ is each independently F, Cl, Br, or I; each represents a separate embodiment according to this invention. In some embodiments, R₁ is C(O)-R₁₀, wherein R₁₀ is as defined below. In some embodiments, R₁ is C(O)-CH₃. In some embodiments, R₁ is each independently C₁-C₅ linear or branched, substituted or unsubstituted alkyl. In some embodiments, R₁ is methyl. In some embodiments,

R₁ is ethyl. In some embodiments, R₁ is iso-propyl. In some embodiments, R₁ is each independently methyl, 2, 3, or 4-CH₂-C₆H₄-Cl, ethyl, propyl, iso-propyl, t-Bu, iso-butyl, pentyl, benzyl; each represents a separate embodiment according to this invention. In some embodiments, R₁ is each independently OH, SH, R₈-OH, CH₂-OH, R₈-SH, -R₈-O-R₁₀, -CH₂-O-CH₃, R₈-(C₃-C₈ cycloalkyl), CH₂-cyclohexyl, R₈-(C₃-C₈ heterocyclic ring), CH₂-imidazole, CH₂-indazole, CD₃, OCD₃, CN, NO₂, -CH₂CN, -R₈CN, NH₂, NHR, N(R)₂, R₈-N(R₁₀)(R₁₁), CH₂-NH₂, CH₂-N(CH₃)₂, R₉-R₈-N(R₁₀)(R₁₁), C≡C-CH₂-NH₂, B(OH)₂, -OC(O)CF₃, -OCH₂Ph, NHC(O)-R₁₀, NHC(O)CH₃, NHC(O)-N(R₁₀)(R₁₁), NHC(O)N(CH₃)₂, COOH, -C(O)Ph, C(O)O-R₁₀, C(O)O-CH₃, C(O)O-CH(CH₃)₂, C(O)O-CH₂CH₃, R₈-C(O)-R₁₀, CH₂C(O)CH₃, C(O)H, C(O)-R₁₀, C(O)-CH₃, C(O)-CH₂CH₃, C(O)-CH₂CH₂CH₃, C₁-C₅ linear or branched C(O)-haloalkyl, C(O)-CF₃, -C(O)NH₂, C(O)NHR, C(O)N(R₁₀)(R₁₁), C(O)N(CH₃)₂, SO₂R, SO₂N(R₁₀)(R₁₁), SO₂N(CH₃)₂, SO₂NHC(O)CH₃, C₁-C₅ linear or branched, substituted or unsubstituted alkyl, methyl, 2, 3, or 4-CH₂-C₆H₄-Cl, ethyl, propyl, iso-propyl, t-Bu, iso-butyl, pentyl, benzyl, C₁-C₅ linear or branched, substituted or unsubstituted alkenyl, CH=C(Ph)₂, C₁-C₅ linear or branched or C₃-C₈ cyclic haloalkyl, CF₃, CF₂CH₃, CH₂CF₃, CF₂CH₂CH₃, CH₂CH₂CF₃, CF₂CH(CH₃)₂, CF(CH₃)-CH(CH₃)₂, C₁-C₅ linear or branched or C₃-C₈ cyclic alkoxy, methoxy, ethoxy, propoxy, isopropoxy, O-CH₂-cyclopropyl, O-cyclobutyl, O-cyclopentyl, O-cyclohexyl, 1-butoxy, 2-butoxy, O-tBu, C₁-C₅ linear or branched or C₃-C₈ cyclic alkoxy wherein at least one methylene group (CH₂) in the alkoxy is replaced with an oxygen atom, O-1-oxacyclobutyl, O-2-oxacyclobutyl, C₁-C₅ linear or branched thioalkoxy, C₁-C₅ linear or branched haloalkoxy, OCF₃, OCHF₂, C₁-C₅ linear or branched alkoxyalkyl, substituted or unsubstituted C₃-C₈ cycloalkyl (e.g., cyclopropyl, cyclopentyl), substituted or unsubstituted C₃-C₈ heterocyclic ring, 3-methyl-4H-1,2,4-triazole, 5-methyl-1,2,4-oxadiazole, thiophene, oxazole, oxadiazole, imidazole, furane, triazole, tetrazole, pyridine (2, 3, or 4-pyridine), 3-methyl-2-pyridine, pyrimidine, pyrazine, oxacyclobutane (1 or 2-oxacyclobutane), indole, protonated or deprotonated pyridine oxide), substituted or unsubstituted aryl, phenyl, substituted or unsubstituted benzyl, 4-Cl-benzyl, 4-OH-benzyl; each represents a separate embodiment according to this invention. In some embodiments, R₁ is H, C₁-C₅ linear or branched, unsubstituted alkyl, methyl, ethyl, propyl, iso-propyl, t-Bu, iso-butyl, pentyl, benzyl, C(O)-R₁₀ or C(O)-CH₃; each represents a separate embodiment according to this invention. In some embodiments each R₁ may be further substituted with at least one substitution selected from: F, Cl, Br, I, OH, SH, C₁-C₅ linear or branched alkyl (e.g. methyl, ethyl), C₂-C₅ linear or branched alkenyl, C₂-C₅ linear or branched alkynyl (e.g. CCH), C₃-C₈ cycloalkyl, linear, branched or cyclic alkoxy, COOH, COO(R), NH₂, N(R)₂, CF₃, aryl, phenyl, heteroaryl (e.g., imidazole), C₃-C₈ cycloalkyl (e.g., cyclohexyl), CN and NO₂; each represents a separate embodiment according to this invention.

[00108] In some embodiments, R₁' of compound of formula **I** and/or **I(a)-I(h)** is each independently H. In some embodiments, R₁' is each independently F, Cl, Br, or I; each represents a separate embodiment according to this invention. In some embodiments, R₁' is C(O)-R₁₀, wherein R₁₀ is as defined below. In some embodiments, R₁' is C(O)-CH₃. In some embodiments, R₁' is each independently C₁-C₅ linear or branched, substituted or unsubstituted alkyl. In some embodiments, R₁' is methyl. In some embodiments, R₁' is ethyl. In some embodiments, R₁' is iso-propyl. In some embodiments, R₁' is methyl, 2, 3, or 4-CH₂-C₆H₄-Cl, ethyl, propyl, iso-propyl, t-Bu, iso-butyl, pentyl, benzyl; each represents a

separate embodiment according to this invention. In some embodiments, R_1' is each independently OH, SH, R_8 -OH, CH_2 -OH, R_8 -SH, $-R_8$ -O- R_{10} , $-CH_2$ -O- CH_3 , R_8 -(C_3 - C_8 cycloalkyl), CH_2 -cyclohexyl, R_8 -(C_3 - C_8 heterocyclic ring), CH_2 -imidazole, CH_2 -indazole, CD_3 , OCD_3 , CN, NO_2 , $-CH_2CN$, $-R_8CN$, NH_2 , NHR, $N(R)_2$, R_8 - $N(R_{10})(R_{11})$, CH_2 - NH_2 , CH_2 - $N(CH_3)_2$, R_9 - R_8 - $N(R_{10})(R_{11})$, $C\equiv C$ - CH_2 - NH_2 , $B(OH)_2$, $-OC(O)CF_3$, $-OCH_2Ph$, $NHC(O)-R_{10}$, $NHC(O)CH_3$, $NHC(O)-N(R_{10})(R_{11})$, $NHC(O)N(CH_3)_2$, $COOH$, $-C(O)Ph$, $C(O)O-R_{10}$, $C(O)O-CH_3$, $C(O)O-CH(CH_3)_2$, $C(O)O-CH_2CH_3$, $R_8-C(O)-R_{10}$, $CH_2C(O)CH_3$, $C(O)H$, $C(O)-R_{10}$, $C(O)-CH_3$, $C(O)-CH_2CH_3$, $C(O)-CH_2CH_2CH_3$, C_1 - C_5 linear or branched $C(O)$ -haloalkyl, $C(O)-CF_3$, $-C(O)NH_2$, $C(O)NHR$, $C(O)N(R_{10})(R_{11})$, $C(O)N(CH_3)_2$, SO_2R , $SO_2N(R_{10})(R_{11})$, $SO_2N(CH_3)_2$, $SO_2NHC(O)CH_3$, C_1 - C_5 linear or branched, substituted or unsubstituted alkyl, methyl, 2, 3, or 4- CH_2 - C_6H_4 -Cl, ethyl, propyl, iso-propyl, t-Bu, iso-butyl, pentyl, benzyl, C_1 - C_5 linear or branched, substituted or unsubstituted alkenyl, $CH=C(Ph)_2$, C_1 - C_5 linear or branched or C_3 - C_8 cyclic haloalkyl, CF_3 , CF_2CH_3 , CH_2CF_3 , $CF_2CH_2CH_3$, $CH_2CH_2CF_3$, $CF_2CH(CH_3)_2$, $CF(CH_3)-CH(CH_3)_2$, C_1 - C_5 linear or branched or C_3 - C_8 cyclic alkoxy, methoxy, ethoxy, propoxy, isopropoxy, O- CH_2 -cyclopropyl, O-cyclobutyl, O-cyclopentyl, O-cyclohexyl, 1-butoxy, 2-butoxy, O-tBu, C_1 - C_5 linear or branched or C_3 - C_8 cyclic alkoxy wherein at least one methylene group (CH_2) in the alkoxy is replaced with an oxygen atom, O-1-oxacyclobutyl, O-2-oxacyclobutyl, C_1 - C_5 linear or branched thioalkoxy, C_1 - C_5 linear or branched haloalkoxy, OCF_3 , $OCHF_2$, C_1 - C_5 linear or branched alkoxyalkyl, substituted or unsubstituted C_3 - C_8 cycloalkyl (e.g., cyclopropyl, cyclopentyl), substituted or unsubstituted C_3 - C_8 heterocyclic ring, 3-methyl-4H-1,2,4-triazole, 5-methyl-1,2,4-oxadiazole, thiophene, oxazole, oxadiazole, imidazole, furane, triazole, tetrazole, pyridine (2, 3, or 4-pyridine), 3-methyl-2-pyridine, pyrimidine, pyrazine, oxacyclobutane (1 or 2-oxacyclobutane), indole, protonated or deprotonated pyridine oxide), substituted or unsubstituted aryl, phenyl, substituted or unsubstituted benzyl, 4-Cl-benzyl, 4-OH-benzyl; each represents a separate embodiment according to this invention. In some embodiments, R_1' is H, C_1 - C_5 linear or branched, unsubstituted alkyl, methyl, ethyl, propyl, iso-propyl, t-Bu, iso-butyl, pentyl, benzyl, $C(O)-R_{10}$ or $C(O)-CH_3$; each represents a separate embodiment according to this invention. In some embodiments each R_1' may be further substituted with at least one substitution selected from: F, Cl, Br, I, OH, SH, C_1 - C_5 linear or branched alkyl (e.g. methyl, ethyl), C_2 - C_5 linear or branched alkenyl, C_2 - C_5 linear or branched alkynyl (e.g. CCH), C_3 - C_8 cycloalkyl, linear, branched or cyclic alkoxy, COOH, COO(R), NH_2 , $N(R)_2$, CF_3 , aryl, phenyl, heteroaryl (e.g., imidazole), C_3 - C_8 cycloalkyl (e.g., cyclohexyl), CN and NO_2 ; each represents a separate embodiment according to this invention.

[00109] In some embodiments, R_2 of compound of formula **I** and/or **I(a)**-**I(h)** is H. In some embodiments, R_2 is F, Cl, Br, or I; each represents a separate embodiment according to this invention. In some embodiments, R_2 is $C(O)-R_{10}$, wherein R_{10} is as defined below. In some embodiments, R_2 is $C(O)-CH_3$. In some embodiments, R_2 is C_1 - C_5 linear or branched, substituted or unsubstituted alkyl. In some embodiments, R_2 is C_1 - C_5 linear or branched, unsubstituted alkyl. In some embodiments, R_2 is methyl. In some embodiments, R_2 is ethyl. In some embodiments, R_2 is iso-propyl. In some embodiments, R_2 is methyl, 2, 3, or 4- CH_2 - C_6H_4 -Cl, ethyl, propyl, iso-propyl, t-Bu, iso-butyl, pentyl, benzyl; each represents a separate embodiment according to this invention. In some embodiments, R_2 is OH, SH, R_8 -OH, CH_2 -OH, R_8 -SH, $-R_8$ -O- R_{10} , $-CH_2$ -O- CH_3 , R_8 -(C_3 - C_8 cycloalkyl), CH_2 -cyclohexyl, R_8 -

(C₃-C₈ heterocyclic ring), CH₂-imidazole, CH₂-indazole, CD₃, OCD₃, CN, NO₂, -CH₂CN, -R₈CN, NH₂, NHR, N(R)₂, R₈-N(R₁₀)(R₁₁), CH₂-NH₂, CH₂-N(CH₃)₂, R₉-R₈-N(R₁₀)(R₁₁), C≡C-CH₂-NH₂, B(OH)₂, -OC(O)CF₃, -OCH₂Ph, NHC(O)-R₁₀, NHC(O)CH₃, NHC(O)-N(R₁₀)(R₁₁), NHC(O)N(CH₃)₂, COOH, -C(O)Ph, C(O)O-R₁₀, C(O)O-CH₃, C(O)O-CH(CH₃)₂, C(O)O-CH₂CH₃, R₈-C(O)-R₁₀, CH₂C(O)CH₃,
 5 C(O)H, C(O)-R₁₀, C(O)-CH₃, C(O)-CH₂CH₃, C(O)-CH₂CH₂CH₃, C₁-C₅ linear or branched C(O)-haloalkyl, C(O)-CF₃, -C(O)NH₂, C(O)NHR, C(O)N(R₁₀)(R₁₁), C(O)N(CH₃)₂, SO₂R, SO₂N(R₁₀)(R₁₁), SO₂N(CH₃)₂, SO₂NHC(O)CH₃, C₁-C₅ linear or branched, substituted or unsubstituted alkyl, methyl, 2, 3, or 4-CH₂-C₆H₄-Cl, ethyl, propyl, iso-propyl, t-Bu, iso-butyl, pentyl, benzyl, C₁-C₅ linear or branched, substituted or unsubstituted alkenyl, CH=C(Ph)₂, C₁-C₅ linear or branched or C₃-C₈ cyclic haloalkyl,
 10 CF₃, CF₂CH₃, CH₂CF₃, CF₂CH₂CH₃, CH₂CH₂CF₃, CF₂CH(CH₃)₂, CF(CH₃)-CH(CH₃)₂, C₁-C₅ linear or branched or C₃-C₈ cyclic alkoxy, methoxy, ethoxy, propoxy, isopropoxy, O-CH₂-cyclopropyl, O-cyclobutyl, O-cyclopentyl, O-cyclohexyl, 1-butoxy, 2-butoxy, O-tBu, C₁-C₅ linear or branched or C₃-C₈ cyclic alkoxy wherein at least one methylene group (CH₂) in the alkoxy is replaced with an oxygen atom, O-1-oxacyclobutyl, O-2-oxacyclobutyl, C₁-C₅ linear or branched thioalkoxy, C₁-C₅ linear or
 15 branched haloalkoxy, OCF₃, OCHF₂, C₁-C₅ linear or branched alkoxyalkyl, substituted or unsubstituted C₃-C₈ cycloalkyl (e.g., cyclopropyl, cyclopentyl), substituted or unsubstituted C₃-C₈ heterocyclic ring, 3-methyl-4H-1,2,4-triazole, 5-methyl-1,2,4-oxadiazole, thiophene, oxazole, oxadiazole, imidazole, furane, triazole, tetrazole, pyridine (2, 3, or 4-pyridine), 3-methyl-2-pyridine, pyrimidine, pyrazine, oxacyclobutane (1 or 2-oxacyclobutane), indole, protonated or deprotonated pyridine oxide), substituted
 20 or unsubstituted aryl, phenyl, substituted or unsubstituted benzyl, 4-Cl-benzyl, 4-OH-benzyl; each represents a separate embodiment according to this invention. In some embodiments, R₂ is H, C₁-C₅ linear or branched, unsubstituted alkyl, methyl, ethyl, propyl, iso-propyl, t-Bu, iso-butyl, pentyl, benzyl, C(O)-R₁₀ or C(O)-CH₃; each represents a separate embodiment according to this invention. In some embodiments R₂ may be further substituted with at least one substitution selected from: F, Cl, Br, I, OH,
 25 SH, C₁-C₅ linear or branched alkyl (e.g. methyl, ethyl), C₂-C₅ linear or branched alkenyl, C₂-C₅ linear or branched alkynyl (e.g. CCH), C₃-C₈ cycloalkyl, linear, branched or cyclic alkoxy, COOH, COO(R), NH₂, N(R)₂, CF₃, aryl, phenyl, heteroaryl (e.g., imidazole), C₃-C₈ cycloalkyl (e.g., cyclohexyl), CN and NO₂; each represents a separate embodiment according to this invention.

[00110] In some embodiments, **R₂'** of compound of **I** and/or **I(a)-I(h)** is H. In some embodiments, R₂' is F, Cl, Br, or I; each represents a separate embodiment according to this invention. In some
 30 embodiments, R₂' is C(O)-R₁₀, wherein R₁₀ is as defined below. In some embodiments, R₂' is C(O)-CH₃. In some embodiments, R₂' is C₁-C₅ linear or branched, substituted or unsubstituted alkyl. In some embodiments, R₂' is C₁-C₅ linear or branched, unsubstituted alkyl. In some embodiments, R₂' is methyl. In some embodiments, R₂' is ethyl. In some embodiments, R₂' is iso-propyl. In some embodiments, R₂'
 35 is methyl, 2, 3, or 4-CH₂-C₆H₄-Cl, ethyl, propyl, iso-propyl, t-Bu, iso-butyl, pentyl, benzyl; each represents a separate embodiment according to this invention. In some embodiments, R₂' is OH, SH, R₈-OH, CH₂-OH, R₈-SH, -R₈-O-R₁₀, -CH₂-O-CH₃, R₈-(C₃-C₈ cycloalkyl), CH₂-cyclohexyl, R₈-(C₃-C₈ heterocyclic ring), CH₂-imidazole, CH₂-indazole, CD₃, OCD₃, CN, NO₂, -CH₂CN, -R₈CN, NH₂, NHR, N(R)₂, R₈-N(R₁₀)(R₁₁), CH₂-NH₂, CH₂-N(CH₃)₂, R₉-R₈-N(R₁₀)(R₁₁), C≡C-CH₂-NH₂, B(OH)₂, -

OC(O)CF₃, -OCH₂Ph, NHC(O)-R₁₀, NHC(O)CH₃, NHC(O)-N(R₁₀)(R₁₁), NHC(O)N(CH₃)₂, COOH, -C(O)Ph, C(O)O-R₁₀, C(O)O-CH₃, C(O)O-CH(CH₃)₂, C(O)O-CH₂CH₃, R₈-C(O)-R₁₀, CH₂C(O)CH₃, C(O)H, C(O)-R₁₀, C(O)-CH₃, C(O)-CH₂CH₃, C(O)-CH₂CH₂CH₃, C₁-C₅ linear or branched C(O)-haloalkyl, C(O)-CF₃, -C(O)NH₂, C(O)NHR, C(O)N(R₁₀)(R₁₁), C(O)N(CH₃)₂, SO₂R, SO₂N(R₁₀)(R₁₁),
 5 SO₂N(CH₃)₂, SO₂NHC(O)CH₃, C₁-C₅ linear or branched, substituted or unsubstituted alkyl, methyl, 2, 3, or 4-CH₂-C₆H₄-Cl, ethyl, propyl, iso-propyl, t-Bu, iso-butyl, pentyl, benzyl, C₁-C₅ linear or branched, substituted or unsubstituted alkenyl, CH=C(Ph)₂, C₁-C₅ linear or branched or C₃-C₈ cyclic haloalkyl, CF₃, CF₂CH₃, CH₂CF₃, CF₂CH₂CH₃, CH₂CH₂CF₃, CF₂CH(CH₃)₂, CF(CH₃)-CH(CH₃)₂, C₁-C₅ linear or branched or C₃-C₈ cyclic alkoxy, methoxy, ethoxy, propoxy, isopropoxy, O-CH₂-cyclopropyl, O-cyclobutyl, O-cyclopentyl, O-cyclohexyl, 1-butoxy, 2-butoxy, O-tBu, C₁-C₅ linear or branched or C₃-C₈ cyclic alkoxy wherein at least one methylene group (CH₂) in the alkoxy is replaced with an oxygen atom, O-1-oxacyclobutyl, O-2-oxacyclobutyl, C₁-C₅ linear or branched thioalkoxy, C₁-C₅ linear or branched haloalkoxy, OCF₃, OCHF₂, C₁-C₅ linear or branched alkoxyalkyl, substituted or unsubstituted C₃-C₈ cycloalkyl (e.g., cyclopropyl, cyclopentyl), substituted or unsubstituted C₃-C₈ heterocyclic ring,
 15 3-methyl-4H-1,2,4-triazole, 5-methyl-1,2,4-oxadiazole, thiophene, oxazole, oxadiazole, imidazole, furane, triazole, tetrazole, pyridine (2, 3, or 4-pyridine), 3-methyl-2-pyridine, pyrimidine, pyrazine, oxacyclobutane (1 or 2-oxacyclobutane), indole, protonated or deprotonated pyridine oxide), substituted or unsubstituted aryl, phenyl, substituted or unsubstituted benzyl, 4-Cl-benzyl, 4-OH-benzyl; each represents a separate embodiment according to this invention. In some embodiments, R₂' is H, C₁-C₅ linear or branched, unsubstituted alkyl, methyl, ethyl, propyl, iso-propyl, t-Bu, iso-butyl, pentyl, benzyl, C(O)-R₁₀ or C(O)-CH₃; each represents a separate embodiment according to this invention. In some embodiments R₂' may be further substituted with at least one substitution selected from: F, Cl, Br, I, OH, SH, C₁-C₅ linear or branched alkyl (e.g. methyl, ethyl), C₂-C₅ linear or branched alkenyl, C₂-C₅ linear or branched alkynyl (e.g. CCH), C₃-C₈ cycloalkyl, linear, branched or cyclic alkoxy, COOH, COO(R), NH₂, N(R)₂, CF₃, aryl, phenyl, heteroaryl (e.g., imidazole), C₃-C₈ cycloalkyl (e.g., cyclohexyl), CN and NO₂; each represents a separate embodiment according to this invention.

[00111] In some embodiments, **R**₄₀ of compound of **I**, **I(a)**-**I(c)**, **I(g)** and/or **I(ga)** is H. In some embodiments, **R**₄₀ is F, Cl, Br, or I; each represents a separate embodiment according to this invention. In some embodiments, **R**₄₀ is C(O)-R₁₀, wherein R₁₀ is as defined below. In some embodiments, **R**₄₀ is C(O)-CH₃. In some embodiments, **R**₄₀ is C₁-C₅ linear or branched, substituted or unsubstituted alkyl. In some embodiments, **R**₄₀ is C₁-C₅ linear or branched, unsubstituted alkyl. In some embodiments, **R**₄₀ is methyl. In some embodiments, **R**₄₀ is ethyl. In some embodiments, **R**₄₀ is iso-propyl. In some embodiments, **R**₄₀ is methyl, 2, 3, or 4-CH₂-C₆H₄-Cl, ethyl, propyl, iso-propyl, t-Bu, iso-butyl, pentyl, benzyl; each represents a separate embodiment according to this invention. In some embodiments, **R**₄₀ is OH, SH, R₈-OH, CH₂-OH, R₈-SH, -R₈-O-R₁₀, -CH₂-O-CH₃, R₈-(C₃-C₈ cycloalkyl), CH₂-cyclohexyl, R₈-(C₃-C₈ heterocyclic ring), CH₂-imidazole, CH₂-indazole, CD₃, OCD₃, CN, NO₂, -CH₂CN, -R₈CN, NH₂, NHR, N(R)₂, R₈-N(R₁₀)(R₁₁), CH₂-NH₂, CH₂-N(CH₃)₂, R₉-R₈-N(R₁₀)(R₁₁), C≡C-CH₂-NH₂, B(OH)₂, -OC(O)CF₃, -OCH₂Ph, NHC(O)-R₁₀, NHC(O)CH₃, NHC(O)-N(R₁₀)(R₁₁), NHC(O)N(CH₃)₂, COOH, -C(O)Ph, C(O)O-R₁₀, C(O)O-CH₃, C(O)O-CH(CH₃)₂, C(O)O-CH₂CH₃, R₈-C(O)-R₁₀,

CH₂C(O)CH₃, C(O)H, C(O)-R₁₀, C(O)-CH₃, C(O)-CH₂CH₃, C(O)-CH₂CH₂CH₃, C₁-C₅ linear or branched C(O)-haloalkyl, C(O)-CF₃, -C(O)NH₂, C(O)NHR, C(O)N(R₁₀)(R₁₁), C(O)N(CH₃)₂, SO₂R, SO₂N(R₁₀)(R₁₁), SO₂N(CH₃)₂, SO₂NHC(O)CH₃, C₁-C₅ linear or branched, substituted or unsubstituted alkyl, methyl, 2, 3, or 4-CH₂-C₆H₄-Cl, ethyl, propyl, iso-propyl, t-Bu, iso-butyl, pentyl, benzyl, C₁-C₅ linear or branched, substituted or unsubstituted alkenyl, CH=C(Ph)₂, C₁-C₅ linear or branched or C₃-C₈ cyclic haloalkyl, CF₃, CF₂CH₃, CH₂CF₃, CF₂CH₂CH₃, CH₂CH₂CF₃, CF₂CH(CH₃)₂, CF(CH₃)-CH(CH₃)₂, C₁-C₅ linear or branched or C₃-C₈ cyclic alkoxy, methoxy, ethoxy, propoxy, isopropoxy, O-CH₂-cyclopropyl, O-cyclobutyl, O-cyclopentyl, O-cyclohexyl, 1-butoxy, 2-butoxy, O-tBu, C₁-C₅ linear or branched or C₃-C₈ cyclic alkoxy wherein at least one methylene group (CH₂) in the alkoxy is replaced with an oxygen atom, O-1-oxacyclobutyl, O-2-oxacyclobutyl, C₁-C₅ linear or branched thioalkoxy, C₁-C₅ linear or branched haloalkoxy, OCF₃, OCHF₂, C₁-C₅ linear or branched alkoxyalkyl, substituted or unsubstituted C₃-C₈ cycloalkyl (e.g., cyclopropyl, cyclopentyl), substituted or unsubstituted C₃-C₈ heterocyclic ring, 3-methyl-4H-1,2,4-triazole, 5-methyl-1,2,4-oxadiazole, thiophene, oxazole, oxadiazole, imidazole, furane, triazole, tetrazole, pyridine (2, 3, or 4-pyridine), 3-methyl-2-pyridine, pyrimidine, pyrazine, oxacyclobutane (1 or 2-oxacyclobutane), indole, protonated or deprotonated pyridine oxide), substituted or unsubstituted aryl, phenyl, substituted or unsubstituted benzyl, 4-Cl-benzyl, 4-OH-benzyl; each represents a separate embodiment according to this invention. In some embodiments, R₂' is H, C₁-C₅ linear or branched, unsubstituted alkyl, methyl, ethyl, propyl, iso-propyl, t-Bu, iso-butyl, pentyl, benzyl, C(O)-R₁₀ or C(O)-CH₃; each represents a separate embodiment according to this invention. In some embodiments R₄₀ may be further substituted with at least one substitution selected from: F, Cl, Br, I, OH, SH, C₁-C₅ linear or branched alkyl (e.g. methyl, ethyl), C₂-C₅ linear or branched alkenyl, C₂-C₅ linear or branched alkynyl (e.g. CCH), C₃-C₈ cycloalkyl, linear, branched or cyclic alkoxy, COOH, COO(R), NH₂, N(R)₂, CF₃, aryl, phenyl, heteroaryl (e.g., imidazole), C₃-C₈ cycloalkyl (e.g., cyclohexyl), CN and NO₂; each represents a separate embodiment according to this invention.

[00112] In some embodiments, R₂ and R₂' of compound of formula I, and I(a)-I(h) are identical. In some embodiments, R₂ and R₂' are both H. In some embodiments, R₂ and R₂' are both methyl. In some embodiments, R₂ and R₂' are different. In some embodiments, R₂ is H and R₂' is a C₁-C₅ linear or branched, substituted or unsubstituted alkyl. In some embodiments, R₂ is H and R₂' is a methyl. In some embodiments, R₂ is H and R₂' is an ethyl.

[00113] In some embodiments, R₂ and R₁ of formula I and I(a)-I(h) (in some embodiments, where n is 1), are joined to form a C₃-C₈ substituted or unsubstituted, carbocyclic or heterocyclic ring. In some embodiments, R₂ and R₁ are joined to form a C₃-C₈ carbocyclic ring (e.g., cyclopropane, cyclopentane, cyclohexane). In some embodiments, R₂ and R₁ are joined to form a C₃-C₈ heterocyclic ring.

[00114] In some embodiments, R₃ of compound of formula I, I(a), and I(h) is H, F, Cl, Br, I, OH, SH, =O, R₈-OH, CH₂-OH, R₈-SH, -R₈-O-R₁₀, CH₂-O-CH₃, CD₃, OCD₃, CN, NO₂, -CH₂CN, -R₈CN, NH₂, NHNH₂, NHR, N(R)₂, R₈-N(R₁₀)(R₁₁), CH₂-NH₂, CH₂-N(CH₃)₂, R₉-R₈-N(R₁₀)(R₁₁), B(OH)₂, -OC(O)CF₃, -OCH₂Ph, NHC(O)OBz, -NHC(O)-R₁₀, NHC(O)CH₃, NHC(O)-N(R₁₀)(R₁₁), NHC(O)N(CH₃)₂, COOH, -C(O)Ph, C(O)O-R₁₀, C(O)O-CH₃, C(O)O-CH₂CH₃, R₈-C(O)-R₁₀,

CH₂C(O)CH₃, C(O)H, C(O)-R₁₀, C(O)-CH₃, C(O)-CH₂CH₃, C(O)-CH₂CH₂CH₃, C₁-C₅ linear or branched C(O)-haloalkyl, C(O)-CF₃, -C(O)NH₂, C(O)NHR, C(O)N(R₁₀)(R₁₁), C(O)N(CH₃)₂, SO₂R, SO₂N(R₁₀)(R₁₁), SO₂N(CH₃)₂, C₁-C₅ linear or branched, substituted or unsubstituted alkyl, methyl, C(OH)(CH₃)(Ph), ethyl, propyl, iso-propyl, t-Bu, iso-butyl, pentyl, C₁-C₅ linear or branched or C₃-C₈ cyclic haloalkyl, CF₃, CF₂CH₃, CF₂-cyclobutyl, CH₂CF₃, CF₂CH₂CH₃, CH₂CH₂CF₃, CF₂CH(CH₃)₂, CF(CH₃)-CH(CH₃)₂, C₁-C₅ linear or branched or C₃-C₈ cyclic alkoxy, methoxy, ethoxy, propoxy, isopropoxy, O-CH₂-cyclopropyl, C₁-C₅ linear or branched thioalkoxy, C₁-C₅ linear or branched haloalkoxy, C₁-C₅ linear or branched alkoxyalkyl, substituted or unsubstituted C₃-C₈ cycloalkyl, cyclopropyl, cyclopentyl, substituted or unsubstituted C₃-C₈ heterocyclic ring, 3-methyl-4H-1,2,4-triazole, 5-methyl-1,2,4-oxadiazole, thiophene, oxazole, isoxazole, imidazole, furane, triazole, pyridine (2, 3, or 4-pyridine), pyrimidine, pyrazine, oxacyclobutane (1 or 2-oxacyclobutane), indole, substituted or unsubstituted aryl, or phenyl; each represents a separate embodiment according to this invention.

[00115] In some embodiments, **R₃** of compound of formula **I**, **I(a)**, **I(b)**, **I(c)**, **I(e)**, **I(g)**, **I(ga)** and **I(h)** is OH, F, SH, NH₂, NHNH₂, NHR, N(R)₂, NHC(O)OBz, -NHC(O)-R₁₀ (e.g., NHC(O)CH₃), C₁-C₅ linear or branched, substituted or unsubstituted alkyl (e.g., methyl, ethyl, propyl, iso-propyl, t-Bu, iso-butyl, pentyl), substituted or unsubstituted C₃-C₈ cycloalkyl, C₁-C₅ linear or branched or C₃-C₈ cyclic haloalkyl, C₁-C₅ linear or branched or C₃-C₈ cyclic alkoxy, substituted or unsubstituted C₃-C₈ heterocyclic ring or substituted or unsubstituted aryl; each represents a separate embodiment according to this invention. In some embodiments, **R₃** is OH. In some embodiments, **R₃** is F. In some embodiments, **R₃** is F, Cl, Br or I; each represents a separate embodiment. In some embodiments, **R₃** is SH. In some embodiments, **R₃** is NH₂. In some embodiments, **R₃** is NHNH₂. In some embodiments, **R₃** is NHR. In some embodiments, **R₃** is N(R)₂. In some embodiments, **R₃** is NHC(O)OBz. In some embodiments, **R₃** is -NHC(O)-R₁₀. In some embodiments, **R₃** is NHC(O)CH₃. In some embodiments, **R₃** is C₁-C₅ linear or branched, substituted or unsubstituted alkyl. In some embodiments, **R₃** is methyl. In some embodiments, **R₃** is ethyl. In some embodiments, **R₃** is propyl. In some embodiments, **R₃** is iso-propyl. In some embodiments, **R₃** is t-Bu. In some embodiments, **R₃** is iso-butyl. In some embodiments, **R₃** is pentyl. In some embodiments, **R₃** is substituted or unsubstituted C₃-C₈ cycloalkyl. In some embodiments, **R₃** is C₁-C₅ linear or branched or C₃-C₈ cyclic haloalkyl. In some embodiments, **R₃** is C₁-C₅ linear or branched or C₃-C₈ cyclic alkoxy. In some embodiments, **R₃** is substituted or unsubstituted C₃-C₈ heterocyclic ring.

In some embodiments, **R₃** is substituted or unsubstituted aryl. In some embodiments, **R₃** of compound of formula **I(f)** is SH, NHNH₂, or NHC(O)OBz; each represents a separate embodiment according to this invention. In some embodiments, **R₃** may be further substituted with at least one substitution selected from: F, Cl, Br, I, OH, SH, C₁-C₅ linear or branched alkyl (e.g. methyl, ethyl), C₂-C₅ linear or branched alkenyl, C₂-C₅ linear or branched alkynyl (e.g. CCH), C₃-C₈ cycloalkyl, linear, branched or cyclic alkoxy, COOH, COO(R), NH₂, N(R)₂, CF₃, aryl, phenyl, heteroaryl (e.g., imidazole), C₃-C₈ cycloalkyl (e.g., cyclohexyl), CN and NO₂; each represents a separate embodiment according to this invention. In some embodiments, **R₃** of compound of formula **I**, **I(a)**, **I(b)**, **I(c)**, **I(e)**, **I(g)**, **I(ga)** and/or **I(h)** is OH or NH₂.

[00116] In some embodiments, **R**₃ and **R**₂ of compound of any one of formula **I-I(h)** are joined to form a C₃-C₈ substituted or unsubstituted, carbocyclic or heterocyclic ring. In some embodiments, **R**₃ and **R**₂ are joined to form a C₃-C₈ carbocyclic ring. In some embodiments, **R**₃ and **R**₂ are joined to form a cyclopropyl. In some embodiments, **R**₃ and **R**₂ are joined to form a heterocyclic ring.

5 [00117] In some embodiments, **R**₄ of compound of formula **I**, **I(a)**, **I(b)**, and/or **I(f)** is H, F, Cl, Br, I, OH, SH, =O, =NH-OH, R₈-OH, CH₂-OH, R₈-SH, -R₈-O-R₁₀, CH₂-O-CH₃, CD₃, OCD₃, CN, NO₂, -CH₂CN, -R₈CN, NH₂, NHNH₂, NHR, N(R)₂, R₈-N(R₁₀)(R₁₁), CH₂-NH₂, CH₂-N(CH₃)₂, R₉-R₈-N(R₁₀)(R₁₁), B(OH)₂, -OC(O)CF₃, -OCH₂Ph, NHC(O)OBz, -NHC(O)-R₁₀, NHC(O)H, NHC(O)CH₃, NHC(O)-N(R₁₀)(R₁₁), NHC(O)N(CH₃)₂, COOH, -C(O)Ph, C(O)O-R₁₀, C(O)O-CH₃, C(O)O-CH₂CH₃,
 10 R₈-C(O)-R₁₀, CH₂C(O)CH₃, C(O)H, C(O)-R₁₀, C(O)-CH₃, C(O)-CH₂CH₃, C(O)-CH₂CH₂CH₃, C₁-C₅ linear or branched C(O)-haloalkyl, C(O)-CF₃, -C(O)NH₂, C(O)NHR, C(O)N(R₁₀)(R₁₁), C(O)N(CH₃)₂, SO₂R, SO₂N(R₁₀)(R₁₁), SO₂N(CH₃)₂, C₁-C₅ linear or branched, substituted or unsubstituted alkyl, methyl, C(OH)(CH₃)(Ph), ethyl, propyl, iso-propyl, t-Bu, iso-butyl, pentyl, C₁-C₅ linear or branched or C₃-C₈ cyclic haloalkyl, CF₃, CF₂CH₃, CF₂-cyclobutyl, CH₂CF₃, CF₂CH₂CH₃, CH₂CH₂CF₃,
 15 CF₂CH(CH₃)₂, CF(CH₃)-CH(CH₃)₂, C₁-C₅ linear or branched or C₃-C₈ cyclic alkoxy, methoxy, ethoxy, propoxy, isopropoxy, O-CH₂-cyclopropyl, C₁-C₅ linear or branched thioalkoxy, C₁-C₅ linear or branched haloalkoxy, C₁-C₅ linear or branched alkoxyalkyl, substituted or unsubstituted C₃-C₈ cycloalkyl, cyclopropyl, cyclopentyl, substituted or unsubstituted C₃-C₈ heterocyclic ring, 3-methyl-4H-1,2,4-triazole, 5-methyl-1,2,4-oxadiazole, thiophene, oxazole, isoxazole, imidazole, furane, triazole, pyridine
 20 (2, 3, or 4-pyridine), pyrimidine, pyrazine, oxacyclobutane (1 or 2-oxacyclobutane), indole, substituted or unsubstituted aryl, or phenyl; each represents a separate embodiment according to this invention. In some embodiments, **R**₄ is H. In some embodiments, **R**₄ is =O. In some embodiments, **R**₄ is =NH-OH. In some embodiments, **R**₄ is NH₂. In some embodiments, **R**₄ is OH. In some embodiments, **R**₄ is -NHC(O)-R₁₀. In some embodiments, **R**₄ is NHC(O)H. In some embodiments, **R**₄ is NHC(O)CH₃. In some
 25 embodiments, **R**₄ of compound is NHNH₂. In some embodiments, **R**₄ is alkyl. In some embodiments, **R**₄ is methyl. In some embodiments, **R**₄ may be further substituted with at least one substitution selected from: F, Cl, Br, I, OH, SH, C₁-C₅ linear or branched alkyl (e.g. methyl, ethyl), C₂-C₅ linear or branched alkenyl, C₂-C₅ linear or branched alkynyl (e.g. CCH), C₃-C₈ cycloalkyl, linear, branched or cyclic alkoxy, COOH, COO(R), NH₂, N(R)₂, CF₃, aryl, phenyl, heteroaryl (e.g., imidazole), C₃-C₈ cycloalkyl
 30 (e.g., cyclohexyl), CN and NO₂; each represents a separate embodiment according to this invention.

[00118] In some embodiments, **R**₄ of compound of formula **I-I(c)** and/or **I(e)-I(ga)** is NH₂, OH, NHNH₂, NHR, N(R)₂, -NHC(O)-R₁₀, NHC(O)H, NHC(O)CH₃, C₁-C₅ linear or branched, substituted or unsubstituted alkyl, C₁-C₅ linear or branched or C₃-C₈ cyclic haloalkyl, C₁-C₅ linear or branched or C₃-C₈ cyclic alkoxy, substituted or unsubstituted C₃-C₈ cycloalkyl, substituted or unsubstituted C₃-C₈
 35 heterocyclic ring, or substituted or unsubstituted aryl; each represents a separate embodiment according to this invention. In some embodiments, **R**₄ is NH₂. In some embodiments, **R**₄ is OH. In some embodiments, **R**₄ is alkyl. In some embodiments, **R**₄ is methyl. In some embodiments, **R**₄ may be further substituted with at least one substitution selected from: F, Cl, Br, I, OH, SH, C₁-C₅ linear or branched alkyl (e.g. methyl, ethyl), C₂-C₅ linear or branched alkenyl, C₂-C₅ linear or branched alkynyl (e.g. CCH),


C₃-C₈ cycloalkyl, linear, branched or cyclic alkoxy, COOH, COO(R), NH₂, N(R)₂, CF₃, aryl, phenyl, heteroaryl (e.g., imidazole), C₃-C₈ cycloalkyl (e.g., cyclohexyl), CN and NO₂; each represents a separate embodiment according to this invention. In some embodiments, **R₄** of compound of formula **I**, **I(a)**, **I(b)**, **I(c)**, **I(e)**, **I(g)**, **I(ga)** and/or **I(f)** is OH or NH₂.

5 [00119] In some embodiments, if **R₃** of compound of formula **I**, **I(a)**, **I(b)**, **I(c)**, **I(e)**, **I(g)**, **I(ga)** is OH then **R₄** is NH₂ and if **R₃** is NH₂ then **R₄** is OH. In some embodiments, if **R₃** is OH and **R₄** is NH₂, then **n + m** cannot be equal to 3. In some embodiments, **R₃** and **R₄** cannot both be NH₂.

[00120] In some embodiments, **R₃** and **R₄** of compound of formula **I**, **I(a)**-**I(c)**, **I(e)**, **I(g)** and/or **I(ga)** are joined together to form ring **A**. In some embodiments, ring **A** has two chiral centers.

10 [00121] In some embodiments, ring **A** of formula **I** and/or **I(e)** is a substituted aryl. In some embodiments, ring **A** is 2-amino-phenyl. In some embodiments, ring **A** is methyloxazolidin-2-one. In some embodiments, ring **A** is a substituted or unsubstituted cycloalkyl. In some embodiments, ring **A** is cyclopentyl. In some embodiments, ring **A** is cyclohexyl. In some embodiments, ring **A** is a substituted cycloalkyl. In some embodiment, the substitution is at least one selected from: F, Cl, Br, I, OH, SH, C₁-
15 C₅ linear or branched alkyl, C₃-C₈ cycloalkyl, linear, branched or cyclic alkoxy, COOH, COO(R), NH₂, N(R)₂, CF₃, aryl, phenyl, heteroaryl, C₃-C₈ cycloalkyl, CN and NO₂; each represents a separate embodiment according to this invention. In some embodiments, ring **A** is substituted cyclopropyl. In some embodiments, ring **A** is substituted cyclobutyl. In some embodiments, ring **A** is substituted cyclopentyl. In some embodiments, ring **A** is substituted cyclohexyl. In some embodiments, ring **A** is substituted at least with NH₂. In some embodiments, ring **A** is substituted with NH₂. In some
20 embodiments, ring **A** is a cycloalkyl, substituted at least with NH₂. In some embodiments, ring **A** is a cycloalkyl, substituted with NH₂. In some embodiments, ring **A** is a 5 or 6 membered nitrogen containing heterocyclic ring. In some embodiments, ring **A** is 1, 2, or 3-piperidine, oxazolidin-2-one, tetrahydropyrimidine, pyridine, dihydro-thiazine, dihydro-imidazole, tetrahydropyridine, or pyrrolidine,
25 which may be substituted or unsubstituted; each is a separate embodiment according to this invention. In some embodiments, ring **A** is substituted at least with NH₂. In some embodiment, **R₃** and **R₄** of formula **I** and/or **I(e)** are joined to form a 5 or 6 membered substituted or unsubstituted, aliphatic or aromatic, carbocyclic or heterocyclic ring. In some embodiment, **R₃** and **R₄** are joined to form a 5 membered unsubstituted, aliphatic heterocyclic ring. In some embodiment, **R₃** and **R₄** are joined to form a 6
30 membered unsubstituted, aliphatic heterocyclic ring. In some embodiment, **R₃** and **R₄** are joined to form a 5 membered substituted, aliphatic heterocyclic ring. In some embodiment, **R₃** and **R₄** are joined to form a 5 membered substituted, aliphatic heterocyclic ring. In some embodiment, **R₃** and **R₄** are joined to form a 6 membered substituted, aliphatic heterocyclic ring. In some embodiment, **R₃** and **R₄** are joined to form a 5 membered unsubstituted, aromatic heterocyclic ring. In some embodiment, **R₃** and **R₄** are joined to form a 6
35 membered unsubstituted, aromatic heterocyclic ring. In some embodiment, **R₃** and **R₄** are joined to form a 5 membered substituted, aromatic heterocyclic ring. In some embodiment, **R₃** and **R₄** are joined to form a 6 membered substituted, aromatic heterocyclic ring. In some embodiment, **R₃** and **R₄** are joined to form an arene.

[00122] In some embodiment, R_3 and R_4 of compound of formula **I**, **I(a)**, **I(b)**, **I(e)**, **I(g)**, and/or **I(ga)** are joined to form ring **A** represented by the following structure, wherein R_2 and R_2' are as defined above:



5 [00123] It is understood that the wavy line in ring **A** above, represents the connection point of **A** to the rest of the molecule (i.e., to $-(C(R_1)(R_1'))_n$ in formula **I-I(h)**).

[00124] In some embodiment, R_5 of compound of formula **I**, **I(a)**-**I(h)** and/or **X-X(d)**, is H, C_1 - C_5 linear or branched, substituted or unsubstituted alkyl, methyl, CH_2SH , ethyl, iso-propyl, butyl, CH_2-CCH , $CH_2-C(O)-OCH_3$, C_2 - C_5 linear or branched, substituted or unsubstituted alkenyl, C_2 - C_5 linear or branched, substituted or unsubstituted alkynyl, CCH , CH_2-CCH , C_1 - C_5 linear or branched haloalkyl, CF_3 , CF_2CH_3 , CH_2CF_3 , $CF_2CH_2CH_3$, $CH_2CH_2CF_3$, $CF_2CH(CH_3)_2$, $CF(CH_3)-CH(CH_3)_2$, R_8 -aryl, CH_2-Ph , $C(=CH_2)-R_{10}$, $C(=CH_2)-C(O)-OCH_3$, $C(=CH_2)-CN$, substituted or unsubstituted alkyl sulfone, SO_2-CH_2 -cyclopentyl, substituted or unsubstituted aryl, phenyl, substituted or unsubstituted heteroaryl, or pyridine (2, 3, and 4-pyridine); each represents a separate embodiment according to this invention. In some

15 embodiments, R_5 is H, C_1 - C_5 linear or branched, substituted or unsubstituted alkyl, methyl, ethyl, iso-propyl, butyl, CH_2-CCH , $CH_2-C(O)-OCH_3$, C_2 - C_5 linear or branched, substituted or unsubstituted alkenyl, C_2 - C_5 linear or branched, substituted or unsubstituted alkynyl, CCH , CH_2-CCH , C_1 - C_5 linear or branched haloalkyl, substituted or unsubstituted alkyl sulfone, SO_2-CH_2 -cyclopentyl, or substituted or unsubstituted aryl. In some embodiments, R_5 is H. In some embodiments, R_5 is C_1 - C_5 linear or branched,

20 substituted or unsubstituted alkyl. In some embodiments, R_5 is C_1 - C_5 linear alkyl. In some embodiments, R_5 is branched C_1 - C_5 alkyl. In some embodiments, R_5 is methyl. In some embodiments, R_5 is ethyl. In some embodiments, R_5 is butyl. In some embodiments, R_5 is a substituted C_1 - C_5 alkyl. In some embodiments, R_5 is a C_1 - C_5 alkyl substituted with a C_2 - C_5 linear or branched alkynyl. In some embodiments, R_5 is CH_2-CCH . In some embodiments, R_5 is a C_1 - C_5 alkyl substituted with CCH . In some

25 embodiments, R_5 is $CH_2-C(O)-OCH_3$. In some embodiments, R_5 is substituted or unsubstituted alkyl sulfone. In some embodiments, R_5 of compound of formula **I**, **I(a)**-**I(h)** is substituted alkyl sulfone. In some embodiments, R_5 of compound of formula **I**, **I(a)**-**I(h)** is SO_2-CH_2 -cyclopentyl. In some embodiments, R_5 may be further substituted with at least one substitution selected from: F, Cl, Br, I, OH, SH, C_1 - C_5 linear or branched alkyl, C_2 - C_5 linear or branched alkenyl, C_2 - C_5 linear or branched alkynyl (e.g. CCH), C_3 - C_8 cycloalkyl, linear, branched or cyclic alkoxy, $COOH$, $COO(R)$, NH_2 , $N(R)_2$, CF_3 , aryl, phenyl, heteroaryl, C_3 - C_8 cycloalkyl, CN and NO_2 ; each represents a separate embodiment according to this invention.

[00125] In some embodiment, R_{60} of compound of formula **I(a)**, **I(b)**, **I(d)**, **I(e)**, **I(f)** and/or **I(h)** is H, F, Cl, Br, I, OH, SH, R_8-OH , R_8-SH , $-R_8-O-R_{10}$, CF_3 , CN, NO_2 , NH_2 , NHR , $N(R)_2$, $R_8-N(R_{10})(R_{11})$, -

35 $OC(O)CF_3$, $-OCH_2Ph$, $NHC(O)OBz$, $-NHC(O)-R_{10}$, $COOH$, $-C(O)Ph$, $C(O)O-R_{10}$, $C(O)H$, $C(O)-R_{10}$,

C₁-C₅ linear or branched C(O)-haloalkyl, -C(O)NH₂, C(O)NHR, C(O)N(R₁₀)(R₁₁), SO₂R, SO₂N(R₁₀)(R₁₁), C₁-C₅ linear or branched, substituted or unsubstituted alkyl, C₁-C₅ linear or branched or C₃-C₈ cyclic haloalkyl, C₁-C₅ linear or branched or C₃-C₈ cyclic alkoxy, C₁-C₅ linear or branched thioalkoxy, C₁-C₅ linear or branched haloalkoxy, C₁-C₅ linear or branched alkoxyalkyl, substituted or unsubstituted C₃-C₈ cycloalkyl, substituted or unsubstituted C₃-C₈ heterocyclic ring, or substituted or unsubstituted aryl; each is a separate embodiment according to this invention. In some embodiment, **R₆₀** is H. In some embodiment, **R₆₀** is COOH. In some embodiment, **R₆₀** is absent (e.g., when X₄ is N). In some embodiments, **R₆₀** may be further substituted with at least one substitution selected from: F, Cl, Br, I, OH, SH, C₁-C₅ linear or branched alkyl, C₂-C₅ linear or branched alkenyl, C₂-C₅ linear or branched alkynyl (e.g. CCH), C₃-C₈ cycloalkyl, linear, branched or cyclic alkoxy, COOH, COO(R), NH₂, N(R)₂, CF₃, aryl, phenyl, heteroaryl, C₃-C₈ cycloalkyl, CN and NO₂; each represents a separate embodiment according to this invention.

[00126] In some embodiment, **R₇₀** of compound of formula **I(a), I(b), I(d), I(e), I(f)** and/or **I(h)** is H, F, Cl, Br, I, OH, SH, R₈-OH, R₈-SH, -R₈-O-R₁₀, CF₃, CN, NO₂, NH₂, NHR, N(R)₂, R₈-N(R₁₀)(R₁₁), -OC(O)CF₃, -OCH₂Ph, NHC(O)OBz, -NHC(O)-R₁₀, COOH, -C(O)Ph, C(O)O-R₁₀, C(O)H, C(O)-R₁₀, C₁-C₅ linear or branched C(O)-haloalkyl, -C(O)NH₂, C(O)NHR, C(O)N(R₁₀)(R₁₁), SO₂R, SO₂N(R₁₀)(R₁₁), C₁-C₅ linear or branched or C₃-C₈ cyclic haloalkyl, C₁-C₅ linear or branched or C₃-C₈ cyclic alkoxy, C₁-C₅ linear or branched thioalkoxy, C₁-C₅ linear or branched haloalkoxy, C₁-C₅ linear or branched alkoxyalkyl, substituted or unsubstituted C₃-C₈ cycloalkyl, substituted or unsubstituted C₃-C₈ heterocyclic ring, or substituted or unsubstituted aryl; each is a separate embodiment according to this invention. In some embodiments, **R₇₀** is not an alkyl. In some embodiments, **R₇₀** may be further substituted with at least one substitution selected from: F, Cl, Br, I, OH, SH, C₁-C₅ linear or branched alkyl, C₂-C₅ linear or branched alkenyl, C₂-C₅ linear or branched alkynyl (e.g. CCH), C₃-C₈ cycloalkyl, linear, branched or cyclic alkoxy, COOH, COO(R), NH₂, N(R)₂, CF₃, aryl, phenyl, heteroaryl, C₃-C₈ cycloalkyl, CN and NO₂; each represents a separate embodiment according to this invention.

[00127] In some embodiment, **R₈₀** of compound of formula **I(a), I(b), I(d), I(e), I(f)** and/or **I(h)** is H, F, Cl, Br, I, OH, SH, R₈-OH, R₈-SH, -R₈-O-R₁₀, CF₃, CN, NO₂, NH₂, NHR, N(R)₂, R₈-N(R₁₀)(R₁₁), -OC(O)CF₃, -OCH₂Ph, NHC(O)OBz, -NHC(O)-R₁₀, COOH, -C(O)Ph, C(O)O-R₁₀, C(O)H, C(O)-R₁₀, C₁-C₅ linear or branched C(O)-haloalkyl, -C(O)NH₂, C(O)NHR, C(O)N(R₁₀)(R₁₁), SO₂R, SO₂N(R₁₀)(R₁₁), C₁-C₅ linear or branched, substituted or unsubstituted alkyl, C₁-C₅ linear or branched or C₃-C₈ cyclic haloalkyl, C₁-C₅ linear or branched or C₃-C₈ cyclic alkoxy, C₁-C₅ linear or branched thioalkoxy, C₁-C₅ linear or branched haloalkoxy, C₁-C₅ linear or branched alkoxyalkyl, substituted or unsubstituted C₃-C₈ cycloalkyl, substituted or unsubstituted C₃-C₈ heterocyclic ring, or substituted or unsubstituted aryl; each is a separate embodiment according to this invention. In some embodiment, **R₈₀** is H. In some embodiment, **R₈₀** is absent (e.g., when X₆ is N). In some embodiments, **R₈₀** may be further substituted with at least one substitution selected from: F, Cl, Br, I, OH, SH, C₁-C₅ linear or branched alkyl, C₂-C₅ linear or branched alkenyl, C₂-C₅ linear or branched alkynyl (e.g. CCH), C₃-C₈ cycloalkyl, linear, branched or cyclic alkoxy, COOH, COO(R), NH₂, N(R)₂, CF₃, aryl, phenyl, heteroaryl, C₃-C₈ cycloalkyl, CN and NO₂; each represents a separate embodiment according to this invention.

[00128] In some embodiment, R_{90} of compound of formula **I(a)**, **I(b)**, **I(d)**, **I(e)**, **I(f)** and/or **I(h)** is H, F, Cl, Br, I, OH, SH, R_8 -OH, R_8 -SH, $-R_8-O-R_{10}$, CF_3 , CN, NO_2 , NH_2 , NHR, $N(R)_2$, $R_8-N(R_{10})(R_{11})$, $-OC(O)CF_3$, $-OCH_2Ph$, $NHC(O)OBz$, $-NHC(O)-R_{10}$, $COOH$, $-C(O)Ph$, $C(O)O-R_{10}$, $C(O)H$, $C(O)-R_{10}$, C_1-C_5 linear or branched $C(O)$ -haloalkyl, $-C(O)NH_2$, $C(O)NHR$, $C(O)N(R_{10})(R_{11})$, SO_2R , $SO_2N(R_{10})(R_{11})$, C_1-C_5 linear or branched, substituted or unsubstituted alkyl, C_1-C_5 linear or branched or C_3-C_8 cyclic haloalkyl, C_1-C_5 linear or branched or C_3-C_8 cyclic alkoxy, C_1-C_5 linear or branched thioalkoxy, C_1-C_5 linear or branched haloalkoxy, C_1-C_5 linear or branched alkoxyalkyl, substituted or unsubstituted C_3-C_8 cycloalkyl, substituted or unsubstituted C_3-C_8 heterocyclic ring, or substituted or unsubstituted aryl; each is a separate embodiment according to this invention. In some embodiment, R_{90} is H. In some embodiment, R_{90} is absent (e.g., when X_7 is N). In some embodiments, R_{90} may be further substituted with at least one substitution selected from: F, Cl, Br, I, OH, SH, C_1-C_5 linear or branched alkyl, C_2-C_5 linear or branched alkenyl, C_2-C_5 linear or branched alkynyl (e.g. CCH), C_3-C_8 cycloalkyl, linear, branched or cyclic alkoxy, $COOH$, $COO(R)$, NH_2 , $N(R)_2$, CF_3 , aryl, phenyl, heteroaryl, C_3-C_8 cycloalkyl, CN and NO_2 ; each represents a separate embodiment according to this invention.

[00129] In some embodiments, if ring **B** of formula **I-I(b)**, **I(e)**, **I(f)** or **I(h)** exists, and R_4 is NH_2 , then R_3 cannot be OH. In some embodiments, if ring **B** of formula **I-I(b)**, **I(e)**, **I(f)** or **I(h)** exists, R_3 is OH and R_4 is NH_2 , n cannot be 1. In some embodiments, if R_3 is OH and R_4 is NH_2 , X_1 or X_7-R_{90} of compound of formula **I-I(b)**, **I(e)**, **I(f)** or **I(h)** cannot be CH.

[00130] In some embodiments, R_8 of formula **I**, **I(a)-I(h)** and/or **X-X(d)** is CH_2 . In other embodiments, R_8 is CH_2CH_2 . In other embodiments, R_8 is $CH_2CH_2CH_2$.

[00131] In some embodiments, p of formula **I**, **I(a)-I(h)** and/or **X-X(d)** is 1. In other embodiments, p is 2. In other embodiments, p is 3.

[00132] In some embodiments, R_9 of formula **I**, **I(a)-I(h)** and/or **X-X(d)** is $C\equiv C$.

[00133] In some embodiments, q of formula **I**, **I(a)-I(h)** and/or **X-X(d)** is 2.

[00134] In some embodiments, R_{10} of formula **I**, **I(a)-I(h)** and/or **X-X(d)** is H. In other embodiments, R_{10} is H, CN, C_1-C_5 linear or branched alkyl, methyl, ethyl, $C(O)R$, $C(O)(OCH_3)$ or $S(O)_2R$; each represents a separate embodiment according to this invention. In other embodiments, R_{10} is C_1-C_5 linear or branched alkyl. In other embodiments, R_{10} is CH_3 . In other embodiments, R_{10} is CH_2CH_3 . In other embodiments, R_{10} is $CH_2CH_2CH_3$. In other embodiments, R_{10} is CN. In other embodiments, R_{10} is $C(O)R$. In other embodiments, R_{10} is $S(O)_2R$. In other embodiments, R_{10} is $C(O)(OCH_3)$.

[00135] In some embodiments, R_{11} of formula **I**, **I(a)-I(h)** and/or **X-X(d)** is C_1-C_5 linear or branched alkyl. In other embodiments, R_{11} is H. In other embodiments, R_{11} is H, CN, C_1-C_5 linear or branched alkyl, methyl, ethyl, $C(O)R$, $C(O)(OCH_3)$ or $S(O)_2R$; each represents a separate embodiment according to this invention. In other embodiments, R_{11} is CH_3 . In other embodiments, R_{11} is CH_2CH_3 . In other embodiments, R_{11} is $CH_2CH_2CH_3$. In other embodiments, R_{11} is CN. In other embodiments, R_{11} is $C(O)R$. In other embodiments, R_{11} is $S(O)_2R$. In other embodiments, R_{11} is $C(O)(OCH_3)$.

[00136] In some embodiments, R_{10} and R_{11} of formula **I**, **I(a)-I(h)** and/or **X-X(d)** are joined to form a substituted or unsubstituted C_3-C_8 heterocyclic ring. In other embodiments, R_{10} and R_{11} are joined to form a piperazine ring. In other embodiments, R_{10} and R_{11} are joined to form a piperidine ring. In some

embodiments, the rings may be further substituted with at least one substitution selected from: F, Cl, Br, I, OH, SH, C₁-C₅ linear or branched alkyl, C₂-C₅ linear or branched alkenyl, C₂-C₅ linear or branched alkynyl (e.g. CCH), C₃-C₈ cycloalkyl, linear, branched or cyclic alkoxy, COOH, COO(R), NH₂, N(R)₂, CF₃, aryl, phenyl, heteroaryl, C₃-C₈ cycloalkyl, CN and NO₂; each represents a separate embodiment according to this invention.

[00137] In some embodiments, **R** of formula **I**, **I(a)**-**I(h)** and/or **X-X(d)** is H. In other embodiments, **R** is C₁-C₅ linear or branched alkyl, C₁-C₅ linear or branched alkoxy, phenyl, aryl or heteroaryl. In other embodiments, **R** is not H. In other embodiments, **R** is C₁-C₅ linear or branched alkyl. In other embodiments, **R** is methyl. In other embodiments, **R** is ethyl. In other embodiments, **R** is C₁-C₅ linear or branched alkoxy. In other embodiments, **R** is methoxy. In other embodiments, **R** is phenyl. In other embodiments, **R** is aryl. In other embodiments, **R** is heteroaryl. In other embodiments, two gem **R** substituents are joined together to form a 5 or 6 membered heterocyclic ring.

[00138] In some embodiments, **m** of formula **I**, **I(a)**-**I(d)** and **I(f)**-**I(h)** is 1. In some embodiments, **m** is 2. In some embodiments, **m** is 3, 4 or 5; each is a separate embodiment according to this invention.

[00139] In some embodiments, **m** of formula **X(a)**-**X(d)** is 1. In some embodiments, **m** is 2. In some embodiments, **m** is 3. In some embodiments, **m** is 1 or 2. In some embodiments, **m** is 1 or 3.

[00140] In some embodiments, **n** of formula **I** and/or **I(a)**-**I(h)** is 0. In other embodiments, **n** is 1. In other embodiments, **n** is 2. In other embodiments, **n** is 3. In some embodiments, **n** is 4. In some embodiments, **n** is 5.

[00141] In some embodiments, **n** of formula **X** and/or **X(a)**-**X(d)** is 1. In some embodiments, **n** is 2. In some embodiments, **n** is 0. In some embodiments, **n** is 3. In some embodiments, **n** is 1 or 2. In some embodiments, **n** is 0 or 1. In some embodiments, **n** is between 0 and 2.

[00142] In some embodiments, **o** of formula **X(a)** is 1. In some embodiments, **o** is 2. In some embodiments, **o** is 0. In some embodiments, **o** is 3. In some embodiments, **o** is 1 or 2. In some embodiments, **o** is 0 or 1. In some embodiments, **o** is between 0 and 2.

[00143] In some embodiments, **X₁** of formula **I**, **I(a)**-**I(d)** and **I(f)**-**I(h)** is S. In other embodiments, **X₁** is O. In other embodiments, **X₁** is CH₂. In some embodiments, **X₁** of formula **I**, **I(a)**, **I(c)**, **I(d)** and **I(f)**-**I(h)** is N-OH. In some embodiments, **X₁** is C(R)₂. In some embodiments, **X₁** is N-OMe.

[00144] In some embodiments, **X₁** of formula **X-X(c)** is S. In other embodiments, **X₁** is O. In other embodiments, **X₁** is CH₂. In some embodiments, **X₁** is CH(R). In some embodiments, **X₁** is C(R)₂. In some embodiments, **X₁** is S, O, or CH₂.


[00145] In some embodiments, **X₂** of formula **I**, **I(a)**-**I(d)** and **I(g)** is S. In other embodiments, **X₂** is O. In other embodiments, **X₂** is CH₂. In some embodiments, **X₂** of formula **I**, **I(a)**, **I(c)**, **I(d)** and **I(f)**-**I(h)** is N-OH. In some embodiments, **X₂** is C(R)₂. In some embodiments, **X₂** is N-OMe.

[00146] In some embodiments, **X₂** of formula **I** is O, and **R₃** or **R₄** are NH₂.

[00147] In some embodiments, **X₂** of formula **X**, **X(b)**, **X(c)** and/or **X(d)** is S. In other embodiments, **X₂** is O. In other embodiments, **X₂** is CH₂. In some embodiments, **X₂** is CH(R). In some embodiments, **X₂** is C(R)₂. In some embodiments, **X₂** is S, O, or CH₂.

[00148] In some embodiments, **X₁** and **X₂** of compound of formula **X-X(d)** are both CH₂.

[00149] In some embodiments, X_2 together with the carbon next to X_1 in compound of formula **I**, **I(a)**, **I(b)**, **I(f)** and/or **I(h)** are joined to form ring **B**, represented by the following structure (in such case, X_1 is X_7):

**B**

wherein X_4 , X_5 , X_6 and X_7 are each independently C or N, wherein if any of X_4 , X_5 , X_6 and X_7 is N, then the respective substitution R_{90} , R_{60} , R_{70} or R_{80} is absent, and wherein R_{90} , R_{60} , R_{70} or R_{80} are as defined above.

[00150] In some embodiments, X_3 of formula **I** and/or **I(a)**-**I(h)** is O. In other embodiments, X_3 is NH.

10 In other embodiments, X_3 is N- R_{50} . In other embodiments, X_3 is S.

[00151] In some embodiments, R_{50} of formula **I** and/or **I(a)**-**I(h)** is H. In some embodiments, R_{50} is C₁-C₅ linear or branched, substituted or unsubstituted alkyl. In some embodiments, R_{50} is methyl. In some embodiments, R_{50} is ethyl. In some embodiments, R_{50} is butyl. In some embodiments, R_{50} is i-propyl. In some embodiments, R_{50} be further substituted with at least one substitution selected from: F, Cl, Br, I, OH, SH, C₁-C₅ linear or branched alkyl, C₂-C₅ linear or branched alkenyl, C₂-C₅ linear or branched alkynyl (e.g. CCH), C₃-C₈ cycloalkyl, linear, branched or cyclic alkoxy, COOH, COO(R), NH₂, N(R)₂, CF₃, aryl, phenyl, heteroaryl, C₃-C₈ cycloalkyl, CN and NO₂; each represents a separate embodiment according to this invention.

20 [00152] In some embodiments, X_4 of formula **I**, **I(a)**-**I(d)** and **I(f)**-**I(h)** is C. In other embodiments, X_4 is N.

[00153] In some embodiments, X_5 of formula **I**, **I(a)**-**I(d)** and **I(f)**-**I(h)** is C. In other embodiments, X_5 is N.

[00154] In some embodiments, X_6 of formula **I**, **I(a)**-**I(d)** and **I(f)**-**I(h)** is C. In other embodiments, X_6 is N.

25 [00155] In some embodiments, X_7 of formula **I**, **I(a)**-**I(d)** and **I(f)**-**I(h)** is C. In other embodiments, X_7 is N.

[00156] In some embodiments, if any of X_4 , X_5 , X_6 and X_7 is N, then the respective substitution R_{90} , R_{60} , R_{70} or R_{80} is absent.

30 [00157] In some embodiments, ring **A** of compound of formula **X** is a C₅-C₇ cycloalkyl. In some embodiments, ring **A** is cyclohexyl. In some embodiments, ring **A** is cyclopropyl. In some embodiments, ring **A** is cyclobutyl. In some embodiments, ring **A** is absent.

[00158] In some embodiments, ring **B** of compound of formula **X** is a 5-7 membered nitrogen-containing heterocyclic ring. In some embodiments, ring **B** is pyrrolidine. In some embodiments, ring **B** is piperidine. In some embodiments, ring **B** is absent.

[00159] In some embodiments, ring **C** of compound of formula **X** is a C₅-C₇ substituted or unsubstituted cycloalkyl. In some embodiments, ring **C** is an unsubstituted cycloalkyl. In some embodiments, ring **C** is a substituted cycloalkyl. In some embodiments, ring **C** is a cyclopentyl. In some embodiments, ring **C** is a cyclohexyl. In some embodiments, ring **C** is aromatic ring. In some
5 embodiments, ring **C** is a phenyl. In some embodiments, ring **C** is absent.

[00160] In some embodiments, ring **D** of compound of formula **X** is a C₅-C₇ cycloalkyl. In some embodiments, ring **D** is cyclopentyl. In some embodiments, ring **D** is absent.

[00161] In some embodiments, ring **E** of compound of formula **X** is a substituted or unsubstituted 5-7 membered nitrogen-containing heterocyclic ring. In some embodiments, ring **E** is a substituted 5-7
10 membered nitrogen-containing heterocyclic ring. In some embodiments, ring **E** is an unsubstituted 5-7 membered nitrogen-containing heterocyclic ring. In some embodiments, ring **E** is pyrrolidine. In some embodiments, ring **E** is piperidine. In some embodiments, ring **E** is oxazolidin-2-one. In some embodiments, **E** is substituted oxazolidin-2-one. In some embodiments, ring **E** is absent.

[00162] In some embodiments, at least one of rings **A** – **E** of compound of formula (**X**) is not absent. In
15 some embodiments, only one of rings **A** – **E** is not absent. In some embodiments, all rings **A** – **E** of compound of formula (**X**) are absent.

[00163] In various embodiments, this invention is directed to any one of the compounds presented in Table 1 as described herein above, agrochemical compositions and/or method of use thereof in
controlling the growth of undesired plants.

[00164] In various embodiments, this invention is directed to the use of any one of the compounds
20 presented in Table 2 as described herein above, and/or agrochemical compositions thereof, in controlling the growth of undesired plants.

[00165] It is well understood that in structures presented in this invention wherein the carbon atom has less than 4 bonds, H atoms are present to complete the valence of the carbon. It is well understood that
25 in structures presented in this invention wherein the nitrogen atom has less than 3 bonds, H atoms are present to complete the valence of the nitrogen.

[00166] In some embodiments, this invention is directed to the compounds listed hereinabove, agrochemical compositions and/or method of use thereof, wherein the compound is agrochemically
30 acceptable salt, stereoisomer, optical isomer, tautomer, hydrate, *N*-oxide, reverse amide analog, isotopic variant (deuterated analog), or any combination thereof. In some embodiments, the compounds are herbicides. In some embodiments, the compounds control the growth of undesired plants.

[00167] As used herein, “single or fused aromatic or heteroaromatic ring systems” can be any such ring, including but not limited to phenyl, naphthyl, pyridinyl, (2-, 3-, and 4-pyridinyl), quinolinyl,
35 pyrimidinyl, pyridazinyl, pyrazinyl, triazinyl, tetrazinyl, thiazolyl, isothiazolyl, oxazolyl, isoxazolyl, imidazolyl, 1-methylimidazole, pyrazolyl, pyrrolyl, furanyl, thiophene-yl, quinolinyl, isoquinolinyl, 2,3-dihydroindenyl, indenyl, tetrahydronaphthyl, 3,4-dihydro-2H-benzo[b][1,4]dioxepine, benzodioxolyl, benzo[d][1,3]dioxole, tetrahydronaphthyl, indolyl, 1H-indole, isoindolyl, anthracenyl, benzimidazolyl, 2,3-dihydro-1H-benzo[d]imidazolyl, indazolyl, 2H-indazole, triazolyl, 4,5,6,7-

tetrahydro-2H-indazole, 3H-indol-3-one, purinyl, benzoxazolyl, 1,3-benzoxazolyl, benzisoxazolyl, benzothiazolyl, 1,3-benzothiazole, 4,5,6,7-tetrahydro-1,3-benzothiazole, quinazoliny, quinoxaliny, 1,2,3,4-tetrahydroquinoxaline, 1-(pyridin-1(2H)-yl)ethanone, cinnoliny, phthalaziny, quinoliny, isoquinoliny, acridiny, benzofurany, 1-benzofuran, isobenzofurany, benzofuran-2(3H)-one, 5 benzothiophenyl, benzoxadiazole, benzo[c][1,2,5]oxadiazolyl, benzo[c]thiophenyl, benzodioxolyl, thiadiazolyl, [1,3]oxazolo[4,5-b]pyridine, oxadiazolyl, imidazo[2,1-b][1,3]thiazole, 4H,5H,6H-cyclopenta[d][1,3]thiazole, 5H,6H,7H,8H-imidazo[1,2-a]pyridine, 7-oxo-6H,7H-[1,3]thiazolo[4,5-d]pyrimidine, [1,3]thiazolo[5,4-b]pyridine, 2H,3H-imidazo[2,1-b][1,3]thiazole, thieno[3,2-d]pyrimidin-4(3H)-one, 4-oxo-4H-thieno[3,2-d][1,3]thiazin, imidazo[1,2-a]pyridine, 1H-imidazo[4,5-b]pyridine, 10 b]pyridine, 1H-imidazo[4,5-c]pyridine, 3H-imidazo[4,5-c]pyridine, pyrazolo[1,5-a]pyridine, imidazo[1,2-a]pyrazine, imidazo[1,2-a]pyrimidine, 1H-pyrrolo[2,3-b]pyridine, pyrido[2,3-b]pyrazine, pyrido[2,3-b]pyrazin-3(4H)-one, 4H-thieno[3,2-b]pyrrole, quinoxalin-2(1H)-one, 1H-pyrrolo[3,2-b]pyridine, 7H-pyrrolo[2,3-d]pyrimidine, oxazolo[5,4-b]pyridine, thiazolo[5,4-b]pyridine, thieno[3,2-c]pyridine, 3-methyl-4H-1,2,4-triazole, 5-methyl-1,2,4-oxadiazole, methyloxazolidin-2-one, etc; each 15 represents a separate embodiment according to this invention.

[00168] As used herein, the term “alkyl” can be any linear- or branched-chain alkyl group containing up to about 30 carbons unless otherwise specified. In various embodiments, an alkyl includes C₁-C₅ carbons. In some embodiments, an alkyl includes C₁-C₆ carbons. In some embodiments, an alkyl includes C₁-C₈ carbons. In some embodiments, an alkyl includes C₂-C₅ carbons. In some embodiments, 20 an alkyl includes C₂-C₈ carbons. In some embodiments, an alkyl includes C₁-C₁₀ carbons. In some embodiments, an alkyl is a C₁-C₁₂ carbons. In some embodiments, an alkyl is a C₁-C₂₀ carbons. In some embodiments, branched alkyl is an alkyl substituted by alkyl side chains of 1 to 5 carbons. In various embodiments, the alkyl group may be unsubstituted. In some embodiments, the alkyl group may be substituted by a halogen, haloalkyl, hydroxyl, alkoxy, carbonyl, amido, alkylamido, dialkylamido, 25 cyano, nitro, CO₂H, amino, alkylamino, dialkylamino, carboxyl, thio, thioalkyl, C₁-C₅ linear or branched haloalkoxy, CF₃, phenyl, halophenyl, (benzyloxy)phenyl, -CH₂CN, NH₂, NH-alkyl, N(alkyl)₂, -OC(O)CF₃, -OCH₂Ph, -NHC(O)-alkyl, -C(O)Ph, C(O)O-alkyl, C(O)H, -C(O)NH₂ or any combination thereof.

[00169] The alkyl group can be a sole substituent, or it can be a component of a larger substituent, such 30 as in an alkoxy, alkoxyalkyl, haloalkyl, arylalkyl, alkylamino, dialkylamino, alkylamido, alkylurea, etc. Preferred alkyl groups are methyl, ethyl, and propyl, and thus halomethyl, dihalomethyl, trihalomethyl, haloethyl, dihaloethyl, trihaloethyl, halopropyl, dihalopropyl, trihalopropyl, methoxy, ethoxy, propoxy, arylmethyl, arylethyl, arylpropyl, methylamino, ethylamino, propylamino, dimethylamino, diethylamino, methylamido, acetamido, propylamido, halomethylamido, haloethylamido, 35 halopropylamido, methyl-urea, ethyl-urea, propyl-urea, 2, 3, or 4-CH₂-C₆H₄-Cl, C(OH)(CH₃)(Ph), etc.

[00170] As used herein, the term “alkenyl” can be any linear- or branched-chain alkenyl group containing up to about 30 carbons as defined hereinabove for the term “alkyl” and at least one carbon-carbon double bond. Accordingly, the term alkenyl as defined herein includes also alkadienes, alkatrienes, alkatetraenes, and so on. In some embodiments, the alkenyl group contains one carbon-

carbon double bond. In some embodiments, the alkenyl group contains two, three, four, five, six, seven or eight carbon-carbon double bonds; each represents a separate embodiment according to this invention. Non limiting examples of alkenyl groups include: Ethenyl, Propenyl, Butenyl (i.e., 1-Butenyl, *trans*-2-Butenyl, *cis*-2-Butenyl, and Isobutylenyl), Pentene (i.e., 1-Pentenyl, *cis*-2-Pentenyl, and *trans*-2-Pentenyl), Hexene (e.g., 1-Hexenyl, (*E*)-2-Hexenyl, (*Z*)-2-Hexenyl, (*E*)-3-Hexenyl, (*Z*)-3-Hexenyl, 2-Methyl-1-Pentene, etc.), which may all be substituted as defined herein above for the term “alkyl”.

[00171] As used herein, the term “alkynyl” can be any linear- or branched-chain alkynyl group containing up to about 30 carbons as defined hereinabove for the term “alkyl” and at least one carbon-carbon triple bond. Accordingly, the term alkynyl as defined herein includes also alkadiynes, alkatriynes, alkateaynes, and so on. In some embodiments, the alkynyl group contains one carbon-carbon triple bond. In some embodiments, the alkynyl group contains two, three, four, five, six, seven or eight carbon-carbon triple bonds; each represents a separate embodiment according to this invention. Non limiting examples of alkynyl groups include: acetylenyl, Propynyl, Butynyl (i.e., 1-Butynyl, 2-Butynyl, and Isobutylynyl), Pentyne (i.e., 1-Pentynyl, 2-Pentynyl), Hexyne (e.g., 1-Hexynyl, 2-Hexynyl, 3-Hexynyl, etc.), which may all be substituted as defined herein above for the term “alkyl”.

[00172] As used herein, the term “aryl” refers to any aromatic ring that is directly bound to another group and can be either substituted or unsubstituted. The aryl group can be a sole substituent, or the aryl group can be a component of a larger substituent, such as in an arylalkyl, arylamino, arylamido, etc. Exemplary aryl groups include, without limitation, phenyl, tolyl, xylyl, furanyl, naphthyl, pyridinyl, pyrimidinyl, pyridazinyl, pyrazinyl, triazinyl, thiazolyl, oxazolyl, isooxazolyl, pyrazolyl, imidazolyl, thiophene-yl, pyrrolyl, indolyl, phenylmethyl, phenylethyl, phenylamino, phenylamido, 3-methyl-4H-1,2,4-triazolyl, 5-methyl-1,2,4-oxadiazolyl, etc. Substitutions include but are not limited to: F, Cl, Br, I, C₁-C₅ linear or branched alkyl, C₁-C₅ linear or branched haloalkyl, C₁-C₅ linear or branched alkoxy, C₁-C₅ linear or branched haloalkoxy, CF₃, phenyl, halophenyl, (benzyloxy)phenyl, CN, NO₂, -CH₂CN, NH₂, NH-alkyl, N(alkyl)₂, hydroxyl, -OC(O)CF₃, -OCH₂Ph, -NHC(O)-alkyl, COOH, -C(O)Ph, C(O)O-alkyl, C(O)H, -C(O)NH₂ or any combination thereof.

[00173] As used herein, the term “alkoxy” refers to an ether group substituted by an alkyl group as defined above. Alkoxy refers both to linear and to branched alkoxy groups. Nonlimiting examples of alkoxy groups are methoxy, ethoxy, propoxy, *iso*-propoxy, *tert*-butoxy.

[00174] A “haloalkyl” group refers, in some embodiments, to an alkyl group as defined above, which is substituted by one or more halogen atoms, e.g. by F, Cl, Br or I. The term “haloalkyl” include but is not limited to fluoroalkyl, i.e., to an alkyl group bearing at least one fluorine atom. Nonlimiting examples of haloalkyl groups are CF₃, CF₂CF₃, CF₂CH₃, CH₂CF₃, CF₂CH₂CH₃, CH₂CH₂CF₃, CF₂CH(CH₃)₂ and CF(CH₃)-CH(CH₃)₂.

[00175] A “halophenyl” group refers, in some embodiments, to a phenyl substituent which is substituted by one or more halogen atoms, e.g. by F, Cl, Br or I. In one embodiment, the halophenyl is 4-chlorophenyl.

[00176] An "alkoxyalkyl" group refers, in some embodiments, to an alkyl group as defined above, which is substituted by alkoxy group as defined above, e.g. by methoxy, ethoxy, propoxy, i-propoxy, t-butoxy etc. Nonlimiting examples of alkoxyalkyl groups are $-\text{CH}_2\text{-O-CH}_3$, $-\text{CH}_2\text{-O-CH}(\text{CH}_3)_2$, $-\text{CH}_2\text{-O-C}(\text{CH}_3)_3$, $-\text{CH}_2\text{-CH}_2\text{-O-CH}_3$, $-\text{CH}_2\text{-CH}_2\text{-O-CH}(\text{CH}_3)_2$, $-\text{CH}_2\text{-CH}_2\text{-O-C}(\text{CH}_3)_3$.

5 [00177] A "cycloalkyl" or "carbocyclic" group refers, in various embodiments, to a ring structure comprising carbon atoms as ring atoms, which may be either saturated or unsaturated, substituted or unsubstituted, single or fused. In some embodiments the cycloalkyl is a 3-10 membered ring. In some embodiments the cycloalkyl is a 3-12 membered ring. In some embodiments the cycloalkyl is a 6 membered ring. In some embodiments the cycloalkyl is a 5-7 membered ring. In some embodiments the cycloalkyl is a 3-8 membered ring. In some embodiments, the cycloalkyl group may be unsubstituted or substituted by a
10 halogen, alkyl, haloalkyl, hydroxyl, alkoxy, carbonyl, amido, alkylamido, dialkylamido, cyano, nitro, CO_2H , amino, alkylamino, dialkylamino, carboxyl, thio, thioalkyl, $\text{C}_1\text{-C}_5$ linear or branched haloalkoxy, CF_3 , phenyl, halophenyl, (benzyloxy)phenyl, $-\text{CH}_2\text{CN}$, NH_2 , NH-alkyl , N(alkyl)_2 , $-\text{OC(O)CF}_3$, $-\text{OCH}_2\text{Ph}$, $-\text{NHC(O)-alkyl}$, $-\text{C(O)Ph}$, C(O)O-alkyl , C(O)H , $-\text{C(O)NH}_2$ or any combination thereof. In some
15 embodiments, the cycloalkyl ring may be fused to another saturated or unsaturated cycloalkyl or heterocyclic 3-8 membered ring. In some embodiments, the cycloalkyl ring is a saturated ring. In some embodiments, the cycloalkyl ring is an unsaturated ring. Non limiting examples of a cycloalkyl group comprise cyclohexyl, cyclohexenyl, cyclopropyl, cyclopropenyl, cyclopentyl, cyclopentenyl, cyclobutyl, cyclobutenyl, cyclooctyl, cyclooctadienyl (COD), cyclooctane (COE) etc.

20 [00178] A "heterocycle" or "heterocyclic" group refers, in various embodiments, to a ring structure comprising in addition to carbon atoms, sulfur, oxygen, nitrogen or any combination thereof, as part of the ring. A "heteroaromatic ring" refers in various embodiments, to an aromatic ring structure comprising in addition to carbon atoms, sulfur, oxygen, nitrogen, selenium or any combination thereof, as part of the ring. In some embodiments the heterocycle or heteroaromatic ring is a 3-10 membered ring. In some
25 embodiments the heterocycle or heteroaromatic ring is a 3-12 membered ring. In some embodiments the heterocycle or heteroaromatic ring is a 6 membered ring. In some embodiments the heterocycle or heteroaromatic ring is a 5-7 membered ring. In some embodiments the heterocycle or heteroaromatic ring is a 3-8 membered ring. In some embodiments, the heterocycle group or heteroaromatic ring may be unsubstituted or substituted by a halogen, alkyl, haloalkyl, hydroxyl, alkoxy, carbonyl, amido, alkylamido,
30 dialkylamido, cyano, nitro, CO_2H , amino, alkylamino, dialkylamino, carboxyl, thiol, thioalkyl, $\text{C}_1\text{-C}_5$ linear or branched haloalkoxy, CF_3 , phenyl, halophenyl, (benzyloxy)phenyl, $-\text{CH}_2\text{CN}$, NH_2 , NH-alkyl , N(alkyl)_2 , $-\text{OC(O)CF}_3$, $-\text{OCH}_2\text{Ph}$, $-\text{NHC(O)-alkyl}$, $-\text{C(O)Ph}$, C(O)O-alkyl , C(O)H , $-\text{C(O)NH}_2$ or any combination thereof. In some embodiments, the heterocycle ring or heteroaromatic ring may be fused to another saturated or unsaturated cycloalkyl or heterocyclic 3-8 membered ring. In some embodiments, the heterocyclic ring
35 is a saturated ring. In some embodiments, the heterocyclic ring is an unsaturated ring. Non limiting examples of a heterocyclic ring or heteroaromatic ring systems comprise pyridine, piperidine, morpholine, piperazine, thiophene, pyrrole, benzodioxole, benzofuran-2(3H)-one, benzo[d][1,3]dioxole, indole, oxazole, isoxazole, imidazole and 1-methylimidazole, furane, triazole, pyrimidine, pyrazine, oxacyclobutane (1 or 2-oxacyclobutane), naphthalene, tetrahydrothiophene 1,1-dioxide, thiazole, benzimidazole, piperidine, 1-

methylpiperidine, isoquinoline, 1,3-dihydroisobenzofuran, benzofuran, 3-methyl-4H-1,2,4-triazole, 5-methyl-1,2,4-oxadiazole, oxazolidin-2-one, methyloxazolidin-2-one or indole; each is a separate embodiment according to this invention.

5 [00179] In various embodiments, this invention provides a compound of this invention or its agrochemically acceptable salt, stereoisomer, optical isomer, tautomer, hydrate, *N*-oxide, reverse amide analog, isotopic variants (e.g., deuterated analog), or any combination thereof. In various embodiments, this invention provides a single stereoisomer of the compound of this invention. In some embodiments, this invention provides an optical isomer of the compound of this invention. In some embodiments, this invention provides an agrochemically acceptable salt of the compound of this invention. In some
10 embodiments, this invention provides a tautomer of the compound of this invention. In some embodiments, this invention provides a hydrate of the compound of this invention. In some embodiments, this invention provides an *N*-oxide of the compound of this invention. In some embodiments, this invention provides a reverse amide analog of the compound of this invention. In some embodiments, this invention provides an isotopic variant (including but not limited to deuterated analog) of the compound of this invention. In some
15 embodiments, this invention provides a polymorph of the compound of this invention. In some embodiments, this invention provides a crystal of the compound of this invention. In some embodiments, this invention provides an agrochemical composition comprising a compound of this invention, as described herein, or, in some embodiments, any combination of a stereoisomer, optical isomer, agrochemically acceptable salt, tautomer, hydrate, *N*-oxide, isotopic variant (deuterated analog), polymorph, or crystal of
20 the compound of this invention.

[00180] In various embodiments, the term "isomer" includes, but is not limited to, stereoisomers including optical isomers and analogs, structural isomers and analogs, conformational isomers and analogs, and the like. In some embodiments, the isomer is a stereoisomer. In another embodiment, the isomer is an
25 optical isomer.

[00181] In various embodiments, this invention encompasses the use of various stereoisomers of the compounds of the invention. It will be appreciated by those skilled in the art that the compounds of the present invention may contain at least one chiral center. Accordingly, the compounds used in the methods of the present invention may exist in, and be isolated in, optically-active or racemic forms. The
30 compounds according to this invention may further exist as stereoisomers which may be also optically-active isomers (e.g., enantiomers such as (*R*) or (*S*)), as enantiomerically enriched mixtures, racemic mixtures, or as single diastereomers, diastereomeric mixtures, or any other stereoisomers, including but not limited to: (*R*)(*R*), (*R*)(*S*), (*S*)(*S*), (*S*)(*R*), (*R*)(*R*)(*R*), (*R*)(*R*)(*S*), (*R*)(*S*)(*R*), (*S*)(*R*)(*R*), (*R*)(*S*)(*S*), (*S*)(*R*)(*S*), (*S*)(*S*)(*R*) or (*S*)(*S*)(*S*) stereoisomers. Some compounds may also exhibit polymorphism. It is
35 to be understood that the present invention encompasses any racemic, optically-active, polymorphic, or

stereoisomeric form, or mixtures thereof, which form possesses properties useful in controlling the growth of various undesired plants as described herein.

[00182] It is well known in the art how to prepare optically-active forms (for example, by resolution of the racemic form by recrystallization techniques, by synthesis from optically-active starting materials, by chiral synthesis, or by chromatographic separation using a chiral stationary phase).

[00183] The compounds of the present invention can also be present in the form of a racemic mixture, containing substantially equivalent amounts of stereoisomers. In some embodiments, the compounds of the present invention can be prepared or otherwise isolated, using known procedures, to obtain a stereoisomer substantially free of its corresponding stereoisomer (i.e., substantially pure). By substantially pure, it is intended that a stereoisomer is at least about 95% pure, more preferably at least about 98% pure, most preferably at least about 99% pure.

[00184] Compounds of the present invention can also be in the form of a hydrate, which means that the compound further includes a stoichiometric or non-stoichiometric amount of water bound by non-covalent intermolecular forces.


[00185] As used herein, when some chemical functional group (e.g. alkyl or aryl) is said to be "substituted", it is herein defined that one or more substitutions are possible.

[00186] Compounds of the present invention may exist in the form of one or more of the possible tautomers and depending on the conditions it may be possible to separate some or all of the tautomers into individual and distinct entities. It is to be understood that all of the possible tautomers, including all additional enol and keto tautomers and/or isomers are hereby covered. For example, the following tautomers, but not limited to these, are included:

Tautomerization of the imidazole ring



Tautomerization of the pyrazolone ring:



[00187] The invention includes "agrochemically acceptable salts" of the compounds of this invention, which may be produced, by reaction of a compound of this invention with an acid or base. Certain compounds, particularly those possessing acidic or basic groups, can also be in the form of a salt, preferably an agrochemically acceptable salt. The term "agrochemically acceptable salt" refers to those salts that retain the agrochemical effectiveness and properties of the free bases or free acids, which are not agrochemically or otherwise undesirable. The salts are formed with inorganic acids such as hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid and the like, and organic acids such as acetic acid, propionic acid, glycolic acid, pyruvic acid, oxalic acid, maleic acid, malonic acid, succinic acid, fumaric acid, tartaric acid, citric acid, benzoic acid, cinnamic acid, mandelic acid, methanesulfonic acid, ethanesulfonic acid, *p*-toluenesulfonic acid, salicylic acid, *N*-acetylcysteine and

the like. Other salts are known to those of skill in the art and can readily be adapted for use in accordance with the present invention.

[00188] Suitable agrochemically-acceptable salts of amines of the compounds of this invention may be prepared from an inorganic acid or from an organic acid. In various embodiments, examples of inorganic salts of amines are bisulfates, borates, bromides, chlorides, hemisulfates, hydrobromates, hydrochlorates, 2-hydroxyethylsulfonates (hydroxyethanesulfonates), iodates, iodides, isothionates, nitrates, persulfates, phosphates, sulfates, sulfamates, sulfanilates, sulfonic acids (alkylsulfonates, arylsulfonates, halogen substituted alkylsulfonates, halogen substituted arylsulfonates), sulfonates and thiocyanates.

[00189] In various embodiments, examples of organic salts of amines may be selected from aliphatic, cycloaliphatic, aromatic, araliphatic, heterocyclic, carboxylic and sulfonic classes of organic acids, examples of which are acetates, aspartates, ascorbates, adipates, anthranilates, alginates, alkane carboxylates, substituted alkane carboxylates, alginates, benzenesulfonates, benzoates, bisulfates, butyrates, bicarbonates, bitartrates, citrates, camphorates, camphorsulfonates, cyclohexylsulfamates, cyclopentanepropionates, calcium edetates, camsylates, carbonates, clavulanates, cinnamates, dicarboxylates, digluconates, dodecylsulfonates, dihydrochlorides, decanoates, enanthuates, ethanesulfonates, edetates, edisylates, estolates, esylates, fumarates, formates, fluorides, galacturonates gluconates, glutamates, glycolates, gluconate, glucoheptanoates, glycerophosphates, gluceptates, glycolylarsanilates, glutarates, glutamate, heptanoates, hexanoates, hydroxymaleates, hydroxycarboxylic acids, hexylresorcinates, hydroxybenzoates, hydroxynaphthoates, hydrofluorates, lactates, lactobionates, laurates, malates, maleates, methylenebis(beta-oxynaphthoate), malonates, mandelates, mesylates, methane sulfonates, methylbromides, methylnitrates, methylsulfonates, monopotassium maleates, mucates, monocarboxylates, naphthalenesulfonates, 2-naphthalenesulfonates, nicotines, nitrates, napsylates, *N*-methylglucamines, oxalates, octanoates, oleates, pamoates, phenylacetates, picrates, phenylbenzoates, pivalates, propionates, phthalates, phenylacetate, pectinates, phenylpropionates, palmitates, pantothenates, polygalacturates, pyruvates, quinate, salicylates, succinates, stearates, sulfanilate, subacetates, tartrates, theophyllineacetates, *p*-toluenesulfonates (tosylates), trifluoroacetates, terephthalates, tannates, teoclates, trihaloacetates, triethiodide, tricarboxylates, undecanoates and valerates.

[00190] In various embodiments, examples of inorganic salts of carboxylic acids or hydroxyls may be selected from ammonium, alkali metals to include lithium, sodium, potassium, cesium; alkaline earth metals to include calcium, magnesium, aluminum; zinc, barium, cholines, quaternary ammoniums.

[00191] In some embodiments, examples of organic salts of carboxylic acids or hydroxyl may be selected from arginine, organic amines to include aliphatic organic amines, alicyclic organic amines, aromatic organic amines, benzathines, *t*-butylamines, benethamines (*N*-benzylphenethylamine), dicyclohexylamines, dimethylamines, diethanolamines, ethanolamines, ethylenediamines, hydrabamines, imidazoles, lysines, methylamines, meglamines, *N*-methyl-*D*-glucamines, *N,N'*-dibenzylethylenediamines, nicotinamides, organic amines, ornithines, pyridines, picolines, piperazines,

procain, tris(hydroxymethyl)methylamines, triethylamines, triethanolamines, trimethylamines, tromethamines and ureas.

[00192] In various embodiments, the salts may be formed by conventional means, such as by reacting the free base or free acid form of the product with one or more equivalents of the appropriate acid or base in a solvent or medium in which the salt is insoluble or in a solvent such as water, which is removed in vacuo or by freeze drying or by exchanging the ions of an existing salt for another ion or suitable ion-exchange resin.

Agrochemical composition

[00193] Another aspect of the present invention relates to an agrochemical composition including an agrochemically acceptable carrier or diluent and a compound according to the aspects of the present invention. The agrochemical composition can contain one or more of the above-identified compounds of the present invention. Typically, the agrochemical composition of the present invention will include a compound of the present invention or its agrochemically acceptable salt, as well as an agrochemically acceptable carrier or diluent. The term "agrochemically acceptable carrier" refers to any suitable adjuvants, carriers, excipients, or stabilizers, and can be in solid or liquid form such as sprays, aerosols, powders, solutions, suspensions, or emulsions.

[00194] The compounds according to the invention can be used as herbicidal agents in unmodified form, but they are generally formulated into compositions in various ways using formulation adjuvants, such as carriers, solvents and surface-active substances. The formulations can be in various physical forms, e.g. in the form of dusting powders, gels, wettable powders, water-dispersible granules, water-dispersible tablets, effervescent pellets, emulsifiable concentrates, microemulsifiable concentrates, oil-in-water emulsions, oil-flowables, aqueous dispersions, oily dispersions, suspo-emulsions, capsule suspensions, emulsifiable granules, soluble liquids, water-soluble concentrates (with water or a water-miscible organic solvent as carrier), impregnated polymer films or in other forms known. Such formulations can either be used directly or diluted prior to use. The dilutions can be made, for example, with water, liquid fertilizers, micronutrients, biological organisms, oil or solvents.

[00195] Typically, the composition will contain from about 0.01 to 99 percent, preferably from about 20 to 75 percent of active compound(s), together with the adjuvants, carriers and/or excipients. While individual needs may vary, determination of optimal ranges of effective amounts of each component is within the skill of the art.

[00196] The formulations can be prepared e.g. by mixing the active ingredient with the formulation adjuvants in order to obtain compositions in the form of finely divided solids, granules, solutions, dispersions or emulsions. The active ingredients can also be formulated with other adjuvants, such as finely divided solids, mineral oils, oils of vegetable or animal origin, modified oils of vegetable or animal origin, organic solvents, water, surface-active substances or combinations thereof.

[00197] The active ingredients can also be contained in very fine microcapsules. Microcapsules contain the active ingredients in a porous carrier. This enables the active ingredients to be released into the environment in controlled amounts (e.g. slow-release). Microcapsules usually have a diameter of from 0.1

to 500 microns. They contain active ingredients in an amount of about from 25 to 95 % by weight of the capsule weight. The active ingredients can be in the form of a monolithic solid, in the form of fine particles in solid or liquid dispersion or in the form of a suitable solution. The encapsulating membranes can comprise, for example, natural or synthetic rubbers, cellulose, styrene/butadiene copolymers, polyacrylonitrile, polyacrylate, polyesters, polyamides, polyureas, polyurethane or chemically modified polymers and starch xanthates or other polymers that are known to the person skilled in the art. Alternatively, very fine microcapsules can be formed in which the active ingredient is contained in the form of finely divided particles in a solid matrix of base substance, but the microcapsules are not themselves encapsulated.

10 [00198] The formulation adjuvants that are suitable for the preparation of the compositions according to the invention are known per se. As liquid carriers there may be used: water, toluene, xylene, petroleum ether, vegetable oils, acetone, methyl ethyl ketone, cyclohexanone, acid anhydrides, acetonitrile, acetophenone, amyl acetate, 2-butanone, butylene carbonate, chlorobenzene, cyclohexane, cyclohexanol, alkyl esters of acetic acid, diacetone alcohol, 1,2-dichloropropane, diethanolamine, p-diethylbenzene, diethylene glycol, diethylene glycol abietate, diethylene glycol butyl ether, diethylene glycol ethyl ether, diethylene glycol methyl ether, dimethylformamide (DMF), dimethyl sulfoxide (DMSO), 1,4-dioxane, dipropylene glycol, dipropylene glycol methyl ether, dipropylene glycol dibenzoate, diproxitol, alkyldipyrrolidone, ethyl acetate, 2-ethylhexanol, ethylene carbonate, 1,1,1-trichloroethane, 2-heptanone, alpha-pinene, d-limonene, ethyl lactate, ethylene glycol, ethylene glycol butyl ether, ethylene glycol methyl ether, gamma-butyrolactone, glycerol, glycerol acetate, glycerol diacetate, glycerol triacetate, hexadecane, hexylene glycol, isoamyl acetate, isobornyl acetate, isooctane, isophorone, isopropylbenzene, isopropyl myristate, lactic acid, laurylamine, mesityl oxide, methoxypropanol, methyl isoamyl ketone, methyl isobutyl ketone, methyl laurate, methyl octanoate, methyl oleate, methylene chloride, m-xylene, n-hexane, n-octylamine, octadecanoic acid, octylamine acetate, oleic acid, oleylamine, o-xylene, phenol, polyethylene glycol, propionic acid, propyl lactate, propylene carbonate, propylene glycol, propylene glycol methyl ether, p-xylene, toluene, triethyl phosphate, triethylene glycol, xylenesulfonic acid, paraffin, mineral oil, trichloroethylene, perchloroethylene, ethyl acetate, amyl acetate, butyl acetate, propylene glycol methyl ether, diethylene glycol methyl ether, methanol, ethanol, isopropanol, and alcohols of higher molecular weight, such as amyl alcohol, tetrahydrofurfuryl alcohol, hexanol, octanol, ethylene glycol, propylene glycol, glycerol, methyl-2-pyrrolidone and the like.

[00199] Suitable solid carriers are, for example, talc, titanium dioxide, pyrophyllite clay, silica, attapulgite clay, kieselguhr, limestone, calcium carbonate, bentonite, calcium montmorillonite, cottonseed husks, wheat flour, soybean flour, pumice, wood flour, ground walnut shells, lignin and similar substances.

35 [00200] A large number of surface-active substances can advantageously be used in both solid and liquid formulations, especially in those formulations which can be diluted with a carrier prior to use. Surface-active substances may be anionic, cationic, non-ionic or polymeric and they can be used as emulsifiers, wetting agents or suspending agents or for other purposes. Typical surface-active substances include, for example, salts of alkyl sulfates, such as diethanolammonium lauryl sulfate; salts of alkylarylsulfonates, such as calcium dodecylbenzenesulfonate; alkylphenol/alkylene oxide addition products, such as nonylphenol

ethoxylate; alcohol/alkylene oxide addition products, such as tridecylalcohol ethoxylate; soaps, such as sodium stearate; salts of alkylnaphthalenesulfonates, such as sodium dibutylnaphthalenesulfonate; dialkyl esters of sulfosuccinate salts, such as sodium di(2-ethylhexyl)sulfosuccinate; sorbitol esters, such as sorbitol oleate; quaternary amines, such as lauryltrimethylammonium chloride, polyethylene glycol esters of fatty acids, such as polyethylene glycol stearate; block copolymers of ethylene oxide and propylene oxide; and salts of mono- and di-alkylphosphate esters; as well as further substances known to the skilled in the arts.

[00201] Further adjuvants that can be used in herbicidal formulations include crystallization inhibitors, viscosity modifiers, suspending agents, dyes, anti-oxidants, foaming agents, light absorbers, mixing auxiliaries, antifoams, complexing agents, neutralizing or pH-modifying substances and buffers, corrosion inhibitors, fragrances, wetting agents, take-up enhancers, micronutrients, plasticizers, glidants, lubricants, dispersants, thickeners, antifreezes, microbicides, and liquid and solid fertilizers.

[00202] The compositions according to the invention can include an additive comprising an oil of vegetable or animal origin, a mineral oil, alkyl esters of such oils or mixtures of such oils and oil derivatives. The amount of oil additive in the composition according to the invention is generally from 0.01 to 10 %, based on the mixture to be applied. For example, the oil additive can be added to a spray tank in the desired concentration after a spray mixture has been prepared. Preferred oil additives comprise mineral oils or an oil of vegetable origin, for example rapeseed oil, olive oil or sunflower oil, emulsified vegetable oil, alkyl esters of oils of vegetable origin, for example the methyl derivatives, or an oil of animal origin, such as fish oil or beef tallow. Preferred oil additives comprise alkyl esters of C₈-C₂₂ fatty acids, especially the methyl derivatives of C₁₂-C₁₈ fatty acids, for example the methyl esters of lauric acid, palmitic acid and oleic acid (methyl laurate, methyl palmitate and methyl oleate, respectively). Other oil derivatives are known to the skilled in the arts, for examples from the Compendium of Herbicide Adjuvants, 10th Edition, Southern Illinois University, 2010.

[00203] The herbicidal compositions generally comprise from 0.1 to 99 % by weight, especially from 0.1 to 95 % by weight, compounds according to this invention and from 1 to 99.9 % by weight of a formulation adjuvant which preferably includes from 0 to 25 % by weight of a surface-active substance. The inventive compositions generally comprise from 0.1 to 99 % by weight, especially from 0.1 to 95 % by weight, of compounds of the present invention and from 1 to 99.9 % by weight of a formulation adjuvant which preferably includes from 0 to 25 % by weight of a surface-active substance. Whereas commercial products may preferably be formulated as concentrates, the end user will normally employ dilute formulations.

[00204] The rates of application vary within wide limits and depend on the nature of the soil, the method of application, the crop plant, the pest to be controlled, the prevailing climatic conditions, and other factors governed by the method of application, the time of application and the target crop. As a general guideline, compounds may be applied at a rate of from 1 to 2000 l/ha, especially from 10 to 1000 l/ha. Preferred formulations can have the following compositions (weight %):

[00205] Emulsifiable concentrates:

- active ingredient: 1 to 95 %, preferably 60 to 90 %

- surface-active agent: 1 to 30 %, preferably 5 to 20 %
- liquid carrier: 1 to 80 %, preferably 1 to 35 %

[00206] Dusts:

- active ingredient: 0.1 to 10 %, preferably 0.1 to 5 %
- solid carrier: 90 to 99.9 %, preferably 99 to 99.9 %

[00207] Suspension concentrates:

- active ingredient: 5 to 75 %, preferably 10 to 50 %
- water: 24 to 94 %, preferably 30 to 88 %
- surface-active agent: 1 to 40 %, preferably 2 to 30 %

[00208] Wettable powders:

- active ingredient: 0.5 to 90 %, preferably 1 to 80 %
- surface-active agent 0.5 to 20 %, preferably 1 to 15 %
- solid carrier: 5 to 95 %, preferably 15 to 90 %

[00209] Granules:

- active ingredient: 0.1 to 30 %, preferably 0.1 to 15 %
- solid carrier: 70 to 99.5 %, preferably 85 to 97 %

[00210] When the compounds or agrochemical compositions of the present invention are administered to control the growth of undesired plants, the agrochemical composition can also contain, or can be administered in conjunction with, other agrochemical agents or treatment regimen presently known or hereafter developed for the growth control of various types of plants.

[00211] Accordingly, the composition of the present invention may further comprise at least one additional pesticide including but not limited to herbicide. For example, the compounds according to the invention can also be used in combination with other herbicides or plant growth regulators. In a preferred embodiment the additional pesticide is an herbicide and/or herbicide safener.

[00212] Examples of herbicides that can be used in combination with the compounds of the invention, include but are not limited to: acetochlor, acifluorfen (including acifluorfen-sodium), aclonifen, alachlor, alloxymid, ametryn, amicarbazone, amidosulfuron, aminocyclopyrachlor, aminopyralid, amitrole, asulam, atrazine, bensulfuron (including bensulfuron-methyl), bentazone, bicyclopyrone, bilanafos, bifenox, bispyribac-sodium, bixlozone, bromacil, bromoxynil, butachlor, butafenacil, cafenstrole, carfentrazone (including carfentrazone-ethyl); cloransulam (including cloransulam-methyl), chlorimuron (including chlorimuron-ethyl), chlorotoluron, cinosulfuron, chlorsulfuron, cinmethylin, clacyfos, clethodim, clodinafop (including clodinafop-propargyl), clomazone, clopyralid, cyclopyranil, cyclopyrimorate, cyclosulfamuron, cyhalofop (including cyhalofop-butyl), 2,4-D (including the choline salt and 2-ethylhexyl ester thereof), 2,4-DB, daimuron, desmedipham, dicamba (including the aluminum, aminopropyl, bis-aminopropylmethyl, choline, dichloroprop, diglycolamine, dimethylamine, dimethylammonium, potassium and sodium salts thereof), diclofop-methyl, diclosulam, diflufenican, difenzoquat, diflufenican, diflufenzopyr, dimethachlor, dimethenamid-P, diquat dibromide, diuron, esprocarb, ethalfluralin, ethofumesate, fenoxaprop (including fenoxaprop-P-ethyl), fenoxasulfone, fenquinotrione, fentrazamide,

flazasulfuron, florasulam, florypyrauxifen, fluzifop (including fluzifop-P-butyl), flucarbazone (including flucarbazone-sodium);, flufenacet, flumetralin, flumetsulam, flumioxazin, flupyrsulfuron (including flupyrsulfuron-methyl-sodium);, fluoxypyr (including fluoxypyr-meptyl);, fluthiacet-methyl, fomesafen, foramsulfuron, glufosinate (including the ammonium salt thereof), glyphosate (including the diammonium, isopropylammonium and potassium salts thereof), halauxifen (including halauxifen-methyl), halosulfuron-methyl, haloxyfop (including haloxyfop-methyl), hexazinone, hydantocidin, imazamox, imazapic, imazapyr, imazaquin, imazethapyr, indaziflam, iodosulfuron (including iodosulfuron-methyl-sodium), iofensulfuron, iofensulfuron-sodium, ioxynil, ipfencarbazone, isoproturon, isoxaben, isoxaflutole, lactofen, lancotrione, linuron, MCPA, MCPB, mecoprop-P, mefenacet, mesosulfuron, mesosulfuron-methyl, mesotrione, metamitron, metazachlor, methiozolin, metobromuron, metolachlor, metosulam, metoxuron, metribuzin, metsulfuron, molinate, napropamide, nicosulfuron, norflurazon, orthosulfamuron, oxadiargyl, oxadiazon, oxasulfuron, oxyfluorfen, paraquat dichloride, pendimethalin, penoxsulam, phenmedipham, picloram, picolinafen, pinoxaden, pretilachlor, primisulfuron-methyl, prodiamine, prometryn, propachlor, propanil, propaquizafop, propham, propyrisulfuron, propyzamide, prosulfocarb, prosulfuron, pyraclonil, pyraflufen (including pyraflufen-ethyl), pyrasulfotole, pyrazolynate, pyrazosulfuron-ethyl, pyribenzoxim, pyridate, pyriftalid, pyrimisulfan, pyrithiobac-sodium, pyroxasulfone, pyroxsulam, quinclorac, quinmerac, quizalofop (including quizalofop-P-ethyl and quizalofop-P-tefuryl), rimsulfuron, saflufenacil, sethoxydim, simazine, S-metolachlor, sulcotrione, sulfentrazone, sulfosulfuron, tebuthiuron, tefuryltrione, tembotrione, terbuthylazine, terbutryn, thiencarbazone, thifensulfuron, tiafenacil, tolpyralate, topramezone, tralkoxydim, triafamone, triallate, triasulfuron, tribenuron (including tribenuron-methyl), triclopyr, trifloxysulfuron (including trifloxysulfuron-sodium), trifludimoxazin, trifluralin, triflusulfuron, tritosulfuron, 4-hydroxy-1-methoxy-5-methyl-3-[4-(trifluoromethyl)-2-pyridyl]imidazolidin-2-one, 4-hydroxy-1,5-dimethyl-3-[4-(trifluoromethyl)-2-pyridyl]imidazolidin-2-one, 5-ethoxy-4-hydroxy-1-methyl-3-[4-(trifluoromethyl)-2-pyridyl]imidazolidin-2-one, 4-hydroxy-1-methyl-3-[4-(trifluoromethyl)-2-pyridyl]imidazolidin-2-one, 4-hydroxy-1,5-dimethyl-3-[1-methyl-5-(trifluoromethyl)pyrazol-3-yl]imidazolidin-2-one, (4R)-1-(5-tert-butylisoxazol-3-yl)-4-ethoxy-5-hydroxy-3-methyl-imidazolidin-2-one, 3-[2-(3,4-dimethoxyphenyl)-6-methyl-3-oxo-pyridazine-4-carbonyl]bicyclo[3.2.1]octane-2,4-dione, 2-[2-(3,4-dimethoxyphenyl)-6-methyl-3-oxo-pyridazine-4-carbonyl]-5-methyl-cyclohexane-1,3-dione, 2-[2-(3,4-dimethoxyphenyl)-6-methyl-3-oxo-pyridazine-4-carbonyl]cyclohexane-1,3-dione, 2-[2-(3,4-dimethoxyphenyl)-6-methyl-3-oxo-pyridazine-4-carbonyl]-5,5-dimethyl-cyclohexane-1,3-dione, 6-[2-(3,4-dimethoxyphenyl)-6-methyl-3-oxo-pyridazine-4-carbonyl]-2,2,4,4-tetramethyl-cyclohexane-1,3,5-trione, 2-[2-(3,4-dimethoxyphenyl)-6-methyl-3-oxo-pyridazine-4-carbonyl]-5-ethyl-cyclohexane-1,3-dione, 2-[2-(3,4-dimethoxyphenyl)-6-methyl-3-oxo-pyridazine-4-carbonyl]-4,4,6,6-tetramethyl-cyclohexane-1,3-dione, 2-[6-cyclopropyl-2-(3,4-dimethoxyphenyl)-3-oxo-pyridazine-4-carbonyl]-5-methyl-cyclohexane-1,3-dione, 3-[6-cyclopropyl-2-(3,4-dimethoxyphenyl)-3-oxo-pyridazine-4-carbonyl]bicyclo[3.2.1]octane-2,4-dione, 2-[6-cyclopropyl-2-(3,4-dimethoxyphenyl)-3-oxo-pyridazine-4-carbonyl]-5,5-dimethyl-cyclohexane-1,3-dione, 6-[6-cyclopropyl-2-(3,4-dimethoxyphenyl)-3-oxo-pyridazine-4-carbonyl]-2,2,4,4-tetramethyl-cyclohexane-1,3,5-trione, 2-[6-cyclopropyl-2-(3,4-dimethoxyphenyl)-3-oxo-pyridazine-4-carbonyl]cyclohexane-1,3-dione, 4-[2-(3,4-dimethoxyphenyl)-6-methyl-3-oxo-pyridazine-4-carbonyl]-

2,2,6,6-tetramethyl-tetrahydropyran-3,5-dione and I + 4-[6-cyclopropyl-2-(3,4-dimethoxyphenyl)-3-oxo-pyridazine-4-carbonyl]-2,2,6,6-tetramethyl-tetrahydropyran-3,5-dione. These additional agents may also be present in the form of esters or salts thereof.

[00213] Compounds of the present invention may also be combined with herbicide safeners. Example of herbicide safeners include but are not limited to: benoxacor, cloquintocet (including cloquintocet-mexyl), cyprosulfamide, dichlormid, fenclorazole (including fenclorazole-ethyl), fenclorim, fluxofenim, furilazole, isoxadifen (including isoxadifen-ethyl), mefenpyr (including mefenpyr-diethyl), metcamifen, N-(2-methoxybenzoyl)-4-[(methylaminocarbonyl)amino] benzenesulfonamide and oxabetrinil; all of which may be in the form of esters or salts thereof.

[00214] The compound of the invention can also be used in mixtures with other agrochemicals such as fungicides, nematicides or insecticides, examples of which are known to the skilled in the art.

[00215] The mixing ratio of compound of the invention and the additional agent, is preferably from 1:100 to 1000:1. Preferably the mixing ratio of compound of the invention to safener is from 100:1 to 1:10, especially from 20:1 to 1:1.

[00216] The mixtures can advantageously be used in the above-mentioned formulations (in which case "active ingredient" relates to the respective mixture of compound of the invention with the additional agent.

Herbicidal Activity

[00217] In various embodiments, the invention provides compounds and compositions, including any embodiment described herein, for use in any of the methods of this invention. In various embodiments, use of a compound of this invention or a composition comprising the same, will have utility in inhibiting, suppressing, enhancing, or stimulating a desired response, as will be understood by one skilled in the art. In some embodiments, the compositions may further comprise additional active ingredients, whose activity is useful for the particular application for which the compound of this invention is being administered.

[00218] The compounds of this invention are useful as herbicides or herbicidal compounds. The present invention therefore further comprises a method for controlling the growth of undesired plants, comprising applying to the plants or a locus comprising them, an effective amount of a compound according to this invention, or an agrochemical composition thereof, under conditions effective to control the growth of the undesired plants, in particular the growth of weeds, in crops of useful plants.

[00219] In some embodiments, "Controlling" according to this invention refers to killing, reducing or retarding growth or preventing or reducing germination. Generally, the plants to be controlled are unwanted plants (weeds).

[00220] In some embodiments, "Locus" refers to the area in which the plants are growing or will grow.

[00221] The rates of application of compounds of the invention may vary within wide limits and depend on the nature of the soil, the method of application (for example: pre-plant, pre-emergence; post-emergence; application to the seed furrow; no tillage application etc.), the crop plant, the weed(s) to be controlled, the prevailing climatic conditions, and other factors governed by the method of application, the time of

application and the target crop. The compounds of the invention are generally applied at a rate of from 10 to 2000 g/ha, especially from 50 to 1000 g/ha.

[00222] In some embodiments, the application is made by spraying the composition, typically by tractor mounted sprayer for large areas, but other methods such as dusting (for powders), drip or drench can also be used.

[00223] In some embodiments, useful plants in which the composition according to the invention can be used upon include crops such as cereals including but not limited to barley and wheat, cotton, oilseed rape, sunflower, maize, rice, soybeans, sugar beet, sugar cane and turf.

[00224] In some embodiments, crop plants also include trees, such as fruit trees, palm trees, coconut trees or other nuts. Also included are vines such as grapes, fruit bushes, fruit plants and vegetables.

[00225] In some embodiments, the crops are resistant crops. Therefore, according to some embodiments, crops also include those crops which have been rendered tolerant to herbicides or classes of herbicides (including but not limited to: ALS-, GS-, EPSPS-, PPO-, ACCase- and HPPD-inhibitors) by conventional methods of breeding or by genetic engineering. Examples of crops that have been rendered tolerant to herbicides by genetic engineering methods include but not limited to glyphosate- and glufosinate-resistant maize varieties commercially available under the trade names RoundupReady® and LibertyLink®. According to other embodiments, crops also include those which have been rendered resistant to harmful insects by genetic engineering methods, examples of such crops include but not limited to: Bt maize (resistant to European corn borer), Bt cotton (resistant to cotton boll weevil) and Bt potatoes (resistant to Colorado beetle). Non limiting examples of Bt maize include the Bt 176 maize hybrids of NK® (Syngenta Seeds). Non limiting examples of transgenic plants comprising one or more genes that code for an insecticidal resistance and express one or more toxins are: KnockOut® (maize), Yield Gard® (maize), NuCOTIN33B® (cotton), Bollgard® (cotton), NewLeaf® (potatoes), NatureGard® and Protexcta®. Plant crops or seed material thereof can be both resistant to herbicides and, at the same time, resistant to insect feeding ("stacked" transgenic events).

[00226] In some embodiments, crops include those which are obtained by conventional methods of breeding or genetic engineering and contain so-called output traits (e.g. improved storage stability, higher nutritional value and improved flavour). Other useful plants include turf grass for example in golf-courses, lawns, parks and roadsides, or grown commercially for sod, and ornamental plants such as flowers or bushes.

[00227] Herbicidal compounds, or chemically active herbicides, may be broken down into pre-plant herbicides, pre-emergent herbicides and post-emergent herbicides. Pre-plant and pre-emergent herbicides typically interfere with germination of weed seeds and are applied before and after planting or sowing, respectively, but before seed germination, whereas post-emergent herbicides kill the weeds after the weed seeds have germinated and weed growth has begun.

[00228] When administering the compounds of the present invention, they can be administered in pre-plant or pre-emergence treatments, in post-emergence treatments, or both.

[00229] In various embodiments, this invention is directed to a method of controlling the growth of undesired plants, comprising applying a compound according to this invention, or an agrochemical

composition thereof, to crop fields. In some embodiments, the compound is a pre-plant herbicide. In some embodiments, the compound is a pre-emergent herbicide. In some embodiments, the compound is a post-emergent herbicide. Therefore, in some embodiment, the compound is applied to crop fields before the undesired plants emerge (i.e. pre-emergent or pre-plant herbicide). In some embodiments, the compound is applied to crop fields after the undesired plants emerge (i.e. post-emergent herbicide).

[00230] In various embodiments, compounds according to this invention, and agrochemical compositions thereof, are used to control undesired plants, which include a wide variety of monocotyledonous and dicotyledonous weed species.

[00231] In some embodiments, the undesired plant is a weed. In some embodiments, the undesired plant is a eudicot (dicotyledonous or dicot). In some embodiments, the undesired plant is a monocotyledon (monocotyledonous or monocot).

[00232] Non limiting examples of monocotyledonous species that can typically be controlled include *Alopecurus myosuroides*, *Avena fatua*, *Brachiaria plantaginea*, *Bromus tectorum*, *Cyperus esculentus*, *Digitaria sanguinalis*, *Echinochloa crus-galli*, *Lolium perenne*, *Lolium multiflorum*, *Panicum miliaceum*, *Poa annua*, *Setaria viridis*, *Setaria faberi* and *Sorghum bicolor*; each represents a separate embodiment according to this invention.

[00233] Non limiting examples of dicotyledonous species that can be controlled include *Abutilon theophrasti*, *Amaranthus retroflexus*, *Bidens pilosa*, *Chenopodium album*, *Euphorbia heterophylla*, *Galium aparine*, *Ipomoea hederacea*, *Kochia scoparia*, *Polygonum convolvulus*, *Sida spinosa*, *Sinapis arvensis*, *Solanum nigrum*, *Stellaria media*, *Veronica persica* and *Xanthium strumarium*; each represents a separate embodiment according to this invention.

[00234] In some embodiments, the undesired plant is *Abutilon theophrasti*, *Amaranthus palmeri*, *Ambrosia artemisiifolia*, *Alopecurus myosuroides*, *Avena sterilis*, *Chenopodium album*, *Conyza Canadensis*, *Digitaria sanguinalis*, *Echinochloa colona*, *Euphorbia heterophylla*, *Lolium perenne*, *Lolium rigidum*, *Matricaria chamomilla*, *Phalaris paradoxa*, *Poa annua*, *Portulaca oleracea*, *Setaria viridis*, *Solanum nigrum* or any combination thereof. In some embodiments, the compound is any one of the compounds listed in Table 1 and 2; each compound represents a separate embodiment according to this invention.

[00235] In some embodiments, compounds, and compositions according to this invention are utilized to control undesirable vegetation in rice. In certain embodiments, the undesirable vegetation is *Brachiaria platyphylla* (Groseb.) Nash (broadleaf signalgrass, BRAPP), *Digitaria sanguinalis* (L.) Scop, (large crabgrass, DIGSA), *Echinochloa crus-galli* (L.) P. Beauv. (barnyardgrass, ECHCG), *Echinochloa colonum* (L.) LINK (junglerice, ECHCO), *Echinochloa oryzoides* (Ard.) Fritsch (early watergrass, ECHOR), *Echinochloa oryzicola* (Vasinger) Vasinger (late watergrass, ECHPH), *Ischaemum rugosum* Salisb. (saramollagrass, ISCRU), *Leptochloa chinensis* (L.) Nees (Chinese sprangletop, LEFCH), *Leptochloa fascicularis* (Lam.) Gray (bearded sprangletop, LEFFA), *Leptochloa panicoides* (Presl.) Hitchc. (Amazon sprangletop, LEFPA), *Panicum dichotomiflorum* (L.) Michx. (fall panicum, PANDI), *Paspalum dilatatum* Poir. (dallisgrass, PASDI), *Cyperus difformis* L. (smallflower flatsedge, CYPDI), *Cyperus esculentus* L. (yellow nutsedge, CYPES), *Cyperus iria* L. (rice flatsedge, CYPPI), *Cyperus*

rotundus L. (purple nutsedge, CYPRO), *Eleocharis* species (ELOSS), *Fimbristylis miliacea* (L.) Vahl (globe fringerush, FIMMI), *Schoenoplectus juncooides* Roxb. (Japanese bulrush, SCPJU), *Schoenoplectus maritimus* L. (sea clubrush, SCPMA), *Schoenoplectus mucronatus* L. (ricefield bulrush, SCPMU), *Aeschynomene* species, (jointvetch, AESSS), *Alternanthera philoxeroides* (Mart.) Griseb. (alligatorweed, ALRPH), *Alisma plantago-aquatica* L. (common waterplantain, ALSPA), *Amaranthus* species, (pigweeds and amaranths, AMASS), *Ammannia coccinea* Rottb. (redstem, AMMCO), *Eclipta alba* (L.) Hassk. (American false daisy, ECLAL), *Heteranthera limosa* (SW.) Willd./Vahl (ducksalad, HETLI), *Heteranthera reniformis* R. & P. (roundleaf mudplantain, HETRE), *Ipomoea hederacea* (L.) Jacq. (ivyleaf momingglory, IPOHE), *Lindernia dubia* (L.) Pennell (low false pimpernel, LIDDU),
 5 *Monochoria korsakowii* Regel & Maack (monochoria, MOOKA), *Monochoria vaginalis* (Burm. F.) C. Presl ex Kuhth, (monochoria, MOOVA), *Murdannia nudiflora* (L.) Brenan (doveweed, MUDNU), *Polygonum pennsylvanicum* L. (Pennsylvania smartweed, POLPY), *Polygonum persicaria* L. (ladysthumb, POLPE), *Polygonum hydropiperoides* Michx. (POLHP, mild smartweed), *Rotala indica* (Willd.) Koehne (Indian toothcup, ROTIN), *Sagittaria* species, (arrowhead, SAGSS), *Sesbania exaltata*
 10 (Raf) Cory/Rydb. Ex Hill (hemp sesbania, SEBEX), or *Sphenoclea zeylanica* Gaertn. (gooseweed, SPDZE); each is a separate embodiment according to this invention. In some embodiments, the compound is any one of the compounds listed in Table 1 and 2; each compound represents a separate embodiment according to this invention.

[00236] In some embodiments, the compounds and compositions according to this invention are utilized
 20 to control undesirable vegetation in cereals. In certain embodiments, the undesirable vegetation is *Alopecurus myosuroides* Huds. (blackgrass, ALOMY), *Apera spica-venti* (L.) Beauv. (windgrass, APESV), *Avena fatua* L. (wild oat, AVEFA), *Bromus tectorum* L. (downy brome, BROTE), *Lolium multiflorum* Lam. (Italian ryegrass, LOLMU), *Phalaris minor* Retz. (littleseed canarygrass, PHAMI), *Poa annua* L. (annual bluegrass, POAAN), *Setaria pumila* (Poir.) Roemer & J.A. Schultes (yellow
 25 foxtail, SETLU), *Setaria viridis* (L.) Beauv. (green foxtail, SETVI), *Cirsium arvense* (L.) Scop. (Canada thistle, CIRARy), *Galium aparine* L. (catchweed bedstraw, GALAP), *Kochia scoparia* (L.) Schrad. (kochia, KCHSC), *Lamium purpureum* L. (purple deadnettle, LAMP), *Matricaria recutita* L. (wild chamomile, MATCH), *Matricaria matricarioides* (Less.) Porter (pineappleweed, MATMT), *Papaver rhoeas* L. (common poppy, PAPRH), *Polygonum convolvulus* L. (wild buckwheat, POLCO), *Salsola tragus* L. (Russian thistle, SASKR), *Stellaria media* (L.) Vill (common chickweed, STEME), *Veronica persica* Poir. (Persian speedwell, VERPE), *Viola arvensis* Murr. (field violet, VIOAR), or *Viola tricolor*
 30 L. (wild violet, VIOTR); each is a separate embodiment according to this invention. In some embodiments, the compound is any one of the compounds listed in Table 1 and 2; each compound represents a separate embodiment according to this invention.

[00237] In some embodiments, the compounds and compositions according to this invention are utilized
 35 to control undesirable vegetation in range and pasture. In certain embodiments, the undesirable vegetation is *Ambrosia artemisiifolia* L. (common ragweed, AMBEL), *Cassia obtusifolia* (sickle pod, CASOB), *Centaurea maculosa* auct. non Lam. (spotted knapweed, CENMA), *Cirsium arvense* (L.) Scop. (Canada thistle, CIRAR), *Convolvulus arvensis* L. (field bindweed, CONAR), *Euphorbia esula*

L. (leafy spurge, EPHES), *Lactuca serriola* L./Tom. (prickly lettuce, LACSE), *Plantago lanceolata* L. (buckhorn plantain, PLALA), *Rumex obtusifolius* L. (broadleaf dock, RUMOB), *Sida spinosa* L. (prickly sida, SIDSP), *Sinapis arvensis* L. (wild mustard, SINAR), *Sonchus arvensis* L. (perennial sowthistle, SONAR), *Solidago* species (goldenrod, SOOSS), *Taraxacum officinale* G.H. Weber ex Wiggers (dandelion, TAROF), *Trifolium repens* L. (white clover, TRFRE), or *Urtica dioica* L. (common nettle, URTDI); each is a separate embodiment according to this invention. In some embodiments, the compound is any one of the compounds listed in Table 1 and 2; each compound represents a separate embodiment according to this invention.


[00238] In some embodiments, the compounds and compositions according to this invention are utilized to control undesirable vegetation found in row crops. In certain embodiments, the undesirable vegetation is *Alopecurus myosuroides* Huds. (blackgrass, ALOMY), *Avena fatua* L. (wild oat, AVEFA), *Brachiaria platyphylla* (Groseb.) Nash (broadleaf signalgrass, BRAPP), *Digitaria sanguinalis* (L.) Scop. (large crabgrass, DIGSA), *Echinochloa crus-galli* (L.) P. Beauv. (barnyardgrass, ECHCG), *Echinochloa colonum* (L.) Link (junglerice, ECHCO), *Lolium multiflorum* Lam. (Italian ryegrass, LOLMU), *Panicum dichotomiflorum* Michx. (fall panicum, PANDI), *Panicum miliaceum* L. (wild-proso millet, PANMI), *Setaria faberi* Herrm. (giant foxtail, SETFA), *Setaria viridis* (L.) Beauv. (green foxtail, SETVI), *Sorghum halepense* (L.) Pers. (Johnsongrass, SORHA), *Sorghum bicolor* (L.) Moench ssp. *Arundinaceum* (shattercane, SORVU), *Cyperus esculentus* L. (yellow nutsedge, CYPES), *Cyperus rotundus* L. (purple nutsedge, CYPRO), *Abutilon theophrasti* Medik. (velvetleaf, ABUTH), *Amaranthus* species (pigweeds and amaranths, AMASS), *Ambrosia artemisiifolia* L. (common ragweed, AMBEL), *Ambrosia psilostachya* DC. (western ragweed, AMBPS), *Ambrosia trifida* L. (giant ragweed, AMBTR), *Asclepias syriaca* L. (common milkweed, ASCSY), *Chenopodium album* L. (common lambsquarters, CHEAL), *Cirsium arvense* (L.) Scop. (Canada thistle, CIRAR), *Commelina benghalensis* L. (tropical spiderwort, COMBE), *Datura stramonium* L. (jimsonweed, DATST), *Daucus carota* L. (wild carrot, DAUCA), *Euphorbia heterophylla* L. (wild poinsettia, EPHHL), *Erigeron bonariensis* L. (hairy fleabane, ERIBO), *Erigeron canadensis* L. (Canadian fleabane, ERICA), *Helianthus annuus* L. (common sunflower, HELAN), *Jacquemontia tamnifolia* (L.) Griseb. (smallflower morningglory, IAQTA), *Ipomoea hederacea* (L.) Jacq. (ivyleaf morningglory, IPOHE), *Ipomoea lacunosa* L. (white morningglory, IPOLA), *Lactuca serriola* L./Tom. (prickly lettuce, LACSE), *Portulaca oleracea* L. (common purslane, POROL), *Sida spinosa* L. (prickly sida, SIDSP), *Sinapis arvensis* L. (wild mustard, SINAR), *Solanum ptychanthum* Dunal (eastern black nightshade, SOLPT), or *Xanthium strumarium* L. (common cocklebur, XANST); each is a separate embodiment according to this invention. In some embodiments, the compound is any one of the compounds listed in Table 1 and 2; each compound represents a separate embodiment according to this invention.

[00239] The following examples are presented in order to more fully illustrate the preferred embodiments of the invention. They should in no way, however, be construed as limiting the broad scope of the invention.

EXAMPLES

EXAMPLE 1

Synthetic Details for Compounds of the Invention



5

Scheme 1. Synthetic route and procedures for Compounds 104, 105, 106, 114 and 115.

[00240] Compounds 106, 115 and 114 were prepared using the same route, with the suitable enantiomer of threonine.

10 **Step M:**

[00241] To solution of compound 1 (40 g, 336 mmol) in methanol (400 mL) was added dropwise SOCl_2 (39.4 mL, 543 mmol) at 0°C . The reaction mixture was refluxed, concentrated under reduced pressure to obtain crude product which was used in the next step without further purification.

Step N:

15 [00242] To solution from the previous step were added MeCN (300 mL) and methanol (250 mL). Then to obtain mixture at 0°C were added dropwise Boc_2O (92.2 mL, 401 mmol) and triethylamine (105 mL, 753 mmol), and it was left stirring overnight. After that it was concentrated under reduced pressure, dissolved in dichloromethane (400 mL) and water (400 mL), extracted with dichloromethane (2×200 mL).

Combined organic layers was washed with 2N HCl (200 mL), K_2CO_3 , dried over Na_2SO_4 , and concentrated to obtained 74.6 g of compound **3** (320 mmol, 95% yield for 2 steps).

Step O:

5 [00243] Compound **3** (74.6 g, 320 mmol) was dissolved in MeCN (1200 mL) and dimethoxy propane (358 mL), and after cooling to 10-15°C was added $BF_3 \cdot Et_2O$ (2.39 mL, 19.4 mmol), the solution was left stirring overnight. Then triethylamine (16 mL, 115 mmol), concentrated under reduced pressure, dissolved with water (300 mL), extracted with dichloromethane (3×300 mL), dried over Na_2SO_4 , and concentrated to obtained crude product **4** which was used in the next step without further purification.

Step P:

10 [00244] To crude product **4** dissolved in THF (200 mL) was added dropwise solution $LiAlH_4$ (13 g, 343 mmol) in THF (700 mL) and it was stirred overnight. After that it was quench with solution of KOH (6.7 g, 119 mmol) in water (90 mL), filtered, the obtain precipitate was washed with THF, filtrate was concentrated under reduced pressure. The crude residue was purified by column chromatography to obtained 39 g of compound **5** (159 mmol, 50% yield for 2 steps).

Step Q:

15 [00245] To solution of compound **5** (24.05 g, 98.0 mmol) in dichloromethane (420 mL) was added in portions Dess-Martin periodinane (50.4 g, 119 mmol) and the reaction mixture was stirred overnight. The solution of $NaHCO_3$ (42.2 g, 502 mmol) and Na_2SO_4 (189.2 g, 1332 mmol) in water (700 mL) were added and then it had been stirred by the time the whole precipitate was dissolved. The reaction mixture was
20 separated, water layer was extracted with dichloromethane (2×200 mL), organic layer was dried over Na_2SO_4 , and concentrated to obtained 20.3 g of compound **6** (83.4 mmol, 85% yield).

Step R:

[00246] Ethyl 4-bromocrotonate (**7**) (20.00 g, 93.24 mmol) was added in one portion to triethyl phosphine (17.71 g, 106.56 mmol) at 120-130°C and the solution was stirred for 1 h. Distillation of the
25 resulting reaction mixture provided 20.39 g of compound **8** as a light yellow oil (81.5 mmol, 88% yield).

Step S:

[00247] K_2CO_3 (57.7 g, 418 mmol) and 18-crown-6 (36.9 g, 140 mmol) were dissolved in toluene and it was stirred for 1 h. To the solution was cooled to -20°C and then were added dropwise compound **8** (20.39 g, 81.5 mmol) and compound **6** (16.9 g, 69.5 mmol). Then the reaction mixture was heated to r.t.,
30 stirred, and monitored by TLC (for 1-2 days). After reaction was completed, water (300 mL) was added there, the solution was extracted with hexane (2×200 mL), combined organic layers were washed with water (2×150 mL), 5% citric acid (100 mL), dried over Na_2SO_4 , and concentrated under reduced pressure. The crude residue was purified by column chromatography to obtained 17 g of compound **9** (50.1 mmol, 72% yield).

Step T:

[00248] To solution of compound **9** (16.5 g, 48.6 mmol) in ethanol (200 mL) was added 10% Pd/C (1 g). The reaction flask was vacuumed and the solution was stirred overnight under H_2 atmosphere. Then it was filtered and concentrated under reduced pressure to obtain 16 g of compound **10** (46.6 mmol, 96% yield).

Step U:

[00249] Compound **10** was dissolved in methanol (150 mL) and p-toluenesulfonic acid (0.1 g, 0.581 mmol) was added. The reaction mixture was stirred overnight and then it was concentrated under reduced pressure. Then was added the solution of K₂CO₃ (1 g, 7.24 mmol) in water (30 mL), extracted with
5 MTBE (3×100 mL), dried over Na₂SO₄, and concentrated to obtained 14.1 g of mixture of esters **11** (the esterification took place under this reaction Me:Et=2:1) which was used in next step.

Step V1:

[00250] The mixture of esters **11** (2.69 g) was dissolved in solution of THF (80 mL), methanol (25 mL), and water (25 mL), then LiOH·H₂O (1.1 g, 26.2 mmol) was added at 0°C. The solution was heated to r.t.
10 and left stirring. After that THF and methanol were evaporated under reduced pressure, the resultant water solution was cooled, acidified 1.2N solution of NaHSO₄, extracted with MTBE (3×50 mL), dried over Na₂SO₄, concentrated in vacuum, and purified by reversed-phase chromatography. On Horner stage took place partial racemization (85:15), but on stage of reverse-phase chromatography the diastereomers was separated and obtained 0.6 g of compound **14** (2.18 mmol, 25% yield).

Step V2:

[00251] Compound **14** (0.6 g, 2.18 mmol) was dissolved in MTBE (4 mL) and Di/HCl (2.5 mL) was added and the resultant mixture was stirred overnight. The occurred precipitate was filtered, washed with MTBE and dried under reduced pressure to obtain 0.36 g of hydrochloride target compound **104**
20 (2.05 mmol, 94%).

m/z = 176.2

¹H NMR (400 MHz, dmsO) δ (ppm): 12.04 (bs, 1H, COOH), 7.89 (bs, 3H, NH₃), 5.25 (s, 1H, OH), 3.85 (m, 1H, CH), 2.97 (m, 1H, CH), 2.19 (t, J=7.12 Hz, 2H, CH₂-COOH), 1.48-1.30 (m, 6H, CH₂-CH₂-CH₂), 1.06 (d, J=6.36 Hz, CH₃).

Step W:

[00252] To solution of NaH (1.4 g, 35.0 mmol) in THF (300 mL) at 0°C was added dropwise the solution of compound **11** (7.05 g, 28.7 mmol) in THF (30 mL). The solution was heated to r.t. and stirred overnight. Then it was poured into mixture of ice water (200 mL) and NH₄Cl (6 g, 112 mmol), the layers were separated, water layer was extracted with MTBE (3×100 mL), dried over Na₂SO₄, and concentrated to obtained 4.5 g of crude product compound **12** which was used in the next step without further
30 purification.

Step X:

[00253] The mixture of esters **12** (1.5 g) was dissolved in solution of THF (40 mL), methanol (15 mL), and water (15 mL), then LiOH·H₂O (0.96 g, 22.9 mmol) was added at 0°C. The solution was heated to r.t. and left stirring. After that THF and methanol were evaporated under reduced pressure, the resultant water solution was cooled, acidified 1.2N solution of NaHSO₄, extracted with MTBE (3×50 mL), dried over Na₂SO₄, concentrated in vacuum to obtain 0.87 g of crude product compound **105** (62%). On Horner stage took place partial racemization (85:15).


Step Y:

[00254] To solution of diastereomeric acid (0.87 g, 4.96 mmol) from the previous step in DMF (10 mL) was added K_2CO_3 (1.19 g) and benzyl bromide (0.56 mL, 4.71 mmol), and the reaction mixture was stirred overnight. Then water (40 mL) was added there, extracted with MTBE (3×50 mL), organic layer was washed with water (5×30 mL), dried over Na_2SO_4 , concentrated under reduced pressure, and purified by reversed-phase chromatography to obtain 0.6 g (48%) crude product which was purified by chiral column chromatography to give 0.36 g of required diastereomer **13** (0.012 mmol, 60%).

Step Z:


[00255] To solution of compound **13** (0.36 g, 1.31 mmol) in methanol (5 mL) was added 10% Pd(C) (0.1 g). The flask was degassed, and reaction carried out under H_2 atmosphere and left stirring overnight. After that it was filtered, and filtrate concentrated under reduced pressure. The residue was purified by column chromatography to obtain 0.16 g of target compound **105** (0.795 mmol, 96%).

[00256] From compound (**14**) in above scheme, the following are obtained:



15 **Scheme 2.** Procedure adapted from Adibi, Hadi; *et al.*, *Synthetic Communications* (2010), 40(18), 2753-2766.


Preparation of Compounds 123 and 148



20 **Scheme 3.** Synthetic route and procedure for the preparation of Compound **123**.


[00257] Compound **148** was prepared using the same route, starting with tert-butyl 2-formylpyrrolidine-1-carboxylate (instead of tert-butyl 3-formylpyrrolidine-1-carboxylate), as described below.

Synthesis of Compound 123




[00258] To a solution of compound **1** (754 mg, 3.01 mmol, 1.2 *eq*) in THF (20 mL) NaH was added (130 mg, 3.26 mmol, 60% purity, 1.3 *eq*) at 0 ~ 5 °C and stirred at 25 °C for 0.5 hr. Then
 5 Compound **123_1** (500 mg, 2.51 mmol, 1 *eq*) was added and the mixture was stirred at 25 °C for 1.5 hrs. LCMS showed the reaction was completed. The residue was poured into saturated aqueous NH₄Cl solution (50 mL). The aqueous phase was extracted with ethyl acetate (50 mL*2). The combined organic phase was washed with brine (50 mL), dried with anhydrous MgSO₄, filtered and concentrated in vacuum. The residue was purified by flash silica gel chromatography
 10 (ISCO®; 20 g SepaFlash® Silica Flash Column, Eluent of 0~15% Ethyl acetate/Petroleum ether gradient @ 100 mL/min) to give Compound **123_2** (420 mg, 1.41 mmol, 56.0% yield, 98.9% purity) as a colorless oil.

[00259] ¹H NMR: (400 MHz, CDCl₃) δ 7.19 - 7.13 (m, 1H), 6.22 - 6.05 (m, 1H), 6.02 - 5.91 (m, 1H), 5.89 - 5.72 (m, 1H), 4.23 - 4.03 (m, 2H), 3.57 - 3.47 (m, 1H), 3.42 (ddd, *J* = 3.4, 8.0, 10.9 Hz, 1H), 3.34 - 3.18 (m, 1H), 3.08 - 2.92 (m, 1H), 2.90 - 2.75 (m, 1H), 2.05 - 1.93 (m, 1H), 1.75 - 1.59 (m, 1H), 1.39 (s, 9H), 1.26 - 1.20 (m, 3H)




[00260] To a solution of Compound **123_2** (420 mg, 1.42 mmol, 1 *eq*) in EtOH (20 mL) Pd/C (200 mg, 1.42 mmol, 10% purity, 1 *eq*) was added under N₂. The suspension was degassed under vacuum and purged with H₂ several times. The mixture was stirred under H₂ (40 psi) at 25 °C for
 20 2 hours. LCMS showed the reaction was completed. The reaction mixture was filtered and the filtrate was concentrated to give crude compound **123_3** (327 mg, 1.09 mmol, 76.8% yield) as a colorless oil, which used into the next step without further purification.

[00261] ¹H NMR: (400 MHz, CDCl₃) δ 4.12 - 3.99 (m, 2H), 3.54 - 3.27 (m, 2H), 3.24 - 3.07 (m, 1H), 2.85 - 2.67 (m, 1H), 2.23 (dt, *J* = 3.6, 7.2 Hz, 2H), 2.10 - 1.95 (m, 1H), 1.89 (ddt, *J* = 2.9, 6.6, 9.3 Hz, 1H), 1.62 - 1.50 (m, 3H), 1.39 (s, 9H), 1.35 - 1.22 (m, 4H), 1.22 - 1.16 (m, 3H)



[00262] To a solution of compound **123_3** (327 mg, 1.09 mmol) in EtOH (3.00 mL) and H₂O (3.00 mL) NaOH was added (87.4 mg, 2.18 mmol), the mixture was stirred at 25 °C for 2 hrs. LCMS showed the starting material was consumed. It was diluted with ethyl acetate (4 mL) and the mixture was adjusted pH to 4, the organic layer was dried over MgSO₄, filtered and concentrated. It was used as is in the next step without additional purification. Compound **123_4** (240 mg, 776 μmol, 71.0 % yield, 87.7 % purity) was obtained as a light yellow oil.


[00263] ¹H NMR: (400 MHz, DMSO-*d*₆) δ 11.99 (br s, 1H), 3.45 - 3.37 (m, 1H), 3.32 - 3.27 (m, 1H), 3.20 - 3.07 (m, 1H), 2.79 - 2.64 (m, 1H), 2.21 (t, *J* = 7.3 Hz, 2H), 2.13 - 1.98 (m, 1H), 1.91 (s, 1H), 1.56 - 1.43 (m, 2H), 1.39 (s, 9H), 1.36 - 1.21 (m, 4H)



[00264] To a solution of compound **123_4** (240 mg, 884 μmol) in dioxane (2.50 mL), HCl/dioxane was added (2.50 mL) and stirred at 25 °C for 2 hrs. LCMS showed the starting material was consumed. The solvent was removed under reduced pressure to give the product. Compound **123** (150 mg, 858 μmol, 97.1 % yield, 98.0 % purity) was obtained as a light yellow solid.

[00265] ¹H NMR: (400 MHz, DMSO-*d*₆) δ 12.05 (br d, *J* = 1.9 Hz, 1H), 9.45 (br d, *J* = 1.6 Hz, 1H), 8.79 (br s, 1H), 3.25 - 3.01 (m, 2H), 2.23 (t, *J* = 7.3 Hz, 2H), 2.12 - 2.01 (m, 1H), 1.98 - 1.42 (m, 8H), 1.41 - 1.26 (m, 2H)


20 Synthesis of Compound 148



[00266] To a solution of compound **1** (1.51 g, 6.02 mmol, 1.2 *eq*) in THF (20 mL) NaH was added (261 mg, 6.52 mmol, 60% purity, 1.3 *eq*) at 0 ~ 5 °C and stirred at 25 °C for 0.5 hr. Then compound **148_1** (1.00 g, 5.02 mmol, 1 *eq*) was added and stirred 25 °C for 1.5 hrs. LCMS showed the reaction was completed. The residue was poured into saturated aqueous NH₄Cl solution (50 mL). The aqueous phase was extracted with ethyl acetate (50 mL*2). The combined organic phase was washed with brine (50 mL), dried with anhydrous MgSO₄, filtered and concentrated in vacuum. The residue was purified by flash silica gel chromatography (ISCO®; 20 g SepaFlash® Silica Flash Column, Eluent of 0~15% Ethyl acetate/Petroleum ether gradient


@ 100 mL/min) to give compound **148_2** (640 mg, 2.15 mmol, 42.9% yield, 99.3% purity) as a colorless oil.

[00267] $^1\text{H NMR}$: (400 MHz, CDCl_3) δ 7.65 - 7.12 (m, 2H), 6.20 - 5.50 (m, 3H), 4.81 - 3.92 (m, 3H), 3.64 - 3.06 (m, 2H), 2.17 - 1.90 (m, 1H), 1.90 - 1.71 (m, 2H), 1.71 - 1.55 (m, 1H), 1.41 - 1.27 (m, 9H), 1.23 (dt, $J = 1.9, 7.1$ Hz, 3H)




[00268] To a solution of compound **148_2** (640 mg, 2.17 mmol, 1 *eq*) in EtOH (20 mL) Pd/C (200 mg, 2.17 mmol, 10% purity, 1 *eq*) was added under N_2 . The suspension was degassed under vacuum and purged with H_2 several times. The mixture was stirred under H_2 (50 psi) at 25 °C for 2 hours. LCMS showed the reaction was completed. The reaction mixture was filtered and the filtrate was concentrated to give crude compound **148_3** (540 mg, 1.80 mmol, 83.2% yield) as a colorless oil which used into the next step without further purification.

[00269] $^1\text{H NMR}$: (400 MHz, CDCl_3) δ 4.05 (q, $J = 7.1$ Hz, 2H), 3.65 (q, $J = 7.0$ Hz, 1H), 3.39 - 3.14 (m, 2H), 2.23 (t, $J = 7.5$ Hz, 2H), 1.90 - 1.62 (m, 4H), 1.62 - 1.48 (m, 4H), 1.41 - 1.35 (m, 9H), 1.28 - 1.14 (m, 6H).



[00270] To a solution of compound **148_3** (540 mg, 1.80 mmol) in EtOH (5.00 mL) and H_2O (5.00 mL) NaOH was added (144 mg, 3.61 mmol), the mixture was stirred at 25 °C for 2 hrs. LCMS showed the starting material was consumed. The solvent was removed under reduced pressure to give the crude product. It was used as is in the next step without additional purification. Compound **148_4** (450 mg, 1.49 mmol, 82.8 % yield, 90.0 % purity) was obtained as a white solid.

[00271] $^1\text{H NMR}$: (400 MHz, $\text{DMSO}-d_6$) δ 11.98 (br s, 2H), 3.71 - 3.54 (m, 1H), 3.28 - 3.10 (m, 2H), 2.27 - 2.13 (m, 2H), 1.90 - 1.67 (m, 3H), 1.66 - 1.43 (m, 4H), 1.42 - 1.35 (m, 9H), 1.34 - 1.12 (m, 4H)



[00272] To a solution of compound **148_4** (450 mg, 1.66 mmol) in dioxane (5.00 mL) HCl/dioxane (5.00 mL) was added and the solution was stirred at 25 °C for 2 hrs. LCMS showed the starting material was consumed. The solvent was removed under reduced pressure to give the

product. Compound **148** (200 mg, 1.14 mmol, 69.0 % yield, 98.0 % purity) was obtained as a yellow solid.

[00273] ¹H NMR: (400 MHz, DMSO-*d*₆) δ 12.18 - 11.95 (m, 1H), 9.45 (br d, *J* = 1.6 Hz, 1H), 8.79 (br s, 1H), 3.20 - 3.02 (m, 2H), 2.23 (t, *J* = 7.3 Hz, 2H), 2.12 - 1.99 (m, 1H), 1.97 - 1.45 (m, 8H), 1.40 - 1.27 (m, 2H).

EXAMPLE 2

Herbicidal Activity Data

10 Applications on a weed panel

[00274] Herbicidal activity of compounds (active ingredient; A.I.) was demonstrated by the following greenhouse experiments:

In-planta Low through-put screens (LTP) results

15 Post-emergence treatments

[00275] A basic panel of eight weed species (Table 3) sowed in 4X4X7cm plastic pots containing a garden mix (klasmann). Each specie was sowed in a separate pot. In each pot, 10-15 seeds were sowed according to the specie viability. Timing of application determined at a 1-2 true leaf stage. The plants grew for 30 days in a controlled greenhouse (26±2°C day, 20±2°C night). Flood irrigation (tap water + Shefer 5:3:8 8mM) was given at a 50% water content by weight. Two days before application, the tested plants thinned down to three plants per pot. Compounds were soluble in water (DDW), and commercial herbicide control was soluble in formulation B (Table 4). Before application, 1% (v/v) crop oil and 0.02% (v/v) surfactant (Tergitol™ 15-S-7) were added to the solution. Application was conducted with an industrial sprayer (TeeJet 6502E nozzle) at a rate of 2kg/ha and spray volume of 480l/ha. Plants were evaluated at 3 time points (4, 8, 12 days after application (DAA)). At each time point, visual phenotyping was recorded using a scale of 0-6 (0: no visible effect, 6: maximum effect). At 12 DAA, plants foliage was harvested, dried and weighed for dry weight analysis.

[00276] **Post-emergence advanced** dose response experiment included 4 species: SETVI, ECHCO, AMAPA, ABUTH (Tables 3, 5). Application was conducted at six rates between 0.6–0.0187kg/ha and spray volume of 480l/ha. Plants were evaluated at 4 time points (6, 12, 18 and 26 DAA). At each time point, visual phenotyping was recorded using a scale of 0-6 (0: no visible effect, 6: maximum effect). At 26 DAA, plants foliage was harvested, dried and weighed for dry weight analysis.

[00277] **Post-emergence advanced wide panel** experiment included 24 weed species (Table 5). Application was conducted at 2 rates of 2kg/ha and 0.25kg/ha and spray volume of 480l/ha. Visual phenotyping was recorded at 4, 11, 17 and 20 DAA using a scale of 0-6 (0: no visible effect, 6: maximum effect). At 21 DAA, plants foliage was harvested, dried, and weighed for dry weight analysis. All experiments included an untreated control, a solvent control, and a positive control (commercial herbicide A.I.). Statistical analysis for visual phenotyping determined by a median value of ≥3.5 and Fisher test

($p \leq 0.05$). Statistical analysis for dry weight determined by % inhibition ≥ 50 and T test ($p \leq 0.05$), as well as Wilcoxon test ($p \leq 0.05$).

Pre-emergence treatments

- 5 [00278] A basic panel of 8 weed species (Table 3) were sowed in 4X4X7 cm plastic pots containing inert sand (Sweet sand), intensively washed using osmosis water. Each specie was sowed in a separate pot. In each pot 10-15 seeds were sowed according to the specie viability. Sowing was performed one day before application. The plants were grown for 21 days in a controlled greenhouse ($26 \pm 2^\circ\text{C}$ day $20 \pm 2^\circ\text{C}$ night). Flood irrigation (tap water + Shefer 5:3:8 8mM) was given at a 50% water content by weight.
- 10 Compounds were soluble in water (DDW), and commercial herbicide control was soluble in formulation B (Table 4). Application was conducted with an industrial sprayer (TeeJet 6502E nozzle) at a rate of 2kg/ha and spray volume of 480l/ha.
- [00279] **Pre-emergence advanced** dose response experiment application was conducted at six rates between 1–0.0312kg/ha and spray volume of 480l/ha. Percentage of emergence was evaluated at 15 DAA.
- 15 Visual phenotyping was recorded using a scale of 0-6 (0: no visible effect, 6: maximum effect) at 18 DAA. All experiments included an untreated control, a solvent control and a positive control (commercial herbicide A.I.). Statistical analysis for visual phenotyping determined by a median value of ≥ 3.5 and Fisher test ($p \leq 0.05$). Statistical analysis for plant emergence determined by % emergence ≥ 50 and T test ($p \leq 0.05$), as well as Wilcoxon test ($p \leq 0.05$).

20

Table 3: Basic weed panel species

Bayer code	Scientific name
ABUTH	<i>Abutilon theophrasti</i>
AMBEL	<i>Ambrosia artemisiifolia</i>
AMAPA	<i>Amaranthus palmeri</i>
MATCH	<i>Matricaria chamomilla</i>
ALOMY	<i>Alopecurus myosuroides</i>
POAAN	<i>Poa annua</i>
LOLPE	<i>Lolium perenne</i>
SETVI	<i>Setaria viridis</i>

Table 4: Formulations

Formulation	Ingredients
A	100% DDW
B	48.5% acetone 40% DDW 10% isopropyl alcohol 1.5% DMSO

25 **Table 5:** Advanced weed panel species

Bayer code	Scientific name
ABUTH	<i>Abutilon theophrasti</i>
AMBEL	<i>Ambrosia artemisiifolia</i>
AMAPA	<i>Amaranthus palmeri</i>
MATCH	<i>Matricaria chamomilla</i>
ERICA	<i>Conyza Canadensis</i>

EPHHL	<i>Euphorbia heterophylla</i>
AMARE	<i>Amaranthus retroflexus</i>
SOLNI	<i>Solanum nigrum</i>
CHEAL	<i>Chenopodium album</i>
POROL	<i>Portulaca oleracea</i>
GLXMA	<i>Glycine max</i>
BRSNN	<i>Brassica napus</i>
ALOMY	<i>Alopecurus myosuroides</i>
POAAN	<i>Poa annua</i>
LOLPE	<i>Lolium perenne</i>
SETVI	<i>Setaria viridis</i>
DIGSA	<i>Digitaria sanguinalis</i>
ECHCO	<i>Echinochloa colona</i>
LOLRI	<i>Lolium rigidum</i>
PHAPA	<i>Phalaris paradoxa</i>
ZEAMX	<i>Zea mays</i>
AVEST	<i>Avena sterilis</i>
TRZAX	<i>Triticum aestivum</i>
ORYSA	<i>Oryza sativa</i>

Results:

5 [00280] The *In-planta* low throughput (LTP) results for compounds **104**, **109**, **148**, **146** and **145** are presented in Table 6 below:

Table 6. Herbicidal activity (*In-planta*) for compounds according to this invention.

<i>LTP-Pre emergence</i>	<i>Compound 104</i>	<i>Compound 109</i>	<i>Compound 148</i>	<i>Compound 146</i>
Dosage (g/ha)	2000	2000	2000	2000
<i>Setaria viridis</i>	100	0	40	60
<i>Alopecurus myosuroides</i>	40	0	0	0
<i>Lolium perenne</i>	80	0	0	40
<i>Poa annua</i>	100	0	60-80	0
<i>Echinochloa colona</i>	N/A	N/A	N/A	N/A
<i>Phalaris paradoxa</i>	60	N/A	N/A	N/A
<i>Eragrostis tef</i>	100	N/A	N/A	N/A
<i>Amaranthus palmeri</i>	80	40	60-80	60-80
<i>Matricaria chamomilla</i>	80	0	60-80	0
<i>Abutilon theophrasti</i>	60-80	0	40-60	0
<i>Ambrosia artemisiifolia</i>	0	0	60	0
<i>Sinapis alba</i>	40-60	N/A	N/A	N/A
<i>Arabidopsis thaliana</i>	40-80	N/A	N/A	N/A
<i>LTP-Post emergence</i>	<i>Compound 104</i>	<i>Compound 109</i>	<i>Compound 145</i>	
Dosage (g/ha)	2000	2000	2000	
<i>Phalaris minor</i>	20-30	N/A	0	
<i>Lolium multiflorum</i>	80-90	N/A	20-40	
<i>Echinochloa crus-galli</i>	N/A	N/A	60-80	
<i>Digitaria sanguinalis</i>	95	N/A	60-80	

<i>Setaria viridis</i>	90-95	0	60
<i>Alopecurus myosuroides</i>	95	0	60-80
<i>Lolium perenne</i>	85-90	0	40-60
<i>Poa annua</i>	90	0	20-40
<i>Echinochloa colon</i>	95	N/A	20-40
<i>Phalaris paradoxa</i>	80	N/A	N/A
<i>Amaranthus tuberculatus</i>	80	N/A	40-60
<i>Chenopodium album</i>	90	N/A	40-60
<i>Conyza canadensis</i>	95	N/A	40-60
<i>Galium aparine</i>	70	N/A	40-60
<i>Stellaria media</i>	95	N/A	20-40
<i>Solanum nigrum</i>	80-90	N/A	40-60
<i>Eragrostis tef</i>	60-100	N/A	N/A
<i>Amaranthus palmeri</i>	90	40	20-40
<i>Matricaria chamomilla</i>	70-80	0	0
<i>Abutilon theophrasti</i>	80	0	20-40
<i>Ambrosia artemisiifolia</i>	70-80	40	20-40
<i>Sinapis alba</i>	60	N/A	N/A
<i>Arabidopsis thaliana</i>	60	N/A	N/A

N/A: not available

[00281] The results show excellent control of compound **104** on both monocot and eudicot weeds species in both post-emergence and pre-emergence applications modes for most of the species tested. Compound **148** shows good to moderate activity on dicots and monocots when applied pre-emergence. In addition, compounds **109** and **146** displayed moderate growth inhibition in dicot species, in either pre- or post-emergence applications. Compound **145** displayed good to moderate activity when applied post-emergence to both monocot and dicot species.

10

***In-planta* high through-put screens (HTPS) results**

Post-emergence applications on miniature dicot model plants

[00282] *Arabidopsis thaliana* seeds were sown in 96 well plates filled with irrigated Sweet sand (10% > clay) that is washed from salts and minerals using tap water. 5-10 seeds were sown in the center of each well. 7-8 days post sowing, at 2 true leaves stage, thinning out was performed to ensure that compound application is done on a single plant per well. The plates were placed in a controlled greenhouse in a random order inside a bath which allows flooding irrigation with tap water supplemented with a fertilizer. Applied compounds were dissolved into a final solution of 50% acetone, 49.9% DDW, 0.1% tween-20.

[00283] A 96 well plate was used as a stock plate for preparing application solutions for 8 repeats. Each row contained different concentrations per chemical. Maximal concentration for application was 1.5Kg/Ha and dilution factor is 2.5. Chemical application was performed one day after thinning out in a chemical hood. 5µL applied on first two true leaves in each well using 12 channel pipettes. Data collection: RGB (red, green, blue) data for green area per well was documented using camera. Data is collected at a few

20

time points during the experiment: one day after thinning and before chemical application, two-, six- or nine-days post application. During the last two documentations, visual phenotyping was performed. Data analysis: Given RGB results and visual phenotype scores, a student *t*-test conducted to compare between treatment and control performance for continuous data (RGB) and Fisher exact test to analyze the non-continuous data (phenotypic scores). Dose-Response curves are generated for each treatment to infer ED₅₀ and max. inhibition parameters, using treatment's log concentration range as the dependent variable and normalized green area.

Pre-emergence applications on dicot and monocot model plants

[00284] These experiments are conducted similarly except that either *Arabidopsis thaliana* or *Eragrostis teff* seeds were sowed (5-10 or 5-7, respectively), plant thinning out was not performed, compound dissolved into a final solution of 50% acetone, 49.9% DDW, compound application was done at 30µL volume per well directly on sowed soil before plant emergence and data was collected 10 or 7 days after chemical application on dicot or monocot plants, respectively.

Imaging, RGB and statistical analysis

[00285] Plate imaging performed at 11 and 18 DAA. RGB data for green area per well was documented and used to extract % inhibition. Dose-Response curves were built for each treatment to infer EC₅₀, EC₇₅ and EC₉₀ parameters, using treatment's log concentration range as the independent variable and normalized green area as the dependent variable. Treatments were compared to controls performance given RGB results and using student's *t*-test (p-val≤0.05).

Results:

[00286] Compounds **104-154** were applied in either or both pre- and post-emergence mode to either or both monocot and dicot model plants, following the methods described above. The compounds were applied as a gradient of concentrations as listed above, from which the ED₅₀ was calculated.

[00287] The *in-planta* high through put screens (HTPS) results for compounds **101-154** are presented in Table 7 below:

Table 7. Herbicidal activity for compounds according to this invention.


System	Compound	Max Inhibition (%)	ED ₅₀ (g/ha)
HTPS Pre Monocot	104	100	35
	145	100	16.8
HTPS Pre Dicot	104	94	6
	109	100	24
	108	100	175
	147	100	N/A
HTPS Post Dicot	101	100	29
	102	90	2

104	100	0.2
105	94	430
106	82	>1000
109	100	200
107	100	200
154	100	8
152	100	4
149	100	90
123	100	140
115	100	13
114	100	40
145	100	N/A
113	100	26
143	100	61
147	50	807
148	100	30
146	100	80

[00288] The results in the table show excellent to very good control of most compounds, (particularly compound **104**) in the systems where they were tested. Compound **149** displayed very good activity also on monocots. Some compounds show weak to median activities in dicot systems.

WHAT IS CLAIMED:

1. A compound represented by the structure of formula **I(g)**:



5

I(g)

wherein

R₁, **R₁'**, **R₂**, **R₂'** and **R₄₀** are each independently H, C₁-C₅ linear or branched, unsubstituted alkyl, methyl, ethyl, propyl, iso-propyl, t-Bu, iso-butyl, pentyl, benzyl, C(O)-R₁₀, or C(O)-CH₃;


10 **R₃** is OH, or NH₂;

R₄ is NH₂, or OH;

wherein if **R₃** is OH then **R₄** is NH₂ and if **R₃** is NH₂ then **R₄** is OH;

wherein if **R₃** is OH and **R₄** is NH₂, then **n + m** cannot be equal to 3;

or **R₃** and **R₄** are joined together to form ring **A**, represented by the following structure:



15

R₅ is H, C₁-C₅ linear or branched, substituted or unsubstituted alkyl (e.g., methyl, CH₂SH, ethyl, butyl, CH₂-CCH, iso-propyl, CH₂-C(O)-OCH₃), C₂-C₅ linear or branched, substituted or unsubstituted alkenyl, C₂-C₅ linear or branched, substituted or unsubstituted alkynyl (e.g., CCH, CH₂-CCH), C₁-C₅ linear or branched haloalkyl (e.g., CF₃, CF₂CH₃, CH₂CF₃, CF₂CH₂CH₃, CH₂CH₂CF₃, CF₂CH(CH₃)₂, CF(CH₃)-CH(CH₃)₂), R₈-aryl (e.g., CH₂-Ph), C(=CH₂)-R₁₀ (e.g., C(=CH₂)-C(O)-OCH₃, C(=CH₂)-CN), substituted or unsubstituted alkyl sulfone (e.g., SO₂-CH₂-cyclopentyl), substituted or unsubstituted aryl (e.g., phenyl), substituted or unsubstituted heteroaryl (e.g., pyridine (2, 3, and 4-pyridine));

20

R₈ is [CH₂]_p

25

wherein **p** is between 1 and 10;

R₁₀ is H, CN, C₁-C₅ linear or branched alkyl, C(O)R, or S(O)₂R;

R is C₁-C₅ linear or branched alkyl, C₁-C₅ linear or branched alkoxy, phenyl, aryl or heteroaryl;

m is 1 or 2;

30

n is 0, 1, 2 or 3;

X_1 is S, O, or CH_2 ;

X_2 is S, O, or CH_2 ;

or its agrochemically acceptable salt, stereoisomer, tautomer, hydrate, *N*-oxide, reverse amide analog, isotopic variant (e.g., deuterated analog), or any combination thereof.

- 5 2. The compound of claim 1, wherein ring A has two chiral centers.
3. The compound of claim 1, wherein the compound is not (6*R*,7*S*)-6-amino-7-hydroxyoctanoic acid or 5-((4*R*,5*S*)-5-methyl-2-oxooxazolidin-4-yl)pentanoic acid.
4. The compound of claim 1 or 2, wherein the compound is a substantially pure single stereoisomer.
5. The compound of claim 1 or 2, wherein the compound is a mixture of stereoisomers.
- 10 6. The compounds of any one of claims 1-3, wherein the compound is the substantially pure *SR* stereoisomer, *RS* stereoisomer, *RR* stereoisomer, or *SS* diastereomer.
7. The compound of claim 4 or 6, wherein the substantially pure stereoisomer has a purity higher than 90%, preferably higher than 95%, most preferably higher than 98%.
8. The compound of claim 1, wherein R_1 and R_1' are both H, R_2 is CH_3 , R_2' is H or CH_3 , R_{40} is H or CH_3 , X_1 is CH_2 , X_2 is CH_2 , X_3 is O, NH or N- CH_3 , R_5 is H, ethyl, butyl, CH_2 -CCH, CH_2 -C(O)-OCH₃ or SO_2 - CH_2 -cyclopentyl, R_3 is OH, R_4 is NH_2 or R_3 and R_4 are joined to form ring A, n is 1, m is 1, or any combination thereof.
- 15 9. The compound of any one of claims 1-8, represented by any one of the following structures:

Compound	Structure
101	
102	
104	
105	
113	
114	
115	
116	
117	


118	
119	
120	

10. A compound, represented by any one of the following structures:

Compound	Structure
101	
102	
104	
105	
113	
114	
115	
123	
116	
117	
118	
119	
124	
125	
126	

120	
127	
128	
129	
130	
131	
132	
133	
134	
137	
138	
139	
140	
141	
142	


11. An herbicidal compound, represented by the structure of formula **I(ga)**:



I(ga)

5 wherein

C_A and C_B are both chiral carbon centers, or C_A and C_B together with R_3 and R_4 are joined to form ring **A**, represented by the following structure:



R_1 , R_1' , R_2 , R_2' and R_{40} are each independently H, C₁-C₅ linear or branched, unsubstituted alkyl, methyl, ethyl, propyl, iso-propyl, t-Bu, iso-butyl, pentyl, benzyl, C(O)-R₁₀, or C(O)-CH₃;

R_3 is OH, SH, NH₂, NHNH₂, NHR, N(R)₂, NHC(O)OBz, -NHC(O)-R₁₀, NHC(O)CH₃, C₁-C₅ linear or branched, substituted or unsubstituted alkyl, methyl, ethyl, propyl, iso-propyl, t-Bu, iso-butyl, pentyl, substituted or unsubstituted C₃-C₈ cycloalkyl, C₁-C₅ linear or branched or C₃-C₈ cyclic haloalkyl, C₁-C₅ linear or branched or C₃-C₈ cyclic alkoxy, substituted or unsubstituted C₃-C₈ heterocyclic ring, or substituted or unsubstituted aryl;

R_4 is NH₂, NHNH₂, N(R)₂, -NHC(O)-R₁₀, NHC(O)H, NHC(O)CH₃, C₁-C₅ linear or branched, substituted or unsubstituted alkyl, C₁-C₅ linear or branched or C₃-C₈ cyclic haloalkyl, C₁-C₅ linear or branched or C₃-C₈ cyclic alkoxy, substituted or unsubstituted C₃-C₈ cycloalkyl, substituted or unsubstituted C₃-C₈ heterocyclic ring, or substituted or unsubstituted aryl;

or R_3 and R_4 are joined together to form ring **A** as described above;

wherein R_3 and R_4 cannot both be NH₂, and

wherein if R_3 is OH and R_4 is NH₂, then $n + m$ cannot be equal to 3;

R_5 is H, C₁-C₅ linear or branched, substituted or unsubstituted alkyl (e.g., methyl, CH₂SH, ethyl, butyl, CH₂-CCH, iso-propyl, CH₂-C(O)-OCH₃), C₂-C₅ linear or branched, substituted or unsubstituted alkenyl, C₂-C₅ linear or branched, substituted or unsubstituted alkynyl (e.g., CCH, CH₂-CCH), C₁-C₅ linear or branched haloalkyl (e.g., CF₃, CF₂CH₃, CH₂CF₃, CF₂CH₂CH₃, CH₂CH₂CF₃, CF₂CH(CH₃)₂, CF(CH₃)-CH(CH₃)₂), R₈-aryl (e.g., CH₂-Ph), C(=CH₂)-R₁₀ (e.g., C(=CH₂)-C(O)-OCH₃, C(=CH₂)-CN), substituted or unsubstituted alkyl sulfone (e.g., SO₂-CH₂-cyclopentyl), substituted or unsubstituted aryl (e.g., phenyl), substituted or unsubstituted heteroaryl (e.g., pyridine (2, 3, and 4-pyridine));

R_8 is [CH₂]_p

wherein p is between 1 and 10;

R_{10} and R_{11} are each independently H, CN, C₁-C₅ linear or branched alkyl, C(O)R, or S(O)₂R;

or R_{10} and R_{11} are joined to form a substituted or unsubstituted C₃-C₈ heterocyclic ring;

R is C₁-C₅ linear or branched alkyl, C₁-C₅ linear or branched alkoxy, phenyl, aryl or heteroaryl, or two gem R substituents are joined together to form a 5 or 6 membered heterocyclic ring;

m is 1 or 2;

n is 0, 1, 2 or 3;





















X_1 is S, O, or CH₂;























X_2 is S, O, or CH₂;

or its agrochemically acceptable salt, stereoisomer, tautomer, hydrate, *N*-oxide, reverse amide analog, isotopic variant (e.g., deuterated analog), or any combination thereof.

12. An herbicidal compound, represented by any one of the following structures:


Compound	Structure
101	
102	
104	
105	
106	
113	
114	
115	
123	
116	
117	
118	
119	
124	
125	
126	
120	

127	
128	
129	
130	
131	
132	
133	
134	
137	
138	
139	
140	
141	
142	
107	
112	
143	
145	
146	
147	

148	
150	
151	
152	
153	
154	
155	
156	
157	
158	
159	
160	
161	
162	
163	
164	
165	
166	
167	
168	
169	
170	

171	
108	
109	
172	
173	
174	
175	
176	
177	
178	

13. An herbicidal compound represented by the structure of formula **X(a)**:



5 wherein

R_5 is H, C_1 - C_5 linear or branched, substituted or unsubstituted alkyl (e.g., methyl, CH_2SH , ethyl, butyl, CH_2-CCH , iso-propyl, $CH_2-C(O)-OCH_3$), C_2 - C_5 linear or branched, substituted or unsubstituted alkenyl, C_2 - C_5 linear or branched, substituted or unsubstituted alkynyl (e.g., CCH , CH_2-CCH), C_1 - C_5 linear or branched haloalkyl (e.g., CF_3 , CF_2CH_3 , CH_2CF_3 , $CF_2CH_2CH_3$, $CH_2CH_2CF_3$, $CF_2CH(CH_3)_2$, $CF(CH_3)-CH(CH_3)_2$), R_8 -aryl (e.g., CH_2-Ph), $C(=CH_2)-R_{10}$ (e.g., $C(=CH_2)-C(O)-OCH_3$, $C(=CH_2)-CN$), substituted or unsubstituted aryl (e.g., phenyl), substituted or unsubstituted heteroaryl (e.g., pyridine (2, 3, and 4-pyridine));

R_8 is $[CH_2]_p$

wherein p is between 1 and 10;

R_{10} is H, CN, C₁-C₅ linear or branched alkyl (e.g., methyl, ethyl), C(O)R (e.g., C(O)(OCH₃)), or S(O)₂R;

R is H, C₁-C₅ linear or branched alkyl (e.g., methyl, ethyl), C₁-C₅ linear or branched alkoxy (e.g., methoxy), phenyl, aryl or heteroaryl, or two gem R substituents are joined together to form a 5 or 6 membered heterocyclic ring;

X_1 is S, O, CH₂, CH(R) or C(R)₂;

n and o are each independently an integer number between 0 and 2;


m is an integer number between 1 and 3;

or its agrochemically acceptable salt, stereoisomer, tautomer, hydrate, *N*-oxide, reverse amide analog, isotopic variant (e.g., deuterated analog), or any combination thereof.

14. The compound of claim 13, represented by any one of the following structures:

Compound	Structure
143	
144	
145	
124	
125	
126	
139	

15. A compound represented by the structure of formula **X(a)**:



15 wherein

R_5 is H, C₁-C₅ linear or branched, substituted or unsubstituted alkyl (e.g., methyl, CH₂SH, ethyl, butyl, CH₂-CCH, iso-propyl, CH₂-C(O)-OCH₃), C₂-C₅ linear or branched, substituted or unsubstituted alkenyl, C₂-C₅ linear or branched, substituted or unsubstituted alkynyl (e.g., CCH, CH₂-CCH), C₁-C₅ linear or branched haloalkyl (e.g., CF₃, CF₂CH₃, CH₂CF₃, CF₂CH₂CH₃, CH₂CH₂CF₃, CF₂CH(CH₃)₂, CF(CH₃)-CH(CH₃)₂), R₈-aryl (e.g., CH₂-Ph), C(=CH₂)-R₁₀ (e.g., C(=CH₂)-C(O)-OCH₃, C(=CH₂)-CN), substituted or unsubstituted aryl (e.g., phenyl), substituted or unsubstituted heteroaryl (e.g., pyridine (2, 3, and 4-pyridine));

R_8 is [CH₂]_p

wherein **p** is between 1 and 10;

R₁₀ is H, CN, C₁-C₅ linear or branched alkyl (e.g., methyl, ethyl), C(O)R (e.g., C(O)(OCH₃)), or S(O)₂R;

R is H, C₁-C₅ linear or branched alkyl (e.g., methyl, ethyl), C₁-C₅ linear or branched alkoxy (e.g., methoxy), phenyl, aryl or heteroaryl, or two gem R substituents are joined together to form a 5 or 6 membered heterocyclic ring;

X₁ is S, O, CH₂, CH(R) or C(R)₂;

wherein if **X₁** is CH₂, then **R₅** cannot be H;

n is 2;

10 **o** is 0;

m is an integer number between 1 and 3;


or its agrochemically acceptable salt, stereoisomer, tautomer, hydrate, *N*-oxide, reverse amide analog, isotopic variant (e.g., deuterated analog), or any combination thereof.

16. The compound of claim 15, represented by any one of the following structures:

Compound	Structure
124	
125	
126	
139	

15

17. An herbicidal compound represented by the structure of formula **X(b)**:



wherein

20

R₅ is H, C₁-C₅ linear or branched, substituted or unsubstituted alkyl (e.g., methyl, CH₂SH, ethyl, butyl, CH₂-CCH, iso-propyl, CH₂-C(O)-OCH₃), C₂-C₅ linear or branched, substituted or unsubstituted alkenyl, C₂-C₅ linear or branched, substituted or unsubstituted alkynyl (e.g., CCH, CH₂-CCH), C₁-C₅ linear or branched haloalkyl (e.g., CF₃, CF₂CH₃, CH₂CF₃, CF₂CH₂CH₃, CH₂CH₂CF₃, CF₂CH(CH₃)₂, CF(CH₃)-CH(CH₃)₂), R₈-aryl (e.g., CH₂-Ph), C(=CH₂)-R₁₀ (e.g., C(=CH₂)-C(O)-OCH₃, C(=CH₂)-CN), substituted or unsubstituted aryl (e.g., phenyl), substituted or unsubstituted heteroaryl (e.g., pyridine (2, 3, and 4-pyridine));

25

R_8 is $[CH_2]_p$

wherein p is between 1 and 10;

R_{10} is H, CN, C_1 - C_5 linear or branched alkyl (e.g., methyl, ethyl), $C(O)R$ (e.g., $C(O)(OCH_3)$), or $S(O)_2R$;

5 R is H, C_1 - C_5 linear or branched alkyl (e.g., methyl, ethyl), C_1 - C_5 linear or branched alkoxy (e.g., methoxy), phenyl, aryl or heteroaryl, or two gem R substituents are joined together to form a 5 or 6 membered heterocyclic ring;

X_1 is S, O, CH_2 , $CH(R)$ or $C(R)_2$;

X_2 is S, O, CH_2 , $CH(R)$ or $C(R)_2$;

10 n is an integer number between 0 and 2;


m is an integer number between 1 and 3;

or its agrochemically acceptable salt, stereoisomer, tautomer, hydrate, *N*-oxide, reverse amide analog, isotopic variant (e.g., deuterated analog), or any combination thereof.

18. The compound of claim 17, represented by the following structures:

Compound	Structure
148	
155	
166	

15 19. An herbicidal compound represented by the structure of formula **X(c)**:



X(c)

wherein

20 R_5 is H, C_1 - C_5 linear or branched, substituted or unsubstituted alkyl (e.g., methyl, CH_2SH , ethyl, butyl, CH_2-CCH , iso-propyl, $CH_2-C(O)-OCH_3$), C_2 - C_5 linear or branched, substituted or unsubstituted alkenyl, C_2 - C_5 linear or branched, substituted or unsubstituted alkynyl (e.g., CCH , CH_2-CCH), C_1 - C_5 linear or branched haloalkyl (e.g., CF_3 , CF_2CH_3 , CH_2CF_3 , $CF_2CH_2CH_3$, $CH_2CH_2CF_3$, $CF_2CH(CH_3)_2$, $CF(CH_3)-CH(CH_3)_2$), R_8 -aryl (e.g., CH_2-Ph), $C(=CH_2)-R_{10}$ (e.g., $C(=CH_2)-C(O)-OCH_3$, $C(=CH_2)-CN$), substituted or unsubstituted aryl (e.g., phenyl), substituted or unsubstituted heteroaryl (e.g., pyridine (2, 3, and 4-pyridine));

R_8 is $[CH_2]_p$

wherein **p** is between 1 and 10;

R₁₀ is H, CN, C₁-C₅ linear or branched alkyl (e.g., methyl, ethyl), C(O)R (e.g., C(O)(OCH₃)), or S(O)₂R;

R is H, C₁-C₅ linear or branched alkyl (e.g., methyl, ethyl), C₁-C₅ linear or branched alkoxy (e.g., methoxy), phenyl, aryl or heteroaryl, or two gem R substituents are joined together to form a 5 or 6 membered heterocyclic ring;

X₁ is S, O, CH₂, CH(R) or C(R)₂;








X₂ is S, O, CH₂, CH(R) or C(R)₂;

n is an integer number between 0 and 2;


m is an integer number between 1 and 3;

or its agrochemically acceptable salt, stereoisomer, tautomer, hydrate, *N*-oxide, reverse amide analog, isotopic variant (e.g., deuterated analog), or any combination thereof.

20. The compound of claim 19, represented by the following structures:

Compound	Structure
146	
147	
152	
153	
131	
132	
171	

21. A compound represented by the structure of formula **X(c)**:



X(c)

15

wherein

R₅ is H, C₁-C₅ linear or branched, substituted or unsubstituted alkyl (e.g., methyl, CH₂SH, ethyl, butyl, CH₂-CCH, iso-propyl, CH₂-C(O)-OCH₃), C₂-C₅ linear or branched, substituted or unsubstituted alkenyl, C₂-C₅ linear or branched, substituted or unsubstituted alkynyl (e.g., CCH, CH₂-CCH), C₁-C₅ linear or branched haloalkyl (e.g., CF₃, CF₂CH₃, CH₂CF₃, CF₂CH₂CH₃, CH₂CH₂CF₃, CF₂CH(CH₃)₂, CF(CH₃)-CH(CH₃)₂), R₈-aryl (e.g., CH₂-Ph), C(=CH₂)-R₁₀ (e.g., C(=CH₂)-C(O)-OCH₃, C(=CH₂)-CN), substituted or unsubstituted aryl (e.g., phenyl), substituted or unsubstituted heteroaryl (e.g., pyridine (2, 3, and 4-pyridine));

R₈ is [CH₂]_p

wherein **p** is between 1 and 10;

R₁₀ is H, CN, C₁-C₅ linear or branched alkyl (e.g., methyl, ethyl), C(O)R (e.g., C(O)(OCH₃)), or S(O)₂R;

R is H, C₁-C₅ linear or branched alkyl (e.g., methyl, ethyl), C₁-C₅ linear or branched alkoxy (e.g., methoxy), phenyl, aryl or heteroaryl, or two gem R substituents are joined together to form a 5 or 6 membered heterocyclic ring;

X₁ is S, O, CH₂, CH(R) or C(R)₂;



X₂ is O, CH₂, CH(R) or C(R)₂;

n is an integer number between 1 and 2;


m is an integer number between 1 and 3;

or its agrochemically acceptable salt, stereoisomer, tautomer, hydrate, *N*-oxide, reverse amide analog, isotopic variant (e.g., deuterated analog), or any combination thereof.

22. The compound of claim 21, represented by any one of the following structures:

Compound	Structure
131	
132	

23. An herbicidal compound represented by the structure of formula **X(d)**:



X(d)

25 wherein

R₅ is H, C₁-C₅ linear or branched, substituted or unsubstituted alkyl (e.g., methyl, CH₂SH, ethyl, butyl, CH₂-CCH, iso-propyl, CH₂-C(O)-OCH₃), C₂-C₅ linear or branched, substituted or unsubstituted alkenyl, C₂-C₅ linear or branched, substituted or unsubstituted alkynyl (e.g., CCH, CH₂-CCH), C₁-C₅ linear or branched haloalkyl (e.g., CF₃, CF₂CH₃, CH₂CF₃, CF₂CH₂CH₃, CH₂CH₂CF₃,

CF₂CH(CH₃)₂, CF(CH₃)-CH(CH₃)₂, R₈-aryl (e.g., CH₂-Ph), C(=CH₂)-R₁₀ (e.g., C(=CH₂)-C(O)-OCH₃, C(=CH₂)-CN), substituted or unsubstituted aryl (e.g., phenyl), substituted or unsubstituted heteroaryl (e.g., pyridine (2, 3, and 4-pyridine));

R₈ is [CH₂]_p

5 wherein **p** is between 1 and 10;

R₁₀ is H, CN, C₁-C₅ linear or branched alkyl (e.g., methyl, ethyl), C(O)R (e.g., C(O)(OCH₃)), or S(O)₂R;

R is H, C₁-C₅ linear or branched alkyl (e.g., methyl, ethyl), C₁-C₅ linear or branched alkoxy (e.g., methoxy), phenyl, aryl or heteroaryl, or two gem R substituents are joined together to form a 5 or
10 6 membered heterocyclic ring;



X₂ is S, O, CH₂, CH(R) or C(R)₂;

n is an integer number between 0 and 2;


m is an integer number between 1 and 3;

or its agrochemically acceptable salt, stereoisomer, tautomer, hydrate, *N*-oxide, reverse amide analog,
15 isotopic variant (e.g., deuterated analog), or any combination thereof.

24. The compound of claim 23, represented by any one of the following structures:

Compound	Structure
123	
154	

25. A compound represented by the structure of formula **X(d)**:



20 wherein

R₅ is H, C₁-C₅ linear or branched, substituted or unsubstituted alkyl (e.g., methyl, CH₂SH, ethyl, butyl, CH₂-CCH, iso-propyl, CH₂-C(O)-OCH₃), C₂-C₅ linear or branched, substituted or unsubstituted alkenyl, C₂-C₅ linear or branched, substituted or unsubstituted alkynyl (e.g., CCH, CH₂-CCH), C₁-C₅ linear or branched haloalkyl (e.g., CF₃, CF₂CH₃, CH₂CF₃, CF₂CH₂CH₃, CH₂CH₂CF₃,
25 CF₂CH(CH₃)₂, CF(CH₃)-CH(CH₃)₂), R₈-aryl (e.g., CH₂-Ph), C(=CH₂)-R₁₀ (e.g., C(=CH₂)-C(O)-OCH₃, C(=CH₂)-CN), substituted or unsubstituted aryl (e.g., phenyl), substituted or unsubstituted heteroaryl (e.g., pyridine (2, 3, and 4-pyridine));

R₈ is [CH₂]_p

wherein **p** is between 1 and 10;

R₁₀ is H, CN, C₁-C₅ linear or branched alkyl (e.g., methyl, ethyl), C(O)R (e.g., C(O)(OCH₃)), or S(O)₂R;

R is H, C₁-C₅ linear or branched alkyl (e.g., methyl, ethyl), C₁-C₅ linear or branched alkoxy (e.g., methoxy), phenyl, aryl or heteroaryl, or two gem R substituents are joined together to form a 5 or 6 membered heterocyclic ring;


X₂ is S, O, CH₂, CH(R) or C(R)₂;

n is an integer number between 0 and 2;

m is 1 or 3;

or its agrochemically acceptable salt, stereoisomer, tautomer, hydrate, *N*-oxide, reverse amide analog, isotopic variant (e.g., deuterated analog), or any combination thereof.

26. The compound of claim 25, represented by the following structure:

Compound	Structure
123	

27. The compound of any one of claims 10-26, wherein the compound is a substantially pure single stereoisomer.

28. The compound of any one of claims 1-27, wherein the substantial pure stereoisomer has a purity higher than 90%, preferably higher than 95%, most preferably higher than 98%.

29. The compound of any one of claims 1-28, for use in controlling the growth of an undesired plant.

30. The compound of claim 29, wherein the plant is a eudicot (dicot) or a monocotyledon (monocot).

31. The compound of claim 29 or 30, wherein said plant is a weed.

32. The compound of claim 31, wherein said weed comprises: *Abutilon theophrasti*, *Amaranthus palmeri*, *Ambrosia artemisiifolia*, *Alopecurus myosuroides*, *Avena sterilis*, *Chenopodium album*, *Conyza Canadensis*, *Digitaria sanguinalis*, *Echinochloa colona*, *Euphorbia heterophylla*, *Lolium perenne*, *Lolium rigidum*, *Matricaria chamomilla*, *Phalaris paradoxa*, *Poa annua*, *Portulaca oleracea*, *Setaria viridis*, *Solanum nigrum* or any combination thereof.

33. The compound of claim 30, wherein the dicot plant is *Arabidopsis thaliana*, and/or the monocot plant is *Dactyloctenium aegyptium* or *Eragrostis teff*.

34. The compound of any one of claims 29-33, for use in pre-plant treatments, pre-emergence treatments, post-emergence treatments, or any combination thereof.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/IL2021/051150

A. CLASSIFICATION OF SUBJECT MATTER

(PC(8) - A01N 31/02; A01N 33/08; A01N 37/02; C07C 53/00; C07C 215/08; C07C 321/14 (2022.01)
 CPC - A01N 31/02; A01N 33/08; A01N 37/02; C07C 53/00; C07C 215/08; C07C 321/14 (2022.01)

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
 see Search History document

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched
 see Search History document

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
 see Search History document

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A ✓	PUBCHEM, Substance Record for SID 326612545, Available Date: 25 January 2017 [retrieved on 30 November 2021]. Retrieved from the Internet: <URL: https://pubchem.ncbi.nlm.nih.gov/substance/326612545 >. entire document	1, 3, 4, 6
A ✓	PUBCHEM, Substance Record for SID 341526259, Available Date: 13 September 2017 [retrieved on 30 November 2021]. Retrieved from the Internet: <URL: https://pubchem.ncbi.nlm.nih.gov/substance/341526259 >. entire document	1, 3, 4, 6
A ✓	PUBCHEM, Substance Record for SID 319544054, Available Date: 02 December 2016 [retrieved on 30 November 2021]. Retrieved from the Internet: <URL: https://pubchem.ncbi.nlm.nih.gov/substance/319544054 >. entire document	1, 3, 4, 6
A ✓	PUBCHEM, Substance Record for SID 349320386, Available Date: 18 December 2017 [retrieved on 30 November 2021]. Retrieved from the Internet: <URL: https://pubchem.ncbi.nlm.nih.gov/substance/349320386 >. entire document	1, 3, 4, 6

Further documents are listed in the continuation of Box C.

See patent family annex.

* Special categories of cited documents:	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"A" document defining the general state of the art which is not considered to be of particular relevance	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"D" document cited by the applicant in the international application	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"E" earlier application or patent but published on or after the international filing date	"&" document member of the same patent family
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	
"O" document referring to an oral disclosure, use, exhibition or other means	
"P" document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search
 27 January 2022

Date of mailing of the international search report

FEB 22 2022

Name and mailing address of the ISA/US
 Mail Stop PCT, Attn: ISA/US, Commissioner for Patents
 P.O. Box 1450, Alexandria, VA 22313-1450
 Facsimile No. 571-273-8300

Authorized officer

Harry Kim

Telephone No. PCT Helpdesk: 571-272-4300

INTERNATIONAL SEARCH REPORT

International application No.

PCT/IL2021/051150

Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

3. Claims Nos.: 7, 9, 28-34
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

See extra sheet(s).

1. As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. As all searchable claims could be searched without effort justifying additional fees, this Authority did not invite payment of additional fees.
3. As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:

4. No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:
1, 3, 4, 6

Remark on Protest

- The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.
- The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.
- No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/IL2021/051150

Continued from Box No. III Observations where unity of invention is lacking

Claims 1, 3, 4, and 6 have been analyzed subject to the restriction that the claims read on a compound represented by the structure of formula I(g) wherein R1 and R1' are absent; R2, R2' and R40 are each independently H; R3 is OH; R4 is NH2; R5 is H; m is 1; n is 0; X1 is S; X2 is S; X3 is O; wherein the compound is the substantially pure SR stereoisomer; or its agrochemically acceptable salt, stereoisomer, tautomer, hydrate, N-oxide, reverse amide analog, isotopic variant, or any combination thereof.

This application contains the following inventions or groups of inventions which are not so linked as to form a single general inventive concept under PCT Rule 13.1. In order for all inventions to be examined, the appropriate additional examination fees need to be paid.

Group I+: claims 1-6, 8, and 10-27 are drawn to compounds represented by the structure of formula I(g) or its agrochemically acceptable salt, stereoisomer, tautomer, hydrate, N-oxide, reverse amide analog, isotopic variant, or any combination thereof, herbicidal compounds, represented by the structure of formula I(ga) or its agrochemically acceptable salt, stereoisomer, tautomer, hydrate, N-oxide, reverse amide analog, isotopic variant, or any combination thereof, herbicidal compounds represented by the structure of formula X(a) or its agrochemically acceptable salt, stereoisomer, tautomer, hydrate, N-oxide, reverse amide analog, isotopic variant, or any combination thereof, herbicidal compounds represented by the structure of formula X(b) or its agrochemically acceptable salt, stereoisomer, tautomer, hydrate, N-oxide, reverse amide analog, isotopic variant, or any combination thereof, herbicidal compounds represented by the structure of formula X(c) or its agrochemically acceptable salt, stereoisomer, tautomer, hydrate, N-oxide, reverse amide analog, isotopic variant, or any combination thereof, and herbicidal compounds represented by the structure of formula X(d) or its agrochemically acceptable salt, stereoisomer, tautomer, hydrate, N-oxide, reverse amide analog, isotopic variant, or any combination thereof.

The first invention of Group I+ is restricted based on the proviso if R3 is OH then R4 is NH2 and if R3 is NH2 then R4 is OH, if R3 is OH and R4 is NH2, then $n + m$ cannot be equal to 3; and is restricted to a compound represented by the structure of formula I(g) wherein R1 and R1' are absent; R2, R2' and R40 are each independently H; R3 is OH; R4 is NH2; R5 is H; m is 1; n is 0; X1 is S; X2 is S; X3 is O; wherein the compound is the substantially pure SR stereoisomer; or its agrochemically acceptable salt, stereoisomer, tautomer, hydrate, N-oxide, reverse amide analog, isotopic variant, or any combination thereof; herbicidal compounds, represented by the structure of formula I(ga) or its agrochemically acceptable salt, stereoisomer, tautomer, hydrate, N-oxide, reverse amide analog, isotopic variant, or any combination thereof, herbicidal compounds represented by the structure of formula X(a) or its agrochemically acceptable salt, stereoisomer, tautomer, hydrate, N-oxide, reverse amide analog, isotopic variant, or any combination thereof, herbicidal compounds represented by the structure of formula X(b) or its agrochemically acceptable salt, stereoisomer, tautomer, hydrate, N-oxide, reverse amide analog, isotopic variant, or any combination thereof, herbicidal compounds represented by the structure of formula X(c) or its agrochemically acceptable salt, stereoisomer, tautomer, hydrate, N-oxide, reverse amide analog, isotopic variant, or any combination thereof, and herbicidal compounds represented by the structure of formula X(d) or its agrochemically acceptable salt, stereoisomer, tautomer, hydrate, N-oxide, reverse amide analog, isotopic variant, or any combination thereof. It is believed that claims 1, 3, 4, and 6 read on this first named invention and thus these claims will be searched without fee to the extent that they read on the above embodiment.

Applicant is invited to elect additional formula(e) for each additional compound to be searched in a specific combination by paying an additional fee for each set of election. Each additional elected formula(e) requires the selection of a single definition for each compound variable. An exemplary election would be a compound represented by the structure of formula I(g) wherein R1, R1', R2, R2' and R40 are each independently H; R3 is OH; R4 is NH2; R5 is H; m is 1; n is 1; X1 is S; X2 is S; X3 is O; wherein the compound is the substantially pure SR stereoisomer; or its agrochemically acceptable salt, stereoisomer, tautomer, hydrate, N-oxide, reverse amide analog, isotopic variant, or any combination thereof; herbicidal compounds, represented by the structure of formula I(ga) or its agrochemically acceptable salt, stereoisomer, tautomer, hydrate, N-oxide, reverse amide analog, isotopic variant, or any combination thereof, herbicidal compounds represented by the structure of formula X(a) or its agrochemically acceptable salt, stereoisomer, tautomer, hydrate, N-oxide, reverse amide analog, isotopic variant, or any combination thereof, herbicidal compounds represented by the structure of formula X(b) or its agrochemically acceptable salt, stereoisomer, tautomer, hydrate, N-oxide, reverse amide analog, isotopic variant, or any combination thereof, herbicidal compounds represented by the structure of formula X(c) or its agrochemically acceptable salt, stereoisomer, tautomer, hydrate, N-oxide, reverse amide analog, isotopic variant, or any combination thereof, and herbicidal compounds represented by the structure of formula X(d) or its agrochemically acceptable salt, stereoisomer, tautomer, hydrate, N-oxide, reverse amide analog, isotopic variant, or any combination thereof. Additional formula(e) will be searched upon the payment of additional fees. Applicants must specify the claims that read on any additional elected inventions. Applicants must further indicate, if applicable, the claims which read on the first named invention if different than what was indicated above for this group. Failure to clearly identify how any paid additional invention fees are to be applied to the "+" group(s) will result in only the first claimed invention to be searched/examined.

The inventions listed in Groups I+ do not relate to a single general inventive concept under PCT Rule 13.1, because under PCT Rule 13.2 they lack the same or corresponding special technical features for the following reasons:

The Groups I+ formulae do not share a significant structural element requiring the selection of alternatives for the compound variables, R1, R1', R2, R2', R40, R3, R4, X1, X2, X3, R5, m, n, CA, CB, o, and accordingly these groups lack unity a priori.

Additionally, even if Groups I+ were considered to share the technical features of a compound represented by the core structure of formula I(g) or its agrochemically acceptable salt, stereoisomer, tautomer, hydrate, N-oxide, reverse amide analog, isotopic variant, or any combination thereof; a herbicidal compound, represented by the core structure of formulae I(ga), X(a), X(b), X(c), and X(d) or its agrochemically acceptable salt, stereoisomer, tautomer, hydrate, N-oxide, reverse amide analog, isotopic variant, or any combination thereof, these shared technical features do not represent a contribution over the prior art as disclosed Substance Record for PubChem SID 326612545, Substance Record for PubChem SID 341526259, Substance Record for PubChem SID 319544054, and Substance Record for PubChem SID 349320386.

Substance Record for PubChem SID 326612545 teaches a compound represented by the core structure of formula I(g) (Pg. 2, see shown structure); a herbicidal compound, represented by the core structure of formulae I(ga) (Pg. 2, see shown structure; i.e., the compound has the same core structure as instant invention; thus the compound must inherently exhibit the same properties, such as herbicidal activity.); a herbicidal compound, represented by the core structure of formulae X(a) (Pg. 2, see shown structure; i.e., the compound has the same core structure as instant invention; thus the compound must inherently exhibit the same properties, such as herbicidal activity.).

Substance Record for PubChem SID 341526259 teaches a herbicidal compound, represented by the core structure of formulae X(b) (Pg. 2, see shown structure; i.e., the compound has the same core structure as instant invention; thus the compound must inherently exhibit the same properties, such as herbicidal activity.).

INTERNATIONAL SEARCH REPORT

International application No.

PCT/IL2021/051150

Substance Record for PubChem SID 319544054 teaches a herbicidal compound, represented by the core structure of formulae X(c) (Pg. 2, see shown structure; i.e., the compound has the same core structure as instant invention; thus the compound must inherently exhibit the same properties, such as herbicidal activity.).

Substance Record for PubChem SID 349320386 teaches a herbicidal compound, represented by the core structure of formulae X(d) (Pg. 2, see shown structure; i.e., the compound has the same core structure as instant invention; thus the compound must inherently exhibit the same properties, such as herbicidal activity.).

The inventions listed in Groups I+ therefore lack unity under Rule 13 because they do not share a same or corresponding special technical feature.

WRITTEN OPINION OF THE
INTERNATIONAL SEARCHING AUTHORITY

International application No.

PCT/IL2021/051150

Supplemental Box

In case the space in any of the preceding boxes is not sufficient.

Continuation of:

the same properties, such as herbicidal activity.).

Substance Record for PubChem SID 319544054 teaches a herbicidal compound, represented by the core structure of formulae X(c) (Pg. 2, see shown structure; i.e., the compound has the same core structure as instant invention; thus the compound must inherently exhibit the same properties, such as herbicidal activity.).

Substance Record for PubChem SID 349320386 teaches a herbicidal compound, represented by the core structure of formulae X(d) (Pg. 2, see shown structure; i.e., the compound has the same core structure as instant invention; thus the compound must inherently exhibit the same properties, such as herbicidal activity.).

The inventions listed in Groups I+ therefore lack unity under Rule 13 because they do not share a same or corresponding special technical feature.