



(51) International Patent Classification:

C07D 471/10 (2006.01) A61P 9/00 (2006.01)
C07D 487/10 (2006.01) A61P 29/00 (2006.01)
A61K 31/435 (2006.01) A61P 35/00 (2006.01)

(21) International Application Number:

PCT/US2018/058611

(22) International Filing Date:

01 November 2018 (01.11.2018)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/581,116 03 November 2017 (03.11.2017) US

(71) Applicant: **BRISTOL-MYERS SQUIBB COMPANY**
[US/US]; Route 206 and Province Line Road, Princeton,
New Jersey 08543 (US).

(72) Inventors: **ZHU, Yeheng**; c/o Bristol-Myers Squibb Com-
pany, Route 206 and Province Line Road, Princeton,
New Jersey 08543 (US). **DEWNANI, Sunita, V.**; c/o
Bristol-Myers Squibb Company, Route 206 and Province
Line Road, Princeton, New Jersey 08543 (US). **EWING,
William, R.**; c/o Bristol-Myers Squibb Company, Route
206 and Province Line Road, Princeton, New Jersey 08543
(US).

(74) Agent: **LIU, Hong** et al.; Bristol-Myers Squibb Company,
Route 206 and Province Line Road, Princeton, New Jersey
08543 (US).

(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ,
CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO,
DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN,
HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP,

KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME,
MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ,
OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA,
SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN,
TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ,
UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ,
TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK,
EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV,
MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM,
TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW,
KM, ML, MR, NE, SN, TD, TG).


Declarations under Rule 4.17:

- as to applicant's entitlement to apply for and be granted a
patent (Rule 4.17(ii))
- as to the applicant's entitlement to claim the priority of the
earlier application (Rule 4.17(iii))
- of inventorship (Rule 4.17(iv))

Published:

- with international search report (Art. 21(3))

(54) Title: DIAZASPIRO ROCK INHIBITORS



(57) Abstract: The present invention provides compounds of Formula (I): or stereoisomers, tautomers, or pharmaceutically acceptable salts thereof, wherein all the variables are as defined herein. These compounds are selective ROCK inhibitors. This invention also relates to pharmaceutical compositions comprising these compounds and methods of treating cardiovascular, smooth muscle, oncologic, neuropathologic, autoimmune, fibrotic, and/or inflammatory disorders using the same.

DIAZASPIRO ROCK INHIBITORS

CROSS REFERENCE TO RELATED APPLICATIONS

This application is entitled to priority pursuant to 35 U.S.C. §119(e) to U.S.
5 provisional patent application No. 62/581,116, filed November 3, 2017, which is
incorporated herein in its entirety.

FIELD OF THE INVENTION

The present invention relates generally to novel diazaspiro compounds and their
10 analogues thereof, which are inhibitors of Rho kinases, compositions containing them,
and methods of using them, for example, for the treatment or prophylaxis of disorders
associated with aberrant Rho kinase activity.

BACKGROUND OF THE INVENTION

15 Rho-Kinase (ROCK) is a member of the serine-threonine protein kinase family.
ROCK exists in two isoforms, ROCK1 and ROCK2 (Ishizaki, T. et al., *EMBO J.*,
15:1885-1893 (1996)). ROCK has been identified as an effector molecule of RhoA, a
small GTP-binding protein (G protein) that plays a key role in multiple cellular signaling
pathways. ROCK and RhoA are ubiquitously expressed across tissues. The RhoA/ROCK
20 signaling pathway is involved in a number of cellular functions, such as ACTIN®
organization, cell adhesion, cell migration, and cytokinesis (Riento, K. et al., *Nat. Rev.*
Mol. Cell Biol., 4:446-456 (2003)). It is also directly involved in regulating smooth
muscle contraction (Somlyo, A.P., *Nature*, 389:908-911 (1997)). Upon activation of its
receptor, RhoA is activated, and, in turn, it activates ROCK. Activated ROCK
25 phosphorylates the myosin-binding subunit of myosin light chain phosphatase, which
inhibits activity of the phosphatase and leads to contraction. Contraction of the smooth
muscle in the vasculature increases blood pressure, leading to hypertension.

There is considerable evidence in the literature that the Rho A/ROCK signaling
pathway plays an important role in signal transduction initiated by several vasoactive
30 factors, for example angiotensin II (Yamakawa, T. et al., *Hypertension*, 35:313-318
(2000)), urotension II (Sauzeau, V. et al., *Circ. Res.*, 88:1102-1104 (2001)), endothelin-1
(Tangkijvanich, P. et al., *Hepatology*, 33:74-80 (2001)), serotonin (Shimokawa, H., *Jpn.*

Circ. J., 64:1-12 (2000)), norepinephrine (Martinez, M.C. et al., *Am. J. Physiol.*, 279:H1228-H1238 (2000)) and platelet-derived growth factor (PDGF) (Kishi, H. et al., *J. Biochem.*, 128:719-722 (2000)). Many of these factors are implicated in the pathogenesis of cardiovascular disease.

5 Additional studies in the literature, some using the known ROCK inhibitors fasudil (Asano, T. et al., *J. Pharmacol. Exp. Ther.*, 241:1033-1040 (1987)) or Y-27632 (Uehata, M. et al., *Nature*, 389:990-994 (1997)) further illustrate the link between ROCK and cardiovascular disease. For example, ROCK expression and activity have been shown to be elevated in spontaneously hypertensive rats, suggesting a link to the development of
10 hypertension in these animals (Mukai, Y. et al., *FASEB J.*, 15:1062-1064 (2001)). The ROCK inhibitor Y-27632 (Uehata, M. et al., *Nature, ibid.*) was shown to significantly decrease blood pressure in three rat models of hypertension, including the spontaneously hypertensive rat, renal hypertensive rat and deoxycortisone acetate salt hypertensive rat models, while having only a minor effect on blood pressure in control rats. This
15 reinforces the link between ROCK and hypertension.

Other studies suggest a link between ROCK and atherosclerosis. For example, gene transfer of a dominant negative form of ROCK suppressed neointimal formation following balloon injury in porcine femoral arteries (Eto, Y. et al., *Am. J. Physiol. Heart Circ. Physiol.*, 278:H1744-H1750 (2000)). In a similar model, ROCK inhibitor Y-27632
20 also inhibited neointimal formation in rats (Sawada, N. et al., *Circulation*, 101:2030-2033 (2000)). In a porcine model of IL-1 beta-induced coronary stenosis, long term treatment with the ROCK inhibitor fasudil was shown to progressively reduce coronary stenosis, as well as promote a regression of coronary constrictive remodeling (Shimokawa, H. et al., *Cardiovascular Res.*, 51:169-177 (2001)).

25 Additional investigations suggest that a ROCK inhibitor would be useful in treating other cardiovascular diseases. For example, in a rat stroke model, fasudil was shown to reduce both the infarct size and neurologic deficit (Toshima, Y., *Stroke*, 31:2245-2250 (2000)). The ROCK inhibitor Y-27632 was shown to improve ventricular hypertrophy, fibrosis and function in a model of congestive heart failure in Dahl salt-
30 sensitive rats (Kobayashi, N. et al., *Cardiovascular Res.*, 55:757-767 (2002)).

Other animal or clinical studies have implicated ROCK in additional diseases including coronary vasospasm (Shimokawa, H. et al., *Cardiovasc. Res.*, 43:1029-1039

(1999)), cerebral vasospasm (Sato, M. et al., *Circ. Res.*, 87:195-200 (2000)), ischemia/reperfusion injury (Yada, T. et al., *J. Am. Coll. Cardiol.*, 45:599-607 (2005)), pulmonary hypertension (Fukumoto, Y. et al., *Heart*, 91:391-392 (2005)), angina (Shimokawa, H. et al., *J. Cardiovasc. Pharmacol.*, 39:319-327 (2002)), renal disease
 5 (Sato, S. et al., *Eur. J. Pharmacol.*, 455:169-174 (2002)) and erectile dysfunction (Gonzalez-Cadavid, N.F. et al., *Endocrine*, 23:167-176 (2004)).

In another study, it has been demonstrated that inhibition of the RhoA/ROCK signaling pathway allows formation of multiple competing lamellipodia that disrupt the productive migration of monocytes (Worthylake, R.A. et al., *J. Biol. Chem.*, 278:13578-
 10 13584 (2003)). It has also been reported that small molecule inhibitors of Rho Kinase are capable of inhibiting MCP-1 mediated chemotaxis *in vitro* (Iijima, H., *Bioorg. Med. Chem.*, 15:1022-1033 (2007)). Due to the dependence of immune cell migration upon the RhoA/ROCK signaling pathway one would anticipate inhibition of Rho Kinase should also provide benefit for diseases such as rheumatoid arthritis, psoriasis, and inflammatory
 15 bowel disease.

The above studies provide evidence for a link between ROCK and cardiovascular diseases including hypertension, atherosclerosis, restenosis, stroke, heart failure, coronary vasospasm, cerebral vasospasm, ischemia/reperfusion injury, pulmonary hypertension and angina, as well as renal disease and erectile dysfunction. Given the demonstrated effect of
 20 ROCK on smooth muscle, ROCK inhibitors may also be useful in other diseases involving smooth muscle hyper-reactivity, including asthma and glaucoma (Shimokawa, H. et al., *Arterioscler. Thromb. Vase. Biol.*, 25:1767-1775 (2005)). Furthermore, Rho-kinase has been indicated as a drug target for the treatment of various other diseases, including airway inflammation and hyperresponsiveness (Henry, P.J. et al., *Pulm.*
 25 *Pharmacol Ther.*, 18:67-74 (2005)), cancer (Rattan, R. et al., *J. Neurosci. Res.*, 83:243-255 (2006); Lepley, D. et al., *Cancer Res.*, 65:3788-3795 (2005)), fibrotic diseases (Jiang, C. et al., *Int. J. Mol. Sci.*, 13:8293-8307 (2012); Zhou, L. et al., *Am. J. Nephrol.*, 34:468-475 (2011)), as well as neurological disorders, such as spinal-cord injury, Alzheimer's disease, multiple sclerosis, stroke and neuropathic pain (Mueller, B.K. et al., *Nat. Rev.*
 30 *Drug Disc.*, 4:387-398 (2005); Sun, X. et al., *J. Neuroimmunol.*, 180:126-134 (2006)).

There remains an unmet medical need for new drugs to treat cardiovascular disease. In the 2012 update of Heart Disease and Stroke Statistics from the American

Heart Association (*Circulation*, 125:e2-e220 (2012)), it was reported that cardiovascular disease accounted for 32.8% of all deaths in the U.S., with coronary heart disease accounting for ~1 in 6 deaths overall in the U.S.. Contributing to these numbers, it was found that ~33.5% of the adult U.S. population was hypertensive, and it was estimated
5 that in 2010 ~6.6 million U.S. adults would have heart failure. Therefore, despite the number of medications available to treat cardiovascular diseases (CVD), including diuretics, beta blockers, angiotensin converting enzyme inhibitors, angiotensin blockers and calcium channel blockers, CVD remains poorly controlled or resistant to current medication for many patients.

10 There are many reports of ROCK inhibitors under investigation (see, for example, US 2012/0122842 A1, US 2010/0041645 A1, US 2008/0161297 A1, and Hu, E. et al., *Exp. Opin. Ther. Targets*, 9:715-736 (2005)). Reports also include WO2014/113620, WO 2014/134388, WO 2014/134391, WO2015/002915, WO2015/002926, WO2016/010950, WO2016/028971, WO2016/112236, WO2016/144936, and
15 WO2017/123860, all of which are assigned to the present applicant. However, fasudil is the only marketed ROCK inhibitor at this time. An i.v. formulation was approved in Japan for treatment of cerebral vasospasm. Thus, there remains a need for new therapeutics, including ROCK inhibitors, for the treatment of cardiovascular diseases, cancer, neurological diseases, renal diseases, fibrotic diseases, bronchial asthma, erectile
20 dysfunction, and glaucoma.

SUMMARY OF THE INVENTION

The present invention provides novel diazaspiro compounds and their analogues, including stereoisomers, tautomers, pharmaceutically acceptable salts, or solvates thereof,
25 which are useful as selective inhibitors of Rho kinases.

The present invention also provides processes and intermediates for making the compounds of the present invention.

The present invention also provides pharmaceutical compositions comprising a pharmaceutically acceptable carrier and at least one of the compounds of the present
30 invention or stereoisomers, tautomers, pharmaceutically acceptable salts, or solvates thereof.

The compounds of the invention may be used in the treatment and/or prophylaxis of conditions associated with aberrant ROCK activity.

The compounds of the present invention may be used in therapy.

5 The compounds of the present invention may be used for the manufacture of a medicament for the treatment and/or prophylaxis of a condition associated with aberrant ROCK activity.

In another aspect, the present invention is directed to a method of treating a cardiovascular or related disease which method comprises administering to a patient in need of such treatment a compound of the present invention as described above.

10 Examples of such diseases that may be treated include, for example, hypertension, atherosclerosis, restenosis, stroke, heart failure, renal failure, coronary artery disease, peripheral artery disease, coronary vasospasm, cerebral vasospasm, ischemia/reperfusion injury, pulmonary hypertension, angina, erectile dysfunction and renal disease.

In another aspect, the present invention is directed to a method of treating diseases
15 involving smooth muscle hyper reactivity including asthma, erectile dysfunction and glaucoma, which method comprises administering to a patient in need of such treatment a compound of the present invention as described above.

In another aspect, the present invention is directed to a method of treating diseases mediated at least partially by Rho kinase including fibrotic diseases, oncology, spinal-
20 cord injury, Alzheimer's disease, multiple sclerosis, stroke, neuropathic pain, rheumatoid arthritis, psoriasis and inflammatory bowel disease, which method comprises administering to a patient in need of such treatment a compound of the present invention as described above.

In yet additional aspects, the present invention is directed at pharmaceutical
25 compositions comprising the above-mentioned compounds, processes for preparing the above-mentioned compounds and intermediates used in these processes.

The compounds of the invention can be used alone, in combination with other compounds of the present invention, or in combination with one or more, preferably one to two other agent(s).


30 These and other features of the invention will be set forth in expanded form as the disclosure continues.

DETAILED DESCRIPTION OF THE INVENTION

I. COMPOUNDS OF THE INVENTION


In one aspect, the present invention provides, *inter alia*, compounds of Formula

5 (I):



or stereoisomers, tautomers, pharmaceutically acceptable salts, solvates, or prodrugs thereof, wherein

Ring A is selected from



J₁, J₂, J₃, and J₄ are independently selected from N, CR₃, and CR₄; provided no more than two of J₁, J₂, J₃, and J₄ are N;

L is selected from -C(O)-, -C(O)NH-, and -S(O)_p-;

R₁, at each occurrence, is independently selected from H, F, Cl, Br, OH, CN, NR_aR_a,
15 -OC₁₋₄ alkyl substituted with 0-3 R_e, and C₁₋₄ alkyl substituted with 0-3 R_e;

R₂, at each occurrence, is independently selected from H, -(CH₂)_rOR_b, (CH₂)_rS(O)_pR_c,
-(CH₂)_rC(=O)R_b, -(CH₂)_rNR_aR_a, -(CH₂)_rCN, -(CH₂)_rC(=O)NR_aR_a,
-(CH₂)_rNR_aC(=O)R_b, -(CH₂)_rNR_aC(=O)NR_aR_a, -(CH₂)_rNR_aC(=O)OR_b,
-(CH₂)_rOC(=O)NR_aR_a, -(CH₂)_rC(=O)OR_b, -(CH₂)_rS(O)_pNR_aR_a,
20 -(CH₂)_rNR_aS(O)_pNR_aR_a, -(CH₂)_rNR_aS(O)_pR_c, (CH₂)_r-C₃₋₆ carbocyclyl substituted
with 0-3 R_e, and -(CH₂)_r-heterocyclyl substituted with 0-3 R_e;

R₃, at each occurrence, is independently selected from H, F, Cl, Br, CN, C₁₋₄ alkyl
substituted with 0-3 R_e, -(CH₂)_rOR_b, (CH₂)_rS(O)_pR_c, -(CH₂)_rC(=O)R_b,
-(CH₂)_rNR_aR_a, -(CH₂)_rC(=O)NR_aR_a, -(CH₂)_rC(=O)(CH₂)_rNR_aR_a, -(CH₂)_rCN,
25 -(CH₂)_rNR_aC(=O)R_b, -(CH₂)_rNR_aC(=O)OR_b, -(CH₂)_rOC(=O)NR_aR_a,

- $-(CH_2)_rNR_aC(=O)NR_aR_a$, $-(CH_2)_rC(=O)OR_b$, $-(CH_2)_rS(O)_pNR_aR_a$,
 $-(CH_2)_rNR_aS(O)_pNR_aR_a$, $-(CH_2)_rNR_aS(O)_pR_c$, $(CH_2)_r$ -C₃₋₆ carbocyclyl substituted
 with 0-3 R_e, and $-(CH_2)_r$ -heterocyclyl substituted with 0-3 R_e;
 R₄, at each occurrence, is independently selected from H, F, Cl, Br, OH, CN, OC₁₋₄ alkyl
 5 substituted with 0-3 R_e, NR_aR_a, and C₁₋₄ alkyl substituted with 0-3 R_e;
 R₅ is selected from C₃₋₁₀ carbocyclyl substituted with 0-5 R₆ and heterocyclyl comprising
 carbon atoms and 1-6 heteroatoms selected from N, NR_{6a}, S, and O, and
 substituted with 0-6 R₆;
 R₆, at each occurrence, is independently selected from F, Cl, Br, =O, C₁₋₄alkyl,
 10 C₂₋₄alkenyl, C₂₋₄alkynyl, nitro, $-(CHR_d)_rS(O)_pR_c$, $-(CHR_d)_rS(O)_pNR_aR_a$,
 $-(CHR_d)_rNR_aS(O)_pR_c$, $-(CHR_d)_rOR_b$, $-(CHR_d)_rCN$, $-(CHR_d)_rNR_7R_7$,
 $-(CHR_d)_rNR_aC(=O)R_b$, $-(CHR_d)_rNR_aC(=O)NR_aR_a$, $-(CHR_d)_rC(=O)OR_b$,
 $-(CHR_d)_rC(=O)R_b$, $-(CHR_d)_rC(=O)NR_aR_a$, $-(CHR_d)_rOC(=O)R_b$, $-(CHR_d)_r$ -C₃₋₆
 cycloalkyl, $-(CHR_d)_r$ -heterocyclyl, $-(CHR_d)_r$ -aryl, and $-(CHR_d)_r$ -heteroaryl,
 15 wherein said alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclyl, aryl, or heteroaryl is
 substituted with 0-4 R_e;
 R_{6a} is selected from H, C₁₋₄alkyl, C₂₋₄alkenyl, C₂₋₄alkynyl, $-S(O)_pR_c$, $-S(O)_pNR_aR_a$, $-$
 $C(=O)OR_b$, $-C(=O)R_b$, $-C(=O)NR_aR_a$, $-(CHR_d)_r$ -C₃₋₆cycloalkyl,
 $-(CHR_d)_r$ -heterocyclyl, $-(CHR_d)_r$ -aryl, and $-(CHR_d)_r$ -heteroaryl, wherein said
 20 alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclyl, aryl, or heteroaryl is substituted
 with 0-4 R_e;
 R₇, at each occurrence, is independently selected from H, C₁₋₆ alkyl substituted with 1-5
 R_a, C₂₋₆ alkenyl substituted with 1-5 R_a, C₂₋₆ alkynyl substituted with 1-5 R_a,
 $-(CH_2)_r$ -C₃₋₁₀carbocyclyl substituted with 1-5 R_a, and $-(CH_2)_r$ -heterocyclyl
 25 substituted with 1-5 R_a; or R₇ and R₇ together with the nitrogen atom to which
 they are both attached form a heterocyclic ring substituted with 1-5 R_a;
 R_a, at each occurrence, is independently selected from H, C(=O)OR_b, C₁₋₆ alkyl
 substituted with 0-5 R_e, C₂₋₆ alkenyl substituted with 0-5 R_e, C₂₋₆ alkynyl
 substituted with 0-5 R_e, $-(CH_2)_r$ -C₃₋₁₀carbocyclyl substituted with 0-5 R_e, and
 30 $-(CH_2)_r$ -heterocyclyl substituted with 0-5 R_e; or R_a and R_a together with the
 nitrogen atom to which they are both attached form a heterocyclic ring substituted
 with 0-5 R_e;

- R_b , at each occurrence, is independently selected from H, C_{1-6} alkyl substituted with 0-5 R_e , C_{2-6} alkenyl substituted with 0-5 R_e , C_{2-6} alkynyl substituted with 0-5 R_e , $-(CH_2)_r-C_{3-10}$ carbocyclyl substituted with 0-5 R_e , and $-(CH_2)_r$ -heterocyclyl substituted with 0-5 R_e ;
- 5 R_c , at each occurrence, is independently selected from C_{1-6} alkyl substituted with 0-5 R_e , C_{2-6} alkenyl substituted with 0-5 R_e , C_{2-6} alkynyl substituted with 0-5 R_e , C_{3-6} carbocyclyl, and heterocyclyl;
- R_d , at each occurrence, is independently selected from H and C_{1-4} alkyl substituted with 0-5 R_e ;
- 10 R_e , at each occurrence, is independently selected from F, Cl, Br, CN, NO_2 , $=O$, C_{1-6} alkyl substituted with 1-5 R_f , C_{2-6} alkenyl, C_{2-6} alkynyl, $-(CH_2)_rOR_f$, $S(O)_pR_f$, $C(=O)NR_fR_f$, $NR_fC(=O)R_d$, $S(O)_pNR_fR_f$, $NR_fS(O)_pR_d$, $NR_fC(=O)OR_d$, $OC(=O)NR_fR_f$, $-(CH_2)_rNR_fR_f$, $-(CH_2)_r-C_{3-6}$ cycloalkyl substituted with 1-5 R_f , $-O(CH_2)_r-C_{3-6}$ cycloalkyl substituted with 1-5 R_f , $-(CH_2)_r$ -heterocyclyl substituted with 1-5 R_f , $-O(CH_2)_r$ -heterocyclyl substituted with 1-5 R_f , $-(CH_2)_r$ -aryl substituted with 1-5 R_f , $-O(CH_2)_r$ -aryl substituted with 1-5 R_f , $-(CH_2)_r$ -heteroaryl substituted with 1-5 R_f , and $-O(CH_2)_r$ -heteroaryl substituted with 1-5 R_f ;
- 15 R_f , at each occurrence, is independently selected from H, F, Cl, Br, CN, NO_2 , $=O$, C_{1-5} alkyl (optionally substituted with F, Cl, Br, OH, OC_{1-4} alkyl, NR_gR_g), $-(CH_2)_r-C_{3-10}$ carbocyclyl and $-(CH_2)_r$ -heterocyclyl, or R_f and R_f together with the nitrogen atom to which they are both attached form a heterocyclic ring optionally substituted with C_{1-4} alkyl;
- R_g , at each occurrence, is independently selected from H and C_{1-5} alkyl;
- m is an integer of zero, 1 or 2;
- 25 p is an integer of zero, 1 or 2;
- q is an integer of zero or 1;
- r is an integer of zero, 1, 2, 3 or 4; and
- t is an integer of zero, 1 or 2.

- 30 In another aspect, the present invention provides compounds of Formula (I), or stereoisomers, tautomers, pharmaceutically acceptable salts, solvates, or prodrugs thereof, wherein

R₁, at each occurrence, is independently selected from H, F, Cl, Br, OH, CN, NR_aR_a,

-OC₁₋₄ alkyl substituted with 0-3 R_e, and C₁₋₄ alkyl substituted with 0-3 R_e;

R_2 , at each occurrence, is independently selected from H, $-(CH_2)_bOR_b$, $(CH_2)_bS(O)_pR_c$,

$$-(\text{CH}_2)_r\text{C}(=\text{O})\text{R}_b, -(\text{CH}_2)_r\text{NR}_a\text{R}_a, -(\text{CH}_2)_r\text{CN}, -(\text{CH}_2)_r\text{NR}_a\text{C}(=\text{O})\text{OR}_b,$$

5 $-(\text{CH}_2)_r\text{OC}(=\text{O})\text{NR}_a\text{R}_a$, $-(\text{CH}_2)_r\text{C}(=\text{O})\text{OR}_b$, $-(\text{CH}_2)_r\text{NR}_a\text{S}(\text{O})_p\text{NR}_a\text{R}_a$, and

$$-(\text{CH}_2)_r\text{NR}_a\text{S}(\text{O})_p\text{R}_c;$$

R₃, at each occurrence, is independently selected from H, F, Cl, Br, CN, C₁₋₄ alkyl

substituted with 0-3 R_e, -(CH₂)_iOR_b, -(CH₂)_iS(O)_pR_c, -(CH₂)_iC(=O)R_b,
$$-(\text{CH}_2)_r\text{NR}_a\text{R}_a, -(\text{CH}_2)_r\text{C}(=\text{O})\text{NR}_a\text{R}_a, -(\text{CH}_2)_r\text{NR}_a\text{C}(=\text{O})\text{R}_b, -(\text{CH}_2)_r\text{OC}(=\text{O})\text{NR}_a\text{R}_a,$$

10 $-(\text{CH}_2)_r\text{NR}_a\text{C}(=\text{O})\text{NR}_a\text{R}_a$, $-(\text{CH}_2)_r\text{C}(=\text{O})\text{OR}_b$, $-(\text{CH}_2)_r\text{S}(\text{O})_p\text{NR}_a\text{R}_a$, $-(\text{CH}_2)_r\text{-C}_{3-6}$


carbocyclyl substituted with 0-3 R_e, and -(CH₂)_r-heterocyclyl substituted with 0-3

 $\mathbb{R}_e$


R₄, at each occurrence, is independently selected from H, F, Cl, Br, OH, CN, and C₁₋₄

alkyl substituted with 0-3 R_e:

15 R_5 is selected from



The image displays 24 chemical structures for the selection of R_5 , arranged in three rows of eight. Each structure is connected to a wavy line representing a bond. The structures include various rings and heterocycles with substituents (R_6) and R_{6a} indicated by superscripts and labels. The structures are: Row 1: 1. Cyclopropane with $(R_6)_{0-2}$. 2. Cyclohexane with $(R_6)_{0-4}$. 3. 2,5-dihydrofuran with $(R_6)_{0-1}$. 4. 2,5-dihydrothiophene with $(R_6)_{0-1}$. 5. 2,5-dihydro-1H-pyridine with $(R_6)_{0-1}$. Row 2: 6. 2,5-dihydro-1H-pyridine with $(R_6)_{0-1}$. 7. 2,5-dihydrothiophene with $(R_6)_{0-1}$. 8. 2,5-dihydro-1H-pyridine with $(R_6)_{0-1}$. 9. 2,5-dihydro-1H-pyridine with $(R_6)_{0-1}$. 10. 2,5-dihydro-1H-pyridine with $(R_6)_{0-1}$. 11. 2,5-dihydro-1H-pyridine with $(R_6)_{0-1}$. Row 3: 12. Benzene with $(R_6)_{0-4}$. 13. Pyridine with $(R_6)_{0-4}$. 14. Pyridine with $(R_6)_{0-3}$. 15. Pyridine with $(R_6)_{0-3}$. 16. Pyridine with $(R_6)_{0-2}$. 17. Pyridine with $(R_6)_{0-4}$. Row 4: 18. Naphthalene with $(R_6)_{0-3}$. 19. Naphthalene with $(R_6)_{0-3}$. 20. Naphthalene with $(R_6)_{0-3}$. 21. Naphthalene with $(R_6)_{0-3}$. 22. Naphthalene with $(R_6)_{0-4}$. 23. Naphthalene with $(R_6)_{0-3}$. 24. Naphthalene with $(R_6)_{0-3}$.




R_6 , at each occurrence, is independently selected from F, Cl, Br, =O, C₁₋₄alkyl, C₂₋₄alkenyl, C₂₋₄alkynyl, nitro, $-(CHR_d)_rS(O)_pR_c$, $-(CHR_d)_rS(O)_pNR_aR_a$, $-(CHR_d)_rNR_aS(O)_pR_c$, $-(CHR_d)_rOR_b$, $-(CHR_d)_rCN$, $-(CHR_d)_rNR_aR_a$, $-(CHR_d)_rNR_aC(=O)R_b$, $-(CHR_d)_rNR_aC(=O)NR_aR_a$, $-(CHR_d)_rC(=O)OR_b$, $-(CHR_d)_rC(=O)R_b$, $-C(=O)NR_aR_a$, $-(CHR_d)_rOC(=O)R_b$, $-(CHR_d)_r$ -C₃₋₆ cycloalkyl, $-(CHR_d)_r$ -heterocyclyl, $-(CHR_d)_r$ -aryl, and $-(CHR_d)_r$ -heteroaryl, wherein said alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclyl, aryl, or heteroaryl is substituted with 0-4 R_e ;

- R_{6a} , at each occurrence, is independently selected from H, C_{1-4} alkyl, $-S(O)_pR_c$, $-S(O)_pNR_aR_a$, $-C(=O)OR_b$, $-C(=O)R_b$, $-C(=O)NR_aR_a$, $-(CHR_d)_r-C_{3-6}$ cycloalkyl, $-(CHR_d)_r$ -heterocyclyl, $-(CHR_d)_r$ -aryl, and $-(CHR_d)_r$ -heteroaryl, wherein said alkyl, cycloalkyl, heterocyclyl, aryl, or heteroaryl is substituted with 0-4 R_e ;
- 5 R_7 , at each occurrence, is independently selected from H, C_{1-6} alkyl substituted with 1-5 R_a , C_{2-6} alkenyl substituted with 1-5 R_a , C_{2-6} alkynyl substituted with 1-5 R_a , $-(CH_2)_r-C_{3-10}$ carbocyclyl substituted with 1-5 R_a , and $-(CH_2)_r$ -heterocyclyl substituted with 1-5 R_a ; or R_7 and R_7 together with the nitrogen atom to which they are both attached form a heterocyclic ring substituted with 1-5 R_a ;
- 10 R_a , at each occurrence, is independently selected from H, $rC(=O)OR_b$, C_{1-6} alkyl substituted with 0-5 R_e , C_{2-6} alkenyl substituted with 0-5 R_e , C_{2-6} alkynyl substituted with 0-5 R_e , $-(CH_2)_r-C_{3-10}$ carbocyclyl substituted with 0-5 R_e , and $-(CH_2)_r$ -heterocyclyl substituted with 0-5 R_e ; or R_a and R_a together with the nitrogen atom to which they are both attached form a heterocyclic ring substituted
- 15 with 0-5 R_e ;
- R_b , at each occurrence, is independently selected from H, C_{1-6} alkyl substituted with 0-5 R_e , C_{2-6} alkenyl substituted with 0-5 R_e , C_{2-6} alkynyl substituted with 0-5 R_e , $-(CH_2)_r-C_{3-10}$ carbocyclyl substituted with 0-5 R_e , and $-(CH_2)_r$ -heterocyclyl substituted with 0-5 R_e ;
- 20 R_c , at each occurrence, is independently selected from C_{1-6} alkyl substituted with 0-5 R_e , C_{2-6} alkenyl substituted with 0-5 R_e , C_{2-6} alkynyl substituted with 0-5 R_e , C_{3-6} carbocyclyl, and heterocyclyl;
- R_d , at each occurrence, is independently selected from H and C_{1-4} alkyl substituted with 0-5 R_e ;
- 25 R_e , at each occurrence, is independently selected from F, Cl, Br, CN, NO_2 , $=O$, C_{1-6} alkyl substituted with 1-5 R_f , C_{2-6} alkenyl, C_{2-6} alkynyl, $-(CH_2)_rOR_f$, $S(O)_pR_f$, $C(=O)NR_fR_f$, $NR_fC(=O)R_d$, $S(O)_pNR_fR_f$, $NR_fS(O)_pR_d$, $NR_fC(=O)OR_d$, $OC(=O)NR_fR_f$, $-(CH_2)_rNR_fR_f$, $-(CH_2)_r-C_{3-6}$ cycloalkyl substituted with 1-5 R_f , $-O(CH_2)_r-C_{3-6}$ cycloalkyl substituted with 1-5 R_f , $-(CH_2)_r$ -heterocyclyl substituted
- 30 with 1-5 R_f , $-O(CH_2)_r$ -heterocyclyl substituted with 1-5 R_f , $-(CH_2)_r$ -aryl substituted with 1-5 R_f , $-O(CH_2)_r$ -aryl substituted with 1-5 R_f , $-(CH_2)_r$ -heteroaryl substituted with 1-5 R_f , and $-O(CH_2)_r$ -heteroaryl substituted with 1-5 R_f ;

R_f , at each occurrence, is independently selected from H, F, Cl, Br, CN, OH, C_{1-5} alkyl (optionally substituted with F, Cl, Br, OH, and OC_{1-4} alkyl, $-NR_gR_g$), $-(CH_2)_r-C_{3-10}$ carbocyclyl, and $-(CH_2)_r$ -heterocyclyl, or R_f and R_f together with the nitrogen atom to which they are both attached form a heterocyclic ring optionally substituted with C_{1-4} alkyl;


R_g , at each occurrence, is independently selected from H and C_{1-5} alkyl; and other variables are as defined in Formula (I).

In another aspect, the present invention provides compounds of Formula (II):



or stereoisomers, tautomers, pharmaceutically acceptable salts, solvates, or prodrugs thereof, wherein

Ring A is selected from



J_2 and J_4 are independently selected from N, CR_3 , and CR_4 ;

L is selected from $-C(O)-$ and $-S(O)_p-$;


R_1 , at each occurrence, is independently selected from H, F, Cl, Br, CN, NR_aR_a , and C_{1-4} alkyl substituted with 0-4 R_e ;


R_2 , at each occurrence, is independently selected from H, F, Cl, Br, OH, CN, NR_aR_a , and C_{1-4} alkyl substituted with 0-4 R_e ;

R_3 , at each occurrence, is independently selected from H, F, Cl, Br, CN, C_{1-4} alkyl substituted with 0-3 R_e , $-(CH_2)_rOR_b$, and $-C_{3-6}$ cycloalkyl;

R_4 , at each occurrence, is independently selected from H, F, Cl, Br, OH, CN, OC_{1-4} alkyl substituted with 0-3 R_e , and C_{1-4} alkyl substituted with 0-3 R_e ;

R_5 is selected from






- 5 R_6 , at each occurrence, is independently selected from F, Cl, Br, C_{1-4} alkyl, C_{2-4} alkenyl, C_{2-4} alkynyl, nitro, $-(CH_2)_rS(O)_pR_c$, $-(CH_2)_rS(O)_pNR_aR_a$, $-(CH_2)_rNR_aS(O)_pR_c$, $-(CH_2)_rOR_b$, $-(CH_2)_rCN$, $-(CH_2)_rNR_aR_a$, $-(CH_2)_rNR_aC(=O)R_b$, $-(CH_2)_rNR_aC(=O)NR_aR_a$, $-(CH_2)_rC(=O)OR_b$, $-(CH_2)_rC(=O)R_b$, $-C(=O)NR_aR_a$, $-(CH_2)_rOC(=O)R_b$, $-(CH_2)_r-C_{3-6}$ cycloalkyl, $-(CH_2)_r$ -heterocyclyl, $-(CH_2)_r$ -aryl, and $-(CH_2)_r$ -heteroaryl, wherein said alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclyl, aryl, or heteroaryl is substituted with 0-4 R_e ;
- 10 R_{6a} , at each occurrence, is independently selected from H, C_{1-4} alkyl, $-S(O)_pR_c$, $-S(O)_pNR_aR_a$, $-C(=O)OR_b$, $-C(=O)R_b$, $-C(=O)NR_aR_a$, $-(CH_2)_r-C_{3-6}$ cycloalkyl, $-(CH_2)_r$ -heterocyclyl, $-(CH_2)_r$ -aryl, and $-(CH_2)_r$ -heteroaryl, wherein said alkyl, cycloalkyl, heterocyclyl, aryl, or heteroaryl is substituted with 0-4 R_e ;
- 15 R_7 , at each occurrence, is independently selected from H, C_{1-6} alkyl substituted with 0-5 R_a , C_{2-6} alkenyl substituted with 1-5 R_a , C_{2-6} alkynyl substituted with 1-5 R_a , $-(CH_2)_r-C_{3-10}$ carbocyclyl substituted with 1-5 R_a , and $-(CH_2)_r$ -heterocyclyl substituted with 1-5 R_a ; or R_7 and R_7 together with the nitrogen atom to which they are both attached form a heterocyclic ring substituted with 1-5 R_a ;
- 20


- R_a , at each occurrence, is independently selected from H, $C(=O)OR_b$, C_{1-6} alkyl substituted with 0-5 R_e , C_{2-6} alkenyl substituted with 0-5 R_e , C_{2-6} alkynyl substituted with 0-5 R_e , $-(CH_2)_r-C_{3-10}$ cycloalkyl substituted with 0-5 R_e , $-(CH_2)_r$ -aryl substituted with 0-5 R_e , and $-(CH_2)_r$ -heterocyclyl substituted with 0-5 R_e ; or R_a and R_a together with the nitrogen atom to which they are both attached form a heterocyclic ring substituted with 0-5 R_e ;
- R_b , at each occurrence, is independently selected from H, C_{1-6} alkyl substituted with 0-5 R_e , C_{2-6} alkenyl substituted with 0-5 R_e , C_{2-6} alkynyl substituted with 0-5 R_e , $-(CH_2)_r-C_{3-10}$ carbocyclyl substituted with 0-5 R_e , and $-(CH_2)_r$ -heterocyclyl substituted with 0-5 R_e ;
- R_c , at each occurrence, is independently selected from C_{1-6} alkyl substituted with 0-5 R_e , C_{2-6} alkenyl substituted with 0-5 R_e , C_{2-6} alkynyl substituted with 0-5 R_e , C_{3-6} carbocyclyl, and heterocyclyl;
- R_e , at each occurrence, is independently selected from F, Cl, Br, CN, NO_2 , $=O$, C_{1-6} alkyl substituted with 1-5 R_f , C_{2-6} alkenyl, C_{2-6} alkynyl, CO_2H , $-(CH_2)_rOR_f$, $S(O)_pR_f$, $S(O)_pNR_fR_f$, $NR_fS(O)_pR_d$, $-(CH_2)_rNR_fR_f$, $-(CH_2)_r-C_{3-6}$ cycloalkyl, $-O(CH_2)_r-C_{3-6}$ cycloalkyl, $-(CH_2)_r$ -heterocyclyl, $-O(CH_2)_r$ -heterocyclyl, $-(CH_2)_r$ -aryl, and $-O(CH_2)_r$ -aryl;
- R_f , at each occurrence, is independently selected from H, F, Cl, Br, CN, OH, C_{1-5} alkyl (optionally substituted with F, Cl, Br, OH, and OC_{1-4} alkyl, $-NR_gR_g$), $-(CH_2)_r-C_{3-6}$ cycloalkyl, and $-(CH_2)_r$ -phenyl;
- R_g , at each occurrence, is independently selected from H and C_{1-5} alkyl; and other variables are as defined in Formula (I).

In another aspect, the present invention provides compounds of Formula (III):



or stereoisomers, tautomers, pharmaceutically acceptable salts, solvates, or prodrugs thereof, wherein

Ring A is selected from



J₂ is selected from N, CR₃, and CR₄;

L is selected from -C(O)- and -S(O)_p-;

5 R₁, at each occurrence, is independently selected from H, F, Cl, Br, CN, NR_aR_a, and C₁₋₄alkyl substituted with 0-4 R_e;

R₂, at each occurrence, is independently selected from H and C₁₋₄alkyl substituted with 0-4 R_e;

10 R₃, at each occurrence, is independently selected from H, F, Cl, Br, CN, C₁₋₄ alkyl substituted with 0-3 R_e, -(CH₂)_rOR_b, and -C₃₋₆cycloalkyl;

R₄, at each occurrence, is independently selected from H, F, Cl, Br, OH, CN, OC₁₋₄ alkyl substituted with 0-3 R_e, and C₁₋₄ alkyl substituted with 0-3 R_e;

15 R₆, at each occurrence, is independently selected from F, Cl, Br, C₁₋₄ alkyl, -(CH₂)_rOR_b, -(CH₂)_rCN, -(CH₂)_rNR₇R₇, -(CH₂)_rNR_aC(=O)R_b, -(CH₂)_rNR_aC(=O)NR_aR_a, -(CH₂)_rC(=O)OR_b, -(CH₂)_rC(=O)R_b, -C(=O)NR_aR_a, -(CH₂)_rOC(=O)R_b, -(CH₂)_rC(=O)NR_aR_a, -(CH₂)_r-C₃₋₆cycloalkyl, -(CH₂)_r-heterocyclyl, -(CH₂)_r-aryl, and -(CH₂)_r-heteroaryl, wherein said alkyl, cycloalkyl, heterocyclyl, aryl, or heteroaryl is substituted with 0-4 R_e;


20 R₇, at each occurrence, is independently selected from H, C₁₋₆ alkyl substituted with 1-5 R_a, C₂₋₆ alkenyl substituted with 1-5 R_a, C₂₋₆ alkynyl substituted with 1-5 R_a, -(CH₂)_r-C₃₋₁₀carbocyclyl substituted with 1-5 R_a, and -(CH₂)_r-heterocyclyl substituted with 1-5 R_a; or R₇ and R₇ together with the nitrogen atom to which they are both attached form a heterocyclic ring substituted with 1-5 R_a;

25 R_a, at each occurrence, is independently selected from H, C(=O)OR_b, C₁₋₆ alkyl substituted with 0-5 R_e, C₂₋₆ alkenyl substituted with 0-5 R_e, C₂₋₆ alkynyl substituted with 0-5 R_e, -(CH₂)_r-C₃₋₁₀cycloalkyl substituted with 0-5 R_e, -(CH₂)_r-aryl substituted with 0-5 R_e, and -(CH₂)_r-heterocyclyl substituted with 0-5

- R_e ; or R_a and R_a together with the nitrogen atom to which they are both attached form a heterocyclic ring substituted with 0-5 R_e ;
- R_b , at each occurrence, is independently selected from H, C_{1-6} alkyl substituted with 0-5 R_e , C_{2-6} alkenyl substituted with 0-5 R_e , C_{2-6} alkynyl substituted with 0-5 R_e ,
- 5 $-(CH_2)_r-C_{3-10}$ carbocyclyl substituted with 0-5 R_e , and $-(CH_2)_r$ -heterocyclyl substituted with 0-5 R_e ;
- R_c , at each occurrence, is independently selected from C_{1-6} alkyl substituted with 0-5 R_e , C_{2-6} alkenyl substituted with 0-5 R_e , C_{2-6} alkynyl substituted with 0-5 R_e , C_{3-6} carbocyclyl, and heterocyclyl;
- 10 R_e , at each occurrence, is independently selected from F, Cl, Br, CN, NO_2 , $=O$, C_{1-6} alkyl substituted with 1-5 R_f , C_{2-6} alkenyl, C_{2-6} alkynyl, CO_2H , $-(CH_2)_rOR_f$, $S(O)_pR_f$, $S(O)_pNR_fR_f$, $NR_fS(O)_pR_d$, $-(CH_2)_rNR_fR_f$, $-(CH_2)_r-C_{3-6}$ cycloalkyl, $-O(CH_2)_r-C_{3-6}$ cycloalkyl, $-(CH_2)_r$ -heterocyclyl, $-O(CH_2)_r$ -heterocyclyl, $-(CH_2)_r$ -aryl, and $-O(CH_2)_r$ -aryl;
- 15 R_f , at each occurrence, is independently selected from H, F, Cl, Br, CN, OH, C_{1-5} alkyl (optionally substituted with F, Cl, Br, OH, OC_{1-4} alkyl, NR_gR_g), $-(CH_2)_r-C_{3-6}$ cycloalkyl, and $-(CH_2)_r$ -phenyl;
- R_g , at each occurrence, is independently selected from H and C_{1-5} alkyl; and other variables are as defined in Formula (I).

20

In another aspect, the present invention provides compounds of Formula (IV):




or stereoisomers, tautomers, pharmaceutically acceptable salts, solvates, or prodrugs thereof, wherein

- 25 J_2 is selected from N and CH;
- R_3 is selected from H, CN, C_{1-4} alkyl, $-OC_{1-3}$ alkyl, and $-C_{3-6}$ cycloalkyl;
- R_6 , at each occurrence, is independently selected from F, Cl, Br, C_{1-4} alkyl, $-(CH_2)_rOR_b$, $-(CH_2)_rNR_7R_7$, $-(CH_2)_rNR_aC(=O)R_b$, $-(CH_2)_rNR_aC(=O)NR_aR_a$, $-(CH_2)_rC(=O)OR_b$,


- $-(CH_2)_rC(=O)R_b$, $-C(=O)NR_aR_a$, $-(CH_2)_rOC(=O)R_b$, $-(CH_2)_rC(=O)NR_aR_a$, $-(CH_2)_r$ - C_{3-6} cycloalkyl, $-(CH_2)_r$ -heterocyclyl, $-(CH_2)_r$ -aryl, and $-(CH_2)_r$ -heteroaryl, wherein said alkyl, cycloalkyl, heterocyclyl, aryl, or heteroaryl is substituted with 0-4 R_e ;
- R_7 , at each occurrence, is independently selected from H, C_{1-6} alkyl substituted with 1-5 R_a , C_{2-6} alkenyl substituted with 1-5 R_a , C_{2-6} alkynyl substituted with 1-5 R_a , $-(CH_2)_r$ - C_{3-10} carbocyclyl substituted with 1-5 R_a , and $-(CH_2)_r$ -heterocyclyl substituted with 1-5 R_a ; or R_7 and R_7 together with the nitrogen atom to which they are both attached form a heterocyclic ring substituted with 1-5 R_a ;
- R_a , at each occurrence, is independently selected from H, $C(=O)OR_b$, C_{1-6} alkyl substituted with 0-5 R_e , $-(CH_2)_r$ - C_{3-6} cycloalkyl substituted with 0-5 R_e , $-(CH_2)_r$ -aryl substituted with 0-5 R_e , and $-(CH_2)_r$ -heterocyclyl substituted with 0-5 R_e ; or R_a and R_a together with the nitrogen atom to which they are both attached form a heterocyclic ring substituted with 0-5 R_e ;
- R_b , at each occurrence, is independently selected from H, C_{1-6} alkyl substituted with 0-5 R_e , $-(CH_2)_r$ - C_{3-10} carbocyclyl substituted with 0-5 R_e , and $-(CH_2)_r$ -heterocyclyl substituted with 0-5 R_e ;
- R_e , at each occurrence, is independently selected from F, Cl, Br, CN, NO_2 , $=O$, C_{1-6} alkyl substituted with 1-5 R_f , CO_2H , OR_f , $NHS(O)_pC_{1-4}$ alkyl, NR_fR_f , $-(CH_2)_r$ - C_{3-6} cycloalkyl, $-O(CH_2)_r$ - C_{3-6} cycloalkyl, $-(CH_2)_r$ -heterocyclyl, $-O(CH_2)_r$ -heterocyclyl, $-(CH_2)_r$ -aryl, and $-O(CH_2)_r$ -aryl;
- R_f , at each occurrence, is independently selected from H, F, Cl, Br, CN, $=O$, C_{1-5} alkyl (optionally substituted with F, Cl, Br, OH, OC_{1-4} alkyl, NR_gR_g), $-(CH_2)_r$ - C_{3-6} cycloalkyl, and $-(CH_2)_r$ -phenyl;
- R_g , at each occurrence, is independently selected from H and C_{1-5} alkyl; and
- r is an integer of zero, 1, 2 or 3.

In another aspect, the present invention provides compounds of Formula (V):



or stereoisomers, tautomers, pharmaceutically acceptable salts, solvates, or prodrugs thereof, wherein


Ring A is selected from



5 R_1 is selected from H, F, Cl, Br, NR_aR_a , and C_{1-4} alkyl;

R_3 is $-OC_{1-3}$ alkyl;

R_7 , at each occurrence, is independently selected from H, C_{1-6} alkyl substituted with 1-5 R_a ; or R_7 and R_7 together with the nitrogen atom to which they are both attached form a heterocyclic ring selected from



15 R_a , at each occurrence, is independently selected from H, $C(=O)OR_b$, C_{1-6} alkyl substituted with 0-5 R_e , $-(CH_2)_r-C_{3-6}$ cycloalkyl substituted with 0-5 R_e , $-(CH_2)_r$ -aryl substituted with 0-5 R_e , and $-(CH_2)_r$ -heterocyclyl substituted with 0-5 R_e ;

R_b , at each occurrence, is independently selected from H, C_{1-6} alkyl substituted with 0-5 R_e , $-(CH_2)_r-C_{3-10}$ carbocyclyl substituted with 0-5 R_e , and $-(CH_2)_r$ -heterocyclyl substituted with 0-5 R_e ;


20 R_e , at each occurrence, is independently selected from F, Cl, Br, CN, NO_2 , $=O$, C_{1-6} alkyl substituted with 1-4 R_f , CO_2H , OR_f , $NHS(O)_pC_{1-4}$ alkyl, NR_fR_f , $-(CH_2)_r-C_{3-6}$ cycloalkyl, $-(CH_2)_r$ -heterocyclyl, $-(CH_2)_r$ -aryl, $-O(CH_2)_r-C_{3-6}$ cycloalkyl, $-O(CH_2)_r$ -heterocyclyl, and $-O(CH_2)_r$ -aryl;

R_f , at each occurrence, is independently selected from H, F, Cl, Br, CN, OH, C_{1-5} alkyl (optionally substituted with F, Cl, Br, OH, OC_{1-4} alkyl, NR_gR_g), $-(CH_2)_r-$ C_{3-6} cycloalkyl, and $-(CH_2)_r$ -phenyl;

R_g , at each occurrence, is independently selected from H and C_{1-5} alkyl; and

5 r is an integer of zero, 1, 2 or 3.


In another aspect, the present invention provides compounds of Formula (VI):



or stereoisomers, tautomers, pharmaceutically acceptable salts, solvates, or prodrugs

10 thereof, wherein

Ring A is selected from



R_1 is selected from H, F, Cl, Br, NR_aR_a , and C_{1-4} alkyl;

R_3 is $-OC_{1-4}$ alkyl;

15 R_a is selected from H, $C(=O)OC_{1-4}$ alkyl, C_{1-6} alkyl substituted with 0-5 R_e , $-(CH_2)_r-C_{3-6}$ cycloalkyl substituted with 0-5 R_e , $-(CH_2)_r$ -aryl substituted with 0-5 R_e , and $-(CH_2)_r$ -heterocyclyl substituted with 0-5 R_e ;

R_e , at each occurrence, is independently selected from F, Cl, Br, CN, NO_2 , $=O$, C_{1-4} alkyl

CO_2H , OR_f , $NHS(O)_pC_{1-4}$ alkyl, NR_fR_f , $-(CH_2)_r-C_{3-6}$ cycloalkyl,

20 $-(CH_2)_r$ -heterocyclyl, $-(CH_2)_r$ -aryl, $-O(CH_2)_r-C_{3-6}$ cycloalkyl,

$-O(CH_2)_r$ -heterocyclyl, and $-O(CH_2)_r$ -aryl;

R_f , at each occurrence, is independently selected from H, F, Cl, Br, CN, OH, C_{1-5} alkyl


(optionally substituted with F, Cl, Br, OH, OC_{1-4} alkyl, NR_gR_g), and $-(CH_2)_r$ -phenyl;

25 R_g , at each occurrence, is independently selected from H and C_{1-5} alkyl; and

r is an integer of zero, 1, 2 or 3.

In another aspect, the present invention provides compounds of Formula (VI), or stereoisomers, tautomers, pharmaceutically acceptable salts, solvates, or prodrugs thereof, wherein

R_a is selected from H, C(=O)OC₁₋₄alkyl, C₁₋₆ alkyl substituted with 0-5 R_e, -(CH₂)_r-C₃₋₆cycloalkyl substituted with 0-5 R_e, -(CH₂)_r-phenyl substituted with 0-5 R_e, and




R_e, at each occurrence, is independently selected from F, Cl, Br, CN, NO₂, =O, C₁₋₆ alkyl substituted with 1-4 R_f, OR_f NHS(O)_pC₁₋₄alkyl, NR_fR_f, -(CH₂)_r-C₃₋₆cycloalkyl, -(CH₂)_r-heterocyclyl, -(CH₂)_r-aryl, -O(CH₂)_r-C₃₋₆cycloalkyl, -O(CH₂)_r-heterocyclyl, and -O(CH₂)_r-aryl;

15 R_f, at each occurrence, is independently selected from H, F, Cl, Br, CN, OH, and C₁₋₅ alkyl (optionally substituted with F, Cl, Br, OH, OC₁₋₄ alkyl, and NR_gR_g), and -(CH₂)_r-phenyl;


R_g, at each occurrence, is independently selected from H and C₁₋₄alkyl; and


r is an integer of zero, 1, 2 or 3.

20 In another aspect, the present invention provides compounds of Formula (VII):



or stereoisomers, tautomers, pharmaceutically acceptable salts, solvates, or prodrugs thereof, wherein .





5 R₆, at each occurrence, is independently selected from F, Cl, Br, C₁₋₄alkyl, C₂₋₄alkenyl, C₂₋₄alkynyl, nitro, -(CH₂)_rS(O)_pR_c, -(CH₂)_rS(O)_pNR_aR_a, -(CH₂)_rNR_aS(O)_pR_c, -(CH₂)_rOR_b, -(CH₂)_rCN, -(CH₂)_rNR_aR_a, -(CH₂)_rNR_aC(=O)R_b, -(CH₂)_rNR_aC(=O)NR_aR_a, -(CH₂)_rC(=O)OR_b, -(CH₂)_rC(=O)R_b, -C(=O)NR_aR_a, -(CH₂)_rOC(=O)R_b, -(CH₂)_r-C₃₋₆cycloalkyl, -(CH₂)_r-heterocyclyl, -(CH₂)_r-aryl, and

10 -(CH₂)_r-heteroaryl, wherein said alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclyl, aryl, or heteroaryl is substituted with 0-4 R_e;

R_{6a}, at each occurrence, is independently selected from H, C₁₋₄alkyl, -S(O)_pR_c, -S(O)_pNR_aR_a, -C(=O)OR_b, -C(=O)R_b, -C(=O)NR_aR_a, -(CH₂)_r-C₃₋₆cycloalkyl, -(CH₂)_r-heterocyclyl, -(CH₂)_r-aryl, and -(CH₂)_r-heteroaryl, wherein said alkyl, cycloalkyl, heterocyclyl, aryl, or heteroaryl is substituted with 0-4 R_e;


15

p is an integer of zero, 1 or 2;


r is an integer of zero, 1, 2 or 3; and

other variables are as defined in Formula (I).

In another aspect, the present invention provides compounds of Formula (VII), or stereoisomers, tautomers, pharmaceutically acceptable salts, solvates, or prodrugs thereof, wherein




In another aspect, the present invention provides compounds of Formula (VIII):



15 or stereoisomers, tautomers, pharmaceutically acceptable salts, solvates, or prodrugs thereof, wherein

L is selected from $-C(O)-$ and $-S(O)_p-$;



R_6 , at each occurrence, is independently selected from F, Cl, Br, C_{1-4} alkyl, C_{2-4} alkenyl,

- 5 C_{2-4} alkynyl, nitro, $-(CH_2)_rS(O)_pR_c$, $-(CH_2)_rS(O)_pNR_aR_a$, $-(CH_2)_rNR_aS(O)_pR_c$, $-(CH_2)_rOR_b$, $-(CH_2)_rCN$, and $-(CH_2)_rNR_aR_a$;

R_{6a} , at each occurrence, is independently selected from H and C_{1-4} alkyl;

R_a , at each occurrence, is independently selected from H and C_{1-6} alkyl substituted with 0-5 R_e ;

- 10 R_b , at each occurrence, is independently selected from H and C_{1-6} alkyl substituted with 0-5 R_e ;

R_c , at each occurrence, is independently selected from C_{1-6} alkyl substituted with 0-5 R_e , C_{2-6} alkenyl substituted with 0-5 R_e , C_{2-6} alkynyl substituted with 0-5 R_e , C_{3-6} carbocyclyl, and heterocyclyl;

- 15 R_e , at each occurrence, is independently selected from F, Cl, Br, CN, NO_2 , $=O$, C_{1-6} alkyl substituted with 1-5 R_f , C_{2-6} alkenyl, C_{2-6} alkynyl, CO_2H , $-(CH_2)_rOR_f$, $S(O)_pR_f$, $S(O)_pNR_fR_f$, $NR_fS(O)_pC_{1-4}$ alkyl, $-(CH_2)_rNR_fR_f$, $-(CH_2)_r-C_{3-6}$ cycloalkyl, $-(CH_2)_r$ -heterocyclyl, and $-(CH_2)_r$ -aryl;

R_f , at each occurrence, is independently selected from H, F, Cl, Br, CN, OH, C_{1-5} alkyl

- 20 (optionally substituted with F, Cl, Br, OH, and OC_{1-4} alkyl, NR_gR_g), C_{3-6} cycloalkyl, and phenyl;

R_g , at each occurrence, is independently selected from H and C_{1-5} alkyl;

p is an integer of zero, 1 or 2; and

r is an integer of zero, 1, 2 or 3.

The invention may be embodied in other specific forms without departing from the spirit or essential attributes thereof. This invention also encompasses all combinations of alternative aspects of the invention noted herein. It is understood that any and all embodiments of the present invention may be taken in conjunction with any
5 other embodiment to describe additional embodiments of the present invention. Furthermore, any elements (including individual variable definitions) of an embodiment are meant to be combined with any and all other elements from any of the embodiments to describe additional embodiments.

In another aspect, the present invention provides a compound selected from any
10 subset list of compounds exemplified in the present application.

In another embodiment, the compounds of the present invention have ROCK IC₅₀ values $\leq 10 \mu\text{M}$.

In another embodiment, the compounds of the present invention have ROCK IC₅₀ values $\leq 1 \mu\text{M}$.

15 In another embodiment, the compounds of the present invention have ROCK IC₅₀ values $\leq 0.1 \mu\text{M}$.

In another embodiment, the compounds of the present invention have ROCK IC₅₀ values $\leq 0.05 \mu\text{M}$.

In another embodiment, the compounds of the present invention have ROCK IC₅₀
20 values $\leq 0.01 \mu\text{M}$.

II. OTHER EMBODIMENTS OF THE INVENTION

In another embodiment, the present invention provides a composition comprising at least one of the compounds of the present invention or a stereoisomer, a tautomer, a pharmaceutically acceptable salt, or a solvate thereof.

25 In another embodiment, the present invention provides a pharmaceutical composition comprising a pharmaceutically acceptable carrier and at least one of the compounds of the present invention or a stereoisomer, a tautomer, a pharmaceutically acceptable salt, or a solvate, thereof.

In another embodiment, the present invention provides a pharmaceutical
30 composition, comprising: a pharmaceutically acceptable carrier and a therapeutically effective amount of at least one of the compounds of the present invention or a stereoisomer, a tautomer, a pharmaceutically acceptable salt, or a solvate thereof.

In another embodiment, the present invention provides a process for making a compound of the present invention.

In another embodiment, the present invention provides an intermediate for making a compound of the present invention.

5 In another embodiment, the present invention provides a pharmaceutical composition further comprising additional therapeutic agent(s).

In another embodiment, the present invention provides a method for the treatment and/or prophylaxis of a condition associated with aberrant ROCK activity comprising administering to a patient in need of such treatment and/or prophylaxis a therapeutically
10 effective amount of at least one of the compounds of the present invention or a stereoisomer, a tautomer, a pharmaceutically acceptable salt, or a solvate thereof. As used herein, the term "patient" encompasses all mammalian species.

As used herein, "treating" or "treatment" cover the treatment of a disease-state in a mammal, particularly in a human, and include: (a) inhibiting the disease-state, *i.e.*,
15 arresting its development; and/or (b) relieving the disease-state, *i.e.*, causing regression of the disease state.

As used herein, "prophylaxis" or "prevention" covers the preventive treatment of a subclinical disease-state in a mammal, particularly in a human, aimed at reducing the probability of the occurrence of a clinical disease-state. Patients are selected for
20 preventative therapy based on factors that are known to increase risk of suffering a clinical disease state compared to the general population. "Prophylaxis" therapies can be divided into (a) primary prevention and (b) secondary prevention. Primary prevention is defined as treatment in a patient that has not yet presented with a clinical disease state, whereas secondary prevention is defined as preventing a second occurrence of the same
25 or similar clinical disease state. In another embodiment, the present invention provides a combined preparation of a compound of the present invention and additional therapeutic agent(s) for simultaneous, separate or sequential use in therapy.

The present invention may be embodied in other specific forms without departing from the spirit or essential attributes thereof. This invention encompasses all
30 combinations of preferred aspects of the invention noted herein. It is understood that any and all embodiments of the present invention may be taken in conjunction with any other embodiment or embodiments to describe additional embodiments. It is also to be

understood that each individual element of the embodiments is its own independent embodiment. Furthermore, any element of an embodiment is meant to be combined with any and all other elements from any embodiment to describe an additional embodiment.

5 III. CHEMISTRY

Throughout the specification and the appended claims, a given chemical formula or name shall encompass all stereo and optical isomers and racemates thereof where such isomers exist. Unless otherwise indicated, all chiral (enantiomeric and diastereomeric) and racemic forms are within the scope of the invention. Many geometric isomers of C=C
10 double bonds, C=N double bonds, ring systems, and the like can also be present in the compounds, and all such stable isomers are contemplated in the present invention. *Cis*- and *trans*- (or *E*- and *Z*-) geometric isomers of the compounds of the present invention are described and may be isolated as a mixture of isomers or as separated isomeric forms. The present compounds can be isolated in optically active or racemic forms. Optically
15 active forms may be prepared by resolution of racemic forms or by synthesis from optically active starting materials. All processes used to prepare compounds of the present invention and intermediates made therein are considered to be part of the present invention. When enantiomeric or diastereomeric products are prepared, they may be separated by conventional methods, for example, by chromatography or fractional
20 crystallization. Depending on the process conditions the end products of the present invention are obtained either in free (neutral) or salt form. Both the free form and the salts of these end products are within the scope of the invention. If so desired, one form of a compound may be converted into another form. A free base or acid may be converted into a salt; a salt may be converted into the free compound or another salt; a mixture of
25 isomeric compounds of the present invention may be separated into the individual isomers. Compounds of the present invention, free form and salts thereof, may exist in multiple tautomeric forms, in which hydrogen atoms are transposed to other parts of the molecules and the chemical bonds between the atoms of the molecules are consequently rearranged. It should be understood that all tautomeric forms, insofar as they may exist,
30 are included within the invention.

The term "stereoisomer" refers to isomers of identical constitution that differ in the arrangement of their atoms in space. Enantiomers and diastereomers are examples of

stereoisomers. The term "enantiomer" refers to one of a pair of molecular species that are mirror images of each other and are not superimposable. The term "diastereomer" refers to stereoisomers that are not mirror images. The term "racemate" or "racemic mixture" refers to a composition composed of equimolar quantities of two enantiomeric species,
5 wherein the composition is devoid of optical activity.

The symbols "R" and "S" represent the configuration of substituents around a chiral carbon atom(s). The isomeric descriptors "R" and "S" are used as described herein for indicating atom configuration(s) relative to a core molecule and are intended to be used as defined in the literature (IUPAC Recommendations 1996, *Pure and Applied*
10 *Chemistry*, 68:2193-2222 (1996)).

The term "chiral" refers to the structural characteristic of a molecule that makes it impossible to superimpose it on its mirror image. The term "homochiral" refers to a state of enantiomeric purity. The term "optical activity" refers to the degree to which a homochiral molecule or nonracemic mixture of chiral molecules rotates a plane of
15 polarized light.

As used herein, the term "alkyl" or "alkylene" is intended to include both branched and straight-chain saturated aliphatic hydrocarbon groups having the specified number of carbon atoms. For example, "C₁ to C₁₀ alkyl" or "C₁₋₁₀ alkyl" (or alkylene), is intended to include C₁, C₂, C₃, C₄, C₅, C₆, C₇, C₈, C₉, and C₁₀ alkyl groups. Additionally,
20 for example, "C₁ to C₆ alkyl" or "C₁₋₆ alkyl" denotes alkyl having 1 to 6 carbon atoms. Alkyl group can be unsubstituted or substituted with at least one hydrogen being replaced by another chemical group. Example alkyl groups include, but are not limited to, methyl (Me), ethyl (Et), propyl (*e.g.*, *n*-propyl and isopropyl), butyl (*e.g.*, *n*-butyl, isobutyl, *t*-butyl), and pentyl (*e.g.*, *n*-pentyl, isopentyl, neopentyl). When "C₀ alkyl" or "C₀
25 alkylene" is used, it is intended to denote a direct bond.

"Alkenyl" or "alkenylene" is intended to include hydrocarbon chains of either straight or branched configuration having the specified number of carbon atoms and one or more, preferably one to two, carbon-carbon double bonds that may occur in any stable point along the chain. For example, "C₂ to C₆ alkenyl" or "C₂₋₆ alkenyl" (or alkenylene), is
30 intended to include C₂, C₃, C₄, C₅, and C₆ alkenyl groups. Examples of alkenyl include, but are not limited to, ethenyl, 1-propenyl, 2-propenyl, 2-butenyl, 3-butenyl, 2-pentenyl,

3-pentenyl, 4-pentenyl, 2-hexenyl, 3-hexenyl, 4-hexenyl, 5-hexenyl, 2-methyl-2-propenyl, and 4-methyl-3-pentenyl.

"Alkynyl" or "alkynylene" is intended to include hydrocarbon chains of either straight or branched configuration having one or more, preferably one to three, carbon-carbon triple bonds that may occur in any stable point along the chain. For example, "C₂ to C₆ alkynyl" or "C₂₋₆ alkynyl" (or alkynylene), is intended to include C₂, C₃, C₄, C₅, and C₆ alkynyl groups; such as ethynyl, propynyl, butynyl, pentynyl, and hexynyl.

The term "alkoxy" or "alkyloxy" refers to an -O-alkyl group. "C₁ to C₆ alkoxy" or "C₁₋₆ alkoxy" (or alkyloxy), is intended to include C₁, C₂, C₃, C₄, C₅, and C₆ alkoxy groups. Example alkoxy groups include, but are not limited to, methoxy, ethoxy, propoxy (*e.g.*, *n*-propoxy and isopropoxy), and *t*-butoxy. Similarly, "alkylthio" or "thioalkoxy" represents an alkyl group as defined above with the indicated number of carbon atoms attached through a sulphur bridge; for example methyl-S- and ethyl-S-.

"Halo" or "halogen" includes fluoro (F), chloro (Cl), bromo (Br), and iodo (I). "Haloalkyl" is intended to include both branched and straight-chain saturated aliphatic hydrocarbon groups having the specified number of carbon atoms, substituted with 1 or more halogens. Examples of haloalkyl include, but are not limited to, fluoromethyl, difluoromethyl, trifluoromethyl, trichloromethyl, pentafluoroethyl, pentachloroethyl, 2,2,2-trifluoroethyl, heptafluoropropyl, and heptachloropropyl. Examples of haloalkyl also include "fluoroalkyl" that is intended to include both branched and straight-chain saturated aliphatic hydrocarbon groups having the specified number of carbon atoms, substituted with 1 or more fluorine atoms.

"Haloalkoxy" or "haloalkyloxy" represents a haloalkyl group as defined above with the indicated number of carbon atoms attached through an oxygen bridge. For example, "C₁ to C₆ haloalkoxy" or "C₁₋₆ haloalkoxy", is intended to include C₁, C₂, C₃, C₄, C₅, and C₆ haloalkoxy groups. Examples of haloalkoxy include, but are not limited to, trifluoromethoxy, 2,2,2-trifluoroethoxy, and pentafluoroethoxy. Similarly, "haloalkylthio" or "thiohaloalkoxy" represents a haloalkyl group as defined above with the indicated number of carbon atoms attached through a sulphur bridge; for example trifluoromethyl-S-, and pentafluoroethyl-S-.

The term "cycloalkyl" refers to cyclized alkyl groups, including mono-, bi- or poly-cyclic ring systems. "C₃ to C₇ cycloalkyl" or "C₃₋₇ cycloalkyl" is intended to include C₃, C₄, C₅, C₆, and C₇ cycloalkyl groups. Example cycloalkyl groups include, but are not limited to, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, and norbornyl. Branched cycloalkyl groups such as 1-methylcyclopropyl and 2-methylcyclopropyl are included in the definition of "cycloalkyl".

As used herein, "carbocycle" or "carbocyclic residue" is intended to mean any stable 3-, 4-, 5-, 6-, 7-, or 8-membered monocyclic or bicyclic or 7-, 8-, 9-, 10-, 11-, 12-, or 13-membered bicyclic or tricyclic hydrocarbon ring, any of which may be saturated, partially unsaturated, unsaturated or aromatic. Examples of such carbocycles include, but are not limited to, cyclopropyl, cyclobutyl, cyclobutenyl, cyclopentyl, cyclopentenyl, cyclohexyl, cycloheptenyl, cycloheptyl, cycloheptenyl, adamantyl, cyclooctyl, cyclooctenyl, cyclooctadienyl, [3.3.0]bicyclooctane, [4.3.0]bicyclononane, [4.4.0]bicyclodecane (decalin), [2.2.2]bicyclooctane, fluorenyl, phenyl, naphthyl, indanyl, adamantyl, anthracenyl, and tetrahydronaphthyl (tetralin). As shown above, bridged rings are also included in the definition of carbocycle (*e.g.*, [2.2.2]bicyclooctane). Preferred carbocycles, unless otherwise specified, are cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, phenyl, and indanyl. When the term "carbocycle" is used, it is intended to include "aryl". A bridged ring occurs when one or more carbon atoms link two non-adjacent carbon atoms. Preferred bridges are one or two carbon atoms. It is noted that a bridge always converts a monocyclic ring into a tricyclic ring. When a ring is bridged, the substituents recited for the ring may also be present on the bridge.

As used herein, the term "bicyclic carbocycle" or "bicyclic carbocyclic group" is intended to mean a stable 9- or 10-membered carbocyclic ring system that contains two fused rings and consists of carbon atoms. Of the two fused rings, one ring is a benzo ring fused to a second ring; and the second ring is a 5- or 6-membered carbon ring which is saturated, partially unsaturated, or unsaturated. The bicyclic carbocyclic group may be attached to its pendant group at any carbon atom which results in a stable structure. The bicyclic carbocyclic group described herein may be substituted on any carbon if the resulting compound is stable. Examples of a bicyclic carbocyclic group are, but not limited to, naphthyl, 1,2-dihydronaphthyl, 1,2,3,4-tetrahydronaphthyl, and indanyl.

"Aryl" groups refer to monocyclic or polycyclic aromatic hydrocarbons, including, for example, phenyl, naphthyl, and phenanthranyl. Aryl moieties are well known and described, for example, in Lewis, R.J., ed., *Hawley's Condensed Chemical Dictionary*, 13th Edition, John Wiley & Sons, Inc., New York (1997). "C₆ or C₁₀ aryl" or "C₆₋₁₀ aryl" refers to phenyl and naphthyl. Unless otherwise specified, "aryl", "C₆ or C₁₀ aryl" or "C₆₋₁₀ aryl" or "aromatic residue" may be unsubstituted or substituted with 1 to 5 groups, preferably 1 to 3 groups, OH, OCH₃, Cl, F, Br, I, CN, NO₂, NH₂, N(CH₃)H, N(CH₃)₂, CF₃, OCF₃, C(=O)CH₃, SCH₃, S(=O)CH₃, S(=O)₂CH₃, CH₃, CH₂CH₃, CO₂H, and CO₂CH₃.

The term "benzyl", as used herein, refers to a methyl group on which one of the hydrogen atoms is replaced by a phenyl group, wherein said phenyl group may optionally be substituted with 1 to 5 groups, preferably 1 to 3 groups, OH, OCH₃, Cl, F, Br, I, CN, NO₂, NH₂, N(CH₃)H, N(CH₃)₂, CF₃, OCF₃, C(=O)CH₃, SCH₃, S(=O)CH₃, S(=O)₂CH₃, CH₃, CH₂CH₃, CO₂H, and CO₂CH₃.

As used herein, the term "heterocycle" or "heterocyclic group" is intended to mean a stable 3-, 4-, 5-, 6-, or 7-membered monocyclic or bicyclic or 7-, 8-, 9-, 10-, 11-, 12-, 13-, or 14-membered polycyclic heterocyclic ring that is saturated, partially unsaturated, or fully unsaturated, and that contains carbon atoms and 1, 2, 3 or 4 heteroatoms independently selected from the group consisting of N, O and S; and including any polycyclic group in which any of the above-defined heterocyclic rings is fused to a benzene ring. The nitrogen and sulfur heteroatoms may optionally be oxidized (*i.e.*, N→O and S(O)_p, wherein p is 0, 1 or 2). The nitrogen atom may be substituted or unsubstituted (*i.e.*, N or NR wherein R is H or another substituent, if defined). The heterocyclic ring may be attached to its pendant group at any heteroatom or carbon atom that results in a stable structure. The heterocyclic rings described herein may be substituted on carbon or on a nitrogen atom if the resulting compound is stable. A nitrogen in the heterocycle may optionally be quaternized. It is preferred that when the total number of S and O atoms in the heterocycle exceeds 1, then these heteroatoms are not adjacent to one another. It is preferred that the total number of S and O atoms in the heterocycle is not more than 1. When the term "heterocycle" is used, it is intended to include heteroaryl.

Examples of heterocycles include, but are not limited to, acridinyl, azetidiny, azocinyl, benzimidazolyl, benzofuranyl, benzothiofuranyl, benzothiophenyl,

benzoxazolyl, benzoxazoliny, benzthiazolyl, benztriazolyl, benztetrazolyl,
 benzisoxazolyl, benzisothiazolyl, benzimidazoliny, carbazolyl, 4a*H*-carbazolyl,
 carboliny, chromanyl, chromenyl, cinnoliny, decahydroquinoliny, 2*H*,6*H*-1,5,2-
 dithiaziny, dihydrofuro[2,3-*b*]tetrahydrofuran, furanyl, furazanyl, imidazolidiny,
 5 imidazoliny, imidazolyl, 1*H*-indazolyl, imidazolopyridiny, indolenyl, indoliny,
 indoliziny, indolyl, 3*H*-indolyl, isatinoyl, isobenzofuranyl, isochromanyl, isoindazolyl,
 isoindoliny, isoindolyl, isoquinoliny, isothiazolyl, isothiazolopyridiny, isoxazolyl,
 isoxazolopyridiny, methylenedioxyphenyl, morpholiny, naphthyridiny,
 octahydroisoquinoliny, oxadiazolyl, 1,2,3-oxadiazolyl, 1,2,4-oxadiazolyl, 1,2,5-
 10 oxadiazolyl, 1,3,4-oxadiazolyl, oxazolidiny, oxazolyl, oxazolopyridiny,
 oxazolidinyperimidiny, oxindolyl, pyrimidiny, phenanthridiny, phenanthroliny,
 phenaziny, phenothiaziny, phenoxathiiny, phenoxaziny, phthalaziny, piperaziny,
 piperidiny, piperidonyl, 4-piperidonyl, piperonyl, pteridiny, puriny, pyranly, pyraziny,
 pyrazolidiny, pyrazoliny, pyrazolopyridiny, pyrazolyl, pyridaziny, pyridooxazolyl,
 15 pyridoimidazolyl, pyridothiazolyl, pyridiny, pyrimidiny, pyrrolidiny, pyrroliny, 2-
 pyrrolidonyl, 2*H*-pyrrolyl, pyrrolyl, quinazoliny, quinoliny, 4*H*-quinoliziny,
 quinoxaliny, quinuclidiny, tetrazolyl, tetrahydrofuranyl, tetrahydroisoquinoliny,
 tetrahydroquinoliny, 6*H*-1,2,5-thiadiaziny, 1,2,3-thiadiazolyl, 1,2,4-thiadiazolyl, 1,2,5-
 thiadiazolyl, 1,3,4-thiadiazolyl, thianthrenyl, thiazolyl, thienyl, thiazolopyridiny,
 20 thienothiazolyl, thienooxazolyl, thienoimidazolyl, thiophenyl, triaziny, 1,2,3-triazolyl,
 1,2,4-triazolyl, 1,2,5-triazolyl, 1,3,4-triazolyl, and xanthenyl. Also included are fused ring
 and spiro compounds containing, for example, the above heterocycles.

Examples of 5- to 10-membered heterocycles include, but are not limited to,
 pyridiny, furanyl, thienyl, pyrrolyl, pyrazolyl, pyraziny, piperaziny, piperidiny,
 25 imidazolyl, imidazolidiny, indolyl, tetrazolyl, isoxazolyl, morpholiny, oxazolyl,
 oxadiazolyl, oxazolidiny, tetrahydrofuranyl, thiadiaziny, thiadiazolyl, thiazolyl,
 triaziny, triazolyl, benzimidazolyl, 1*H*-indazolyl, benzofuranyl, benzothiofuranyl,
 benztetrazolyl, benzotriazolyl, benzisoxazolyl, benzoxazolyl, oxindolyl, benzoxazoliny,
 benzthiazolyl, benzisothiazolyl, isatinoyl, isoquinoliny, octahydroisoquinoliny,
 30 tetrahydroisoquinoliny, tetrahydroquinoliny, isoxazolopyridiny, quinazoliny,
 quinoliny, isothiazolopyridiny, thiazolopyridiny, oxazolopyridiny, imidazolopyridiny,
 and pyrazolopyridiny.

Examples of 5- to 6-membered heterocycles include, but are not limited to, pyridinyl, furanyl, thienyl, pyrrolyl, pyrazolyl, pyrazinyl, piperazinyl, piperidinyl, imidazolyl, imidazolidinyl, indolyl, tetrazolyl, isoxazolyl, morpholinyl, oxazolyl, oxadiazolyl, oxazolidinyl, tetrahydrofuranyl, thiadiazinyl, thiadiazolyl, thiazolyl, triazinyl, and triazolyl. Also included are fused ring and spiro compounds containing, for example, the above heterocycles.

As used herein, the term "bicyclic heterocycle" or "bicyclic heterocyclic group" is intended to mean a stable 9- or 10-membered heterocyclic ring system which contains two fused rings and consists of carbon atoms and 1, 2, 3, or 4 heteroatoms independently selected from the group consisting of N, O and S. Of the two fused rings, one ring is a 5- or 6-membered monocyclic aromatic ring comprising a 5-membered heteroaryl ring, a 6-membered heteroaryl ring or a benzo ring, each fused to a second ring. The second ring is a 5- or 6-membered monocyclic ring which is saturated, partially unsaturated, or unsaturated, and comprises a 5-membered heterocycle, a 6-membered heterocycle or a carbocycle (provided the first ring is not benzo when the second ring is a carbocycle).

The bicyclic heterocyclic group may be attached to its pendant group at any heteroatom or carbon atom which results in a stable structure. The bicyclic heterocyclic group described herein may be substituted on carbon or on a nitrogen atom if the resulting compound is stable. It is preferred that when the total number of S and O atoms in the heterocycle exceeds 1, then these heteroatoms are not adjacent to one another. It is preferred that the total number of S and O atoms in the heterocycle is not more than 1.

Examples of a bicyclic heterocyclic group are, but not limited to, quinolinyl, isoquinolinyl, phthalazinyl, quinazolinyl, indolyl, isoindolyl, indolinyl, 1H-indazolyl, benzimidazolyl, 1,2,3,4-tetrahydroquinolinyl, 1,2,3,4-tetrahydroisoquinolinyl, 5,6,7,8-tetrahydro-quinolinyl, 2,3-dihydro-benzofuranyl, chromanyl, 1,2,3,4-tetrahydro-quinoxalanyl, and 1,2,3,4-tetrahydro-quinazolinyl.

As used herein, the term "aromatic heterocyclic group" or "heteroaryl" is intended to mean stable monocyclic and polycyclic aromatic hydrocarbons that include at least one heteroatom ring member such as sulfur, oxygen, or nitrogen. Heteroaryl groups include, without limitation, pyridyl, pyrimidinyl, pyrazinyl, pyridazinyl, triazinyl, furyl, quinolyl, isoquinolyl, thienyl, imidazolyl, thiazolyl, indolyl, pyrrolyl, oxazolyl, benzofuryl, benzothienyl, benzthiazolyl, isoxazolyl, pyrazolyl, triazolyl, tetrazolyl, indazolyl, 1,2,4-

thiadiazolyl, isothiazolyl, purinyl, carbazolyl, benzimidazolyl, indolinyl, benzodioxolanyl, and benzodioxane. Heteroaryl groups are substituted or unsubstituted. The nitrogen atom is substituted or unsubstituted (*i.e.*, N or NR wherein R is H or another substituent, if defined). The nitrogen and sulfur heteroatoms may optionally be oxidized
5 (*i.e.*, N $\rightarrow$ O and S(O)_p, wherein p is 0, 1 or 2).

Bridged rings are also included in the definition of heterocycle. A bridged ring occurs when one or more atoms (*i.e.*, C, O, N, or S) link two non-adjacent carbon or nitrogen atoms. Examples of bridged rings include, but are not limited to, one carbon atom, two carbon atoms, one nitrogen atom, two nitrogen atoms, and a carbon-nitrogen
10 group. It is noted that a bridge always converts a monocyclic ring into a tricyclic ring. When a ring is bridged, the substituents recited for the ring may also be present on the bridge.

The term "counterion" is used to represent a negatively charged species such as chloride, bromide, hydroxide, acetate, and sulfate.

15 When a dotted ring is used within a ring structure, this indicates that the ring structure may be saturated, partially saturated or unsaturated.

As referred to herein, the term "substituted" means that at least one hydrogen atom is replaced with a non-hydrogen group, provided that normal valencies are maintained and that the substitution results in a stable compound. When a substituent is keto (*i.e.*,
20 =O), then 2 hydrogens on the atom are replaced. Keto substituents are not present on aromatic moieties. When a ring system (*e.g.*, carbocyclic or heterocyclic) is said to be substituted with a carbonyl group or a double bond, it is intended that the carbonyl group or double bond be part (*i.e.*, within) of the ring. Ring double bonds, as used herein, are double bonds that are formed between two adjacent ring atoms (*e.g.*, C=C, C=N, or
25 N=N).

In cases wherein there are nitrogen atoms (*e.g.*, amines) on compounds of the present invention, these may be converted to N-oxides by treatment with an oxidizing agent (*e.g.*, mCPBA and/or hydrogen peroxides) to afford other compounds of this invention. Thus, shown and claimed nitrogen atoms are considered to cover both the
30 shown nitrogen and its N-oxide (N $\rightarrow$ O) derivative.

When any variable occurs more than one time in any constituent or formula for a compound, its definition at each occurrence is independent of its definition at every other

occurrence. Thus, for example, if a group is shown to be substituted with 0-3 R groups, then said group may optionally be substituted with up to three R groups, and at each occurrence R is selected independently from the definition of R. Also, combinations of substituents and/or variables are permissible only if such combinations result in stable
5 compounds.

When a bond to a substituent is shown to cross a bond connecting two atoms in a ring, then such substituent may be bonded to any atom on the ring. When a substituent is listed without indicating the atom in which such substituent is bonded to the rest of the compound of a given formula, then such substituent may be bonded via any atom in such
10 substituent. Combinations of substituents and/or variables are permissible only if such combinations result in stable compounds.

The phrase "pharmaceutically acceptable" is employed herein to refer to those compounds, materials, compositions, and/or dosage forms that are, within the scope of sound medical judgment, suitable for use in contact with the tissues of human beings and
15 animals without excessive toxicity, irritation, allergic response, and/or other problem or complication, commensurate with a reasonable benefit/risk ratio.

As used herein, "pharmaceutically acceptable salts" refer to derivatives of the disclosed compounds wherein the parent compound is modified by making acid or base salts thereof. Examples of pharmaceutically acceptable salts include, but are not limited
20 to, mineral or organic acid salts of basic groups such as amines; and alkali or organic salts of acidic groups such as carboxylic acids. The pharmaceutically acceptable salts include the conventional non-toxic salts or the quaternary ammonium salts of the parent compound formed, for example, from non-toxic inorganic or organic acids. For example, such conventional non-toxic salts include those derived from inorganic acids such as
25 hydrochloric, hydrobromic, sulfuric, sulfamic, phosphoric, and nitric; and the salts prepared from organic acids such as acetic, propionic, succinic, glycolic, stearic, lactic, malic, tartaric, citric, ascorbic, pantoic, maleic, hydroxymaleic, phenylacetic, glutamic, benzoic, salicylic, sulfanilic, 2-acetoxybenzoic, fumaric, toluenesulfonic, methanesulfonic, ethane disulfonic, oxalic, and isethionic.

30 The pharmaceutically acceptable salts of the present invention can be synthesized from the parent compound that contains a basic or acidic moiety by conventional chemical methods. Generally, such salts can be prepared by reacting the free acid or base

forms of these compounds with a stoichiometric amount of the appropriate base or acid in water or in an organic solvent, or in a mixture of the two; generally, nonaqueous media like ether, ethyl acetate, ethanol, isopropanol, or acetonitrile are preferred. Lists of suitable salts are found in *Remington's Pharmaceutical Sciences*, 18th Edition, Mack
5 Publishing Company, Easton, PA (1990), the disclosure of which is hereby incorporated by reference.

In addition, compounds of formula I may have prodrug forms. Any compound that will be converted *in vivo* to provide the bioactive agent (*i.e.*, a compound of formula I) is a prodrug within the scope and spirit of the invention. Various forms of prodrugs are well
10 known in the art. For examples of such prodrug derivatives, see:

- a) Bundgaard, H., ed., *Design of Prodrugs*, Elsevier (1985), and Widder, K. et al., eds., *Methods in Enzymology*, 112:309-396, Academic Press (1985);
- b) Bundgaard, H., Chapter 5, "Design and Application of Prodrugs", Krosgaard-Larsen, P. et al., eds., *A Textbook of Drug Design and Development*, pp. 113-
15 191, Harwood Academic Publishers (1991);
- c) Bundgaard, H., *Adv. Drug Deliv. Rev.*, 8:1-38 (1992);
- d) Bundgaard, H. et al., *J. Pharm. Sci.*, 77:285 (1988); and
- e) Kakeya, N. et al., *Chem. Pharm. Bull.*, 32:692 (1984).

20 Compounds containing a carboxy group can form physiologically hydrolyzable esters that serve as prodrugs by being hydrolyzed in the body to yield formula I compounds *per se*. Such prodrugs are preferably administered orally since hydrolysis in many instances occurs principally under the influence of the digestive enzymes. Parenteral administration may be used where the ester *per se* is active, or in those
25 instances where hydrolysis occurs in the blood. Examples of physiologically hydrolyzable esters of compounds of formula I include C₁₋₆alkyl, C₁₋₆alkylbenzyl, 4-methoxybenzyl, indanyl, phthalyl, methoxymethyl, C₁₋₆ alkanoyloxy-C₁₋₆alkyl (*e.g.*, acetoxymethyl, pivaloyloxymethyl or propionyloxymethyl), C₁₋₆alkoxycarbonyloxy-C₁₋₆alkyl (*e.g.*, methoxycarbonyl-oxyethyl or ethoxycarbonyloxymethyl, glycyloxymethyl,
30 phenylglycyloxymethyl, (5-methyl-2-oxo-1,3-dioxolen-4-yl)-methyl), and other well known physiologically hydrolyzable esters used, for example, in the penicillin and

cephalosporin arts. Such esters may be prepared by conventional techniques known in the art.

Preparation of prodrugs is well known in the art and described in, for example, King, F.D., ed., *Medicinal Chemistry: Principles and Practice*, The Royal Society of
5 Chemistry, Cambridge, UK (1994); Testa, B. et al., *Hydrolysis in Drug and Prodrug Metabolism. Chemistry, Biochemistry and Enzymology*, VCHA and Wiley-VCH, Zurich, Switzerland (2003); Wermuth, C.G., ed., *The Practice of Medicinal Chemistry*, Academic Press, San Diego, CA (1999).

The present invention is intended to include all isotopes of atoms occurring in the
10 present compounds. Isotopes include those atoms having the same atomic number but different mass numbers. By way of general example and without limitation, isotopes of hydrogen include deuterium and tritium. Deuterium has one proton and one neutron in its nucleus and that has twice the mass of ordinary hydrogen. Deuterium can be represented by symbols such as "²H" or "D". The term "deuterated" herein, by itself or used to modify
15 a compound or group, refers to replacement of one or more hydrogen atom(s), which is attached to carbon(s), with a deuterium atom. Isotopes of carbon include ¹³C and ¹⁴C.

Isotopically-labeled compounds of the invention can generally be prepared by conventional techniques known to those skilled in the art or by processes analogous to those described herein, using an appropriate isotopically-labeled reagent in place of the
20 non-labeled reagent otherwise employed. Such compounds have a variety of potential uses, e.g., as standards and reagents in determining the ability of a potential pharmaceutical compound to bind to target proteins or receptors, or for imaging compounds of this invention bound to biological receptors *in vivo* or *in vitro*.

"Stable compound" and "stable structure" are meant to indicate a compound that is
25 sufficiently robust to survive isolation to a useful degree of purity from a reaction mixture, and formulation into an efficacious therapeutic agent. It is preferred that compounds of the present invention do not contain a N-halo, S(O)₂H, or S(O)H group.

The term "solvate" means a physical association of a compound of this invention with one or more solvent molecules, whether organic or inorganic. This physical
30 association includes hydrogen bonding. In certain instances the solvate will be capable of isolation, for example when one or more solvent molecules are incorporated in the crystal lattice of the crystalline solid. The solvent molecules in the solvate may be present in a

regular arrangement and/or a non-ordered arrangement. The solvate may comprise either a stoichiometric or nonstoichiometric amount of the solvent molecules. "Solvate" encompasses both solution-phase and isolable solvates. Exemplary solvates include, but are not limited to, hydrates, ethanlates, methanlates, and isopropanolates. Methods of solvation are generally known in the art.

Abbreviations as used herein, are defined as follows: "1 x" for once, "2 x" for twice, "3 x" for thrice, "°C" for degrees Celsius, "eq" for equivalent or equivalents, "g" for gram or grams, "mg" for milligram or milligrams, "L" for liter or liters, "mL" for milliliter or milliliters, "μL" for microliter or microliters, "N" for normal, "M" for molar, "mmol" for millimole or millimoles, "min" for minute or minutes, "h" for hour or hours, "rt" for room temperature, "RT" for retention time, "atm" for atmosphere, "psi" for pounds per square inch, "conc." for concentrate, "sat" or "saturated" for saturated, "MW" for molecular weight, "mp" for melting point, "ee" for enantiomeric excess, "MS" or "Mass Spec" for mass spectrometry, "ESI" for electrospray ionization mass spectroscopy, "HR" for high resolution, "HRMS" for high resolution mass spectrometry, "LCMS" for liquid chromatography mass spectrometry, "HPLC" for high pressure liquid chromatography, "RP HPLC" for reverse phase HPLC, "TLC" or "tlc" for thin layer chromatography, "NMR" for nuclear magnetic resonance spectroscopy, "nOe" for nuclear Overhauser effect spectroscopy, "¹H" for proton, "δ" for delta, "s" for singlet, "d" for doublet, "t" for triplet, "q" for quartet, "m" for multiplet, "br" for broad, "Hz" for hertz, and "α", "β", "R", "S", "E", and "Z" are stereochemical designations familiar to one skilled in the art.

Me	methyl
Et	ethyl
Pr	propyl
<i>i</i> -Pr	isopropyl
Bu	butyl
<i>i</i> -Bu	isobutyl
<i>t</i> -Bu	<i>tert</i> -butyl
Ph	phenyl
Bn	Benzyl
Boc	<i>tert</i> -butoxycarbonyl

AcOH or HOAc	acetic acid
BOP reagent	benzotriazol-1-yloxytris(dimethylamino)phosphonium hexafluorophosphate
CBz	carbobenzyloxy
CH ₂ Cl ₂	dichloromethane
CH ₃ CN or ACN	acetonitrile
CDCl ₃	deutero-chloroform
CHCl ₃	chloroform
DCM	dichloromethane
DEA	diethylamine
DIC or DIPCDI	diisopropylcarbodiimide
DIEA, DIPEA or Hunig's base	diisopropylethylamine
DMAP	4-dimethylaminopyridine
DME	1,2-dimethoxyethane
DMF	dimethyl formamide
DMSO	dimethyl sulfoxide
cDNA	complimentary DNA
EDTA	ethylenediaminetetraacetic acid
Et ₃ N or TEA	triethylamine
EtOAc	ethyl acetate
Et ₂ O	diethyl ether
EtOH	ethanol
HCl	hydrochloric acid
Hex	Hexane
K ₂ CO ₃	potassium carbonate
KOAc	potassium acetate
K ₃ PO ₄	potassium phosphate
LAH	lithium aluminum hydride
LG	leaving group
MeOH	methanol
MgSO ₄	magnesium sulfate

MsOH or MSA	methanesulfonic acid
NaOAc	sodium acetate
NaCl	sodium chloride
NaH	sodium hydride
NaHCO ₃	sodium bicarbonate
Na ₂ CO ₃	sodium carbonate
Na ₂ SO ₄	sodium sulfate
NH ₄ Cl	ammonium chloride
OTf	triflate or trifluoromethanesulfonate
Pd/C	palladium on carbon
PG	protecting group
i-PrOH or IPA	Isopropanol
rt	Room temperature
RT or t _R	retention time
SiO ₂	silica oxide
TEA	triethylamine
TFA	trifluoroacetic acid
THF	tetrahydrofuran
T3P®	propane phosphonic acid anhydride

IV. BIOLOGY

In Vitro Assays

The effectiveness of compounds of the present invention as ROCK inhibitors can be determined in a 30 μ L assay containing 20 mM HEPES, pH 7.5, 20 mM MgCl₂, 0.015% Brij-35, 4 mM DTT, 5 μ M ATP and 1.5 μ M peptide substrate (FITC-AHA-AKRRRLSSLRA-OH) (SEQ ID No. 1). Compounds were dissolved in DMSO so that the final concentration of DMSO was < 2%, and the reaction was initiated with Rho kinase variants. After incubation, the reaction was terminated by the addition of EDTA and the phosphorylated and non-phosphorylated peptides separated using a LABCHIP® 3000 Reader (Caliper Life Sciences). Controls consisted of assays that did not contain compound, and backgrounds consisted of assays that contained enzyme and substrate but had EDTA from the beginning of the reaction to inhibit kinase activity. Compounds were

tested in dose-response format, and the inhibition of kinase activity was calculated at each concentration of compound. The inhibition data were fit using a curve-fitting program to determine the IC₅₀; *i.e.*, the concentration of compound required to inhibit 50% of kinase activity.

5

V. PHARMACEUTICAL COMPOSITIONS, FORMULATIONS AND COMBINATIONS

The compounds of this invention can be administered in such oral dosage forms as tablets, capsules (each of which includes sustained release or timed release formulations), pills, powders, granules, elixirs, tinctures, suspensions, syrups, and emulsions. They may also be administered in intravenous (bolus or infusion), intraperitoneal, subcutaneous, or intramuscular form, all using dosage forms well known to those of ordinary skill in the pharmaceutical arts. They can be administered alone, but generally will be administered with a pharmaceutical carrier selected on the basis of the chosen route of administration and standard pharmaceutical practice.

The term "pharmaceutical composition" means a composition comprising a compound of the invention in combination with at least one additional pharmaceutically acceptable carrier. A "pharmaceutically acceptable carrier" refers to media generally accepted in the art for the delivery of biologically active agents to animals, in particular, mammals, including, *i.e.*, adjuvant, excipient or vehicle, such as diluents, preserving agents, fillers, flow regulating agents, disintegrating agents, wetting agents, emulsifying agents, suspending agents, sweetening agents, flavoring agents, perfuming agents, antibacterial agents, antifungal agents, lubricating agents and dispensing agents, depending on the nature of the mode of administration and dosage forms.

Pharmaceutically acceptable carriers are formulated according to a number of factors well within the purview of those of ordinary skill in the art. These include, without limitation: the type and nature of the active agent being formulated; the patient to which the agent-containing composition is to be administered; the intended route of administration of the composition; and the therapeutic indication being targeted. Pharmaceutically acceptable carriers include both aqueous and non-aqueous liquid media, as well as a variety of solid and semi-solid dosage forms. Such carriers can include a number of different ingredients and additives in addition to the active agent, such additional ingredients being included in

the formulation for a variety of reasons, *e.g.*, stabilization of the active agent, binders, etc., well known to those of ordinary skill in the art. Descriptions of suitable pharmaceutically acceptable carriers, and factors involved in their selection, are found in a variety of readily available sources such as, for example, *Remington's Pharmaceutical Sciences*, 18th Edition (1990).

The dosage regimen for the compounds of the present invention will, of course, vary depending upon known factors, such as the pharmacodynamic characteristics of the particular agent and its mode and route of administration; the species, age, sex, health, medical condition, and weight of the recipient; the nature and extent of the symptoms; the kind of concurrent treatment; the frequency of treatment; the route of administration, the renal and hepatic function of the patient, and the effect desired. A physician or veterinarian can determine and prescribe the effective amount of the drug required to prevent, counter, or arrest the progress of the disorder.

By way of general guidance, the daily oral dosage of each active ingredient, when used for the indicated effects, will range between about 0.001 to about 1000 mg/kg of body weight, preferably between about 0.01 to about 100 mg/kg of body weight per day, and most preferably between about 0.1 to about 20 mg/kg/day. Intravenously, the most preferred doses will range from about 0.001 to about 10 mg/kg/minute during a constant rate infusion. Compounds of this invention may be administered in a single daily dose, or the total daily dosage may be administered in divided doses of two, three, or four times daily.

Compounds of this invention can also be administered by parenteral administration (*e.g.*, intra-venous, intra-arterial, intramuscularly, or subcutaneously). When administered intra-venous or intra-arterial, the dose can be given continuously or intermittent. Furthermore, formulation can be developed for intramuscularly and subcutaneous delivery that ensure a gradual release of the active pharmaceutical ingredient.

Compounds of this invention can be administered in intranasal form via topical use of suitable intranasal vehicles, or via transdermal routes, using transdermal skin patches. When administered in the form of a transdermal delivery system, the dosage administration will, of course, be continuous rather than intermittent throughout the dosage regimen.

The compounds are typically administered in admixture with suitable pharmaceutical diluents, excipients, or carriers (collectively referred to herein as pharmaceutical carriers) suitably selected with respect to the intended form of administration, *e.g.*, oral tablets, capsules, elixirs, and syrups, and consistent with
5 conventional pharmaceutical practices.

For instance, for oral administration in the form of a tablet or capsule, the active drug component can be combined with an oral, non-toxic, pharmaceutically acceptable, inert carrier such as lactose, starch, sucrose, glucose, methyl cellulose, magnesium stearate, dicalcium phosphate, calcium sulfate, mannitol, sorbitol and the like; for oral
10 administration in liquid form, the oral drug components can be combined with any oral, non-toxic, pharmaceutically acceptable inert carrier such as ethanol, glycerol, water, and the like. Moreover, when desired or necessary, suitable binders, lubricants, disintegrating agents, and coloring agents can also be incorporated into the mixture. Suitable binders include starch, gelatin, natural sugars such as glucose or beta-lactose, corn sweeteners,
15 natural and synthetic gums such as acacia, tragacanth, or sodium alginate, carboxymethylcellulose, polyethylene glycol, waxes, and the like. Lubricants used in these dosage forms include sodium oleate, sodium stearate, magnesium stearate, sodium benzoate, sodium acetate, sodium chloride, and the like. Disintegrators include, without limitation, starch, methyl cellulose, agar, bentonite, xanthan gum, and the like.

20 The compounds of the present invention can also be administered in the form of liposome delivery systems, such as small unilamellar vesicles, large unilamellar vesicles, and multilamellar vesicles. Liposomes can be formed from a variety of phospholipids, such as cholesterol, stearylamine, or phosphatidylcholines.

Compounds of the present invention may also be coupled with soluble polymers
25 as targetable drug carriers. Such polymers can include polyvinylpyrrolidone, pyran copolymer, polyhydroxypropylmethacrylamide-phenol, polyhydroxyethylaspartamidephenol, or polyethyleneoxide-polylysine substituted with palmitoyl residues. Furthermore, the compounds of the present invention may be coupled to a class of biodegradable polymers useful in achieving controlled release of a drug, for
30 example, polylactic acid, polyglycolic acid, copolymers of polylactic and polyglycolic acid, polyepsilon caprolactone, polyhydroxy butyric acid, polyorthoesters, polyacetals,

polydihydropyrans, polycyanoacylates, and crosslinked or amphipathic block copolymers of hydrogels.

Dosage forms (pharmaceutical compositions) suitable for administration may contain from about 1 milligram to about 1000 milligrams of active ingredient per dosage unit. In these pharmaceutical compositions the active ingredient will ordinarily be present in an amount of about 0.1-95% by weight based on the total weight of the composition.

Gelatin capsules may contain the active ingredient and powdered carriers, such as lactose, starch, cellulose derivatives, magnesium stearate, stearic acid, and the like. Similar diluents can be used to make compressed tablets. Both tablets and capsules can be manufactured as sustained release products to provide for continuous release of medication over a period of hours. Compressed tablets can be sugar coated or film coated to mask any unpleasant taste and protect the tablet from the atmosphere, or enteric coated for selective disintegration in the gastrointestinal tract.

Liquid dosage forms for oral administration can contain coloring and flavoring to increase patient acceptance.

In general, water, a suitable oil, saline, aqueous dextrose (glucose), and related sugar solutions and glycols such as propylene glycol or polyethylene glycols are suitable carriers for parenteral solutions. Solutions for parenteral administration preferably contain a water soluble salt of the active ingredient, suitable stabilizing agents, and if necessary, buffer substances. Antioxidizing agents such as sodium bisulfite, sodium sulfite, or ascorbic acid, either alone or combined, are suitable stabilizing agents. Also used are citric acid and its salts and sodium EDTA. In addition, parenteral solutions can contain preservatives, such as benzalkonium chloride, methyl- or propyl-paraben, and chlorobutanol.

The compounds of the present invention can be administered alone or in combination with one or more additional therapeutic agents. By "administered in combination" or "combination therapy" it is meant that the compound of the present invention and one or more additional therapeutic agents are administered concurrently to the mammal being treated. When administered in combination, each component may be administered at the same time or sequentially in any order at different points in time. Thus, each component may be administered separately but sufficiently closely in time so as to provide the desired therapeutic effect.

The compounds of the present invention are also useful as standard or reference compounds, for example as a quality standard or control, in tests or assays involving the inhibition of ROCK. Such compounds may be provided in a commercial kit, for example, for use in pharmaceutical research involving ROCK. For example, a compound of the present invention could be used as a reference in an assay to compare its known activity to a compound with an unknown activity. This would ensure the experimenter that the assay was being performed properly and provide a basis for comparison, especially if the test compound was a derivative of the reference compound. When developing new assays or protocols, compounds according to the present invention could be used to test their effectiveness.

The present invention also encompasses an article of manufacture. As used herein, article of manufacture is intended to include, but not be limited to, kits and packages. The article of manufacture of the present invention, comprises: (a) a first container; (b) a pharmaceutical composition located within the first container, wherein the composition, comprises: a first therapeutic agent, comprising: a compound of the present invention or a pharmaceutically acceptable salt form thereof; and, (c) a package insert stating that the pharmaceutical composition can be used for the treatment of a cardiovascular and/or inflammatory disorder (as defined previously). In another embodiment, the package insert states that the pharmaceutical composition can be used in combination (as defined previously) with a second therapeutic agent to treat cardiovascular and/or inflammatory disorder. The article of manufacture can further comprise: (d) a second container, wherein components (a) and (b) are located within the second container and component (c) is located within or outside of the second container. Located within the first and second containers means that the respective container holds the item within its boundaries.

The first container is a receptacle used to hold a pharmaceutical composition. This container can be for manufacturing, storing, shipping, and/or individual/bulk selling. First container is intended to cover a bottle, jar, vial, flask, syringe, tube (*e.g.*, for a cream preparation), or any other container used to manufacture, hold, store, or distribute a pharmaceutical product.

The second container is one used to hold the first container and, optionally, the package insert. Examples of the second container include, but are not limited to, boxes (*e.g.*, cardboard or plastic), crates, cartons, bags (*e.g.*, paper or plastic bags), pouches, and

sacks. The package insert can be physically attached to the outside of the first container via tape, glue, staple, or another method of attachment, or it can rest inside the second container without any physical means of attachment to the first container. Alternatively, the package insert is located on the outside of the second container. When located on the outside of the second container, it is preferable that the package insert is physically attached via tape, glue, staple, or another method of attachment. Alternatively, it can be adjacent to or touching the outside of the second container without being physically attached.

The package insert is a label, tag, marker, etc. that recites information relating to the pharmaceutical composition located within the first container. The information recited will usually be determined by the regulatory agency governing the area in which the article of manufacture is to be sold (e.g., the United States Food and Drug Administration). Preferably, the package insert specifically recites the indications for which the pharmaceutical composition has been approved. The package insert may be made of any material on which a person can read information contained therein or thereon. Preferably, the package insert is a printable material (e.g., paper, plastic, cardboard, foil, adhesive-backed paper or plastic, etc.) on which the desired information has been formed (e.g., printed or applied).

Other features of the invention will become apparent in the course of the following descriptions of exemplary embodiments that are given for illustration of the invention and are not intended to be limiting thereof. The following Examples have been prepared, isolated and characterized using the methods disclosed herein.

VI. GENERAL SYNTHESIS INCLUDING SCHEMES

The compounds of the present invention may be synthesized by methods available to those skilled in the art of organic chemistry (Maffrand, J.P. et al., *Heterocycles*, 16(1):35-37 (1981)). General synthetic schemes for preparing compounds of the present invention are described below. These schemes are illustrative and are not meant to limit the possible techniques one skilled in the art may use to prepare the compounds disclosed herein. Different methods to prepare the compounds of the present invention will be evident to those skilled in the art. Additionally, the various steps in the synthesis may be performed in an alternate sequence in order to give the desired compound or compounds.

Examples of compounds of the present invention prepared by methods described in the general schemes are given in the intermediates and examples section set out hereinafter. Preparation of homochiral examples may be carried out by techniques known to one skilled in the art. For example, homochiral compounds may be prepared by separation of racemic products by chiral phase preparative HPLC. Alternatively, the example compounds may be prepared by methods known to give enantiomerically enriched products. These include, but are not limited to, the incorporation of chiral auxiliary functionalities into racemic intermediates which serve to control the diastereoselectivity of transformations, providing enantio-enriched products upon cleavage of the chiral auxiliary.


The compounds of the present invention can be prepared in a number of ways known to one skilled in the art of organic synthesis. The compounds of the present invention can be synthesized using the methods described below, together with synthetic methods known in the art of synthetic organic chemistry, or by variations thereon as appreciated by those skilled in the art. Preferred methods include, but are not limited to, those described below. The reactions are performed in a solvent or solvent mixture appropriate to the reagents and materials employed and suitable for the transformations being effected. It will be understood by those skilled in the art of organic synthesis that the functionality present on the molecule should be consistent with the transformations proposed. This will sometimes require a judgment to modify the order of the synthetic steps or to select one particular process scheme over another in order to obtain a desired compound of the invention.

It will also be recognized that another major consideration in the planning of any synthetic route in this field is the judicious choice of the protecting group used for protection of the reactive functional groups present in the compounds described in this invention. An authoritative account describing the many alternatives to the trained practitioner is Greene et al., (*Protective Groups in Organic Synthesis*, 4th Edition, Wiley-Interscience (2006)).

Representative 2,8-diazaspiro[4.5]decan-1-one compounds **1g** of this invention can be prepared as shown in Scheme 1. Starting from tert-butyl 1-oxo-2,8-diazaspiro[4.5]decane-8-carboxylate **1a**, coupling with **1b** using Xanphos, Pd₂(dba)₃ and


Cs_2CO_3 gives **1c**. Suzuki coupling with 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazole using XPhos Generation 3 catalyst gives **1d**. Then the protecting Boc group is selectively removed followed by coupling with either acetyl chlorides or carboxylic acids to yield **1f**. The protecting SEM group is then removed to give **1g**.

Scheme 1



Representative 2,8-diazaspiro[4.5]decan-1-one compounds **2c**, **2d** and **2e** of this invention can also be prepared as shown in Scheme 2 in which the Suzuki coupling reaction was performed first. Starting from **1c** shown in Scheme 1, Suzuki coupling with 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-pyrazole using XPhos Generation 3 catalyst gives **2a**, after which protecting Boc group is removed to yield **2b**. Then **2b** can be coupled with either acetyl chlorides or carboxylic acids to produce amides **2c**, or reacted with isocyanides to give urea **2d**, or coupled with sulfonyl chlorides to give sulfonamides **2e**.

Scheme 2




5

Representative 2,8-diazaspiro[4.5]decan-1-one compounds **3c** of this invention can also be prepared as shown in Scheme 3 in which the Suzuki coupling of the hinge aryl group was performed in the last step of the synthesis. Deprotecting Boc group of **1c**

yields **3a**, which can be coupled with either acetyl chlorides or carboxylic acids to give **3b**. Suzuki coupling of **3b** with aryl boronic acids or boronic esters to give **3c**.


Scheme 3

5



Representative 2-(3-methoxy-4-(1H-pyrazol-4-yl)phenyl)-8-(4-(piperazin-1-yl)benzoyl)-2,8-diazaspiro[4.5]decan-1-one compounds **4e** of this invention can be prepared as shown in Scheme 4. Coupling of **4a** and **4b** followed by deprotecting Boc group produces **4d**. Then **4e** can be synthesized via reductive-amination of **4d** with aldehydes and ketones.

Scheme 4



5 Purification method A:

The crude material was purified via preparative LC/MS with the following conditions:

Column: XBridge C18, 19 x 200 mm, 5- μ m particles;

Mobile Phase A: 5:95 acetonitrile: water with 10-mM ammonium acetate; Mobile Phase

B: 95:5 acetonitrile: water with 10-mM ammonium

10 acetate; Gradient: 15-55% B over 20 minutes, then a 5-minute hold at 100% B; Flow: 20 mL/min. Fractions containing the desired product were combined and dried via centrifugal evaporation.

Purification method B:

15 The crude material was purified via preparative LC/MS with the following conditions:

Column: XBridge C18, 19 x 200 mm, 5- μ m particles;

Mobile Phase A: 5:95 acetonitrile: water with 10-mM ammonium acetate; Mobile Phase B: 95:5 acetonitrile: water with 10-mM ammonium acetate; Gradient: 15-100% B over 20 minutes, then a 2-minute hold at 100% B; Flow: 20 mL/min. Fractions containing the desired

5 product were combined and dried via centrifugal evaporation.

Purification method C:

The crude material was purified via preparative LC/MS with the following conditions:

Column: XBridge C18, 19 x 200 mm, 5- μ m particles;

10 Mobile Phase A: 5:95 acetonitrile: water with 10-mM ammonium acetate; Mobile Phase B: 95:5 acetonitrile: water with 10-mM ammonium acetate; Gradient: 0-20% B over 19 minutes, then a 5-minute hold at 100% B; Flow: 20 mL/min. Fractions containing the desired product were combined and dried via centrifugal evaporation.

15


Analytical Retention Time Method 1: Column: Waters Acquity UPLC BEH C18, 2.1 x 50 mm, 1.7- μ m particles; Mobile Phase A: 5:95 acetonitrile:water with 10 mM ammonium acetate; Mobile Phase B: 95:5 acetonitrile:water with 10 mM ammonium acetate; Temperature: 50 °C; Gradient: 0-100% B over 3 minutes, then a 0.75-minute hold at 100% B; Flow: 1.0 mL/min; Detection: UV at 220 nm.

20

Analytical Retention Time Method 2: Column: Waters Acquity UPLC BEH C18, 2.1 x 50 mm, 1.7- μ m particles; Mobile Phase A: 5:95 acetonitrile:water with 0.1% trifluoroacetic acid; Mobile Phase B: 95:5 acetonitrile:water with 0.1% trifluoroacetic acid; Temperature: 50 °C; Gradient: 0-100% B over 3 minutes, then a 0.75-minute hold at 100% B; Flow: 1.0 mL/min; Detection: UV at 220 nm.


25

Intermediate 1: Preparation of tert-butyl 2-(4-bromophenyl)-1-oxo-2,8-diazaspiro[4.5]decane-8-carboxylate.



In a sealable reaction tube was added 1-bromo-4-iodobenzene (1112 mg, 3.93 mmol), tert-butyl 1-oxo-2,8-diazaspiro[4.5]decane-8-carboxylate (500 mg, 1.966 mmol), cesium carbonate (1281 mg, 3.93 mmol) and dioxane (3932 μ l). The reaction was purged
 5 with nitrogen. Then 9,9-dimethyl-4,5-bis(diphenylphosphino)xanthene (171 mg, 0.295 mmol) and $\text{Pd}_2(\text{dba})_3$ (90 mg, 0.098 mmol) was added and nitrogen gas was bubbled through reaction for 1 min. The reaction was sealed and stirred at 100 °C overnight. The reaction was partitioned between EtOAc (50 ml) and water (30 ml). The organic layer was separated, washed with water (2 x 30 ml) and brine (30 ml), dried over MgSO_4 ,
 10 filtered and concentrated. The residue was purified using ISCO system (0-100% EtOAc/Hex gradient) to give tert-butyl 2-(4-bromophenyl)-1-oxo-2,8-diazaspiro[4.5]decane-8-carboxylate (740 mg, 1.663 mmol, 85 % yield) as a beige solid. ^1H NMR (500MHz, DMSO-d_6) δ 7.72 - 7.64 (m, 2H), 7.61 - 7.51 (m, 2H), 3.91 - 3.83 (m, 2H), 3.80 (t, $J=7.0$ Hz, 2H), 2.98 (br. s., 2H), 2.09 (t, $J=7.0$ Hz, 2H), 1.62 (td, $J=12.4$,
 15 4.4 Hz, 2H), 1.54 - 1.47 (m, 2H), 1.42 (s, 9H). ^{13}C NMR (500MHz, DMSO-d_6) δ 177.4, 154.4, 139.4, 131.9, 121.8, 116.3, 79.2, 44.9, 44.7, 32.1, 28.6, 28.0 MS (ESI) m/z : 411.0 ($\text{M}+\text{H}$) $^+$.


20 **Intermediate 2 : Preparation of tert-butyl 2-(4-bromo-3-methoxyphenyl)-1-oxo-2,8-diazaspiro[4.5]decane-8-carboxylate.**




In a sealable reaction tube was added 1-bromo-4-iodo-2-methoxybenzene (2461 mg, 7.86 mmol), tert-butyl 1-oxo-2,8-diazaspiro[4.5]decane-8-carboxylate (1000 mg,

3.93 mmol), cesium carbonate (2562 mg, 7.86 mmol) and dioxane (7864 μ l). The reaction was purged with nitrogen. Then 9,9-dimethyl-4,5-bis(diphenylphosphino)xanthene (341 mg, 0.590 mmol), $\text{Pd}_2(\text{dba})_3$ (180 mg, 0.197 mmol) was added and nitrogen was bubbled through reaction for 1 min. The reaction was sealed and stirred at 100 °C overnight. The reaction was partitioned between EtOAc (50 ml) and water (30 ml). The organic layer was separated, washed with water (2 x 30 ml) and brine (30 ml), dried over MgSO_4 , filtered and concentrated. The residue was purified using ISCO system (0-100% EtOAc/Hex gradient) to give tert-butyl 2-(4-bromo-3-methoxyphenyl)-1-oxo-2,8-diazaspiro[4.5]decane-8-carboxylate (1.6 g, 3.64 mmol, 93 % yield) as a beige solid. ^1H NMR (500 MHz, $\text{DMSO}-d_6$) δ 7.67 (d, $J=2.2$ Hz, 1H), 7.55 (d, $J=8.5$ Hz, 1H), 7.12 (dd, $J=8.8, 2.5$ Hz, 1H), 3.96 - 3.78 (m, 7H), 3.10 - 2.84 (m, 2H), 2.10 (t, $J=6.9$ Hz, 2H), 1.62 (dd, $J=12.0, 4.0$ Hz, 2H), 1.51 (d, $J=13.5$ Hz, 2H), 1.42 (s, 9H). MS (ESI) m/z : 439.0, 441.0 ($\text{M}+\text{H}$) $^+$.

15 **Intermediate 3 : Preparation of 2-(4-(1H-pyrazol-4-yl)phenyl)-2,8-diazaspiro[4.5]decan-1-one, TFA**




3A. Preparation of tert-butyl 1-oxo-2-(4-(1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazol-4-yl)phenyl)-2,8-diazaspiro[4.5]decane-8-carboxylate.




To a microwave vial was added Intermediate 1 (1000 mg, 2.443 mmol), 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazole (951 mg, 2.93 mmol), 3M tripotassium phosphate (2.443 mL, 7.33 mmol), dioxane (10 mL) and methanesulfonato(2-dicyclohexylphosphino-2',4',6'-tri-*i*-propyl-1,1'-biphenyl)(2'-amino-1,1'-biphenyl-2-yl)palladium(II) (103 mg, 0.122 mmol). The reaction was purged with argon and sealed. The reaction was then subjected to microwave oven and stirred at 120 °C for 45 min. The reaction was partitioned between EtOAc (30 ml) and water (20 ml). The organic layer was separated, washed with water (15 ml) and brine (15 ml), dried over MgSO₄, filtered and concentrated. The residue was purified using ISCO system (0-100% EtOAc/Hex gradient) to give tert-butyl 1-oxo-2-(4-(1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazol-4-yl)phenyl)-2,8-diazaspiro[4.5]decane-8-carboxylate (760 mg, 1.443 mmol, 59.1 % yield) as an off-white solid. MS (ESI) *m/z*: 527.3 (M+H)⁺.

3B. Preparation of 2-(4-(1H-pyrazol-4-yl)phenyl)-2,8-diazaspiro[4.5]decan-1-one, TFA




To a round bottom flask was added intermediate 3A (20 mg, 0.038 mmol), CH₂Cl₂ (0.5 mL) and TFA (0.5 mL). The reaction was stirred at rt for 3 hr. The reaction was concentrated and purified using prep-HPLC. The desired fraction was concentrated to give 2-(4-(1H-pyrazol-4-yl)phenyl)-2,8-diazaspiro[4.5]decan-1-one, TFA (10 mg, 0.024 mmol, 62.9 % yield) as a beige solid. ¹H NMR (500MHz, METHANOL-d₄) δ 7.99 (br. s., 2H), 7.64 (s, 4H), 3.94 (q, *J*=6.2 Hz, 2H), 3.61 - 3.50 (m, 2H), 3.26 - 3.15 (m, 2H), 2.28 - 2.21 (m, 2H), 2.19 - 2.07 (m, 2H), 1.95 - 1.76 (m, 2H); MS (ESI) *m/z*: 297.1 (M+H)⁺. ROCK2 IC₅₀=1010nM

Intermediate 4 : Preparation of 2-(3-methoxy-4-(1H-pyrazol-4-yl)phenyl)-2,8-diazaspiro[4.5]decan-1-one, 2 HCl



5

4A. Preparation of tert-butyl 2-(4-(1-(tert-butoxycarbonyl)-1H-pyrazol-4-yl)-3-methoxyphenyl)-1-oxo-2,8-diazaspiro[4.5]decane-8-carboxylate.




10 To a round bottle flask was added Intermediate 2 (1.8 g, 4.10 mmol), tert-butyl 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-pyrazole-1-carboxylate (1.808 g, 6.15 mmol), 3M tripotassium phosphate (4.10 mL, 12.29 mmol), dioxane (10 mL) and chloro(2-dicyclohexylphosphino-2',4',6'-triisopropyl-1,1'-biphenyl)[2-(2'-amino-1,1'-biphenyl)]palladium(II) (0.226 g, 0.287 mmol). The reaction was purged with nitrogen

15 then stirred at 80 °C for 4hr. The reaction was partitioned between EtOAc (50 mL) and water (20 mL). The organic layer was separated, washed with water (20 mL) and brine (30 mL), dried over MgSO₄, filtered and concentrated. The residue was purified using ISCO system (0-100% EtOAc/Hex gradient) to give tert-butyl 2-(4-(1-(tert-butoxycarbonyl)-1H-pyrazol-4-yl)-3-methoxyphenyl)-1-oxo-2,8-diazaspiro[4.5]decane-8-carboxylate (1.98 g, 3.76 mmol, 92 % yield) as an off-white solid. ¹H NMR (500MHz, DMSO-d₆) δ 8.54 (d, *J*=0.6 Hz, 1H), 8.30 (d, *J*=0.8 Hz, 1H), 7.74 (d, *J*=8.5 Hz, 1H), 7.67 (d, *J*=2.2 Hz, 1H), 7.21 (dd, *J*=8.5, 2.2 Hz, 1H), 3.91 (s, 3H), 3.88 - 3.83 (m, 3H), 3.10 - 2.88 (m, 2H), 2.11

20


(t, $J=7.0$ Hz, 2H), 1.68 - 1.62 (m, 2H), 1.61 (s, 9H), 1.51 (d, $J=13.2$ Hz, 2H), 1.42 (s, 9H);
MS (ESI) m/z : 527.2 (M+H)⁺.

5 4B. Preparation of 2-(3-methoxy-4-(1H-pyrazol-4-yl)phenyl)-2,8-diazaspiro[4.5]decan-1-one, 2 HCl




To a round bottom flask was added Intermediate 4A (1.98 g, 3.76 mmol), dioxane (10 mL) and 4N HCl (2.285 mL, 75 mmol) in dioxane. The reaction was stirred at rt overnight. The reaction was concentrated to give 2-(3-methoxy-4-(1H-pyrazol-4-yl)phenyl)-2,8-diazaspiro[4.5]decan-1-one, 2 HCl (1.5 g, 3.76 mmol, 100 % yield) as an
10 off-white solid. MS (ESI) m/z : 327.0 (M+H)⁺.

Intermediate 5 : Preparation of 2-(3-methoxy-4-(1H-pyrazol-4-yl)phenyl)-8-(4-(piperazin-1-yl)benzoyl)-2,8-diazaspiro[4.5]decan-1-one, 2 HCl




15

5A. Preparation of tert-butyl 4-(4-(2-(3-methoxy-4-(1H-pyrazol-4-yl)phenyl)-1-oxo-2,8-diazaspiro[4.5]decane-8-carbonyl)phenyl)piperazine-1-carboxylate.




To a 1-dram vial was added 4-(4-(tert-butoxycarbonyl)piperazin-1-yl)benzoic acid (345 mg, 1.127 mmol), THF (5 mL), HATU (514 mg, 1.352 mmol) and Et₃N (0.785 mL, 5.63 mmol). The reaction was stirred at rt for 10 min and then Intermediate 4 (450 mg, 1.127 mmol) was added and the reaction was continued for 4 hr. The reaction was partitioned between EtOAc (20 ml) and water (15 ml). The organic layer was separated, washed with brine (15 ml), dried over MgSO₄, filtered and concentrated. The residue was purified using ISCO system (0-100% EtOAc/Hex gradient, then 0-20% MeOH/CH₂Cl₂ gradient) to give tert-butyl 4-(4-(2-(3-methoxy-4-(1H-pyrazol-4-yl)phenyl)-1-oxo-2,8-diazaspiro[4.5]decan-8-carbonyl)phenyl)piperazine-1-carboxylate (500 mg, 0.813 mmol, 72.2 % yield) as a light beige solid. ¹H NMR (500MHz, DMSO-d₆) δ 8.09 (br. s., 1H), 7.93 (br. s., 1H), 7.64 - 7.57 (m, 2H), 7.31 (d, *J*=8.8 Hz, 2H), 7.15 (dd, *J*=8.4, 2.1 Hz, 1H), 6.98 (d, *J*=8.8 Hz, 2H), 3.92 - 3.83 (m, 5H), 3.52 - 3.42 (m, 4H), 3.28 (s, 4H), 3.22 - 3.19 (m, 4H), 2.15 (t, *J*=6.9 Hz, 2H), 1.78 - 1.67 (m, 2H), 1.57 (d, *J*=12.9 Hz, 2H), 1.43 (s, 9H); MS (ESI) *m/z*: 615.1 (M+H)⁺.

5B. Preparation of 2-(3-methoxy-4-(1H-pyrazol-4-yl)phenyl)-8-(4-(piperazin-1-yl)benzoyl)-2,8-diazaspiro[4.5]decan-1-one, 2 HCl




To a round bottom flask was added Intermediate 5A (380 mg, 0.618 mmol), dioxane (3 mL) and 4N HCl (0.019 mL, 0.618 mmol) in dioxane. The reaction was stirred at rt for 4hr. The reaction was concentrated to give 2-(3-methoxy-4-(1H-pyrazol-4-yl)phenyl)-8-(4-(piperazin-1-yl)benzoyl)-2,8-diazaspiro[4.5]decan-1-one, 2 HCl (350 mg, 0.596 mmol, 96 % yield) as a white solid. MS (ESI) m/z : 515.0 (M+H)⁺.

Intermediate 6 : Preparation of 2-(4-bromophenyl)-8-(4-(4-methylpiperazin-1-yl)benzoyl)-2,8-diazaspiro[4.5]decan-1-one.




10

6A. Preparation of 2-(4-bromophenyl)-2,8-diazaspiro[4.5]decan-1-one, HCl.




To a round bottom flask was added Intermediate 1 (630 mg, 1.539 mmol), dioxane (5 ml) and 4N HCl (7696 μ l, 30.8 mmol) in dioxane. The reaction was stirred at rt for 3 hr. The reaction was concentrated to give 2-(4-bromophenyl)-2,8-diazaspiro[4.5]decan-1-one, HCl (530 mg, 1.533 mmol, 100 % yield) as a light colored solid. MS (ESI) m/z : 308.8, 310.8 ($M+H$)⁺.

6B. Preparation of 2-(4-bromophenyl)-8-(4-(4-methylpiperazin-1-yl)benzoyl)-2,8-diazaspiro[4.5]decan-1-one.




To a round bottom flask was added 4-(4-methylpiperazin-1-yl)benzoic acid (147 mg, 0.665 mmol), THF (10 mL), HATU (304 mg, 0.798 mmol) and Et₃N (0.464 mL, 3.33 mmol). The reaction was stirred at rt for 10 min and then 6A (230 mg, 0.665 mmol) was added and the reaction was continued for 2 hr. The reaction was concentrated and the residue was purified using ISCO system (0-15% MeOH/CH₂Cl₂ gradient) to give 2-(4-bromophenyl)-8-(4-(4-methylpiperazin-1-yl)benzoyl)-2,8-diazaspiro[4.5]decan-1-one (305 mg, 0.596 mmol, 90 % yield) as an off-white solid. ¹H NMR (500MHz, DMSO-d₆) δ 7.71 - 7.63 (m, 2H), 7.59 - 7.53 (m, 2H), 7.31 (d, J =8.8 Hz, 2H), 6.99 (d, J =8.8 Hz, 2H), 4.17 - 3.85 (m, 2H), 3.82 (t, J =6.9 Hz, 2H), 3.35 - 3.28 (m, 7H), 3.18 (br. s., 2H), 2.84 - 2.66 (m, 2H), 2.44 (br. s., 2H), 2.14 (t, J =6.9 Hz, 2H), 1.77 - 1.65 (m, 2H), 1.57 (d, J =12.7 Hz, 2H); MS (ESI) m/z : 511.0, 513.0 ($M+H$)⁺.

Intermediate 7 : Preparation of 2-(4-(1H-pyrazol-4-yl)phenyl)-2,7-diazaspiro[3.5]nonan-1-one, 2 HCl



- 2-(4-(1H-pyrazol-4-yl)phenyl)-2,7-diazaspiro[3.5]nonan-1-one, 2 HCl (20 mg, 0.056) was prepared in a similar manner as the procedure described in Intermediate 3, using tert-butyl 1-oxo-2,7-diazaspiro[3.5]nonane-7-carboxylate (150mg, 0.624 mmol);
- 5 MS (ESI) m/z : 283.0 (M+H)⁺.


Intermediate 8 : Preparation of 2-(3-methoxy-4-(1H-pyrazol-4-yl)phenyl)-2,8-diazaspiro[4.5]decan-3-one, 2 HCl




- 10 2-(3-methoxy-4-(1H-pyrazol-4-yl)phenyl)-2,8-diazaspiro[4.5]decan-3-one, 2 HCl (70 mg, 0.167 mmol) was prepared in a similar manner as the procedure described in Intermediate 4, using tert-butyl 3-oxo-2,8-diazaspiro[4.5]decane-8-carboxylate (150mg, 0.59 mmol). MS (ESI) m/z : 327.1 (M+H)⁺.

15 **Examples:**

Example 1: Preparation of 8-benzoyl-2-[4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one




1A: Preparation of 8-benzoyl-2-(4-bromophenyl)-2,8-diazaspiro[4.5]decan-1-one




To a round bottom flask was added Intermediate 3 (38 mg, 0.110 mmol), THF (2 mL), Hunig's base (0.115 mL, 0.660 mmol) and benzoyl chloride (23.18 mg, 0.165 mmol). The reaction was stirred at rt for 15 min. The reaction was partitioned between EtOAc (30 ml) and water (20 ml). The organic layer was separated, washed with water (20 ml) and brine (20 ml), dried over MgSO₄, filtered and concentrated. The residue was purified using ISCO system (0-100% EtOAc/Hex gradient) to give 8-benzoyl-2-(4-bromophenyl)-2,8-diazaspiro[4.5]decan-1-one (45 mg, 0.109 mmol, 99 % yield) as an off-white solid. ¹H NMR (500MHz, DMSO-d₆) δ 7.72 - 7.65 (m, 2H), 7.60 - 7.54 (m, 2H), 7.50 - 7.43 (m, 3H), 7.42 - 7.37 (m, 2H), 3.82 (br. s., 2H), 3.27 - 3.10 (m, 2H), 2.14 (br. s., 2H), 1.81 - 1.45 (m, 4H); MS (ESI) *m/z*: 412.9, 414.9 (M+H)⁺.

1B: Preparation of 8-benzoyl-2-[4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one




To a microwave vial was added 1A (45 mg, 0.109 mmol), 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-pyrazole (42.3 mg, 0.218 mmol), tripotassium phosphate (0.145 mL, 0.436 mmol), Dioxane (2 mL) and methanesulfonato(2-dicyclohexylphosphino-2',4',6'-tri-*i*-propyl-1,1'-biphenyl)(2'-amino-1,1'-biphenyl-2-yl)palladium(II) (9.22 mg, 10.89 μ mol). The reaction was purged with argon and sealed. The reaction was then subjected to microwave oven and stirred at 120 °C for 35 min. The reaction was partitioned between EtOAc(10 ml) and water (5 ml). The organic layer was separated, washed with brine (5 ml), dried and concentrated. The residue was purified using purification Method A to give 8-benzoyl-2-[4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one (6.6 mg, 0.016mmol, 14.8% yield). ¹H NMR (500MHz, DMSO-*d*₆) δ 8.11 (br. s., 1H), 7.91 (br. s., 1H), 7.69 - 7.56 (m, 4H), 7.51 - 7.36 (m, 5H), 3.82 (br. s., 2H), 3.52 (br. s., 2H), 3.33 - 3.05 (m, 2H), 2.14 (br. s., 2H), 1.90 - 1.40 (m, 4H); MS (ESI) *m/z*: 400.8 (M+H)⁺; Anal. HPLC Retention time: 1.3(Method 1); ROCK 2 IC₅₀=326nM.

Example 2: Preparation of 8-(benzenesulfonyl)-2-[4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one




2A: Preparation of 2-(4-bromophenyl)-8-(phenylsulfonyl)-2,8-diazaspiro[4.5]decan-1-one



5 To a round bottom flask was added Intermediate 3 (38 mg, 0.110 mmol), THF (2 mL), Hunig's Base (0.115 mL, 0.660 mmol) and benzenesulfonyl chloride (29.1 mg, 0.165 mmol). The reaction was stirred at rt for 15 min. The reaction was partitioned between EtOAc (30 ml) and water (20 ml). The organic layer was separated, washed with water (20 ml) and brine (20 ml), dried over MgSO₄, filtered and concentrated. The
10 residue was purified using ISCO system (0-100% EtOAc/Hex gradient) to give 2-(4-bromophenyl)-8-(phenylsulfonyl)-2,8-diazaspiro[4.5]decan-1-one (38 mg, 0.085 mmol, 77 % yield) as an off-white solid. MS (ESI) *m/z*: 448.9, 450.9 (M+H)⁺.


2B: Preparation of 8-(benzenesulfonyl)-2-[4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one

15




To a microwave vial was added 2A (37 mg, 0.082 mmol), 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-pyrazole (32.0 mg, 0.165 mmol), 3M tripotassium phosphate (0.110 mL, 0.329 mmol), Dioxane (2 mL) and methanesulfonato(2-dicyclohexylphosphino-2',4',6'-tri-*i*-propyl-1,1'-biphenyl)(2'-amino-1,1'-biphenyl-2-yl)palladium(II) (6.97 mg, 8.23 μ mol). The reaction was purged with argon and sealed. The reaction was then subjected to microwave oven and stirred at 120 °C for 35 min. The reaction was partitioned between EtOAc(10 ml) and water (5 ml). The organic layer was separated, washed with brine (5 ml), dried over MgSO₄ and concentrated. The residue purified using purification Method A to give 8-(benzenesulfonyl)-2-[4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one (6.8 mg, 0.015 mmol, 18.8% yield). ¹H NMR (500 MHz, DMSO-*d*₆) δ 8.16 - 8.01 (m, 1H), 8.00 - 7.84 (m, 1H), 7.79 - 7.71 (m, 3H), 7.69 - 7.64 (m, 2H), 7.58 (s, 4H), 3.72 (br t, *J*=6.9 Hz, 2H), 3.51 (br d, *J*=12.1 Hz, 2H), 2.66 - 2.54 (m, 2H), 1.86 (br t, *J*=6.8 Hz, 2H), 1.82 - 1.73 (m, 2H), 1.61 (br d, *J*=13.4 Hz, 2H); MS (ESI) *m/z*: 437.2 (M+H)⁺; Anal. HPLC Retention time: 1.51 (Method 1); ROCK2 IC₅₀=330nM.

Example 3: Preparation of 8-(2-methoxybenzoyl)-2-[4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one.




3A: Preparation of 2-(4-(1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazol-4-yl)phenyl)-2,8-diazaspiro[4.5]decan-1-one




- 5 To a microwave vial was added Intermediate 3A (595 mg, 1.130 mmol), MeOH (8 mL) and H₂O (45 mL). The reaction was stirred at 150 °C for 90 min. The reaction was extraced with EtOAc (3x 20 ml). The combined EtOAc was dried over MgSO₄, filtered and concentrated to give 2-(4-(1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazol-4-yl)phenyl)-2,8-diazaspiro[4.5]decan-1-one (295 mg, 0.691 mmol, 61.2 % yield) as an off-
10 white solid. (ESI) *m/z*: 427.1 (M+H)⁺.

3B: Preparation of 8-(2-methoxybenzoyl)-2-[4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one.




To a round bottom flask was added 3A (25 mg, 0.059 mmol), THF (1 mL), 2-methoxybenzoyl chloride (11.00 mg, 0.064 mmol) and Et₃N (0.012 mL, 0.088 mmol). The reaction was stirred at rt for 1 hr. The reaction was concentrated and then CH₂Cl₂ (0.5 mL) and TFA (0.5 mL) was added and then the reaction was stirred at rt overnight. The reaction was concentrated and the residue was purified using purification Method A to give 8-(2-methoxybenzoyl)-2-[4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one (11.6 mg, 0.026 mmol, 45.1% yield). ¹H NMR (500 MHz, DMSO-d₆) δ 8.24 - 7.85 (m, 2H), 7.75 - 7.55 (m, 4H), 7.40 (br s, 1H), 7.29 - 6.92 (m, 3H), 4.49 - 4.28 (m, 1H), 3.91 - 3.76 (m, 4H), 3.33 (br d, *J*=8.8 Hz, 1H), 3.21 - 2.99 (m, 2H), 2.24 - 2.01 (m, 2H), 1.83 - 1.39 (m, 4H); MS (ESI) *m/z*: 431.3 (M+H)⁺; Anal. HPLC Retention time: 1.43 (Method 1); ROCK2 IC₅₀=160nM

Example 4: Preparation of 8-(4-methoxybenzoyl)-2-[4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one



8-(4-methoxybenzoyl)-2-[4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one (12.2 mg, 0.027mmol, 46.9% yield) was prepared in a similar manner as the procedure described in Example 3, using 4-methoxybenzoyl chloride (11.00 mg, 0.064 mmol). ¹H NMR (500 MHz, DMSO-d₆) δ 8.24 - 7.82 (m, 2H), 7.68 - 7.57 (m, 4H), 7.37 (d, J=8.5 Hz, 2H), 6.99 (d, J=8.4 Hz, 2H), 3.82 (br. s., 2H), 3.78 (s, 3H), 3.69 - 3.60 (m, 2H), 2.55 (m, 2H), 2.13 (br. s., 2H), 1.70 (br. s., 2H), 1.56 (br. s., 2H); MS (ESI) *m/z*: 431.3 (M+H)⁺; Anal. HPLC Retention time: 1.32 (Method 2); ROCK2 IC₅₀=21nM.

10 **Example 5: Preparation of 8-(3-methoxybenzoyl)-2-[4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one**




8-(3-methoxybenzoyl)-2-[4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one (9.6 mg, 0.021mmol, 36.5% yield) was prepared in a similar manner as the procedure described in Example 3, using 3-methoxybenzoyl chloride (11.00 mg, 0.064 mmol). ¹H NMR (500 MHz, DMSO-d₆) δ 8.11 (br. s., 1H), 7.94 (br. s., 1H), 7.72 - 7.55 (m, 4H),

7.37 (t, $J=7.9$ Hz, 1H), 7.02 (dd, $J=8.2, 2.1$ Hz, 1H), 6.96 - 6.88 (m, 2H), 3.82 (br. s., 2H), 3.78 (s, 3H), 3.58 - 3.52 (m, 2H), 3.31 - 3.03 (m, 2H), 2.13 (br. s., 2H), 1.66 (br. s., 3H), 1.51 (br. s., 1H); MS (ESI) m/z : 431.3 ($M+H$)⁺; Anal. HPLC Retention time: 1.49 (Method 1); ROCK2 IC₅₀=85nM.


5

Example 6: Preparation of 8-(1-methyl-1H-indazole-3-carbonyl)-2-[4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one




To a round bottom flask was added 3A (25 mg, 0.059 mmol), THF (1 mL), 1-methyl-1H-indazole-3-carboxylic acid (10.32 mg, 0.059 mmol), HATU (26.7 mg, 0.070 mmol) and Et₃N (0.012 mL, 0.088 mmol). The reaction was stirred at rt overnight. The reaction was partitioned between EtOAc (20 ml) and water (15 ml). The organic layer was separated, washed with brine (15 ml), dried over MgSO₄, filtered and concentrated. To the residue was added CH₂Cl₂ (0.5 mL) and TFA (0.5 mL). The reaction was stirred at rt for 4 hr. The reaction was concentrated and purified using Purification Method A to give 8-(1-methyl-1H-indazole-3-carbonyl)-2-[4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one (10.2 mg, 0.022 mmol, 37.5% yield). ¹H NMR (500 MHz, DMSO-d₆) δ 8.814 (br. s., 1H), 7.93 (d, $J=8.1$ Hz, 2H), 7.74 - 7.57 (m, 5H), 7.47 (t, $J=7.6$ Hz, 1H), 7.26 (t, $J=7.4$ Hz, 1H), 4.55 (br. s., 1H), 4.47 (br. s., 1H), 4.10 (s, 3H), 3.85 (br. s., 2H), 3.65 - 3.56 (m, 1H), 3.16 (d, $J=5.7$ Hz, 1H), 2.19 (d, $J=6.0$ Hz, 2H), 1.86 - 1.73 (m, 2H), 1.68 (br. s., 1H), 1.60 (br. s., 1H); MS (ESI) m/z : 455.0 ($M+H$)⁺; Anal. HPLC Retention time: 1.34 (Method 2); ROCK2 IC₅₀=143nM.

Example 7: Preparation of 8-cyclopropanecarbonyl-2-[4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one




8-Cyclopropanecarbonyl-2-[4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-
 5 1-one (9.9 mg, 0.027 mmol, 45.4% yield) was prepared in a similar manner as the
 procedure described in Example 3, using cyclopropanecarbonyl chloride (6.13 mg, 0.059
 mmol). ¹H NMR (500 MHz, DMSO-d₆) δ 8.15 (br. s., 1H), 7.91 (br. s., 1H), 7.72 - 7.54
 (m, 4H), 4.21 (br. s., 2H), 3.84 (t, J=6.9 Hz, 2H), 3.33 (br. s., 1H), 2.90 (br. s., 1H), 2.14
 (t, J=6.9 Hz, 2H), 2.00 (t, J=6.1 Hz, 1H), 1.73 (br. s., 1H), 1.63 - 1.44 (m, 3H), 0.72 (br.
 10 s., 4H); MS (ESI) *m/z*: 365.2 (M+H)⁺; Anal. HPLC Retention time: 1.11 Method 1;
 ROCK2 IC₅₀=875nM.

Example 8: Preparation of 8-(3-fluoro-2-methoxybenzoyl)-2-[4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one




8-(3-Fluoro-2-methoxybenzoyl)-2-[4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one (7.6 mg, 0.017 mmol, 47.2% yield) was prepared in a similar manner as the procedure described in Example 6, using 3-fluoro-2-methoxybenzoic acid (5.98 mg, 0.035 mmol). ¹H NMR (500 MHz, DMSO-d₆) δ 8.15 (br. s., 1H), 7.90 (br. s., 1H), 7.76 - 7.54 (m, 4H), 7.40 - 7.27 (m, 1H), 7.17 (br. s., 1H), 7.12 - 6.96 (m, 1H), 4.37 (br. s., 1H), 3.99 - 3.71 (m, 5H), 3.42 (s, 1H), 3.21 - 3.04 (m, 2H), 2.24 - 2.04 (m, 2H), 1.80 - 1.57 (m, 3H), 1.48 (d, J=12.5 Hz, 1H); MS (ESI) *m/z*: 449.3 (M+H)⁺; Anal. HPLC Retention time: 1.48 (Method 1); ROCK2 IC₅₀=112nM.

10 **Example 9: Preparation of 8-(4-fluoro-2-methoxybenzoyl)-2-[4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one**



8-(4-Fluoro-2-methoxybenzoyl)-2-[4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one (7.6 mg, 0.017 mmol, 28.3% yield) was prepared in a similar manner as the procedure described in Example 6, using 4-fluoro-2-methoxybenzoic acid (9.97 mg, 0.059 mmol). ¹H NMR (500 MHz, DMSO-d₆) δ 8.16 (br. s., 1H), 7.99 - 7.79 (m, 1H), 7.72 - 7.54 (m, 4H), 7.34 - 7.13 (m, 1H), 7.01 (d, J=11.0 Hz, 1H), 6.84 (br. s., 1H), 4.33 (d, J=13.7 Hz, 1H), 3.91 - 3.69 (m, 5H), 3.47 - 3.31 (m, 1H), 3.23 - 3.00 (m, 2H), 2.26 - 2.00 (m, 2H), 1.82 - 1.52 (m, 3H), 1.46 (d, J=12.8 Hz, 1H); MS (ESI) *m/z*: 449.1 (M+H)⁺; Anal. HPLC Retention time: 1.47 (Method 1); ROCK2 IC₅₀=241nM.

Example 10: Preparation of 2-[4-(1H-pyrazol-4-yl)phenyl]-8-[4-(trifluoromethoxy)benzoyl]-2,8-diazaspiro[4.5]decan-1-one




2-[4-(1H-pyrazol-4-yl)phenyl]-8-[4-(trifluoromethoxy)benzoyl]-2,8-diazaspiro[4.5]decan-1-one (14.5 mg, 0.030 mmol, 36.1% yield) was prepared in a similar manner as the procedure described in Example 6, using 4-

5 (trifluoromethoxy)benzoic acid (16.9 mg, 0.082 mmol). ¹H NMR (500 MHz, DMSO-d₆) δ 8.16 (br. s., 1H), 7.93 (br. s., 1H), 7.70 - 7.64 (m, 2H), 7.63 - 7.59 (m, 2H), 7.56 (d, J=8.5 Hz, 2H), 7.45 (d, J=8.2 Hz, 2H), 4.40 - 4.21 (m, 1H), 3.83 (br. s., 2H), 3.30 - 3.07 (m, 1H), 2.60 - 2.45 (m, 2H), 2.23 - 2.06 (m, 2H), 1.87 - 1.60 (m, 3H), 1.53 (br. s., 1H); MS (ESI) *m/z*: 485.2 (M+H)⁺; Anal. HPLC Retention time: 1.74 (Method 2); ROCK2

10 IC₅₀=95nM.

Example 11: Preparation of 8-(3-fluoro-4-methoxybenzoyl)-2-[4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one




15 8-(3-fluoro-4-methoxybenzoyl)-2-[4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one (15.8 mg, 0.035 mmol, 58.9% yield) was prepared in a


similar manner as the procedure described in Example 6, using 3-fluoro-4-methoxybenzoic acid (9.97 mg, 0.059 mmol). ¹H NMR (500 MHz, DMSO-d₆) δ 8.13 (br. s., 1H), 7.91 (br. s., 1H), 7.71 - 7.55 (m, 4H), 7.28 (d, J=11.6 Hz, 1H), 7.23 (br. s., 2H), 4.23 (q, J=5.2 Hz, 2H), 3.87 (s, 3H), 3.84 - 3.80 (m, 2H), 3.68 - 3.63 (m, 2H), 2.13 (br. s., 2H), 1.72 (br. s., 2H), 1.57 (br. s., 2H); MS (ESI) *m/z*: 449.2 (M+H)⁺; Anal. HPLC Retention time: 1.53 (Method 1); ROCK2 EC₅₀=16nM.

Example 12: Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-(2-methoxybenzoyl)-2,8-diazaspiro[4.5]decan-1-one

10



12A: Preparation of 2-(4-bromo-3-methoxyphenyl)-8-(2-methoxybenzoyl)-2,8-diazaspiro[4.5]decan-1-one




15

To a round bottom flask was added Intermediate 2 (50 mg, 0.114 mmol), and 4N HCl (1.423 mL, 5.69 mmol) in Dioxane (0.5 mL). The reaction was stirred at rt for 30 min. The reaction was concentrated. Then the reaction was dissolved in THF (1.5 mL)

and Et₃N (0.095 mL, 0.683 mmol) and 2-methoxybenzoyl chloride (23.30 mg, 0.137 mmol) was added to the reaction. The reaction was stirred at rt for 15 min. The reaction was then partitioned between EtOAc (20 ml) and water (10 ml). The organic layer was separated, washed with water (10 ml) and brine (10 ml), dried over MgSO₄, filtered and concentrated. The residue was purified using ISCO system (0-100% EtOAc/Hex gradient) to give 2-(4-bromo-3-methoxyphenyl)-8-(2-methoxybenzoyl)-2,8-diazaspiro[4.5]decan-1-one (50 mg, 0.106 mmol, 93 % yield) as a beige solid. ¹H NMR (500 MHz, DMSO-d₆) δ 7.67 (dd, *J*=10.5, 1.9 Hz, 1H), 7.55 (d, *J*=8.5 Hz, 1H), 7.43 - 7.36 (m, 1H), 7.24 - 7.14 (m, 1H), 7.14 - 7.07 (m, 2H), 7.01 (q, *J*=7.4 Hz, 1H), 4.36 (dd, *J*=17.6, 13.8 Hz, 1H), 3.85 (d, *J*=5.0 Hz, 4H), 3.82 - 3.78 (m, 3H), 3.39 - 3.32 (m, 1H), 3.21 - 3.01 (m, 2H), 2.24 - 2.04 (m, 2H), 1.83 - 1.53 (m, 3H), 1.46 (d, *J*=12.9 Hz, 1H), 1.33 - 1.24 (m, 1H); MS (ESI) *m/z*: 473.0, 475.0 (M+H)⁺.

12B: Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-(2-methoxybenzoyl)-2,8-diazaspiro[4.5]decan-1-one




To a microwave vial was added 2-(4-bromo-3-methoxyphenyl)-8-(2-methoxybenzoyl)-2,8-diazaspiro[4.5]decan-1-one (60 mg, 0.127 mmol), 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-pyrazole (73.8 mg, 0.380 mmol), tripotassium phosphate (0.169 mL, 0.507 mmol), Dioxane (3 mL) and methanesulfonato(2-dicyclohexylphosphino-2',4',6'-tri-*i*-propyl-1,1'-biphenyl)(2'-amino-1,1'-biphenyl-2-yl)palladium(II) (10.73 mg, 0.013 mmol). The reaction was purged with argon and sealed. The reaction was then subjected to microwave oven and stirred at 120°C for 30 min. The reaction was partitioned between EtOAc (50 ml) and water (20 ml). The organic

layer was separated, washed with water (20 ml) and brine (30 ml), dried over MgSO_4 , filtered and concentrated. The residue was purified using Purification Method A to give 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-(2-methoxybenzoyl)-2,8-diazaspiro[4.5]decan-1-one (21.6 mg, 0.046 mmol, 36.3% yield). ^1H NMR (500 MHz, DMSO- d_6) δ 8.30 - 7.78 (m, 2H), 7.61 (d, $J=9.7$ Hz, 2H), 7.40 (d, $J=4.0$ Hz, 1H), 7.26 - 7.06 (m, 3H), 7.01 (q, $J=7.6$ Hz, 1H), 4.51 - 4.25 (m, 1H), 3.98 - 3.66 (m, 7H), 3.54 - 3.27 (m, 2H), 3.22 - 2.97 (m, 2H), 2.28 - 2.00 (m, 2H), 1.83 - 1.52 (m, 3H), 1.46 (d, $J=12.5$ Hz, 1H); MS (ESI) m/z : 461.0 ($\text{M}+\text{H}^+$); Anal. HPLC Retention time: 1.5 (Method 1); ROCK2 $\text{IC}_{50}=54\text{nM}$.


10

Example 13: Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-(4-methoxybenzoyl)-2,8-diazaspiro[4.5]decan-1-one




2-[3-Methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-(4-methoxybenzoyl)-2,8-diazaspiro[4.5]decan-1-one (19.6 mg, 0.042 mmol, 43.9% yield) was prepared in a similar manner as the procedure described in Example 10, using 4-methoxybenzoyl chloride (23.3mg, 0.137 mmol). ^1H NMR (500 MHz, DMSO- d_6) δ 8.02 (br. s., 2H), 7.64 - 7.52 (m, 2H), 7.37 (d, $J=8.5$ Hz, 2H), 7.19 - 7.10 (m, 1H), 6.99 (d, $J=8.5$ Hz, 2H), 3.86 - 3.81 (m, 4H), 3.79 (s, 3H), 3.60 (br. s., 1H), 3.17 (s, 2H), 2.53 (d, $J=19.2$ Hz, 2H), 2.14 (br. s., 2H), 1.72 (br. s., 2H), 1.58 (br. s., 2H); MS (ESI) m/z : 461.0 ($\text{M}+\text{H}^+$). Anal; HPLC Retention time: 1.52 (Method 1); ROCK2 $\text{IC}_{50}=8\text{nM}$.

Example 14: Preparation of 8-(3-fluoro-4-methoxybenzoyl)-2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one



To a round bottom flask was added 3-fluoro-4-methoxybenzoic acid (8.52 mg, 0.050 mmol), THF (1 mL), HATU (22.85 mg, 0.060 mmol) and Et₃N (0.035 mL, 0.250 mmol). The reaction was stirred at rt for 10 min and then Intermediate 4 (20 mg, 0.050 mmol) was added and the reaction. The reaction was continued for 2 hr. The reaction was partitioned between EtOAc (20 ml) and water (15 ml). The organic layer was separated, washed with brine (15 ml), dried over MgSO₄, filtered and concentrated. The residue was purified using Purification method A to give 8-(3-fluoro-4-methoxybenzoyl)-2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one (4.1 mg, 0.0081 mmol, 16.1% yield). ¹H NMR (500 MHz, DMSO-d₆) δ 8.10 - 7.93 (m, 2H), 7.62 - 7.58 (m, 2H), 7.30 (d, J=11.6 Hz, 1H), 7.23 (br. s., 2H), 7.15 (d, J=8.5 Hz, 1H), 3.91 - 3.84 (m, 8H), 3.42 (br. s., 2H), 3.17 (d, J=5.2 Hz, 2H), 2.14 (br. s., 2H), 1.75 (d, J=13.1 Hz, 2H), 1.59 (br. s., 2H); MS (ESI) *m/z*: 479.1 (M+H)⁺; Anal. HPLC Retention time: 1.53 (Method 1); ROCK2 IC₅₀=8nM.

Example 15: Preparation of 8-[4-(hydroxymethyl)benzoyl]-2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one




8-[4-(Hydroxymethyl)benzoyl]-2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one (12.4 mg, 0.026 mmol, 52.7% yield) was prepared in a similar manner as the procedure described in Example 14, using 4-

- 5 (hydroxymethyl)benzoic acid (7.62 mg, 0.050 mmol). ¹H NMR (500 MHz, DMSO-d₆) δ 8.01 (br. s., 2H), 7.62 - 7.54 (m, 2H), 7.42 - 7.33 (m, 4H), 7.14 (d, J=8.5 Hz, 1H), 4.53 (d, J=5.2 Hz, 2H), 3.93 - 3.82 (m, 5H), 3.60 (br. s., 2H), 3.17 (d, J=5.2 Hz, 2H), 2.14 (br. s., 2H), 1.71 (br. s., 4H); MS (ESI) *m/z*: 461.1 (M+H)⁺; Anal. HPLC Retention time: 1.22 (Method 1); ROCK2 IC₅₀=53nM.

10


Example 16: Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-(5-methoxypyridine-2-carbonyl)-2,8-diazaspiro[4.5]decan-1-one



- 15 2-[3-Methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-(5-methoxypyridine-2-carbonyl)-2,8-diazaspiro[4.5]decan-1-one (15.8 mg, 0.034 mmol, 67% yield) was prepared in a similar


manner as the procedure described in Example 14, using 5-methoxypicolinic acid (7.67 mg, 0.050 mmol). ¹H NMR (500 MHz, DMSO-d₆) δ 8.29 (d, J=2.4 Hz, 1H), 8.02 (br. s., 2H), 7.70 - 7.56 (m, 3H), 7.51 (dd, J=8.6, 2.5 Hz, 1H), 7.15 (d, J=8.4 Hz, 1H), 4.36 (d, J=13.2 Hz, 1H), 3.87 (d, J=6.1 Hz, 9H), 3.36 - 3.22 (m, 1H), 3.13 (t, J=11.0 Hz, 1H), 2.16 (dd, J=17.7, 6.6 Hz, 2H), 1.84 - 1.71 (m, 2H), 1.65 (d, J=12.5 Hz, 1H), 1.52 (d, J=11.6 Hz, 1H); MS (ESI) *m/z*: 461.9 (M+H)⁺; Anal. HPLC Retention time: 1.41 (Method 1); ROCK2 IC₅₀=13nM.

Example 17: Preparation of 8-(2-fluoro-4-methoxybenzoyl)-2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one




8-(2-Fluoro-4-methoxybenzoyl)-2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one (9.1 mg, 0.019 mmol, 37.2 % yield) was prepared in a similar manner as the procedure described in Example 14, using 2-fluoro-4-methoxybenzoic acid (8.52 mg, 0.050 mmol). ¹H NMR (500 MHz, DMSO-d₆) δ 8.17 - 7.77 (m, 2H), 7.67 - 7.56 (m, 2H), 7.32 (t, J=8.2 Hz, 1H), 7.14 (d, J=8.5 Hz, 1H), 6.98 - 6.78 (m, 3H), 4.19 (br. s., 1H), 3.92 - 3.82 (m, 6H), 3.80 (s, 3H), 3.12 (br. s., 2H), 2.13 (dd, J=13.9, 7.2 Hz, 3H), 1.83 - 1.59 (m, 3H), 1.51 (d, J=11.9 Hz, 1H); MS (ESI) *m/z*: 478.9 (M+H)⁺; Anal. HPLC Retention time: 1.47 (Method 1); ROCK2 IC₅₀=8nM.

Example 18: Preparation of 8-(4-chlorobenzoyl)-2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one



8-(4-Chlorobenzoyl)-2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one (21.1 mg, 0.044 mmol, 88% yield) was prepared in a similar manner as the procedure described in Example 14, using 4-chlorobenzoic acid (7.84 mg, 0.050 mmol). ¹H NMR (500 MHz, DMSO-d₆) δ 8.18 - 7.92 (m, 2H), 7.65 - 7.56 (m, 2H), 7.53 (d, J=8.2 Hz, 2H), 7.44 (d, J=8.2 Hz, 2H), 7.30 - 7.00 (m, 2H), 3.94 - 3.76 (m, 5H), 3.34 - 3.03 (m, 2H), 2.55 (s, 2H), 2.25 - 2.09 (m, 2H), 1.81 - 1.44 (m, 4H); MS (ESI) *m/z*: 464.9 (M+H)⁺; Anal. HPLC Retention time: 1.7 (Method 1); ROCK2 IC₅₀=62nM.


10 **Example 19: Preparation of 8-(4-cyclopropylbenzoyl)-2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one**



8-(4-Cyclopropylbenzoyl)-2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one (11.2 mg, 0.023 mmol, 46.6% yield) was prepared in a

similar manner as the procedure described in Example 14, using 4-cyclopropylbenzoic acid (8.12 mg, 0.050 mmol). ¹H NMR (500 MHz, DMSO-d₆) δ 7.99 (br. s., 2H), 7.67 - 7.53 (m, 2H), 7.28 (d, J=8.0 Hz, 2H), 7.14 (d, J=8.0 Hz, 3H), 3.86 (s, 5H), 3.51 - 3.38 (m, 1H), 3.17 (br. s., 1H), 2.55 (s, 2H), 2.14 (br. s., 2H), 1.99 - 1.90 (m, 1H), 1.83 - 1.38 (m, 4H), 0.99 (d, J=6.6 Hz, 2H), 0.71 (d, J=5.1 Hz, 2H); MS (ESI) *m/z*: 471.3 (M+H)⁺; Anal. HPLC Retention time: 1.58 (Method 1); ROCK2 IC₅₀=10nM


Example 20: Preparation of 8-(4-ethylbenzoyl)-2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one



10

8-(4-Ethylbenzoyl)-2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one (21.3 mg, 0.046 mmol, 91% yield) was prepared in a similar manner as the procedure described in Example 14, using 4-ethylbenzoic acid (7.52 mg, 0.050 mmol). ¹H NMR (500 MHz, DMSO-d₆) δ 8.02 (br. s., 2H), 7.66 - 7.53 (m, 2H), 7.31 (q, J=7.9 Hz, 4H), 7.14 (d, J=8.1 Hz, 1H), 3.86 (s, 5H), 3.35 - 3.02 (m, 1H), 2.64 (q, J=7.6 Hz, 2H), 2.55 (s, 2H), 2.14 (br. s., 2H), 1.83 - 1.45 (m, 4H), 1.20 (t, J=7.5 Hz, 3H); MS (ESI) *m/z*: 458.9 (M+H)⁺; Anal. HPLC Retention time: 1.79 (Method 1); ROCK2 IC₅₀=50nM


20 **Example 21: Preparation of 8-[4-(dimethylamino)benzoyl]-2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one**



8-[4-(Dimethylamino)benzoyl]-2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one (18.4 mg, 0.038 mmol, 76% yield) was prepared in a similar manner as the procedure described in Example 14, using 4-(dimethylamino)benzoic acid (8.27 mg, 0.050 mmol). ¹H NMR (500 MHz, DMSO-d₆) δ 8.23 - 7.79 (m, 2H), 7.69 - 7.55 (m, 2H), 7.28 (d, J=8.6 Hz, 2H), 7.14 (d, J=6.9 Hz, 1H), 6.72 (d, J=8.7 Hz, 2H), 3.93 - 3.77 (m, 4H), 3.52 (d, J=5.6 Hz, 2H), 3.17 (d, J=5.0 Hz, 2H), 3.00 - 2.85 (m, 6H), 2.25 - 2.06 (m, 2H), 1.78 - 1.64 (m, 2H), 1.56 (d, J=12.9 Hz, 2H); MS (ESI) *m/z*: 474.4 (M+H)⁺; Anal. HPLC Retention time: 1.43 (Method 1); ROCK2 IC₅₀=16nM.


10

Example 22: Preparation of 8-(3-chloro-4-methoxybenzoyl)-2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one



8-(3-Chloro-4-methoxybenzoyl)-2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one (21.2 mg, 0.042 mmol, 84% yield) was prepared in a similar manner as the procedure described in Example 14, using 3-chloro-4-methoxybenzoic acid (9.35 mg, 0.050 mmol). ¹H NMR (500 MHz, DMSO-d₆) δ 8.16 - 7.83 (m, 2H), 7.62 (d, J=5.3 Hz, 2H), 7.49 (s, 1H), 7.40 (d, J=8.4 Hz, 1H), 7.22 (d, J=8.5 Hz, 1H), 7.15 (d, J=8.5 Hz, 1H), 3.90 (s, 3H), 3.87 (s, 5H), 3.44 - 3.15 (m, 2H), 2.55 (s, 2H), 2.22 - 2.09 (m, 2H), 1.73 (br. s., 2H), 1.59 (br. s., 2H); MS (ESI) *m/z*: 495.2 (M+H)⁺; Anal. HPLC Retention time: 1.45 (Method 1); ROCK2 IC₅₀=6nM.


10 Example 23: Preparation of 4-{2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-1-oxo-2,8-diazaspiro[4.5]decane-8-carbonyl}benzamide



4-{2-[3-Methoxy-4-(1H-pyrazol-4-yl)phenyl]-1-oxo-2,8-diazaspiro[4.5]decane-8-carbonyl}benzamide (10.5 mg, 0.022 mmol, 57.8% yield) was prepared in a similar manner as the procedure described in Example 14, using 4-carbamoylbenzoic acid (6.20 mg, 0.038 mmol). ¹H NMR (500 MHz, DMSO-d₆) δ 8.16 - 7.87 (m, 5H), 7.67 - 7.56 (m, 2H), 7.48 (d, J=7.8 Hz, 3H), 7.14 (d, J=8.4 Hz, 2H), 3.86 (s, 2H), 3.31 - 3.10 (m, 1H), 2.51 (br. s., 5H), 2.14 (br. s., 2H), 1.85 - 1.38 (m, 4H); MS (ESI) *m/z*: 474.3 (M+H)⁺; ROCK2 IC₅₀=36nM.

20

Example 24: Preparation of methyl 4-{2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-1-oxo-2,8-diazaspiro[4.5]decane-8-carbonyl}benzoate




Methyl 4-{2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-1-oxo-2,8-diazaspiro[4.5]decane-8-carbonyl}benzoate (40 mg, 0.078 mmol) was prepared in a similar manner as the procedure described in Example 14, using 4-

- 5 (methoxycarbonyl)benzoic acid (22.56 mg, 0.125 mmol). ¹H NMR (500 MHz, DMSO-d₆) δ 8.10 - 7.93 (m, 2H), 7.62 - 7.58 (m, 2H), 7.30 (d, J=11.6 Hz, 1H), 7.23 (br. s., 2H), 7.15 (d, J=8.5 Hz, 1H), 3.91 - 3.84 (m, 8H), 3.42 (br. s., 2H), 3.17 (d, J=5.2 Hz, 2H), 2.14 (br. s., 2H), 1.75 (d, J=13.1 Hz, 2H), 1.59 (br. s., 2H); MS (ESI) *m/z*: 479.1 (M+H)⁺; ROCK2 IC₅₀=23nM.

10


Example 25: Preparation of 4-{2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-1-oxo-2,8-diazaspiro[4.5]decane-8-carbonyl}benzoic acid



15

To a round bottom flask was added Example 24 (55 mg, 0.113 mmol), THF (1 mL), water (0.1 mL) and LiOH (2.70 mg, 0.113 mmol). The reaction was stirred at rt overnight. The reaction was concentrated to give 4-(2-(3-methoxy-4-(1H-pyrazol-4-yl)phenyl)-1-oxo-2,8-diazaspiro[4.5]decane-8-carbonyl)benzoic acid, lithium salt (48 mg, 0.090 mmol, 80 % yield) as a white solid. MS (ESI) m/z : 475.1 (M+H)⁺; ROCK2 IC₅₀=1430nM

Example 26: Preparation of N-(cyclopropylmethyl)-4-{2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-1-oxo-2,8-diazaspiro[4.5]decane-8-carbonyl}benzamide




10

To a round bottom flask was added cyclopropylmethanamine (2.70 mg, 0.038 mmol), HATU (21.64 mg, 0.057 mmol), DMF (0.5 mL), Et₃N (0.016 mL, 0.114 mmol) followed by Example 25 (18 mg, 0.038 mmol). The reaction was stirred at rt overnight. The reaction was filtered using Purification Method A to give N-(cyclopropylmethyl)-4-{2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-1-oxo-2,8-diazaspiro[4.5]decane-8-carbonyl}benzamide (4.4 mg, 0.0077 mmol, 20.2% yield). ¹H NMR (500 MHz, DMSO-d₆) δ 8.43 (br t, J =5.3 Hz, 1H), 7.79 (br s, 2H), 7.69 (br d, J =8.2 Hz, 2H), 7.41 - 7.33 (m, 2H), 7.26 (br d, J =8.2 Hz, 2H), 6.92 (dd, J =8.5, 1.8 Hz, 1H), 3.75 - 3.54 (m, 5H), 3.16 - 2.86 (m, 6H), 1.92 (br s, 2H), 1.59 - 1.38 (m, 3H), 1.30 (br s, 1H), 0.87 - 0.70 (m, 1H), 0.21 (br d, J =6.7 Hz, 2H), 0.01 (br d, J =4.6 Hz, 2H); MS (ESI) m/z : 528.3 (M+H)⁺; Anal. HPLC Retention time: 1.28 (Method 1); ROCK2 IC₅₀=119nM


20

Example 27: Preparation of 8-[2-chloro-4-(4-methylpiperazin-1-yl)benzoyl]-2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one




8-[2-Chloro-4-(4-methylpiperazin-1-yl)benzoyl]-2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one was prepared in a similar manner as the procedure described in Example 14, using 2-chloro-4-(4-methylpiperazin-1-yl)benzoic acid (9.57 mg, 0.038 mmol). ¹H NMR (500 MHz, DMSO-d₆) δ 8.09 - 7.92 (m, 2H), 8.02 (br. s., 2H), 7.67 - 7.55 (m, 2H), 7.65 - 7.54 (m, 2H), 7.21 - 7.07 (m, 2H), 7.22 - 7.07 (m, 2H), 7.02 - 6.88 (m, 2H), 6.97 (d, J=14.4 Hz, 2H), 4.28 - 4.25 (m, 1H), 3.92 - 3.79 (m, 6H), 3.95 - 3.79 (m, 6H), 3.31 - 3.02 (m, 4H), 2.43 (br. s., 4H), 2.46 - 2.38 (m, 4H), 2.21 (s, 3H), 2.18 - 2.04 (m, 2H), 2.17 - 2.03 (m, 2H), 1.83 - 1.69 (m, 2H), 1.75 (d, J=12.9 Hz, 2H), 1.64 (d, J=12.4 Hz, 1H), 1.49 (d, J=11.6 Hz, 1H), 1.67 - 1.43 (m, 2H); MS (ESI) *m/z*: 563.2 (M+H)⁺; Anal. HPLC Retention time: 1.03 (Method 2); ROCK2 IC₅₀=0.8nM.

Example 28: Preparation of 8-(2,6-difluoro-4-methoxybenzoyl)-2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one




8-(2,6-Difluoro-4-methoxybenzoyl)-2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one (17.6 mg, 0.035 mmol, 92% yield) was prepared in a similar manner as the procedure described in Example 14, using 2,6-difluoro-4-methoxybenzoic acid (7.07 mg, 0.038 mmol). ¹H NMR (500 MHz, DMSO-d₆) δ 8.00 (br. s., 2H), 7.64 - 7.55 (m, 2H), 7.13 (d, J=8.2 Hz, 1H), 6.84 (t, J=12.2 Hz, 2H), 4.37 (d, J=13.4 Hz, 2H), 3.92 - 3.75 (m, 8H), 3.32 - 3.08 (m, 2H), 2.22 - 2.05 (m, 2H), 1.75 - 1.45 (m, 4H); MS (ESI) *m/z*: 497.0 (M+H)⁺; ROCK2 IC₅₀=5nM,

Example 29: Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-(2,4,6-trimethoxybenzoyl)-2,8-diazaspiro[4.5]decan-1-one



2-[3-Methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-(2,4,6-trimethoxybenzoyl)-2,8-diazaspiro[4.5]decan-1-one (13.9 mg, 0.026 mol, 69.7% yield) was prepared in a similar manner as the procedure described in Example 14, using 2,4,6-trimethoxybenzoic acid (7.97 mg, 0.038 mmol). ¹H NMR (500 MHz, DMSO-d₆) δ 8.01 (br. s., 2H), 7.60 (d, J=3.6 Hz, 2H), 7.12 (d, J=8.2 Hz, 1H), 6.26 (d, J=5.3 Hz, 2H), 4.37 (d, J=13.5 Hz, 1H), 3.89 - 3.67 (m, 11H), 3.11 - 2.92 (m, 2H), 2.22 - 2.00 (m, 2H), 1.67 (d, J=11.8 Hz, 2H), 1.57 (d, J=12.8 Hz, 1H), 1.42 (d, J=12.4 Hz, 1H); MS (ESI) *m/z*: 521.1 (M+H)⁺; ROCK2 IC₅₀=336nM.


Example 30: Preparation of 4-{2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-1-oxo-2,8-diazaspiro[4.5]decane-8-carbonyl}-3,5-dimethylbenzamide



4-{2-[3-Methoxy-4-(1H-pyrazol-4-yl)phenyl]-1-oxo-2,8-diazaspiro[4.5]decane-8-carbonyl}-3,5-dimethylbenzamide (10.2 mg, 0.020 mmol, 52.5% yield) was prepared in a similar manner as the procedure described in Example 14, using 4-carbamoyl-2,6-dimethylbenzoic acid (7.26 mg, 0.038 mmol). ¹H NMR (500 MHz, DMSO-d₆) δ 8.11 - 7.91 (m, 3H), 7.67 - 7.54 (m, 4H), 7.32 (br. s., 1H), 7.11 (d, J=7.9 Hz, 1H), 4.46 (d, J=13.1 Hz, 1H), 3.90 - 3.78 (m, 5H), 3.55 (br. s., 1H), 3.19 - 3.12 (m, 2H), 2.26 (s, 3H), 2.21 - 2.09 (m, 5H), 1.72 - 1.57 (m, 3H), 1.48 (d, J=13.4 Hz, 1H); MS (ESI) *m/z*: 502.2 (M+H)⁺; ROCK2 IC₅₀=130nM.


10

Example 31: Preparation of 8-[4-(4-methylpiperazin-1-yl)benzoyl]-2-[4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one



15


31A. Preparation of 2-(4-bromophenyl)-8-(4-(4-methylpiperazin-1-yl)benzoyl)-2,8-diazaspiro[4.5]decan-1-one



To a round bottom flask was added 4-(4-methylpiperazin-1-yl)benzoic acid (147
 5 mg, 0.665 mmol), THF (10 mL), HATU (304 mg, 0.798 mmol) and Et₃N (0.464 mL, 3.33
 mmol). The reaction was stirred at rt for 10 min and then Intermediate 3 (230 mg, 0.665
 mmol) was added and the reaction was continued for 2 hr. The reaction was concentrated
 and the residue was purified using ISCO system (0-15% MeOH/CH₂Cl₂ gradient) to give
 2-(4-bromophenyl)-8-(4-(4-methylpiperazin-1-yl)benzoyl)-2,8-diazaspiro[4.5]decan-1-
 10 one (305 mg, 0.596 mmol, 90 % yield) as an off-white solid. ¹H NMR (500 MHz,
 DMSO-d₆) δ 7.71 - 7.63 (m, 2H), 7.59 - 7.53 (m, 2H), 7.31 (d, *J*=8.8 Hz, 2H), 6.99 (d,
J=8.8 Hz, 2H), 4.17 - 3.85 (m, 2H), 3.82 (t, *J*=6.9 Hz, 2H), 3.35 - 3.28 (m, 7H), 3.18 (br.
 s., 2H), 2.84 - 2.66 (m, 2H), 2.44 (br. s., 2H), 2.14 (t, *J*=6.9 Hz, 2H), 1.77 - 1.65 (m, 2H),
 1.57 (d, *J*=12.7 Hz, 2H); MS (ESI) *m/z*: 511.0, 513.0 (M+H)⁺.


15

31B. Preparation of 8-[4-(4-methylpiperazin-1-yl)benzoyl]-2-[4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one



8-[4-(4-Methylpiperazin-1-yl)benzoyl]-2-[4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one was prepared in a similar manner as the procedure described in Example 1, using 31A (20 mg, 0.039 mmol). ¹H NMR (500 MHz, DMSO-d₆) δ 8.04 (br. s., 2H), 7.71 - 7.56 (m, 4H), 7.29 (d, J=8.3 Hz, 2H), 6.96 (d, J=8.5 Hz, 2H), 3.83 (t, J=6.6 Hz, 2H), 3.19 (d, J=19.4 Hz, 4H), 2.55 (s, 2H), 2.44 (br. s., 4H), 2.14 (br. s., 2H), 1.91 (s, 2H), 1.78 - 1.64 (m, 2H), 1.57 (br. s., 2H); MS (ESI) *m/z*: 499.4 (M+H)⁺; Anal. HPLC Retention time: 1.22 Method 1; ROCK2 IC₅₀=0.7nM.

10 **Example 32: Preparation of 2-[4-(2-hydroxypyridin-4-yl)phenyl]-8-[4-(4-methylpiperazin-1-yl)benzoyl]-2,8-diazaspiro[4.5]decan-1-one**




To a microwave vial was added Intermediate 6 (20 mg, 0.039 mmol), 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyridin-2-ol (17.29 mg, 0.078 mmol), tripotassium phosphate (0.052 mL, 0.156 mmol), Dioxane (3 mL) and methanesulfonato(2-dicyclohexylphosphino-2',4',6'-tri-*i*-propyl-1,1'-biphenyl)(2'-amino-1,1'-biphenyl-2-yl)palladium(II) (3.31 mg, 3.91 μmol). The reaction was purged with argon and sealed.

The reaction was then subjected to microwave oven and stirred at 120 °C for 30 min. The reaction was partitioned between EtOAc (50 ml) and water (20 ml). The organic layer was separated, washed with water (20 ml) and brine (30 ml), dried over MgSO₄, filtered and concentrated. The residue was purified using Purification Method A to give

5 2-[4-(2-hydroxypyridin-4-yl)phenyl]-8-[4-(4-methylpiperazin-1-yl)benzoyl]-2,8-diazaspiro[4.5]decan-1-one (1.6 mg, 0.0027 mmol, 6.7% yield). ¹H NMR (500 MHz, DMSO-d₆) δ 7.84 - 7.78 (m, 2H), 7.78 - 7.72 (m, 2H), 7.44 (d, *J*=6.7 Hz, 1H), 7.35 (d, *J*=8.2 Hz, 2H), 7.05 (d, *J*=8.5 Hz, 2H), 6.60 (s, 1H), 6.55 (d, *J*=6.7 Hz, 1H), 3.91 - 3.85 (m, 2H), 3.44 (d, *J*=6.1 Hz, 2H), 3.25 - 3.07 (m, 2H), 2.85 (s, 3H), 2.55 (s, 8H), 2.17 (br.

10 s., 2H), 1.72 (br. s., 2H), 1.60 (br. s., 2H); MS (ESI) *m/z*: 526.4 (M+H)⁺; Anal. HPLC Retention time: 0.88 (Method 2); ROCK2 IC₅₀=54nM.

Example 33: Preparation of 2-[4-(3-fluoropyridin-4-yl)phenyl]-8-[4-(4-methylpiperazin-1-yl)benzoyl]-2,8-diazaspiro[4.5]decan-1-one




15

2-[4-(3-Fluoropyridin-4-yl)phenyl]-8-[4-(4-methylpiperazin-1-yl)benzoyl]-2,8-diazaspiro[4.5]decan-1-one (2.3 mg, 4.3 mmol, 10.9% yield) was prepared in a similar manner as the procedure described in Example 32, using 3-fluoro-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyridine (17.44 mg, 0.078 mmol). ¹H NMR (500 MHz, DMSO-


20 d₆) δ 8.66 (br. s., 1H), 8.50 (br. s., 1H), 7.88 (d, *J*=8.6 Hz, 2H), 7.73 (d, *J*=8.2 Hz, 2H), 7.66 (br. s., 1H), 7.30 (d, *J*=8.5 Hz, 2H), 6.96 (d, *J*=8.6 Hz, 2H), 3.89 (t, *J*=6.7 Hz, 2H), 3.44 - 3.09 (m, 4H), 2.55 - 2.51 (m, 4H), 2.44 (br. s., 4H), 2.22 (s, 3H), 2.17 (t, *J*=6.4 Hz, 2H), 1.80 - 1.68 (m, 2H), 1.60 (br. s., 2H); MS (ESI) *m/z*: 528.4 (M+H)⁺; Anal. HPLC Retention time: 1.01 (Method 2); ROCK2 IC₅₀=1.3nM.

Example 34: Preparation of 2-[4-(1H-indazol-6-yl)phenyl]-8-[4-(4-methylpiperazin-1-yl)benzoyl]-2,8-diazaspiro[4.5]decan-1-one




5 2-[4-(1H-indazol-6-yl)phenyl]-8-[4-(4-methylpiperazin-1-yl)benzoyl]-2,8-diazaspiro[4.5]decan-1-one (9.1 mg, 0.016 mmol, 41.6% yield) was prepared in a similar manner as the procedure described in Example 32, using 6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-indazole (19.09 mg, 0.078 mmol). ¹H NMR (500 MHz, DMSO-d₆) δ 13.13 (s, 1H), 8.09 (s, 1H), 7.86 - 7.70 (m, 6H), 7.43 (d, J=8.5 Hz, 1H), 7.31 (d, J=8.4 Hz, 2H), 6.97 (d, J=8.6 Hz, 2H), 3.89 (t, J=6.7 Hz, 2H), 3.23 (br. s., 2H), 2.58 - 2.48 (m, 10H), 2.27 (s, 3H), 2.17 (t, J=6.4 Hz, 2H), 1.78 - 1.68 (m, 2H), 1.59 (d, J=11.6 Hz, 2H); MS (ESI) *m/z*: 549.2 (M+H)⁺; Anal. HPLC Retention time: 1.45 (Method 1); ROCK2 IC₅₀=106nM.

15 **Example 35: Preparation of 8-[4-(4-methylpiperazin-1-yl)benzoyl]-2-(4-{1H-pyrrolo[2,3-b]pyridin-4-yl}phenyl)-2,8-diazaspiro[4.5]decan-1-one**




8-[4-(4-Methylpiperazin-1-yl)benzoyl]-2-(4-{1H-pyrrolo[2,3-b]pyridin-4-yl}phenyl)-2,8-diazaspiro[4.5]decan-1-one (8.9 mg, 0.016 mmol, 40.7% yield) was prepared in a similar manner as the procedure described in Example 32, using 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-pyrrolo[2,3-b]pyridine (19.09 mg, 0.078 mmol). ¹H NMR (500 MHz, DMSO-d₆) δ 11.77 (br. s., 1H), 8.28 (d, J=4.9 Hz, 1H), 7.92 - 7.85 (m, 2H), 7.81 (d, J=8.2 Hz, 2H), 7.54 (br. s., 1H), 7.31 (d, J=8.2 Hz, 2H), 7.19 (d, J=4.9 Hz, 1H), 6.97 (d, J=8.5 Hz, 2H), 6.63 (br. s., 1H), 4.31 - 3.96 (m, 2H), 3.91 (t, J=6.6 Hz, 2H), 3.24 - 3.14 (m, 4H), 2.51 - 2.44 (m, 6H), 2.25 (s, 3H), 2.18 (t, J=6.3 Hz, 2H), 1.82 - 1.68 (m, 2H), 1.60 (d, J=11.3 Hz, 2H); MS (ESI) *m/z*: 549.1 (M+H)⁺; Anal. HPLC Retention time: 1.42 (Method 1); ROCK2 IC₅₀=0.3nM.

Example 36: Preparation of 2-[4-(2-aminopyridin-4-yl)phenyl]-8-[4-(4-methylpiperazin-1-yl)benzoyl]-2,8-diazaspiro[4.5]decan-1-one



2-[4-(2-Aminopyridin-4-yl)phenyl]-8-[4-(4-methylpiperazin-1-yl)benzoyl]-2,8-diazaspiro[4.5]decan-1-one (1.4 mg, 0.0026 mmol, 6.6% yield) was prepared in a similar manner as the procedure described in Example 32, using 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyridin-2-amine (17.21 mg, 0.078 mmol). ¹H NMR (500 MHz, DMSO-d₆) δ 7.95 (d, J=5.2 Hz, 1H), 7.81 (d, J=8.5 Hz, 2H), 7.67 (d, J=8.2 Hz, 2H), 7.29 (d, J=8.2 Hz, 2H), 6.96 (d, J=8.2 Hz, 2H), 6.79 (d, J=5.2 Hz, 1H), 6.71 (s, 1H), 5.93 (s, 2H), 4.28 - 3.73 (m, 4H), 3.20 - 3.12 (m, 2H), 2.57 - 2.47 (m, 6H), 2.22 (s, 3H), 2.16 (t, J=6.6 Hz, 2H), 1.77 - 1.65 (m, 2H), 1.58 (d, J=12.2 Hz, 2H); MS (ESI) *m/z*: 525.1 (M+H)⁺; Anal. HPLC Retention time: 0.76 (Method 2); ROCK2 IC₅₀=2.0nM.


Example 37: Preparation of 2-[4-(1H-indazol-5-yl)phenyl]-8-[4-(4-methylpiperazin-1-yl)benzoyl]-2,8-diazaspiro[4.5]decan-1-one



2-[4-(1H-indazol-5-yl)phenyl]-8-[4-(4-methylpiperazin-1-yl)benzoyl]-2,8-diazaspiro[4.5]decan-1-one (4.3 mg, 0.0076 mmol, 19.4% yield) was prepared in a similar manner as the procedure described in Example 32, using (1H-indazol-5-yl)boronic acid (12.67 mg, 0.078 mmol). ¹H NMR (500 MHz, DMSO-d₆) δ 8.13 (s, 1H), 8.03 (s, 1H), 7.81 - 7.76 (m, 2H), 7.72 (d, J=8.5 Hz, 2H), 7.70 - 7.65 (m, 1H), 7.64 - 7.58 (m, 1H), 7.31 (d, J=8.5 Hz, 2H), 6.97 (d, J=8.5 Hz, 2H), 3.92 - 3.82 (m, 2H), 3.23 (br. s., 4H), 2.55 (s, 2H), 2.50 - 2.48 (m, 4H), 2.26 (s, 3H), 2.17 (t, J=6.6 Hz, 2H), 1.79 - 1.68 (m, 2H), 1.59 (d, J=11.3 Hz, 2H); MS (ESI) *m/z*: 549.0 (M+H)⁺; Anal. HPLC Retention time: 1.29 (Method 1); ROCK2 IC₅₀=0.5nM.

10

Example 38: Preparation of 2-[4-(1H-1,2,3-benzotriazol-6-yl)phenyl]-8-[4-(4-methylpiperazin-1-yl)benzoyl]-2,8-diazaspiro[4.5]decan-1-one




2-[4-(1H-1,2,3-benzotriazol-6-yl)phenyl]-8-[4-(4-methylpiperazin-1-yl)benzoyl]-2,8-diazaspiro[4.5]decan-1-one (5.1 mg, 0.0089 mmol, 22.8% yield) was prepared in a similar manner as the procedure described in Example 32, using 6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-benzo[d][1,2,3]triazole (19.17 mg, 0.078 mmol). ¹H NMR (500 MHz, DMSO-d₆) δ 8.34 (s, 1H), 8.08 (s, 1H), 7.94 (d, J=8.9 Hz, 1H), 7.80 (d, J=4.6 Hz, 4H), 7.70 (d, J=8.5 Hz, 1H), 7.30 (d, J=8.5 Hz, 2H), 6.96 (d, J=8.9 Hz, 2H), 3.88 (br. s., 2H), 3.47 (br. s., 2H), 3.24 - 3.05 (m, 6H), 2.44 (br. s., 4H), 2.21 (s, 3H), 2.17 (br. s., 2H), 1.72 (d, J=11.9 Hz, 2H), 1.58 (d, J=12.2 Hz, 2H); MS (ESI) *m/z*: 550.1 (M+H)⁺; Anal. HPLC Retention time: 1.26 (Method 1); ROCK2 IC₅₀=116nM.


20

Example 39: Preparation of 2-[6-methoxy-5-(1H-pyrazol-4-yl)pyridin-2-yl]-8-(4-methoxybenzoyl)-2,8-diazaspiro[4.5]decan-1-one

5



39A: Preparation of tert-butyl 2-(5-bromo-6-methoxypyridin-2-yl)-1-oxo-2,8-diazaspiro[4.5]decane-8-carboxylate




- 10 In a sealable reaction tube was mixed 3,6-dibromo-2-methoxypyridine (283 mg, 1.062 mmol), tert-butyl 1-oxo-2,8-diazaspiro[4.5]decane-8-carboxylate (135 mg, 0.531 mmol), cesium carbonate (346 mg, 1.062 mmol) and dioxane (1062 μ l). The reaction was purged with N_2 . Then 9,9-Dimethyl-4,5-bis(diphenylphosphino)xanthene (46.1 mg, 0.080 mmol), $Pd_2(dba)_3$ (24.30 mg, 0.027 mmol) was added and N_2 was bubbled through
- 15 reaction for 1 min. The reaction was sealed and stirred at 100 $^{\circ}C$ overnight. The reaction was partitioned between EtOAc (50 ml) and water (30 ml). The organic layer was separated, washed with water (2 x 30 ml) and brine (30 ml). dried over $MgSO_4$, filtered

and concentrated. The residue was purified using ISCO system (0-100% EtOAc/Hex gradient) to give tert-butyl 2-(5-bromo-6-methoxypyridin-2-yl)-1-oxo-2,8-diazaspiro[4.5]decane-8-carboxylate (225 mg, 0.511 mmol, 96 % yield) as a beige solid. MS (ESI) m/z : 441.9 (M+H)⁺.

5


39B: Preparation of tert-butyl 2-(6-methoxy-5-(1H-pyrazol-4-yl)pyridin-2-yl)-1-oxo-2,8-diazaspiro[4.5]decane-8-carboxylate



To a microwave vial was added 39A (230 mg, 0.522 mmol), 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-pyrazole (304 mg, 1.567 mmol), tripotassium phosphate (0.696 mL, 2.089 mmol), Dioxane (3 mL) and methanesulfonato(2-dicyclohexylphosphino-2',4',6'-tri-*i*-propyl-1,1'-biphenyl)(2'-amino-1,1'-biphenyl-2-yl)palladium(II) (22.11 mg, 0.026 mmol). The reaction was purged with nitrogen and sealed. The reaction was then subjected to microwave oven and stirred at 120 °C for 40 min. The reaction was partitioned between EtOAc (50 mL) and water (20 mL). The organic layer was separated, washed with water (20 mL) and brine (30 mL), dried over MgSO₄, filtered and concentrated. The residue was purified using ISCO system (0-100% EtOAc/Hex gradient) to give tert-butyl 2-(6-methoxy-5-(1H-pyrazol-4-yl)pyridin-2-yl)-1-oxo-2,8-diazaspiro[4.5]decane-8-carboxylate (200 mg, 0.468 mmol, 90 % yield) as an off-white solid. ¹H NMR (500 MHz, DMSO-*d*₆) δ 8.21 - 8.10 (m, 1H), 8.06 (d, *J*=8.0 Hz, 1H), 8.04 - 7.95 (m, 1H), 7.90 (d, *J*=8.3 Hz, 1H), 4.04 (t, *J*=7.2 Hz, 2H), 3.99 (s, 3H), 2.99 (br. s., 2H), 2.09 (t, *J*=7.0 Hz, 2H), 1.74 - 1.59 (m, 2H), 1.53 (d, *J*=13.2 Hz, 2H), 1.43 (s, 9H); MS (ESI) m/z : 427.2 (M+H)⁺.


25

39C:Preparation of 2-(6-methoxy-5-(1H-pyrazol-4-yl)pyridin-2-yl)-2,8-diazaspiro[4.5]decan-1-one, 3 HCl



To a round bottom flask was added 39B (100 mg, 0.234 mmol), MeOH. 4N HCl (0.178 mL, 5.85 mmol) in Dioxane (1 mL). The reaction was stirred at rt for 1 hr. The reaction was concentrated to give 2-(6-methoxy-5-(1H-pyrazol-4-yl)pyridin-2-yl)-2,8-diazaspiro[4.5]decan-1-one, 3 HCl (95 mg, 0.218 mmol, 93 % yield) as an off-white solid. MS (ESI) m/z : 328.1 (M+H)⁺.


39D:Preparation of 2-[6-methoxy-5-(1H-pyrazol-4-yl)pyridin-2-yl]-8-(4-methoxybenzoyl)-2,8-diazaspiro[4.5]decan-1-one



To a round bottom flask was added 39C (20 mg, 0.046 mmol) THF (1 mL), 4-methoxybenzoyl chloride (7.81 mg, 0.046 mmol) and Et₃N (0.038 mL, 0.275 mmol). The reaction was stirred at rt for 2 hr. The reaction was concentrated and purified using Purification Method A to give 2-[6-methoxy-5-(1H-pyrazol-4-yl)pyridin-2-yl]-8-(4-

methoxybenzoyl)-2,8-diazaspiro[4.5]decan-1-one (13.9 mg, 0.029 mmol, 63.1% yield).
¹H NMR (500 MHz, DMSO-d₆) δ 8.05 (d, J=8.1 Hz, 3H), 7.88 (d, J=8.2 Hz, 1H), 7.38 (d, J=8.4 Hz, 2H), 6.99 (d, J=8.4 Hz, 2H), 7.14 - 6.85 (m, 1H), 4.03 (t, J=6.7 Hz, 2H), 3.97 (s, 3H), 3.79 (s, 3H), 3.66 - 3.52 (m, 2H), 2.55 (s, 2H), 2.19 - 2.03 (m, 2H), 1.71 (br. s., 2H), 1.59 (br. s., 2H); MS (ESI) *m/z*: 462.1 (M+H)⁺; Anal. HPLC Retention time: 1.58 (Method 2); ROCK2 IC₅₀=4nM.

Example 40: Preparation of 7-(4-methoxybenzoyl)-2-[4-(1H-pyrazol-4-yl)phenyl]-2,7-diazaspiro[3.5]nonan-1-one




10

To a round bottom flask was added Intermediate 7 (10 mg, 0.028 mmol). THF (1 mL), 4-methoxybenzoyl chloride (4.80 mg, 0.028 mmol) and Et₃N (0.020 mL, 0.141 mmol). The reaction was stirred at rt for 2 hr. The reaction was concentrated and purified using Purification Method A to give 7-(4-methoxybenzoyl)-2-[4-(1H-pyrazol-4-yl)phenyl]-2,7-diazaspiro[3.5]nonan-1-one (6.7 mg, 0.015 mmol, 53.7% yield).
¹H NMR (500 MHz, DMSO-d₆) δ 8.14 (br. s., 1H), 7.90 (br. s., 1H), 7.62 (d, J=8.3 Hz, 2H), 7.43 - 7.27 (m, 4H), 6.99 (d, J=8.5 Hz, 2H), 3.79 (s, 3H), 3.69 - 3.37 (m, 3H), 3.17 (d, J=5.2 Hz, 1H), 2.58 - 2.50 (m, 2H), 1.88 (br. s., 4H); MS (ESI) *m/z*: 416.9 (M+H)⁺; Anal. HPLC Retention time: 1.51 (Method 1); ROCK2 IC₅₀=132nM.


20

Example 41: Preparation of 7-(3-methoxybenzoyl)-2-[4-(1H-pyrazol-4-yl)phenyl]-2,7-diazaspiro[3.5]nonan-1-one




7-(3-Methoxybenzoyl)-2-[4-(1H-pyrazol-4-yl)phenyl]-2,7-diazaspiro[3.5]nonan-1-one (6.8 mg, 0.016 mmol, 56.3% yield) was prepared in a similar manner as the procedure described in Example 40, using 3-methoxybenzoyl chloride (4.80 mg, 0.028 mmol). ¹H NMR (500 MHz, DMSO-d₆) δ 8.36 - 7.75 (m, 2H), 7.62 (d, J=8.3 Hz, 2H), 7.45 - 7.22 (m, 3H), 7.10 - 6.89 (m, 3H), 3.79 (s, 3H), 3.69 - 3.37 (m, 2H), 2.58 - 2.50 (m, 4H), 1.99 - 1.67 (m, 4H); MS (ESI) *m/z*: 417.1 (M+H)⁺; Anal. HPLC Retention time: 1.44 (Method 1); ROCK2 IC₅₀=285nM.

10 Example 42: Preparation of 7-benzoyl-2-[4-(1H-pyrazol-4-yl)phenyl]-2,7-diazaspiro[4.4]nonan-1-one




42A: Preparation of tert-butyl 7-(4-bromophenyl)-6-oxo-2,7-diazaspiro[4.4]nonane-2-carboxylate.



In a sealable reaction tube was mixed 1-bromo-4-iodobenzene (1177 mg, 4.16 mmol), tert-butyl 6-oxo-2,7-diazaspiro[4.4]nonane-2-carboxylate (500 mg, 2.081 mmol), cesium carbonate (1356 mg, 4.16 mmol) and dioxane (4161 μ l). The reaction was purged with N₂. Then 9,9-Dimethyl-4,5-bis(diphenylphosphino)xanthene (120 mg, 0.208 mmol), Pd₂(dba)₃ (95 mg, 0.104 mmol) was added and N₂ was bubbled through reaction for 1 min. The reaction was sealed and stirred at 100 °C overnight. The reaction was partitioned between EtOAc(50 ml) and water (30 ml). The organic layer was separated, washed with water (2 x 30 ml) and brine (30 ml). dried over MgSO₄, filtered and concentrated. The residue was purified using ISCO system (0-100% EtOAc/Hex gradient) to give tert-butyl 7-(4-bromophenyl)-6-oxo-2,7-diazaspiro[4.4]nonane-2-carboxylate (740 mg, 1.872 mmol, 90 % yield) as a beige solid. ¹H NMR (500 MHz, DMSO-d₆) δ 7.71 - 7.65 (m, 2H), 7.60 - 7.54 (m, 2H), 3.82 (t, *J*=6.9 Hz, 2H), 3.50 (ddd, *J*=10.6, 8.1, 4.4 Hz, 1H), 3.46 - 3.40 (m, 1H), 3.37 - 3.32 (m, 2H), 2.13 (t, *J*=6.7 Hz, 2H), 2.09 - 1.99 (m, 1H), 1.92 (d, *J*=7.4 Hz, 1H), 1.41 (s, 10H); MS (ESI) *m/z*: 397.0 (M+H)⁺.

42B: Preparation of tert-butyl 6-oxo-7-(4-(1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazol-4-yl)phenyl)-2,7-diazaspiro[4.4]nonane-2-carboxylate




To a microwave vial was added 42A (730 mg, 1.847 mmol), 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazole (1198 mg, 3.69 mmol), 3M tripotassium phosphate (2.462 mL, 7.39 mmol), Dioxane (10

mL) and methanesulfonato(2-dicyclohexylphosphino-2',4',6'-tri-*i*-propyl-1,1'-biphenyl)(2'-amino-1,1'-biphenyl-2-yl)palladium(II) (156 mg, 0.185 mmol). The reaction was purged with argon and sealed. The reaction was then subjected to microwave oven and stirred at 120 °C for 45 min. The reaction was partitioned between EtOAc(50 ml) and water (30 ml). The organic layer was separated, washed with water (30 ml) and brine (30 ml), dried over MgSO₄, filtered and concentrated. The residue was purified using ISCO system (0-100% EtOAc/Hex gradient) to give tert-butyl 6-oxo-7-(4-(1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazol-4-yl)phenyl)-2,7-diazaspiro[4.4]nonane-2-carboxylate (845 mg, 1.648 mmol, 89 % yield) as an off-white solid. ¹H NMR (500 MHz, DMSO-*d*₆) δ 8.36 (d, *J*=0.6 Hz, 1H), 8.01 (d, *J*=0.6 Hz, 1H), 7.76 - 7.70 (m, 2H), 7.69 - 7.57 (m, 2H), 5.45 (s, 2H), 3.88 (t, *J*=7.0 Hz, 2H), 3.67 - 3.58 (m, 2H), 3.57 - 3.51 (m, 1H), 3.48 (t, *J*=11.1 Hz, 1H), 3.37 (d, *J*=11.0 Hz, 2H), 2.17 (t, *J*=6.6 Hz, 2H), 2.13 - 2.02 (m, 1H), 1.96 (d, *J*=7.4 Hz, 1H), 1.45 (s, 9H), 0.94 - 0.82 (m, 2H), 0.03 - -0.07 (m, 9H); MS (ESI) *m/z*: 513.3 (M+H)⁺.

15


42C: Preparation of 2-(4-(1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazol-4-yl)phenyl)-2,7-diazaspiro[4.4]nonan-1-one



To a microwave vial was 42B (275 mg, 0.536 mmol), MeOH (2 mL) and H₂O (8 mL). The reaction was subjected to microwave oven and stirred at 150 °C for 90 min. The reaction was partitioned between EtOAc (40 ml) and water (10 ml). The organic layer was separated, washed with brine (20 ml), dried over MgSO₄, filtered and concentrated to give 2-(4-(1-((2-(trimethylsilyl)ethoxy)methyl)-1H-pyrazol-4-yl)phenyl)-2,7-diazaspiro[4.4]nonan-1-one (160 mg, 0.388 mmol, 72.3 % yield) as a beige solid. It was used without further purification. MS (ESI) *m/z*: 413.2 (M+H)⁺.

25


42D: Preparation of 7-benzoyl-2-[4-(1H-pyrazol-4-yl)phenyl]-2,7-diazaspiro[4.4]nonan-1-one.



To a round bottom flask was added 42C (28 mg, 0.068 mmol), THF (1.5 mL),
 5 benzoyl chloride (14.31 mg, 0.102 mmol) and Hunig's Base (0.047 mL, 0.271 mmol).
 The reaction was stirred at rt overnight. The reaction was concentrated. Then to the
 residue was added CH₂Cl₂(1 ml) and TFA(1 ml). The reaction was stirred at rt for 6 hr.
 The reaction was concentrated and purified using purification Method A to give 7-
 benzoyl-2-[4-(1H-pyrazol-4-yl)phenyl]-2,7-diazaspiro[4.4]nonan-1-one (12.3 mg, 0.032
 10 mmol, 46.4% yield). ¹H NMR (500MHz, DMSO-d₆) δ 8.27 - 7.80 (m, 2H), 7.73 - 7.66
 (m, 1H), 7.65 - 7.58 (m, 3H), 7.55 (br. s., 2H), 7.50 - 7.37 (m, 3H), 3.88 (t, J=6.6 Hz, 1H),
 3.84 - 3.71 (m, 2H), 3.69 - 3.52 (m, 2H), 3.42 (d, J=9.1 Hz, 1H), 2.27 - 1.92 (m, 4H); MS
 (ESI) *m/z*: 387.1 (M+H)⁺; Anal. HPLC Retention time: 1.34 (Method 1); ROCK2
 IC₅₀=1222nM.


15

Example 43:Preparation of 7-(2-methoxybenzoyl)-2-[4-(1H-pyrazol-4-yl)phenyl]-2,7-diazaspiro[4.4]nonan-1-one



7-(2-Methoxybenzoyl)-2-[4-(1H-pyrazol-4-yl)phenyl]-2,7-diazaspiro[4.4]nonan-1-one (9.1 mg, 0.021 mmol, 44.2% yield) was prepared in a similar manner as the procedure described in Example 42, using 2-methoxybenzoyl chloride (12.40 mg, 0.073 mmol). ¹H NMR (500 MHz, DMSO-d₆) δ 8.18 - 7.89 (m, 2H), 7.74 - 7.54 (m, 4H), 7.40 (q, *J*=7.7 Hz, 1H), 7.22 (br t, *J*=6.6 Hz, 1H), 7.09 (br t, *J*=9.2 Hz, 1H), 7.00 (q, *J*=7.5 Hz, 1H), 3.94 - 3.68 (m, 5H), 2.55 (s, 2H), 2.25 - 1.98 (m, 4H); MS (ESI) *m/z*: 416.9 (M+H)⁺; Anal. HPLC Retention time: 1.44 (Method 1); ROCK2 IC₅₀=2560nM.


10 **Example 44: Preparation of 7-(3-methoxybenzoyl)-2-[4-(1H-pyrazol-4-yl)phenyl]-2,7-diazaspiro[4.4]nonan-1-one**



7-(3-Methoxybenzoyl)-2-[4-(1H-pyrazol-4-yl)phenyl]-2,7-diazaspiro[4.4]nonan-1-one (6.6 mg, 0.016 mmol, 32.0% yield) was prepared in a similar manner as the procedure described in Example 42, using 3-methoxybenzoyl chloride (12.40 mg, 0.073 mmol). ¹H NMR (500 MHz, DMSO-d₆) δ 8.17 (br s, 1H), 7.91 (br d, *J*=6.1 Hz, 1H),

7.72 - 7.55 (m, 4H), 7.37 (q, $J=8.0$ Hz, 1H), 7.17 - 6.96 (m, 3H), 3.88 (br t, $J=6.5$ Hz, 1H), 3.83 - 3.69 (m, 4H), 3.67 - 3.54 (m, 2H), 2.28 - 1.91 (m, 4H); MS (ESI) m/z : 417.2 ($M+H$)⁺; Anal. HPLC Retention time: 1.41 (Method 1); ROCK2 IC₅₀=444nM.


5 **Example 45: Preparation of 7-(4-methoxybenzoyl)-2-[4-(1H-pyrazol-4-yl)phenyl]-2,7-diazaspiro[4.4]nonan-1-one**



7-(4-Methoxybenzoyl)-2-[4-(1H-pyrazol-4-yl)phenyl]-2,7-diazaspiro[4.4]nonan-1-one (10.6 mg, 0.025 mmol, 53.6% yield) was prepared in a similar manner as the procedure described in Example 42, using 4-methoxybenzoyl chloride (12.40 mg, 0.073 mmol). ¹H NMR (500 MHz, DMSO-d₆) δ 8.27 - 7.80 (m, 2H), 7.73 - 7.66 (m, 1H), 7.65 - 7.58 (m, 3H), 7.55 (br. s., 2H), 7.50 - 7.37 (m, 3H), 3.88 (t, $J=6.6$ Hz, 1H), 3.84 - 3.71 (m, 2H), 3.69 - 3.52 (m, 2H), 3.42 (d, $J=9.1$ Hz, 1H), 2.27 - 1.92 (m, 4H); MS (ESI) m/z : 417.2 ($M+H$)⁺; Anal. HPLC Retention time: 1.39 (Method 1), ROCK2 IC₅₀=180nM.


15

Example 46: Preparation of 7-(benzenesulfonyl)-2-[4-(1H-pyrazol-4-yl)phenyl]-2,7-diazaspiro[4.4]nonan-1-one



To a round bottom flask was added 42C (28 mg, 0.068 mmol), THF (2 mL), benzenesulfonyl chloride (17.98 mg, 0.102 mmol) and Hunig's Base (0.047 mL, 0.271 mmol). The reaction was stirred at rt overnight. The reaction was concentrated. Then
 5 the residue was dissolved in CH₂Cl₂ (1 mL) and TFA (1 mL). The reaction was stirred at rt for 6 hr. The reaction was concentrated and the residue was purified using purification Method B to give 7-(benzenesulfonyl)-2-[4-(1H-pyrazol-4-yl)phenyl]-2,7-diazaspiro[4.4]nonan-1-one (14.1 mg, 0.033 mmol, 48.7% yield). ¹H NMR (500MHz, DMSO-d₆) δ 8.02 (br. s., 2H), 7.86 (d, J=7.6 Hz, 2H), 7.78 - 7.73 (m, 1H), 7.71 - 7.64
 10 (m, 2H), 7.59 (s, 4H), 3.73 (t, J=6.7 Hz, 2H), 3.33 - 3.20 (m, 2H), 2.55 (s, 2H), 1.97 - 1.68 (m, 4H); MS (ESI) *m/z*: 423.2 (M+H)⁺; Anal. HPLC Retention time: 1.55 (Method 1); ROCK2 IC₅₀=750nM.


15 Example 47: Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-(4-methoxybenzoyl)-2,8-diazaspiro[4.5]decan-3-one




To a round bottom flask was added Intermediate 8, 2-(3-methoxy-4-(1H-pyrazol-4-yl)phenyl)-2,8-diazaspiro[4.5]decan-3-one, 2 HCl (20 mg, 0.050 mmol), THF (1 mL), 4-methoxybenzoyl chloride (8.54 mg, 0.050 mmol) and Et₃N (0.035 mL, 0.250 mmol). The reaction was stirred at rt for 15 min. The reaction was concentrated and purified using Purification Method B to give

2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-(4-methoxybenzoyl)-2,8-diazaspiro[4.5]decan-3-one (8.3 mg, 0.018 mmol, 35.3 % yield). ¹H NMR (500MHz, DMSO-d₆) δ 8.02 (br. s., 2H), 7.60 (d, J=8.3 Hz, 1H), 7.49 (s, 1H), 7.36 (d, J=8.3 Hz, 2H), 7.16 (d, J=8.8 Hz, 1H), 6.99 (d, J=8.4 Hz, 2H), 3.86 (s, 3H), 3.79 (s, 3H), 3.75 (br. s., 2H), 3.48 (d, J=12.9 Hz, 2H), 1.65 (br. s., 4H); MS (ESI) *m/z*: 461.1 (M+H)⁺; Anal. HPLC Retention time: 1.38 (Method 1); ROCK2 IC₅₀=306nM.

48: Preparation of 8-[4-(4-methylpiperazin-1-yl)benzoyl]-2-[3-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one




48A: Preparation of tert-butyl 2-(3-bromophenyl)-1-oxo-2,8-diazaspiro[4.5]decane-8-carboxylate



In a sealable reaction tube was mixed 1-bromo-3-iodobenzene (445 mg, 1.573 mmol), tert-butyl 1-oxo-2,8-diazaspiro[4.5]decane-8-carboxylate (200 mg, 0.786 mmol), Cesium carbonate (512 mg, 1.573 mmol) and dioxane (1573 μ l). The reaction was purged with N₂. Then 9,9-Dimethyl-4,5-bis(diphenylphosphino)xanthene (68.3 mg, 0.118 mmol), Pd₂(dba)₃ (36.0 mg, 0.039 mmol) was added and N₂ was bubbled through reaction for 1 min. The reaction was sealed and stirred at 100 °C overnight. The reaction was partitioned between EtOAc (50 ml) and water (30 ml). The organic layer was separated, washed with water (2 x 30 ml) and brine (30 ml), dried over MgSO₄, filtered and concentrated. The residue was purified using ISCO system (0-100% EtOAc/Hex gradient) to give tert-butyl 2-(3-bromophenyl)-1-oxo-2,8-diazaspiro[4.5]decane-8-carboxylate (300 mg, 0.733 mmol, 93 % yield) as a beige solid. ¹H NMR (500 MHz, DMSO-d₆) 8.13 - 7.93 (m, 1H), 7.61 (dt, *J*=6.7, 2.3 Hz, 1H), 7.42 - 7.24 (m, 2H), 3.93 - 3.74 (m, 4H), 3.09 - 2.82 (m, 2H), 2.09 (t, *J*=7.0 Hz, 2H), 1.62 (td, *J*=12.4, 4.1 Hz, 2H), 1.55 - 1.46 (m, 2H), 1.42 (s, 9H); MS (ESI) *m/z*: 409.0, 411.0 (M+H)⁺.


48B: Preparation of tert-butyl 2-(3-(1-(tert-butoxycarbonyl)-1H-pyrazol-4-yl)phenyl)-1-oxo-2,8-diazaspiro[4.5]decane-8-carboxylate



To a round bottle flask was added 48A (300 mg, 0.733 mmol), tert-butyl 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-pyrazole-1-carboxylate (431 mg, 1.466


mmol), tripotassium phosphate (0.733 mL, 2.199 mmol), Dioxane (3 mL) and chloro(2-dicyclohexylphosphino-2',4',6'-triisopropyl-1,1'-biphenyl)[2-(2'-amino-1,1'-biphenyl)]palladium(II) (40.4 mg, 0.051 mmol). The reaction was purged with nitrogen then stirred at 80 °C for 4hr. The reaction was partitioned between EtOAc (50 ml) and
 5 water (20 ml). The organic layer was separated, washed with water (20 ml) and brine (30 ml), dried over MgSO₄, filtered and concentrated. The residue was purified using ISCO system (0-100% EtOAc/Hex gradient) to give tert-butyl 2-(3-(1-(tert-butoxycarbonyl)-1H-pyrazol-4-yl)phenyl)-1-oxo-2,8-diazaspiro[4.5]decane-8-carboxylate (320 mg, 0.644 mmol, 88 % yield) as an off-white solid. ¹H NMR (500MHz, DMSO-d₆) 8.73 (s, 1H),
 10 8.29 (s, 1H), 7.89 (d, *J*=1.7 Hz, 1H), 7.71 (dd, *J*=8.3, 1.4 Hz, 1H), 7.53 (d, *J*=7.7 Hz, 1H), 7.46 - 7.32 (m, 1H), 3.91 - 3.86 (m, 4H), 3.00 (br. s., 2H), 2.11 (t, *J*=6.9 Hz, 2H), 1.68 - 1.60 (m, 11H), 1.51 (d, *J*=13.2 Hz, 2H), 1.42 (s, 9H); MS (ESI) *m/z*: 497.1 (M+H)⁺.

15 **48C: Preparation of 2-(3-(1H-pyrazol-4-yl)phenyl)-2,8-diazaspiro[4.5]decan-1-one, HCl**




To a round bottom flask was added 48B (310 mg, 0.624 mmol), Dioxane (10 mL) and 4N HCl (0.379 mL, 12.48 mmol) in dioxane. The reaction was stirred at rt overnight. The reaction was concentrated to give 2-(3-(1H-pyrazol-4-yl)phenyl)-2,8-diazaspiro[4.5]decan-1-one, HCl (221 mg, 0.631 mmol, 100 % yield) as an off-white
 20 solid. MS (ESI) *m/z*: 296.9 (M+H)⁺.

48D: Preparation of 8-[4-(4-methylpiperazin-1-yl)benzoyl]-2-[3-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one




To a round bottom flask was added 4-(4-methylpiperazin-1-yl)benzoic acid (8.95 mg, 0.041 mmol), DMF (0.5 mL), HATU (18.53 mg, 0.049 mmol) and Et₃N (0.028 mL, 0.203 mmol). The reaction was stirred at rt for 10 min and then 48C (15 mg, 0.041 mmol) was added and the reaction was continued for 2 hr. The reaction was purified using Purification Method A to give 8-[4-(4-methylpiperazin-1-yl)benzoyl]-2-[3-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one (18.9 mg, 0.037 mmol, 91% yield). ¹H NMR (500MHz, DMSO-d₆) 8.05 (br. s., 2H), 7.82 (s, 1H), 7.56 (d, *J*=7.8 Hz, 1H), 7.43 - 7.25 (m, 4H), 6.96 (d, *J*=8.5 Hz, 2H), 3.96 - 3.73 (m, 2H), 3.21 (br. s., 2H), 2.55 (s, 4H), 2.45 (br. s., 4H), 2.22 (s, 3H), 2.15 (t, *J*=6.4 Hz, 2H), 1.92 (br. s., 2H), 1.78 - 1.66 (m, 2H), 1.58 (br. s., 2H); MS (ESI) *m/z*: 499.1 (M+H)⁺; Anal. HPLC Retention time: 1.14 (Method 1); ROCK2 IC₅₀=1095nM.

Example 49: Preparation of 2-[3-(1H-indazol-6-yl)phenyl]-8-[4-(4-methylpiperazin-1-yl)benzoyl]-2,8-diazaspiro[4.5]decan-1-one




49A: Preparation of tert-butyl 2-(3-(1H-indazol-6-yl)phenyl)-1-oxo-2,8-diazaspiro[4.5]decane-8-carboxylate



To a round bottle flask was added 48A (140 mg, 0.342 mmol), (1H-indazol-6-yl)boronic acid (111 mg, 0.684 mmol), tripotassium phosphate (0.342 mL, 1.026 mmol), Dioxane (3 mL) and chloro(2-dicyclohexylphosphino-2',4',6'-triisopropyl-1,1'-biphenyl)[2-(2'-amino-1,1'-biphenyl)]palladium(II) (18.84 mg, 0.024 mmol). The reaction was purged with nitrogen then stirred at 80 °C for 4hr. The reaction was partitioned between EtOAc (50 mL) and water (20 mL). The organic layer was separated, washed with water (20 mL) and brine (30 mL), dried over MgSO₄, filtered and concentrated. The residue was purified using ISCO system (0-100% EtOAc/Hex gradient) to give tert-butyl 2-(3-(1H-indazol-6-yl)phenyl)-1-oxo-2,8-diazaspiro[4.5]decane-8-carboxylate (120 mg, 0.269 mmol, 79 % yield) as an off-white solid. ¹H NMR (500MHz, DMSO-d₆) 13.14 (s, 1H), 8.12 (d, *J*=1.1 Hz, 2H), 7.87 (d, *J*=8.6 Hz, 1H), 7.75 (s, 1H), 7.70 - 7.63 (m, 1H), 7.57 - 7.48 (m, 2H), 7.43 (dd, *J*=8.5, 1.4 Hz, 1H), 3.97 - 3.81 (m, 4H), 3.02 (br. s., 2H), 2.13 (t, *J*=6.9 Hz, 2H), 1.74 - 1.62 (m, 2H), 1.59 - 1.50 (m, 2H), 1.44 (s, 9H); MS (ESI) *m/z*: 447.0 (M+H)⁺.


49B: Preparation of 2-(3-(1H-indazol-6-yl)phenyl)-2,8-diazaspiro[4.5]decan-1-one,

HCl




To a round bottom flask was added 49A (120 mg, 0.269 mmol), Dioxane (10 mL) and 4N HCl (1.344 mL, 5.37 mmol) in dioxane. The reaction was stirred at rt overnight. The reaction was concentrated to give 2-(3-(1H-indazol-6-yl)phenyl)-2,8-diazaspiro[4.5]decan-1-one, HCl (100 mg, 0.261 mmol, 97 % yield) as an off-white solid. MS (ESI) m/z : 347.0 (M+H)⁺.

49C: Preparation of 2-[3-(1H-indazol-6-yl)phenyl]-8-[4-(4-methylpiperazin-1-yl)benzoyl]-2,8-diazaspiro[4.5]decan-1-one



To a 1 dram vial was added 4-(4-methylpiperazin-1-yl)benzoic acid (11.51 mg, 0.052 mmol), DMF (0.5 mL), HATU (23.83 mg, 0.063 mmol) and Et₃N (0.036 mL, 0.261 mmol). The reaction was stirred at rt for 10 min and then 49B (20 mg, 0.052 mmol) was added and the reaction was continued for 4 hr. The reaction was partitioned between EtOAc (20 ml) and water (15 ml). The organic layer was separated, washed with brine (15 ml), dried over MgSO₄, filtered and concentrated. The residue was purified using Purification Method A to give 2-[3-(1H-indazol-6-yl)phenyl]-8-[4-(4-methylpiperazin-1-yl)benzoyl]-2,8-diazaspiro[4.5]decan-1-one (19.3 mg, 0.034 mmol, 66.0% yield). ¹H NMR (500MHz, DMSO-d₆) 8.18 - 8.03 (m, 2H), 7.86 (d, J=8.4 Hz, 1H), 7.75 (s, 1H), 7.64 (br. s., 1H), 7.51 (br. s., 2H), 7.42 (d, J=8.4 Hz, 1H), 7.30 (d, J=8.4 Hz, 2H), 6.96 (d, J=8.5 Hz, 2H), 4.02 - 3.78 (m, 2H), 3.21 (br. s., 4H), 2.45 (br. s., 4H), 2.22 (s, 2H), 2.17 (t, J=6.6 Hz, 2H), 1.91 (s, 3H), 1.78 - 1.65 (m, 2H), 1.60 (br. s., 2H); MS (ESI) m/z : 549.2 (M+H)⁺; Anal. HPLC Retention time: 1.32 (Method 2); ROCK2 IC₅₀=6.8nM.


Example 50: Preparation of 2-[3-(1H-indazol-6-yl)phenyl]-8-(4-methoxybenzoyl)-2,8-diazaspiro[4.5]decan-1-one




To a round bottom flask was added 49B (20 mg, 0.052 mmol), DMF (0.5 mL), Et₃N (0.022 mL, 0.157 mmol) and 4-methoxybenzoyl chloride (8.91 mg, 0.052 mmol). The reaction was stirred at rt for 15 min. The reaction was purified using Purification
 5 Method A to give 2-[3-(1H-indazol-6-yl)phenyl]-8-(4-methoxybenzoyl)-2,8-diazaspiro[4.5]decan-1-one (14.6 mg, 0.029 mmol, 54.7% yield). ¹H NMR (500MHz, DMSO-d₆) 8.11 (s, 1H), 8.07 (br s, 1H), 7.86 (d, *J*=8.4 Hz, 1H), 7.75 (s, 1H), 7.64 (br s, 1H), 7.53 - 7.48 (m, 2H), 7.42 (br d, *J*=8.3 Hz, 1H), 7.38 (br d, *J*=8.3 Hz, 2H), 7.00 (d, *J*=8.5 Hz, 2H), 3.91 (br d, *J*=5.8 Hz, 2H), 3.79 (s, 3H), 3.57 - 3.53 (m, 3H), 3.27 - 3.09
 10 (m, 2H), 2.16 (br s, 2H), 1.73 (br s, 2H), 1.60 (br s, 2H); MS (ESI) *m/z*: 480.1 (M+H)⁺; Anal. HPLC Retention time: 1.85 (Method 1); ROCK2 IC₅₀=1680nM.

Example 51: Preparation of 2-[3-(1H-indazol-5-yl)phenyl]-8-[4-(4-methylpiperazin-1-yl)benzoyl]-2,8-diazaspiro[4.5]decan-1-one

15




51A: Preparation of tert-butyl 2-(3-(1H-indazol-5-yl)phenyl)-1-oxo-2,8-diazaspiro[4.5]decane-3-carboxylate



To a round bottle flask was added 48A (140 mg, 0.342 mmol), 5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1H-indazole (167 mg, 0.684 mmol), tripotassium phosphate (0.342 mL, 1.026 mmol), Dioxane (3 mL) and chloro(2-dicyclohexylphosphino-2',4',6'-triisopropyl-1,1'-biphenyl)[2-(2'-amino-1,1'-biphenyl)]palladium(II) (18.84 mg, 0.024 mmol). The reaction was purged with nitrogen then stirred at 100 °C for 4hr. The reaction was partitioned between EtOAc (50 ml) and water (20 ml). The organic layer was separated, washed with water (20 ml) and brine (30 ml), dried over MgSO₄, filtered and concentrated. The residue was purified using ISCO system (0-100% EtOAc/Hex gradient) to give tert-butyl 2-(3-(1H-indazol-5-yl)phenyl)-1-oxo-2,8-diazaspiro[4.5]decane-8-carboxylate (70 mg, 0.157 mmol, 45.8 % yield) as an off-white solid. ¹H NMR (500MHz, DMSO-d₆) 13.13 (s, 1H), 8.15 (s, 1H), 8.04 (d, *J*=7.2 Hz, 2H), 7.70 - 7.59 (m, 3H), 7.54 - 7.42 (m, 2H), 3.96 - 3.81 (m, 4H), 3.00 (br. s., 2H), 2.13 (t, *J*=6.9 Hz, 2H), 1.66 (td, *J*=12.4, 4.3 Hz, 2H), 1.53 (d, *J*=12.9 Hz, 2H), 1.43 (s, 9H); MS (ESI) *m/z*: 447.0 (M+H)⁺.


51B: Preparation of 2-(3-(1H-indazol-5-yl)phenyl)-2,8-diazaspiro[4.5]decan-1-one, HCl



To a round bottom flask was added 51A (70 mg, 0.157 mmol), Dioxane (1) and 4N HCl (784 µl, 3.14 mmol) in dioxane. The reaction was stirred at rt overnight. The reaction was concentrated to give 2-(3-(1H-indazol-5-yl)phenyl)-2,8-


diazaspiro[4.5]decan-1-one, HCl (60 mg, 0.157 mmol, 100 % yield) as an off-white solid. MS (ESI) m/z : 347.0 (M+H)⁺.

51C: Preparation of 2-[3-(1H-indazol-5-yl)phenyl]-8-[4-(4-methylpiperazin-1-yl)benzoyl]-2,8-diazaspiro[4.5]decan-1-one



To a 1 dram vial was added 4-(4-methylpiperazin-1-yl)benzoic acid (11.51 mg, 0.052 mmol), DMF (0.5 mL), HATU (23.83 mg, 0.063 mmol) and Et₃N (0.036 mL, 0.261 mmol). The reaction was stirred at rt for 10 min and then 51B (20 mg, 0.052 mmol) was added and the reaction was continued for 4 hr. The reaction was partitioned between EtOAc (20 ml) and water (15 ml). The organic layer was separated, washed with brine (15 ml), dried over MgSO₄, filtered and concentrated. The residue was purified using Purification Method A to give 2-[3-(1H-indazol-5-yl)phenyl]-8-[4-(4-methylpiperazin-1-yl)benzoyl]-2,8-diazaspiro[4.5]decan-1-one (19.1 mg, 0.034 mmol, 64.6% yield). ¹H NMR (500MHz, DMSO-d₆) 8.14 (s, 1H), 8.02 (br. s., 2H), 7.70 - 7.57 (m, 3H), 7.47 (d, J=4.6 Hz, 2H), 7.34 (d, J=8.5 Hz, 2H), 7.04 (d, J=8.5 Hz, 2H), 4.08 - 3.81 (m, 4H), 3.64 - 3.41 (m, 6H), 3.17 (s, 4H), 2.16 (br. s., 2H), 1.72 (d, J=10.7 Hz, 2H), 1.60 (br. s., 2H), 1.22 (s, 2H); MS (ESI) m/z : 549.1 (M+H)⁺; Anal. HPLC Retention time: 1.53 (Method 1); ROCK2 IC₅₀=14nM.

Example 52: Preparation of 2-[3-(1H-indazol-5-yl)phenyl]-8-(4-methoxybenzoyl)-2,8-diazaspiro[4.5]decan-1-one



2-[3-(1H-indazol-5-yl)phenyl]-8-(4-methoxybenzoyl)-2,8-diazaspiro[4.5]decan-1-one (10.1 mg, 0.021 mmol, 39.4% yield) was prepared in a similar manner as the procedure described in Example 51, using 51B (20 mg, 0.052 mmol). ¹H NMR (500
 5 MHz, DMSO-d₆) δ 8.15 (s, 1H), 8.03 (br. s., 2H), 7.72 - 7.57 (m, 3H), 7.47 (d, J=4.7 Hz, 2H), 7.38 (d, J=8.4 Hz, 2H), 7.00 (d, J=8.4 Hz, 2H), 3.92 (br. s., 2H), 3.79 (s, 3H), 3.17 (d, J=5.0 Hz, 2H), 2.16 (br. s., 2H), 1.73 (br. s., 2H), 1.60 (br. s., 2H); MS (ESI) *m/z*: 481.1 (M+H)⁺; Anal. HPLC Retention time: 1.66 (Method 2); ROCK2 IC₅₀=1875nM.

10 Preparation of Examples 53 - 205

The following compounds were made in a parallel manner using the following procedure: Reagents were weighed into stubby tubes. Stock solutions were made for reagent addition : Dissolved 2305.8 mg Intermediate 4 in 45.75 mL DMF and DIEA(2 eq). Dissolved 3131.85 mg HATU in 75 mL DMF. To each vial containing reagent acid
 15 (0.063 mmol) was added 0.25 ml of the HATU solution in DMF followed by DIEA (0.028 mL, 0.158 mmol). After stirring for 10 min, 0.25 ml of the Intermediate 4 solution was added and the resulting reaction mixture vials were placed in a Bohdan Miniblock XT and were agitated overnight at 400rpm.


Upon completion, reaction mixtures were purified via preparative LC/MS with the
 20 following conditions: Column: XBridge C18, 19 x 100 mm, 5-μm particles; Mobile Phase A: 5:95 acetonitrile: water with 10 mM ammonium acetate; Mobile Phase B: 95:5 acetonitrile: water with 10 mM ammonium acetate; Gradient: 10-50% B over 20 minutes, then a 2-minute hold at 100% B; Flow: 20 mL/min. Fractions containing the desired product were combined and dried via centrifugal evaporation. **Gradient varied for each
 25 reaction depending on polarity of compound.

Compound purity was assigned based on the methods below.

Method 1: Column: Waters Acquity UPLC BEH C18, 2.1 x 50 mm, 1.7- μ m particles; Mobile Phase A: 5:95 acetonitrile:water with 10 mM ammonium acetate; Mobile Phase B: 95:5 acetonitrile:water with 10 mM ammonium acetate; Temperature: 50 °C; Gradient: 0-100% B over 3 minutes, then a 0.75-minute hold at 100% B; Flow: 1.0 mL/min; Detection: UV at 220 nm.


Method 2: Column: Waters Acquity UPLC BEH C18, 2.1 x 50 mm, 1.7- μ m particles; Mobile Phase A: 5:95 acetonitrile:water with 0.1% trifluoroacetic acid; Mobile Phase B: 95:5 acetonitrile:water with 0.1% trifluoroacetic acid; Temperature: 50 °C; Gradient: 0-100% B over 3 minutes, then a 0.75-minute hold at 100% B; Flow: 1.0 mL/min; Detection: UV at 220 nm.

Example 53: Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-[4-(1H-pyrazol-3-yl)benzoyl]-2,8-diazaspiro[4.5]decan-1-one




¹H NMR (500 MHz, DMSO-*d*₆) δ 8.10 (br s, 1H), 7.94 (br s, 1H), 7.90 (br d, *J*=6.9 Hz, 2H), 7.81 (br s, 1H), 7.69 - 7.55 (m, 2H), 7.44 (br d, *J*=7.0 Hz, 2H), 7.15 (br d, *J*=7.0 Hz, 1H), 6.78 (br s, 1H), 4.41 - 4.29 (m, 1H), 3.87 (s, 5H), 3.69 (br d, *J*=9.3 Hz, 1H), 3.35 - 3.23 (m, 1H), 3.17 - 3.07 (m, 1H), 2.15 (br s, 2H), 1.85 - 1.48 (m, 4H); MS ESI *m/z* 497.2 (M+H); Anal. HPLC Retention time: 1.2 (Method 2); ROCK2 IC₅₀=16.5nM.

Example 54: Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-[5-(pyrrolidin-1-yl)pyridine-2-carbonyl]-2,8-diazaspiro[4.5]decan-1-one




¹H NMR (500 MHz, DMSO-d₆) δ 8.01 (br s, 2H), 7.89 (d, *J*=2.1 Hz, 1H), 7.65 - 7.57 (m, 2H), 7.50 (d, *J*=8.9 Hz, 1H), 7.15 (br d, *J*=8.2 Hz, 1H), 6.96 (dd, *J*=8.5, 2.4 Hz, 1H), 4.33 (br s, 1H), 4.20 (br d, *J*=7.0 Hz, 1H), 3.91 - 3.81 (m, 5H), 3.31 (br s, 4H), 3.17 (br s, 1H), 3.08 (br s, 1H), 2.16 (br s, 2H), 1.97 (br s, 4H), 1.83 - 1.71 (m, 2H), 1.68 - 1.43 (m, 2H); MS ESI *m/z* 501.3 (M+H); Anal. HPLC Retention time: 1.3 (Method 2); ROCK2 IC₅₀=2.1nM.

Example 55: Preparation of 4-{2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-1-oxo-2,8-diazaspiro[4.5]decane-8-carbonyl}benzene-1-sulfonamide



¹H NMR (500 MHz, DMSO-d₆) δ 8.02 (br s, 2H), 7.90 (d, *J*=8.2 Hz, 2H), 7.61 (br d, *J*=6.5 Hz, 4H), 7.15 (dd, *J*=8.5, 1.6 Hz, 1H), 4.34 (br d, *J*=12.8 Hz, 1H), 3.92 - 3.79 (m, 5H), 3.17 (br d, *J*=4.5 Hz, 1H), 2.21 - 2.08 (m, 2H), 1.83 - 1.63 (m, 3H), 1.51 (br s, 1H); MS ESI *m/z* 510 (M+H); Anal. HPLC Retention time: 1.07 (Method 1); ROCK2 IC₅₀=96nM.


Example 56: Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-(pyridine-4-carbonyl)-2,8-diazaspiro[4.5]decan-1-one



5 ^1H NMR (500 MHz, DMSO- d_6) δ 8.68 (br d, $J=5.0$ Hz, 2H), 8.02 (br s, 2H), 7.63 - 7.57 (m, 2H), 7.41 (br d, $J=5.5$ Hz, 2H), 7.15 (br d, $J=8.4$ Hz, 1H), 4.32 (br d, $J=13.4$ Hz, 1H), 3.87 (s, 5H), 3.38 (br s, 1H), 3.28 - 3.12 (m, 2H), 2.22 - 2.03 (m, 2H), 1.83 - 1.64 (m, 3H), 1.52 (br d, $J=14.0$ Hz, 1H); MS ESI m/z 432.1 (M+H); Anal. HPLC Retention time: 0.86 (Method 1); ROCK2 $\text{IC}_{50}=433\text{nM}$.

10


Example 57: Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-(3-phenoxybenzoyl)-2,8-diazaspiro[4.5]decan-1-one



15 ^1H NMR (500 MHz, DMSO- d_6) δ 7.66 - 7.58 (m, 2H), 7.51 - 7.40 (m, 4H), 7.23 - 7.05 (m, 7H), 6.95 (s, 1H), 4.35 - 4.22 (m, 1H), 3.86 (s, 5H), 3.40 (br s, 3H), 2.12 (br d,


$J=8.8$ Hz, 2H), 1.78 - 1.48 (m, 4H); Anal. HPLC Retention time: 1.84 Method 2; ROCK2
 $IC_{50}=118$ nM.

5 Example 58: Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-[3-(trifluoromethyl)benzoyl]-2,8-diazaspiro[4.5]decan-1-one



^1H NMR (500 MHz, DMSO- d_6) δ 8.01 (br s, 2H), 7.84 (br d, $J=6.8$ Hz, 1H), 7.78 - 7.68 (m, 3H), 7.63 - 7.55 (m, 2H), 7.14 (dd, $J=8.4$, 1.7 Hz, 1H), 4.38 - 4.20 (m, 1H),
 3.93 - 3.77 (m, 5H), 3.58 - 3.46 (m, 3H), 2.14 (br d, $J=5.5$ Hz, 2H), 1.83 - 1.61 (m, 3H),
 10 1.52 (br d, $J=9.6$ Hz, 1H); Anal. HPLC Retention time: 1.61 (Method 1); ROCK2
 $IC_{50}=144$ nM.


15 Example 59: Preparation of 8-(4-chlorobenzoyl)-2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one



¹H NMR (500 MHz, DMSO-d₆) δ 8.02 (br s, 2H), 7.65 - 7.58 (m, 2H), 7.56 - 7.50 (m, 2H), 7.44 (br d, *J*=8.3 Hz, 2H), 7.15 (dd, *J*=8.5, 1.6 Hz, 1H), 4.36 - 4.25 (m, 1H), 3.93 - 3.79 (m, 5H), 3.31 - 3.09 (m, 2H), 2.14 (br s, 2H), 1.80 - 1.46 (m, 4H); Anal. HPLC Retention time: 1.56 (Method 2); ROCK2 IC₅₀=67nM.


5

Example 60: Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-(4-phenoxybenzoyl)-2,8-diazaspiro[4.5]decan-1-one




¹H NMR (500 MHz, DMSO-d₆) δ 8.02 (br s, 2H), 7.64 - 7.57 (m, 2H), 7.48 - 7.39 (m, 3H), 7.33 (s, 1H), 7.21 (t, *J*=7.4 Hz, 1H), 7.15 (br d, *J*=8.6 Hz, 1H), 7.10 (br d, *J*=7.9 Hz, 2H), 7.03 (d, *J*=8.5 Hz, 2H), 4.36 - 4.18 (m, 1H), 3.91 - 3.81 (m, 5H), 3.50 - 3.39 (m, 1H), 3.34 - 3.07 (m, 1H), 3.02 - 2.87 (m, 1H), 2.14 (br s, 2H), 1.79 - 1.52 (m, 4H); Anal. HPLC Retention time: 1.8 (Method 1); ROCK2 IC₅₀=54nM.

Example 61: Preparation of 8-[[1,1'-biphenyl]-4-carbonyl]-2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one



¹H NMR (500 MHz, DMSO-d₆) δ 8.02 (br s, 2H), 7.76 (d, *J*=8.2 Hz, 2H), 7.72 (br d, *J*=7.5 Hz, 2H), 7.64 - 7.59 (m, 2H), 7.55 - 7.46 (m, 4H), 7.44 - 7.37 (m, 1H), 7.16 (dd, *J*=8.5, 1.7 Hz, 1H), 4.42 - 4.29 (m, 1H), 3.87 (s, 5H), 3.75 - 3.65 (m, 1H), 2.16 (br s, 2H),
 5 1.76 (br s, 2H), 1.56 (br d, *J*=5.6 Hz, 1H); MS ESI *m/z* 507.2 (M+H); Anal. HPLC Retention time: 1.77 (Method 1); ROCK2 IC₅₀=18nM.


Example 62: Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-(1-methyl-1H-pyrrole-2-carbonyl)-2,8-diazaspiro[4.5]decan-1-one



10


¹H NMR (500 MHz, DMSO-d₆) δ 8.02 (br s, 2H), 7.66 - 7.56 (m, 2H), 7.15 (dd, *J*=8.4, 1.6 Hz, 1H), 6.89 (s, 1H), 6.31 (br d, *J*=2.3 Hz, 1H), 6.04 (t, *J*=3.0 Hz, 1H), 4.22 (br d, *J*=13.0 Hz, 2H), 3.90 - 3.82 (m, 5H), 3.67 (s, 2H), 2.55 (s, 2H), 2.16 (br t, *J*=6.8 Hz, 2H), 1.79 - 1.67 (m, 2H), 1.60 (br d, *J*=13.0 Hz, 2H); MS ESI *m/z* 867.2 (M+H); Anal.
 15 HPLC Retention time: 1.35 (Method 2); ROCK2 IC₅₀=1430nM.

Example 63: Preparation of 8-(furan-2-carbonyl)-2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one




¹H NMR (500 MHz, DMSO-d₆) δ 8.02 (s, 2H), 7.85 (s, 1H), 7.62 (dd, *J*=5.0, 3.2 Hz, 2H), 7.16 (dd, *J*=8.5, 1.8 Hz, 1H), 7.00 (d, *J*=3.3 Hz, 1H), 6.64 (dd, *J*=3.2, 1.6 Hz, 1H), 4.23 (br d, *J*=13.3 Hz, 2H), 3.76 - 3.09 (m, 4H), 2.17 (br t, *J*=6.8 Hz, 2H), 1.76 (s, 2H), 1.63 (br d, *J*=13.6 Hz, 2H); MS ESI *m/z* 421.2 (M+H); Anal. HPLC Retention time: 1.24 Method 2; ROCK2 IC₅₀=1690nM.

10 **Example 64: Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-(naphthalene-2-carbonyl)-2,8-diazaspiro[4.5]decan-1-one**




15 ¹H NMR (500 MHz, DMSO-d₆) δ 8.07 - 7.94 (m, 6H), 7.65 - 7.57 (m, 4H), 7.53 (d, *J*=8.8 Hz, 1H), 7.16 (dd, *J*=8.5, 1.6 Hz, 1H), 4.44 - 4.35 (m, 1H), 4.27 (s, 1H), 3.93 - 3.81 (m, 5H), 2.16 (br s, 2H), 1.86 - 1.61 (m, 4H), 1.53 (br d, *J*=6.4 Hz, 1H); Anal. HPLC Retention time: 1.62 Method 1; ROCK2 IC₅₀=104nM.

Example 65: Preparation of 8-(furan-3-carbonyl)-2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one




¹H NMR (500 MHz, DMSO-d₆) δ 8.17 - 7.93 (m, 5H), 7.78 - 7.68 (m, 3H), 7.63 - 7.56 (m, 2H), 7.15 (br d, *J*=8.2 Hz, 1H), 4.52 (br d, *J*=6.1 Hz, 1H), 4.26 - 4.16 (m, 1H), 3.86 (br s, 5H), 3.48 (br s, 2H), 2.15 (br s, 2H), 1.77 - 1.67 (m, 2H), 1.59 (br d, *J*=12.1 Hz, 2H); MS ESI *m/z* 421.2 (M+H); Anal. HPLC Retention time: 1.31 (Method 1); ROCK2 IC₅₀=752nM.

Example 66: Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-(5-methylthiophene-2-carbonyl)-2,8-diazaspiro[4.5]decan-1-one




¹H NMR (500 MHz, DMSO-d₆) δ 8.02 (br s, 2H), 7.66 - 7.57 (m, 2H), 7.23 (d, *J*=3.4 Hz, 1H), 7.16 (br d, *J*=8.2 Hz, 1H), 6.83 (br d, *J*=2.8 Hz, 1H), 4.19 (br d, *J*=11.1 Hz, 2H), 3.93 - 3.80 (m, 5H), 2.55 (s, 2H), 2.48 (s, 3H), 2.16 (br t, *J*=6.6 Hz, 2H), 1.79 - 1.71 (m, 2H), 1.62 (br d, *J*=13.5 Hz, 2H); MS ESI *m/z* 451.2 (M+H); Anal. HPLC Retention time: 1.49 (Method 2); ROCK2 IC₅₀=372nM.

Example 67: Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-(thiophene-3-carbonyl)-2,8-diazaspiro[4.5]decan-1-one




¹H NMR (500 MHz, DMSO-d₆) δ 8.02 (br s, 2H), 7.80 (d, *J*=1.7 Hz, 1H), 7.68 - 7.56 (m, 3H), 7.22 (d, *J*=4.9 Hz, 1H), 7.15 (dd, *J*=8.4, 1.7 Hz, 1H), 4.41 - 4.21 (m, 2H), 3.87 (s, 5H), 3.44 - 3.11 (m, 2H), 2.15 (br s, 2H), 1.75 - 1.52 (m, 4H); MS ESI *m/z* 437 (M+H); Anal. HPLC Retention time: 1.3 (Method 2); ROCK2 IC₅₀=382nM.

Example 68: Preparation of 8-(1H-indole-2-carbonyl)-2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one



¹H NMR (500 MHz, DMSO-d₆) δ 11.56 (br s, 1H), 8.02 (br s, 2H), 7.61 (br dd, *J*=9.3, 6.1 Hz, 3H), 7.44 (br d, *J*=8.2 Hz, 1H), 7.24 - 7.11 (m, 2H), 7.05 (br t, *J*=7.3 Hz, 1H), 6.81 (s, 1H), 4.36 (br d, *J*=13.2 Hz, 2H), 3.92 - 3.80 (m, 4H), 3.74 - 3.58 (m, 3H), 2.18 (br t, *J*=6.1 Hz, 2H), 1.84 - 1.75 (m, 2H), 1.65 (br d, *J*=12.7 Hz, 2H); MS ESI *m/z* 470.3 (M+H); Anal. HPLC Retention time: 1.55 (Method 1); ROCK2 IC₅₀=467nM.


Example 69 : Preparation of 8-(2H-1,3-benzodioxole-5-carbonyl)-2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one



5 ^1H NMR (500 MHz, DMSO- d_6) δ 8.02 (br s, 2H), 7.67 - 7.57 (m, 2H), 7.15 (br d, $J=8.2$ Hz, 1H), 7.03 - 6.95 (m, 2H), 6.93 - 6.89 (m, 1H), 6.08 (s, 2H), 4.07 - 3.99 (m, 1H), 3.94 - 3.81 (m, 6H), 3.51 - 3.32 (m, 1H), 3.26 - 3.09 (m, 1H), 2.14 (br s, 2H), 1.78 - 1.66 (m, 2H), 1.63 - 1.46 (m, 2H); MS ESI m/z 474.9 ($M+H$); Anal. HPLC Retention time: 1.35 (Method 2); ROCK2 $\text{IC}_{50}=93\text{nM}$.

10


Example 70 : Preparation of 8-(1-benzofuran-2-carbonyl)-2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one



15 ^1H NMR (500 MHz, DMSO- d_6) δ 8.17 - 7.90 (m, 2H), 7.76 (d, $J=7.6$ Hz, 1H), 7.67 (d, $J=8.2$ Hz, 1H), 7.63 - 7.58 (m, 2H), 7.45 (t, $J=7.5$ Hz, 1H), 7.42 (s, 1H), 7.37 - 7.30 (m, 1H), 7.16 (br d, $J=8.4$ Hz, 1H), 4.93 (d, $J=5.0$ Hz, 1H), 4.70 (t, $J=5.6$ Hz, 1H), 3.91 - 3.87 (m, 5H), 3.17 (d, $J=5.2$ Hz, 2H), 2.19 (br t, $J=6.3$ Hz, 2H), 1.67 (br d, $J=13.5$


Hz, 2H), 1.50 (br t, $J=6.9$ Hz, 2H); MS ESI m/z 471.2 (M+H); Anal. HPLC Retention time: 1.57 Method 1; ROCK2 $IC_{50}=529$ nM.

5 Example 71 : Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-(pyrazine-2-carbonyl)-2,8-diazaspiro[4.5]decan-1-one



^1H NMR (500 MHz, DMSO- d_6) δ 8.86 (s, 1H), 8.75 (d, $J=2.3$ Hz, 1H), 8.69 (s, 1H), 8.02 (br s, 2H), 7.66 - 7.53 (m, 3H), 7.16 (br d, $J=8.4$ Hz, 1H), 4.37 (br d, $J=13.3$ Hz, 1H), 3.93 - 3.81 (m, 6H), 3.35 - 3.10 (m, 2H), 2.16 (dt, $J=12.9, 6.3$ Hz, 2H), 1.86 - 1.74 (m, 2H), 1.69 (br d, $J=13.7$ Hz, 1H), 1.54 (br d, $J=13.5$ Hz, 1H); MS ESI m/z 433 (M+H); Anal. HPLC Retention time: 1.03 Method 2; ROCK2 $IC_{50}=289$ nM.


10 Example 72 : Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-(6-methylpyridine-3-carbonyl)-2,8-diazaspiro[4.5]decan-1-one



15


¹H NMR (500 MHz, DMSO-d₆) δ 8.47 (s, 1H), 8.01 (br s, 2H), 7.74 (dd, *J*=8.0, 1.9 Hz, 1H), 7.64 - 7.56 (m, 2H), 7.35 (d, *J*=8.0 Hz, 1H), 7.17 - 7.10 (m, 1H), 4.03 (dd, *J*=11.2, 4.2 Hz, 1H), 3.90 - 3.80 (m, 5H), 3.61 (br s, 1H), 3.37 - 3.22 (m, 2H), 2.14 (br d, *J*=6.2 Hz, 2H), 1.89 (s, 3H), 1.80 - 1.61 (m, 3H), 1.57 - 1.44 (m, 2H); MS ESI *m/z* 446 (M+H); Anal. HPLC Retention time: 1.12 (Method 2); ROCK2 IC₅₀=219nM.

Example 73 : Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-(pyridine-3-carbonyl)-2,8-diazaspiro[4.5]decan-1-one



¹H NMR (500 MHz, DMSO-d₆) δ 8.69 - 8.64 (m, 1H), 8.62 (s, 1H), 8.02 (br s, 2H), 7.86 (br d, *J*=7.8 Hz, 1H), 7.65 - 7.57 (m, 2H), 7.50 (dd, *J*=7.6, 5.0 Hz, 1H), 7.15 (br d, *J*=8.4 Hz, 1H), 4.38 - 4.25 (m, 1H), 4.06 - 3.99 (m, 1H), 3.87 (s, 5H), 3.65 - 3.54 (m, 1H), 3.41 (br s, 1H), 2.15 (br d, *J*=6.0 Hz, 2H), 1.81 - 1.65 (m, 3H), 1.54 (br d, *J*=12.4 Hz, 1H); MS ESI *m/z* 432 (M+H); Anal. HPLC Retention time: 0.89 (Method 1); ROCK2 IC₅₀=414nM.


Example 74 : Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-(quinoline-3-carbonyl)-2,8-diazaspiro[4.5]decan-1-one



¹H NMR (500 MHz, DMSO-d₆) δ 8.92 (d, *J*=1.7 Hz, 1H), 8.50 (s, 1H), 8.09 (dd, *J*=11.8, 8.5 Hz, 2H), 8.02 (br s, 2H), 7.86 (t, *J*=7.3 Hz, 1H), 7.70 (t, *J*=7.4 Hz, 1H), 7.65 - 7.58 (m, 2H), 7.20 - 7.10 (m, 1H), 4.39 (br d, *J*=2.9 Hz, 1H), 3.92 - 3.82 (m, 5H), 3.71 (br d, *J*=9.3 Hz, 1H), 3.39 - 3.20 (m, 2H), 2.16 (br s, 2H), 1.82 (br s, 2H), 1.71 (br d, *J*=3.5 Hz, 1H), 1.61 - 1.51 (m, 1H)

MS ESI *m/z* 482.3 (M+H); Anal. HPLC Retention time: 1.16 (Method 1); ROCK2 IC₅₀=78nM.


10 **Example 75 : Preparation of 8-(4-chloro-2-fluorobenzoyl)-2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one**



15 ¹H NMR (500 MHz, DMSO-d₆) δ 8.02 (br s, 2H), 7.65 - 7.56 (m, 3H), 7.47 (br s, 1H), 7.43 - 7.38 (m, 1H), 7.14 (br d, *J*=8.6 Hz, 1H), 4.35 (br d, *J*=13.5 Hz, 1H), 3.92 -


3.81 (m, 5H), 3.29 - 3.11 (m, 1H), 2.55 (s, 2H), 2.22 - 2.08 (m, 2H), 1.79 - 1.61 (m, 3H), 1.52 (br d, $J=10.7$ Hz, 1H); MS ESI m/z 483.18, 483.06 ($M+H$); ROCK2 $IC_{50}=50$ nM.

5 Example 76 : Preparation of 8-(1-benzothiophene-2-carbonyl)-2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one



^1H NMR (500 MHz, DMSO- d_6) δ 8.07 - 7.98 (m, 3H), 7.97 - 7.91 (m, 1H), 7.75 (s, 1H), 7.66 - 7.58 (m, 2H), 7.50 - 7.41 (m, 2H), 7.16 (dd, $J=8.4, 1.8$ Hz, 1H), 4.36 - 4.10 (m, 2H), 3.92 - 3.83 (m, 5H), 3.56 - 3.41 (m, 2H), 2.18 (br t, $J=6.6$ Hz, 2H), 1.85 - 1.75 (m, 2H), 1.66 (br d, $J=13.4$ Hz, 2H); MS ESI m/z 487.1 ($M+H$); Anal. HPLC Retention time: 1.68 (Method 2); ROCK2 $IC_{50}=27$ nM.


15 Example 77 : Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-(1,2,3-thiadiazole-4-carbonyl)-2,8-diazaspiro[4.5]decan-1-one



^1H NMR (500 MHz, DMSO- d_6) δ 9.57 (s, 1H), 8.02 (br s, 2H), 7.67 - 7.53 (m, 2H), 7.16 (dd, $J=8.4, 1.6$ Hz, 1H), 4.42 (br d, $J=13.5$ Hz, 1H), 3.93 - 3.84 (m, 5H), 3.42 - 3.21 (m, 2H), 3.17 (d, $J=5.1$ Hz, 1H), 2.22 - 2.10 (m, 2H), 1.85 - 1.75 (m, 2H), 1.71 (br d, $J=13.0$ Hz, 1H), 1.59 (br d, $J=12.6$ Hz, 1H); MS ESI m/z 439.2 (M+H); Anal. HPLC

5 Retention time: 1.13 (Method 1); ROCK2 $\text{IC}_{50}=453\text{nM}$.


Example 78 : Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-[2-(pyridin-3-yl)-1,3-thiazole-4-carbonyl]-2,8-diazaspiro[4.5]decan-1-one



10 ^1H NMR (500 MHz, DMSO- d_6) δ 9.17 (s, 1H), 8.70 (br d, $J=3.9$ Hz, 1H), 8.34 (br d, $J=7.9$ Hz, 1H), 8.26 (s, 1H), 8.02 (br s, 2H), 7.69 - 7.52 (m, 3H), 7.16 (br d, $J=8.3$ Hz, 1H), 4.39 (br d, $J=10.4$ Hz, 1H), 4.18 (br d, $J=11.9$ Hz, 1H), 3.95 - 3.81 (m, 5H), 3.74 (br d, $J=4.1$ Hz, 1H), 3.17 (br d, $J=4.3$ Hz, 1H), 2.19 (br d, $J=6.1$ Hz, 2H), 1.88 - 1.73 (m, 2H), 1.72 - 1.55 (m, 2H)


15 MS ESI m/z 515 (M+H); Anal. HPLC Retention time: 1.26 Method 2; ROCK2 $\text{IC}_{50}=76\text{nM}$.

Example 79 : Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-(5-methylpyrazine-2-carbonyl)-2,8-diazaspiro[4.5]decan-1-one




¹H NMR (500 MHz, DMSO-d₆) δ 8.72 (s, 1H), 8.57 (s, 1H), 8.02 (br s, 2H), 7.65 - 7.55 (m, 2H), 7.15 (dd, *J*=8.4, 1.6 Hz, 1H), 4.36 (br d, *J*=13.0 Hz, 1H), 3.96 - 3.80 (m, 5H), 3.74 (br d, *J*=13.3 Hz, 1H), 3.42 (br s, 1H), 3.21 - 3.12 (m, 1H), 2.51 (br s, 3H), 2.24 - 2.09 (m, 2H), 1.84 - 1.73 (m, 2H), 1.71 - 1.62 (m, 1H), 1.53 (br d, *J*=13.5 Hz, 1H); MS ESI *m/z* 447.1 (M+H); Anal. HPLC Retention time: 1.11 (Method 2); ROCK2 IC₅₀=110nM.

Example 30 : Preparation of 8-(2,6-dimethoxypyridine-3-carbonyl)-2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one



¹H NMR (500 MHz, DMSO-d₆) δ 8.02 (br s, 2H), 7.61 (br d, *J*=8.4 Hz, 3H), 7.14 (br d, *J*=8.3 Hz, 1H), 6.45 (d, *J*=8.0 Hz, 1H), 4.34 (br d, *J*=13.4 Hz, 1H), 4.10 - 3.99 (m, 1H), 3.94 - 3.82 (m, 11H), 3.17 (s, 1H), 3.12 - 3.03 (m, 1H), 2.14 (br d, *J*=5.5 Hz, 2H), 1.77 - 1.68 (m, 2H), 1.64 - 1.61 (m, 1H), 1.52 - 1.43 (m, 1H); MS ESI *m/z* 492.2 (M+H); Anal. HPLC Retention time: 1.44 (Method 1); ROCK2 IC₅₀=99nM.

Example 81 : Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-{5-oxo-2H,3H,5H-[1,3]thiazolo[3,2-a]pyrimidine-6-carbonyl}-2,8-diazaspiro[4.5]decan-1-one




5

^1H NMR (500 MHz, DMSO- d_6) δ 8.02 (br s, 2H), 7.85 (s, 1H), 7.65 - 7.57 (m, 2H), 7.15 (br d, $J=8.3$ Hz, 1H), 4.40 (br t, $J=7.9$ Hz, 2H), 4.29 (br d, $J=13.3$ Hz, 1H), 4.03 (dd, $J=11.1, 4.0$ Hz, 1H), 3.90 - 3.81 (m, 5H), 3.61 - 3.52 (m, 1H), 3.06 (br t, $J=11.0$ Hz, 1H), 2.19 - 2.09 (m, 2H), 1.81 - 1.66 (m, 2H), 1.60 (br d, $J=13.9$ Hz, 1H), 1.65 - 1.57 (m, 1H), 1.54 - 1.49 (m, 1H), 1.51 (br d, $J=13.0$ Hz, 1H); MS ESI m/z 1013.2 (M+H); Anal. HPLC Retention time: 1.01 (Method 2); ROCK2 $\text{IC}_{50}=57\text{nM}$.

10


Example 82 : Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-(4-methylthiophene-2-carbonyl)-2,8-diazaspiro[4.5]decan-1-one



15


¹H NMR (500 MHz, DMSO-d₆) δ 8.02 (br s, 2H), 7.66 - 7.58 (m, 2H), 7.34 (s, 1H), 7.25 (s, 1H), 7.16 (dd, *J*=8.5, 1.6 Hz, 1H), 4.19 (br d, *J*=11.4 Hz, 2H), 3.94 - 3.84 (m, 5H), 3.62 (br d, *J*=5.1 Hz, 1H), 3.37 (br s, 1H), 2.24 (s, 3H), 2.16 (br t, *J*=6.8 Hz, 2H), 1.81 - 1.70 (m, 2H), 1.69 - 1.55 (m, 3H); MS ESI *m/z* 451.2 (M+H); Anal. HPLC Retention time: 1.47 (Method 1); ROCK2 IC₅₀=109nM.

Example 83 : Preparation of 8-(1,3-benzothiazole-6-carbonyl)-2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one




¹H NMR (500 MHz, DMSO-d₆) δ 9.47 (s, 1H), 8.28 (s, 1H), 8.15 (d, *J*=8.3 Hz, 1H), 8.02 (br s, 2H), 7.64 - 7.53 (m, 3H), 7.15 (br d, *J*=8.4 Hz, 1H), 4.44 - 4.32 (m, 1H), 3.86 (s, 5H), 3.52 (br s, 3H), 2.15 (br s, 2H), 1.72 - 1.62 (m, 1H), 1.57 - 1.44 (m, 1H), 1.25 - 1.05 (m, 2H); MS ESI *m/z* 488.1 (M+H); Anal. HPLC Retention time: 1.23 (Method 1); ROCK2 IC₅₀=26nM.

Example 84 : Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-(2-phenyl-1,3-thiazole-4-carbonyl)-2,8-diazaspiro[4.5]decan-1-one




¹H NMR (500 MHz, DMSO-d₆) δ 8.55 - 8.46 (m, 1H), 8.15 (s, 1H), 8.06 (br s, 1H), 8.00 - 7.93 (m, 2H), 7.65 - 7.58 (m, 2H), 7.53 (br d, *J*=5.4 Hz, 3H), 7.15 (br d, *J*=6.9 Hz, 1H), 4.38 (br d, *J*=9.8 Hz, 1H), 4.20 (br d, *J*=11.4 Hz, 1H), 3.86 (s, 5H), 3.61 - 3.48 (m, 1H), 3.46 - 3.34 (m, 1H), 2.18 (br d, *J*=4.0 Hz, 2H), 1.91 - 1.73 (m, 2H), 1.72 - 1.53 (m, 2H); MS ESI *m/z* 514.3 (M+H); Anal. HPLC Retention time: 1.65 (Method 1); ROCK2 IC₅₀=138nM.

Example 85 : Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-(5-phenylfuran-2-carbonyl)-2,8-diazaspiro[4.5]decan-1-one




¹H NMR (500 MHz, DMSO-d₆) δ 8.09 - 7.95 (m, 2H), 7.78 (br d, *J*=7.7 Hz, 2H), 7.62 (br d, *J*=6.1 Hz, 2H), 7.48 (t, *J*=7.6 Hz, 2H), 7.38 (br d, *J*=7.3 Hz, 1H), 7.20 - 7.10 (m, 3H), 4.32 (br d, *J*=13.4 Hz, 2H), 4.23 - 4.14 (m, 1H), 4.03 (dd, *J*=11.3, 4.0 Hz, 1H), 3.92 - 3.86 (m, 5H), 2.19 (br t, *J*=6.8 Hz, 2H), 1.91 - 1.77 (m, 2H), 1.68 (br d, *J*=12.8 Hz, 2H); MS ESI *m/z* 497.1 (M+H); Anal. HPLC Retention time: 1.69 (Method 1); ROCK2 IC₅₀=851nM.

Example 86 : Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-(1,3-thiazole-4-carbonyl)-2,8-diazaspiro[4.5]decan-1-one



5 ^1H NMR (500 MHz, DMSO- d_6) δ 9.14 (s, 1H), 8.13 (d, $J=1.4$ Hz, 1H), 8.08 - 7.91 (m, 2H), 7.66 - 7.53 (m, 2H), 7.14 (br d, $J=8.3$ Hz, 1H), 4.35 (br d, $J=11.6$ Hz, 1H), 4.11 - 3.99 (m, 1H), 3.92 - 3.79 (m, 3H), 3.66 (br s, 2H), 3.33 (br s, 1H), 3.14 - 3.04 (m, 1H), 2.16 (br d, $J=7.2$ Hz, 2H), 1.75 (br t, $J=10.9$ Hz, 2H), 1.65 (br d, $J=13.7$ Hz, 1H), 1.55 (br d, $J=11.8$ Hz, 1H); MS ESI m/z 438 ($M+H$); Anal. HPLC Retention time: 1.09 (Method
10 2); ROCK2 $\text{IC}_{50}=443\text{nM}$.


Example 87 : Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-[6-(trifluoromethyl)pyridine-3-carbonyl]-2,8-diazaspiro[4.5]decan-1-one



15 ^1H NMR (500 MHz, DMSO- d_6) δ 8.83 (s, 1H), 8.16 (br d, $J=7.4$ Hz, 1H), 8.01 (br d, $J=8.0$ Hz, 3H), 7.65 - 7.56 (m, 2H), 7.20 - 7.09 (m, 1H), 4.34 (br d, $J=11.2$ Hz, 1H),


4.03 (dd, $J=11.1, 4.2$ Hz, 1H), 3.94 - 3.79 (m, 5H), 3.37 - 3.17 (m, 2H), 2.14 (dt, $J=13.1, 6.8$ Hz, 2H), 1.86 - 1.65 (m, 3H), 1.54 (br d, $J=13.3$ Hz, 1H); MS ESI m/z 500.2 ($M+H$); Anal. HPLC Retention time: 1.42 (Method 1); ROCK2 $IC_{50}=557$ nM.

5 **Example 88 : Preparation of N-(4-{2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-1-oxo-2,8-diazaspiro[4.5]decane-8-carbonyl}phenyl)acetamide**




^1H NMR (500 MHz, DMSO- d_6) δ 10.16 (s, 1H), 8.02 (br s, 2H), 7.70 - 7.56 (m, 4H), 7.36 (d, $J=8.4$ Hz, 2H), 7.14 (br d, $J=8.5$ Hz, 1H), 4.37 - 4.22 (m, 2H), 3.86 (s, 5H),
 10 3.50 (br s, 1H), 3.32 - 3.04 (m, 1H), 2.14 (br s, 2H), 2.06 (s, 3H), 1.78 - 1.65 (m, 2H),
 1.65 - 1.47 (m, 2H); MS ESI m/z 488 ($M+H$); Anal. HPLC Retention time: 1.1 (Method 2); ROCK2 $IC_{50}=73$ nM.

15 **Example 89 : Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-(thiophene-2-carbonyl)-2,8-diazaspiro[4.5]decan-1-one**



¹H NMR (500 MHz, DMSO-d₆) δ 8.19 - 7.88 (m, 2H), 7.76 (d, *J*=5.0 Hz, 1H), 7.66 - 7.57 (m, 2H), 7.42 (d, *J*=3.4 Hz, 1H), 7.19 - 7.09 (m, 2H), 4.19 (br s, 2H), 3.91 - 3.83 (m, 5H), 3.52 - 3.37 (m, 2H), 2.16 (br t, *J*=6.8 Hz, 2H), 1.82 - 1.70 (m, 2H), 1.62 (br d, *J*=13.3 Hz, 2H); MS ESI *m/z* 437.2 (M+H); Anal. HPLC Retention time: 1.33 (Method 1); ROCK2 IC₅₀=117nM.

Example 90: Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-(4-methoxyisquinoline-1-carbonyl)-2,8-diazaspiro[4.5]decan-1-one




10

¹H NMR (500 MHz, DMSO-d₆) δ 8.27 - 8.16 (m, 2H), 8.01 (br s, 2H), 7.93 (br d, *J*=8.2 Hz, 1H), 7.84 (br d, *J*=7.9 Hz, 1H), 7.78 (br d, *J*=7.6 Hz, 1H), 7.60 (br d, *J*=5.2 Hz, 2H), 7.14 (br d, *J*=7.6 Hz, 1H), 4.52 (br d, *J*=13.1 Hz, 1H), 4.11 (s, 3H), 3.93 - 3.80 (m, 5H), 3.35 - 3.23 (m, 2H), 3.16 (br t, *J*=11.3 Hz, 1H), 2.26 - 2.17 (m, 1H), 2.16 - 2.05 (m, 1H), 1.94 - 1.81 (m, 1H), 1.74 (br d, *J*=13.1 Hz, 1H), 1.67 (br t, *J*=10.2 Hz, 1H), 1.43 (br d, *J*=13.1 Hz, 1H); MS ESI *m/z* 512 (M+H); Anal. HPLC Retention time: 1.48 (Method 1); ROCK2 IC₅₀=127nM.


15

Example 91 : Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-[(1R,2R)-2-phenylcyclopropanecarbonyl]-2,8-diazaspiro[4.5]decan-1-one



- 5 ¹H NMR (500 MHz, DMSO-d₆) δ 8.02 (br s, 2H), 7.63 - 7.53 (m, 2H), 7.30 - 7.23 (m, 2H), 7.19 (br d, *J*=7.4 Hz, 3H), 7.13 (br t, *J*=7.1 Hz, 1H), 4.24 (br s, 1H), 4.13 (br t, *J*=14.2 Hz, 1H), 3.88 - 3.80 (m, 5H), 3.56 - 3.44 (m, 2H), 3.39 - 3.25 (m, 1H), 2.93 (br d, *J*=9.9 Hz, 1H), 2.30 (dt, *J*=19.3, 7.7 Hz, 2H), 2.13 (br s, 2H), 1.68 - 1.54 (m, 3H), 1.43 (br dd, *J*=9.0, 4.3 Hz, 1H); MS ESI *m/z* 471 (M+H); Anal. HPLC Retention time: 1.62
- 10 (Method 1); ROCK2 IC₅₀=436nM.


Example 92 : Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-[4-(propan-2-yl)benzoyl]-2,8-diazaspiro[4.5]decan-1-one



- 15 ¹H NMR (500 MHz, DMSO-d₆) δ 8.02 (br s, 2H), 7.61 (dd, *J*=4.9, 3.3 Hz, 2H), 7.33 (s, 3H), 7.15 (dd, *J*=8.5, 1.4 Hz, 1H), 4.37 - 4.29 (m, 1H), 4.21 (br d, *J*=11.8 Hz, 1H), 3.87 (s, 4H), 3.50 - 3.38 (m, 1H), 3.27 - 3.09 (m, 1H), 2.93 (quin, *J*=6.9 Hz, 1H),


2.15 (br d, $J=6.3$ Hz, 2H), 1.80 - 1.47 (m, 4H), 1.22 (d, $J=7.0$ Hz, 6H); MS ESI m/z 473.1 (M+H); Anal. HPLC Retention time: 1.76 Method 2; ROCK2 $IC_{50}=12$ nM.

5 Example 93 : Preparation of 8-{6-bromopyrazolo[1,5-a]pyrimidine-2-carbonyl}-2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one



^1H NMR (500 MHz, DMSO- d_6) δ 9.62 (d, $J=1.2$ Hz, 1H), 8.70 (d, $J=2.0$ Hz, 1H), 8.19 - 7.88 (m, 2H), 7.70 - 7.53 (m, 2H), 7.15 (dd, $J=8.4, 1.6$ Hz, 1H), 7.02 (s, 1H), 4.39 (br d, $J=13.0$ Hz, 1H), 4.13 (br d, $J=13.8$ Hz, 1H), 3.94 - 3.77 (m, 5H), 3.42 - 3.31 (m, 1H), 3.22 - 3.16 (m, 1H), 2.21 - 2.10 (m, 2H), 1.82 - 1.72 (m, 2H), 1.71 - 1.64 (m, 1H), 1.56 (br d, $J=13.0$ Hz, 1H); MS ESI m/z 550.1 (M+H); Anal. HPLC Retention time: 1.35 (Method 1); ROCK2 $IC_{50}=5$ nM.


15 Example 94 : Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-[4-(morpholin-4-yl)benzoyl]-2,8-diazaspiro[4.5]decan-1-one



¹H NMR (500 MHz, DMSO-d₆) δ 8.02 (br s, 2H), 7.68 - 7.56 (m, 2H), 7.31 (br d, *J*=8.6 Hz, 2H), 7.20 - 7.10 (m, 1H), 6.98 (br d, *J*=8.7 Hz, 2H), 4.08 - 3.99 (m, 2H), 3.92 - 3.81 (m, 5H), 3.78 - 3.66 (m, 4H), 3.37 (br s, 2H), 3.22 - 3.11 (m, 4H), 2.14 (br t, *J*=6.6 Hz, 2H), 1.71 (br d, *J*=9.6 Hz, 2H), 1.59 (br s, 3H); MS ESI *m/z* 516 (M+H); Anal. HPLC Retention time: 1.33 (Method 2); ROCK2 IC₅₀=2nM.

Example 95 : Preparation of 8-[4-(1H-imidazol-1-yl)benzoyl]-2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one


10



¹H NMR (500 MHz, DMSO-d₆) δ 8.35 (s, 1H), 8.02 (br s, 2H), 7.83 (s, 1H), 7.76 (br d, *J*=8.3 Hz, 2H), 7.61 (br d, *J*=7.3 Hz, 2H), 7.57 (br d, *J*=8.4 Hz, 2H), 7.18 - 7.11 (m, 2H), 4.35 (br d, *J*=11.6 Hz, 2H), 3.87 (s, 5H), 3.72 - 3.60 (m, 2H), 2.16 (br s, 2H), 1.89 -


1.72 (m, 2H), 1.67 (br d, $J=8.6$ Hz, 1H), 1.55 (br d, $J=6.0$ Hz, 1H); MS ESI m/z 497 (M+H); Anal. HPLC Retention time: 1.16 (Method 2); ROCK2 $IC_{50}=13$ nM.

5 Example 96 : Preparation of 4-{2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-1-oxo-2,8-diazaspiro[4.5]decan-8-carbonyl}pyridin-1-ium-1-olate




^1H NMR (500 MHz, DMSO- d_6) δ 8.27 (d, $J=6.7$ Hz, 2H), 8.02 (br s, 2H), 7.60 (d, $J=8.4$ Hz, 1H), 7.57 (d, $J=1.4$ Hz, 1H), 7.49 (d, $J=6.7$ Hz, 2H), 7.14 (dd, $J=8.5, 1.6$ Hz, 1H), 4.02 (dd, $J=11.1, 4.0$ Hz, 2H), 3.89 - 3.81 (m, 5H), 3.31 (dt, $J=17.1, 6.3$ Hz, 1H),
 10 3.16 (br d, $J=4.9$ Hz, 1H), 2.13 (br d, $J=4.6$ Hz, 2H), 1.75 (br d, $J=14.5$ Hz, 2H), 1.69 - 1.59 (m, 1H), 1.59 - 1.44 (m, 1H); MS ESI m/z 448 (M+H); Anal. HPLC Retention time: 0.87 (Method 1); ROCK2 $IC_{50}=1095$ nM.

15 Example 97 : Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-(1,2-oxazole-5-carbonyl)-2,8-diazaspiro[4.5]decan-1-one




¹H NMR (500 MHz, DMSO-d₆) δ 8.01 (br s, 2H), 7.60 (br d, *J*=8.4 Hz, 1H), 7.15 (br dd, *J*=16.7, 8.0 Hz, 2H), 6.86 (br d, *J*=8.5 Hz, 1H), 4.46 (br d, *J*=6.1 Hz, 1H), 4.35 (br d, *J*=3.2 Hz, 1H), 3.75 (br s, 5H), 3.43 - 3.27 (m, 2H), 2.15 (br d, *J*=5.8 Hz, 2H), 1.80 - 1.70 (m, 3H), 1.63 (br s, 1H); MS ESI *m/z* 422.2 (M+H); Anal. HPLC Retention time: 1.13 (Method 1); ROCK2 IC₅₀=427nM.

Example 98 : Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-[5-(trifluoromethyl)pyridine-2-carbonyl]-2,8-diazaspiro[4.5]decan-1-one




¹H NMR (500 MHz, DMSO-d₆) δ 9.02 (s, 1H), 8.37 (br d, *J*=7.9 Hz, 1H), 8.02 (br s, 2H), 7.82 (d, *J*=8.2 Hz, 1H), 7.65 - 7.56 (m, 2H), 7.16 (br d, *J*=8.3 Hz, 1H), 4.37 (br d, *J*=13.1 Hz, 1H), 3.96 - 3.80 (m, 5H), 3.59 (br d, *J*=12.7 Hz, 1H), 3.27 - 3.16 (m, 2H), 2.25 - 2.09 (m, 2H), 1.87 - 1.74 (m, 2H), 1.68 - 1.65 (m, 1H), 1.52 (br d, *J*=13.8 Hz, 1H); MS ESI *m/z* 500.2 (M+H); Anal. HPLC Retention time: 1.43 (Method 1); ROCK2 IC₅₀=259nM.

Example 99 : Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-(2-methoxypyrimidine-5-carbonyl)-2,8-diazaspiro[4.5]decan-1-one



¹H NMR (500 MHz, DMSO-d₆) δ 8.72 (s, 2H), 8.16 - 7.89 (m, 2H), 7.66 - 7.55 (m, 2H), 7.15 (dd, *J*=8.4, 1.6 Hz, 1H), 4.30 (br d, *J*=8.2 Hz, 1H), 3.97 (s, 3H), 3.87 (s, 5H), 3.73 - 3.62 (m, 1H), 3.49 - 3.32 (m, 1H), 3.17 (br d, *J*=5.2 Hz, 1H), 2.14 (br s, 2H), 1.77 (br s, 2H), 1.69 - 1.49 (m, 2H); MS ESI *m/z* 463.2 (M+H); Anal. HPLC Retention time: 1.1 (Method 2); ROCK2 IC₅₀=464nM.

Example 100 : Preparation of 8-(5-cyclopropyl-1,2-oxazole-3-carbonyl)-2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one




10

¹H NMR (500 MHz, DMSO-d₆) δ 8.02 (br s, 2H), 7.69 - 7.56 (m, 2H), 7.15 (br d, *J*=8.3 Hz, 1H), 6.44 (s, 1H), 4.31 (br d, *J*=13.1 Hz, 1H), 3.96 - 3.75 (m, 7H), 3.47 - 3.09 (m, 2H), 2.30 - 2.05 (m, 4H), 1.67 - 1.53 (m, 2H); MS ESI *m/z* 462.2 (M+H); Anal. HPLC Retention time: 1.42 (Method 1); ROCK2 IC₅₀=6nM.


15

Example 101 : Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-(5-methoxypyrazine-2-carbonyl)-2,8-diazaspiro[4.5]decan-1-one



¹H NMR (500 MHz, DMSO-d₆) δ 8.47 (s, 1H), 8.31 (s, 1H), 8.02 (br s, 2H), 7.71 - 7.55 (m, 2H), 7.15 (dd, *J*=8.4, 1.6 Hz, 1H), 4.35 (br d, *J*=12.8 Hz, 1H), 3.96 (s, 3H), 3.86 (s, 6H), 3.44 (br s, 2H), 3.38 - 3.36 (m, 1H), 2.20 - 2.06 (m, 2H), 1.83 - 1.72 (m, 2H), 1.66 (br d, *J*=13.5 Hz, 1H), 1.53 (br d, *J*=12.5 Hz, 1H); MS ESI *m/z* 463 (M+H); Anal. HPLC Retention time: 1.22 (Method 1); ROCK2 IC₅₀=86nM.

Example 102 : Preparation of 8-(2,3-dihydro-1-benzofuran-2-carbonyl)-2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one




10

¹H NMR (500 MHz, DMSO-d₆) δ 8.02 (br s, 2H), 7.69 - 7.54 (m, 2H), 7.24 (br t, *J*=7.9 Hz, 1H), 7.16 (br d, *J*=8.2 Hz, 1H), 7.11 (s, 1H), 6.85 (s, 1H), 6.80 (s, 1H), 5.76 - 5.57 (m, 1H), 4.23 (br d, *J*=4.7 Hz, 1H), 4.01 (br d, *J*=13.4 Hz, 1H), 3.93 - 3.79 (m, 5H), 3.48 - 3.30 (m, 1H), 3.07 - 2.91 (m, 1H), 2.17 (br t, *J*=6.8 Hz, 2H), 1.83 (br s, 2H), 1.73 - 1.50 (m, 4H); MS ESI *m/z* 473.2 (M+H); Anal. HPLC Retention time: 1.48 (Method 2); ROCK2 IC₅₀=1966nM.


15

Example 103 : Preparation of 8-(4-fluoro-3-methylbenzoyl)-2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one




¹H NMR (500 MHz, DMSO-d₆) δ 8.02 (br s, 2H), 7.66 - 7.55 (m, 2H), 7.36 (br d, *J*=6.7 Hz, 1H), 7.31 - 7.25 (m, 1H), 7.23 - 7.18 (m, 1H), 7.15 (dd, *J*=8.5, 1.5 Hz, 1H), 4.40 - 4.22 (m, 1H), 3.87 (s, 5H), 3.63 - 3.43 (m, 2H), 3.36 - 3.33 (m, 1H), 2.27 (s, 3H), 2.14 (br s, 2H), 1.73 (br s, 2H), 1.67 - 1.48 (m, 2H); MS ESI *m/z* 463 (M+H); Anal. HPLC Retention time: 1.51 (Method 1); ROCK2 IC₅₀=96nM.

Example 104 : Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-(5-methyl-1,2-oxazole-3-carbonyl)-2,8-diazaspiro[4.5]decan-1-one



¹H NMR (500 MHz, DMSO-d₆) δ 8.02 (br s, 2H), 7.67 - 7.53 (m, 2H), 7.15 (dd, *J*=8.5, 1.6 Hz, 1H), 6.47 (s, 1H), 4.32 (br d, *J*=13.0 Hz, 1H), 3.96 - 3.78 (m, 6H), 3.35 (br t, *J*=11.3 Hz, 1H), 3.22 - 3.09 (m, 1H), 2.46 (s, 3H), 2.16 (q, *J*=6.6 Hz, 2H), 1.74 - 1.55 (m, 4H); MS ESI *m/z* 436 (M+H); Anal. HPLC Retention time: 1.23 (Method 1); ROCK2 IC₅₀=72nM.

Example 105 : Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-[4-(pyrimidin-2-yloxy)benzoyl]-2,8-diazaspiro[4.5]decan-1-one




5

¹H NMR (500 MHz, DMSO-d₆) δ 8.67 (d, *J*=4.7 Hz, 2H), 8.02 (br s, 2H), 7.67 - 7.57 (m, 2H), 7.50 (br d, *J*=8.3 Hz, 2H), 7.35 - 7.23 (m, 4H), 7.16 (br d, *J*=7.6 Hz, 2H), 4.43 - 4.25 (m, 1H), 3.87 (s, 5H), 3.74 - 3.63 (m, 1H), 3.40 (br s, 1H), 3.17 (br s, 1H), 2.16 (br s, 2H), 1.76 (br s, 2H), 1.69 - 1.51 (m, 2H); MS ESI *m/z* 525.1 (M+H); Anal.

10 HPLC Retention time: 1.22 (Method 1); ROCK2 IC₅₀=78nM.


Example 106 : Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-[4-(1H-1,2,4-triazol-1-yl)benzoyl]-2,8-diazaspiro[4.5]decan-1-one



^1H NMR (500 MHz, DMSO- d_6) δ 9.35 (s, 1H), 8.27 (s, 1H), 7.95 (br d, $J=8.5$ Hz, 4H), 7.68 - 7.57 (m, 4H), 7.15 (br d, $J=8.5$ Hz, 1H), 4.34 (br d, $J=6.4$ Hz, 1H), 3.87 (s, 5H), 3.54 - 3.47 (m, 3H), 2.16 (br s, 2H), 1.84 - 1.50 (m, 4H); MS ESI m/z 498.2 (M+H); Anal. HPLC Retention time: 1.14 (Method 1); ROCK2 $\text{IC}_{50}=25\text{nM}$.

5


Example 107 : Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-[1-methyl-5-(trifluoromethyl)-1H-pyrazole-3-carbonyl]-2,8-diazaspiro[4.5]decan-1-one



10 ^1H NMR (500 MHz, DMSO- d_6) δ 8.02 (br s, 2H), 7.61 (br d, $J=5.8$ Hz, 2H), 7.25 - 7.10 (m, 3H), 7.01 (s, 1H), 4.33 (br s, 2H), 4.02 (s, 3H), 3.87 (s, 5H), 3.44 (br d, $J=11.0$ Hz, 1H), 3.20 - 3.10 (m, 1H), 2.17 (br d, $J=7.0$ Hz, 2H), 1.81 - 1.70 (m, 2H), 1.68 - 1.56 (m, 2H); MS ESI m/z 503.2 (M+H); Anal. HPLC Retention time: 1.52 (Method 2); ROCK2 $\text{IC}_{50}=67\text{nM}$.


15

Example 108 : Preparation of 8-(5-ethylpyridine-2-carbonyl)-2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one




¹H NMR (500 MHz, DMSO-d₆) δ 8.46 (s, 1H), 8.02 (br s, 2H), 7.79 (br d, *J*=7.6 Hz, 1H), 7.66 - 7.57 (m, 2H), 7.51 (d, *J*=7.9 Hz, 1H), 7.15 (br d, *J*=8.5 Hz, 1H), 4.36 (br d, *J*=13.3 Hz, 1H), 3.91 - 3.80 (m, 5H), 3.74 (br d, *J*=13.2 Hz, 1H), 3.25 (br t, *J*=12.8 Hz, 1H), 3.18 - 3.08 (m, 1H), 2.68 (q, *J*=7.5 Hz, 2H), 2.23 - 2.06 (m, 2H), 1.81 - 1.71 (m, 2H), 1.66 (br d, *J*=13.8 Hz, 1H), 1.51 (br d, *J*=13.5 Hz, 1H), 1.22 (t, *J*=7.6 Hz, 3H); MS ESI *m/z* 460.2 (M+H); Anal. HPLC Retention time: 1.32 (Method 1); ROCK2 IC₅₀=12nM.

Example 109 : Preparation of 8-(3-chloro-1,2-oxazole-5-carbonyl)-2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one




¹H NMR (500 MHz, DMSO-d₆) δ 8.02 (br s, 2H), 7.67 - 7.59 (m, 2H), 7.35 (s, 1H), 7.20 - 7.09 (m, 1H), 4.26 (br d, *J*=12.8 Hz, 1H), 3.87 (s, 6H), 2.25 - 2.10 (m, 2H), 1.86 - 1.74 (m, 2H), 1.63 (br d, *J*=13.5 Hz, 2H); MS ESI *m/z* 456.2 (M+H); Anal. HPLC Retention time: 1.4 (Method 2); ROCK2 IC₅₀=362nM.

Example 110 : Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-[3-(propan-2-yl)-1H-pyrazole-5-carbonyl]-2,8-diazaspiro[4.5]decan-1-one



¹H NMR (500 MHz, DMSO-d₆) δ 8.26 - 7.88 (m, 2H), 7.69 - 7.53 (m, 2H), 7.25 - 7.02 (m, 1H), 6.32 (br s, 1H), 4.63 (br d, *J*=13.7 Hz, 1H), 4.36 (br d, *J*=12.2 Hz, 1H), 3.87 (s, 5H), 3.42 (br s, 1H), 3.11 - 3.02 (m, 1H), 3.04 - 2.91 (m, 1H), 2.16 (br d, *J*=5.3 Hz, 2H), 1.87 - 1.67 (m, 2H), 1.65 - 1.50 (m, 2H), 1.23 (br d, *J*=6.8 Hz, 6H); MS ESI *m/z* 463.3 (M+H); Anal. HPLC Retention time: 1.31 (Method 1); ROCK2 IC₅₀=34nM.

Example 111 : Preparation of 8-(3-cyclopropyl-1H-pyrazole-5-carbonyl)-2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one




¹H NMR (500 MHz, DMSO-d₆) δ 8.02 (br s, 2H), 7.67 - 7.56 (m, 2H), 7.15 (br d, *J*=8.4 Hz, 1H), 6.22 (br s, 1H), 4.58 (br d, *J*=3.3 Hz, 1H), 4.34 (br s, 1H), 3.87 (s, 5H), 3.17 (br d, *J*=4.4 Hz, 1H), 3.08 - 2.96 (m, 1H), 2.16 (br d, *J*=4.5 Hz, 2H), 1.91 (br s, 1H),

1.71 (br d, $J=11.2$ Hz, 2H), 1.60 (br d, $J=19.9$ Hz, 2H), 0.94 (br s, 2H), 0.71 (br d, $J=3.0$ Hz, 2H); MS ESI m/z 461.1 (M+H);


Anal. HPLC Retention time: 1.21 (Method 2); ROCK2 $IC_{50}=17$ nM.

5 **Example 112 : Preparation of 8-[4-(4-chlorophenoxy)benzoyl]-2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one**




^1H NMR (500 MHz, DMSO- d_6) δ 8.01 (br s, 2H), 7.65 - 7.57 (m, 2H), 7.46 (br t, $J=9.2$ Hz, 5H), 7.12 (br d, $J=8.9$ Hz, 3H), 7.07 (br d, $J=8.5$ Hz, 2H), 4.40 - 4.20 (m, 1H),
 10 3.86 (s, 6H), 3.34 (br s, 2H), 2.15 (br s, 2H), 1.74 (br s, 2H), 1.59 (br d, $J=3.4$ Hz, 2H);
 MS ESI m/z 557.2 (M+H); Anal. HPLC Retention time: 1.93 (Method 2); ROCK2
 $IC_{50}=39$ nM.

15 **Example 113 : Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-[3-(4-methoxyphenyl)-1,2-oxazole-5-carbonyl]-2,8-diazaspiro[4.5]decan-1-one**




¹H NMR (500 MHz, DMSO-d₆) δ 8.09 (br s, 1H), 7.95 (br s, 1H), 7.88 (br d, *J*=8.5 Hz, 2H), 7.63 - 7.58 (m, 2H), 7.16 (s, 2H), 7.11 (br d, *J*=8.5 Hz, 2H), 4.36 (br d, *J*=12.8 Hz, 1H), 3.99 (br d, *J*=13.4 Hz, 1H), 3.87 (s, 5H), 3.84 (s, 3H), 3.50 - 3.40 (m, 2H), 2.22 - 2.14 (m, 2H), 1.78 (br d, *J*=12.5 Hz, 2H), 1.73 - 1.60 (m, 2H); MS ESI *m/z* 528.1 (M+H); Anal. HPLC Retention time: 1.68 (Method 2), ROCK2 IC₅₀=17nM.

Example 114 : Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-[4-oxo-4H,5H,6H,7H-pyrazolo[1,5-a]pyrazine-2-carbonyl]-2,8-diazaspiro[4.5]decan-1-one




¹H NMR (500 MHz, DMSO-d₆) δ 8.33 (br s, 1H), 8.14 - 7.90 (m, 2H), 7.61 (dd, *J*=4.7, 3.2 Hz, 2H), 7.16 (dd, *J*=8.4, 1.7 Hz, 1H), 6.94 (s, 1H), 4.36 (br t, *J*=6.0 Hz, 4H), 3.87 (s, 5H), 3.65 (br s, 2H), 3.42 (br d, *J*=10.7 Hz, 2H), 2.25 - 2.11 (m, 2H), 1.75 (br d, *J*=13.1 Hz, 2H), 1.68 - 1.53 (m, 2H); MS ESI *m/z* 489.9 (M+H); Anal. HPLC Retention time: 1.17 (Method 1); ROCK2 IC₅₀=6nM.

Example 115 : Preparation of 8-(5-chloropyrazine-2-carbonyl)-2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one




¹H NMR (500 MHz, DMSO-d₆) δ 8.84 (s, 1H), 8.74 (s, 1H), 8.02 (br s, 2H), 7.67 - 7.55 (m, 2H), 7.16 (br d, *J*=8.2 Hz, 1H), 4.35 (br d, *J*=13.5 Hz, 1H), 3.87 (s, 5H), 3.73 (br d, *J*=13.6 Hz, 1H), 3.34 - 3.12 (m, 2H), 2.26 - 2.09 (m, 2H), 1.84 - 1.74 (m, 2H), 1.68 (br d, *J*=13.7 Hz, 1H), 1.53 (br d, *J*=13.0 Hz, 1H); MS ESI *m/z* 467.2 (M+H); Anal. HPLC Retention time: 1.28 (Method 1); ROCK2 IC₅₀=200nM.

Example 116 : Preparation of 8-(5-bromopyrimidine-2-carbonyl)-2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one




¹H NMR (500 MHz, DMSO-d₆) δ 9.11 (s, 2H), 8.02 (br s, 2H), 7.67 - 7.54 (m, 2H), 7.15 (dd, *J*=8.4, 1.6 Hz, 1H), 4.33 (br d, *J*=12.9 Hz, 1H), 3.99 - 3.79 (m, 5H), 3.59 - 3.46 (m, 3H), 2.24 - 2.10 (m, 2H), 1.86 - 1.62 (m, 5H), 1.49 (br d, *J*=13.0 Hz, 1H); MS ESI *m/z* 511 (M+H); Anal. HPLC Retention time: 1.21 (Method 1); ROCK2 IC₅₀=88nM.

Example 117 : Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-[4-(2-oxo-1,2-dihydropyridin-1-yl)benzoyl]-2,8-diazaspiro[4.5]decan-1-one




- 5 ¹H NMR (500 MHz, DMSO-d₆) δ 8.02 (br d, *J*=5.5 Hz, 2H), 7.75 - 7.67 (m, 1H), 7.62 (s, 2H), 7.53 (br d, *J*=17.7 Hz, 5H), 7.16 (dd, *J*=8.5, 1.5 Hz, 1H), 6.51 (br d, *J*=9.5 Hz, 1H), 6.35 (br t, *J*=6.4 Hz, 1H), 4.35 (br d, *J*=4.6 Hz, 1H), 3.87 (s, 5H), 3.65 (br s, 1H), 3.42 (br d, *J*=6.1 Hz, 2H), 2.16 (br s, 2H), 1.77 (br s, 2H), 1.67 (br s, 1H), 1.57 (br s, 1H); MS ESI *m/z* 524.3 (M+H); Anal. HPLC Retention time: 1.13 (Method 2); ROCK2
- 10 IC₅₀=7nM.

Example 118 : Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-[4-(1H-1,2,3,4-tetrazol-1-yl)benzoyl]-2,8-diazaspiro[4.5]decan-1-one



¹H NMR (600 MHz, DMSO-d₆) δ 10.14 (s, 1H), 8.10 (br s, 1H), 8.01 (br d, *J*=8.1 Hz, 2H), 7.94 (br s, 1H), 7.70 (br d, *J*=8.3 Hz, 2H), 7.64 - 7.56 (m, 2H), 7.15 (br d, *J*=8.5 Hz, 1H), 4.66 (br d, *J*=5.0 Hz, 1H), 4.35 (br d, *J*=7.3 Hz, 1H), 3.94 - 3.86 (m, 5H), 3.70 - 3.57 (m, 1H), 3.32 (br dd, *J*=10.7, 5.7 Hz, 1H), 2.21 - 2.10 (m, 2H), 1.86 - 1.72 (m, 2H),
 5 1.68 (br d, *J*=10.1 Hz, 1H), 1.58 - 1.47 (m, 1H); MS ESI *m/z* 499 (M+H); Anal. HPLC Retention time: 1.16 (Method 1); ROCK2 IC₅₀=57nM.


Example 119 : Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-(2-methyl-1,3-benzoxazole-5-carbonyl)-2,8-diazaspiro[4.5]decan-1-one



10


¹H NMR (500 MHz, DMSO-d₆) δ 8.02 (s, 2H), 7.73 (d, *J*=8.3 Hz, 1H), 7.68 (s, 1H), 7.63 - 7.57 (m, 2H), 7.39 (d, *J*=8.2 Hz, 1H), 7.14 (dd, *J*=8.5, 1.6 Hz, 1H), 4.03 (dd, *J*=11.1, 3.9 Hz, 1H), 3.93 - 3.81 (m, 5H), 3.62 (br s, 1H), 3.47 (br s, 1H), 3.33 (br s, 1H), 2.63 (s, 3H), 2.14 (br s, 2H), 1.81 - 1.61 (m, 3H), 1.51 (br d, *J*=5.6 Hz, 1H); MS ESI *m/z*
 15 486.3 (M+H); Anal. HPLC Retention time: 1.27 (Method 2); ROCK2 IC₅₀=70nM.

Example 120 : Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-[4-(4-methylpiperazin-1-yl)benzoyl]-2,8-diazaspiro[4.5]decan-1-one



¹H NMR (500 MHz, DMSO-d₆) δ 8.18 - 7.91 (m, 2H), 7.61 (dd, *J*=5.0, 3.3 Hz, 2H), 7.29 (br d, *J*=8.6 Hz, 2H), 7.15 (dd, *J*=8.5, 1.6 Hz, 1H), 6.96 (br d, *J*=8.8 Hz, 2H), 4.03 (br dd, *J*=11.0, 4.0 Hz, 1H), 3.94 - 3.77 (m, 5H), 3.63 (br d, *J*=5.6 Hz, 1H), 3.40 (br s, 1H), 3.21 (br s, 2H), 3.17 (br d, *J*=4.4 Hz, 1H), 2.51 (br s, 8H), 2.46 (br d, *J*=4.4 Hz, 4H), 2.22 (s, 3H), 2.14 (br t, *J*=6.6 Hz, 2H), 1.77 - 1.66 (m, 2H), 1.57 (br d, *J*=10.7 Hz, 2H); MS ESI *m/z* 529.2 (M+H); Anal. HPLC Retention time: 0.98 (Method 2); ROCK2 IC₅₀=0.3nM.

10 **Example 121: Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-(2-methylpyrimidine-5-carbonyl)-2,8-diazaspiro[4.5]decan-1-one**




¹H NMR (500 MHz, DMSO-d₆) δ 8.78 (s, 2H), 8.02 (br s, 2H), 7.65 - 7.57 (m, 2H), 7.15 (br d, *J*=8.5 Hz, 1H), 4.32 (br d, *J*=11.0 Hz, 1H), 4.07 - 3.99 (m, 1H), 3.91 - 3.82 (m, 5H), 3.63 (br s, 1H), 3.39 - 3.27 (m, 1H), 2.67 (s, 3H), 2.14 (br d, *J*=6.6 Hz, 2H), 1.77 (br d, *J*=7.7 Hz, 2H), 1.67 (br d, *J*=10.9 Hz, 1H), 1.55 (br d, *J*=11.9 Hz, 1H); MS

ESI m/z 447.2 (M+H); Anal. HPLC Retention time: 0.99 (Method 1); ROCK2


IC₅₀=585nM.

5 Example 122: Preparation of 8-(2-chloro-4-methoxybenzoyl)-2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one




¹H NMR (500 MHz, DMSO-d₆) δ 8.16 - 7.88 (m, 2H), 7.67 - 7.58 (m, 2H), 7.36 - 7.24 (m, 1H), 7.12 (br d, *J*=9.3 Hz, 2H), 7.05 - 6.95 (m, 1H), 4.37 (br d, *J*=4.5 Hz, 1H), 4.03 (dd, *J*=11.1, 4.2 Hz, 1H), 3.86 (s, 4H), 3.81 (s, 3H), 3.17 - 3.07 (m, 2H), 2.12 (br dd, *J*=13.1, 6.9 Hz, 2H), 1.86 - 1.71 (m, 2H), 1.65 (br d, *J*=13.0 Hz, 1H), 1.49 (br d, *J*=13.4 Hz, 1H); MS ESI m/z 495 (M+H); Anal. HPLC Retention time: 1.47 (Method 1); ROCK2
10 IC₅₀=11nM.

15 Example 123: Preparation of 8-(5-chloropyridine-2-carbonyl)-2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one




¹H NMR (500 MHz, DMSO-d₆) δ 8.67 (d, *J*=2.0 Hz, 1H), 8.08 (dd, *J*=8.4, 2.2 Hz, 1H), 8.02 (br s, 2H), 7.69 - 7.56 (m, 3H), 7.15 (dd, *J*=8.4, 1.4 Hz, 1H), 4.35 (br d, *J*=13.1 Hz, 1H), 3.94 - 3.80 (m, 5H), 3.67 (br d, *J*=13.3 Hz, 1H), 3.34 - 3.08 (m, 2H), 2.24 - 2.08 (m, 2H), 1.85 - 1.72 (m, 2H), 1.66 (br d, *J*=12.9 Hz, 1H), 1.52 (br d, *J*=13.0 Hz, 1H); MS ESI *m/z* 466 (M+H); Anal. HPLC Retention time: 1.33 Method 2; ROCK2 IC₅₀=65nM.

Example 124: Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-[4-(1H-pyrazol-1-yl)benzoyl]-2,8-diazaspiro[4.5]decan-1-one




¹H NMR (500 MHz, DMSO-d₆) δ 8.58 (br s, 1H), 8.11 (br d, *J*=5.0 Hz, 1H), 8.01 (br d, *J*=9.0 Hz, 1H), 7.94 (br d, *J*=8.2 Hz, 2H), 7.79 (s, 1H), 7.62 (br s, 1H), 7.55 (br d, *J*=8.2 Hz, 1H), 7.16 (br d, *J*=8.4 Hz, 1H), 6.59 (br s, 1H), 4.57 (br d, *J*=8.6 Hz, 1H), 4.47 - 4.38 (m, 1H), 4.03 (br s, 1H), 3.87 (s, 3H), 3.73 - 3.59 (m, 1H), 3.39 - 3.31 (m, 2H), 2.15 (br s, 2H), 1.76 (br s, 2H), 1.66 - 1.45 (m, 2H); MS ESI *m/z* 496.9 (M+H); Anal. HPLC Retention time: 1.48 (Method 1); ROCK2 IC₅₀=62nM.

Example 125: Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-[5-oxo-3-(pyridin-2-yl)-4,5-dihydro-1,2,4-triazine-6-carbonyl]-2,8-diazaspiro[4.5]decan-1-one



¹H NMR (500 MHz, DMSO-d₆) δ 8.82 (br d, *J*=4.3 Hz, 1H), 8.33 (br d, *J*=7.9 Hz, 1H), 8.12 (br t, *J*=7.8 Hz, 1H), 8.09 - 7.89 (m, 2H), 7.78 - 7.70 (m, 1H), 7.63 - 7.55 (m, 2H), 7.15 (br d, *J*=8.2 Hz, 1H), 4.31 (br d, *J*=13.4 Hz, 1H), 3.92 - 3.80 (m, 5H), 3.66 (br d, *J*=14.6 Hz, 1H), 3.35 - 3.22 (m, 1H), 3.13 (br s, 1H), 2.23 - 2.11 (m, 2H), 1.74 (br d, *J*=7.0 Hz, 2H), 1.66 (br d, *J*=11.9 Hz, 1H), 1.54 (br d, *J*=12.5 Hz, 1H); MS ESI *m/z* 527.1 (M+H); Anal. HPLC Retention time: 0.86 (Method 1); ROCK2 IC₅₀=26nM.


Example 126: Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-[4-(2-oxo-1,3-diazinan-1-yl)benzoyl]-2,8-diazaspiro[4.5]decan-1-one



¹H NMR (500 MHz, DMSO-d₆) δ 8.02 (br s, 2H), 7.68 - 7.55 (m, 2H), 7.42 - 7.28 (m, 5H), 7.24 (br d, *J*=8.2 Hz, 1H), 7.14 (br d, *J*=7.9 Hz, 1H), 6.66 (br s, 1H), 3.86 (s, 5H), 3.65 (br t, *J*=5.3 Hz, 1H), 3.55 (br s, 1H), 3.49 (br s, 2H), 3.23 (br s, 4H), 2.14 (br s,


2H), 1.94 (br d, $J=5.3$ Hz, 2H), 1.73 (br s, 2H), 1.65 - 1.48 (m, 2H), 1.12 (br s, 2H); MS ESI m/z 529 (M+H); Anal. HPLC Retention time: 1.22 (Method 1); ROCK2 $IC_{50}=7.7$ nM.

Example 127: Preparation of 8-[3-fluoro-4-(morpholin-4-yl)benzoyl]-2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one



^1H NMR (500 MHz, DMSO- d_6) δ 8.14 - 7.85 (m, 2H), 7.60 (br d, $J=6.7$ Hz, 2H), 7.26 - 7.18 (m, 2H), 7.15 (br d, $J=9.2$ Hz, 1H), 7.08 (br t, $J=8.5$ Hz, 1H), 3.90 (s, 5H), 3.87 (s, 4H), 3.75 (br s, 4H), 3.18 (br s, 2H), 3.07 (br s, 4H), 2.14 (br s, 2H), 1.82 - 1.67 (m, 2H), 1.59 (br s, 2H); MS ESI m/z 534.3 (M+H); Anal. HPLC Retention time: 1.47 (Method 1); ROCK2 $IC_{50}=11.2$ nM.

Example 128: Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-[4-(2-oxopyrrolidin-1-yl)benzoyl]-2,8-diazaspiro[4.5]decan-1-one




¹H NMR (500 MHz, DMSO-d₆) δ 8.17 - 7.87 (m, 2H), 7.75 (br d, *J*=8.4 Hz, 2H), 7.65 - 7.58 (m, 2H), 7.44 (br d, *J*=8.3 Hz, 2H), 7.15 (br d, *J*=8.7 Hz, 1H), 4.39 - 4.24 (m, 1H), 3.95 - 3.77 (m, 7H), 3.75 - 3.61 (m, 1H), 3.31 (br s, 1H), 3.18 (br dd, *J*=19.3, 4.2 Hz, 1H), 2.15 (br s, 2H), 2.08 (br t, *J*=7.4 Hz, 2H), 1.81 - 1.72 (m, 2H), 1.67 - 1.48 (m, 2H);

5 MS ESI *m/z* 514.2 (M+H); Anal. HPLC Retention time: 1.27 (Method 1); ROCK2 IC₅₀=13nM.

Example 129: Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-[7-oxo-4H,7H-[1,2,3,4]tetrazolo[1,5-a]pyrimidine-5-carbonyl]-2,8-diazaspiro[4.5]decan-1-one

10 **one**




¹H NMR (500 MHz, DMSO-d₆) δ 8.02 (br s, 2H), 7.60 (s, 2H), 7.22 (s, 1H), 7.14 (br s, 1H), 7.12 (s, 1H), 7.02 (s, 1H), 5.63 (s, 1H), 4.30 (br d, *J*=13.3 Hz, 1H), 3.93 - 3.78 (m, 4H), 3.66 (br d, *J*=12.3 Hz, 1H), 3.61 (br s, 1H), 3.37 (br d, *J*=5.6 Hz, 1H), 3.25 -

15 3.17 (m, 1H), 3.11 (br t, *J*=11.1 Hz, 1H), 2.21 - 2.08 (m, 2H), 1.82 - 1.68 (m, 2H), 1.64 (br d, *J*=13.3 Hz, 1H), 1.51 (br d, *J*=12.5 Hz, 1H); MS ESI *m/z* 490.18, 490.18 (M+H); Anal. HPLC Retention time: (Method 2); ROCK2 IC₅₀=292nM.


Example 130: Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-[4-(piperazin-1-yl)benzoyl]-2,8-diazaspiro[4.5]decan-1-one

20 **one**



¹H NMR (500 MHz, DMSO-d₆) δ 8.01 (br s, 2H), 7.67 - 7.56 (m, 2H), 7.34 (br d, *J*=8.5 Hz, 2H), 7.15 (br d, *J*=8.5 Hz, 1H), 7.03 (br d, *J*=8.5 Hz, 2H), 4.21 - 4.11 (m, 1H), 3.94 - 3.82 (m, 5H), 3.73 - 3.64 (m, 1H), 3.43 (br s, 2H), 3.29 - 3.12 (m, 8H), 2.14 (br t, *J*=6.4 Hz, 2H), 1.79 - 1.66 (m, 2H), 1.57 (br d, *J*=10.1 Hz, 2H); MS ESI *m/z* 515.2 (M+H); Anal. HPLC Retention time: 1.08 (Method 1); ROCK2 IC₅₀=0.4nM.

Example 131: Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-{5-methyl-2-oxo-1H,2H-pyrazolo[1,5-a]pyrimidine-6-carbonyl}-2,8-diazaspiro[4.5]decan-1-one




10

¹H NMR (500 MHz, DMSO-d₆) δ 8.25 (br s, 1H), 8.02 (br s, 2H), 7.70 - 7.55 (m, 2H), 7.14 (br d, *J*=8.5 Hz, 1H), 5.95 (s, 1H), 4.38 (br d, *J*=11.3 Hz, 1H), 3.87 (s, 5H), 3.62 (br s, 1H), 3.35 - 3.14 (m, 2H), 2.61 (br s, 3H), 2.15 (br d, *J*=5.8 Hz, 2H), 1.80 (br s, 1H), 1.70 (br s, 2H), 1.54 (br s, 1H); MS ESI *m/z* 502 (M+H); Anal. HPLC Retention time: 1.06 (Method 1); ROCK2 IC₅₀=2087nM.

15


Example 132: Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-[7-methyl-4-oxo-3H,4H-pyrrolo[2,1-f][1,2,4]triazine-5-carbonyl]-2,8-diazaspiro[4.5]decan-1-one



5 ^1H NMR (500 MHz, DMSO- d_6) δ 11.91 - 11.25 (m, 1H), 8.21 - 7.92 (m, 2H), 7.87 (br s, 1H), 7.67 - 7.55 (m, 2H), 7.14 (br d, $J=8.2$ Hz, 1H), 6.42 (s, 1H), 4.38 (br d, $J=13.1$ Hz, 1H), 3.89 (d, $J=18.6$ Hz, 6H), 3.61 (br d, $J=14.0$ Hz, 1H), 3.18 - 3.05 (m, 2H), 2.40 (s, 3H), 2.21 - 2.05 (m, 2H), 1.83 - 1.57 (m, 3H), 1.42 (br d, $J=13.1$ Hz, 1H); MS ESI m/z (M+H); Anal. HPLC Retention time: 1.25 (Method 1); ROCK2 $\text{IC}_{50}=3\text{nM}$.

10


Example 133: Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-[5-methyl-4-oxo-3H,4H-pyrrolo[2,1-f][1,2,4]triazine-6-carbonyl]-2,8-diazaspiro[4.5]decan-1-one



15 ^1H NMR (500 MHz, DMSO- d_6) δ 8.01 (br s, 2H), 7.78 (s, 1H), 7.66 (s, 1H), 7.63 - 7.56 (m, 2H), 7.14 (br d, $J=6.8$ Hz, 1H), 3.86 (s, 5H), 3.63 - 3.42 (m, 4H), 2.40 (s, 3H),


2.14 (br s, 2H), 1.71 (br s, 2H), 1.58 (br s, 2H); MS ESI m/z 502.2 (M+H); Anal. HPLC Retention time: 1.18 (Method 1); ROCK2 IC₅₀=88nM.

Example 134: Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-[4-[5-(propan-2-yl)-1,2,4-oxadiazol-3-yl]benzoyl]-2,8-diazaspiro[4.5]decan-1-one



¹H NMR (500 MHz, DMSO-d₆) δ 8.12 (s, 1H), 8.08 (br d, *J*=7.0 Hz, 2H), 8.03 (br s, 2H), 7.62 - 7.56 (m, 4H), 7.15 (br d, *J*=8.5 Hz, 1H), 4.41 - 4.30 (m, 1H), 3.94 - 3.78 (m, 5H), 3.51 (br s, 1H), 3.42 - 3.32 (m, 1H), 3.29 (br d, *J*=11.0 Hz, 1H), 3.18 (br s, 1H), 2.15 (br s, 2H), 1.84 - 1.65 (m, 3H), 1.53 (br d, *J*=7.6 Hz, 1H), 1.39 (br d, *J*=6.7 Hz, 6H); MS ESI m/z 541.1 (M+H); Anal. HPLC Retention time: 1.76 (Method 1); ROCK2 IC₅₀=75nM.


Example 135: Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-[4-(pyrimidin-5-yl)benzoyl]-2,8-diazaspiro[4.5]decan-1-one



¹H NMR (500 MHz, DMSO-d₆) δ 9.26 - 9.14 (m, 3H), 8.01 (br s, 2H), 7.91 (br d, *J*=8.0 Hz, 2H), 7.66 - 7.52 (m, 4H), 7.14 (br d, *J*=8.3 Hz, 1H), 4.43 - 4.27 (m, 1H), 3.86 (s, 4H), 3.58 (br s, 2H), 3.31 (br d, *J*=10.2 Hz, 1H), 3.17 (br d, *J*=5.0 Hz, 1H), 2.16 (br s, 2H), 1.88 - 1.61 (m, 3H), 1.53 (br d, *J*=10.1 Hz, 1H); MS ESI *m/z* 509 (M+H); Anal. HPLC Retention time: 1.29 (Method 1); ROCK2 IC₅₀=10nM.

Example 136: Preparation of 8-[2-fluoro-4-(2-oxo-1,2-dihydropyrazin-1-yl)benzoyl]-2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one

10




¹H NMR (500 MHz, DMSO-d₆) δ 8.19 - 7.87 (m, 2H), 7.70 (br d, *J*=6.1 Hz, 1H), 7.64 - 7.48 (m, 5H), 7.37 (br d, *J*=8.2 Hz, 1H), 7.15 (br d, *J*=7.9 Hz, 1H), 6.52 (br d, *J*=9.2 Hz, 1H), 6.59 - 6.48 (m, 1H), 6.37 (br t, *J*=6.4 Hz, 1H), 4.39 (br d, *J*=12.5 Hz, 1H), 3.87 (s, 5H), 3.74 - 3.65 (m, 1H), 3.30 (br d, *J*=9.2 Hz, 1H), 3.23 - 3.10 (m, 1H), 2.27 -

15


2.08 (m, 2H), 1.89 - 1.65 (m, 3H), 1.56 (br d, $J=13.1$ Hz, 1H); MS ESI m/z 543.2 (M+H);
 Anal. HPLC Retention time: 1.23 (Method 1); ROCK2 $IC_{50}=4.2$ nM.

5 Example 137: Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-[4-(2-oxo-
1,3-oxazinan-3-yl)benzoyl]-2,8-diazaspiro[4.5]decan-1-one




^1H NMR (500 MHz, DMSO- d_6) δ 8.01 (br s, 2H), 7.65 - 7.57 (m, 2H), 7.43 (br d, $J=3.4$ Hz, 4H), 7.14 (br d, $J=8.5$ Hz, 1H), 4.35 (br t, $J=5.2$ Hz, 2H), 3.86 (s, 5H), 3.70 (br
 10 t, $J=6.0$ Hz, 2H), 3.57 - 3.49 (m, 2H), 3.36 - 3.09 (m, 2H), 2.20 - 2.06 (m, 4H), 1.74 (br s, 2H), 1.67 - 1.50 (m, 2H); MS ESI m/z 530.4 (M+H); Anal. HPLC Retention time: 1.2 (Method 2); ROCK2 $IC_{50}=7.1$ nM.

15 Example 138: Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-[2-
(trifluoromethyl)quinoxaline-6-carbonyl]-2,8-diazaspiro[4.5]decan-1-one




MS ESI m/z 551.2 (M+H); Anal. HPLC Retention time: 1.58 Method 1; ROCK2
 IC_{50} =63nM.

5 Example 139: Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-[4-(2-oxoimidazolidin-1-yl)benzoyl]-2,8-diazaspiro[4.5]decan-1-one



^1H NMR (500 MHz, DMSO- d_6) δ 8.02 (br s, 2H), 7.68 - 7.56 (m, 4H), 7.38 (br d, J =8.5 Hz, 2H), 7.14 (br d, J =8.2 Hz, 1H), 7.04 (s, 1H), 4.01 (br s, 1H), 3.92 - 3.81 (m, 7H), 3.76 - 3.68 (m, 1H), 3.47 (br s, 2H), 3.42 (br t, J =7.9 Hz, 1H), 3.17 (br d, J =4.6 Hz, 1H), 2.14 (br s, 2H), 1.73 (br s, 2H), 1.58 (br d, J =4.6 Hz, 2H); MS ESI m/z 515.3
 10 (M+H); Anal. HPLC Retention time: 1.11 (Method 2); ROCK2 IC_{50} =3.7nM.


Example 140: Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-[1-(pyrimidin-2-yl)piperidine-4-carbonyl]-2,8-diazaspiro[4.5]decan-1-one



¹H NMR (500 MHz, DMSO-d₆) δ 8.34 (br d, *J*=4.5 Hz, 2H), 8.21 - 7.88 (m, 2H),
 5 7.70 - 7.53 (m, 2H), 7.15 (br d, *J*=8.6 Hz, 1H), 6.59 (br t, *J*=4.5 Hz, 1H), 4.66 (br d,
J=10.7 Hz, 2H), 4.25 (br d, *J*=12.5 Hz, 1H), 4.01 (br d, *J*=13.0 Hz, 1H), 3.87 (s, 5H), 3.28
 (br t, *J*=11.4 Hz, 1H), 3.06 - 2.92 (m, 4H), 2.87 (br t, *J*=10.4 Hz, 1H), 2.14 (br t, *J*=6.5
 Hz, 2H), 1.78 - 1.39 (m, 9H); MS ESI *m/z* 516.2 (M+H); Anal. HPLC Retention time:
 1.41 (Method 1); ROCK2 IC₅₀=1240nM.

10


Example 141: Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-[4-(2-oxo-1,3-oxazolidin-3-yl)benzoyl]-2,8-diazaspiro[4.5]decan-1-one



¹H NMR (500 MHz, DMSO-d₆) δ 8.01 (br s, 2H), 7.66 - 7.54 (m, 4H), 7.46 (br d,
 15 *J*=8.5 Hz, 2H), 7.14 (br d, *J*=7.3 Hz, 1H), 4.46 (br t, *J*=7.9 Hz, 2H), 4.09 (br t, *J*=7.8 Hz,


2H), 3.86 (s, 5H), 3.56 - 3.45 (m, 2H), 3.32 - 3.17 (m, 2H), 2.14 (br s, 2H), 1.73 (br s, 2H), 1.65 - 1.49 (m, 2H); MS ESI m/z 515.9 (M+H); Anal. HPLC Retention time: 1.28 (Method 1); ROCK2 IC₅₀=3nM.

5 **Example 142: Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-[4-(2-oxopiperidin-1-yl)benzoyl]-2,8-diazaspiro[4.5]decan-1-one**




¹H NMR (500 MHz, DMSO-d₆) δ 8.19 - 7.86 (m, 2H), 7.67 - 7.57 (m, 2H), 7.46 - 7.40 (m, 2H), 7.39 - 7.33 (m, 2H), 7.15 (br d, J=8.2 Hz, 1H), 4.42 - 4.25 (m, 1H), 3.87 (s, 5H), 3.63 (br t, J=5.5 Hz, 2H), 3.51 - 3.47 (m, 1H), 3.32 - 3.11 (m, 2H), 2.41 (br t, J=6.3 Hz, 2H), 2.15 (br s, 2H), 1.93 - 1.80 (m, 4H), 1.74 (br s, 2H), 1.68 - 1.50 (m, 2H); MS ESI m/z 528.2 (M+H); Anal. HPLC Retention time: 1.26 (Method 1); ROCK2 IC₅₀=8.8nM.

15 **Example 143: Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-[4-(2H-1,2,3,4-tetrazol-5-yl)benzoyl]-2,8-diazaspiro[4.5]decan-1-one**



¹H NMR (500 MHz, DMSO-d₆) δ 8.07 (br d, *J*=7.9 Hz, 2H), 8.01 (br s, 2H), 7.64 - 7.56 (m, 2H), 7.48 (d, *J*=7.9 Hz, 2H), 7.18 - 7.09 (m, 1H), 4.41 - 4.27 (m, 1H), 3.87 (s, 5H), 3.75 - 3.61 (m, 1H), 3.33 - 3.10 (m, 2H), 2.15 (br s, 2H), 1.76 (br d, *J*=8.2 Hz, 2H), 1.68 - 1.50 (m, 2H); MS ESI *m/z* 499.3 (M+H); Anal. HPLC Retention time: 1.19 (Method 2); ROCK2 IC₅₀=1018nM.

Example 144: Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-(1-oxo-2,3-dihydro-1H-isoindole-5-carbonyl)-2,8-diazaspiro[4.5]decan-1-one




10

¹H NMR (500 MHz, DMSO-d₆) δ 8.67 (s, 1H), 8.02 (br s, 2H), 7.75 (d, *J*=7.6 Hz, 1H), 7.61 (br d, *J*=5.5 Hz, 3H), 7.49 (br d, *J*=7.6 Hz, 1H), 7.15 (br d, *J*=9.2 Hz, 1H), 4.43 (s, 2H), 4.38 - 4.27 (m, 1H), 3.90 (s, 5H), 3.65 - 3.52 (m, 1H), 3.40 (br s, 1H), 3.28 - 3.21 (m, 1H), 2.25 - 2.07 (m, 2H), 1.83 - 1.60 (m, 3H), 1.53 (br s, 1H); MS ESI *m/z* 486.02, 486.02 (M+H); ROCK2 IC₅₀=94nM.


15

Example 145: Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-[6-(1H-1,2,4-triazol-1-yl)pyridine-3-carbonyl]-2,8-diazaspiro[4.5]decan-1-one



5 ¹H NMR (500 MHz, DMSO-d₆) δ 9.42 (s, 1H), 8.61 (s, 1H), 8.34 (s, 1H), 8.14 (br d, *J*=8.5 Hz, 1H), 8.09 - 7.91 (m, 3H), 7.63 - 7.55 (m, 2H), 7.15 (br d, *J*=8.2 Hz, 1H), 4.34 (br d, *J*=8.5 Hz, 1H), 3.87 (s, 5H), 3.72 - 3.62 (m, 1H), 3.34 (br d, *J*=4.6 Hz, 1H), 3.23 (br d, *J*=12.2 Hz, 1H), 2.15 (br d, *J*=4.6 Hz, 2H), 1.79 (br s, 2H), 1.72 - 1.63 (m, 1H), 1.56 (br s, 1H); MS ESI *m/z* 499.1 (M+H); Anal. HPLC Retention time: 1.2
10 (Method 2); ROCK2 IC₅₀=54nM.


Example 146: Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-[4-(2-oxo-1,2-dihydropyrazin-1-yl)benzoyl]-2,8-diazaspiro[4.5]decan-1-one



^1H NMR (500 MHz, DMSO- d_6) δ 8.15 (s, 1H), 8.04 (br d, $J=12.5$ Hz, 2H), 7.71 (br d, $J=4.3$ Hz, 1H), 7.63 - 7.56 (m, 6H), 7.42 (br d, $J=4.3$ Hz, 1H), 7.16 (br d, $J=8.2$ Hz, 1H), 4.42 - 4.30 (m, 1H), 4.13 (br d, $J=4.9$ Hz, 1H), 3.87 (s, 5H), 3.39 (br d, $J=4.0$ Hz, 1H), 3.33 - 3.26 (m, 1H), 2.15 (br s, 2H), 1.77 (br s, 2H), 1.72 - 1.63 (m, 1H), 1.57 (br s, 1H); MS ESI m/z 525.3 (M+H);

Anal. HPLC Retention time: 1.16 (Method 2); ROCK2 $\text{IC}_{50}=12\text{nM}$.

Example 147: Preparation of 1-(4-{2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-1-oxo-2,8-diazaspiro[4.5]decane-8-carbonyl}phenyl)imidazolidine-2,4-dione




10

^1H NMR (500 MHz, DMSO- d_6) δ 8.18 - 7.90 (m, 2H), 7.68 (br d, $J=8.5$ Hz, 2H), 7.61 (br d, $J=5.5$ Hz, 2H), 7.44 (br d, $J=8.5$ Hz, 2H), 7.15 (br d, $J=7.3$ Hz, 1H), 4.48 (s, 2H), 4.37 - 4.19 (m, 1H), 3.87 (s, 5H), 3.74 - 3.60 (m, 1H), 3.44 - 3.34 (m, 1H), 3.29 - 3.21 (m, 1H), 2.15 (br s, 2H), 1.74 (br s, 2H), 1.57 (br d, $J=8.2$ Hz, 2H); MS ESI m/z 529.4 (M+H); Anal. HPLC Retention time: 1.18 (Method 1); ROCK2 $\text{IC}_{50}=12\text{nM}$.


15

Example 148: Preparation of 4-{2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-1-oxo-2,8-diazaspiro[4.5]decane-8-carbonyl}cyclohexane-1-carboxamide



¹H NMR (500 MHz, DMSO-d₆) δ 8.00 (br s, 2H), 7.59 (d, *J*=8.4 Hz, 1H), 7.55 (s, 1H), 7.34 (br s, 1H), 7.12 (br d, *J*=8.3 Hz, 1H), 6.69 (br s, 1H), 4.24 (br d, *J*=13.7 Hz, 1H), 3.91 (br d, *J*=12.4 Hz, 1H), 3.80 (br s, 6H), 3.29 - 3.15 (m, 1H), 2.81 (br t, *J*=10.3 Hz, 1H), 2.59 (br s, 1H), 2.11 (br t, *J*=6.6 Hz, 4H), 1.74 (br s, 2H), 1.67 (br s, 3H), 1.60 - 1.50 (m, 3H), 1.44 - 1.34 (m, 4H); MS ESI *m/z* 480.4 (M+H); Anal. HPLC Retention time: 0.94 (Method 2); ROCK2 IC₅₀=2165nM.


Example 149: Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-(quinoline-8-carbonyl)-2,8-diazaspiro[4.5]decan-1-one



¹H NMR (500 MHz, DMSO-d₆) δ 8.96 (br dd, *J*=10.7, 3.1 Hz, 1H), 8.50 - 8.39 (m, 1H), 8.11 - 7.95 (m, 3H), 7.80 - 7.63 (m, 2H), 7.63 - 7.56 (m, 3H), 7.13 (br t, *J*=6.4 Hz, 1H), 4.63 - 4.39 (m, 1H), 3.94 - 3.71 (m, 5H), 3.33 - 2.97 (m, 3H), 2.19 (br dd, *J*=12.8, 5.2 Hz, 1H), 2.12 - 2.00 (m, 1H), 1.83 - 1.75 (m, 1H), 1.74 - 1.52 (m, 2H), 1.45 -


1.28 (m, 1H); MS ESI m/z 481.9 (M+H); Anal. HPLC Retention time: 1.42 (Method 1); ROCK2 IC₅₀=251nM.

5 Example 150: Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-[4-methyl-1-oxo-1H,2H,3H,4H-pyrazino[1,2-a]indole-7-carbonyl]-2,8-diazaspiro[4.5]decan-1-one




¹H NMR (500 MHz, DMSO-d₆) δ 8.13 (br d, *J*=4.6 Hz, 1H), 8.01 (br s, 2H), 7.75 (d, *J*=8.2 Hz, 1H), 7.63 (s, 1H), 7.60 (br d, *J*=8.4 Hz, 1H), 7.56 (s, 1H), 7.17 - 7.06 (m, 3H), 4.98 - 4.86 (m, 1H), 3.85 (s, 2H), 3.74 (br s, 5H), 3.39 (br dd, *J*=12.5, 4.8 Hz, 1H), 3.32 - 3.22 (m, 1H), 3.15 (br dd, *J*=16.5, 4.2 Hz, 1H), 2.15 (br s, 2H), 1.84 - 1.60 (m, 3H), 1.50 (br d, *J*=9.3 Hz, 1H), 1.30 (br d, *J*=6.4 Hz, 3H); MS ESI m/z 553.3 (M+H); Anal. HPLC Retention time: 1.16 (Method 2); ROCK2 IC₅₀=144nM.

15 Example 151: Preparation of 8-[1-[(4-fluoro-3-methoxyphenyl)methyl]piperidine-4-carbonyl]-2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one



¹H NMR (500 MHz, DMSO-d₆) δ 8.02 (br s, 2H), 7.65 - 7.56 (m, 2H), 7.17 - 7.10 (m, 2H), 7.07 (br d, *J*=8.6 Hz, 1H), 6.84 (br d, *J*=6.2 Hz, 1H), 4.25 (br d, *J*=13.2 Hz, 1H), 3.89 (br s, 1H), 3.88 - 3.83 (m, 5H), 3.82 (s, 3H), 3.22 (br t, *J*=12.3 Hz, 1H), 2.91 - 2.74 (m, 3H), 2.60 (br d, *J*=4.3 Hz, 1H), 2.12 (br t, *J*=6.7 Hz, 2H), 1.99 (br d, *J*=8.5 Hz, 2H), 1.85 (br s, 1H), 1.74 - 1.42 (m, 9H); MS ESI *m/z* 576.2 (M+H); Anal. HPLC Retention time: 1.46 (Method 1); ROCK2 IC₅₀=2472nM.


Example 152: Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-(1-methyl-1H-indazole-5-carbonyl)-2,8-diazaspiro[4.5]decan-1-one



¹H NMR (500 MHz, DMSO-d₆) δ 8.14 (s, 1H), 8.07 - 7.94 (m, 2H), 7.86 (s, 1H), 7.71 (d, *J*=8.6 Hz, 1H), 7.64 - 7.58 (m, 2H), 7.44 (d, *J*=8.7 Hz, 1H), 7.15 (br d, *J*=8.5 Hz,


1H), 4.35 (br d, $J=8.7$ Hz, 1H), 4.25 - 4.17 (m, 1H), 4.07 (s, 3H), 3.87 (s, 5H), 3.43 (br s, 1H), 3.31 - 3.12 (m, 1H), 2.15 (br s, 2H), 1.85 - 1.69 (m, 2H), 1.59 (br s, 2H); MS ESI m/z 485.2 (M+H); Anal. HPLC Retention time: 1.21 (Method 2); ROCK2 $IC_{50}=6.2$ nM.

5 **Example 153: Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-(1-methyl-1H-indole-5-carbonyl)-2,8-diazaspiro[4.5]decan-1-one**




^1H NMR (500 MHz, DMSO- d_6) δ 8.01 (br s, 2H), 7.66 - 7.54 (m, 3H), 7.49 (br d, $J=8.4$ Hz, 1H), 7.38 (br d, $J=2.2$ Hz, 1H), 7.20 (br d, $J=8.3$ Hz, 1H), 7.13 (br d, $J=8.7$ Hz, 1H), 6.51 (br d, $J=2.1$ Hz, 1H), 3.85 (s, 2H), 3.67 (br s, 9H), 3.19 (br d, $J=18.9$ Hz, 2H), 2.14 (br s, 2H), 1.72 (br s, 2H), 1.56 (br d, $J=4.5$ Hz, 2H); MS ESI m/z 484.3 (M+H); Anal. HPLC Retention time: 1.42 Method 2; ROCK2 $IC_{50}=4.5$ nM.

15 **Example 154: Preparation of 8-(1-methanesulfonylpiperidine-4-carbonyl)-2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one**




¹H NMR (500 MHz, DMSO-d₆) δ 8.01 (br s, 2H), 7.65 - 7.55 (m, 2H), 7.14 (br d, *J*=8.5 Hz, 1H), 4.25 (br d, *J*=13.0 Hz, 1H), 3.93 (br d, *J*=13.8 Hz, 1H), 3.86 (s, 4H), 3.58 (br s, 5H), 3.25 (br t, *J*=11.6 Hz, 1H), 2.90 - 2.74 (m, 7H), 2.13 (br t, *J*=6.7 Hz, 2H), 1.73 (br s, 3H), 1.65 - 1.51 (m, 5H); MS ESI *m/z* 516.2 (M+H); Anal. HPLC Retention time: 1.07 (Method 2); ROCK2 IC₅₀=1995nM.

Example 155: Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-(1-methyl-1H-indole-6-carbonyl)-2,8-diazaspiro[4.5]decan-1-one




¹H NMR (500 MHz, DMSO-d₆) δ 8.01 (br s, 2H), 7.63 - 7.54 (m, 3H), 7.50 (s, 1H), 7.41 (br d, *J*=2.6 Hz, 1H), 7.13 (br d, *J*=8.0 Hz, 1H), 7.06 (br d, *J*=8.2 Hz, 1H), 6.48 (br d, *J*=2.3 Hz, 1H), 3.85 (s, 2H), 3.69 (br s, 8H), 3.35 - 3.08 (m, 2H), 2.15 (br s, 2H), 1.73 (br d, *J*=2.0 Hz, 2H), 1.66 - 1.44 (m, 2H); MS ESI *m/z* 484.2 (M+H); Anal. HPLC Retention time: 1.47 (Method 2); ROCK2 IC₅₀=40nM.

Example 156: Preparation of 8-(5-methoxy-1-methyl-1H-indole-2-carbonyl)-2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one



¹H NMR (500 MHz, DMSO-d₆) δ 8.02 (br s, 2H), 7.67 - 7.57 (m, 2H), 7.42 (d, *J*=8.9 Hz, 1H), 7.15 (br d, *J*=8.5 Hz, 1H), 7.10 (d, *J*=1.7 Hz, 1H), 6.89 (dd, *J*=8.9, 2.0 Hz, 1H), 6.57 (s, 1H), 4.48 - 4.29 (m, 1H), 4.17 - 3.96 (m, 1H), 3.87 (s, 5H), 3.76 (s, 3H),
 5 3.73 (s, 3H), 3.42 (br s, 1H), 3.17 (br s, 1H), 2.17 (br s, 2H), 1.83 - 1.71 (m, 2H), 1.63 (br s, 2H); MS ESI *m/z* 514.4 (M+H); Anal. HPLC Retention time: 1.61 (Method 1); ROCK2 IC₅₀=54nM.


Example 157: Preparation of 7-{2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-1-oxo-2,8-diazaspiro[4.5]decane-8-carbonyl}-3,4-dihydro-2H-1,4-benzoxazin-3-one
 10



¹H NMR (500 MHz, DMSO-d₆) δ 8.02 (br s, 2H), 7.68 - 7.56 (m, 2H), 7.14 (br d, *J*=8.7 Hz, 1H), 7.05 - 6.91 (m, 3H), 4.62 (s, 2H), 4.34 - 4.19 (m, 1H), 4.01 (s, 1H), 3.86


(s, 5H), 3.42 (br s, 1H), 3.31 - 3.06 (m, 1H), 2.13 (br s, 2H), 1.71 (br s, 2H), 1.57 (br d, $J=2.6$ Hz, 2H); MS ESI m/z 502.1 (M+H); Anal. HPLC Retention time: 1.12 (Method 1); ROCK2 $IC_{50}=22$ nM.

5 **Example 158: Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-[4-(pyridin-3-yl)benzoyl]-2,8-diazaspiro[4.5]decan-1-one**




^1H NMR (500 MHz, DMSO- d_6) δ 8.94 (br s, 1H), 8.60 (br d, $J=3.8$ Hz, 1H), 8.13 (br d, $J=7.7$ Hz, 1H), 8.11 - 7.91 (m, 2H), 7.83 (br d, $J=7.8$ Hz, 2H), 7.61 (br d, $J=2.9$ Hz, 2H), 7.55 (br d, $J=8.2$ Hz, 3H), 7.15 (br d, $J=8.5$ Hz, 1H), 4.35 (br d, $J=7.0$ Hz, 1H), 3.87 (s, 5H), 3.73 - 3.60 (m, 1H), 3.34 - 3.26 (m, 1H), 3.17 (br d, $J=4.9$ Hz, 1H), 2.16 (br s, 2H), 1.85 - 1.60 (m, 3H), 1.55 (br s, 1H); MS ESI m/z 508.2 (M+H); Anal. HPLC Retention time: 0.97 (Method 2); ROCK2 $IC_{50}=4.1$ nM.

15 **Example 159: Preparation of 8-[2-methoxy-4-(1H-pyrazol-4-yl)benzoyl]-2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one**



¹H NMR (500 MHz, DMSO-d₆) δ 8.41 - 8.19 (m, 1H), 8.16 - 7.98 (m, 2H), 7.61 (br d, *J*=7.3 Hz, 2H), 7.35 (br d, *J*=7.0 Hz, 1H), 7.29 (s, 1H), 7.25 (br s, 1H), 7.14 (br d, *J*=7.9 Hz, 2H), 4.92 - 4.80 (m, 1H), 4.48 - 4.31 (m, 1H), 3.96 (br d, *J*=6.4 Hz, 1H), 3.92 - 3.80 (m, 7H), 3.15 - 3.05 (m, 1H), 2.93 (br d, *J*=7.0 Hz, 1H), 2.21 - 2.09 (m, 2H), 1.84 - 1.69 (m, 2H), 1.63 (br d, *J*=12.5 Hz, 1H), 1.48 (br d, *J*=12.2 Hz, 1H); MS ESI *m/z* 527.1 (M+H); Anal. HPLC Retention time: 1.24 (Method 1); ROCK2 IC₅₀=29nM.


Example 160: Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-[4-(1,3-oxazol-5-yl)benzoyl]-2,8-diazaspiro[4.5]decan-1-one



¹H NMR (500 MHz, DMSO-d₆) δ 8.50 (s, 1H), 8.21 - 7.90 (m, 2H), 7.86 - 7.74 (m, 3H), 7.67 - 7.57 (m, 2H), 7.52 (br d, *J*=7.8 Hz, 2H), 7.15 (br d, *J*=8.4 Hz, 1H), 4.33 (br d, *J*=6.6 Hz, 1H), 3.87 (s, 5H), 3.68 - 3.58 (m, 1H), 3.35 - 3.23 (m, 1H), 3.22 - 3.09


(m, 1H), 2.15 (br s, 2H), 1.75 (br d, $J=4.1$ Hz, 3H), 1.55 (br s, 1H); MS ESI m/z 497.9 (M+H); Anal. HPLC Retention time: 1.36 (Method 1); ROCK2 $IC_{50}=23$ nM.

5 Example 161: Preparation of 8-(2-bromo-4-methoxybenzoyl)-2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one



^1H NMR (500 MHz, DMSO- d_6) δ 8.01 (br s, 2H), 7.61 - 7.52 (m, 2H), 7.29 - 7.19 (m, 2H), 7.11 (br t, $J=9.8$ Hz, 1H), 7.03 (br t, $J=9.0$ Hz, 1H), 4.35 (br d, $J=8.3$ Hz, 1H), 3.90 - 3.63 (m, 8H), 3.31 (br d, $J=9.4$ Hz, 1H), 3.23 - 3.03 (m, 2H), 2.21 - 2.06 (m, 2H), 1.93 - 1.70 (m, 2H), 1.63 (br d, $J=13.9$ Hz, 1H), 1.48 (br d, $J=9.6$ Hz, 1H); MS ESI m/z 539.3 (M+H); Anal. HPLC Retention time: 1.47 (Method 2); ROCK2 $IC_{50}=17$ nM.


15 Example 162: Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-[4-(5-methyl-1,3,4-oxadiazol-2-yl)benzoyl]-2,8-diazaspiro[4.5]decan-1-one



15


¹H NMR (500 MHz, DMSO-d₆) δ 8.05 (br d, *J*=8.1 Hz, 4H), 7.68 - 7.56 (m, 4H), 7.15 (br d, *J*=8.5 Hz, 1H), 4.41 - 4.26 (m, 1H), 3.87 (s, 5H), 3.63 - 3.54 (m, 1H), 3.27 (br d, *J*=11.0 Hz, 1H), 3.17 (br d, *J*=5.0 Hz, 1H), 2.60 (s, 3H), 2.16 (br d, *J*=5.2 Hz, 2H), 1.84 - 1.63 (m, 3H), 1.54 (br s, 1H); MS ESI *m/z* 513.3 (M+H); Anal. HPLC Retention time: 1.18 (Method 2); ROCK2 IC₅₀=27nM.

Example 163: Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-[4-(1,3,4-oxadiazol-2-yl)benzoyl]-2,8-diazaspiro[4.5]decan-1-one




¹H NMR (500 MHz, DMSO-d₆) δ 9.26 (s, 1H), 8.11 (br d, *J*=8.1 Hz, 2H), 8.01 (br s, 2H), 7.62 (br d, *J*=8.1 Hz, 3H), 7.55 (s, 1H), 7.12 (br d, *J*=8.4 Hz, 1H), 4.35 (br d, *J*=11.2 Hz, 1H), 3.80 (br s, 5H), 3.57 (br d, *J*=13.1 Hz, 1H), 3.32 - 3.21 (m, 1H), 3.18 - 3.08 (m, 1H), 2.24 - 2.07 (m, 2H), 1.82 - 1.61 (m, 3H), 1.51 (br d, *J*=10.7 Hz, 1H); MS ESI *m/z* 499.1 (M+H); Anal. HPLC Retention time: 1.23 (Method 1); ROCK2 IC₅₀=16nM.

Example 164: Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-(1-methyl-1H-1,3-benzodiazole-5-carbonyl)-2,8-diazaspiro[4.5]decan-1-one



¹H NMR (500 MHz, DMSO-d₆) δ 8.29 (s, 1H), 8.02 (br d, *J*=1.8 Hz, 2H), 7.71 - 7.56 (m, 4H), 7.34 (br d, *J*=8.2 Hz, 1H), 7.15 (br d, *J*=8.2 Hz, 1H), 4.44 - 4.27 (m, 1H), 3.95 - 3.78 (m, 8H), 3.74 - 3.62 (m, 1H), 3.37 (br s, 1H), 3.17 (d, *J*=5.0 Hz, 1H), 2.15 (br s, 2H), 1.75 (br s, 2H), 1.59 (br d, *J*=9.9 Hz, 2H); MS ESI *m/z* 485 (M+H); Anal. HPLC Retention time: 1.17 (Method 1); ROCK2 IC₅₀=2.6nM.

Example 165: Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-(2-methyl-1,6-naphthyridine-3-carbonyl)-2,8-diazaspiro[4.5]decan-1-one




10

¹H NMR (500 MHz, DMSO-d₆) δ 9.39 (br s, 1H), 8.74 (br d, *J*=5.8 Hz, 1H), 8.54 - 8.39 (m, 1H), 8.02 (br d, *J*=5.5 Hz, 2H), 7.87 (br d, *J*=5.8 Hz, 1H), 7.61 (br d, *J*=8.2 Hz, 2H), 7.15 (br s, 1H), 4.54 - 4.35 (m, 1H), 3.96 - 3.80 (m, 5H), 3.46 (s, 1H), 3.27 (br t, *J*=11.3 Hz, 2H), 2.66 (br s, 3H), 2.25 - 2.08 (m, 2H), 1.90 - 1.64 (m, 3H), 1.49 (br d,


$J=5.5$ Hz, 1H); MS ESI m/z 496.9 (M+H); Anal. HPLC Retention time: 1.07 (Method 2); ROCK2 $IC_{50}=896$ nM.

5 Example 166: Preparation of 8-(isoquinoline-6-carbonyl)-2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one




^1H NMR (500 MHz, DMSO- d_6) δ 9.38 (s, 1H), 8.57 (d, $J=5.7$ Hz, 1H), 8.22 (d, $J=8.3$ Hz, 1H), 8.03 (br d, $J=12.2$ Hz, 3H), 7.94 (br d, $J=5.6$ Hz, 1H), 7.69 (br d, $J=8.2$ Hz, 1H), 7.64 - 7.58 (m, 2H), 7.15 (br d, $J=8.6$ Hz, 1H), 4.39 (br s, 1H), 3.96 - 3.78 (m, 5H), 3.60 (br d, $J=9.8$ Hz, 1H), 3.26 (br dd, $J=19.9, 12.8$ Hz, 1H), 3.17 (s, 1H), 2.30 - 2.07 (m, 2H), 1.87 - 1.66 (m, 3H), 1.52 (br d, $J=10.4$ Hz, 1H); MS ESI m/z 482.2 (M+H); Anal. HPLC Retention time: 0.89 (Method 2); ROCK2 $IC_{50}=33$ nM.

15 Example 167: Preparation of 8-[4-(hydroxymethyl)benzoyl]-2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one




¹H NMR (500 MHz, DMSO-d₆) δ 8.01 (br s, 2H), 7.59 (d, *J*=8.4 Hz, 1H), 7.56 (s, 1H), 7.43 - 7.29 (m, 4H), 7.12 (br d, *J*=8.3 Hz, 1H), 4.53 (br s, 2H), 4.38 - 4.28 (m, 1H), 3.84 (s, 2H), 3.75 (br s, 3H), 3.64 - 3.55 (m, 1H), 3.23 (br s, 1H), 3.09 (br d, *J*=2.6 Hz, 1H), 2.13 (br s, 2H), 1.80 - 1.58 (m, 3H), 1.50 (br d, *J*=7.5 Hz, 1H); MS ESI *m/z* 460.9 (M+H); Anal. HPLC Retention time: 1.2 (Method 1); ROCK2 IC₅₀=53nM.

Example 168: Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-[4-[(pyrrolidin-1-yl)methyl]benzoyl]-2,8-diazaspiro[4.5]decan-1-one




¹H NMR (500 MHz, DMSO-d₆) δ 8.02 (s, 2H), 7.66 - 7.55 (m, 2H), 7.44 - 7.31 (m, 4H), 7.14 (br d, *J*=8.8 Hz, 1H), 4.38 - 4.26 (m, 1H), 3.93 - 3.78 (m, 5H), 3.60 (s, 1H), 3.44 (br d, *J*=11.8 Hz, 1H), 3.23 (br d, *J*=10.0 Hz, 1H), 3.17 (s, 1H), 3.15 - 3.06 (m, 1H), 2.43 (br s, 4H), 2.14 (br s, 2H), 1.89 (s, 4H), 1.69 (br s, 7H), 1.52 (br s, 1H); MS ESI *m/z* 514.2 (M+H); Anal. HPLC Retention time: 1.07 (Method 1); ROCK2 IC₅₀=17nM.

Example 169: Preparation of 8-[4-(methanesulfonylmethyl)benzoyl]-2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one



¹H NMR (500 MHz, DMSO-d₆) δ 8.02 (br s, 2H), 7.66 - 7.57 (m, 2H), 7.53 - 7.39 (m, 4H), 7.15 (br d, *J*=8.5 Hz, 1H), 4.55 (s, 2H), 4.34 (br s, 1H), 3.87 (s, 5H), 3.59 (br d, *J*=9.6 Hz, 1H), 3.25 (br d, *J*=10.9 Hz, 1H), 3.19 - 3.11 (m, 1H), 2.95 (s, 3H), 2.15 (br s, 2H), 1.83 - 1.61 (m, 3H), 1.53 (br s, 1H); MS ESI *m/z* 523.1 (M+H); Anal. HPLC Retention time: 1.19 (Method 1); ROCK2 IC₅₀=13nM.

Example 170: Preparation of 8-(6-methoxy-1-methyl-1H-indole-2-carbonyl)-2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one




10

¹H NMR (500 MHz, DMSO-d₆) δ 8.18 - 7.91 (m, 2H), 7.61 (br d, *J*=6.1 Hz, 2H), 7.48 (d, *J*=8.7 Hz, 1H), 7.15 (br d, *J*=8.1 Hz, 1H), 7.02 (s, 1H), 6.75 (br d, *J*=8.8 Hz, 1H), 6.61 (s, 1H), 4.39 - 4.06 (m, 1H), 3.87 (s, 5H), 3.82 (s, 3H), 3.73 (s, 2H), 3.47 (br s, 5H), 2.25 - 2.13 (m, 2H), 1.83 - 1.70 (m, 2H), 1.63 (br d, *J*=10.2 Hz, 2H); MS ESI *m/z* 514.1 (M+H); Anal. HPLC Retention time: 1.62 (Method 1); ROCK2 IC₅₀=118nM.

15


Example 171: Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-[4-(pyridin-4-yl)benzoyl]-2,8-diazaspiro[4.5]decan-1-one



5 ^1H NMR (500 MHz, DMSO- d_6) δ 8.66 (br d, $J=5.3$ Hz, 2H), 8.01 (br s, 2H), 7.89 (br d, $J=8.0$ Hz, 2H), 7.76 (br d, $J=5.5$ Hz, 2H), 7.65 - 7.52 (m, 4H), 7.14 (br d, $J=8.6$ Hz, 1H), 4.43 - 4.29 (m, 1H), 3.86 (s, 4H), 3.54 (br s, 2H), 3.29 (br s, 1H), 3.17 (br d, $J=5.0$ Hz, 1H), 2.15 (br s, 2H), 1.83 - 1.61 (m, 3H), 1.54 (br s, 1H); MS ESI m/z 508.1 (M+H); Anal. HPLC Retention time: 1.36 (Method 1); ROCK2 $\text{IC}_{50}=5.5\text{nM}$.

10


Example 172: Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-(1-methylpiperidine-4-carbonyl)-2,8-diazaspiro[4.5]decan-1-one



15 ^1H NMR (500 MHz, DMSO- d_6) δ 8.01 (br s, 2H), 7.66 - 7.56 (m, 2H), 7.15 (br d, $J=8.5$ Hz, 1H), 4.25 (br d, $J=12.2$ Hz, 1H), 3.97 - 3.82 (m, 6H), 3.53 - 3.46 (m, 1H), 3.24 (br s, 1H), 2.86 (br s, 3H), 2.67 - 2.57 (m, 1H), 2.23 (s, 3H), 2.17 - 2.03 (m, 4H), 1.74 -


1.47 (m, 8H); MS ESI m/z 452.2 (M+H); Anal. HPLC Retention time: 1.01 (Method 2); ROCK2 IC_{50} =2068nM.

5 Example 173: Preparation of 8-{4'-methanesulfonyl-[1,1'-biphenyl]-4-carbonyl}-2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one




^1H NMR (500 MHz, DMSO- d_6) δ 8.20 - 7.90 (m, 6H), 7.85 (br d, J =8.0 Hz, 2H), 7.67 - 7.59 (m, 2H), 7.56 (br d, J =8.0 Hz, 2H), 7.15 (br d, J =8.4 Hz, 1H), 4.42 - 4.29 (m, 1H), 3.87 (s, 4H), 3.73 - 3.61 (m, 1H), 3.53 - 3.48 (m, 1H), 3.26 (s, 3H), 3.21 - 3.14 (m, 1H), 2.15 (br s, 2H), 1.87 - 1.63 (m, 3H), 1.55 (br s, 1H); MS ESI m/z 585.1 (M+H); Anal. HPLC Retention time: 1.46 (Method 2); ROCK2 IC_{50} =3.7nM.

15 Example 174: Preparation of 8-(2,4-dimethoxybenzoyl)-2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one



¹H NMR (500 MHz, DMSO-d₆) δ 8.02 (s, 2H), 7.60 (br d, *J*=7.7 Hz, 2H), 7.27 (s, 1H), 7.17 (s, 1H), 7.13 (br d, *J*=8.0 Hz, 2H), 7.06 (s, 1H), 6.66 - 6.53 (m, 2H), 4.36 (br s, 1H), 3.95 - 3.74 (m, 9H), 3.59 (br s, 2H), 3.37 (br d, *J*=11.3 Hz, 1H), 3.12 - 2.89 (m, 2H),
 5 2.11 (br d, *J*=17.0 Hz, 2H), 1.70 (br d, *J*=11.4 Hz, 2H), 1.60 (br d, *J*=13.0 Hz, 1H), 1.45 (br d, *J*=13.0 Hz, 1H); MS ESI *m/z* 490.9 (M+H); Anal. HPLC Retention time: 1.44 (Method 2); ROCK2 IC₅₀=29nM.


Example 175: Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-[1-(pyridin-4-yl)piperidin-4-carbonyl]-2,8-diazaspiro[4.5]decan-1-one



¹H NMR (500 MHz, DMSO-d₆) δ 8.11 (br d, *J*=4.2 Hz, 2H), 8.02 (br s, 2H), 7.66 - 7.55 (m, 2H), 7.14 (br d, *J*=8.4 Hz, 1H), 6.81 (br d, *J*=5.1 Hz, 2H), 4.25 (br d, *J*=13.4 Hz, 1H), 3.94 (br d, *J*=11.4 Hz, 3H), 3.86 (s, 4H), 3.26 (br t, *J*=11.5 Hz, 1H), 2.96 (br d, *J*=13.5 Hz, 4H), 2.14 (br t, *J*=6.6 Hz, 2H), 1.89 (s, 2H), 1.69 (br s, 3H), 1.57 (br d, *J*=14.6
 15


Hz, 5H); MS ESI m/z 515.2 (M+H); Anal. HPLC Retention time: 1.02 (Method 1); ROCK2 IC₅₀=195nM.

5 Example 176: Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-(6-methoxynaphthalene-2-carbonyl)-2,8-diazaspiro[4.5]decan-1-one



¹H NMR (500 MHz, DMSO-d₆) δ 8.01 (br s, 2H), 7.95 - 7.86 (m, 3H), 7.66 - 7.56 (m, 2H), 7.47 (br d, J =8.5 Hz, 1H), 7.37 (s, 1H), 7.22 (dd, J =9.0, 2.0 Hz, 1H), 7.14 (br d, J =8.4 Hz, 1H), 3.87 (br d, J =12.4 Hz, 6H), 3.64 - 3.62 (m, 1H), 3.35 - 3.23 (m, 1H), 3.20 - 3.11 (m, 1H), 2.15 (br s, 2H), 1.84 - 1.49 (m, 4H); MS ESI m/z 511.2 (M+H); Anal. HPLC Retention time: 1.69 (Method 1); ROCK2 IC₅₀=20nM.

10 Example 177: Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-(2-oxo-2,3-dihydro-1H-indole-5-carbonyl)-2,8-diazaspiro[4.5]decan-1-one




15

¹H NMR (500 MHz, DMSO-d₆) δ 8.01 (br s, 2H), 7.65 - 7.52 (m, 2H), 7.32 - 7.22 (m, 2H), 7.13 (br d, *J*=8.2 Hz, 1H), 6.89 (d, *J*=7.9 Hz, 1H), 3.85 (s, 3H), 3.71 - 3.51 (m, 8H), 2.14 (br s, 2H), 1.71 (br s, 2H), 1.57 (br s, 2H); MS ESI *m/z* 485.9 (M+H); Anal. HPLC Retention time: 1.13 (Method 2); ROCK2 IC₅₀=78nM.


5

Example 178: Preparation of 8-(5,6-dimethoxy-1-methyl-1H-indole-2-carbonyl)-2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one




¹H NMR (500 MHz, DMSO-d₆) δ 8.02 (br s, 2H), 7.67 - 7.56 (m, 2H), 7.15 (br d, *J*=8.5 Hz, 1H), 7.08 (d, *J*=14.4 Hz, 2H), 6.55 (s, 1H), 4.35 - 4.11 (m, 1H), 3.87 (s, 5H), 3.84 (s, 3H), 3.75 (d, *J*=9.7 Hz, 6H), 2.17 (br t, *J*=6.4 Hz, 2H), 1.81 - 1.69 (m, 2H), 1.62 (br d, *J*=13.0 Hz, 2H); MS ESI *m/z* 544.1 (M+H); Anal. HPLC Retention time: 1.4 (Method 2); ROCK2 IC₅₀=14nM.

Example 179: Preparation of 8-(1,2-dimethyl-1H-1,3-benzodiazole-5-carbonyl)-2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one




¹H NMR (500 MHz, DMSO-d₆) δ 8.19 - 7.87 (m, 2H), 7.66 - 7.58 (m, 2H), 7.56 - 7.50 (m, 2H), 7.25 (br d, *J*=8.5 Hz, 1H), 7.14 (br d, *J*=8.5 Hz, 1H), 3.86 (s, 7H), 3.75 (s, 4H), 3.42 (br s, 4H), 3.28 - 3.12 (m, 1H), 2.15 (br s, 2H), 1.83 - 1.43 (m, 4H); MS ESI m/z 499.1 (M+H); Anal. HPLC Retention time: 1.22 (Method 1); ROCK2 IC₅₀=2.3nM.

Example 180: Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-{4-[(morpholin-4-yl)methyl]benzoyl}-2,8-diazaspiro[4.5]decan-1-one




¹H NMR (500 MHz, DMSO-d₆) δ 8.02 (br s, 2H), 7.64 - 7.58 (m, 2H), 7.45 - 7.32 (m, 4H), 7.15 (br d, *J*=8.5 Hz, 1H), 4.34 (br d, *J*=10.1 Hz, 1H), 3.87 (s, 5H), 3.58 (br d, *J*=4.0 Hz, 4H), 3.50 (s, 2H), 3.40 (br s, 1H), 3.22 (br s, 1H), 3.15 - 3.09 (m, 1H), 2.37 (br s, 4H), 2.14 (br s, 2H), 1.80 - 1.47 (m, 4H); MS ESI m/z 530.2 (M+H); Anal. HPLC Retention time: 1.31 (Method 1); ROCK2 IC₅₀=27nM.

Example 181: Preparation of 4-{2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-1-oxo-2,8-diazaspiro[4.5]decane-8-carbonyl}benzamide




5 ^1H NMR (500 MHz, DMSO- d_6) δ 8.13 (br s, 1H), 8.01 (br s, 2H), 7.91 (br d, $J=8.0$ Hz, 2H), 7.59 (d, $J=8.4$ Hz, 1H), 7.56 (s, 1H), 7.47 (br d, $J=7.7$ Hz, 3H), 7.12 (br d, $J=8.1$ Hz, 1H), 4.34 (br d, $J=11.8$ Hz, 1H), 3.74 (br s, 5H), 3.53 (br d, $J=11.1$ Hz, 1H), 3.23 (br d, $J=10.0$ Hz, 1H), 3.18 - 3.05 (m, 1H), 2.22 - 2.06 (m, 2H), 1.81 - 1.61 (m, 3H), 1.50 (br d, $J=12.4$ Hz, 1H); MS ESI m/z 474.1 (M+H); Anal. HPLC Retention time: 0.97
10 (Method 2); ROCK2 $\text{IC}_{50}=35\text{nM}$.

Example 182: Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-[2-(pyridin-4-yl)-1H-1,3-benzodiazole-6-carbonyl]-2,8-diazaspiro[4.5]decan-1-one



¹H NMR (500 MHz, DMSO-d₆) δ 8.79 (br d, *J*=5.1 Hz, 2H), 8.11 (br d, *J*=5.1 Hz, 2H), 8.02 (br s, 2H), 7.72 (br s, 2H), 7.65 - 7.57 (m, 2H), 7.34 (br d, *J*=7.9 Hz, 1H), 7.16 (br d, *J*=7.9 Hz, 1H), 4.44 - 4.24 (m, 1H), 3.96 - 3.79 (m, 5H), 3.31 - 3.22 (m, 1H), 3.17 (br s, 1H), 2.89 (s, 1H), 2.16 (br s, 2H), 1.76 (br s, 2H), 1.68 - 1.48 (m, 2H); MS ESI *m/z* 548.2 (M+H); Anal. HPLC Retention time: 1.21 (Method 1); ROCK2 IC₅₀=16nM.


Example 183: Preparation of 8-{4-[(1H-imidazol-1-yl)methyl]benzoyl}-2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one



¹H NMR (500 MHz, DMSO-d₆) δ 8.01 (s, 2H), 7.80 (br s, 1H), 7.59 (d, *J*=8.4 Hz, 1H), 7.55 (s, 1H), 7.42 - 7.36 (m, 2H), 7.32 (br d, *J*=7.9 Hz, 2H), 7.22 (br s, 1H), 7.12 (br d, *J*=8.2 Hz, 1H), 6.93 (br s, 1H), 5.23 (s, 2H), 4.36 - 4.24 (m, 1H), 3.84 (s, 1H), 3.79 - 3.67 (m, 4H), 3.61 - 3.47 (m, 1H), 3.28 - 3.17 (m, 1H), 3.10 (br d, *J*=9.7 Hz, 1H), 2.12 (br


d, $J=6.3$ Hz, 2H), 1.76 - 1.56 (m, 3H), 1.48 (br d, $J=9.6$ Hz, 1H); MS ESI m/z 511.2 (M+H); Anal. HPLC Retention time: 1.24 (Method 1); ROCK2 $IC_{50}=23$ nM.

5 Example 184: Preparation of 8-[2-(benzyloxy)-4-methoxybenzoyl]-2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one




^1H NMR (500 MHz, DMSO- d_6) δ 8.02 (br s, 2H), 7.69 - 7.54 (m, 2H), 7.53 - 7.38 (m, 4H), 7.37 - 7.29 (m, 1H), 7.21 - 6.99 (m, 2H), 6.70 (br s, 1H), 6.64 - 6.53 (m, 1H), 5.16 (br s, 2H), 4.29 (br d, $J=8.8$ Hz, 1H), 3.92 - 3.72 (m, 8H), 3.46 (br s, 1H), 3.21 - 3.05 (m, 2H), 2.16 - 1.92 (m, 2H), 1.84 - 1.60 (m, 2H), 1.55 (br s, 1H), 1.50 - 1.37 (m, 1H); MS ESI m/z 567 (M+H); Anal. HPLC Retention time: 1.78 (Method 1); ROCK2 $IC_{50}=12$ nM.

15 Example 185: Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-[4-(3-methyl-1,2,4-oxadiazol-5-yl)benzoyl]-2,8-diazaspiro[4.5]decan-1-one



¹H NMR (500 MHz, DMSO-d₆) δ 8.17 (br d, *J*=7.9 Hz, 2H), 8.12 - 7.85 (m, 2H), 7.72 - 7.54 (m, 4H), 7.15 (br d, *J*=8.4 Hz, 1H), 4.35 (br d, *J*=12.1 Hz, 1H), 3.86 (s, 5H), 3.63 - 3.52 (m, 1H), 3.28 (br s, 1H), 3.22 - 3.12 (m, 1H), 2.43 (s, 3H), 2.24 - 2.07 (m, 2H), 1.88 - 1.61 (m, 3H), 1.53 (br d, *J*=9.4 Hz, 1H); MS ESI *m/z* 513.3 (M+H); Anal. HPLC Retention time: 1.37 (Method 2); ROCK2 IC₅₀=37nM.

Example 186: Preparation of 8-(4-{4-[(3-chlorophenyl)methyl]piperazin-1-yl}benzoyl)-2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one




10

¹H NMR (500 MHz, DMSO-d₆) δ 8.02 (br d, *J*=2.3 Hz, 2H), 7.70 - 7.53 (m, 2H), 7.43 - 7.23 (m, 6H), 7.14 (br d, *J*=8.5 Hz, 1H), 6.95 (br d, *J*=8.5 Hz, 2H), 4.11 - 3.97 (m, 1H), 3.95 - 3.72 (m, 5H), 3.53 (s, 1H), 3.45 (br s, 2H), 3.30 - 3.07 (m, 4H), 2.51 (br s,


10H), 2.14 (br t, $J=6.2$ Hz, 2H), 1.79 - 1.65 (m, 2H), 1.56 (br d, $J=10.2$ Hz, 2H); MS ESI m/z 639.3 (M+H); Anal. HPLC Retention time: 1.28 (Method 2); ROCK2 $IC_{50}=1.2$ nM.

Example 187: Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-(5-methyl-1H-pyrazole-3-carbonyl)-2,8-diazaspiro[4.5]decan-1-one



^1H NMR (500 MHz, DMSO- d_6) δ 8.02 (br s, 2H), 7.66 - 7.51 (m, 2H), 7.15 (br d, $J=8.5$ Hz, 1H), 6.30 (s, 1H), 4.57 (br s, 1H), 4.35 (br d, $J=11.8$ Hz, 1H), 3.86 (s, 5H), 3.41 - 3.29 (m, 1H), 3.04 (br d, $J=10.6$ Hz, 1H), 2.25 (s, 3H), 2.16 (br d, $J=5.6$ Hz, 2H), 1.71 (br d, $J=10.4$ Hz, 2H), 1.63 - 1.51 (m, 2H); MS ESI m/z 434.9 (M+H); Anal. HPLC Retention time: 1.18 (Method 1); ROCK2 $IC_{50}=27$ nM.


Example 188: Preparation of 8-(1,5-dimethyl-1H-pyrazole-3-carbonyl)-2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one



^1H NMR (500 MHz, DMSO- d_6) δ 8.02 (br s, 2H), 7.66 - 7.54 (m, 2H), 7.15 (br d, $J=8.4$ Hz, 1H), 6.33 (s, 1H), 4.55 (br d, $J=12.7$ Hz, 1H), 4.34 (br d, $J=11.2$ Hz, 1H), 3.86


(s, 5H), 3.75 (s, 3H), 3.40 - 3.32 (m, 1H), 3.09 - 2.98 (m, 1H), 2.26 (s, 3H), 2.16 (br d, $J=5.6$ Hz, 2H), 1.69 (br s, 2H), 1.63 - 1.48 (m, 2H); MS ESI m/z 449.1 (M+H); Anal. HPLC Retention time: 1.23 (Method 1); ROCK2 $IC_{50}=31$ nM.

5 **Example 189: Preparation of 8-(4-amino-2-chlorobenzoyl)-2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one**




^{1} H NMR (500 MHz, DMSO- d_6) δ 8.02 (br s, 2H), 7.70 - 7.52 (m, 2H), 7.12 (br s, 1H), 7.04 - 6.89 (m, 1H), 6.62 (br s, 1H), 6.54 (br s, 1H), 5.63 (br s, 2H), 4.40 - 4.26 (m, 1H), 3.86 (s, 5H), 3.52 - 3.45 (m, 1H), 3.29 - 3.01 (m, 2H), 2.13 (br s, 2H), 1.86 - 1.57 (m, 3H), 1.48 (br d, $J=12.3$ Hz, 1H); MS ESI m/z 480.2 (M+H); Anal. HPLC Retention time: 1.12 (Method 2); ROCK2 $IC_{50}=38$ nM.

15 **Example 190: Preparation of 8-[4-(dimethylamino)benzoyl]-2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one**




¹H NMR (500 MHz, DMSO-d₆) δ 8.19 - 7.85 (m, 2H), 7.65 - 7.54 (m, 2H), 7.29 (br d, *J*=8.4 Hz, 2H), 7.14 (br d, *J*=8.4 Hz, 1H), 6.72 (br d, *J*=8.4 Hz, 2H), 4.14 - 3.93 (m, 1H), 3.91 - 3.73 (m, 5H), 3.45 (br s, 2H), 3.17 (br d, *J*=4.3 Hz, 1H), 2.94 (s, 6H), 2.14 (br t, *J*=6.6 Hz, 2H), 1.70 (br d, *J*=9.2 Hz, 2H), 1.56 (br d, *J*=12.1 Hz, 2H); MS ESI *m/z* 474.4 (M+H); Anal. HPLC Retention time: 1.12 (Method 2); ROCK2 IC₅₀=16nM.

Example 191: Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-[4-(pyrrolidin-1-yl)benzoyl]-2,8-diazaspiro[4.5]decan-1-one




¹H NMR (500 MHz, DMSO-d₆) δ 8.02 (br s, 2H), 7.61 (br d, *J*=5.2 Hz, 2H), 7.28 (br d, *J*=7.9 Hz, 2H), 7.15 (br d, *J*=8.5 Hz, 1H), 6.55 (br d, *J*=8.2 Hz, 2H), 4.05 (br s, 1H), 3.90 - 3.79 (m, 5H), 3.41 (br s, 1H), 3.26 (br s, 4H), 3.18 (br d, *J*=4.0 Hz, 2H), 3.02 (br d, *J*=3.4 Hz, 1H), 2.15 (br t, *J*=6.6 Hz, 2H), 1.96 (br s, 4H), 1.78 - 1.67 (m, 3H), 1.56 (br d, *J*=12.5 Hz, 2H); MS ESI *m/z* 499.9 (M+H); Anal. HPLC Retention time: 1.62 (Method 2); ROCK2 IC₅₀=4nM.

Example 192: Preparation of 8-(4-(1,1-dioxido-1,2-thiazinan-2-yl)benzoyl)-2-(3-methoxy-4-(1H-pyrazol-4-yl)phenyl)-2,8-diazaspiro[4.5]decan-1-one



¹H NMR (500 MHz, DMSO-d₆) δ 8.02 (br s, 2H), 7.65 - 7.56 (m, 2H), 7.48 - 7.35 (m, 4H), 7.15 (br d, *J*=8.1 Hz, 1H), 4.38 - 4.28 (m, 1H), 3.86 (s, 4H), 3.78 - 3.62 (m, 2H), 3.45 (br s, 1H), 3.36 - 3.24 (m, 2H), 3.21 - 3.17 (m, 1H), 3.14 (br s, 1H), 2.16 (br d, *J*=5.3 Hz, 4H), 1.89 - 1.46 (m, 6H); MS ESI *m/z* 564.1 (M+H); Anal. HPLC Retention time: 1.25 (Method 2); ROCK2 IC₅₀=3.3nM.

Example 193: Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-[4-(1H-pyrrol-1-yl)benzoyl]-2,8-diazaspiro[4.5]decan-1-one




10

¹H NMR (500 MHz, DMSO-d₆) δ 8.01 (br s, 2H), 7.65 (br d, *J*=8.2 Hz, 2H), 7.59 (br d, *J*=5.4 Hz, 2H), 7.50 (br d, *J*=8.2 Hz, 2H), 7.42 (br s, 2H), 7.14 (br d, *J*=8.4 Hz, 1H), 6.30 (br s, 2H), 4.09 - 4.04 (m, 1H), 3.86 (s, 5H), 3.61 (br s, 1H), 3.37 - 3.23 (m, 1H), 3.16 (br s, 1H), 2.15 (br s, 2H), 1.73 (br d, *J*=6.3 Hz, 2H), 1.68 - 1.47 (m, 2H); MS ESI *m/z* 496.1 (M+H); Anal. HPLC Retention time: 1.61 (Method 2); ROCK2 IC₅₀=12nM.

15


Example 194: Preparation of 8-[4-(diethylamino)benzoyl]-2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one



5 ^1H NMR (500 MHz, DMSO- d_6) δ 8.02 (br s, 2H), 7.66 - 7.52 (m, 2H), 7.27 (br d, $J=8.5$ Hz, 2H), 7.15 (br d, $J=8.2$ Hz, 1H), 6.66 (br d, $J=8.5$ Hz, 2H), 4.06 (br s, 2H), 3.91 - 3.79 (m, 5H), 3.47 - 3.29 (m, 3H), 3.18 (br d, $J=4.3$ Hz, 3H), 2.15 (br t, $J=6.7$ Hz, 2H), 1.71 (br d, $J=9.2$ Hz, 2H), 1.57 (br d, $J=12.8$ Hz, 2H), 1.11 (br t, $J=7.0$ Hz, 6H); MS ESI m/z 502.2 (M+H); Anal. HPLC Retention time: 1.76 (Method 1); ROCK2 $\text{IC}_{50}=5.2\text{nM}$.

10


Example 195: Preparation of 8-(4-bromobenzoyl)-2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one



15 ^1H NMR (500 MHz, DMSO- d_6) δ 8.02 (s, 2H), 7.67 (br d, $J=8.2$ Hz, 2H), 7.64 - 7.57 (m, 2H), 7.38 (br d, $J=8.2$ Hz, 2H), 7.15 (br d, $J=8.2$ Hz, 1H), 4.37 - 4.26 (m, 1H), 3.87 (s, 5H), 3.51 (br s, 1H), 3.30 - 3.21 (m, 1H), 3.18 (s, 1H), 2.14 (br s, 2H), 1.85 - 1.59


(m, 3H), 1.53 (br s, 1H); MS ESI m/z 509.1 (M+H); Anal. HPLC Retention time: 1.59 (Method 1); ROCK2 IC₅₀=33nM.

5 Example 196: Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-(1-methyl-1H-1,2,3-benzotriazole-5-carbonyl)-2,8-diazaspiro[4.5]decan-1-one



¹H NMR (500 MHz, DMSO-d₆) δ 8.08 (s, 1H), 8.01 (br s, 2H), 7.94 (d, *J*=8.9 Hz, 1H), 7.60 (br s, 3H), 7.15 (br d, *J*=8.5 Hz, 1H), 4.34 (s, 3H), 4.11 - 4.02 (m, 1H), 3.87 (s, 5H), 3.68 - 3.60 (m, 1H), 3.29 (br s, 1H), 3.24 (br s, 1H), 2.16 (br s, 2H), 1.84 - 1.50 (m, 4H); MS ESI m/z 486.2 (M+H); Anal. HPLC Retention time: 1.21 (Method 1); ROCK2 IC₅₀=9.1nM.


15 Example 197: Preparation of 8-(1H-1,3-benzodiazole-4-carbonyl)-2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one



^1H NMR (500 MHz, DMSO- d_6) δ 8.27 (s, 1H), 8.01 (s, 2H), 7.68 - 7.54 (m, 3H), 7.33 - 7.10 (m, 3H), 4.52 - 4.40 (m, 1H), 3.87 (s, 5H), 3.51 (br s, 1H), 3.31 - 3.10 (m, 2H), 2.16 (br s, 2H), 1.87 - 1.74 (m, 2H), 1.69 (br d, $J=3.1$ Hz, 1H), 1.43 (br s, 1H); MS ESI m/z 470.9 (M+H); Anal. HPLC Retention time: 1.27 (Method 1); ROCK2

5 $\text{IC}_{50}=61\text{nM}$.

Example 198: Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-[pyrazolo[1,5-a]pyrimidine-3-carbonyl]-2,8-diazaspiro[4.5]decan-1-one




10

^1H NMR (500 MHz, DMSO- d_6) δ 9.18 (dd, $J=7.0$, 1.5 Hz, 1H), 8.72 (br d, $J=2.4$ Hz, 1H), 8.44 (s, 1H), 8.15 - 7.87 (m, 2H), 7.61 (dd, $J=4.7$, 3.2 Hz, 2H), 7.25 - 7.09 (m, 2H), 4.46 - 4.29 (m, 1H), 3.87 (s, 5H), 3.38 - 3.25 (m, 1H), 3.21 - 3.09 (m, 1H), 2.17 (br s, 2H), 1.83 (br s, 2H), 1.72 - 1.48 (m, 2H); MS ESI m/z 472.2 (M+H); Anal. HPLC


15 Retention time: 1.23 (Method 1); ROCK2 $\text{IC}_{50}=1060\text{nM}$.

Example 199: Preparation of 8-(4,7-dimethoxynaphthalene-1-carbonyl)-2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one



¹H NMR (500 MHz, DMSO-d₆) δ 8.13 (br d, *J*=9.3 Hz, 1H), 8.01 (br s, 2H), 7.60 (br d, *J*=8.2 Hz, 2H), 7.42 - 7.27 (m, 1H), 7.24 - 7.12 (m, 2H), 7.11 - 6.96 (m, 1H), 6.88 (br d, *J*=7.2 Hz, 1H), 4.60 (br d, *J*=10.2 Hz, 1H), 3.98 (s, 3H), 3.93 (br s, 3H), 3.86 (s, 5H), 3.56 - 3.49 (m, 1H), 3.31 (br d, *J*=10.8 Hz, 1H), 3.15 - 3.09 (m, 1H), 2.18 (br d, *J*=4.9 Hz, 1H), 2.11 (br s, 1H), 1.89 - 1.80 (m, 1H), 1.77 - 1.68 (m, 1H), 1.58 (br s, 1H), 1.38 (br d, *J*=10.6 Hz, 1H); MS ESI *m/z* 541 (M+H); Anal. HPLC Retention time: 1.74 (Method 2); ROCK2 IC₅₀=186nM.

10 **Example 200: Preparation of 8-[4-hydroxy-7-(propan-2-yl)pyrrolo[2,1-f][1,2,4]triazine-5-carbonyl]-2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one**




¹H NMR (500 MHz, DMSO-d₆) δ 8.08 (br s, 1H), 7.94 (br s, 1H), 7.85 (s, 1H), 7.66 - 7.53 (m, 2H), 7.13 (br d, *J*=8.2 Hz, 1H), 6.41 (s, 1H), 4.39 (br d, *J*=12.8 Hz, 1H), 3.88 -

3.80 (m, 5H), 3.60 (br d, $J=13.7$ Hz, 1H), 3.37 - 3.26 (m, 1H), 3.20 - 3.02 (m, 2H), 2.20 - 2.00 (m, 2H), 1.81 - 1.66 (m, 2H), 1.62 (br d, $J=13.1$ Hz, 1H), 1.42 (br d, $J=11.9$ Hz, 1H), 1.28 (br d, $J=6.7$ Hz, 6H); MS ESI m/z 530 (M+H); Anal. HPLC Retention time: 1.51 (Method 1); ROCK2 $IC_{50}=2.1$ nM.

5

Example 201: Preparation of 3-{2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-1-oxo-2,8-diazaspiro[4.5]decan-8-carbonyl}-1H-indole-5-carbonitrile




10

^1H NMR (500 MHz, DMSO- d_6) δ 8.15 (s, 1H), 8.02 (br s, 2H), 7.95 (s, 1H), 7.68 - 7.58 (m, 3H), 7.54 (d, $J=7.9$ Hz, 1H), 7.16 (br d, $J=8.2$ Hz, 1H), 4.21 (br d, $J=11.6$ Hz, 2H), 3.94 - 3.82 (m, 5H), 3.44 (br s, 1H), 3.36 - 3.23 (m, 1H), 2.19 (br t, $J=6.9$ Hz, 2H), 1.83 - 1.72 (m, 2H), 1.61 (br d, $J=12.5$ Hz, 2H); MS ESI m/z 495.2 (M+H); Anal. HPLC

15


Retention time: 1.42 (Method 1); ROCK2 $IC_{50}=718$ nM.

Example 202: Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-{4-oxo-3H,4H-pyrrolo[2,1-f][1,2,4]triazine-5-carbonyl}-2,8-diazaspiro[4.5]decan-1-one



¹H NMR (500 MHz, DMSO-d₆) δ 8.02 (br s, 2H), 7.86 (br d, *J*=2.9 Hz, 1H), 7.69 - 7.57 (m, 3H), 7.13 (br d, *J*=8.1 Hz, 1H), 6.59 (br s, 1H), 4.39 (br d, *J*=12.8 Hz, 1H), 3.93 - 3.76 (m, 5H), 3.59 (br d, *J*=13.5 Hz, 1H), 3.22 - 3.04 (m, 2H), 2.22 - 2.01 (m, 2H), 1.82 - 1.66 (m, 2H), 1.62 (br d, *J*=13.4 Hz, 1H), 1.42 (br d, *J*=12.4 Hz, 1H); MS ESI *m/z* 487.9 (M+H); Anal. HPLC Retention time: 1.11 (Method 1); ROCK2 IC₅₀=5.4nM.

Example 203: Preparation of 8-(isoquinoline-8-carbonyl)-2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one




10

¹H NMR (500 MHz, DMSO-d₆) δ 9.26 - 9.08 (m, 1H), 8.57 (br d, *J*=5.2 Hz, 1H), 8.05 (br d, *J*=8.2 Hz, 1H), 8.00 (br s, 2H), 7.93 (d, *J*=5.8 Hz, 1H), 7.84 (br t, *J*=7.6 Hz, 1H), 7.68 - 7.55 (m, 3H), 7.12 (br s, 1H), 4.64 - 4.41 (m, 1H), 3.97 - 3.74 (m, 4H), 3.41 - 3.09 (m, 3H), 2.19 (br d, *J*=5.8 Hz, 1H), 2.15 - 2.05 (m, 1H), 1.86 (s, 2H), 1.80 - 1.71 (m, 1H), 1.42 (br d, *J*=12.5 Hz, 1H); MS ESI *m/z* 482.1 (M+H); Anal. HPLC Retention time: 1.26 (Method 1); ROCK2 IC₅₀=326nM.

15

Example 204: Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-[2-(2-methoxyethyl)-3-oxo-2,3-dihydro-1H-isoindole-4-carbonyl]-2,8-diazaspiro[4.5]decan-1-one




5

¹H NMR (500 MHz, DMSO-d₆) δ 8.13 - 7.88 (m, 2H), 7.66 - 7.54 (m, 4H), 7.38 - 7.25 (m, 1H), 7.17 - 7.06 (m, 1H), 4.55 (br s, 2H), 4.44 - 4.31 (m, 1H), 3.94 - 3.67 (m, 7H), 3.64 - 3.56 (m, 1H), 3.53 - 3.49 (m, 3H), 3.27 (br d, *J*=3.4 Hz, 4H), 3.16 - 3.00 (m, 2H), 2.14 (br dd, *J*=13.0, 6.9 Hz, 1H), 2.07 (br d, *J*=6.4 Hz, 1H), 1.78 (br d, *J*=11.6 Hz, 1H), 1.71 (br s, 1H), 1.64 - 1.54 (m, 1H), 1.50 - 1.30 (m, 1H); MS ESI *m/z* 544 (M+H); Anal. HPLC Retention time: 1.32 (Method 1); ROCK2 IC₅₀=50nM.

10

Example 205: Preparation of tert-butyl 4-(4-[2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-1-oxo-2,8-diazaspiro[4.5]decane-8-carbonyl}phenyl)piperazine-1-carboxylate

15



¹H NMR (500 MHz, DMSO-d₆) δ 8.02 (br d, *J*=2.4 Hz, 2H), 7.60 (br d, *J*=1.2 Hz, 2H), 7.31 (br d, *J*=8.5 Hz, 2H), 7.18 - 7.11 (m, 1H), 6.98 (br d, *J*=8.5 Hz, 2H), 4.01 (br d, *J*=3.1 Hz, 1H), 3.91 - 3.82 (m, 9H), 3.50 - 3.44 (m, 1H), 3.32 - 3.26 (m, 1H), 3.24 - 3.11 (m, 5H), 2.14 (br t, *J*=6.6 Hz, 2H), 1.79 - 1.68 (m, 2H), 1.57 (br d, *J*=11.9 Hz, 2H), 1.42 (s, 9H); MS ESI *m/z* 615.3 (M+H); Anal. HPLC Retention time: 1.76 (Method 1); ROCK2 IC₅₀=8.9nM.

10 Preparation of Examples 206 - 208

The following compounds were made in a parallel manner using the following procedure: Reagents were weighed into stubby tubes. Stock solutions were made for reagent addition : Dissolved 63.0 mg 2-(3-methoxy-4-(1H-pyrazol-4-yl)phenyl)-2,8-diazaspiro[4.5]decan-1-one, 2 HCl in 2.5 ml of THF. To each vial containing sulfonyl chloride reagents was added 0.5 ml of the 2-(3-methoxy-4-(1H-pyrazol-4-yl)phenyl)-2,8-diazaspiro[4.5]decan-1-one, 2 HCl solution followed by DIEA (0.028 ml, 0.150 mmol). The reaction mixture vials were placed in a Bohdan Miniblock XT and were agitated overnight at 400rpm.

Upon completion, reaction mixtures were purified via preparative LC/MS with the following conditions: Column: XBridge C18, 19 x 100 mm, 5-μm particles; Mobile Phase A: 5:95 acetonitrile: water with 10 mM ammonium acetate; Mobile Phase B: 95:5 acetonitrile: water with 10 mM ammonium acetate; Gradient: 10-50% B over 20 minutes,

then a 2-minute hold at 100% B; Flow: 20 mL/min. Fractions containing the desired product were combined and dried via centrifugal evaporation. **Gradient varied for each reaction depending on polarity of compound.


Compound purity was assigned based on the methods below.

5 **Method 1:** Column: Waters Acquity UPLC BEH C18, 2.1 x 50 mm, 1.7- μ m particles; Mobile Phase A: 5:95 acetonitrile:water with 10 mM ammonium acetate; Mobile Phase B: 95:5 acetonitrile:water with 10 mM ammonium acetate; Temperature: 50 °C; Gradient: 0-100% B over 3 minutes, then a 0.75-minute hold at 100% B; Flow: 1.0 mL/min; Detection: UV at 220 nm.

10 **Method 2:** Column: Waters Acquity UPLC BEH C18, 2.1 x 50 mm, 1.7- μ m particles; Mobile Phase A: 5:95 acetonitrile:water with 0.1% trifluoroacetic acid; Mobile Phase B: 95:5 acetonitrile:water with 0.1% trifluoroacetic acid; Temperature: 50 °C; Gradient: 0-100% B over 3 minutes, then a 0.75-minute hold at 100% B; Flow: 1.0 mL/min; Detection: UV at 220 nm.

15


Example 206: Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-(4-methoxybenzenesulfonyl)-2,8-diazaspiro[4.5]decan-1-one



¹H NMR (500 MHz, DMSO-d₆) δ 8.09 (br s, 1H), 7.93 (br s, 1H), 7.71 (d, J =8.8 Hz, 2H),
 20 7.58 (d, J =8.5 Hz, 1H), 7.53 (s, 1H), 7.18 (d, J =8.8 Hz, 2H), 7.12 - 7.06 (m, 1H), 4.22 -
 4.14 (m, 1H), 3.86 (d, J =11.8 Hz, 6H), 3.76 (br t, J =6.9 Hz, 2H), 3.51 (br s, 1H), 3.43 (br
 s, 1H), 3.17 (d, J =5.2 Hz, 1H), 1.88 (br t, J =6.8 Hz, 2H), 1.83 - 1.75 (m, 2H), 1.62 (br d,


$J=13.4$ Hz, 2H); MS ESI m/z 497.2 (M+H); Anal. HPLC Retention time: 1.55 (Method 1); ROCK2 $IC_{50}=497$ nM.

5 Example 207: Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-(3-methoxybenzenesulfonyl)-2,8-diazaspiro[4.5]decan-1-one



^1H NMR (500 MHz, DMSO- d_6) δ 8.08 (br s, 1H), 7.93 (br s, 1H), 7.63 - 7.56 (m, 2H), 7.54 (s, 1H), 7.38 - 7.28 (m, 2H), 7.22 (s, 1H), 7.10 (br d, $J=8.5$ Hz, 1H), 4.68 (t, $J=5.8$ Hz, 1H), 4.17 (q, $J=5.0$ Hz, 1H), 3.85 (d, $J=5.5$ Hz, 6H), 3.76 (br t, $J=6.8$ Hz, 2H), 3.54 (br d, $J=12.3$ Hz, 1H), 3.17 (d, $J=5.3$ Hz, 1H), 1.89 (br t, $J=6.8$ Hz, 2H), 1.84 - 1.73 (m, 2H), 1.63 (br d, $J=13.5$ Hz, 2H); MS ESI m/z 497.3 (M+H); Anal. HPLC Retention time: 1.64 (Method 1); ROCK2 $IC_{50}=266$ nM.

15 Example 208: Preparation of methyl 5-({2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-1-oxo-2,8-diazaspiro[4.5]decan-8-yl}sulfonyl)furan-2-carboxylate



¹H NMR (500 MHz, DMSO-d₆) δ 8.01 (br s, 2H), 7.60 (d, *J*=8.5 Hz, 1H), 7.55 (d, *J*=1.4 Hz, 1H), 7.51 (d, *J*=3.7 Hz, 1H), 7.37 (d, *J*=3.6 Hz, 1H), 7.11 (dd, *J*=8.5, 1.7 Hz, 1H), 3.86 (d, *J*=11.5 Hz, 6H), 3.80 (br t, *J*=6.8 Hz, 2H), 3.65 (br d, *J*=12.7 Hz, 2H), 2.97 (br t, *J*=10.6 Hz, 2H), 2.00 (br t, *J*=6.8 Hz, 2H), 1.81 - 1.70 (m, 2H), 1.63 (br d, *J*=13.7 Hz, 2H); MS ESI *m/z* 515.2 (M+H); Anal. HPLC Retention time: 1.5 (Method 1); ROCK2 IC₅₀=2072nM.

Preparation of Examples 209 - 218

The following compounds were made in a parallel manner using the following procedure: Reagents were weighed into stubby tubes. Stock solutions were made for reagent addition : Dissolved 360 mg 2-(3-methoxy-4-(1H-pyrazol-4-yl)phenyl)-2,8-diazaspiro[4.5]decan-1-one, 2 HCl in 12 ml of DCM. To each vial containing sulfonyl chloride reagents or isocyanide reagents was added 0.5 ml of the 2-(3-methoxy-4-(1H-pyrazol-4-yl)phenyl)-2,8-diazaspiro[4.5]decan-1-one, 2 HCl solution followed by DIEA (0.033 ml, 0.188 mmol). The reaction mixture vials were placed in a Bohdan Miniblock XT and were agitated overnight at 400rpm.


Upon completion, reaction mixtures were purified via preparative LC/MS with the following conditions: Column: XBridge C18, 19 x 100 mm, 5-μm particles; Mobile Phase A: 5:95 acetonitrile: water with 10 mM ammonium acetate; Mobile Phase B: 95:5 acetonitrile: water with 10 mM ammonium acetate; Gradient: 10-50% B over 20 minutes, then a 2-minute hold at 100% B; Flow: 20 mL/min. Fractions containing the desired product were combined and dried via centrifugal evaporation. **Gradient varied for each reaction depending on polarity of compound.

Compound purity was assigned based on the methods below.

Method 1: Column: Waters Acquity UPLC BEH C18, 2.1 x 50 mm, 1.7- μ m particles; Mobile Phase A: 5:95 acetonitrile:water with 10 mM ammonium acetate; Mobile Phase B: 95:5 acetonitrile:water with 10 mM ammonium acetate; Temperature: 50 °C; Gradient: 0-100% B over 3 minutes, then a 0.75-minute hold at 100% B; Flow: 1.0 mL/min; Detection: UV at 220 nm.


Method 2: Column: Waters Acquity UPLC BEH C18, 2.1 x 50 mm, 1.7- μ m particles; Mobile Phase A: 5:95 acetonitrile:water with 0.1% trifluoroacetic acid; Mobile Phase B: 95:5 acetonitrile:water with 0.1% trifluoroacetic acid; Temperature: 50 °C; Gradient: 0-100% B over 3 minutes, then a 0.75-minute hold at 100% B; Flow: 1.0 mL/min; Detection: UV at 220 nm.

Example 209: Preparation of 6-({2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-1-oxo-2,8-diazaspiro[4.5]decan-8-yl}sulfonyl)-1,2,3,4-tetrahydroquinazoline-2,4-dione




¹H NMR (500 MHz, DMSO-*d*₆) δ 8.14 (d, *J*=1.5 Hz, 1H), 8.05 - 7.92 (m, 3H), 7.57 (d, *J*=8.2 Hz, 1H), 7.51 (s, 1H), 7.35 (d, *J*=8.5 Hz, 1H), 7.09 (br d, *J*=8.2 Hz, 1H), 3.83 (s, 3H), 3.75 (br t, *J*=6.7 Hz, 2H), 3.50 (br d, *J*=15.3 Hz, 2H), 2.60 (br t, *J*=11.0 Hz, 2H), 1.89 (br t, *J*=6.6 Hz, 2H), 1.83 - 1.72 (m, 2H), 1.63 (br d, *J*=13.1 Hz, 2H); MS ESI *m/z* 551.2 (M+H); Anal. HPLC Retention time: 1.15 (Method 1); ROCK2 IC₅₀=410nM.

Example 210: Preparation of 5-({2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-1-oxo-2,8-diazaspiro[4.5]decan-8-yl}sulfonyl)-2,3-dihydro-1H-isoindole-1,3-dione



¹H NMR (500 MHz, DMSO-d₆) δ 8.00 (br d, *J*=4.6 Hz, 3H), 7.78 (br d, *J*=11.0 Hz, 2H), 7.61 - 7.50 (m, 2H), 7.10 (br d, *J*=7.6 Hz, 1H), 3.88 - 3.79 (m, 3H), 3.76 (br t, *J*=6.1 Hz, 2H), 3.64 - 3.52 (m, 2H), 2.66 - 2.59 (m, 2H), 1.94 - 1.86 (m, 2H), 1.84 - 1.78 (m, 2H), 1.64 (br d, *J*=13.7 Hz, 2H); MS ESI *m/z* 536.1 (M+H); Anal. HPLC Retention time: 1.37 (Method 1); ROCK2 IC₅₀=2305nM.

Example 211: Preparation of 8-(1,3-benzothiazole-6-sulfonyl)-2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one




10

¹H NMR (500 MHz, DMSO-d₆) δ 9.65 (s, 1H), 8.74 (s, 1H), 8.33 (d, *J*=8.5 Hz, 1H), 7.99 (br s, 2H), 7.90 (br d, *J*=8.5 Hz, 1H), 7.57 (d, *J*=8.5 Hz, 1H), 7.50 (s, 1H), 7.14 - 7.01 (m, 1H), 3.83 (s, 3H), 3.74 (br t, *J*=6.7 Hz, 2H), 3.64 - 3.53 (m, 2H), 2.66 (br t, *J*=10.4 Hz,


2H), 1.87 (br t, $J=6.9$ Hz, 2H), 1.83 - 1.76 (m, 2H), 1.63 (br d, $J=13.4$ Hz, 2H); MS ESI m/z 524 (M+H); Anal. HPLC Retention time: 1.42 (Method 1); ROCK2 $IC_{50}=46$ nM.

Example 212: Preparation of 8-(3-fluoro-4-methoxybenzenesulfonyl)-2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one



^1H NMR (500 MHz, DMSO- d_6) δ 7.99 (br s, 2H), 7.65 - 7.53 (m, 3H), 7.50 (s, 1H), 7.41 (br t, $J=8.7$ Hz, 1H), 7.09 (br d, $J=8.2$ Hz, 1H), 3.94 (s, 3H), 3.83 (s, 3H), 3.75 (br t, $J=6.7$ Hz, 2H), 3.51 (br d, $J=11.9$ Hz, 2H), 2.65 - 2.56 (m, 2H), 1.89 (br t, $J=6.9$ Hz, 2H), 1.78 (br t, $J=10.4$ Hz, 2H), 1.62 (br d, $J=13.4$ Hz, 2H); MS ESI m/z 515 (M+H); Anal. HPLC Retention time: 1.58 (Method 1); ROCK2 $IC_{50}=285$ nM.


Example 213: Preparation of 4-({2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-1-oxo-2,8-diazaspiro[4.5]decan-8-yl}sulfonyl)benzene-1-sulfonamide



^1H NMR (500 MHz, DMSO- d_6) δ 8.07 (br d, $J=8.2$ Hz, 3H), 7.98 (br d, $J=8.5$ Hz, 2H), 7.66 (br s, 1H), 7.57 (d, $J=8.5$ Hz, 1H), 7.50 (s, 1H), 7.09 (br d, $J=8.5$ Hz, 1H), 3.83 (s, 3H), 3.76 (br t, $J=6.6$ Hz, 2H), 3.52 - 3.50 (m, 2H), 2.69 (br t, $J=10.4$ Hz, 2H), 1.90 (br t,


$J=6.7$ Hz, 2H), 1.79 (br d, $J=10.1$ Hz, 2H), 1.64 (br d, $J=13.4$ Hz, 2H); MS ESI m/z 546.2 (M+H); Anal. HPLC Retention time: 1.34 (Method 2); ROCK2 $IC_{50}=456$ nM.

5 Example 214: Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-1-oxo-N-(pyridin-3-yl)-2,8-diazaspiro[4.5]decane-8-carboxamide




^1H NMR (500 MHz, DMSO- d_6) δ 8.76 (s, 1H), 8.62 (br s, 1H), 8.14 (br d, $J=4.0$ Hz, 1H), 8.09 - 7.90 (m, 2H), 7.87 (br d, $J=8.2$ Hz, 1H), 7.64 - 7.54 (m, 2H), 7.28 (br dd, $J=8.2, 4.6$ Hz, 1H), 7.15 (br d, $J=8.2$ Hz, 1H), 4.05 (br d, $J=13.7$ Hz, 2H), 3.86 (s, 5H), 3.08 (br t, $J=12.2$ Hz, 2H), 2.14 (br t, $J=6.7$ Hz, 2H), 1.81 - 1.65 (m, 2H), 1.57 (br d, $J=12.8$ Hz, 2H); MS ESI m/z 447 (M+H); Anal. HPLC Retention time: 0.86 (Method 1); ROCK2 $IC_{50}=1483$ nM.

15 Example 215: Preparation of N-[4-(dimethylamino)phenyl]-2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-1-oxo-2,8-diazaspiro[4.5]decane-8-carboxamide



¹H NMR (500 MHz, DMSO-d₆) δ 8.23 (s, 1H), 8.01 (br s, 2H), 7.66 - 7.53 (m, 2H), 7.23 (br d, *J*=8.9 Hz, 2H), 7.15 (br d, *J*=8.2 Hz, 1H), 6.66 (br d, *J*=8.5 Hz, 2H), 4.02 (br d, *J*=13.7 Hz, 2H), 3.91 - 3.81 (m, 5H), 3.00 (br d, *J*=9.8 Hz, 2H), 2.82 (s, 6H), 2.13 (br t, *J*=6.6 Hz, 2H), 1.74 - 1.66 (m, 2H), 1.53 (br d, *J*=12.8 Hz, 2H); MS ESI *m/z* 489.2 (M+H); Anal. HPLC Retention time: 1.29 (Method 1); ROCK2 IC₅₀=791nM.

Example 216: Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-N-(3-methoxyphenyl)-1-oxo-2,8-diazaspiro[4.5]decane-8-carboxamide




10

¹H NMR (500 MHz, DMSO-d₆) δ 8.52 (s, 1H), 8.02 (br s, 2H), 7.70 - 7.55 (m, 2H), 7.21 - 7.03 (m, 4H), 6.51 (br d, *J*=7.0 Hz, 1H), 4.05 (br d, *J*=13.4 Hz, 3H), 3.88 - 3.83 (m, 5H),


3.71 (s, 3H), 3.09 - 3.03 (m, 2H), 2.14 (br t, $J=6.9$ Hz, 2H), 1.72 (br d, $J=11.3$ Hz, 2H), 1.56 (br d, $J=12.5$ Hz, 2H); MS ESI m/z 476.1 (M+H); Anal. HPLC Retention time: 1.33 (Method 1); ROCK2 $IC_{50}=2252$ nM.

5 **Example 217: Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-N-(4-methoxyphenyl)-1-oxo-2,8-diazaspiro[4.5]decane-8-carboxamide**



^1H NMR (500 MHz, DMSO- d_6) δ 8.37 (s, 1H), 7.68 - 7.56 (m, 2H), 7.35 (br d, $J=8.5$ Hz, 2H), 7.15 (br d, $J=7.6$ Hz, 1H), 6.82 (br d, $J=8.9$ Hz, 2H), 4.04 (br d, $J=13.4$ Hz, 2H),
 10 3.94 - 3.81 (m, 5H), 3.74 - 3.68 (m, 3H), 3.03 (br t, $J=11.9$ Hz, 2H), 2.13 (br t, $J=6.7$ Hz, 2H), 1.72 (br t, $J=10.7$ Hz, 2H), 1.54 (br d, $J=13.1$ Hz, 2H); MS ESI m/z 476.11, 476.11 (M+H); ROCK2 $IC_{50}=2460$ nM.

15 **Example 218: Preparation of 8-(4-methanesulfonylbenzenesulfonyl)-2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one**



¹H NMR (500 MHz, DMSO-d₆) δ 8.20 (br d, *J*=8.2 Hz, 2H), 8.05 (br d, *J*=8.2 Hz, 4H), 7.58 (br d, *J*=8.5 Hz, 1H), 7.51 (s, 1H), 7.09 (br d, *J*=8.2 Hz, 1H), 3.83 (s, 3H), 3.76 (br t, *J*=6.6 Hz, 2H), 3.31 (s, 3H), 3.17 (d, *J*=4.9 Hz, 2H), 2.71 (br t, *J*=10.8 Hz, 2H), 1.92 (br t, *J*=6.7 Hz, 2H), 1.79 (br t, *J*=10.2 Hz, 2H), 1.63 (br d, *J*=13.1 Hz, 2H); MS ESI *m/z* 545 (M+H); Anal. HPLC Retention time: 1.42 (Method 1); ROCK2 IC₅₀=843nM.

Preparation of Examples 219 - 246

The following compounds were made in a parallel manner using the following procedure: Reagents were weighed into stubby tubes. Stock solutions were made for reagent addition : Dissolved 510 mg 2-(3-methoxy-4-(1H-pyrazol-4-yl)phenyl)-8-(4-(piperazin-1-yl)benzoyl)-2,8-diazaspiro[4.5]decan-1-one, HCl in 14.50 ml of DCE. To each stubby tube containing aldehyde/ketone reagent (0.072 mmol) was added 0.25 ml of the 2-(3-methoxy-4-(1H-pyrazol-4-yl)phenyl)-8-(4-(piperazin-1-yl)benzoyl)-2,8-diazaspiro[4.5]decan-1-one, HCl solution and trimethyl orthoformate (0.020 ml, 0.180 mmol) followed by acetic acid (1.039 ul, 0.018 mmol). The reaction mixture vials were placed in a Bohdan Miniblock XT and were agitated at 400rpm on an Innova platform shaker at rt for 1 hr. To each vial was then added MP-cyanoborohydride (30.1 mg, 0.072 mmol) and continued agitaion overnight.


Upon completion, reaction mixtures were purified via preparative LC/MS with the following conditions: Column: XBridge C18, 19 x 100 mm, 5-μm particles; Mobile Phase A: 5:95 acetonitrile: water with 10 mM ammonium acetate; Mobile Phase B: 95:5 acetonitrile: water with 10 mM ammonium acetate; Gradient: 10-50% B over 20 minutes, then a 2-minute hold at 100% B; Flow: 20 mL/min. Fractions containing the desired product were combined and dried via centrifugal evaporation. **Gradient varied for each reaction depending on polarity of compound.

Compound purity was assigned based on the methods below.

Method 1: Column: Waters Acquity UPLC BEH C18, 2.1 x 50 mm, 1.7-μm particles; Mobile Phase A: 5:95 acetonitrile:water with 10 mM ammonium acetate; Mobile Phase B: 95:5 acetonitrile:water with 10 mM ammonium acetate; Temperature: 50 °C; Gradient: 0-100% B over 3 minutes, then a 0.75-minute hold at 100% B; Flow: 1.0 mL/min; Detection: UV at 220 nm.


Method 2: Column: Waters Acquity UPLC BEH C18, 2.1 x 50 mm, 1.7- μ m particles; Mobile Phase A: 5:95 acetonitrile:water with 0.1% trifluoroacetic acid; Mobile Phase B: 95:5 acetonitrile:water with 0.1% trifluoroacetic acid; Temperature: 50 °C; Gradient: 0-100% B over 3 minutes, then a 0.75-minute hold at 100% B; Flow: 1.0 mL/min; Detection: UV at 220 nm.

Example 219: Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-(4-{4-[(quinoxalin-2-yl)methyl]piperazin-1-yl}benzoyl)-2,8-diazaspiro[4.5]decan-1-one




¹H NMR (500 MHz, DMSO-*d*₆) δ 9.05 (s, 1H), 8.41 - 8.26 (m, 1H), 8.23 - 8.11 (m, 2H), 7.97 - 7.89 (m, 2H), 7.65 - 7.54 (m, 2H), 7.36 - 7.26 (m, 3H), 7.13 (br d, *J*=8.7 Hz, 1H), 7.03 (br d, *J*=8.2 Hz, 2H), 4.63 (br s, 2H), 3.85 (s, 5H), 3.74 (br s, 6H), 3.33 (br d, *J*=6.6 Hz, 2H), 3.25 - 3.20 (m, 1H), 2.97 - 2.86 (m, 3H), 2.14 (br s, 2H), 1.70 (br s, 2H), 1.61 - 1.47 (m, 2H); MS ESI *m/z* 657.3 (M+H); Anal. HPLC Retention time: 1.62 (Method 2); ROCK2 IC₅₀=1.9 nM.

Example 220: Preparation of 8-[4-(4-{4-(1H-imidazol-1-yl)phenyl}methyl)piperazin-1-yl}benzoyl]-2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one




¹H NMR (500 MHz, DMSO-d₆) δ 8.17 (br s, 2H), 8.02 (s, 2H), 7.91 (br d, *J*=8.1 Hz, 2H), 7.76 (br d, *J*=8.2 Hz, 2H), 7.63 - 7.58 (m, 2H), 7.35 (br d, *J*=8.4 Hz, 2H), 7.19 - 7.12 (m, 2H), 7.08 - 7.01 (m, 2H), 4.47 (br s, 2H), 3.86 (s, 6H), 3.59 - 3.47 (m, 1H), 3.36 - 3.25 (m, 1H), 3.21 - 3.12 (m, 1H), 2.51 (br s, 6H), 2.15 (br s, 2H), 1.71 (br d, *J*=9.5 Hz, 2H), 1.58 (br s, 2H); MS ESI *m/z* 671.5 (M+H); Anal. HPLC Retention time: 0.93 (Method 1); ROCK2 IC₅₀=0.3 nM.

Example 221: Preparation of 8-(4-{4-[2-(benzyloxy)ethyl]piperazin-1-yl}benzoyl)-2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one




¹H NMR (500 MHz, DMSO-d₆) δ 8.02 (br s, 2H), 7.64 - 7.58 (m, 2H), 7.39 - 7.31 (m, 8H), 7.15 (br d, *J*=8.4 Hz, 1H), 7.05 (br d, *J*=8.4 Hz, 1H), 4.72 (br d, *J*=4.5 Hz, 1H), 4.57 (br d, *J*=5.6 Hz, 3H), 3.86 (s, 4H), 3.78 (br s, 2H), 3.60 (br d, *J*=5.6 Hz, 2H), 3.57 - 3.46 (m, 8H), 3.39 (br s, 1H), 3.33 (br d, *J*=7.0 Hz, 1H), 2.15 (br s, 2H), 1.70 (br d, *J*=9.2 Hz, 2H), 1.58 (br s, 2H); MS ESI *m/z* 649.2 (M+H); Anal. HPLC Retention time: 1.68 (Method 2); ROCK2 IC₅₀=0.4 nM.

Example 222: Preparation of 8-{4-[4-(2,2-diphenylethyl)piperazin-1-yl]benzoyl}-2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one




¹H NMR (500 MHz, DMSO-d₆) δ 8.01 (br s, 2H), 7.66 - 7.56 (m, 2H), 7.39 - 7.24 (m, 15H), 7.21 - 7.09 (m, 4H), 6.92 (br d, *J*=8.5 Hz, 2H), 4.30 (br s, 2H), 3.86 (s, 6H), 3.48 (br s, 10H), 3.25 (s, 1H), 3.20 - 3.16 (m, 1H), 3.11 (br d, *J*=4.6 Hz, 4H), 3.00 (br d, *J*=7.6 Hz, 2H), 2.58 (br s, 4H), 2.14 (br s, 2H), 1.73 - 1.65 (m, 2H), 1.56 (br d, *J*=12.5 Hz, 2H); MS ESI *m/z* 695.3 (M+H); Anal. HPLC Retention time: 1.51 (Method 1); ROCK2 IC₅₀=9.3 nM.

10 Example 223: Preparation of 8-(4-{4-[(2-chlorophenyl)methyl]piperazin-1-yl}benzoyl)-2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one



¹H NMR (500 MHz, DMSO-d₆) δ 8.08 - 7.88 (m, 2H), 7.65 - 7.56 (m, 2H), 7.53 (br d, *J*=7.4 Hz, 1H), 7.44 (br d, *J*=7.7 Hz, 1H), 7.38 - 7.26 (m, 4H), 7.14 (br d, *J*=8.5 Hz, 1H), 6.96 (br d, *J*=8.6 Hz, 2H), 3.92 - 3.80 (m, 5H), 3.69 - 3.63 (m, 4H), 3.23 (br s, 4H), 2.57 (br s, 4H), 2.14 (br t, *J*=6.3 Hz, 2H), 1.81 - 1.66 (m, 4H), 1.58 (br s, 2H); MS ESI *m/z* 639.1 (M+H); Anal. HPLC Retention time: 1.28 (Method 1); ROCK2 IC₅₀=1.1nM.


Example 224: Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-(4-{4-[(1,2,3-thiadiazol-4-yl)methyl]piperazin-1-yl}benzoyl)-2,8-diazaspiro[4.5]decan-1-one



¹H NMR (500 MHz, DMSO-d₆) δ 9.10 (s, 1H), 8.02 (br d, *J*=3.4 Hz, 2H), 7.64 - 7.54 (m, 2H), 7.30 (br d, *J*=8.5 Hz, 2H), 7.16 (br d, *J*=7.3 Hz, 1H), 6.96 (br d, *J*=8.5 Hz, 2H), 4.11 (s, 3H), 3.93 - 3.81 (m, 5H), 3.57 (br d, *J*=6.7 Hz, 1H), 3.38 (s, 1H), 3.24 (br s, 5H), 2.61 (br s, 4H), 2.15 (br t, *J*=6.6 Hz, 2H), 1.72 (br d, *J*=9.5 Hz, 2H), 1.57 (br d, *J*=11.3 Hz, 2H); MS ESI *m/z* 612.9 (M+H);

Anal. HPLC Retention time: 1.37 (Method 2); ROCK2 IC₅₀=1.5nM.

Example 225: Preparation of 8-[4-[4-([1,1'-biphenyl]-4-yl)methyl]piperazin-1-yl]benzoyl]-2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one




10

¹H NMR (500 MHz, DMSO-d₆) δ 8.02 (br s, 2H), 7.80 (br d, *J*=7.8 Hz, 2H), 7.72 (br d, *J*=7.6 Hz, 2H), 7.67 - 7.57 (m, 4H), 7.50 (t, *J*=7.6 Hz, 2H), 7.45 - 7.38 (m, 1H), 7.34 (br d, *J*=8.3 Hz, 2H), 7.18 - 7.12 (m, 1H), 7.10 - 6.99 (m, 2H), 4.43 (br s, 2H), 3.86 (s, 5H), 3.63 - 3.42 (m, 6H), 3.38 - 3.29 (m, 1H), 3.26 (br s, 1H), 2.59 - 2.55 (m, 4H), 2.14 (br s, 2H), 1.70 (br d, *J*=10.5 Hz, 2H), 1.58 (br s, 2H); MS ESI *m/z* 681.3 (M+H); Anal. HPLC Retention time: 1.52 (Method 1); ROCK2 IC₅₀=4.5nM.


15

Example 226: Preparation of 8-[4-(4-cyclohexylpiperazin-1-yl)benzoyl]-2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one




¹H NMR (500 MHz, DMSO-d₆) δ 8.01 (br s, 2H), 7.65 - 7.54 (m, 2H), 7.34 (br d, *J*=8.5 Hz, 2H), 7.14 (br d, *J*=8.5 Hz, 1H), 7.04 (br d, *J*=8.5 Hz, 2H), 3.86 (s, 5H), 3.69 - 3.60 (m, 13H), 2.14 (br s, 2H), 2.09 - 2.03 (m, 2H), 1.83 (br d, *J*=12.5 Hz, 2H), 1.70 (br d, *J*=11.0 Hz, 2H), 1.60 (br s, 3H), 1.46 - 1.33 (m, 2H), 1.27 (br d, *J*=12.8 Hz, 2H), 1.18 - 1.08 (m, 1H); MS ESI *m/z* 597.2 (M+H); Anal. HPLC Retention time: 1.17 (Method 1); ROCK2 IC₅₀=0.7nM.

Example 227: Preparation of 8-(4-{4-[(4-chlorophenyl)methyl]piperazin-1-yl}benzoyl)-2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one




¹H NMR (500 MHz, DMSO-d₆) δ 8.01 (br s, 2H), 7.65 - 7.55 (m, 2H), 7.42 - 7.34 (m, 4H), 7.28 (br d, *J*=8.2 Hz, 2H), 7.14 (br d, *J*=7.6 Hz, 1H), 6.95 (br d, *J*=8.5 Hz, 2H), 3.85 (s, 5H), 3.67 - 3.63 (m, 8H), 3.55 - 3.50 (m, 1H), 3.29 - 3.11 (m, 7H), 2.14 (br t, *J*=6.3 Hz, 2H), 1.70 (br s, 2H), 1.56 (br d, *J*=12.8 Hz, 2H); MS ESI *m/z* 639.1 (M+H); Anal. HPLC Retention time: 1.96 (Method 2); ROCK2 IC₅₀=2.2nM.

Example 228: Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-(4-{4-[(1-methyl-1H-pyrrol-2-yl)methyl]piperazin-1-yl}benzoyl)-2,8-diazaspiro[4.5]decan-1-one



¹H NMR (500 MHz, DMSO-d₆) δ 8.02 (s, 2H), 7.66 - 7.57 (m, 2H), 7.29 (br d, *J*=8.5 Hz, 2H), 7.16 (br d, *J*=8.2 Hz, 1H), 6.96 (br d, *J*=8.5 Hz, 2H), 6.68 (br s, 1H), 5.90 (br d, *J*=9.5 Hz, 2H), 3.91 (s, 2H), 3.87 (s, 5H), 3.61 (s, 3H), 3.45 (br s, 6H), 3.24 - 3.18 (m, 6H), 2.15 (br s, 2H), 1.72 (br d, *J*=9.8 Hz, 2H), 1.57 (br d, *J*=11.9 Hz, 2H); MS ESI *m/z* 608.2 (M+H); Anal. HPLC Retention time: 1.66 (Method 2); ROCK2 IC₅₀=0.9nM.


Example 229: Preparation of 8-(4-{4-[(1,3-benzothiazol-2-yl)methyl]piperazin-1-yl}benzoyl)-2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one



¹H NMR (500 MHz, DMSO-d₆) δ 8.06 (br d, *J*=7.6 Hz, 1H), 8.01 (br s, 2H), 7.94 (br d, *J*=8.2 Hz, 1H), 7.65 - 7.57 (m, 2H), 7.49 (br d, *J*=7.6 Hz, 1H), 7.45 - 7.39 (m, 1H), 7.30 (br d, *J*=8.5 Hz, 2H), 7.14 (br d, *J*=7.9 Hz, 1H), 6.98 (br d, *J*=8.5 Hz, 2H), 4.02 (s, 2H),
 5 3.86 (s, 5H), 3.49 - 3.44 (m, 2H), 3.28 (br s, 4H), 3.17 (br s, 2H), 2.72 (br s, 4H), 2.14 (br t, *J*=6.3 Hz, 2H), 1.76 - 1.69 (m, 2H), 1.57 (br d, *J*=12.8 Hz, 2H); MS ESI *m/z* 662.1 (M+H); Anal. HPLC Retention time: 1.36 (Method 1); ROCK2 IC₅₀=1.9nM.


Example 230: Preparation of 8-[4-[4-(cyclopentylmethyl)piperazin-1-yl]benzoyl]-2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one

10




¹H NMR (500 MHz, DMSO-d₆) δ 8.24 (s, 1H), 7.98 (s, 1H), 7.68 - 7.59 (m, 2H), 7.34 (br d, *J*=8.4 Hz, 2H), 7.16 (br d, *J*=8.8 Hz, 1H), 7.04 (br d, *J*=8.3 Hz, 2H), 3.89 - 3.84 (m, 5H), 3.21 - 3.09 (m, 6H), 2.31 - 2.20 (m, 2H), 2.15 (br s, 2H), 1.83 (br d, *J*=6.1 Hz, 3H),
 5 1.70 (br d, *J*=9.5 Hz, 2H), 1.65 - 1.44 (m, 12H), 1.19 - 1.13 (m, 1H), 1.09 - 0.98 (m, 1H);
 MS ESI *m/z* 597.1 (M+H); Anal. HPLC Retention time: 2.56 (Method 2); ROCK2
 IC₅₀=2.2nM.

Example 231: Preparation of 8-{4-[4-(3,3-dimethylbutyl)piperazin-1-yl]benzoyl}-2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one



- ¹H NMR (500 MHz, DMSO-d₆) δ 8.02 (br s, 2H), 7.65 - 7.58 (m, 2H), 7.30 (br d, *J*=8.2 Hz, 2H), 7.16 (br d, *J*=10.4 Hz, 1H), 6.96 (br d, *J*=8.5 Hz, 2H), 3.87 (s, 5H), 3.35 (br s, 1H), 3.19 (br d, *J*=15.3 Hz, 5H), 2.35 - 2.26 (m, 2H), 2.15 (br s, 2H), 1.70 (br s, 4H), 1.58 (br d, *J*=9.8 Hz, 2H), 1.44 - 1.35 (m, 2H), 0.94 - 0.85 (m, 9H); MS ESI *m/z* 599.2 (M+H);
- 5 Anal. HPLC Retention time: 1.27 (Method 1); ROCK2 IC₅₀=2.2nM.


Example 232: Preparation of 8-(4-{4-[1-(dimethylamino)propan-2-yl]piperazin-1-yl}benzoyl)-2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one



- 10 ¹H NMR (500 MHz, DMSO-d₆) δ 8.02 (br s, 2H), 7.66 - 7.58 (m, 2H), 7.34 - 7.27 (m, 3H), 7.19 (s, 1H), 6.98 (br d, *J*=8.5 Hz, 1H), 3.87 (s, 5H), 3.33 - 3.11 (m, 4H), 2.93 (br s, 2H), 2.81 (s, 6H), 2.15 (br t, *J*=6.4 Hz, 2H), 1.71 (br d, *J*=11.8 Hz, 2H), 1.57 (br d, *J*=10.4 Hz, 2H), 1.24 (s, 1H), 1.17 (br d, *J*=6.6 Hz, 1H), 0.93 (br d, *J*=5.5 Hz, 3H); Anal. HPLC Retention time: 1.18 (Method 2); MS ESI *m/z* 600.4 (M+H); ROCK2 IC₅₀=3.1 nM.


15

Example 233: Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-{4-[4-(propan-2-yl)piperazin-1-yl]benzoyl}-2,8-diazaspiro[4.5]decan-1-one



¹H NMR (500 MHz, DMSO-d₆) δ 8.01 (br s, 2H), 7.66 - 7.53 (m, 2H), 7.29 (br d, *J*=8.4 Hz, 2H), 7.14 (br d, *J*=8.3 Hz, 1H), 6.95 (br d, *J*=8.5 Hz, 2H), 3.93 - 3.79 (m, 5H), 3.26 - 3.12 (m, 5H), 2.65 (br d, *J*=5.9 Hz, 1H), 2.55 (s, 5H), 2.19 - 2.10 (m, 2H), 1.81 - 1.65 (m, 4H), 1.56 (br d, *J*=11.4 Hz, 2H), 1.00 (d, *J*=6.4 Hz, 6H); MS ESI *m/z* 557.4 (*M*+*H*); Anal. HPLC Retention time: 1.05 (Method 1); ROCK2 IC₅₀=0.2nM.


Example 234: Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-[4-(4-{4-(trifluoromethyl)phenyl}methyl)piperazin-1-yl]benzoyl]-2,8-diazaspiro[4.5]decan-1-one



¹H NMR (500 MHz, DMSO-d₆) δ 8.03 (br s, 2H), 7.88 (br d, *J*=7.9 Hz, 2H), 7.77 (br d, *J*=7.6 Hz, 2H), 7.64 - 7.55 (m, 2H), 7.34 (br d, *J*=8.2 Hz, 2H), 7.19 - 7.13 (m, 2H), 7.03 (br d, *J*=8.5 Hz, 1H), 4.49 - 3.24 (m, 19H), 2.15 (br s, 2H), 1.72 (br s, 2H), 1.63 - 1.49 (m, 2H); MS ESI *m/z* 673.2 (M+H); Anal. HPLC Retention time: 1.39 (Method 1); ROCK2

5 IC₅₀=4.1 nM.

Example 235: Preparation of N-(3-{[4-(4-{2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-1-oxo-2,8-diazaspiro[4.5]decane-8-carbonyl}phenyl)piperazin-1-yl]methyl}phenyl)methanesulfonamide




10

¹H NMR (500 MHz, DMSO-d₆) δ 9.98 (br s, 1H), 8.09 - 7.88 (m, 2H), 7.65 - 7.55 (m, 2H), 7.49 - 7.42 (m, 1H), 7.38 (br s, 1H), 7.34 (br d, *J*=8.2 Hz, 1H), 7.29 - 7.24 (m, 2H), 7.20 - 7.11 (m, 2H), 7.09 - 7.00 (m, 2H), 4.36 (br s, 2H), 3.86 (s, 5H), 3.60 (br s, 12H), 3.04 (s, 3H), 2.14 (br s, 2H), 1.71 (br s, 2H), 1.57 (br d, *J*=6.9 Hz, 2H); MS ESI *m/z*


15 698.2 (M+H); Anal. HPLC Retention time: 1.13 (Method 1); ROCK2 IC₅₀=0.2 nM.

Example 236: Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-(4-{4-[(1,3-thiazol-2-yl)methyl]piperazin-1-yl}benzoyl)-2,8-diazaspiro[4.5]decan-1-one



¹H NMR (500 MHz, DMSO-d₆) δ 8.02 (br s, 2H), 7.74 (br d, *J*=3.0 Hz, 1H), 7.68 (br s, 1H), 7.63 - 7.57 (m, 2H), 7.29 (br d, *J*=8.3 Hz, 2H), 7.14 (br d, *J*=8.2 Hz, 1H), 6.97 (br d, *J*=8.4 Hz, 2H), 3.95 - 3.79 (m, 7H), 3.52 (br s, 2H), 3.36 - 3.11 (m, 6H), 2.64 (br s, 4H), 2.14 (br s, 2H), 1.70 (br d, *J*=10.3 Hz, 2H), 1.57 (br d, *J*=8.5 Hz, 2H); MS ESI *m/z* 612.2 (M+H); Anal. HPLC Retention time: 1.16 (Method 1); ROCK2 IC₅₀=2.7 nM.


Example 237: Preparation of 8-{4-[4-({4-[3-(dimethylamino)propoxy]phenyl}methyl)piperazin-1-yl]benzoyl}-2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one



¹H NMR (500 MHz, DMSO-d₆) δ 8.01 (br s, 2H), 7.62 - 7.55 (m, 2H), 7.28 (br d, *J*=8.2 Hz, 2H), 7.22 (br d, *J*=8.2 Hz, 2H), 7.14 (br d, *J*=7.6 Hz, 1H), 6.94 (br d, *J*=8.5 Hz, 2H), 6.87 (br d, *J*=8.2 Hz, 2H), 4.00 - 3.91 (m, 2H), 3.89 - 3.79 (m, 5H), 3.58 - 3.33 (m, 6H),
 5 3.25 - 3.16 (m, 4H), 2.49 - 2.42 (m, 6H), 2.21 (s, 6H), 2.13 (br t, *J*=6.4 Hz, 2H), 1.88 - 1.80 (m, 2H), 1.70 (br d, *J*=10.7 Hz, 2H), 1.56 (br d, *J*=11.3 Hz, 2H); MS ESI *m/z* 706.3 (M+H); Anal. HPLC Retention time: 0.92 (Method 1); ROCK2 IC₅₀=0.1 nM.


Example 238: Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-{4-[4-(3-methylbutyl)piperazin-1-yl]benzoyl}-2,8-diazaspiro[4.5]decan-1-one

10




¹H NMR (500 MHz, DMSO-d₆) δ 8.01 (br s, 2H), 7.65 - 7.56 (m, 2H), 7.29 (br d, *J*=8.5 Hz, 2H), 7.15 (br d, *J*=8.5 Hz, 1H), 6.95 (br d, *J*=8.5 Hz, 2H), 3.93 - 3.80 (m, 5H), 3.60 (br s, 1H), 3.57 - 3.52 (m, 1H), 3.35 (br s, 1H), 3.29 - 3.09 (m, 5H), 2.32 (br t, *J*=7.5 Hz, 2H), 2.14 (br s, 2H), 2.05 (br d, *J*=4.3 Hz, 1H), 1.70 (br s, 1H), 1.63 - 1.52 (m, 3H), 1.35 (br d, *J*=7.6 Hz, 2H), 1.00 (d, *J*=6.1 Hz, 2H), 0.89 (br d, *J*=6.7 Hz, 8H); MS ESI *m/z* 585.2 (M+H); Anal. HPLC Retention time: 1.53 (Method 2); ROCK2 IC₅₀=1.6 nM.

Example 239: Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-{4-[4-(2-phenylethyl)piperazin-1-yl]benzoyl}-2,8-diazaspiro[4.5]decan-1-one



¹H NMR (500 MHz, DMSO-d₆) δ 8.01 (br s, 2H), 7.65 - 7.55 (m, 2H), 7.35 (br d, *J*=7.9 Hz, 4H), 7.29 (br d, *J*=7.9 Hz, 3H), 7.14 (br d, *J*=8.9 Hz, 1H), 7.06 (br d, *J*=8.5 Hz, 2H), 3.86 (s, 6H), 3.46 (br s, 5H), 3.40 - 3.31 (m, 1H), 3.23 (br d, *J*=13.4 Hz, 1H), 3.06 - 2.98 (m, 2H), 2.50 (br s, 6H), 2.14 (br s, 2H), 1.71 (br d, *J*=9.8 Hz, 2H), 1.57 (br d, *J*=10.1 Hz, 2H); MS ESI *m/z* 619.2 (M+H); Anal. HPLC Retention time: 1.79 (Method 2); ROCK2 IC₅₀=0.6 nM.

Example 240: Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-[4-(3,3,3-trifluoropropyl)piperazin-1-yl]benzoyl}-2,8-diazaspiro[4.5]decan-1-one




10

¹H NMR (500 MHz, DMSO-d₆) δ 8.01 (br s, 2H), 7.64 - 7.57 (m, 2H), 7.30 (br d, *J*=8.2 Hz, 2H), 7.15 (br d, *J*=8.5 Hz, 1H), 6.97 (br d, *J*=8.5 Hz, 2H), 3.92 - 3.80 (m, 5H), 3.55 - 3.47 (m, 4H), 3.27 - 3.13 (m, 6H), 2.51 - 2.44 (m, 4H), 2.14 (br t, *J*=6.4 Hz, 2H), 1.80 - 1.64 (m, 4H), 1.57 (br d, *J*=11.6 Hz, 2H); MS ESI *m/z* 611.1 (M+H); Anal. HPLC Retention time: 1.12 (Method 1); ROCK2 IC₅₀=1.0 nM.


15

Example 241: Preparation of 8-[4-[4-(2,2-dimethylpropyl)piperazin-1-yl]benzoyl]-2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one



¹H NMR (500 MHz, DMSO-d₆) δ 8.01 (br s, 2H), 7.64 - 7.55 (m, 2H), 7.28 (br d, *J*=8.4 Hz, 2H), 7.14 (br d, *J*=8.2 Hz, 1H), 6.94 (br d, *J*=8.5 Hz, 2H), 3.86 (s, 5H), 3.55 (br s, 2H), 3.24 - 3.13 (m, 4H), 2.63 - 2.53 (m, 6H), 2.19 - 2.11 (m, 2H), 2.09 (s, 2H), 1.73 - 1.66 (m, 2H), 1.57 (br s, 2H), 0.87 (s, 9H); MS ESI *m/z* 293.2 (M+H); Anal. HPLC Retention time: 2.18 (Method 2); ROCK2 IC₅₀=0.4 nM.

Example 242: Preparation of 8-{4-[4-(1-hydroxypropan-2-yl)piperazin-1-yl]benzoyl}-2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one




10

¹H NMR (500 MHz, DMSO-d₆) δ 8.01 (br s, 2H), 7.65 - 7.57 (m, 2H), 7.29 (br d, *J*=8.2 Hz, 2H), 7.15 (br d, *J*=7.6 Hz, 1H), 6.95 (br d, *J*=8.2 Hz, 2H), 3.87 (s, 5H), 3.49 (br s,

1H), 3.43 (br d, $J=8.9$ Hz, 1H), 3.32 (br d, $J=4.0$ Hz, 1H), 3.20 (br s, 5H), 3.00 (s, 1H), 2.72 - 2.58 (m, 4H), 2.45 (br s, 1H), 2.22 (s, 1H), 2.15 (br t, $J=6.4$ Hz, 2H), 1.78 - 1.67 (m, 2H), 1.57 (br d, $J=11.3$ Hz, 2H), 0.96 (br d, $J=6.7$ Hz, 3H); MS ESI m/z 573.3 (M+H); Anal. HPLC Retention time: 1.09 (Method 2); ROCK2 $IC_{50}=0.7$ nM.


5

Example 243: Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-(4-{4-[(pyridin-3-yl)methyl]piperazin-1-yl}benzoyl)-2,8-diazaspiro[4.5]decan-1-one




^1H NMR (500 MHz, DMSO- d_6) δ 8.51 (s, 1H), 8.47 (br d, $J=4.3$ Hz, 1H), 8.00 (s, 2H),
 10 7.75 (br d, $J=7.6$ Hz, 1H), 7.62 - 7.55 (m, 2H), 7.38 (dd, $J=7.3, 4.9$ Hz, 1H), 7.28 (br d, $J=8.5$ Hz, 2H), 7.13 (br d, $J=9.5$ Hz, 1H), 6.95 (br d, $J=8.9$ Hz, 2H), 4.02 (br dd, $J=11.0, 4.0$ Hz, 1H), 3.91 - 3.78 (m, 5H), 3.57 (br d, $J=10.4$ Hz, 4H), 3.34 (br s, 1H), 3.29 - 3.11 (m, 6H), 2.27 (br t, $J=7.5$ Hz, 1H), 2.13 (br t, $J=6.4$ Hz, 2H), 1.69 (br d, $J=9.5$ Hz, 2H), 1.56 (br d, $J=10.4$ Hz, 2H); MS ESI m/z 606 (M+H); Anal. HPLC Retention time: 1.44
 15 (Method 2); ROCK2 $IC_{50}=1.0$ nM.

Example 244: Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-{4-[4-(1-methylpiperidin-4-yl)piperazin-1-yl]benzoyl}-2,8-diazaspiro[4.5]decan-1-one




¹H NMR (500 MHz, DMSO-d₆) δ 8.02 (s, 2H), 7.64 - 7.55 (m, 2H), 7.35 (br d, *J*=8.4 Hz, 2H), 7.14 (br d, *J*=8.5 Hz, 1H), 7.06 (br d, *J*=8.4 Hz, 2H), 3.86 (s, 5H), 3.62 (br s, 6H), 3.46 - 3.40 (m, 1H), 3.34 (br d, *J*=6.3 Hz, 1H), 3.25 - 3.09 (m, 2H), 3.06 - 2.93 (m, 2H), 2.78 (br s, 3H), 2.51 - 2.47 (m, 5H), 2.35 (br d, *J*=12.2 Hz, 2H), 2.14 (br s, 2H), 1.96 - 1.82 (m, 2H), 1.71 (br t, *J*=9.5 Hz, 2H), 1.58 (br s, 2H); MS ESI *m/z* 612.1 (M+H); Anal. HPLC Retention time: 1.06 Method 1; ROCK2 IC₅₀=0.9 nM.

Example 245: Preparation of 8-[4-(4-{[4-(dimethylamino)phenyl]methyl}piperazin-1-yl)benzoyl]-2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-2,8-diazaspiro[4.5]decan-1-one



¹H NMR (500 MHz, DMSO-d₆) δ 8.02 (br s, 2H), 7.60 (s, 2H), 7.28 (br d, *J*=8.3 Hz, 2H), 7.12 (br d, *J*=8.5 Hz, 3H), 6.94 (br d, *J*=8.4 Hz, 2H), 6.69 (br d, *J*=8.2 Hz, 2H), 3.86 (s, 5H), 3.59 - 3.54 (m, 5H), 3.25 - 3.11 (m, 5H), 2.87 (s, 6H), 2.46 (br s, 4H), 2.14 (br s, 2H), 1.70 (br d, *J*=9.3 Hz, 2H), 1.56 (br d, *J*=10.9 Hz, 2H); MS ESI *m/z* 648.2 (M+H); Anal. HPLC Retention time: 1.75 (Method 2); ROCK2 IC₅₀=0.1 nM.


Example 246: Preparation of 2-[3-methoxy-4-(1H-pyrazol-4-yl)phenyl]-8-(4-{4-[(oxolan-3-yl)methyl]piperazin-1-yl}benzoyl)-2,8-diazaspiro[4.5]decan-1-one



¹H NMR (500 MHz, DMSO-d₆) δ 8.02 (br s, 2H), 7.66 - 7.55 (m, 2H), 7.29 (br d, *J*=8.5 Hz, 2H), 7.14 (br d, *J*=8.4 Hz, 1H), 6.95 (br d, *J*=8.6 Hz, 2H), 4.03 (br dd, *J*=11.3, 4.0 Hz, 1H), 3.91 - 3.79 (m, 5H), 3.77 - 3.66 (m, 3H), 3.64 - 3.58 (m, 5H), 3.42 - 3.33 (m, 1H), 3.25 - 3.16 (m, 4H), 2.49 - 2.43 (m, 3H), 2.30 (br d, *J*=7.3 Hz, 2H), 2.14 (br t, *J*=6.3 Hz, 2H), 1.95 (br dd, *J*=12.3, 5.0 Hz, 1H), 1.78 - 1.65 (m, 3H), 1.62 - 1.47 (m, 3H); MS ESI *m/z* 599.2 (M+H); Anal. HPLC Retention time: 1.38 (Method 2); ROCK2 IC₅₀=0.7 nM.


WHAT IS CLAIMED IS:

1. A compound according to Formula (I):



- 5 or an enantiomer, a diastereomer, a stereoisomer, a tautomer, a pharmaceutically acceptable salt thereof, wherein

Ring A is selected from



J₁, J₂, J₃, and J₄ are independently selected from N, CR₃, and CR₄; provided no more than

- 10 two of J₁, J₂, J₃, and J₄ are N;

L is selected from -C(O)-, -C(O)NH-, and -S(O)_p-;


R₁, at each occurrence, is independently selected from H, F, Cl, Br, OH, CN, NR_aR_a,
-OC₁₋₄ alkyl substituted with 0-3 R_e, and C₁₋₄ alkyl substituted with 0-3 R_e;

- 15 R₂, at each occurrence, is independently selected from H, -(CH₂)_rOR_b, (CH₂)_rS(O)_pR_c,
-(CH₂)_rC(=O)R_b, -(CH₂)_rNR_aR_a, -(CH₂)_rCN, -(CH₂)_rC(=O)NR_aR_a,
-(CH₂)_rNR_aC(=O)R_b, -(CH₂)_rNR_aC(=O)NR_aR_a, -(CH₂)_rNR_aC(=O)OR_b,
-(CH₂)_rOC(=O)NR_aR_a, -(CH₂)_rC(=O)OR_b, -(CH₂)_rS(O)_pNR_aR_a,
-(CH₂)_rNR_aS(O)_pNR_aR_a, -(CH₂)_rNR_aS(O)_pR_c, (CH₂)_r-C₃₋₆ carbocyclyl substituted
with 0-3 R_e, and -(CH₂)_r-heterocyclyl substituted with 0-3 R_e;

- 20 R₃, at each occurrence, is independently selected from H, F, Cl, Br, CN, C₁₋₄ alkyl
substituted with 0-3 R_e, -(CH₂)_rOR_b, (CH₂)_rS(O)_pR_c, -(CH₂)_rC(=O)R_b,
-(CH₂)_rNR_aR_a, -(CH₂)_rC(=O)NR_aR_a, -(CH₂)_rC(=O)(CH₂)_rNR_aR_a, -(CH₂)_rCN,
-(CH₂)_rNR_aC(=O)R_b, -(CH₂)_rNR_aC(=O)OR_b, -(CH₂)_rOC(=O)NR_aR_a,
-(CH₂)_rNR_aC(=O)NR_aR_a, -(CH₂)_rC(=O)OR_b, -(CH₂)_rS(O)_pNR_aR_a,

- $-(CH_2)_rNR_aS(O)_pNR_aR_a$, $-(CH_2)_rNR_aS(O)_pR_c$, $(CH_2)_r$ -C₃₋₆ carbocyclyl substituted with 0-3 R_e, and $-(CH_2)_r$ -heterocyclyl substituted with 0-3 R_e;
- R₄, at each occurrence, is independently selected from H, F, Cl, Br, OH, CN, OC₁₋₄ alkyl substituted with 0-3 R_e, NR_aR_a, and C₁₋₄ alkyl substituted with 0-3 R_e;
- 5 R₅ is selected from C₃₋₁₀ carbocyclyl substituted with 0-5 R₆ and heterocyclyl comprising carbon atoms and 1-6 heteroatoms selected from N, NR_{6a}, S, and O, and substituted with 0-6 R₆;
- R₆, at each occurrence, is independently selected from F, Cl, Br, =O, C₁₋₄alkyl, C₂₋₄alkenyl, C₂₋₄alkynyl, nitro, $-(CHR_d)_rS(O)_pR_c$, $-(CHR_d)_rS(O)_pNR_aR_a$,
- 10 $-(CHR_d)_rNR_aS(O)_pR_c$, $-(CHR_d)_rOR_b$, $-(CHR_d)_rCN$, $-(CHR_d)_rNR_7R_7$, $-(CHR_d)_rNR_aC(=O)R_b$, $-(CHR_d)_rNR_aC(=O)NR_aR_a$, $-(CHR_d)_rC(=O)OR_b$, $-(CHR_d)_rC(=O)R_b$, $-(CHR_d)_rC(=O)NR_aR_a$, $-(CHR_d)_rOC(=O)R_b$, $-(CHR_d)_r$ -C₃₋₆cycloalkyl, $-(CHR_d)_r$ -heterocyclyl, $-(CHR_d)_r$ -aryl, and $-(CHR_d)_r$ -heteroaryl, wherein said alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclyl, aryl, or heteroaryl is
- 15 substituted with 0-4 R_e;
- R_{6a} is selected from H, C₁₋₄alkyl, C₂₋₄alkenyl, C₂₋₄alkynyl, $-S(O)_pR_c$, $-S(O)_pNR_aR_a$, $-C(=O)OR_b$, $-C(=O)R_b$, $-C(=O)NR_aR_a$, $-(CHR_d)_r$ -C₃₋₆cycloalkyl, $-(CHR_d)_r$ -heterocyclyl, $-(CHR_d)_r$ -aryl, and $-(CHR_d)_r$ -heteroaryl, wherein said alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclyl, aryl, or heteroaryl is substituted
- 20 with 0-4 R_e;
- R₇, at each occurrence, is independently selected from H, C₁₋₆ alkyl substituted with 1-5 R_a, C₂₋₆ alkenyl substituted with 1-5 R_a, C₂₋₆ alkynyl substituted with 1-5 R_a, $-(CH_2)_r$ -C₃₋₁₀carbocyclyl substituted with 1-5 R_a, and $-(CH_2)_r$ -heterocyclyl substituted with 1-5 R_a; or R₇ and R₇ together with the nitrogen atom to which
- 25 they are both attached form a heterocyclic ring substituted with 1-5 R_a;
- R_a, at each occurrence, is independently selected from H, C(=O)OR_b, C₁₋₆ alkyl substituted with 0-5 R_e, C₂₋₆ alkenyl substituted with 0-5 R_e, C₂₋₆ alkynyl substituted with 0-5 R_e, $-(CH_2)_r$ -C₃₋₁₀carbocyclyl substituted with 0-5 R_e, and $-(CH_2)_r$ -heterocyclyl substituted with 0-5 R_e; or R_a and R_a together with the
- 30 nitrogen atom to which they are both attached form a heterocyclic ring substituted with 0-5 R_e;

- R_b , at each occurrence, is independently selected from H, C_{1-6} alkyl substituted with 0-5 R_e , C_{2-6} alkenyl substituted with 0-5 R_e , C_{2-6} alkynyl substituted with 0-5 R_e , $-(CH_2)_r-C_{3-10}$ carbocyclyl substituted with 0-5 R_e , and $-(CH_2)_r$ -heterocyclyl substituted with 0-5 R_e ;
- 5 R_c , at each occurrence, is independently selected from C_{1-6} alkyl substituted with 0-5 R_e , C_{2-6} alkenyl substituted with 0-5 R_e , C_{2-6} alkynyl substituted with 0-5 R_e , C_{3-6} carbocyclyl, and heterocyclyl;
- R_d , at each occurrence, is independently selected from H and C_{1-4} alkyl substituted with 0-5 R_e ;
- 10 R_e , at each occurrence, is independently selected from F, Cl, Br, CN, NO_2 , $=O$, C_{1-6} alkyl substituted with 1-5 R_f , C_{2-6} alkenyl, C_{2-6} alkynyl, $-(CH_2)_rOR_f$, $S(O)_pR_f$, $C(=O)NR_fR_f$, $NR_fC(=O)R_d$, $S(O)_pNR_fR_f$, $NR_fS(O)_pR_d$, $NR_fC(=O)OR_d$, $OC(=O)NR_fR_f$, $-(CH_2)_rNR_fR_f$, $-(CH_2)_r-C_{3-6}$ cycloalkyl substituted with 1-5 R_f , $-O(CH_2)_r-C_{3-6}$ cycloalkyl substituted with 1-5 R_f , $-(CH_2)_r$ -heterocyclyl substituted with 1-5 R_f , $-O(CH_2)_r$ -heterocyclyl substituted with 1-5 R_f , $-(CH_2)_r$ -aryl substituted with 1-5 R_f , $-O(CH_2)_r$ -aryl substituted with 1-5 R_f , $-(CH_2)_r$ -heteroaryl substituted with 1-5 R_f , and $-O(CH_2)_r$ -heteroaryl substituted with 1-5 R_f ;
- 15 R_f , at each occurrence, is independently selected from H, F, Cl, Br, CN, NO_2 , $=O$, C_{1-5} alkyl (optionally substituted with F, Cl, Br, OH, OC_{1-4} alkyl, NR_gR_g), $-(CH_2)_r-C_{3-10}$ carbocyclyl and $-(CH_2)_r$ -heterocyclyl, or R_f and R_f together with the nitrogen atom to which they are both attached form a heterocyclic ring optionally substituted with C_{1-4} alkyl;
- 20 R_g , at each occurrence, is independently selected from H and C_{1-5} alkyl;
- m is an integer of zero, 1 or 2;
- 25 p is an integer of zero, 1 or 2;
- q is an integer of zero or 1;
- r is an integer of zero, 1, 2, 3 or 4; and
- t is an integer of zero, 1 or 2.
- 30 2. The compound of claim 1 or an enantiomer, a diastereomer, a stereoisomer, a tautomer, a pharmaceutically acceptable salt thereof, wherein




R_6 , at each occurrence, is independently selected from F, Cl, Br, $=O$, C_{1-4} alkyl, C_{2-4} alkenyl, C_{2-4} alkynyl, nitro, $-(CHR_d)_rS(O)_pR_c$, $-(CHR_d)_rS(O)_pNR_aR_a$, $-(CHR_d)_rNR_aS(O)_pR_c$, $-(CHR_d)_rOR_b$, $-(CHR_d)_rCN$, $-(CHR_d)_rNR_aR_a$, $-(CHR_d)_rNR_aC(=O)R_b$, $-(CHR_d)_rNR_aC(=O)NR_aR_a$, $-(CHR_d)_rC(=O)OR_b$, $-(CHR_d)_rC(=O)R_b$, $-C(=O)NR_aR_a$, $-(CHR_d)_rOC(=O)R_b$, $-(CHR_d)_r-C_{3-6}$ cycloalkyl, $-(CHR_d)_r$ -heterocyclyl, $-(CHR_d)_r$ -aryl, and $-(CHR_d)_r$ -heteroaryl, wherein said alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclyl, aryl, or heteroaryl is substituted with 0-4 R_c ;

- R_{6a} , at each occurrence, is independently selected from H, C_{1-4} alkyl, $-S(O)_pR_c$, $-S(O)_pNR_aR_a$, $-C(=O)OR_b$, $-C(=O)R_b$, $-C(=O)NR_aR_a$, $-(CHR_d)_r-C_{3-6}$ cycloalkyl, $-(CHR_d)_r$ -heterocyclyl, $-(CHR_d)_r$ -aryl, and $-(CHR_d)_r$ -heteroaryl, wherein said alkyl, cycloalkyl, heterocyclyl, aryl, or heteroaryl is substituted with 0-4 R_e ;
- 5 R_7 , at each occurrence, is independently selected from H, C_{1-6} alkyl substituted with 1-5 R_a , C_{2-6} alkenyl substituted with 1-5 R_a , C_{2-6} alkynyl substituted with 1-5 R_a , $-(CH_2)_r-C_{3-10}$ carbocyclyl substituted with 1-5 R_a , and $-(CH_2)_r$ -heterocyclyl substituted with 1-5 R_a ; or R_7 and R_7 together with the nitrogen atom to which they are both attached form a heterocyclic ring substituted with 1-5 R_a ;
- 10 R_a , at each occurrence, is independently selected from H, $rC(=O)OR_b$, C_{1-6} alkyl substituted with 0-5 R_e , C_{2-6} alkenyl substituted with 0-5 R_e , C_{2-6} alkynyl substituted with 0-5 R_e , $-(CH_2)_r-C_{3-10}$ carbocyclyl substituted with 0-5 R_e , and $-(CH_2)_r$ -heterocyclyl substituted with 0-5 R_e ; or R_a and R_a together with the nitrogen atom to which they are both attached form a heterocyclic ring substituted
- 15 with 0-5 R_e ;
- R_b , at each occurrence, is independently selected from H, C_{1-6} alkyl substituted with 0-5 R_e , C_{2-6} alkenyl substituted with 0-5 R_e , C_{2-6} alkynyl substituted with 0-5 R_e , $-(CH_2)_r-C_{3-10}$ carbocyclyl substituted with 0-5 R_e , and $-(CH_2)_r$ -heterocyclyl substituted with 0-5 R_e ;
- 20 R_c , at each occurrence, is independently selected from C_{1-6} alkyl substituted with 0-5 R_e , C_{2-6} alkenyl substituted with 0-5 R_e , C_{2-6} alkynyl substituted with 0-5 R_e , C_{3-6} carbocyclyl, and heterocyclyl;
- R_d , at each occurrence, is independently selected from H and C_{1-4} alkyl substituted with 0-5 R_e ;
- 25 R_e , at each occurrence, is independently selected from F, Cl, Br, CN, NO_2 , $=O$, C_{1-6} alkyl substituted with 1-5 R_f , C_{2-6} alkenyl, C_{2-6} alkynyl, $-(CH_2)_rOR_f$, $S(O)_pR_f$, $C(=O)NR_fR_f$, $NR_fC(=O)R_d$, $S(O)_pNR_fR_f$, $NR_fS(O)_pR_d$, $NR_fC(=O)OR_d$, $OC(=O)NR_fR_f$, $-(CH_2)_rNR_fR_f$, $-(CH_2)_r-C_{3-6}$ cycloalkyl substituted with 1-5 R_f , $-O(CH_2)_r-C_{3-6}$ cycloalkyl substituted with 1-5 R_f , $-(CH_2)_r$ -heterocyclyl substituted
- 30 with 1-5 R_f , $-O(CH_2)_r$ -heterocyclyl substituted with 1-5 R_f , $-(CH_2)_r$ -aryl substituted with 1-5 R_f , $-O(CH_2)_r$ -aryl substituted with 1-5 R_f , $-(CH_2)_r$ -heteroaryl substituted with 1-5 R_f , and $-O(CH_2)_r$ -heteroaryl substituted with 1-5 R_f ;

R_f , at each occurrence, is independently selected from H, F, Cl, Br, CN, OH, C_{1-5} alkyl (optionally substituted with F, Cl, Br, OH, and OC_{1-4} alkyl, $-NR_gR_g$), $-(CH_2)_r-C_{3-10}$ carbocyclyl, and $-(CH_2)_r$ -heterocyclyl, or R_f and R_f together with the nitrogen atom to which they are both attached form a heterocyclic ring optionally substituted with C_{1-4} alkyl; and

R_g , at each occurrence, is independently selected from H and C_{1-5} alkyl.


3. The compound of claim 2, having Formula (II):



(II)

or an enantiomer, a diastereomer, a stereoisomer, a tautomer, a pharmaceutically acceptable salt thereof, wherein

Ring A is selected from



J_2 and J_4 are independently selected from N, CR_3 , and CR_4 ;

L is selected from $-C(O)-$ and $-S(O)_p-$;


R_1 , at each occurrence, is independently selected from H, F, Cl, Br, CN, NR_aR_a , and C_{1-4} alkyl substituted with 0-4 R_e ;


R_2 , at each occurrence, is independently selected from H, F, Cl, Br, OH, CN, NR_aR_a , and C_{1-4} alkyl substituted with 0-4 R_e ;

R_3 , at each occurrence, is independently selected from H, F, Cl, Br, CN, C_{1-4} alkyl substituted with 0-3 R_e , $-(CH_2)_rOR_b$, and $-C_{3-6}$ cycloalkyl;

R_4 , at each occurrence, is independently selected from H, F, Cl, Br, OH, CN, OC_{1-4} alkyl substituted with 0-3 R_e , and C_{1-4} alkyl substituted with 0-3 R_e ;

R_5 is selected from






- 5 R_6 , at each occurrence, is independently selected from F, Cl, Br, C_{1-4} alkyl, C_{2-4} alkenyl, C_{2-4} alkynyl, nitro, $-(CH_2)_rS(O)_pR_c$, $-(CH_2)_rS(O)_pNR_aR_a$, $-(CH_2)_rNR_aS(O)_pR_c$, $-(CH_2)_rOR_b$, $-(CH_2)_rCN$, $-(CH_2)_rNR_aR_a$, $-(CH_2)_rNR_aC(=O)R_b$, $-(CH_2)_rNR_aC(=O)NR_aR_a$, $-(CH_2)_rC(=O)OR_b$, $-(CH_2)_rC(=O)R_b$, $-C(=O)NR_aR_a$, $-(CH_2)_rOC(=O)R_b$, $-(CH_2)_r-C_{3-6}$ cycloalkyl, $-(CH_2)_r$ -heterocyclyl, $-(CH_2)_r$ -aryl, and $-(CH_2)_r$ -heteroaryl, wherein said alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclyl, aryl, or heteroaryl is substituted with 0-4 R_e ;
- 10 R_{6a} , at each occurrence, is independently selected from H, C_{1-4} alkyl, $-S(O)_pR_c$, $-S(O)_pNR_aR_a$, $-C(=O)OR_b$, $-C(=O)R_b$, $-C(=O)NR_aR_a$, $-(CH_2)_r-C_{3-6}$ cycloalkyl, $-(CH_2)_r$ -heterocyclyl, $-(CH_2)_r$ -aryl, and $-(CH_2)_r$ -heteroaryl, wherein said alkyl, cycloalkyl, heterocyclyl, aryl, or heteroaryl is substituted with 0-4 R_e ;
- 15 R_7 , at each occurrence, is independently selected from H, C_{1-6} alkyl substituted with 0-5 R_a , C_{2-6} alkenyl substituted with 1-5 R_a , C_{2-6} alkynyl substituted with 1-5 R_a , $-(CH_2)_r-C_{3-10}$ carbocyclyl substituted with 1-5 R_a , and $-(CH_2)_r$ -heterocyclyl substituted with 1-5 R_a ; or R_7 and R_7 together with the nitrogen atom to which they are both attached form a heterocyclic ring substituted with 1-5 R_a ;
- 20


- R_a , at each occurrence, is independently selected from H, $C(=O)OR_b$, C_{1-6} alkyl substituted with 0-5 R_e , C_{2-6} alkenyl substituted with 0-5 R_e , C_{2-6} alkynyl substituted with 0-5 R_e , $-(CH_2)_r-C_{3-10}$ cycloalkyl substituted with 0-5 R_e , $-(CH_2)_r$ -aryl substituted with 0-5 R_e , and $-(CH_2)_r$ -heterocyclyl substituted with 0-5 R_e ; or R_a and R_a together with the nitrogen atom to which they are both attached form a heterocyclic ring substituted with 0-5 R_e ;
- R_b , at each occurrence, is independently selected from H, C_{1-6} alkyl substituted with 0-5 R_e , C_{2-6} alkenyl substituted with 0-5 R_e , C_{2-6} alkynyl substituted with 0-5 R_e , $-(CH_2)_r-C_{3-10}$ carbocyclyl substituted with 0-5 R_e , and $-(CH_2)_r$ -heterocyclyl substituted with 0-5 R_e ;
- R_c , at each occurrence, is independently selected from C_{1-6} alkyl substituted with 0-5 R_e , C_{2-6} alkenyl substituted with 0-5 R_e , C_{2-6} alkynyl substituted with 0-5 R_e , C_{3-6} carbocyclyl, and heterocyclyl;
- R_e , at each occurrence, is independently selected from F, Cl, Br, CN, NO_2 , $=O$, C_{1-6} alkyl substituted with 1-5 R_f , C_{2-6} alkenyl, C_{2-6} alkynyl, CO_2H , $-(CH_2)_rOR_f$, $S(O)_pR_f$, $S(O)_pNR_fR_f$, $NR_fS(O)_pR_d$, $-(CH_2)_rNR_fR_f$, $-(CH_2)_r-C_{3-6}$ cycloalkyl, $-O(CH_2)_r-C_{3-6}$ cycloalkyl, $-(CH_2)_r$ -heterocyclyl, $-O(CH_2)_r$ -heterocyclyl, $-(CH_2)_r$ -aryl, and $-O(CH_2)_r$ -aryl;
- R_f , at each occurrence, is independently selected from H, F, Cl, Br, CN, OH, C_{1-5} alkyl (optionally substituted with F, Cl, Br, OH, and OC_{1-4} alkyl, $-NR_gR_g$), $-(CH_2)_r-C_{3-6}$ cycloalkyl, and $-(CH_2)_r$ -phenyl; and
- R_g , at each occurrence, is independently selected from H and C_{1-5} alkyl.

4. The compound of claim 3, having Formula (III):



or an enantiomer, a diastereomer, a stereoisomer, a tautomer, a pharmaceutically acceptable salt thereof, wherein

Ring A is selected from



J₂ is selected from N, CR₃, and CR₄;

L is selected from -C(O)- and -S(O)_p-;

R₁, at each occurrence, is independently selected from H, F, Cl, Br, CN, NR_aR_a, and

5 C₁₋₄alkyl substituted with 0-4 R_e;

R₂, at each occurrence, is independently selected from H and C₁₋₄alkyl substituted with 0-4 R_e;

R₃, at each occurrence, is independently selected from H, F, Cl, Br, CN, C₁₋₄ alkyl substituted with 0-3 R_e, -(CH₂)_rOR_b, and -C₃₋₆cycloalkyl;

10 R₄, at each occurrence, is independently selected from H, F, Cl, Br, OH, CN, OC₁₋₄ alkyl substituted with 0-3 R_e, and C₁₋₄ alkyl substituted with 0-3 R_e;

R₆, at each occurrence, is independently selected from F, Cl, Br, C₁₋₄ alkyl, -(CH₂)_rOR_b, -(CH₂)_rCN, -(CH₂)_rNR₇R₇, -(CH₂)_rNR_aC(=O)R_b, -(CH₂)_rNR_aC(=O)NR_aR_a, -(CH₂)_rC(=O)OR_b, -(CH₂)_rC(=O)R_b, -C(=O)NR_aR_a, -(CH₂)_rOC(=O)R_b, 15 -(CH₂)_rC(=O)NR_aR_a, -(CH₂)_r-C₃₋₆cycloalkyl, -(CH₂)_r-heterocyclyl, -(CH₂)_r-aryl, and -(CH₂)_r-heteroaryl, wherein said alkyl, cycloalkyl, heterocyclyl, aryl, or heteroaryl is substituted with 0-4 R_e;

R₇, at each occurrence, is independently selected from H, C₁₋₆ alkyl substituted with 1-5 R_a, C₂₋₆ alkenyl substituted with 1-5 R_a, C₂₋₆ alkynyl substituted with 1-5 R_a, 20 -(CH₂)_r-C₃₋₁₀carbocyclyl substituted with 1-5 R_a, and -(CH₂)_r-heterocyclyl substituted with 1-5 R_a; or R₇ and R₇ together with the nitrogen atom to which they are both attached form a heterocyclic ring substituted with 1-5 R_a;

R_a, at each occurrence, is independently selected from H, C(=O)OR_b, C₁₋₆ alkyl substituted with 0-5 R_e, C₂₋₆ alkenyl substituted with 0-5 R_e, C₂₋₆ alkynyl substituted with 0-5 R_e, -(CH₂)_r-C₃₋₁₀cycloalkyl substituted with 0-5 R_e, 25 -(CH₂)_r-aryl substituted with 0-5 R_e, and -(CH₂)_r-heterocyclyl substituted with 0-5 R_e; or R_a and R_a together with the nitrogen atom to which they are both attached form a heterocyclic ring substituted with 0-5 R_e;

R_b , at each occurrence, is independently selected from H, C_{1-6} alkyl substituted with 0-5 R_e , C_{2-6} alkenyl substituted with 0-5 R_e , C_{2-6} alkynyl substituted with 0-5 R_e , $-(CH_2)_r-C_{3-10}$ carbocyclyl substituted with 0-5 R_e , and $-(CH_2)_r$ -heterocyclyl substituted with 0-5 R_e ;


- 5 R_c , at each occurrence, is independently selected from C_{1-6} alkyl substituted with 0-5 R_e , C_{2-6} alkenyl substituted with 0-5 R_e , C_{2-6} alkynyl substituted with 0-5 R_e , C_{3-6} carbocyclyl, and heterocyclyl;

R_e , at each occurrence, is independently selected from F, Cl, Br, CN, NO_2 , $=O$, C_{1-6} alkyl substituted with 1-5 R_f , C_{2-6} alkenyl, C_{2-6} alkynyl, CO_2H , $-(CH_2)_rOR_f$, $S(O)_pR_f$,
 10 $S(O)_pNR_fR_f$, $NR_fS(O)_pR_d$, $-(CH_2)_rNR_fR_f$, $-(CH_2)_r-C_{3-6}$ cycloalkyl, $-O(CH_2)_r-C_{3-6}$ cycloalkyl, $-(CH_2)_r$ -heterocyclyl, $-O(CH_2)_r$ -heterocyclyl, $-(CH_2)_r$ -aryl, and $-O(CH_2)_r$ -aryl;

R_f , at each occurrence, is independently selected from H, F, Cl, Br, CN, OH, C_{1-5} alkyl (optionally substituted with F, Cl, Br, OH, OC_{1-4} alkyl, NR_gR_g), $-(CH_2)_r$ -
 15 C_{3-6} cycloalkyl, and $-(CH_2)_r$ -phenyl; and

R_g , at each occurrence, is independently selected from H and C_{1-5} alkyl.

5. The compound of claim 4, having Formula (IV):



20 or enantiomer, a diastereomer, a stereoisomer, a tautomer, a pharmaceutically acceptable salt thereof, wherein


J_2 is selected from N and CH;

R_3 is selected from H, CN, C_{1-4} alkyl, $-OC_{1-3}$ alkyl, and $-C_{3-6}$ cycloalkyl;

R_6 , at each occurrence, is independently selected from F, Cl, Br, C_{1-4} alkyl, $-(CH_2)_rOR_b$,
 25 $-(CH_2)_rNR_7R_7$, $-(CH_2)_rNR_aC(=O)R_b$, $-(CH_2)_rNR_aC(=O)NR_aR_a$, $-(CH_2)_rC(=O)OR_b$, $-(CH_2)_rC(=O)R_b$, $-C(=O)NR_aR_a$, $-(CH_2)_rOC(=O)R_b$, $-(CH_2)_rC(=O)NR_aR_a$, $-(CH_2)_r$ - C_{3-6} cycloalkyl, $-(CH_2)_r$ -heterocyclyl, $-(CH_2)_r$ -aryl, and $-(CH_2)_r$ -heteroaryl, wherein said alkyl, cycloalkyl, heterocyclyl, aryl, or heteroaryl is substituted with 0-4 R_e ;


- R_7 , at each occurrence, is independently selected from H, C_{1-6} alkyl substituted with 1-5 R_a , C_{2-6} alkenyl substituted with 1-5 R_a , C_{2-6} alkynyl substituted with 1-5 R_a , $-(CH_2)_r-C_{3-10}$ carbocyclyl substituted with 1-5 R_a , and $-(CH_2)_r$ -heterocyclyl substituted with 1-5 R_a ; or R_7 and R_7 together with the nitrogen atom to which they are both attached form a heterocyclic ring substituted with 1-5 R_a ;
- R_a , at each occurrence, is independently selected from H, $C(=O)OR_b$, C_{1-6} alkyl substituted with 0-5 R_e , $-(CH_2)_r-C_{3-6}$ cycloalkyl substituted with 0-5 R_e , $-(CH_2)_r$ -aryl substituted with 0-5 R_e , and $-(CH_2)_r$ -heterocyclyl substituted with 0-5 R_e ; or R_a and R_a together with the nitrogen atom to which they are both attached form a heterocyclic ring substituted with 0-5 R_e ;
- R_b , at each occurrence, is independently selected from H, C_{1-6} alkyl substituted with 0-5 R_e , $-(CH_2)_r-C_{3-10}$ carbocyclyl substituted with 0-5 R_e , and $-(CH_2)_r$ -heterocyclyl substituted with 0-5 R_e ;
- R_e , at each occurrence, is independently selected from F, Cl, Br, CN, NO_2 , $=O$, C_{1-6} alkyl substituted with 1-5 R_f , CO_2H , OR_f , $NHS(O)_pC_{1-4}$ alkyl, NR_fR_f , $-(CH_2)_r-C_{3-6}$ cycloalkyl, $-O(CH_2)_r-C_{3-6}$ cycloalkyl, $-(CH_2)_r$ -heterocyclyl, $-O(CH_2)_r$ -heterocyclyl, $-(CH_2)_r$ -aryl, and $-O(CH_2)_r$ -aryl;
- R_f , at each occurrence, is independently selected from H, F, Cl, Br, CN, $=O$, C_{1-5} alkyl (optionally substituted with F, Cl, Br, OH, OC_{1-4} alkyl, NR_gR_g), $-(CH_2)_r-C_{3-6}$ cycloalkyl, and $-(CH_2)_r$ -phenyl;
- R_g , at each occurrence, is independently selected from H and C_{1-5} alkyl; and
- r is an integer of zero, 1, 2 or 3.

6. The compound of claim 4, having Formula (V):



or enantiomer, a diastereomer, a stereoisomer, a tautomer, a pharmaceutically acceptable salt thereof, wherein

Ring A is selected from




R_1 is selected from H, F, Cl, Br, NR_aR_a , and C_{1-4} alkyl;

R_3 is $-OC_{1-3}$ alkyl;

R_7 , at each occurrence, is independently selected from H, C_{1-6} alkyl substituted with 1-5

5 R_a ; or R_7 and R_7 together with the nitrogen atom to which they are both attached form a heterocyclic ring selected from



R_a , at each occurrence, is independently selected from H, $C(=O)OR_b$, C_{1-6} alkyl

10 substituted with 0-5 R_e , $-(CH_2)_r-C_{3-6}$ cycloalkyl substituted with 0-5 R_e , $-(CH_2)_r$ -aryl substituted with 0-5 R_e , and $-(CH_2)_r$ -heterocyclyl substituted with 0-5 R_e ;

R_b , at each occurrence, is independently selected from H, C_{1-6} alkyl substituted with 0-5 R_e , $-(CH_2)_r-C_{3-10}$ carbocyclyl substituted with 0-5 R_e , and $-(CH_2)_r$ -heterocyclyl

15 substituted with 0-5 R_e ;

R_e , at each occurrence, is independently selected from F, Cl, Br, CN, NO_2 , $=O$, C_{1-6} alkyl substituted with 1-4 R_f , CO_2H , OR_f , $NHS(O)_pC_{1-4}$ alkyl, NR_fR_f ,

$-(CH_2)_r-C_{3-6}$ cycloalkyl, $-(CH_2)_r$ -heterocyclyl, $-(CH_2)_r$ -aryl,


$-O(CH_2)_r-C_{3-6}$ cycloalkyl, $-O(CH_2)_r$ -heterocyclyl, and $-O(CH_2)_r$ -aryl;

20 R_f , at each occurrence, is independently selected from H, H, F, Cl, Br, CN, OH, C_{1-5} alkyl (optionally substituted with F, Cl, Br, OH, OC_{1-4} alkyl, NR_gR_g), $-(CH_2)_r-C_{3-6}$ cycloalkyl, and $-(CH_2)_r$ -phenyl;

R_g , at each occurrence, is independently selected from H and C_{1-5} alkyl; and

r is an integer of zero, 1, 2 or 3.


7. The compound of claim 6, having Formula (VI):



or enantiomer, a diastereomer, a stereoisomer, a tautomer, a pharmaceutically

5 acceptable salt thereof, wherein

Ring A is selected from



R₁ is selected from H, F, Cl, Br, NR_aR_a, and C₁₋₄alkyl;

R₃ is -OC₁₋₄ alkyl;

10 R_a is selected from H, C(=O)OC₁₋₄alkyl, C₁₋₆alkyl substituted with 0-5 R_e,

-(CH₂)_r-C₃₋₆cycloalkyl substituted with 0-5 R_e, -(CH₂)_r-aryl substituted with 0-5 R_e, and -(CH₂)_r-heterocyclyl substituted with 0-5 R_e;

R_e, at each occurrence, is independently selected from F, Cl, Br, CN, NO₂, =O, C₁₋₄ alkyl

CO₂H, OR_f, NHS(O)_pC₁₋₄alkyl, NR_fR_f, -(CH₂)_r-C₃₋₆cycloalkyl,

15 -(CH₂)_r-heterocyclyl, -(CH₂)_r-aryl, -O(CH₂)_r-C₃₋₆cycloalkyl,

-O(CH₂)_r-heterocyclyl, and -O(CH₂)_r-aryl;

R_f, at each occurrence, is independently selected from H, F, Cl, Br, CN, OH, C₁₋₅ alkyl

(optionally substituted with F, Cl, Br, OH, OC₁₋₄ alkyl, NR_gR_g), and -(CH₂)_r-phenyl; and


20 R_g, at each occurrence, is independently selected from H and C₁₋₅alkyl.

8. The compound of claim 7 or an enantiomer, a diastereomer, a

stereoisomer, a tautomer, a pharmaceutically acceptable salt thereof, wherein

R_a is selected from H, C(=O)OC₁₋₄alkyl, C₁₋₆ alkyl substituted with 0-5 R_e, -(CH₂)_r-C₃₋

25 ₆cycloalkyl substituted with 0-5 R_e, -(CH₂)_r-phenyl substituted with 0-5 R_e, and




R_e , at each occurrence, is independently selected from F, Cl, Br, CN, NO_2 , $=O$, C_{1-6} alkyl
 5 substituted with 1-4 R_f , OR_f , $NHS(O)_pC_{1-4}$ alkyl, NR_fR_f , $-(CH_2)_r-C_{3-6}$ cycloalkyl,
 $-(CH_2)_r$ -heterocyclyl, $-(CH_2)_r$ -aryl, $-O(CH_2)_r-C_{3-6}$ cycloalkyl,
 $-O(CH_2)_r$ -heterocyclyl, and $-O(CH_2)_r$ -aryl;


R_f , at each occurrence, is independently selected from H, F, Cl, Br, CN, OH, and C_{1-5}
 alkyl (optionally substituted with F, Cl, Br, OH, OC_{1-4} alkyl, and NR_gR_g), and
 10 $-(CH_2)_r$ -phenyl;


R_g , at each occurrence, is independently selected from H and C_{1-4} alkyl.


9. The compound of claim 3, having Formula (VII):



15 or enantiomer, a diastereomer, a stereoisomer, a tautomer, a pharmaceutically acceptable
 salt thereof, wherein







R_6 , at each occurrence, is independently selected from F, Cl, Br, C_{1-4} alkyl, C_{2-4} alkenyl,

C_{2-4} alkynyl, nitro, $-(CH_2)_rS(O)_pR_c$, $-(CH_2)_rS(O)_pNR_aR_a$, $-(CH_2)_rNR_aS(O)_pR_c$,

5 $-(CH_2)_rOR_b$, $-(CH_2)_rCN$, $-(CH_2)_rNR_aR_a$, $-(CH_2)_rNR_aC(=O)R_b$,

$-(CH_2)_rNR_aC(=O)NR_aR_a$, $-(CH_2)_rC(=O)OR_b$, $-(CH_2)_rC(=O)R_b$, $-C(=O)NR_aR_a$

$-(CH_2)_rOC(=O)R_b$, $-(CH_2)_r-C_{3-6}$ cycloalkyl, $-(CH_2)_r$ -heterocyclyl, $-(CH_2)_r$ -aryl, and

$-(CH_2)_r$ -heteroaryl, wherein said alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclyl,

aryl, or heteroaryl is substituted with 0-4 R_e ; and


10 R_{6a} , at each occurrence, is independently selected from H, C_{1-4} alkyl, $-S(O)_pR_c$,

$-S(O)_pNR_aR_a$, $-C(=O)OR_b$, $-C(=O)R_b$, $-C(=O)NR_aR_a$, $-(CH_2)_r-C_{3-6}$ cycloalkyl,

$-(CH_2)_r$ -heterocyclyl, $-(CH_2)_r$ -aryl, and $-(CH_2)_r$ -heteroaryl, wherein said alkyl,


cycloalkyl, heterocyclyl, aryl, or heteroaryl is substituted with 0-4 R_e .

15 10. The compound of claim 9, or enantiomer, a diastereomer, a stereoisomer, a tautomer, a pharmaceutically acceptable salt thereof, wherein




- R₆, at each occurrence, is independently selected from F, Cl, Br, =O, C₁₋₄alkyl, OR_b, C₃₋₆cycloalkyl, heterocyclyl, aryl and heteroaryl wherein said alkyl, cycloalkyl, heterocyclyl, aryl, or heteroaryl is substituted with 0-4 R_e; and
- R_{6a}, at each occurrence, is independently selected from H and C₁₋₄ alkyl.

11. The compound of claim 2, having Formula (VIII):



or an enantiomer, a diastereomer, a stereoisomer, a tautomer, a pharmaceutically acceptable salt thereof, wherein

L is selected from -C(O)- and -S(O)_p-;



R_6 , at each occurrence, is independently selected from F, Cl, Br, C_{1-4} alkyl, C_{2-4} alkenyl,

- 5 C_{2-4} alkynyl, nitro, $-(CH_2)_rS(O)_pR_c$, $-(CH_2)_rS(O)_pNR_aR_a$, $-(CH_2)_rNR_aS(O)_pR_c$, $-(CH_2)_rOR_b$, $-(CH_2)_rCN$, and $-(CH_2)_rNR_aR_a$;

R_{6a} , at each occurrence, is independently selected from H and C_{1-4} alkyl;

R_a , at each occurrence, is independently selected from H and C_{1-6} alkyl substituted with 0-5 R_e ;

- 10 R_b , at each occurrence, is independently selected from H and C_{1-6} alkyl substituted with 0-5 R_e ;

R_c , at each occurrence, is independently selected from C_{1-6} alkyl substituted with 0-5 R_e , C_{2-6} alkenyl substituted with 0-5 R_e , C_{2-6} alkynyl substituted with 0-5 R_e , C_{3-6} carbocyclyl, and heterocyclyl;

- 15 R_e , at each occurrence, is independently selected from F, Cl, Br, CN, NO_2 , $=O$, C_{1-6} alkyl substituted with 1-5 R_f , C_{2-6} alkenyl, C_{2-6} alkynyl, CO_2H , $-(CH_2)_rOR_f$, $S(O)_pR_f$, $S(O)_pNR_fR_f$, $NR_fS(O)_pC_{1-4}$ alkyl, $-(CH_2)_rNR_fR_f$, $-(CH_2)_r-C_{3-6}$ cycloalkyl, $-(CH_2)_r$ -heterocyclyl, and $-(CH_2)_r$ -aryl;

R_f , at each occurrence, is independently selected from H, F, Cl, Br, CN, OH, C_{1-5} alkyl

- 20 (optionally substituted with F, Cl, Br, OH, and OC_{1-4} alkyl, NR_gR_g), C_{3-6} cycloalkyl, and phenyl;

R_g , at each occurrence, is independently selected from H and C_{1-5} alkyl; and

r is an integer of zero, 1, 2 or 3.

12. A pharmaceutical composition comprising one or more compounds according to any one of claims 1-11 and a pharmaceutically acceptable carrier or diluent.

13. Use of a compound according to any one of claims 1-11 for prophylaxis
5 and/or treatment of a disorder associated with aberrant Rho kinase activity.

14. The use of claim 13, wherein said disorder is selected from the group consisting of a cardiovascular disorder, a smooth muscle related disorder, a fibrotic disease, an inflammatory disease, a neuropathic disorder, an oncologic disorder, and an
10 autoimmune disorder.

15. The use of claim 14, wherein said cardiovascular disorder is selected from the group consisting of angina, atherosclerosis, stroke, cerebrovascular disease, heart failure, coronary artery disease, myocardial infarction, peripheral vascular disease,
15 stenosis, vasospasm, hypertension and pulmonary hypertension.

INTERNATIONAL SEARCH REPORT

International application No
PCT/US2018/058611

A. CLASSIFICATION OF SUBJECT MATTER		
INV.	C07D471/10	C07D487/10
	A61K31/435	A61P9/00
	A61P35/00	A61P29/00
ADD.		
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED		
Minimum documentation searched (classification system followed by classification symbols)		
C07D		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)		
EPO-Internal, WPI Data		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A,P	WO 2018/009627 A1 (BRISTOL-MYERS SQUIBB COMPANY [US]) 11 January 2018 (2018-01-11) claims 1,14	1-15
A,P	WO 2018/009625 A1 (BRISTOL-MYERS SQUIBB COMPANY [US]) 11 January 2018 (2018-01-11) claims 1,13	1-15
A	WO 2016/112236 A1 (SQUIBB BRISTOL MYERS CO [US]) 14 July 2016 (2016-07-14) claims 1,14	1-15
A	WO 2011/143148 A1 (SANOFI SA [FR]; GAO ZHONGLI [US]; HALL DANIEL [US]; STEFANY DAVID [US]) 17 November 2011 (2011-11-17) claim 1	1-15
	- / - -	
☒ Further documents are listed in the continuation of Box C. ☒ See patent family annex.		
* Special categories of cited documents :		
"A" document defining the general state of the art which is not considered to be of particular relevance		
"E" earlier application or patent but published on or after the international filing date		
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)		
"O" document referring to an oral disclosure, use, exhibition or other means		
"P" document published prior to the international filing date but later than the priority date claimed		
"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention		
"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone		
"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art		
"&" document member of the same patent family		
Date of the actual completion of the international search		Date of mailing of the international search report
21 January 2019		04/02/2019
Name and mailing address of the ISA/ European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Fax: (+31-70) 340-3016		Authorized officer Gettins, Marc

INTERNATIONAL SEARCH REPORT

International application No
PCT/US2018/058611

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 2011/143150 A1 (SANOFI SA [FR]; GAO ZHONGLI [US]) 17 November 2011 (2011-11-17) claim 1 -----	1-15
A	WO 2010/065803 A1 (SANOFI AVENTIS [FR]; GAO ZHONGLI [US]; HALL DANIEL [US]; HARTUNG RYAN) 10 June 2010 (2010-06-10) claim 1 -----	1-15

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/US2018/058611

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 2018009627	A1	11-01-2018	NONE

WO 2018009625	A1	11-01-2018	NONE

WO 2016112236	A1	14-07-2016	AR 103990 A1 21-06-2017
		CN 107406426 A 28-11-2017	
		EP 3242873 A1 15-11-2017	
		JP 2018501285 A 18-01-2018	
		TW 201630899 A 01-09-2016	
		US 2018000788 A1 04-01-2018	
		WO 2016112236 A1 14-07-2016	

WO 2011143148	A1	17-11-2011	AR 081908 A1 31-10-2012
		EP 2569304 A1 20-03-2013	
		JP 5833105 B2 16-12-2015	
		JP 2013526526 A 24-06-2013	
		TW 201202251 A 16-01-2012	
		US 2013059863 A1 07-03-2013	
		US 2014315923 A1 23-10-2014	
		WO 2011143148 A1 17-11-2011	

WO 2011143150	A1	17-11-2011	AR 083718 A1 20-03-2013
		EP 2569280 A1 20-03-2013	
		JP 5968307 B2 10-08-2016	
		JP 2013526527 A 24-06-2013	
		TW 201206939 A 16-02-2012	
		US 2013065877 A1 14-03-2013	
		US 2014275005 A1 18-09-2014	
		WO 2011143150 A1 17-11-2011	

WO 2010065803	A1	10-06-2010	AR 074466 A1 19-01-2011
		AU 2009322248 A1 23-06-2011	
		CA 2745721 A1 10-06-2010	
		CN 102239168 A 09-11-2011	
		CO 6390058 A2 29-02-2012	
		CR 20110294 A 21-09-2011	
		DK 2373652 T3 20-01-2014	
		EC SP11011108 A 30-11-2011	
		EP 2373652 A1 12-10-2011	
		ES 2441221 T3 03-02-2014	
		IL 213290 A 30-01-2014	
		JP 5538421 B2 02-07-2014	
		JP 2012511012 A 17-05-2012	
		KR 20110100639 A 14-09-2011	
		MA 32943 B1 02-01-2012	
		NZ 593151 A 30-11-2012	
		PE 02612012 A1 04-04-2012	
		PT 2373652 E 16-01-2014	
		RU 2011127401 A 10-01-2013	
		TN 2011000233 A1 17-12-2012	
		TW 201036974 A 16-10-2010	
		US 2011257213 A1 20-10-2011	
		US 2012258979 A1 11-10-2012	
		WO 2010065803 A1 10-06-2010	
		ZA 201103448 B 25-01-2012	
