

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
29 November 2001 (29.11.2001)

PCT


(10) International Publication Number
WO 01/90147 A2

- (51) International Patent Classification⁷: **C07K 14/00** (72) Inventors: **WACHTER, Rebekka**; 33465 Irish Lane, Creswell, OR 97426 (US). **REMINGTON, S., James**; 2100 River Loop, Eugene, OR 97404 (US).
- (21) International Application Number: PCT/US01/16149
- (22) International Filing Date: 17 May 2001 (17.05.2001) (74) Agent: **HAILE, Lisa, A.**; Gray Cary Ware & Freidenrich, LLP, Suite 1600, 4365 Executive Drive, San Diego, CA 92121-2189 (US).
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
09/575,847 19 May 2000 (19.05.2000) US
- (71) Applicant: **THE STATE OF OREGON**, acting by and through **THE STATE BOARD OF HIGHER EDUCATION** on behalf of **THE UNIVERSITY OF OREGON** [US/US]; 1238 University of Oregon, Eugene, OR 97403 (US).
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European


[Continued on next page]

(54) Title: LONG WAVELENGTH ENGINEERED FLUORESCENT PROTEINS

(57) Abstract: Engineered fluorescent proteins, nucleic acids encoding them and methods of use.



A



B

WO 01/90147 A2



patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

Published:

— *without international search report and to be republished upon receipt of that report*

LONG WAVELENGTH ENGINEERED FLUORESCENT PROTEINS

5

BACKGROUND OF THE INVENTION

This a continuation in part of Application No 08/974,737, filed 11/19/97, now allowed, which is a continuation of Application No 08/911,825, filed 8/15/97, now issued as US patent No. 6,054,321, which is a continuation in part of Application No. 08/706,408, filed 8/30/96, now allowed, which claims the benefit of the earlier filing
10 date of a United States provisional patent application serial number 60/024,050 filed on August 16, 1996 each of which are herein incorporated by reference.

This invention was made in part with Government support under grant no. MCB 9418479 awarded by the National Science Foundation. The Government may have rights in this invention.

15

Fluorescent molecules are attractive as reporter molecules in many assay systems because of their high sensitivity and ease of quantification. Recently, fluorescent proteins have been the focus of much attention because they can be produced *in vivo* by biological systems, and can be used to trace intracellular events without the need to be introduced into the cell through microinjection or
20 permeabilization. The green fluorescent protein of *Aequorea victoria* is particularly interesting as a fluorescent protein. A cDNA for the protein has been cloned. (D.C. Prasher et al., "Primary structure of the *Aequorea victoria* green-fluorescent protein," *Gene* (1992) 111:229-33.) Not only can the primary amino acid sequence of the protein be expressed from the cDNA, but the expressed protein can fluoresce. This
25 indicates that the protein can undergo the cyclization and oxidation believed to be necessary for fluorescence. *Aequorea* green fluorescent protein ("GFP") is a stable, proteolysis-resistant single chain of 238 residues and has two absorption maxima at around 395 and 475 nm. The relative amplitudes of these two peaks is sensitive to environmental factors (W. W. Ward. *Bioluminescence and Chemiluminescence* (M.

A. DeLuca and W. D. McElroy, eds) Academic Press pp. 235-242 (1981); W. W. Ward & S. H. Bokman *Biochemistry* 21:4535-4540 (1982); W. W. Ward et al. *Photochem. Photobiol.* 35:803-808 (1982)) and illumination history (A. B. Cubitt et al. *Trends Biochem. Sci.* 20:448-455 (1995)), presumably reflecting two or more ground states. Excitation at the primary absorption peak of 395 nm yields an emission maximum at 508 nm with a quantum yield of 0.72-0.85 (O. Shimomura and F.H. Johnson *J. Cell. Comp. Physiol.* 59:223 (1962); J. G. Morin and J. W. Hastings, *J. Cell. Physiol.* 77:313 (1971); H. Morise et al. *Biochemistry* 13:2656 (1974); W. W. Ward *Photochem. Photobiol. Reviews* (Smith, K. C. ed.) 4:1 (1979); A. B. Cubitt et al. *Trends Biochem. Sci.* 20:448-455 (1995); D. C. Prasher *Trends Genet.* 11:320-323 (1995); M. Chalfie *Photochem. Photobiol.* 62:651-656 (1995); W. W. Ward. *Bioluminescence and Chemiluminescence* (M. A. DeLuca and W. D. McElroy, eds) Academic Press pp. 235-242 (1981); W. W. Ward & S. H. Bokman *Biochemistry* 21:4535-4540 (1982); W. W. Ward et al. *Photochem. Photobiol.* 35:803-808 (1982)).

The fluorophore results from the autocatalytic cyclization of the polypeptide backbone between residues Ser⁶⁵ and Gly⁶⁷ and oxidation of the β bond of Tyr⁶⁶ (A. B. Cubitt et al. *Trends Biochem. Sci.* 20:448-455 (1995); C. W. Cody et al. *Biochemistry* 32:1212-1218 (1993); R. Heim et al. *Proc. Natl. Acad. Sci. USA* 91:12501-12504 (1994)). Mutation of Ser⁶⁵ to Thr (S65T) simplifies the excitation spectrum to a single peak at 488 nm of enhanced amplitude (R. Heim et al. *Nature* 373:664-665 (1995)), which no longer gives signs of conformational isomers (A. B. Cubitt et al. *Trends Biochem. Sci.* 20:448-455 (1995)).

Fluorescent proteins have been used as markers of gene expression, tracers of cell lineage and as fusion tags to monitor protein localization within living cells. (M. Chalfie et al., "Green fluorescent protein as a marker for gene expression," *Science* 263:802-805; A.B. Cubitt et al., "Understanding, improving and using green fluorescent proteins," *TIBS* 20, November 1995, pp. 448-455. U.S. patent 5,491,084, M. Chalfie and D. Prasher. Furthermore, engineered versions of *Aequorea* green fluorescent protein have been identified that exhibit altered fluorescence characteristics, including altered excitation and emission maxima, as well as excitation and emission spectra of different shapes. (R. Heim et al., "Wavelength

mutations and posttranslational autoxidation of green fluorescent protein," *Prôc. Natl. Acad. Sci. USA*, (1994) 91:12501-04; R. Heim et al., "Improved green fluorescence," *Nature* (1995) 373:663-665.)

A second class of applications rely on GFP as a specific indicator of some cellular property, and hence depend on the particular spectral characteristics of the variant employed. For recent reviews on GFP variants and their applications, see (Palm & Wlodawer, 1999; Tsien, 1998) , and for a review volume on specialized applications, see (Sullivan & Kay, 1999) . Biosensor applications include the use of differently colored GFPs for fluorescence resonance energy transfer (FRET) to monitor protein-protein interactions (Heim, 1999) or Ca²⁺ concentrations (Miyawaki et al., 1999) , and receptor insertions within GFP surface loops to monitor ligand binding (Baird et al., 1999; Doi & Yanagawa, 1999) .

The fluorescence emission of a number of variants is highly sensitive to the acidity of the environment (Elslinger et al., 1999; Wachter et al., 1998) . Hence, one particularly successful application of green fluorescent protein (GFP) as a visual reporter in live cells has been the determination of organelle or cytosol pH (Kneen et al., 1998; Llopis et al., 1998; Miesenbock et al., 1998; Robey et al., 1998) . The two chromophore charge states have been found to be relevant to the pH sensitivity of the intact protein, and have been characterized crystallographically in terms of conformational changes in the vicinity of the phenolic end (Elslinger et al., 1999) , and spectroscopically using Raman studies (Bell et al., 2000) . The neutral form of the chromophore, band A, absorbs around 400 nm in most variants, whereas the chromophore anion with the phenolic end deprotonated (band B) absorbs in the blue to green, depending on the particular mutations in the vicinity of the chromophore. WT GFP exhibits spectral characteristics that are consistent with two ground states characterized by a combination of bands A and B, the ratio of which is relatively invariant between pH 6 and 10 (Palm & Wlodawer, 1999; Ward et al., 1982) . It has been suggested that an internal equilibrium exists where a proton is shared between the chromophore phenolate and the carboxylate of Glu222 over a broad range of pH (Brejc et al., 1997; Palm et al., 1997) . Recent electrostatic calculations support this model (Scharnagl et al., 1999) , and estimate the theoretical pK_a for complete

chromophore deprotonation to be about 13, consistent with the observation of a doubling of emission intensity at pH 11-12 (Bokman & Ward, 1981; Palm & Wlodawer, 1999) .

In contrast to WT GFP, the chromophore of most variants titrates with a single
5 pK_a. The color emission and the chromophore pK_a are strongly modulated by the protein surroundings (Llopis *et al.*, 1998) . Glu222 is completely conserved among GFP homologs (Matz *et al.*, 1999) , and its substitution by a glutamine has been shown to dramatically reduce efficiency of chromophore generation (Elslinger *et al.*, 1999) . Protonation of Glu222 in S65T and in GFPs containing the T203Y mutation
10 (YFPs) is generally thought to be responsible for lowering the chromophore pK_a from that of WT to about 5.9 in GFP S65T (Elslinger *et al.*, 1999; Kneen *et al.*, 1998) , and 5.2 -5.4 in YFP (GFP S65G/V68L/S72A/T203Y) (Ormo *et al.*, 1996; Wachter & Remington, 1999). In the YFPs, it is thought that the crystallographically identified stacking interaction of the chromophore with Tyr203 is largely responsible for the
15 spectral red-shift (Wachter *et al.*, 1998) .

Unlike other variants, we have discovered that the YFP chromophore pK_a shows a strong dependence on the concentration of certain small anions such as chloride (Wachter & Remington, 1999) , and increases in pK_a from about 5.2 to 7.0 in the presence of 140 mM NaCl (Elslinger *et al.*, 1999) . This sensitivity can be
20 exploited to enable the creation of novel GFPs as biosensors to measure ions present both in the cytoplasm or in cellular compartments (Wachter & Remington, 1999) within living cells. The present invention includes the creation and use of novel GFP variants that permit the fluorescent measurement of a variety of ions, including halides such as chloride and iodide. These properties add variety and utility to the
25 arsenal of biologically based fluorescent indicators. There is a need for engineered fluorescent proteins with varied fluorescent properties and with the ability to respond to ion concentrations via a change in fluorescence characteristics.

BRIEF DESCRIPTION OF THE DRAWINGS

Figs. 1A-1B. (A) Schematic drawing of the backbone of GFP produced by Molscript (J.P. Kraulis, *J. Appl. Cryst.*, 24:946 (1991)). The chromophore is shown as a ball and stick model. (B) Schematic drawing of the overall fold of GFP. Approximate residue numbers mark the beginning and ending of the secondary structure elements.

Figs. 2A-2C. (A) Stereo drawing of the chromophore and residues in the immediate vicinity. Carbon atoms are drawn as open circles, oxygen is filled and nitrogen is shaded. Solvent molecules are shown as isolated filled circles. (B) Portion of the final $2F_o - F_c$ electron density map contoured at 1.0σ , showing the electron density surrounding the chromophore. (C) Schematic diagram showing the first and second spheres of coordination of the chromophore. Hydrogen bonds are shown as dashed lines and have the indicated lengths in Å. Inset: proposed structure of the carbinolamine intermediate that is presumably formed during generation of the chromophore.

Fig. 3 depicts the nucleotide sequence (SEQ ID NO:1) and deduced amino acid sequence (SEQ ID NO:2) of an *Aequorea* green fluorescent protein.

Figs. 4A-B depict the nucleotide sequence (SEQ ID NO:3) and deduced amino acid sequence (SEQ ID NO:4) of the engineered *Aequorea*-related fluorescent protein S65G/S72A/T203Y utilizing preferred mammalian codons and optimal Kozak sequence.

Figs. 5A to 5AT present the coordinates for the crystal structure of *Aequorea*-related green fluorescent protein S65T.

Fig. 6 shows the fluorescence excitation and emission spectra for engineered fluorescent proteins 20A and 10C (Table F). The vertical line at 528 nm compares the emission maxima of 10C, to the left of the line, and 20A, to the right of the line.

Fig 7. Shows absorbance scans of YFP at varying NaCl concentration and constant pH 6.4, buffered with 20 mM MES (—O— 0 mM NaCl, —∇— 15 mM NaCl, —□— 50mM NaCl, —◇— 100 mM NaCl, and —Δ— 400 mM NaCl). Band A

corresponds to the neutral form of the chromophore ($\lambda_{\text{max}}=392\text{nm}$), and band B corresponds to the chromophore anion ($\lambda_{\text{max}}=514\text{nm}$).

Fig 8. Shows normalized fluorescence emission of (a) YFP and (b) YFP-H148Q, as a function of pH and $[\text{Cl}^-]$ at constant ionic strength of 150 mM. The pH was controlled with 20 mM TAPS pH 8.0 (O), 20 mM HEPES pH 7.5 (Δ), 20 mM PIPES pH 7.0 ($\diamond$), and 20 mM MES pH 6.5 (V) and pH 6.0 (). Figure 8(b) also includes the fluorescence emission of YFP-H148Q as a function of $[\text{I}^-]$ at pH 7.5 (*). Potassium chloride (or iodide) was added to the indicated concentration, and the ionic strength was adjusted to 150 mM with potassium gluconate. The samples, containing approximately 0.01 mg/ml protein, were excited at 514 nm, and emission intensity was determined at 528 nm.

Fig. 9 Shows a stereoview of the $2F_o-F_c$ electron-density map of the YFP-H148Q chromophore, Tyr203, Arg96, Gln69, and the buried iodide after refinement. The 2.1 Å resolution map was contoured at +1 standard deviation. This figure was drawn by the program BOBSCRIPT.

Fig. 10 Shows a schematic diagram showing all residues that contain atoms within 5 Å of the buried iodide in the crystal structure of YFP-H148Q (iodide soak).

Fig. 11 Shows a stereoview of an overlay of a subset of residues lining the anion binding cavity of YFP-H148Q, with and without iodide (iodide-bound structure, grey; apo-structure, black). The iodide is represented by the center sphere. This figure was drawn by the program MOLSCRIPT (Kraulis, 1991).

Fig 12. Shows a schematic diagram of the immediate chromophore environment of YFP-H148Q in the (a) apo-structure, and (b) iodide-bound structure.

Fig. 13 Shows a stereoview of the solvent-accessible surface of the iodide-bound YFP-H148Q structure, calculated using a 1.4 Å probe radius. The surface was calculated after deleting all water molecules and the iodide. The chromophore and all surface segments in contact with the chromophore are also shown. The outer surface

of the protein is along the left edge of the figure. This figure was generated using the program MidasPlus™ (UCSF, 1994).

Fig. 14 Shows the backbone atom trace of β -strands 7 and 8 of YFP, the apo-structure of YFP-H148Q, and the iodide-bound structure of YFP-H148Q. The side chain of His148 (YFP) and Gln148 (YFP-H148Q), and a few water molecules are also shown. The dashed lines represent possible hydrogen bonds.

Fig. 15 Shows YFP chromophore pK_a as a function of halide concentration (— $\diamond$ — fluoride, — ∇ — iodide, — Δ — chloride, —O— bromide). The chromophore pK_a was estimated from absorbance scans at varying halide concentrations (see Materials and Methods). The data were curve-fit to equation 1 (see text).

SUMMARY OF THE INVENTION

This invention provides functional engineered fluorescent proteins with varied fluorescence characteristics that can be easily distinguished from currently existing green and blue fluorescent proteins. Such engineered fluorescent proteins enable the simultaneous measurement of two or more processes within cells and can be used as fluorescence energy donors or acceptors, as well as biosensors for detecting anions. Longer wavelength engineered fluorescent proteins are particularly useful because photodynamic toxicity and auto-fluorescence of cells are significantly reduced at longer wavelengths. In particular, the introduction of the substitution T203X, wherein X is an aromatic amino acid, results in an increase in the excitation and emission wavelength maxima of *Aequorea*-related fluorescent proteins.

In one aspect, this invention provides a nucleic acid molecule comprising a nucleotide sequence encoding a functional engineered fluorescent protein whose amino acid sequence is substantially identical to the amino acid sequence of *Aequorea* green fluorescent protein (SEQ ID NO:2) and which differs from SEQ ID NO:2 by at least an amino acid substitution located no more than about 0.5 nm from the chromophore of the engineered fluorescent protein, wherein the substitution alters the electronic environment of the chromophore, whereby the functional engineered

fluorescent protein has a different fluorescent property than *Aequorea* green fluorescent protein.

In one aspect this invention provides a nucleic acid molecule comprising a nucleotide sequence encoding a functional engineered fluorescent protein whose amino acid sequence is substantially identical to the amino acid sequence of *Aequorea* green fluorescent protein (SEQ ID NO:2) and which differs from SEQ ID NO:2 by at least a substitution at T203 and, in particular, T203X, wherein X is an aromatic amino acid selected from H, Y, W or F, said functional engineered fluorescent protein having a different fluorescent property than *Aequorea* green fluorescent protein. In one embodiment, the amino acid sequence further comprises a substitution at S65, wherein the substitution is selected from S65G, S65T, S65A, S65L, S65C, S65V and S65I. In another embodiment, the amino acid sequence differs by no more than the substitutions S65T/T203H; S65T/T203Y; S72A/F64L/S65G/T203Y; S65G/V68L/Q69K/S72A/T203Y; S72A/S65G/V68L/T203Y; S65G/S72A/T203Y; or S65G/S72A/T203W. In another embodiment, the amino acid sequence further comprises a substitution at Y66, wherein the substitution is selected from Y66H, Y66F, and Y66W. In another embodiment, the amino acid sequence further comprises a mutation from Table A. In another embodiment, the amino acid sequence further comprises a folding mutation. In another embodiment, the nucleotide sequence encoding the protein differs from the nucleotide sequence of SEQ ID NO:1 by the substitution of at least one codon by a preferred mammalian codon. In another embodiment, the nucleic acid molecule encodes a fusion protein wherein the fusion protein comprises a polypeptide of interest and the functional engineered fluorescent protein.

In another aspect, this invention provides a nucleic acid molecule comprising a nucleotide sequence encoding a functional engineered fluorescent protein whose amino acid sequence is substantially identical to the amino acid sequence of *Aequorea* green fluorescent protein (SEQ ID NO:2) and which differs from SEQ ID NO:2 by at least an amino acid substitution at L42, V61, T62, V68, Q69, Q94, N121, Y145, H148, V150, F165, I167, Q183, N185, L220, E222 (not E222G), or V224, said

functional engineered fluorescent protein having a different fluorescent property than *Aequorea* green fluorescent protein. In one embodiment, amino acid substitution is:

- L42X, wherein X is selected from C, F, H, W and Y,
V61X, wherein X is selected from F, Y, H and C,
5 T62X, wherein X is selected from A, V, F, S, D, N, Q, Y, H and C,
V68X, wherein X is selected from F, Y and H;
Q69X, wherein X is selected from K, R, E and G,
Q94X, wherein X is selected from D, E, H, K and N,
N121X, wherein X is selected from F, H, W and Y,
10 Y145X, wherein X is selected from W, C, F, L, E, H, K and Q,
H148X, wherein X is selected from F, Y, N, K, Q and R,
V150X, wherein X is selected from F, Y and H,
F165X, wherein X is selected from H, Q, W and Y,
I167X, wherein X is selected from F, Y and H,
15 Q183X, wherein X is selected from H, Y, E and K,
N185X, wherein X is selected from D, E, H, K and Q,
L220X, wherein X is selected from H, N, Q and T,
E222X, wherein X is selected from N and Q, or
V224X, wherein X is selected from H, N, Q, T, F, W and Y.

20

In a further aspect, this invention provides an expression vector comprising expression control sequences operatively linked to any of the aforementioned nucleic acid molecules. In a further aspect, this invention provides a recombinant host cell comprising the aforementioned expression vector.

- 25 In another aspect, this invention provides a functional engineered fluorescent protein whose amino acid sequence is substantially identical to the amino acid sequence of *Aequorea* green fluorescent protein (SEQ ID NO:2) and which differs from SEQ ID NO:2 by at least an amino acid substitution located no more than about 0.5 nm from the chromophore of the engineered fluorescent protein, wherein the
30 substitution alters the electronic environment of the chromophore, whereby the

functional engineered fluorescent protein has a different fluorescent property than *Aequorea* green fluorescent protein.

In another aspect, this invention provides a functional engineered fluorescent protein whose amino acid sequence is substantially identical to the amino acid sequence of *Aequorea* green fluorescent protein (SEQ ID NO:2) and which differs from SEQ ID NO:2 by at least the amino acid substitution at T203, and in particular, T203X, wherein X is an aromatic amino acid selected from H, Y, W or F, said functional engineered fluorescent protein having a different fluorescent property than *Aequorea* green fluorescent protein. In one embodiment, the amino acid sequence further comprises a substitution at S65, wherein the substitution is selected from S65G, S65T, S65A, S65L, S65C, S65V and S65I. In another embodiment, the amino acid sequence differs by no more than the substitutions S65T/T203H; S65T/T203Y; S72A/F64L/S65G/T203Y; S72A/S65G/V68L/T203Y; S65G/V68L/Q69K/S72A/T203Y; S65G/S72A/T203Y; or S65G/S72A/T203W. In another embodiment, the amino acid sequence further comprises a substitution at Y66, wherein the substitution is selected from Y66H, Y66F, and Y66W. In another embodiment, the amino acid sequence further comprises a folding mutation. In another embodiment, the engineered fluorescent protein is part of a fusion protein wherein the fusion protein comprises a polypeptide of interest and the functional engineered fluorescent protein.

In another aspect this invention provides a functional engineered fluorescent protein whose amino acid sequence is substantially identical to the amino acid sequence of *Aequorea* green fluorescent protein (SEQ ID NO:2) and which differs from SEQ ID NO:2 by at least an amino acid substitution at L42, V61, T62, V68, Q69, Q94, N121, Y145, H148, V150, F165, I167, Q183, N185, L220, E222, or V224, said functional engineered fluorescent protein having a different fluorescent property than *Aequorea* green fluorescent protein.

In another aspect, this invention provides a fluorescently labelled antibody comprising an antibody coupled to any of the aforementioned functional engineered fluorescent proteins. In one embodiment, the fluorescently labelled antibody is a

fusion protein wherein the fusion protein comprises the antibody fused to the functional engineered fluorescent protein.

In another aspect, this invention provides a nucleic acid molecule comprising a nucleotide sequence encoding an antibody fused to a nucleotide sequence encoding a functional engineered fluorescent protein of this invention.

In another aspect, this invention provides a fluorescently labelled nucleic acid probe comprising a nucleic acid probe coupled to a functional engineered fluorescent protein whose amino acid sequence of this invention. The fusion can be through a linker peptide.

In another aspect, this invention provides a method for determining whether a mixture contains a target comprising contacting the mixture with a fluorescently labelled probe comprising a probe and a functional engineered fluorescent protein of this invention; and determining whether the target has bound to the probe. In one embodiment, the target molecule is captured on a solid matrix.

In another aspect, this invention provides a method for engineering a functional engineered fluorescent protein having a fluorescent property different than *Aequorea* green fluorescent protein, comprising substituting an amino acid that is located no more than 0.5 nm from any atom in the chromophore of an *Aequorea*-related green fluorescent protein with another amino acid; whereby the substitution alters a fluorescent property of the protein. In one embodiment, the amino acid substitution alters the electronic environment of the chromophore.

In another aspect, this invention provides a method for engineering a functional engineered fluorescent protein having a different fluorescent property than *Aequorea* green fluorescent protein comprising substituting amino acids in a loop domain of an *Aequorea*-related green fluorescent protein with amino acids so as to create a consensus sequence for phosphorylation or for proteolysis.

In another aspect, this invention provides a method for producing fluorescence resonance energy transfer comprising providing a donor molecule comprising a functional engineered fluorescent protein this invention; providing an appropriate acceptor molecule for the fluorescent protein; and bringing the donor molecule and

the acceptor molecule into sufficiently close contact to allow fluorescence resonance energy transfer.

In another aspect, this invention provides a method for producing fluorescence resonance energy transfer comprising providing an acceptor molecule comprising a functional engineered fluorescent protein of this invention; providing an appropriate donor molecule for the fluorescent protein; and bringing the donor molecule and the acceptor molecule into sufficiently close contact to allow fluorescence resonance energy transfer. In one embodiment, the donor molecule is a engineered fluorescent protein whose amino acid sequence comprises the substitution T203I and the acceptor molecule is an engineered fluorescent protein whose amino acid sequence comprises the substitution T203X, wherein X is an aromatic amino acid selected from H, Y, W or F, said functional engineered fluorescent protein having a different fluorescent property than *Aequorea* green fluorescent protein.

In another aspect, this invention provides a crystal of a protein comprising a fluorescent protein with an amino acid sequence substantially identical to SEQ ID NO: 2, wherein said crystal diffracts with at least a 2.0 to 3.0 angstrom resolution.

In another embodiment, this invention provides computational method of designing a fluorescent protein comprising determining from a three dimensional model of a crystallized fluorescent protein comprising a fluorescent protein with a bound ligand, at least one interacting amino acid of the fluorescent protein that interacts with at least one first chemical moiety of the ligand, and selecting at least one chemical modification of the first chemical moiety to produce a second chemical moiety with a structure to either decrease or increase an interaction between the interacting amino acid and the second chemical moiety compared to the interaction between the interacting amino acid and the first chemical moiety.

In another embodiment, this invention provides a computational method of modeling the three dimensional structure of a fluorescent protein comprising determining a three dimensional relationship between at least two atoms listed in the atomic coordinates of Figs. 5-1 to 5-28.

In another embodiment, this invention provides a device comprising a storage device and, stored in the device, at least 10 atomic coordinates selected from the

atomic coordinates listed in Figs. 5-1 to 5-28. In one embodiment, the storage device is a computer readable device that stores code that receives as input the atomic coordinates. In another embodiment, the computer readable device is a floppy disk or a hard drive.

5 In another embodiment this invention provides a nucleic acid molecule comprising a nucleotide sequence encoding a functional engineered fluorescent protein whose amino acid sequence is substantially identical to the amino acid sequence of *Aequorea* green fluorescent protein (SEQ ID NO:2) and which differs from SEQ ID NO:2 by at least one first substitution at position T203, wherein the
10 substitution selected from the group consisting H, Y, W or F, and at least one second substitution at position H148.

 In another aspect the present invention includes a method of determining the presence of an anion of interest in a sample, comprising the steps of introducing an engineered green fluorescent protein into a sample, said engineered green fluorescent
15 protein comprising an amino acid sequence substantially identical to the amino acid sequence of *Aequorea* green fluorescent protein (SEQ ID NO:2) and which differs from SEQ ID NO:2 by at least one first substitution at position T203, wherein the substitution selected from the group consisting H, Y, W or F, and determining the fluorescence of said engineered green fluorescent protein in said sample.

20 In another embodiment, the invention includes a functional engineered fluorescent protein whose amino acid sequence is substantially identical to the amino acid sequence of *Aequorea* green fluorescent protein (SEQ ID NO:2) and which differs from SEQ ID NO:2 by at least one first substitution at position T203, wherein the substitution selected from the group consisting H, Y, W or F, and at least one
25 second substitution at position H148, wherein said functional engineered fluorescent protein has a different fluorescent property than *Aequorea* green fluorescent protein.

 In another embodiment the invention includes a host cell comprising a functional engineered fluorescent protein whose amino acid sequence is substantially identical to the amino acid sequence of *Aequorea* green fluorescent protein (SEQ ID
30 NO:2) and which differs from SEQ ID NO:2 by at least one first substitution at position T203, wherein the substitution selected from the group consisting H, Y, W or

F, and at least one second substitution at position H148, wherein said functional engineered fluorescent protein has a different fluorescent property than *Aequorea* green fluorescent protein.

DETAILED DESCRIPTION OF THE INVENTION**I. DEFINITIONS**

Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by those of ordinary skill in the art to which this invention belongs. Although any methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention, the preferred methods and materials are described. For purposes of the present invention, the following terms are defined below.

“Binding pair” refers to two moieties (e.g. chemical or biochemical) that have an affinity for one another. Examples of binding pairs include antigen/antibodies, lectin/avidin, target polynucleotide/probe oligonucleotide, antibody/anti-antibody, receptor/ligand, enzyme/ligand and the like. “One member of a binding pair” refers to one moiety of the pair, such as an antigen or ligand.

“Nucleic acid” refers to a deoxyribonucleotide or ribonucleotide polymer in either single- or double-stranded form, and, unless otherwise limited, encompasses known analogs of natural nucleotides that can function in a similar manner as naturally occurring nucleotides. It will be understood that when a nucleic acid molecule is represented by a DNA sequence, this also includes RNA molecules having the corresponding RNA sequence in which “U” replaces “T.”

“Recombinant nucleic acid molecule” refers to a nucleic acid molecule which is not naturally occurring, and which comprises two nucleotide sequences which are not naturally joined together. Recombinant nucleic acid molecules are produced by artificial recombination, e.g., genetic engineering techniques or chemical synthesis.

Reference to a nucleotide sequence “encoding” a polypeptide means that the sequence, upon transcription and translation of mRNA, produces the polypeptide. This includes both the coding strand, whose nucleotide sequence is identical to mRNA and whose sequence is usually provided in the sequence listing, as well as its complementary strand, which is used as the template for transcription. As any person skilled in the art recognizes, this also includes all degenerate nucleotide sequences encoding the same amino acid sequence. Nucleotide sequences encoding a polypeptide include sequences containing introns.

"Expression control sequences" refers to nucleotide sequences that regulate the expression of a nucleotide sequence to which they are operatively linked. Expression control sequences are "operatively linked" to a nucleotide sequence when the expression control sequences control and regulate the transcription and, as appropriate, translation of the nucleotide sequence. Thus, expression control sequences can include appropriate promoters, enhancers, transcription terminators, a start codon (i.e., ATG) in front of a protein-encoding gene, splicing signals for introns, maintenance of the correct reading frame of that gene to permit proper translation of the mRNA, and stop codons.

10 "Naturally-occurring" as used herein, as applied to an object, refers to the fact that an object can be found in nature. For example, a polypeptide or polynucleotide sequence that is present in an organism (including viruses) that can be isolated from a source in nature and which has not been intentionally modified by man in the laboratory is naturally-occurring.

15 "Operably linked" refers to a juxtaposition wherein the components so described are in a relationship permitting them to function in their intended manner. A control sequence "operably linked" to a coding sequence is ligated in such a way that expression of the coding sequence is achieved under conditions compatible with the control sequences, such as when the appropriate molecules (e.g., inducers and polymerases) are bound to the control or regulatory sequence(s).

20 "Control sequence" refers to polynucleotide sequences which are necessary to effect the expression of coding and non-coding sequences to which they are ligated. The nature of such control sequences differs depending upon the host organism; in prokaryotes, such control sequences generally include promoter, ribosomal binding site, and transcription termination sequence; in eukaryotes, generally, such control sequences include promoters and transcription termination sequence. The term "control sequences" is intended to include, at a minimum, components whose presence can influence expression, and can also include additional components whose presence is advantageous, for example, leader sequences and fusion partner sequences.

30

"Isolated polynucleotide" refers a polynucleotide of genomic, cDNA, or synthetic origin or some combination there of, which by virtue of its origin the "isolated polynucleotide" (1) is not associated with the cell in which the "isolated polynucleotide" is found in nature, or (2) is operably linked to a polynucleotide which it is not linked to in nature.

"Polynucleotide" refers to a polymeric form of nucleotides of at least 10 bases in length, either ribonucleotides or deoxynucleotides or a modified form of either type of nucleotide. The term includes single and double stranded forms of DNA.

The term "probe" refers to a substance that specifically binds to another substance (a "target"). Probes include, for example, antibodies, nucleic acids, receptors and their ligands.

"Modulation" refers to the capacity to either enhance or inhibit a functional property of biological activity or process (e.g., enzyme activity or receptor binding); such enhancement or inhibition may be contingent on the occurrence of a specific event, such as activation of a signal transduction pathway, and/or may be manifest only in particular cell types.

The term "modulator" refers to a chemical (naturally occurring or non-naturally occurring), such as a synthetic molecule (e.g., nucleic acid, protein, non-peptide, or organic molecule), or an extract made from biological materials such as bacteria, plants, fungi, or animal (particularly mammalian) cells or tissues. Modulators can be evaluated for potential activity as inhibitors or activators (directly or indirectly) of a biological process or processes (e.g., agonist, partial antagonist, partial agonist, inverse agonist, antagonist, antineoplastic agents, cytotoxic agents, inhibitors of neoplastic transformation or cell proliferation, cell proliferation-promoting agents, and the like) by inclusion in screening assays described herein. The activity of a modulator may be known, unknown or partially known.

The term "test chemical" refers to a chemical to be tested by one or more screening method(s) of the invention as a putative modulator. A test chemical is usually not known to bind to the target of interest. The term "control test chemical" refers to a chemical known to bind to the target (e.g., a known agonist, antagonist,

partial agonist or inverse agonist). Usually, various predetermined concentrations of test chemicals are used for screening, such as .01 μM , .1 μM , 1.0 μM , and 10.0 μM .

The term "target" refers to a biochemical entity involved a biological process. Targets are typically proteins that play a useful role in the physiology or biology of an organism. A therapeutic chemical binds to target to alter or modulate its function. As
5 used herein targets can include cell surface receptors, G-proteins, kinases, ion channels, phospholipases and other proteins mentioned herein.

The term "label" refers to a composition detectable by spectroscopic, photochemical, biochemical, immunochemical, or chemical means. For example,
10 useful labels include ^{32}P , fluorescent dyes, fluorescent proteins, electron-dense reagents, enzymes (e.g., as commonly used in an ELISA), biotin, dioxigenin, or haptens and proteins for which antisera or monoclonal antibodies are available. For example, polypeptides of this invention can be made as detectable labels, by e.g., incorporating a them as into a polypeptide, and used to label antibodies specifically
15 reactive with the polypeptide. A label often generates a measurable signal, such as radioactivity, fluorescent light or enzyme activity, which can be used to quantitate the amount of bound label.

The term "nucleic acid probe" refers to a nucleic acid molecule that binds to a specific sequence or sub-sequence of another nucleic acid molecule. A probe is
20 preferably a nucleic acid molecule that binds through complementary base pairing to the full sequence or to a sub-sequence of a target nucleic acid. It will be understood that probes may bind target sequences lacking complete complementarity with the probe sequence depending upon the stringency of the hybridization conditions. Probes are preferably directly labelled as with isotopes, chromophores, lumiphores,
25 chromogens, fluorescent proteins, or indirectly labelled such as with biotin to which a streptavidin complex may later bind. By assaying for the presence or absence of the probe, one can detect the presence or absence of the select sequence or sub-sequence.

A "labeled nucleic acid probe" is a nucleic acid probe that is bound, either covalently, through a linker, or through ionic, van der Waals or hydrogen bonds to a
30 label such that the presence of the probe may be detected by detecting the presence of the label bound to the probe.

The terms "polypeptide" and "protein" refers to a polymer of amino acid residues. The terms apply to amino acid polymers in which one or more amino acid residue is an artificial chemical analogue of a corresponding naturally occurring amino acid, as well as to naturally occurring amino acid polymers. The term

5 "recombinant protein" refers to a protein that is produced by expression of a nucleotide sequence encoding the amino acid sequence of the protein from a recombinant DNA molecule.

The term "recombinant host cell" refers to a cell that comprises a recombinant nucleic acid molecule. Thus, for example, recombinant host cells can express genes

10 that are not found within the native (non-recombinant) form of the cell.

The terms "isolated" "purified" or "biologically pure" refer to material which is substantially or essentially free from components which normally accompany it as found in its native state. Purity and homogeneity are typically determined using analytical chemistry techniques such as polyacrylamide gel electrophoresis or high

15 performance liquid chromatography. A protein or nucleic acid molecule which is the predominant protein or nucleic acid species present in a preparation is substantially purified. Generally, an isolated protein or nucleic acid molecule will comprise more than 80% of all macromolecular species present in the preparation. Preferably, the protein is purified to represent greater than 90% of all macromolecular species

20 present. More preferably the protein is purified to greater than 95%, and most preferably the protein is purified to essential homogeneity, wherein other macromolecular species are not detected by conventional techniques.

The term "naturally-occurring" as applied to an object refers to the fact that an object can be found in nature. For example, a polypeptide or polynucleotide sequence

25 that is present in an organism (including viruses) that can be isolated from a source in nature and which has not been intentionally modified by man in the laboratory is naturally-occurring.

The term "antibody" refers to a polypeptide substantially encoded by an immunoglobulin gene or immunoglobulin genes, or fragments thereof, which

30 specifically bind and recognize an analyte (antigen). The recognized immunoglobulin genes include the kappa, lambda, alpha, gamma, delta, epsilon and mu constant region

genes, as well as the myriad immunoglobulin variable region genes. Antibodies exist, e.g., as intact immunoglobulins or as a number of well characterized fragments produced by digestion with various peptidases. This includes, e.g., Fab' and F(ab)'₂ fragments. The term "antibody," as used herein, also includes antibody fragments
5 either produced by the modification of whole antibodies or those synthesized *de novo* using recombinant DNA methodologies.

The term "immunoassay" refers to an assay that utilizes an antibody to specifically bind an analyte. The immunoassay is characterized by the use of specific binding properties of a particular antibody to isolate, target, and/or quantify the
10 analyte.

The term "identical" in the context of two nucleic acid or polypeptide sequences refers to the residues in the two sequences which are the same when aligned for maximum correspondence. When percentage of sequence identity is used in reference to proteins or peptides it is recognized that residue positions which are
15 not identical often differ by conservative amino acid substitutions, where amino acids residues are substituted for other amino acid residues with similar chemical properties (e.g. charge or hydrophobicity) and therefore do not change the functional properties of the molecule. Where sequences differ in conservative substitutions, the percent sequence identity may be adjusted upwards to correct for the conservative nature of
20 the substitution. Means for making this adjustment are well known to those of skill in the art. Typically this involves scoring a conservative substitution as a partial rather than a full mismatch, thereby increasing the percentage sequence identity. Thus, for example, where an identical amino acid is given a score of 1 and a non-conservative substitution is given a score of zero, a conservative substitution is given a score
25 between zero and 1. The scoring of conservative substitutions is calculated, e.g., according to known algorithm. See, e.g., Meyers and Miller, *Computer Applic. Biol. Sci.*, 4: 11-17 (1988); Smith and Waterman (1981) *Adv. Appl. Math.* 2: 482; Needleman and Wunsch (1970) *J. Mol. Biol.* 48: 443; Pearson and Lipman (1988) *Proc. Natl. Acad. Sci. USA* 85: 2444; Higgins and Sharp (1988) *Gene*, 73: 237-244
30 and Higgins and Sharp (1989) *CABIOS* 5: 151-153; Corpet, *et al.* (1988) *Nucleic Acids Research* 16, 10881-90; Huang, *et al.* (1992) *Computer Applications in the*

Biosciences 8, 155-65, and Pearson, *et al.* (1994) *Methods in Molecular Biology* 24, 307-31. Alignment is also often performed by inspection and manual alignment.

"Conservatively modified variations" of a particular nucleic acid sequence refers to those nucleic acids which encode identical or essentially identical amino acid sequences, or where the nucleic acid does not encode an amino acid sequence, to essentially identical sequences. Because of the degeneracy of the genetic code, a large number of functionally identical nucleic acids encode any given polypeptide. For instance, the codons CGU, CGC, CGA, CGG, AGA, and AGG all encode the amino acid arginine. Thus, at every position where an arginine is specified by a codon, the codon can be altered to any of the corresponding codons described without altering the encoded polypeptide. Such nucleic acid variations are "silent variations," which are one species of "conservatively modified variations." Every nucleic acid sequence herein which encodes a polypeptide also describes every possible silent variation. One of skill will recognize that each codon in a nucleic acid (except AUG, which is ordinarily the only codon for methionine) can be modified to yield a functionally identical molecule by standard techniques. Accordingly, each "silent variation" of a nucleic acid which encodes a polypeptide is implicit in each described sequence. Furthermore, one of skill will recognize that individual substitutions, deletions or additions which alter, add or delete a single amino acid or a small percentage of amino acids (typically less than 5%, more typically less than 1%) in an encoded sequence are "conservatively modified variations" where the alterations result in the substitution of an amino acid with a chemically similar amino acid. Conservative amino acid substitutions providing functionally similar amino acids are well known in the art. The following six groups each contain amino acids that are conservative substitutions for one another:

- 1) Alanine (A), Serine (S), Threonine (T);
- 2) Aspartic acid (D), Glutamic acid (E);
- 3) Asparagine (N), Glutamine (Q);
- 4) Arginine (R), Lysine (K);
- 5) Isoleucine (I), Leucine (L), Methionine (M), Valine (V); and
- 6) Phenylalanine (F), Tyrosine (Y), Tryptophan (W).

The term "complementary" means that one nucleic acid molecule has the sequence of the binding partner of another nucleic acid molecule. Thus, the sequence 5'-ATGC-3' is complementary to the sequence 5'-GCAT-3'.

An amino acid sequence or a nucleotide sequence is "substantially identical" or "substantially similar" to a reference sequence if the amino acid sequence or nucleotide sequence has at least 80% sequence identity with the reference sequence over a given comparison window. Thus, substantially similar sequences include those having, for example, at least 85% sequence identity, at least 90% sequence identity, at least 95% sequence identity or at least 99% sequence identity. Two sequences that are identical to each other are, of course, also substantially identical.

A subject nucleotide sequence is "substantially complementary" to a reference nucleotide sequence if the complement of the subject nucleotide sequence is substantially identical to the reference nucleotide sequence.

The term "stringent conditions" refers to a temperature and ionic conditions used in nucleic acid hybridization. Stringent conditions are sequence dependent and are different under different environmental parameters. Generally, stringent conditions are selected to be about 5°C to 20°C lower than the thermal melting point (T_m) for the specific sequence at a defined ionic strength and pH. The T_m is the temperature (under defined ionic strength and pH) at which 50% of the target sequence hybridizes to a perfectly matched probe.

The term "allelic variants" refers to polymorphic forms of a gene at a particular genetic locus, as well as cDNAs derived from mRNA transcripts of the genes and the polypeptides encoded by them.

The term "preferred mammalian codon" refers to the subset of codons from among the set of codons encoding an amino acid that are most frequently used in proteins expressed in mammalian cells as chosen from the following list:

	Amino Acid	Preferred codons for high level mammalian expression
	Gly	GGC,GGG
	Glu	GAG
	Asp	GAC
5	Val	GUG,GUC
	Ala	GCC,GCU
	Ser	AGC,UCC
	Lys	AAG
	Asn	AAC
10	Met	AUG
	Ile	AUC
	Thr	ACC
	Trp	UGG
	Cys	UGC
15	Tyr	UAU,UAC
	Leu	CUG
	Phe	UUC
	Arg	CGC,AGG,AGA
	Gln	CAG
20	His	CAC
	Pro	CCC

Fluorescent molecules are useful in fluorescence resonance energy transfer ("FRET"). FRET involves a donor molecule and an acceptor molecule. To optimize the efficiency and detectability of FRET between a donor and acceptor molecule, several factors need to be balanced. The emission spectrum of the donor should overlap as much as possible with the excitation spectrum of the acceptor to maximize the overlap integral. Also, the quantum yield of the donor moiety and the extinction coefficient of the acceptor should likewise be as high as possible to maximize R_0 , the distance at which energy transfer efficiency is 50%. However, the excitation spectra of the donor and acceptor should overlap as little as possible so that a wavelength

region can be found at which the donor can be excited efficiently without directly exciting the acceptor. Fluorescence arising from direct excitation of the acceptor is difficult to distinguish from fluorescence arising from FRET. Similarly, the emission spectra of the donor and acceptor should overlap as little as possible so that the two emissions can be clearly distinguished. High fluorescence quantum yield of the acceptor moiety is desirable if the emission from the acceptor is to be measured either as the sole readout or as part of an emission ratio. One factor to be considered in choosing the donor and acceptor pair is the efficiency of fluorescence resonance energy transfer between them. Preferably, the efficiency of FRET between the donor and acceptor is at least 10%, more preferably at least 50% and even more preferably at least 80%.

The term "fluorescent property" refers to the molar extinction coefficient at an appropriate excitation wavelength, the fluorescence quantum efficiency, the shape of the excitation spectrum or emission spectrum, the excitation wavelength maximum and emission wavelength maximum, the ratio of excitation amplitudes at two different wavelengths, the ratio of emission amplitudes at two different wavelengths, the excited state lifetime, or the fluorescence anisotropy. A measurable difference in any one of these properties between wild-type *Aequorea* GFP and the mutant form is useful. A measurable difference can be determined by determining the amount of any quantitative fluorescent property, e.g., the amount of fluorescence at a particular wavelength, or the integral of fluorescence over the emission spectrum. Determining ratios of excitation amplitude or emission amplitude at two different wavelengths ("excitation amplitude ratioing" and "emission amplitude ratioing", respectively) are particularly advantageous because the ratioing process provides an internal reference and cancels out variations in the absolute brightness of the excitation source, the sensitivity of the detector, and light scattering or quenching by the sample.

II. LONG WAVELENGTH ENGINEERED FLUORESCENT PROTEINS

A. Fluorescent Proteins

As used herein, the term "fluorescent protein" refers to any protein capable of fluorescence when excited with appropriate electromagnetic radiation. This includes fluorescent proteins whose amino acid sequences are either naturally occurring or engineered (i.e., analogs or mutants). Many cnidarians use green fluorescent proteins ("GFPs") as energy-transfer acceptors in bioluminescence. A "green fluorescent protein," as used herein, is a protein that fluoresces green light. Similarly, "blue fluorescent proteins" fluoresce blue light and "red fluorescent proteins" fluoresce red light. GFPs have been isolated from the Pacific Northwest jellyfish, *Aequorea victoria*, the sea pansy, *Renilla reniformis*, and *Phialidium gregarium*. W.W. Ward et al., *Photochem. Photobiol.*, 35:803-808 (1982); L.D. Levine et al., *Comp. Biochem. Physiol.*, 72B:77-85 (1982).

A variety of *Aequorea*-related fluorescent proteins having useful excitation and emission spectra have been engineered by modifying the amino acid sequence of a naturally occurring GFP from *Aequorea victoria*. (D.C. Prasher et al., *Gene*, 111:229-233 (1992); R. Heim et al., *Proc. Natl. Acad. Sci., USA*, 91:12501-04 (1994); U.S. patent application 08/337,915, filed November 10, 1994; International application PCT/US95/14692, filed 11/10/95.)

As used herein, a fluorescent protein is an "*Aequorea*-related fluorescent protein" if any contiguous sequence of 150 amino acids of the fluorescent protein has at least 85% sequence identity with an amino acid sequence, either contiguous or non-contiguous, from the 238 amino-acid wild-type *Aequorea* green fluorescent protein of Fig. 3 (SEQ ID NO:2). More preferably, a fluorescent protein is an *Aequorea*-related fluorescent protein if any contiguous sequence of 200 amino acids of the fluorescent protein has at least 95% sequence identity with an amino acid sequence, either contiguous or non-contiguous, from the wild type *Aequorea* green fluorescent protein of Fig. 3 (SEQ ID NO:2). Similarly, the fluorescent protein may be related to *Renilla* or *Phialidium* wild-type fluorescent proteins using the same standards.

Aequorea-related fluorescent proteins include, for example and without limitation, wild-type (native) *Aequorea victoria* GFP (D.C. Prasher et al., "Primary

structure of the *Aequorea victoria* green fluorescent protein," *Gene*, (1992) 111:229-33), whose nucleotide sequence (SEQ ID NO:1) and deduced amino acid sequence (SEQ ID NO:2) are presented in Fig. 3; allelic variants of this sequence, e.g., Q80R, which has the glutamine residue at position 80 substituted with arginine (M. Chalfie et al., *Science*, (1994) 263:802-805); those engineered *Aequorea*-related fluorescent proteins described herein, e.g., in Table A or Table F, variants that include one or more folding mutations and fragments of these proteins that are fluorescent, such as *Aequorea* green fluorescent protein from which the two amino-terminal amino acids have been removed. Several of these contain different aromatic amino acids within the central chromophore and fluoresce at a distinctly shorter wavelength than wild type species. For example, engineered proteins P4 and P4-3 contain (in addition to other mutations) the substitution Y66H, whereas W2 and W7 contain (in addition to other mutations) Y66W. Other mutations both close to the chromophore region of the protein and remote from it in primary sequence may affect the spectral properties of GFP and are listed in the first part of the table below.

TABLE A

Clone	Mutation(s)	Excitation max (nm)	Emission max (nm)	Extinct. Coeff. (M ⁻¹ cm ⁻¹)	Quantum yield
Wild type	None	395 (475)	508	21,000 (7,150)	0.77
P4	Y66H	383	447	13,500	0.21
P4-3	Y66H Y145F	381	445	14,000	0.38
P4-3E	Y66H Y145F V163A	384	448	22,000	0.27
W7	Y66W N146I M153T V163A N212K	433 (453)	475 (501)	18,000 (17,100)	0.67
W2	Y66W I123V Y145H H148R M153T V163A N212K	432 (453)	480	10,000 (9,600)	0.72

W1C	S65A Y66W S72A N146I M153T V163A	435	495	21,200	0.39
W1B	F64L S65T Y66W N146I M153T V163A	434 (452)	476 (505)	32,500	0.4
S65T	S65T	489	511	39,200	0.68
P4-1	S65T M153A K238E	504 (396)	514	14,500 (8,600)	0.53
Emerald	S65T, S72A, N149K, M153T, I167T	487	509	57,500	0.68
EGFP	F64L, S65T	488	507	55,900	0.64
S65A	S65A	471	504		
S65C	S65C	479	507		
S65L	S65L	484	510		
Y66F	Y66F	360	442		
Y66W	Y66W	458	480		
Topaz	S65G S72A K79R T203Y	514	527	94,500	0.6
10C YFP	S65G V68L S72A T203Y	514	527	83,400	0.61
Sapphire	S72A, Y145F T203I	399	511	29,000	0.64

Additional mutations in *Aequorea*-related fluorescent proteins, referred to as "folding mutations," improve the ability of fluorescent proteins to fold at higher temperatures, and to be more fluorescent when expressed in mammalian cells, but have little or no effect on the peak wavelengths of excitation and emission. It should be noted that these may be combined with mutations that influence the spectral properties of GFP to produce proteins with altered spectral and folding properties. Folding mutations include: F64L, V68L, S72A, and also T44A, F99S, Y145F, N146I, M153T or A, V163A, I167T, S175G, S205T and N212K.

As used herein, the term "loop domain" refers to an amino acid sequence of an *Aequorea*-related fluorescent protein that connects the amino acids involved in the secondary structure of the eleven strands of the β -barrel or the central α -helix (residues 56-72) (see Fig. 1A and 1B).

As used herein, the "fluorescent protein moiety" of a fluorescent protein is that portion of the amino acid sequence of a fluorescent protein which, when the amino acid sequence of the fluorescent protein substrate is optimally aligned with the amino acid sequence of a naturally occurring fluorescent protein, lies between the amino terminal and carboxy terminal amino acids, inclusive, of the amino acid sequence of the naturally occurring fluorescent protein.

It has been found that fluorescent proteins can be genetically fused to other target proteins and used as markers to identify the location and amount of the target protein produced. Accordingly, this invention provides fusion proteins comprising a fluorescent protein moiety and additional amino acid sequences. Such sequences can be, for example, up to about 15, up to about 50, up to about 150 or up to about 1000 amino acids long. The fusion proteins possess the ability to fluoresce when excited by electromagnetic radiation. In one embodiment, the fusion protein comprises a polyhistidine tag to aid in purification of the protein.

B. Use Of The Crystal Structure Of Green Fluorescent Protein To Design Mutants Having Altered Fluorescent Characteristics

Using X-ray crystallography and computer processing, we have created a
5 model of the crystal structure of *Aequorea* green fluorescent protein showing the relative location of the atoms in the molecule. This information is useful in identifying amino acids whose substitution alters fluorescent properties of the protein.

Fluorescent characteristics of *Aequorea*-related fluorescent proteins depend, in part, on the electronic environment of the chromophore. In general, amino acids that
10 are within about 0.5 nm of the chromophore influence the electronic environment of the chromophore. Therefore, substitution of such amino acids can produce fluorescent proteins with altered fluorescent characteristics. In the excited state, electron density tends to shift from the phenolate towards the carbonyl end of the chromophore. Therefore, placement of increasing positive charge near the carbonyl
15 end of the chromophore tends to decrease the energy of the excited state and cause a red-shift in the absorbance and emission wavelength maximum of the protein. Decreasing positive charge near the carbonyl end of the chromophore tends to have the opposite effect, causing a blue-shift in the protein's wavelengths.

Amino acids with charged (ionized D, E, K, and R), dipolar (H, N, Q, S, T,
20 and uncharged D, E and K), and polarizable side groups (e.g., C, F, H, M, W and Y) are useful for altering the electronic environment of the chromophore, especially when they substitute an amino acid with an uncharged, nonpolar or non-polarizable side chain. In general, amino acids with polarizable side groups alter the electronic environment least, and, consequently, are expected to cause a comparatively smaller
25 change in a fluorescent property. Amino acids with charged side groups alter the environment most, and, consequently, are expected to cause a comparatively larger change in a fluorescent property. However, amino acids with charged side groups are more likely to disrupt the structure of the protein and to prevent proper folding if buried next to the chromophore without any additional solvation or salt bridging.
30 Therefore charged amino acids are most likely to be tolerated and to give useful effects when they replace other charged or highly polar amino acids that are already

solvated or involved in salt bridges. In certain cases, where substitution with a polarizable amino acid is chosen, the structure of the protein may make selection of a larger amino acid, e.g., W, less appropriate. Alternatively, positions occupied by amino acids with charged or polar side groups that are unfavorably oriented may be substituted with amino acids that have less charged or polar side groups. In another alternative, an amino acid whose side group has a dipole oriented in one direction in the protein can be substituted with an amino acid having a dipole oriented in a different direction.

More particularly, Table B lists several amino acids located within about 0.5 nm from the chromophore whose substitution can result in altered fluorescent characteristics. The table indicates, underlined, preferred amino acid substitutions at the indicated location to alter a fluorescent characteristic of the protein. In order to introduce such substitutions, the table also provides codons for primers used in site-directed mutagenesis involving amplification. These primers have been selected to encode economically the preferred amino acids, but they encode other amino acids as well, as indicated, or even a stop codon, denoted by Z. In introducing substitutions using such degenerate primers the most efficient strategy is to screen the collection to identify mutants with the desired properties and then sequence their DNA to find out which of the possible substitutions is responsible. Codons are shown in double-stranded form with sense strand above, antisense strand below. In nucleic acid sequences, R=(A or g); Y=(C or T); M=(A or C); K=(g or T); S=(g or C); W=(A or T); H=(A, T, or C); B=(g, T, or C); V=(g, A, or C); D=(g, A, or T); N=(A, C, g, or T).

TABLE B

Original position and presumed role	Change to	Codon
L42 Aliphatic residue near C=N of chromophore	<u>CFHLQRWYZ</u>	5'YDS 3' 3'RHS 5'
V61 Aliphatic residue near central -CH= of chromophore	<u>FYHCLR</u>	YDC RHg

	T62	Almost directly above center of chromophore bridge	<u>AVFS</u>	KYC MRg
5			<u>DEHKNO</u>	VAS BTS
			<u>FYHCLR</u>	YDC RHg
10	V68	Aliphatic residue near carbonyl and G67	<u>FYHL</u>	YWC RWg
	N121	Near C-N site of ring closure between T65 and G67	<u>CFHLQRWYZ</u>	YDS RHS
15	Y145	Packs near tyrosine ring of chromophore	<u>WCFL</u>	TKS AMS
			<u>DEHNKO</u>	VAS BTS
20	H148	H-bonds to phenolate oxygen	<u>FYNI</u>	WWC WWg
			<u>KQR</u>	MRg KYC
25	V150	Aliphatic residue near tyrosine ring of chromophore	<u>FYHL</u>	YWC RWg
30	F165	Packs near tyrosine ring	<u>CHQRWYZ</u>	YRS RYS
	I167	Aliphatic residue near phenolate; I167T has effects	<u>FYHL</u>	YWC RWg
35	T203	H-bonds to phenolic oxygen of chromophore	<u>FHLQRWYZ</u>	YDS RHS
40	E222	Protonation regulates ionization of chromophore	<u>HKNQ</u>	MAS KTS

Examples of amino acids with polar side groups that can be substituted with polarizable side groups include, for example, those in Table C.

32

TABLE C

	Original position and presumed role	Change to	Codon
5	Q69 Terminates chain of H-bonding waters	<u>KREG</u>	RRg YYC
	Q94 H-bonds to carbonyl terminus of chromophore	<u>DEHKNQ</u>	VAS BTS
10	Q183 Bridges Arg96 and center of chromophore bridge	<u>HY</u>	YAC RTG
		<u>EK</u>	RAg YTC
15	N185 Part of H-bond network near carbonyl of chromophore	<u>DEHNKQ</u>	VAS BTS

20 In another embodiment, an amino acid that is close to a second amino acid within about 0.5 nm of the chromophore can, upon substitution, alter the electronic properties of the second amino acid, in turn altering the electronic environment of the chromophore. Table D presents two such amino acids. The amino acids, L220 and V224, are close to E222 and oriented in the same direction in the β pleated sheet.

25

TABLE D

	Original position and presumed role	Change to	Codon
30	L220 Packs next to Glu222; to make GFP pH sensitive	<u>HKNPQT</u>	MMS KKS
	V224 Packs next to Glu222; to make GFP pH sensitive	<u>HKNPQT</u>	MMS KKS
35		<u>CFHLQRWYZ</u>	YDS RHS

One embodiment of the invention includes a nucleic acid molecule comprising a nucleotide sequence encoding a functional engineered fluorescent protein whose amino acid sequence is substantially identical to the amino acid sequence of *Aequorea* green fluorescent protein (SEQ ID NO:2) and which differs from SEQ ID NO:2 by at least a substitution at Q69, wherein the functional engineered fluorescent protein has a different fluorescent property than *Aequorea* green fluorescent protein. Preferably, the substitution at Q69 is selected from the group of K, R, E and G. The Q69 substitution can be combined with other mutations to improve the properties of the protein, such as a functional mutation at S65.

One embodiment of the invention includes a nucleic acid molecule comprising a nucleotide sequence encoding a functional engineered fluorescent protein whose amino acid sequence is substantially identical to the amino acid sequence of *Aequorea* green fluorescent protein (SEQ ID NO:2) and which differs from SEQ ID NO:2 by at least a substitution at E222, but not including E222G, wherein the functional engineered fluorescent protein has a different fluorescent property than *Aequorea* green fluorescent protein. Preferably, the substitution at E222 is selected from the group of N and Q. The E222 substitution can be combined with other mutations to improve the properties of the protein, such as a functional mutation at F64.

One embodiment of the invention includes a nucleic acid molecule comprising a nucleotide sequence encoding a functional engineered fluorescent protein whose amino acid sequence is substantially identical to the amino acid sequence of *Aequorea* green fluorescent protein (SEQ ID NO:2) and which differs from SEQ ID NO:2 by at least a substitution at Y145, wherein the functional engineered fluorescent protein has a different fluorescent property than *Aequorea* green fluorescent protein. Preferably, the substitution at Y145 is selected from the group of W, C, F, L, E, H, K and Q. The Y145 substitution can be combined with other mutations to improve the properties of the protein, such as a Y66.

The invention also includes computer related embodiments, including computational methods of using the crystal coordinates for designing new fluorescent protein mutations and devices for storing the crystal data, including coordinates. For instance the invention includes a device comprising a storage device and, stored in the device, at least 10 atomic coordinates selected from the atomic coordinates listed in Figs. 5-1 to 5-28. More coordinates can be stored depending of the complexity of the calculations or the objective of using the coordinates (e.g. about 100, 1,000, or more coordinates). For example, larger numbers of coordinates will be desirable for more detailed representations of fluorescent protein structure. Typically, the storage device is a computer readable device that stores code that it receives as input the atomic coordinates. Although, other storage means as known in the art are contemplated. The computer readable device can be a floppy disk or a hard drive.

C. Use Of The Crystal Structure Of YFP To Design Mutants Having Altered Anion Binding Characteristics

In another aspect the invention includes the use of X-ray crystallography and computer processing, to create a model of the crystal structure of YFP showing the relative location, and amino acids that interact with bound ions. This information is useful in identifying amino acids whose substitution alters the specificity and affinity of the binding site to various anions. Because the binding of the anion is close to the chromophore of YFP, binding results in a modulation of the fluorescent properties of YFP that can be used to monitor anion binding and therefore the concentration of the anion.

The anion binding site found in YFP-H148Q exhibits many of the characteristics generally found in halide binding sites in other proteins. In the case of the anion-containing cavity in YFP-H148Q, the binding site is amphiphilic in nature, with one side lined with polar and charged groups (Tyr203, the chromophore, Arg96, Gln69, and Gln183), and the other with hydrophobic residues (Ile152, Leu201, Val163, Val150, and Phe165).

The design of engineered fluorescent proteins with altered anion binding specificities requires consideration of a number of factors. For example, one of the most significant factors contributing to the anion affinity and selectivity is the electrostatic configuration and make up of the binding pocket. In YFP these include
 5 the groups listed in Table E below.

TABLE E

Original position and presumed role		
	S65, Y66, G67	Forms chromophore, aromatic edge interaction with ion
	Q69	Hydrogen bonds to ion
10	R96	Charge interaction with ion
	Q183	Charge interaction with ion
	Y203	Hydrogen bonds to ion (Y)

In general, anion binding can be improved by creating more and or tighter
 15 binding interactions between the anion of interest and polar groups within the binding pocket. For example either directly substituting the polar residues above with more polar residues, or by substituting residues of different sizes, that may interact more effectively with the anion, can improve ion binding. For example the size and position of the chromophore may be altered by the substitution of S65 to G, A, C, V, L, I or T;
 20 Y66 may be altered by substitution to H, F or W; Q69 may be substituted to N or K; R96 to K; Q183 to N or K.

Hydration energy

Additionally the binding of an anion in a buried cavity near the chromophore requires replacement of ion-solvent interactions with ion-protein interactions.
 25 Relative binding energies of monovalent anions to YFP (Table J) and YFP-H148Q (Jayaraman *et al.*, 2000) in relation to their hydration energy indicate that hydration forces make important contributions towards binding. In the following series of

monoanions, the hydration energies are ordered from weak to strong: $\text{CO}_4^- < \text{I}^- < \text{NO}_3^- < \text{SCN}^- < \text{Br}^- < \text{Cl}^- < \text{F}^-$ (Wright & Diamond, 1977). Polyatomic monoanions and iodide have relatively weak hydration energies, whereas the other halides interact more strongly with water. In case of the spherically symmetric halides, hydration energy increases with decreasing atomic volume (Born, 1920), which is why larger halides are easier to bury in the more hydrophobic environment of a protein's interior. The trend observed for anion binding to the YFPs roughly follows the above series (Table J). Protein interaction generally increases with decreasing hydration energy, with the exception of fluoride, which may not completely dehydrate upon protein binding due to its small size.

The development of higher affinity anion binding sites therefore requires the creation of sufficient ion-protein interactions for example by the substitution of hydrophobic residues that line the ion binding pocket with more polar residues with more hydrogen bonding potential. Examples for these type of substitutions for improving the ion binding for larger and smaller anions are presented in Table F

TABLE F

i) Mutation of amino acids around the ion binding pocket to increase binding affinity for smaller anions than iodide.

	Original position and presumed role		Change to
20	V150	Lines binding pocket	S, T, Q, N
	I152	Lines binding pocket	L, V, F, S, T, Q, N
	V163	Lines binding pocket	S, T, Q, N
	F165	Lines binding pocket	Y, W
	H181	Lines binding pocket	F, W
25	Q183	Lines binding pocket	K, R, N
	L201	Lines binding pocket	S, T, Q, N, V, I

- ii) Mutation of amino acids around the ion binding pocket to increase binding affinity for larger anions than iodide.

	V150	Lines binding pocket	A, C, M, G, S, L
5	I152	Lines binding pocket	A, C, M, G, S
	V163	Lines binding pocket	A, C, M, G, S, L
	F165	Lines binding pocket	Y, L
	H181	Lines binding pocket	K, R
	Q183	Lines binding pocket	N, S, C
10	L201	Lines binding pocket	A, C, M, G, S

Size of the binding pocket

The size and shape of the binding pocket may also be of particular importance due to the buried nature of the binding site for larger anions. TCA, with a mean
 15 geometric diameter of 6.2 Å (Halm & Frizzell, 1992), is apparently too large to interact with YFP to a measurable extent, whereas the somewhat smaller TFA does show weak binding (Table J). Improvements in the binding affinity of larger anions could thus be achieved via the substitution of amino acids lining the binding pocket with smaller residues, as outlined in Table F above, as well as increasing solvent
 20 accessibility as discussed below.

Conformational changes upon anion binding

A series of conformational changes of various side chains lining the binding pocket in YFP are necessary for halide binding. The largest movements are observed
 25 for Gln69, and Gln183 although the apolar side chains of Leu201, Ile152, Val150, and Val163 (Fig. 11) all undergo movements to increase the cavity size in the presence of a bound halide. Another approach towards tighter anion binding therefore is the

substitution of the residues that undergo the most dramatic conformational change upon binding, for smaller residues. These changes may reduce the need for structural rearrangements upon binding thereby making anion binding, more energetically favorable. These changes include those listed in Table F above as well as the
5 substitution of Q69 for N.

Solvent Accessibility

The results from the structural determinations of various mutations at His148 suggests that specific mutations at this position can result in overall structural
10 adjustments in the beta barrel that can directly affect both solvent accessibility and the volume of the binding pocket. Substitution of His148 for example to smaller amino acids such Q, N, G, A, L, V and I would therefore be predicted to increase solvent access to the chromophore and therefore improve binding of larger anions. Likewise substitution of His 148 with larger amino acids such as F or W would be likely to
15 reduce anion access to the chromophore. Similarly more subtle changes could be achieved by substituting positions 147 and 149 with smaller or larger amino acids.

These mutations will typically be introduced in the YFP template protein via oligo-mediated site directed mutagenesis to create libraries of mutant proteins that typically have a 10 % probability of containing the wild-type amino acid residue and a
20 90% probability of containing one of the various mutant residues. Using this approach it is possible to rapidly screen libraries containing various combinations of mutants to identify the best combinations for a specific anion of interest. Typically this process can be repeated iteratively to ensure that sequence space around the binding pocket has been completely explored for any specific anion of interest.

D. Production Of Engineered Fluorescent Proteins

Recombinant production of a fluorescent protein involves expressing a nucleic acid molecule having sequences that encode the protein.

In one embodiment, the nucleic acid encodes a fusion protein in which a single
5 polypeptide includes the fluorescent protein moiety within a longer polypeptide. The longer polypeptide can include a second functional protein, such as FRET partner or a protein having a second function (e.g., an enzyme, antibody or other binding protein). Nucleic acids that encode fluorescent proteins are useful as starting materials.

The fluorescent proteins can be produced as fusion proteins by recombinant
10 DNA technology. Recombinant production of fluorescent proteins involves expressing nucleic acids having sequences that encode the proteins. Nucleic acids encoding fluorescent proteins can be obtained by methods known in the art. Fluorescent proteins can be made by site-specific mutagenesis of other nucleic acids encoding fluorescent proteins, or by random mutagenesis caused by increasing the
15 error rate of PCR of the original polynucleotide with 0.1 mM MnCl₂ and unbalanced nucleotide concentrations. See, e.g., U.S. patent application 08/337,915, filed November 10, 1994 or International application PCT/US95/14692, filed 11/10/95. The nucleic acid encoding a green fluorescent protein can be isolated by polymerase chain reaction of cDNA from *A. victoria* using primers based on the DNA sequence
20 of *A. victoria* green fluorescent protein, as presented in Fig. 3. PCR methods are described in, for example, U.S. Pat. No. 4,683,195; Mullis et al. (1987) *Cold Spring Harbor Symp. Quant. Biol.* 51:263; and Erlich, ed., *PCR Technology*, (Stockton Press, NY, 1989).

The construction of expression vectors and the expression of genes in
25 transfected cells involves the use of molecular cloning techniques also well known in the art. Sambrook et al., *Molecular Cloning -- A Laboratory Manual*, Cold Spring Harbor Laboratory, Cold Spring Harbor, NY, (1989) and *Current Protocols in Molecular Biology*, F.M. Ausubel et al., eds., (Current Protocols, a joint venture between Greene Publishing Associates, Inc. and John Wiley & Sons, Inc.). The
30 expression vector can be adapted for function in prokaryotes or eukaryotes by inclusion of appropriate promoters, replication sequences, markers, etc.

Nucleic acids used to transfect cells with sequences coding for expression of the polypeptide of interest generally will be in the form of an expression vector including expression control sequences operatively linked to a nucleotide sequence coding for expression of the polypeptide. As used, the term "nucleotide sequence
5 coding for expression of" a polypeptide refers to a sequence that, upon transcription and translation of mRNA, produces the polypeptide. This can include sequences containing, *e.g.*, introns. Expression control sequences are operatively linked to a nucleic acid sequence when the expression control sequences control and regulate the transcription and, as appropriate, translation of the nucleic acid sequence. Thus,
10 expression control sequences can include appropriate promoters, enhancers, transcription terminators, a start codon (*i.e.*, ATG) in front of a protein-encoding gene, splicing signals for introns, maintenance of the correct reading frame of that gene to permit proper translation of the mRNA, and stop codons.

Methods which are well known to those skilled in the art can be used to
15 construct expression vectors containing the fluorescent protein coding sequence and appropriate transcriptional/translational control signals. These methods include *in vitro* recombinant DNA techniques, synthetic techniques and *in vivo* recombination/genetic recombination. (See, for example, the techniques described in Maniatis, *et al.*, *Molecular Cloning A Laboratory Manual*, Cold Spring Harbor
20 Laboratory, N.Y., 1989).

Transformation of a host cell with recombinant DNA may be carried out by conventional techniques as are well known to those skilled in the art. Where the host is prokaryotic, such as *E. coli*, competent cells which are capable of DNA uptake can be prepared from cells harvested after exponential growth phase and subsequently
25 treated by the CaCl_2 method by procedures well known in the art. Alternatively, MgCl_2 or RbCl can be used. Transformation can also be performed after forming a protoplast of the host cell or by electroporation.

When the host is a eukaryote, such methods of transfection of DNA as calcium phosphate co-precipitates, conventional mechanical procedures such as
30 microinjection, electroporation, insertion of a plasmid encased in liposomes, or virus vectors may be used. Eukaryotic cells can also be co-transfected with DNA

sequences encoding the fusion polypeptide of the invention, and a second foreign DNA molecule encoding a selectable phenotype, such as the herpes simplex thymidine kinase gene. Another method is to use a eukaryotic viral vector, such as simian virus 40 (SV40) or bovine papilloma virus, to transiently infect or transform eukaryotic cells and express the protein. (*Eukaryotic Viral Vectors*, Cold Spring Harbor Laboratory, Gluzman ed., 1982). Preferably, a eukaryotic host is utilized as the host cell as described herein.

Techniques for the isolation and purification of either microbially or eukaryotically expressed polypeptides of the invention may be by any conventional means such as, for example, preparative chromatographic separations and immunological separations such as those involving the use of monoclonal or polyclonal antibodies or antigen. In one embodiment recombinant fluorescent proteins can be produced by expression of nucleic acid encoding for the protein in *E. coli*. *Aequorea*-related fluorescent proteins are best expressed by cells cultured between about 15°C and 30°C but higher temperatures (e.g. 37°C) are possible. After synthesis, these enzymes are stable at higher temperatures (e.g., 37°C) and can be used in assays at those temperatures.

A variety of host-expression vector systems may be utilized to express fluorescent protein coding sequence. These include but are not limited to microorganisms such as bacteria transformed with recombinant bacteriophage DNA, plasmid DNA or cosmid DNA expression vectors containing a fluorescent protein coding sequence; yeast transformed with recombinant yeast expression vectors containing the fluorescent protein coding sequence; plant cell systems infected with recombinant virus expression vectors (e.g., cauliflower mosaic virus, CaMV; tobacco mosaic virus, TMV) or transformed with recombinant plasmid expression vectors (e.g., Ti plasmid) containing a fluorescent protein coding sequence; insect cell systems infected with recombinant virus expression vectors (e.g., baculovirus) containing a fluorescent protein coding sequence; or animal cell systems infected with recombinant virus expression vectors (e.g., retroviruses, adenovirus, vaccinia virus) containing a fluorescent protein coding sequence, or transformed animal cell systems engineered for stable expression.

Depending on the host/vector system utilized, any of a number of suitable transcription and translation elements, including constitutive and inducible promoters, transcription enhancer elements, transcription terminators, *etc.* may be used in the expression vector (see, *e.g.*, Bitter, *et al.*, Methods in Enzymology 153:516-544, 1987). For example, when cloning in bacterial systems, inducible promoters such as pL of bacteriophage Σ , plac, ptrp, ptac (ptrp-lac hybrid promoter) and the like may be used. When cloning in mammalian cell systems, promoters derived from the genome of mammalian cells (*e.g.*, metallothionein promoter) or from mammalian viruses (*e.g.*, the retrovirus long terminal repeat; the adenovirus late promoter; the vaccinia virus 7.5K promoter) may be used. Promoters produced by recombinant DNA or synthetic techniques may also be used to provide for transcription of the inserted fluorescent protein coding sequence.

In bacterial systems a number of expression vectors may be advantageously selected depending upon the use intended for the fluorescent protein expressed. For example, when large quantities of the fluorescent protein are to be produced, vectors which direct the expression of high levels of fusion protein products that are readily purified may be desirable. Those which are engineered to contain a cleavage site to aid in recovering fluorescent protein are preferred.

In yeast, a number of vectors containing constitutive or inducible promoters may be used. For a review see, Current Protocols in Molecular Biology, Vol. 2, Ed. Ausubel, *et al.*, Greene Publish. Assoc. & Wiley Interscience, Ch. 13, 1988; Grant, *et al.*, Expression and Secretion Vectors for Yeast, in Methods in Enzymology, Eds. Wu & Grossman, 31987, Acad. Press, N.Y., Vol. 153, pp.516-544, 1987; Glover, DNA Cloning, Vol. II, IRL Press, Wash., D.C., Ch. 3, 1986; and Bitter, Heterologous Gene Expression in Yeast, Methods in Enzymology, Eds. Berger & Kimmel, Acad. Press, N.Y., Vol. 152, pp. 673-684, 1987; and The Molecular Biology of the Yeast *Saccharomyces*, Eds. Strathern *et al.*, Cold Spring Harbor Press, Vols. I and II, 1982. A constitutive yeast promoter such as ADH or LEU2 or an inducible promoter such as GAL may be used (Cloning in Yeast, Ch. 3, R. Rothstein In: DNA Cloning Vol.11, A Practical Approach, Ed. DM Glover, IRL Press, Wash., D.C., 1986). Alternatively,

vectors may be used which promote integration of foreign DNA sequences into the yeast chromosome.

In cases where plant expression vectors are used, the expression of a fluorescent protein coding sequence may be driven by any of a number of promoters. For example, viral promoters such as the 35S RNA and 19S RNA promoters of CaMV (Brisson, *et al.*, *Nature* 310:511-514, 1984), or the coat protein promoter to TMV (Takamatsu, *et al.*, *EMBO J.* 6:307-311, 1987) may be used; alternatively, plant promoters such as the small subunit of RUBISCO (Coruzzi, *et al.*, 1984, *EMBO J.* 3:1671-1680; Broglie, *et al.*, *Science* 224:838-843, 1984); or heat shock promoters, *e.g.*, soybean hsp17.5-E or hsp17.3-B (Gurley, *et al.*, *Mol. Cell. Biol.* 6:559-565, 1986) may be used. These constructs can be introduced into plant cells using Ti plasmids, Ri plasmids, plant virus vectors, direct DNA transformation, microinjection, electroporation, *etc.* For reviews of such techniques see, for example, Weissbach & Weissbach, *Methods for Plant Molecular Biology*, Academic Press, NY, Section VIII, pp. 421-463, 1988; and Grierson & Corey, *Plant Molecular Biology*, 2d Ed., Blackie, London, Ch. 7-9, 1988.

An alternative expression system which could be used to express fluorescent protein is an insect system. In one such system, *Autographa californica* nuclear polyhedrosis virus (AcNPV) is used as a vector to express foreign genes. The virus grows in *Spodoptera frugiperda* cells. The fluorescent protein coding sequence may be cloned into non-essential regions (for example, the polyhedrin gene) of the virus and placed under control of an AcNPV promoter (for example the polyhedrin promoter). Successful insertion of the fluorescent protein coding sequence will result in inactivation of the polyhedrin gene and production of non-occluded recombinant virus (*i.e.*, virus lacking the proteinaceous coat coded for by the polyhedrin gene). These recombinant viruses are then used to infect *Spodoptera frugiperda* cells in which the inserted gene is expressed, see Smith, *et al.*, *J. Virol.* 46:584, 1983; Smith, U.S. Patent No. 4,215,051.

Eukaryotic systems, and preferably mammalian expression systems, allow for proper post-translational modifications of expressed mammalian proteins to occur. Eukaryotic cells which possess the cellular machinery for proper processing of the

primary transcript, glycosylation, phosphorylation, and, advantageously secretion of the gene product should be used as host cells for the expression of fluorescent protein. Such host cell lines may include but are not limited to CHO, VERO, BHK, HeLa, COS, MDCK, Jurkat, HEK-293, and WI38.

5 Mammalian cell systems which utilize recombinant viruses or viral elements to direct expression may be engineered. For example, when using adenovirus expression vectors, the fluorescent protein coding sequence may be ligated to an adenovirus transcription/translation control complex, *e.g.*, the late promoter and tripartite leader sequence. This chimeric gene may then be inserted in the adenovirus
10 genome by *in vitro* or *in vivo* recombination. Insertion in a non-essential region of the viral genome (*e.g.*, region E1 or E3) will result in a recombinant virus that is viable and capable of expressing the fluorescent protein in infected hosts (*e.g.*, see Logan & Shenk, *Proc. Natl. Acad. Sci. USA*, 81: 3655-3659, 1984). Alternatively, the vaccinia virus 7.5K promoter may be used. (*e.g.*, see, Mackett, *et al.*, *Proc. Natl. Acad. Sci.*
15 *USA*, 79: 7415-7419, 1982; Mackett, *et al.*, *J. Virol.* 49: 857-864, 1984; Panicali, *et al.*, *Proc. Natl. Acad. Sci. USA* 79: 4927-4931, 1982). Of particular interest are vectors based on bovine papilloma virus which have the ability to replicate as extrachromosomal elements (Sarver, *et al.*, *Mol. Cell. Biol.* 1: 486, 1981). Shortly after entry of this DNA into mouse cells, the plasmid replicates to about 100 to 200
20 copies per cell. Transcription of the inserted cDNA does not require integration of the plasmid into the host's chromosome, thereby yielding a high level of expression. These vectors can be used for stable expression by including a selectable marker in the plasmid, such as the *neo* gene. Alternatively, the retroviral genome can be modified for use as a vector capable of introducing and directing the expression of the
25 fluorescent protein gene in host cells (Cone & Mulligan, *Proc. Natl. Acad. Sci. USA*, 81:6349-6353, 1984). High level expression may also be achieved using inducible promoters, including, but not limited to, the metallothionine IIA promoter and heat shock promoters.

The invention can also include a localization sequence, such as a nuclear
30 localization sequence, an endoplasmic reticulum localization sequence, a peroxisome localization sequence, a mitochondrial localization sequence, or a localized protein.

Localization sequences can be targeting sequences which are described, for example, in "Protein Targeting", chapter 35 of Stryer, L., *Biochemistry* (4th ed.). W.H.

Freeman, 1995. The localization sequence can also be a localized protein. Some important localization sequences include those targeting the nucleus (KKKRRK),
5 mitochondrion (amino terminal MLRTSSLFTRRVQPSLFRNLRRLQST-),
endoplasmic reticulum (KDEL at C-terminus, assuming a signal sequence present at N-terminus), peroxisome (SKF at C-terminus), prenylation or insertion into plasma membrane (CaaX, CC, CXC, or CCXX at C-terminus), cytoplasmic side of plasma membrane (fusion to SNAP-25), or the Golgi apparatus (fusion to furin).

10 For long-term, high-yield production of recombinant proteins, stable expression is preferred. Rather than using expression vectors which contain viral origins of replication, host cells can be transformed with the fluorescent protein cDNA controlled by appropriate expression control elements (*e.g.*, promoter, enhancer, sequences, transcription terminators, polyadenylation sites, *etc.*), and a
15 selectable marker. The selectable marker in the recombinant plasmid confers resistance to the selection and allows cells to stably integrate the plasmid into their chromosomes and grow to form foci which in turn can be cloned and expanded into cell lines. For example, following the introduction of foreign DNA, engineered cells may be allowed to grow for 1-2 days in an enriched media, and then are switched to a
20 selective media. A number of selection systems may be used, including but not limited to the herpes simplex virus thymidine kinase (Wigler, *et al.*, *Cell*, 11: 223, 1977), hypoxanthine-guanine phosphoribosyltransferase (Szybalska & Szybalski, *Proc. Natl. Acad. Sci. USA*, 48:2026, 1962), and adenine phosphoribosyltransferase (Lowy, *et al.*, *Cell*, 22: 817, 1980) genes can be employed in *tk⁻*, *hgp^rt⁻* or *ap^rt⁻* cells
25 respectively. Also, antimetabolite resistance can be used as the basis of selection for *dhfr*, which confers resistance to methotrexate (Wigler, *et al.*, *Proc. Natl. Acad. Sci. USA*, 77: 3567, 1980; O'Hare, *et al.*, *Proc. Natl. Acad. Sci. USA*, 8: 1527, 1981); *gpt*, which confers resistance to mycophenolic acid (Mulligan & Berg, *Proc. Natl. Acad. Sci. USA*, 78: 2072, 1981; neo, which confers resistance to the aminoglycoside G-418
30 (Colberre-Garapin, *et al.*, *J. Mol. Biol.*, 150:1, 1981); and *hygro*, which confers resistance to hygromycin (Santerre, *et al.*, *Gene*, 30: 147, 1984) genes. Recently,

additional selectable genes have been described, namely *trpB*, which allows cells to utilize indole in place of tryptophan; *hisD*, which allows cells to utilize histinol in place of histidine (Hartman & Mulligan, *Proc. Natl. Acad. Sci. USA*, 85:8047, 1988); and ODC (ornithine decarboxylase) which confers resistance to the ornithine decarboxylase inhibitor, 2-(difluoromethyl)-DL-ornithine, DFMO (McConlogue L., In: *Current Communications in Molecular Biology*, Cold Spring Harbor Laboratory, ed., 1987).

DNA sequences encoding the fluorescence protein polypeptide of the invention can be expressed *in vitro* by DNA transfer into a suitable host cell. "Host cells" are cells in which a vector can be propagated and its DNA expressed. The term also includes any progeny of the subject host cell. It is understood that all progeny may not be identical to the parental cell since there may be mutations that occur during replication. However, such progeny are included when the term "host cell" is used. Methods of stable transfer, in other words when the foreign DNA is continuously maintained in the host, are known in the art.

The expression vector can be transfected into a host cell for expression of the recombinant nucleic acid. Host cells can be selected for high levels of expression in order to purify the fluorescent protein fusion protein. *E. coli* is useful for this purpose. Alternatively, the host cell can be a prokaryotic or eukaryotic cell selected to study the activity of an enzyme produced by the cell. In this case, the linker peptide is selected to include an amino acid sequence recognized by the protease. The cell can be, *e.g.*, a cultured cell or a cell *in vivo*.

A primary advantage of fluorescent protein fusion proteins is that they are prepared by normal protein biosynthesis, thus completely avoiding organic synthesis and the requirement for customized unnatural amino acid analogs. The constructs can be expressed in *E. coli* in large scale for *in vitro* assays. Purification from bacteria is simplified when the sequences include polyhistidine tags for one-step purification by nickel-chelate chromatography. Alternatively, the substrates can be expressed directly in a desired host cell for assays *in situ*.

In another embodiment, the invention provides a transgenic non-human animal that expresses a nucleic acid sequence which encodes the fluorescent protein.

The "non-human animals" of the invention comprise any non-human animal having nucleic acid sequence which encodes a fluorescent protein. Such non-human animals include vertebrates such as rodents, non-human primates, sheep, dog, cow, pig, amphibians, and reptiles. Preferred non-human animals are selected from the rodent family including rat and mouse, most preferably mouse. The "transgenic non-human animals" of the invention are produced by introducing "transgenes" into the germline of the non-human animal. Embryonal target cells at various developmental stages can be used to introduce transgenes. Different methods are used depending on the stage of development of the embryonic target cell. The zygote is the best target for micro-injection. In the mouse, the male pronucleus reaches the size of approximately 20 micrometers in diameter which allows reproducible injection of 1-2 pl of DNA solution. The use of zygotes as a target for gene transfer has a major advantage in that in most cases the injected DNA will be incorporated into the host gene before the first cleavage (Brinster *et al.*, *Proc. Natl. Acad. Sci. USA* 82:4438-4442, 1985). As a consequence, all cells of the transgenic non-human animal will carry the incorporated transgene. This will in general also be reflected in the efficient transmission of the transgene to offspring of the founder since 50% of the germ cells will harbor the transgene. Microinjection of zygotes is the preferred method for incorporating transgenes in practicing the invention.

The term "transgenic" is used to describe an animal which includes exogenous genetic material within all of its cells. A "transgenic" animal can be produced by cross-breeding two chimeric animals which include exogenous genetic material within cells used in reproduction. Twenty-five percent of the resulting offspring will be transgenic *i.e.*, animals which include the exogenous genetic material within all of their cells in both alleles. 50% of the resulting animals will include the exogenous genetic material within one allele and 25% will include no exogenous genetic material.

Retroviral infection can also be used to introduce transgene into a non-human animal. The developing non-human embryo can be cultured *in vitro* to the blastocyst stage. During this time, the blastomeres can be targets for retro viral infection (Jaenich, R., *Proc. Natl. Acad. Sci USA* 73:1260-1264, 1976). Efficient infection of

the blastomeres is obtained by enzymatic treatment to remove the zona pellucida (Hogan, *et al.* (1986) in *Manipulating the Mouse Embryo*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.). The viral vector system used to introduce the transgene is typically a replication-defective retro virus carrying the transgene (Jahner, *et al.*, *Proc. Natl. Acad. Sci. USA* 82:6927-6931, 1985; Van der Putten, *et al.*, *Proc. Natl. Acad. Sci. USA* 82:6148-6152, 1985). Transfection is easily and efficiently obtained by culturing the blastomeres on a monolayer of virus-producing cells (Van der Putten, *supra*; Stewart, *et al.*, *EMBO J.* 6:383-388, 1987). Alternatively, infection can be performed at a later stage. Virus or virus-producing cells can be injected into the blastocoele (D. Jahner *et al.*, *Nature* 298:623-628, 1982). Most of the founders will be mosaic for the transgene since incorporation occurs only in a subset of the cells which formed the transgenic nonhuman animal. Further, the founder may contain various retro viral insertions of the transgene at different positions in the genome which generally will segregate in the offspring. In addition, it is also possible to introduce transgenes into the germ line, albeit with low efficiency, by intrauterine retro viral infection of the midgestation embryo (D. Jahner *et al.*, *supra*).

A third type of target cell for transgene introduction is the embryonal stem cell (ES). ES cells are obtained from pre-implantation embryos cultured *in vitro* and fused with embryos (M. J. Evans *et al.* *Nature* 292:154-156, 1981; M.O. Bradley *et al.*, *Nature* 309: 255-258, 1984; Gossler, *et al.*, *Proc. Natl. Acad. Sci. USA* 83: 9065-9069, 1986; and Robertson *et al.*, *Nature* 322:445-448, 1986). Transgenes can be efficiently introduced into the ES cells by DNA transfection or by retro virus-mediated transduction. Such transformed ES cells can thereafter be combined with blastocysts from a nonhuman animal. The ES cells thereafter colonize the embryo and contribute to the germ line of the resulting chimeric animal. (For review see Jaenisch, R., *Science* 240: 1468-1474, 1988).

"Transformed" means a cell into which (or into an ancestor of which) has been introduced, by means of recombinant nucleic acid techniques, a heterologous nucleic acid molecule. "Heterologous" refers to a nucleic acid sequence that either originates

from another species or is modified from either its original form or the form primarily expressed in the cell.

"Transgene" means any piece of DNA which is inserted by artifice into a cell, and becomes part of the genome of the organism (*i.e.*, either stably integrated or as a stable extrachromosomal element) which develops from that cell. Such a transgene may include a gene which is partly or entirely heterologous (*i.e.*, foreign) to the transgenic organism, or may represent a gene homologous to an endogenous gene of the organism. Included within this definition is a transgene created by the providing of an RNA sequence which is transcribed into DNA and then incorporated into the genome. The transgenes of the invention include DNA sequences which encode which encodes the fluorescent protein which may be expressed in a transgenic non-human animal. The term "transgenic" as used herein additionally includes any organism whose genome has been altered by *in vitro* manipulation of the early embryo or fertilized egg or by any transgenic technology to induce a specific gene knockout. The term "gene knockout" as used herein, refers to the targeted disruption of a gene *in vivo* with complete loss of function that has been achieved by any transgenic technology familiar to those in the art. In one embodiment, transgenic animals having gene knockouts are those in which the target gene has been rendered nonfunctional by an insertion targeted to the gene to be rendered non-functional by homologous recombination. As used herein, the term "transgenic" includes any transgenic technology familiar to those in the art which can produce an organism carrying an introduced transgene or one in which an endogenous gene has been rendered non-functional or "knocked out."

25 III. USES OF ENGINEERED FLUORESCENT PROTEINS

The proteins of this invention are useful in any methods that employ fluorescent proteins.

The engineered fluorescent proteins of this invention are useful as fluorescent markers in the many ways fluorescent markers already are used. This includes, for example, coupling engineered fluorescent proteins to antibodies, nucleic acids or

other receptors for use in detection assays, such as immunoassays or hybridization assays.

The engineered fluorescent proteins of this invention are useful to track the movement of proteins in cells. In this embodiment, a nucleic acid molecule encoding the fluorescent protein is fused to a nucleic acid molecule encoding the protein of interest in an expression vector. Upon expression inside the cell, the protein of interest can be localized based on fluorescence. In another version, two proteins of interest are fused with two engineered fluorescent proteins having different fluorescent characteristics.

The engineered fluorescent proteins of this invention are useful in systems to detect induction of transcription. In certain embodiments, a nucleotide sequence encoding the engineered fluorescent protein is fused to expression control sequences of interest and the expression vector is transfected into a cell. Induction of the promoter can be measured by detecting the expression and/or quantity of fluorescence. Such constructs can be used to follow signaling pathways from receptor to promoter.

The engineered fluorescent proteins of this invention are useful in applications involving FRET. Such applications can detect events as a function of the movement of fluorescent donors and acceptor towards or away from each other. One or both of the donor/acceptor pair can be a fluorescent protein. A preferred donor and receptor pair for FRET based assays is a donor with a T203I mutation and an acceptor with the mutation T203X, wherein X is an aromatic amino acid-39, especially T203Y, T203W, or T203H. In a particularly useful pair the donor contains the following mutations: S72A, K79R, Y145F, M153A and T203I (with a excitation peak of 395 nm and an emission peak of 511 nm) and the acceptor contains the following mutations S65G, S72A, K79R, and T203Y. This particular pair provides a wide separation between the excitation and emission peaks of the donor and provides good overlap between the donor emission spectrum and the acceptor excitation spectrum. Other red-shifted mutants, such as those described herein, can also be used as the acceptor in such a pair.

In one aspect, FRET is used to detect the cleavage of a substrate having the donor and acceptor coupled to the substrate on opposite sides of the cleavage site. Upon cleavage of the substrate, the donor/acceptor pair physically separate, eliminating FRET. Assays involve contacting the substrate with a sample, and
5 determining a qualitative or quantitative change in FRET. In one embodiment, the engineered fluorescent protein is used in a substrate for β -lactamase. Examples of such substrates are described in United States patent applications 08/407,544, filed March 20, 1995 and International Application PCT/US96/04059, filed March 20, 1996. In another embodiment, an engineered fluorescent protein donor/acceptor pair
10 are part of a fusion protein coupled by a peptide having a proteolytic cleavage site. Such tandem fluorescent proteins are described in United States patent application 08/594,575, filed January 31, 1996.

In another aspect, FRET is used to detect changes in potential across a membrane. A donor and acceptor are placed on opposite sides of a membrane such
15 that one translates across the membrane in response to a voltage change. This creates a measurable FRET. Such a method is described in United States patent application 08/481,977, filed June 7, 1995 and International Application PCT/US96/09652, filed June 6, 1996.

The engineered proteins of this invention are useful in the creation of
20 biosensors for determining the concentrations of ions within samples and living cells and transgenic organisms. Upon binding of an ion to the fluorescent protein, a change in at least one measurable fluorescent property of the engineered fluorescent protein occurs that provides the basis for determining the presence of the ion of interest.

The engineered protein of this invention are useful in the creation of
25 fluorescent substrates for protein kinases. Such substrates incorporate an amino acid sequence recognizable by protein kinases. Upon phosphorylation, the engineered fluorescent protein undergoes a change in a fluorescent property. Such substrates are useful in detecting and measuring protein kinase activity in a sample of a cell, upon transfection and expression of the substrate. Preferably, the kinase recognition site is
30 placed within about 20 amino acids of a terminus of the engineered fluorescent protein. The kinase recognition site also can be placed in a loop domain of the

protein. (See, e.g. Figure 1B.) Methods for making fluorescent substrates for protein kinases are described in United States patent application 08/680,877, filed July 16, 1996.

5 A protease recognition site also can be introduced into a loop domain. Upon cleavage, fluorescent property changes in a measurable fashion.

The invention also includes a method of identifying a test chemical. Typically, the method includes contacting a test chemical a sample containing a biological entity labeled with a functional, engineered fluorescent protein or a polynucleotide encoding said functional, engineered fluorescent protein. By
10 monitoring fluorescence (i.e. a fluorescent property) from the sample containing the functional engineered fluorescent protein it can be determined whether a test chemical is active. Controls can be included to insure the specificity of the signal. Such controls include measurements of a fluorescent property in the absence of the test chemical, in the presence of a chemical with an expected activity (e.g., a known
15 modulator) or engineered controls (e.g., absence of engineered fluorescent protein, absence of engineered fluorescent protein polynucleotide or the absence of operably linkage of the engineered fluorescent protein).

The fluorescence in the presence of a test chemical can be greater or less than in the absence of said test chemical. For instance if the engineered fluorescent protein
20 is used a reporter of gene expression, the test chemical may up or down regulate gene expression. For such types of screening, the polynucleotide encoding the functional, engineered fluorescent protein is operatively linked to a genomic polynucleotide or a re. Alternatively, the functional, engineered fluorescent protein is fused to second functional protein. This embodiment can be used to track localization of the second
25 protein or to track protein-protein interactions using energy transfer.

IV. PROCEDURES

Fluorescence in a sample is measured using a fluorimeter. In general, excitation radiation from an excitation source having a first wavelength, passes
30 through excitation optics. The excitation optics cause the excitation radiation to

excite the sample. In response, fluorescent proteins in the sample emit radiation which has a wavelength that is different from the excitation wavelength. Collection optics then collect the emission from the sample. The device can include a temperature controller to maintain the sample at a specific temperature while it is being scanned. According to one embodiment, a multi-axis translation stage moves a microtiter plate holding a plurality of samples in order to position different wells to be exposed. The multi-axis translation stage, temperature controller, auto-focusing feature, and electronics associated with imaging and data collection can be managed by an appropriately programmed digital computer. The computer also can transform the data collected during the assay into another format for presentation. This process can be miniaturized and automated to enable screening many thousands of compounds.

Methods of performing assays on fluorescent materials are well known in the art and are described in, e.g., Lakowicz, J.R., *Principles of Fluorescence Spectroscopy*, New York: Plenum Press (1983); Herman, B., Resonance energy transfer microscopy, in: *Fluorescence Microscopy of Living Cells in Culture, Part B, Methods in Cell Biology*, vol. 30, ed. Taylor, D.L. & Wang, Y.-L., San Diego: Academic Press (1989), pp. 219-243; Turro, N.J., *Modern Molecular Photochemistry*, Menlo Park: Benjamin/Cummings Publishing Co., Inc. (1978), pp. 296-361.

Mutagenesis and protein preparation

YFP variants and revertants were prepared using the PCR-based QuikChange™ Site-Directed Mutagenesis Kit (Stratagene, La Jolla, CA), according to the manufacturer's directions and using the YFP clone 10c as a template (Ormo *et al.*, 1996). Mutations were verified by sequencing the entire gene, and all GFP variants were expressed and purified as described (Ormo *et al.*, 1996).

Fluorescence measurements

Small aliquots of concentrated protein were diluted 25-fold into a series of buffers (20 mM MES pH 6.0, MES pH 6.5, PIPES 7.0, HEPES 7.5, and TAPS 8.0) with constant ionic strength. The buffers contained varying concentrations of either potassium chloride or potassium iodide, and the ionic strength was adjusted to 150 mM with potassium D-gluconate. Fluorescence measurements as a function of pH and halide concentration were carried out on a Hitachi F4500 fluorescence spectrophotometer at room temperature ($\lambda_{\text{ex}} = 514 \text{ nm}$), scanning the emission between 520 and 550 nm three times at a rate of 60 nm/min. Maximum emission at 528 nm was averaged, and normalized with respect to fluorescence in the absence of halides.

Crystal growth and data collection

YFP-H148Q was concentrated to 15 mg/ml in 20 mM TRIS pH 7.9, and crystals were grown in hanging drops containing 5 μl protein and 5 μl mother liquor. The mother liquor contained 22% PEG 1550 at pH 5.5 in 100 mM sodium acetate and 90 mM MgCl_2 . The rod-shaped crystals were approximately 0.04 mm across and up to 1.0 mm long, and grew within 1.5 to 2 years at 4 $^{\circ}\text{C}$. One crystal was soaked in synthetic mother liquor containing the above ingredients without MgCl_2 but 100 mM potassium iodide, and 20% ethylene glycol for cryo-protection (referred to as iodide soak). Another crystal was soaked in the above mother liquor containing 100 mM MgCl_2 and 20% ethylene glycol (referred to as chloride soak). Both soaks were carried out at pH 4.6 for 4 hours at room temperature, and data collection proceeded immediately thereafter. The crystals were flash-frozen, and X-ray diffraction data were collected at 100 K using a RAXIS-IIc image plate mounted on a Rigaku RUH3 rotating anode generator equipped with mirrors.

Structure determination of YFP-H148Q, and identification of iodide binding sites

5 The two data sets were processed with Denzo v1.9 and scaled using ScalePack (Otwinowski & Minor, 1997). The spacegroup is P2₁2₁2₁, with unit cell parameters $a = 51.2$, $b = 62.8$, and $c = 68.7$ Å for the iodide soak, and $a = 51.7$, $b = 62.6$, and $c = 66.2$ Å for the chloride soak. The crystals are nearly isomorphous to YFP-H148G (Wachter et al., 1998) and GFP S65T crystals (Ormo *et al.*, 1996) previously
10 described, and the YFP-H148G coordinate file 2yfp (Wachter *et al.*, 1998) was used as a model for phasing. A model for the anionic chromophore was obtained by semi-empirical molecular orbital calculations using AM1 in the program SPARTAN version 4.1 (Wavefunction Inc., Irvine, CA).

 An anomalous difference map was calculated from the data set derived from
15 the iodide soak (anomalous data 65% complete), since iodine exhibits a significant anomalous signal at the in-house CuK α wavelength of 1.54 Å. Heavy atom phases were approximated by subtraction of 90° from calculated protein phases using the program scaleit in the CCP4 program suite (Collaborative Computational Project N. 4, 1994). The anomalous difference map identified two iodide positions, one buried
20 in the protein interior and one on the protein surface.

Refinement of YFP-H148Q with and without bound iodide

The two datasets, derived from the iodide and from the chloride soak, were refined in a similar manner. After initial rigid body refinement to 4.0 Å, positional refinement was carried out using the data to 3.0 Å, then to the limit of resolution (Table 1), using the program TNT (Tronrud *et al.*, 1987). During early cycles of refinement, bound halides were not modeled, and the glutamine in position 148 was modeled as a glycine. Electron density maps ($2F_o - F_c$ and $F_o - F_c$) were inspected intermittently using O (Jones *et al.*, 1991). The $F_o - F_c$ maps clearly indicated the positions of the buried and surface iodides, at 11 and 5.5 rms deviations respectively, located in the centers of the two anomalous difference density peaks, though no positive difference density consistent with buried chloride binding was observed. Density for the Gln148 side chain was clearly visible early on, allowing for the modeling of the glutamine as a rotamer different from the original histidine.

B-factors were refined using the default TNT B-factor correlation library. B-factor correlation values derived from His and Phe were used to model the chromophore atoms. Bound solvent molecules were added to the model where appropriate as judged from difference density and proximity of hydrogen bond partners. Before refining the occupancy of the bound halides, the B-factors for these halides were fixed. The thermal factor of the buried iodide was set to the average B-factor of the twelve atoms closest to it, 30 Å², (Figure 5), and the thermal factor for the surface iodide was set to the average B-factor of the six closest solvent molecules bound to the protein surface, 39 Å². The last step in refinement was the refinement of the occupancy of the two bound halides.

Determination of chromophore pK_a and iodide binding constants by absorbance

The chromophore pK_a was determined from absorbance scans at varying anion concentrations. Absorbance scans were collected at room temperature between 250 and 600 nm (Shimadzu 2101 spectrophotometer) on 0.05 mg/ml YFP under two different pH conditions appropriate for the particular anion, chosen from a series of buffers (20 mM malic acid pH 5.8, malic acid pH 6.1, MES pH 6.4, HEPES pH 7.1). The optical density of the chromophore anion (514 to 515 nm for YFP and YFP-H148Q) at the buffer pH, as well as the optical density at pH 9 in the absence of interacting anions, were used in the Henderson-Hasselbalch equation to estimate the chromophore pK_a for each condition examined. Microscopic binding constants for anion binding to the protein were extracted by curve fitting of the chromophore pK_a to the anion concentration, using an expression for a linked equilibrium involving two different ligands.

The following examples are provided by way of illustration, not by way of limitation.

EXAMPLES

As a step in understanding the properties of GFP, and to aid in the tailoring of GFPs with altered characteristics, we have determined the three dimensional structure at 1.9 Å resolution of the S65T mutant (R. Heim et al. *Nature* 373:664-665 (1995)) of *A. victoria* GFP. This mutant also contains the ubiquitous Q80R substitution, which accidentally occurred in the early distribution of the GFP cDNA and is not known to have any effect on the protein properties (M. Chalfie et al. *Science* 263:802-805 (1994)).

Histidine-tagged S65T GFP (R. Heim et al. *Nature* 373:664-665 (1995)) was overexpressed in JM109/pRSET_B in 4 l YT broth plus ampicillin at 37°C, 450 rpm and 5 l/min air flow. The temperature was reduced to 25°C at A₅₉₅ = 0.3, followed by induction with 1mM isopropylthiogalactoside for 5h. Cell paste was stored at -80°C

overnight, then was resuspended in 50 mM HEPES pH 7.9, 0.3 M NaCl, 5 mM 2-mercaptoethanol, 0.1 mM phenylmethyl-sulfonylfluoride (PMSF), passed once through a French press at 10,000 psi, then centrifuged at 20 K rpm for 45 min. The supernatant was applied to a Ni-NTA-agarose column (Qiagen), followed by a wash
5 with 20 mM imidazole, then eluted with 100 mM imidazole. Green fractions were pooled and subjected to chymotryptic (Sigma) proteolysis (1:50 w/w) for 22 h at RT. After addition of 0.5 mM PMSF, the digest was reapplied to the Ni column. N-terminal sequencing verified the presence of the correct N-terminal methionine. After dialysis against 20 mM HEPES, pH 7.5 and concentration to $A_{490} = 20$, rod-shaped
10 crystals were obtained at RT in hanging drops containing 5 μ l protein and 5 μ l well solution, 22-26% PEG 4000 (Serva), 50 mM HEPES pH 8.0-8.5, 50 mM $MgCl_2$ and 10 mM 2-mercapto-ethanol within 5 days. Crystals were 0.05 mm across and up to 1.0 mm long. The space group is $P2_12_12_1$ with $a = 51.8$, $b = 62.8$, $c = 70.7$ Å, $Z=4$. Two crystal forms of wild-type GFP, unrelated to the present form, have been
15 described by M. A. Perrozo, K. B. Ward, R. B. Thompson, & W. W. Ward. *J. Biol. Chem.* 203, 7713-7716 (1988).

The structure of GFP was determined by multiple isomorphous replacement and anomalous scattering (Table E), solvent flattening, phase combination and crystallographic refinement. The most remarkable feature of the fold of GFP is an
20 eleven stranded β -barrel wrapped around a single central helix (Fig. 1A and 1B), where each strand consists of approximately 9-13 residues. The barrel forms a nearly perfect cylinder 42 Å long and 24 Å in diameter. The N-terminal half of the polypeptide comprises three anti-parallel strands, the central helix, and then 3 more anti-parallel strands, the latter of which (residues 118-123) is parallel to the N-
25 terminal strand (residues 11-23). The polypeptide backbone then crosses the "bottom" of the molecule to form the second half of the barrel in a five-strand Greek Key motif. The top end of the cylinder is capped by three short, distorted helical segments, while one short, very distorted helical segment caps the bottom of the cylinder. The main-chain hydrogen bonding lacing the surface of the cylinder very
30 likely accounts for the unusual stability of the protein towards denaturation and proteolysis. There are no large segments of the polypeptide that could be excised

while preserving the intactness of the shell around the chromophore. Thus it would seem difficult to re-engineer GFP to reduce its molecular weight (J. Dopf & T.M. Horiagon *Gene* 173:39-43 (1996)) by a large percentage.

The *p*-hydroxybenzylideneimidazolidinone chromophore (C. W. Cody et al. *Biochemistry* 32:1212-1218 (1993)) is completely protected from bulk solvent and centrally located in the molecule. The total and presumably rigid encapsulation is probably responsible for the small Stokes' shift (i.e. wavelength difference between excitation and emission maxima), high quantum yield of fluorescence, inability of O₂ to quench the excited state (B.D. Nageswara Rao et al. *Biophys. J.* 32:630-632 (1980)), and resistance of the chromophore to titration of the external pH (W. W. Ward. *Bioluminescence and Chemiluminescence* (M. A. DeLuca and W. D. McElroy, eds) Academic Press pp. 235-242 (1981); W. W. Ward & S. H. Bokman. *Biochemistry* 21:4535-4540 (1982); W. W. Ward et al. *Photochem. Photobiol.* 35:803-808 (1982)). It also allows one to rationalize why fluorophore formation should be a spontaneous intramolecular process (R. Heim et al. *Proc. Natl. Acad. Sci. USA* 91:12501-12504 (1994)), as it is difficult to imagine how an enzyme could gain access to the substrate. The plane of the chromophore is roughly perpendicular (60°) to the symmetry axis of the surrounding barrel. One side of the chromophore faces a surprisingly large cavity, that occupies a volume of approximately 135 Å³ (B. Lee & F. M. Richards. *J. Mol. Biol.* 55:379-400 (1971)). The atomic radii were those of Lee & Richards, calculated using the program MS with a probe radius of 1.4 Å. (M. L. Connolly, *Science* 221:709-713 (1983)). The cavity does not open out to bulk solvent. Four water molecules are located in the cavity, forming a chain of hydrogen bonds linking the buried side chains of Glu²²² and Gln⁶⁹. Unless occupied, such a large cavity would be expected to de-stabilize the protein by several kcal/mol (S. J. Hubbard et al., *Protein Engineering* 7:613-626 (1994); A. E. Eriksson et al. *Science* 255:178-183 (1992)). Part of the volume of the cavity might be the consequence of the compaction resulting from cyclization and dehydration reactions. The cavity might also temporarily accommodate the oxidant, most likely O₂ (A. B. Cubitt et al. *Trends Biochem. Sci.* 20:448-455 (1995); R. Heim et al. *Proc. Natl. Acad. Sci. USA* 91:12501-12504 (1994); S. Inouye & F.I. Tsuji. *FEBS Lett.* 351:211-214 (1994)), that

dehydrogenates the α - β bond of Tyr⁶⁶. The chromophore, cavity, and side chains that contact the chromophore are shown in Figure 2A and a portion of the final electron density map in this vicinity in 2B.

The opposite side of the chromophore is packed against several aromatic and polar side chains. Of particular interest is the intricate network of polar interactions with the chromophore (Fig. 2C). His¹⁴⁸, Thr²⁰³ and Ser²⁰⁵ form hydrogen bonds with the phenolic hydroxyl; Arg⁹⁶ and Gln⁹⁴ interact with the carbonyl of the imidazolidinone ring and Glu²²² forms a hydrogen bond with the side chain of Thr⁶⁵. Additional polar interactions, such as hydrogen bonds to Arg⁹⁶ from the carbonyl of Thr⁶², and the side-chain carbonyl of Gln¹⁸³, presumably stabilize the buried Arg⁹⁶ in its protonated form. In turn, this buried charge suggests that a partial negative charge resides on the carbonyl oxygen of the imidazolidinone ring of the deprotonated fluorophore, as has previously been suggested (W. W. Ward. *Bioluminescence and Chemiluminescence* (M. A. DeLuca and W. D. McElroy, eds) Academic Press pp. 235-242 (1981); W. W. Ward & S. H. Bokman. *Biochemistry* 21:4535-4540 (1982); W. W. Ward et al. *Photochem. Photobiol.* 35:803-808 (1982)). Arg⁹⁶ is likely to be essential for the formation of the fluorophore, and may help catalyze the initial ring closure. Finally, Tyr¹⁴⁵ shows a typical stabilizing edge-face interaction with the benzyl ring. Trp⁵⁷, the only tryptophan in GFP, is located 13 Å to 15 Å from the chromophore and the long axes of the two ring systems are nearly parallel. This indicates that efficient energy transfer to the latter should occur, and explains why no separate tryptophan emission is observable (D.C. Prasher et al. *Gene* 111:229-233 (1992). The two cysteines in GFP, Cys⁴⁸ and Cys⁷⁰, are 24 Å apart, too distant to form a disulfide bridge. Cys⁷⁰ is buried, but Cys⁴⁸ should be relatively accessible to sulfhydryl-specific reagents. Such a reagent, 5,5'-dithiobis(2-nitrobenzoic acid), is reported to label GFP and quench its fluorescence (S. Inouye & F.I. Tsuji *FEBS Lett.* 351:211-214 (1994)). This effect was attributed to the necessity for a free sulfhydryl, but could also reflect specific quenching by the 5-thio-2-nitrobenzoate moiety that would be attached to Cys⁴⁸.

Although the electron density map is for the most part consistent with the proposed structure of the chromophore (D.C. Prasher et al. *Gene* 111:229-233 (1992);

C. W. Cody et al. *Biochemistry* 32:1212-1218 (1993)) in the *cis* [Z-] configuration, with no evidence for any substantial fraction of the opposite isomer around the chromophore double bond, difference features are found at $>4 \sigma$ in the final ($F_o - F_c$) electron density map that can be interpreted to represent either the intact, uncyclized polypeptide or a carbinolamine (inset to Fig. 2C). This suggests that a significant fraction, perhaps as much as 30% of the molecules in the crystal, have failed to undergo the final dehydration reaction. Confirmation of incomplete dehydration comes from electrospray mass spectrometry, which consistently shows that the average masses of both wild-type and S65T GFP ($31,086 \pm 4$ and $31,099.5 \pm 4$ Da, respectively) are 6-7 Da higher than predicted (31,079 and 31,093 Da, respectively) for the fully matured proteins. Such a discrepancy could be explained by a 30-35% mole fraction of apoprotein or carbinolamine with 18 or 20 Da higher molecular weight. The natural abundance of ^{13}C and ^2H and the finite resolution of the Hewlett-Packard 5989B electrospray mass spectrometer used to make these measurements do not permit the individual peaks to be resolved, but instead yields an average mass peak with a full width at half maximum of approximately 15 Da. The molecular weights shown include the His-tag, which has the sequence MRGSHHHHHH GMASMTGGQQM GRDL YDDDDK DPPAEF (SEQ ID NO:5). Mutants of GFP that increase the efficiency of fluorophore maturation might yield somewhat brighter preparations. In a model for the apoprotein, the Thr⁶⁵-Tyr⁶⁶ peptide bond is approximately in the α -helical conformation, while the peptide of Tyr⁶⁶-Gly⁶⁷ appears to be tipped almost perpendicular to the helix axis by its interaction with Arg⁹⁶. This further supports the speculation that Arg⁹⁶ is important in generating the conformation required for cyclization, and possibly also for promoting the attack of Gly⁶⁷ on the carbonyl carbon of Thr⁶⁵ (A. B. Cubitt et al. *Trends Biochem. Sci.* 20:448-455 (1995)).

The results of previous random mutagenesis have implicated several amino acid side chains to have substantial effects on the spectra and the atomic model confirms that these residues are close to the chromophore. The mutations T203I and E222G have profound but opposite consequences on the absorption spectrum (T. Ehrig et al. *FEBS Letters* 367:163-166 (1995)). T203I (with wild-type Ser⁶⁵) lacks

the 475 nm absorbance peak usually attributed to the anionic chromophore and shows only the 395 nm peak thought to reflect the neutral chromophore (R. Heim et al. *Proc. Natl. Acad. Sci. USA* 91:12501-12504 (1994); T. Ehrig et al. *FEBS Letters* 367:163-166 (1995)). Indeed, Thr²⁰³ is hydrogen-bonded to the phenolic oxygen of the chromophore, so replacement by Ile should hinder ionization of the phenolic oxygen. Mutation of Glu²²² to Gly (T. Ehrig et al. *FEBS Letters* 367:163-166 (1995)) has much the same spectroscopic effect as replacing Ser⁶⁵ by Gly, Ala, Cys, Val, or Thr, namely to suppress the 395 nm peak in favor of a peak at 470-490 nm (R. Heim et al. *Nature* 373:664-665 (1995); S. Delagrave et al. *Bio/Technology* 13:151-154 (1995)). Indeed Glu²²² and the remnant of Thr⁶⁵ are hydrogen-bonded to each other in the present structure, probably with the uncharged carboxyl of Glu²²² acting as donor to the side chain oxygen of Thr⁶⁵. Mutations E222G, S65G, S65A, and S65V would all suppress such H-bonding. To explain why only wild-type protein has both excitation peaks, Ser⁶⁵, unlike Thr⁶⁵, may adopt a conformation in which its hydroxyl donates a hydrogen bond to and stabilizes Glu²²² as an anion, whose charge then inhibits ionization of the chromophore. The structure also explains why some mutations seem neutral. For example, Gln⁸⁰ is a surface residue far removed from the chromophore, which explains why its accidental and ubiquitous mutation to Arg seems to have no obvious intramolecular spectroscopic effect (M. Chalfie et al. *Science* 263:802-805 (1994)).

The development of GFP mutants with red-shifted excitation and emission maxima is an interesting challenge in protein engineering (A. B. Cubitt et al. *Trends Biochem. Sci.* 20:448-455 (1995); R. Heim et al. *Nature* 373:664-665 (1995); S. Delagrave et al. *Bio/Technology* 13:151-154 (1995)). Such mutants would also be valuable for avoidance of cellular autofluorescence at short wavelengths, for simultaneous multicolor reporting of the activity of two or more cellular processes, and for exploitation of fluorescence resonance energy transfer as a signal of protein-protein interaction (R. Heim & R.Y. Tsien. *Current Biol.* 6:178-182 (1996)). Extensive attempts using random mutagenesis have shifted the emission maximum by at most 6 nm to longer wavelengths, to 514 nm (R. Heim & R.Y. Tsien. *Current Biol.* 6:178-182 (1996)); previously described "red-shifted" mutants merely suppressed the

395 nm excitation peak in favor of the 475 nm peak without any significant reddening of the 505 nm emission (S. Delagrave et al. *Bio/Technology* 13:151-154 (1995)). Because Thr²⁰³ is revealed to be adjacent to the phenolic end of the chromophore, we mutated it to polar aromatic residues such as His, Tyr, and Trp in the hope that the additional polarizability of their systems would lower the energy of the excited state of the adjacent chromophore. All three substitutions did indeed shift the emission peak to greater than 520 nm (Table F). A particularly attractive mutation was T203Y/S65G/V68L/S72A, with excitation and emission peaks at 513 and 527 nm respectively. These wavelengths are sufficiently different from previous GFP mutants to be readily distinguishable by appropriate filter sets on a fluorescence microscope. The extinction coefficient, 36,500 M⁻¹cm⁻¹, and quantum yield, 0.63, are almost as high as those of S65T (R. Heim et al. *Nature* 373:664-665 (1995)).

Comparison of *Aequorea* GFP with other protein pigments is instructive. Unfortunately, its closest characterized homolog, the GFP from the sea pansy *Renilla reniformis* (O. Shimomura and F.H. Johnson *J. Cell. Comp. Physiol.* 59:223 (1962); J. G. Morin and J. W. Hastings, *J. Cell. Physiol.* 77:313 (1971); H. Morise et al. *Biochemistry* 13:2656 (1974); W. W. Ward *Photochem. Photobiol. Reviews* (Smith, K. C. ed.) 4:1 (1979); W. W. Ward. *Bioluminescence and Chemiluminescence* (M. A. DeLuca and W. D. McElroy, eds) Academic Press pp. 235-242 (1981); W. W. Ward & S. H. Bokman *Biochemistry* 21:4535-4540 (1982); W. W. Ward et al. *Photochem. Photobiol.* 35:803-808 (1982)), has not been sequenced or cloned, though its chromophore is derived from the same FSYG sequence as in wild-type *Aequorea* GFP (R. M. San Pietro et al. *Photochem. Photobiol.* 57:63S (1993)). The closest analog for which a three dimensional structure is available is the photoactive yellow protein (PYP, G. E. O. Borgstahl et al. *Biochemistry* 34:6278-6287 (1995)), a 14-kDa photoreceptor from halophilic bacteria. PYP in its native dark state absorbs maximally at 446 nm and transduces light with a quantum yield of 0.64, rather closely matching wild-type GFP's long wavelength absorbance maximum near 475 nm and fluorescence quantum yield of 0.72-0.85. The fundamental chromophore in both proteins is an anionic *p*-hydroxycinnamyl group, which is covalently attached to the protein via a thioester linkage in PYP and a heterocyclic iminolactam in GFP. Both

proteins stabilize the negative charge on the chromophore with the help of buried cationic arginine and neutral glutamic acid groups, Arg⁵² and Glu⁴⁶ in PYP and Arg⁹⁶ and Glu²²² in GFP, though in PYP the residues are close to the oxyphenyl ring whereas in GFP they are nearer the carbonyl end of the chromophore. However, PYP has an overall α/β fold with appropriate flexibility and signal transduction domains to enable it to mediate the cellular phototactic response, whereas GFP is a much more regular and rigid β -barrel to minimize parasitic dissipation of the excited state energy as thermal or conformational motions. GFP is an elegant example of how a visually appealing and extremely useful function, efficient fluorescence, can be spontaneously generated from a cohesive and economical protein structure.

A. Summary Of GFP Structure Determination

Data were collected at room temperature in house using either Molecular Structure Corp. R-axis II or San Diego Multiwire Systems (SDMS) detectors (Cu K α) and later at beamline X4A at the Brookhaven National Laboratory at the selenium absorption edge ($\lambda = 0.979$ Å) using image plates. Data were evaluated using the HKL package (Z. Otwinowski, in *Proceedings of the CCP4 Study Weekend: Data Collection and Processing*, L. Sawyer, N. Issacs, S. Bailey, Eds. (Science and Engineering Research Council (SERC), Daresbury Laboratory, Warrington, UK, (1991)), pp 56-62; W. Minor, XDISPLAYF (Purdue University, West Lafayette, IN, 1993)) or the SDMS software (A. J. Howard et al. *Meth. Enzymol.* 114:452-471 (1985)). Each data set was collected from a single crystal. Heavy atom soaks were 2 mM in mother liquor for 2 days. Initial electron density maps were based on three heavy atom derivatives using in-house data, then later were replaced with the synchrotron data. The EMTS difference Patterson map was solved by inspection, then used to calculate difference Fourier maps of the other derivatives. Lack of closure refinement of the heavy atom parameters was performed using the Protein package (W. Steigemann, in Ph.D. Thesis (Technical University, Munich, 1974)). The MIR maps were much poorer than the overall figure of merit would suggest, and it was clear that the EMTS isomorphous differences dominated the phasing. The enhanced anomalous occupancy for the synchrotron data provided a partial solution to

the problem. Note that the phasing power was reduced for the synchrotron data, but the figure of merit was unchanged. All experimental electron density maps were improved by solvent flattening using the program DM of the CCP4 (CCP4: *A Suite of Programs for Protein Crystallography* (SERC Daresbury Laboratory, Warrington WA4 4AD UK, 1979)) package assuming a solvent content of 38%. Phase combination was performed with PHASCO2 of the Protein package using a weight of 1.0 on the atomic model. Heavy atom parameters were subsequently improved by refinement against combined phases. Model building proceeded with FRODO and O (T. A. Jones et al. *Acta. Crystallogr. Sect. A* 47:110 (1991); T. A. Jones, in *Computational Crystallography* D. Sayre, Ed. (Oxford University Press, Oxford, 1982) pp. 303-317) and crystallographic refinement was performed with the TNT package (D. E. Tronrud et al. *Acta Cryst. A* 43:489-503 (1987)). Bond lengths and angles for the chromophore were estimated using CHEM3D (Cambridge Scientific Computing). Final refinement and model building was performed against the X4A selenomethione data set, using (2F_o-F_c) electron density maps. The data beyond 1.9 Å resolution have not been used at this stage. The final model contains residues 2-229 as the terminal residues are not visible in the electron density map, and the side chains of several disordered surface residues have been omitted. Density is weak for residues 156-158 and coordinates for these residues are unreliable. This disordering is consistent with previous analyses showing that residues 1 and 233-238 are dispensable but that further truncations may prevent fluorescence (J. Dopf & T.M. Horiagon. *Gene* 173:39-43 (1996)). The atomic model has been deposited in the Protein Data Bank (access code 1EMA).

Table GDiffraction Data Statistics

<u>Crystal</u>	<u>Resolution</u> <u>(Å)</u>	<u>Total</u> <u>obs</u>	<u>Unique</u> <u>obs</u>	<u>Compl.</u> <u>(%)</u> ^a	<u>Compl.</u> <u>(shell)</u> ^b	<u>Rmerge</u> <u>(%)</u> ^c	<u>Riso</u> <u>(%)</u> ^d
<u>R-axis II</u>							
Native	2.0	51907	13582	80	69	4.1	5.8
EMTS ^c	2.6	17727	6787	87	87	5.7	20.6
SeMet	2.3	44975	10292	92	88	10.2	9.3
<u>Multiwire</u>							
HGI4-Se	3.0	15380	4332	84	79	7.2	28.8
<u>X4a</u>							
SeMet	1.8	12607	19503	80	55	9.3	9.4
		8					
EMTS	2.3	57812	9204	82	66	7.2	26.3

67

Phasing Statistics

<u>Derivative</u>	<u>Resolution</u> (Å)	<u>Number</u> <u>of sites</u>	<u>Phasing</u> <u>power^f</u>	<u>Phasing</u> <u>Power(shell)</u>	<u>FOM^g</u>	<u>FOM</u> <u>(shell)</u>
-------------------	--------------------------	----------------------------------	--	---------------------------------------	------------------------	------------------------------

In House

EMTS	3.0	2	2.08	2.08	0.77	.072
------	-----	---	------	------	------	------

SeMet	3.0	4	1.66	1.28	-	-
-------	-----	---	------	------	---	---

HGI4-Se	3.0	9	1.77	1.90	-	-
---------	-----	---	------	------	---	---

X4a

EMTS	3.0	2	1.36	1.26	0.77	.072
------	-----	---	------	------	------	------

SeMet	3.0	4	1.31	1.08	-	-
-------	-----	---	------	------	---	---

Atomic Model Statistics

Protein atoms	1790
---------------	------

5 Solvent atoms	94
-----------------	----

Resol. range (Å)	20-1.9
------------------	--------

Number of reflections (F > 0)	17676
-------------------------------	-------

Completeness	84.
--------------	-----

R. factor ^(h)	0.175
--------------------------	-------

10 Mean B-value (Å ²)	24.1
-----------------------------------	------

Deviations from ideality

Bond lengths (Å)	0.014
------------------	-------

Bond angles (°)	1.9
-----------------	-----

Restrained B-values (Å ²)	4.3
---------------------------------------	-----

15 Ramachandran outliers	0
--------------------------	---

Notes:

- (a) Completeness is the ratio of observed reflections to theoretically possible expressed as a percentage.
- (b) Shell indicates the highest resolution shell, typically 0.1-0.4 Å wide.
- 5 (c) $R_{\text{merge}} = \sum |I - \langle I \rangle| / \sum I$, where $\langle I \rangle$ is the mean of individual observations of intensities I .
- (d) $R_{\text{iso}} = \sum |I_{\text{DER}} - I_{\text{NAT}}| / \sum I_{\text{NAT}}$
- (e) Derivatives were EMTS=ethymercurithiosalicylate (residues modified Cys⁴⁸ and Cys⁷⁰), SeMet=selenomethionine substituted protein (Met¹ and Met²³³ could not be located); HgI₄-SeMet = double derivative HgI₄ on SeMet background.
- 10 (f) Phasing power = $\langle F_H \rangle / \langle E \rangle$ where $\langle F_H \rangle$ =r.m.s. heavy atom scattering and $\langle E \rangle$ =lack of closure.
- (g) FOM, mean figure of merit
- 15 (h) Standard crystallographic R-factor, $R = \sum ||F_{\text{obs}}| - |F_{\text{calc}}|| / \sum |F_{\text{obs}}|$

B. Spectral properties of Thr²⁰³ ("T203") mutants compared to S65T

The mutations F64L, V68L and S72A improve the folding of GFP at 37° (B. P. Cormack et al. *Gene* 173:33 (1996)) but do not significantly shift the

20 emission spectra.

TABLE H

Clone	Mutations	Excitation max.(nm)	Extinction coefficient (10 ³ M ⁻¹ cm ⁻¹)	Emission max.(nm)
S65T	S65T	489	39.2	511
5B	T203H/S65T	512	19.4	524
6C	T203Y/S65T	513	14.5	525

10B	T203Y/F64L/S65G/S72A	513	30.8	525
10C	T203Y/F65G/V68L/S72A	513	36.5	527
11	T203W/S65G/S72A	502	33.0	512
12H	T203Y/S65G/S72A	513	36.5	527
20A	T203Y/S65G/V68L/Q69K/S72A	515	46.0	527

C. YFP and YFP-H148Q as halide sensors at acidic and neutral pH

5 The absorbance spectrum of YFP is a function of NaCl concentration (Fig.7), with conversion of band B, the chromophore anion ($\lambda_{\max}$ 514 nm), to band A, the neutral form ($\lambda_{\max}$ 392 nm) upon addition of chloride. Since only the anion is fluorescent in the YFPs, suppression of fluorescence occurs concomitant with increasing [NaCl]. For YFP, a clean isosbestic point is observed (Fig. 7), whereas for

10 YFP-H148Q, the isosbestic point is less well defined (Jayaraman *et al.*, 2000). The effect is fully reversible. In YFP-H148Q and YFP-H148G, the absorbance maximum of band A is blue-shifted by 20 nm (from 415 nm to 395 nm) upon addition of salt, though the absorbance maximum of band B is unaffected. The detailed binding equilibria and anion specificities are discussed below.

15 To further establish the usefulness of the YFPs as halide sensors, in particular for organelles that are more acidic than the cytosol (pH 7.4) where CFTR pumping was assayed (Jayaraman *et al.*, 2000), emission intensity of YFP and YFP-H148Q was measured between 0 and 150 mM NaCl at pH 6.0 to 8.0 (Fig. 8A,B), under conditions of constant ionic strength. We found that YFP constitutes an excellent

20 probe under acidic conditions. At pH 6.0, fluorescence decreased by 39% from 0 to 20 mM NaCl, whereas at the cytosolic pH of 7.5, the decrease is only 3.2% under identical conditions. For YFP-H148Q titrated with NaCl, fluorescence loss is also large at pH 6.0 (48%), and remains fairly significant (11%) at pH 7.5. For measurements of chloride concentrations in the low millimolar range near neutral pH,

YFP-H148Q appears to be the preferred probe. If iodide is substituted for chloride, YFP-H148Q fluorescence loss is much larger, even at pH 7.5 where a loss of 50% is observed (0 to 20 mM NaI) (Fig. 8B). This observation has recently been exploited in vivo, in studies of Cl⁻/I⁻ exchange by the CFTR channel in plasma membranes (Jayaraman *et al.*, 2000). In contrast to the above variant, in the original YFP the magnitude of the iodide effect is more comparable to the chloride effect (see binding data below).

D. Crystallographic identification and description of halide binding sites

10

We determined two crystal structures of YFP-H148Q, one containing two bound iodides (100 mM iodide soak), and the other containing no bound halides at all (200 mM chloride soak). The respective R-factors are 18.8% and 20.4% to a resolution of 2.1 Å, and the geometry is reasonably good. A summary of the relevant crystallographic statistics is presented in Table I. Since iodine has an anomalous signal at the in-house CuKα wavelength, an anomalous difference map was calculated for the iodide soak in order to identify heavy atom positions. We found two distinct electron density peaks at 7.7 and 5.5 rms deviations respectively, one located close to the chromophore and buried in the protein interior, the other in a small indentation on the protein surface near Trp57 at the cap of the barrel (Data not shown).

20

The buried iodide refined to an occupancy of 0.60 with the thermal factor fixed at 30 Å², indicating that binding in the crystal at 100 mM iodide is not nearly as tight as in solution, where the binding constant is 2.7 mM (see below). This iodide is located 4.3 Å away from the chromophore heterocyclic carbonyl oxygen, and is involved in a charge interaction with Arg96, with a distance of 4.1 Å to NE2 of the guanidinium group (Fig. 9 and 10), a buried positive charge that is likely providing a large fraction of the anion binding energy. Furthermore, the iodide is hydrogen bonded to both the phenolic hydroxyl of Tyr203 and the side chain amide nitrogen of Gln69, with hydrogen bonding distances of 3.3 and 3.2 Å respectively (Fig. 9 and 10). These distances are within the range of hydrogen bonding distances expected for iodide interacting with oxygen or nitrogen. A statistical database analysis of small

30

molecule crystal structures found that the mean distance between iodide and a phenolic hydroxyl is 3.47 Å, and between iodide and a sp²-hybridized nitrogen is 3.66 Å (Steiner, 1998). In a crystal structure of haloalkane dehalogenase with bound iodide, it was found that the iodide is 3.4 and 3.6 Å away from the indole nitrogens of two tryptophans, 3.3 Å from a solvent molecule, and 4.2 Å from the phenolic oxygen of a tyrosine (Verschuere *et al.*, 1993).

The buried halide also interacts with the aromatic rings of the chromophore and Tyr203 (Fig. 9). Anions are often preferentially located in or near the plane of aromatic rings, since aromatic ring hydrogens carry a partial positive charge (Burley & Petsko, 1988). In YFP-H148Q, the iodide is not quite in the plane of either of the two π systems, but is located roughly equidistant from the 2 planes, offset from the center of the stacking interaction, 4.1 Å from the aromatic CE1 of Tyr203 and 4.5 Å from the aromatic CD2 of the chromophore (Fig. 9). On the opposite side of the binding site, a series of hydrophobic residues line the halide binding site, consisting of Ile152, Leu201, Val163, Val150, and Phe165, all near van der Waals contact with the iodide. Both aromatic edge interactions with tyrosines and tryptophans, as well as apolar interactions with hydrophobic side chains are commonly found in other halide binding sites in proteins, such as in haloalkane dehalogenases (Pikkemaat *et al.*, 1999).

The second, surface-bound iodide is hydrogen-bonded to the amide nitrogen of Trp 57 and several ordered solvent molecules. This exposed anion is 16 Å from the chromophore phenolic oxygen, indicating that its influence on the chromophore charge state is negligible. The occupancy of this halide refines to 0.41 with the B-factor fixed at 39 Å², consistent weak binding as compared to the primary iodide adjacent to the chromophore.

E. Conformational changes adjacent to the buried iodide

The anion-binding pocket near the chromophore appears to be empty in the apo-structure of YFP-H148Q, in spite of the fact that the crystals were grown in the presence of 180 mM chloride, followed by soaking in 200 mM chloride. The solution

binding constant of 28 mM for chloride predicts that most of the pocket is occupied by chloride at the pH of the crystal mother liquor, 4.6 (see below). As has been found with the iodide soak, the molecules in the crystals do not appear to bind anions as tightly as in solution, possibly due to crystal packing forces. The volume of the
5 internal cavity in YFP-H148Q is $55 \text{ \AA}^3$, calculated using a probe with a radius of $1.2 \text{ \AA}$ (Connolly, 1985).

The iodide-containing cavity in the bound structure of YFP-H148Q is larger, $91 \text{ \AA}^3$, to accommodate the rather large iodide which has a van der Waals volume of $42 \text{ \AA}^3$. A series of conformational changes of various side chains lining the pocket
10 are observed (Fig. 11). The largest movement is observed by Gln69, where the side chain amide has swung out from the center of the cavity, resulting in a $2.6 \text{ \AA}$ movement of the NE2 which is hydrogen-bonded to the halide (Fig. 11 and 12). Gln183 NE2 has moved out by $1.0 \text{ \AA}$, though it is not clear whether it is a hydrogen bond donor to the iodide or Gln94 (NE2 and OE may be assigned oppositely). The
15 apolar side chains of Leu201, Ile152, Val150, and Val163 (Fig. 11) all undergo movements to increase the cavity size in the presence of iodide, with their terminal carbons (CD1 for leucine and isoleucine, CG1 for valines) shifting by $2.4 \text{ \AA}$, $1.9 \text{ \AA}$, $1.6 \text{ \AA}$, and $1.2 \text{ \AA}$ respectively. The aromatic ring plane of Phe165 has rotated by about 25° .

20 The phenolic hydroxyl of Tyr203 has shifted towards the iodide-containing cavity by $0.6 \text{ \AA}$, likely to improve the hydrogen bonding interaction with the halide (Fig.12). There appears to be some flexibility in positioning the Tyr203 side chain next to the chromophore, presumably since it protrudes into a large water-filled cavity originally identified in the structure of GFP S65T (Ormo *et al.*, 1996). An C_α -
25 carbon overlay of 5 structures of YFP and its variants (Wachter *et al.*, 1998) shows that the C_β s of Tyr203 overlay quite well, whereas the phenolic oxygen varies by up to $1.4 \text{ \AA}$. The hydrogen bond between Tyr203 and the halide appears to be of major importance in the generation of a halide binding site with reasonably tight affinity (see mutational analysis below). The chromophore shift toward the halide may also
30 serve to improve aromatic edge interactions with the anion. As a consequence, the carboxylate of Glu222 has rotated away from the chromophore ring nitrogen (distance

increases from 3.3 to 3.6 Å), and is now involved in a tight hydrogen bond to Ser205 (Fig.12).

F. Relationship between anion binding and cavity size

5

The buried iodide site in YFP-H148Q identifies a small cavity that is present in a number of structures examined and does not vary much in size (Figure 13). Calculating van der Waals volumes using a sphere with a probe radius of 1.2 Å (Connolly, 1985), the volume of this cavity is 21 Å³ in WT GFP (Brejc *et al.*, 1997), 10 19 Å³ in GFP S65T (Ormo *et al.*, 1996), 16 Å³ in YFP and YFP-H148G (Wachter *et al.*, 1998), and 21 Å³ in YFP-H148G soaked in 500 mM KBr, where the crystallographic analysis shows that the binding site is also empty (unpublished data). The position of these cavities is essentially the same in the GFPs listed above, with its center close to Val150, Val163, Leu201, Ile152, Gln183, and Gln69, but about 6.6 Å 15 distant from the chromophore methylene bridge and 6.1 Å from Arg96. WT (see below) and S65T GFP (Wachter & Remington, 1999) do not appear to interact with NaCl. On the other hand, all YFPs examined show anion interactions, with tightest Cl⁻ binding observed for YFP (see below). Clearly, cavity size and position are not directly correlated with Cl⁻ binding.

20

The cavities described above are too small to bind chloride, bromide, or iodide, whose van der Waals volumes range from 24.8 to 54 Å³. Conformational changes are clearly necessary to allow for the interaction with any anions. In the apo-structure of YFP-H148Q, the cavity is somewhat larger, even in the absence of bound anions, with a volume of 55 Å³. In this variant, the cavity is extended towards the 25 chromophore, and the volume is increased by small movements of side chain atoms (0.4 Å and 1.2 Å) lining the binding site (Gln69, Tyr203, Val150, Val163, Phe165, Arg96, His181), and the chromophore itself. Many of these residues undergo further shifts upon iodide binding, as described above and figure 6. Compensating movements of the terminal side-chain carbons of Ile152 (2.1 Å) and Leu201 (2.3 Å) 30 lead to some repacking of the hydrophobic core, without changing the adjacent cavity

surface much. The larger cavity size of YFP-H148Q may in part be responsible for the unexpectedly tight binding of iodide compared to chloride.

5 G. Relaxation of the β -barrel in response to the H148Q substitution and iodide binding

Both the introduction of H148Q in YFP background in the absence of halides, and the binding of iodide to YFP-H148Q lead to structural adjustments of β -strands 7
10 (residues 143 to 154) and 8 (residues 160 to 171). These adjustments are evident from C_{α} overlays of YFP and YFP-H148Q, with an rms deviation of 0.42 Å, and of YFP-H148Q with and without I^{-} , with an rms deviation of 0.47 Å (Figure 14). At one cap of the barrel, strands 7 and 8 are connected via a turn centered on residue 158, whereas near the other cap of the barrel, strand 7 forms a β -bulge around residue 148,
15 and main chain β -sheet interactions are disrupted (Ormo et al., 1996; Wachter et al., 1998). Instead, several ordered solvent molecules and side-chain contacts (His148 in YFP, Gln148 in YFP-H148Q) form a hydrogen bond network between the strands (Figure 14). Upon substitution of His148 with Gln, the α -carbon of residue 166 is pulled in towards the center of the barrel by 0.94 Å, and the α -carbon of residue 148
20 is pushed out by 0.94 Å. These movements are compensated for by structural adjustments within the adjacent loop regions (1.4 Å shift by the α -carbon of residue 172, and 0.94 Å by the α -carbon of residue 157). None of these residues are involved in crystal contacts in either of the two structures.

Upon binding of iodide to YFP-H148Q, Lys166 is pushed back out from the
25 center of the barrel by 1.0 Å, and is located near its original position in YFP (Figure 14). Likely, this shift in position occurs in response to the expansion of the buried cavity. Lys166 is not involved in a crystal contact in either of the two structures, whereas nearby Arg168 forms an intermolecular salt bridge with Asp149 upon iodide binding, but not without. The hydrogen bond of the Lys166 backbone oxygen to the
30 side chain of Gln148 is not disrupted by this movement. Compensating shifts are

again observed at the end of this strand, where the backbone loop residues 172 and 173 are pulled in by up to 1.7 Å, though density in this area is less well defined.

Whether halide binding to YFP has a similar effect on the β -bulge region as in YFP-H148Q is not known. Backbone movements in position 148 have been observed

5 previously in YFP-H148G (Wachter *et al.*, 1998), consistent with increased flexibility in that part of the barrel.

H. Solvent accessibility of the chromophore in YFP-H148Q

10 The structure of YFP-H148Q shows that the Gln148 side chain is swung out towards the protein exterior (Figure 9), unlike the original histidine imidazole that is hydrogen bonded to the chromophore hydroxyl (Wachter *et al.*, 1998), and constitutes a barrier to bulk solvent. Even before a structure was available, we predicted that Gln148 may be flipped out into the solvent (Elslinger *et al.*, 1999), since

15 partial chromophore exposure to exterior solvent may explain the higher pK_a of YFP-H148Q as compared to YFP (see Table K). Both in the apo and iodide-bound structure of YFP-H148Q, the Gln148 side chain amide nitrogen NE2 is hydrogen-bonded to the backbone carbonyl oxygen of Lys166, and the amide oxygen OE1 to the backbone nitrogen of Asn149 (Fig. 12 and 14), well away from the chromophore.

20 Calculations of solvent-accessible surface (Connolly, 1983) using a probe sphere radius of 1.4 Å, as implemented by MidasPlus™, show that a shallow invagination on the protein surface is formed where the wild-type imidazole of His148 was located (Fig. 13). This solvent pocket is nearly in contact with the chromophore van der Waals surface. If one considers protein breathing motions, some solvent access that is

25 not observable in the crystal structure is likely to occur. As compared to YFP-H148G (Wachter *et al.*, 1998), where the solvent channel is directly in contact with the chromophore cavity, the channel of YFP-H148Q is truncated, consistent with triplet state photobleaching experiments which suggested that the chromophore is not exposed to aqueous-phase quenchers (Jayaraman *et al.*, 2000).

I. Energetic analysis of linkage between anion and proton binding

The strong dependence of chromophore pK_a on specific anion binding can be described by a linked binding equilibrium that considers the interaction between two different ligands, the anion that binds adjacent to Arg96, and the proton that binds to the phenolic end of the chromophore. Positive cooperativity is indicated by the fact that binding of the anion facilitates binding of the proton, raising the pK_a of the chromophore. The binding constant for anion binding is therefore influenced by the amount of proton binding, and vice versa. Hence, in a simple system with one binding site each for two different ligands, one can define two microscopic binding constants, k₁ for anion binding when the proton is on and k₂ for anion binding when the proton is off. Our crystallographic analysis for YFP-H148Q is consistent with one relevant binding site for the anion, and a previous crystallographic analysis on S65T is consistent with one proton binding site on the chromophore (Elslinger *et al.*, 1999). The observed extent of anion binding is a function of pH, hence the macroscopic binding constants lie somewhere between the limiting values of k₁ and k₂.

A mathematical description has been developed by J. Wyman (1964) and is presented by Cantor and Schimmel in Biophysical Chemistry, Part III, (Cantor & Schimmel, 1980). Here, we apply the general equation 15-79 to the special case of having one binding site for each ligand, with pK_a^o representing the chromophore pK_a in the absence of any bound anions:

$$pK_a = \log \{(k_1 + [\text{chloride}]) / k_1\} - \log \{(k_2 + [\text{chloride}]) / k_2\} + pK_a^o$$

equation (1)

25

Using absorbance measurements at both pH 6.5 and pH 7.0, the pK_a of YFP was determined as a function of anion concentration for a large number of different ions. The concentration of the particular anion of interest was varied between 0 and at least 150 mM (for the halides as high as 400 mM, Fig. 15), and ionic strength was controlled by the addition of potassium gluconate which does not interact with the YFPs (Wachter & Remington, 1999). Results for interacting anions were fit to

30

equation (1), and the microscopic binding constants k_1 , and k_2 where possible, were extracted from the curve fit (Figure 15 and Table J). In general, small monovalent anions appear to show some interaction with YFP. Binding is tightest for fluoride, with $k_1 = 0.214$ mM. Other anions that were found to interact, including the other
 5 halides, have microscopic binding constants in the low millimolar range, with trifluoroacetic acid (TFA) giving the weakest interaction in this series ($k_1 = 21.2$ mM).

There does not appear to be any particular molecular shape dependence for this interaction, since triatomic linear (e.g. thiocyanate), square planar molecules (e.g.
 10 perchlorate), trigonal (e.g. nitrite), and spherical (e.g. halides) molecules are also found to bind. Formate modulated the chromophore pK_a as well ($k_1 = 7.47$ mM), though a previous, somewhat preliminary experiment by fluorescence indicated no interaction (Wachter & Remington, 1999). As expected, anion binding to the anionic chromophore is unfavorable, with k_2 in the high millimolar or in the molar range,
 15 often outside the range of measurement (Table J).

Table J. Anion binding to the YFP chromophore in order of decreasing interaction strength.^a

Interacting anions	k_1 (mM) ^b	k_2 (mM) ^b
Fluoride F ⁻	0.214 (0.009)	301 (64)
Thiocyanate SCN ⁻	1.37 (0.02)	large ^c
Perchlorate ClO ₄ ⁻	1.46 (0.36)	175 (11)
Nitrite NO ₂ ⁻	2.12 (0.40)	273 (200)
Iodide I ⁻	2.46 (0.11)	325 (64)
Nitrate NO ₃ ⁻	4.44 (0.25)	large ^c
Chloride Cl ⁻	4.69 (0.17)	288 (40)

Formate HCOO^-	7.47 (0.004)	large ^c
Bromide Br^-	7.76 (1.00)	280 (126)
TFA CF_3COO^-	21.2 (3.7)	large ^c

^aConjugate bases (prevalent ion at pH 6 to 7) are listed in order of decreasing interaction strength.

^bThe numbers in parenthesis are a lower estimate of the standard deviation as determined by Kaleidagraph™.

5 ^cThese binding constants could not be determined since they fall outside the range of measurement, and are likely in the molar range.

Divalent anions such as phosphate and sulfate, and larger monovalent anions such as gluconate, Good buffers (e.g. HEPES, PIPES), isethionate (2-
10 hydroxyethanesulfonic acid), and TCA (trichloroacetic acid), do not interact (Table J), as indicated by a constant pK_a of about 5.4 for YFP, essentially the same as when measurements are carried out in low ionic strength buffers without addition of salts (Wachter & Remington, 1999). Somewhat smaller monovalent anions that do not
15 interact include phosphoric acid, bicarbonate, and acetate. The hydration energy may be of importance in discrimination of anions, since acetate is strongly solvated, whereas TCA is only weakly hydrated in aqueous solvents (March, 1992). The series presented for YFP in Table J is very similar to the one determined for YFP-H148Q by fluorescence at pH 7.5 (Jayaraman *et al.*, 2000), with only minor
20 differences in ordering. For example, YFP-H148Q binds Br^- more strongly than Cl^- , whereas for YFP, the order is reversed, likely due to the larger binding site in YFP-H148Q.

J. Identification of key residues for anion binding by mutational analysis

To identify which substitutions in YFP (S65G/V68L/S72A/T203Y) are contributors to specific anion binding near the chromophore, we carried out a mutational analysis, converting the four substitutions back to wild-type one-by-one. We then determined the pK_a of these revertants in the absence of interacting anions, and measured their affinity to chloride and iodide by pK_a determination as a function of halide concentration, followed by curve fitting to equation (1). Revertant 1 (S65G/S72A/T203Y) and revertant 2 (S65G/V68L/S72A) exhibited well-behaved pH and halide titration behavior as is observed for the YFPs, and their pK_a and k_1 for chloride and iodide binding are compared with those obtained for the YFPs in Table K. Reversion of residue 68 or residue 203 raises the chromophore pK_a to 5.8 and 6.4 respectively. Reversion of residue 68 leads to a slight loss of chloride affinity ($k_1 = 13.2$ mM, as compared to 4.69 mM in YFP), whereas reversion of residue 203 dramatically weakens the interaction ($k_1 = 153$ mM). As is evident from Table K, chloride affinity is strongly coupled to chromophore pK_a , with a weakening of the anion interaction with increasing pK_a .

Table K Microscopic dissociation constants for chloride and iodide binding to the YFPs and its revertants.

variant	substitutions	k_1 (mM) ^a for Cl ⁻	k_1 (mM) ^a for I ⁻	pK_a ^b
YFP	S65G/V68L/S72A/T203Y	4.69 (0.17)	2.46 (0.11)	5.4
revertant 1	S65G/S72A/T203Y	13.2 (0.34)	3.04 (0.11)	5.8
YFP-H148Q	S65G/V68L/S72A/H148Q/T203Y	28.4 (5.1)	2.68 (0.11)	6.7
YFP-H148G	S65G/V68L/S72A/H148G/T203Y	82.8 (18.3)	15.73 (2.6)	7.5
revertant 2	S65G/V68L/S72A	153 (26)	117 (16)	6.4

^aA lower estimate of the standard deviation (as reported by Kaleidagraph™) is given in parenthesis.

^bThe chromophore pK_a determined by absorbance in the absence of any interfering anions, such as chloride (buffered with either HEPES or PIPES, 150 mM gluconate).

5

Since the only exception to this rule is revertant 2, it appears that the correlation is intact only in the presence of T203Y. This substitution appears to be indispensable for strong anion interactions.

Iodide binding appears to be considerably tighter than chloride binding for all variants tested (Table K). Any correlation with chromophore pK_a is weak at best. The relative selectivity of iodide over chloride is strongest for YFP-H148Q, followed by YFP-H148G. This may reflect the fact that iodide is a larger, softer ion than chloride, more difficult to fit into a small cavity unless the particular variant allows for structural relaxation of the β -barrel (see above).

15 Revertants 3 (S72A/T203Y) and 4 (T203Y) were more difficult to analyze, since their titration behavior is similar to WT GFP. Their absorbance spectra exhibit a mixed ground state of bands A and B, and are nearly pH-independent above pH 6.5. Excitation of either band A or B leads to green fluorescence in these revertants, reminiscent of the excited-state deprotonation described for WT GFP (Chattoraj *et al.*,
20 1996). Addition of NaCl to 250 mM to revertants 3 and 4 at pH 6.5 changes the ratio of the two absorbance bands only to a small degree, resulting in roughly a 20% decrease of band B in favor of band A. In WT GFP at pH 6.5, no spectral change is observed upon addition of 250 mM NaCl under conditions of constant ionic strength, consistent with a sensitivity towards ionic strength (Ward *et al.*, 1982) but not
25 specific anion binding.

The present invention provides novel long wavelength engineered fluorescent proteins. While specific examples have been provided, the above description is illustrative and not restrictive. Many variations of the invention will become apparent to those skilled in the art upon review of this specification. The scope of the
5 invention should, therefore, be determined not with reference to the above description, but instead should be determined with reference to the appended claims along with their full scope of equivalents.

Publications

- Baird, G. S., Zacharias, D. A. & Tsien, R. Y. (1999). Circular permutation and
10 receptor insertion within green fluorescent proteins. *Proc. Natl. Acad. Sci. USA* **96**, 11241-11246.
- Bell, A. F., He, X., Wachter, R. M. & Tonge, P. J. (2000). Probing the ground state structure of the green fluorescent protein chromophore using Raman spectroscopy. *Biochemistry* (in press).
- 15 Bokman, S. H. & Ward, W. W. (1981). Renaturation of *Aequorea* green fluorescent protein. *Biochem. Biophys. Res. Commun.* **101**, 1372-1380.
- Born, M. (1920). Volumen und hydrationswaerme der ionen. *Z. Physik* **1**, 45-48.
- Brejc, K., Sixma, T. K., Kitts, P. A., Kain, S. R., Tsien, R. Y., Ormö, M. & Remington, S. J. (1997). Structural basis for dual excitation and
20 photoisomerization of the *Aequorea victoria* Green Fluorescent Protein. *Proc. Natl. Acad. Sci. USA* **94**, 2306-2311.
- Burley, S. K. & Petsko, G. A. (1988). Weakly polar interactions in proteins. *Adv. Protein Chem.* **39**, 125-189.
- Cantor, C. R. & Schimmel, P. R. (1980). *Biophysical Chemistry Part III: The behavior of biological macromolecules*, W. H. Freeman and Company, New
25 York, pp 866-878.

- Chattoraj, M., King, B. A., Bublit, G. U. & Boxer, S. G. (1996). Ultra-fast excited state dynamics in Green Fluorescent Protein: Multiple states and proton transfer. *Proc. Natl. Acad. Sci. USA* **93**, 8362-8367.
- Collaborative Computational Project N. 4 (1994). The CCP4 Suite: Programs for protein crystallography. *Acta Cryst.* **D50**, 760-763.
- Connolly, M. L. (1983). Solvent-accessible surfaces of proteins and nucleic acids. *Science* **221**(4612), 709-713.
- Connolly, M. L. (1985). Computation of molecular volume. *J. Am. Chem. Soc.* **107**, 1118-1124.
- De Giorgi, F., Ahmed, Z., Bastianutto, C., Brini, M., Jouvaille, L. S., Marsault, L. S., Murgia, M., Pinton, P., Pozzan, T. & Rizzutto, R. (1999). *Meth. Cell Biol.* **58**, 75-85.
- Doi, N. & Yanagawa, H. (1999). Design of generic biosensors based on green fluorescent proteins with allosteric sites by directed evolution. *FEBS Lett.* **453**, 305-307.
- Elsiger, M.-A., Wachter, R. M., Hanson, G. T., Kallio, K. & Remington, S. J. (1999). Structural and spectral response of green fluorescent protein variants to changes in pH. *Biochemistry* **38**, 5296-5301.
- Feller, G., le Bussy, O., Houssier, C. & Gerday, C. (1996). Structural and functional aspects of chloride binding to *Alteromonas haloplanctis* alpha-amylase. *J. Biol. Chem.* **271**, 23836-23841.
- Fersht, A. (1985). *Enzyme structure and mechanism*, W. H. Freeman and Company, New York, NY, pg 298.
- Halm, D. R. & Frizzell, R. A. (1992). Anion permeation in an apical membrane chloride channel of a secretory epithelial cell. *J. Gen. Physiol.* **99**, 339-366.
- Heim, R. (1999). Green fluorescent protein forms for energy transfer. *Methods Enzymol.* **302**, 408-423.

- Heim, R., Prasher, D. C. & Tsien, R. Y. (1994). Wavelength mutations and posttranslational autooxidation of green fluorescent protein. *Proc. Natl. Acad. Sci. USA* **91**, 12501-12504.
- Jayaraman, S., Haggie, P., Wachter, R. M., Remington, S. J. & Verkman, A. S. (2000). Mechanism and cellular applications of a green fluorescent protein-based halide sensor. *J. Biol. Chem.* (in press).
- Jones, T. A., Zou, J.-Y., Cowan, S. W. & Kjeldgaard, M. (1991). Improved methods for building protein models in electron density maps and the location of errors in these models. *Acta Crystallogr. Sect. A* **47**, 110-119.
- 10 Kneen, M., Farinas, J., Li, Y. & Verkman, A. S. (1998). Green fluorescent protein as a noninvasive intracellular pH indicator. *Biophys. J.* **74**, 1591-1599.
- Kraulis, P. (1991). MOLSCRIPT: A program to produce both detailed and schematic plots of protein structures. *J. Appl. Crystallog.* **24**, 946-950.
- Llopis, J., McCaffery, J. M., Miyawaki, A., Farquhar, M. & Tsien, R. Y. (1998).
15 Measurement of cytosolic, mitochondrial, and Golgi pH in single living cells with green fluorescent proteins. *Proc. Natl. Acad. Sci. USA* **95**, 6803-6808.
- March, J. (1992). *Advanced Organic Chemistry*, John Wiley & Sons, New York, pg 272.
- Matz, M. V., Fradkov, A. F., Labas, Y. A., Savitsky, A. P., Zaraisky, A. G.,
20 Markelov, M. L. & Lukyanov, S. A. (1999). Fluorescent proteins from nonbioluminescent Anthozoa species. *Nature Biotechnol.* **17**, 969-973.
- Miesenbock, G., De Angelis, D. A. & Rothman, J. E. (1998). Visualizing secretion and synaptic transmission with pH-sensitive green fluorescent proteins. *Nature* **394**, 192-195.
- 25 Miyawaki, A., Griesbeck, O., Heim, R. & Tsien, R. Y. (1999). Dynamic and quantitative Ca²⁺ measurements using improved cameleons. *Proc. Natl. Acad. Sci. USA* **96**, 2135-2140.

- Ormö, M., Cubitt, A. B., Kallio, K., Gross, L. A., Tsien, R. Y. & Remington, S. J. (1996). Crystal structure of the *Aequorea victoria* Green Fluorescent Protein. *Science* **273**, 1392-1395.
- Otwinowski, Z. & Minor, W. (1997). Processing of X-ray diffraction data collected in oscillation mode. *Methods Enzymol.* **276**, 307-326.
- Palm, G. J. & Wlodawer, A. (1999). Spectral variants of green fluorescent protein. *Methods Enzymol.* **302**, 378-394.
- Palm, G. J., Zdanov, A., Gaitanaris, G. A., Stauber, R., Pavlakis, G. N. & Wlodawer, A. (1997). The structural basis for spectral variations in green fluorescent protein. *Nat. Struct. Biol.* **4**(5), 361-365.
- Pikkemaat, M. G., Ridder, I. S., Rozeboom, H. J., Kalk, K. H., Dijkstra, B. W. & Janssen, D. B. (1999). Crystallographic and kinetic evidence of a collision complex formed during halide import in haloalkane dehalogenase. *Biochemistry* **38**, 12052-12061.
- Robey, R. B., Ruiz, O., Santos, A. V. P., Ma, J., Kear, F., Wang, L.-J., Li, C.-J., Bernardo, A. A. & Arruda, J. A. L. (1998). pH-Dependent fluorescence of a heterologously expressed *Aequorea* Green Fluorescent Protein mutant: In situ spectral characteristics and applicability to intracellular pH estimation. *Biochemistry* **37**, 9894-9901.
- Scharnagl, C., Raupp-Kossmann, R. & Fischer, S. F. (1999). Molecular basis for pH sensitivity and proton transfer in Green Fluorescent Protein: Protonation and Conformational substates from electrostatic calculations. *Biophys. J.* **77**, 1839-1857.
- Steiner, T. (1998). Hydrogen-bond distances to halide ions in organic and organometallic crystal structures: Up-to-date database study. *Acta Cryst.* **B54**, 456-463.

- Sullivan, K. F. & Kay, S. A. E., eds (1999). Green fluorescent proteins. *Methods Cell Biol.* **58**, Academic Press, San Diego.
- Topell, S., Hennecke, J. & Glockshuber, R. (1999). Circularly permuted variants of the green fluorescent protein. *FEBS Lett.* **457**, 283-289.
- 5 Tronrud, D. E., Ten Eyck, L. F. & Matthews, B. W. (1987). An efficient general-purpose least-squares refinement program for macromolecular structures. *Acta Crystallogr. Sect. A* **43**, 489-501.
- Tsien, R. Y. (1998). The green fluorescent protein. *Ann. Rev. Biochem.* **67**, 509-544.
- Verschueren, K. H. G., Kingma, J., Rozeboom, H. J., Kalk, K. H., Janssen, D. B. & Dijkstra, B. W. (1993). Crystallographic and fluorescence studies of the
10 interaction of haloalkane dehalogenase with halide ions. Studies with halide compounds reveal a halide binding site in the active site. *Biochemistry* **32**, 9031-9037.
- Wachter, R. M., Elsliger, M.-A., Kallio, K., Hanson, G. T. & Remington, S. J. (1998).
15 Structural basis of spectral shifts in the yellow-emission variants of green fluorescent protein. *Structure* **6**(10), 1267-1277.
- Wachter, R. M. & Remington, S. J. (1999). Sensitivity of the yellow variant of green fluorescent protein to halides and nitrate. *Curr. Biol.* **9**(17), R628-R629.
- Wang, Z., Asenjo, A. B. & Oprian, D. D. (1993). Identification of the Cl⁻ binding site
20 in the human red and green color vision pigments. *Biochemistry* **32**, 2125-2130.
- Ward, W. W., Prentice, H. J., Roth, A. F., Cody, C. W. & Reeves, S. C. (1982). Spectral perturbations of the *Aequorea* green-fluorescent protein. *Photochem. Photobiol.* **35**, 803-808.
- Wright, E. M. & Diamond, J. M. (1977). Anion selectivity in biological systems.
25 *Physiological Reviews* **57**, 109-156.
- Yang, F., Moss, L. G. & Phillips, G. N. (1996). The molecular structure of green fluorescent protein. *Nature Biotech.* **14**, 1246-1251.

Yuan, C., Kuwata, O., Liang, J., Misra, S., Balashov, S. P. & Ebrey, T. G. (1999). Chloride binding regulates the Schiff base pK in Gecko P521 cone-type visual pigment. *Biochemistry* **38**, 4649-4654.

All publications and patent documents cited in this application are
5 incorporated by reference in their entirety for all purposes to the same extent as if
each individual publication or patent document were so individually denoted.

WHAT IS CLAIMED IS:

1. A nucleic acid molecule comprising a nucleotide sequence encoding a functional engineered fluorescent protein whose amino acid sequence is substantially identical to the amino acid sequence of *Aequorea* green fluorescent protein (SEQ ID NO:2) and which differs from SEQ ID NO:2 by at least the substitution T203X, wherein X is an aromatic amino acid selected from H, Y, W or F, said functional engineered fluorescent protein having a different fluorescent property than *Aequorea* green fluorescent protein.
2. The nucleic acid molecule of claim 1 wherein the amino acid sequence further comprises a substitution at S65, wherein the substitution is selected from S65G, S65T, S65A, S65L, S65C, S65V and S65I.
3. The nucleic acid molecule of claim 1 wherein the amino acid sequence differs by no more than the substitutions S65T/T203H; S65T/T203Y; S72A/F64L/S65G/T203Y; S72A/S65G/V68L/T203Y; S65G/V68L/Q69K/S72A/T203Y; S65G/S72A/T203Y; or S65G/S72A/T203W.
4. The nucleic acid molecule of claim 1 or 2 wherein the amino acid sequence further comprises a substitution at Y66, wherein the substitution is selected from Y66H, Y66F, and Y66W.
5. The nucleic acid molecule of claim 1 or 2 wherein the amino acid sequence further comprises a mutation from Table A.
6. The nucleic acid molecule of claim 1 or 2 wherein the amino acid sequence further comprises a folding mutation.
7. The nucleic acid molecule of any of claims 1-3 wherein the nucleotide sequence encoding the protein differs from the nucleotide sequence of SEQ ID NO:1 by the substitution of at least one codon by a preferred mammalian codon.

8. The nucleic acid molecule of any of claims 1-3 encoding a fusion protein wherein the fusion protein comprises a polypeptide of interest and the functional engineered fluorescent protein.
9. An expression vector comprising expression control sequences operatively linked to a nucleic acid molecule comprising a nucleotide sequence encoding a functional engineered fluorescent protein whose amino acid sequence is substantially identical to the amino acid sequence of *Aequorea* green fluorescent protein (SEQ ID NO:2) and which differs from SEQ ID NO:2 by at least the amino acid substitution T203X, wherein X is an aromatic amino acid selected from H, Y, W or F, said functional engineered fluorescent protein having a different fluorescent property than *Aequorea* green fluorescent protein.
10. The expression vector of claim 9 wherein the amino acid sequence further comprises a substitution at S65, wherein the substitution is selected from S65G, S65T, S65A, S65L, S65C, S65V and S65I.
11. The expression vector of claim 9 wherein the amino acid sequence differs by no more than the substitutions S65T/T203H; S65T/T203Y; S72A/F64L/S65G/T203Y; S72A/S65G/V68L/T203Y; S65G/V68L/Q69K/S72A/T203Y, S65G/S72A/T203Y; or S65G/S72A/T203W.
12. The expression vector of claim 10 or 11 wherein the amino acid sequence further comprises a substitution at Y66, wherein the substitution is selected from Y66H, Y66F, and Y66W.
13. The expression vector of claim 10 or 11 wherein the amino acid sequence further comprises a mutation from Table A.
14. The expression vector of claim 9 or 10 wherein the amino acid sequence further comprises a folding mutation.

15. The expression vector of any of claims 9-11 wherein the nucleotide sequence encoding the protein differs from the nucleotide sequence of SEQ ID NO:1 by the substitution of at least one codon by a preferred mammalian codon.
16. The expression vector of any of claims 9-11 encoding a fusion protein wherein the fusion protein comprises a polypeptide of interest and the functional engineered fluorescent protein.
17. A recombinant host cell comprising an expression vector that comprises expression control sequences operatively linked to a nucleic acid molecule comprising a nucleotide sequence encoding a functional engineered fluorescent protein whose amino acid sequence is substantially identical to the amino acid sequence of *Aequorea* green fluorescent protein (SEQ ID NO:2) and which differs from SEQ ID NO:2 by at least the amino acid substitution T203X, wherein X is an aromatic amino acid selected from H, Y, W or F, said functional engineered fluorescent protein having a different fluorescent property than *Aequorea* green fluorescent protein.
18. The recombinant host cell of claim 17 wherein the amino acid sequence further comprises a substitution at S65, wherein the substitution is selected from S65G, S65T, S65A, S65L, S65C, S65V and S65I.
19. The recombinant host cell of claim 17 wherein the amino acid sequence differs by no more than the substitutions S65T/T203H; S65T/T203Y; S72A/F64L/S65G/T203Y; S72A/S65G/V68L/T203Y; S65G/V68L/Q69K/S72A/T203Y; S65G/S72A/T203Y; or S65G/S72A/T203W.
20. The recombinant host cell of claim 17 or 18 wherein the amino acid sequence further comprises a substitution at Y66, wherein the substitution is selected from Y66H, Y66F, and Y66W.

21. The recombinant host cell of claim 17 or 18 wherein the amino acid sequence further comprises a mutation from Table A.
22. The recombinant host cell of claim 17 or 18 wherein the amino acid sequence further comprises a folding mutation.
23. The recombinant host cell of any of claims 17-19 wherein the nucleotide sequence encoding the protein differs from the nucleotide sequence of SEQ ID NO:1 by the substitution of at least one codon by a preferred mammalian codon.
24. The recombinant host cell of any of claims 17-19 encoding a fusion protein wherein the fusion protein comprises a polypeptide of interest and the functional engineered fluorescent protein.
25. The recombinant host cell of any of claims 17-19 which is a prokaryotic cell.
26. The recombinant host cell of any of claims 17-19 which is a eukaryotic cell.
27. A functional engineered fluorescent protein whose amino acid sequence is substantially identical to the amino acid sequence of *Aequorea* green fluorescent protein (SEQ ID NO:2) and which differs from SEQ ID NO:2 by at least the amino acid substitution T203X, wherein X is an aromatic amino acid selected from H, Y, W or F, said functional engineered fluorescent protein having a different fluorescent property than *Aequorea* green fluorescent protein.
28. The protein of claim 27 wherein the amino acid sequence further comprises a substitution at S65, wherein the substitution is selected from S65G, S65T, S65A, S65L, S65C, S65V and S65I.

29. The protein of claim 27 wherein the amino acid sequence differs by no more than the substitutions S65T/T203H; S65T/T203Y; S72A/F64L/S65G/T203Y; S72A/S65G/V68L/T203Y; S65G/V68L/Q69K/S72A/T203Y; S65G/S72A/T203Y; or S65G/S72A/T203W.
30. The protein of claim 27 or 28 wherein the amino acid sequence further comprises a substitution at Y66, wherein the substitution is selected from Y66H, Y66F, and Y66W.
31. The protein of claim 27 or 28 wherein the amino acid sequence further comprises a folding mutation.
32. The protein of any of claims 27-29 which is a fusion protein wherein the fusion protein comprises a polypeptide of interest and the functional engineered fluorescent protein.
33. A fluorescently labelled antibody comprising an antibody coupled to a functional engineered fluorescent protein whose amino acid sequence is substantially identical to the amino acid sequence of *Aequorea* green fluorescent protein (SEQ ID NO:2) and which differs from SEQ ID NO:2 by at least the amino acid substitution T203X, wherein X is an aromatic amino acid selected from H, Y, W or F, said functional engineered fluorescent protein having a different fluorescent property than *Aequorea* green fluorescent protein.
34. The fluorescently labelled antibody of claim 33 wherein the amino acid sequence further comprises a substitution at S65, wherein the substitution is selected from S65G, S65T, S65A, S65L, S65C, S65V and S65I.
35. The fluorescently labelled antibody of claim 33 wherein the amino acid sequence differs by no more than the substitutions S65T/T203H; S65T/T203Y; S72A/F64L/S65G/T203Y; S72A/S65G/V68L/T203Y; S65G/V68L/Q69K/S72A/T203Y; S65G/S72A/T203Y; or S65G/S72A/T203W.

36. The fluorescently labelled antibody of claim 33 or 34 wherein the amino acid sequence further comprises a substitution at Y66, wherein the substitution is selected from Y66H, Y66F, and Y66W.

37. The fluorescently labelled antibody of any of claims 33-35 which is a fusion protein wherein the fusion protein comprises the antibody fused to the functional engineered fluorescent protein.

38. A nucleic acid molecule comprising a nucleotide sequence encoding an antibody fused to a nucleotide sequence encoding a functional engineered fluorescent protein whose amino acid sequence is substantially identical to the amino acid sequence of *Aequorea* green fluorescent protein (SEQ ID NO:2) and which differs from SEQ ID NO:2 by at least the amino acid substitution T203X, wherein X is an aromatic amino acid selected from H, Y, W or F, said functional engineered fluorescent protein having a different fluorescent property than *Aequorea* green fluorescent protein.

39. The nucleic acid molecule of claim 38 wherein the amino acid sequence further comprises a substitution at S65, wherein the substitution is selected from S65G, S65T, S65A, S65L, S65C, S65V and S65I.

40. The nucleic acid molecule of claim 38 wherein the amino acid sequence differs by no more than the substitutions S65T/T203H; S65T/T203Y; S72A/F64L/S65G/T203Y; S72A/S65G/V68L/T203Y; S65G/V68L/Q69K/S72A/T203Y; S65G/S72A/T203Y; or S65G/S72A/T203W.

41. The nucleic acid molecule of claim 38 or 39 wherein the amino acid sequence further comprises a substitution at Y66, wherein the substitution is selected from Y66H, Y66F, and Y66W.

42. A fluorescently labelled nucleic acid probe comprising a nucleic acid probe coupled to a functional engineered fluorescent protein whose amino acid sequence is substantially identical to the amino acid sequence of *Aequorea* green fluorescent protein (SEQ ID NO:2) and which differs from SEQ ID NO:2 by at least the amino acid substitution T203X, wherein X is an aromatic amino acid selected from H, Y, W or F, said functional engineered fluorescent protein having a different fluorescent property than *Aequorea* green fluorescent protein.

43. The fluorescently labelled nucleic acid probe of claim 42 wherein the amino acid sequence further comprises a substitution at S65, wherein the substitution is selected from S65G, S65T, S65A, S65L, S65C, S65V and S65I.

44. The fluorescently labelled nucleic acid probe of claim 42 wherein the amino acid sequence differs by no more than the substitutions S65T/T203H; S65T/T203Y; S72A/F64L/S65G/T203Y; S72A/S65G/V68L/T203Y; S65G/V68L/Q69K/S72A/T203Y; S65G/S72A/T203Y; or S65G/S72A/T203W.

45. The nucleic acid molecule of claim 42 or 43 wherein the amino acid sequence further comprises a substitution at Y66, wherein the substitution is selected from Y66H, Y66F, and Y66W.

46. A nucleic acid molecule comprising a nucleotide sequence encoding a functional engineered fluorescent protein whose amino acid sequence is substantially identical to the amino acid sequence of *Aequorea* green fluorescent protein (SEQ ID NO:2) and which differs from SEQ ID NO:2 by at least an amino acid substitution at L42, V61, T62, V68, Q69, Q94, N121, Y145, H148, V150, F165, I167, Q183, N185, L220, E222 (not E222G), or V224, said functional engineered fluorescent protein having a different fluorescent property than *Aequorea* green fluorescent protein.

47. The nucleic acid molecule of claim 46 wherein the amino acid substitution is:

L42X, wherein X is selected from C, F, H, W and Y,

V61X, wherein X is selected from F, Y, H and C,

T62X, wherein X is selected from A, V, F, S, D, N, Q, Y, H and C,

V68X, wherein X is selected from F, Y and H,

Q69X, wherein X is selected from K, R, E and G,

Q94X, wherein X is selected from D, E, H, K and N,

N121X, wherein X is selected from F, H, W and Y,

Y145X, wherein X is selected from W, C, F, L, E, H, K and Q,

H148X, wherein X is selected from F, Y, N, K, Q and R,

V150X, wherein X is selected from F, Y and H,

F165X, wherein X is selected from H, Q, W and Y,

I167X, wherein X is selected from F, Y and H,

Q183X, wherein X is selected from H, Y, E and K,

N185X, wherein X is selected from D, E, H, K and Q,

L220X, wherein X is selected from H, N, Q and T,

E222X, wherein X is selected from N and Q or

V224X, wherein X is selected from H, N, Q, T, F, W and Y.

48. An expression vector comprising expression control sequences operatively linked to a nucleic acid molecule of comprising a nucleotide sequence encoding a functional engineered fluorescent protein whose amino acid sequence is substantially identical to the amino acid sequence of *Aequorea* green fluorescent protein (SEQ ID NO:2) and which differs from SEQ ID NO:2 by at least an amino acid substitution at L42, V61, T62, V68, Q69, Q94, N121, Y145, H148, V150, F165, I167, Q183, N185, L220, E222 (not E222G), or V224, said functional engineered fluorescent protein having a different fluorescent property than *Aequorea* green fluorescent protein.

49. The expression vector of claim 48 wherein the amino acid substitution is:

L42X, wherein X is selected from C, F, H, W and Y,

V61X, wherein X is selected from F, Y, H and C,

T62X, wherein X is selected from A, V, F, S, D, N, Q, Y, H and C,

V68X, wherein X is selected from F, Y and H,

Q69X, wherein X is selected from K, R, E and G,

Q94X, wherein X is selected from D, E, H, K and N,

N121X, wherein X is selected from F, H, W and Y,

Y145X, wherein X is selected from W, C, F, L, E, H, K and Q,

H148X, wherein X is selected from F, Y, N, K, Q and R,

V150X, wherein X is selected from F, Y and H,

F165X, wherein X is selected from H, Q, W and Y,

I167X, wherein X is selected from F, Y and H,

Q183X, wherein X is selected from H, Y, E and K,

N185X, wherein X is selected from D, E, H, K and Q,

L220X, wherein X is selected from H, N, Q and T,

E222X, wherein X is selected from N and Q or

V224X, wherein X is selected from H, N, Q, T, F, W and Y.

50. A recombinant host cell comprising an expression vector that comprises expression control sequences operatively linked to a nucleic acid molecule comprising a nucleotide sequence encoding a functional engineered fluorescent protein whose amino acid sequence is substantially identical to the amino acid sequence of *Aequorea* green fluorescent protein (SEQ ID NO:2) and which differs from SEQ ID NO:2 by at least an amino acid substitution at L42, V61, T62, V68, Q69, Q94, N121, Y145, H148, V150, F165, I167, Q183, N185, L220, E222 (not E222G), or V224, said functional engineered fluorescent protein having a different fluorescent property than *Aequorea* green fluorescent protein.

51. The recombinant host cell of claim 50 wherein the amino acid substitution is:
L42X, wherein X is selected from C, F, H, W and Y,
V61X, wherein X is selected from F, Y, H and C,
T62X, wherein X is selected from A, V, F, S, D, N, Q, Y, H and C,
V68X, wherein X is selected from F, Y and H,
Q69X, wherein X is selected from K, R, E and G,
Q94X, wherein X is selected from D, E, H, K and N,
N121X, wherein X is selected from F, H, W and Y,
Y145X, wherein X is selected from W, C, F, L, E, H, K and Q,
H148X, wherein X is selected from F, Y, N, K, Q and R,
V150X, wherein X is selected from F, Y and H,
F165X, wherein X is selected from H, Q, W and Y,
I167X, wherein X is selected from F, Y and H,
Q183X, wherein X is selected from H, Y, E and K,
N185X, wherein X is selected from D, E, H, K and Q,
L220X, wherein X is selected from H, N, Q and T,
E222X, wherein X is selected from N and Q or
V224X, wherein X is selected from H, N, Q, T, F, W and Y.
52. A functional engineered fluorescent protein whose amino acid sequence is substantially identical to the amino acid sequence of *Aequorea* green fluorescent protein (SEQ ID NO:2) and which differs from SEQ ID NO:2 by at least an amino acid substitution at L42, V61, T62, V68, Q69, Q94, N121, Y145, H148, V150, F165, I167, Q183, N185, L220, E222 (E222G), or V224, said functional engineered fluorescent protein having a different fluorescent property than *Aequorea* green fluorescent protein.

53. The functional engineered fluorescent protein of claim 52 wherein the amino acid substitution is:

L42X, wherein X is selected from C, F, H, W and Y,
V61X, wherein X is selected from F, Y, H and C,
T62X, wherein X is selected from A, V, F, S, D, N, Q, Y, H and C,
V68X, wherein X is selected from F, Y and H,
Q69X, wherein X is selected from K, R, E and G,
Q94X, wherein X is selected from D, E, H, K and N,
N121X, wherein X is selected from F, H, W and Y,
Y145X, wherein X is selected from W, C, F, L, E, H, K and Q,
H148X, wherein X is selected from F, Y, N, K, Q and R,
V150X, wherein X is selected from F, Y and H,
F165X, wherein X is selected from H, Q, W and Y,
I167X, wherein X is selected from F, Y and H,
Q183X, wherein X is selected from H, Y, E and K,
N185X, wherein X is selected from D, E, H, K and Q,
L220X, wherein X is selected from H, N, Q and T,
E222X, wherein X is selected from N and Q or
V224X, wherein X is selected from H, N, Q, T, F, W and Y.

54. A fluorescently labelled antibody comprising an antibody coupled to a functional engineered fluorescent protein whose amino acid sequence is substantially identical to the amino acid sequence of *Aequorea* green fluorescent protein (SEQ ID NO:2) and which differs from SEQ ID NO:2 by at least an amino acid substitution at L42, V61, T62, V68, Q69, Q94, N121, Y145, H148, V150, F165, I167, Q183, N185, L220, E222 (not E222G), or V224, said functional engineered fluorescent protein having a different fluorescent property than *Aequorea* green fluorescent protein.

55. The antibody of claim 54 wherein the amino acid substitution is:

L42X, wherein X is selected from C, F, H, W and Y,

V61X, wherein X is selected from F, Y, H and C,

T62X, wherein X is selected from A, V, F, S, D, N, Q, Y, H and C,

V68X, wherein X is selected from F, Y and H,

Q69X, wherein X is selected from K, R, E and G,

Q94X, wherein X is selected from D, E, H, K and N,

N121X, wherein X is selected from F, H, W and Y,

Y145X, wherein X is selected from W, C, F, L, E, H, K and Q,

H148X, wherein X is selected from F, Y, N, K, Q and R,

V150X, wherein X is selected from F, Y and H,

F165X, wherein X is selected from H, Q, W and Y,

I167X, wherein X is selected from F, Y and H,

Q183X, wherein X is selected from H, Y, E and K,

N185X, wherein X is selected from D, E, H, K and Q,

L220X, wherein X is selected from H, N, Q and T,

E222X, wherein X is selected from N and Q or

V224X, wherein X is selected from H, N, Q, T, F, W and Y.

56. A nucleic acid molecule comprising a nucleotide sequence encoding an antibody fused to a nucleotide sequence encoding a functional engineered fluorescent protein whose amino acid sequence is substantially identical to the amino acid sequence of *Aequorea* green fluorescent protein (SEQ ID NO:2) and which differs from SEQ ID NO:2 by at least an amino acid substitution at L42, V61, T62, V68, Q69, Q94, N121, Y145, H148, V150, F165, I167, Q183, N185, L220, E222 (not E222G), or V224, said functional engineered fluorescent protein having a different fluorescent property than *Aequorea* green fluorescent protein.

57. The nucleic acid molecule of claim 56 wherein the amino acid substitution is:

L42X, wherein X is selected from C, F, H, W and Y,

V61X, wherein X is selected from F, Y, H and C,

T62X, wherein X is selected from A, V, F, S, D, N, Q, Y, H and C,

V68X, wherein X is selected from F, Y and H,

Q69X, wherein X is selected from K, R, E and G,

Q94X, wherein X is selected from D, E, H, K and N,

N121X, wherein X is selected from F, H, W and Y,

Y145X, wherein X is selected from W, C, F, L, E, H, K and Q,

H148X, wherein X is selected from F, Y, N, K, Q and R,

V150X, wherein X is selected from F, Y and H,

F165X, wherein X is selected from H, Q, W and Y,

I167X, wherein X is selected from F, Y and H,

Q183X, wherein X is selected from H, Y, E and K,

N185X, wherein X is selected from D, E, H, K and Q,

L220X, wherein X is selected from H, N, Q and T,

E222X, wherein X is selected from N and Q or

V224X, wherein X is selected from H, N, Q, T, F, W and Y.

58. A fluorescently labelled nucleic acid probe comprising a nucleic acid probe coupled to a functional engineered fluorescent protein whose amino acid sequence is substantially identical to the amino acid sequence of *Aequorea* green fluorescent protein (SEQ ID NO:2) and which differs from SEQ ID NO:2 by at least an amino acid substitution at L42, V61, T62, V68, Q69, Q94, N121, Y145, H148, V150, F165, I167, Q183, N185, L220, E222 (E222G), or V224, said functional engineered fluorescent protein having a different fluorescent property than *Aequorea* green fluorescent protein.

59. The probe of claim 58 wherein the amino acid substitution is:
L42X, wherein X is selected from C, F, H, W and Y,
V61X, wherein X is selected from F, Y, H and C,
T62X, wherein X is selected from A, V, F, S, D, N, Q, Y, H and C,
V68X, wherein X is selected from F, Y and H,
Q69X, wherein X is selected from K, R, E and G,
Q94X, wherein X is selected from D, E, H, K and N,
N121X, wherein X is selected from F, H, W and Y,
Y145X, wherein X is selected from W, C, F, L, E, H, K and Q,
H148X, wherein X is selected from F, Y, N, K, Q and R,
V150X, wherein X is selected from F, Y and H,
F165X, wherein X is selected from H, Q, W and Y,
I167X, wherein X is selected from F, Y and H,
Q183X, wherein X is selected from H, Y, E and K,
N185X, wherein X is selected from D, E, H, K and Q,
L220X, wherein X is selected from H, N, Q and T,
E222X, wherein X is selected from N and Q or
V224X, wherein X is selected from H, N, Q, T, F, W and Y.
60. A method for determining whether a mixture contains a target comprising:
 contacting the mixture with a fluorescently labelled probe comprising a
 probe and a functional engineered fluorescent protein of claim 27 or claim 52; and
 determining whether the target has bound to the probe.
61. The method of any of claim 60 the target is bound to a solid matrix.

62. A method for engineering a functional engineered fluorescent protein having a fluorescent property different than *Aequorea* green fluorescent protein, comprising substituting an amino acid that is located no more than 0.5 nm from any atom in the chromophore of an *Aequorea*-related green fluorescent protein with another amino acid; whereby the substitution alters a fluorescent property of the protein.

63. The method of claim 62 wherein the amino acid substitution alters the electronic environment of the chromophore.

64. A method for engineering a functional engineered fluorescent protein having a different fluorescent property than *Aequorea* green fluorescent protein comprising substituting amino acids in a loop domain of an *Aequorea*-related green fluorescent protein with amino acids so as to create a consensus sequence for phosphorylation or for proteolysis.

65. A method for producing fluorescence resonance energy transfer comprising:
providing a donor molecule comprising a functional engineered fluorescent protein of claim 27 or claim 52;
providing an appropriate acceptor molecule for the fluorescent protein; and
bringing the donor molecule and the acceptor molecule into sufficiently close contact to allow fluorescence resonance energy transfer.

66. A method for producing fluorescence resonance energy transfer comprising:
providing an acceptor molecule comprising a functional engineered fluorescent protein of claim 27 or claim 52;
providing an appropriate donor molecule for the fluorescent protein; and
bringing the donor molecule and the acceptor molecule into sufficiently close contact to allow fluorescence resonance energy transfer.

67. The method of claim 66 wherein the donor molecule is a engineered fluorescent protein whose amino acid sequence comprises the substitution T203I and the acceptor molecule is a mutant fluorescent protein whose amino acid sequence comprises the substitution T203X, wherein X is an aromatic amino acid selected from H, Y, W or F, said functional engineered fluorescent protein having a different fluorescent property than *Aequorea* green fluorescent protein.

68. A nucleic acid molecule comprising a nucleotide sequence encoding a functional engineered fluorescent protein whose amino acid sequence is substantially identical to the amino acid sequence of *Aequorea* green fluorescent protein (SEQ ID NO:2) and which differs from SEQ ID NO:2 by at least an amino acid substitution located no more than about 0.5 nm from the chromophore of the engineered fluorescent protein, wherein the substitution alters the electronic environment of the chromophore, whereby the functional engineered fluorescent protein has a different fluorescent property than *Aequorea* green fluorescent protein.

69. An expression vector comprising expression control sequences operatively linked to a nucleotide sequence encoding a functional engineered fluorescent protein whose amino acid sequence is substantially identical to the amino acid sequence of *Aequorea* green fluorescent protein (SEQ ID NO:2) and which differs from SEQ ID NO:2 by at least an amino acid substitution located no more than about 0.5 nm from the chromophore of the engineered fluorescent protein, wherein the substitution alters the electronic environment of the chromophore, whereby the functional engineered fluorescent protein has a different fluorescent property than *Aequorea* green fluorescent protein.

70. A functional engineered fluorescent protein whose amino acid sequence is substantially identical to the amino acid sequence of *Aequorea* green fluorescent protein (SEQ ID NO:2) and which differs from SEQ ID NO:2 by at least an amino acid substitution located no more than about 0.5 nm from the chromophore of the engineered fluorescent protein, wherein the substitution alters the electronic environment of the chromophore, whereby the functional engineered fluorescent protein has a different fluorescent property than *Aequorea* green fluorescent protein.

71. A crystal of a protein comprising a fluorescent protein with an amino acid sequence substantially identical to SEQ ID NO: 2, wherein said crystal diffracts with at least a 2.0 to 3.0 angstrom resolution.

72. The crystal of claim 71, wherein the fluorescent protein has at least 200 amino acids, a completeness value of at least 80% and has a crystal stability within 0.5% of its unit cell dimensions.

73. The crystal of claim 71, wherein the amino acid sequence comprises a substitution at S65, wherein the substitution is selected from S65G, S65T, S65A, S65L, S65C, S65V and S65I.

74. The crystal of claim 71, wherein said crystal has the following unit cell dimensions in angstroms: $a = 51.8$, $b = 62.8$ and $c = 70.7$ with a space group of P2 2 2 and an α angle of 90.00° , a β angle of 90.00° and a γ angle of 90.00° and the crystal has a diffraction limit where 90% or greater of the potential reflections can be used to determine the coordinates of the atoms.

75. A computational method of designing a fluorescent protein comprising:

determining from a three dimensional model of a crystallized fluorescent protein comprising a fluorescent protein with a bound ligand, at least one interacting amino acid of the fluorescent protein that interacts with at least one first chemical moiety of the ligand, and

selecting at least one chemical modification of the first chemical moiety to produce a second chemical moiety with a structure to either decrease or increase an interaction between the interacting amino acid and the second chemical moiety compared to the interaction between the interacting amino acid and the first chemical moiety.

76. The computational method of claim 75, further comprising generating the three dimensional model of the crystallized protein comprising a fluorescent protein with an amino acid sequence substantially identical to SEQ ID NO:2.

77. The computational method of claim 75, wherein the selecting selects the first chemical moiety that interacts with at least one of the amino acids listed in Figs. 5-1 to 5-28.

78. The computational method of claim 75, wherein the chemical modification enhances hydrogen bonding interaction, charge interaction, hydrophobic interaction, Van Der Waals interaction or dipole interaction between the second chemical moiety and the interacting amino acid compared to the first chemical moiety and the interacting amino acid.

79. A computational method of modeling the three dimensional structure of a fluorescent protein comprising determining a three dimensional relationship between at least two atoms listed in the atomic coordinates of Figs. 5-1 to 5-28.

80. The computational method of claim 79, wherein the determining comprises determining the three dimensional structure of a fluorescent protein with an amino acid sequence at least 80% identical to SEQ ID NO:2.

81. The computational method of claim 79, wherein the determining comprises determining the three dimensional structure of a fluorescent protein with an amino acid sequence at least 95% identical to SEQ ID NO:2.

82. The computational method of claim 79, wherein the determining comprises determining the three dimensional relationship of at least 1500 atoms listed in Figs. 5-1 to 5-28.

83. A device comprising a storage device and, stored in the device, at least 10 atomic coordinates selected from the atomic coordinates listed in Figs. 5-1 to 5-28.

84. The device of claim 83, wherein the storage device is a computer readable device that stores code that receives as input the atomic coordinates.
85. The device of claim 84, wherein computer readable device is a floppy disk or a hard drive.
86. A nucleic acid molecule comprising a nucleotide sequence encoding a functional engineered fluorescent protein whose amino acid sequence is substantially identical to the amino acid sequence of *Aequorea* green fluorescent protein (SEQ ID NO:2) and which differs from SEQ ID NO:2 by at least a substitution at Q69, wherein said functional engineered fluorescent protein has a different fluorescent property than *Aequorea* green fluorescent protein.
87. The nucleic acid molecule of claim 86, wherein said substitution at Q69 is selected from the group of K, R, E and G.
88. The nucleic acid molecule of claim 86, wherein said amino acid sequence further comprises a function mutation at S65.
89. A nucleic acid molecule comprising a nucleotide sequence encoding a functional engineered fluorescent protein whose amino acid sequence is substantially identical to the amino acid sequence of *Aequorea* green fluorescent protein (SEQ ID NO:2) and which differs from SEQ ID NO:2 by at least a substitution at E222, but not including E222G, wherein said functional engineered fluorescent protein has a different fluorescent property than *Aequorea* green fluorescent protein.
90. The nucleic acid molecule of claim 89, wherein said substitution at E222 is selected from the group of N and Q.

91. The nucleic acid molecule of claim 89, wherein said amino acid sequence further comprises a function mutation at F64.
92. A nucleic acid molecule comprising a nucleotide sequence encoding a functional engineered fluorescent protein whose amino acid sequence is substantially identical to the amino acid sequence of *Aequorea* green fluorescent protein (SEQ ID NO:2) and which differs from SEQ ID NO:2 by at least a substitution at Y145, wherein said functional engineered fluorescent protein has a different fluorescent property than *Aequorea* green fluorescent protein.
93. The nucleic acid molecule of claim 92, wherein said substitution at Y145 is selected from the group of W, C, F, L, E, H, K and Q.
94. The nucleic acid molecule of claim 92, wherein said amino acid sequence further comprises a function mutation at Y66.
95. A method of identifying a test chemical, comprising:
contacting a test chemical a sample containing a biological entity labeled with a functional, engineered fluorescent protein or a polynucleotide encoding said functional, engineered fluorescent protein, and
detecting fluorescence of said functional engineered fluorescent protein.
96. The method of claim 95, wherein said fluorescence in the presence of a test chemical is greater than in the absence of said test chemical.
97. The method of claim 96, wherein said polynucleotide encoding said functional, engineered fluorescent protein is operatively linked to a genomic polynucleotide.
98. The method of claim 95, wherein said functional, engineered fluorescent protein is fused to second functional protein.

99. The method of claim 96, wherein said polynucleotide encoding said functional, engineered fluorescent protein is operatively linked to a response element.
100. The method of claim 96, wherein said polynucleotide encoding said functional, engineered fluorescent protein is operatively linked to a response element in a mammalian cell.
101. A nucleic acid molecule comprising a nucleotide sequence encoding a functional engineered fluorescent protein whose amino acid sequence is substantially identical to the amino acid sequence of *Aequorea* green fluorescent protein (SEQ ID NO:2) and which differs from SEQ ID NO:2 by
- i) at least one first substitution at position T203, wherein the substitution selected from the group consisting of H, Y, W or F, and
 - ii) at least one second substitution at position H148,
- wherein said functional engineered fluorescent protein has a different fluorescent property than *Aequorea* green fluorescent protein.
102. The nucleic acid molecule of claim 101, wherein said at least one second substitution at position H148 is selected from the group consisting of H148R, H148G, H148Q, H148A, H148N, and H148K.
103. The nucleic acid of claim 101, wherein said at least one second substitution at position H148 is H148Q.
104. The nucleic acid of claim 101, wherein said at least one second substitution at position H148 is H148G.
105. The nucleic acid of claim 101, wherein said at least one second substitution at position H148 is H148R.

106. The nucleic acid of claim 101, wherein said functional engineered fluorescent protein further comprises at least one third substitution at position V150.
107. The nucleic acid of claim 106, wherein said at least one third substitution at position V150 is selected from the group consisting of A, C, M, G, L, Q, S, T and N.
108. The nucleic acid of claim 101, wherein said functional engineered fluorescent protein further comprises at least one third substitution at position V163.
109. The nucleic acid of claim 108, wherein said at least one third substitution at position V163 is selected from the group consisting of A, C, M, G, L, Q, S, T and N.
110. The nucleic acid of claim 101, wherein said functional engineered fluorescent protein further comprises at least one third substitution at position Q69.
111. The nucleic acid of claim 110, wherein said at least one third substitution at position Q69 is selected from the group consisting of N, S, T and V.
112. The nucleic acid of claim 101, wherein said functional engineered fluorescent protein further comprises at least one third substitution at position V152.
113. The nucleic acid of claim 112, wherein said at least one third substitution at position V152 is selected from the group consisting of A, C, M, G, L, V, F, S, T, Q and N.
114. The nucleic acid of claim 101, wherein said functional engineered fluorescent protein further comprises at least one third substitution at position F165.
115. The nucleic acid of claim 114, wherein said at least one third substitution at position F165 is selected from the group consisting of Y, L and W.

116. The nucleic acid of claim 101, wherein said functional engineered fluorescent protein further comprises at least one third substitution at position H181.
117. The nucleic acid of claim 116, wherein said at least one third substitution at position H181 is selected from the group consisting of K, R, F, Y and W.
118. The nucleic acid of claim 101, wherein said functional engineered fluorescent protein further comprises at least one third substitution at position L201.
119. The nucleic acid of claim 118, wherein said at least one third substitution at position L201 is selected from the group consisting of A, C, M, G, S, T, Q, N, V and I.
120. A method of determining the presence of an anion of interest in a sample, comprising the steps of,
- 1) introducing an engineered green fluorescent protein into a sample, said engineered green fluorescent protein comprising an amino acid sequence substantially identical to the amino acid sequence of *Aequorea* green fluorescent protein (SEQ ID NO:2) and which differs from SEQ ID NO:2 by at least one first substitution at position T203, wherein the substitution selected from the group consisting H, Y, W or F,
 - 2) determining the fluorescence of said engineered green fluorescent protein in said sample.
121. The method of claim 120, further comprising the step of comparing the fluorescence of said engineered green fluorescent protein in said sample to the fluorescence of a control engineered green fluorescent protein introduced into a control sample comprising said anion of interest at a known concentration.
122. The method of claim 120, wherein said sample comprises at least one living cell.

123. The method of claim 120, wherein said anion of interest is a halide.
124. The method of claim 120, wherein said engineered fluorescent protein comprises at least one second substitution at position H148.
125. The method of claim 124, wherein, wherein said at least one second substitution at position H148 is selected from the group consisting of H148R, H148G, H148Q, H148A, H148N, and H148K.
126. The method of claim 125, wherein, wherein said at least one second substitution at position H148 is H148Q.
127. The method of claim 125, wherein, wherein said at least one second substitution at position H148 is H148G.
128. The method of claim 125, wherein, wherein said at least one second substitution at position H148 is H148R.
129. The method of claim 120, wherein, wherein said functional engineered fluorescent protein further comprises at least one third substitution at position V150.
130. The method of claim 129, wherein, wherein said at least one third substitution at position V150 is selected from the group consisting of A, C, M, G, L, Q, S, T and N.
131. The method of claim 120, wherein, wherein said functional engineered fluorescent protein further comprises at least one third substitution at position V163.
132. The method of claim 131, wherein, wherein said at least one third substitution at position V163 is selected from the group consisting of Q, S, T and N.

133. The method of claim 120, wherein, wherein said functional engineered fluorescent protein further comprises at least one third substitution at position Q69.
134. The method of claim 133, wherein, wherein said at least one third substitution at position Q69 is selected from the group consisting of N, S, T and V.
135. The method of claim 120, wherein said functional engineered fluorescent protein further comprises at least one third substitution at position V152.
136. The method of claim 135, wherein said at least one third substitution at position V152 is selected from the group consisting of A, C, M, G, L, V, F, S, T, Q and N.
137. The method of claim 120, wherein said functional engineered fluorescent protein further comprises at least one third substitution at position F165.
138. The method of claim 137, wherein said at least one third substitution at position F165 is selected from the group consisting of Y, L and W.
139. The method of claim 120, wherein said functional engineered fluorescent protein further comprises at least one third substitution at position H181.
140. The method of claim 139, wherein said at least one third substitution at position H181 is selected from the group consisting of K, R, F, Y and W.
141. The method of claim 120, wherein said functional engineered fluorescent protein further comprises at least one third substitution at position L201.
142. The method of claim 141, wherein said at least one third substitution at position L201 is selected from the group consisting of A, C, M, G, S, T, Q, N, V and I.

143. A functional engineered fluorescent protein whose amino acid sequence is substantially identical to the amino acid sequence of *Aequorea* green fluorescent protein (SEQ ID NO:2) and which differs from SEQ ID NO:2 by
- i) at least one first substitution at position T203, wherein the substitution selected from the group consisting H, Y, W or F, and
 - ii) at least one second substitution at position H148,
- wherein said functional engineered fluorescent protein has a different fluorescent property than *Aequorea* green fluorescent protein.
144. The functional engineered fluorescent protein of claim 143, wherein said at least one second substitution at position H148 is selected from the group consisting of H148R, H148G, H148Q, H148A, H148N, and H148K.
145. The functional engineered fluorescent protein of claim 144, wherein said at least one second substitution at position H148 is H148Q.
146. The functional engineered fluorescent protein of claim 144, wherein said at least one second substitution at position H148 is H148G.
147. The functional engineered fluorescent protein of claim 144, wherein said at least one second substitution at position H148 is H148R.
148. The functional engineered fluorescent protein of claim 143, wherein said functional engineered fluorescent protein further comprises at least one third substitution at position V150.
149. The functional engineered fluorescent protein of claim 148, wherein said at least one third substitution at position V150 is selected from the group consisting of Q, S, T and N.

150. The functional engineered fluorescent protein of claim 143, wherein said functional engineered fluorescent protein further comprises at least one third substitution at position V163.
151. The functional engineered fluorescent protein of claim 150, wherein said at least one third substitution at position V163 is selected from the group consisting of Q, S, T and N.
152. The functional engineered fluorescent protein of claim 143, wherein said functional engineered fluorescent protein further comprises at least one third substitution at position Q69.
153. The functional engineered fluorescent protein of claim 152, wherein said at least one third substitution at position Q69 is selected from the group consisting of N, S, T and V.
154. A host cell comprising a functional engineered fluorescent protein whose amino acid sequence is substantially identical to the amino acid sequence of *Aequorea* green fluorescent protein (SEQ ID NO:2) and which differs from SEQ ID NO:2 by
- i) at least one first substitution at position T203, wherein the substitution selected from the group consisting H, Y, W or F, and
 - ii) at least one second substitution at position H148,
- wherein said functional engineered fluorescent protein has a different fluorescent property than *Aequorea* green fluorescent protein.
155. The host cell of claim 154, wherein said at least one second substitution at position H148 is selected from the group consisting of H148R, H148G, H148Q, H148A, H148N, and H148K.
156. The host cell of claim 155, wherein said at least one second substitution at position H148 is H148Q.

157. The host cell of claim 155, wherein said at least one second substitution at position H148 is H148G.
158. The host cell of claim 155, wherein said at least one second substitution at position H148 is H148R.
159. The host cell of claim 143, wherein said functional engineered fluorescent protein further comprises at least one third substitution at position V150.
160. The host cell of claim 159, wherein said at least one third substitution at position V150 is selected from the group consisting of Q, S, T and N.
161. The host cell of claim 143, wherein said functional engineered fluorescent protein further comprises at least one third substitution at position V163.
162. The host cell of claim 161, wherein said at least one third substitution at position V163 is selected from the group consisting of Q, S, T and N.
163. The host cell of claim 143, wherein said functional engineered fluorescent protein further comprises at least one third substitution at position Q69.
164. The host cell of claim 163, wherein said at least one third substitution at position Q69 is selected from the group consisting of N, S, T and V.

165. A method for screening the effects of test compounds on ion channel activity comprising the steps of,

- i) providing a cell comprising
 - a) an engineered green fluorescent protein said engineered green fluorescent protein comprising an amino acid sequence substantially identical to the amino acid sequence of *Aequorea* green fluorescent protein (SEQ ID NO:2) and which differs from SEQ ID NO:2 by at least one first substitution at position T203, wherein the substitution selected from the group consisting H, Y, W or F,
 - b) an ion channel of interest,
- ii) contacting said cell with a test compound,
- iii) determining fluorescence from said engineered green fluorescent protein.

166. The method of claim 165 wherein said method further comprises the step of contacting said cell with a known activator of said ion channel of interest.

167. The method of claim 165, further comprising the step of comparing the fluorescence of said engineered green fluorescent protein in said cell to the fluorescence of a control engineered green fluorescent protein introduced into a control cell..

168. The method of claim 165, wherein said ion channel of interest transports halides.

169. The method of claim 165, wherein said engineered fluorescent protein comprises at least one second substitution at position H148.

170. The method of claim 169, wherein, wherein said at least one second substitution at position H148 is selected from the group consisting of H148R, H148G, H148Q, H148A, H148N, and H148K.
171. The method of claim 170, wherein, wherein said at least one second substitution at position H148 is H148Q.
172. The method of claim 170, wherein, wherein said at least one second substitution at position H148 is H148G.
173. The method of claim 170, wherein, wherein said at least one second substitution at position H148 is H148R.
174. The method of claim 165, wherein, wherein said functional engineered fluorescent protein further comprises at least one third substitution at position V150.
175. The method of claim 174, wherein, wherein said at least one third substitution at position V150 is selected from the group consisting of A, C, M, G, L, Q, S, T and N.
176. The method of claim 165, wherein, wherein said functional engineered fluorescent protein further comprises at least one third substitution at position V163.
177. The method of claim 176, wherein, wherein said at least one third substitution at position V163 is selected from the group consisting of Q, S, T and N.
178. The method of claim 165, wherein, wherein said functional engineered fluorescent protein further comprises at least one third substitution at position Q69.
179. The method of claim 178, wherein, wherein said at least one third substitution at position Q69 is selected from the group consisting of N, S, T and V.

180. The method of claim 165, wherein said functional engineered fluorescent protein further comprises at least one third substitution at position V152.
181. The method of claim 180, wherein said at least one third substitution at position V152 is selected from the group consisting of A, C, M, G, L, V, F, S, T, Q and N.
182. The method of claim 165, wherein said functional engineered fluorescent protein further comprises at least one third substitution at position F165.
183. The method of claim 182, wherein said at least one third substitution at position F165 is selected from the group consisting of Y, L and W.
184. The method of claim 165, wherein said functional engineered fluorescent protein further comprises at least one third substitution at position H181.
185. The method of claim 184, wherein said at least one third substitution at position H181 is selected from the group consisting of K, R, F, Y and W.
186. The method of claim 165, wherein said functional engineered fluorescent protein further comprises at least one third substitution at position L201.
187. The method of claim 186, wherein said at least one third substitution at position L201 is selected from the group consisting of A, C, M, G, S, T, Q, N, V and I.




FIG. 1A






FIG. 1B

**FIG. 2A****FIG. 2B**




FIG. 2C

(xi) SEQUENCE DESCRIPTION:

SEQ ID NO:1:

SEQ ID NO:2:

ATG	AGT	AAA	GGA	GAA	GAA	CTT	TTC	ACT	GGA	GTT	GTC	CCA	ATT	CTT	GTT	48
Met	Sen	Lys	Gly	Glu	Glu	Leu	Phe	Thr	Gly	Val	Val	Pro	Ile	Leu	Val	
1			5						10					15		
GAA	TTA	GAT	GGT	GAT	GTT	AAT	GGG	CAC	AAA	TTT	TCT	GTC	AGT	GGA	GAG	96
Glu	Leu	Asp	Gly	Asp	Val	Asn	Gly	His	Lys	Phe	Ser	Val	Ser	Gly	Glu	
		20					25						30			
GGT	GAA	GGT	GAT	GCA	ACA	TAC	GGA	AAA	CTT	ACC	CTT	AAA	TTT	ATT	TGC	144
Gly	Glu	Gly	Asp	Ala	Thr	Tyr	Gly	Lys	Leu	Thr	Leu	Lys	Phe	Ile	Cys	
		35					40					45				
ACT	ACT	GGA	AAA	CTA	CCT	GTT	CCA	TGG	CCA	ACA	CTT	GTC	ACT	ACT	TTC	192
Thr	Thr	Gly	Lys	Leu	Pro	Val	Pro	Trp	Pro	Thr	Leu	Val	Thr	Thr	Phe	
	50					55						60				
TCT	TAT	GGT	GTT	CAA	TGC	TTT	TCA	AGA	TAC	CCA	GAT	CAT	ATG	AAA	CGG	240
Sen	Tyr	Gly	Val	Gln	Cys	Phe	Ser	Arg	Tyr	Pro	Asp	His	Met	Lys	Arg	
65				70						75				80		
CAT	GAC	TTT	TTC	AAG	AGT	GCC	ATG	CCC	GAA	GGT	TAT	GTA	CAG	GAA	AGA	288
His	Asp	Phe	Phe	Lys	Sen	Ala	Met	Pro	Glu	Gly	Tyr	Val	Gln	Glu	Arg	
			85					90					95			
ACT	ATA	TTT	TTC	AAA	GAT	GAC	GGG	AAC	TAC	AAG	ACA	CGT	GCT	GAA	GTC	336
Thr	Ile	Phe	Phe	Lys	Asp	Asp	Gly	Asn	Tyr	Lys	Thr	Arg	Ala	Glu	Val	
		100					105					110				
AAG	TTT	GAA	GGT	GAT	ACC	CTT	GTT	AAT	AGA	ATC	GAG	TTA	AAA	GGT	ATT	384
Lys	Phe	Glu	Gly	Asp	Thr	Leu	Val	Asn	Arg	Ile	Glu	Leu	Lys	Gly	Ile	
	115					120						125				
GAT	TTT	AAA	GAA	GAT	GGA	AAC	ATT	CTT	GGA	CAC	AAA	TTG	GAA	TAC	AAC	432
Asp	Phe	Lys	Glu	Asp	Gly	Asn	Ile	Leu	Gly	His	Lys	Leu	Glu	Tyr	Asn	
	130				135						140					
TAT	AAC	TCA	CAC	AAT	GTA	TAC	ATC	ATG	GCA	GAC	AAA	CAA	AAG	AAT	GGA	480
Tyr	Asn	Ser	His	Asn	Val	Tyr	Ile	Met	Ala	Asp	Lys	Gln	Lys	Asn	Gly	
145				150						155					160	
ATC	AAA	GTT	AAC	TTC	AAA	ATT	AGA	CAC	AAC	ATT	GAA	GAT	GGA	AGC	GTT	528
Ile	Lys	Val	Asn	Phe	Lys	Ile	Arg	His	Asn	Ile	Glu	Asp	Gly	Ser	Val	
			165						170				175			
CAA	CTA	GCA	GAC	CAT	TAT	CAA	CAA	AAT	ACT	COA	ATT	GGC	GAT	GGC	CCT	576
Gln	Leu	Ala	Asp	His	Tyr	Gln	Gln	Asn	Thr	Pro	Ile	Gly	Asp	Gly	Pro	
		180						185				190				
GTC	CTT	TTA	CCA	GAC	AAC	CAT	TAC	CTG	TCC	ACA	CAA	TCT	GCC	CTT	TCG	624
Val	Leu	Leu	Pro	Asp	Asn	His	Tyr	Leu	Sen	Thr	Gln	Ser	Ala	Leu	Ser	
	195					200						205				
AAA	GAT	CCC	AAC	GAA	AAG	AGA	GAC	CAC	ATG	GTC	CTT	CTT	GAG	TTT	GTA	672
Lys	Asp	Pro	Asn	Glu	Lys	Arg	Asp	His	Met	Val	Leu	Leu	Glu	Phe	Val	
	210					215					220					
ACA	GCT	GCT	GGG	ATT	ACA	CAT	GGC	ATG	GAT	GAA	CTA	TAC	AAA	TA		717
Thr	Ala	Ala	Gly	Ile	Thr	His	Gly	Met	Asp	Glu	Leu	Tyr	Lys			
225					230					235						

FIG. 3

T203Y, S65G, S72A humanized codon usage, with an additional amino acid
ater the start met to provide optimal kozak sequence

	9		18		27		36		45		54						
ATG	GTG	AGC	AAG	GCC	GAG	GAG	CTG	TTC	ACC	GGG	GTG	GTG	CCC	ATC	CTG	GTC	GAG
Met	Val	Ser	Lys	Gly	Glu	Glu	Leu	Phe	Thr	Gly	Val	Val	Pro	Ile	Leu	Val	Glu
	63		72		81		90		99		108						
CTG	GAG	GGC	GAC	GTA	AAC	GGC	CAC	AAG	TTC	AGC	GTG	TCC	GGC	GAG	GGC	GAG	GGC
Leu	Asp	Gly	Asp	Val	Asn	Gly	His	Lys	Phe	Ser	Val	Ser	Gly	Glu	Gly	Glu	Gly
	117		126		135		144		153		162						
GAT	GCC	ACC	TAC	GGC	AAG	CTG	ACC	CTG	AAG	TTC	ATC	TGC	ACC	ACC	GGC	AAG	CTG
Asp	Ala	Thr	Tyr	Gly	Lys	Leu	Thr	Leu	Lys	Phe	Ile	Cys	Thr	Thr	Gly	Lys	Leu
	171		180		189		198		207		216						
CCC	GTG	CCC	TGG	CCC	ACC	CTC	GTG	ACC	ACC	TTC	GGC	TAC	GGC	GTG	CAG	TGC	TTC
Pro	Val	Pro	Trp	Pro	Thr	Leu	Val	Thr	Thr	Phe	Gly	Tyr	Gly	Val	Gln	Cys	Phe
	225		234		243		252		261		270						
GCC	CGC	TAC	CCC	GAC	CAC	ATG	AAG	OAG	CAC	GAC	TTC	TTC	AAG	TCC	GCC	ATG	CCC
Ala	Arg	Tyr	Pro	Asp	His	Met	Lys	Gln	His	Asp	Phe	Phe	Lys	Ser	Ala	Met	Pro
	279		288		297		306		315		324						
GAA	GGC	TAC	GTC	CAG	GAG	CGG	ACC	ATC	TTC	TTC	AAG	GAC	GAC	GGC	AAC	TAC	AAG
Glu	Gly	Tyr	Val	Gln	Glu	Arg	Thr	Ile	Phe	Phe	Lys	Asp	Asp	Gly	Asn	Tyr	Lys
	333		342		351		360		369		378						
ACC	CGC	GCC	GAG	GTG	AAG	TTC	GAG	GGC	GAC	ACC	CTG	GTG	AAC	CGC	ATC	GAG	CTG
Thr	Arg	Ala	Glu	Val	Lys	Phe	Glu	Gly	Asp	Thr	Leu	Val	Asn	Arg	Ile	Glu	Leu
	387		396		405		414		423		432						
AAG	GGC	ATC	GAC	TTC	AAG	GAG	GAC	GGC	AAC	ATC	CTG	GGG	CAC	AAG	CTG	GAG	TAC
Lys	Gly	Ile	Asp	Phe	Lys	Glu	Asp	Gly	Asn	Ile	Leu	Gly	His	Lys	Leu	Glu	Tyr
	441		450		459		468		477		486						
AAC	TAC	AAC	AGC	CAC	AAC	GTC	TAT	ATC	ATG	GCC	GAC	AAG	CAG	AAG	AAC	GGC	ATC
Asn	Tyr	Asn	Ser	His	Asn	Val	Tyr	Ile	Met	Ala	Asp	Lys	Gln	Lys	Asn	Gly	Ile

FIG. 4A

		495			504			513		522			531		540		
AAG	GTG	AAC	TTC	AAG	ATC	CGC	CAC	AAC	ATC	GAG	GAC	GGC	AGC	GTG	CAG	CTC	GCC
---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
Lys	Val	Asn	Phe	Lys	Ile	Arg	His	Asn	Ile	Glu	Asp	Gly	Ser	Val	Gln	Leu	Ala
		549			558			567		576			585		594		
GAC	CAC	TAC	CAG	CAG	AAC	ACC	CCC	ATC	GGC	GAC	GGC	CCC	GTG	CTG	CTG	CCC	GAC
---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
Asp	His	Tyr	Gln	Gln	Asn	Thr	Pro	Ile	Gly	Asp	Gly	Pro	Val	Leu	Leu	Pro	Asp
		603			612			621		630			639		648		
AAC	CAC	TAC	CTG	AGC	TAC	CAG	TCC	GCC	CTG	AGC	AAA	GAC	CCC	AAC	GAG	AAG	CGC
---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
Asn	His	Tyr	Leu	Ser	Tyr	Gln	Ser	Ala	Leu	Ser	Lys	Asp	Pro	Asn	Glu	Lys	Arg
		657			666			675		684			693		702		
GAT	CAC	ATG	GTC	CTG	CTG	GAG	TTC	GTG	ACC	GCC	GCC	GGG	ATC	ACT	CAC	GGC	ATG
---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---	---
Asp	His	Met	Val	Leu	Leu	Glu	Phe	Val	Thr	Ala	Ala	Gly	Ile	Thr	His	Gly	Met
					711												
GAC	GAG	CTG	TAC	AAG	TAA	3'											
---	---	---	---	---	---	---											
Asp	Glu	Leu	Tyr	Lys	***												

FIG. 4B

CRYST1	51.767	62.845	70.666	90.00	90.00	90.00		
ORIGX1	1.000000	0.000000	0.000000			0.000000		
ORIGX2	0.000000	1.000000	0.000000			0.000000		
ORIGX3	0.000000	0.000000	1.000000			0.000000		
SCALE1	0.019317	0.000000	0.000000			0.000000		
SCALE2	0.000000	0.015912	0.000000			0.000000		
SCALE3	0.000000	0.000000	0.014151			0.000000		
ATOM	1	N	SER	2	28.888	9.409	52.301	1.00 85.05
ATOM	2	CA	SER	2	27.638	10.125	52.516	1.00 80.05
ATOM	3	C	SER	2	26.499	9.639	51.644	1.00 85.36
ATOM	4	O	SER	2	26.606	8.656	50.915	1.00 84.56
ATOM	5	CB	SER	2	27.783	11.635	52.378	1.00 70.97
ATOM	6	OG	SER	2	27.690	12.033	51.012	1.00 44.08
ATOM	7	N	LYS	3	25.418	10.403	51.731	1.00 87.71
ATOM	8	CA	LYS	3	24.141	10.191	51.036	1.00 87.15
ATOM	9	C	LYS	3	24.214	10.266	49.497	1.00 76.86
ATOM	10	O	LYS	3	24.107	9.258	48.774	1.00 78.27
ATOM	11	CB	LYS	3	23.127	11.240	51.521	1.00 89.44
ATOM	12	CG	LYS	3	21.768	10.697	51.949	1.00 75.06
ATOM	13	CD	LYS	3	20.681	11.781	51.987	1.00 76.58
ATOM	14	CE	LYS	3	20.711	12.655	53.243	1.00 68.55
ATOM	15	NZ	LYS	3	20.816	14.103	52.953	1.00 46.24
ATOM	16	N	GLY	4	24.318	11.495	49.015	1.00 53.62
ATOM	17	CA	GLY	4	24.297	11.798	47.605	1.00 45.97
ATOM	18	C	GLY	4	25.425	11.206	46.796	1.00 31.90
ATOM	19	O	GLY	4	25.234	10.923	45.619	1.00 33.63
ATOM	20	N	GLU	5	26.606	11.082	47.420	1.00 32.54
ATOM	21	CA	GLU	5	27.821	10.598	46.726	1.00 32.57
ATOM	22	C	GLU	5	27.523	9.590	45.616	1.00 28.40
ATOM	23	O	GLU	5	27.850	9.803	44.444	1.00 26.12
ATOM	24	CB	GLU	5	28.873	10.053	47.718	1.00 38.53
ATOM	25	CG	GLU	5	30.337	10.461	47.425	1.00 41.36
ATOM	26	CD	GLU	5	31.311	9.584	48.170	1.00 90.82
ATOM	27	OE1	GLU	5	31.508	9.677	49.381	1.00 74.80
ATOM	28	OE2	GLU	5	31.839	8.653	47.403	1.00 100.00
ATOM	29	N	GLU	6	26.883	8.499	46.017	1.00 28.57
ATOM	30	CA	GLU	6	26.479	7.410	45.150	1.00 31.50
ATOM	31	C	GLU	6	25.561	7.837	43.979	1.00 31.10
ATOM	32	O	GLU	6	25.479	7.142	42.955	1.00 30.96
ATOM	33	CB	GLU	6	25.780	6.330	45.992	1.00 35.64
ATOM	34	CG	GLU	6	25.260	6.893	47.338	1.00 55.53
ATOM	35	N	LEU	7	24.864	8.966	44.138	1.00 22.26

FIG. 5A

ATOM	36	CA	LEU	7	23.954	9.456	43.089	1.00	21.61
ATOM	37	C	LEU	7	24.693	10.061	41.917	1.00	16.90
ATOM	38	O	LEU	7	24.152	10.250	40.836	1.00	18.38
ATOM	39	CB	LEU	7	23.050	10.548	43.665	1.00	22.41
ATOM	40	CG	LEU	7	21.672	10.058	44.098	1.00	32.84
ATOM	41	CD1	LEU	7	21.597	8.536	44.074	1.00	31.64
ATOM	42	CD2	LEU	7	21.332	10.591	45.485	1.00	33.14
ATOM	43	N	PHE	8	25.944	10.407	42.157	1.00	20.75
ATOM	44	CA	PHE	8	26.740	11.132	41.159	1.00	21.64
ATOM	45	C	PHE	8	27.818	10.333	40.427	1.00	30.59
ATOM	46	O	PHE	8	28.590	10.856	39.600	1.00	30.05
ATOM	47	CB	PHE	8	27.309	12.376	41.820	1.00	16.95
ATOM	48	CG	PHE	8	26.222	13.355	42.163	1.00	13.29
ATOM	49	CD1	PHE	8	25.672	13.378	43.447	1.00	17.27
ATOM	50	CD2	PHE	8	25.726	14.227	41.189	1.00	13.12
ATOM	51	CE1	PHE	8	24.661	14.290	43.772	1.00	15.14
ATOM	52	CE2	PHE	8	24.712	15.137	41.499	1.00	13.19
ATOM	53	CZ	PHE	8	24.192	15.170	42.794	1.00	5.69
ATOM	54	N	THR	9	27.798	9.074	40.699	1.00	27.35
ATOM	55	CA	THR	9	28.704	8.122	40.175	1.00	34.93
ATOM	56	C	THR	9	28.709	7.998	38.636	1.00	45.22
ATOM	57	O	THR	9	29.642	7.452	38.062	1.00	50.55
ATOM	58	CB	THR	9	28.447	6.795	40.892	1.00	44.60
ATOM	59	OG1	THR	9	29.629	6.330	41.527	1.00	40.40
ATOM	60	CG2	THR	9	27.801	5.779	39.959	1.00	29.76
ATOM	61	N	GLY	10	27.690	8.510	37.956	1.00	30.53
ATOM	62	CA	GLY	10	27.689	8.458	36.507	1.00	23.21
ATOM	63	C	GLY	10	27.144	9.746	35.914	1.00	16.55
ATOM	64	O	GLY	10	27.011	10.729	36.617	1.00	25.70
ATOM	65	N	VAL	11	26.835	9.719	34.629	1.00	16.39
ATOM	66	CA	VAL	11	26.209	10.863	33.971	1.00	22.28
ATOM	67	C	VAL	11	24.758	11.020	34.479	1.00	29.60
ATOM	68	O	VAL	11	23.972	10.062	34.456	1.00	20.43
ATOM	69	CB	VAL	11	26.173	10.664	32.467	1.00	30.87
ATOM	70	CG1	VAL	11	25.912	11.980	31.734	1.00	31.75
ATOM	71	CG2	VAL	11	27.480	10.048	32.015	1.00	33.85
ATOM	72	N	VAL	12	24.417	12.227	34.931	1.00	20.12
ATOM	73	CA	VAL	12	23.080	12.561	35.433	1.00	12.88
ATOM	74	C	VAL	12	22.407	13.624	34.516	1.00	14.37
ATOM	75	O	VAL	12	23.007	14.639	34.179	1.00	13.42
ATOM	76	CB	VAL	12	23.270	13.077	36.839	1.00	15.01

FIG. 5B

ATOM	77	CG1VAL	12	22.000	13.662	37.422	1.00	17.57
ATOM	78	CG2VAL	12	23.781	11.936	37.728	1.00	16.55
ATOM	79	N PRO	13	21.180	13.382	34.066	1.00	14.72
ATOM	80	CA PRO	13	20.493	14.382	33.265	1.00	10.76
ATOM	81	C PRO	13	20.116	15.589	34.141	1.00	7.65
ATOM	82	O PRO	13	19.797	15.468	35.337	1.00	15.14
ATOM	83	CB PRO	13	19.225	13.707	32.745	1.00	17.36
ATOM	84	CG PRO	13	19.043	12.422	33.550	1.00	19.69
ATOM	85	CD PRO	13	20.315	12.195	34.340	1.00	15.41
ATOM	86	N ILE	14	20.196	16.766	33.557	1.00	14.91
ATOM	87	CA ILE	14	19.893	17.991	34.266	1.00	12.93
ATOM	88	C ILE	14	18.768	18.760	33.596	1.00	12.08
ATOM	89	O ILE	14	18.724	18.878	32.399	1.00	11.04
ATOM	90	CB ILE	14	21.109	18.905	34.325	1.00	16.54
ATOM	91	CG1ILE	14	22.271	18.169	35.015	1.00	18.08
ATOM	92	CG2ILE	14	20.783	20.207	35.084	1.00	11.56
ATOM	93	CD1ILE	14	23.642	18.836	34.738	1.00	16.15
ATOM	94	N LEU	15	17.899	19.307	34.421	1.00	13.85
ATOM	95	CA LEU	15	16.811	20.136	33.955	1.00	14.82
ATOM	96	C LEU	15	16.915	21.474	34.685	1.00	3.62
ATOM	97	O LEU	15	17.080	21.509	35.901	1.00	10.00
ATOM	98	CB LEU	15	15.462	19.450	34.285	1.00	21.25
ATOM	99	CG LEU	15	14.412	19.541	33.199	1.00	40.50
ATOM	100	CD1LEU	15	13.279	20.440	33.679	1.00	46.97
ATOM	101	CD2LEU	15	15.008	20.098	31.913	1.00	49.22
ATOM	102	N VAL	16	16.885	22.556	33.919	1.00	10.56
ATOM	103	CA VAL	16	16.964	23.905	34.479	1.00	10.23
ATOM	104	C VAL	16	15.716	24.727	34.063	1.00	9.47
ATOM	105	O VAL	16	15.347	24.748	32.904	1.00	16.72
ATOM	106	CB VAL	16	18.273	24.668	34.098	1.00	12.85
ATOM	107	CG1VAL	16	18.226	26.075	34.691	1.00	12.58
ATOM	108	CG2VAL	16	19.520	23.945	34.628	1.00	14.24
ATOM	109	N GLU	17	15.059	25.317	35.060	1.00	14.43
ATOM	110	CA GLU	17	13.904	26.144	34.870	1.00	13.61
ATOM	111	C GLU	17	14.086	27.474	35.571	1.00	9.38
ATOM	112	O GLU	17	14.331	27.524	36.765	1.00	15.74
ATOM	113	CB GLU	17	12.650	25.402	35.344	1.00	14.15
ATOM	114	CG GLU	17	12.436	24.178	34.447	1.00	15.37
ATOM	115	CD GLU	17	11.865	24.573	33.105	1.00	49.50
ATOM	116	OE1 GLU	17	11.160	25.557	32.950	1.00	83.46
ATOM	117	OE2 GLU	17	12.220	23.766	32.127	1.00	38.75

FIG. 5C

ATOM	118	N	LEU	18	13.990	28.571	34.805	1.00	17.82
ATOM	119	CA	LEU	18	14.116	29.914	35.401	1.00	16.61
ATOM	120	C	LEU	18	12.962	30.855	35.057	1.00	14.91
ATOM	121	O	LEU	18	12.585	30.978	33.917	1.00	14.31
ATOM	122	CB	LEU	18	15.426	30.630	35.005	1.00	13.56
ATOM	123	CG	LEU	18	15.533	32.049	35.579	1.00	19.27
ATOM	124	CD1	LEU	18	16.740	32.182	36.489	1.00	21.40
ATOM	125	CD2	LEU	18	15.682	33.033	34.438	1.00	18.38
ATOM	126	N	ASP	19	12.480	31.551	36.082	1.00	17.88
ATOM	127	CA	ASP	19	11.476	32.577	35.940	1.00	19.57
ATOM	128	C	ASP	19	12.098	33.896	36.360	1.00	11.65
ATOM	129	O	ASP	19	12.486	34.044	37.493	1.00	16.82
ATOM	130	CB	ASP	19	10.234	32.305	36.847	1.00	24.92
ATOM	131	CG	ASP	19	9.305	31.262	36.282	1.00	38.46
ATOM	132	OD1	ASP	19	8.572	30.587	36.989	1.00	61.49
ATOM	133	OD2	ASP	19	9.337	31.189	34.949	1.00	22.44
ATOM	134	N	GLY	20	12.178	34.863	35.471	1.00	16.82
ATOM	135	CA	GLY	20	12.784	36.101	35.908	1.00	19.52
ATOM	136	C	GLY	20	12.048	37.385	35.538	1.00	19.35
ATOM	137	O	GLY	20	11.240	37.443	34.628	1.00	18.22
ATOM	138	N	ASP	21	12.401	38.407	36.286	1.00	13.19
ATOM	139	CA	ASP	21	11.908	39.737	36.112	1.00	16.36
ATOM	140	C	ASP	21	13.039	40.683	36.424	1.00	12.77
ATOM	141	O	ASP	21	13.517	40.742	37.569	1.00	15.18
ATOM	142	CB	ASP	21	10.701	40.036	37.040	1.00	22.26
ATOM	143	CG	ASP	21	10.230	41.491	37.022	1.00	30.80
ATOM	144	OD1	ASP	21	10.878	42.407	36.557	1.00	27.40
ATOM	145	OD2	ASP	21	9.062	41.658	37.604	1.00	45.92
ATOM	146	N	VAL	22	13.464	41.393	35.397	1.00	19.66
ATOM	147	CA	VAL	22	14.524	42.388	35.542	1.00	25.10
ATOM	148	C	VAL	22	14.010	43.780	35.154	1.00	18.26
ATOM	149	O	VAL	22	13.769	44.062	33.955	1.00	15.10
ATOM	150	CB	VAL	22	15.803	42.012	34.750	1.00	26.57
ATOM	151	CG1	VAL	22	16.861	43.127	34.896	1.00	24.27
ATOM	152	CG2	VAL	22	16.365	40.710	35.297	1.00	22.98
ATOM	153	N	ASN	23	13.823	44.641	36.166	1.00	25.32
ATOM	154	CA	ASN	23	13.319	45.993	35.908	1.00	32.81
ATOM	155	C	ASN	23	11.987	45.958	35.142	1.00	32.77
ATOM	156	O	ASN	23	11.774	46.730	34.187	1.00	30.47
ATOM	157	CB	ASN	23	14.344	46.831	35.096	1.00	31.26
ATOM	158	CG	ASN	23	15.374	47.607	35.938	1.00	24.72

FIG. 5D

ATOM	159	OD1ASN	23	15.795	47.183	37.024	1.00	27.22
ATOM	160	ND2ASN	23	15.829	48.723	35.389	1.00	41.15
ATOM	161	N GLY	24	11.118	45.024	35.519	1.00	24.95
ATOM	162	CA GLY	24	9.831	44.919	34.848	1.00	23.22
ATOM	163	C GLY	24	9.832	44.111	33.573	1.00	23.31
ATOM	164	O GLY	24	8.780	43.868	33.024	1.00	28.37
ATOM	165	N HIS	25	11.000	43.691	33.071	1.00	20.89
ATOM	166	CA HIS	25	11.042	42.840	31.877	1.00	19.30
ATOM	167	C HIS	25	10.981	41.373	32.316	1.00	27.26
ATOM	168	O HIS	25	11.898	40.850	32.951	1.00	26.47
ATOM	169	CB HIS	25	12.268	43.060	30.958	1.00	24.20
ATOM	170	CG HIS	25	12.313	44.382	30.218	1.00	33.04
ATOM	171	ND1HIS	25	12.917	45.514	30.758	1.00	37.58
ATOM	172	CD2HIS	25	11.876	44.716	28.971	1.00	42.76
ATOM	173	CE1HIS	25	12.801	46.497	29.867	1.00	39.14
ATOM	174	NE2HIS	25	12.185	46.050	28.778	1.00	42.80
ATOM	175	N LYS	26	9.872	40.728	32.028	1.00	25.90
ATOM	176	CA LYS	26	9.675	39.355	32.446	1.00	26.27
ATOM	177	C LYS	26	10.154	38.361	31.429	1.00	27.09
ATOM	178	O LYS	26	10.027	38.576	30.232	1.00	25.75
ATOM	179	CB LYS	26	8.230	39.069	32.863	1.00	27.58
ATOM	180	CG LYS	26	7.873	39.770	34.166	1.00	44.94
ATOM	181	CD LYS	26	6.369	39.914	34.400	1.00	71.44
ATOM	182	CE LYS	26	6.008	41.000	35.421	1.00	45.29
ATOM	183	N PHE	27	10.703	37.250	31.910	1.00	22.04
ATOM	184	CA PHE	27	11.164	36.236	30.978	1.00	18.78
ATOM	185	C PHE	27	11.273	34.863	31.619	1.00	14.75
ATOM	186	O PHE	27	11.293	34.722	32.842	1.00	15.94
ATOM	187	CB PHE	27	12.495	36.638	30.287	1.00	21.58
ATOM	188	CG PHE	27	13.599	36.826	31.311	1.00	22.06
ATOM	189	CD1PHE	27	14.490	35.791	31.612	1.00	23.61
ATOM	190	CD2PHE	27	13.722	38.029	32.005	1.00	17.55
ATOM	191	CE1PHE	27	15.487	35.963	32.579	1.00	16.61
ATOM	192	CE2PHE	27	14.747	38.234	32.931	1.00	19.75
ATOM	193	CZ PHE	27	15.621	37.187	33.234	1.00	13.83
ATOM	194	N SER	28	11.370	33.857	30.752	1.00	12.40
ATOM	195	CA SER	28	11.492	32.479	31.186	1.00	15.59
ATOM	196	C SER	28	12.579	31.749	30.379	1.00	15.96
ATOM	197	O SER	28	12.699	31.933	29.167	1.00	18.99
ATOM	198	CB SER	28	10.143	31.702	31.086	1.00	14.48
ATOM	199	OG SER	28	9.510	31.678	32.353	1.00	31.95

FIG. 5E

ATOM	200	N	VAL	29	13.335	30.902	31.073	1.00	16.73
ATOM	201	CA	VAL	29	14.361	30.093	30.435	1.00	14.06
ATOM	202	C	VAL	29	14.258	28.614	30.187	1.00	6.80
ATOM	203	O	VAL	29	14.058	28.266	31.987	1.00	10.85
ATOM	204	CB	VAL	29	15.768	30.570	30.839	1.00	17.96
ATOM	205	CG1	VAL	29	16.826	29.599	30.234	1.00	15.30
ATOM	206	CG2	VAL	29	15.989	32.001	30.357	1.00	16.37
ATOM	207	N	SER	30	14.462	27.781	29.824	1.00	11.31
ATOM	208	CA	SER	30	14.535	26.351	30.011	1.00	17.96
ATOM	209	C	SER	30	15.917	25.818	29.571	1.00	11.26
ATOM	210	O	SER	30	16.398	26.157	28.513	1.00	13.17
ATOM	211	CB	SER	30	13.471	25.603	29.202	1.00	19.91
ATOM	212	OG	SER	30	12.249	25.667	29.882	1.00	48.74
ATOM	213	N	GLY	31	16.480	24.926	30.364	1.00	9.88
ATOM	214	CA	GLY	31	17.718	24.321	29.977	1.00	12.44
ATOM	215	C	GLY	31	17.737	22.816	30.249	1.00	13.16
ATOM	216	O	GLY	31	17.149	22.324	31.176	1.00	12.41
ATOM	217	N	GLU	32	18.459	22.112	29.433	1.00	13.44
ATOM	218	CA	GLU	32	18.622	20.670	29.570	1.00	13.73
ATOM	219	C	GLU	32	20.079	20.297	29.262	1.00	17.33
ATOM	220	O	GLU	32	20.734	20.946	28.456	1.00	15.56
ATOM	221	CB	GLU	32	17.761	19.893	28.543	1.00	12.67
ATOM	222	CG	GLU	32	16.264	20.187	28.618	1.00	26.43
ATOM	223	CD	GLU	32	15.501	19.547	27.468	1.00	21.13
ATOM	224	OE1	GLU	32	15.996	18.767	26.698	1.00	23.45
ATOM	225	OE2	GLU	32	14.292	20.022	27.337	1.00	30.63
ATOM	226	N	GLY	33	20.534	19.207	29.822	1.00	15.36
ATOM	227	CA	GLY	33	21.860	18.687	29.518	1.00	12.84
ATOM	228	C	GLY	33	22.236	17.602	30.467	1.00	14.69
ATOM	229	O	GLY	33	21.390	16.919	31.011	1.00	13.56
ATOM	230	N	GLU	34	23.525	17.453	30.702	1.00	15.15
ATOM	231	CA	GLU	34	23.971	16.450	31.621	1.00	18.14
ATOM	232	C	GLU	34	25.220	16.874	32.367	1.00	16.26
ATOM	233	O	GLU	34	25.926	17.760	31.944	1.00	18.67
ATOM	234	CB	GLU	34	24.180	15.114	30.927	1.00	22.53
ATOM	235	CG	GLU	34	24.948	15.261	29.624	1.00	33.78
ATOM	236	CD	GLU	34	24.879	14.020	28.796	1.00	55.15
ATOM	237	OE1	GLU	34	25.861	13.352	28.534	1.00	45.39
ATOM	238	OE2	GLU	34	23.653	13.719	28.430	1.00	56.26
ATOM	239	N	GLY	35	25.461	16.222	33.485	1.00	11.20
ATOM	240	CA	GLY	35	26.611	16.502	34.315	1.00	10.62

FIG. 5F

ATOM	241	C	GLY	35	27.293	15.192	34.662	1.00	19.92
ATOM	242	O	GLY	35	26.650	14.161	34.750	1.00	16.69
ATOM	243	N	ASP	36	28.594	15.238	34.860	1.00	16.92
ATOM	244	CA	ASP	36	29.367	14.061	35.221	1.00	16.19
ATOM	245	C	ASP	36	30.396	14.505	36.233	1.00	13.94
ATOM	246	O	ASP	36	31.469	15.004	35.879	1.00	15.77
ATOM	247	CB	ASP	36	30.032	13.457	33.948	1.00	19.98
ATOM	248	CG	ASP	36	30.681	12.066	34.075	1.00	31.92
ATOM	249	OD1	ASP	36	31.236	11.519	33.141	1.00	30.97
ATOM	250	OD2	ASP	36	30.587	11.515	35.248	1.00	25.32
ATOM	251	N	ALA	37	30.015	14.402	37.490	1.00	13.40
ATOM	252	CA	ALA	37	30.818	14.846	38.582	1.00	12.98
ATOM	253	C	ALA	37	32.181	14.145	38.637	1.00	21.94
ATOM	254	O	ALA	37	33.084	14.604	39.331	1.00	13.61
ATOM	255	CB	ALA	37	30.070	14.741	39.916	1.00	11.49
ATOM	256	N	THR	38	32.307	13.016	37.945	1.00	15.63
ATOM	257	CA	THR	38	33.581	12.280	37.943	1.00	19.94
ATOM	258	C	THR	38	34.705	13.114	37.335	1.00	25.61
ATOM	259	O	THR	38	35.850	13.069	37.775	1.00	17.89
ATOM	260	CB	THR	38	33.462	10.898	37.299	1.00	22.57
ATOM	261	OG1	THR	38	32.543	10.146	38.067	1.00	29.86
ATOM	262	CG2	THR	38	34.821	10.213	37.355	1.00	22.90
ATOM	263	N	TYR	39	34.323	13.920	36.347	1.00	18.45
ATOM	264	CA	TYR	39	35.210	14.837	35.675	1.00	9.39
ATOM	265	C	TYR	39	34.874	16.291	35.991	1.00	14.41
ATOM	266	O	TYR	39	35.454	17.177	35.410	1.00	16.24
ATOM	267	CB	TYR	39	35.156	14.582	34.180	1.00	11.82
ATOM	268	CG	TYR	39	35.426	13.137	33.929	1.00	28.73
ATOM	269	CD1	TYR	39	36.715	12.633	34.065	1.00	33.75
ATOM	270	CD2	TYR	39	34.392	12.249	33.642	1.00	39.19
ATOM	271	CE1	TYR	39	36.982	11.276	33.828	1.00	29.75
ATOM	272	CE2	TYR	39	34.635	10.885	33.435	1.00	45.41
ATOM	273	CZ	TYR	39	35.943	10.410	33.570	1.00	57.62
ATOM	274	OH	TYR	39	36.199	9.070	33.364	1.00	70.77
ATOM	275	N	GLY	40	33.935	16.525	36.929	1.00	9.94
ATOM	276	CA	GLY	40	33.474	17.879	37.266	1.00	7.02
ATOM	277	C	GLY	40	32.952	18.600	36.004	1.00	9.45
ATOM	278	O	GLY	40	33.068	19.830	35.829	1.00	12.63
ATOM	279	N	LYS	41	32.380	17.823	35.092	1.00	5.44
ATOM	280	CA	LYS	41	31.954	18.335	33.842	1.00	6.63
ATOM	281	C	LYS	41	30.414	18.554	33.703	1.00	20.92

FIG. 5G

ATOM	282	O	LYS	41	29.617	17.693	34.085	1.00	12.94
ATOM	283	CB	LYS	41	32.360	17.357	32.827	1.00	8.27
ATOM	284	CG	LYS	41	32.099	17.771	31.419	1.00	13.19
ATOM	285	CD	LYS	41	32.521	16.644	30.481	1.00	20.20
ATOM	286	CE	LYS	41	32.690	17.068	29.032	1.00	35.79
ATOM	287	NZ	LYS	41	33.113	15.954	28.147	1.00	47.56
ATOM	288	N	LEU	42	30.049	19.684	33.069	1.00	18.31
ATOM	289	CA	LEU	42	28.643	20.064	32.794	1.00	16.08
ATOM	290	C	LEU	42	28.456	20.422	31.330	1.00	14.23
ATOM	291	O	LEU	42	29.240	21.168	30.787	1.00	14.79
ATOM	292	CB	LEU	42	28.223	21.300	33.621	1.00	13.22
ATOM	293	CG	LEU	42	28.007	21.061	35.082	1.00	16.70
ATOM	294	CD1	LEU	42	27.894	22.406	35.782	1.00	13.79
ATOM	295	CD2	LEU	42	26.732	20.243	35.295	1.00	18.70
ATOM	296	N	THR	43	27.395	19.914	30.672	1.00	8.04
ATOM	297	CA	THR	43	27.103	20.275	29.282	1.00	4.87
ATOM	298	C	THR	43	25.636	20.666	29.186	1.00	17.23
ATOM	299	O	THR	43	24.811	19.818	29.442	1.00	14.38
ATOM	300	CB	THR	43	27.351	19.140	28.317	1.00	21.59
ATOM	301	OG1	THR	43	28.692	18.743	28.415	1.00	42.74
ATOM	302	CG2	THR	43	27.073	19.675	26.917	1.00	31.23
ATOM	303	N	LEU	44	25.327	21.934	28.830	1.00	11.83
ATOM	304	CA	LEU	44	23.944	22.409	28.847	1.00	13.81
ATOM	305	C	LEU	44	23.589	23.307	27.668	1.00	18.19
ATOM	306	O	LEU	44	24.416	23.989	27.107	1.00	13.86
ATOM	307	CB	LEU	44	23.725	23.275	30.125	1.00	15.37
ATOM	308	CG	LEU	44	23.369	22.584	31.456	1.00	24.69
ATOM	309	CD1	LEU	44	21.869	22.381	31.601	1.00	23.20
ATOM	310	CD2	LEU	44	24.083	21.286	31.650	1.00	46.18
ATOM	311	N	LYS	45	22.294	23.331	27.339	1.00	10.29
ATOM	312	CA	LYS	45	21.752	24.224	26.358	1.00	11.94
ATOM	313	C	LYS	45	20.534	24.913	26.957	1.00	19.35
ATOM	314	O	LYS	45	19.665	24.248	27.530	1.00	18.43
ATOM	315	CB	LYS	45	21.409	23.560	25.060	1.00	13.75
ATOM	316	CG	LYS	45	20.878	24.556	24.045	1.00	8.83
ATOM	317	CD	LYS	45	20.486	23.863	22.746	1.00	26.87
ATOM	318	CE	LYS	45	19.574	24.688	21.842	1.00	16.58
ATOM	319	NZ	LYS	45	19.318	24.024	20.555	1.00	18.33
ATOM	320	N	PHE	46	20.535	26.236	26.910	1.00	12.34
ATOM	321	CA	PHE	46	19.463	27.048	27.451	1.00	13.32
ATOM	322	C	PHE	46	18.759	27.718	26.343	1.00	18.26

FIG. 5H

ATOM	323	O	PHE	46	19.386	28.093	25.360	1.00	16.83
ATOM	324	CB	PHE	46	19.934	28.101	28.473	1.00	15.29
ATOM	325	CG	PHE	46	20.773	27.495	29.552	1.00	13.81
ATOM	326	CD1	PHE	46	22.132	27.268	29.337	1.00	17.06
ATOM	327	CD2	PHE	46	20.209	27.121	30.774	1.00	8.24
ATOM	328	CE1	PHE	46	22.924	26.693	30.331	1.00	15.95
ATOM	329	CE2	PHE	46	20.979	26.524	31.767	1.00	11.90
ATOM	330	CZ	PHE	46	22.340	26.309	31.540	1.00	8.84
ATOM	331	N	ILE	47	17.440	27.845	26.498	1.00	13.24
ATOM	332	CA	ILE	47	16.588	28.453	25.479	1.00	18.02
ATOM	333	C	ILE	47	15.645	29.460	26.118	1.00	20.14
ATOM	334	O	ILE	47	15.039	29.162	27.148	1.00	17.67
ATOM	335	CB	ILE	47	15.737	27.386	24.801	1.00	22.67
ATOM	336	CG1	ILE	47	16.585	26.271	24.291	1.00	20.66
ATOM	337	CG2	ILE	47	15.024	28.002	23.641	1.00	33.79
ATOM	338	CD1	ILE	47	16.639	26.293	22.805	1.00	23.69
ATOM	339	N	CYS	48	15.564	30.653	25.561	1.00	14.68
ATOM	340	CA	CYS	48	14.681	31.635	26.170	1.00	16.93
ATOM	341	C	CYS	48	13.323	31.352	25.628	1.00	24.18
ATOM	342	O	CYS	48	13.122	31.513	24.453	1.00	20.63
ATOM	343	CB	CYS	48	15.063	33.116	25.885	1.00	16.85
ATOM	344	SG	CYS	48	13.913	34.268	26.712	1.00	22.06
ATOM	345	N	THR	49	12.424	30.871	26.484	1.00	27.31
ATOM	346	CA	THR	49	11.101	30.458	26.042	1.00	32.18
ATOM	347	C	THR	49	10.106	31.572	25.803	1.00	37.51
ATOM	348	O	THR	49	9.150	31.407	25.061	1.00	35.71
ATOM	349	CB	THR	49	10.537	29.417	26.972	1.00	23.66
ATOM	350	OG1	THR	49	10.387	29.989	28.258	1.00	30.10
ATOM	351	CG2	THR	49	11.512	28.226	27.022	1.00	29.98
ATOM	352	N	THR	50	10.314	32.693	26.447	1.00	32.34
ATOM	353	CA	THR	50	9.416	33.810	26.283	1.00	28.67
ATOM	354	C	THR	50	9.836	34.711	25.126	1.00	37.98
ATOM	355	O	THR	50	9.228	35.763	24.904	1.00	39.17
ATOM	356	CB	THR	50	9.251	34.611	27.589	1.00	36.23
ATOM	357	OG1	THR	50	10.512	34.980	28.118	1.00	35.37
ATOM	358	CG2	THR	50	8.507	33.773	28.602	1.00	27.78
ATOM	359	N	GLY	51	10.881	34.282	24.372	1.00	31.04
ATOM	360	CA	GLY	51	11.394	35.059	23.239	1.00	32.42
ATOM	361	C	GLY	51	12.865	35.542	23.427	1.00	48.45
ATOM	362	O	GLY	51	13.779	34.737	23.701	1.00	57.11
ATOM	363	N	LYS	52	13.087	36.862	23.282	1.00	36.08

FIG. 5I

ATOM	364	CA	LYS	52	14.416	37.460	23.416	1.00	35.75
ATOM	365	C	LYS	52	14.827	37.726	24.861	1.00	29.65
ATOM	366	O	LYS	52	14.140	38.420	25.620	1.00	25.70
ATOM	367	CB	LYS	52	14.577	38.714	22.582	1.00	43.37
ATOM	368	CG	LYS	52	15.772	38.649	21.644	1.00	78.17
ATOM	369	N	LEU	53	15.983	37.190	25.250	1.00	19.22
ATOM	370	CA	LEU	53	16.439	37.430	26.596	1.00	13.52
ATOM	371	C	LEU	53	16.717	38.932	26.775	1.00	17.76
ATOM	372	O	LEU	53	17.392	39.539	25.973	1.00	21.59
ATOM	373	CB	LEU	53	17.705	36.567	26.845	1.00	17.39
ATOM	374	CG	LEU	53	18.100	36.435	28.302	1.00	17.43
ATOM	375	CD1	LEU	53	17.048	35.621	29.053	1.00	20.12
ATOM	376	CD2	LEU	53	19.440	35.718	28.368	1.00	16.11
ATOM	377	N	PRO	54	16.197	39.525	27.817	1.00	16.69
ATOM	378	CA	PRO	54	16.324	40.962	28.092	1.00	18.60
ATOM	379	C	PRO	54	17.638	41.414	28.707	1.00	25.39
ATOM	380	O	PRO	54	17.865	42.609	28.861	1.00	18.88
ATOM	381	CB	PRO	54	15.268	41.265	29.139	1.00	22.52
ATOM	382	CG	PRO	54	14.832	39.933	29.720	1.00	26.02
ATOM	383	CD	PRO	54	15.318	38.855	28.779	1.00	21.26
ATOM	384	N	VAL	55	18.435	40.455	29.161	1.00	23.32
ATOM	385	CA	VAL	55	19.746	40.716	29.711	1.00	15.83
ATOM	386	C	VAL	55	20.688	39.868	28.973	1.00	19.38
ATOM	387	O	VAL	55	20.268	39.035	28.219	1.00	20.34
ATOM	388	CB	VAL	55	19.814	40.409	31.147	1.00	17.67
ATOM	389	CG1	VAL	55	18.864	41.340	31.851	1.00	22.52
ATOM	390	CG2	VAL	55	19.402	38.959	31.397	1.00	19.11
ATOM	391	N	PRO	56	21.963	40.070	29.167	1.00	19.37
ATOM	392	CA	PRO	56	22.911	39.258	28.447	1.00	13.09
ATOM	393	C	PRO	56	23.059	37.834	29.038	1.00	5.83
ATOM	394	O	PRO	56	23.067	37.631	30.254	1.00	12.35
ATOM	395	CB	PRO	56	24.231	40.062	28.420	1.00	18.34
ATOM	396	CG	PRO	56	23.851	41.478	28.849	1.00	20.73
ATOM	397	CD	PRO	56	22.525	41.379	29.578	1.00	18.66
ATOM	398	N	TRP	57	23.202	36.848	28.158	1.00	11.12
ATOM	399	CA	TRP	57	23.354	35.458	28.595	1.00	12.55
ATOM	400	C	TRP	57	24.411	35.239	29.700	1.00	14.13
ATOM	401	O	TRP	57	24.178	34.586	30.709	1.00	11.49
ATOM	402	CB	TRP	57	23.604	34.535	27.406	1.00	10.56
ATOM	403	CG	TRP	57	22.335	34.237	26.641	1.00	12.65
ATOM	404	CD1	TRP	57	21.999	34.714	25.426	1.00	16.24

FIG. 5J

ATOM	405	CD2TRP	57	21.281	33.327	27.013	1.00	12.50
ATOM	406	NE1TRP	57	20.784	34.200	25.018	1.00	14.25
ATOM	407	CE2TRP	57	20.315	22.354	25.963	1.00	14.65
ATOM	408	CE3TRP	57	21.052	32.521	28.129	1.00	12.01
ATOM	409	CZ2TRP	57	19.148	32.583	26.007	1.00	14.36
ATOM	410	CZ3TRP	57	19.887	31.767	28.170	1.00	14.23
ATOM	411	CH2TRP	57	18.945	31.818	27.128	1.00	10.01
ATOM	412	N PRO	58	25.594	35.800	29.518	1.00	15.78
ATOM	413	CA PRO	58	26.629	35.616	30.503	1.00	9.53
ATOM	414	C PRO	58	26.241	36.010	31.878	1.00	9.71
ATOM	415	O PRO	58	26.760	35.467	32.825	1.00	11.70
ATOM	416	CB PRO	58	27.833	36.441	30.040	1.00	10.83
ATOM	417	CG PRO	58	27.597	36.748	28.582	1.00	18.50
ATOM	418	CD PRO	58	26.137	36.432	28.278	1.00	15.82
ATOM	419	N THR	59	25.336	36.977	32.021	1.00	7.54
ATOM	420	CA THR	59	24.976	37.366	33.357	1.00	4.53
ATOM	421	C THR	59	24.228	36.258	34.137	1.00	8.41
ATOM	422	O THR	59	24.174	36.261	35.367	1.00	10.57
ATOM	423	CB THR	59	24.187	38.691	33.384	1.00	16.64
ATOM	424	OG1THR	59	22.895	38.480	32.844	1.00	15.51
ATOM	425	CG2THR	59	24.917	39.731	32.542	1.00	15.76
ATOM	426	N LEU	60	23.686	35.304	33.427	1.00	11.99
ATOM	427	CA LEU	60	22.899	34.248	34.073	1.00	9.15
ATOM	428	C LEU	60	23.657	32.944	34.385	1.00	15.62
ATOM	429	O LEU	60	23.118	32.027	35.042	1.00	11.99
ATOM	430	CB LEU	60	21.645	33.914	33.203	1.00	7.67
ATOM	431	CG LEU	60	20.728	35.111	33.042	1.00	14.06
ATOM	432	CD1LEU	60	19.620	34.775	32.062	1.00	14.54
ATOM	433	CD2LEU	60	20.142	35.456	34.394	1.00	10.67
ATOM	434	N VAL	61	24.893	32.837	33.917	1.00	11.27
ATOM	435	CA VAL	61	25.656	31.587	34.094	1.00	4.37
ATOM	436	C VAL	61	25.678	31.013	35.496	1.00	6.02
ATOM	437	O VAL	61	25.385	29.805	35.743	1.00	10.75
ATOM	438	CB VAL	61	27.050	31.643	33.406	1.00	7.14
ATOM	439	CG1VAL	61	27.888	30.396	33.805	1.00	6.47
ATOM	440	CG2VAL	61	26.890	31.745	31.876	1.00	6.63
ATOM	441	N THR	62	26.053	31.843	36.442	1.00	7.02
ATOM	442	CA THR	62	26.178	31.421	37.808	1.00	6.51
ATOM	443	C THR	62	24.862	30.954	38.410	1.00	9.22
ATOM	444	O THR	62	24.801	30.163	39.352	1.00	6.99
ATOM	445	CB THR	62	26.816	32.520	38.660	1.00	16.97

FIG. 5K

ATOM	446	OG1THR	62	26.103	33.744	38.453	1.00	12.00
ATOM	447	CG2THR	62	28.297	32.708	39.225	1.00	8.86
ATOM	448	N THR	63	23.814	31.547	37.910	1.00	9.98
ATOM	449	CA THR	63	22.457	31.212	38.388	1.00	6.69
ATOM	450	C THR	63	22.033	29.830	37.865	1.00	8.14
ATOM	451	O THR	63	21.499	28.984	38.604	1.00	13.48
ATOM	452	CB THR	63	21.458	32.312	37.295	1.00	11.14
ATOM	453	OG1THR	63	21.785	33.498	38.602	1.00	11.75
ATOM	454	OG2THR	63	20.024	31.897	38.296	1.00	9.31
ATOM	455	N PHE	64	22.250	29.620	36.583	1.00	10.19
ATOM	456	CA PHE	64	21.895	28.371	35.995	1.00	8.00
ATOM	457	C PHE	64	22.774	27.253	36.518	1.00	25.26
ATOM	458	O PHE	64	23.313	26.147	36.761	1.00	9.64
ATOM	459	CB PHE	64	22.114	28.438	34.513	1.00	6.88
ATOM	460	CG PHE	64	21.233	29.357	33.750	1.00	10.96
ATOM	461	CD1PHE	64	21.724	29.954	32.593	1.00	9.15
ATOM	462	CD2PHE	64	19.899	29.563	34.106	1.00	14.43
ATOM	463	CE1PHE	64	20.936	30.792	31.805	1.00	14.20
ATOM	464	CE2PHE	64	19.077	30.375	33.317	1.00	13.95
ATOM	465	CZ PHE	64	19.597	30.983	32.171	1.00	16.35
HETATM	466	N1 CRO	66	24.077	27.513	36.610	1.00	11.86
HETATM	467	CG1CRO	66	25.155	25.422	34.796	1.00	16.67
HETATM	468	OG1CRO	66	26.679	27.129	35.461	1.00	14.22
HETATM	469	CB1CRO	66	25.931	26.035	35.930	1.00	10.77
HETATM	470	CA1CRO	66	25.011	26.478	37.078	1.00	7.34
HETATM	471	C1 CRO	66	25.718	26.991	38.253	1.00	17.70
HETATM	472	N2 CRO	66	26.975	27.732	38.216	1.00	9.21
HETATM	473	OH CRO	66	32.894	30.804	36.971	1.00	13.84
HETATM	474	CD2CRO	66	30.487	30.110	39.805	1.00	10.79
HETATM	475	CE2CRO	66	31.614	30.563	39.085	1.00	10.01
HETATM	476	CZ CRO	66	31.718	30.300	37.721	1.00	9.48
HETATM	477	CE1CRO	66	30.707	29.546	37.033	1.00	17.44
HETATM	478	CD1CRO	66	29.541	29.103	37.742	1.00	11.31
HETATM	479	CG2CRO	66	29.437	29.370	39.124	1.00	7.67
HETATM	480	CB2CRO	66	28.329	28.822	39.960	1.00	10.75
HETATM	481	CA2CRO	66	27.197	28.245	39.512	1.00	16.08
HETATM	482	C2 CRO	66	26.043	27.875	40.370	1.00	5.46
HETATM	483	O2 CRO	66	26.022	27.962	41.566	1.00	13.20
HETATM	484	N3 CRO	66	25.240	26.978	39.517	1.00	18.43
HETATM	485	CA3CRO	66	23.840	26.511	39.734	1.00	10.40
HETATM	486	C3 CRO	66	23.413	25.550	40.817	1.00	11.96

FIG. 5L

HETATM	487	O3	CRO	66	22.747	26.014	41.764	1.00	100.00
ATOM	488	N	VAL	68	23.737	24.208	41.005	1.00	29.95
ATOM	489	CA	VAL	68	24.209	22.972	40.304	1.00	17.16
ATOM	490	C	VAL	68	25.692	22.550	40.734	1.00	14.88
ATOM	491	O	VAL	68	26.378	21.821	40.026	1.00	9.03
ATOM	492	CB	VAL	68	23.870	22.899	38.831	1.00	18.94
ATOM	493	CG1	VAL	68	24.685	22.088	37.942	1.00	17.17
ATOM	494	CG2	VAL	68	22.396	22.538	38.680	1.00	18.80
ATOM	495	N	GLN	69	26.129	22.965	41.914	1.00	11.04
ATOM	496	CA	GLN	69	27.465	22.764	42.394	1.00	15.00
ATOM	497	C	GLN	69	27.749	21.366	42.893	1.00	22.46
ATOM	498	O	GLN	69	28.876	21.026	43.154	1.00	15.84
ATOM	499	CB	GLN	69	27.929	23.852	43.414	1.00	10.93
ATOM	500	CG	GLN	69	28.202	25.174	42.615	1.00	14.13
ATOM	501	CD	GLN	69	28.216	26.385	43.520	1.00	17.01
ATOM	502	OE1	GLN	69	27.433	26.476	44.448	1.00	18.94
ATOM	503	NE2	GLN	69	29.151	27.300	43.241	1.00	8.52
ATOM	504	N	CYS	70	26.703	20.540	42.906	1.00	12.10
ATOM	505	CA	CYS	70	26.862	19.171	43.287	1.00	11.84
ATOM	506	C	CYS	70	27.611	18.391	42.175	1.00	10.54
ATOM	507	O	CYS	70	28.036	17.242	42.367	1.00	14.70
ATOM	508	CB	CYS	70	25.476	18.584	43.596	1.00	14.52
ATOM	509	SG	CYS	70	24.325	19.012	42.251	1.00	15.61
ATOM	510	N	PHE	71	27.801	19.029	41.005	1.00	8.64
ATOM	511	CA	PHE	71	28.525	18.419	39.883	1.00	6.59
ATOM	512	C	PHE	71	30.041	18.754	39.876	1.00	16.43
ATOM	513	O	PHE	71	30.753	18.481	38.916	1.00	13.05
ATOM	514	CB	PHE	71	27.951	18.771	38.523	1.00	7.61
ATOM	515	CG	PHE	71	26.669	18.016	38.303	1.00	14.73
ATOM	516	CD1	PHE	71	26.693	16.642	38.050	1.00	10.34
ATOM	517	CD2	PHE	71	25.434	18.660	38.453	1.00	17.14
ATOM	518	CE1	PHE	71	25.506	15.931	37.866	1.00	15.09
ATOM	519	CE2	PHE	71	24.238	17.961	38.300	1.00	20.92
ATOM	520	CZ	PHE	71	24.282	16.598	37.990	1.00	18.49
ATOM	521	N	SER	72	30.500	19.370	40.938	1.00	13.13
ATOM	522	CA	SER	72	31.889	19.715	41.075	1.00	11.65
ATOM	523	C	SER	72	32.689	18.446	41.357	1.00	14.56
ATOM	524	O	SER	72	32.256	17.566	42.122	1.00	10.90
ATOM	525	CB	SER	72	32.075	20.672	42.257	1.00	8.65
ATOM	526	OG	SER	72	31.361	21.874	42.038	1.00	19.29
ATOM	527	N	ARG	73	33.905	18.358	40.794	1.00	16.27

FIG. 5M

ATOM	528	CA	ARG	73	34.695	17.212	41.117	1.00	13.56
ATOM	529	C	ARG	73	35.414	17.426	42.443	1.00	19.96
ATOM	530	O	ARG	73	36.182	18.376	42.599	1.00	16.14
ATOM	531	CB	ARG	73	35.694	16.817	40.013	1.00	16.80
ATOM	532	CG	ARG	73	36.549	15.616	40.460	1.00	20.13
ATOM	533	CD	ARG	73	37.489	15.093	39.381	1.00	28.47
ATOM	534	NE	ARG	73	38.743	15.859	39.260	1.00	25.48
ATOM	535	CZ	ARG	73	39.756	15.777	40.127	1.00	28.04
ATOM	536	NH1	ARG	73	39.688	15.004	41.195	1.00	28.76
ATOM	537	NH2	ARG	73	40.865	16.504	39.918	1.00	39.65
ATOM	538	N	TYR	74	35.151	16.561	43.424	1.00	12.05
ATOM	539	CA	TYR	74	35.861	16.659	44.690	1.00	11.57
ATOM	540	C	TYR	74	36.946	15.566	44.721	1.00	25.02
ATOM	541	O	TYR	74	36.658	14.387	44.558	1.00	19.71
ATOM	542	CB	TYR	74	34.978	16.528	45.934	1.00	15.51
ATOM	543	CG	TYR	74	34.395	17.850	46.402	1.00	16.59
ATOM	544	CD1	TYR	74	33.455	18.546	45.631	1.00	14.44
ATOM	545	CD2	TYR	74	34.799	18.399	47.618	1.00	15.94
ATOM	546	CE1	TYR	74	32.901	19.756	46.059	1.00	7.99
ATOM	547	CE2	TYR	74	34.261	19.612	48.058	1.00	18.29
ATOM	548	CZ	TYR	74	33.294	20.276	47.298	1.00	13.87
ATOM	549	OH	TYR	74	32.829	21.507	47.738	1.00	18.39
ATOM	550	N	PRO	75	38.181	15.947	44.902	1.00	19.20
ATOM	551	CA	PRO	75	39.213	14.940	44.995	1.00	18.42
ATOM	552	C	PRO	75	38.958	13.993	46.175	1.00	15.60
ATOM	553	O	PRO	75	39.373	14.361	47.174	1.00	11.99
ATOM	554	CB	PRO	75	40.514	15.681	45.196	1.00	18.31
ATOM	555	CG	PRO	75	40.242	17.158	44.868	1.00	24.81
ATOM	556	CD	PRO	75	38.742	17.306	44.694	1.00	15.41
ATOM	557	N	ASP	76	39.433	12.756	46.038	1.00	18.63
ATOM	558	CA	ASP	76	39.269	11.770	47.062	1.00	16.19
ATOM	559	C	ASP	76	39.581	12.280	48.431	1.00	15.92
ATOM	560	O	ASP	76	38.862	12.042	49.389	1.00	17.35
ATOM	561	CB	ASP	76	40.083	10.507	46.790	1.00	18.69
ATOM	562	CG	ASP	76	39.826	9.432	47.825	1.00	24.04
ATOM	563	OD1	ASP	76	40.523	9.268	48.817	1.00	29.72
ATOM	564	OD2	ASP	76	38.732	8.743	47.584	1.00	40.96
ATOM	565	N	HIS	77	40.647	12.984	48.561	1.00	18.79
ATOM	566	CA	HIS	77	40.978	13.418	49.877	1.00	19.36
ATOM	567	C	HIS	77	40.117	14.507	50.397	1.00	24.57
ATOM	568	O	HIS	77	40.205	14.826	51.551	1.00	27.15

FIG. 5N

ATOM	569	CB	HIS	77	42.435	13.806	50.042	1.00	19.84
ATOM	570	CG	HIS	77	42.743	15.035	49.322	1.00	17.31
ATOM	571	ND1	HIS	77	42.925	15.028	47.953	1.00	21.86
ATOM	572	CD2	HIS	77	42.925	16.295	49.774	1.00	18.70
ATOM	573	CE1	HIS	77	43.203	16.289	47.593	1.00	17.49
ATOM	574	NE2	HIS	77	43.213	17.069	48.668	1.00	18.11
ATOM	575	N	MSE	78	39.277	15.069	49.565	1.00	25.36
ATOM	576	CA	MSE	78	38.412	16.140	50.026	1.00	24.65
ATOM	577	C	MSE	78	36.920	15.774	50.066	1.00	26.47
ATOM	578	O	MSE	78	36.070	16.636	50.260	1.00	28.16
ATOM	579	CB	MSE	78	38.596	17.331	49.121	1.00	26.38
ATOM	580	CG	MSE	78	39.803	18.177	49.406	1.00	27.01
ATOM	581	SE	MSE	78	39.987	19.608	48.177	1.00	43.09
ATOM	582	CE	MSE	78	38.874	20.873	49.044	1.00	27.11
ATOM	583	N	LYS	79	36.606	14.509	49.856	1.00	18.68
ATOM	584	CA	LYS	79	35.216	14.061	49.853	1.00	21.54
ATOM	585	C	LYS	79	34.406	14.449	51.082	1.00	20.21
ATOM	586	O	LYS	79	33.186	14.652	51.025	1.00	21.08
ATOM	587	CB	LYS	79	35.152	12.581	49.612	1.00	23.48
ATOM	588	CG	LYS	79	35.859	12.225	48.317	1.00	41.09
ATOM	589	CD	LYS	79	35.159	11.134	47.535	1.00	34.66
ATOM	590	CE	LYS	79	35.796	10.881	46.181	1.00	53.46
ATOM	591	NZ	LYS	79	35.084	11.549	45.080	1.00	49.53
ATOM	592	N	ARG	80	35.069	14.542	52.213	1.00	19.77
ATOM	593	CA	ARG	80	34.365	14.874	53.434	1.00	20.13
ATOM	594	C	ARG	80	33.898	16.311	53.481	1.00	26.42
ATOM	595	O	ARG	80	33.251	16.717	54.467	1.00	23.51
ATOM	596	CB	ARG	80	35.155	14.549	54.700	1.00	24.58
ATOM	597	CG	ARG	80	36.204	15.620	55.034	1.00	29.71
ATOM	598	CD	ARG	80	36.964	15.344	56.335	1.00	61.30
ATOM	599	NE	ARG	80	36.551	16.230	57.415	1.00	71.14
ATOM	600	CZ	ARG	80	37.398	16.882	58.192	1.00	100.00
ATOM	601	NH1	ARG	80	38.714	16.758	48.040	1.00	100.00
ATOM	602	NH2	ARG	80	36.917	17.679	59.155	1.00	99.06
ATOM	603	N	HIS	81	34.275	17.121	52.473	1.00	18.77
ATOM	604	CA	HIS	81	33.903	18.547	52.499	1.00	19.60
ATOM	605	C	HIS	81	32.841	18.883	51.486	1.00	18.62
ATOM	606	O	HIS	81	32.557	20.043	51.295	1.00	17.76
ATOM	607	CB	HIS	81	35.129	19.472	52.283	1.00	20.39
ATOM	608	CG	HIS	81	36.221	19.224	53.305	1.00	28.02
ATOM	609	ND1	HIS	81	36.127	19.701	54.618	1.00	30.59

FIG. 50

ATOM	610	CD2HIS	81	37.392	18.535	53.202	1.00	29.02
ATOM	611	CE1HIS	81	37.218	19.308	55.265	1.00	26.24
ATOM	612	NE2HIS	81	37.991	18.603	54.452	1.00	28.18
ATOM	613	N ASP	82	32.298	17.843	50.841	1.00	12.20
ATOM	614	CA ASP	82	31.358	18.011	49.769	1.00	13.24
ATOM	615	C ASP	82	29.922	18.148	50.259	1.00	24.30
ATOM	616	O ASP	82	29.175	17.195	50.243	1.00	16.55
ATOM	617	CB ASP	82	31.480	16.917	48.730	1.00	12.23
ATOM	618	CG ASP	82	30.642	17.209	47.518	1.00	9.92
ATOM	619	OD1ASP	82	29.870	18.134	47.459	1.00	20.31
ATOM	620	OD2ASP	82	30.938	16.466	46.507	1.00	11.12
ATOM	621	N PHE	83	29.566	19.353	50.705	1.00	23.66
ATOM	622	CA PHE	83	28.220	19.634	51.201	1.00	20.23
ATOM	623	C PHE	83	27.154	19.333	50.168	1.00	20.93
ATOM	624	O PHE	83	26.116	18.733	50.503	1.00	15.97
ATOM	625	CB PHE	83	28.077	21.106	51.666	1.00	19.59
ATOM	626	CG PHE	83	26.624	21.613	51.805	1.00	16.91
ATOM	627	CD1PHE	83	25.946	21.498	53.021	1.00	17.76
ATOM	628	CD2PHE	83	25.968	22.236	50.734	1.00	18.88
ATOM	629	CE1PHE	83	24.635	21.960	53.156	1.00	24.13
ATOM	630	CE2PHE	83	24.650	22.690	50.840	1.00	19.24
ATOM	631	CZ PHE	83	24.001	22.575	52.068	1.00	20.67
ATOM	632	N PHE	84	27.432	19.784	48.921	1.00	14.06
ATOM	633	CA PHE	84	26.515	19.693	47.809	1.00	12.96
ATOM	634	C PHE	84	25.893	18.332	47.602	1.00	24.96
ATOM	635	O PHE	84	24.674	18.200	47.534	1.00	21.55
ATOM	636	CB PHE	84	27.085	20.265	46.513	1.00	13.44
ATOM	637	CG PHE	84	27.630	21.645	46.721	1.00	14.27
ATOM	638	CD1PHE	84	29.001	21.845	46.890	1.00	15.17
ATOM	639	CD2PHE	84	26.781	22.753	46.752	1.00	13.48
ATOM	640	CE1PHE	84	29.520	23.129	47.073	1.00	14.63
ATOM	641	CE2PHE	84	27.276	24.041	46.969	1.00	16.34
ATOM	642	CZ PHE	84	28.650	24.221	47.137	1.00	15.77
ATOM	643	N LYS	85	26.738	17.330	47.482	1.00	14.07
ATOM	644	CA LYS	85	26.294	15.985	47.283	1.00	13.30
ATOM	645	C LYS	85	25.657	15.371	48.547	1.00	13.43
ATOM	646	O LYS	85	24.773	14.509	48.429	1.00	18.46
ATOM	647	CB LYS	85	27.434	15.089	46.757	1.00	17.38
ATOM	648	CG LYS	85	27.873	15.372	45.323	1.00	13.93
ATOM	649	CD LYS	85	28.969	14.381	44.888	1.00	13.23
ATOM	650	CE LYS	85	29.766	14.819	43.662	1.00	10.36

FIG. 5P

ATOM	651	NZ	LYS	85	30.319	16.185	43.773	1.00	12.92
ATOM	652	N	SER	86	26.119	15.795	49.752	1.00	11.03
ATOM	653	CA	SER	86	25.610	15.267	50.998	1.00	12.09
ATOM	654	C	SER	86	24.156	15.639	51.240	1.00	21.58
ATOM	655	O	SER	86	23.452	14.979	52.013	1.00	19.89
ATOM	656	CB	SER	86	26.448	15.661	52.208	1.00	16.45
ATOM	657	OG	SER	86	26.308	17.042	52.495	1.00	22.05
ATOM	658	N	ALA	87	23.705	16.698	50.582	1.00	15.09
ATOM	659	CA	ALA	87	22.333	17.138	50.762	1.00	19.52
ATOM	660	C	ALA	87	21.337	16.399	49.870	1.00	18.60
ATOM	661	O	ALA	87	20.162	16.557	50.040	1.00	19.55
ATOM	662	CB	ALA	87	22.204	18.647	50.632	1.00	19.23
ATOM	663	N	MSE	88	21.835	15.536	48.976	1.00	14.05
ATOM	664	CA	MSE	88	21.007	14.796	48.035	1.00	15.32
ATOM	665	C	MSE	88	20.496	13.448	48.579	1.00	21.48
ATOM	666	O	MSE	88	21.109	12.876	49.457	1.00	23.03
ATOM	667	CB	MSE	88	21.848	14.593	46.791	1.00	16.98
ATOM	668	CG	MSE	88	22.263	15.891	46.131	1.00	10.66
ATOM	669	SE	MSE	88	20.737	16.894	45.394	1.00	31.99
ATOM	670	CE	MSE	88	21.318	18.684	45.748	1.00	28.86
ATOM	671	N	PRO	89	19.363	12.930	48.084	1.00	14.78
ATOM	672	CA	PRO	89	18.552	13.475	47.008	1.00	14.80
ATOM	673	C	PRO	89	17.572	14.611	47.385	1.00	12.10
ATOM	674	O	PRO	89	17.085	15.301	46.493	1.00	18.06
ATOM	675	CB	PRO	89	17.733	12.294	46.494	1.00	17.00
ATOM	676	CG	PRO	89	17.726	11.261	47.607	1.00	15.83
ATOM	677	CD	PRO	89	18.844	11.642	48.560	1.00	17.16
ATOM	678	N	GLU	90	17.278	14.795	48.695	1.00	14.63
ATOM	679	CA	GLU	90	16.348	15.838	49.157	1.00	20.68
ATOM	680	C	GLU	90	16.701	17.229	48.645	1.00	25.59
ATOM	681	O	GLU	90	15.833	18.042	48.368	1.00	21.57
ATOM	682	CB	GLU	90	16.031	15.816	50.682	1.00	22.21
ATOM	683	CG	GLU	90	15.782	14.403	51.228	1.00	37.69
ATOM	684	CD	GLU	90	17.071	13.641	51.447	1.00	83.49
ATOM	685	OE1	GLU	90	18.179	14.151	51.342	1.00	54.80
ATOM	686	OE2	GLU	90	16.875	12.373	51.749	1.00	64.65
ATOM	687	N	GLY	91	17.977	17.509	48.510	1.00	21.39
ATOM	688	CA	GLY	91	18.394	18.769	47.906	1.00	17.77
ATOM	689	C	GLY	91	18.673	19.911	48.839	1.00	12.17
ATOM	690	O	GLY	91	18.769	19.764	50.055	1.00	16.81
ATOM	691	N	TYR	92	18.861	21.086	48.225	1.00	13.02

FIG. 5Q

ATOM	692	CA	TYR	92	19.143	22.266	48.994	1.00	10.33
ATOM	693	C	TYR	92	18.575	23.478	48.347	1.00	9.87
ATOM	694	O	TYR	92	18.270	23.483	47.144	1.00	15.89
ATOM	695	CB	TYR	92	20.678	22.488	49.278	1.00	15.40
ATOM	696	CG	TYR	92	21.546	22.468	48.012	1.00	15.13
ATOM	697	CD1	TYR	92	21.620	23.576	47.166	1.00	14.75
ATOM	698	CD2	TYR	92	22.317	21.350	47.683	1.00	16.09
ATOM	699	CE1	TYR	92	22.404	23.561	46.006	1.00	6.50
ATOM	700	CE2	TYR	92	23.067	21.300	46.504	1.00	15.12
ATOM	701	CZ	TYR	92	23.156	22.424	45.683	1.00	18.13
ATOM	702	OH	TYR	92	23.944	22.393	44.517	1.00	13.37
ATOM	703	N	VAL	93	18.447	24.504	49.189	1.00	11.93
ATOM	704	CA	VAL	93	18.025	25.822	48.778	1.00	14.74
ATOM	705	C	VAL	93	19.281	26.666	48.625	1.00	16.00
ATOM	706	O	VAL	93	20.172	26.625	49.451	1.00	16.16
ATOM	707	CB	VAL	93	17.073	26.480	49.791	1.00	23.45
ATOM	708	CG1	VAL	93	16.855	27.937	49.413	1.00	26.05
ATOM	709	CG2	VAL	93	15.716	25.764	49.771	1.00	22.90
ATOM	710	N	GLN	94	19.361	27.345	47.521	1.00	13.78
ATOM	711	CA	GLN	94	20.480	28.195	47.227	1.00	10.53
ATOM	712	C	GLN	94	19.948	29.583	46.998	1.00	12.23
ATOM	713	O	GLN	94	19.153	29.788	46.061	1.00	15.52
ATOM	714	CB	GLN	94	21.232	27.727	45.934	1.00	7.95
ATOM	715	CG	GLN	94	22.361	28.708	45.469	1.00	11.87
ATOM	716	CD	GLN	94	23.431	27.999	44.632	1.00	12.04
ATOM	717	OE1	GLN	94	23.805	26.879	44.946	1.00	13.60
ATOM	718	NE2	GLN	94	23.719	28.527	43.449	1.00	7.98
ATOM	719	N	GLU	95	20.396	30.531	47.820	1.00	11.78
ATOM	720	CA	GLU	95	19.974	31.899	47.643	1.00	13.47
ATOM	721	C	GLU	95	21.149	32.804	47.398	1.00	18.42
ATOM	722	O	GLU	95	22.206	32.623	47.985	1.00	19.23
ATOM	723	CB	GLU	95	19.277	32.427	48.878	1.00	13.52
ATOM	724	CG	GLU	95	18.009	31.684	49.215	1.00	28.46
ATOM	725	CD	GLU	95	17.657	32.016	50.622	1.00	45.93
ATOM	726	OE1	GLU	95	17.574	33.166	51.011	1.00	100.00
ATOM	727	OE2	GLU	95	17.764	30.987	51.423	1.00	61.33
ATOM	728	N	ARG	96	20.929	33.838	46.601	1.00	16.51
ATOM	729	CA	ARG	96	21.978	34.783	46.342	1.00	16.87
ATOM	730	C	ARG	96	21.510	36.195	46.206	1.00	15.84
ATOM	731	O	ARG	96	20.389	36.488	45.806	1.00	15.01
ATOM	732	CB	ARG	96	22.582	34.463	44.967	1.00	16.19

FIG. 5R

ATOM	733	CG	ARG	96	23.495	33.247	44.929	1.00	17.61
ATOM	734	CD	ARG	96	24.615	33.453	43.908	1.00	9.06
ATOM	735	NE	ARG	96	25.411	32.277	43.766	1.00	9.88
ATOM	736	CZ	ARG	96	25.434	31.493	42.693	1.00	20.03
ATOM	737	NH1	ARG	96	24.684	31.709	41.615	1.00	15.29
ATOM	738	NH2	ARG	96	26.236	30.430	42.714	1.00	11.03
ATOM	739	N	THR	97	22.470	37.068	46.344	1.00	13.39
ATOM	740	CA	THR	97	22.368	38.424	45.935	1.00	13.12
ATOM	741	C	THR	97	23.593	38.688	45.084	1.00	16.81
ATOM	742	O	THR	97	24.686	38.347	45.485	1.00	19.25
ATOM	743	CB	THR	97	22.282	39.442	47.066	1.00	26.27
ATOM	744	OG1	THR	97	21.225	39.101	47.945	1.00	31.43
ATOM	745	CG2	THR	97	22.038	40.804	46.445	1.00	15.90
ATOM	746	N	ILE	98	23.396	39.219	43.899	1.00	16.23
ATOM	747	CA	ILE	98	24.486	39.526	42.977	1.00	16.70
ATOM	748	C	ILE	98	24.533	41.017	42.686	1.00	21.10
ATOM	749	O	ILE	98	23.628	41.566	42.075	1.00	14.58
ATOM	750	CB	ILE	98	24.385	38.752	41.660	1.00	13.47
ATOM	751	CG1	ILE	98	24.480	37.236	41.890	1.00	16.09
ATOM	752	CG2	ILE	98	25.457	39.231	40.679	1.00	13.30
ATOM	753	CD1	ILE	98	23.875	36.431	40.738	1.00	13.93
ATOM	754	N	PHE	99	25.613	41.678	43.110	1.00	14.86
ATOM	755	CA	PHE	99	25.719	43.098	42.896	1.00	12.44
ATOM	756	C	PHE	99	26.514	43.441	41.699	1.00	20.37
ATOM	757	O	PHE	99	27.696	43.164	41.700	1.00	20.07
ATOM	758	CB	PHE	99	26.401	43.770	44.084	1.00	15.96
ATOM	759	CG	PHE	99	25.638	43.624	45.356	1.00	21.41
ATOM	760	CD1	PHE	99	25.863	42.524	46.189	1.00	24.98
ATOM	761	CD2	PHE	99	24.698	44.585	45.743	1.00	22.94
ATOM	762	CE1	PHE	99	25.176	42.400	47.400	1.00	32.06
ATOM	763	CE2	PHE	99	23.992	44.469	46.946	1.00	24.26
ATOM	764	CZ	PHE	99	24.235	43.369	47.771	1.00	28.19
ATOM	765	N	PHE	100	25.906	44.085	40.704	1.00	12.53
ATOM	766	CA	PHE	100	26.679	44.522	39.554	1.00	8.75
ATOM	767	C	PHE	100	27.294	45.855	39.872	1.00	21.81
ATOM	768	O	PHE	100	26.599	46.775	40.308	1.00	20.31
ATOM	769	CB	PHE	100	25.927	44.572	38.226	1.00	5.94
ATOM	770	CG	PHE	100	25.537	43.183	37.764	1.00	12.75
ATOM	771	CD1	PHE	100	24.426	42.538	38.325	1.00	16.31
ATOM	772	CD2	PHE	100	26.317	42.484	36.843	1.00	15.27
ATOM	773	CE1	PHE	100	24.087	41.230	37.975	1.00	13.50

FIG. 5S

ATOM	774	CE2 PHE	100	25.965	41.192	36.435	1.00	21.25
ATOM	775	CZ PHE	100	24.852	40.567	37.014	1.00	21.06
ATOM	776	N LYS	101	28.603	45.946	39.737	1.00	15.49
ATOM	777	CA LYS	101	29.270	47.179	40.085	1.00	17.93
ATOM	778	C LYS	101	28.732	48.349	39.287	1.00	13.71
ATOM	779	O LYS	101	28.658	48.304	38.072	1.00	17.18
ATOM	780	CB LYS	101	30.784	47.069	39.950	1.00	17.13
ATOM	781	CG LYS	101	31.518	48.252	40.551	1.00	18.01
ATOM	782	CD LYS	101	33.036	48.060	40.534	1.00	26.70
ATOM	783	CE LYS	101	33.797	49.116	41.332	1.00	41.58
ATOM	784	N ASP	102	28.353	49.403	39.997	1.00	18.09
ATOM	785	CA ASP	102	27.805	50.618	39.368	1.00	23.08
ATOM	786	C ASP	102	26.559	50.356	38.549	1.00	25.42
ATOM	787	O ASP	102	26.292	51.061	37.586	1.00	23.34
ATOM	788	CB ASP	102	28.840	51.369	38.516	1.00	26.27
ATOM	789	CG ASP	102	30.109	51.629	39.296	1.00	57.01
ATOM	790	OD1ASP	102	31.206	51.233	38.931	1.00	63.33
ATOM	791	OD2ASP	102	29.886	52.200	40.464	1.00	47.66
ATOM	792	N ASP	103	25.813	49.328	38.933	1.00	20.17
ATOM	793	CA ASP	103	24.602	48.949	38.233	1.00	15.70
ATOM	794	C ASP	103	23.608	48.284	39.189	1.00	18.47
ATOM	795	O ASP	103	23.749	48.431	40.409	1.00	17.72
ATOM	796	CB ASP	103	24.899	48.085	36.995	1.00	19.89
ATOM	797	CG ASP	103	23.946	48.387	35.860	1.00	23.93
ATOM	798	OD1ASP	103	24.238	48.274	34.688	1.00	19.05
ATOM	799	OD2ASP	103	22.774	48.809	36.283	1.00	23.89
ATOM	800	N GLY	104	22.612	47.542	38.646	1.00	20.17
ATOM	801	CA GLY	104	21.598	46.900	39.498	1.00	20.22
ATOM	802	C GLY	104	22.055	45.619	40.180	1.00	24.68
ATOM	803	O GLY	104	23.202	45.211	40.085	1.00	18.06
ATOM	804	N ASN	105	21.125	44.967	40.872	1.00	15.71
ATOM	805	CA ASN	105	21.425	43.703	41.510	1.00	8.89
ATOM	806	C ASN	105	20.399	42.620	41.181	1.00	21.85
ATOM	807	O ASN	105	19.255	42.911	40.824	1.00	15.17
ATOM	808	CB ASN	105	21.605	43.840	43.001	1.00	8.58
ATOM	809	CG ASN	105	20.359	44.366	43.697	1.00	43.57
ATOM	810	OD1ASN	105	19.565	43.601	44.259	1.00	36.67
ATOM	811	ND2ASN	105	20.178	45.674	43.659	1.00	36.47
ATOM	812	N TYR	106	20.826	41.365	41.328	1.00	16.80
ATOM	813	CA TYR	106	19.966	40.219	41.156	1.00	13.90
ATOM	814	C TYR	106	19.763	39.543	42.475	1.00	11.05

FIG. 5T

ATOM	815	O	TYR	106	20.678	39.404	43.281	1.00	13.86
ATOM	816	CB	TYR	106	20.547	39.128	40.246	1.00	15.88
ATOM	817	CG	TYR	106	20.619	39.398	38.793	1.00	15.57
ATOM	818	CD1	TYR	106	19.952	40.458	38.178	1.00	13.14
ATOM	819	CD2	TYR	106	21.373	38.524	38.006	1.00	13.35
ATOM	820	CE1	TYR	106	20.038	40.632	36.793	1.00	13.44
ATOM	821	CE2	TYR	106	21.481	38.692	36.628	1.00	10.87
ATOM	822	CZ	TYR	106	20.814	39.751	36.025	1.00	15.93
ATOM	823	OH	TYR	106	20.970	39.931	34.670	1.00	17.32
ATOM	824	N	LYS	107	18.538	39.115	42.709	1.00	12.39
ATOM	825	CA	LYS	107	18.194	38.349	43.897	1.00	11.51
ATOM	826	C	LYS	107	17.619	37.037	43.397	1.00	17.25
ATOM	827	O	LYS	107	16.704	37.010	42.562	1.00	13.14
ATOM	828	CB	LYS	107	17.217	39.063	44.823	1.00	14.82
ATOM	829	CG	LYS	107	17.860	39.631	46.060	1.00	40.71
ATOM	830	CD	LYS	107	18.528	40.974	45.793	1.00	43.48
ATOM	831	N	THR	108	18.205	35.951	43.835	1.00	14.95
ATOM	832	CA	THR	108	17.774	34.658	43.352	1.00	11.97
ATOM	833	C	THR	108	17.463	33.696	44.468	1.00	15.81
ATOM	834	O	THR	108	18.043	33.734	45.582	1.00	18.68
ATOM	835	CB	THR	108	18.847	34.034	42.410	1.00	23.81
ATOM	836	OG1	THR	108	20.064	33.791	43.137	1.00	13.88
ATOM	837	CG2	THR	108	19.123	34.968	41.264	1.00	13.04
ATOM	838	N	ARG	109	16.560	32.804	44.154	1.00	13.57
ATOM	839	CA	ARG	109	16.212	31.751	45.048	1.00	12.56
ATOM	840	C	ARG	109	15.939	30.498	44.254	1.00	13.07
ATOM	841	O	ARG	109	15.239	30.509	43.249	1.00	12.52
ATOM	842	CB	ARG	109	15.069	32.100	45.959	1.00	17.32
ATOM	843	CG	ARG	109	14.767	30.995	46.932	1.00	17.92
ATOM	844	CD	ARG	109	13.400	31.160	47.610	1.00	19.99
ATOM	845	NE	ARG	109	12.821	29.854	47.883	1.00	36.05
ATOM	846	CZ	ARG	109	12.968	29.244	49.035	1.00	55.71
ATOM	847	NH1	ARG	109	13.630	29.815	50.046	1.00	44.11
ATOM	848	NH2	ARG	109	12.432	28.041	49.195	1.00	94.34
ATOM	849	N	ALA	110	16.577	29.414	44.635	1.00	13.26
ATOM	850	CA	ALA	110	16.377	28.207	43.870	1.00	12.68
ATOM	851	C	ALA	110	16.346	26.979	44.734	1.00	13.15
ATOM	852	O	ALA	110	16.829	26.965	45.869	1.00	16.75
ATOM	853	CB	ALA	110	17.465	28.059	42.822	1.00	17.31
ATOM	854	N	GLU	111	15.770	25.939	44.176	1.00	15.39
ATOM	855	CA	GLU	111	15.741	24.655	44.823	1.00	15.24

FIG. 5U

ATOM	856	C	GLU	111	16.438	23.678	43.926	1.00	12.08
ATOM	857	O	GLU	111	16.086	23.545	42.771	1.00	15.70
ATOM	858	CB	GLU	111	14.303	24.123	44.993	1.00	19.20
ATOM	859	CG	GLU	111	13.744	24.242	46.399	1.00	38.62
ATOM	860	CD	GLU	111	12.247	24.280	46.372	1.00	60.99
ATOM	861	OE1	GLU	111	11.589	23.843	45.432	1.00	76.05
ATOM	862	OE2	GLU	111	11.742	24.956	47.380	1.00	54.87
ATOM	863	N	VAL	112	17.438	22.965	44.457	1.00	10.78
ATOM	864	CA	VAL	112	18.063	21.978	43.631	1.00	10.98
ATOM	865	C	VAL	112	17.968	20.630	44.261	1.00	8.62
ATOM	866	O	VAL	112	18.271	20.438	45.432	1.00	15.63
ATOM	867	CB	VAL	112	19.428	22.358	43.012	1.00	22.75
ATOM	868	CG1	VAL	112	19.966	23.704	43.487	1.00	16.69
ATOM	869	CG2	VAL	112	20.452	21.232	43.078	1.00	18.47
ATOM	870	N	LYS	113	17.415	19.732	43.516	1.00	14.67
ATOM	871	CA	LYS	113	17.175	18.421	44.045	1.00	16.41
ATOM	872	C	LYS	113	16.822	17.485	42.931	1.00	7.11
ATOM	873	O	LYS	113	16.695	17.893	41.808	1.00	16.27
ATOM	874	CB	LYS	113	16.032	18.497	45.036	1.00	22.50
ATOM	875	CG	LYS	113	14.792	19.084	44.376	1.00	20.40
ATOM	876	CD	LYS	113	13.509	18.321	44.703	1.00	44.65
ATOM	877	CE	LYS	113	12.526	19.134	45.528	1.00	54.02
ATOM	878	NZ	LYS	113	12.379	20.518	45.036	1.00	100.00
ATOM	879	N	PHE	114	16.683	16.208	43.267	1.00	10.09
ATOM	880	CA	PHE	114	16.325	15.175	42.317	1.00	11.41
ATOM	881	C	PHE	114	14.806	14.975	42.181	1.00	14.18
ATOM	882	O	PHE	114	14.110	14.878	43.160	1.00	15.03
ATOM	883	CB	PHE	114	16.866	13.838	42.838	1.00	12.89
ATOM	884	CG	PHE	114	18.231	13.536	42.338	1.00	16.80
ATOM	885	CD1	PHE	114	19.344	13.795	43.139	1.00	18.61
ATOM	886	CD2	PHE	114	18.403	13.009	41.056	1.00	19.50
ATOM	887	CE1	PHE	114	20.627	13.500	42.665	1.00	22.78
ATOM	888	CE2	PHE	114	19.673	12.708	40.572	1.00	25.36
ATOM	889	CZ	PHE	114	20.780	12.953	41.387	1.00	23.99
ATOM	890	N	GLU	115	14.354	14.819	40.966	1.00	15.29
ATOM	891	CA	GLU	115	12.978	14.473	40.642	1.00	11.40
ATOM	892	C	GLU	115	13.121	13.193	39.906	1.00	13.30
ATOM	893	O	GLU	115	13.434	13.207	38.730	1.00	18.72
ATOM	894	CB	GLU	115	12.348	15.481	39.667	1.00	9.68
ATOM	895	CG	GLU	115	11.856	16.747	40.376	1.00	19.54
ATOM	896	CD	GLU	115	10.742	16.460	41.342	1.00	38.12

FIG. 5V

ATOM	897	OE1 GLU	115	10.181	15.395	41.431	1.00	34.84
ATOM	898	OE2 GLU	115	10.460	17.461	42.079	1.00	27.88
ATOM	899	N GLY	116	13.005	12.087	40.585	1.00	14.51
ATOM	900	CA GLY	116	13.225	10.861	39.869	1.00	15.91
ATOM	901	C GLY	116	14.727	10.767	39.641	1.00	23.59
ATOM	902	O GLY	116	15.516	10.922	40.570	1.00	19.35
ATOM	903	N ASP	117	15.137	10.564	38.439	1.00	20.26
ATOM	904	CA ASP	117	16.572	10.462	38.233	1.00	28.00
ATOM	905	C ASP	117	17.237	11.677	37.598	1.00	22.39
ATOM	906	O ASP	117	18.423	11.672	37.265	1.00	21.38
ATOM	907	CB ASP	117	17.055	9.074	37.733	1.00	33.06
ATOM	908	CG ASP	117	16.624	8.677	36.348	1.00	55.04
ATOM	909	OD1ASP	117	16.230	9.468	35.495	1.00	59.57
ATOM	910	OD2ASP	117	16.805	7.391	36.130	1.00	82.48
ATOM	911	N THR	118	16.463	12.729	37.493	1.00	19.62
ATOM	912	CA THR	118	16.889	13.981	36.910	1.00	18.21
ATOM	913	C THR	118	17.186	14.988	37.976	1.00	18.92
ATOM	914	O THR	118	16.498	15.064	38.996	1.00	15.94
ATOM	915	CB THR	118	15.806	14.497	35.952	1.00	19.03
ATOM	916	OG1THR	118	15.552	13.508	34.990	1.00	21.42
ATOM	917	CG2THR	118	16.217	15.793	35.275	1.00	15.49
ATOM	918	N LEU	119	18.284	15.681	37.805	1.00	13.66
ATOM	919	CA LEU	119	18.679	16.706	38.759	1.00	13.50
ATOM	920	C LEU	119	18.036	17.992	38.269	1.00	8.81
ATOM	921	O LEU	119	18.194	18.368	37.091	1.00	12.49
ATOM	922	CB LEU	119	20.243	16.815	38.839	1.00	12.25
ATOM	923	CG LEU	119	20.845	17.678	39.951	1.00	3.90
ATOM	924	CD1LEU	119	20.701	19.167	39.669	1.00	10.11
ATOM	925	CD2LEU	119	20.366	17.311	41.333	1.00	7.86
ATOM	926	N VAL	120	17.230	18.595	39.170	1.00	13.34
ATOM	927	CA VAL	120	16.466	19.797	38.859	1.00	13.77
ATOM	928	C VAL	120	16.929	21.039	39.587	1.00	8.56
ATOM	929	O VAL	120	17.135	21.039	40.762	1.00	13.32
ATOM	930	CB VAL	120	14.939	19.566	39.082	1.00	17.60
ATOM	931	CG1VAL	120	14.133	20.790	38.642	1.00	17.58
ATOM	932	CG2VAL	120	14.501	18.351	38.246	1.00	15.35
ATOM	933	N ASN	121	17.067	22.111	38.839	1.00	12.24
ATOM	934	CA ASN	121	17.424	23.405	39.400	1.00	11.78
ATOM	935	C ASN	121	16.301	24.382	39.060	1.00	11.18
ATOM	936	O ASN	121	16.195	24.802	37.934	1.00	11.09
ATOM	937	CB ASN	121	18.753	23.928	38.791	1.00	11.41

FIG. 5W

ATOM	938	CG	ASN	121	19.201	25.261	39.367	1.00	11.07
ATOM	939	OD	ASN	121	18.773	25.654	40.461	1.00	12.06
ATOM	940	ND2	ASN	121	20.124	25.938	38.670	1.00	11.90
ATOM	941	N	ARG	122	15.470	24.706	40.029	1.00	13.69
ATOM	942	CA	ARG	122	14.348	25.610	39.825	1.00	12.99
ATOM	943	C	ARG	122	14.622	26.946	40.498	1.00	5.89
ATOM	944	O	ARG	122	14.749	27.011	41.723	1.00	14.47
ATOM	945	CB	ARG	122	13.068	25.025	40.417	1.00	15.99
ATOM	946	CG	ARG	122	12.478	23.921	39.589	1.00	30.23
ATOM	947	CD	ARG	122	11.282	23.244	40.281	1.00	60.61
ATOM	948	N	ILE	123	14.663	27.992	39.680	1.00	11.46
ATOM	949	CA	ILE	123	15.030	29.340	40.095	1.00	11.86
ATOM	950	C	ILE	123	13.991	30.450	39.835	1.00	10.54
ATOM	951	O	ILE	123	13.370	30.535	38.765	1.00	12.83
ATOM	952	CB	ILE	123	16.296	29.757	39.292	1.00	15.41
ATOM	953	CG1	ILE	123	17.316	28.585	39.180	1.00	12.27
ATOM	954	CG2	ILE	123	16.944	30.993	39.918	1.00	14.01
ATOM	955	CD1	ILE	123	17.652	28.242	37.743	1.00	7.74
ATOM	956	N	GLU	124	13.953	31.358	40.793	1.00	11.36
ATOM	957	CA	GLU	124	13.189	32.752	40.700	1.00	15.20
ATOM	958	C	GLU	124	14.168	33.713	40.811	1.00	11.93
ATOM	959	O	GLU	124	14.919	33.797	41.780	1.00	15.61
ATOM	960	CB	GLU	124	12.028	32.677	41.751	1.00	19.74
ATOM	961	CG	GLU	124	12.387	33.337	43.089	1.00	72.94
ATOM	962	N	LEU	125	14.183	34.550	39.808	1.00	12.19
ATOM	963	CA	LEU	125	15.092	35.654	39.767	1.00	15.00
ATOM	964	C	LEU	125	14.420	37.011	39.722	1.00	19.35
ATOM	965	O	LEU	125	13.563	37.267	38.893	1.00	18.41
ATOM	966	CB	LEU	125	15.976	35.533	38.510	1.00	14.29
ATOM	967	CG	LEU	125	17.003	36.683	38.375	1.00	17.65
ATOM	968	CD1	LEU	125	18.302	36.083	37.849	1.00	13.46
ATOM	969	CD2	LEU	125	16.511	37.732	37.367	1.00	12.09
ATOM	970	N	LYS	126	14.890	37.897	40.554	1.00	12.73
ATOM	971	CA	LYS	126	14.391	39.260	40.579	1.00	15.92
ATOM	972	C	LYS	126	15.563	40.276	40.445	1.00	18.53
ATOM	973	O	LYS	126	16.489	40.246	41.246	1.00	19.86
ATOM	974	CB	LYS	126	13.611	39.487	41.877	1.00	17.31
ATOM	975	CG	LYS	126	12.853	40.786	41.923	1.00	33.94
ATOM	976	CD	LYS	126	11.366	40.601	41.675	1.00	60.87
ATOM	977	CE	LYS	126	10.652	41.929	41.521	1.00	52.70
ATOM	978	NZ	LYS	126	11.229	42.988	42.367	1.00	47.22

FIG. 5X

ATOM	979	N	GLY	127	15.514	41.127	39.411	1.00	18.71
ATOM	980	CA	GLY	127	16.551	42.151	39.121	1.00	17.32
ATOM	981	C	GLY	127	16.012	43.572	39.272	1.00	25.32
ATOM	982	O	GLY	127	14.981	43.908	38.693	1.00	20.14
ATOM	983	N	ILE	128	16.706	44.404	40.070	1.00	18.42
ATOM	984	CA	ILE	128	16.282	45.787	40.243	1.00	21.04
ATOM	985	C	ILE	128	17.405	46.789	40.196	1.00	25.93
ATOM	986	O	ILE	128	18.562	46.496	40.429	1.00	19.37
ATOM	987	CB	ILE	128	15.482	46.052	41.504	1.00	23.82
ATOM	988	CG1	ILE	128	16.408	45.888	42.701	1.00	23.86
ATOM	989	CG2	ILE	128	14.272	45.120	41.577	1.00	28.95
ATOM	990	CD1	ILE	128	15.824	46.391	44.013	1.00	29.89
ATOM	991	N	ASP	129	16.999	48.002	39.918	1.00	20.26
ATOM	992	CA	ASP	129	17.861	49.124	39.882	1.00	18.53
ATOM	993	C	ASP	129	18.864	49.086	38.801	1.00	20.36
ATOM	994	O	ASP	129	19.949	49.632	38.953	1.00	24.28
ATOM	995	CB	ASP	129	18.498	49.407	41.253	1.00	20.57
ATOM	996	CG	ASP	129	17.545	50.077	42.226	1.00	43.70
ATOM	997	OD1	ASP	129	16.653	50.842	41.883	1.00	49.42
ATOM	998	OD2	ASP	129	17.770	49.740	43.475	1.00	38.07
ATOM	999	N	PHE	130	18.510	48.493	37.693	1.00	16.40
ATOM	1000	CA	PHE	130	19.433	48.459	36.563	1.00	16.99
ATOM	1001	C	PHE	130	19.330	49.732	35.576	1.00	35.37
ATOM	1002	O	PHE	130	18.242	50.318	35.623	1.00	27.34
ATOM	1003	CB	PHE	130	19.248	47.223	35.657	1.00	18.07
ATOM	1004	CG	PHE	130	19.809	45.980	36.312	1.00	19.10
ATOM	1005	CD1	PHE	130	19.021	45.210	37.171	1.00	16.15
ATOM	1006	CD2	PHE	130	21.126	45.572	36.073	1.00	19.17
ATOM	1007	CE1	PHE	130	19.536	44.074	37.801	1.00	23.37
ATOM	1008	CE2	PHE	130	21.665	44.445	36.703	1.00	21.11
ATOM	1009	CZ	PHE	130	20.867	43.703	37.575	1.00	22.13
ATOM	1010	N	LYS	131	20.464	50.169	35.218	1.00	31.09
ATOM	1011	CA	LYS	131	20.477	51.371	34.400	1.00	27.52
ATOM	1012	C	LYS	131	20.105	51.045	32.992	1.00	25.57
ATOM	1013	O	LYS	131	20.695	50.169	32.343	1.00	22.97
ATOM	1014	CB	LYS	131	21.796	52.109	34.438	1.00	32.64
ATOM	1015	CG	LYS	131	22.153	52.633	35.813	1.00	38.34
ATOM	1016	CD	LYS	131	23.646	52.886	35.975	1.00	75.76
ATOM	1017	N	GLU	132	19.116	51.751	32.509	1.00	26.88
ATOM	1018	CA	GLU	132	18.623	51.484	31.189	1.00	28.42
ATOM	1019	C	GLU	132	19.710	51.514	30.140	1.00	36.19

FIG. 5Y

ATOM	1020 O	GLU	132	19.617	50.862	29.101	1.00	39.24
ATOM	1021 CB	GLU	132	17.374	52.331	30.830	1.00	29.04
ATOM	1022 N	ASP	133	20.752	52.254	30.438	1.00	40.08
ATOM	1023 CA	ASP	133	21.883	52.442	29.525	1.00	45.36
ATOM	1024 C	ASP	133	23.224	51.861	30.049	1.00	50.61
ATOM	1025 O	ASP	133	24.299	52.243	29.572	1.00	52.14
ATOM	1026 CB	ASP	133	22.063	53.946	29.332	1.00	50.45
ATOM	1027 CG	ASP	133	22.109	54.642	30.670	1.00	87.10
ATOM	1028 OD1	ASP	133	21.408	54.314	31.624	1.00	91.27
ATOM	1029 OD2	ASP	133	23.047	55.552	30.739	1.00	100.00
ATOM	1030 N	GLY	134	23.159	50.970	31.053	1.00	37.06
ATOM	1031 CA	GLY	134	24.349	50.376	31.639	1.00	30.22
ATOM	1032 C	GLY	134	24.845	49.228	30.803	1.00	23.10
ATOM	1033 O	GLY	134	24.360	48.990	29.685	1.00	19.23
ATOM	1034 N	ASN	135	25.807	48.486	31.341	1.00	18.66
ATOM	1035 CA	ASN	135	26.339	47.370	30.563	1.00	18.03
ATOM	1036 C	ASN	135	25.372	46.199	30.406	1.00	15.75
ATOM	1037 O	ASN	135	25.485	45.430	29.461	1.00	16.03
ATOM	1038 CB	ASN	135	27.665	46.883	31.139	1.00	19.27
ATOM	1039 CG	ASN	135	28.743	47.943	31.108	1.00	20.99
ATOM	1040 OD1	ASN	135	28.969	48.595	30.078	1.00	25.69
ATOM	1041 ND2	ASN	135	29.423	48.095	32.239	1.00	22.57
ATOM	1042 N	ILE	136	24.444	46.052	31.362	1.00	18.14
ATOM	1043 CA	ILE	136	23.494	44.924	31.368	1.00	19.78
ATOM	1044 C	ILE	136	22.331	45.086	30.384	1.00	23.76
ATOM	1045 O	ILE	136	22.178	44.313	29.395	1.00	22.53
ATOM	1046 CB	ILE	136	23.078	44.500	32.804	1.00	21.24
ATOM	1047 CG1	ILE	136	24.230	43.728	33.423	1.00	28.44
ATOM	1048 CG2	ILE	136	21.899	43.543	32.770	1.00	22.77
ATOM	1049 CD1	ILE	136	25.346	44.596	33.935	1.00	12.39
ATOM	1050 N	LEU	137	21.543	46.117	30.640	1.00	18.21
ATOM	1051 CA	LEU	137	20.394	46.415	29.815	1.00	23.30
ATOM	1052 C	LEU	137	20.828	46.875	28.470	1.00	27.26
ATOM	1053 O	LEU	137	20.181	46.619	27.488	1.00	27.00
ATOM	1054 CB	LEU	137	19.442	47.430	30.490	1.00	21.74
ATOM	1055 CG	LEU	137	18.828	46.852	31.762	1.00	22.56
ATOM	1056 CD1	LEU	137	17.856	47.837	32.415	1.00	22.27
ATOM	1057 CD2	LEU	137	18.118	45.554	31.424	1.00	37.52
ATOM	1058 N	GLY	138	21.979	47.527	28.432	1.00	22.14
ATOM	1059 CA	GLY	138	22.510	48.033	27.187	1.00	20.03
ATOM	1060 C	GLY	138	23.157	46.959	26.368	1.00	20.16

FIG. 5Z

ATOM	1061 O	GLY	138	23.600	47.202	25.264	1.00 22.44
ATOM	1062 N	HIS	139	23.246	45.756	26.903	1.00 18.27
ATOM	1063 CA	HIS	139	23.859	44.655	26.148	1.00 20.24
ATOM	1064 C	HIS	139	25.301	44.929	25.616	1.00 20.13
ATOM	1065 O	HIS	139	25.605	44.745	24.439	1.00 17.97
ATOM	1066 CB	HIS	139	22.931	44.207	25.018	1.00 22.20
ATOM	1067 CG	HIS	139	21.708	43.551	25.550	1.00 25.52
ATOM	1068 ND1	HIS	139	21.666	42.182	25.785	1.00 25.67
ATOM	1069 CD2	HIS	139	20.525	44.092	25.927	1.00 28.09
ATOM	1070 CE1	HIS	139	20.474	41.918	26.275	1.00 27.50
ATOM	1071 NE2	HIS	139	19.766	43.044	26.382	1.00 29.53
ATOM	1072 N	LYS	140	26.187	45.311	26.525	1.00 23.51
ATOM	1073 CA	LYS	140	27.569	45.638	26.197	1.00 25.82
ATOM	1074 C	LYS	140	28.600	44.537	26.560	1.00 26.28
ATOM	1075 O	LYS	140	29.824	44.730	26.391	1.00 22.29
ATOM	1076 CB	LYS	140	27.977	46.937	26.911	1.00 27.56
ATOM	1077 CG	LYS	140	27.269	48.217	26.445	1.00 31.19
ATOM	1078 CD	LYS	140	27.234	49.254	27.582	1.00 51.32
ATOM	1079 CE	LYS	140	26.924	50.696	27.169	1.00 47.92
ATOM	1080 NZ	LYS	140	27.112	51.663	28.284	1.00 73.76
ATOM	1081 N	LEU	141	28.116	43.403	27.115	1.00 19.33
ATOM	1082 CA	LEU	141	28.987	42.296	27.559	1.00 14.32
ATOM	1083 C	LEU	141	29.366	41.401	26.427	1.00 20.75
ATOM	1084 O	LEU	141	28.526	41.087	25.620	1.00 19.01
ATOM	1085 CB	LEU	141	28.313	41.488	28.676	1.00 12.53
ATOM	1086 CG	LEU	141	27.979	42.352	29.875	1.00 17.54
ATOM	1087 CD1	LEU	141	27.700	41.469	31.070	1.00 24.81
ATOM	1088 CD2	LEU	141	29.116	43.310	30.182	1.00 27.50
ATOM	1089 N	GLU	142	30.644	40.987	26.346	1.00 14.76
ATOM	1090 CA	GLU	142	31.040	40.059	25.311	1.00 13.43
ATOM	1091 C	GLU	142	30.462	38.691	25.641	1.00 15.69
ATOM	1092 O	GLU	142	30.175	38.393	26.787	1.00 16.43
ATOM	1093 CB	GLU	142	32.558	39.866	25.204	1.00 14.73
ATOM	1094 CG	GLU	142	33.290	41.077	24.624	1.00 29.30
ATOM	1095 CD	GLU	142	34.787	41.003	24.825	1.00 56.32
ATOM	1096 OE1	GLU	142	35.340	40.098	25.420	1.00 31.70
ATOM	1097 OE2	GLU	142	35.430	42.015	24.321	1.00 34.10
ATOM	1098 N	TYR	143	30.365	37.873	24.632	1.00 16.30
ATOM	1099 CA	TYR	143	29.837	36.542	24.764	1.00 20.04
ATOM	1100 C	TYR	143	30.925	35.559	25.049	1.00 12.46
ATOM	1101 O	TYR	143	31.327	34.792	24.193	1.00 16.99

FIG. 5AA

ATOM	1102 CB TYR	143	29.035	36.118	23.498	1.00	20.96
ATOM	1103 CG TYR	143	28.187	34.857	23.674	1.00	16.12
ATOM	1104 CD1TYR	143	27.040	34.859	24.472	1.00	18.24
ATOM	1105 CD2TYR	143	28.512	33.684	22.986	1.00	12.87
ATOM	1106 CE1 TYR	143	26.257	33.708	24.615	1.00	17.91
ATOM	1107 CE2 TYR	143	27.735	32.530	23.104	1.00	16.58
ATOM	1108 CZ TYR	143	26.603	32.551	23.914	1.00	17.35
ATOM	1109 OH TYR	143	25.861	31.432	24.035	1.00	23.40
ATOM	1110 N ASN	144	31.392	35.597	26.251	1.00	12.40
ATOM	1111 CA ASN	144	32.428	34.703	26.689	1.00	12.05
ATOM	1112 C ASN	144	32.433	34.675	28.193	1.00	15.75
ATOM	1113 O ASN	144	31.637	35.369	28.837	1.00	14.58
ATOM	1114 CB ASN	144	33.823	35.038	26.068	1.00	18.45
ATOM	1115 CG ASN	144	34.310	36.445	26.374	1.00	18.98
ATOM	1116 OD1ASN	144	34.150	36.951	27.488	1.00	20.34
ATOM	1117 ND2ASN	144	34.891	37.085	25.382	1.00	23.02
ATOM	1118 N TYR	145	33.311	33.876	28.773	1.00	12.16
ATOM	1119 CA TYR	145	33.343	33.765	30.195	1.00	10.63
ATOM	1120 C TYR	145	34.765	33.458	30.730	1.00	14.58
ATOM	1121 O TYR	145	35.510	32.751	30.090	1.00	18.83
ATOM	1122 CB TYR	145	32.404	32.627	30.571	1.00	9.76
ATOM	1123 CG TYR	145	31.698	32.916	31.826	1.00	11.86
ATOM	1124 CD1TYR	145	30.515	33.658	31.808	1.00	9.04
ATOM	1125 CD2TYR	145	32.188	32.419	33.030	1.00	10.07
ATOM	1126 CE1 TYR	145	29.860	33.948	32.999	1.00	8.36
ATOM	1127 CE2 TYR	145	31.544	32.707	34.235	1.00	15.32
ATOM	1128 CZ TYR	145	30.375	33.469	34.206	1.00	11.69
ATOM	1129 OH TYR	145	29.730	33.735	35.376	1.00	15.23
ATOM	1130 N ASN	146	35.086	33.931	31.933	1.00	15.36
ATOM	1131 CA ASN	146	36.415	33.737	32.560	1.00	17.00
ATOM	1132 C ASN	146	36.426	32.618	33.589	1.00	19.68
ATOM	1133 O ASN	146	35.395	32.043	33.848	1.00	14.71
ATOM	1134 CB ASN	146	36.844	35.062	33.235	1.00	11.89
ATOM	1135 CG ASN	146	37.013	36.147	32.215	1.00	35.45
ATOM	1136 OD1ASN	146	37.533	35.890	31.105	1.00	31.63
ATOM	1137 ND2ASN	146	36.547	37.349	32.553	1.00	19.74
ATOM	1138 N SER	147	37.630	32.338	34.201	1.00	12.09
ATOM	1139 CA SER	147	37.804	31.320	35.266	1.00	8.55
ATOM	1140 C SER	147	37.769	31.999	36.575	1.00	11.70
ATOM	1141 O SER	147	38.219	33.125	36.671	1.00	16.56
ATOM	1142 CB SER	147	39.148	30.540	35.129	1.00	9.87
ATOM	1143 OG SER	147	39.212	29.980	33.828	1.00	33.20

FIG. 5AB

ATOM	1144 N	HIS	148	37.195	31.365	37.583	1.00	5.53
ATOM	1145 CA	HIS	148	37.090	31.998	38.850	1.00	8.06
ATOM	1146 C	HIS	148	37.346	31.038	39.949	1.00	11.30
ATOM	1147 O	HIS	148	37.328	29.844	39.754	1.00	16.87
ATOM	1148 CB	HIS	148	35.648	32.608	39.067	1.00	11.29
ATOM	1149 CG	HIS	148	35.215	33.554	37.972	1.00	10.84
ATOM	1150 ND1	HIS	148	34.548	33.121	36.836	1.00	12.77
ATOM	1151 CD2	HIS	148	35.403	34.887	37.851	1.00	8.82
ATOM	1152 CE1	HIS	148	34.389	34.178	36.060	1.00	8.84
ATOM	1153 NE2	HIS	148	34.882	35.242	36.647	1.00	8.82
ATOM	1154 N	ASN	149	37.534	31.579	41.125	1.00	10.80
ATOM	1155 CA	ASN	149	37.626	30.805	42.345	1.00	13.35
ATOM	1156 C	ASN	149	36.409	31.157	43.205	1.00	14.47
ATOM	1157 O	ASN	149	36.099	32.320	43.387	1.00	18.17
ATOM	1158 CB	ASN	149	38.890	31.093	43.184	1.00	12.67
ATOM	1159 CG	ASN	149	40.148	30.822	42.424	1.00	20.21
ATOM	1160 OD1	ASN	149	40.993	31.713	42.281	1.00	56.34
ATOM	1161 ND2	ASN	149	40.210	29.641	41.818	1.00	16.44
ATOM	1162 N	VAL	150	35.773	30.144	43.741	1.00	14.65
ATOM	1163 CA	VAL	150	34.588	30.262	44.552	1.00	12.92
ATOM	1164 C	VAL	150	34.910	29.806	45.943	1.00	16.30
ATOM	1165 O	VAL	150	35.257	28.665	46.147	1.00	17.83
ATOM	1166 CB	VAL	150	33.482	29.382	43.914	1.00	15.22
ATOM	1167 CG1	VAL	150	32.252	29.297	44.765	1.00	14.09
ATOM	1168 CG2	VAL	150	33.172	29.791	42.464	1.00	10.94
ATOM	1169 N	TYR	151	34.796	30.716	46.900	1.00	17.64
ATOM	1170 CA	TYR	151	35.139	30.440	48.275	1.00	18.31
ATOM	1171 C	TYR	151	34.003	29.917	49.117	1.00	24.35
ATOM	1172 O	TYR	151	32.963	30.536	49.239	1.00	20.83
ATOM	1173 CB	TYR	151	35.793	31.681	48.920	1.00	20.15
ATOM	1174 CG	TYR	151	37.025	32.033	48.141	1.00	25.86
ATOM	1175 CD1	TYR	151	37.003	32.989	47.127	1.00	26.00
ATOM	1176 CD2	TYR	151	38.200	31.315	48.355	1.00	28.66
ATOM	1177 CE1	TYR	151	38.151	33.234	46.369	1.00	33.73
ATOM	1178 CE2	TYR	151	39.360	31.550	47.619	1.00	29.01
ATOM	1179 CZ	TYR	151	39.325	32.512	46.618	1.00	29.55
ATOM	1180 OH	TYR	151	40.449	32.737	45.877	1.00	38.69
ATOM	1181 N	ILE	152	34.250	28.791	49.753	1.00	17.71
ATOM	1182 CA	ILE	152	33.255	28.159	50.572	1.00	14.12
ATOM	1183 C	ILE	152	33.619	28.056	52.000	1.00	18.51
ATOM	1184 O	ILE	152	34.728	27.703	52.336	1.00	22.05

FIG. 5AC

ATOM	1185 CB	ILE	152	32.979	26.776	50.060	1.00	16.66
ATOM	1186 CG1	ILE	152	32.431	26.875	48.638	1.00	11.30
ATOM	1187 CG2	ILE	152	32.017	26.078	51.021	1.00	17.96
ATOM	1188 CD1	ILE	152	32.377	25.559	47.949	1.00	13.48
ATOM	1189 N	MSE	153	32.623	28.278	52.841	1.00	17.41
ATOM	1190 AC	MSE	153	32.789	28.162	54.269	1.00	22.61
ATOM	1191 C	MSE	153	31.534	27.648	54.916	1.00	27.31
ATOM	1192 O	MSE	153	30.433	27.831	54.396	1.00	20.50
ATOM	1193 CB	MSE	153	33.145	29.490	54.855	1.00	19.11
ATOM	1194 CG	MSE	153	34.010	30.302	53.957	1.00	100.00
ATOM	1195 SE	MSE	153	34.060	32.117	54.524	1.00	100.00
ATOM	1196 CE	MSE	153	33.463	31.798	56.330	1.00	30.27
ATOM	1197 N	ALA	154	31.733	26.983	56.053	1.00	22.29
ATOM	1198 CA	ALA	154	30.669	26.389	56.796	1.00	22.66
ATOM	1199 C	ALA	154	29.820	27.401	57.552	1.00	29.00
ATOM	1200 O	ALA	154	30.274	28.457	57.960	1.00	27.02
ATOM	1201 CB	ALA	154	31.224	25.336	57.744	1.00	19.78
ATOM	1202 N	ASP	155	28.566	27.063	57.726	1.00	29.48
ATOM	1203 CA	ASP	155	27.669	27.887	58.484	1.00	32.18
ATOM	1204 C	ASP	155	26.976	27.019	59.511	1.00	44.51
ATOM	1205 O	ASP	155	25.898	26.492	59.274	1.00	39.56
ATOM	1206 CB	ASP	155	26.659	28.617	57.597	1.00	31.70
ATOM	1207 CG	ASP	155	26.140	29.851	58.247	1.00	49.89
ATOM	1208 OD1	ASP	155	26.595	30.297	59.277	1.00	46.67
ATOM	1209 OD2	ASP	155	25.187	30.422	57.565	1.00	76.07
ATOM	1210 N	LYS	156	27.646	26.816	60.629	1.00	46.37
ATOM	1211 CA	LYS	156	27.116	25.954	61.654	1.00	53.23
ATOM	1212 C	LYS	156	25.750	26.369	62.224	1.00	65.62
ATOM	1213 O	LYS	156	25.012	25.520	62.703	1.00	65.54
ATOM	1214 CB	LYS	156	28.147	25.612	62.725	1.00	59.51
ATOM	1215 N	GLN	157	25.398	27.655	62.138	1.00	68.32
ATOM	1216 CA	GLN	157	24.119	28.135	62.670	1.00	73.00
ATOM	1217 C	GLN	157	22.891	27.767	61.817	1.00	87.53
ATOM	1218 O	GLN	157	21.778	27.547	62.325	1.00	96.16
ATOM	1219 N	LYS	158	23.095	27.725	60.506	1.00	72.49
ATOM	1220 CA	LYS	158	22.040	27.386	59.593	1.00	66.19
ATOM	1221 C	LYS	158	22.235	25.985	59.040	1.00	58.21
ATOM	1222 O	LYS	158	21.447	25.524	58.226	1.00	59.85
ATOM	1223 N	ASN	159	23.303	25.294	59.502	1.00	40.00
ATOM	1224 CA	ASN	159	23.582	23.944	59.012	1.00	36.67
ATOM	1225 C	ASN	159	23.755	24.002	57.500	1.00	34.11

FIG. 5AD

ATOM	1226 CO	ASN	159	23.223	23.167	56.754	1.00	31.69
ATOM	1227 CB	ASN	159	22.431	22.952	59.367	1.00	46.42
ATOM	1228 CG	ASN	159	22.842	21.485	59.428	1.00	80.46
ATOM	1229 OD1	ASN	159	23.850	21.121	60.054	1.00	100.00
ATOM	1230 ND2	ASN	159	22.003	20.620	58.854	1.00	58.09
ATOM	1231 N	GLY	160	24.474	25.044	57.062	1.00	22.34
ATOM	1232 CA	GLY	160	24.686	25.247	55.663	1.00	17.58
ATOM	1233 C	GLY	160	26.055	25.791	55.433	1.00	26.75
ATOM	1234 O	GLY	160	26.960	25.664	56.271	1.00	25.57
ATOM	1235 N	ILE	161	26.200	26.395	54.277	1.00	23.28
ATOM	1236 CA	ILE	161	27.442	26.975	53.909	1.00	16.45
ATOM	1237 C	ILE	161	27.200	28.354	53.395	1.00	15.77
ATOM	1238 O	ILE	161	26.118	28.680	52.962	1.00	15.95
ATOM	1239 CB	ILE	161	28.129	26.117	52.864	1.00	19.27
ATOM	1240 CG1	ILE	161	27.237	26.016	51.619	1.00	18.53
ATOM	1241 CG2	ILE	161	28.351	24.735	53.445	1.00	21.96
ATOM	1242 CD1	ILE	161	28.009	25.614	50.350	1.00	14.44
ATOM	1243 N	LYS	162	28.226	29.169	53.471	1.00	17.86
ATOM	1244 CA	LYS	162	28.187	30.508	52.948	1.00	14.42
ATOM	1245 C	LYS	162	29.216	30.524	51.857	1.00	17.73
ATOM	1246 O	LYS	162	30.249	29.875	51.991	1.00	19.16
ATOM	1247 CB	LYS	162	28.480	31.540	54.055	1.00	18.15
ATOM	1248 CG	LYS	162	27.221	31.963	54.796	1.00	42.08
ATOM	1249 CD	LYS	162	27.493	32.787	56.039	1.00	70.42
ATOM	1250 N	VAL	163	28.911	31.176	50.759	1.00	13.74
ATOM	1251 CA	VAL	163	29.798	31.201	49.629	1.00	11.95
ATOM	1252 C	VAL	163	29.928	32.610	49.103	1.00	19.30
ATOM	1253 O	VAL	163	28.944	33.318	48.983	1.00	19.84
ATOM	1254 CB	VAL	163	29.249	30.268	48.532	1.00	15.89
ATOM	1255 CG1	VAL	163	30.105	30.277	47.261	1.00	12.09
ATOM	1256 CG2	VAL	163	29.029	28.852	49.077	1.00	15.86
ATOM	1257 N	ASN	164	31.146	32.999	48.733	1.00	14.03
ATOM	1258 CA	ASN	164	31.382	34.310	48.195	1.00	15.58
ATOM	1259 C	ASN	164	32.396	34.271	47.050	1.00	20.08
ATOM	1260 O	ASN	164	33.268	33.386	46.988	1.00	23.49
ATOM	1261 CB	ASN	164	31.732	35.325	49.308	1.00	20.52
ATOM	1262 CG	ASN	164	33.196	35.697	49.330	1.00	89.21
ATOM	1263 OD1	ASN	164	34.020	34.987	49.929	1.00	100.00
ATOM	1264 ND2	ASN	164	33.515	36.831	48.700	1.00	91.46
ATOM	1265 N	PHE	165	32.244	35.207	46.109	1.00	17.37
ATOM	1266 CA	PHE	165	33.133	35.301	44.953	1.00	10.86

FIG. 5AE

ATOM	1267 C	PHE	165	32.751	36.445	44.071	1.00 15.53
ATOM	1268 O	PHE	165	31.686	37.020	44.251	1.00 17.16
ATOM	1269 CB	PHE	165	33.207	33.960	44.187	1.00 12.86
ATOM	1270 CG	PHE	165	31.862	33.486	43.622	1.00 14.35
ATOM	1271 CD1	PHE	165	31.510	33.749	42.293	1.00 14.61
ATOM	1272 CD2	PHE	165	30.978	32.757	44.413	1.00 13.56
ATOM	1273 CE1	PHE	165	30.300	33.297	41.759	1.00 22.67
ATOM	1274 CE2	PHE	165	29.774	32.282	43.893	1.00 15.78
ATOM	1275 CZ	PHE	165	29.426	32.572	42.573	1.00 16.20
ATOM	1276 N	LYS	166	33.641	36.799	43.132	1.00 10.79
ATOM	1277 CA	LYS	166	33.417	37.864	42.162	1.00 10.74
ATOM	1278 C	LYS	166	33.603	37.344	40.774	1.00 15.95
ATOM	1279 O	LYS	166	34.602	36.727	40.470	1.00 22.80
ATOM	1280 CB	LYS	166	34.387	39.055	42.249	1.00 16.61
ATOM	1281 CG	LYS	166	34.573	39.688	43.573	1.00 18.11
ATOM	1282 CD	LYS	166	35.540	40.875	43.454	1.00 32.56
ATOM	1283 CE	LYS	166	35.272	41.966	44.476	1.00 48.19
ATOM	1284 NZ	LYS	166	34.823	41.435	45.782	1.00 85.81
ATOM	1285 N	ILE	167	32.703	37.704	39.911	1.00 9.75
ATOM	1286 CA	ILE	167	32.768	37.340	38.558	1.00 9.35
ATOM	1287 C	ILE	167	33.203	38.542	37.823	1.00 14.36
ATOM	1288 O	ILE	167	32.811	39.640	38.170	1.00 16.22
ATOM	1289 CB	ILE	167	31.379	36.929	38.005	1.00 13.16
ATOM	1290 CG1	ILE	167	30.909	35.624	38.669	1.00 13.02
ATOM	1291 CG2	ILE	167	31.423	36.786	36.472	1.00 7.91
ATOM	1292 CD1	ILE	167	31.773	34.415	38.344	1.00 19.57
ATOM	1293 N	ARG	168	34.005	38.299	36.815	1.00 12.19
ATOM	1294 CA	ARG	168	34.500	39.308	35.945	1.00 15.07
ATOM	1295 C	ARG	168	33.948	39.122	34.528	1.00 16.64
ATOM	1296 O	ARG	168	34.278	38.156	33.836	1.00 17.70
ATOM	1297 CB	ARG	168	36.024	39.287	35.944	1.00 16.54
ATOM	1298 CG	ARG	168	36.580	39.632	37.321	1.00 25.54
ATOM	1299 CD	ARG	168	37.894	38.910	37.601	1.00 63.52
ATOM	1300 NE	ARG	168	38.380	38.191	36.416	1.00 73.52
ATOM	1301 CZ	ARG	168	38.764	36.926	36.416	1.00 67.92
ATOM	1302 NH1	ARG	168	38.795	36.192	37.527	1.00 57.44
ATOM	1303 NH2	ARG	168	39.192	36.375	35.271	1.00 59.15
ATOM	1304 N	HIS	169	33.090	40.064	34.098	1.00 14.88
ATOM	1305 CA	HIS	169	32.505	40.025	32.758	1.00 13.24
ATOM	1306 C	HIS	169	33.214	41.001	31.839	1.00 12.64
ATOM	1307 O	HIS	169	33.306	42.203	32.121	1.00 14.99

FIG. 5AF

ATOM	1308 CB HIS	169	30.970	40.374	32.760	1.00 10.46
ATOM	1309 CG HIS	169	30.097	39.474	33.573	1.00 6.54
ATOM	1310 ND1HIS	169	29.724	38.246	33.111	1.00 12.63
ATOM	1311 CD2HIS	169	29.474	39.695	34.764	1.00 10.21
ATOM	1312 CE1 HIS	169	28.892	37.718	34.031	1.00 10.53
ATOM	1313 NE2HIS	169	28.734	38.566	35.063	1.00 11.84
ATOM	1314 N ASN	170	33.691	40.513	30.737	1.00 10.66
ATOM	1315 CA ASN	170	34.349	41.368	29.812	1.00 15.87
ATOM	1316 C ASN	170	33.356	42.224	29.067	1.00 25.06
ATOM	1317 O ASN	170	32.386	41.701	28.537	1.00 16.60
ATOM	1318 CB ASN	170	35.110	40.550	28.755	1.00 19.60
ATOM	1319 CG ASN	170	36.245	39.717	29.312	1.00 18.70
ATOM	1320 OD1ASN	170	36.702	38.752	28.684	1.00 48.29
ATOM	1321 ND2ASN	170	36.695	40.073	30.480	1.00 19.13
ATOM	1322 N ILE	171	33.662	43.527	28.947	1.00 18.75
ATOM	1323 CA ILE	171	32.848	44.460	28.168	1.00 16.74
ATOM	1324 C ILE	171	33.459	44.638	26.791	1.00 19.51
ATOM	1325 O ILE	171	34.643	44.596	26.642	1.00 21.06
ATOM	1326 CB ILE	171	32.713	45.804	28.842	1.00 20.46
ATOM	1327 CG1ILE	171	32.089	45.617	30.193	1.00 24.79
ATOM	1328 CG2ILE	171	31.852	46.727	27.997	1.00 19.03
ATOM	1329 CD1ILE	171	32.630	46.599	31.229	1.00 41.65
ATOM	1330 N GLU	172	32.632	44.818	25.804	1.00 16.54
ATOM	1331 CA GLU	172	33.034	44.933	24.420	1.00 17.00
ATOM	1332 C GLU	172	34.110	45.967	24.147	1.00 26.80
ATOM	1333 O GLU	172	34.776	45.898	23.125	1.00 29.20
ATOM	1334 CB GLU	172	31.813	45.165	23.509	1.00 22.46
ATOM	1335 CG GLU	172	31.122	46.531	23.786	1.00 58.53
ATOM	1336 CD GLU	172	29.871	46.783	22.933	1.00 100.00
ATOM	1337 OE1 GLU	172	29.415	45.970	22.156	1.00 100.00
ATOM	1338 OE2 GLU	172	29.370	47.982	23.149	1.00 100.00
ATOM	1339 N ASP	173	34.277	46.934	25.034	1.00 24.41
ATOM	1340 CA ASP	173	35.292	47.978	24.852	1.00 25.03
ATOM	1341 C ASP	173	36.651	47.624	25.455	1.00 33.40
ATOM	1342 O ASP	173	37.561	48.451	25.518	1.00 30.42
ATOM	1343 CB ASP	173	34.822	49.319	25.401	1.00 23.30
ATOM	1344 CG ASP	173	34.743	49.358	26.912	1.00 32.47
ATOM	1345 OD1ASP	173	34.406	50.355	27.513	1.00 37.58
ATOM	1346 OD2ASP	173	34.949	48.196	27.504	1.00 49.22
ATOM	1347 N GLY	174	36.766	46.410	25.956	1.00 23.87
ATOM	1348 CA GLY	174	38.019	45.994	26.537	1.00 21.30

FIG. 5AG

ATOM	1349 C	GLY	174	38.012	46.090	28.044	1.00	19.99
ATOM	1350 O	GLY	174	38.927	45.585	28.709	1.00	20.45
ATOM	1351 N	SER	175	36.972	46.767	28.598	1.00	13.88
ATOM	1352 CA	SER	175	36.898	46.931	30.034	1.00	8.70
ATOM	1353 C	SER	175	36.296	45.728	30.765	1.00	17.30
ATOM	1354 O	SER	175	36.136	44.655	30.175	1.00	18.77
ATOM	1355 CB	SER	175	36.288	48.235	30.450	1.00	14.07
ATOM	1356 OG	SER	175	36.360	48.316	31.865	1.00	24.79
ATOM	1357 N	VAL	176	35.963	45.912	32.051	1.00	13.74
ATOM	1358 CA	VAL	176	35.415	44.826	32.864	1.00	16.46
ATOM	1359 C	VAL	176	34.191	45.204	33.703	1.00	22.46
ATOM	1360 O	VAL	176	34.159	46.254	34.334	1.00	21.31
ATOM	1361 CB	VAL	176	36.477	44.285	33.818	1.00	24.43
ATOM	1362 CG1	VAL	176	35.847	43.344	34.827	1.00	27.45
ATOM	1363 CG2	VAL	176	37.532	43.536	33.035	1.00	25.65
ATOM	1364 N	GLN	177	33.234	44.269	33.787	1.00	15.47
ATOM	1365 CA	GLN	177	32.048	44.430	34.647	1.00	15.40
ATOM	1366 C	GLN	177	32.102	43.457	35.813	1.00	10.60
ATOM	1367 O	GLN	177	32.027	42.243	35.634	1.00	13.65
ATOM	1368 CB	GLN	177	30.709	44.283	33.872	1.00	15.57
ATOM	1369 CG	GLN	177	29.468	44.294	34.828	1.00	19.13
ATOM	1370 CD	GLN	177	29.108	45.678	35.361	1.00	14.91
ATOM	1371 OE1	GLN	177	28.759	46.588	34.574	1.00	20.17
ATOM	1372 NE2	GLN	177	29.128	45.821	36.690	1.00	17.28
ATOM	1373 N	LEU	178	32.227	43.993	37.018	1.00	8.17
ATOM	1374 CA	LEU	178	32.313	43.180	38.181	1.00	16.66
ATOM	1375 C	LEU	178	30.954	42.786	38.712	1.00	20.93
ATOM	1376 O	LEU	178	30.033	43.608	38.753	1.00	14.66
ATOM	1377 CB	LEU	178	33.089	43.896	39.293	1.00	20.63
ATOM	1378 CG	LEU	178	34.286	43.110	39.815	1.00	39.28
ATOM	1379 CD1	LEU	178	33.831	42.087	40.852	1.00	45.14
ATOM	1380 CD2	LEU	178	35.018	42.426	38.648	1.00	39.52
ATOM	1381 N	ALA	179	30.869	41.550	39.171	1.00	16.72
ATOM	1382 CA	ALA	179	29.652	41.033	39.754	1.00	15.55
ATOM	1383 C	ALA	179	29.932	40.277	41.040	1.00	15.70
ATOM	1384 O	ALA	179	30.337	39.119	41.028	1.00	15.91
ATOM	1385 CB	ALA	179	28.853	40.197	38.731	1.00	14.08
ATOM	1386 N	ASP	180	29.694	40.946	42.155	1.00	8.88
ATOM	1387 CA	ASP	180	29.897	40.407	43.480	1.00	7.18
ATOM	1388 C	ASP	180	28.802	39.460	43.891	1.00	17.07
ATOM	1389 O	ASP	180	27.651	39.844	43.987	1.00	18.22

FIG. 5AH

ATOM	1390 CB ASP	180	29.934	41.509	44.509	1.00	13.06
ATOM	1391 CG ASP	180	31.285	41.902	44.935	1.00	46.28
ATOM	1392 OD1ASP	180	31.981	41.206	45.655	1.00	60.46
ATOM	1393 OD2ASP	180	31.574	43.121	44.560	1.00	46.61
ATOM	1394 N HIS	181	29.173	38.242	44.197	1.00	14.51
ATOM	1395 CA HIS	181	28.213	37.223	44.575	1.00	10.49
ATOM	1396 C HIS	181	28.218	36.897	46.049	1.00	14.28
ATOM	1397 O HIS	181	29.255	36.580	46.607	1.00	17.40
ATOM	1398 CB HIS	181	28.450	35.915	43.769	1.00	9.89
ATOM	1399 CG HIS	181	28.077	35.972	42.328	1.00	10.38
ATOM	1400 ND1HIS	181	28.606	36.926	41.455	1.00	12.24
ATOM	1401 CD2HIS	181	27.279	35.146	41.606	1.00	10.42
ATOM	1402 CE1 HIS	181	28.093	36.678	40.269	1.00	9.97
ATOM	1403 NE2 HIS	181	27.314	35.594	40.316	1.00	9.38
ATOM	1404 N TYR	182	27.029	36.897	46.668	1.00	10.40
ATOM	1405 CA TYR	182	26.848	36.518	48.062	1.00	13.86
ATOM	1406 C TYR	182	25.871	35.393	48.089	1.00	20.61
ATOM	1407 O TYR	182	24.819	35.520	47.532	1.00	16.35
ATOM	1408 CB TYR	182	26.359	37.664	48.934	1.00	21.12
ATOM	1409 CG TYR	182	27.421	38.693	49.062	1.00	34.16
ATOM	1410 CD1TYR	182	27.521	39.715	48.120	1.00	46.06
ATOM	1411 CD2TYR	182	28.389	38.616	50.064	1.00	38.56
ATOM	1412 CE1 TYR	182	28.532	40.674	48.197	1.00	57.53
ATOM	1413 CE2 TYR	182	29.418	39.559	50.147	1.00	40.76
ATOM	1414 CZ TYR	182	29.480	40.594	49.216	1.00	54.61
ATOM	1415 OH TYR	182	30.461	41.534	49.308	1.00	61.92
ATOM	1416 N GLN	183	26.246	34.277	48.686	1.00	17.63
ATOM	1417 CA GLN	183	25.410	33.104	48.583	1.00	16.37
ATOM	1418 C GLN	183	25.289	32.311	49.863	1.00	21.39
ATOM	1419 O GLN	183	26.260	32.174	50.623	1.00	19.86
ATOM	1420 CB GLN	183	25.984	32.219	47.422	1.00	13.33
ATOM	1421 CG GLN	183	25.651	30.688	47.457	1.00	17.38
ATOM	1422 CD GLN	183	26.411	29.884	46.389	1.00	17.27
ATOM	1423 OE1 GLN	183	26.975	30.454	45.456	1.00	13.80
ATOM	1424 NE2 GLN	183	26.361	28.553	46.473	1.00	13.94
ATOM	1425 N GLN	184	24.080	31.739	50.055	1.00	19.74
ATOM	1426 CA GLN	184	23.760	30.829	51.168	1.00	16.55
ATOM	1427 C GLN	184	23.033	29.582	50.658	1.00	13.60
ATOM	1428 O GLN	184	22.219	29.640	49.747	1.00	18.01
ATOM	1429 CB GLN	184	22.949	31.444	52.330	1.00	20.11
ATOM	1430 CG GLN	184	23.364	32.855	52.768	1.00	74.84

FIG. 5A1

ATOM	1431	CD	GLN	184	22.312	33.517	53.657	1.00	100.00
ATOM	1432	OE1	GLN	184	21.159	33.054	53.752	1.00	97.99
ATOM	1433	NE2	GLN	184	22.689	34.625	54.286	1.00	100.00
ATOM	1434	N	ASN	185	23.418	28.446	51.207	1.00	14.76
ATOM	1435	CA	ASN	185	22.831	27.155	50.887	1.00	13.86
ATOM	1436	C	ASN	185	22.421	26.463	52.166	1.00	16.06
ATOM	1437	O	ASN	185	23.176	26.402	53.172	1.00	17.39
ATOM	1438	CB	ASN	185	23.761	26.212	50.119	1.00	15.20
ATOM	1439	CG	ASN	185	24.110	26.696	48.748	1.00	12.75
ATOM	1440	OD1	ASN	185	24.704	27.758	48.592	1.00	22.56
ATOM	1441	ND2	ASN	185	23.830	25.868	47.763	1.00	17.70
ATOM	1442	N	THR	186	21.227	25.941	52.139	1.00	18.01
ATOM	1443	CA	THR	186	20.707	25.227	53.288	1.00	17.40
ATOM	1444	C	THR	186	19.976	24.010	52.824	1.00	23.63
ATOM	1445	O	THR	186	19.389	23.991	51.730	1.00	24.57
ATOM	1446	CB	THR	186	19.856	26.100	54.206	1.00	28.82
ATOM	1447	OG1	THR	186	18.874	26.752	53.446	1.00	35.65
ATOM	1448	CG2	THR	186	20.753	27.121	54.903	1.00	28.86
ATOM	1449	N	PRO	187	20.101	22.951	53.620	1.00	22.40
ATOM	1450	CA	PRO	187	19.504	21.683	53.269	1.00	20.28
ATOM	1451	C	PRO	187	17.988	21.757	53.288	1.00	22.41
ATOM	1452	O	PRO	187	17.390	22.518	54.071	1.00	25.07
ATOM	1453	CB	PRO	187	19.977	20.682	54.337	1.00	19.79
ATOM	1454	CG	PRO	187	20.840	21.449	55.338	1.00	26.98
ATOM	1455	CD	PRO	187	20.786	22.918	54.949	1.00	22.04
ATOM	1456	N	ILE	188	17.382	20.957	52.453	1.00	18.77
ATOM	1457	CA	ILE	188	15.907	20.855	52.407	1.00	20.12
ATOM	1458	C	ILE	188	15.470	19.766	53.389	1.00	31.58
ATOM	1459	O	ILE	188	14.596	19.966	54.202	1.00	38.58
ATOM	1460	CB	ILE	188	15.385	20.574	50.991	1.00	21.52
ATOM	1461	CG1	ILE	188	15.555	21.775	50.102	1.00	16.10
ATOM	1462	CG2	ILE	188	13.916	20.141	50.981	1.00	28.85
ATOM	1463	CD1	ILE	188	15.139	21.471	48.660	1.00	15.31
ATOM	1464	N	GLY	189	16.142	18.618	53.352	1.00	32.39
ATOM	1465	CA	GLY	189	15.833	17.531	54.283	1.00	32.94
ATOM	1466	C	GLY	189	16.339	17.817	55.702	1.00	40.20
ATOM	1467	O	GLY	189	17.016	18.810	55.967	1.00	35.57
ATOM	1468	N	ASP	190	16.003	19.928	56.617	1.00	49.41
ATOM	1469	CA	ASP	190	16.392	17.047	58.021	1.00	55.01
ATOM	1470	C	ASP	190	17.556	16.115	58.338	1.00	56.16
ATOM	1471	O	ASP	190	18.083	16.100	59.463	1.00	58.30

FIG. 5AJ

ATOM	1472 CB ASP	190	15.195	16.734	58.955	1.00	63.89
ATOM	1473 CG ASP	190	14.592	15.365	58.686	1.00	99.67
ATOM	1474 OD1ASP	190	14.599	14.466	59.514	1.00	100.00
ATOM	1475 OD2ASP	190	14.088	15.240	57.470	1.00	100.00
ATOM	1476 N GLY	191	17.921	15.312	57.323	1.00	47.20
ATOM	1477 CA GLY	191	19.015	14.347	57.419	1.00	44.96
ATOM	1478 C GLY	191	20.359	15.044	57.587	1.00	34.43
ATOM	1479 O GLY	191	20.452	16.266	57.438	1.00	29.96
ATOM	1480 N PRO	192	21.402	14.264	57.905	1.00	27.26
ATOM	1481 CA PRO	192	22.737	14.834	58.100	1.00	24.01
ATOM	1482 C PRO	192	23.444	15.274	56.787	1.00	20.55
ATOM	1483 O PRO	192	23.323	14.648	55.740	1.00	23.84
ATOM	1484 CB PRO	192	23.583	13.764	58.825	1.00	21.00
ATOM	1485 CG PRO	192	22.739	12.501	58.915	1.00	27.49
ATOM	1486 CD PRO	192	21.330	12.863	58.448	1.00	27.26
ATOM	1487 N VAL	193	24.193	16.363	56.892	1.00	17.87
ATOM	1488 CA VAL	193	24.964	16.902	55.792	1.00	19.51
ATOM	1489 C VAL	193	26.380	17.108	56.249	1.00	22.37
ATOM	1490 O VAL	193	26.663	17.189	57.443	1.00	23.84
ATOM	1491 CB VAL	193	24.449	18.245	55.256	1.00	25.24
ATOM	1492 CG1VAL	193	23.059	18.118	54.632	1.00	21.90
ATOM	1493 CG2VAL	193	24.497	19.322	56.346	1.00	24.81
ATOM	1494 N LEU	194	27.253	17.241	55.277	1.00	19.04
ATOM	1495 CA LEU	194	28.654	17.438	55.516	1.00	20.29
ATOM	1496 C LEU	194	29.006	18.930	55.571	1.00	18.71
ATOM	1497 O LEU	194	28.907	19.615	54.591	1.00	20.13
ATOM	1498 CB LEU	194	29.412	16.806	54.327	1.00	22.92
ATOM	1499 CG LEU	194	29.994	15.423	54.542	1.00	30.60
ATOM	1500 CD1LEU	194	29.227	14.642	55.595	1.00	35.19
ATOM	1501 CD2LEU	194	30.048	14.672	53.211	1.00	25.61
ATOM	1502 N LEU	195	29.453	19.430	56.713	1.00	17.39
ATOM	1503 CA LEU	195	29.881	20.808	56.785	1.00	18.83
ATOM	1504 C LEU	195	31.389	20.837	56.579	1.00	28.32
ATOM	1505 O LEU	195	32.161	20.152	57.281	1.00	21.98
ATOM	1506 CB LEU	195	29.489	21.525	58.072	1.00	22.20
ATOM	1507 CG LEU	195	28.055	21.349	58.444	1.00	26.40
ATOM	1508 CD1LEU	195	27.937	21.508	59.941	1.00	31.99
ATOM	1509 CD2LEU	195	27.225	22.395	57.726	1.00	26.90
ATOM	1510 N PRO	196	31.789	21.610	55.597	1.00	21.58
ATOM	1511 CA PRO	196	33.177	21.666	55.154	1.00	22.17
ATOM	1512 C PRO	196	34.080	22.623	55.892	1.00	29.56

FIG. 5AK

ATOM	1513 O	PRO	196	33.635	23.588	56.490	1.00	29.04
ATOM	1514 CB	PRO	196	33.054	22.265	53.752	1.00	22.77
ATOM	1515 CG	PRO	196	31.761	23.104	53.735	1.00	18.99
ATOM	1516 CD	PRO	196	30.910	22.567	54.861	1.00	16.42
ATOM	1517 N	ASP	197	35.379	22.410	55.716	1.00	22.95
ATOM	1518 CA	ASP	197	36.364	23.370	56.134	1.00	19.71
ATOM	1519 C	ASP	197	36.556	24.295	54.931	1.00	24.74
ATOM	1520 O	ASP	197	36.251	23.913	53.800	1.00	24.88
ATOM	1521 CB	ASP	197	37.711	22.730	56.446	1.00	22.28
ATOM	1522 CG	ASP	197	37.690	21.913	57.687	1.00	43.93
ATOM	1523 OD1	ASP	197	36.912	22.117	58.608	1.00	53.47
ATOM	1524 OD2	ASP	197	38.634	21.006	57.694	1.00	31.58
ATOM	1525 N	ASN	198	37.062	25.501	55.168	1.00	19.74
ATOM	1526 CA	ASN	198	37.254	26.470	54.118	1.00	15.38
ATOM	1527 C	ASN	198	37.974	25.889	52.971	1.00	19.61
ATOM	1528 O	ASN	198	38.958	25.236	53.134	1.00	22.69
ATOM	1529 CB	ASN	198	38.013	27.704	54.614	1.00	24.48
ATOM	1530 CG	ASN	198	37.236	28.504	55.632	1.00	52.21
ATOM	1531 OD1	ASN	198	36.107	28.174	55.961	1.00	34.54
ATOM	1532 ND2	ASN	198	37.854	29.556	56.150	1.00	55.11
ATOM	1533 N	HIS	199	37.462	26.125	51.801	1.00	16.30
ATOM	1534 CA	HIS	199	38.071	25.627	50.616	1.00	15.80
ATOM	1535 C	HIS	199	37.496	26.357	49.450	1.00	14.85
ATOM	1536 O	HIS	199	36.757	27.295	49.643	1.00	16.45
ATOM	1537 CB	HIS	199	37.988	24.103	50.471	1.00	16.53
ATOM	1538 CG	HIS	199	36.597	23.628	50.218	1.00	16.65
ATOM	1539 ND1	HIS	199	35.695	23.491	51.244	1.00	17.85
ATOM	1540 CD2	HIS	199	35.987	23.282	49.048	1.00	18.67
ATOM	1541 CE1	HIS	199	34.561	23.052	50.688	1.00	19.45
ATOM	1542 NE2	HIS	199	34.716	22.905	49.364	1.00	18.74
ATOM	1543 N	TYR	200	37.879	25.998	48.247	1.00	12.56
ATOM	1544 CA	TYR	200	37.334	26.689	47.100	1.00	14.01
ATOM	1545 C	TYR	200	37.207	25.824	45.870	1.00	15.57
ATOM	1546 O	TYR	200	37.793	24.751	45.768	1.00	20.20
ATOM	1547 CB	TYR	200	38.030	28.011	46.779	1.00	19.79
ATOM	1548 CG	TYR	200	39.382	27.745	46.202	1.00	22.25
ATOM	1549 CD1	TYR	200	39.543	27.526	44.835	1.00	22.53
ATOM	1550 CD2	TYR	200	40.473	27.605	47.057	1.00	25.73
ATOM	1551 CE1	TYR	200	40.800	27.222	44.317	1.00	35.51
ATOM	1552 CE2	TYR	200	41.739	27.314	46.559	1.00	29.34
ATOM	1553 CZ	TYR	200	41.896	27.132	45.186	1.00	54.14

FIG. 5AL

ATOM	1554 OH	TYR	200	43.153	26.820	44.703	1.00	62.66
ATOM	1555 N	LEU	201	36.393	26.309	44.946	1.00	15.07
ATOM	1556 CA	LEU	201	36.147	25.680	43.678	1.00	11.01
ATOM	1557 C	LEU	201	36.753	26.532	42.593	1.00	17.30
ATOM	1558 O	LEU	201	36.619	27.753	42.610	1.00	20.19
ATOM	1559 CB	LEU	201	34.628	25.518	43.354	1.00	10.09
ATOM	1560 CG	LEU	201	33.749	25.027	44.480	1.00	13.41
ATOM	1561 CD1	LEU	201	32.293	24.938	43.954	1.00	17.11
ATOM	1562 CD2	LEU	201	34.196	23.635	44.927	1.00	23.03
ATOM	1563 N	SER	202	37.407	25.868	41.651	1.00	10.75
ATOM	1564 CA	SER	202	38.047	26.490	40.528	1.00	8.51
ATOM	1565 C	SER	202	37.222	26.189	39.294	1.00	11.56
ATOM	1566 O	SER	202	36.919	25.038	38.996	1.00	14.58
ATOM	1567 CB	SER	202	39.485	25.987	40.442	1.00	15.68
ATOM	1568 OG	SER	202	40.067	26.353	39.228	1.00	36.44
ATOM	1569 N	THR	203	36.798	27.241	38.601	1.00	12.36
ATOM	1570 CA	THR	203	35.879	27.067	37.499	1.00	15.60
ATOM	1571 C	THR	203	36.417	27.521	36.195	1.00	20.19
ATOM	1572 O	THR	203	37.192	28.472	36.114	1.00	18.29
ATOM	1573 CB	THR	203	34.565	27.892	37.757	1.00	20.51
ATOM	1574 OG1	THR	203	34.911	29.260	37.780	1.00	20.39
ATOM	1575 CG2	THR	203	33.935	27.557	39.093	1.00	6.80
ATOM	1576 N	GLN	204	35.913	26.883	35.164	1.00	10.30
ATOM	1577 CA	GLN	204	36.173	27.271	33.807	1.00	14.85
ATOM	1578 C	GLN	204	34.956	26.980	32.921	1.00	23.14
ATOM	1579 O	GLN	204	34.334	25.932	33.056	1.00	21.66
ATOM	1580 CB	GLN	204	37.475	26.696	33.237	1.00	20.33
ATOM	1581 CG	GLN	204	37.271	25.371	32.518	1.00	40.16
ATOM	1582 CD	GLN	204	38.588	24.722	32.193	1.00	59.76
ATOM	1583 OE1	GLN	204	39.011	24.716	31.035	1.00	41.80
ATOM	1584 NE2	GLN	204	39.276	24.241	33.235	1.00	34.80
ATOM	1585 N	SER	205	34.619	27.913	32.021	1.00	15.83
ATOM	1586 CA	SER	205	33.447	27.762	31.172	1.00	14.60
ATOM	1587 C	SER	205	33.654	28.307	29.783	1.00	20.21
ATOM	1588 O	SER	205	34.282	29.337	29.581	1.00	17.82
ATOM	1589 CB	SER	205	32.197	28.445	31.758	1.00	11.88
ATOM	1590 OG	SER	205	32.121	28.406	33.177	1.00	15.45
ATOM	1591 N	ALA	206	33.065	27.630	28.827	1.00	13.00
ATOM	1592 CA	ALA	206	33.079	28.029	27.426	1.00	9.99
ATOM	1593 C	ALA	206	31.623	28.192	26.924	1.00	21.23
ATOM	1594 O	ALA	206	30.809	27.306	27.139	1.00	14.10

FIG. 5AM

ATOM	1595	CB	ALA	206	33.751	26.936	26.596	1.00	13.45
ATOM	1596	N	LEU	207	31.335	29.320	26.263	1.00	16.09
ATOM	1597	CA	LEU	207	30.036	29.167	25.706	1.00	12.07
ATOM	1598	C	LEU	207	30.070	29.445	24.235	1.00	19.76
ATOM	1599	O	LEU	207	31.014	29.840	23.576	1.00	20.82
ATOM	1600	CB	LEU	207	29.580	31.057	26.004	1.00	8.24
ATOM	1601	CG	LEU	207	29.744	31.493	27.457	1.00	16.35
ATOM	1602	CD1	LEU	207	28.955	32.790	27.707	1.00	13.78
ATOM	1603	CD2	LEU	207	29.268	30.406	28.400	1.00	18.79
ATOM	1604	N	SER	208	29.011	28.863	23.698	1.00	15.35
ATOM	1605	CA	SER	208	28.914	28.692	22.270	1.00	13.74
ATOM	1606	C	SER	208	27.449	28.852	21.794	1.00	20.16
ATOM	1607	O	SER	208	26.548	29.085	22.594	1.00	15.81
ATOM	1608	CB	SER	208	29.495	27.367	21.822	1.00	17.82
ATOM	1609	OG	SER	208	28.769	26.311	22.431	1.00	31.45
ATOM	1610	N	LYS	209	27.242	28.738	20.485	1.00	16.50
ATOM	1611	CA	LYS	209	25.907	28.828	19.906	1.00	18.02
ATOM	1612	C	LYS	209	25.637	27.610	19.031	1.00	29.99
ATOM	1613	O	LYS	209	26.578	27.004	18.502	1.00	32.55
ATOM	1614	CB	LYS	209	25.783	30.100	19.082	1.00	20.96
ATOM	1615	CG	LYS	209	24.746	31.055	19.606	1.00	34.50
ATOM	1616	CD	LYS	209	25.262	31.964	20.666	1.00	22.72
ATOM	1617	CE	LYS	209	24.370	33.159	290.896	1.00	18.96
ATOM	1618	NZ	LYS	209	23.565	33.067	22.116	1.00	27.39
ATOM	1619	N	ASP	210	24.347	27.241	18.912	1.00	27.01
ATOM	1620	CA	ASP	210	23.890	26.159	18.038	1.00	24.62
ATOM	1621	C	ASP	210	23.465	26.793	16.705	1.00	26.77
ATOM	1622	O	ASP	210	22.468	27.514	16.605	1.00	23.00
ATOM	1623	CB	ASP	210	22.744	25.361	18.691	1.00	24.43
ATOM	1624	CG	ASP	210	22.197	24.249	17.839	1.00	35.55
ATOM	1625	OD1	ASP	210	22.333	24.185	16.631	1.00	36.53
ATOM	1626	OD2	ASP	210	21.499	23.400	18.535	1.00	45.51
ATOM	1627	N	PRO	211	24.306	26.618	15.708	1.00	30.25
ATOM	1628	CA	PRO	211	24.120	27.224	14.397	1.00	30.30
ATOM	1629	C	PRO	211	22.733	26.982	13.770	1.00	39.72
ATOM	1630	O	PRO	211	22.253	27.782	12.959	1.00	37.65
ATOM	1631	CB	PRO	211	25.197	26.620	13.500	1.00	29.99
ATOM	1632	CG	PRO	211	25.782	25.418	14.255	1.00	38.59
ATOM	1633	CD	PRO	211	25.158	25.405	15.647	1.00	35.05
ATOM	1634	N	ASN	212	22.102	25.868	14.140	1.00	39.64
ATOM	1635	CA	ASN	212	20.808	25.515	13.592	1.00	39.60

FIG. 5AN

ATOM	1636 C	ASN	212	19.642	25.894	14.497	1.00	41.92
ATOM	1637 O	ASN	212	18.485	25.518	14.263	1.00	42.30
ATOM	1638 CB	ASN	212	20.788	24.028	13.236	1.00	48.64
ATOM	1639 CG	ASN	212	21.883	23.678	12.230	1.00	53.61
ATOM	1640 N	GLU	213	19.947	26.675	15.520	1.00	27.84
ATOM	1641 CA	GLU	213	18.953	27.080	16.478	1.00	20.43
ATOM	1642 C	GLU	213	18.485	28.527	16.241	1.00	29.95
ATOM	1643 O	GLU	213	19.247	29.475	16.324	1.00	32.77
ATOM	1644 CB	GLU	213	19.535	26.878	17.894	1.00	16.45
ATOM	1645 CG	GLU	213	18.594	27.326	18.995	1.00	18.29
ATOM	1646 CD	GLU	213	17.229	26.703	18.853	1.00	38.01
ATOM	1647 OE1	GLU	213	16.238	27.334	18.508	1.00	25.07
ATOM	1648 OE2	GLU	213	17.223	25.423	19.122	1.00	19.17
ATOM	1649 N	LYS	214	17.223	28.713	15.963	1.00	22.99
ATOM	1650 CA	LYS	214	16.721	30.081	15.726	1.00	22.84
ATOM	1651 C	LYS	214	16.252	30.778	16.982	1.00	21.50
ATOM	1652 O	LYS	214	16.130	32.016	17.032	1.00	28.15
ATOM	1653 CB	LYS	214	15.653	30.197	14.606	1.00	27.58
ATOM	1654 CG	LYS	214	16.153	29.816	13.209	1.00	32.71
ATOM	1655 CD	LYS	214	16.752	30.979	12.431	1.00	55.31
ATOM	1656 N	ARG	215	15.947	30.028	18.014	1.00	14.52
ATOM	1657 CA	ARG	215	15.518	30.726	19.209	1.00	15.58
ATOM	1658 C	ARG	215	16.719	31.382	19.892	1.00	21.87
ATOM	1659 O	ARG	215	17.848	31.075	19.572	1.00	26.69
ATOM	1660 CB	ARG	215	14.808	29.804	20.159	1.00	18.82
ATOM	1661 CG	ARG	215	13.660	29.067	19.475	1.00	23.30
ATOM	1662 CD	ARG	215	13.220	27.806	20.205	1.00	15.45
ATOM	1663 NE	ARG	215	14.107	26.668	19.929	1.00	28.08
ATOM	1664 CZ	ARG	215	14.022	25.473	20.543	1.00	21.38
ATOM	1665 NH1	ARG	215	13.074	25.215	21.455	1.00	23.92
ATOM	1666 NH2	ARG	215	14.893	24.514	20.225	1.00	20.46
ATOM	1667 N	ASP	216	16.466	32.275	20.830	1.00	16.72
ATOM	1668 CA	ASP	216	17.556	32.895	21.617	1.00	19.06
ATOM	1669 C	ASP	216	18.047	31.817	22.607	1.00	20.02
ATOM	1670 O	ASP	216	17.261	31.214	23.350	1.00	18.45
ATOM	1671 CB	ASP	216	17.066	34.169	22.383	1.00	21.33
ATOM	1672 CG	ASP	216	18.138	35.140	22.893	1.00	20.97
ATOM	1673 OD1	ASP	216	17.869	36.079	23.620	1.00	28.46
ATOM	1674 OD2	ASP	216	19.342	34.900	22.441	1.00	20.37
ATOM	1675 N	HIS	217	19.332	31.537	22.589	1.00	13.18
ATOM	1676 CA	HIS	217	19.813	30.482	23.433	1.00	11.21

FIG. 5A0

ATOM	1677 C	HIS	217	21.313	30.614	23.723	1.00	21.35
ATOM	1678 O	HIS	217	22.014	31.471	23.163	1.00	15.03
ATOM	1679 CB	HIS	217	19.587	29.168	22.690	1.00	13.03
ATOM	1680 CG	HIS	217	20.525	29.025	21.542	1.00	15.49
ATOM	1681 ND1	HIS	217	20.463	29.871	20.449	1.00	17.88
ATOM	1682 CD2	HIS	217	21.589	28.172	21.361	1.00	17.51
ATOM	1683 CE1	HIS	217	21.457	29.524	19.635	1.00	17.94
ATOM	1684 NE2	HIS	217	22.152	28.501	20.151	1.00	17.59
ATOM	1685 N	MSE	218	21.794	29.725	24.576	1.00	11.26
ATOM	1686 CA	MSE	218	23.186	29.642	24.887	1.00	11.49
ATOM	1687 C	MSE	218	23.560	28.198	25.094	1.00	24.15
ATOM	1688 O	MSE	218	22.822	27.446	25.751	1.00	20.70
ATOM	1689 CB	MSE	218	23.539	30.421	26.172	1.00	12.84
ATOM	1690 CG	MSE	218	24.809	30.004	26.907	1.00	12.59
ATOM	1691 SE	MSE	218	25.267	31.128	28.434	1.00	29.94
ATOM	1692 CE	MSE	218	24.039	30.502	29.781	1.00	13.54
ATOM	1693 N	VAL	219	24.727	27.824	24.558	1.00	15.62
ATOM	1694 CA	VAL	219	25.309	26.518	24.782	1.00	10.58
ATOM	1695 C	VAL	219	26.473	26.689	25.753	1.00	16.54
ATOM	1696 O	VAL	219	27.280	27.604	25.585	1.00	15.54
ATOM	1697 CB	VAL	219	25.774	25.883	23.498	1.00	15.08
ATOM	1698 CG1	VAL	219	26.330	24.495	23.824	1.00	14.34
ATOM	1699 CG2	VAL	219	24.599	25.766	22.512	1.00	15.78
ATOM	1700 N	LEU	220	26.523	25.836	26.783	1.00	10.95
ATOM	1701 CA	LEU	220	27.490	25.939	27.850	1.00	11.01
ATOM	1702 C	LEU	220	28.206	24.643	28.184	1.00	21.26
ATOM	1703 O	LEU	220	27.592	23.577	28.324	1.00	15.94
ATOM	1704 CB	LEU	220	26.807	26.545	29.100	1.00	13.75
ATOM	1705 CG	LEU	220	27.624	26.578	30.402	1.00	21.10
ATOM	1706 CD1	LEU	220	28.433	27.875	30.483	1.00	23.53
ATOM	1707 CD2	LEU	220	26.663	26.556	31.586	1.00	22.04
ATOM	1708 N	LEU	221	29.570	24.758	28.273	1.00	19.04
ATOM	1709 CA	LEU	221	30.498	23.666	28.697	1.00	13.22
ATOM	1710 C	LEU	221	31.309	24.178	29.887	1.00	10.73
ATOM	1711 O	LEU	221	31.846	25.267	29.857	1.00	12.98
ATOM	1712 CB	LEU	221	31.382	23.102	27.549	1.00	13.74
ATOM	1713 CG	LEU	221	32.580	22.257	28.045	1.00	18.64
ATOM	1714 CD1	LEU	221	32.149	20.868	28.496	1.00	17.38
ATOM	1715 CD2	LEU	221	33.571	22.109	26.911	1.00	26.97
ATOM	1716 N	GLU	222	31.316	23.446	30.963	1.00	9.31
ATOM	1717 CA	GLU	222	31.936	23.929	32.144	1.00	9.97

FIG. 5AP

ATOM	1718 C	GLU	222	32.548	22.803	32.951	1.00	12.94
ATOM	1719 O	GLU	222	32.072	21.662	32.966	1.00	13.38
ATOM	1720 CB	GLU	222	30.836	24.762	32.896	1.00	12.14
ATOM	1721 CG	GLU	222	31.092	25.119	34.364	1.00	13.88
ATOM	1722 CD	GLU	222	29.895	25.891	34.934	1.00	13.57
ATOM	1723 OE1	GLU	222	29.128	26.477	34.240	1.00	19.47
ATOM	1724 OE2	GLU	222	29.752	25.789	36.207	1.00	18.51
ATOM	1725 N	PHE	223	33.687	23.123	33.542	1.00	15.86
ATOM	1726 CA	PHE	223	34.476	22.227	34.373	1.00	9.34
ATOM	1727 C	PHE	223	34.711	22.864	35.722	1.00	11.08
ATOM	1728 O	PHE	223	35.028	24.055	35.828	1.00	19.86
ATOM	1729 CB	PHE	223	35.847	21.919	33.684	1.00	8.30
ATOM	1730 CG	PHE	223	35.703	21.134	32.431	1.00	10.50
ATOM	1731 CD1	PHE	223	35.570	19.747	32.469	1.00	13.56
ATOM	1732 CD2	PHE	223	35.750	21.750	31.184	1.00	11.32
ATOM	1733 CE1	PHE	223	35.481	19.010	31.287	1.00	12.58
ATOM	1734 CE2	PHE	223	35.667	21.032	29.995	1.00	12.17
ATOM	1735 CZ	PHE	223	35.521	19.648	30.050	1.00	10.87
ATOM	1736 N	VAL	224	34.542	22.081	36.765	1.00	9.28
ATOM	1737 CA	VAL	224	34.708	22.587	38.080	1.00	11.18
ATOM	1738 C	VAL	224	35.324	21.553	39.010	1.00	17.52
ATOM	1739 O	VAL	224	34.848	20.418	39.137	1.00	13.17
ATOM	1740 CB	VAL	224	33.370	23.078	38.662	1.00	16.61
ATOM	1741 CG1	VAL	224	33.622	23.736	40.022	1.00	13.90
ATOM	1742 CG2	VAL	224	32.674	24.048	37.697	1.00	13.85
ATOM	1743 N	THR	225	36.380	21.965	39.676	1.00	11.71
ATOM	1744 CA	THR	225	37.026	21.099	40.617	1.00	11.61
ATOM	1745 C	THR	225	37.366	21.798	41.927	1.00	14.76
ATOM	1746 O	THR	225	37.702	23.002	41.962	1.00	16.64
ATOM	1747 CB	THR	225	38.162	20.279	40.014	1.00	20.38
ATOM	1748 OG1	THR	225	39.288	20.337	40.822	1.00	30.44
ATOM	1749 CG2	THR	225	38.468	20.722	38.631	1.00	10.89
ATOM	1750 N	ALA	226	37.222	21.065	43.011	1.00	7.89
ATOM	1751 CA	ALA	226	37.478	21.595	44.352	1.00	11.63
ATOM	1752 C	ALA	226	38.969	21.558	44.677	1.00	16.61
ATOM	1753 O	ALA	226	39.687	20.699	44.199	1.00	15.60
ATOM	1754 CB	ALA	226	36.695	20.847	45.444	1.00	12.17
ATOM	1755 N	ALA	227	39.395	22.490	45.479	1.00	13.95
ATOM	1756 CA	ALA	227	40.789	22.550	45.871	1.00	19.64
ATOM	1757 C	ALA	227	40.987	23.299	47.170	1.00	26.33
ATOM	1758 O	ALA	227	40.042	23.715	47.840	1.00	25.39

FIG. 5AQ

ATOM	1759	CB	ALA	227	41.557	23.246	44.760	1.00	18.42
ATOM	1760	N	GLY	228	42.245	23.476	47.523	1.00	23.28
ATOM	1761	CA	GLY	228	42.616	24.292	48.658	1.00	21.61
ATOM	1762	C	GLY	228	42.805	23.562	49.939	1.00	32.93
ATOM	1763	O	GLY	228	42.948	24.201	51.009	1.00	32.53
ATOM	1764	N	ILE	229	42.803	22.231	49.842	1.00	33.59
ATOM	1765	CA	ILE	229	43.006	21.375	50.998	1.00	31.81
ATOM	1766	C	ILE	229	44.016	20.291	50.633	1.00	28.78
ATOM	1767	O	ILE	229	45.090	20.176	51.246	1.00	96.02
ATOM	1768	CB	ILE	229	41.691	20.772	51.519	1.00	35.70
ATOM	1769	CG1	ILE	229	40.890	21.807	52.325	1.00	30.66
ATOM	1770	CG2	ILE	229	41.990	19.549	52.392	1.00	33.37
ATOM	1771	CD1	ILE	229	39.386	21.715	52.092	1.00	38.74
ATOM	1772	O	HOH	301	27.530	12.735	38.010	1.00	15.09
ATOM	1773	O	HOH	302	23.919	34.589	37.331	1.00	10.29
ATOM	1774	O	HOH	303	27.229	34.816	35.487	1.00	11.12
ATOM	1775	O	HOH	304	29.914	18.943	44.692	1.00	16.10
ATOM	1776	O	HOH	305	30.956	21.886	49.900	1.00	21.47
ATOM	1777	O	HOH	306	20.072	31.196	43.592	1.00	16.85
ATOM	1778	O	HOH	307	26.660	48.630	33.797	1.00	24.67
ATOM	1779	O	HOH	308	22.329	33.239	41.399	1.00	14.11
ATOM	1780	O	HOH	309	22.465	48.025	32.810	1.00	18.51
ATOM	1781	O	HOH	310	31.012	39.126	29.118	1.00	16.01
ATOM	1782	O	HOH	311	33.067	35.809	33.010	1.00	19.92
ATOM	1783	O	HOH	312	31.130	37.076	30.841	1.00	12.68
ATOM	1784	O	HOH	313	40.304	30.058	38.616	1.00	56.07
ATOM	1785	O	HOH	314	34.166	26.379	57.222	1.00	22.58
ATOM	1786	O	HOH	315	36.215	35.320	43.598	1.00	22.30
ATOM	1787	O	HOH	316	33.866	29.786	34.671	1.00	12.21
ATOM	1865	O	HOH	317	42.341	20.166	43.534	1.00	26.67
ATOM	1788	O	HOH	318	10.270	28.684	30.403	1.00	43.66
ATOM	1789	O	HOH	319	28.448	16.822	30.655	1.00	25.44
ATOM	1790	O	HOH	320	30.612	20.922	37.231	1.00	21.57
ATOM	1791	O	HOH	321	11.639	37.421	26.801	1.00	34.12
ATOM	1792	O	HOH	322	27.030	37.308	36.869	1.00	13.10
ATOM	1793	O	HOH	323	33.119	14.524	43.070	1.00	30.93
ATOM	1794	O	HOH	324	37.973	14.036	53.352	1.00	35.39
ATOM	1795	O	HOH	235	32.015	49.100	37.028	1.00	59.37
ATOM	1796	O	HOH	326	11.959	12.020	43.429	1.00	29.06
ATOM	1797	O	HOH	327	36.760	29.941	31.666	1.00	22.03
ATOM	1864	O	HOH	328	15.305	26.513	15.694	1.00	39.62

FIG. 5AR

ATOM	1798 O	HOH	329	33.005	46.924	36.994	1.00	22.07
ATOM	1863 O	HOH	330	23.801	36.134	22.715	1.00	45.30
ATOM	1799 O	HOH	331	33.609	31.296	26.261	1.00	23.65
ATOM	1862 O	HOH	332	34.942	24.780	29.532	1.00	38.93
ATOM	1800 O	HOH	333	25.235	12.919	54.611	1.00	36.20
ATOM	1861 O	HOH	334	38.048	23.467	36.645	1.00	37.73
ATOM	1801 O	HOH	335	12.284	43.511	38.338	1.00	33.79
ATOM	1802 O	HOH	336	9.826	47.020	32.568	1.00	46.67
ATOM	1803 O	HOH	337	7.671	41.532	29.806	1.00	40.88
ATOM	1804 O	HOH	338	15.430	23.713	26.808	1.00	34.73
ATOM	1805 O	HOH	339	24.344	20.385	25.121	1.00	53.42
ATOM	1806 O	HOH	340	31.550	10.656	40.819	1.00	47.85
ATOM	1807 O	HOH	341	17.569	23.030	25.796	1.00	28.17
ATOM	1808 O	HOH	342	19.174	38.552	23.965	1.00	45.54
ATOM	1809 O	HOH	343	24.268	37.527	25.415	1.00	30.97
ATOM	1810 O	HOH	344	21.266	29.482	41.551	1.00	19.69
ATOM	1811 O	HOH	345	20.668	26.999	41.933	1.00	11.81
ATOM	1812 O	HOH	346	24.780	24.795	43.460	1.00	20.95
ATOM	1813 O	HOH	347	42.962	13.170	46.312	1.00	31.00
ATOM	1814 O	HOH	348	32.322	14.088	47.013	1.00	28.20
ATOM	1815 O	HOH	349	31.708	13.186	49.679	1.00	35.57
ATOM	1816 O	HOH	350	22.408	35.801	50.514	1.00	40.71
ATOM	1817 O	HOH	351	25.366	47.090	42.583	1.00	38.15
ATOM	1818 O	HOH	352	27.243	47.647	43.977	1.00	41.55
ATOM	1819 O	HOH	353	29.868	45.076	42.906	1.00	29.32
ATOM	1820 O	HOH	354	14.175	22.269	42.680	1.00	74.11
ATOM	1821 O	HOH	355	13.414	10.739	35.791	1.00	29.92
ATOM	1822 O	HOH	356	20.338	9.974	37.765	1.00	30.46
ATOM	1823 O	HOH	357	23.520	40.420	24.953	1.00	29.75
ATOM	1824 O	HOH	358	25.718	41.692	26.023	1.00	30.43
ATOM	1825 O	HOH	359	26.826	38.466	25.345	1.00	31.72
ATOM	1826 O	HOH	360	37.768	42.373	25.123	1.00	41.53
ATOM	1827 O	HOH	361	40.078	42.268	25.852	1.00	37.12
ATOM	1828 O	HOH	362	31.483	38.677	22.083	1.00	54.21
ATOM	1829 O	HOH	363	33.891	37.723	30.126	1.00	23.35
ATOM	1860 O	HOH	364	39.936	26.543	36.329	1.00	47.93
ATOM	1830 O	HOH	365	36.631	34.210	41.636	1.00	62.74
ATOM	1831 O	HOH	366	37.038	29.783	52.197	1.00	40.07
ATOM	1832 O	HOH	367	37.289	37.407	40.231	1.00	37.59
ATOM	1833 O	HOH	368	18.930	17.517	52.472	1.00	35.80
ATOM	1834 O	HOH	369	19.506	18.914	57.913	1.00	45.72

FIG. 5AS

ATOM	1835 O	HOH	370	30.903	26.708	41.139	1.00	21.54
ATOM	1836 O	HOH	371	30.369	25.678	24.583	1.00	22.46
ATOM	1837 O	HOH	372	21.000	33.705	20.826	1.00	26.00
ATOM	1838 O	HOH	373	13.648	32.794	21.329	1.00	27.98
ATOM	1839 O	HOH	374	29.735	25.683	38.707	1.00	21.00
ATOM	1859 O	HOH	375	33.670	24.419	60.503	1.00	50.04
ATOM	1840 O	HOH	376	30.034	11.047	37.420	1.00	43.28
ATOM	1841 O	HOH	377	8.662	35.846	35.068	1.00	51.94
ATOM	1842 O	HOH	378	10.847	36.466	39.503	1.00	42.32
ATOM	1843 O	HOH	379	14.395	48.943	39.085	1.00	29.72
ATOM	1844 O	HOH	380	36.676	11.660	40.172	1.00	39.81
ATOM	1845 O	HOH	381	35.968	7.212	34.763	1.00	58.66
ATOM	1846 O	HOH	382	17.426	21.988	21.077	1.00	41.69
ATOM	1847 O	HOH	383	29.837	22.623	39.378	1.00	32.82
ATOM	1848 O	HOH	384	23.855	29.386	55.164	1.00	55.00
ATOM	1849 O	HOH	385	17.408	35.360	47.495	1.00	61.61
ATOM	1850 O	HOH	386	27.900	49.720	42.445	1.00	47.70
ATOM	1851 O	HOH	387	13.932	36.230	44.385	1.00	45.08
ATOM	1852 O	HOH	388	12.650	28.021	43.288	1.00	49.86
ATOM	1853 O	HOH	389	16.974	42.367	43.435	1.00	34.38
ATOM	1854 O	HOH	390	37.335	42.653	28.295	1.00	64.46
ATOM	1855 O	HOH	391	29.701	49.856	35.323	1.00	62.61
ATOM	1856 O	HOH	392	27.267	50.835	33.976	1.00	66.60
ATOM	1857 O	HOH	393	19.661	29.181	51.537	1.00	34.01
ATOM	1858 O	HOH	394	29.412	17.505	59.089	1.00	51.78
TER								
END								

FIG. 5AT







FIG. 6

FIG. 7

FIG. 8






FIG. 9

FIG. 10

FIG. 11





FIG. 12




FIG. 13





FIG. 14

FIG. 15