(12) STANDARD PATENT (11) Application No. **AU 2013361540 B2**
(19) AUSTRALIAN PATENT OFFICE

(54) Title
Pesticidal compositions and processes related thereto

(51) International Patent Classification(s)
C07C 211/27 (2006.01) ***A01N 25/26*** (2006.01)

(21) Application No: **2013361540** (22) Date of Filing: **2013.12.18**

(87) WIPO No: **WO14/100190**

(30) Priority Data

(31) Number	(32) Date	(33) Country
61/739,025	**2012.12.19**	**US**

(43) Publication Date: **2014.06.26**
(44) Accepted Journal Date: **2017.02.02**

(71) Applicant(s)
Dow AgroSciences LLC

(72) Inventor(s)
Lo, William C.;Hunter, James E.;Watson, Gerald B.;Patny, Akshay;Iyer, Pravin S.;Boruwa, Joshodeep

(74) Agent / Attorney
FPA Patent Attorneys Pty Ltd, Level 43 101 Collins Street, Melbourne, VIC, 3000

(56) Related Art
WO 2012/177813 A1

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau

(43) International Publication Date
26 June 2014 (26.06.2014)

WIPO | PCT

(10) International Publication Number
WO 2014/100190 A1

(51) International Patent Classification:
A01N 25/26 (2006.01)

(21) International Application Number:
PCT/US2013/076142

(22) International Filing Date:
18 December 2013 (18.12.2013)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
61/739,025 19 December 2012 (19.12.2012) US

(71) Applicant: **DOW AGROSCIENCES LLC** [US/US]; 9330 Zionsville Road, Indianapolis, IN 46268 (US).

(72) Inventors: **LO, William C.**; 12027 Weathered Edge Drive, Fishers, IN 46037 (US). **HUNTER, James E.**; 5953 North Evanston Avenue, Indianapolis, IN 46220 (US). **WATSON, Gerald B.**; 535 Quail Valley Drive, Zionsville, IN 46077 (US). **PATNY, Akshay**; 35 Crescent Street, Apt 410, Waltham, MA 02453 (US). **IYER, Pravin S.**; 302 Manbhum Millenium Residency, St. 10 West Marredpally, Secunderabad 500 026 (IN). **BORUWA, Joshodeep**; 2-5/87 Navaratna House, Habsiguda Street No 2, Hyderabad 500 007 (IN).

(74) Agent: **CORVIN, Carl**; Dow AgroSciences LLC, 9330 Zionsville Rd, Indianapolis, Indiana 46268 (US).

(81) Designated States *(unless otherwise indicated, for every kind of national protection available)*: AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States *(unless otherwise indicated, for every kind of regional protection available)*: ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:


— *as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))*

Published:

— *with international search report (Art. 21(3))*

— *before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))*

(54) Title: PESTICIDAL COMPOSITIONS AND PROCESSES RELATED THERETO



Formula One

(57) Abstract: This document discloses molecules having the following formula ("Formula One"): and processes associated therewith.

WO 2014/100190 A1

2013361540 12 Dec 2016

PESTICIDAL COMPOSITIONS AND PROCESSES RELATED THERETO

CROSS REFERENCE TO RELATED APPLICATIONS

This application claims the benefit of U.S. Provisional Patent Application Serial No. 61/739,025 filed December 19, 2012, the disclosure of which is expressly incorporated herein by reference.

FIELD OF THE DISCLOSURE

The invention disclosed in this document is related to the field of processes to produce molecules that are useful as pesticides (*e.g.*, acaricides, insecticides, molluscicides, and nematicides), such molecules, and processes of using such molecules to control pests.

BACKGROUND OF THE DISCLOSURE

Reference to any prior art in the specification is not, and should not be taken as, an acknowledgment or any form of suggestion that this prior art forms part of the common general knowledge in Australia or any other jurisdiction.

Pests cause millions of human deaths around the world each year. Furthermore, there are more than ten thousand species of pests that cause losses in agriculture. The world-wide agricultural losses amount to billions of U.S. dollars each year.

Termites cause damage to all kinds of private and public structures. The world-wide termite damage losses amount to billions of U.S. dollars each year.

Stored food pests eat and adulterate stored food. The world-wide stored food losses amount to billions of U.S. dollars each year, but more importantly, deprive people of needed food.

There is an acute need for new pesticides. Certain pests are developing resistance to pesticides in current use. Hundreds of pest species are resistant to one or more pesticides. The development of resistance to some of the older pesticides, such as DDT, the carbamates, and the organophosphates, is well known. But resistance has even developed to some of the newer pesticides, for example, imidacloprid.

Therefore, for many reasons, including the above reasons, a need exists for new pesticides.

DEFINITIONS

The examples given in the definitions are generally non-exhaustive and must not be construed as limiting the invention disclosed in this document. It is understood that a substituent should comply with chemical bonding rules and steric compatibility constraints in relation to the particular molecule to which it is attached.

201336154O 12 Dec 2016

"**Alkenyl**" means an acyclic, unsaturated (at least one carbon-carbon double bond), branched or unbranched, substituent consisting of carbon and hydrogen, for example, vinyl, allyl, butenyl, pentenyl, and hexenyl.

"**Alkenyloxy**" means an alkenyl further consisting of a carbon-oxygen single bond, for example, allyloxy, butenyloxy, pentenyloxy, hexenyloxy.

"**Alkoxy**" means an alkyl further consisting of a carbon-oxygen single bond, for example, methoxy, ethoxy, propoxy, isopropoxy, butoxy, isobutoxy, and *tert*-butoxy.

"**Alkyl**" means an acyclic, saturated, branched or unbranched, substituent consisting of carbon and hydrogen, for example, methyl, ethyl, (C_3)alkyl which represents *n*-propyl and isopropyl), (C_4)alkyl which represents *n*-butyl, *sec*-butyl, isobutyl, and *tert*-butyl.

"**Alkynyl**" means an acyclic, unsaturated (at least one carbon-carbon triple bond), branched or unbranched, substituent consisting of carbon and hydrogen, for example, ethynyl, propargyl, butynyl, and pentynyl.

"**Alkynyloxy**" means an alkynyl further consisting of a carbon-oxygen single bond, for example, pentynyloxy, hexynyloxy, heptynyloxy, and octynyloxy.

"**Aryl**" means a cyclic, aromatic substituent consisting of hydrogen and carbon, for example, phenyl, naphthyl, and biphenyl.

"**(C_x-C_y)**" where the subscripts "x" and "y" are integers such as 1, 2, or 3, means the range of carbon atoms for a substituent – for example, (C_1-C_4)alkyl means methyl, ethyl, *n*-propyl, isopropyl, *n*-butyl, *sec*-butyl, isobutyl, and *tert*-butyl, each individually.

"**Cycloalkenyl**" means a monocyclic or polycyclic, unsaturated (at least one carbon-carbon double bond) substituent consisting of carbon and hydrogen, for example, cyclobutenyl, cyclopentenyl, cyclohexenyl, norbornenyl, bicyclo[2.2.2]octenyl, tetrahydronaphthyl, hexahydronaphthyl, and octahydronaphthyl.

"**Cycloalkenyloxy**" means a cycloalkenyl further consisting of a carbon-oxygen single bond, for example, cyclobutenyloxy, cyclopentenyloxy, norbornenyloxy, and bicyclo[2.2.2]octenyloxy.

"**Cycloalkyl**" means a monocyclic or polycyclic, saturated substituent consisting of carbon and hydrogen, for example, cyclopropyl, cyclobutyl, cyclopentyl, norbornyl, bicyclo[2.2.2]octyl, and decahydronaphthyl.

"**Cycloalkoxy**" means a cycloalkyl further consisting of a carbon-oxygen single bond, for example, cyclopropyloxy, cyclobutyloxy, cyclopentyloxy, norbornyloxy, and bicyclo[2.2.2]octyloxy.

"**Halo**" means fluoro, chloro, bromo, and iodo.

"**Haloalkoxy**" means an alkoxy further consisting of, from one to the maximum possible number of identical or different, halos, for example, fluoromethoxy,

12 Dec 2016 2013361540

trifluoromethoxy, 2,2-difluoropropoxy, chloromethoxy, trichloromethoxy, 1,1,2,2-tetrafluoroethoxy, and pentafluoroethoxy.


"**Haloalkyl**" means an alkyl further consisting of, from one to the maximum possible number of, identical or different, halos, for example, fluoromethyl, trifluoromethyl, 2,2-difluoropropyl, chloromethyl, trichloromethyl, and 1,1,2,2-tetrafluoroethyl.

"**Heterocyclyl**" means a cyclic substituent that may be fully saturated, partially unsaturated, or fully unsaturated, where the cyclic structure contains at least one carbon and at least one heteroatom, where said heteroatom is nitrogen, sulfur, or oxygen. In the case of sulfur, that atom can be in other oxidation states such as a sulfoxide and sulfone. Examples of aromatic heterocyclyls include, but are not limited to, benzofuranyl, benzoisothiazolyl, benzoisoxazolyl, benzoxazolyl, benzothienyl, benzothiazolyl, cinnolinyl, furanyl, imidazolyl, indazolyl, indolyl, isoindolyl, isoquinolinyl, isothiazolyl, isoxazolyl, oxadiazolyl, oxazolinyl, oxazolyl, phthalazinyl, pyrazinyl, pyrazolinyl, pyrazolyl, pyridazinyl, pyridyl, pyrimidinyl, pyrrolyl, quinazolinyl, quinolinyl, quinoxalinyl, tetrazolyl, thiazolinyl, thiazolyl, thienyl, triazinyl, and triazolyl. Examples of fully saturated heterocyclyls include, but are not limited to, piperazinyl, piperidinyl, morpholinyl, pyrrolidinyl, oxetanyl, tetrahydrofuranyl, tetrahydrothienyl and tetrahydropyranyl. Examples of partially unsaturated heterocyclyls include, but are not limited to, 1,2,3,4-tetrahydroquinolinyl, 4,5-dihydro-oxazolyl, 4,5-dihydro-1*H*-pyrazolyl, 4,5-dihydro-isoxazolyl, and 2,3-dihydro-[1,3,4]-oxadiazolyl.


Additional examples include the following



thietanyl



thietanyl-oxide




thietanyl-dioxide.

DETAILED DESCRIPTION OF THE DISCLOSURE

As used herein, the term "comprise" and variations of the term, such as "comprising", "comprises" and "comprised", are not intended to exclude other additives, components, integers or steps.

2013361540 12 Dec 2016

This document discloses molecules having the following formula ("**Formula One**"):



Formula One

wherein:

(a) **R1** is selected from

(1) H, F, Cl, Br, I, CN, NO_2, $(C_1$-$C_8)$alkyl, halo$(C_1$-$C_8)$alkyl, $(C_1$-$C_8)$alkoxy, halo$(C_1$-$C_8)$alkoxy, S$(C_1$-$C_8)$alkyl, S(halo$(C_1$-$C_8)$alkyl), S(O)$(C_1$-$C_8)$alkyl, S(O)(halo$(C_1$-$C_8)$alkyl), $S(O)_2(C_1$-$C_8)$alkyl, $S(O)_2$(halo$(C_1$-$C_8)$alkyl), N(R14)(R15),

(2) **substituted $(C_1$-$C_8)$alkyl**, wherein said substituted $(C_1$-$C_8)$alkyl has one or more substituents selected from CN and NO_2,

(3) **substituted halo$(C_1$-$C_8)$alkyl**, wherein said substituted halo$(C_1$-$C_8)$alkyl, has one or more substituents selected from CN and NO_2,

(4) **substituted $(C_1$-$C_8)$alkoxy**, wherein said substituted $(C_1$-$C_8)$alkoxy has one or more substituents selected from CN and NO_2, and

(5) **substituted halo$(C_1$-$C_8)$alkoxy**, wherein said substituted halo$(C_1$-$C_8)$alkoxy has one or more substituents selected from CN and NO_2;

(b) **R2** is selected from

(1) H, F, Cl, Br, I, CN, NO_2, $(C_1$-$C_8)$alkyl, halo$(C_1$-$C_8)$alkyl, $(C_1$-$C_8)$alkoxy, halo$(C_1$-$C_8)$alkoxy, S$(C_1$-$C_8)$alkyl, S(halo$(C_1$-$C_8)$alkyl), S(O)$(C_1$-$C_8)$alkyl, S(O)(halo$(C_1$-$C_8)$alkyl), $S(O)_2(C_1$-$C_8)$alkyl, $S(O)_2$(halo$(C_1$-$C_8)$alkyl), N(R14)(R15),

(2) **substituted $(C_1$-$C_8)$alkyl**, wherein said substituted $(C_1$-$C_8)$alkyl has one or more substituents selected from CN and NO_2,

(3) **substituted halo$(C_1$-$C_8)$alkyl**, wherein said substituted halo$(C_1$-$C_8)$alkyl, has one or more substituents selected from CN and NO_2,

(4) **substituted $(C_1$-$C_8)$alkoxy**, wherein said substituted $(C_1$-$C_8)$alkoxy has one or more substituents selected from CN and NO_2, and

(5) **substituted halo$(C_1$-$C_8)$alkoxy**, wherein said substituted halo$(C_1$-$C_8)$alkoxy has one or more substituents selected from CN and NO_2;

(c) **R3** is selected from

(1) H, F, Cl, Br, I, CN, NO_2, $(C_1$-$C_8)$alkyl, halo$(C_1$-$C_8)$alkyl, $(C_1$-$C_8)$alkoxy, halo$(C_1$-$C_8)$alkoxy, S$(C_1$-$C_8)$alkyl, S(halo$(C_1$-$C_8)$alkyl), S(O)$(C_1$-$C_8)$alkyl, S(O)(halo$(C_1$-$C_8)$alkyl), $S(O)_2(C_1$-$C_8)$alkyl, $S(O)_2$(halo$(C_1$-$C_8)$alkyl), N(R14)(R15),

(2) **substituted $(C_1$-$C_8)$alkyl**, wherein said substituted $(C_1$-$C_8)$alkyl has one or more substituents selected from CN and NO_2,

(3) **substituted halo$(C_1$-$C_8)$alkyl**, wherein said substituted halo$(C_1$-$C_8)$alkyl, has one or more substituents selected from CN and NO_2,

(4) **substituted (C_1-C_8)alkoxy**, wherein said substituted (C_1-C_8)alkoxy has one or more substituents selected from CN and NO_2, and

(5) **substituted halo(C_1-C_8)alkoxy**, wherein said substituted halo(C_1-C_8)alkoxy has one or more substituents selected from CN and NO_2;

(d) **R4** is selected from

(1) H, F, Cl, Br, I, CN, NO_2, (C_1-C_8)alkyl, halo(C_1-C_8)alkyl, (C_1-C_8)alkoxy, halo(C_1-C_8)alkoxy, S(C_1-C_8)alkyl, S(halo(C_1-C_8)alkyl), S(O)(C_1-C_8)alkyl, S(O)(halo(C_1-C_8)alkyl), $S(O)_2$(C_1-C_8)alkyl, $S(O)_2$(halo(C_1-C_8)alkyl), N(R14)(R15),

(2) **substituted (C_1-C_8)alkyl**, wherein said substituted (C_1-C_8)alkyl has one or more substituents selected from CN and NO_2,

(3) **substituted halo(C_1-C_8)alkyl**, wherein said substituted halo(C_1-C_8)alkyl, has one or more substituents selected from CN and NO_2,

(4) **substituted (C_1-C_8)alkoxy**, wherein said substituted (C_1-C_8)alkoxy has one or more substituents selected from CN and NO_2, and

(5) **substituted halo(C_1-C_8)alkoxy**, wherein said substituted halo(C_1-C_8)alkoxy has one or more substituents selected from CN and NO_2;

(e) **R5** is selected from

(1) H, F, Cl, Br, I, CN, NO_2, (C_1-C_8)alkyl, halo(C_1-C_8)alkyl, (C_1-C_8)alkoxy, halo(C_1-C_8)alkoxy, S(C_1-C_8)alkyl, S(halo(C_1-C_8)alkyl), S(O)(C_1-C_8)alkyl, S(O)(halo(C_1-C_8)alkyl), $S(O)_2$(C_1-C_8)alkyl, $S(O)_2$(halo(C_1-C_8)alkyl), N(R14)(R15),

(2) **substituted (C_1-C_8)alkyl**, wherein said substituted (C_1-C_8)alkyl has one or more substituents selected from CN and NO_2,

(3) **substituted halo(C_1-C_8)alkyl**, wherein said substituted halo(C_1-C_8)alkyl, has one or more substituents selected from CN and NO_2,

(4) **substituted (C_1-C_8)alkoxy**, wherein said substituted (C_1-C_8)alkoxy has one or more substituents selected from CN and NO_2, and

(5) **substituted halo(C_1-C_8)alkoxy**, wherein said substituted halo(C_1-C_8)alkoxy has one or more substituents selected from CN and NO_2;

(f) **R6** is a (C_1-C_8)haloalkyl;

(g) **R7** is selected from H, F, Cl, Br, I, OH, (C_1-C_8)alkoxy, and halo(C_1-C_8)alkoxy;

(h) **R8** is selected from H, (C_1-C_8)alkyl, halo(C_1-C_8)alkyl, OR14, and N(R14)(R15);

(i) **R9** is selected from H, F, Cl, Br, I, $(C_1$-$C_8)$alkyl, halo$(C_1$-$C_8)$alkyl, $(C_1$-$C_8)$alkoxy, halo$(C_1$-$C_8)$alkoxy, OR14, and N(R14)(R15);

(j) **R10** is selected from

(1) H, F, Cl, Br, I, CN, NO_2, $(C_1$-$C_8)$alkyl, halo$(C_1$-$C_8)$alkyl, $(C_1$-$C_8)$alkoxy, halo$(C_1$-$C_8)$alkoxy, cyclo$(C_3$-$C_6)$alkyl, S$(C_1$-$C_8)$alkyl, S(halo$(C_1$-$C_8)$alkyl), S(O)$(C_1$-$C_8)$alkyl, S(O)(halo$(C_1$-$C_8)$alkyl), $S(O)_2(C_1$-$C_8)$alkyl, $S(O)_2$(halo$(C_1$-$C_8)$alkyl), NR14R15, C(=O)H, C(=O)N(R14)(R15), CN(R14)(R15)(=NOH), (C=O)O$(C_1$-$C_8)$alkyl, (C=O)OH, heterocyclyl, $(C_2$-$C_8)$alkenyl, halo$(C_2$-$C_8)$alkenyl, $(C_2$-$C_8)$alkynyl,

(2) **substituted $(C_1$-$C_8)$alkyl**, wherein said substituted $(C_1$-$C_8)$alkyl has one or more substituents selected from OH, $(C_1$-$C_8)$alkoxy, S$(C_1$-$C_8)$alkyl, S(O)$(C_1$-$C_8)$alkyl, $S(O)_2(C_1$-$C_8)$alkyl, NR14R15, and

(3) **substituted halo$(C_1$-$C_8)$alkyl**, wherein said substituted halo$(C_1$-$C_8)$alkyl, has one or more substituents selected from $(C_1$-$C_8)$alkoxy, S$(C_1$-$C_8)$alkyl, S(O)$(C_1$-$C_8)$alkyl, $S(O)_2(C_1$-$C_8)$alkyl, and N(R14)(R15);

(k) **R11** is selected from C(=X5)N(R14)((C_1-$C_8)$alkylC(=X5)N(R14)(R15)) wherein each X5 is independently selected from O, or S;

(l) **R12** is selected from **(v)**, H, F, Cl, Br, I, CN, $(C_1$-$C_8)$alkyl, halo$(C_1$-$C_8)$alkyl, $(C_1$-$C_8)$alkoxy, halo$(C_1$-$C_8)$alkoxy, and cyclo$(C_3$-$C_6)$alkyl;

(m) **R13** is selected from **(v)**, H, F, Cl, Br, I, CN, $(C_1$-$C_8)$alkyl, halo$(C_1$-$C_8)$alkyl, $(C_1$-$C_8)$alkoxy, and halo$(C_1$-$C_8)$alkoxy;

(n) **each R14** is independently selected from H, $(C_1$-$C_8)$alkyl, $(C_2$-$C_8)$alkenyl, substituted $(C_1$-$C_8)$alkyl, halo$(C_1$-$C_8)$alkyl, substituted halo$(C_1$-$C_8)$alkyl), $(C_1$-$C_8)$alkoxy, cyclo$(C_3$-$C_6)$alkyl, aryl, substituted-aryl, $(C_1$-$C_8)$alkyl-aryl, $(C_1$-$C_8)$alkyl-(substituted-aryl), O-$(C_1$-$C_8)$alkyl-aryl, O-$(C_1$-$C_8)$alkyl-(substituted-aryl), heterocyclyl, substituted-heterocyclyl, $(C_1$-$C_8)$alkyl-heterocyclyl, $(C_1$-$C_8)$alkyl-(substituted-heterocyclyl), O-$(C_1$-$C_8)$alkyl-heterocyclyl, O-$(C_1$-$C_8)$alkyl-(substituted-heterocyclyl), N(R16)(R17), $(C_1$-$C_8)$alkyl-C(=O)N(R16)(R17), C(=O)$(C_1$-$C_8)$alkyl, C(=O)(halo$(C_1$-$C_8)$alkyl),C(=O)$(C_3$-$C_6)$cycloalkyl, $(C_1$-$C_8)$alkyl-C(=O)O$(C_1$-$C_8)$alkyl, C(=O)H

wherein each said substituted $(C_1$-$C_8)$alkyl has one or more substituents selected from CN, and NO_2,

wherein each said substituted halo$(C_1$-$C_8)$alkyl), has one or more substituents selected from CN, and NO_2,

wherein each said substituted-aryl has one or more substituents selected from F, Cl, Br, I, CN, NO_2, $(C_1$-$C_8)$alkyl, halo$(C_1$-$C_8)$alkyl, $(C_1$-$C_8)$alkoxy, halo$(C_1$-$C_8)$alkoxy,

$S(C_1\text{-}C_8)$alkyl, S(halo$(C_1\text{-}C_8)$alkyl), N($(C_1\text{-}C_8)$alkyl)$_2$ (wherein each $(C_1\text{-}C_8)$alkyl is independently selected), and oxo, and

wherein each said substituted-heterocyclyl has one or more substituents selected from F, Cl, Br, I, CN, NO_2, $(C_1\text{-}C_8)$alkyl, halo$(C_1\text{-}C_8)$alkyl, $(C_1\text{-}C_8)$alkoxy, halo$(C_1\text{-}C_8)$alkoxy, $(C_3\text{-}C_6)$cycloalkyl S$(C_1\text{-}C_8)$alkyl, S(halo$(C_1\text{-}C_8)$alkyl), N($(C_1\text{-}C_8)$alkyl)$_2$ (wherein each $(C_1\text{-}C_8)$alkyl is independently selected), heterocyclyl, C(=O)$(C_1\text{-}C_8)$alkyl, C(=O)O$(C_1\text{-}C_8)$alkyl, and oxo, (wherein said alkyl, alkoxy, and heterocyclyl, may be further substituted with one or more of F, Cl, Br, I, CN, and NO_2);

(o) each R15 is independently selected from H, $(C_1\text{-}C_8)$alkyl, $(C_2\text{-}C_8)$alkenyl, substituted $(C_1\text{-}C_8)$alkyl, halo$(C_1\text{-}C_8)$alkyl, substituted halo$(C_1\text{-}C_8)$alkyl), $(C_1\text{-}C_8)$alkoxy, cyclo$(C_3\text{-}C_6)$alkyl, aryl, substituted-aryl, $(C_1\text{-}C_8)$alkyl-aryl, $(C_1\text{-}C_8)$alkyl-(substituted-aryl), O-$(C_1\text{-}C_8)$alkyl-aryl, O-$(C_1\text{-}C_8)$alkyl-(substituted-aryl), heterocyclyl, substituted-heterocyclyl, $(C_1\text{-}C_8)$alkyl-heterocyclyl, $(C_1\text{-}C_8)$alkyl-(substituted-heterocyclyl), O-$(C_1\text{-}C_8)$alkyl-heterocyclyl, O-$(C_1\text{-}C_8)$alkyl-(substituted-heterocyclyl), N(R16)(R17), $(C_1\text{-}C_8)$alkyl-C(=O)N(R16)(R17), C(=O)$(C_1\text{-}C_8)$alkyl, C(=O)(halo$(C_1\text{-}C_8)$alkyl), C(=O)$(C_3\text{-}C_6)$cycloalkyl, $(C_1\text{-}C_8)$alkyl-C(=O)O$(C_1\text{-}C_8)$alkyl, C(=O)H

wherein each said substituted $(C_1\text{-}C_8)$alkyl has one or more substituents selected from CN, and NO_2,

wherein each said substituted halo$(C_1\text{-}C_8)$alkyl), has one or more substituents selected from CN, and NO_2,

wherein each said substituted-aryl has one or more substituents selected from F, Cl, Br, I, CN, NO_2, $(C_1\text{-}C_8)$alkyl, halo$(C_1\text{-}C_8)$alkyl, $(C_1\text{-}C_8)$alkoxy, halo$(C_1\text{-}C_8)$alkoxy, S$(C_1\text{-}C_8)$alkyl, S(halo$(C_1\text{-}C_8)$alkyl), N($(C_1\text{-}C_8)$alkyl)$_2$ (wherein each $(C_1\text{-}C_8)$alkyl is independently selected), and oxo, and

wherein each said substituted-heterocyclyl has one or more substituents selected from F, Cl, Br, I, CN, NO_2, $(C_1\text{-}C_8)$alkyl, halo$(C_1\text{-}C_8)$alkyl, $(C_1\text{-}C_8)$alkoxy, halo$(C_1\text{-}C_8)$alkoxy, $(C_3\text{-}C_6)$cycloalkyl S$(C_1\text{-}C_8)$alkyl, S(halo$(C_1\text{-}C_8)$alkyl), N($(C_1\text{-}C_8)$alkyl)$_2$ (wherein each $(C_1\text{-}C_8)$alkyl is independently selected), heterocyclyl, C(=O)$(C_1\text{-}C_8)$alkyl, C(=O)O$(C_1\text{-}C_8)$alkyl, and oxo, (wherein said alkyl, alkoxy, and heterocyclyl, may be further substituted with one or more of F, Cl, Br, I, CN, and NO_2);

(p) each R16 is independently selected from H, $(C_1\text{-}C_8)$alkyl, substituted-$(C_1\text{-}C_8)$alkyl, halo$(C_1\text{-}C_8)$alkyl, substituted-halo$(C_1\text{-}C_8)$alkyl, cyclo$(C_3\text{-}C_6)$alkyl, aryl, substituted-aryl, $(C_1\text{-}C_8)$alkyl-aryl, $(C_1\text{-}C_8)$alkyl-(substituted-aryl), O-$(C_1\text{-}C_8)$alkyl-aryl, O-$(C_1\text{-}C_8)$alkyl-(substituted-aryl), heterocyclyl, substituted-heterocyclyl, $(C_1\text{-}C_8)$alkyl-heterocyclyl, $(C_1$-

C_8)alkyl-(substituted-heterocyclyl), O-(C_1-C_8)alkyl-heterocyclyl, O-(C_1-C_8)alkyl-(substituted-heterocyclyl), O-(C_1-C_8)alkyl

wherein each said substituted (C_1-C_8)alkyl has one or more substituents selected from CN, and NO_2,

wherein each said substituted halo(C_1-C_8)alkyl), has one or more substituents selected from CN, and NO_2,

wherein each said substituted-aryl has one or more substituents selected from F, Cl, Br, I, CN, NO_2, (C_1-C_8)alkyl, halo(C_1-C_8)alkyl, (C_1-C_8)alkoxy, halo(C_1-C_8)alkoxy, S(C_1-C_8)alkyl, S(halo(C_1-C_8)alkyl), N((C_1-C_8)alkyl)$_2$ (wherein each (C_1-C_8)alkyl is independently selected), and oxo, and

wherein each said substituted-heterocyclyl has one or more substituents selected from F, Cl, Br, I, CN, NO_2, (C_1-C_8)alkyl, halo(C_1-C_8)alkyl, (C_1-C_8)alkoxy, halo(C_1-C_8)alkoxy, S(C_1-C_8)alkyl, S(halo(C_1-C_8)alkyl), N((C_1-C_8)alkyl)$_2$ (wherein each (C_1-C_8)alkyl is independently selected), and oxo;

(q) **each R17** is independently selected from H, (C_1-C_8)alkyl, substituted-(C_1-C_8)alkyl, halo(C_1-C_8)alkyl, substituted-halo(C_1-C_8)alkyl, cyclo(C_3-C_6)alkyl, aryl, substituted-aryl, (C_1-C_8)alkyl-aryl, (C_1-C_8)alkyl-(substituted-aryl), O-(C_1-C_8)alkyl-aryl, O-(C_1-C_8)alkyl-(substituted-aryl), heterocyclyl, substituted-heterocyclyl, (C_1-C_8)alkyl-heterocyclyl, (C_1-C_8)alkyl-(substituted-heterocyclyl), O-(C_1-C_8)alkyl-heterocyclyl, O-(C_1-C_8)alkyl-(substituted-heterocyclyl), O-(C_1-C_8)alkyl

wherein each said substituted (C_1-C_8)alkyl has one or more substituents selected from CN, and NO_2,

wherein each said substituted halo(C_1-C_8)alkyl), has one or more substituents selected from CN, and NO_2,

wherein each said substituted-aryl has one or more substituents selected from F, Cl, Br, I, CN, NO_2, (C_1-C_8)alkyl, halo(C_1-C_8)alkyl, (C_1-C_8)alkoxy, halo(C_1-C_8)alkoxy, S(C_1-C_8)alkyl, S(halo(C_1-C_8)alkyl), N((C_1-C_8)alkyl)$_2$ (wherein each (C_1-C_8)alkyl is independently selected), and oxo, and

wherein each said substituted-heterocyclyl has one or more substituents selected from F, Cl, Br, I, CN, NO_2, (C_1-C_8)alkyl, halo(C_1-C_8)alkyl, (C_1-C_8)alkoxy, halo(C_1-C_8)alkoxy, S(C_1-C_8)alkyl, S(halo(C_1-C_8)alkyl), N((C_1-C_8)alkyl)$_2$ (wherein each (C_1-C_8)alkyl is independently selected), and oxo;

(r) **X1** is selected from N and CR12;

(s) **X2** is selected from N, CR9, and CR13;

(t) **X3** is selected from N and CR9; and

(v) R12 and R13 together form a linkage containing 3 to 4 atoms selected from C, N, O, and S, wherein said linkage connects back to the ring to form a 5 to 6 member saturated or unsaturated cyclic ring, wherein said linkage has at least one substituent X4 wherein X4 is selected from R14, N(R14)(R15), N(R14)(C(=O)R14), N(R14)(C(=S)R14), N(R14)(C(=O)N(R14)(R14)), N(R14)(C(=S)N(R14)(R14)), N(R14)(C(=O)N(R14)((C_2-C_8)alkenyl)), N(R14)(C(=S)N(R14)((C_2-C_8)alkenyl)), wherein each R14 is independently selected.

In another embodiment of this invention R1 may be selected from any combination of one or more of the following - H, F, Cl, Br, I, CN, NO_2, methyl, ethyl, (C_3)alkyl, (C_4)alkyl, (C_5)alkyl, (C_6)alkyl, (C_7)alkyl, (C_8)alkyl, halomethyl, haloethyl, halo(C_3)alkyl, halo(C_4)alkyl, halo(C_5)alkyl, halo(C_6)alkyl, halo(C_7)alkyl, halo(C_8)alkyl, methoxy, ethoxy, (C_3)alkoxy, (C_4)alkoxy, (C_5)alkoxy, (C_6)alkoxy, (C_7)alkoxy, (C_8)alkoxy, halomethoxy, haloethoxy, halo(C_3)alkoxy, halo(C_4)alkoxy, halo(C_5)alkoxy, halo(C_6)alkoxy, halo(C_7)alkoxy, and halo(C_8)alkoxy.

In another embodiment of this invention R2 may be selected from any combination of one or more of the following - H, F, Cl, Br, I, CN, NO_2, methyl, ethyl, (C_3)alkyl, (C_4)alkyl, (C_5)alkyl, (C_6)alkyl, (C_7)alkyl, (C_8)alkyl, halomethyl, haloethyl, halo(C_3)alkyl, halo(C_4)alkyl, halo(C_5)alkyl, halo(C_6)alkyl, halo(C_7)alkyl, halo(C_8)alkyl, methoxy, ethoxy, (C_3)alkoxy, (C_4)alkoxy, (C_5)alkoxy, (C_6)alkoxy, (C_7)alkoxy, (C_8)alkoxy, halomethoxy, haloethoxy, halo(C_3)alkoxy, halo(C_4)alkoxy, halo(C_5)alkoxy, halo(C_6)alkoxy, halo(C_7)alkoxy, and halo(C_8)alkoxy.

In another embodiment of this invention R3 may be selected from any combination of one or more of the following - H, F, Cl, Br, I, CN, NO_2, methyl, ethyl, (C_3)alkyl, (C_4)alkyl, (C_5)alkyl, (C_6)alkyl, (C_7)alkyl, (C_8)alkyl, halomethyl, haloethyl, halo(C_3)alkyl, halo(C_4)alkyl, halo(C_5)alkyl, halo(C_6)alkyl, halo(C_7)alkyl, halo(C_8)alkyl, methoxy, ethoxy, (C_3)alkoxy, (C_4)alkoxy, (C_5)alkoxy, (C_6)alkoxy, (C_7)alkoxy, (C_8)alkoxy, halomethoxy, haloethoxy, halo(C_3)alkoxy, halo(C_4)alkoxy, halo(C_5)alkoxy, halo(C_6)alkoxy, halo(C_7)alkoxy, and halo(C_8)alkoxy.

In another embodiment of this invention R4 may be selected from any combination of one or more of the following - H, F, Cl, Br, I, CN, NO_2, methyl, ethyl, (C_3)alkyl, (C_4)alkyl, (C_5)alkyl, (C_6)alkyl, (C_7)alkyl, (C_8)alkyl, halomethyl, haloethyl, halo(C_3)alkyl, halo(C_4)alkyl, halo(C_5)alkyl, halo(C_6)alkyl, halo(C_7)alkyl, halo(C_8)alkyl, methoxy, ethoxy, (C_3)alkoxy, (C_4)alkoxy, (C_5)alkoxy, (C_6)alkoxy, (C_7)alkoxy, (C_8)alkoxy, halomethoxy, haloethoxy,

halo(C_3)alkoxy, halo(C_4)alkoxy, halo(C_5)alkoxy, halo(C_6)alkoxy, halo(C_7)alkoxy, and halo(C_8)alkoxy.

In another embodiment of this invention R5 may be selected from any combination of one or more of the following - H, F, Cl, Br, I, CN, NO_2, methyl, ethyl, (C_3)alkyl, (C_4)alkyl, (C_5)alkyl, (C_6)alkyl, (C_7)alkyl, (C_8)alkyl, halomethyl, haloethyl, halo(C_3)alkyl, halo(C_4)alkyl, halo(C_5)alkyl, halo(C_6)alkyl, halo(C_7)alkyl, halo(C_8)alkyl, methoxy, ethoxy, (C_3)alkoxy, (C_4)alkoxy, (C_5)alkoxy, (C_6)alkoxy, (C_7)alkoxy, (C_8)alkoxy, halomethoxy, haloethoxy, halo(C_3)alkoxy, halo(C_4)alkoxy, halo(C_5)alkoxy, halo(C_6)alkoxy, halo(C_7)alkoxy, and halo(C_8)alkoxy.

In another embodiment of this invention R2 and R4 are selected from F, Cl, Br, I, CN, and NO_2 and R1, R3, and R5 are H.

In another embodiment of this invention R2, R3, and R4 are selected from F, Cl, Br, I, CN, and NO_2 and R1, and R5 are H.

In another embodiment of this invention R2, R3, and R4 are independently selected from F and Cl and R1and R5 are H.

In another embodiment of this invention R1 is selected from Cl and H.

In another embodiment of this invention R2 is selected from CF_3, CH_3, Cl, F, and H.

In another embodiment of this invention R3 is selected from OCH_3, CH_3, F, Cl, or H.

In another embodiment of this invention R4 is selected from CF_3, CH_3, Cl, F, and H.

In another embodiment of this invention R5 is selected from F, Cl, and H.

In another embodiment of this invention R6 may be selected from any combination of one or more of the following - halomethyl, haloethyl, halo(C_3)alkyl, halo(C_4)alkyl, halo(C_5)alkyl, halo(C_6)alkyl, halo(C_7)alkyl, and halo(C_8)alkyl.

In another embodiment of this invention R6 is trifluoromethyl.

In another embodiment of this invention R7 may be selected from any combination of one or more of the following - H, F, Cl, Br, and I.

In another embodiment of this invention R7 is selected from H, OCH_3, and OH.

In another embodiment of this invention R8 may be selected from any combination of one or more of the following - H, methyl, ethyl, (C_3)alkyl, (C_4)alkyl, (C_5)alkyl, (C_6)alkyl, (C_7)alkyl, (C_8)alkyl, halomethyl, haloethyl, halo(C_3)alkyl, halo(C_4)alkyl, halo(C_5)alkyl, halo(C_6)alkyl, halo(C_7)alkyl, and halo(C_8)alkyl.

In another embodiment of this invention R8 is selected from CH_3 and H.

In another embodiment of this invention R9 may be selected from any combination of one or more of the following - H, F, Cl, Br, I, methyl, ethyl, (C_3)alkyl, (C_4)alkyl, (C_5)alkyl,

(C_6)alkyl, (C_7)alkyl, (C_8)alkyl, halomethyl, haloethyl, halo(C_3)alkyl, halo(C_4)alkyl, halo(C_5)alkyl, halo(C_6)alkyl, halo(C_7)alkyl, halo(C_8)alkyl, methoxy, ethoxy, (C_3)alkoxy, (C_4)alkoxy, (C_5)alkoxy, (C_6)alkoxy, (C_7)alkoxy, (C_8)alkoxy, halomethoxy, haloethoxy, halo(C_3)alkoxy, halo(C_4)alkoxy, halo(C_5)alkoxy, halo(C_6)alkoxy, halo(C_7)alkoxy, and halo(C_8)alkoxy.

In another embodiment of this invention R10 may be selected from any combination of one or more of the following – H, F, Cl, Br, I, CN, methyl, ethyl, (C_3)alkyl, (C_4)alkyl, (C_5)alkyl, (C_6)alkyl, (C_7)alkyl, (C_8)alkyl, halomethyl, haloethyl, halo(C_3)alkyl, halo(C_4)alkyl, halo(C_5)alkyl, halo(C_6)alkyl, halo(C_7)alkyl, halo(C_8)alkyl, methoxy, ethoxy, (C_3)alkoxy, (C_4)alkoxy, (C_5)alkoxy, (C_6)alkoxy, (C_7)alkoxy, (C_8)alkoxy, halomethoxy, haloethoxy, halo(C_3)alkoxy, halo(C_4)alkoxy, halo(C_5)alkoxy, halo(C_6)alkoxy, halo(C_7)alkoxy, halo(C_8)alkoxy, cyclopropyl, cyclobutyl, cyclopentyl, and cyclohexyl.

In another embodiment of this invention R10 may be selected from any combination of one or more of the following - H, Cl, Br, CH_3, and CF_3.

In another embodiment of this invention R10 is selected from Br, C(=NOH)NH_2, C(=O)H, C(=O)NH_2, C(=O)OCH_2CH_3, C(=O)OH, CF_3, CH_2CH_3, CH_2OH, CH3, Cl, CN, F, H, NH_2, NHC(=O)H, $NHCH_3$, NO_2, OCH_3, $OCHF_2$, and pyridyl.

In another embodiment of this invention R11 may be selected from any combination of one or more of the following – C(=O)N(H)(C((CH_3)$_2$)C(=O)N(H)(CH_2CF_3)), C(=O)N(H)(CH(CH_3)C(=O)N(H)(CH_2CF_3)), C(=O)N(H)(CH(CH_2CH_3)C(=O)N(H)(CH_2CF_3)), C(=O)N(H)(CH(CH_3)C(=S)N(H)(CH_2CF_3)), C(=O)N(H)(C((CH_3)$_2$)C(=S)N(H)(CH_2CF_3)), and C(=S)N(H)(C((CH_3)$_2$)C(=S)N(H)(CH_2CF_3)).

In another embodiment of this invention R11 is C(=(O or S))N(H)((((C_1-C_8)alkyl)C(=(O or S))N(H)(halo(C_1-C_8)alkyl)), which may be used in combination with any embodiment of R1 through R10 and X1 through X3.

In another embodiment of this invention R12 may be selected from any combination of one or more of the following - H, F, Cl, Br, I, methyl, ethyl, (C_3)alkyl, (C_4)alkyl, (C_5)alkyl, (C_6)alkyl, (C_7)alkyl, (C_8)alkyl, halomethyl, haloethyl, halo(C_3)alkyl, halo(C_4)alkyl, halo(C_5)alkyl, halo(C_6)alkyl, halo(C_7)alkyl, halo(C_8)alkyl, halomethoxy, haloethoxy, halo(C_3)alkoxy, halo(C_4)alkoxy, halo(C_5)alkoxy, halo(C_6)alkoxy, halo(C_7)alkoxy, and halo(C_8)alkoxy.

In another embodiment of this invention R12 is selected from CH3, and H.

In another embodiment of this invention R13 may be selected from any combination of one or more of the following - H, F, Cl, Br, I, methyl, ethyl, (C_3)alkyl, (C_4)alkyl, (C_5)alkyl, (C_6)alkyl, (C_7)alkyl, (C_8)alkyl, halomethyl, haloethyl, halo(C_3)alkyl, halo(C_4)alkyl, halo(C_5)alkyl, halo(C_6)alkyl, halo(C_7)alkyl, halo(C_8)alkyl, halomethoxy, haloethoxy, halo(C_3)alkoxy, halo(C_4)alkoxy, halo(C_5)alkoxy, halo(C_6)alkoxy, halo(C_7)alkoxy, and halo(C_8)alkoxy.

In another embodiment of this invention R13 is selected from CH_3, Cl and H.

In another embodiment of this invention R12-R13 are a hydrocarbyl linkage containing CH=CHCH=CH.

In another embodiment of this invention R14 may be selected from any combination of one or more of the following – H, methyl, ethyl, (C_3)alkyl, (C_4)alkyl, (C_5)alkyl, (C_6)alkyl, (C_7)alkyl, (C_8)alkyl, halomethyl, haloethyl, halo(C_3)alkyl, halo(C_4)alkyl, halo(C_5)alkyl, halo(C_6)alkyl, halo(C_7)alkyl, halo(C_8)alkyl, methyl-aryl, ethyl-aryl, (C_3)alkyl-aryl, (C_4)alkyl-aryl, (C_5)alkyl-aryl, (C_6)alkyl-aryl, (C_7)alkyl-aryl, (C_8)alkyl-aryl, methyl-(substituted-aryl), ethyl-(substituted-aryl), (C_3)alkyl-(substituted-aryl), (C_4)alkyl-(substituted-aryl), (C_5)alkyl-(substituted-aryl), (C_6)alkyl-(substituted-aryl), (C_7)alkyl-(substituted-aryl), (C_8)alkyl-(substituted-aryl), O-methyl-aryl, O-ethyl-aryl, O-(C_3)alkyl-aryl, O-(C_4)alkyl-aryl, O-(C_5)alkyl-aryl, O-(C_6)alkyl-aryl, O-(C_7)alkyl-aryl, O-(C_8)alkyl-aryl, O-methyl-(substituted-aryl), O-ethyl-(substituted-aryl), O-(C_3)alkyl-(substituted-aryl), O-(C_4)alkyl-(substituted-aryl), O-(C_5)alkyl-(substituted-aryl), O-(C_6)alkyl-(substituted-aryl), O-(C_7)alkyl-(substituted-aryl), O-(C_8)alkyl-(substituted-aryl), methyl-heterocyclyl, ethyl-heterocyclyl, (C_3)alkyl-heterocyclyl, (C_4)alkyl-heterocyclyl, (C_5)alkyl-heterocyclyl, (C_6)alkyl-heterocyclyl, (C_7)alkyl-heterocyclyl, (C_8)alkyl-heterocyclyl, methyl-(substituted-heterocyclyl), ethyl-(substituted-heterocyclyl), (C_3)alkyl-(substituted-heterocyclyl), (C_4)alkyl-(substituted-heterocyclyl), (C_5)alkyl-(substituted-heterocyclyl), (C_6)alkyl-(substituted-heterocyclyl), (C_7)alkyl-(substituted-heterocyclyl), (C_8)alkyl-(substituted-heterocyclyl), O-methyl-heterocyclyl, O-ethyl-heterocyclyl, O-(C_3)alkyl-heterocyclyl, O-(C_4)alkyl-heterocyclyl, O-(C_5)alkyl-heterocyclyl, O-(C_6)alkyl-heterocyclyl, O-(C_7)alkyl-heterocyclyl, O-(C_8)alkyl-heterocyclyl, O-methyl-(substituted-heterocyclyl), O-ethyl-(substituted-heterocyclyl), O-(C_3)alkyl-(substituted-heterocyclyl), O-(C_4)alkyl-(substituted-heterocyclyl), O-(C_5)alkyl-(substituted-heterocyclyl), O-(C_6)alkyl-(substituted-heterocyclyl), O-(C_7)alkyl-(substituted-heterocyclyl), O-(C_8)alkyl-(substituted-heterocyclyl), methyl-C(=O)N(R16)(R17), ethyl-C(=O)N(R16)(R17), (C_3)alkyl-C(=O)N(R16)(R17), (C_4)alkyl-C(=O)N(R16)(R17), (C_5)alkyl-

C(=O)N(R16)(R17), (C_6)alkyl-C(=O)N(R16)(R17), (C_7)alkyl-C(=O)N(R16)(R17), and (C_8)alkyl-C(=O)N(R16)(R17).

In another embodiment of this invention R14 may be selected from any combination of one or more of the following – H, CH_3, CH_2CF_3, CH_2-halopyridyl, oxo-pyrrolidinyl, halophenyl, thietanyl, CH_2-phenyl, CH_2-pyridyl, thietanyl-dioxide, CH_2-halothiazolyl, C((CH_3)$_2$)-pyridyl, N(H)(halophenyl), CH_2-pyrimidinyl, CH_2-tetrahydrofuranyl, CH_2-furanyl, O-CH_2-halopyridyl, and CH_2C(=O)N(H)(CH_2CF_3).

In another embodiment of this invention R15 may be selected from any combination of one or more of the following - H, methyl, ethyl, (C_3)alkyl, (C_4)alkyl, (C_5)alkyl, (C_6)alkyl, (C_7)alkyl, (C_8)alkyl, halomethyl, haloethyl, halo(C_3)alkyl, halo(C_4)alkyl, halo(C_5)alkyl, halo(C_6)alkyl, halo(C_7)alkyl, halo(C_8)alkyl, methyl-aryl, ethyl-aryl, (C_3)alkyl-aryl, (C_4)alkyl-aryl, (C_5)alkyl-aryl, (C_6)alkyl-aryl, (C_7)alkyl-aryl, (C_8)alkyl-aryl, methyl-(substituted-aryl), ethyl-(substituted-aryl), (C_3)alkyl-(substituted-aryl), (C_4)alkyl-(substituted-aryl), (C_5)alkyl-(substituted-aryl), (C_6)alkyl-(substituted-aryl), (C_7)alkyl-(substituted-aryl), (C_8)alkyl-(substituted-aryl), O-methyl-aryl, O-ethyl-aryl, O-(C_3)alkyl-aryl, O-(C_4)alkyl-aryl, O-(C_5)alkyl-aryl, O-(C_6)alkyl-aryl, O-(C_7)alkyl-aryl, O-(C_8)alkyl-aryl, O-methyl-(substituted-aryl), O-ethyl-(substituted-aryl), O-(C_3)alkyl-(substituted-aryl), O-(C_4)alkyl-(substituted-aryl), O-(C_5)alkyl-(substituted-aryl), O-(C_6)alkyl-(substituted-aryl), O-(C_7)alkyl-(substituted-aryl), O-(C_8)alkyl-(substituted-aryl), methyl-heterocyclyl, ethyl-heterocyclyl, (C_3)alkyl-heterocyclyl, (C_4)alkyl-heterocyclyl, (C_5)alkyl-heterocyclyl, (C_6)alkyl-heterocyclyl, (C_7)alkyl-heterocyclyl, (C_8)alkyl-heterocyclyl, methyl-(substituted-heterocyclyl), ethyl-(substituted-heterocyclyl), (C_3)alkyl-(substituted-heterocyclyl), (C_4)alkyl-(substituted-heterocyclyl), (C_5)alkyl-(substituted-heterocyclyl), (C_6)alkyl-(substituted-heterocyclyl), (C_7)alkyl-(substituted-heterocyclyl), (C_8)alkyl-(substituted-heterocyclyl), O-methyl-heterocyclyl, O-ethyl-heterocyclyl, O-(C_3)alkyl-heterocyclyl, O-(C_4)alkyl-heterocyclyl, O-(C_5)alkyl-heterocyclyl, O-(C_6)alkyl-heterocyclyl, O-(C_7)alkyl-heterocyclyl, O-(C_8)alkyl-heterocyclyl, O-methyl-(substituted-heterocyclyl), O-ethyl-(substituted-heterocyclyl), O-(C_3)alkyl-(substituted-heterocyclyl), O-(C_4)alkyl-(substituted-heterocyclyl), O-(C_5)alkyl-(substituted-heterocyclyl), O-(C_6)alkyl-(substituted-heterocyclyl), O-(C_7)alkyl-(substituted-heterocyclyl), O-(C_8)alkyl-(substituted-heterocyclyl), methyl-C(=O)N(R16)(R17), ethyl-C(=O)N(R16)(R17), (C_3)alkyl-C(=O)N(R16)(R17), (C_4)alkyl-C(=O)N(R16)(R17), (C_5)alkyl-C(=O)N(R16)(R17), (C_6)alkyl-C(=O)N(R16)(R17), (C_7)alkyl-C(=O)N(R16)(R17), and (C_8)alkyl-C(=O)N(R16)(R17).

In another embodiment of this invention R15 may be selected from any combination of one or more of the following – H, CH_3, CH_2CF_3, CH_2-halopyridyl, oxo-pyrrolidinyl, halophenyl, thietanyl, CH_2-phenyl, CH_2-pyridyl, thietanyl-dioxide, CH_2-halothiazolyl, $C((CH_3)_2)$-pyridyl, N(H)(halophenyl), CH_2-pyrimidinyl, CH_2-tetrahydrofuranyl, CH_2-furanyl, O-CH_2-halopyridyl, and $CH_2C(=O)N(H)(CH_2CF_3)$.

In another embodiment of this invention R16 may be selected from any combination of one or more of the following – H, methyl, ethyl, (C_3)alkyl, (C_4)alkyl, (C_5)alkyl, (C_6)alkyl, (C_7)alkyl, (C_8)alkyl, halomethyl, haloethyl, halo(C_3)alkyl, halo(C_4)alkyl, halo(C_5)alkyl, halo(C_6)alkyl, halo(C_7)alkyl, halo(C_8)alkyl, methyl-aryl, ethyl-aryl, (C_3)alkyl-aryl, (C_4)alkyl-aryl, (C_5)alkyl-aryl, (C_6)alkyl-aryl, (C_7)alkyl-aryl, (C_8)alkyl-aryl, methyl-(substituted-aryl), ethyl-(substituted-aryl), (C_3)alkyl-(substituted-aryl), (C_4)alkyl-(substituted-aryl), (C_5)alkyl-(substituted-aryl), (C_6)alkyl-(substituted-aryl), (C_7)alkyl-(substituted-aryl), (C_8)alkyl-(substituted-aryl), O-methyl-aryl, O-ethyl-aryl, O-(C_3)alkyl-aryl, O-(C_4)alkyl-aryl, O-(C_5)alkyl-aryl, O-(C_6)alkyl-aryl, O-(C_7)alkyl-aryl, O-(C_8)alkyl-aryl, O-methyl-(substituted-aryl), O-ethyl-(substituted-aryl), O-(C_3)alkyl-(substituted-aryl), O-(C_4)alkyl-(substituted-aryl), O-(C_5)alkyl-(substituted-aryl), O-(C_6)alkyl-(substituted-aryl), O-(C_7)alkyl-(substituted-aryl), O-(C_8)alkyl-(substituted-aryl), methyl-heterocyclyl, ethyl-heterocyclyl, (C_3)alkyl-heterocyclyl, (C_4)alkyl-heterocyclyl, (C_5)alkyl-heterocyclyl, (C_6)alkyl-heterocyclyl, (C_7)alkyl-heterocyclyl, (C_8)alkyl-heterocyclyl, methyl-(substituted-heterocyclyl), ethyl-(substituted-heterocyclyl), (C_3)alkyl-(substituted-heterocyclyl), (C_4)alkyl-(substituted-heterocyclyl), (C_5)alkyl-(substituted-heterocyclyl), (C_6)alkyl-(substituted-heterocyclyl), (C_7)alkyl-(substituted-heterocyclyl), (C_8)alkyl-(substituted-heterocyclyl), O-methyl-heterocyclyl, O-ethyl-heterocyclyl, O-(C_3)alkyl-heterocyclyl, O-(C_4)alkyl-heterocyclyl, O-(C_5)alkyl-heterocyclyl, O-(C_6)alkyl-heterocyclyl, O-(C_7)alkyl-heterocyclyl, O-(C_8)alkyl-heterocyclyl, O-methyl-(substituted-heterocyclyl), O-ethyl-(substituted-heterocyclyl), O-(C_3)alkyl-(substituted-heterocyclyl), O-(C_4)alkyl-(substituted-heterocyclyl), O-(C_5)alkyl-(substituted-heterocyclyl), O-(C_6)alkyl-(substituted-heterocyclyl), O-(C_7)alkyl-(substituted-heterocyclyl), and O-(C_8)alkyl-(substituted-heterocyclyl).

In another embodiment of this invention R16 may be selected from any combination of one or more of the following – H, CH_2CF_3, cyclopropyl, thietanyl, thietanyl dioxide, and halophenyl.

In another embodiment of this invention R17 may be selected from any combination of one or more of the following - H, methyl, ethyl, (C_3)alkyl, (C_4)alkyl, (C_5)alkyl, (C_6)alkyl, (C_7)alkyl, (C_8)alkyl, halomethyl, haloethyl, halo(C_3)alkyl, halo(C_4)alkyl, halo(C_5)alkyl,

halo(C_6)alkyl, halo(C_7)alkyl, halo(C_8)alkyl, methyl-aryl, ethyl-aryl, (C_3)alkyl-aryl, (C_4)alkyl-aryl, (C_5)alkyl-aryl, (C_6)alkyl-aryl, (C_7)alkyl-aryl, (C_8)alkyl-aryl, methyl-(substituted-aryl), ethyl-(substituted-aryl), (C_3)alkyl-(substituted-aryl), (C_4)alkyl-(substituted-aryl), (C_5)alkyl-(substituted-aryl), (C_6)alkyl-(substituted-aryl), (C_7)alkyl-(substituted-aryl), (C_8)alkyl-(substituted-aryl), O-methyl-aryl, O-ethyl-aryl, O-(C_3)alkyl-aryl, O-(C_4)alkyl-aryl, O-(C_5)alkyl-aryl, O-(C_6)alkyl-aryl, O-(C_7)alkyl-aryl, O-(C_8)alkyl-aryl, O-methyl-(substituted-aryl), O-ethyl-(substituted-aryl), O-(C_3)alkyl-(substituted-aryl), O-(C_4)alkyl-(substituted-aryl), O-(C_5)alkyl-(substituted-aryl), O-(C_6)alkyl-(substituted-aryl), O-(C_7)alkyl-(substituted-aryl), O-(C_8)alkyl-(substituted-aryl), methyl-heterocyclyl, ethyl-heterocyclyl, (C_3)alkyl-heterocyclyl, (C_4)alkyl-heterocyclyl, (C_5)alkyl-heterocyclyl, (C_6)alkyl-heterocyclyl, (C_7)alkyl-heterocyclyl, (C_8)alkyl-heterocyclyl, methyl-(substituted-heterocyclyl), ethyl-(substituted-heterocyclyl), (C_3)alkyl-(substituted-heterocyclyl), (C_4)alkyl-(substituted-heterocyclyl), (C_5)alkyl-(substituted-heterocyclyl), (C_6)alkyl-(substituted-heterocyclyl), (C_7)alkyl-(substituted-heterocyclyl), (C_8)alkyl-(substituted-heterocyclyl), O-methyl-heterocyclyl, O-ethyl-heterocyclyl, O-(C_3)alkyl-heterocyclyl, O-(C_4)alkyl-heterocyclyl, O-(C_5)alkyl-heterocyclyl, O-(C_6)alkyl-heterocyclyl, O-(C_7)alkyl-heterocyclyl, O-(C_8)alkyl-heterocyclyl, O-methyl-(substituted-heterocyclyl), O-ethyl-(substituted-heterocyclyl), O-(C_3)alkyl-(substituted-heterocyclyl), O-(C_4)alkyl-(substituted-heterocyclyl), O-(C_5)alkyl-(substituted-heterocyclyl), O-(C_6)alkyl-(substituted-heterocyclyl), O-(C_7)alkyl-(substituted-heterocyclyl), and O-(C_8)alkyl-(substituted-heterocyclyl).

In another embodiment of this invention R17 may be selected from any combination of one or more of the following – H, CH_2CF_3, cyclopropyl, thietanyl, thietanyl dioxide, and halophenyl.

In another embodiment of this invention X1 is CR12, X2 is CR13, and X3 is CR9.


In another embodiment of this invention a heterocyclyl has preferably about 6 to 10 atoms in the ring structure, more preferably, 6 to 8 atoms.

The molecules of Formula One will generally have a molecular mass of about 100 Daltons to about 1200 Daltons. However, it is generally preferred if the molecular mass is from about 120 Daltons to about 900 Daltons, and it is even more generally preferred if the molecular mass is from about 140 Daltons to about 600 Daltons.

The benzyl alcohol of Formula IV, wherein R1, R2, R3, R4, R5, R6, and R7 are as previously disclosed, can be synthesized in two ways. One way, disclosed in step ***a*** of Scheme I, is by treatment of the ketone of Formula II, wherein R1, R2, R3, R4, R5, and R6 are as previously disclosed, with a reducing agent, such as sodium borohydride ($NaBH_4$),

under basic conditions, such as aqueous sodium hydroxide (NaOH), in a polar protic solvent, such as methyl alcohol (MeOH) at 0 °C. Alternatively, an aldehyde of Formula III, wherein R1, R2, R3, R4, R5, and R7 are as previously disclosed, is allowed to react with trifluorotrimethylsilane in the presence of a catalytic amount of tetrabutylammonium fluoride (TBAF) in a polar aprotic solvent, such as tetrahydrofuran (THF), as in step ***b*** of Scheme I. The compound of Formula IV can be transformed into the compound of Formula V, wherein Y is selected from Br, Cl or I, and R1, R2, R3, R4, R5, R6, and R7 are as previously disclosed, by reaction with a halogenating reagent, such as *N*-bromosuccinimide (NBS) and triethyl phosphite in a non-reactive solvent, such as dichloromethane (CH_2Cl_2) at reflux temperature to provide Y = Br, or such as thionyl chloride and pyridine in a hydrocarbon solvent, such as toluene at reflux temperature to provide Y = Cl, as in step ***c*** of Scheme I.

Scheme I




Formation of the styrene coupling partners can be accomplished as in Schemes II, III IV and V.

In Scheme II, a vinylbenzoic acid of Formula VI, wherein R11 is (C=O)OH and R8, R9, R10, R12, R13, X1, X2, and X3 are as previously disclosed, can be converted in two steps to the vinylbenzamide of Formula VIIa, wherein R11 is (C=O)N(R14)(R15), and R8, R9, R10, R12, R13, R14, R15, and X are as previously disclosed. As in step ***d*** of Scheme II, the benzoic acid of Formula VI is treated with oxalyl chloride in the presence of a catalytic amount of *N,N*-dimethylformamide (DMF) in a non-reactive solvent such as CH_2Cl_2 to form the acid chloride, which is subsequently allowed to react with an amine (HN(R14)(R15)), wherein R14 and R15 are as previously disclosed, in the presence of a base, such as

triethylamine (TEA), in a polar aprotic solvent, such as THF, to provide the vinyl benzamide of Formula VIIa, wherein R11 is (C=O)N(R14)(R15), and R8, R9, R10, R12, R13, R14, R15, X1, X2, and X3 are as previously disclosed, as in step ***e*** of Scheme II.


Scheme II



VI VIIa

In Schemes III and IV, a halobenzoic acid of Formula VIII, wherein R18 is Br or I, R11 is (C=O)OH and R9, R10, R12, R13, X1, X2, and X3 are as previously disclosed can be converted to a vinylbenzoic acid ester of Formula VIIb1 or Formula VIIb2, wherein R18 is Br or I, R11 is (C=O)O(C_1-C_6 alkyl), and R8, R9, R10, R12, R13, X1, X2, and X3 are as previously disclosed. In step ***f*** of Scheme III, the halobenzoic acid of Formula VIII, wherein R18 is Br, is treated with a base, such as *n*-butyllithium (*n*-BuLi), and DMF in a polar, aprotic solvent, such as THF, at a temperature of about -78 °C. The resulting formyl benzoic acid is allowed to react with an acid, such as sulfuric acid (H_2SO_4), in the presence of an alcohol, such as ethyl alcohol (EtOH), as in step ***g***, to provide the formyl benzoic acid ethyl ester of Formula IX, wherein R11 is (C=O)O(C_1-C_6 alkyl), and R9, R10, R12, R13, X1, X2, and X3 are as previously disclosed. The vinyl benzoic acid ester of Formula VIIb1 is accessed via reaction of the compounds of Formula IX, with a base, such as potassium carbonate (K_2CO_3), and methyl triphenyl phosphonium bromide in a polar aprotic solvent, such as 1,4-dioxane, at ambient temperature, as in step ***h*** of Scheme III.


Scheme III



VIII IX VIIb1

In step ***i*** of Scheme IV, the halobenzoic acid of Formula VIII, wherein R18 is Br, R11 is (C=O)OH, and R8, R9, R10, R12, R13, X1, X2, and X3 are as previously disclosed, is treated with di-*tert*-butyl dicarbonate in the presence of a base, such as TEA and a catalytic amount of 4-(dimethylamino)pyridine (DMAP) in a polar aprotic solvent, such as THF, at ambient temperature. The resulting benzoic acid *tert*-butyl ester is allowed to react with vinyl boronic anhydride pyridine complex in the presence of a palladium catalyst, such a tetrakis(triphenylphospine)palladium(0) ($Pd(PPh_3)_4$), and a base, such as K_2CO_3, in a non-reactive solvent such as toluene at reflux temperature, as in step ***j***, to provide the vinyl benzoic acid ester of Formula VIIb2, wherein R11 is (C=O)O(C_1-C_6 alkyl), and R8, R9, R10, R12, R13, X1, X2, and X3 are as previously disclosed.


Scheme IV



VIII VIIb2

In step ***k*** of Scheme V, the vinyl benzoic acid ester of Formula VIIb2, wherein R10 is Br, R11 is (C=O)O(C_1-C_6 alkyl), and R8, R9, R12, R13, X1, X2, and X3 are as previously defined, can be further transformed into the corresponding vinyl benzoic acid ester of Formula VIIb3, wherein R10 is CN, R11 is (C=O)O(C_1-C_6 alkyl), and R8, R9, R12, R13, X1, X2, and X3 are as previously disclosed, by reaction with copper(I) cyanide (CuCN) in a polar aprotic solvent, such as DMF, at 140 °C.

Scheme V




VIIb2 VIIb3

Coupling of the compounds of Formula V with the compounds of Formula VIIa, VIIb1, VIIb2 and VIIb3 can be accomplished as in Schemes VI, VII, and VIII. In step ***l*** of Scheme VI, a compound of Formula V, wherein Y, R1, R2, R3, R4, R5, R6, and R7 are as previously disclosed, and the vinylbenzamide of Formula VIIa, wherein R11 is

(C=O)N(R14)(R15), and R8, R9, R10, R12, R13, R14, R15, X1, X2, and X3 are as previously disclosed, are allowed to react in the presence of copper(I) chloride (CuCl) and 2,2-bipyridyl in a solvent, such as 1,2-dichlorobenzene, at a temperature of about 180 °C to provide the molecules of Formula One, wherein R11 is (C=O)N(R14)(R15), and R1, R2, R3, R4, R5, R6, R7, R8, R9, R10, R12, R13, R14, R15, X1, X2, and X3 are as previously disclosed.

Scheme VI




V VIIa Formula One

In step ***l*** of Scheme VII, the compound of Formula V, wherein Y, R1, R2, R3, R4, R5, R6, and R7 are as previously disclosed, and the vinylbenzoic acid ester of Formula VIIb1, wherein R11 is (C=O)O(C_1-C_6 alkyl), and R8, R9, R10, R12, R13, X1, X2, and X3 are as previously disclosed, are allowed to react in the presence of CuCl and 2,2-bipyridyl in a solvent, such as 1,2-dichlorobenzene, at a temperature of about 180 °C to provide the compounds of Formula Xa, wherein R11 is (C=O)O(C_1-C_6 alkyl), and R1, R2, R3, R4, R5, R6, R7, R8, R9, R10, R12, R13, X1, X2, and X3 are as previously disclosed. The compounds of Formula Xa are then converted to the molecules of Formula One, wherein R11 is (C=O)N(R14)(R15), and R1, R2, R3, R4, R5, R6, R7, R8, R9, R10, R12, R13, R14, R15, X1, X2, and X3 are as previously disclosed, by either a two-step process as disclosed in steps ***m*** and ***n*** or in one step as disclosed in step ***o***. In step ***m*** of Scheme VII, the ester of Formula Xa is saponified to the corresponding acid under acidic conditions, such as about 11 Normal (N) hydrochloric acid (HCl), in a polar aprotic solvent, such as 1,4-dioxane, at about 100 °C. The acid can subsequently be coupled to an amine (HN(R14)(R15)), wherein R14 and R15 are as previously disclosed using peptide coupling reagents, such as 1-hydroxybenzotriazole (HOBt), *N*-(3-dimethylaminopropyl)-*N'*-ethyl-carbodiimide hydrochloride (EDC•HCl), benzotriazol-1-yl-oxytripyrrolidinophosphonium hexafluorophosphate (PyBOP), 2-chloro-1,3-dimethylimidazolidinium hexafluorophosphate (CIP), 1-hydroxy-7-azabenzotriazole (HOAt), or *O*-benzotriazole-*N,N,N',N'*-tetramethyl-uronium-hexafluoro-phosphate (HBTU)

in the presence of a base, such as *N,N*-diisopropylethylamine (DIPEA) or DMAP to give the molecules of Formula One, wherein R11 is (C=O)N(R14)(R15), and R1, R2, R3, R4, R5, R6, R7, R8, R9, R10, R12, R13, R14, R15, X1, X2, and X3 are as previously disclosed. Alternatively, the ester of Formula Xa is allowed to react with an amine (HN(R14)(R15)) in the presence of a solution of trimethylaluminum in toluene in a non-reactive solvent, such as CH_2Cl_2, at ambient temperature, as in step ***o*** of Scheme VII, to access the molecules of Formula One, wherein R11 is (C=O)N(R14)(R15), and R1, R2, R3, R4, R5, R6, R7, R8, R9, R10, R12, R13, R14, R15, X1, X2, and X3 are as previously disclosed.


Scheme VII




In step ***l*** of Scheme VIII, the compound of Formula V, wherein Y, R1, R2, R3, R4, R5, R6, and R7 are as previously disclosed, and the vinylbenzoic acid ester of Formula VIIb2 or VIIb3, wherein R11 is (C=O)O(C_1-C_6 alkyl), and R8, R9, R10, R12, R13, X1, X2, and X3 are as previously disclosed, are allowed to react in the presence of CuCl and 2,2-bipyridyl in a solvent, such as 1,2-dichlorobenzene, at a temperature of about 180 °C to provide the compounds of Formula Xb, wherein R11 is (C=O)OH, and R1, R2, R3, R4, R5, R6, R7, R8, R9, R10, R12, R13, R14, R15, X1, X2, and X3 are as previously disclosed. The compounds of Formula Xb are then converted to the molecules of Formula One, wherein R11 is (C=O)N(R14)(R15), and R1, R2, R3, R4, R5, R6, R7, R8, R9, R10, R12, R13, R14, R15, X1, X2, and X3 are as previously disclosed, in one step as disclosed in step ***n***. In step ***n*** of Scheme VIII, the acid of Formula Xb can be coupled to an amine (HN(R14)(R15)), wherein

R14 and R15 are as previously disclosed, using peptide coupling reagents, such as HOBt, EDC•HCl, PyBOP, CIP, HOAt, or HBTU in the presence of a base, such as DIPEA or DMAP to give the molecules of Formula One, wherein R11 is (C=O)N(R14)(R15), and R1, R2, R3, R4, R5, R6, R7, R8, R9, R10, R12, R13, R14, R15, X1, X2, and X3 are as previously disclosed.

Scheme VIII




V VIIb2 or VIIb3 Xb



Formula One

In step *t* of Scheme XIII, the vinyl benzyl chloride of Formula XIa, wherein R11 is -CH_2Cl and R8, R9, R10, R12, R13, X1, X2, and X3 are as previously defined, can be transformed into the corresponding phthalimide-protected benzyl amine of Formula XIIa, wherein R11 is CH_2N(Phthalimide), and R8, R9, R10, R12, R13, X1, X2, and X3 are as previously disclosed, by reaction with potassium phthalimide in a polar aprotic solvent, such as DMF, at 70 °C.

Scheme XIII




XIa XIIa

In step *u* of Scheme XIV, the 4-methylbenzonitrile of Formula XIIIa, wherein R11 is CH_3 and R9, R10, R12, R13, X1, X2, and X3 are as previously defined, can be transformed

into the corresponding benzyl bromide of Formula XIVa, wherein R11 is CH_2Br and R8, R9, R10, R12, R13, X1, X2, and X3 are as previously disclosed, by reaction with NBS and azobisisobutyronitrile (AIBN) in a non-reactive solvent, such as carbon tetrachloride (CCl_4) at 77 °C. The nitrile group (CN) of Formula XIVa can be reduced to the corresponding aldehyde of Formula XVa, wherein R11 is CH_2Br and R9, R10, R12, R13, X1, X2, and X3 are as previously defined via reaction with diisobutylaluminum hydride (DIBAL-H) in an aprotic solvent, such as toluene, at 0 °C, followed by quenching with 1.0 M HCl as in step ***v*** of Scheme XIV. The compound of Formula XVa can be further transformed to the corresponding phthalimide-protected benzyl amine of Formula XVIa, wherein R11 is CH_2N(Phthalimide) and R9, R10, R12, R13, X1, X2, and X3 are as previously disclosed, by reaction with potassium phthalimide in a polar aprotic solvent, such as DMF, at 60 °C as in step ***t*** of Scheme XIV. In step ***w*** of Scheme XIV, the aldehyde of Formula XVIa can be converted to the olefin of Formula XIIb, wherein R11 is CH_2N(Phthalimide) and R8, R9, R10, R12, R13, X1, X2, and X3 are as previously disclosed, by reaction with methyl triphenyl phosphonium bromide in a polar aprotic solvent, such as 1,4-dioxane, in the presence of a base, such as K_2CO_3, at ambient temperature.

Scheme XIV




XIIIa XIVa XVa

XVIa XIIb

The aldehyde of Formula XVa, wherein R11 is CH_2Br and R9, R10, R12, R13, X1, X2, and X3 are as previously defined, can be reacted with a nucleophile, such as 2-aminopyridine, in a polar aprotic solvent, such as *N,N*-dimethylacetamide (DMA), in the presence of a base, such as K_2CO_3, at ambient temperature to provide the compound of Formula XVII, wherein R11 is CH_2NH(2-pyridine) and R9, R10, R12, R13, X1, X2, and X3 are as previously disclosed, as in step ***x*** of Scheme XV. In step ***w*** of Scheme XV, the

compound of Formula XVII can be converted to the olefin of Formula XVIII, wherein R11 is CH_2NH(2-pyridine) and R8, R9, R10, R12, R13, X1, X2, and X3 are as previously disclosed.


Scheme XV



XVa XVII XVIII

In a two-step, one-pot reaction as in steps ***y*** and ***z*** of Scheme XVI, the compound of Formula XIX can be reacted with the compounds of Formula XX, wherein R10 and R11 are Cl, X1 is N, and R9, R13, X2, and X3 are as previously disclosed, in the presence of a base, such as sodium hydride (NaH), and a polar aprotic solvent, such as DMF, at ambient temperature to provide the compounds of Formula XXI, wherein R10 is Cl, R11 is $(CH)NH_2CO_2CH_2CH_3$, X1 is N, and R9, R13, X2, and X3 are as previously defined. Hydrolysis and decarboxylation of the compounds of Formula XXI can be accomplished by reaction under acidic conditions, such as with 3 N HCl, at reflux temperature, to afford the compounds of Formula XXII, wherein R10 is Cl, R11 is CH_2NH_2•HCl, X1 is N, and R9, R13, X2, and X3 are as previously disclosed, as in step ***aa*** in Scheme XVI. The compounds of Formula XXII can be further transformed to the corresponding phthalimide-protected benzyl amines of Formula XXIIIa, wherein R10 is Cl, R11 is CH_2N(Phthalimide), X1 is N, and R9, R13, X1, X2, and X3 are as previously disclosed, by reaction with phthalic anhydride in the presence of a base, such as TEA, and an aprotic solvent, such as toluene, at reflux temperature as in step ***ab*** of Scheme XVI. The bromide of Formula XXIIIa can be converted to the olefin of Formula XIIc, wherein R10 is Cl, R11 is CH_2N(Phthalimide), X1 is N, and R8, R9, R13, X2 and X3 are as previously disclosed, by reaction with vinyl boronic anhydride pyridine complex in the presence of a palladium catalyst, such as $Pd(PPh_3)_4$, and a base, such as K_2CO_3, in a non-reactive solvent such as toluene at reflux temperature, as in step ***ac*** of Scheme XVI.

Scheme XVI




XIX XX XXI

XXII XXIIIa XIIc

In step ***u*** of Scheme XVII, the 4-methylnaphthonitrile of Formula XIIIb, wherein X3 is CR9, R10 and X3 together form a linkage having 4 carbon atoms and with the ring carbon atoms form a 6-membered aromatic ring, R11 is CH_3, and R12, R13, X1 and X2 are as previously defined, can be transformed into the corresponding naphthyl bromide of Formula XIVb, wherein X3 is CR9, R10 and X3 together form a linkage having 4 carbon atoms and with the ring carbon atoms form a 6-membered aromatic ring, R11 is CH_2Br, and R12, R13, X1 and X2 are as previously disclosed, by reaction with NBS and AIBN in a non-reactive solvent, such as CCl_4 at 77 °C. The nitrile group (CN) of Formula XIVb can be reduced to the corresponding aldehyde of Formula XVb, wherein X3 is CR9, R10 and X3 together form a linkage having 4 carbon atoms and with the ring carbon atoms form a 6-membered aromatic ring (or if desired a non-aromatic ring), R11 is CH_2Br, and R12, R13, X1 and X2 are as previously defined via reaction with diisobutylaluminum hydride (DIBAL-H) in an aprotic solvent, such as toluene, at 0 °C, followed by quenching with 1.0 M HCl as in step ***v*** of Scheme XVII. The compound of Formula XVb can be further transformed to the corresponding phthalimide-protected benzyl amine of Formula XVIb, wherein X3 is CR9, R10 and X3 together form a linkage having 4 carbon atoms and with the ring carbon atoms form a 6-membered aromatic ring, R11 is CH_2N(Phthalimide), and R12, R13, X1 and X2 are as previously disclosed, by reaction with potassium phthalimide in a polar aprotic solvent, such as DMF, at 60 °C as in step ***t*** of Scheme XVII. In step ***w*** of Scheme XVII, the aldehyde of Formula XVIb can be converted to the olefin of Formula XIId, wherein X3 is CR9, R10 and X3 together form a linkage having 4 carbon atoms and with the ring carbon atoms form a 6-membered aromatic ring, R11 is CH_2N(Phthalimide), and R8, R12, R13, X1 and X2 are as previously disclosed, by reaction with methyl triphenyl phosphonium bromide in a polar


aprotic solvent, such as 1,4-dioxane, in the presence of a base, such as K_2CO_3, at ambient temperature.

Scheme XVII




The compound of Formula XXIV, wherein R11 is $NHNH_2$•HCl and R9, R10, R12, R13, X1, X2, and X3 are as previously disclosed, can be transformed into the corresponding phthalimide-protected hydrazine of Formula XXV, wherein R11 is NHN(Phthalimide) and R9, R10, R12, R13, X1, X2, and X3 are as previously disclosed, by reaction with phthalic anhydride in glacial acetic acid (AcOH) at reflux temperature as in step ***ad*** of Scheme XVIII. The bromide of Formula XXV can be converted to the olefin of Formula XIIe, wherein R11 is NHN(Phthalimide) and R8, R9, R10, R13, X1, X2 and X3 are as previously disclosed, by reaction with vinyl boronic anhydride pyridine complex in the presence of a palladium catalyst, such as $Pd(PPh_3)_4$, and a base, such as K_2CO_3, in a polar aprotic solvent such as 1,2-dimethoxyethane at 150 °C under microwave conditions, as in step ***ae*** of Scheme XVIII.

Scheme XVIII



In step ***af*** of Scheme XIX, the compound of Formula XXVI, wherein R11 is $B(OH)_2$, and R8, R9, R10, R12, R13, X1, X2, and X3 are as previously disclosed, are allowed to react with 2-hydroxyisoindoline-1,3-dione in the presence of CuCl and pyridine in a solvent, such as 1,2-dichlorobenzene, at ambient temperature to provide the compound of Formula XIIf, wherein R11 is ON(Phthalimide) and R8, R9, R10, R12, R13, X1, X2, and X3 are as previously disclosed.


Scheme XIX



XXVI XIIf

In step ***l*** of Scheme XX, the compound of Formula V, wherein Y, R1, R2, R3, R4, R5, R6, and R7 are as previously disclosed, and the compounds of Formula XIIa, wherein R11 is CH_2N(Phthalimide) and R8, R9, R10, R12, R13, X1, X2, and X3 are as previously disclosed, are allowed to react in the presence of CuCl and 2,2-bipyridyl in a solvent, such as 1,2-dichlorobenzene, at a temperature of about 180 °C to provide the corresponding compounds of Formula XXVIIa, wherein R11 is CH_2N(Phthalimide) and R1, R2, R3, R4, R5, R6, R7, R8, R9, R10, R12, R13, X1, X2, and X3 are as previously disclosed. The phthalimide protecting group in the compounds of Formula XXVIIa is removed as in step ***ag*** of Scheme XX by reaction with hydrazine hydrate in a polar protic solvent such as EtOH at 90 °C to provide the compounds of Formula XXVIIIa, wherein R11 is CH_2NH_2 and R1, R2, R3, R4, R5, R6, R7, R8, R9, R10, R12, R13, X1, X2, and X3 are as previously disclosed. The compounds of Formula XXVIIIa can be transformed into the compounds of Formula One, wherein R11 is $CH_2N(C{=}O)(R14)$ and R1, R2, R3, R4, R5, R6, R7, R8, R9, R10, R12, R13, X1, X2, and X3 are as previously disclosed, by acylation with an anhydride, such as acetic anhydride, and a base, such as TEA, in a non-reactive solvent such as CH_2Cl_2 at 0 °C as in step ***ah$_1$*** of Scheme XX.

Scheme XX



V XIIa XXVIIa

XXVIIIa Formula One

In step ***l*** of Scheme XXI, the compound of Formula V, wherein Y, R1, R2, R3, R4, R5, R6, and R7 are as previously disclosed, and the compounds of Formula XIIb, wherein R11 is CH_2N(Phthalimide) and R8, R9, R10, R12, R13, X1, X2, and X3 are as previously disclosed, are allowed to react in the presence of CuCl and 2,2-bipyridyl in a solvent, such as 1,2-dichlorobenzene, at a temperature of about 180 °C to provide the corresponding compounds of Formula XXVIIb, wherein R11 is CH_2N(Phthalimide) and R1, R2, R3, R4, R5, R6, R7, R8, R9, R10, R12, R13, X1, X2, and X3 are as previously disclosed. The phthalimide protecting group in the compounds of Formula XXVIIb is removed as in step ***ag*** of Scheme XXI by reaction with hydrazine hydrate in a polar protic solvent such as EtOH at 90 °C to provide the compounds of Formula XXVIIIb, wherein R11 is CH_2NH_2 and R1, R2, R3, R4, R5, R6, R7, R8, R9, R10, R12, R13, X1, X2, and X3 are as previously disclosed. The compounds of Formula XXVIIIb can be transformed into the compounds of Formula One, wherein R11 is $CH_2N(C=O)(R14)$ and R1, R2, R3, R4, R5, R6, R7, R8, R9, R10, R12, R13, X1, X2, and X3 are as previously disclosed, by reaction with an acid in the presence of $HOBt•H_2O$, EDC•HCl and a base, such as DIPEA, in a polar aprotic solvent, such as DMF, as in step ***ah$_{2a}$*** of Scheme XXI.

In another embodiment, the compounds of Formula XXVIIIb can be transformed into the compounds of Formula One, wherein R11 is $CH_2N(C=S)(R14)$ and R1, R2, R3, R4, R5, R6, R7, R8, R9, R10, R12, R13, X1, X2, and X3 are as previously disclosed, by reaction with a thioacid in the presence of $HOBt•H_2O$, EDC•HCl and a base, such as DIPEA, in a polar aprotic solvent, such as DMF, as in step ***ah$_2$*** of Scheme XXI.

In another embodiment, the compounds of Formula XXVIIIb can be transformed into the compounds of Formula One, wherein R11 is $CH_2N(C=O)N(R14)(R15)$ and R1, R2, R3, R4, R5, R6, R7, R8, R9, R10, R12, R13, X1, X2, and X3 are as previously disclosed, in two steps. The first step (step ***ah$_{3a}$*** of Scheme XXI) involves reaction with an aldehyde in a polar protic solvent such as MeOH, followed by reaction with $NaBH_4$. The second step (step ***ah$_{3b}$*** of Scheme XXI) involves acylation with an acid chloride, such as cyclopropylcarbonyl chloride, and a base, such as TEA, in a non-reactive solvent such as CH_2Cl_2 at ambient temperature of Scheme XXI.

In another embodiment, the compounds of Formula XXVIIIb can be transformed into the compounds of Formula One, wherein R11 is $CH_2N(C=O)N(R14)(R15)$ and R1, R2, R3, R4, R5, R6, R7, R8, R9, R10, R12, R13, X1, X2, and X3 are as previously disclosed, by reaction with an isocyanate (step ***ai$_1$*** of Scheme XXI) or a carbamoyl chloride (step ***ai$_2$*** of


Scheme XXI) in the presence of a base such as TEA and in a non-reactive solvent such as CH_2Cl_2 at 0 °C.

In another embodiment, the compounds of Formula XXVIIIb can be transformed into the compounds of Formula One, wherein R11 is $CH_2N(C=S)N(R14)(R15)$ and R1, R2, R3, R4, R5, R6, R7, R8, R9, R10, R12, R13, X1, X2, and X3 are as previously disclosed, by reaction with an isothiocyanate in the presence of a base such as TEA and in a non-reactive solvent such as CH_2Cl_2 at 0 °C, as in steps ***aj*** of Scheme XXI.

In another embodiment, the compounds of Formula XXVIIIb can be transformed into the compounds of Formula One, wherein R11 is $CH_2N(C=O)O(R14)$ and R1, R2, R3, R4, R5, R6, R7, R8, R9, R10, R12, R13, X1, X2, and X3 are as previously disclosed, by reaction with a dicarbonate, such as di-*tert*-butyl dicarbonate in the presence of a base such as TEA and in a non-reactive solvent such as CH_2Cl_2 at ambient temperature, as in steps ***ak*** of Scheme XXI.

In yet another embodiment, the compounds of Formula XXVIIIb can be transformed into the compounds of Formula One, wherein R11 is $CH_2N(C=O)(C=O)O(R14)$ and R1, R2, R3, R4, R5, R6, R7, R8, R9, R10, R12, R13, X1, X2, and X3 are as previously disclosed, by reaction with a chlorooxalic acid ester, such as 2-chloro-2-oxoacetate in the presence of a base such as TEA and in a non-reactive solvent such as CH_2Cl_2 at 0 °C, as in steps ***al*** of Scheme XXI.


Scheme XXI



In step ***l*** of Scheme XXII, the compound of Formula V, wherein Y, R1, R2, R3, R4, R5, R6, and R7 are as previously disclosed, and the compounds of Formula XIIc, wherein R10 is Cl, R11 is $CH_2N(Phthalimide)$, X1 is N, and R8, R9, R12, R13, X2, and X3 are as

previously disclosed, are allowed to react in the presence of CuCl and 2,2-bipyridyl in a solvent, such as 1,2-dichlorobenzene, at a temperature of about 180 °C to provide the corresponding compounds of Formula XXVIIc, wherein R10 is Cl, R11 is CH_2N(Phthalimide), X1 is N, and R1, R2, R3, R4, R5, R6, R7, R8, R9, R12, R13, X2, and X3 are as previously disclosed. The phthalimide protecting group in the compounds of Formula XXVIIc is removed as in step ***ag*** of Scheme XXII by reaction with hydrazine hydrate in a polar protic solvent such as EtOH at 90 °C to provide the compounds of Formula XXVIIIc, wherein R10 is Cl, R11 is CH_2NH_2, X1 is N, and R1, R2, R3, R4, R5, R6, R7, R8, R9, R12, R13, X2, and X3 are as previously disclosed. The compounds of Formula XXVIIIc can be transformed into the compounds of Formula One, wherein R10 is Cl, R11 is $CH_2N(C{=}O)(R14)$, X1 is N, and R1, R2, R3, R4, R5, R6, R7, R8, R9, R12, R13, X2, and X3 are as previously disclosed, by reaction with an acid in the presence of $HOBt{\bullet}H_2O$, EDC•HCl and a base, such as DIPEA, in a polar aprotic solvent, such as CH_2Cl_2, as in step ***ah$_{2b}$*** of Scheme XXII.

Scheme XXII




In step ***l*** of Scheme XXIII, the compound of Formula V, wherein Y, R1, R2, R3, R4, R5, R6, and R7 are as previously disclosed, and the compounds of Formula XIId, wherein X3 is CR9, R10 and X3 together form a linkage having 4 carbon atoms and with the ring carbon atoms form a 6-membered aromatic ring (or if desired a non-aromatic ring), R11 is CH_2N(Phthalimide) and R8, R9, R12, R13, X1 and X2 are as previously disclosed, are allowed to react in the presence of CuCl and 2,2-bipyridyl in a solvent, such as 1,2-dichlorobenzene, at a temperature of about 180 °C to provide the corresponding compounds of Formula XXVIId, wherein X3 is CR9, R10 and X3 together form a linkage having 4

carbon atoms and with the ring carbon atoms form a 6-membered aromatic ring, R11 is CH_2N(Phthalimide) and R1, R2, R3, R4, R5, R6, R7, R8, R9, R12, R13, X1 and X2 are as previously disclosed. The phthalimide protecting group in the compounds of Formula XXVIId is removed as in step ***ag*** of Scheme XXIII by reaction with hydrazine hydrate in a polar protic solvent such as EtOH at 90 °C to provide the compounds of Formula XXVIIId, wherein X3 is CR9, R10 and X3 together form a linkage having 4 carbon atoms and with the ring carbon atoms form a 6-membered aromatic ring, R11 is CH_2NH_2 and R1, R2, R3, R4, R5, R6, R7, R8, R9, R12, R13, X1 and X2 are as previously disclosed. The compounds of Formula XXVIIId can be transformed into the compounds of Formula One, wherein X3 is CR9, R10 and X3 together form a linkage having 4 carbon atoms and with the ring carbon atoms form a 6-membered aromatic ring, R11 is $CH_2N(C=O)(R14)$ and R1, R2, R3, R4, R5, R6, R7, R8, R9, R12, R13, X1 and X2 are as previously disclosed, by reaction with an acid in the presence of $HOBt•H_2O$, EDC•HCl and a base, such as DIPEA, in a polar aprotic solvent, such as CH_2Cl_2, as in step ***ah_{2b}*** of Scheme XXIII.


In another embodiment, the compounds of Formula XXVIIId can be transformed into the compounds of Formula One, wherein X3 is CR9, R10 and X3 together form a linkage having 4 carbon atoms and with the ring carbon atoms form a 6-membered aromatic ring, R11 is $CH_2N(C=O)N(R14)(R15)$ and R1, R2, R3, R4, R5, R6, R7, R8, R9, R10, R12, R13, X1 and X2 are as previously disclosed, by reaction with an isocyanate in the presence of a base such as TEA and in a non-reactive solvent such as CH_2Cl_2 at 0 °C as in step ***ai_1*** of Scheme XXIII.

Scheme XXIII



In step ***l*** of Scheme XXIV, the compound of Formula V, wherein Y, R1, R2, R3, R4, R5, R6, and R7 are as previously disclosed, and the compounds of Formula XIIe, wherein R11 is NHN(Phthalimide) and R8, R9, R12, R13, X1, X2, and X3 are as previously disclosed, are allowed to react in the presence of CuCl and 2,2-bipyridyl in a solvent, such as 1,2-dichlorobenzene, at a temperature of about 180 °C to provide the corresponding compounds of Formula XXVIIe, wherein R11 is NHN(Phthalimide) and R1, R2, R3, R4, R5, R6, R7, R8, R9, R12, R13, X1, X2, and X3 are as previously disclosed. The phthalimide protecting group in the compounds of Formula XXVIIe is removed as in step ***ag*** of Scheme XXIV by reaction with hydrazine hydrate in a polar protic solvent such as EtOH at 90 °C to provide the compounds of Formula XXVIIIe, wherein R11 is $NHNH_2$ and R1, R2, R3, R4, R5, R6, R7, R8, R9, R12, R13, X1, X2, and X3 are as previously disclosed. The compounds of Formula XXVIIIe can be transformed into the compounds of Formula One, wherein R11 is NHN(C=O)(R14) and R1, R2, R3, R4, R5, R6, R7, R8, R9, R12, R13, X1, X2, and X3 are as previously disclosed, by reaction with an acid in the presence of $HOBt•H_2O$, EDC•HCl and a base, such as DIPEA, in a polar aprotic solvent, such as CH_2Cl_2, as in step ***ah2b*** of Scheme XXIV.


Scheme XXIV



In step ***l*** of Scheme XXV, the compound of Formula V, wherein Y, R1, R2, R3, R4, R5, R6, and R7 are as previously disclosed, and the compounds of Formula XIIf, wherein R11 is ON(Phthalimide) and R8, R9, R10, R12, R13, X1, X2, and X3 are as previously disclosed, are allowed to react in the presence of CuCl and 2,2-bipyridyl in a solvent, such as 1,2-dichlorobenzene, at a temperature of about 180 °C to provide the corresponding compounds of Formula XXVIIf, wherein R11 is ON(Phthalimide) and R1, R2, R3, R4, R5,

R6, R7, R8, R9, R10, R12, R13, X1, X2, and X3 are as previously disclosed. The phthalimide protecting group in the compounds of Formula XXVIIf is removed as in step ***ag*** of Scheme XXV by reaction with hydrazine hydrate in a polar protic solvent such as EtOH at 90 °C to provide the compounds of Formula XXVIIIf, wherein R11 is ONH_2 and R1, R2, R3, R4, R5, R6, R7, R8, R9, R10, R12, R13, X1, X2, and X3 are as previously disclosed. The compounds of Formula XXVIIIf can be transformed into the compounds of Formula One, wherein R11 is ON(C=O)(R14) and R1, R2, R3, R4, R5, R6, R7, R8, R9, R10, R12, R13, X1, X2, and X3 are as previously disclosed, by reaction with an acid in the presence of $HOBt•H_2O$, EDC•HCl and a base, such as DIPEA, in a polar aprotic solvent, such as CH_2Cl_2, as in step ***ah$_{2b}$*** of Scheme XXV.

Scheme XXV




In step ***l*** of Scheme XXVI, the compound of Formula V, wherein Y, R1, R2, R3, R4, R5, R6, and R7 are as previously disclosed, and the compounds of Formula XVIII, wherein R11 is CH_2NH(2-pyridine) and R8, R9, R10, R12, R13, X1, X2, and X3 are as previously disclosed, are allowed to react in the presence of CuCl and 2,2-bipyridyl in a solvent, such as 1,2-dichlorobenzene, at a temperature of about 180 °C to provide the corresponding compounds of Formula One, wherein R11 is CH_2NH(2-pyridine), and R1, R2, R3, R4, R5, R6, R7, R8, R9, R10, R12, R13, X1, X2, and X3 are as previously disclosed.

The compounds of Formula One can be further elaborated by standard methods. For example, when R11 contains a thioether, the thioether can be oxidized to the sulfone by treatment with oxone in the presence of an acetone:water mixture at ambient temperature. When R11 contains an oxalate ester, the compound of Formula One can be transformed into

the corresponding oxalamide by reaction with an amine hydrochloride and a solution of trimethylaluminum in toluene in a non-reactive solvent such as CH_2Cl_2.


Scheme XXVI



V XVIII Formula One

In Scheme XXVII, a fluorobenzaldehyde of Formula XXIX, wherein R10, X1, X2, and X3 are as previously disclosed can be converted to a (1,2,4-triazol-1-yl)benzaldehyde of Formula XXX, wherein R11 is a substituted or unsubstituted 1,2,4-triazol-1-yl group, and R10, X1, X2, and X3 are as previously disclosed by reaction with a substituted or unsubstituted 1,2,4-triazole in the presence of a base, such as K_2CO_3, in a solvent such as DMF as in step ***aj***. In step ***ak***, the (1,2,4-triazol-1-yl)benzaldehyde of Formula XXX is converted to a (1,2,4-triazol-1-yl)vinyl benzene of Formula XXXIa wherein R11 is a substituted or unsubstituted 1,2,4-triazol-1-yl group, and R8, R10, X1, X2, and X3 are as previously disclosed by reaction with triphenyl phosphonium bromide in the presence of a base, such as K_2CO_3, in an aprotic solvent, such as 1,4-dioxane.

Scheme XXVII




XXIX XXX XXXIa

In Scheme XXVIII, a bromofluorobenzene of Formula XXXII, wherein R10, X1, X2, and X3 are as previously disclosed can be converted to a (1,2,4-triazol-1-yl)vinylbenzene of Formula XXXIb, wherein R11 is a substituted or unsubstituted 1,2,4-triazol-1-yl group, and R8, R10, X1, X2, and X3 are as previously disclosed in two steps. In step ***al***, the bromofluorobenzene is reacted with a substituted or unsubstituted 1,2,4-triazole in the presence of a base, such as K_2CO_3, in a solvent such as DMF to generate the (1,2,4-triazol-1-yl)bromobenzene. In step ***cl***, the (1,2,4-triazol-1-yl)bromobenzene is reacted with vinyl

boronic anhydride pyridine complex in the presence of a catalyst, such as $Pd(PPh_3)_4$, and a base, such as K_2CO_3 in a solvent such as toluene.


Scheme XXVIII



XXXII XXXIb

Coupling of the compounds of Formula V with compounds of Formula XXXIa and XXXIb can be accomplished as in Schemes XXIX. In step ***l***, a compound of Formula V, wherein Y is Br, R1, R2, R3, R4, R5, R6, and R7 are as previously disclosed, and a vinylbenzene of Formula XXXIa or XXXIb, wherein R11 is a substituted or unsubstituted 1,2,4-triazol-1-yl group, and R8, R9, R10, X1, X2, and X3 are as previously disclosed, are allowed to react in the presence of CuCl and 2,2-bipyridyl in a solvent, such as 1,2-dichlorobenzene, at a temperature of about 180 °C to provide the molecules of Formula One, wherein R11 is a substituted or unsubstituted 1,2,4-triazol-1-yl group, and R1, R2, R3, R4, R5, R6, R7, R8, R10, X1, X2, and X3 are as previously disclosed.

Scheme XXIX




V XXXIa or XXXIb Formula One

In Scheme XXX, compounds of Formula XXXIII wherein R11 is a 3-nitro-1,2,4-triazol-1-yl group, and R1, R2, R3, R4, R5, R6, R7, R8, R10, X1, X2, and X3 are as previously disclosed can be converted to compounds of Formula One, wherein R11 is a 3-amido-1,2,4-triazol-1-yl group, and R1, R2, R3, R4, R5, R6, R7, R8, R10, X1, X2, and X3 are as previously disclosed by a two-step process. In step ***am***, the 3-nitro-1,2,4-triazol-1-yl group is reduced to a 3-amino-1,2,4-triazol-1-yl group in the presence of zinc dust and ammonium chloride (NH_4Cl) in a protic solvent, such as MeOH. In step ***an***, the 3-amino-1,2,4-triazol-1-yl group is acylated with an acid chloride, such as cyclopropylcarbonyl

chloride or acetyl chloride, in the presence of a base, such as TEA, in a solvent such as CH_2Cl_2.


Scheme XXX



XXXIII Formula One


In step ***ao*** of Scheme XXXI, a bromophenyl methyl ketone of Formula XXXIV wherein R10, X1, X2, and X3 are as previously disclosed is converted to an phenyl methyl ketone of the Formula XXXV wherein R11 is a 1,2,4-triazol-1-yl group, and R10, X1, X2, and X3 are as previously disclosed by treatment with 1,2,4-triazole in the presence of a base, such as cesium carbonate (Cs_2CO_3), and a catalyst, such as copper iodide (CuI), in a solvent, such as DMF. In step ***ap***, the 1,2,4-triazolylacetophenone of Formula XXXV is converted to the trimethylsilyl enol ether of Formula XXXVI by treatment with trimethylsilyl triflluoromethanesulfonate in the presence of a base, such as TEA, in an aprotic solvent, such as CH_2Cl_2. In step ***aq***, the silyl enol ether is reacted with a compound of Formula V, wherein Y is Br, R1, R2, R3, R4, R5, R6, and R7 are as previously disclosed in the presence of CuCl and 2,2-bipyridyl in a solvent, such as 1,2-dichlorobenzene at a temperature of about 180 °C to generate a ketone of the Formula XXXVII, wherein R11 is a 1,2,4-triazol-1-yl group, and R1, R2, R3, R4, R5, R6, R7, R10, X1, X2, and X3 are as previously disclosed. In step ***ar***, the ketone of the Formula XXXVII is treated with methylmagnesium bromide in an aprotic solvent, such as THF to generate the tertiary alcohol. The tertiary alcohol then undergoes an elimination reaction when treated with a catalytic amount of *p*-toluenesulfonic acid in a solvent, such as toluene, when heated to a temperature to allow azeotropic removal of water to produce compounds of Formula One wherein R11 is a 1,2,4-triazol-1-yl group, R8 is methyl, and R1, R2, R3, R4, R5, R6, R7, R10, X1, X2, and X3 are as previously disclosed, as in step ***as***.

Scheme XXXI



In Scheme XXXII, a compound of Formula XXXVIII, wherein R10 and R11 together form a linkage, having 3-4 carbon atoms and an oxo substituent and with the ring carbon atoms form a 5- or 6-membered cyclic ring, and R1, R2, R3, R4, R5, R6, R7, R8, X1, X2, and X3 are as previously disclosed is converted to a molecule of Formula One, wherein R10 and R11 together form a linkage, having 3-4 carbon atoms and an alkylamine substituent with the ring carbon atoms form a 5- or 6-membered cyclic ring and R1, R2, R3, R4, R5, R6, R7, R8, X1, X2, and X3 are as previously disclosed, by treatment with an alkylamine, such as 3,3,3-trifluoropropylamine, in the presence of a reducing agent, such as sodium cyanoborohydride ($NaBH_3CN$), in a solvent, such as 1,2-dichloroethane (DCE).


Scheme XXXII



XXXVIII Formula One

In Scheme XXXIII, a compound of Formula XXXIX, wherein X1, X2, and X3 are as previously disclosed is converted to a molecule of Formula XL, wherein X1, X2, and X3 are as previously disclosed, by treatment with a reducing agent, such as $NaBH_3CN$, in a solvent, such as AcOH, as in step ***au***. In step ***av***, the nitrogen atom is protected with a *tert*-butyloxycarbonyl (BOC) group by reaction with di-*tert*-butyl dicarbonate in the presence of a catalyst, such as DMAP, in a solvent, such as acetonitrile (MeCN). The bromide of Formula XL can be converted to the olefin of Formula XLI, wherein R8, X1, X2 and X3 are as previously disclosed, by reaction with potassium vinyl trifluoroborate in the presence of a palladium catalyst, such as $PdCl_2(dppf)$, and a base, such as K_2CO_3, in a polar aprotic solvent such as dimethylsulfoxide (DMSO) at 100 °C, as in step ***aw***.


Scheme XXXIII



XXXIX XL XLI

In Scheme XXXIV, a compound of Formula XXXIX, wherein X1, X2, and X3 are as previously disclosed is converted to a molecule of Formula XLII, wherein X1, X2, and X3 are as previously disclosed in two steps. In step ***ax***, the olefin is formed by treatment of the bromide with potassium vinyl trifluoroborate in the presence of a palladium catalyst, such as $PdCl_2$, and a ligand, such as triphenylphosphine, and a base, such as Cs_2CO_3, in a solvent mixture such as THF/water. In step ***ay***, the nitrogen atom is protected with a BOC group by reaction with di-*tert*-butyl dicarbonate in the presence of a catalyst, such as DMAP, in a solvent, such as MeCN.


Scheme XXXIV



XXXIX XLII

In step ***l*** of Scheme XXXV, the compound of Formula V, wherein Y, R1, R2, R3, R4, R5, R6, and R7 are as previously disclosed, and the compounds of Formula XLI or XLII, wherein R8, X1, X2 and X3 are as previously disclosed, are allowed to react in the presence of CuCl and 2,2-bipyridyl in a solvent, such as 1,2-dichlorobenzene, at a temperature of about 150 °C to provide the corresponding compounds of Formula XLIIIa or XLIIIb, wherein R1, R2, R3, R4, R5, R6, R7, R8, X1, X2, and X3 are as previously disclosed.

Scheme XXXV




V XLI or XLII XLIIIa or XLIIIb

In Scheme XXXVI, a compound of Formula XLIIIa, wherein R1, R2, R3, R4, R5, R6, R7, R8, X1, X2, and X3 are as previously disclosed is converted to a molecule of Formula XLIV, wherein R1, R2, R3, R4, R5, R6, R7, R8, X1, X2, and X3 are as previously disclosed by treatment with trifluoroacetic acid (TFA), in a solvent such as CH_2Cl_2, as in step ***az***. Compounds of the Formula XLIV can then be transformed into compounds of the Formula XLV wherein R1, R2, R3, R4, R5, R6, R7, R8, X1, X2, and X3 are as previously disclosed, in two steps. In step ***ba***, the indoline is treated with sodium nitrite ($NaNO_2$), in an acid, such as concentrated HCl, at a temperature around 5 °C, to form the nitrosoindole. In step ***bb***, the nitrosoindole is reacted with NH_4Cl in the presence of zinc powder in a protic solvent, such as MeOH. In step ***bc***, compounds of the Formula XLV are transformed into compounds of the Formula XLVI, wherein X4 is N(R14)(C(=O)R14) and R1, R2, R3, R4, R5, R6, R7, R8, X1, X2, and X3 are as previously disclosed, by treatment with and acid, such as 3,3,3-


trifluoropropanoic acid, PyBOP, and a base, such as DIPEA, in a polar aprotic solvent, such as CH_2Cl_2.

Scheme XXXVI




In Scheme XXXVII, a compound of Formula XLIIIb, wherein R1, R2, R3, R4, R5, R6, R7, R8, X1, X2, and X3 are as previously disclosed is converted to an indole of Formula XLVII, wherein R1, R2, R3, R4, R5, R6, R7, R8, X1, X2, and X3 are as previously disclosed by treatment with TFA, in a solvent such as CH_2Cl_2, as in step ***bd***. Compounds of the Formula XLVII can be transformed into compounds of the Formula XLVIII wherein R1, R2, R3, R4, R5, R6, R7, R8, X1, X2, and X3 are as previously disclosed, by reaction with 4-nitrophenyl-2-((*tert*-butoxycarbonyl)amino)acetate in the presence of potassium fluoride (KF) and a crown ether, such as 18-crown-6-ether, in a solvent, such as MeCN, as in step ***be***. Compounds of the Formula XLVIII can be transformed into compounds of the Formula XLIX, wherein R1, R2, R3, R4, R5, R6, R7, R8, X1, X2, and X3 are as previously disclosed in two steps. In step ***bf***, the Boc group is removed by treatment with TFA, in a solvent such as CH_2Cl_2. In step ***bg***, the amine is treated with 3,3,3-trifluoropropanoic acid, PyBOP, and a base, such as DIPEA, in a polar aprotic solvent, such as CH_2Cl_2.

Scheme XXXVII



In Scheme XXXVIII, a compound of Formula L, wherein X1, X2, and X3 are as previously disclosed is converted to a compound of the Formula LI, wherein X1, X2, and X3 are as previously disclosed by treatment with copper (II) sulfate pentahydrate and Zn powder in a base, such as NaOH as in step ***bh***. Compounds of the Formula LI can be transformed into compounds of the Formula LII wherein X1, X2, and X3 are as previously disclosed, by reaction with hydrazine, in a solvent such as water, at a temperature around 95 °C, as in step ***bi***. In step ***bj***, the olefin of the Formula LIII wherein X1, X2, and X3 are as previously disclosed is formed by treatment of the bromide with potassium vinyl trifluoroborate in the presence of a palladium catalyst, such as $PdCl_2$(dppf), and a base, such as K_2CO_3, in a solvent mixture such as DMSO. Compounds of the Formula LIV, wherein X1, X2, and X3 are as previously disclosed, can be formed from compounds of the Formula LIII by reaction with ethyl bromoacetate, in the presence of a base, such as Cs_2CO_3, in a solvent, such as DMF.


Scheme XXXVIII




L LI LII LIII LIV

In step ***l*** of Scheme XXXIX, the compound of Formula V, wherein Y, R1, R2, R3, R4, R5, R6, and R7 are as previously disclosed, and the compound of Formula LIV, wherein R8, X1, X2 and X3 are as previously disclosed, are allowed to react in the presence of CuCl and 2,2-bipyridyl in a solvent, such as 1,2-dichlorobenzene, at a temperature of about 180 °C to provide the corresponding compound of Formula LV, wherein R1, R2, R3, R4, R5, R6, R7, R8, X1, X2, and X3 are as previously disclosed. The compound of Formula LV can be further transformed into a compound of the Formula LVI, wherein R1, R2, R3, R4, R5, R6, R7, R8, X1, X2, and X3 are as previously disclosed, in two steps. In step ***bl***, the ester is hydrolyzed to the acid in the presence of HCl and AcOH, at a temperature of about 100 °C. In step ***bm***, the acid is treated with an amine, such as 2,2,2-trifluoroethylamine, PyBOP, and a base, such as DIPEA, in a polar aprotic solvent, such as CH_2Cl_2.


Scheme XXXIX



V LIV




LV



LVI

In step ***bn*** of Scheme XL, carboxylic acids of the Formula LVII, wherein R11 is C(=O)OH and R8, R10, X1, X2, and X3 are as previously disclosed and compounds of the Formula V, wherein Y is Br and R1, R2, R3, R4, R5, R6, and R7 are as previously disclosed are allowed to react in the presence of CuCl and 2,2-bipyridyl in a solvent, such as *N*-methyl pyrrolidine, at a temperature of about 150 °C to afford compounds of Formula LVIII, wherein R11 is (C=O)OH and R1, R2, R3, R4, R5, R6, R7, R8, R9, R10, X1, X2, and X3 are as previously disclosed. Compounds of the Formula LVIII can be further transformed to the corresponding benzamides of Formula LIX, wherein R11 is (C=O)N(R14)(R15), and R1, R2, R3, R4, R5, R6, R7, R8, R9, R10, X1, X2, and X3 are as previously disclosed, by treatment with an amine, such as 2-amino-*N*-(2,2,2-trifluoroethyl)acetamide, PyBOP, and a base, such as DIPEA, in a polar aprotic solvent, such as CH_2Cl_2, as in step ***bo***.

Scheme XL




EXAMPLES

The examples are for illustration purposes and are not to be construed as limiting the invention disclosed in this document to only the embodiments disclosed in these examples.

Starting materials, reagents, and solvents that were obtained from commercial sources were used without further purification. Anhydrous solvents were purchased as Sure/Seal™ from Aldrich and were used as received. Melting points were obtained on a Thomas Hoover Unimelt capillary melting point apparatus or an OptiMelt Automated Melting Point System from Stanford Research Systems and are uncorrected. Molecules are given their known names, named according to naming programs within ISIS Draw, ChemDraw, or ACD Name Pro. If such programs are unable to name a molecule, the molecule is named using conventional naming rules. ^{1}H NMR spectral data are in ppm (δ) and were recorded at 300, 400, or 600 MHz, and ^{13}C NMR spectral data are in ppm (δ) and were recorded at 75, 100, or 150 MHz, unless otherwise stated.

Example 1: Preparation of 1-(1-Bromo-2,2,2-trifluoroethyl)-3,5-dichlorobenzene (AI1)




Step 1 Method A. 1-(3,5-Dichlorophenyl)-2,2,2-trifluoroethanol (AI2). To a stirred solution of 1-(3,5-dichlorophenyl)-2,2,2-trifluoroethanone (procured from Rieke Metals, UK; 5.0 grams (g), 20.5 millimoles (mmol)) in MeOH (100 milliliters (mL)) at 0 °C were added $NaBH_4$ (3.33 g, 92.5 mL) and 1 N aqueous NaOH solution (10 mL). The reaction mixture was warmed to 25 °C and stirred for 2 hours (h). After the reaction was deemed complete by thin layer chromatography (TLC), saturated aqueous NH_4Cl solution was added to the reaction mixture, and the mixture was concentrated under reduced pressure. The residue was diluted with diethyl ether (Et_2O) and washed with water (3 x 50 mL). The organic layer was dried over sodium sulfate (Na_2SO_4) and concentrated under reduced pressure to afford the title compound as a liquid (4.0 g, 79%): ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.41 (m, 3H), 5.00 (m, 2H), 2.74 (s, 1H); ESIMS *m/z* 242.97 ([M-H]$^-$).

Step 1 Method B. 1-(3,5-Dichlorophenyl)-2,2,2-trifluoroethanol (AI2). To a stirred solution of 3,5-dichlorobenzaldehyde (10 g, 57 mmol) in THF (250 mL) were added trifluoromethyltrimethylsilane (9.79 g, 69.2 mmol) and a catalytic amount of TBAF. The reaction mixture was stirred at 25 °C for 8 h. After the reaction was deemed complete by TLC, the reaction mixture was diluted with 3 N HCl and then was stirred for 16 h. The reaction mixture was diluted with water and was extracted with ethyl acetate (EtOAc; 3 x). The combined organic extracts were washed with brine, dried over Na_2SO_4, and concentrated under reduced pressure to afford the title compound as a liquid (8.41 g, 60%).


The following compounds were made in accordance with the procedures disclosed in **Step 1 Method A** of **Example 1** above.

2,6-Difluoro-4-(2,2,2-trifluoro-1-hydroxyethyl)benzonitrile




The product was isolated as a brown solid: mp 83-87 °C; ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.26 (d, *J* = 9.0 Hz, 2H), 5.12 (d, *J* = 6.0 Hz, 1H), 3.06 (s, 1H); ESIMS *m/z* 237.1 ([M+H]$^+$).

1-(3,5-Difluoro-4-methoxyphenyl)-2,2,2-trifluoroethanol



The product was isolated as a pale yellow liquid: ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.06 (d, *J* = 8.4 Hz, 2H), 4.97-4.94 (m, 1H), 4.03 (s, 3H), 2.64 (s, 1H); EIMS *m/z* 242.1 ($[M]^+$); IR (thinfilm) 3459, 1135 cm^{-1}.


1-(3,4-Dichlorophenyl)-2,2-difluoropropan-1-ol



The product was isolated as a colorless liquid: ^{1}H NMR (300 MHz, DMSO-d_6) δ 7.65 - 7.62 (m, 2H), 7.41 (d, *J* = 8.4 Hz, 1H), 6.49 (d, *J* = 5.1 Hz, 1H), 4.87 - 4.78 (m, 1H), 1.53 (t, *J* = 18.9 Hz, 3H); EIMS *m/z* 240.0 ($[M]^+$); IR (thinfilm) 3434, 1131, 801, 512 cm^{-1}.


The following compounds were made in accordance with the procedures disclosed in **Step 1 Method B** of **Example 1** above.

2,2,2-Trifluoro-1-(3,4,5-trichlorophenyl)ethanol (AI3)




The product was isolated as a pale yellow liquid (500 mg, 65%): ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.45 (s, 2H), 5.00 (m, 1H), 2.80 (s, 1H); ESIMS *m/z* 278 ($[M+H]^+$); IR (thin film) 3420, 1133, 718 cm^{-1}.

1-(3,5-Dichloro-4-fluorophenyl)-2,2,2-trifluoroethanol (AI4)




The product was isolated as a pale yellow liquid (500 mg, 65%): ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.41 (s, 2H), 5.00 (m, 1H), 2.80 (s, 1H); ESIMS *m/z* 262 ($[M+H]^+$); IR (thin film) 3420, 1133, 718 cm^{-1}.

1-(3,4-Dichlorophenyl)-2,2,2-trifluoroethanol (AI5)




The product was isolated as a pale yellow liquid (500 mg, 65%): ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.60 (s, 1H), 7.51 (m, 1H), 7.35 (m, 1H), 5.01 (m, 1H), 2.60 (s, 1H); EIMS *m/z* 244 ($[M]^+$).

1-(3,5-Dibromophenyl)-2,2,2-trifluoroethanol




The title molecule was isolated as a colorless liquid: ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.67 (s, 1H), 7.58 (s, 2H), 5.08-5.02 (m, 1H), 4.42 (bs, 1H); EIMS *m/z* 333.7 ($[M]^+$); IR (thin film) 3417, 2966, 1128, 531 cm^{-1}.

2,2,2-Trifluoro-1-(3-fluoro-5-(trifluoromethyl)phenyl)ethanol




The title molecule was isolated as a clear, colorless oil: ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.56 (s, 1H), 7.45 - 7.37 (m, 2H), 5.11 (q, *J* = 6.4 Hz, 1H), 3.22 (bs, 1H); ^{13}C NMR (101 MHz, $CDCl_3$) δ 162.42 (d, *J* = 249.5 Hz), 137.46 (d, *J* = 7.8 Hz), 132.89 (qd, *J* = 33.5, 7.9 Hz), 123.67 (q, *J* = 283.8 Hz), 122.92 (q, *J* = 270.68 Hz), 120.10 (t, *J* = 4.1 Hz), 118.13 (d, *J* = 23.0 Hz), 113.94 (dq, *J* = 24.2, 3.9 Hz), 71.57 (q, *J* = 32.4 Hz); EIMS *m/z* 262 ($[M]^+$).

1-(3-Chloro-5-(trifluoromethyl)phenyl)-2,2,2-trifluoroethanol




The product was isolated as a white solid (4.98 g, 77%): mp 42-46 °C; ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.83 - 7.50 (m, 3H), 5.10 (p, *J* = 6.2 Hz, 1H), 2.88 (d, *J* = 4.3 Hz, 1H); ^{13}C NMR (101 MHz, $CDCl_3$) δ 137.12, 135.84, 131.4, 133.03 (q, *J* = 33.3 Hz), 127.15 (q, *J* = 3.8 Hz), 124.50 (q, *J* = 308.0 Hz), 123.45 (q, *J* = 301.8 Hz),123.04, 72.06 (q, *J* = 32.5 Hz); ^{19}F NMR (376 MHz, $CDCl_3$) δ -62.93, -78.43; EIMS *m/z* 278 ($[M]^+$).

2,2,2-Trifluoro-1-(4-fluoro-3-(trifluoromethyl)phenyl)ethanol




The product was isolated as a brown liquid: ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.76 (d, *J* = 6.8 Hz, 1H), 7.69-7.67 (m, 1H), 7.28-7.23 (m, 1H), 5.05-5.02 (m, 1H); ESIMS *m/z* 261.1 ([M-H]$^-$); IR (thin film) 3418, 1131 cm^{-1}.

2,2,2-Trifluoro-1-(3,4,5-trifluorophenyl)ethanol




The product was isolated as a colorless liquid: ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.19-7.10 (m, 2H), 5.03-4.96 (m, 1H), 2.85 (bs, 1H); EIMS *m/z* 230.1 ([M]$^+$).

2,2,2-Trifluoro-1-(2,3,4-trifluorophenyl)ethanol




The product was isolated as a clear colorless liquid (4.61 g 66%): ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.23 (qd, *J* = 7.4, 6.1, 4.2 Hz, 1H), 6.93 (tdd, *J* = 9.2, 6.9, 2.2 Hz, 1H), 5.25 (q, *J* = 6.3 Hz, 1H), 3.02 - 2.74 (m, 1H); ^{13}C NMR (101 MHz, $CDCl_3$) δ 151.79 (ddd, *J* = 254.5, 9.8, 3.4 Hz), 149.52 (ddd, *J* = 253.5,11.0, 3.5 Hz), 139.67 (dt, *J* = 252.5, 15.3 Hz), 123.68 (q, *J* = 282.2 Hz), 122.48 (dt, *J* = 8.2, 4.1 Hz), 118.95 (dd, *J* = 10.6, 3.6 Hz), 112.73 (dd, *J* = 17.7, 3.9 Hz), 66.58 - 64.42 (m); ^{19}F NMR (376 MHz, $CDCl_3$) δ -78.95 (d, *J* = 6.2 Hz),-132.02 (dd, *J* = 20.0, 8.2 Hz), -137.89 (m), 159.84 (t, *J* = 20.3 Hz); EIMS *m/z* 230 ([M]$^+$).

2,2,2-Trifluoro-1-(2,4,5-trichlorophenyl)ethanol




The product was isolated as a white solid (3.37 g, 73%) : mp 70-73 °C; ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.63 (d, *J* = 2.5 Hz, 1H), 7.54 (d, *J* = 2.5 Hz, 1H), 5.72 - 5.57 (m, 1H), 2.85 (d, *J* = 4.8 Hz, 1H); ^{19}F NMR (376 MHz, $CDCl_3$) δ -77.84.

1-(4-Chloro-3-nitrophenyl)-2,2,2-trifluoroethanol



The product was isolated as a yellow oil (6.52 g, 73%): ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.04 (d, *J* = 2.0 Hz, 1H), 7.75 - 7.51 (m, 2H), 5.16 (m, 1H), 3.41 (d, *J* = 4.3 Hz, 1H); ^{13}C NMR (101 MHz, CDCl3) δ 147.65, 134.44, 132.23, 132.17, 128.11, 124.66, 123.60 (q, *J* = 283.8), 70.99 (q, *J* = 32.6 Hz); ^{19}F NMR (376 MHz, $CDCl_3$) δ -78.47; EIMS *m/z* 230 ($[M]^+$).


2,2,2-Trifluoro-1-(4-fluoro-3,5-dimethylphenyl)ethanol



The product was isolated as a white solid (6.49 g, 84%) : mp 45-49 °C; ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.10 (d, *J* = 6.8 Hz, 2H), 4.89 (m, 1H), 2.63 (d, *J* = 4.3 Hz, 1H), 2.27 (d, *J* = 2.2 Hz, 6H); ^{13}C NMR (101 MHz, $CDCl_3$) δ 160.45 (d, *J* = 246.0 Hz), 128.73, 127.97, 124.92 (d, *J* = 18.6 Hz), 124.19 (q, *J* = 279.1 Hz), 72.36 (q, *J* = 32.0 Hz), 14.61 (d, *J* = 4.1 Hz).


; ^{19}F NMR (376 MHz, $CDCl_3$) δ -78.48, -120.14; EIMS *m/z* 222 ($[M]^+$).

2,2,2-Trifluoro-1-(4-fluoro-3-methylphenyl)ethanol




The product was isolated as a white solid (2.12 g, 33%): mp 40-46 °C; ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.28 (d, *J* = 7.4 Hz, 1H), 7.25 - 7.14 (m, 1H), 7.01 (t, *J* = 8.9 Hz, 1H), 5.05 - 4.63 (m, 1H), 3.03 (d, *J* = 4.2 Hz, 1H); ^{13}C NMR (101 MHz, $CDCl_3$) δ 161.91 (d, *J* = 247.0 Hz), 130.62 (d, *J* = 5.6 Hz), 129.41 (d, *J* = 3.5 Hz), 126.55 (d, *J* = 8.5 Hz), 115.19 (d, *J* = 22.9 Hz), 72.23 (q, *J* = 32.1 Hz), 14.44 (d, *J* = 3.6 Hz); ^{19}F NMR (376 MHz, $CDCl_3$) δ -78.57, -116.15; EIMS *m/z* 208 ($[M]^+$).

1-(3-Chloro-4-methylphenyl)-2,2,2-trifluoroethanol




The product was isolated as a clear colorless oil (4.99 g, 75%): ^{1}H NMR (400 MHz, CDCl3) δ 7.31 (s, 1H), 7.10 (m, 2H), 4.79 (q, *J* = 6.1 Hz, 1H), 2.89 (bs, 1H), 2.25 (s, 3H); ^{13}C NMR (101 MHz, CDCl3) δ 137.64, 134.67, 132.99, 131.09, 128.01, 125.58, 124.02 (q, *J* = 284.8 Hz), 72.08 (q, *J* = 32.3 Hz); ^{19}F NMR (376 MHz, $CDCl_3$) δ -78.39; EIMS *m/z* 224.5 ([M]$^+$).

1-(3,4-Dibromophenyl)-2,2,2-trifluoroethanol




The product was isolated as a clear colorless oil (5.92 g, 88%): ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.76 (d, *J* = 2.0 Hz, 1H), 7.66 (d, *J* = 8.3 Hz, 1H), 7.29 (dd, *J* = 8.3, 2.0 Hz, 1H), 4.99 (qd, *J* = 6.4, 4.2 Hz, 1H), 2.75 (d, *J* = 4.3 Hz, 1H); ^{13}C NMR (101 MHz, $CDCl_3$) δ 134.52, 133.81, 132.60, 127.45, 126.19, 125.16, 123.71 (q, *J* = 283.8 Hz)., 71.57 (q, *J* = 32.5 Hz); ^{19}F NMR (376 MHz, $CDCl_3$) δ -78.44; EIMS *m/z* 334 ([M]$^+$).

2,2,2-Trifluoro-1-(3-(trifluoromethoxy)phenyl)ethanol




The product was isolated as a clear colorless oil (20.9 g, 79%): ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.55 - 7.36 (m, 3H), 7.33 - 7.14 (m, 1H), 5.06 (m, 1H), 2.80 (br m, 1H); ^{13}C NMR (101 MHz, $CDCl_3$) δ 149.36 (q, *J* = 2.0 Hz), 136.04, 129.99,125.78, 123.91 (q, *J* = 282.8 Hz), 121.90, 120.31 (q, *J* = 258.6 Hz),120.12, 72.04 (q, *J* = 32.3 Hz); ^{19}F NMR (376 MHz, $CDCl_3$) δ -57.92, -78.49; EIMS *m/z* 260 ([M]$^+$).

2-Fluoro-5-(2,2,2-trifluoro-1-hydroxyethyl)benzonitrile




The product was isolated as a clear colorless oil (5.47 g, 58%): ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.80 (dd, *J* = 5.9, 2.2 Hz, 1H), 7.76 (ddd, *J* = 7.8, 5.0, 2.3 Hz, 1H), 7.30 (d, *J* = 8.6 Hz, 1H), δ 5.09 (qd, *J* = 6.3, 4.2 Hz, 1H), 3.12 (bm, 1H); ^{13}C NMR (101 MHz, $CDCl_3$) δ 163.49 (d, *J* = 261.7 Hz), 134.23 (d, *J* = 8.6 Hz), 132.67 , 131.17 , 123.66 (q, *J* = 282.4 Hz), 116.79 (d, *J* = 20.1 Hz), 113.39 , 100.96 (d, *J* = 194.9), 71.07 (q, *J* = 32.5 Hz); ^{19}F NMR (376 MHz, $CDCl_3$) δ -78.70, -105.22; EIMS *m/z* 219 ([M]$^+$).

1-(3-Bromo-5-chlorophenyl)-2,2,2-trifluoroethanol




The product was isolated as a yellow liquid: ^{1}H NMR (300 MHz, DMSO-d_6) δ 7.78 (s, 1H), 7.67 (s, 1H), 7.57 (s, 1H), 7.15 (d, J = 5.7 Hz, 1H); EIMS m/z 288 ([M]$^+$); IR (thin film) 3435, 1175, 750 cm^{-1}.

1-(3-Bromo-5-fluorophenyl)-2,2,2-trifluoroethanol




The product was isolated as a pale yellow liquid: ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.43 (s, 1H), 7.29 - 7.26 (m, 1H), 7.18 (d, J = 8.8 Hz, 1H), 5.03 - 4.98 (m, 1H), 3.60 (bs, 1H). ; EIMS m/z 272.0 ([M]$^+$); IR (thin film) 3400, 1176, 520 cm^{-1}.

1-(3,5-Dichlorophenyl)-2,2,3,3,3-pentafluoropropan-1-ol




Using pentafluoroethyltrimethylsilane, the product was isolated as a white solid (6.22 g, 88%): mp 71-73 °C; ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.42 (t, J = 1.9 Hz, 1H), 7.37 (d, J = 1.8 Hz, 2H), 5.11 (dt, J = 16.2, 5.7 Hz, 1H), 2.62 (d, J = 4.9 Hz, 1H); ^{13}C NMR (101 MHz, $CDCl_3$) δ 136.90, 135.31, 129.84, 126.38, 70.94 (dd, J = 28.2, 23.1 Hz); ^{19}F NMR (376 MHz, $CDCl_3$) δ -81.06, -120.94 (d, J = 277.5 Hz), -129.18 (d, J = 277.5 Hz); EIMS m/z 295 ([M]$^+$).

2,2,3,3,3-Pentafluoro-1-(3,4,5-trichlorophenyl)propan-1-ol




Using pentafluoroethyltrimethylsilane, the product was isolated as an off white semi solid: ^{1}H NMR (300 MHz, DMSO-d_6) δ 7.78 (s, 2H), 7.29 (d, J = 5.4 Hz,), 5.50 - 5.40 (m, 1H); EIMS m/z 328.0 ([M]$^+$); IR (thin film) 3459, 1188, 797 cm^{-1}.

2,2,2-Trifluoro-1-(3-(trifluoromethyl)phenyl)ethanol




The product was isolated as a light yellow (13.8 g, 89%): ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.77 (s, 1H), 7.70-7.67 (m, 2H), 7.55 (t, *J* = 7.8 Hz, 1H), 5.12 (q, *J* = 6.6 Hz, 1H), 2.76 (s, 1H); ^{19}F NMR (376 MHz, $CDCl_3$) δ -62.8, -78.5; EIMS *m/z* 244 ($[M]^+$).

1-(3,4-Dichloro-5-methylphenyl)-2,2,2-trifluoroethanol




The product was isolated as an off pale yellow solid: ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.44 (s, 1H), 7.26 (s, 1H), 4.98 - 4.95 (m, 1H), 2.61 (d, *J* = 4.4 Hz, 1H), 2.44 (s, 3H).
; EIMS *m/z* 258.1 ($[M]^+$); IR (thin film) 3421, 2926, 1129, 748 cm^{-1}.

1-(3-Chloro-5-ethylphenyl)-2,2,2-trifluoroethanol




The product was isolated as an off brown liquid (0.43 g, 85%): ^{1}H NMR (300 MHz, DMSO-d_6) δ 7.34 (s, 1H), 7.31 - 7.30 (m, 2H), 6.99 (d, *J* = 5.7 Hz, 1H), 5.23 - 5.16 (m, 1H), 2.67 (m, 2H), 1.19 (t, *J* = 7.8 Hz, 3H); EIMS *m/z* 238.0 ($[M]^+$); IR (thin film) 3361, 1172, 749 cm^{-1}.

1-(4-Bromo-3,5-dichlorophenyl)-2,2,2-trifluoroethanol



The product was isolated as a colorless liquid: ^{1}H NMR (300 MHz, DMSO-d_6) δ 7.75 (s, 2H), 7.24 (d, *J* = 6.0 Hz, 1H), 5.34 - 5.29 (m, 1H); EIMS *m/z* 321.88 ($[M]^+$); IR (thin film) 3420, 1706, 1267, 804, 679 cm^{-1}.

1-(3,5-Dibromo-4-chlorophenyl)-2,2,2-trifluoroethanol




The product was isolated as a pale yellow gum: ^{1}H NMR (300 MHz, DMSO-d_6) δ 7.89 (s, 2H), 7.20 (d, J = 6.0 Hz, 1H) 5.34 - 5.30 (m, 1H); EIMS m/z 366.0 ([M]$^+$).

Step 2. 1-(1-Bromo-2,2,2-trifluoroethyl)-3,5-dichlorobenzene (AI1). To a stirred solution of 1-(3,5-dichlorophenyl)-2,2,2-trifluoroethanol (4.0 g, 16.3 mmol) in CH_2Cl_2 (50 mL), were added NBS (2.9 g, 16.3 mmol) and triphenyl phosphite (5.06 g, 16.3 mmol), and the resultant reaction mixture was heated at reflux for 18 h. After the reaction was deemed complete by TLC, the reaction mixture was cooled to 25 °C and was concentrated under reduced pressure. Purification by flash column chromatography (SiO_2, 100-200 mesh; eluting with 100% pentane) afforded the title compound as a liquid (2.0 g, 40%): ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.41 (s, 3H), 5.00 (m, 1H); EIMS m/z 306 ([M]$^+$).


The following compounds were made in accordance with the procedures disclosed in **Step 2** of **Example 1**.

5-(1-Bromo-2,2,2-trifluoroethyl)-1,2,3-trichlorobenzene (AI6)




The product was isolated as a colorless oil (300 mg, 60%): ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.59 (s, 2H), 5.00 (m, 1H); EIMS m/z 340.00 ([M]$^+$).

5-(1-Bromo-2,2,2-trifluoroethyl)-1,3-dichloro-2-fluorobenzene (AI7)




The product was isolated as a colorless oil (320 mg, 60%): ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.45 (s, 2H), 5.00 (m, 2H); EIMS m/z 324.00 ([M]$^+$).

4-(1-Bromo-2,2,2-trifluoroethyl)-1,2-dichlorobenzene (AI8)




The product was isolated as a colorless oil (300 mg, 60%): ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.63 (s, 1H), 7.51 (m, 1H), 7.35 (m, 1H), 5.01 (m, 1H); EIMS *m/z* 306.00 ($[M]^+$).

1,3-Dibromo-5-(1-bromo-2,2,2-trifluoroethyl)benzene




The title molecule was isolated as a colorless liquid: ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.71 (s, 1H), 7.59 (s, 2H), 5.04-4.97 (m, 1H); EIMS *m/z* 394.6 ($[M]^+$); IR (thin film) 1114, 535 cm^{-1}.

1-(1-Bromo-2,2,2-trifluoroethyl)-3-fluoro-5-(trifluoromethyl)benzene




The title molecule was isolated as a colorless liquid: ^{1}H NMR (400 MHz, DMSO-d_6) δ 7.90 (d, *J* = 8.4 Hz, 1H), 7.79-7.77 (m, 2H), 6.40-6.34 (m, 1H); EIMS *m/z* 324.00 ($[M]^+$); IR (thin film) 1175, 525 cm^{-1}.

1-(1-Bromo-2,2,2-trifluoroethyl)-3-chloro-5-(trifluoromethyl)benzene




The title molecule was isolated as a colorless liquid: ^{1}H NMR (400 MHz, CDCl3) δ 7.71 (s, 1H), 7.67 (s, 1H), 7.64 (s, 1H), 5.15-5.09 (m, 1H); EIMS *m/z* 340.00 ($[M]^+$); IR (thin film) 1178, 750, 540 cm^{-1}.

4-(1-Bromo-2,2,2-trifluoroethyl)-1-fluoro-2-(trifluoromethyl)benzene




The title molecule was isolated as a colorless liquid: ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.75-7.72 (m, 2H), 7.28-7.24 (m, 1H), 5.19-5.16 (m, 1H); EIMS *m/z* 326.0 ($[M]^+$); IR (thin film) 1114, 571 cm^{-1}.

5-(1-Bromo-2,2,2-trifluoroethyl)-1,2,3-trifluorobenzene




The title molecule was isolated as a brown liquid: ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.23-7.12 (m, 2H), 5.05-4.98 (m, 1H); EIMS *m/z* 292.0 ([M]$^+$); IR (thin film) 1116, 505 cm^{-1}.

1-(1-Bromo-2,2,2-trifluoroethyl)-2,3,4-trifluorobenzene




The title molecule was isolated as a colorless oil: ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.44 (qd, *J* = m, 1H), 7.11-7.03 (m,1H), 5.53-5.45 (m, 1H).

1-(1-Bromo-2,2,2-trifluoroethyl)-2,4,5-trichlorobenzene




The title molecule was isolated as an off white solid: ^{1}H NMR (300 MHz, DMSO-d_6) δ 8.06 (d, *J* = 2.1 Hz, 1H), 7.71 (s, 1H), 6.45 - 6.37 (m, 1H); EIMS *m/z* 340.0 ([M]$^+$); IR (thin film) 1186, 764, 576 cm^{-1}.

4-(1-Bromo-2,2,2-trifluoroethyl)-1-chloro-2-nitrobenzene




The title molecule was isolated as an off white solid: ^{1}H NMR (300 MHz, DMSO-d_6) δ 8.30 (s, 1H), 7.92 (d, *J* = 9.0 Hz, 1H), 6.43 - 6.35 (m, 1H); EIMS *m/z* 317.0 ([M]$^+$); IR (thin film) 2927, 1540, 1353, 1177, 766, 530 cm^{-1}.

5-(1-Bromo-2,2,2-trifluoroethyl)-2-fluoro-1,3-dimethylbenzene




The title molecule was isolated as a colorless liquid: ^{1}H NMR (300 MHz, DMSO-d_6) δ 7.32 (d, J = 7.2 Hz, 2H), 6.15-6.07 (m, 1H), 3.23 (s, 6H); ESIMS m/z 284.1([M+H]$^+$); IR (thin film) 2962, 1112, 500 cm^{-1}.

4-(1-Bromo-2,2,2-trifluoroethyl)-1-fluoro-2-methylbenzene




The title molecule was isolated as a colorless liquid: ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.34-7.28 (m, 2H), 7.04-6.98 (m, 1H), 5.10-5.03 (m, 1H), 2.29 (s, 3H); EIMS m/z 270.1([M]$^+$); IR (thin film) 2989, 1163 cm^{-1}.

1-(1-Bromo-2,2,3,3,3-pentafluoropropyl)-3,5-dichlorobenzene




The title molecule was isolated as a colorless liquid: ^{1}H NMR (400 MHz, DMSO-d_6) δ 7.79 (t, J = 2.0 Hz, 1H), 7.63 (S, 2H), 6.37-6.29 (m, 1H); EIMS m/z 356([M]$^+$); IR (thin film) 1673, 1130, 715, 518 cm^{-1}.

4-(1-Bromo-2,2,2-trifluoroethyl)-2-chloro-1-methylbenzene




The title molecule was isolated as a colorless liquid : ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.55 - 7.50 (m, 2H), 7.44 (d, J = 8.4 Hz, 1H), 6.24 - 6.16 (m, 1H); IR (thin film) 2983, 1112, 749, 564 cm^{-1}.

1,2-Dibromo-4-(1-bromo-2,2,2-trifluoroethyl)benzene




The title molecule was isolated as a colorless liquid: ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.75 (s, 1H), 7.67 (d, J = 8.4 Hz, 1H), 7.33-7.30 (m, 1H), 5.07-5.00 (m, 1H); EIMS m/z 393.8 ([M]$^+$); IR (thin film) 2981, 1644, 1165 cm^{-1}.

1-(1-Bromo-2,2,2-trifluoroethyl)-3-(trifluoromethoxy)benzene




The title molecule was isolated as a colorless liquid: ^{1}H NMR (300 MHz, DMSO-d_6) δ 7.65-7.60 (m, 2H), 7.56-7.50 (m, 2H), 6.35-6.27 (m, 1H); EIMS *m/z* 322 ([M]$^+$); IR (thin film) 3413, 1161, 564 cm^{-1}.

5-(1-Bromo-2,2,2-trifluoroethyl)-2-fluorobenzonitrile




The title molecule was isolated as a pale yellow liquid: ^{1}H NMR (300 MHz, $CDCl_3$) δ 8.15 - 8.12 (m, 1H), 8.00 - 7.98 (m, 1H), 7.69 - 7.63 (m, 1H), 6.31 - 6.26 (m, 1H); EIMS *m/z* 280.9 ([M]$^+$).

1-Bromo-3-(1-bromo-2,2,2-trifluoroethyl)-5-chlorobenzene




The title molecule was isolated as a pale yellow liquid: ^{1}H NMR (400 MHz, DMSO-d_6) δ 7.90 (s, 1H), 7.74 (s, 1H), 7.65 (s, 1H), 6.26 - 6.20 (m, 1H); EIMS *m/z* 349.9 ([M]$^+$); IR (thin film) 1114, 764 cm^{-1}.

1-Bromo-3-(1-bromo-2,2,2-trifluoroethyl)-5-fluorobenzene




The title molecule was isolated as a colorless liquid: ^{1}H NMR(400 MHz, $CDCl_3$) δ 7.43 (s, 1H), 7.32 - 7.29 (m, 1H), 7.22 (d, *J* = 8.8 Hz, 1H), 1.06 (q, 1H); EIMS *m/z* 334.0 ([M]$^+$); IR (thin film) 3087, 1168, 533 cm^{-1}.

5-(1-Bromo-2,2,3,3,3-pentafluoropropyl)-1,2,3-trichlorobenzene




The title molecule was isolated as a colorless liquid: ^{1}H NMR (300 MHz, DMSO-d_6) δ 7.85 (s, 2H), 6.38 - 6.29 (m, 1H); EIMS *m/z* 389.9 ([M]$^+$); IR (thin film) 1208, 798, 560 cm^{-1}.

4-(1-Bromo-2,2,2-trifluoroethyl)-2,6-difluorobenzonitrile




The title molecule was isolated as a purple solid: mp 59-63 °C; ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.25 (s, 2H), 5.11-5.07 (m, 1H); ESIMS *m/z* 299.0 ([M+H]$^+$).

1-(1-Bromo-2,2,2-trifluoroethyl)-3-(trifluoromethyl)benzene




The title molecule was isolated as a colorless liquid: mp 59-63 °C; ^{1}H NMR (300 MHz, CDCl3) δ 7.75-7.67 (m, 3H), 7.57-7.52 (m, 1H), 5.20-5.13 (m, 1H); ESIMS *m/z* 306.0 ([M]$^+$); IR (thinfilm) 3436, 2925, 1265, 749 cm^{-1}.

5-(1-Bromo-2,2,2-trifluoroethyl)-1,3-difluoro-2-methoxybenzene




The title molecule was isolated as a pale yellow liquid: ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.08 (d, *J* = 8.4 Hz, 2H), 5.03-4.98 (m, 1H), 4.04 (s, 3H); ESIMS *m/z* 304.1 ([M+H]$^+$); IR (thinfilm) 1114, 613 cm^{-1}.

5-(1-Bromo-2,2,2-trifluoroethyl)-1,2-dichloro-3-methylbenzene




The title molecule was isolated as a colorless liquid: ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.46 (s, 1H), 7.27 (s, 1H), 5.04 - 4.99 (m, 1H), 2.44 (s, 3H); EIMS *m/z* 320.0 ([M]$^+$); IR (thinfilm) 2925, 1112, 752, 580 cm^{-1}.

4-(1-Bromo-2,2-difluoropropyl)-1,2-dichlorobenzene




The title molecule was isolated as a colorless liquid: ^{1}H NMR (300 MHz, DMSO-d_6) δ 7.76 - 7.70 (m, 2H), 7.54 (dd, *J* = 8.4 1.8 Hz, 1H), 5.81 - 5.73 (m, 1H), 1.67 (d, *J* = 18.9 Hz, 3H); EIMS *m/z* 304.0 ([M]$^+$); IR (thinfilm) 1118, 800, 499 cm^{-1}.

1-(1-Bromo-2,2,2-trifluoroethyl)-3-chloro-5-ethylbenzene




The title molecule was isolated as a colorless liquid: ^{1}H NMR (400 MHz, DMSO-d_6) δ 7.43 (d, *J* = 5.6 Hz, 2H), 7.39 (s, 1H), 6.20-6.16 (m, 1H), 2.68 - 2.62 (m, 2H), 1.19 (t, *J* = 7.6 Hz, 3H); EIMS *m/z* 300.0 ([M]$^+$); IR (thinfilm) 2970, 1167, 716, 539 cm^{-1}.

2-Bromo-5-(1-bromo-2,2,2-trifluoroethyl)-1,3-dichlorobenzene




The title molecule was isolated as a colorless liquid: ^{1}H NMR (400 MHz, DMSO-d_6) δ 7.79 (s, 2H), 6.27 - 6.21 (m, 1H); EIMS *m/z* 383.9 ([M]$^+$); IR (thinfilm) 2924, 1114, 749, 534 cm^{-1}.

1,3-Dibromo-5-(1-bromo-2,2,2-trifluoroethyl)-2-chlorobenzene



The title molecule was isolated as a pale yellow liquid: ^{1}H NMR (300 MHz, DMSO-d_6) δ 7.97 (s, 2H), 6.27 - 6.19 (m, 1H); EIMS *m/z* 428.0 ([M]$^+$).

Example 2: Preparation of *N*-Methyl-4-vinylbenzamide (AI9)




Step 1. 4-Vinylbenzoyl chloride (AI10). To a stirred solution of 4-vinylbenzoic acid (1 g, 6.75 mmol) in CH_2Cl_2 (20 mL) at 0 °C were added a catalytic amount of DMF and oxalyl chloride (1.27 g, 10.12 mmol) dropwise over a period of 15 minutes (min). The reaction mixture was stirred at 25 °C for 6 h. After the reaction was deemed complete by TLC, the reaction mixture was concentrated under reduced pressure to give the crude acid chloride.

Step 2. *N*-Methyl-4-vinylbenzamide (AI9). To 1 M *N*-methylamine in THF (13.5 mL, 13.5 mmol) at 0 °C were added TEA (1.34 mL, 10.12 mmol) and the acid chloride from Step 1 above in THF (10 mL), and the reaction mixture was stirred at 25 °C for 3 h. After the reaction was deemed complete by TLC, the reaction mixture was quenched with water and then was extracted with EtOAc (3x). The combined EtOAc layer was washed with brine and dried over Na_2SO_4 and concentrated under reduced pressure to afford the title compound as an off-white solid (650 mg, 60%): ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.76 (d, J = 8.0 Hz, 2H), 7.45 (d, J = 8.0 Hz, 2H), 6.79 (m, 1H), 6.20 (br s,1H), 5.82 (d, J = 17.6 Hz, 1H), 5.39 (d, J = 10.8 Hz, 1H); ESIMS *m/z* 161.95 ($[M+H]^+$).


The following compounds were made in accordance with the procedures disclosed in accordance with **Example 2**.

***N,N*-Dimethyl-4-vinylbenzamide (AI11)**




The product was isolated as an off-white solid (650 mg, 60%): ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.42 (m, 4H), 6.71 (m, 1H), 5.80 (d, J = 17.6 Hz, 1H), 5.31 (d, J = 10.8 Hz, 1H), 3.05 (s, 3H), 3.00 (s, 3H); ESIMS *m/z* 176.01 ($[M+H]^+$).

***N*-(2,2,3-Trifluoromethyl)-4-vinylbenzamide (AI12)**




The product was isolated as an off-white solid (900 mg, 60%): ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.76 (d, J = 8.0 Hz, 2H), 7.45 (d, J = 8.0 Hz, 2H), 6.79 (m, 1H), 6.20 (br s,1H), 5.82 (d, J = 17.6 Hz, 1H), 5.39 (d, J = 10.8 Hz, 1H), 4.19 (m, 2H); ESIMS *m/z* 230.06 ($[M+H]^+$).

Morpholino(4-vinylphenyl)methanone (AI13)



The product was isolated as a white solid (850 mg, 60%): ESIMS *m/z* 218.12 ($[M+H]^+$).

Example 3: Preparation of Ethyl 2-methyl-4-vinylbenzoate (AI14)




Step 1. 4-Formyl-2-methylbenzoic acid (AI15). To a stirred solution of 4-bromo-2-methylbenzoic acid (10 g, 46.4 mmol) in dry THF (360 mL) at -78 °C was added *n*-BuLi (1.6 M solution in hexanes; 58.17 mL, 93.0 mmol) and DMF (8 mL). The reaction mixture was stirred at -78 °C for 1 h then was warmed to 25 °C and stirred for 1 h. The reaction mixture was quenched with 1 N HCl solution and extracted with EtOAc. The combined EtOAc extracts were washed with brine and dried over Na_2SO_4 and concentrated under reduced pressure. The residue was washed with *n*-hexane to afford the title compound as a solid (3.0 g, 40%): mp 196–198 °C; ^{1}H NMR (400 MHz, DMSO-d_6) δ 13.32 (br s, 1H), 10.05 (s, 1H), 7.98 (m, 1H), 7.84 (m, 2H), 2.61 (s, 3H); ESIMS *m/z* 163.00 ($[M-H]^-$).

Step 2. Ethyl 4-formyl-2-methylbenzoate (AI16). To a stirred solution of 4-formyl-2-methylbenzoic acid (3 g, 18.2 mmol) in EtOH (30 mL) was added H_2SO_4 and the reaction mixture was heated at 80 °C for 18 h. The reaction mixture was cooled to 25 °C and concentrated under reduced pressure. The residue was diluted with EtOAc and washed with water. The combined EtOAc extracts were washed with brine, dried over Na_2SO_4 and concentrated under reduced pressure to afford the title compound as a solid (2.8 g, 80%): ^{1}H NMR (400 MHz, $CDCl_3$) δ 10.05 (s, 1H), 8.04 (m, 1H), 7.75 (m, 2H), 4.43 (m, 2H), 2.65 (s, 3H), 1.42 (m, 3H).

Step 3. Ethyl 2-methyl-4-vinylbenzoate (AI14). To a stirred solution of ethyl 4-formyl-2-methylbenzoate (2.8 g, 4 mmol) in 1,4-dioxane (20 mL) were added K_2CO_3 (3.01 g, 21.87 mmol) and methyltriphenyl phosphonium bromide (7.8 g, 21.87 mmol) at 25 °C. Then the reaction mixture was heated at 100 °C for 18 h. After the reaction was deemed complete by TLC, the reaction mixture was cooled to 25 °C and filtered, and the filtrate was concentrated under reduced pressure. The crude compound was purified by flash chromatography (SiO_2, 100-200 mesh; eluting with 25–30% EtOAc in *n*-Hexane) to afford the title compound as a solid (2.0 g, 72%): ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.86 (m, 1H), 7.27

(m, 2H), 6.68 (dd, J = 17.6, 10.8 Hz , 1H), 5.84 (d, J = 17.6 Hz, 1H), 5.39 (d, J = 10.8 Hz, 1H), 4.39 (m, 2H), 2.60 (s, 3H), 1.40 (m, 3H); ESIMS *m/z* 191.10 ([M-H]$^-$); IR (thin film) 2980, 1716, 1257 cm^{-1}.


Example 4: Preparation of *tert*-Butyl 2-chloro-4-vinylbenzoate (AI17)



Step 1. *tert*-Butyl 4-bromo-2-chlorobenzoate (AI18). To a stirred solution of 4-bromo-2-chlorobenzoic acid (5 g, 21.37 mmol) in THF (30 mL) was added di-*tert*-butyl dicarbonate (25.5 g, 25.58 mmol), TEA (3.2 g, 31.98 mmol) and DMAP (0.78 g, 6.398 mmol), and the reaction mixture was stirred at 25 °C for 18 h. The reaction mixture was diluted with EtOAc and washed with water. The combined organic layer was washed with brine, dried over Na_2SO_4 and concentrated under reduced pressure. The residue was purified by flash chromatography (SiO_2, 100-200 mesh; eluting with 2–3% EtOAc in *n*-hexane) to afford the title compound as a liquid (3.2 g, 51%): ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.62 (m, 2H), 7.44 (d, J = 8.4 Hz, 1H), 1.59 (s, 9H); ESIMS *m/z* 290.10 ([M+H]$^+$); IR(thin film) 1728 cm^{-1}.


The following compounds were made in accordance with the procedures disclosed in **Step 1** of **Example 4**.

***tert*-Butyl 2-bromo-4-iodobenzoate (AI19)**



The product was isolated as a colorless oil (1.2 g, 50%): ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.01 (s, 1H), 7.68 (d, J = 8.4 Hz, 1H), 7.41 (d, J = 8.0 Hz, 1H), 1.59 (s, 9H); ESIMS *m/z* 382.10 ([M+H]$^+$); IR(thin film) 1727 cm^{-1}.

***tert*-Butyl 4-bromo-2-(trifluoromethyl)benzoate(AI20)**




The product was isolated as a colorless oil (1 g, 52%): ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.85 (s, 1H), 7.73 (d, J = 8.4 Hz, 1H), 7.62 (d, J = 8.4 Hz, 1H), 1.57 (s, 9H); ESIMS *m/z* 324.10 ([M+H]$^+$); IR (thin film) 1725 cm^{-1}.

Step 2. *tert*-Butyl 2-chloro-4-vinylbenzoate (AI17). To a stirred solution of *tert*-butyl 4-bromo-2-chlorobenzoate (1.6 g, 5.50 mmol) in toluene (20 mL) was added $Pd(PPh_3)_4$ (0.31 mg, 0.27 mmol), K_2CO_3 (2.27 g, 16.5 mmol) and vinylboronic anhydride pyridine complex (2.0 g, 8.3 mmol) and the reaction mixture was heated to reflux for 16 h. The reaction mixture was filtered, and the filtrate was washed with water and brine, dried over Na_2SO_4 and concentrated under reduced pressure. Purification by flash column chromatography (SiO_2, 100-200 mesh; eluting with 5–6% EtOAc in *n*-hexane) afforded the title compound as a liquid (0.6 g, 46%): ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.72 (d, *J* = 8.1 Hz, 1H), 7.44 (m, 1H), 7.31 (d, *J* = 8.0 Hz, 1H), 6.69 (dd, *J* =17.6, 10.8 Hz , 1H) , 5.85 (d, *J* = 17.6 Hz, 1H), 5.40 (d, *J* = 10.8 Hz, 1H), 1.60 (s, 9H); ESIMS *m/z* 238.95 ($[M+H]^+$); IR (thin film) 2931, 1725, 1134 cm^{-1}.


The following compounds were made in accordance with the procedures disclosed in **Step 2** of **Example 4**.

***tert*-Butyl 2-bromo-4-vinylbenzoate (AI21)**




The product was isolated as a colorless oil (1g, 52%): ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.68 (m, 2H), 7.36 (d, *J* = 8.0 Hz, 1H), 6.68 (dd, *J* = 17.6, 10.8 Hz , 1H), 5.84 (d, *J* = 17.6 Hz, 1H), 5.39 (d, *J* = 10.8 Hz, 1H), 1.60 (s, 9H); ESIMS *m/z* 282.10 ($[M+H]^+$); IR (thin film) 2978, 1724, 1130 cm^{-1}.

***tert*-Butyl 2-(trifluoromethyl)-4-vinylbenzoate (AI22)**




The product was isolated as a colorless oil (1.2 g, 50%): ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.71 (d, *J* = 6.4 Hz, 2H), 7.59 (d, *J* = 7.6 Hz, 1H), 6.77 (dd, *J* = 17.6, 10.8 Hz , 1H), 5.89 (d, *J* = 17.6 Hz, 1H), 5.44 (d, *J* = 10.8 Hz, 1H), 1.58 (s, 9H); ESIMS *m/z* 272.20 ($[M+H]^+$); IR (thin film) 2982, 1727, 1159 cm^{-1}.

Example 5: Preparation of *tert*-Butyl 2-cyano-4-vinylbenzoate (AI23)



To a stirred solution of *tert*-butyl 2-bromo-4-vinylbenzoate (0.5 g, 1.77 mmol) in DMF (20 mL) was added CuCN (0.23 g, 2.65 mmol), and the reaction mixture was heated at 140 °C for 3 h. The reaction mixture was cooled to 25 °C, diluted with water, and extracted with EtOAc. The combined organic layer was washed with brine, dried over Na_2SO_4, and concentrated under reduced pressure. The residue was purified by flash chromatography (SiO_2, 100-200 mesh; eluting with 15% EtOAc in *n*-hexane) to afford the title compound as a white solid (0.3 g, 72%): mp 51–53 °C; ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.03 (s, 1H), 7.77 (s, 1H), 7.64 (d, *J* = 8.4 Hz, 1H), 6.75 (dd, *J* = 17.6, 10.8 Hz , 1H), 5.93 (d, *J* = 17.6 Hz, 1H), 5.51 (d, *J* = 10.8 Hz, 1H), 1.65 (s, 9H); ESIMS *m/z* 229.84 ($[M+H]^+$); IR (thin film) 2370, 1709, 1142 cm^{-1}.


Example 6: Preparation of Ethyl 2-bromo-4-iodobenzoate (AI46)



To a stirred solution of 4-iodo-2-bromobenzoic acid (5 g, 15.29 mmol) in EtOH (100 mL) was added H_2SO_4 (5 mL), and the reaction mixture was heated at 80 °C for 18 h. The reaction mixture was cooled to 25 °C and concentrated under reduced pressure. The residue was diluted with EtOAc (2x100 mL) and washed with water (100 mL). The combined EtOAc extracts were washed with brine, dried over Na_2SO_4 and concentrated under reduced pressure to afford the compound as a pale yellow solid (5 g, 92%): ^{1}H NMR (400 MHz, DMSO-d_6) δ 8.04 (d, *J* = 1.2 Hz, 1H), 7.71 (d, *J* = 7.6 Hz, 1H), 7.51 (d, *J* = 8.4 Hz, 1H), 4.41 (q, *J* = 7.2 Hz, 2H), 1.41 (t, *J* = 7.2 Hz, 3H).


The following compounds were made in accordance with the procedures disclosed in **Example 6**.

Ethyl 4-bromo-2-chlorobenzoate (AI47)




The title compound was isolated as an off-white solid (2.0 g, 80 %): ^{1}H NMR (400 MHz, DMSO-d_6) δ 8.25 (d, *J* = 1.2 Hz, 1H), 7.79 (d, *J* = 7.6 Hz, 1H), 7.65 (d, *J* = 8.4 Hz, 1H), 4.65 (q, *J* = 7.2 Hz, 2H), 1.56 (t, *J* = 7.2 Hz, 3H).

Ethyl 4-bromo-2-methylbenzoate (AI48)




The title compound was isolated as a pale yellow liquid (3.0 g, 83%): ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.79 (d, J = 8.4 Hz, 1H), 7.41 (s, 1H), 7.39 (d, J = 8.4 Hz, 1H), 4.42 (q, J = 7.2 Hz, 2H), 2.60 (s, 3H), 1.40 (t, J = 7.2 Hz, 3H)ESIMS m/z 229.11 ($[M+H]^+$); IR (thin film) 1725 cm^{-1}.

Ethyl 4-bromo-2-fluorolbenzoate (AI49)



The title compound was isolated as a colorless liquid (9.0 g, 79%): ^{1}H NMR (400 MHz, DMSO-d_6) δ 7.84 (t, J = 8.4 Hz, 1H), 7.76 (d, J = 2.0 Hz, 1H), 7.58 (d, J = 1.6 Hz, 1H), 4.34 (q, J = 7.2 Hz, 2H), 1.32 (t, J = 7.2 Hz, 3H); ESIMS m/z 246.99 ($[M+H]^+$), IR (thin film) 1734 cm^{-1}.

Example 7: Preparation of Ethyl 4-bromo-2-ethylbenzoate (AI50)




To a stirred solution of 4-bromo-2-fluorobenzoic acid (2.0 g, 9.17 mmol) in THF (16 mL), was added 1.0 M ethyl magnesium bromide in THF (32 mL, 32.0 mmol) dropwise at 0°C and the resultant reaction mixture was stirred at ambient temperature for 18h. The reaction mixture was quenched with 2 N HCl and extracted with EtOAc. The combined EtOAc layer was dried over anhydrous Na_2SO_4 and concentrated under reduced pressure to afford crude 4-bromo-2-ethylbenzoic acid as a colorless liquid that was used in the next step without purification (0.4 g): ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.64 (d, J = 8.4 Hz, 1H), 7.47 (m, 1H), 7.43 (m, 1H), 2.95 (q, J = 4.0 Hz, 2H), 1.32 (t, J = 4.0 Hz, 3H); ESIMS m/z 228.97 ($[M+H]^+$).

The title compound was synthesized from 4-bromo-2-ethylbenzoic acid in accordance to the procedure in Example 6, isolated as a colorless liquid (0.15 g, 68%): ^{1}H NMR (400 MHz, DMSO-d_6)δ 7.90 (d, J = 8.4 Hz, 1H), 7.47 (m, 2H), 4.40 (q, J = 7.2 Hz, 2H), 3.06 (q, J

= 7.6 Hz, 2H), 1.42 (t, *J* = 7.2 Hz, 3H), 1.26 (t, *J* = 7.6 Hz, 3H); ESIMS *m/z* 226.96 ([M-H]$^-$); IR (thin film) 3443, 1686, 568 cm^{-1}.


Example 8: Preparation of Ethyl 2-bromo-4-vinylbenzoate (AI51)



To a stirred solution of ethyl 2-bromo-4-iodobenzoate (5 g, 14.3 mmol) in THF/water (100 mL, 9:1) was added potassium vinyltrifluoroborate (1.89 g, 14.3 mmol), Cs_2CO_3 (18.27 g, 56.07 mmol) and triphenylphosphine (0.22 g, 0.85 mmol) and the reaction mixture was degassed with argon for 20 min, then charged with $PdCl_2$ (0.05 g,0.28 mmol). The reaction mixture was heated to reflux for 16 h. The reaction mixture was cooled to ambient temperature and filtered through a Celite® bed and washed with EtOAc. The filtrate was again extracted with EtOAc and the combined organic layers washed with water and brine, dried over Na_2SO_4 and concentrated under reduced pressure to afford crude compound. The crude compound was purified by column chromatography (SiO_2, 100-200 mesh; eluting with 2% EtOAc/ petroleum ether) to afford the title compound as a light brown gummy material (2 g, 56%): ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.78 (d, *J* = 8.4 Hz, 1H), 7.71 (d, *J* = 1.2 Hz, 1H), 7.51 (d, *J* = 8.4 Hz, 1H), 6.69 (dd, *J* = 17.6, 10.8 Hz, 1H), 5.86 (d, *J* = 17.6 Hz, 1H), 5.42 (d, *J* = 11.2 Hz, 1H), 4.42 (q, *J* = 7.2Hz, 2H), 1.43 (t, *J* = 3.6 Hz, 3H); ESIMS *m/z* 255.18 ([M+H]$^+$); IR (thin film) 1729 cm^{-1}.


The following compounds were made in accordance with the procedures disclosed in **Example 8**.

Ethyl 2-methyl-4-vinylbenzoate (AI52)




The title compound was isolated as a colorless liquid (0.8 g, 80 %): ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.89 (d, *J* = 8.4 Hz, 1H), 7.27 (m, 2H), 6.79 (dd, *J* = 17.6, 10.8 Hz, 1H), 5.86 (d, *J* = 17.6 Hz, 1H), 5.42 (d, *J* = 11.2 Hz, 1H), 4.42 (q, *J* = 7.2 Hz, 2H), 2.60 (s, 3H), 1.43 (t, *J* = 7.2 Hz, 3H); ESIMS *m/z* 191.10 ([M+H]$^+$); IR (thin film) 1717, 1257 cm^{-1}.

Ethyl 2-fluoro-4-vinylbenzoate (AI53)



The title compound was isolated as a pale yellow liquid (2.0 g, 50 %): ^{1}H NMR (400 MHz, DMSO-d_6) δ 7.87 (t, J = 8.0 Hz, 1H), 7.51(d, J = 16.0 Hz, 1H), 7.48 (d, J = 16.0 Hz, 1H), 6.82 (dd, J = 17.6, 10.8 Hz, 1H), 6.09 (d, J = 17.6 Hz, 1H), 5.50 (d, J = 10.8 Hz, 1H), 4.35 (q, J = 7.2 Hz, 2H), 1.35 (t, J = 7.2 Hz, 3H); ESIMS m/z 195.19 ([M+H]$^+$); IR (thin film) 1728 cm^{-1}.


Example 9: Preparation of Ethyl 2-chloro-4-vinylbenzoate (AI54)



To a stirred solution of ethyl 2-chloro-4-bromobenzoate (2 g, 7.63 mmol) in DMSO (20 mL) was added potassium vinyltrifluoroborate (3.06 g, 22.9 mmol) and K_2CO_3 (3.16 g, 22.9 mmol). The reaction mixture was degassed with argon for 30 min. Bistriphenylphosphine(diphenylphosphinoferrocene)palladium dichloride (0.27 g, 0.38 mmol) was added and the reaction mixture was heated to 80 °C for 1 h. The reaction mixture was diluted with water (100 mL), extracted with EtOAc (2 x 50 mL), washed with brine, dried over Na_2SO_4 and concentrated under reduced pressure to obtain the compound as brown gummy material (1.1 g, 69%): ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.81 (d, J = 8.4 Hz, 1H), 7.46 (s, 1H), 7.33 (d, J = 8.4 Hz, 1H), 6.70 (dd, J = 17.6, 11.2 Hz, 1H), 5.87 (d, J = 17.6 Hz, 1H), 5.42 (d, J = 10.8 Hz, 1H), 4.41 (q, J = 7.2 Hz,2H), 1.43 (t, J = 7.2 Hz, 3H); ESIMS m/z 211.22 ([M+H]$^+$); IR (thin film) 1729, 886 cm^{-1}.


The following compounds were made in accordance with the procedures disclosed in **Example 9**.

Ethyl 2-ethyl-4-vinylbenzoate (AI55)




The title compound was isolated as a color less liquid (1.0 g, 66 %): ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.85 (m, 1H), 7.29 (m, 2H), 6.76 (d, J = 10.8 Hz, 1H), 5.86 (d, J = 17.6 Hz, 1H), 5.36 (d, J = 10.5 Hz, 1H), 4.41 (q, J = 7.2 Hz, 2H), 3.10 (q, J = 7.2 Hz, 2H), 1.40 (t, J = 7.2 Hz, 3H), 1.30 (t, J = 7.2 Hz, 3H); ESIMS m/z 205.26 ([M+H]$^+$); IR (thin film) 1720, 1607, 1263 cm^{-1}.

Methyl 2-methoxy-4-vinylbenzoate (AI56)




The title compound was isolated as a pale yellow liquid (1.2 g, 75 %): ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.79 (d, *J* = 8.0 Hz, 1H), 7.04 (d, *J* = 1.2 Hz, 1H), 6.97 (s, 1H), 6.74 (dd, *J* = 11.2, 11.2 Hz, 1H), 5.86 (d, *J* = 17.6 Hz, 1H), 5.39 (d, *J* = 17.6 Hz, 1H) 3.93 (s, 3H), 3.91 (s, 3H); ESIMS *m/z* 193.18 ([M+H]$^+$); IR (thin film) 1732 cm^{-1}.

Ethyl 2-(methylthio)-4-vinylbenzoate



The title compound was isolated as a brown liquid: ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.98 (d, *J* = 8.4 Hz, 1H), 7.23 - 7.18 (m, 2H), 6.78 (dd, *J* = 17.7, 10.8, Hz, 1H), 5.89 (d, *J* = 17.4 Hz, 1H), 5.42 (d, *J* = 10.8 Hz, 1H), 4.39 - 4.36 (m, 2H), 2.48 (s,3H), 1.39 (t, *J* = 6.9 Hz, 3H); ESIMS *m/z* 221.9 ([M+H]$^+$); IR (thin film) 1708 cm^{-1}.


Example 10: Preparation of (*E*)-Ethyl 4-(3-(3,5-dichlorophenyl)-4,4,4-trifluorobut-1-enyl)-2-methylbenzoate (AI24)



To a stirred solution of ethyl 2-methyl-4-vinylbenzoate (2.0 g, 10.5 mmol) in 1,2-dichlorobenzene (25 mL) were added 1-(1-bromo-2,2,2-trifluoroethyl)-3,5-dichlorobenzene (6.44 g, 21.0 mmol), CuCl (208 mg, 21 mmol) and 2,2bipyridyl (0.65 g, 4.1 mmol). The reaction mixture was degassed with argon for 30 min and then stirred at 180 °C for 24 h. After the reaction was deemed complete by TLC, the reaction mixture was cooled to 25 °C and filtered, and the filtrate was concentrated under reduced pressure. Purification by flash chromatography (SiO_2, 100-200 mesh; eluting with 25–30% EtOAc in petroleum ether) afforded the title compound as a solid (1.7 g, 40%): ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.91 (d, *J* = 8.0 Hz, 1H), 7.37 (m, 1H), 7.27–7.24 (m, 4H), 6.59 (d, *J* = 16.0 Hz, 1H), 6.59 (dd, *J* = 16.0, 8.0 Hz, 1H), 4.38 (q, *J* = 7.2 Hz, 2H), 4.08 (m, 1H), 2.62 (s, 3H), 1.42 (t, *J* = 7.2 Hz, 3H); ESIMS *m/z* 415.06 ([M-H]$^-$); IR (thin film) 1717, 1255, 1114 cm^{-1}.


Compounds **AI25, AI57-AI68** and **AC1–AC5** (Table 1) were made in accordance with the procedures disclosed in **Example 10**.

(*E*)-Ethyl 4-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-enyl)-2-(trifluoromethyl)-benzoic acid (AI25)




The product was isolated as a pale brown gummy liquid (500 mg, 40%): ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.79 (d, *J* = 8.0 Hz, 1H), , 7.71 (m, 1H), 7.61 (d, *J* = 7.6 Hz, 1H),7.42 (s, 2H), 6.70 (d, *J* = 16.0 Hz, 1H), 6.57 (dd, *J* = 16.0, 8.0 Hz, 1H), 4.42 (q, *J* = 7.2 Hz, 2H), 4.19 (m, 1H), 1.40 (t, *J* = 7.6 Hz, 3H),; ESIMS *m/z* 502.99 ([M-H]$^-$); IR (thin film) 1730, 1201, 1120, 749 cm^{-1}.

(*E*)-Ethyl 4-(3-(3,5-dichlorophenyl)-4,4,4-trifluorobut-1-enyl)-2-fluorobenzoate (AI57)




^{1}H NMR (400 MHz, $CDCl_3$) δ 7.38 (s, 1H), 7.26 (s, 3H), 7.21 (d, *J* = 8.4 Hz, 1H), 7.16 (d, *J* = 11.6 Hz, 1H), 6.59 (d, *J* = 16.0 Hz, 1H), 6.47 (dd, *J* = ,16.0, 8.0 Hz, 1H), 4.41 (q, *J* = 6.8 Hz, 2H), 4.18 (m, 1H), 1.41 (t, *J* = 6.8 Hz, 3H); ESIMS *m/z* 419.33 ([M-H]$^-$); IR (thin film) 1723, 1115, 802 cm^{-1}.

(*E*)-Ethyl 4-(3-(3,5-dichlorophenyl)-4,4,4-trifluorobut-1-enyl)-2-bromobenzoate (AI58)




^{1}H NMR (400 MHz, $CDCl_3$) δ 7.79 (d, *J* = 8.0 Hz, 1H), 7.67 (s, 1H), 7.38 (m, 2H), 7.26 (m, 2H), 6.56 (d, *J* = 16.0 Hz, 1H), 6.45 (dd, *J* = 16.0, 7.6 Hz, 1H), 4.42 (q, *J* = 7.2 Hz, 2H), 4.39 (m, 1H), 1.42 (t, *J* = 7.2 Hz, 3H); ESIMS *m/z* 481.22 ([M-H]$^-$); IR (thin film) 1727, 1114, 801, 685 cm^{-1}.

(*E*)-Ethyl 2-bromo-4-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl) but-1-enyl)benzoate (AI59)




^{1}H NMR (400 MHz, $CDCl_3$) δ 7.79 (d, *J* = 8.0 Hz, 1H), 7.67 (d, *J* = 1.6 Hz, 1H), 7.40 (s, 2H), 7.36 (d, *J* = 1.6 Hz, 1H), 6.56 (d, *J* = 16.0 Hz, 1H), 6.44 (dd, *J* = 16.0, 7.6 Hz, 1H), 4.42 (q, *J* = 6.8 Hz, 2H), 4.15 (m, 1H), 1.42 (t, *J* = 6.8 Hz, 3H); ESIMS *m/z* 514.74 ([M-H]$^-$); IR (thin film) 1726, 1115, 808, 620 cm^{-1}.

(*E*)-Ethyl 2-methyl-4-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl) but-1-enyl)benzoate (AI60)




The title compound was isolated as a light brown gummy material: ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.90 (d, *J* = 8.8 Hz, 1H), 7.34 (d, *J* = 6.0 Hz, 2H), 7.25 (d, *J* = 7.2 Hz, 2H), 6.59 (d, *J* = 16.0 Hz, 1H), 6.42 (dd, *J* = 16.0, 8.0 Hz, 1H), 4.38 (q, *J* = 7.2 Hz, 2H), 4.19 (m, 1H), 2.63 (s, 3H), 1.41 (t, *J* = 7.2 Hz, 3H).

(*E*)-Ethyl 2-chloro-4-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl) but-1-enyl)benzoate (AI61)




^{1}H NMR (400 MHz, $CDCl_3$) δ 7.87 (d, *J* = 8.0 Hz, 1H), 7.46 (d, *J* = 1.6 Hz, 1H), 7.40 (s, 2H), 7.31 (d, *J* = 1.6 Hz, 1H), 6.57 (d, *J* = 16.0 Hz, 1H), 6.44 (dd, *J* = 16.0 Hz,, 8.0 Hz, 1H), 4.42 (q, *J* = 6.8 Hz, 2H), 4.15 (m, 1H), 1.42 (t, *J* = 6.8 Hz, 3H); ESIMS *m/z* 470.73 ([M-H]$^-$); IR (thin film) 1726, 1115, 809, 3072 cm^{-1}.

(*E*)-Ethyl 4-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-enyl)-2-(trifluoromethyl)benzoate (AI62)




The title compound was isolated as a pale brown liquid (1.0 g, 46.3 %): ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.79 (d, *J* = 8.0 Hz, 1H), 7.71 (s, 1H), 7.61 (d, *J* = 7.6 Hz, 1H), 7.41 (s, 2H) 6.65 (d, *J* = 16.0 Hz, 1H), 6.49 (dd, *J* = 16.0, 8.0 Hz, 1H), 4.42 (q, *J* = 7.6 Hz, 2H), 4.15 (m, 1H), 1.42 (t, *J* = 7.6 Hz, 3H); ESIMS *m/z* 502.99 ([M-H]$^-$); IR (thin film) 1730, 1202, 1120, 750 cm^{-1}.

(*E*)-Ethyl 2-chloro-4-(3-(3,5-dichloro-4-fluorophenyl)-4,4,4-trifluorobut-1-enyl)benzoate (AI63)




^{1}H NMR (400 MHz, $CDCl_3$) δ 7.85 (d, *J* = 6.0 Hz, 1H), 7.46 (d, *J* = 1.8 Hz, 2H), 7.34 (m, 1H), 7.24 (m, 1H), 6.57 (d, *J* = 16.2 Hz, 1H), 6.45 (dd, *J* = 16.2, 7.2 Hz, 1H), 4.43 (q, *J* = 7.2 Hz, 2H), 4.13 (m, 1H), 1.41 (t, *J* = 7.2 Hz, 3H); ESIMS *m/z* 455.0 ([M+H]$^+$); IR (thin film) 1728, 1115, 817 cm^{-1}.

(*E*)-Ethyl 2-fluoro-4-(3-(3,5-dichloro-4-fluorophenyl)-4,4,4-trifluorobut-1-enyl)benzoate (AI64)




^{1}H NMR (400 MHz, $CDCl_3$) δ 7.93 (t, *J* = 7.6 Hz, 1H), 7.34 (d, *J* = 5.6 Hz, 2H), 7.21 (d, *J* = 8.0 Hz, 1H), 7.16 (d, *J* = 11.6 Hz, 1H), 6.59 (d, *J* = 16.0 Hz, 1H), 6.49 (dd, *J* = 16.0, 7.6 Hz, 1H), 4.42 (q, *J* = 7.6 Hz, 2H), 4.13 (m, 1H), 1.41 (t, *J* = 7.6 Hz, 3H); ESIMS *m/z* 436.81([M-H]$^-$); IR (thin film) 1725 cm^{-1}.

(*E*)-Ethyl 2-bromo-4-(3-(3,5-dichloro-4-fluorophenyl)-4,4,4-trifluorobut-1-enyl)benzoate (AI65)




^{1}H NMR (400 MHz, $CDCl_3$) δ 7.94 (d, *J* = 8.0 Hz, 1H), 7.67 (s, 1H), 7.36 (m, 3H), 6.56 (d, *J* = 15.6 Hz, 1H), 6.44 (dd, *J* = 15.6, 8.0 Hz, 1H), 4.42 (q, *J* = 6.8 Hz, 2H), 4.10 (m, 1H), 1.42 (t, *J* = 6.8 Hz, 3H); ESIMS *m/z* 498.74 ([M-H]$^-$); IR (thin film) 1726, 1114, 820, 623 cm^{-1}.

(*E*)-Ethyl 2-methyl-4-(3-(3,5-dichloro-4-fluorophenyl)-4,4,4-trifluorobut-1-enyl)benzoate (AI66)




The title compound was isolated as a brown semi-solid: ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.90 (d, *J* = 8.8 Hz, 1H), 7.34 (d, *J* = 6.0 Hz, 2H), 7.25 (d, *J* = 7.2 Hz, 2H), 6.59 (d, *J* = 16.0 Hz, 1H), 6.42 (dd, *J* = 16.0 Hz, 8.0 Hz, 1H), 4.38 (q, *J* = 7.2 Hz, 2H), 4.19 (m, 1H), 2.63 (s, 3H), 1.41 (t, *J* = 7.2 Hz, 3H); ESIMS *m/z* 432.90 ([M-H]$^-$); IR (thin film) 1715 cm^{-1}.

(*E*)-Methyl 2-methoxy-4-(3-(3,5-dichloro-4-fluorophenyl)-4,4,4-trifluorobut-1-enyl)benzoate (AI67)




^{1}H NMR (400 MHz, $CDCl_3$) δ 7.80 (d, *J* = 8.4 Hz, 1H), 7.35 (d, *J* = 6.0 Hz, 2H), 7.03 (d, *J* = 1.2 Hz, 1H), 6.92 (s, 1H), 6.59 (d, *J* = 15.6 Hz, 1H), 6.42 (dd, *J* = 15.6, 8.0 Hz, 1H), 4.13 (m, 1H), 3.93 (s, 3H), 3.88 (s, 3H); ESIMS *m/z* 437.29 ([M+H]$^+$); IR (thin film) 1724 cm^{-1}.

(*E*)-Ethyl 2-ethyl-4-(3-(3,5-dichloro-4-fluorophenyl)-4,4,4-trifluorobut-1-enyl)benzoate (AI68)




^{1}H NMR (400 MHz, $CDCl_3$) δ 7.85 (d, *J* = 8.0 Hz, 1H), 7.35 (d, *J* = 9.6 Hz, 2H), 7.26 (m, 1H), 7.24 (m, 1H), 6.60 (d, *J* = 15.6 Hz, 1H), 6.42 (dd, *J* = 15.6, 8.0 Hz, 1H), 4.38 (q, *J* = 7.2 Hz, 2H), 4.14 (m, 1H), 3.01 (q, *J* = 7.6 Hz 2H), 1.41 (t, *J* = 7.2 Hz, 3H), 1.26 (t, *J* = 7.6 Hz, 3H); ESIMS *m/z* 447.05 ([M-H]$^-$); IR (thin film) 1715, 1115, 817 cm^{-1}.

(*E*)-Ethyl 4-(3-(3,5-dichloro-4-fluorophenyl)-4,4,4-trifluorobut-1-en-1-yl)-2-(methylthio)benzoate




Isolated as a brown liquid: ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.99 (d, J = 8.1 Hz, 2H), 7.35 - 7.32 (m, 2H), 7.21 - 7.16 (m, 2H), 6.63 (d, J = 15.8 Hz, 1H), 6.45 (dd, J = 15.9, 7.8 Hz, 1H), 4.41 - 4.31 (m, 2H), 4.30 - 4.10 (m, 1H), 2.47 (s, 3H), 1.40 (t, J = 7.5 Hz, 3H); ESIMS *m/z* 466.88 ([M+H]$^+$); IR (thin film) 1705, 1114 cm^{-1}.

(*E*)-Ethyl 2-bromo-4-(3-(3,5-difluoro-4-methoxyphenyl)-4,4,4-trifluorobut-1-en-1-yl)benzoate



The product was isolated as a pale yellow liquid: ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.78 (d, J = 8.0 Hz, 1H), 7.66 (d, J = 1.6 Hz, 1H), 7.35 - 7.33 (m, 1H), 6.96 - 6.90 (m, 2H), 6.54 (d, J = 15.6 Hz, 1H), 6.43 (dd, J = 15.6, 8.0 Hz, 1H), 4.39 (q, J = 6.8 Hz, 2H), 4.09 - 4.05 (m, 1H), 4.02 (s, 3H), 1.40 (t, J = 7.2 Hz, 3H); EIMS *m/z* 478.2 ([M]$^+$); IR (thin film) 1727, 1113 cm^{-1}.

Example 11: Preparation of (*E*)-4-(3-(3,5-Dichlorophenyl)-4,4,4-trifluorobut-1-enyl)-2-methylbenzoic acid (AI32)




To a stirred solution of (*E*)-ethyl 4-(3-(3,5-dichlorophenyl)-4,4,4-trifluorobut-1-enyl)-2-methylbenzoate (1.7 g, 4.0 mmol) in 1,4-dioxane (10 mL) was added 11 N HCl (30 mL), and the reaction mixture was heated at 100 °C for 48 h. The reaction mixture was cooled to 25 °C and concentrated under reduced pressure. The residue was diluted with water and extracted with chloroform ($CHCl_3$). The combined organic layer was dried over Na_2SO_4 and concentrated under reduced pressure, and the crude compound was washed with *n*-hexane to afford the title compound as a white solid (0.7 g, 50%): mp 142–143 °C; ^{1}H NMR (400 MHz, DMSO-d_6) δ 12.62 (br s, 1H), 7.81 (d, J = 8.0 Hz, 1H), 7.66 (s, 3H), 7.52–7.44 (m, 2H), 6.89

(dd, *J* = 16.0, 8.0 Hz, 1H), 6.78–6.74 (d, *J* = 16.0 Hz, 1H), 4.84 (m, 1H), 2.50 (s, 3H); ESIMS *m/z* 387.05 ([M-H]$^-$); IR (thin film) 3448, 1701, 1109, 777 cm^{-1}.


The following compounds were made in accordance with the procedures disclosed in **Example 11**.

(*E*)-2-Methyl-4-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-enyl)benzoic acid (AI26)




The product was isolated as a pale brown gummy liquid (1 g, 46%): ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.97 (d, *J* = 8.0 Hz, 1H), 7.77 (s, 1H), 7.65 (m, 1H), 7.41 (s, 2H), 6.68 (d, *J* = 16.0 Hz, 1H), 6.53 (dd, *J* = 16.0, 8.0 Hz, 1H), 4.16 (m, 1H), 2.50 (s, 3H); ESIMS *m/z* 422.67 ([M-H]$^-$).

(*E*)-2-Chloro-4-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-enyl)benzoic acid (AI27)




The product was isolated as an off-white semi-solid (1 g, 45%): ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.99 (d, *J* = 8.4 Hz, 1H), 7.50 (m, 1H), 7.40 (s, 1H), 7.36 (m, 2H), 6.59 (d, *J* = 15.6 Hz, 1H), 6.48 (dd, *J* = 15.6, 7.6 Hz, 1H), 4.14 (m, 1H); ESIMS *m/z* 442.72 ([M-H]$^-$); IR (thin film) 3472, 1704, 1113, 808 cm^{-1}.

(*E*)-2-Bromo-4-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-enyl)benzoic acid (AI28)




The product was isolated as a brown solid (1 g, 45%): mp 70–71 °C; ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.99 (d, *J* = 8.0 Hz, 1H), 7.72 (s, 1H), 7.40 (m, 3H), 6.58 (d, *J* = 16.0 Hz, 1H), 6.48 (dd, *J* = 16.0, 8.0 Hz, 1H), 4.14 (m, 1H); ESIMS *m/z* 484.75 ([M-H]$^-$); IR (thin film) 3468, 1700 cm^{-1}.

(*E*)-2-Cyano-4-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-enyl)benzoic acid (AI29)




The product was isolated as an off-white solid (500 mg, 45%): mp 100–101 °C; ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.90 (s, 1H), 7.85 (d, *J* = 7.6 Hz, 1H), 7.72 (d, *J* = 8.0 Hz, 1H), 7.65 (br s, 1H), 7.42 (s, 2H), 6.73 (d, *J* = 16.0 Hz, 1H), 6.58 (dd, *J* = 16.0, 8.0 Hz, 1H), 4.19 (m, 1H); ESIMS *m/z* 431.93 ([M-H]$^-$).

(*E*)-4-(3-(3,4-Dichlorophenyl)-4,4,4-trifluorobut-1-enyl)-2-methylbenzoic acid (AI30)




The product was isolated as a pale brown liquid (500 mg, 46%): ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.03 (m, 1H), 7.49 (m, 2H), 7.29 (m, 1H), 7.22 (m, 2H), 6.73 (d, *J* = 16.0 Hz, 1H), 6.58 (dd, *J* = 16.0, 7.8 Hz, 1H), 4.16 (m, 1H), 2.64 (s, 3H); ESIMS *m/z* 386.84 ([M-H]$^-$); IR (thin film) 3428, 1690, 1113, 780 cm^{-1}.

(*E*)-4-(3-(3,5-Dichloro-4-fluorophenyl)-4,4,4-trifluorobut-1-enyl)-2-methylbenzoic acid (AI31)




The product was isolated as a white solid (500 mg, 50%): mp 91–93°C; ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.02 (d, *J* = 8.0 Hz, 1H), 7.35 (d, *J* = 5.6 Hz, 1H), 7.30 (m, 3H), 6.61 (d, *J* = 16.0 Hz, 1H), 6.48 (dd, *J* = 16.0, 8.0 Hz, 1H), 4.13 (m, 1H), 2.65 (s, 3H); ESIMS *m/z* 406.87 ([M-H]$^-$).

(*E*)-4-(4,4,4-Trifluoro-3-(3,4,5-trichlorophenyl)but-1-enyl)-2-(trifluoromethyl)benzoic acid (AI33)




The product was isolated as a white solid (500 mg, 45%): mp 142–143 °C; ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.97 (d, *J* = 8.0 Hz, 1H), 7.77 (s, 1H), 7.65 (m, 1H), 7.41 (s, 2H), 6.68 (d, *J* = 16.0 Hz, 1H), 6.53 (dd, *J* = 16.0, 8.0 Hz, 1H), 4.16 (m, 1H); ESIMS *m/z* 474.87 ([M-H]$^-$).

(*E*)-2-Bromo-4-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-enyl)benzoic acid (AI69)




The title compound was isolated as a brown solid (0.8 g, 28%): ^{1}H NMR (400 MHz, $CDCl_3$) δ 13.42 (br, 1H), 7.98 (d, *J* = 1.5 Hz, 1H), 7.94 (m, 2H), 7.75 (d, *J* = 8.1 Hz, 1H), 7.65 (m, 1H), 7.06 (dd, *J* = 15.9, 9.0 Hz, 1H), 6.80 (d, *J* = 15.9 Hz, 1H), 4.91 (m, 1H); ESIMS *m/z* 484.75 ([M-H]$^-$); IR (thin film) 3469, 1700 cm^{-1}.

(*E*)-2-Bromo-4-(3-(3,5-dichloro-4-fluorophenyl)-4,4,4-trifluo robut-1-enyl)benzoic acid (AI70)




The title compound was isolated as a yellow liquid (0.3 g, crude): ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.79 (d, *J* = 8.1 Hz, 1H), 7.67 (s, 1H), 7.34 (m, 3H), 6.56 (d, *J* = 15.9 Hz, 1H), 6.45 (dd, *J* = 15.9, 7.6 Hz, 1H), 4.43 (m, 1H); ESIMS *m/z* 471.0 ([M-H]$^-$).

(*E*)-4-(3-(3,5-Dichloro-4-fluorophenyl)-4,4,4-trifluorobut-1-enyl)-2-ethylbenzoic acid (AI71)




The title compound was isolated as a brown gummy material (0.2 g, crude): ^{1}H NMR (300 MHz, DMSO-d_6) δ 12.5 (br, 1H), 7.85 (d, *J* = 6.3 Hz, 2H), 7.75 (d, *J* = 8.1 Hz, 1H), 7.52 (m, 2H), 6.96 (dd, *J* = 8.7, 8.7 Hz, 1H), 6.78 (d, *J* = 15.6 Hz, 1H), 4.80 (m, 1H), 4.06 (q, *J* = 7.2 Hz, 2H), 1.33 (t, *J* = 7.2 Hz, 3H); ESIMS *m/z* 419.06 ([M-H]$^-$).

(*E*)-2-Chloro-4-(3-(3,5-dichloro-4-fluorophenyl)-4,4,4-trifluorobut-1-enyl)benzoic acid (AI72)




The title compound was isolated as a yellow liquid (0.7 g, 95%): ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.85 (d, *J* = 6.0 Hz, 1H), 7.46 (d, *J* = 1.8 Hz, 1H), 7.41 (s, 3H), 6.57 (d, *J* = 16.0 Hz, 1H), 6.45 (dd, *J* = 16.0, 8.0 Hz, 1H), 4.16 (m, 1H); ESIMS *m/z* 455.0 ($[M+H]^+$); IR (thin film) 1728, 1115, 817 cm^{-1}.

(*E*)-4-(3-(3,5-Dichloro-4-fluorophenyl)-4,4,4-trifluorobut-1-enyl)-2-methylbenzoic acid (AI73)




The title compound was isolated as a light brown gummy material (0.7 g, 38%): mp 91-93 °C; ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.02 (d, *J* = 8.0 Hz, 1H), 7.35 (d, *J* = 5.6 Hz, 1H), 7.30 (m, 3H), 6.10 (d, *J* = 16.0 Hz, 1H), 6.46 (dd, *J* = 16.0, 8.0 Hz, 1H), 4.03 (m, 1H), 2.65 (s, 3H); ESIMS *m/z* 406.87 ($[M-H]^-$).

(*E*)-4-(3-(3,5-Dichlorophenyl)-4,4,4-trifluorobut-1-enyl)-2-fluorobenzoic acid (AI74)




The title compound was isolated as a light brown liquid (0.3 g, crude): ESIMS *m/z* 393.15 ($[M-H]^-$).

(*E*)-2-Bromo-4-(3-(3,5-dichlorophenyl)-4,4,4-trifluorobut-1-enyl)benzoic acid (AI75)




The title compound was isolated as a light brown liquid (0.35 g, crude): ESIMS *m/z* 451.91 ($[M-H]^-$).

(*E*)-4-(3-(3,5-Dichloro-4-fluorophenyl)-4,4,4-trifluorobut-1-en-1-yl)-2-(methylthio)benzoic acid



^{1}H NMR (400 MHz, $CDCl_3$) δ 7.88 - 7.85 (m, 3H), 7.46 (d, *J* = 6.8 Hz, 1H), 7.37 (s,1H), 6.99 (dd, *J* = 15.6, 8.8 Hz, 1H), 6.85 (d, *J* = 16.0 Hz, 1H), 4.85 - 4.81 (m, 2H), 2.45 (s, 3H); ESIMS *m/z* 436.89 [(M-H)$^-$]; IR (thinfilm) 3469, 1686, 1259, 714 cm^{-1}.


(*E*)-2-Bromo-4-(3-(3,5-difluoro-4-methoxyphenyl)-4,4,4-trifluorobut-1-en-1-yl)benzoic acid



The title molecule was isolated as a brown liquid: ^{1}H NMR (300 MHz, DMSO-d_6) δ 13.48 (bs, 1H), 8.03 (s, 1H), 7.81 (d, *J* = 7.8 Hz, 1H), 7.69 (d, *J* = 8.1 Hz, 1H), 7.48 (d, *J* = 9.3 Hz, 2H), 7.05 (dd, *J* = 15.6, 9.0 Hz, 1H), 6.83 (d, *J* = 15.9 Hz, 1H), 4.86 - 4.74 (m, 1H), 4.00 (s, 3H); EIMS *m/z* 451.18 ([M]$^+$); IR (thin film) 3431, 1132 cm^{-1}.

Prophetically, compounds **AI34, AI36-AI41, AI44-AI45** (Table 1) could be made in accordance with the procedures disclosed in **Example 10**, or **Examples 10 and 11**.

Example 12: Preparation of (*E*)-4-(3-(3,5-Dichlorophenyl)-4,4,4-trifluorobut-1-enyl)-2-methyl-*N*-(2,2,2-trifluoroethyl)benzamide (AC6)




To a stirred solution of (*E*)-4-(3-(3,5-dichlorophenyl)-4,4,4-trifluorobut-1-enyl)-2-methylbenzoic acid in DMF was added 2,2,2-trifluoroethylamine, HOBt•H_2O, EDC•HCl and DIPEA, and the reaction mixture was stirred at 25 °C for 18 h. The reaction mixture was diluted with water and extracted with EtOAc. The combined organic layer was washed with brine, dried over Na_2SO_4 and concentrated under reduced pressure. Purification by flash column chromatography (SiO_2, 100-200 mesh; eluting with hexane:EtOAc afforded a white

semi-solid (110 mg, 50%): ^{1}H NMR (400 MHz, $CDCl_3$) 7.40 (m, 2H), 7.26 (m, 3H), 6.56 (d, J = 16.0 Hz, 1H), 6.48 (dd, J = 16.0, 8.0 Hz, 1H), 5.82 (br s, 1H), 4.08 (m, 3H), 2.52 (s, 3H); ESIMS m/z 468.40 ([M-H]$^-$); IR (thin film) 1657, 1113, 804 cm^{-1}.


Compounds **AC7–AC38, AC40-AC58, AC110-AC112, AC117,** and **AC118** (Table 1) were made in accordance with the procedures disclosed in **Example 12**.

Example 13: Preparation of 4-((*E*)-3-(3,5-Dichlorophenyl)-4,4,4-trifluorobut-1-enyl)-2-methyl-*N*-((pyrimidin-5-yl)methyl)benzamide (AC39)



To a stirred solution of (pyrimidin-5-yl)methanamine (0.15 g, 1.43 mmol) in CH_2Cl_2 (10 mL) was added drop wise trimethylaluminum (2 M solution in toluene; 0.71 mL, 1.43 mmol), and the reaction mixture was stirred at 25 °C for 30 min. A solution of ethyl 4-((*E*)-3-(3,5-dichlorophenyl)-4,4,4-trifluorobut-1-enyl)-2-methylbenzoate (0.3 g, 0.71 mmol) in CH_2Cl_2 was added drop wise to the reaction mixture at 25 °C. The reaction mixture was stirred at reflux for 18 h, cooled to 25 °C, quenched with 0.5 N HCl solution (50 mL) and extracted with EtOAc (2 x 50 mL). The combined organic extracts were washed with brine, dried over Na_2SO_4, and concentrated under reduced pressure. The crude compound was purified by flash chromatography (SiO_2, 100-200 mesh; eluting with 40% EtOAc in *n*-hexane) to afford the title compound (0.18 g, 55%): mp 141–144 °C; ^{1}H (400 MHz, $CDCl_3$) δ 9.19 (s, 1H), 8.79 (s, 2H), 7.37 (m, 2H), 7.23 (m, 2H),7.21 (m, 1H), 6.57 (d, J = 16.0 Hz, 1H), 6.40 (dd, J = 16.0, 7.6 Hz 1H), 6.21 (m, 1H), 4.65 (s, 2H), 4.11 (m, 1H), 2.46 (s, 3H); ESIMS m/z 477.83 ([M-H]$^-$).

Example 14: Preparation of (*E*)-2-Chloro-*N*-(2-oxo-2-((2,2,2-trifluoroethyl)amino)ethyl)-4-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-en-1-yl)benzamide (AC64)




To a stirred solution of glycine amide (0.15 g, 0.58 mmol) in CH_2Cl_2 (5 mL) was added trimethylaluminum (2 M solution in toluene; 1.45 mL, 2.91 mmol) dropwise, and the reaction mixture was stirred at 28 °C for 30 min. A solution of (*E*)-ethyl 2-chloro-4-(4,4,4-

trifluoro-3-(3,4,5-trichlorophenyl)but-1-enyl)benzoate (0.3 g, 0.58 mmol) in CH_2Cl_2 (5 mL) was added drop wise to the reaction mixture at 28 °C. The reaction mixture was stirred at reflux for 18 h, cooled to 25 °C, quenched with 1N HCl solution (50 mL) and extracted with CH_2Cl_2 (2 x 50 mL). The combined organic extracts were washed with brine, dried over Na_2SO_4, and concentrated under reduced pressure. The crude compound was purified by flash chromatography (SiO_2, 100-200 mesh; eluting with 40% EtOAc in *n*-hexane) to afford the title compound as yellow solid (0.15 g, 50%): mp 83-85 °C; ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.72 (d, *J* = 8.0 Hz, 1H), 7.44 (s, 1H), 7.40 (s, 2H), 7.36 (d, *J* = 6.8 Hz, 1H), 7.05 (t, *J* = 5.2 Hz, 1H), 6.70 (t, *J* = 5.2 Hz, 1H), 6.57 (d, *J* = 15.6 Hz, 1H), 6.44 (dd, *J* = 15.6, 8.0 Hz, 1H), 4.23 (d, *J* = 5.6 Hz, 2H), 4.15 (m, 1H), 4.01 (m, 2H); ESIMS *m/z* 580.72 ([M-H]$^-$).

Compounds **AC59-AC75** (Table 1) were made in accordance with the procedures disclosed in **Example 14**.


Example 15: Preparation of (*E*)-2-Bromo-4-(3-(3,5-dichloro-4-fluorophenyl)-4,4,4-trifluorobut-1-en-1-yl)-*N*-(2-oxo-2-((2,2,2-trifluoroethyl)amino)ethyl)benzamide (AC79)



To a stirred solution of (*E*)-2-bromo-4-(3-(3,5-dichloro-4-fluorophenyl)-4,4,4-trifluorobut-1-enyl)benzoic acid (300 mg, 0.638 mmol) in CH_2Cl_2 (5.0 mL) was added 2-amino-*N*-(2,2,2-trifluoroethyl)acetamide (172. mg, 0.638 mmol) followed by PyBOP (364.5 mg, 0.701 mmol) and DIPEA (0.32 mL, 1.914 mmol), and the resultant reaction mixture was stirred at ambient temperature for 18 h. The reaction mixture was diluted with water and extracted with CH_2Cl_2. The combined CH_2Cl_2 layer was washed with brine, dried over Na_2SO_4 and concentrated under reduced pressure. Purification by flash column chromatography (SiO_2, 100-200 mesh; eluting with 40% EtOAc/ petroleum ether) afforded the title compound as an off-white solid (121 mg, 31 %): ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.69 (t, *J* = 6.0 Hz, 1H), 8.58 (t, *J* = 6.0 Hz, 1H), 7.92 (s, 1H), 7.87 (d, *J* = 6.4 Hz, 2H), 7.62 (d, *J* = 8.4 Hz, 1H), 7.45 (d, *J* = 8.4 Hz, 1H), 7.0 (m, 1H), 6.76 (d, *J* = 15.6 Hz, 1H), 4.83 (t, *J* = 8.0 Hz, 1H), 3.98 (m, 4H); ESIMS *m/z* 610.97 ([M+H]$^+$); IR (thin film) 3303, 1658, 1166, 817 cm^{-1}.

Compounds **AC76-AC80, AC96-AC102,** and **AC113** (Table 1) were made in accordance with the procedures disclosed in **Example 15**.


Example 16: Preparation of (*E*)-4-(3-(3,5-Dichlorophenyl)-4,4,4-trifluorobut-1-en-1-yl)-*N*-(1,1-dioxidothietan-3-yl)-2-fluorobenzamide (AC83)



To a stirred solution of (*E*)-4-(3-(3,5-dichlorophenyl)-4,4,4-trifluorobut-1-enyl)-2-fluoro-*N*-(thietan-3-yl)benzamide (100 mg, 0.2159 mmol) in acetone/water (1:1, 5.0 mL) was added oxone (266 mg, 0.4319 mmol) and the resultant reaction mixture was stirred at ambient temperature for 4h. The reaction mixture was diluted with water and extracted with EtOAc. The combined EtOAc layer was dried over anhydrous Na_2SO_4 and concentrated under reduced pressure. Purification by flash column chromatography (SiO_2, 100-200 mesh; eluting with 30% EtOAc/ pet ether) afforded the title compound as an off white solid (70.0 mg, 66 %): ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.07 (t, *J* = 8.4 Hz, 1H), 7.39 (t, *J* = 1.6 Hz, 1H), 7.31 (d, *J* = 1.2 Hz, 1H), 7.26 (m, 2H), 7.23 (m, 2H), 7.19 (d, *J* = 1.6 Hz, 1H), 6.60 (d, *J* = 16.8 Hz, 1H), 6.49 (dd, *J* = 16.8, 7.6 Hz, 1H), 4.90 (m, 1H), 4.64 (m, 2H), 4.14 (m, 2H),; ESIMS *m/z* 493.83 ([M-H]$^-$); IR (thin film) 1527, 1113, 801, 1167, 1321 cm^{-1}.

Compounds **AC81-AC87** (Table 1) were made in accordance with the procedures disclosed in **Example 16**.


Example 17: Preparation of (E)-*N*-((5-Cyclopropyl-1,3,4-oxadiazol-2-yl)methyl)-4-(3-(3,5-dichlorophenyl)-4,4,4-trifluorobut-1-en-1-yl)-2-methylbenzamide (AC89)



A solution of (E)-*N*-(2-(2-(cyclopropanecarbonyl)hydrazinyl)-2-oxoethyl)-4-(3-(3,5-dichlorophenyl)-4,4,4-trifluorobut-1-enyl)-2-methylbenzamide (200 mg, 0.379 mmol) in phosphoryl chloride ($POCl_3$, 2.0 mL) was stirred at ambient temperature for 10 min, then the resultant reaction mixture was heated to 50 °C for 1h. The reaction mixture was quenched with ice water at 0 °C and extracted with EtOAc. The combined EtOAc layer was washed with saturated sodium bicarbonate ($NaHCO_3$) solution and brine solution, dried over anhydrous Na_2SO_4, and concentrated under reduced pressure. Purification by flash column chromatography (SiO_2, 100-200 mesh; eluting with 50% EtOAc/ pet ether) afforded the title compound as a light brown gummy material (70.0 mg, 36 %): ^{1}H NMR (400 MHz, $CDCl_3$) δ


7.43 (m, 2H), 7.27 (m, 2H), 7.23 (m, 2H), 6.58 (d, J = 16.0 Hz, 1H), 6.41 (dd, J = 16.0, 7.6 Hz, 1H), 4.79 (d, J = 5.6 Hz, 2H), 4.14 (m, 1H), 2.48 (s, 3H), 2.18 (m, 1H), 1.16 (m, 4H); ESIMS m/z 509.89 ($[M+H]^+$); IR (thin film) 1666, 1166, 1112, 800 cm^{-1}.

Example 18: Preparation of (*E*)-2-Bromo-*N*-(2-thioxo-2-((2,2,2-trifluoroethyl)amino)ethyl)-4-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-en-1-yl)benzothioamide (AC90)



To a stirred solution of (*E*)-2-bromo-*N*-(2-oxo-2-((2,2,2-trifluoroethyl)amino)ethyl)-4-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-en-1-yl)benzamide (400 mg, 0.638 mmol) in 5 mL of THF at ambient temperature was added 2,4-bis(4-methoxyphenyl)-1,3,2,4-dithiadiphosphetane-2,4-disulfide (Lawesson's reagent) (336 mg, 0.830 mmol) in one portion. The resulting reaction mixture was stirred for 18 h. TLC showed the reaction was not complete, therefore additional Lawesson's reagent (168 mg, 0.415 mmol) was added and reaction stirred for 48 h. After the reaction was deemed complete by TLC, the reaction mixture was concentrated under reduced pressure. Purification by flash chromatography (SiO_2, 230-400 mesh; eluting with 20% EtOAc in hexanes) afforded the title compound as a yellow glassy oil (188 mg, 44.7%): ^1H NMR (400 MHz, $CDCl_3$) δ 8.34 (m, 1H), 8.27 (m, 1H), 7.60 (d, J = 1.6 Hz, 1H), 7.49 (d, J = 8.0 Hz, 2H), 7.40 (s, 2H), 7.36 (dd, J = 8.2, 1.7 Hz, 1H), 6.53 (d, J = 16.0 Hz, 1H), 6.38 (dd, J = 15.9, 7.9 Hz, 1H), 4.89 (d, J = 8.4, 5.5 Hz, 2H), 4.48 (qd, J = 9.0, 6.0 Hz, 2H), 4.11 (m, 1H); ESIMS m/z 656.9 ($[M-H]^-$).

Example 19: Preparation of (*E*)-2-(2-Bromo-4-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-en-1-yl)phenylthioamido)-*N*-(2,2,2-trifluoroethyl)acetamide (AC91)




To a stirred solution of (E)-2-bromo-*N*-(2-oxo-2-((2,2,2-trifluoroethyl)amino)ethyl)-4-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-en-1-yl)benzamide (400 mg, 0.638mmol) in 5 mL of THF at ambient temperature was added Lawesson's reagent (64.5 mg, 0.160 mmol) in one portion. The resulting reaction mixture was stirred for 18 h, after which time, the

reaction mixture was concentrated under reduced pressure. Purification by flash chromatography (SiO_2, 230-400 mesh; eluting with 20% EtOAc in hexanes) afforded the title compounds as a yellow oil (18.5 mg, 4.51%): ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.18 (t, *J* = 5.0 Hz, 1H), 7.58 (d, *J* = 1.6 Hz, 1H), 7.47 (d, *J* = 8.0 Hz, 1H), 7.40 (s, 2H), 7.34 (dd, *J* = 8.1, 1.6 Hz, 1H), 6.52 (m, 2H), 6.37 (dd, *J* = 15.9, 7.9 Hz, 1H), 4.54 (d, *J* = 4.9 Hz, 2H), 4.12 (m, 1H), 3.99 (qd, *J* = 8.9, 6.5 Hz, 2H); ESIMS *m/z* 640.9 ([M-H]$^-$).


The following compound was made in accordance with the procedures disclosed in **Example 19**.

(*E*)-2-Bromo-*N*-(2-thioxo-2-((2,2,2-trifluoroethyl)amino)ethyl)-4-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-en-1-yl)benzamide (AC92)




The product was isolated as a colorless oil (17.9 mg, 4.36%): ^{1}H NMR (400 MHz, $CDCl_3$) δ 9.16 (d, *J* = 6.1 Hz, 1H), 7.65 (d, *J* = 1.6 Hz, 1H), 7.57 (d, *J* = 8.0 Hz, 1H), 7.41 (m, 3H), 7.21 (t, *J* = 5.6 Hz, 1H), 6.55 (d, *J* = 15.9 Hz, 1H), 6.41 (dd, *J* = 15.9, 7.8 Hz, 1H), 4.59 (d, *J* = 5.6 Hz, 2H), 4.45 (qd, *J* = 9.0, 6.0 Hz, 2H), 4.12 (q, *J* = 7.2 Hz, 1H); ESIMS *m/z* 640.9 ([M-H]$^-$).

Example 106: Preparation of Ethyl (*Z*)-2-Bromo-4-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-en-1-yl)benzoate (AI76)




The title compound was made in accordance with the procedure disclosed in Example 88 and was isolated as a yellow viscous oil (416 mg, 23%): ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.80 (d, *J* = 8.0 Hz, 1H), 7.40 (d, *J* = 1.7 Hz, 1H), 7.35 (s, 2H), 7.12 (dd, *J* = 8.0, 1.7 Hz, 1H), 6.86 (d, *J* = 11.4 Hz, 1H), 6.23 - 5.91 (m, 1H), 4.42 (q, *J* = 7.1 Hz, 2H), 4.33 - 4.10 (m, 1H), 1.42 (t, *J* = 7.2 Hz, 3H); ^{19}F NMR (376 MHz, $CDCl_3$) δ -69.34 (d, *J* = 8.3 Hz); EIMS *m/z* 514.10 ([M]$^-$); IR (thin film) 2983, 1727, 1247, 1204, 1116 cm^{-1}.

Example 107 : Preparation of (*Z*)-2-Bromo-4-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-en-1-yl)benzoic acid (AI77)



To a stirred solution of (*Z*)-ethyl 2-bromo-4-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-en-1-yl)benzoate (360 mg, 0.70 mmol) in CH_3CN (1.0 mL) was added iodotrimethylsilane (0.28 mL, 2.8 mmol). The reaction mixture was heated to reflux for 20 h, allowed to cool to ambient temperature and partitioned between CH_2Cl_2 and aqueous 10 % sodium thiosulfate ($Na_2S_2O_3$). The organic phase was washed once with aqueous 10% $Na_2S_2O_3$ and dried over magnesium sulfate ($MgSO_4$) and concentrated in vacuo. Passing the material through a silica plug with 10% EtOAc in hexanes, followed by 20% MeOH in CH_2Cl_2) as the eluting solvents afforded the title compound as a yellow foam (143 mg, 42%): mp 54-64°C; ^{1}H NMR (400 MHz, $CDCl_3$) δ 11.36 (s, 1H), 7.99 (d, *J* = 8.0 Hz, 1H), 7.43 (s, 1H), 7.30 (s, 2H), 7.14 (d, *J* = 7.9 Hz, 1H), 6.85 (d, *J* = 11.4 Hz, 1H), 6.15 (t, *J* = 10.9 Hz, 1H), 4.36 - 4.09 (m, 1H);^{19}F NMR (376 MHz, $CDCl_3$) δ -69.30.


Example 108 : Preparation of (*Z*)-2-Bromo-*N*-(2-oxo-2-((2,2,2-trifluoroethyl)amino)ethyl)-4-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-en-1-yl)benzamide (AC95)



To a stirred solution of (*Z*)-2-bromo-4-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-en-1-yl)benzoic acid (200 mg, 0.41 mmol) in anhydrous THF (5.0 mL) was added carbonyldiimidazole (82 mg, 0.51 mmol). The mixture was heated in a 50 °C oil bath for 1.5 h, treated with 2-amino-*N*-(2,2,2-trifluoroethyl)acetamide hydrochloride (109 mg, .057 mmol) and the resulting mixture heated to reflux for 8 h. After cooling to ambient temperature, the mixture was taken up in Et_2O and washed twice with aqueous 5% sodium bisulfate ($NaHSO_4$) (2X) and once with saturated NaCl (1X). After dying over $MgSO_4$, concentration in vacuo and purification by medium pressure chromatography on silica with EtOAc/Hexanes as the eluents, the title compound was obtained as a white foam (160 mg,


41%) mp 48-61°C: ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.58 (d, *J* = 7.9 Hz, 1H), 7.44 - 7.29 (m, 3H), 7.14 (dd, *J* = 7.9, 1.6 Hz, 1H), 6.86 (d, *J* = 11.4 Hz, 1H), 6.76 (t, *J* = 5.9 Hz, 1H), 6.59 (br s, 1H), 6.21 - 6.04 (m, 1H), 4.23 (d, *J* = 5.5 Hz, 1H), 3.98 (qd, *J* = 9.0, 6.5 Hz, 2H); ^{19}F NMR (376 MHz, $CDCl_3$) δ -69.31, -72.3; EIMS *m/z* 626.9 ($[M+1]^+$).

Example 109a: Preparation of (*E*)-2-Bromo-*N*-(piperidin-4-yl)-4-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-en-1-yl)benzamide (AC114)




(*E*)-*tert*-Butyl 4-(2-bromo-4-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-enyl)benzamido)piperidine-1-carboxylate (0.75 g, 1.11 mmol) was added to dioxane HCl (10 mL) at 0 °C and was stirred for 18 h. The reaction mixture was concentrated under reduced pressure and triturated with diethylether to afford the compound as a light brown solid (0.6 g, 88%).

Example 109b: Preparation of (*E*)-*N*-(1-Acetylpiperidin-4-yl)-2-bromo-4-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-en-1-yl)benzamide (AC103)




To a stirred solution of (*E*)-2-bromo-*N*-(piperidin-4-yl)-4-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-enyl)benzamide (0.1 g, 0.16 mmol) in CH_2Cl_2 (10.0 mL) was added TEA (0.046 mL, 0.35 mmol) and stirred for 10 min. Then acetyl chloride (0.014, 0.18 mmol) was added and stirred for 16 h at ambient temperature. The reaction mixture was diluted with CH_2Cl_2 and washed with saturated $NaHCO_3$ solution and brine solution. The combined CH_2Cl_2 layer was dried over Na_2SO_4 and concentrated under reduced pressure to afford crude compound. The crude compound was washed with 5% Et_2O / *n*-pentane to afford the title compound as a white solid (0.054 g, 50%).

Example 110: Preparation of (*E*)-2-Bromo-4-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-en-1-yl)-*N*-(1-(3,3,3-trifluoropropanoyl)piperidin-4-yl)benzamide (AC104)




To a stirred solution of 3,3,3-trifluoropropanoic acid (0.02g, 0.16 mmol) in CH_2Cl_2 (10.0 mL), (*E*)-2-bromo-N-(piperidin-4-yl)-4-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-enyl)benzamide (0.1 g, 0.16 mmol), PYBOP (0.09 g, 0.17 mmol), and DIPEA (0.06 g, 0.48 mmol) were added at ambient temperature. The reaction mixture was stirred at ambient temperature for 5 h. The reaction mixture was diluted with CH_2Cl_2. The combined CH_2Cl_2 layer was washed with 3N HCl and saturated $NaHCO_3$ solution, the separated CH_2Cl_2 layer was dried over anhydrous Na_2SO_4 and concentrated under reduced pressure to afford crude compound. The crude compound was purified by column chromatography (SiO_2, 100-200 mesh; eluting with 2% MeOH in CH_2Cl_2) to afford the title compound as an off white gummy material (0.035 g, 29.%).

Example 111: Preparation of (E)-2-Bromo-4-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-en-1-yl)-N-(1-(2,2,2-trifluoroethyl)piperidin-4-yl)benzamide (AC105)




To a stirred solution of (*E*)-2-bromo-*N*-(piperidin-4-yl)-4-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-enyl)benzamide (0.1 g, 0.16 mmol) in THF (5.0 mL) was added TEA (0.06 mL, 0.64 mmol) and stirred for 10 min. Then 2,2,2-trifluoroethyl triflluoromethanesulfonate (0.03, 0.16 mmol) was added and stirred for 16 h at ambient temperature. The reaction mixture was diluted with EtOAc and washed with saturated $NaHCO_3$ solution and brine solution. The combined EtOAc layer was dried over Na_2SO_4 and concentrated under reduced pressure to afford the title compound as a brown solid (0.05 g, 44%).

Example 112: Preparation of (*E*)-2-Bromo-*N*-(1-methylpiperidin-4-yl)-4-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-en-1-yl)benzamide (AC106)




A solution of (*E*)-2-bromo-*N*-(piperidin-4-yl)-4-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-enyl)benzamide (0.1 g, 0.16 mmol), formaldehyde (30% in water) (0.1 mL, 0.16 mmol) and AcOH (0.01 mL) in MeOH (5.0 mL) was stirred at ambient temperature for 30 min. After that $NaBH_3CN$ (0.01 g, 0.16 mmol) was added at 0°C and the reaction was stirred for 8 h at ambient temperature. The solvent was removed under reduced pressure to obtain residue which was diluted with EtOAc and washed with saturated aqueous $NaHCO_3$ solution and brine solution. The combined EtOAc layer was dried over Na_2SO_4 and concentrated under reduced pressure to obtain a residue, which was triturated with Et_2O/ pentane to afford the title compound as a pale yellow gummy material (0.06 g, 59%).

Example 113: Preparation of ((*E*)-2-Bromo-*N*-(1-(cyanomethyl)piperidin-4-yl)-4-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-en-1-yl)benzamide (AC107)




To a stirred solution of (*E*)-2-bromo-*N*-(piperidin-4-yl)-4-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-enyl)benzamide (0.25 g, 0.43 mmol) in THF (10.0 mL) was added TEA (0.16 mL, 1.29 mmol) and the reaction was stirred for 10 min. Then 2-bromoacetonitrile (0.07, 0.65 mmol) was added and the reaction was stirred for 8 h at ambient temperature. The reaction mixture was diluted with EtOAc and washed with saturated brine solution. The combined EtOAc layer was dried over Na_2SO_4 and concentrated under reduced pressure to afford the title compound as an off-white solid (0.125 g, 46.8%).

Example 114: Preparation of (*E*)-2-Bromo-*N*-(1-(oxetan-3-yl)piperidin-4-yl)-4-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-en-1-yl)benzamide (AC108)




A solution of (*E*)-2-bromo-*N*-(piperidin-4-yl)-4-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-enyl)benzamide (0.2 g, 0.35 mmol), oxetan-3-one (0.027 g, 0.38 mmol) and AcOH (0.01 mL) in MeOH (5.0 mL) was stirred at ambient temperature for 30 min. After that $NaBH_3CN$ (0.022 g, 0.35 mmol) was added at 0 °C slowly lot wise over the period of 10 min and the reaction was stirred for 8 h at ambient temperature. The solvent was removed under reduced pressure to obtain a residue which was diluted with EtOAc and washed with saturated $NaHCO_3$ solution and brine solution. The combined EtOAc layer was dried over Na_2SO_4 and concentrated under reduced pressure to obtain a residue, which was triturated with Et_2O/ pentane to afford the title compound as an off-white solid (0.05 g, 23%).

Example 115: Preparation of (*E*)-2-Bromo-*N*-(1-(2-hydroxyethyl)piperidin-4-yl)-4-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-en-1-yl)benzamide (AC109)




To a stirred solution of (*E*)-2-bromo-*N*-(piperidin-4-yl)-4-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-enyl)benzamide (0.25 g, 0.43 mmol) in THF (10.0 mL) was added TEA (0.16 mL, 1.29 mmol) and the reaction was stirred for 10 min. Then 2-chloroethanol (0.05, 0.65 mmol) was added and the reaction was stirred for 8 h at ambient temperature. The reaction mixture was diluted with EtOAc and washed with saturated brine solution. The combined EtOAc layer was dried over Na_2SO_4 and concentrated under reduced pressure to afford the title compound as an off-white solid (0.09 g, 34%).

Example 116: Preparation of (*E*)-2-(2-Bromo-4-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-en-1-yl)benzamido)acetic acid (AI78)




To a stirred solution of (*E*)-*tert*-butyl 2-(2-bromo-4-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-enyl)benzamido)acetate (440 mg, 0.734 mmol) in CH_2Cl_2 (36.0 ml), was added TFA (4.0 mL) and the reaction mixture was stirred at ambient temperature for 1 h. The reaction mixture was concentrated under reduced pressure to obtain residue which was washed with *n*-pentane to afford the title compound as an off-white solid (310 mg, 78%): ^{1}H NMR (400 MHz, $CDCl_3$) δ 13.0 (s, 1H), 8.75 (t, *J* = 5.7 Hz, 1H), 7.93 (m, 2H), 7.62 (d, *J* = 7.5 Hz, 1H), 7.40 (d, *J* = 8.1 Hz, 1H), 6.96 (dd, *J* = 15.3, 9.3 Hz, 1H), 6.78 (d, *J* = 15.3 Hz, 1H), 4.83 (m, 1H), 3.90 (d, *J* = 5.7 Hz, 2H); ESIMS *m/z* 543.61([M+H]$^+$); IR (thin film) 3429, 1635, 1114, 772 cm^{-1}.

Example 117: Preparation of (*E*)-*N*-((6-Chloropyridin-3-yl)methyl)-4-(3-(3,5-dichlorophenyl)-4,4,4-trifluorobut-1-en-1-yl)-2-methylbenzothioamide (AC115)



To the stirred solution of (*E*)-*N*-((6-chloropyridin-3-yl)methyl)-4-(3-(3,5-dichlorophenyl)-4,4,4-trifluorobut-1-enyl)-2-methylbenzamide (0.06 g, 0.117 mmol) in toluene (3 mL) was added Lawesson's reagent (0.14 g, 0.351 mmol) and the reaction was irradiated at 100 °C for 1 h, then cooled to ambient temperature and concentrated under reduced pressure to provide crude compound. The crude product was purified by preparative HPLC to afford the product as yellow color solid (0.03 g, 49%).

Example 118: Preparation of (*E*)-4-(3-(3,5-Dichloro-4-fluorophenyl)-4,4,4-trifluorobut-1-en-1-yl)-*N*-(2-oxo-2-((2,2,2-trifluoroethyl)amino)ethyl)-2-(trifluoromethoxy)benzamide (AC116)




Step 1. 2-(Trifluoromethoxy)-4-vinylbenzoic acid (AI79): To a stirred solution of 4-bromo-2-(trifluoromethoxy)benzoic acid (1 g, 3.67 mmol) in DMSO (20 mL) was added potassium vinyltrifluoroborate (1.47 g, 11.02 mmol) and K_2CO_3 (1.52 g, 11.02 mmol). The reaction mixture was degassed with argon for 30 min. Bistriphenylphosphine(diphenylphosphinoferrocene)palladium dichloride (0.13 g, 0.18 mmol) was added and the reaction mixture was heated to 80 °C for 1 h. The reaction mixture was diluted with water (100mL), extracted with EtOAc (2 x 50 mL), washed with brine, and dried over Na_2SO_4. Concentration under reduced pressure furnished the crude compound which was purified by flash column chromatography to afford the product as pale yellow gummy material (0.4 g, 47%): ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.05 (d, *J* = 8.1 Hz, 1H), 7.44 (d, *J* = 1.8 Hz, 1H), 7.35 (s, 1H), 6.78 (dd, *J* =17.4.1, 11.1 Hz, 1H), 5.92 (d, *J* = 17.4 Hz, 1H), 5.51 (d, *J* = 10.8 Hz, 1H); ESIMS *m/z* 232.97 ($[M+H]^+$).

Step 2. (*E*)-4-(3-(3,5-Dichloro-4-fluorophenyl)-4,4,4-trifluorobut-1-enyl)-2-(trifluoromethoxy)benzoic acid (AI80): To a stirred solution of 2-(trifluoromethoxy)-4-vinylbenzoic acid (0.356 g, 1.53 mmol) in 1N methyl pyrrolidine (5.0 mL) was added 1-(1-bromo-2,2,2-trifluoroethyl)-3,5-dichloro 4-fluorobenzene (1.0 g, 3.07 mmol), CuCl (0.03 g, 0.307 mmol) and 2,2 bipyridyl (0.095 g, 0.614 mmol). The reaction mixture was stirred at 150 °C for 1 h. After the reaction was complete by TLC, the reaction mixture was diluted with water (100mL) and extracted with EtOAc (2X 50 mL). The combined organic layers were washed with brine, dried over Na_2SO_4 and concentrated under reduced pressure to obtain the crude compound which was purified by flash column chromatography to afford the product as pale yellow gummy material (0.3 g, 21%): ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.08 (d, *J* = 8.0 Hz, 1H), 7.45 (d, *J* = 1.6 Hz, 1H), 7.35 (s, 3H), 6.63 (d, *J* = 16.0 Hz, 1H), 6.50 (dd, *J* = 16.0, 8.0 Hz, 1H), 4.15 (m, 1H); ESIMS *m/z* 474.81 ($[M-H]^-$).

Step 3. (*E*)-4-(3-(3,5-Dichloro-4-fluorophenyl)-4,4,4-trifluorobut-1-enyl)-*N*-(2-oxo-2-(2,2,2-trifluoroethylamino)ethyl)-2-(trifluoromethoxy)benzamide (AC116) : A mixture of (*E*)-4-(3-(3,5-dichloro-4-fluorophenyl)-4,4,4-trifluorobut-1-enyl)-2-(trifluoromethoxy)benzoic acid (0.25 g, 0.52 mmol), 2-amino-*N*-(2,2,2-trifluoroethyl)acetamide (0.158 g, 0.62 mmol), PyBOP (0.40 g, 0.78 mmol) and DIPEA (0.134 g, 1.04 mmol) in CH_2Cl_2 (10.0 mL) were stirred at ambient temperature for 16 h. The reaction mixture was diluted with water and extracted with CH_2Cl_2. The combined CH_2Cl_2 layer was washed with brine, dried over Na_2SO_4 and concentrated under reduced pressure. Purification by flash column chromatography (SiO_2, 100-200 mesh; eluting with 20% EtOAc/ pet ether) afforded the title compound as a pale yellow gummy material (0.15 g, 47 %).


The following molecules were made in accordance with the procedures disclosed in **Example 118, Step 2**:

(*E*)-4-(3-(3,5-Dibromophenyl)-4,4,4-trifluorobut-1-en-1-yl)-2-methylbenzoic acid




The title molecule was isolated as a brown solid: ^{1}H NMR (400 MHz, DMSO-d_6) δ 12.90 (bs, 1H), 7.85 (s, 1H), 7.78-7.75 (m, 3H), 7.47-7.41 (m, 2H), 6.89 (dd, J = 15.6, 9.2 Hz, 1H), 6.72 (d, J = 15.6 Hz, 1H), 4.80-4.75 (m, 1H), 2.33 (s, 3H); ESIMS m/z 474.90 ([M-H]$^-$); IR (thin film) 3437, 1689, 1165, 579 cm^{-1}.

(*E*)-4-(3-(3,5-Dibromophenyl)-4,4,4-trifluorobut-1-en-1-yl)-2-(trifluoromethyl)benzoic acid




The title molecule was isolated as a brown solid: ^{1}H NMR (300 MHz, DMSO-d_6) δ 13.5 (bs, 1H), 8.03 (s, 1H), 7.95-7.85 (m, 4H), 7.81 (d, J = 7.8 Hz, 1H), 7.14 (dd, J = 15.6, 9.6 Hz, 1H), 6.90 (d, J = 15.9 Hz, 1H), 4.86-4.79 (m, 1H); ESIMS m/z 528.82 ([M-H]$^+$); IR (thin film) 3437, 1707, 1153, 555 cm^{-1}.

(*E*)-2-Bromo-4-(3-(3,5-dibromophenyl)-4,4,4-trifluorobut-1-en-1-yl)benzoic acid




The title molecule was isolated as a brown liquid: ^{1}H NMR (400 MHz, DMSO-d_6) δ 13.90 (bs, 1H), 7.98 (s, 1H), 7.88 (s, 1H), 7.84 (s, 2H), 7.74 (d, J = 7.6 Hz, 1H), 7.64 (d, J = 8.8 Hz, 1H), 7.04 (dd, J = 15.6, 8.8 Hz, 1H), 6.78 (d, J = 15.6 Hz, 1H), 4.80-4.78 (m, 1H); ESIMS m/z 538.74 ([M-H]$^-$); IR (thin film) 3424, 1695, 1168, 578 cm^{-1}.

(*E*)-2-Bromo-4-(4,4,4-trifluoro-3-(3-fluoro-5-(trifluoromethyl)phenyl)but-1-en-1-yl)benzoic acid




The title molecule was isolated as a brown liquid: ^{1}H NMR (400 MHz, DMSO-d_6) δ 13.3 (bs, 1H), 7.93 (s, 1H), 7.82-7.77 (m, 2H), 7.72-7.66 (m, 2H), 7.59 (d, J = 8.0 Hz, 1H), 7.03 (dd, J = 15.6, 9.2 Hz, 1H), 6.76 (d, J = 15.6 Hz, 1H), 4.94-4.90 (m, 1H); ESIMS *m/z* 469.02 ([M-H]$^-$); IR (thin film) 3444, 1704, 1172, 513 cm^{-1}.

(*E*)-4-(3-(3,5-Bis(trifluoromethyl)phenyl)-4,4,4-trifluorobut-1-en-1-yl)-2-bromobenzoic acid




The title molecule was isolated as a brown solid: ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.98 (d, J = 7.6 Hz, 1H), 7.92 (s, 1H), 7.83 (s, 2H), 7.73 (d, J = 1.6 Hz, 1H), 7.42-7.40 (m, 1H), 6.62 (d, J = 16.4 Hz, 1H), 6.55 (dd, J = 16.0, 8.0 Hz, 1H), 4.40-4.30 (m, 1H); ESIMS *m/z* 518.94 ([M-H]$^-$); IR (thin film) 3447, 1705, 1171, 526 cm^{-1}.

(*E*)-2-Bromo-4-(4,4,4-trifluoro-3-(3-(trifluoromethyl)phenyl)but-1-en-1-yl)benzoic acid



The title molecule was isolated as a brown liquid: ^{1}H NMR (300 MHz, DMSO-d_6) δ 13.50 (bs, 1H), 7.97-7.87 (m, 3H), 7.78-7.61 (m, 4H), 7.08 (dd, J = 15.9, 9.3 Hz, 1H), 6.81 (d, J = 15.9 Hz, 1H), 4.97-4.84 (m, 1H); ESIMS *m/z* 518.94 ([M-H]$^-$); IR (thin film) 3447, 1705, 1171, 526 cm^{-1}.


(*E*)-2-Bromo-4-(3-(3-chloro-5-(trifluoromethyl)phenyl)-4,4,4-trifluorobut-1-en-1-yl)benzoic acid



The title molecule was isolated as a pale yellow gum: ^{1}H NMR (300 MHz, DMSO-d_6) δ 13.9 (s, 1H), 8.03 (s, 1H), 7.96 - 7.91 (m, 3H), 7.72 (d, J = 8.1 Hz, 1H), 7.63 - 7.60 (m, 1H),


7.11 (dd, *J* = 15.9, 9.6 Hz, 1H), 6.79 (d, *J* = 15.9 Hz, 1H), 4.98 - 4.91 (m, 1H); ESIMS *m/z* 484.94 ([M-H]$^-$); IR (thin film) 3444, 1705, 1171, 764 cm^{-1}.

(*E*)-2-Bromo-4-(4,4,4-trifluoro-3-(4-fluoro-3-(trifluoromethyl)phenyl)but-1-en-1-yl)benzoic acid




The title molecule was isolated as a brown liquid: ^{1}H NMR (300 MHz, $CDCl_3$) δ 8.00 (d, *J* = 8.1 Hz, 1H), 7.71 (s, 1H), 7.61-7.59 (m, 2H), 7.41 (d, *J* = 8.1 Hz, 1H), 7.30-7.24 (m, 1H), 6.59 (dd, *J* = 16.2, 6.0 Hz, 1H), 6.48 (d, *J* = 16.5 Hz, 1H), 4.26-4.21 (m, 1H); ESIMS *m/z* 469.0 ([M-H]$^-$); IR (thin film) 3444, 1699, 1327 cm^{-1}.

(*E*)-2-Bromo-4-(4,4,4-trifluoro-3-(3,4,5-trifluorophenyl)but-1-en-1-yl)benzoic acid




The title molecule was isolated as a brown gum: ^{1}H NMR (300 MHz, DMSO-d_6) δ 13.60 (bs, 1H), 7.97 (s, 2H), 7.72 (d, *J* = 7.2 Hz, 1H), 7.41 - 7.31 (m, 2H), 7.04 (dd, *J* = 15.6, 9.0 Hz, 1H), 6.71 (d, *J* = 15.9 Hz, 1H), 4.15 - 4 .11 (m, 1H); ESIMS *m/z* 438.8 ([M+H]$^+$).

(*E*)-4-(4,4,4-Trifluoro-3-(2,3,4-trifluorophenyl)but-1-en-1-yl)-2-(trifluoromethyl)benzoic acid




The title molecule was isolated as a brown gum: ^{1}H NMR (300 MHz, DMSO-d_6) δ 8.00 (s, 1H), 7.93 (d, *J* = 8.4 Hz, 1H), 7.81 (d, *J* = 8.1 Hz, 1H), 7.63 -7.60 (m, 1H), 7.47 - 7.44 (m, 1H), 7.02 - 7.01 (m, 1H), 5.10 - 4.90 (m, 1H).

(*E*)-2-Bromo-4-(4,4,4-trifluoro-3-(2,3,4-trifluorophenyl)but-1-en-1-yl)benzoic acid




The title molecule was isolated as a brown gum and the crude acid was taken on directly to the next step: ^{1}H NMR (300 MHz, DMSO-d_6) δ 13.65 (bs, 1H), 7.95 (s, 1H), 7.75 (d, J = 7.8 Hz, 1H), 7.62 - 7.59 (m, 2H), 7.50 (dd, J = 15.6, 9.0 Hz, 1H), 6.95 (d, J = 15.9 Hz, 1H), 4.86 - 4 .74 (m, 1H); ESIMS m/z 436.92 ([M-H]$^-$); IR (thin film) 3445, 1641, 1116 cm^{-1}.

(*E*)-4-(4,4,4-Trifluoro-3-(2,4,5-trichlorophenyl)but-1-en-1-yl)-2-(trifluoromethyl)benzoic acid




The title molecule was isolated as a brown gum: ^{1}H NMR (300 MHz, DMSO-d_6) δ 13.6 (s, 1H), 8.04 (s, 1H), 7.96 (d, J = 8.4 Hz, 3H), 7.83 (d, J = 8.1 Hz, 1H), 7.17 - 7.03 (m, 2H), 5.16 - 5.05 (m, 1H); ESIMS m/z 476.9 ([M-H]$^-$); IR (thin film) 3436, 1651, 1116, 661 cm^{-1}.

(*E*)-2-Bromo-4-(4,4,4-trifluoro-3-(2,4,5-trichlorophenyl)but-1-en-1-yl)benzoic acid




The title molecule was isolated as a brown gum: ^{1}H NMR (300 MHz, DMSO-d_6) δ (300 MHz, DMSO-d_6) δ 13.4 (s, 1H), 7.99 (d, J = 10.2 Hz, 3H), 7.76 (d, J = 8.1 Hz, 1H), 7.65 (d, J = 7.8 Hz, 1H), 7.09 - 6.91 (m, 2H), 5.11 - 5.05 (m, 1H); ESIMS m/z 486.8 ([M-H]$^-$) ; IR (thin film) 3436, 1651, 1115, 737 cm^{-1}.

(*E*)-4-(3-(4-Chloro-3-nitrophenyl)-4,4,4-trifluorobut-1-en-1-yl)-2-(trifluoromethyl)benzoic acid



The title molecule was isolated as a brown gum and the crude acid was taken on directly to the next step: ^{1}H NMR (300 MHz, DMSO-d_6) 13.80 (bs, 1H), 8.33 (s,1H), 7.94-7.81 (m, 5H), 7.75-7.72 (m, 1H), 7.06 (dd, J = 15.9, 8.7 Hz, 1H), 6.90 (d, J = 15.9 Hz, 1H), 5.02-4.81 (m, 1H).


(*E*)-2-Bromo-4-(3-(4-chloro-3-nitrophenyl)-4,4,4-trifluorobut-1-en-1-yl)benzoic acid



The title molecule was isolated as a brown gum: ^{1}H NMR (300 MHz, DMSO-d_6) 13.50 (bs, 1H), 8.31 (s, 1H), 8.00 - 7.77 (m, 3H), 7.75 -7.72 (m, 1H), 7.63 - 7.55 (m, 1H), 7.03 (dd, J = 15.9, 9.0 Hz, 1H), 6.81 (d, J = 15.9 Hz, 1H), 5.04 - 4.91 (m, 1H).


; ESIMS m/z 462.16 ([M-H]$^-$); IR (thin film) 3428, 1697, 1113, 749 cm^{-1}.

(*E*)-4-(4,4,4-Trifluoro-3-(4-fluoro-3,5-dimethylphenyl)but-1-en-1-yl)-2-(trifluoromethyl)benzoic acid




The title molecule was isolated as a brown gum: ^{1}H NMR (300 MHz, DMSO-d_6) δ 7.96 (s, 1H), 7.92 (d, J = 8.4 Hz, 1H), 7.80 - 7.75 (m, 1H), 7.27 (d, J = 6.9 Hz, 2H), 6.96 (dd, J = 15.6, 8.7 Hz, 1H), 6.87 (d, J = 15.6 Hz, 1H), 4.68-4.56 (m, 1H), 2.23 (s, 6H); ESIMS m/z 419.03 ([M-H]$^-$); IR (thin film) 3445, 2928, 1713, 1146 cm^{-1}.

(*E*)-2-Bromo-4-(4,4,4-trifluoro-3-(4-fluoro-3,5-dimethylphenyl)but-1-en-1-yl)benzoic acid



The title molecule was isolated as a brown gum: ^{1}H NMR (300 MHz, DMSO-d_6) δ 7.91 (s, 1H), 7.74 (d, J = 7.8 Hz, 1H), 7.61-7.58 (m, 1H), 7.26 (d, J = 6.6 Hz, 2H), 6.93 (dd, J = 15.9, 8.7 Hz, 1H), 6.87 (d, J = 15.9 Hz, 1H), 4.59-4.53 (m, 1H), 2.23 (s, 6H); ESIMS m/z 428.97 ([M-H]$^-$); IR (thin film) 3473, 1701, 1111, 581 cm^{-1}.


(*E*)-4-(4,4,4-Trifluoro-3-(4-fluoro-3-methylphenyl)but-1-en-1-yl)-2-(trifluoromethyl)benzoic acid



The title molecule was isolated as a brown liquid: ^{1}H NMR (300 MHz, DMSO-d_6) δ 13.58 (bs, 1H), 7.98 (s, 1H), 7.92 - 7.90 (m, 1H), 7.80 (d, J = 8.1 Hz, 1H), 7.48 - 7.45 (m,


1H), 7.42 - 7.37 (m, 1H), 7.22 - 7.16 (m, 1H), 7.04 (dd, *J* = 15.9, 8.7 Hz, 1H), 6.88 (d, *J* = 15.9 Hz, 1H), 4.70 - 4.60 (m, 1H), 4.04 - 3.99 (m, 1H), 2.26 (s, 3H); ESIMS *m/z* 405.05 ([M-H]$^-$); IR (thin film) 3437, 1710, 1145 cm^{-1}.

(*E*)-2-Bromo-4-(4,4,4-trifluoro-3-(4-fluoro-3-methylphenyl)but-1-en-1-yl)benzoic acid




The title molecule was isolated as a brown liquid: ^{1}H NMR (300 MHz, DMSO-d_6) δ 13.39 (bs, 1H), 7.91(s, 1H), 7.72 (d, *J* = 8.1 Hz, 1H), 7.61 - 7.58 (m 1H), 7.47 - 7.44 (m, 1H), 7.38 - 7.36 (m, 1H), 7.18 (t, *J* = 9.6 Hz, 1H), 6.95 (dd, *J* = 15.6, 8.7 Hz, 1H), 6.76 (d, *J* = 15.9 Hz, 1H), 4.67 - 4.61(m, 1H), 2.25 (s, 3H); ESIMS *m/z* 415.0 ([M-H]$^-$); IR (thin film) 3435, 2989, 1700, 1260 cm^{-1}.

(*E*)-4-(3-(3,5-Dichlorophenyl)-4,4,5,5,5-pentafluoropent-1-en-1-yl)-2-(trifluoromethyl)benzoic acid




The title molecule was isolated as a brown semi solid: ^{1}H NMR (400 MHz, DMSO-d_6) δ 13.70 (bs, 1H), 8.01 (s, 1H), 7.91(s, 1H), 7.80 (d, *J* = 8.4 Hz, 1H), 7.72 (*J* = 1.6 Hz, 2H), 7.66 (t, *J* = 3.2 Hz, 1H), 7.15 (dd, *J* = 15.6, 9.6 Hz, 1H), 6.91 (d, *J* = 15.6 Hz, 1H), 4.86-4.78 (m, 1H); ESIMS *m/z* 491.0 ([M-H]$^-$); IR (thin film) 3446, 1712,1141,749 cm^{-1}.

(*E*)-2-Bromo-4-(3-(3,5-dichlorophenyl)-4,4,5,5,5-pentafluoropent-1-en-1-yl)benzoic acid




The title molecule was isolated as a brown gum: ^{1}H NMR (400 MHz, DMSO-d_6) δ 7.85 (s, 1H), 7.70 (s, 2H), 7.65-7.64 (m, 1H), 7.56-7.52 (m, 2H), 6.94 (d, *J* = 9.2 Hz, 1H), 6.76 (d, *J* = 16 Hz, 1H), 4.82-4.80 (m, 1H); ESIMS *m/z* 500.8 ([M-H]$^-$); IR (thin film) 3422, 1683, 1184, 750, 575 cm^{-1}.

(*E*)-4-(3-(3,4-Dibromophenyl)-4,4,4-trifluorobut-1-en-1-yl)-2-(trifluoromethyl)benzoic acid




The title molecule was isolated as a brown gum: ^{1}H NMR (300 MHz, DMSO-d_6) δ 13.5 (bs, 1H), 8.01-7.99 (m, 2H), 7.94-7.91 (m, 1H), 7.85-7.78 (m, 2H), 7.53-7.50 (m, 1H), 7.09 (dd, J = 15.6, 8.7 Hz, 1H), 6.89 (d, J = 15.9 Hz, 1H), 4.85-4.78 (m, 1H); ESIMS m/z 528.8 ([M-H]$^-$); IR (thin film) 3437, 1722, 1168 cm^{-1}.

(*E*)-2-Bromo-4-(3-(3,4-dibromophenyl)-4,4,4-trifluorobut-1-en-1-yl)benzoic acid




The title molecule was isolated as a brown gum: ^{1}H NMR (300 MHz, DMSO-d_6) δ 13.38 (bs, 1H), 7.98-7.96 (m, 2H), 7.84 (d, J = 8.4 Hz, 1H), 7.74 (d, J = 8.1 Hz, 1H), 7.63-7.61 (m, 1H), 7.51-7.49 (m, 1H), 7.01 (dd, J = 15.9, 9.0 Hz, 1H), 6.78 (d, J = 15.6 Hz, 1H), 4.82-4.76 (m, 1H); ESIMS m/z 538.8 ([(M-H]$^-$); IR (thin film) 3446, 1699, 1166, 581 cm^{-1}.

(*E*)-4-(4,4,4-Trifluoro-3-(3-(trifluoromethoxy)phenyl)but-1-en-1-yl)-2-(trifluoromethyl)benzoic acid




The title molecule was isolated as a brown semi solid: ^{1}H NMR (300 MHz, DMSO-d_6) δ 8.01(s, 1H), 7.94 (d, J = 8.7 Hz, 1H), 7.80 (d, J = 8.1 Hz, 1H), 7.63-7.55 (m, 3H), 7.41(d, J = 7.5 Hz, 1H), 7.11(dd, J = 15.6, 9.0 Hz, 1H), 6.92 (d, J = 15.9 Hz, 1H), 4.89-4.82 (m, 1H); ESIMS m/z 456.98 ([M-H]$^-$); IR (thin film) 3413, 1668, 1161 cm^{-1}.

(*E*)-2-Bromo-4-(4,4,4-trifluoro-3-(3-(trifluoromethoxy)phenyl)but-1-en-1-yl)benzoic acid




The title molecule was isolated as a brown solid: ^{1}H NMR (300 MHz, DMSO-d_6) δ 7.73 (s, 1H), 7.59 (m, 3H), 7.44 (s, 1H), 7.40 (d, J = 7.6 Hz, 2H), 6.88 (dd, J = 15.6, 9.0 Hz, 1H), 6.73 (d, J = 15.9 Hz, 1H), 4.85-4.82 (m, 1H); ESIMS m/z 466.93 ([M-H]$^-$); IR (thin film) 3437, 1703, 1111 cm^{-1}.

(*E*)-4-(3-(3-Cyano-4-fluorophenyl)-4,4,4-trifluorobut-1-en-1-yl)-2-(trifluoromethyl)benzoic acid




The title molecule was isolated as a brown liquid: ^{1}H NMR (300 MHz, DMSO-d_6) δ 13.60 (bs, 1H), 8.21-8.19(m, 1H), 8.01-7.91(m, 3H), 7.81 (d, J = 8.4 Hz, 1H), 7.12 (dd, J = 15.9,8.1 Hz, 1H), 6.91 (d, J = 15.6 Hz, 1H), 4.92-4.86 (m, 1H); ESIMS m/z 416.27 ([M-H]$^-$); IR (thin film) 3429, 2238, 1713, 1116 cm^{-1}.

(*E*)-2-Bromo-4-(3-(3-cyano-4-fluorophenyl)-4,4,4-trifluorobut-1-en-1-yl)benzoic acid




The title molecule was isolated as a brown gum: ^{1}H NMR (300 MHz, DMSO-d_6) δ 13.56 (bs, 1H), 8.21 - 8.18 (m, 1H), 8.00 -7 .95(m, 2H), 7.73 - 7.59 (m, 3H), 7.03 (dd, J = 15.9, 9.3 Hz, 1H), 6.79 (d, J = 15.3 Hz, 1H), 4.87 - 4.84 (m, 1H); ESIMS m/z 426.0 ([M-H]$^-$).

(*E*)-2-Bromo-4-(3-(3,4-dichlorophenyl)-4,4,4-trifluorobut-1-en-1-yl)benzoic acid




The title molecule was isolated as a brown gum: ^{1}H NMR (300 MHz, DMSO-d_6) δ 13.4 (s, 1H),7.96 (d, J = 1.2 Hz, 1H), 7.88 (d, J = 1.8 Hz, 1H), 7.74 - 7.68 (m, 2H), 7.63 (dd, J = 8.1, 1.2 Hz, 1H), 7.57 (dd, J = 8.4, 1.8 Hz, 1H), 7.02 (dd, J = 15.9, 9.3 Hz, 1H), 6.78 (dd, J = 5.9 Hz, 1H), 4.84 - 4.78 (m, 1H); ESIMS m/z 451.0 ([M-H]$^-$); IR (thin film) 3445, 1704, 1113, 740 cm^{-1}.

(*E*)-4-(3-(3-Bromo-5-chlorophenyl)-4,4,4-trifluorobut-1-en-1-yl)-2-(trifluoromethyl)benzoic acid




The title molecule was isolated as a brown solid: ^{1}H NMR (300 MHz, DMSO-d_6) δ 13.50 (bs, 1H), 7.91 (s,1H), 7.86 - 7.64 (m, 5H), 7.06 (dd, J = 15.9, 9.0 Hz, 1H), 6.87 (d, J = 15.9 Hz, 1H), 4.85 - 4.78 (m, 1H); ESIMS m/z 485.17 ([M-H]$^-$); IR (thin film) 3438, 1708, 1114, 774, 516 cm^{-1}.

(*E*)-2-Bromo-4-(3-(3-bromo-5-chlorophenyl)-4,4,4-trifluorobut-1-en-1-yl)benzoic acid




The title molecule was isolated as a brown gum: ^{1}H NMR (300 MHz, DMSO-d_6) δ 13.38 (bs, 1H), 7.98 (s, 1H), 7.80 - 7.72 (m, 4H), 7.64 - 7.61 (m, 1H), 7.06 (dd, J = 15.9, 9.3 Hz, 1H), 6.79 (d, J = 15.6 Hz, 1H), 4.88 - 4.80 (m, 1H); ESIMS m/z 495.05 ([M-H]$^-$); IR (thin film) 3436, 1699 ,1116, 750, 531 cm^{-1}.

(*E*)-4-(3-(3-Bromo-5-fluorophenyl)-4,4,4-trifluorobut-1-en-1-yl)-2-(trifluoromethyl)benzoic acid




The title molecule was isolated as a brown liquid: ^{1}H NMR (300 MHz, DMSO-d_6) δ 13.6 (bs, 1H), 8.02 (s, 1H), 7.91 - 7.89 (m, 1H), 7.81 (d, J = 8.0 Hz, 1H), 7.69 (s, 1H), 7.63 - 7.59 (m, 1H), 7.55 (d, J = 9.3 Hz, 1H), 7.11 (dd, J = 15.9, 9.0 Hz, 1H), 6.91 (d, J = 15.9 Hz, 1H), 4.87 - 4.80 (m, 1H); ESIMS m/z 469.07 ([M-H]$^-$); IR (thin film) 3428, 1712, 1171, 523 cm^{-1}.

(*E*)-4-(4,4,4-Trifluoro-3-(3,4,5-trichlorophenyl)but-1-en-1-yl)benzoic acid




The title molecule was isolated as a yellow solid: ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.18 – 8.03 (m, 2H), 7.49 (d, J = 8.3 Hz, 2H), 7.42 (s, 2H), 6.66 (d, J = 15.9 Hz, 1H), 6.47 (dd, J = 15.9, 8.0 Hz, 1H), 4.13 (*p*, J = 8.6 Hz, 1H); ^{19}F NMR (376 MHz, $CDCl_3$) δ -68.65; ESIMS m/z 409.1 ([M-H]$^-$).

(*E*)-2-Bromo-4-(3-(3-chloro-4-methylphenyl)-4,4,4-trifluorobut-1-en-1-yl)benzoic acid




The title molecule was isolated as a brown liquid: ^{1}H NMR (300 MHz, DMSO-d_6) δ 13.30 (bs, 1H), 7.93 (d, J = 1.2 Hz, 1H), 7.42 (d, J = 8.1 Hz, 1H), 7.62 (dd, J = 1.5, 8.1 Hz, 1H), 7.53 (s, 1H), 7.48 (d, J = 7.8 Hz, 1H), 7.39 (d, J = 7.8 Hz, 1H), 6.96 (dd, J = 15.6, 8.7 Hz, 1H), 6.77 (d, J = 15.6 Hz, 1H), 4.73 - 4.61 (m, 1H), 2.35 (s, 3H); ESIMS m/z 431.77 ([M-H]$^-$); IR (thin film) 3435, 1701, 1111, 750 cm^{-1}.

(*E*)-4-(3-(3-Chloro-4-methylphenyl)-4,4,4-trifluorobut-1-en-1-yl)-2-(trifluoromethyl)benzoic acid




The title molecule was isolated as a brown gum: ^{1}H NMR (300 MHz, DMSO-d_6) δ 13.50 (bs, 1H), 7.98 (s, 1H), 7.92 (d, J = 8.1 Hz, 1H), 7.80 (d, J = 8.1 Hz, 1H), 7.53 (s, 1H), 7.48 (d, J = 8.1 Hz, 1H), 7.40 (d, J = 8.4 Hz, 1H), 7.04 (dd, J = 15.6, 8.4 Hz, 1H), 6.88 (d, J = 15.6 Hz, 1H), 4.72 - 4.66 (m, 1H), 2.35 (s, 3H); ESIMS m/z 421.82 ([M-H]$^-$); IR (thin film) 3460, 2926, 1712, 1170, 750 cm^{-1}.

(*E*)-4-(4,4,5,5,5-Pentafluoro-3-(3,4,5-trichlorophenyl)pent-1-en-1-yl)-2-(trifluoromethyl)benzoic acid




The title molecule was isolated as a dark brown gum: ^{1}H NMR (300 MHz, DMSO-d_6) δ 13.6 (bs, 1H), 8.03 (s, 1H), 7.95 - 7.86 (m, 3H), 7.81 (d, J = 8.1 Hz, 1H), 7.16 (dd, J = 15.3, 9.3 Hz, 1H), 6.92 (d, J = 15.6 Hz, 1H), 4.95 - 4.88 (m, 1H); ^{19}F NMR (300 MHz, DMSO-d_6) δ -80.35, -58.02; ESIMS m/z 526.8 ([M+H]$^+$).

(*E*)-2-Bromo-4-(4,4,5,5,5-pentafluoro-3-(3,4,5-trichlorophenyl)pent-1-en-1-yl)benzoic acid




The title molecule was isolated as a dark brown gum: ^{1}H NMR (300 MHz, DMSO-d_6) δ 13.6 (bs,1H), 7.94 (s, 2H), 7.78 (d, J = 7.8 Hz, 1H), 7.71 (d, J = 7.8 Hz, 1H), 7.60 (d, J = 7.5 Hz 1H), 7.07 (dd, J = 15.0, 8.7 Hz, 1H), 6.79 (d, J = 15.6 Hz, 1H), 4.93 - 4.78 (m, 1H); ESIMS m/z 538.9 ([M+H]$^+$); IR (thinfilm) 3420, 1602, 1123, 746 cm^{-1}.

(*E*)-2-Bromo-4-(3-(4-cyano-3,5-difluorophenyl)-4,4,4-trifluorobut-1-en-1-yl)benzoic acid




The title molecule was isolated as a brown gum: ESIMS m/z 443.91 ([M-H]$^-$) ; IR (thin film) 3447, 2244, 1703, 1114 cm^{-1}.

(*E*)-2-Chloro-4-(3-(3,5-dibromophenyl)-4,4,4-trifluorobut-1-en-1-yl)benzoic acid




The title molecule was isolated as a brown liquid: ^{1}H NMR (300 MHz, DMSO-d_6) δ 13.39 (bs, 1H), 7.95-7.70 (m, 5H), 7.61 (d, J = 8.1 Hz, 1H), 7.07 (dd, J = 15.6, 9.3 Hz, 1H), 6.80 (d, J = 15.6 Hz, 1H), 4.84-4.78 (m, 1H); ESIMS m/z 496.77 ([M-H]$^-$); ; IR (thin film) 3439, 2920, 1707, 1165 cm^{-1}.

(*E*)-4-(3-(3,5-Dichloro-4-fluorophenyl)-4,4,4-trifluorobut-1-en-1-yl)-2-(trifluoromethyl)benzoic acid




The title molecule was isolated as an off white solid: mp 140-143 °C; ^{1}H NMR (400 MHz, DMSO) δ13.60 (bs, 1H), 8.02 (s, 1H), 7.94 - 7.90 (m, 1H), 7.88 -7.86 (m, 2H), 7.81 - 7.79 (m, 1H), 7.12 (dd, J = 15.6, 8.8 Hz, 1H), 6.89 (d, J = 15.6 Hz, 1H), 4.86 - 4.81 (m, 2H); ESIMS m/z 458.88 ([M-H]$^-$).

(*E*)-4-(3-(3,4-Dichlorophenyl)-4,4,4-trifluorobut-1-en-1-yl)benzoic acid




The title molecule was isolated as a light orange crystalline solid (875 mg, 88%): ^{1}H NMR (400 MHz, $CDCl_3$) δ 12.35 (s, 1H), 8.08 (d, *J* = 8.4 Hz, 2H), 7.55 - 7.41 (m, 4H), 7.24 (dd, *J* = 8.3, 2.1 Hz, 1H), 6.64 (d, *J* = 15.8 Hz, 1H), 6.51 (dd, *J* = 15.9, 7.7 Hz, 1H), 4.15 (p, *J* = 8.7 Hz, 1H); ^{19}F NMR 376 MHz, $CDCl_3$) δ -68.75; ESIMS *m/z* 375 ($[M+H]^+$).

(*E*)-4-(3-(3,4-Dichlorophenyl)-4,4,4-trifluorobut-1-en-1-yl)-2-(trifluoromethyl)benzoic acid




The title molecule was isolated was isolated as a brown gum: ^{1}H NMR (400 MHz, DMSO-d_6) δ 13.6 (s, 1H), 8.02 (s, 1H), 7.93 - 7.89 (m, 2H), 7.80 (d, *J* = 7.6 Hz, 1H), 7.73 (d, *J* = 8.4, Hz, 1H), 7.58 (dd, *J* = 8.4, 2.0 Hz, 1H), 7.09 (dd, *J* = 15.6, 8.8, Hz, 1H), 6.89 (d, *J* = 15.6, Hz, 1H), 4.86 - 4.81 (m, 1H); ESIMS *m/z* 441.0 ($[M-H]^-$); IR (thinfilm) 3447, 1710, 1169, 749 cm^{-1}.

(*E*)-4-(3-(3,5-Dichlorophenyl)-4,4,4-trifluorobut-1-en-1-yl)-2-(trifluoromethyl)benzoic acid




The title molecule was isolated was isolated as a brown gum: ^{1}H NMR (300 MHz, DMSO-d_6) δ 13.6 (bs, 1H), 7.98 (s, 1H), 7.91 (d, *J* = 7.8 Hz 1H), 7.75 - 7.66 (m, 1H), 7.10 (dd, *J* = 15.6, 9.0 Hz, 1H), 6.89 (d, *J* = 15.9 Hz 1H), 4.86 - 4.80 (m, 1H); ESIMS *m/z* 441.1 ($[M-H]^-$); IR (thinfilm) 3460, 2928, 1721, 1170, 764 cm^{-1}.

(*E*)-4-(3-(3,4-Dichloro-5-methylphenyl)-4,4,4-trifluorobut-1-en-1-yl)-2-(trifluoromethyl)benzoic acid




The title molecule was isolated as a pale yellow semi solid: ^{1}H NMR (400 MHz, DMSO-d_6) δ 13.58 (bs, 1H), 8.00 (s, 1H), 7.93 (d, *J* = 8.4 Hz, 1H), 7.80 (d, *J* = 8.4 Hz, 1H), 7.72 (s, 1H), 7.55 (s, 1H), 7.07 (dd, *J* = 16.4, 9.6 Hz, 1H), 6.89 (d, *J* = 15.6 Hz, 1H), 4.78 - 4.73 (m, 1H), 2.42 (s, 3H); ESIMS *m/z* 455.0 ($[M-H]^-$); IR (thin film) 1713, 1170, 750 cm^{-1}.

(*E*)-2-Bromo-4-(3-(3,4-dichlorophenyl)-4,4-difluoropent-1-en-1-yl)benzoic acid




The title molecule was isolated as a brown gum: ^{1}H NMR (400 MHz, DMSO-d_6) δ 13.3 (s, 1H), 7.92 (s, 1H), 7.77 - 7.71 (m, 2H), 7.68 - 7.63 (m, 1H), 7.61 - 7.60 (m, 1H), 7.60 - 7.58 (m, 1H), 6.98 (dd, *J* = 15.6, 9.2 Hz, 1H), 6.65 (d, *J* = 15.6 Hz, 1H), 4.83 - 4.80 (m, 1H), 1.59 – 1.54 (m, 3H); ESIMS *m/z* 448.8 ([M-H]$^-$).

(*E*)-2-Bromo-4-(3-(3-chloro-5-ethylphenyl)-4,4,4-trifluorobut-1-en-1-yl)benzoic acid




The title molecule was isolated as a brown liquid: ^{1}H NMR (400 MHz, DMSO-d_6) δ 13.4 (bs, 1H), 7.97 (s, 2H), 7.91 (s, 1H), 7.74 (d, *J* = 8.4 Hz, 2H), 7.66 -7.61 (m, 1H), 7.03 (dd, *J* = 16.0, 8.4 Hz , 1H), 6.8 (d, *J* = 15.6 Hz, 1H), 4.89 - 4.84 (m, 1H), 2.66 - 2.65 (m, 2H), 1.25 (t, *J* = 9.2 Hz, 3H); ESIMS *m/z* 446.8 ([M+H]$^+$).

(*E*)-2,6-Dimethyl-4-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-enyl)benzoic acid.




The title molecule was isolated as a brown gum: ^{1}H NMR (300 MHz, DMSO-d_6) δ 13.1 (s, 1H), 7.87 (s, 2H), 7.27 (s, 2H), 6.81 (dd, *J* = 15.6, 8.7 Hz, 1H), 6.69 (d, *J* = 15.3 Hz, 1H), 4.85 - 4.79 (m, 1H), 2.27 (s, 6H); ESIMS *m/z* 437.01 ([M-H]$^-$); IR (thin film) 3285, 1621, 1162, 954 cm^{-1}.

(*E*)-2-Bromo-4-(3-(3,5-dibromo-4-chlorophenyl)-4,4,4-trifluorobut-1-en-1-yl)benzoic acid




The title molecule was isolated as a brown gum: ^{1}H NMR (300 MHz, DMSO-d_6) δ 13.40 (bs, 1H), 8.07(d, *J* = 7.5 Hz, 1H), 7.94-7.89 (m, 2H), 7.66 -7.60 (m, 2H), 7.10 (dd, *J* = 8.7, 16.0 Hz, 1H), 6.96 (d, *J* = 15.6 Hz, 1H), 4.82-4.80 (m, 1H); ESIMS *m/z* 574.7 ([M+H]$^+$).

(*E*)-4-(3-(3,5-Dibromo-4-chlorophenyl)-4,4,4-trifluorobut-1-en-1-yl)-2-(trifluoromethyl)benzoic acid




The title molecule was isolated as a brown gum: ^{1}H NMR (300 MHz, DMSO-d_6) δ 13.36 (bs, 1H) 8.05(s, 2H), 7.95 (d, *J* = 8.1 Hz, 1H), 7.87-7.67 (m, 2H), 7.14 (dd, *J* = 9.0, 15.6 Hz, 1H), 6.96 (d, *J* = 15.6 Hz, 1H), 4.88-4.82 (m, 1H); ESIMS *m/z* 564.58 ([M+H]$^+$).

(*E*)-2-Bromo-4-(3-(4-bromo-3,5-dichlorophenyl)-4,4,4-trifluorobut-1-en-1-yl)benzoic acid




The title molecule was isolated as a brown gum: ^{1}H NMR (300 MHz, DMSO-d_6) δ 13.40 (bs, 1H), 7.98 (s, 1H), 7.87 (s, 2H), 7.75 (d, *J* = 8.1 Hz, 1H), 7.65 - 7.62 (m, 1H), 7.06 (dd, *J* = 15.9, 9.3 Hz, 1H), 6.80 (d, *J* = 15.9 Hz, 1H), 4.87 - 4.80 (m, 1H); ESIMS *m/z* 518.9 ([M-H]$^-$).

(*E*)-4-(3-(4-Bromo-3,5-dichlorophenyl)-4,4,4-trifluorobut-1-en-1-yl)-2-(trifluoromethyl)benzoic acid




The title molecule was isolated as a brown gum: ^{1}H NMR (300 MHz, DMSO-d_6) δ 13.6 (bs, 1H) 8.03 (s, 1H), 7.95 (d, *J* = 8.4 Hz, 1H), 7.88 (s, 2H), 7.81 (d, *J* = 8.1 Hz, 1H), 7.13 (dd, *J* = 16.2, 7.5 Hz, 1H), 6.91 (d, *J* = 15.9 Hz, 1H), 4.89 - 4.83 (m, 1H); ESIMS *m/z* 532.0 ([M+H]$^+$).

(*E*)-2-Bromo-4-(3-(3-chloro-4-(trifluoromethoxy)phenyl)-4,4,4-trifluorobut-1-en-1-yl)benzoic acid



The title molecule was isolated as a brown gum: ^{1}H NMR (400 MHz, DMSO-d_6) δ 13.36 (bs, 1H) 7.95 (s, 1H), 7.73 (d, *J* = 7.6 Hz, 1H), 7.63 (d, *J* = 8.1 Hz, 1H), 7.46 (s, 1H) 7.35-7.31(m, 2H), 7.04 (dd, *J* = 16.0, 8.8 Hz, 1H), 6.78 (d, *J* = 16.4 Hz, 1H), 4.71 - 4.68 (m, 1H); ESIMS *m/z* 500.8 ([M-H]$^-$).


Example 20: Preparation of 5-vinyl-2,3-dihydro-1*H*-inden-1-one (BI1)



To a stirred solution of 5-bromo-2,3-dihydro-1*H*-inden-1-one (5 g, 23.7 mmol) in toluene were added vinylboronic anhydride pyridine complex (8.55 g, 35.54 mmol), $Pd(PPh_3)_4$ (0.1 g, 0.094 mmol), K_2CO_3 (22.88 g, 165.83 mmol). The resultant reaction mixture was heated at reflux for 16 h. The reaction mixture was cooled to 25 °C and filtered, and the filtrate was concentrated under reduced pressure. The residue was diluted with EtOAc and washed with water and brine. The combined organic extracts were dried over anhydrous Na_2SO_4 and concentrated under reduced pressure. The obtained residue was purified by flash column chromatography (SiO_2, 5% EtOAc in petroleum ether) afforded the title compound as a solid (1.8 g, 48%): ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.74 (d, *J* = 7.2 Hz, 1H), 7.49 (br s, 1H), 7.44 (d, *J* = 7.2 Hz, 1H), 6.82 (m, 1H), 5.90 (d, *J* = 7.4 Hz, 1H), 5.42 (d, *J* = 6.4 Hz, 1H), 3.20 (m, 2H), 2.70 (m, 2H); ESIMS *m/z* 159.06 ([M+H]$^-$).

The following compound was made in accordance with the procedures disclosed in **Example 20**.


6-Vinyl-3,4-dihydronaphthalen-1(2*H*)-one (BI2)



The product was isolated as an off-white solid (5 g, 48%): ^{1}H NMR (400 MHz, DMSO-d_6) δ 7.85 (d, *J* = 8.4 Hz, 1H), 7.48 (m, 2H), 6.82 (m, 1H), 6.02 (d, *J* = 7.4 Hz, 1H),

5.44 (d, J = 6.4 Hz, 1H), 2.95 (m, 2H), 2.60 (m, 2H), 2.00 (m, 2H); ESIMS m/z 173.14 ([M-H]$^-$); IR (thin film) 1681 cm^{-1}.


Example 21: Preparation of (*E*)-5-(4,4,4-Trifluoro-3-(3,4,5-trichlorophenyl)but-1-enyl)-2,3-dihydro-1*H*-inden-1-one (BI3)



5-(1-Bromo-2,2,2-trifluoroethyl)-1,2,3-trichlorobenzene (4 g, 11.7 mmol), 5-vinyl-2,3-dihydro-1*H*-inden-1-one (0.92 g, 5.8 mmol), CuCl (0.115 g, 1.171 mmol) and 2,2-bipyridyl (0.053 g, 0.34 mmol) in 1,2-dichlorobenzene (25 mL) were heated at 180 °C for 16 h. The reaction mixture was cooled to 25 °C and concentrated under reduced pressure. The residue was purified by flash column chromatography (SiO_2, 5% EtOAc in petroleum ether) to afford the title compound as a liquid (1.28 g, 25%): ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.76 (d, J = 7.4 Hz, 1H), 7.52 (m, 3H), 6.68 (d, J = 7.4 Hz, 1H), 6.52 (m, 1H), 4.18 (m, 1H), 3.18 (m, 2H), 2.75 (m, 2H); ESIMS m/z 419.14 ([M+H]$^-$); IR (thin film) 1708.94, 1113.60, 807.77 cm^{-1}.


The following compound was made in accordance with the procedures disclosed in **Example 21**.

(E)-5-(3-(3,5-Dichloro-4-fluorophenyl)-4,4,4-trifluorobut-1-en-1-yl)-2,3-dihydro-1*H*-inden-1-one (BI4)




The product was isolated as a brown semi-solid (1.2 g, 16%): ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.76 (d, J = 7.4 Hz, 1H), 7.54 (m, 3H), 7.30 (s, 1H), 6.68 (d, J = 7.4 Hz, 1H), 6.52 (m, 1H), 4.18 (m, 1H), 3.18 (m, 2H), 2.75 (m, 2H); ESIMS m/z 400.84 ([M-H]$^-$); IR (thin film) 815, 1113, 1709 cm^{-1}.

(*E*)-6-(4,4,4-Trifluoro-3-(3,4,5-trichlorophenyl)but-1-enyl)-3,4-dihydronaphthalen-1(2*H*)-one (BI5)




The product was isolated as a pale yellow semi solid (1.2 g, 30%): ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.20 (d, *J* = 8.0 Hz, 1H), 7.42 (s, 2H), 7.35 (m, 1H), 7.24 (m, 2H), 6.62 (d, *J* = 16 Hz, 1H), 6.46 (m, 1H), 4.18 (m, 1H), 2.95 (m, 2H), 2.65 (m, 2H), 2.19 (m, 2H); ESIMS *m/z* 432.94 ([M-H]$^-$); IR (thin film) 1680, 1113, 808 cm^{-1}.

Example 22: Preparation of (E)-5-(3-(3,5-Dichloro-4-fluorophenyl)-4,4,4-trifluorobut-1-en-1-yl)-2-fluoro-2,3-dihydro-1*H*-inden-1-one (BI6)



To a stirred solution of (E)-5-(3-(3,5-dichloro-4-fluorophenyl)-4,4,4-trifluorobut-1-enyl)-2,3-dihydro-1*H*-inden-1-one (0.5 g, 1.24 mmol) in MeCN (20 mL), was added Selectfluor® (0.52 g, 1.48 mmol) and the reaction was heated to reflux temperature for 16 h. The reaction mixture was cooled to room temperature, concentrated under reduced pressure and diluted with CH_2Cl_2. The solution was washed with water and brine, dried over anhydrous Na_2SO_4 and concentrated under reduced pressure to give the crude product which was purified by flash column chromatography (SiO_2, 100-200 mesh; 15% EtOAc in petroleum ether) to afford the title compound as a pale yellow semi solid (0.1g, 24%): ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.80 (m, 1H), 7.48 (m, 2H), 7.32 (m, 2H), 6.65 (d, *J* = 16.0 Hz, 1H), 6.54 (dd, *J* = 16.0, 8.0 Hz, 1H), 5.38 (m, 1H), 4.18 (m, 1H), 3.62 (m, 1H), 3.32 (m, 1H); ESIMS *m/z* 419.06 ([M-H]$^-$); IR (thin film) 1728, 1114, 817 cm^{-1}.


Example 23: Preparation of (E)-5-(3-(3,5-Dichloro-4-fluorophenyl)-4,4,4-trifluorobut-1-en-1-yl)-*N*-(3,3,3-trifluoropropyl)-2,3-dihydro-1H-inden-1-amine (BC10)



To a stirred solution of (E)-5-(3-(3,5-dichloro-4-fluorophenyl)-4,4,4-trifluorobut-1-enyl)-2,3-dihydro-1*H*-inden-1-one (0.15 g, 0.35 mmol) in DCE (10 mL), was added


trifluoropropyl amine (0.048 g, 0.42 mmol) and $NaBH_3CN$ (0.055 g, 0.875 mmol) in cooling and the reaction mixture was stirred at room temperature for 16 h. The reaction mixture was diluted with DCE, was washed with water and brine and dried over anhydrous Na_2SO_4. Concentration under reduced pressure gave the crude compound, which was purified by flash column chromatography (SiO_2, 100-200 mesh; 10-15% EtOAc in petroleum ether) to afford the title compound as a colorless gummy material (0.042g, 24%): ^1H NMR (400 MHz, $CDCl_3$) δ 7.38-7.20 (m, 5H), 6.62 (d, *J* = 16.0 Hz, 1H), 6.34 (dd, *J* = 16.0, 8.0 Hz, 1H), 5.83 (br, 1H), 5.52 (m, 1H), 4.12 (m, 1H), 3.02 (m, 3H), 2.82 (m, 1H), 2.50 (m, 2H), 1.82 (m, 1H), 1.42 (m, 1H); ESIMS *m/z* 497.98 ($[M-H]^-$); IR (thin film) 3027, 1654, 815 cm^{-1}.

Example 24: Preparation of 6-((*E*)-4,4,4-Trifluoro-3-(3,4,5-trichlorophenyl)but-1-enyl)-3,4-dihydronaphthalen-1(2*H*)-one oxime (BI5a)



To a stirred solution of ((*E*)-6-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-enyl)-3,4-dihydronaphthalen-1(2*H*)-one (0.4 g, 0.92 mmol) in EtOH (50 mL) were added hydroxylamine hydrochloride (0.128 g, 1.85 mmol) and sodium acetate (NaOAc, 0.23 g, 2.77 mmol), and the reaction mixture was heated at reflux for 3 h. The reaction mixture was concentrated under reduced pressure, and the residue was diluted with water and extracted with EtOAc. The combined organic extracts were washed with brine, dried over anhydrous Na_2SO_4 and concentrated under reduced pressure to give the crude compound, which was purified by flash column chromatography (SiO_2, 100-200 mesh; 10-15% EtOAc in petroleum ether). The title compound was isolated as a solid (0.3 g, 73%): mp 155–158 °C; ^1H NMR (400 MHz, $CDCl_3$) δ 7.89 (d, *J* = 8.4 Hz, 1H), 7.41 (s, 2H), 7.24 (m, 1H), 7.17 (m, 1H), 6.57 (d, *J* = 16 Hz, 1H), 6.46 (dd, *J* = 16.0, 8.0 Hz, 1H), 4.13 (m, 1H), 2.82 (m, 4H), 2.04 (m, 2H); ESIMS *m/z* 445.95 ($[M-H]^-$).


Example 25: Preparation of (*E*)-5-(4,4,4-Trifluoro-3-(3,4,5-trichlorophenyl)but-1-enyl)-2,3-dihydro-1*H*-inden-1-amine (BI5b)



To a stirred solution of (*E*)-5-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-enyl)-2,3-dihydro-1*H*-inden-1-one (1 g, 2.39 mmol) in MeOH (10 mL) were added ammonium acetate (NH_4OAc, 1.84 g, 23.9 mmol) and $NaBH_3CN$ (0.44 g, 7.17 mmol,) and the reaction mixture was heated at reflux for 16 h. The reaction mixture was concentrated under reduced pressure, and the residue was diluted with water and extracted with EtOAc . The combined organic extracts were washed with water and saturated aqueous $NaHCO_3$ solution, dried over anhydrous Na_2SO_4, and concentrated under reduced pressure to afford the title compound as a liquid (500 mg, crude): ^{1}H NMR (400 MHz, DMSO-d_6) δ 7.85 (s, 2H), 7.40 (s, 1H), 7.30 (s, 2H), 6.71 (s, 2H), 4.78 (m, 1H), 4.2 (m, 1H), 2.80 (m, 1H), 2.73 (m, 1H), 1.60 (m, 2H); ESIMS *m/z* 419.02 ($[M+H]^+$); IR (thin film) 2924, 1552, 1112, 807 cm^{-1}.


The following compound was made in accordance with the procedures disclosed in **Example 25**.

(*E*)-5-(3-(3,5-Dichloro-4-fluorophenyl)-4,4,4-trifluorobut-1-en-1-yl)-2,3-dihydro-1*H*-inden-1-amine (BI7)




The product was isolated as a light brown gummy material, taken as such to the next step (0.15 g, crude compound): ESIMS *m/z* 401.97 ($[M-H]^-$).

(*E*)-5-(3-(3,5-Dichloro-4-fluorophenyl)-4,4,4-trifluorobut-1-en-1-yl)-2-fluoro-2,3-dihydro-1*H*-inden-1-amine (BI8)




The product was isolated as a light brown gummy material, taken as such to the next step (0.15 g, crude compound): ESIMS *m/z* 420.15 ($[M-H]^-$).

(*E*)-6-(4,4,4-Trifluoro-3-(3,4,5-trichlorophenyl)but-1-enyl)-1,2,3,4-tetrahydronaphthalen-1-amine (BI9)



The product was isolated as a pale yellow liquid (500 mg crude).


Example 26: Preparation of (*E*)-1-Methyl-3-(5-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)-but-1-enyl)-2,3-dihydro-1*H*-inden-1-yl)thiourea (BC1)



To a stirred solution of (*E*)-5-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-enyl)-2,3-dihydro-1*H*-inden-1-amine (0.1 g, 0.23 mmol) in Et_2O (5 mL) was added methylisothiocyanate (0.026 g, 0.35 mmol), and the mixture was stirred for 2 h at 25 °C. The reaction mixture was concentrated under reduced pressure, and the residue was purified by flash column chromatography (SiO_2, 20% EtOAc in petroleum ether). The title compound was isolated as a liquid (65 mg, 50%): ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.39 (s, 2H), 7.25 – 7.18 (m, 3H), 6.58 (d, *J* = 16.0 Hz, 1H), 6.30 (dd, *J* = 16.0, 8.4 Hz, 1H), 5.91 – 5.70 (br, 2H), 4.05 (m, 1H), 3.05 – 2.80 (m, 6H), 2.70 (m, 1H), 1.81 (m, 1H); ESIMS *m/z* 492.17 ($[M+H]^+$); IR (thin film) 3211, 1569, 1113, 806 cm^{-1}.

Compounds **BC2** – **BC3** in Table 1 were made in accordance with the procedures disclosed in **Example 26**.

Example 27: Preparation of (*E*)-3,3,3-Trifluoro-*N*-(5-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-enyl)-2,3-dihydro-1*H*-inden-1-yl)propanamide (BC4)




To a stirred solution of (*E*)-5-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-enyl)-2,3-dihydro-1*H*-inden-1-amine (0.1 g, 0.23 mmol) in CH_2Cl_2 (10 mL) were added

trifluoropropionic acid (0.044 g, 0.34 mmol), EDC•HCl (0.038 g, 0.35 mmol), HOBt•H_2O (0.07 g, 0.46 mmol) and DIPEA (0.074 g, 0.57 mmol), and the reaction mixture was stirred for 16 h at 25 °C. The reaction mixture was diluted with CH_2Cl_2 and washed with water. The combined organic layer was washed with brine, dried over anhydrous Na_2SO_4, and concentrated under reduced pressure. The crude material was purified by flash column chromatography (SiO_2, 15% EtOAc in petroleum ether) to afford the title compound as a liquid (65 mg, 65%): ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.39 (s, 2H), 7.25-7.20 (m, 3H), 6.34 (d, J = 16.0 Hz, 1H), 6.30 (dd, J = 16.0, 8.0 Hz, 1H), 5.81 (br, 1H), 5.48 (m, 1H), 4.10 (m, 1H), 3.10 (m, 2H), 2.86-3.07 (m, 2H), 2.86 (m, 1H), 1.81 (m, 1H); ESIMS *m/z* 529.02 ([M+H]$^+$); IR (thin film) 3283, 1652, 1241, 811 cm^{-1}.

Compounds **BC5 – BC9, BC11** in Table 1 were made in accordance with the procedures disclosed in **Example 27**.

Example 28: Preparation of *tert*-Butyl 5-vinylindoline-1-carboxylate (BI10)




Step 1. 5-Bromo-indoline (BI11): To 5-Bromo-1*H*-indole (2.5 g, 12.82 mmol) in AcOH (10.0 mL), $NaBH_3CN$ (2.38 g, 38.46 mmol) was added portion wise at 10 °C over the period of 20 min. After that the reaction mixture was stirred at ambient temperature for 3 h. The reaction mixture was diluted with water and extracted with Et_2O. The organic layer was washed with saturated $NaHCO_3$, water and brine solution. The combined ether layer was dried over anhydrous Na_2SO_4 and concentrated under reduced pressure to afford title compound as a pale yellow semi-solid (1.8 g, 71%).

Step 2. *tert*-Butyl-5-bromoindoline-1-carboxylate (BI12): To a stirred solution of 5-bromo-indoline (3.0 g , 15mmol) in MeCN (100 ml), was added DMAP (0.185 g , 1.522 mmol) and di-*tert*-butyl dicarbonate (3.98 g, 18.3 mmol) and the reaction was stirred at ambient temperature for 16 h. The reaction mixture was concentrated on reduced pressure to obtain a residue which was diluted with Et_2O and washed with water and brine solution (2X). The combined ether layer was dried over anhydrous Na_2SO_4 and concentrated under reduced pressure to afford the crude product as an off-white solid, which was used in the next step without further purification (3.0 g).

Step 3. *tert*-Butyl-5-vinylindoline-1-carboxylate (BI10): A stirred solution of *tert*-butyl-5-bromoindoline-1-carboxylate (2.0 g, 6.73 mmol), potassium vinyl trifluoroborate (2.6


g, 20.20 mmol) and K_2CO_3 (2.78 g, 20.2 mmol) in DMSO (50.0 mL) was degassed with argon for 20 min at ambient temperature. $PdCl_2$(dppf) (0.49 g, 0.67mmol) was added at ambient temperature, then the reaction mixture was heated to 100 °C for 3 h. The reaction mixture was cooled to ambient temperature and filtered through a Celite® bed under vacuum and washed with Et_2O. The reaction mixture was extracted with Et_2O. The combined Et_2O layer was dried over Na_2SO_4 and concentrated under reduced pressure to afford crude product. The crude compound was purified by column chromatography (SiO_2, 100-200 mesh; eluting with 2% EtOAc/ petroleum ether) to afford the title compound as an off-white solid (1.2 g, 73%): mp 85.5 -88.6 °C; ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.23 (m, 3H), 6.69 (dd, *J* = 17.4, 10.8 Hz, 1H), 5.64 (d, *J* = 10.5 Hz, 1H), 5.13 (d, *J* = 10.5 Hz, 1H), 4.00 (t, *J* = 9.0 Hz, 2H), 3.10 (t, *J* = 9.0 Hz, 2H), 1.55 (bs, 9H).

Example 29: Preparation of (*E*)-*tert*-Butyl 5-(3-(3,5-dichloro-4-fluorophenyl)-4,4,4-trifluorobut-1-en-1-yl)indoline-1-carboxylate (BI13)



To a stirred solution of *tert*-butyl-5-vinylindoline-1-carboxylate (1.28 g, 5.23mmol) in1,2-dichlorobenzene (10.0 mL), was added 5-(1-bromo-2,2,2-trifluoroethyl)-1,3-dichloro-2-fluorobenzene (3.4 g ,10 mmol), CuCl (103 mg, 1.05 mmol) and 2,2-bipyridyl (0.326 g, 2.092 mmol) and the resultant reaction mixture was degassed with argon for 30 min and heated to 150 °C for 1 h. The reaction mixture was cooled to ambient temperature and filtered and the filtrate was concentrated under reduced pressure. The crude compound was purified by column chromatography (SiO_2, 100-200 mesh; 2% EtOAc/ petroleum ether) to afford the title compound as a pale yellow gummy solid (0.3 g, 61%): ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.34 (d, *J* = 6.0 Hz, 2H), 7.22 (s, 2H), 7.16 (d, *J* = 8.4 Hz, 1H), 6.52 (d, *J* = 16.0 Hz, 1H), 6.21 (dd, *J* = 16.0, 7.6 Hz, 1H), 4.07 (m, 3H), 3.10 (t, *J* = 8.4 Hz, 2H), 1.55 (s, 9H); ESIMS *m/z* 433.79 ([M-H]$^-$); IR (thin film) 1168, 858 cm^{-1}.

Example 30: Preparation of (*E*)-5-(3-(3,5-Dichloro-4-fluorophenyl)-4,4,4-trifluorobut-1-en-1-yl)indolin-1-amine (BI14)




Step 1. (*E*)- 5-(3-(3,5-Dichloro-4-fluorophenyl)-4,4,4-trifluorobut-1-enyl)indoline (BI15) To a stirred solution of (*E*)-*tert*-butyl-5-(3-(3,5-dichloro-4-fluorophenyl)-4,4,4-trifluorobut-1-enyl)indoline-1-carboxylate (0.2 g, 0.4 mmol) in CH_2Cl_2 (10.0 mL) was added TFA (0.6 mL) and the reaction was stirred at ambient temperature for 2 h. The reaction mixture was diluted with CH_2Cl_2, washed with saturated aq $NaHCO_3$, water and brine solution. The separated CH_2Cl_2 layer was dried over anhydrous Na_2SO_4 and concentrated under reduced pressure to afford the crude product as a light brown gummy material which was used in the next step without further purification (0.12 g): ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.33 (d, *J* = 6.4 Hz, 2H), 7.21 (s, 1H), 7.02 (d, *J* = 8.0 Hz, 1H), 6.57 (d, *J* = 8.4 Hz, 1H), 6.49 (d, *J* = 15.6 Hz, 1H), 6.21(dd, *J* = 15.6, 8.4 Hz, 1H), 4.07 (m, 1H), 3.61 (t, *J* = 8.4 Hz, 2H), 3.05 (t, *J* = 8.4 Hz, 2H); ESIMS *m/z* 389.89 ($[M+H]^+$); IR (thin film) 3385, 1112, 816 cm^{-1}.

Step 2. 5-(3-(3,5-Dichloro-4-fluorophenyl)-4,4,4-trifluorobut-1-enyl)-1-nitrosoindoline (BI16): To (*E*)- 5-(3-(3,5-dichloro-4-fluorophenyl)-4,4,4-trifluorobut-1-enyl)indoline (0.2 g, 0.5 mmol) in concentrated HCl (5.0 ml) at 5 °C, was added slowly $NaNO_2$ in water and the reaction was allowed to stir at ambient temperature for 2 h. The reaction mixture was diluted with CH_2Cl_2, and the CH_2Cl_2 layer washed with water and brine solution. The separated CH_2Cl_2 layer was dried over anhydrous Na_2SO_4 and concentrated under reduced pressure to afford the crude product as a pale yellow solid that was used in the next step without further purification (0.2 g): ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.33 (d, *J* = 8.4 Hz, 1H), 7.39 (m, 4H), 6.61 (d, *J* = 16.0 Hz, 1H), 6.35 (dd, *J* =16.0, 8.4 Hz, 1H), 4.07 (m, 3H), 3.23 (t, *J* = 8.4 Hz, 2H); ESIMS *m/z* 418.82 ($[M+H]^+$); IR (thin film) 1488, 1112, 860 cm^{-1}.


Step 3. (*E*)-5-(3-(3,5-Dichloro-4-fluorophenyl)-4,4,4-trifluorobut-1-en-1-yl)indolin-1-amine (BI14): To (*E*)- 5-(3-(3,5-dichloro-4-fluorophenyl)-4,4,4-trifluorobut-1-enyl)-1-nitrosoindoline (0.1 g, 0.2 mmol) in MeOH(10.0 mL) was added zinc powder (77.5 mg) and NH_4Cl (36.9 mg, 0.69 mmol) in water (2.0 mL). The reaction mixture was stirred at ambient temperature for 3 h. The reaction mixture was diluted with CH_2Cl_2 and the CH_2Cl_2 layer was washed with water and brine solution. The separated CH_2Cl_2 layer was dried over anhydrous Na_2SO_4 and concentrated under reduced pressure to afford the crude compound, which was purified by column chromatography (SiO_2, 100-200 mesh; eluting with 2% EtOAc/ petroleum ether) to afford the title compound as a light brown gummy material (0.08 g): ESIMS *m/z* 404.86 ($[M+H]^+$).

Example 31: Preparation of (*E*)-N-(5-(3-(3,5-Dichloro-4-fluorophenyl)-4,4,4-trifluorobut-1-en-1-yl)indolin-1-yl)-3,3,3-trifluoropropanamide (BC12)



To a stirred solution of (*E*)-5-(3-(3,5-dichloro-4-fluorophenyl)-4,4,4-trifluorobut-1-enyl)indoline-1-amine (0.1 g, 0.247 mmol) in CH_2Cl_2 (10.0 ml) was added 3,3,3-trifluoropropanoic acid (0.038 g, 0.297 mmol), PyBOP (0.192 g, 0.370 mmol) and DIPEA (0.047 g, 0.370 mmol) and the reaction was stirred at ambient temperature for 18 h. The reaction mixture was diluted with CH_2Cl_2, and the separated CH_2Cl_2 layer dried over anhydrous Na_2SO_4 and concentrated under reduced pressure to afford the crude compound. The crude compound was purified by column chromatography (SiO_2, 100-200 mesh; 20-25% EtOAc/ petroleum ether) to afford the title compound as a light brown gummy material (0.12 g, 33%): ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.32, (d, *J* = 6.0 Hz, 2H) 7.28 (m, 1H), 7.20 (d, *J* = 8.0, 1H), 7.14 (d, *J* = 8.8, 1H), 6.70 (d, *J* = 8.0 Hz, 1H), 6.60 (m, 2H), 4.15 (m, 1H), 3.85 (m, 1H), 3.65 (m, 1H), 3.46 (m, 2H), 3.19 (m, 2H); ESIMS *m/z* 514.86 ($[M+H]^+$); IR (thin film) 3428, 1112, 857 cm^{-1}.


Example 32: Preparation of *tert*-Butyl-5-vinyl-1*H*-indole-1-carboxylate (BI17)



Step 1. 5-Vinyl-1*H*-indole (BI18): A mixture of 5-bromo-1*H*-indole (2.5 g, 12.82 mmol), potassium vinyltrifluoroborate (2.57 g ,19.2 mmol), Cs_2CO_3 (12.53 g, 38.46 mmol) and triphenylphosphine (201 mg, 0.769 mmol) in THF/water (9:1, 75 ml) was degassed with argon for 20 min, then charged with $PdCl_2$ (45.3 mg,0.256 mmol). The reaction mixture was heated to reflux for 16 h, then cooled to ambient temperature, filtered through Celite® bed and washed with EtOAc. The filtrate was again extracted with EtOAc, and the combined organic layer washed with water and brine, dried over Na_2SO_4 and concentrated under reduced pressure to afford the crude compound. The crude compound was purified by column chromatography (SiO_2, 100-200 mesh; 2% EtOAc/ petroleum ether) to afford the title compound as a light brown gummy material (1.5 g, 83%): ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.20 (br, 1H), 7.68 (s, 1H), 7.45 (s, 2H), 7.21 (m, 1H), 6.90 (dd, *J* =16.0, 10.8 Hz, 1H), 6.55 (m, 1H), 5.75 (d, *J* = 10.5 Hz, 1H), 5.21 (d, *J* = 10.5 Hz, 1H); ESIMS *m/z* 142.05 ($[M-H]^-$).


Step 2. *tert*-Butyl-5-vinyl-1*H*-indole-1-carboxylate (BI17): To a stirred solution of 5-vinyl-1*H*-indole (0.7 g, 4.89 mmol) in MeCN (20 ml) was added DMAP (59.65 mg, 0.489 mmol) and di-*tert*-butyl dicarbonate (1.38 g, 6.36 mmol), and the reaction was stirred at ambient temperature for 3 h. The reaction mixture was concentrated under reduced pressure to obtain a residue which was diluted with CH_2Cl_2 and washed with water and brine solution. The combined CH_2Cl_2 layer was dried over anhydrous Na_2SO_4 and concentrated under reduced pressure to afford the crude compound. The crude compound was purified by column chromatography (SiO_2, 100-200 mesh; 2% EtOAc/ petroleum ether) to afford the title compound as an off-white semi-solid (0.7 g, 59%): ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.15 (d, *J* = 8.0 Hz, 1H), 7.60 (s, 2H), 7.30 (d, *J* = 8.4 Hz, 1H), 7.21 (m, 1H), 6.90 (dd, *J* =16.0, 10.8 Hz, 1H), 6.59 (s, 1H), 5.75 (d, *J* = 10.5 Hz, 1H), 5.21 (d, *J* = 10.5 Hz, 1H), 1.65 (s, 9H); ESIMS *m/z* 242.10 ([M-H]$^-$); IR (thin film) 1630 cm^{-1}.

Example 33: Preparation of (*E*)-*tert*-Butyl 5-(3-(3,5-dichloro-4-fluorophenyl)-4,4,4-trifluorobut-1-en-1-yl)-1*H*-indole-1-carboxylate (BI19)




To a stirred solution of *tert*-butyl 5-vinyl-1*H*-indole-1-carboxylate (0.65 g, 2.67 mmol), in 1,2-dichlorobenzene (10.0 mL) was added 5-(1-bromo-2,2,2-trifluoroethyl)-1,3-dichloro-2-fluorobenzene (1.74 g, 5.37 mmol), CuCl (53 mg, 0.537 mmol) and 2,2-bipyridyl (167 mg, 1.07 mmol). The resultant reaction mixture was degassed with argon for 30 min and heated to 150 °C for 2 h. The reaction mixture was cooled to ambient temperature and filtered, and the filtrate concentrated under reduced pressure. The crude compound was purified by column chromatography (SiO_2, 100-200 mesh; 2% EtOAc/ petroleum ether) to afford the title compound as a light brown gummy material (0.25 g, 10%): ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.20 (d, *J* = 8.0 Hz, 1H), 7.60 (m, 2H), 7.39 (m, 3H), 6.69 (d, *J* = 16.0 Hz, 1H), 6.55 (d, *J* = 10.5 Hz, 1H), 6.36 (dd, *J* = 16.0, 8.0 Hz, 1H), 4.10 (m, 1H), 1.65 (s, 9H); ESIMS *m/z* 485.91 ([M-H]$^-$); IR (thin film) 1165, 854 cm^{-1}.

Example 34: Preparation of (*E*)-5-(3-(3,5-Dichloro-4-fluorophenyl)-4,4,4-trifluorobut-1-en-1-yl)-1*H*-indole (BI20)




To a stirred solution of (*E*)-*tert*-butyl 5-(3-(3,5-dichloro-4-fluorophenyl)-4,4,4-trifluorobut-1-enyl)-1*H*-indole-1-carboxylate (0.2 g, 0.40 mmol) in CH_2Cl_2 (10.0 mL) was added TFA (70 mg, 0.61 mmol) and the reaction was stirred at ambient temperature for 2 h. The reaction mixture was diluted with CH_2Cl_2 and washed with saturated $NaHCO_3$ solution, water and brine solution. The separated CH_2Cl_2 layer was dried over anhydrous Na_2SO_4 and concentrated under reduced pressure to afford the title compound as a light brown solid (0.2 g, 97%): mp 132.9-138.8 °C; ^{1}H NMR (400 MHz, $CDCl_3$) δ 11.19 (br, 1H), 8.20 (d, *J* = 8.0 Hz, 1H), 7.60 (m, 2H), 7.39 (m, 3H), 6.69 (d, *J* = 16.0 Hz, 1H), 6.55 (d, *J* = 10.5 Hz, 1H), 6.36 (dd, *J* =16.0, 8.0 Hz, 1H), 4.82 (m, 1H); ESIMS *m/z* 387.98 ([M+H]$^+$).

Example 35: Preparation of 4-Nitrophenyl 2-((*tert*-butoxycarbonyl)amino)acetate (BI21)



To a stirred solution of 4-nitrophenol (1.0 g, 7.19 mmol) in CH_2Cl_2 (20.0 mL) was added *N*-Boc glycine (1.38 g, 7.91 mmol) and EDC HCl (2.05 g,10.785 mmol) and the reaction was stirred at ambient temperature for 24 h. The reaction mixture was diluted with CH_2Cl_2 and washed with water and saturated brine solution. The separated CH_2Cl_2 layer was dried over anhydrous Na_2SO_4 and concentrated under reduced pressure to afford the title compound as a light brown gummy material that was used in the next step without further purification (1.1 g): ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.29 (d, *J* = 9.2 Hz, 2H), 7.33 (d, *J* = 8.8 Hz, 2H), 5.07 (br, 1H), 4.20 (s, 2H), 1.47 (s, 9H); ESIMS *m/z* 296.27 ([M+H]$^+$).


Example 36: Preparation of (*E*)-*tert*-Butyl (2-(5-(3-(3,5-dichloro-4-fluorophenyl)-4,4,4-trifluorobut-1-en-1-yl)-1*H*-indol-1-yl)-2-oxoethyl)carbamate (BI22)



To a stirred solution of (*E*)-5-(3-(3,5-dichloro-4-fluorophenyl)-4,4,4-trifluorobut-1-enyl)-1*H*-indole (0.1 g, 0.258 mmol) in MeCN (5.0 mL) was added 4-nitrophenyl 2-(*tert*-butoxycarbonylamino) acetate (0.114 g, 0.387 mmol), KF (0.03 g, 0.516 mmol), 18-crown-6-

ether (0.075 g, 0.283 mmol) and DIPEA (0.0332 g, 0.258 mmol) and the reaction was stirred at ambient temperature for 16 h. The reaction mixture was concentrated to obtain a residue which was diluted with CH_2Cl_2 and washed with water and brine solution. The separated CH_2Cl_2 layer was dried over anhydrous Na_2SO_4 and concentrated under reduced pressure to afford the crude title compound as a light brown gummy material which was used in the next step without further purification (0.1 g): ESIMS *m/z* 545.23 ($[M+H]^+$).


Example 37: Preparation of (*E*)-*N*-(2-(5-(3-(3,5-Dichloro-4-fluorophenyl)-4,4,4-trifluorobut-1-en-1-yl)-1*H*-indol-1-yl)-2-oxoethyl)-3,3,3-trifluoropropanamide (BC13)



Step 1. (*E*)-2-Amino-1-(5-(3-(3,5-dichloro-4-fluorophenyl)-4,4,4-trifluorobut-1-enyl)-1*H*-indol-1-yl)ethanone (BI23): To a stirred solution of (*E*)-*tert*-butyl 2-(5-(3-(3,5-dichloro-4-fluorophenyl)-4,4,4-trifluorobut-1-enyl)-1*H*-indol-1-yl)-2-oxoethylcarbamate (0.05 g, 0.09 mmol) in CH_2Cl_2 (5.0 mL) was added TFA (0.01 mL) and the reaction was stirred at ambient temperature for 16 h. The reaction mixture was diluted with CH_2Cl_2 and washed with saturated $NaHCO_3$ solution, water and brine solution. The separated CH_2Cl_2 layer was dried over anhydrous Na_2SO_4 and concentrated under reduced pressure to afford the crude title compound which was used in the next step without further purification (50 mg).

Step 2. (*E*)-*N*-(2-(5-(3-(3,5-Dichloro-4-fluorophenyl)-4,4,4-trifluorobut-1-en-1-yl)-1*H*-indol-1-yl)-2-oxoethyl)-3,3,3-trifluoropropanamide (BC13): To a stirred solution of (*E*)-2-amino-1-(5-(3-(3,5-dichloro-4-fluorophenyl)-4,4,4-trifluorobut-1-enyl)-1*H*-indol-1-yl) ethanone (0.04 g, 0.09 mmol) in CH_2Cl_2 (5.0 ml) was added 3,3,3-trifluoropropanoic acid (17.5 mg, 0.136 mmol), PyBOP (70 mg, 0.135 mmol) and DIPEA (29 mg, 0.225 mmol) and the reaction was stirred at ambient temperature for 16 h. The reaction mixture was diluted with CH_2Cl_2, and the CH_2Cl_2 layer was washed with water and saturated brine solution .The separated CH_2Cl_2 layer was dried over anhydrous Na_2SO_4 and concentrated under reduced pressure to afford the crude compound, which was purified by column chromatography (SiO_2, 100-200 mesh; 10% EtOAc/ petroleum ether) to afford the title compound as an off-white solid (30 mg, 60%): mp 121-126 °C; ^1H NMR (400 MHz, $CDCl_3$) δ 8.33 (br, 1H), 7.59 (s, 1H), 7.45 (m, 4H), 6.72 (d, *J* = 3.6 Hz, 3H) , 6.39 (m, 1H), 4.71 (t, *J* = 7.2 Hz, 2H), 4.15 (m, 1H), 3.51 (m, 1H), 3.28 (m, 1H); ESIMS *m/z* 553.06 ($[M-H]^-$).

Example 38: Preparation of Ethyl 2-(1-oxo-6-vinylphthalazin-2(1*H*)-yl)acetate (BI24)




Step 1. 5-Bromo-3-hydroxyisoindoline-1-one (BI25): A mixture of Zn powder (1.73 g, 26.154 mmol), copper (II) sulfate pentahydrate (0.02 g ,0.08 mmol) and 2M aq NaOH (27 mL) were cooled to 0 °C. 5-Bromoisoindoline-1,3-dione (5 g, 22mmol) was added at the same temperature over the period of 30 min. The reaction mixture was stirred at 0 °C for 30 min and 3 h at ambient temperature. The reaction mixture was filtered and the filtrate was neutralized with concentrated HCl. The reaction mixture was diluted with ethanol and extracted with EtOAc. The combined EtOAc layer was dried over Na_2SO_4 and concentrated under reduced pressure to afford the crude title compound as a brown solid, which was used in the next step without further purification (1.3 g): mp 258-261°C; ^{1}H NMR (400 MHz, DMSO-d_6) δ 9.03 (br, 1H), 7.81 (m, 2H), 7.69 (m, 1H), 6.44 (m, 1H), 5.88 (d, J = 9.3 Hz, 1H); ESIMS m/z 225.83 ([M-H]$^-$); IR (thin film) 1684, 3246, 606 cm^{-1}.

Step 2. 6-Bromophthalazine-1(2*H*)-one (BI26): To a stirred solution of 5-bromo-3-hydroxyisoindoline-1-one (1.0 g, 4.40 mmol) in water, was added hydrazine hydrate (0.45 g , 8.80 mmol) and heated to 95°C for 5 h. The reaction mixture was cooled to ambient temperature, filtered and washed with Et_2O and pentane (1:1) to afford the title compound as a white solid that was used in the next step without further purification (0.5 g): ESIMS m/z 225.15 ([M+H]$^+$).

Step 3. 6-Vinylphthalazine-1(2*H*)-one (BI27): A solution of 6-bromophthalazine-1(2*H*)-one (0.25 g, 1.11 mmol), potassium vinyl trifluoroborate (0.446 g, 3.33 mmol) and K_2CO_3 (0.46 g, 3.33 mmol) in DMSO (2 mL) was degassed with argon for 20 min at ambient temperature. $PdCl_2$(dppf) (0.04 g, 0.055 mmol) was added at ambient temperature, and the reaction mixture was heated to 80 °C for 2 h. The reaction mixture was cooled to ambient temperature and filtered through Celite® bed under vacuum and washed with EtOAc. The reaction mixture was extracted with EtOAc and the combined EtOAc layer dried over Na_2SO_4 and concentrated under reduced pressure to afford the crude product. The crude compound was purified by column chromatography (SiO_2, 100-200 mesh; 50% EtOAc/ petroleum ether) to afford the title compound as a brown solid (0.12 g, 63%): ^{1}H NMR (400 MHz, DMSO-d_6) δ 13.61 (br, 1H), 8.33 (m, 1H), 8.19 (m, 1H), 8.01 (m, 2H), 6.97 (m, 1H), 6.15 (m, 1H), 5.56 (d, J = 10.8 Hz, 1H); ESIMS m/z 172.93 ([M+H]$^+$); IR (thin film) 1748, 1655, 3241 cm^{-1}.


Step 4. Ethyl-2-(1-oxo-6-vinylphthalazine-2(1H)-yl acetate (BI24): To a stirred solution of 6-vinylphthalazine-1(2*H*)-one (0.5 g, 2.90 mmol) in DMF (5.0 mL) was added Cs_2CO_3 (0.94 g, 2.90 mmol) and the reaction was stirred for 10 min. Ethyl bromoacetate (0.48 g,2.90 mmol) was added to the reaction mixture at ambient temperature and the reaction was stirred for 8 h at ambient temperature. The reaction mixture was diluted and extracted with EtOAc, and the EtOAc layer was washed with water and brine solution (2X). The separated EtOAc layer was dried over anhydrous Na_2SO_4 and concentrated under reduced pressure to afford crude product. The crude compound was purified by column chromatography (SiO_2, 100-200 mesh; 25% EtOAc/ petroleum ether) to afford the title compound as a brown solid (0.34 g, 45%): ^{1}H NMR (400 MHz, DMSO-d_6) δ 8.45 (m, 1H), 8.24 (m, 1H), 8.04 (m, 2H), 7.01 (m, 1H), 6.17 (d, J = 2.1 Hz, 1H), 5.56 (d, J = 10.8 Hz, 1H), 4.92 (s, 2H), 4.19 (m, 2H), 1.23 (m, 3H). ESIMS *m/z* 259.10 ([M+H]$^+$); IR (thin film) 1750, 1660 cm^{-1}.

Example 39: Preparation of (*E*)-Ethyl 2-(6-(3-(3,5-dichloro-4-fluorophenyl)-4,4,4-trifluorobut-1-en-1-yl)-1-oxophthalazin-2(1H)-yl)acetate (BI28)




To a stirred solution of ethyl-2-(1-oxo-6-vinylphthalazine-2(1*H*)-yl acetate (0.07 g, 0.27 mmol) in 1,2-dichlorobenzene (1.0 mL) was added 5-(1-bromo-2,2,2-trifluoroethyl)-1,3-dichloro-2fluorobenzene (0.17 g, 0.54 mmol), CuCl (0.005 g, 0.05 mmol) and 2,2-bipyridyl (0.016 g, 0.10 mmol) and the resultant reaction mixture was degassed with argon for 30 min and heated to 180 °C for 12 h. The reaction mixture was cooled to ambient temperature and filtered and the filtrated was concentrated under reduced pressure. The crude compound was purified by column chromatography (SiO_2, 100-200 mesh; 10-15% EtOAc/ petroleum ether) to afford the title compound as a brown solid (40 mg, 29%): ^{1}H NMR (400 MHz, DMSO-d_6) δ 8.40 (d, J = 8.4 Hz, 1H), 7.84 (d, J = 1.5 Hz, 1H), 7.65 (s, 1H), 7.37 (d, J = 6.3 Hz, 2H), 6.76 (d, J = 16.0 Hz, 1H), 6.59 (dd, J =16.0, 8.0 Hz, 1H), 4.96 (s, 2H), 4.29 (m, 3H), 1.31 (t, J = 7.2 Hz, 3H); ESIMS *m/z* 503.0 ([M+H]$^+$); IR (thin film) 1660, 1114, 817 cm^{-1}.

Example 40: Preparation of (*E*)-2-(6-(3-(3,5-Dichloro-4-fluorophenyl)-4,4,4-trifluorobut-1-en-1-yl)-1-oxophthalazin-2(1*H*)-yl)acetic acid (BI29)




A solution of (*E*)-ethyl-2-(6-(3-(3,5-dichloro-4-fluorophenyl)-4,4,4-trifluorobut-1-enyl)-1-oxophthalazin-2(1*H*)-yl) acetate (0.04 g, 0.07mmol) in HCl (0.5 mL) and AcOH (0.5 mL) was heated to 100 °C for 3 h. The solvent was removed under reduced pressure and the residue diluted with water. The aqueous layer was extracted with EtOAc and the separated EtOAc layer dried over anhydrous Na_2SO_4 and concentrated under reduced pressure to afford the crude compound. The crude compound was triturated with Et_2O-pentane mixture to afford the title compound as a brown solid (0.03 g): ^{1}H NMR (400 MHz, DMSO-d_6) δ 13.0 (br s, 1H), 8.43 (m, 1H), 8.23 (d, *J* = 8.1 Hz, 1H), 8.14 (m, 2H), 7.91 (m, 2H), 7.16 (dd, *J* =16.0, 8.0 Hz, 1H), 6.99 (d, *J* = 16.0 Hz, 1H), 4.96 (m, 3H),; ESIMS *m/z* 473.0 ([M-H]$^-$); IR (thin film) 1629, 1168, 817 cm^{-1}.

Example 41: Preparation of (*E*)-2-(6-(3-(3,5-Dichloro-4-fluorophenyl)-4,4,4-trifluorobut-1-en-1-yl)-1-oxophthalazin-2(1*H*)-yl)-N-(2,2,2-trifluoroethyl)acetamide (BC14)




To a stirred solution of (*E*)-2-(6-(3-(3,5-dichloro-4-fluorophenyl)-4,4,4-trifluorobut-1-enyl)-1-oxophthalazin-2(1*H*)-yl)acetic acid (0.15 g, 0.31 mmol) in CH_2Cl_2 (20.0 ml) was added 2,2,2,-trifluoroethanamine (0.03 g, 0.31mmol), PyBOP (0.17 g, 0.34 mmol) and DIPEA (0.15 ml, 0.93 mmol) at ambient temperature, and the reaction was stirred for 18 h. The reaction mixture was diluted with CH_2Cl_2 and washed with 3N HCl (2 x 20 mL), $NaHCO_3$ (2 x 20 mL) and brine solution (2x).The separated CH_2Cl_2 layer was dried over anhydrous Na_2SO_4 and concentrated under reduced pressure to afford the crude compound. The crude compound was purified by column chromatography (SiO_2, 100-200 mesh; 20-25% EtOAc/ petroleum ether) to afford the title compound as a brown solid (0.11 g): mp 172-175 °C; ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.83 (t, *J* = 6.6 Hz, 1H), 8.42 (t, *J* = 14.7 Hz, 1H), 8.22 (d, *J* = 8.1 Hz, 1H), 8.13 (t, *J* = 6.3 Hz, 1H), 7.98-7.86 (m, 2H), 7.16 - 7.07 (m, 1H), 7.01 - 6.93 (m, 1H), 4.96 - 4.81 (m, 3H), 4.00 - 3.88 (m, 2H); ESIMS *m/z* 554.0 ([M-H]$^-$).

Example 42: Preparation of 2-(4-Vinylbenzyl)isoindoline-1,3-dione (CI1)



To a stirred solution of 1-(chloromethyl)-4-vinylbenzene (10 g, 66 mmol) in DMF (100 mL) was added potassium phthalimide (13.3 g, 72.1 mmol), and the resultant reaction mixture was heated at 70 °C for 16 h. The reaction mixture was diluted with water and extracted with $CHCl_3$. The combined $CHCl_3$ layer was washed with brine, dried over Na_2SO_4 and concentrated under reduced pressure. Recrystallization from MeOH afforded the title compound as an off-white solid (8 g, 46%): ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.83 (m, 2H), 7.71 (m, 2H), 7.39 (m, 4H), 6.65 (dd, *J* = 17.6, 10.8 Hz, 1H), 5.72 (d, *J* = 17.6 Hz, 1H), 5.21 (d, *J* = 10.8 Hz , 1H), 4.82 (s, 2H); GCMS *m/z* 263.2 ($[M]^+$); IR (thin film) 3420, 1133, 718 cm^{-1}.


Example 43: Preparation of (*E*)-2-(4-(3-(3,5-Dichlorophenyl)-4,4,4-trifluorobut-1-en-1-yl)benzyl)isoindoline-1,3-dione (CI2)



Using the procedure of Example 10 with 2-(4-vinylbenzyl)isoindoline-1,3-dione and 1-(1-bromoethyl)-3,5-dichlorobenzene as the starting materials, the title compound was isolated as an off-white solid (0.3 g, 40-50%): mp 142–145 °C; ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.86 (m, 2H), 7.74 (m, 2H), 7.42 (m, 2H), 7.36 (m,3H), 7.27 (m, 2H), 6.58 (d, *J* = 16.0 Hz, 1H), 6.32 (dd, *J* = 16.0, 8.0 Hz, 1H), 4.82 (s, 2H), 4.05 (m, 1H); ESIMS *m/z* 488.17 ($[M-H]^-$).

The following compound was made in accordance with the procedures disclosed in **Example 43**.

(*E*)-2-(4-(4,4,4-Trifluoro-3-(3,4,5-trichlorophenyl)but-1-en-1-yl)benzyl)isoindoline-1,3-dione (CI3)




The title compound was isolated as an off white solid (0.3 g, 56%): mp 145–146 °C; ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.86 (m, 2H), 7.74 (m, 2H), 7.42-7.31 (m, 6H), , 6.58 (d, *J* =

16.0 Hz, 1H), 6.53 (dd, J = 16.0, 8.0 Hz, 1H), 4.82 (s, 2H), 4.05 (m, 1H); ESIMS m/z 522.2 ([M-H]$^-$); IR (thin film) 1716, 1110, 712 cm^{-1}.

Prophetically, compounds **CI4**–**CI5** (Table 1) could be made in accordance with the procedures disclosed in **Example 43**.


Example 44: Preparation of (*E*)-(4-(3,5-Dichlorophenyl)-4,4,4-trifluorobut-1-en-1-yl)phenyl)methanamine (CI6)



To a stirred solution of (*E*)-2-(4-(3-(3,5-dichlorophenyl)but-1-en-1-yl)benzyl)-isoindoline-1,3-dione (1.2 g, 2.45 mmol) in EtOH was added hydrazine hydrate (0.61 g, 12 mmol), and the resultant reaction mixture was heated at 90 °C for 1 h. The reaction mixture was filtered, and the filtrate was concentrated. The residue was dissolved in CH_2Cl_2, washed with brine, dried over Na_2SO_4, and concentrated under reduced pressure to afford the crude title compound as a gummy liquid (0.9 g) which was used without further purification.

The following compounds were made in accordance with the procedures disclosed in **Example 44**.


(*E*)-(4-(4,4,4-Trifluoro-3-(3,4,5-trichlorophenyl)but-1-en-1-yl)phenyl)methanamine (CI7)



The title compound was isolated and used without further purification.

Prophetically, compounds **CI8**–**CI9** (Table 1) could be made in accordance with the procedures disclosed in **Example 44**.

Example 45: Preparation of 4-(Bromomethyl)-3-chlorobenzonitrile (CI10)




To a stirred solution of 3-chloro-4-methylbenzonitrile (5 g, 25.4 mmol) in CCl_4 (50 mL) under an argon atmosphere was added NBS (5.16 g, 29 mmol), and the mixture was degassed for 30 min. To this was added AIBN (0.3 g, 1.8 mmol), and the resultant reaction

mixture was heated at reflux for 4 h. The reaction mixture was cooled to ambient temperature, washed with water, and extracted with CH_2Cl_2. The combined CH_2Cl_2 layer was washed with brine, dried over Na_2SO_4, and concentrated under reduced pressure. The crude compound was purified by flash column chromatography (SiO_2, 100-200 mesh; 5% EtOAc in *n*-Hexane) to afford the title compound as a white solid (4.8 g, 68%): mp 87–88 °C; ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.71 (s, 1H), 7.59 (s, 2H), 4.60 (s, 2H); ESIMS *m/z* 229.77 ($[M+H]^+$); IR (thin film) 2235, 752, 621 cm^{-1}.


The following compounds were made in accordance with the procedures disclosed in **Example 45**.

4-(Bromomethyl)-3-(trifluoromethyl)benzonitrile (CI11)




The title compound was isolated as an off-white gummy material (5 g, 66%): ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.96 (s, 1H), 7.86 (d, *J* = 8.0 Hz, 1H), 7.76 (d, *J* = 8.0 Hz, 1H), 4.62 (s, 2H); ESIMS *m/z* 262.11 ($[M-H]^-$); IR (thin film) 2236, 1132, 617 cm^{-1}.

3-Bromo-4-(bromomethyl)benzonitrile (CI12)




The title compound was isolated as an off-white solid(5 g, 67%): mp 82–83 °C; ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.90 (s, 1H), 7.61 (m, 2H), 4.62 (s, 2H); EIMS *m/z* 272.90; IR (thin film) 2229, 618 cm^{-1}.

4-(Bromomethyl)-3-fluorobenzonitrile (CI13)



The title compound was isolated as an off-white solid (2 g, 60%): mp 79–81 °C; ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.54 (t, *J* = 8.0 Hz, 1H), 7.48 (dd, *J* = 8.0 Hz, 8.0, 1H), 7.38 (dd, *J* = 5 Hz, 1H), 4.5 (s, 2H); EIMS *m/z* 215.


Example 46: Preparation of 4-(Bromomethyl)-3-chlorobenzaldehyde (CI14)



To a stirred solution of 4-(bromomethyl)-3-chlorobenzonitrile (4.8 g, 17 mmol) in toluene (50 mL) at 0 °C was added dropwise DIBAl-H (1.0 M solution in toluene; 23.9 mL), and the reaction mixture was stirred at 0 °C for 1 h. 10 M HCl in water (5 mL) was added until the reaction mixture turned to a white slurry and then additional 1 N HCl (20 mL) was added. The organic layer was collected and the aqueous layer was extracted with $CHCl_3$. The combined organic layer was dried over Na_2SO_4 and concentrated under reduced pressure. The crude compound was purified by flash column chromatography (SiO_2, 100-200 mesh; 5% EtOAc in *n*-Hexane) to afford the title compound as a white solid (3.8 g, 80%): mp 64–66 °C; ^{1}H NMR (400 MHz, $CDCl_3$) δ 10.00 (s, 1H), 7.92 (s, 1H), 7.78 (d, *J* = 8.0 Hz, 1H), 7.64 (d, *J* = 8.0 Hz, 1H), 4.60 (s, 2H); ESIMS *m/z* 232.78 ($[M+H]^+$).


The following compounds were made in accordance with the procedures disclosed in **Example 46**.

4-(Bromomethyl)-3-(trifluoromethyl)benzaldehyde (CI15)




The title compound was isolated as a pale yellow low-melting solid (5 g, 60%): ^{1}H NMR (400 MHz, $CDCl_3$) δ 10.09 (s, 1H), 8.19 (s, 1H), 8.09 (m, 1H), 7.81 (m, 1H), 4.61 (s, 2H); ESIMS *m/z* 265.04 ($[M-H]^-$); IR (thin film) 1709, 1126, 649 cm^{-1}.

3-Bromo-4-(bromomethyl)benzaldehyde (CI16)




The title compound was isolated as a pale yellow solid (5 g, 62%): mp 94–95 °C; ^{1}H NMR (400 MHz, $CDCl_3$) δ 9.96 (s, 1H), 8.05 (s, 1H), 7.81 (d, *J* = 8.0 Hz, 1H), 7.62 (d, *J* = 8.0 Hz, 1H), 4.60 (s, 2H); EIMS *m/z* 275.90 ($[M]^+$).

4-(Bromomethyl)-3-fluorobenzaldehyde (CI17)



The title compound was isolated as an off-white solid (5 g, 61%): mp 43–45 °C; ^{1}H NMR (400 MHz, $CDCl_3$) δ 9.1 (s, 1H), 7.54 (t, *J* = 8 Hz, 1H), 7.48 (d, *J* = 8 Hz, 1H), 7.38 (d, *J* = 5 Hz, 1H), 4.5 (s, 2H); EIMS *m/z* 216 ($[M]^+$).


Example 47: Preparation of 3-Chloro-4-((1,3-dioxoisoindolin-2-yl)methyl)benzaldehyde (CI18)



To a stirred solution of 4-(bromomethyl)-3-chlorobenzaldehyde (3.8 g, 14 mmol) in DMF (40 mL) was added potassium pthalimide (3.54 g, 19.14 mmol), and the mixture was heated at 60°C for 6 h. The reaction mixture was cooled to ambient temperature and diluted with water (100 mL). The solid obtained was separated by filtration and dried under vacuum to afford the title compound as a white solid (2.8 g, 60%): mp 123–126 °C; ^{1}H NMR (400 MHz, $CDCl_3$) δ 9.95 (s, 1H), 8.21 (s, 1H), 7.91 (m, 3H), 7.80 (m, 2H), 7.20 (m, 1H), 5.05 (s, 2H); ESIMS *m/z* 298.03 ($[M-H]^-$).


The following compounds were made in accordance with the procedures disclosed in **Example 47**.

4-((1,3-Dioxoisoindolin-2-yl)-3-(trifluoromethyl)benzaldehyde (CI19)




The title compound was isolated as an off white solid (1 g, 62%): mp 142–143 °C; ^{1}H NMR (400 MHz, $CDCl_3$) δ 10.05 (s, 1H), 8.15 (s, 1H), 7.91 (m, 2H), 7.80 (m, 3H), 7.27 (m, 1H), 5.19 (s, 2H); ESIMS *m/z* 332.03 ($[M-H]^-$).

3-Bromo-4-((1,3-dioxoisoindolin-2-yl)methyl)benzaldehyde (CI20)




The title compound was isolated as an off-white solid (0.5 g, 64%): mp 159–161 °C; ^{1}H NMR (400 MHz, $CDCl_3$) δ 9.95 (s, 1H), 8.21 (s, 1H), 7.91 (m, 3H), 7.80 (m, 2H), 7.20 (m, 1H), 5.05 (s, 2H); ESIMS *m/z* 314.00 ([M-CHO]$^-$).

4-((1,3-Dioxoisoindolin-2-yl)-3-fluorobenzaldehyde (CI21)



The title compound was isolated as a white solid (2 g, 60%): mp 154–156 °C; ^{1}H NMR (400 MHz, $CDCl_3$) δ 9.95 (s, 1H), 7.9 (m, 2H), 7.75 (m, 2H), 7.6 (m, 2H), 7.5 (t, *J* = 7.6 Hz, 1H), 5.05 (s, 2H); EIMS *m/z* 283.1 ([M]$^+$).

Example 48: Preparation of 2-(2-Chloro-4-vinylbenzyl)isoindoline-1,3-dione (CI22)




To a stirred solution of 3-chloro-4-((1,3-dioxoisoindolin-2-yl)methyl)benzaldehyde (2.8 g, 8.2 mmol) in 1,4-dioxane (30 mL) were added K_2CO_3 (1.68 g, 12.24 mmol) and methyl triphenyl phosphonium bromide (4.37 g, 12.24 mmol) at ambient temperature. Then the resultant reaction mixture was heated at 100 °C for 18 h. After the reaction was deemed complete by TLC, the reaction mixture was cooled to ambient temperature and filtered, and the obtained filtrate was concentrated under reduced pressure. The residue was purified by flash chromatography (SiO_2, 100-200 mesh; 20% EtOAc in *n*-Hexane) to afford the title compound as a white solid (1.94 g, 70%): mp 141–143 °C; ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.85 (m, 2H), 7.70 (m, 2H), 7.41 (m, 1H), 7.21 (m, 2H), 6.71 (dd, *J* = 17.6, 10.8 Hz, 1H),

5.72 (d, J = 17.6 Hz, 1H), 5.23 (d, J = 10.8 Hz, 1H), 4.92 (s, 2H); ESIMS m/z 298.10 ($[M-H]^-$).


The following compounds were made in accordance with the procedures disclosed in **Example 48**.

2-(2-(Trifluoromethyl)-4-vinylbenzyl)isoindoline-1,3-dione (CI23)




The title compound was isolated as a light brown solid (0.5 g, 60%): mp 134–135 °C; ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.92 (m, 2H), 7.80 (m, 2H), 7.71 (s, 1H), 7.46 (d, J = 8.0 Hz, 1H), 7.16 (d, J = 8.0 Hz, 1H), 6.65 (m, 1H), 5.80 (d, J = 17.8 Hz, 1H), 5.19 (d, J = 10.8 Hz, 1H), 5.09 (s, 2H); ESIMS m/z 332.10 ($[M+H]^+$).

2-(2-Bromo-4-vinylbenzyl)isoindoline-1,3-dione (CI24)




The title compound was isolated as an off white solid (0.5 g, 62%): mp 126–128 °C; ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.92 (m, 2H), 7.79 (m, 2H), 7.62 (s, 1H), 7.21 (m, 1H), 7.16 (d, J = 8.0 Hz, 1H), 6.62 (m, 1H), 5.72 (d, J = 17.8 Hz, 1H), 5.15 (d, J = 10.8 Hz, 1H), 4.95 (s, 2H); EIMS m/z 341.10 ($[M]^+$).

2-(2-Fluoro-4-vinylbenzyl)isoindoline-1,3-dione (CI25)



The title compound was isolated as a white solid (0.5 g, 61%): mp 140–142 °C; ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.85 (m, 2H), 7.72 (m, 2H), 7.25 (m, 1H), 7.11 (m, 2H), 6.63 (m, 1H), 5.80 (d, J = 17.6 Hz, 1H), 5.28 (d, J = 10.8 Hz, 1H), 4.92 (s, 2H); EIMS m/z 282.08.


Example 49: Preparation of (*E*)-2-(2-Chloro-4-(3-(3,5-dichlorophenyl)-4,4,4-trifluorobut-1-en-1-yl)benzyl)isoindoline-1,3-dione (CI26)



To a stirred solution of 2-(2-chloro-4-vinylbenzyl)isoindoline-1,3-dione (2.0 g, 6.51 mmol) in 1,2-dichlorobenzene (25 mL) were added 1-(1-bromo-2,2,2-trifluoroethyl)-3,5-dichlorobenzene (3.48 g, 11.36 mmol), CuCl (112 mg, 1.13 mmol) and 2,2- bipyridyl (0.35 g). The resultant reaction mixture was degassed with argon for 30 min and then was stirred at 180 °C for 24 h. After the reaction was deemed complete by TLC, the reaction mixture was cooled to ambient temperature and filtered, and the filtrate was concentrated under reduced pressure. The residue was purified by flash chromatography (SiO_2, 100-200 mesh; 25–30% EtOAc in *n*-hexane) to afford the title compound as solid (1.3 g, 50%): mp 141–143 °C; ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.92 (m, 2H), 7.79 (m, 2H), 7.42 (m, 2H), 7.24 (m, 2H), 7.20 (m, 2H), 6.54 (d, *J* = 16.0 Hz, 1H), 6.34 (dd, *J* = 16.0, 8.0 Hz, 1H), 5.00 (s, 2H), 4.10 (m, 1H); ESIMS *m/z* 524.07 ($[M+H]^+$).


The following compounds were made in accordance with the procedures disclosed in **Example 49**.

(*E*)-2-(2-Chloro-4-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-en-1-yl)benzyl)isoindoline-1,3-dione (CI27)




The title compound was isolated as a pale white solid (0.2 g, 55%): mp 128–129 °C; ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.92 (m, 2H), 7.79 (m, 2H), 7.42 (m, 3H), 7.22 (m, 2H), 6.52 (d, *J* = 16.0 Hz, 1H), 6.32 (dd, *J* = 16.0, 8.0 Hz, 1H), 5.00 (s, 2H), 4.05 (m, 1H); ESIMS *m/z* 557.99 ($[M+H]^+$).

(*E*)-2-(2-Chloro-4-(3-(3,5-dichloro-4-fluorophenyl)-4,4,4-trifluorobut-1-en-1-yl)benzyl)isoindoline-1,3-dione (CI28)




The title compound was isolated as an off white solid (0.2 g, 54%): mp 177–180 °C; ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.90 (m, 2H), 7.77 (m, 2H), 7.42 (s, 1H), 7.32 (d, *J* = 8.0 Hz, 2H), 7.21 (m, 2H), 6.52 (d, *J* = 16.0 Hz, 1H), 6.32 (dd, *J* = 16.0, 8.0 Hz, 1H), 5.00 (s, 2H), 4.05 (m, 1H); ESIMS *m/z* 540.08 ([M-H]$^-$); IR (thin film) 1716 cm^{-1}.

(*E*)-2-(2-Chloro-4-(3-(3,4-dichlorophenyl)-4,4,4-trifluorobut-1-en-1-yl)benzyl)isoindoline-1,3-dione (CI29)



The title compound was isolated as an off-white solid (0.2 g, 59%): ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.89 (m, 2H), 7.76 (m, 2H), 7.47 (m, 3H), 7.21 (m, 3H), 6.50 (d, *J* = 16.0 Hz, 1H), 6.32 (dd, *J* = 16.0, 7.6 Hz, 1H), 4.97 (s, 2H), 4.11 (m, 1H); ESIMS *m/z* 522.27 ([M-H]$^-$); IR (thin film) 3064, 1717, 1111, 715 cm^{-1}.


(*E*)-2-(4-(3-(3,5-Dichlorophenyl)-4,4,4-trifluorobut-1-en-1-yl)-2-(trifluoromethyl)-benzyl)isoindoline-1,3-dione (CI30)



The title compound was isolated as an off-white solid (0.2 g, 54%): mp 141–142 °C; ^{1}H NMR (400 MHz, $CDCl_3$) 7.94 (m, 2H), 7.80 (m, 2H), 7.69 (s, 1H), 7.44 (m, 1H), 7.38 (m,


1H), 7.24 (m, 2H), 7.19 (m, 1H), 6.60 (d, *J* = 16.0 Hz, 1H), 6.39 (dd, *J* = 16.0, 7.6 Hz, 1H), 5.10 (s, 2H), 4.11 (m, 1H); ESIMS *m/z* 556.00 ([M-H]$^-$).

(*E*)-2-(4-(4,4,4-Trifluoro-3-(3,4,5-trichlorophenyl)but-1-en-1-yl)-2-(trifluoromethyl)-benzyl)isoindoline-1,3-dione (CI31)




The title compound was isolated as an off-white solid (0.2 g, 56%): mp 130–132 °C; ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.94 (m, 2H), 7.80 (m, 2H), 7.69 (s, 1H), 7.44 (m, 3H), 7.19 (m, 1H), 6.61 (d, *J* = 16.0 Hz, 1H), 6.38 (dd, *J* = 16.0, 7.6 Hz, 1H), 5.10 (s, 2H), 4.12 (m, 1H); ESIMS *m/z* 589.57 ([M-2H]$^-$).

(*E*)-2-(2-Bromo-4-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-en-1-yl)benzyl)-isoindoline-1,3-dione (CI32)



The title compound was isolated as a pale yellow solid (0.2 g, 55%): mp 160–162 °C; ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.92 (m, 2H), 7.80 (m, 2H), 7.62 (s, 1H), 7.39 (s, 2H), 7.24 (m, 1H), 7.16 (m, 1H), 6.52 (d, *J* = 16.0 Hz, 1H), 6.32 (dd, *J* = 16.0, 8.0 Hz, 1H), 4.98 (s, 2H), 4.12 (m, 1H); ESIMS *m/z* 599.78 ([M-H]$^-$).


(*E*)-2-(2-Fluoro-4-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-en-1-yl)benzyl)-isoindoline-1,3-dione (CI33)



The title compound was isolated as an off-white solid (0.2 g, 55%): mp 72–74 °C; ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.88 (m, 2H), 7.74 (m, 2H), 7.38 (s, 2H), 7.34 (m, 1H), 7.18 (m, 2H), 6.54 (d, *J* = 16.0 Hz, 1H), 6.32 (dd, *J* = 16.0, 8.0 Hz, 1H), 4.91 (s, 2H), 4.08 (m, 1H); ESIMS *m/z* 539.89 ([M-H]$^-$); IR (thin film)1773 cm^{-1}.

Prophetically, compounds **CI34**–**CI41** (Table 1) could be made in accordance with the procedures disclosed in **Example 49**.


Example 50: Preparation of (*E*)-(2-Chloro-4-(3-(3,5-dichlorophenyl)-4,4,4-trifluorobut-1-en-1-yl)phenyl)methanamine (CI42)



To a stirred solution of (*E*)-2-(2-chloro-4-(3-(3,5-dichlorophenyl)-4,4,4-trifluorobut-1-en-1-yl)benzyl)isoindoline-1,3-dione (0.4 g, 0.76 mmol) in EtOH was added hydrazine hydrate (0.38 g, 7.6 mmol), and the resultant reaction mixture was heated at 80 °C for 2 h. The reaction mixture was filtered, and the filtrate was concentrated. The residue was dissolved in CH_2Cl_2, washed with brine, dried over Na_2SO_4, and concentrated under reduced pressure to afford the title compound as a gummy liquid (0.3 g), which was carried on to the next step without further purification.


The following compounds were made in accordance with the procedures disclosed in **Example 50**.

(*E*)-(2-Chloro-4-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-en-1-yl)phenyl)-methanamine (CI43)




The product obtained in this reaction was carried on to the next step without further purification.

(*E*)-(2-Chloro-4-(3-(3,4-dichlorophenyl)-4,4,4-trifluorobut-1-en-1-yl)phenyl)-methanamine (CI44)




The product obtained in this reaction was carried on to the next step without further purification: ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.48 (d, J = 8.4 Hz, 2H), 7.39 (m, 2H), 7.23 (m, 2H), 6.52 (d, J = 16.0 Hz, 1H), 6.38 (dd, J = 16.0, 7.6 Hz, 1H), 4.12 (m, 1H), 3.90 (s, 2H); ESIMS m/z 391.90 ([M-H]$^-$); IR (thin film) 3370, 3280, 1111, 817 cm^{-1}.

(*E*)-(4-(4,4,4-Trifluoro-3-(3,4,5-trichlorophenyl)but-1-en-1-yl)-2-(trifluoromethyl)-phenyl)methanamine (CI45)




The title compound was isolated as a gummy material. The product obtained in this reaction was carried on to the next step without further purification.

(*E*)-(2-Bromo-4-(3-(3,5-dichlorophenyl)-4,4,4-trifluorobut-1-en-1-yl)phenyl)-methanamine (CI46)




The title compound was isolated as a gummy material: The product obtained in this reaction was carried on to the next step without further purification.

(*E*)-(2-Bromo-4-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-en-1-yl)phenyl)-methanamine (CI47)



The title compound was isolated as a gummy material. The product obtained in this reaction was carried on to the next step without further purification.


(*E*)-(2-Fluoro-4-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-en-1-yl)phenyl)-methanamine (CI48)



The title compound was isolated as a gummy material: ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.40 (s, 2H), 7.33 (t, *J* = 7.6 Hz, 1H), 7.13 (m, 2H), 6.56 (d, *J* = 16.0 Hz, 1H), 6.33 (dd, *J* = 16.0, 7.6 Hz, 1H), 4.08 (m, 1H), 3.90 (s, 2H); ESIMS *m/z* 413.84 ($[M+H]^+$); IR (thin film) 3368, 3274, 1114, 808 cm^{-1}.


Prophetically, compounds **CI49**–**CI57** (Table 1) could be made in accordance with the procedures disclosed in **Example 50**.

Example 51: Preparation of 3-Chloro-4-((pyridin-2-ylamino)methyl)benzaldehyde (CI58)



To a stirred solution of 4-(bromomethyl)-3-chlorobenzaldehyde (2 g, 9 mmol) in *N,N*-dimethylacetamide (DMA; 20 mL) was added K_2CO_3 (2.36 g, 17.16 mmol) and 2-aminopyridine (0.84 g, 8.58 mmol), and the reaction mixture was stirred at ambient temperature for 4 h. The reaction mixture was diluted with water and extracted with EtOAc. The combined organic layer was washed with brine, dried over Na_2SO_4, and concentrated under reduced pressure. The residue was purified by flash column chromatography (SiO_2, 100-200 mesh; 20% EtOAc in *n*-Hexane) to afford the title compound as off-white solid (1.05 g, 50%): mp 122–123 °C; ^{1}H NMR (400 MHz, $CDCl_3$) δ 9.94 (s, 1H), 8.11 (s, 1H), 7.88 (s, 1H), 7.72 (d, *J* = 4.8 Hz, 1H), 7.62 (d, *J* = 5.7 Hz, 1H), 7.4 (m, 1H), 6.64 (d, *J* = 3.9 Hz, 1H), 6.38 (d, *J* = 6.3 Hz, 1H), 5.04 (br s, 1H), 4.71 (s, 2H); ESIMS *m/z* 246.97 ($[M+H]^+$).


Example 52: Preparation of *N*-(2-Chloro-4-vinylbenzyl)pyridin-2-amine (CI59)



To a stirred solution of 3-chloro-4-((pyridin-2-ylamino)methyl)benzaldehyde (1 g, 4. mmol) in 1,4-dioxane (20 mL) were added K_2CO_3 (0.84 g, 6.09 mmol) and methyl triphenyl phosphonium bromide (2.17 g, 6.09 mmol) at ambient temperature. Then the resultant reaction mixture was heated at 100 °C for 18 h. After the reaction was deemed complete by TLC, the reaction mixture was cooled to ambient temperature and filtered, and the obtained filtrate was concentrated under reduced pressure. The residue was purified by flash chromatography (SiO_2, 100-200 mesh; 10% EtOAc in *n*-Hexane) to afford the title compound as a white solid (0.5 g, 50%): mp 119–121 °C; ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.12 (s, 1H),


7.42 – 7.40 (m, 3H), 7.26 (s, 1H), 6.66 (m, 2H), 6.36 (d, J = 6.3 Hz, 1H), 5.75 (d, J = 13.2 Hz, 1H), 4.92 (br s, 1H), 4.60 (s, 2H); ESIMS m/z 245.05 ($[M+H]^+$).

Example 53: Preparation of Ethyl 2-amino-2-(5-bromo-3-chloropyridin-2-yl)acetate (CI60)




Ethyl 2-(diphenylmethyleneamino)acetate (10.2 g, 38.2 mmol) was added to sodium hydride (NaH; 3.18 g, 133.52 mmol) in DMF (50 mL) at 0 °C, and the mixture was stirred for 30 min. To this was added 5-bromo-2,3-dichloropyridine (12.9 g, 57.23 mmol), and the reaction mixture was stirred for 3 h at ambient temperature. The reaction mixture was quenched with 2 N HCl solution and then stirred for 4 h at ambient temperature. The mixture was extracted with EtOAc. The combined EtOAc layer was washed with brine, dried over anhydrous Na_2SO_4, and concentrated under reduced pressure. Purification by flash column chromatography (20–30% EtOAc in hexane) afforded the title compound as a liquid (1.3 g, 20%): ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.52 (s, 1H), 7.89 (s, 1H), 5.09 (s1H), 4.23 (m, 2H), 2.27 (br s, 2H), 1.26 (m, 3H); ESIMS m/z 293.05 ($[M+H]^+$); IR (thin film) 3381, 3306, 1742, 759, 523 cm^{-1}.

Example 54: Preparation of (5-Bromo-3-chloropyridin-2-yl)methanamine hydrochloride (CI61)




A stirred solution of ethyl 2-amino-2-(5-bromo-3-chloropyridin-2-yl)acetate (0.5 g, 1.7 mmol) in 3 N HCl (25 mL) was heated at reflux for 4 h. The reaction mixture was washed with Et_2O and water. The combined ether layer was concentrated under reduced pressure to afford the title compound as an off-white solid (400 mg, 65%): ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.78 (s, 1H), 8.70 (br s, 2H), 8.45 (s, 1H), 4.56 (m, 2H); ESIMS m/z 221.15 ($[M+H]^+$).

Example 55: Preparation of 2-((5-Bromo-3-chloropyridin-2-yl)methyl)isoindoline-1,3-dione (CI62)




To a stirred solution of (5-bromo-3-chloropyridin-2-yl)methanamine hydrochloride (0.3 g, 1.4 mmol) in toluene (40 mL) was added TEA (0.41 g, 4.08 mmol) and phthalic anhydride (0.24 g, 1.63 mmol), and the reaction mixture was heated at reflux for 2 h. The reaction mixture was concentrated under reduced pressure, and the residue was diluted with water and extracted with EtOAc . The combined EtOAc layer was washed with brine, dried over anhydrous Na_2SO_4, and concentrated under reduced pressure. The residue was purified by column chromatography (20–30% EtOAc in hexane) to afford the title compound as a white solid (0.25 g, 65%): ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.78 (s, 1H), 8.45 (s, 1H), 7.88 (m, 2H), 7.74 (m, 2H), 4.56 (m, 2H); ESIMS *m/z* 349 ($[M-H]^-$); IR (thin film) 3307, 1665, 1114, 813 cm^{-1}.

Example 56: Preparation of 2-((3-Chloro-5-vinylpyridin-2-yl)methyl)isoindoline-1,3-dione (CI63)




To a stirred solution of 2-((5-bromo-3-chloropyridin-2-yl)methyl)isoindoline-1,3-dione (0.23 g, 0.65 mmol) in toluene (10 mL) were added $Pd(PPh_3)_4$ (3.7 mg, 0.003 mmol), K_2CO_3 (0.269 g, 1.95 mmol) and vinyl boronic anhydride pyridine complex (0.78 g, 3.28 mmol), and the reaction mixture was heated at reflux for 16 h. The reaction mixture was filtered, and the filtrate was washed with water and brine, dried over anhydrous Na_2SO_4, and concentrated under reduced pressure. Purification by flash column chromatography (20–30% EtOAc in hexane) afforded the title compound as an off-white solid (0.2 g, 65%): ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.30 (s, 1H), 7.91 (m, 2H), 7.77 (m, 3H), 7.72 (m, 1H), 6.63 (m, 1H), 5.79 (d, J = 16.0 Hz, 1H), 5.39 (d, J = 16.0 Hz, 1H), 5.12 (s, 2H); ESIMS *m/z* 299.20 ($[M+H]^+$).

Example 57: Preparation of (*E*)-2-((3-Chloro-5-(4,4,4-trifluoro-3-(3,4,5-trichloro-phenyl)but-1-en-1-yl)pyridin-2-yl)methyl)isoindoline-1,3-dione (CI64)




To a stirred solution of 2-((3-chloro-5-vinylpyridin-2-yl)methyl)isoindoline-1,3-dione (0.35 g, 1.17 mmol) in 1,2-dichlorobenzene (10 mL) were added 5-(1-bromo-2,2,2-trifluoroethyl)-1,2,3-trichlorobenzene (0.8 g, 2.3 mmol), CuCl (23 mg, 0.12 mmol), 2,2-bipyridyl (0.073 g, 0.234 mmol), and the reaction mixture was heated at 180 °C for 16 h. The reaction mixture was concentrated under reduced pressure and purified by column chromatography (20–30% EtOAc in hexane) to afford the title compound as a liquid (0.4 g, 50%): mp 79–82 °C; ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.27 (s, 1H), 7.91 (m, 2H), 7.77 (m, 3H), 7.36 (s, 2H), 6.51 (d, *J* = 15.6 Hz, 1H), 6.32 (dd, *J* = 15.6, 8.0 Hz, 1H), 5.30 (s, 2H), 4.13 (m, 1H); ESIMS *m/z* 559 ($[M+H]^+$).

Example 58: Preparation of (*E*)-(3-Chloro-5-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-en-1-yl)pyridin-2-yl)methanamine (CI65)




To a stirred solution of (*E*)-2-((3-chloro-5-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-en-1-yl)pyridin-2-yl)methyl)isoindoline-1,3-dione (200 mg, 0.358 mmol) in EtOH (5 mL) was added hydrazine hydrate (89.6 mg, 1.79 mmol), and the reaction mixture was heated at reflux for 2 h. The reaction mixture was concentrated under reduced pressure, and the residue was dissolved in CH_2Cl_2. The organic layer was washed with water and brine, dried over anhydrous Na_2SO_4, and concentrated under reduced pressure to afford the title compound as a solid (100 mg). The product obtained in this reaction was carried on to the next step without further purification.

Example 59: Preparation of 4-(Bromomethyl)-1-naphthonitrile (CI66)




To a stirred solution of 4-methyl-1-naphthonitrile (5 g, 30 mmol) in CCl_4 (50 mL) under argon atmosphere was added NBS (6.06 g, 34.09 mmol), and the reaction mixture was degassed for 30 min. AIBN (0.3 g, 2.1 mmol) was added, and the resultant reaction mixture was heated at reflux for 4 h. The reaction mixture was cooled to ambient temperature, diluted with water and extracted with CH_2Cl_2 (3 x 100 mL). The combined CH_2Cl_2 layer was washed with brine, dried over Na_2SO_4, and concentrated under reduced pressure. The residue was purified by flash column chromatography (SiO_2, 100-200 mesh; 5% EtOAc in *n*-Hexane) to afford the title compound as a white solid (3.8 g, 52%): mp 131–133 °C; ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.33 (m, 1H), 8.24 (m, 1H), 7.88 (d, J = 8.0 Hz, 1H), 7.78 (m, 2H), 7.62 (d, J = 8.0 Hz, 1H), 4.95 (s, 2H); ESIMS *m/z* 245.92 ($[M+H]^+$); IR (thin film) 2217 cm^{-1}.

Example 60: Preparation of 4-(Bromomethyl)-1-naphthaldehyde (CI67)




To a stirred solution of 4-(bromomethyl)-1-naphthonitrile (8 g, 33mmol) in toluene (100 mL) at 0 °C was added dropwise DIBAL-H (1.0 M solution in toluene; 43 mL), and the reaction mixture was stirred at 0 °C for 1 h. 3 N HCl in water (50 mL) was added to the mixture until it became a white slurry and then additional 1 N HCl (20 mL) was added. The organic layer was collected and the aqueous layer was extracted with EtOAc (3 x100 mL). The combined organic layer was dried over Na_2SO_4 and concentrated under reduced pressure. Purification by flash column chromatography (SiO_2, 100-200 mesh; 5% EtOAc in petroleum ether) afforded the title compound as a white solid (7 g, 88%): mp 115–116 °C; ^{1}H NMR (400 MHz, $CDCl_3$) δ 10.41 (s, 1H), 9.35 (m, 1H), 8.22 (m, 1H), 7.90 (d, J = 8.0 Hz, 1H), 7.75 (m, 3H), 4.95 (s, 2H); ESIMS *m/z* 248.88 ($[M+H]^+$).

Example 61: Preparation of 4-((1,3-Dioxoisoindolin-2-yl)methyl)-1-naphthaldehyde (CI68)




To a stirred solution of 4-(bromomethyl)-1-naphthaldehyde (7 g, 28. mmol) in DMF (100 mL) was added potassium phthalimide (7.3 g, 39.5 mmol), and the mixture was heated at 85 °C for 2 h. The reaction mixture was cooled to ambient temperature and diluted with water (100 mL). The obtained solid was separated by filtration and dried under vacuum to afford the title compound as a white solid (8.8 g, 98%): mp 190–192 °C; ^{1}H NMR (400 MHz, $CDCl_3$) δ 10.39 (s, 1H), 9.25 (m, 1H), 8.41 (m, 1H), 8.10 (d, J = 8.0 Hz, 1H), 7.95 (m, 4H), 7.80 (m, 4H), 7.61 (m, 4H), 5.39 (s, 2H); ESIMS *m/z* 316.09 ([M+H]$^+$); IR (thin film) 1708 cm^{-1}.

Example 62: Preparation of 2-((4-Vinylnaphthalen-1-yl)methyl) isoindoline-1,3-dione (CI69)




To a stirred solution of 4-((1,3-dioxoisoindolin-2-yl)methyl)-1-naphthaldehyde (9 g, 28.5 mmol) in 1,4-dioxane (100 mL) were added K_2CO_3 (6 g, 42.8 mmol) and methyl triphenyl phosphonium bromide (15.3 g, 35.7 mmol) at ambient temperature. The reaction mixture was heated at 100 °C for 14 h and then was cooled to ambient temperature. The reaction mixture was filtered, and the obtained filtrate was concentrated under reduced pressure. Purification by flash chromatography (SiO_2, 100-200 mesh; 20% EtOAc in petroleum ether) afforded the title compound as a white solid (6 g, 67%): mp 146–147 °C; ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.35 (m, 2H), 7.95 (m, 4H), 7.65 (m, 4H), 7.39 (m, 1H), 5.81 (m, 1H), 5.45 (m, 1H), 5.21 (s, 2H); ESIMS *m/z* 314.13 ([M+H]$^+$).

Example 63: Preparation of (*E*)-2-((4-(4,4,4-Trifluoro-3-(3,4,5-trichlorophenyl)but-1-en-1-yl)naphthalen-1-yl)methyl)isoindoline-1,3-dione (CI70)




To a stirred solution of 2-((4-vinylnaphthalen-1-yl)methyl)isoindoline-1,3-dione (1.5 g, 4.79 mmol) in 1,2-dichlorobenzene (15 mL) were added 1-(1-bromo-2,2,2-trifluoroethyl)-3,4,5-trichlorobenzene (3.2 g, 9.5 mmol), CuCl (24 mg, 0.24 mmol) and 2,2-bipyridyl (0.149 g, 0.95 mmol), and the resultant reaction mixture was degassed with argon for 30 min and then stirred at 180 °C for 14 h. After the reaction was deemed complete by TLC, the reaction mixture was cooled to ambient temperature and filtered, and the filtrate was concentrated under reduced pressure. Purification by flash chromatography (SiO_2, 100-200 mesh; 25–30% EtOAc in petroleum ether) afforded the title compound as an off-white solid (1.5 g, 56%): mp 158–160 °C; ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.40 (m, 1H), 7.89 (m, 2H), 7.74 (m, 2H), 7.64 (m, 2H), 7.58 (m, 2H), 7.46 (s, 2H), 7.36 (m, 2H), 6.31 (m, 1H), 5.30 (s, 2H), 4.21 (m, 1H); ESIMS *m/z* 572.08 ([M-H]$^-$).

Example 64: Preparation of (*E*)-(4-(4,4,4-Trifluoro-3-(3,4,5-trichlorophenyl)but-1-en-1-yl)naphthalen-1-yl)methanamine (CI71)




To a stirred solution of (*E*)-2-((4-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-en-1-yl)naphthalen-1-yl)methyl)isoindoline-1,3-dione (0.4 g, 0.7 mmol) in EtOH was added hydrazine hydrate (0.18 g, 3.5 mmol), and the resultant reaction mixture was heated at 80 °C for 2 h. The reaction mixture was filtered, and the filtrate was concentrated. The residue was dissolved in CH_2Cl_2, and the solution was washed with brine, dried over Na_2SO_4, and concentrated under reduced pressure. The title compound was isolated as a gummy liquid (150 mg, 50%). The product obtained in this reaction was carried on to the next step without further purification.

Example 65: Preparation of 2-((4-Bromophenyl)amino)isoindoline-1,3-dione (CI72)




To a stirred solution of (4-bromophenyl)hydrazine hydrochloride (0.5 g, 2.2 mmol) in AcOH (8 mL) was added phthalic anhydride (0.398 g, 2.690 mmol), and the reaction mixture was stirred at 130 °C for 1 h under a nitrogen atmosphere. The reaction mixture was quenched with satd aqueous $NaHCO_3$ solution and filtered to give a solid. Purification by column chromatography (SiO_2, 0–10% EtOAc in petroleum ether) afforded the title compound as a solid (60 mg, 84%): mp 205–206 °C; ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.71 (s, 1H), 7.99 (m, 4H), 7.32 (d, J = 8.8 Hz, 2H), 6.79 (d, J = 8.8 Hz, 2H); ESIMS *m/z* 314.95 ([M-H]$^-$).

Example 66: Preparation of 2-((4-Vinylphenyl)amino)isoindoline-1,3-dione (CI73)




To a solution of 2-(4-bromophenylamino)isoindoline-1,3-dione (2 g, 6. mmol) in 1,2-dimethoxyethane (20 mL) and water (4 mL) were added vinyl boronic anhydride pyridine complex (4.57 g, 18.98 mmol) and K_2CO_3 (1.3 g, 9.5 mmol) followed by $Pd(PPh_3)_4$ (0.219 g, 0.189 mmol). The resultant reaction mixture was heated at 150 °C in a microwave for 30 min and then was concentrated under reduced pressure. Purification by column chromatography (SiO_2, 15% EtOAc in petroleum ether) afforded the title compound as a solid (200 mg, 13%): mp 174–176 °C; ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.65 (s, 1H), 7.94 (m, 4H), 7.29 (d, J = 8.4 Hz, 2H), 6.72 (d, J = 8.4 Hz, 2H), 6.61 (m, 1H), 5.61 (d, J = 17.6 Hz, 1H), 5.05 (d, J = 11.2 Hz, 1H); ESIMS *m/z* 263.18 ([M-H]$^-$).

Example 67: Preparation of (*E*)-2-((4-(4,4,4-Trifluoro-3-(3,4,5-trichlorophenyl)but-1-en-1-yl)phenyl)amino)isoindoline-1,3-dione (CI74)




To a stirred solution of 2-(4-vinylphenylamino)isoindoline-1,3-dione (0.3 g, 1.1 mmol) in 1,2-dichlorobenzene (5 mL) were added CuCl (0.022 g, 0.273 mmol), 2,2-bipyridyl (0.07 g, 0.46 mmol) and 5-(1-bromo-2,2,2-trifluoroethyl)-1,2,3-trichlorobenzene (0.77 g, 2.27 mmol). The reaction mixture was degassed with argon for 30 min and was heated at 180 °C for 2 h. The reaction mixture was then concentrated under reduced pressure, and the residue was purified by column chromatography (SiO_2, 0–30% EtOAc in petroleum ether) to afford the title compound as a solid (450 mg, 75%): mp 187–189 °C; ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.75 (s, 1H), 7.96 (m, 4H), 7.82 (s, 2H), 7.37 (d, J = 8.8 Hz, 1H), 6.73 (d, J = 8.4 Hz, 2H), 6.61 (m, 2H), 6.58 (m, 1H), 4.59 (m, 1H); ESIMS m/z 523.05 ([M-H]$^-$).

Example 68: Preparation of (*E*)-(4-(4,4,4-Trifluoro-3-(3,4,5-trichlorophenyl)but-1-en-1-yl)phenyl)hydrazine (CI75)



To a stirred solution of (*E*)-2-(4-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-enyl)phenylamino)isoindoline-1,3-dione (0.16 g, 0.31 mmol) in EtOH (5 mL), was added hydrazine hydrate (0.076 g, 1.52 mmol), and the reaction mixture was heated at 85 °C for 1 h. The reaction mixture was cooled to ambient temperature and filtered, and the filtrate was concentrated under reduced pressure to afford the title compound as a solid (0.08 g, 66%) which was carried on to the next step without further purification.


Example 69: Preparation of 2-(4-Vinylphenoxy)isoindoline-1,3-dione (CI76)



To a stirred solution of 4-vinylphenylboronic acid (2 g, 13 mmol), 2-hydroxyisoindoline-1,3-dione (3.63 g, 24.53 mmol), and CuCl (1.214 g 12.26 mmol) in 1,2-dichloroethane (50 mL) was added pyridine (1.065 g, 13.48 mmol), and the resultant reaction mixture was stirred at ambient temperature for 48 h. The reaction mixture was diluted with water and extracted with $CHCl_3$. The combined $CHCl_3$ layer was washed with brine, dried over Na_2SO_4 and concentrated under reduced pressure. Purification by flash column chromatography (SiO_2; 20% EtOAc in petroleum ether) afforded the title compound as a


white solid (2 g, 63%): mp 129–131 °C; ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.93 (d, *J* = 2.0 Hz, 2H), 7.82 (d, *J* = 3.2 Hz, 2H), 7.38 (d, *J* = 2.0 Hz, 2H), 7.14 (d, *J* = 2.0 Hz, 2H), 6.70 (m, 1H), 5.83 (d, *J* = 16.0 Hz, 1H), 5.22 (d, *J* = 10.8 Hz, 1H); ESIMS *m/z* 266.12 ([M+H]$^+$).

Example 70: Preparation of (*E*)-2-(4-(4,4,4-Trifluoro-3-(3,4,5-trichlorophenyl)but-1-en-1-yl)phenoxy)isoindoline-1,3-dione (CI77)




To a stirred solution of 2-(4-vinylphenoxy)isoindoline-1,3-dione (0.3g, 1.1 mmol) in 1,2-dichlorobenzene (10 mL) was added 1-(1-bromoethyl)-3,4,5-trichlorobenzene (769 mg, 2.26 mmol), CuCl (22 mg, 0.22mmol) and 2,2-bipyridyl (35 mg, 0.44 mmol), and the resultant reaction mixture was degassed with argon for 30 min and heated to 180 °C for 24 h. The reaction mixture was cooled to ambient temperature and filtered, and the filtrate was concentrated under reduced pressure. The crude material was purified by column chromatography (SiO_2, 100-200 mesh; 20% EtOAc in petroleum ether) to afford the title compound as a solid (0.29 g, 50%): ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.90 (m, 1H), 7.62 (m, 2H), 7.50 (m, 1H), 7.40 (s, 2H), 7.12 (s, 1H), 6.90 (m, 2H), 6.60 (m, 2H), 6.20 (m,1H), 4.08 (m, 1H); ESIMS *m/z* 524.09 ([M-H]$^-$).

Example 71: Preparation of (*E*)-*O*-(4-(4,4,4-Trifluoro-3-(3,4,5-trichlorophenyl)but-1-en-1-yl)phenyl)hydroxylamine (CI78)



To a stirred solution of (*E*)-2-(4-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-enyl)phenoxy)isoindoline-1,3-dione (0.2 g, 0.4 mmol) in EtOH was added hydrazine hydrate (0.1 g, 1.9 mmol), and the resultant reaction mixture was heated at 90 °C for 1 h. The reaction mixture was filtered, and the filtrate was concentrated. The residue was dissolved in CH_2Cl_2. washed with brine, dried over Na_2SO_4 and concentrated under reduced pressure to afford the crude title compound as a gummy liquid (0.08 g, 53%): ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.40 (s, 2H), 6.98 (s, 1H), 6.82 (s, 2H), 6.48 (m, 1H), 6.20 (m, 1H), 5.02 (s, 1H), 4.08 (m, 1H); ESIMS *m/z* 394.94 ([M-H]$^-$).


Example 72: Preparation of (*E*)-*N*-(4-(3-(3,5-Dichlorophenyl)-4,4,4-trifluorobut-1-enyl)benzyl)acetamide (CC1)



To a stirred solution of (*E*)-(2-chloro-4-(3-(3,5-dichlorophenyl)-4,4,4-trifluorobut-1-en-1-yl)phenyl)methanamine (0.3 g, 0.8 mmol) in CH_2Cl_2 (10 mL) was added acetic anhydride (0.12 mL, 1.14 mmol), and TEA (0.217 mL, 1.52 mmol), and the resultant reaction mixture was stirred at ambient temperature for 6 h. The reaction mixture was diluted with water and extracted with CH_2Cl_2. The combined CH_2Cl_2 layer was washed with brine, dried over Na_2SO_4, and concentrated under reduced pressure. Purification by flash column chromatography (SiO_2, 100-200 mesh; 30–50% EtOAc in hexane) afforded the title compound as an off-white solid (0.2 g, 60%) mp 107–109 °C; ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.37 (m, 3H), 7.28 (m, 4H), 6.60 (d, J = 16.0 Hz, 1H), 6.36 (dd, J = 16.0, 8.0 Hz, 1H), 5.75 (br s, 1H), 4.46 (d, J = 6 Hz, 2H), 4.01 (m, 1H), 2.11 (s, 3H); ESIMS *m/z* 402.00 ($[M+H]^+$).

Compounds **CC2** – **CC6** in Table 1 were made in accordance with the procedures disclosed in **Example 72**. In addition, compound **DC56** in Table 1 was made from compound **DC55** in accordance with the procedures disclosed in **Example 72**.

Example 73: Preparation of (*E*)-*N*-(2-Chloro-4-(3-(3,5-dichlorophenyl)-4,4,4-trifluorobut-1-en-1-yl)benzyl)acetamide (CC7)




To a stirred solution of (*E*)-(2-chloro-4-(3-(3,5-dichlorophenyl)-4,4,4-trifluorobut-1-en-1-yl)phenyl)methanamine (0.3 g, 0.8 mmol) in DMF (5 mL) was added 2,2,2-trifluoro-propanoic acid (97 mg, 0.76 mmol), HOBt•H_2O (174 mg, 1.14 mmol) and EDC•HCl (217 mg, 1.14 mmol) and DIPEA (196 mg, 1.52 mmol), and the resultant reaction mixture was stirred at ambient temperature for 18 h. The reaction mixture was diluted with water and extracted with EtOAc. The combined EtOAc layer was washed with brine, dried over Na_2SO_4, and concentrated under reduced pressure. Purification by flash column chromatography (SiO_2, 100-200 mesh; EtOAc in hexane (30–50% afforded the title compound as an off-white solid (0.2 g, 60%): mp 127–128 °C; ^{1}H NMR (400 MHz, $CDCl_3$)

δ 7.42 (m, 4H), 7.24 (m, 2H), 6.53 (d, *J* = 16.0 Hz, 1H), 6.36 (dd, *J* = 16.0, 8.0 Hz, 1H) , 5.86 (br s, 1H), 4.51 (d, *J* = 6.0 Hz, 2H), 4.05 (m, 1H), 2.02 (s, 3H); ESIMS *m/z* 436.03 ($[M+H]^+$).

Compounds **CC8** – **CC28** in Table 1 were made in accordance with the procedures disclosed in **Example 73**.

Example 74: Preparation of (*E*)-*N*-(Pyridin-2-ylmethyl)-*N*-(4-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-enyl)-2-(trifluoromethyl)benzyl)cyclopropanecarboxamide (CC29)




Step 1: (*E*)-1-(Pyridin-2-yl)-*N*-(4-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-enyl)-2-(trifluoromethyl)benzyl)methanamine. (*E*)-(4-(4,4,4-Trifluoro-3-(3,4,5-trichlorophenyl)but-1-en-1-yl)-2-(trifluoromethyl)phenyl)methanamine (0.46 g, 1 mmol) was dissolved in MeOH (3 mL). To this was added pyridine-2-carbaldehyde (0.107 g, 1 mmol). The reaction mixture was stirred for 1 h. After 1 h, $NaBH_4$ (0.076 g, 2 mmol) was added and left at ambient temperature for 3 h. The reaction mixture was concentrated to give an oily residue. Purification by flash column chromatography (SiO_2, 100-200 mesh; 30–50% EtOAc in hexane) afforded the title compound as a pale yellow liquid (0.22 g, 40%): ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.58 (d, *J*= 4.8 Hz, 1H), 7.74 (m, 1H), 7.62 (m, 2H), 7.52 (m, 1H), 7.4 (s, 2H), 7.3 (m, 1H), 7.2 (m, 2H), 6.60 (d, *J* = 16.0 Hz, 1H), 6.38 (dd, *J* = 16.0, 8.0 Hz, 1H), 4.10 (m, 1H), 4.02 (s, 2H), 3.96 (s, 2H); ESIMS *m/z* 552.95 ($[M+H]^+$); IR (thin film) 3338, 1114, 808 cm^{-1}.

Step 2: (*E*)-*N*-(Pyridin-2-ylmethyl)-*N*-(4-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-enyl)-2-(trifluoromethyl)benzyl)cyclopropanecarboxamide. (*E*)-1-(Pyridin-2-yl)-*N*-(4-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-en-1-yl)-2-(trifluoromethyl)benzyl)methanamine (0.27 g, 0.05 mmol) was taken up in CH_2Cl_2 (3 mL). To this was added TEA (0.14 mL, 0.1 mmol). The reaction mixture was stirred for 10 min. After 10 min, the reaction mixture was cooled to 0 °C, and cyclopropylcarbonyl chloride (0.08 mL, 0.075 mmol) was added. The reaction mixture was stirred at ambient temperature for 1 h and then was washed with water and satd aq $NaHCO_3$ solution. The organic layer was dried over anhydrous Na_2SO_4 and evaporated to obtain pale yellow gummy material (0.15 g,


50%): ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.58 (d, *J* = 4.6 Hz, 1H), 7.74 (m, 1H), 7.62 (m, 2H), 7.52 (m, 1H), 7.4 (s, 2H), 7.3 (m, 1H), 7.2 (m, 2H), 6.60 (d, *J* = 16.0 Hz, 1H), 6.38 (dd, *J* = 16.0, 8.0 Hz, 1H), 5.02 (s, 1H), 4.8 (s, 1H), 4.8 (d, *J* = 10 Hz, 2H), 4.10 (m, 1H), 1.8 (m, 1H), 1.2 (m, 2H), 0.6 (m, 2H); ESIMS *m/z* 620.86 ([M-H]$^-$); IR (thin film) 1645, 1115, 808 cm^{-1}.

Example 75: Preparation of (*E*)-*N*-(2-Chloro-4-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-en-1-yl)benzyl)-3-(methylsulfonyl)propanamide (CC30)



(*E*)-*N*-(2-Chloro-4-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-en-1-yl)benzyl)-3-(methylthio)propanamide (0.15 g, 0.28 mmol) was treated with oxone (0.175 g, 0.569 mmol) in 1:1 acetone:water (20mL) for 4 h at ambient temperature. The acetone was evaporated to obtain a white solid (0.095 g, 60%): mp 101–104 °C; ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.41 (m, 4H), 7.24 (m, 1H), 6.53 (d, *J* = 16.0 Hz, 1H), 6.35 (dd, *J* = 16.0, 8.0 Hz, 1H), 6.12 (br s, 1H), 4.53 (m, 2H), 4.10 (m, 1H), 3.42 (m, 2H), 2.91 (s, 3H), 2.78 (m, 2H); ESIMS *m/z* 559.75 ([M-H]$^-$).


Example 76: Preparation of (*E*)-1-(2-Chloro-4-(3-(3,5-dichlorophenyl)-4,4,4-trifluorobut-1-en-1-yl)benzyl)-3-ethylurea (CC31)



To a stirred solution of (*E*)-(2-chloro-4-(3-(3,5-dichlorophenyl)-4,4,4-trifluorobut-1-en-1-yl)phenyl)methanamine (0.2 g, 0.5 mmol) in CH_2Cl_2 (5 mL) at 0 °C were added TEA (0.141 mL, 1 mmol) and ethylisocyanate (0.053 g, 0.75 mmol), and the reaction mixture was stirred for 1 h at 0 °C. The reaction mixture was diluted with CH_2Cl_2. The organic layer was washed with water and brine, dried over Na_2SO_4, and concentrated under reduced pressure. Purification by column chromatography (SiO_2, 100-200 mesh; 30–50% EtOAc in hexane) afforded the title compound as a solid (0.141 g, 60%): mp 177–178 °C; ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.58 (m, 2H), 7.41 (m, 3H), 7.24 (m, 1H), 6.53 (d, *J* = 16.0 Hz, 1H), 6.35 (dd, *J* = 16.0, 8.0 Hz, 1H), 4.70 (br s, 1H), 4.43 (s, 2H), 4.08 (m, 1H), 3.21 (m, 2H), 1.25 (m, 3H); ESIMS *m/z* 463 ([M-H]$^-$).


Compounds **CC32** – **CC35** in Table 1 were made in accordance with the procedures disclosed in **Example 76**.

Example 77: Preparation of (*E*)-3-(2-Chloro-4-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-en-1-yl)benzyl)-1,1-dimethylurea (CC36)



To a stirred solution of (*E*)-(2-chloro-4-(3-(3,4,5-trichlorophenyl)-4,4,4-trifluorobut-1-en-1-yl)phenyl)methanamine (0.2 g, 0.5 mmol) in CH_2Cl_2 (5 mL) at 0 °C were added TEA (0.141 mL, 1 mmol) and *N,N*-dimethylcarbamoyl chloride (0.08 g, 0.075 mmol), and the reaction mixture was stirred for 1 h at 0 °C. The reaction mixture was diluted with CH_2Cl_2. The organic layer was washed with water and brine, dried over Na_2SO_4, and concentrated under reduced pressure. Purification by column chromatography (SiO_2, 100-200 mesh; 30–50% EtOAc in hexane) afforded the title compound as a solid (0.15 g, 60%): ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.39 (m, 4H), 7.28 (m, 1H), 6.54 (d, *J* = 16.0 Hz, 1H), 6.34 (dd, *J* = 16.0, 8.0 Hz, 1H), 4.97 (br s, 1H), 4.38 (d, *J* = 6.0 Hz, 2H), 4.10 (m, 1H), 2.9 (s, 3H), 2.7 (s, 3H); ESIMS *m/z* 497 ([M-H]$^-$); IR (thin film) 3350, 1705, 1114, 808 cm^{-1}.


Example 78: Preparation of (*E*)-1-(2-Chloro-4-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-en-1-yl)benzyl)-3-ethylthiourea (CC37)



To a stirred solution of (*E*)-(2-chloro-4-(3-(3,4,5-trichlorophenyl)-4,4,4-trifluorobut-1-en-1-yl)phenyl)methanamine (0.2 g, 0.5 mmol) in CH_2Cl_2 (5 mL) at 0 °C were added TEA (0.141 mL, 1 mmol) and ethyl isothicyanate (0.053 g, 0.75 mmol), and the reaction mixture was stirred for 1 h at 0 °C. The reaction mixture was diluted with CH_2Cl_2. The organic layer was washed with water and brine, dried over Na_2SO_4, and concentrated under reduced pressure. Purification by column chromatography (SiO_2, 100-200 mesh; 30–50% EtOAc in hexane) afforded the title compound as a solid (0.14 g, 60%): mp 88–91 °C; ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.49 (d, *J* = 8 Hz, 1H), 7.41 (d, *J* = 7.2 Hz, 2H), 7.26 (m, 2H), 6.50 (d, *J* = 16 Hz, 1H), 6.35 (dd, *J* = 16.0, 8.0 Hz, 1H), 6.0 (br s, 1H), 5.73 (br s, 1H), 4.80 (br s, 2H), 4.09 (m, 1H), 1.23 (m, 3H); ESIMS *m/z* 515.01 ([M+H]$^+$).

Compound **CC38** in Table 1 was made in accordance with the procedures disclosed in **Example 78**.


Example 79: Preparation of (*E*)-*tert*-Butyl (2-chloro-4-(3-(3,5-dichlorophenyl)-4,4,4-trifluorobut-1-en-1-yl)benzyl)-3-ethylurea (CC39)



To a stirred solution of (*E*)-(2-chloro-4-(3-(3,4,5-trichlorophenyl)-4,4,4-trifluorobut-1-en-1-yl)phenyl)methanamine (0.2 g, 0.5 mmol in CH_2Cl_2 (5 mL) at 0 °C were added TEA (0.141 mL, 1 mmol) and di-*tert*-butyl dicarbonate (0.163 mL, 0.75 mmol), and the reaction mixture was stirred for 4 h at ambient temperature. The reaction mixture was diluted with CH_2Cl_2. The organic layer was washed with water and brine, dried over Na_2SO_4, and concentrated under reduced pressure. Purification by column chromatography (SiO_2, 100-200 mesh; 10–20% EtOAc in hexane) afforded the title compound as a white solid (0.147 g, 60%): ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.39 (m, 4H), 7.28 (m, 1H), 6.54 (d, *J* = 16.0 Hz, 1H), 6.34 (dd, *J* = 16.0, 8.0 Hz, 1H), 4.97 (br s, 1H), 4.38 (d, *J* = 6.0 Hz, 2H), 4.10 (m, 1H), 1.53 (s, 9H); ESIMS *m/z* 526.09 ([M-H]$^-$); IR (thin film) 3350, 1705, 1114, 808 cm^{-1}.

Compound **CC40** in Table 1 was made in accordance with the procedures disclosed in **Example 79**.


Example 80: Preparation of (*E*)-Methyl 2-((2-chloro-4-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-en-1-yl)benzyl)amino)-2-oxoacetate (CC41)



To a stirred solution of (*E*)-(2-chloro-4-(3-(3,4,5-trichlorophenyl)-4,4,4-trifluorobut-1-en-1-yl)phenyl)methanamine (0.2 g, 0.5 mmol) in CH_2Cl_2 (5 mL) at 0 °C were added TEA (0.141 mL, 1 mmol) and methyl 2-chloro-2-oxoacetate (0.09 g, 0.75 mmol), and the reaction mixture was stirred for 1 h at 0 °C. The reaction mixture was diluted with CH_2Cl_2. The organic layer was washed with water and brine, dried over Na_2SO_4, and concentrated under reduced pressure. Purification by column chromatography (SiO_2, 100-200 mesh; 20% EtOAc in hexane) afforded the title compound as a solid (0.12 g, 50%): ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.48 (m, 1H). 7.43 (m, 3H), 7.38 (m, 1H), 7.23 (s, 1H), 6.55 (d, *J* = 16.0 Hz, 1H), 6.36 (dd,


J = 16.0, 8.0 Hz, 1H), 4.60 (d, J = 4.4 Hz, 2H), 4.18 (m, 1H), 3.85 (s, 3H); ESIMS m/z 512.22 ([M-H]$^-$); IR (thin film) 1740, 1701, 1114, 808 cm^{-1}.

Example 81: Preparation of (*E*)-*N*^{1}-(2-Chloro-4-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-en-1-yl)benzyl)-*N*^{2}-(2,2,2-trifluoroethyl)oxalamide (CC42)



To a stirred solution of 2,2,2-trifluoroethylamine hydrochloride (0.1 g, 0.77 mmol) in CH_2Cl_2 (10 mL) was added dropwise trimethylaluminum (2 M solution in toluene; 0.39 mL, 0.77 mmol), and the reaction mixture was stirred at 25 °C for 30 min. A solution of (*E*)-methyl 2-((2-chloro-4-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-en-1-yl)benzyl)-2-oxoacetate (0.2 g, 0.38 mmol) in CH_2Cl_2 (5 mL) was added dropwise to the reaction mixture at 25 °C. The reaction mixture was stirred at reflux for 18 h, cooled to 25 °C, quenched with 0.5 N HCl solution (50 mL) and extracted with EtOAc (2 x 50 mL). The combined organic extracts were washed with brine, dried over Na_2SO_4, and concentrated under reduced pressure. The crude compound was purified by flash chromatography (SiO_2, 100-200 mesh; 20%–40% EtOAc in *n*-hexane) to afford the title compound (0.13 g, 60%): mp 161–163 °C; ^{1}H NMR (400 MHz, DMSO-d_6) δ 9.45 (br s, 2H), 7.90 (s, 2H), 7.75 (s, 1H), 7.46 (s, 1H), 7.28 (s, 1H), 6.93 (m, 1H), 6.75 (m, 1H), 4.80 (m, 1H), 4.40 (s, 2H), 3.90 (s, 2H); ESIMS m/z 578.96 ([M-H]$^-$).


Example 82: Preparation of (*E*)-*N*-(2-Chloro-4-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-en-1-yl)benzyl)pyridin-2-amine (CC43)



To a stirred solution of *N*-(2-chloro-4-vinylbenzyl)pyridin-2-amine (0.3 g, 1.22 mmol) in 1,2-dichlorobenzene (5 mL) were added 5-(1-bromo-2,2,2-trifluoroethyl)-1,2,3-trichlorobenzene (0.83 g, 2.44 mmol), CuCl (24 mg, 0.24 mmol) and 2,2-bipyridyl (76 mg, 0.48 mmol). The resultant reaction mixture was degassed with argon for 30 min and then stirred at 180 °C for 24 h. After the reaction was deemed complete by TLC, the reaction mixture was cooled to ambient temperature and filtered, and the filtrate was concentrated under reduced pressure. Purification by flash chromatography (SiO_2, 100-200 mesh; 15%

EtOAc in *n*-hexane) afforded the title compound as an off-white solid (0.2 g, 35%): mp 140–142 °C; ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.11 (d, *J* = 4.0 Hz, 1H), 7.40 (m, 5H), 7.22 (m, 1H), 6.61 (m, 2H), 6.35 (m, 2H), 4.94 (br s, 1H), 4.61 (d, *J* = 6.4 Hz, 2H), 4.11 (m, 1H); ESIMS *m/z* 505.39 ([M+H]$^+$).


Example 83: Preparation of (*E*)-*N*-((3-Chloro-5-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)-but-1-en-1-yl)pyridin-2-yl)methyl)-3,3,3-trifluoropropanamide (CC44)



To a stirred solution of (*E*)-(3-chloro-5-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-en-1-yl)pyridin-2-yl)methanamine (0.1 g, 0.2 mmol) in CH_2Cl_2 (5 mL) were added 3,3,3-trifluoropropanoic acid (45 mg, 0.350 mmol), EDC•HCl (67 mg, 0.350 mmol), HOBt•H_2O (71 mg, 0.467 mmol) and DIPEA (60.2 mg, 0.467 mmol), and the reaction mixture was stirred at ambient temperature for 18 h. The reaction mixture was diluted with CH_2Cl_2 and washed with water. The combined CH_2Cl_2 layer was washed with brine, dried over anhydrous Na_2SO_4, and concentrated under reduced pressure. Purification by flash column chromatography (SiO_2, 100-200 mesh; 15% EtOAc in petroleum ether) afforded the title compound as a pale yellow liquid (30 mg, 35%): ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.41 (s, 1H), 7.77 (s, 1H), 7.47 (br s, 1H), 7.40 (s, 2H), 6.58 (d, *J* = 16.0 Hz, 1H), 6.45 (dd, *J* = 16.0, 8.0 Hz, 1H), 4.68 (d, *J* = 4.0 Hz, 2H), 4.14 (m, 1H), 3.24 (q, *J* = 10.8 Hz, 2H); ESIMS *m/z* 536.88 ([M-H]$^-$); IR (thin film) 3320, 1674, 1114, 808.

Compound **CC45** in Table 1 was made in accordance with the procedures disclosed in **Example 83**.

Example 84: Preparation of (*E*)-3,3,3-Trifluoro-*N*-((4-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-en-1-yl)naphthalen-1-yl)methyl)propanamide (CC46)




To a stirred solution of (*E*)-(4-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-en-1-yl)naphthalen-1-yl)methanamine (0.1 g, 0.22 mmol) in CH_2Cl_2 (8 mL) were added 3,3,3-trifluoropropanoic acid (0.032 g, 0.24 mmol), HOBt•H_2O (52 mg, 0.33 mmol), EDC•HCl (0.065 g, 0.33 mmol) and DIPEA (0.044 g, 0.45 mmol), and the resultant reaction mixture

was stirred at ambient temperature for 18 h. The reaction mixture was diluted with water and extracted with EtOAc (3 x30 mL). The combined EtOAc layer was washed with brine, dried over Na_2SO_4, and concentrated under reduced pressure. Purification by flash column chromatography (SiO_2, 100-200 mesh; 15% EtOAc in *n*-hexane) afforded the title compound as a gummy material (60 mg, 50%): mp 151–153 °C; ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.06 (m, 1H), 7.61 (m, 4H), 7.48 (s, 2H), 7.44 (d, J = 8.0 Hz, 1H), 7.38 (m, 1H), 6.42 (m, 1H), 5.92 (br s, 1H), 4.92 (m, 2H), 4.24 (m, 1H), 3.12 (m, 2H); ESIMS *m/z* 554.04 ([M-H]$^-$).


Compounds **CC47** – **CC48** in Table 1 were made in accordance with the procedures disclosed in **Example 84**.

Example 85: Preparation of (*E*)-1-Ethyl-3-((4-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-en-1-yl)naphthalen-1-yl)methyl)urea (CC49)



To a stirred solution of (*E*)-(4-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-en-1-yl)naphthalen-1-yl)methanamine (0.1 g, 0.22 mmol) in CH_2Cl_2 at 0 °C were added TEA (0.064 mL, 0.44 mmol) and ethylisocyanate (0.023 mL, 0.33 mmol), and the reaction mixture was stirred for 1 h at 0 °C. The reaction mixture was diluted with CH_2Cl_2. The organic layer was washed with water and brine, dried over Na_2SO_4, and concentrated under reduced pressure. Purification by column chromatography (SiO_2, 100-200 mesh; 30% EtOAc in hexane) afforded the title compound as a solid (0.07 g, 60%): mp 84–87 °C; ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.06 (m, 1H), 7.98 (m, 1H), 7.61 (m, 3H), 7.48 (s, 2H), 7.44 (d, J = 8.0 Hz, 1H), 7.38 (m, 2H), 6.42 (m, 1H), 4.92 (s, 2H), 4.6 (br s, 1H), 4.24 (m, 1H), 3.21 (m, 2H), 1.2 (t, J = 4.6 Hz, 3H); ESIMS *m/z* 515.33 ([M+H]$^+$).

Example 86: Preparation of (*E*)-*N*'-(4-(4,4,4-Trifluoro-3-(3,4,5-trichlorophenyl)but-1-en-1-yl)phenyl)cyclopropanecarbohydrazide (CC50)




To a stirred solution of (*E*)-(4-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-en-1-yl)phenyl)hydrazine (0.1 g, 0. 3 mmol) in CH_2Cl_2 (10 mL) was added DIPEA (65 mg, 0.51 mmol), HOBt•H_2O (59 mg, 0.38 mmol), EDC•HCl (73 mg, 0.38 mmol) and

cyclopropanecarbonyl chloride (0.024 g, 0.28 mmol), and the reaction mixture was stirred at ambient temperature for 1 h. The reaction mixture was diluted with satd aq $NaHCO_3$ solution and extracted with CH_2Cl_2. The combined CH_2Cl_2 layer was washed with brine, dried over anhydrous Na_2SO_4, and concentrated under reduced pressure. Purification by flash column chromatography (SiO_2; 5–25% EtOAc in petroleum ether) afforded the title compound as a solid (65 mg, 55%): mp 138–140 °C; ^{1}H NMR (400 MHz, $CDCl_3$) δ 9.81 (s, 1H), 7.90 (s, 1H), 7.84 (s, 2H), 7.34 (d, *J* = 8.4 Hz, 2H), 6.65 (d, *J* = 15.6 Hz, 1H), 6.61 (m, 1H), 6.57 (s, 1H), 6.48 (dd, *J* = 15.6, 8.8 Hz, 1H), 4.74 (m, 1H), 1.64 (m, 1H), 0.75 (m, 4H); ESIMS *m/z* 461.32 ([M-H]$^-$).

Compound **CC51** in Table 1 was made in accordance with the procedures disclosed in **Example 86**.


Example 87: Preparation of (*E*)-*N*-(4-(4,4,4-Trifluoro-3-(3,4,5-trichlorophenyl)but-1-en-1-yl)phenoxy)cyclopropanecarboxamide (CC52)



To a stirred solution of (*E*)-*O*-(4-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-en-1-yl)phenyl)hydroxylamine (0.15 g, 0.38 mmol) in CH_2Cl_2 (5 mL) was added EDC•HCl (0.109 g, 0.569 mmol), HOBt•H_2O (0.087 g, 0.569 mmol), DIPEA (0.097 g, 0.758 mmol) and cyclopropanecarboxylic acid (0.049 g, 0.569 mmol). The resultant reaction mixture was stirred at ambient temperature for 18 h. The reaction mixture was diluted with water and extracted with $CHCl_3$ (35 mL) The combined $CHCl_3$ layer was washed with brine, dried over Na_2SO_4 and concentrated under reduced pressure. Purification by flash column chromatography (SiO_2; 20% EtOAc in hexane) afforded the title compound as a brown liquid (0.06 g, 34%): ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.40 (s, 2H), 7.18 (s, 1H), 7.08 (s, 1H), 6.85 (m, 1H), 6.45 (m, 1H), 6.65 (m, 1H), 6.20 (m, 1H), 5.55 (s, 1H), 4.08 (m, 1H), 1.90 (m, 1H), 1.30 – 1.10 (m, 4H); ESIMS *m/z* 464.87 ([M-H]$^-$).

Compound **CC53** in Table 1 was made in accordance with the procedures disclosed in **Example 87**.


Example 88: Preparation of (*Z*)-3,3,3-Trifluoro-*N*-(4-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-en-1-yl)benzyl)propanamide (CC54)



A silicon borate vial was charged with (*E*)-3,3,3-trifluoro-*N*-(4-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-en-1-yl)benzyl)propanamide (133 mg, 0.269 mmol) and dimethyl sulfoxide (DMSO; 10 mL). The mixture was placed within 0.6 to 1 meter (m) of a bank of eight 115 watt Sylvania FR48T12/350BL/VHO/180 Fluorescent Tube Black Lights and four 115 watt Sylvania (daylight) F48T12/D/VHO Straight T12 Fluorescent Tube Lights for 72 h. The mixture was concentrated *in vacuo* and purified by reverse phase chromatography to give the title compound as a colorless oil (11 mg, 8%): ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.28 (s, 2H), 7.25 (m, 2H), 7.10 (d, *J* = 8.0 Hz, 2H), 6.89 (d, *J* = 11.4 Hz, 1H), 6.07 (br s, 1H), 6.01 (m, 1H), 4.51 (d, *J* = 5.8 Hz, 2H), 4.34 (m, 1H), 3.12 (q, *J* = 7.5 Hz, 2H); ^{13}C NMR (101 MHz, $CDCl_3$) δ 162.44, 137.20, 135.38, 135.23, 134.82, 134.68, 131.71, 129.00, 128.80, 128.69, 128.10, 127.96, 122.63,76.70, 47.33 (q, *J* = 28 Hz), 43.59, 42.12 (q, *J* = 30 Hz); ESIMS *m/z* 504 ([M+H]$^+$).

Compounds **DC46, AC93. AC94** in Table 1 were made in accordance with the procedures disclosed in **Example 88**.

Example 89: Preparation of 1-(1-Bromo-2,2,2-trifluoroethyl)-3-chlorobenzene (DI2)




DI1 **DI2**

The title compound was synthesized in two steps via 1-(3-chlorophenyl)-2,2,2-trifluoroethanol (**DI1**, prepared as in Step 1, Method B in Example 1); isolated as a colorless viscous oil (1.5 g, 75%): ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.50 (s, 1H), 7.42-7.35 (m, 3H), 5.02 (m, 1H), 2.65 (br s, 1H)) and Step 2 in Example 1 and isolated (0.14 g, 22%): ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.50 (br s, 1H), 7,42-7.35 (m, 3H), 5.07 (m, 1H).

The following compounds were made in accordance with the procedures disclosed in **Example 89**.


(1-Bromo-2,2,2-trifluoroethyl)benzene (DI4)



DI3 **DI4**

2,2,2-Trifluoro-1-phenylethanol (**DI3**) was isolated (10 g, 80%): ^{1}H NMR (300 MHz, $CDCl_3$) δ 7.48 (m, 2H), 7.40 (m, 3H), 5.02 (m, 1H), 2.65 (d, J = 7.1 Hz, 1H). The title compound (**DI4**) was isolated as a liquid (8.0 g, 60%): ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.50 (m, 2H), 7.40 (m, 3H), 5.00 (q, J = 7.5 Hz, 1H).


1-(1-Bromo-2,2,2-trifluoroethyl)-3,5-dimethylbenzene (DI20)



DI19 **DI20**

1-(3,5-Dimethylphenyl)-2,2,2-trifluoroethanol (**DI19**) was isolated an off white solid : ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.05 (s, 2H), 7.02 (s, 1H), 4.95 (m, 1H), 2.32 (s, 6H); ESIMS m/z 204 ($[M]^-$). The title compound (**DI20**) was isolated (3.0 g, 51%).


1-(1-Bromo-2,2,2-trifluoroethyl)-2,4-dichlorobenzene (DI22)



DI21 **DI22**

1-(2,4-Dichlorophenyl)-2,2,2-trifluoroethanol (**DI21**) was isolated as an off white powder (5.3 g, 61%): mp 49-51 °C; ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.62-7.66 (d, 1H), 7.42-7.44 (d, 1H), 7.32-7.36 (d, 1H), 5.6 (m, 1H), 2.7 (s, 1H); ESIMS m/z 244 ($[M]^+$). The title compound (**DI22**) was isolated (3.2 g, 50%): ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.62-7.72 (m, 1H), 7.4-7.42 (m, 1H), 7.3-7.38 (m, 1H), 5.7-5.8 (m, 1H).


1-(1-Bromo-2,2,2-trifluoroethyl)-2,3-dichlorobenzene (DI24)



DI23 **DI24**


1-(2,3-Dichlorophenyl)-2,2,2-trifluoroethanol (**DI23**) was isolated as a pale yellow oil (5.2 g, 60%): ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.62-7.64 (d, 1H), 7.52-7.54 (m, 1H), 7.29-7.33 (t, 1H), 5.6-5.76 (m, 1H), 2.7 (s, 1H); ESIMS *m/z* 243.9 ([M]$^+$). The title compound (**DI24**) was isolated as an oil (8.7 g, 60%): ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.62-7.71 (m, 1H), 7.44-7.52 (m, 1H), 7.27-7.3 (s, 1H), 5.81-5.91 (m, 1H).

2-(1-Bromo-2,2,2-trifluoroethyl)-1,4-dichlorobenzene (DI26)




1-(2,5-Dichlorophenyl)-2,2,2-trifluoroethanol (**DI25**) was isolated as a yellow oil (4.1 g, 60%): ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.68-7.7 (s, 1H), 7.3-7.37 (m, 2H), 5.51-5.6 (m, 1H), 2.7 (s, 1H); ESIMS *m/z* 244 ([M]$^+$)). The title compound (**DI26**) was isolated (3.0 g, 60%): ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.7-7.78 (m, 1H), 7.3-7.4 (m, 2H), 5.7-5.8 (m, 1H).

1-(1-Bromo-2,2,2-trifluoroethyl)-3,5-bis(trifluoromethyl)benzene (DI28)




1-(3,5-Bis(trifluoromethyl)phenyl)-2,2,2-trifluoroethanol (**DI27**) was isolated (3.8 g, 60%): ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.98 (m, 3H), 5.25 (m, 1H), 3.2 (br, 1H); ESIMS *m/z* 312.2 ([M]$^+$). The title compound (**DI28**) was prepared and carried on crude.

1-(1-Bromo-2,2,2-trifluoroethyl)-2,3,5-trichlorobenzene (DI30)




2,2,2-Trifluoro-1-(2,3,5-trichlorophenyl)ethanol (**DI29**) was isolated as a white solid (4.0 g, 60%): mp 113-115 °C; ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.62 (d, 1H), 7.50 (d, 1H), 5.60-5.70 (m, 1H), 2.75 (s, 1H); ESIMS *m/z* 278.0 ([M$^+$]). The title compound (**DI30**) was isolated (2.9 g, 60%): ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.70 (d, 1H), 7.50 (d, 1H), 5.72-5.82 (m, 1H).

1-(1-Bromo-2,2,2-trifluoroethyl)-3-chloro-5-(trifluoromethyl)benzene (DI32)




1-(3-Chloro-5-(trifluoromethyl)phenyl)-2,2,2-trifluoroethanol (**DI31**) was isolated as a pale yellow oil (2.0 g, 50%): ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.51 (m, 3H), 5.08 (m, 1H), 2.81 (s, 1H); ESIMS *m/z* 278.1 ($[M]^+$). The title compound (**DI32**) was isolated oil (2.0 g, 40%): ESIMS *m/z* 342 ($[M]^+$).

5-(1-Bromo-2,2,2-trifluoroethyl)-1,3-dichloro-2-methoxybenzene (DI34)



1-(3,5-Dichloro-4-methoxyphenyl)-2,2,2-trifluoroethanol (**DI33**) was isolated as an off white solid (0.8 g, 60%); mp 92-95 °C: ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.41 (s, 2H), 5.00 (m, 1H), 3.89 (s, 3H), 2.64 (m, 1H); ESIMS *m/z* 274 ($[M]^+$). The title compound (**DI34**) was isolated as a colorless liquid (0.6 g, 57%).


Example 90: Preparation of 1-(1-Bromo-2,2,2-trifluoroethyl)-3,5-difluorobenzene (DI36)



The title compound was synthesized in two steps via 1-(3,5-difluorophenyl)-2,2,2-trifluoroethanol (**DI35**, prepared as in Step 1, Method A in Example 1; isolated as a colorless oil (0.2 g, 75%): ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.05 (m, 2H), 6.88 (m, 1H), 5.06 (m, 1H), 2.66 (s, 1H); ESIMS *m/z* 212 ($[M]^+$) and Step 2 in Example 1 and isolated (3.2 g, 50%); ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.05 (m, 2H), 6.86 (m, 1H), 5.03 (q, *J* = 7.4 Hz, 1H).

The following compounds were made in accordance with the procedures disclosed in **Example 90**.


1-(1-Bromo-2,2,2-trifluoroethyl)-4-chlorobenzene (DI38)



DI37 **DI38**

1-(4-Chlorophenyl)-2,2,2-trifluoroethanol (**DI37**) was isolated as a colorless oil (5.0 g, 99%): ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.44-7.38 (m, 4H), 5.05 (m, 1H), 2.55 (s, 1H); ESIMS *m/z* 210 ($[M]^+$). The title compound (**DI38**) was isolated (3.0 g, 46 %): ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.45 (d, *J* = 8.2 Hz, 2H), 7.37 (d, *J* = 8.2 Hz, 2H), 5.10 (q, *J* = 7.2 Hz, 1H).


1-(1-Bromo-2,2,2-trifluoroethyl)-4-methoxybenzene (DI40)



DI39 **DI40**

2,2,2-Trifluoro-1-(4-methoxyphenyl)ethanol (**DI39**) was isolated as a pale yellow liquid: ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.41 (d, *J* = 8.8 Hz, 2H), 6.95 (m, *J* = 8.8 Hz, 2H), 5.00 (m, 1H), 3.82 (s, 3H), 2.44 (s, 1H); ESIMS *m/z* 206.1 ($[M]^+$). The title compound (**DI40**) was isolated (3.8 g, 62%).


1-(1-Bromo-2,2,2-trifluoroethyl)-4-fluorobenzene (DI42)



DI41 **DI42**

2,2,2-Trifluoro-1-(4-fluorophenyl)ethanol (**DI41**) was isolated as a colorless oil (5 g, 99%): ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.48-7.45 (m, 2H), 7.13-7.07 (m, 2H), 5.06 (m, 1H), 2.53 (s, 1H); ESIMS *m/z* 194 ($[M]^+$). The title compound (**DI42**) was prepared and carried on as crude intermediate.


1-(1-Bromo-2,2,2-trifluoroethyl)-4-methylbenzene (DI44)



DI43 **DI44**

2,2,2-Trifluoro-1-(*p*-tolyl)ethanol (**DI43**) was isolated as colorless oil (5.0 g, 99%): ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.37 (d, *J* = 8.0 Hz, 2H), 7.23 (d, *J* = 8.0 Hz, 2H), 5.02 (m, 1H), 2.46 (m, 1H), 2.37 (s, 3H); ESIMS *m/z* 190 ([M]$^+$). The title compound (**DI44**) was isolated (3.0 g, 45%).


1-(1-Bromo-2,2,2-trifluoroethyl)-3-fluorobenzene (DI46)



DI45 **DI46**

2,2,2-Trifluoro-1-(3-fluorophenyl)ethanol (**DI45**) was isolated as a colorless viscous oil (2.8 g, 93%): ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.41 (m, 1H), 7.25 (m, 2H), 7.14 (m, 1H), 5.06 (m, 1H), 2.60 (s, 1H); ESIMS *m/z* 194 ([M]$^+$). The title compound (**DI46**) was isolated (2.0 g, 61%).


1-(1-Bromo-2,2,2-trifluoroethyl)-2-fluorobenzene (DI48)



DI47 **DI48**

2,2,2-Trifluoro-1-(2-fluorophenyl)ethanol (**DI47**) was isolated as a colorless oil (2.5 g, 99%): ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.40 (m, 1H), 7.43 (m,1H), 7.24 (m, 1H), 7.13 (m, 1H), 5.42 (m, 1H), 2.65 (s, 1H); ESIMS *m/z* 194 ([M]$^+$). The title compound (**DI48**) was isolated (2.0 g, 61%): ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.61 (m, 1H), 7.40 (m, 1H), 7.23 (m, 1H), 7.10 (m, 1H), 5.40 (m, 1H); GCMS *m/z* 255 ([M-H]$^-$).

Example 91: Preparation of 4-(1*H*-1,2,4-triazol-1-yl)benzaldehyde (DI5)




To a stirring solution of 4-fluorobenzaldehyde (10.0 g, 80.6 mmol) in DMF (150 mL) were added K_2CO_3 (13.3 g, 96.7 mmol) and 1,2,4- triazole (6.67 g, 96.7 mmol) and the resultant reaction mixture was stirred at 120 °C for 6 h. After completion of reaction (by TLC), the reaction mixture was diluted with water and extracted with EtOAc (3 x100 mL). The combined EtOAc layer was washed with water and brine, dried over Na_2SO_4, and

concentrated under reduced pressure to afford the title compound as a solid (9.0 g, 65%): mp 145-149 °C: ^{1}H NMR (400 MHz, $CDCl_3$) δ 10.08 (s, 1H), 8.70 (s, 1H), 8.16 (s, 1H), 8.06 (d, *J* = 8.0 Hz, 2H), 7.92 (d, *J* = 8.0 Hz, 2H); ESIMS *m/z* 173.9 ([M+H]$^+$).


The following compound was made in accordance with the procedures disclosed in **Example 91**.

5-Formyl-2-(1*H*-1,2,4-triazol-1-yl)benzonitrile (DI49)




The title compound was isolated (2.8 g, 60%); ^{1}H NMR (400 MHz, $CDCl_3$) δ 10.10 (s, 1H), 8.98 (s, 1H), 8.35 (s, 1H), 8.30 (d, 1H), 8.22 (s, 1H), 8.07 (d, 1H); IR (thin film) 3433, 3120, 1702, 1599, 1510 cm^{-1}.

2-Chloro-4-(1*H*-1,2,4-triazol-1-yl)benzaldehyde (DI50)




The title compound was isolated as an off white solid (3.0 g, 40%): mp 149-151 °C; ^{1}H NMR (400 MHz, $CDCl_3$) δ 10.05 (s, 1H), 8.74 (s, 1H), 8.17 (s, 1H), 8.10 (s, 1H), 7.90 (m, 2H) ; ESIMS *m/z* 208.10 ([M+H]$^+$).

5-Methyl-4-(1*H*-1,2,4-triazol-1-yl)benzaldehyde (DI51)




The title compound was isolated as a white solid (0.5 g, 74 %): mp 109-111 °C; ^{1}H NMR (400 MHz, D_6-DMSO) δ 10.06 (s, 1H), 9.00 (s, 1H), 8.30 (s, 1H), 7.99 (s, 1H), 7.92 (d, *J* = 9.2 Hz, 1H), 7.69 (d, *J* = 9.2 Hz, 1H), 2.30 (s, 3H); ESIMS *m/z* 188.13 ([M+H]$^+$).

Example 92: Preparation of 5-Formyl-2-(3-nitro-1*H*-1,2,4-triazol-1-yl)benzonitrile (DI52)



To a stirring solution of 2-fluoro-5-formylbenzonitrile (0.5 g, 3.3 mmol) in DMF (25 mL) were added K_2CO_3 (0.68 g, 4.95 mmol) and 3-nitro-1,2,4 triazole (0.45 g, 4.2 mmol) and the resultant reaction mixture was stirred at ambient temperature for 14 h. After completion of reaction (TLC), the reaction mixture was diluted with water and extracted with EtOAc. The combined EtOAc layer was washed with water and brine then dried over Na_2SO_4 and concentrated under reduced pressure to afforded the title compound as a pale yellow solid (0.36 g, 45%): mp 170-172 °C; ^{1}H NMR (300 MHz, DMSO-d_6) δ 10.12 (s, 1H), 9.61 (s, 1H), 8.69 (s, 1H), 8.45 (d, J = 9.3 Hz, 1H), 8.23 (d, J = 9.3 Hz, 1H); ESIMS m/z 242.3 ([M-H]$^-$); IR (thin film) 2238, 1705, 1551, 1314 cm^{-1}.


Example 93: Preparation of 4-(3-Methyl-1*H*-1,2,4-triazol-1-yl)benzaldehyde (DI53)



To a stirring solution of 4-fluorobenzaldehyde (5.0 g, 40.32 mmol) in DMF (50 mL), were added K_2CO_3 (3.34 g, 40.32 mmol) and 3-methyl-1,2,4-trizole (3.34 g, 40.32 mmol) and the resultant reaction mixture was stirred at ambient temperature for 4 h. After completion of the reaction (TLC), the reaction mixture was diluted with water and extracted with EtOAc (3x). The combined EtOAc layer was washed with water and brine then dried over Na_2SO_4 and concentrated under reduced pressure to afforded the title compound as a white solid (4.1 g, 60%): mp 125-128°C; ^{1}H NMR (400 MHz, $CDCl_3$) δ 10.05 (s, 1H), 8.76 (s, 1H), 8.02 (d, 2H), 7.85 (d, 2H), 2.50 (s, 3H); ESIMS m/z 188.04 ([M+H]$^+$).


The following compound was made in accordance with the procedures disclosed in **Example 93**.

4-(1*H*-1,2,4-triazol-1-yl)-3-(trifluoromethyl)benzaldehyde (DI54)




The title compound was isolated as white solid (1.05 g, 60%): mp 81-83 °C; ^{1}H NMR (400 MHz, $CDCl_3$) δ 10.15 (s, 1H), 8.43 (s, 1H), 8.37 (s, 1H), 8.25 (d, *J* = 7.2 Hz, 1H), 8.18 (s, 1H), 7.79 (d, *J* = 7.2 Hz, 1H); ESIMS *m/z* 241.0 ([M]$^+$).

4-(3-Nitro-1*H*-1,2,4-triazol-1-yl)benzaldehyde (DI55)




The title compound was isolated as pale yellow solid (0.10 g, 23%): mp 159-161 °C; ^{1}H NMR (400 MHz, $CDCl_3$) δ 10.10 (s, 1H), 8.89 (s, 1H), 8.15 (m, 2H), 8.00 (m, 2H); ESIMS *m/z* 217.11 ([M-H]$^-$).

3-Bromo-4-(1*H*-1,2,4-triazol-1-yl)benzaldehyde (DI56)




The title compound was isolated as white solid (3.2 g, 51%): mp 126-128 °C; ^{1}H NMR (400 MHz, $CDCl_3$) δ 10.04 (s, 1H), 8.69 (s, 1H), 8.27 (M, 1H, 8.18 (s, 1H) 7.99 (d, *J* = 9.2 Hz, 1H), 7.76 (d, *J* = 9.2 Hz, 1H); ESIMS *m/z* 250.9 ([M]$^+$).

5-Formyl-2-(3-methyl-1H-1,2,4-triazol-1-yl)benzonitrile (DI57)




The title compound was isolated as white solid (0.13 g, 30%): mp 147-149 °C; ^{1}H NMR (400 MHz, $CDCl_3$) δ 10.07 (s, 1H), 8.89 (s, 1H), 8.32 (d, *J* = 1.8 Hz, 1H), 8.24 (dd, *J* = 8.6, 1.3 Hz, 1H), 8.06 (d, *J* = 8.6 Hz, 1H), 2.54 (s, 3H); ESIMS *m/z* 213.09 ([M+H]$^+$); IR (thin film) 2239, 1697 cm^{-1}.

3-Nitro-4-(1*H*-1,2,4-triazol-1-yl)benzaldehyde (DI58)



The title compound was isolated as pale yellow solid (3.0 g, 60 %): mp 116-118 °C; ^{1}H NMR (400 MHz, $CDCl_3$) δ 10.15 (s, 1H), 8.48 (s, 1H), 8.46 (s, 1H), 8.26 (d, *J* = 6.9 Hz, 1H), 8.16 (s, 1H), 7.83 (d, *J* = 6.9 Hz, 1H); ESIMS *m/z* 219.00 ([M+H]$^+$).


Example 94: Preparation of 1-(4-Vinylphenyl)-1*H*-1,2,4-triazole (DI59)



To a stirred solution of 4-[1,2,4]triazol-1-yl-benzaldehyde (9.0 g, 52 mmol) in 1,4-dioxane (100 mL), were added K_2CO_3 (10.76 g, 78 mmol) and methyl triphenyl phosphonium bromide (22.2 g, 62.4 mmol) at room temperature. The resultant reaction mixture was heated to 70 °C for 18 h. After completion of the reaction (TLC), the reaction mixture was cooled to room temperature and filtered and the obtained filtrate was concentrated under reduced pressure. Purification by flash chromatography (SiO_2, 100-200 mesh; 25-30% EtOAc in petroleum ether) to afforded the title compound as a white solid (5.6 g, 63%): ESIMS *m/z* 172.09 ([M+H]$^+$).


The following compound was made in accordance with the procedures disclosed in **Example 94**.

1-(2-Methyl-4-vinylphenyl)-1*H*-1,2,4-triazole (DI60)




The title compound was isolated as an off white solid (1.5 g, 76%): ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.25 (s, 1H), 8.11 (s, 1H), 7.35 (m, 2H), 7.27 (d, *J* = 8.7 Hz, 1H), 6.74 (m, 1H), 5.82 (d, *J* = 17.3 Hz, 1H), 5.36 (d, *J* = 10.0 Hz, 1H), 2.25 (s, 3H); ESIMS *m/z* 186.14 ([M+H]$^+$).

2-(1*H*-1,2,4-Triazol-1-yl)-5-vinylbenzonitrile (DI61)



The title compound was isolated as an off-white solid (1.40 g, 71%): mp 126-129 °C; ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.76 (s, 1H), 8.18 (s, 1H), 7.82-7.84 (m, 1H), 7.72-7.80 (m, 2H), 6.70-6.80 (dd, *J* = 17.6, 10.8 Hz, 1H), 5.90-5.95 (d, *J* = 17.6 Hz, 1H), 5.50-5.70 (d, *J* = 10.8 Hz, 1H); ESIMS *m/z* 197.03 ([M+H]$^+$).


Example 95: Preparation of 2-(3-Nitro-1*H*-1,2,4-triazol-1-yl)-5-vinylbenzonitrile (DI62)



To a stirred solution of 5-formyl-2-(3-nitro-1*H*-1,2,4-triazol-1-yl)benzonitrile (0.36 g, 1.49 mmol) in 1,4-dioxane (25 mL), were added K_2CO_3 (0.3 g, 2.2 mmol) and methyl triphenyl phosphonium bromide (0.63 g, 1.79 mmol). The resultant reaction mixture was heated to 100 °C for 18 h. After completion of the reaction (TLC), the reaction mixture was cooled to room temperature and filtered and the obtained filtrate was concentrated under reduced pressure. Purification by flash chromatography (SiO_2, 100-200 mesh; 25-30% EtOAc in petroleum ether) to afford the title compound as a solid (0.25 g, 70%): mp 103-105 °C; ^{1}H NMR (400 MHz, DMSO-d_6) δ 9.50 (s, 1H), 8.34 (m, 1H), 7.98 (d, J = 7.8 Hz, 1H), 7.68 (d, J = 7.8 Hz, 1H), 6.87 (m, 1H), 6.20 (d, J = 15.7 Hz, 1H), 5.56 (d, J = 11.8 Hz, 1H); ESIMS m/z 240.27 ([M-H]$^-$); IR (thin film) 2240, 1514, 1312 cm^{-1}.


The following compound was made in accordance with the procedures disclosed in **Example 95**.

1-(3-Chloro-4-vinylphenyl)-1*H*-1,2,4-triazole (DI63)




The title compound was isolated as an off-white solid (2.3 g, 80%): mp 134-137 °C; ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.56 (s, 1H), 8.11 (s, 1H), 7.76 (s, 1H), 7.70 (d, J = 9.0 Hz, 1H), 7.57 (d, J = 9.0 Hz, 1H), 7.10 (m, 1H), 5.80 (d, J = 17.2 Hz, 1H), 5.47 (d, J = 12.4 Hz, 1H); ESIMS m/z 206.04 ([M+H]$^+$.

3-Methyl-1-(4-vinylphenyl)-1H-1,2,4-triazole (DI64)




The title compound was isolated as a white solid (0.6 g, 60%): mp 109-111 °C; ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.42 (s, 1H), 7.40-7.60 (m, 4H), 6.70-7.00 (dd, J = 17.6, 10.8 Hz, 1H), 5.80 (d, J = 17.6 Hz, 1H), 5.30 (d, J = 17.6 Hz,1H), 2.50 (s, 3H); ESIMS m/z 186.20 ([M+H]$^+$).

1-(2-(Trifluoromethyl)-4-vinylphenyl)-1H-1,2,4-triazole (DI65)




The title compound was isolated as a colorless oil (0.6 g, 60%): ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.32 (s, 1H), 8.14 (s, 1H), 7.84 (s, 1H), 7.72 (d, *J* = 8.0 Hz, 1H), 7.50 (d, *J* = 7.6 Hz, 1H), 6.70-6.90 (dd, *J* = 17.6, 10.8 Hz, 1H), 5.90-6.00 (d, *J* = 17.6 Hz, 1H), 5.50-5.80 (d, *J* = 10.8 Hz 1H); ESIMS *m/z* 240.16 ($[M+H]^+$).

3-Nitro-1-(4-vinylphenyl)-1*H*-1,2,4-triazole (DI66)




The title compound was isolated as a pale yellow solid (61 mg, 20%): mp 137-139 °C; ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.60 (s, 1H), 7.68 (d, *J* = 7.7 Hz, 2H),7.60 (d, *J* = 8.3 Hz, 2H), 6.77 (dd, *J* = 17.7, 10.8, 1H), 5.87 (d, *J* = 17.7 Hz, 1H), 5.42 (d, *J* = 10.8 Hz, 1H); ESIMS *m/z* 217.28 ($[M+H]^+$).

1-(2-Bromo-4-vinylphenyl)-1*H*-1,2,4-triazole (DI67)




The title compound was isolated as a white solid (1.2 g, 40%): mp 75-77 °C; ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.48 (s, 1H), 8.12 (s, 1H), 7.75 (s, 1H) 7.42 (s, 2H), 6.70 (m, 1H), 5.83 (d, *J* = 18 Hz, 1H), 5.42 (d, *J* = 12 Hz, 1H); ESIMS *m/z* 249.1 ($[M]^+$).

2-(3-Methyl-1*H*-1,2,4-triazol-1-yl)-5-vinylbenzonitrile (DI68)




The title compound was isolated as an off-white solid (0.6 g, 60%): mp 96-97 °C; ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.66 (s, 1H), 7.80 (s, 1H), 7.74 (m, 2H), 6.73 (dd, *J* = 17.6 Hz, 10.8 Hz, 1H), 5.88 (d, *J* = 17.6 Hz, 1H), 5.49 (d, *J* = 10.8 Hz, 1H), 2.52 (s, 3H); ESIMS *m/z* 211.10 ($[M+H]^+$); IR (thin film) 2229 cm^{-1}.

1-(2-Nitro-4-vinylphenyl)-1*H*-1,2,4-triazole (DI69)



The title compound was isolated as a yellow solid (1.78 g, 60%): mp 102-104 °C; ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.40 (s, 1H), 8.12 (s, 1H), 8.02 (s, 1H), 7.72-7.76 (d, *J* = 8.0 Hz, 1H), 7.52-7.56 (d, *J* = 17.6 Hz, 1H), 6.70-6.82 (dd, *J* = 17.6, 10.8 Hz, 1H), 5.85-6.00 (d, *J* = 17.6 Hz, 1H), 5.50-5.60 (d, *J* = 10.8, Hz 1H); ESIMS *m/z* 217.0 ([M+H]$^+$).

Example 96: Preparation of 3-Methyl-2-(1*H*-1,2,4-triazol-1-yl)-5-vinylbenzonitrile (DI70)



Step 1. 5-Bromo-2-fluoro-3-methylbenzaldehyde: To a stirred solution of di-isopropyl amine (4.01 g, 39.88 mmol) in THF (20 mL) was added *n*-BuLi (1.6 M in hexane) (19.9 mL, 31.91 mmol) at -78 °C slowly dropwise over the period of 10 min, the reaction mixture was stirred at -78°C for 30 min. A solution of 4-bromo-1-fluoro-2-methylbenzene (5.0 g, 26.6 mmol) in THF (30.0 mL) was added at -78°C, and the reaction mixture was stirred for 1h at the same temperature. DMF (5.0 mL) was added and stirred at -78°C for another 30 min. The reaction was monitored by TLC; then the reaction mixture was quenched with 1N HCl solution (aq) at 0°C. The aqueous layer was extracted with Et_2O, washed with water and saturated brine solution. The combined organic layer was dried over anhydrous Na_2SO_4 and concentrated under reduced pressure to obtain the crude compound purified by flash column chromatography (SiO_2, 100-200 mesh; eluting with 5% EtOAc/ pet ether) to afford the title compound as a white solid (3.6 g, 64 %); mp 48-50°C: ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.33 (s, 1H), 8.22 (s, 1H), 7.67 (s, 1H), 7.60 (s, 1H), 6.75 (dd, *J* = 17.6, 10.8 Hz, 1H), 5.92 (dd, *J* = 17.6, 10.8 Hz, 1H), 5.52 (d, *J* = 17.6 Hz, 1H), 2.21 (s, 3H); ESIMS *m/z* 211.35 ([M-H]$^-$).

Step 2. ((*E*)-5-Bromo-2-fluoro-3-methylbenzaldehyde oxime: To a stirred solution of 5-bromo-2-fluoro-3-methylbenzaldehyde (3.5 g, 16.2 mmol) in ethanol (50.0 mL) were added NaOAc (2.0 g, 24.3 mmol) and hydroxylamine hydrochloride (1.69 g, 24.3 mmol) at ambient temperature. The reaction mixture was stirred at ambient temperature for 3 h. The reaction mixture was concentrated on rotavapour to obtain crude compound, which was

washed with water filtered and dried under vacuum to afford the title compound as a white solid: mp 126-127 °C; ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.32 (s, 1H), 7.73 (d, *J* = 2.4 Hz, 1H), 7.51 (s, 1H), 7.34 (d, *J* = 2.4 Hz, 1H), 2.25 (s, 3H); ESIMS *m/z* 232.10 ($[M+H]^+$).


Step 3. 5-Bromo-2-fluoro-3-methylbenzonitrile: A stirred solution of (*E*)-5-bromo-2-fluoro-3-methylbenzaldehyde oxime (0.5 g, 2.2 mmol) in acetic anhydride (5.0 mL) was heated to reflux for 18 h. The reaction mixture was diluted with water and extracted with EtOAc. The combined EtOAc layer was washed with brine and dried over Na_2SO_4 and concentrated under reduced pressure to afford the crude compound as a light brown gummy material (0.4 g, crude): ESIMS *m/z* 213.82 ($[M+H]^+$).

Step 4. 5-Bromo-3-methyl-2-(1H-1,2,4-triazol-1-yl)benzonitrile (DI71) : To a stirred solution of 5-bromo-2-fluoro-3-methylbenzonitrile (1.0 g, 47.716 mmol), in DMF (10.0 mL) was added K_2CO_3 (1.95 g, 14.14 mmol) followed by 1*H*-1,2,4-triazole (0.811 g, 9.433 mmol) at ambient temperature. The reaction mixture was heated to 140 °C for 18 h. The reaction mixture was cooled to ambient temperature, diluted with water and extracted with EtOAc (2 x 100 mL). The combined EtOAc layer was washed with brine and dried over Na_2SO_4 and concentrated under reduced pressure to afford the crude compound purified by flash column chromatography (SiO_2, 100-200 mesh; eluting with 30% EtOAc/ pet ether) to afford the title compound as a pink solid (0.6 g, 49 %): ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.39 (s, 1H), 8.23 (s, 1H), 7.91 (d, *J* = 2.4 Hz, 2H), 2.21 (s, 3H), ESIMS *m/z* 262.57 ($[M+H]^+$); IR (thin film) 2231, 554 cm^{-1}.

Step 5. 3-Methyl-2-(1H-1,2,4-triazol-1-yl)-5-vinylbenzonitrile (DI70) : A mixture of 5-bromo-3-methyl-2-(1H-1,2,4-triazol-1-yl)benzonitrile (0.6 g, 2.3 mmol), K_2CO_3 (0.95 g, 6.87 mmol), vinyl boronic anhydride (0.82 g, 3.43 mmol) and triphenylphosphine (0.13 g, 0.114 mmol) in toluene (20.0 mL) were stirred and degassed with argon for 30 min. The reaction mixture was heated to reflux for 18 h. The reaction mixture was cooled to ambient temperature, diluted with water and extracted with EtOAc (2 x 100 mL). The combined EtOAc layer was washed with brine, dried over Na_2SO_4 and concentrated under reduced pressure to afford the crude compound that was purified by flash column chromatography (SiO_2, 100-200 mesh; eluting with 30% EtOAc/ pet ether) to afford the title compound as a pink solid (0.25 g, 52 %): ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.33 (s, 1H), 8.22 (s, 1H), 7.67 (s, 1H), 7.60 (s, 1H), 6.75 (dd, *J* = 17.6, 10.8 Hz, 1H), 5.92 (d, *J* = 17.6, 1H), 5.52 (d, *J* = 10.8 Hz, 1H), 2.21 (s, 3H), ESIMS *m/z* 211.35 ($[M+H]^+$); IR (thin film) 2236, 1511 cm^{-1}.


The following compound was made in accordance with the procedures disclosed in Steps 4 and 5 of **Example 96**.

1-(2-Fluoro-4-vinylphenyl)-1H-1,2,4-triazole (DI72)




1-(4-Bromo-2-fluorophenyl)-1H-1,2,4-triazole (**DI73**) was isolated as a pale yellow solid (3.0 g, 75%): mp 113-116 °C; ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.69 (s, 1H), 8.13 (m, 2H), 7.50 (m, 1H), 7.21 (m, 1H); ESIMS *m/z* 241.93 ([M]$^+$). The title compound (**DI72**) was isolated as a yellow solid (1.0 g, 71%): mp 67-70 °C; ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.67 (s, 1H), 8.13 (s, 1H), 7.94 (m, 1H), 7.41 (m, 1H), 7.24 (s, 1H), 6.75 (dd, *J* = 17.6, 10.8 Hz, 1H), 5.81 (d, *J* = 17.6 Hz, 1H), 5.37 (d, *J* = 10.8 Hz, 1H); ESIMS *m/z* 190.00 ([M+H]$^+$).

Example 119: Preparation of 1-(1-(4-Vinylphenyl)-1*H*-1,2,4-triazol-5-yl)ethanone (DI78)



To a stirred solution of 1-(4-vinyl-phenyl)-1*H*-[1,2,4]triazole (1 g, 5.8 mmol) in 25 mL of THF, was added *n*-BuLi (0.37 g, 5.8 mmol) at -78 °C and stirred for 30 min. To this *N*-methoxy-*N*-methyl acetamide in THF (0.66 g, 6.4 mmol) was added and the resultant reaction mixture was stirred at ambient temperature for 16 h. The reaction mixture was quenched with a saturated aqueous NH_4Cl solution and extracted with EtOAc (3 x50 mL). The combined EtOAc layer was washed with brine and dried over sodium sulphate and concentrated under reduced pressure. The crude compound was purified by flash chromatography (SiO_2, 100-200 mesh, 40% EtOAc in Pet ether) to afford the title compound as an off white solid (280 mg, 23%): mp 97-98 °C; ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.10 (s, 1H), 7.50 (d, 2H), 7.38 (d, 2H), 6.68 (dd, 1H), 5.85 (d, 1H), 5.38 (d, 1H), 2.75 (s, 3H); ESIMS *m/z* 214.14 ([M+H]$^+$).


Example 120: Preparation of Cyclopropyl(1-(4-vinylphenyl)-1*H*-1,2,4-triazol-5-yl)methanone (DI79)



To a stirred solution of 1-(4-vinyl-phenyl)-1*H*-[1,2,4]triazole (1 g, 5.8 mmol) in 25 mL of THF, was added *n*-BuLi (0.37 g, 5.8 mmol) at -78 °C and stirred for 30 min. To this *N*-methoxy *N*-methylcyclopropoxide in THF (0.82 g, 6.4 mmol) was added and the resultant reaction mixture was stirred at ambient temperature for 16 h. The reaction mixture was


quenched with a saturated aqueous NH_4Cl solution and extracted with EtOAc (3 x25 mL). The combined EtOAc layer was washed with brine and dried over sodium sulphate and concentrated under reduced pressure. The crude compound was purified by flash chromatography (SiO_2, 100-200 mesh, 40% EtOAc in Pet ether) to afford the title compound as an off white solid (420 mg, 30%): mp 90-91 °C; ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.12 (s, 1H), 7.50 (d, *J* = 7.8 Hz, 2H), 7.38 (d, *J* = 7.8 Hz, 2H), 6.75 (dd, *J* = 16.3, 10.7 Hz, 1H), 5.81 (d, *J* = 16.3 Hz, 1H), 5.35 (d, *J* = 10.7 Hz, 1H), 3.22 (m, 1H), 1.27(m, 2H), 1.18 (m, 2H); ESIMS *m/z* 240.18 ([M+H]$^+$); IR (thin film) 2922, 1630 cm^{-1}.

Example 121: Preparation of 5-(Methylthio)-1-(4-vinylphenyl)-1*H*-1,2,4-triazole (DI80)



To a stirred solution of 1-(4-vinyl-phenyl)-1*H*-[1,2,4]triazole (1 g, 5.8 mmol) in 50 mL of THF, was added *n*-BuLi (0.41 g, 6.4 mmol) at -78 °C and stirred for 30 min. To this dimethyldisulfide in THF (0.6 g, 6.43 mmol) was added and the resultant reaction mixture was stirred at ambient temperature for 16 h. The reaction mixture was quenched with a saturated aqueous NH_4Cl solution and extracted with EtOAc (3 x 25 mL). The combined EtOAc layer was washed with brine and dried over sodium sulphate and concentrated under reduced pressure. The crude compound was purified by flash chromatography (SiO_2, 100-200 mesh, 40% EtOAc in Pet ether) to afford the title compound as an off white solid (0.6 g, 48%): mp 68-70 °C; ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.96 (s, 1H), 7.05 (m, 4H), 6.75 (dd, *J* = 16.4, 10.7 Hz, 1H), 5.81 (d, *J* = 16.4 Hz, 1H), 5.35 (d, *J* = 10.7 Hz, 1H), 2.73 (s, 3H); ESIMS *m/z* 218.09 ([M+H]$^+$).


Example 122: Preparation of 5-Methyl-1-(4-vinylphenyl)-1*H*-1,2,4-triazole (DI81)



To a stirred solution of 1-(4-vinyl-phenyl)-1*H*-[1,2,4]triazole (0.5 g, 2.9 mmol) in 10 mL of THF, was added *n*-BuLi (0.22 g, 3.5 mmol) at -78 °C and stirred for 30 min. To this methyl iodide in THF (0.50 g, 3.5 mmol) was added and the resultant reaction mixture was stirred at ambient temperature for 16 h. The reaction mixture was quenched with a saturated aqueous NH_4Cl solution and extracted with EtOAc (3 x 25 mL). The combined EtOAc layer was washed with brine and dried over sodium sulphate and concentrated under reduced pressure The crude compound was purified by flash chromatography (SiO_2, 100-200 mesh,

40% EtOAc in Pet ether) afford the title compound as a pale brown liquid (250 mg, 46%): ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.93 (s, 1H), 7.55 (d, *J* = 9 Hz, 2H), 7.42 (d, *J* = 9 Hz, 2H), 6.76 (dd, *J* = 18, 11 Hz, 1H), 5.83 (d, *J* = 18 Hz, 1H), 5.38 (d, *J* = 11 Hz, 1H), 2.55 (s, 3H); ESIMS *m/z* 186.13 ($[M+H]^+$); IR (thin film) 1517, 1386, 1182, 847 cm^{-1}.


Example 97: Preparation of (E)-1-(4-(3-(3,5-Dichlorophenyl)-4,4,4-trifluorobut-1-en-1-yl)phenyl)-1*H*-1,2,4-triazole (DC1)



To a stirred solution of 1-(1-bromo-2,2,2-trifluoro-ethyl)-3,5-dichloro-benzene (2.0 g, 6.51 mmol) in 1,2-dichlorobenzene (25 mL), were added 1-(4-vinyl-phenyl)-1*H*-[1,2,4]triazole (2.22 g, 13.0 mmol), CuCl (64 mg, 0.65 mmol) and 2,2 -bipyridyl (0.2 g, 1.3 mmol). The resultant reaction mixture was degassed with argon for 30 min, then stirred at 180 °C for 24 h. After completion of reaction (TLC), the reaction mixture was cooled to ambient temperature and filtered and the filtrate concentrated under reduced pressure. Purification by flash chromatography (SiO_2, 100-200 mesh; 25-30% EtOAc in petroleum ether) afforded the title compound as an off-white solid (0.8 g, 32%): mp 93–97 °C; ^{1}H NMR (300 MHz, $CDCl_3$) δ 8.56 (s, 1H), 8.11 (s, 1H), 7.68 (d, *J* = 8.4 Hz, 2H), 7.54 (d, *J* = 8.4 Hz, 2H), 7.38 (t, *J* = 1.8 Hz, 1H), 7.29 (s, 2H), 6.62 (d, *J* = 15.6 Hz, 1H), 6.42 (dd, *J* = 15.6, 8.2 Hz, 1H), 4.15 (m, 1H); ESIMS *m/z* 398.05 ($[M+H]^+$).

Compounds **DC2-DC37, DC44, DC45, DC47-49, DC50, DC51, DC54, DC58, DC60, DC62,** and **DC63-DC67** in Table 1 were made in accordance with the procedures disclosed in **Example 97**.


Example 98: Preparation of (E)-2-(3-Nitro-1*H*-1,2,4-triazol-1-yl)-5-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-en-1-yl)benzonitrile (DC40)



To a stirred solution of 2-(3-nitro-1*H*-1,2,4-triazol-1-yl)-5-vinylbenzonitrile (0.9 g, 3.7 mmol) in 1,2- dichlorobenzene (10 mL), were added 5-(1-bromo-2,2,2-trifluoroethyl)-1,2,3-trichlorobenzene (2.5 g, 7.5 mmol), CuCl (73 mg, 0.74 mmol) and 2,2-bipyridyl (0.23 g, 1.49 mmol) and the resultant reaction mixture was degassed with argon for 30 min and

then stirred at 180°C for 14 h. After completion of the reaction (TLC), the reaction mixture was cooled to ambient temperature and filtered and the filtrate was concentrated under reduced pressure. Purification by flash chromatography (SiO_2, 100-200 mesh, 25-30% EtOAc in Pet ether) afforded the title compound as an off white solid (0.9 g, 50%): mp 70–73 °C; ^{1}H NMR (300 MHz, $CDCl_3$) δ 8.86 (s, 1H), 7.88 (m, 3H), 7.44 (s, 2H), 6.67 (d, J = 16.0 Hz, 1H), 6.56 (dd, J = 16.0, 7.6 Hz, 1H), 4.19 (m, 1H); ESIMS *m/z* 436.11 ([M-2H]$^-$).


Example 99: Preparation of (E)-2-(3-Amino-1*H*-1,2,4-triazol-1-yl)-5-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-en-1-yl)benzonitrile (DC41)



To a stirred solution of (E)-2-(3-nitro-1H-1,2,4-triazol-1-yl)-5-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-enyl)benzonitrile (0.6 g, 1.2 mmol) in MeOH (10 mL), were added Zn dust (0.39g, 5.98 mmol) and saturated aqueous NH_4Cl solution (5 mL) and the resultant reaction mixture was stirred at ambient temperature for 2 h. After completion of the reaction (TLC), the reaction mass was concentrated under reduced pressure. The reaction mass was diluted with CH_2Cl_2, filtered through a Celite® bed, and the obtained filtrate concentrated under reduced pressure to afford the title compound as a solid (0.5 g, 89%): mp 72-75 °C; ^{1}H NMR (300 MHz, DMSO-d_6) δ 8.72 (s, 1H), 8.26 (s, 1H), 8.01 (d, J = 8.4 Hz, 1H), 7.91 (s, 2H), 7.77 (d, J = 8.4 Hz, 1H), 6.42 (dd, J = 15.6, 9.2 Hz, 1H), 6.83 (d, J = 15.6 Hz, 1H), 5.87 (s, 2H), 4.89 (m, 1H); ESIMS *m/z* 469.95 ([M-H]$^-$).


Compound **DC38** in Table 1 was made in accordance with the procedures disclosed in **Example 99**. Also, compound **DC55** in Table 1 was made from compound **DC54** in accordance with the procedures disclosed in **Example 99**, with the exception of using ammonium formate in place of NH_4Cl.

Example 100: Preparation of (*E*)-*N*-(1-(2-Cyano-4-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-en-1-yl)phenyl)-1H-1,2,4-triazol-3-yl)-N-(cyclopropanecarbonyl)cyclopropanecarboxamide (DC42)



To a stirred solution of (*E*)-2-(3-amino-1*H*-1,2,4-triazol-1-yl)-5-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-enyl)benzonitrile (0.1 g, 0.21 mmol) in CH_2Cl_2 at ambient temperature, was added cyclopropylcarbonyl chloride (0.045 g, 0.42 mmol) and the reaction mixture was stirred for 2 h at ambient temperature. The reaction mixture was diluted with CH_2Cl_2 and washed with water and brine and dried over Na_2SO_4. Concentration under reduced pressure and purification by preparative HPLC afforded the title compound as a solid (0.09g, 79%): mp 104-107 °C; ^{1}H NMR (300 MHz, $CDCl_3$) δ 8.78 (s, 2H), 7.83 (s, 1H), 7.80 (m, 2H), 7.42 (s, 2H) , 6.65 (d, *J* = 16.4 Hz, 1H), 6.51 (dd, *J* = 7.6, 8.0 Hz, 1H), 4.17 (m, 1H), 2.16 (m, 2H), 1.25 (m, 4H), 1.00 (m, 4H); ESIMS *m/z* 609.98 ($[M+H]^+$); IR (thin film) 2234, 1714, 1114, 807 cm^{-1}.


Example 101: Preparation of (E)-*N*-(1-(2-Cyano-4-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-en-1-yl)phenyl)-1*H*-1,2,4-triazol-3-yl)cyclopropanecarboxamide (DC43)



To a stirred solution of (E)-2-(3-amino-1H-1,2,4-triazol-1-yl)-5-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-enyl)benzonitrile (0.15 g,0.31 mmol) in CH_2Cl_2 at 0 °C, were added TEA (0.1 g, 1 mmol) and cyclopropylcarbonyl chloride (0.04 g, 0.38mmol) and the reaction mixture was stirred for 1 h at 0 °C. The reaction mixture was diluted with CH_2Cl_2 and washed with water and brine and dried over Na_2SO_4. Concentration under reduced pressure and purification by column chromatography (SiO_2, 100-200 mesh) afforded the title compound as a solid (66 mg, 34%): mp 109-112 °C; ^{1}H NMR (300 MHz, DMSO-d_6) δ 10.94 (br s, 1H), 8.36 (s, 1H), 8.08 (m, *J* = 8.4 Hz, 1H), 7.91 (s, 2H), 7.84 (d, *J* = 8.4 Hz, 1H), 7.13 (dd, *J* = 15.6, 9.2 Hz, 1H), 6.87 (d, *J* = 15.6 Hz, 1H), 4.92 (m, 1H), 1.99 (br s, 1H), 0.82 (s, 4H); ESIMS *m/z* 540.04 ($[M+H]^+$); IR (thin film) 3233, 2233, 1699, 1114, 807 cm^{-1}.


Compound **DC39** in Table 1 was made in accordance with the procedures disclosed in **Example 101**.

Example 102: Preparation of 1-(4-(1*H*-1,2,4-triazol-1-yl)phenyl)ethanone (DI74)



To a stirred solution of 4-bromoacetophenone (10 g, 50 mmol) in DMF (100 mL), were added 1,2,4-triazole (5 g, 75mmol), Cs_2CO_3 (32.6 g, 100.5 mmol) and CuI (1.4 g, 10.1 mmol) and the resultant reaction mixture was refluxed for 48 h. After completion of the reaction (by TLC), the reaction mixture was cooled to ambient temperature and diluted with water (200 mL) and extracted with EtOAc. The combined organic layer was washed with brine and dried over Na_2SO_4 and concentrated under reduced pressure. Purification by washing with Et_2O afforded the title compound as a solid (5 g, 96%): ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.71 (s, 1H), 8.16, (s, 1H), 8.13 (d, J = 8.6 Hz, 2H), 7.83 (d, J = 8.6 Hz, 2H), 2.66 (s, 3H); ESIMS *m/z* 186.02 ([M-H]$^-$).


Example 103: Preparation of 1-(4-(1*H*-1,2,4-triazol-1-yl)phenyl)-3-(3,5-dichlorophenyl)-4,4,4-trifluorobutan-1-one (DI75)



Step 1. 1-(4-(1-(Trimethylsilyloxy)vinyl)phenyl)-1H-1,2,4-triazole (DI76) To a stirred solution of 1-(4-(1*H*-1,2,4-triazol-1-yl)phenyl)ethanone (4.5 g, 24.0 mmol) in CH_2Cl_2 at 0 °C, were added TEA (3.7 g, 36.1 mmol) and trimethylsilyl triflluoromethanesulfonate (8 g, 36 mmol) and the resultant reaction mixture was stirred for 1 h. The reaction mixture was quenched with a mixture of sat aqueous $NaHCO_3$ solution and ether. The ether layer and was separated, washed with brine, dried over Na_2SO_4 and concentrated under reduced pressure to afford the title compound (5.5 g) which was taken directly to next step.


Step 2. 1-(4-(1*H*-1,2,4-Triazol-1-yl)phenyl)-3-(3,5-dichlorophenyl)-4,4,4-trifluorobutan-1-one (DI75): To a stirred solution of 1-(4-(1-(trimethylsilyloxy)vinyl)phenyl)-1*H*-1,2,4-triazole (6g, 23 mmol) and 1-(1-bromo-2,2,2-trifluoro-ethyl)-3,5-dichlorobenzene (7.1 g, 34.7 mmol) in 1,2-dichlorobenzene (30 mL) was degassed with argon. To this CuCl (0.23g, 2.31 mmol) and 2,2-bipyridyl (0.73g, 4.63 mmol) was added to the above reaction mixture and the resultant reaction mixture was heated to 180 °C for 18 h. After completion of the reaction (by TLC), the reaction mixture was absorbed onto silica gel and purified by column chromatography (SiO2; 10% EtOAc in petroleum ether) to afford title compound as a solid (3 g, 31%): ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.67 (s, 1H), 8.15 (s, 1H), 8.10 (d, J = 8.3 Hz, 2H), 7.82 (d, J = 8.3 Hz, 2H), 7.33 (m, 1H), 7.30 (m, 2H), 4.20 (m, 1H), 3.63 (m, 2H); ESIMS *m/z* 412. 14 ([M-H]$^-$).

Example 104: Preparation of 2-(4-(1*H*-1,2,4-triazol-1-yl)phenyl)-4-(3,5-dichlorophenyl)-5,5,5-trifluoropentan-2-ol (DI77)




To a solution of 1-(4-(1*H*-1,2,4-triazol-1-yl)phenyl)-3-(3,5-dichlorophenyl)-4,4,4-trifluorobutan-1-one (300 mg, 0.726 mmol) in THF cooled to 0 °C was added methylmagnesium bromide (450 mg, 5 mmol) drop wise. The reaction was stirred for 3 h at 0 °C, then the reaction mixture was quenched with sat aqueous NH_4Cl solution and extracted with EtOAc. The combined EtOAc layer was washed with water and brine, dried over Na_2SO_4 and concentrated under reduced pressure. Purification by column chromatography (SiO_2, 100-200 mesh; 20%-25% EtOAc in petroleum ether) afforded the title compound as a solid (100 mg, 32%): ^1H NMR (400 MHz, $CDCl_3$) δ two diastereoisomers 8.58 (s, 1H, minor), 8.48 (s, 1H, major), 8.13 (s, 1H, minor), 8.09 (s, 1H, major), 7.70 (d, J = 9.0 Hz, 2H, minor), 7.53 (d, J = 9.0 Hz, 2H, minor), 7.40 (d, J = 9.0 Hz, 2H, major), 7.31 (m, 1H, minor), 7.27 (d, J = 9.0 Hz, 2H, major), 7.20 (m, 2H, minor), 7.01 (m, 1H, major), 6.75 (m, 2H, major), 350 (m, 1H), 2.50 (m, 2H), 1.56 (s, 3H, major), 1.54 (s, 3H, minor); ESIMS *m/z* 430.05 ($[M+H]^+$).

Example 105: Preparation of (*E*)-1-(4-(4-(3,5-Dichlorophenyl)-5,5,5-trifluoropent-2-en-2-yl)phenyl)-1*H*-1,2,4-triazole (DC68)



To a solution of 2-(4-(1*H*-1,2,4-triazol-1-yl)phenyl)-4-(3,5-dichlorophenyl)-5,5,5-trifluoropentan-2-ol (100 mg, 0.233 mmol) in toluene was added a catalytic amount of *p*-toluenesulfonic acid and the water was removed by azeotropic distillation over the course of 12 h. The reaction mixture was cooled to room temperature and dissolved in EtOAc. The solution was washed with sat aqueous $NaHCO_3$ solution and brine, dried over Na_2SO_4 and concentrated under reduced pressure. Purification by column chromatography (SiO_2, 100-200 mesh; 20%-25% EtOAc in petroleum ether) afforded the title compound as a solid (30 mg, 31%).


Example 123: Preparation of (*E*)-5-(3-(3,5-Dichlorophenyl)-4,4,4-trifluorobut-1-en-1-yl)-2-(1H-1,2,4-triazol-1-yl)benzaldehyde (DC52)



To a stirred solution of (*E*)-5-(3-(3,5-dichlorophenyl)-4,4,4-trifluorobut-1-en-1-yl)-2-(1*H*-1,2,4-triazol-1-yl)benzonitrile (0.3 g, 0.71 mmol) in toluene (10 mL) at -78 °C was added dropwise diisobutylaluminum hydride (DIBAL-H, 1.0 M solution in toluene; 0.85 mL), and the reaction mixture was stirred at -78 °C for 20 min. The reaction mixture was quenched with the addition of 1 N HCl solution, then the aqueous layer was extracted with EtOAc (2x). The combined organic layers were washed with brine, dried over Na_2SO_4 and concentrated under reduced pressure. The crude compound was purified by flash column chromatography (SiO_2; 50% EtOAc/ Pet ether) to afford the title compound as a yellow oil.


Compound **DC53** in Table 1 was made in accordance with the procedures disclosed in **Example 123**.

Example 124: Preparation of (*E*)-5-(3-(3,5-Dichlorophenyl)-4,4,4-trifluorobut-1-en-1-yl)-*N*-methyl-2-(1*H*-1,2,4-triazol-1-yl)aniline (DC57)




To a stirred solution of (*E*)-5-(3-(3,5-dichlorophenyl)-4,4,4-trifluorobut-1-en-1-yl)-2-(1*H*-1,2,4-triazol-1-yl)aniline (0.3 g, 0.7 mmol) in CH_2Cl_2 (10 mL) was added TEA (0.155 mL, 1.09 mmol) and methyl iodide (0.124 g, 0.873 mmol). The reaction was stirred at ambient temperature for 18 h. The CH_2Cl_2 layer was washed with water and brine, dried over Na_2SO_4 and concentrated under reduced pressure. The crude compound was purified by flash column chromatography (SiO_2; 50% EtOAc/ Pet ether) to afford the title compound as a yellow semi-solid (0.07 g, 70%).

Example 125: Preparation of (*E*)-5-(3-(3,5-Dichlorophenyl)-4,4,4-trifluorobut-1-en-1-yl)-2-(1*H*-1,2,4-triazol-1-yl)benzoic acid (DC61)




A solution of (*E*)-ethyl 5-(3-(3,5-dichlorophenyl)-4,4,4-trifluorobut-1-en-1-yl)-2-(1*H*-1,2,4-triazol-1-yl)benzoate (0.2 g, 0.4 mmol) in 6 N HCl (10 mL) was stirred at 100 °C for 18 h. The reaction was cooled to ambient temperature, resulting in a white solid precipitate. The precipitate was filtered to afford the title compound as a white solid (0.12 g, 60%).

Example 126: Preparation of (*Z*)-5-((*E*)-3-(3,5-Dichlorophenyl)-4,4,4-trifluorobut-1-en-1-yl)-*N*'-hydroxy-2-(1H-1,2,4-triazol-1-yl)benzimidamide (DC59)



A solution of (*E*)-5-(3-(3,5-dichlorophenyl)-4,4,4-trifluorobut-1-en-1-yl)-2-(1*H*-1,2,4-triazol-1-yl)benzonitrile (0.3 g, 0.71 mmol), NaOAc (0.087 g, 1.065 mmol) and hydroxylammonium chloride (0.072 g, 1.065 mmol) in 9:1 ethanol/water mixture (10 mL) was stirred at 70 °C for 8 h. The reaction was cooled to ambient temperature, and the ethanol was evaporated. The residue was dissolved in water and extracted with EtOAc (2x). The combined organic layers were washed with brine, dried over Na_2SO_4 and concentrated under reduced pressure to afford the title compound as an off white solid.

Example 127: Preparation of (*E*)-1-(4-(3-(3,5-Dichlorophenyl)-4,4,4-trifluoro-3-methoxybut-1-en-1-yl)phenyl)-1*H*-1,2,4-triazole (DC70)




Step 1. (*E*)-3-(4-(1*H*-1,2,4-triazol-1-yl)phenyl)-1-(3,5-dichlorophenyl)prop-2-en-1-one: To a solution of 1-(3,5-dichlorophenyl)ethanone (0.5 g, 2.6 mmol) in ethanol (20 mL) was added 4-(1*H*-1,2,4-triazol-1-yl)benzaldehyde (0.46 g, 2.65 mmol) and the reaction was cooled to 0 °C. NaOH (0.22 g, 5.29 mmol) in water (10 mL) was then added and the reaction was allowed to stir for 2 h at 0 °C. The reaction was extracted with EtOAc and the combined organic layers were dried over Na_2SO_4 and concentrated under reduced pressure to afford the title compound (0.149 g, 17%):); ESIMS *m/z* 430.05 ($[M+H]^+$) 344.08

Step 2. (*E*)-4-(4-(1*H*-1,2,4-triazol-1-yl)phenyl)-2-(3,5-dichlorophenyl)-1,1,1-trifluorobut-3-en-2-ol (DC69): To a solution of (*E*)-3-(4-(1*H*-1,2,4-triazol-1-yl)phenyl)-1-(3,5-dichlorophenyl)prop-2-en-1-one (1 g, 3 mmol) in THF (150 mL) was added trifluoromethyltrimethylsilane (0.517 g, 3.644 mmol) and tetra-*n*-butylammonium fluoride (TBAF) (1.0 M, 1 mL) at 0 °C. The reaction was slowly warmed to ambient temperature and allowed to stir for 2 h. The reaction was then cooled to 0 °C and 5 M HCl solution was added and the reaction was stirred for an additional 4 h at ambient temperature. The reaction was extracted with CH_2Cl_2 and the combined organic layers were dried over Na_2SO_4 and concentrated under reduced pressure. The crude compound was purified by flash column chromatography (SiO_2; 25% EtOAc/ hexanes) to afford the title compound as an off-white solid (0.3 g, 25%).

Step 3. (*E*)-1-(4-(3-(3,5-Dichlorophenyl)-4,4,4-trifluoro-3-methoxybut-1-en-1-yl)phenyl)-1*H*-1,2,4-triazole (DC70): To a solution of (*E*)-4-(4-(1*H*-1,2,4-triazol-1-yl)phenyl)-2-(3,5-dichlorophenyl)-1,1,1-trifluorobut-3-en-2-ol (0.15 g, 0.36 mmol) in THF (5 mL) was added NaH (60%, 10 mg, 0.44 mmol) at 0 °C. The reaction was allowed to stir at 0 °C for 30 min, then methyl iodide (61 mg, 0.44 mmol) was added slowly and the reaction was warmed to ambient temperature and allowed to stir for 4 h. The reaction was quenched with aqueous NH_4Cl solution and extracted with CH_2Cl_2. The combined organic layers were dried over Na_2SO_4 and concentrated under reduced pressure to afford the title compound as an off-white solid (55 mg, 35%).


Example 128: Preparation of *tert*-Butyl (2-methyl-1-oxo-1-((2,2,2-trifluoroethyl)amino)propan-2-yl)carbamate



To a stirred solution of 2-((*tert*-butoxycarbonyl)amino)-2-methylpropanoic acid (4.58 g, 22.6 mmol) in methylene chloride (50 mL) was added EDC·HCl (4.75 g, 24.8 mmol) followed by 2,2,2-trifluoroethylamine (2.67 g, 27.0 mmol) and DMAP (3.03 g, 24.8 mmol). The reaction mixture was stirred at ambient temperature for 18 h, then washed with aqueous 5% $NaHSO_4$ (2x), aqueous 10 % HCl (1x) and aqueous saturated $NaHCO_3$ (2x). The organic phase was dried ($MgSO_4$) and concentrated in vacuo to afford the title compound as a white solid (2.97 g, 46%).


The following molecules were made in accordance with the procedures disclosed in **Example 128**:

(S)-*tert*-Butyl (1-oxo-1-((2,2,2-trifluoroethyl)amino)butan-2-yl)carbamate




The title molecule was isolated as a white solid: mp 108-111 °C; ^{1}H NMR (400 MHz, $CDCl_3$) δ 6.90 (s, 1H), 5.04 (m, 1H), 4.07 (m, 1H), 3.92 (m, 3H), 1.87 (m, 1H), 1.66 (m, 1H), 1.44 (s, 9H), 0.96 (t, J = 7.4 Hz, 3H); ^{19}F NMR (376 MHz, $CDCl_3$) δ -72.54; ^{13}C NMR (101 MHz, $CDCl_3$) δ 173.05, 156.04, 124.03 (q, J = 278.5 Hz), 80.30, 55.56, 40.43 (q, J = 34.7 Hz), 28.19, 25.63, 9.80; $[\alpha]_D$ = - 33.3 (c, 10.1 mg/mL in CH_2Cl_2).

***tert*-Butyl (1-oxo-1-((2,2,2-trifluoroethyl)amino)butan-2-yl)carbamate**




The title molecule was isolated as a white solid: mp 113-116 °C; ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.36 (d, J = 8.4 Hz, 1H), 5.43 - 5.25 (m, 1H), 4.16 (m, 1H), 3.98 (m, 1H), 3.82 (m, 1H), 1.84 (dt, J = 14.0, 7.0 Hz, 1H), 1.66 (dt, J = 14.2, 7.3 Hz, 1H), 1.44 (s, 9H), 0.95 (t, J = 7.3 Hz, 3H); ^{19}F NMR (376 MHz, $CDCl_3$) δ -72.51; ^{13}C NMR (101 MHz, $CDCl_3$) δ 172.94, 156.02, 124.47 (q, J = 380.8 Hz), 80.33, 55.54, 40.46 (q, J = 34.8 Hz), 28.19, 25.61, 9.79.

***tert*-Butyl (2-oxo-2-((1,1,1-trifluoropropan-2-yl)amino)ethyl)carbamate**



The title molecule was isolated as a white solid: mp 84-88 °C; ^{1}H NMR (400 MHz, $CDCl_3$) δ 6.89 (s, 1H), 5.44 (t, J = 5.8 Hz, 1H), 4.77 - 4.48 (m, 1H), 3.83 (d, J = 5.9 Hz, 2H), 1.45 (s, 9H), 1.33 (d, J = 7.0 Hz, 3H); ^{19}F NMR (376 MHz, $CDCl_3$) δ -77.63; ^{13}C NMR (101 MHz, $CDCl_3$) δ 169.84, 156.33, 125.19 (q, J = 280.9 Hz), 80.29, 46.20 (q, J = 31.7 Hz), 44.15, 28.11, 13.88; EIMS *m/z* 270 ([M]$^+$).


(*R*)-*tert*-Butyl (1-((2-fluoroethyl)amino)-1-oxopropan-2-yl)carbamate



The title molecule was isolated as a white solid: mp 91-94 °C; ^{1}H NMR (300 MHz, DMSO-d_6) δ 7.98 (bs, 1H), 6.87 (t, J = 7.2 Hz, 1H), 4.47 (t, J = 4.8 Hz, 1H), 4.32 (t, J = 5.1


Hz, 1H), 3.97 - 3.92 (m, 1H), 3.41- 3.37 (m, 1H), 3.33 - 3.28 (m, 1H), 1.37 (s, 9H), 1.16 (d, *J* = 7.3 Hz, 3H); ESIMS *m/z* 235.0 ([M+H]$^+$).

***tert*-Butyl (3-oxo-3-((2,2,2-trifluoroethyl)amino)propyl)carbamate**




The title molecule was isolated as a white solid: mp 123-125 °C; ^{1}H NMR (400 MHz, $CDCl_3$) δ 6.42 - 6.22 (m, 1H), 5.07 (s, 1H), 3.92 (qd, *J* = 9.1, 6.4 Hz, 2H), 3.43 (q, *J* = 6.2 Hz, 2H), 2.50 (t, *J* = 6.0 Hz, 2H), 1.43 (s, 9H); ^{19}F NMR (376 MHz, $CDCl_3$) δ -72.50; ^{13}C NMR (101 MHz, $CDCl_3$) δ 171.76, 156.30, 124.02 (q, *J* = 278.5 Hz), 79.67, 40.53 (q, *J* = 34.8 Hz), 36.41, 36.27, 28.31.

(*R*)-*tert*-Butyl (1-(ethylamino)-1-oxopropan-2-yl)carbamate




The title molecule was isolated as a white solid: mp 88-93 °C; ^{1}H NMR (400 MHz, $CDCl_3$) δ 6.35 (s, 1H), 5.25 - 5.04 (m, 1H), 4.21 - 3.99 (m, 1H), 3.29 (dd, *J* = 7.5, 5.9 Hz, 2H), 1.45 (s, 8H), 1.35 (d, *J* = 7.0 Hz, 3H), 1.13 (t, *J* = 7.3 Hz, 3H); ^{13}C NMR (101 MHz, $CDCl_3$) δ 172.79, 155.51, 79.59, 49.97, 34.16, 28.25, 18.77, 14.60.

(*R*)-*tert*-Butyl (1-oxo-1-((3,3,3-trifluoropropyl)amino)propan-2-yl)carbamate



The title molecule was isolated as an off white solid: mp 101-105 °C; ^{1}H NMR (300 MHz, DMSO-d_6) δ 7.96 (bs, 1H), 6.90 (d, *J* = 6.9 Hz, 1H), 3.91 - 3.86 (m, 1H), 3.34 - 3.19 (m, 2H), 2.50 - 2.32 (m, 2H), 1.37 (s, 9H), 1.15 (d, *J* = 7.2 Hz, 3H).


(*R*)-*tert*-Butyl (1-oxo-1-((2,2,2-trifluoroethyl)amino)pentan-2-yl)carbamate



The title molecule was isolated as a white solid: mp 107-122 °C; ^{1}H NMR (400 MHz, CDCl3)) δ 6.90 (s, 1H), 5.00 (d, *J* = 8.0 Hz, 1H), 4.12 (d, *J* = 7.3 Hz, 1H), 3.99 - 3.76 (m, 2H), 1.87 - 1.73 (m, 1H), 1.65 - 1.52 (m, 1H), 1.44 (s, 9H), 1.38 (m, 2H), 0.94 (t, *J* = 7.3 Hz, 3H); ^{19}F NMR (376 MHz, $CDCl_3$) δ rotomer -72.55, -73.27; ^{13}C NMR (101 MHz, CD_3OD) δ


rotomers 176.08, 157.86, minor 126.13 (q, J = 279.8 Hz), major 125.83 (q, J = 278.8 Hz), 80.65, 55.90, minor 42.27 (q, J = 35.4 Hz), major 41.24 (q, J = 35.4 Hz), 35.50, 28.73, 20.04, 14.03.

(*R*)-Benzyl (3-methyl-1-oxo-1-((2,2,2-trifluoroethyl)amino)butan-2-yl)carbamate



The title molecule was isolated as a white solid: mp 157-161 °C; ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.46 - 7.31 (m, 5H), 6.57 (d, J = 8.3 Hz, 1H), 5.34 (d, J = 8.9 Hz, 1H), 5.11 (s, 2H), 4.02 (dq, J = 16.1, 8.8, 7.7 Hz, 2H), 3.78 (td, J = 9.0, 4.7 Hz, 1H), 2.15 (q, J = 6.7 Hz, 1H), 0.97 (d, J = 6.8 Hz, 3H), 0.94 (d, J = 6.8 Hz, 3H); ^{19}F NMR (376 MHz, $CDCl_3$) δ -72.44.


Example 129: Preparation of *N*-(2,2,2-Trifluoroethyl) 1-amino-2-methylpropanecarboxamide hydrochloride



To *tert*-butyl (2-methyl-1-oxo-1-((2,2,2-trifluoroethyl)amino)propan-2-yl)carbamate (2.61 g, 9.18 mmol) in methylene chloride (20 mL) was added 4 M HCl in dioxane (20 mL). The solution was stirred for 6 h at ambient temperature. The reaction mixture was concentrated in vacuo to afford the title compound as a white solid (2.18 g).


The following molecules were made in accordance with the procedures disclosed in **Example 129**:

(*R*)-1-Oxo-1-((2,2,2-trifluoroethyl)amino)propan-2-aminium chloride




The title molecule was isolated as a white solid: mp 210-213 °C; ^{1}H NMR (400 MHz, DMSO-d_6) δ 9.22 (t, J = 6.3 Hz, 1H), 8.37 - 8.27 (m, 3H), 4.07 - 3.95 (m, 2H), 3.95 - 3.84 (m, 1H), 1.38 (d, J = 7.0 Hz, 3H); ^{19}F NMR (376 MHz, DMSO-d_6) δ -70.75; $[\alpha]_D$ = - 6.6 (c, 5.0 mg/mL in MeOH).

1-Oxo-1-((2,2,2-trifluoroethyl)amino)butan-2-aminium chloride




The title molecule was isolated as a white solid: ^{1}H NMR (300 MHz, DMSO-d_6) δ 9.12 (t, J = 5.7 Hz, 1H), 8.19 (s, 3H), 4.14-3.93 (m, 2H), 3.78 (t, J = 6.0 Hz, 1H), 1.81-1.71(m, 2H), 0.88 (t, J = 7.2 Hz, 3H); ESIMS m/z 184.90 ([(M-TFA)+H]$^+$); IR (thinfilm) 3269, 1681, 1158 cm^{-1}.

2-Oxo-2-((1,1,1-trifluoropropan-2-yl)amino)ethanaminium chloride




The title molecule was isolated as a white solid: ^{1}H NMR (300 MHz, DMSO-d_6) δ 8.96 (d, J = 8.7 Hz, 1H), 8.09 (bs, 3H), 4.71 - 4.59 (m, 1H), 3.64 - 3.62 (m, 2H), 1.27 (d, J = 6.9 Hz, 3H); EIMS m/z 170.1 ([M]$^+$).

(*R*)-1-((2-Fluoroethyl)amino)-1-oxopropan-2-aminium chloride




The title molecule was isolated as a white solid: ^{1}H NMR (300 MHz, DMSO-d_6) δ 8.76 (t, J = 5.1 Hz, 1H), 8.21 (bs, 3H), 4.54 (t, J = 5.1 Hz, 1H), 4.38 (t, J = 4.8 Hz, 1H), 3.85 - 3.79 (m, 1H), 3.50 - 3.45 (m, 1H), 3.41 - 3.36 (m, 1H), 1.36 (d, J = 7.2 Hz, 3H); ESIMS m/z 135.1 ([M+H]$^+$); IR (thinfilm) 3331, 2983, ,1660, 1161, 597 cm^{-1}.

3-Oxo-3-((2,2,2-trifluoroethyl)amino)propan-1-aminium chloride



The title molecule was isolated as a white solid: mp 193-197 °C; ^{1}H NMR (400 MHz, DMSO-d_6) δ 8.94 (t, J = 6.4 Hz, 1H), 8.16 (s, 3H), 3.99 - 3.79 (m, 2H), 2.98 (t, J = 7.3 Hz, 2H), 2.64 (t, J = 7.3 Hz, 2H); ^{19}F NMR (376 MHz, DMSO-d_6) δ -70.74.


(*R*)-1-(Ethylamino)-1-oxopropan-2-aminium chloride



The title molecule was isolated as a white solid: mp 223-236 °C; ^{1}H NMR (400 MHz, DMSO-d_6) δ 8.65 (t, J = 5.4 Hz, 1H), 8.32 (s, 3H), 3.89 - 3.66 (m, 1H), 3.12 (p, J = 7.0 Hz,


2H), 1.35 (d, *J* = 6.9 Hz, 3H), 1.05 (t, *J* = 7.2 Hz, 3H); ^{13}C NMR (101 MHz, DMSO-d_6) δ 168.98, 48.08, 33.54, 17.16, 14.43.

(*R*)-1-Oxo-1-((3,3,3-trifluoropropyl)amino)propan-2-aminium chloride




The title molecule was isolated as an off white solid: mp 128-131 °C; ^{1}H NMR (300 MHz, DMSO-d_6) δ 8.62 (bs, 1H), 8.10 (bs, 3H), 3.82 - 3.79 (m, 1H), 3.50 - 3.38 (m, 2H), 2.50 - 2.37 (m, 2H), 1.34 (d, *J* = 6.9 Hz, 3H).

(*R*)-1-Oxo-1-((2,2,2-trifluoroethyl)amino)pentan-2-aminium chloride




The title molecule was isolated as a white solid: mp 204-210 °C; ^{1}H NMR (400 MHz, CD_3OD) δ 4.17 - 4.00 (m, 1H), 3.98 - 3.69 (m, 2H), 1.83 (dt, *J* = 15.0, 7.5 Hz, 2H), 1.50 - 1.35 (m, 2H), 0.99 (t, *J* = 7.3 Hz, 3H); ^{19}F NMR (376 MHz, CD_3OD) δ -73.90; ^{13}C NMR (101 MHz, CD_3OD) δ 170.97, 125.72 (q, *J* = 277.9 Hz), 54.37, 41.30 (q, *J* = 34.7 Hz), 34.65, 19.00, 13.94.

Example 130: Preparation of (*R*)-*tert*-Butyl 1-thioxo-1-(2,2,2-trifluoroethylamino)propan-2-ylcarbamate



To a stirred solution of (*R*)-*tert*-butyl 1-oxo-1-(2,2,2-trifluoroethylamino)propan-2-ylcarbamate (100 mg, 0.37 mmol) in CH_2Cl_2 (10 mL) was added P_2S_5 (24 mg, 0.11 mmol) and hexamethyldisiloxane (HMDO) (0.13 mL, 0.59 mmol) at room temperature and the mixture was refluxed for 3 h. The reaction mixture was cooled to room temperature and another portion of P_2S_5 (24 mg, 0.11 mmol) was added and the resulting mixture was refluxed for 18 h. The volatiles were evaporated, pentane (25 mL) was added to the residue and stirred for 10-15 min. The pentane layer was decanted, concentrated in vacuo and the residue was passed through a short silica pad eluting with pentane followed by CH_2Cl_2 to give the title compound as colorless liquid (30 mg, 30%): ^{1}H NMR (400 MHz, DMSO-d_6) δ 10.27 (t, *J* = 5.4 Hz, 1H), 7.00 (d, *J* = 6.8 Hz, 1H), 4.57 - 4.35 (m, 3H), 1.32 (s, 9H), 1.25 (d, *J* = 7.6 Hz, 3H); ESIMS *m/z* 286.2 ([M+H]$^+$); IR (thin film) 3233, 1683, 1257 cm^{-1}.


Example 131: Preparation of (*R*)-1-Thioxo-1-((2,2,2-trifluoroethyl)amino)propan-2-aminium 2,2,2-trifluoroacetate



To a stirred solution of (*R*)-*tert*-butyl 1-thioxo-1-(2,2,2-trifluoroethylamino)propan-2-ylcarbamate (200 mg, 0.69 mmol) in CH_2Cl_2 (5 mL) was added TFA (0.5 mL) dropwise and the reaction mixture was stirred for 18 h. The volatiles were evaporated and the residue was triturated with pentane to give the title compound as colorless gum, which was taken to next step without further purification (200 mg): ^{1}H NMR (300 MHz, DMSO-d_6) δ 10.99 (bs, 1H), 8.23 (bs, 2H), 4.62-4.55 (m, 2H), 4.23-4.19 (m, 1H), 1.43 (d, J = 8.4 Hz, 3H); ESIMS *m/z* 186.2 ([M+H]$^+$); IR (thin film) 3445, 2967, 1168 cm^{-1}.


The following molecule was made in accordance with the procedures disclosed in **Example 131**:

(*S*)-1-Oxo-1-((2,2,2-trifluoroethyl)amino)butan-2-aminium 2,2,2-trifluoroacetate



The title molecule was isolated as a colorless gum: ^{1}H NMR (300 MHz, DMSO-d_6) δ 9.12 (t, J = 5.7 Hz, 1H), 8.19 (bs, 2H), 4.14-3.93 (m, 2H), 3.80 (t, J = 6.0 Hz, 1H), 1.81-1.71 (m, 2H), 0.88 (t, J = 7.2 Hz, 3H); ESIMS *m/z* 185.00 ([M+H]$^+$); IR (thin film) 3459, 1674, 1169 cm^{-1}.


Example 132: Preparation of 2-Bromo-*N*-((S)-1-oxo-1-((2,2,2-trifluoroethyl)amino)propan-2-yl)-4-((*E*)-4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-en-1-yl)benzamide (F10 and F11)



The title molecule was prepared as described in Example 15. The diastereomeric pairs were separated by chiral HPLC using Chiralpak® IA (4.6 x 250 mm) 5μm column using 0.1% TFA in hexane and isopropanol as the mobile phase (isocratic 70:30) with a flow rate 1.0 mL/min at ambient temperature. Diastereomer F10 was collected at a retention time of 4.55 min and possessed an optical rotation of $[\alpha]_D^{30}$ = + 35.6 (*c*, 0.5% in CH_2Cl_2).

Diastereomer F11 was collected at 8.71 min and possessed an optical rotation of $[\alpha]_D^{30}$ = -82.0 (*c*, 0.5% in CH_2Cl_2). Characterization data for these molecules are listed in Table 2.


Example 133: Preparation of 2-Bromo-*N*-((R)-1-oxo-1-((2,2,2-trifluoroethyl)amino)propan-2-yl)-4-((*E*)-4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-en-1-yl)benzamide (F12 and F13)



The title molecule was prepared as described in Example 15. The diastereomeric pairs were separated by chiral HPLC using Chiralpak® IA (4.6 x 250 mm) 5μm column using 0.1% TFA in hexane and isopropanol as the mobile phase (isocratic 70:30) with a flow rate 1.0 mL/min at ambient temperature. Diastereomer F12 was collected at a retention time of 5.62 min and possessed an optical rotation of $[\alpha]_D^{30}$ = +59.4 (*c*, 1% in CH_2Cl_2). Diastereomer F13 was collected at 8.85 min and possessed an optical rotation of $[\alpha]_D^{30}$ = - 44.0 (*c*, 1% in CH_2Cl_2). Characterization data for these molecules are listed in Table 2.

The following molecules was prepared in accordance with the procedures disclosed in **Example 133**:

***N*-((*R*)-1-Oxo-1-((2,2,2-trifluoroethyl)amino)propan-2-yl)-4-((*E*)-4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-en-1-yl)-2-(trifluoromethyl)benzamide (F20A and F20B)**




Diastereomer F20A (isomer 1) was collected at a retention time of 4.13 min and possessed an optical rotation of $[\alpha]_D^{25}$ = +49.2 (*c*, 1.0% in CH_2Cl_2). Diastereomer F20B was collected at 4.88 min and possessed an optical rotation of $[\alpha]_D^{25}$ = -38.8 (*c*, 1.0% in CH_2Cl_2). Characterization data for these molecules are listed in Table 2.

F20A and F20B stereochemical assignment F20A and F20B were dissolved in $CDCl_3$ and placed in a 100 μm path length cell with BaF_2 windows. IR and vibrational circular dichroism (VCD) spectra were recorded on a IR-2XTM VCD spectrometer (BioTools, Inc.) equipped with dual PEM accessory, with 4 cm^{-1} resolution. The sample and $CDCl_3$ spectra were acquired for 21 h on an instrument optimized at 1400 cm^{-1}. The solvent-subtracted IR and VCD spectra were collected.

Theoretical Calculations: F20 with *R,R*- and *S,R*-configurations were built with Maestro (Schrodinger, LLC. New York, NY). The conformational search was carried out with MacroModel (Schrodinger, LLC. New York, NY) with MMFF94x force field to generate low-energy conformers. Single point calculation (SPE), geometry, frequency, and IR and VCD calculations were performed at the DFT level (B3LYP/lacvp**) in Jaguar (Schrodinger, LLC. New York, NY). A scaling factor of 0.96 was applied to the frequency calculation.*Analysis*: for F20 with *R,R*- and *S,R*-configurations, the top 100 low-energy conformers generated with MacroModel were selected for DFT SPE calculations. These calculations resulted in the 8 and 4 conformers that have energies within 1 kcal/mol higher than the lowest energy conformer for *R,R*- and *S,R*-configurations, respectively. The frequency calculations were performed on these conformers to determine the IR and VCD spectra. The Boltzmann-weighted IR and VCD spectra of these conformers were compared with the observed IR and VCD spectra. Based on the overall agreement in VCD pattern between the observed and calculated spectra, the absolute configuration of F20Ais assigned as R,R- configuration. The assignment was evaluated by CompareVOA program (BioTools). The confidence level of the assignment is 88% based on a database that includes 105 previous correct assignments for different chiral structures. However, the observed spectrum for F20B does not agree well with the calculated spectrum for S,R- configuration with a confidence level of 65%. But considering that the compound has only one chiral center, two possible configurations and F20A is of *R,R*- configuration, F20B can be with high confidence assigned as the *S,R*-configuration.


Example 134: Preparation of (*E*)-2-Methyl-*N*-(2-methyl-1-thioxo-1-(2,2,2-trifluoroethylamino)propan-2-yl)-4-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-enyl)benzothioamide (F31)



To a stirred solution of (*E*)-2-methyl-*N*-(2-methyl-1-oxo-1-(2,2,2-trifluoroethylamino)propan-2-yl)-4-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-enyl)benzamide (400 mg, 0.68 mmol) in CH_2Cl_2 (50 mL) was added P_2S_5 (75 mg, 0.34 mmol) and HMDO (0.25 mL, 1.12 mmol) at room temperature and the mixture was refluxed for 3 h. The reaction mixture was cooled to room temperature and another portion of P_2S_5 (75 mg, 0.34 mmol) was added and the resulting mixture was refluxed for 18 h. The volatiles


were evaporated and the residue was purified by prep TLC to give the title compound as pale yellow gum (47 mg, 11%). Characterization data for this molecule is listed in Table 2.

Example 135: Preparation of (*E*)-2-Bromo-*N*-(2-methyl-1-oxo-1-((2,2,2-trifluoroethyl)amino)propan-2-yl)-4-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-en-1-yl)benzamide (F1)



To (*E*)-2-bromo-4-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-enyl)benzoic acid (200 mg, 0.409 mmol) in MeCN (5 mL) was added 1*H*-benzo[d][1,2,3]triazol-1-ol hydrate (63 mg, 0.411 mmol), HBTU (155 mg, 0.409 mmol), *N*-(2,2,2-trifluoroethyl) 1-amino-2-methyl-propanecarboxamide hydrochloride (180 mg, 0.816 mmol) and diisopropylethylamine (0.24 mL, 1.38 mmol). After 24 h the material was concentrated in vacuo. The crude product was purified by passing the crude reaction mixture through a silica frit and eluting with EtOAc/hexane (1:2). The recovered material was further purified by medium pressure chromatography on silica with EtOAc/hexane as the eluent to afford the title as a white foam (147 mg, 55%). Characterization data for this molecule is listed in Table 2.

Example 136: Preparation of 1-(3,5-Difluoro-4-methoxyphenyl)-2,2,2-trifluoroethanone




Isopropyl magnesium chloride lithium chloride complex (22.0 mL, 28.02 mmol) was added dropwise to a stirred solution of 5-bromo-1,3-difluoro-2-methoxybenzene (5.0 g, 22.42 mmol) at -5 °C in THF (100 mL) and the reaction mixture was stirred at same temperature for 30 min. Methyl trifluoroacetate (3.67g, 28.69 mmol) was added dropwise and the reaction mixture was stirred at ambient temperature for 2 h. A 2 N HCl solution (200 mL) was added to quench the reaction and then it was extracted with diethylether. The combined organic layers were washed with brine, dried (Na_2SO_4) filtered and concentrated to afford the title compound (5.4 g, crude) as a yellow liquid. The material was taken to next step without

further purification. ^{1}H NMR(400 MHz, $CDCl_3$) δ 7.68 - 7.60 (m, 2H) 4.19 (s, 3H); ESIMS *m/z* 240.1 ($[M]^+$).


The following molecule was prepared in accordance with the procedures disclosed in **Example 136**:

2,6-Difluoro-4-(2,2,2-trifluoroacetyl)benzonitrile



^{1}H NMR (400 MHz, CDCl3) δ 7.45 (d, *J* = 8.4 Hz, 1H), 7.37 (d, *J* = 8.4 Hz, 1H). ; EIMS *m/z* 235.1 ($[M]^+$).

Example 137: Preparation of (*E*)-*N*-(1-((2-Fluoroethyl)amino)-1-oxopropan-2-yl)-4-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-en-1-yl)-2-(trifluoromethyl)benzamide (P1618A)



Step 1: (*2R*)-*tert*-Butyl 2-(4-((*E*)-4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-enyl)-2-(trifluoromethyl)benzamido)propanoate: The title compound was prepared according to procedures outlined in Example 15: ^{1}HNMR (400 MHz, DMSO d_6) δ 8.73 (d, *J* = 6.8 Hz, 1H), 7.92 - 7.90 (m, 3H), 7.61 (d, *J* = 7.2 Hz, 1H), 7.36 (d, *J* = 7.6 Hz, 1H), 6.99 (dd, *J* = 15.2, 9.2 Hz, 1H), 6.77 (d, *J* = 15.2 Hz, 1H), 4.85 - 4.80 (m, 1H), 4.30 - 4.26 (m, 1H), 1.43 (s, 9H), 1.33 (d, *J* = 6.8 Hz, 3H); ESIMS *m/z* 601.9 ($[M-H]^-$); IR (KBr) 3414, 1732, 1661, 1170, 748 cm^{-1}.


Step 2: (*2R*)-(4-((*E*)-4,4,4-Trifluoro-3-(3,4,5-trichlorophenyl)but-1-enyl)-2-(trifluoromethyl)benzamido)propanoic acid: TFA (1 mL) was added to a stirred solution of *tert*-butyl 2-(2-bromo-4-((*E*)-4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-enyl)benzamido)propanoate (1.0 g, 1.63 mmol) in CH_2Cl_2 (20 mL) at 0 °C and the reaction mixture was stirred at ambient temperature for 18 h. The volatiles were evaporated under vacuum and the residue was triturated with pentane to afford the title compound as brown solid (0.65 g, 67 %): ^{1}HNMR (400 MHz, DMSO-d_6) δ 12.60 (bs, 1H), 8.82 (d, *J* = 8.0 Hz, 1H), 7.99 (s, 1H), 7.92 - 7.89 (m, 3H), 7.51 (d, *J* = 8.0 Hz, 1H), 7.08 (dd, *J* = 15.6, 8.8 Hz,

1H), 6.88 (d, *J* = 15.6 Hz, 1H), 4.88 - 4.83 (m, 1H), 4.41 - 4.34 (m, 1H), 1.34 (d, *J* = 7.2 Hz, 3H); ESIMS: *m/z* 545.7 ([M-H]$^+$); IR (KBr) 3410, 3281, 2928, 1728, 1172, 744 cm^{-1}.

Step 3. (*E*)-*N*-(1-((2-Fluoroethyl)amino)-1-oxopropan-2-yl)-4-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-en-1-yl)-2-(trifluoromethyl)benzamide (P1618): DIPEA (0.60 mL, 1.08 mmol), PyBOP (180 mg, 0.36 mmol) 2-(4-((*E*)-4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-enyl)-2-(trifluoromethyl)benzamido)propanoic acid (35 mg, 0.36 mmol) were added to a stirred solution of compound 1 (200 mg, 0.36 mmol) in CH_2Cl_2 (10 mL) at ambient temperature and the reaction mixture was stirred for 18h. The reaction mixture was diluted CH_2Cl_2, washed with 1N HCl, followed by a saturated $NaHCO_3$ solution, water and brine. The organic phase was dried (Na_2SO_4), filtered, concentrated and the residue was purified by column chromatography on silica (100-200 mesh) eluting with 10% EtOAc in petroleum ether to afford the title compound as a brown solid(85 mg, 39%).


The following molecule was prepared in accordance with the procedures disclosed in **Example 137, Step 1**:

***tert*-Butyl 2-(2-bromo-4-((*E*)-4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-enyl)benzamido)propanoate**



^{1}H NMR (400 MHz, DMSO-d_6): δ 8.82 (d, *J* = 7.6 Hz, 1H), 7.99 (s, 1H), 7.91 - 7.90 (m, 3H), 7.50 (d, *J* = 7.6 Hz, 1H), 7.07 (dd, *J* = 16.0, 8.8 Hz, 1H), 6.88 (d, *J* = 15.2 Hz, 1H), 4.88 - 4.83 (m, 1H), 4.31 - 4.27 (m, 1H), 1.42 (s, 9H), 1.32 (d, *J* = 7.6 Hz, 3H); ESIMS *m/z* 611.7 ([M-H]$^-$); IR (KBr) 3296, 2932, 1732, 1162, 743, 556 cm^{-1}.

(*E*)-*tert*-Butyl 2-(2-bromo-4-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-enyl)benzamido)acetate




Isolated as a (620 mg, 72 %) pale yellow solid. ^{1}H NMR (400 MHz, DMSO-d_6) δ 8.77 (t, *J* = 6.0 Hz, 1H), 7.93 - 7.91 (m, 3H), 7.62 (d, *J* = 6.8 Hz, 1H), 7.39 (d, *J* =8.0 Hz,

1H), 7.00 (dd, *J* = 15.6, 9.2 Hz, 1H), 6.77 (d, *J* = 16.0 Hz, 1H), 4.86 - 4.81 (m, 1H), 3.86 - 3.85 (m, 2H), 1.33 (s, 9H). ESIMS *m/z* 599.87 ([M+H]$^+$).


The following molecule was prepared in accordance with the procedures disclosed in **Example 137, Step 2**:

2-(2-Bromo-4-((*E*)-4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-enyl)benzamido)propanoic acid




^{1}HNMR (400 MHz, DMSO-d_6): δ 12.62 (bs, 1H), 8.73 (d, *J* = 9.6 Hz, 1H), 7.93 - 7.91 (m, 3H), 7.61 (d, *J* = 8.1 Hz, 1H), 7.37 (d, *J* = 7.8 Hz, 1H), 7.01 (dd, *J* = 15.6, 9.0 Hz, 1H), 6.78 (d, *J* = 15.9 Hz, 1H), 4.89 - 4.79 (m, 1H), 4.42 - 4.32 (m, 1H), 1.36 (d, *J* = 7.2 Hz, 3H); ESIMS: *m/z* 558.0 ([M+H]$^+$); IR (KBr) 3418, 1650, 1115, 747, 560 cm^{-1}.

(*E*)-2-(2-Bromo-4-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-enyl)benzamido)acetic acid



Isolated as an (550 mg, 97 %) off white solid. ^{1}H NMR (300 MHz, DMSO-d6) δ 12.56 (bs, 1H), 8.73 (t, *J* = 5.4 Hz, 1H), 7.93 - 7.91 (m, 3H), 7.62 (d, *J* = 9.3 Hz, 1H), 7.40 (d, *J* = 8.1 Hz, 1H), 7.01 (dd, *J* = 15.9, 9.0 Hz, 1H), 6.78 (d, *J* = 15.9 Hz, 1H), 4.89 - 4.80 (m, 1H), 3.90 - 3.88 (m, 2H). ESIMS *m/z* 541.82 ([M-H]$^-$).

Example 138: Preparation of (*E*)-2-Chloro-5-hydroxy-*N*-(2-oxo-2-((2,2,2-trifluoroethyl)amino)ethyl)-4-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-en-1-yl)benzamide



Step 1. **Methyl 4-bromo-2-chloro-5-methoxybenzoate:** A 25 mL round bottomed flask equipped with a magnetic stir bar was charged with 4-bromo-2-chloro-5-methoxybenzoic acid (*JACS*, **1963**, 730-2; 1.25 g, 4.72 mmol), 20% MeOH/EtOAc (25 mL)

and cooled in an ice-water bath. Trimethylsilyldiazomethane ($TMSCHN_2$ 2 M in hexanes, 2.6 mL, 5.20 mmol) was added dropwise via an addition funnel. The reaction continued to stir for 1 h then it was concentrated to afford the title compound as a white solid (1.31 g, 100%): mp 78-79 °C; ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.65 (s, 1H), 7.36 (s, 1H), 3.94 (s, 3H), 3.93 (s, 3H); EIMS *m/z* 280 ($[M]^+$).


Step 2. 2-Chloro-5-methoxy-4-vinylbenzoic acid: A 25 mL round bottomed flask was charged with methyl 4-bromo-2-chloro-5-methoxybenzoate (640 mg, 2.29 mmol), K_2CO_3 (665 mg, 4.81 mmol), potassium trifluoro(vinyl)borate (920 mg, 6.87 mmol), $PdCl_2$(dppf) (84 mg, 0.11 mmol) and anhydrous DMSO (15 mL) and stirred at 80 °C for 2 h. The reaction was allowed to cool, water (150 mL) was added and then extracted several times with Et_2O. The organic layer was washed with brine, dried over $MgSO_4$, filtered and concentrated to give a brown residue. The crude product was purified via flash chromatography eluting with 15% Et_2O/hexanes to give methyl 2-chloro-5-methoxy-4-vinylbenzoate as a yellow oil (440 mg, 85%). To a 25 mL round bottomed flask containing methyl 2-chloro-5-methoxy-4-vinylbenzoate (440 mg, 1.94 mmol) and MeOH (10 mL) was added 1N NaOH (2 mL, 2.04 mmol) and reaction stirred at ambient temperature for 18 h. The reaction mixture was concentrated to give a solid residue. The residue was dissolved in water and extracted 1x with 50% Et_2O/hexanes. The aqueous layer was made acidic with 2N HCl and extracted 2x with CH_2Cl_2, dried over $MgSO_4$, filtered and concentrated to afford the title compound as a white solid (0.39 g, 94%): ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.54 (d, J = 0.5 Hz, 1H), 7.52 (s, 1H), 6.98 (ddd, J = 17.7, 11.2, 0.6 Hz, 1H), 5.87 (dd, J = 17.7, 1.1 Hz, 1H), 5.45 (dd, J = 11.2, 1.1 Hz, 1H), 3.90 (s, 3H).

Step 3. (*E*)-2-Chloro-5-methoxy-4-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-en-1-yl)benzoic acid: A 50 mL 3 neck round bottomed flask was charged with 2-chloro-5-methoxy-4-vinylbenzoic acid (390 mg, 1.84 mmol)), 5-(1-bromo-2,2,2-trifluoroethyl)-1,2,3-trichlorobenzene (754 mg, 2.20 mmol) and anhydrous *N*-methyl pyrrolidinone (10 mL). Nitrogen was bubbled into the reaction mixture for 15 min. After which time, 2,2'-dipyridyl (57.3 mg, 0.37 mmol) and CuBr (11.7 mg, 0.18 mmol) were added and reaction mixture stirred at 150 °C for 1 h. Reaction mixture was allowed to cool, water (300 mL) was added and extracted several times with Et_2O. Organic layer was washed repeatedly with water, dried over $MgSO_4$, filtered and concentrated to afford- the title compound as a light brown foam (870 mg, 100%). This material was 95% pure by LC/MS; ESIMS *m/z* 473 ($[M-H]^-$). This material was used without further purification.

Step 4. (*E*)-2-Chloro-5-methoxy-N-(2-oxo-2-((2,2,2-trifluoroethyl)amino)ethyl)-4-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-en-1-yl)benzamide: A 50 mL round bottomed flask was charged with (*E*)-2-chloro-5-methoxy-4-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-en-1-yl)benzoic acid (780 mg, 1.65 mmol)), di(1H-imidazol-1-yl)methanone (267 mg, 1.65 mmol) and anhydrous THF (30 mL). The resulting mixture was heated at reflux until it ceased giving off gas. 2-Amino-*N*-(2,2,2-trifluoroethyl)acetamide HCl (257 mg, 1.65 mmol) was added in one portion and the reaction mixture continued to stir at reflux for 18 h. The reaction mixture was concentrated to dryness and the residue was taken up in Et_2O (50 mL) and 0.1N HCl (10 mL). The layers were separated. The aqueous layer was extracted 2x with Et_2O. The Et_2O layers were combined and washed 1x with aqueous $NaHCO_3$, 1x with brine, dried over $MgSO_4$, filtered and concentrated to give a brown oil. The crude product was purified via flash chromatography eluting with 30-40% EtOAc/hexanes to afford the title compound - as an off white foam (280 mg, 28%): ^1H NMR (400 MHz, $CDCl_3$) δ 7.45 (s, 1H), 7.41 (s, 2H), 7.29 (s, 1H), 7.28 (s, 1H), 6.84 (m, 2H), 6.43 (dd, *J* = 16.0, 8.3 Hz, 1H), 4.25 (d, *J* = 5.4 Hz, 2H), 4.10 (m, 1H), 3.97 (qd, *J* = 9.0, 6.4 Hz, 2H), 3.87 (s, 3H); ESIMS *m/z* 613.1 ($[M+H]^+$).


Step 5. (*E*)-2-Chloro-5-hydroxy-*N*-(2-oxo-2-((2,2,2-trifluoroethyl)amino)ethyl)-4-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-en-1-yl)benzamide: A 25 mL round bottomed flask was charged with (*E*)-2-chloro-5-methoxy-*N*-(2-oxo-2-((2,2,2-trifluoroethyl)amino)ethyl)-4-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-en-1-yl)benzamide (57 mg, 0.093 mmol) and CH_2Cl_2 (3 mL). The reaction mixture was cooled to -78° C and boron tribromide (BBr_3, 1.0M solution in CH_2Cl_2, 0.33 mL, 0.326 mmol) was added slowly via syringe. The reaction allowed to warm to ambient temperature and stirred for 18 h. An additional 0.3-0.4 mL of BBr_3 was added at ambient temperature and continued to stir for 3 h. The reaction mixture was added to aqueous $NaHCO_3$ and extracted 3x with CH_2Cl_2. The CH_2Cl_2 layers were combined and dried over $MgSO_4$, filtered and concentrated to give an oil. The crude material was purified via flash chromatography eluting with 50% EtOAc/hexanes to afford the title compound as a white solid (18 mg, 33%): mp 190 °C (dec.); ^1H NMR (400 MHz, $CDCl_3$) δ 9.81 (s, 1H), 8.06 (d, *J* = 7.0 Hz, 1H), 7.71 (m, 1H), 7.44 (d, *J* = 2.8 Hz, 2H), 7.37 (s, 1H), 7.20 (s, 1H), 6.82 (d, *J* = 15.9 Hz, 1H), 6.50 (m, 1H), 4.14 (m, 3H), 3.89 (m, 2H); ESIMS *m/z* 599 ($[M+H]^+$).

Example 139: Preparation of 1-(3,4-Dichlorophenyl)-2,2-difluoropropan-1-one




To a magnetically stirred solution of 4-bromo-1,2-dichlorobenzene (5.64 g, 24.98 mmol) in dry Et_2O (109 mL) was added *n*-BuLi (10.86 mL, 24.98 mmol) via an addition funnel under a nitrogen atmosphere. The reaction mixture was stirred at -78 °C for 30 min, A solution of ethyl 2,2-difluoropropanoate (3.0 g, 21.7 mmol) in Et_2O (10 mL) was added dropwise over 15 min and allowed to stir for 1 h. The reaction was then carefully quenched with 1 N HCl (4 mL) and allowed to warm to 23 °C. The solution was dilute with Et_2O and washed with water. The combined organic layers were dried over Na_2SO_4, concentrated under reduced pressure and the resulting material was purified via flash column chromatography using 100% hexanes to 5% acetone/95% hexanes as eluent. The relevant fractions were concentrated under reduced pressure to afford the title compound as a colorless oil (3.89 g, 71%): ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.21 – 8.18 (m, 1H), 7.99 – 7.93 (m, 1H), 7.59 (dd, *J* = 8.4, 4.2 Hz, 1H), 1.89 (t, *J* = 19.6 Hz, 3H); ^{19}F NMR (376 MHz, $CDCl_3$) δ -92.08 – -93.21 (m); EISMS *m/z* 240 ([M-H]$^+$).

Example 140: (*E*)-4-(4,4,4-Trifluoro-3-(3,4,5-trichlorophenyl)but-1-enyl)-2-vinylbenzoic acid




To a stirred solution of (*E*)-2-bromo-4-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-enyl)benzoic acid (600 mg, 1.23 mmol) in dry toluene (10 mL) was added tributyl(vinyl)stannane (470 mg, 1.48 mmol) and the mixture was degassed with argon for 15 min. $Pd(PPh_3)_4$ (72 mg, 0.06 mmol) was added and the reaction mixture was refluxed for 2 h. The reaction mixture was brought to ambient temperature, water was added and the mixture extracted with EtOAc. The organic layer was washed with 2N HCl and brine, dried (Na_2SO_4), filtered, and concentrated. The residue was purified by column chromatography on silica eluting with 30% EtOAc in petroleum ether to afford the title compound as brown solid (295 mg, 55%): ^{1}H NMR (300 MHz, DMSO-d_6) δ 13.05 (bs, 1H), 7.91 (s, 2H), 7.81 - 7.75 (m, 2H), 7.59 (d, *J* = 8.1 Hz, 1H), 7.48 (dd, *J* = 17.4, 10.8 Hz, 1H), 7.03 (dd, *J* = 15.9, 8.7 Hz, 1H), 6.84 (d, *J* = 15.6 Hz, 1H), 5.88 (d, *J* = 16.5 Hz, 1H), 5.39 (d, *J* = 12.3 Hz, 1H), 4.89 - 4.82 (m, 1H); ESIMS *m/z* 432.18 ([M-H]$^-$); IR (thinfilm) 3418, 1689, 1114, 747 cm^{-1}.

Example 141: (*E*)-2-Iodo-4-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-enyl)benzoic acid



Per Buchwald, et al.; *JACS*, **2002**, *124*, 14844-14845, potassium iodide (KI, 273 mg, 1.64 mmol), CuI (31 mg, 0.16 mmol) and *trans-N,N'*-dimethylcyclohexane-1,2-diamine (catalytic amount) were added to a solution of (*E*)-2-bromo-4-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-enyl)benzoic acid (400 mg, 0.82 mmol) in 1,4-dioxane (8 mL). The mixture in an Ace pressure tube was heated at 100 °C for 3 h. The reaction mixture was brought to ambient temperature and filtered through a Celite® pad. The filtrate was concentrated and residue was diluted with EtOAc and washed with 1N HCl followed by brine. The organic layer was dried (Na_2SO_4), filtered, and concentrated. The residue was purified by column chromatography on silica eluting with 25% EtOAc in petroleum ether to afford the title compound as brown semi solid (240 mg, 55%): ^{1}H NMR (400 MHz, DMSO-d_6) δ 13.3 (bs, 1H), 8.21 (s, 1H), 7.91 (s, 2H), 7.71 - 7.64 (m, 2H), 7.01 (dd, J = 15.6, 9.2 Hz, 1H), 6.75 (d, J = 15.6 Hz, 1H), 4.85 - 4.81 (m, 1H); ESIMS *m/z* 532.8 ([M-H]$^-$); IR (thinfilm) 3436, 1699, 1113, 750 cm^{-1}.

Example 142: Preparation of (*E*)-2-Bromo-*N*-(2-methyl-1-(neopentylamino)-1-oxopropan-2-yl)-4-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-en-1-yl)benzamide




Step 1. **(*E*)-2-(2-Bromo-4-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-en-1-yl)benzamido)-2-methylpropanoic acid:** A 25 mL round bottomed flask equipped with a magnetic stir bar and reflux condenser was charged with (*E*)-2-bromo-4-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-en-1-yl)benzoic acid (400 mg, 0.82 mmol) and 1,2-dichloroethane (DCE) (5 mL). Thionyl chloride (0.12 mL, 1.64 mmol) was added neat in one portion and the resulting reaction mixture was heated at reflux for 2 h. After which time, reaction mixture was allowed to cool and concentrated to give the crude acid chloride which was used without further purification. To a solution containing $NaHCO_3$ (68.8 mg, 0.82 mmol), 2-amino-2-methylpropanoic acid (84 mg, 0.82 mmol) and

dodecyltrimethylammonium bromide (2.52 mg, 8.19 μmol) in 10 mL of THF was added to the acid chloride in THF (1 mL). The resulting mixture was heated at reflux for 18 h. Reaction mixture was allowed to cool and added to water, made acidic with 0.1N HCl, extracted (3x) with Et_2O, washed (1x) with brine, dried over $MgSO_4$, filtered and evaporated to afford the title compound as a light brown foam (400 mg, 85%) : ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.60 (d, *J* = 1.6 Hz, 1H), 7.59 (d, *J* = 8.0 Hz, 1H), 7.40 (s, 2H), 7.37 (dd, *J* = 8.1, 1.6 Hz, 1H), 6.63 (s, 1H), 6.53 (d, *J* = 15.9 Hz, 1H), 6.38 (dd, *J* = 15.9, 7.9 Hz, 1H), 4.10 (p, *J* = 8.3 Hz, 1H), 1.73 (s, 6H). This material is used without further purification.

Step 2. (*E*)-2-(2-Bromo-4-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-en-1-yl)phenyl)-4,4-dimethyloxazol-5(4H)-one: A 25 mL round bottomed flask was charged with (*E*)-2-(2-bromo-4-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-en-1-yl)benzamido)-2-methylpropanoic acid (400 mg, 0.70 mmol), CH_2Cl_2 (10 mL) and stirred at 0 °C. EDC·HCl (134 mg, 0.70 mmol) was added in one portion as a solid and the reaction mixture was allowed to warm toward ambient temperature and continued to stir for 1 h. The reaction was diluted with CH_2Cl_2, washed with brine, dried over $MgSO_4$, filtered and evaporated to give a dark oil. The crude product was purified via flash chromatography eluting with 50% hexanes/CH_2Cl_2 to afford the title compound as an off white foam (220 mg, 57): ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.76 (d, *J* = 8.1 Hz, 1H), 7.73 (d, *J* = 1.6 Hz, 1H), 7.42 (d, *J* = 7.8 Hz, 3H), 6.56 (d, *J* = 15.9 Hz, 1H), 6.45 (dd, *J* = 15.9, 7.7 Hz, 1H), 4.12 (p, *J* = 8.5 Hz, 1H), 1.57 (s, 6H); ^{19}F NMR (376 MHz, $CDCl_3$) δ -68.55; ESIMS *m/z* 554 ([M-H]$^-$).

Step 3. (E)-2-Bromo-*N*-(2-methyl-1-(neopentylamino)-1-oxopropan-2-yl)-4-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-en-1-yl)benzamide: A 10 mL round bottomed flask was charged with (*E*)-2-(2-bromo-4-(4,4,4-trifluoro-3-(3,4,5-trichlorophenyl)but-1-en-1-yl)phenyl)-4,4-dimethyloxazol-5(4H)-one (90 mg, 0.16 mmol) and CH_2Cl_2 (2 mL). 2,2-Dimethylpropan-1-amine (28.2 mg, 0.324 mmol) was added neat via pipette and the reaction stirred at ambient temperature for 18 h. The reaction mixture was evaporated to give an oil. The crude product was purified via flash chromatography eluting with 20% EtOAc/hexanes to afford the title compound as a white foam (90 mg, 86%): ^{1}H NMR (400 MHz, $CDCl_3$) δ 7.60 (d, *J* = 1.6 Hz, 1H), 7.51 (d, *J* = 8.0 Hz, 1H), 7.40 (s, 2H), 7.36 (dd, *J* = 8.1, 1.6 Hz, 1H), 6.66 (d, *J* = 10.0 Hz, 2H), 6.53 (d, *J* = 15.9 Hz, 1H), 6.38 (dd, *J* = 15.9, 7.8 Hz, 1H), 4.11 (m, 1H), 3.12 (d, *J* = 6.2 Hz, 2H), 1.72 (s, 6H), 0.93 (s, 9H); 19F NMR (376 MHz, $CDCl_3$) δ -68.62; ESIMS *m/z* 643.19 ([M+H]$^+$).

Example 143: Preparation of 3,5-Dibromo-4-chlorobenzaldehyde



Step 1. Methyl 4-amino-3,5-dibromobenzoate: conc. H_2SO_4 (1.35 mL, 25.48 mmol) was added dropwise to a stirred solution of 4-amino-3,5-dibromobenzoic acid (5.0 g, 16.99 mmol) in MeOH (50 mL) at ambient temperature and the reaction mixture was then stirred at 80 °C for 8 h. The reaction mixture was brought to ambient temperature, volatiles were evaporated and ice cold water was added to the residue and which was then extracted with EtOAc. The organic layer was washed with an aqueous $NaHCO_3$ solution followed by brine and water. The solution was then dried (Na_2SO_4), filtered and concentrated to afford the title compound as an off white solid (5.0 g, 95%): ^{1}H NMR (300 MHz, DMSO-d_6) δ 7.91 (s, 2H), 6.20 (bs, 2H), 3.78 (s, 3H); ESIMS *m/z* 307.0 ([M]$^+$); IR (thin film) 3312 , 2953, 1726, 595 cm^{-1}.


Step 2. Methyl 3,5-dibromo-4-chlorobenzoate: $CuCl_2$ (2.82 g, 21.0 mmol) in MeCN (30 mL) was stirred at 80 °C for 30 min. To this mixture *tert*-butylnitrite (2.7 mL, 23 mmol) was then added dropwise at same temperature and the mixture was stirred for another 10 min. Methyl 4-amino-3,5-dibromobenzoate (5.0 g, 16 mmol) in MeCN (30 mL) was added dropwise to the reaction mixture and then stirred at 80 °C for 30 min. The reaction mixture was brought to ambient temperature and an aqueous ammonia solution (20 mL) was added to the reaction mixture and extracted with petroleum ether. The organic layer was washed with brine followed by water, dried (Na_2SO_4), filtered and concentrated to afford the title compound as an off white solid (4.5 g, 84%). ^{1}H NMR (300 MHz, DMSO-d_6) δ 8.21 (s, 2H), 3.94 (s, 3H); ESIMS *m/z* 326 ([M]$^+$); IR (thin film) 1732, 746 cm^{-1}.

Step 3. (3,5-Dibromo-4-chlorophenyl)methanol: $NaBH_4$ (1.53 g, 40.65 mmol) was added portionwise to a stirred solution of methyl 3,5-dibromo-4-chlorobenzoate (4.45 g, 13.6 mmol) in MeOH (50 mL) at 0 °C. The reaction mixture was then stirred at ambient temperature for 8 h. The volatiles were evaporated and the residue was diluted with CH_2Cl_2 and washed with brine followed by water. The organic layer was dried (Na_2SO_4), filtered and concentrated to afford the title compound as an off white solid (3.3 g, 80%): ^{1}H NMR (300 MHz, DMSO-d_6) δ 7.71 (s, 2H), 5.49 (bs, 1H), 4.48 (d, J = 4.5 Hz, 2H); ESIMS *m/z* 297.9 ([M]$^+$); IR (thin film) 3460, 747, 534 cm^{-1}.

Step 4. 3,5-Dibromo-4-chlorobenzaldehyde: Pyridinium chlorochormate (PCC, 3.44 g, 15.9 mmol) was added in one portion to a stirred solution of (3,5-dibromo-4-

chlorophenyl)methanol (3.2 g, 11.0 mmol) in $CHCl_3$ (40 mL) at ambient temperature and the reaction mixture was stirred overnight. The reaction mixture was filtered through Celite®, the Celite® pad was washed with $CHCl_3$ and the filtrate was concentrated to afford the title compound as an off white solid (2.0 g, 62%): mp 110-113 °C; ^{1}H NMR (300 MHz, DMSO-d_6) δ 9.93 (s, 1H), 8.27 (s, 2H); ESIMS *m/z* 297.0 ($[M]^+$).

Example 144: Preparation of 4-Bromo-3,5-dichlorobenzaldehyde



Step 1. Methyl 4-amino-3, 5-dichlorobenzoate: conc. H_2SO_4 (2.5 mL, 97.04 mmol) was added drop wise to a stirred solution of 4-amino-3,5-dichlorobenzoic acid (10.0 g, 48.54 mmol) in MeOH (150 mL) at 0 °C and the reaction mixture was then stirred at 80 °C for 8 h. The volatiles were evaporated; ice cold water was added to the residue and which was then extracted with EtOAc. The combined organic layers were washed with brine, dried (Na_2SO_4), filtered and concentrated under reduced pressure to afford the title compound as a white solid (7.5 g, 70%): ^{1}H NMR (300 MHz, DMSO-d_6) δ 8.05 (s, 2H), 3.96 (s, 3H); ESIMS *m/z* 282 ($[M]^+$); IR (KBr): 1733, 762, 514 cm^{-1}.


Step 2. Methyl 4-bromo-3, 5-dichlorobenzoate: $CuBr_2$ (7.5 g, 34.08 mmol) in MeCN (50 mL) was stirred at 80 °C for 30 min. To this solution *tert*-butylnitrite (6.5 mL, 54.55 mmol) was added dropwise at the same temperature and the mixture was stirred for another 10 min. Methyl 4-amino-3,5-dichlorobenzoate in MeCN (30 mL) was added dropwise to the reaction mixture which was then stirred at 80 °C for 30 min. The reaction mixture was brought to ambient temperature. Aqueous ammonia solution (20 mL) was added and extracted with petroleum ether. The organic layer was washed with brine followed by water. The organic solution was dried (Na_2SO_4), filtered and concentrated to afford the title compound as an off white solid (7.5 g, 77 %): ^{1}H NMR (300 MHz, DMSO-d_6) δ 8.02 (s, 2H), 3.94 (s, 3H); ESIMS *m/z* 282 ($[M]^+$); IR (thin film) 1733, 762, 514 cm^{-1}.

Step 3. (4-Bromo-3,5-dichlorophenyl)methanol: DIBAL-H (1M in toluene, 66 mL, and 66.0 mmol) was added dropwise to a stirred solution of methyl 4-bromo-3, 5-dichlorobenzoate (7.5 g, 26.0 mmol) in THF (50 mL) at -78 °C. The reaction mixture was brought to ambient temperature and stirred for 6 h. The reaction mixture was poured into ice-water and extracted with CH_2Cl_2. The organic layer was washed with brine followed by water, dried (Na_2SO_4), filtered and concentrated to afford a mixture of (4-bromo-3,5-

dichlorophenyl)methanol and 4-bromo-3,5-dichlorobenzaldehyde (6.0 g) as an off white solid which was taken to next step without purification.


Step 4. 4-Bromo-3, 5-dichlorobenzaldehyde: PCC (7.5 g, 35.16 mmol) was added in one portion to a stirred solution containing a mixture of (4-bromo-3,5-dichlorophenyl)methanol and 4-bromo-3,5-dichlorobenzaldehyde (6.0 g) in $CHCl_3$ (40 mL) at ambient temperature and the reaction mixture was stirred overnight. The reaction mixture was filtered through celite. The celite pad was washed with $CHCl_3$. The filtrate was concentrated to afford the title compound as an off white solid (3.5 g, 67%): mp 125-128 °C; ^{1}H NMR (300 MHz, DMSO-d_6) δ 9.96 (s, 1H), 8.10 (s, 2H); ESIMS *m/z* 252 ([M]$^+$).

Example 145: 3-Chloro-5-ethylbenzaldehyde



$PdCl_2$(dppf)(37 mg, 0.046 mmol), potassium phosphate (1.93 g, 9.11 mmol) and triethylborane (1M in hexane, 0.45 g, 4.56 mmol) were added to a solution of 3-bromo-5-chloro-benzaldehyde (1.0 g, 4.56 mmol) in THF (20 mL) at ambient temperature and the mixture was refluxed for 12 h. The reaction mixture was brought to ambient temperature, diluted with EtOAc and washed with water. The organic layer was dried (Na_2SO_4), filtered, concentrated and the residue was purified by column chromatography on silica (100-200 mesh) eluting with 2% EtOAc in petroleum ether to afford the title compound (330 mg, 41%) as a pale yellow liquid: ^{1}H NMR (400 MHz, DMSO-d_6) δ 9.97 (s, 1H), 7.75 (d, J = 1.6 Hz 1H), 7.73 (s, 1H), 7.65 (s, 1H), 2.74 -2.68 (m, 2H), 1.23 (t, J = 7.6 Hz, 3H); ESIMS *m/z* 168.0 ([M]$^+$); IR (thin film) 3071, 1699, 692 cm^{-1}.

Example 146: (*E*)-2-Amino-4-(3-(3,5-dichloro-4-fluorophenyl)-4,4,4-trifluorobut-1-enyl)-*N*-(2-oxo-2-(2,2,2-trifluoroethylamino)ethyl)benzamide



Step 1. (*E*)-Ethyl 4-(3-(3,5-dichloro-4-fluorophenyl)-4,4,4-trifluorobut-1-enyl)-2-nitrobenzoate: 5-(1-Bromo-2,2,2-trifluoroethyl)-1,3-dichloro-2-fluorobenzene (3.5 g, 10.8 mmol), CuCl (54 mg, 0.54 mmol) and 2,2-bipyridyl (169 mg, 1.08 mmol) were added to a stirred solution of ethyl 2-nitro-4-vinylbenzoate (1.2 g, 5.42 mmol) in 1,2–dichlorobenzene

(12 mL) at ambient temperature and the mixture was then stirred at 180 °C for 18 h. The reaction mixture was then cooled to ambient temperature, adsorbed on silica gel and purified by column chromatography eluting with 10% EtOAc in petroleum ether to afford the title compound (1.3 g, 53%) as a brown liquid: ^{1}H NMR (DMSO-d_6, 300 MHz) δ 8.31 (s, 1H), 8.02 - 7.95 (m, 1H), 7.88 - 7.85 (m, 3H), 7.20 (dd, J = 15.9, 9.3 Hz, 1H), 6.91 (d, J = 15.6 Hz, 1H), 4.91- 4.85 (m, 1H), 4.34 - 4.27 (m, 2H), 1.27 (t, J = 6.6 Hz, 3H); ESIMS m/z 463.8 ([M-H]$^-$); IR (KBr) 3439, 2985, 1731, 1251 cm^{-1}.


Step 2. (*E*)-4-(3-(3,5-Dichloro-4-fluorophenyl)-4,4,4-trifluorobut-1-enyl)-2-nitrobenzoic acid: Concentrated HCl (16.0 mL) was added dropwise to a stirred solution of (*E*)-ethyl 4-(3-(3,5-dichloro-4-fluorophenyl)-4,4,4-trifluorobut-1-enyl)-2-nitrobenzoate (800 mg, 1.72 mmol) in 1,4-dioxane (8.0 mL) at 0 °C and the reaction mixture was refluxed for 36 h. The volatiles were evaporated; the residue was diluted with EtOAc and washed with brine and water. The organic layer was dried (Na_2SO_4), filtered, concentrated and the residue was purified by column chromatography on silica (100-200 mesh) eluting with 30% EtOAc in petroleum ether to afford the title compound as a yellow solid (390 mg, 52%): ^{1}H NMR (400MHz, DMSO-d_6) δ 13.9 (bs, 1H), 8.22 (s, 1H), 7.93 - 7.91 (m,1H), 7.86 - 7.84 (m, 3H), 7.16 (dd, J = 15.6, 9.2 Hz, 1H), 6.89 (d, J = 15.6 Hz, 1H), 4.89 - 4.85 (m, 1H). ESIMS m/z 435.9 ([M-H]$^-$); IR (KBr) 3445, 2924, 1708, 1541, 817 cm^{-1}.

Step 3. (*E*)-4-(3-(3,5-Dichloro-4-fluorophenyl)-4,4,4-trifluorobut-1-enyl)-2-nitro-*N*-(2-oxo-2-(2,2,2-trifluoroethylamino)ethyl)benzamide: 2-Amino-*N*-(2,2,2-trifluoroethyl)acetamide hydrochloride (96 mg, 0.35 mmol), PyBOP (165 mg, 0.32 mmol) and DIPEA (0.1 mL, 0.57 mmol) were added to a stirred solution of (*E*)-4-(3-(3,5-dichloro-4-fluorophenyl)-4,4,4-trifluorobut-1-enyl)-2-nitrobenzoic acid (130 mg, 0.29 mmol) in CH_2Cl_2 (3 mL) and the reaction mixture was stirred at ambient temperature for 8h. Water was added to reaction mixture and extracted with CH_2Cl_2. The organic layer was washed with brine, dried (Na_2SO_4), filtered, concentrated and the residue was purified by column chromatography on silica (100-200 mesh) eluting with 30% EtOAc in petroleum ether to afford the title compound as a yellow solid (120 mg, 74%): ^{1}H NMR (300 MHz, DMSO-d_6) δ 9.04 (t, J = 5.7 Hz, 1H), 8.60 (t, J = 6.0 Hz, 1H), 8.25 (s, 1H), 7.97 - 7.94 (m, 1H), 7.87 (d, J = 6.3 Hz, 2H), 7.69 (d, J = 7.5 Hz, 1H), 7.15 (dd, J= 15.9, 9.3 Hz, 1H), 6.89 (d, J = 15.9 Hz, 1H), 4.88 - 4.83 (m, 1H), 3.98 - 3.89 (m, 4H); ESIMS m/z 575.87 ([M+H]$^+$); IR (KBr) 3430, 2925, 1663, 1168, 832 cm^{-1}.

Step 4. (*E*)-2-Amino-4-(3-(3,5-dichloro-4-fluorophenyl)-4,4,4-trifluorobut-1-enyl)-*N*-(2-oxo-2-(2,2,2-trifluoroethylamino)ethyl)benzamide: Iron powder (81.8 mg, 1.46


mmol) and NH_4Cl (104 mg, 1.94 mmol) was added to a stirred solution of (*E*)-4-(3-(3,5-dichloro-4-fluorophenyl)-4,4,4-trifluorobut-1-enyl)-2-nitro-*N*-(2-oxo-2-(2,2,2-trifluoroethylamino)ethyl)benzamide (280 mg, 0.486 mmol) in EtOH : water (6 mL, 1:1) at ambient temperature and the mixture was stirred at reflux for 90 min. The reaction mixture was cooled to ambient temperature and filtered through celite. The filtrate was concentrated and the residue was dissolved in EtOAc and washed with saturated $NaHCO_3$ solution, brine and water. The organic layer was dried (Na_2SO_4), filtered, concentrated and the residue was purified by column chromatography on silica (10-200 mesh) eluting with 35% EtOAc in petroleum ether to afford the titled compound as a yellow solid (215 mg, 81%).

Example 147: (*E*)-2-Bromo-4-(3-(3,5-dichloro-4-hydroxyphenyl)-4,4,4-trifluorobut-1-enyl)-*N*-(2-oxo-2-(2,2,2-trifluoroethylamino)ethyl)benzamide



DIPEA (0.20 mL, 1.26 mmol), PyBOP (245 mg, 0.47 mmol) and 2-amino-*N*-(2,2,2-trifluoroethyl)acetamide (90 mg, 0.47 mmol) were added to a stirred solution of (*E*)-2-bromo-4-(3-(3,5-dichloro-4-hydroxyphenyl)-4,4,4-trifluorobut-1-enyl)benzoic acid (200 mg, 0.42 mmol, 66% purity) in CH_2Cl_2 (5 mL) at ambient temperature. The resulting mixture was then stirred for 12 h. The reaction mixture was diluted with CH_2Cl_2, washed with 1N HCl, followed by saturated sodium bicarbonate solution, brine solution and water. The organic layer was dried (Na_2SO_4), filtered, concentrated under vacuum. The residue was purified by column chromatography on silica (100-200 mesh) eluting with 30% EtOAc in petroleum ether to give the title compound as light green solid (130 mg, 52 %).


Example 148: Preparation of (*R*)-2-Amino-3-methyl-*N*-(2,2,2-trifluoroethyl)butanamide



A Parr shaker flask charged with (*R*)-benzyl (3-methyl-1-oxo-1-((2,2,2-trifluoroethyl)amino)-butan-2-yl)carbamate (4.95 g, 14.9 mmol) in 100 mL of EtOAc and 25 mL of EtOH was treated with 175 mg of 10% Pd/C. The mixture was placed under 40 psi of hydrogen and shaken for 6 h. An additional 65 mg of 10% Pd/C was added to the reaction mixture which was then placed under 40 psi of hydrogen and shaken for 3.5 h. The reaction solution was then filtered through a pad of celite, concentrated in vacuo and purified by



sublimation (75 to 85 °C, 200-230 millibar) to afford the title compound as a white solid (1.92 g, 65%): mp 43-47 °C; ^{1}H NMR (400 MHz, $CDCl_3$) δ 8.04 - 7.74 (m, 1H), 3.92 (ddd, *J* = 16.0, 9.3, 6.8 Hz, 1H), 3.84 - 3.60 (m, 1H), 3.23 (d, *J* = 3.8 Hz, 1H), 2.23 (ddd, *J* = 10.7, 7.0, 3.5 Hz, 1H), 0.92 (*d*, *J* = 7.0 Hz, 3H), 0.75 (d, *J* = 6.9 Hz, 3H); ^{19}F NMR (376 MHz, $CDCl_3$) δ -72.70.











Example 149: Preparation of (*E*)-4-(3-(3,5-Dichloro-4-fluorophenyl)-4,4,4-trifluorobut-1-enyl)-2-(methylsulfonyl)-*N*-(2-oxo-2-(2,2,2-trifluoroethylamino)ethyl)benzamide























Oxone (320 mg, 0.50 mmol) was added to a stirred solution of (*E*)-4-(3-(3,5-dichloro-4-fluorophenyl)-4,4,4-trifluorobut-1-enyl)-2-(methylthio)-*N*-(2-oxo-2-(2,2,2-trifluoroethylamino)ethyl)benzamide (150 mg, 0.25 mmol) in acetone –water (10 mL, 1:1) at ambient temperature and the reaction mixture was stirred for 18 h. The reaction mixture was then extracted with CH_2Cl_2. The organic layer was washed with brine and water, dried (Na_2SO_4), filtered, concentrated and the residue was triturated with pentane-Et_2O (1:1) to afford the title compound as an off white solid (85 mg, 56%): ^{1}H NMR (300 MHz, DMSO-d_6) δ 9.01 (t, *J* = 6.0, 1H), 8.37 (t, *J* = 6.4 Hz, 1H), 8.08 (s, 1H), 8.01 - 7.96 (m, 1H), 7.88 - 7.86 (m, 2H), 7.64 (d, *J* = 7.6 Hz, 1H), 7.05 (dd, *J* = 16.4, 8.8 Hz,1H), 6.91 (d, *J* = 15.6 Hz, 1H), 4.87 - 4.80 (m, 1H), 3.98 - 3.91 (m, 4H), 3.38 (s, 3H); ESIMS *m/z* 608.85 ([M+H]$^+$); IR (thin film) 337, 416, 721, 164, 768 cm^{-1}.











The following prophetic molecules could be made in accordance with the procedures disclosed in this application:




Compound Number	Structure
P1	
P2	

P3	
P4	
P5	
P6	
P7	
P8	
P9	
P10	
P11	
P12	


P13	
P14	
P15	
P16	
P17	
P18	
P19	
P20	
P21	
P22	

P23	
P24	
P25	
P26	
P27	
P28	
P29	
P30	
P31	
P32	

P33	
P34	
P35	
P36	
P37	
P38	
P39	
P40	
P41	
P42	

P43	
P44	
P45	

The following prophetic molecules could be made in accordance with the procedures disclosed in this application:



5

Cmpd No.	R 1	R2	R3	R4	R6	R8	R10	W 1	(C1-C8) alkyl	W 2	R15
P46	F	F	F	H	CF_3	H	Br	O	CH_2	O	CH_2CF_3
P47	F	F	F	H	CF_3	H	Cl	O	CH_2	O	CH_2CF_3
P48	F	F	F	H	CF_3	H	CF_3	O	CH_2	O	CH_2CF_3
P49	F	F	F	H	CF_3	H	CH_3	O	CH_2	O	CH_2CF_3
P50	F	F	F	H	CF_3	H	Br	O	CH_2	S	CH_2CF_3
P51	F	F	F	H	CF_3	H	Cl	O	CH_2	S	CH_2CF_3
P52	F	F	F	H	CF_3	H	CF_3	O	CH_2	S	CH_2CF_3
P53	F	F	F	H	CF_3	H	CH_3	O	CH_2	S	CH_2CF_3
P54	F	F	F	H	CF_3	H	Br	S	CH_2	O	CH_2CF_3
P55	F	F	F	H	CF_3	H	Cl	S	CH_2	O	CH_2CF_3

P56	F	F	F	H	CF_3	H	CF_3	S	CH_2	O	CH_2CF_3
P57	F	F	F	H	CF_3	H	CH_3	S	CH_2	O	CH_2CF_3
P58	F	F	F	H	CF_3	H	Br	O	CH_2	O	CH_2CHF_2
P59	F	F	F	H	CF_3	H	Cl	O	CH_2	O	CH_2CHF_2
P60	F	F	F	H	CF_3	H	CF_3	O	CH_2	O	CH_2CHF_2
P61	F	F	F	H	CF_3	H	CH_3	O	CH_2	O	CH_2CHF_2
P62	F	F	F	H	CF_3	CF_3	Br	O	CH_2	O	CH_2CF_3
P63	F	F	F	H	CF_3	CF_3	Cl	O	CH_2	O	CH_2CF_3
P64	F	F	F	H	CF_3	CF_3	CF_3	O	CH_2	O	CH_2CF_3
P65	F	F	F	H	CF_3	CF_3	CH_3	O	CH_2	O	CH_2CF_3
P66	F	F	F	H	CF_2CF_3	H	Br	O	CH_2	O	CH_2CF_3
P67	F	F	F	H	CF_2CF_3	H	Cl	O	CH_2	O	CH_2CF_3
P68	F	F	F	H	CF_2CF_3	H	CF_3	O	CH_2	O	CH_2CF_3
P69	F	F	F	H	CF_2CF_3	H	CH_3	O	CH_2	O	CH_2CF_3
P70	F	F	F	H	CF_3	H	Br	O	CH_2	O	$CH(CH_3)CF_3$
P71	F	F	F	H	CF_3	H	Cl	O	CH_2	O	$CH(CH_3)CF_3$
P72	F	F	F	H	CF_3	H	CF_3	O	CH_2	O	$CH(CH_3)CF_3$
P73	F	F	F	H	CF_3	H	CH_3	O	CH_2	O	$CH(CH_3)CF_3$
P74	F	F	F	H	CF_3	CF_3	Br	O	$CH(CH_3)$	O	CH_2CF_3
P75	F	F	F	H	CF_3	CF_3	Cl	O	$CH(CH_3)$	O	CH_2CF_3
P76	F	F	F	H	CF_3	CF_3	CF_3	O	$CH(CH_3)$	O	CH_2CF_3
P77	F	F	F	H	CF_3	CF_3	CH_3	O	$CH(CH_3)$	O	CH_2CF_3
P78	F	F	F	H	CF_2CF_3	H	Br	O	$CH(CH_3)$	O	CH_2CF_3
P79	F	F	F	H	CF_2CF_3	H	Cl	O	$CH(CH_3)$	O	CH_2CF_3
P80	F	F	F	H	CF_2CF_3	H	CF_3	O	$CH(CH_3)$	O	CH_2CF_3
P81	F	F	F	H	CF_2CF_3	H	CH_3	O	$CH(CH_3)$	O	CH_2CF_3
P82	F	F	F	H	CF_3	H	Br	O	$CH(CH_3)$	O	CH_2CF_3

P83	F	F	F	H	CF_3	H	Cl	O	$CH(CH_3)$	O	CH_2CF_3
P84	F	F	F	H	CF_3	H	CF_3	O	$CH(CH_3)$	O	CH_2CF_3
P85	F	F	F	H	CF_3	H	CH_3	O	$CH(CH_3)$	O	CH_2CF_3
P86	F	F	F	H	CF_3	H	Br	O	$CH(CH_3)$	S	CH_2CF_3
P87	F	F	F	H	CF_3	H	Cl	O	$CH(CH_3)$	S	CH_2CF_3
P88	F	F	F	H	CF_3	H	CF_3	O	$CH(CH_3)$	S	CH_2CF_3
P89	F	F	F	H	CF_3	H	CH_3	O	$CH(CH_3)$	S	CH_2CF_3
P90	F	F	F	H	CF_3	H	Br	S	$CH(CH_3)$	O	CH_2CF_3
P91	F	F	F	H	CF_3	H	Cl	S	$CH(CH_3)$	O	CH_2CF_3
P92	F	F	F	H	CF_3	H	CF_3	S	$CH(CH_3)$	O	CH_2CF_3
P93	F	F	F	H	CF_3	H	CH_3	S	$CH(CH_3)$	O	CH_2CF_3
P94	F	F	F	H	CF_3	H	Br	O	$CH(CH_3)$	O	CH_2CHF_2
P95	F	F	F	H	CF_3	H	Cl	O	$CH(CH_3)$	O	CH_2CHF_2
P96	F	F	F	H	CF_3	H	CF_3	O	$CH(CH_3)$	O	CH_2CHF_2
P97	F	F	F	H	CF_3	H	CH_3	O	$CH(CH_3)$	O	CH_2CHF_2
P98	F	F	F	H	CF_3	H	Br	O	$CH(CH_3)$	O	$CH(CH_3)CF_3$
P99	F	F	F	H	CF_3	H	Cl	O	$CH(CH_3)$	O	$CH(CH_3)CF_3$
P100	F	F	F	H	CF_3	H	CF_3	O	$CH(CH_3)$	O	$CH(CH_3)CF_3$
P101	F	F	F	H	CF_3	H	CH_3	O	$CH(CH_3)$	O	$CH(CH_3)CF_3$
P102	F	F	F	H	CF_3	H	Br	O	$CH(CH_3)$	O	$CH_2CH_2CF_3$
P103	F	F	F	H	CF_3	H	Cl	O	$CH(CH_3)$	O	$CH_2CH_2CF_3$
P104	F	F	F	H	CF_3	H	CF_3	O	$CH(CH_3)$	O	$CH_2CH_2CF_3$
P105	F	F	F	H	CF_3	H	CH_3	O	$CH(CH_3)$	O	$CH_2CH_2CF_3$
P106	F	F	F	H	CF_3	H	Br	O	$CH(CH_2CH_3)$	O	CH_2CF_3
P107	F	F	F	H	CF_3	H	Cl	O	$CH(CH_2CH_3)$	O	CH_2CF_3
P108	F	F	F	H	CF_3	H	CF_3	O	$CH(CH_2CH_3)$	O	CH_2CF_3
P109	F	F	F	H	CF_3	H	CH_3	O	$CH(CH_2CH_3)$	O	CH_2CF_3

P110	F	F	F	H	CF_3	H	Br	O	$C(CH_3)_2$	O	CH_2CF_3
P111	F	F	F	H	CF_3	H	Cl	O	$C(CH_3)_2$	O	CH_2CF_3
P112	F	F	F	H	CF_3	H	CF_3	O	$C(CH_3)_2$	O	CH_2CF_3
P113	F	F	F	H	CF_3	H	CH_3	O	$C(CH_3)_2$	O	CH_2CF_3
P114	F	F	F	H	CF_3	H	Br	O	CH_2CH_2	O	CH_2CF_3
P115	F	F	F	H	CF_3	H	Cl	O	CH_2CH_2	O	CH_2CF_3
P116	F	F	F	H	CF_3	H	CF_3	O	CH_2CH_2	O	CH_2CF_3
P117	F	F	F	H	CF_3	H	CH_3	O	CH_2CH_2	O	CH_2CF_3
P118	Cl	Cl	H	Cl	CF_3	H	Br	O	CH_2	O	CH_2CF_3
P119	Cl	Cl	H	Cl	CF_3	H	Cl	O	CH_2	O	CH_2CF_3
P120	Cl	Cl	H	Cl	CF_3	H	CF_3	O	CH_2	O	CH_2CF_3
P121	Cl	Cl	H	Cl	CF_3	H	CH_3	O	CH_2	O	CH_2CF_3
P122	Cl	Cl	H	Cl	CF_3	H	Br	O	CH_2	S	CH_2CF_3
P123	Cl	Cl	H	Cl	CF_3	H	Cl	O	CH_2	S	CH_2CF_3
P124	Cl	Cl	H	Cl	CF_3	H	CF_3	O	CH_2	S	CH_2CF_3
P125	Cl	Cl	H	Cl	CF_3	H	CH_3	O	CH_2	S	CH_2CF_3
P126	Cl	Cl	H	Cl	CF_3	H	Br	S	CH_2	O	CH_2CF_3
P127	Cl	Cl	H	Cl	CF_3	H	Cl	S	CH_2	O	CH_2CF_3
P128	Cl	Cl	H	Cl	CF_3	H	CF_3	S	CH_2	O	CH_2CF_3
P129	Cl	Cl	H	Cl	CF_3	H	CH_3	S	CH_2	O	CH_2CF_3
P130	Cl	Cl	H	Cl	CF_3	H	Br	O	CH_2	O	CH_2CHF_2
P131	Cl	Cl	H	Cl	CF_3	H	Cl	O	CH_2	O	CH_2CHF_2
P132	Cl	Cl	H	Cl	CF_3	H	CF_3	O	CH_2	O	CH_2CHF_2
P133	Cl	Cl	H	Cl	CF_3	H	CH_3	O	CH_2	O	CH_2CHF_2
P134	Cl	Cl	H	Cl	CF_3	CF_3	Br	O	CH_2	O	CH_2CF_3
P135	Cl	Cl	H	Cl	CF_3	CF_3	Cl	O	CH_2	O	CH_2CF_3
P136	Cl	Cl	H	Cl	CF_3	CF_3	CF_3	O	CH_2	O	CH_2CF_3

P137	Cl	Cl	H	Cl	CF_3	CF_3	CH_3	O	CH_2	O	CH_2CF_3
P138	Cl	Cl	H	Cl	CF_2CF_3	H	Br	O	CH_2	O	CH_2CF_3
P139	Cl	Cl	H	Cl	CF_2CF_3	H	Cl	O	CH_2	O	CH_2CF_3
P140	Cl	Cl	H	Cl	CF_2CF_3	H	CF_3	O	CH_2	O	CH_2CF_3
P141	Cl	Cl	H	Cl	CF_2CF_3	H	CH_3	O	CH_2	O	CH_2CF_3
P142	Cl	Cl	H	Cl	CF_3	H	Br	O	CH_2	O	$CH(CH_3)CF_3$
P143	Cl	Cl	H	Cl	CF_3	H	Cl	O	CH_2	O	$CH(CH_3)CF_3$
P144	Cl	Cl	H	Cl	CF_3	H	CF_3	O	CH_2	O	$CH(CH_3)CF_3$
P145	Cl	Cl	H	Cl	CF_3	H	CH_3	O	CH_2	O	$CH(CH_3)CF_3$
P146	Cl	Cl	H	Cl	CF_3	CF_3	Br	O	$CH(CH_3)$	O	CH_2CF_3
P147	Cl	Cl	H	Cl	CF_3	CF_3	Cl	O	$CH(CH_3)$	O	CH_2CF_3
P148	Cl	Cl	H	Cl	CF_3	CF_3	CF_3	O	$CH(CH_3)$	O	CH_2CF_3
P149	Cl	Cl	H	Cl	CF_3	CF_3	CH_3	O	$CH(CH_3)$	O	CH_2CF_3
P150	Cl	Cl	H	Cl	CF_2CF_3	H	Br	O	$CH(CH_3)$	O	CH_2CF_3
P151	Cl	Cl	H	Cl	CF_2CF_3	H	Cl	O	$CH(CH_3)$	O	CH_2CF_3
P152	Cl	Cl	H	Cl	CF_2CF_3	H	CF_3	O	$CH(CH_3)$	O	CH_2CF_3
P153	Cl	Cl	H	Cl	CF_2CF_3	H	CH_3	O	$CH(CH_3)$	O	CH_2CF_3
P154	Cl	Cl	H	Cl	CF_3	H	Br	O	$CH(CH_3)$	O	CH_2CF_3
P155	Cl	Cl	H	Cl	CF_3	H	Cl	O	$CH(CH_3)$	O	CH_2CF_3
P156	Cl	Cl	H	Cl	CF_3	H	CF_3	O	$CH(CH_3)$	O	CH_2CF_3
P157	Cl	Cl	H	Cl	CF_3	H	CH_3	O	$CH(CH_3)$	O	CH_2CF_3
P158	Cl	Cl	H	Cl	CF_3	H	Br	O	$CH(CH_3)$	S	CH_2CF_3
P159	Cl	Cl	H	Cl	CF_3	H	Cl	O	$CH(CH_3)$	S	CH_2CF_3
P160	Cl	Cl	H	Cl	CF_3	H	CF_3	O	$CH(CH_3)$	S	CH_2CF_3
P161	Cl	Cl	H	Cl	CF_3	H	CH_3	O	$CH(CH_3)$	S	CH_2CF_3
P162	Cl	Cl	H	Cl	CF_3	H	Br	S	$CH(CH_3)$	O	CH_2CF_3
P163	Cl	Cl	H	Cl	CF_3	H	Cl	S	$CH(CH_3)$	O	CH_2CF_3

P164	Cl	Cl	H	Cl	CF_3	H	CF_3	S	$CH(CH_3)$	O	CH_2CF_3
P165	Cl	Cl	H	Cl	CF_3	H	CH_3	S	$CH(CH_3)$	O	CH_2CF_3
P166	Cl	Cl	H	Cl	CF_3	H	Br	O	$CH(CH_3)$	O	CH_2CHF_2
P167	Cl	Cl	H	Cl	CF_3	H	Cl	O	$CH(CH_3)$	O	CH_2CHF_2
P168	Cl	Cl	H	Cl	CF_3	H	CF_3	O	$CH(CH_3)$	O	CH_2CHF_2
P169	Cl	Cl	H	Cl	CF_3	H	CH_3	O	$CH(CH_3)$	O	CH_2CHF_2
P170	Cl	Cl	H	Cl	CF_3	H	Br	O	$CH(CH_3)$	O	$CH(CH_3)CF_3$
P171	Cl	Cl	H	Cl	CF_3	H	Cl	O	$CH(CH_3)$	O	$CH(CH_3)CF_3$
P172	Cl	Cl	H	Cl	CF_3	H	CF_3	O	$CH(CH_3)$	O	$CH(CH_3)CF_3$
P173	Cl	Cl	H	Cl	CF_3	H	CH_3	O	$CH(CH_3)$	O	$CH(CH_3)CF_3$
P174	Cl	Cl	H	Cl	CF_3	H	Br	O	$CH(CH_3)$	O	$CH_2CH_2CF_3$
P175	Cl	Cl	H	Cl	CF_3	H	Cl	O	$CH(CH_3)$	O	$CH_2CH_2CF_3$
P176	Cl	Cl	H	Cl	CF_3	H	CF_3	O	$CH(CH_3)$	O	$CH_2CH_2CF_3$
P177	Cl	Cl	H	Cl	CF_3	H	CH_3	O	$CH(CH_3)$	O	$CH_2CH_2CF_3$
P178	Cl	Cl	H	Cl	CF_3	H	Br	O	$CH(CH_2CH_3)$	O	CH_2CF_3
P179	Cl	Cl	H	Cl	CF_3	H	Cl	O	$CH(CH_2CH_3)$	O	CH_2CF_3
P180	Cl	Cl	H	Cl	CF_3	H	CF_3	O	$CH(CH_2CH_3)$	O	CH_2CF_3
P181	Cl	Cl	H	Cl	CF_3	H	CH_3	O	$CH(CH_2CH_3)$	O	CH_2CF_3
P182	Cl	Cl	H	Cl	CF_3	H	Br	O	$C(CH_3)_2$	O	CH_2CF_3
P183	Cl	Cl	H	Cl	CF_3	H	Cl	O	$C(CH_3)_2$	O	CH_2CF_3
P184	Cl	Cl	H	Cl	CF_3	H	CF_3	O	$C(CH_3)_2$	O	CH_2CF_3
P185	Cl	Cl	H	Cl	CF_3	H	CH_3	O	$C(CH_3)_2$	O	CH_2CF_3
P186	Cl	Cl	H	Cl	CF_3	H	Br	O	CH_2CH_2	O	CH_2CF_3
P187	Cl	Cl	H	Cl	CF_3	H	Cl	O	CH_2CH_2	O	CH_2CF_3
P188	Cl	Cl	H	Cl	CF_3	H	CF_3	O	CH_2CH_2	O	CH_2CF_3
P189	Cl	Cl	H	Cl	CF_3	H	CH_3	O	CH_2CH_2	O	CH_2CF_3

P190	H	H	H	OCF_3	CF_3	H	Br	O	CH_2	O	CH_2CF_3
P191	H	H	H	OCF_3	CF_3	H	Cl	O	CH_2	O	CH_2CF_3
P192	H	H	H	OCF_3	CF_3	H	CF_3	O	CH_2	O	CH_2CF_3
P193	H	H	H	OCF_3	CF_3	H	CH_3	O	CH_2	O	CH_2CF_3
P194	H	H	H	OCF_3	CF_3	H	Br	O	CH_2	S	CH_2CF_3
P195	H	H	H	OCF_3	CF_3	H	Cl	O	CH_2	S	CH_2CF_3
P196	H	H	H	OCF_3	CF_3	H	CF_3	O	CH_2	S	CH_2CF_3
P197	H	H	H	OCF_3	CF_3	H	CH_3	O	CH_2	S	CH_2CF_3
P198	H	H	H	OCF_3	CF_3	H	Br	S	CH_2	O	CH_2CF_3
P199	H	H	H	OCF_3	CF_3	H	Cl	S	CH_2	O	CH_2CF_3
P200	H	H	H	OCF_3	CF_3	H	CF_3	S	CH_2	O	CH_2CF_3
P201	H	H	H	OCF_3	CF_3	H	CH_3	S	CH_2	O	CH_2CF_3
P202	H	H	H	OCF_3	CF_3	H	Br	O	CH_2	O	CH_2CHF_2
P203	H	H	H	OCF_3	CF_3	H	Cl	O	CH_2	O	CH_2CHF_2
P204	H	H	H	OCF_3	CF_3	H	CF_3	O	CH_2	O	CH_2CHF_2
P205	H	H	H	OCF_3	CF_3	H	CH_3	O	CH_2	O	CH_2CHF_2
P206	H	H	H	OCF_3	CF_3	CF_3	Br	O	CH_2	O	CH_2CF_3

P207	H	H	H	OCF_3	CF_3	CF_3	Cl	O	CH_2	O	CH_2CF_3
P208	H	H	H	OCF_3	CF_3	CF_3	CF_3	O	CH_2	O	CH_2CF_3
P209	H	H	H	OCF_3	CF_3	CF_3	CH_3	O	CH_2	O	CH_2CF_3
P210	H	H	H	OCF_3	CF_2CF_3	H	Br	O	CH_2	O	CH_2CF_3
P211	H	H	H	OCF_3	CF_2CF_3	H	Cl	O	CH_2	O	CH_2CF_3
P212	H	H	H	OCF_3	CF_2CF_3	H	CF_3	O	CH_2	O	CH_2CF_3
P213	H	H	H	OCF_3	CF_2CF_3	H	CH_3	O	CH_2	O	CH_2CF_3
P214	H	H	H	OCF_3	CF_3	H	Br	O	CH_2	O	$CH(CH_3)CF_3$
P215	H	H	H	OCF_3	CF_3	H	Cl	O	CH_2	O	$CH(CH_3)CF_3$
P216	H	H	H	OCF_3	CF_3	H	CF_3	O	CH_2	O	$CH(CH_3)CF_3$
P217	H	H	H	OCF_3	CF_3	H	CH_3	O	CH_2	O	$CH(CH_3)CF_3$
P218	H	H	H	OCF_3	CF_3	CF_3	Br	O	$CH(CH_3)$	O	CH_2CF_3
P219	H	H	H	OCF_3	CF_3	CF_3	Cl	O	$CH(CH_3)$	O	CH_2CF_3
P220	H	H	H	OCF_3	CF_3	CF_3	CF_3	O	$CH(CH_3)$	O	CH_2CF_3
P221	H	H	H	OCF_3	CF_3	CF_3	CH_3	O	$CH(CH_3)$	O	CH_2CF_3
P222	H	H	H	OCF_3	CF_2CF_3	H	Br	O	$CH(CH_3)$	O	CH_2CF_3
P223	H	H	H	OCF_3	CF_2CF_3	H	Cl	O	$CH(CH_3)$	O	CH_2CF_3

P224	H	H	H	OCF_3	CF_2CF_3	H	CF_3	O	CH(CH$_3$)	O	CH_2CF_3
P225	H	H	H	OCF_3	CF_2CF_3	H	CH_3	O	CH(CH$_3$)	O	CH_2CF_3
P226	H	H	H	OCF_3	CF_3	H	Br	O	CH(CH$_3$)	O	CH_2CF_3
P227	H	H	H	OCF_3	CF_3	H	Cl	O	CH(CH$_3$)	O	CH_2CF_3
P228	H	H	H	OCF_3	CF_3	H	CF_3	O	CH(CH$_3$)	O	CH_2CF_3
P229	H	H	H	OCF_3	CF_3	H	CH_3	O	CH(CH$_3$)	O	CH_2CF_3
P230	H	H	H	OCF_3	CF_3	H	Br	O	CH(CH$_3$)	S	CH_2CF_3
P231	H	H	H	OCF_3	CF_3	H	Cl	O	CH(CH$_3$)	S	CH_2CF_3
P232	H	H	H	OCF_3	CF_3	H	CF_3	O	CH(CH$_3$)	S	CH_2CF_3
P233	H	H	H	OCF_3	CF_3	H	CH_3	O	CH(CH$_3$)	S	CH_2CF_3
P234	H	H	H	OCF_3	CF_3	H	Br	S	CH(CH$_3$)	O	CH_2CF_3
P235	H	H	H	OCF_3	CF_3	H	Cl	S	CH(CH$_3$)	O	CH_2CF_3
P236	H	H	H	OCF_3	CF_3	H	CF_3	S	CH(CH$_3$)	O	CH_2CF_3
P237	H	H	H	OCF_3	CF_3	H	CH_3	S	CH(CH$_3$)	O	CH_2CF_3
P238	H	H	H	OCF_3	CF_3	H	Br	O	CH(CH$_3$)	O	CH_2CHF_2
P239	H	H	H	OCF_3	CF_3	H	Cl	O	CH(CH$_3$)	O	CH_2CHF_2
P240	H	H	H	OCF_3	CF_3	H	CF_3	O	CH(CH$_3$)	O	CH_2CHF_2

P241	H	H	H	OCF_3	CF_3	H	CH_3	O	$CH(CH_3)$	O	CH_2CHF_2
P242	H	H	H	OCF_3	CF_3	H	Br	O	$CH(CH_3)$	O	$CH(CH_3)CF_3$
P243	H	H	H	OCF_3	CF_3	H	Cl	O	$CH(CH_3)$	O	$CH(CH_3)CF_3$
P244	H	H	H	OCF_3	CF_3	H	CF_3	O	$CH(CH_3)$	O	$CH(CH_3)CF_3$
P245	H	H	H	OCF_3	CF_3	H	CH_3	O	$CH(CH_3)$	O	$CH(CH_3)CF_3$
P246	H	H	H	OCF_3	CF_3	H	Br	O	$CH(CH_3)$	O	$CH_2CH_2CF_3$
P247	H	H	H	OCF_3	CF_3	H	Cl	O	$CH(CH_3)$	O	$CH_2CH_2CF_3$
P248	H	H	H	OCF_3	CF_3	H	CF_3	O	$CH(CH_3)$	O	$CH_2CH_2CF_3$
P249	H	H	H	OCF_3	CF_3	H	CH_3	O	$CH(CH_3)$	O	$CH_2CH_2CF_3$
P250	H	H	H	OCF_3	CF_3	H	Br	O	$CH(CH_2CH_3)$	O	CH_2CF_3
P251	H	H	H	OCF_3	CF_3	H	Cl	O	$CH(CH_2CH_3)$	O	CH_2CF_3
P252	H	H	H	OCF_3	CF_3	H	CF_3	O	$CH(CH_2CH_3)$	O	CH_2CF_3
P253	H	H	H	OCF_3	CF_3	H	CH_3	O	$CH(CH_2CH_3)$	O	CH_2CF_3
P254	H	H	H	OCF_3	CF_3	H	Br	O	$C(CH_3)_2$	O	CH_2CF_3
P255	H	H	H	OCF_3	CF_3	H	Cl	O	$C(CH_3)_2$	O	CH_2CF_3
P256	H	H	H	OCF_3	CF_3	H	CF_3	O	$C(CH_3)_2$	O	CH_2CF_3
P257	H	H	H	OCF_3	CF_3	H	CH_3	O	$C(CH_3)_2$	O	CH_2CF_3

P258	H	H	H	OCF_3	CF_3	H	Br	O	CH_2CH_2	O	CH_2CF_3
P259	H	H	H	OCF_3	CF_3	H	Cl	O	CH_2CH_2	O	CH_2CF_3
P260	H	H	H	OCF_3	CF_3	H	CF_3	O	CH_2CH_2	O	CH_2CF_3
P261	H	H	H	OCF_3	CF_3	H	CH_3	O	CH_2CH_2	O	CH_2CF_3
P262	H	F	H	Br	CF_3	H	Br	O	CH_2	O	CH_2CF_3
P263	H	F	H	Br	CF_3	H	Cl	O	CH_2	O	CH_2CF_3
P264	H	F	H	Br	CF_3	H	CF_3	O	CH_2	O	CH_2CF_3
P265	H	F	H	Br	CF_3	H	CH_3	O	CH_2	O	CH_2CF_3
P266	H	F	H	Br	CF_3	H	Br	O	CH_2	S	CH_2CF_3
P267	H	F	H	Br	CF_3	H	Cl	O	CH_2	S	CH_2CF_3
P268	H	F	H	Br	CF_3	H	CF_3	O	CH_2	S	CH_2CF_3
P269	H	F	H	Br	CF_3	H	CH_3	O	CH_2	S	CH_2CF_3
P270	H	F	H	Br	CF_3	H	Br	S	CH_2	O	CH_2CF_3
P271	H	F	H	Br	CF_3	H	Cl	S	CH_2	O	CH_2CF_3
P272	H	F	H	Br	CF_3	H	CF_3	S	CH_2	O	CH_2CF_3
P273	H	F	H	Br	CF_3	H	CH_3	S	CH_2	O	CH_2CF_3
P274	H	F	H	Br	CF_3	H	Br	O	CH_2	O	CH_2CHF_2
P275	H	F	H	Br	CF_3	H	Cl	O	CH_2	O	CH_2CHF_2
P276	H	F	H	Br	CF_3	H	CF_3	O	CH_2	O	CH_2CHF_2
P277	H	F	H	Br	CF_3	H	CH_3	O	CH_2	O	CH_2CHF_2
P278	H	F	H	Br	CF_3	CF_3	Br	O	CH_2	O	CH_2CF_3
P279	H	F	H	Br	CF_3	CF_3	Cl	O	CH_2	O	CH_2CF_3
P280	H	F	H	Br	CF_3	CF_3	CF_3	O	CH_2	O	CH_2CF_3
P281	H	F	H	Br	CF_3	CF_3	CH_3	O	CH_2	O	CH_2CF_3
P282	H	F	H	Br	CF_2CF_3	H	Br	O	CH_2	O	CH_2CF_3

P283	H	F	H	Br	CF_2CF_3	H	Cl	O	CH_2	O	CH_2CF_3
P284	H	F	H	Br	CF_2CF_3	H	CF_3	O	CH_2	O	CH_2CF_3
P285	H	F	H	Br	CF_2CF_3	H	CH_3	O	CH_2	O	CH_2CF_3
P286	H	F	H	Br	CF_3	H	Br	O	CH_2	O	$CH(CH_3)CF_3$
P287	H	F	H	Br	CF_3	H	Cl	O	CH_2	O	$CH(CH_3)CF_3$
P288	H	F	H	Br	CF_3	H	CF_3	O	CH_2	O	$CH(CH_3)CF_3$
P289	H	F	H	Br	CF_3	H	CH_3	O	CH_2	O	$CH(CH_3)CF_3$
P290	H	F	H	Br	CF_3	CF_3	Br	O	$CH(CH_3)$	O	CH_2CF_3
P291	H	F	H	Br	CF_3	CF_3	Cl	O	$CH(CH_3)$	O	CH_2CF_3
P292	H	F	H	Br	CF_3	CF_3	CF_3	O	$CH(CH_3)$	O	CH_2CF_3
P293	H	F	H	Br	CF_3	CF_3	CH_3	O	$CH(CH_3)$	O	CH_2CF_3
P294	H	F	H	Br	CF_2CF_3	H	Br	O	$CH(CH_3)$	O	CH_2CF_3
P295	H	F	H	Br	CF_2CF_3	H	Cl	O	$CH(CH_3)$	O	CH_2CF_3
P296	H	F	H	Br	CF_2CF_3	H	CF_3	O	$CH(CH_3)$	O	CH_2CF_3
P297	H	F	H	Br	CF_2CF_3	H	CH_3	O	$CH(CH_3)$	O	CH_2CF_3
P298	H	F	H	Br	CF_3	H	Br	O	$CH(CH_3)$	O	CH_2CF_3
P299	H	F	H	Br	CF_3	H	Cl	O	$CH(CH_3)$	O	CH_2CF_3
P300	H	F	H	Br	CF_3	H	CF_3	O	$CH(CH_3)$	O	CH_2CF_3
P301	H	F	H	Br	CF_3	H	CH_3	O	$CH(CH_3)$	O	CH_2CF_3
P302	H	F	H	Br	CF_3	H	Br	O	$CH(CH_3)$	S	CH_2CF_3
P303	H	F	H	Br	CF_3	H	Cl	O	$CH(CH_3)$	S	CH_2CF_3
P304	H	F	H	Br	CF_3	H	CF_3	O	$CH(CH_3)$	S	CH_2CF_3
P305	H	F	H	Br	CF_3	H	CH_3	O	$CH(CH_3)$	S	CH_2CF_3
P306	H	F	H	Br	CF_3	H	Br	S	$CH(CH_3)$	O	CH_2CF_3
P307	H	F	H	Br	CF_3	H	Cl	S	$CH(CH_3)$	O	CH_2CF_3
P308	H	F	H	Br	CF_3	H	CF_3	S	$CH(CH_3)$	O	CH_2CF_3
P309	H	F	H	Br	CF_3	H	CH_3	S	$CH(CH_3)$	O	CH_2CF_3

P310	H	F	H	Br	CF_3	H	Br	O	$CH(CH_3)$	O	CH_2CHF_2
P311	H	F	H	Br	CF_3	H	Cl	O	$CH(CH_3)$	O	CH_2CHF_2
P312	H	F	H	Br	CF_3	H	CF_3	O	$CH(CH_3)$	O	CH_2CHF_2
P313	H	F	H	Br	CF_3	H	CH_3	O	$CH(CH_3)$	O	CH_2CHF_2
P314	H	F	H	Br	CF_3	H	Br	O	$CH(CH_3)$	O	$CH(CH_3)CF_3$
P315	H	F	H	Br	CF_3	H	Cl	O	$CH(CH_3)$	O	$CH(CH_3)CF_3$
P316	H	F	H	Br	CF_3	H	CF_3	O	$CH(CH_3)$	O	$CH(CH_3)CF_3$
P317	H	F	H	Br	CF_3	H	CH_3	O	$CH(CH_3)$	O	$CH(CH_3)CF_3$
P318	H	F	H	Br	CF_3	H	Br	O	$CH(CH_3)$	O	$CH_2CH_2CF_3$
P319	H	F	H	Br	CF_3	H	Cl	O	$CH(CH_3)$	O	$CH_2CH_2CF_3$
P320	H	F	H	Br	CF_3	H	CF_3	O	$CH(CH_3)$	O	$CH_2CH_2CF_3$
P321	H	F	H	Br	CF_3	H	CH_3	O	$CH(CH_3)$	O	$CH_2CH_2CF_3$
P322	H	F	H	Br	CF_3	H	Br	O	$CH(CH_2CH_3)$	O	CH_2CF_3
P323	H	F	H	Br	CF_3	H	Cl	O	$CH(CH_2CH_3)$	O	CH_2CF_3
P324	H	F	H	Br	CF_3	H	CF_3	O	$CH(CH_2CH_3)$	O	CH_2CF_3
P325	H	F	H	Br	CF_3	H	CH_3	O	$CH(CH_2CH_3)$	O	CH_2CF_3
P326	H	F	H	Br	CF_3	H	Br	O	$C(CH_3)_2$	O	CH_2CF_3
P327	H	F	H	Br	CF_3	H	Cl	O	$C(CH_3)_2$	O	CH_2CF_3
P328	H	F	H	Br	CF_3	H	CF_3	O	$C(CH_3)_2$	O	CH_2CF_3
P329	H	F	H	Br	CF_3	H	CH_3	O	$C(CH_3)_2$	O	CH_2CF_3
P330	H	F	H	Br	CF_3	H	Br	O	CH_2CH_2	O	CH_2CF_3
P331	H	F	H	Br	CF_3	H	Cl	O	CH_2CH_2	O	CH_2CF_3
P332	H	F	H	Br	CF_3	H	CF_3	O	CH_2CH_2	O	CH_2CF_3
P333	H	F	H	Br	CF_3	H	CH_3	O	CH_2CH_2	O	CH_2CF_3
P334	H	CH_3	Cl	H	CF_3	H	Br	O	CH_2	O	CH_2CF_3
P335	H	CH_3	Cl	H	CF_3	H	Cl	O	CH_2	O	CH_2CF_3
P336	H	CH_3	Cl	H	CF_3	H	CF_3	O	CH_2	O	CH_2CF_3

P337	H	CH_3	Cl	H	CF_3	H	CH_3	O	CH_2	O	CH_2CF_3
P338	H	CH_3	Cl	H	CF_3	H	Br	O	CH_2	S	CH_2CF_3
P339	H	CH_3	Cl	H	CF_3	H	Cl	O	CH_2	S	CH_2CF_3
P340	H	CH_3	Cl	H	CF_3	H	CF_3	O	CH_2	S	CH_2CF_3
P341	H	CH_3	Cl	H	CF_3	H	CH_3	O	CH_2	S	CH_2CF_3
P342	H	CH_3	Cl	H	CF_3	H	Br	S	CH_2	O	CH_2CF_3
P343	H	CH_3	Cl	H	CF_3	H	Cl	S	CH_2	O	CH_2CF_3
P344	H	CH_3	Cl	H	CF_3	H	CF_3	S	CH_2	O	CH_2CF_3
P345	H	CH_3	Cl	H	CF_3	H	CH_3	S	CH_2	O	CH_2CF_3
P346	H	CH_3	Cl	H	CF_3	H	Br	O	CH_2	O	CH_2CHF_2
P347	H	CH_3	Cl	H	CF_3	H	Cl	O	CH_2	O	CH_2CHF_2
P348	H	CH_3	Cl	H	CF_3	H	CF_3	O	CH_2	O	CH_2CHF_2
P349	H	CH_3	Cl	H	CF_3	H	CH_3	O	CH_2	O	CH_2CHF_2
P350	H	CH_3	Cl	H	CF_3	CF_3	Br	O	CH_2	O	CH_2CF_3
P351	H	CH_3	Cl	H	CF_3	CF_3	Cl	O	CH_2	O	CH_2CF_3
P352	H	CH_3	Cl	H	CF_3	CF_3	CF_3	O	CH_2	O	CH_2CF_3
P353	H	CH_3	Cl	H	CF_3	CF_3	CH_3	O	CH_2	O	CH_2CF_3
P354	H	CH_3	Cl	H	CF_2CF_3	H	Br	O	CH_2	O	CH_2CF_3
P355	H	CH_3	Cl	H	CF_2CF_3	H	Cl	O	CH_2	O	CH_2CF_3
P356	H	CH_3	Cl	H	CF_2CF_3	H	CF_3	O	CH_2	O	CH_2CF_3
P357	H	CH_3	Cl	H	CF_2CF_3	H	CH_3	O	CH_2	O	CH_2CF_3
P358	H	CH_3	Cl	H	CF_3	H	Br	O	CH_2	O	$CH(CH_3)CF_3$
P359	H	CH_3	Cl	H	CF_3	H	Cl	O	CH_2	O	$CH(CH_3)CF_3$
P360	H	CH_3	Cl	H	CF_3	H	CF_3	O	CH_2	O	$CH(CH_3)CF_3$
P361	H	CH_3	Cl	H	CF_3	H	CH_3	O	CH_2	O	$CH(CH_3)CF_3$
P362	H	CH_3	Cl	H	CF_3	CF_3	Br	O	$CH(CH_3)$	O	CH_2CF_3
P363	H	CH_3	Cl	H	CF_3	CF_3	Cl	O	$CH(CH_3)$	O	CH_2CF_3

P364	H	CH_3	Cl	H	CF_3	CF_3	CF_3	O	$CH(CH_3)$	O	CH_2CF_3
P365	H	CH_3	Cl	H	CF_3	CF_3	CH_3	O	$CH(CH_3)$	O	CH_2CF_3
P366	H	CH_3	Cl	H	CF_2CF_3	H	Br	O	$CH(CH_3)$	O	CH_2CF_3
P367	H	CH_3	Cl	H	CF_2CF_3	H	Cl	O	$CH(CH_3)$	O	CH_2CF_3
P368	H	CH_3	Cl	H	CF_2CF_3	H	CF_3	O	$CH(CH_3)$	O	CH_2CF_3
P369	H	CH_3	Cl	H	CF_2CF_3	H	CH_3	O	$CH(CH_3)$	O	CH_2CF_3
P370	H	CH_3	Cl	H	CF_3	H	Br	O	$CH(CH_3)$	O	CH_2CF_3
P371	H	CH_3	Cl	H	CF_3	H	Cl	O	$CH(CH_3)$	O	CH_2CF_3
P372	H	CH_3	Cl	H	CF_3	H	CF_3	O	$CH(CH_3)$	O	CH_2CF_3
P373	H	CH_3	Cl	H	CF_3	H	CH_3	O	$CH(CH_3)$	O	CH_2CF_3
P374	H	CH_3	Cl	H	CF_3	H	Br	O	$CH(CH_3)$	S	CH_2CF_3
P375	H	CH_3	Cl	H	CF_3	H	Cl	O	$CH(CH_3)$	S	CH_2CF_3
P376	H	CH_3	Cl	H	CF_3	H	CF_3	O	$CH(CH_3)$	S	CH_2CF_3
P377	H	CH_3	Cl	H	CF_3	H	CH_3	O	$CH(CH_3)$	S	CH_2CF_3
P378	H	CH_3	Cl	H	CF_3	H	Br	S	$CH(CH_3)$	O	CH_2CF_3
P379	H	CH_3	Cl	H	CF_3	H	Cl	S	$CH(CH_3)$	O	CH_2CF_3
P380	H	CH_3	Cl	H	CF_3	H	CF_3	S	$CH(CH_3)$	O	CH_2CF_3
P381	H	CH_3	Cl	H	CF_3	H	CH_3	S	$CH(CH_3)$	O	CH_2CF_3
P382	H	CH_3	Cl	H	CF_3	H	Br	O	$CH(CH_3)$	O	CH_2CHF_2
P383	H	CH_3	Cl	H	CF_3	H	Cl	O	$CH(CH_3)$	O	CH_2CHF_2
P384	H	CH_3	Cl	H	CF_3	H	CF_3	O	$CH(CH_3)$	O	CH_2CHF_2
P385	H	CH_3	Cl	H	CF_3	H	CH_3	O	$CH(CH_3)$	O	CH_2CHF_2
P386	H	CH_3	Cl	H	CF_3	H	Br	O	$CH(CH_3)$	O	$CH(CH_3)CF_3$
P387	H	CH_3	Cl	H	CF_3	H	Cl	O	$CH(CH_3)$	O	$CH(CH_3)CF_3$
P388	H	CH_3	Cl	H	CF_3	H	CF_3	O	$CH(CH_3)$	O	$CH(CH_3)CF_3$
P389	H	CH_3	Cl	H	CF_3	H	CH_3	O	$CH(CH_3)$	O	$CH(CH_3)CF_3$
P390	H	CH_3	Cl	H	CF_3	H	Br	O	$CH(CH_3)$	O	$CH_2CH_2CF_3$

P391	H	CH_3	Cl	H	CF_3	H	Cl	O	CH (CH_3)	O	$CH_2CH_2CF_3$
P392	H	CH_3	Cl	H	CF_3	H	CF_3	O	CH(CH_3)	O	$CH_2CH_2CF_3$
P393	H	CH_3	Cl	H	CF_3	H	CH_3	O	CH (CH_3)	O	$CH_2CH_2CF_3$
P394	H	CH_3	Cl	H	CF_3	H	Br	O	CH(CH_2CH_3)	O	CH_2CF_3
P395	H	CH_3	Cl	H	CF_3	H	Cl	O	CH(CH_2CH_3)	O	CH_2CF_3
P396	H	CH_3	Cl	H	CF_3	H	CF_3	O	CH(CH_2CH_3)	O	CH_2CF_3
P397	H	CH_3	Cl	H	CF_3	H	CH_3	O	CH(CH_2CH_3)	O	CH_2CF_3
P398	H	CH_3	Cl	H	CF_3	H	Br	O	$C(CH_3)_2$	O	CH_2CF_3
P399	H	CH_3	Cl	H	CF_3	H	Cl	O	$C(CH_3)_2$	O	CH_2CF_3
P400	H	CH_3	Cl	H	CF_3	H	CF_3	O	$C(CH_3)_2$	O	CH_2CF_3
P401	H	CH_3	Cl	H	CF_3	H	CH_3	O	$C(CH_3)_2$	O	CH_2CF_3
P402	H	CH_3	Cl	H	CF_3	H	Br	O	CH_2CH_2	O	CH_2CF_3
P403	H	CH_3	Cl	H	CF_3	H	Cl	O	CH_2CH_2	O	CH_2CF_3
P404	H	CH_3	Cl	H	CF_3	H	CF_3	O	CH_2CH_2	O	CH_2CF_3
P405	H	CH_3	Cl	H	CF_3	H	CH_3	O	CH_2CH_2	O	CH_2CF_3
P406	H	Cl	CH_3	H	CF_3	H	Br	O	CH_2	O	CH_2CF_3
P407	H	Cl	CH_3	H	CF_3	H	Cl	O	CH_2	O	CH_2CF_3
P408	H	Cl	CH_3	H	CF_3	H	CF_3	O	CH_2	O	CH_2CF_3
P409	H	Cl	CH_3	H	CF_3	H	CH_3	O	CH_2	O	CH_2CF_3
P410	H	Cl	CH_3	H	CF_3	H	Br	O	CH_2	S	CH_2CF_3
P411	H	Cl	CH_3	H	CF_3	H	Cl	O	CH_2	S	CH_2CF_3
P412	H	Cl	CH_3	H	CF_3	H	CF_3	O	CH_2	S	CH_2CF_3
P413	H	Cl	CH_3	H	CF_3	H	CH_3	O	CH_2	S	CH_2CF_3
P414	H	Cl	CH_3	H	CF_3	H	Br	S	CH_2	O	CH_2CF_3
P415	H	Cl	CH_3	H	CF_3	H	Cl	S	CH_2	O	CH_2CF_3
P416	H	Cl	CH_3	H	CF_3	H	CF_3	S	CH_2	O	CH_2CF_3
P417	H	Cl	CH_3	H	CF_3	H	CH_3	S	CH_2	O	CH_2CF_3

P418	H	Cl	CH_3	H	CF_3	H	Br	O	CH_2	O	CH_2CHF_2
P419	H	Cl	CH_3	H	CF_3	H	Cl	O	CH_2	O	CH_2CHF_2
P420	H	Cl	CH_3	H	CF_3	H	CF_3	O	CH_2	O	CH_2CHF_2
P421	H	Cl	CH_3	H	CF_3	H	CH_3	O	CH_2	O	CH_2CHF_2
P422	H	Cl	CH_3	H	CF_3	CF_3	Br	O	CH_2	O	CH_2CF_3
P423	H	Cl	CH_3	H	CF_3	CF_3	Cl	O	CH_2	O	CH_2CF_3
P424	H	Cl	CH_3	H	CF_3	CF_3	CF_3	O	CH_2	O	CH_2CF_3
P425	H	Cl	CH_3	H	CF_3	CF_3	CH_3	O	CH_2	O	CH_2CF_3
P426	H	Cl	CH_3	H	CF_2CF_3	H	Br	O	CH_2	O	CH_2CF_3
P427	H	Cl	CH_3	H	CF_2CF_3	H	Cl	O	CH_2	O	CH_2CF_3
P428	H	Cl	CH_3	H	CF_2CF_3	H	CF_3	O	CH_2	O	CH_2CF_3
P429	H	Cl	CH_3	H	CF_2CF_3	H	CH_3	O	CH_2	O	CH_2CF_3
P430	H	Cl	CH_3	H	CF_3	H	Br	O	CH_2	O	$CH(CH_3)CF_3$
P431	H	Cl	CH_3	H	CF_3	H	Cl	O	CH_2	O	$CH(CH_3)CF_3$
P432	H	Cl	CH_3	H	CF_3	H	CF_3	O	CH_2	O	$CH(CH_3)CF_3$
P433	H	Cl	CH_3	H	CF_3	H	CH_3	O	CH_2	O	$CH(CH_3)CF_3$
P434	H	Cl	CH_3	H	CF_3	CF_3	Br	O	$CH(CH_3)$	O	CH_2CF_3
P435	H	Cl	CH_3	H	CF_3	CF_3	Cl	O	$CH(CH_3)$	O	CH_2CF_3
P436	H	Cl	CH_3	H	CF_3	CF_3	CF_3	O	$CH(CH_3)$	O	CH_2CF_3
P437	H	Cl	CH_3	H	CF_3	CF_3	CH_3	O	$CH(CH_3)$	O	CH_2CF_3
P438	H	Cl	CH_3	H	CF_2CF_3	H	Br	O	$CH(CH_3)$	O	CH_2CF_3
P439	H	Cl	CH_3	H	CF_2CF_3	H	Cl	O	$CH(CH_3)$	O	CH_2CF_3
P440	H	Cl	CH_3	H	CF_2CF_3	H	CF_3	O	$CH(CH_3)$	O	CH_2CF_3
P441	H	Cl	CH_3	H	CF_2CF_3	H	CH_3	O	$CH(CH_3)$	O	CH_2CF_3
P442	H	Cl	CH_3	H	CF_3	H	Br	O	$CH(CH_3)$	O	CH_2CF_3
P443	H	Cl	CH_3	H	CF_3	H	Cl	O	$CH(CH_3)$	O	CH_2CF_3
P444	H	Cl	CH_3	H	CF_3	H	CF_3	O	$CH(CH_3)$	O	CH_2CF_3

P445	H	Cl	CH_3	H	CF_3	H	CH_3	O	CH (CH_3)	O	CH_2CF_3
P446	H	Cl	CH_3	H	CF_3	H	Br	O	CH(CH_3)	S	CH_2CF_3
P447	H	Cl	CH_3	H	CF_3	H	Cl	O	CH (CH_3)	S	CH_2CF_3
P448	H	Cl	CH_3	H	CF_3	H	CF_3	O	CH(CH_3)	S	CH_2CF_3
P449	H	Cl	CH_3	H	CF_3	H	CH_3	O	CH (CH_3)	S	CH_2CF_3
P450	H	Cl	CH_3	H	CF_3	H	Br	S	CH(CH_3)	O	CH_2CF_3
P451	H	Cl	CH_3	H	CF_3	H	Cl	S	CH (CH_3)	O	CH_2CF_3
P452	H	Cl	CH_3	H	CF_3	H	CF_3	S	CH(CH_3)	O	CH_2CF_3
P453	H	Cl	CH_3	H	CF_3	H	CH_3	S	CH (CH_3)	O	CH_2CF_3
P454	H	Cl	CH_3	H	CF_3	H	Br	O	CH(CH_3)	O	CH_2CHF_2
P455	H	Cl	CH_3	H	CF_3	H	Cl	O	CH (CH_3)	O	CH_2CHF_2
P456	H	Cl	CH_3	H	CF_3	H	CF_3	O	CH(CH_3)	O	CH_2CHF_2
P457	H	Cl	CH_3	H	CF_3	H	CH_3	O	CH (CH_3)	O	CH_2CHF_2
P458	H	Cl	CH_3	H	CF_3	H	Br	O	CH(CH_3)	O	$CH(CH_3)CF_3$
P459	H	Cl	CH_3	H	CF_3	H	Cl	O	CH (CH_3)	O	$CH(CH_3)CF_3$
P460	H	Cl	CH_3	H	CF_3	H	CF_3	O	CH(CH_3)	O	$CH(CH_3)CF_3$
P461	H	Cl	CH_3	H	CF_3	H	CH_3	O	CH (CH_3)	O	$CH(CH_3)CF_3$
P462	H	Cl	CH_3	H	CF_3	H	Br	O	CH(CH_3)	O	$CH_2CH_2CF_3$
P463	H	Cl	CH_3	H	CF_3	H	Cl	O	CH (CH_3)	O	$CH_2CH_2CF_3$
P464	H	Cl	CH_3	H	CF_3	H	CF_3	O	CH(CH_3)	O	$CH_2CH_2CF_3$
P465	H	Cl	CH_3	H	CF_3	H	CH_3	O	CH (CH_3)	O	$CH_2CH_2CF_3$
P466	H	Cl	CH_3	H	CF_3	H	Br	O	CH(CH_2CH_3)	O	CH_2CF_3
P467	H	Cl	CH_3	H	CF_3	H	Cl	O	CH(CH_2CH_3)	O	CH_2CF_3
P468	H	Cl	CH_3	H	CF_3	H	CF_3	O	CH(CH_2CH_3)	O	CH_2CF_3
P469	H	Cl	CH_3	H	CF_3	H	CH_3	O	CH(CH_2CH_3)	O	CH_2CF_3
P470	H	Cl	CH_3	H	CF_3	H	Br	O	$C(CH_3)_2$	O	CH_2CF_3
P471	H	Cl	CH_3	H	CF_3	H	Cl	O	$C(CH_3)_2$	O	CH_2CF_3

P472	H	Cl	CH_3	H	CF_3	H	CF_3	O	$C(CH_3)_2$	O	CH_2CF_3
P473	H	Cl	CH_3	H	CF_3	H	CH_3	O	$C(CH_3)_2$	O	CH_2CF_3
P474	H	Cl	CH_3	H	CF_3	H	Br	O	CH_2CH_2	O	CH_2CF_3
P475	H	Cl	CH_3	H	CF_3	H	Cl	O	CH_2CH_2	O	CH_2CF_3
P476	H	Cl	CH_3	H	CF_3	H	CF_3	O	CH_2CH_2	O	CH_2CF_3
P477	H	Cl	CH_3	H	CF_3	H	CH_3	O	CH_2CH_2	O	CH_2CF_3
P478	H	CH_3	F	CH_3	CF_3	H	Br	O	CH_2	O	CH_2CF_3
P479	H	CH_3	F	CH_3	CF_3	H	Cl	O	CH_2	O	CH_2CF_3
P480	H	CH_3	F	CH_3	CF_3	H	CF_3	O	CH_2	O	CH_2CF_3
P481	H	CH_3	F	CH_3	CF_3	H	CH_3	O	CH_2	O	CH_2CF_3
P482	H	CH_3	F	CH_3	CF_3	H	Br	O	CH_2	S	CH_2CF_3
P483	H	CH_3	F	CH_3	CF_3	H	Cl	O	CH_2	S	CH_2CF_3
P484	H	CH_3	F	CH_3	CF_3	H	CF_3	O	CH_2	S	CH_2CF_3
P485	H	CH_3	F	CH_3	CF_3	H	CH_3	O	CH_2	S	CH_2CF_3
P486	H	CH_3	F	CH_3	CF_3	H	Br	S	CH_2	O	CH_2CF_3
P487	H	CH_3	F	CH_3	CF_3	H	Cl	S	CH_2	O	CH_2CF_3
P488	H	CH_3	F	CH_3	CF_3	H	CF_3	S	CH_2	O	CH_2CF_3
P489	H	CH_3	F	CH_3	CF_3	H	CH_3	S	CH_2	O	CH_2CF_3
P490	H	CH_3	F	CH_3	CF_3	H	Br	O	CH_2	O	CH_2CHF_2
P491	H	CH_3	F	CH_3	CF_3	H	Cl	O	CH_2	O	CH_2CHF_2
P492	H	CH_3	F	CH_3	CF_3	H	CF_3	O	CH_2	O	CH_2CHF_2
P493	H	CH_3	F	CH_3	CF_3	H	CH_3	O	CH_2	O	CH_2CHF_2
P494	H	CH_3	F	CH_3	CF_3	CF_3	Br	O	CH_2	O	CH_2CF_3
P495	H	CH_3	F	CH_3	CF_3	CF_3	Cl	O	CH_2	O	CH_2CF_3
P496	H	CH_3	F	CH_3	CF_3	CF_3	CF_3	O	CH_2	O	CH_2CF_3
P497	H	CH_3	F	CH_3	CF_3	CF_3	CH_3	O	CH_2	O	CH_2CF_3
P498	H	CH_3	F	CH_3	CF_2CF_3	H	Br	O	CH_2	O	CH_2CF_3

P499	H	CH_3	F	CH_3	CF_2CF_3	H	Cl	O	CH_2	O	CH_2CF_3
P500	H	CH_3	F	CH_3	CF_2CF_3	H	CF_3	O	CH_2	O	CH_2CF_3
P501	H	CH_3	F	CH_3	CF_2CF_3	H	CH_3	O	CH_2	O	CH_2CF_3
P502	H	CH_3	F	CH_3	CF_3	H	Br	O	CH_2	O	$CH(CH_3)CF_3$
P503	H	CH_3	F	CH_3	CF_3	H	Cl	O	CH_2	O	$CH(CH_3)CF_3$
P504	H	CH_3	F	CH_3	CF_3	H	CF_3	O	CH_2	O	$CH(CH_3)CF_3$
P505	H	CH_3	F	CH_3	CF_3	H	CH_3	O	CH_2	O	$CH(CH_3)CF_3$
P506	H	CH_3	F	CH_3	CF_3	CF_3	Br	O	$CH(CH_3)$	O	CH_2CF_3
P507	H	CH_3	F	CH_3	CF_3	CF_3	Cl	O	$CH(CH_3)$	O	CH_2CF_3
P508	H	CH_3	F	CH_3	CF_3	CF_3	CF_3	O	$CH(CH_3)$	O	CH_2CF_3
P509	H	CH_3	F	CH_3	CF_3	CF_3	CH_3	O	$CH(CH_3)$	O	CH_2CF_3
P510	H	CH_3	F	CH_3	CF_2CF_3	H	Br	O	$CH(CH_3)$	O	CH_2CF_3
P511	H	CH_3	F	CH_3	CF_2CF_3	H	Cl	O	$CH(CH_3)$	O	CH_2CF_3
P512	H	CH_3	F	CH_3	CF_2CF_3	H	CF_3	O	$CH(CH_3)$	O	CH_2CF_3
P513	H	CH_3	F	CH_3	CF_2CF_3	H	CH_3	O	$CH(CH_3)$	O	CH_2CF_3
P514	H	CH_3	F	CH_3	CF_3	H	Br	O	$CH(CH_3)$	O	CH_2CF_3
P515	H	CH_3	F	CH_3	CF_3	H	Cl	O	$CH(CH_3)$	O	CH_2CF_3
P516	H	CH_3	F	CH_3	CF_3	H	CF_3	O	$CH(CH_3)$	O	CH_2CF_3
P517	H	CH_3	F	CH_3	CF_3	H	CH_3	O	$CH(CH_3)$	O	CH_2CF_3
P518	H	CH_3	F	CH_3	CF_3	H	Br	O	$CH(CH_3)$	S	CH_2CF_3
P519	H	CH_3	F	CH_3	CF_3	H	Cl	O	$CH(CH_3)$	S	CH_2CF_3
P520	H	CH_3	F	CH_3	CF_3	H	CF_3	O	$CH(CH_3)$	S	CH_2CF_3
P521	H	CH_3	F	CH_3	CF_3	H	CH_3	O	$CH(CH_3)$	S	CH_2CF_3
P522	H	CH_3	F	CH_3	CF_3	H	Br	S	$CH(CH_3)$	O	CH_2CF_3
P523	H	CH_3	F	CH_3	CF_3	H	Cl	S	$CH(CH_3)$	O	CH_2CF_3
P524	H	CH_3	F	CH_3	CF_3	H	CF_3	S	$CH(CH_3)$	O	CH_2CF_3
P525	H	CH_3	F	CH_3	CF_3	H	CH_3	S	$CH(CH_3)$	O	CH_2CF_3

P526	H	CH_3	F	CH_3	CF_3	H	Br	O	$CH(CH_3)$	O	CH_2CHF_2
P527	H	CH_3	F	CH_3	CF_3	H	Cl	O	$CH(CH_3)$	O	CH_2CHF_2
P528	H	CH_3	F	CH_3	CF_3	H	CF_3	O	$CH(CH_3)$	O	CH_2CHF_2
P529	H	CH_3	F	CH_3	CF_3	H	CH_3	O	$CH(CH_3)$	O	CH_2CHF_2
P530	H	CH_3	F	CH_3	CF_3	H	Br	O	$CH(CH_3)$	O	$CH(CH_3)CF_3$
P531	H	CH_3	F	CH_3	CF_3	H	Cl	O	$CH(CH_3)$	O	$CH(CH_3)CF_3$
P532	H	CH_3	F	CH_3	CF_3	H	CF_3	O	$CH(CH_3)$	O	$CH(CH_3)CF_3$
P533	H	CH_3	F	CH_3	CF_3	H	CH_3	O	$CH(CH_3)$	O	$CH(CH_3)CF_3$
P534	H	CH_3	F	CH_3	CF_3	H	Br	O	$CH(CH_3)$	O	$CH_2CH_2CF_3$
P535	H	CH_3	F	CH_3	CF_3	H	Cl	O	$CH(CH_3)$	O	$CH_2CH_2CF_3$
P536	H	CH_3	F	CH_3	CF_3	H	CF_3	O	$CH(CH_3)$	O	$CH_2CH_2CF_3$
P537	H	CH_3	F	CH_3	CF_3	H	CH_3	O	$CH(CH_3)$	O	$CH_2CH_2CF_3$
P538	H	CH_3	F	CH_3	CF_3	H	Br	O	$CH(CH_2CH_3)$	O	CH_2CF_3
P539	H	CH_3	F	CH_3	CF_3	H	Cl	O	$CH(CH_2CH_3)$	O	CH_2CF_3
P540	H	CH_3	F	CH_3	CF_3	H	CF_3	O	$CH(CH_2CH_3)$	O	CH_2CF_3
P541	H	CH_3	F	CH_3	CF_3	H	CH_3	O	$CH(CH_2CH_3)$	O	CH_2CF_3
P542	H	CH_3	F	CH_3	CF_3	H	Br	O	$C(CH_3)_2$	O	CH_2CF_3
P543	H	CH_3	F	CH_3	CF_3	H	Cl	O	$C(CH_3)_2$	O	CH_2CF_3
P544	H	CH_3	F	CH_3	CF_3	H	CF_3	O	$C(CH_3)_2$	O	CH_2CF_3
P545	H	CH_3	F	CH_3	CF_3	H	CH_3	O	$C(CH_3)_2$	O	CH_2CF_3
P546	H	CH_3	F	CH_3	CF_3	H	Br	O	CH_2CH_2	O	CH_2CF_3
P547	H	CH_3	F	CH_3	CF_3	H	Cl	O	CH_2CH_2	O	CH_2CF_3
P548	H	CH_3	F	CH_3	CF_3	H	CF_3	O	CH_2CH_2	O	CH_2CF_3
P549	H	CH_3	F	CH_3	CF_3	H	CH_3	O	CH_2CH_2	O	CH_2CF_3
P550	H	Cl	H	Br	CF_3	H	Br	O	CH_2	O	CH_2CF_3
P551	H	Cl	H	Br	CF_3	H	Cl	O	CH_2	O	CH_2CF_3
P552	H	Cl	H	Br	CF_3	H	CF_3	O	CH_2	O	CH_2CF_3

P553	H	Cl	H	Br	CF_3	H	CH_3	O	CH_2	O	CH_2CF_3
P554	H	Cl	H	Br	CF_3	H	Br	O	CH_2	S	CH_2CF_3
P555	H	Cl	H	Br	CF_3	H	Cl	O	CH_2	S	CH_2CF_3
P556	H	Cl	H	Br	CF_3	H	CF_3	O	CH_2	S	CH_2CF_3
P557	H	Cl	H	Br	CF_3	H	CH_3	O	CH_2	S	CH_2CF_3
P558	H	Cl	H	Br	CF_3	H	Br	S	CH_2	O	CH_2CF_3
P559	H	Cl	H	Br	CF_3	H	Cl	S	CH_2	O	CH_2CF_3
P560	H	Cl	H	Br	CF_3	H	CF_3	S	CH_2	O	CH_2CF_3
P561	H	Cl	H	Br	CF_3	H	CH_3	S	CH_2	O	CH_2CF_3
P562	H	Cl	H	Br	CF_3	H	Br	O	CH_2	O	CH_2CHF_2
P563	H	Cl	H	Br	CF_3	H	Cl	O	CH_2	O	CH_2CHF_2
P564	H	Cl	H	Br	CF_3	H	CF_3	O	CH_2	O	CH_2CHF_2
P565	H	Cl	H	Br	CF_3	H	CH_3	O	CH_2	O	CH_2CHF_2
P566	H	Cl	H	Br	CF_3	CF_3	Br	O	CH_2	O	CH_2CF_3
P567	H	Cl	H	Br	CF_3	CF_3	Cl	O	CH_2	O	CH_2CF_3
P568	H	Cl	H	Br	CF_3	CF_3	CF_3	O	CH_2	O	CH_2CF_3
P569	H	Cl	H	Br	CF_3	CF_3	CH_3	O	CH_2	O	CH_2CF_3
P570	H	Cl	H	Br	CF_2CF_3	H	Br	O	CH_2	O	CH_2CF_3
P571	H	Cl	H	Br	CF_2CF_3	H	Cl	O	CH_2	O	CH_2CF_3
P572	H	Cl	H	Br	CF_2CF_3	H	CF_3	O	CH_2	O	CH_2CF_3
P573	H	Cl	H	Br	CF_2CF_3	H	CH_3	O	CH_2	O	CH_2CF_3
P574	H	Cl	H	Br	CF_3	H	Br	O	CH_2	O	$CH(CH_3)CF_3$
P575	H	Cl	H	Br	CF_3	H	Cl	O	CH_2	O	$CH(CH_3)CF_3$
P576	H	Cl	H	Br	CF_3	H	CF_3	O	CH_2	O	$CH(CH_3)CF_3$
P577	H	Cl	H	Br	CF_3	H	CH_3	O	CH_2	O	$CH(CH_3)CF_3$
P578	H	Cl	H	Br	CF_3	CF_3	Br	O	$CH(CH_3)$	O	CH_2CF_3
P579	H	Cl	H	Br	CF_3	CF_3	Cl	O	$CH(CH_3)$	O	CH_2CF_3

P580	H	Cl	H	Br	CF_3	CF_3	CF_3	O	$CH(CH_3)$	O	CH_2CF_3
P581	H	Cl	H	Br	CF_3	CF_3	CH_3	O	$CH(CH_3)$	O	CH_2CF_3
P582	H	Cl	H	Br	CF_2CF_3	H	Br	O	$CH(CH_3)$	O	CH_2CF_3
P583	H	Cl	H	Br	CF_2CF_3	H	Cl	O	$CH(CH_3)$	O	CH_2CF_3
P584	H	Cl	H	Br	CF_2CF_3	H	CF_3	O	$CH(CH_3)$	O	CH_2CF_3
P585	H	Cl	H	Br	CF_2CF_3	H	CH_3	O	$CH(CH_3)$	O	CH_2CF_3
P586	H	Cl	H	Br	CF_3	H	Br	O	$CH(CH_3)$	O	CH_2CF_3
P587	H	Cl	H	Br	CF_3	H	Cl	O	$CH(CH_3)$	O	CH_2CF_3
P588	H	Cl	H	Br	CF_3	H	CF_3	O	$CH(CH_3)$	O	CH_2CF_3
P589	H	Cl	H	Br	CF_3	H	CH_3	O	$CH(CH_3)$	O	CH_2CF_3
P590	H	Cl	H	Br	CF_3	H	Br	O	$CH(CH_3)$	S	CH_2CF_3
P591	H	Cl	H	Br	CF_3	H	Cl	O	$CH(CH_3)$	S	CH_2CF_3
P592	H	Cl	H	Br	CF_3	H	CF_3	O	$CH(CH_3)$	S	CH_2CF_3
P593	H	Cl	H	Br	CF_3	H	CH_3	O	$CH(CH_3)$	S	CH_2CF_3
P594	H	Cl	H	Br	CF_3	H	Br	S	$CH(CH_3)$	O	CH_2CF_3
P595	H	Cl	H	Br	CF_3	H	Cl	S	$CH(CH_3)$	O	CH_2CF_3
P596	H	Cl	H	Br	CF_3	H	CF_3	S	$CH(CH_3)$	O	CH_2CF_3
P597	H	Cl	H	Br	CF_3	H	CH_3	S	$CH(CH_3)$	O	CH_2CF_3
P598	H	Cl	H	Br	CF_3	H	Br	O	$CH(CH_3)$	O	CH_2CHF_2
P599	H	Cl	H	Br	CF_3	H	Cl	O	$CH(CH_3)$	O	CH_2CHF_2
P600	H	Cl	H	Br	CF_3	H	CF_3	O	$CH(CH_3)$	O	CH_2CHF_2
P601	H	Cl	H	Br	CF_3	H	CH_3	O	$CH(CH_3)$	O	CH_2CHF_2
P602	H	Cl	H	Br	CF_3	H	Br	O	$CH(CH_3)$	O	$CH(CH_3)CF_3$
P603	H	Cl	H	Br	CF_3	H	Cl	O	$CH(CH_3)$	O	$CH(CH_3)CF_3$
P604	H	Cl	H	Br	CF_3	H	CF_3	O	$CH(CH_3)$	O	$CH(CH_3)CF_3$
P605	H	Cl	H	Br	CF_3	H	CH_3	O	$CH(CH_3)$	O	$CH(CH_3)CF_3$
P606	H	Cl	H	Br	CF_3	H	Br	O	$CH(CH_3)$	O	$CH_2CH_2CF_3$

P607	H	Cl	H	Br	CF_3	H	Cl	O	$CH(CH_3)$	O	$CH_2CH_2CF_3$
P608	H	Cl	H	Br	CF_3	H	CF_3	O	$CH(CH_3)$	O	$CH_2CH_2CF_3$
P609	H	Cl	H	Br	CF_3	H	CH_3	O	$CH(CH_3)$	O	$CH_2CH_2CF_3$
P610	H	Cl	H	Br	CF_3	H	Br	O	$CH(CH_2CH_3)$	O	CH_2CF_3
P611	H	Cl	H	Br	CF_3	H	Cl	O	$CH(CH_2CH_3)$	O	CH_2CF_3
P612	H	Cl	H	Br	CF_3	H	CF_3	O	$CH(CH_2CH_3)$	O	CH_2CF_3
P613	H	Cl	H	Br	CF_3	H	CH_3	O	$CH(CH_2CH_3)$	O	CH_2CF_3
P614	H	Cl	H	Br	CF_3	H	Br	O	$C(CH_3)_2$	O	CH_2CF_3
P615	H	Cl	H	Br	CF_3	H	Cl	O	$C(CH_3)_2$	O	CH_2CF_3
P616	H	Cl	H	Br	CF_3	H	CF_3	O	$C(CH_3)_2$	O	CH_2CF_3
P617	H	Cl	H	Br	CF_3	H	CH_3	O	$C(CH_3)_2$	O	CH_2CF_3
P618	H	Cl	H	Br	CF_3	H	Br	O	CH_2CH_2	O	CH_2CF_3
P619	H	Cl	H	Br	CF_3	H	Cl	O	CH_2CH_2	O	CH_2CF_3
P620	H	Cl	H	Br	CF_3	H	CF_3	O	CH_2CH_2	O	CH_2CF_3
P621	H	Cl	H	Br	CF_3	H	CH_3	O	CH_2CH_2	O	CH_2CF_3
P622	H	H	Br	Br	CF_3	H	Br	O	CH_2	O	CH_2CF_3
P623	H	H	Br	Br	CF_3	H	Cl	O	CH_2	O	CH_2CF_3
P624	H	H	Br	Br	CF_3	H	CF_3	O	CH_2	O	CH_2CF_3
P625	H	H	Br	Br	CF_3	H	CH_3	O	CH_2	O	CH_2CF_3
P626	H	H	Br	Br	CF_3	H	Br	O	CH_2	S	CH_2CF_3
P627	H	H	Br	Br	CF_3	H	Cl	O	CH_2	S	CH_2CF_3
P628	H	H	Br	Br	CF_3	H	CF_3	O	CH_2	S	CH_2CF_3
P629	H	H	Br	Br	CF_3	H	CH_3	O	CH_2	S	CH_2CF_3
P630	H	H	Br	Br	CF_3	H	Br	S	CH_2	O	CH_2CF_3
P631	H	H	Br	Br	CF_3	H	Cl	S	CH_2	O	CH_2CF_3
P632	H	H	Br	Br	CF_3	H	CF_3	S	CH_2	O	CH_2CF_3
P633	H	H	Br	Br	CF_3	H	CH_3	S	CH_2	O	CH_2CF_3

P634	H	H	Br	Br	CF_3	H	Br	O	CH_2	O	CH_2CHF_2
P635	H	H	Br	Br	CF_3	H	Cl	O	CH_2	O	CH_2CHF_2
P636	H	H	Br	Br	CF_3	H	CF_3	O	CH_2	O	CH_2CHF_2
P637	H	H	Br	Br	CF_3	H	CH_3	O	CH_2	O	CH_2CHF_2
P638	H	H	Br	Br	CF_3	CF_3	Br	O	CH_2	O	CH_2CF_3
P639	H	H	Br	Br	CF_3	CF_3	Cl	O	CH_2	O	CH_2CF_3
P640	H	H	Br	Br	CF_3	CF_3	CF_3	O	CH_2	O	CH_2CF_3
P641	H	H	Br	Br	CF_3	CF_3	CH_3	O	CH_2	O	CH_2CF_3
P642	H	H	Br	Br	CF_2CF_3	H	Br	O	CH_2	O	CH_2CF_3
P643	H	H	Br	Br	CF_2CF_3	H	Cl	O	CH_2	O	CH_2CF_3
P644	H	H	Br	Br	CF_2CF_3	H	CF_3	O	CH_2	O	CH_2CF_3
P645	H	H	Br	Br	CF_2CF_3	H	CH_3	O	CH_2	O	CH_2CF_3
P646	H	H	Br	Br	CF_3	H	Br	O	CH_2	O	$CH(CH_3)CF_3$
P647	H	H	Br	Br	CF_3	H	Cl	O	CH_2	O	$CH(CH_3)CF_3$
P648	H	H	Br	Br	CF_3	H	CF_3	O	CH_2	O	$CH(CH_3)CF_3$
P649	H	H	Br	Br	CF_3	H	CH_3	O	CH_2	O	$CH(CH_3)CF_3$
P650	H	H	Br	Br	CF_3	CF_3	Br	O	$CH(CH_3)$	O	CH_2CF_3
P651	H	H	Br	Br	CF_3	CF_3	Cl	O	$CH(CH_3)$	O	CH_2CF_3
P652	H	H	Br	Br	CF_3	CF_3	CF_3	O	$CH(CH_3)$	O	CH_2CF_3
P653	H	H	Br	Br	CF_3	CF_3	CH_3	O	$CH(CH_3)$	O	CH_2CF_3
P654	H	H	Br	Br	CF_2CF_3	H	Br	O	$CH(CH_3)$	O	CH_2CF_3
P655	H	H	Br	Br	CF_2CF_3	H	Cl	O	$CH(CH_3)$	O	CH_2CF_3
P656	H	H	Br	Br	CF_2CF_3	H	CF_3	O	$CH(CH_3)$	O	CH_2CF_3
P657	H	H	Br	Br	CF_2CF_3	H	CH_3	O	$CH(CH_3)$	O	CH_2CF_3
P658	H	H	Br	Br	CF_3	H	Br	O	$CH(CH_3)$	O	CH_2CF_3
P659	H	H	Br	Br	CF_3	H	Cl	O	$CH(CH_3)$	O	CH_2CF_3
P660	H	H	Br	Br	CF_3	H	CF_3	O	$CH(CH_3)$	O	CH_2CF_3

P661	H	H	Br	Br	CF_3	H	CH_3	O	$CH(CH_3)$	O	CH_2CF_3
P662	H	H	Br	Br	CF_3	H	Br	O	$CH(CH_3)$	S	CH_2CF_3
P663	H	H	Br	Br	CF_3	H	Cl	O	$CH(CH_3)$	S	CH_2CF_3
P664	H	H	Br	Br	CF_3	H	CF_3	O	$CH(CH_3)$	S	CH_2CF_3
P665	H	H	Br	Br	CF_3	H	CH_3	O	$CH(CH_3)$	S	CH_2CF_3
P666	H	H	Br	Br	CF_3	H	Br	S	$CH(CH_3)$	O	CH_2CF_3
P667	H	H	Br	Br	CF_3	H	Cl	S	$CH(CH_3)$	O	CH_2CF_3
P668	H	H	Br	Br	CF_3	H	CF_3	S	$CH(CH_3)$	O	CH_2CF_3
P669	H	H	Br	Br	CF_3	H	CH_3	S	$CH(CH_3)$	O	CH_2CF_3
P670	H	H	Br	Br	CF_3	H	Br	O	$CH(CH_3)$	O	CH_2CHF_2
P671	H	H	Br	Br	CF_3	H	Cl	O	$CH(CH_3)$	O	CH_2CHF_2
P672	H	H	Br	Br	CF_3	H	CF_3	O	$CH(CH_3)$	O	CH_2CHF_2
P673	H	H	Br	Br	CF_3	H	CH_3	O	$CH(CH_3)$	O	CH_2CHF_2
P674	H	H	Br	Br	CF_3	H	Br	O	$CH(CH_3)$	O	$CH(CH_3)CF_3$
P675	H	H	Br	Br	CF_3	H	Cl	O	$CH(CH_3)$	O	$CH(CH_3)CF_3$
P676	H	H	Br	Br	CF_3	H	CF_3	O	$CH(CH_3)$	O	$CH(CH_3)CF_3$
P677	H	H	Br	Br	CF_3	H	CH_3	O	$CH(CH_3)$	O	$CH(CH_3)CF_3$
P678	H	H	Br	Br	CF_3	H	Br	O	$CH(CH_3)$	O	$CH_2CH_2CF_3$
P679	H	H	Br	Br	CF_3	H	Cl	O	$CH(CH_3)$	O	$CH_2CH_2CF_3$
P680	H	H	Br	Br	CF_3	H	CF_3	O	$CH(CH_3)$	O	$CH_2CH_2CF_3$
P681	H	H	Br	Br	CF_3	H	CH_3	O	$CH(CH_3)$	O	$CH_2CH_2CF_3$
P682	H	H	Br	Br	CF_3	H	Br	O	$CH(CH_2CH_3)$	O	CH_2CF_3
P683	H	H	Br	Br	CF_3	H	Cl	O	$CH(CH_2CH_3)$	O	CH_2CF_3
P684	H	H	Br	Br	CF_3	H	CF_3	O	$CH(CH_2CH_3)$	O	CH_2CF_3
P685	H	H	Br	Br	CF_3	H	CH_3	O	$CH(CH_2CH_3)$	O	CH_2CF_3
P686	H	H	Br	Br	CF_3	H	Br	O	$C(CH_3)_2$	O	CH_2CF_3
P687	H	H	Br	Br	CF_3	H	Cl	O	$C(CH_3)_2$	O	CH_2CF_3

P688	H	H	Br	Br	CF_3	H	CF_3	O	$C(CH_3)_2$	O	CH_2CF_3
P689	H	H	Br	Br	CF_3	H	CH_3	O	$C(CH_3)_2$	O	CH_2CF_3
P690	H	H	Br	Br	CF_3	H	Br	O	CH_2CH_2	O	CH_2CF_3
P691	H	H	Br	Br	CF_3	H	Cl	O	CH_2CH_2	O	CH_2CF_3
P692	H	H	Br	Br	CF_3	H	CF_3	O	CH_2CH_2	O	CH_2CF_3
P693	H	H	Br	Br	CF_3	H	CH_3	O	CH_2CH_2	O	CH_2CF_3
P694	H	H	Cl	NO_2	CF_3	H	Br	O	CH_2	O	CH_2CF_3
P695	H	H	Cl	NO_2	CF_3	H	Cl	O	CH_2	O	CH_2CF_3
P696	H	H	Cl	NO_2	CF_3	H	CF_3	O	CH_2	O	CH_2CF_3
P697	H	H	Cl	NO_2	CF_3	H	CH_3	O	CH_2	O	CH_2CF_3
P698	H	H	Cl	NO_2	CF_3	H	Br	O	CH_2	S	CH_2CF_3
P699	H	H	Cl	NO_2	CF_3	H	Cl	O	CH_2	S	CH_2CF_3
P700	H	H	Cl	NO_2	CF_3	H	CF_3	O	CH_2	S	CH_2CF_3
P701	H	H	Cl	NO_2	CF_3	H	CH_3	O	CH_2	S	CH_2CF_3
P702	H	H	Cl	NO_2	CF_3	H	Br	S	CH_2	O	CH_2CF_3
P703	H	H	Cl	NO_2	CF_3	H	Cl	S	CH_2	O	CH_2CF_3
P704	H	H	Cl	NO_2	CF_3	H	CF_3	S	CH_2	O	CH_2CF_3
P705	H	H	Cl	NO_2	CF_3	H	CH_3	S	CH_2	O	CH_2CF_3
P706	H	H	Cl	NO_2	CF_3	H	Br	O	CH_2	O	CH_2CHF_2
P707	H	H	Cl	NO_2	CF_3	H	Cl	O	CH_2	O	CH_2CHF_2
P708	H	H	Cl	NO_2	CF_3	H	CF_3	O	CH_2	O	CH_2CHF_2
P709	H	H	Cl	NO_2	CF_3	H	CH_3	O	CH_2	O	CH_2CHF_2
P710	H	H	Cl	NO_2	CF_3	CF_3	Br	O	CH_2	O	CH_2CF_3
P711	H	H	Cl	NO_2	CF_3	CF_3	Cl	O	CH_2	O	CH_2CF_3
P712	H	H	Cl	NO_2	CF_3	CF_3	CF_3	O	CH_2	O	CH_2CF_3
P713	H	H	Cl	NO_2	CF_3	CF_3	CH_3	O	CH_2	O	CH_2CF_3
P714	H	H	Cl	NO_2	CF_2CF_3	H	Br	O	CH_2	O	CH_2CF_3

P715	H	H	Cl	NO_2	CF_2CF_3	H	Cl	O	CH_2	O	CH_2CF_3
P716	H	H	Cl	NO_2	CF_2CF_3	H	CF_3	O	CH_2	O	CH_2CF_3
P717	H	H	Cl	NO_2	CF_2CF_3	H	CH_3	O	CH_2	O	CH_2CF_3
P718	H	H	Cl	NO_2	CF_3	H	Br	O	CH_2	O	$CH(CH_3)CF_3$
P719	H	H	Cl	NO_2	CF_3	H	Cl	O	CH_2	O	$CH(CH_3)CF_3$
P720	H	H	Cl	NO_2	CF_3	H	CF_3	O	CH_2	O	$CH(CH_3)CF_3$
P721	H	H	Cl	NO_2	CF_3	H	CH_3	O	CH_2	O	$CH(CH_3)CF_3$
P722	H	H	Cl	NO_2	CF_3	CF_3	Br	O	$CH(CH_3)$	O	CH_2CF_3
P723	H	H	Cl	NO_2	CF_3	CF_3	Cl	O	$CH(CH_3)$	O	CH_2CF_3
P724	H	H	Cl	NO_2	CF_3	CF_3	CF_3	O	$CH(CH_3)$	O	CH_2CF_3
P725	H	H	Cl	NO_2	CF_3	CF_3	CH_3	O	$CH(CH_3)$	O	CH_2CF_3
P726	H	H	Cl	NO_2	CF_2CF_3	H	Br	O	$CH(CH_3)$	O	CH_2CF_3
P727	H	H	Cl	NO_2	CF_2CF_3	H	Cl	O	$CH(CH_3)$	O	CH_2CF_3
P728	H	H	Cl	NO_2	CF_2CF_3	H	CF_3	O	$CH(CH_3)$	O	CH_2CF_3
P729	H	H	Cl	NO_2	CF_2CF_3	H	CH_3	O	$CH(CH_3)$	O	CH_2CF_3
P730	H	H	Cl	NO_2	CF_3	H	Br	O	$CH(CH_3)$	O	CH_2CF_3
P731	H	H	Cl	NO_2	CF_3	H	Cl	O	$CH(CH_3)$	O	CH_2CF_3
P732	H	H	Cl	NO_2	CF_3	H	CF_3	O	$CH(CH_3)$	O	CH_2CF_3
P733	H	H	Cl	NO_2	CF_3	H	CH_3	O	$CH(CH_3)$	O	CH_2CF_3
P734	H	H	Cl	NO_2	CF_3	H	Br	O	$CH(CH_3)$	S	CH_2CF_3
P735	H	H	Cl	NO_2	CF_3	H	Cl	O	$CH(CH_3)$	S	CH_2CF_3
P736	H	H	Cl	NO_2	CF_3	H	CF_3	O	$CH(CH_3)$	S	CH_2CF_3
P737	H	H	Cl	NO_2	CF_3	H	CH_3	O	$CH(CH_3)$	S	CH_2CF_3
P738	H	H	Cl	NO_2	CF_3	H	Br	S	$CH(CH_3)$	O	CH_2CF_3
P739	H	H	Cl	NO_2	CF_3	H	Cl	S	$CH(CH_3)$	O	CH_2CF_3
P740	H	H	Cl	NO_2	CF_3	H	CF_3	S	$CH(CH_3)$	O	CH_2CF_3
P741	H	H	Cl	NO_2	CF_3	H	CH_3	S	$CH(CH_3)$	O	CH_2CF_3

P742	H	H	Cl	NO_2	CF_3	H	Br	O	$CH(CH_3)$	O	CH_2CHF_2
P743	H	H	Cl	NO_2	CF_3	H	Cl	O	$CH(CH_3)$	O	CH_2CHF_2
P744	H	H	Cl	NO_2	CF_3	H	CF_3	O	$CH(CH_3)$	O	CH_2CHF_2
P745	H	H	Cl	NO_2	CF_3	H	CH_3	O	$CH(CH_3)$	O	CH_2CHF_2
P746	H	H	Cl	NO_2	CF_3	H	Br	O	$CH(CH_3)$	O	$CH(CH_3)CF_3$
P747	H	H	Cl	NO_2	CF_3	H	Cl	O	$CH(CH_3)$	O	$CH(CH_3)CF_3$
P748	H	H	Cl	NO_2	CF_3	H	CF_3	O	$CH(CH_3)$	O	$CH(CH_3)CF_3$
P749	H	H	Cl	NO_2	CF_3	H	CH_3	O	$CH(CH_3)$	O	$CH(CH_3)CF_3$
P750	H	H	Cl	NO_2	CF_3	H	Br	O	$CH(CH_3)$	O	$CH_2CH_2CF_3$
P751	H	H	Cl	NO_2	CF_3	H	Cl	O	$CH(CH_3)$	O	$CH_2CH_2CF_3$
P752	H	H	Cl	NO_2	CF_3	H	CF_3	O	$CH(CH_3)$	O	$CH_2CH_2CF_3$
P753	H	H	Cl	NO_2	CF_3	H	CH_3	O	$CH(CH_3)$	O	$CH_2CH_2CF_3$
P754	H	H	Cl	NO_2	CF_3	H	Br	O	$CH(CH_2CH_3)$	O	CH_2CF_3
P755	H	H	Cl	NO_2	CF_3	H	Cl	O	$CH(CH_2CH_3)$	O	CH_2CF_3
P756	H	H	Cl	NO_2	CF_3	H	CF_3	O	$CH(CH_2CH_3)$	O	CH_2CF_3
P757	H	H	Cl	NO_2	CF_3	H	CH_3	O	$CH(CH_2CH_3)$	O	CH_2CF_3
P758	H	H	Cl	NO_2	CF_3	H	Br	O	$C(CH_3)_2$	O	CH_2CF_3
P759	H	H	Cl	NO_2	CF_3	H	Cl	O	$C(CH_3)_2$	O	CH_2CF_3
P760	H	H	Cl	NO_2	CF_3	H	CF_3	O	$C(CH_3)_2$	O	CH_2CF_3
P761	H	H	Cl	NO_2	CF_3	H	CH_3	O	$C(CH_3)_2$	O	CH_2CF_3
P762	H	H	Cl	NO_2	CF_3	H	Br	O	CH_2CH_2	O	CH_2CF_3
P763	H	H	Cl	NO_2	CF_3	H	Cl	O	CH_2CH_2	O	CH_2CF_3
P764	H	H	Cl	NO_2	CF_3	H	CF_3	O	CH_2CH_2	O	CH_2CF_3
P765	H	H	Cl	NO_2	CF_3	H	CH_3	O	CH_2CH_2	O	CH_2CF_3
P766	H	H	F	CN	CF_3	H	Br	O	CH_2	O	CH_2CF_3
P767	H	H	F	CN	CF_3	H	Cl	O	CH_2	O	CH_2CF_3
P768	H	H	F	CN	CF_3	H	CF_3	O	CH_2	O	CH_2CF_3

P769	H	H	F	CN	CF_3	H	CH_3	O	CH_2	O	CH_2CF_3
P770	H	H	F	CN	CF_3	H	Br	O	CH_2	S	CH_2CF_3
P771	H	H	F	CN	CF_3	H	Cl	O	CH_2	S	CH_2CF_3
P772	H	H	F	CN	CF_3	H	CF_3	O	CH_2	S	CH_2CF_3
P773	H	H	F	CN	CF_3	H	CH_3	O	CH_2	S	CH_2CF_3
P774	H	H	F	CN	CF_3	H	Br	S	CH_2	O	CH_2CF_3
P775	H	H	F	CN	CF_3	H	Cl	S	CH_2	O	CH_2CF_3
P776	H	H	F	CN	CF_3	H	CF_3	S	CH_2	O	CH_2CF_3
P777	H	H	F	CN	CF_3	H	CH_3	S	CH_2	O	CH_2CF_3
P778	H	H	F	CN	CF_3	H	Br	O	CH_2	O	CH_2CHF_2
P779	H	H	F	CN	CF_3	H	Cl	O	CH_2	O	CH_2CHF_2
P780	H	H	F	CN	CF_3	H	CF_3	O	CH_2	O	CH_2CHF_2
P781	H	H	F	CN	CF_3	H	CH_3	O	CH_2	O	CH_2CHF_2
P782	H	H	F	CN	CF_3	CF_3	Br	O	CH_2	O	CH_2CF_3
P783	H	H	F	CN	CF_3	CF_3	Cl	O	CH_2	O	CH_2CF_3
P784	H	H	F	CN	CF_3	CF_3	CF_3	O	CH_2	O	CH_2CF_3
P785	H	H	F	CN	CF_3	CF_3	CH_3	O	CH_2	O	CH_2CF_3
P786	H	H	F	CN	CF_2CF_3	H	Br	O	CH_2	O	CH_2CF_3
P787	H	H	F	CN	CF_2CF_3	H	Cl	O	CH_2	O	CH_2CF_3
P788	H	H	F	CN	CF_2CF_3	H	CF_3	O	CH_2	O	CH_2CF_3
P789	H	H	F	CN	CF_2CF_3	H	CH_3	O	CH_2	O	CH_2CF_3
P790	H	H	F	CN	CF_3	H	Br	O	CH_2	O	$CH(CH_3)CF_3$
P791	H	H	F	CN	CF_3	H	Cl	O	CH_2	O	$CH(CH_3)CF_3$
P792	H	H	F	CN	CF_3	H	CF_3	O	CH_2	O	$CH(CH_3)CF_3$
P793	H	H	F	CN	CF_3	H	CH_3	O	CH_2	O	$CH(CH_3)CF_3$
P794	H	H	F	CN	CF_3	CF_3	Br	O	$CH(CH_3)$	O	CH_2CF_3
P795	H	H	F	CN	CF_3	CF_3	Cl	O	$CH(CH_3)$	O	CH_2CF_3

P796	H	H	F	CN	CF_3	CF_3	CF_3	O	$CH(CH_3)$	O	CH_2CF_3
P797	H	H	F	CN	CF_3	CF_3	CH_3	O	$CH(CH_3)$	O	CH_2CF_3
P798	H	H	F	CN	CF_2CF_3	H	Br	O	$CH(CH_3)$	O	CH_2CF_3
P799	H	H	F	CN	CF_2CF_3	H	Cl	O	$CH(CH_3)$	O	CH_2CF_3
P800	H	H	F	CN	CF_2CF_3	H	CF_3	O	$CH(CH_3)$	O	CH_2CF_3
P801	H	H	F	CN	CF_2CF_3	H	CH_3	O	$CH(CH_3)$	O	CH_2CF_3
P802	H	H	F	CN	CF_3	H	Br	O	$CH(CH_3)$	O	CH_2CF_3
P803	H	H	F	CN	CF_3	H	Cl	O	$CH(CH_3)$	O	CH_2CF_3
P804	H	H	F	CN	CF_3	H	CF_3	O	$CH(CH_3)$	O	CH_2CF_3
P805	H	H	F	CN	CF_3	H	CH_3	O	$CH(CH_3)$	O	CH_2CF_3
P806	H	H	F	CN	CF_3	H	Br	O	$CH(CH_3)$	S	CH_2CF_3
P807	H	H	F	CN	CF_3	H	Cl	O	$CH(CH_3)$	S	CH_2CF_3
P808	H	H	F	CN	CF_3	H	CF_3	O	$CH(CH_3)$	S	CH_2CF_3
P809	H	H	F	CN	CF_3	H	CH_3	O	$CH(CH_3)$	S	CH_2CF_3
P810	H	H	F	CN	CF_3	H	Br	S	$CH(CH_3)$	O	CH_2CF_3
P811	H	H	F	CN	CF_3	H	Cl	S	$CH(CH_3)$	O	CH_2CF_3
P812	H	H	F	CN	CF_3	H	CF_3	S	$CH(CH_3)$	O	CH_2CF_3
P813	H	H	F	CN	CF_3	H	CH_3	S	$CH(CH_3)$	O	CH_2CF_3
P814	H	H	F	CN	CF_3	H	Br	O	$CH(CH_3)$	O	CH_2CHF_2
P815	H	H	F	CN	CF_3	H	Cl	O	$CH(CH_3)$	O	CH_2CHF_2
P816	H	H	F	CN	CF_3	H	CF_3	O	$CH(CH_3)$	O	CH_2CHF_2
P817	H	H	F	CN	CF_3	H	CH_3	O	$CH(CH_3)$	O	CH_2CHF_2
P818	H	H	F	CN	CF_3	H	Br	O	$CH(CH_3)$	O	$CH(CH_3)CF_3$
P819	H	H	F	CN	CF_3	H	Cl	O	$CH(CH_3)$	O	$CH(CH_3)CF_3$
P820	H	H	F	CN	CF_3	H	CF_3	O	$CH(CH_3)$	O	$CH(CH_3)CF_3$
P821	H	H	F	CN	CF_3	H	CH_3	O	$CH(CH_3)$	O	$CH(CH_3)CF_3$
P822	H	H	F	CN	CF_3	H	Br	O	$CH(CH_3)$	O	$CH_2CH_2CF_3$

P823	H	H	F	CN	CF_3	H	Cl	O	$CH(CH_3)$	O	$CH_2CH_2CF_3$
P824	H	H	F	CN	CF_3	H	CF_3	O	$CH(CH_3)$	O	$CH_2CH_2CF_3$
P825	H	H	F	CN	CF_3	H	CH_3	O	$CH(CH_3)$	O	$CH_2CH_2CF_3$
P826	H	H	F	CN	CF_3	H	Br	O	$CH(CH_2CH_3)$	O	CH_2CF_3
P827	H	H	F	CN	CF_3	H	Cl	O	$CH(CH_2CH_3)$	O	CH_2CF_3
P828	H	H	F	CN	CF_3	H	CF_3	O	$CH(CH_2CH_3)$	O	CH_2CF_3
P829	H	H	F	CN	CF_3	H	CH_3	O	$CH(CH_2CH_3)$	O	CH_2CF_3
P830	H	H	F	CN	CF_3	H	Br	O	$C(CH_3)_2$	O	CH_2CF_3
P831	H	H	F	CN	CF_3	H	Cl	O	$C(CH_3)_2$	O	CH_2CF_3
P832	H	H	F	CN	CF_3	H	CF_3	O	$C(CH_3)_2$	O	CH_2CF_3
P833	H	H	F	CN	CF_3	H	CH_3	O	$C(CH_3)_2$	O	CH_2CF_3
P834	H	H	F	CN	CF_3	H	Br	O	CH_2CH_2	O	CH_2CF_3
P835	H	H	F	CN	CF_3	H	Cl	O	CH_2CH_2	O	CH_2CF_3
P836	H	H	F	CN	CF_3	H	CF_3	O	CH_2CH_2	O	CH_2CF_3
P837	H	H	F	CN	CF_3	H	CH_3	O	CH_2CH_2	O	CH_2CF_3
P838	H	Cl	OCF_3	Cl	CF_3	H	Br	O	CH_2	O	CH_2CF_3
P839	H	Cl	OCF_3	Cl	CF_3	H	Cl	O	CH_2	O	CH_2CF_3
P840	H	Cl	OCF_3	Cl	CF_3	H	CF_3	O	CH_2	O	CH_2CF_3
P841	H	Cl	OCF_3	Cl	CF_3	H	CH_3	O	CH_2	O	CH_2CF_3
P842	H	Cl	OCF_3	Cl	CF_3	H	Br	O	CH_2	S	CH_2CF_3
P843	H	Cl	OCF_3	Cl	CF_3	H	Cl	O	CH_2	S	CH_2CF_3
P844	H	Cl	OCF_3	Cl	CF_3	H	CF_3	O	CH_2	S	CH_2CF_3

P845	H	Cl	OCF_3	Cl	CF_3	H	CH_3	O	CH_2	S	CH_2CF_3
P846	H	Cl	OCF_3	Cl	CF_3	H	Br	S	CH_2	O	CH_2CF_3
P847	H	Cl	OCF_3	Cl	CF_3	H	Cl	S	CH_2	O	CH_2CF_3
P848	H	Cl	OCF_3	Cl	CF_3	H	CF_3	S	CH_2	O	CH_2CF_3
P849	H	Cl	OCF_3	Cl	CF_3	H	CH_3	S	CH_2	O	CH_2CF_3
P850	H	Cl	OCF_3	Cl	CF_3	H	Br	O	CH_2	O	CH_2CHF_2
P851	H	Cl	OCF_3	Cl	CF_3	H	Cl	O	CH_2	O	CH_2CHF_2
P852	H	Cl	OCF_3	Cl	CF_3	H	CF_3	O	CH_2	O	CH_2CHF_2
P853	H	Cl	OCF_3	Cl	CF_3	H	CH_3	O	CH_2	O	CH_2CHF_2
P854	H	Cl	OCF_3	Cl	CF_3	CF_3	Br	O	CH_2	O	CH_2CF_3
P855	H	Cl	OCF_3	Cl	CF_3	CF_3	Cl	O	CH_2	O	CH_2CF_3
P856	H	Cl	OCF_3	Cl	CF_3	CF_3	CF_3	O	CH_2	O	CH_2CF_3
P857	H	Cl	OCF_3	Cl	CF_3	CF_3	CH_3	O	CH_2	O	CH_2CF_3
P858	H	Cl	OCF_3	Cl	CF_2CF_3	H	Br	O	CH_2	O	CH_2CF_3
P859	H	Cl	OCF_3	Cl	CF_2CF_3	H	Cl	O	CH_2	O	CH_2CF_3
P860	H	Cl	OCF_3	Cl	CF_2CF_3	H	CF_3	O	CH_2	O	CH_2CF_3
P861	H	Cl	OCF_3	Cl	CF_2CF_3	H	CH_3	O	CH_2	O	CH_2CF_3

P862	H	Cl	OCF_3	Cl	CF_3	H	Br	O	CH_2	O	$CH(CH_3)CF_3$
P863	H	Cl	OCF_3	Cl	CF_3	H	Cl	O	CH_2	O	$CH(CH_3)CF_3$
P864	H	Cl	OCF_3	Cl	CF_3	H	CF_3	O	CH_2	O	$CH(CH_3)CF_3$
P865	H	Cl	OCF_3	Cl	CF_3	H	CH_3	O	CH_2	O	$CH(CH_3)CF_3$
P866	H	Cl	OCF_3	Cl	CF_3	CF_3	Br	O	$CH(CH_3)$	O	CH_2CF_3
P867	H	Cl	OCF_3	Cl	CF_3	CF_3	Cl	O	$CH(CH_3)$	O	CH_2CF_3
P868	H	Cl	OCF_3	Cl	CF_3	CF_3	CF_3	O	$CH(CH_3)$	O	CH_2CF_3
P869	H	Cl	OCF_3	Cl	CF_3	CF_3	CH_3	O	$CH(CH_3)$	O	CH_2CF_3
P870	H	Cl	OCF_3	Cl	CF_2CF_3	H	Br	O	$CH(CH_3)$	O	CH_2CF_3
P871	H	Cl	OCF_3	Cl	CF_2CF_3	H	Cl	O	$CH(CH_3)$	O	CH_2CF_3
P872	H	Cl	OCF_3	Cl	CF_2CF_3	H	CF_3	O	$CH(CH_3)$	O	CH_2CF_3
P873	H	Cl	OCF_3	Cl	CF_2CF_3	H	CH_3	O	$CH(CH_3)$	O	CH_2CF_3
P874	H	Cl	OCF_3	Cl	CF_3	H	Br	O	$CH(CH_3)$	O	CH_2CF_3
P875	H	Cl	OCF_3	Cl	CF_3	H	Cl	O	$CH(CH_3)$	O	CH_2CF_3
P876	H	Cl	OCF_3	Cl	CF_3	H	CF_3	O	$CH(CH_3)$	O	CH_2CF_3
P877	H	Cl	OCF_3	Cl	CF_3	H	CH_3	O	$CH(CH_3)$	O	CH_2CF_3
P878	H	Cl	OCF_3	Cl	CF_3	H	Br	O	$CH(CH_3)$	S	CH_2CF_3

P879	H	Cl	OCF_3	Cl	CF_3	H	Cl	O	$CH(CH_3)$	S	CH_2CF_3
P880	H	Cl	OCF_3	Cl	CF_3	H	CF_3	O	$CH(CH_3)$	S	CH_2CF_3
P881	H	Cl	OCF_3	Cl	CF_3	H	CH_3	O	$CH(CH_3)$	S	CH_2CF_3
P882	H	Cl	OCF_3	Cl	CF_3	H	Br	S	$CH(CH_3)$	O	CH_2CF_3
P883	H	Cl	OCF_3	Cl	CF_3	H	Cl	S	$CH(CH_3)$	O	CH_2CF_3
P884	H	Cl	OCF_3	Cl	CF_3	H	CF_3	S	$CH(CH_3)$	O	CH_2CF_3
P885	H	Cl	OCF_3	Cl	CF_3	H	CH_3	S	$CH(CH_3)$	O	CH_2CF_3
P886	H	Cl	OCF_3	Cl	CF_3	H	Br	O	$CH(CH_3)$	O	CH_2CHF_2
P887	H	Cl	OCF_3	Cl	CF_3	H	Cl	O	$CH(CH_3)$	O	CH_2CHF_2
P888	H	Cl	OCF_3	Cl	CF_3	H	CF_3	O	$CH(CH_3)$	O	CH_2CHF_2
P889	H	Cl	OCF_3	Cl	CF_3	H	CH_3	O	$CH(CH_3)$	O	CH_2CHF_2
P890	H	Cl	OCF_3	Cl	CF_3	H	Br	O	$CH(CH_3)$	O	$CH(CH_3)CF_3$
P891	H	Cl	OCF_3	Cl	CF_3	H	Cl	O	$CH(CH_3)$	O	$CH(CH_3)CF_3$
P892	H	Cl	OCF_3	Cl	CF_3	H	CF_3	O	$CH(CH_3)$	O	$CH(CH_3)CF_3$
P893	H	Cl	OCF_3	Cl	CF_3	H	CH_3	O	$CH(CH_3)$	O	$CH(CH_3)CF_3$
P894	H	Cl	OCF_3	Cl	CF_3	H	Br	O	$CH(CH_3)$	O	$CH_2CH_2CF_3$
P895	H	Cl	OCF_3	Cl	CF_3	H	Cl	O	$CH(CH_3)$	O	$CH_2CH_2CF_3$

P896	H	Cl	OCF_3	Cl	CF_3	H	CF_3	O	$CH(CH_3)$	O	$CH_2CH_2CF_3$
P897	H	Cl	OCF_3	Cl	CF_3	H	CH_3	O	$CH(CH_3)$	O	$CH_2CH_2CF_3$
P898	H	Cl	OCF_3	Cl	CF_3	H	Br	O	$CH(CH_2CH_3)$	O	CH_2CF_3
P899	H	Cl	OCF_3	Cl	CF_3	H	Cl	O	$CH(CH_2CH_3)$	O	CH_2CF_3
P900	H	Cl	OCF_3	Cl	CF_3	H	CF_3	O	$CH(CH_2CH_3)$	O	CH_2CF_3
P901	H	Cl	OCF_3	Cl	CF_3	H	CH_3	O	$CH(CH_2CH_3)$	O	CH_2CF_3
P902	H	Cl	OCF_3	Cl	CF_3	H	Br	O	$C(CH_3)_2$	O	CH_2CF_3
P903	H	Cl	OCF_3	Cl	CF_3	H	Cl	O	$C(CH_3)_2$	O	CH_2CF_3
P904	H	Cl	OCF_3	Cl	CF_3	H	CF_3	O	$C(CH_3)_2$	O	CH_2CF_3
P905	H	Cl	OCF_3	Cl	CF_3	H	CH_3	O	$C(CH_3)_2$	O	CH_2CF_3
P906	H	Cl	OCF_3	Cl	CF_3	H	Br	O	CH_2CH_2	O	CH_2CF_3
P907	H	Cl	OCF_3	Cl	CF_3	H	Cl	O	CH_2CH_2	O	CH_2CF_3
P908	H	Cl	OCF_3	Cl	CF_3	H	CF_3	O	CH_2CH_2	O	CH_2CF_3
P909	H	Cl	OCF_3	Cl	CF_3	H	CH_3	O	CH_2CH_2	O	CH_2CF_3
P910	H	Cl	CN	Cl	CF_3	H	Br	O	CH_2	O	CH_2CF_3
P911	H	Cl	CN	Cl	CF_3	H	Cl	O	CH_2	O	CH_2CF_3
P912	H	Cl	CN	Cl	CF_3	H	CF_3	O	CH_2	O	CH_2CF_3
P913	H	Cl	CN	Cl	CF_3	H	CH_3	O	CH_2	O	CH_2CF_3
P914	H	Cl	CN	Cl	CF_3	H	Br	O	CH_2	S	CH_2CF_3

P915	H	Cl	CN	Cl	CF_3	H	Cl	O	CH_2	S	CH_2CF_3
P916	H	Cl	CN	Cl	CF_3	H	CF_3	O	CH_2	S	CH_2CF_3
P917	H	Cl	CN	Cl	CF_3	H	CH_3	O	CH_2	S	CH_2CF_3
P918	H	Cl	CN	Cl	CF_3	H	Br	S	CH_2	O	CH_2CF_3
P919	H	Cl	CN	Cl	CF_3	H	Cl	S	CH_2	O	CH_2CF_3
P920	H	Cl	CN	Cl	CF_3	H	CF_3	S	CH_2	O	CH_2CF_3
P921	H	Cl	CN	Cl	CF_3	H	CH_3	S	CH_2	O	CH_2CF_3
P922	H	Cl	CN	Cl	CF_3	H	Br	O	CH_2	O	CH_2CHF_2
P923	H	Cl	CN	Cl	CF_3	H	Cl	O	CH_2	O	CH_2CHF_2
P924	H	Cl	CN	Cl	CF_3	H	CF_3	O	CH_2	O	CH_2CHF_2
P925	H	Cl	CN	Cl	CF_3	H	CH_3	O	CH_2	O	CH_2CHF_2
P926	H	Cl	CN	Cl	CF_3	CF_3	Br	O	CH_2	O	CH_2CF_3
P927	H	Cl	CN	Cl	CF_3	CF_3	Cl	O	CH_2	O	CH_2CF_3
P928	H	Cl	CN	Cl	CF_3	CF_3	CF_3	O	CH_2	O	CH_2CF_3
P929	H	Cl	CN	Cl	CF_3	CF_3	CH_3	O	CH_2	O	CH_2CF_3
P930	H	Cl	CN	Cl	CF_2CF_3	H	Br	O	CH_2	O	CH_2CF_3
P931	H	Cl	CN	Cl	CF_2CF_3	H	Cl	O	CH_2	O	CH_2CF_3
P932	H	Cl	CN	Cl	CF_2CF_3	H	CF_3	O	CH_2	O	CH_2CF_3
P933	H	Cl	CN	Cl	CF_2CF_3	H	CH_3	O	CH_2	O	CH_2CF_3
P934	H	Cl	CN	Cl	CF_3	H	Br	O	CH_2	O	$CH(CH_3)CF_3$
P935	H	Cl	CN	Cl	CF_3	H	Cl	O	CH_2	O	$CH(CH_3)CF_3$
P936	H	Cl	CN	Cl	CF_3	H	CF_3	O	CH_2	O	$CH(CH_3)CF_3$
P937	H	Cl	CN	Cl	CF_3	H	CH_3	O	CH_2	O	$CH(CH_3)CF_3$
P938	H	Cl	CN	Cl	CF_3	CF_3	Br	O	$CH(CH_3)$	O	CH_2CF_3
P939	H	Cl	CN	Cl	CF_3	CF_3	Cl	O	$CH(CH_3)$	O	CH_2CF_3
P940	H	Cl	CN	Cl	CF_3	CF_3	CF_3	O	$CH(CH_3)$	O	CH_2CF_3
P941	H	Cl	CN	Cl	CF_3	CF_3	CH_3	O	$CH(CH_3)$	O	CH_2CF_3

P942	H	Cl	CN	Cl	CF_2CF_3	H	Br	O	CH(CH$_3$)	O	CH_2CF_3
P943	H	Cl	CN	Cl	CF_2CF_3	H	Cl	O	CH(CH$_3$)	O	CH_2CF_3
P944	H	Cl	CN	Cl	CF_2CF_3	H	CF_3	O	CH(CH$_3$)	O	CH_2CF_3
P945	H	Cl	CN	Cl	CF_2CF_3	H	CH_3	O	CH(CH$_3$)	O	CH_2CF_3
P946	H	Cl	CN	Cl	CF_3	H	Br	O	CH(CH$_3$)	O	CH_2CF_3
P947	H	Cl	CN	Cl	CF_3	H	Cl	O	CH(CH$_3$)	O	CH_2CF_3
P948	H	Cl	CN	Cl	CF_3	H	CF_3	O	CH(CH$_3$)	O	CH_2CF_3
P949	H	Cl	CN	Cl	CF_3	H	CH_3	O	CH(CH$_3$)	O	CH_2CF_3
P950	H	Cl	CN	Cl	CF_3	H	Br	O	CH(CH$_3$)	S	CH_2CF_3
P951	H	Cl	CN	Cl	CF_3	H	Cl	O	CH(CH$_3$)	S	CH_2CF_3
P952	H	Cl	CN	Cl	CF_3	H	CF_3	O	CH(CH$_3$)	S	CH_2CF_3
P953	H	Cl	CN	Cl	CF_3	H	CH_3	O	CH(CH$_3$)	S	CH_2CF_3
P954	H	Cl	CN	Cl	CF_3	H	Br	S	CH(CH$_3$)	O	CH_2CF_3
P955	H	Cl	CN	Cl	CF_3	H	Cl	S	CH(CH$_3$)	O	CH_2CF_3
P956	H	Cl	CN	Cl	CF_3	H	CF_3	S	CH(CH$_3$)	O	CH_2CF_3
P957	H	Cl	CN	Cl	CF_3	H	CH_3	S	CH(CH$_3$)	O	CH_2CF_3
P958	H	Cl	CN	Cl	CF_3	H	Br	O	CH(CH$_3$)	O	CH_2CHF_2
P959	H	Cl	CN	Cl	CF_3	H	Cl	O	CH(CH$_3$)	O	CH_2CHF_2
P960	H	Cl	CN	Cl	CF_3	H	CF_3	O	CH(CH$_3$)	O	CH_2CHF_2
P961	H	Cl	CN	Cl	CF_3	H	CH_3	O	CH(CH$_3$)	O	CH_2CHF_2
P962	H	Cl	CN	Cl	CF_3	H	Br	O	CH(CH$_3$)	O	$CH(CH_3)CF_3$
P963	H	Cl	CN	Cl	CF_3	H	Cl	O	CH(CH$_3$)	O	$CH(CH_3)CF_3$
P964	H	Cl	CN	Cl	CF_3	H	CF_3	O	CH(CH$_3$)	O	$CH(CH_3)CF_3$
P965	H	Cl	CN	Cl	CF_3	H	CH_3	O	CH(CH$_3$)	O	$CH(CH_3)CF_3$
P966	H	Cl	CN	Cl	CF_3	H	Br	O	CH(CH$_3$)	O	$CH_2CH_2CF_3$
P967	H	Cl	CN	Cl	CF_3	H	Cl	O	CH(CH$_3$)	O	$CH_2CH_2CF_3$
P968	H	Cl	CN	Cl	CF_3	H	CF_3	O	CH(CH$_3$)	O	$CH_2CH_2CF_3$

P969	H	Cl	CN	Cl	CF_3	H	CH_3	O	CH (CH_3)	O	$CH_2CH_2CF_3$
P970	H	Cl	CN	Cl	CF_3	H	Br	O	$CH(CH_2CH_3)$	O	CH_2CF_3
P971	H	Cl	CN	Cl	CF_3	H	Cl	O	$CH(CH_2CH_3)$	O	CH_2CF_3
P972	H	Cl	CN	Cl	CF_3	H	CF_3	O	$CH(CH_2CH_3)$	O	CH_2CF_3
P973	H	Cl	CN	Cl	CF_3	H	CH_3	O	$CH(CH_2CH_3)$	O	CH_2CF_3
P974	H	Cl	CN	Cl	CF_3	H	Br	O	$C(CH_3)_2$	O	CH_2CF_3
P975	H	Cl	CN	Cl	CF_3	H	Cl	O	$C(CH_3)_2$	O	CH_2CF_3
P976	H	Cl	CN	Cl	CF_3	H	CF_3	O	$C(CH_3)_2$	O	CH_2CF_3
P977	H	Cl	CN	Cl	CF_3	H	CH_3	O	$C(CH_3)_2$	O	CH_2CF_3
P978	H	Cl	CN	Cl	CF_3	H	Br	O	CH_2CH_2	O	CH_2CF_3
P979	H	Cl	CN	Cl	CF_3	H	Cl	O	CH_2CH_2	O	CH_2CF_3
P980	H	Cl	CN	Cl	CF_3	H	CF_3	O	CH_2CH_2	O	CH_2CF_3
P981	H	Cl	CN	Cl	CF_3	H	CH_3	O	CH_2CH_2	O	CH_2CF_3
P982	H	CH_3	H	Br	CF_3	H	Br	O	CH_2	O	CH_2CF_3
P983	H	CH_3	H	Br	CF_3	H	Cl	O	CH_2	O	CH_2CF_3
P984	H	CH_3	H	Br	CF_3	H	CF_3	O	CH_2	O	CH_2CF_3
P985	H	CH_3	H	Br	CF_3	H	CH_3	O	CH_2	O	CH_2CF_3
P986	H	CH_3	H	Br	CF_3	H	Br	O	CH_2	S	CH_2CF_3
P987	H	CH_3	H	Br	CF_3	H	Cl	O	CH_2	S	CH_2CF_3
P988	H	CH_3	H	Br	CF_3	H	CF_3	O	CH_2	S	CH_2CF_3
P989	H	CH_3	H	Br	CF_3	H	CH_3	O	CH_2	S	CH_2CF_3
P990	H	CH_3	H	Br	CF_3	H	Br	S	CH_2	O	CH_2CF_3
P991	H	CH_3	H	Br	CF_3	H	Cl	S	CH_2	O	CH_2CF_3
P992	H	CH_3	H	Br	CF_3	H	CF_3	S	CH_2	O	CH_2CF_3
P993	H	CH_3	H	Br	CF_3	H	CH_3	S	CH_2	O	CH_2CF_3
P994	H	CH_3	H	Br	CF_3	H	Br	O	CH_2	O	CH_2CHF_2
P995	H	CH_3	H	Br	CF_3	H	Cl	O	CH_2	O	CH_2CHF_2

P996	H	CH_3	H	Br	CF_3	H	CF_3	O	CH_2	O	CH_2CHF_2
P997	H	CH_3	H	Br	CF_3	H	CH_3	O	CH_2	O	CH_2CHF_2
P998	H	CH_3	H	Br	CF_3	CF_3	Br	O	CH_2	O	CH_2CF_3
P999	H	CH_3	H	Br	CF_3	CF_3	Cl	O	CH_2	O	CH_2CF_3
P1000	H	CH_3	H	Br	CF_3	CF_3	CF_3	O	CH_2	O	CH_2CF_3
P1001	H	CH_3	H	Br	CF_3	CF_3	CH_3	O	CH_2	O	CH_2CF_3
P1002	H	CH_3	H	Br	CF_2CF_3	H	Br	O	CH_2	O	CH_2CF_3
P1003	H	CH_3	H	Br	CF_2CF_3	H	Cl	O	CH_2	O	CH_2CF_3
P1004	H	CH_3	H	Br	CF_2CF_3	H	CF_3	O	CH_2	O	CH_2CF_3
P1005	H	CH_3	H	Br	CF_2CF_3	H	CH_3	O	CH_2	O	CH_2CF_3
P1006	H	CH_3	H	Br	CF_3	H	Br	O	CH_2	O	$CH(CH_3)CF_3$
P1007	H	CH_3	H	Br	CF_3	H	Cl	O	CH_2	O	$CH(CH_3)CF_3$
P1008	H	CH_3	H	Br	CF_3	H	CF_3	O	CH_2	O	$CH(CH_3)CF_3$
P1009	H	CH_3	H	Br	CF_3	H	CH_3	O	CH_2	O	$CH(CH_3)CF_3$
P1010	H	CH_3	H	Br	CF_3	CF_3	Br	O	$CH(CH_3)$	O	CH_2CF_3
P1011	H	CH_3	H	Br	CF_3	CF_3	Cl	O	$CH(CH_3)$	O	CH_2CF_3
P1012	H	CH_3	H	Br	CF_3	CF_3	CF_3	O	$CH(CH_3)$	O	CH_2CF_3
P1013	H	CH_3	H	Br	CF_3	CF_3	CH_3	O	$CH(CH_3)$	O	CH_2CF_3
P1014	H	CH_3	H	Br	CF_2CF_3	H	Br	O	$CH(CH_3)$	O	CH_2CF_3
P1015	H	CH_3	H	Br	CF_2CF_3	H	Cl	O	$CH(CH_3)$	O	CH_2CF_3
P1016	H	CH_3	H	Br	CF_2CF_3	H	CF_3	O	$CH(CH_3)$	O	CH_2CF_3
P1017	H	CH_3	H	Br	CF_2CF_3	H	CH_3	O	$CH(CH_3)$	O	CH_2CF_3
P1018	H	CH_3	H	Br	CF_3	H	Br	O	$CH(CH_3)$	O	CH_2CF_3
P1019	H	CH_3	H	Br	CF_3	H	Cl	O	$CH(CH_3)$	O	CH_2CF_3
P1020	H	CH_3	H	Br	CF_3	H	CF_3	O	$CH(CH_3)$	O	CH_2CF_3
P1021	H	CH_3	H	Br	CF_3	H	CH_3	O	$CH(CH_3)$	O	CH_2CF_3
P1022	H	CH_3	H	Br	CF_3	H	Br	O	$CH(CH_3)$	S	CH_2CF_3

P1023	H	CH_3	H	Br	CF_3	H	Cl	O	CH (CH_3)	S	CH_2CF_3
P1024	H	CH_3	H	Br	CF_3	H	CF_3	O	CH(CH_3)	S	CH_2CF_3
P1025	H	CH_3	H	Br	CF_3	H	CH_3	O	CH (CH_3)	S	CH_2CF_3
P1026	H	CH_3	H	Br	CF_3	H	Br	S	CH(CH_3)	O	CH_2CF_3
P1027	H	CH_3	H	Br	CF_3	H	Cl	S	CH (CH_3)	O	CH_2CF_3
P1028	H	CH_3	H	Br	CF_3	H	CF_3	S	CH(CH_3)	O	CH_2CF_3
P1029	H	CH_3	H	Br	CF_3	H	CH_3	S	CH (CH_3)	O	CH_2CF_3
P1030	H	CH_3	H	Br	CF_3	H	Br	O	CH(CH_3)	O	CH_2CHF_2
P1031	H	CH_3	H	Br	CF_3	H	Cl	O	CH (CH_3)	O	CH_2CHF_2
P1032	H	CH_3	H	Br	CF_3	H	CF_3	O	CH(CH_3)	O	CH_2CHF_2
P1033	H	CH_3	H	Br	CF_3	H	CH_3	O	CH (CH_3)	O	CH_2CHF_2
P1034	H	CH_3	H	Br	CF_3	H	Br	O	CH(CH_3)	O	$CH(CH_3)CF_3$
P1035	H	CH_3	H	Br	CF_3	H	Cl	O	CH (CH_3)	O	$CH(CH_3)CF_3$
P1036	H	CH_3	H	Br	CF_3	H	CF_3	O	CH(CH_3)	O	$CH(CH_3)CF_3$
P1037	H	CH_3	H	Br	CF_3	H	CH_3	O	CH (CH_3)	O	$CH(CH_3)CF_3$
P1038	H	CH_3	H	Br	CF_3	H	Br	O	CH(CH_3)	O	$CH_2CH_2CF_3$
P1039	H	CH_3	H	Br	CF_3	H	Cl	O	CH (CH_3)	O	$CH_2CH_2CF_3$
P1040	H	CH_3	H	Br	CF_3	H	CF_3	O	CH(CH_3)	O	$CH_2CH_2CF_3$
P1041	H	CH_3	H	Br	CF_3	H	CH_3	O	CH (CH_3)	O	$CH_2CH_2CF_3$
P1042	H	CH_3	H	Br	CF_3	H	Br	O	CH(CH_2CH_3)	O	CH_2CF_3
P1043	H	CH_3	H	Br	CF_3	H	Cl	O	CH(CH_2CH_3)	O	CH_2CF_3
P1044	H	CH_3	H	Br	CF_3	H	CF_3	O	CH(CH_2CH_3)	O	CH_2CF_3
P1045	H	CH_3	H	Br	CF_3	H	CH_3	O	CH(CH_2CH_3)	O	CH_2CF_3
P1046	H	CH_3	H	Br	CF_3	H	Br	O	$C(CH_3)_2$	O	CH_2CF_3
P1047	H	CH_3	H	Br	CF_3	H	Cl	O	$C(CH_3)_2$	O	CH_2CF_3
P1048	H	CH_3	H	Br	CF_3	H	CF_3	O	$C(CH_3)_2$	O	CH_2CF_3
P1049	H	CH_3	H	Br	CF_3	H	CH_3	O	$C(CH_3)_2$	O	CH_2CF_3

P1050	H	CH_3	H	Br	CF_3	H	Br	O	CH_2CH_2	O	CH_2CF_3
P1051	H	CH_3	H	Br	CF_3	H	Cl	O	CH_2CH_2	O	CH_2CF_3
P1052	H	CH_3	H	Br	CF_3	H	CF_3	O	CH_2CH_2	O	CH_2CF_3
P1053	H	CH_3	H	Br	CF_3	H	CH_3	O	CH_2CH_2	O	CH_2CF_3
P1054	H	H	F	CH_3	CF_3	H	Br	O	CH_2	O	CH_2CF_3
P1055	H	H	F	CH_3	CF_3	H	Cl	O	CH_2	O	CH_2CF_3
P1056	H	H	F	CH_3	CF_3	H	CF_3	O	CH_2	O	CH_2CF_3
P1057	H	H	F	CH_3	CF_3	H	CH_3	O	CH_2	O	CH_2CF_3
P1058	H	H	F	CH_3	CF_3	H	Br	O	CH_2	S	CH_2CF_3
P1059	H	H	F	CH_3	CF_3	H	Cl	O	CH_2	S	CH_2CF_3
P1060	H	H	F	CH_3	CF_3	H	CF_3	O	CH_2	S	CH_2CF_3
P1061	H	H	F	CH_3	CF_3	H	CH_3	O	CH_2	S	CH_2CF_3
P1062	H	H	F	CH_3	CF_3	H	Br	S	CH_2	O	CH_2CF_3
P1063	H	H	F	CH_3	CF_3	H	Cl	S	CH_2	O	CH_2CF_3
P1064	H	H	F	CH_3	CF_3	H	CF_3	S	CH_2	O	CH_2CF_3
P1065	H	H	F	CH_3	CF_3	H	CH_3	S	CH_2	O	CH_2CF_3
P1066	H	H	F	CH_3	CF_3	H	Br	O	CH_2	O	CH_2CHF_2
P1067	H	H	F	CH_3	CF_3	H	Cl	O	CH_2	O	CH_2CHF_2
P1068	H	H	F	CH_3	CF_3	H	CF_3	O	CH_2	O	CH_2CHF_2
P1069	H	H	F	CH_3	CF_3	H	CH_3	O	CH_2	O	CH_2CHF_2
P1070	H	H	F	CH_3	CF_3	CF_3	Br	O	CH_2	O	CH_2CF_3
P1071	H	H	F	CH_3	CF_3	CF_3	Cl	O	CH_2	O	CH_2CF_3
P1072	H	H	F	CH_3	CF_3	CF_3	CF_3	O	CH_2	O	CH_2CF_3
P1073	H	H	F	CH_3	CF_3	CF_3	CH_3	O	CH_2	O	CH_2CF_3
P1074	H	H	F	CH_3	CF_2CF_3	H	Br	O	CH_2	O	CH_2CF_3
P1075	H	H	F	CH_3	CF_2CF_3	H	Cl	O	CH_2	O	CH_2CF_3
P1076	H	H	F	CH_3	CF_2CF_3	H	CF_3	O	CH_2	O	CH_2CF_3

P1077	H	H	F	CH_3	CF_2CF_3	H	CH_3	O	CH_2	O	CH_2CF_3
P1078	H	H	F	CH_3	CF_3	H	Br	O	CH_2	O	$CH(CH_3)CF_3$
P1079	H	H	F	CH_3	CF_3	H	Cl	O	CH_2	O	$CH(CH_3)CF_3$
P1080	H	H	F	CH_3	CF_3	H	CF_3	O	CH_2	O	$CH(CH_3)CF_3$
P1081	H	H	F	CH_3	CF_3	H	CH_3	O	CH_2	O	$CH(CH_3)CF_3$
P1082	H	H	F	CH_3	CF_3	CF_3	Br	O	$CH(CH_3)$	O	CH_2CF_3
P1083	H	H	F	CH_3	CF_3	CF_3	Cl	O	$CH(CH_3)$	O	CH_2CF_3
P1084	H	H	F	CH_3	CF_3	CF_3	CF_3	O	$CH(CH_3)$	O	CH_2CF_3
P1085	H	H	F	CH_3	CF_3	CF_3	CH_3	O	$CH(CH_3)$	O	CH_2CF_3
P1086	H	H	F	CH_3	CF_2CF_3	H	Br	O	$CH(CH_3)$	O	CH_2CF_3
P1087	H	H	F	CH_3	CF_2CF_3	H	Cl	O	$CH(CH_3)$	O	CH_2CF_3
P1088	H	H	F	CH_3	CF_2CF_3	H	CF_3	O	$CH(CH_3)$	O	CH_2CF_3
P1089	H	H	F	CH_3	CF_2CF_3	H	CH_3	O	$CH(CH_3)$	O	CH_2CF_3
P1090	H	H	F	CH_3	CF_3	H	Br	O	$CH(CH_3)$	O	CH_2CF_3
P1091	H	H	F	CH_3	CF_3	H	Cl	O	$CH(CH_3)$	O	CH_2CF_3
P1092	H	H	F	CH_3	CF_3	H	CF_3	O	$CH(CH_3)$	O	CH_2CF_3
P1093	H	H	F	CH_3	CF_3	H	CH_3	O	$CH(CH_3)$	O	CH_2CF_3
P1094	H	H	F	CH_3	CF_3	H	Br	O	$CH(CH_3)$	S	CH_2CF_3
P1095	H	H	F	CH_3	CF_3	H	Cl	O	$CH(CH_3)$	S	CH_2CF_3
P1096	H	H	F	CH_3	CF_3	H	CF_3	O	$CH(CH_3)$	S	CH_2CF_3
P1097	H	H	F	CH_3	CF_3	H	CH_3	O	$CH(CH_3)$	S	CH_2CF_3
P1098	H	H	F	CH_3	CF_3	H	Br	S	$CH(CH_3)$	O	CH_2CF_3
P1099	H	H	F	CH_3	CF_3	H	Cl	S	$CH(CH_3)$	O	CH_2CF_3
P1100	H	H	F	CH_3	CF_3	H	CF_3	S	$CH(CH_3)$	O	CH_2CF_3
P1101	H	H	F	CH_3	CF_3	H	CH_3	S	$CH(CH_3)$	O	CH_2CF_3
P1102	H	H	F	CH_3	CF_3	H	Br	O	$CH(CH_3)$	O	CH_2CHF_2
P1103	H	H	F	CH_3	CF_3	H	Cl	O	$CH(CH_3)$	O	CH_2CHF_2

P1104	H	H	F	CH_3	CF_3	H	CF_3	O	$CH(CH_3)$	O	CH_2CHF_2
P1105	H	H	F	CH_3	CF_3	H	CH_3	O	$CH(CH_3)$	O	CH_2CHF_2
P1106	H	H	F	CH_3	CF_3	H	Br	O	$CH(CH_3)$	O	$CH(CH_3)CF_3$
P1107	H	H	F	CH_3	CF_3	H	Cl	O	$CH(CH_3)$	O	$CH(CH_3)CF_3$
P1108	H	H	F	CH_3	CF_3	H	CF_3	O	$CH(CH_3)$	O	$CH(CH_3)CF_3$
P1109	H	H	F	CH_3	CF_3	H	CH_3	O	$CH(CH_3)$	O	$CH(CH_3)CF_3$
P1110	H	H	F	CH_3	CF_3	H	Br	O	$CH(CH_3)$	O	$CH_2CH_2CF_3$
P1111	H	H	F	CH_3	CF_3	H	Cl	O	$CH(CH_3)$	O	$CH_2CH_2CF_3$
P1112	H	H	F	CH_3	CF_3	H	CF_3	O	$CH(CH_3)$	O	$CH_2CH_2CF_3$
P1113	H	H	F	CH_3	CF_3	H	CH_3	O	$CH(CH_3)$	O	$CH_2CH_2CF_3$
P1114	H	H	F	CH_3	CF_3	H	Br	O	$CH(CH_2CH_3)$	O	CH_2CF_3
P1115	H	H	F	CH_3	CF_3	H	Cl	O	$CH(CH_2CH_3)$	O	CH_2CF_3
P1116	H	H	F	CH_3	CF_3	H	CF_3	O	$CH(CH_2CH_3)$	O	CH_2CF_3
P1117	H	H	F	CH_3	CF_3	H	CH_3	O	$CH(CH_2CH_3)$	O	CH_2CF_3
P1118	H	H	F	CH_3	CF_3	H	Br	O	$C(CH_3)_2$	O	CH_2CF_3
P1119	H	H	F	CH_3	CF_3	H	Cl	O	$C(CH_3)_2$	O	CH_2CF_3
P1120	H	H	F	CH_3	CF_3	H	CF_3	O	$C(CH_3)_2$	O	CH_2CF_3
P1121	H	H	F	CH_3	CF_3	H	CH_3	O	$C(CH_3)_2$	O	CH_2CF_3
P1122	H	H	F	CH_3	CF_3	H	Br	O	CH_2CH_2	O	CH_2CF_3
P1123	H	H	F	CH_3	CF_3	H	Cl	O	CH_2CH_2	O	CH_2CF_3
P1124	H	H	F	CH_3	CF_3	H	CF_3	O	CH_2CH_2	O	CH_2CF_3
P1125	H	H	F	CH_3	CF_3	H	CH_3	O	CH_2CH_2	O	CH_2CF_3
P1126	H	H	F	Cl	CF_3	H	Cl	O	CH_2	O	CH_2CF_3
P1127	H	H	F	Cl	CF_3	H	CF_3	O	CH_2	O	CH_2CF_3
P1128	H	H	F	Cl	CF_3	H	CH_3	O	CH_2	O	CH_2CF_3
P1129	H	H	F	Cl	CF_3	H	Br	O	CH_2	S	CH_2CF_3
P1130	H	H	F	Cl	CF_3	H	Cl	O	CH_2	S	CH_2CF_3

P1131	H	H	F	Cl	CF_3	H	CF_3	O	CH_2	S	CH_2CF_3
P1132	H	H	F	Cl	CF_3	H	CH_3	O	CH_2	S	CH_2CF_3
P1133	H	H	F	Cl	CF_3	H	Br	S	CH_2	O	CH_2CF_3
P1134	H	H	F	Cl	CF_3	H	Cl	S	CH_2	O	CH_2CF_3
P1135	H	H	F	Cl	CF_3	H	CF_3	S	CH_2	O	CH_2CF_3
P1136	H	H	F	Cl	CF_3	H	CH_3	S	CH_2	O	CH_2CF_3
P1137	H	H	F	Cl	CF_3	H	Br	O	CH_2	O	CH_2CHF_2
P1138	H	H	F	Cl	CF_3	H	Cl	O	CH_2	O	CH_2CHF_2
P1139	H	H	F	Cl	CF_3	H	CF_3	O	CH_2	O	CH_2CHF_2
P1140	H	H	F	Cl	CF_3	H	CH_3	O	CH_2	O	CH_2CHF_2
P1141	H	H	F	Cl	CF_3	CF_3	Br	O	CH_2	O	CH_2CF_3
P1142	H	H	F	Cl	CF_3	CF_3	Cl	O	CH_2	O	CH_2CF_3
P1143	H	H	F	Cl	CF_3	CF_3	CF_3	O	CH_2	O	CH_2CF_3
P1144	H	H	F	Cl	CF_3	CF_3	CH_3	O	CH_2	O	CH_2CF_3
P1145	H	H	F	Cl	CF_2CF_3	H	Br	O	CH_2	O	CH_2CF_3
P1146	H	H	F	Cl	CF_2CF_3	H	Cl	O	CH_2	O	CH_2CF_3
P1147	H	H	F	Cl	CF_2CF_3	H	CF_3	O	CH_2	O	CH_2CF_3
P1148	H	H	F	Cl	CF_2CF_3	H	CH_3	O	CH_2	O	CH_2CF_3
P1149	H	H	F	Cl	CF_3	H	Br	O	CH_2	O	$CH(CH_3)CF_3$
P1150	H	H	F	Cl	CF_3	H	Cl	O	CH_2	O	$CH(CH_3)CF_3$
P1151	H	H	F	Cl	CF_3	H	CF_3	O	CH_2	O	$CH(CH_3)CF_3$
P1152	H	H	F	Cl	CF_3	H	CH_3	O	CH_2	O	$CH(CH_3)CF_3$
P1153	H	H	F	Cl	CF_3	CF_3	Br	O	$CH(CH_3)$	O	CH_2CF_3
P1154	H	H	F	Cl	CF_3	CF_3	Cl	O	$CH(CH_3)$	O	CH_2CF_3
P1155	H	H	F	Cl	CF_3	CF_3	CF_3	O	$CH(CH_3)$	O	CH_2CF_3
P1156	H	H	F	Cl	CF_3	CF_3	CH_3	O	$CH(CH_3)$	O	CH_2CF_3
P1157	H	H	F	Cl	CF_2CF_3	H	Br	O	$CH(CH_3)$	O	CH_2CF_3

P1158	H	H	F	Cl	CF_2CF_3	H	Cl	O	CH (CH_3)	O	CH_2CF_3
P1159	H	H	F	Cl	CF_2CF_3	H	CF_3	O	CH (CH_3)	O	CH_2CF_3
P1160	H	H	F	Cl	CF_2CF_3	H	CH_3	O	CH (CH_3)	O	CH_2CF_3
P1161	H	H	F	Cl	CF_3	H	Br	O	$CH(CH_3)$	O	CH_2CF_3
P1162	H	H	F	Cl	CF_3	H	Cl	O	CH (CH_3)	O	CH_2CF_3
P1163	H	H	F	Cl	CF_3	H	CF_3	O	$CH(CH_3)$	O	CH_2CF_3
P1164	H	H	F	Cl	CF_3	H	CH_3	O	CH (CH_3)	O	CH_2CF_3
P1165	H	H	F	Cl	CF_3	H	Br	O	$CH(CH_3)$	S	CH_2CF_3
P1166	H	H	F	Cl	CF_3	H	Cl	O	CH (CH_3)	S	CH_2CF_3
P1167	H	H	F	Cl	CF_3	H	CF_3	O	$CH(CH_3)$	S	CH_2CF_3
P1168	H	H	F	Cl	CF_3	H	CH_3	O	CH (CH_3)	S	CH_2CF_3
P1169	H	H	F	Cl	CF_3	H	Br	S	$CH(CH_3)$	O	CH_2CF_3
P1170	H	H	F	Cl	CF_3	H	Cl	S	CH (CH_3)	O	CH_2CF_3
P1171	H	H	F	Cl	CF_3	H	CF_3	S	$CH(CH_3)$	O	CH_2CF_3
P1172	H	H	F	Cl	CF_3	H	CH_3	S	CH (CH_3)	O	CH_2CF_3
P1173	H	H	F	Cl	CF_3	H	Br	O	$CH(CH_3)$	O	CH_2CHF_2
P1174	H	H	F	Cl	CF_3	H	Cl	O	CH (CH_3)	O	CH_2CHF_2
P1175	H	H	F	Cl	CF_3	H	CF_3	O	$CH(CH_3)$	O	CH_2CHF_2
P1176	H	H	F	Cl	CF_3	H	CH_3	O	CH (CH_3)	O	CH_2CHF_2
P1177	H	H	F	Cl	CF_3	H	Br	O	$CH(CH_3)$	O	$CH(CH_3)CF_3$
P1178	H	H	F	Cl	CF_3	H	Cl	O	CH (CH_3)	O	$CH(CH_3)CF_3$
P1179	H	H	F	Cl	CF_3	H	CF_3	O	$CH(CH_3)$	O	$CH(CH_3)CF_3$
P1180	H	H	F	Cl	CF_3	H	CH_3	O	CH (CH_3)	O	$CH(CH_3)CF_3$
P1181	H	H	F	Cl	CF_3	H	Br	O	$CH(CH_3)$	O	$CH_2CH_2CF_3$
P1182	H	H	F	Cl	CF_3	H	Cl	O	CH (CH_3)	O	$CH_2CH_2CF_3$
P1183	H	H	F	Cl	CF_3	H	CF_3	O	$CH(CH_3)$	O	$CH_2CH_2CF_3$
P1184	H	H	F	Cl	CF_3	H	CH_3	O	CH (CH_3)	O	$CH_2CH_2CF_3$

P1185	H	H	F	Cl	CF_3	H	Br	O	$CH(CH_2CH_3)$	O	CH_2CF_3
P1186	H	H	F	Cl	CF_3	H	Cl	O	$CH(CH_2CH_3)$	O	CH_2CF_3
P1187	H	H	F	Cl	CF_3	H	CF_3	O	$CH(CH_2CH_3)$	O	CH_2CF_3
P1188	H	H	F	Cl	CF_3	H	CH_3	O	$CH(CH_2CH_3)$	O	CH_2CF_3
P1189	H	H	F	Cl	CF_3	H	Br	O	$C(CH_3)_2$	O	CH_2CF_3
P1190	H	H	F	Cl	CF_3	H	Cl	O	$C(CH_3)_2$	O	CH_2CF_3
P1191	H	H	F	Cl	CF_3	H	CF_3	O	$C(CH_3)_2$	O	CH_2CF_3
P1192	H	H	F	Cl	CF_3	H	CH_3	O	$C(CH_3)_2$	O	CH_2CF_3
P1193	H	H	F	Cl	CF_3	H	Br	O	CH_2CH_2	O	CH_2CF_3
P1194	H	H	F	Cl	CF_3	H	Cl	O	CH_2CH_2	O	CH_2CF_3
P1195	H	H	F	Cl	CF_3	H	CF_3	O	CH_2CH_2	O	CH_2CF_3
P1196	H	H	F	Cl	CF_3	H	CH_3	O	CH_2CH_2	O	CH_2CF_3
P1197	H	F	F	F	CF_3	H	Br	O	CH_2	O	CH_2CF_3
P1198	H	F	F	F	CF_3	H	Cl	O	CH_2	O	CH_2CF_3
P1199	H	F	F	F	CF_3	H	CF_3	O	CH_2	O	CH_2CF_3
P1200	H	F	F	F	CF_3	H	CH_3	O	CH_2	O	CH_2CF_3
P1201	H	F	F	F	CF_3	H	Br	O	CH_2	S	CH_2CF_3
P1202	H	F	F	F	CF_3	H	Cl	O	CH_2	S	CH_2CF_3
P1203	H	F	F	F	CF_3	H	CF_3	O	CH_2	S	CH_2CF_3
P1204	H	F	F	F	CF_3	H	CH_3	O	CH_2	S	CH_2CF_3
P1205	H	F	F	F	CF_3	H	Br	S	CH_2	O	CH_2CF_3
P1206	H	F	F	F	CF_3	H	Cl	S	CH_2	O	CH_2CF_3
P1207	H	F	F	F	CF_3	H	CF_3	S	CH_2	O	CH_2CF_3
P1208	H	F	F	F	CF_3	H	CH_3	S	CH_2	O	CH_2CF_3
P1209	H	F	F	F	CF_3	H	Br	O	CH_2	O	CH_2CHF_2
P1210	H	F	F	F	CF_3	H	Cl	O	CH_2	O	CH_2CHF_2
P1211	H	F	F	F	CF_3	H	CF_3	O	CH_2	O	CH_2CHF_2

P1212	H	F	F	F	CF_3	H	CH_3	O	CH_2	O	CH_2CHF_2
P1213	H	F	F	F	CF_3	CF_3	Br	O	CH_2	O	CH_2CF_3
P1214	H	F	F	F	CF_3	CF_3	Cl	O	CH_2	O	CH_2CF_3
P1215	H	F	F	F	CF_3	CF_3	CF_3	O	CH_2	O	CH_2CF_3
P1216	H	F	F	F	CF_3	CF_3	CH_3	O	CH_2	O	CH_2CF_3
P1217	H	F	F	F	CF_2CF_3	H	Br	O	CH_2	O	CH_2CF_3
P1218	H	F	F	F	CF_2CF_3	H	Cl	O	CH_2	O	CH_2CF_3
P1219	H	F	F	F	CF_2CF_3	H	CF_3	O	CH_2	O	CH_2CF_3
P1220	H	F	F	F	CF_2CF_3	H	CH_3	O	CH_2	O	CH_2CF_3
P1221	H	F	F	F	CF_3	H	Br	O	CH_2	O	$CH(CH_3)CF_3$
P1222	H	F	F	F	CF_3	H	Cl	O	CH_2	O	$CH(CH_3)CF_3$
P1223	H	F	F	F	CF_3	H	CF_3	O	CH_2	O	$CH(CH_3)CF_3$
P1224	H	F	F	F	CF_3	H	CH_3	O	CH_2	O	$CH(CH_3)CF_3$
P1225	H	F	F	F	CF_3	CF_3	Br	O	$CH(CH_3)$	O	CH_2CF_3
P1226	H	F	F	F	CF_3	CF_3	Cl	O	$CH(CH_3)$	O	CH_2CF_3
P1227	H	F	F	F	CF_3	CF_3	CF_3	O	$CH(CH_3)$	O	CH_2CF_3
P1228	H	F	F	F	CF_3	CF_3	CH_3	O	$CH(CH_3)$	O	CH_2CF_3
P1229	H	F	F	F	CF_2CF_3	H	Br	O	$CH(CH_3)$	O	CH_2CF_3
P1230	H	F	F	F	CF_2CF_3	H	Cl	O	$CH(CH_3)$	O	CH_2CF_3
P1231	H	F	F	F	CF_2CF_3	H	CF_3	O	$CH(CH_3)$	O	CH_2CF_3
P1232	H	F	F	F	CF_2CF_3	H	CH_3	O	$CH(CH_3)$	O	CH_2CF_3
P1233	H	F	F	F	CF_3	H	Br	O	$CH(CH_3)$	O	CH_2CF_3
P1234	H	F	F	F	CF_3	H	Cl	O	$CH(CH_3)$	O	CH_2CF_3
P1235	H	F	F	F	CF_3	H	CF_3	O	$CH(CH_3)$	O	CH_2CF_3
P1236	H	F	F	F	CF_3	H	CH_3	O	$CH(CH_3)$	O	CH_2CF_3
P1237	H	F	F	F	CF_3	H	Br	O	$CH(CH_3)$	S	CH_2CF_3
P1238	H	F	F	F	CF_3	H	Cl	O	$CH(CH_3)$	S	CH_2CF_3

P1239	H	F	F	F	CF_3	H	CF_3	O	$CH(CH_3)$	S	CH_2CF_3
P1240	H	F	F	F	CF_3	H	CH_3	O	$CH(CH_3)$	S	CH_2CF_3
P1241	H	F	F	F	CF_3	H	Br	S	$CH(CH_3)$	O	CH_2CF_3
P1242	H	F	F	F	CF_3	H	Cl	S	$CH(CH_3)$	O	CH_2CF_3
P1243	H	F	F	F	CF_3	H	CF_3	S	$CH(CH_3)$	O	CH_2CF_3
P1244	H	F	F	F	CF_3	H	CH_3	S	$CH(CH_3)$	O	CH_2CF_3
P1245	H	F	F	F	CF_3	H	Br	O	$CH(CH_3)$	O	CH_2CHF_2
P1246	H	F	F	F	CF_3	H	Cl	O	$CH(CH_3)$	O	CH_2CHF_2
P1247	H	F	F	F	CF_3	H	CF_3	O	$CH(CH_3)$	O	CH_2CHF_2
P1248	H	F	F	F	CF_3	H	CH_3	O	$CH(CH_3)$	O	CH_2CHF_2
P1249	H	F	F	F	CF_3	H	Br	O	$CH(CH_3)$	O	$CH(CH_3)CF_3$
P1250	H	F	F	F	CF_3	H	Cl	O	$CH(CH_3)$	O	$CH(CH_3)CF_3$
P1251	H	F	F	F	CF_3	H	CF_3	O	$CH(CH_3)$	O	$CH(CH_3)CF_3$
P1252	H	F	F	F	CF_3	H	CH_3	O	$CH(CH_3)$	O	$CH(CH_3)CF_3$
P1253	H	F	F	F	CF_3	H	Br	O	$CH(CH_3)$	O	$CH_2CH_2CF_3$
P1254	H	F	F	F	CF_3	H	Cl	O	$CH(CH_3)$	O	$CH_2CH_2CF_3$
P1255	H	F	F	F	CF_3	H	CF_3	O	$CH(CH_3)$	O	$CH_2CH_2CF_3$
P1256	H	F	F	F	CF_3	H	CH_3	O	$CH(CH_3)$	O	$CH_2CH_2CF_3$
P1257	H	F	F	F	CF_3	H	Br	O	$CH(CH_2CH_3)$	O	CH_2CF_3
P1258	H	F	F	F	CF_3	H	Cl	O	$CH(CH_2CH_3)$	O	CH_2CF_3
P1259	H	F	F	F	CF_3	H	CF_3	O	$CH(CH_2CH_3)$	O	CH_2CF_3
P1260	H	F	F	F	CF_3	H	CH_3	O	$CH(CH_2CH_3)$	O	CH_2CF_3
P1261	H	F	F	F	CF_3	H	Br	O	$C(CH_3)_2$	O	CH_2CF_3
P1262	H	F	F	F	CF_3	H	Cl	O	$C(CH_3)_2$	O	CH_2CF_3
P1263	H	F	F	F	CF_3	H	CF_3	O	$C(CH_3)_2$	O	CH_2CF_3
P1264	H	F	F	F	CF_3	H	CH_3	O	$C(CH_3)_2$	O	CH_2CF_3
P1265	H	F	F	F	CF_3	H	Br	O	CH_2CH_2	O	CH_2CF_3

P1266	H	F	F	F	CF_3	H	Cl	O	CH_2CH_2	O	CH_2CF_3
P1267	H	F	F	F	CF_3	H	CF_3	O	CH_2CH_2	O	CH_2CF_3
P1268	H	F	F	F	CF_3	H	CH_3	O	CH_2CH_2	O	CH_2CF_3
P1269	H	CF_3	H	CF_3	CF_3	H	Br	O	CH_2	O	CH_2CF_3
P1270	H	CF_3	H	CF_3	CF_3	H	Cl	O	CH_2	O	CH_2CF_3
P1271	H	CF_3	H	CF_3	CF_3	H	CF_3	O	CH_2	O	CH_2CF_3
P1272	H	CF_3	H	CF_3	CF_3	H	CH_3	O	CH_2	O	CH_2CF_3
P1273	H	CF_3	H	CF_3	CF_3	H	Br	O	CH_2	S	CH_2CF_3
P1274	H	CF_3	H	CF_3	CF_3	H	Cl	O	CH_2	S	CH_2CF_3
P1275	H	CF_3	H	CF_3	CF_3	H	CF_3	O	CH_2	S	CH_2CF_3
P1276	H	CF_3	H	CF_3	CF_3	H	CH_3	O	CH_2	S	CH_2CF_3
P1277	H	CF_3	H	CF_3	CF_3	H	Br	S	CH_2	O	CH_2CF_3
P1278	H	CF_3	H	CF_3	CF_3	H	Cl	S	CH_2	O	CH_2CF_3
P1279	H	CF_3	H	CF_3	CF_3	H	CF_3	S	CH_2	O	CH_2CF_3
P1280	H	CF_3	H	CF_3	CF_3	H	CH_3	S	CH_2	O	CH_2CF_3
P1281	H	CF_3	H	CF_3	CF_3	H	Br	O	CH_2	O	CH_2CHF_2
P1282	H	CF_3	H	CF_3	CF_3	H	Cl	O	CH_2	O	CH_2CHF_2
P1283	H	CF_3	H	CF_3	CF_3	H	CF_3	O	CH_2	O	CH_2CHF_2
P1284	H	CF_3	H	CF_3	CF_3	H	CH_3	O	CH_2	O	CH_2CHF_2
P1285	H	CF_3	H	CF_3	CF_3	CF_3	Br	O	CH_2	O	CH_2CF_3
P1286	H	CF_3	H	CF_3	CF_3	CF_3	Cl	O	CH_2	O	CH_2CF_3
P1287	H	CF_3	H	CF_3	CF_3	CF_3	CF_3	O	CH_2	O	CH_2CF_3
P1288	H	CF_3	H	CF_3	CF_3	CF_3	CH_3	O	CH_2	O	CH_2CF_3
P1289	H	CF_3	H	CF_3	CF_2CF_3	H	Br	O	CH_2	O	CH_2CF_3
P1290	H	CF_3	H	CF_3	CF_2CF_3	H	Cl	O	CH_2	O	CH_2CF_3
P1291	H	CF_3	H	CF_3	CF_2CF_3	H	CF_3	O	CH_2	O	CH_2CF_3
P1292	H	CF_3	H	CF_3	CF_2CF_3	H	CH_3	O	CH_2	O	CH_2CF_3

P1293	H	CF_3	H	CF_3	CF_3	H	Br	O	CH_2	O	$CH(CH_3)CF_3$
P1294	H	CF_3	H	CF_3	CF_3	H	Cl	O	CH_2	O	$CH(CH_3)CF_3$
P1295	H	CF_3	H	CF_3	CF_3	H	CF_3	O	CH_2	O	$CH(CH_3)CF_3$
P1296	H	CF_3	H	CF_3	CF_3	H	CH_3	O	CH_2	O	$CH(CH_3)CF_3$
P1297	H	CF_3	H	CF_3	CF_3	CF_3	Br	O	$CH(CH_3)$	O	CH_2CF_3
P1298	H	CF_3	H	CF_3	CF_3	CF_3	Cl	O	$CH(CH_3)$	O	CH_2CF_3
P1299	H	CF_3	H	CF_3	CF_3	CF_3	CF_3	O	$CH(CH_3)$	O	CH_2CF_3
P1300	H	CF_3	H	CF_3	CF_3	CF_3	CH_3	O	$CH(CH_3)$	O	CH_2CF_3
P1301	H	CF_3	H	CF_3	CF_2CF_3	H	Br	O	$CH(CH_3)$	O	CH_2CF_3
P1302	H	CF_3	H	CF_3	CF_2CF_3	H	Cl	O	$CH(CH_3)$	O	CH_2CF_3
P1303	H	CF_3	H	CF_3	CF_2CF_3	H	CF_3	O	$CH(CH_3)$	O	CH_2CF_3
P1304	H	CF_3	H	CF_3	CF_2CF_3	H	CH_3	O	$CH(CH_3)$	O	CH_2CF_3
P1305	H	CF_3	H	CF_3	CF_3	H	Cl	O	$CH(CH_3)$	O	CH_2CF_3
P1306	H	CF_3	H	CF_3	CF_3	H	CF_3	O	$CH(CH_3)$	O	CH_2CF_3
P1307	H	CF_3	H	CF_3	CF_3	H	CH_3	O	$CH(CH_3)$	O	CH_2CF_3
P1308	H	CF_3	H	CF_3	CF_3	H	Br	O	$CH(CH_3)$	S	CH_2CF_3
P1309	H	CF_3	H	CF_3	CF_3	H	Cl	O	$CH(CH_3)$	S	CH_2CF_3
P1310	H	CF_3	H	CF_3	CF_3	H	CF_3	O	$CH(CH_3)$	S	CH_2CF_3
P1311	H	CF_3	H	CF_3	CF_3	H	CH_3	O	$CH(CH_3)$	S	CH_2CF_3
P1312	H	CF_3	H	CF_3	CF_3	H	Br	S	$CH(CH_3)$	O	CH_2CF_3
P1313	H	CF_3	H	CF_3	CF_3	H	Cl	S	$CH(CH_3)$	O	CH_2CF_3
P1314	H	CF_3	H	CF_3	CF_3	H	CF_3	S	$CH(CH_3)$	O	CH_2CF_3
P1315	H	CF_3	H	CF_3	CF_3	H	CH_3	S	$CH(CH_3)$	O	CH_2CF_3
P1316	H	CF_3	H	CF_3	CF_3	H	Br	O	$CH(CH_3)$	O	CH_2CHF_2
P1317	H	CF_3	H	CF_3	CF_3	H	Cl	O	$CH(CH_3)$	O	CH_2CHF_2
P1318	H	CF_3	H	CF_3	CF_3	H	CF_3	O	$CH(CH_3)$	O	CH_2CHF_2
P1319	H	CF_3	H	CF_3	CF_3	H	CH_3	O	$CH(CH_3)$	O	CH_2CHF_2

P1320	H	CF_3	H	CF_3	CF_3	H	Br	O	$CH(CH_3)$	O	$CH(CH_3)CF_3$
P1321	H	CF_3	H	CF_3	CF_3	H	Cl	O	$CH(CH_3)$	O	$CH(CH_3)CF_3$
P1322	H	CF_3	H	CF_3	CF_3	H	CF_3	O	$CH(CH_3)$	O	$CH(CH_3)CF_3$
P1323	H	CF_3	H	CF_3	CF_3	H	CH_3	O	$CH(CH_3)$	O	$CH(CH_3)CF_3$
P1324	H	CF_3	H	CF_3	CF_3	H	Br	O	$CH(CH_3)$	O	$CH_2CH_2CF_3$
P1325	H	CF_3	H	CF_3	CF_3	H	Cl	O	$CH(CH_3)$	O	$CH_2CH_2CF_3$
P1326	H	CF_3	H	CF_3	CF_3	H	CF_3	O	$CH(CH_3)$	O	$CH_2CH_2CF_3$
P1327	H	CF_3	H	CF_3	CF_3	H	CH_3	O	$CH(CH_3)$	O	$CH_2CH_2CF_3$
P1328	H	CF_3	H	CF_3	CF_3	H	Br	O	$CH(CH_2CH_3)$	O	CH_2CF_3
P1329	H	CF_3	H	CF_3	CF_3	H	Cl	O	$CH(CH_2CH_3)$	O	CH_2CF_3
P1330	H	CF_3	H	CF_3	CF_3	H	CF_3	O	$CH(CH_2CH_3)$	O	CH_2CF_3
P1331	H	CF_3	H	CF_3	CF_3	H	CH_3	O	$CH(CH_2CH_3)$	O	CH_2CF_3
P1332	H	CF_3	H	CF_3	CF_3	H	Br	O	$C(CH_3)_2$	O	CH_2CF_3
P1333	H	CF_3	H	CF_3	CF_3	H	Cl	O	$C(CH_3)_2$	O	CH_2CF_3
P1334	H	CF_3	H	CF_3	CF_3	H	CF_3	O	$C(CH_3)_2$	O	CH_2CF_3
P1335	H	CF_3	H	CF_3	CF_3	H	CH_3	O	$C(CH_3)_2$	O	CH_2CF_3
P1336	H	CF_3	H	CF_3	CF_3	H	Br	O	CH_2CH_2	O	CH_2CF_3
P1337	H	CF_3	H	CF_3	CF_3	H	Cl	O	CH_2CH_2	O	CH_2CF_3
P1338	H	CF_3	H	CF_3	CF_3	H	CF_3	O	CH_2CH_2	O	CH_2CF_3
P1339	H	CF_3	H	CF_3	CF_3	H	CH_3	O	CH_2CH_2	O	CH_2CF_3
P1340	H	F	H	CF_3	CF_3	H	Br	O	CH_2	O	CH_2CF_3
P1341	H	F	H	CF_3	CF_3	H	Cl	O	CH_2	O	CH_2CF_3
P1342	H	F	H	CF_3	CF_3	H	CF_3	O	CH_2	O	CH_2CF_3
P1343	H	F	H	CF_3	CF_3	H	CH_3	O	CH_2	O	CH_2CF_3
P1344	H	F	H	CF_3	CF_3	H	Br	O	CH_2	S	CH_2CF_3
P1345	H	F	H	CF_3	CF_3	H	Cl	O	CH_2	S	CH_2CF_3
P1346	H	F	H	CF_3	CF_3	H	CF_3	O	CH_2	S	CH_2CF_3

P1347	H	F	H	CF_3	CF_3	H	CH_3	O	CH_2	S	CH_2CF_3
P1348	H	F	H	CF_3	CF_3	H	Br	S	CH_2	O	CH_2CF_3
P1349	H	F	H	CF_3	CF_3	H	Cl	S	CH_2	O	CH_2CF_3
P1350	H	F	H	CF_3	CF_3	H	CF_3	S	CH_2	O	CH_2CF_3
P1351	H	F	H	CF_3	CF_3	H	CH_3	S	CH_2	O	CH_2CF_3
P1352	H	F	H	CF_3	CF_3	H	Br	O	CH_2	O	CH_2CHF_2
P1353	H	F	H	CF_3	CF_3	H	Cl	O	CH_2	O	CH_2CHF_2
P1354	H	F	H	CF_3	CF_3	H	CF_3	O	CH_2	O	CH_2CHF_2
P1355	H	F	H	CF_3	CF_3	H	CH_3	O	CH_2	O	CH_2CHF_2
P1356	H	F	H	CF_3	CF_3	CF_3	Br	O	CH_2	O	CH_2CF_3
P1357	H	F	H	CF_3	CF_3	CF_3	Cl	O	CH_2	O	CH_2CF_3
P1358	H	F	H	CF_3	CF_3	CF_3	CF_3	O	CH_2	O	CH_2CF_3
P1359	H	F	H	CF_3	CF_3	CF_3	CH_3	O	CH_2	O	CH_2CF_3
P1360	H	F	H	CF_3	CF_2CF_3	H	Br	O	CH_2	O	CH_2CF_3
P1361	H	F	H	CF_3	CF_2CF_3	H	Cl	O	CH_2	O	CH_2CF_3
P1362	H	F	H	CF_3	CF_2CF_3	H	CF_3	O	CH_2	O	CH_2CF_3
P1363	H	F	H	CF_3	CF_2CF_3	H	CH_3	O	CH_2	O	CH_2CF_3
P1364	H	F	H	CF_3	CF_3	H	Br	O	CH_2	O	$CH(CH_3)CF_3$
P1365	H	F	H	CF_3	CF_3	H	Cl	O	CH_2	O	$CH(CH_3)CF_3$
P1366	H	F	H	CF_3	CF_3	H	CF_3	O	CH_2	O	$CH(CH_3)CF_3$
P1367	H	F	H	CF_3	CF_3	H	CH_3	O	CH_2	O	$CH(CH_3)CF_3$
P1368	H	F	H	CF_3	CF_3	CF_3	Br	O	$CH(CH_3)$	O	CH_2CF_3
P1369	H	F	H	CF_3	CF_3	CF_3	Cl	O	$CH(CH_3)$	O	CH_2CF_3
P1370	H	F	H	CF_3	CF_3	CF_3	CF_3	O	$CH(CH_3)$	O	CH_2CF_3
P1371	H	F	H	CF_3	CF_3	CF_3	CH_3	O	$CH(CH_3)$	O	CH_2CF_3
P1372	H	F	H	CF_3	CF_2CF_3	H	Br	O	$CH(CH_3)$	O	CH_2CF_3
P1373	H	F	H	CF_3	CF_2CF_3	H	Cl	O	$CH(CH_3)$	O	CH_2CF_3

P1374	H	F	H	CF_3	CF_2CF_3	H	CF_3	O	CH (CH_3)	O	CH_2CF_3
P1375	H	F	H	CF_3	CF_2CF_3	H	CH_3	O	CH (CH_3)	O	CH_2CF_3
P1376	H	F	H	CF_3	CF_3	H	Cl	O	CH (CH_3)	O	CH_2CF_3
P1377	H	F	H	CF_3	CF_3	H	CF_3	O	CH(CH_3)	O	CH_2CF_3
P1378	H	F	H	CF_3	CF_3	H	CH_3	O	CH (CH_3)	O	CH_2CF_3
P1379	H	F	H	CF_3	CF_3	H	Br	O	CH(CH_3)	S	CH_2CF_3
P1380	H	F	H	CF_3	CF_3	H	Cl	O	CH (CH_3)	S	CH_2CF_3
P1381	H	F	H	CF_3	CF_3	H	CF_3	O	CH(CH_3)	S	CH_2CF_3
P1382	H	F	H	CF_3	CF_3	H	CH_3	O	CH (CH_3)	S	CH_2CF_3
P1383	H	F	H	CF_3	CF_3	H	Br	S	CH(CH_3)	O	CH_2CF_3
P1384	H	F	H	CF_3	CF_3	H	Cl	S	CH (CH_3)	O	CH_2CF_3
P1385	H	F	H	CF_3	CF_3	H	CF_3	S	CH(CH_3)	O	CH_2CF_3
P1386	H	F	H	CF_3	CF_3	H	CH_3	S	CH (CH_3)	O	CH_2CF_3
P1387	H	F	H	CF_3	CF_3	H	Br	O	CH(CH_3)	O	CH_2CHF_2
P1388	H	F	H	CF_3	CF_3	H	Cl	O	CH (CH_3)	O	CH_2CHF_2
P1389	H	F	H	CF_3	CF_3	H	CF_3	O	CH(CH_3)	O	CH_2CHF_2
P1390	H	F	H	CF_3	CF_3	H	CH_3	O	CH (CH_3)	O	CH_2CHF_2
P1391	H	F	H	CF_3	CF_3	H	Br	O	CH(CH_3)	O	$CH(CH_3)CF_3$
P1392	H	F	H	CF_3	CF_3	H	Cl	O	CH (CH_3)	O	$CH(CH_3)CF_3$
P1393	H	F	H	CF_3	CF_3	H	CF_3	O	CH(CH_3)	O	$CH(CH_3)CF_3$
P1394	H	F	H	CF_3	CF_3	H	CH_3	O	CH (CH_3)	O	$CH(CH_3)CF_3$
P1395	H	F	H	CF_3	CF_3	H	Br	O	CH(CH_3)	O	$CH_2CH_2CF_3$
P1396	H	F	H	CF_3	CF_3	H	Cl	O	CH (CH_3)	O	$CH_2CH_2CF_3$
P1397	H	F	H	CF_3	CF_3	H	CF_3	O	CH(CH_3)	O	$CH_2CH_2CF_3$
P1398	H	F	H	CF_3	CF_3	H	CH_3	O	CH (CH_3)	O	$CH_2CH_2CF_3$
P1399	H	F	H	CF_3	CF_3	H	Br	O	CH(CH_2CH_3)	O	CH_2CF_3
P1400	H	F	H	CF_3	CF_3	H	Cl	O	CH(CH_2CH_3)	O	CH_2CF_3

P1401	H	F	H	CF_3	CF_3	H	CF_3	O	$CH(CH_2CH_3)$	O	CH_2CF_3
P1402	H	F	H	CF_3	CF_3	H	CH_3	O	$CH(CH_2CH_3)$	O	CH_2CF_3
P1403	H	F	H	CF_3	CF_3	H	Br	O	$C(CH_3)_2$	O	CH_2CF_3
P1404	H	F	H	CF_3	CF_3	H	Cl	O	$C(CH_3)_2$	O	CH_2CF_3
P1405	H	F	H	CF_3	CF_3	H	CF_3	O	$C(CH_3)_2$	O	CH_2CF_3
P1406	H	F	H	CF_3	CF_3	H	CH_3	O	$C(CH_3)_2$	O	CH_2CF_3
P1407	H	F	H	CF_3	CF_3	H	Br	O	CH_2CH_2	O	CH_2CF_3
P1408	H	F	H	CF_3	CF_3	H	Cl	O	CH_2CH_2	O	CH_2CF_3
P1409	H	F	H	CF_3	CF_3	H	CF_3	O	CH_2CH_2	O	CH_2CF_3
P1410	H	F	H	CF_3	CF_3	H	CH_3	O	CH_2CH_2	O	CH_2CF_3
P1411	H	Cl	H	CF_3	CF_3	H	Br	O	CH_2	O	CH_2CF_3
P1412	H	Cl	H	CF_3	CF_3	H	Cl	O	CH_2	O	CH_2CF_3
P1413	H	Cl	H	CF_3	CF_3	H	CF_3	O	CH_2	O	CH_2CF_3
P1414	H	Cl	H	CF_3	CF_3	H	CH_3	O	CH_2	O	CH_2CF_3
P1415	H	Cl	H	CF_3	CF_3	H	Br	O	CH_2	S	CH_2CF_3
P1416	H	Cl	H	CF_3	CF_3	H	Cl	O	CH_2	S	CH_2CF_3
P1417	H	Cl	H	CF_3	CF_3	H	CF_3	O	CH_2	S	CH_2CF_3
P1418	H	Cl	H	CF_3	CF_3	H	CH_3	O	CH_2	S	CH_2CF_3
P1419	H	Cl	H	CF_3	CF_3	H	Br	S	CH_2	O	CH_2CF_3
P1420	H	Cl	H	CF_3	CF_3	H	Cl	S	CH_2	O	CH_2CF_3
P1421	H	Cl	H	CF_3	CF_3	H	CF_3	S	CH_2	O	CH_2CF_3
P1422	H	Cl	H	CF_3	CF_3	H	CH_3	S	CH_2	O	CH_2CF_3
P1423	H	Cl	H	CF_3	CF_3	H	Br	O	CH_2	O	CH_2CHF_2
P1424	H	Cl	H	CF_3	CF_3	H	Cl	O	CH_2	O	CH_2CHF_2
P1425	H	Cl	H	CF_3	CF_3	H	CF_3	O	CH_2	O	CH_2CHF_2
P1426	H	Cl	H	CF_3	CF_3	H	CH_3	O	CH_2	O	CH_2CHF_2
P1427	H	Cl	H	CF_3	CF_3	CF_3	Br	O	CH_2	O	CH_2CF_3

P1428	H	Cl	H	CF_3	CF_3	CF_3	Cl	O	CH_2	O	CH_2CF_3
P1429	H	Cl	H	CF_3	CF_3	CF_3	CF_3	O	CH_2	O	CH_2CF_3
P1430	H	Cl	H	CF_3	CF_3	CF_3	CH_3	O	CH_2	O	CH_2CF_3
P1431	H	Cl	H	CF_3	CF_2CF_3	H	Br	O	CH_2	O	CH_2CF_3
P1432	H	Cl	H	CF_3	CF_2CF_3	H	Cl	O	CH_2	O	CH_2CF_3
P1433	H	Cl	H	CF_3	CF_2CF_3	H	CF_3	O	CH_2	O	CH_2CF_3
P1434	H	Cl	H	CF_3	CF_2CF_3	H	CH_3	O	CH_2	O	CH_2CF_3
P1435	H	Cl	H	CF_3	CF_3	H	Br	O	CH_2	O	$CH(CH_3)CF_3$
P1436	H	Cl	H	CF_3	CF_3	H	Cl	O	CH_2	O	$CH(CH_3)CF_3$
P1437	H	Cl	H	CF_3	CF_3	H	CF_3	O	CH_2	O	$CH(CH_3)CF_3$
P1438	H	Cl	H	CF_3	CF_3	H	CH_3	O	CH_2	O	$CH(CH_3)CF_3$
P1439	H	Cl	H	CF_3	CF_3	CF_3	Br	O	$CH(CH_3)$	O	CH_2CF_3
P1440	H	Cl	H	CF_3	CF_3	CF_3	Cl	O	$CH(CH_3)$	O	CH_2CF_3
P1441	H	Cl	H	CF_3	CF_3	CF_3	CF_3	O	$CH(CH_3)$	O	CH_2CF_3
P1442	H	Cl	H	CF_3	CF_3	CF_3	CH_3	O	$CH(CH_3)$	O	CH_2CF_3
P1443	H	Cl	H	CF_3	CF_2CF_3	H	Br	O	$CH(CH_3)$	O	CH_2CF_3
P1444	H	Cl	H	CF_3	CF_2CF_3	H	Cl	O	$CH(CH_3)$	O	CH_2CF_3
P1445	H	Cl	H	CF_3	CF_2CF_3	H	CF_3	O	$CH(CH_3)$	O	CH_2CF_3
P1446	H	Cl	H	CF_3	CF_2CF_3	H	CH_3	O	$CH(CH_3)$	O	CH_2CF_3
P1447	H	Cl	H	CF_3	CF_3	H	Br	O	$CH(CH_3)$	O	CH_2CF_3
P1448	H	Cl	H	CF_3	CF_3	H	Cl	O	$CH(CH_3)$	O	CH_2CF_3
P1449	H	Cl	H	CF_3	CF_3	H	CF_3	O	$CH(CH_3)$	O	CH_2CF_3
P1450	H	Cl	H	CF_3	CF_3	H	CH_3	O	$CH(CH_3)$	O	CH_2CF_3
P1451	H	Cl	H	CF_3	CF_3	H	Br	O	$CH(CH_3)$	S	CH_2CF_3
P1452	H	Cl	H	CF_3	CF_3	H	Cl	O	$CH(CH_3)$	S	CH_2CF_3
P1453	H	Cl	H	CF_3	CF_3	H	CF_3	O	$CH(CH_3)$	S	CH_2CF_3
P1454	H	Cl	H	CF_3	CF_3	H	CH_3	O	$CH(CH_3)$	S	CH_2CF_3

P1455	H	Cl	H	CF_3	CF_3	H	Br	S	$CH(CH_3)$	O	CH_2CF_3
P1456	H	Cl	H	CF_3	CF_3	H	Cl	S	$CH(CH_3)$	O	CH_2CF_3
P1457	H	Cl	H	CF_3	CF_3	H	CF_3	S	$CH(CH_3)$	O	CH_2CF_3
P1458	H	Cl	H	CF_3	CF_3	H	CH_3	S	$CH(CH_3)$	O	CH_2CF_3
P1459	H	Cl	H	CF_3	CF_3	H	Br	O	$CH(CH_3)$	O	CH_2CHF_2
P1460	H	Cl	H	CF_3	CF_3	H	Cl	O	$CH(CH_3)$	O	CH_2CHF_2
P1461	H	Cl	H	CF_3	CF_3	H	CF_3	O	$CH(CH_3)$	O	CH_2CHF_2
P1462	H	Cl	H	CF_3	CF_3	H	CH_3	O	$CH(CH_3)$	O	CH_2CHF_2
P1463	H	Cl	H	CF_3	CF_3	H	Br	O	$CH(CH_3)$	O	$CH(CH_3)CF_3$
P1464	H	Cl	H	CF_3	CF_3	H	Cl	O	$CH(CH_3)$	O	$CH(CH_3)CF_3$
P1465	H	Cl	H	CF_3	CF_3	H	CF_3	O	$CH(CH_3)$	O	$CH(CH_3)CF_3$
P1466	H	Cl	H	CF_3	CF_3	H	CH_3	O	$CH(CH_3)$	O	$CH(CH_3)CF_3$
P1467	H	Cl	H	CF_3	CF_3	H	Br	O	$CH(CH_3)$	O	$CH_2CH_2CF_3$
P1468	H	Cl	H	CF_3	CF_3	H	Cl	O	$CH(CH_3)$	O	$CH_2CH_2CF_3$
P1469	H	Cl	H	CF_3	CF_3	H	CF_3	O	$CH(CH_3)$	O	$CH_2CH_2CF_3$
P1470	H	Cl	H	CF_3	CF_3	H	CH_3	O	$CH(CH_3)$	O	$CH_2CH_2CF_3$
P1471	H	Cl	H	CF_3	CF_3	H	Br	O	$CH(CH_2CH_3)$	O	CH_2CF_3
P1472	H	Cl	H	CF_3	CF_3	H	Cl	O	$CH(CH_2CH_3)$	O	CH_2CF_3
P1473	H	Cl	H	CF_3	CF_3	H	CF_3	O	$CH(CH_2CH_3)$	O	CH_2CF_3
P1474	H	Cl	H	CF_3	CF_3	H	CH_3	O	$CH(CH_2CH_3)$	O	CH_2CF_3
P1475	H	Cl	H	CF_3	CF_3	H	Br	O	$C(CH_3)_2$	O	CH_2CF_3
P1476	H	Cl	H	CF_3	CF_3	H	Cl	O	$C(CH_3)_2$	O	CH_2CF_3
P1477	H	Cl	H	CF_3	CF_3	H	CF_3	O	$C(CH_3)_2$	O	CH_2CF_3
P1478	H	Cl	H	CF_3	CF_3	H	CH_3	O	$C(CH_3)_2$	O	CH_2CF_3
P1479	H	Cl	H	CF_3	CF_3	H	Br	O	CH_2CH_2	O	CH_2CF_3
P1480	H	Cl	H	CF_3	CF_3	H	Cl	O	CH_2CH_2	O	CH_2CF_3
P1481	H	Cl	H	CF_3	CF_3	H	CF_3	O	CH_2CH_2	O	CH_2CF_3

P1482	H	Cl	H	CF_3	CF_3	H	CH_3	O	CH_2CH_2	O	CH_2CF_3
P1483	H	H	F	CF_3	CF_3	H	Br	O	CH_2	O	CH_2CF_3
P1484	H	H	F	CF_3	CF_3	H	Cl	O	CH_2	O	CH_2CF_3
P1485	H	H	F	CF_3	CF_3	H	CF_3	O	CH_2	O	CH_2CF_3
P1486	H	H	F	CF_3	CF_3	H	CH_3	O	CH_2	O	CH_2CF_3
P1487	H	H	F	CF_3	CF_3	H	Br	O	CH_2	S	CH_2CF_3
P1488	H	H	F	CF_3	CF_3	H	Cl	O	CH_2	S	CH_2CF_3
P1489	H	H	F	CF_3	CF_3	H	CF_3	O	CH_2	S	CH_2CF_3
P1490	H	H	F	CF_3	CF_3	H	CH_3	O	CH_2	S	CH_2CF_3
P1491	H	H	F	CF_3	CF_3	H	Br	S	CH_2	O	CH_2CF_3
P1492	H	H	F	CF_3	CF_3	H	Cl	S	CH_2	O	CH_2CF_3
P1493	H	H	F	CF_3	CF_3	H	CF_3	S	CH_2	O	CH_2CF_3
P1494	H	H	F	CF_3	CF_3	H	CH_3	S	CH_2	O	CH_2CF_3
P1495	H	H	F	CF_3	CF_3	H	Br	O	CH_2	O	CH_2CHF_2
P1496	H	H	F	CF_3	CF_3	H	Cl	O	CH_2	O	CH_2CHF_2
P1497	H	H	F	CF_3	CF_3	H	CF_3	O	CH_2	O	CH_2CHF_2
P1498	H	H	F	CF_3	CF_3	H	CH_3	O	CH_2	O	CH_2CHF_2
P1499	H	H	F	CF_3	CF_3	CF_3	Br	O	CH_2	O	CH_2CF_3
P1500	H	H	F	CF_3	CF_3	CF_3	Cl	O	CH_2	O	CH_2CF_3
P1501	H	H	F	CF_3	CF_3	CF_3	CF_3	O	CH_2	O	CH_2CF_3
P1502	H	H	F	CF_3	CF_3	CF_3	CH_3	O	CH_2	O	CH_2CF_3
P1503	H	H	F	CF_3	CF_2CF_3	H	Br	O	CH_2	O	CH_2CF_3
P1504	H	H	F	CF_3	CF_2CF_3	H	Cl	O	CH_2	O	CH_2CF_3
P1505	H	H	F	CF_3	CF_2CF_3	H	CF_3	O	CH_2	O	CH_2CF_3
P1506	H	H	F	CF_3	CF_2CF_3	H	CH_3	O	CH_2	O	CH_2CF_3
P1507	H	H	F	CF_3	CF_3	H	Br	O	CH_2	O	$CH(CH_3)CF_3$
P1508	H	H	F	CF_3	CF_3	H	Cl	O	CH_2	O	$CH(CH_3)CF_3$

P1509	H	H	F	CF_3	CF_3	H	CF_3	O	CH_2	O	$CH(CH_3)CF_3$
P1510	H	H	F	CF_3	CF_3	H	CH_3	O	CH_2	O	$CH(CH_3)CF_3$
P1511	H	H	F	CF_3	CF_3	CF_3	Br	O	$CH(CH_3)$	O	CH_2CF_3
P1512	H	H	F	CF_3	CF_3	CF_3	Cl	O	$CH(CH_3)$	O	CH_2CF_3
P1513	H	H	F	CF_3	CF_3	CF_3	CF_3	O	$CH(CH_3)$	O	CH_2CF_3
P1514	H	H	F	CF_3	CF_3	CF_3	CH_3	O	$CH(CH_3)$	O	CH_2CF_3
P1515	H	H	F	CF_3	CF_2CF_3	H	Br	O	$CH(CH_3)$	O	CH_2CF_3
P1516	H	H	F	CF_3	CF_2CF_3	H	Cl	O	$CH(CH_3)$	O	CH_2CF_3
P1517	H	H	F	CF_3	CF_2CF_3	H	CF_3	O	$CH(CH_3)$	O	CH_2CF_3
P1518	H	H	F	CF_3	CF_2CF_3	H	CH_3	O	$CH(CH_3)$	O	CH_2CF_3
P1519	H	H	F	CF_3	CF_3	H	Br	O	$CH(CH_3)$	O	CH_2CF_3
P1520	H	H	F	CF_3	CF_3	H	Cl	O	$CH(CH_3)$	O	CH_2CF_3
P1521	H	H	F	CF_3	CF_3	H	CF_3	O	$CH(CH_3)$	O	CH_2CF_3
P1522	H	H	F	CF_3	CF_3	H	CH_3	O	$CH(CH_3)$	O	CH_2CF_3
P1523	H	H	F	CF_3	CF_3	H	Br	O	$CH(CH_3)$	S	CH_2CF_3
P1524	H	H	F	CF_3	CF_3	H	Cl	O	$CH(CH_3)$	S	CH_2CF_3
P1525	H	H	F	CF_3	CF_3	H	CF_3	O	$CH(CH_3)$	S	CH_2CF_3
P1526	H	H	F	CF_3	CF_3	H	CH_3	O	$CH(CH_3)$	S	CH_2CF_3
P1527	H	H	F	CF_3	CF_3	H	Br	S	$CH(CH_3)$	O	CH_2CF_3
P1528	H	H	F	CF_3	CF_3	H	Cl	S	$CH(CH_3)$	O	CH_2CF_3
P1529	H	H	F	CF_3	CF_3	H	CF_3	S	$CH(CH_3)$	O	CH_2CF_3
P1530	H	H	F	CF_3	CF_3	H	CH_3	S	$CH(CH_3)$	O	CH_2CF_3
P1531	H	H	F	CF_3	CF_3	H	Br	O	$CH(CH_3)$	O	CH_2CHF_2
P1532	H	H	F	CF_3	CF_3	H	Cl	O	$CH(CH_3)$	O	CH_2CHF_2
P1533	H	H	F	CF_3	CF_3	H	CF_3	O	$CH(CH_3)$	O	CH_2CHF_2
P1534	H	H	F	CF_3	CF_3	H	CH_3	O	$CH(CH_3)$	O	CH_2CHF_2
P1535	H	H	F	CF_3	CF_3	H	Br	O	$CH(CH_3)$	O	$CH(CH_3)CF_3$

P1536	H	H	F	CF_3	CF_3	H	Cl	O	CH (CH_3)	O	$CH(CH_3)CF_3$
P1537	H	H	F	CF_3	CF_3	H	CF_3	O	$CH(CH_3)$	O	$CH(CH_3)CF_3$
P1538	H	H	F	CF_3	CF_3	H	CH_3	O	CH (CH_3)	O	$CH(CH_3)CF_3$
P1539	H	H	F	CF_3	CF_3	H	Br	O	$CH(CH_3)$	O	$CH_2CH_2CF_3$
P1540	H	H	F	CF_3	CF_3	H	Cl	O	CH (CH_3)	O	$CH_2CH_2CF_3$
P1541	H	H	F	CF_3	CF_3	H	CF_3	O	$CH(CH_3)$	O	$CH_2CH_2CF_3$
P1542	H	H	F	CF_3	CF_3	H	CH_3	O	CH (CH_3)	O	$CH_2CH_2CF_3$
P1543	H	H	F	CF_3	CF_3	H	Br	O	$CH(CH_2CH_3)$	O	CH_2CF_3
P1544	H	H	F	CF_3	CF_3	H	Cl	O	$CH(CH_2CH_3)$	O	CH_2CF_3
P1545	H	H	F	CF_3	CF_3	H	CF_3	O	$CH(CH_2CH_3)$	O	CH_2CF_3
P1546	H	H	F	CF_3	CF_3	H	CH_3	O	$CH(CH_2CH_3)$	O	CH_2CF_3
P1547	H	H	F	CF_3	CF_3	H	Br	O	$C(CH_3)_2$	O	CH_2CF_3
P1548	H	H	F	CF_3	CF_3	H	Cl	O	$C(CH_3)_2$	O	CH_2CF_3
P1549	H	H	F	CF_3	CF_3	H	CF_3	O	$C(CH_3)_2$	O	CH_2CF_3
P1550	H	H	F	CF_3	CF_3	H	CH_3	O	$C(CH_3)_2$	O	CH_2CF_3
P1551	H	H	F	CF_3	CF_3	H	Br	O	CH_2CH_2	O	CH_2CF_3
P1552	H	H	F	CF_3	CF_3	H	Cl	O	CH_2CH_2	O	CH_2CF_3
P1553	H	H	F	CF_3	CF_3	H	CF_3	O	CH_2CH_2	O	CH_2CF_3
P1554	H	H	F	CF_3	CF_3	H	CH_3	O	CH_2CH_2	O	CH_2CF_3
P1555	H	Cl	Cl	Cl	CF_3	H	H	O	CH_2	O	CH_2CF_3
P1556	H	Cl	Cl	Cl	CF_3	H	CF_3	O	CH_2	O	CH_2CF_3
P1557	H	Cl	Cl	Cl	CF_3	H	H	O	CH_2	S	CH_2CF_3
P1558	H	Cl	Cl	Cl	CF_3	H	Cl	O	CH_2	S	CH_2CF_3
P1559	H	Cl	Cl	Cl	CF_3	H	CF_3	O	CH_2	S	CH_2CF_3
P1560	H	Cl	Cl	Cl	CF_3	H	CH_3	O	CH_2	S	CH_2CF_3
P1561	H	Cl	Cl	Cl	CF_3	H	H	S	CH_2	O	CH_2CF_3
P1562	H	Cl	Cl	Cl	CF_3	H	Cl	S	CH_2	O	CH_2CF_3

P1563	H	Cl	Cl	Cl	CF_3	H	CF_3	S	CH_2	O	CH_2CF_3
P1564	H	Cl	Cl	Cl	CF_3	H	CH_3	S	CH_2	O	CH_2CF_3
P1565	H	Cl	Cl	Cl	CF_3	H	H	O	CH_2	O	CH_2CHF_2
P1566	H	Cl	Cl	Cl	CF_3	H	CF_3	O	CH_2	O	CH_2CHF_2
P1567	H	Cl	Cl	Cl	CF_3	H	CH_3	O	CH_2	O	CH_2CHF_2
P1568	H	Cl	Cl	Cl	CF_3	H	H	O	CH_2	O	CH_2CH_2F
P1569	H	Cl	Cl	Cl	CF_3	H	Br	O	CH_2	O	CH_2CH_2F
P1570	H	Cl	Cl	Cl	CF_3	H	Cl	O	CH_2	O	CH_2CH_2F
P1571	H	Cl	Cl	Cl	CF_3	H	CF_3	O	CH_2	O	CH_2CH_2F
P1572	H	Cl	Cl	Cl	CF_3	H	CH_3	O	CH_2	O	CH_2CH_2F
P1573	H	Cl	Cl	Cl	CF_3	H	H	O	CH_2	O	CH_2CH_3
P1574	H	Cl	Cl	Cl	CF_3	H	Br	O	CH_2	O	CH_2CH_3
P1575	H	Cl	Cl	Cl	CF_3	H	Cl	O	CH_2	O	CH_2CH_3
P1576	H	Cl	Cl	Cl	CF_3	H	CF_3	O	CH_2	O	CH_2CH_3
P1577	H	Cl	Cl	Cl	CF_3	H	CH_3	O	CH_2	O	CH_2CH_3
P1578	H	Cl	Cl	Cl	CF_3	CF_3	H	O	CH_2	O	CH_2CF_3
P1579	H	Cl	Cl	Cl	CF_3	CF_3	Br	O	CH_2	O	CH_2CF_3
P1580	H	Cl	Cl	Cl	CF_3	CF_3	Cl	O	CH_2	O	CH_2CF_3
P1581	H	Cl	Cl	Cl	CF_3	CF_3	CF_3	O	CH_2	O	CH_2CF_3
P1582	H	Cl	Cl	Cl	CF_3	CF_3	CH_3	O	CH_2	O	CH_2CF_3
P1583	H	Cl	Cl	Cl	CF_2CF_3	H	H	O	CH_2	O	CH_2CF_3
P1584	H	Cl	Cl	Cl	CF_2CF_3	H	Br	O	CH_2	O	CH_2CF_3
P1585	H	Cl	Cl	Cl	CF_2CF_3	H	Cl	O	CH_2	O	CH_2CF_3
P1586	H	Cl	Cl	Cl	CF_2CF_3	H	CF_3	O	CH_2	O	CH_2CF_3
P1587	H	Cl	Cl	Cl	CF_2CF_3	H	CH_3	O	CH_2	O	CH_2CF_3
P1588	H	Cl	Cl	Cl	CF_3	H	H	O	CH_2	O	$CH(CH_3)CF_3$
P1589	H	Cl	Cl	Cl	CF_3	H	Br	O	CH_2	O	$CH(CH_3)CF_3$

P1590	H	Cl	Cl	Cl	CF_3	H	Cl	O	CH_2	O	$CH(CH_3)CF_3$
P1591	H	Cl	Cl	Cl	CF_3	H	CF_3	O	CH_2	O	$CH(CH_3)CF_3$
P1592	H	Cl	Cl	Cl	CF_3	H	CH_3	O	CH_2	O	$CH(CH_3)CF_3$
P1593	H	Cl	Cl	Cl	CF_3	CF_3	H	O	$CH(CH_3)$	O	CH_2CF_3
P1594	H	Cl	Cl	Cl	CF_3	CF_3	Br	O	$CH(CH_3)$	O	CH_2CF_3
P1595	H	Cl	Cl	Cl	CF_3	CF_3	Cl	O	$CH(CH_3)$	O	CH_2CF_3
P1596	H	Cl	Cl	Cl	CF_3	CF_3	CF_3	O	$CH(CH_3)$	O	CH_2CF_3
P1597	H	Cl	Cl	Cl	CF_3	CF_3	CH_3	O	$CH(CH_3)$	O	CH_2CF_3
P1598	H	Cl	Cl	Cl	CF_2CF_3	H	H	O	$CH(CH_3)$	O	CH_2CF_3
P1599	H	Cl	Cl	Cl	CF_2CF_3	H	Br	O	$CH(CH_3)$	O	CH_2CF_3
P1600	H	Cl	Cl	Cl	CF_2CF_3	H	Cl	O	$CH(CH_3)$	O	CH_2CF_3
P1601	H	Cl	Cl	Cl	CF_2CF_3	H	CF_3	O	$CH(CH_3)$	O	CH_2CF_3
P1602	H	Cl	Cl	Cl	CF_2CF_3	H	CH_3	O	$CH(CH_3)$	O	CH_2CF_3
P1603	H	Cl	Cl	Cl	CF_3	H	H	O	$CH(CH_3)$	O	CH_2CF_3
P1604	H	Cl	Cl	Cl	CF_3	H	H	O	$CH(CH_3)$	S	CH_2CF_3
P1605	H	Cl	Cl	Cl	CF_3	H	H	S	$CH(CH_3)$	O	CH_2CF_3
P1606	H	Cl	Cl	Cl	CF_3	H	Br	S	$CH(CH_3)$	O	CH_2CF_3
P1607	H	Cl	Cl	Cl	CF_3	H	Cl	S	$CH(CH_3)$	O	CH_2CF_3
P1608	H	Cl	Cl	Cl	CF_3	H	CF_3	S	$CH(CH_3)$	O	CH_2CF_3
P1609	H	Cl	Cl	Cl	CF_3	H	CH_3	S	$CH(CH_3)$	O	CH_2CF_3
P1610	H	Cl	Cl	Cl	CF_3	H	H	O	$CH(CH_3)$	O	CH_2CHF_2
P1611	H	Cl	Cl	Cl	CF_3	H	Br	O	$CH(CH_3)$	O	CH_2CHF_2
P1612	H	Cl	Cl	Cl	CF_3	H	Cl	O	$CH(CH_3)$	O	CH_2CHF_2
P1613	H	Cl	Cl	Cl	CF_3	H	CF_3	O	$CH(CH_3)$	O	CH_2CHF_2
P1614	H	Cl	Cl	Cl	CF_3	H	CH_3	O	$CH(CH_3)$	O	CH_2CHF_2
P1615	H	Cl	Cl	Cl	CF_3	H	H	O	$CH(CH_3)$	O	CH_2CH_2F
P1616	H	Cl	Cl	Cl	CF_3	H	Br	O	$CH(CH_3)$	O	CH_2CH_2F

P1617	H	Cl	Cl	Cl	CF_3	H	Cl	O	$CH(CH_3)$	O	CH_2CH_2F
P1618	H	Cl	Cl	Cl	CF_3	H	CF_3	O	$CH(CH_3)$	O	CH_2CH_2F
P1619	H	Cl	Cl	Cl	CF_3	H	CH_3	O	$CH(CH_3)$	O	CH_2CH_2F
P1620	H	Cl	Cl	Cl	CF_3	H	H	O	$CH(CH_3)$	O	CH_2CH_3
P1621	H	Cl	Cl	Cl	CF_3	H	Br	O	$CH(CH_3)$	O	CH_2CH_3
P1622	H	Cl	Cl	Cl	CF_3	H	Cl	O	$CH(CH_3)$	O	CH_2CH_3
P1623	H	Cl	Cl	Cl	CF_3	H	CF_3	O	$CH(CH_3)$	O	CH_2CH_3
P1624	H	Cl	Cl	Cl	CF_3	H	CH_3	O	$CH(CH_3)$	O	CH_2CH_3
P1625	H	Cl	Cl	Cl	CF_3	H	H	O	$CH(CH_3)$	O	$CH(CH_3)CF_3$
P1626	H	Cl	Cl	Cl	CF_3	H	Br	O	$CH(CH_3)$	O	$CH(CH_3)CF_3$
P1627	H	Cl	Cl	Cl	CF_3	H	Cl	O	$CH(CH_3)$	O	$CH(CH_3)CF_3$
P1628	H	Cl	Cl	Cl	CF_3	H	CF_3	O	$CH(CH_3)$	O	$CH(CH_3)CF_3$
P1629	H	Cl	Cl	Cl	CF_3	H	CH_3	O	$CH(CH_3)$	O	$CH(CH_3)CF_3$
P1630	H	Cl	Cl	Cl	CF_3	H	H	O	$CH(CH_3)$	O	$CH_2CH_2CF_3$
P1631	H	Cl	Cl	Cl	CF_3	H	Br	O	$CH(CH_3)$	O	$CH_2CH_2CF_3$
P1632	H	Cl	Cl	Cl	CF_3	H	Cl	O	$CH(CH_3)$	O	$CH_2CH_2CF_3$
P1633	H	Cl	Cl	Cl	CF_3	H	CF_3	O	$CH(CH_3)$	O	$CH_2CH_2CF_3$
P1634	H	Cl	Cl	Cl	CF_3	H	CH_3	O	$CH(CH_3)$	O	$CH_2CH_2CF_3$
P1635	H	Cl	Cl	Cl	CF_3	H	H	O	$CH(CH_2CH_3)$	O	CH_2CF_3
P1636	H	Cl	Cl	Cl	CF_3	H	H	O	$C(CH_3)_2$	O	CH_2CF_3
P1637	H	Cl	Cl	Cl	CF_3	H	H	O	CH_2CH_2	O	CH_2CF_3
P1638	H	Cl	Cl	Cl	CF_3	H	Br	O	CH_2CH_2	O	CH_2CF_3
P1639	H	Cl	Cl	Cl	CF_3	H	Cl	O	CH_2CH_2	O	CH_2CF_3
P1640	H	Cl	Cl	Cl	CF_3	H	CF_3	O	CH_2CH_2	O	CH_2CF_3
P1641	H	Cl	Cl	Cl	CF_3	H	CH_3	O	CH_2CH_2	O	CH_2CF_3
P1642	H	Cl	H	Cl	CF_3	H	H	O	CH_2	O	CH_2CF_3
P1643	H	Cl	H	Cl	CF_3	H	Br	O	CH_2	O	CH_2CF_3

P1644	H	Cl	H	Cl	CF_3	H	CF_3	O	CH_2	O	CH_2CF_3
P1645	H	Cl	H	Cl	CF_3	H	H	O	CH_2	S	CH_2CF_3
P1646	H	Cl	H	Cl	CF_3	H	Br	O	CH_2	S	CH_2CF_3
P1647	H	Cl	H	Cl	CF_3	H	Cl	O	CH_2	S	CH_2CF_3
P1648	H	Cl	H	Cl	CF_3	H	CF_3	O	CH_2	S	CH_2CF_3
P1649	H	Cl	H	Cl	CF_3	H	CH_3	O	CH_2	S	CH_2CF_3
P1650	H	Cl	H	Cl	CF_3	H	H	S	CH_2	O	CH_2CF_3
P1651	H	Cl	H	Cl	CF_3	H	Br	S	CH_2	O	CH_2CF_3
P1652	H	Cl	H	Cl	CF_3	H	Cl	S	CH_2	O	CH_2CF_3
P1653	H	Cl	H	Cl	CF_3	H	CF_3	S	CH_2	O	CH_2CF_3
P1654	H	Cl	H	Cl	CF_3	H	CH_3	S	CH_2	O	CH_2CF_3
P1655	H	Cl	H	Cl	CF_3	H	H	O	CH_2	O	CH_2CHF_2
P1656	H	Cl	H	Cl	CF_3	H	Br	O	CH_2	O	CH_2CHF_2
P1657	H	Cl	H	Cl	CF_3	H	Cl	O	CH_2	O	CH_2CHF_2
P1658	H	Cl	H	Cl	CF_3	H	CF_3	O	CH_2	O	CH_2CHF_2
P1659	H	Cl	H	Cl	CF_3	H	CH_3	O	CH_2	O	CH_2CHF_2
P1660	H	Cl	H	Cl	CF_3	H	H	O	CH_2	O	CH_2CH_2F
P1661	H	Cl	H	Cl	CF_3	H	Br	O	CH_2	O	CH_2CH_2F
P1662	H	Cl	H	Cl	CF_3	H	Cl	O	CH_2	O	CH_2CH_2F
P1663	H	Cl	H	Cl	CF_3	H	CF_3	O	CH_2	O	CH_2CH_2F
P1664	H	Cl	H	Cl	CF_3	H	CH_3	O	CH_2	O	CH_2CH_2F
P1665	H	Cl	H	Cl	CF_3	H	H	O	CH_2	O	CH_2CH_3
P1666	H	Cl	H	Cl	CF_3	H	Br	O	CH_2	O	CH_2CH_3
P1667	H	Cl	H	Cl	CF_3	H	Cl	O	CH_2	O	CH_2CH_3
P1668	H	Cl	H	Cl	CF_3	H	CF_3	O	CH_2	O	CH_2CH_3
P1669	H	Cl	H	Cl	CF_3	H	CH_3	O	CH_2	O	CH_2CH_3
P1670	H	Cl	H	Cl	CF_3	CF_3	H	O	CH_2	O	CH_2CF_3

P1671	H	Cl	H	Cl	CF_3	CF_3	Br	O	CH_2	O	CH_2CF_3
P1672	H	Cl	H	Cl	CF_3	CF_3	Cl	O	CH_2	O	CH_2CF_3
P1673	H	Cl	H	Cl	CF_3	CF_3	CF_3	O	CH_2	O	CH_2CF_3
P1674	H	Cl	H	Cl	CF_3	CF_3	CH_3	O	CH_2	O	CH_2CF_3
P1675	H	Cl	H	Cl	CF_2CF_3	H	H	O	CH_2	O	CH_2CF_3
P1676	H	Cl	H	Cl	CF_2CF_3	H	Br	O	CH_2	O	CH_2CF_3
P1677	H	Cl	H	Cl	CF_2CF_3	H	Cl	O	CH_2	O	CH_2CF_3
P1678	H	Cl	H	Cl	CF_2CF_3	H	CF_3	O	CH_2	O	CH_2CF_3
P1679	H	Cl	H	Cl	CF_2CF_3	H	CH_3	O	CH_2	O	CH_2CF_3
P1680	H	Cl	H	Cl	CF_3	H	H	O	CH_2	O	$CH(CH_3)CF_3$
P1681	H	Cl	H	Cl	CF_3	H	Br	O	CH_2	O	$CH(CH_3)CF_3$
P1682	H	Cl	H	Cl	CF_3	H	Cl	O	CH_2	O	$CH(CH_3)CF_3$
P1683	H	Cl	H	Cl	CF_3	H	CF_3	O	CH_2	O	$CH(CH_3)CF_3$
P1684	H	Cl	H	Cl	CF_3	H	CH_3	O	CH_2	O	$CH(CH_3)CF_3$
P1685	H	Cl	H	Cl	CF_3	CF_3	H	O	$CH(CH_3)$	O	CH_2CF_3
P1686	H	Cl	H	Cl	CF_3	CF_3	Br	O	$CH(CH_3)$	O	CH_2CF_3
P1687	H	Cl	H	Cl	CF_3	CF_3	Cl	O	$CH(CH_3)$	O	CH_2CF_3
P1688	H	Cl	H	Cl	CF_3	CF_3	CF_3	O	$CH(CH_3)$	O	CH_2CF_3
P1689	H	Cl	H	Cl	CF_3	CF_3	CH_3	O	$CH(CH_3)$	O	CH_2CF_3
P1690	H	Cl	H	Cl	CF_2CF_3	H	H	O	$CH(CH_3)$	O	CH_2CF_3
P1691	H	Cl	H	Cl	CF_2CF_3	H	Br	O	$CH(CH_3)$	O	CH_2CF_3
P1692	H	Cl	H	Cl	CF_2CF_3	H	Cl	O	$CH(CH_3)$	O	CH_2CF_3
P1693	H	Cl	H	Cl	CF_2CF_3	H	CF_3	O	$CH(CH_3)$	O	CH_2CF_3
P1694	H	Cl	H	Cl	CF_2CF_3	H	CH_3	O	$CH(CH_3)$	O	CH_2CF_3
P1695	H	Cl	H	Cl	CF_3	H	H	O	$CH(CH_3)$	O	CH_2CF_3
P1696	H	Cl	H	Cl	CF_3	H	Br	O	$CH(CH_3)$	O	CH_2CF_3
P1697	H	Cl	H	Cl	CF_3	H	Cl	O	$CH(CH_3)$	O	CH_2CF_3

P1698	H	Cl	H	Cl	CF_3	H	CF_3	O	$CH(CH_3)$	O	CH_2CF_3
P1699	H	Cl	H	Cl	CF_3	H	CH_3	O	$CH(CH_3)$	O	CH_2CF_3
P1700	H	Cl	H	Cl	CF_3	H	H	O	$CH(CH_3)$	S	CH_2CF_3
P1701	H	Cl	H	Cl	CF_3	H	Br	O	$CH(CH_3)$	S	CH_2CF_3
P1702	H	Cl	H	Cl	CF_3	H	Cl	O	$CH(CH_3)$	S	CH_2CF_3
P1703	H	Cl	H	Cl	CF_3	H	CF_3	O	$CH(CH_3)$	S	CH_2CF_3
P1704	H	Cl	H	Cl	CF_3	H	CH_3	O	$CH(CH_3)$	S	CH_2CF_3
P1705	H	Cl	H	Cl	CF_3	H	H	S	$CH(CH_3)$	O	CH_2CF_3
P1706	H	Cl	H	Cl	CF_3	H	Br	S	$CH(CH_3)$	O	CH_2CF_3
P1707	H	Cl	H	Cl	CF_3	H	Cl	S	$CH(CH_3)$	O	CH_2CF_3
P1708	H	Cl	H	Cl	CF_3	H	CF_3	S	$CH(CH_3)$	O	CH_2CF_3
P1709	H	Cl	H	Cl	CF_3	H	CH_3	S	$CH(CH_3)$	O	CH_2CF_3
P1710	H	Cl	H	Cl	CF_3	H	H	O	$CH(CH_3)$	O	CH_2CHF_2
P1711	H	Cl	H	Cl	CF_3	H	Br	O	$CH(CH_3)$	O	CH_2CHF_2
P1712	H	Cl	H	Cl	CF_3	H	Cl	O	$CH(CH_3)$	O	CH_2CHF_2
P1713	H	Cl	H	Cl	CF_3	H	CF_3	O	$CH(CH_3)$	O	CH_2CHF_2
P1714	H	Cl	H	Cl	CF_3	H	CH_3	O	$CH(CH_3)$	O	CH_2CHF_2
P1715	H	Cl	H	Cl	CF_3	H	H	O	$CH(CH_3)$	O	CH_2CH_2F
P1716	H	Cl	H	Cl	CF_3	H	Br	O	$CH(CH_3)$	O	CH_2CH_2F
P1717	H	Cl	H	Cl	CF_3	H	Cl	O	$CH(CH_3)$	O	CH_2CH_2F
P1718	H	Cl	H	Cl	CF_3	H	CF_3	O	$CH(CH_3)$	O	CH_2CH_2F
P1719	H	Cl	H	Cl	CF_3	H	CH_3	O	$CH(CH_3)$	O	CH_2CH_2F
P1720	H	Cl	H	Cl	CF_3	H	H	O	$CH(CH_3)$	O	CH_2CH_3
P1721	H	Cl	H	Cl	CF_3	H	Br	O	$CH(CH_3)$	O	CH_2CH_3
P1722	H	Cl	H	Cl	CF_3	H	Cl	O	$CH(CH_3)$	O	CH_2CH_3
P1723	H	Cl	H	Cl	CF_3	H	CF_3	O	$CH(CH_3)$	O	CH_2CH_3
P1724	H	Cl	H	Cl	CF_3	H	CH_3	O	$CH(CH_3)$	O	CH_2CH_3

P1725	H	Cl	H	Cl	CF_3	H	H	O	$CH(CH_3)$	O	$CH(CH_3)CF_3$
P1726	H	Cl	H	Cl	CF_3	H	Br	O	$CH(CH_3)$	O	$CH(CH_3)CF_3$
P1727	H	Cl	H	Cl	CF_3	H	Cl	O	$CH(CH_3)$	O	$CH(CH_3)CF_3$
P1728	H	Cl	H	Cl	CF_3	H	CF_3	O	$CH(CH_3)$	O	$CH(CH_3)CF_3$
P1729	H	Cl	H	Cl	CF_3	H	CH_3	O	$CH(CH_3)$	O	$CH(CH_3)CF_3$
P1730	H	Cl	H	Cl	CF_3	H	H	O	$CH(CH_3)$	O	$CH_2CH_2CF_3$
P1731	H	Cl	H	Cl	CF_3	H	Br	O	$CH(CH_3)$	O	$CH_2CH_2CF_3$
P1732	H	Cl	H	Cl	CF_3	H	Cl	O	$CH(CH_3)$	O	$CH_2CH_2CF_3$
P1733	H	Cl	H	Cl	CF_3	H	CF_3	O	$CH(CH_3)$	O	$CH_2CH_2CF_3$
P1734	H	Cl	H	Cl	CF_3	H	CH_3	O	$CH(CH_3)$	O	$CH_2CH_2CF_3$
P1735	H	Cl	H	Cl	CF_3	H	H	O	$CH(CH_2CH_3)$	O	CH_2CF_3
P1736	H	Cl	H	Cl	CF_3	H	Br	O	$CH(CH_2CH_3)$	O	CH_2CF_3
P1737	H	Cl	H	Cl	CF_3	H	Cl	O	$CH(CH_2CH_3)$	O	CH_2CF_3
P1738	H	Cl	H	Cl	CF_3	H	CF_3	O	$CH(CH_2CH_3)$	O	CH_2CF_3
P1739	H	Cl	H	Cl	CF_3	H	CH_3	O	$CH(CH_2CH_3)$	O	CH_2CF_3
P1740	H	Cl	H	Cl	CF_3	H	H	O	$C(CH_3)_2$	O	CH_2CF_3
P1741	H	Cl	H	Cl	CF_3	H	Br	O	$C(CH_3)_2$	O	CH_2CF_3
P1742	H	Cl	H	Cl	CF_3	H	Cl	O	$C(CH_3)_2$	O	CH_2CF_3
P1743	H	Cl	H	Cl	CF_3	H	CF_3	O	$C(CH_3)_2$	O	CH_2CF_3
P1744	H	Cl	H	Cl	CF_3	H	CH_3	O	$C(CH_3)_2$	O	CH_2CF_3
P1745	H	Cl	H	Cl	CF_3	H	H	O	CH_2CH_2	O	CH_2CF_3
P1746	H	Cl	H	Cl	CF_3	H	Br	O	CH_2CH_2	O	CH_2CF_3
P1747	H	Cl	H	Cl	CF_3	H	Cl	O	CH_2CH_2	O	CH_2CF_3
P1748	H	Cl	H	Cl	CF_3	H	CF_3	O	CH_2CH_2	O	CH_2CF_3
P1749	H	Cl	H	Cl	CF_3	H	CH_3	O	CH_2CH_2	O	CH_2CF_3
P1750	H	H	Cl	Cl	CF_3	H	H	O	CH_2	O	CH_2CF_3
P1751	H	H	Cl	Cl	CF_3	H	Br	O	CH_2	O	CH_2CF_3

P1752	H	H	Cl	Cl	CF_3	H	Cl	O	CH_2	O	CH_2CF_3
P1753	H	H	Cl	Cl	CF_3	H	CF_3	O	CH_2	O	CH_2CF_3
P1754	H	H	Cl	Cl	CF_3	H	CH_3	O	CH_2	O	CH_2CF_3
P1755	H	H	Cl	Cl	CF_3	H	H	O	CH_2	S	CH_2CF_3
P1756	H	H	Cl	Cl	CF_3	H	Br	O	CH_2	S	CH_2CF_3
P1757	H	H	Cl	Cl	CF_3	H	Cl	O	CH_2	S	CH_2CF_3
P1758	H	H	Cl	Cl	CF_3	H	CF_3	O	CH_2	S	CH_2CF_3
P1759	H	H	Cl	Cl	CF_3	H	CH_3	O	CH_2	S	CH_2CF_3
P1760	H	H	Cl	Cl	CF_3	H	H	S	CH_2	O	CH_2CF_3
P1761	H	H	Cl	Cl	CF_3	H	Br	S	CH_2	O	CH_2CF_3
P1762	H	H	Cl	Cl	CF_3	H	Cl	S	CH_2	O	CH_2CF_3
P1763	H	H	Cl	Cl	CF_3	H	CF_3	S	CH_2	O	CH_2CF_3
P1764	H	H	Cl	Cl	CF_3	H	CH_3	S	CH_2	O	CH_2CF_3
P1765	H	H	Cl	Cl	CF_3	H	H	O	CH_2	O	CH_2CHF_2
P1766	H	H	Cl	Cl	CF_3	H	Br	O	CH_2	O	CH_2CHF_2
P1767	H	H	Cl	Cl	CF_3	H	Cl	O	CH_2	O	CH_2CHF_2
P1768	H	H	Cl	Cl	CF_3	H	CF_3	O	CH_2	O	CH_2CHF_2
P1769	H	H	Cl	Cl	CF_3	H	CH_3	O	CH_2	O	CH_2CHF_2
P1770	H	H	Cl	Cl	CF_3	H	H	O	CH_2	O	CH_2CH_2F
P1771	H	H	Cl	Cl	CF_3	H	Br	O	CH_2	O	CH_2CH_2F
P1772	H	H	Cl	Cl	CF_3	H	Cl	O	CH_2	O	CH_2CH_2F
P1773	H	H	Cl	Cl	CF_3	H	CF_3	O	CH_2	O	CH_2CH_2F
P1774	H	H	Cl	Cl	CF_3	H	CH_3	O	CH_2	O	CH_2CH_2F
P1775	H	H	Cl	Cl	CF_3	H	H	O	CH_2	O	CH_2CH_3
P1776	H	H	Cl	Cl	CF_3	H	Br	O	CH_2	O	CH_2CH_3
P1777	H	H	Cl	Cl	CF_3	H	Cl	O	CH_2	O	CH_2CH_3
P1778	H	H	Cl	Cl	CF_3	H	CF_3	O	CH_2	O	CH_2CH_3

P1779	H	H	Cl	Cl	CF_3	H	CH_3	O	CH_2	O	CH_2CH_3
P1780	H	H	Cl	Cl	CF_3	CF_3	H	O	CH_2	O	CH_2CF_3
P1781	H	H	Cl	Cl	CF_3	CF_3	Br	O	CH_2	O	CH_2CF_3
P1782	H	H	Cl	Cl	CF_3	CF_3	Cl	O	CH_2	O	CH_2CF_3
P1783	H	H	Cl	Cl	CF_3	CF_3	CF_3	O	CH_2	O	CH_2CF_3
P1784	H	H	Cl	Cl	CF_3	CF_3	CH_3	O	CH_2	O	CH_2CF_3
P1785	H	H	Cl	Cl	CF_2CF_3	H	H	O	CH_2	O	CH_2CF_3
P1786	H	H	Cl	Cl	CF_2CF_3	H	Br	O	CH_2	O	CH_2CF_3
P1787	H	H	Cl	Cl	CF_2CF_3	H	Cl	O	CH_2	O	CH_2CF_3
P1788	H	H	Cl	Cl	CF_2CF_3	H	CF_3	O	CH_2	O	CH_2CF_3
P1789	H	H	Cl	Cl	CF_2CF_3	H	CH_3	O	CH_2	O	CH_2CF_3
P1790	H	H	Cl	Cl	CF_3	H	H	O	CH_2	O	$CH(CH_3)CF_3$
P1791	H	H	Cl	Cl	CF_3	H	Br	O	CH_2	O	$CH(CH_3)CF_3$
P1792	H	H	Cl	Cl	CF_3	H	Cl	O	CH_2	O	$CH(CH_3)CF_3$
P1793	H	H	Cl	Cl	CF_3	H	CF_3	O	CH_2	O	$CH(CH_3)CF_3$
P1794	H	H	Cl	Cl	CF_3	H	CH_3	O	CH_2	O	$CH(CH_3)CF_3$
P1795	H	H	Cl	Cl	CF_3	CF_3	H	O	$CH(CH_3)$	O	CH_2CF_3
P1796	H	H	Cl	Cl	CF_3	CF_3	Br	O	$CH(CH_3)$	O	CH_2CF_3
P1797	H	H	Cl	Cl	CF_3	CF_3	Cl	O	$CH(CH_3)$	O	CH_2CF_3
P1798	H	H	Cl	Cl	CF_3	CF_3	CF_3	O	$CH(CH_3)$	O	CH_2CF_3
P1799	H	H	Cl	Cl	CF_3	CF_3	CH_3	O	$CH(CH_3)$	O	CH_2CF_3
P1800	H	H	Cl	Cl	CF_2CF_3	H	H	O	$CH(CH_3)$	O	CH_2CF_3
P1801	H	H	Cl	Cl	CF_2CF_3	H	Br	O	$CH(CH_3)$	O	CH_2CF_3
P1802	H	H	Cl	Cl	CF_2CF_3	H	Cl	O	$CH(CH_3)$	O	CH_2CF_3
P1803	H	H	Cl	Cl	CF_2CF_3	H	CF_3	O	$CH(CH_3)$	O	CH_2CF_3
P1804	H	H	Cl	Cl	CF_2CF_3	H	CH_3	O	$CH(CH_3)$	O	CH_2CF_3
P1805	H	H	Cl	Cl	CF_3	H	H	O	$CH(CH_3)$	O	CH_2CF_3

P1806	H	H	Cl	Cl	CF_3	H	Br	O	$CH(CH_3)$	O	CH_2CF_3
P1807	H	H	Cl	Cl	CF_3	H	Cl	O	$CH(CH_3)$	O	CH_2CF_3
P1808	H	H	Cl	Cl	CF_3	H	CF_3	O	$CH(CH_3)$	O	CH_2CF_3
P1809	H	H	Cl	Cl	CF_3	H	CH_3	O	$CH(CH_3)$	O	CH_2CF_3
P1810	H	H	Cl	Cl	CF_3	H	H	O	$CH(CH_3)$	S	CH_2CF_3
P1811	H	H	Cl	Cl	CF_3	H	Br	O	$CH(CH_3)$	S	CH_2CF_3
P1812	H	H	Cl	Cl	CF_3	H	Cl	O	$CH(CH_3)$	S	CH_2CF_3
P1813	H	H	Cl	Cl	CF_3	H	CF_3	O	$CH(CH_3)$	S	CH_2CF_3
P1814	H	H	Cl	Cl	CF_3	H	CH_3	O	$CH(CH_3)$	S	CH_2CF_3
P1815	H	H	Cl	Cl	CF_3	H	H	S	$CH(CH_3)$	O	CH_2CF_3
P1816	H	H	Cl	Cl	CF_3	H	Br	S	$CH(CH_3)$	O	CH_2CF_3
P1817	H	H	Cl	Cl	CF_3	H	Cl	S	$CH(CH_3)$	O	CH_2CF_3
P1818	H	H	Cl	Cl	CF_3	H	CF_3	S	$CH(CH_3)$	O	CH_2CF_3
P1819	H	H	Cl	Cl	CF_3	H	CH_3	S	$CH(CH_3)$	O	CH_2CF_3
P1820	H	H	Cl	Cl	CF_3	H	H	O	$CH(CH_3)$	O	CH_2CHF_2
P1821	H	H	Cl	Cl	CF_3	H	Br	O	$CH(CH_3)$	O	CH_2CHF_2
P1822	H	H	Cl	Cl	CF_3	H	Cl	O	$CH(CH_3)$	O	CH_2CHF_2
P1823	H	H	Cl	Cl	CF_3	H	CF_3	O	$CH(CH_3)$	O	CH_2CHF_2
P1824	H	H	Cl	Cl	CF_3	H	CH_3	O	$CH(CH_3)$	O	CH_2CHF_2
P1825	H	H	Cl	Cl	CF_3	H	H	O	$CH(CH_3)$	O	CH_2CH_2F
P1826	H	H	Cl	Cl	CF_3	H	Br	O	$CH(CH_3)$	O	CH_2CH_2F
P1827	H	H	Cl	Cl	CF_3	H	Cl	O	$CH(CH_3)$	O	CH_2CH_2F
P1828	H	H	Cl	Cl	CF_3	H	CF_3	O	$CH(CH_3)$	O	CH_2CH_2F
P1829	H	H	Cl	Cl	CF_3	H	CH_3	O	$CH(CH_3)$	O	CH_2CH_2F
P1830	H	H	Cl	Cl	CF_3	H	H	O	$CH(CH_3)$	O	CH_2CH_3
P1831	H	H	Cl	Cl	CF_3	H	Br	O	$CH(CH_3)$	O	CH_2CH_3
P1832	H	H	Cl	Cl	CF_3	H	Cl	O	$CH(CH_3)$	O	CH_2CH_3

P1833	H	H	Cl	Cl	CF_3	H	CF_3	O	$CH(CH_3)$	O	CH_2CH_3
P1834	H	H	Cl	Cl	CF_3	H	CH_3	O	$CH(CH_3)$	O	CH_2CH_3
P1835	H	H	Cl	Cl	CF_3	H	H	O	$CH(CH_3)$	O	$CH(CH_3)CF_3$
P1836	H	H	Cl	Cl	CF_3	H	Br	O	$CH(CH_3)$	O	$CH(CH_3)CF_3$
P1837	H	H	Cl	Cl	CF_3	H	Cl	O	$CH(CH_3)$	O	$CH(CH_3)CF_3$
P1838	H	H	Cl	Cl	CF_3	H	CF_3	O	$CH(CH_3)$	O	$CH(CH_3)CF_3$
P1839	H	H	Cl	Cl	CF_3	H	CH_3	O	$CH(CH_3)$	O	$CH(CH_3)CF_3$
P1840	H	H	Cl	Cl	CF_3	H	H	O	$CH(CH_3)$	O	$CH_2CH_2CF_3$
P1841	H	H	Cl	Cl	CF_3	H	Br	O	$CH(CH_3)$	O	$CH_2CH_2CF_3$
P1842	H	H	Cl	Cl	CF_3	H	Cl	O	$CH(CH_3)$	O	$CH_2CH_2CF_3$
P1843	H	H	Cl	Cl	CF_3	H	CF_3	O	$CH(CH_3)$	O	$CH_2CH_2CF_3$
P1844	H	H	Cl	Cl	CF_3	H	CH_3	O	$CH(CH_3)$	O	$CH_2CH_2CF_3$
P1845	H	H	Cl	Cl	CF_3	H	H	O	$CH(CH_2CH_3)$	O	CH_2CF_3
P1846	H	H	Cl	Cl	CF_3	H	Br	O	$CH(CH_2CH_3)$	O	CH_2CF_3
P1847	H	H	Cl	Cl	CF_3	H	Cl	O	$CH(CH_2CH_3)$	O	CH_2CF_3
P1848	H	H	Cl	Cl	CF_3	H	CF_3	O	$CH(CH_2CH_3)$	O	CH_2CF_3
P1849	H	H	Cl	Cl	CF_3	H	CH_3	O	$CH(CH_2CH_3)$	O	CH_2CF_3
P1850	v	H	Cl	Cl	CF_3	H	H	O	$C(CH_3)_2$	O	CH_2CF_3
P1851	H	H	Cl	Cl	CF_3	H	Br	O	$C(CH_3)_2$	O	CH_2CF_3
P1852	H	H	Cl	Cl	CF_3	H	Cl	O	$C(CH_3)_2$	O	CH_2CF_3
P1853	H	H	Cl	Cl	CF_3	H	CF_3	O	$C(CH_3)_2$	O	CH_2CF_3
P1854	H	H	Cl	Cl	CF_3	H	CH_3	O	$C(CH_3)_2$	O	CH_2CF_3
P1855	H	H	Cl	Cl	CF_3	H	H	O	CH_2CH_2	O	CH_2CF_3
P1856	H	H	Cl	Cl	CF_3	H	Br	O	CH_2CH_2	O	CH_2CF_3
P1857	H	H	Cl	Cl	CF_3	H	Cl	O	CH_2CH_2	O	CH_2CF_3
P1858	H	H	Cl	Cl	CF_3	H	CF_3	O	CH_2CH_2	O	CH_2CF_3
P1859	H	H	Cl	Cl	CF_3	H	CH_3	O	CH_2CH_2	O	CH_2CF_3

P1860	H	Cl	F	Cl	CF_3	H	H	O	CH_2	O	CH_2CF_3
P1861	H	Cl	F	Cl	CF_3	H	Cl	O	CH_2	O	CH_2CF_3
P1862	H	Cl	F	Cl	CF_3	H	CF_3	O	CH_2	O	CH_2CF_3
P1863	H	Cl	F	Cl	CF_3	H	H	O	CH_2	S	CH_2CF_3
P1864	H	Cl	F	Cl	CF_3	H	Br	O	CH_2	S	CH_2CF_3
P1865	H	Cl	F	Cl	CF_3	H	Cl	O	CH_2	S	CH_2CF_3
P1866	H	Cl	F	Cl	CF_3	H	CF_3	O	CH_2	S	CH_2CF_3
P1867	H	Cl	F	Cl	CF_3	H	CH_3	O	CH_2	S	CH_2CF_3
P1868	H	Cl	F	Cl	CF_3	H	H	S	CH_2	O	CH_2CF_3
P1869	H	Cl	F	Cl	CF_3	H	Br	S	CH_2	O	CH_2CF_3
P1870	H	Cl	F	Cl	CF_3	H	Cl	S	CH_2	O	CH_2CF_3
P1871	H	Cl	F	Cl	CF_3	H	CF_3	S	CH_2	O	CH_2CF_3
P1872	H	Cl	F	Cl	CF_3	H	CH_3	S	CH_2	O	CH_2CF_3
P1873	H	Cl	F	Cl	CF_3	H	H	O	CH_2	O	CH_2CHF_2
P1874	H	Cl	F	Cl	CF_3	H	Br	O	CH_2	O	CH_2CHF_2
P1875	H	Cl	F	Cl	CF_3	H	Cl	O	CH_2	O	CH_2CHF_2
P1876	H	Cl	F	Cl	CF_3	H	CF_3	O	CH_2	O	CH_2CHF_2
P1877	H	Cl	F	Cl	CF_3	H	CH_3	O	CH_2	O	CH_2CHF_2
P1878	H	Cl	F	Cl	CF_3	H	H	O	CH_2	O	CH_2CH_2F
P1879	H	Cl	F	Cl	CF_3	H	Br	O	CH_2	O	CH_2CH_2F
P1880	H	Cl	F	Cl	CF_3	H	Cl	O	CH_2	O	CH_2CH_2F
P1881	H	Cl	F	Cl	CF_3	H	CF_3	O	CH_2	O	CH_2CH_2F
P1882	H	Cl	F	Cl	CF_3	H	CH_3	O	CH_2	O	CH_2CH_2F
P1883	H	Cl	F	Cl	CF_3	H	H	O	CH_2	O	CH_2CH_3
P1884	H	Cl	F	Cl	CF_3	H	Br	O	CH_2	O	CH_2CH_3
P1885	H	Cl	F	Cl	CF_3	H	Cl	O	CH_2	O	CH_2CH_3
P1886	H	Cl	F	Cl	CF_3	H	CF_3	O	CH_2	O	CH_2CH_3

P1887	H	Cl	F	Cl	CF_3	H	CH_3	O	CH_2	O	CH_2CH_3
P1888	H	Cl	F	Cl	CF_3	CF_3	H	O	CH_2	O	CH_2CF_3
P1889	H	Cl	F	Cl	CF_3	CF_3	Br	O	CH_2	O	CH_2CF_3
P1890	H	Cl	F	Cl	CF_3	CF_3	Cl	O	CH_2	O	CH_2CF_3
P1891	H	Cl	F	Cl	CF_3	CF_3	CF_3	O	CH_2	O	CH_2CF_3
P1892	H	Cl	F	Cl	CF_3	CF_3	CH_3	O	CH_2	O	CH_2CF_3
P1893	H	Cl	F	Cl	CF_2CF_3	H	H	O	CH_2	O	CH_2CF_3
P1894	H	Cl	F	Cl	CF_2CF_3	H	Br	O	CH_2	O	CH_2CF_3
P1895	H	Cl	F	Cl	CF_2CF_3	H	Cl	O	CH_2	O	CH_2CF_3
P1896	H	Cl	F	Cl	CF_2CF_3	H	CF_3	O	CH_2	O	CH_2CF_3
P1897	H	Cl	F	Cl	CF_2CF_3	H	CH_3	O	CH_2	O	CH_2CF_3
P1898	H	Cl	F	Cl	CF_3	H	H	O	CH_2	O	$CH(CH_3)CF_3$
P1899	H	Cl	F	Cl	CF_3	H	Br	O	CH_2	O	$CH(CH_3)CF_3$
P1900	H	Cl	F	Cl	CF_3	H	Cl	O	CH_2	O	$CH(CH_3)CF_3$
P1901	H	Cl	F	Cl	CF_3	H	CF_3	O	CH_2	O	$CH(CH_3)CF_3$
P1902	H	Cl	F	Cl	CF_3	H	CH_3	O	CH_2	O	$CH(CH_3)CF_3$
P1903	H	Cl	F	Cl	CF_3	CF_3	H	O	$CH(CH_3)$	O	CH_2CF_3
P1904	H	Cl	F	Cl	CF_3	CF_3	Br	O	$CH(CH_3)$	O	CH_2CF_3
P1905	H	Cl	F	Cl	CF_3	CF_3	Cl	O	$CH(CH_3)$	O	CH_2CF_3
P1906	H	Cl	F	Cl	CF_3	CF_3	CF_3	O	$CH(CH_3)$	O	CH_2CF_3
P1907	H	Cl	F	Cl	CF_3	CF_3	CH_3	O	$CH(CH_3)$	O	CH_2CF_3
P1908	H	Cl	F	Cl	CF_2CF_3	H	H	O	$CH(CH_3)$	O	CH_2CF_3
P1909	H	Cl	F	Cl	CF_2CF_3	H	Br	O	$CH(CH_3)$	O	CH_2CF_3
P1910	H	Cl	F	Cl	CF_2CF_3	H	Cl	O	$CH(CH_3)$	O	CH_2CF_3
P1911	H	Cl	F	Cl	CF_2CF_3	H	CF_3	O	$CH(CH_3)$	O	CH_2CF_3
P1912	H	Cl	F	Cl	CF_2CF_3	H	CH_3	O	$CH(CH_3)$	O	CH_2CF_3
P1913	H	Cl	F	Cl	CF_3	H	H	O	$CH(CH_3)$	O	CH_2CF_3

P1914	H	Cl	F	Cl	CF_3	H	Br	O	$CH(CH_3)$	O	CH_2CF_3
P1915	H	Cl	F	Cl	CF_3	H	Cl	O	$CH(CH_3)$	O	CH_2CF_3
P1916	H	Cl	F	Cl	CF_3	H	CF_3	O	$CH(CH_3)$	O	CH_2CF_3
P1917	H	Cl	F	Cl	CF_3	H	CH_3	O	$CH(CH_3)$	O	CH_2CF_3
P1918	H	Cl	F	Cl	CF_3	H	H	O	$CH(CH_3)$	S	CH_2CF_3
P1919	H	Cl	F	Cl	CF_3	H	Br	O	$CH(CH_3)$	S	CH_2CF_3
P1920	H	Cl	F	Cl	CF_3	H	Cl	O	$CH(CH_3)$	S	CH_2CF_3
P1921	H	Cl	F	Cl	CF_3	H	CF_3	O	$CH(CH_3)$	S	CH_2CF_3
P1922	H	Cl	F	Cl	CF_3	H	CH_3	O	$CH(CH_3)$	S	CH_2CF_3
P1923	H	Cl	F	Cl	CF_3	H	H	S	$CH(CH_3)$	O	CH_2CF_3
P1924	H	Cl	F	Cl	CF_3	H	Br	S	$CH(CH_3)$	O	CH_2CF_3
P1925	H	Cl	F	Cl	CF_3	H	Cl	S	$CH(CH_3)$	O	CH_2CF_3
P1926	H	Cl	F	Cl	CF_3	H	CF_3	S	$CH(CH_3)$	O	CH_2CF_3
P1927	H	Cl	F	Cl	CF_3	H	CH_3	S	$CH(CH_3)$	O	CH_2CF_3
P1928	H	Cl	F	Cl	CF_3	H	H	O	$CH(CH_3)$	O	CH_2CHF_2
P1929	H	Cl	F	Cl	CF_3	H	Br	O	$CH(CH_3)$	O	CH_2CHF_2
P1930	H	Cl	F	Cl	CF_3	H	Cl	O	$CH(CH_3)$	O	CH_2CHF_2
P1931	H	Cl	F	Cl	CF_3	H	CF_3	O	$CH(CH_3)$	O	CH_2CHF_2
P1932	H	Cl	F	Cl	CF_3	H	CH_3	O	$CH(CH_3)$	O	CH_2CHF_2
P1933	H	Cl	F	Cl	CF_3	H	H	O	$CH(CH_3)$	O	CH_2CH_2F
P1934	H	Cl	F	Cl	CF_3	H	Br	O	$CH(CH_3)$	O	CH_2CH_2F
P1935	H	Cl	F	Cl	CF_3	H	Cl	O	$CH(CH_3)$	O	CH_2CH_2F
P1936	H	Cl	F	Cl	CF_3	H	CF_3	O	$CH(CH_3)$	O	CH_2CH_2F
P1937	H	Cl	F	Cl	CF_3	H	CH_3	O	$CH(CH_3)$	O	CH_2CH_2F
P1938	H	Cl	F	Cl	CF_3	H	H	O	$CH(CH_3)$	O	CH_2CH_3
P1939	H	Cl	F	Cl	CF_3	H	Br	O	$CH(CH_3)$	O	CH_2CH_3
P1940	H	Cl	F	Cl	CF_3	H	Cl	O	$CH(CH_3)$	O	CH_2CH_3

P1941	H	Cl	F	Cl	CF_3	H	CF_3	O	$CH(CH_3)$	O	CH_2CH_3
P1942	H	Cl	F	Cl	CF_3	H	CH_3	O	$CH(CH_3)$	O	CH_2CH_3
P1943	H	Cl	F	Cl	CF_3	H	H	O	$CH(CH_3)$	O	$CH(CH_3)CF_3$
P1944	H	Cl	F	Cl	CF_3	H	Br	O	$CH(CH_3)$	O	$CH(CH_3)CF_3$
P1945	H	Cl	F	Cl	CF_3	H	Cl	O	$CH(CH_3)$	O	$CH(CH_3)CF_3$
P1946	H	Cl	F	Cl	CF_3	H	CF_3	O	$CH(CH_3)$	O	$CH(CH_3)CF_3$
P1947	H	Cl	F	Cl	CF_3	H	CH_3	O	$CH(CH_3)$	O	$CH(CH_3)CF_3$
P1948	H	Cl	F	Cl	CF_3	H	H	O	$CH(CH_3)$	O	$CH_2CH_2CF_3$
P1949	H	Cl	F	Cl	CF_3	H	Br	O	$CH(CH_3)$	O	$CH_2CH_2CF_3$
P1950	H	Cl	F	Cl	CF_3	H	Cl	O	$CH(CH_3)$	O	$CH_2CH_2CF_3$
P1951	H	Cl	F	Cl	CF_3	H	CF_3	O	$CH(CH_3)$	O	$CH_2CH_2CF_3$
P1952	H	Cl	F	Cl	CF_3	H	CH_3	O	$CH(CH_3)$	O	$CH_2CH_2CF_3$
P1953	H	Cl	F	Cl	CF_3	H	H	O	$CH(CH_2CH_3)$	O	CH_2CF_3
P1954	H	Cl	F	Cl	CF_3	H	Br	O	$CH(CH_2CH_3)$	O	CH_2CF_3
P1955	H	Cl	F	Cl	CF_3	H	Cl	O	$CH(CH_2CH_3)$	O	CH_2CF_3
P1956	H	Cl	F	Cl	CF_3	H	CF_3	O	$CH(CH_2CH_3)$	O	CH_2CF_3
P1957	H	Cl	F	Cl	CF_3	H	CH_3	O	$CH(CH_2CH_3)$	O	CH_2CF_3
P1958	H	Cl	F	Cl	CF_3	H	H	O	$C(CH_3)_2$	O	CH_2CF_3
P1959	H	Cl	F	Cl	CF_3	H	Br	O	$C(CH_3)_2$	O	CH_2CF_3
P1960	H	Cl	F	Cl	CF_3	H	Cl	O	$C(CH_3)_2$	O	CH_2CF_3
P1961	H	Cl	F	Cl	CF_3	H	CF_3	O	$C(CH_3)_2$	O	CH_2CF_3
P1962	H	Cl	F	Cl	CF_3	H	CH_3	O	$C(CH_3)_2$	O	CH_2CF_3
P1963	H	Cl	F	Cl	CF_3	H	H	O	CH_2CH_2	O	CH_2CF_3
P1964	H	Cl	F	Cl	CF_3	H	Br	O	CH_2CH_2	O	CH_2CF_3
P1965	H	Cl	F	Cl	CF_3	H	Cl	O	CH_2CH_2	O	CH_2CF_3
P1966	H	Cl	F	Cl	CF_3	H	CF_3	O	CH_2CH_2	O	CH_2CF_3
P1967	H	Cl	F	Cl	CF_3	H	CH_3	O	CH_2CH_2	O	CH_2CF_3

P1968	H	Br	H	Br	CF_3	H	H	O	CH_2	O	CH_2CF_3
P1969	H	Br	H	Br	CF_3	H	Br	O	CH_2	O	CH_2CF_3
P1970	H	Br	H	Br	CF_3	H	Cl	O	CH_2	O	CH_2CF_3
P1971	H	Br	H	Br	CF_3	H	CF_3	O	CH_2	O	CH_2CF_3
P1972	H	Br	H	Br	CF_3	H	CH_3	O	CH_2	O	CH_2CF_3
P1973	H	Br	H	Br	CF_3	H	H	O	CH_2	S	CH_2CF_3
P1974	H	Br	H	Br	CF_3	H	Br	O	CH_2	S	CH_2CF_3
P1975	H	Br	H	Br	CF_3	H	Cl	O	CH_2	S	CH_2CF_3
P1976	H	Br	H	Br	CF_3	H	CF_3	O	CH_2	S	CH_2CF_3
P1977	H	Br	H	Br	CF_3	H	CH_3	O	CH_2	S	CH_2CF_3
P1978	H	Br	H	Br	CF_3	H	H	S	CH_2	O	CH_2CF_3
P1979	H	Br	H	Br	CF_3	H	Br	S	CH_2	O	CH_2CF_3
P1980	H	Br	H	Br	CF_3	H	Cl	S	CH_2	O	CH_2CF_3
P1981	H	Br	H	Br	CF_3	H	CF_3	S	CH_2	O	CH_2CF_3
P1982	H	Br	H	Br	CF_3	H	CH_3	S	CH_2	O	CH_2CF_3
P1983	H	Br	H	Br	CF_3	H	H	O	CH_2	O	CH_2CHF_2
P1984	H	Br	H	Br	CF_3	H	Br	O	CH_2	O	CH_2CHF_2
P1985	H	Br	H	Br	CF_3	H	Cl	O	CH_2	O	CH_2CHF_2
P1986	H	Br	H	Br	CF_3	H	CF_3	O	CH_2	O	CH_2CHF_2
P1987	H	Br	H	Br	CF_3	H	CH_3	O	CH_2	O	CH_2CHF_2
P1988	H	Br	H	Br	CF_3	H	H	O	CH_2	O	CH_2CH_2F
P1989	H	Br	H	Br	CF_3	H	Br	O	CH_2	O	CH_2CH_2F
P1990	H	Br	H	Br	CF_3	H	Cl	O	CH_2	O	CH_2CH_2F
P1991	H	Br	H	Br	CF_3	H	CF_3	O	CH_2	O	CH_2CH_2F
P1992	H	Br	H	Br	CF_3	H	CH_3	O	CH_2	O	CH_2CH_2F
P1993	H	Br	H	Br	CF_3	H	H	O	CH_2	O	CH_2CH_3
P1994	H	Br	H	Br	CF_3	H	Br	O	CH_2	O	CH_2CH_3

P1995	H	Br	H	Br	CF_3	H	Cl	O	CH_2	O	CH_2CH_3
P1996	H	Br	H	Br	CF_3	H	CF_3	O	CH_2	O	CH_2CH_3
P1997	H	Br	H	Br	CF_3	H	CH_3	O	CH_2	O	CH_2CH_3
P1998	H	Br	H	Br	CF_3	CF_3	H	O	CH_2	O	CH_2CF_3
P1999	H	Br	H	Br	CF_3	CF_3	Br	O	CH_2	O	CH_2CF_3
P2000	H	Br	H	Br	CF_3	CF_3	Cl	O	CH_2	O	CH_2CF_3
P2001	H	Br	H	Br	CF_3	CF_3	CF_3	O	CH_2	O	CH_2CF_3
P2002	H	Br	H	Br	CF_3	CF_3	CH_3	O	CH_2	O	CH_2CF_3
P2003	H	Br	H	Br	CF_2CF_3	H	H	O	CH_2	O	CH_2CF_3
P2004	H	Br	H	Br	CF_2CF_3	H	Br	O	CH_2	O	CH_2CF_3
P2005	H	Br	H	Br	CF_2CF_3	H	Cl	O	CH_2	O	CH_2CF_3
P2006	H	Br	H	Br	CF_2CF_3	H	CF_3	O	CH_2	O	CH_2CF_3
P2007	H	Br	H	Br	CF_2CF_3	H	CH_3	O	CH_2	O	CH_2CF_3
P2008	H	Br	H	Br	CF_3	H	H	O	CH_2	O	$CH(CH_3)CF_3$
P2009	H	Br	H	Br	CF_3	H	Br	O	CH_2	O	$CH(CH_3)CF_3$
P2010	H	Br	H	Br	CF_3	H	Cl	O	CH_2	O	$CH(CH_3)CF_3$
P2011	H	Br	H	Br	CF_3	H	CF_3	O	CH_2	O	$CH(CH_3)CF_3$
P2012	H	Br	H	Br	CF_3	H	CH_3	O	CH_2	O	$CH(CH_3)CF_3$
P2013	H	Br	H	Br	CF_3	CF_3	H	O	$CH(CH_3)$	O	CH_2CF_3
P2014	H	Br	H	Br	CF_3	CF_3	Br	O	$CH(CH_3)$	O	CH_2CF_3
P2015	H	Br	H	Br	CF_3	CF_3	Cl	O	$CH(CH_3)$	O	CH_2CF_3
P2016	H	Br	H	Br	CF_3	CF_3	CF_3	O	$CH(CH_3)$	O	CH_2CF_3
P2017	H	Br	H	Br	CF_3	CF_3	CH_3	O	$CH(CH_3)$	O	CH_2CF_3
P2018	H	Br	H	Br	CF_2CF_3	H	H	O	$CH(CH_3)$	O	CH_2CF_3
P2019	H	Br	H	Br	CF_2CF_3	H	Br	O	$CH(CH_3)$	O	CH_2CF_3
P2020	H	Br	H	Br	CF_2CF_3	H	Cl	O	$CH(CH_3)$	O	CH_2CF_3
P2021	H	Br	H	Br	CF_2CF_3	H	CF_3	O	$CH(CH_3)$	O	CH_2CF_3

P2022	H	Br	H	Br	CF_2CF_3	H	CH_3	O	CH (CH_3)	O	CH_2CF_3
P2023	H	Br	H	Br	CF_3	H	H	O	CH(CH_3)	O	CH_2CF_3
P2024	H	Br	H	Br	CF_3	H	Cl	O	CH (CH_3)	O	CH_2CF_3
P2025	H	Br	H	Br	CF_3	H	H	O	CH(CH_3)	S	CH_2CF_3
P2026	H	Br	H	Br	CF_3	H	Br	O	CH(CH_3)	S	CH_2CF_3
P2027	H	Br	H	Br	CF_3	H	Cl	O	CH (CH_3)	S	CH_2CF_3
P2028	H	Br	H	Br	CF_3	H	CF_3	O	CH(CH_3)	S	CH_2CF_3
P2029	H	Br	H	Br	CF_3	H	CH_3	O	CH (CH_3)	S	CH_2CF_3
P2030	H	Br	H	Br	CF_3	H	H	S	CH(CH_3)	O	CH_2CF_3
P2031	H	Br	H	Br	CF_3	H	Br	S	CH(CH_3)	O	CH_2CF_3
P2032	H	Br	H	Br	CF_3	H	Cl	S	CH (CH_3)	O	CH_2CF_3
P2033	H	Br	H	Br	CF_3	H	CF_3	S	CH(CH_3)	O	CH_2CF_3
P2034	H	Br	H	Br	CF_3	H	CH_3	S	CH (CH_3)	O	CH_2CF_3
P2035	H	Br	H	Br	CF_3	H	H	O	CH(CH_3)	O	CH_2CHF_2
P2036	H	Br	H	Br	CF_3	H	Br	O	CH(CH_3)	O	CH_2CHF_2
P2037	H	Br	H	Br	CF_3	H	Cl	O	CH (CH_3)	O	CH_2CHF_2
P2038	H	Br	H	Br	CF_3	H	CF_3	O	CH(CH_3)	O	CH_2CHF_2
P2039	H	Br	H	Br	CF_3	H	CH_3	O	CH (CH_3)	O	CH_2CHF_2
P2040	H	Br	H	Br	CF_3	H	H	O	CH(CH_3)	O	CH_2CH_2F
P2041	H	Br	H	Br	CF_3	H	Br	O	CH(CH_3)	O	CH_2CH_2F
P2042	H	Br	H	Br	CF_3	H	Cl	O	CH(CH_3)	O	CH_2CH_2F
P2043	H	Br	H	Br	CF_3	H	CF_3	O	CH(CH_3)	O	CH_2CH_2F
P2044	H	Br	H	Br	CF_3	H	CH_3	O	CH(CH_3)	O	CH_2CH_2F
P2045	H	Br	H	Br	CF_3	H	H	O	CH(CH_3)	O	CH_2CH_3
P2046	H	Br	H	Br	CF_3	H	Br	O	CH(CH_3)	O	CH_2CH_3
P2047	H	Br	H	Br	CF_3	H	Cl	O	CH(CH_3)	O	CH_2CH_3
P2048	H	Br	H	Br	CF_3	H	CF_3	O	CH(CH_3)	O	CH_2CH_3

P2049	H	Br	H	Br	CF_3	H	CH_3	O	$CH(CH_3)$	O	CH_2CH_3
P2050	H	Br	H	Br	CF_3	H	H	O	$CH(CH_3)$	O	$CH(CH_3)CF_3$
P2051	H	Br	H	Br	CF_3	H	Br	O	$CH(CH_3)$	O	$CH(CH_3)CF_3$
P2052	H	Br	H	Br	CF_3	H	Cl	O	$CH(CH_3)$	O	$CH(CH_3)CF_3$
P2053	H	Br	H	Br	CF_3	H	CF_3	O	$CH(CH_3)$	O	$CH(CH_3)CF_3$
P2054	H	Br	H	Br	CF_3	H	CH_3	O	$CH(CH_3)$	O	$CH(CH_3)CF_3$
P2055	H	Br	H	Br	CF_3	H	H	O	$CH(CH_3)$	O	$CH_2CH_2CF_3$
P2056	H	Br	H	Br	CF_3	H	Br	O	$CH(CH_3)$	O	$CH_2CH_2CF_3$
P2057	H	Br	H	Br	CF_3	H	Cl	O	$CH(CH_3)$	O	$CH_2CH_2CF_3$
P2058	H	Br	H	Br	CF_3	H	CF_3	O	$CH(CH_3)$	O	$CH_2CH_2CF_3$
P2059	H	Br	H	Br	CF_3	H	CH_3	O	$CH(CH_3)$	O	$CH_2CH_2CF_3$
P2060	H	Br	H	Br	CF_3	H	H	O	$CH(CH_2CH_3)$	O	CH_2CF_3
P2061	H	Br	H	Br	CF_3	H	Br	O	$CH(CH_2CH_3)$	O	CH_2CF_3
P2062	H	Br	H	Br	CF_3	H	Cl	O	$CH(CH_2CH_3)$	O	CH_2CF_3
P2063	H	Br	H	Br	CF_3	H	CF_3	O	$CH(CH_2CH_3)$	O	CH_2CF_3
P2064	H	Br	H	Br	CF_3	H	CH_3	O	$CH(CH_2CH_3)$	O	CH_2CF_3
P2065	H	Br	H	Br	CF_3	H	H	O	$C(CH_3)_2$	O	CH_2CF_3
P2066	H	Br	H	Br	CF_3	H	Br	O	$C(CH_3)_2$	O	CH_2CF_3
P2067	H	Br	H	Br	CF_3	H	Cl	O	$C(CH_3)_2$	O	CH_2CF_3
P2068	H	Br	H	Br	CF_3	H	CF_3	O	$C(CH_3)_2$	O	CH_2CF_3
P2069	H	Br	H	Br	CF_3	H	CH_3	O	$C(CH_3)_2$	O	CH_2CF_3
P2070	H	Br	H	Br	CF_3	H	H	O	CH_2CH_2	O	CH_2CF_3
P2071	H	Br	H	Br	CF_3	H	Br	O	CH_2CH_2	O	CH_2CF_3
P2072	H	Br	H	Br	CF_3	H	Cl	O	CH_2CH_2	O	CH_2CF_3
P2073	H	Br	H	Br	CF_3	H	CF_3	O	CH_2CH_2	O	CH_2CF_3
P2074	H	Br	H	Br	CF_3	H	CH_3	O	CH_2CH_2	O	CH_2CF_3

Example A: BIOASSAYS ON BEET ARMYWORM ("BAW") AND CORN EARWORM ("CEW") AND CABBAGE LOOPER ("CL")

BAW has few effective parasites, diseases, or predators to lower its population. BAW infests many weeds, trees, grasses, legumes, and field crops. In various places, it is of economic concern upon asparagus, cotton, corn, soybeans, tobacco, alfalfa, sugar beets, peppers, tomatoes, potatoes, onions, peas, sunflowers, and citrus, among other plants. CEW is known to attack corn and tomatoes, but it also attacks artichoke, asparagus, cabbage, cantaloupe, collards, cowpeas, cucumbers, eggplant, lettuce, lima beans, melon, okra, peas, peppers, potatoes, pumpkin, snap beans, spinach, squash, sweet potatoes, and watermelon, among other plants. CEW is also known to be resistant to certain insecticides. CL is also known to be resistant to certain insecticides. Consequently, because of the above factors control of these pests is important. Furthermore, molecules that control these pests are useful in controlling other pests.

Certain molecules disclosed in this document were tested against BAW, CEW and CL using procedures described in the following examples. In the reporting of the results, the "**BAW & CEW & CL Rating Table**" was used (See Table Section).

BIOASSAYS ON BAW (*Spodoptera exigua*)

Bioassays on BAW were conducted using a 128-well diet tray assay. One to five second instar BAW larvae were placed in each well (3 mL) of the diet tray that had been previously filled with 1 mL of artificial diet to which 50 μg/cm^2 of the test compound (dissolved in 50 μL of 90:10 acetone-water mixture) had been applied (to each of eight wells) and then allowed to dry. Trays were covered with a clear self-adhesive cover, and held at 25 °C, 14:10 light-dark for five to seven days. Percent mortality was recorded for the larvae in each well; activity in the eight wells was then averaged. The results are indicated in the tables entitled "**Table 3: Assay Results Part 1**" and "**Table 4: Assay Results Part 2**" (See Table Section).

BIOASSAYS ON CEW (*Helicoverpa zea*)

Bioassays on CEW were conducted using a 128-well diet tray assay. One to five second instar CEW larvae were placed in each well (3 mL) of the diet tray that had been previously filled with 1 mL of artificial diet to which 50 μg /cm^2 of the test compound (dissolved in 50 μL of 90:10 acetone–water mixture) had been applied (to each of eight wells) and then allowed to dry. Trays were covered with a clear self-adhesive cover, and held at 25 °C, 14:10 light-dark for five to seven days. Percent mortality was recorded for the

larvae in each well; activity in the eight wells was then averaged. The results are indicated in the table entitled "**Table 3: Assay Results Part 1**" (See Table Section).

Bioassays on CL (*Trichoplusia ni*)

Bioassays on CL were conducted using a 128-well diet tray assay. One to five second instar CL larvae were placed in each well (3 mL) of the diet tray that had been previously filled with 1 mL of artificial diet to which 50 μg /cm^2 of the test compound (dissolved in 50 μL of 90:10 acetone–water mixture) had been applied (to each of eight wells) and then allowed to dry. Trays were covered with a clear self-adhesive cover, and held at 25 °C, 14:10 light-dark for five to seven days. Percent mortality was recorded for the larvae in each well; activity in the eight wells was then averaged. The results are indicated in the table entitled "**Table 4: Assay Results Part 2**" (See Table Section).

Example B: BIOASSAYS ON GREEN PEACH APHID ("GPA") (*Myzus persicae*).

GPA is the most significant aphid pest of peach trees, causing decreased growth, shriveling of the leaves, and the death of various tissues. It is also hazardous because it acts as a vector for the transport of plant viruses, such as potato virus Y and potato leafroll virus to members of the nightshade/potato family *Solanaceae*, and various mosaic viruses to many other food crops. GPA attacks such plants as broccoli, burdock, cabbage, carrot, cauliflower, daikon, eggplant, green beans, lettuce, macadamia, papaya, peppers, sweet potatoes, tomatoes, watercress, and zucchini, among other plants. GPA also attacks many ornamental crops such as carnation, chrysanthemum, flowering white cabbage, poinsettia, and roses. GPA has developed resistance to many pesticides.

Certain molecules disclosed in this document were tested against GPA using procedures described in the following example. In the reporting of the results, the "**GPA Rating Table**" was used (See Table Section).

Cabbage seedlings grown in 3-inch pots, with 2-3 small (3-5 cm) true leaves, were used as test substrate. The seedlings were infested with 20-50 GPA (wingless adult and nymph stages) one day prior to chemical application. Four pots with individual seedlings were used for each treatment. Test compounds (2 mg) were dissolved in 2 mL of acetone/MeOH (1:1) solvent, forming stock solutions of 1000 ppm test compound. The stock solutions were diluted 5X with 0.025% Tween 20 in water to obtain the solution at 200 ppm test compound. A hand-held aspirator-type sprayer was used for spraying a solution to both sides of cabbage leaves until runoff. Reference plants (solvent check) were sprayed with the diluent only containing 20% by volume of acetone/MeOH (1:1) solvent. Treated plants were held in a holding room for three days at approximately 25 °C and ambient relative humidity

(RH) prior to grading. Evaluation was conducted by counting the number of live aphids per plant under a microscope. Percent Control was measured by using Abbott's correction formula (W.S. Abbott, "A Method of Computing the Effectiveness of an Insecticide" J. Econ. Entomol. 18 (1925), pp.265-267) as follows.

Corrected % Control = 100 * (X - Y) / X

where

X = No. of live aphids on solvent check plants and

Y = No. of live aphids on treated plants

The results are indicated in the tables entitled "**Table 3: Assay Results Part 1**" and "**Table 4: Assay Results Part 2**" (See Table Section).

PESTICIDALLY ACCEPTABLE ACID ADDITION SALTS, SALT DERIVATIVES, SOLVATES, ESTER DERIVATIVES, POLYMORPHS, ISOTOPES AND RADIONUCLIDES

Molecules of Formula One may be formulated into pesticidally acceptable acid addition salts. By way of a non-limiting example, an amine function can form salts with hydrochloric, hydrobromic, sulfuric, phosphoric, acetic, benzoic, citric, malonic, salicylic, malic, fumaric, oxalic, succinic, tartaric, lactic, gluconic, ascorbic, maleic, aspartic, benzenesulfonic, methanesulfonic, ethanesulfonic, hydroxymethanesulfonic, and hydroxyethanesulfonic acids. Additionally, by way of a non-limiting example, an acid function can form salts including those derived from alkali or alkaline earth metals and those derived from ammonia and amines. Examples of preferred cations include sodium, potassium, and magnesium.

Molecules of Formula One may be formulated into salt derivatives. By way of a non-limiting example, a salt derivative can be prepared by contacting a free base with a sufficient amount of the desired acid to produce a salt. A free base may be regenerated by treating the salt with a suitable dilute aqueous base solution such as dilute aqueous NaOH, potassium carbonate, ammonia, and sodium bicarbonate. As an example, in many cases, a pesticide, such as 2,4-D, is made more water-soluble by converting it to its dimethylamine salt..

Molecules of Formula One may be formulated into stable complexes with a solvent, such that the complex remains intact after the non-complexed solvent is removed. These complexes are often referred to as "solvates." However, it is particularly desirable to form stable hydrates with water as the solvent.

Molecules of Formula One may be made into ester derivatives. These ester derivatives can then be applied in the same manner as the invention disclosed in this document is applied.

Molecules of Formula One may be made as various crystal polymorphs. Polymorphism is important in the development of agrochemicals since different crystal polymorphs or structures of the same molecule can have vastly different physical properties and biological performances.

Molecules of Formula One may be made with different isotopes. Of particular importance are molecules having ^{2}H (also known as deuterium) in place of ^{1}H.

Molecules of Formula One may be made with different radionuclides. Of particular importance are molecules having ^{14}C.

STEREOISOMERS

Molecules of Formula One may exist as one or more stereoisomers. Thus, certain molecules can be produced as racemic mixtures. It will be appreciated by those skilled in the art that one stereoisomer may be more active than the other stereoisomers. Individual stereoisomers may be obtained by known selective synthetic procedures, by conventional synthetic procedures using resolved starting materials, or by conventional resolution procedures. Certain molecules disclosed in this document can exist as two or more isomers. The various isomers include geometric isomers, diastereomers, and enantiomers. Thus, the molecules disclosed in this document include geometric isomers, racemic mixtures, individual stereoisomers, and optically active mixtures. It will be appreciated by those skilled in the art that one isomer may be more active than the others. The structures disclosed in the present disclosure are drawn in only one geometric form for clarity, but are intended to represent all geometric forms of the molecule.

COMBINATIONS

Molecules of Formula One may also be used in combination (such as, in a compositional mixture, or a simultaneous or sequential application) with one or more compounds having acaricidal, algicidal, avicidal, bactericidal, fungicidal, herbicidal, insecticidal, molluscicidal, nematicidal, rodenticidal, or virucidal properties. Additionally, the molecules of Formula One may also be used in combination (such as, in a compositional mixture, or a simultaneous or sequential application) with compounds that are antifeedants, bird repellents, chemosterilants, herbicide safeners, insect attractants, insect repellents, mammal repellents, mating disrupters, plant activators, plant growth regulators, or synergists. Examples of such compounds in the above groups that may be used with the Molecules of

Formula One are - (3-ethoxypropyl)mercury bromide, 1,2-dichloropropane, 1,3-dichloropropene, 1-methylcyclopropene, 1-naphthol, 2-(octylthio)ethanol, 2,3,5-tri-iodobenzoic acid, 2,3,6-TBA, 2,3,6-TBA-dimethylammonium, 2,3,6-TBA-lithium, 2,3,6-TBA-potassium, 2,3,6-TBA-sodium, 2,4,5-T, 2,4,5-T-2-butoxypropyl, 2,4,5-T-2-ethylhexyl, 2,4,5-T-3-butoxypropyl, 2,4,5-TB, 2,4,5-T-butometyl, 2,4,5-T-butotyl, 2,4,5-T-butyl, 2,4,5-T-isobutyl, 2,4,5-T-isoctyl, 2,4,5-T-isopropyl, 2,4,5-T-methyl, 2,4,5-T-pentyl, 2,4,5-T-sodium, 2,4,5-T-triethylammonium, 2,4,5-T-trolamine, 2,4-D, 2,4-D-2-butoxypropyl, 2,4-D-2-ethylhexyl, 2,4-D-3-butoxypropyl, 2,4-D-ammonium, 2,4-DB, 2,4-DB-butyl, 2,4-DB-dimethylammonium, 2,4-DB-isoctyl, 2,4-DB-potassium, 2,4-DB-sodium, 2,4-D-butotyl, 2,4-D-butyl, 2,4-D-diethylammonium, 2,4-D-dimethylammonium, 2,4-D-diolamine, 2,4-D-dodecylammonium, 2,4-DEB, 2,4-DEP, 2,4-D-ethyl, 2,4-D-heptylammonium, 2,4-D-isobutyl, 2,4-D-isoctyl, 2,4-D-isopropyl, 2,4-D-isopropylammonium, 2,4-D-lithium, 2,4-D-meptyl, 2,4-D-methyl, 2,4-D-octyl, 2,4-D-pentyl, 2,4-D-potassium, 2,4-D-propyl, 2,4-D-sodium, 2,4-D-tefuryl, 2,4-D-tetradecylammonium, 2,4-D-triethylammonium, 2,4-D-tris(2-hydroxypropyl)ammonium, 2,4-D-trolamine, 2iP, 2-methoxyethylmercury chloride, 2-phenylphenol, 3,4-DA, 3,4-DB, 3,4-DP, 4-aminopyridine, 4-CPA, 4-CPA-potassium, 4-CPA-sodium, 4-CPB, 4-CPP, 4-hydroxyphenethyl alcohol, 8-hydroxyquinoline sulfate, 8-phenylmercurioxyquinoline, abamectin, abscisic acid, ACC, acephate, acequinocyl, acetamiprid, acethion, acetochlor, acetophos, acetoprole, acibenzolar, acibenzolar-S-methyl, acifluorfen, acifluorfen-methyl, acifluorfen-sodium, aclonifen, acrep, acrinathrin, acrolein, acrylonitrile, acypetacs, acypetacs-copper, acypetacs-zinc, alachlor, alanycarb, albendazole, aldicarb, aldimorph, aldoxycarb, aldrin, allethrin, allicin, allidochlor, allosamidin, alloxydim, alloxydim-sodium, allyl alcohol, allyxycarb, alorac, *alpha*-cypermethrin, *alpha*-endosulfan, ametoctradin, ametridione, ametryn, amibuzin, amicarbazone, amicarthiazol, amidithion, amidoflumet, amidosulfuron, aminocarb, aminocyclopyrachlor, aminocyclopyrachlor-methyl, aminocyclopyrachlor-potassium, aminopyralid, aminopyralid-potassium, aminopyralid-tris(2-hydroxypropyl)ammonium, amiprofos-methyl, amiprophos, amisulbrom, amiton, amiton oxalate, amitraz, amitrole, ammonium sulfamate, ammonium α-naphthaleneacetate, amobam, ampropylfos, anabasine, ancymidol, anilazine, anilofos, anisuron, anthraquinone, antu, apholate, aramite, arsenous oxide, asomate, aspirin, asulam, asulam-potassium, asulam-sodium, athidathion, atraton, atrazine, aureofungin, aviglycine, aviglycine hydrochloride, azaconazole, azadirachtin, azafenidin, azamethiphos, azimsulfuron, azinphos-ethyl, azinphos-methyl, aziprotryne, azithiram, azobenzene, azocyclotin, azothoate, azoxystrobin, bachmedesh, barban, barium hexafluorosilicate, barium polysulfide, barthrin, BCPC,

beflubutamid, benalaxyl, benalaxyl-M, benazolin, benazolin-dimethylammonium, benazolin-ethyl, benazolin-potassium, bencarbazone, benclothiaz, bendiocarb, benfluralin, benfuracarb, benfuresate, benodanil, benomyl, benoxacor, benoxafos, benquinox, bensulfuron, bensulfuron-methyl, bensulide, bensultap, bentaluron, bentazone, bentazone-sodium, benthiavalicarb, benthiavalicarb-isopropyl, benthiazole, bentranil, benzadox, benzadox-ammonium, benzalkonium chloride, benzamacril, benzamacril-isobutyl, benzamorf, benzfendizone, benzipram, benzobicyclon, benzofenap, benzofluor, benzohydroxamic acid, benzoximate, benzoylprop, benzoylprop-ethyl, benzthiazuron, benzyl benzoate, benzyladenine, berberine, berberine chloride, *beta*-cyfluthrin, *beta*-cypermethrin, bethoxazin, bicyclopyrone, bifenazate, bifenox, bifenthrin, bifujunzhi, bilanafos, bilanafos-sodium, binapacryl, bingqingxiao, bioallethrin, bioethanomethrin, biopermethrin, bioresmethrin, biphenyl, bisazir, bismerthiazol, bispyribac, bispyribac-sodium, bistrifluron, bitertanol, bithionol, bixafen, blasticidin-S, borax, Bordeaux mixture, boric acid, boscalid, brassinolide, brassinolide-ethyl, brevicomin, brodifacoum, brofenvalerate, brofluthrinate, bromacil, bromacil-lithium, bromacil-sodium, bromadiolone, bromethalin, bromethrin, bromfenvinfos, bromoacetamide, bromobonil, bromobutide, bromocyclen, bromo-DDT, bromofenoxim, bromophos, bromophos-ethyl, bromopropylate, bromothalonil, bromoxynil, bromoxynil butyrate, bromoxynil heptanoate, bromoxynil octanoate, bromoxynil-potassium, brompyrazon, bromuconazole, bronopol, bucarpolate, bufencarb, buminafos, bupirimate, buprofezin, Burgundy mixture, busulfan, butacarb, butachlor, butafenacil, butamifos, butathiofos, butenachlor, butethrin, buthidazole, buthiobate, buthiuron, butocarboxim, butonate, butopyronoxyl, butoxycarboxim, butralin, butroxydim, buturon, butylamine, butylate, cacodylic acid, cadusafos, cafenstrole, calcium arsenate, calcium chlorate, calcium cyanamide, calcium polysulfide, calvinphos, cambendichlor, camphechlor, camphor, captafol, captan, carbamorph, carbanolate, carbaryl, carbasulam, carbendazim, carbendazim benzenesulfonate, carbendazim sulfite, carbetamide, carbofuran, carbon disulfide, carbon tetrachloride, carbophenothion, carbosulfan, carboxazole, carboxide, carboxin, carfentrazone, carfentrazone-ethyl, carpropamid, cartap, cartap hydrochloride, carvacrol, carvone, CDEA, cellocidin, CEPC, ceralure, Cheshunt mixture, chinomethionat, chitosan, chlobenthiazone, chlomethoxyfen, chloralose, chloramben, chloramben-ammonium, chloramben-diolamine, chloramben-methyl, chloramben-methylammonium, chloramben-sodium, chloramine phosphorus, chloramphenicol, chloraniformethan, chloranil, chloranocryl, chlorantraniliprole, chlorazifop, chlorazifop-propargyl, chlorazine, chlorbenside, chlorbenzuron, chlorbicyclen, chlorbromuron, chlorbufam, chlordane, chlordecone, chlordimeform, chlordimeform

hydrochloride, chlorempenthrin, chlorethoxyfos, chloreturon, chlorfenac, chlorfenac-ammonium, chlorfenac-sodium, chlorfenapyr, chlorfenazole, chlorfenethol, chlorfenprop, chlorfenson, chlorfensulphide, chlorfenvinphos, chlorfluazuron, chlorflurazole, chlorfluren, chlorfluren-methyl, chlorflurenol, chlorflurenol-methyl, chloridazon, chlorimuron, chlorimuron-ethyl, chlormephos, chlormequat, chlormequat chloride, chlornidine, chlornitrofen, chlorobenzilate, chlorodinitronaphthalenes, chloroform, chloromebuform, chloromethiuron, chloroneb, chlorophacinone, chlorophacinone-sodium, chloropicrin, chloropon, chloropropylate, chlorothalonil, chlorotoluron, chloroxuron, chloroxynil, chlorphonium, chlorphonium chloride, chlorphoxim, chlorprazophos, chlorprocarb, chlorpropham, chlorpyrifos, chlorpyrifos-methyl, chlorquinox, chlorsulfuron, chlorthal, chlorthal-dimethyl, chlorthal-monomethyl, chlorthiamid, chlorthiophos, chlozolinate, choline chloride, chromafenozide, cinerin I, cinerin II, cinerins, cinidon-ethyl, cinmethylin, cinosulfuron, ciobutide, cisanilide, cismethrin, clethodim, climbazole, cliodinate, clodinafop, clodinafop-propargyl, cloethocarb, clofencet, clofencet-potassium, clofentezine, clofibric acid, clofop, clofop-isobutyl, clomazone, clomeprop, cloprop, cloproxydim, clopyralid, clopyralid-methyl, clopyralid-olamine, clopyralid-potassium, clopyralid-tris(2-hydroxypropyl)ammonium, cloquintocet, cloquintocet-mexyl, cloransulam, cloransulam-methyl, closantel, clothianidin, clotrimazole, cloxyfonac, cloxyfonac-sodium, CMA, codlelure, colophonate, copper acetate, copper acetoarsenite, copper arsenate, copper carbonate, basic, copper hydroxide, copper naphthenate, copper oleate, copper oxychloride, copper silicate, copper sulfate, copper zinc chromate, coumachlor, coumafuryl, coumaphos, coumatetralyl, coumithoate, coumoxystrobin, CPMC, CPMF, CPPC, credazine, cresol, crimidine, crotamiton, crotoxyphos, crufomate, cryolite, cue-lure, cufraneb, cumyluron, cuprobam, cuprous oxide, curcumenol, cyanamide, cyanatryn, cyanazine, cyanofenphos, cyanophos, cyanthoate, cyantraniliprole, cyazofamid, cybutryne, cyclafuramid, cyclanilide, cyclethrin, cycloate, cycloheximide, cycloprate, cycloprothrin, cyclosulfamuron, cycloxaprid, cycloxydim, cycluron, cyenopyrafen, cyflufenamid, cyflumetofen, cyfluthrin, cyhalofop, cyhalofop-butyl, cyhalothrin, cyhexatin, cymiazole, cymiazole hydrochloride, cymoxanil, cyometrinil, cypendazole, cypermethrin, cyperquat, cyperquat chloride, cyphenothrin, cyprazine, cyprazole, cyproconazole, cyprodinil, cyprofuram, cypromid, cyprosulfamide, cyromazine, cythioate, daimuron, dalapon, dalapon-calcium, dalapon-magnesium, dalapon-sodium, daminozide, dayoutong, dazomet, dazomet-sodium, DBCP, *d*-camphor, DCIP, DCPTA, DDT, debacarb, decafentin, decarbofuran, dehydroacetic acid, delachlor, deltamethrin, demephion, demephion-O, demephion-S, demeton, demeton-methyl, demeton-

O, demeton-O-methyl, demeton-S, demeton-S-methyl, demeton-S-methylsulphon, desmedipham, desmetryn, *d*-fanshiluquebingjuzhi, diafenthiuron, dialifos, di-allate, diamidafos, diatomaceous earth, diazinon, dibutyl phthalate, dibutyl succinate, dicamba, dicamba-diglycolamine, dicamba-dimethylammonium, dicamba-diolamine, dicamba-isopropylammonium, dicamba-methyl, dicamba-olamine, dicamba-potassium, dicamba-sodium, dicamba-trolamine, dicapthon, dichlobenil, dichlofenthion, dichlofluanid, dichlone, dichloralurea, dichlorbenzuron, dichlorflurenol, dichlorflurenol-methyl, dichlormate, dichlormid, dichlorophen, dichlorprop, dichlorprop-2-ethylhexyl, dichlorprop-butotyl, dichlorprop-dimethylammonium, dichlorprop-ethylammonium, dichlorprop-isoctyl, dichlorprop-methyl, dichlorprop-P, dichlorprop-P-2-ethylhexyl, dichlorprop-P-dimethylammonium, dichlorprop-potassium, dichlorprop-sodium, dichlorvos, dichlozoline, diclobutrazol, diclocymet, diclofop, diclofop-methyl, diclomezine, diclomezine-sodium, dicloran, diclosulam, dicofol, dicoumarol, dicresyl, dicrotophos, dicyclanil, dicyclonon, dieldrin, dienochlor, diethamquat, diethamquat dichloride, diethatyl, diethatyl-ethyl, diethofencarb, dietholate, diethyl pyrocarbonate, diethyltoluamide, difenacoum, difenoconazole, difenopenten, difenopenten-ethyl, difenoxuron, difenzoquat, difenzoquat metilsulfate, difethialone, diflovidazin, diflubenzuron, diflufenican, diflufenzopyr, diflufenzopyr-sodium, diflumetorim, dikegulac, dikegulac-sodium, dilor, dimatif, dimefluthrin, dimefox, dimefuron, dimepiperate, dimetachlone, dimetan, dimethacarb, dimethachlor, dimethametryn, dimethenamid, dimethenamid-P, dimethipin, dimethirimol, dimethoate, dimethomorph, dimethrin, dimethyl carbate, dimethyl phthalate, dimethylvinphos, dimetilan, dimexano, dimidazon, dimoxystrobin, dinex, dinex-diclexine, dingjunezuo, diniconazole, diniconazole-M, dinitramine, dinobuton, dinocap, dinocap-4, dinocap-6, dinocton, dinofenate, dinopenton, dinoprop, dinosam, dinoseb, dinoseb acetate, dinoseb-ammonium, dinoseb-diolamine, dinoseb-sodium, dinoseb-trolamine, dinosulfon, dinotefuran, dinoterb, dinoterb acetate, dinoterbon, diofenolan, dioxabenzofos, dioxacarb, dioxathion, diphacinone, diphacinone-sodium, diphenamid, diphenyl sulfone, diphenylamine, dipropalin, dipropetryn, dipyrithione, diquat, diquat dibromide, disparlure, disul, disulfiram, disulfoton, disul-sodium, ditalimfos, dithianon, dithicrofos, dithioether, dithiopyr, diuron, d-limonene, DMPA, DNOC, DNOC-ammonium, DNOC-potassium, DNOC-sodium, dodemorph, dodemorph acetate, dodemorph benzoate, dodicin, dodicin hydrochloride, dodicin-sodium, dodine, dofenapyn, dominicalure, doramectin, drazoxolon, DSMA, dufulin, EBEP, EBP, ecdysterone, edifenphos, eglinazine, eglinazine-ethyl, emamectin, emamectin benzoate, EMPC, empenthrin, endosulfan, endothal, endothal-diammonium, endothal-

dipotassium, endothal-disodium, endothion, endrin, enestroburin, EPN, epocholeone, epofenonane, epoxiconazole, eprinomectin, epronaz, EPTC, erbon, ergocalciferol, erlujixiancaoan, esdépalléthrine, esfenvalerate, esprocarb, etacelasil, etaconazole, etaphos, etem, ethaboxam, ethachlor, ethalfluralin, ethametsulfuron, ethametsulfuron-methyl, ethaprochlor, ethephon, ethidimuron, ethiofencarb, ethiolate, ethion, ethiozin, ethiprole, ethirimol, ethoate-methyl, ethofumesate, ethohexadiol, ethoprophos, ethoxyfen, ethoxyfen-ethyl, ethoxyquin, ethoxysulfuron, ethychlozate, ethyl formate, ethyl α-naphthaleneacetate, ethyl-DDD, ethylene, ethylene dibromide, ethylene dichloride, ethylene oxide, ethylicin, ethylmercury 2,3-dihydroxypropyl mercaptide, ethylmercury acetate, ethylmercury bromide, ethylmercury chloride, ethylmercury phosphate, etinofen, etnipromid, etobenzanid, etofenprox, etoxazole, etridiazole, etrimfos, eugenol, EXD, famoxadone, famphur, fenamidone, fenaminosulf, fenamiphos, fenapanil, fenarimol, fenasulam, fenazaflor, fenazaquin, fenbuconazole, fenbutatin oxide, fenchlorazole, fenchlorazole-ethyl, fenchlorphos, fenclorim, fenethacarb, fenfluthrin, fenfuram, fenhexamid, fenitropan, fenitrothion, fenjuntong, fenobucarb, fenoprop, fenoprop-3-butoxypropyl, fenoprop-butometyl, fenoprop-butotyl, fenoprop-butyl, fenoprop-isoctyl, fenoprop-methyl, fenoprop-potassium, fenothiocarb, fenoxacrim, fenoxanil, fenoxaprop, fenoxaprop-ethyl, fenoxaprop-P, fenoxaprop-P-ethyl, fenoxasulfone, fenoxycarb, fenpiclonil, fenpirithrin, fenpropathrin, fenpropidin, fenpropimorph, fenpyrazamine, fenpyroximate, fenridazon, fenridazon-potassium, fenridazon-propyl, fenson, fensulfothion, fenteracol, fenthiaprop, fenthiaprop-ethyl, fenthion, fenthion-ethyl, fentin, fentin acetate, fentin chloride, fentin hydroxide, fentrazamide, fentrifanil, fenuron, fenuron TCA, fenvalerate, ferbam, ferimzone, ferrous sulfate, fipronil, flamprop, flamprop-isopropyl, flamprop-M, flamprop-methyl, flamprop-M-isopropyl, flamprop-M-methyl, flazasulfuron, flocoumafen, flometoquin, flonicamid, florasulam, fluacrypyrim, fluazifop, fluazifop-butyl, fluazifop-methyl, fluazifop-P, fluazifop-P-butyl, fluazinam, fluazolate, fluazuron, flubendiamide, flubenzimine, flucarbazone, flucarbazone-sodium, flucetosulfuron, fluchloralin, flucofuron, flucycloxuron, flucythrinate, fludioxonil, fluenetil, fluensulfone, flufenacet, flufenerim, flufenican, flufenoxuron, flufenprox, flufenpyr, flufenpyr-ethyl, flufiprole, flumethrin, flumetover, flumetralin, flumetsulam, flumezin, flumiclorac, flumiclorac-pentyl, flumioxazin, flumipropyn, flumorph, fluometuron, fluopicolide, fluopyram, fluorbenside, fluoridamid, fluoroacetamide, fluorodifen, fluoroglycofen, fluoroglycofen-ethyl, fluoroimide, fluoromidine, fluoronitrofen, fluothiuron, fluotrimazole, fluoxastrobin, flupoxam, flupropacil, flupropadine, flupropanate, flupropanate-sodium, flupyradifurone, flupyrsulfuron, flupyrsulfuron-methyl, flupyrsulfuron-

methyl-sodium, fluquinconazole, flurazole, flurenol, flurenol-butyl, flurenol-methyl, fluridone, flurochloridone, fluroxypyr, fluroxypyr-butometyl, fluroxypyr-meptyl, flurprimidol, flursulamid, flurtamone, flusilazole, flusulfamide, fluthiacet, fluthiacet-methyl, flutianil, flutolanil, flutriafol, fluvalinate, fluxapyroxad, fluxofenim, folpet, fomesafen, fomesafen-sodium, fonofos, foramsulfuron, forchlorfenuron, formaldehyde, formetanate, formetanate hydrochloride, formothion, formparanate, formparanate hydrochloride, fosamine, fosamine-ammonium, fosetyl, fosetyl-aluminium, fosmethilan, fospirate, fosthiazate, fosthietan, frontalin, fuberidazole, fucaojing, fucaomi, funaihecaoling, fuphenthiourea, furalane, furalaxyl, furamethrin, furametpyr, furathiocarb, furcarbanil, furconazole, furconazole-cis, furethrin, furfural, furilazole, furmecyclox, furophanate, furyloxyfen, *gamma*-cyhalothrin, *gamma*-HCH, genit, gibberellic acid, gibberellins, gliftor, glufosinate, glufosinate-ammonium, glufosinate-P, glufosinate-P-ammonium, glufosinate-P-sodium, glyodin, glyoxime, glyphosate, glyphosate-diammonium, glyphosate-dimethylammonium, glyphosate-isopropylammonium, glyphosate-monoammonium, glyphosate-potassium, glyphosate-sesquisodium, glyphosate-trimesium, glyphosine, gossyplure, grandlure, griseofulvin, guazatine, guazatine acetates, halacrinate, halfenprox, halofenozide, halosafen, halosulfuron, halosulfuron-methyl, haloxydine, haloxyfop, haloxyfop-etotyl, haloxyfop-methyl, haloxyfop-P, haloxyfop-P-etotyl, haloxyfop-P-methyl, haloxyfop-sodium, HCH, hemel, hempa, HEOD, heptachlor, heptenophos, heptopargil, heterophos, hexachloroacetone, hexachlorobenzene, hexachlorobutadiene, hexachlorophene, hexaconazole, hexaflumuron, hexaflurate, hexalure, hexamide, hexazinone, hexylthiofos, hexythiazox, HHDN, holosulf, huancaiwo, huangcaoling, huanjunzuo, hydramethylnon, hydrargaphen, hydrated lime, hydrogen cyanide, hydroprene, hymexazol, hyquincarb, IAA, IBA, icaridin, imazalil, imazalil nitrate, imazalil sulfate, imazamethabenz, imazamethabenz-methyl, imazamox, imazamox-ammonium, imazapic, imazapic-ammonium, imazapyr, imazapyr-isopropylammonium, imazaquin, imazaquin-ammonium, imazaquin-methyl, imazaquin-sodium, imazethapyr, imazethapyr-ammonium, imazosulfuron, imibenconazole, imicyafos, imidacloprid, imidaclothiz, iminoctadine, iminoctadine triacetate, iminoctadine trialbesilate, imiprothrin, inabenfide, indanofan, indaziflam, indoxacarb, inezin, iodobonil, iodocarb, iodomethane, iodosulfuron, iodosulfuron-methyl, iodosulfuron-methyl-sodium, iofensulfuron, iofensulfuron-sodium, ioxynil, ioxynil octanoate, ioxynil-lithium, ioxynil-sodium, ipazine, ipconazole, ipfencarbazone, iprobenfos, iprodione, iprovalicarb, iprymidam, ipsdienol, ipsenol, IPSP, isamidofos, isazofos, isobenzan, isocarbamid, isocarbophos, isocil, isodrin, isofenphos, isofenphos-methyl, isolan, isomethiozin, isonoruron, isopolinate, isoprocarb,

isopropalin, isoprothiolane, isoproturon, isopyrazam, isopyrimol, isothioate, isotianil, isouron, isovaledione, isoxaben, isoxachlortole, isoxadifen, isoxadifen-ethyl, isoxaflutole, isoxapyrifop, isoxathion, ivermectin, izopamfos, japonilure, japothrins, jasmolin I, jasmolin II, jasmonic acid, jiahuangchongzong, jiajizengxiaolin, jiaxiangjunzhi, jiecaowan, jiecaoxi, jodfenphos, juvenile hormone I, juvenile hormone II, juvenile hormone III, kadethrin, karbutilate, karetazan, karetazan-potassium, kasugamycin, kasugamycin hydrochloride, kejunlin, kelevan, ketospiradox, ketospiradox-potassium, kinetin, kinoprene, kresoxim-methyl, kuicaoxi, lactofen, lambda-cyhalothrin, latilure, lead arsenate, lenacil, lepimectin, leptophos, lindane, lineatin, linuron, lirimfos, litlure, looplure, lufenuron, lvdingjunzhi, lvxiancaolin, lythidathion, MAA, malathion, maleic hydrazide, malonoben, maltodextrin, MAMA, mancopper, mancozeb, mandipropamid, maneb, matrine, mazidox, MCPA, MCPA-2-ethylhexyl, MCPA-butotyl, MCPA-butyl, MCPA-dimethylammonium, MCPA-diolamine, MCPA-ethyl, MCPA-isobutyl, MCPA-isoctyl, MCPA-isopropyl, MCPA-methyl, MCPA-olamine, MCPA-potassium, MCPA-sodium, MCPA-thioethyl, MCPA-trolamine, MCPB, MCPB-ethyl, MCPB-methyl, MCPB-sodium, mebenil, mecarbam, mecarbinzid, mecarphon, mecoprop, mecoprop-2-ethylhexyl, mecoprop-dimethylammonium, mecoprop-diolamine, mecoprop-ethadyl, mecoprop-isoctyl, mecoprop-methyl, mecoprop-P, mecoprop-P-2-ethylhexyl, mecoprop-P-dimethylammonium, mecoprop-P-isobutyl, mecoprop-potassium, mecoprop-P-potassium, mecoprop-sodium, mecoprop-trolamine, medimeform, medinoterb, medinoterb acetate, medlure, mefenacet, mefenpyr, mefenpyr-diethyl, mefluidide, mefluidide-diolamine, mefluidide-potassium, megatomoic acid, menazon, mepanipyrim, meperfluthrin, mephenate, mephosfolan, mepiquat, mepiquat chloride, mepiquat pentaborate, mepronil, meptyldinocap, mercuric chloride, mercuric oxide, mercurous chloride, merphos, mesoprazine, mesosulfuron, mesosulfuron-methyl, mesotrione, mesulfen, mesulfenfos, metaflumizone, metalaxyl, metalaxyl-M, metaldehyde, metam, metam-ammonium, metamifop, metamitron, metam-potassium, metam-sodium, metazachlor, metazosulfuron, metazoxolon, metconazole, metepa, metflurazon, methabenzthiazuron, methacrifos, methalpropalin, methamidophos, methasulfocarb, methazole, methfuroxam, methidathion, methiobencarb, methiocarb, methiopyrisulfuron, methiotepa, methiozolin, methiuron, methocrotophos, methometon, methomyl, methoprene, methoprotryne, methoquin-butyl, methothrin, methoxychlor, methoxyfenozide, methoxyphenone, methyl apholate, methyl bromide, methyl eugenol, methyl iodide, methyl isothiocyanate, methylacetophos, methylchloroform, methyldymron, methylene chloride, methylmercury benzoate, methylmercury dicyandiamide, methylmercury pentachlorophenoxide,

methylneodecanamide, metiram, metobenzuron, metobromuron, metofluthrin, metolachlor, metolcarb, metominostrobin, metosulam, metoxadiazone, metoxuron, metrafenone, metribuzin, metsulfovax, metsulfuron, metsulfuron-methyl, mevinphos, mexacarbate, mieshuan, milbemectin, milbemycin oxime, milneb, mipafox, mirex, MNAF, moguchun, molinate, molosultap, monalide, monisouron, monochloroacetic acid, monocrotophos, monolinuron, monosulfuron, monosulfuron-ester, monuron, monuron TCA, morfamquat, morfamquat dichloride, moroxydine, moroxydine hydrochloride, morphothion, morzid, moxidectin, MSMA, muscalure, myclobutanil, myclozolin, N-(ethylmercury)-p-toluenesulphonanilide, nabam, naftalofos, naled, naphthalene, naphthaleneacetamide, naphthalic anhydride, naphthoxyacetic acids, naproanilide, napropamide, naptalam, naptalam-sodium, natamycin, neburon, niclosamide, niclosamide-olamine, nicosulfuron, nicotine, nifluridide, nipyraclofen, nitenpyram, nithiazine, nitralin, nitrapyrin, nitrilacarb, nitrofen, nitrofluorfen, nitrostyrene, nitrothal-isopropyl, norbormide, norflurazon, nornicotine, noruron, novaluron, noviflumuron, nuarimol, OCH, octachlorodipropyl ether, octhilinone, ofurace, omethoate, orbencarb, orfralure, ortho-dichlorobenzene, orthosulfamuron, oryctalure, orysastrobin, oryzalin, osthol, ostramone, oxabetrinil, oxadiargyl, oxadiazon, oxadixyl, oxamate, oxamyl, oxapyrazon, oxapyrazon-dimolamine, oxapyrazon-sodium, oxasulfuron, oxaziclomefone, oxine-copper, oxolinic acid, oxpoconazole, oxpoconazole fumarate, oxycarboxin, oxydemeton-methyl, oxydeprofos, oxydisulfoton, oxyfluorfen, oxymatrine, oxytetracycline, oxytetracycline hydrochloride, paclobutrazol, paichongding, para-dichlorobenzene, parafluron, paraquat, paraquat dichloride, paraquat dimetilsulfate, parathion, parathion-methyl, parinol, pebulate, pefurazoate, pelargonic acid, penconazole, pencycuron, pendimethalin, penflufen, penfluron, penoxsulam, pentachlorophenol, pentanochlor, penthiopyrad, pentmethrin, pentoxazone, perfluidone, permethrin, pethoxamid, phenamacril, phenazine oxide, phenisopham, phenkapton, phenmedipham, phenmedipham-ethyl, phenobenzuron, phenothrin, phenproxide, phenthoate, phenylmercuriurea, phenylmercury acetate, phenylmercury chloride, phenylmercury derivative of pyrocatechol, phenylmercury nitrate, phenylmercury salicylate, phorate, phosacetim, phosalone, phosdiphen, phosfolan, phosfolan-methyl, phosglycin, phosmet, phosnichlor, phosphamidon, phosphine, phosphocarb, phosphorus, phostin, phoxim, phoxim-methyl, phthalide, picloram, picloram-2-ethylhexyl, picloram-isoctyl, picloram-methyl, picloram-olamine, picloram-potassium, picloram-triethylammonium, picloram-tris(2-hydroxypropyl)ammonium, picolinafen, picoxystrobin, pindone, pindone-sodium, pinoxaden, piperalin, piperonyl butoxide, piperonyl cyclonene, piperophos, piproctanyl, piproctanyl

bromide, piprotal, pirimetaphos, pirimicarb, pirimioxyphos, pirimiphos-ethyl, pirimiphos-methyl, plifenate, polycarbamate, polyoxins, polyoxorim, polyoxorim-zinc, polythialan, potassium arsenite, potassium azide, potassium cyanate, potassium gibberellate, potassium naphthenate, potassium polysulfide, potassium thiocyanate, potassium α-naphthaleneacetate, *pp'*-DDT, prallethrin, precocene I, precocene II, precocene III, pretilachlor, primidophos, primisulfuron, primisulfuron-methyl, probenazole, prochloraz, prochloraz-manganese, proclonol, procyazine, procymidone, prodiamine, profenofos, profluazol, profluralin, profluthrin, profoxydim, proglinazine, proglinazine-ethyl, prohexadione, prohexadione-calcium, prohydrojasmon, promacyl, promecarb, prometon, prometryn, promurit, propachlor, propamidine, propamidine dihydrochloride, propamocarb, propamocarb hydrochloride, propanil, propaphos, propaquizafop, propargite, proparthrin, propazine, propetamphos, propham, propiconazole, propineb, propisochlor, propoxur, propoxycarbazone, propoxycarbazone-sodium, propyl isome, propyrisulfuron, propyzamide, proquinazid, prosuler, prosulfalin, prosulfocarb, prosulfuron, prothidathion, prothiocarb, prothiocarb hydrochloride, prothioconazole, prothiofos, prothoate, protrifenbute, proxan, proxan-sodium, prynachlor, pydanon, pymetrozine, pyracarbolid, pyraclofos, pyraclonil, pyraclostrobin, pyraflufen, pyraflufen-ethyl, pyrafluprole, pyramat, pyrametostrobin, pyraoxystrobin, pyrasulfotole, pyrazolynate, pyrazophos, pyrazosulfuron, pyrazosulfuron-ethyl, pyrazothion, pyrazoxyfen, pyresmethrin, pyrethrin I, pyrethrin II, pyrethrins, pyribambenz-isopropyl, pyribambenz-propyl, pyribencarb, pyribenzoxim, pyributicarb, pyriclor, pyridaben, pyridafol, pyridalyl, pyridaphenthion, pyridate, pyridinitril, pyrifenox, pyrifluquinazon, pyriftalid, pyrimethanil, pyrimidifen, pyriminobac, pyriminobac-methyl, pyrimisulfan, pyrimitate, pyrinuron, pyriofenone, pyriprole, pyripropanol, pyriproxyfen, pyrithiobac, pyrithiobac-sodium, pyrolan, pyroquilon, pyroxasulfone, pyroxsulam, pyroxychlor, pyroxyfur, quassia, quinacetol, quinacetol sulfate, quinalphos, quinalphos-methyl, quinazamid, quinclorac, quinconazole, quinmerac, quinoclamine, quinonamid, quinothion, quinoxyfen, quintiofos, quintozene, quizalofop, quizalofop-ethyl, quizalofop-P, quizalofop-P-ethyl, quizalofop-P-tefuryl, quwenzhi, quyingding, rabenzazole, rafoxanide, rebemide, resmethrin, rhodethanil, rhodojaponin-III, ribavirin, rimsulfuron, rotenone, ryania, saflufenacil, saijunmao, saisentong, salicylanilide, sanguinarine, santonin, schradan, scilliroside, sebuthylazine, secbumeton, sedaxane, selamectin, semiamitraz, semiamitraz chloride, sesamex, sesamolin, sethoxydim, shuangjiaancaolin, siduron, siglure, silafluofen, silatrane, silica gel, silthiofam, simazine, simeconazole, simeton, simetryn, sintofen, SMA, S-metolachlor, sodium arsenite, sodium azide, sodium chlorate, sodium fluoride, sodium fluoroacetate, sodium hexafluorosilicate,

sodium naphthenate, sodium orthophenylphenoxide, sodium pentachlorophenoxide, sodium polysulfide, sodium thiocyanate, sodium α-naphthaleneacetate, sophamide, spinetoram, spinosad, spirodiclofen, spiromesifen, spirotetramat, spiroxamine, streptomycin, streptomycin sesquisulfate, strychnine, sulcatol, sulcofuron, sulcofuron-sodium, sulcotrione, sulfallate, sulfentrazone, sulfiram, sulfluramid, sulfometuron, sulfometuron-methyl, sulfosulfuron, sulfotep, sulfoxaflor, sulfoxide, sulfoxime, sulfur, sulfuric acid, sulfuryl fluoride, sulglycapin, sulprofos, sultropen, swep, *tau*-fluvalinate, tavron, tazimcarb, TCA, TCA-ammonium, TCA-calcium, TCA-ethadyl, TCA-magnesium, TCA-sodium, TDE, tebuconazole, tebufenozide, tebufenpyrad, tebufloquin, tebupirimfos, tebutam, tebuthiuron, tecloftalam, tecnazene, tecoram, teflubenzuron, tefluthrin, tefuryltrione, tembotrione, temephos, tepa, TEPP, tepraloxydim, terallethrin, terbacil, terbucarb, terbuchlor, terbufos, terbumeton, terbuthylazine, terbutryn, tetcyclacis, tetrachloroethane, tetrachlorvinphos, tetraconazole, tetradifon, tetrafluron, tetramethrin, tetramethylfluthrin, tetramine, tetranactin, tetrasul, thallium sulfate, thenylchlor, theta-cypermethrin, thiabendazole, thiacloprid, thiadifluor, thiamethoxam, thiapronil, thiazafluron, thiazopyr, thicrofos, thicyofen, thidiazimin, thidiazuron, thiencarbazone, thiencarbazone-methyl, thifensulfuron, thifensulfuron-methyl, thifluzamide, thiobencarb, thiocarboxime, thiochlorfenphim, thiocyclam, thiocyclam hydrochloride, thiocyclam oxalate, thiodiazole-copper, thiodicarb, thiofanox, thiofluoximate, thiohempa, thiomersal, thiometon, thionazin, thiophanate, thiophanate-methyl, thioquinox, thiosemicarbazide, thiosultap, thiosultap-diammonium, thiosultap-disodium, thiosultap-monosodium, thiotepa, thiram, thuringiensin, tiadinil, tiaojiean, tiocarbazil, tioclorim, tioxymid, tirpate, tolclofos-methyl, tolfenpyrad, tolylfluanid, tolylmercury acetate, topramezone, tralkoxydim, tralocythrin, tralomethrin, tralopyril, transfluthrin, transpermethrin, tretamine, triacontanol, triadimefon, triadimenol, triafamone, tri-allate, triamiphos, triapenthenol, triarathene, triarimol, triasulfuron, triazamate, triazbutil, triaziflam, triazophos, triazoxide, tribenuron, tribenuron-methyl, tribufos, tributyltin oxide, tricamba, trichlamide, trichlorfon, trichlormetaphos-3, trichloronat, triclopyr, triclopyr-butotyl, triclopyr-ethyl, triclopyr-triethylammonium, tricyclazole, tridemorph, tridiphane, trietazine, trifenmorph, trifenofos, trifloxystrobin, trifloxysulfuron, trifloxysulfuron-sodium, triflumizole, triflumuron, trifluralin, triflusulfuron, triflusulfuron-methyl, trifop, trifop-methyl, trifopsime, triforine, trihydroxytriazine, trimedlure, trimethacarb, trimeturon, trinexapac, trinexapac-ethyl, triprene, tripropindan, triptolide, tritac, triticonazole, tritosulfuron, trunc-call, uniconazole, uniconazole-P, urbacide, uredepa, valerate, validamycin, valifenalate, valone, vamidothion, vangard, vaniliprole, vernolate, vinclozolin,

warfarin, warfarin-potassium, warfarin-sodium, xiaochongliulin, xinjunan, xiwojunan, XMC, xylachlor, xylenols, xylylcarb, yishijing, zarilamid, zeatin, zengxiaoan, zeta-cypermethrin, zinc naphthenate, zinc phosphide, zinc thiazole, zineb, ziram, zolaprofos, zoxamide, zuomihuanglong, α-chlorohydrin, α-ecdysone, α-multistriatin, and α-naphthaleneacetic acid. For more information consult the "**COMPENDIUM OF PESTICIDE COMMON NAMES**" located at http://www.alanwood.net/pesticides/index.html. Also consult "**THE PESTICIDE MANUAL**" 14th Edition, edited by C D S Tomlin, copyright 2006 by British Crop Production Council, or its prior or more recent editions.

BIOPESTICIDES

Molecules of Formula One may also be used in combination (such as in a compositional mixture, or a simultaneous or sequential application) with one or more biopesticides. The term "biopesticide" is used for microbial biological pest control agents that are applied in a similar manner to chemical pesticides. Commonly these are bacterial, but there are also examples of fungal control agents, including *Trichoderma* spp. and *Ampelomyces quisqualis* (a control agent for grape powdery mildew). *Bacillus subtilis* are used to control plant pathogens. Weeds and rodents have also been controlled with microbial agents. One well-known insecticide example is *Bacillus thuringiensis*, a bacterial disease of Lepidoptera, Coleoptera, and Diptera. Because it has little effect on other organisms, it is considered more environmentally friendly than synthetic pesticides. Biological insecticides include products based on:

1. entomopathogenic fungi (*e.g. Metarhizium anisopliae*);
2. entomopathogenic nematodes (*e.g. Steinernema feltiae*); and
3. entomopathogenic viruses (*e.g. Cydia pomonella* granulovirus).

Other examples of entomopathogenic organisms include, but are not limited to, baculoviruses, bacteria and other prokaryotic organisms, fungi, protozoa and Microsproridia. Biologically derived insecticides include, but not limited to, rotenone, veratridine, as well as microbial toxins; insect tolerant or resistant plant varieties; and organisms modified by recombinant DNA technology to either produce insecticides or to convey an insect resistant property to the genetically modified organism. In one embodiment, the molecules of Formula One may be used with one or more biopesticides in the area of seed treatments and soil amendments. *The Manual of Biocontrol Agents* gives a review of the available biological insecticide (and other biology-based control) products. Copping L.G. (ed.) (2004). *The Manual of Biocontrol Agents* (formerly the *Biopesticide Manual*) 3rd Edition. British Crop Production Council (BCPC), Farnham, Surrey UK.

OTHER ACTIVE COMPOUNDS

Molecules of Formula One may also be used in combination (such as in a compositional mixture, or a simultaneous or sequential application) with one or more of the following:

1. 3-(4-chloro-2,6-dimethylphenyl)-4-hydroxy-8-oxa-1-azaspiro[4,5]dec-3-en-2-one;
2. 3-(4'-chloro-2,4-dimethyl[1,1'-biphenyl]-3-yl)-4-hydroxy-8-oxa-1-azaspiro[4,5]dec-3-en-2-one;
3. 4-[[(6-chloro-3-pyridinyl)methyl]methylamino]-2(5*H*)-furanone;
4. 4-[[(6-chloro-3-pyridinyl)methyl]cyclopropylamino]-2(5*H*)-furanone;
5. 3-chloro-*N*2-[(1*S*)-1-methyl-2-(methylsulfonyl)ethyl]-*N*1-[2-methyl-4-[1,2,2,2-tetrafluoro-1-(trifluoromethyl)ethyl]phenyl]-1,2-benzenedicarboxamide;
6. 2-cyano-*N*-ethyl-4-fluoro-3-methoxy-benenesulfonamide;
7. 2-cyano-*N*-ethyl-3-methoxy-benzenesulfonamide;
8. 2-cyano-3-difluoromethoxy-*N*-ethyl-4-fluoro-benzenesulfonamide;
9. 2-cyano-3-fluoromethoxy-*N*-ethyl-benzenesulfonamide;
10. 2-cyano-6-fluoro-3-methoxy-*N,N*-dimethyl-benzenesulfonamide;
11. 2-cyano-*N*-ethyl-6-fluoro-3-methoxy-*N*-methyl-benzenesulfonamide;
12. 2-cyano-3-difluoromethoxy-*N,N*-dimethylbenzenesulfon-amide;
13. 3-(difluoromethyl)-*N*-[2-(3,3-dimethylbutyl)phenyl]-1-methyl-1*H*-pyrazole-4-carboxamide;
14. *N*-ethyl-2,2-dimethylpropionamide-2-(2,6-dichloro-α,α,α-trifluoro-*p*-tolyl) hydrazone;
15. *N*-ethyl-2,2-dichloro-1-methylcyclopropane-carboxamide-2-(2,6-dichloro-α,α,α-trifluoro-*p*-tolyl) hydrazone nicotine;
16. O-{(E-)-[2-(4-chloro-phenyl)-2-cyano-1-(2-trifluoromethylphenyl)-vinyl]} S-methyl thiocarbonate;
17. (E)-N1-[(2-chloro-1,3-thiazol-5-ylmethyl)]-N2-cyano-N1-methylacetamidine;
18. 1-(6-chloropyridin-3-ylmethyl)-7-methyl-8-nitro-1,2,3,5,6,7-hexahydro-imidazo[1,2-a]pyridin-5-ol;
19. 4-[4-chlorophenyl-(2-butylidine-hydrazono)methyl)]phenyl mesylate; and
20. N-Ethyl-2,2-dichloro-1-methylcyclopropanecarboxamide-2-(2,6-dichloro-*alpha,alpha,alpha*-trifluoro-p-tolyl)hydrazone.

SYNERGISTIC MIXTURES

Molecules of Formula One may be used with certain active compounds to form synergistic mixtures where the mode of action of such compounds compared to the mode of

action of the molecules of Formula One are the same, similar, or different. Examples of modes of action include, but are not limited to: acetylcholinesterase inhibitor; sodium channel modulator; chitin biosynthesis inhibitor; GABA and glutamate-gated chloride channel antagonist; GABA and glutamate-gated chloride channel agonist; acetylcholine receptor agonist; acetylcholine receptor antagonist; MET I inhibitor; Mg-stimulated ATPase inhibitor; nicotinic acetylcholine receptor; Midgut membrane disrupter; oxidative phosphorylation disrupter, and ryanodine receptor (RyRs). Generally, weight ratios of the molecules of Formula One in a synergistic mixture with another compound are from about 10:1 to about 1:10, in another embodiment from about 5:1 to about 1:5, and in another embodiment from about 3:1, and in another embodiment about 1:1.

FORMULATIONS

A pesticide is rarely suitable for application in its pure form. It is usually necessary to add other substances so that the pesticide can be used at the required concentration and in an appropriate form, permitting ease of application, handling, transportation, storage, and maximum pesticide activity. Thus, pesticides are formulated into, for example, baits, concentrated emulsions, dusts, emulsifiable concentrates, fumigants, gels, granules, microencapsulations, seed treatments, suspension concentrates, suspoemulsions, tablets, water soluble liquids, water dispersible granules or dry flowables, wettable powders, and ultra-low volume solutions. For further information on formulation types see "Catalogue of Pesticide Formulation Types and International Coding System" Technical Monograph n°2, 5th Edition by CropLife International (2002).

Pesticides are applied most often as aqueous suspensions or emulsions prepared from concentrated formulations of such pesticides. Such water-soluble, water-suspendable, or emulsifiable formulations are either solids, usually known as wettable powders, or water dispersible granules, or liquids usually known as emulsifiable concentrates, or aqueous suspensions. Wettable powders, which may be compacted to form water dispersible granules, comprise an intimate mixture of the pesticide, a carrier, and surfactants. The concentration of the pesticide is usually from about 10% to about 90% by weight. The carrier is usually selected from among the attapulgite clays, the montmorillonite clays, the diatomaceous earths, or the purified silicates. Effective surfactants, comprising from about 0.5% to about 10% of the wettable powder, are found among sulfonated lignins, condensed naphthalenesulfonates, naphthalenesulfonates, alkylbenzenesulfonates, alkyl sulfates, and non-ionic surfactants such as ethylene oxide adducts of alkyl phenols.

Emulsifiable concentrates of pesticides comprise a convenient concentration of a pesticide, such as from about 50 to about 500 grams per liter of liquid dissolved in a carrier that is either a water miscible solvent or a mixture of water-immiscible organic solvent and emulsifiers. Useful organic solvents include aromatics, especially xylenes and petroleum fractions, especially the high-boiling naphthalenic and olefinic portions of petroleum such as heavy aromatic naphtha. Other organic solvents may also be used, such as the terpenic solvents including rosin derivatives, aliphatic ketones such as cyclohexanone, and complex alcohols such as 2-ethoxyethanol. Suitable emulsifiers for emulsifiable concentrates are selected from conventional anionic and non-ionic surfactants.

Aqueous suspensions comprise suspensions of water-insoluble pesticides dispersed in an aqueous carrier at a concentration in the range from about 5% to about 50% by weight. Suspensions are prepared by finely grinding the pesticide and vigorously mixing it into a carrier comprised of water and surfactants. Ingredients, such as inorganic salts and synthetic or natural gums may also be added, to increase the density and viscosity of the aqueous carrier. It is often most effective to grind and mix the pesticide at the same time by preparing the aqueous mixture and homogenizing it in an implement such as a sand mill, ball mill, or piston-type homogenizer.

Pesticides may also be applied as granular compositions that are particularly useful for applications to the soil. Granular compositions usually contain from about 0.5% to about 10% by weight of the pesticide, dispersed in a carrier that comprises clay or a similar substance. Such compositions are usually prepared by dissolving the pesticide in a suitable solvent and applying it to a granular carrier which has been pre-formed to the appropriate particle size, in the range of from about 0.5 to about 3 mm. Such compositions may also be formulated by making a dough or paste of the carrier and compound and crushing and drying to obtain the desired granular particle size.

Dusts containing a pesticide are prepared by intimately mixing the pesticide in powdered form with a suitable dusty agricultural carrier, such as kaolin clay, ground volcanic rock, and the like. Dusts can suitably contain from about 1% to about 10% of the pesticide. They can be applied as a seed dressing or as a foliage application with a dust blower machine.

It is equally practical to apply a pesticide in the form of a solution in an appropriate organic solvent, usually petroleum oil, such as the spray oils, which are widely used in agricultural chemistry.

Pesticides can also be applied in the form of an aerosol composition. In such compositions the pesticide is dissolved or dispersed in a carrier, which is a pressure-

generating propellant mixture. The aerosol composition is packaged in a container from which the mixture is dispensed through an atomizing valve.

Pesticide baits are formed when the pesticide is mixed with food or an attractant or both. When the pests eat the bait they also consume the pesticide. Baits may take the form of granules, gels, flowable powders, liquids, or solids. They can be used in pest harborages.

Fumigants are pesticides that have a relatively high vapor pressure and hence can exist as a gas in sufficient concentrations to kill pests in soil or enclosed spaces. The toxicity of the fumigant is proportional to its concentration and the exposure time. They are characterized by a good capacity for diffusion and act by penetrating the pest's respiratory system or being absorbed through the pest's cuticle. Fumigants are applied to control stored product pests under gas proof sheets, in gas sealed rooms or buildings or in special chambers.

Pesticides can be microencapsulated by suspending the pesticide particles or droplets in plastic polymers of various types. By altering the chemistry of the polymer or by changing factors in the processing, microcapsules can be formed of various sizes, solubility, wall thicknesses, and degrees of penetrability. These factors govern the speed with which the active ingredient within is released, which in turn, affects the residual performance, speed of action, and odor of the product.

Oil solution concentrates are made by dissolving pesticide in a solvent that will hold the pesticide in solution. Oil solutions of a pesticide usually provide faster knockdown and kill of pests than other formulations due to the solvents themselves having pesticidal action and the dissolution of the waxy covering of the integument increasing the speed of uptake of the pesticide. Other advantages of oil solutions include better storage stability, better penetration of crevices, and better adhesion to greasy surfaces.

Another embodiment is an oil-in-water emulsion, wherein the emulsion comprises oily globules which are each provided with a lamellar liquid crystal coating and are dispersed in an aqueous phase, wherein each oily globule comprises at least one compound which is agriculturally active, and is individually coated with a monolamellar or oligolamellar layer comprising: (1) at least one non-ionic lipophilic surface-active agent, (2) at least one non-ionic hydrophilic surface-active agent and (3) at least one ionic surface-active agent, wherein the globules having a mean particle diameter of less than 800 nanometers. Further information on the embodiment is disclosed in U.S. patent publication 20070027034 published February 1, 2007, having Patent Application serial number 11/495,228. For ease of use, this embodiment will be referred to as "OIWE".

For further information consult "Insect Pest Management" 2nd Edition by D. Dent, copyright CAB International (2000). Additionally, for more detailed information consult "Handbook of Pest Control – The Behavior, Life History, and Control of Household Pests" by Arnold Mallis, 9th Edition, copyright 2004 by GIE Media Inc.

OTHER FORMULATION COMPONENTS

Generally, when the molecules disclosed in Formula One are used in a formulation, such formulation can also contain other components. These components include, but are not limited to, (this is a non-exhaustive and non-mutually exclusive list) wetters, spreaders, stickers, penetrants, buffers, sequestering agents, drift reduction agents, compatibility agents, anti-foam agents, cleaning agents, and emulsifiers. A few components are described forthwith.

A wetting agent is a substance that when added to a liquid increases the spreading or penetration power of the liquid by reducing the interfacial tension between the liquid and the surface on which it is spreading. Wetting agents are used for two main functions in agrochemical formulations: during processing and manufacture to increase the rate of wetting of powders in water to make concentrates for soluble liquids or suspension concentrates; and during mixing of a product with water in a spray tank to reduce the wetting time of wettable powders and to improve the penetration of water into water-dispersible granules. Examples of wetting agents used in wettable powder, suspension concentrate, and water-dispersible granule formulations are: sodium lauryl sulfate; sodium dioctyl sulfosuccinate; alkyl phenol ethoxylates; and aliphatic alcohol ethoxylates.

A dispersing agent is a substance which adsorbs onto the surface of particles and helps to preserve the state of dispersion of the particles and prevents them from reaggregating. Dispersing agents are added to agrochemical formulations to facilitate dispersion and suspension during manufacture, and to ensure the particles redisperse into water in a spray tank. They are widely used in wettable powders, suspension concentrates and water-dispersible granules. Surfactants that are used as dispersing agents have the ability to adsorb strongly onto a particle surface and provide a charged or steric barrier to reaggregation of particles. The most commonly used surfactants are anionic, non-ionic, or mixtures of the two types. For wettable powder formulations, the most common dispersing agents are sodium lignosulfonates. For suspension concentrates, very good adsorption and stabilization are obtained using polyelectrolytes, such as sodium naphthalene sulfonate formaldehyde condensates. Tristyrylphenol ethoxylate phosphate esters are also used. Non-ionics such as alkylarylethylene oxide condensates and EO-PO block copolymers are sometimes combined

with anionics as dispersing agents for suspension concentrates. In recent years, new types of very high molecular weight polymeric surfactants have been developed as dispersing agents. These have very long hydrophobic 'backbones' and a large number of ethylene oxide chains forming the 'teeth' of a 'comb' surfactant. These high molecular weight polymers can give very good long-term stability to suspension concentrates because the hydrophobic backbones have many anchoring points onto the particle surfaces. Examples of dispersing agents used in agrochemical formulations are: sodium lignosulfonates; sodium naphthalene sulfonate formaldehyde condensates; tristyrylphenol ethoxylate phosphate esters; aliphatic alcohol ethoxylates; alkyl ethoxylates; EO-PO block copolymers; and graft copolymers.

An emulsifying agent is a substance which stabilizes a suspension of droplets of one liquid phase in another liquid phase. Without the emulsifying agent the two liquids would separate into two immiscible liquid phases. The most commonly used emulsifier blends contain alkylphenol or aliphatic alcohol with twelve or more ethylene oxide units and the oil-soluble calcium salt of dodecylbenzenesulfonic acid. A range of hydrophile-lipophile balance ("HLB") values from 8 to 18 will normally provide good stable emulsions. Emulsion stability can sometimes be improved by the addition of a small amount of an EO-PO block copolymer surfactant.

A solubilizing agent is a surfactant which will form micelles in water at concentrations above the critical micelle concentration. The micelles are then able to dissolve or solubilize water-insoluble materials inside the hydrophobic part of the micelle. The types of surfactants usually used for solubilization are non-ionics, sorbitan monooleates, sorbitan monooleate ethoxylates, and methyl oleate esters.

Surfactants are sometimes used, either alone or with other additives such as mineral or vegetable oils as adjuvants to spray-tank mixes to improve the biological performance of the pesticide on the target. The types of surfactants used for bioenhancement depend generally on the nature and mode of action of the pesticide. However, they are often non-ionics such as: alkyl ethoxylates; linear aliphatic alcohol ethoxylates; aliphatic amine ethoxylates.

A carrier or diluent in an agricultural formulation is a material added to the pesticide to give a product of the required strength. Carriers are usually materials with high absorptive capacities, while diluents are usually materials with low absorptive capacities. Carriers and diluents are used in the formulation of dusts, wettable powders, granules and water-dispersible granules.

Organic solvents are used mainly in the formulation of emulsifiable concentrates, oil-in-water emulsions, suspoemulsions, and ultra-low volume formulations, and to a lesser

extent, granular formulations. Sometimes mixtures of solvents are used. The first main groups of solvents are aliphatic paraffinic oils such as kerosene or refined paraffins. The second main group (and the most common) comprises the aromatic solvents such as xylene and higher molecular weight fractions of C9 and C10 aromatic solvents. Chlorinated hydrocarbons are useful as cosolvents to prevent crystallization of pesticides when the formulation is emulsified into water. Alcohols are sometimes used as cosolvents to increase solvent power. Other solvents may include vegetable oils, seed oils, and esters of vegetable and seed oils.

Thickeners or gelling agents are used mainly in the formulation of suspension concentrates, emulsions and suspoemulsions to modify the rheology or flow properties of the liquid and to prevent separation and settling of the dispersed particles or droplets. Thickening, gelling, and anti-settling agents generally fall into two categories, namely water-insoluble particulates and water-soluble polymers. It is possible to produce suspension concentrate formulations using clays and silicas. Examples of these types of materials, include, but are not limited to, montmorillonite, bentonite, magnesium aluminum silicate, and attapulgite. Water-soluble polysaccharides have been used as thickening-gelling agents for many years. The types of polysaccharides most commonly used are natural extracts of seeds and seaweeds or are synthetic derivatives of cellulose. Examples of these types of materials include, but are not limited to, guar gum; locust bean gum; carrageenam; alginates; methyl cellulose; sodium carboxymethyl cellulose (SCMC); hydroxyethyl cellulose (HEC). Other types of anti-settling agents are based on modified starches, polyacrylates, polyvinyl alcohol and polyethylene oxide. Another good anti-settling agent is xanthan gum.

Microorganisms can cause spoilage of formulated products. Therefore preservation agents are used to eliminate or reduce their effect. Examples of such agents include, but are not limited to: propionic acid and its sodium salt; sorbic acid and its sodium or potassium salts; benzoic acid and its sodium salt; *p*-hydroxybenzoic acid sodium salt; methyl *p*-hydroxybenzoate; and 1,2-benzisothiazolin-3-one (BIT).

The presence of surfactants often causes water-based formulations to foam during mixing operations in production and in application through a spray tank. In order to reduce the tendency to foam, anti-foam agents are often added either during the production stage or before filling into bottles. Generally, there are two types of anti-foam agents, namely silicones and non-silicones. Silicones are usually aqueous emulsions of dimethyl polysiloxane, while the non-silicone anti-foam agents are water-insoluble oils, such as

octanol and nonanol, or silica. In both cases, the function of the anti-foam agent is to displace the surfactant from the air-water interface.

"Green" agents (*e.g.*, adjuvants, surfactants, solvents) can reduce the overall environmental footprint of crop protection formulations. Green agents are biodegradable and generally derived from natural and/or sustainable sources, *e.g.* plant and animal sources. Specific examples are: vegetable oils, seed oils, and esters thereof, also alkoxylated alkyl polyglucosides.

For further information, see "Chemistry and Technology of Agrochemical Formulations" edited by D.A. Knowles, copyright 1998 by Kluwer Academic Publishers. Also see "Insecticides in Agriculture and Environment – Retrospects and Prospects" by A.S. Perry, I. Yamamoto, I. Ishaaya, and R. Perry, copyright 1998 by Springer-Verlag.

PESTS

In general, the molecules of Formula One may be used to control pests *e.g.* beetles, earwigs, cockroaches, flies. aphids, scales, whiteflies, leafhoppers, ants, wasps, termites, moths, butterflies, lice, grasshoppers, locusts, crickets, fleas, thrips, bristletails, mites, ticks, nematodes, and symphylans.

In another embodiment, the molecules of Formula One may be used to control pests in the **Phyla Nematoda** and/or **Arthropoda**.

In another embodiment, the molecules of Formula One may be used to control pests in the **Subphyla Chelicerata**, **Myriapoda**, and/or **Hexapoda**.

In another embodiment, the molecules of Formula One may be used to control pests in the **Classes** of **Arachnida, Symphyla**, and/or **Insecta**.

In another embodiment, the molecules of Formula One may be used to control pests of the **Order Anoplura**. A non-exhaustive list of particular genera includes, but is not limited to, *Haematopinus* spp., *Hoplopleura* spp., *Linognathus* spp., *Pediculus* spp., and *Polyplax* spp. A non-exhaustive list of particular species includes, but is not limited to, *Haematopinus asini, Haematopinus suis*, *Linognathus setosus*, *Linognathus ovillus*, *Pediculus humanus capitis*, *Pediculus humanus humanus*, and *Pthirus pubis.*

In another embodiment, the molecules of Formula One may be used to control pests in the **Order Coleoptera**. A non-exhaustive list of particular genera includes, but is not limited to, *Acanthoscelides* spp., *Agriotes* spp., *Anthonomus* spp., *Apion* spp., *Apogonia* spp., *Aulacophora* spp., *Bruchus* spp., *Cerosterna* spp., *Cerotoma* spp., *Ceutorhynchus* spp., *Chaetocnema* spp., *Colaspis* spp., *Ctenicera* spp., *Curculio* spp., *Cyclocephala* spp., *Diabrotica* spp., *Hypera* spp., *Ips* spp., *Lyctus* spp., *Megascelis* spp., *Meligethes* spp.,

Otiorhynchus spp., *Pantomorus* spp., *Phyllophaga* spp., *Phyllotreta* spp., *Rhizotrogus* spp., *Rhynchites* spp., *Rhynchophorus* spp., *Scolytus* spp., *Sphenophorus* spp., *Sitophilus* spp., and *Tribolium* spp. A non-exhaustive list of particular species includes, but is not limited to, *Acanthoscelides obtectus, Agrilus planipennis, Anoplophora glabripennis, Anthonomus grandis, Ataenius spretulus, Atomaria linearis, Bothynoderes punctiventris, Bruchus pisorum, Callosobruchus maculatus, Carpophilus hemipterus, Cassida vittata, Cerotoma trifurcata, Ceutorhynchus assimilis, Ceutorhynchus napi, Conoderus scalaris, Conoderus stigmosus, Conotrachelus nenuphar, Cotinis nitida, Crioceris asparagi, Cryptolestes ferrugineus, Cryptolestes pusillus, Cryptolestes turcicus, Cylindrocopturus adspersus, Deporaus marginatus, Dermestes lardarius, Dermestes maculatus, Epilachna varivestis, Faustinus cubae, Hylobius pales, Hypera postica, Hypothenemus hampei, Lasioderma serricorne, Leptinotarsa decemlineata, Liogenys fuscus, Liogenys suturalis, Lissorhoptrus oryzophilus, Maecolaspis joliveti, Melanotus communis, Meligethes aeneus, Melolontha melolontha, Oberea brevis, Oberea linearis, Oryctes rhinoceros, Oryzaephilus mercator, Oryzaephilus surinamensis, Oulema melanopus, Oulema oryzae, Phyllophaga cuyabana, Popillia japonica, Prostephanus truncatus, Rhyzopertha dominica,, Sitona lineatus, Sitophilus granarius, Sitophilus oryzae, Sitophilus zeamais, Stegobium paniceum, Tribolium castaneum, Tribolium confusum, Trogoderma variabile,* and *Zabrus tenebrioides.*

In another embodiment, the molecules of Formula One may be used to control pests of the **Order Dermaptera**.

In another embodiment, the molecules of Formula One may be used to control pests of the **Order Blattaria**. A non-exhaustive list of particular species includes, but is not limited to, *Blattella germanica, Blatta orientalis, Parcoblatta pennsylvanica, Periplaneta americana, Periplaneta australasiae, Periplaneta brunnea, Periplaneta fuliginosa, Pycnoscelus surinamensis,* and *Supella longipalpa.*

In another embodiment, the molecules of Formula One may be used to control pests of the **Order Diptera**. A non-exhaustive list of particular genera includes, but is not limited to, *Aedes* spp., *Agromyza* spp., *Anastrepha* spp., *Anopheles* spp., *Bactrocera* spp., *Ceratitis* spp., *Chrysops* spp., *Cochliomyia* spp., *Contarinia* spp., *Culex* spp., *Dasineura* spp., *Delia* spp., *Drosophila* spp., *Fannia* spp., *Hylemyia* spp., *Liriomyza* spp., *Musca* spp., *Phorbia* spp., *Tabanus* spp., and *Tipula* spp. A non-exhaustive list of particular species includes, but is not limited to, *Agromyza frontella, Anastrepha suspensa, Anastrepha ludens, Anastrepha obliqa, Bactrocera cucurbitae, Bactrocera dorsalis, Bactrocera invadens, Bactrocera zonata, Ceratitis capitata, Dasineura brassicae, Delia platura, Fannia canicularis, Fannia scalaris,*

Gasterophilus intestinalis, *Gracillia perseae*, *Haematobia irritans*, *Hypoderma lineatum*, *Liriomyza brassicae*, *Melophagus ovinus*, *Musca autumnalis*, *Musca domestica*, *Oestrus ovis*, *Oscinella frit*, *Pegomya betae*, *Psila rosae*, *Rhagoletis cerasi*, *Rhagoletis pomonella*, *Rhagoletis mendax*, *Sitodiplosis mosellana*, and *Stomoxys calcitrans*.

In another embodiment, the molecules of Formula One may be used to control pests of the **Order Hemiptera**. A non-exhaustive list of particular genera includes, but is not limited to, *Adelges* spp., *Aulacaspis* spp., *Aphrophora* spp., *Aphis* spp., *Bemisia* spp., *Ceroplastes* spp., *Chionaspis* spp., *Chrysomphalus* spp., *Coccus* spp., *Empoasca* spp., *Lepidosaphes* spp., *Lagynotomus* spp., *Lygus* spp., *Macrosiphum* spp., *Nephotettix* spp., *Nezara* spp., *Philaenus* spp., *Phytocoris* spp., *Piezodorus* spp., *Planococcus* spp., *Pseudococcus* spp., *Rhopalosiphum* spp., *Saissetia* spp., *Therioaphis* spp., *Toumeyella* spp., *Toxoptera* spp., *Trialeurodes* spp., *Triatoma* spp. and *Unaspis* spp. A non-exhaustive list of particular species includes, but is not limited to, *Acrosternum hilare*, *Acyrthosiphon pisum*, *Aleyrodes proletella*, *Aleurodicus dispersus*, *Aleurothrixus floccosus*, *Amrasca biguttula biguttula*, *Aonidiella aurantii*, *Aphis gossypii*, *Aphis glycines*, *Aphis pomi*, *Aulacorthum solani*, *Bemisia argentifolii*, *Bemisia tabaci*, *Blissus leucopterus*, *Brachycorynella asparagi*, *Brevennia rehi*, *Brevicoryne brassicae*, *Calocoris norvegicus*, *Ceroplastes rubens*, *Cimex hemipterus*, *Cimex lectularius*, *Dagbertus fasciatus*, *Dichelops furcatus*, *Diuraphis noxia*, *Diaphorina citri*, *Dysaphis plantaginea*, *Dysdercus suturellus*, *Edessa meditabunda*, *Eriosoma lanigerum*, *Eurygaster maura*, *Euschistus heros*, *Euschistus servus*, *Helopeltis antonii*, *Helopeltis theivora*, *Icerya purchasi*, *Idioscopus nitidulus*, *Laodelphax striatellus*, *Leptocorisa oratorius*, *Leptocorisa varicornis*, *Lygus hesperus*, *Maconellicoccus hirsutus*, *Macrosiphum euphorbiae*, *Macrosiphum granarium*, *Macrosiphum rosae*, *Macrosteles quadrilineatus*, *Mahanarva frimbiolata*, *Metopolophium dirhodum*, *Mictis longicornis*, *Myzus persicae*, *Nephotettix cinctipes*, *Neurocolpus longirostris*, *Nezara viridula*, *Nilaparvata lugens*, *Parlatoria pergandii*, *Parlatoria ziziphi*, *Peregrinus maidis*, *Phylloxera vitifoliae*, *Physokermes piceae*,, *Phytocoris californicus*, *Phytocoris relativus*, *Piezodorus guildinii*, *Poecilocapsus lineatus*, *Psallus vaccinicola*, *Pseudacysta perseae*, *Pseudococcus brevipes*, *Quadraspidiotus perniciosus*, *Rhopalosiphum maidis*, *Rhopalosiphum padi*, *Saissetia oleae*, *Scaptocoris castanea*, *Schizaphis graminum*, *Sitobion avenae*, *Sogatella furcifera*, *Trialeurodes vaporariorum*, *Trialeurodes abutiloneus*, *Unaspis yanonensis*, and *Zulia entrerriana*.

In another embodiment, the molecules of Formula One may be used to control pests of the **Order Hymenoptera**. A non-exhaustive list of particular genera includes, but is not

limited to, *Acromyrmex* spp., *Atta* spp., *Camponotus* spp., *Diprion* spp., *Formica* spp., *Monomorium* spp., *Neodiprion* spp., *Pogonomyrmex* spp., *Polistes* spp., *Solenopsis* spp., *Vespula* spp., and *Xylocopa* spp. A non-exhaustive list of particular species includes, but is not limited to, *Athalia rosae, Atta texana, Iridomyrmex humilis*, *Monomorium minimum, Monomorium pharaonis, Solenopsis invicta, Solenopsis geminata, Solenopsis molesta, Solenopsis richtery, Solenopsis xyloni,* and *Tapinoma sessile*.

In another embodiment, the molecules of Formula One may be used to control pests of the **Order Isoptera**. A non-exhaustive list of particular genera includes, but is not limited to, *Coptotermes* spp., *Cornitermes* spp., *Cryptotermes* spp., *Heterotermes* spp., *Kalotermes* spp., *Incisitermes* spp., *Macrotermes* spp., *Marginitermes* spp., *Microcerotermes* spp., *Procornitermes* spp., *Reticulitermes* spp., *Schedorhinotermes* spp., and *Zootermopsis* spp. A non-exhaustive list of particular species includes, but is not limited to, *Coptotermes curvignathus, Coptotermes frenchi, Coptotermes formosanus*, *Heterotermes aureus, Microtermes obesi, Reticulitermes banyulensis, Reticulitermes grassei, Reticulitermes flavipes*, *Reticulitermes hageni, Reticulitermes hesperus*, *Reticulitermes santonensis, Reticulitermes speratus, Reticulitermes tibialis,* and *Reticulitermes virginicus.*

In another embodiment, the molecules of Formula One may be used to control pests of the **Order Lepidoptera**. A non-exhaustive list of particular genera includes, but is not limited to, *Adoxophyes* spp., *Agrotis* spp., *Argyrotaenia* spp., *Cacoecia* spp., *Caloptilia* spp., *Chilo* spp., *Chrysodeixis* spp., *Colias* spp., *Crambus* spp., *Diaphania* spp., *Diatraea* spp., *Earias* spp., *Ephestia* spp., *Epimecis* spp., *Feltia* spp., *Gortyna* spp., *Helicoverpa* spp., *Heliothis* spp., *Indarbela* spp., *Lithocolletis* spp., *Loxagrotis* spp., *Malacosoma* spp., *Peridroma* spp., *Phyllonorycter* spp., *Pseudaletia* spp., *Sesamia* spp., *Spodoptera* spp., *Synanthedon* spp., and *Yponomeuta* spp. A non-exhaustive list of particular species includes, but is not limited to, *Achaea janata, Adoxophyes orana, Agrotis ipsilon*, *Alabama argillacea, Amorbia cuneana, Amyelois transitella, Anacamptodes defectaria, Anarsia lineatella*, *Anomis sabulifera*, *Anticarsia gemmatalis*, *Archips argyrospila*, *Archips rosana*, *Argyrotaenia citrana*, *Autographa gamma, Bonagota cranaodes, Borbo cinnara, Bucculatrix thurberiella, Capua reticulana, Carposina niponensis*, *Chlumetia transversa*, *Choristoneura rosaceana, Cnaphalocrocis medinalis, Conopomorpha cramerella, Cossus cossus*, *Cydia caryana, Cydia funebrana*, *Cydia molesta*, *Cydia nigricana, Cydia pomonella*, *Darna diducta, Diatraea saccharalis*, *Diatraea grandiosella, Earias insulana, Earias vittella, Ecdytolopha aurantianum, Elasmopalpus lignosellus*, *Ephestia cautella*, *Ephestia elutella*, *Ephestia kuehniella*, *Epinotia aporema, Epiphyas postvittana, Erionota thrax*, *Eupoecilia ambiguella*,

Euxoa auxiliaris, *Grapholita molesta*, *Hedylepta indicata*, *Helicoverpa armigera*, *Helicoverpa zea*, *Heliothis virescens*, *Hellula undalis*, *Keiferia lycopersicella*, *Leucinodes orbonalis*, *Leucoptera coffeella*, *Leucoptera malifoliella*, *Lobesia botrana*, *Loxagrotis albicosta*, *Lymantria dispar*, *Lyonetia clerkella*, *Mahasena corbetti*, *Mamestra brassicae*, *Maruca testulalis*, *Metisa plana*, *Mythimna unipuncta*, *Neoleucinodes elegantalis*, *Nymphula depunctalis*, *Operophtera brumata*, *Ostrinia nubilalis*, *Oxydia vesulia*, *Pandemis cerasana*, *Pandemis heparana*, *Papilio demodocus*, *Pectinophora gossypiella*, *Peridroma saucia*, *Perileucoptera coffeella*, *Phthorimaea operculella*, *Phyllocnistis citrella*, *Pieris rapae*, *Plathypena scabra*, *Plodia interpunctella*, *Plutella xylostella*, *Polychrosis viteana*, *Prays endocarpa*, *Prays oleae*, *Pseudaletia unipuncta*, *Pseudoplusia includens*, *Rachiplusia nu*, *Scirpophaga incertulas*, *Sesamia inferens*, *Sesamia nonagrioides*, *Setora nitens*, *Sitotroga cerealella*, *Sparganothis pilleriana*, *Spodoptera exigua*, *Spodoptera frugiperda*, *Spodoptera eridania*, *Thecla basilides*, *Tineola bisselliella*, *Trichoplusia ni*, *Tuta absoluta*, *Zeuzera coffeae*, and *Zeuzera pyrina*.

In another embodiment, the molecules of Formula One may be used to control pests of the **Order Mallophaga**. A non-exhaustive list of particular genera includes, but is not limited to, *Anaticola* spp., *Bovicola* spp., *Chelopistes* spp., *Goniodes* spp., *Menacanthus* spp., and *Trichodectes* spp. A non-exhaustive list of particular species includes, but is not limited to, *Bovicola bovis*, *Bovicola caprae*, *Bovicola ovis*, *Chelopistes meleagridis*, *Goniodes dissimilis*, *Goniodes gigas*, *Menacanthus stramineus*, *Menopon gallinae*, and *Trichodectes canis*.

In another embodiment, the molecules of Formula One may be used to control pests of the **Order Orthoptera**. A non-exhaustive list of particular genera includes, but is not limited to, *Melanoplus* spp., and *Pterophylla* spp. A non-exhaustive list of particular species includes, but is not limited to, *Anabrus simplex*, *Gryllotalpa africana*, *Gryllotalpa australis*, *Gryllotalpa brachyptera*, *Gryllotalpa hexadactyla*, *Locusta migratoria*, *Microcentrum retinerve*, *Schistocerca gregaria*, and *Scudderia furcata*.

In another embodiment, the molecules of Formula One may be used to control pests of the **Order Siphonaptera**. A non-exhaustive list of particular species includes, but is not limited to, *Ceratophyllus gallinae*, *Ceratophyllus niger*, *Ctenocephalides canis*, *Ctenocephalides felis*, and *Pulex irritans*.

In another embodiment, the molecules of Formula One may be used to control pests of the **Order Thysanoptera**. A non-exhaustive list of particular genera includes, but is not limited to, Caliothrips spp., Frankliniella spp., *Scirtothrips* spp., and *Thrips* spp. A non-

exhaustive list of particular sp. includes, but is not limited to, *Frankliniella fusca*, *Frankliniella occidentalis*, *Frankliniella schultzei, Frankliniella williamsi*, *Heliothrips haemorrhoidalis, Rhipiphorothrips cruentatus, Scirtothrips citri*, *Scirtothrips dorsalis*, and *Taeniothrips rhopalantennalis, Thrips hawaiiensis, Thrips nigropilosus, Thrips orientalis, Thrips tabaci*.

In another embodiment, the molecules of Formula One may be used to control pests of the **Order Thysanura**. A non-exhaustive list of particular genera includes, but is not limited to, *Lepisma* spp. and *Thermobia* spp.

In another embodiment, the molecules of Formula One may be used to control pests of the **Order Acarina**. A non-exhaustive list of particular genera includes, but is not limited to, *Acarus* spp., *Aculops* spp., *Boophilus* spp., *Demodex* spp., *Dermacentor* spp., *Epitrimerus* spp., *Eriophyes* spp., *Ixodes* spp., *Oligonychus* spp., *Panonychus* spp., *Rhizoglyphus* spp., and *Tetranychus* spp. A non-exhaustive list of particular species includes, but is not limited to, *Acarapis woodi, Acarus siro*, *Aceria mangiferae*, *Aculops lycopersici*, *Aculus pelekassi, Aculus schlechtendali*, *Amblyomma americanum*, *Brevipalpus obovatus*, *Brevipalpus phoenicis*, *Dermacentor variabilis*, *Dermatophagoides pteronyssinus*, *Eotetranychus carpini*, *Notoedres cati, Oligonychus coffeae, Oligonychus ilicis, Panonychus citri*, *Panonychus ulmi*, *Phyllocoptruta oleivora*, *Polyphagotarsonemus latus, Rhipicephalus sanguineus*, *Sarcoptes scabiei*, *Tegolophus perseaflorae, Tetranychus urticae*, and *Varroa destructor.*

In another embodiment, the molecules of Formula One may be used to control pest of the **Order Symphyla**. A non-exhaustive list of particular sp. includes, but is not limited to, *Scutigerella immaculata.*

In another embodiment, the molecules of Formula One may be used to control pests of the **Phylum Nematoda**. A non-exhaustive list of particular genera includes, but is not limited to, *Aphelenchoides* spp., *Belonolaimus* spp., *Criconemella* spp., *Ditylenchus* spp., *Heterodera* spp., *Hirschmanniella* spp., *Hoplolaimus* spp., *Meloidogyne* spp., *Pratylenchus* spp., and *Radopholus* spp. A non-exhaustive list of particular sp. includes, but is not limited to, *Dirofilaria immitis*, *Heterodera zeae, Meloidogyne incognita, Meloidogyne javanica, Onchocerca volvulus*, *Radopholus similis, and Rotylenchulus reniformis.*

For additional information consult "**HANDBOOK OF PEST CONTROL – THE BEHAVIOR, LIFE HISTORY, AND CONTROL OF HOUSEHOLD PESTS**" by Arnold Mallis, 9th Edition, copyright 2004 by GIE Media Inc.

APPLICATIONS

Molecules of Formula One are generally used in amounts from about 0.01 grams per hectare to about 5000 grams per hectare to provide control. Amounts from about 0.1 grams per hectare to about 500 grams per hectare are generally preferred, and amounts from about 1 gram per hectare to about 50 grams per hectare are generally more preferred.

The area to which a molecule of Formula One is applied can be any area inhabited (or maybe inhabited, or traversed by) a pest, for example: where crops, trees, fruits, cereals, fodder species, vines, turf and ornamental plants, are growing; where domesticated animals are residing; the interior or exterior surfaces of buildings (such as places where grains are stored), the materials of construction used in building (such as impregnated wood), and the soil around buildings. Particular crop areas to use a molecule of Formula One include areas where apples, corn, sunflowers, cotton, soybeans, canola, wheat, rice, sorghum, barley, oats, potatoes, oranges, alfalfa, lettuce, strawberries, tomatoes, peppers, crucifers, pears, tobacco, almonds, sugar beets, beans and other valuable crops are growing or the seeds thereof are going to be planted. It is also advantageous to use ammonium sulfate with a molecule of Formula One when growing various plants.

Controlling pests generally means that pest populations, pest activity, or both, are reduced in an area. This can come about when: pest populations are repulsed from an area; when pests are incapacitated in or around an area; or pests are exterminated, in whole, or in part, in or around an area. Of course, a combination of these results can occur. Generally, pest populations, activity, or both are desirably reduced more than fifty percent, preferably more than 90 percent. Generally, the area is not in or on a human; consequently, the locus is generally a non-human area.

The molecules of Formula One may be used in mixtures, applied simultaneously or sequentially, alone or with other compounds to enhance plant vigor (*e.g.* to grow a better root system, to better withstand stressful growing conditions). Such other compounds are, for example, compounds that modulate plant ethylene receptors, most notably 1-methylcyclopropene (also known as 1-MCP). Furthermore, such molecules may be used during times when pest activity is low, such as before the plants that are growing begin to produce valuable agricultural commodities. Such times include the early planting season when pest pressure is usually low.

The molecules of Formula One can be applied to the foliar and fruiting portions of plants to control pests. The molecules will either come in direct contact with the pest, or the pest will consume the pesticide when eating leaf, fruit mass, or extracting sap, that contains the pesticide. The molecules of Formula One can also be applied to the soil, and when

applied in this manner, root and stem feeding pests can be controlled. The roots can absorb a molecule taking it up into the foliar portions of the plant to control above ground chewing and sap feeding pests.

Generally, with baits, the baits are placed in the ground where, for example, termites can come into contact with, and/or be attracted to, the bait. Baits can also be applied to a surface of a building, (horizontal, vertical, or slant surface) where, for example, ants, termites, cockroaches, and flies, can come into contact with, and/or be attracted to, the bait. Baits can comprise a molecule of Formula One.

The molecules of Formula One can be encapsulated inside, or placed on the surface of a capsule. The size of the capsules can range from nanometer size (about 100-900 nanometers in diameter) to micrometer size (about 10-900 microns in diameter).

Because of the unique ability of the eggs of some pests to resist certain pesticides, repeated applications of the molecules of Formula One may be desirable to control newly emerged larvae.

Systemic movement of pesticides in plants may be utilized to control pests on one portion of the plant by applying (for example by spraying an area) the molecules of Formula One to a different portion of the plant. For example, control of foliar-feeding insects can be achieved by drip irrigation or furrow application, by treating the soil with for example pre- or post-planting soil drench, or by treating the seeds of a plant before planting.

Seed treatment can be applied to all types of seeds, including those from which plants genetically modified to express specialized traits will germinate. Representative examples include those expressing proteins toxic to invertebrate pests, such as *Bacillus thuringiensis* or other insecticidal toxins, those expressing herbicide resistance, such as "Roundup Ready" seed, or those with "stacked" foreign genes expressing insecticidal toxins, herbicide resistance, nutrition-enhancement, drought resistance, or any other beneficial traits. Furthermore, such seed treatments with the molecules of Formula One may further enhance the ability of a plant to better withstand stressful growing conditions. This results in a healthier, more vigorous plant, which can lead to higher yields at harvest time. Generally, about 1 gram of the molecules of Formula One to about 500 grams per 100,000 seeds is expected to provide good benefits, amounts from about 10 grams to about 100 grams per 100,000 seeds is expected to provide better benefits, and amounts from about 25 grams to about 75 grams per 100,000 seeds is expected to provide even better benefits.

It should be readily apparent that the molecules of Formula One may be used on, in, or around plants genetically modified to express specialized traits, such as *Bacillus*

thuringiensis or other insecticidal toxins, or those expressing herbicide resistance, or those with "stacked" foreign genes expressing insecticidal toxins, herbicide resistance, nutrition-enhancement, or any other beneficial traits.

The molecules of Formula One may be used for controlling endoparasites and ectoparasites in the veterinary medicine sector or in the field of non-human animal keeping. The molecules of Formula One are applied, such as by oral administration in the form of, for example, tablets, capsules, drinks, granules, by dermal application in the form of, for example, dipping, spraying, pouring on, spotting on, and dusting, and by parenteral administration in the form of, for example, an injection.

The molecules of Formula One may also be employed advantageously in livestock keeping, for example, cattle, sheep, pigs, chickens, and geese. They may also be employed advantageously in pets such as, horses, dogs, and cats. Particular pests to control would be fleas and ticks that are bothersome to such animals. Suitable formulations are administered orally to the animals with the drinking water or feed. The dosages and formulations that are suitable depend on the species.

The molecules of Formula One may also be used for controlling parasitic worms, especially of the intestine, in the animals listed above.

The molecules of Formula One may also be employed in therapeutic methods for human health care. Such methods include, but are limited to, oral administration in the form of, for example, tablets, capsules, drinks, granules, and by dermal application.

Pests around the world have been migrating to new environments (for such pest) and thereafter becoming a new invasive species in such new environment. The molecules of Formula One may also be used on such new invasive species to control them in such new environment.

The molecules of Formula One may also be used in an area where plants, such as crops, are growing (*e.g.* pre-planting, planting, pre-harvesting) and where there are low levels (even no actual presence) of pests that can commercially damage such plants. The use of such molecules in such area is to benefit the plants being grown in the area. Such benefits, may include, but are not limited to, improving the health of a plant, improving the yield of a plant (*e.g.* increased biomass and/or increased content of valuable ingredients), improving the vigor of a plant (*e.g.* improved plant growth and/or greener leaves), improving the quality of a plant (*e.g.* improved content or composition of certain ingredients), and improving the tolerance to abiotic and/or biotic stress of the plant.

Before a pesticide can be used or sold commercially, such pesticide undergoes lengthy evaluation processes by various governmental authorities (local, regional, state, national, and international). Voluminous data requirements are specified by regulatory authorities and must be addressed through data generation and submission by the product registrant or by a third party on the product registrant's behalf, often using a computer with a connection to the World Wide Web. These governmental authorities then review such data and if a determination of safety is concluded, provide the potential user or seller with product registration approval. Thereafter, in that locality where the product registration is granted and supported, such user or seller may use or sell such pesticide.

A molecule according to Formula One can be tested to determine its efficacy against pests. Furthermore, mode of action studies can be conducted to determine if said molecule has a different mode of action than other pesticides. Thereafter, such acquired data can be disseminated, such as by the internet, to third parties.






The headings in this document are for convenience only and must not be used to interpret any portion hereof.









TABLE SECTION









BAW, CEW & CL Rating Table	
% Control (or Mortality)	**Rating**
50-100	A
More than 0 – Less than 50	B
Not Tested	C
No activity noticed in this bioassay	D









GPA Rating Table	
% Control (or Mortality)	**Rating**
80-100	A
More than 0 – Less than 80	B
Not Tested	C
No activity noticed in this bioassay	D









Table 1: Structures for Compounds









Compound Number	Structure
AI34	
AI36	
AI37	
AI38	
AI39	








AI40	
AI41	
AI44	
AI45	
AC1	
AC2	
AC3	
AC4	







AC5	
AC6	
AC7	
AC8	
AC9	
AC10	
AC11	
AC12	







AC13	
AC14	
AC15	
AC16	
AC17	
AC18	
AC19	
AC20	








AC21	
AC22	
AC23	
AC24	
AC25	
AC26	
AC27	
AC28	








AC29	
AC30	
AC31	
AC32	
AC33	
AC34	
AC35	
AC36	







AC37	
AC38	
AC39	
AC40	
AC41	
AC42	
AC43	







AC44	
AC45	
AC46	
AC47	
AC48	
AC49	








AC50	
AC51	
AC52	
AC53	
AC54	
AC57	








AC58	
AC59	
AC60	
AC61	
AC62	
AC63	
AC64	







AC65	
AC66	
AC67	
AC68	
AC69	
AC70	
AC71	







AC72	
AC75	
AC76	
AC77	
AC78	
AC79	








AC80	
AC81	
AC82	
AC83	
AC84	
AC85	






AC86	
AC87	
AC89	
AC90	
AC91	
AC92	
AC93	






AC94	
AC95	
AC96	
AC97	
AC98	
AC99	
AC100	








AC101	
AC102	
AC103	
AC104	
AC105	
AC106	






AC107	
AC108	
AC109	
AC110	
AC111	
AC112	







AC113	
AC114	
AC115	
AC116	
AC117	
AC118	
BC1	










BC2	
BC3	
BC4	
BC5	
BC6	









BC7	
BC8	
BC9	
BC10	
BC11	









BC12	
BC13	
BC14	
CI4	
CI5	
CI8	
CI9	









CI34	
CI35	
CI36	
CI37	
CI38	








CI39	
CI40	
CI41	
CI49	
CI50	
CI51	









CI52	
CI53	
CI54	
CI55	
CI56	
CI57	
CC1	
CC2	
CC3	









CC4	
CC5	
CC6	
CC7	
CC8	
CC9	
CC10	
CC11	









CC12	
CC13	
CC14	
CC15	
CC16	
CC17	
CC18	
CC19	









CC20	
CC21	
CC22	
CC23	
CC24	
CC25	
CC26	
CC27	









CC28	
CC29	
CC30	
CC31	
CC32	
CC33	
CC34	








CC35	
CC36	
CC37	
CC38	
CC39	
CC40	
CC41	
CC42	









CC43	
CC44	
CC45	
CC46	
CC47	
CC48	
CC49	
CC50	









CC51	
CC52	
CC53	
CC54	
DC1	
DC2	
DC3	
DC4	








DC5	
DC6	
DC7	
DC8	
DC9	
DC10	
DC11	
DC12	









DC13	
DC14	
DC15	
DC16	
DC17	
DC18	
DC19	
DC20	









DC21	
DC22	
DC23	
DC24	
DC25	
DC26	
DC27	

DC28	
DC29	
DC30	
DC31	
DC32	
DC33	
DC34	
DC35	

DC36	
DC37	
DC38	
DC39	
DC40	
DC41	
DC42	
DC43	

DC44	
DC45	
DC46	
DC47	
DC48	
DC49	
DC50	

DC51	
DC52	
DC53	
DC54	
DC55	
DC56	
DC57	
DC58	

DC59	
DC60	
DC61	
DC62	
DC63	
DC64	
DC65	
DC66	





















DC67	
DC68	
DC69	
DC70	






Table 1A: Structures for F Compounds







Compound Number	Structure	Appearance	Prepared as in Example:
F1		white foam	135






F2		brown gum	15
F3		pale yellow gum	15
F4		yellow solid	15
F5			108
F6			108






F7		yellow solid	15
F8		off-white solid	15
F8A		pale yellow solid	15
F9		yellow solid	15
F10		off-white solid	132

F11	 (–)-	off-white solid	132
F12	 (+)-	off-white solid	133
F13	 (–)-	off-white solid	133
F14		pale yellow solid	15
F15		yellow solid	15

F15A		yellow solid	15
F16		pale yellow solid	15
F16A		yellow solid	15
F17		off-white solid	15
F18		yellow solid	15

F19		yellow gum	15
F20		yellow solid	15
F20A		off-white solid	133
F20B		off-white solid	133
F20C		white solid	88
F21		yellow gum	15

F22		light brown gum	15
F23		pale yellow liquid	15
F23A		yellow solid	15
F24		pale yellow liquid	15
F25		yellow solid	15

F26		brown gum	15
F27		pale yellow gum	15
F28		brown gum	15
F29		pale yellow gum	19
F30		brown gum	15
































F31		pale yellow gum	134









Table 1B: Structures of Prophetic Compounds Subsequently Exemplified









Compound Number	Structure	Appearance	Prepared as in Example:
P1		Yellow solid	15
P2		Brown gum	15
P12		Off white solid	15
P14		Light yellow gum	19
P15		Light yellow gum	19
P82		Brown semi solid	15









P84		Pale yellow solid	15
P156		Green liquid	15
P226		Brown semi solid	15
P228		Brown semi solid	15
P298		Yellow solid	15
P300		Brown gum	15
P442		Pale yellow solid	15
P444		Pale yellow solid	15









P514		Off white solid	15
P516		Brown solid	15
P568		Brown semi solid	15
P586		Brown semi solid	15
P588		Brown semi solid	15
P660		Pale yellow solid	15
P730		Yellow solid	15
P732		Brown semi solid	15









P802		Brown gum	15
P804		Brown gum	15
P1090		Light brown solid	15
P1092		Brown semi solid	15
P1197		Off white solid	15
P1269		Pale yellow solid	15
P1340		Pale yellow liquid	15
P1411		Pale yellow liquid	15









P1483		Off white solid	15
P1556		Brown solid	15
P1558		Brown solid	134
P1559		Brown gum	134
P1560		Off white solid	19
P1564		Yellow solid	19
P1566		Brown solid	15
P1589		Pale yellow gum	15

P1591		Brown gum	15
P1592		Pale yellow gum	15
P1599		Brown semi solid	12
P1601		Brown viscous liquid	15
P1603		Off white solid	135
P1611A		Pale yellow solid	137
P1613A		Brown solid	137
P1616		Brown gum	15

P1616A		Pale yellow semi solid	137
P1618A		Brown solid	137
P1621		Brown viscous liquid	12
P1623		Brown liquid	12
P1624		Brown semi solid	12
P1631A		Brown viscous liquid	137
P1633A		Brown solid	137
P1636		Off white solid	135

P1638		Pale yellow solid	15
P1640		Brown semi solid	15
P1641		Brown solid	15
P1691		Brown gum	15
P1693		Pale yellow semi solid	15
P1696		Yellow solid	15
P1698		Yellow semi solid	15
P1776		White solid	15

P1781		White solid	15
P1806		Brown gum	15
P1808		Pale yellow gum	15
P1862		Off white gum	15
P1864		Brown gum	19
P1866		Brown liquid	19
P1969		Brown solid	15
P1970		Brown gum	15

P1971		Pale yellow solid	15
P1972		Light brown solid	15
P2009		Yellow solid	15
P2010		Yellow gum	15
P2011		Yellow solid	15
P2012		Off white solid	15
P2036		Brown solid	15
P2038		Yellow liquid	15
































P2041		Brown solid	15
P2043		Dark green solid	15
P2056		Brown solid	15
P2058		Pale yellow semi solid	15

Table 1C: Structures for FA Compounds

Compound Number	Structure	Appearance	Prepared as in Example:
FA1		Pale yellow gum	15
FA2		Light green liquid	15
FA3		Brown gum	15

FA4		White foam	138
FA5		White foam	138
FA6		Yellow solid	146
FA7		Pale yellow solid	137
FA8		Pale yellow gum	15
FA9		Yellow gum	134
FA10		Yellow gum	134
FA11		Brown semi solid	134

FA12		Brown semi solid	134
FA13		Light green solid	147
FA14		White foam	142
FA15		White foam	142
FA16		Pale yellow solid	15
FA17		Off white solid	15
FA18		Yellow semi solid	15
FA19		Pale yellow solid	15

FA20		Pale yellow solid	15
FA21		Brown gum	15
FA22		Yellow gum	15
FA23		Light yellow oil	142
FA24		Off white solid	15
FA25		Yellow solid	15
FA26		Light green solid	12
FA27		Pale green solid	12








FA28		Brown solid	15
FA29		Brown liquid	15
FA30		Brown gum	12
FA31		Pale yellow solid	12
FA32		Yellow solid	15
FA33		Yellow solid	135
FA34		Off white solid	149

Table 2: Analytical Data for Compounds in Table 1.

Compound Number	mp (°C)	ESIMS	^{1}H NMR (δ)[a]	IR (cm^{-1})

Compound Number	mp (°C)	ESIMS	^{1}H NMR (δ)[a]	IR (cm^{-1})
AC1	156–161	386.09 ([M-H]$^-$)	7.83 (m, 2H), 7.68-7.63 (m, 5H), 6.93 (dd, *J* = 15.6, 8.0 Hz, 1H), 6.81 (d *J* = 15.6 Hz, 1H,), 4.15 (m, 1H), 2.80 (s, 3H)	
AC2	110–112	374 ([M+H]$^+$)	7.80 (d, *J* = 8.4 Hz, 2H), 7.48 (d, *J* = 8.0 Hz, 2H), 7.38 (m, 1H), 7.30 (s, 2H), 6.65 (d, *J* = 16.0 Hz, 1H), 6.46 (dd, *J* = 16.0, 8.0 Hz, 1H), 4.15 (m, 1H)	
AC3	162–166	402.24 ([M+H]$^+$)	7.42 (m, 4H), 7.37 (t, *J* = 1.8 Hz, 1H), 7.28 (s, 2H), 6.63 (d, *J* = 16.0 Hz, 1H), 6.41 (dd, *J* = 16.0, 8.4 Hz, 1H), 4.15 (m, 1H), 3.20 (s, 3H), 3.00 (s, 3H)	
AC4	122–126	454 ([M-H]$^-$)	7.79 (d, *J* = 1.2 Hz, 2H), 7.48 (d, *J* = 8.4 Hz, 2H), 7.38 (t, *J* = 1.8 Hz, 1H), 7.30 (s, 2H), 6.64 (d, *J* = 15.6 Hz, 1H), 6.40 (dd, *J* = 15.6, 8.0 Hz, 1H), 6.30 (m, 1H), 4.15 (m, 3H)	
AC5		444.12 ([M+H]$^+$)	7.67 (s, 3H), 7.64 (d, *J* = 8.0 Hz, 2H), 7.42 (d, *J* = 8.0 Hz, 2H), 6.91 (dd, *J* = 15.6, 8.0 Hz, 1H), 6.80 (d, *J* = 15.6 Hz, 1H), 4.80 (m, 1H), 3.60 (br s, 8H)	
AC6		468.40 ([M-H]$^-$)	7.40 (m, 2H), 7.26 (m, 3H), 6.56 (d, *J* = 16.0 Hz, 1H), 6.48 (dd, *J* = 16.0, 8.0 Hz, 1H), 5.82 (br s, 1H), 4.08 (m, 3H), 2.52 (s, 3H)	1657, 1113, 804

Compound Number	mp (°C)	ESIMS	^{1}H NMR (δ)[a]	IR (cm^{-1})
AC7		511.02 ([M-H]$^-$)	8.39 (s, 1H), 7.74 (m, 1H), 7.39 (m, 3H), 7.24 (m, 4H), 6.58 (d, *J* = 16.0 Hz, 1H), 6.38 (dd, *J* = 16.0, 8.0 Hz, 1H), 6.16 (br s, 1H), 4.63 (m, 2H), 4.12 (m, 1H), 2.41 (s, 3H)	3276, 1645, 1111, 801
AC8		454.11 ([M-H]$^-$)	7.39 (s, 1H), 7.22 (m, 2H), 7.19 (m, 3H), 6.53 (d, *J* = 16.0 Hz, 1H), 6.39-6.34 (dd, *J* = 16.0, 8.0 Hz, 1H), 4.22 (m, 1H), 3.95 (t, *J* = 7.0 Hz, 2H), 2.62 (t, *J* = 8.0 Hz, 2H), 2.30 (s, 3H), 2.18 (m, 2H)	1748, 1112, 801
AC9		494.02 ([M-H]$^-$)	7.45 (t, *J* = 7.6 Hz, 1H), 7.36 (m, 2H), 7.21 (m, 3H), 7.15 (m, 4H), 6.56 (d, *J* = 16.0 Hz, 1H), 6.38 (dd, *J* = 16.0, 8.4 Hz, 1H), 6.08 (br s, 1H), 4.68 (d, *J* = 5.6 Hz, 2H), 4.11 (m, 1H), 2.44 (s, 3H)	3276, 1645, 1112, 801
A10	140–143	458.00 ([M-H]$^-$)	7.38 (t, *J* = 1.6 Hz, 1H), 7.34 (d, *J* = 7.6 Hz, 1H), 7.27 (m, 2H), 7.24 (m, 2H), 6.57 (d, *J* = 16.0 Hz, 1H), 6.40 (dd, *J* = 16.0, 8.0 Hz, 1H), 6.16 (m 1H), 5.44 (m, 1H), 4.12 (m, 1H), 3.51 (m, 2H), 3.40 (m, 2H), 2.44 (s, 3H)	
AC11		476.17 ([M-H]$^-$)	7.39-7.29 (m, 9H), 7.24 (m, 2H), 6.56 (d, *J* = 16.0 Hz, 1H), 6.38 (dd, *J* = 16.0, 8.0 Hz, 1H), 5.99 (br s, 1H), 4.63 (d, *J* = 6.0 Hz, 1H), 4.11 (m, 1H), 2.47 (s, 3H)	3287, 1644, 1112, 801

Compound Number	mp (°C)	ESIMS	^{1}H NMR (δ)[a]	IR (cm^{-1})
AC12		479.30 ([M+H]$^+$)	8.63 (d, *J* = 4.4 Hz, 1H), 7.71 (m, 1H), 7.47 (d, *J* = 8.4 Hz, 1H), 7.37 (m, 2H), 7.32 (m, 2H), 7.23 (m, 2H), 7.13 (m, 1H), 6.58 (d, *J* = 16.0 Hz, 1H), 6.40 (dd, *J* = 16.0, 8.0 Hz, 1H), 4.75 (d, *J* = 4.8 Hz, 2H), 4.12 (m, 1H), 2.49 (s, 3H)	3293, 1653, 1112, 800
AC13	75–78	490.04 ([M-H]$^-$)	7.38 (m, 2H), 7.27 (m, 3H), 7.23 (br s, 1H), 6.58 (d, *J* = 16.0 Hz, 1H), 6.45 (m 1H), 6.42 (dd, *J* = 16.0, 8.4 Hz, 1H), 4.91 (m 1H), 4.64 (m, 2H), 4.14 (m, 1H), 4.04 (m, 2H), 2.46 (s, 3H)	
AC14		480.99 ([M+2H]$^+$)	8.63 (s, 2H), 7.76 (d, *J* = 8.0 Hz, 1H), 7.36 (m, 3H), 7.22 (m, 1H), 7.13 (m, 2H), 6.57 (d, *J* = 16.0 Hz, 1H), 6.39 (dd, *J* = 16.0, 8.0 Hz, 1H), 6.13 (br s, 1H), 4.66 (d, *J* = 5.6 Hz, 2H), 4.11 (m, 1H), 2.46 (s, 3H)	3293, 1645, 1113, 800
AC15	59–61	516.86 ([M-H]$^-$)	7.45 (s, 1H), 7.37 (m, 1H), 7.34 (m, 1H), 7.26 (m, 3H), 7.22 (m, 1H), 6.57 (d, *J* = 16.0 Hz, 1H), 6.40 (dd, *J* = 16.0, 8.0 Hz, 1H), 6.18 (m, 1H), 4.71 (d, *J* = 6.4 Hz, 2H), 4.11 (m, 1H), 2.46 (s, 3H)	3246, 1635, 1112, 801

Compound Number	mp (°C)	ESIMS	^{1}H NMR (δ)[a]	IR (cm^{-1})
AC16		506.93 ([M+H]$^+$)	8.47 (m, 1H), 8.19 (s, 1H), 7.76 (m, 1H), 7.47 (m, 2H), 7.37 (m, 1H), 7.28 (m, 2H), 7.24 (m, 1H), 7.21 (m, 1H), 6.59 (d, *J* = 16.0 Hz, 1H), 6.39 (dd, *J* = 16.0, 8.4 Hz, 1H), 4.12 (m, 1H), 2.48 (s, 3H), 1.88 (s, 6H)	1657, 1113, 801
AC17	70–73	494.98 ([M-H]$^-$)	7.49 (m, 2H), 7.38 (m, 1H), 7.29 (m, 4H), 7.08 (m, 3H), 6.91 (m, 1H), 6.61 (d, *J* = 16.0 Hz, 1H), 6.48 (m, 1H), 6.43 (dd, *J* = 16.0, 8.0 Hz, 1H), 4.13 (m, 1H), 2.49 (s, 3H)	
AC18	155–158	480.44 ([M+H]$^+$)	8.73 (d, *J* = 4.8 Hz, 2H), 7.53 (d, *J* = 8.4 Hz, 1H), 7.37 (m, 1H), 7.27 (m, 4H), 7.23 (m, 1H), 7.11 (m, 1H), 6.60 (d, *J* = 16.0 Hz, 1H), 6.41 (dd, *J* = 16.0, 8.0 Hz, 1H), 4.90 (d, *J* = 4.8 Hz, 2H), 4.13 (m, 1H), 2.52 (s, 3H)	
AC19	55–57	471.66 ([M+H]$^+$)	7.37 (m, 1H), 7.33 (d, *J* = 7.6 Hz, 1H), 7.27 (m, 2H), 7.22 (m, 2H), 6.57 (d, *J* = 16.0 Hz, 1H), 6.39 (dd, *J* = 16.0, 8.0 Hz, 1H), 6.10 (brs, 1H), 4.13 (m, 2H), 3.94 (m, 1H), 3.79 (m, 2H), 3.35 (m, 1H), 2.45 (s, 3H), 2.14 (m, 1H), 1.71 (m, 2H), 1.65 (m, 1H).	

Compound Number	mp (°C)	ESIMS	^{1}H NMR (δ)[a]	IR (cm^{-1})
AC20		467.68 ([M+H]$^+$)	7.37 (m, 2H), 7.27 (m, 2H), 7.23 (m, 2H), 6.57 (d, *J* = 16.0 Hz, 1H), 6.38 (m, 3H), 6.01 (m, 1H), 4.63 (d, *J* = 5.6 Hz, 2H), 4.13 (m, 1H), 2.45 (s, 3H)	3437, 1664, 1265, 1114, 746
AC21	61–64	528.78 ([M+H]$^+$)	8.44 (s, 1H), 8.18 (s, 1H), 7.83 (br s, 1H), 7.38 (m, 2H), 7.27 (m, 2H), 7.25 (m, 2H), 7.21 (m, 1H), 6.57 (d, *J* = 16.0 Hz, 1H), 6.40 (dd, *J* = 16.0, 8.0 Hz, 1H), 5.01 (s, 2H), 4.11 (m, 1H), 2.43 (s, 3H)	
AC22		545.08 ([M-H]$^-$)	8.39 (s, 1H), 7.73 (m, 1H), 7.40 (s, 1H), 7.35 (m, 2H), 7.22 (m, 3H), 6.57 (d, *J* = 16.0 Hz, 1H), 6.38 (dd, *J* = 16.0, 7.6 Hz, 1H), 6.14 (br s, 1H), 4.62 (d, *J* = 6.0 Hz, 2H), 4.13 (m, 1H), 2.45 (s, 3H)	3270, 1642, 1111, 809
AC23		492.35 ([M-H]$^-$)	7.42 (s, 2H), 7.36 (m, 1H), 7.24 (m, 2H), 6.59 (d, *J* = 16.0 Hz, 1H), 6.40 (dd, *J* = 16.0, 8.0 Hz, 1H), 6.20 (br s, 1H), 5.46 (m, 1H), 4.15 (m, 1H), 3.52 (m, 2H), 3.41 (m, 2H), 2.45 (s, 3H)	3273, 1641, 1250, 1113, 807
AC24	129–132	526.98 ([M+H]$^+$)	7.40 (m, 2H), 7.27 (m, 2H), 7.25 (m, 2H), 6.92 (br s, 2H), 6.60 (m, 1H), 6.48(dd, *J* = 16.0, 8.0 Hz, 1H), 4.19 (d, *J* = 5.2, 2H), 4.08 (m, 1H), 3.99 (m, 2H), 2.46 (s, 3H)	3298, 1664, 1113, 803

Compound Number	mp (°C)	ESIMS	^{1}H NMR (δ)[a]	IR (cm^{-1})
AC25		542.24 ([M-H]$^-$)	7.41 (m, 3H), 7.27 (m, 2H), 6.58 (d, *J* = 15.6 Hz, 1H), 6.42 (m, 2H), 4.92 (m, 1H), 4.65 (m, 2H), 4.14 (m, 1H), 4.09 (m, 2H), 2.46 (s, 3H)	3257, 1652, 1316, 1109, 807
AC26		550.69 ([M-H]$^-$)	7.45 (s, 1H), 7.40 (s, 2H), 7.34 (d, *J* = 8.0 Hz, 1H), 7.22 (m, 2H), 6.54 (d, *J* = 16.0 Hz, 1H), 6.38 (dd, *J* = 16.0, 8.0 Hz, 1H), 4.71 (d, *J* = 6.0 Hz, 2H), 4.11 (m, 1H), 2.46 (s, 3H)	3255, 1638, 1113, 809
AC27		541.00 ([M-H]$^-$)	8.46 (d, *J* = 4.0 Hz, 1H), 8.20 (s, 1H), 7.76 (m, 1H), 7.47 (m, 2H), 7.41 (s, 2H), 7.23 (m, 2H), 7.21 (m, 1H), 6.59 (d, *J* = 16.0 Hz, 1H), 6.37 (dd, *J* = 16.0, 8.4 Hz, 1H), 4.11 (m, 1H), 2.48 (s, 3H), 1.88 (s, 6H)	1653, 1113, 809
AC28	65–67	564.84 ([M-H]$^-$)	8.40 (s, 1H), 7.74 (m, 2H), 7.42 (m, 3H), 7.36 (m, 2H), 6.72 (br s, 1H), 6.52 (d, *J* = 16.0 Hz, 1H), 6.43 (dd, *J* = 16.0, 8.0 Hz, 1H), 4.66 (d, *J* = 6.4 Hz, 2H), 4.12 (m, 1H)	3267,1650, 1112, 809
AC29	75–78	511.78 ([M-H]$^-$)	7.71 (d, *J* = 8.4 Hz, 1H), 7.42 (m, 3H), 7.35 (m, 1H), 6.75 (br s, 1H), 6.56 (d, *J* = 16.0 Hz, 1H), 6.43 (dd, *J* = 16.0, 8.0 Hz, 1H), 5.49 (m, 1H), 4.14 (m, 1H), 3.50 (m, 4H)	

Compound Number	mp (°C)	ESIMS	^{1}H NMR (δ)[a]	IR (cm^{-1})
AC30	110–113	543.72 ([M-H]$^-$)	7.42 (d, *J* = 8.4 Hz, 1H), 7.44 (s, 1H), 7.40 (s, 1H), 7.38 (m, 1H), 7.06 (br s, 1H), 6.58 (d, *J* = 15.6 Hz, 1H), 6.45 (dd, *J* = 15.6, 8.0 Hz, 1H), 4.93 (m, 1H), 4.65 (m, 2H), 4.13 (m, 3H)	
AC31	68–70	610.73 ([M+H]$^+$)	8.42 (s, 1H), 7.76 (m, 1H), 7.61 (m, 2H), 7.39 (m, 4H), 6.54 -6.39 (m, 3H), 4.66 (d, *J* = 6.0 Hz, 2H), 4.12 (m, 1H)	
AC32	78–80	555.89 ([M-H]$^-$)	7.61 (m, 2H), 7.40 (m, 3H), 6.54 (m, 2H), 6.40 (dd, *J* = 16.0, 8.0 Hz, 1H), 5.46 (m, 1H), 4.14 (m, 1H), 3.50 (m, 4H)	
AC33	182–184	587.68 ([M-H]$^-$)	7.62 (s, 1H), 7.58 (d, *J* = 8.0 Hz, 1H), 7.40 (m, 3H), 6.84 (br s, 1H), 6.55 (d, *J* = 15.6 Hz, 1H), 6.45 (dd, *J* = 15.6, 7.6 Hz, 1H), 4.93 (m 1H), 4.65 (m, 2H), 4.13 (m, 4H)	
AC34	151–153	545.83 ([M-H]$^-$)	7.67 (s, 1H), 7.61 (d, *J* = 6.0 Hz, 1H), 7.53 (m, 1H), 7.41 (s, 2H), 6.64 (d, *J* = 16.0 Hz, 1H), 6.40 (dd, *J* = 16.0, 8.0 Hz, 1H), 6.18 (br s, 1H), 5.44 (m, 1H), 4.14 (m, 1H), 3.50 (m, 2H), 3.40 (m, 2H)	
AC35	100–102	577.71 ([M-H]$^-$)	7.70 (s, 1H), 7.63 (m, 1H), 7.53 (d, *J* = 7.6 Hz, 1H), 7.41 (s, 2H), 6.53 (d, *J* = 16.0 Hz, 1H), 6.49 (m, 2H), 4.93 (m, 1H), 4.64 (m, 2H), 4.13 (m, 1H), 4.03 (m, 2H)	3257, 1655, 1113, 808

Compound Number	mp (°C)	ESIMS	^{1}H NMR (δ)[a]	IR (cm^{-1})
AC36	81–83	600.83 ([M+H]$^+$)	8.40 (s, 1H), 7.73 (m, 2H), 7.61 (d, *J* = 8.4 Hz, 1H), 7.52 (d, *J* = 8.0 Hz, 1H), 7.40 (s, 2H), 7.35 (d, *J* = 8.0 Hz, 1H), 6.63 (d, *J* = 16.0 Hz, 1H), 6.46 (dd, *J* = 16.0, 7.6 Hz, 1H), 6.14 (m, 1H), 4.63 (d, *J* = 6.0 Hz, 2H), 4.14 (m, 1H)	
AC37		512.68 ([M+H]$^+$)	8.39 (s, 1H), 7.73 (m, 1H), 7.48 (m, 2H), 7.34 (d, *J* = 7.6 Hz, 1H), 7.24 (m, 3H), 6.55 (d, *J* = 16.0 Hz, 1H), 6.41 (dd, *J* = 16.0, 7.6 Hz, 1H), 6.12 (m, 1H), 4.62 (d, *J* = 6.0 Hz, 2H), 4.13 (m, 1H), 2.45 (s, 3H)	3268, 1644, 1109, 820
AC38	79–80	528.85 ([M-H]$^-$)	8.46 (m, 1H), 7.73 (m, 1H), 7.35 (m, 4H), 7.22 (m, 2H), 6.56 (d, *J* = 16.0 Hz, 1H), 6.38 (dd, *J* = 16.0, 8.0 Hz, 1H), 4.62 (d, *J* = 6.0 Hz, 2H), 4.10 (m, 1H), 2.45 (s, 3H)	
AC39	141–144	477.83 ([M-H]$^-$)	9.19 (s, 1H), 8.79 (s, 2H), 7.37 (m, 2H), 7.23 (m, 2H), 7.21 (m, 1H), 6.57 (d, *J* = 16.0 Hz, 1H), 6.40 (dd, *J* = 16.0, 7.6 Hz 1H), 6.21 (m, 1H), 4.65 (s, 2H), 4.11 (m, 1H), 2.46 (s, 3H)	

Compound Number	mp (°C)	ESIMS	^{1}H NMR (δ)[a]	IR (cm^{-1})
AC40	69-72	484.67 ([M+H]$^+$)	8.33 (t, *J* = 5.6 Hz, 1H), 8.61 (m, 1H), 7.68 (m, 3H), 7.48 (m, 2H), 6.86 (dd, *J* = 15.6, 8.2 Hz 1H), 6.74 (d, *J* = 15.6 Hz, 1H), 4.44 (m, 1H), 3.76 (d, *J* = 6.0 Hz, 2H), 2.54 (m, 1H), 2.67 (s, 3H), 0.59 (m, 2H), 0.54 (m, 2H)	
AC41	196-199	515.00 ([M-H]$^-$)	8.66 (d, *J* = 7.6 Hz, 1H), 8.39 (t, *J* = 5.6 Hz, 1H), 7.65 (s, 3H), 7.45 (m, 3H), 6.86 (dd, *J* = 15.6, 8.8 Hz, 1H), 6.74 (d, *J* = 15.6 Hz, 1H), 5.01 (m, 1H), 4.99 (m, 1H), 3.78 (d, *J* = 6.0 Hz, 2H), 3.40 (m, 2H), 3.22 (m, 2H), 2.37 (m, 3H)	
AC42	79-82	534.72 ([M+H]$^+$)	7.99 (d, *J* = 8.0 Hz, 1H), 7.89 (d, *J* = 8.0 Hz, 1H), 7.51 (m, 2H), 7.44 (m, 2H), 7.27 (m, 4H), 6.71 (t, *J* = 5.2 Hz, 1H), 6.59 (d, *J* = 16.0 Hz, 1H), 6.41 (dd, *J* = 16.0, 8.0 Hz, 1H), 5.05 (d, *J* = 1.6 Hz, 2H), 4.12 (m, 1H), 2.52 (m, 3H)	
AC43		481.75 ([M+H]$^+$)	8.69 (s, 1H), 8.52 (s, 2H), 7.45 (d, *J* = 7.6 Hz, 1H), 7.37 (d, *J* = 2.0 Hz, 1H), 7.26 (m, 2H), 7.21 (m, 1H), 6.83 (s, 1H), 6.58 (d, *J* = 16.0 Hz, 1H), 6.40 (dd, *J* = 16.0, 8.4 Hz, 1H), 4.81 (d, *J* = 5.6 Hz, 2H), 4.12 (t, *J* = 8.4 Hz 1H), 2.45 (s, 3H)	1663, 1608,1168, 1114, 801

Compound Number	mp (°C)	ESIMS	^{1}H NMR (δ)[a]	IR (cm^{-1})
AC44		528.01 ([M+H]$^+$)	8.44 (d, *J* = 2.4 Hz, 1H), 7.69 (d, *J* = 2.4 Hz, 1H), 7.37 (m, 1H), 7.33 (s, 1H), 7.31 (s, 1H), 7.26 (m, 1H), 7.24 (m, 3H), 6.57 (d, *J* = 16.0 Hz, 1H), 6.39 (dd, *J* = 16.0, 8.0 Hz, 1H), 5.96 (d, *J* = 7.2 Hz, 1H), 5.32 (t, *J* = 7.2 Hz, 1H), 4.11 (t, *J* = 8.4 Hz, 1H), 2.41 (s, 3H), 1.61 (d, *J* = 7.2 Hz, 3H)	1640, 1166, 1112, 800
AC45		512.88 ([M+H]$^+$)	7.66 (s, 1H), 7.37 (d, *J* = 6.8 Hz, 2H), 7.26 (m, 3H), 7.18 (m, 1H), 7.11 (m, 2H), 6.99 (m, 1H), 6.57 (d, *J* = 15.6 Hz, 1H), 6.39 (dd, *J* = 15.6, 8.0 Hz, 1H), 4.11 (t, *J* = 8.4 Hz, 1H), 3.36 (s, 3H), 2.43 (s, 3H)	1657, 1167, 1106, 800
AC46	61-64	575.93 ([M+H]$^+$)	8.42 (d, *J* = 2.0 Hz, 1H), 7.76 (d, *J* = 2.4 Hz, 1H), 7.61 (m, 2H), 7.39 (m, 3H), 7.26 (s, 2H), 6.54 (d, *J* = 16.0 Hz, 1H), 6.42 (dd, *J* = 16.0, 7.6 Hz, 1H), 4.65 (d, *J* = 6.0 Hz, 2H), 4.14 (m, 1H)	
AC47		525.89 ([M-H]$^-$)	10.02 (s, 1H), 9.87 (s, 1H), 8.47 (t, *J* = 6.0 Hz, 1H), 7.66 (s, 3H), 7.44 (s, 1H), 7.40 (d, *J* = 3.6 Hz, 2H), 6.86 (dd, *J* = 15.6, 9.2 Hz, 1H), 6.74 (d, *J* = 15.6 Hz, 1H), 4.82 (t, *J* = 9.6 Hz, 2H), 3.88 (d, *J* = 6.0 Hz, 2H), 2.36 (s, 3H), 1.63 (m, 1H), 0.76 (m, 4H)	3280, 1640

Compound Number	mp (°C)	ESIMS	^{1}H NMR (δ)[a]	IR (cm^{-1})
AC48		509.96 ([M-H]$^-$)	7.37 (m, 7H), 7.34 (m, 3H), , 6.57 (d, *J* = 16.0 Hz, 1H), 6.39 (dd, *J* = 16.0, 8.0 Hz, 1H), 6.01 (m, 1H), 4.60 (d, *J* = 6.0 Hz, 2H), 4.13 (m, 1H), 2.46 (s, 3H)	3275, 1642
AC49		518.85 ([M+H]$^+$)	8.39 (d, *J* = 2.0 Hz, 1H), 8.11 (m, 1H), 7.71 (d, *J* = 2.4 Hz, 1H), 7.41 (m, 3H), 7.17 (m, 3H), 6.59 (d, *J* = 16.0 Hz, 1H), 6.47 (dd, *J* = 16.0, 8.0 Hz, 1H), 4.66 (d, *J* = 5.6 Hz, 2H), 4.14 (m, 1H)	1658, 1112, 1025, 2219
AC50		481.88 ([M+H]$^+$)	8.72 (m, 1H), 7.67 (s, 3H), 7.46 (s, 1H), 7.40 (m, 2H), 7.08 (s, 1H), 6.82 (m, 2H), 6.55 (d, *J* = 7.6 Hz, 1H), 4.82 (m, 1H), 4.48 (s, 2H), 3.65 (s, 3H), 2.38 (s, 3H)	1654, 1112, 800, 3069
AC51		540.83 ([M+H]$^+$)	7.45 (d, *J* = 7.6 Hz, 1H), 7.38 (m, 1H), 7.27 (m, 2H), 7.22 (m, 2H), 6.85 (m, 1H), 6.58 (d, *J* = 16.0 Hz, 1H), 6.40 (dd, *J* = 16.0, 8.0 Hz, 1H), 4.33 (m, 2H), 4.14 (m, 3H), 3.18 (s, 3H), 2.48 (s, 3H)	1652, 1571, 802, 1114, 2926
AC52		488.29 ([M-H]$^-$)	7.33 (m, 2H), 7.25 (m, 3H), 6.56 (d, *J* = 15.6 Hz, 1H), 6.37 (dd, *J* = 15.6, 8.0 Hz, 1H), 5.61 (d, *J* = 8.0 Hz, 1H), 4.21 (m, 1H), 4.01 (m, 1H), 4.08 (m, 2H), 3.56 (t, *J* = 10.0 Hz, 2H), 2.48 (m, 2H), 2.08 (m, 2H), 1.5 (m, 3H)	1635, 11134, 813, 2927

Compound Number	mp (°C)	ESIMS	^{1}H NMR (δ)[a]	IR (cm^{-1})
AC53		532.92 ([M+H]$^+$)	8.49 (d, *J* = 2.0 Hz, 1H), 7.69 (d, *J* = 2.4 Hz, 1H), 7.43 (d, *J* = 8.0 Hz, 1H), 7.34 (m, 3H), 7.26 (m, 2H), 6.95 (m, 1H), 6.58 (d, *J* = 16.0 Hz, 1H), 6.38 (dd, *J* = 16.0, 8.0 Hz, 1H), 4.72 (d, *J* = 5.2 Hz, 2H), 4.09 (m, 1H), 2.47 (s, 3H)	1651, 3027, 815, 1113
AC54		529.06 ([M-H]$^-$)	8.37 (d, *J* = 5.2 Hz, 1H), 7.41 (d, *J* = 8.0 Hz, 1H), 7.36 (m, 3H), 7.31 (m, 1H), 7.26 (m, 2H), 6.58 (d, *J* = 16.0 Hz, 1H), 6.40 (dd, *J* = 16.0, 7.6 Hz, 1H), 5.20 (t, *J* = 5.6 Hz, 1H), 4.63 (d, *J* = 6.0 Hz, 2H), 4.13 (m, 1H), 2.18 (s, 3H)	1654, 3434, 814, 1112
AC57		464.96 ([M+H]$^+$)	8.69 (t, *J* = 6.0 Hz, 1H), 8.58 (t, *J* = 6.0 Hz, 1H), 7.92 (s, 1H), 7.87 (d, *J* = 6.4 Hz, 2H), 7.62 (d, *J* = 8.4 Hz, 1H), 7.45 (d, *J* = 8.4 Hz, 1H), 7.0 (m, 1H), 6.76 (d, *J* = 15.6 Hz, 1H), 6.76 (dd, *J* = 15.6, 8.0 Hz, 1H), 4.01 (m, *J* = 8.0 Hz, 1H), 3.71 (m, 2H), 3.49 (m, 2H)	3417, 1658, 1165, 817
AC58	124.4-126.9	599.76 ([M+H]$^+$)	7.62 (m, 2H), 7.40 (s, 2H), 7.37 (d, *J* = 1.6 Hz, 1H), 6.61 (t, *J* = 4.8 Hz, 1H), 6.55 (d, *J* = 16.0 Hz, 1H), 6.41 (dd, *J* = 16.0, 7.6 Hz, 1H), 4.16 (d, *J* = 6.0 Hz, 2H), 4.01 (m, 1H), 1.56 (s, 9H)	

Compound Number	mp (°C)	ESIMS	^{1}H NMR (δ)[a]	IR (cm^{-1})
AC59	80-83	497.40 ([M-H]$^-$)	8.42 (d, *J* = 2.1 Hz, 1H), 8.29 (d, *J* = 7.5 Hz, 1H), 7.51 (m, 2H), 7.39 (m, 1H), 7.36 (m, 4H), 7.28 (m, 1H), 6.61 (d, *J* = 15.9 Hz, 1H), 6.45 (dd, *J* = 15.9, 7.8 Hz 1H), 4.14 (t, *J* = 8.4 Hz, 1H), 2.51 (s, 3H)	
AC60		515.09 ([M+H]$^-$)	8.52 (s, 1H), 8.39 (d, *J* = 1.8 Hz, 2H), 7.70 (d, *J* = 2.1 Hz, 1H), 7.62 (s, 1H), 7.43 (s, 1H), 7.35 (m, 3H), 6.62 (d, *J* = 16.2 Hz, 1H), 6.52 (dd, *J* = 16.2, 7.5 Hz, 1H), 4.62 (d, *J* = 6.3 Hz, 2H), 4.19 (m, 1H), 2.76 (s, 3H)	1668, 1589, 1167, 1113, 802
AC61		461.90 ([M-H]$^-$)	8.07 (t, *J* = 8.0 Hz, 1H), 7.39 (t, *J* = 2.0 Hz, 1H), 7.28 (d, *J* = 1.2 Hz, 3H), 7.17 (d, *J* = 1.6 Hz, 1H), 7.11 (m, 1H), 6.59 (d, *J* = 15.6 Hz, 1H), 6.47 (dd, *J* = 15.6, 7.6 Hz, 1H), 5.49 (m, 1H), 4.14 (t, *J* = 8.4 Hz, 1H), 3.48 (m, 4H)	1658, 1114, 801
AC62	105-108	528.88 ([M-H]$^-$)	8.62 (t, *J* = 6.4 Hz, 1H), 8.46 (m, 1H), 7.73 (m, 5H), 7.48 (d, *J* = 7.6 Hz, 1H), 7.03 (dd, *J* = 15.6, 9.2 Hz, 1H), 6.81 (d, *J* = 15.6 Hz, 1H), 4.86 (m, 1H), 3.97 (m, 4H)	

Compound Number	mp (°C)	ESIMS	^{1}H NMR (δ)[a]	IR (cm^{-1})
AC63	77-80	594.67 ([M+H]$^+$)	8.43 (s, 1H), 7.76 (d, *J* = 2.4 Hz, 1H), 7.60 (m, 2H), 7.38 (d, *J* = 7.6 Hz, 1H), 7.33 (d, *J* = 6.4 Hz, 3H), 6.54 (d, *J* = 16.0 Hz, 1H), 6.46 (m, 1H), 6.41 (dd, *J* = 16.0 8.0 Hz, 1H), 4.65 (d, *J* = 6.0 Hz, 2H), 4.15 (m, 1H)	3257, 1653
AC64	83-85	580.72 ([M-H]$^-$)	7.72 (d, *J* = 8.0 Hz, 1H), 7.44 (s, 1H), 7.40 (s, 2H), 7.36 (d, *J* = 6.8 Hz, 1H), 7.05 (t, *J* = 5.2 Hz, 1H), 6.70 (t, *J* = 5.2 Hz, 1H), 6.57 (d, *J* = 15.6 Hz, 1H), 6.44 (dd, *J* = 15.6, 8.0 Hz, 1H), 4.23 (d, *J* = 5.6 Hz, 2H), 4.15 (m, 1H), 4.01 (m, 2H)	
AC65		534.72 ([M-H]$^-$)	8.39 (d, *J* = 2.0 Hz, 1H), 8.12 (t, *J* = 8.4 Hz, 1H), 7.71 (d, *J* = 2.4 Hz, 1H), 7.34 (m, 3H), 7.26 (m, 1H), 7.11 (m, 2H), 6.59 (d, *J* = 16.0 Hz, 1H), 6.46 (dd, *J* = 16.0, 8.0 Hz, 1H), 4.66 (d, *J* = 5.2 Hz, 2H), 4.13 (m, 1H)	1658, 1113, 817, 2925
AC66	73-75	624.61 ([M-H]$^-$)	7.88 (s, 1H), 7.63 (d, *J* = 1.6 Hz, 1H), 7.57 (d, *J* = 8.0 Hz, 1H), 7.40 (m, 2H), 6.80 (t, *J* = 5.6 Hz, 1H), 6.70 (t, *J* = 5.6 Hz, 1H), 6.56 (d, *J* = 16.0 Hz, 1H), 6.44 (dd, *J* = 16.0, 8.0 Hz, 1H), 4.22 (m, 2H), 4.12 (m, 1H), 4.01 (m, 2H)	

Compound Number	mp (°C)	ESIMS	^{1}H NMR (δ)[a]	IR (cm^{-1})
AC67		479.82 ([M-H]$^-$)	8.07 (t, *J* = 8.0 Hz, 1H), 7.34 (d, *J* = 6.0 Hz, 2H), 7.28 (s, 1H), 7.17(s, 2H), 6.59 (d, *J* = 15.6 Hz, 1H), 6.46 (dd, *J* = 15.6, 8.0 Hz, 1H), 5.49 (m, 1H), , 4.12 (m, 1H), 3.49 (m, 4H).	3272, 1644
AC68	90-93	546.80 ([M-H]$^-$)	8.6 (t, *J* = 6.4 Hz, 1H), 8.45 (m, 1H), 7.86 (d, *J* = 6.4 Hz, 2H), 7.75 (t, *J* = 8.0 Hz, 1H), 7.63 (d, *J* = 12.0 Hz, 1H), 7.48 (d, *J* = 8.0 Hz, 1H), 7.03 (dd, *J* = 15.6, 9.6 Hz, 1H), 6.80 (d, *J* = 15.6 Hz, 1H), 4.88 (m, 1H), 3.96 (m, 4H)	3315, 1684
AC69		542.82 ([M-H]$^-$)	7.41 (d, *J* = 8.0 Hz, 1H), 7.34 (d, *J* = 5.6 Hz, 2H), 7.26 (m, 1H), 7.23 (m, 1H), 6.81 (s, 1H), 6.57 (d, *J* = 15.6 Hz, 1H), 6.55 (s, 1H), 6.39 (dd, *J* = 15.6, 8.0 Hz, 1H), 4.18 (m, 2H), 4.13 (m, 1H), 3.97 (m, 2H), 2.46 (s, 3H)	3294, 1685
AC70	176-178	545.23 ([M-H]$^-$)	8.38 (d, *J* = 2.4 Hz, 1H), 8.22 (d, *J* = 6.8 Hz, 2H), 7.71 (d, *J* = 2.4 Hz, 1H), 7.35 (d, *J* = 6.0 Hz, 2H), 7.30 (d, *J* = 7.6 Hz, 1H), 7.15 (d, *J* = 1.6 Hz, 1H), 6.93 (d, *J* = 1.2 Hz, 1H), 6.60 (d, *J* = 15.6 Hz, 1H), 6.43 (dd, *J* = 15.6, 7.6 Hz, 1H), 4.66 (d, *J* = 6.0 Hz, 2H), 4.13 (m, 1H), 3.98 (s, 3H)	

Compound Number	mp (°C)	ESIMS	^{1}H NMR (δ)[a]	IR (cm^{-1})
AC71		492.20 ([M-H]$^-$)	8.24 (d, *J* = 7.6 Hz, 1H), 8.15 (d, *J* = 8.4 Hz, 1H), 7.35 (d, *J* = 6.0 Hz, 2H), 7.13 (d, *J* = 1.2 Hz, 1H), 6.92 (s, 1H), 6.61 (d, *J* = 16.0 Hz, 1H), 6.43 (dd, *J* = 16.0, 7.6 Hz, 1H), 5.48 (m, 1H), 4.13 (m, 1H), 4.03 (s, 3H), 3.48 (m, 4H)	1639, 3079, 858
AC72		543.05 ([M-H]$^-$)	8.42 (d, *J* = 2.4 Hz, 1H), 7.75 (d, *J* = 2.4 Hz, 1H), 7.34 (m, 4H), 7.20 (m, 2H), 6.60 (d, *J* = 16.0 Hz, 1H), 6.36 (dd, *J* = 16.0, 8.0 Hz, 1H), 6.12 (t, *J* = 5.6 Hz, 1H), 4.62 (d, *J* = 6.0 Hz, 2H), 4.20 (m, 1H), 2.82 (m, 2H), 1.45 (t, *J* = 5.6 Hz, 3H)	1642, 3246, 814, 1113
AC75		644.78 ([M+H]$^+$)	8.72 (s, 1H), 7.97 (d, *J* = 7.2 Hz, 1H), 7.70 (d, *J* = 8.4 Hz, 1H), 7.61 (m, 2H), 7.40 (m, 2H), 6.55 (m, 2H), 6.42 (dd, *J* = 16.0, 8.0 Hz, 1H), 4.76 (d, *J* = 6.0 Hz, 2H), 4.12 (m, 1H)	3431, 1652, 1171, 809
AC76		531.34 ([M+H]$^+$)	8.87 (t, *J* = 6.0 Hz, 1H), 8.34 (d, *J* = 2.1 Hz, 1H), 7.85 (d, *J* = 6.3 Hz, 3H), 7.48 (m, 4H), 6.57 (d, *J* = 15.6 Hz, 1H), 6.45 (dd, *J* = 15.6, 9.0 Hz, 1H), 4.84 (m, 1H), 4.49 (d, *J* = 5.7 Hz, 2H), 2.82 (m, 2H), 2.36 (t, *J* = 5.6 Hz , 3H)	3120, 1708, 1171

Compound Number	mp (°C)	ESIMS	^{1}H NMR (δ)[a]	IR (cm^{-1})
AC77		531.1 ([M+H]$^+$)	8.87 (t, *J* = 6.0 Hz, 1H), 8.34 (d, *J* = 2.1 Hz, 1H), 7.85 (d, *J* = 6.3 Hz, 3H), 7.48 (m, 4H), 6.57 (d, *J* = 15.6 Hz, 1H), 6.45 (dd, *J* = 15.6, 8.0 Hz, 1H), 4.84 (m, 1H), 4.49 (d, *J* = 5.7 Hz, 2H), 2.36 (s, 3H)	3444, 1648, 1114, 814
AC78		561.06 ([M+H]$^+$)	8.59 (t, *J* = 6.4 Hz, 1H), 8.47 (t, *J* = 5.6 Hz, 1H), 7.89 (s, 2H), 7.45 (m, 3H), 6.87 (m, 1H), 6.75 (d, *J* = 15.6 Hz, 1H), 4.85 (t, *J* = 8.0 Hz 1H), 3.98 (m, 4H), 2.58 (s, 3H)	3432, 1631, 1161, 840
AC79		610.97 ([M+H]$^+$)	8.69 (t, *J* = 6.0 Hz, 1H), 8.58 (t, *J* = 6.0 Hz, 1H), 7.92 (s, 1H), 7.87 (d, *J* = 6.4 Hz, 2H), 7.62 (d, *J* = 8.4 Hz, 1H), 7.45 (d, *J* = 8.4 Hz, 1H), 7.0 (m, 1H), 6.76 (d, *J* = 15.6 Hz, 1H) 4.83 (t, *J* = 8.0 Hz, 1H), 3.98 (m, 4H)	3303, 1658, 1166, 817
AC80		561.06 ([M+H]$^+$)	7.37 (m, 3H), 7.26 (m, 1H), 7.24 (m, 1H), 6.59 (d, *J* = 15.6 Hz, 1H), 6.39 (dd, *J* = 15.6, 8.0 Hz, 1H), 4.24 (m, 4H), 3.90 (m, 1H), 2.83 (m, 2H), 1.26 (m, 3H)	3412, 1624, 1157, 825
AC81	9-92	546.93 ([M-H]$^-$)	8.73 (d, *J* = 5.6 Hz, 1H), 8.45 (t, *J* = 6.0 Hz, 1H), 7.76 (s, 3H), 7.45 (m, 3H), 6.86 (dd, *J* = 16.0, 9.2 Hz, 1H), 4.83 (m, 1H), 4.56 (m, 2H), 4.51 (m, 1H), 4.10 (m, 2H), 3.85 (d, *J* = 6.0 Hz, 2H), 2.50 (m, 3H)	

Compound Number	mp (°C)	ESIMS	^{1}H NMR (δ)[a]	IR (cm^{-1})
AC82		477.69 ([M+H]$^+$)	7.38 (d, *J* = 1.8 Hz, 2H), 7.33 (s, 1H), 7.27 (s, 3H), 6.58 (d, *J* = 16.0 Hz, 1H), 6.42 (d, *J* = 8.1 Hz, 1H), 6.36 (dd, *J* = 16.0, 7.8 Hz, 1H), 4.71 (m, 1H), 4.23 (m, 3H), 3.26 (m, 2H), 2.45 (s, 3H)	1646, 1353, 1196, 1112, 800
AC83		493.83 ([M-H]$^-$)	8.07 (t, *J* = 8.4 Hz, 1H), 7.39 (t, *J* = 1.6 Hz, 1H), 7.31 (d, *J* = 1.2 Hz, 1H), 7.26 (m, 2H), 7.23 (m, 1H), 7.19 (d, *J* = 1.6 Hz, 1H), 6.60 (d, *J* = 16.8 Hz, 1H), 6.49 (dd, *J* = 16.8, 7.6 Hz, 1H), 4.90 (m, 1H), 4.64 (m, 2H), 4.14 (m, 2H), 4.10 (m, 1H)	1527, 1113, 801, 1167, 1321
AC84		511.75 ([M-H]$^-$)	8.07 (t, *J* = 8.0 Hz, 1H), 7.34 (m, 3H), 7.19 (d, *J* = 13.2 Hz, 1H), 6.60 (d, *J* = 16.4 Hz, 1H), 6.48 (dd, *J* = 16.4, 8.0 Hz, 1H), 4.88 (m, 1H), 4.62 (m, 2H), 4.12 (m, 3H)	1645, 1113, 804, 3030, 1245
AC85		523.83 ([M-H]$^-$)	8.60 (d, *J* = 6.8 Hz, 1H), 8.15 (d, *J* = 8.4 Hz, 1H), 7.35 (d, *J* = 6.0 Hz, 1H), 7.15 (d, *J* = 7.2 Hz, 1H), 6.94 (s, 1H), 6.60 (d, *J* = 15.6 Hz, 1H), 6.44 (dd, *J* = 7.6, 7.6 Hz, 1H), 4.93 (m, 1H), 4.62 (m, 2H), 4.13 (m, 6H)	1652, 3039, 802, 1114

Compound Number	mp (°C)	ESIMS	^{1}H NMR (δ)[a]	IR (cm^{-1})
AC86		524.36 ([M+H]$^+$)	7.35 (d, *J* = 6.3 Hz, 3H), 7.26 (m, 2H), 7.20 (m, 1H), 6.60 (d, *J* = 15.9 Hz, 1H), 6.47 (dd, *J* = 15.9, 6.6 Hz, 1H), 4.86 (m, 1H), 4.65 (m, 2H), 4.13 (m, 3H), 2.84 (q, 2.8 Hz, 2H), 1.26 (m, 3H)	3333, 1651, 815
AC87		495.82 ([M-H]$^-$)	8.07 (t, *J* = 8.0 Hz, 1H), 7.52 (m, 3H), 7.19 (d, *J* = 13.2 Hz, 1H), 6.59 (d, *J* = 16.4 Hz, 1H), 6.47 (dd, *J* = 16.4, 8.0 Hz, 1H), 4.69 (m, 1H), 4.23 (m, 3H), 3.29 (m, 2H)	1623, 1114, 816
AC89		509.89 ([M+H]$^+$)	7.43 (m, 2H), 7.27 (m, 2H), 7.23 (m, 2H), 6.58 (d, *J* = 16.0 Hz, 1H), 6.41 (dd, *J* = 16.0, 7.6 Hz, 1H), 4.79 (d, *J* = 5.6 Hz, 2H), 4.14 (m, 1H), 2.48 (s, 3H), 2.18 (m, 1H), 1.16 (m, 4H)	1666, 1166, 1112, 800
AC90		656.9 ([M-H]$^-$)	8.34 (m, 1H), 8.27 (m, 1H), 7.60 (d, *J* = 1.6 Hz, 1H), 7.49 (d, *J* = 8.0 Hz, 2H), 7.40 (s, 2H), 7.36 (dd, *J* = 8.2, 1.7 Hz, 1H), 6.53 (d, *J* = 16.0 Hz, 1H), 6.38 (dd, *J* = 15.9, 7.9 Hz, 1H), 4.89 (d, *J* = 8.4 Hz, 2H), 4.48 (d, *J* = 9.0 Hz, 2H), 4.11 (m, 1H)	

Compound Number	mp (°C)	ESIMS	^{1}H NMR (δ)[a]	IR (cm^{-1})
AC91		640.9 ([M-H]$^-$)	8.18 (t, *J* = 5.0 Hz, 1H), 7.58 (d, *J* = 1.6 Hz, 1H), 7.47 (d, *J* = 8.0 Hz, 1H), 7.40 (s, 2H), 7.34 (dd, *J* = 8.1, 1.6 Hz, 1H), 6.52 (m, 2H), 6.37 (dd, *J* = 15.9, 7.9 Hz, 1H), 4.54 (d, *J* = 4.9 Hz, 2H), 4.12 (m, 1H), 3.99 (qd, *J* = 8.9, 6.5 Hz, 2H)	
AC92		640.9 ([M-H]$^-$)	9.16 (d, *J* = 6.1 Hz, 1H), 7.65 (d, *J* = 1.6 Hz, 1H), 7.57 (d, *J* = 8.0 Hz, 1H), 7.41 (m, 3H), 7.21 (t, *J* = 5.6 Hz, 1H), 6.55 (d, *J* = 15.9 Hz, 1H), 6.41 (dd, *J* = 15.9, 7.8 Hz, 1H), 4.59 (d, *J* = 5.6 Hz, 2H), 4.45 (qd, *J* = 9.0, 6.0 Hz, 2H), 4.12 (q, *J* = 7.2 Hz, 1H)	
AC93		485.5 ([M+H]$^+$)	7.52-7.41 (d, *J* = 8.2 Hz, 1H), 7.39-7.34 (m, 1H), 7.24-7.17 (d, *J* = 1.8 Hz, 2H), 7.02-6.92 (m, 2H), 6.90-6.83 (d, *J* = 11.4 Hz, 1H), 6.71 (br s, 1H), 6.17 (br s, 1H), 6.12-6.01 (dd, *J* = 11.4, 10.3 Hz, 1H), 4.44-4.38 (d, *J* = 4.2 Hz, 1H), 4.35-4.27 (m, 1H), 4.10-3.99 (d, *J* = 5.1 Hz, 2H), 2.78-2.67 (m, 1H), 2.44 (s, 3H), 0.88-0.78 (m, 2H), 0.60-0.45 (m, 2H)	^{13}C NMR (δ)[3] 169.91, 169.84, 138.23, 137.41, 136.84, 134.79, 134.69, 131.07, 128.69, 127.49, 127.43, 126.72, 126.61 (q, *J* = 212.10 Hz), 125.61, 123.76, 47.89 (q, *J* = 28.28 Hz), 43.46, 22.65, 19.97, 8.21

Compound Number	mp (°C)	ESIMS	^{1}H NMR (δ)[a]	IR (cm^{-1})
AC94		511.6 ([M]$^-$)	8.36 - 8.24 (d, *J* = 2.4 Hz, 1H), 7.75 - 7.64 (m, 1H), 7.38 - 7.24 (m, 3H), 7.24 - 7.09 (d, *J* = 1.8 Hz, 2H), 6.99 - 6.90 (m, 2H), 6.89 - 6.74 (d, *J* = 11.4 Hz, 1H), 6.63 - 6.43 (m, 1H), 6.14 - 5.98 (m, 1H), 4.69 - 4.51 (d, *J* = 6.1 Hz, 2H), 4.37 - 4.20 (m, 1H), 2.46 - 2.31 (s, 3H)	3262, 1607, 1247, 1164, 1111
AC95	48-61	626.9 ([M+H]$^+$)	7.58 (d, *J* = 7.9 Hz, 1H), 7.44 - 7.29 (m, 3H), 7.14 (dd, *J* = 7.9, 1.6 Hz, 1H), 6.86 (d, *J* = 11.4 Hz, 1H), 6.76 (t, *J* = 5.9 Hz, 1H), 6.59 (br s, 1H), 6.21 - 6.04 (m, 1H), 4.23 (d, *J* = 5.5 Hz, 1H), 3.98 (qd, *J* = 9.0, 6.5 Hz, 2H)	
AC96		619.6 ([M+H]$^+$)	8.83 (s, 1H), 8.06 (br, 1H), 7.90 (s, 2H), 7.63 (d, *J* = 8.1 Hz, 2H), 7.53 (m, 1H), 6.94 (m, 1H), 6.77 (d, *J* = 15.3 Hz, 1H), 6.63 (d, *J* = 9.3 Hz, 1H), 4.84 (m, 1H), 4.30 (d, *J* = 5.6 Hz, 2H), 2.99 (s, 6H)	1616, 1114
AC97		606.6 ([M+H]$^+$)	8.20 (d, *J* = 2.1 Hz, 1H), 7.73 (d, *J* = 2.7 Hz, 1H), 7.60 (m, 2H), 7.39 (s, 2H), 7.29 (m, 1H), 6.79 (d, *J* = 8.4 Hz, 1H), 6.55 (d, *J* = 15.9 Hz, 1H), 6.40 (m, 2H), 4.60 (d, *J* = 2.7 Hz, 2H), 4.13 (m, 1H), 3.95 (s, 3H)	1644, 1113

Compound Number	mp (°C)	ESIMS	^{1}H NMR (δ)[a]	IR (cm^{-1})
AC98		577.87 ([M+H]$^+$)	9.04 (t, *J* = 6.0 Hz, 1H), 8.60 (t, *J* = 6.6 Hz, 1H), 8.25 (s, 1H), 7.97 (d, *J* = 8.1 Hz, 1H), 7.87 (d, *J* = 6.3 Hz, 2H), 7.69 (d, *J* = 7.5 Hz, 1H), 7.15 (dd, *J* = 15.9, 9.3 Hz, 1H), 6.89 (d, *J* = 15.9 Hz, 1H), 4.86 (m, 1H), 3.98 (m, 4H).	1663, 1168
AC99		574.81 ([M+H]$^+$)	8.69 (t, *J* = 6.0 Hz, 1H), 8.58 (t, *J* = 6.6 Hz, 1H), 7.91 (s, 1H), 7.85 (m, 1H), 7.61 (m, 2H), 7.52 (m, 2H), 6.98 (dd, *J* = 15.3, 9.0 Hz, 1H), 6.76 (d, *J* = 15.3 Hz, 1H), 4.81 (m, 1H), 4.01 (m, 4H)	1650, 1164
AC100		673.80 ([M+H]$^+$)	8.29 (s, 1H), 8.22 (d, *J* = 8.1 Hz, 1H), 7.93 (d, *J* = 7.8 Hz, 1H), 7.72 (m, 1H), 7.65 (m, 2H), 7.40 (s, 2H), 7.18 (br, 1H), 6.59 (d, *J* = 16.0 Hz, 1H), 6.43 (dd, *J* = 16.0, 7.6 Hz, 1H), 5.02 (d, *J* = 1.2 Hz, 2H), 4.12 (m, 1H)	3403, 1659
AC101		636.83 ([M+H]$^+$)	7.56 (d, *J* = 9.0 Hz, 1H), 7.39 (d, *J* = 6.0 Hz, 2H), 7.26 (m, 2H), 6.54 (d, *J* = 15.9 Hz, 1H), 6.37 (dd, *J* = 8.0, 15.9 Hz, 1H), 4.01 (m, 1H), 3.84 (m, 2H), 3.33 (m, 2H), 3.04 (m, 2H), 2.84 (m, 3H), 2.62 (m, 1H)	1637, 1113

Compound Number	mp (°C)	ESIMS	^{1}H NMR (δ)[a]	IR (cm^{-1})
AC102		592.84 ($[M+H]^+$)	7.60 (m, 2H), 7.32 (m, 1H), 7.03 (d, *J* = 7.2 Hz, 2H), 6.74 (br, 1H), 6.62 (br, 1H), 6.56 (d, *J* = 16.2 Hz, 1H), 6.41 (dd, *J* = 16.2, 7.8 Hz, 1H), 4.22 (d, *J* = 5.4 Hz, 2H), 4.14 (m, 1H), 4.01 (m, 2H)	1668, 1167
AC103	99.2-105.0	612.7 ($[M+H]^+$)	8.40 (d, *J* = 8.0 Hz, 1H), 7.92 (d, *J* = 5.2 Hz, 1H), 7.59 (d, *J* = 8.0 Hz, 1H), 7.35 (d, *J* = 8.0 Hz, 1H), 6.99 (dd, *J* = 16.0, 7.6 Hz, 1H), 6.76 (d, *J* = 16.0 Hz, 1H), 4.84 (m, 1H), 4.23 (d, *J* = 13.2 Hz, 1H), 3.97 (m, 1H), 3.79 (d, *J* = 13.6 Hz, 1H), 3.16 (t, *J* = 11.2 Hz, 1H), 2.77 (t, *J* = 11.2 Hz, 1H), 1.99 (s, 3H), 1.88 (m, 2H), 1.45 (m, 2H)	1634, 1113, 809
AC104		680.97 ($[M+H]^+$)	7.60 (m, 2H), 7.40 (m 3H), 6.55 (d, *J* = 15.6 Hz, 1H), 6.41 (dd, *J* = 15.6, 7.8 Hz, 1H), 4.24 (m, 1H), 3.34 (m, 2H), 2.90 (m, 1H), 2.24 (m, 2H), 1.52(m, 2H), 1.34 (m, 4H)	3437, 1644, 1113, 807, 511
AC105		609.9 ($[M+H]^+$)	7.59 (s, 1H), 7.55 (m, 1H), 7.50 (m, 1H),7.40 (m, 2H), 6.54(d, *J* = 16.0 Hz, 1H), 6.50 (*J* = 16.0, 8.0 Hz, 1H), 4.14 (m, 2H), 3.08 (m, 4H), 2.67 (m, 2H), 2.12 (m, 2H), 1.70 (m, 2H).	3303, 1649, 1115, 2242, 809, 506

Compound Number	mp (°C)	ESIMS	^{1}H NMR (δ)[a]	IR (cm^{-1})
AC106		584.95 ([M+H]$^+$)	7.59 (s, 1H), 7.51 (d, *J* = 8.4 Hz, 1H), 7.40 (s, 2H), 7.36 (d, *J* = 6.8 Hz, 1H), 6.54 (d, *J* = 16.0 Hz, 1H), 6 .40 (dd, *J* = 16.0, 8.0 Hz, 1H), 6.03 (d, *J* = 8.0 Hz, 1H), 4.11 (m, 2H), 3.10 (m, 2H), 2.50 (m, 2H), 2.50 (s, 3H) (m, 2H), 1.94 (m, 2H)	3417, 1648, 1112, 805, 555
AC107		609.9 ([M+H]$^+$)	8.41 (d, *J* = 7.8 Hz, 1H), 7.90 (s, 2H), 7.62 (m, 2H), 7.51(m, 1H), 6.92 (dd, *J* = 15.9, 9.0 Hz, 1H), 6.77 (d, *J* = 15.9 Hz, 1H), 4.81 (m, 1H), 3.73 (s, 2H), 3.31 (m, 1H), 3.28 (m, 1H), 2.82 (t, *J* = 11.4 Hz, 2H), 2.82 (m, 2H), 2.30 (m, 2H), 1.88 (m, 2H), 1.57 (m, 2H)	3303, 1645, 1115, 2243, 810, 507
AC108		626.9 ([M+H]$^+$)	7.60 (m, 2H) 7.39 (s, 2H), 7.28 (m, 1H), 6.56 (d, *J* = 15.6 Hz, 1H), 6.40 (dd, *J* = 15.6, 7.8 Hz, 1H), 5.91 (m, 1H), 4.65 (m, 2H), 4.10 (m, 1H), 4.07 (m, 2H), 3.59 (m, 1H), 2.74 (m, 2H), 2.13 (m, 4H), 2.07 (m, 1H)	3420, 1649, 1113, 809, 554
AC109		614.6 ([M+H]$^+$)	7.56 (m, 2H), 7.39 (s, 2H), 7.29 (s, 1H), 6.50 (d, *J* = 15.9 Hz, 1H), 6.41 (dd, *J* = 15.9, 8.0 Hz 1H), 4.09 (m, 1H), 3.88 (m, 2H), 3.49 (m, 2H), 2.92 (m, 2H), 2.81 (m, 1H), 2.74 (m, 2H), 2.25 (m, 4H)	1647, 1113

Compound Number	mp (°C)	ESIMS	^{1}H NMR (δ)[a]	IR (cm^{-1})
AC110		572.6 ([M+H]$^+$)	11.20 (s, 1H), 8.66 (br, 1H), 7.92 (m, 3H), 7.62 (d, *J* = 8.0 Hz, 1H), 7.45 (d, *J* = 8.0 Hz, 1H), 6.77 (dd, *J* = 15.6, 9.2 Hz, 1H), 6.77 (d, *J* = 15.6 Hz, 1H), 4.85 (m, 1H), 3.74 (d, *J* = 5.2 Hz, 2H), 3.61 (s, 3H)	3412, 1690, 1114, 846, 559
AC111		582.79 ([M+H]$^+$)	8.63 (t, *J* = 6.0 Hz, 1H), 8.04 (t, *J* = 6.0 Hz, 1H), 7.92 (m, 3H), 7.62 (d, *J* = 1.2 Hz, 1H), 7.47 (d, *J* = 7.6 Hz, 1H), 7.00 (dd, *J* = 15.6, 8.8 Hz, 1H), 6.77 (d, *J* = 15.6 Hz, 1H), 5.19 (d, *J* = 1.6 Hz, 1H), 5.01 (d, *J* = 1.2 Hz, 1H), 4.85 (m, 1H), 3.86 (d, *J* = 5.6 Hz, 2H), 3.75 (t, *J* = 5.6 Hz, 2H)	3419, 1659, 843, 557
AC112		582.79 ([M+H]$^+$)	8.84 (br, 1H), 8.58 (m, 1H), 8.30 (m, 1H), 7.91 (s, 2H), 7.61 (d, *J* = 8.1 Hz, 1H), 7.42 (d, *J* = 7.8 Hz, 1H), 7.00 (dd, *J* = 15.6, 9.3 Hz, 1H), 6.77 (d, *J* = 15.6 Hz, 1H), 4.85 (m, 1H), 4.11 (d, *J* = 5.6 Hz, 1H), 3.73 (d, *J* = 5.6 Hz, 1H), 3.04 (s, 6H)	3399, 1662, 1114, 807, 582
AC113		626.88 ([M+H]$^+$)	8.48 (t, *J* = 5.2 Hz, 1H), 8.3 (s, 1H), 7.90 (s, 2H), 7.79 (dd, *J* = 2.0, 2.0 Hz 2H), 7.58 (d, *J* = 8.4 Hz, 1H) 7.46 (d, *J* = 7.6 Hz, 1H) 7.26 (d, *J* = 7.6 Hz, 1H), 6.98 (m, 1H), 6.75 (d, *J* = 15.6 Hz, 1H), 4.85 (m, 1H), 3.49 (d, *J* = 6.4 Hz, 2H) 2.87 (t, *J* = 6.4 Hz, 2H)	3431, 1651, 1113, 808, 554

Compound Number	mp (°C)	ESIMS	^{1}H NMR (δ)[a]	IR (cm^{-1})
AC114	113.7-117.5	570.7 ([M+H]$^+$)	8.77 (s, 1H), 8.58 (d, *J* = 7.2 Hz, 2H), 7.93 (d, *J* = 7.2 Hz, 2H), 7.60 (dd, *J* = 1.2, 0.8 Hz, 1H), 7.37 (d, *J* = 7.6 Hz, 1H), 6.99 (m, 1H), 6.77 (d, *J* = 16 Hz, 1H), 4.85 (m, 1H), 4.10 (m, 1H) 3.29 (m, 2H), 3.05 (m, 2H), 2.0 (m, 2H), 1.76 (m, 2H)	
AC115		529.00 ([M+H]$^+$)	8.43 (s, 1H), 7.79 (d, *J* = 8.0 Hz, 1H), 7.51 (m, 1H), 7.36 (d, *J* = 8.4 Hz, 3H), 7.21 (m, 3H), 6.55 (d, *J* = 15.6 Hz, 1H), 6.36 (dd, *J* = 15.6, 8.0 Hz, 1H), 5.04 (d, *J* = 5.6 Hz, 2H), 4.10 (m, 1H), 2.35 (s, 3H)	1589, 3459, 801, 1110
AC116		614.87 ([M+H]$^+$)	7.99 (d, *J* = 8.4 Hz, 1H), 7.46 (d, *J* = 1.6 Hz, 1H), 7.34 (d, *J* = 6.4 Hz, 2H), 7.28 (m, 2H), 6.62 (m, 2H), 6.47 (dd, *J* = 16.0, 7.2 Hz, 1H), 4.23 (m, 2H), 4.12 (m, 1H), 4.00 (m, 2H)	3424, 1657, 1165
AC117		525.42 ([M-H]$^-$)	8.39 (br, 1H), 7.85 (br, 1H), 7.62 (m, 3H), 7.53 (d, *J* = 8.0 Hz, 1H), 7.46 (s, 1H), 7.40 (d, *J* = 8.0 Hz, 1H), 7.17 (m, 1H), 6.78 (dd, *J* = 16.0, 8.8 Hz, 1H), 6.70 (m, 1H), 4.77 (m, 1H), 4.66 (s, 1H), 4.32 (s, 1H), 2.97 (s, 3H), 2.16 (s, 3H)	3401, 1636, 1113, 750

Compound Number	mp (°C)	ESIMS	^{1}H NMR (δ)[a]	IR (cm^{-1})
AC118		471.79 ([M+H]$^+$)	7.36 (d, *J* = 8.0 Hz, 2H), 7.27 (m, 2H), 7.22 (m, 2H), 6.57 (d, *J* = 16.0 Hz, 1H), 6.38 (dd, *J* = 16.0, 8.0 Hz, 1H), 6.10 (br, 1H), 4.15 (m, 2H), 3.89 (m, 1H), 3.80 (m, 2H), 3.35 (m, 1H), 2.46 (s, 3H), 2.06 (s,1H), 1.96 (m, 2H), 1.65 (m, 1H)	3437, 1655, 1262, 1105, 802
BC1		492.17 ([M+H]$^+$)	7.39 (s, 2H), 7.25 – 7.18 (m, 3H), 6.58 (d, *J* = 16.0 Hz, 1H), 6.30 (dd, *J* = 16.0, 8.4 Hz, 1H), 5.91 – 5.70 (br, 2H), 4.05 (m, 1H), 3.05 – 2.80 (m, 6H), 2.70 (m, 1H), 1.81 (m, 1H)	3211, 1569, 1113, 806
BC2		506.4 ([M+H]$^+$)	8.80 (s, 1H), 8.20 (s, 1H), 7.82 (m, 3H), 7.4 (s, 2H), 6.62 (d, *J* = 16.0 Hz, 1H), 6.52 (dd, *J* = 16.0, 8.0 Hz, 1H), 4.18(m, 1H), 3.38 (m, 2H), 2.98 (m, 2H), 2.71 (m, 1H), 2.04 (m, 2H), 1.54 (s, 3H).	2923, 1542, 1033, 805
BC3		518.04 ([M-H]$^-$)	7.40 (s, 2H), 7.33 – 7.22 (m, 3H), 6.61 (d, *J* = 16.0 Hz, 1H), 6.34 – 6.28 (dd, *J* = 16.0, 8.0 Hz, 1H), 5.96 – 5.80 (m, 3H), 5.22 (m, 4H), 4.01 (m, 2H), 2.84 – 2.99 (m, 2H), 2.71 (m, 1H), 1.86 (m, 1H)	3120, 1592, 1146, 895

Compound Number	mp (°C)	ESIMS	^{1}H NMR (δ)[a]	IR (cm^{-1})
BC4		529.02 ([M+H]$^+$)	7.39 (s, 2H), 7.25-7.20 (m, 3H), 6.34 (d, *J* = 16.0 Hz, 1H), 6.30 (dd, *J* = 16.0, 8.0 Hz, 1H), 5.81 (br, 1H), 5.48 (m, 1H), 4.10 (m, 1H), 3.10 (m, 2H), 2.86-3.07 (m, 2H), 2.86 (m, 1H), 1.81 (m, 1H);	3283, 1652, 1241, 811
BC5		544.25 ([M-H]$^-$)	7.40 (s, 2H), 7.21 (s, 1H), 7.12 (m, 1H), 6.56 (d, *J* = 16.0 Hz, 1H), 6.32 (dd, *J* = 16.0, 8.4 Hz, 1H), 5.85 (br s, 1H), 5.23 (br s, 1H), 4.12 (m, 1H), 3.18 (m, 3H), 2.80 (m, 3H), 2.08 (m, 2H), 1.83 (m, 5H), 1.25 (m, 2H), 1.01 (m, 3H), 0.78 (m, 2H)	3489, 3291, 1655, 1112, 808
BC6		485.96 ([M-H]$^-$)	7.40 (s, 2H), 7.31 – 7.18 (m, 3H), 6.58 (d, *J* = 16.0 Hz, 1H), 6.24 – 6.28 (dd, *J* = 16.0, 8.0 Hz, 1H), 5.40 (br, 1H), 4.01 (m, 2H), 2.78 – 3.01 (m, 2H), 2.51 (s, 1H), 1.86 (m, 1H), 1.20 (m, 2H), 1.01 (m, 2H), 0.78 (m, 2H)	3429, 1114, 804
BC7		500.01 ([M-H]$^-$)	7.40 (s, 2H), 7.31 (s, 1H), 7.18 (m, 1H), 7.18 (s, 1H), 6.58 (d, *J* = 16.0 Hz, 1H), 6.32 (dd, *J* = 16.0, 8.0 Hz, 1H), 5.78 (br s, 1H), 5.21 (br s, 1H), 4.01 (m, 1H), 2.78 (m, 2H), 2.01 (m, 1H), 1.86 (m, 4H), 1.25 (m, 2H), 1.01 (m, 3H), 0.78 (m, 2H)	3296, 1115, 806

Compound Number	mp (°C)	ESIMS	^{1}H NMR (δ)[a]	IR (cm^{-1})
BC8		511.88 ([M-H]$^-$)	7.38-7.20 (m, 5H), 6.62 (d, *J* = 16.0 Hz, 1H), 6.34 (dd, *J* = 16.0, 8.0 Hz, 1H), 5.83 (br, 1H), 5.52 (m, 1H), 4.12 (m, 1H), 3.12 (m, 2H), 3.06-2.82 (m, 2H), 2.75 (m, 1H), 1.85 (m, 1H)	1657, 1113, 855
BC9	179-181	556.83 ([M-H]$^-$)	8.30 (s, 1H), 7.68 (d, *J* = 6.4 Hz, 1H), 7.38-7.20 (m, 5H), 6.60 (d, *J* = 16.0 Hz, 1H), 6.34 (dd, *J* = 16.0, 8.0 Hz, 1H), 5.63 (br, 1H), 5.52 (m, 1H), 4.12 (m, 1H), 3.56 (s, 2H), 3.06-2.82 (m, 2H), 2.70 (m, 1H), 1.82 (m, 1H)	
BC10		497.98 ([M-H]$^-$)	7.38-7.20 (m, 5H), 6.62 (d, *J* = 16.0 Hz, 1H), 6.34 (dd, *J* = 16.0, 8.0 Hz, 1H), 5.83 (br, 1H), 5.52 (m, 1H), 4.12 (m, 1H), 3.02 (m, 3H), 2.82 (m, 1H), 2.50 (m, 3H), 1.82 (m, 1H), 1.42 (m, 1H)	3027, 1654, 815
BC11		530.09 ([M-H]$^-$)	7.80 (m, 1H), 7.48 (m, 2H), 7.32 6.65 (d, *J* = 16.0 Hz, 1H), 6.54 (dd, *J* = 16.0, 8.0 Hz, 1H), 5.38 (m, 1H), 4.18 (m, 1H), 3.62 (m, 1H), 3.32 (m, 1H), 2.86 (m, 1H), 1.81 (m, 1H)	1715, 1113, 816

Compound Number	mp (°C)	ESIMS	^{1}H NMR (δ)[a]	IR (cm^{-1})
BC12		514.86 ([M+H]$^+$)	7.32, (d, *J* = 6.0 Hz, 2H) 7.28 (m, 1H), 7.20 (d, *J* = 8.0, 1H), 7.14 (d, *J* = 8.8, 1H), 6.70 (d, *J* = 8.0 Hz, 1H), 6.60 (m, 2H), 4.15 (m, 1H), 3.85 (m, 1H), 3.65 (m, 1H), 3.46 (m, 2H), 3.19 (m, 2H);	3428, 1112, 857
BC13	121-126	553.06 ([M-H]$^-$)	8.33 (br, 1H), 7.59 (s, 1H), 7.45 (m, 3H), 6.72 (d, *J* = 3.6, 1H) , 6.39 (m, 1H), 4.71 (t, *J* = 7.2 Hz, 2H), 4.15 (m, 2H)	
BC14	172-175	554.0 ([M-H]$^-$)	8.83 (t, *J* = 6.6 Hz, 1H), 8.42 (t, *J* = 14.7 Hz, 1H), 8.22 (d, *J* = 8.1 Hz, 1H), 8.13 (t, *J* = 6.3 Hz, 1H), 7.98-7.86 (m, 2H), 7.16 - 7.07 (m, 1H), 7.01 - 6.93 (m, 1H), 4.96 - 4.81 (m, 3H), 4.00 - 3.88 (m, 2H)	
CC1	107–109	402.00 ([M+H]$^+$)	7.37 (m, 3H), 7.28 (m, 4H), 6.60 (d, *J* = 16.0 Hz, 1H), 6.36 (dd, *J* = 16.0, 8.0 Hz, 1H), 5.75 (br s, 1H), 4.46 (d, *J* = 6 Hz, 2H), 4.01 (m, 1H), 2.11 (s, 3H)	
CC2	118–120	428.11 ([M+H]$^+$)	7.37 (m, 3H), 7.28 (m, 4H), 6.60 (d, *J* = 16.0 Hz, 1H), 6.35 (dd, *J* = 16.0, 8.0 Hz, 1H), 5.83 (br s, 1H), 4.46 (d, *J* = 6.0 Hz, 2H), 4.11 (m, 1H), 1.40 (m, 1H), 1.02 (m, 2H), 0.77 (m, 2H)	

Compound Number	mp (°C)	ESIMS	^{1}H NMR (δ)[a]	IR (cm^{-1})
CC3	119–122	468.20 ([M-H]$^-$)	7.38 (m, 3H), 7.27 (m, 3H), 6.60 (d, *J* = 16.0 Hz, 1H), 6.36 (dd, *J* = 16.0, 8.4 Hz, 1H), 5.00 (br s, 1H), 4.48 (d, *J* = 5.6 Hz, 2H), 4.11 (m, 1H), 3.15 (q, *J* = 10.4 Hz, 2H)	
CC4		414.16 ([M-H]$^-$)	7.37 (m, 3H), 7.28 (m, 3H), 6.60 (d, *J* = 16.0 Hz, 1H), 6.35 (dd, *J* = 16.0, 8.0 Hz, 1H), 5.69 (br s, 1H), 4.46 (d, *J* = 6.0 Hz, 2H), 4.21 (m, 1H), 2.29 (q, *J* = 5.8 Hz, 2H), 1.30 (t, *J* = 7.2 Hz, 3H)	
CC5		460.28 ([M-H]$^-$)	7.40 (m, 3H), 7.28 (m, 2H), 6.60 (d, *J* = 15.6 Hz, 1H), 6.33 (dd, *J* = 15.6, 8.0 Hz, 1H), 5.84 (br s, 1H), 4.46 (d, *J* = 5.6 Hz, 2H), 4.10 (m, 1H), 1.36 (m, 1H), 1.02 (m, 2H), 0.77 (m, 2H)	
CC6	106–108	504.08 ([M-H]$^-$)	7.40 (m, 3H), 7.26 (m, 1H), 6.60 (d, *J* = 16.0 Hz, 1H), 6.34 (dd, *J* = 16.0, 8.0 Hz, 1H), 5.96 (br s, 1H), 4.49 (d, *J* = 5.6 Hz, 2H), 4.10 (m, 1H), 3.15 (q, *J* = 10.8 Hz, 2H)	
CC7	127–128	436.03 ([M+H]$^+$)	7.42 (m, 4H), 7.24 (m, 2H), 6.53 (d, *J* = 16.0 Hz, 1H), 6.36 (dd, *J* = 16.0, 8.0 Hz, 1H) , 5.86 (br s, 1H), 4.51 (d, *J* = 6.0 Hz, 2H), 4.05 (m, 1H), 2.02 (s, 3H)	

Compound Number	mp (°C)	ESIMS	^{1}H NMR (δ)[a]	IR (cm^{-1})
CC8	129–131	462.15 ([M+H]$^+$)	8.58 (t, *J* = 5.6 Hz, 1H), 7.72 (m, 1H), 7.66 (m, 3H), 7.49 (d, *J* = 8.0 Hz, 1H), 7.30 (d, *J* = 8.0 Hz, 1H), 6.90 (dd, *J* = 16.0, 8.0 Hz, 1H), 6.73 (d, *J* = 16 Hz, 1H), 4.81 (m, 1H), 4.33 (d, *J* = 6.0 Hz, 1H), 1.64 (m, 1H), 0.68 (m, 4H)	
CC9	132–134	504.25 ([M+H]$^+$)	7.41 (m, 3H), 7.26 (m, 3H), 6.54 (d, *J* = 16.0 Hz, 1H), 6.37 (dd, *J* = 16.0, 8.0 Hz, 1H), 6.13 (br s, 1H), 4.56 (d, *J* = 6.0 Hz, 2H), 4.11 (m, 1H), 3.13 (m, 2H)	
CC10		538.03 ([M+2H]$^+$)	7.38 (m, 4H), 6.56 (d, *J* = 16.0 Hz, 1H), 6.38 (dd, *J* = 16.0, 8.0 Hz, 1H), 6.18 (m, 1H), 4.58 (m, 2H), 4.08 (m, 1H), 3.08 (m, 2H)	1651, 1112, 807
CC11	111–112	494.12 ([M-H]$^-$)	7.42 (m, 3H), 7.24 (m, 1H), 6.54 (d, *J* = 15.6 Hz, 1H), 6.34 (dd, *J* = 16.0, 8.0 Hz, 1H), 6.03 (m, 1H), 4.53 (d, *J* = 6.0 Hz, 1H), 4.10 (m, 1H), 1.39 (m, 1H), 1.00 (m, 2H), 0.77 (m, 2H)	
CC12	76–78	510.07 ([M-H]$^-$)	7.39 (s, 4H), 7.34 (d, *J* = 8.0 Hz, 1H), 7.26 (m, 1H), 6.57 (d, *J* = 16.0 Hz, 1H), 6.35 (dd, *J* = 16.0, 8.0 Hz, 1H), 6.10 (br s, 1H), 4.49 (d, *J* = 6.0 Hz, 2H), 4.10 (m, 1H), 1.20 (s, 9H)	

Compound Number	mp (°C)	ESIMS	^{1}H NMR (δ)[a]	IR (cm^{-1})
CC13	73–76	563.37 ([M-H]$^-$)	8.51 (d, *J* = 5.2 Hz, 1H), 7.63 (s, 1H), 7.51 (m, 1H), 7.45 (m, 2H), 7.39 (s, 2H), 7.28 (m, 1H), 6.58 (m, 2H), 6.37 (dd, *J* = 16.0, 8.0 Hz, 1H), 4.71 (d, *J* = 6.0 Hz, 1H), 4.11 (m, 1H)	
CC14		581.45 ([M+1H]$^+$)	8.51 (m, 1H), 8.30 (d, *J* = 2.4 Hz, 1H), 7.73 (m, 1H), 7.61 (s, 2H), 7.51 (s, 1H), 7.32 (m, 3H), 6.66 (d, *J* = 16.0 Hz, 1H), 6.56 (dd, *J* = 16.0, 8.4 Hz, 1H), 4.50 (m, 1H), 4.45 (d, *J* = 5.6 Hz, 1H), 3.56 (s, 2H)	3430, 1656, 1109, 806
CC15		480.24 ([M+H]$^+$)	7.40 (m, 3H), 7.33 (m, 1H), 7.22 (m, 2H), 6.54 (d, *J* = 15.6 Hz, 1H), 6.34 (dd, *J* = 16.0, 8.0 Hz, 1H), 6.03 (br s, 1H), 4.53 (d, *J* = 6.0 Hz, 2H), 4.13 (m, 1H), 1.41 (m, 1H), 1.00 (m, 2H), 0.77 (m, 2H)	3293, 1651, 1543, 1114, 812
CC16		520.33 ([M-H]$^-$)	7.42 (s, 1H), 7.37 (m, 3H), 7.22 (m, 1H), 6.54 (d, *J* = 16.0 Hz, 1H), 6.36 (dd, *J* = 16.0, 8.0 Hz, 1H), 6.19 (br s, 1H), 4.51 (d, *J* = 6.0 Hz, 2H), 4.21 (m, 1H), 3.33 (m, 2H)	3307, 1665, 1114, 813
CC17	117–119	459.83 ([M-H]$^-$)	7.51 (m, 2H), 7.39 (m, 2H), 7.24 (m, 2H), 6.52 (d, *J* = 15.6 Hz, 1H), 6.38 (dd, *J* = 15.6, 7.6 Hz, 1H), 6.02 (br s, 1H), 4.53 (d, *J* = 6.0 Hz, 2H), 4.14 (m, 1H), 1.38 (m, 1H)), 1.00 (m, 2H), 0.77 (m, 2H)	3293, 1633, 1110, 820

Compound Number	mp (°C)	ESIMS	^{1}H NMR (δ)[a]	IR (cm^{-1})
CC18	119–123	501.88 ([M-H]$^-$)	7.48 (m, 2H), 7.41 (s, 1H), 7.36 (d, *J* = 8.0 Hz, 1H), 7.23 (m, 2H), 6.52 (d, *J* = 16.0 Hz, 1H), 6.39 (dd, *J* = 16.0, 8.0 Hz, 1H), 6.13 (br s, 1H), 4.56 (d, *J* = 6.0 Hz, 2H), 4.15 (m, 1H), 3.13 (m, 2H)	3435, 1644, 1111, 817
CC19		530 ([M+H]$^+$)	7.41 (m, 2H), 7.24 (m, 1H), 6.53 (d, *J* = 16.0 Hz, 1H), 6.35 (dd, *J* = 16.0, 8.0 Hz, 1H), 4.53 (m, 2H), 4.10 (m, 1H), 3.42 (m, 2H), 2.97 (s, 3H), 2.78 (m, 2H)	3435, 1644, 1111, 817
CC20		512 ([M+H]$^+$)	7.42 (m, 3H), 7.24 (m, 1H), 6.54 (d, *J* = 15.6 Hz, 1H), 6.34 (dd, *J* = 15.6, 8.0 Hz, 1H), 6.03 (m 1H), 4.53 (d, *J* = 6.0 Hz, 1H), 4.10 (m, 1H), 1.19 (m, 1H), 1.00 (m, 2H), 0.77 (m, 2H)	3293, 1633, 1110, 820
CC21	55–58	493.99 ([M-H]$^-$)	(DMSO-d_6) 8.62 (m, 1H), 7.95 (s, 1H), 7.85 (m, 1H), 7.66 (m, 3H), 7.47 (d, *J* = 8.0 Hz, 1H), 6.98 (dd, *J* =16.0, 8.0 Hz, 1H), 6.84 (d, *J* = 16.0 Hz, 1H), 4.83 (m, 1H), 4.44 (s, 2H), 1.68 (m, 1H), 0.71 (m, 4H)	
CC22	67–69	530.01 ([M+H]$^+$)	8.62 (m, 1H), 7.90 (s, 3H), 7.82 (m, 1H), 7.45 (m, 1H), 6.98 (m, 1H), 6.84 (d, *J* = 16.0 Hz, 1H), 4.82 (m, 1H), 4.4 (s, 2H), 1.66 (m, 1H), 0.72 (m, 4H)	

Compound Number	mp (°C)	ESIMS	^{1}H NMR (δ)[a]	IR (cm^{-1})
CC23	69–71	564.99 ([M-H]$^-$)	9.02 (br s, 1H), 8.54 (br s, 1H), 8.26 (br s, 1H), 7.48 – 7.54 (m, 3H), 7.22 – 7.42 (m, 3H), 6.59 – 6.62 (m, 2H), 6.38 – 6.42 (m, 1H), 4.82 (m, 2H), 4.19 (s, 1H)	
CC24	125–127	570.26 ([M-H]$^-$)	7.64 (s, 1H), 7.54 (s, 2H), 7.46 (s, 2H), 6.62 (d, *J* = 16.0 Hz, 1H), 6.41 (dd, *J* = 16.0, 8.4 Hz, 1H), 6.03 (m, 1H), 4.65 (d, *J* = 6.4 Hz, 2H), 4.14 (m, 1H,), 3.13 (q, *J* = 10.6 Hz, 2H)	
CC25		579.86 ([M-H]$^-$)	7.60 (s, 1H), 7.40 (s, 2H), 7.37 (d, *J* = 8.0 Hz, 1H), 7.31 (d, *J* = 8.0 Hz, 1H), 6.53 (d, 1H, *J* = 16.0 Hz), 6.35 (dd, *J* = 16.0, 8.0 Hz, 1H), 6.17 (br s, 1H), 4.56 (d, *J* = 6.4 Hz, 2H), 4.12 (m, 1H), 3.15 (q, *J* = 10.6 Hz, 2H)	3297, 1663, 1114, 809
CC26	129–131	539.89 ([M+H]$^+$)	7.59 (s, 1H), 7.39 (m, 2H), 7.30 (s, 1H), 6.53 (d, *J* = 16.0 Hz, 1H), 6.35 (dd, *J* = 16.0, 8.0 Hz, 1H), 6.06 (br s, 1H), 4.42 (d, *J* = 4.4 Hz, 2H), 4.12 (m, 1H), 1.35 (br s, 1H), 0.95 (br s, 2H), 0.75 (m, 2H)	

Compound Number	mp (°C)	ESIMS	^{1}H NMR (δ)[a]	IR (cm^{-1})
CC27		519.95 ([M-H]$^-$)	7.39 (s, 2H), 7.33 (t, *J* = 7.6 Hz, 1H), 7.14 (m, 2H), 6.56 (d, *J* = 16.0 Hz, 1H), 6.35 (dd, *J* = 16.0, 7.6 Hz, 1H), 6.06 (br s, 1H), 4.52 (d, *J* = 16.0 Hz, 2H), 4.08 (m, 1H), 3.90 (s, 2H), 3.13 (m, 2H)	3306, 1786
CC28		477.93 ([M-H]$^-$)	7.39 (s, 2H), 7.35 (m, 1H), 7.14 (m, 2H), 6.55 (d, *J* = 15.6 Hz, 1H), 6.33 (dd, *J* = 15.6, 8.0 Hz, 1H), 5.93 (br s, 1H), 4.49 (d, *J* = 16.0 Hz, 2H), 4.10 (m, 1H), 1.36 (m, 1H), 1.00 (m, 2H), 0.77 (m, 2H)	3625, 1747
CC29		620.86 ([M-H]$^-$)	8.58 (d, *J* = 4.6 Hz, 1H), 7.74 (m, 1H), 7.62 (m, 2H), 7.52 (m, 1H), 7.4 (s, 2H), 7.3 (m, 1H), 7.2 (m, 2H), 6.60 (d, *J* = 16.0 Hz, 1H), 6.38 (dd, *J* = 16.0, 8.0 Hz, 1H), 5.02 (s, 1H), 4.8 (s, 1H), 4.8 (d, *J* = 10 Hz, 2H), 4.10 (m, 1H), 1.8 (m, 1H), 1.2 (m, 2H), 0.6 (m, 2H)	1645, 1115, 808
CC30	101–104	559.75 ([M-H]$^-$)	7.41 (m, 4H), 7.24 (m, 1H), 6.53 (d, *J* = 16.0 Hz, 1H), 6.35 (dd, *J* = 16.0, 8.0 Hz, 1H), 6.12 (br s, 1H), 4.53 (m, 2H), 4.10 (m, 1H), 3.42 (m, 2H), 2.91 (s, 3H), 2.78 (m, 2H)	

Compound Number	mp (°C)	ESIMS	^{1}H NMR (δ)[a]	IR (cm^{-1})
CC31	177–178	463 ([M-H]$^-$)	7.58 (m, 2H), 7.41 (m, 3H), 7.24 (m, 1H), 6.53 (d, *J* = 16.0 Hz, 1H), 6.35 (dd, *J* = 16.0, 8.0 Hz, 1H), 4.70 (br s, 1H), 4.43 (s, 2H), 4.08 (m, 1H), 3.21 (m, 2H), 1.25 (m, 3H);	
CC32	141–142	532.99 ([M+H]$^+$)	7.66 (m, 2H), 7.54 (m, 1H), 7.41 (s, 2H), 6.62 (d, *J* = 16.0 Hz, 1H), 6.40 (dd, *J* = 16.0, 8.0 Hz, 1H), 4.59 (s, 3H), 4.19 (m, 1H), 3.25 (m, 2H), 1.15 (m, 2H)	
CC33		540.88 ([M-H]$^-$)	7.57 (s, 1H), 7.40 (m, 2H), 7.30 (s, 1H), 7.20 (br s, 1H), 6.53 (d, *J* = 16.0 Hz, 1H), 6.33 (dd, *J* = 16.0, 8.0 Hz, 1H), 6.06 (br s, 1H), 4.75 (br s, 1H), 4.42 (s, 2H), 4.20 (br s, 1H), 4.15 (m, 2H), 3.20 (m, 2H), 1.15 (m, 3H)	3338, 1631, 1578, 1114, 809
CC34	118–120	541.40 ([M+H]$^+$)	7.42 (m, 3H), 7.28 (m, 2H), 6.54 (d, *J* = 16.0 Hz, 1H), 6.36 (dd, *J* = 16.0, 8.0 Hz, 1H), 4.96 (m, 1H), 4.51 (d, *J* = 5.6 Hz, 2H), 4.12 (m, 1H), 3.69 (t, *J* = 4.8 Hz, 4H), 3.35 (t, *J* = 4.8 Hz, 1H)	

Compound Number	mp (°C)	ESIMS	^{1}H NMR (δ)[a]	IR (cm^{-1})
CC35	78–79	547.82 ([M+H]$^+$)	9.95 (br s, 1H), 8.17 (d, *J* = 4.8 Hz, 1H), 7.61 (d, *J* = 6.4 Hz), 7.43 (m, 3H), 7.24 (m, 2H), 6.90 (t, *J* = 5.6 Hz, 1H), 6.66 (d, *J* = 8.4 Hz, 1H), 6.54 (d, *J* = 16.0 Hz, 1H), 6.33 (dd, *J* = 16.0, 8.0 Hz, 1H), 4.65 (d, *J* = 6.0 Hz, 1H), 4.09 (m, 1H)	
CC36		497 ([M-H]$^-$)	7.39 (m, 4H), 7.28 (m, 1H), 6.54 (d, *J* = 16.0 Hz, 1H), 6.34 (dd, *J* = 16.0, 8.0 Hz, 1H), 4.97 (br s, 1H), 4.38 (d, *J* = 6.0 Hz, 2H), 4.10 (m, 1H), 2.9 (s, 3H), 2.7 (s, 3H)	3350, 1705, 1114, 808
CC37	88–91	515.01 ([M+H]$^+$)	7.49 (d, *J* = 8 Hz, 1H), 7.41 (d, *J* = 7.2 Hz, 2H), 7.26 (m, 2H), 6.50 (d, *J* = 16 Hz, 1H), 6.35 (dd, *J* = 16.0, 8.0 Hz, 1H), 6.0 (brs, 1H), 5.73 (br s, 1H), 4.80 (br s, 2H), 4.09 (m, 1H), 1.23 (m, 3H)	
CC38	63–66	526.97 ([M+H]$^+$)	7.48 (d, *J* = 8 Hz, 1H), 7.39 (m, 3H), 7.27 (m, 1H), 6.54 (d, *J* = 16 Hz, 1H), 6.33 (dd, *J* = 6.0, 8.0 Hz, 1H), 6.17 (br s, 1H), 5.92 (br s, 1H), 5.83 (m, 2H), 5.29 (t, *J* = 15.4 Hz, 2H), 4.80 (br s, 2H), 4.12 (m, 1H), 4.02 (br s, 2H)	

Compound Number	mp (°C)	ESIMS	^{1}H NMR (δ)[a]	IR (cm^{-1})
CC39		526.09 ([M-H]$^-$)	7.39 (m, 4H), 7.28 (m, 1H), 6.54 (d, *J* = 16.0 Hz, 1H), 6.34 (dd, *J* = 16.0, 8.0 Hz, 1H), 4.97 (br s, 1H), 4.38 (d, *J* = 6.0 Hz, 2H), 4.10 (m, 1H), 1.53 (s, 9H)	3350, 1705, 1114, 808
CC40	159–160	580.25 ([M-H]$^-$)	7.46 (m, 5H), 7.29 (m, 1H), 7.20 (m, 3H), 6.55 (d, *J* = 16.0 Hz, 1H), 6.37 (dd, *J* = 16.0, 8.0 Hz, 1H), 5.62 (br s, 1H), 4.55 (d, *J* = 6.4 Hz, 2H), 4.11 (m, 1H)	
CC41		512.22 ([M-H]$^-$)	7.48 (m, 1H), 7.43 (m, 3H), 7.38 (m, 1H), 7.23 (s, 1H), 6.55 (d, *J* = 16.0 Hz, 1H), 6.36 (d, *J* = 16.0 Hz, 1H), 4.60 (d, 2H), 4.18 (m, 1H), 3.85 (s, 3H)	1740, 1701, 1114, 808
CC42	161–163	578.96 ([M-H]$^-$)	(DMSO-d_6) 9.45 (br s, 2H), 7.90 (s, 2H), 7.75 (s, 1H), 7.46 (br s, 1H), 7.28 (br s, 1H), 6.93 (m, 1H), 6.75 (br s, 1H), 4.80 (m, 1H), 4.40 (br s, 2H), 3.90 (br s, 2H)	
CC43	140–142	505.39 ([M+H]$^+$)	8.11 (d, *J* = 4.0 Hz, 1H), 7.40 (m, 5H), 7.22 (m, 1H), 6.61 (m, 2H), 6.35 (m, 2H), 4.94 (br s, 1H) 4.61 (d, *J* = 6.4 Hz, 2H), 4.11 (m, 1H)	
CC44		536.88 ([M-H]$^-$)	8.41 (s, 1H), 7.77 (s, 1H), 7.47 (br s, 1H), 7.40 (s, 2H), 6.58 (d, *J* = 16.0 Hz, 1H), 6.45 (dd, *J* = 16.0, 8.0 Hz, 1H), 4.68 (d, *J* = 4.0 Hz, 2H), 4.14 (m, 1H), 3.24 (q, *J* = 10.8 Hz, 2H)	3320, 1674, 1114, 808

Compound Number	mp (°C)	ESIMS	^{1}H NMR (δ)[a]	IR (cm^{-1})
CC45		494.88 ([M-H]$^-$)	8.41 (s, 1H), 7.76 (s, 1H), 7.40 (s, 2H), 7.15 (br s, 1H), 6.58 (d, *J* = 16.0 Hz, 1H), 6.44 (dd, *J* = 16.0, 8.0 Hz, 1H), 4.67 (d, *J* = 4.4 Hz, 2H), 4.16 (m, 1H), 1.57 (m, 1H), 1.04 (m, 2H), 0.87 (m, 2H)	3309, 1659, 1115, 808
CC46	151–153	554.04 ([M-H]$^-$)	8.06 (m, 1H), 7.61 (m, 4H), 7.48 (s, 2H), 7.44 (d, *J* = 8.0 Hz, 1H), 7.38 (m, 1H), 6.42 (m, 1H), 5.92 (br s, 1H), 4.92 (m, 2H), 4.24 (m, 1H), 3.12 (m, 2H)	
CC47		478.09 ([M+H]$^+$)	8.06 (m, 2H), 7.61 (m, 4H), 7.48 (s, 2H), 7.44 (d, *J* = 8.0 Hz, 1H), 7.38 (m, 2H), 6.42 (m, 1H), 4.92 (s, 2H), 1.36 (m, 1H), 1.00 (m, 2H), 0.77 (m, 2H)	3309, 1659, 1115, 808
CC48		511.05 ([M+H]$^+$)	8.06 (m, 2H), 7.61 (m, 3H), 7.48 (s, 2H), 7.44 (d, *J* = 8.0 Hz, 1H), 7.38 (m, 2H), 6.42 (m, 1H), 4.92 (s, 2H), 1.36 (m, 1H), 1.00 (m, 2H), 0.77 (m, 2H)	3309, 1659, 1115, 808
CC49	84–87	515.33 ([M+H]$^+$).	8.06 (m, 1H), 7.98 (m, 1H), 7.61 (m, 3H), 7.48 (s, 2H), 7.44 (d, *J* = 8.0 Hz, 1H), 7.38 (m, 2H), 6.42 (m, 1H), 4.92 (s, 2H), 4.6 (br s, 1H), 4.24 (m, 1H), 3.21 (m, 2H), 1.2 (t, *J* = 4.6 Hz, 3H)	

Compound Number	mp (°C)	ESIMS	^{1}H NMR (δ)[a]	IR (cm^{-1})
CC50	138–140	461.32 ([M-1H]$^-$)	9.81 (s, 1H), 7.90 (s, 1H), 7.84 (s, 2H), 7.34 (d, *J* = 8.4 Hz, 2H), 6.65 (d, *J* = 15.6 Hz, 1H), 6.61 (m, 1H), 6.57 (s, 1H), 6.48 (dd, *J* = 15.6, 8.8 Hz, 1H), 4.74 (m, 1H), 1.64 (m, 1H), 0.75 (m, 4H);	
CC51	149–150	505.31 ([M-H]$^-$)	7.56 (br s, 1H), 7.4 (s, 3H), 7.3 (m, 3H), 7.05 (br s, 1H), 6.8 (d, *J* = 6 Hz, 2H), 6.57 (m, 2H), 6.20 (m, 2H), 4.05 (m, 1H), 3.2 (q, *J* = 10.4 Hz, 2H)	
CC52		464.87 ([M-H]$^-$)	7.40 (s, 2H), 7.18 (s, 1H), 7.08 (s, 1H), 6.85 (m, 1H), 6.45 (m, 1H), 6.20 (m, 1H), 5.55 (s,1H), 4.08 (m, 1H), 1.30 – 1.10 (m, 4H), 1.90 (m,1H)	3309, 1659, 1115, 808
CC53		506 ([M+H]$^+$)	7.40 (s, 2H), 7.18 (s, 1H), 7.08 (s, 1H), 6.85 (m, 1H), 6.45 (m, 1H), 6.20 (m, 1H), 5.55 (s,1H), 4.08 (m, 1H), 3.21 (m, 2H)	3309, 1659, 1115, 808
CC54		504 ([M+H]$^+$)	7.28 (s, 2H), 7.25 (m, 2H), 7.10 (d, *J* = 8.0 Hz, 2H), 6.89 (d, *J* = 11.4 Hz, 1H), 6.07 (br s, 1H), 6.01 (m, 1H), 4.51 (d, *J* = 5.8 Hz, 2H), 4.34 (m, 1H), 3.12 (q, *J* = 7.5 Hz, 2H)	

Compound Number	mp (°C)	ESIMS	^{1}H NMR (δ)[a]	IR (cm^{-1})
DC1	93–97	398.05 ([M+H]$^+$)	8.56 (s, 1H), 8.11 (s, 1H), 7.68 (d, *J* = 8.4 Hz, 2H), 7.54 (d, *J* = 8.4 Hz, 2H), 7.38 (t, *J* = 1.8 Hz, 1H), 7.29 (s, 2H), 6.62 (d, *J* = 15.6 Hz, 1H), 6.42 (dd, *J* = 15.6, 8.2 Hz, 1H), 4.15 (m, 1H)	
DC2		363.0746 (363.075)	8.59 (s, 1H), 8.13 (s, 1H), 7.69 (d, *J* = 8.5 Hz, 2H), 7.55 (d, *J* = 8.5 Hz, 2H), 7.41 – 7.29 (m, 4H), 6.64 (d, *J* = 15.7 Hz, 1H), 6.47 (dd, *J* = 15.9, 8.0 Hz, 1H), 4.17 (m, 1H)	3121, 1524, 1251, 1165, 1119
DC3		329.1144 (329.114)	8.56 (s, 1H), 8.11 (s, 1H), 7.65 (d, *J* = 8.4 Hz, 2H), 7.52 (d, *J* = 8.3 Hz, 2H), 7.40 (m, 5H), 6.61 (d, *J* = 15.8 Hz, 1H), 6.51 (dd, *J* = 15.9, 7.7 Hz, 1H), 4.18 (m, 1H)	1521, 1246, 1219, 1162, 1152, 1107
DC4		364.11 ([M+H]$^+$)	8.56 (s, 1H), 8.10 (s, 1H), 7.66 (d, *J* = 2.0 Hz, 2H), 7.52 (d, *J* = 8.8 Hz, 2H), 7.38 (d, *J* = 2.4 Hz, 2H), 7.34 (d, *J* = 8.4 Hz, 2H), 6.61 (d, *J* = 16.0 Hz, 1H), 6.40 (dd, *J* = 16.0, 7.6 Hz, 1H), 4.15 (m, 1H)	3147, 1528, 1494, 1246, 1165, 1108
DC5		344.25 ([M+H]$^+$)	8.54 (s, 1H), 8.10 (s, 1H), 7.62 (d, *J* = 8.3 Hz, 2H), 7.50 (d, *J* = 8.4 Hz, 2H), 7.25 (d, *J* = 8.3 Hz, 2H), 7.20 (d, *J* = 8.0 Hz, 2H), 6.60 (d, *J* = 16.0 Hz, 1H), 6.51 (dd, *J* = 16.0, 8.0 Hz, 1H), 4.15 (m, 1H), 2.37 (s, 3H)	3122, 3047, 1523, 1252, 1160, 1107

Compound Number	mp (°C)	ESIMS	^{1}H NMR (δ)[a]	IR (cm^{-1})
DC6		360.28 ([M+H]$^+$)	8.55 (s, 1H), 8.10 (s, 1H), 7.65 (d, *J* = 8.8 Hz, 2H), 7.52 (d, *J* = 8.8 Hz, 2H), 7.32 (d, *J* = 8.8 Hz, 2H), 6.95 (d, *J* = 8.8 Hz, 2H), 6.60 (d, *J* = 16.0 Hz, 1H), 6.56 (dd, *J* = 16.0, 7.4 Hz, 1H), 4.15 (m, 1H), 3.82 (s, 3H)	3124, 2936, 1522, 1249, 1160
DC7		348 ([M+H]$^+$)	8.55 (s, 1H), 8.10 (s, 1H), 7.62 (d, *J* = 8.8 Hz, 2H), 7.5 (d, *J* = 8.4 Hz, 2H), 7.38 (m, 2H), 7.12 (m, 2H), 6.61 (d, *J* = 16.0 Hz, 1H), 6.40 (dd, *J* = 16.0, 7.6 Hz, 1H), 4.15 (m, 1H)	3141, 1512, 1246, 1118
DC8		366.13 ([M+H]$^+$)	8.57 (s, 1H), 8.11 (s, 1H), 7.65 (d, *J* = 7.2 Hz, 2H), 7.52 (d, *J* = 8.0 Hz, 2H), 6.95 (m, 2H), 6.82 (m, 1H), 6.65 (d, *J* = 16.0 Hz, 1H), 6.50 (dd, *J* = 16.0, 8.0 Hz, 1H), 4.15 (m, 1H)	3116, 1628, 1524, 1252, 1168, 1118
DC9		348.11 ([M+H]$^+$)	8.71 (s, 1H), 8.20 (s, 1H), 7.70 (d, *J* = 8.0 Hz, 2H), 7.57 (d, *J* = 8.0 Hz, 2H), 7.40 (m, 1H), 7.19 (m, 3H), 6.60 (d, *J* = 16.0 Hz, 1H), 6.40 (dd, *J* = 16.0, 8.4 Hz, 1H), 4.15 (m, 1H)	3115, 1525, 1248, 1174
DC10		348.11 ([M+H]$^+$)	8.75 (s, 1H), 8.20 (s, 1H), 7.72 (d, *J* = 8.4 Hz, 2H), 7.6 (d, *J* = 8.4 Hz, 2H), 7.20 – 7.40 (m, 4H), 6.60 (d, *J* = 16.0 Hz, 1H), 6.40 (dd, *J* = 16.0, 8.0 Hz, 1H,), 4.60 (m, 1H)	3114, 1526, 1259, 1238, 1193, 1114

Compound Number	mp (°C)	ESIMS	^{1}H NMR (δ)[a]	IR (cm^{-1})
DC11	75.5–78.5	358.14 ([M+H]$^+$)	8.55 (s, 1H), 8.10 (s, 1H), 7.65 (d, *J* = 8.8 Hz, 2H), 7.52 (d, *J* = 8.4 Hz, 2H), 7.01 (s, 3H), 6.60 (d, *J* = 16.0 Hz, 1H), 6.51 (dd, *J* = 16.0, 7.8 Hz, 1H), 4.15 (m, 1H), 2.34 (s, 6H)	
DC12		398.05 ([M+H]$^+$)	8.58 (s, 1H), 8.10 (s, 1H), 7.68 (d, *J* = 8.4 Hz, 2H), 7.53 (m, 4H), 7.2 (s, 1H) 6.62 (d, *J* = 15.6 Hz, 1H), 6.44 (dd, *J* = 15.6, 8.0 Hz, 1H), 4.15 (m, 1H)	3055, 2930, 1523, 1250, 1165
DC13		396.16 ([M+H]$^+$)	8.58 (s, 1H), 8.10 (s, 1H), 7.62 (d, *J* = 8.4 Hz, 2H), 7.55 (m, 4H), 7.25 (m, 1H), 6.64 (d, *J* = 16.0 Hz, 1H), 6.40 (dd, *J* = 16.0, 8.0 Hz, 1H), 4.90 (m, 1H)	3108, 1523, 1249, 1166, 1127
DC14		398.05 ([M+H]$^+$)	8.58 (s, 1H), 8.10 (s, 1H), 7.62 (d, *J* = 8.4 Hz, 2H), 7.55 (m, 4H), 7.25 (m, 1H), 6.67 (d, *J* = 16.0 Hz, 1H), 6.40 (dd, *J* = 16.0, 8.0 Hz, 1H), 5.00 (m, 1H)	3117, 2925, 1526, 1246, 1172, 1117
DC15		397.95 ([M+H]$^+$)	8.58 (s, 1H), 8.10 (s, 1H), 7.66 (d, *J* = 8.0 Hz, 2H), 7.52 (m, 3H), 7.40 (d, *J* = 8.0 Hz, 1H), 7.30 (dd, *J* = 8.4, 2.9 Hz, 1H), 6.64 (d, *J* = 16.0 Hz, 1H), 6.40 (dd, *J* = 16.0, 8.0 Hz, 1H), 4.90 (m, 1H)	3120, 1524, 1267, 1176, 1112

Compound Number	mp (°C)	ESIMS	^{1}H NMR (δ)[a]	IR (cm^{-1})
DC16		466 ([M+H]$^+$)	8.61 (s, 1H), 8.13 (s, 1H), 7.92 (s, 1H), 7.86 (s, 2H), 7.70 (d, *J* = 7.0 Hz, 2H), 7.54 (d, *J* = 7.0 Hz, 2H), 6.67 (d, *J* = 16.0 Hz, 1H), 6.46 (dd, *J* = 16.0, 8.0 Hz, 1H), 4.35 (m, 1H)	
DC17		430.06 ([M+H]$^+$)	8.58 (s, 1H), 8.1 (s, 1H), 7.68 (d, *J* = 8.4 Hz, 2H), 7.54 (d, *J* = 8.4 Hz, 2H), 7.51 (s, 1H), 7.42 (s, 1H), 6.68 (d, *J* = 16.0 Hz, 1H), 6.35 (dd, *J* = 16.0, 8.0, Hz, 1H), 4.98 (m, 1H)	3122, 3076, 2929, 1523, 1250, 1168, 1114
DC18	92–95	429.91 ([M+H]$^+$)	8.57 (s, 1H), 8.11 (s, 1H), 7.69 (d, *J* = 8.8 Hz, 2H), 7.54 (d, *J* = 8.4 Hz, 2H), 7.42 (s, 2H), 6.65 (d, *J* = 16.0 Hz, 1H), 6.40 (dd, *J* = 16.0, 8.0 Hz, 1H), 4.10 (m, 1H)	
DC19	97–99	430.321 ([M+H]$^+$)	8.58 (s, 1H), 8.12 (s, 1H), 7.68 (d, *J* = 8.0 Hz, 2H), 7.64 (s, 1H), 7.59 (s, 1H), 7.55 (m, 3H), 6.60 (d, *J* = 16.0 Hz, 1H), 6.40 (dd, *J* = 16.0, 8.0 Hz, 1H), 4.22 (m, 1H)	
DC20		427.0463 (427.0466)	8.58 (s, 1H), 8.15 (s, 1H), 7.70 (d, *J* = 8.4 Hz, 2H), 7.58 (d, *J* = 8.4 Hz, 2H), 7.36 (s, 2H), 6.62 (d, *J* = 16.0 Hz, 1H), 6.43 (dd, *J* = 16.0, 8.0 Hz, 1H), 4.12 (m, 1H), 3.88 (s, 3H)	2937, 1524, 1482, 1278, 1249, 1166, 1112

Compound Number	mp (°C)	ESIMS	^{1}H NMR (δ)[a]	IR (cm^{-1})
DC21		412.04 ([M+H]$^+$)	8.42 (s, 1H), 7.60 (d, *J* = 8.0 Hz, 2H), 7.50 (d, *J* = 8.0 Hz, 2H), 7.40 (s, 1H), 7.22 (s, 2H), 6.60 (d, *J* = 16.0 Hz, 1H), 6.42 (dd, *J* = 16.0, 8.0 Hz, 1H), 4.15 (m, 1H), 2.5 (s, 3H)	3108, 1572, 1531, 1242, 1172, 1104
DC22	147–149	441.01 ([M-H]$^-$)	8.62 (s, 1H), 7.78 (d, *J* = 8.0 Hz, 2H), 7.60 (d, *J* = 8.0 Hz, 2H), 7.40 (s, 1H), 7.30 (s, 2H), 6.67 (d, *J* = 16.0 Hz, 1H), 6.48 (dd, *J* = 16.0, 8.0 Hz, 1H), 4.15 (m, 1H)	
DC23		412.05 ([M+H]$^+$)	7.95 (s, 1H), 7.35 (d, *J* = 8.0 Hz, 2H), 7.46 (d, *J* = 8.0 Hz, 2H), 7.39 (s, 1H), 7.29 (s, 2H), 6.67 (d, *J* = 16.0 Hz, 1H), 6.45 (dd, *J* = 16.0, 8.0 Hz, 1H), 4.12 (m, 1H), 2.51 (s, 3H)	1112, 799
DC24	133–134	440.03 ([M+H]$^+$)	8.10 (s, 1H), 7.52 (d, *J* = 8.0 Hz, 2H), 7.42-7.38 (m, 3H), 7.28 (s, 2H), 6.67 (d, *J* = 16.0 Hz, 1H), 6.45 (dd, *J* = 16.0, 8.0 Hz, 1H), 4.16 (m, 1H), 2.79 (s, 3H)	
DC25		442.02 ([M-H]$^-$)	7.97 (s, 1H), 7.59 (d, *J* = 8.0 Hz, 2H), 7.53 (d, *J* = 8.0 Hz, 2H), 7.38 (m, 1H), 7.29 (s, 2H), 6.65 (d, *J* = 16.0 Hz, 1H), 6.42 (dd, *J* = 16.0, 8.0 Hz, 1H), 4.17 (m, 1H), 2.74 (s, 3H)	1167, 1114, 800

Compound Number	mp (°C)	ESIMS	^{1}H NMR (δ)[a]	IR (cm^{-1})
DC26		464.03 ([M-H]$^-$)	8.12 (s, 1H), 7.49 (d, *J* = 8.0 Hz, 2H), 7.40-7.37 (m 3H), 7.28 (s, 2H), 6.66 (d, *J* = 16.0 Hz, 1H), 6.44 (dd, *J* = 16.0, 8.0 Hz, 1H), 4.14 (m, 1H), 3.22 (m, 1H), 1.09 – 1.16 (m, 4H)	1689, 1253, 1166, 1114, 979, 964
DC27		473.94 ([M-H]$^-$)	8.19 (s, 1H), 7.64 (d, *J* = 7.2 Hz, 2H), 7.55 (d, 7.2 Hz, 2H), 7.39 (s, 1H), 7.30 (s, 2H), 6.62 (d, *J* = 16.0 Hz, 1H), 6.42 (dd, *J* = 8.0, 16.0 Hz, 1H), 4.18 (m, 1H), 3.58 (s, 3H)	1571, 1331, 1170, 1113, 764
DC28		421.22 ([M+H]$^+$)	8.79 (s, 1H), 8.18 (s, 1H), 7.80 (m, 3H), 7.52 (m, 2H), 7.24 (m, 1H), 6.63 (d, *J* = 16.0 Hz, 1H), 6.54 (d, *J* = 16.0, 7.6 Hz, 1H), 4.19 (m, 1H)	3126, 2233, 1516, 1250, 1165, 1109
DC29		421.22 ([M+H]$^+$)	8.80 (s, 1H), 8.2 (s, 1H), 7.75 – 7.82 (m, 3H), 7.41 (t, *J* = 2 Hz, 1H), 7.26 (m, 2H), 6.65 (d, *J* = 16.0 Hz, 1H), 6.52 (dd, *J* = 16.0, 7.6 Hz, 1H), 4.16 (m, 1H)	3005, 1716, 1363, 1223
DC30		489.17 ([M+H]$^+$)	8.81 (s, 1H), 8.20 (s, 1H), 7.94 (s, 1H), 7.85 (m, 3H), 7.79 (m, 2H), 6.70 (d, *J* = 16.0 Hz, 1H), 6.58 (dd, *J* = 16.0, 8.0 Hz, 1H), 4.35 (m, 1H)	2964, 2234, 1289, 1166, 1136

Compound Number	mp (°C)	ESIMS	^{1}H NMR (δ)[a]	IR (cm^{-1})
DC31	117–118	455.27 ([M+H]$^+$)	8.80 (s, 1H), 8.20 (s, 1H), 7.82 (m, 3H), 7.4 (s, 2H), 6.62 (d, *J* = 16.0 Hz, 1H), 6.52 (dd, *J* = 16.0, 8.0 Hz, 1H), 4.18 (m, 1H)	
DC32		388.0705 (388.0703)	8.82 (s, 1H), 8.22 (s, 1H), 7.82-7.78 (m, 3H), 7.38-7.30 (m, 3H), 6.62 (d, *J* = 16.1 Hz, 1H), 6.56 (dd, *J* = 16.1, 6.8 Hz, 1H), 4.18 (m, 1H)	3126, 2234, 1520, 1280, 1164, 1112
DC33		455.22 ([M-H]$^-$)	8.80 (s, 1H), 8.20 (s, 1H), 7.82-7.80 (m, 3H), 7.70-7.50 (m, 3H), 6.65 (d, *J* = 16.9 Hz, 1H), 6.54 (dd, *J* = 16.9, 6.8 Hz, 1H), 4.25 (m, 1H)	3122, 3086, 2234, 1517, 1327, 1168, 1113
DC34		452.0412 (452.0419)	8.85 (s, 1H), 8.23 (br s, 1H), 7.83-7.78 (m, 3H), 7.33 (s, 2H), 6.69 (d, *J* = 14.9 Hz, 1H), 6.50 (dd, *J* = 14.9, 7.2 Hz, 1H), 4.15 (m, 1H), 3.90 (s, 3H)	3122, 2934, 2231, 1516, 1480, 1248, 1211, 1165, 1111
DC35		439.01 ([M-H]$^-$)	8.60 (s, 1H), 8.20 (s, 1H), 7.82 (m, 3H), 7.28 (m, 2H), 6.65 (d, *J* = 16.0 Hz, 1H), 6.48 (dd, *J* = 16.0, 8.0 Hz, 1H), 4.20 (m, 1H)	2233, 1518, 1250, 1169, 1035, 817
DC36		437.25 ([M+H]$^+$)	8.70 (s, 1H), 7.80 (m, 3H), 7.40 (s, 1H), 7.28 (s, 2H), 6.63 (d, *J* = 16.0 Hz, 1H), 6.50 (dd, *J* = 16.0, 8.0 Hz, 1H), 4.18 (m, 1H), 2.50 (s, 1H)	2927, 2233, 1572, 1531, 1248, 1166, 1112

Compound Number	mp (°C)	ESIMS	^{1}H NMR (δ)[a]	IR (cm^{-1})
DC37	109–111	466.10 ([M-H]$^-$)	8.86 (s, 1H), 7.89 (m, 3H), 7.40 (s, 1H), 7.30 (s, 2H), 6.68 (d, *J* = 16.0 Hz, 1H), 6.57 (dd, *J* = 16.0, 8.0 Hz, 1H), 4.18 (m, 1H)	
DC38	96–98	436.11 ([M-H]$^-$)	8.58 (s, 1H), 7.75 (m, 3H), 7.40 (s, 1H), 7.28 (s, 2H), 6.61 (d, *J* = 16.0 Hz, 1H), 6.42 (dd, *J* = 16.0, 8.2 Hz, 1H), 4.40 (br s, 2H), 4.15 (m, 1H)	
DC39	224–226	480.30 ([M+H]$^+$)	8.65 (s, 1H), 8.18 (br s, 1H), 7.80-7.70 (m, 3H), 7.40 (s, 1H), 7.27 (s, 2H), 7.36 (m, 1H), 7.28 (m, 2H), 6.60 (d, *J* = 16.8 Hz, 1H), 6.47 (m, 1H), 4.16 (m, 1H), 2.40 (br s, 3H)	3352, 2237, 1707, 1163, 841
DC40	70–73	436.11 ([M-2H]$^-$)	8.86 (s, 1H), 7.88 (m, 3H), 7.44 (s, 2H), 6.67 (d, *J* = 16.0 Hz, 1H), 6.56 (dd, *J* = 16.0 7.6 Hz, 1H), 4.19 (m, 1H)	
DC41	72–75	469.95 ([M-H]$^-$)	(DMSO-d_6) 8.72 (s, 1H), 8.26 (s, 1H), 8.01 (d, *J* = 8.4 Hz, 1H), 7.91 (s, 2H), 7.77 (d, *J* = 8.4 Hz, 1H), 6.42 (dd, *J* = 15.6, 9.2 Hz, 1H), 6.83 (d, *J* = 15.6 Hz, 1H), 5.87 (s, 2H), 4.89 (m, 1H)	
DC42	104–107	609.98 ([M+H]$^+$)	8.78 (s, 2H), 7.83 (s, 1H), 7.80 (m, 2H), 7.42 (s, 2H), 6.65 (d, *J* = 16.4 Hz, 1H), 6.51 (dd, *J* = 16.4, 7.8 Hz, 1H), 4.17 (m, 1H), 4 2.16 (m, 2H), 1.25 (m, 4H), 1.00 (m, 4H),	2234, 1714, 1114, 807

Compound Number	mp (°C)	ESIMS	^{1}H NMR (δ)[a]	IR (cm^{-1})
DC43	109–112	540.04 ([M+H]$^+$)	(DMSO-d_6) 10.94 (br s, 1H), 8.36 (s, 1H), 8.08 (m, *J* = 8.4 Hz, 1H), 7.91 (s, 2H), 7.84 (d, *J* = 8.4 Hz, 1H), 7.13 (dd, *J* = 15.6, 9.2 Hz, 1H), 6.87 (d, *J* = 15.6 Hz, 1H), 4.92 (m, 1H), 1.99 (br s, 1H), 0.82 (s, 4H)	3233, 2233, 1699, 1114, 807
DC44		435.26 [M-H]$^-$	8.33 (s, 1H), 8.23 (s, 1H), 7.66 (s, 1H), 7.60 (s, 1H), 7.41 (m, 1H), 7.28 (m, 2H), 6.62 (d, *J* = 16.0 Hz, 1H), 6.51 (dd, *J* = 16.0, 7.8 Hz, 1H), 4.16 (m, 1H), 2.20 (s, 3H)	2236,1510, 1114, 801
DC45	75-78	468.87 [M-H]$^-$	8.36 (s, 1H), 8.23 (s, 1H), 7.66 (s, 1H), 7.60 (s, 1H), 7.41 (s, 2H), 6.62 (d, *J* = 16.4 Hz, 1H), 6.51 (dd, *J* = 16.4, 7.6 Hz, 1H), 4.16 (m, 1H), 2.20 (s, 3H)	
DC46		411.4 ([M]$^+$)	8.83 (s, 1H), 8.21 (s, 1H), 7.83 (d, *J* = 8.5 Hz, 1H), 7.61 (d, *J* = 1.9 Hz, 1H), 7.52 (dd, *J* = 8.4, 1.9 Hz, 1H), 7.28 (d, *J* = 3.8 Hz, 2H), 6.93 (d, *J* = 11.5 Hz, 1H), 6.26 - 6.20 (m, 1H), 4.22 (m, 1H)	^{13}C NMR (δ)[3] 155.63, 153.27, 153.12, 143.01, 137.89, 136.25, 134.03, 133.88, 132.23, 131.23, 131.18, 129.20, 126.17, 125.04, 124.99

Compound Number	mp (°C)	ESIMS	^{1}H NMR (δ)[a]	IR (cm^{-1})
DC47	139–141	474.16 ([M-H]$^-$)	8.51 (s, 1H), 8.14 (s, 1H), 7.75 (s, 1H), 7.5 (m, 2H), 7.4 (s, 1H), 7.30 (m, 2H), 6.60 (d, *J* = 16.0 Hz, 1H), 6.50 (dd, *J* = 16.0, 8.0 Hz, 1H), 4.15 (m, 1H)	
DC48	124-126	414.05 [M-H]$^-$	8.69 (s, 1H), 8.14 (s, 1H), 7.96 (d, *J* = 4.8 Hz, 1H), 7.39-7.27 (m, 5H), 6.95 (d, *J* = 16.0 Hz, 1H), 6.51 (dd, *J* = 16.0, 7.6 Hz, 1H), 4.13 (m, 1H)	
DC49	81-83	463.96 [M-H]$^-$	8.57 (s, 1H), 8.14 (s, 1H), 7.60 (m, 2H), 7.44 (m, 3H), 6.95 (d, *J* = 16.0 Hz, 1H), 6.51 (dd, *J* = 16.0, 7.6 Hz, 1H), 4.13 (m, 1H)	
DC50	140-143	430.07 [M-H]$^-$)	8.56 (s, 1H), 8.13 (s, 1H), 7.59 (d, *J* = 1.2 Hz, 2H), 7.44 (m, 2H), 7.28 (m, 2H), 6.61 (d, *J* = 16.0 Hz, 1H), 6.47 (dd, *J* = 16.0, 8.0 Hz, 1H), 4.15 (m, 1H)	1110, 803
DC51	118–121	464.22 ([M-H]$^-$)	8.32 (s, 1H), 8.15 (s, 1H), 7.82 (s, 1H), 7.73 (d, *J* = 8.4 Hz, 1H), 7.53 (d, *J* = 8.4 Hz, 1H), 7.41 (s, 1H), 7.29 (s, 2H), 6.70 (d, *J* = 15.6 Hz, 1H), 6.50 (dd, *J* = 15.6, 8.0 Hz, 1H), 4.20 (m, 1H)	

Compound Number	mp (°C)	ESIMS	^{1}H NMR (δ)[a]	IR (cm^{-1})
DC52			9.99 (s, 1H), 8.42 (s, 1H), 8.12 (s, 1H), 8.01 (s, 1H), 7.68 (m, 1H), 7.44 (m, 1H), 7.33 (m, 1H), 7.22 (s, 2H), 6.62 (d, *J* = 16.7 Hz, 1H), 6.45 (dd, *J* = 16.7, 9.3 Hz, 1H), 4.10 (m, 1H)	3123, 3079, 2925, 1692, 1571, 1512, 1253, 1164, 1111
DC53			8.30 (m, 1H), 8.00 (br s, 1H), 7.75 (m, 1H),7.68 (m, 1H), 7.55 (m, 1H), 7.36 (m, 1H), 7.28 (m, 2H), 6.70 (m, 1H), 6.58 (br s, 1H), 6.33 (m, 1H), 5.88 (m, 2H), 4.10 (m, 1H)	3250, 3043, 1683, 1116
DC54	56–58	441.07 ([M-H]$^-$)	8.40 (s, 1H), 8.13 (s, 1H), 8.02 (s, 1H), 7.76 (d, *J* = 8.4 Hz, 1H), 7.59 (d, *J* = 8.0 Hz, 1H), 7.4 (s, 1H), 7.29 (m, 2H), 6.69 (d, *J* = 15.6 Hz, 1H), 6.57 (dd, *J* = 15.6, 7.8 Hz, 1H), 4.15 (m, 1H)	
DC55		412.97 ([M+H]$^+$)	8.37 (s, 1H), 8.18 (s, 1H), 7.39 (s, 1H), 7.30 (m, 2H), 7.19 (d, *J* = 8.0 Hz, 1H), 6.90 (m, 2H), 6.55 (d, *J* = 15.6 Hz, 1H), 6.38 (dd, *J* = 15.6, 8.2 Hz, 1H), 4.20 (m, 1H), 2.50 (br s, 2H)	
DC56	175–177	453 ([M-H]$^-$)	9.59 (br s, 1H), 8.55 (s, 1H), 8.47 (s, 2H), 8.23 (s, 1H), 7.30 (m, 4H), 6.62 (d, *J* = 16.0 Hz, 1H), 6.40 (dd, *J* = 16.0, 8.0 Hz, 1H), 4.15 (m, 1H), 2.20 (s, 3H)	

Compound Number	mp (°C)	ESIMS	^{1}H NMR (δ)[a]	IR (cm^{-1})
DC57		426.0627 (426.0626)	8.33 (s, 1H), 8.16 (s, 1H), 7.38 (s, 1H), 7.29 (s, 2H), 7.15 (d, *J* = 7.6 Hz, 1H), 6.80 (d, *J* = 7.6 Hz, 1H), 6.74 (m, 1H), 6.60 (d, *J* = 15.6 Hz, 1H), 6.35 (dd, *J* = 15.6, 8.4 Hz, 1H), 5.40 (br s, 1H), 4.15 (m, 1H), 2.90 (s, 3H)	3342, 3112, 2931, 1606, 1583, 1574, 1528, 1153
DC58	94–97	440.0424 (440.0419)	(DMSO-d_6) 8.76 (s, 1H), 8.16 (s, 1H), 7.90 (br s, 1H), 7.83 (s, 1H), 7.70 (d, *J* = 7.9 Hz, 1H), 7.71-7.67 (m, 3H), 7.58 (d, *J* = 7.9 Hz, 1H), 7.52 (br s, 1H), 7.00 (dd, *J* = 15.8, 8.7 Hz, 1H), 6.85 (d, *J* = 15.8 Hz, 1H), 4.85 (m, 1H)	3403, 3304, 3178, 1674, 1571, 1169, 1108
DC59	87–90		(DMSO-d_6) 9.00 (s, 1H), 8.63 (s, 1H), 8.17 (s, 1H), 7.70-7.59 (m, 5H), 7.00 (dd, *J* = 16.2, 9.7 Hz, 1H), 6.85 (d, *J* = 16.2 Hz, 1H), 5.90 (br s 2H), 4.83 (m, 1H)	
DC60		469.0577 (469.0572)	8.32 (s, 1H), 8.10 (s, 1H), 7.97 (s, 1H), 7.65 (d, *J* = 8.1 Hz, 1H), 7.47 (d, *J* = 8.1 Hz, 1H), 7.40 (m, 1H), 7.28 (s, 2H), 6.62 (d, *J* = 16.5 Hz, 1H), 6.49 (dd, *J* = 16.5, 7.7 Hz, 1H), 4.23-4.04 (m, 3H), 1.15 (t, *J* = 8.0 Hz, 3H)	2987, 1725, 1518, 1275, 1166, 1113

Compound Number	mp (°C)	ESIMS	^{1}H NMR (δ)[a]	IR (cm^{-1})
DC61	130–132	442.15 ([M+H]$^+$)	(DMSO-d_6) 9.90 (s, 1H), 8.17 (s, 1H), 8.15 (m, 1H), 7.90 (m, 1H), 7.71 (m, 2H), 7.67 (m, 1H), 7.62 (d, *J* = 7.3 Hz, 1H), 7.03 (dd, *J* = 16.5, 8.3 Hz, 1H), 6.62 (d, *J* = 16.5 Hz, 1H), 4.87 (m, 1H)	
DC62		412.10 ([M+H]$^+$)	8.27 (s, 1H), 8.23 (s, 1H), 7.40 (m, 3H), 7.30 (m, 3H), 6.64 (d, *J* = 16.0 Hz, 1H), 6.45 (dd, *J* = 16.0, 8.0 Hz, 1H), 4.19 (m, 1H), 2.21 (s, 3H)	1513, 1252, 1166, 1112, 801
DC63		446.01 ([M+H]$^+$)	8.26 (s, 1H), 8.12 (s, 1H), 7.42 (s, 2H), 7.18-7.28 (m, 3H), 6.62 (d, *J* = 15.6 Hz, 1H), 6.39 (dd, *J* = 15.6, 9.4 Hz, 1H), 4.10 (m, 1H), 2.25 (s, 3H)	2928, 2525,1249, 1169, 1114, 809
DC64		475.03 ([M+H]$^+$)	8.84 (d, *J* = 5.8 Hz, 2H), 8.33 (s, 1H), 8.20 (s, 1H), 7.75 (m, 1H), 7.60 (d, *J* = 28.6 Hz, 1H), 7.58-7.48 (m, 3H), 7.42 (m, 1H), 7.28 (s, 2H), 6.71 (d, *J* = 16.9 Hz, 1H), 6.39 (dd, *J* = 16.9, 8.2 Hz, 1H), 4.15 (m, 1H)	1683, 1167, 650, 479
DC65		412.05 ([M+H]$^+$)	8.55 (s, 1H), 8.12 (s, 1H), 7.55 (m, 3H), 7.39 (m, 1H), 7.30 (d, *J* = 1.6 Hz, 1H), 6.85 (d, *J* = 16.0 Hz, 1H), 6.41 (dd, *J* = 16.0, 8.0 Hz, 1H), 4.17 (m, 1H), 2.40 (s, 3H)	722, 111

Compound Number	mp (°C)	ESIMS	^{1}H NMR (δ)[a]	IR (cm^{-1})
DC66	60–61	468.26 ([M+H]$^+$)	8.59 (s, 1H), 8.14 (s, 1H), 7.94 (s, 1H), 7.70 (d, *J* = 8.0 Hz, 1H), 7.61 (d, *J* = 8.0 Hz, 1H), 7.43 (s, 2H), 7.23 (d, *J* = 16.0 Hz, 1H), 6.41 (dd, *J* = 16.0, 8.0 Hz, 1H), 4.20 (m, 1H)	
DC67	133–134	432.30 ([M+H]$^+$)	8.59 (s, 1H), 8.12 (s, 1H), 7.78 (br s, 1H), 7.71 (m, 1H), 7.62 (m, 1H), 7.39 (s, 1H), 7.32 (s, 2H), 7.03 (d, *J* = 16.0 Hz, 1H), 6.43 (dd, *J* = 16.0, 8.0 Hz, 1H), 0.21 (m, 1H)	800, 114
DC68		412.03 ([M+H]$^+$)	8.71 (s, 1H), 8.18 (s, 1H), 7.71 (d, *J* = 8.0 Hz, 2H), 7.55 (d, *J* = 8.0 Hz, 2H), 7.37 (s, 1H), 7.28 (m, 2H), 6.08 (d, *J* = 16.0 Hz, 1H), 4.26 (m, 1H), 2.05 (s, 3H)	
DC69	162–168	414.03 ([M+H]$^+$)	8.56 (s, 1H), 8.11 (s, 1H), 7.70 (d, *J* = 8.5 Hz, 2H), 7.56 (d, *J* = 8.5 Hz, 2H), 7.54 (m, 2H), 7.40 (m, 1H), 6.91 (d, *J* = 16.5 Hz, 1H), 6.66 (d, *J* = 16.5 Hz, 1H)	
DC70	99–103	428.05 ([M+H]$^+$)	8.58 (s, 1H), 8.13 (s, 1H), 7.73 (d, *J* = 8.7 Hz, 2H), 7.60 (d, *J* = 8.7 Hz, 2H), 7.46 (m, 2H), 7.42 (m, 1H), 6.85 (d, *J* = 16.2 Hz, 1H), 6.40 (d, *J* = 16.2 Hz, 1H), 3.42 (s, 3H)	

[a] ^{1}H NMR spectral data were acquired using a 400 MHz instrument in $CDCl_3$ except where noted. HRMS data are noted observed value (theoretical value).

Table 2A: Analytical Data for Compounds in Table 1A.

Compound Number	mp (°C); $[\alpha]_D^{25}$	ESIMS	^{1}H NMR (δ)[a]	IR (cm^{-1}); ^{19}F NMR (δ)
F1	53-64	655 ($[M+H]^+$)	rotomers δ 7.61 (d, *J* = 1.6 Hz, 1H), 7.50 (d, *J* = 7.9 Hz, 1H), 7.45 (dd, *J* = 6.4, 3.0 Hz, 0.5H), 7.41 (s, 2H), 7.37 (dd, *J* = 8.0, 1.6 Hz, 1H), 7.33 (t, *J* = 6.2 Hz, 0.5H), 6.53 (d, *J* = 15.9 Hz, 1H), 6.45 (s, 1H), 6.39 (dd, *J* = 15.9, 7.8 Hz, 1H), 4.21 - 4.01 (m, 1H), 3.96 (qd, *J* = 9.1, 6.4 Hz, 1.5H), 3.85 (td, *J* = 9.2, 6.5 Hz, 0.5 H) 1.69 (s, 6H)	
F2		608.92 ($[M+H]^+$)	(300 MHz, $CDCl_3$) δ 7.65 (d, *J* = 7.6 Hz, 1H), 7.43-7.40 (m, 2H), 7.36 (d, *J* = 8.4 Hz, 1H), 7.29 (m, 1H), 6.56 (d, *J* = 15.6 Hz, 1H), 6.43 (dd, *J* = 15.6, 7.2 Hz, 1H), 4.12-4.08 (m, 1H), 3.97-3.94 (m, 2H), 1.70 (s, 6H)	3368, 1682, 1162, 808
F3		588.90 ($[M+H]^+$)	(300 MHz, DMSO-d_6) δ 8.23 (s, 1H), 8.17 (broad s, 1H), 7.89 (s, 2H), 7.48-7.38 (m, 3H), 6.82 (dd, *J* = 15.6, 7.8 Hz, 1H), 6.74 (d, *J* = 15.6 Hz, 1H), 4.90-4.80 (m, 1H), 3.89-3.84 (m, 2H), 2.30 (s, 3H), 1.42 (s, 6H)	3394, 1678, 1163, 807

Compound Number	mp (°C); $[\alpha]_D^{25}$	ESIMS	^{1}H NMR (δ)[a]	IR (cm^{-1}); ^{19}F NMR (δ)
F4		642.99 ($[M+H]^+$)	(300 MHz, DMSO-d_6) δ 8.58 (s, 1H), 8.20 (t, *J* = 6.6 Hz, 1H), 7.98-7.89 (m, 4H), 7.80 (d, *J* = 8.1 Hz, 1H), 7.04 (dd, *J* = 15.6, 8.7 Hz, 1H), 6.88 (d, *J* = 15.6 Hz, 1H), 4.88-4.78 (m, 1H), 3.89-3.82 (m, 2H), 1.40 (s, 6H)	3460, 1677, 1165, 557
F5		640.9 ($[M+H]^+$)	7.62 (d, *J* = 1.7 Hz, 1H), 7.53 (d, *J* = 7.8 Hz, 1H), 7.42 - 7.29 (m, 3H), 6.91 (t, *J* = 6.7 Hz, 1H), 6.54 (bs, 1H) 6.53 (d, *J* = 15.9 Hz, 1H), 6.39 (dd, *J* = 15.9, 7.8 Hz, 1H), 4.74 (q, *J* = 7.1 Hz, 1H), 4.10 (m, 1H), 4.05 - 3.72 (m, 2H), 1.53 (d, *J* = 7.0 Hz, 3H)	3288, 1644, 1162
F6		640.9 ($[M+H]^+$)	7.62 (d, *J* = 1.6 Hz, 1H), 7.53 (d, *J* = 8.0 Hz, 1H), 7.41 (s, 2H), 7.38 (dd, *J* = 8.0, 1.7 Hz, 1H), 6.86 (t, *J* = 6.2 Hz, 1H), 6.57 - 6.49 (m, 2H), 6.40 (dd, *J* = 15.9, 7.8 Hz, 1H), 4.74 (m, 1H), 4.15 - 4.04 (m, 1H), 4.00 - 3.81 (m, 2H), 1.53 (d, *J* = 7.1 Hz, 3H)	3288, 1645, 1164

Compound Number	mp (°C); $[\alpha]_D^{25}$	ESIMS	^{1}H NMR (δ)[a]	IR (cm^{-1}); ^{19}F NMR (δ)
F7		574.92 ($[M+H]^+$)	(300 MHz, DMSO-d_6) δ 8.56 (t, *J* = 6.3 Hz, 1H), 8.43 (d, *J* = 4.5 Hz, 1H), 7.89 (s, 2H), 7.44-7.34 (m, 3H), 6.88 (dd, *J* = 15.6, 8.4 Hz, 1H), 6.75 (d, *J* = 15.6 Hz, 1H), 4.85-4.79 (m, 1H), 4.49-4.44 (m, 1H), 3.99-3.83 (m, 2H), 2.33 (s, 3H), 1.34 (d, *J* = 7.2 Hz, 3H)	3412, 1685, 1163, 809
F8		652.95 ($[M+H]^+$)	(400 MHz, DMSO-d_6) δ 8.62 (t, *J* = 6.0 Hz, 1H), 8.59 (d, *J* = 7.6 Hz, 1H), 7.92-7.91 (m, 3H), 7.60 (d, *J* = 7.6 Hz, 1H), 7.38 (d, *J* = 8.1 Hz, 1H), 6.99 (dd, *J* = 15.6, 9.2 Hz, 1H), 6.77 (d, *J* = 15.6 Hz, 1H), 4.85-4.80 (m, 1H), 4.41-4.36 (m, 1H), 4.05-3.99 (m, 1H), 3.91-3.84 (m, 1H), 1.73-1.63 (m, 2H), 0.99 (t, *J* = 7.6 Hz, 3H)	3291, 1647, 1165, 808, 565
F8A		650.99 ($[M-H]^-$)	(300 MHz, DMSO-d_6) δ 8.64-8.57 (m, 2H), 7.92-7.91 (m, 3H), 7.60 (d, *J* =7 .5 Hz, 1H), 7.38 (d, *J* = 7.5 Hz, 1H),7.01 (dd, *J* =1 5.6, 9.0 Hz, 1H), 6.78 (d, *J* = 15.6 Hz, 1H), 4.86-4.80 (m, 1H), 4.42-4.35 (m, 1H), 4.06-3.84 (m, 2H), 1.76-1.62 (m, 2H), 0.93 (t, *J* = 7.2 Hz, 3H).	3289, 1646, 1164, 808. 725, 649

Compound Number	mp (°C); $[\alpha]_D^{25}$	ESIMS	^{1}H NMR (δ)[a]	IR (cm^{-1}); ^{19}F NMR (δ)
F9		575.04 ($[M+H]^+$)	(400 MHz, DMSO-d_6) δ 8.56 (t, J = 6.0 Hz, 1H), 8.44 (d, J = 7.2 Hz, 1H), 7.89 (s, 2H), 7.45 (s, 1H), 7.41-7.35 (m, 2H), 6.87 (dd, J = 16.0, 9.2 Hz, 1H), 6.74 (d, J = 15.2 Hz, 1H), 4.85-4.80 (m, 1H), 4.48-4.44 (m, 1H), 4.03-3.85 (m, 2H), 2.33 (s, 3H), 1.31 (d, J = 7.2 Hz, 3H)	3407, 1685, 1161, 808
F10		638.84 ($[M+H]^+$)	(400 MHz, DMSO-d_6) δ 8.61 (d, J = 7.2 Hz, 1H), 8.52 (t, J = 6.0 Hz, 1H), 7.88-7.87 (m, 3H), 7.56 (d, J = 7.6 Hz, 1H), 7.38 (d, J = 8.0 Hz, 1H), 6.96 (dd, J = 16.6, 9.2 Hz, 1H), 6.73 (d, J = 15.6 Hz, 1H), 4.81-4.77 (m, 1H), 4.46-4.42 (m, 1H), 4.00-3.81 (m, 2H), 1.27 (d, J = 7.2 Hz, 3H)	3415, 1652, 1162, 807, 561
F11		638.90 ($[M+H]^+$)	(400 MHz, DMSO-d_6) δ 8.61 (d, J = 7.2 Hz, 1H), 8.52 (t, J = 6.0 Hz, 1H), 7.88-7.87 (m, 3H), 7.56 (d, J = 7.6 Hz, 1H), 7.38 (d, J = 8.0 Hz, 1H), 6.96 (dd, J = 16.6, 9.2 Hz, 1H), 6.73 (d, J = 15.6 Hz, 1H), 4.81-4.77 (m, 1H), 4.46-4.42 (m, 1H), 4.00-3.81 (m, 2H), 1.27 (d, J = 7.2 Hz, 3H)	3415, 1652, 1162, 807, 561

Compound Number	mp (°C); $[\alpha]_D^{25}$	ESIMS	^{1}H NMR (δ)[a]	IR (cm^{-1}); ^{19}F NMR (δ)
F12		638.90 ($[M+H]^+$)	(400 MHz, DMSO-d_6) δ 8.65 (d, J = 7.2 Hz, 1H), 8.56 (t, J = 8.0 Hz, 1H), 7.92 (s, 1H), 7.91 (s, 2H), 7.60-7.58 (m, 1H), 7.42 (d, J = 7.6 Hz, 1H), 6.99 (dd, J = 15.6, 8.0 Hz, 1H), 6.77 (d, J = 15.6 Hz, 1H), 4.83-4.76 (m, 1H), 4.52-4.45 (m, 1H), 4.06-3.82 (m, 2H), 1.33 (d, J = 8.0 Hz, 3H)	3418, 1646, 1163, 808, 564
F13		638.96 ($[M+H]^+$)	(400 MHz, DMSO-d_6) δ 8.65 (d, J = 7.2 Hz, 1H), 8.56 (t, J = 8.0 Hz, 1H), 7.92 (s, 1H), 7.91 (s, 2H), 7.60-7.58 (m, 1H), 7.42 (d, J = 7.6 Hz, 1H), 6.99 (dd, J = 15.6, 8.0 Hz, 1H), 6.77 (d, J = 15.6 Hz, 1H), 4.83-4.76 (m, 1H), 4.52-4.45 (m, 1H), 4.06-3.82 (m, 2H), 1.33 (d, J = 8.0 Hz, 3H)	3418, 1646, 1163, 808, 564
F14		673.31 ($[M+H]^+$)	(300 MHz, $CDCl_3$) δ 7.91 (s, 1H), 7.82 (s, 2H), 7.62 (s, 1H), 7.53 (d, J = 7.5 Hz, 1H), 7.40 (d, J = 8.1 Hz,1H), 6.80 (bs, 1H), 6.62 (s, 1H), 6.55 (d, J = 16.0 Hz, 1H), 6.50 (dd, J = 16.0, 8.0 Hz, 1H), 4.73 -4.70 (m, 1H), 4.40 - 4.25 (m, 1H), 3.95-3.92 (m, 2H), 1.56 (d, J = 7.5 Hz, 3H)	3422, 1640, 1169, 528

Compound Number	mp (°C); $[\alpha]_D^{25}$	ESIMS	^{1}H NMR (δ)[a]	IR (cm^{-1}); ^{19}F NMR (δ)
F15		589.00 ($[M+H]^+$)	(400 MHz, DMSO-d_6) δ 8.58 (t. J = 8.8 Hz, 1H), 8.32 (d, J = 8.0 Hz, 1H), 7.85 (s, 2H), 7.42 (s, 1H), 7.37 (d, J = 8.0 Hz, 1H), 7.31 (d, J = 7.6 Hz, 1H), 6.83 (dd, J = 15.6 Hz, 8.8 Hz, 1H), 6.71 (d, J = 15.6 Hz, 1H), 4.77-4.81 (m, 1H), 4.30-4.35 (m, 1H), 3.94-4.00 (m, 1H), 3.80-3.86 (m, 1H), 2.29 (s, 3H), 1.71-1.60 (m, 2H), 0.88 (t, J = 7.6 Hz, 3H)	3295, 1682, 1164, 80
F15A		588.9 ($[M+H]^+$)	(300 MHz, DMSO-d_6) δ 8.61 (t, J = 6.0 Hz, 1H), 8.35 (d, J = 7.5 Hz, 1H), 7.89 (s, 2H), 7.45 (s, 1H), 7.41 (d, J = 8.4 Hz, 1H), 7.34 (d, J = 7.8 Hz, 1H), 6.88 (dd, J = 15.9, 8.7 Hz, 1H), 6.75 (d, J = 15.9 Hz, 1H), 4.85-4.79 (m, 1H), 4.40-4.32 (m, 1H), 4.04-3.82 (m, 2H), 2.33 (s, 3H),1.76-1.62 (m, 2H), 0.91 (t, J = 7.5 Hz, 3H)	3290, 1646, 1165, 808. 725, 651

Compound Number	mp (°C); $[\alpha]_D^{25}$	ESIMS	^{1}H NMR (δ)[a]	IR (cm^{-1}); ^{19}F NMR (δ)
F16		643.06 ($[M+H]^+$)	(400 MHz, DMSO-d_6) δ 8.65-8.60 (m, 2H), 7.95 (s, 1H), 7.89-7.82 (m, 3H), 7.47 (d, *J* = 8.0 Hz, 1H), 7.04 (dd, *J* = 15.6, 9.2 Hz, 1H), 6.85 (d, *J* = 15.9 Hz, 1H), 4.85-4.80 (m, 1H), 4.38-4.33 (m, 1H), 4.01 (m, 2H), 1.72-1.58 (m, 2H), 0.86 (t, *J* = 7.6 Hz, 3H)	3292, 1652, 1167, 809
F16A		640.9 ($[M-H]^-$)	(300 MHz, DMSO-d_6) δ 8.66-8.62 (m, 2H), 7.98 (s,1H), 7.92-7.87 (m, 3H), 7.51 (d, *J* = 7.8 Hz, 1H),7.04 (dd, *J* = 15.6, 9.0 Hz, 1H), 6.89 (d, *J* = 15.6 Hz, 1H), 4.86-4.79 (m, 1H), 4.41-4.37 (m, 1H), 4.05-3.87 (m, 2H), 1.72-1.63 (m, 2H), 0.90 (t, *J* = 6.9 Hz, 3H)	3290, 1651, 1166, 809
F17		628.95 ($[M+H]^+$)	(400 MHz, DMSO-d_6) δ 8.55 (t, *J* = 6.4 Hz, 1H), 8.42 (d, *J* = 7.6 Hz, 1H), 7.89 (s, 1H), 7.82 (s, 2H), 7.45 (s, 1H), 7.41-7.35 (m, 2H), 6.85 (dd, *J* = 16.0 Hz, 8.8 Hz, 1H), 6.74 (d, *J* = 15.6 Hz, 1H), 4.81-4.76 (m, 1H), 4.48-4.44 (m, 1H), 3.99-3.84 (m, 2H), 2.33 (s, 3H), 1.30 (d, *J* = 7.6 Hz, 3H)	3299, 1686, 1165, 568

Compound Number	mp (°C); $[\alpha]_D^{25}$	ESIMS	^{1}H NMR (δ)[a]	IR (cm^{-1}); ^{19}F NMR (δ)
F18		629.0 ($[M+H]^+$)	(300 MHz, DMSO-d_6) δ 8.76 (d, J = 7.8 Hz, 1H), 8.61 (t, J = 12.9 Hz, 1H), 7.98 (s, 1 H), 7.92-7.88 (m, 3H), 7.56 (d, J = 8.1 Hz, 1H), 7.10 (dd, J = 16.2, 9.3 Hz, 1H), 6.89 (d, J = 16.0 Hz, 1H), 4.89-4.83 (m, 1H), 4.51-4.47 (m, 1H), 4.01-3.88 (m, 2H), 1.29 (d, J = 7.2 Hz, 3H)	3407, 1713, 1160, 807
F19		692.88 ($[M+H]^+$)	(400 MHz, DMSO-d_6) δ 8.61 (d, J = 7.2 Hz, 1H), 8.52 (t, J = 12.8 Hz, 1H), 7.88 (s, 1H), 7.85 (s, 1H), 7.80 (s, 2H), 7.56 (d, J = 6.8 Hz, 1H), 7.38 (d, J = 8.0 Hz, 1H), 6.95 (dd, J = 15.6, 9.2 Hz, 1H), 6.72 (d, J = 15.6 Hz, 1H), 4.77-4.72 (m, 1H), 4.46-4.42 (m, 1H), 3.98-3.82 (m, 2H), 1.27 (d, J = 6.8 Hz, 3H)	3404, 1646, 1165, 565
F20		629.29 ($[M+H]^+$)	(400 MHz, DMSO-d_6) δ 8.75 (d, J = 7.8 Hz, 1H), 8.59 (t, J = 6.6 Hz, 1H), 7.98-7.88 (m, 4H), 7.56 (d, J = 8.1 Hz, 1H), 7.09 (dd, J = 15.6, 8.7 Hz, 1H), 6.89 (d, J = 15.9 Hz, 1H), 4.89-4.83 (m, 1H), 4.52-4.47 (m, 1H), 4.01-3.88 (m, 2H), 1.30 (d, J = 6.9 Hz, 3H)	3407, 1679, 1165, 808

Compound Number	mp (°C); $[\alpha]_D^{25}$	ESIMS	^{1}H NMR (δ)[a]	IR (cm^{-1}); ^{19}F NMR (δ)
F20A	$[\alpha]_D^{25}$ = +49.2 (*c*, 1% in CH_2Cl_2)	628.89 ($[M+H]^+$)	(400 MHz, DMSO-d_6) δ 8.74 (d, *J* = 7.6 Hz, 1H), 7.98 (s, 1H), 7.92 - 7.89 (m, 3H), 7.56 (d, *J* = 8.4 Hz, 1H), 7.08 (dd, *J* = 15.6, 8.8 Hz, 1H), 6.88 (d, *J* = 15.6 Hz, 1H), 4.88 - 4.83 (m, 1H), 4.51 - 4.48 (m, 1H), 3.99 - 3.86 (m, 2H), 1.49 (d, *J* = 6.8 Hz, 3H)	3315, 1657, 1167, 700
F20B	$[\alpha]_D^{25}$ = -38.8 (*c*, 1% in CH_2Cl_2)	628.89 ($[M+H]^+$)	(400 MHz, DMSO-d_6) δ 8.74 (d, *J* = 7.6 Hz, 1H), 7.98 (s, 1H), 7.92 - 7.89 (m, 3H), 7.56 (d, *J* = 8.4 Hz, 1H), 7.08 (dd, *J* = 15.6, 8.8 Hz, 1H), 6.88 (d, *J* = 15.6 Hz, 1H), 4.88 - 4.83 (m, 1H), 4.51 - 4.48 (m, 1H), 3.99 - 3.86 (m, 2H), 1.49 (d, *J* = 6.8 Hz, 3H)	3315, 1657, 1167, 700
F20C	61- 70	629 ($[M+H]^+$)	7.64 (t, *J* = 6.4 Hz, 1H), 7.52 (d, *J* = 7.9 Hz, 1H), 7.42 (s, 1H), 7.34 (d, *J* = 2.9 Hz, 3H), 6.64 (m, 1H), 6.91 (d, *J* = 11.4 Hz, 1H), 6.19 (dd, *J* = 11.4, 10.3 Hz, 1H), 4.94 (p, *J* = 7.1 Hz, 1H), 4.18 (q, *J* = 8.8 Hz, 1H), 3.95 - 3.71 (m, 2H), 1.52 (d, *J* = 6.9 Hz, 3H)	**^{19}F NMR** (376 MHz, CDCl3) δ -59.22, amide rotamers -69.43 and -69.45, -72.60

Compound Number	mp (°C); $[\alpha]_D^{25}$	ESIMS	^{1}H NMR (δ)[a]	IR (cm^{-1}); ^{19}F NMR (δ)
F21		682.98 ($[M+H]^+$)	(400 MHz, DMSO-d_6) δ 8.70 (d, J = 7.2 Hz, 1H), 8.55 (t, J = 6.4 Hz, 1H), 7.94 (s, 1H), 7.87-7.81 (m, 4H), 7.52 (d, J = 8.4 Hz, 1H), 7.04 (dd, J = 15.6, 9.2 Hz, 1H), 6.84 (d, J = 15.6 Hz, 1H), 4.80-4.75 (m, 1H), 4.47-4.44 (m, 1H), 3.97-3.78 (m, 2H), 1.29 (d, J = 7.6 Hz, 3H)	3302, 1656, 1166, 557
F22		605.32 ($[M+H]^+$)	(300 MHz, DMSO-d_6) δ 8.65 (d, J = 7.5 Hz, 1H), 8.55 (t, J = 8.1 Hz, 1H), 7.93-7.87 (m, 3H), 7.78 (d, J = 8.1 Hz, 1H), 7.71-7.66 (m, 1H), 7.61 (d, J = 8.1 Hz, 1H), 7.41 (d, J = 6.0 Hz, 1H), 7.03 (dd, J = 15.6, 9.2 Hz, 1H), 6.79 (d, J = 15.9 Hz, 1H), 4.93-4.86 (m, 1H), 4.50-4.45 (m, 1H), 4.00-3.85 (m, 2H), 1.33 (d, J = 7.2 Hz, 3H)	3412, 1645, 1164, 597
F23		609.00 ($[M+H]^+$)	7.66 (d, J = 8.4 Hz, 1H), 7.43 (s, 1H), 7.40 (s, 2H), 7.34 (d, J = 7.6 Hz, 1H), 6.79 (s, 1H), 6.73 (s, 1H), 6.56 (d, J = 16.4 Hz, 1H), 6.43 (dd, J = 16.4, 8.0 Hz, 1H), 4.64-4.60 (m, 1H), 4.13-4.06 (m, 1H), 4.00-3.85 (m, 2H), 2.09 (m, 1H), 1.87 -1.78 (m, 2H), 1.37 (t, J = 7.2 Hz, 3H)	3291, 1645, 1165, 749

Compound Number	mp (°C); $[\alpha]_D^{25}$	ESIMS	^{1}H NMR (δ)[a]	IR (cm^{-1}); ^{19}F NMR (δ)
F23A		608.99 ($[M+H]^+$)	(400 MHz, DMSO-d_6) δ 8.63 (t, J = 6.4 Hz, 1H), 8.59 (d, J = 8.0 Hz, 1H), 7.91 (s, 2H), 7.77 (s, 1H), 7.56 (d, J = 8.0 Hz, 1H), 7.42 (d, J = 8.0 Hz, 1H), 7.00 (dd, J = 15.6 , 9.2 Hz, 1H), 6.78 (d, J = 15.6 Hz, 1H), 4.86 - 4.81 (m, 1H), 4.42 - 4.37 (m, 1H), 4.05 - 3.84 (m, 2H), 1.75 - 1.61 (m, 2H), 0.92 (t, J = 7.2 Hz, 3H)	3292, 1646, 1165, 808
F24		622.97 ($[M+H]^+$)	(300 MHz, DMSO-d_6) δ 8.65 (d, J = 7.5 Hz, 1H), 8.56 (bs, 1H), 7.92-7.85 (m, 3H), 7.76-7.73 (m, 1H), 7.61 (d, J = 7.8 Hz, 1H), 7.42 (d, J = 7.8 Hz, 1H), 7.04 (dd, J = 15.6, 9.0 Hz, 1H), 6.08 (d, J = 15.9 Hz, 1H), 4.98-4.92 (m, 1H), 4.50-4.46 (m, 1H), 4.00-3.88 (m, 2H), 1.31 (d, J = 7.2 Hz, 3H)	3421, 1646, 1168
F25		594.94 ($[M+H]^+$)	(300 MHz, DMSO-d_6) δ 8.66 (d, J = 7.2 Hz, 1H), 8.57 (t, J = 6.0 Hz, 1H), 7.91 (s, 2H), 7.77 (s, 1H), 7.57 (d, J = 6.9 Hz, 1H), 7.46 (d, J = 7.8 Hz, 1H), 7.02 (dd, J = 15.6, 9.0 Hz, 1H), 6.78 (d, J = 15.6 Hz, 1H), 4.87-4.80 (m, 1H), 4.51-4.47 (m, 1H), 4.04-3.88 (m, 2H), 1.31 (d, J = 7.2 Hz, 3H)	3299, 1687, 1164, 808

Compound Number	mp (°C); $[\alpha]_D^{25}$	ESIMS	^{1}H NMR (δ)[a]	IR (cm^{-1}); ^{19}F NMR (δ)
F26		588.96 ($[M-H]^-$)	(300 MHz, DMSO-d_6) δ 10.54 (bs, 1H), 8.53 (d, J = 6.9 Hz, 1H), 7.89 (s, 2H), 7.45-7.38 (m, 3H), 6.88 (dd, J = 15.9, 8.4 Hz, 1H), 6.75 (d, J = 15.9 Hz, 1H), 4.87-4.79 (m, 2H), 4.67-4.65 (m, 1H), 4.53-4.50 (m, 1H), 2.33 (s, 3H), 1.37 (d, J = 6.9 Hz, 3H)	3429, 1645, 1165, 750
F27		655.27 ($[M+H]^+$)	(300 MHz, DMSO-d_6) δ 10.52 (bs, 1H), 8.74 (d, J = 5.7 Hz, 1H), 7.93-7.91 (m, 3H), 7.62 (d, J = 8.7 Hz, 1H), 7.48 (d, J = 7.8 Hz, 1H), 6.98 (dd, J = 15.0, 6.3 Hz, 1H), 6.77 (d, J = 15.6 Hz, 1H), 4.88-4.80 (m, 2H), 4.52-4.43 (m, 2H), 1.37 (d, J = 6.9 Hz, 3H)	3422, 1657, 1161, 806, 557
F28		644.94 ($[M+H]^+$)	(300 MHz, DMSO-d_6) δ 10.57 (bs, 1H), 8.85 (d, J = 7.5 Hz, 1H), 7.99-7.70 (m, 4H), 7.65 (d, J = 8.1 Hz, 1H), 7.07 (dd, J = 15.9, 6.9 Hz, 1H), 6.89 (d, J = 15.6 Hz, 1H), 4.88-4.84 (m, 2H), 4.67-4.50 (m, 2H), 1.36 (d, J = 6.9 Hz, 3H)	3257, 1663, 1170, 809, 552

Compound Number	mp (°C); $[\alpha]_D^{25}$	ESIMS	^{1}H NMR (δ)[a]	IR (cm^{-1}); ^{19}F NMR (δ)
F29		602.94 ($[M-H]^-$)	(300 MHz, DMSO-d_6) δ 9.87 (bs, 1H), 8.40 (bs, 1H), 7.89 (s, 2H), 7.54 (d, J = 8.1 Hz, 1H), 7.45 (s, 1H), 7.41 (d, J = 7.8 Hz, 1H), 6.88 (dd, J = 15.9, 8.7 Hz, 1H), 6.75 (d, J = 15.3 Hz, 1H), 4.85-4.79 (m, 1H), 4.55-4.50 (m, 2H), 2.32 (s, 3H), 1.60 (s, 6H)	3420, 1645, 1164 , 750
F30		611.0 ($[M+H]^+$)	(300 MHz, DMSO-d_6) δ 10.65 (bs, 1H), 8.75 (d, J = 6.9 Hz, 1H), 7.90 (s, 2H), 7.78 (s, 1H), 7.58-7.49 (m, 2H), 6.98 (dd, J = 14.7 Hz, 8.0 Hz, 1H), 6.78 (d, J = 15.3 Hz, 1H), 4.89-4.84 (m, 2H), 4.78-4.60 (m, 1H), 4.59-4.49 (m, 1H), 1.37 (d, J = 6.9 Hz, 3H)	3432, 1651, 1259, 750
F31		618.87 ($[M-H]^-$)	(300 MHz, DMSO-d_6) δ 10.14 (s, 1H), 9.72 (broad s, 1H), 7.88 (s, 2H), 7.40-7.34 (m, 2H), 7.24 (d, J = 7.8 Hz, 1H), 6.81 (dd, J = 16.0, 8.0 Hz, 1H) 6.69 (d, J = 16.0 Hz, 1H), 4.84-4.78 (m, 1H), 4.55-4.50 (m, 2H), 2.29 (s, 3H), 1.76 (s, 6H)	3436, 1261, 750

[a] ^{1}H NMR spectral data were acquired using a 400 MHz instrument in $CDCl_3$ except where noted. HRMS data are noted observed value (theoretical value).

Table 2B: Analytical Data for Compounds in Table 1B.

Compound Number	mp (°C)	ESIMS	^{1}H NMR (δ)[a]	IR (cm^{-1}); ^{19}F NMR (δ)
P1		590.91 ([M+H]$^+$)	(400 MHz, DMSO-d_6) δ 8.69 (t, J = 5.6 Hz, 1H), 8.56 (t, J = 6.4 Hz, 1H), 7.93-7.88 (m, 3H), 7.77-7.75 (m, 1H), 7.70-7.66 (m, 1H), 7.62-7.60 (m, 1H), 7.45 (d, J = 8.0 Hz, 1H), 7.03 (dd, J = 15.6, 8.8 Hz, 1H), 6.78 (d, J = 15.6 Hz, 1H), 4.92-4.87 (m, 1H), 3.97-3.90 (m, 4H)	3327, 1703, 1164, 595
P2		581.91 ([M-H]$^-$)	7.63-7.62 (m, 1H), 7.58 (d, J = 8.4 Hz, 1H), 7.39-7.37 (m, 1H), 7.12 (d, J = 8.0 Hz, 2H), 6.78 (t, J = 4.8 Hz, 1H), 6.65 (bs, 1H), 6.59 (d, J = 16.0 Hz, 1H), 6.39 (dd, J = 15.6, 7.6 Hz, 1H), 4.24-4.20 (m, 3H), 4.00-3.92 (m, 2H)	3280, 2243, 1637, 1166
P12		577.07 ([M+H]$^+$)	(400 MHz, DMSO-d_6) δ 8.61-8.54 (m, 2H), 7.86 (d, J = 6.4 Hz, 2H), 7.53 (d, J = 8.0 Hz, 1H), 7.45-7.42 (m, 2H), 6.95 (dd, J = 15.6, 9.6 Hz, 1H), 6.80 (d, J = 15.6 Hz, 1H), 4.86-4.76 (m, 1H), 3.98-3.89 (m, 4H), 2.43 (s, 3H)	3311, 1646, 1164, 714

P14		638.80 ([M-H]$^-$)	(300 MHz, DMSO-d_6) δ 10.55 (t, J = 6.0 Hz, 1H), 10.53 (t, J = 5.2 Hz, 1H), 7.90 (s, 1H), 7.87 (d, J = 8.8 Hz, 2H), 7.59-7.56 (m, 1H), 7.37-7.35 (m, 1H), 6.98 (dd, J = 15.6, 9.0 Hz, 1H), 6.76 (d, J = 15.9 Hz, 1H), 4.84-4.77 (m, 1H), 4.70-4.58 (m, 4H)	3435, 1166, 749, 597
P15		590.8 ([M-H]$^-$)	(400 MHz, DMSO-d_6) δ 10.62 (t, J = 6.0 Hz, 1H), 10.54 (t, J = 5.2 Hz, 1H), 7.89 (s, 2H), 7.42 (s, 1H), 7.39-7.37 (m, 1H), 7.24 (d, J = 7.6 Hz, 1H), 6.84 (dd, J = 15.6, 8.8 Hz, 1H), 6.74 (d, J = 15.6, 1H), 4.84-4.80 (m, 1H), 4.71 (d, J = 6.0 Hz, 2H), 4.65-4.56 (m, 2H), 2.35 (s, 3H)	3243, 2923, 1106, 614
P82		590.9 ([M+H]$^+$)	(300 MHz, DMSO-d_6) δ 8.65 (d, J = 7.2 Hz, 1H), 8.56 (t, J = 6.6 Hz, 1H), 7.88 (s, 1H), 7.59 - 7.56 (m, 2H), 7.47 - 7.40 (m, 2H), 6.90 - 6.88 (m, 2H), 4.97- 4.94 (m, 1H), 4.50 - 4.46 (m, 1H), 4.00 - 3.88 (m, 2H), 1.31 (d, J = 7.2 Hz, 3H)	3298, 1643, 1162
P84	82-85	581.1 ([M+H]$^+$)	(300 MHz, DMSO-d_6) δ 8.74 (d, J = 7.2 Hz, 1H), 8.59 (t, J = 6.6 Hz, 1H), 7.96 (s, 1H), 7.89 (d, J = 7.5 Hz, 1H), 7.61 - 7.44 (m, 3H) 6.99 (m, 2H), 4.98 - 4.95 (m, 1H), 4.51 - 4.47 (m, 1H), 3.99 - 3.88 (m, 2H), 1.29 (d, J = 6.8 Hz, 3H)	

P156		639.0 ($[M+H]^+$)	(400 MHz, DMSO-d_6) δ 8.66 (d, J = 8.0 Hz, 1H), 8.56 (t, J = 5.6 Hz, 1H), 7.95 - 7.93 (m, 2H), 7.60 (d, J = 7.6 Hz, 1H), 7.42 (d, J = 8.0 Hz, 1H), 7.03 (dd, J = 15.6, 8.8 Hz, 1H), 6.94 (d, J = 15.6 Hz, 1H), 5.09 - 5.04 (m, 1H), 4.53 - 4.43 (m, 1H), 4.02 - 3.86 (m, 2H), 1.31 (t, J = 6.8 Hz, 3H)	3436, 2924, 1662, 1162, 750
P226		620.95 ($[M+H]^+$)	(300 MHz, DMSO-d_6) δ 8.65 (d, J = 7.5 Hz 1H), 8.57 (t, J = 6.3 Hz, 1H), 7.90 (s, 1H), 7.60-7.55 (m, 4H),7.41(d, J = 7.8 Hz, 2H), 6.99 (dd, J = 15.6, 9.0 Hz, 1H), 6.78 (d, J = 15.9 Hz, 1H), 4.85-4.79 (m, 1H), 4.50-4.43 (m,1H), 4.00-3.85 (m, 2H), 1.33 (d, J = 7.8 Hz, 3H)	3413, 1668, 1161
P228		610.94 ($[M+H]^+$)	(300 MHz, DMSO-d_6) δ 8.74 (d, J = 7.5 Hz, 1H), 8.61 (t, J = 6.6 Hz, 1H), 7.96 (s, 1H), 7.91 (d, J = 8.1 Hz,1H), 7.63 - 7.53 (m, 4H), 7.41 (d, J = 7.5 Hz, 1H),7.08 (dd, J = 15.6, 8.7 Hz, 1H), 6.89 (d, J = 15.6 Hz, 1H), 4.84 - 4.81 (m, 1H), 4.51 - 4.43 (m, 1H), 3.99 - 3.85 (m, 2H), 1.29 (d, J = 7.5 Hz, 3H)	3413, 1668, 1161, 564

P298		634.8 ([M+H]$^+$)	(400 MHz, DMSO-d_6) δ 8.65 (d, J = 7.6 Hz, 1H), 8.55 (t, J = 6.4 Hz, 1H), 7.92 - 7.91 (d, J = 1.6 Hz, 1H), 7.67 (s, 1H), 7.62 - 7.58 (m, 2H), 7.54 (d, J = 9.6 Hz, 1H), 7.41 (d, J = 8.0 Hz, 1H), 6.97 (dd, J = 15.6, 9.2 Hz, 1H), 6.77 (d, J = 15.6 Hz, 1H), 4.82 - 4.77 (m, 1H), 4.50 - 4.46 (m, 1H), 4.00 - 3.82 (m, 2H), 1.31 (t, J = 7.2 Hz, 3H)	3307, 2925, 1652, 1164
P300		623.2 ([M+H]$^+$)	(400 MHz, DMSO-d_6) δ 8.74 (d, J = 7.6 Hz, 1H), 8.60 (d, J = 16.0 Hz, 1H), 7.97 (s, 1H), 7.90 (d, J = 8.0 Hz, 1H), 7.68 (s, 1H), 7.62 - 7.59 (m, 1H), 7.56 - 7.53 (m, 2H), 7.06 (dd, J = 15.6, 9.2 Hz, 1H), 6.88 (d, J = 15.6 Hz, 1H), 4.84 - 4.80 (m, 1H), 4.51 - 4.47 (m, 1H), 4.03 - 3.85 (m, 2H), 1.29 (d, J = 7.2 Hz, 3H)	3296, 1652, 1167
P442		584.87 ([M+H]$^+$)	(400 MHz, DMSO-d_6) δ 8.64 (d, J = 7.2 Hz, 1H), 8.57 (t, J = 12.8 Hz, 1H), 7.87 (d, J = 1.2 Hz, 1H), 7.58 - 7.52 (m, 2H), 7.48 (d, J = 8.4 Hz, 1H), 7.41 - 7.37 (m, 2H), 6.90 (dd, J = 16.0, 8.8 Hz, 1H), 6.74 (d, J = 15.6 Hz, 1H), 4.68 - 4.63 (m, 1H), 4.49 - 4.46 (m, 1H), 3.99 - 3.86 (m, 2H), 2.35 (s, 3H), 1.28 (d, J = 7.2 Hz, 3H)	3410, 1692, 1163, 769, 565

P444		574.98 ($[M+H]^+$)	(400 MHz, DMSO-d_6) δ 8.73 (d, J = 8.0 Hz, 1H), 8.60 (t, J = 6.0 Hz, 1H), 7.92 (s, 1H), 7.88 (d, J = 8.4 Hz, 1H), 7.55 - 7.53 (m, 2H), 7.48 (d, J = 8.4 Hz, 1H), 7.40 (d, J = 8.4 Hz, 1H), 6.98 (dd, J = 15.6, 8.8 Hz, 1H), 6.86 (d, J = 15.6 Hz, 1H), 4.70 - 4.66 (m, 1H), 4.50 - 4.47 (m, 1H), 3.99 - 3.82 (m, 2H), 2.35 (s, 3H), 1.27 (d, J = 7.2 Hz, 3H)	3292, 1661, 1158, 741
P514		583.0 ($[M+H]^+$)	(300 MHz, DMSO-d_6) δ 8.64 (d, J = 7.2 Hz, 1H), 8.57 (t, J = 6.3 Hz, 1H), 7.86 (s,1H), 7.58 (d, J = 7.5 Hz,1H), 7.41 (d, J = 8.1 Hz, 1H), 7.26 (d, J = 6.6 Hz, 2H), 6.88 (dd, J = 15.9, 8.4 Hz, 1H), 6.73 (d, J = 15.9 Hz, 1H), 4.58 - 4.46 (m, 2H), 4.00 - 3.85 (m, 2H), 2.24 (s, 6H), 1.31 (d, J = 7.5 Hz, 3H)	3276 , 1638, 1167, 598
P516		573.37 ($[M+H]^+$)	(300 MHz, DMSO-d_6) δ 8.73 (d, J = 7.8 Hz, 1H), 8.61 (t, J = 6.3 Hz, 1H), 7.88 (d, J = 8.4 Hz, 2H), 7.55 (d, J = 8.1 Hz, 1H), 7.27 (d, J = 6.9 Hz, 2H), 6.98 (dd, J = 15.6, 8.4 Hz, 1H), 6.85 (d, J = 15.6 Hz, 1H), 4.57 - 4.46 (m, 2H), 3.99 - 3.85 (m, 2H), 2.24 (s, 6H), 1.29 (d, J = 7.2 Hz, 3H)	3299, 1654, 1165

P568		692.88 ([M+H]$^+$)	(300 MHz, DMSO-d_6) δ 8.65 (d, J = 7.8 Hz, 1H), 8.55 (t, J = 7.8 Hz, 1H), 7.98 (s, 1H), 7.90 (s, 1H), 7.85 (d, J = 8.7 Hz, 1H), 7.60 - 7.57 (m, 1H), 7.51-7.48 (m, 1H), 7.41 - 7.39 (m, 1H), 6.96 (dd, J = 15.6, 8.7 Hz, 1H), 6.76 (d, J = 15.9 Hz, 1H), 4.81 - 4.72 (m, 1H), 4.50 - 4.46 (m, 1H), 4.01 - 3.82 (m, 2H), 1.31 (d, J = 6.9 Hz, 3H)	3306, 1646, 1164
P586		646.9 ([M-H]$^-$)	7.64 (s, 1H), 7.56 - 7.54 (m, 2H), 7.43 - 7.39 (m, 2H), 7.33 (s, 1H), 6.81 (bs, 1H), 6.57 (d, J = 16.0 Hz, 1H), 6.50 (d, J = 8.0 Hz, 1H), 6.45 (dd, J = 16.0, 7.6 Hz, 1H), 4.75 - 4.72 (m, 1H), 4.14 - 4.10 (m, 1H), 4.00 - 3.90 (m, 2H), 1.30 (d, J = 7.2 Hz, 3H)	3384, 1647, 1164, 749, 566
P588		638.84 ([M+H]$^+$)	(300 MHz, $CDCl_3$) δ 7.69 (s, 1H), 7.62-7.59 (m, 1H), 7.55 - 7.49 (m, 2H), 7.42 (s, 1H), 7.32 (s, 1H), 6.88 (bs, 1H), 6.65 (d, J = 16.2 Hz, 1H), 6.49 (dd, J = 16.2, 8.1 Hz, 1H), 6.34 (d, J = 7.2 Hz, 1H), 4.76 - 4.71 (m, 1H), 4.15 - 4.12 (m, 1H), 3.96 - 3.89 (m, 2H), 1.49 (d, J = 7.3 Hz, 3H)	3304, 2928, 1650, 1165, 673, 558

P660		682.8 ([M+H]$^+$)	(400 MHz, DMSO-d_6) δ 8.74 (d, J = 8.0 Hz, 1H), 8.60 (t, J = 8.8 Hz, 1H), 7.99 - 7.96 (m, 2H), 7.90 (d, J = 8.3 Hz, 1H), 7.85 (d, J = 8.3 Hz, 1H), 7.55-7.50 (m, 2H), 7.04 (dd, J = 15.6, 8.8 Hz, 1H), 6.86 (d, J = 16.0 Hz, 1H), 4.83 - 4.78 (m, 1H), 4.51 - 4.47 (m, 1H), 3.99 - 3.86 (m, 2H), 1.29 (d, J = 5.1 Hz, 3H)	3310, 1650, 1166, 558
P730		615.85 ([M+H]$^+$)	(300 MHz, DMSO-d_6) δ 8.67 (d, J = 7.8 Hz, 1H), 8.55 (t, J = 6.6 Hz, 1H), 8.31 (s, 1H), 7.90 - 7.86 (m, 3H), 7.60 (d, J = 9.2 Hz, 1H), 7.42 (d, J = 8.1 Hz, 1H), 6.98 (dd, J = 15.6, 8.7 Hz, 1H), 6.79 (d, J = 15.6 Hz, 1H), 4.96 - 4.95 (m, 1H), 4.50 - 4.45 (m, 1H), 3.96 - 3.88 (m, 2H), 1.31 (d, J = 9.3 Hz, 3H)	3299, 1651, 1166, 739
P732		605.96 ([M+H]$^+$)	(300 MHz, $CDCl_3$) δ 7.91 (s, 1H), 7.69 (s, 1H), 7.64 - 7.50 (m, 4H), 6.71 (bs,1H), 6.67 (d, J = 16.2 Hz, 1H), 6.52 (dd, J = 15.9, 7.8 Hz, 1H), 6.30 (d, J = 6.9 Hz, 1H), 4.73 - 4.69 (m, 1H), 4.30 - 4.24 (m, 1H), 3.96 - 3.91 (m, 2H), 1.49 (d, J = 7.2 Hz, 3H)	3297, 1651, 1167, 749

P802		578.1 ([M-H]$^-$)	(300 MHz, DMSO-d_6) δ 8.65 (d, J = 7.2 Hz, 1H), 8.56 (t, J = 6.3 Hz, 1H), 8.20 - 8.18 (m, 1H), 8.00 - 7.90 (m, 2H), 7.66 - 7.54 (m, 2H), 7.42 (d, J = 8.1 Hz, 1H), 6.99 (dd, J = 15.9, 9.3 Hz, 1H), 6.77 (d, J = 15.9 Hz, 1H), 4.88 - 4.82 (m, 1H), 4.51 - 4.46 (m, 1H), 4.00 - 3.88 (m, 2H), 1.29 (d, J = 7.2 Hz, 3H)	3307, 2927, 2238, 1659, 1166
P804		570.3 ([M+H]$^+$)	(300 MHz, DMSO-d_6) δ 8.74 (d, J = 7.8 Hz, 1H), 8.59 (t, J = 6.6 Hz, 1H), 8.21 - 8.19 (m, 1H), 8.01 - 7.96 (m, 2H), 7.90 (d, J = 8.1Hz, 1H), 7.66 - 7.60 (m, 1H), 7.56 (d, J = 8.1 Hz, 1H), 7.08 (dd, J = 15.9, 8.7 Hz, 1H), 6.88 (d, J = 15.6 Hz, 1H), 4.91 - 4.85 (m, 1H), 4.52 - 4.47 (m, 1H), 3.99 - 3.88 (m, 2H), 1.29 (d, J = 6.9 Hz, 3H)	3301, 3078, 2239, 1657, 1167
P1090		569.89 ([M+H]$^+$)	(300 MHz, DMSO-d_6) δ 8.64 (d, J = 7.5 Hz, 1H), 8.55 (t, J = 6.3 Hz, 1H), 7.86 (s, 1H), 7.58 (d, J = 7.2 Hz, 1H), 7.47 (d, J = 6.9 Hz, 1H), 7.41 - 7.36 (m, 3H), 7.21 - 7.15 (m, 1H), 6.91 (dd, J = 15.6, 8.7 Hz, 1H), 6.74 (d, J = 15.9 Hz, 1H), 4.66 - 4.60 (m, 1H), 4.50 -4.45 (m, 1H), 4.03 - 3.85 (m, 2H), 2.26 (s, 3H), 1.31 (d, J = 7.2 Hz, 3H)	3277, 1698, 1167, 518

P1092		559.05 ([M+H]$^+$)	(300 MHz, DMSO-d_6) δ 8.74 (d, J = 7.8 Hz, 1H), 8.59 (t, J = 6.3 Hz, 1H), 7.92 (s, 1H), 7.89 (d, J = 8.1 Hz, 1H), 7.56 (d, J = 8.1 Hz, 1H), 7.48 (d, J = 7.2 Hz, 1H), 7.42 - 7.38 (m, 1H), 7.19 (t, J = 9.3 Hz, 1H), 7.00 (dd, J = 15.9, 8.7 Hz, 1H), 6.85 (d, J = 15.9 Hz, 1H), 4.69 - 4.62 (m, 1H), 4.51 - 4.47 (m, 1H), 4.02 - 3.85 (m, 2H), 2.26 (s, 3H), 1.29 (d, J = 7.2 Hz, 3H)	3437, 1643, 1165
P1197		576.83 ([M+H]$^+$)	(300 MHz, $CDCl_3$) δ 7.62-7.56 (m, 2H), 7.40-7.37 (m, 1H), 7.05-7.00 (m, 2H), 6.72 (bs, 1H), 6.56 (bs, 1H), 6.51 (d, J = 16.2 Hz, 1H), 6.52 (dd, J = 15.9, 7.5 Hz, 1H), 4.21 (d, J = 5.4 Hz, 2H), 4.13-4.08 (m, 1H), 4.02-3.91 (m, 2H)	3311, 1655, 1166
P1269		659.07 ([M+H]$^+$)	(300 MHz, $CDCl_3$) δ 7.91 (s, 1H), 7.83 (s, 2H), 7.64-7.56 (m, 2H), 7.42-7.39 (m, 1H), 6.74 (bs, 1H), 6.62 (bs, 1H), 6.61 (d, J = 15.9 Hz, 1H), 6.50 (dd, J = 15.7, 7.8 Hz, 1H), 4.35-4.30 (m, 1H), 4.21 (d, J = 6.0 Hz, 2H), 4.02-3.90 (m, 2H)	3317, 1706, 1175, 559

P1340		607.00 ([M-H]$^-$)	(300 MHz, DMSO-d_6) δ 8.68 (t, J = 5.7 Hz, 1H), 8.56 (t, J = 6.3 Hz, 1H), 7.93 (s, 1H), 7.87-7.83 (m, 2H), 7.76-7.75 (m, 1H), 7.62-7.55 (m, 1H), 7.45 (d, J = 7.8 Hz, 1H), 7.05 (dd, J = 15.6, 9.0 Hz, 1H), 6.88 (d, J - 15.3 Hz, 1H), 4.99-4.92 (m, 1H), 4.01-3.90 (m, 4H)	3411, 1652, 1166
P1411		624.92 ([M+H]$^+$)	(300 MHz, $CDCl_3$) δ 7.64 (s, 1H), 7.58-7.55 (m, 3H), 7.51 (s, 1H), 7.41-7.38 (m, 1H), 6.77 (bs, 1H), 6.72 (bs, 1H), 6.59 (d, J = 15.9 Hz, 1H), 6.48 (dd, J = 15.9, 7.5 Hz, 1H), 4.24-4.19 (m, 4H), 4.01-3.90 (m, 1H)	3098, 1721, 1214, 723, 513
P1483		608.78 ([M+H]$^+$)	(300 MHz, $CDCl_3$) δ 7.62-7.54 (m, 3H), 7.39 (d, J = 8.1 Hz, 1H), 7.29-7.24 (m, 1H), 6.84-6.82 (m, 1H), 6.56 (d, J = 15.6 Hz, 1H), 6.49 (dd, J = 15.6, 6.6 Hz, 1H), 4.23-4.19 (m, 2H), 4.01-3.93 (m, 3H)	3300, 1657, 1164, 560
P1556		614.84 ([M+H]$^+$)	7.70 (s, 1H), 7.63 - 7.61 (m, 1H), 7.56 - 7.51 (m, 1H), 7.41 (s, 2H), 6.65 (d, J = 15.6 Hz, 1H), 6.60 (bs, 2H), 6.50 (dd, J = 15.2, 6.8 Hz, 1H), 4.19 - 4.13 (m, 3H), 3.98 - 3.94 (m, 2H)	3369, 1719, 1164, 807.

P1558		595.00 ([M-H]$^-$)	(300 MHz, $CDCl_3$) δ 8.89 (bs, 1H), 7.72-7.69 (d, *J* = 8.1 Hz, 1H), 7.45-7.35 (m, 4H), 7.25-7.19 (m, 1H), 6.58 (d, *J* = 15.9 Hz, 1H), 6,45 (dd, *J* = 15.9, 7.8 Hz, 1H), 4.57 (d, *J* = 5.7 Hz, 2H), 4.49-4.38 (m, 2H), 4.16-4.03 (m, 1H)	3351, 1660, 1161, 700
P1559		628.95 ([M-H]$^-$)	(300 MHz, DMSO-d_6) δ 10.45 (t, *J* = 6.4 Hz, 1H), 8.97 (t, *J* = 6.0 Hz, 1H), 8.02-7.92 (m, 4H), 7.77 (d, *J* = 7.5 Hz, 1H), 7.10 (dd, *J* = 15.6, 9.0 Hz, 1H), 6.90 (d, *J* = 15.6 Hz, 1H), 4.90-4.82 (m, 1H), 4.63-4.58 (m, 2H), 4.23 (d, *J* = 6.0 Hz, 2H)	3398, 1664, 1163, 808
P1560		574.88 ([M-H]$^-$)	(400 MHz, DMSO-d_6) δ 10.38 (t, *J* = 6.0 Hz, 1H), 8.65 (t, *J* = 5.6 Hz, 1H), 7.88 (s, 2H), 7.50-7.42 (m, 3H), 6.88 (dd, *J* = 15.6, 8.8 Hz, 1H), 6.75 (d, *J* = 15.6 Hz, 1H), 4.85-4.81 (m, 1H), 4.63-4.54 (m, 2H), 4.24 (d, *J* = 6.2 Hz, 2H), 2.36 (s, 3H)	3246, 1646, 1161, 808
P1564		576.6 ([M+H]$^+$)	(400 MHz, DMSO-d_6) δ 10.48 (t, *J* = 6.0 Hz, 1H), 8.74 (t, *J* = 6.2 Hz, 1H), 7.89 (s, 2H), 7.41-7.36 (m, 2H), 7.19 (d, *J* = 8.0 Hz, 1H), 6.84 (dd, *J* = 15.6, 8.8 Hz, 1H), 6.73 (d, *J* = 15.6 Hz, 1H), 4.84-4.79 (m, 1H), 4.42 (d, *J* = 6.0 Hz, 2H), 3.98-3.92 (m, 2H), 2.32 (s, 3H)	3351, 1684, 1163, 808

P1566		597.00 ($[M+H]^+$)	(400 MHz, DMSO-d_6) δ 8.74 (t, J = 5.6 Hz, 1H), 8.30 (t, J = 6.0 Hz, 1H), 8.03 - 7.84 (m, 4H), 7.61 (d, J = 8.4 Hz, 1H), 7.09 (dd, J = 15.6, 8.8 Hz, 1H), 6.89 (d, J = 16.4 Hz, 1H), 4.88 -4.81 (m, 1H), 3.89 (d, J - 5.6 Hz, 2H), 3.59 - 3.48 (m, 3H)	3323, 1654, 1115, 808
P1589		638.9 ($[M+H]^+$)	(300 MHz, DMSO-d_6) δ 8.63 (t, J = 5.7 Hz, 1H), 8.50 (d, J = 9.0 Hz, 1H), 7.93-7.91 (m, 3H), 7.62-7.60 (m, 1H), 7.42 (d, J = 7.8 Hz, 1H), 7.01 (dd, J = 15.6, 9.6 Hz, 1H), 6.77 (d, J = 15.6 Hz, 1H), 4.88-4.80 (m, 1H), 4.66-4.60 (m, 1H), 4.00-3.97 (m, 1H), 3.87-3.80 (m, 1H), 1.27 (d, J = 6.9 Hz, 3H)	3412, 1657, 1169, 749, 565
P1591		628.95 ($[M+H]^+$)	(300 MHz, DMSO-d_6) δ 8.74 (t, J = 6.0 Hz, 1H), 8.51 (d, J = 8.7 Hz, 1H), 7.99 (s, 1H), 7.92-7.90 (m, 3H), 7.58 (d, J = 8.1 Hz, 1H), 7.10 (dd, J = 15.6, 8.7 Hz, 1H), 6.89 (d, J = 15.9 Hz, 1H), 4.89-4.86 (m, 1H), 4.66-4.58 (m, 1H), 4.00-3.92 (m, 1H), 3.88-3.80 (m, 1H), 1.27 (d, J = 7.2 Hz, 3H)	3436, 1667, 1261, 749

P1592		574.98 ($[M+H]^+$)	(300 MHz, DMSO-d_6) δ 8.51 (d, J = 8.7 Hz, 1H), 8.41 (t, J = 6.0 Hz, 1H), 7.89 (s, 2H), 7.45-7.35 (m, 3H), 6.83 (dd, J = 15.6, 8.5 Hz, 1H), 6.75 (d, J = 15.6 Hz, 1H), 4.86-4.79 (m, 1H), 4.65-4.60 (m, 1H), 3.98-3.91 (m, 1H), 3.85-3.78 (m, 1H), 2.36 (s, 3H), 1.26 (d, J = 6.9 Hz, 3H)	3418, 1651, 1163, 748
P1599		689.9 ($[M+H]^+$)	(300 MHz, DMSO-d_6) δ 8.67 (bs, 1H), 8.60 (t, J = 6.0 Hz, 1H), 7.95 (d, J = 9.9 Hz, 2H), 7.91 (s, 1H), 7.78 (d, J = 6.6 Hz, 1H), 7.42 (d, J = 7.8 Hz, 1H) 7.12 (dd, J = 15.6, 9.9 Hz, 1H), 6.78 (d, J = 15.3 Hz, 1H), 4.94 - 4.91 (m, 1H), 4.52 - 4.45 (m, 1H), 4.00 - 3.85 (m, 2H), 1.33 (d, J = 6.9 Hz, 3H)	3415, 1599, 1162, 748
P1601		678.6 ($[M+H]^+$)	(300 MHz, DMSO-d_6) δ 8.73 (bs, 1H), 8.59 (t, J = 6.0 Hz, 1H), 7.95 (d, J = 9.9 Hz, 2H), 7.88 - 7.83 (m, 2H), 7.56 (d, J = 7.2 Hz, 1H), 7.12 (dd, J = 15.6, 10.5 Hz, 1H), 6.78 (d, J = 15.3 Hz, 1H), 4.94 - 4.91 (m, 1H), 4.51 - 4.43 (m, 1H), 3.99 - 3.88 (m, 2H), 1.31 (d, J = 6.9 Hz, 3H)	3423, 1646, 1141, 807

P1603	113-117	563 ($[M+H]^+$)	7.80 – 7.72 (m, 2H), 7.48 – 7.44 (m, 2H), 7.42 (s, 2H), 6.92 (t, *J* = 6.4 Hz, 1H), 6.62 (m, 1H), 6.42 (dd, *J* = 15.9, 8.0 Hz, 1H), 4.75 (p, *J* = 7.1 Hz, 1H), 4.11 (p, *J* = 8.5 Hz, 1H), 4.07 – 3.79 (m, 2H), 1.54 (d, *J* = 7.0 Hz, 3H)	**^{19}F NMR** (376 MHz, $CDCl_3$) δ -68.66, -72.52
P1611A		621.0 ($[M+H]^+$)	(300 MHz, DMSO-d_6) δ 8.60 (d, *J* = 7.8 Hz, 1H), 8.28 (t, *J* = 5.7 Hz, 1H), 7.91 (s, 3H), 7.61 - 7.58 (m, 1H), 7.43 (d, *J* = 8.1 Hz, 1H), 7.01 (dd, *J* = 15.6, 9.0 Hz, 1H), 6.77 (d, *J* = 15.9 Hz, 1H), 4.86 - 4.79 (m, 1H), 4.49 - 4.44 (m, 1H), 3.55 - 3.31 (m, 2H), 1.30 (d, *J* = 6.9 Hz, 3H)	3410, 2925, 1645, 1115, 748, 561
P1613A		611.1 ($[M+H]^+$)	(300 MHz, DMSO-d_6) δ 8.70 (d, *J* = 7.8 Hz, 1H), 8.31 (t, *J* = 5.7 Hz, 1H), 7.98 (s, 1H), 7.92 - 7.88 (m, 3H), 7.57 -7 .55 (m, 1H),), 7.09 (dd, *J* = 15.3, 9.0 Hz, 1H), 6.89 (d, *J* = 15.9 Hz, 1H), 4.89 - 4.82 (m, 1H), 4.52 - 4.43 (m, 1H), 3.57 - 3.47 (m, 2H), 1.29 (d, *J* =7.5 Hz, 3H)	3297, 1651, 1115, 808

P1616		602.8 ($[M+H]^+$)	(300 MHz, DMSO-d_6) δ 8.64 (d, J = 7.5 Hz, 1H), 8.11 (t, J = 5.4 Hz, 1H), 7.91 (s, 3H), 7.60 - 7.58 (m, 1H), 7.43 (d, J = 8.2 Hz, 1H), 7.00 (dd, J = 15.6, 9.0 Hz, 1H), 6.77 (d, J = 15.9 Hz, 1H), 4.86 - 4.53 (m, 1H), 4.51 - 4.42 (m, 2H), 4.35 (t, J = 5.1 Hz, 1H), 3.45 - 3.31 (m, 2H), 1.31 (d, J = 7.2 Hz, 3H)	3305, 1650, 1169, 749, 559
P1616A		602.8 ($[M+H]^+$)	(400 MHz, DMSO-d_6) δ 8.53 (d, J = 7.60 Hz, 1H), 8.08 (t, J = 6.0 Hz, 1H), 7.91 - 7.90 (m 3H), 7.59 (d, J = 7.6 Hz, 1H), 7.43 (d, J = 8.0 Hz, 1H), 6.98 (dd, J = 16.0, 9.2 Hz, 1H), 6.77 (d, J = 16.0 Hz, 1H), 4.84 - 4.80 (m, 1H), 4.50 - 4.36 (m, 3H), 3.44 - 3.24 (m, 2H), 1.30 (d, J = 7.2 Hz, 3H)	3407, 1651, 1169, 744, 559
P1618A		593.1 ($[M+H]^+$)	(300 MHz, DMSO-d_6) δ 8.64 (d, J = 7.8 Hz, 1H), 8.17 (t, J = 5.4 Hz, 1H), 8.03 - 7.87 (m, 3H), 7.60 (d, J = 7.8 Hz, 1H), 7.09 (dd, J = 15.9, 9.0 Hz, 1H), 6.89 (d, J = 15.9 Hz, 1H), 4.89 - 4.53 (m, 1H), 4.53 - 4.42 (m, 2H), 4.35 (t, J = 5.1 Hz, 1H), 3.44 - 3.42 (m, 2H), 1.28 (d, J = 6.9 Hz, 3H)	3411, 1651, 1115, 809

P1621		585.0 ($[M+H]^+$)	(300 MHz, DMSO-d_6) δ 8.50 (bs, 1H), 7.91 (s, 3H), 7.83 - 7.81 (m, 1H), 7.60 (d, J = 8.1 Hz, 1H), 7.43. (d, J = 8.1 Hz, 1H), 7.00 (dd, J = 15.9, 9.9 Hz, 1H), 6.77 (d, J = 15.9 Hz, 1H), 4.86 - 4.80 (m, 1H), 4.40 - 4.36 (m, 1H), 3.14 - 3.06 (m, 2H), 1.28 (d, J = 6.9 Hz, 3H), 1.03 (t, J = 6.0 Hz, 3H)	3411, 1649, 1168, 806, 559
P1623		575.1 ($[M+H]^+$)	(300 MHz, DMSO-d_6) δ 8.59 (bs, 1H), 7.97 (s, 1H), 7.92 - 7.85 (m, 4H), 7.57 (d, J = 7.8 Hz, 1H), 7.09 (dd, J = 16.2, 9.3 Hz, 1H), 6.88 (d, J = 15.9 Hz, 1H), 4.89 - 4.82 (m, 1H), 4.43 - 4.36 (m, 1H), 3.12 - 3.05 (m, 2H), 1.26 (d, J = 7.2 Hz, 3H), 1.05 (t, J = 7.5 Hz, 3H)	3412, 1645, 1115, 749
P1624		521.2 ($[M+H]^+$)	(300 MHz, DMSO-d_6) δ 8.26 (bs, 1H), 7.89 (s, 1H), 7.85-7.81 (m, 2H), 7.47 - 7.37 (m, 2H), 7.34 (d, J = 7.5 Hz, 1H), 6.88 (dd, J = 15.9, 8.7 Hz, 1H), 6.75 (d, J = 15.9 Hz, 1H), 4.85 - 4.79 (m, 1H), 4.39 - 4.34 (m, 1H), 3.13 - 3.04 (m, 2H), 2.48 (s, 3H), 1.28 (d, J = 7.5 Hz, 3H), 1.04 (t, J = 7.5 Hz, 3H)	3307, 1642, 1114, 749

P1631A		652.8 ([M+H]$^+$)	(400 MHz, DMSO-d_6) δ 8.54 (d, J = 7.60 Hz, 1H), 8.07 (t, J = 5.4 Hz, 1H), 7.91 - 7.90 (m 3H), 7.60 (d, J = 9.2 Hz, 1H), 7.43 (d, J = 8.0 Hz, 1H), 6.98 (dd, J = 16.0, 9.0 Hz, 1H), 6.77 (d, J = 16.0 Hz, 1H), 4.85 - 4.80 (m, 1H), 4.41 - 4.37 (m, 1H), 3.40 - 3.26 (m, 2H), 2.50 - 2.38 (m, 2H), 1.28 (d, J = 7.6 Hz, 3H)	3297, 1646, 1161, 742, 555
P1633A		640.7 ([M-H]$^-$)	(300 MHz, DMSO-d_6) δ 8.66 (d, J = 7.2 Hz, 1H), 8.13 (t, J = 5.4 Hz, 1H), 7.97 (s, 1H), 7.92 - 7.88 (m, 3H), 7.58 (d, J = 8.2 Hz, 1H), 7.09 (dd, J = 16.3, 9.0 Hz, 1H), 6.89 (d, J = 15.9 Hz, 1H), 4.89 - 4.82 (m, 1H), 4.42 - 4.37 (m, 1H), 3.38 - 3.24 (m, 2H), 2.44 - 2.36 (m, 2H), 1.27 (d, J = 7.2 Hz, 3H)	3299, 1651, 1139, 808
P1636	176-184	575 ([M-H]$^-$)	7.76 - 7.72 (m, 2H), 7.48 - 7.44 (m, 2H), 7.42 (s, 2H), 6.62 (d, J = 15.9 Hz, 1H), 6.54 (s, 1H), 6.42 (dd, J = 15.9, 7.9 Hz, 1H), 4.89 (s, 1H), 4.11 (p, J = 8.7 Hz, 1H), 3.92 (dqd, J = 22.9, 9.0, 6.5 Hz, 2H), 1.71 (s, 6H)	**^{19}F NMR** (376 MHz, $CDCl_3$) rotomers δ -68.65, -72.57, -73.24

P1638		638.64 ([M+H]$^+$)	(300 MHz, DMSO-d_6) δ 8.60 (bs, 1H), 8.44 (t, J = 6.0 Hz, 1H), 7.90 (s, 3H), 7.59 (d, J = 6.9 Hz,1H),7.33 (d, J = 7.8 Hz, 1H),7.00 (dd, J = 16.2, 9.6 Hz, 1H), 6.76 (d, J = 15.3 Hz, 1H), 4.88 - 4.4.80 (m, 1H), 3.92 - 3.87 (m, 2H), 3.42 - 3.40 (m, 2H), 2.50 - 2.49 (m, 2H)	3427, 2924, 1651, 1162, 680
P1640		628.9 ([M+H]$^+$)	(300 MHz, DMSO-d_6) δ 8.61 (bs, 1H), 8.53 (t, J = 6.0 Hz, 1H), 7.96 (s, 1H), 7.92 - 7.86 (m, 3H), 7.47 (d, J = 9.0 Hz, 1H), 7.18 (dd, J = 15.6, 8.1 Hz, 1H), 6.88 (d, J = 15.6 Hz, 1H), 4.91 -4.78 (m, 1H), 3.96 - 3.84 (m, 2H), 3.45 - 3.38 (m, 2H), 2.50 - 2.43 (m, 2H)	3445, 1644, 1163, 749
P1641		574.8 ([M+H]$^+$)	(300 MHz, DMSO-d_6) δ 8.61 (bs, 1H), 8.27 (t, J = 5.4 Hz, 1H), 7.89 (s, 2H), 7.47 - 7.37 (m, 2H), 7.28 (d, J = 8.1 Hz, 1H), 6.88 (dd, J = 15.6, 9.0 Hz, 1H), 6.74 (d, J = 15.9 Hz, 1H), 4.88 - 4.81 (m, 1H), 3.92- 3.86 (m, 2H), 3.45- 3.38 (m, 2H), 2.51 - 2.44 (m, 2H), 2.32 (s, 3H)	3427, 2926, 1645, 1162, 809

P1691		654.8 ([M+H]$^+$)	(300 MHz, DMSO-d_6) δ 8.64 (d, J = 7.2 Hz, 1H), 8.57 (t, J = 6.3 Hz, 1H), 7.91 (s, 1H), 7.71 (s, 2H), 7.66 (s, 1H), 7.57 (d, J = 8.1 Hz, 1H), 7.41 (d, J = 7.5 Hz, 1H), 7.03 (dd, J = 15.6, 9.6 Hz, 1H), 6.78 (d, J = 15.6 Hz, 1H), 4.91 - 4.71 (m, 1H), 4.50 - 4.45 (m, 1H), 4.02 - 3.82 (m, 2H), 1.30 (d, J = 7.2 Hz, 3H)	3301, 1647, 1164, 746
P1693		645.1 ([M+H]$^+$)	(300 MHz, DMSO-d_6) δ 8.73 (d, J = 7.2 Hz, 1H), 8.61 (d, J = 6.3 Hz, 1H), 7.96 (s, 1H), 7.86 - 7.80 (m, 1H), 7.71 (s, 2H), 7.67 (d, J = 8.1 Hz, 1H), 7.56 (d, J = 7.8 Hz, 1H), 7.00 (dd, J = 15.6, 9.3 Hz, 1H), 6.90 (d, J = 15.9 Hz, 1H), 4.92 - 4.71 (m, 1H), 4.51 - 4.49 (m, 1H), 3.99 - 3.85 (m, 2H), 1.29 (d, J = 7.2 Hz, 3H)	3309, 1650, 1165, 677
P1696		604.2 ([M+H]$^+$)	(300 MHz, DMSO-d_6) δ 8.65 (d, J = 7.2 Hz, 1H), 8.55 (t, J = 6.0 Hz, 1H), 7.92 (d , J = 1.5 Hz, 1H), 7.68 (t, J = 3.3 Hz, 3H), 7.60 (d, J = 6.9Hz, 1H), 7.42 (d, J = 7.8 Hz, 1H), 7.00 (dd, J = 15.3, 8.7 Hz, 1H), 6.76 (d, J = 15.6 Hz, 1H), 4.83 (t, J = 7.2 Hz, 1H), 4.51 (t, J = 7.2 Hz, 1H), 4.01 - 3.88 (m, 2H), 1.29 (d, J = 7.2 Hz, 3H)	3418, 1645, 1165, 750, 562

P1698		595.1 ([M+H]$^+$)	(300 MHz, DMSO-d_6) δ 8.74 (d J = 7.5 Hz, 1H), 8.59 (t, J = 6.6 Hz, 1H), 7.98 (s , 1H), 7.91 (d, J = 8.4 Hz, 1H), 7.69 - 7.62 (m, 2H), 7.56 (d, J = 7.5 Hz, 1H), 7.09 (dd, J = 15.3, 8.7 Hz , 1H), 6.88 (d, J = 15.6 Hz, 1H), 4.83 - 4.80 (m, 1H), 4.51 - 4.47 (m, 1H), 3.99 - 3.86 (m, 2H), 1.29 (d, J = 7.2 Hz, 3H)	3294, 2928, 1646, 1166, 750
P1776		593.2 ([M+H]$^+$)	(400 MHz, DMSO-d_6) δ 8.69 (t, J = 6.0 Hz, 1H), 8.57 (t, J = 6.4 Hz, 1H), 7.90 (dd, J = 13.3, 1.8 Hz, 2H), 7.72 (d, J = 8.4 Hz, 1H), 7.58 (ddd, J = 18.0, 8.2, 1.8 Hz, 2H), 7.44 (d, J = 7.9 Hz, 1H), 6.94 (dd, J = 15.7, 9.0 Hz, 1H), 6.74 (d, J = 15.7 Hz, 1H), 4.81 (p, J = 9.4 Hz, 1H), 4.02 - 3.87 (m, 4H)	**^{19}F NMR** (376 MHz, DMSO-d_6) δ -67.98, -70.71
P1781		513.2 ([M+H]$^+$)	7.85 - 7.74 (m, 2H), 7.53 - 7.41 (m, 4H), 7.30 (t, J = 6.2 Hz, 1H), 7.23 (dd, J = 8.4, 2.0 Hz, 1H), 7.19 (t, J = 5.2 Hz, 1H), 6.61 (d, J = 15.9 Hz, 1H), 6.46 (dd, J = 15.9, 7.8 Hz, 1H), 4.26 (d, J = 5.1 Hz, 2H), 4.21 - 4.07 (m, 1H), 3.95 (qd, J = 9.1, 6.4 Hz, 2H)	**^{19}F NMR** (376 MHz, $CDCl_3$) δ -68.78, -72.44

P1806		605.0 ([M+H]$^+$)	(300 MHz, DMSO-d_6) δ 8.67 (d, J = 7.2 Hz, 1H), 8.57 (t, J = 6.3 Hz, 1H), 7.91 - 7.88 (m, 2H), 7.74 (d, J = 8.7 Hz, 1H), 7.64 – 7.59 (m, 2H), 7.41 (d, J = 8.1 Hz, 1H), 6.98 (dd, J = 15.9, 9.3 Hz, 1H), 6.76 (d, J = 15.9 Hz, 1H), 4.83 - 4.77 (m, 1H), 4.52 - 4.43 (m, 1H), 4.05 - 3.83 (m, 2H), 1.30 (d, J = 7.2 Hz, 3H)	3411, 2925, 1650, 1163, 747
P1808		695.1 ([M+H]$^+$)	(400 MHz, DMSO-d_6) δ 8.74 (d, J = 7.2 Hz, 1H), 8.59 (s, 1H), 7.96 - 7.89 (m, 3H), 7.73 (d, J = 8.4 Hz, 1H), 7.55 (t, J = 8.8 Hz, 2H), 7.05 (dd, J = 16.0, 8.8 Hz, 1H), 6.87 (d, J = 16.0 Hz, 1H), 4.85 - 4.80 (m, 1H), 4.51 - 4.47 (m, 1H), 3.99 - 3.88 (m, 2H), 1.28 (d, J = 6.8 Hz, 3H)	3297, 1652, 1165, 745
P1862		598.89 ([M+H]$^+$)	7.70 (s, 1H), 7.62-7.60 (m, 1H), 7.55-7.53 (m, 1H), 7.35 (d, J = 5.6 Hz, 2H), 7.89 (t, J = 6.0 Hz, 1H), 6.67 (t, J = 5.6 Hz, 1H), 6.64 (d, J = 16.0 Hz, 1H), 6.48 (dd, J = 16.4, 8.0 Hz, 1H), 4.23 (d, J = 5.6 Hz, 2H), 4.17-4.08 (m, 1H), 3.97-3.89 (m, 2H)	3324, 1673, 1164, 772

P1864		622.88 ([M-H]$^-$)	(400 MHz, DMSO-d_6) δ 10.39 (bs, 1H), 8.85 (t, J = 6.0 Hz, 1H), 7.94 (s, 1H), 7.87 (d, J = 6.4 Hz, 2H), 7.64 (d, J = 8.4 Hz, 1H), 7.58 (d, J = 8.0 Hz, 1H), 7.00 (dd, J = 15.6, 8.8 Hz, 1H), 6.77 (d, J = 15.6 Hz, 1H), 4.81-4.79 (m, 1H), 4.63-4.59 (m, 2H), 4.24 (d, J = 6.0 Hz, 2H)	3256, 1657, 1166, 749, 591
P1866		614.91 ([M+H]$^+$)	(400 MHz, DMSO-d_6) δ 10.43 (bs, 1H), 8.96 (t, J = 6.0 Hz, 1H), 8.00 (s, 1H), 7.94 (d, J = 7.6 Hz, 1H), 7.88 (d, J = 6.4 Hz, 2H), 7.76 (d, J = 8.0 Hz, 1H), 7.09 (dd, J = 15.6, 9.2 Hz, 1H), 6.88 (d, J = 16.0 Hz, 1H), 4.86-4.78 (m, 1H), 4.62-4.58 (m, 2H), 4.23 (d, J = 5.6 Hz, 2H)	3447, 1655, 1167, 816
P1969		678.77 ([M+H]$^+$)	(300 MHz, DMSO-d_6) δ 8.65 (t, J = 5.4 Hz, 1H), 8.54 (t, J = 6.3 Hz, 1H), 7.90-7.86 (m, 2H), 7.82 (s, 2H), 7.60 (d, J = 7.2 Hz, 1H), 7.43 (d, J = 7.8 Hz, 1H), 6.99 (dd, J = 15.6, 9.0 Hz, 1H), 6.75 (d, J = 15.6 Hz, 1H), 4.83-4.76 (m, 1H), 4.01-3.91 (m, 4H)	3296, 1697, 1164, 596
P1970		634.87 ([M+H]$^+$)	(300 MHz, DMSO-d_6) δ 8.68 (t, J = 6.0 Hz, 1H), 8.57 (t, J = 6.0 Hz, 1H), 7.89 - 7.78 (m, 4H), 7.59 - 7.56 (m, 1H), 7.49 - 7.46 (m, 1H), 7.02 (dd, J = 15.9, 8.1 Hz, 1H), 6,78 (d, J = 15.6 Hz, 1H), 4.83 - 4.76 (m, 1H), 4.01 - 3.91 (m, 4H)	3299, 1658, 1164, 745, 543

P1971		668.87 ($[M+H]^+$)	(300 MHz, DMSO-d_6) δ 8.77 (t, J = 6.0 Hz, 1H), 8.59 (t, J = 6.3 Hz, 1H), 7.99 (s, 1H), 7.93-7.85 (m, 4H), 7.60 (d, J = 8.1 Hz, 1H), 7.10 (dd, J = 15.9, 9.3 Hz, 1H), 6.89 (d, J = 15.9 Hz, 1H), 4.85-4.79 (m, 1H), 3.98-3.90 (m, 4H)	3351, 1705, 1165, 551
P1972		614.84 ($[M+H]^+$)	(400 MHz, DMSO-d_6) δ 8.57 (t, J = 6.4 Hz, 1H), 8.47 (t, J = 6.0 Hz, 1H), 7.88-7.80 (m, 3H), 7.45 (s, 1H), 7.43-7.37 (m, 2H), 6.86 (dd, J = 15.6, 8.8 Hz, 1H), 6.74 (d, J = 15.6 Hz, 1H), 4.83-4.74 (m, 1H), 3.98-3.88 (m, 4H), 2.37 (s, 3H)	3311, 1650, 1163, 543
P2009		692.88 ($[M+H]^+$)	(300 MHz, DMSO-d_6) δ 8.62 (t, J = 5.7 Hz, 1H), 8.50 (d, J = 8.7 Hz, 1H), 7.93 - 7.92 (m, 1H), 7.89 - 7.88 (m, 1H), 7.84 (s, 2H), 7.62-7.59 (m, 1H), 7.42 (d, J = 8.1 Hz, 1H), 7.01 (dd, J = 15.9, 9.2 Hz, 1H), 6.77 (d, J = 15.6 Hz, 1H), 4.82 - 4.76 (m, 1H), 4.65 -4.58 (m, 1H), 4.00 - 3.92 (m, 1H), 3.87 - 3.80 (m, 1H), 1.27 (d, J = 6.9 Hz, 3H)	3402, 1659, 1165, 560

P2010		646.84 ([M-H]$^-$)	(300 MHz, DMSO-d_6) δ 8.63 (t, J = 6.0 Hz, 1H), 8.50 (d, J = 8.7 Hz, 1H), 7.89 - 7.88 (m, 1H), 7.84 (s, 2H), 7.78 (s, 1H), 7.59 - 7.56 (m, 1H), 7.46 (d, J = 7.5 Hz, 1H), 7.00 (dd, J = 15.6, 9.0 Hz, 1H), 6.78 (d, J = 15.3 Hz, 1H), 4.79 - 4.76 (m, 1H), 4.63 - 4.60 (m, 1H), 3.95 - 3.86 (m, 2H), 1.27 (d, J = 7.2 Hz, 3H)	3406, 1658, 1165, 746, 583
P2011		682.93 ([M+H]$^+$)	(400 MHz, DMSO-d_6) δ 8.73 (t, J = 5.6 Hz, 1H), 8.50 (d, J = 8.8 Hz, 1H), 7.98 (s, 1H), 7.90 - 7.89 (m, 2H), 7.85 (s, 2H), 7.57 (d, J = 8.4 Hz, 1H), 7.09 (dd, J = 15.6, 9.2 Hz, 1H), 6.88 (d, J = 15.6 Hz, 1H), 4.84 - 4.79 (m, 1H), 4.65 - 4.59 (m, 1H), 4.03 - 3.95 (m, 1H), 3.87 - 3.81 (m, 1H), 1.25 (d, J = 6.0 Hz, 3H)	3417, 1666, 1115, 558
P2012		628.89 ([M+H]$^+$)	(300 MHz, DMSO-d_6) δ 8.51 (d, J = 8.7 Hz, 1H), 8.41 (t, J = 6.0 Hz, 1H), 7.88 (s, 1H), 7.82 (s, 1H), 7.45-7.34 (m, 3H), 6.87 (dd, J = 15.9, 8.7 Hz, 1H), 6.74 (d, J = 15.6 Hz, 1H), 4.82-4.75 (m, 1H), 4.65-4.57 (m, 1H), 3.98-3.78 (m, 2H), 2.36 (s, 3H), 1.26 (d, J = 7.2 Hz, 3H)	3406, 1643, 1163, 583

P2036		674.7 ([M+H]$^+$)	(300 MHz, DMSO-d_6) δ 8.61 (d, J = 7.8 Hz, 1H), 8.26 (t, J = 5.7 Hz, 1H), 7.92 - 7.88 (m, 2H), 7.84 (s, 2H), 7.61 (d, J = 7.8 Hz, 1H), 7.42 (d, J = 7.8 Hz, 1H), 7.00 (dd, J = 15.9, 9.3 Hz, 1H), 6.77 (d, J = 15.9 Hz, 1H), 6.20 - 5.80 (m, 1H), 4.83 - 4.74 (m, 1H), 4.51 - 4.42 (m, 1H), 3.61 - 3.31 (m, 2H), 1.31 (d, J = 7.5 Hz, 3H)	3296, 1651, 1113, 534
P2038		662.7 ([M-H]$^+$)	(300 MHz, DMSO-d_6) δ 8.69 (d, J = 7.2 Hz, 1H), 8.30 (bs, 1H), 7.98 (s, 1H), 7.89 - 7.85 (m, 4H), 7.57 (d, J = 8.1 Hz, 1H), 7.42 (d, J = 7.8 Hz, 1H), 7.09 (dd, J = 15.9, 9.6 Hz, 1H), 6.88 (d, J = 15.9 Hz, 1H), 6.18 - 5.81 (m, 1H), 4.84 - 4.78 (m, 1H), 4.50 - 4.45 (m, 1H), 3.50 - 3.45 (m, 2H), 1.29 (d, J = 7.2 Hz, 3H)	3292, 1650, 1115, 747, 681
P2041		656.6 ([M+H]$^+$)	(400 MHz, DMSO-d_6) δ 8.56 (d, J = 7.2 Hz, 1H), 8.10 (t, J = 5.2 Hz, 1H), 7.92 - 7.88 (m, 2H), 7.84 (s, 2H), 7.60 (d, J = 8.0 Hz, 1H), 7.42 (d, J = 8.0 Hz, 1H), 6.99 (dd, J = 16.0, 9.6 Hz, 1H), 6.76 (d, J = 16.0 Hz, 1H), 4.81 - 4.74 (m, 1H), 4.50 - 4.36 (m, 3H), 3.56 - 3.36 (m, 2H), 1.29 (d, J = 6.8 Hz, 3H)	3305, 1645, 1167, 559

P2043		646.9 ($[M+H]^+$)	(300 MHz, DMSO-d_6) δ 8.64 (d, J = 7.8 Hz, 1H), 8.15 (t, J = 5.7 Hz, 1H), 7.97 (s, 1H) 7.90 - 7.85 (m, 4H), 7.57 (d, J = 8.0 Hz, 1H), 7.09 (dd, J = 15.9, 9.3 Hz, 1H), 6.88 (d, J = 15.6 Hz, 1H), 4.84 - 4.78 (m, 1H), 4.51- 4.42 (m, 2H), 4.37 - 4.33 (m, 1H), 3.45 - 3.42 (m, 2H), 1.28 (d, J = 7.8 Hz, 3H)	3291, 1651, 1115, 588
P2056	91-94	706.8 ($[M+H]^+$)	(400 MHz, DMSO-d_6) δ 8.57 (d, J = 7.2 Hz, 1H), 8.09 (t, J = 5.2 Hz, 1H), 7.92 - 7.88 (m, 2H), 7.84 (s ,2H), 7.60 (d, J = 7.6 Hz, 1H), 7.43 (d, J = 7.6 Hz, 1H), 6.99 (dd, J = 15.6, 9.1 Hz, 1H), 6.76 (d, J = 15.6 Hz, 1H), 4.81 - 4.77 (m, 1H), 4.40 - 4.36 (m, 1H), 3.40 - 3.25 (m, 2H) 2.44 - 2.32 (m, 2H), 1.28 (d, J = 6.8 Hz, 3H)	
P2058		696.8 ($[M+H]^+$)	(300 MHz, DMSO-d_6) δ 8.66 (s, 1H), 8.14 (s, 2H), 7.97 (s, 1H), 7.89 - 7.85 (m, 3H) 7.57 (d, J = 8.1 1H), 7.06 (dd, J = 16.2, 9.0 Hz, 1H), 6.88 (d, J = 16.2 Hz, 1H), 4.82 - 4.81 (m, 1H), 4.41 - 4.37 (m, 1H) 3.38 - 2.99 (m, 2H), 2.44 - 2.36 (m, 2H) 1.27 (d, J = 7.2 Hz, 3H)	3309, 1650, 1141

[a] ^{1}H NMR spectral data were acquired using a 400 MHz instrument in $CDCl_3$ except where noted. HRMS data are noted observed value (theoretical value).

Table 2C: Analytical Data for Compounds in Table 1C.

Compound Number	mp (°C)	ESIMS	^{1}H NMR (δ)[a]	IR (cm^{-1}); ^{19}F NMR (δ)
FA1		562.89 ([M+H]$^+$)	(400 MHz, DMSO-d$_6$) δ 8.65 (t, *J* = 5.2 Hz, 1H), 8.28 (t, *J* = 6.0 Hz, 1H), 7.91 (s, 2H), 7.78 (s, 1H), 7.58 (d, *J* = 8.0 Hz, 1H), 7.50 (d, *J* = 8.0 Hz, 1H), 7.01 (dd, *J* = 15.6, 9.2 Hz, 1H), 6.78 (d, *J* = 15.6 Hz, 1H), 4.86 - 4.79 (m, 1H), 3.90 (d, *J* = 6.0 Hz, 2H), 3.59 - 3.48 (m, 3H)	3418, 1658, 1114, 749
FA2		542.93 ([M+H]$^+$)	(400 MHz, DMSO-d$_6$) δ 8.45 (t, *J* = 6.0 Hz, 1H), 8.29 (t, *J* = 4.5 Hz, 1H), 7.89 (s, 2H), 7.45 - 7.34 (m, 3H), 6.87 (dd, *J* = 15.7, 6.9 Hz, 1H), 6.75 (d, *J* = 15.8 Hz, 1H), 4.85 - 4.80 (m, 1H), 3.87 (d, *J* = 4.8 Hz, 2H), 3.57 - 3.47 (m, 3H), 2.42 (s, 3H)	3430, 1646, 1113, 749
FA3		606.78 ([M+H]$^+$)	(400 MHz, DMSO-d$_6$) δ 8.65 (t, *J* = 5.6 Hz, 1H), 8.27 (t, *J* = 5.6 Hz, 1H), 7.93 - 7.91 (m, 2H), 7.62 - 7.58 (m, 2H), 7.46 (d, *J* = 8.0 Hz, 1H), 7.00 (dd, *J* = 15.2, 9.2 Hz, 1H), 6.77 (d, *J* = 15.6 Hz, 1H), 4.85 - 4.81 (m, 1H), 3.89 (d, *J* = 6.0 Hz, 1H), 3.59 - 3.48 (m, 2H), 3.17 - 3.08 (m, 2H)	3420, 1654, 114, 749, 558

FA4		599.00 ($[M+H]^+$)	(400 MHz, $CDCl_3$) δ 9.81 (s, 1H), 8.06 (d, *J* = 7.1 Hz, 1H), 7.70 (d, *J* = 5.5 Hz, 1H), 7.44 (s, 1H), 7.37 (t, *J* = 2.7 Hz, 2H), 7.20 (s, 1H), 6.82 (d, *J* = 15.9 Hz, 1H), 6.50 (dd, *J* = 16.0, 8.5 Hz, 1H), 4.14 (m, 3H), 3.89 (m, 2H)	
FA5		613.10 ($[M+H]^+$)	(400 MHz, $CDCl_3$) δ 7.45 (s, 1H), 7.41 (s, 2H), 7.29 (s, 1H), 7.28 (s, 1H), 6.84 (m, 2H), 6.43 (dd, *J* = 16.0, 8.3 Hz, 1H), 4.25 (d, *J* = 5.4 Hz, 2H), 4.10 (m, 1H), 3.97 (qd, *J* = 9.0, 6.4 Hz, 2H), 3.87 (s, 3H)	
FA6		545.92 ($[M+H]^+$)	(400 MHz, DMSO-d_6) δ 8.56 (t, *J* = 6.4 Hz, 1H), 8.51 (t, *J* = 5.6 Hz, 1H), 7.82 (d, *J* = 6.4 Hz, 2H), 7.56 (d, *J* = 8.0 Hz, 1H), 6.81 (d, *J* = 8.4 Hz, 1H), 6.75 (s, 1H), 6.72 (dd, *J* = 15.6, 6.8 Hz, 1H), 6.62 (d, *J* = 15.6 Hz, 1H), 6.51 (bs, 2H), 4.82 - 4.78 (m, 1H), 3.96 - 3.85 (m, 4H)	3369, 1694, 1164, 713
FA7		638.78 ($[M+H]^+$)	(300 MHz, $CDCl_3$) δ 7.62-7.56 (m, 2H), 7.46-7.37 (m, 2H), 6.77-6.76 (m, 1H), 6.56 (d, *J* = 14.1 Hz, 1H), 6.43 (dd, *J* = 15.9, 7.8 Hz, 1H), 4.14-4.07 (m, 3H), 3.61-3.54 (m, 2H), 2.42-2.34 (m, 2H)	3284, 1667, 1166, 744, 591

FA8		588.87 ([M+H]$^+$)	(300 MHz,DMSO-d_6) δ 8.68 (t, *J* = 5.7 Hz, 1H), 8.56 (t, *J* = 6.6 Hz, 1H), 7.91 (s, 1H), 7.61-7.58 (m, 1H), 7.45-7.39 (m, 3H), 6.95 (dd, *J* = 15.9, 9.3 Hz, 1H), 6.75 (d, *J* = 15.9 Hz, 1H), 4.76-4.64 (m, 1H), 4.06-3.92 (m, 4H), 3.90 (s, 3H)	3297, 1658, 1165
FA9		624.92 ([M+H]$^+$)	(300 MHz, DMSO-d_6) δ 9.81 (bs, 1H), 8.69 (s, 1H), 7.90 (s, 2H), 7.78 (s, 1H),7.59 - 7.58 (m, 2H), 7.02 (dd, *J* = 15.8, 9.3 Hz, 1H), 6.78 (d, *J* = 15.8 Hz, 1H), 4.86 - 4.79 (m, 1H), 4.57 - 4.52 (m, 2H), 1.60 (s, 6H)	3430, 1651, 1166, 750
FA10		640.9 ([M+H]$^+$)	(300 MHz, DMSO-d_6) δ 10.41 (s, 1H), 9.68 (bs, 1H), 7.90 (s, 2H), 7.76 - 7.67 (m, 1H), 7.53 (d, *J* = 8.1 Hz, 1H), 7.42 (d, *J* = 7.8 Hz, 1H), 6.97 (dd, *J* = 15.6, 8.7 Hz, 1H), 6.77 (d, *J* = 15.6, 1H), 4.86 (t, *J* = 9.3 Hz, 1H), 4.55 - 4.47 (m, 2H), 1.75 (s, 6H)	3434, 1260, 750
FA11		668.8 ([M+H]$^+$)	(300 MHz, DMSO-d_6) δ 9.80 (bs, 1H), 8.69 (s, 1H), 7.92 - 7.90 (m, 3H), 7.60 - 7.55 (m, 2H), 7.01 (dd, *J* = 15.5, 9.1 Hz, 1H), 6.77 (d, *J* = 15.6 Hz, 1H), 4.84 - 4.30 (m, 1H), 4.57 - 4.52 (m, 2H), 1.60 (s, 6H)	3425, 1651, 1261, 750

FA12		660.8 ([M+2H]$^+$)	(300 MHz, DMSO-d_6) δ 9.87 (bs, 1H), 8.76 (bs, 1H), 7.98 (s, 1H), 7.95 - 7.91 (m, 3H), 7.83 (d, *J* = 8.1 Hz, 1H), 7.09 (dd, *J* = 15.9, 9.3 Hz, 1H), 6.89 (d, *J* = 15.9, 1H), 4.89 (t, *J* = 9.6 Hz, 1H), 4.57 - 4.51 (m, 2H), 1.58 (s, 6H)	3430, 1651, 1261, 750
FA13		606.8 ([M+H]$^+$)	(300 MHz, DMSO-d_6) δ 10.36 (s, 1H), 8.67 (bs, 1H), 8.56 (bs, 1H), 7.92 (s, 1H), 7.61 - 7.58 (m, 3H), 7.44 (d, *J* = 7.8 Hz, 1H), 6.96 (dd, *J* = 15.4, 8.8 Hz, 1H), 6.74 (d, *J* = 15.7 Hz, 1H), 4.61 - 4.70 (m, 1H), 4.04 - 3.90 (m, 4H)	3418, 1651, 1164, 749
FA14		611.9 ([M+H]$^+$)	(400 MHz, $CDCl_3$) δ 7.69 (s, 1H), 7.62 (d, *J* = 1.6 Hz, 1H), 7.53 (d, *J* = 8.0 Hz, 1H), 7.39 (m, 3H), 6.54 (d, *J* = 15.9 Hz, 1H), 6.40 (dd, *J* = 15.9, 7.8 Hz, 1H), 6.28 (s, 1H), 4.21 (d, J = 5.7 Hz, 2H), 4.12 (m, 1H), 1.69 (s, 6H)	**^{19}F NMR** (376 MHz, $CDCl_3$) δ -68.59
FA15		643.19 ([M+H]$^+$)	(400 MHz, CDCl3) δ 7.60 (d, *J* = 1.6 Hz, 1H), 7.51 (d, *J* = 8.0 Hz, 1H), 7.40 (s, 2H), 7.36 (dd, *J* = 8.1, 1.6 Hz, 1H), 6.66 (d, *J* = 10.0 Hz, 2H), 6.53 (d, *J* = 15.9 Hz, 1H), 6.38 (dd, *J* = 15.9, 7.8 Hz, 1H), 4.11 (m, 1H), 3.12 (d, *J* = 6.2 Hz, 2H), 1.72 (s, 6H), 0.93 (s, 9H)	**^{19}F NMR** (376 MHz, $CDCl_3$) δ -68.62

FA16		687.0 ([M+H]$^+$)	(300 MHz, DMSO-d_6) δ 8.61 - 8.53 (m, 2H), 8.11 (s, 1H), 7.91 (s, 2H), 7.62 (d, *J* = 8.4 Hz, 1H), 7.36 (d, *J* = 8.1 Hz, 1H), 6.98 (dd, *J* = 15.9, 9.3 Hz, 1H), 6.74 (d, *J* = 15.9 Hz, 1H), 4.85 - 4.82 (m, 1H), 4.50 - 4.45 (m, 1H), 4.00 - 3.88 (m, 2H), 1.36 (d, *J* = 7.5 Hz, 3H)	3433, 1640, 1163, 749
FA17		587.1 ([M+H]$^+$)	(300 MHz, DMSO-d_6) δ 8.60 - 8.53 (m, 2H), 7.91 - 7.85 (m, 3H), 7.54 (d, *J* = 8.1 Hz, 1H), 7.40 (d, *J* = 8.1 Hz, 1H), 7.02 (dd, *J* = 17.4, 11.1 Hz, 1H), 6.92 (dd, *J* = 15.9, 9.0 Hz, 1H), 6.81 (d, *J* = 15.9 Hz, 1H), 5.95 (d, *J* = 17.1 Hz, 1H), 5.33 (d, *J* = 11.7 Hz, 1H), 4.87 - 4.80 (m, 1H), 4.47 - 4.43 (m, 1H), 4.01 - 3.87 (m, 2H), 1.31 (d, *J* = 6.9 Hz, 3H)	3419, 1644, 1162, 748
FA18		573.2 ([M+H]$^+$)	(300 MHz, DMSO-d_6) δ 8.62 - 8.56 (m, 2H), 7.91 - 7.86 (m, 3H), 7.56 (d, *J* = 9.3 Hz, 1H), 7.42 (d, *J* = 8.1 Hz, 1H), 7.11 (dd, *J* = 17.4, 11.1 Hz, 1H), 6.98 (dd, *J* = 15.9, 9.3 Hz, 1H), 6.82 (d, *J* = 15.9 Hz, 1H), 5.96 (d, *J* = 16.8 Hz, 1H), 5.34 (d, *J* = 12.0 Hz, 1H), 4.87 - 4.81 (m, 1H), 4.01 - 3.89 (m, 4H)	3422, 1649, 1164, 809

FA19		667.0 ($[M+H]^+$)	(300 MHz, DMSO-d_6) δ 8.64 - 8.58 (m, 2H), 7.92 - 7.89 (m, 3H), 7.60 (d, *J* = 8.1 Hz, 1H), 7.37 (d, *J* = 7.8 Hz, 1H), 7.00 (dd, *J* = 15.3, 8.7 Hz, 1H), 6.77 (d, *J* = 15.6 Hz, 1H), 4.86 - 4.79 (m, 1H), 4.49 - 4.42 (m, 1H), 4.07 - 3.81 (m, 2H), 1.65 - 1.59 (m, 2H), 1.40 - 1.30 (m, 2H), 0.89 (t, *J* = 6.9 Hz, 3H)	3288, 1645, 1164, 743, 563
FA20		609.1 ($[M+H]^+$)	(400 MHz, DMSO-d_6) δ 8.74 (d *J* = 7.6 Hz, 1H), 8.59 (t, *J* = 6.4 Hz, 1H), 7.95 (s , 1H), 7.89 (d, *J* = 7.2 Hz, 1H), 7.72 (s, 1H), 7.55 (d, *J* = 7.2 Hz, 2H), 7.02 (dd, *J* = 16.0, 8.8 Hz , 1H), 6.87 (d, *J* = 15.6 Hz, 1H), 4.76 - 4.72 (m, 1H), 4.51 - 4.47 (m, 1H), 3.99 - 3.88 (m, 2H), 2.42 (s, 3H) , 1.29 (d, *J* = 7.2 Hz, 3H)	3422, 2926, 1651, 1165, 783
FA21		673.0 ($[M+H]^+$)	(300 MHz, DMSO-d_6) δ 8.74 (d, *J* = 7.5 Hz, 1H), 8.59 (t, *J* = 6.3 Hz, 1H), 7.98 (s, 1H), 7.91 - 7.85 (m, 3H), 7.56 (d, *J* = 8.1 Hz, 1H), 7.09 (dd, *J* = 15.9, 9.3 Hz, 1H), 6.88 (d, *J* = 15.6 Hz, 1H), 4.88 - 4.81 (m, 1H), 4.52 - 4.57 (m, 1H), 4.04 - 3.85 (m, 2H), 1.29 (t, *J* = 6.9 Hz, 3H)	3429, 1650, 1165, 804

FA22		681.0 ($[M-H]^-$)	(300 MHz, DMSO-d_6) δ 8.65 (d, *J* = 7.5 Hz, 1H), 8.55 (t, *J* = 5.7 Hz, 1H), 7.91 - 7.85 (m, 2H), 7.48 - 7.39 (m, 2H), 7.31 - 7.26 (m, 1H), 7.00 (dd, *J* = 15.9, 9.3 Hz, 1H), 6.77 (d, *J* = 15.6 Hz, 1H), 4.86 - 4.82 (m, 1H), 4.50 - 4.46 (m, 1H), 4.00 - 3.88 (m, 2H), 1.31 (t, *J* = 7.2 Hz, 3H)	3298, 1651, 1164, 689, 536
FA23		651.2 ($[M+H]^+$)	(400 MHz, $CDCl_3$) δ 7.61 (d, *J* = 1.6 Hz, 1H), 7.51 (d, *J* = 8.0 Hz, 1H), 7.40 (s, 2H), 7.37 (dd, *J* = 8.1, 1.6 Hz, 1H), 6.85 (t, *J* = 6.2 Hz, 1H), 6.53 (d, *J* = 15.9 Hz, 1H), 6.42 (s, 1H), 6.38 (dd, *J* = 15.9, 7.8 Hz, 1H), 4.11 (m, 1H), 3.70 (td, *J* = 13.3, 6.3 Hz, 2H), 1.67 (m, 9H)	**^{19}F NMR** (376 MHz, $CDCl_3$) δ -68.61 , -96.22
FA24		672.9 ($[M+H]^+$)	(400 MHz, DMSO-d_6) δ 8.49 (t, *J* = 6.0 Hz, 1H), 8.42 (t, *J* = 6.4 Hz, 1H), 7.98 (s, 1H), 7.76 (s, 2H), 7.49 - 7.47 (m, 1H), 7.24 (d, *J* = 8.0 Hz, 1H), 6.82 (dd, *J* = 15.6, 9.2 Hz, 1H), 6.59 (d, *J* = 16.4 Hz, 1H), 4.70 - 4.65 (m, 1H), 3.89 - 3.75 (m, 4H)	3410, 1653, 1164, 812

FA25		603.0 ($[M+H]^+$)	(400 MHz, DMSO-d_6) δ 8.61 (t, *J* = 6.4 Hz, 1H), 8.37 (d, *J* = 8.0 Hz, 1H), 7.89 (s, 2H), 7.45 (s, 1H), 7.40 (d, *J* = 8.4 Hz, 1H), 7.33(d, *J* = 8.0 Hz, 1H), 6.86 (dd, *J* = 16.0, 8.8 Hz, 1H), 6.75 (d, *J* = 15.6 Hz, 1H), 4.85 - 4.80 (m, 1H), 4.47 - 4.41(m, 1H), 4.02 - 3.96 (m, 1H), 3.90 - 3.83 (m, 1H), 2.32 (s, 3H), 1.67 - 1.61 (m, 2H), 1.42 - 1.28 (m, 2H), 0.89 (t, *J* = 7.6 Hz, 3H)	3288, 1676, 1164, 809
FA26	108-111	726.9 ($[M+H]^+$)	(300 MHz, DMSO-d_6) δ 8.65 (d, *J* = 7.2 Hz, 1H), 8.55 (t, *J* = 6.3 Hz, 1H), 8.04 (s, 1H), 7.92 (s, 1H), 7.84 - 7.76 (m, 1H), 7.61 (d, *J* = 7.8 Hz, 1H), 7.42 (d, *J* = 7.8 Hz, 1H), 7.01 (dd, *J* = 15.6, 9.0 Hz, 1H), 6.77 (d, *J* = 15.9 Hz, 1H), 4.84 - 4.78 (m, 1H), 4.51 - 4.46 (m, 1H), 4.00 - 3.88 (m, 2H), 1.31 (d, *J* = 7.5 Hz, 3H)	
FA27	103-107	717.1 ($[M+H]^+$)	(300 MHz, DMSO-d_6) δ 8.74 (d, *J* = 6.9 Hz, 1H), 8.59 (t, *J* = 6.0 Hz, 1H), 8.06 (s, 1H), 7.91-7.79 (m, 3H), 7.56 (d, *J* = 8.1 Hz, 1H), 7.09 (dd, *J* = 15.9, 8.6 Hz, 1H), 6.88 (d, *J* = 15.3 Hz, 1H), 4.84 - 4 .81 (m, 1H), 4.52 - 4.47 (m, 1H), 3.99 - 3.88 (m, 2H), 1.29 (d, *J* = 6.9 Hz, 3H)	

FA28		588.9 ($[M+H]^+$)	(300 MHz, DMSO-d_6) δ 8.59 (t, *J* = 6.6 Hz, 1H), 8.51 (d, *J* = 6.9 Hz, 1H), 7.88 (s, 2H), 7.22 (s, 2H), 6.75 - 6.63 (m, 2H), 4.83 - 4.77 (m, 1H), 4.50 - 4.49 (m, 1H), 4.00 - 3.87 (m, 2H), 2.21 (s, 6H), 1.33 - 1.22 (m, 3H)	3361, 2960, 1690, 1166, 819
FA29		600.8 ($[M+H]^+$)	(400 MHz, DMSO-d_6) δ 8.63 (d, *J* = 7.2 Hz, 1H), 8.55 (t, *J* = 6.4 Hz, 1H), 7.86 (s, 2H), 7.77 (d, *J* = 2.0 Hz, 1H), 7.66 (d, *J* = 8.4 Hz, 1H), 7.54 (d, *J* = 7.2 Hz, 1H), 7.47 (dd, *J* = 10.0, 8.0 Hz, 1H), 7.39 (d, *J* = 7.6 Hz, 1H), 6.94 (dd, *J* = 15.6, 9.3 Hz, 1H), 6.63 (d, *J* = 15.6 Hz, 1H), 4.48 - 4.43 (m, 1H), 4.05 - 3.92 (m, 2H), 1.64 (t, *J* = 19.2 Hz, 3H), 1.30 (d, *J* = 6.8 Hz, 3H)	3288, 1687, 1164, 748, 568
FA30		655.0 ($[M+H]^+$)	(300 MHz, DMSO-d_6) δ 8.64 (d, *J* = 7.5 Hz, 1H), 8.58 (t, *J* = 6.3 Hz, 1H), 7.95 - 7.91 (m, 2H), 7.66 (s, 2H), 7.60 (d, *J* = 8.1 Hz, 1H), 7.42 (d, *J* = 7.8 Hz, 1H), 6.99 (dd, *J* = 15.9, 9.3 Hz, 1H), 6.78 (d, *J* = 15.9 Hz, 1H), 4.88 - 4.82 (m, 1H), 4.50 - 4.46 (m, 1H), 4.00 - 3.85 (m, 2H), 1.31 (d, *J* = 7.5 Hz, 3H)	3418, 1717, 1165, 748, 525

FA31		598.9 ($[M+H]^+$)	(300 MHz, DMSO-d_6) δ 8.65 (d, *J* = 7.5 Hz, 1H), 8.55 (t, *J* = 5.7 Hz, 1H), 7.89 (s, 1H), 7.60 (d, *J* = 8.1 Hz, 1H), 7.46 (s, 1H), 7.41 (d, *J* = 7.8 Hz, 2H), 7.31 (s,1H), 6.96 (dd, *J* = 15.9, 8.7 Hz, 1H), 6.76 (d, *J* = 15.9 Hz, 1H), 4.71 (t, *J* = 9.3 Hz, 1H), 4.50 (t, *J* = 7.5 Hz, 1H), 4.00 - 3.88 (m, 2H), 2.67 - 2.60 (m, 2H), 1.31 (d, *J* = 7.2 Hz, 3H), 1.21 (t, *J* = 7.5 Hz, 3H)	3406, 1645, 1163, 749, 597
FA32		666.9 ($[M+H]^+$)	(300 MHz, DMSO-d_6) δ 8.68 (s, 1H), 8.54 (d, *J* = 8.4 Hz, 1H), 7.96 - 7.91 (m, 3H), 7.59 (d, *J* = 8.1 Hz, 1H), 7.33 (d, *J* = 6.6 Hz, 1H), 6.98 (dd, *J* = 15.6, 8.0 Hz, 1H), 6.77 (d, *J* = 15.9, 1H), 4.88 - 4.72 (m, 1H), 4.36 - 4.31(m, 1H), 4.07 - 3.81 (m, 2H), 2.06 - 1.99 (m, 1H), 0.95 - 0.85 (m, 6H)	3288, 1645, 1165, 750
FA33		656.9 ($[M+H]^+$)	(300 MHz, DMSO-d_6) δ 8.71 (d, *J* = 6.3 Hz 1H), 8.65 (d, *J* = 8.4 Hz, 1H), 7.98 (s, 1H), 7.92-7.87 (m, 3H), 7.45 (d, *J* = 8.1 Hz, 1H), 7.03 (dd, *J* = 1 6 Hz, 8.0 Hz, 1H), 6.89 (d, *J* = 15.9 Hz, 1H), 4.88 - 4.82 (m, 1H), 4.35 - 4.10 (m, 1H), 4.07 - 3.81 (m, 2H), 2.06 - 1.99 (m, 1H), 0.95 - 0.85 (m, 6H)	3294, 1650, 1165, 810

FA34		608.85 [(M+H)$^+$]	(300 MHz, DMSO-d_6) δ 9.01 (t, *J* = 6.0, 1H), 8.37 (t, *J* = 6.4 Hz, 1H), 8.08 (s, 1H), 8.01 - 7.96 (m, 1H), 7.88 - 7.86 (m, 2H), 7.64 (d, *J* = 7.6 Hz, 1H), 7.05 (dd, *J* = 16.4, 8.8 Hz, 1H), 6.91 (d, *J* = 15.6 Hz, 1H), 4.87 - 4.80 (m, 1H), 3.98 - 3.91 (m, 4H), 3.38 (s, 3H)	768, 721, 416, 337, 164

[a] ^{1}H NMR spectral data were acquired using a 400 MHz instrument in $CDCl_3$ except where noted. HRMS data are noted observed value (theoretical value).

Table 3: Assays Results Part 1

Compound Number	BAW Rating	CEW Rating	GPA Rating
AC1	D	D	B
AC2	C	C	C
AC3	D	D	B
AC4	D	A	B
AC5	D	D	B
AC6	D	A	B
AC7	A	A	B
AC8	D	B	B
AC9	A	A	B
AC10	A	A	B
AC11	A	A	D
AC12	A	A	D
AC13	A	A	B
AC14	A	B	D

AC15	A	A	B
AC16	A	A	C
AC17	A	A	B
AC18	A	A	B
AC19	D	D	B
AC20	A	A	C
AC21	D	D	C
AC22	A	A	D
AC23	A	A	B
AC24	A	A	D
AC25	A	A	D
AC26	A	A	B
AC27	A	A	B
AC28	A	A	B
AC29	A	A	B
AC30	A	A	B
AC31	A	A	B
AC32	A	A	B
AC33	A	A	B
AC34	A	A	B
AC35	A	A	C
AC36	A	A	B
AC37	A	A	B
AC38	A	A	C
AC39	A	A	C
AC40	A	A	D

AC41	A	D	D
AC42	A	D	D
AC43	A	A	B
AC44	A	A	B
AC45	A	A	D
AC46	A	A	D
AC47	D	D	B
AC48	A	A	B
AC49	A	A	B
AC50	A	D	B
AC51	A	A	B
AC52	A	A	B
AC53	A	A	B
AC54	A	A	B
AC57	A	A	B
AC58	A	A	B
AC59	A	A	B
AC60	A	A	B
AC61	A	A	B
AC62	A	A	D
AC63	A	A	B
AC64	A	A	B
AC65	A	A	B
AC66	A	A	B
AC67	A	A	B
AC68	A	A	D

AC69	A	A	A
AC70	D	D	B
AC71	A	A	B
AC72	A	A	B
AC75	A	A	B
AC76	A	A	D
AC77	A	A	B
AC78	A	A	A
AC79	A	A	A
AC80	A	A	B
AC81	A	D	D
AC82	A	A	B
AC83	A	A	B
AC84	A	A	D
AC85	A	A	B
AC86	A	A	D
AC87	A	A	B
AC89	A	A	B
AC90	A	A	C
AC91	A	A	C
AC92	A	A	C
AC93	A	D	C
AC94	D	B	B
AC95	A	A	C
AC96	D	D	C
AC97	D	D	C

AC98	A	A	C
AC99	A	A	C
AC100	C	C	C
AC101	D	D	C
AC102	D	A	C
AC103	A	A	D
AC104	A	A	B
AC105	A	A	D
AC106	A	A	B
AC107	B	A	D
AC108	B	D	D
AC109	D	D	C
AC110	A	A	C
AC111	A	A	C
AC112	A	A	C
AC113	B	A	D
AC114	A	B	D
AC115	A	A	D
AC116	C	C	C
AC117	A	D	B
AC118	A	D	D
BC1	A	A	D
BC2	A	A	D
BC3	A	A	D
BC4	A	A	B
BC5	A	A	B

BC6	A	A	D
BC7	A	A	D
BC8	A	A	B
BC9	A	A	D
BC10	A	A	B
BC11	C	C	C
BC12	C	C	C
BC13	A	A	D
BC14	A	D	D
CC1	D	D	D
CC2	A	A	B
CC3	A	A	D
CC4	A	B	B
CC5	A	A	B
CC6	A	A	B
CC7	A	A	B
CC8	A	A	D
CC9	A	A	B
CC10	A	A	B
CC11	A	A	B
CC12	D	D	B
CC13	A	A	B
CC14	A	D	D
CC15	A	A	B
CC16	A	A	B
CC17	A	A	B

CC18	A	A	B
CC19	A	A	B
CC20	A	A	D
CC21	A	A	D
CC22	A	A	B
CC23	A	A	B
CC24	A	A	D
CC25	A	A	B
CC26	A	D	B
CC27	A	A	D
CC28	A	A	D
CC29	A	A	B
CC30	A	A	D
CC31	B	D	C
CC32	A	A	B
CC33	A	A	B
CC34	A	A	B
CC35	D	D	D
CC36	A	A	D
CC37	A	A	D
CC38	A	A	D
CC39	D	D	B
CC40	D	A	D
CC41	D	D	B
CC42	D	D	D
CC43	A	B	B

CC44	A	A	B
CC45	A	A	D
CC46	D	A	C
CC47	D	D	C
CC48	D	D	C
CC49	D	D	D
CC50	A	A	D
CC51	A	A	D
CC52	A	D	D
CC53	D	D	B
CC54	A	A	C
DC1	A	A	D
DC2	D	D	C
DC3	B	D	C
DC4	A	D	C
DC5	D	D	C
DC6	D	D	C
DC7	A	D	C
DC8	A	D	C
DC9	D	D	C
DC10	D	D	C
DC11	A	D	C
DC12	A	A	B
DC13	A	A	C
DC14	D	D	C
DC15	D	D	C

DC16	A	A	C
DC17	A	A	C
DC18	A	A	C
DC19	A	A	C
DC20	A	D	C
DC21	D	D	C
DC22	D	D	C
DC23	D	A	C
DC24	D	D	C
DC25	D	D	C
DC26	D	D	C
DC27	D	D	C
DC28	A	A	B
DC29	A	A	C
DC30	A	A	C
DC31	A	A	B
DC32	D	D	C
DC33	A	A	C
DC34	A	A	B
DC35	A	A	B
DC36	D	D	C
DC37	A	A	C
DC38	A	A	C
DC39	A	A	C
DC40	A	A	C
DC41	A	A	C

DC42	A	A	C
DC43	A	A	C
DC44	A	A	C
DC45	A	A	C
DC46	A	A	C
DC47	A	A	C
DC48	A	A	C
DC49	A	A	C
DC50	A	A	C
DC51	A	A	C
DC52	D	D	C
DC53	D	A	C
DC54	D	D	C
DC55	D	D	C
DC56	D	D	C
DC57	A	A	C
DC58	D	D	C
DC59	D	D	C
DC60	A	A	C
DC61	D	D	C
DC62	A	A	C
DC63	A	A	C
DC64	D	D	C
DC65	D	A	C
DC66	A	A	C
DC67	A	A	C

DC68	A	A	C
DC69	D	D	C
DC70	A	A	C

Table 4: Assays Results F Compounds

Compound Number	BAW Rating	CL Rating	GPA Rating
F1	A	A	B
F2	A	A	C
F3	A	A	C
F4	A	A	C
F5	A	A	C
F6	C	C	C
F7	A	A	C
F8	A	A	C
F8A	A	A	C
F9	A	A	C
F10	A	A	C
F11	A	A	C
F12	A	A	C
F13	A	A	C
F14	A	A	C
F15	A	A	C
F15A	A	A	C
F16	A	A	C
F16A	A	A	C
F17	A	A	C

F18	A	A	C
F19	A	A	C
F20	A	A	C
F20A	A	A	A
F20B	A	A	C
F20C	A	A	C
F21	A	A	C
F22	A	A	C
F23	D	A	C
F23A	A	A	C
F24	A	A	C
F25	A	A	C
F26	A	A	C
F27	A	A	C
F28	A	A	C
F29	A	A	C
F30	A	A	C
F31	A	A	C

Table 5: Assay Results Prophetic Compounds Subsequently Exemplified

Compound Number	BAW Rating	CL Rating	GPA Rating
P1	A	A	C
P2	D	A	C
P12	A	A	C
P14	B	A	C
P15	A	A	C

P82	A	A	C
P84	A	A	C
P156	A	A	C
P226	A	A	C
P228	A	A	C
P298	D	A	B
P300	A	A	C
P442	A	A	C
P444	A	A	C
P514	A	A	C
P516	A	A	C
P568	A	A	C
P586	A	A	C
P588	A	A	C
P660	A	A	C
P730	A	A	C
P732	A	A	C
P802	A	A	C
P804	A	A	C
P1090	A	A	C
P1092	A	A	C
P1197	A	A	C
P1269	A	A	C
P1340	A	A	C
P1411	A	A	B
P1483	A	A	C

P1556	A	A	C
P1558	A	A	C
P1559	A	A	C
P1560	A	A	C
P1564	A	A	C
P1566	A	A	A
P1589	A	A	C
P1591	A	A	C
P1592	A	A	C
P1599	A	A	C
P1601	A	A	B
P1603	A	A	C
P1611A	A	A	C
P1613A	A	A	C
P1616	A	A	C
P1616A	A	A	C
P1618A	A	A	C
P1621	A	A	C
P1623	A	A	C
P1624	A	A	A
P1631A	A	A	B
P1633A	A	A	C
P1636	A	A	C
P1638	A	A	C
P1640	D	A	D
P1641	A	A	C

P1691	A	A	C
P1693	A	A	C
P1696	A	A	C
P1698	A	A	C
P1776	A	A	C
P1781	A	A	C
P1806	A	A	C
P1808	A	A	C
P1862	A	A	C
P1864	A	A	C
P1866	A	A	C
P1969*	A	A	C
P1970	A	A	C
P1971	A	A	C
P1972	A	A	C
P2009	A	A	C
P2010	A	A	C
P2011	A	A	C
P2012	A	A	C
P2036	A	A	C
P2038	A	A	C
P2041	A	A	C
P2043	A	A	C
P2056	A	A	C
P2058	A	A	C

*Performed at 12.5 ug/cm^2


Table 6: Assay Results FA Compounds

Compound Number	BAW Rating	CL Rating	GPA Rating
FA1	A	A	C
FA2	A	A	C
FA3	A	A	C
FA4	A	A	C
FA5	A	A	C
FA6	A	A	C
FA7	A	A	C
FA8	B	A	C
FA9	A	A	C
FA10	B	B	C
FA11	B	A	D
FA12	A	A	C
FA13	D	A	C
FA14	A	A	C
FA15	A	A	C
FA16	A	A	A
FA17	A	A	C
FA18	A	A	C
FA19	A	A	C
FA20	A	A	C
FA21	A	A	C
FA22	A	A	C
FA23	A	A	C

FA24	A	A	C
FA25	A	A	C
FA26	A	A	C
FA27	A	A	C
FA28	A	A	C
FA29	A	A	C
FA30	A	A	C
FA31	A	A	C
FA32	A	A	C
FA33	A	A	C
FA34	D	A	C

WE CLAIM

1. A composition comprising a molecule according to Formula One:



Formula One

wherein:

(a) **R1** is selected from

(1) H, F, Cl, Br, I, CN, NO_2, $(C_1\text{-}C_8)$alkyl, halo$(C_1\text{-}C_8)$alkyl, $(C_1\text{-}C_8)$alkoxy, halo$(C_1\text{-}C_8)$alkoxy, S$(C_1\text{-}C_8)$alkyl, S(halo$(C_1\text{-}C_8)$alkyl), S(O)$(C_1\text{-}C_8)$alkyl, S(O)(halo$(C_1\text{-}C_8)$alkyl), $S(O)_2(C_1\text{-}C_8)$alkyl, $S(O)_2$(halo$(C_1\text{-}C_8)$alkyl), N(R14)(R15),

(2) **substituted $(C_1\text{-}C_8)$alkyl**, wherein said substituted $(C_1\text{-}C_8)$alkyl has one or more substituents selected from CN and NO_2,

(3) **substituted halo$(C_1\text{-}C_8)$alkyl**, wherein said substituted halo$(C_1\text{-}C_8)$alkyl, has one or more substituents selected from CN and NO_2,

(4) **substituted $(C_1\text{-}C_8)$alkoxy**, wherein said substituted $(C_1\text{-}C_8)$alkoxy has one or more substituents selected from CN and NO_2, and

(5) **substituted halo$(C_1\text{-}C_8)$alkoxy**, wherein said substituted halo$(C_1\text{-}C_8)$alkoxy has one or more substituents selected from CN and NO_2;

(b) **R2** is selected from

(1) H, F, Cl, Br, I, CN, NO_2, $(C_1\text{-}C_8)$alkyl, halo$(C_1\text{-}C_8)$alkyl, $(C_1\text{-}C_8)$alkoxy, halo$(C_1\text{-}C_8)$alkoxy, S$(C_1\text{-}C_8)$alkyl, S(halo$(C_1\text{-}C_8)$alkyl), S(O)$(C_1\text{-}C_8)$alkyl, S(O)(halo$(C_1\text{-}C_8)$alkyl), $S(O)_2(C_1\text{-}C_8)$alkyl, $S(O)_2$(halo$(C_1\text{-}C_8)$alkyl), N(R14)(R15),

(2) **substituted $(C_1\text{-}C_8)$alkyl**, wherein said substituted $(C_1\text{-}C_8)$alkyl has one or more substituents selected from CN and NO_2,

(3) **substituted halo$(C_1\text{-}C_8)$alkyl**, wherein said substituted halo$(C_1\text{-}C_8)$alkyl, has one or more substituents selected from CN and NO_2,

(4) **substituted $(C_1\text{-}C_8)$alkoxy**, wherein said substituted $(C_1\text{-}C_8)$alkoxy has one or more substituents selected from CN and NO_2, and

(5) **substituted halo(C_1-C_8)alkoxy**, wherein said substituted halo(C_1-C_8)alkoxy has one or more substituents selected from CN and NO_2;

(c) **R3** is selected from

(1) H, F, Cl, Br, I, CN, NO_2, (C_1-C_8)alkyl, halo(C_1-C_8)alkyl, (C_1-C_8)alkoxy, halo(C_1-C_8)alkoxy, S(C_1-C_8)alkyl, S(halo(C_1-C_8)alkyl), S(O)(C_1-C_8)alkyl, S(O)(halo(C_1-C_8)alkyl), $S(O)_2$(C_1-C_8)alkyl, $S(O)_2$(halo(C_1-C_8)alkyl), N(R14)(R15),

(2) **substituted (C_1-C_8)alkyl**, wherein said substituted (C_1-C_8)alkyl has one or more substituents selected from CN and NO_2,

(3) **substituted halo(C_1-C_8)alkyl**, wherein said substituted halo(C_1-C_8)alkyl, has one or more substituents selected from CN and NO_2,

(4) **substituted (C_1-C_8)alkoxy**, wherein said substituted (C_1-C_8)alkoxy has one or more substituents selected from CN and NO_2, and

(5) **substituted halo(C_1-C_8)alkoxy**, wherein said substituted halo(C_1-C_8)alkoxy has one or more substituents selected from CN and NO_2;

(d) **R4** is selected from

(1) H, F, Cl, Br, I, CN, NO_2, (C_1-C_8)alkyl, halo(C_1-C_8)alkyl, (C_1-C_8)alkoxy, halo(C_1-C_8)alkoxy, S(C_1-C_8)alkyl, S(halo(C_1-C_8)alkyl), S(O)(C_1-C_8)alkyl, S(O)(halo(C_1-C_8)alkyl), $S(O)_2$(C_1-C_8)alkyl, $S(O)_2$(halo(C_1-C_8)alkyl), N(R14)(R15),

(2) **substituted (C_1-C_8)alkyl**, wherein said substituted (C_1-C_8)alkyl has one or more substituents selected from CN and NO_2,

(3) **substituted halo(C_1-C_8)alkyl**, wherein said substituted halo(C_1-C_8)alkyl, has one or more substituents selected from CN and NO_2,

(4) **substituted (C_1-C_8)alkoxy**, wherein said substituted (C_1-C_8)alkoxy has one or more substituents selected from CN and NO_2, and

(5) **substituted halo(C_1-C_8)alkoxy**, wherein said substituted halo(C_1-C_8)alkoxy has one or more substituents selected from CN and NO_2;

(e) **R5** is selected from

(1) H, F, Cl, Br, I, CN, NO_2, (C_1-C_8)alkyl, halo(C_1-C_8)alkyl, (C_1-C_8)alkoxy, halo(C_1-C_8)alkoxy, S(C_1-C_8)alkyl, S(halo(C_1-C_8)alkyl), S(O)(C_1-C_8)alkyl, S(O)(halo(C_1-C_8)alkyl), $S(O)_2$(C_1-C_8)alkyl, $S(O)_2$(halo(C_1-C_8)alkyl), N(R14)(R15),

(2) **substituted (C_1-C_8)alkyl**, wherein said substituted (C_1-C_8)alkyl has one or more substituents selected from CN and NO_2,

(3) **substituted halo(C_1-C_8)alkyl**, wherein said substituted halo(C_1-C_8)alkyl, has one or more substituents selected from CN and NO_2,

(4) **substituted (C_1-C_8)alkoxy**, wherein said substituted (C_1-C_8)alkoxy has one or more substituents selected from CN and NO_2, and

(5) **substituted halo(C_1-C_8)alkoxy**, wherein said substituted halo(C_1-C_8)alkoxy has one or more substituents selected from CN and NO_2;

(f) **R6** is a (C_1-C_8)haloalkyl;

(g) **R7** is selected from H, F, Cl, Br, I, OH, (C_1-C_8)alkoxy, and halo(C_1-C_8)alkoxy;

(h) **R8** is selected from H, (C_1-C_8)alkyl, halo(C_1-C_8)alkyl, OR14, and N(R14)(R15);

(i) **R9** is selected from H, F, Cl, Br, I, (C_1-C_8)alkyl, halo(C_1-C_8)alkyl, (C_1-C_8)alkoxy, halo(C_1-C_8)alkoxy, OR14, and N(R14)(R15);

(j) **R10** is selected from

(1) H, F, Cl, Br, I, CN, NO_2, (C_1-C_8)alkyl, halo(C_1-C_8)alkyl, (C_1-C_8)alkoxy, halo(C_1-C_8)alkoxy, cyclo(C_3-C_6)alkyl, S(C_1-C_8)alkyl, S(halo(C_1-C_8)alkyl), S(O)(C_1-C_8)alkyl, S(O)(halo(C_1-C_8)alkyl), S(O)$_2$(C_1-C_8)alkyl, S(O)$_2$(halo(C_1-C_8)alkyl), NR14R15, C(=O)H, C(=O)N(R14)(R15), CN(R14)(R15)(=NOH), (C=O)O(C_1-C_8)alkyl, (C=O)OH, heterocyclyl, (C_2-C_8)alkenyl, halo(C_2-C_8)alkenyl, (C_2-C_8)alkynyl,

(2) **substituted (C_1-C_8)alkyl**, wherein said substituted (C_1-C_8)alkyl has one or more substituents selected from OH, (C_1-C_8)alkoxy, S(C_1-C_8)alkyl, S(O)(C_1-C_8)alkyl, S(O)$_2$(C_1-C_8)alkyl, NR14R15, and

(3) **substituted halo(C_1-C_8)alkyl**, wherein said substituted halo(C_1-C_8)alkyl, has one or more substituents selected from (C_1-C_8)alkoxy, S(C_1-C_8)alkyl, S(O)(C_1-C_8)alkyl, S(O)$_2$(C_1-C_8)alkyl, and N(R14)(R15);

(k) **R11** is selected from C(=X5)N(R14)((C_1-C_8)alkylC(=X5)N(R14)(R15)) wherein each X5 is independently selected from O, or S;

(l) **R12** is selected from **(v)**, H, F, Cl, Br, I, CN, (C_1-C_8)alkyl, halo(C_1-C_8)alkyl, (C_1-C_8)alkoxy, halo(C_1-C_8)alkoxy, and cyclo(C_3-C_6)alkyl;

(m) **R13** is selected from **(v)**, H, F, Cl, Br, I, CN, (C_1-C_8)alkyl, halo(C_1-C_8)alkyl, (C_1-C_8)alkoxy, and halo(C_1-C_8)alkoxy;

(n) **each R14** is independently selected from H, (C_1-C_8)alkyl, (C_2-C_8)alkenyl, substituted (C_1-C_8)alkyl, halo(C_1-C_8)alkyl, substituted halo(C_1-C_8)alkyl), (C_1-C_8)alkoxy, cyclo(C_3-C_6)alkyl, aryl, substituted-aryl, (C_1-C_8)alkyl-aryl, (C_1-C_8)alkyl-(substituted-aryl), O-(C_1-C_8)alkyl-aryl, O-(C_1-C_8)alkyl-(substituted-aryl), heterocyclyl, substituted-heterocyclyl, (C_1-C_8)alkyl-heterocyclyl, (C_1-C_8)alkyl-(substituted-heterocyclyl), O-(C_1-C_8)alkyl-

heterocyclyl, O-(C_1-C_8)alkyl-(substituted-heterocyclyl), N(R16)(R17), (C_1-C_8)alkyl-C(=O)N(R16)(R17), C(=O)(C_1-C_8)alkyl, C(=O)(halo(C_1-C_8)alkyl),C(=O)(C_3-C_6)cycloalkyl, (C_1-C_8)alkyl-C(=O)O(C_1-C_8)alkyl, C(=O)H

wherein each said substituted (C_1-C_8)alkyl has one or more substituents selected from CN, and NO_2,

wherein each said substituted halo(C_1-C_8)alkyl), has one or more substituents selected from CN, and NO_2,

wherein each said substituted-aryl has one or more substituents selected from F, Cl, Br, I, CN, NO_2, (C_1-C_8)alkyl, halo(C_1-C_8)alkyl, (C_1-C_8)alkoxy, halo(C_1-C_8)alkoxy, S(C_1-C_8)alkyl, S(halo(C_1-C_8)alkyl), N((C_1-C_8)alkyl)$_2$ (wherein each (C_1-C_8)alkyl is independently selected), and oxo, and

wherein each said substituted-heterocyclyl has one or more substituents selected from F, Cl, Br, I, CN, NO_2, (C_1-C_8)alkyl, halo(C_1-C_8)alkyl, (C_1-C_8)alkoxy, halo(C_1-C_8)alkoxy, (C_3-C_6)cycloalkyl S(C_1-C_8)alkyl, S(halo(C_1-C_8)alkyl), N((C_1-C_8)alkyl)$_2$ (wherein each (C_1-C_8)alkyl is independently selected), heterocyclyl, C(=O)(C_1-C_8)alkyl, C(=O)O(C_1-C_8)alkyl, and oxo, (wherein said alkyl, alkoxy, and heterocyclyl, may be further substituted with one or more of F, Cl, Br, I, CN, and NO_2);

(o) each R15 is independently selected from H, (C_1-C_8)alkyl, (C_2-C_8)alkenyl, substituted (C_1-C_8)alkyl, halo(C_1-C_8)alkyl, substituted halo(C_1-C_8)alkyl), (C_1-C_8)alkoxy, cyclo(C_3-C_6)alkyl, aryl, substituted-aryl, (C_1-C_8)alkyl-aryl, (C_1-C_8)alkyl-(substituted-aryl), O-(C_1-C_8)alkyl-aryl, O-(C_1-C_8)alkyl-(substituted-aryl), heterocyclyl, substituted-heterocyclyl, (C_1-C_8)alkyl-heterocyclyl, (C_1-C_8)alkyl-(substituted-heterocyclyl), O-(C_1-C_8)alkyl-heterocyclyl, O-(C_1-C_8)alkyl-(substituted-heterocyclyl), N(R16)(R17), (C_1-C_8)alkyl-C(=O)N(R16)(R17), C(=O)(C_1-C_8)alkyl, C(=O)(halo(C_1-C_8)alkyl), C(=O)(C_3-C_6)cycloalkyl, (C_1-C_8)alkyl-C(=O)O(C_1-C_8)alkyl, C(=O)H

wherein each said substituted (C_1-C_8)alkyl has one or more substituents selected from CN, and NO_2,

wherein each said substituted halo(C_1-C_8)alkyl), has one or more substituents selected from CN, and NO_2,

wherein each said substituted-aryl has one or more substituents selected from F, Cl, Br, I, CN, NO_2, (C_1-C_8)alkyl, halo(C_1-C_8)alkyl, (C_1-C_8)alkoxy, halo(C_1-C_8)alkoxy, S(C_1-C_8)alkyl, S(halo(C_1-C_8)alkyl), N((C_1-C_8)alkyl)$_2$ (wherein each (C_1-C_8)alkyl is independently selected), and oxo, and

wherein each said substituted-heterocyclyl has one or more substituents selected from F, Cl, Br, I, CN, NO_2, $(C_1$-$C_8)$alkyl, halo$(C_1$-$C_8)$alkyl, $(C_1$-$C_8)$alkoxy, halo$(C_1$-$C_8)$alkoxy, $(C_3$-$C_6)$cycloalkyl S$(C_1$-$C_8)$alkyl, S(halo$(C_1$-$C_8)$alkyl), N$((C_1$-$C_8)$alkyl$)_2$ (wherein each $(C_1$-$C_8)$alkyl is independently selected), heterocyclyl, C(=O)$(C_1$-$C_8)$alkyl, C(=O)O$(C_1$-$C_8)$alkyl, and oxo, (wherein said alkyl, alkoxy, and heterocyclyl, may be further substituted with one or more of F, Cl, Br, I, CN, and NO_2);

(p) each R16 is independently selected from H, $(C_1$-$C_8)$alkyl, substituted-$(C_1$-$C_8)$alkyl, halo$(C_1$-$C_8)$alkyl, substituted-halo$(C_1$-$C_8)$alkyl, cyclo$(C_3$-$C_6)$alkyl, aryl, substituted-aryl, $(C_1$-$C_8)$alkyl-aryl, $(C_1$-$C_8)$alkyl-(substituted-aryl), O-$(C_1$-$C_8)$alkyl-aryl, O-$(C_1$-$C_8)$alkyl-(substituted-aryl), heterocyclyl, substituted-heterocyclyl, $(C_1$-$C_8)$alkyl-heterocyclyl, $(C_1$-$C_8)$alkyl-(substituted-heterocyclyl), O-$(C_1$-$C_8)$alkyl-heterocyclyl, O-$(C_1$-$C_8)$alkyl-(substituted-heterocyclyl), O-$(C_1$-$C_8)$alkyl

wherein each said substituted $(C_1$-$C_8)$alkyl has one or more substituents selected from CN, and NO_2,

wherein each said substituted halo$(C_1$-$C_8)$alkyl), has one or more substituents selected from CN, and NO_2,

wherein each said substituted-aryl has one or more substituents selected from F, Cl, Br, I, CN, NO_2, $(C_1$-$C_8)$alkyl, halo$(C_1$-$C_8)$alkyl, $(C_1$-$C_8)$alkoxy, halo$(C_1$-$C_8)$alkoxy, S$(C_1$-$C_8)$alkyl, S(halo$(C_1$-$C_8)$alkyl), N$((C_1$-$C_8)$alkyl$)_2$ (wherein each $(C_1$-$C_8)$alkyl is independently selected), and oxo, and

wherein each said substituted-heterocyclyl has one or more substituents selected from F, Cl, Br, I, CN, NO_2, $(C_1$-$C_8)$alkyl, halo$(C_1$-$C_8)$alkyl, $(C_1$-$C_8)$alkoxy, halo$(C_1$-$C_8)$alkoxy, S$(C_1$-$C_8)$alkyl, S(halo$(C_1$-$C_8)$alkyl), N$((C_1$-$C_8)$alkyl$)_2$ (wherein each $(C_1$-$C_8)$alkyl is independently selected), and oxo;

(q) each R17 is independently selected from H, $(C_1$-$C_8)$alkyl, substituted-$(C_1$-$C_8)$alkyl, halo$(C_1$-$C_8)$alkyl, substituted-halo$(C_1$-$C_8)$alkyl, cyclo$(C_3$-$C_6)$alkyl, aryl, substituted-aryl, $(C_1$-$C_8)$alkyl-aryl, $(C_1$-$C_8)$alkyl-(substituted-aryl), O-$(C_1$-$C_8)$alkyl-aryl, O-$(C_1$-$C_8)$alkyl-(substituted-aryl), heterocyclyl, substituted-heterocyclyl, $(C_1$-$C_8)$alkyl-heterocyclyl, $(C_1$-$C_8)$alkyl-(substituted-heterocyclyl), O-$(C_1$-$C_8)$alkyl-heterocyclyl, O-$(C_1$-$C_8)$alkyl-(substituted-heterocyclyl), O-$(C_1$-$C_8)$alkyl

wherein each said substituted $(C_1$-$C_8)$alkyl has one or more substituents selected from CN, and NO_2,

wherein each said substituted halo$(C_1$-$C_8)$alkyl), has one or more substituents selected from CN, and NO_2,

2013361540 12 Dec 2016

wherein each said substituted-aryl has one or more substituents selected from F, Cl, Br, I, CN, NO_2, $(C_1$-$C_8)$alkyl, halo$(C_1$-$C_8)$alkyl, $(C_1$-$C_8)$alkoxy, halo$(C_1$-$C_8)$alkoxy, S$(C_1$-$C_8)$alkyl, S(halo$(C_1$-$C_8)$alkyl), N$((C_1$-$C_8)$alkyl$)_2$ (wherein each $(C_1$-$C_8)$alkyl is independently selected), and oxo, and

wherein each said substituted-heterocyclyl has one or more substituents selected from F, Cl, Br, I, CN, NO_2, $(C_1$-$C_8)$alkyl, halo$(C_1$-$C_8)$alkyl, $(C_1$-$C_8)$alkoxy, halo$(C_1$-$C_8)$alkoxy, S$(C_1$-$C_8)$alkyl, S(halo$(C_1$-$C_8)$alkyl), N$((C_1$-$C_8)$alkyl$)_2$ (wherein each $(C_1$-$C_8)$alkyl is independently selected), and oxo;

(r) **X1** is selected from N and CR12;

(s) **X2** is selected from N, CR9, and CR13;

(t) **X3** is selected from N and CR9; or

(v) R12 and R13 together form a linkage containing 3 to 4 atoms selected from C, N, O, and S, wherein said linkage connects back to the ring to form a 5 to 6 member saturated or unsaturated cyclic ring, wherein said linkage has at least one substituent X4 wherein X4 is selected from R14, N(R14)(R15), N(R14)(C(=O)R14), N(R14)(C(=S)R14), N(R14)(C(=O)N(R14)(R14)), N(R14)(C(=S)N(R14)(R14)), N(R14)(C(=O)N(R14)$((C_2$-$C_8)$alkenyl)), N(R14)(C(=S)N(R14)$((C_2$-$C_8)$alkenyl)), wherein each R14 is independently selected.

2. A composition according to claim 1, wherein R1 is selected from H, F, Cl, Br, I, CN, NO_2, methyl, ethyl, (C_3)alkyl, (C_4)alkyl, (C_5)alkyl, (C_6)alkyl, (C_7)alkyl, (C_8)alkyl, halomethyl, haloethyl, halo(C_3)alkyl, halo(C_4)alkyl, halo(C_5)alkyl, halo(C_6)alkyl, halo(C_7)alkyl, halo(C_8)alkyl, methoxy, ethoxy, (C_3)alkoxy, (C_4)alkoxy, (C_5)alkoxy, (C_6)alkoxy, (C_7)alkoxy, (C_8)alkoxy, halomethoxy, haloethoxy, halo(C_3)alkoxy, halo(C_4)alkoxy, halo(C_5)alkoxy, halo(C_6)alkoxy, halo(C_7)alkoxy, and halo(C_8)alkoxy; or R1 is selected from Cl and H;

or

wherein R2 is selected from H, F, Cl, Br, I, CN, NO_2, methyl, ethyl, (C_3)alkyl, (C_4)alkyl, (C_5)alkyl, (C_6)alkyl, (C_7)alkyl, (C_8)alkyl, halomethyl, haloethyl, halo(C_3)alkyl, halo(C_4)alkyl, halo(C_5)alkyl, halo(C_6)alkyl, halo(C_7)alkyl, halo(C_8)alkyl, methoxy, ethoxy, (C_3)alkoxy, (C_4)alkoxy, (C_5)alkoxy, (C_6)alkoxy, (C_7)alkoxy, (C_8)alkoxy, halomethoxy, haloethoxy, halo(C_3)alkoxy, halo(C_4)alkoxy, halo(C_5)alkoxy, halo(C_6)alkoxy, halo(C_7)alkoxy, and halo(C_8)alkoxy; or R2 is selected from CF_3, CH_3, Cl, F, and H;

or

2013361540 12 Dec 2016

wherein R3 is selected from H, F, Cl, Br, I, CN, NO_2, methyl, ethyl, (C_3)alkyl, (C_4)alkyl, (C_5)alkyl, (C_6)alkyl, (C_7)alkyl, (C_8)alkyl, halomethyl, haloethyl, halo(C_3)alkyl, halo(C_4)alkyl, halo(C_5)alkyl, halo(C_6)alkyl, halo(C_7)alkyl, halo(C_8)alkyl, methoxy, ethoxy, (C_3)alkoxy, (C_4)alkoxy, (C_5)alkoxy, (C_6)alkoxy, (C_7)alkoxy, (C_8)alkoxy, halomethoxy, haloethoxy, halo(C_3)alkoxy, halo(C_4)alkoxy, halo(C_5)alkoxy, halo(C_6)alkoxy, halo(C_7)alkoxy, and halo(C_8)alkoxy; or R3 is selected from OCH_3, CH_3, F, Cl, or H;

or

wherein R4 is selected from H, F, Cl, Br, I, CN, NO_2, methyl, ethyl, (C_3)alkyl, (C_4)alkyl, (C_5)alkyl, (C_6)alkyl, (C_7)alkyl, (C_8)alkyl, halomethyl, haloethyl, halo(C_3)alkyl, halo(C_4)alkyl, halo(C_5)alkyl, halo(C_6)alkyl, halo(C_7)alkyl, halo(C_8)alkyl, methoxy, ethoxy, (C_3)alkoxy, (C_4)alkoxy, (C_5)alkoxy, (C_6)alkoxy, (C_7)alkoxy, (C_8)alkoxy, halomethoxy, haloethoxy, halo(C_3)alkoxy, halo(C_4)alkoxy, halo(C_5)alkoxy, halo(C_6)alkoxy, halo(C_7)alkoxy, and halo(C_8)alkoxy; or R4 is selected from CF_3, CH_3, Cl, F, and H;

or

wherein R5 is selected from H, F, Cl, Br, I, CN, NO_2, methyl, ethyl, (C_3)alkyl, (C_4)alkyl, (C_5)alkyl, (C_6)alkyl, (C_7)alkyl, (C_8)alkyl, halomethyl, haloethyl, halo(C_3)alkyl, halo(C_4)alkyl, halo(C_5)alkyl, halo(C_6)alkyl, halo(C_7)alkyl, halo(C_8)alkyl, methoxy, ethoxy, (C_3)alkoxy, (C_4)alkoxy, (C_5)alkoxy, (C_6)alkoxy, (C_7)alkoxy, (C_8)alkoxy, halomethoxy, haloethoxy, halo(C_3)alkoxy, halo(C_4)alkoxy, halo(C_5)alkoxy, halo(C_6)alkoxy, halo(C_7)alkoxy, and halo(C_8)alkoxy; or R5 is selected from F, Cl, and H;

or

wherein R6 is selected from halomethyl, haloethyl, halo(C_3)alkyl, halo(C_4)alkyl, halo(C_5)alkyl, halo(C_6)alkyl, halo(C_7)alkyl, and halo(C_8)alkyl; or R6 is trifluoromethyl;

or

wherein R7 is selected from H, F, Cl, Br, and I; or R7 is selected from H, OCH_3, and OH;

or

wherein R8 is selected from H, methyl, ethyl, (C_3)alkyl, (C_4)alkyl, (C_5)alkyl, (C_6)alkyl, (C_7)alkyl, (C_8)alkyl, halomethyl, haloethyl, halo(C_3)alkyl, halo(C_4)alkyl, halo(C_5)alkyl, halo(C_6)alkyl, halo(C_7)alkyl, and halo(C_8)alkyl; or R8 is selected from CH_3 and H;

or

wherein R9 is selected from H, F, Cl, Br, I, methyl, ethyl, (C_3)alkyl, (C_4)alkyl, (C_5)alkyl, (C_6)alkyl, (C_7)alkyl, (C_8)alkyl, halomethyl, haloethyl, halo(C_3)alkyl, halo(C_4)alkyl, halo(C_5)alkyl, halo(C_6)alkyl, halo(C_7)alkyl, halo(C_8)alkyl, methoxy, ethoxy, (C_3)alkoxy, (C_4)alkoxy, (C_5)alkoxy, (C_6)alkoxy, (C_7)alkoxy, (C_8)alkoxy, halomethoxy, haloethoxy,

2013361540 12 Dec 2016

halo(C_3)alkoxy, halo(C_4)alkoxy, halo(C_5)alkoxy, halo(C_6)alkoxy, halo(C_7)alkoxy, and halo(C_8)alkoxy;

or

wherein R10 is selected from H, F, Cl, Br, I, CN, methyl, ethyl, (C_3)alkyl, (C_4)alkyl, (C_5)alkyl, (C_6)alkyl, (C_7)alkyl, (C_8)alkyl, halomethyl, haloethyl, halo(C_3)alkyl, halo(C_4)alkyl, halo(C_5)alkyl, halo(C_6)alkyl, halo(C_7)alkyl, halo(C_8)alkyl, methoxy, ethoxy, (C_3)alkoxy, (C_4)alkoxy, (C_5)alkoxy, (C_6)alkoxy, (C_7)alkoxy, (C_8)alkoxy, halomethoxy, haloethoxy, halo(C_3)alkoxy, halo(C_4)alkoxy, halo(C_5)alkoxy, halo(C_6)alkoxy, halo(C_7)alkoxy, halo(C_8)alkoxy, cyclopropyl, cyclobutyl, cyclopentyl, and cyclohexyl; or wherein R10 is selected from H, Cl, Br, CH_3, and CF_3; or wherein R10 is selected from Br, $C(=NOH)NH_2$, C(=O)H, $C(=O)NH_2$, $C(=O)OCH_2CH_3$, C(=O)OH, CF_3, CH_2CH_3, CH_2OH, CH3, Cl, CN, F, H, NH_2, NHC(=O)H, $NHCH_3$, NO_2, OCH_3, $OCHF_2$, and pyridyl;

or

wherein R11 is selected from $C(=O)N(H)(C((CH_3)_2)C(=O)N(H)(CH_2CF_3))$, $C(=O)N(H)(CH(CH_3)C(=O)N(H)(CH_2CF_3))$, $C(=O)N(H)(CH(CH_2CH_3)C(=O)N(H)(CH_2CF_3))$, $C(=O)N(H)(CH(CH_3)C(=S)N(H)(CH_2CF_3))$, $C(=O)N(H)(C((CH_3)_2)C(=S)N(H)(CH_2CF_3))$, and $C(=S)N(H)(C((CH_3)_2)C(=S)N(H)(CH_2CF_3))$; or wherein R11 is C(=(O or S))N(H)(((C$_1$-C$_8$)alkyl)C(=(O or S))N(H)(halo(C$_1$-C$_8$)alkyl)); or wherein R11 is selected from $C(=O)N(H)(C((CH_3)_2)C(=O)N(H)(CH_2CF_3))$, $C(=O)N(H)(CH(CH_3)C(=O)N(H)(CH_2CF_3))$, $C(=O)N(H)(CH(CH_3)C(=S)N(H)(CH_2CF_3))$, $C(=O)N(H)(C((CH_3)_2)C(=S)N(H)(CH_2CF_3))$, and $C(=S)N(H)(C((CH_3)_2)C(=S)N(H)(CH_2CF_3))$; or wherein R11 is selected from $C(=O)N(H)(C((CH_3)_2)C(=O)N(H)(CH_2CF_3))$;

or

wherein R12 is selected from H, F, Cl, Br, I, methyl, ethyl, (C_3)alkyl, (C_4)alkyl, (C_5)alkyl, (C_6)alkyl, (C_7)alkyl, (C_8)alkyl, halomethyl, haloethyl, halo(C_3)alkyl, halo(C_4)alkyl, halo(C_5)alkyl, halo(C_6)alkyl, halo(C_7)alkyl, halo(C_8)alkyl, halomethoxy, haloethoxy, halo(C_3)alkoxy, halo(C_4)alkoxy, halo(C_5)alkoxy, halo(C_6)alkoxy, halo(C_7)alkoxy, and halo(C_8)alkoxy; or wherein R12 is selected from CH_3 and H;

or

wherein R13 is selected from H, F, Cl, Br, I, methyl, ethyl, (C_3)alkyl, (C_4)alkyl, (C_5)alkyl, (C_6)alkyl, (C_7)alkyl, (C_8)alkyl, halomethyl, haloethyl, halo(C_3)alkyl, halo(C_4)alkyl, halo(C_5)alkyl, halo(C_6)alkyl, halo(C_7)alkyl, halo(C_8)alkyl, halomethoxy, haloethoxy,

2013361540 12 Dec 2016

halo(C_3)alkoxy, halo(C_4)alkoxy, halo(C_5)alkoxy, halo(C_6)alkoxy, halo(C_7)alkoxy, and halo(C_8)alkoxy; or wherein R13 is selected from CH_3, Cl, and H;

or

wherein R14 and R15 are independently selected from H, methyl, ethyl, (C_3)alkyl, (C_4)alkyl, (C_5)alkyl, (C_6)alkyl, (C_7)alkyl, (C_8)alkyl, halomethyl, haloethyl, halo(C_3)alkyl, halo(C_4)alkyl, halo(C_5)alkyl, halo(C_6)alkyl, halo(C_7)alkyl, halo(C_8)alkyl, methyl-aryl, ethyl-aryl, (C_3)alkyl-aryl, (C_4)alkyl-aryl, (C_5)alkyl-aryl, (C_6)alkyl-aryl, (C_7)alkyl-aryl, (C_8)alkyl-aryl, methyl-(substituted-aryl), ethyl-(substituted-aryl), (C_3)alkyl-(substituted-aryl), (C_4)alkyl-(substituted-aryl), (C_5)alkyl-(substituted-aryl), (C_6)alkyl-(substituted-aryl), (C_7)alkyl-(substituted-aryl), (C_8)alkyl-(substituted-aryl), O-methyl-aryl, O-ethyl-aryl, O-(C_3)alkyl-aryl, O-(C_4)alkyl-aryl, O-(C_5)alkyl-aryl, O-(C_6)alkyl-aryl, O-(C_7)alkyl-aryl, O-(C_8)alkyl-aryl, O-methyl-(substituted-aryl), O-ethyl-(substituted-aryl), O-(C_3)alkyl-(substituted-aryl), O-(C_4)alkyl-(substituted-aryl), O-(C_5)alkyl-(substituted-aryl), O-(C_6)alkyl-(substituted-aryl), O-(C_7)alkyl-(substituted-aryl), O-(C_8)alkyl-(substituted-aryl), methyl-heterocyclyl, ethyl-heterocyclyl, (C_3)alkyl-heterocyclyl, (C_4)alkyl-heterocyclyl, (C_5)alkyl-heterocyclyl, (C_6)alkyl-heterocyclyl, (C_7)alkyl-heterocyclyl, (C_8)alkyl-heterocyclyl, methyl-(substituted-heterocyclyl), ethyl-(substituted-heterocyclyl), (C_3)alkyl-(substituted-heterocyclyl), (C_4)alkyl-(substituted-heterocyclyl), (C_5)alkyl-(substituted-heterocyclyl), (C_6)alkyl-(substituted-heterocyclyl), (C_7)alkyl-(substituted-heterocyclyl), (C_8)alkyl-(substituted-heterocyclyl), O-methyl-heterocyclyl, O-ethyl-heterocyclyl, O-(C_3)alkyl-heterocyclyl, O-(C_4)alkyl-heterocyclyl, O-(C_5)alkyl-heterocyclyl, O-(C_6)alkyl-heterocyclyl, O-(C_7)alkyl-heterocyclyl, O-(C_8)alkyl-heterocyclyl, O-methyl-(substituted-heterocyclyl), O-ethyl-(substituted-heterocyclyl), O-(C_3)alkyl-(substituted-heterocyclyl), O-(C_4)alkyl-(substituted-heterocyclyl), O-(C_5)alkyl-(substituted-heterocyclyl), O-(C_6)alkyl-(substituted-heterocyclyl), O-(C_7)alkyl-(substituted-heterocyclyl), O-(C_8)alkyl-(substituted-heterocyclyl), methyl-C(=O)N(R16)(R17), ethyl-C(=O)N(R16)(R17), (C_3)alkyl-C(=O)N(R16)(R17), (C_4)alkyl-C(=O)N(R16)(R17), (C_5)alkyl-C(=O)N(R16)(R17), (C_6)alkyl-C(=O)N(R16)(R17), (C_7)alkyl-C(=O)N(R16)(R17), and (C_8)alkyl-C(=O)N(R16)(R17); or wherein R14 and R15 are independently selected from H, CH_3, CH_2CF_3, CH_2-halopyridyl, oxo-pyrrolidinyl, halophenyl, thietanyl, CH_2-phenyl, CH_2-pyridyl, thietanyl-dioxide, CH_2-halothiazolyl, C((CH_3)$_2$)-pyridyl, N(H)(halophenyl), CH_2-pyrimidinyl, CH_2-tetrahydrofuranyl, CH_2-furanyl, O-CH_2-halopyridyl, and $CH_2C(=O)N(H)(CH_2CF_3)$;

or

2013361540 12 Dec 2016

wherein R16 and R17 are independently selected from H, methyl, ethyl, (C_3)alkyl, (C_4)alkyl, (C_5)alkyl, (C_6)alkyl, (C_7)alkyl, (C_8)alkyl, halomethyl, haloethyl, halo(C_3)alkyl, halo(C_4)alkyl, halo(C_5)alkyl, halo(C_6)alkyl, halo(C_7)alkyl, halo(C_8)alkyl, methyl-aryl, ethyl-aryl, (C_3)alkyl-aryl, (C_4)alkyl-aryl, (C_5)alkyl-aryl, (C_6)alkyl-aryl, (C_7)alkyl-aryl, (C_8)alkyl-aryl, methyl-(substituted-aryl), ethyl-(substituted-aryl), (C_3)alkyl-(substituted-aryl), (C_4)alkyl-(substituted-aryl), (C_5)alkyl-(substituted-aryl), (C_6)alkyl-(substituted-aryl), (C_7)alkyl-(substituted-aryl), (C_8)alkyl-(substituted-aryl), O-methyl-aryl, O-ethyl-aryl, O-(C_3)alkyl-aryl, O-(C_4)alkyl-aryl, O-(C_5)alkyl-aryl, O-(C_6)alkyl-aryl, O-(C_7)alkyl-aryl, O-(C_8)alkyl-aryl, O-methyl-(substituted-aryl), O-ethyl-(substituted-aryl), O-(C_3)alkyl-(substituted-aryl), O-(C_4)alkyl-(substituted-aryl), O-(C_5)alkyl-(substituted-aryl), O-(C_6)alkyl-(substituted-aryl), O-(C_7)alkyl-(substituted-aryl), O-(C_8)alkyl-(substituted-aryl), methyl-heterocyclyl, ethyl-heterocyclyl, (C_3)alkyl-heterocyclyl, (C_4)alkyl-heterocyclyl, (C_5)alkyl-heterocyclyl, (C_6)alkyl-heterocyclyl, (C_7)alkyl-heterocyclyl, (C_8)alkyl-heterocyclyl, methyl-(substituted-heterocyclyl), ethyl-(substituted-heterocyclyl), (C_3)alkyl-(substituted-heterocyclyl), (C_4)alkyl-(substituted-heterocyclyl), (C_5)alkyl-(substituted-heterocyclyl), (C_6)alkyl-(substituted-heterocyclyl), (C_7)alkyl-(substituted-heterocyclyl), (C_8)alkyl-(substituted-heterocyclyl), O-methyl-heterocyclyl, O-ethyl-heterocyclyl, O-(C_3)alkyl-heterocyclyl, O-(C_4)alkyl-heterocyclyl, O-(C_5)alkyl-heterocyclyl, O-(C_6)alkyl-heterocyclyl, O-(C_7)alkyl-heterocyclyl, O-(C_8)alkyl-heterocyclyl, O-methyl-(substituted-heterocyclyl), O-ethyl-(substituted-heterocyclyl), O-(C_3)alkyl-(substituted-heterocyclyl), O-(C_4)alkyl-(substituted-heterocyclyl), O-(C_5)alkyl-(substituted-heterocyclyl), O-(C_6)alkyl-(substituted-heterocyclyl), O-(C_7)alkyl-(substituted-heterocyclyl), and O-(C_8)alkyl-(substituted-heterocyclyl); wherein R16 and R17 are independently selected from H, CH_2CF_3, cyclopropyl, thietanyl, thietanyl dioxide, and halophenyl; or

wherein X1 is CR12, X2 is CR13, and X3 is CR9;

or






wherein R12-R13 is the hydrocarbyl linkage CH=CHCH=CH.






3. A composition according to claim 1 or 2, wherein R2 and R4 are selected from F, Cl, Br, I, CN, and NO_2 and R1, R3, and R5 are H; or






wherein R2, R3, and R4 are selected from F, Cl, Br, I, CN, and NO_2 and R1, and R5 are H; or






wherein R2, R3, and R4 are independently selected from F and Cl and R1and R5 are H.






4. A composition according to claim 1 wherein said molecule has the following structure






Compound Number	Structure
F1	
F2	
F3	
F4	
F5	






F6	
F7	
F8	
F8A	
F9	




F10	 (+)-
F11	 (−)-
F12	 (+)-
F13	 (−)-
F14	




F15	
F15A	
F16	
F16A	
F17	









F18	
F19	
F20	
F20A	
F20B	









F20C	
F21	
F22	
F23	
F23A	









F24	
F25	
F26	
F27	
F28	









F29	
F30	
F31	









Compound Number	Structure
P1	
P2	
P12	









P14	
P15	
P82	
P84	
P156	
P226	
P228	
P298	









P300	
P442	
P444	
P514	
P516	
P568	
P586	
P588	









P660	
P730	
P732	
P802	
P804	
P1090	
P1092	
P1197	









P1269	
P1340	
P1411	
P1483	
P1556	
P1558	
P1559	
P1560	








P1564	
P1566	
P1589	
P1591	
P1592	
P1599	
P1601	
P1603	

P1611A	
P1613A	
P1616	
P1616A	
P1618A	
P1621	
P1623	
P1624	









P1631A	
P1633A	
P1636	
P1638	
P1640	
P1641	
P1691	
P1693	









P1696	
P1698	
P1776	
P1781	
P1806	
P1808	
P1862	
P1864	









P1866	
P1969	
P1970	
P1971	
P1972	
P2009	
P2010	
P2011	








P2012	
P2036	
P2038	
P2041	
P2043	
P2056	
P2058	

Compound Number	Structure




FA1	
FA2	
FA3	
FA4	
FA5	
FA6	
FA7	
FA8	

FA9	
FA10	
FA11	
FA12	
FA13	
FA14	
FA15	
FA16	




FA17	
FA18	
FA19	
FA20	
FA21	
FA22	
FA23	
FA24	

FA25	
FA26	
FA27	
FA28	
FA29	
FA30	
FA31	


2013361540 12 Dec 2016

FA32	
FA33	
FA34	

or a composition according to claim 1 having one of the following structures

F20	
F20A	
F20B	

12 Dec 2016 2013361540

F20C	

5. A composition according to any one of the previous claims, further comprising:

(a) one or more compounds having acaricidal, algicidal, avicidal, bactericidal, fungicidal, herbicidal, insecticidal, molluscicidal, nematicidal, rodenticidal, or virucidal properties; or

(b) one or more compounds that are antifeedants, bird repellents, chemosterilants, herbicide safeners, insect attractants, insect repellents, mammal repellents, mating disrupters, plant activators, plant growth regulators, or synergists; or

(c) both (a) and (b); or

(d) one or more compounds selected from: (3-ethoxypropyl)mercury bromide, 1,2-dichloropropane, 1,3-dichloropropene, 1-methylcyclopropene, 1-naphthol, 2-(octylthio)ethanol, 2,3,5-tri-iodobenzoic acid, 2,3,6-TBA, 2,3,6-TBA-dimethylammonium, 2,3,6-TBA-lithium, 2,3,6-TBA-potassium, 2,3,6-TBA-sodium, 2,4,5-T, 2,4,5-T-2-butoxypropyl, 2,4,5-T-2-ethylhexyl, 2,4,5-T-3-butoxypropyl, 2,4,5-TB, 2,4,5-T-butometyl, 2,4,5-T-butotyl, 2,4,5-T-butyl, 2,4,5-T-isobutyl, 2,4,5-T-isoctyl, 2,4,5-T-isopropyl, 2,4,5-T-methyl, 2,4,5-T-pentyl, 2,4,5-T-sodium, 2,4,5-T-triethylammonium, 2,4,5-T-trolamine, 2,4-D, 2,4-D-2-butoxypropyl, 2,4-D-2-ethylhexyl, 2,4-D-3-butoxypropyl, 2,4-D-ammonium, 2,4-DB, 2,4-DB-butyl, 2,4-DB-dimethylammonium, 2,4-DB-isoctyl, 2,4-DB-potassium, 2,4-DB-sodium, 2,4-D-butotyl, 2,4-D-butyl, 2,4-D-diethylammonium, 2,4-D-dimethylammonium, 2,4-D-diolamine, 2,4-D-dodecylammonium, 2,4-DEB, 2,4-DEP, 2,4-D-ethyl, 2,4-D-heptylammonium, 2,4-D-isobutyl, 2,4-D-isoctyl, 2,4-D-isopropyl, 2,4-D-isopropylammonium, 2,4-D-lithium, 2,4-D-meptyl, 2,4-D-methyl, 2,4-D-octyl, 2,4-D-pentyl, 2,4-D-potassium, 2,4-D-propyl, 2,4-D-sodium, 2,4-D-tefuryl, 2,4-D-tetradecylammonium, 2,4-D-triethylammonium, 2,4-D-tris(2-hydroxypropyl)ammonium, 2,4-D-trolamine, 2iP, 2-methoxyethylmercury chloride, 2-phenylphenol, 3,4-DA, 3,4-DB, 3,4-DP, 4-aminopyridine, 4-CPA, 4-CPA-potassium, 4-CPA-sodium, 4-CPB, 4-CPP, 4-hydroxyphenethyl alcohol, 8-hydroxyquinoline sulfate, 8-phenylmercurioxyquinoline, abamectin, abscisic acid, ACC, acephate, acequinocyl, acetamiprid, acethion, acetochlor, acetophos, acetoprole, acibenzolar,

2013361540 12 Dec 2016

acibenzolar-S-methyl, acifluorfen, acifluorfen-methyl, acifluorfen-sodium, aclonifen, acrep, acrinathrin, acrolein, acrylonitrile, acypetacs, acypetacs-copper, acypetacs-zinc, alachlor, alanycarb, albendazole, aldicarb, aldimorph, aldoxycarb, aldrin, allethrin, allicin, allidochlor, allosamidin, alloxydim, alloxydim-sodium, allyl alcohol, allyxycarb, alorac, *alpha*-cypermethrin, *alpha*-endosulfan, ametoctradin, ametridione, ametryn, amibuzin, amicarbazone, amicarthiazol, amidithion, amidoflumet, amidosulfuron, aminocarb, aminocyclopyrachlor, aminocyclopyrachlor-methyl, aminocyclopyrachlor-potassium, aminopyralid, aminopyralid-potassium, aminopyralid-tris(2-hydroxypropyl)ammonium, amiprofos-methyl, amiprophos, amisulbrom, amiton, amiton oxalate, amitraz, amitrole, ammonium sulfamate, ammonium α-naphthaleneacetate, amobam, ampropylfos, anabasine, ancymidol, anilazine, anilofos, anisuron, anthraquinone, antu, apholate, aramite, arsenous oxide, asomate, aspirin, asulam, asulam-potassium, asulam-sodium, athidathion, atraton, atrazine, aureofungin, aviglycine, aviglycine hydrochloride, azaconazole, azadirachtin, azafenidin, azamethiphos, azimsulfuron, azinphos-ethyl, azinphos-methyl, aziprotryne, azithiram, azobenzene, azocyclotin, azothoate, azoxystrobin, bachmedesh, barban, barium hexafluorosilicate, barium polysulfide, barthrin, BCPC, beflubutamid, benalaxyl, benalaxyl-M, benazolin, benazolin-dimethylammonium, benazolin-ethyl, benazolin-potassium, bencarbazone, benclothiaz, bendiocarb, benfluralin, benfuracarb, benfuresate, benodanil, benomyl, benoxacor, benoxafos, benquinox, bensulfuron, bensulfuron-methyl, bensulide, bensultap, bentaluron, bentazone, bentazone-sodium, benthiavalicarb, benthiavalicarb-isopropyl, benthiazole, bentranil, benzadox, benzadox-ammonium, benzalkonium chloride, benzamacril, benzamacril-isobutyl, benzamorf, benzfendizone, benzipram, benzobicyclon, benzofenap, benzofluor, benzohydroxamic acid, benzoximate, benzoylprop, benzoylprop-ethyl, benzthiazuron, benzyl benzoate, benzyladenine, berberine, berberine chloride, *beta*-cyfluthrin, *beta*-cypermethrin, bethoxazin, bicyclopyrone, bifenazate, bifenox, bifenthrin, bifujunzhi, bilanafos, bilanafos-sodium, binapacryl, bingqingxiao, bioallethrin, bioethanomethrin, biopermethrin, bioresmethrin, biphenyl, bisazir, bismerthiazol, bispyribac, bispyribac-sodium, bistrifluron, bitertanol, bithionol, bixafen, blasticidin-S, borax, Bordeaux mixture, boric acid, boscalid, brassinolide, brassinolide-ethyl, brevicomin, brodifacoum, brofenvalerate, brofluthrinate, bromacil, bromacil-lithium, bromacil-sodium, bromadiolone, bromethalin, bromethrin, bromfenvinfos, bromoacetamide, bromobonil, bromobutide, bromocyclen, bromo-DDT, bromofenoxim, bromophos, bromophos-ethyl, bromopropylate, bromothalonil, bromoxynil, bromoxynil butyrate, bromoxynil heptanoate, bromoxynil octanoate, bromoxynil-potassium, brompyrazon, bromuconazole, bronopol, bucarpolate,

2013361540 12 Dec 2016

bufencarb, buminafos, bupirimate, buprofezin, Burgundy mixture, busulfan, butacarb, butachlor, butafenacil, butamifos, butathiofos, butenachlor, butethrin, buthidazole, buthiobate, buthiuron, butocarboxim, butonate, butopyronoxyl, butoxycarboxim, butralin, butroxydim, buturon, butylamine, butylate, cacodylic acid, cadusafos, cafenstrole, calcium arsenate, calcium chlorate, calcium cyanamide, calcium polysulfide, calvinphos, cambendichlor, camphechlor, camphor, captafol, captan, carbamorph, carbanolate, carbaryl, carbasulam, carbendazim, carbendazim benzenesulfonate, carbendazim sulfite, carbetamide, carbofuran, carbon disulfide, carbon tetrachloride, carbophenothion, carbosulfan, carboxazole, carboxide, carboxin, carfentrazone, carfentrazone-ethyl, carpropamid, cartap, cartap hydrochloride, carvacrol, carvone, CDEA, cellocidin, CEPC, ceralure, Cheshunt mixture, chinomethionat, chitosan, chlobenthiazone, chlomethoxyfen, chloralose, chloramben, chloramben-ammonium, chloramben-diolamine, chloramben-methyl, chloramben-methylammonium, chloramben-sodium, chloramine phosphorus, chloramphenicol, chloraniformethan, chloranil, chloranocryl, chlorantraniliprole, chlorazifop, chlorazifop-propargyl, chlorazine, chlorbenside, chlorbenzuron, chlorbicyclen, chlorbromuron, chlorbufam, chlordane, chlordecone, chlordimeform, chlordimeform hydrochloride, chlorempenthrin, chlorethoxyfos, chloreturon, chlorfenac, chlorfenac-ammonium, chlorfenac-sodium, chlorfenapyr, chlorfenazole, chlorfenethol, chlorfenprop, chlorfenson, chlorfensulphide, chlorfenvinphos, chlorfluazuron, chlorflurazole, chlorfluren, chlorfluren-methyl, chlorflurenol, chlorflurenol-methyl, chloridazon, chlorimuron, chlorimuron-ethyl, chlormephos, chlormequat, chlormequat chloride, chlornidine, chlornitrofen, chlorobenzilate, chlorodinitronaphthalenes, chloroform, chloromebuform, chloromethiuron, chloroneb, chlorophacinone, chlorophacinone-sodium, chloropicrin, chloropon, chloropropylate, chlorothalonil, chlorotoluron, chloroxuron, chloroxynil, chlorphonium, chlorphonium chloride, chlorphoxim, chlorprazophos, chlorprocarb, chlorpropham, chlorpyrifos, chlorpyrifos-methyl, chlorquinox, chlorsulfuron, chlorthal, chlorthal-dimethyl, chlorthal-monomethyl, chlorthiamid, chlorthiophos, chlozolinate, choline chloride, chromafenozide, cinerin I, cinerin II, cinerins, cinidon-ethyl, cinmethylin, cinosulfuron, ciobutide, cisanilide, cismethrin, clethodim, climbazole, cliodinate, clodinafop, clodinafop-propargyl, cloethocarb, clofencet, clofencet-potassium, clofentezine, clofibric acid, clofop, clofop-isobutyl, clomazone, clomeprop, cloprop, cloproxydim, clopyralid, clopyralid-methyl, clopyralid-olamine, clopyralid-potassium, clopyralid-tris(2-hydroxypropyl)ammonium, cloquintocet, cloquintocet-mexyl, cloransulam, cloransulam-methyl, closantel, clothianidin, clotrimazole, cloxyfonac, cloxyfonac-sodium, CMA,

2013361540 12 Dec 2016

codlelure, colophonate, copper acetate, copper acetoarsenite, copper arsenate, copper carbonate, basic, copper hydroxide, copper naphthenate, copper oleate, copper oxychloride, copper silicate, copper sulfate, copper zinc chromate, coumachlor, coumafuryl, coumaphos, coumatetralyl, coumithoate, coumoxystrobin, CPMC, CPMF, CPPC, credazine, cresol, crimidine, crotamiton, crotoxyphos, crufomate, cryolite, cue-lure, cufraneb, cumyluron, cuprobam, cuprous oxide, curcumenol, cyanamide, cyanatryn, cyanazine, cyanofenphos, cyanophos, cyanthoate, cyantraniliprole, cyazofamid, cybutryne, cyclafuramid, cyclanilide, cyclethrin, cycloate, cycloheximide, cycloprate, cycloprothrin, cyclosulfamuron, cycloxaprid, cycloxydim, cycluron, cyenopyrafen, cyflufenamid, cyflumetofen, cyfluthrin, cyhalofop, cyhalofop-butyl, cyhalothrin, cyhexatin, cymiazole, cymiazole hydrochloride, cymoxanil, cyometrinil, cypendazole, cypermethrin, cyperquat, cyperquat chloride, cyphenothrin, cyprazine, cyprazole, cyproconazole, cyprodinil, cyprofuram, cypromid, cyprosulfamide, cyromazine, cythioate, daimuron, dalapon, dalapon-calcium, dalapon-magnesium, dalapon-sodium, daminozide, dayoutong, dazomet, dazomet-sodium, DBCP, *d*-camphor, DCIP, DCPTA, DDT, debacarb, decafentin, decarbofuran, dehydroacetic acid, delachlor, deltamethrin, demephion, demephion-O, demephion-S, demeton, demeton-methyl, demeton-O, demeton-O-methyl, demeton-S, demeton-S-methyl, demeton-S-methylsulphon, desmedipham, desmetryn, *d*-fanshiluquebingjuzhi, diafenthiuron, dialifos, di-allate, diamidafos, diatomaceous earth, diazinon, dibutyl phthalate, dibutyl succinate, dicamba, dicamba-diglycolamine, dicamba-dimethylammonium, dicamba-diolamine, dicamba-isopropylammonium, dicamba-methyl, dicamba-olamine, dicamba-potassium, dicamba-sodium, dicamba-trolamine, dicapthon, dichlobenil, dichlofenthion, dichlofluanid, dichlone, dichloralurea, dichlorbenzuron, dichlorflurenol, dichlorflurenol-methyl, dichlormate, dichlormid, dichlorophen, dichlorprop, dichlorprop-2-ethylhexyl, dichlorprop-butotyl, dichlorprop-dimethylammonium, dichlorprop-ethylammonium, dichlorprop-isoctyl, dichlorprop-methyl, dichlorprop-P, dichlorprop-P-2-ethylhexyl, dichlorprop-P-dimethylammonium, dichlorprop-potassium, dichlorprop-sodium, dichlorvos, dichlozoline, diclobutrazol, diclocymet, diclofop, diclofop-methyl, diclomezine, diclomezine-sodium, dicloran, diclosulam, dicofol, dicoumarol, dicresyl, dicrotophos, dicyclanil, dicyclonon, dieldrin, dienochlor, diethamquat, diethamquat dichloride, diethatyl, diethatyl-ethyl, diethofencarb, dietholate, diethyl pyrocarbonate, diethyltoluamide, difenacoum, difenoconazole, difenopenten, difenopenten-ethyl, difenoxuron, difenzoquat, difenzoquat metilsulfate, difethialone, diflovidazin, diflubenzuron, diflufenican, diflufenzopyr, diflufenzopyr-sodium, diflumetorim, dikegulac, dikegulac-sodium, dilor, dimatif,

2013361540 12 Dec 2016

dimefluthrin, dimefox, dimefuron, dimepiperate, dimetachlone, dimetan, dimethacarb, dimethachlor, dimethametryn, dimethenamid, dimethenamid-P, dimethipin, dimethirimol, dimethoate, dimethomorph, dimethrin, dimethyl carbate, dimethyl phthalate, dimethylvinphos, dimetilan, dimexano, dimidazon, dimoxystrobin, dinex, dinex-diclexine, dingjunezuo, diniconazole, diniconazole-M, dinitramine, dinobuton, dinocap, dinocap-4, dinocap-6, dinocton, dinofenate, dinopenton, dinoprop, dinosam, dinoseb, dinoseb acetate, dinoseb-ammonium, dinoseb-diolamine, dinoseb-sodium, dinoseb-trolamine, dinosulfon, dinotefuran, dinoterb, dinoterb acetate, dinoterbon, diofenolan, dioxabenzofos, dioxacarb, dioxathion, diphacinone, diphacinone-sodium, diphenamid, diphenyl sulfone, diphenylamine, dipropalin, dipropetryn, dipyrithione, diquat, diquat dibromide, disparlure, disul, disulfiram, disulfoton, disul-sodium, ditalimfos, dithianon, dithicrofos, dithioether, dithiopyr, diuron, d-limonene, DMPA, DNOC, DNOC-ammonium, DNOC-potassium, DNOC-sodium, dodemorph, dodemorph acetate, dodemorph benzoate, dodicin, dodicin hydrochloride, dodicin-sodium, dodine, dofenapyn, dominicalure, doramectin, drazoxolon, DSMA, dufulin, EBEP, EBP, ecdysterone, edifenphos, eglinazine, eglinazine-ethyl, emamectin, emamectin benzoate, EMPC, empenthrin, endosulfan, endothal, endothal-diammonium, endothal-dipotassium, endothal-disodium, endothion, endrin, enestroburin, EPN, epocholeone, epofenonane, epoxiconazole, eprinomectin, epronaz, EPTC, erbon, ergocalciferol, erlujixiancaoan, esdépalléthrine, esfenvalerate, esprocarb, etacelasil, etaconazole, etaphos, etem, ethaboxam, ethachlor, ethalfluralin, ethametsulfuron, ethametsulfuron-methyl, ethaprochlor, ethephon, ethidimuron, ethiofencarb, ethiolate, ethion, ethiozin, ethiprole, ethirimol, ethoate-methyl, ethofumesate, ethohexadiol, ethoprophos, ethoxyfen, ethoxyfen-ethyl, ethoxyquin, ethoxysulfuron, ethychlozate, ethyl formate, ethyl α-naphthaleneacetate, ethyl-DDD, ethylene, ethylene dibromide, ethylene dichloride, ethylene oxide, ethylicin, ethylmercury 2,3-dihydroxypropyl mercaptide, ethylmercury acetate, ethylmercury bromide, ethylmercury chloride, ethylmercury phosphate, etinofen, etnipromid, etobenzanid, etofenprox, etoxazole, etridiazole, etrimfos, eugenol, EXD, famoxadone, famphur, fenamidone, fenaminosulf, fenamiphos, fenapanil, fenarimol, fenasulam, fenazaflor, fenazaquin, fenbuconazole, fenbutatin oxide, fenchlorazole, fenchlorazole-ethyl, fenchlorphos, fenclorim, fenethacarb, fenfluthrin, fenfuram, fenhexamid, fenitropan, fenitrothion, fenjuntong, fenobucarb, fenoprop, fenoprop-3-butoxypropyl, fenoprop-butometyl, fenoprop-butotyl, fenoprop-butyl, fenoprop-isoctyl, fenoprop-methyl, fenoprop-potassium, fenothiocarb, fenoxacrim, fenoxanil, fenoxaprop, fenoxaprop-ethyl, fenoxaprop-P, fenoxaprop-P-ethyl, fenoxasulfone, fenoxycarb, fenpiclonil, fenpirithrin, fenpropathrin,

2013361540 12 Dec 2016

fenpropidin, fenpropimorph, fenpyrazamine, fenpyroximate, fenridazon, fenridazon-potassium, fenridazon-propyl, fenson, fensulfothion, fenteracol, fenthiaprop, fenthiaprop-ethyl, fenthion, fenthion-ethyl, fentin, fentin acetate, fentin chloride, fentin hydroxide, fentrazamide, fentrifanil, fenuron, fenuron TCA, fenvalerate, ferbam, ferimzone, ferrous sulfate, fipronil, flamprop, flamprop-isopropyl, flamprop-M, flamprop-methyl, flamprop-M-isopropyl, flamprop-M-methyl, flazasulfuron, flocoumafen, flometoquin, flonicamid, florasulam, fluacrypyrim, fluazifop, fluazifop-butyl, fluazifop-methyl, fluazifop-P, fluazifop-P-butyl, fluazinam, fluazolate, fluazuron, flubendiamide, flubenzimine, flucarbazone, flucarbazone-sodium, flucetosulfuron, fluchloralin, flucofuron, flucycloxuron, flucythrinate, fludioxonil, fluenetil, fluensulfone, flufenacet, flufenerim, flufenican, flufenoxuron, flufenprox, flufenpyr, flufenpyr-ethyl, flufiprole, flumethrin, flumetover, flumetralin, flumetsulam, flumezin, flumiclorac, flumiclorac-pentyl, flumioxazin, flumipropyn, flumorph, fluometuron, fluopicolide, fluopyram, fluorbenside, fluoridamid, fluoroacetamide, fluorodifen, fluoroglycofen, fluoroglycofen-ethyl, fluoroimide, fluoromidine, fluoronitrofen, fluothiuron, fluotrimazole, fluoxastrobin, flupoxam, flupropacil, flupropadine, flupropanate, flupropanate-sodium, flupyradifurone, flupyrsulfuron, flupyrsulfuron-methyl, flupyrsulfuron-methyl-sodium, fluquinconazole, flurazole, flurenol, flurenol-butyl, flurenol-methyl, fluridone, flurochloridone, fluroxypyr, fluroxypyr-butometyl, fluroxypyr-meptyl, flurprimidol, flursulamid, flurtamone, flusilazole, flusulfamide, fluthiacet, fluthiacet-methyl, flutianil, flutolanil, flutriafol, fluvalinate, fluxapyroxad, fluxofenim, folpet, fomesafen, fomesafen-sodium, fonofos, foramsulfuron, forchlorfenuron, formaldehyde, formetanate, formetanate hydrochloride, formothion, formparanate, formparanate hydrochloride, fosamine, fosamine-ammonium, fosetyl, fosetyl-aluminium, fosmethilan, fospirate, fosthiazate, fosthietan, frontalin, fuberidazole, fucaojing, fucaomi, funaihecaoling, fuphenthiourea, furalane, furalaxyl, furamethrin, furametpyr, furathiocarb, furcarbanil, furconazole, furconazole-cis, furethrin, furfural, furilazole, furmecyclox, furophanate, furyloxyfen, *gamma*-cyhalothrin, *gamma*-HCH, genit, gibberellic acid, gibberellins, gliftor, glufosinate, glufosinate-ammonium, glufosinate-P, glufosinate-P-ammonium, glufosinate-P-sodium, glyodin, glyoxime, glyphosate, glyphosate-diammonium, glyphosate-dimethylammonium, glyphosate-isopropylammonium, glyphosate-monoammonium, glyphosate-potassium, glyphosate-sesquisodium, glyphosate-trimesium, glyphosine, gossyplure, grandlure, griseofulvin, guazatine, guazatine acetates, halacrinate, halfenprox, halofenozide, halosafen, halosulfuron, halosulfuron-methyl, haloxydine, haloxyfop, haloxyfop-etotyl, haloxyfop-methyl, haloxyfop-P, haloxyfop-P-etotyl, haloxyfop-P-methyl, haloxyfop-sodium, HCH,

2013361540 12 Dec 2016

hemel, hempa, HEOD, heptachlor, heptenophos, heptopargil, heterophos, hexachloroacetone, hexachlorobenzene, hexachlorobutadiene, hexachlorophene, hexaconazole, hexaflumuron, hexaflurate, hexalure, hexamide, hexazinone, hexylthiofos, hexythiazox, HHDN, holosulf, huancaiwo, huangcaoling, huanjunzuo, hydramethylnon, hydrargaphen, hydrated lime, hydrogen cyanide, hydroprene, hymexazol, hyquincarb, IAA, IBA, icaridin, imazalil, imazalil nitrate, imazalil sulfate, imazamethabenz, imazamethabenz-methyl, imazamox, imazamox-ammonium, imazapic, imazapic-ammonium, imazapyr, imazapyr-isopropylammonium, imazaquin, imazaquin-ammonium, imazaquin-methyl, imazaquin-sodium, imazethapyr, imazethapyr-ammonium, imazosulfuron, imibenconazole, imicyafos, imidacloprid, imidaclothiz, iminoctadine, iminoctadine triacetate, iminoctadine trialbesilate, imiprothrin, inabenfide, indanofan, indaziflam, indoxacarb, inezin, iodobonil, iodocarb, iodomethane, iodosulfuron, iodosulfuron-methyl, iodosulfuron-methyl-sodium, iofensulfuron, iofensulfuron-sodium, ioxynil, ioxynil octanoate, ioxynil-lithium, ioxynil-sodium, ipazine, ipconazole, ipfencarbazone, iprobenfos, iprodione, iprovalicarb, iprymidam, ipsdienol, ipsenol, IPSP, isamidofos, isazofos, isobenzan, isocarbamid, isocarbophos, isocil, isodrin, isofenphos, isofenphos-methyl, isolan, isomethiozin, isonoruron, isopolinate, isoprocarb, isopropalin, isoprothiolane, isoproturon, isopyrazam, isopyrimol, isothioate, isotianil, isouron, isovaledione, isoxaben, isoxachlortole, isoxadifen, isoxadifen-ethyl, isoxaflutole, isoxapyrifop, isoxathion, ivermectin, izopamfos, japonilure, japothrins, jasmolin I, jasmolin II, jasmonic acid, jiahuangchongzong, jiajizengxiaolin, jiaxiangjunzhi, jiecaowan, jiecaoxi, jodfenphos, juvenile hormone I, juvenile hormone II, juvenile hormone III, kadethrin, karbutilate, karetazan, karetazan-potassium, kasugamycin, kasugamycin hydrochloride, kejunlin, kelevan, ketospiradox, ketospiradox-potassium, kinetin, kinoprene, kresoxim-methyl, kuicaoxi, lactofen, lambda-cyhalothrin, latilure, lead arsenate, lenacil, lepimectin, leptophos, lindane, lineatin, linuron, lirimfos, litlure, looplure, lufenuron, lvdingjunzhi, lvxiancaolin, lythidathion, MAA, malathion, maleic hydrazide, malonoben, maltodextrin, MAMA, mancopper, mancozeb, mandipropamid, maneb, matrine, mazidox, MCPA, MCPA-2-ethylhexyl, MCPA-butotyl, MCPA-butyl, MCPA-dimethylammonium, MCPA-diolamine, MCPA-ethyl, MCPA-isobutyl, MCPA-isoctyl, MCPA-isopropyl, MCPA-methyl, MCPA-olamine, MCPA-potassium, MCPA-sodium, MCPA-thioethyl, MCPA-trolamine, MCPB, MCPB-ethyl, MCPB-methyl, MCPB-sodium, mebenil, mecarbam, mecarbinzid, mecarphon, mecoprop, mecoprop-2-ethylhexyl, mecoprop-dimethylammonium, mecoprop-diolamine, mecoprop-ethadyl, mecoprop-isoctyl, mecoprop-methyl, mecoprop-P, mecoprop-P-2-ethylhexyl, mecoprop-P-dimethylammonium, mecoprop-P-isobutyl, mecoprop-potassium,

2013361540 12 Dec 2016

mecoprop-P-potassium, mecoprop-sodium, mecoprop-trolamine, medimeform, medinoterb, medinoterb acetate, medlure, mefenacet, mefenpyr, mefenpyr-diethyl, mefluidide, mefluidide-diolamine, mefluidide-potassium, megatomoic acid, menazon, mepanipyrim, meperfluthrin, mephenate, mephosfolan, mepiquat, mepiquat chloride, mepiquat pentaborate, mepronil, meptyldinocap, mercuric chloride, mercuric oxide, mercurous chloride, merphos, mesoprazine, mesosulfuron, mesosulfuron-methyl, mesotrione, mesulfen, mesulfenfos, metaflumizone, metalaxyl, metalaxyl-M, metaldehyde, metam, metam-ammonium, metamifop, metamitron, metam-potassium, metam-sodium, metazachlor, metazosulfuron, metazoxolon, metconazole, metepa, metflurazon, methabenzthiazuron, methacrifos, methalpropalin, methamidophos, methasulfocarb, methazole, methfuroxam, methidathion, methiobencarb, methiocarb, methiopyrisulfuron, methiotepa, methiozolin, methiuron, methocrotophos, methometon, methomyl, methoprene, methoprotryne, methoquin-butyl, methothrin, methoxychlor, methoxyfenozide, methoxyphenone, methyl apholate, methyl bromide, methyl eugenol, methyl iodide, methyl isothiocyanate, methylacetophos, methylchloroform, methyldymron, methylene chloride, methylmercury benzoate, methylmercury dicyandiamide, methylmercury pentachlorophenoxide, methylneodecanamide, metiram, metobenzuron, metobromuron, metofluthrin, metolachlor, metolcarb, metominostrobin, metosulam, metoxadiazone, metoxuron, metrafenone, metribuzin, metsulfovax, metsulfuron, metsulfuron-methyl, mevinphos, mexacarbate, mieshuan, milbemectin, milbemycin oxime, milneb, mipafox, mirex, MNAF, moguchun, molinate, molosultap, monalide, monisouron, monochloroacetic acid, monocrotophos, monolinuron, monosulfuron, monosulfuron-ester, monuron, monuron TCA, morfamquat, morfamquat dichloride, moroxydine, moroxydine hydrochloride, morphothion, morzid, moxidectin, MSMA, muscalure, myclobutanil, myclozolin, N-(ethylmercury)-p-toluenesulphonanilide, nabam, naftalofos, naled, naphthalene, naphthaleneacetamide, naphthalic anhydride, naphthoxyacetic acids, naproanilide, napropamide, naptalam, naptalam-sodium, natamycin, neburon, niclosamide, niclosamide-olamine, nicosulfuron, nicotine, nifluridide, nipyraclofen, nitenpyram, nithiazine, nitralin, nitrapyrin, nitrilacarb, nitrofen, nitrofluorfen, nitrostyrene, nitrothal-isopropyl, norbormide, norflurazon, nornicotine, noruron, novaluron, noviflumuron, nuarimol, OCH, octachlorodipropyl ether, octhilinone, ofurace, omethoate, orbencarb, orfralure, ortho-dichlorobenzene, orthosulfamuron, oryctalure, orysastrobin, oryzalin, osthol, ostramone, oxabetrinil, oxadiargyl, oxadiazon, oxadixyl, oxamate, oxamyl, oxapyrazon, oxapyrazon-dimolamine, oxapyrazon-sodium, oxasulfuron, oxaziclomefone, oxine-copper, oxolinic acid,

2013361540 12 Dec 2016

oxpoconazole, oxpoconazole fumarate, oxycarboxin, oxydemeton-methyl, oxydeprofos, oxydisulfoton, oxyfluorfen, oxymatrine, oxytetracycline, oxytetracycline hydrochloride, paclobutrazol, paichongding, para-dichlorobenzene, parafluron, paraquat, paraquat dichloride, paraquat dimetilsulfate, parathion, parathion-methyl, parinol, pebulate, pefurazoate, pelargonic acid, penconazole, pencycuron, pendimethalin, penflufen, penfluron, penoxsulam, pentachlorophenol, pentanochlor, penthiopyrad, pentmethrin, pentoxazone, perfluidone, permethrin, pethoxamid, phenamacril, phenazine oxide, phenisopham, phenkapton, phenmedipham, phenmedipham-ethyl, phenobenzuron, phenothrin, phenproxide, phenthoate, phenylmercuriurea, phenylmercury acetate, phenylmercury chloride, phenylmercury derivative of pyrocatechol, phenylmercury nitrate, phenylmercury salicylate, phorate, phosacetim, phosalone, phosdiphen, phosfolan, phosfolan-methyl, phosglycin, phosmet, phosnichlor, phosphamidon, phosphine, phosphocarb, phosphorus, phostin, phoxim, phoxim-methyl, phthalide, picloram, picloram-2-ethylhexyl, picloram-isoctyl, picloram-methyl, picloram-olamine, picloram-potassium, picloram-triethylammonium, picloram-tris(2-hydroxypropyl)ammonium, picolinafen, picoxystrobin, pindone, pindone-sodium, pinoxaden, piperalin, piperonyl butoxide, piperonyl cyclonene, piperophos, piproctanyl, piproctanyl bromide, piprotal, pirimetaphos, pirimicarb, pirimioxyphos, pirimiphos-ethyl, pirimiphos-methyl, plifenate, polycarbamate, polyoxins, polyoxorim, polyoxorim-zinc, polythialan, potassium arsenite, potassium azide, potassium cyanate, potassium gibberellate, potassium naphthenate, potassium polysulfide, potassium thiocyanate, potassium α-naphthaleneacetate, *pp'*-DDT, prallethrin, precocene I, precocene II, precocene III, pretilachlor, primidophos, primisulfuron, primisulfuron-methyl, probenazole, prochloraz, prochloraz-manganese, proclonol, procyazine, procymidone, prodiamine, profenofos, profluazol, profluralin, profluthrin, profoxydim, proglinazine, proglinazine-ethyl, prohexadione, prohexadione-calcium, prohydrojasmon, promacyl, promecarb, prometon, prometryn, promurit, propachlor, propamidine, propamidine dihydrochloride, propamocarb, propamocarb hydrochloride, propanil, propaphos, propaquizafop, propargite, proparthrin, propazine, propetamphos, propham, propiconazole, propineb, propisochlor, propoxur, propoxycarbazone, propoxycarbazone-sodium, propyl isome, propyrisulfuron, propyzamide, proquinazid, prosuler, prosulfalin, prosulfocarb, prosulfuron, prothidathion, prothiocarb, prothiocarb hydrochloride, prothioconazole, prothiofos, prothoate, protrifenbute, proxan, proxan-sodium, prynachlor, pydanon, pymetrozine, pyracarbolid, pyraclofos, pyraclonil, pyraclostrobin, pyraflufen, pyraflufen-ethyl, pyrafluprole, pyramat, pyrametostrobin, pyraoxystrobin, pyrasulfotole, pyrazolynate, pyrazophos, pyrazosulfuron, pyrazosulfuron-ethyl, pyrazothion,

2013361540 12 Dec 2016

pyrazoxyfen, pyresmethrin, pyrethrin I, pyrethrin II, pyrethrins, pyribambenz-isopropyl, pyribambenz-propyl, pyribencarb, pyribenzoxim, pyributicarb, pyriclor, pyridaben, pyridafol, pyridalyl, pyridaphenthion, pyridate, pyridinitril, pyrifenox, pyrifluquinazon, pyriftalid, pyrimethanil, pyrimidifen, pyriminobac, pyriminobac-methyl, pyrimisulfan, pyrimitate, pyrinuron, pyriofenone, pyriprole, pyripropanol, pyriproxyfen, pyrithiobac, pyrithiobac-sodium, pyrolan, pyroquilon, pyroxasulfone, pyroxsulam, pyroxychlor, pyroxyfur, quassia, quinacetol, quinacetol sulfate, quinalphos, quinalphos-methyl, quinazamid, quinclorac, quinconazole, quinmerac, quinoclamine, quinonamid, quinothion, quinoxyfen, quintiofos, quintozene, quizalofop, quizalofop-ethyl, quizalofop-P, quizalofop-P-ethyl, quizalofop-P-tefuryl, quwenzhi, quyingding, rabenzazole, rafoxanide, rebemide, resmethrin, rhodethanil, rhodojaponin-III, ribavirin, rimsulfuron, rotenone, ryania, saflufenacil, saijunmao, saisentong, salicylanilide, sanguinarine, santonin, schradan, scilliroside, sebuthylazine, secbumeton, sedaxane, selamectin, semiamitraz, semiamitraz chloride, sesamex, sesamolin, sethoxydim, shuangjiaancaolin, siduron, siglure, silafluofen, silatrane, silica gel, silthiofam, simazine, simeconazole, simeton, simetryn, sintofen, SMA, S-metolachlor, sodium arsenite, sodium azide, sodium chlorate, sodium fluoride, sodium fluoroacetate, sodium hexafluorosilicate, sodium naphthenate, sodium orthophenylphenoxide, sodium pentachlorophenoxide, sodium polysulfide, sodium thiocyanate, sodium α-naphthaleneacetate, sophamide, spinetoram, spinosad, spirodiclofen, spiromesifen, spirotetramat, spiroxamine, streptomycin, streptomycin sesquisulfate, strychnine, sulcatol, sulcofuron, sulcofuron-sodium, sulcotrione, sulfallate, sulfentrazone, sulfiram, sulfluramid, sulfometuron, sulfometuron-methyl, sulfosulfuron, sulfotep, sulfoxaflor, sulfoxide, sulfoxime, sulfur, sulfuric acid, sulfuryl fluoride, sulglycapin, sulprofos, sultropen, swep, *tau*-fluvalinate, tavron, tazimcarb, TCA, TCA-ammonium, TCA-calcium, TCA-ethadyl, TCA-magnesium, TCA-sodium, TDE, tebuconazole, tebufenozide, tebufenpyrad, tebufloquin, tebupirimfos, tebutam, tebuthiuron, tecloftalam, tecnazene, tecoram, teflubenzuron, tefluthrin, tefuryltrione, tembotrione, temephos, tepa, TEPP, tepraloxydim, terallethrin, terbacil, terbucarb, terbuchlor, terbufos, terbumeton, terbuthylazine, terbutryn, tetcyclacis, tetrachloroethane, tetrachlorvinphos, tetraconazole, tetradifon, tetrafluron, tetramethrin, tetramethylfluthrin, tetramine, tetranactin, tetrasul, thallium sulfate, thenylchlor, theta-cypermethrin, thiabendazole, thiacloprid, thiadifluor, thiamethoxam, thiapronil, thiazafluron, thiazopyr, thicrofos, thicyofen, thidiazimin, thidiazuron, thiencarbazone, thiencarbazone-methyl, thifensulfuron, thifensulfuron-methyl, thifluzamide, thiobencarb, thiocarboxime, thiochlorfenphim, thiocyclam, thiocyclam hydrochloride, thiocyclam oxalate, thiodiazole-copper, thiodicarb, thiofanox, thiofluoximate,

2013361540 12 Dec 2016

thiohempa, thiomersal, thiometon, thionazin, thiophanate, thiophanate-methyl, thioquinox, thiosemicarbazide, thiosultap, thiosultap-diammonium, thiosultap-disodium, thiosultap-monosodium, thiotepa, thiram, thuringiensin, tiadinil, tiaojiean, tiocarbazil, tioclorim, tioxymid, tirpate, tolclofos-methyl, tolfenpyrad, tolylfluanid, tolylmercury acetate, topramezone, tralkoxydim, tralocythrin, tralomethrin, tralopyril, transfluthrin, transpermethrin, tretamine, triacontanol, triadimefon, triadimenol, triafamone, tri-allate, triamiphos, triapenthenol, triarathene, triarimol, triasulfuron, triazamate, triazbutil, triaziflam, triazophos, triazoxide, tribenuron, tribenuron-methyl, tribufos, tributyltin oxide, tricamba, trichlamide, trichlorfon, trichlormetaphos-3, trichloronat, triclopyr, triclopyr-butotyl, triclopyr-ethyl, triclopyr-triethylammonium, tricyclazole, tridemorph, tridiphane, trietazine, trifenmorph, trifenofos, trifloxystrobin, trifloxysulfuron, trifloxysulfuron-sodium, triflumizole, triflumuron, trifluralin, triflusulfuron, triflusulfuron-methyl, trifop, trifop-methyl, trifopsime, triforine, trihydroxytriazine, trimedlure, trimethacarb, trimeturon, trinexapac, trinexapac-ethyl, triprene, tripropindan, triptolide, tritac, triticonazole, tritosulfuron, trunc-call, uniconazole, uniconazole-P, urbacide, uredepa, valerate, validamycin, valifenalate, valone, vamidothion, vangard, vaniliprole, vernolate, vinclozolin, warfarin, warfarin-potassium, warfarin-sodium, xiaochongliulin, xinjunan, xiwojunan, XMC, xylachlor, xylenols, xylylcarb, yishijing, zarilamid, zeatin, zengxiaoan, zeta-cypermethrin, zinc naphthenate, zinc phosphide, zinc thiazole, zineb, ziram, zolaprofos, zoxamide, zuomihuanglong, α-chlorohydrin, α-ecdysone, α-multistriatin, and α-naphthaleneacetic acid.

6. A composition according to any one of the previous claims, further comprising an agriculturally acceptable carrier, a biopesticide; or one or more of the following compounds:

(a) 3-(4-chloro-2,6-dimethylphenyl)-4-hydroxy-8-oxa-1-azaspiro[4,5]dec-3-en-2-one;

(b) 3-(4'-chloro-2,4-dimethyl[1,1'-biphenyl]-3-yl)-4-hydroxy-8-oxa-1-azaspiro[4,5]dec-3-en-2-one;

(c) 4-[[(6-chloro-3-pyridinyl)methyl]methylamino]-2(5*H*)-furanone;

(d) 4-[[(6-chloro-3-pyridinyl)methyl]cyclopropylamino]-2(5*H*)-furanone;

(e) 3-chloro-*N*2-[(1*S*)-1-methyl-2-(methylsulfonyl)ethyl]-*N*1-[2-methyl-4-[1,2,2,2-tetrafluoro-1-(trifluoromethyl)ethyl]phenyl]-1,2-benzenedicarboxamide;

(f) 2-cyano-*N*-ethyl-4-fluoro-3-methoxy-benenesulfonamide;

(g) 2-cyano-*N*-ethyl-3-methoxy-benzenesulfonamide;

(h) 2-cyano-3-difluoromethoxy-*N*-ethyl-4-fluoro-benzenesulfonamide;

(i) 2-cyano-3-fluoromethoxy-*N*-ethyl-benzenesulfonamide;

2013361540 12 Dec 2016

(j) 2-cyano-6-fluoro-3-methoxy-*N,N*-dimethyl-benzenesulfonamide;

(k) 2-cyano-*N*-ethyl-6-fluoro-3-methoxy-*N*-methyl-benzenesulfonamide;

(l) 2-cyano-3-difluoromethoxy-*N,N*-dimethylbenzenesulfon-amide;

(m) 3-(difluoromethyl)-*N*-[2-(3,3-dimethylbutyl)phenyl]-1-methyl-1*H*-pyrazole-4-carboxamide;

(n) *N*-ethyl-2,2-dimethylpropionamide-2-(2,6-dichloro-α,α,α-trifluoro-*p*-tolyl) hydrazone;

(o) *N*-ethyl-2,2-dichloro-1-methylcyclopropane-carboxamide-2-(2,6-dichloro-α,α,α-trifluoro-*p*-tolyl) hydrazone nicotine;

(p) O-{(E-)-[2-(4-chloro-phenyl)-2-cyano-1-(2-trifluoromethylphenyl)-vinyl]} S-methyl thiocarbonate;

(q) (E)-N1-[(2-chloro-1,3-thiazol-5-ylmethyl)]-N2-cyano-N1-methylacetamidine;

(r) 1-(6-chloropyridin-3-ylmethyl)-7-methyl-8-nitro-1,2,3,5,6,7-hexahydro-imidazo[1,2-a]pyridin-5-ol;

(s) 4-[4-chlorophenyl-(2-butylidine-hydrazono)methyl)]phenyl mesylate; and

(t) N-Ethyl-2,2-dichloro-1-methylcyclopropanecarboxamide-2-(2,6-dichloro-*alpha,alpha,alpha*-trifluoro-p-tolyl)hydrazone.

7. A composition according to any one of the previous claims, wherein said molecule is in the form of a pesticidally acceptable acid addition salt; or

wherein said molecule is in the form of a salt derivative; or

wherein said molecule is in the form a hydrate; or

wherein said molecule is in the form an ester derivative; or

wherein said molecule is in the form a crystal polymorph; or

wherein said molecule has a ^{2}H in place of ^{1}H; or wherein said molecule has a ^{14}C in place of a ^{12}C.

8. A composition according to any one of the previous claims, further comprising a compound having one or more of the following modes of action: acetylcholinesterase inhibitor; sodium channel modulator; chitin biosynthesis inhibitor; GABA and glutamate-gated chloride channel antagonist; GABA and glutamate-gated chloride channel agonist; acetylcholine receptor agonist; acetylcholine receptor antagonist; MET I inhibitor; Mg-stimulated ATPase inhibitor; nicotinic acetylcholine receptor; Midgut membrane disrupter; oxidative phosphorylation disrupter, and ryanodine receptor (RyRs);

or

further comprising a seed; or

2013361540 12 Dec 2016

further comprising a seed that has been genetically modified to express one or more specialized traits;

or

wherein said composition is encapsulated inside, or placed on the surface of, a capsule; or

wherein said composition is encapsulated inside, or placed on the surface of, a capsule, wherein said capsule has a diameter of about 100-900 nanometers or about 10-900 microns.

9. A process comprising applying a composition according to any one of the previous claims, to an area to control a pest, in an amount sufficient to control such pest.

10. A process according to claim 9, wherein said pest is selected from beetles, earwigs, cockroaches, flies, aphids, scales, whiteflies, leafhoppers, ants, wasps, termites, moths, butterflies, lice, grasshoppers, locusts, crickets, fleas, thrips, bristletails, mites, ticks, nematodes, and symphylans; or wherein

said pest is from the Phyla Nematoda or Arthropoda; or

wherein said pest is from the Subphyla Chelicerata, Myriapoda, or Hexapoda; or wherein said pest is from the Class of Arachnida, Symphyla, or Insecta; or

said pest is from the Order Anoplura, Order Coleoptera, Order Dermaptera, Order Blattaria, Order Diptera, Order Hemiptera, Order Hymenoptera, Order Isoptera, Order Lepidoptera, Order Mallophaga, Order Orthoptera, Order Siphonaptera, Order Thysanoptera, Order Thysanura, Order Acarina, or Order Symphyla; or wherein

said pest is Beet Armyworm (BAW), Corn Earworm (CEW), or Green Peach Aphid (GPA).

11. A process according to claim 9 or 10, wherein said amount is from about 0.01 grams per hectare to about 5000 grams per hectare; or

from about 0.1 grams per hectare to about 500 grams per hectare; or

from about 1 gram per hectare to about 50 grams per hectare.

12. A process according to any one of claims 9 to 11, wherein said area is an area where apples, corn, cotton, soybeans, canola, wheat, rice, sorghum, barley, oats, potatoes, oranges, alfalfa, lettuce, strawberries, tomatoes, peppers, crucifers, pears, tobacco, almonds, sugar beets, or beans, are growing, or the seeds thereof are going to be planted.

13. A process according to any one of claims 9 to 12, further comprising applying said composition to a genetically modified plant that has been genetically modified to express one or more specialized traits.

14. A process according to any one of claims 9 to 13, where said composition further comprises ammonium sulfate.

2013361540 12 Dec 2016

15. A process comprising: orally administering; or topically applying; a composition according to any one of claims 1 to 8, to a non-human animal, to control endoparasites, ectoparasites, or both;

or

a process comprising applying a composition according to any one of claims 1 to 8, to a plant to enhance the plant's health, yield, vigor, quality, or tolerance, at a time when pest activity is low.